



US 20250277227A1

(19) **United States**

(12) **Patent Application Publication**
CIGAN et al.

(10) **Pub. No.: US 2025/0277227 A1**

(43) **Pub. Date: Sep. 4, 2025**

(54) **METHODS FOR PRODUCING A COMPLEX
TRANSGENIC TRAIT LOCUS**

(71) Applicants: **PIONEER HI-BRED
INTERNATIONAL, INC.,**
JOHNSTON, IA (US); **CORTEVA
AGRISCIENCE LLC,**
INDIANAPOLIS, IN (US)

(72) Inventors: **ANDREW MARK CIGAN,**
MADISON, WI (US); **SAVERIO
CARL FALCO,** WILMINGTON, DE
(US); **HUIRONG GAO,** JOHNSTON,
IA (US); **MICHAEL LASSNER,**
PORTLAND, OR (US); **DEREK
JANTZ,** DURHAM, NC (US);
ZHONGSEN LI, HOCKESSIN, DE
(US); **ZHAN-BIN LIU,** CLIVE, IA
(US); **SERGEI SVITASHEV,**
JOHNSTON, IA (US); **JAMES
JEFFERSON SMITH,** DURHAM, NC
(US)

(73) Assignees: **PIONEER HI-BRED
INTERNATIONAL, INC.,**
JOHNSTON, IA (US); **CORTEVA
AGRISCIENCE LLC,**
INDIANAPOLIS, IN (US)

(21) Appl. No.: **19/210,051**

(22) Filed: **May 16, 2025**

Related U.S. Application Data

(63) Continuation of application No. 17/821,528, filed on
Aug. 23, 2022, now Pat. No. 12,338,444, which is a

continuation of application No. 16/925,973, filed on
Jul. 10, 2020, now Pat. No. 11,459,576, which is a
continuation of application No. 16/007,529, filed on
Jun. 13, 2018, now Pat. No. 10,822,610, which is a
continuation of application No. 13/427,138, filed on
Mar. 22, 2012, now Pat. No. 10,030,245.

(60) Provisional application No. 61/499,443, filed on Jun.
21, 2011, provisional application No. 61/466,602,
filed on Mar. 23, 2011.

Publication Classification

(51) **Int. Cl.**
C12N 15/82 (2006.01)
A01H 1/06 (2006.01)
C12N 9/22 (2006.01)
(52) **U.S. Cl.**
CPC ***C12N 15/8202*** (2013.01); ***A01H 1/06***
(2013.01); ***C12N 9/22*** (2013.01); ***C12N***
15/8213 (2013.01)

(57) **ABSTRACT**

Methods for producing in a plant a complex transgenic trait locus comprising at least two altered target sequences in a genomic region of interest are disclosed. The methods involve the use of two or more double-strand-break-inducing agents, each of which can cause a double-strand break in a target sequence in the genomic region of interest which results in an alteration in the target sequence. Also disclosed are complex transgenic trait loci in plants. A complex transgenic trait locus comprises at least two altered target sequences that are genetically linked to a polynucleotide of interest. Plants, plant cells, plant parts, and seeds comprising one or more complex transgenic trait loci are also disclosed.

Specification includes a Sequence Listing.

DNA Double-Strand-Break-Induced DNA Alteration of an Endogenous Target Site




FIG. 1
DNA Double-Strand-Break-Induced DNA Alteration of an Endogenous Target Site




FIG. 1B Nuclease gene



FIG. 1C Donor DNA



FIG. 1D Donor DNA




FIG. 1E Gene targeting




FIG. 1F Target site deletion



FIG. 1G Target site insertion




FIG. 2

Genetic / physical Distance Between Target Sites and Transgene of Interest




FIG. 3

FIG. 3A: Genomic PCR assay to identify target site modification or transgene integration




FIG. 3B: Alignment of sequences of selected TS21 transgenic event (TS 21 target sites underlined, cut sites in bold)

SEQ ID		BssSI
NO:	Deletion/insertion event	
121	WT	TCTTAAAGAAGATACACTCTGTAAATGTTAAATGGCACTGGCACTCTCTGTTGGTGAATTAAAGTCATATATGGTTTAAGATACCTTTTTTTTATAAAGATAGTAGTG
122	Pic11	TCTTAAAGAAGATACACTCTGTAAATGTTAAATGGCACTGGCACTCTCTGTTGGTGAATTAAAGTCATATATGGTTTAAGATACCTTTTTTTTATAAAGATAGTAGTG
123	2637-45	221bp deletion flanking the target site/no insertion.....GATACTTTTTTTTATAAAGATAGTAGTG

FIG. 4

FIG. 4A: Alignment of sequences of TS5 transgenic events (TS5 target sites underlined)

SEQ ID			BssSI
NO:	Deletion/insertion event		
124	WT	TAATGATCAGATTTTTTTTCTCAGACTCACCCTAAGTGCAGAGTACACACGTAAGTCTTAGGTTAAAGTTTATGCCCCCCCCCCCCCCCCCCAAA	
125	P2A6	TAATGATCAGATTTTTTTTCTCAGACTCACCCTAAGTGCAGAGTACACACGTT.....326bp deletion with 42bp filler DNA.....	
126	P1B9	TAATGATCAGATTTTTTTTCTCAGACTCACCCTAAGTGCAGAGTACACAC▼GTAAAGTCTTAGGTTAAAGTTTATGCCCCCCCCCCCCCCCCCCAAA	
127	P2C5	TAATGATCAGATTTTTTTTCTCAGACTCACCCTAAGTGCAGAG.....94bp deletion with 193bp filler DNA.....	

▼ indicates a 161 bp insertion into the target site

FIG. 4B: Alignment of sequences of selected TS14 transgenic event (TS5 target sites underlined)

SEQ ID			BsiWI
NO:	Deletion/insertion event		
128	WT	TAATGATCAGATTTTTTTTCTCAGACTCACCCTAAGTGCAGACGTAAGCAAGTAGCTTTTCTTACCTTCGTATTGACAATCAAAATCGTCCTTTATTTTATT	
129	P4G10	TAATGATCAGATTTTTTTTCTCAGACTCACCCTAAGTGCAGACGTA▼CAAGTAGCTTTTCTTACCTTCGTATTGACAATCAAAATCGTCCTTTATTTTATT	

HO Mega14 target sites are underlined with the cut sites in bold

▼ indicates a 115 bp insertion into the target site

FIG. 5

Gene Integration by Homologous Recombination Enabled by Double-Strand Breaks with Meganuclease Near a Transgenic Event (TE) of Interest




FIG. 6

Location of Target Sites Near a Herbicide-Resistant Transgenic Event in Soybean





FIG. 7

Cluster of Meganuclease Target Sites for Trait Stacking






FIG. 9

FIG. 9A




FIG. 9B




FIG. 10

FIG. 10A




FIG. 10B



FIG. 11A

	70	80	90	100	110	120
SEQIDNO-9.seq	CTCTACCTGGCCGGCTTCGTGGACGGCGACGGCTCCATCATGGCGCAGATCAAGCCGCGAG	120				
SEQIDNO-10.seq	CTCTACCTGGCCGGCTTCGTGGACGGCGACGGCTCCATCATCGCGTCCATCAAGCCGGAG	120				
SEQIDNO-11.seq	CTCTACCTGGCCGGCTTCGTGGACGGCGACGGCTCCATCATGGCGAAGATCAAGCCGGAG	120				
SEQIDNO-12.seq	CTCTACCTGGCCGGCTTCGTGGACGGCGACGGTTCATCATCGCGCAGATCAAGCCGGAG	120				
SEQIDNO-13.seq	CTCTACCTGGCCGGCTTCGTGGACGGCGACGGCTCCATCATCGCGCAGATCAAGCCGGAG	120				
SEQIDNO-14.seq	CTCTACCTGGCCGGCTTCGTGGACGGCGACGGCTCCATCATCGCGCAGATCAAGCCGGAG	120				
SEQIDNO-15.seq	CTCTACCTGGCCGGCTTCGTGGACGGCGACGGCTCCATCATCGCGCAGATCAAGCCGGAG	120				
SEQIDNO-16.seq	CTCTACCTGGCCGGCTTCGTGGACGGCGACGGCTCCATCATCGCGTCCATCAAGCCGGAG	120				
SEQIDNO-80.seq	CTCTACCTGGCCGGCTTCGTGGACGGCGACGGATCCATCATCGCGCAGATCAAGCCGAAC	120				
SEQIDNO-78.seq	CTCTACCTGGCCGGCTTCGTGGACGGCGACGGCTCCATCATCGCGCAGATCAAGCCGAAC	120				
SEQIDNO-79.seq	CTCTACCTGGCCGGCTTCGTGGACGGCGACGGCTCCATCATCGCGCAGATCAAGCCGAAC	120				
SEQIDNO-81.seq	CTCTACCTGGCCGGCTTCGTGGACGGCGACGGCTCCATCATCGCGCAGATCAAGCCGAAC	120				
SEQIDNO-82.seq	CTCTACCTGGCCGGCTTCGTGGACGGCGACGGCTCCATCATCGCGCAGATCAAGCCGAAC	120				
SEQIDNO-83.seq	CTCTACCTGGCCGGCTTCGTGGACGGCGACGGCTCCATCATCGCGCAGATCAAGCCGGAG	120				

	130	140	150	160	170	180
SEQIDNO-9.seq	CAGTCCTGCAAGTTCAAGCACCAGCTCCA-GCTGACCTTCACCGTGACCCAGAAGACGCA	179				
SEQIDNO-10.seq	CAGTCCTACAAGTTCAAGCACCAGCTCCTCCTGACCTTCACCGTGACCCAGAAGACGCA	179				
SEQIDNO-11.seq	CAGTCCTACAAGTTCAAGCACCAGCTCCTCCTGACCTTCACCGTGACCCAGAAGACGCA	179				
SEQIDNO-12.seq	CAGTCCTACAAGTTCAAGCACCAGCTCCTCCTGACCTTCACCGTGACCCAGAAGACGCA	179				
SEQIDNO-13.seq	CAGTCCTACAAGTTCAAGCACCAGCTCCTCCTGACCTTCACCGTGACCCAGAAGACGCA	179				
SEQIDNO-14.seq	CAGTCCTACAAGTTCAAGCACCAGCTCCTCCTGACCTTCACCGTGACCCAGAAGACGCA	179				
SEQIDNO-15.seq	CAGTCCTACAAGTTCAAGCACCAGCTCCTCCTGACCTTCACCGTGACCCAGAAGACGCA	179				
SEQIDNO-16.seq	CAGTCCTACAAGTTCAAGCACCAGCTCCTCCTGACCTTCACCGTGACCCAGAAGACGCA	179				
SEQIDNO-80.seq	CAGTCCTGCAAGTTCAAGCACCAGCTCCTCCTGACCTTCACCGTGACCCAGAAGACGCA	179				
SEQIDNO-78.seq	CAGTCCTACAAGTTCAAGCACCAGCTCCTCCTGACCTTCACCGTGACCCAGAAGACGCA	179				
SEQIDNO-79.seq	CAGTCCTACAAGTTCAAGCACCAGCTCCTCCTGACCTTCACCGTGACCCAGAAGACGCA	179				
SEQIDNO-81.seq	CAGTCCTGCAAGTTCAAGCACCAGCTCCTCCTGACCTTCACCGTGACCCAGAAGACGCA	179				
SEQIDNO-82.seq	CAGTCCTGCAAGTTCAAGCACCAGCTCCTCCTGACCTTCACCGTGACCCAGAAGACGCA	179				
SEQIDNO-83.seq	CAGTCCTACAAGTTCAAGCACCAGCTCCTCCTGACCTTCACCGTGACCCAGAAGACGCA	179				

FIG. 11B

	190	200	210	220	230	240	
SEQIDNO-9.seq	GAGGCGCT	GGTT CCT	CGACAAGCT	GGT CGACGAGAT	CGGGGT GGGCAAGGT	CTACGACCG	239
SEQIDNO-10.seq	GAGGCGCT	GGTT CCT	CGACAAGCT	GGT CGACGAGAT	CGGGGT GGGCAAGGT	CTACGACCG	239
SEQIDNO-11.seq	GAGGCGCT	GGTT CCT	CGACAAGCT	GGT CGACGAGAT	CGGGGT GGGCT	ACGTCTACGACCG	239
SEQIDNO-12.seq	GAGGCGCT	GGTT CCT	CGACAAGCT	GGT CGACGAGAT	CGGGGT GGGCT	ACGTCTACGACCG	239
SEQIDNO-13.seq	GAGGCGCT	GGTT CCT	CGACAAGCT	GGT CGACGAGAT	CGGGGT GGGCAAGGT	CCGCGACCG	239
SEQIDNO-14.seq	GAGGCGCT	GGTT CCT	CGACAAGCT	GGT CGACGAGAT	CGGGGT GGGCT	ACGTCTACGACCG	239
SEQIDNO-15.seq	GAGGCGCT	GGTT CCT	CGACAAGCT	GGT CGACGAGAT	CGGGGT GGGCAAGGT	CTACGACCG	239
SEQIDNO-16.seq	GAGGCGCT	GGTT CCT	CGACAAGCT	GGT CGACGAGAT	CGGGGT GGGCAAGGT	CCGCGACCG	239
SEQIDNO-80.seq	GAGGCGCT	GGTT CCT	CGACAAGCT	GGT CGACGAGAT	CGGGGT GGGCT	ACGTCCGCGACCG	239
SEQIDNO-78.seq	GAGGCGCT	GGTT CCT	CGACAAGCT	GGT CGACGAGAT	CGGGGT GGGCT	ACGTCCGCGACCG	239
SEQIDNO-79.seq	GAGGCGCT	GGTT CCT	CGACAAGCT	GGT CGACGAGAT	CGGGGT GGGCAAGGT	CCGCGACCG	239
SEQIDNO-81.seq	GAGGCGCT	GGTT CCT	CGACAAGCT	GGT CGACGAGAT	CGGGGT GGGCT	ACGTCCGCGACCG	239
SEQIDNO-82.seq	GAGGCGCT	GGTT CCT	CGACAAGCT	GGT CGACGAGAT	CGGGGT GGGCT	ACGTCCGCGACCG	239
SEQIDNO-83.seq	GAGGCGCT	GGTT CCT	CGACAAGCT	GGT CGACGAGAT	CGGGGT GGGCT	ACGTCCGCGACCG	239
	250	260	270	280	290	300	
SEQIDNO-9.seq	CGGGT CGGT	GTCCGACT	ACATCCT	CTCCCAGAT	CAAGCCCCT	GCACAACTT	CCTCACCCTA 299
SEQIDNO-10.seq	CGGGT CGGT	GTCCGACT	ACCGCCT	CTCCCAGAT	CAAGCCCCT	GCACAACTT	CCTCACCCTA 299
SEQIDNO-11.seq	CGGGT CGGT	GTCCGACT	ACATCCT	CTCCCAGAT	CAAGCCCCT	GCACAACTT	CCTCACCCTA 299
SEQIDNO-12.seq	CGGGT CGGT	GTCCGACT	ACCGCCT	CTCCCAGAT	CAAGCCCCT	GCACAACTT	CCTCACCCTA 299
SEQIDNO-13.seq	CGGGT CGGT	GTCCGACT	ACATCCT	CTCCCAGAT	CAAGCCCCT	GCACAACTT	CCTCACCCTA 299
SEQIDNO-14.seq	CGGGT CGGT	GTCCGACT	ACATCCT	CTCCCAGAT	CAAGCCCCT	GCACAACTT	CCTCACCCTA 299
SEQIDNO-15.seq	CGGGT CGGT	GTCCGACT	ACCGCCT	CTCCCAGAT	CAAGCCCCT	GCACAACTT	CCTCACCCTA 299
SEQIDNO-16.seq	CGGGT CGGT	GTCCGACT	ACCGCCT	CTCCCAGAT	CAAGCCCCT	GCACAACTT	CCTCACCCTA 299
SEQIDNO-80.seq	CGGGT CGGT	GTCCGACT	ACATCCT	CTCCCAGAT	CAAGCCCCT	GCACAACTT	CCTCACCCTA 299
SEQIDNO-78.seq	CGGGT CGGT	GTCCGACT	ACATCCT	CTCCCAGAT	CAAGCCCCT	GCACAACTT	CCTCACCCTA 299
SEQIDNO-79.seq	CGGGT CGGT	GTCCGACT	ACATCCT	CTCCCAGAT	CAAGCCCCT	GCACAACTT	CCTCACCCTA 299
SEQIDNO-81.seq	CGGGT CGGT	GTCCGACT	ACATCCT	CTCCCAGAT	CAAGCCCCT	GCACAACTT	CCTCACCCTA 299
SEQIDNO-82.seq	CGGGT CGGT	GTCCGACT	ACATCCT	CTCCCAGAT	CAAGCCCCT	GCACAACTT	CCTCACCCTA 299
SEQIDNO-83.seq	CGGGT CGGT	GTCCGACT	ACATCCT	CTCCCAGAT	CAAGCCCCT	GCACAACTT	CCTCACCCTA 299

FIG. 11C

	850	860	870	880	890	900	
SEQIDNO-9.seq	CGGCTCCATCAAGGCGCAGATCAAGCCGCGCCAGTCCCGCAAGTTCAAGCACGAGCTCTC						899
SEQIDNO-10.seq	CGGCTCCATCATCGCGAAGATCAACCCGAACCAAGTCTTCAAGTTCAAGCACCAGCTCCA						899
SEQIDNO-11.seq	CGGCTCCATCAAGGCGTCCATCAACCCGCGCAGCAGTCTTCAAGTTCAAGCACGCGCTCCA						899
SEQIDNO-12.seq	CGGCTCCATCATCGCGTCCATCTCCCGCGCGCAGTCTTCAAGTTCAAGCACGAGCTCCG						899
SEQIDNO-13.seq	CGGCTCCATCATCGCGCAGATCTCCCGCGCAGCAGTCCCGCAAGTTCAAGCACATCTCTC						899
SEQIDNO-14.seq	CGGCTCCATCATCGCGCAGATCAAGCCGAACCAAGTCTTCAAGTTCAAGCACCAGCTCAT						899
SEQIDNO-15.seq	CGGCTCCATCAAGGCGCAGATCAAGCCGCGCAGTCTTCAAGTTCAAGCACGCGCTCAT						899
SEQIDNO-16.seq	CGGCTCCATCAAGGCGCAGATCAACCCGAACCAAGTCTTCAAGTTCAAGCACCAGCTCCG						899
SEQIDNO-80.seq	CGGCTCCATCATCGCGTCCATCAAGCCGCGCAGTCCCGCAAGTTCAAGCACCAGCTCCA						899
SEQIDNO-78.seq	CGGCTCCATCATCGCGCGCATCAAGCCGAACCAAGTCTTCAAGTTCAAGCACCAGCTCTC						710
SEQIDNO-79.seq	CGGCTCCATCATCGCGCGCATCAAGCCGAACCAAGTCTTCAAGTTCAAGCACCAGCTCTC						710
SEQIDNO-81.seq	CGGCTCCATCATCGCGTCCATCAAGCCGCGCAGTCCCGCAAGTTCAAGCACCAGCTCCA						710
SEQIDNO-82.seq	CGGCTCCATCATCGCGTCCATCAAGCCGCGCAGTCCCGCAAGTTCAAGCACCAGCTCCA						710
SEQIDNO-83.seq	CGGCTCCATCATCGCGTCCATCCGCGCGCAGTCCCGCAAGTTCAAGCACCAGCTCCA						710
	910	920	930	940	950	960	
SEQIDNO-9.seq	CCTGACCTTCCAGGTGACCCAGAAGACGCAGAGGCGCTGGTTCTCGACAAGCTGGTCGA						959
SEQIDNO-10.seq	GCTGCGCTTCAACCGTGACCCAGAAGACGCAGAGGCGCTGGTTCTCGACAAGCTGGTCGA						959
SEQIDNO-11.seq	GCTGACCTTCCAGGTGACCCAGAAGACGCAGAGGCGCTGGTTCTCGACAAGCTGGTCGA						959
SEQIDNO-12.seq	CCTGACCTTCAACCGTGACCCAGAAGACGCAGAGGCGCTGGTTCTCGACAAGCTGGTCGA						959
SEQIDNO-13.seq	CCTGACCTTCCAGGTGACCCAGAAGACGCAGAGGCGCTGGTTCTCGACAAGCTGGTCGA						959
SEQIDNO-14.seq	GCTGACCTTCCAGGTGACCCAGAAGACGCAGAGGCGCTGGTTCTCGACAAGCTGGTCGA						959
SEQIDNO-15.seq	GCTGACCTTCCAGGTGACCCAGAAGACGCAGAGGCGCTGGTTCTCGACAAGCTGGTCGA						959
SEQIDNO-16.seq	CCTGACCTTCCAGGTGACCCAGAAGACGCAGAGGCGCTGGTTCTCGACAAGCTGGTCGA						959
SEQIDNO-80.seq	GCTGACCTTCCAGGTGACCCAGAAGACGCAGAGGCGCTGGTTCTCGACAAGCTGGTCGA						959
SEQIDNO-78.seq	CCTGACCTTCAACCGTGACCCAGAAGACGCAGAGGCGCTGGTTCTCGACAAGCTGGTCGA						770
SEQIDNO-79.seq	CCTGACCTTCAACCGTGACCCAGAAGACGCAGAGGCGCTGGTTCTCGACAAGCTGGTCGA						770
SEQIDNO-81.seq	GCTGACCTTCCAGGTGACCCAGAAGACGCAGAGGCGCTGGTTCTCGACAAGCTGGTCGA						770
SEQIDNO-82.seq	GCTGACCTTCCAGGTGACCCAGAAGACGCAGAGGCGCTGGTTCTCGACAAGCTGGTCGA						770
SEQIDNO-83.seq	GCTGCGCTTCAACCGTGACCCAGAAGACGCAGAGGCGCTGGTTCTCGACAAGCTGGTCGA						770

FIG. 11D

	970	980	990	1000	1010	1020	
SEQIDNO-9.seq	CGAGATCGGGGTGGGCTACGTCTACGACCGCGGGTCGGTGTCCGACTACATCCTCTCCCA						1019
SEQIDNO-10.seq	CGAGATCGGGGTGGGCAAGGTCTACGACCGCGGGTCGGTGTCCGACTACATCCTCTCCCA						1019
SEQIDNO-11.seq	CGAGATCGGGGTGGGCAAGGTCTACGACCGCGGGTCGGTGTCCGACTACCGCCTCTCCCA						1019
SEQIDNO-12.seq	CGAGATCGGGGTGGGCTACGTCTACGACCGCGGGTCGGTGTCCGACTACCGCCTCTCCCA						1019
SEQIDNO-13.seq	CGAGATCGGGGTGGGCAAGGTCTACGACCGCGGGTCGGTGTCCGACTACCGCCTCTCCCA						1019
SEQIDNO-14.seq	CGAGATCGGGGTGGGCTACGTCCGCGACCGCGGGTCGGTGTCCGACTACATCCTCTCCCA						1019
SEQIDNO-15.seq	CGAGATCGGGGTGGGCTACGTCTACGACCGCGGGTCGGTGTCCGACTACATCCTCTCCCA						1019
SEQIDNO-16.seq	CGAGATCGGGGTGGGCAAGGTCTACGACCGCGGGTCGGTGTCCGACTACATCCTCTCCCA						1019
SEQIDNO-80.seq	CGAGATCGGGGTGGGCTACGTCCGCGACCGCGGGTCGGTGTCCGACTACCGCCTCTCCCA						1019
SEQIDNO-78.seq	CGAGATCGGGGTGGGCTACGTCCGCGACCGGGGTTCGGTGTCCGACTACCGCTCTCCCA						830
SEQIDNO-79.seq	CGAGATCGGGGTGGGCTACGTCCGCGACCGGGGTTCGGTGTCCGACTACCGCTCTCCCA						830
SEQIDNO-81.seq	CGAGATCGGGGTGGGCTACGTCCGCGACCGCGGGTCGGTGTCCGACTACCGCCTCTCCCA						830
SEQIDNO-82.seq	CGAGATCGGGGTGGGCTACGTCCGCGACCGCGGGTCGGTGTCCGACTACCGCCTCTCCCA						830
SEQIDNO-83.seq	CGAGATCGGGGTGGGCTACGTCTACGACCGGGGTTCGGTGTCCGACTACCGCCTCTCCCA						830

METHODS FOR PRODUCING A COMPLEX TRANSGENIC TRAIT LOCUS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application is a continuation of U.S. patent application Ser. No. 17/821,528 filed on Aug. 23, 2022 (published as U.S. patent application No. 20230235342), U.S. patent application Ser. No. 16/925,973 filed on Jul. 10, 2020 (published as U.S. Pat. No. 11,459,576), which is a continuation of U.S. patent application Ser. No. 16/007,529 filed on Jun. 13, 2018 (now allowed as U.S. Pat. No. 10,822,610), which is a continuation of U.S. patent application Ser. No. 13/427,138 (now allowed as U.S. Pat. No. 10,030,245), which itself claims the benefit of U.S. Provisional Patent Application No. 61/499,443, filed Jun. 11, 2011 and U.S. Provisional Patent Application No. 61/466,602, filed Mar. 23, 2011; all of which are hereby incorporated herein in their entirety by reference.

REFERENCE TO SEQUENCE LISTING

[0002] The Sequence Listing is submitted as a text file named BB1990-US-CNT[3]_SeqListing_ST25 created on Aug. 5 2022 having a size of 150 KB, and is hereby incorporated by reference pursuant to 37 C.F.R. § 1.52 (e) (5).

FIELD OF INVENTION

[0003] The invention relates to the field of plant molecular biology, in particular, to methods for altering the genome of a plant cell.

BACKGROUND

[0004] Recombinant DNA technology has made it possible to insert foreign DNA sequences into the genome of an organism, thus, altering the organism's phenotype. The most commonly used plant transformation methods are *Agrobacterium* infection and biolistic particle bombardment in which transgenes integrate into a plant genome in a random fashion and in an unpredictable copy number. Thus, efforts are undertaken to control transgene integration in plants.

[0005] One method for inserting or modifying a DNA sequence involves homologous DNA recombination by introducing a transgenic DNA sequence flanked by sequences homologous to the genomic target. U.S. Pat. No. 5,527,695 describes transforming eukaryotic cells with DNA sequences that are targeted to a predetermined sequence of the eukaryote's DNA. Specifically, the use of site-specific recombination is discussed. Transformed cells are identified through use of a selectable marker included as a part of the introduced DNA sequences.

[0006] It was shown that artificially induced site-specific genomic double-stranded breaks in plant cells were repaired by homologous recombination with exogenously supplied DNA using two different pathways. (Puchta et al., (1996) *Proc. Natl. Acad. Sci. USA* 93:5055-5060; U.S. Patent Application Publication No. 2005/0172365A1 published Aug. 4, 2005; U.S. Patent Application Publication No. 2006/0282914 published Dec. 14, 2006; WO 2005/028942 published Jun. 2, 2005).

[0007] Since the isolation, cloning, transfer and recombination of DNA segments, including coding sequences and non-coding sequences, is most conveniently carried out

using restriction endonuclease enzymes. Much research has focused on studying and designing endonucleases such as WO 2004/067736 published Aug. 12, 2004; U.S. Pat. No. 5,792,632 issued to Dujon et al., Aug. 11, 1998; U.S. Pat. No. 6,610,545 B2 issued to Dujon et al., Aug. 26, 2003; Chevalier et al., (2002) *Mol Cell* 10:895-905; Chevalier et al., (2001) *Nucleic Acids Res* 29:3757-3774; Seligman et al., (2002) *Nucleic Acids Res* 30:3870-3879.

[0008] Although a plethora of approaches have been developed to target a specific site for modification in the genome of a plant, there still remains a need for methods for producing a fertile plant, having an altered genome comprising two or more site-specific modifications in defined region of the genome of the plant.

BRIEF SUMMARY OF THE INVENTION

[0009] The present invention provides methods for producing in a plant a complex transgenic trait locus comprising at least two altered target sequences in a genomic region of interest. The methods involve selecting a genomic region in a plant that comprises a first target sequence and a second target sequence and then providing a first double-strand-break-inducing agent and a second double-strand-break-inducing agent. The first double-strand-break-inducing agent is capable of inducing a first double-strand break in DNA comprising the first target sequence, and the second double-strand-break-inducing agent is capable of inducing a second double-strand break in DNA comprising the second target sequence. The methods further involve contacting at least one plant cell with the first double-strand-break-inducing agent, identifying a cell comprising a first alteration at the first target sequence, and then recovering a first fertile plant from the cell comprising the first alteration. The first fertile plant also comprises the first alteration. Additionally, the methods involve contacting at least one plant cell with the second double-strand-break-inducing agent, identifying a cell comprising a second alteration at the second target sequence, and then recovering a second fertile plant from the cell comprising the second alteration. The methods further involve obtaining a fertile progeny plant from the second fertile plant, wherein the fertile progeny plant comprises both the first and second alterations in physical linkage.

[0010] In a first embodiment of the methods for producing in a plant a complex transgenic trait locus, the fertile progeny plant is obtained by crossing the first fertile plant and the second fertile plant and selecting the fertile progeny plant comprising both the first and second alterations in physical linkage.

[0011] In second embodiment, a cell of the first fertile plant, or progeny thereof comprising the first alteration, is contacted with the second double-strand-break-inducing agent.

[0012] In third embodiment, the complex transgenic trait locus further comprises at least one polynucleotide of interest in the genomic region of interest. Such a polynucleotide of interest can be, for example, a transgene, a native gene, and a gene that was a native gene prior to a targeted mutation therein.

[0013] In a fourth embodiment, the first alteration comprises insertion of a first DNA sequence of interest, or part thereof, into the first target sequence, and/or the second alteration comprises insertion of a second DNA sequence of interest, or part thereof, into the second target sequence. Such a first and/or a second DNA sequence of interest can

be, for example, a DNA for gene silencing, a DNA encoding a phenotypic marker and a DNA encoding a protein providing an agronomic advantage.

[0014] In a fifth embodiment, the first and second double-strand-break-inducing agents are selected from the group consisting of an endonuclease, a zinc finger nuclease, or a TAL effector nuclease.

[0015] In a sixth embodiment, the endonuclease is modified to specifically cut at the first target sequence or at the second target sequence and no longer cuts at its wild-type endonuclease target sequence.

[0016] In a seventh embodiment, the first target sequence and the second target sequence are separated from each other by about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, or 5 centimorgans (CM) in the genome of the plant.

[0017] In an eighth embodiment, the methods can involve crossing the fertile progeny plant with an additional fertile plant that comprises at least a third altered target sequence in the genomic region of interest and then selecting from the crossing a fertile progeny plant comprising the first alteration, the second and the at least third alteration in physical linkage. Like the first and second altered target sequences, the third altered target sequence originated from a third target sequence that is recognized and cleaved by a third double-strand-break-inducing agent.

[0018] Additionally provided are complex trait loci in plants produced by the methods of the invention and plants, plant cells, plant parts, and seeds thereof comprising at least one complex transgenic trait locus of the invention.

[0019] The present invention further provides a complex transgenic trait locus comprising at least two altered target sequences that are genetically linked in the genome of a plant to a polynucleotide of interest. Such altered target sequences originated from a corresponding target sequence that is recognized and cleaved by a double-strand-break-inducing agent. The altered target sequences comprise an alteration such as, for example, replacement of at least one nucleotide in the target sequence, a deletion of at least one nucleotide in the target sequence, an insertion of at least one nucleotide in the target sequence, or any combination thereof. The polynucleotide interest can be, for example, a transgene, a native gene, and a mutated gene. The present invention further provides plants, plant parts, plant cells, and seeds comprising at least one complex transgenic trait locus of the invention.

[0020] In an embodiment of the complex transgenic trait locus of the invention, at least one altered target sequence comprises a recombinant DNA molecule. Recombinant DNA molecules include, but are not limited to, a DNA for gene silencing, a DNA encoding a phenotypic marker, and a DNA encoding a protein providing an agronomic advantage.

[0021] In another embodiment, the two altered target sequences of the complex transgenic trait locus are located within about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or up to 21 centimorgan (cM) of the polynucleotide of interest.

[0022] The invention provides plants, plant parts, plant cells, and seeds comprising at least one complex transgenic trait locus of the invention.

[0023] Additionally provided is an alternative method for producing in a plant a complex transgenic trait locus comprising at least two altered target sequences in a genomic region of interest. This method involves obtaining a first

fertile plant comprising a first altered target sequence at the genomic region of interest and a second fertile plant comprising a second altered target sequence at the genomic region of interest. In this method, the first altered target sequence originated from a first target sequence that is recognized and cleaved by a first double-strand-break-inducing agent, and the second altered target sequence originated from a second target sequence that is recognized and cleaved by a second double-strand-break-inducing agent. The alternative method further involves crossing the first fertile plant and the second fertile plant, and then selecting from the crossing a fertile progeny plant comprising the first alteration and the second alteration in physical linkage.

[0024] Also provided are plants produced by the second method of the invention and plant cells, plant parts, and seeds thereof comprising at least one complex transgenic trait locus.

[0025] In another embodiment, the present invention provides a plant comprising an expression construct, which comprises a promoter operably linked to a nucleotide sequence encoding an endonuclease. The endonuclease is capable of specifically binding to and creating a double strand break in a target sequence selected from the group consisting of SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 68, 69, 70, 71, 72, 73, 74, 75, 76, and 77, wherein the promoter is capable of driving expression of an operably linked nucleotide sequence in a plant cell. The nucleotide sequence encoding the endonuclease can comprise a coding sequence of a DNA binding domain of an endonuclease, wherein the coding sequence comprises nucleotides 100-261 and nucleotides 850-1011 of SEQ ID NO: 9, 10, 11, 12, 13, 14, 15, 16, 78, 79, 80, 81, 82 or 83; or a degenerate coding sequence thereof. Preferably, the nucleotide sequence encoding the endonuclease is a nucleotide sequence selected from the group consisting of SEQ ID NO: 9, 10, 11, 12, 13, 14, 15, 16, 78, 79, 80, 81, 82, and 83.

[0026] In yet another embodiment of the invention, a plant of the invention comprises at least one altered target sequence, wherein the at least one altered target sequence originated from a corresponding target sequence that was recognized and cleaved by a double-strand break-inducing agent. In this embodiment, the altered target sequence is in a genomic region of interest that extends from: the target sequence set forth in SEQ ID NO: 4 to the target sequence set forth in SEQ ID NO: 2; the target sequence set forth in SEQ ID NO: 5 to the target sequence set forth in SEQ ID NO: 8; or the target sequence set forth in SEQ ID NO: 68 to the target sequence set forth in SEQ ID NO: 77. Such a plant of the invention can be produced by a method comprising providing at least one double-strand-break-inducing agent that is capable of inducing a double-strand break in DNA comprising a target sequence, wherein the target sequence is in a genomic region of interest that extends from: the target sequence set forth in SEQ ID NO: 4 to the target sequence set forth in SEQ ID NO: 2; the target sequence set forth in SEQ ID NO: 5 to the target sequence set forth in SEQ ID NO: 8; or the target sequence set forth in SEQ ID NO: 68 to the target sequence set forth in SEQ ID NO: 77. The method further comprises contacting at least one plant cell with the double-strand-break-inducing agent, identifying a cell comprising an alteration at the target sequence, and recovering a fertile plant comprising the alteration. In one embodiment of this method, the double-strand-break-inducing agent is encoded by a nucleotide sequence comprising a coding

sequence of a DNA binding domain of an endonuclease, and wherein the coding sequence is selected from the group consisting of nucleotides 100-261 and nucleotides 850-1011 of SEQ ID NO: 9, 10, 11, 12, 13, 14, 15, 16, and 80, and degenerate coding sequences thereof. In another embodiment of this method, the double-strand-break-inducing agent is encoded by a nucleotide sequence comprising a coding sequence of a DNA binding domain of an endonuclease, and wherein the coding sequence is selected from the group consisting of nucleotides 100-261 and nucleotides 661-822 of SEQ ID NO: 78, 79, 81, 82 and 83, and degenerate coding sequences thereof. In another embodiment of this method, the double-strand-break-inducing agent is encoded by a nucleotide sequence is selected from the group consisting of SEQ ID NO: 9, 10, 11, 12, 13, 14, 15, 16, 78, 79, 80, 81, 82, and 83.

[0027] Additional embodiments of the methods and compositions of the present invention are disclosed below.

BRIEF DESCRIPTION OF THE DRAWINGS AND THE SEQUENCE LISTING

[0028] The invention can be more fully understood from the following detailed description and the accompanying drawings and Sequence Listing, which form a part of this application. The sequence descriptions and sequence listing attached hereto comply with the rules governing nucleotide and amino acid sequence disclosures in patent applications as set forth in 37 C.F.R. §§ 1.821-1.825. The sequence descriptions contain the three letter codes for amino acids as defined in 37 C.F.R. §§ 1.821-1.825, which are incorporated herein by reference.

FIGURES

[0029] FIG. 1. DNA double-strand break induced DNA alteration of an endogenous target site. FIG. 1A A generalized endogenous target site with flanking genomic DNA sequences designated as DNA 1 and DNA 2 which can be used as DNA exchange regions by homologous recombination. FIG. 1B A generalized DNA construct that can be used to express a DNA endonuclease to recognize and cleave the endogenous target site. The DNA endonuclease gene can be physically linked to the donor DNA described in FIG. 1C or FIG. 1D, or substituted by other double-strand break inducing agents. FIG. 1C A generalized donor DNA construct having two regions DNA1 and DNA 2 of homology to the genomic target which flank a polynucleotide of interest and/or marker gene. FIG. 1D A generalized donor DNA construct that does not have regions of homology to the genomic target to flank a polynucleotide of interest and/or marker gene. Insertion of the DNA fragment will produce an insertion of the polynucleotide of interest at or near the recognition site. FIG. 1E One expected outcome when the polynucleotide of interest and/or marker gene of donor construct described in FIG. 1C or FIG. 1D is inserted at the endogenous target site by homologous recombination or non-homologous recombination, respectively. FIG. 1F Another outcome when the endogenous target site is altered by a deletion during the repair of the DNA double-strand break cleaved by the DNA endonuclease. The polynucleotide of interest and/or marker gene of donor construct described in FIG. 1C or FIG. 1D can be inserted at unrelated sites by random DNA integration. FIG. 1G Another outcome when the endogenous target site is altered by the insertion of

an unrelated DNA during the repair of the DNA double-strand breaks cleaved by the DNA endonuclease. The polynucleotide of interest and/or marker gene of donor construct described in FIG. 1C or FIG. 1D can be inserted at unrelated sites by random DNA integration.

[0030] FIG. 2. Genetic distance between target sites and transgene of interest.

[0031] FIG. 3. FIG. 3A: Schematic diagram of PCR assays to detect TS21 target site modifications and transgenic integrations. FIG. 3B: Alignment of altered target sequences of selected TS21 transgenic event.

[0032] FIG. 4. FIG. 4A: Alignment of altered target sequences of selected TS5 transgenic events. FIG. 4B: Alignment of altered target sequences of selected TS14 transgenic events

[0033] FIG. 5. Gene integration by homologous recombination enabled by double-strand breaks with custom designed meganuclease.

[0034] FIG. 6. Location of target sites near a herbicide resistant transgenic event in soybean.

[0035] FIG. 7. Use of cluster of meganuclease target sites for stacking of multiple traits either by sequential transformation or genetic crosses.

[0036] FIG. 8. The locations of various MHP target sites surrounding a transgenic DNA of interest integration site in a maize plant. Solid black rectangles represent BAC clones. Names and numbers in each box are the target sites. Arrows from box to BAC indicated the target site affiliated to BAC clones. Numbers and arrows on the bottom of the figure indicate the genetic distance of the target sites relative to the insertion location of the transgenic DNA of interest. As indicated at the top of the figure, the physical distance is about 1.8 Mb nucleotides in this region of the maize chromosome.

[0037] FIG. 9. FIG. 9A: Outline of PCR screening for integration of donor at MHP14 target site (donor was PHP44779) FIG. 9B: PCR of MHP14 events: B1-B12 junction PCR with primers 146773/146775; b1-b12 junction PCR with primers 146772/146778. Two events (B2 and B5) were positive for both junctions PCR. The arrows indicate the locations corresponding to the various primers used.

[0038] FIG. 10. FIG. 10A: Schematic outline of PCR to confirm ubi: mopat: pinII cassette integration at the endogenous MHP14 target. FIG. 10B: Long PCR on TO plants from three events showed integration at the target site. The plant A5 was from event #1, A6-A8 event #2, and C4-C6 event #3. CKP: positive control from callus DNA. FIG. 10B: The left panel shows the results of junction PCR on the HR1 side using a genomic primer (146775) and a moPAT primer (mopatR2). The right panel shows the results of junction PCR on the HR2 side with a moPAT primer (mopatF2) a genomic primer (146772). The arrows on FIG. 10A indicate the locations corresponding to the various primers used.

[0039] FIG. 11A-11D. Alignment of fragments from the plant-optimized nucleotide sequences of meganucleases comprising the nucleotides 100-261 and nucleotides 850-1011 of SEQ ID NO: 9, 10, 11, 12, 13, 14, 15, 16, and 80, and the nucleotides 100-261 and nucleotides 661-822 of SEQ ID NO: 78, 79, 81, 82 and 83. FIG. 11A shows SEQ ID NOs: 9-16 and 78-80 sequence positions 60-180, FIG. 11B shows SEQ ID NOs: 9-16 and 78-80 sequence positions 180-300, FIG. 11C shows SEQ ID NOs: 9-16 and 78-80 sequence positions 840-960, and FIG. 11D shows SEQ ID NOs: 9-16 and 78-80 sequence positions 960-1020.

SEQUENCES

[0040] SEQ ID NO: 1 is the nucleotide sequence of the TS21 target site in soybean genome.

[0041] SEQ ID NO: 2 is the nucleotide sequence of the TS14 target site in soybean genome.

[0042] SEQ ID NO: 3 is the nucleotide sequence of the TS30 target site in soybean genome.

[0043] SEQ ID NO: 4 is the nucleotide sequence of the TS5 target site in soybean genome.

[0044] SEQ ID NO: 5 is the nucleotide sequence of the TS7 target site in soybean genome.

[0045] SEQ ID NO: 6 is the nucleotide sequence of the TS4 target site in soybean genome.

[0046] SEQ ID NO: 7 is the nucleotide sequence of the TS22 target site in soybean genome.

[0047] SEQ ID NO: 8 is the nucleotide sequence of the TS24 target site in soybean genome.

[0048] SEQ ID NO: 9 is the plant-optimized nucleotide sequence of the TS21 meganuclease containing a nuclear target site and an ST-LS1 intron.

[0049] SEQ ID NO: 10 is the plant-optimized nucleotide sequence of the TS14 meganuclease containing a nuclear target site and an ST-LS1 intron.

[0050] SEQ ID NO: 11 is the plant-optimized nucleotide sequence of the TS30 meganuclease containing a nuclear target site and an ST-LS1 intron.

[0051] SEQ ID NO: 12 is the plant-optimized nucleotide sequence of the TS5 meganuclease containing a nuclear target site and an ST-LS1 intron.

[0052] SEQ ID NO: 13 is the plant-optimized nucleotide sequence of the TS7 meganuclease containing a nuclear target site and an ST-LS1 intron.

[0053] SEQ ID NO: 14 is the plant-optimized nucleotide sequence of the TS4 meganuclease containing a nuclear target site and an ST-LS1 intron.

[0054] SEQ ID NO: 15 is the plant-optimized nucleotide sequence of the TS22 meganuclease containing a nuclear target site and an ST-LS1 intron.

[0055] SEQ ID NO: 16 is the plant-optimized nucleotide sequence of the TS24 meganuclease containing a nuclear target site and an ST-LS1 intron.

[0056] SEQ ID NO: 17 is the homologous region 1 (HR1) of the TS21 target site.

[0057] SEQ ID NO: 18 is the homologous region 2 (HR2) of the TS21 target site.

[0058] SEQ ID NO: 19 is the HR1 of the TS14 target site.

[0059] SEQ ID NO: 20 is the homologous region 2 of the TS14 target site.

[0060] SEQ ID NO: 21 is the HR1 of the TS30 target site.

[0061] SEQ ID NO: 22 is the homologous region 2 of the TS30 target site.

[0062] SEQ ID NO: 23 is the HR1 of the TS5 target site.

[0063] SEQ ID NO: 24 is the homologous region 2 of the TS5 target site.

[0064] SEQ ID NO: 25 is the HR1 of the TS7 target site.

[0065] SEQ ID NO: 26 is the homologous region 2 of the TS7 target site.

[0066] SEQ ID NO: 27 is the HR1 of the TS4 target site.

[0067] SEQ ID NO: 28 is the homologous region 2 of the TS4 target site.

[0068] SEQ ID NO: 29 is the HR1 of the TS22 target site.

[0069] SEQ ID NO: 30 is the homologous region 2 of the TS22 target site.

[0070] SEQ ID NO: 31 is the HR1 of the TS24 target site.

[0071] SEQ ID NO: 32 is the homologous region 2 of the TS24 target site.

[0072] SEQ ID NO: 33 is the plant-optimized nucleotide sequence of the TS21 meganuclease without a ST-LS1 intron.

[0073] SEQ ID NO: 34 is the amino acid sequence of the SV40 nuclear localization signal.

[0074] SEQ ID NO: 35 is the nucleotide sequences of expression cassette RTW317, comprising the TS21 meganuclease plant optimized sequence without an intron and operably linked to the soybean EF1A promoter.

[0075] SEQ ID NO: 36 is the nucleotide sequences of expression cassette RTW322, comprising the TS21 meganuclease plant optimized sequence with an intron and operably linked to the soybean EF1A promoter.

[0076] SEQ ID NO: 37 is the nucleotide sequence of RTW328A, which is the repair DNA fragment for TS21 meganuclease.

[0077] SEQ ID NO:38 is the nucleotide sequence of TS21 qPCR forward primer Mega21-190F.

[0078] SEQ ID NO:39 is the nucleotide sequence of TS21 qPCR reverse primer Mega21-301R.

[0079] SEQ ID NO:40 is the nucleotide sequence of TS21 qPCR probe mega21-250T.

[0080] The fluorescent probe is labeled with FAM quenched with MGB.

[0081] SEQ ID NO:41 is the nucleotide sequence of TS14 qPCR forward primer Mega14-13F.

[0082] SEQ ID NO:42 is the nucleotide sequence of TS14 qPCR reverse primer Mega14-128R.

[0083] SEQ ID NO:43 is the nucleotide sequence of TS14 qPCR probe Mega14-85T.

[0084] The fluorescent probe is labeled with FAM quenched with MGB.

[0085] SEQ ID NO:44 is the nucleotide sequence of TS30 qPCR forward primer Mega30-30F.

[0086] SEQ ID NO:45 is the nucleotide sequence of TS30 qPCR reverse primer Mega30-87R.

[0087] SEQ ID NO:46 is the nucleotide sequence of TS30 qPCR probe Mega30-52T. The fluorescent probe is labeled with FAM quenched with MGB.

[0088] SEQ ID NO:47 is the nucleotide sequence of TS5 qPCR forward primer Mega5-F1.

[0089] SEQ ID NO:48 is the nucleotide sequence of TS5 qPCR reverse primer Mega5-R1

[0090] SEQ ID NO:49 is the nucleotide sequence of TS5 qPCR probe Mega5-T1. The fluorescent probe is labeled with FAM quenched with MGB.

[0091] SEQ ID NO:50 is the nucleotide sequence of the sense primer, WOL133, which is upstream of the TS21 target site in the soybean genome.

[0092] SEQ ID NO:51 is the nucleotide sequence of the antisense primer, WOL134, which is downstream of the TS21 target site in the soybean genome.

[0093] SEQ ID NO:52 is the nucleotide sequence of the sense primer, WOL190 which is further upstream of the TS21 target site beyond the TS21 HR1 fragment in the soybean genome.

[0094] SEQ ID NO:53 is the nucleotide sequence of the antisense primer, WOL242, which is specific to the hygromycin coding sequences.

[0095] SEQ ID NO:54 is the nucleotide sequence of the sense primer, WOL153, which is specific to the NOS Terminator.

[0096] SEQ ID NO:55 is the nucleotide sequence of the antisense primer, WOL247, which is further downstream of the TS21 target site beyond the TS21 HR2 fragment in the soybean genome.

[0097] SEQ ID NO:56 is the nucleotide sequence of the sense primer, WOL121, which is upstream of the TS14 target site in the soybean genome.

[0098] SEQ ID NO:57 is the nucleotide sequence of the antisense primer, WOL150, which is downstream of the TS21 target site in the soybean genome.

[0099] SEQ ID NO:58 is the nucleotide sequence of the sense primer, WOL192, which is further upstream of the TS14 target site beyond the TS14 HR1 fragment in the soybean genome.

[0100] SEQ ID NO:59 is the nucleotide sequence of the antisense primer, WOL193, which is further downstream of the TS14 target site beyond the TS14 HR2 fragment in the soybean genome.

[0101] SEQ ID NO:60 is the nucleotide sequence of the sense primer, WOL113, which is upstream of the TS30 target site in the soybean genome.

[0102] SEQ ID NO:61 is the nucleotide sequence of the antisense primer, WOL114, which is downstream of the TS30 target site in the soybean genome.

[0103] SEQ ID NO:62 is the nucleotide sequence of the sense primer, WOL194, which is further upstream of the TS30 target site beyond the TS30 HR1 fragment in the soybean genome.

[0104] SEQ ID NO:63 is the nucleotide sequence of the antisense primer, WOL195, which is further downstream of the TS30 target site beyond the TS30 HR2 fragment in the soybean genome.

[0105] SEQ ID NO:64 is the nucleotide sequence of the sense primer, WOL105, which is upstream of the TS5 target site in the soybean genome.

[0106] SEQ ID NO:65 is the nucleotide sequence of the antisense primer, WOL144, which is downstream of the TS5 target site in the soybean genome.

[0107] SEQ ID NO:66 is the nucleotide sequence of the sense primer, WOL196, which is further upstream of the TS5 target site beyond the TS5 HR1 fragment in the soybean genome.

[0108] SEQ ID NO:67 is the nucleotide sequence of the antisense primer, WOL197, which is further downstream of the TS5 target site beyond the TS5 HR2 fragment in the soybean genome.

[0109] SEQ ID NO:68 is the nucleotide sequence of the MHP1 target site in the maize genome.

[0110] SEQ ID NO:69 is the nucleotide sequence of the MHP14 target site sequence in the maize genome.

[0111] SEQ ID NO:70 is the nucleotide sequence of the MHP32 target site sequence in the maize genome.

[0112] SEQ ID NO:71 is the nucleotide sequence of the MHP42 target site sequence in the maize genome.

[0113] SEQ ID NO:72 is the nucleotide sequence of the MHP55 target site sequence in the maize genome.

[0114] SEQ ID NO:73 is the nucleotide sequence of the MHP67 target site sequence in the maize genome.

[0115] SEQ ID NO:74 is the nucleotide sequence of the MHP77 target site sequence in the maize genome.

[0116] SEQ ID NO:75 is the nucleotide sequence of the MHP98 target site sequence in the maize genome.

[0117] SEQ ID NO:76 is the nucleotide sequence of the MHP107 target site sequence in the maize genome.

[0118] SEQ ID NO: 77 is the nucleotide sequence of the MHP115 target site sequence in the maize genome.

[0119] SEQ ID NO:78 is the plant-optimized nucleotide sequence of MHP14 comprising a nuclear localization signal and lacking an intron.

[0120] SEQ ID NO:79 is the plant-optimized nucleotide sequence of the MHP14+ comprising a nuclear localization signal and lacking an intron.

[0121] SEQ ID NO:80 is the plant-optimized nucleotide sequence of MHP55 comprising a nuclear localization signal and an intron.

[0122] SEQ ID NO:81 is the plant-optimized nucleotide sequence of MHP55 comprising a nuclear localization signal and lacking an intron.

[0123] SEQ ID NO:82 is the plant-optimized nucleotide sequence of MHP55-2 comprising a nuclear localization signal and lacking an intron.

[0124] SEQ ID NO:83 plant-optimized nucleotide sequence of MHP77 comprising a nuclear localization signal and lacking an intron.

[0125] SEQ ID NO:84 is the HR1 of the MHP14 target site.

[0126] SEQ ID NO:85 is the HR2 of the MHP14 target site.

[0127] SEQ ID NO:86 is the HR1 of the MHP55 target site.

[0128] SEQ ID NO:87 is the HR2 of the MHP55 target site.

[0129] SEQ ID NO:88 is the HR1 of the MHP77 target site.

[0130] SEQ ID NO:89 is the HR2 of the MHP77 target site.

[0131] SEQ ID NO: 90 is the HR1 of the MHP1 target site.

[0132] SEQ ID NO:91 is the HR2 of the MHP1 target site.

[0133] SEQ ID NO:92 is the HR1 of the MHP32 target site.

[0134] SEQ ID NO:93 is the HR2 of the MHP32 target site.

[0135] SEQ ID NO:94 is the HR1 of the MHP42 target site.

[0136] SEQ ID NO:95 is the HR2 of the MHP42 target site.

[0137] SEQ ID NO:96 is the HR1 of the MHP67 target site.

[0138] SEQ ID NO:97 is the HR2 of the MHP67 target site.

[0139] SEQ ID NO:98 is the HR1 of the MHP98 target site.

[0140] SEQ ID NO:99 is the HR2 of the MHP98 target site.

[0141] SEQ ID NO: 100 is the HR1 of the MHP107 target site.

[0142] SEQ ID NO:101 is the HR2 of the MHP107 target site.

[0143] SEQ ID NO: 102 is the HR1 of the MHP115 target site.

[0144] SEQ ID NO:103 is the HR2 of the MHP115 target site.

[0145] SEQ ID NO: 104 is the nucleotide sequence of the plasmid PHP 44285 (MHP14 and donor DNA).

[0146] SEQ ID NO: 105 is the nucleotide sequence of the plasmid PHP44779 (MHP14+ and donor DNA).

[0147] SEQ ID NO: 106 is the nucleotide sequence of the MHP14TS probe.

[0148] SEQ ID NO: 107 is the nucleotide sequence of the MHPTS14TS_Forward_MGB primer.

[0149] SEQ ID NO: 108 is the nucleotide sequence of the MHPTS14TS_Reverse_MGB primer.

[0150] SEQ ID NO: 109 is the nucleotide sequence of the primer 146775 on genomic HR1 side.

[0151] SEQ ID NO: 110 is the nucleotide sequence of the primer 146773 on vector HR1 side.

[0152] SEQ ID NO: 111 is the nucleotide sequence of the primer 146772 on genomic HR2 side.

[0153] SEQ ID NO: 112 is the nucleotide sequence of the primer 146778 on vector HR2 side.

[0154] SEQ ID NO: 113 is the nucleotide sequence of the primer mopatF2.

[0155] SEQ ID NO: 114 is the nucleotide sequence of the primer mopatR2.

[0156] SEQ ID NO: 115 is the nucleotide sequence of the MHP55TS probe sequence.

[0157] SEQ ID NO: 116 is the nucleotide sequence of the MHPTS55_Forward_MGB primer.

[0158] SEQ ID NO: 117 is the nucleotide sequence of the MHP55TS_Reverse_MGB primer.

[0159] SEQ ID NO: 118 is the nucleotide sequence of the MHP77TS probe.

[0160] SEQ ID NO: 119 is the nucleotide sequence of the MHP77TS_Forward_MGB primer.

[0161] SEQ ID NO: 120 is the nucleotide sequence of the MHP77TS_Reverse_MGB primer.

DETAILED DESCRIPTION OF THE INVENTION

[0162] All publications and patent applications mentioned in the specification are indicative of the level of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

[0163] As used herein and in the appended claims, the singular forms “a”, “an”, and “the” include plural reference unless the context clearly dictates otherwise. Thus, for example, reference to “a plant” includes a plurality of such plants; reference to “a cell” includes one or more cells and equivalents thereof known to those skilled in the art, and so forth.

[0164] In the context of this disclosure, a number of terms and abbreviations are used. The following definitions are provided.

[0165] As used herein a “complex transgenic trait locus” (plural: “complex transgenic trait loci”) is a chromosomal segment within a genomic region of interest that comprises at least two altered target sequences that are genetically linked to each other and can also comprise one or more polynucleotides of interest as described hereinbelow. Each of the altered target sequences in the complex transgenic trait locus originates from a corresponding target sequence that was altered, for example, by a mechanism involving a double-strand break within the target sequence that was induced by a double-strand break-inducing agent of the invention. In certain embodiments of the invention, the altered target sequences comprise a transgene.

[0166] As used herein, a “genomic region of interest” is a segment of a chromosome in the genome of a plant that is desirable for producing a complex transgenic trait locus or

the segment of a chromosome comprising a complex transgenic trait locus that was produced by the methods of the invention. The genomic region of interest can include, for example, one or more polynucleotides of interest prior to producing a complex transgenic trait locus therein. Generally, a genomic region of interest of the present invention comprises a segment of chromosome that is 0-15 cM.

[0167] The term “recognition sequence” or “recognition site” as used herein refers to a DNA sequence at which a double-strand break is induced in the plant cell genome by a double-strand break inducing agent. The terms “recognition sequence” and “recognition site” are used interchangeably herein.

[0168] The terms “target site”, “target sequence”, “target locus”, “genomic target site”, “genomic target sequence”, and “genomic target locus” as used interchangeably herein refer to a polynucleotide sequence in the genome of a plant cell that comprises a recognition sequence for a double-strand break inducing agent.

[0169] An “artificial target sequence” is a target sequence that has been introduced into the genome of a plant. Such an artificial target sequence can be identical in sequence to an endogenous or native target sequence in the genome of a plant but be located in a different position (i.e., a non-endogenous or non-native position) in the genome of a plant.

[0170] The terms “endogenous target sequence” and “native target sequence” are used interchangeably herein to refer to a target sequence that is endogenous or native to the genome of a plant and is at the endogenous or native position of that target sequence in the genome of the plant.

[0171] An “altered target sequence” refers to a target sequence as disclosed herein that comprises at least one alteration of the invention when compared to non-altered target sequence. Such “alterations” of the invention include, for example: (i) replacement of at least one nucleotide, (ii) a deletion of at least one nucleotide, (iii) an insertion of at least one nucleotide, or (iv) any combination of (i)-(iii).

[0172] The term “double-strand-break-inducing agent” as used herein refers to any nuclease which produces a double-strand break in the target sequence. Producing the double-strand break in a target sequence or other DNA can be referred to herein as “cutting” or “cleaving” the target sequence or other DNA. In some embodiments of the invention, the double-strand-break-inducing agent has been engineered (or modified) to cut a specific endogenous target sequence, wherein the endogenous target sequence prior to being cut by the engineered double-strand-break-inducing agent was not a sequence that would have been recognized by a native (non-engineered or non-modified) double-strand-break-inducing agent.

[0173] As used herein, “physically linked,” “in physical linkage”, and “genetically linked” are used to refer to any two or more genes, transgenes, native genes, mutated genes, alterations, target sites, markers, and the like that are part of the same DNA molecule or chromosome.

[0174] As used herein, a “polynucleotide of interest” within a genomic region of interest is any coding and/or non-coding portion of the genomic region of interest including, but not limited to, a transgene, a native gene, a mutated gene, and a genetic marker such as, for example, a single nucleotide polymorphism (SNP) marker and a simple sequence repeat (SSR) marker.

[0175] “Open reading frame” is abbreviated ORF.

[0176] As used herein, “nucleic acid” means a polynucleotide and includes a single or a double-stranded polymer of deoxyribonucleotide or ribonucleotide bases. Nucleic acids may also include fragments and modified nucleotides. Thus, the terms “polynucleotide”, “nucleic acid sequence”, “nucleotide sequence” and “nucleic acid fragment” are used interchangeably to denote a polymer of RNA and/or DNA that is single- or double-stranded, optionally containing synthetic, non-natural, or altered nucleotide bases. Nucleotides (usually found in their 5'-monophosphate form) are referred to by their single letter designation as follows: “A” for adenosine or deoxyadenosine (for RNA or DNA, respectively), “C” for cytosine or deoxycytosine, “G” for guanosine or deoxyguanosine, “U” for uridine, “T” for deoxythymidine, “R” for purines (A or G), “Y” for pyrimidines (C or T), “K” for G or T, “H” for A or C or T, “I” for inosine, and “N” for any nucleotide.

[0177] The terms “subfragment that is functionally equivalent” and “functionally equivalent subfragment” are used interchangeably herein. These terms refer to a portion or subsequence of an isolated nucleic acid fragment in which the ability to alter gene expression or produce a certain phenotype is retained whether or not the fragment or subfragment encodes an active enzyme. For example, the fragment or subfragment can be used in the design of chimeric genes to produce the desired phenotype in a transformed plant. Chimeric genes can be designed for use in suppression by linking a nucleic acid fragment or subfragment thereof, whether or not it encodes an active enzyme, in the sense or antisense orientation relative to a plant promoter sequence.

[0178] The term “conserved domain” or “motif” means a set of amino acids conserved at specific positions along an aligned sequence of evolutionarily related proteins. While amino acids at other positions can vary between homologous proteins, amino acids that are highly conserved at specific positions indicate amino acids that are essential to the structure, the stability, or the activity of a protein. Because they are identified by their high degree of conservation in aligned sequences of a family of protein homologues, they can be used as identifiers, or “signatures”, to determine if a protein with a newly determined sequence belongs to a previously identified protein family.

[0179] Polynucleotide and polypeptide sequences, variants thereof, and the structural relationships of these sequences can be described by the terms “homology”, “homologous”, “substantially identical”, “substantially similar” and “corresponding substantially” which are used interchangeably herein. These refer to polypeptide or nucleic acid fragments wherein changes in one or more amino acids or nucleotide bases do not affect the function of the molecule, such as the ability to mediate gene expression or to produce a certain phenotype. These terms also refer to modification(s) of nucleic acid fragments that do not substantially alter the functional properties of the resulting nucleic acid fragment relative to the initial, unmodified fragment. These modifications include deletion, substitution, and/or insertion of one or more nucleotides in the nucleic acid fragment.

[0180] Substantially similar nucleic acid sequences encompassed may be defined by their ability to hybridize (under moderately stringent conditions, e.g., 0.5×SSC, 0.1% SDS, 60° C.) with the sequences exemplified herein, or to any portion of the nucleotide sequences disclosed herein and which are functionally equivalent to any of the nucleic acid

sequences disclosed herein. Stringency conditions can be adjusted to screen for moderately similar fragments, such as homologous sequences from distantly related organisms, to highly similar fragments, such as genes that duplicate functional enzymes from closely related organisms. Post-hybridization washes determine stringency conditions.

[0181] The term “selectively hybridizes” includes reference to hybridization, under stringent hybridization conditions, of a nucleic acid sequence to a specified nucleic acid target sequence to a detectably greater degree (e.g., at least 2-fold over background) than its hybridization to non-target nucleic acid sequences and to the substantial exclusion of non-target nucleic acids. Selectively hybridizing sequences typically have about at least 80% sequence identity, or 90% sequence identity, up to and including 100% sequence identity (i.e., fully complementary) with each other.

[0182] The term “stringent conditions” or “stringent hybridization conditions” includes reference to conditions under which a probe will selectively hybridize to its target sequence in an in vitro hybridization assay. Stringent conditions are sequence-dependent and will be different in different circumstances. By controlling the stringency of the hybridization and/or washing conditions, target sequences can be identified which are 100% complementary to the probe (homologous probing). Alternatively, stringency conditions can be adjusted to allow some mismatching in sequences so that lower degrees of similarity are detected (heterologous probing). Generally, a probe is less than about 1000 nucleotides in length, optionally less than 500 nucleotides in length.

[0183] Typically, stringent conditions will be those in which the salt concentration is less than about 1.5 M Na ion, typically about 0.01 to 1.0 M Na ion concentration (or other salt(s)) at pH 7.0 to 8.3, and at least about 30° C. for short probes (e.g., 10 to 50 nucleotides) and at least about 60° C. for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. Exemplary low stringency conditions include hybridization with a buffer solution of 30 to 35% formamide, 1 M NaCl, 1% SDS (sodium dodecyl sulphate) at 37° C., and a wash in 1× to 2×SSC (20×SSC=3.0 M NaCl/0.3 M trisodium citrate) at 50 to 55° C. Exemplary moderate stringency conditions include hybridization in 40 to 45% formamide, 1 M NaCl, 1% SDS at 37° C., and a wash in 0.5× to 1×SSC at 55 to 60° C. Exemplary high stringency conditions include hybridization in 50% formamide, 1 M NaCl, 1% SDS at 37° C., and a wash in 0.1×SSC at 60 to 65° C.

[0184] Specificity is typically the function of post-hybridization washes, the critical factors being the ionic strength and temperature of the final wash solution. For DNA-DNA hybrids, the T_m can be approximated from the equation of Meinkoth et al., (1984) *Anal Biochem* 138:267-284: $T_m = 81.5^\circ \text{C.} + 16.6 (\log M) + 0.41 (\% \text{ GC}) - 0.61 (\% \text{ form}) - 500/L$; where M is the molarity of monovalent cations, % GC is the percentage of guanosine and cytosine nucleotides in the DNA, % form is the percentage of formamide in the hybridization solution, and L is the length of the hybrid in base pairs. The T_m is the temperature (under defined ionic strength and pH) at which 50% of a complementary target sequence hybridizes to a perfectly matched probe. T_m is reduced by about 1° C. for each 1% of mismatching; thus, T_m hybridization and/or wash conditions can be adjusted to hybridize to sequences of the desired identity. For example,

if sequences with 90% identity are sought, the T_m can be decreased 10° C. Generally, stringent conditions are selected to be about 5° C. lower than the thermal melting point (T_m) for the specific sequence and its complement at a defined ionic strength and pH. However, severely stringent conditions can utilize a hybridization and/or wash at 1, 2, 3 or 4° C. lower than the thermal melting point (T_m); moderately stringent conditions can utilize a hybridization and/or wash at 6, 7, 8, 9 or 10° C. lower than the thermal melting point (T_m); low stringency conditions can utilize a hybridization and/or wash at 11, 12, 13, 14, 15 or 20° C. lower than the thermal melting point (T_m). Using the equation, hybridization and wash compositions, and desired T_m , those of ordinary skill will understand that variations in the stringency of hybridization and/or wash solutions are inherently described. If the desired degree of mismatching results in a T_m of less than 45° C. (aqueous solution) or 32° C. (formamide solution) it is preferred to increase the SSC concentration so that a higher temperature can be used. An extensive guide to the hybridization of nucleic acids is found in Tijssen, *Laboratory Techniques in Biochemistry and Molecular Biology—Hybridization with Nucleic Acid Probes*, Part I, Chapter 2 “Overview of principles of hybridization and the strategy of nucleic acid probe assays”, Elsevier, New York (1993); and *Current Protocols in Molecular Biology*, Chapter 2, Ausubel et al., Eds., Greene Publishing and Wiley-Interscience, New York (1995). Hybridization and/or wash conditions can be applied for at least 10, 30, 60, 90, 120 or 240 minutes.

[0185] “Sequence identity” or “identity” in the context of nucleic acid or polypeptide sequences refers to the nucleic acid bases or amino acid residues in two sequences that are the same when aligned for maximum correspondence over a specified comparison window.

[0186] The term “percentage of sequence identity” refers to the value determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide or polypeptide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the results by 100 to yield the percentage of sequence identity. Useful examples of percent sequence identities include, but are not limited to, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90% or 95%, or any integer percentage from 50% to 100%. These identities can be determined using any of the programs described herein.

[0187] Sequence alignments and percent identity or similarity calculations may be determined using a variety of comparison methods designed to detect homologous sequences including, but not limited to, the MegAlign™ program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Within the context of this application it will be understood that where sequence analysis software is used for analysis, that the results of the analysis will be based on the “default values” of the program referenced, unless otherwise specified. As used herein

“default values” will mean any set of values or parameters that originally load with the software when first initialized.

[0188] The “Clustal V method of alignment” corresponds to the alignment method labeled Clustal V (described by Higgins and Sharp, (1989) *CABIOS* 5:151-153; Higgins et al., (1992) *Comput Appl Biosci* 8:189-191) and found in the MegAlign™ program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). For multiple alignments, the default values correspond to GAP PENALTY=10 and GAP LENGTH PENALTY=10. Default parameters for pairwise alignments and calculation of percent identity of protein sequences using the Clustal method are KTUPLE=1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5. For nucleic acids these parameters are KTUPLE=2, GAP PENALTY=5, WINDOW=4 and DIAGONALS SAVED=4. After alignment of the sequences using the Clustal V program, it is possible to obtain a “percent identity” by viewing the “sequence distances” table in the same program.

[0189] The “Clustal W method of alignment” corresponds to the alignment method labeled Clustal W (described by Higgins and Sharp, (1989) *CABIOS* 5:151-153; Higgins et al., (1992) *Comput Appl Biosci* 8:189-191) and found in the MegAlign™ v6.1 program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Default parameters for multiple alignment (GAP PENALTY=10, GAP LENGTH PENALTY=0.2, Delay Divergen Seqs (%)=30, DNA Transition Weight=0.5, Protein Weight Matrix=Gonnet Series, DNA Weight Matrix=IUB). After alignment of the sequences using the Clustal W program, it is possible to obtain a “percent identity” by viewing the “sequence distances” table in the same program.

[0190] Unless otherwise stated, sequence identity/similarity values provided herein refer to the value obtained using GAP Version 10 (GCG, Accelrys, San Diego, CA) using the following parameters: % identity and % similarity for a nucleotide sequence using a gap creation penalty weight of 50 and a gap length extension penalty weight of 3, and the nwsgapdna.cmp scoring matrix; % identity and % similarity for an amino acid sequence using a GAP creation penalty weight of 8 and a gap length extension penalty of 2, and the BLOSUM62 scoring matrix (Henikoff and Henikoff, (1989) *Proc. Natl. Acad. Sci. USA* 89:10915). GAP uses the algorithm of Needleman and Wunsch, (1970) *J Mol Biol* 48:443-53, to find an alignment of two complete sequences that maximizes the number of matches and minimizes the number of gaps. GAP considers all possible alignments and gap positions and creates the alignment with the largest number of matched bases and the fewest gaps, using a gap creation penalty and a gap extension penalty in units of matched bases.

[0191] BLAST® is a searching algorithm provided by the National Center for Biotechnology Information (NCBI) used to find regions of similarity between biological sequences. The program compares nucleotide or protein sequences to sequence databases and calculates the statistical significance of matches to identify sequences having sufficient similarity to a query sequence such that the similarity would not be predicted to have occurred randomly. BLAST® reports the identified sequences and their local alignment to the query sequence.

[0192] It is well understood by one skilled in the art that many levels of sequence identity are useful in identifying polypeptides from other species or modified naturally or

synthetically wherein such polypeptides have the same or similar function or activity. Useful examples of percent identities include, but are not limited to, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90% or 95%, or any integer percentage from 50% to 100%. Indeed, any integer amino acid identity from 50% to 100% may be useful in describing the present invention, such as 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99%.

[0193] “Gene” refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence. “Native gene” refers to a gene as found in nature with its own regulatory sequences. “Chimeric gene” refers to any gene that is not a native gene, comprising regulatory and coding sequences that are not found together in nature. Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that found in nature, or at a different genetic locus than that found in nature. A “foreign” gene refers to a gene not normally found in the host organism, but that is introduced into the host organism by gene transfer. Foreign genes can comprise native genes inserted into a non-native organism, or chimeric genes.

[0194] A “mutated gene” is a native gene that has been altered through human intervention. Such a “mutated gene” has a sequence that differs from the sequence of the corresponding native gene by at least one nucleotide addition, deletion, or substitution. In certain embodiments of the invention, the mutated gene comprises an alteration that results from a double-strand-break-inducing agent as disclosed herein.

[0195] A “transgene” is a gene that has been introduced into the genome by a transformation procedure. A transgene can, for example encode one or more proteins or RNA that is not translated into protein. However, a transgene of the invention need not encode a protein and/or non-translated RNA. In certain embodiments of the invention, the transgene comprises one or more chimeric genes, including chimeric genes comprising, for example, a gene of interest, phenotypic marker, a selectable marker, and a DNA for gene silencing.

[0196] As used herein, a “targeted mutation” is mutation in a native gene that was made by altering a target sequence within the native gene using a method involving a double-strand-break-inducing agent that is capable of inducing a double-strand break in the DNA of the target sequence as disclosed herein or known in the art.

[0197] The term “genome” as it applies to a plant cells encompasses not only chromosomal DNA found within the nucleus, but organelle DNA found within subcellular components (e.g., mitochondria, or plastid) of the cell.

[0198] A “codon-modified gene” or “codon-preferred gene” or “codon-optimized gene” is a gene having its frequency of codon usage designed to mimic the frequency of preferred codon usage of the host cell.

[0199] An “allele” is one of several alternative forms of a gene occupying a given locus on a chromosome. When all

the alleles present at a given locus on a chromosome are the same, that plant is homozygous at that locus. If the alleles present at a given locus on a chromosome differ, that plant is heterozygous at that locus.

[0200] “Coding sequence” refers to a polynucleotide sequence which codes for a specific amino acid sequence. “Regulatory sequences” refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences may include, but are not limited to: promoters, translation leader sequences, 5' untranslated sequences, 3' untranslated sequences, introns, polyadenylation recognition sequences, RNA processing sites, effector binding sites, and stem-loop structures.

[0201] “A plant-optimized nucleotide sequence” is nucleotide sequence that has been optimized for increased expression in plants, particularly for increased expression in plants or in one or more plants of interest. For example, a plant-optimized nucleotide sequence can be synthesized by modifying a nucleotide sequence encoding a protein such as, for example, double-strand-break-inducing agent (e.g., an endonuclease) as disclosed herein, using one or more plant-preferred codons for improved expression. See, for example, Campbell and Gowri (1990) *Plant Physiol.* 92:1-11 for a discussion of host-preferred codon usage. Methods are available in the art for synthesizing plant-preferred genes. See, for example, U.S. Pat. Nos. 5,380,831, and 5,436,391, and Murray et al. (1989) *Nucleic Acids Res.* 17:477-498, herein incorporated by reference. Additional sequence modifications are known to enhance gene expression in a plant host. These include, for example, elimination of: one or more sequences encoding spurious polyadenylation signals, one or more exon-intron splice site signals, one or more transposon-like repeats, and other such well-characterized sequences that may be deleterious to gene expression. The G-C content of the sequence may be adjusted to levels average for a given plant host, as calculated by reference to known genes expressed in the host plant cell. When possible, the sequence is modified to avoid one or more predicted hairpin secondary mRNA structures. Thus, “a plant-optimized nucleotide sequence” of the present invention comprises one or more of such sequence modifications.

[0202] “Promoter” refers to a DNA sequence capable of controlling the expression of a coding sequence or functional RNA. The promoter sequence consists of proximal and more distal upstream elements, the latter elements often referred to as enhancers. An “enhancer” is a DNA sequence that can stimulate promoter activity, and may be an innate element of the promoter or a heterologous element inserted to enhance the level or tissue-specificity of a promoter. Promoters may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, and/or comprise synthetic DNA segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental conditions. It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, DNA fragments of some variation may have identical promoter activity. Promoters that cause a gene to be expressed in most cell types at most times are com-

monly referred to as “constitutive promoters”. New promoters of various types useful in plant cells are constantly being discovered; numerous examples may be found in the compilation by Okamuro and Goldberg, (1989) In *The Biochemistry of Plants*, Vol. 115, Stumpf and Conn, eds (New York, NY: Academic Press), pp. 1-82.

[0203] “Translation leader sequence” refers to a polynucleotide sequence located between the promoter sequence of a gene and the coding sequence. The translation leader sequence is present in the fully processed mRNA upstream of the translation start sequence. The translation leader sequence may affect processing of the primary transcript to mRNA, mRNA stability or translation efficiency. Examples of translation leader sequences have been described (e.g., Turner and Foster, (1995) *Mol Biotechnol* 3:225-236).

[0204] “3' non-coding sequences”, “transcription terminator” or “termination sequences” refer to DNA sequences located downstream of a coding sequence and include polyadenylation recognition sequences and other sequences encoding regulatory signals capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor. The use of different 3' non-coding sequences is exemplified by Ingelbrecht et al., (1989) *Plant Cell* 1:671-680.

[0205] “RNA transcript” refers to the product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When the RNA transcript is a perfect complementary copy of the DNA sequence, it is referred to as the primary transcript. A RNA transcript is referred to as the mature RNA when it is a RNA sequence derived from post-transcriptional processing of the primary transcript. “Messenger RNA” or “mRNA” refers to the RNA that is without introns and that can be translated into protein by the cell. “cDNA” refers to a DNA that is complementary to, and synthesized from, a mRNA template using the enzyme reverse transcriptase. The cDNA can be single-stranded or converted into double-stranded form using the Klenow fragment of DNA polymerase I. “Sense” RNA refers to RNA transcript that includes the mRNA and can be translated into protein within a cell or in vitro. “Antisense RNA” refers to an RNA transcript that is complementary to all or part of a target primary transcript or mRNA, and that blocks the expression of a target gene (see, e.g., U.S. Pat. No. 5,107,065). The complementarity of an antisense RNA may be with any part of the specific gene transcript, i.e., at the 5' non-coding sequence, 3' non-coding sequence, introns, or the coding sequence. “Functional RNA” refers to antisense RNA, ribozyme RNA, or other RNA that may not be translated but yet has an effect on cellular processes. The terms “complement” and “reverse complement” are used interchangeably herein with respect to mRNA transcripts, and are meant to define the antisense RNA of the message.

[0206] The term “operably linked” refers to the association of nucleic acid sequences on a single nucleic acid fragment so that the function of one is regulated by the other. For example, a promoter is operably linked with a coding sequence when it is capable of regulating the expression of that coding sequence (i.e., the coding sequence is under the transcriptional control of the promoter). Coding sequences can be operably linked to regulatory sequences in a sense or antisense orientation. In another example, the complementary RNA regions can be operably linked, either directly or indirectly, 5' to the target mRNA, or 3' to the target mRNA,

or within the target mRNA, or a first complementary region is 5' and its complement is 3' to the target mRNA.

[0207] Standard recombinant DNA and molecular cloning techniques used herein are well known in the art and are described more fully in Sambrook et al., *Molecular Cloning: A Laboratory Manual*; Cold Spring Harbor Laboratory: Cold Spring Harbor, NY (1989). Transformation methods are well known to those skilled in the art and are described infra.

[0208] “PCR” or “polymerase chain reaction” is a technique for the synthesis of specific DNA segments and consists of a series of repetitive denaturation, annealing, and extension cycles. Typically, a double-stranded DNA is heat denatured, and two primers complementary to the 3' boundaries of the target segment are annealed to the DNA at low temperature, and then extended at an intermediate temperature. One set of these three consecutive steps is referred to as a “cycle”.

[0209] The term “recombinant” refers to an artificial combination of two otherwise separated segments of sequence, e.g., by chemical synthesis, or manipulation of isolated segments of nucleic acids by genetic engineering techniques.

[0210] The terms “plasmid”, “vector” and “cassette” refer to an extra chromosomal element often carrying genes that are not part of the central metabolism of the cell, and usually in the form of double-stranded DNA. Such elements may be autonomously replicating sequences, genome integrating sequences, phage, or nucleotide sequences, in linear or circular form, of a single- or double-stranded DNA or RNA, derived from any source, in which a number of nucleotide sequences have been joined or recombined into a unique construction which is capable of introducing a polynucleotide of interest into a cell. “Transformation cassette” refers to a specific vector containing a foreign gene and having elements in addition to the foreign gene that facilitates transformation of a particular host cell. “Expression cassette” refers to a specific vector containing a foreign gene and having elements in addition to the foreign gene that allow for expression of that gene in a foreign host.

[0211] The terms “recombinant DNA molecule”, “recombinant construct”, “expression construct”, “chimeric construct”, “construct”, and “recombinant DNA construct” are used interchangeably herein. A recombinant construct comprises an artificial combination of nucleic acid fragments, e.g., regulatory and coding sequences that are not all found together in nature. For example, a chimeric construct may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that found in nature. Such a construct may be used by itself or may be used in conjunction with a vector. If a vector is used, then the choice of vector is dependent upon the method that will be used to transform host cells as is well known to those skilled in the art. For example, a plasmid vector can be used. The skilled artisan is well aware of the genetic elements that must be present on the vector in order to successfully transform, select and propagate host cells. The skilled artisan will also recognize that different independent transformation events may result in different levels and patterns of expression (Jones et al., (1985) *EMBO J* 4:2411-2418; De Almeida et al., (1989) *Mol Gen Genetics* 218:78-86), and thus that multiple events are typically screened in order to obtain lines

displaying the desired expression level and pattern. Such screening may be accomplished standard molecular biological, biochemical, and other assays including Southern analysis of DNA, Northern analysis of mRNA expression, PCR, real time quantitative PCR (qPCR), reverse transcription PCR (RT-PCR), immunoblotting analysis of protein expression, enzyme or activity assays, and/or phenotypic analysis.

[0212] The term “expression”, as used herein, refers to the production of a functional end-product (e.g., an mRNA or a protein) in either precursor or mature form.

[0213] The term “introduced” means providing a nucleic acid (e.g., expression construct) or protein into a cell. Introduced includes reference to the incorporation of a nucleic acid into a eukaryotic or prokaryotic cell where the nucleic acid may be incorporated into the genome of the cell, and includes reference to the transient provision of a nucleic acid or protein to the cell. Introduced includes reference to stable or transient transformation methods, as well as sexually crossing. Thus, “introduced” in the context of inserting a nucleic acid fragment (e.g., a recombinant DNA construct/ expression construct) into a cell, means “transfection” or “transformation” or “transduction” and includes reference to the incorporation of a nucleic acid fragment into a eukaryotic or prokaryotic cell where the nucleic acid fragment may be incorporated into the genome of the cell (e.g., chromosome, plasmid, plastid, or mitochondrial DNA), converted into an autonomous replicon, or transiently expressed (e.g., transfected mRNA).

[0214] “Mature” protein refers to a post-translationally processed polypeptide (i.e., one from which any pre- or propeptides present in the primary translation product have been removed). “Precursor” protein refers to the primary product of translation of mRNA (i.e., with pre- and propeptides still present). Pre- and propeptides may be but are not limited to intracellular localization signals.

[0215] “Stable transformation” refers to the transfer of a nucleic acid fragment into a genome of a host organism, including both nuclear and organellar genomes, resulting in genetically stable inheritance. In contrast, “transient transformation” refers to the transfer of a nucleic acid fragment into the nucleus, or other DNA-containing organelle, of a host organism resulting in gene expression without integration or stable inheritance. Host organisms containing the transformed nucleic acid fragments are referred to as “transgenic” organisms.

[0216] As used herein, “transgenic” refers to a plant or a cell which comprises within its genome a heterologous polynucleotide. Typically, the heterologous polynucleotide is stably integrated within the genome such that the polynucleotide is passed on to successive generations. The heterologous polynucleotide may be integrated into the genome alone or as part of an expression construct. Transgenic is used herein to include any cell, cell line, callus, tissue, plant part or plant, the genotype of which has been altered by the presence of heterologous nucleic acid including those transgenics initially so altered as well as those created by sexual crosses or asexual propagation from the initial transgenic. The term “transgenic” as used herein does not encompass the alteration of the genome (chromosomal or extra-chromosomal) by conventional plant breeding methods or by naturally occurring events such as random cross-fertilization, non-recombinant viral infection, non-recombinant bacterial transformation, non-recombinant transposition, or spontaneous mutation.

[0217] The term “plant” refers to whole plants, plant organs, plant tissues, seeds, plant cells, seeds and progeny of the same. Plant cells include, without limitation, cells from seeds, suspension cultures, embryos, meristematic regions, callus tissue, leaves, roots, shoots, gametophytes, sporophytes, pollen and microspores. Plant parts include differentiated and undifferentiated tissues including, but not limited to roots, stems, shoots, leaves, pollens, seeds, tumor tissue and various forms of cells and culture (e.g., single cells, protoplasts, embryos, and callus tissue). The plant tissue may be in plant or in a plant organ, tissue or cell culture. The term “plant organ” refers to plant tissue or a group of tissues that constitute a morphologically and functionally distinct part of a plant. The term “genome” refers to the entire complement of genetic material (genes and non-coding sequences) that is present in each cell of an organism, or virus or organelle; and/or a complete set of chromosomes inherited as a (haploid) unit from one parent. “Progeny” comprises any subsequent generation of a plant.

[0218] A “fertile plant” is a plant that is capable of producing a progeny plant. In certain embodiments of the invention, a fertile plant is a plant that produces viable male and female gametes and is self fertile. Such a self-fertile plant can produce a progeny plant without the contribution from any other plant of a gamete and the genetic material contained therein. Other embodiments of the invention can involve the use of a plant that is not self fertile because the plant does not produce male or female gametes that are viable or otherwise capable of fertilization. As used herein, a “male sterile plant” is a plant that does not produce male gametes that are viable or otherwise capable of fertilization. As used herein, a “female sterile plant” is a plant that does not produce female gametes that are viable or otherwise capable of fertilization. It is recognized that male and female sterile plants can be female and male fertile, respectively. It is further recognized that a male fertile (but female sterile) plant can produce viable progeny when crossed with a female fertile plant and that a female fertile (but male sterile) plant can produce viable progeny when crossed with a male fertile plant.

[0219] A “centimorgan” (CM) or “map unit” is the distance between two linked genes, markers, target sites, loci, or any pair thereof, wherein 1% of the products of meiosis are recombinant. Thus, a centimorgan is equivalent to a distance equal to an 1% average recombination frequency between the two linked genes, markers, target sites, loci, or any pair thereof.

[0220] The present invention finds use in the breeding of plants comprising two to more transgenic traits. Currently, transgenic traits are randomly inserted throughout the plant genome as a consequence of transformation systems based on *Agrobacterium*, biolistics, or other commonly used procedures. More recently, gene targeting protocols have been developed that enable directed transgene insertion. One important technology, site-specific integration (SSI) enables the targeting of a transgene to the same chromosomal location as a previously inserted transgene. Custom-designed meganucleases and custom-designed zinc finger meganucleases allow researchers to design nucleases to target specific chromosomal locations, and these reagents allow the targeting of transgenes at the chromosomal site cleaved by these nucleases.

[0221] As disclosed herein, nuclease-mediated gene targeting can be used in methods for producing complex

transgenic trait loci comprising multiple transgenes. In one embodiment of the invention, a complex transgenic trait locus is a locus that has multiple transgenes genetically linked to each other. By inserting independent transgenes within 1, 2 or even 5 centimorgans (cM) from each other, the transgenes can be bred as single genetic locus. FIG. 7 depicts the process of how two traits could be integrated into the genome at a genetic distance of, for example, 0.2 cM from each other in independent transformation runs or in sequential transformations (e.g., transformation and re-transformation). After selecting the events, plants containing the two events can be crossed to form an F1 that contains the events on different chromosomes. In progeny from these F1 (F2 or BC1) 1/500 progeny would have the two different transgenes recombined onto the same chromosome. The complex locus could then be bred as single genetic locus with both transgene traits. This process could be repeated to stack as many traits as desired.

[0222] The present invention provides methods for producing complex transgenic trait loci at selected genomic regions to simplify breeding with multiple transgenes. To initiate the development of a complex transgenic trait locus, a region of the genome is first selected. Second, the sequence of nearby genomic regions is compiled and nuclease reagents designed to facilitate targeting additional transgenes to those closely linked sites. Subsequently, algorithms for nuclease design such as, for example, those described in U.S. Patent Application Publication No. 2007/0117128 A1 are used to select potential target sites. Additional bioinformatic analysis such as, for example, copy number of the site in the target genome, location of the site relative to known gene coding regions and other factors could be used to filter the sites to a subset of preferred sites. Nucleases could then be used to target new transgenes to these preferred sites using published protocols See, for example, Halluin et al. (2008) *Plant Biotechnol. J.* 6:93-102; Shukla et al. (2009) *Nature* doi: 10.1038/nature07992; Wright et al. *Plant J.* (2005) 44:693-705; and WO 2009/006297; all of which are herein incorporated by reference.

[0223] In a first aspect, the present invention provides methods for producing in a plant a complex transgenic trait locus comprising at least two altered target sequences in a genomic region of interest. In one embodiment, the methods involve selecting a genomic region in a plant that comprises a first target sequence and a second target sequence. Generally, the first target sequence and the second target sequence are separated from each other by about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8., 0.9, 1, 2, 3, 4, or 5 centimorgans (cM) in the genome of the plant. In certain embodiments of the invention, the first and second target sequences are physically linked to a polynucleotide of interest such as, for example, a transgene, native gene, or a gene with a targeted mutation, that is within about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8., 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or 21 cM of the first and/or the second target sequence.

[0224] The methods of the invention further involve providing a first double-strand-break-inducing agent and a second double-strand-break-inducing agent. The first double-strand-break-inducing agent is capable of inducing a first double-strand break in DNA comprising the first target sequence, and the second double-strand-break-inducing agent is capable of inducing a second double-strand break in DNA comprising the second target sequence. The methods

of the invention do not depend on a particular double-strand-break-inducing agent but only that the double-strand-break-inducing agent is capable of inducing a double-strand break in DNA in a target sequence of the invention. Any such double-strand-break-inducing agent that is disclosed herein or known in the art can be used in the methods of the present invention.

[0225] Additionally, the methods involve contacting at least one plant cell with the first double-strand-break-inducing agent, identifying a cell comprising a first alteration at the first target sequence, and then recovering a first fertile plant from the cell comprising the first alteration. The first fertile plant also comprises the first alteration. Additionally, the method involves contacting at least one plant cell with the second double-strand-break-inducing agent, identifying a cell comprising a second alteration at the second target sequence, and then recovering a second fertile plant from the cell comprising the second alteration. The method further involves obtaining a fertile progeny plant from the second fertile plant, wherein the fertile progeny plant comprises both the first and second alterations in physical linkage.

[0226] In one embodiment of this method, the fertile progeny plant is obtained by crossing the first fertile plant and the second fertile plant and selecting for a fertile progeny plant comprising both the first and second alterations in physical linkage. In another embodiment, a cell of the first fertile plant, or progeny thereof comprising the first alteration, is contacted with the second double-strand-break-inducing agent, and the second fertile plant comprises both the first and second alterations, which may or may not be physically linked. If necessary, the second fertile plant can be selfed and a fertile progeny plant selected comprising both the first and second alterations in physical linkage.

[0227] The first and second alterations are selected from the group consisting of (i) replacement of at least one nucleotide, (ii) a deletion of at least one nucleotide, (iii) an insertion of at least one nucleotide, and (iv) any combination of (i)-(iii). In one embodiment of the invention, the first and/or the second alterations comprise insertion of a DNA sequence of interest including, but not limited to, a DNA for gene silencing, a DNA encoding a phenotypic marker, and a DNA encoding a protein providing an agronomic advantage. In another embodiment, the first and/or the second alterations comprise a targeted mutation in a native gene.

[0228] In a like manner, the methods disclosed herein can be used to produce in a plant a complex transgenic trait locus comprising 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more altered target sequences in physical linkage in a genomic region of interest comprising 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more target sequences of interest. Each additional target sequence of interest in the genomic region of interest can be recognized and cleaved by a double-strand-break-inducing agent essentially as described above.

[0229] For example, a third DNA sequence of interest is inserted into a third target sequence by contacting at least one cell of a plant with a third double-strand-break-inducing agent and a third DNA molecule comprising the DNA sequence of interest, and then identifying a cell comprising the DNA sequence of interest. The method can further comprise recovering a fertile plant comprising the third DNA sequence of interest. In one embodiment, the cell comprising the third DNA sequence of interest comprises the first alteration, the second alteration, or both the first alteration and the second alteration. The method of the

invention can further comprising producing a fertile plant comprising the first alteration, the second alteration, and the third DNA sequence of interest in physical linkage. In another embodiment, the fertile plant comprising the first alteration, the second alteration, and the third DNA sequence of interest is produced by crossing the fertile plant comprising the first and second alterations with a second fertile plant comprising the third DNA sequence of interest, and selecting a fertile progeny plant from the crossing, wherein the fertile progeny plant comprises the first alteration, the second alteration, and the third DNA sequence of interest in physical linkage.

[0230] The fertile plant comprising the first alteration, the second alteration, and the third DNA sequence of interest can be produced, for example, by: (i) contacting a cell comprising the first alteration and the second alteration with the third double-strand-break-inducing agent; (ii) identifying a cell from (i) comprising the third DNA sequence of interest, wherein the cell comprises the first alteration and the second alteration, and wherein the first alteration, the second alteration, and the third DNA sequence of interest are physically linked; and (iii) recovering a fertile plant comprising in physical linkage the first alteration, the second alteration, and the third DNA sequence of interest.

[0231] In another embodiment of the invention, the methods for producing in a plant a complex transgenic trait locus comprising at least two altered target sequences in a genomic region of interest that involve obtaining a first fertile plant comprising a first altered target site at the genomic region of interest and a second fertile plant comprising a second altered target site at the genomic region of interest. In this method, the first altered target sequence originated from a first target sequence that is recognized and cleaved by a first double-strand-break-inducing agent, and the second altered target sequence originated from a second target sequence that is recognized and cleaved by a second double-strand-break-inducing agent. The second method further involves crossing the first fertile plant and the second fertile plant, and then selecting from the crossing a fertile progeny plant comprising the first alteration and the second alteration in physical linkage.

[0232] The second method can optionally involve crossing the fertile progeny plant with an additional fertile plant that comprises at least a third altered target sequence in the genomic region of interest and then selecting from the crossing a fertile progeny plant comprising the first alteration, the second and the at least third alteration in physical linkage. Like the first and second altered target sequences, the third altered target sequence originated from a third target sequence that is recognized and cleaved by a third double-strand-break-inducing agent. In a like manner, a complex transgenic trait locus can be produced comprising 4, 5, 6, 7, 8, 9, 10, or more altered target sequences in physical linkage in the genomic region of interest.

[0233] In another aspect, the present invention provides complex transgenic trait loci in plants as well as plants, plant parts, plant cells, and seeds comprising at least one complex transgenic trait locus of the invention. A complex transgenic trait locus of the invention comprises at least two altered target sequences that are genetically linked to a polynucleotide of interest. Such altered target sequences originated from a corresponding target sequence that is recognized and cleaved by a double-strand-break-inducing agent using, for example, the methods disclosed herein. The altered target

sequences comprise an alteration such as, for example, replacement of at least one nucleotide in the target sequence, a deletion of at least one nucleotide in the target sequence, an insertion of at least one nucleotide in the target sequence, or any combination thereof. The polynucleotide interest can be, for example, a transgene, a native gene, and a mutated gene. The present invention provides plants, plant parts, plant cells, and seeds comprising at least one complex transgenic trait locus of the invention.

[0234] In one embodiment, a complex transgenic trait locus of the invention comprises at least one altered target sequence comprising a recombinant DNA molecule. Recombinant DNA molecules of the invention include, but are not limited to, a DNA for gene silencing, a DNA encoding a phenotypic marker, and a DNA encoding a protein providing an agronomic advantage.

[0235] Generally, each of the altered target sites of the complex transgenic trait locus are located within about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or 21 centimorgan (CM) of the polynucleotide of interest.

[0236] The methods of the present invention involve the use of one or more double-strand break inducing agents. A double-strand break inducing agent of the present invention is any agent that recognizes and/or binds to a specific polynucleotide recognition sequence to produce a break in the target sequence at or near the recognition sequence. Examples of double-strand break inducing agents include, but are not limited to, endonucleases, TAL effector nucleases, and zinc finger nucleases, and include modified derivatives, variants, and fragments thereof.

[0237] A recognition sequence is any polynucleotide sequence that is specifically recognized and/or bound by a double-strand break inducing agent. The length of the recognition site sequence can vary, and includes, for example, sequences that are at least 4, 6, 8, 10, 12, 14, 16, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70 or more nucleotides in length.

[0238] It is possible that the recognition site could be palindromic, that is, the sequence on one strand reads the same in the opposite direction on the complementary strand. The nick/cleavage site could be within the recognition sequence or the nick/cleavage site could be outside of the recognition sequence. In another variation, the cleavage could occur at nucleotide positions immediately opposite each other to produce a blunt end cut or, in other cases, the incisions could be staggered to produce single-stranded overhangs, also called “sticky ends”, which can be either 5' overhangs, or 3' overhangs. The recognition sequence can be endogenous or exogenous. When the recognition site is an endogenous sequence, it may be a recognition sequence recognized by a naturally-occurring, or native double-strand break inducing agent. Alternatively, an endogenous recognition site could be recognized and/or bound by a modified or engineered double-strand break inducing agent designed or selected to specifically recognize the endogenous recognition sequence to produce a double-strand break. A modified double-strand break inducing agent can be derived from a native, naturally-occurring double-strand break inducing agent or it could be artificially created or synthesized.

[0239] A variety of methods are available to identify those cells having an altered genome at or near the recognition

sequence without using a screenable marker phenotype. Such methods can be viewed as directly analyzing a recognition sequence to detect any change in the recognition sequence, including but not limited to PCR methods, sequencing methods, nuclease digestion, Southern blots, and any combination thereof.

[0240] Proteins may be altered in various ways including amino acid substitutions, deletions, truncations, and insertions. Methods for such manipulations are generally known. For example, amino acid sequence variants of the protein(s) can be prepared by mutations in the DNA. Methods for mutagenesis and nucleotide sequence alterations include, for example, Kunkel, (1985) *Proc. Natl. Acad. Sci. USA* 82:488-92; Kunkel et al., (1987) *Meth Enzymol* 154:367-82; U.S. Pat. No. 4,873,192; Walker and Gaastra, eds. (1983) *Techniques in Molecular Biology* (MacMillan Publishing Company, New York) and the references cited therein. Guidance regarding amino acid substitutions not likely to affect biological activity of the protein is found, for example, in the model of Dayhoff et al., (1978) *Atlas of Protein Sequence and Structure* (Natl Biomed Res Found, Washington, D.C.). Conservative substitutions, such as exchanging one amino acid with another having similar properties, may be preferable. Conservative deletions, insertions, and amino acid substitutions are not expected to produce radical changes in the characteristics of the protein, and the effect of any substitution, deletion, insertion, or combination thereof can be evaluated by routine screening assays. Assays for double-strand-break-inducing activity are known and generally measure the overall activity and specificity of the agent on DNA substrates containing recognition sites.

[0241] Endonucleases are enzymes that cleave the phosphodiester bond within a polynucleotide chain, and include restriction endonucleases that cleave DNA as specific sites without damaging the bases. Restriction endonucleases include Type I, Type II, Type III, and Type IV endonucleases, which further include subtypes. In the Type I and Type III systems, both the methylase and restriction activities are contained in a single complex.

[0242] Type I and Type III restriction endonucleases recognize specific recognition sites, but typically cleave at a variable position from the recognition site, which can be hundreds of base pairs away from the recognition site. In Type II systems the restriction activity is independent of any methylase activity, and cleavage typically occurs at specific sites within or near to the recognition site. Most Type II enzymes cut palindromic sequences, however Type IIa enzymes recognize non-palindromic recognition sites and cleave outside of the recognition site, Type IIb enzymes cut sequences twice with both sites outside of the recognition site, and Type IIc enzymes recognize an asymmetric recognition site and cleave on one side and at a defined distance of about 1-20 nucleotides from the recognition site.

[0243] Type IV restriction enzymes target methylated DNA. Restriction enzymes are further described and classified, for example in the REBASE database (webpage at rebase.neb.com; Roberts et al., (2003) *Nucleic Acids Res* 31:418-20), Roberts et al., (2003) *Nucleic Acids Res* 31:1805-12, and Belfort et al., (2002) in *Mobile DNA II*, pp. 761-783, Eds. Craigie et al., (ASM Press, Washington, DC).

[0244] Endonucleases also include meganucleases, also known as homing endonucleases (HEases), which like restriction endonucleases, bind and cut at a specific recognition sequence, however the recognition sites for mega-

nucleases are typically longer, about 18 bp or more. Meganucleases have been classified into four families based on conserved sequence motifs, the families are the LAGLI-DADG, GIY-YIG, H—N—H, and His-Cys box families. These motifs participate in the coordination of metal ions and hydrolysis of phosphodiester bonds. HEases are notable for their long recognition sites, and for tolerating some sequence polymorphisms in their DNA substrates. The naming convention for meganuclease is similar to the convention for other restriction endonuclease. Meganucleases are also characterized by prefix F-, I-, or PI-for enzymes encoded by free-standing ORFs, introns, and inteins, respectively. For example, intron-, intein-, and freestanding gene encoded meganuclease from *Saccharomyces cerevisiae* are denoted I-SceI, PI-SceI, and F-SceI, respectively. Meganuclease domains, structure and function are known, see for example, Guhan and Muniyappa (2003) *Crit Rev Biochem Mol Biol* 38:199-248; Lucas et al., (2001) *Nucleic Acids Res* 29:960-9; Jurica and Stoddard, (1999) *Cell Mol Life Sci* 55:1304-26; Stoddard, (2006) *Q Rev Biophys* 38:49-95; and Moure et al., (2002) *Nat Struct Biol* 9:764. In some examples a naturally occurring variant, and/or engineered derivative meganuclease is used. Methods for modifying the kinetics, cofactor interactions, expression, optimal conditions, and/or recognition site specificity, and screening for activity are known, see for example, Epinat et al., (2003) *Nucleic Acids Res* 31:2952-62; Chevalier et al., (2002) *Mol Cell* 10:895-905; Gimble et al., (2003) *Mol Biol* 334:993-1008; Seligman et al., (2002) *Nucleic Acids Res* 30:3870-9; Sussman et al., (2004) *J Mol Biol* 342:31-41; Rosen et al., (2006) *Nucleic Acids Res* 34:4791-800; Chames et al., (2005) *Nucleic Acids Res* 33: e178; Smith et al., (2006) *Nucleic Acids Res* 34: e149; Gruen et al., (2002) *Nucleic Acids Res* 30: e29; Chen and Zhao, (2005) *Nucleic Acids Res* 33: e154; WO2005105989; WO2003078619; WO2006097854; WO2006097853; WO2006097784; and WO2004031346.

[0245] Any meganuclease can be used as a double-strand break inducing agent including, but not limited to, I-SceI, I-SceII, I-SceIII, I-SceIV, I-SceV, I-SceVI, I-SceVII, I-CeuI, I-CeuAIP, I-Crel, I-CrepsbIP, I-CrepsbIIP, I-CrepsbIIIP, I-CrepsbIVP, I-TliI, I-PpoI, PI-PspI, F-SceI, F-SceII, F-SuVI, F-TevI, F-TevII, I-Amal, I-Anil, I-Chul, I-Cmoel, I-Cpal, I-Cpall, I-Csml, I-Cvul, I-CvuAIP, I-DdII, I-DdIII, I-Dirl, I-Dmol, I-Hmul, I-HmulI, I-HsNIP, I-Llal, I-Msol, I-Naal, I-NanI, I-NclIP, I-NgrIP, I-NitI, I-Njal, I-Nsp236IP, I-PakI, I-PbolIP, I-PculIP, I-PcuAl, I-PcuVI, I-PgrIP, I-PobIP, I-PorI, I-PorIIP, I-PbpIP, I-SpBetalP, I-Scal, I-SexIP, I-SnelP, I-SpomI, I-SpomCP, I-SpomIP, I-SpomIIP, I-SquIP, I-Ssp68031, I-SthPhiIP, I-SthPhiST3P, I-SthPhiSTe3bP, I-TdelIP, I-TevI, I-TevII, I-TevIII, I-UarAP, I-UarHGPAIP, I-UarHGPA13P, I-VinIP, I-ZbilIP, PI-MtuI, PI-MtuHIIP, PI-MtuHIIIP, PI-Pful, PI-Pfull, PI-Pkol, PI-Pkoll, PI-Rma43812IP, PI-SpBetalP, PI-SceI, PI-Tful, PI-Tfull, PI-Thyl, PI-TliI, PI-TliII, or any variant or derivative thereof.

[0246] The endonuclease can be a modified endonuclease that binds a non-native or exogenous recognition sequence and does not bind a native or endogenous recognition sequence. Modification of the endonuclease can be as little as one nucleotide. A modified endonuclease is not capable of making a double-strand break within a wild-type target sequence. A wild-type (i.e., prior to being modified) endonuclease is capable of making a double-strand break within the wild-type target sequence.

[0247] The endonuclease can be provided via a polynucleotide encoding the endonuclease. Such a polynucleotide encoding an endonuclease can be modified to substitute codons having a higher frequency of usage in a plant, as compared to the naturally occurring polynucleotide sequence. For example the polynucleotide encoding the endonuclease can be modified to substitute codons having a higher frequency of usage in a maize or soybean plant, as compared to the naturally occurring polynucleotide sequence.

[0248] A site-specific recombinase, also referred to as a recombinase, is a polypeptide that catalyzes conservative site-specific recombination between its compatible recombination sites, and includes native polypeptides as well as derivatives, variants and/or fragments that retain activity, and native polynucleotides, derivatives, variants, and/or fragments that encode a recombinase that retains activity.

[0249] One step in the recombination process involves polynucleotide cleavage at or near the recognition site. This cleaving activity can be used to produce a double-strand break. For reviews of site-specific recombinases and their recognition sites, see, Sauer (1994) *Curr Op Biotechnol* 5:521-7; and Sadowski (1993) *FASEB* 7:760-7. In some examples the recombinase is from the Integrase or Resolvase families.

[0250] The Integrase family of recombinases has over one hundred members and includes, for example, FLP, Cre, lambda integrase, and R. The Integrase family has been grouped into two classes based on the structure of the active sites, serine recombinases and tyrosine recombinases. The tyrosine family, which includes Cre, FLP, SSV1, and lambda (2) integrase, uses the catalytic tyrosine's hydroxyl group for a nucleophilic attack on the phosphodiester bond of the DNA. Typically, members of the tyrosine family initially nick the DNA, which later forms a double-strand break. In the serine recombinase family, which includes phiC31 (ΦC31) integrase, a conserved serine residue forms a covalent link to the DNA target site (Grindley et al., (2006) *Ann Rev Biochem* 16:16). For other members of the Integrase family, see for example, Esposito et al., (1997) *Nucleic Acids Res* 25:3605-14 and Abremski et al., (1992) *Protein Eng* 5:87-91.

[0251] Other recombination systems include, for example, the streptomycete bacteriophage phiC31 (Kuhstoss et al., (1991) *J Mol Biol* 20:897-908); the SSV1 site-specific recombination system from *Sulfolobus shibatae* (Maskhlishvili et al., (1993) *Mol Gen Genet* 237:334-42); and a retroviral integrase-based integration system (Tanaka et al., (1998) *Gene* 17:67-76).

[0252] Sometimes the recombinase is one that does not require cofactors or a supercoiled substrate, including but not limited to Cre, FLP, and active derivatives, variants or fragments thereof. FLP recombinase catalyzes a site-specific reaction during DNA replication and amplification of the two-micron plasmid of *S. cerevisiae*. FLP recombinase catalyzes site-specific recombination between two FRT sites. The FLP protein has been cloned and expressed (Cox (1993) *Proc. Natl. Acad. Sci. USA* 80:4223-7). Functional derivatives, variants, and fragments of FLP are known (Buchholz et al., (1998) *Nat Biotechnol* 16:617-8, Hartung et al., (1998) *J Biol Chem* 273:22884-91, Saxena et al., (1997) *Biochim Biophys Acta* 1340:187-204, and Hartley et al., (1980) *Nature* 286:860-4).

[0253] The bacteriophage recombinase Cre catalyzes site-specific recombination between two lox sites (Guo et al., (1997) *Nature* 389:40-6; Abremski et al., (1984) *J Biol Chem* 259:1509-14; Chen et al., (1996) *Somat Cell Mol Genet* 22:477-88; Shaikh et al., (1977) *J Biol Chem* 272:5695-702; and Buchholz et al., (1998) *Nat Biotechnol* 16:617-8). Examples of site-specific recombinases that can be used to produce a double-strand break at a recognition sequence, including for example FLP, Cre, SSV1, lambda Int, phi C31, HK022, and R. Examples of site-specific recombination systems used in plants can be found in U.S. Pat. Nos. 5,929,301; 6,175,056; WO99/25821; U.S. Pat. No. 6,331,661; WO99/25855; WO99/25841, and WO99/25840, the contents of each are herein incorporated by reference.

[0254] Methods for modifying the kinetics, cofactor interaction and requirements, expression, optimal conditions, and/or recognition site specificity, and screening for activity of recombinases and variants are known, see for example Miller et al., (1980) *Cell* 20:721-9; Lange-Gustafson and Nash, (1984) *J Biol Chem* 259:12724-32; Christ et al., (1998) *J Mol Biol* 288:825-36; Lorbach et al., (2000) *J Mol Biol* 296:1175-81; Vergunst et al., (2000) *Science* 290:979-82; Dorgai et al., (1995) *J Mol Biol* 252:178-88; Dorgai et al., (1998) *J Mol Biol* 277:1059-70; Yagu et al., (1995) *J Mol Biol* 252:163-7; Scimiente et al., (2001) *Nucleic Acids Res* 29:5044-51; Santoro and Schultze, (2002) *Proc. Natl. Acad. Sci. USA* 99:4185-90; Buchholz and Stewart, (2001) *Nat Biotechnol* 19:1047-52; Vozizyanov et al., (2002) *Nucleic Acids Res* 30:1656-63; Vozizyanov et al., (2003) *J Mol Biol* 326:65-76; Klippel et al., (1988) *EMBO J* 7:3983-9; Arnold et al., (1999) *EMBO J* 18:1407-14; WO03/08045; WO99/25840; and WO99/25841. The recognition sites range from about 30 nucleotide minimal sites to a few hundred nucleotides.

[0255] Any recognition site for a recombinase can be used, including naturally occurring sites, and variants. Variant recognition sites are known, see for example Hoess et al., (1986) *Nucleic Acids Res* 14:2287-300; Albert et al., (1995) *Plant J* 7:649-59; Thomson et al., (2003) *Genesis* 36:162-7; Huang et al., (1991) *Nucleic Acids Res* 19:443-8; Siebler and Bode, (1997) *Biochemistry* 36:1740-7; Schlake and Bode, (1994) *Biochemistry* 33:12746-51; Thygarajan et al., (2001) *Mol Cell Biol* 21:3926-34; Umlauf and Cox, (1988) *EMBO J* 7:1845-52; Lee and Saito, (1998) *Gene* 216:55-65; WO01/23545; WO99/25821; WO99/25851; WO01/11058; WO01/07572 and U.S. Pat. No. 5,888,732.

[0256] A recombinase can be provided via a polynucleotide that encodes the recombinase or it can be provided via a modified polynucleotide encoding the recombinase. For example, the polynucleotide (encoding a recombinase) can be modified to substitute codons having a higher frequency of usage in a plant, as compared to the naturally occurring polynucleotide sequence or it can be modified to substitute codons having a higher frequency of usage in a maize or soybean plant, as compared to the naturally occurring polynucleotide sequence.

[0257] TAL effector nucleases are a new class of sequence-specific nucleases that can be used to make double-strand breaks at specific target sequences in the genome of a plant or other organism. TAL effector nucleases are created by fusing a native or engineered transcription activator-like (TAL) effector, or functional part thereof, to the catalytic domain of an endonuclease, such as, for example, FokI. The unique, modular TAL effector DNA

binding domain allows for the design of proteins with potentially any given DNA recognition specificity. Thus, the DNA binding domains of the TAL effector nucleases can be engineered to recognize specific DNA target sites and thus, used to make double-strand breaks at desired target sequences. See, WO 2010/079430; Morbitzer et al. (2010) *PNAS* 10.1073/pnas.1013133107; Scholze & Boch (2010) *Virulence* 1:428-432; Christian et al. *Genetics* (2010) 186:757-761; Li et al. (2010) *Nuc. Acids Res.* (2010) doi:10.1093/nar/gkq704; and Miller et al. (2011) *Nature Biotechnology* 29:143-148; all of which are herein incorporated by reference.

[0258] Transposases are polypeptides that mediate transposition of a transposon from one location in the genome to another. Transposases typically induce double-strand breaks to excise the transposon, recognize subterminal repeats, and bring together the ends of the excised transposon, in some systems other proteins are also required to bring together the ends during transposition.

[0259] Examples of transposons and transposases include, but are not limited to, the Ac/Ds, Dt/rdt, Mu-M1/Mn, and Spm (En)/dSpm elements from maize, the Tam elements from snapdragon, the Mu transposon from bacteriophage, bacterial transposons (Tn) and insertion sequences (IS), Ty elements of yeast (retrotransposon), Tal elements from *Arabidopsis* (retrotransposon), the P element transposon from *Drosophila* (Gloor et al., (1991) *Science* 253:1110-1117), the Copia, Mariner and Minos elements from *Drosophila*, the Hermes elements from the housefly, the Piggy-Back elements from *Trichoplusia ni*, Tc1 elements from *C. elegans*, and IAP elements from mice (retrotransposon). In some examples the transposase is provided via a polynucleotide that encodes the transposase.

[0260] It is possible to modify the polynucleotide encoding the transposase by substituting codons having a higher frequency of usage in a plant, as compared to the naturally occurring polynucleotide sequence of by substituting codons having a higher frequency of usage in a maize or soybean plant, as compared to the naturally occurring polynucleotide sequence.

[0261] DNA topoisomerases modulate DNA secondary and higher order structures and functions related primarily to replication, transcription, recombination and repair. Topoisomerases share two characteristics: (i) the ability to cleave and reseal the phosphodiester backbone of DNA in two successive transesterification reactions; and (ii) once a topoisomerase cleaved DNA intermediate is formed, the enzyme allows the severed DNA ends to come apart, allowing the passage of another single- or double-stranded DNA segment. DNA topoisomerases can be classified into three evolutionary independent families: type IA, type IB and type II.

[0262] Those that cleave one strand of DNA and allow single step changes in the linking number of circular DNA are defined as type I DNA topoisomerases. The *Escherichia coli* topoisomerase I and topoisomerase III, *Saccharomyces cerevisiae* topoisomerase III and reverse gyrase belong to the type IA or type I-5' subfamily as the protein link is to a 5' phosphate in the DNA. The prototype of type IB or I-3' enzymes are found in all eukaryotes and also in vaccinia virus topoisomerase I where the protein is attached to a 3' phosphate. Despite differences in mechanism and specificity between the bacterial and eukaryotic enzymes, yeast DNA topoisomerase I can complement a bacterial DNA topoi-

somerase I mutant (Bjornsti et al., (1987) *Proc. Natl. Acad. Sci. USA* 84:8971-5). Type IA topoisomerases relax negatively supercoiled DNA and require magnesium and a single-stranded region of DNA. Topoisomerases IB relax both positively and negatively supercoiled DNA with equal efficiency and do not require a single-stranded region of DNA or metal ions for function.

[0263] The type II family includes *E. coli* DNA gyrase, *E. coli* topoisomerase IV (par E), eukaryotic type II topoisomerases, and archaic topoisomerase VI. Type II enzymes are homodimeric (eukaryotic topoisomerase II) or tetrameric (gyrase), cleaving both strands of a duplex. Preferred cutting sites are known for available topoisomerases.

[0264] Zinc finger nucleases (ZFNs) are engineered double-strand break inducing agents comprised of a zinc finger DNA binding domain and a double-strand-break-inducing agent domain. Recognition site specificity is conferred by the zinc finger domain, which typically comprising two, three, or four zinc fingers, for example having a C2H2 structure, however other zinc finger structures are known and have been engineered. Zinc finger domains are amenable for designing polypeptides which specifically bind a selected polynucleotide recognition sequence. ZFNs consist of an engineered DNA-binding zinc finger domain linked to a non-specific endonuclease domain, for example nuclease domain from a Type IIs endonuclease such as FokI. Additional functionalities can be fused to the zinc-finger binding domain, including transcriptional activator domains, transcription repressor domains, and methylases. In some examples, dimerization of nuclease domain is required for cleavage activity. Each zinc finger recognizes three consecutive base pairs in the target DNA. For example, a 3 finger domain recognized a sequence of 9 contiguous nucleotides, with a dimerization requirement of the nuclease, two sets of zinc finger triplets are used to bind a 18 nucleotide recognition sequence. A recognition sequence of 18 nucleotides is long enough to be unique in a mammalian genome ($4^{18}=6.9 \times 10^{10}$).

[0265] To date, designer zinc finger modules predominantly recognize GNN and ANN triplets (Dreier et al., (2001) *J Biol Chem* 276:29466-78; Dreier et al., (2000) *J Mol Biol* 303:489-502; Liu et al., (2002) *J Biol Chem* 277:3850-6), but examples using CNN or TNN triplets are also known (Dreier et al., (2005) *J Biol Chem* 280:35588-97; Jamieson et al., (2003) *Nature Rev Drug Discov* 2:361-8). See also, Durai et al., (2005) *Nucleic Acids Res* 33:5978-90; Segal, (2002) *Methods* 26:76-83; Porteus and Carroll, (2005) *Nat Biotechnol* 23:967-73; zinc-finger consortium (website at www.zincfinger.org); Pabo et al., (2001) *Ann Rev Biochem* 70:313-40; Wolfe et al., (2000) *Ann Rev Biophys Biomol Struct* 29:183-212; Segal and Barbas, (2001) *Curr Opin Biotechnol* 12:632-7; Segal et al., (2003) *Biochemistry* 42:2137-48; Beerli and Barbas, (2002) *Nat Biotechnol* 20:135-41; Carroll et al., (2006) *Nature Protocols* 1:1329; Ordiz et al., (2002) *Proc. Natl. Acad. Sci. USA* 99:13290-5; Guan et al., (2002) *Proc. Natl. Acad. Sci. USA* 99:13296-301; WO2002099084; WO00/42219; WO02/42459; WO2003062455; U.S. Patent Application Publication No. 20030059767; U.S. Patent Application Publication No. 2003/0108880; U.S. Pat. Nos. 6,140,466, 6,511,808 and 6,453,242.

[0266] Alternatively, engineered zinc finger DNA binding domains can be fused to other double-strand break inducing agents or derivatives thereof that retain DNA nicking/cleav-

ing activity. For example, this type of fusion can be used to direct the double-strand break inducing agent to a different target site, to alter the location of the nick or cleavage site, to direct the inducing agent to a shorter target site, or to direct the inducing agent to a longer target site. In some examples a zinc finger DNA binding domain is fused to a site-specific recombinase, transposase, topoisomerase, or a derivative thereof that retains DNA nicking and/or cleaving activity.

[0267] It is possible to provide a zinc-finger nuclease via a polynucleotide that encodes the zinc-finger nuclease. This polynucleotide encoding the zinc-finger nuclease can be modified by substituting codons having a higher frequency of usage in a plant, as compared to the naturally occurring polynucleotide sequence or by substituting codons having a higher frequency of usage in a maize or soybean plant, as compared to the naturally occurring polynucleotide sequence.

[0268] Sufficient homology or sequence identity indicates that two polynucleotide sequences have sufficient structural similarity to act as substrates for a homologous recombination reaction. The structural similarity includes overall length of each polynucleotide fragment, as well as the sequence similarity of the polynucleotides. Sequence similarity can be described by the percent sequence identity over the whole length of the sequences, and/or by conserved regions comprising localized similarities such as contiguous nucleotides having 100% sequence identity, and percent sequence identity over a portion of the length of the sequences.

[0269] The amount of homology or sequence identity shared by a target and a donor polynucleotide can vary and includes total lengths and/or regions having unit integral values in the ranges of about 1-20 bp, 20-50 bp, 50-100 bp, 75-150 bp, 100-250 bp, 150-300 bp, 200-400 bp, 250-500 bp, 300-600 bp, 350-750 bp, 400-800 bp, 450-900 bp, 500-1000 bp, 600-1250 bp, 700-1500 bp, 800-1750 bp, 900-2000 bp, 1-2.5 kb, 1.5-3 kb, 2-4 kb, 2.5-5 kb, 3-6 kb, 3.5-7 kb, 4-8 kb, 5-10 kb, or up to and including the total length of the target site. These ranges include every integer within the range, for example, the range of 1-20 bp includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 and 20 bp. The amount of homology can also be described by percent sequence identity over the full aligned length of the two polynucleotides which includes percent sequence identity of about at least 50%, 55%, 60%, 65%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100%. Sufficient homology includes any combination of polynucleotide length, global percent sequence identity, and optionally conserved regions of contiguous nucleotides or local percent sequence identity, for example sufficient homology can be described as a region of 75-150 bp having at least 80% sequence identity to a region of the target locus. Sufficient homology can also be described by the predicted ability of two polynucleotides to specifically hybridize under high stringency conditions, see, for example, Sambrook et al., (1989) *Molecular Cloning: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, NY); *Current Protocols in Molecular Biology*, Ausubel et al., Eds (1994) *Current Protocols*, (Greene Publishing Associates, Inc. and John Wiley & Sons, Inc); and, Tijssen (1993) *Laboratory Tech-*

niques in Biochemistry and Molecular Biology—Hybridization with Nucleic Acid Probes, (Elsevier, New York).

[0270] Any means can be used to bring together the various components needed to alter the genome of a dicot plant cell. For example, in in vitro systems, the double-strand-break-inducing agent and the polynucleotide(s) comprising the recognition site(s) can be provided by contacting the components under the appropriate conditions for DNA cleavage.

[0271] Alternatively a variety of methods are known for the introduction of nucleotide sequences and polypeptides into an organism, including, for example, transformation, sexual crossing, and the introduction of the polypeptide, DNA, or mRNA into the cell.

[0272] Methods for contacting, providing, and/or introducing a composition into various organisms are known and include but are not limited to, stable transformation methods, transient transformation methods, virus-mediated methods, and sexual breeding. Stable transformation indicates that the introduced polynucleotide integrates into the genome of the organism and is capable of being inherited by progeny thereof. Transient transformation indicates that the introduced composition is only temporarily expressed or present in the organism.

[0273] Protocols for introducing polynucleotides and polypeptides into plants may vary depending on the type of plant or plant cell targeted for transformation, such as monocot or dicot. Suitable methods of introducing polynucleotides and polypeptides into plant cells and subsequent insertion into the plant genome include microinjection (Crossway et al., (1986) *Biotechniques* 4:320-34 and U.S. Pat. No. 6,300,543), meristem transformation (U.S. Pat. No. 5,736,369), electroporation (Riggs et al., (1986) *Proc. Natl. Acad. Sci. USA* 83:5602-6, *Agrobacterium*-mediated transformation (U.S. Pat. Nos. 5,563,055 and 5,981,840), direct gene transfer (Paszowski et al., (1984) *EMBO J* 3:2717-22), and ballistic particle acceleration (U.S. Pat. Nos. 4,945,050; 5,879,918; 5,886,244; 5,932,782; Tomes et al., (1995) "Direct DNA Transfer into Intact Plant Cells via Microprojectile Bombardment" in *Plant Cell, Tissue, and Organ Culture: Fundamental Methods*, ed. Gamborg & Phillips (Springer-Verlag, Berlin); McCabe et al., (1988) *Biotechnology* 6:923-6; Weissinger et al., (1988) *Ann Rev Genet* 22:421-77; Sanford et al., (1987) *Particulate Science and Technology* 5:27-37 (onion); Christou et al., (1988) *Plant Physiol* 87:671-4 (soybean); Finer and McMullen, (1991) *In Vitro Cell Dev Biol* 27P: 175-82 (soybean); Singh et al., (1998) *Theor Appl Genet* 96:319-24 (soybean); Datta et al., (1990) *Biotechnology* 8:736-40 (rice); Klein et al., (1988) *Proc. Natl. Acad. Sci. USA* 85:4305-9 (maize); Klein et al., (1988) *Biotechnology* 6:559-63 (maize); U.S. Pat. Nos. 5,240,855; 5,322,783 and 5,324,646; Klein et al., (1988) *Plant Physiol* 91:440-4 (maize); Fromm et al., (1990) *Biotechnology* 8:833-9 (maize); Hooykaas-Van Slogteren et al., (1984) *Nature* 311:763-4; U.S. Pat. No. 5,736,369 (cereals); Bytebier et al., (1987) *Proc. Natl. Acad. Sci. USA* 84:5345-9 (Liliaceae); De Wet et al., (1985) in *The Experimental Manipulation of Ovule Tissues*, ed. Chapman et al., (Longman, New York), pp. 197-209 (pollen); Kaeppler et al., (1990) *Plant Cell Rep* 9:415-8; and Kaeppler et al., (1992) *Theor Appl Genet* 84:560-6 (whisker-mediated transformation); D'Halluin et al., (1992) *Plant Cell* 4:1495-505 (electroporation); Li et al., (1993) *Plant Cell Rep* 12:250-5; Christou and Ford (1995) *Annals Botany* 75:407-13 (rice)

and Osjoda et al., (1996) *Nat Biotechnol* 14:745-50 (maize via *Agrobacterium tumefaciens*).

[0274] Alternatively, polynucleotides may be introduced into plants by contacting plants with a virus or viral nucleic acids. Generally, such methods involve incorporating a polynucleotide within a viral DNA or RNA molecule. In some examples a polypeptide of interest may be initially synthesized as part of a viral polyprotein, which is later processed by proteolysis in vivo or in vitro to produce the desired recombinant protein. Methods for introducing polynucleotides into plants and expressing a protein encoded therein, involving viral DNA or RNA molecules, are known, see, for example, U.S. Pat. Nos. 5,889,191, 5,889,190, 5,866,785, 5,589,367 and 5,316,931. Transient transformation methods include, but are not limited to, the introduction of polypeptides, such as a double-strand break inducing agent, directly into the organism, the introduction of polynucleotides such as DNA and/or RNA polynucleotides, and the introduction of the RNA transcript, such as an mRNA encoding a double-strand break inducing agent, into the organism. Such methods include, for example, microinjection or particle bombardment. See, for example Crossway et al., (1986) *Mol Gen Genet* 202:179-85; Nomura et al., (1986) *Plant Sci* 44:53-8; Hepler et al., (1994) *Proc. Natl. Acad. Sci. USA* 91:2176-80; and, Hush et al., (1994) *J Cell Sci* 107:775-84.

[0275] Standard DNA isolation, purification, molecular cloning, vector construction, and verification/characterization methods are well established, see, for example Sambrook et al., (1989) *Molecular Cloning: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, NY). Vectors and constructs include circular plasmids, and linear polynucleotides, comprising a polynucleotide of interest and optionally other components including linkers, adapters, regulatory regions, introns, restriction sites, enhancers, insulators, selectable markers, nucleotide sequences of interest, promoters, and/or other sites that aid in vector construction or analysis. In some examples a recognition site and/or target site can be contained within an intron, coding sequence, 5' UTRs, 3' UTRs, and/or regulatory regions.

[0276] The present invention further provides expression constructs for expressing in a plant, plant cell, or plant part an endonuclease that is capable of binding to and creating a double strand break in a target site. The expression constructs of the invention comprise a promoter operably linked to a nucleotide sequence encoding an endonuclease of the present invention. The promoter is capable of driving expression of an operably linked nucleotide sequence in a plant cell. Any such promoter that is disclosed herein or known in the art can be used in the present invention. In one embodiment, the target site of the endonuclease is selected from the group consisting of TS21, TS14, TS30, TS5, TS7, TS4, TS22, and TS24 target sites of soybean, which have the nucleotide sequences set forth in SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, and 8, respectively. In another embodiment, the target site of the endonuclease is selected from the group consisting of MHP1, MHP14, MHP32, MHP42, MHP55, MHP67, MHP77, MHP98, MHP107, and MHP115 target sites of maize, which have the nucleotide sequences set forth in SEQ ID NO:68, 69, 70, 71, 72, 73, 74, 75, 76, and 77, respectively.

[0277] In certain embodiments, the expression constructs comprise a nucleotide sequence encoding the endonuclease that has been custom designed or engineered to cut at one the

soybean target sites set forth above. Such nucleotide sequences include, for example, the nucleotide sequences set forth in SEQ ID NOS: 9, 10, 11, 12, 13, 14, 15, and 16. Other nucleotide sequences of the invention include, but are not limited to, nucleotide sequences comprising a coding sequence of a DNA binding domain of an endonuclease, wherein the coding sequence is nucleotides 100-261 and nucleotides 850-1011 of SEQ ID NO:9, 10, 11, 12, 13, 14, 15 or 16 and degenerate coding sequences thereof. Such a degenerate coding sequence encodes the same amino acid sequence as that encoded by one of the coding sequences set forth in nucleotides 100-261 and nucleotides 850-1011 of SEQ ID NO:9, 10, 11, 12, 13, 14, 15 or 16 but differs in its nucleotide sequence due to the degeneracy of the genetic code.

[0278] In certain other embodiments, the expression constructs comprise a nucleotide sequence encoding the endonuclease that has been custom designed or engineered to cut at one the maize target sites set forth above. Such nucleotide sequences include, for example, the nucleotide sequences set forth in SEQ ID NOS: 78, 79, 80, 81, 82, and 83. Other nucleotide sequences of the invention include, but are not limited to, nucleotide sequences comprising a coding sequence of a DNA binding domain of an endonuclease, wherein the coding sequence comprises nucleotides 100-261 and nucleotides 850-1011 of SEQ ID NO: 80 and degenerate coding sequences thereof. Such a degenerate coding sequence encodes the same amino acid sequence as that encoded by one of the coding sequences set forth in nucleotides 100-261 and nucleotides 850-1011 of SEQ ID NO: 80 but differs in its nucleotide sequence due to the degeneracy of the genetic code. Other nucleotide sequences of the invention include, but are not limited to, nucleotide sequences comprising a coding sequence of a DNA binding domain of an endonuclease, wherein the coding sequence is nucleotides 100-261 and nucleotides 661-822 of SEQ ID NO: 78, 79, 81, 82 or 83 and degenerate coding sequences thereof. Such a degenerate coding sequence encodes the same amino acid sequence as that encoded by one of the coding sequences set forth in nucleotides 100-261 and nucleotides 661-822 of SEQ ID NO: 78, 79, 81, 82 or 83 but differs in its nucleotide sequence due to the degeneracy of the genetic code.

[0279] Any promoter can be used, and can be selected based on the desired outcome. A promoter is a region of DNA involved in recognition and binding of RNA polymerase and other proteins to initiate transcription. A plant promoter is a promoter capable of initiating transcription in a plant cell, for a review of plant promoters, see, Potenza et al., (2004) *In Vitro Cell Dev Biol* 40:1-22. Constitutive promoters include, for example, the core promoter of the Rsyn7 promoter and other constitutive promoters disclosed in WO99/43838 and U.S. Pat. No. 6,072,050; the core CaMV 35S promoter (Odell et al., (1985) *Nature* 313:810-2); rice actin (McElroy et al., (1990) *Plant Cell* 2:163-71); ubiquitin (Christensen et al., (1989) *Plant Mol Biol* 12:619-32; Christensen et al., (1992) *Plant Mol Biol* 18:675-89); pEMU (Last et al., (1991) *Theor Appl Genet* 81:581-8); MAS (Velten et al., (1984) *EMBO J* 3:2723-30); ALS promoter (U.S. Pat. No. 5,659,026), and the like. Other constitutive promoters are described in, for example, U.S. Pat. Nos. 5,608,149; 5,608,144; 5,604,121; 5,569,597; 5,466,785; 5,399,680; 5,268,463; 5,608,142 and 6,177,611. In some examples an inducible promoter may be used.

Pathogen-inducible promoters induced following infection by a pathogen include, but are not limited to those regulating expression of PR proteins, SAR proteins, beta-1,3-glucanase, chitinase, etc.

[0280] Chemical-regulated promoters can be used to modulate the expression of a gene in a plant through the application of an exogenous chemical regulator. The promoter may be a chemical-inducible promoter, where application of the chemical induces gene expression, or a chemical-repressible promoter, where application of the chemical represses gene expression. Chemical-inducible promoters include, but are not limited to, the maize In2-2 promoter, activated by benzene sulfonamide herbicide safeners (De Veylder et al., (1997) *Plant Cell Physiol* 38:568-77), the maize GST promoter (GST-II-27, WO93/01294), activated by hydrophobic electrophilic compounds used as pre-emergent herbicides, and the tobacco PR-1a promoter (Ono et al., (2004) *Biosci Biotechnol Biochem* 68:803-7) activated by salicylic acid. Other chemical-regulated promoters include steroid-responsive promoters (see, for example, the glucocorticoid-inducible promoter (Scheda et al., (1991) *Proc. Natl. Acad. Sci. USA* 88:10421-5; McNellis et al., (1998) *Plant J* 14:247-257); tetracycline-inducible and tetracycline-repressible promoters (Gatz et al., (1991) *Mol Gen Genet* 227:229-37; U.S. Pat. Nos. 5,814,618 and 5,789,156).

[0281] Tissue-preferred promoters can be utilized to target enhanced expression within a particular plant tissue. Tissue-preferred promoters include, for example, Kawamata et al., (1997) *Plant Cell Physiol* 38:792-803; Hansen et al., (1997) *Mol Gen Genet* 254:337-43; Russell et al., (1997) *Transgenic Res* 6:157-68; Rinehart et al., (1996) *Plant Physiol* 112:1331-41; Van Camp et al., (1996) *Plant Physiol* 112:525-35; Canevascini et al., (1996) *Plant Physiol* 112:513-524; Lam, (1994) *Results Probl Cell Differ* 20:181-96; and Guevara-Garcia et al., (1993) *Plant J* 4:495-505. Leaf-preferred promoters include, for example, Yamamoto et al., (1997) *Plant J* 12:255-65; Kwon et al., (1994) *Plant Physiol* 105:357-67; Yamamoto et al., (1994) *Plant Cell Physiol* 35:773-8; Gotor et al., (1993) *Plant J* 3:509-18; Orozco et al., (1993) *Plant Mol Biol* 23:1129-38; Matsuoka et al., (1993) *Proc. Natl. Acad. Sci. USA* 90:9586-90; Simpson et al., (1958) *EMBO J* 4:2723-9; Timko et al., (1988) *Nature* 318:57-8. Root-preferred promoters include, for example, Hire et al., (1992) *Plant Mol Biol* 20:207-18 (soybean root-specific glutamine synthase gene); Miao et al., (1991) *Plant Cell* 3:11-22 (cytosolic glutamine synthase (GS)); Keller and Baumgartner, (1991) *Plant Cell* 3:1051-61 (root-specific control element in the GRP 1.8 gene of French bean); Sanger et al., (1990) *Plant Mol Biol* 14:433-43 (root-specific promoter of *A. tumefaciens* mannopine synthase (MAS)); Bogusz et al., (1990) *Plant Cell* 2:633-41 (root-specific promoters isolated from *Parasponia andersonii* and *Trema tomentosa*); Leach and Aoyagi, (1991) *Plant Sci* 79:69-76 (*A. rhizogenes* rolC and rolD root-inducing genes); Teeri et al., (1989) *EMBO J* 8:343-50 (*Agrobacterium* wound-induced TR1' and TR2' genes); VtENOD-GRP3 gene promoter (Kuster et al., (1995) *Plant Mol Biol* 29:759-72); and rolB promoter (Capana et al., (1994) *Plant Mol Biol* 25:681-91; phaseolin gene (Murai et al., (1983) *Science* 23:476-82; Sengopta-Gopalan et al., (1988) *Proc. Natl. Acad. Sci. USA* 82:3320-4). See also, U.S. Pat. Nos. 5,837,876; 5,750,386; 5,633,363; 5,459,252; 5,401,836; 5,110,732 and 5,023,179.

[0282] Seed-preferred promoters include both seed-specific promoters active during seed development, as well as seed-germinating promoters active during seed germination. See, Thompson et al., (1989) *BioEssays* 10:108. Seed-preferred promoters include, but are not limited to, Cim1 (cytokinin-induced message); cZ19B1 (maize 19 kDa zein); and milps (myo-inositol-1-phosphate synthase); (WO00/11177; and U.S. Pat. No. 6,225,529). For dicots, seed-preferred promoters include, but are not limited to, bean β -phaseolin, napin, β -conglycinin, soybean lectin, cruciferin, and the like. For monocots, seed-preferred promoters include, but are not limited to, maize 15 kDa zein, 22 kDa zein, 27 kDa gamma zein, waxy, shrunken 1, shrunken 2, globulin 1, oleosin, and nuc1. See also, WO00/12733, where seed-preferred promoters from END1 and END2 genes are disclosed.

[0283] A phenotypic marker is screenable or selectable marker that includes visual markers and selectable markers whether it is a positive or negative selectable marker. Any phenotypic marker can be used. Specifically, a selectable or screenable marker comprises a DNA segment that allows one to identify, or select for or against a molecule or a cell that contains it, often under particular conditions. These markers can encode an activity, such as, but not limited to, production of RNA, peptide, or protein, or can provide a binding site for RNA, peptides, proteins, inorganic and organic compounds or compositions and the like.

[0284] Examples of selectable markers include, but are not limited to, DNA segments that comprise restriction enzyme sites; DNA segments that encode products which provide resistance against otherwise toxic compounds including antibiotics, such as, spectinomycin, ampicillin, kanamycin, tetracycline, Basta, neomycin phosphotransferase II (NEO) and hygromycin phosphotransferase (HPT); DNA segments that encode products which are otherwise lacking in the recipient cell (e.g., tRNA genes, auxotrophic markers); DNA segments that encode products which can be readily identified (e.g., phenotypic markers such as B-galactosidase, GUS; fluorescent proteins such as green fluorescent protein (GFP), cyan (CFP), yellow (YFP), red (RFP), and cell surface proteins); the generation of new primer sites for PCR (e.g., the juxtaposition of two DNA sequence not previously juxtaposed), the inclusion of DNA sequences not acted upon or acted upon by a restriction endonuclease or other DNA modifying enzyme, chemical, etc.; and, the inclusion of a DNA sequences required for a specific modification (e.g., methylation) that allows its identification.

[0285] Additional selectable markers include genes that confer resistance to herbicidal compounds, such as glufosinate ammonium, bromoxynil, imidazolinones, and 2,4-dichlorophenoxyacetate (2,4-D). See for example, Yarranton, (1992) *Curr Opin Biotech* 3:506-11; Christopherson et al., (1992) *Proc. Natl. Acad. Sci. USA* 89:6314-8; Yao et al., (1992) *Cell* 71:63-72; Reznikoff, (1992) *Mol Microbiol* 6:2419-22; Hu et al., (1987) *Cell* 48:555-66; Brown et al., (1987) *Cell* 49:603-12; Figge et al., (1988) *Cell* 52:713-22; Deuschle et al., (1989) *Proc. Natl. Acad. Sci. USA* 86:5400-4; Fuerst et al., (1989) *Proc. Natl. Acad. Sci. USA* 86:2549-53; Deuschle et al., (1990) *Science* 248:480-3; Gossen, (1993) Ph.D. Thesis, University of Heidelberg; Reines et al., (1993) *Proc. Natl. Acad. Sci. USA* 90:1917-21; Labow et al., (1990) *Mol Cell Biol* 10:3343-56; Zambretti et al., (1992) *Proc. Natl. Acad. Sci. USA* 89:3952-6; Baim et al., (1991) *Proc. Natl. Acad. Sci. USA* 88:5072-6; Wyborski et al.,

(1991) *Nucleic Acids Res* 19:4647-53; Hillen and Wissman, (1989) *Topics Mol Struc Biol* 10:143-62; Degenkolb et al., (1991) *Antimicrob Agents Chemother* 35:1591-5; Kleinschmidt et al., (1988) *Biochemistry* 27:1094-104; Bonin, (1993) Ph.D. Thesis, University of Heidelberg; Gossen et al., (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-51; Oliva et al., (1992) *Antimicrob Agents Chemother* 36:913-9; Hlavka et al., (1985) *Handbook of Experimental Pharmacology*, Vol. 78 (Springer-Verlag, Berlin); Gill et al., (1988) *Nature* 334:721-4.

[0286] The cells having the introduced sequence may be grown or regenerated into plants using conventional conditions, see for example, McCormick et al., (1986) *Plant Cell Rep* 5:81-4. These plants may then be grown, and either pollinated with the same transformed strain or with a different transformed or untransformed strain, and the resulting progeny having the desired characteristic and/or comprising the introduced polynucleotide or polypeptide identified. Two or more generations may be grown to ensure that the polynucleotide is stably maintained and inherited, and seeds harvested.

[0287] Any plant can be used, including monocot and dicot plants. Examples of monocot plants that can be used include, but are not limited to, corn (*Zea mays*), rice (*Oryza sativa*), rye (*Secale cereale*), sorghum (*Sorghum bicolor*, *Sorghum vulgare*), millet (e.g., pearl millet (*Pennisetum glaucum*), proso millet (*Panicum miliaceum*), foxtail millet (*Setaria italica*), finger millet (*Eleusine coracana*)), wheat (*Triticum aestivum*), sugarcane (*Saccharum* spp.), oats (*Avena*), barley (*Hordeum*), switchgrass (*Panicum virgatum*), pineapple (*Ananas comosus*), banana (*Musa* spp.), palm, ornamentals, turfgrasses, and other grasses. Examples of dicot plants that can be used include, but are not limited to, soybean (*Glycine max*), canola (*Brassica napus* and *B. campestris*), alfalfa (*Medicago sativa*), tobacco (*Nicotiana tabacum*), *Arabidopsis* (*Arabidopsis thaliana*), sunflower (*Helianthus annuus*), cotton (*Gossypium arboreum*), and peanut (*Arachis hypogaea*), tomato (*Solanum lycopersicum*), potato (*Solanum tuberosum*) etc.

[0288] The transgenes, recombinant DNA molecules, DNA sequences of interest, and polynucleotides of interest can comprise one or more genes of interest. Such genes of interest can encode, for example, a protein that provides agronomic advantage to the plant. Genes of interest, including, but not limited to, those that encode proteins that provide agronomic advantage, can be reflective of the commercial markets and interests of those involved in the development of the crop. Crops and markets of interest change, and as developing nations open up world markets, new crops and technologies will emerge also. In addition, as our understanding of agronomic traits and characteristics such as yield and heterosis increase, the choice of genes for transformation will change accordingly. General categories of genes of interest include, for example, those genes involved in information, such as zinc fingers, those involved in communication, such as kinases, and those involved in housekeeping, such as heat shock proteins. More specific categories of transgenes, for example, include genes encoding important traits for agronomics, insect resistance, disease resistance, herbicide resistance, sterility, grain characteristics, and commercial products. Genes of interest include, generally, those involved in oil, starch, carbohydrate, or nutrient metabolism as well as those affecting kernel size, sucrose loading, and the like.

[0289] Agronomically important traits such as oil, starch, and protein content can be genetically altered in addition to using traditional breeding methods. Modifications include increasing content of oleic acid, saturated and unsaturated oils, increasing levels of lysine and sulfur, providing essential amino acids, and also modification of starch. Hordothionin protein modifications are described in U.S. Pat. Nos. 5,703,049, 5,885,801, 5,885,802, and 5,990,389, herein incorporated by reference. Another example is lysine and/or sulfur rich seed protein encoded by the soybean 2S albumin described in U.S. Pat. No. 5,850,016, and the chymotrypsin inhibitor from barley, described in Williamson et al. (1987) *Eur. J. Biochem.* 165:99-106, the disclosures of which are herein incorporated by reference.

[0290] Derivatives of the coding sequences can be made by site-directed mutagenesis to increase the level of preselected amino acids in the encoded polypeptide. For example, the gene encoding the barley high lysine polypeptide (BHL) is derived from barley chymotrypsin inhibitor, U.S. application Ser. No. 08/740,682, filed Nov. 1, 1996, and WO 98/20133, the disclosures of which are herein incorporated by reference. Other proteins include methionine-rich plant proteins such as from sunflower seed (Lilley et al. (1989) *Proceedings of the World Congress on Vegetable Protein Utilization in Human Foods and Animal Feedstuffs*, ed. Applewhite (American Oil Chemists Society, Champaign, Illinois), pp. 497-502; herein incorporated by reference); corn (Pedersen et al. (1986) *J. Biol. Chem.* 261:6279; Kiriara et al. (1988) *Gene* 71:359; both of which are herein incorporated by reference); and rice (Musumura et al. (1989) *Plant Mol. Biol.* 12:123, herein incorporated by reference). Other agronomically important genes encode latex, Floury 2, growth factors, seed storage factors, and transcription factors.

[0291] Insect resistance genes may encode resistance to pests that have great yield drag such as rootworm, cutworm, European Corn Borer, and the like. Such genes include, for example, *Bacillus thuringiensis* toxic protein genes (U.S. Pat. Nos. 5,366,892; 5,747,450; 5,736,514; 5,723,756; 5,593,881; and Geiser et al. (1986) *Gene* 48:109); and the like.

[0292] Genes encoding disease resistance traits include detoxification genes, such as against fumonisin (U.S. Pat. No. 5,792,931); avirulence (avr) and disease resistance (R) genes (Jones et al. (1994) *Science* 266:789; Martin et al. (1993) *Science* 262:1432; and Mindrinos et al. (1994) *Cell* 78:1089); and the like.

[0293] Herbicide resistance traits may include genes coding for resistance to herbicides that act to inhibit the action of acetolactate synthase (ALS), in particular the sulfonyleurea-type herbicides (e.g., the acetolactate synthase (ALS) gene containing mutations leading to such resistance, in particular the S4 and/or Hra mutations), genes coding for resistance to herbicides that act to inhibit action of glutamine synthase, such as phosphinothricin or basta (e.g., the bar gene); glyphosate (e.g., the EPSPS gene and the GAT gene; see, for example, U.S. Publication No. 20040082770 and WO 03/092360); or other such genes known in the art. The bar gene encodes resistance to the herbicide basta, the nptII gene encodes resistance to the antibiotics kanamycin and geneticin, and the ALS-gene mutants encode resistance to the herbicide chlorsulfuron.

[0294] Sterility genes can also be encoded in an expression cassette and provide an alternative to physical detas-

seling. Examples of genes used in such ways include male tissue-preferred genes and genes with male sterility phenotypes such as QM, described in U.S. Pat. No. 5,583,210. Other genes include kinases and those encoding compounds toxic to either male or female gametophytic development.

[0295] The quality of grain is reflected in traits such as levels and types of oils, saturated and unsaturated, quality and quantity of essential amino acids, and levels of cellulose. In corn, modified hordothionin proteins are described in U.S. Pat. Nos. 5,703,049, 5,885,801, 5,885,802, and 5,990,389.

[0296] Commercial traits can also be encoded on a gene or genes that could increase for example, starch for ethanol production, or provide expression of proteins. Another important commercial use of transformed plants is the production of polymers and bioplastics such as described in U.S. Pat. No. 5,602,321. Genes such as β -Ketothiolase, PHBase (polyhydroxybutyrate synthase), and acetoacetyl-CoA reductase (see Schubert et al. (1988) *J. Bacteriol.* 170:5837-5847) facilitate expression of polyhydroxyalkanoates (PHAs).

[0297] Exogenous products include plant enzymes and products as well as those from other sources including procaryotes and other eukaryotes. Such products include enzymes, cofactors, hormones, and the like. The level of proteins, particularly modified proteins having improved amino acid distribution to improve the nutrient value of the plant, can be increased. This is achieved by the expression of such proteins having enhanced amino acid content.

[0298] The transgenes, recombinant DNA molecules, DNA sequences of interest, and polynucleotides of interest can be comprise one or more DNA sequences for gene silencing. Methods for gene silencing involving the expression of DNA sequences in plant are known in the art include, but are not limited to, cosuppression, antisense suppression, double-stranded RNA (dsRNA) interference, hairpin RNA (hpRNA) interference, intron-containing hairpin RNA (ihpRNA) interference, transcriptional gene silencing, and micro RNA (miRNA) interference

[0299] Cosuppression may be used to inhibit the expression of plant genes to produce plants having undetectable protein levels for the proteins encoded by these genes. See, for example, Broin et al. (2002) *Plant Cell* 14:1417-1432. Cosuppression may also be used to inhibit the expression of multiple proteins in the same plant. See, for example, U.S. Pat. No. 5,942,657. Methods for using cosuppression to inhibit the expression of endogenous genes in plants are described in Flavell et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:3490-3496; Jorgensen et al. (1996) *Plant Mol. Biol.* 31:957-973; Johansen and Carrington (2001) *Plant Physiol.* 126:930-938; Broin et al. (2002) *Plant Cell* 14:1417-1432; Stoutjesdijk et al (2002) *Plant Physiol.* 129:1723-1731; Yu et al. (2003) *Phytochemistry* 63:753-763; and U.S. Pat. Nos. 5,034,323, 5,283,184, and 5,942,657; each of which is herein incorporated by reference. The efficiency of cosuppression may be increased by including a poly-dT region in the expression cassette at a position 3' to the sense sequence and 5' of the polyadenylation signal. See, U.S. Patent Publication No. 20020048814, herein incorporated by reference. Typically, such a nucleotide sequence has substantial sequence identity to the sequence of the transcript of the endogenous gene, optimally greater than about 65% sequence identity, more optimally greater than about 85% sequence identity, most optimally greater than about 95%

sequence identity. See, U.S. Pat. Nos. 5,283,184 and 5,034,323; herein incorporated by reference.

[0300] Antisense suppression may be used to inhibit the expression of multiple proteins in the same plant. See, for example, U.S. Pat. No. 5,942,657. Furthermore, portions of the antisense nucleotides may be used to disrupt the expression of the target gene. Generally, sequences of at least 50 nucleotides, 100 nucleotides, 200 nucleotides, 300, 400, 450, 500, 550, or greater may be used. Methods for using antisense suppression to inhibit the expression of endogenous genes in plants are described, for example, in Liu et al (2002) *Plant Physiol.* 129:1732-1743 and U.S. Pat. Nos. 5,759,829 and 5,942,657, each of which is herein incorporated by reference. Efficiency of antisense suppression may be increased by including a poly-dT region in the expression cassette at a position 3' to the antisense sequence and 5' of the polyadenylation signal. See, U.S. Patent Publication No. 20020048814, herein incorporated by reference.

[0301] Methods for using dsRNA interference to inhibit the expression of endogenous plant genes are described in Waterhouse et al. (1998) *Proc. Natl. Acad. Sci. USA* 95:13959-13964, Liu et al. (2002) *Plant Physiol.* 129:1732-1743, and WO 99/49029, WO 99/53050, WO 99/61631, and WO 00/49035; each of which is herein incorporated by reference.

[0302] Methods of hpRNA interference are described in Waterhouse and Helliwell (2003) *Nat. Rev. Genet.* 4:29-38 and the references cited therein. These methods are highly efficient at inhibiting the expression of endogenous genes. See, for example, Chuang and Meyerowitz (2000) *Proc. Natl. Acad. Sci. USA* 97:4985-4990; Stoutjesdijk et al. (2002) *Plant Physiol.* 129:1723-1731; and Waterhouse and Helliwell (2003) *Nat. Rev. Genet.* 4:29-38. Methods for using hpRNA interference to inhibit or silence the expression of genes are described, for example, in Chuang and Meyerowitz (2000) *Proc. Natl. Acad. Sci. USA* 97:4985-4990; Stoutjesdijk et al. (2002) *Plant Physiol.* 129:1723-1731; Waterhouse and Helliwell (2003) *Nat. Rev. Genet.* 4:29-38; Pandolfini et al. *BMC Biotechnology* 3:7, and U.S. Patent Publication No. 20030175965; each of which is herein incorporated by reference. A transient assay for the efficiency of hpRNA constructs to silence gene expression in vivo has been described by Panstruga et al. (2003) *Mol. Biol. Rep.* 30:135-140, herein incorporated by reference.

[0303] For ihpRNA, the interfering molecules have the same general structure as for hpRNA, but the RNA molecule additionally comprises an intron that is capable of being spliced in the cell in which the ihpRNA is expressed. The use of an intron minimizes the size of the loop in the hairpin RNA molecule following splicing, and this increases the efficiency of interference. See, for example, Smith et al. (2000) *Nature* 407:319-320. In fact, Smith et al. show 100% suppression of endogenous gene expression using ihpRNA-mediated interference. Methods for using ihpRNA interference to inhibit the expression of endogenous plant genes are described, for example, in Smith et al. (2000) *Nature* 407:319-320; Wesley et al. (2001) *Plant J.* 27:581-590; Wang and Waterhouse (2001) *Curr. Opin. Plant Biol.* 5:146-150; Waterhouse and Helliwell (2003) *Nat. Rev. Genet.* 4:29-38; Helliwell and Waterhouse (2003) *Methods* 30:289-295, and U.S. Patent Publication No. 20030180945, each of which is herein incorporated by reference.

[0304] Transcriptional gene silencing (TGS) may be accomplished through use of hpRNA constructs wherein the

inverted repeat of the hairpin shares sequence identity with the promoter region of a gene to be silenced. Processing of the hpRNA into short RNAs which can interact with the homologous promoter region may trigger degradation or methylation to result in silencing (Aufsatz et al. (2002) *PNAS* 99 (Suppl. 4): 16499-16506; Mette et al. (2000) *EMBO J* 19 (19): 5194-5201).

[0305] The inhibition of the expression of a target protein may be obtained by RNA interference by expression of a gene encoding a micro RNA (miRNA). miRNAs are regulatory agents consisting of about 22 ribonucleotides. miRNA are highly efficient at inhibiting the expression of endogenous genes. See, for example Javier et al. (2003) *Nature* 425:257-263, herein incorporated by reference. For miRNA interference, the expression cassette is designed to express an RNA molecule that is modeled on an endogenous miRNA gene. The miRNA gene encodes an RNA that forms a hairpin structure containing a 22-nucleotide sequence that is complementary to another endogenous gene (target sequence). miRNA molecules are highly efficient at inhibiting the expression of endogenous genes, and the RNA interference they induce is inherited by subsequent generations of plants.

[0306] The frequency of homologous recombination is influenced by a number of factors. Different organisms vary with respect to the amount of homologous recombination and the relative proportion of homologous to non-homologous recombination. Generally, the length of the region of homology affects the frequency of homologous recombination events, the longer the region of homology, the greater the frequency. The length of the homology region needed to observe homologous recombination is also species-variable. In many cases, at least 5 kb of homology has been utilized, but homologous recombination has been observed with as little as 25-50 bp of homology. The minimum length of homology needed has been estimated at 20-50 bp in *E. coli* (Singer et al., (1982) *Cell* 31:25-33; Shen and Huang, (1986) *Genetics* 112:441-57; Watt et al., (1985) *Proc. Natl. Acad. Sci. USA* 82:4768-72), 63-89 bp in *Saccharomyces cerevisiae* (Sugawara and Haber, (1992) *Mol Cell Biol* 12:563-75), and 163-300 bp in mammalian cells (Rubnitz and Subramani, (1984) *Mol Cell Biol* 4:2253-8; Ayares et al., (1986) *Proc. Natl. Acad. Sci. USA* 83:5199-203; Liskay et al., (1987) *Genetics* 115:161-7).

[0307] Homologous recombination has been demonstrated in insects. In *Drosophila*, Dray and Gloor found that as little as 3 kb of total template: target homology sufficed to copy a large non-homologous segment of DNA into the target with reasonable efficiency (Dray and Gloor, (1997) *Genetics* 147:689-99). Using FLP-mediated DNA integration at a target FRT in *Drosophila*, Golic et al., showed integration was approximately 10-fold more efficient when the donor and target shared 4.1 kb of homology as compared to 1.1 kb of homology (Golic et al., (1997) *Nucleic Acids Res* 25:3665). Data from *Drosophila* indicates that 2-4 kb of homology is sufficient for efficient targeting, but there is some evidence that much less homology may suffice, on the order of about 30 bp to about 100 bp (Nassif and Engels, (1993) *Proc. Natl. Acad. Sci. USA* 90:1262-6; Keeler and Gloor, (1997) *Mol Cell Biol* 17:627-34).

[0308] Homologous recombination has also been accomplished in other organisms. For example, at least 150-200 bp of homology was required for homologous recombination in the parasitic protozoan *Leishmania* (Papadopoulou and

Dumas, (1997) *Nucleic Acids Res* 25:4278-86). In the filamentous fungus *Aspergillus nidulans*, gene replacement has been accomplished with as little as 50 bp flanking homology (Chaveroche et al., (2000) *Nucleic Acids Res* 28: e97). Targeted gene replacement has also been demonstrated in the ciliate *Tetrahymena thermophila* (Gaertig et al., (1994) *Nucleic Acids Res* 22:5391-8). In mammals, homologous recombination has been most successful in the mouse using pluripotent embryonic stem cell lines (ES) that can be grown in culture, transformed, selected and introduced into a mouse embryo. Embryos bearing inserted transgenic ES cells develop as genetically chimeric offspring. By interbreeding siblings, homozygous mice carrying the selected genes can be obtained. An overview of the process is provided in Watson et al., (1992) *Recombinant DNA*, 2nd Ed., (Scientific American Books distributed by WH Freeman & Co.); Capecchi, (1989) *Trends Genet* 5:70-6; and Bronson, (1994) *J Biol Chem* 269:27155-8. Homologous recombination in mammals other than mouse has been limited by the lack of stem cells capable of being transplanted to oocytes or developing embryos. However, McCreath et al., *Nature* 405:1066-9 (2000) reported successful homologous recombination in sheep by transformation and selection in primary embryo fibroblast cells.

[0309] Error-prone DNA repair mechanisms can produce mutations at double-strand break sites. The nonhomologous end-joining (NHEJ) pathways are the most common repair mechanism to bring the broken ends together (Bleuyard et al., (2006) *DNA Repair* 5:1-12). The structural integrity of chromosomes is typically preserved by the repair, but deletions, insertions, or other rearrangements are possible. The two ends of one double-strand break are the most prevalent substrates of NHEJ (Kirik et al., (2000) *EMBO J* 19:5562-6), however if two different double-strand breaks occur, the free ends from different breaks can be ligated and result in chromosomal deletions (Siebert and Puchta, (2002) *Plant Cell* 14:1121-31), or chromosomal translocations between different chromosomes (Pacher et al., (2007) *Genetics* 175: 21-9).

[0310] Episomal DNA molecules can also be ligated into the double-strand break, for example, integration of T-DNAs into chromosomal double-strand breaks (Chilton and Que, (2003) *Plant Physiol* 133:956-65; Salomon and Puchta, (1998) *EMBO J* 17:6086-95). Once the sequence around the double-strand breaks is altered, for example, by exonuclease activities involved in the maturation of double-strand breaks, gene conversion pathways can restore the original structure if a homologous sequence is available, such as a homologous chromosome in non-dividing somatic cells, or a sister chromatid after DNA replication (Molinier et al., (2004) *Plant Cell* 16:342-52). Ectopic and/or epigenic DNA sequences may also serve as a DNA repair template for homologous recombination (Puchta, (1999) *Genetics* 152: 1173-81).

[0311] Alteration of the genome of a plant cell, for example, through homologous recombination (HR), is a powerful tool for genetic engineering. Despite the low frequency of homologous recombination in higher plants, there are a few examples of successful homologous recombination of plant endogenous genes. The parameters for homologous recombination in plants have primarily been investigated by rescuing introduced truncated selectable marker genes. In these experiments, the homologous DNA fragments were typically between 0.3 kb to 2 kb. Observed

frequencies for homologous recombination were on the order of 10^{-4} to 10^{-5} . See, for example, Halfter et al., (1992) *Mol Gen Genet* 231:186-93; Offringa et al., (1990) *EMBO J* 9:3077-84; Offringa et al., (1993) *Proc. Natl. Acad. Sci. USA* 90:7346-50; Paszkowski et al., (1988) *EMBO J* 7:4021-6; Hourda and Paszkowski, (1994) *Mol Gen Genet* 243:106-11; and Risseuw et al., (1995) *Plant J* 7:109-19.

[0312] An endogenous, non-selectable gene was targeted in *Arabidopsis* using a targeting vector containing a region of about 7 kb homologous to the target gene and the targeting frequency was estimated to be at least 3.9×10^{-4} (Maio and Lam, (1995) *Plant J* 7:359-65). In another example, using a positive-negative selection scheme and a targeting vector containing up to 22.9 kb of sequence homologous to the target, homologous recombination was detected with a frequency less than 5.3×10^{-5} , despite the large flanking sequences available for recombination (Thybjær et al., (1997) *Plant Mol Biol* 35:523-30). In *Arabidopsis*, the AGL5 MADS-box gene was knocked out by homologous recombination using a targeting construct consisting of a kanamycin-resistance cassette inserted into the AGL5 sequence roughly 3 kb from the 5' end and 2 kb from the 3' end. Of the 750 kanamycin-resistant transgenic lines that were generated, one line contained the anticipated insertion (Kempin et al., (1997) *Nature* 389:802-3). Hanin et al., obtained homologous recombination events at a basal frequency of 7×10^{-4} using 3 kb 5'-end and 2 kb 3'-end homology to the *Arabidopsis* PPO gene encoding protoporphyrinogen oxidase (Hanin et al., (2001) *Plant J* 28:671-7). Terada et al., targeted the Waxy locus in rice using an *Agrobacterium*-mediated transformation procedure. Negative selection, in the form of two copies of the dipterin toxin gene placed at both ends of T-DNA, was used to eliminate random integration of T-DNAs, allowing for enrichment of rare homologous recombination events in the selected material, and their transformation system generated thousands of events from just 150 rice seeds. The reported frequency of homologous recombination of the waxy gene in rice was 0.65×10^{-3} , without inclusion of elements to enhance homologous recombination (Terada et al., (2002) *Nat Biotech* 20:1030-4).

[0313] DNA double-strand breaks (DSBs) appear to be an effective factor to stimulate homologous recombination pathways in every organism tested to date (Puchta et al., (1995) *Plant Mol Biol* 28:281-92; Tzfira and White, (2005) *Trends Biotechnol* 23:567-9; Puchta, (2005) *J Exp Bot* 56:1-14). Using DNA-breaking agents, two-to nine-fold increase of homologous recombination was observed between artificially constructed homologous DNA repeats in plants (Puchta et al., (1995) *Plant Mol Biol* 28:281-92). In maize protoplasts, experiments with linear DNA molecules demonstrated enhanced homologous recombination between plasmids (Lyznik et al., (1991) *Mol Gen Genet* 230:209-18).

[0314] The effects of DSBs on homologous recombination have been investigated by using rare-cutting enzymes as well as transposons such as Ac and Mutator (Chiurazzi et al., (1996) *Plant Cell* 8:2057-66; Puchta et al., (1996) *Proc. Natl. Acad. Sci. USA* 93:5055-60; Xiao and Peterson, (2000) *Mol Gen Genet* 263:22-9; and Shalev and Levy (1997) *Genetics* 146:1143-51). Chiurazzi et al., (1996) *Plant Cell* 8:2057-66 introduced DSBs into an *Arabidopsis* chromosome using HO-endonuclease and observed 10-fold increase in the frequency of homologous recombination between

repeats flanking the HO recognition site. Excision of Ac transposable elements also stimulated homologous recombination between repeats flanking the elements at an even higher frequency (Xiao and Peterson (2000) *Mol Gen Genet* 263:22-9).

[0315] Puchta et al. reported that homologous recombination frequency at an artificial target locus was increased by up to two orders of magnitude when DSBs were generated using I-SceI (Puchta et al., (1996) *Proc. Natl. Acad. Sci. USA* 93:5055-60). In the experiment reported in Puchta et al., an I-SceI expression cassette was introduced into transgenic tobacco target lines together with targeting construct by co-inoculation with the two respective *Agrobacterium* strains. Homologous recombination between T-DNA containing the targeting construct and the target site reconstituted the kanamycin-resistance gene (nptII). There was an apparent correlation between frequency of homologous recombination and the amount of I-SceI expression cassette, suggesting that more DSBs yielded higher homologous recombination frequency.

[0316] High frequency of homologous recombination at a pre-introduced artificial target site was obtained using a zinc-finger nuclease (ZFN) in tobacco (Wright et al., (2005) *Plant J* 44:693-705). The zinc-finger nuclease expression cassette and donor DNA were introduced into protoplasts by co-electroporation and targeted modification was monitored by kanamycin resistance and GUS activity. One modified event was observed in approximately every 10 transformants, however, only 20% of the modified events contained the desired homologous recombination products as indicated by Southern blot analysis.

[0317] Zinc finger nucleases are engineered endonucleases with altered specificities, for example by fusion of an engineered DNA binding domain to an endonuclease, for example, FokI (Durai et al., (2005) *Nucleic Acids Res* 33:5978-90; Mani et al., (2005) *Biochem Biophys Res Comm* 335:447-57). Wright et al., and Lloyd et al., reported a high frequency mutagenesis at a DNA target site integrated into tobacco or *Arabidopsis* chromosomal DNA using zinc-finger nucleases (Wright et al., (2005) *Plant J* 44:693-705; Lloyd et al., (2005) *Proc. Natl. Acad. Sci. USA* 102:2232-7). Using a designed zinc-finger nuclease recognizing a tobacco endogenous acetolactate synthase (ALS) gene locus, a mutated ALS gene known to confer resistance to imidazolinone and sulphonylurea herbicides was introduced to replace the endogenous ALS gene at frequencies exceeding 2% of transformed cells (Townsend et al., (2009) *Nature* 459:442-5). The knock-out of an endogenous gene and the expression of a transgene can be achieved simultaneously by gene targeting. The IPK1 gene, which encodes inositol-1,3,4,5,6-pentakisphosphate 2-kinase needed in the final step of phytate biosynthesis in maize seeds, was targeted using a designed zinc-finger nuclease to insert via homologous recombination a PAT gene, which encodes phosphinothricin acetyl transferase tolerance to glufosinate ammonium herbicides such as bialaphos. The disruption of the IPK1 gene with the insertion of the PAT gene resulted in both herbicide tolerance and the expected alteration of the inositol phosphate profile in developing seeds (Shukla et al., (2009) *Nature* 459:437-41).

[0318] Members of the serine family of recombinases produce double-strand breaks at the recombination sites as a part of their catalytic activities (Grindley et al., (2006) *Ann Rev Biochem* 16:16). The R/RS system in sweet orange

appeared to induce mutations of RS sites leading to chromosomal deletions not associated with site-specific recombination reactions per se (Ballester et al., (2006) *Plant Cell Rep* 26:39-45).

[0319] Another approach uses protein engineering of existing homing endonucleases to alter their target specificities. Homing endonucleases, such as I-SceI or I-CreI, bind to and cleave relatively long DNA recognition sequences (18 bp and 22 bp, respectively). These sequences are predicted to naturally occur infrequently in a genome, typically only 1 or 2 sites/genome. The cleavage specificity of a homing endonuclease can be changed by rational design of amino acid substitutions at the DNA binding domain and/or combinatorial assembly and selection of mutated monomers (see, for example, Arnould et al., (2006) *J Mol Biol* 355:443-58; Ashworth et al., (2006) *Nature* 441:656-9; Doyon et al., (2006) *J Am Chem Soc* 128:2477-84; Rosen et al., (2006) *Nucleic Acids Res* 34:4791-800; and Smith et al., (2006) *Nucleic Acids Res* 34: e149; Lyznik et al., (2009) U.S. Patent Application Publication No. 20090133152A1; Smith et al., (2007) U.S. Patent Application Publication No. 20070117128A1). Engineered meganucleases have been demonstrated that can cleave cognate mutant sites without broadening their specificity. An artificial recognition site specific to the wild type yeast I-SceI homing nuclease was introduced in maize genome and mutations of the recognition sequence were detected in 1% of analyzed F1 plants when a transgenic I-SceI was introduced by crossing and activated by gene excision (Yang et al., (2009) *Plant Mol Biol* 70:669-79). More practically, the maize liguleless locus was targeted using an engineered single-chain endonuclease designed based on the I-CreI meganuclease sequence. Mutations of the selected liguleless locus recognition sequence were detected in 3% of the TO transgenic plants when the designed homing nuclease was introduced by *Agrobacterium*-mediated transformation of immature embryos (Gao et al., (2010) *Plant J* 61:176-87).

EXAMPLES

[0320] The present invention is further defined in the following Examples, in which parts and percentages are by weight and degrees are Celsius, unless otherwise stated. It should be understood that these Examples, while indicating embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions. Such modifications are also intended to fall within the scope of the appended claims.

[0321] The meaning of abbreviations is as follows: "sec" means second(s), "min" means minute(s), "h" means hour(s), "d" means day(s), "μL" means microliter(s), "mL" means milliliter(s), "L" means liter(s), "μM" means micromolar, "mM" means millimolar, "M" means molar, "mmol" means millimole(s), "μmole" mean micromole(s), "g" means gram(s), "μg" means microgram(s), "ng" means nanogram(s), "U" means unit(s), "bp" means base pair(s) and "kb" means kilobase(s).

[0322] The DNA repair mechanisms of cells are the basis of transformation to introduce extraneous DNA or induce mutations on endogenous genes. DNA homologous recombination is a specialized way of DNA repair that the cells

repair DNA damages using a homologous sequence. In plants, DNA homologous recombination happens at frequencies too low to be used in transformation until it has been found that the process can be stimulated by DNA double-strand breaks (Bibikova et al., (2001) *Mol. Cell Biol.* 21:289-297; Puchta and Baltimore, (2003) *Science* 300:763; Wright et al., (2005) *Plant J.* 44:693-705).

Example 1

DNA Double-Strand-Break-Induced Alteration of an Endogenous Target Site

[0323] When a DNA double-strand-break-inducing agent recognizes and cleaves the specific recognition sequence at a target site in the genome, a DNA double-strand break is formed triggering the cell DNA repair mechanisms to mobilize to repair the damage that could be fatal to the cell. The process can be utilized in plant transformation to introduce mutations specifically at the target site to knock out the gene residing at the target site or to insert a donor DNA of interest at the target site. Once the DNA double-strand break is formed, depending on the designs of the DNA constructs involved and the actual processes of DNA repair, different outcomes can be obtained serving different transformation purposes.

[0324] For simple site-specific gene mutations, a target site containing a recognition sequence (FIG. 1A) and a DNA double-strand break agent such as an endonuclease (FIG. 1B) that recognizes specifically the recognition sequence have to be present in the same cell. After the endonuclease recognizes and cuts the DNA, the two free ends can be repaired through end joining by the cell DNA repair machinery without the intervention of any external factors. The two ends can be repaired to its original state so no change can be detected or they can be altered before being repaired resulting detectable changes after they are connected again such as the deletion of one or more nucleotides of the recognition sequence and possibly extra surrounding sequences (FIG. 1F). Mutations are introduced at the target site by the latter process.

[0325] To achieve site-specific DNA insertions, a donor DNA containing the DNA of interest has to be simultaneously present in the cell in addition to the target site and the endonuclease. The donor DNA can contain the same DNA sequences that flank the target site to flank the gene of interest, i.e., the homologous sequences (FIG. 1C). The DNA of interest can be inserted at the target site by homologous recombination

[0326] (FIG. 1E), a process that is stimulated by the DNA double-strand break at the target site. The donor DNA can also contain only the DNA of interest without any flanking homologous sequences (FIG. 1D). The DNA of interest can still be inserted at the target site though in a less predictable fashion through non-homologous recombination. Similarly, any unrelated DNA that happens to be present when the DNA ends are repaired can be inserted at the target site (FIG. 1G). The different outcomes (FIGS. 1E-G) can be obtained simultaneously in the same transformation experiment.

[0327] Any means to make a DNA double-strand break in vivo can be used as the DNA double-strand-break-inducing agent such as the most commonly used meganucleases which recognize >18 bp sequences, which are long enough to be unique in most genomes. Even numerous meganucleases have been found and characterized to recognize many

different sequences, but such sequences are often not naturally present in important crops such as soybean or maize and even if similar sequences can be found in crop genomes, the limited numbers of these sequences are still too small to be useful. Certain meganucleases such as I-Crel can be modified by protein engineering in such a way that it will no longer preferentially recognize the recognition sequence of wild type I-Crel and instead will preferentially recognize specifically selected sequences of interest. Taking advantage of the flexibility of the I-Crel endonuclease, one can design and make a modified I-Crel to cleave a target site of our choice in the genome and subsequently introduce mutations or insert genes of interest at the selected target site. The precise genetic engineering that this methodology provides will solve many problems that traditional plant transformation methods such as *Agrobacterium* infection and biolistic bombardment currently face, such as unpredictable integration, unwanted endogenous gene interruption, unpredicted transgene expression, etc.

[0328] In one embodiment of the invention, we used engineered I-Crel-like meganucleases that recognize selected different endogenous target sites in the soybean genome and produced mutations and insertions at the selected target sites.

Example 2

Production of a Complex Trait Locus in the Soybean Genome Near a Transgenic Event for Oil Quality Using Engineered Meganucleases

[0329] Soybean lines comprising an endogenous target recognition sequence in their genome were contacted with a custom designed meganuclease, derived from I Crel, which is designed to specifically recognize and create a double-strand break in the endogenous target sequence. Soybean embryos comprising an endogenous target site were contacted with the components described below, events selected and characterized.

A. TS21, TS14, TS30 and TS5 Target Sites

[0330] Sequence analyses were done for about 500000 bp genomic region in soybean near a transgenic event of interest (event DP-305423-1, U.S. Patent Application Publication No. 2008/0312082 A1, published Dec.-18-2008). A series of soybean genomic endogenous target recognition sequences, referred to as TS21, TS14, TS30 and TS5, were selected for design of custom double-strand break inducing agents derived from I-Crel meganuclease. Each of these target recognition sequences is a unique 22 bp polynucleotide. The target recognition sites have the following sequences:

TS21 target	(SEQ ID NO: 1)
GGCACTCTCGTGT▼GTGATTAAA	
TS14 target	(SEQ ID NO: 2)
CAGACGTACGCAA▼GTAGCTTTG	
TS30 target	(SEQ ID NO: 3)
GAGTCCCACGCAA▼GAGCATAAA	

-continued

TS5 target	(SEQ ID NO: 4)
AAGACTTACGTGT▼GTACTCGTG	

[0331] The double-strand break sites and overhang regions are shown in bold, the enzyme cuts after C13, as indicated by the solid triangle.

[0332] Within the soybean genome, TS5 is about 600 kbp upstream of, and on the same chromosome as, the transgenic event of interest. TS30, TS21 and TS14 are on the same chromosome as TS5 and are 120 kbp, 125 kbp and 500 kbp downstream of the transgenic event of interest (FIG. 2).

B. TS21, TS14, TS30, and TS5 Meganucleases

[0333] The I-Crel meganuclease was modified to produce the TS21, TS14, TS30 and TS5 meganucleases, which are designed to recognize their corresponding target sequences, under a contract with Precision Biosciences (Durham, NC USA). Wild-type I-Crel meganuclease is a homodimer. In order to recognize their target sequences, different substitutions were made to each monomer. The coding sequences for each monomer were joined by a linker sequence to produce single-chain fusion polypeptides. Genes encoding the designed meganucleases were optimized for expression in plants. SEQ ID NO: 9 is the plant-optimized nucleotide sequence of the TS21 meganuclease. SEQ ID NO: 10 is the plant-optimized nucleotide sequence of the TS14 meganuclease. SEQ ID NO: 11 is the plant-optimized nucleotide sequence of the TS30 meganuclease. SEQ ID NO: 12 is the plant-optimized nucleotide sequence of the TS5 meganuclease. These genes include a nucleus localization signal from the SV40 virus (SEQ ID NO: 34) and an intron from the potato ST-LS1 gene. The intron prevents expression of the genes in bacteria during the cloning process, but is not necessary for expression in plant cells. In these plant-optimized nucleotide sequences (SEQ ID NOs: 9-16) nucleotides 1-30 encode an SV40 nucleus localization amino acid sequence, nucleotides 100-261 and nucleotides 850-1011 encode the 1st half and 2nd half target site binding amino acid sequences, respectively, nucleotides 403-591 are the potato ST-LS1 intron, and nucleotides 685-798 encode the amino acid sequence of the polypeptide that links the two re-engineered I-Crel monomers into a single chain.

[0334] Plant optimized nucleotide sequences without the ST-LS1 intron encoding the engineered meganucleases were constructed as well (see, SEQ ID NO: 33 for example).

C. Vector Construction for Plant Expression Vectors of the Meganuclease Genes and Repair DNAs for Transgene Integration by Homologous Recombination

[0335] Vectors comprising expression cassettes for the appropriate meganuclease were constructed using standard molecular biological techniques. All custom designed meganucleases were tested including TS21, TS14, TS30 and TS5. For each of the meganucleases, a plant expression vector comprising a polynucleotide encoding one of the meganuclease genes was operably linked to a soybean constitutive promoter.

[0336] The following meganuclease plant expression vectors were made:

[0337] RTW317 (SEQ ID NO: 35, GM-EF1A pro::TS21::pinII) expression cassette contains the TS21 meganuclease plant optimized sequence without an intron and driven by soybean EF1A promoter.

[0338] RTW322 (SEQ ID NO: 36, GM-EF1A pro::TS21 with ST-LS1 intron2::pinII) expression cassette contains the TS21 meganuclease plant optimized sequence with an intron and driven by soybean EF1A promoter. Other expression cassettes were made in a similar manner as RTW317 and RTW322, but contained a different promoter, or meganuclease, such as: RTW319 (GM-EF1A pro::TS14::pinII), RTW324 (GM-EF1A pro::TS14 with ST-LS1 intron2::pinII), RTW323 (GM-EF1A pro::TS5 with ST-LS1 intron2::pinII), RTW325 (GM-EF1A pro::TS30 with ST-LS1 intron2::pinII), RTW345 (GM-UBQ pro::TS21::pinII), RTW334 (GM-UBQ pro: TS21 with ST-LS1 intron2::pinII), RTW351 (GM-MTH1 pro::TS21::pinII), RTW339 (GM-MTH1 pro::TS21 with ST-LS1 intron2::pinII), wherein GM-ETF1A is the soybean ETF1A promoter, GM-UBQ is the soybean ubiquitin promoter, GM-MTH1 is the soybean MTH1 promoter, and pinII is the pinII terminator.

[0339] To achieve site-specific DNA insertions, a repair DNA (donor DNA) containing the gene of interest has to be simultaneously present in the cell in addition to the target site and the endonuclease. The gene of interest was flanked by two homologous recombination fragments (HR1 and HR2), which were 1 to 3 kb long genomic DNA sequences flanking the meganuclease target sites. The gene of interest can be inserted at the target site by DNA homologous recombination, a process that is stimulated by the DNA double-strand break at the target site.

[0340] A repair DNA (or donor DNA) fragment, Rep-RTW328A (SEQ ID NO: 37) was made for gene integration at TS21 target site in the soybean genome. The RTW328 repair DNA consists of a 1020 bp TS21 HR1 fragment (SEQ ID NO:17), a hygromycin selection marker cassette and a 1000 bp TS21 HR2 fragment (SEQ ID NO:18). The hygromycin selection marker was driven by a SCP1 promoter and a NOS terminator (U.S. Pat. No. 6,072,050; Suzuki et al., Gene (2000) 242 (1-2): 331-336). Similar repair DNA vectors were made for TS14, TS30, and TS5 target sites in soybean genome. The Rep-TS14 repair DNA vector consists of a 1000 bp TS14 HR1 fragment (SEQ ID NO: 19, the same hygromycin selection marker cassette and a 928 bp TS14 HR2 fragment (SEQ ID NO:20). The Rep-TS30 repair DNA vector (consists of a 1000 bp TSO HR1 fragment (SEQ ID NO:21), the same hygromycin selection marker cassette and a 1009 bp TS30 HR2 fragment (SEQ ID NO:22). The Rep-TS5 repair DNA vector consists of a 1006 bp TS5 HR1 fragment (SEQ ID NO:23), the same hygromycin selection marker cassette and a 1007 bp TS5 HR2 fragment (SEQ ID NO:24).

[0341] A DNA double-strand break agent was simultaneously introduced with the repair DNA to facilitate homologous DNA recombination. It is convenient to transiently express the custom designed meganuclease by co-bombardment of a meganuclease expression vector with its corresponding repair DNA in soybean transformation. The presence or absence of an ST-LS1 intron in the DNA nucleotide sequence encoding a meganuclease did not affect the func-

tionality of the meganuclease. Alterations at the target site were observed when expression of the meganuclease with both a DNA sequence that included or excluded the ST-LS1 intron in the expression cassette.

D. Genomic Sequence Modifications and Transgene Integration at Endogenous Target Sites with Custom Designed Meganucleases

[0342] PCR and qPCR assays were done following established protocols using gene-specific primers and probes (Li et al., (2007) *Plant Mol Biol* 65:329-41; Li et al., (2009) *Plant Physiol* 151:1087-95). qPCR assays specific to the TS21, TS14, TS30, and TS5 target sequences were developed to identify sequence changes that happen in the region. The primers and probe were designed as below and tested. TS21 qPCR:

[0343] Mega21-190F (SEQ ID NO:38)

[0344] Mega21-301R (SEQ ID NO:39)

[0345] Mega21-250T (SEQ ID NO:40)

TS14 qPCR:

[0346] Mega14-13F (SEQ ID NO:41)

[0347] Mega14-128R (SEQ ID NO:42)

[0348] Mega14-85T (SEQ ID NO:43)

TS30 qPCR:

[0349] Mega30-30F (SEQ ID NO:44)

[0350] Mega21-87R (SEQ ID NO:45)

[0351] Mega21-52T (SEQ ID NO:46)

TS5 qPCR:

[0352] Mega5-F1 (SEQ ID NO:47)

[0353] Mega5-R1 (SEQ ID NO:48)

[0354] Mega5-T1 (SEQ ID NO:49)

[0355] All hygromycin resistant soybean transgenic events were first analyzed by qPCR assays of the meganuclease target site. Changes in the meganuclease target sequence caused by DNA cleavage and repair result in the copy number reduction of the meganuclease target site from two copies in wild type soybean genome to either one or zero copies in the transgenic events. These “qPCR hit” events with reduced target site copy numbers were chosen for further genomic PCR and sequencing analyses. From qPCR analyses of the TS21, TS14, TS30 and TS5 target sites, it was shown that the copy numbers of the target sites in most of the positive transgenic events were reduced by half, indicating one allele of the target sites in soybean genome was disrupted by meganuclease cutting/DNA repair mechanism.

[0356] Two groups of genomic PCR amplifications were carried out to further characterize these candidate events from qPCR assay to understand the genomic sequence modifications and transgene integrations. The first group of genomic PCRs were designed to identify mutations in the meganuclease target sites, by amplifying genomic fragments containing the TS21 target site using a primer that anneals in HR1 and another primer that anneals in HR2. For example, for TS21, the primer set WOL133 and WOL134 (SEQ ID NO:50 and 51) were used to amplify genomic fragments containing the TS21 target site (FIG. 3A). The PCR products were cloned and sequenced to identify mutations at the TS21 target site. In some cases, a meganuclease in vitro cutting assay to cut the PCR product of an unmodified target site was used to test if the target site had been modified. In the in vitro cutting assay, the PCR products amplified using primers directed to the target site were digested with the meganuclease at 37° C. overnight. Samples with meganuclease enzyme were treated with pro-

teinase K and SDS to denature the protein. The digestion products were separated on a 1.5 to 2% agarose gel. Undigested products indicate that the target site was modified. The undigested PCR products were then cloned and sequenced to verify the genome sequence modification. An example of the soybean genome sequence modification on TS21 target site is shown in FIG. 3B.

[0357] With this approach, soybean genome sequence modifications were detected at TS5, TS14 and TS30 target sites (FIG. 4 and Table 1).

TABLE 1

qPCR copy number analyses of TS30 target sites, pinII (representing the meganuclease cassette) and Hygro (representing the repair DNA cassette)			
Clone ID	TS30 qPCR Copy#	pinII qPCR copy#	Hygro qPCR copy#
A 7052.2.5	0.56	0.00	1.98
A 7052.10.26	0.55	0.00	1.55
A 7052.10.28	0.54	0.00	1.96
A 7034.1.11	0.53	0.00	2.98
A 7034.3.1	0.54	1.70	3.41
A 7034.3.15	0.52	0.96	4.54
WT control	0.96	2.23	5.19

[0358] The copy numbers of the TS30 target sites in positive transgenic events were reduced by half, indicating one allele of the target sites in soybean genome was disrupted by meganuclease cutting/DNA repair mechanism. These results demonstrate that introduction of the meganuclease gene into the plant cell leads to modifications in the genomic region of interest.

[0359] Both wild type soybean and transgenic embryos have been used in the soybean transformation. The target modification rate (qPCR) with TS21 is the same in wild type soybean and the transgenic event. These results demonstrated that we can directly introduce genome modifications in the transgenic event or introduce genome modifications to the same locus in wild type soybean.

[0360] The second group of genomic PCR amplifications was more focused on transgene integration with border specific PCR. For example, for TS21 (FIG. 3A), the primer set WOL190 (SEQ ID NO:52) and WOL242 (SEQ ID NO:53) were designed and used to amplify the left border DNA fragment that results from transgene integration. WOL190 is a sequence specific primer located in soybean genome 5' beyond the TS21 HR1 region and WOL242 is a sequence specific primer to the 5' hygromycin-resistance marker gene coding sequence in the reverse orientation. An 1860 bp PCR product can only be obtained when the RTW328A repair DNA is integrated by homologous recombination facilitated by a double-strand break introduced at the genomic target site by TS21 meganuclease. Another set of primers, WOL153 (SEQ ID NO:54) and WOL247 (SEQ ID NO: 55), was also designed and used to amplify the right border DNA fragment that results from transgene integration. WOL153 is the sense primer from the NOS terminator and the WOL247 is a sequence specific primer located in soybean genome 3' beyond the TS21 HR2 region. A 1727 bp PCR product can only be obtained when the RTW328A repair DNA is integrated by homologous recombination facilitated by a double-strand break introduced at the genomic target site by TS21 meganuclease. Similar genomic PCR primers have been designed and tested for other custom designed meganuclease.

TS21 qPCR

Target Site Primers

[0361] WOL133 (SEQ ID NO:50)
[0362] WOL134 (SEQ ID NO:51)

Left Border Primers

[0363] WOL190 (SEQ ID NO:52)
[0364] WOL242 (SEQ ID NO:53)

Right Border Primers

[0365] WOL153 (SEQ ID NO:54)
[0366] WOL247 (SEQ ID NO:55)

TS14 qPCR

Target Site Primers

[0367] WOL121 (SEQ ID NO:56)
[0368] WOL150 (SEQ ID NO:57)

Left Border Primers

[0369] WOL192 (SEQ ID NO:58)
[0370] WOL242 (SEQ ID NO:53)

Right Border Primers

[0371] WOL153 (SEQ ID NO:54)
[0372] WOL193 (SEQ ID NO:59)

TS30 qPCR

Target Site Primers

[0373] WOL113 (SEQ ID NO:60)
[0374] WOL114 (SEQ ID NO:61)

Left Border Primers

[0375] WOL194 (SEQ ID NO:62)
[0376] WOL242 (SEQ ID NO:53)

Right Border Primers

[0377] WOL153 (SEQ ID NO:54)
[0378] WOL195 (SEQ ID NO:63)

TS5 qPCR

Target Site Primers

[0379] WOL105 (SEQ ID NO:64)
[0380] WOL144 (SEQ ID NO:65)

Left Border Primers

[0381] WOL196 (SEQ ID NO:66)
[0382] WOL242 (SEQ ID NO:53)

Right Border Primers

[0383] WOL153 (SEQ ID NO:54)
[0384] WOL197 (SEQ ID NO:67)

[0385] Primer pairs were designed with one primer capable of annealing to either the 5' or 3' sequence flanking a target site and another primer capable of annealing to a sequence within the potential insert (i.e., the transgene). For the TS14 target site, 18 qPCR positive events were identified from total 68 events by qPCR analyses. Out of the 18 qPCR positive events, three events were confirmed to be perfect

TS14 meganuclease mediated transgene integration events by homologous recombination.

[0386] These results demonstrate that soybean cells possess natural DNA repair machinery that can repair DNA double-strand break ends by simple end joining or by homologous recombination. It is thus expected that similar rates of site-directed mutagenesis and gene insertion via homologous recombination can be achieved at any target sites in the soybean genome using proper double-strand break inducing agents specific to the target recognition sequences. Using a simple PCR screening procedure described herein, it is practical to identify such insertion and mutation events. A perfect transgene integration event can be identified when both left border PCR and right border PCR indicate insertion at the target site. Transgene integration at the pre-defined target sites within a genomic region of interest provides a novel gene stacking technology. FIG. 5 is a schematic example of stacking new trait genes into a single target site in close proximity to a transgenic event of interest.

Example 3

Production of a Complex Trait Locus in the Soybean Genome Near a Herbicide Resistance Transgenic Event Using Engineered Meganucleases

A. TS7, TS4, TS22 and TS24 Target Sites

[0387] The transgene border analyses of a herbicide resistance transgenic event (Event 3560.4.3.5 described in U.S. Patent Application Publication Nos. 2010/0184079, 2009/0036308, and 2008/0051288) showed that the transgene was inserted in a soybean chromosome about 12 cM away from three disease resistance markers based on molecular marker analyses (FIG. 6). Sequence analyses were done for about 400000 bp in this genomic region of interest and four meganuclease target sites (TS7, TS4, TS22 and TS24) were identified with desirable genetic distances between these target sites and nearby disease resistance markers, and a herbicide resistance transgenic event. Each of these target recognition sequences is a unique 22 bp polynucleotide. The target recognition sites have the following sequences:

TS7 target	(SEQ ID NO: 5)
GACATTGTC GTGA ▼GAAAAGAGA	
TS4 target	(SEQ ID NO: 6)
AAATCTGTCT TTGC ▼GAAACGGCA	
TS22 target	(SEQ ID NO: 7)
TATTCTCTC ATAA ▼ATAAACTTT	
TS24 target	(SEQ ID NO: 8)
GGAATGGAC ATAA ▼GAGAACTGT	

[0388] The double-strand break sites and overhang regions are shown in bold, the enzyme cuts after C13, as indicated by the solid triangle.

B. TS7, TS4, TS22 and TS24 Meganucleases

[0389] The I-CreI meganuclease was modified to produce the TS7, TS4, TS22 and TS24 meganucleases, which are

designed to recognize their corresponding target sequences, under a contract with Precision Biosciences (Durham, NC USA). Wild-type I-CreI meganuclease is a homodimer. In order to recognize their target sequences, different substitutions were made to each monomer. The coding sequences for each monomer were joined by a linker sequence to produce single-chain fusion polypeptides. All these target sites are about 1 to 10 cM away from the cluster of the three disease resistance markers.

[0390] The plant optimized nucleotide sequence encoding the TS7 meganuclease (SEQ ID NO: 13), TS4 meganuclease (SEQ ID NO: 14), TS22 meganuclease (SEQ ID NO: 15) and TS24 meganuclease (SEQ ID NO: 16) includes a DNA fragment (from bp 1-30) encoding an SV40 nuclear localization signal (MAPKKKRRKVH; SEQ ID NO: 34) as well as a ST-LS1 intron (from bp 403 to bp 591 of SEQ ID 13-16) in order to eliminate expression in *E. coli* and *Agrobacterium*. Nucleotides 685-798 of SEQ ID NOs: 13-16 encode the amino acid sequence of the polypeptide that links the two engineered I-CreI monomers into a single chain. Nucleotides 100-261 of SEQ ID NOs: 13-16 and nucleotides 850-1011 of SEQ ID NOs: 13-16 encode the first half and the second half target site binding amino acid sequences, respectively.

C. Vector Construction for Plant Expression Vectors of the Meganuclease Genes and Repair DNAs for Transgene Integration by Homologous Recombination

[0391] Vectors comprising expression cassettes for the appropriate meganuclease were constructed using standard molecular biological techniques. All custom designed meganucleases were tested including TS7, TS4, TS22 and TS24. For each of the meganucleases, a plant expression vector comprising a polynucleotide encoding one of the meganuclease genes was operably linked to a soybean constitutive promoter.

[0392] To achieve site-specific DNA insertions, a repair DNA (donor DNA) containing the DNA of interest has to be simultaneously present in the cell in addition to the target site and the endonuclease. The DNA of interest was flanked by two homologous recombination fragments (HR1 and HR2), which were 1 to 3 kb long genomic DNA sequences flanking the meganuclease target sites. The DNA of interest can be inserted at the target site by DNA homologous recombination, a process that is stimulated by the DNA double-strand break at the target site.

[0393] The HR1 and HR2 domains for TS7, TS4, TS22 and TS24 are SEQ ID NOs: 25 and 26, SEQ ID NOs: 27 and 28, SEQ ID NOs: 29 and 30 and SEQ ID NOs: 31 and 32, respectively.

[0394] Repair DNA vectors were made as described in Example 2 C.

[0395] A DNA double-strand break agent was simultaneously introduced with the repair DNA to facilitate homologous DNA recombination. It is convenient to transiently express the custom designed meganuclease by co-bombardment of a meganuclease expression vector with its corresponding repair DNA in soybean transformation.

Example 4

Cluster of Meganuclease Target Sites in a Short Region of the Soybean Genome for Stacking of Multiple Trait Genes

[0396] As shown in FIG. 7, a series of meganuclease target sites can be identified with desirable genetic distances

between these target sites. Custom designed meganucleases can be used to target a series of trait genes into this defined genome locus either by sequential transformation or by genetic crosses with individual trait genes. Using this method depicted in FIG. 7, multiple traits can be stacked in a genomic region of interest that comprises, for example, a transgene or native gene of interest, and other transgenic traits or native trait loci such as disease resistance markers.

Example 5

Production of a Complex Trait Locus at a Maize Endogenous Locus by Engineered Meganucleases

a. MHP Target Sites

[0397] A genomic region encompassing about 1.8 million nucleotides and representing a genetic region of approximately 4.3 centimorgans (CM) on a maize chromosome was chosen as a target region for generation of a complex trait locus. The genomic region was scanned for 22-mer sequences that could serve as target sites containing recognition sequences for double-strand-break inducing meganucleases and be useful for insertion of additional transgenes in order to create a complex trait locus. A series of 35 putative target sites (SEQ ID NOs: 68-77) were selected in a 2 cM region (FIG. 8) in close proximity of the transgene insertion site for design of custom double-strand break inducing agents derived from I-Crel meganuclease. FIG. 8 show the genetic and physical location of the MHP target sites relative to each other and the transgene of interest.

B. MHP Meganucleases

[0398] The I-Crel meganuclease was modified to produce endonucleases, which were designed to recognize their corresponding target sequences, (SEQ ID NOs: 68-77). The design of custom made meganucleases has been described in United States Patent Application Publication No. US 2007/0117128 A1.

[0399] Genes encoding the designed meganucleases were optimized for expression in plants. The engineered endonuclease expression cassettes contained the maize codon-optimized nucleotide sequences for better performance in maize cells. The endonuclease gene sequences were also supplemented with DNA sequences encoding a SV40 nuclear localization signal (SEQ ID NO: 34). The maize ubiquitin promoter and the potato proteinase inhibitor II gene terminator sequences completed the endonuclease gene designs. The MHP55 (SEQ ID NO:80) expression cassette was additionally modified by addition of the ST-LS1 intron to the coding sequence of the first monomer in order to eliminate its expression in *E. coli* and *Agrobacterium*. SEQ ID NO: 82 is the plant-optimized nucleotide sequence of MHP55-2 containing a nuclear localization signal and without an intron. SEQ ID NO: 78 is the plant-optimized nucleotide sequence of the MHP14 meganuclease. A custom designed meganuclease, referred to as MHP14+ was made as well. SEQ ID NO: 79 is the plant-optimized nucleotide sequence of the MHP14+ meganuclease. SEQ ID NO: 83 is the plant-optimized nucleotide sequence of the MHP77 meganuclease

C. Vector Construction for Plant Expression Vectors of the Meganuclease Genes and Repair (Donor) DNAs for Transgene Integration by Homologous Recombination

[0400] Vectors comprising expression cassettes for the appropriate meganuclease were constructed using standard

molecular biological techniques. For each of the meganucleases, a plant expression vector comprising a polynucleotide encoding one of the meganuclease genes was operably linked to a maize constitutive promoter.

[0401] To achieve site-specific DNA insertions, a repair DNA (donor DNA) containing the gene of interest has to be simultaneously present in the cell in addition to the target site and the meganuclease. A vector (PHP44285, SEQ ID NO: 104), or PHP44779, SEQ ID NO: 105) containing a polynucleotide encoding the engineered meganuclease MHP14, or the optimized meganuclease MHP14+, and a donor DNA was constructed using standard molecular biology techniques. The donor DNA contained an herbicide resistance gene used as the selection marker for transformation. The herbicide resistance gene MoPAT encodes a phosphinothricin acetyltransferase, and was flanked by two homologous recombination fragments, HR1 (SEQ ID NO: 84) and HR2 (SEQ ID NO: 85), which were about 1 kb long genomic DNA sequences flanking the meganuclease target sites. Each vector PHP44285 or PHP44779 contained the meganuclease cassette, the donor DNA and the homology sequences HR1 and HR2.

[0402] Maize immature embryos 9-12 DAP (days after pollination, approximately 1.5-2.0 mm in size) from a maize transformable line were used for gene transformation by bombardment (Example 6). The immature embryos were placed on 560Y medium for 4 hours at 26° C. or alternatively, immature embryos were incubated at temperatures ranging from 26° C. to 37° C. for 8 to 24 hours prior to placing on 560Y preceding bombardment (as described in Example 6). Developmental genes ODP2 (AP2 domain transcription factor ODP2 (Ovule development protein 2); US20090328252 A1) and Wushel were included in the experiments through co-bombardment (Example 7). Maize immature embryos were transformed with the vectors PHP44285 or PHP44779.

D. Genomic Sequence Modifications and Transgene Integration at Endogenous Target Sites with Custom Designed Meganuclease

[0403] Successful delivery of the MHP14 donor vector (PHP 44285 or PHP44779) conferred bialaphos herbicide resistance, and was used to identify putative events by callus selection on herbicide containing media. Callus tissues and/or plants regenerated from stable transformants using standard culture and regeneration conditions were screened for modification of the endogenous MHP14 target site.

[0404] Real time PCR (qPCR) was used to determine the target site copy number. Two copies of the target site indicate that both alleles are wild type and that no modification occurred at the target site. One copy means one allele of the target site has changed during repair of the double strand break generated by the MHP14 or MHP14+, while absence of the target site (null) is the result of both alleles modified. The copy number can also be in between 1 and 2 due to chimeric nature of callus samples. The probe sequence for qPCR of MHP14 target site was CAGATTCACGTCA-GATTT (SEQ ID NO: 106), the MHPTS14_forward primer was AGCGACATAGTGGTGTATAAAAGGAA (SEQ ID NO: 107) and MHPTS14_reverse primer was TGGAT-TGTAATATGTGTACCTCATGCT (SEQ ID NO: 108). The amplicon was approximately 100 bp.

[0405] To examine whether increased temperature would increase the rate of target site modification, maize embryos were incubated at different temperatures following bom-

bombardment with several meganucleases. Table 2 shows the effect of temperature on the meganuclease activity of MHP14 as determined by target site modification. Table 2 indicates that increased temperature results in increased target site mutation rate.

TABLE 2

Effect of incubating maize embryos at increased temperature post-bombardment on target site mutation rate of meganucleases		
Meganuclease	Temperature (° C.)	Target Site Mutation Rate
MHP14	28	14%
MHP14	32	46%

[0406] Following bombardment, embryos were incubated on 560P (maintenance medium) for 12 to 48 hours at 28° C. or 32° C. and then placed at 28° C. Herbicide-resistant events were screened for modification at the target site by measuring target site copy-number using qPCR. Target site mutation rate indirectly measures the meganuclease activity. TSMutRate (target site mutation rate) indicated the modification rate of the MHP14 or LIG3/4 target site (#events with modification/#events*100%). As shown in Table 2, target site mutation rate for both MHP14 and LIG34 was approximately 3× higher when embryos were placed at 32° C. for 48 hours after bombardment compared to no temperature elevation treatment.

[0407] Maize calli were also screened for integration of the transgene cassette from the donor DNA (PHP44285 or PHP44779) at the MHP14 target site through junction PCR and selected callus events were regenerated into TO plants. FIG. 9A shows an outline of PCR screening for integration of the donor DNA fragment via homologous recombination at MHP14 target site (PHP44779 donor). Arrows indicate primer locations. FIG. 9B shows PCR of MHP14 callus events: B1-B12 Junction PCR with primers 146773/146775; b1-b12 Junction PCR with primers 146772/146778. Two events (B2 and B5) yielded the predicted 1-1.2 kb PCR fragments that result from integration by homologous recombination for both junctions. PCR products from TO plants derived from these callus events were sequenced to verify the callus results. PCR screening revealed integration of the herbicide resistance transgene cassette at MHP14 target site. Primers were from the genomic region outside of the homology of donor vector and from the transgene cassette close to the end of the homology.

[0408] FIG. 10A shows a schematic outline of long fragment PCR reactions used to confirm UBI: moPAT: PinII cassette integration at the endogenous MHP14 target. FIG. 10B: shows the results of long fragment PCR on TO plants from three events where integration occurred at the target site. The plant A5 was from event #1, A6-A8 event #2, and C4-C6 event #3. 10B-left shows the long junction fragment PCR on the HR1 side using genomic primer (146775) and moPAT primer (mopatR2); 10B-right shows the long junction fragment PCR on HR2 side (mopatF2/146772). Arrows indicated PCR primer locations. Primer set 146772/mopatF2 amplified a 4 kb fragment, spanning from moPAT gene through the UBI intron, UBI promoter, and the HR2 sequence to the adjacent genomic region. Primer set 146775/mopatR2 amplified a 2.2 kb fragment, spanning from the moPAT gene through the HR1 to the adjacent genomic region. These two fragments overlapped and covered the

whole insert at MHP14 target site. The sizes of the two long PCR products indicate a perfect integration of the donor gene cassette at MHP 14 target site

[0409] To determine the segregation pattern of the integration events in progeny, T1 seeds from selfed TO plants were planted in flats and T1 plants genotyped by using PCR and/or qPCR. The segregation ratio of integration genotypes fit 1:2:1 for wild type (no integration), heterozygous (one allele having integration and the other wild-type) and homozygous integration of the transgene at the MHP14 target site, demonstrating Mendelian inheritance. No visible phenotype was observed in the homozygous or heterozygous integration plants.

[0410] The entire inserted fragment of UBI: moPAT: PinII was obtained by using PCR on DNA from homozygous T1 plants with primers in the genomic region outside of the HR1 and HR2 (146772/146775). A PCR product of 5 kb was amplified from homozygous plants as expected. A 2 kb PCR product was amplified from the unmodified intact genomic sequence from wild-type plants.

[0411] Trait gene cassettes can be introduced at other target sites of the complex trait locus through homologous recombination mediated by engineered meganucleases. Engineered meganucleases were designed to direct double strand breaks a two other MHP target sites, MHP55 (SEQ ID NO: 72) and MHP77 (SEQ ID NO: 74) within the complex trait locus. Target site modification was determined using qPCR. The probe sequence for qPCR screening of the MHP55 target site was AACCGTCGTGAGACCT (SEQ ID NO: 115), the MHPTS55_Forward_MGBprimer sequence was AAGGCGCAGCCGTTGAG (SEQ ID NO: 116), and MHP55_reverse_MGB primer was CTACCGGTTTCGCGTGCTCT (SEQ ID NO: 117). The probe sequence for qPCR of MHP77 target site was TAGTATGACATACATACCGCC (SEQ ID NO: 118), the MHPTS77_Forward_MGB primer sequence was TCCT-TAGGGCGGTATGTATGTCA (SEQ ID NO: 119), and MHP77_reverse_MGB primer was CATCGGT-CAAAAAACACATAAACTTT (SEQ ID NO: 120). The trait gene cassettes encoding MHP14, MHP55 and MHP77 were introduced into maize somatic embryos via transformation techniques using bombardment and following bombardment, embryos were incubated on 560P (maintenance medium) for 48 hours at. As shown in Table 3, maize callus containing the MHP55 target site bombarded with PHP45782 or PHP46924 which include genes encoding MHP55 or MHP55.2 meganucleases, respectively, also lead to an observed increase in the target site mutation rate modified MHP55.2 variant. In addition, maize callus containing a MHP77 target site bombarded with vectors PHP45970 or PHP50238 which include genes encoding MHP77 or MHP77.3 meganucleases, respectively, showed a higher frequency of mutated target sites from callus bombarded with the modified variant MHP77.3. Taken together, like MHP14, these meganucleases directed mutations to their corresponding target sites and modified versions lead to an increase in the target site mutation rate (approx 2 to 10-fold increase when compared to their original versions) suggesting the newly designed versions of the meganucleases were more active than the original nucleases.

TABLE 3

Meganuclease activity (defined as target site mutation rate) of original and modified meganucleases	
Meganuclease	Target Site Mutation Rate
MHP55	0%
MHP55-2	5%
MHP77	1%
MHP77-3	11%
MHP14	29%
MHP14+	40%

[0412] The mutations observed at these target sites indicated that the engineered meganucleases were functional and that the target sites can be used for integration of additional trait genes.

E. Production of a Complex Trait Locus at a Maize Endogenous Locus by Crossing

[0413] A maize event obtained through random integration containing a transgene DNA of interest was identified and MHP14, MHP55 and MHP77 target sites surrounding the transgenic DNA of interest were identified as described above. Other maize events containing a modification at the MHP14, MHP55 and MHP77 target site (through addition of herbicide resistance gene as described above) were also identified.

[0414] Plants homozygous for the integration of a herbicide resistance gene at the MHP14 target site were crossed with homozygous maize plants containing the transgene DNA of interest. The cross resulted in fertile plants producing F1 seeds. The F1 seeds were planted and out-crossed with Elite inbred line plants and screened for the stacked phenotype. Additional trait genes can be added to the complex trait locus by crossing one transgenic event containing n-transgenes with other transgenic events containing the additional trait gene at the additional target site, and progeny can be screened for the presence of n+1 transgenes. This process can be repeated as many times as the amount of target sites are present in the complex trait locus.

F. Production of a Complex Trait Locus at a Maize Endogenous Locus by Serial Transformation

[0415] A complex trait locus can be also be created by serial transformation. A first transformed line containing a first trait gene integrated at a first MHP target site can be used to supply embryos. The first transformed line can be retransformed with a second trait gene and a vector encoding a second engineered meganuclease; resulting in the second trait gene being integrated at a second MHP target site through homologous recombination mediated by the second engineered meganuclease. The homozygous integration plants containing a selectable marker at the MHP14 target site can be used to supply embryos. Two rounds of transformations will create two traits at the MHP locus. A transformed line that is homozygous for integration events with two trait genes at MHP target sites can be used to supply embryos for another retransformation, and a third trait gene can be introduced to a third target site.

Example 6

Transformation of Maize Immature Embryos

[0416] Transformation can be accomplished by various methods known to be effective in plants, including particle-

mediated delivery, *Agrobacterium*-mediated transformation, PEG-mediated delivery, and electroporation.

A. Particle-Mediated Delivery

[0417] Transformation of maize immature embryos using particle delivery is performed as follows. Media recipes follow below.

[0418] The ears are husked and surface sterilized in 30% Clorox bleach plus 0.5% Micro detergent for 20 minutes, and rinsed two times with sterile water. The immature embryos are isolated and placed embryo axis side down (scutellum side up), 25 embryos per plate, on 560Y medium for 4 hours and then aligned within the 2.5-cm target zone in preparation for bombardment. Alternatively, isolated embryos are placed on 560L (Initiation medium) and placed in the dark at temperatures ranging from 26° C. to 37° C. for 8 to 24 hours prior to placing on 560Y for 4 hours at 26° C. prior to bombardment as described above.

[0419] A plasmid comprising the Zm-BBM (also referred to as Zm-ODP2) coding sequence (set forth in SEQ ID NO: 9) operably linked to a promoter is constructed. This could be a weak promoter such as nos, a tissue-specific promoter, such as globulin-1 or oleosin, an inducible promoter such as In2, or a strong promoter such as ubiquitin plus a plasmid containing the selectable marker gene phosphinothricin N-acetyltransferase (PAT; Wohlleben et al. (1988) *Gene* 70:25-37) that confers resistance to the herbicide bialaphos. Furthermore, plasmids containing the double strand brake inducing agent and donor DNA such as PHP44285 or PHP44779 are constructed as described above and co-bombarded with the plasmids containing the developmental genes ODP2 (AP2 domain transcription factor ODP2 (Ovule development protein 2); US20090328252 A1) and Wushel.

[0420] The plasmids are precipitated onto 1.1 µm (average diameter) tungsten pellets using a calcium chloride (CaCl₂) precipitation procedure by mixing 100 µl prepared tungsten particles in water, 10 µl (1 µg) DNA in Tris EDTA buffer (1 µg total DNA), 100 µl 2.5 M CaCl₂, and 10 µl 0.1 M spermidine. Each reagent is added sequentially to the tungsten particle suspension, with mixing. The final mixture is sonicated briefly and allowed to incubate under constant vortexing for 10 minutes. After the precipitation period, the tubes are centrifuged briefly, liquid is removed, and the particles are washed with 500 ml 100% ethanol, followed by a 30 second centrifugation. Again, the liquid is removed, and 105 µl 100% ethanol is added to the final tungsten particle pellet. For particle gun bombardment, the tungsten/DNA particles are briefly sonicated. 10 µl of the tungsten/DNA particles is spotted onto the center of each macrocarrier, after which the spotted particles are allowed to dry about 2 minutes before bombardment.

[0421] The sample plates are bombarded at level #4 with a Biorad Helium Gun. All samples receive a single shot at 450 PSI, with a total of ten aliquots taken from each tube of prepared particles/DNA.

[0422] Following bombardment, the embryos are incubated on 560P (maintenance medium) for 12 to 48 hours at temperatures ranging from 26 C to 37 C, and then placed at 26 C. After 5 to 7 days the embryos are transferred to 560R selection medium containing 3 mg/liter Bialaphos, and subcultured every 2 weeks at 26 C. After approximately 10 weeks of selection, selection-resistant callus clones are transferred to 288J medium to initiate plant regeneration.

Following somatic embryo maturation (2-4 weeks), well-developed somatic embryos are transferred to medium for germination and transferred to a lighted culture room. Approximately 7-10 days later, developing plantlets are transferred to 272V hormone-free medium in tubes for 7-10 days until plantlets are well established. Plants are then transferred to inserts in flats (equivalent to a 2.5" pot) containing potting soil and grown for 1 week in a growth chamber, subsequently grown an additional 1-2 weeks in the greenhouse, then transferred to Classic 600 pots (1.6 gallon) and grown to maturity. Plants are monitored and scored for transformation efficiency, and/or modification of regenerative capabilities.

[0423] Initiation medium (560L) comprises 4.0 g/l N6 basal salts (SIGMA C-1416), 1.0 ml/l Eriksson's Vitamin Mix (1000x SIGMA-1511), 0.5 mg/l thiamine HCl, 20.0 g/l sucrose, 1.0 mg/l 2,4-D, and 2.88 g/l L-proline (brought to volume with D-I H₂O following adjustment to pH 5.8 with KOH); 2.0 g/l Gelrite (added after bringing to volume with D-I H₂O); and 8.5 mg/l silver nitrate (added after sterilizing the medium and cooling to room temperature).

[0424] Maintenance medium (560P) comprises 4.0 g/l N6 basal salts (SIGMA C-1416), 1.0 ml/l Eriksson's Vitamin Mix (1000x SIGMA-1511), 0.5 mg/l thiamine HCl, 30.0 g/l sucrose, 2.0 mg/l 2,4-D, and 0.69 g/l L-proline (brought to volume with D-I H₂O following adjustment to pH 5.8 with KOH); 3.0 g/l Gelrite (added after bringing to volume with D-I H₂O); and 0.85 mg/l silver nitrate (added after sterilizing the medium and cooling to room temperature).

[0425] Bombardment medium (560Y) comprises 4.0 g/l N6 basal salts (SIGMA C-1416), 1.0 ml/l Eriksson's Vitamin Mix (1000x SIGMA-1511), 0.5 mg/l thiamine HCl, 120.0 g/l sucrose, 1.0 mg/l 2,4-D, and 2.88 g/l L-proline (brought to volume with D-I H₂O following adjustment to pH 5.8 with KOH); 2.0 g/l Gelrite (added after bringing to volume with D-I H₂O); and 8.5 mg/l silver nitrate (added after sterilizing the medium and cooling to room temperature).

Selection medium (560R) comprises 4.0 g/l N6 basal salts (SIGMA C-1416), 1.0 ml/l Eriksson's Vitamin Mix (1000x SIGMA-1511), 0.5 mg/l thiamine HCl, 30.0 g/l sucrose, and 2.0 mg/l 2,4-D (brought to volume with D-I H₂O following adjustment to pH 5.8 with KOH); 3.0 g/l Gelrite (added after bringing to volume with D-I H₂O); and 0.85 mg/l silver nitrate and 3.0 mg/l bialaphos (both added after sterilizing the medium and cooling to room temperature).

[0426] Plant regeneration medium (288J) comprises 4.3 g/l MS salts (GIBCO 11117-074), 5.0 ml/l MS vitamins stock solution (0.100 g nicotinic acid, 0.02 g/l thiamine HCL, 0.10 g/l pyridoxine HCL, and 0.40 g/l glycine brought to volume with polished D-I H₂O) (Murashige and Skoog (1962) *Physiol. Plant.* 15:473), 100 mg/l myo-inositol, 0.5 mg/l zeatin, 60 g/l sucrose, and 1.0 ml/l of 0.1 mM abscisic acid (brought to volume with polished D-I H₂O after adjusting to pH 5.6); 3.0 g/l Gelrite (added after bringing to volume with D-I H₂O); and 1.0 mg/l indoleacetic acid and 3.0 mg/l bialaphos (added after sterilizing the medium and cooling to 60° C.).

Hormone-free medium (272V) comprises 4.3 g/l MS salts (GIBCO 11117-074), 5.0 ml/l MS vitamins stock solution (0.100 g/l nicotinic acid, 0.02 g/l thiamine HCL, 0.10 g/l pyridoxine HCL, and 0.40 g/l glycine brought to volume with polished D-I H₂O), 0.1 g/l myo-inositol, and 40.0 g/l sucrose (brought to volume with polished D-I H₂O after

adjusting pH to 5.6); and 6 g/l bacto-agar (added after bringing to volume with polished D-I H₂O), sterilized and cooled to 60° C.

B. *Agrobacterium*-Mediated Transformation

[0427] *Agrobacterium*-mediated transformation was performed essentially as described in Djukanovic et al. (2006) *Plant Biotech J* 4:345-57. Briefly, 10-12 day old immature embryos (0.8-2.5 mm in size) were dissected from sterilized kernels and placed into liquid medium (4.0 g/l N6 Basal Salts (Sigma C-1416), 1.0 ml/l Eriksson's Vitamin Mix (Sigma E-1511), 1.0 mg/l thiamine HCl, 1.5 mg/l 2, 4-D, 0.690 g/l L-proline, 68.5 g/l sucrose, 36.0 g/l glucose, pH 5.2). After embryo collection, the medium was replaced with 1 ml *Agrobacterium* at a concentration of 0.35-0.45 OD₅₅₀. Maize embryos were incubated with *Agrobacterium* for 5 min at room temperature, then the mixture was poured onto a media plate containing 4.0 g/l N6 Basal Salts (Sigma C-1416), 1.0 ml/l Eriksson's Vitamin Mix (Sigma E-1511), 1.0 mg/l thiamine HCl, 1.5 mg/l 2, 4-D, 0.690 g/l L-proline, 30.0 g/l sucrose, 0.85 mg/l silver nitrate, 0.1 nM acetosyringone, and 3.0 g/l Gelrite, pH 5.8. Embryos were incubated axis down, in the dark for 3 days at 20° C., then incubated 4 days in the dark at 28° C., then transferred onto new media plates containing 4.0 g/l N6 Basal Salts (Sigma C-1416), 1.0 ml/l Eriksson's Vitamin Mix (Sigma E-1511), 1.0 mg/l thiamine HCl, 1.5 mg/l 2, 4-D, 0.69 g/l L-proline, 30.0 g/l sucrose, 0.5 g/l MES buffer, 0.85 mg/l silver nitrate, 3.0 mg/l Bialaphos, 100 mg/l carbenicillin, and 6.0 g/l agar, pH 5.8. Embryos were subcultured every three weeks until transgenic events were identified. Somatic embryogenesis was induced by transferring a small amount of tissue onto regeneration medium (4.3 g/l MS salts (Gibco 11117), 5.0 ml/l MS Vitamins Stock Solution, 100 mg/l myo-inositol, 0.1 μM ABA, 1 mg/l IAA, 0.5 mg/l zeatin, 60.0 g/l sucrose, 1.5 mg/l Bialaphos, 100 mg/l carbenicillin, 3.0 g/l Gelrite, pH 5.6) and incubation in the dark for two weeks at 28° C. All material with visible shoots and roots were transferred onto media containing 4.3 g/l MS salts (Gibco 11117), 5.0 ml/l MS Vitamins Stock Solution, 100 mg/l myo-inositol, 40.0 g/l sucrose, 1.5 g/l Gelrite, pH 5.6, and incubated under artificial light at 28° C. One week later, plantlets were moved into glass tubes containing the same medium and grown until they were sampled and/or transplanted into soil.

Example 7

Transient Expression of BBM Enhances Transformation

[0428] Parameters of the transformation protocol can be modified to ensure that the BBM activity is transient. One such method involves precipitating the BBM-containing plasmid in a manner that allows for transcription and expression, but precludes subsequent release of the DNA, for example, by using the chemical PEI. In one example, the BBM plasmid is precipitated onto gold particles with PEI, while the transgenic expression cassette (UBI::moPAT~GFPm::PinII; moPAT is the maize optimized PAT gene) to be integrated is precipitated onto gold particles using the standard calcium chloride method.

[0429] Briefly, gold particles were coated with PEI as follows. First, the gold particles were washed. Thirty-five mg of gold particles, 1.0 in average diameter (A.S.I. #162-

0010), were weighed out in a microcentrifuge tube, and 1.2 ml absolute EtOH was added and vortexed for one minute. The tube was incubated for 15 minutes at room temperature and then centrifuged at high speed using a microfuge for 15 minutes at 4° C. The supernatant was discarded and a fresh 1.2 ml aliquot of ethanol (EtOH) was added, vortexed for one minute, centrifuged for one minute, and the supernatant again discarded (this is repeated twice). A fresh 1.2 ml aliquot of EtOH was added, and this suspension (gold particles in EtOH) was stored at -20° C. for weeks. To coat particles with polyethylimine (PEI; Sigma #P3143), 250 µl of the washed gold particle/EtOH mix was centrifuged and the EtOH discarded. The particles were washed once in 100 µl ddH₂O to remove residual ethanol, 250 µl of 0.25 mM PEI was added, followed by a pulse-sonication to suspend the particles and then the tube was plunged into a dry ice/EtOH bath to flash-freeze the suspension, which was then lyophilized overnight. At this point, dry, coated particles could be stored at -80° C. for at least 3 weeks. Before use, the particles were rinsed 3 times with 250 µl aliquots of 2.5 mM HEPES buffer, pH 7.1, with 1× pulse-sonication, and then a quick vortex before each centrifugation. The particles were then suspended in a final volume of 250 µl HEPES buffer. A 25 µl aliquot of the particles was added to fresh tubes before attaching DNA. To attach uncoated DNA, the particles were pulse-sonicated, then 1 µg of DNA (in 5 µl water) was added, followed by mixing by pipetting up and down a few times with a Pipetteman and incubated for 10 minutes. The particles were spun briefly (i.e. 10 seconds), the supernatant removed, and 60 µl EtOH added. The particles with PEI-precipitated DNA-1 were washed twice in 60 µl of EtOH. The particles were centrifuged, the supernatant discarded, and the particles were resuspended in 45 µl water. To attach the second DNA (DNA-2), precipitation using TFX-50 was used. The 45 µl of particles/DNA-1 suspension was briefly sonicated, and then 5 µl of 100 ng/µl of DNA-2 and 2.5 µl of TFX-50 were added. The solution was placed on a rotary shaker for 10 minutes, centrifuged at 10,000g for 1 minute. The supernatant was removed, and the particles resuspended in 60 µl of EtOH. The solution was spotted onto macrocarriers and the gold particles onto which DNA-1 and DNA-2 had been sequentially attached were delivered into scutellar cells of 10 DAP Hi-II immature embryos using a standard protocol for the PDS-1000. For this experiment, the DNA-1 plasmid contained a UBI::RFP::pinII expression cassette, and DNA-2 contained a UBI::CFP::pinII expression cassette. Two days after bombardment, transient expression of both the CFP and RFP fluorescent markers was observed as numerous red & blue cells on the surface of the immature embryo. The embryos were then placed on non-selective culture medium and allowed to grow for 3 weeks before scoring for stable colonies. After this 3-week period, 10 multicellular, stably-expressing blue colonies were observed, in comparison to only one red colony. This demonstrated that PEI-precipitation could be used to effectively introduce DNA for transient expression while dramatically reducing integration of the PEI-introduced DNA and thus reducing the recovery of RFP-expressing transgenic events. In this manner, PEI-precipitation can be used to deliver transient expression of BBM and/or WUS2.

[0430] For example, the particles are first coated with UBI::BBM::pinII using PEI, then coated with UBI::moPAT~YFP using TFX-50, and then bombarded into

scutellar cells on the surface of immature embryos. PEI-mediated precipitation results in a high frequency of transiently expressing cells on the surface of the immature embryo and extremely low frequencies of recovery of stable transformants (relative to the TFX-50 method). Thus, it is expected that the PEI-precipitated BBM cassette expresses transiently and stimulates a burst of embryogenic growth on the bombarded surface of the tissue (i.e. the scutellar surface), but this plasmid will not integrate. The PAT~GFP plasmid released from the Ca⁺⁺/gold particles is expected to integrate and express the selectable marker at a frequency that results in substantially improved recovery of transgenic events. As a control treatment, PEI-precipitated particles containing a UBI::GUS::pinII (instead of BBM) are mixed with the PAT~GFP/Ca⁺⁺ particles. Immature embryos from both treatments are moved onto culture medium containing 3 mg/l bialaphos. After 6-8 weeks, it is expected that GFP+, bialaphos-resistant calli will be observed in the PEI/BBM treatment at a much higher frequency relative to the control treatment (PEI/GUS).

[0431] As an alternative method, the BBM plasmid is precipitated onto gold particles with PEI, and then introduced into scutellar cells on the surface of immature embryos, and subsequent transient expression of the BBM gene elicits a rapid proliferation of embryogenic growth. During this period of induced growth, the explants are treated with *Agrobacterium* using standard methods for maize (see Example 1), with T-DNA delivery into the cell introducing a transgenic expression cassette such as UBI::moPAT~GFPm::pinII. After co-cultivation, explants are allowed to recover on normal culture medium, and then are moved onto culture medium containing 3 mg/l bialaphos. After 6-8 weeks, it is expected that GFP+, bialaphos-resistant calli will be observed in the PEI/BBM treatment at a much higher frequency relative to the control treatment (PEI/GUS).

[0432] It may be desirable to “kick start” callus growth by transiently expressing the BBM and/or WUS2 polynucleotide products. This can be done by delivering BBM and WUS2 5'-capped polyadenylated RNA, expression cassettes containing BBM and WUS2 DNA, or BBM and/or WUS2 proteins. All of these molecules can be delivered using a biolistics particle gun. For example 5'-capped polyadenylated BBM and/or WUS2 RNA can easily be made in vitro using Ambion's mMessage mMachine kit. RNA is co-delivered along with DNA containing a polynucleotide of interest and a marker used for selection/screening such as Ubi::moPAT~GFPm::PinII. It is expected that the cells receiving the RNA will immediately begin dividing more rapidly and a large portion of these will have integrated the agronomic gene. These events can further be validated as being transgenic clonal colonies because they will also express the PAT~GFP fusion protein (and thus will display green fluorescence under appropriate illumination). Plants regenerated from these embryos can then be screened for the presence of the polynucleotide of interest.

[0433] Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.


```
SEQ ID NO: 1          moltype = DNA  length = 22
FEATURE               Location/Qualifiers
source                1..22
                     mol_type = genomic DNA
                     organism = Glycine max
```

```
SEQ ID NO: 2          moltype = DNA  length = 22
FEATURE               Location/Qualifiers
source                1..22
                     mol_type = genomic DNA
                     organism = Glycine max
```

```
SEQ ID NO: 3          moltype = DNA  length = 22
FEATURE               Location/Qualifiers
source                1..22
                     mol_type = genomic DNA
                     organism = Glycine max
```

```
SEQ ID NO: 4      moltype = DNA  length = 22
FEATURE
source            Location/Qualifiers
                  1..22
                  mol_type = genomic DNA
                  organism = Glycine max
```

```
SEQ ID NO: 5          moltype = DNA  length = 22
FEATURE               Location/Qualifiers
source                1..22
                     mol_type = genomic DNA
                     organism = Glycine max
```

```
SEQ ID NO: 6          moltype = DNA  length = 22
FEATURE               Location/Qualifiers
source                1..22
                     mol_type = genomic DNA
                     organism = Glycine max
```

```
SEQ ID NO: 7          moltype = DNA  length = 22
FEATURE
source                1..22
                      mol_type = genomic DNA
                      organism = Glycine max
```

```
SEQ ID NO: 8          moltype = DNA  length = 22
FEATURE               Location/Qualifiers
source                1..22
                     mol_type = genomic DNA
                     organism = Glycine max
```

```

SEQ ID NO: 9      multype = DNA   length = 1272
FEATURE           Location/Qualifiers
misc_feature      1..1272
                  note = synthetic construct; plant optimized nucleotide
                  sequence of TS21 meganuclease
source            1..1272
                  mol_type = other DNA
                  organism = synthetic construct

```

SEQUENCE: 9
atqgcaccqa aqaagaacqg caaggtgcac atgaacacca aqtacaacaa qqagttctctg 60

-continued

```

ctctacctgg cgggtctcgt ggaaggcgac ggctccatca tggcgagat caagccgcag 120
cagtcctgca agttcaagca cgcgctccag ctgaccttca ccgtgaccca gaagacgcag 180
aggcgctggt tcctcgacaa gctgggtcgac gagatcgggg tgggcaaggt ctacgaccgc 240
gggtcggtgt ccgactacat cctctcccag atcaagcccc tgcacaactt cctcaccag 300
ctccagccgt tcctcaagct gaagcagaag caggcgaacc tcgtctgaa gatcatcgag 360
cagctccccct cggccaagga gtccccggac aagttcctgg aggttaagttt ctgctcttac 420
ctttgatata tatataataa ttatcattaa ttagtagtaa tataatattt caaatatttt 480
tttcaaaata aaagaatgta gtatatagca attgcttttc ttagtattat aagtgtgtat 540
attttaattt ataacttttc taatatatga ccaaaacatg gtgatgtgca ggtgtgcacg 600
tgggtcgacc agatcgcggc cctcaacgac agcaagaccc gcaagacgac ctcgagagcg 660
gtgcggggcg tcctggactc cctcccagga tccgtgggag gtctatcgcc atctcaggca 720
tccagcgccg catcctcggc ttctcaagc ccgggttcag ggatctccga agcactcaga 780
gctggagcaa ctaagtccaa ggaattcctg ctctacctgg ccggtctcgt ggaaggcgac 840
ggctccatca aggcgcgac caagccgac agtcccgca agttcaagca cgagctctcc 900
ctgaccttcc aggtgaccca gaagacgcag aggcgctggt tcctcgacaa gctgggtcgac 960
gagatcgggg tgggtacgtt ctacgaccgc ggggtcggtt ccgactacat cctctcccag 1020
atcaagcccc tgcacaactt cctcaacgac ctccagccgt tcctcaagct gaagcagaag 1080
caggcgaacc tcgtctgaa gatcatcgag cagctccccct cggccaagga gtccccggac 1140
aagttcctgg aggtgtgcac gtgggtcgac cagatcgcg ccctcaacga cagcaagacc 1200
cgcaagacga cctcgagac ggtgcgggcg gttctagact ccctcagcga gaagaagaag 1260
tcgtccccct ga 1272

SEQ ID NO: 10      moltype = DNA length = 1272
FEATURE            Location/Qualifiers
misc_feature        1..1272
                    note = synthetic construct; plant optimized nucleotide
                    sequence of TS14 meganuclease
source              1..1272
                    mol_type = other DNA
                    organism = synthetic construct

SEQUENCE: 10
atggcaccga agaagaagcg caaggtgcat atgaacacca agtacaacaa ggagttcctg 60
ctctacctgg cgggtctcgt ggaaggcgac ggctccatca tcggtccat caagccggag 120
cagtcctaca agttcaagca cgcctctcc ctgaccttca ccgtgaccca gaagacgcag 180
aggcgctggt tcctcgacaa gctgggtcgac gagatcgggg tgggcaaggt ctacgaccgc 240
gggtcggtgt ccgactaccg cctctcccag atcaagcccc tgcacaactt cctcaccag 300
ctccagccgt tcctgaagct caagcagaag caggcgaacc tcgtctgaa gatcatcgag 360
cagctgcccc cggccaagga atccccggac aagttcctgg aggttaagttt ctgctcttac 420
ctttgatata tatataataa ttatcattaa ttagtagtaa tataatattt caaatatttt 480
tttcaaaata aaagaatgta gtatatagca attgcttttc ttagtattat aagtgtgtat 540
attttaattt ataacttttc taatatatga ccaaaacatg gtgatgtgca ggtgtgcacg 600
tgggtggacc agatcgcggc cctcaacgac agcaagaccc gcaagacgac ctcgagagcg 660
gtgcggggcg tcctggactc cctcccagga tccgtgggag gtctatcgcc atctcaggca 720
tccagcgccg catcctcggc ttctcaagc ccgggttcag ggatctccga agcactcaga 780
gctggagcaa ctaagtccaa ggaattcctg ctctacctgg ccggtctcgt ggaaggcgac 840
ggctccatca tcgcaagat caccgcgaac cagtcctaca agttcaagca ccagctccag 900
ctgaccttca ccgtgaccca gaagacgcag aggcgctggt tcctcgacaa gctgggtcgac 960
gagatcgggg tgggcaaggt ctacgaccgc ggggtcggtt ccgactacat cctctcccag 1020
atcaagcccc tgcacaactt cctcaccag tcctcagccgt tcctcaagct gaagcagaag 1080
caggcgaacc tcgtctgaa gatcatcgag cagctccccct cggccaagga gtccccggac 1140
aagttcctgg aggtgtgcac gtgggtcgac cagatcgcg ccctcaacga cagcaagacc 1200
cgcaagacga cctcgagac ggtgcgggcg gttctagact ccctcagcga gaagaagaag 1260
tcgtccccct ga 1272

SEQ ID NO: 11      moltype = DNA length = 1272
FEATURE            Location/Qualifiers
misc_feature        1..1272
                    note = synthetic construct; plant optimized nucleotide
                    sequence of TS30 meganuclease
source              1..1272
                    mol_type = other DNA
                    organism = synthetic construct

SEQUENCE: 11
atggcaccga agaagaagcg caaggtgcat atgaacacca agtacaacaa ggagttcctg 60
ctctacctgg cgggtctcgt ggaaggcgac ggctccatca tggcgagat caagccggag 120
cagtcctaca agttcaagca cgcctctatg ctgaccttca ccgtgaccca gaagacgcag 180
aggcgctggt tcctcgacaa gctgggtcgac gagatcgggg tgggtctacgt ctacgaccgc 240
gggtcggtgt cctctcccag atcaagcccc tgcacaactt cctcaccag 300
ctccagccgt tcctcaagct gaagcagaag caggcgaacc tcgtctgaa gatcatcgag 360
cagctccccct cggccaagga gtccccggac aagttcctgg aggttaagttt ctgctcttac 420
ctttgatata tatataataa ttatcattaa ttagtagtaa tataatattt caaatatttt 480
tttcaaaata aaagaatgta gtatatagca attgcttttc ttagtattat aagtgtgtat 540
attttaattt ataacttttc taatatatga ccaaaacatg gtgatgtgca ggtgtgcacg 600
tgggtcgacc agatcgcggc cctcaacgac agcaagaccc gcaagacgac ctcgagagcg 660
gtgcggggcg tcctggactc cctcccagga tccgtgggag gtctatcgcc atctcaggca 720
tccagcgccg catcctcggc ttctcaagc ccgggttcag ggatctccga agcactcaga 780

```

-continued

```

gctggagcaa ctaagtccaa ggaattctg ctctacctg cgggtctcg ggaaggcgac 840
ggctccatca aggcgtccat caccgcgag cagtcctgca agttcaagca cgcgtccag 900
ctgaccttcc aggtgaccca gaagacgcag aggcgtcgtg tctcgcacaa gctggtcgac 960
gagatcgggg tgggcaaggt ctacgaccgc gggtcggtg cgaactaccg cctctcccag 1020
atcaagcccc tgcacaactt cctcaccag ctcacgccg tctcaagct gaagcagaag 1080
caggcgaacc tcgtcctgaa gatcatcgag cagctccct cggccaagga gtcccggac 1140
aagttcctgg aggtgtgcac gtgggtcgac cagatcgcg ccctcaacga cagcaagacc 1200
cgcaagacga cctcggagac ggtgcgggag gttctagact ccctcagcga gaagaagaag 1260
tcgtccccct ga 1272

SEQ ID NO: 12      moltype = DNA length = 1272
FEATURE            Location/Qualifiers
misc_feature        1..1272
                    note = synthetic construct; Plant optimized nucleotide
                    sequence of TS5 meganuclease
source              1..1272
                    mol_type = other DNA
                    organism = synthetic construct

SEQUENCE: 12
atggcaccga agaagaagcg caaggtgcat atgaacacca agtacaacaa ggagttcctg 60
ctctacctgg ccggcttcgt ggaaggcgac gggtccatca tcgcgcagat caagccggag 120
cagtcctaca agttcaagca cgcctctccc ctgaccttca cgtgaccca gaagacgcag 180
aggcgtcgtg tctcgcacaa gctggtcgac gagatcgggg tgggctacgt ctacgaccgc 240
gggtcgggtg ccgactaccg cctctcccag atcaagcccc tgcacaactt cctcaccag 300
ctccagccgt tcctgaagct caagcagaag caggccaacc tcgtgctgaa gatcatcgag 360
cagctgccct ccgccaagga atcccggac aagttcctgg aggttaagttt ctgcttctac 420
ctttgatata tatataataa ttatcattaa ttagtagtaa tataatattt caaatatttt 480
tttcaaaata aaagaatgta gtatatagca attgcttttc tgtagtttat aagtggtgat 540
attttaattt ataacttttc taatatatga ccaaaacatg gtgatgtgca ggtgtgcacg 600
tgggtcgacc agatcgcggc cctcaacgac agcaagaccc gcaagacgac ctcgagagcg 660
gtgcggggcg tcctggactc cctcccagga tccgtgggag gtctatcgcc atctcaggca 720
tccagcgccg catcctcggc ttctcaagc ccgggttcag ggatctccga agcactcaga 780
gctggagcaa ctaagtccaa ggaattctg ctctacctg cgggtctcg ggaaggcgac 840
ggctccatca tcgcgcagat ctcccgcgag cagtcctaca agttcaagca cgcgtccag 900
ctgaccttca ccgtgaccca gaagacgcag aggcgtcgtg tctcgcacaa gctggtcgac 960
gagatcgggg tgggtcagct ctacgaccgc gggtcggtg cgaactaccg cctctcccag 1020
atcaagcccc tgcacaactt cctcaccag ctcacgccg tctcaagct gaagcagaag 1080
caggcgaacc tcgtcctgaa gatcatcgag cagctccct cggccaagga gtcccggac 1140
aagttcctgg aggtgtgcac gtgggtcgac cagatcgcg ccctcaacga cagcaagacc 1200
cgcaagacga cctcggagac ggtgcgggag gttctagact ccctcagcga gaagaagaag 1260
tcgtccccct ga 1272

SEQ ID NO: 13      moltype = DNA length = 1272
FEATURE            Location/Qualifiers
misc_feature        1..1272
                    note = synthetic construct; Plant optimized nucleotide
                    sequence of TS7 meganuclease
source              1..1272
                    mol_type = other DNA
                    organism = synthetic construct

SEQUENCE: 13
atggcaccga agaagaagcg caaggtgcat atgaacacca agtacaacaa ggagttcctg 60
ctctacctgg ccggcttcgt ggaaggcgac gggtccatca tcgcgcagat caccgcgag 120
cagtcctaca agttcaagca ctcctcccag ctgaccttcc aggtgaccca gaagacgcag 180
aggcgtcgtg tctcgcacaa gctggtcgac gagatcgggg tgggcaaggt ccgacaccgc 240
gggtcgggtg ccgactacat cctctcccag atcaagcccc tgcacaactt cctcaccag 300
ctccagccgt tctcaagct gaagcagaag caggcgaacc tcgtcctgaa gatcatcgag 360
cagctccctt cggccaagga gtcccggac aagttcctgg aggttaagttt ctgcttctac 420
ctttgatata tatataataa ttatcattaa ttagtagtaa tataatattt caaatatttt 480
tttcaaaata aaagaatgta gtatatagca attgcttttc tgtagtttat aagtggtgat 540
attttaattt ataacttttc taatatatga ccaaaacatg gtgatgtgca ggtgtgcacg 600
tgggtcgacc agatcgcggc cctcaacgac agcaagaccc gcaagacgac ctcgagagcg 660
gtgcggggcg tcctggactc cctcccagga tccgtgggag gtctatcgcc atctcaggca 720
tccagcgccg catcctcggc ttctcaagc ccgggttcag ggatctccga agcactcaga 780
gctggagcaa ctaagtccaa ggaattctg ctctacctg ccggcttcgt ggaaggcgac 840
ggctccatca tcgcgcagat ctcccgcgag cagtcgcgca agttcaagca catcctctcc 900
ctgaccttcc aggtgaccca gaagacgcag aggcgtcgtg tctcgcacaa gctggtcgac 960
gagatcgggg tgggcaaggt ctacgaccgc gggtcggtg ccgactaccg cctctcccag 1020
atcaagcccc tgcacaactt cctcaccag ctcacgccg tctcaagct gaagcagaag 1080
caggcgaacc tcgtcctgaa gatcatcgag cagctccct cggccaagga gtcccggac 1140
aagttcctgg aggtgtgcac gtgggtcgac cagatcgcg ccctcaacga cagcaagacc 1200
cgcaagacga cctcggagac ggtgcgggag gttctagact ccctcagcga gaagaagaag 1260
tcgtccccct ga 1272

SEQ ID NO: 14      moltype = DNA length = 1272
FEATURE            Location/Qualifiers

```

-continued

```

misc_feature      1..1272
                  note = synthetic construct; Plant optimized nucleotide
                  sequence of TS4 meganuclease
source            1..1272
                  mol_type = other DNA
                  organism = synthetic construct

```

SEQUENCE: 14

```

atggcaccga agaagaagcg caaggtgcat atgaacacca agtacaacaa ggagttcctg 60
ctctacctgg ccggtctcgt ggacggcgac ggctccatca tcgcgcagat ccgcccgccg 120
cagtcgccga agttcaagca cgagctcgag ctgcgcttcc aggtgaccca gaagacgcag 180
agggcgtggt tctcgacaaa gctggtcgac gagatcgggg tgggtacagt ctacgaccgc 240
gggtcggtgt ccgactacat cctctcccag atcaagcccc tgcacaactt cctcaccagg 300
ctccagccgt tctcaagct gaagcagaag caggcgaacc tcgtcctgaa gatcatcgag 360
cagctcccc cggccaagga gtccccggac aagttcctgg aggttaagttt ctgcttctac 420
ctttgatata tatataataa ttatcattaa ttagtagtaa tataatat t caaatat ttt 480
tttcaaaata aaagaatgta gtatatagca attgcttttc ttagtttat aagtggtgat 540
attttaattt ataacttttc taatatatga ccaaaacatg gtgatgtgca ggtgtgcacg 600
tgggtcgacc agatcgcggc cctcaacgac agcaagaccc gcaagacgac ctcgagagcg 660
gtgcggcgcg tctcggaact cctcccagga tccgtgggag gtctatcgcc atctcaggca 720
tccagcgccg catcctcggc ttctcaagc ccgggttcag ggatctcgca agcactcaga 780
gctggagcaa ctaagtccaa ggaattctct ctctacctgg cgggcttcgt ggacggcgac 840
ggctccatca tcgcgcagat caagccgaac cagtcctaca agttcaagca ccagctcatg 900
ctgaccttcc aggtgaaccca gaagacgcag aggcgctggt tctcgacaaa gctggtcgac 960
gagatcgggg tgggtacagt ccgcgaccgc gggtcggtgt ccgactacat cctctcccag 1020
atcaagcccc tgcacaactt cctcaccagg ctccagccgt tctcaagct gaagcagaag 1080
caggcgaacc tcgtcctgaa gatcatcgag cagctcccc cggccaagga gtccccggac 1140
aagttcctgg aggtgtgcac gtgggtcgac cagatcgcgg ccctcaacga cagcaagacc 1200
cgcaagacga cctcgagac ggtgcgggcg gttctagact ccctcagcga gaagaagaag 1260
tcgtccccct ga                                     1272

```

```

SEQ ID NO: 15      moltype = DNA length = 1272
FEATURE            Location/Qualifiers
misc_feature        1..1272
                  note = synthetic construct; Plant optimized nucleotide
                  sequence of TS22 meganuclease
source              1..1272
                  mol_type = other DNA
                  organism = synthetic construct

```

SEQUENCE: 15

```

atggcaccga agaagaagcg caaggtgcat atgaacacca agtacaacaa ggagttcctg 60
ctctacctgg ccgggtctcgt ggacggcgac ggctccatca tcgcgcagat ctccccgaac 120
cagtcctaca agttcaagca ccagctccgc ctgaccttca ccgtgaccca gaagacgcag 180
agggcgtggt tctcgacaaa gctggtcgac gagatcgggg tgggcagtgt ctacgaccag 240
gggtcggtgt cccactaccg cctctcccag atcaagcccc tgcacaactt cctcaccagg 300
ctccagccgt tctcaagct gaagcagaag caggcgaacc tcgtcctgaa gatcatcgag 360
cagctcccc cggccaagga gtccccggac aagttcctgg aggttaagttt ctgcttctac 420
ctttgatata tatataataa ttatcattaa ttagtagtaa tataatat t caaatat ttt 480
tttcaaaata aaagaatgta gtatatagca attgcttttc ttagtttat aagtggtgat 540
attttaattt ataacttttc taatatatga ccaaaacatg gtgatgtgca ggtgtgcacg 600
tgggtcgacc agatcgcggc cctcaacgac agcaagaccc gcaagacgac ctcgagagcg 660
gtgcggcgcg tctcggaact cctcccagga tccgtgggag gtctatcgcc atctcaggca 720
tccagcgccg catcctcggc ttctcaagc ccgggttcag ggatctcgca agcactcaga 780
gctggagcaa ctaagtccaa ggaattctct ctctacctgg cgggcttcgt ggacggcgac 840
ggctccatca aggcgcagat caagccgcag cagtgctaca agttcaagca ccgctcatg 900
ctgaccttcc aggtgaaccca gaagacgcag aggcgctggt tctcgacaaa gctggtcgac 960
gagatcgggg tgggtacagt ctacgaccgc gggtcggtgt ccgactacat cctctcccag 1020
atcaagcccc tgcacaactt cctcaccagg ctccagccgt tctcaagct gaagcagaag 1080
caggcgaacc tcgtcctgaa gatcatcgag cagctcccc cggccaagga gtccccggac 1140
aagttcctgg aggtgtgcac gtgggtcgac cagatcgcgg ccctcaacga cagcaagacc 1200
cgcaagacga cctcgagac ggtgcgggcg gttctagact ccctcagcga gaagaagaag 1260
tcgtccccct ga                                     1272

```

```

SEQ ID NO: 16      moltype = DNA length = 1272
FEATURE            Location/Qualifiers
misc_feature        1..1272
                  note = synthetic construct; Plant optimized nucleotide
                  sequence of TS24 meganuclease
source              1..1272
                  mol_type = other DNA
                  organism = synthetic construct

```

SEQUENCE: 16

```

atggcaccga agaagaagcg caaggtgcat atgaacacca agtacaacaa ggagttcctg 60
ctctacctgg ccgggtctcgt ggacggcgac ggctccatca tcgcgtccat caccgccgaac 120
cagtcgccga agttcaagca ccagctccag ctgaccttca ccgtgaccca gaagacgcag 180
agggcgtggt tctcgacaaa gctggtcgac gagatcgggg tgggcaaggt ccgcgaccgc 240
gggtcggtgt ccgactaccg cctctcccag atcaagcccc tgcacaactt cctcaccagg 300

```

-continued

```

ctccagccgt tectcaagct gaagcagaag caggcgaacc tcgtcctgaa gatcatcgag 360
cagctccctt cggccaagga gtccccggac aagttcctgg aggttaagttt ctgcttttac 420
ttttgatata tatataataa ttatcattaa ttagtagtaa tataatattt caaatatttt 480
tttcaaaata aaagaatgta gtatatagca attgcttttc ttagtattat aagtgtgtat 540
attttaattt ataacttttc taatatatga ccaaaacatg gtgatgtgca ggtgtgcacg 600
tgggtcgacc agatcgcggc cctcaacgac agcaagaccc gcaagacgac ctgggagacg 660
gtgcggggcg tcttggaact cctcccagga tccgtgggag gtctatcgcc atctcaggca 720
tccagcgccg catcctcggc ttctccaagc cggggttcag ggatctcga agcactcaga 780
gtctggagca ctaagtccaa ggaattcctg ctctacctgg cgggcttcgt ggacggcgac 840
ggctccatca aggcgcagat cccccgaa cagtcctgca agttcaagca ccagctccgc 900
ctgaccttcc aggtgaccca gaagacgcag aggcgctggt tctcgacaa gctggtcgac 960
gagatcgggg tgggcaaggt ctacgaccgc gggtcgggtg ccgactacat cctctcccag 1020
atcaagcccc tgcacaactt cctcaccag ctccagccgt tctcaagct gaagcagaag 1080
caggcgaaac tctgctctga catcatcgag cagctccctt cggccaagga gtccccggac 1140
aagttccttg aggtgtgcac gtgggtcgac cagatcgcg ccctcaacga cagcaagacc 1200
cgcaagacga ctcggagac ggtgcggcg gttctagact ccctcagcga gaagaagaag 1260
tcgtccctct ga 1272

```

```

SEQ ID NO: 17      moltype = DNA length = 1020
FEATURE           Location/Qualifiers
source            1..1020
                  mol_type = genomic DNA
                  organism = Glycine max

```

```

SEQUENCE: 17
ttatttatcc cctataaagg gcaccagtta gttcaatctg atgtctaacc taatttggat 60
acatgccttt tattgcagct gccgtccgtg cacagaggag tcttaggagg aacaactgta 120
gagaaaagga tctgccaatt tcgctagaaa attcaccaga aacaccaccc gttatccaat 180
taaaccaagat tttggatca cttgtgaagt tgaattgcta tccaactgct attccattt 240
ctaaaccttg ttacacgagc atcttgatca atgggtctga aagggaataa gcagtgagtg 300
ggtgcttcaa cgataagtta ttggatttag tatttatctt agcctgtttt cgtgtacttt 360
gttttgccgg atggaggtat gtgattttgt ctatgattct taatacaata acctacatt 420
actctcattg atagtttgtg cagatctaata agctatgaag caccgatacc ggacatgaca 480
cggtcaggtg gacacatgta atgtctaaaa tattaataa tagaacgtag tacgagtgtc 540
gtgtcgggtg tagatactga tagggacgcg tgtcggacac cggacatgac aaaggactga 600
agtgtctaga tattgtttatg tttagatctt tgttgatgag aggcagatag aggtcaactt 660
gccaagataa cctacagttc tatattagat gctttgtgca aaaacgatca tccaaaggct 720
attggattat tcaagaaaac taaagaccaa ggagttcaaa accgctatg tacacatgca 780
ctatacttat ggaaggattg tgcgaagtgg aagacttcag aatgcaaaaa tgatttttca 840
ggatctactg attaaaggct atcaactaag tgtctgtctg tactctgtat aatgttatga 900
ttcataggct ttgtaaaagg ggattttttg atgaagcatt gatctagaaa tctaaattgg 960
aaaacagatc ttaagaagaa tacactgtgt aaatgtgtaa tggcactggc actctcgtgt 1020

```

```

SEQ ID NO: 18      moltype = DNA length = 1000
FEATURE           Location/Qualifiers
source            1..1000
                  mol_type = genomic DNA
                  organism = Glycine max

```

```

SEQUENCE: 18
gtgtgtgatt aaaagtcata tatggtttaa gatacttttt tttataaaga tagtagtggt 60
caatttttct atattacaca agtgtttctt tttcttctca ttgtactgta gatctgattt 120
actttcaatg attgtttaag tcaactgggt aattgtttgt gtttcaataa tcaaaccaag 180
ctgaaactga gatgatgatg atttgaaatg ctttatctca tgtagtcgac tcaattttcc 240
tgttatattc ttgttctttt taagaaaca ggagctttta agatttaaaa caccagcata 300
ttttgtttgc ataatacaaa ttgtcttagg tgtaaagtgt ctgacatttc ccttgatgtc 360
attgctgcat aattaatttg agccttttca aaacctatgg tttattttgt tggggattat 420
tcaagggaacg cgtgtctcag tctcaagtgt tatgattgct gatatacagt atatatgtct 480
gcacaatgaa gtggaactat tttaaatttc aattgatgat tctgcatcca atttatcatc 540
tgaccttttt atcttttacc tcatctggca ttttagtctt ttaccagata aaaggaccaa 600
acacatgaga tataatcacc aaatgaaaag aatgaaagac gagatataaa gatgtgggtt 660
ttctttttat tcttggaaga tttagatgat gttttcaatt aagttgtttg tggatgcttt 720
tagatgattt tgttttgcat acatatgttt acttttttgt tctcaacttc tcattcattt 780
tccatgattt catcccgtag aaaagtgatt tagcagaaaa cgtttttccc ctgttgtctt 840
tgtctaaac ttttggaattc taagtttttt tatatgaaaa ttagatcatt tggcacatgg 900
ttttccaaag acacaagtag actctttcta tgaaatcaat cttaaatccc ttttagagga 960
aaaacatttt aaaggagggtg aacatgttgt ggagtgaggaa 1000

```

```

SEQ ID NO: 19      moltype = DNA length = 1000
FEATURE           Location/Qualifiers
source            1..1000
                  mol_type = genomic DNA
                  organism = Glycine max

```

```

SEQUENCE: 19
attttgtcag tottgtaact tttgaaaact ttttttcttt tttttataga ccaataatat 60
aatatattat attaaaaaaa ccaactttat aacaacatgt aacacgttag caaacagtag 120
atctcaacca aacgttcgaa aacttttgga tattatatat gtggctgttg gcaactgctaa 180
actcagcagt atatctccat tattgatgag tctctcctaa aattatcttt ccaagtctta 240

```

-continued

```

ttttttttt aattgggttag atattaaatt gaaaaataaa ataaaagtgt tgttgttgtg 300
tagttttcgt cactttttact cataagaaaa tatatatact acgttttagca tcttttaaact 360
gaaaactttt cagttggaat gcataacaaa atattggcca agtaattagt acacaaaatc 420
ttgctcaaa gttttgccac catagattta ggttgtgttt aggaacgatta cttaaaatat 480
cattaattga taattgaaac ttcaaatata atttaaaagt ttaaaagtgt aatagttaaa 540
aatgaaagct gaaaataaat aagctaattg attcaatttg aagtatttaa tagtatcaac 600
tagtgaaatt tattcataaa ttctctttta aatatatacc gattttatta gttaatatata 660
aaaaaaaata gtatgaacta ataaaattga tcaaaagtaa attaatataa atataaaatt 720
ttatatgatg aataatcagt agaaaataa aaaaagttag ctctagaaaa gataaaattga 780
tttaattagg gtcattgacaa aattttgcta gcttctattt tagtctgctt tgcttttagaa 840
tatttacatt caaatagctc ttttatagca taacaaacat aaaaaaagct attgattcta 900
cataaaaaaa aaagattaat tatgctattc ttggggacaa aactttttaga tgaatgccaa 960
tttaaaataa ttattaaggt attcaagcag acgtacgcaa 1000

```

```

SEQ ID NO: 20      moltype = DNA length = 928
FEATURE           Location/Qualifiers
source            1..928
                  mol_type = genomic DNA
                  organism = Glycine max

```

```

SEQUENCE: 20
gcaagtagct ttgttacttt cgtattgaca attcaaaatc gtcttttatt tttattttgt 60
ttgttttaaa tagaggactt ttgaagtcg tccatcatgt gtttcttatt ttgtcagttt 120
tgtcactttat gaacactttt ttacagaca aataatatat tatattaaaa aaaccatact 180
tataacaaca acatgtaaca cgttggcaaa cagttaatct caacccaaacg ctgaaaaact 240
tttgatatt atatatatat atgcatggct attggcagtg ctaaaagtcatt cattatcatt 300
ctaaagtcatt cagtatcatt ctaattctca tattgagtggt attcatttca tcaatcactt 360
tgcttttctc atcataacca ccaaaatgcc aaccattaat ccagtgtggt tgaatttcatt 420
ggaaggcata ataacattta tgatgatgat gttgcaggtt gttgtttctg ctcaagacca 480
tattatgtgc attcagactg agagagaagc actcctccaa ttcaaggctg cacttctgga 540
tcactatggc atgctctctt ctgggaccac ttctgattgc tgccaatggc aagggattcg 600
ctgctccaac ctccagccgc atgttctaat gctcgacctt cacagttagt gcctcagagg 660
agagatccac cagtcgttga tggagttgca acaattaaac tattttaaacc tcagtgggaa 720
ttcttttcaa ggcagaggaa tcccagagtt tcttggttct ctccaccaact tgagatacct 780
tgatctgtca cattctgatt ttgaaggaaa aattccaact cagtttggtt ctctttctca 840
tttgaataac ttaaatcttg ctgggaatta ttatctggag ggttcaatcc cacgtcaact 900
tggaatctc tccagttgc agcatctt 928

```

```

SEQ ID NO: 21      moltype = DNA length = 1000
FEATURE           Location/Qualifiers
source            1..1000
                  mol_type = genomic DNA
                  organism = Glycine max

```

```

SEQUENCE: 21
tcatgccagc ctcagccagc ttccaaatca tttccctcgt attggtagaa ggtccaatgg 60
tgcacactat cttcgtcttg cgcctaaacg ttggttaga ccacattcca acagagttct 120
caccaaatgg ctgcacccca cgtaaatggt gcaaatgctc ctcaatctaa acacaaaac 180
acaaaaaggg tcacaagaat tatcccttaa aaactcaaaa aatgcaaaaa acacgaactt 240
ttggatcatc ttggggtgac atggggatga cttcagaagg ggcagatttc cttgcactga 300
tctgaaggct tctgagcctg agtttggagc gtttgttatt gttctctccc aaagggaaca 360
ccttgggaagc aaaagatgga ggttttaaca ggttttgggt tctgtcacgt gcagatccag 420
aagtggggca caatgggggt ctttgaatgg atcgtgaagc cagcacctga gccattgtta 480
gaaagagaga gaaatggggg ggaatgaggaa gaagagagtg tgaggggtat aagaagaagg 540
tgagggggga aatgggaagt gggaaaaatcg ccgctaagtt tggcggaggt tctgagaagg 600
aagccttgtt cgtatcgaaa cacaaaggac actactgtgt ttgaattctg ttcaacgtgt 660
ttgttgttgt aattttattg aaatggactg tacttctttt ctgttttttt ttttcacagt 720
aaaaatgcac tgtatttctt taaatctgct cataaacaat tacacatatt ttattagcta 780
aaatttaata taaattacaa aatatttaca aatattgtga tcaaatataa agtgaacac 840
ataattttat tatttttaac aaatttatct tatgataaag agtgtatttg aaagattatt 900
attaagggac aattttctgt gttgtggaca actttcataa gtgatccatg aaacaccaca 960
ttttatagtc accagattga tctcagattt atgctcttgc 1000

```

```

SEQ ID NO: 22      moltype = DNA length = 1009
FEATURE           Location/Qualifiers
source            1..1009
                  mol_type = genomic DNA
                  organism = Glycine max

```

```

SEQUENCE: 22
ttgcgtggga ctcagatcct gagggaggac atggaagatg tgtcgaaccc aacaagtgtg 60
gctactcatg gcagctccaa caagaagagc tttaaaacta agttcatgca ggaacaaaag 120
gacaggatga aagatgcacc accagagtgt cctgcaggag ttttgttccc aaagtgcagt 180
tcacgcgccac gtcatgagtt tgaatgctca acacaactaa cacacccttg ctaagaagtc 240
ctagaaaaat aaaaaatctaa ggttagagaa tggactaatg atgtacatgt aaaaaataa 300
tgctgaagcc cctgagttta aaagatgtgg attctaacga ctttgataat ttttaattgg 360
attttttata agttaattta ttgattattt ttaataattt tcttatattt ttttattcac 420
aaaacttaaa tctaaaatct tattttatag aataaggaaat tgagttaaat aacaccgata 480
tgttgataat gatttaaatc atgggaatct gttgtttata atagagaaaa aaaaaccctt 540

```

-continued

```

atgatataaa accttcctta gtctaaact cctagctttg tgttaatttg attgtccaaa 600
aggaggctag ctagtttggt ccttcctcgt ccttacttct aaatgcatgg attaaggaaa 660
gaagcttctt tttgtttggt gttatatggt tttgggggtca ttatagtatg ggcaacttaa 720
cactcactct aagagggtgt accttgggcta ttcttagatg gctagacatc aaaactttga 780
atacaaaaatt attaaagatc aataaaaat gattttatta taatactgag attaaattgt 840
aattttaatc tctctttaat tcatgggatg caattttaga ttttattttt tcatgtaatt 900
ttaatcatca cattttaaaa aattcataat tttaaattga tttttaattt tgtatatggt 960
ttatttttaa tttttatcta gttaaaactgt atatttaaca tattgattt 1009

```

SEQ ID NO: 23 moltype = DNA length = 1006

FEATURE Location/Qualifiers

source 1..1006

mol_type = genomic DNA

organism = Glycine max

SEQUENCE: 23

```

attgaaagat tctattttgc ttgtttggtt gtcatttgga taaatgaatg agtactataa 60
acctttgaaa acatataatac tttagagcagt tctatgacct aattttgctt gtttgattag 120
aattttgaat gaatacaaac gaattgcttt gataaatgtg ttatgaattg aacagataat 180
ttcaatgaaa atagatcaaaa attagatcgg ttcaatttta tatatatata tatatatata 240
tatatatata tatatatata tatattggta taaaatattt ttacacaaaa ttaataaaaag 300
ttttaacatt ttataatatt attttgttca ttaataataag gtaatacagt ataaattcct 360
attatgtggt tatataaaatt tctattttta gtccttaatt ttgataattg tcaattaatt 420
ataattttagt cttcaaaaatt tgatattact agtcaactta aacttaataa ttaataaatt 480
agtcaattta atttaaaaat ttgactatat atatatataa atcaaaaggg ttaacaatt 540
catttatcat aaaaactcggg ttaagatcac atgatagcaa agcaattcgt ttgacaattt 600
aaaaattcgg maaatatagt cttagtcata aaacaaaaatc aaaagggtta agattaaatc 660
ataaaactat atttttttta aatgatatcr tgtgatcaat taaaaagac aactttaatt 720
ataatmatct attcactaaa aaaaactaac tcatttgatt gagtagaata tatatrttat 780
tgtacttttr ttactcttga ttctaccaca taattaaaaa caaataatca tctatctatt 840
ttatatagtc tagttttatt cttctcataa cactaaaaaa ttatttaatg atgatatgat 900
cacttaaaaa aattacataa ttatatattc tatatcgtaa ccattcatgt gatataatga 960
tcacattttt tttttctcac actcacctaa gtgcacgagt acacac 1006

```

SEQ ID NO: 24 moltype = DNA length = 1007

FEATURE Location/Qualifiers

source 1..1007

mol_type = genomic DNA

organism = Glycine max

SEQUENCE: 24

```

acacgtaagt cttaggttaa agtttcatgc cccccccccc cccccccccc aaaattacat 60
ttttttccat taaaataact ccaaaactact ctacttctct cgctttggcg gaaccaccat 120
tgctctatga aaaggccatt ttattttcaa ccgcatgtgt ttggaactat acaacgcaaa 180
agccttccat cctcatcatc tgagctactt caaatcttgg ttctctctcg tctctacca 240
gttatgtaag ctttcttctt ttctcttttt tgggtgggca gcacgaaatt atttttcttc 300
ttgttattag ctagaagcac tattctagaa caagcttgca aaaaggactc aagtattctt 360
tggtaaggga agcttttagac ctcaagtcta gcttggagac ttttgatttt gaagctttgt 420
attttgtatc ttggctaaag aatatatggt ggaaaaagtc ttcttgaaga gctcttaaaag 480
tgttgatttt gatgaaagtt cgttcaaaca taaattgttg atcttgaata tttttctat 540
ttatttgcac caaaaacgctt atgtttatta tgttccacta taattttatt ttttgaggac 600
gggaaaagga tcggagtttg ttaagcttga tcttgagaat atatgtcttt tgtatttgaa 660
gtctttctga tgggtattct agatgacatg tcttatgggt ggagtaatat cgtaagtatc 720
tagagtatgt gatttgtaat gatctctaaa aatactcgtg gaagtaatga cttttacttg 780
aagaaaagac tattatgtga aagagactta tacttgaaag agattattga aatacaagtg 840
tggagtaaaag ttttacttta aataaaaaaa aaaaagttaa atacaagtaa aaaaatactc 900
ataaatttaa ctttaaaaatt ttaaaattaag atgtaatgta atattcactt atatggttac 960
tcataattca tgaatataac tctcctcggg tacatagtcg ttaaatg 1007

```

SEQ ID NO: 25 moltype = DNA length = 993

FEATURE Location/Qualifiers

source 1..993

mol_type = genomic DNA

organism = Glycine max

SEQUENCE: 25

```

agccgataac ttatagataa cgttacagat aacattaact taaagataat tgtaccttgt 60
agataatgtg tagtctgtga gataattgaa tatatatcaa gagataaagg gatgacaaca 120
tattcaaaata aaaaaagtta gagataactt gtgggtttggg gagtgcactt gcgaagggtt 180
ggacgtctgt gctcctacac caggattgac atggaggatt gacgtgtgtc ttggagtgtc 240
acatgggatg atacatgtat ttgtgggatt atgaacaaca caattgctta aagttctact 300
caatttactt attacttcag gtgatgtctt ggtagttcac agatataagt ttttgtctgc 360
tatctttcat gtggacacac aagtatgtgt aatatagatg ttttttgaa agtttgagat 420
ccaggggcgc accaatgtat aggggagggg accttggcgg ttaaatcac cataaaaatt 480
taaaaatctt ttaaaaaaat ttaagccaaa caaattttga cttttttaca tcacctaaaa 540
atgaaccact agaaagtata atattgtcag atcctaattc tatttgggca aaaaaaaca 600
aaaaaaaaaaggaaagaaa aagtattaag aaaaagaaaa caaaaaataa aacaaaaaaa 660
caaaagaata aaaaacaaaa aagagagaaga aaacaaaaaa aaaaaaaaag aagaaaaaat 720
aaaaacaaaa agtattaaga aaaaaaaga acagtaaaaa aacaaaagaa agaaaatgta 780

```

-continued

```

aaaaaaagaa aaaaatagaa gaaaaaagga aaagttaaaa aaaaagattt tgtgacctat 840
tggcttctca aggagagccc attaggtcaa gaggagaaca ttgtataaaa aaataaagaa 900
ggaaagtctg tgcaattaag gcacatagga ggcaacatga atcccaagga gaacaatgga 960
ccaatctgtt ggcgtcattt gacattgtcg tga 993

SEQ ID NO: 26      moltype = DNA length = 1118
FEATURE          Location/Qualifiers
source           1..1118
                 mol_type = genomic DNA
                 organism = Glycine max

SEQUENCE: 26
gtgagaaaaag agagagggat cactaggttaa cccctcattt cctcactctc tcagtttccc 60
tctagtctttt tctcttttct ttcttttctt ccccttttct ttcttcttct tttatgtttt 120
caatctactg cttcgttatg tcccatctcg tcgggtgggca ccttggcgtg cggcgatgtt 180
tttgtgaatt gaattgggat ttcttttctg ttgggttttc acgcattcct tcactctcct 240
tgttcttctt ctttttttgt ttggccacc gtctgtgcct cttcgcatca tcgctatcgt 300
ggctgtgcca tcgtgttccc cgtggcggcc tcgcaccgtt ggatcctgga tcaatgggtg 360
cgaggacggg gcgccaccct ctgtgtctgt tcaccccttt atcgtgtcgt ttggaggcta 420
ggacatctag ggtttttcaa cctgtttgtc taattgcggg ttgggtcagg tcaccctgac 480
cgagttccaa cccacaaaaa aatggaaatt ttttttacta tttacaccac cttttcaaat 540
atgcaccatt ttctcatttt gggcttagcc cgtttttatg aagtatgaaa taaaaataaa 600
aacactattt gcaccaattt tttacacatc accttcttct atgttatgcc tagcccgttt 660
ttgtgaagtc taaagtaaaa taataaccgt tattacactt tttctttaa tacaagcacc 720
ttctctattt ttgggcataa catgtatttt ttagtctgaa agaaaaataa aagtgtctact 780
cacagctgct ttttcaacac atgcaccttc ttttgttttg ggcctagcgt gtgttttttt 840
tattattaag tccgaagtaa aataaaaactg atgattacac cacttttttt atatatgcac 900
ccctgaaact taggatgatg actaggtcca ccatgtctgc actccgttag tgttaattaa 960
gtcaaaagta atccttttga ctttgaaaaa aaatataaat attagtggat gaatctttat 1020
tttatttaaa ttcttttatt tttatatcat ttatttcatt cttcaatgtg attttatttt 1080
tattattgcc tagttagtta gtttaattaa taatgtat 1118

SEQ ID NO: 27      moltype = DNA length = 1002
FEATURE          Location/Qualifiers
source           1..1002
                 mol_type = genomic DNA
                 organism = Glycine max

SEQUENCE: 27
cctctcacga aacggaagcc tctatcagac ttagcttgcc tgataactct ttgtatatat 60
attctcaggg gttaaaaatta aaatagtgtt ttgatacttg ttttgtttct catgttggtt 120
tttatattga taggggaacc aatcttggtg aaaatgtggg agatatatgc catgggtagt 180
tttgtgaaat taagtttgctt acaatgtgtt taacgaaatg ccttgaatg cacttctctg 240
ttagtcattt aaattatgag ctatcatgaa ttttagttta aagttcattt aaattatgct 300
atttgttaca gactttaatt taaattatgc taagagtctt ttgttatgct ataattgggtg 360
tggtttctaa cacttatgtt ggaatagaag taatgtcaat tataattagc ttaatcagcc 420
caaaacattc actgtatttg ttgtgcaaat aagtgtgact aacctgggtt ttgtgaatga 480
tatacctttac ctctaactat gttaaaaatg aacctaaagt tagattattt aaatcctaag 540
ttctacatat taatacacta tgcaagttaa gcaatatagg tgcttatata tgttgcgtga 600
aacatgttat actctgtctc ttgtgataca gctgcattgt gattgacgca gatcttagtt 660
tttaatgacc tgtggaattg ttgtaggctt gtgtgtactc tttgtgaaac aattgatccc 720
atttcatttc aatttttgta cattttttta tgatttttat agtgtaacat gctctgctga 780
gtgatttact gtgtggctgc tgaataaaag caataaaatg ctatgttttg aatatcttga 840
cacttggttt gtatttttag tgaaaaatga cttaaaaagg gttctataga atttggaact 900
actcaaatgg cttttatctt caatttatac caatgtcacc ttttaaggcat gttaatcata 960
tatcttaaca agcgggtaac tatatcttga aatctgtctt gc 1002

SEQ ID NO: 28      moltype = DNA length = 1001
FEATURE          Location/Qualifiers
source           1..1001
                 mol_type = genomic DNA
                 organism = Glycine max

SEQUENCE: 28
ttgcgaaaac gcagataaaa tttgctttgt cttagaacaa ttcatgaaaa cagccccctag 60
tgagtcaaaag tatccccctt tgtttcaatt taaaaaatta atttctcact attcttattg 120
tgtaaatttt taaaggaact tgtgtaactt ttattcatat gttaatgtta atttgatata 180
taaatattaa tagaattata ttgttgatgt cataatgtta atcatgctaa tatagatgtt 240
ttaatcttaa tttatttatt aataaatggt aaatgttaat tattgttagc aaagacaaat 300
tcaaggaagg acaagaaagg atcttgact cccctcctaa ggatccctta tatacatgtg 360
aaaaaagaaa aaataataga agaaaatgaa ttaaaagaaat aagttgttga atttatgttt 420
gtttaatatt ttttattcta gtaatggatc tatcttaatt tttcatacaa atttttctca 480
caaaattaat aatgtttttt tttataaaaa cttattattt attaagggtt agatataaat 540
aatgacacaa aaaaagaaag aaaaatagtt ccccttaaaa atgttttttg atttgttctt 600
cattgttagt aagaggattt gaaccataaa tttttttcct ttttttatta ctaagttaat 660
cttataagtc ttagttaact tgaatgtcaa tcataccaat agagtattat gtgaatattt 720
ttcataaaat attaattatc aagtctatgg atgtagaaaa atagtttaat taaaaaaatg 780
acgataataa aatgtgtcaa ttatgtgttg atttactac acctcaacta ttaaaaaaaa 840
ataccacacc tcacattttt ttcgtttaat tgacatcaag aatgagaatg caaacaacaaa 900

```


-continued

```

atatgaatta gaccagaaaa caatccatcc atcgtatgcc atatagatca tctcataaac 960
cacctgtgta aggaaaattt ttattgtcaa ttgggcttag c 1001

```

```

SEQ ID NO: 29      moltype = DNA length = 1002
FEATURE           Location/Qualifiers
source            1..1002
                  mol_type = genomic DNA
                  organism = Glycine max

```

```

SEQUENCE: 29
acaaatacga atcatgtacc tgcattggaag aaataaccac acaacacaca atgatcagaa 60
taagcaaatg catataatta agcatgatac aatatcaata ttcattggaag taacaatgac 120
ttgtcaaaaa tttggatgaa attcaatatg taaatcaaaag ctttgtccct gaaaccctct 180
atataaatca aagctttgtc cctgaaaccc tctatgtaaa tgtgacagcc tcatgtctcc 240
cttcctgaaa acccactaaa aactgcctaa ccccccctgt gttactccat aatttattct 300
acaataactg cttaaggcag ttacatatgg tcctaaatca ctacacattc agttacgatt 360
aaccccttgt gcctaactac ggtttcgaaa catcacaca gagacagacc attgaacaat 420
ggattttcat cattaacata caacagagac ataccttcga tggaaagcgt gacacgaact 480
ccacaaacgc gaactcgaca atgtggttgc agttacagaa gcatagccca gtttgcgaca 540
aacacgaact caggcagaag gagaaacaac aataaagccc tgggttaaaa cgacgaacgc 600
ctaattgtta aacgacgaac gcctaattgt aaaaacgaag gacgtacctc aatggataag 660
tgccaaagac gatctccaca aacacaatct ccacaatgtg gttgcagtc cagaagcaca 720
agatcaagag atcaagagaa aagattcacg ttagtccatt atttgttaaa agaattctct 780
aatggttgtaa aagggtttggc cttaaaaata atttgttaaa cgattatcaa 840
agatctataa tcgattacta atgagaaaaa ttcaaaaaata acttgaaaaa gtcacatccc 900
tttatgagtt tttgaaaagc caccaaaagg ctatatatat gtgaattgtg ttcgaaaatc 960
tttagaattt tttcaaaact tctttgtctt attctctcat aa 1002

```

```

SEQ ID NO: 30      moltype = DNA length = 1000
FEATURE           Location/Qualifiers
source            1..1000
                  mol_type = genomic DNA
                  organism = Glycine max

```

```

SEQUENCE: 30
ataaataaac ttttggctcaa acacttgcaa atcaattaaa gattctttta agtttttcaa 60
tttgtattat acttctctag aagagagaaa aacttttgta cttcaaaaag aaaactatta 120
tttgtatcaa gaggtagtga gtctcttgat ttgtgagttt ttctgaacac aagagaaatg 180
tatccctagg tgggtcagaa gttgtaaagg aatttacaag aacaatagaa atctcaaatg 240
agttgcttga agattgaacg taaactgagt ttgcaattct ctctttcctt aattatctca 300
ttacataaat tgcaatttaa ttttgtcttg tgcattttaa gagtgtaaat taaattgttc 360
gttgtttctt attctgcata ttaagtttgc atatatcatt taaagagaga attaaaattt 420
gttaggggaa aatttttaaa cttaattcac ctgcctctta aattattgat gccacttggt 480
taaccatatt ttatcaattg aaataaatta attttttaat agaaatatga aaataattat 540
ataacaaaaa aaacccaaaa aaagaagaaa ataatacaata tattatcgac aactactatt 600
atcatattat taacataaaa aatatcaaca tattattaac aactatttta aattaatatg 660
aaaacaatta tataactaaa aaaataataa gaaagtaatg aaaaaaatca aaatattata 720
aactaatcca atattattaa actactattt tgactgatca acataagaga caacaaaaaa 780
tttcatatta ttaacaagag tgtggctcaa gttgtcaatc cataatttat agaataattt 840
attaaagcaa actacgaatg aatatggttt aagcagatag taacataagt ttacaaaact 900
taacaaaatg catgcatttt ctttactcta gaaatataaa agcctatttt aaaagacagt 960
aataaaaatta ctaagaacta caccattcaa aatagtgcgg 1000

```

```

SEQ ID NO: 31      moltype = DNA length = 1002
FEATURE           Location/Qualifiers
source            1..1002
                  mol_type = genomic DNA
                  organism = Glycine max

```

```

SEQUENCE: 31
agctttaatt ttgttcgcta catagtggcc acctattcta tccatgtcat aaccaataag 60
gtccaacata attgacagat aactgaattt tttaaatata ttaggagttc gattactgat 120
catgtgcata cagaagattt tgagaaagac aaaactcact ttagtgatat ctatgtttcg 180
aaagaaatta attttcgact acccattaga agatatcttt ggtacaaaca aaaaagaaga 240
aaaaaattct atcaatatca tagatattat actcaaatta taagactcat atattccacc 300
attcatccac ttgcttttgt gatgccctct gaaaagaga ttggttgcaa tctcttatgt 360
tgttctgatt cctactcgga agacgtcatg tgctcctcac gagactaaga aaaggtcatt 420
aagataaagg atctatactc taaattgcta accaatgttg tgggttgtaa attctgttct 480
tatataatgt tagatatttc tgaagagtca tgggtcagaa tcggggatcc atcacatgcc 540
cctatgtcaa taggttatat acaatgcttt tcgttgagga tgcttacaat attttacgta 600
agagcacaat taaaataaaa aagtaatttg ttagaagatg catagttaa gttaaagtat 660
gaggacagac aacatggata aatactcctt ctgtgtccgc acaaatgcag agcatcgaat 720
atagaacctc ggttgatatt actaatgtat gtgaacaca tactaagtaa taatagatta 780
gtctaagttt gattgatttt tttttattca taggaaatga aaataggtgc aggagattta 840
taataataa gttttatgtat tctacttaac catttaaat agatctcttg acaagttaca 900
atagttaaga gaacaaaact ctctcatgt tttttattt ttttatttac atacaagatt 960
cggacaagac aacttaaaaga aaaagccttg gaatggacat aa 1002

```

```

SEQ ID NO: 32      moltype = DNA length = 1002

```

-continued

```

FEATURE                               Location/Qualifiers
source                               1..1002
                                   mol_type = genomic DNA
                                   organism = Glycine max

SEQUENCE: 32
ataagagaac  tggttttttat  attattatta  ttctctttct  aattcataga  gaaccacagta  60
tggttgaccat  atcacccctga  aatgaaattt  aaatacccta  accaataatt  actaaaataa  120
ccttgaaatg  gctaaattgc  tagctactct  tttatatgta  actgaggtgt  tacttcatca  180
aaaggatata  tatgagatc  actcactaat  tactagtaca  ttgtaacatt  gtaacactat  240
ggatgtgttt  gaataagtat  ttttcatcta  gtttcacttg  acaatgggaa  cagcagcatt  300
gagttgtatt  gattggatga  gatcagagtc  tagcaagttg  agtgggtggc  gagagaaagg  360
gctctctgtg  gaactacctt  ttccggattt  ttgcattgta  agcaaacgag  caactgtgtt  420
tctgaaaatc  cctctcctg  ccacagttat  tgcactcttt  gcttctcttg  ctctctcagc  480
ttcttctct  cccgggaaca  ctgcacgat  tatgctctca  cactccttca  ccagctttga  540
gatcaaatct  gttgtgaaaa  atggctgctc  caaacctttc  tggatgaatg  gcaaacgcag  600
aagcccccct  gtctctctgt  cgtatttctt  cagaattttt  gccaacctgt  caaaattcag  660
tggtgtcact  tgttcaaggt  agcatcaaat  tctgtagaga  aagctgcac  aaagtctatg  720
gcaatgcatt  tcttagcctt  tttatcaata  gcaaacctat  ttttaacatt  ggagttaa  780
tttcatgcac  tgtaagagct  tttatgggtg  aaacacgaga  aattattaga  tggtttaaga  840
ctgttacatt  catgggtccc  attagtttct  acattatata  tataattga  tttttctgaa  900
aactcaatta  tgtctcgtgt  aaaaattatt  aatttgagac  tgtaattctta  tattatgtaa  960
taatttcacg  tcaatcaatt  agaattacc  ttagatataa  ag  1002

SEQ ID NO: 33      moltype = DNA  length = 1083
FEATURE            Location/Qualifiers
misc_feature       1..1083
note = synthetic construct; Plant optimized nucleotide
sequence of TS21 meganuclease
source             1..1083
mol_type = other DNA
organism = synthetic construct

SEQUENCE: 33
atggcaccga  agaagaagcg  caaggtgcat  atgaacacca  agtacaacaa  ggagtctctg  60
ctctacctgg  ccggtctcgt  ggaacggcga  ggctccatca  tggcgagat  caagccgcag  120
cagtcctgca  agttcaagca  cgcgtccag  ctgaccttca  ccgtgaccca  gaagacgcag  180
aggcgctggt  tcttcgacaa  gctggctgac  gagatcgagg  tgggcaagg  ctacgaccgc  240
gggtcggtgt  ccgactacat  cctctcccag  atcaagcccc  tgcacaactt  cctcaccag  300
ctccagccgt  tcctcaagct  gaagcagaag  caggcgaacc  tcgtcctgaa  gatcatcgag  360
cagctcccc  cggccaagga  gtccccggac  aagttcctgg  aggtgtgcac  gtgggtcgac  420
cagatcgagg  cctcaacga  cagcaagacc  cgcaagacga  cctcgagac  ggtgcgggag  480
gtcttgga  cctctccagg  atccgtggga  ggtctatcgc  catctcaggc  atccagcgcc  540
gcatcctcgg  cttctcaag  cccgggttca  gggatctccg  aagcactcag  agctggagca  600
actaagtcca  aggaattcct  gctctacctg  gccggcttcg  tggacggcga  cggctccatc  660
aaggcgagca  tcaagccgag  ccagtcctcc  aagttcaagc  acgagctctc  cctgaccttc  720
caggtgaccc  agaagacgca  gaggcgctgg  ttctctcgaca  agctggctga  cgagatcgag  780
gtgggctacg  tctacgacgc  cgggtcggtg  tccgactaca  tcctctccca  gatcaagccc  840
ctgcacaact  tcctcaccga  gctccagccg  ttcctcaagc  tgaagcagaa  gcaggcgaac  900
ctcgtcctga  agatcctcga  gcagctcccc  tcggccaagg  agtccccgga  caagtctctg  960
gaggtgtgca  cgtgggtcga  ccagatcgcg  gccctcaacg  acagcaagac  ccgcaagacg  1020
acctcgagga  cggtgcgggc  ggttctagac  tccctcagcg  agaagaagaa  gtctgcccc  1080
tga  1083

SEQ ID NO: 34      moltype = AA  length = 10
FEATURE            Location/Qualifiers
source             1..10
mol_type = protein
organism = Simian virus 40

SEQUENCE: 34
MAPKKRKRVH  10

SEQ ID NO: 35      moltype = DNA  length = 5937
FEATURE            Location/Qualifiers
misc_feature       1..5937
note = synthetic construct; Expression cassette RTW317,
comprising the TS21 meganuclease plant optimized
nucleotide sequence without an intron and operably linked
to the soybean EF1A promoter
source             1..5937
mol_type = other DNA
organism = synthetic construct

SEQUENCE: 35
cacatgggta  acctagactt  gtccatcttc  tggattggcc  aacttaatta  atgtatgaaa  60
taaaaggatg  cacacatagt  gacatgctaa  tcaactataat  gtgggcatca  aagttgtgtg  120
ttatgtgtaa  ttactagtta  tctgaataaa  agagaaagag  atcatccata  tttcttatcc  180
taaatgaatg  tcacgtgtct  ttataattct  ttgatgaacc  agatgcattt  cattaaccaa  240
atccatatac  atataaatat  taatcatata  taattaatat  caattggggt  agcaaaaaca  300

```

-continued

atctagtcta	ggtgtgtttt	gcgaatgcgg	ccgccaccgc	ggtggagctc	gaattctagt	360
ggccggccca	gctgatatcc	atcacactgg	cggccgcact	cgactgaatt	ggttccggcg	420
ccagcctgct	tttttgtaca	aagttggcat	tataaaaaag	cattgcttat	caatttggtg	480
caacgaacag	gtcactatct	gtcaaaataa	aatcattatt	tggggcccca	gcttaagtaa	540
ctaactaaca	ggaagagttt	gtagaaacgc	aaaaaggcca	tccgtcagga	tggccttctg	600
cttagtttga	tgcctggcag	tttatggcgg	gcgtcctgcc	cgccaccctc	cgggcggttg	660
cttcacaacg	tcaaatccg	ctcccggcgg	atttgtccta	ctcaggagag	cgttcaccga	720
caaaacaacg	ataaaaacga	aggcccagtc	ttccgactga	gcctttcgtt	ttatttgatg	780
cctggcagtt	ccctactctc	gcttagtagt	tagacgtccc	cgagatccat	gctagcggta	840
atacggttat	ccacagaatc	aggggataac	gcaggaaaga	acatgtgagc	aaaaggccag	900
caaaaggcca	ggaaccgtta	aaaggccgcg	ttgctggcgt	ttttccatag	gctccgcccc	960
cctgacgagc	atcacaaaaa	tcgacgtcca	agtcagaggt	ggcgaaaccc	gacaggacta	1020
taaagatacc	aggcgtttcc	ccctggaagc	tcctctgtgc	gctctcctgt	tccgaccctg	1080
ccgcttaccg	gataccctgc	cgcctttctc	ccttcgggaa	gcgtggcgct	ttctcatagc	1140
tcacgctgta	ggtagctcag	ttcgggtgtag	gtcgttcgct	ccaagctggg	ctgtgtgcac	1200
gaaccccccg	ttcagcccca	cgcgtgcgcc	ttatccggta	actatcgtct	tgagtcacga	1260
ccggttaagac	acgaacttatc	gccactggca	gcagccactg	gtaacaggat	tagcagagcg	1320
aggtagtagt	gcggtgtctac	agagttcttg	aagtggtggc	ctaactaccg	ctacactaga	1380
agaaacagat	ttggtatctg	cgctctgctg	aagccagtta	ccttcggaaa	aagagttggt	1440
agctcttgat	ccggcaaaac	aaccacgcgt	ggtagcgggt	gtttttttgt	ttgcaagcag	1500
cagattacgc	gcagaaaaaa	aggatctcaa	gaagatcctt	tgatcttttc	tacgggggtct	1560
gacgctcagt	ggaacggggc	ccaatctgaa	taatgttaca	accaattaac	caattctgat	1620
tagaaaaact	catacgagcat	caaatgaaac	tgcaatttat	tcatatcagg	attatcaata	1680
ccatattttt	gaaaaagccg	ttctgttaat	gaaggagaaa	actcaccgag	gcagttccat	1740
aggatggcaa	gatcctggta	tcggctcgcg	attccgactc	gtccaaacac	aatacaacct	1800
attaatcttc	cctcgtaaaa	aataagggtta	tcaagtgaga	aatcaccatg	agtgacgact	1860
gaatccggtg	agaatggcaa	aagtttatgc	atttctttcc	agacttggtc	aacaggccag	1920
ccattacgct	cgtcatcaaa	atcactcgca	tcaaccaaac	cgttattcat	tcgtgattgc	1980
gcoctgagca	gacgaaatac	gcgatcgctg	ttaaaaggac	aattacaaac	aggaatcgaa	2040
tgcaaccggc	gcaggaacac	tgccagcgca	tcaacaatat	ttcacctga	atcaggatat	2100
ttcttctaata	cttggaatgc	gttttttccg	gggatcgag	tggtagtaaa	ccatgcatca	2160
tcaggagtag	ggataaaatg	ttgatgggtc	ggaagaggca	taaattccgt	cagccagttt	2220
agttctgacca	tctcatctgt	aacatcattg	gcaacgctac	ctttgccatg	tttcagaaac	2280
aaactctggc	cctcgggctt	cccatacaag	cgatagattg	tcgcacctga	ttgcccgaca	2340
ttatcgcgag	cccatttata	cccataataa	tcagcatcca	tggttggaat	taatccgggc	2400
ctcgacgttt	cccggtgaat	atggctcata	acaccccttg	tattactggt	tatgtaagca	2460
gacagtttta	ttgttcatga	tgatatattt	ttatcttgtg	caatgtaaca	tcagagattt	2520
tgagacacgg	gccagagctg	cagctggatg	gcaataatg	attttatttt	gactgatagt	2580
gacctgttcg	ttgcaacaaa	ttgataagca	atgctttctt	ataatgccaa	ctttgtacaa	2640
gaaagctggg	ctagatatgc	tgaccocggg	caatcaaat	atatatgtaa	agcaattaca	2700
gtttatcaaa	ctttatttat	ggaaataaatt	tattatcaca	ttttatttgg	tttataaatt	2760
ttaaattaaa	ataatcccta	aataaaaaata	atttttaaca	tgacttattg	tcctaataaa	2820
attattttccg	taatttaaat	aaaatgaagt	ttttttcttt	caaagaatct	aaatggctcat	2880
aatgagaatt	ctctaaaaaa	atacataatg	agaataatta	tggaatttat	ttattataaa	2940
aaattaatag	catttttgata	gacaattaat	aaaatttttaa	aaataaccat	atagaataaa	3000
taattttttt	actatcggtt	ccaattaaaa	taatgataaa	aaataaaaata	gatttattaat	3060
tgatatgtat	atgaaattta	aataaagaat	ataatcatat	attttattga	tatatgatat	3120
gatataagatt	aattgatatt	gatttttgata	tggaaatttaa	aaataatata	ataattgttt	3180
ttattttatta	atacgtgttaa	tcaaaataatt	ctcacttttt	gaatcaatca	gtgtacttaa	3240
agataaatatc	agtgaaatat	tttttatcct	tttacgtgtg	ctgtgagaca	ttatcatcaa	3300
ttgtgttgta	tatgatatat	agatatagat	atataaatat	atagattgag	tgatataata	3360
tatttaaaat	ataaatatata	tatatgtttt	aatatatatt	tgcatatata	tatatatttg	3420
taaaaaactag	aagtattttt	tcatgagata	attattatcg	agttgaataa	gtctattatt	3480
tgtgagagcc	aacctataat	atatatgtga	ttaaatttta	tctttgtgaa	attaaaaata	3540
ataaaaaata	ccttaaaatc	ataataatag	aaaaacttat	attataaatt	taccattata	3600
cttaaaaaaa	attaaaaata	tattataaat	ataaatacta	tcgagtaatg	gccgcgctag	3660
ggtttttgag	aaaaaatctt	cccacgcact	caactgcact	gtacggcgctc	gttttcacag	3720
ccgcataata	gaagcccgct	tcccacaacc	ttcctcacaa	cattctcgga	ccctccagca	3780
ccgtcaccca	aacaaatctc	cacgcggtag	taggcgcgtg	aaacaaactc	taatccgaac	3840
tacgagacgt	gagaagcacg	cgccttagcg	agcgtttcaa	ttgtcgctac	gaaagcagag	3900
aaggatacaa	acggaactag	ggtaaattag	taagggtaat	ttcgttaaca	gaagaaaaga	3960
gtttagtagta	taaaataaac	ctctaaccct	cgtcgcatta	cttctcttca	cacctttggt	4020
cactcttctt	ctcttgcggc	taggggttta	gcgcagcttc	ttctagggtc	gttatctacc	4080
accgttctat	ggatttttat	ccttctattc	gtgtttattc	tattggttta	tgtgcttgcc	4140
aatatgtttt	ttctgaaatc	gtcgtcggtg	tcttcaattt	tatccatggt	tcagagatca	4200
attttggttg	tgtagtagt	gcttattctt	cttcttttcc	ttcgagttgt	taataacggt	4260
gctatgggtg	tttcaaaagt	gtttttttta	ttacttttga	tttaaaagtt	ttttggtaag	4320
gcttttattt	gcttggtata	ttcaaatctt	tggatccaga	tcttatataa	gtttttggtt	4380
caagaaagt	tttggttact	gatgaataga	tctattaact	gttactttta	tcgattccaag	4440
ctaaagtttt	ttggttactg	atgaatagat	ctattatctg	ttacttttaa	tcgggtccaag	4500
ctcaagtttt	ttggttactg	atgaatagat	ctatatacgt	cacagtgtgc	taaacatgcc	4560
cttggtttat	ctcgatctta	tgtatgggag	tgccataaat	tttgttatgt	ctattttttt	4620
atctgttgga	atcactctga	gtttgatgcg	ttacgattga	gcataacctat	ttttgggctt	4680
gttgtagtgt	gggtatttag	atcttaactc	ttttatgctt	atgaaagggt	ttgtaatgac	4740
aaaggctcta	atgttggttaa	acttttattt	ttactttata	tgggtgtgtg	atgtgttatg	4800
gttttgacaa	cttttttttt	ttctggattt	ttgcagattt	aaggaaagcca	tggcaccgaa	4860

-continued

```

gaagaagcgc aaggtgcata tgaacaccaa gtacaacaag gagttcctgc tctacctggc 4920
cggttcctg gacggcgacg gctccatcat ggccgagatc aagccgcagc agtcctgcaa 4980
gttcaagcac gcgctccagc tgaccttcac cgtgacccag aagacgcaga ggccgtgggt 5040
cctcgacaag ctggtgcagc agatcggggg gggcaagggtc tacgaccgcg ggtcggtgtc 5100
cgactacatc ctctccaga tcaagccctt gcacaacttc ctcacccagc tccagccgtt 5160
cctcaagctg aagcagaagc aggcgaacct cgtcctgaag atcatcgagc agctcccttc 5220
ggccaaggag tccccggaca agttcctgga ggtgtgcacg tgggtcgacc agatcgcggc 5280
cctcaacgac agcaagaccc gcaagacgac ctccgagacg gtgcgggcgg tcttggaactc 5340
cctcccagga tccgtgggag gtctatcgcc atctcaggca tccagcgcg catcctcggc 5400
ttctcaagc cccgggttcag ggaatctcca agcactcaga gctggagcaa ctaagtccaa 5460
ggaattcctg ctctacctgg ccggcttcgt ggacggcgac ggctccatca aggcgcagat 5520
caagccgcgc cagtcgccca agttcaagca cgagctctcc ctgaccttc aggtgaccca 5580
gaagacgcag aggcgctggt tctcgcacaa gctggtcgac gagatcgggg tgggtacagt 5640
ctacgacgcg ggtcggtgtt ccgactacat cctctccag atcaagcccc tgcacaactt 5700
cctcaccagc ctcagccggt tctcaagct gaagcagaag caggcgaacc tctcctgaa 5760
gatcatcgag cagctccctc cggccaagga gtccccggac aagttcctgg aggtgtgcac 5820
gtgggtcgac cagatcgcgg cctcaacga cagcaagacc cgcaagacga cctcggagac 5880
ggtgcgggcg gttctagact cctcagcga gaagaagaag tcgtccccc gaggtac 5937

```

```

SEQ ID NO: 36      moltype = DNA length = 6126
FEATURE           Location/Qualifiers
misc_feature       1..6126
                  note = synthetic construct; Expression cassette RTW322,
                  comprising the TS21 meganuclease plant optimized
                  nucleotide sequence ST-LS1 intron2 and operably linked to
                  the soybean EF1A promoter
source            1..6126
                  mol_type = other DNA
                  organism = synthetic construct

```

```

SEQUENCE: 36
cacatggtta acctagactt gtccatcttc tggattggcc aacttaatta atgtatgaaa 60
taaaaggatg cacacatagt gacatgctaa tcaactataat gtgggcatca aagttgtgtg 120
ttatgtgtta ttactagtta tctgaataaa agagaaagag atcatccata ttcttatcc 180
taaatgaatg tcacgtgtct ttataattct ttgatgaacc agatgcattt cattaacca 240
atccatatac atataaatat taatcatata taattaatat caattgggtt agcaaaaaaa 300
atctagtcta ggtgtgtttt gcgaatgcgg ccgccaccgc ggtggagctc gaattctagt 360
ggccggccca tttgtatctc atcacactgg ccggccgact cgactgaatt ggttcggcg 420
ccagcctgct tttttgtaca aagttggcat tataaaaaag cattgcttat caatttgttg 480
caacgaacag gtcactatca gtcaaaataa aatcattatt tggggcccca gcttaagtaa 540
ctaactaaca ggaagagtgt ttatgaacgc aaaaaggcca tccgtcagga tggccttctg 600
cttagtttga tgcctggcag ttatggcggt gcgtcctgcc cgccaccctc cgggcccgtg 660
cttcacacag ttcaaatccg ctcccgccgg atttgtccta ctacaggagag cgttcaccca 720
caaaacaacg ataaaacgaa agcccagtc ttccgactga gcctttcgtt ttatttgatg 780
cctggcagtt ccctactctc gcttagtagt tagacgtccc cgagatccat gctagcggta 840
atacggttat ccacagaatc aggggataac gcagaaaaga acatgtgagc aaaaggccag 900
caaaaggcca ggaaccgtta aaaggccgcg ttgctggcgt tttccatag gctccgccc 960
cctgacgagc atacaaaaaa tcgacgctca agtcagaggt ggccgaaacc gacaggacta 1020
taaagatacc aggcgtttcc ccttgaagc tccctcgtgc gctcctctgt tccgaccctg 1080
ccgcttaccg gatcctgtc cgcctttctc ccttcgggaa gcgtggcgct ttctcatagc 1140
tcacgctgta ggtatctcag ttccgtgtag gtcgttcgct ccaagctggg ctgtgtgcac 1200
gaaccccccg ttacgcccga ccgctcgccc ttatccgcta actatcgctc tagtcccaac 1260
ccggtaaagc acgactatc gccactggca gcagccactg gtaacaggat tagcagagcg 1320
aggtagtagc gcggtgtctc agagttcttg aagtggtggc ctaactacgg ctacactaga 1380
agaacagtat ttggtatctg cgctctgctg aagccagtta ccttcggaaa aagagttggt 1440
agctcttgat ccggcaaaaca aaccaccgct ggtagcgggt gtttttttgt ttgcaagcag 1500
cagattacgc gcagaaaaaa aggatctcaa gaagatcctt tgatcttttc tacgggggtc 1560
gacgctcagt ggaacggggc ccaatctgaa taatgttaca accaattaac caattctgat 1620
tagaaaaact catcgacat caaatgaaac tgcaatttat tcatatcagg attatcaata 1680
ccatattttt gaaaaagccg tttctgtaat gaaggagaaa actcaccgag gcagttccat 1740
aggatggcaa gatcctggta tcgggtctgc attccgactc gtccaacatc aatacaacct 1800
attaatttcc cctcgtaaaa aataaggtta tcaagtgaag aatcaccatg agtgacgact 1860
gaatccgggtg agaatggcaa aagtttatgc atttctttcc agacttggtc aacaggccag 1920
ccattacgct cgtcatcaaa atcactcgca tcaaccaaac cgttattcat tctgtattgc 1980
gcctgagcga gacgaaatc gcgatcgctg ttaaaaggac aattacaaac aggaatcgaa 2040
tgcaaccggc gcagggaacac tgccagcgca tcaacaatat ttccacctga atcaggatat 2100
tcttctaata cctggaatgc tgtttttccg gggatcgagc tgggtagtaa ccatgcatca 2160
tcaggagtag ggataaaatg cttgatggtc ggaagaggca taaattccgt cagccagttt 2220
agtcgacca tctcatctgt aacatcattg gcaacgctac ctttgccatg ttccagaaac 2280
aactctggcg catcgggctt cccatacaag cgatagattg tcgcacctga ttgcccgaca 2340
ttatcgcgag cccatttata cccatataaa tcagcatcca tgttggaatt taatcgcgcc 2400
ctcgacgttt ccggttgaa atggctcata acacccttg tattactgtt tatgtaagca 2460
gacagtttta ttgttcatga tgatatattt ttatcttgtg caatgtaaca tcagagattt 2520
tgagacacgg gccagagctg cagctggatg gcaataatg attttattt gactgatagt 2580
gacctgttcg ttgcaacaaa ttgataagca atgctttctt ataatgcaa ctttgtacaa 2640
gaaagctggg tctagatata tcgaccggcg caatcaaat atatatgtaa agcaattaca 2700
gtttatcaaa ctttatttat ggaataaatt tattatcaca tttatttttg tttataaatt 2760

```

ttaaattaaa	atatcaccta	aataaaaaa	atttttaaca	tgactttatt	tctaataata	2820
attatttcgc	taaatataat	aaaatgaagt	ttttttcttt	caaagaattc	aaatggtcat	2880
aatgagaatt	ctctaataaa	atacataagt	agaataatta	tggaatttat	tattaataaa	2940
aaattaatag	cattttgata	gacaattaat	aaaattttaa	aaataaacct	atagaataaa	3000
taattttttt	actatcggtt	coaatataaa	taatgatataa	aaaataaaaa	gattattaat	3060
tgatatgtat	atgaaattta	ataaagaat	ataatcatat	attttattga	tatatgatat	3120
gatatagatt	aaatgtatatt	gatttttgata	tggaatttaa	aaataatata	ataatgtgtt	3180
ttattttatta	ataactgttaa	tcaaatatatt	ctcacttttt	gaatcaatca	gtgtacttaa	3240
agataaatat	agttgaaatt	tttttatcct	tttacgtgtg	ctgtgagaca	ttatcatcaa	3300
ttgtgttgta	tatgatatat	agatatagat	atataaatat	atagattgag	tgatataata	3360
tatttaaaat	ataaattata	tatatgtttt	aaatattttt	tgcatatata	tatatatttg	3420
taaaaactag	aagtattttt	tcatgagata	attattattc	agttgaataa	gtctattatt	3480
tgtgagagcc	aaccatattt	atatatgtga	ttaaatttta	tcttttgtga	attaaaaata	3540
ataaaaaata	ccttaaaact	ataaatagat	aaaaaactat	attataattt	taccattata	3600
cttaaaaaaa	atttaataaa	tattataaat	ataaatacta	tcgagtaagt	gccgcgtag	3660
gggttttgag	aaaaaatctt	cccacgcact	caactgcact	gtacggcgct	gttttcacag	3720
ccgcataata	gaagcgcgct	tccccaaccc	tcttcacaaa	catctccgga	ccctccagca	3780
cctgcaccca	aaacaaatct	cacgcggtag	taggcgcgtg	aaacaaactc	taactccgaac	3840
tacgagacgt	gagaagcacg	cgtcttagcg	agcgtttcaa	ttgtcgctac	gaaagcagag	3900
aaggatacaa	acgggaactag	ggtaaaattag	taagggtaat	ttcgttaaca	gaagaaaaga	3960
gtgtgtagcta	taaaataaac	ctctaaccct	cgtcgcattc	cttcttccca	caccttttgt	4020
cactcttctt	ctcttgccgc	taggggttta	gcgcagcttc	ttctaggttc	gttattctac	4080
accggttctat	ggatttttatt	ccctcttatto	gtgtttattc	tatttggttta	tgttgcttgc	4140
aatatgtttt	ttctgaaatc	ctcgtcggtg	tcttcaattt	tctcatggtt	tcagagatca	4200
attttgtttg	tgtagtatgt	gcttattctt	cttcttttcg	ttcagagttg	taataacggt	4260
gctatgtgtg	tttcaaaagt	gtttttttta	ttacttttga	tttaaaagtt	ttttggttaag	4320
gcttttttgt	gctgttgtata	tcaaatctt	tggaatcaga	tcttataata	gttttatggt	4380
caagaaagtt	tttggttact	gatgaataga	tctattaact	gttactttaa	tcgatccaag	4440
ctaaagtttt	ttgggtactg	atgaatatag	ctattatctg	ttacttttaa	tcgggtcaag	4500
cctaagtttt	ttggttactg	atgaatatag	ctataactg	cacagtggtc	taaacatgcc	4560
cttgttttat	ctcgatctta	tgtatgggag	tgccataaat	tttgttatgt	ctattttttt	4620
atctgttgga	atcatactga	gttttgatcg	ttacgatgtg	gcataacctat	ttttgggctt	4680
gtgtgatggt	gcttattttag	attctaactc	ttttatgctt	atgaagagct	ttgtaatgac	4740
aaagggtcta	atgttgttaa	acttttattt	ttactttata	tggtgtgttg	atgtgttatg	4800
gttttgacia	cttttttgtt	tcttggtatt	ttgcagattt	aaggaaagcca	tgggcacggaa	4860
gaagaaagcg	aaaggtgcata	gtacacccaa	gtacacaaag	gagttctctg	tctacctggc	4920
cggtctcggt	gacggcgacg	gctccatcat	ggcgcagatc	aagccgcagc	agtcctgcaa	4980
tttcaagcac	gcgctccagc	tgaccttcac	cgtgaccagc	atagcagca	ggcgctgggt	5040
cctcgacaag	ctggctgacg	agatcggggt	gggcaaggtc	aacgcgcgag	ggtcggtgtc	5100
cgactacatc	ctctcccaga	tcaagccctc	gcacaacttc	ctcaccacgc	tcagcccggt	5160
cctcaagctg	aagcagaagc	aggcgaaact	cgtcctgaag	atcatcgagc	agctcccttc	5220
ggccaagggag	tcccgggaca	agttcctgga	ggtaagtttc	gtcttctacc	tttgatatat	5280
atataataat	tatcataat	tagtagtaat	ataatttttc	aaatattttt	tcaaaaaata	5340
aaagaatgtg	tatatagcaa	ttgctttttc	gtagtttata	agtggtgata	ttttaattta	5400
tactttttct	aatatagcac	caaaaactag	tgatgtcgag	gtgtgcacgt	gggtgcacca	5460
gatcgcggcg	ctcaacgaca	gcaagaccgc	caagacgacc	tcggagacgg	tgccggcggt	5520
ctctggactc	ctcccaggat	cgggtggagg	tctatcgcca	tctcaggcat	ccagcgccgc	5580
acccctcggt	tctccaagcc	ccgttccagg	gatctccgaa	gcactcagat	ctggagcaac	5640
taagtccaag	gaattcctgc	tctacctggc	cggtctcggt	gacggcgacg	gctccatcaa	5700
ggcgccagat	aagccgcgcc	agtcctcgcaa	gttcaagcac	gagctctccc	tgacctttca	5760
gggcaccagc	aaagcgcgca	ggctctgggt	ctcgcagacg	agatcgcgag	agatcggtgt	5820
gggtcactgc	tacgaccgcg	ggtcgggtgc	cgactacatc	ctctccacga	tcaagccctc	5880
gcacaaactc	ctcaccagca	ttccagccgt	cctcaagctg	aagcagaagc	aggcgaaact	5940
cgctcctgaag	atcatcgacg	agctcccttc	ggccaaggag	tccccggaca	agttcctgga	6000
ggtgtgcacg	tgggtcgacc	agatcgcggc	cctcaacgac	agcaagacc	gcaagacgac	6060
ctcggagacg	gtgcggggcg	ttctagactc	cctcagcgag	aagaagaagt	cgtccccctg	6120
agggtac						6126

SEQ ID NO: 37

FEATURE

misc_feature

source

-continued

ctatccaact	gctattccca	tttctaacc	ttgttacacg	agcatcttga	tcaatggctc	600
agaaagggaa	atagcagttg	agtgggtgctt	caacgataag	ttattggatt	tagtattttat	660
cttagcctgt	tttcgtgtac	tttggtttgc	cggatggagg	tatgtgattt	tgtctatgat	720
tcttaataca	ataacctaca	cttactctca	ttgatagttt	gtgcagatct	aatagctatg	780
aagcaccgat	acgggacatg	acacgggtcag	gtggacacat	gtaatgtcta	aaatattaaa	840
atatagaacg	tagtacagag	gtcgtgtcgg	gtttagatag	tgatagggac	gcgtgtcggg	900
caccggacat	gacaaaggac	tgaagtgcct	agaattgttt	atgtttgaga	tcttgttgat	960
gagaggcgga	tagagggtcaa	cttgccaaga	taacctacag	ttctatatta	gatgctttgt	1020
gcaaaaacga	tcattccaaag	gctattggat	tattcaagaa	aactaaagac	caaggagttc	1080
aaaaccgcct	atgtacacat	gcactatact	tatggatgga	ttgtgcgaag	tggaagactt	1140
cagaatgcga	aaatgatgtt	tcaggatcta	ctgattaaag	gctatcaact	aagtgtctgt	1200
ctgtactctg	tataatgtta	tgattcatag	gctttgtaaa	gagggatgtt	ttgatgaagc	1260
attgatctag	aaactcaaaa	tggaaaacag	atcttaaaag	agatacactg	tgtaaatgtg	1320
taatggcact	ggcactctcg	tgtactagt	gtcacctaag	tgactagggt	cacgtgaccc	1380
tagtcactta	ttcccaacag	aagttcctat	tccgaagttc	ctattctcta	gaaagatag	1440
gaacttccac	tagtaccaca	caagcttgca	tgccctgcag	tttaaacagt	cgactctaga	1500
gatccgtcaa	catgggtggag	cacgacactc	tcgtctactc	caagaatatt	aaagatacac	1560
tctcagaaga	ccaagggtct	attgagactt	ttcaacaaag	ggtaataatc	ggaacacctc	1620
tcggattcca	ttgcccagct	atctgtcact	tcatacaaa	gacagtagaa	aaggaagggtg	1680
gcacctacaa	atgccactcat	tgcgataaa	gaaaggctat	cggtcaagat	gcctctgcgc	1740
acagtggtcc	caaagatgga	ccccaccaca	caggagagcat	cggtgaaaaa	gaagacgttc	1800
caaccacgtc	ttcaaaagca	gtggattgat	gtgatgatcc	tatgcgtatg	gtatgacgtg	1860
tgttcaagat	gatgacttca	aacctacctc	tgacgtatgg	tatgacgtgt	gtcgactgat	1920
gacttagatc	cactcgagcg	gctataaata	cgtaacctac	cacctgtcgc	tacctccctc	1980
agagctgcag	cttattttta	caacaattac	caacaacaac	aaacaacaaa	caacattaca	2040
attactattt	acaattacag	tcgacctcta	gtccatgaaa	aagcctgaac	tcaccgcgac	2100
gtctgtcgag	aagtttctga	tcgaaaagtt	cgacagcgct	tccgacctga	tcgagctctc	2160
ggaggggcga	gaatctcgtg	cttccagctt	cgatgtagga	gggctgtggat	atgtctgcgc	2220
ggtaaatagc	tgcgccgatg	gtttctacaa	agatcggtat	gtttatcggc	actttgcac	2280
ggccgcgctc	ccgattccgg	aagtgtctga	cattggggaa	ttcagcgaga	gcctgacctc	2340
ttgcactctc	cgccgtgcac	agggtgtcac	gttgcaagac	ctgcctgaaa	ccgaactgcc	2400
cgtgttctg	cacccggtcg	cggaggccat	ggatgcgac	gctgcggccg	atcttagcca	2460
gacgagcggg	ttcgggccat	tcgggaccga	aggaatcggt	caatacacta	catggcggtg	2520
tttcatatgc	gcgattctgt	atcccatgt	gtatcactgg	caaatctgtg	tggacgacac	2580
cgtagctg	tcctgtcgcg	aggctctcga	tgagctgatg	ctttgggccc	aggactgccc	2640
cgaagtccgg	cacctcgtgc	acgcggattt	cggctccaac	aatgtcctga	cggacaatgg	2700
ccgcataaca	gcggtcattg	actggagcga	ggcgatgttc	ggggattccc	aatacagagt	2760
cgccaaacac	ttcttctgga	ggccgtgggt	ggcttgtatg	gagcagcaga	cgcgctactt	2820
cgagcggagg	catccggagc	ttgcaggatc	gcgcgggctc	cgggcgatga	tgctccgcat	2880
tggtcttgac	caactcttat	agagcttggt	tgacggcaat	ttcgatgatg	cagcttgggc	2940
gcagggtcga	tgcgacgcaa	tcgtccgac	cggagccggg	actgtcgggc	gtacacaaat	3000
cgcccgacga	agccgcccgc	tctggaccga	tggctgtgta	gaagtactcg	ccgatagtgg	3060
aaaccgacgc	cccagcactc	gtccgagggc	aaaggaaatg	tgaggtaacct	aaagaaggag	3120
tgctgcgaag	cagatcggtc	aaacattttg	caataaaagt	ttctaaagatt	gaatcctggt	3180
gcgggtcttg	cgatgattat	catataattt	ctgttggaatt	acgttaagca	tgtaataatt	3240
aaacatgaa	gcattgacgt	atttatgaga	tggtgtttta	tgattagagt	cccgaattta	3300
tacattta	acgcgataga	aaacaaaata	tagcgcgcaa	actaggataa	attatcgccg	3360
gcgggtgcat	ctatgttact	agatcgatgt	cgacccggga	tcatggctag	cgaagtccct	3420
attccgaagt	tcctattctc	tagaaagtat	aggaacttca	gatcccttag	agtcgacctg	3480
caggcatgcc	cgccgatatc	gatgggcccc	ggccgaagct	tcaagtttgt	acaaaaaagc	3540
aggctccggc	cagaatccgg	taagtgaact	gggtcacgtg	accctagtca	cttaaatctg	3600
gccagaatgg	ccatctggat	tcagcaggcc	tagaaggccc	ggaccgatta	aactttaatt	3660
cggtccggaa	gcttggatcc	gtcgacgaat	tcactagtgt	taccagagct	gggtcacctaa	3720
gtgactaggg	tcacgtgacc	ctagtcaact	attcccgggc	accagacttt	cttgtacaaa	3780
ctggccggtta	acggatccgg	cagaatccgg	taagtgaact	gggtcacgtg	accctagtca	3840
cttaaatctg	gccagaatgg	ccatctggat	tcagcaggcc	tagaaggccc	ggaccgatta	3900
aactttaatt	cggtccgggt	tacctctaga	aagcttgtcg	accctgcagg	gtgtgattaa	3960
aagtcatata	tggttttaaga	tacttttttt	tataaagata	gtagtgttca	atttttcgat	4020
attacacaa	tggtttcttt	tcttctcatt	gtactgtaga	tctgatttac	tttcaatgat	4080
tggttaagtc	actgggtgta	ttgtttgtgt	ttcaaatatc	aaaccaagct	gaaactgaga	4140
tgatgatgat	ttgaaatgct	ttatctcatg	tagtcgactc	aattttctcg	tatatctctt	4200
gtttctttta	aagaacagg	agcttttaag	attttaaaca	ccagcatatt	ttgtttgcat	4260
aatccaaatt	gtcttaggtg	taagttgtct	gacattttcc	ttgatgtcat	tgctgcataa	4320
ttaattggag	ccttttcaaa	acctatgggt	tattttgttg	gggattatct	aagggaacgcg	4380
tgctcagtc	tcaagtgtta	tgattgctga	tatcagtgat	atattgctgc	acaatgaagt	4440
ggaactattt	taaatttcaa	ttgatgattc	tgcatccaat	ttatcatctg	acccttttat	4500
ctttttacct	atctggcatt	ttagtctttt	accagataaa	aggaccaaac	acatgagata	4560
taatacccaa	atgaaaagaa	tgaagaagca	gatataaaga	tggtgttttt	ctttttattc	4620
ctggaagatt	tagatgatgt	tttcaattaa	gttgtttgtg	gatgctttta	gatgattttg	4680
ttttgcatac	atatgtttac	ttttttgttc	tcaacttctc	attcattttc	catgatttca	4740
ttccgtgaaa	aagtgtttta	gcagaaaacg	tttttccctc	gttgcttttg	tcctaaactt	4800
ttggatttcta	agttttttta	tatgaaaatt	agatcatttg	gcacatgggt	ttccaaagac	4860
acaagtagac	tctttctatg	aaatcaatct	taaatccctt	tttagaggaa	aacattttta	4920
aggaggtgaa	catgttgttg	agtggaagg	atcccggtc	ctaagtgaat	agggtcacgt	4980
gacctagtc	acttatccc	gggcaacttt	attatacaaa	gttgatagat	ctcgaattca	5040
ttccgattaa	tcgtgg					5056

SEQ ID NO: 38	moltype = DNA	length = 24
FEATURE	Location/Qualifiers	
misc_feature	1..24	
source	note = synthetic construct; Primer Mega21-190F 1..24 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 38		
ggcactctcg tgtgtgatta aaag		24
SEQ ID NO: 39	moltype = DNA	length = 27
FEATURE	Location/Qualifiers	
misc_feature	1..27	
source	note = synthetic construct; Primer Mega21-301R 1..27 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 39		
caatgagaag aaaaagaaac acttgtg		27
SEQ ID NO: 40	moltype = DNA	length = 16
FEATURE	Location/Qualifiers	
misc_feature	1..16	
source	note = synthetic construct; Probe mega21-250T 1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 40		
agtagtggtc aatttt		16
SEQ ID NO: 41	moltype = DNA	length = 23
FEATURE	Location/Qualifiers	
misc_feature	1..23	
source	note = synthetic construct; Primer Mega14-13F 1..23 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 41		
aacacatgat ggacgacttc aaa		23
SEQ ID NO: 42	moltype = DNA	length = 23
FEATURE	Location/Qualifiers	
misc_feature	1..23	
source	note = synthetic construct; Primer Mega14-128R 1..23 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 42		
caagcagacg tacgcaagta gct		23
SEQ ID NO: 43	moltype = DNA	length = 19
FEATURE	Location/Qualifiers	
misc_feature	1..19	
source	note = synthetic construct; Probe Mega14-85T 1..19 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 43		
ttgtcaatac gaaagtaac		19
SEQ ID NO: 44	moltype = DNA	length = 21
FEATURE	Location/Qualifiers	
misc_feature	1..21	
source	note = synthetic construct; Primer Mega30-30F 1..21 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 44		
tgccatgagt agcaccactt g		21
SEQ ID NO: 45	moltype = DNA	length = 23
FEATURE	Location/Qualifiers	
misc_feature	1..23	
source	note = synthetic construct; Primer Mega30-87R 1..23	

-continued

	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 45		
ctcagattta tgctcttgcg tgg		23
SEQ ID NO: 46	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = synthetic construct; Probe Mega30-52T	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 46		
tgggttcgac acatct		16
SEQ ID NO: 47	moltype = DNA length = 28	
FEATURE	Location/Qualifiers	
misc_feature	1..28	
	note = synthetic construct; Primer Mega5-F1	
source	1..28	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 47		
tcgtaaccat tcatgtgata taatgac		28
SEQ ID NO: 48	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
misc_feature	1..22	
	note = synthetic construct; Primer Mega5-R1	
source	1..22	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 48		
tgcttacgtg tgtactcgtg ca		22
SEQ ID NO: 49	moltype = DNA length = 18	
FEATURE	Location/Qualifiers	
misc_feature	1..18	
	note = synthetic construct; Probe Mega5-T1	
source	1..18	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 49		
ttctcacact cacctaag		18
SEQ ID NO: 50	moltype = DNA length = 32	
FEATURE	Location/Qualifiers	
misc_feature	1..32	
	note = synthetic construct; Primer WOL133	
source	1..32	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 50		
gttgatgaga ggcagataga ggtcaacttg cc		32
SEQ ID NO: 51	moltype = DNA length = 33	
FEATURE	Location/Qualifiers	
misc_feature	1..33	
	note = synthetic construct; Primer WOL134	
source	1..33	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 51		
ttatgcagca atgacatcaa gggtaatgtc agc		33
SEQ ID NO: 52	moltype = DNA length = 27	
FEATURE	Location/Qualifiers	
misc_feature	1..27	
	note = synthetic construct; Primer WOL190	
source	1..27	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 52		
ctaagacac gtgtatcaag taactgg		27
SEQ ID NO: 53	moltype = DNA length = 27	

-continued

FEATURE	Location/Qualifiers	
misc_feature	1..27	
	note = synthetic construct; Primer WOL242	
source	1..27	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 53		
tcgaactttt cgatcagaaa cttctcg		27
SEQ ID NO: 54	moltype = DNA length = 34	
FEATURE	Location/Qualifiers	
misc_feature	1..34	
	note = synthetic construct; Primer WOL153	
source	1..34	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 54		
gattagagtc ccgcaattat acatttaata cgcg		34
SEQ ID NO: 55	moltype = DNA length = 27	
FEATURE	Location/Qualifiers	
misc_feature	1..27	
	note = synthetic construct; Primer WOL247	
source	1..27	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 55		
aactgagaga ctgagcgaca atcacag		27
SEQ ID NO: 56	moltype = DNA length = 31	
FEATURE	Location/Qualifiers	
misc_feature	1..31	
	note = synthetic construct; Primer WOL121	
source	1..31	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 56		
gctaattggat tcaatttgaa gtatttaata g		31
SEQ ID NO: 57	moltype = DNA length = 34	
FEATURE	Location/Qualifiers	
misc_feature	1..34	
	note = synthetic construct; Primer WOL150	
source	1..34	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 57		
actttagaat gataatgatg actttagcac tgcc		34
SEQ ID NO: 58	moltype = DNA length = 27	
FEATURE	Location/Qualifiers	
misc_feature	1..27	
	note = synthetic construct; Primer WOL192	
source	1..27	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 58		
gtacgcaaac agcttggtta cctttcg		27
SEQ ID NO: 59	moltype = DNA length = 28	
FEATURE	Location/Qualifiers	
misc_feature	1..28	
	note = synthetic construct; Primer WOL193	
source	1..28	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 59		
ttccaatttg agagggtata tttccttc		28
SEQ ID NO: 60	moltype = DNA length = 30	
FEATURE	Location/Qualifiers	
misc_feature	1..30	
	note = synthetic construct; Primer WOL113	
source	1..30	
	mol_type = other DNA	
	organism = synthetic construct	

-continued

SEQUENCE: 60
 ggaagccttg ttcgtatcga aacacaaagg 30

SEQ ID NO: 61 moltype = DNA length = 29
 FEATURE Location/Qualifiers
 misc_feature 1..29
 note = synthetic construct; Primer WOL114
 source 1..29
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 61
 ccacatcttt taactcaagg ggcttcagc 29

SEQ ID NO: 62 moltype = DNA length = 28
 FEATURE Location/Qualifiers
 misc_feature 1..28
 note = synthetic construct; Primer WOL194
 source 1..28
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 62
 ccaagtcaat aactttctga tgagaagc 28

SEQ ID NO: 63 moltype = DNA length = 29
 FEATURE Location/Qualifiers
 misc_feature 1..29
 note = synthetic construct; Primer WOL195
 source 1..29
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 63
 ggtaggcaa attagatagt gtttgattt 29

SEQ ID NO: 64 moltype = DNA length = 29
 FEATURE Location/Qualifiers
 misc_feature 1..29
 note = synthetic construct; Primer WOL105
 source 1..29
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 64
 ctcggttaa gatcacatga tagcaaagc 29

SEQ ID NO: 65 moltype = DNA length = 31
 FEATURE Location/Qualifiers
 misc_feature 1..31
 note = synthetic construct; Primer WOL144
 source 1..31
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 65
 ctaaccaac tccgatcctt ttcccgctcct c 31

SEQ ID NO: 66 moltype = DNA length = 27
 FEATURE Location/Qualifiers
 misc_feature 1..27
 note = synthetic construct; Primer WOL196
 source 1..27
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 66
 attctatgaa aaggatgtct tgtggcg 27

SEQ ID NO: 67 moltype = DNA length = 25
 FEATURE Location/Qualifiers
 misc_feature 1..25
 note = synthetic construct; Primer WOL197
 source 1..25
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 67
 acaccaagcc caatcgccat acatc 25

SEQ ID NO: 68 moltype = DNA length = 22
 FEATURE Location/Qualifiers
 source 1..22

-continued

	mol_type = genomic DNA	
	organism = Zea mays	
SEQUENCE: 68		
tggttgact tgcgagataa ac		22
SEQ ID NO: 69	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = genomic DNA	
	organism = Zea mays	
SEQUENCE: 69		
caaacagatt cacgtcagat tt		22
SEQ ID NO: 70	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = genomic DNA	
	organism = Zea mays	
SEQUENCE: 70		
ttacatgacg taggacatta cg		22
SEQ ID NO: 71	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = genomic DNA	
	organism = Zea mays	
SEQUENCE: 71		
gtttctcacg cgtgagagcc tt		22
SEQ ID NO: 72	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = genomic DNA	
	organism = Zea mays	
SEQUENCE: 72		
ccaaccgtcg tgagacctgc cc		22
SEQ ID NO: 73	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = genomic DNA	
	organism = Zea mays	
SEQUENCE: 73		
agatcggacg caagagggtt ta		22
SEQ ID NO: 74	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = genomic DNA	
	organism = Zea mays	
SEQUENCE: 74		
gggcggtatg tatgtcatat ta		22
SEQ ID NO: 75	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = genomic DNA	
	organism = Zea mays	
SEQUENCE: 75		
caagctctcg cgaaaagggc ag		22
SEQ ID NO: 76	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = genomic DNA	
	organism = Zea mays	
SEQUENCE: 76		
ctagtatacg tgagagacct tg		22
SEQ ID NO: 77	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = genomic DNA	
	organism = Zea mays	
SEQUENCE: 77		
aagaaataca tgcgagccag tc		22

-continued

```

SEQ ID NO: 78      moltype = DNA length = 1083
FEATURE           Location/Qualifiers
misc_feature       1..1083
                   note = synthetic construct; Plant optimized nucleotide
                   sequence of MHP14 containing a nuclear localization signal
                   and no intron
source             1..1083
                   mol_type = other DNA
                   organism = synthetic construct

```

```

SEQUENCE: 78
atggcaccga agaagaagcg caaggtgcat atgaacacca agtacaacaa ggagttcctg 60
ctctacctgg ccggcttcgt ggacggcgac ggctccatca tcgcgcagat caagccgaac 120
cagtcctaca agttcaagca ccagctcatg ctgaccttca ccgtgaccca gaagacgcag 180
agggcgtggt tcctcgacaa gctggtcgac gagatcgggg tgggctacgt ccgcgaccgc 240
gggtcgggtg ccgactacat cctctcccag atcaagcccc tgcacaactt cctcaccag 300
ctccagccgt tcctcaagct gaagcagaag caggcgaacc tcgtcctgaa gatcatcgag 360
cagctccccc cggccaagga gtccccggac aagttcctgg aggtgtgcac gtgggtcgac 420
cagatcgcgg ccctcaacga cagcaagacc cgcaagacga cctcggagac ggtgcgggcg 480
gtcctggact cctctccagg atccgtggga ggtctatcgc catctcaggg atccagcgcc 540
gcatcctcgg ctctctcaag cccgggttca gggatctccg aagcactcag agctggagca 600
actaagtcca aggaattcct gctctacctg gccggcttcg tggacggcga cggtccatc 660
atcgcgcgga tcaagccgaa ccagtcctac aagttcaagc accagctctc cctgaccttc 720
accgtgaccc agaagacgca gaggcgctgg ttctcgcaca agctggtcga cgagatcggg 780
gtgggctacg tccgcgacca ggggtcgggt tcccactacc agctctccca gatcaagccc 840
ctgcacaaat tcctcaccga gctccagccg ttctcgaagc tgaagcagaa gcaggcgaac 900
ctcgtcctga agatcatcga gcagctcccc tcggccaagg agtccccgga caagttcctg 960
gaggtgtgca cgtgggtcga ccagatcgcg gccctcaacg acagcaagac ccgcaagacg 1020
acctcgga ga cggtgcgggc ggttctagac tccctcagcg agaagaagaa gtctgcccc 1080
tga 1083

```

```

SEQ ID NO: 79      moltype = DNA length = 1083
FEATURE           Location/Qualifiers
misc_feature       1..1083
                   note = synthetic construct; Plant plant optimized
                   nucleotide sequence of MHP14+ containing a nuclear
                   localization signal and no intron
source             1..1083
                   mol_type = other DNA
                   organism = synthetic construct

```

```

SEQUENCE: 79
atggcaccga agaagaagcg caaggtgcat atgaacacca agtacaacaa ggagttcctg 60
ctctacctgg ccggcttcgt ggacggcgac ggctccatca tcgcgcagat caagccgaac 120
cagtcctaca agttcaagca ccagctcatg ctgaccttca ccgtgaccca gaagacgcag 180
agggcgtggt tcctcgacaa gctggtcgac gagatcgggg tgggcaagggt ccgcgaccgc 240
gggtcgggtg ccgactacat cctctcccag atcaagcccc tgcacaactt cctcaccag 300
ctccagccgt tcctcaagct gaagcagaag caggcgaacc tcgtcctgaa gatcatcgag 360
cagctccccc cggccaagga gtccccggac aagttcctgg aggtgtgcac gtgggtcgac 420
cagatcgcgg ccctcaacga cagcaagacc cgcaagacga cctcggagac ggtgcgggcg 480
gtcctggact cctctccagg atccgtggga ggtctatcgc catctcaggg atccagcgcc 540
gcatcctcgg ctctctcaag cccgggttca gggatctccg aagcactcag agctggagca 600
actaagtcca aggaattcct gctctacctg gccggcttcg tggacggcga cggtccatc 660
atcgcgcgga tcaagccgaa ccagtcctac aagttcaagc accagctctc cctgaccttc 720
accgtgaccc agaagacgca gaggcgctgg ttctcgcaca agctggtcga cgagatcggg 780
gtgggctacg tccgcgacca ggggtcgggt tcccactacc agctctccca gatcaagccc 840
ctgcacaaat tcctcaccga gctccagccg ttctcgaagc tgaagcagaa gcaggcgaac 900
ctcgtcctga agatcatcga gcagctcccc tcggccaagg agtccccgga caagttcctg 960
gaggtgtgca cgtgggtcga ccagatcgcg gccctcaacg acagcaagac ccgcaagacg 1020
acctcgga ga cggtgcgggc ggttctagac tccctcagcg agaagaagaa gtctgcccc 1080
tga 1083

```

```

SEQ ID NO: 80      moltype = DNA length = 1273
FEATURE           Location/Qualifiers
misc_feature       1..1273
                   note = synthetic construct; Plant optimized nucleotide
                   sequence of MHP55 containing a nuclear localization signal
                   and an intron
source             1..1273
                   mol_type = other DNA
                   organism = synthetic construct

```

```

SEQUENCE: 80
atggcaccga agaagaagcg caaggtgcat atgaacacca agtacaacaa ggagttcctg 60
ctctacctgg ccggcttcgt ggacggcgac ggatccatca tcgcgcagat caagccgaac 120
cagtcctgca agttcaagca ccagctctcc ctgaccttcc aggtgaccca gaagacgcag 180
agggcgtggt tcctcgacaa gctggtcgac gagatcgggg tgggctacgt ccgcgaccgc 240
gggtcgggtg ccgactacat cctctcccag atcaagcccc tgcacaactt cctcaccag 300

```

-continued

```

ctgcagccgt tctccaagct gaagcagaag caggcgaacc tcgtcctgaa gatcatcgag 360
cagctccctt cggccaagga gtcccgggac aagttcctgg aggttaagttt ctgcttttac 420
ctttgatata tatataataa ttatcattaa ttagtagtaa tataatattt caaatatttt 480
tttcaaaata aaagaatgta gtatatagca attgcttttc ttagtattat aagtgtgtat 540
attttaattt ataacttttc taatatatga ccaaaacatg gtgatgtgca ggtgtgcacg 600
tgggtcgacc agatcgcggc cctcaacgac agcaagaccc gcaagacgac ctcgagacg 660
gtgcggggcg tctcgactc cctcccagga tccgtgggag gtctatcgcc atctcaggca 720
tccagcgccg catcctcggc ttctcaagc cggggttcag ggatctcga agcactcaga 780
gctggagcaa ctaagtccaa ggaattcctg ctctacctgg ccggttcctg ggacggcgac 840
ggctccatca tcgctccat caagccggag cagtcgccga agttcaagca ccgctcagc 900
ctgaccttcc aggtgaccca gaagacgac aggcgtcgtt tctcgacaa gctggtcgac 960
gagatcgggg tgggtactgt ccgacaccgc gggtcgggtg ccgactaccg cctctcccag 1020
atcaagcccc tgcacaactt cctcaccag ctccagccgt tctcaagct taagcagaag 1080
caggcgaaac tcgtcctgaa gatcatcgag cagctccctt cggccaagga gtcccgggac 1140
aagttcctgg aggtgtgcac gtgggtcgac cagatcgcg ccctcaacga cagcaagacc 1200
cgcaagacga cctcgagac ggtgcgggag gttctagact ccctcagcga gaagaagaag 1260
tcgtccctt gaa 1273

```

```

SEQ ID NO: 81      moltype = DNA length = 1083
FEATURE            Location/Qualifiers
misc_feature       1..1083
                    note = synthetic construct; Plant optimized nucleotide
                    sequence of MHP55 containing a nuclear localization signal
                    and without an intron
source             1..1083
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 81
atggcaccga agaagaagcg caaggtgcat atgaacacca agtacaacaa ggagttcctg 60
ctctacctgg ccggttcctg ggacggcgac ggctccatca tcgcgagat caagccgaac 120
cagtcctgca agttcaagca ccagctctcc ctgaccttcc aggtgaccca gaagacgacg 180
aggcgctggt tctcgacaa gctggtcgac gagatcgggg tgggtactgt ccgacaccgc 240
gggtcggtgt ccgactacat cctctcccag atcaagcccc tgcacaactt cctcaccag 300
ctccagccgt tctccaagct gaagcagaag caggcgaacc tcgtcctgaa gatcatcgag 360
cagctccctt cggccaagga gtcccgggac aagttcctgg aggtgtgcac gtgggtcgac 420
cagatcgcgg ccctcaacga cagcaagacc cgcaagacga cctcgagac ggtgcgggag 480
gtcctggact cctcccagg atccgtggga ggtctatcgc catctcaggc atccagcgcc 540
gcatcctcgg ctctctcaag cccgggttca gggatctcgg aagcactcag agctggagca 600
actaagtcca aggaattcct gctctacctg gccgggttcg tggacggcga ccggtccatc 660
atcgcgctca tcaagccgga gcagtcctgc aagttcaagc accgcctcga gctgaccttc 720
caggtgaccc agaagacgca gaggcgctgg ttctcgcga agctggtcga cgagatcggg 780
gtgggctacg tccgcgaccg cgggtcggtg tccgactacc gcctctccca gatcaagccc 840
ctgcacaact tctcaccga gctccagccg ttctcgaagc tgaagcagaa gcaggcgaac 900
ctcgtcctga agatcatcga gcagctcccc tcggccaagg agtccccgga caagttcctg 960
gaggtgtgca cgtgggtcga ccagatcgcg gccctcaacg acagcaagac ccgcaagacg 1020
acctcgagga cgggtcgggc ggttctagac tccctcagcg agaagaagaa gtcgtccccc 1080
tga 1083

```

```

SEQ ID NO: 82      moltype = DNA length = 1083
FEATURE            Location/Qualifiers
misc_feature       1..1083
                    note = synthetic construct; Plant optimized nucleotide
                    sequence of MHP55-2 containing a nuclear localization
                    signal and without an intron
source             1..1083
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 82
atggcaccga agaagaagcg caaggtgcat atgaacacca agtacaacaa ggagttcctg 60
ctctacctgg ccggttcctg ggacggcgac ggctccatca tcgcgagat caagccgaac 120
cagtcctgca agttcaagca ccagctctcc ctgaccttcc aggtgaccca gaagacgacg 180
aggcgctggt tctcgacaa gctggtcgac gagatcgggg tgggtactgt ccgacaccgc 240
gggtcggtgt ccgactacat cctctcccag atcaagcccc tgcacaactt cctcaccag 300
ctccagccgt tctccaagct gaagcagaag caggcgaacc tcgtcctgaa gatcatcgag 360
cagctccctt cggccaagga gtcccgggac aagttcctgg aggtgtgcac gtgggtcgac 420
cagatcgcgg ccctcaacga cagcaagacc cgcaagacga cctcgagac ggtgcgggag 480
gtcctggact cctcccagg atccgtggga ggtctatcgc catctcaggc atccagcgcc 540
gcatcctcgg ctctctcaag cccgggttca gggatctcgg aagcactcag agctggagca 600
actaagtcca aggaattcct gctctacctg gccgggttcg tggacggcga ccggtccatc 660
atcgcgctca tcaagccgga gcagtcctgc aagttcaagc accgcctcga gctgaccttc 720
caggtgaccc agaagacgca gaggcgctgg ttctcgcga agctggtcga cgagatcggg 780
gtgggctacg tccgcgaccg cgggtcggtg tccgactacc gcctctccca gatcaagccc 840
ctgcacaact tctcaccga gctccagccg ttctcgaagc tgaagcagaa gcaggcgaac 900
ctcgtcctga agatcatcga gcagctcccc tcggccaagg agtccccgga caagttcctg 960
gaggtgtgca cgtgggtcga ccagatcgcg gccctcaacg acagcaagac ccgcaagacg 1020
acctcgagga cgggtcgggc ggttctagac tccctcagcg agaagaagaa gtcgtccccc 1080

```

-continued

```

tga                                     1083

SEQ ID NO: 83      moltype = DNA  length = 1083
FEATURE
misc_feature      1..1083
                  note = synthetic construct; Plant ooptimized nucleotide
                  sequence of MHP77 containing a nuclear localization signal
                  and without an intron
source            1..1083
                  mol_type = other DNA
                  organism = synthetic construct

```

```

SEQUENCE: 83
atggcaccga agaagaagcg caagggtgcat atgaacacca agtacaacaa ggagttcctg 60
ctctacctgg ccggtctcgt ggacggcgac ggctccatca tcgcgcgat caagccggag 120
cagtgctaca agttcaagca ccgcctcatg ctgaccttca ccgtgaccca gaagacgcag 180
aggcgctggt tcctcgacaa gctggtcgac gagatcgggg tgggctacgt ccgcgaccgc 240
gggtcggtgt ccgaactacat cctctcccag atcaagcccc tgcacaactt cctcaccocag 300
ctccagccgt tcctcaagct gaagcagaag caggcgaacc tcgtcctgaa gatcatcgag 360
cagctccctc cggccaagga gtcccgggac aagttcctgg aggtgtgcac gtgggtcgac 420
cagatcgcgg cctcaacga cagcaagacc cgcaagacga cctcggagac ggtgcggggcg 480
gtcctggact cctcccagg atccgtggga ggtctatcgc catctcaggc atccagcgcc 540
gcatcctcgg ctctctcaag cccgggttca gggatctccg aagcactcag agctggagca 600
actaagtcca aggaattcct gctctacctg gccggcttcg tggacggcga cggctccatc 660
atcgcgctca tccgcccgga gcagtcctgc aagttcaagc accgcctcga gctgcgcttc 720
accgtgacct agaagacgca gaggcgctgg ttctcgcga agctggtcga cgagatcggg 780
gtgggctacg tctacgacca ggggtcggtg tcccactacc gcctctccca gatcaagccc 840
ctgcacaact tcctcaccga gctccagccg ttctcctcag tgaagcagaa gcaggcgaac 900
ctcgtctcga agatcctccc gcagtcctcc tcggccaagg agtccccgga caagttcctg 960
gaggtgtgca cgtgggtcga ccagatcgcg gccctcaacg acagcaagac ccgcaagacg 1020
acctcgagga cggtgccggc ggttctagac tccctcagcg agaagaagaa gtcgtccccc 1080
tga                                     1083

```

```

SEQ ID NO: 84      moltype = DNA  length = 1041
FEATURE
source            1..1041
                  mol_type = genomic DNA
                  organism = Zea mays

SEQUENCE: 84
gtgaatctgt ttggaattga aaaacaagtg ctctctttta tacaccacta tgcgcttca 60
atgtttgcga accaaggttaa agaaatgtaa aatcttacaa ttccgtgca tccgacataa 120
atctgtggtc acatagctat tgttaaacgg ttgcaaatcc taaggaggac cattattgtg 180
caacaactac atatggtaga agcgcttgtt ttgatgtgtg cacattttgt tgctaaaagg 240
atcacgatgc ccaagagggg ggtgaattgg gcttttctaa aaatcaacac taattaaaac 300
ctaagcaaga gcccaacttc accccgacaa cttagcaataa gagaatatga aagggaaata 360
ggatcaaac ttttctctaa tgattttggt ggttgaattg ccaacacaa ataattggac 420
taactagttt gctctagatc atacattcta cagggtgccaa aggttcaaca caaaccaatc 480
aaaagaacaa gttaggcttc aaaagaaagg agcaaaaagg aaaccgaagt gtgcttggtc 540
tggcgccacc gggtgtccgg tgtgccacca gacagtgtcc ggtgcaccag ggtgaatcag 600
ctcaagctcc tcaactctcg gtttccaga cgcagctcca ctataattca ttggactgtc 660
cgggtcacc cgcagagcaac ggtacttgc gcgcaacggt cgactctgca aagtgaacag 720
tgcaattcag aagtcagagc agatgttcag aggggcaccg gattgtccgg ttagcaccg 780
gactgtccgg tgccgcataa ggacaaagcc tccaacggtc gaccagctcc aagccctaac 840
tacaagatga cgtggcggtg caccggacac tgtccgggtg tgcaccggac tgttcgggtg 900
gcccacgcgc agtagccttc tccaacgggt acaatttggg ttggtggctat aaataccacc 960
ccaaccggcc actttaaggt gtgggagccc aagcaacatt ccaagtcata tagttgacat 1020
attcaagcca tccaaccac c                                     1041

```

```

SEQ ID NO: 85      moltype = DNA  length = 906
FEATURE
source            1..906
                  mol_type = genomic DNA
                  organism = Zea mays

SEQUENCE: 85
tcactttccc cctattttt ctccctattt ttctcatctc cgcagcggtt cccctaaat 60
actctatat accccaatac aactataaaa tatcattttc tatatcaact atcaattttt 120
tatctactaa caattactcg tggacccaca tcacaatgtt tagggtgatg aacagtgaac 180
cgctagatct gaggggagag agaaaaaggt cggcgcgtag ggggcgctgt agggggcacc 240
gctgcggctg tggagtgcct cctacagccc ccatgcaagg ggagggggat actgaggggg 300
ctgcgttgcg tacagcctga caggctctcc ttgcgatttg cgcgggacag aaatgacttg 360
ccgaggatgg aagcagagag acggatttgg ccgagcgcac agcagctcgc caaagacggc 420
gtcgaagcag cagtgcacgc ggtcgagtga gggagtcatc ctggattcgc ggtttatcga 480
ctcggcacgg gggcaaccat ggcgttgaag gtaggcaaca tgaggagcca tcgattgaca 540
ccggtcttcg gaatcgcgcg atctcgacga tggtgacaag gaggaggcca cgaagcgctg 600
tcgagcagag cgcgacaagc aaatcgagtc ggccacgagc gtggatttgg atctgacccc 660
caagtttttg tatggatcct attcccacat ttgtagatct tcaatttctt tactttaatt 720
ttccatagca caaacgatgt ttgcatgcac gattcggaca atcttgactt gttcgtccac 780

```

-continued

```

ggttgaggatt taggggttgga atgtgtaaaa cacgtgataa actgtgtaca actcgagaac 840
tagataaatc attttgatt gtaatatgtg tacctcatgc tatagttttg gttaaatctg 900
acgtga 906

```

```

SEQ ID NO: 86      moltype = DNA length = 979
FEATURE           Location/Qualifiers
source            1..979
                  mol_type = genomic DNA
                  organism = Zea mays

```

```

SEQUENCE: 86
tcacgacggt tggggtggag agccggctgg tagggggagga cctcaacggc tgcgccttcg 60
tctactgctc cctcagcttc ctggacaaga tcttctgcgg gatcgccctg tttgccctgg 120
aatcgtagca aggtaagtga cgccgagatc ggcagagatg cttgaaaatt tgtgtttttc 180
cttctgtgct gaaggccggc aactgatcgg gggggcgcaa caccgtcgct ttcgctgca 240
gatacgatga gctcgcgcca gacggggggc ctcaacacgg tgagcaggta cggaacgggc 300
ctgatcccc ctctgcttc gccgtcctta ctccgagagc cactgactg gaagcccgcc 360
tgcttgtctg acggactgaa gaaacgggtg gatgtcgatc gggttgggag aagaaatgcg 420
catcctcttc aattagattt gatttgaaga ggaacatgtc actcgctttt tttttcaatt 480
agaatcctct tcgattaggt ttgaagaggg gggaaatgcc actagttttt ttccaatta 540
gatttgaagc caggccactt tgtaataata ttgccatgc cgtcgtgttg gcacatcaca 600
tatgcatagt tttggtgtgc taatagatga cattaagttg ttgtacgtat aactcgaatt 660
tctgcgaagt ttgtgtgcat gtcacagat tattgtacta agagcaggaa cagcatatgg 720
tcgagggtga aacagagtaa tagtatacac atccgtgtag gaaagaaaaa aaaaactagc 780
tttgaaacag ctgaaaacga cctggacact gaatgcaaac atccccgcc gcggcgggct 840
ctctcacag ctcgctctcc gactccgacc ggtacttgct cactccgcc ctccggtgct 900
tccccctgcc gtcgacggcg ggcacgtcgg gccacgcgac cctctccttc acctcctcca 960
gcctctcctt ggcgctgtc 979

```

```

SEQ ID NO: 87      moltype = DNA length = 734
FEATURE           Location/Qualifiers
source            1..734
                  mol_type = genomic DNA
                  organism = Zea mays

```

```

SEQUENCE: 87
tagagtagta gtcctgtgct ggtgcgtgga gtcgcttgag agcgttggcg gcgagaagtc 60
cgccaacggc actgtggatc tcggccacgg aagagaaaaa gaaaaggcaa aaattgcatt 120
gtcgaatacg tgaacaggaa aatccaattt tcgtatcatg acctctgtat atgtatccat 180
atatataaaa aaaattctaa tatataaaca gactcaatat tttgtaaaaa atgccatttt 240
aaatttgtat taatatatgt tggaaaatgt aaagaatgag atatagagaa cgaaattttag 300
agaaggttgc tgaagatata aaagattaaa tcttttagag tgtgctataa aggatagaga 360
atatattgtt aatggatgaa atttagagaa cgttattgga gataggctaa aaaaataact 420
gattgcaaaa attcagcctt ccttccactc acccatctct ggaactgcct gctgcctcg 480
aacgtaggag atcaagtggg acgacggcgg cctcaagtcc ctctcacagg tcggcgcgac 540
gctctgggtc atctccggcg tcaccgtctt cgtcttcccg agccagatgc acaacacctt 600
ctcgccatgg tcacggcgcg tgccaacggc ctcgctcatg taaaacggcg gcgcctagct 660
agcacgcccac gctgcacgtc caaatcctac cggtttcgcg tgctctggct ttacattaca 720
tgggcaggtc tcac 734

```

```

SEQ ID NO: 88      moltype = DNA length = 1016
FEATURE           Location/Qualifiers
source            1..1016
                  mol_type = genomic DNA
                  organism = Zea mays

```

```

SEQUENCE: 88
atacataccg ccctaaggac ctagatagtg tgttggtcca aaaaaccaac atagaaacat 60
caaagttttt gccattaata gaaaagattg atggacaact catttccaaa gccatacagt 120
caattgtaat catcttcaac ccaatcaaaa actcatagtt tataaccggt ctttgccag 180
cttcacggat agtctgtccc tctgagactg aaagtgcctt agaagtgggt gaataggcta 240
aacctaaaaa ttccaccaca aactttgaaa taatgtcaaa taagcagttc aactggtgca 300
ggccagtttc accgctatta aggcgggttg aaccactcta aaccggccca accgtgagag 360
aggatcaagg ctatgaacaa cgagagactt aatgagagat tcttttaaga aaagagctca 420
cgggataaat taggcataaa tagggaaaat ttgtgtggat aagatccaca cacaagacaa 480
ctcgatcgat gtcttctttg ctaaaagataa ttcacaacga tttgaattaa agcaaaagaca 540
caaagacgca aggatttacc ctgagggttc gccacaccat aaagggtgcc tactccctgt 600
tgaggagccc acaaaggacc aagtcttttc caactctaat cctccacaaa tcgaccacaa 660
aggatcaagg aaactcttct tcaactttgc tcaacgagtg agtgaacaaa acttcttggg 720
gtcgtccaca aatttggaga ctcccaagca acctcaaaact gccaaaggaa tcgaagggtc 780
caagggcaac aaactctcac aagaagtgtt tgcagtggcg tcaagagatg aaaaaggggg 840
gggagagaaa actaagtcta aaagtgaaaa actcaaaactt tacaccaagg gcccttcaat 900
caagcgatga gggagcgatt tggggtgtga gagagttggg agcttttatc tcaagttagg 960
tcagcaatga atgcgtggag caaccataat gaatgaggag agagacatga ggggggt 1016

```

```

SEQ ID NO: 89      moltype = DNA length = 898
FEATURE           Location/Qualifiers
source            1..898
                  mol_type = genomic DNA

```

-continued

```

                                organism = Zea mays
SEQUENCE: 89
ctttggcaag caattgcatg cgagtaaaca agtaattaag agtaaggttc accgggttagg 60
ttccttacct cgggtgtaaag ctggaacat ggttgttgag gttagttagg ttccttacc 120
acaagtcaca ctctoccaca tgggtgtgctc atacctaatg tatacttgat cagcctagac 180
cacttggcgc tcttcacacc ccactctact aatgtgtctc tctgtgtctc tgtggggcgga 240
gcacggtagc ccttacaatg cctcctttag agccacacac gatttcacgc aggattccat 300
ggagccataa cctccaaggc acctaggagg tggaaacctc taaaagtaac aagacaatga 360
tcttcctagt gataacttga taatgtgagt tagtaagagg tttggggcgga aggctcaagc 420
atgctcaaca agtgcctcta ttgctcagct tagggagcac acatttacac tcctactttt 480
tatagcccca cttccacaaa ctagacacta taacactttt tgagaaaact acacattagt 540
ggacactcca taatacaacc caccgatagc ccatatttga attccgatga ctatatttca 600
attaaatgcg tgttagtcgt catagaaagt gtttagtgaa cagtctatct gttattttt 660
aacatgtcta taaacttctc aatttatgtc cctttttaag aatgtgcgac agatagtctg 720
cctttgaggc ccatatagta caccgaccaa atatttgcac tcaccgaaac tcccaagttt 780
ctatccacta ctataaacag gttagagaca gtctacatga ggggcccaca tagtccatcg 840
gtcaaaaaac acataaactt taaagttttt gtccatcact tgaattagta tgacatac 898

SEQ ID NO: 90                moltype = DNA length = 260
FEATURE                      Location/Qualifiers
source                       1..260
                               mol_type = genomic DNA
                               organism = Zea mays
SEQUENCE: 90
agggcaagtc aatccatgca acaccactca cgaatatgtt acgttaatac caaaaccata 60
tattgacaca catgcacaac atcatatatt atgttaatca ctttataaat ccaattttta 120
tctaaaacaa tgttttatca cacacgattt cgaatatata atcgggtgata aagatacgcc 180
ggttgaccaa gtaagtcacac aaagggtcga taacgtctgt acacttaaaa ggaggcgagt 240
cacacatcta tatgggtgct 260

SEQ ID NO: 91                moltype = DNA length = 900
FEATURE                      Location/Qualifiers
source                       1..900
                               mol_type = genomic DNA
                               organism = Zea mays
SEQUENCE: 91
cgtacaactt taggaatcac accagccgcc tactctaagc ttgggacctca cttctgtggt 60
ccctatcagg ttctggagga aattgggaca gtatcttaca ggcttgccct gccatctcac 120
gcccccatcc ataattgtgt tcatgtctct ttactaaaga aatatattgg ggctcccccg 180
gctgcaatag ttctcttacc cccaatatta catggcagag tctgcctca gcctgagaag 240
gttactaagg cacgcaagaa ccgaggcgtg tgggaactgc ttgtgcaatg gctgggacaa 300
tcagccgctg atgcaacgtg ggttcagctg gaggacttcc gtcgcccgtt tctgtgtgtc 360
caggctgcgg acgacttggt tttgggggag gggggaaatg ataccgatgc atttgtagga 420
aaggtatacc agagaaggaa tcgccaggaa taaagggaac aacagataag gaataaaaaga 480
gacacagatg aattttctat atttagtcag tcagcagatt aggaataaaa gagacagcag 540
ataagtttct atatttagtc agtctatttt ctacgaagtt gagagtgata tgatttggtt 600
ctatattaac ctgggctcag tctataagag accaggggta gtttgacta gggattatca 660
aaagaagaaa atctcctagt cctaggaggt tgcctggggc cctgggtgac actggaggaa 720
ctctccagcg tccgggaacgc caccaggaat cctcctcccc cttccactc ctatttctg 780
cgttcattgt ccacaacctc ctgctgagcc cccaacgaaa gcaggggagt tgcgtcactc 840
gaccccaaac tgataaagggt ttaaggtcgg gaaatctcac ccgtgaagtt tatctcgcaa 900

SEQ ID NO: 92                moltype = DNA length = 931
FEATURE                      Location/Qualifiers
source                       1..931
                               mol_type = genomic DNA
                               organism = Zea mays
SEQUENCE: 92
ctacgtcatg taagtttcta gtggttgatg tgctctgggt ttgaagatta taggtgattg 60
ttgaaggtag aatatgaaag tagcctacga gtctaatgga gtctcccgtt tcagcattat 120
atggccaaat gaagagataa cctgtatag atcataatcc tatatgcac taaccacttt 180
cactatagac gcaacgcatt ttccggcggt cgggttggtt cctgcgaggc ttgtgtacgg 240
ctgcatgcat gggacatgcc ttccaagatc cccttgactc ggtgtgtgac cgctttactc 300
ggcttcgggt gcatcttgca gagggcccac atgcacccta tggacgaccc atcaggggac 360
ttataagcgt cacatggagc gcatccatgc atgggtgacct aggggatatc catgtaacac 420
cccagggtgt agctagaagt aataacccaa ccacttgga ccatatcaca tgtggataac 480
ttaaggtaaa agtcactaaa attaatgacc atattcctaa taagggtgaa aacaccctag 540
aagaattaac ttaccacccc catggtgatc aaaggaaagg ggagtaacca accccctaaa 600
cctactctct tgagcccaag agcaccaata caaagtgtca agagaaagt aacccaaatc 660
cttaaccaca agtggaccct taacaaaagt tatagctaac taaataccta acaaaagttc 720
ttgaggggta agcaccaaa ggggtgctag agtcccaatc aagtcacaca tgtgggagaa 780
ggggagagaa atcaagattt ttccataaat ccaaaaacag ccctatccca aaaacataaa 840
atctccaatt atgaaatgtg tgctaatgt tctagggaac accctcgtaa agtttgaact 900
cgagccctca ctgtttgaca tgacaagtca t 931

SEQ ID NO: 93                moltype = DNA length = 922

```


-continued

FEATURE
source

Location/Qualifiers
1..922
mol_type = genomic DNA
organism = Zea mays

SEQUENCE: 93

```

cgccgcatta aatgcgcggc agagagccgt tggcgcggag atgaccgttg cttcggagtc 60
gcaccggaca gtcgcggtaa ttatagcggg cttagccgttg gagtttccc aagctggcga 120
gttcctgagg ccgtcctccc ttggcgcacc ggacactgtc cgggtgacac cggacagtcc 180
ggtgaattat agccgagtcg cctctgcgaa ttcccgaagg tgacgagatt gagtctgagt 240
ccccctagtg caccggacat gtcgcggggc gcaccggaca gtcgcggtcg ccagaccagg 300
ggtgccttcg gttgcccctt tgcttctttg ttgaatccaa aactcggctc ttttattggc 360
tgagtgtgaa ccttttacac ctgtataatc tatacacttg ggcaaacctag ttagtccaaa 420
gatttgtgtt gggcaactca accacaaaaa ttatttagga actaggtgta agcctaattc 480
cctttcaagg cttcacttcg gaccactcta gaagtctatg gatggcttag cctcttagca 540
tgaacgatcc acgacaatga tacttagccc actttccaaa acacgctttt gaaaatattt 600
taactcagca attcagaaga attgttaata atcttgctaa tgcacatctt aaaagctcta 660
tgaggcatta agtttcacat aagaaattgt cattgactcc tcttgacagt atggctatct 720
atccgactaa cccagacaat tttcttctct aaacaccttg tgactggtgt cgggtgtttg 780
taccaatggc gcatctatgg atataccatg tagtgctttt gggaggatag cgaatgcgat 840
caaaacttga ggttctatgc caggcacgat ggaacagagc agattatata ggtttgaacc 900
acctagaggc gtaatgtcct ac 922

```

SEQ ID NO: 94

FEATURE
source

Location/Qualifiers
1..994
mol_type = genomic DNA
organism = Zea mays

SEQUENCE: 94

```

acgctgagga aactgagaat cgctgcggg ccgcgaatgg cgagcgagca cggcttgaga 60
atctgacgat ggaggaagct tggatagtg aagggatatt ttggagatca attttagat 120
gctggtggaa gcgttttctt tctatactg gccttctcga aacctgtagg agttgctcta 180
acaggtaaag atcatcagcc cctcctgtc cctcagttg taatacagca atttcagata 240
tgagcataga aaaactctag gtcccaccat gatagtaatc ctctaccatc aaaatatatt 300
ttatttgttt tcgaaaataa ataaagacaat gttaagagta catgtagaac cttctaaaca 360
tctgaacttc agattcaaca ccaaatcaac atgatgagtt tcaacatgtg agttaaata 420
caaagtgggt tgcttttagaa agcaacacaa gttacctagt tagggcttaa gcagacaatt 480
attttttggg tagtgattta caaaataatt tattttccga ttgcaactat gtgttacact 540
caattttaaa aaatatgttt tataatcaga ccacacatcg aagtacaggt gtgtattatc 600
gaggtacatg attatttcaa tatttgagag agccttttca acttggtaca attgggacac 660
ccaaatggaa agaaacagta tgatcaaaag acctgaatag gtgggcacaa taactgaagt 720
tatctggcca attattagtt aacacttttt tagaattcct ggggcctcgt cagcatgtac 780
gattgacctt aaattgttct ggtcagcata gttattgaca actccggaaa ctatctgata 840
agacatactt gggacctgtt tttttgagat tataaatatg tttagattat aaccccaaac 900
aaacaaaccc ttcataattc taaaacaagt ctaaaattta attaaatata taataaatgt 960
tacatatgct atttgtcacc taggtgcgac tggg 994

```

SEQ ID NO: 95

FEATURE
source

Location/Qualifiers
1..1048
mol_type = genomic DNA
organism = Zea mays

SEQUENCE: 95

```

ctgccgatgc tatataagtt gactgcacct gccgatcgtt gacgctcgaa cgtcgacct 60
cctgaagaat aggtcctctg tgcgtgtctc ctaaccacgg ccacctcgac ttcggctaac 120
tcggcatcca ggggctatct ccttatcaga atccacaccg gtctcttctc cagccacaac 180
attggcacc ctcacgacgt gcgaccgcag ggggggtcaa cccgtcgact cctaccttcg 240
gcctctactc cagtttcatc gtgtgtggtg ccccggttgc gactgcgggg atgttagact 300
gtgtgtgtag gccggcacc cgtttgggct gccggcccat taggggttag gttgtgagtc 360
tatatattat ccccatctc ttatcaatac aaccaccact tgatacttct acatagagga 420
tagaggtagg agcagccctt aatcttcagc tttcatagcc aactgcccac gaatatccat 480
aaacctagcc aattcacttc tccaatgcgc ttagtctagt aaaagcaaat gccctatgca 540
tgtaacttta ccttgcactt tcttttccac ttctgcactt ccatccatca tcttcacatg 600
ttgagcactt gcacttcatg gtcttgccta tctccacttc acggttctat atatgtggct 660
caactatctt gtacactaaa tcgcctattc atctcactat aaataaatta gtctggcatt 720
caattatcaa agccaaatca ggtctttcac tccagagctc ctgcttgact agttgcgcgt 780
cttcctgtat gtttgccacc tcttccaccc ccagactgca gtggtatact tcccccaat 840
ctatttttag tgctaaaaat ggggctttct caaattgaat tatttgcctc tccatatgcg 900
catcttactg taatcgcgg tgggtgccta tgaccgccag gatgtgtcta caacgcagta 960
cctatggcta taagtgtcag ctctctccac aactggtagg ccaatctccc ccatgcaggc 1020
gcgcacagga gaggaaggc tctcacgc 1048

```

SEQ ID NO: 96

FEATURE
source

Location/Qualifiers
1..901
mol_type = genomic DNA
organism = Zea mays

-continued

SEQUENCE: 96

ttgcgtccga	tctcatccac	ccgctcctga	tccaaacgacc	cagatccttg	gataccgggt	60
cgagcgccgc	ccccacccc	taggccccac	acgttgccgc	ctgcgcccc	gatgctaggc	120
ctgaactgtc	agtcacccct	ccaccctggc	cgtgaccgc	tctgtcaccg	cttgctcgcg	180
ccccgctgc	cttgcccgc	gatctgatct	cggcagttga	tctgtgatcg	gatggccgag	240
agcgcccgat	acccttctat	ttggaatttt	tgtaaagag	atccccgggt	tcttagaaat	300
caaccgcag	tctagtttta	ttgcgcctga	gtccctgggt	ttttgcagag	agaccagta	360
actttatttt	tatcacaaaa	attgggttaa	tttaggggtt	tgaattccaa	aacttgtaaa	420
tttcatatct	tttgcatatg	aactccaaat	tgggtgggtc	aaattgcaaa	atgttcataa	480
tgttattctc	tatgtgttta	aattatattc	atttactatt	ttcatgtctc	aattttgtgg	540
ctaattcccta	ggtaatttta	aagtgataga	atatttatta	aagggtaaaa	taaaaggtaa	600
agccctaata	aatgtccatg	tgcttaactt	tgtaaaccta	atttcattta	atgtaatccc	660
atccctagaa	tctgttttatt	taagtaagta	atttattgag	atagacttag	ttagaaaata	720
gtagaccttt	aaacatagtg	atctacccta	atacctagag	ttcacttggt	tgtttgtact	780
tttctactga	acctttgttt	gatcggttgc	acatgtttgg	tgtgctgttc	tttgtgttgc	840
cccaagtgtg	ttgaatgaat	gattgctttg	cgtacacaac	gagcaatccg	aggttccgag	900
t						901

SEQ ID NO: 97 moltype = DNA length = 1083
 FEATURE Location/Qualifiers
 source 1..1083
 mol_type = genomic DNA
 organism = Zea mays

SEQUENCE: 97

atggcaccga	agaagaagcg	caagggtgcat	atgaacacca	agtacaacaa	ggagtctctg	60
ctctacctgg	ccggcttcgt	ggacggcgac	ggctccatca	tcgctcccat	caagccgaac	120
cagtcgccga	agttcaagca	ccagctcatg	ctgaccttca	ccgtgaccca	gaagacgcag	180
agggcgtggt	tctctgcaca	gctggctcgac	gagatcgggg	tgggcaagggt	ccgcgaccgc	240
gggtcggtgt	cgactaccg	cctctgccag	atcaagcccc	tgcaacaact	cctcaccacc	300
ctccagccgt	tcctcaagct	gaagcagaag	caggcgaaac	tcgtcctgaa	gatcatcgag	360
caagctcccc	cggccaagga	gtccccggac	aagttcctgg	aggtgtgcac	gtgggtcgac	420
cagatcgccg	ccctcaacga	cagcaagacc	cgcaagacga	cctcgagagc	gggtcgggcg	480
gtcctggact	ccctccagg	atccgtggga	ggtctatcgc	catctcaggc	atccagcgcc	540
gcatacctcg	cttcctcaag	cccggttcca	gggatctccg	aagcaactcag	agctggagca	600
actaagtcca	aggaattcct	gctctacctg	gccggcttcg	tggacggcga	cggctccatc	660
aaggcgacga	tcaagccgaa	ccagtcctac	aagttcaagc	accagctctc	cctgaccttc	720
caagtgaccc	agaagacgca	gaggcgctgg	ttcctcgaca	agctggtcga	cgagatcggg	780
gtgggctacg	tctacgaccg	cgggtcggtg	tccgactaca	tcctctccca	gatcaagccc	840
ctgcacaact	tcctcaccga	gtccagcccg	ttcctcaagc	tgaagcagaa	gcaggcgaa	900
ctcgtctcga	agatcatcga	gcagctcccc	tcggccaagg	agtcctccga	caagttctctg	960
gaggtgtgca	cgtgggtcga	ccagatcgcg	gccctcaacg	acagcaagac	ccgcaagacg	1020
acctcgagga	cgggtcgggc	ggttcttagac	tcctcagcgc	agaagaagaa	gtcgtccccc	1080
tga						1083

SEQ ID NO: 98 moltype = DNA length = 1026
 FEATURE Location/Qualifiers
 source 1..1026
 mol_type = genomic DNA
 organism = Zea mays

SEQUENCE: 98

tcgcgagagc	ttgggggggc	ttgacgactg	agtgagtgtc	tttcgtgtct	ggccttctctg	60
ctctcgtttg	cgtcttcgcg	caaggaaacg	ggaagagaaa	aagagggaac	gtcccgtccg	120
tgccgacgtt	ctctgtgcatg	gggtgggtctt	gcatgatttg	tgctgtccgt	cctgtgggtg	180
acgggaagcg	acggcgagtc	ggcgaaagcc	agctggagcg	tagagccaag	agcccggtgt	240
gtgcgcctgt	tgtgtctgtg	ctgtgccgtt	gcagttgcag	ttgcagttgc	agttgcagcg	300
acgggtttct	acttgatcac	ttcggagttc	aggcaaaagt	ctcgtgggtg	ctgccatgcc	360
accactggct	gatagcgtgt	ggacccattc	caggcccata	cccactttac	ctaccggggc	420
accaaaagcg	cgaagcctgc	tattgtagta	ttgtcggcct	gcgcagcaga	gcgctgagtg	480
tctactgatt	ataccgctga	aattaaatgc	ggtatttcgt	tttcagacca	aaccagacca	540
gatcagccag	tgcaaaagccc	gcagtgggat	ccaggcaaac	gtttttctcca	ctgcaatcga	600
tctgtctgta	cgtagaggcc	gttagtctac	tgagcgcaac	gcgtacaagt	tgctgttctg	660
ggatcgctag	ctcacatacc	tctcgacgca	ctcggttgtt	ggcttcacat	gcattggccg	720
ccaccacctt	ctcggtgacc	acctacatgg	tctctctagc	agaccccgct	agtgcgcgcg	780
gcattcggtg	catgcattgc	tgtatggaca	tgacgtgctg	tctcgagcaa	taattagatc	840
catgtttgca	ccagagatgg	gtgacctcgg	cgtgcacgaa	ataactggta	ccatcagtg	900
acaaaacaag	cattcctcgt	ggcgttcatg	gcgggtgatg	cagtgaggaa	gtacaactga	960
cttcaggggc	cgtgacggac	cgtgggataa	agacgcagta	gagcaggaaa	gatacttctc	1020
accaaa						1026

SEQ ID NO: 99 moltype = DNA length = 856
 FEATURE Location/Qualifiers
 source 1..856
 mol_type = genomic DNA
 organism = Zea mays

SEQUENCE: 99

caaaggagtc	aaggacagac	actgtcgtatg	atatagagaca	aacaacaaca	aagcacactc	60
------------	------------	-------------	-------------	------------	------------	----

-continued

```

cgtttcaata aaaagaagat tatcggactc cctcatacca gccctaccta cgacacgtgt 120
aacgcctcag gggaggaacg ggacacaccc atgcatgcgg atgcggggcg agcatatcct 180
gacacgcgcg gcgtagcccg cctgtcagtg agtctgcacc gtccgatcga gccgcgcggg 240
tcacgtcgag gccgcgctgg tcggctcctt tttcttcgtt ccccggttcc ctctgcccc 300
tggatgtaa tttttttctt gccctcgtgg agtaccgacc gcgagaaagg aacggccgtg 360
ggaagaaagc gacgggaggg ggcgggcgct tggatcgcg gcactctgag aaagatggga 420
atcctcctgc aggacaacta gacaagtgtc caccggaaca gaagacctta tctagtagta 480
gcagaagagt ggtagcagta cacctttcta aagtttgatt taaaaaaaat tgaagtatc 540
aaatatctat ttaacaatat gtaatgcttc tactacaaag tttttttgt aatggagaat 600
ttagtaatac tcattttatt tataagtact aaaatttcat ataaattagg ttaaacctta 660
aaataacttg actgagtcga cgctctatta taaaattttt tttctcagac cgaggagta 720
aataaaacta aacaaagcga ggagctgcgc ggaggacatg tcaacgaggg agacgacgac 780
gagagaaacc aacgagattt tttgccagac atgaaaagcga gagaggcctt gtatgttcct 840
cgctcgccct tttcgc

```

```

SEQ ID NO: 100      moltype = DNA length = 1068
FEATURE            Location/Qualifiers
source              1..1068
                    mol_type = genomic DNA
                    organism = Zea mays

```

```

SEQUENCE: 100
tcacgtatag tagtgaccc atggagccca ttgccagtag ctgtccccc tttacgtctg 60
gatcagggtga ctcaatagaa tccggctgag atcctcgcta gtgaccgtta cttctctcta 120
ggggaacctg cctctctatg tatataagaa accacgccc ctgcgcctct aatcctaccc 180
atgttgagcg tcaccgaaag ctgagagcct cgccatcgag tgaggggggg ggggggagat 240
tgccgccacg gtcaaccat tagtgtgcc catccccgc ttcggagcat ggtctagggg 300
catccacggg tcgtggcgga gctatccgtg gcattatcaa cgaggatggg tccctggacc 360
aatgggaatt tctgcgcgta gtccctcgcc atagatccgc ctgcaccgtg gtcggaggac 420
atcacctact tcactctggg tacataaccc ctataccgtt agctagggtc cctatccat 480
gtagctctcc tccgatcaag atctaaagca tcaaaagcgt aacctgaaag ggaatgggt 540
aaccatttcc tataatcgat ttgtgtgtt gacgactatc acaaacata tggactaact 600
agtttgccca gtcaatattt ttccttaggt gcataaagtt catatacaca ttgtcgggta 660
ccgtaattag gggtaacccc aacactccta aacacgacta gtaaacacct tcaaaagcaa 720
ccatgaagac caacagttcg ggtcaaaagc aaagcttcgt ctaccaaggg acacgatctc 780
gcctcggccg aacccgaccc caggcgggaa cagtagtccc ggacgggatt acgtctcgcc 840
cgagggtctc ctacggcagt gagcacaccc tcggctcagc caaaggcaag ccttgtcgtg 900
caagcgaccc tgcccacaaat gccttaccag tcgacgctat tgcagcgca ttaatgctg 960
ggatcgccgt acaccttatc ctgacacgcy tgctcagtt gacaaggtcg aagtgcaccg 1020
agtcgcttcg cccttcact gaccgatctg acagaaaaat agcaccgc 1068

```

```

SEQ ID NO: 101      moltype = DNA length = 982
FEATURE            Location/Qualifiers
source              1..982
                    mol_type = genomic DNA
                    organism = Zea mays

```

```

SEQUENCE: 101
tttctgacag cccaacgaat ttcgttgatg aaatatgcaa ctttcatgat ctctgaagtc 60
gtaaatgcat cttcatatat cttcaagcct ttcacgacat agcaaaaggg tattgtcggg 120
acctataaac taagtatcct cttctactgt attaagatag agacctccgc ggttaacttt 180
aaacgcgtac taaggtaatg agtggccggg cctacgggac tggacctgtc tgtcaggacc 240
cctggcttca agtccgctc ttcacctcga atttgtaaca ccaaaaaatc ataatttttg 300
gatttagaaa aaatatatta ttcgtgaaatc taaataagaa ttttttctca taaaagattt 360
aagtaaaagt atttcataaa agagattata tattagaaag ttttctctg taaaacaaaa 420
caagtaataa atattaaagt ttttttaagt aacttttaat gtgactacac attcaaaata 480
ttttcgtaaa ataaatatta tgtgtgttgc atattgaaaa cattgcctaa ataaataaat 540
aaggtaataa attaatgaat aaacttaata cacaaacctt gcattcatgc tggatatatt 600
tttgtgcaaa ttagaacttt gtttgaatct aaatctaatt ggaattggaa aatagaaaat 660
agaaaaagaa taaaaaagga aaaggaaact ttacatgcat cgtgggcgga gtaacgcagc 720
acgtccaccc cgcggtctct tttcttttc tactagtcat tgacacgcga gcccacagt 780
gcagcctcac catctcgcac tcgaatgggc tctagaagtc gctcccaaat ggggccaagg 840
tgccaacacc ttcttctcta aatcaatggg gatcacaatg aatcctttcg taaccgcat 900
gtaacggcct ccatagatcc cgaccaccag tactttctct attcgtgcgc gcttgcccga 960
gggacggtg tggcaaggtc tc
982

```

```

SEQ ID NO: 102      moltype = DNA length = 1052
FEATURE            Location/Qualifiers
source              1..1052
                    mol_type = genomic DNA
                    organism = Zea mays

```

```

SEQUENCE: 102
agaatgacga ttgctggact gggagacatt gggttggcgt aactgttcag gacttcgggt 60
acccgtgggt caccgtggg tgagtgaga aaccgggacc cgaaccgaa aagggtcggg 120
tcgggttcgg gtcacaccg tgggtgaaaa aacacaccgc cgccgcacc cgtcgggtcg 180
ggtaccgcag gggtaccgca acctgtgggt gaaattgccg tgctacagc tggctgtggc 240
gtcaggctcg ggagcccatg ccaattgggc taaaagccca aaattttcca atgcacgcca 300
ttggggctgc acagttctag cactagactg gcttttcttc agtgaaactg agcgtgcact 360

```

-continued

```

ttgcaactgc ttcttcaggc aaacactaca tatgattgga cgtccagctc gtgggcgcaa 420
ggctcgcggc ggctcggaag cgcagggatg ctccctcttc gtatctccat gcgtacaaac 480
tgacacacaa aaaagccatt gagttgcacg ggtgcggtgca ggctcgtctc cattcataca 540
gcgcccacca atcacccggt aagtgcggtc aacggaacgt ggagagcctg gacgcacgcg 600
cctagggtac caatcacgcg gactactgtg tccagccgaa gccttcacac gcggccgagg 660
accgaggtat agttcaacgg aacgttgccg tgcctgatcc ggcagtgaga ctccggcctg 720
catcgttgtc gtccttgctt cagaactgac aaaacatgga tgtggacctg gcctagaagg 780
tcatacaata caaaactaga attattttag aattgacagt ggcagagtat taggaatcat 840
cagctgcgat aacataacgg acagttaata ctccatccat ttcaatttat aattcactta 900
tcttttttat cctaaatttg ataggttcgt cttattcaaa aaaaattata attatcatta 960
atttttactg tgatatagtt taacataata tacattttta gcgtgggttt caatttttta 1020
tttttcacaa aaacatggta agaaatacat gc 1052

```

SEQ ID NO: 103 moltype = DNA length = 792

FEATURE Location/Qualifiers

source 1..792

mol_type = genomic DNA

organism = Zea mays

SEQUENCE: 103

```

atgcgagcca gtcaaaacttg ttaagaaaaa tcaaacgaat tagaaattag gacagtggga 60
gtatacaaac gggagtatat aaacgactat cactattaat ttagtggtga gtgagcaatt 120
atcggttcgt gtctatcact atcactcatg atatgaatca acggtgataa ctgctaagca 180
taaatgggtt ataatttttc atatgagatc gaacgaagat atactctata ttgaaattat 240
ttagtttaaa ggaatctaaa actttacagt cgataaactt ttaattaat cagagtgggt 300
taccggtgta aatattcaat atgatttttg aatcatattt gacatgatta acaatgtcaa 360
attcgcaata tcgaatgaag acaaaactca cattaaagtt gtgctagtat aatgctcgtg 420
tgttgtgaca aaacataaat atttgatagt ataacgatta catgaaaatg aacaatagat 480
atattacatc cgatcgacct taatatctga caaattattt gtcaacaacc aatacaaaac 540
taaaacttga attcagaacg cctcctcctc tcccttagct gtccagtcac gcatcacggg 600
tggcagagcc tcctcctcct atccagggaa ggtcgatggt ctttatgggt gtgcggactg 660
cggaagaatc tgggtgaaaga aggtggagac gagaagaga tgagaagtta gagggttaat 720
aaactaatat acgatgacaa gagcgatgaa gagggaggatg aagcgcttat ggtgggtgcat 780
gattgggtgc at 792

```

SEQ ID NO: 104 moltype = DNA length = 15574

FEATURE Location/Qualifiers

misc_feature 1..15574

note = synthetic construct; plasmid PHP44285

source 1..15574

mol_type = other DNA

organism = synthetic construct

SEQUENCE: 104

```

gtttaccgc caatatatcc tgtcaaacac tgatagttaa aactgaaggc gggaaacgac 60
aatctgatca tgagcggaga attaaggagag tcacgttatg accccgcgcg atgacgcggg 120
acaagccggt ttacgttttg aactgacaga accgcaacgt tgaaggagcc actcagcaag 180
ctgggtacgt tgaatacaga ctcaatatag ggcgaattga gcgctgttta aacgctcttc 240
aactggaaga gcggttacta ccggtctggat ggcggggcct tgatcgtgca ccgccggcgt 300
ccgactaac taactagtcg agctagttag cctatgaggt gacatgaagc gctcacgggt 360
actatgacgg ttatgctcac gactgttggt ggcagtagcg tacgacttag ctatagttcc 420
ggacttaccc ttaagataac ttcgtatagc atacattata cgaagttatg ggcccaccgg 480
tggtagccag ctcgtttaaa cgtctctcaa ctggaagagc gggtaccaga gctgggtacc 540
tttgtccacc aagatgggaa tggcgcgccct cattaatata gtcagcggcc gctctagtgt 600
aagacacggt catgtcttca tcgtaagaag acactcagta gtcttcggcc agaattggcc 660
tctggattca gcaggcctag aaggccattt aaatcctgag gatctggctc tccaaaggac 720
ccgggatagc ttgcatgctg gcagtgcagc gtgacccggt cgtgcccttc tctagagata 780
atgagcattg catgtctaag ttataaaaaa ttaccacata tttttttgt cactctgtt 840
tgaagtgcag tttatctatc tttatacata tatttaact ttactctacg aataataata 900
tctatagtac tacaataata tcagtgtttt agagaatcat ataatgaac agttagacat 960
ggtctaaagg acaattgagt attttgacaa caggactcta cagttttatc tttttagtgt 1020
gcatgtgttc tccttttttt ttgcaaatag cttcacctat ataatacttc atccatttta 1080
ttagtacatc catttagggg tttaggttaa tgggttttat agactaattt ttttagtaca 1140
tctattttat tctatttttag cctctaaatt aagaaaaact aaactctatt ttagtttttt 1200
tatttaataa tttagatata aaatagaata aaataaagtg actaaaaatt aaacaaatac 1260
cctttaagaa attaaaaaaa ctaaggaaac atttttcttg ttccgagtag ataagtcag 1320
cctgttaaac gccgtcgacg agtctaacgg acaccaacca gcgaaccagc agcgtcgctg 1380
cgggccaagc gaagcagacg gcacggcatc tctgtcgctg cctctggacc cctctcgaga 1440
gttcgcgtcc accgttggac ttgctccgct gtccgcatcc agaaattgag tggcggagcg 1500
gcagacgtga gccggcagcg caggcggcct cctcctcctc tcacggcacc ggcagctacg 1560
ggggattcct tccccaccgc tccttcgctt tcccttcttc gcccgccgta ataaatagac 1620
acccctcca caccctcttt ccccaacctc gtgtgtgtcg gagcgacac acacacaacc 1680
agatctcccc caaatccacc ctcggcacc tccgcttcaa ggtacgcccgc tcgtcctccc 1740
ccccccccct ctctaccttc tctagatcgg cgttcgggtc catgcatggt tagggcccgg 1800
tagttctact tctgttcagt ttgtgttag atccgtgttt gtgttagatc cgtgctgcta 1860
gcgttcgtac accgatgcga cctgtacgtc agacacgttc tgattgctaa cttgccagtg 1920
ttctctttg gggaaacctg ggatggctct agccgttccg cagacgggat cgatttcact 1980
attttttttg tttcgttgca taggggtttg tttgcccttt tcctttattt caatatatgc 2040

```

-continued

cgtgcacttg	tttgcgggt	catcttttca	tgettttttt	tgtcttggtt	gtgatgatgt	2100
ggctcggttg	ggcggtcggt	ctagatcgga	gtagaattct	gtttcaaac	acctgggtgga	2160
tttattaatt	ttggatctgt	atgtgtgtgc	catacatatt	catagttacg	aatatgaagat	2220
gatggatgga	aatatcgatc	taggataggt	atacatgttg	atgcgggttt	tactgatgca	2280
tatacagaga	tgccttttgt	tcgcttggtt	gtgatgatgt	gggtggttg	ggcggtcggt	2340
cattcggttc	agatcgga	agaatactgt	ttcaaacac	ctggtgtatt	tattaaat	2400
ggaactgtat	gtgtgtgtca	tacatcttca	tagttacgag	tttaagatgg	atggaatat	2460
cgatctagga	taggtatata	tgttgatgtg	ggttttactg	atgcataata	atgatggcat	2520
atgcagcatc	tattcatatg	ctctaaccct	gagtaacctat	ctattataat	aaacaagtat	2580
gttttataat	tattttgatc	ttgataact	tggatgatgg	catatgcagc	agctatatgt	2640
ggattttttt	agccctgect	tcatacgcta	tttattttgt	tggtaactgt	tcttttgcg	2700
atgctcacc	tggtgttg	tgtaacttct	gcaggtcgac	tctagaggat	ccatggcacc	2760
gaagaagaag	cgcaaggtgc	atatgaacac	caagtacaac	aaggagtctc	tgctctacct	2820
ggccggcttc	gtggacggcg	acggctccat	catcgcgag	atcaagcga	accagctcta	2880
caagttcaag	caccagctca	tgtgacctt	caccgtgacc	cagaagacgc	agaggcgctg	2940
gttctcgac	aaagctgggt	acgagatcgg	gggtgggtac	gtccgcgacc	gcgggtcgg	3000
gtccgactac	atcctctccc	agatcaagcc	cctgcacac	tctctcacc	agctccagcc	3060
gttctcgaag	ctgaagcaga	agcaggcgaa	cctcgtcctg	aagatcatcg	agcagctccc	3120
ctcgcccaag	gagtcgccg	acaagttcct	ggaggtgtgc	acgtgggtcg	accagatcgc	3180
ggccctcaac	gacagcaaga	cccgcaagac	gacctcgag	acggtgcggg	cggtcctgga	3240
ctccctccca	ggatccgtgg	gaggtctatc	gccatctcag	gcacccagcg	ccgcctcctc	3300
ggcttctca	agccgggtt	cagggtatctc	cgaagcactc	agagctggag	caactaagtc	3360
caaggaatc	ctgctctacc	tgcccggtt	cgtggacggc	gacggctcca	tcacgcggc	3420
gatcaagccg	aaccagctct	acaagttcaa	gcaccagctc	tccctgacct	tcaccgtgac	3480
ccagaagacg	cagaggcgct	gggtcctcga	caagctggtc	gacgagatcg	gggtgggcta	3540
cgtccgcgac	caggggtcgg	tgtccacta	ccagctctcc	cagatcaagc	ccctgcacaa	3600
cttctcacc	cagctccagc	cgttctcaca	gctgaagcag	aagcaggcga	acctcgtcct	3660
gaagatcatc	gagcagctcc	cctcgcccaa	ggagtcccg	gacaagtctc	tggaggtgtg	3720
caagtggttc	gacagcagc	cgccctcaca	cgacagcaag	acccgcaaga	cgacctcgga	3780
cagcgtgcgg	gcgggtctag	actccctcag	cgaagaagag	aagtcgtccc	cctgagggtac	3840
cacatggtta	acctagactt	gtccatcttc	tggattggcc	aacttaatta	atgtatgaaa	3900
taaaaggatg	cacacatagt	gacatgctaa	tcactataat	gtgggcatca	aagttgtgtg	3960
tatgtgttaa	ttactagtta	tctgaataaa	agagaaagag	atcatccata	tttcttatcc	4020
taaatgaatg	tcacgtgtct	ttataattct	ttgatgaacc	agatgcattt	cattaacca	4080
atccatatac	atataaatat	taatcatata	taattaatat	caattgggtt	agcaaaaaa	4140
atctagtcta	gggtgtgttt	gcgaatgcgg	ccatcggaac	gattaaactt	taattcggtc	4200
cgataaactc	gtatagcata	cattatacga	agttataacct	gggtggcgta	ctttccccc	4260
tatttttctc	cttatttttt	catctccccc	agcgggtccc	cctaaatact	cctatatacc	4320
ccaatacaac	tataaaatat	cattttctat	atcaactatc	aattttttat	ctactaaca	4380
ttactcgtgg	accacatbca	caatgtttag	gggtgatgaac	agtgacacgc	tagatctgag	4440
gggagagaga	aaagggtcgg	cgcgtagggg	gcgctgtagg	gggcaccgct	gcggctgtgg	4500
agtgccccc	acagccccc	tgcaagggga	gggggatact	gagggggctg	cgttgcgtac	4560
agcctgacag	gctctccttc	gcatttgccg	gggacagaaa	tgacttgccg	aggatggaag	4620
cagagagacg	gatttgccg	agcgcacagc	agctcgccaa	agacggcgct	gaagcagcag	4680
tgaccgcggg	cgagtgaagg	agtcacccct	gattcgcggt	ttatcgactc	ggcagcgggg	4740
caaccatggc	gttgaaggta	ggcaacatga	ggagccatcg	attgacacgc	gtcttcggaa	4800
tcggcggaac	tgacagatgg	tgacaaggag	gagggccacga	agcgtcgtcg	agcagagcgc	4860
gacaagcaaa	tcagagtcgg	cacgagcgtg	gatttggatc	tgacccccaa	gtttttgtat	4920
ggatcctatt	ccccaaattt	tagatcttca	atttctttac	tttaattttc	catagcacaa	4980
acgatgtttg	catgcacgat	tcggacaatc	ttgacttgtt	cgtccacggt	tggagtttag	5040
gggttggatg	tgtaaaacac	gtgataaaat	gtgtacaact	cgagaactag	ataatcatt	5100
ttggattgta	atatgtgtac	ctcatgctat	agttttggtt	aaatctgacg	tgaaggggcg	5160
aattcgccgc	tagcctgcag	tcgagcgtga	cccggtcgtg	ccctctctca	gagataatga	5220
gcattgcatg	tctaagttat	aaaaaattac	cacatatttt	ttttgtcaca	cttgtttgaa	5280
gtgcagttta	tctatcttta	tacatatatt	taaaactttac	tctacgaata	atataatcta	5340
tagtactaca	ataatatcag	tgttttagag	aatcatataa	atgaacagtt	agacatggct	5400
taaaggacaa	ttgagtattt	tgacaacagg	actctacagt	tttatctttt	tagtgtgcat	5460
gtgttctcct	tttttttttc	aaatagcttc	acctatataa	tacttcatcc	attttattag	5520
tacatccatt	taggggttag	gggttaatgg	ttttatagac	taattttttt	agtacatcta	5580
ttttattcta	tttttagctc	taaaatga	aaactaaaac	tctatttttag	tttttttatt	5640
taataattta	gatataaaat	agaataaaat	aaagtgaacta	aaaattaaac	aaataccctt	5700
taagaaatta	aaaaaactaa	ggaacatttt	ttctgtgttc	gagtagataa	tgccagcctg	5760
ttaaacgcgc	tcgacgagtc	taacggacac	caaccagcga	accagcagcg	tcgctgcggg	5820
ccaagcgaag	cagacggcac	ggcatctctg	tcgctgcctc	tggaccctcc	tcgagagttc	5880
cgtccacccg	ttggacttgc	tcgctgtcgc	gcacccagaa	attgcgtggc	ggagcgggag	5940
acgtgagccg	gcacggcagg	cggcctcctc	ctcctctcac	ggcaccggca	gctacggggg	6000
attcctttcc	cacgcgtctc	tcgctttccc	ttcctcggcc	gccgtaataa	atagacaccc	6060
ctctcacacc	ctctttcccc	aacctcgtgt	tgttcggagc	gcacacacac	acaaccagat	6120
ctcccccaaa	tcaccccgtc	ggcacctccg	cttcaaggta	gcgcgctcgt	cctccccccc	6180
ccccctctct	acctctctca	gtcggcggtt	ccggtccatg	catgggttagg	gcccggtagt	6240
tctactctcg	tctacttttg	tgttagatcc	gtgtttgtgt	tagatccgtg	ctgctagcgt	6300
tcgtacacgg	atgcgacctg	tacgtcagac	acgttctgat	tgctaaacttg	ccagtgtttc	6360
tctttgggga	atcctgggat	ggctctagcc	gttccgcaga	cgggatcgat	ttcatgattt	6420
tttttgttcc	gttgcatagg	gtttgggttg	cccttttctc	ttatttcaat	atatgccgtg	6480
caactgtttg	tcgggtcacc	tttctatgct	tttttttgtc	ttggttgtga	tgatgtggtc	6540
tggttgggag	gtcgttctag	atcggagtag	aattctgttt	caaactacct	gggtgattta	6600

-continued

ttaatttttg	atctgtatgt	gtgtgccata	catattcata	gttaacgaatt	gaagatgatg	6660
gatggaaata	tcgatctagg	ataggtatac	atgttgatgc	gggttttact	gatgcataata	6720
cagagatgct	ttttgttcgc	ttggttgtga	tgtatgtgtg	tggttgggcg	gtcgttcatt	6780
cgttctagat	cggagtagaa	tactgtttca	aactacctgg	tgtatttatt	aattttggaa	6840
ctgtatgtgt	gtgtcataca	tcttcatagt	tacgagtta	agatggatgg	aaatatcgat	6900
ctaggatagg	tatacatggt	gatgtgggtt	ttactgatgc	atatacatga	tggcatatgc	6960
agcatctatt	catatgctct	aaccttgagt	acctatctat	tataataaac	aagtatgttt	7020
tataattatt	ttgatcttga	tatacttga	tgtatggcata	tgcagcagct	atatgtggat	7080
tttttttagc	ctgccttcat	acgctattta	tttgcttgg	actgtttctt	ttgtcogatg	7140
tcacctgttt	gtttgggtgt	acttctgcag	gtcgactcta	gaggatcaat	tcgctagcga	7200
agttccctatt	ccgaagtctc	tattctctag	aaagtatagg	aacttcagat	ccaccgggat	7260
ccacacgaca	ccatgtcccc	cgaagccgcg	cccgtcgaga	tcggcccgcc	ccaccgccc	7320
gacatggccg	ccgtgtgcga	catcgtgaac	cactacatcg	agacctccac	cgtgaacttc	7380
cgcacccgagc	cgcagacccc	gcaggagtg	atcgacgacc	tggagcgctc	ccaggaccgc	7440
taccctgtgg	tcgtggccga	gggtggagggc	gtgggtggccg	gcacgcgcta	cgccggcccg	7500
tggaggccgc	gcaacgccta	cgactggacc	gtggagtcca	ccgtgtacgt	gtcccaccgc	7560
caaccagccg	tcggcctcgg	ctccaccctc	tacaccccac	tcctcaagag	catggaggcc	7620
cagggcttca	agtcgctggt	ggcctgatc	ggcctccgca	acgacccgct	cgtgcgctc	7680
caagaggccc	tcggctacac	cgcccgccgc	acctccg	cgccgggcta	caagcacggc	7740
ggctggcacg	acgtcgctt	ctggcagcg	gacttcgagc	tgcggccccc	gcccgcggcg	7800
gtgcgcccgg	tgacgcagat	ctgagtcgaa	acctagactt	gtccatcttc	tggattggcc	7860
aaactaatta	atgtatgaaa	taaaaggatg	cacacatagt	gacatgctaa	tcactataat	7920
gtgggcatca	aagttgtgtg	ttatgtgtaa	ttactagtta	tcgaataaaa	agagaagag	7980
atcatccata	tttcttacc	taaatgaatg	tcacgtgtct	ttataattct	ttgatgaacc	8040
agatgcattt	cattaaccaa	atccatatac	atataaatat	taatcatata	taattaatat	8100
caattgggtt	agcaaaacaa	atctagtcta	gggtgtgttt	gcgaatgcgg	ccctagcgta	8160
tacgaagtcc	ctattccgaa	gttcctattc	tcagaaagt	ataggaaactt	ctgtacacct	8220
gagctgattc	catgacttc	gtaggttcct	agctcaagcc	gctcgtgtcc	aagcgtcact	8280
tacgattagc	taagtattac	ggcatctagg	accgactagc	taactaacta	gtacaattcg	8340
cccttgtgaa	tctgtttgga	attgaaaaac	aagtgtctcc	ttttatacac	cactatgtcg	8400
cttcaatgtt	tgcgaaccaa	ggtaaagaaa	tgtaaaatct	tacaatttcc	gtgcaccga	8460
cataaatctg	tggtcacata	gctattgtta	aacggttgca	aatcctaagg	aggaccatta	8520
ttgtgcaaca	actacatag	gtagaagcgc	ttgttttgat	gtgtgcacat	tttgttgcta	8580
aaaggatcac	gatgcccaag	aggggggtga	attgggcttt	tcaaaaaatc	aacactaatt	8640
aaaaacctag	caagagccca	acttcacccc	gacaactagc	aataagagaa	tatgaagggg	8700
aaataggatc	aaaccttttc	ctaaatgatt	ttgggtgttg	aattgcccac	cacaaataat	8760
tggactaact	agtttgtctc	agatcataca	ttctacaggt	gccaaagggt	caacacaaac	8820
caatcaaaag	aacaaagttag	gcttcaaaag	aaaggagcaa	aaaggaaacc	gaagtgtgcc	8880
tggctcggcg	caccgggctg	tcgggtgtgc	caccagacag	tgtccgggtg	accagggtga	8940
atcagctcaa	gctcctcaac	ttcgggtttc	ccagacgcag	ctccactata	attcatttga	9000
ctgtccgggtg	caccgcgaga	gcaacgggta	cttgccgcga	acgggtcgact	ctgcaaaagt	9060
aacagtgcac	ttcagaagtc	agagcagatg	gtcagagggg	caccggattg	tcgggtgtag	9120
caccggagat	tgtcgtgcgc	catgaggaca	aagcctccaa	cgttcgacca	gctccaagcc	9180
ctaaactaaa	gatgacgtgg	cggcgacacc	gacactgtcc	gggtgtgtcac	cggactgttc	9240
gggtgcgcca	tcgcccagtg	ccttctccaa	cgggtacaat	ttggttggtg	gctataaata	9300
ccaccccaac	cggccacttt	aggggtgtgg	agcccaagca	acattccaag	tcataatagt	9360
gacatattca	agccatccca	accaccgtag	aattaattca	ttccgattaa	tcgtggcctc	9420
ttgctcttca	ggatgaagag	ctatgtttaa	acgtgcaagc	gctactagac	aattcagtag	9480
attaaaaacg	tcggcaactgt	gttattaagt	tgtctaagcg	tcaatttgtt	tacaccacaa	9540
tatatcctgc	caccagccag	ccaacagctc	cccgaccggc	agctcggcac	aaaatcacca	9600
ctcgatacag	gcagcccatc	agtcggggac	ggcgtcagcg	ggagagccgt	tgtgaaggcg	9660
cagactttgc	tcattgttacc	gaggtctattc	ggaagaacgg	caactaagct	gcccgggtttg	9720
aaacacggat	gatctcggcg	agggtagcat	gttgattgta	acgatgacag	agcgttgctg	9780
cctgtgatca	aatatcatct	ccctcgcaga	gatccgaatt	atcagccttc	ttattcattt	9840
ctcgcctaac	cgtgacaggc	gtcagatctt	gagaactatg	ccgacataat	aggaaatcgc	9900
tggataaaag	cgtgagggaa	gctgagtggt	gctatttctt	tagaagtga	cgttgacgat	9960
cgtcgaccgt	accccgatga	attaattcgg	acgtacgttc	tgaacacagc	tggatactta	10020
cttggggcat	tgtcatatcat	gacatcaaca	atgtaccctg	ttgtgtaacc	gtctcttgga	10080
ggttcgtatg	acactagtgg	ttccctcag	cttgcgacta	gatgttgagg	cctaaccattt	10140
tattagagag	caggctagtt	gcttagatac	atgatcttca	ggccgttatc	tgtcagggca	10200
agcgaaaatt	ggccatttat	gacgaccaat	gcccgcgaga	agctcccatc	tttgcgcgca	10260
tagacgcgcg	gccccctttt	tgggggttag	aacatccttt	tgcagatgt	ggaaaagaag	10320
ttcgttgtcc	cattgttggt	aatgacgtag	tagccggcga	aagtgcgaga	cccatattgcg	10380
ctatatataa	gcctacgatt	tcggttgcca	ctattgtcgt	aattggatga	actattatcg	10440
tagttgctct	cagagttgtc	gtaatttgat	ggactattgt	cgttaattgct	tatggagttg	10500
tcgtagttgc	ttggagaaat	gtcgtagttg	gatggggagt	agtcataggg	aagacgagct	10560
tcacccacta	aaacaattgg	cagggtcagca	agtgctcgtc	ccgatgccat	cgcgaagtacg	10620
aggcttagaa	ccaccttcaa	cagatcgccg	atagtccttc	ccagctctct	aacgcttgag	10680
ttaagccgcg	ccgcgaagcg	gcgtcggctt	gaacgaattg	ttagacatta	tttgcggact	10740
accttggtga	tctgcgcttt	cacgtagtga	acaaattctt	ccaactgatc	tgcgcgcgag	10800
gccaaagcat	ctcttggttc	aagataagcc	tgctagctt	caagtatgac	gggctgatac	10860
tgggcgggca	ggcgctccat	tgcccagtcg	gcagcgacat	ccttcggcgc	gatttttgccg	10920
gttactgcgc	tgtacaaat	gcgggacaac	gtaagcacta	catttcgctc	atcgccagcc	10980
cagtcggggc	gcgagtcca	tagcgttaag	gtttcattta	gcgcctcaaa	tagatcctgt	11040
tcaggaaccg	tcgcaaacg	ttcctccg	gctggaccta	ccaaggcaac	gctatgttct	11100
cttgcctttg	tcagcaagat	agccagatca	atgtcgatcg	tggtcggctc	gaagatacct	11160

-continued

gcaagaatgt	cattgcgctg	ccattctcca	aattgcagtt	cgcgcttagc	tggataacgc	11220
cacggaatga	tgtcgtcgtg	cacaacaatg	gtgactttcta	cagcgcgagg	aatctcgctc	11280
tctccagggg	aagccgaagt	ttccaaaagg	tcgttgatca	aagctcgccg	cgttggttca	11340
tcaagcctta	cagtcaccgt	aaccagcaaa	tcaatcac	tgtgtggctt	caggccgcca	11400
tccactgcgg	agccgtacaa	atgtacggcc	agcaacgtcg	gttcgagatg	gcgctcgatg	11460
acgccaaacta	cctctgatag	ttgagtcgat	acttcggcga	tcaccgcttc	cctcatgatg	11520
tttaactcct	gaattaagcc	gcgccgcgaa	gcggtgtcgg	cttgaatgaa	ttgttaggcg	11580
tcatcctgtg	ctcccgagaa	ccagtaccag	tacatcgctg	tttcggtcga	gacttgaggt	11640
ctagttttat	acgtgaacag	gtcaatgccg	ccgagagtaa	agccacattt	tgcgtacaaa	11700
ttgcaggcag	gtacattgtt	cgtttgtgtc	tctaactcgt	tgccaaggag	ctgtctgctt	11760
agtgccact	ttttcgcaaa	ttcgatgaga	ctgtgcgcga	ctcctttgcc	tcgggtcgctg	11820
tgcgacacaa	caatgtgttc	gatagaggct	agatcgttcc	atgttgagtt	gagttcaatc	11880
ttcccgacaa	gctcttggtc	gatgaatgcg	ccatagcaag	cagagttctc	atcagagtca	11940
tcatccgaga	tgtaatcctt	ccggtagggg	ctcacacttc	tggtagatag	ttcaaagcct	12000
tggtcgagata	ggtgcacatc	gaacacttca	cgaacaatga	aatggttctc	agcatccaat	12060
gtttccgcca	cctgctcagg	gatcccgaa	atcttcatat	gacgcctaac	gcctggcaca	12120
gcggatcgca	aacctgctgc	ggcttttggc	acaaaaggcg	tgacaggttt	gcgaatccgt	12180
tgctgccact	tgtaaacctt	tttgcagat	ttggttaacta	taatttatgt	tagaggcgaa	12240
gtcttgggta	aaaactggcc	tataattgct	ggggatttca	ggaaagttaa	catcaccttc	12300
cggctcgatg	tctattgtag	atatatgtag	tgtatctact	tgatcggggg	atctgctgcc	12360
tcgcgcgctt	cggtgatgac	ggtgaaaacc	tctgacacat	gcagctcccc	gagacgggtca	12420
cagcttgtct	gtaagcggat	gccgggagca	gacaagcccg	tcaggggcgg	tcagcgggtg	12480
ttggcgggtg	tcggggcgca	gccatgaccc	agtcacgtag	cgatagcgga	gtgtatactg	12540
gcttaactat	gcggcatcag	agcagattgt	actgagagtg	caccatatgc	ggtgtgaaat	12600
accgcacaga	gcgtgaagga	gaaaataccg	catcaggcgc	tcttcgcgtt	cctcgctcac	12660
tgactcgctg	cgtcggctgc	ttcggctcgc	gcgagcggt	tcagctcact	caaaaggcgt	12720
aatacgggta	tccacagaat	caggggataa	cgcaggaaaag	aacatgtgag	caaaaggcca	12780
gcaaaaggcc	aggaaccgta	aaaaggccgc	gttgctggcg	ttttccata	ggctccgccc	12840
ccctgacgag	catcacaaaa	atcgacgctc	aagtcagagg	tgggcgaaac	cgacaggact	12900
ataaagat	ac	caggcgcttc	ccctcggaag	ctccctcgctg	ctccgacct	12960
gcgcttacc	ggatctctgt	ccgcttctct	cccttcggga	agcgtggcgc	tttctcatag	13020
ctcacgctgt	aggtatctca	gttcgggtga	ggtcgttcgc	tccaagctgg	gctgtgtgca	13080
cgaaccccc	gttcagcccg	accgctgcgc	cttatccggt	aactatcgct	ttgagtccaa	13140
cccggtaa	gta	gcgcactggc	agcagccact	ggtaacagga	ttagcagagc	13200
gaggtatgta	ggcgggtgta	cagagttctt	gaagtggtgg	cctaactacg	gctacactag	13260
aaggacagta	tttggtatct	gcgctctgct	gaagccagtt	accttcggaa	aaagagttgg	13320
tagctcttga	tccggcaaac	aaaccaccgc	tggtagcggg	ggtttttttg	tttgcagaca	13380
gcagattacg	gcgagaaaaa	aaggatctca	agaagatcct	ttgatctttt	ctacgggggtc	13440
tgacgctcag	tggaacgaaa	actcacgtta	agggattttt	gtcatgagat	tatcaaaaag	13500
gatcttcacc	taggtctatt	tataattaaa	atgaagtttt	aaatcaatct	aaagtatata	13560
tgagtaaa	ct	gttaccatg	cttaactcgt	gaggcaccta	tctcagcgat	13620
ctgtctattt	cggtcaatcca	tagttgcctg	actcccgctc	gtgtagataa	ctacgatacg	13680
ggagggtctta	ccatctggcc	ccagtgctgc	aatgataccg	cgagaccac	gctcaccggc	13740
tccagattta	tcagcaataa	accagccagc	cggaaaggcc	gagcgagaa	gtggctctgc	13800
aactttatcc	gcctccatcc	agctctattaa	ttgttcggcg	gaagctagag	taagtgttcc	13860
gccaggtta	at	agtttgccga	cattgtctgca	gggggggggg	ggggggggga	13920
cttccattgt	tcattccacg	gacaaaaaca	gagaaaggaa	acgacagagg	ccaaaaagcc	13980
tcgctttcag	cacctgtcgt	ttcctttctt	ttcagagggg	attttaataa	aaaacattaa	14040
gttatgacga	agaagaacgc	aaaccgccta	ttttcataaa	tagcgaatac		14100
ccgcgaggtc	gccgccccgt	aacctgtcgg	atcacccgaa	aggacccgta	aatgtataat	14160
gattatcatc	tacatatcac	aacgtgcgtg	gaggccatca	aaccacgtca	aataatcaat	14220
tatgacgcag	gtatcgat	aattgatctg	catcaactta	acgtaaaaac	aacttcagac	14280
aatacaaatc	agcgacactg	aatacggggc	aacctcatgt	cccccccccc	ccccccccctg	14340
caggcatcgt	gggtgtcagc	tcgtcggtttg	gtatggcttc	attcagctcc	ggttcccaac	14400
gatcaaggcg	agttacatga	tcctccatgt	tgtgcaaaaa	agcgggttagc	tccttcggtc	14460
ctccgatcgt	tgtcagaagt	aagttggccg	cagtggtatc	actcatgggt	atggcagcac	14520
tgcataaattc	tcttactgtc	atgccatccg	taagatgctt	ttctgtgact	ggtagtagt	14580
caaccaagtc	attctgagaa	tagtgtatgc	ggcgaccgag	ttgctcttgc	ccggcgctca	14640
cacgggataa	taccgcgcc	catagcagaa	ctttaaaagt	gctcatcatt	ggaaaacggt	14700
cttcggggcg	aaaactctca	aggatcttac	cgtctgttag	atccagttcg	atgtaaccca	14760
ctcgtgcacc	caactgatct	tcagcatctt	ttactttcac	cagcgtttct	gggtgagcaa	14820
aaacaggaag	gcaaaatgcc	gcaaaaaagg	gaataaggcg	gacacggaaa	tggtgaatac	14880
tcatactctt	cctttttcaa	tattattgaa	gcatttatca	gggttatgtt	ctcatgagcg	14940
gatacatatt	tgaatgtatt	tagaaaaata	aacaaatagg	ggttccgcgc	acatttcccc	15000
gaaaagtgcc	acctgacgtc	taagaaacca	tattatcat	gacattaacc	tataaaaaata	15060
ggcgtatcac	gaggcccttt	cgtcttcaag	aattgggtcga	cgatcttgct	gcgttcggat	15120
attttcgtgg	agttcccgcc	acagaccggg	attgaaggcg	agatccagca	actcgcgcca	15180
gatcatcctg	tgacggaa	ttggcgcggt	atgactggcc	aggacgtcgg	ccgaaagagc	15240
gacaagcaga	tcacgctttt	cgacacgctc	ggatttgcca	tcgaggattt	ttcggcgctg	15300
cgctacgtcc	gcgaccgcgt	tgagggatca	agccacagca	gcccaactcg	ccttctagcc	15360
gaccagacg	agccaaggga	tcctttttgga	atgctgctcc	gtcgtcaggg	tttccgacgt	15420
ttgggtgggt	gacacagaat	cattatcgta	cggaaatgcca	agcactcccc	aggggaaccc	15480
tgtggttgcc	atgcacat	aaatggacga	acggataaac	cttttcacgc	ccttttaaat	15540
atccgttatt	ctaataaacg	ctctttttct	ttag			15574

SEQ ID NO: 105

moltype = DNA length = 15585

-continued

FEATURE	Location/Qualifiers
misc_feature	1..15585
	note = synthetic construct; plasmid PHP44779
source	1..15585
	mol_type = other DNA
	organism = synthetic construct

SEQUENCE: 105

```

gtttaccgcg caatatatcc tgtcaaacac tgatagtta aactgaaggg gggaaacgac 60
aatctgatca tgagcggaga attaagggag tcacgttatg acccccgccg atgacgcggg 120
acaagccgtt ttacgtttgg aactgacaga accgcaacgt tgaaggagcc actcagcaag 180
ctgggtacgat tgaatacga ctactatag ggcgaattga gcgctgttta aacgctcttc 240
aactggaaga gcggttacta ccggtctggat ggccggggcct tgatcgtgca ccgccggcgt 300
ccggactaac taactagtcg agctagtta cctatgaggt gacatgaagc gctcagcgtt 360
actatgacgg ttagtctcac gactgttggg ggcagttagc tacgacttag ctatagttcc 420
ggacttacc ttaagataac ttcgatagc atacattata cgaagttatg gggccaccgg 480
tggtaaccgag ctggttttaa cgctcttcaa ctggaagagc gggtaccaga gctggtcacc 540
ttgtccacc aagatggaac tggcgcgcct cattaattaa gtcagcgccc gctctagttg 600
aagacacggt catgtcttca tcgtaagaag acactcagta gtcttcggcc agaatggcca 660
tctggattca gcaggctcct aaggccattt aaatcctgag gatctggtct tctaaggac 720
ccgggataac ggaccgaagc ttgcatgcct gcagtgcagc gtgaccgggt cgtgcccctc 780
tctagagata atagcattg catgtctaag ttataaaaa ttaccacata ttttttttgt 840
cacacttggt tgaagtgcag ttatctatc ttataacata tatttaaaact ttactctacg 900
aataataata tctatagtag tacaataata tcagtgtttt agagaatcat ataaatgaac 960
agttagacat ggtctaaagg acaattgagt attttgacaa caggactcta cagtttttatc 1020
tttttagtgt ccatgtgttc tcttttttt ttgcaaatag cttcacctat ataactctc 1080
atccatttta ttagtacatc cattttaggt ttagggttaa tgggttttat agactaatat 1140
ttttagtaca tctattttat tctattttag cctctaaatt aagaaaaact aaactctatt 1200
ttagtttttt tatttaataa tttagatata aaatagaata aaataaagtg actaaaaatt 1260
aaacaaatcc cctttaagaa attaaaaaaa ctaagggaac atttttcttg ttccgagtag 1320
ataatgccag cctgttaaac gccgtcgacg agtctaaccg acaccaacca gcgaaccagc 1380
agcgtcgcgt cgggccaagc gaagcagacg gcacggcctc tctgtcgtg cctctgggac 1440
cctctcgaga gttccgctcc accgttggaac ttgctccgct gtcggcatcc agaaattgcy 1500
tggcggagcg gcagacgtga gccggcacgg caggcgccct cctcctcctc tcacggcacc 1560
ggcagctagc ggggactcct ttcccaccgc tccctcgctt tcccttcctc gcccgccgta 1620
ataaatagac accccctcca caccctcttt ccccaacctc gtgttggtcg gagcgacac 1680
acacacaccc agatctcccc caaatccacc cgtcggcacc tccgcttcaa ggtacgcgcg 1740
tcgtcctccc ccccccctct cctaccttc tctagatcgg cgttcgggtc catgcatggt 1800
tagggcccg tagttctact tctgttcctg tttgtgttag atccgtgttt gtgttagatc 1860
cgtgctgcta gcgttcgtac acggatgcga cctgtacgtc agacacgttc tgattgctaa 1920
cttgccagtg tttctctttg gggaatcctg ggatggctct agccgttcctg cagacgggat 1980
cgatttcatg attttttttg ttctgttgca tagggtttgg tttgcccctt tctcttattt 2040
caatatatgc cgtgcacttg ttgtcgggt catcttttca tgcttttttt tgtctgtggt 2100
gtgatgatgt ggtctggttg ggcggtcgtt ctagatcgga gtagaattct gtttcaaaact 2160
acctgggtga ttatttaatt ttggatctgt atgtgtgtgc catacatatt catagttacg 2220
aattgaagat gatggatgga aatctcgatc taggataggt atacatgttg atcggggttt 2280
tactgatgca tatacagaga tgctttttgt tcgcttggtt gtgatgatgt ggtgtgggtg 2340
ggcgggtcgt cattcgttct agatcggagt agaatactgt ttcaaaactc ctggtgtatt 2400
tattaatttt ggaactgtat gtgtgtgtca tacatcttca tagttacgag tttaagatgg 2460
atggaatatat cgatctagga taggtatata gtgtgatgtg ggttttactg atgcataac 2520
atgatggcat atgcagcatc tattcatatg ctctaacctt gagtacctat ctattataat 2580
aaacaagtat gttttataat tattttgatc ttgatatact tggatgatgg catatgcagc 2640
agctatatgt ggattttttt agcccctgct tcatacgcta tttatttctg tggtagctgt 2700
tcttttgtcg atgtccacc tgttgttttg tgttacttct gcaggctcag tctagaggat 2760
ccatggcacc gaagaagaag cgcaagggtc atatgaacac caagtacaac aaggagttcc 2820
tgctctacct ggcgggcttc gtggacggcg acggctccat catcgcgag atcaagccga 2880
accagtctca caagttcaag caccagctca tgctgacctt caccgtgacc cagaagacgc 2940
agaggcgctg gttcctcgac aagctgggtc acgagatcgg ggtgggcaag gtcgcgac 3000
gcgggtcggg gtcgactac atcctctccc agatcaagcc cctgcacaac ttctcacc 3060
agctccagcc gttcctcaag ctgaagcaga agcaggcgaa cctcgtcctg aagatcatcg 3120
agcagctccc ctccggcaag gagtccccgg acaagttcct ggaggtgtgc acgtgggtcg 3180
accagatcgc ggcctccaac gacagcaaga cccgcaagac gacctcggag acggtgcggg 3240
cggctcctga ctcccctcca ggtaccgtgg gaggtctatc gccatctcag gcatccagcg 3300
ccgcatctcc ggctctctca agcccgggtt cagggatctc cgaagcactc agagctggag 3360
caactaagtc caaggaattc ctgctctacc tggcgggctt cgtggacggc gacggctcca 3420
tcacgcggcg gatcaagccg aaccagtcct acaagttcaa gcaccagctc tccctgacct 3480
tcaccgtgac ccagaagacg cagaggcgct ggttcctcga caagctgggt gacgagatcg 3540
gggtgggcta cgtccgcgac caggggtcgg tgtccacta ccagctctcc cagatcaagc 3600
cctcgacaaa cttcctcacc cagctccagc cgttctcaa gctgaagcag aagcaggcga 3660
acctcgtcct gaagatcatc gacagctcc cctcgcccaa ggagtcccc gacaagttcc 3720
tggaggtgtg caggtgggtc gaccagatcg cggccctcaa cgacagcaag acccgcaaga 3780
cgactcggga cagcgtgcgg gcggttctag actccctcag cgagaagaag aagtcgtccc 3840
cctgaggtac cacatggtta acctagactt gtccatcttc tggattggcc aacttaatta 3900
atgtatgaaa taaaggatg cacacatagt gacatgctaa tcaactaat gtgggcatca 3960
aagttgtgtg ttatgtgtaa ttactagtta tctgaataaa agagaaaag atcatccata 4020
ttctttatcc taaatgaat tcacgtgtct ttataattct ttgatgaac agatgcattt 4080
cattaaccaa atccatatac atataaatat taatcatata taattaatat caattgggtt 4140

```


-continued

agcaaaacaa	atctagtcta	ggtgtgtttt	gcgaatgcgg	ccgccaccgc	ggtggagctc	4200
gaattccggt	ccgataactt	cgtatagcat	acattatacg	aagttatacc	tggtggcgctc	4260
actttccccc	ctatttttct	ccctattttt	tcctctcccg	cagcggttcc	ccctaataac	4320
tcctatatac	cccaatacaa	ctataaaata	tcattttcta	tatcaactat	caatttttta	4380
tctactaaca	attactcgtg	gacccacatc	acaatgttta	gggtgatgaa	cagtgacacg	4440
ctagatctga	ggggagagag	aaaagggtcg	gcgcgtaggg	ggcgtctgag	ggggcaccgc	4500
tgccgctgtg	gagtgccccc	tcacgcccc	atgcaagggg	aggggggatac	tgagggggct	4560
gcgttgcgta	cagctcgaca	ggctctcctt	cgcatttgcg	cgggacagaa	atgacttgcc	4620
gaggatggaa	gcagagagac	ggatttgccc	gagcgcacag	cagctcgcca	aagacggcgt	4680
cgaagcagca	gtgaccgcgg	tcgagtggag	gagtcaccc	ggattcgccg	tttatcgact	4740
cggcacgggg	gcaaccatgg	cgttgaaggt	aggcaacatg	aggagccatc	gattgacacc	4800
ggtcttcgga	atcggcggat	ctcgacgatg	gtgacaagga	ggagggccacg	aagcgtcgct	4860
gagcagagcg	cgcacaagca	atcgagtcgg	ccacgagcgt	ggatttggat	ctgaccccca	4920
agtttttcta	tggtatctat	tccccaattt	gtagatcttc	aatttccctta	ctttaatttt	4980
ccatagcaca	aacgatgttt	gcctgacaga	ttcggacaat	cttgacttgt	tcgtccacgg	5040
ttggagttta	gggttggaa	gtgtataaac	cgtgataaac	tgtgtacaac	tcgagaacta	5100
gataattcat	tttggattgt	aatatgtgta	cctcatgcta	tagttttggt	taaaactgac	5160
gtgaaagggc	gaattcgcgc	ctagcctgca	gtgcagcgtg	accgcgtcgt	gccccctctc	5220
agagataatg	agcattgcct	gtctaagtta	taaaaaatta	ccacatattt	tttttgtcac	5280
acttgtttga	agtgacagtt	atctatcttt	atacatatat	ttaaaacttta	ctctacgaat	5340
aatataatct	atagtactac	aataatatca	gtgtttttaga	gaatcatata	aatgaacagt	5400
tagacatgg	ctaaaggaca	attgagtatt	ttgacaacag	gactctacag	tttatctttt	5460
ttagtgtgca	tgtgttcttc	tttttttttg	caaatagcct	cacctatata	atacttcac	5520
catttttata	gtacatccat	ttagggttta	gggttaaatg	tttttataga	ctaatttttt	5580
tagtacctct	attttattct	attttagcct	ctaaattaag	aaaactaaaa	ctctatttta	5640
gtttttttat	ttaataattt	agataaaaa	tagaataaaa	taaagtgact	aaaaattaaa	5700
caaataccct	ttaagaaatt	aaaaaaacta	aggaaacatt	tttctgtttt	cgaatagata	5760
atgccagcct	gttaaacgcc	gtcgacgagt	ctaaccggca	ccaaccagcg	aaccagcagc	5820
gtccgctcgg	gccaaagcga	gcagacggca	cggcatctct	gtcgtctcct	ctggacccct	5880
ctcgagagtt	ccgctccacc	gttggaactg	ctccgctgtc	ggcatccaga	aatgtcggtg	5940
cggagcggca	gacgtgagcg	ggcacggcag	gcggcctcct	cctcctctca	cggcacggcg	6000
agctacgggg	gattcctttc	ccaccgtctc	ttcgttttcc	cttcctcgcc	cgcgtaata	6060
aatagacacc	ccctccacac	ccctcttccc	caacctcggt	ttgttcggag	cgcacacaca	6120
cacaaccaga	tttcccccaa	atccaccctg	cggcacctcc	gcttcaaggt	acgccgctcg	6180
tcctccccc	ccccctctc	taccttctct	agatcggcgt	tcgggtccat	gcattggttag	6240
ggcccggtag	ttctacttct	gttcatgttt	gtgttagatc	cgtgtttgtg	ttagatccgt	6300
gctgctagcg	ttcgtacaac	gattcgacct	gtacgtcaga	cacgttctga	ttgtcatact	6360
gccagtgttt	ctctttgggg	aatcctggga	tggctctagc	cgttcgcgag	acgggatcga	6420
tttcatgatt	ttttttgttt	cgttgcatag	ggtttgggtt	gccccttttc	tttatttcaa	6480
tatatgccgt	gcacttgttt	gtcgggtcat	cttttcatgc	ttttttttgt	cttgggtgtg	6540
atgatgtggt	ctgggtgggc	ggtcgtttct	gatcggagta	gaattctgtt	tcaaaactacc	6600
tggtgggatt	attaattttg	gatctgtatg	tgtgtgccat	acataatcat	agttacgaat	6660
tgaagatgat	ggatggaaat	atcgatctag	gataggtata	catgttgatg	cgggtttttac	6720
tgatgcata	acagagatgc	tttttgttcg	cttgggtgtg	atgatgtggt	gtgggtgggc	6780
ggtcgttcat	tcgttctaga	tcggagtaga	ataactgttc	aaactacctg	gtgtattttat	6840
taattttgga	actgtatgtg	tgtgtcatac	atcttcatag	ttacgagttt	aagatggatg	6900
gaaatatcga	tctaggatag	gtatacatgt	tgatgtgggt	tttactgatg	catatacatg	6960
atggcatatg	cagcatctat	tcatatgtct	taaccttgag	tacatctcta	ttataataaa	7020
caagtatgtt	ttataattat	tttgatcttg	atatacttgg	atgatggcat	atgcagcagc	7080
tatatgtgga	tttttttagc	cctgccttca	tacgtatatt	atttgcttgg	tactgtttct	7140
tttgtcgatg	ctcaccctgt	tgtttgggtg	tactctcgca	ggtcgactct	agaggatcaa	7200
ttcgtatagc	aagttcctat	tcogaagttc	ctattctcta	gaaagtatag	gaacttcaga	7260
tcacccggga	tcacacgcac	accatgtccc	ccgagcgcgg	cccgcgtcag	atccgcccgg	7320
ccaccgcgcg	cgacatggcc	gccgtgtgcg	acatcgtgaa	ccactacatc	gagacctcca	7380
ccgtgaactt	ccgcaccgag	cgcagacacc	cgcaggagtg	gatcgacgac	ctggagcgcc	7440
tcaggagacc	ctaccctgtg	ctcgtggccg	aggtggaggg	cgtgggtggc	ggcatcgctc	7500
acgccggccc	gtggaaggcc	cgcacacgct	acgactggac	cgtggagtcc	accgtgtacg	7560
tgtcccaccg	ccaccagcgc	ctcggcctcg	gctccaccct	ctacacccac	ctcctcaaga	7620
gcattggagg	ccagggtctc	aagtccgtgg	tggcgtgatg	cggcctcccc	aacgacccgt	7680
ccgtgcgcct	accagagggc	ctcggctaca	ccgccccggg	cacctccgcg	gccgcgggct	7740
acaagcaccg	cggctggcac	cagctcggct	ctcggcagcg	cgaacttcgag	ctgccggccc	7800
cgcgcgcgcc	gggtgcgcgc	gtgacgcaga	tctgagtcga	aaactagact	tgtccatctt	7860
ctggattggc	caacttaatt	aatgtatgaa	ataaaaggat	gcacacatag	tgacatgcta	7920
atcactataa	tgtgggcatc	aaagtgtgtg	gttatgtgta	attactagtt	atctgaataa	7980
aagagaaaga	gatcatccat	atttcttatc	ctaaatgaat	gtcacgtgtc	tttataaatt	8040
tttgatgaac	cagatgcatt	tcattaacca	aatccatata	catataaata	ttaatcatat	8100
ataattaata	tcaattgggt	tagcaaaaac	aatctagtct	aggtgtgttt	tgcaaatgcg	8160
gccctagcgt	atacgaagtt	cctattccga	agttcctatt	ctccagaag	tataggaact	8220
tctgtacacc	tgagctgatt	ccgatgactt	cgtagggtcc	tagctcaagc	cgtcgtgtgc	8280
caagcgtcac	ttacgattag	ctaagtatta	cggcatctag	gaccgactag	ctaactaact	8340
agtaacaatt	gccccttgtga	atctgtttgg	aattgaaaaa	caagtgtctt	cttttataca	8400
ccactatgtc	gcttcaatgt	ttgcgaacca	aggtaaagaa	atgtaaaaat	ttacaatttc	8460
cgtgcatccg	acataaatct	gtggtcacat	agctattgtt	aaacgggtgc	aaatcctaag	8520
gaggaccatt	atttgtcaac	aactacatat	ggtagaagcg	cttgttttga	tgtgtgcaca	8580
ttttgtgtct	aaaaggatca	cgatgcccac	gaggggggtg	aattgggctt	ttctaaaaat	8640
caacactaat	taaaacctaa	gcaagagccc	aacttcaccc	cgacaactag	caataagaga	8700

-continued

atatgaaagg	gaaataggat	caaacccttt	cctaaatgat	tttgggtggt	gaattgcccc	8760
acacaaataa	ttggactaac	tagtttgctc	tagatcatat	attctacagg	tgccaaagg	8820
tcaacacaaa	ccaatcaaaa	gaacaagtta	ggcttcaaaa	gaaaggagca	aaaaggaaac	8880
cgaagtgtgc	ctggtctggc	gcaccgggct	gtccgggtgtg	ccaccagaca	gtgtccgggtg	8940
caccaggggtg	aatcagctca	agctcctcaa	cttcgggttt	cccagacgca	gtccactat	9000
aattcattgg	actgtccggt	gcaccgcgag	agcaacggct	acttgccgcg	aacggtcgac	9060
tctgcaaagt	gaacagtgc	atcagaagt	cagagcagat	ggtcagaggg	gcaccggatt	9120
gtccgggtgt	gcaccggact	gtccgggtgc	gcagaggac	aaagcctcca	acggtcgacc	9180
agctccaagc	cctaactaca	agatgacgtg	gcggcgaccc	ggacactgtc	cgggtgtgtg	9240
ccggactgtt	cgggtgcgcc	atcgccagta	gccttctcca	acggctacaa	tttgggttgt	9300
ggctataaat	accaccccaa	ccggccactt	taagggtgtg	gagcccaagc	aacattccaa	9360
gtcatatagt	tgacatatct	aagccatccc	aaccaccgta	gaattaattc	attccgatta	9420
atcgtggcct	cttgctcttc	aggatgaaga	gctatgttta	aacgtgcaag	cgctactaga	9480
caattcagta	cattaaaaac	gtccgcaatg	tggtattaa	ttgtctaaag	gtcaatttgt	9540
ttacaccaca	atatatcctg	ccaccagcca	gccaacagct	ccccgaccgg	cagctcggca	9600
caaaatcacc	actcgataca	ggcagcccat	cagtcgggga	cggcgctcagc	gggagagccg	9660
ttgtaaaggc	gcagactttg	ctcatgttac	cgatgctatt	cgaagaacg	gcaactaagc	9720
tgccgggttt	gaaacacgga	tgatctcgcg	gagggtagca	tggtgatgtg	aacgatgaca	9780
gagcgttgct	cctgtgtatc	aaatatcatc	tccctcgag	agatccgaat	tatcagcctt	9840
cttattcatt	gtctcgctaa	ccgtgacagg	ctgtcgatct	tgagaactat	gccgacataa	9900
taggaaatcg	ctggataaag	ccgctgagga	agctgagtg	cgctatttct	ttagaagtga	9960
acggtgacga	tcgtcgaccg	taccccgatg	aattaattcg	gacgtacgtt	ctgaacacag	10020
ctggataact	acttgggcga	ttgtcataca	tgacatcaac	aatgtacccg	tttgtgtaac	10080
cgctctcttg	aggttcgtat	gacactagt	gttccctcca	gcttgcgact	agatgttgag	10140
gcctaacatt	tattagaga	gcaggctagt	tgcttagata	catgatcttc	aggccgttat	10200
ctgtcagggc	aagcgaaaa	tgcccatcta	tgacgaccaa	tgcccccgag	aagctcccat	10260
ctttgccgcc	atagacgcgc	cgccccctt	ttgggtgtga	gaacatcctt	ttgccagatg	10320
tggaaaagaa	gttcgtgtgc	ccattgtttg	caatgacgta	gtagccggcg	aaagtgcgag	10380
acccatttgc	gtcatatata	agcctacgat	ttccgttgcg	actattgtcg	taattggatg	10440
aactattatc	gtagtgtgtc	tcagagttgt	cgtaatttga	tggaactattg	tcgtaattgc	10500
ttatggagtt	gtcgtagtgc	cttggagaaa	gttcgtagt	ggatggggag	tagtcatagg	10560
gaagacgcag	ttcatccact	aaaacaattg	gcaggtcagc	aagtgcctgc	cccgatgcca	10620
tcgcaagtac	gaggtctaga	accaccttca	acagatcgcg	catagcttcc	cccagctctc	10680
taacgcttga	tttaagccgc	gcgcggaagc	ggcgtcggt	tgaacgaatt	gttagacatt	10740
atlttgcgcg	taccttggtg	atctcgctt	tcacgtagt	aacaaattct	tccaactgat	10800
ctgcgcgcga	ggccaagcga	tcttcttgtc	caagataaag	ctgcctagct	tcaagtatga	10860
cgggctgata	ctgggcccgc	aggcgctcca	ttgcccaagc	ggcagcgaca	tccttcggcg	10920
cgattttgcc	ggttactcgc	ctgtaccaaa	tgccgggcaa	cgtaaagcact	acatttcgct	10980
catcgccacg	ccagtcgggc	ggcgagttcc	atagcgtaa	ggtttcattt	agcgcctcaa	11040
atagatcctg	ttcaggaacc	ggatcaaa	gttctctcgc	cgttggaact	accaaggcaa	11100
cgctatgttc	tcttgccttt	gtcagcaaga	tagccagatc	aatgtcgatc	gtggctggct	11160
cgaagatacc	tgcaagaatg	tcattgcgct	gccattctcc	aaattgcagt	tcgcgcttag	11220
ctggataacg	ccacggaatg	atgtcgtcgt	gcacaacaat	ggtgacttct	acagcgcgga	11280
gaatctcgct	ctctccagg	gaagccgaag	tttccaaaag	gtcgttgatc	aaagctcgcc	11340
gcgttggttc	atcaagcctt	acagtcaccg	taaccagcaa	atcaatatca	ctgtgtggct	11400
tcaggccgcg	atccactcgc	gagccgtaca	aatgtacggc	cagcaacgct	ggttcgagat	11460
ggcgtcgat	gacgccaa	acctctgata	gttgagtcga	tacttcggcg	atcacgcctt	11520
ccctcatgat	gtttaaactcc	tgaattaagc	cgccgcgcga	agcgggtgtg	gcttgaatga	11580
attgttaggc	gtcatcctgt	gctcccgaga	accagtagca	gtacatcgct	gttctgttgc	11640
agacttgagg	tctagtttta	tacgtgaaca	ggcgaatgcc	cccagagata	aagccacatt	11700
ttgcgtacaa	attgcaggca	ggtaacattgt	tcgtttgtgt	ctctaactgt	atgccaaagga	11760
gctgtctgct	tagtgccca	tttttcgcaa	attcgatgag	actgtgcgcg	actcctttgc	11820
ctcggtgcgt	gtgcgacaca	acaatgtgtt	cgatagagcg	tagatcgctt	catgttgagt	11880
tgagttcaat	cttcccgaca	agctcttggt	cgatgaatgc	gccatagcaa	gcagagttct	11940
catcagagtc	atcatccgag	atgtaactct	tcgggtaggg	gctcacactt	ctggtagata	12000
gttcaaaagc	ttggtccgat	aggtgcacat	cgaacacttc	acgaacaatg	aaatggttct	12060
cagcatccaa	tggttccgcg	acctgtcag	ggatcaccca	aatcttcata	tgacgcctaa	12120
cgccgtggac	agcggatcgc	aaacctggcg	cggtttttgg	cacaaaaggc	gtgacagggt	12180
tgccgaatccg	ttgctgccac	ttgttaaccc	ttttgccaga	tttggttaact	ataatttatg	12240
ttagaggcga	agtcttggtg	aaaaactggc	ctaaaattgc	tggggatttc	aggaaagtaa	12300
acatcacctt	ccggtcgat	gtctattgta	gatataatga	gtgtatctac	ttgatcgggg	12360
gatctgctgc	ctcgcgcgtt	tcgggtgatga	cgggtgaaac	ctctgacaca	tgacagctccc	12420
ggagacggtc	acagcttgtc	tgtaaagcga	tgccggggagc	agacaagccc	gtcagggcgc	12480
gtcagcgggt	gttggcgggt	gtcggggcgc	agccatgacc	cagtcacgta	gcgatagcgg	12540
aggtataact	ggcttaacta	tgccgcatca	gagcagattg	tactgagagt	gcacatata	12600
cgggtgtgaa	taccgcacag	atgcgttaag	agaaaaatcc	gcacaggcgc	ctctccgct	12660
tcctcgctca	ctgactcgct	gcgctcggtc	gttcggctgc	ggcgagcggg	atcagctcac	12720
tcaaggcggg	taatacgggt	atccacagaa	tcaggggata	acgcaggaaa	gaacatgtga	12780
gcaaaaggcc	agcaaaaggc	cagggaaccgt	aaaaaggccg	cgtgtgctggc	gtttttccat	12840
agggtccgcc	ccctcgacga	gcatacacia	aatcgacgct	caagtacagag	gtggcgaaac	12900
ccgacaggac	tataaagata	ccaggcggtt	ccccctggaa	gctccctcgt	gcgctctcct	12960
gttccgaccc	tgccgcttac	cggtatacctg	tcgccttttc	tccttcggg	aagcgtggcg	13020
ctttctcata	gctcacgctg	taggtatctc	agttcgggtg	aggtcgttcg	ctccaaagctg	13080
ggctgtgtgc	acgaaccccc	cgttcagccc	gaccgctgcg	ccttatccgg	taactatcgt	13140
cttgagttca	accggttaag	acacgactta	tcgcccactgg	cagcagccac	tggttaacagg	13200
attagcagag	cgaggatgtg	aggcgggtgct	acagagttct	tgaagtggtg	gcctaactac	13260

-continued

```

ggctacacta gaaggacagt atttgggtatc tgcgctctgc tgaagccagt taccttcgga 13320
aaaaagagttg gtagctcttg atccggcaaaa caaacccaccg ctggtagcgg tggttttttt 13380
gttttcaagc agcagattac gcgcagaaaa aaaggatctc aagaagatcc tttgatcttt 13440
tctacgggggt ctgacgctca gtggaacgaa aactcacgtt aagggaattt ggtcatgaga 13500
ttatcaaaaa ggatcttcac ctagatcctt ttaaattaaa aatgaagttt taaatcaatc 13560
taaagtatat atgagtaaac ttggtctgac agttaccaat gcttaatcag tgaggcacct 13620
atctcagcga tctgtctatt cgtttcatcc atagttgcct gactccccgt cgtgtagata 13680
actacgatac gggaggggctt accatctggc cccagtgcgt caatgatacc gcgagaccca 13740
cgctcaccgg ctccagattt atcagcaata aaccagccag ccggaagggc cgagcgacga 13800
agtggtcctg caactttatc cgctccatc cagtctatta attgttgccg ggaagctaga 13860
gtaagtatgt cgccagttaa tagtttgccg aacgttggtg ccattgctgc aggggggggg 13920
gggggggggg acttccattg ttcatccac ggacaaaaac agagaaaagg aacgacagag 13980
gccccaaagc ctgcgtttca gcacctgtcg ttctctttct ttccagaggg tattttaaat 14040
aaaaacatta agttatgacg aagaagaacg gaaacgcctt aaaccggaaa attttcataa 14100
atagcgaaaa cccgcgaggt cgccgccccg taacctgtcg gatcaccgga aaggacccgt 14160
aaagtgtata tgattatcat ctacatatca caacgtgcgt ggaggccatc aaaccacgtc 14220
aaataatcaa ttatgacgca ggtatcgat taattgatct gcatcaactt aacgtaaaaa 14280
caacttcaga caatacaaat cagcgacact gaatacgggg caacctcatg tcccccccc 14340
ccccccctc gcagggcatc tgggtgtcacg ctcgctggtt ggtagggctt cattcagctc 14400
cgggtcccaa cgcataaggg gagttacatg atccccatg ttgtgcaaaa aagcggttag 14460
ctccttcggt cctccgatcg ttgtcagaag taagttggcc gcagtgttat cactcatggt 14520
tatggcagca ctgcataatt ctcttactgt catgccatcc gtaagatgct tttctgtgac 14580
tgggtgagtag tcaaccaagt cattctgaga atagtgtatg cggcgacgga gttgctcttg 14640
cccgcgctca acacgggata ataccgcgcc acatagcaga actttaaag tgctcatcat 14700
tggaaaacgt tcttcggggc gaaaaactctc aaggatctta ccgctgttga gatccagttc 14760
gatgtaaccc actcgtgcac ccaactgatc ttcagcatct tttactttca ccagcgtttc 14820
tgggtgagca aaaacaggaa ggcaaaatgc cgcaaaaaag ggaataaggg cgacacggaa 14880
atggtgaata ctcatactct tcctttttca atattattga agcatttatc aggggtattg 14940
tctcatgagc ggatacatat ttgaatgtat ttagaaaaat aaacaaatag gggttcccg 15000
cacattttcc cgaaaagtgc cactgacgt ctaagaaacc attattatca tgacattaac 15060
ctataaaaaat aggcgtatca gagggccctt tcgtcttcaa gaattggtcg acgatcttgc 15120
tgcgttcgga tattttcgtg gagttcccg caccagaccg gattgaaggc gagatccagc 15180
aactcgcgcc agatcatcct gtgacggaac ttggcgcggt gatgactggc caggacgtcg 15240
gccgaaagag gcacaagcag atcacgcttt tcgacagcgt cggatttgcg atcgaggatt 15300
tttcggcgct gcgctacgtc cgcgaccgcg ttgagggatc aagccacagc agccactcg 15360
accttctagc cgacccagac ggcaccaagg atctttttgg aatgctgctc cgtcgtcagg 15420
ctttccgacg tttgggtggt tgaacagaag tcattatcgt acggaatgcc aagcactccc 15480
gagggggaacc ctggtggttg catgcacata caaatggacg aacggataaa ccttttcacg 15540
cccttttaaa tatcgttat tctaataaac gctcttttct cttag 15585

```

```

SEQ ID NO: 106      moltype = DNA length = 18
FEATURE
misc_feature       Location/Qualifiers
                    1..18
                    note = synthetic construct; MHP14TS probe
source             1..18
                    mol_type = other DNA
                    organism = synthetic construct

SEQUENCE: 106
cagattcacg tcagattt 18

SEQ ID NO: 107      moltype = DNA length = 26
FEATURE
misc_feature       Location/Qualifiers
                    1..26
                    note = synthetic construct; MHPTS14_Forward_MGB primer
source             1..26
                    mol_type = other DNA
                    organism = synthetic construct

SEQUENCE: 107
agcgacatag tggtgtataa aaggaa 26

SEQ ID NO: 108      moltype = DNA length = 27
FEATURE
misc_feature       Location/Qualifiers
                    1..27
                    note = synthetic construct; MHPTS14_Reverse_MGB primer
source             1..27
                    mol_type = other DNA
                    organism = synthetic construct

SEQUENCE: 108
tggattgtaa tatgtgtacc tcatgct 27

SEQ ID NO: 109      moltype = DNA length = 27
FEATURE
misc_feature       Location/Qualifiers
                    1..27
                    note = synthetic construct; primer 146775
source             1..27
                    mol_type = other DNA

```

-continued

```

                organism = synthetic construct
SEQUENCE: 109
gctttctatt ttgtggcact attgtgg                                27

SEQ ID NO: 110          moltype = DNA  length = 31
FEATURE                Location/Qualifiers
misc_feature            1..31
                        note = synthetic construct; primer 146773
source                  1..31
                        mol_type = other DNA
                        organism = synthetic construct
SEQUENCE: 110
gctcgtgtcc aagcgtcact tacgattagc t                            31

SEQ ID NO: 111          moltype = DNA  length = 23
FEATURE                Location/Qualifiers
misc_feature            1..23
                        note = synthetic construct; primer 146772
source                  1..23
                        mol_type = other DNA
                        organism = synthetic construct
SEQUENCE: 111
accgctacca gcaacaatcg tct                                    23

SEQ ID NO: 112          moltype = DNA  length = 21
FEATURE                Location/Qualifiers
misc_feature            1..21
                        note = synthetic construct; primer 146778
source                  1..21
                        mol_type = other DNA
                        organism = synthetic construct
SEQUENCE: 112
tcacgctgca ctgcaggcta g                                      21

SEQ ID NO: 113          moltype = DNA  length = 27
FEATURE                Location/Qualifiers
misc_feature            1..27
                        note = synthetic construct; primer mopatF2
source                  1..27
                        mol_type = other DNA
                        organism = synthetic construct
SEQUENCE: 113
tcagatctgc gtcaccgggc gcaccgg                                27

SEQ ID NO: 114          moltype = DNA  length = 27
FEATURE                Location/Qualifiers
misc_feature            1..27
                        note = synthetic construct; primer mopatR2
source                  1..27
                        mol_type = other DNA
                        organism = synthetic construct
SEQUENCE: 114
ccgccgtgtg cgacatcgtg aaccact                                27

SEQ ID NO: 115          moltype = DNA  length = 16
FEATURE                Location/Qualifiers
misc_feature            1..16
                        note = synthetic construct; MHP55TS probe
source                  1..16
                        mol_type = other DNA
                        organism = synthetic construct
SEQUENCE: 115
aaccgtcgtg agacct                                           16

SEQ ID NO: 116          moltype = DNA  length = 17
FEATURE                Location/Qualifiers
misc_feature            1..17
                        note = synthetic construct; MHPTS55_Forward_MGB primer
source                  1..17
                        mol_type = other DNA
                        organism = synthetic construct
SEQUENCE: 116
aaggcgacgc cgttgag                                           17

SEQ ID NO: 117          moltype = DNA  length = 20
FEATURE                Location/Qualifiers

```

-continued

```

misc_feature      1..20
                  note = synthetic construct; MHP55TS_Reverse_MGB primer
source            1..20
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 117
ctaccggttt cgcgtgctct                               20

SEQ ID NO: 118      moltype = DNA length = 21
FEATURE            Location/Qualifiers
misc_feature        1..21
                  note = synthetic construct; MHP77TS probe
source              1..21
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 118
tagtatgaca tacataccgc c                               21

SEQ ID NO: 119      moltype = DNA length = 23
FEATURE            Location/Qualifiers
misc_feature        1..23
                  note = synthetic construct; MHP77TS_Forward_MGB primer
source              1..23
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 119
tccttagggc ggtatgtatg tca                             23

SEQ ID NO: 120      moltype = DNA length = 26
FEATURE            Location/Qualifiers
misc_feature        1..26
                  note = synthetic construct; MHP77TS_Reverse_MGB primer
source              1..26
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 120
catcgggtcaa aaaacacata aacttt                         26

SEQ ID NO: 121      moltype = DNA length = 105
FEATURE            Location/Qualifiers
source              1..105
                  mol_type = genomic DNA
                  organism = Glycine max

SEQUENCE: 121
tcttaaagaa gatacactgt gtaaatgtgt aatggcactg gcactctcgt gtgtgattaa 60
agtcatatat ggtttaagat accttttttt ataaagatag tagtg                    105

SEQ ID NO: 122      moltype = DNA length = 52
FEATURE            Location/Qualifiers
source              1..52
                  mol_type = genomic DNA
                  organism = Glycine max

SEQUENCE: 122
tcttaaagaa gatacactgt gtaaatgtgt aatggcactg gcactctcgt gt                    52

SEQ ID NO: 123      moltype = DNA length = 28
FEATURE            Location/Qualifiers
source              1..28
                  mol_type = genomic DNA
                  organism = Glycine max

SEQUENCE: 123
gatacttttt tttataaaga tagtagtg                         28

SEQ ID NO: 124      moltype = DNA length = 100
FEATURE            Location/Qualifiers
source              1..100
                  mol_type = genomic DNA
                  organism = Glycine max

SEQUENCE: 124
taatgatcac attttttttt tctcacactc acctaaagtc acgagtacac acgtaagtct 60
taggttaaag tttcatgccc ccccccccc ccccccaaaa                    100

SEQ ID NO: 125      moltype = DNA length = 54
FEATURE            Location/Qualifiers
source              1..54
                  mol_type = genomic DNA

```

-continued

```

                                organism = Glycine max
SEQUENCE: 125
taatgatcac attttttttt tctcacactc acctaagtgc acgagtacac acgt      54

SEQ ID NO: 126                moltype = DNA  length = 100
FEATURE                      Location/Qualifiers
source                      1..100
                             mol_type = genomic DNA
                             organism = Glycine max
SEQUENCE: 126
taatgatcac attttttttt tctcacactc acctaagtgc acgagtacac acgtaagtct  60
taggttaaag ttcatgccc ccccccccc ccccccaaaa      100

SEQ ID NO: 127                moltype = DNA  length = 45
FEATURE                      Location/Qualifiers
source                      1..45
                             mol_type = genomic DNA
                             organism = Glycine max
SEQUENCE: 127
taatgatcac attttttttt tctcacactc acctaagtgc acgag      45

SEQ ID NO: 128                moltype = DNA  length = 104
FEATURE                      Location/Qualifiers
source                      1..104
                             mol_type = genomic DNA
                             organism = Glycine max
SEQUENCE: 128
taatgatcac attttttttt tctcacactc acctaagtgc agacgtacgc aagtagcttt  60
gttactttcg tattgacaat tcaaaatcgt cttttatttt tatt      104

SEQ ID NO: 129                moltype = DNA  length = 102
FEATURE                      Location/Qualifiers
misc_feature                47..48
                             note = 115 bp insertion between position 47 and position 48
                             omitted below and in the alignment in Figure 4B
source                      1..102
                             mol_type = genomic DNA
                             organism = Glycine max
SEQUENCE: 129
taatgatcac attttttttt tctcacactc acctaagtgc agacgtacaa gtagctttgt  60
tactttcgta ttgacaattc aaaatcgtct tttattttta tt      102

```

1.-18. (canceled)

92. A maize plant comprising a complex transgenic trait locus in a plant, the trait locus comprising at least first and second altered target sequences, wherein the first altered target sequence originated from a first endogenous target sequence that is recognized and cleaved by a first double-strand-break-inducing agent and the second altered target sequence originated from a second endogenous target sequence that is recognized and cleaved by a second double-strand-break-inducing agent, wherein each of said altered target sequences differ from their corresponding endogenous target sequence, wherein the first and second endogenous target sequences are located on the same arm of the same

chromosome, wherein each of the alterations comprises a heterologous polynucleotide, and wherein at least one of the double-strand-break-inducing agents cleaves SEQ ID NO: 68 of SEQ ID NO:71.

93. The maize plant of claim 1 wherein the heterologous polynucleotide is selected from the group consisting of: DNA for gene silencing, DNA encoding a phenotypic marker, and DNA encoding a protein providing an agronomic advantage.

94. A seed of the maize plant of claim 1, comprising said complex transgenic trait locus of claim 1.

* * * * *