



US012404266B2

(12) **United States Patent**
Allasia et al.

(10) **Patent No.: US 12,404,266 B2**

(45) **Date of Patent: *Sep. 2, 2025**

(54) **HETEROCYCLE DERIVATIVES FOR
TREATING TRPM3 MEDIATED DISORDERS**

(71) Applicant: **Katholieke Universiteit Leuven,**
Leuven (BE)

(72) Inventors: **Sara Allasia,** Leuven (BE); **Philippe
Arzel,** Ghent (BE); **Patrick Chaltin,**
Zétrud-Lemay (BE); **Amuri Kilonda,**
Boutersem (BE); **Arnaud Marchand,**
Bierbeek (BE); **Jean-Christophe
Vanherck,** Leuven (BE); **Thomas
Voets,** Kessel-Lo (BE); **Joris Vriens,**
Wakkerzeel (BE); **Melanie Reich,**
Aachen (DE); **André Welbers,** Cologne
(DE)

(73) Assignee: **Katholieke Universiteit Leuven,**
Leuven (BE)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

This patent is subject to a terminal dis-
claimer.

(21) Appl. No.: **18/250,795**

(22) PCT Filed: **Nov. 24, 2021**

(86) PCT No.: **PCT/EP2021/082865**

§ 371 (c)(1),

(2) Date: **Apr. 27, 2023**

(87) PCT Pub. No.: **WO2022/112352**

PCT Pub. Date: **Jun. 2, 2022**

(65) **Prior Publication Data**

US 2024/0051949 A1 Feb. 15, 2024

(30) **Foreign Application Priority Data**

Nov. 24, 2020 (EP) 20209574

(51) **Int. Cl.**

C07D 405/12 (2006.01)
C07D 401/14 (2006.01)
C07D 405/14 (2006.01)
C07D 407/12 (2006.01)
C07D 413/12 (2006.01)
C07D 417/12 (2006.01)
C07D 417/14 (2006.01)
C07D 471/04 (2006.01)

(52) **U.S. Cl.**

CPC **C07D 405/12** (2013.01); **C07D 401/14**
(2013.01); **C07D 405/14** (2013.01); **C07D**
407/12 (2013.01); **C07D 413/12** (2013.01);
C07D 417/12 (2013.01); **C07D 417/14**
(2013.01); **C07D 471/04** (2013.01)

(58) **Field of Classification Search**

CPC .. **C07D 405/12**; **C07D 405/14**; **C07D 401/14**;
C07D 417/12; **C07D 417/14**; **C07D**
417/04; **C07D 471/04**
USPC 514/469
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

12,209,081 B2 * 1/2025 Chaltin C07D 417/12

FOREIGN PATENT DOCUMENTS

WO 2018/104479 A1 6/2018

OTHER PUBLICATIONS

May H. Abdel Aziz et al. "Allosteric Regulators of Thrombin. Monosulfated Benzofuran Dimers Selectively Interact with Arg173 of Exosite 2 to Induce Inhibition", Journal of Medicinal Chemistry, Jul. 25, 2012, vol. 55, No. 15, pp. 6888-6897, XP055777824.
Preetpal Singh Sidhu et al. "Supporting Information to Rational Design of Potent, Small, Synthetic Allosteric Inhibitors of Thrombin", Journal of Medicinal Chemistry, Jun. 29, 2011, pp. S1-S12, XP002792324.
Isabelle Straub et al. "Flavanones That Selectively Inhibit TRPM3 Attenuate Thermal Nociception In Vivo", Molecular Pharmacology, Nov. 7, 2013, vol. 84, No. 5, pp. 736-750, XP055777082.
International Search Report dated Feb. 4, 2022 issued for the corresponding application PCT/EP2021/082865 (4 pages).
Written Opinion dated Feb. 4, 2022 issued for the corresponding application PCT/EP2021/082865 (5 pages).
International Preliminary Report on Patentability dated May 30, 2023 issued for the corresponding application PCT/EP2021/082865 (7 pages).

* cited by examiner

Primary Examiner — Kahsay Habte

(57) **ABSTRACT**

The invention relates to compounds that are useful for the prevention or treatment of TRPM3 mediated disorders, more in particular disorders selected from pain and inflammatory hypersensitivity. The invention also relates to a method for the prevention or treatment of said TRPM3 mediated disorders.

13 Claims, No Drawings

1

HETEROCYCLE DERIVATIVES FOR TREATING TRPM3 MEDIATED DISORDERS

FIELD OF THE INVENTION

The invention relates to compounds that are useful for the prevention or treatment of TRPM3 mediated disorders, more in particular disorders selected from pain and inflammatory hypersensitivity. The invention also relates to a method for the prevention or treatment of said TRPM3 mediated disorders.

BACKGROUND OF THE INVENTION

The TRP superfamily consists of proteins with six transmembrane domains (6TM) that assemble as homo- or heterotetramers to form cation-permeable ion channels. The name TRP originates from the *Drosophila* trp (transient receptor potential) mutant, which is characterized by a transient receptor potential in the fly photoreceptors in the response to sustained light. In the last 15 years, trp-related channels have been identified in yeast, worms, insects, fish and mammals, including 27 TRPs in humans. Based on sequence homology, TRP channels can be divided into seven subfamilies: TRPC, TRPV, TRPM, TRPA, TRPP, TRPML and TRPN.

Members of the TRP superfamily are expressed in probably all mammalian organs and cell types, and in recent years great progress has been made in the understanding of their physiological role. The tailored selectivity of certain TRP channels enables them to play key roles in the cellular uptake and/or transepithelial transport of Ca^{2+} , Mg^{2+} and trace metal ions. Moreover, the sensitivity of TRP channels to a broad array of chemical and physical stimuli, allows them to function as dedicated biological sensors involved in processes ranging from vision to taste, and tactile sensation. In particular, several members of the TRP superfamily exhibit a very high sensitivity to temperature. These so-called thermoTRPs are highly expressed in sensory neurons and/or skin keratinocytes, where they act as primary thermosensors for the detection of innocuous and noxious (painful) temperatures.

It is becoming increasingly clear that TRP channel dysfunction is directly involved in the etiology of various inherited and acquired diseases. Indeed, both loss-of-function and gain-of-function mutations in the TRP channel genes have been identified as the direct cause of inherited diseases, including brachyolmia, hypomagnesemia with secondary hypocalcemia, polycystic kidney disease, mucopolysaccharidosis type IV and familial focal segmental glomerulosclerosis. Moreover, TRP channel function/dysfunction has been directly linked to a wide range of pathological conditions, including chronic pain, hypertension, cancer and neurodegenerative disorders.

TRPM3 (Transient receptor potential melastatin 3) represents a promising pharmacological target. TRPM3 is expressed in a large subset of small-diameter sensory neurons from dorsal root and trigeminal ganglia, and is involved in heat sensing. The neurosteroid pregnenolone sulfate is a potent known activator of TRPM3 (Wagner et al., 2008). The neurosteroid pregnenolone sulfate evoked pain in wild type mice but not in knock-out TRPM3 mice. It was also recently shown that CFA induced inflammation and inflammatory pain are eliminated in TRPM3 knock-out mice. Therefore, TRPM3 antagonists could be used as analgesic drugs to counteract pain, such as inflammatory pain (Vriens J. et al. Neuron, May 2011).

2

A few TRPM3 antagonists are known, but none of them points towards the compounds of the current invention (Straub I et al. Mol Pharmacol, November 2013). For instance, Liquiritigenin, a postulated TRPM3 blocker has been described to decrease mechanical and cold hyperalgesia in a rat pain model (Chen L et al. Scientific reports, July 2014). There is still a great medical need for novel, alternative and/or better therapeutics for the prevention or treatment of TRPM3 mediated disorders, more in particular for pain such as inflammatory pain. Therapeutics with good potency on a certain type of pain, low level or no side-effects (such as no possibilities for addiction as with opiates, no toxicity) and/or good or better pharmacokinetic or -dynamic properties are highly needed.

The invention provides a class of novel compounds which are antagonists of TRPM3 and can be used as modulators of TRPM3 mediated disorders.

SUMMARY OF THE INVENTION

The invention provides benzofuran derivatives and pharmaceutical compositions comprising such benzofuran derivatives. The invention also provides benzofuran derivatives for use as a medicament, more in particular for use in the prevention and/or treatment of TRPM3 mediated disorders, especially for use in the prevention and/or treatment of pain and/or inflammatory hypersensitivity; and/or for counteracting pain and/or inflammatory hypersensitivity.

The invention also provides the use of benzofuran derivatives for the manufacture of pharmaceutical compositions or medicaments for the prevention and/or treatment of TRPM3 mediated disorders, especially for the prevention and/or treatment of pain and/or inflammatory hypersensitivity; and/or for counteracting pain and/or inflammatory hypersensitivity.

The invention also provides a method for the prevention or treatment of a TRPM3 mediated disorder by administering the benzofuran derivatives according to the invention to a subject in need thereof. More in particular, the invention relates to such method for the prevention and/or treatment of pain and/or inflammatory hypersensitivity; and/or for counteracting pain and/or inflammatory hypersensitivity.

The invention further provides a method for the preparation of the benzofuran derivatives of the invention, comprising the steps of:


- reacting a benzoquinone with a suitable β -ketoester or an enamine derivative to obtain 5-hydroxybenzofuran-3-carboxylate ester derivatives,
- substituting previously obtained 5-hydroxybenzofuran-3-carboxylate ester derivatives with suitable derivatives bearing a leaving group or alcohol derivatives under Mitsunobu conditions to obtain 5-O-substituted-benzofuran-3-carboxylate ester derivatives
- converting the previously obtained 5-O-substituted-benzofuran-3-carboxylate ester derivatives in carboxylic acid to obtain the desired benzofuran derivatives of the invention, and
- coupling the previously obtained 5-O-substituted-benzofuran-3-carboxylic acid derivatives with a suitable amine to obtain the desired amide derivatives of the invention.

DETAILED DESCRIPTION OF THE INVENTION

The invention will be further described and in some instances with respect to particular embodiments, but the invention is not limited thereto.

3

The first aspect of the invention is the provision of a compound of formula (I), a stereo-isomeric form, a physiologically acceptable salt, solvate and/or polymorph thereof



preferably for use in the treatment of pain;
wherein

R¹ represents —F, —Cl, —Br, —I, —CN, —R^W, —OR^W,
—OC(=O)R^W, —NR^WR^X, —NR^WC(=O)R^X, —SR^W,
—S(=O)R^W, —S(=O)₂R^W, —C(=O)R^W, —C(=O)
OR^W, or —C(=O)NR^WR^X;

Q represents —OR² or —NR³R⁴;

R² represents —R^Y;

R³ represents —OH or —R^Y;

R⁴ represents —R^Y or —S(=O)₂R^Y;

or R³ and R⁴ together form a 4, 5, 6, 7 or 8 membered
heterocycle containing 1 to 3 heteroatoms selected
from N, O and S, saturated or unsaturated, unsubsti-
tuted or mono- or polysubstituted;

T represents —O— and U represents —CR⁵R⁵ⁱ—; or T
represents —CR⁵R⁵ⁱ— and U represents —O—;

R⁵ and R⁵ⁱ independently of one another represent —R^Y;

R⁶, R⁷ and R⁸ independently of one another represent —F,
—Cl, —Br, —I, —CN, —NO₂, —SF₅, —R^W, —OR^W,
—OC(=O)R^W, —NR^WR^X, —NR^WC(=O)R^X, —SR^W,
—S(=O)R^W, —S(=O)₂R^W, —C(=O)R^W, —C(=O)
OR^W, or —C(=O)NR^WR^X;

V represents 3-14-membered heterocycloalkyl, saturated
or unsaturated; or 5-14-membered heteroaryl; in each
case unsubstituted, mono- or polysubstituted with sub-
stituents independently of one another selected from
—F, —Cl, —Br, —I, —CN, —NO, —NO₂, —O, —S,
—SF₅, —R^Y, —OR^Y, —OC(=O)R^Y, —NR^YR^Z,
—NR^YC(=O)R^Z, —SR^Y, —S(=O)R^Y, —S(=O)₂R^Y,
—C(=O)R^Y, —C(=O)OR^Y, or —C(=O)NR^YR^Z;

wherein

R^W and R^X independently of one another in each case
independently represent

—H;

—C₁-C₆-alkyl, saturated or unsaturated, unsubstituted,
mono- or polysubstituted;

—C₁-C₆-heteroalkyl, saturated or unsaturated, unsubsti-
tuted, mono- or polysubstituted;

3-14-membered cycloalkyl, saturated or unsaturated,
unsubstituted, mono- or polysubstituted; wherein said
3-14-membered cycloalkyl is optionally connected
through —C₁-C₆-alkylene- or —C₁-C₆-heteroal-
kylene-, in each case saturated or unsaturated, unsub-
stituted, mono- or polysubstituted; or

3-14-membered heterocycloalkyl, saturated or unsatu-
rated, unsubstituted, mono- or polysubstituted; wherein
said 3-14-membered heterocycloalkyl is optionally
connected through —C₁-C₆-alkylene- or —C₁-C₆-het-
eroalkylene-, in each case saturated or unsaturated,
unsubstituted, mono- or polysubstituted;

4

R^Y and R^Z independently of one another in each case
independently represent

—H;

—C₁-C₆-alkyl, saturated or unsaturated, unsubstituted,
mono- or polysubstituted;

—C₁-C₆-heteroalkyl, saturated or unsaturated, unsubsti-
tuted, mono- or polysubstituted;

3-14-membered cycloalkyl, saturated or unsaturated,
unsubstituted, mono- or polysubstituted; wherein said
3-14-membered cycloalkyl is optionally connected
through —C₁-C₆-alkylene- or —C₁-C₆-heteroal-
kylene-, in each case saturated or unsaturated, unsub-
stituted, mono- or polysubstituted;

3-14-membered heterocycloalkyl, saturated or unsatu-
rated, unsubstituted, mono- or polysubstituted; wherein
said 3-14-membered heterocycloalkyl is optionally
connected through —C₁-C₆-alkylene- or —C₁-C₆-het-
eroalkylene-, in each case saturated or unsaturated,
unsubstituted, mono- or polysubstituted;

6-14-membered aryl, unsubstituted, mono- or polysubsti-
tuted; wherein said 6-14-membered aryl is optionally
connected through —C₁-C₆-alkylene- or —C₁-C₆-het-
eroalkylene-, in each case saturated or unsaturated,
unsubstituted, mono- or polysubstituted; or

5-14-membered heteroaryl, unsubstituted, mono- or poly-
substituted; wherein said 5-14-membered heteroaryl is
optionally connected through —C₁-C₆-alkylene- or
—C₁-C₆-heteroalkylene-, in each case saturated or
unsaturated, unsubstituted, mono- or polysubstituted;
or R^Y and R^Z together form a 4, 5, 6, 7 or 8 membered
heterocycle containing 1 to 3 heteroatoms selected
from N, O and S, saturated or unsaturated, unsubsti-
tuted or mono- or polysubstituted;

and wherein “mono- or polysubstituted” in each case
independently means substituted with one or more, e.g.
1, 2, 3, 4, or more substituents independently of one
another selected from —F, —Cl, —Br, —I, —CN,
—C₁₋₆-alkyl, —CF₃, —CF₂H, —CFH₂, —CF₂Cl,
—CFC₂, —C₁₋₆-alkylene-CF₃, —C₁₋₆-alkylene-
CF₂H, —C₁₋₆-alkylene-CFH₂, —C₁₋₆-alkylene-O—
CF₃, —C₁₋₆-alkylene-O—CF₂H, —C₁₋₆-alkylene-O—
CFH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃,
—C₁₋₆-alkylene-N(C₁₋₆-alkyl)-C₁₋₆-alkylene-CF₃,
—C(=O)—C₁₋₆-alkyl, —C₁₋₆-alkylene-C(=O)—
C₁₋₆-alkyl, —C(=O)OH, —C₁₋₆-alkylene-C(=O)—
OH, —C(=O)—OC₁₋₆-alkyl, —C₁₋₆-alkylene-C
(=O)—OC₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkylene-CF₃,
—C(=O)—NH₂, —C₁₋₆-alkylene-C(=O)—NH₂,
—C(=O)—NH(C₁₋₆-alkyl), —C₁₋₆-alkylene-C
(=O)—NH(C₁₋₆-alkyl), —C(=O)—N(C₁₋₆-alkyl)₂,
—C₁₋₆-alkylene-C(=O)—N(C₁₋₆-alkyl)₂,
—C(=O)—NH(OH), —C₁₋₆-alkylene-C(=O)—NH
(OH), —OH, —C₁₋₆-alkylene-OH, —O, —OCF₃,
—OCF₂H, —OCFH₂, —OCF₂Cl, —OCFC₂,
—O—C₁₋₆-alkyl, —C₁₋₆-alkylene-O—C₁₋₆-alkyl,
—O—C₁₋₆-alkylene-O—C₁₋₆-alkyl, —O—C₁₋₆-al-
kylene-NH₂, —O—C₁₋₆-alkylene-NH—C₁₋₆-alkyl,
—O—C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —O—C(=O)—
C₁₋₆-alkyl, —C₁₋₆-alkylene-O—C(=O)—C₁₋₆-alkyl,
—O—C(=O)—O—C₁₋₆-alkyl, —C₁₋₆-alkylene-O—
C(=O)—O—C₁₋₆-alkyl, —O—C(=O)—NH(C₁₋₆-al-
kyl), —C₁₋₆-alkylene-O—C(=O)—NH(C₁₋₆-alkyl),
—O—C(=O)—N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-O—
C(=O)—N(C₁₋₆-alkyl)₂, —O—S(=O)₂—NH₂,
—C₁₋₆-alkylene-O—S(=O)₂—NH₂, —O—S(=O)₂
—NH(C₁₋₆-alkyl), —C₁₋₆-alkylene-O—S(=O)₂—NH
(C₁₋₆-alkyl), —O—S(=O)₂—N(C₁₋₆-alkyl)₂,

5

$\text{—C}_{1-6}\text{-alkylene—O—S(=O)}_2\text{—N(C}_{1-6}\text{-alkyl)}_2$,
 —NH_2 , —NO , —NO_2 , $\text{—C}_{1-6}\text{-alkylene—NH}_2$, —NH
 $(\text{C}_{1-6}\text{-alkyl})$, $\text{—N(3-14-membered cycloalkyl)(C}_{1-6}\text{-alkyl)}$,
 $\text{—N(C}_{1-6}\text{-alkyl)—C}_{1-6}\text{-alkylene—OH}$, —N(H)—
 $\text{C}_{1-6}\text{-alkylene—OH}$, $\text{—C}_{1-6}\text{-alkylene—NH(C}_{1-6}\text{-alkyl)}$,
 $\text{—N(C}_{1-6}\text{-alkyl)}_2$, $\text{—C}_{1-6}\text{-alkylene—N(C}_{1-6}\text{-alkyl)}_2$,
 $\text{—NH—C(=O)—C}_{1-6}\text{-alkyl}$, $\text{—C}_{1-6}\text{-alkylene—NH—C}$
 $(=\text{O})\text{—C}_{1-6}\text{-alkyl}$, $\text{—NH—C(=O)—O—C}_{1-6}\text{-alkyl}$,
 $\text{—C}_{1-6}\text{-alkylene—NH—C(=O)—O—C}_{1-6}\text{-alkyl}$,
 —NH—C(=O)—NH_2 , $\text{—C}_{1-6}\text{-alkylene—NH—C}$
 $(=\text{O})\text{—NH}_2$, $\text{—NH—C(=O)—NH(C}_{1-6}\text{-alkyl)}$,
 $\text{—C}_{1-6}\text{-alkylene—NH—C(=O)—NH(C}_{1-6}\text{-alkyl)}$,
 $\text{—NH—C(=O)—N(C}_{1-6}\text{-alkyl)}_2$, $\text{—C}_{1-6}\text{-alkylene—}$
 $\text{NH—C(=O)—N(C}_{1-6}\text{-alkyl)}_2$, $\text{—N(C}_{1-6}\text{-alkyl)—C}$
 $(=\text{O})\text{—C}_{1-6}\text{-alkyl}$, $\text{—C}_{1-6}\text{-alkylene—N(C}_{1-6}\text{-alkyl)—C}$
 $(=\text{O})\text{—C}_{1-6}\text{-alkyl}$, $\text{—N(C}_{1-6}\text{-alkyl)—C(=O)—O—}$
 $\text{C}_{1-6}\text{-alkyl}$, $\text{—C}_{1-6}\text{-alkylene—N(C}_{1-6}\text{-alkyl)—C(=O)—}$
 $\text{O—C}_{1-6}\text{-alkyl}$, $\text{—N(C}_{1-6}\text{-alkyl)—C(=O)—NH}_2$,
 $\text{—C}_{1-6}\text{-alkylene—N(C}_{1-6}\text{-alkyl)—C(=O)—NH}_2$,
 $\text{—N(C}_{1-6}\text{-alkyl)—C(=O)—NH(C}_{1-6}\text{-alkyl)}$,
 $\text{—N(C}_{1-6}\text{-alkyl)—C(=O)—N(C}_{1-6}\text{-alkyl)}_2$, $\text{—C}_{1-6}\text{-alkylene—}$
 $\text{N(C}_{1-6}\text{-alkyl)—C(=O)—N(C}_{1-6}\text{-alkyl)}_2$,
 $\text{—NH—S(=O)}_2\text{OH}$, $\text{—C}_{1-6}\text{-alkylene—NH—S(=O)}_2$
 OH , $\text{—NH—S(=O)}_2\text{—C}_{1-6}\text{-alkyl}$, $\text{—C}_{1-6}\text{-alkylene—}$
 $\text{NH—S(=O)}_2\text{—C}_{1-6}\text{-alkyl}$, $\text{—NH—S(=O)}_2\text{—O—}$
 $\text{C}_{1-6}\text{-alkyl}$, $\text{—C}_{1-6}\text{-alkylene—NH—S(=O)}_2\text{—O—C}_{1-6}\text{-}$
 alkyl , $\text{—NH—S(=O)}_2\text{—NH}_2$, $\text{—C}_{1-6}\text{-alkylene—NH—}$
 $\text{S(=O)}_2\text{—NH}_2$, $\text{—NH—S(=O)}_2\text{—NH(C}_{1-6}\text{-alkyl)}$,
 $\text{—C}_{1-6}\text{-alkylene—NH—S(=O)}_2\text{—NH(C}_{1-6}\text{-alkyl)}$,
 $\text{—NH—S(=O)}_2\text{N(C}_{1-6}\text{-alkyl)}_2$, $\text{—C}_{1-6}\text{-alkylene—}$
 $\text{NH—S(=O)}_2\text{N(C}_{1-6}\text{-alkyl)}_2$, $\text{—N(C}_{1-6}\text{-alkyl)—S}$
 $(=\text{O})_2\text{—OH}$, $\text{—C}_{1-6}\text{-alkylene—N(C}_{1-6}\text{-alkyl)—S(=O)}_2$
 —OH , $\text{—N(C}_{1-6}\text{-alkyl)—S(=O)}_2\text{—C}_{1-6}\text{-alkyl}$, $\text{—C}_{1-6}\text{-alkylene—}$
 $\text{N(C}_{1-6}\text{-alkyl)—S(=O)}_2\text{—C}_{1-6}\text{-alkyl}$,
 $\text{—N(C}_{1-6}\text{-alkyl)—S(=O)}_2\text{—O—C}_{1-6}\text{-alkyl}$, $\text{—C}_{1-6}\text{-alkylene—}$
 $\text{N(C}_{1-6}\text{-alkyl)—S(=O)}_2\text{—O—C}_{1-6}\text{-alkyl}$,
 $\text{—N(C}_{1-6}\text{-alkyl)—S(=O)}_2\text{—NH}_2$, $\text{—C}_{1-6}\text{-alkylene—N}$
 $(\text{C}_{1-6}\text{-alkyl)—S(=O)}_2\text{—NH}_2$, $\text{—N(C}_{1-6}\text{-alkyl)—S(=O)}_2$
 $\text{—NH(C}_{1-6}\text{-alkyl)}$, $\text{—C}_{1-6}\text{-alkylene—N(C}_{1-6}\text{-alkyl)—S}$
 $(=\text{O})_2\text{—NH(C}_{1-6}\text{-alkyl)}$, $\text{—N(C}_{1-6}\text{-alkyl)—S(=O)}_2\text{—}$
 $\text{N(C}_{1-6}\text{-alkyl)}_2$, $\text{—C}_{1-6}\text{-alkylene—N(C}_{1-6}\text{-alkyl)—S}$
 $(=\text{O})_2\text{—N(C}_{1-6}\text{-alkyl)}_2$, —SH , —S , —SF_5 , —SCF_3 ,
 $\text{—SCF}_2\text{H}$, —SCFH_2 , $\text{—S—C}_{1-6}\text{-alkyl}$, $\text{—C}_{1-6}\text{-alkylene—}$
 $\text{S—C}_{1-6}\text{-alkyl}$, $\text{—S(=O)—C}_{1-6}\text{-alkyl}$, $\text{—C}_{1-6}\text{-alkylene—}$
 $\text{S(=O)—C}_{1-6}\text{-alkyl}$, $\text{—S(=O)}_2\text{—C}_{1-6}\text{-alkyl}$,
 $\text{—C}_{1-6}\text{-alkylene—S(=O)}_2\text{—C}_{1-6}\text{-alkyl}$, $\text{—S(=O)}_2\text{—}$
 OH , $\text{—C}_{1-6}\text{-alkylene—S(=O)}_2\text{—OH}$, $\text{—S(=O)}_2\text{—O—}$
 $\text{C}_{1-6}\text{-alkyl}$, $\text{—C}_{1-6}\text{-alkylene—S(=O)}_2\text{—O—C}_{1-6}\text{-alkyl}$,
 $\text{—S(=O)}_2\text{—NH}_2$, $\text{—C}_{1-6}\text{-alkylene—S(=O)}_2\text{—NH}_2$,
 $\text{—S(=O)}_2\text{—NH(C}_{1-6}\text{-alkyl)}$, $\text{—C}_{1-6}\text{-alkylene—S}$
 $(=\text{O})_2\text{—NH(C}_{1-6}\text{-alkyl)}$, $\text{—S(=O)}_2\text{—N(C}_{1-6}\text{-alkyl)}_2$,
 $\text{—C}_{1-6}\text{-alkylene—S(=O)}_2\text{—N(C}_{1-6}\text{-alkyl)}_2$, 3-14-mem-
 bered cycloalkyl, $\text{—C}_{1-6}\text{-alkylene-(3-14-membered}$
 cycloalkyl), 3 to 14-membered heterocycloalkyl,
 $\text{—C}_{1-6}\text{-alkylene-(3 to 14-membered heterocycloalkyl)}$,
 —phenyl , $\text{—C}_{1-6}\text{-alkylene—phenyl}$, 5 to 14-membered
 heteroaryl, $\text{—C}_{1-6}\text{-alkylene-(5 to 14-membered heteroaryl)}$,
 $\text{—O-(3-14-membered cycloalkyl)}$, $\text{—O-(3 to 14-membered heterocycloalkyl)}$,
 —O-phenyl , $\text{—O-(5 to 14-membered heteroaryl)}$,
 $\text{—C(=O)-(3-14-membered cycloalkyl)}$, $\text{—C(=O)-(3 to 14-membered heterocycloalkyl)}$,
 —C(=O)-phenyl , $\text{—C(=O)-(5 to 14-membered heteroaryl)}$,
 $\text{—S(=O)}_2\text{-(3-14-membered heterocycloalkyl)}$,
 $\text{—S(=O)}_2\text{-(3 to 14-membered heterocycloalkyl)}$,
 $\text{—S(=O)}_2\text{-phenyl}$, $\text{—S(=O)}_2\text{-(5 to 14-membered heteroaryl)}$.

6

In preferred embodiments of the benzofuran derivative according to the invention




(a-1) Q represents —OR^2 ; and R^1 represents $\text{—CH}_2\text{F}$, —CHF_2 , or —CF_3 ; and/or

(a-2) Q represents —OR^2 ; and at least one of R^5 and $\text{R}^{5'}$ does not represent —H ; and/or

(a-3) Q represents —OR^2 ; and R^6 does not represent —H ; and/or

(a-4) Q represents —OR^2 ; and R^8 does not represent —H ; or

(b-1) Q represents $\text{—NR}^3\text{R}^4$; with the proviso that the following compounds and their salts are excluded



and/or


(b-2) Q represents $\text{—NR}^3\text{R}^4$; and R^1 represents $\text{—CH}_2\text{F}$, —CHF_2 , —CF_3 , —CN , —propyl or —cyclopropyl ; and/or

(b-3) Q represents $\text{—NR}^3\text{R}^4$; and at least one of R^5 and $\text{R}^{5'}$ does not represent —H ; and/or

(b-4) Q represents $\text{—NR}^3\text{R}^4$; and R^3 represents —H .

In a preferred embodiment of the benzofuran derivative according to the invention T represents —O— and U represents $\text{—CR}^5\text{R}^{5'}$. According to this embodiment, the benzofuran derivative according to the invention is a compound of formula (II), a stereo-isomeric form, a physiologically acceptable salt, solvate and/or polymorph thereof.

7



In another preferred embodiment of the benzofuran derivative according to the invention T represents $-\text{CR}^5\text{R}^{5'}$ and U represents $-\text{O}-$.

In a preferred embodiment of the benzofuran derivative according to the invention Q represents $-\text{NR}^3\text{R}^4$.

In another preferred embodiment of the benzofuran derivative according to the invention Q represents $-\text{OR}^2$.

In a preferred embodiment of the benzofuran derivative according to the invention V represents 5-14-membered heteroaryl, unsubstituted, mono- or polysubstituted with substituents independently of one another selected from $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{CN}$, $-\text{NO}$, $-\text{NO}_2$, $=\text{O}$, $=\text{S}$, $-\text{SF}_5$, $-\text{R}^Y$, $-\text{OR}^Y$, $-\text{OC}(=\text{O})\text{R}^Y$, $-\text{NR}^Y\text{R}^Z$, $-\text{NR}^Y\text{C}(=\text{O})\text{R}^Z$, $-\text{SR}^Y$, $-\text{S}(=\text{O})\text{R}^Y$, $-\text{S}(=\text{O})_2\text{R}^Y$, $-\text{C}(=\text{O})\text{R}^Y$, $-\text{C}(=\text{O})\text{OR}^Y$, or $-\text{C}(=\text{O})\text{NR}^Y\text{R}^Z$.

Preferably, the 5-14-membered heteroaryl within the definition of V is not benzofuran, unsubstituted, mono- or polysubstituted.

Preferably, the 5-14-membered heteroaryl within the definition of V is selected from benzimidazole, benzisoxazole, benzoazole, benzodioxole, benzofuran, benzothiadiazole, benzothiazole, benzothiophene, carbazole, cinnoline, dibenzofuran, furane, furazane, imidazole, imidazopyridine, indazole, indole, indolizine, isobenzofuran, isoindole, isoquinoline, isothiazole, isoxazole, naphthyridine, oxadiazole, oxazole, oxindole, phthalazine, purine, pyrazine, pyrazole, pyridazine, pyridine, pyrimidine, pyrrole, quinazoline, quinoxaline, tetrazole, thiadiazole, thiazole, thiophene, triazine, triazole, and [1,2,4]triazolo[4,3-a]pyrimidine; in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{CN}$, $-\text{NO}$, $-\text{NO}_2$, $=\text{O}$, $=\text{S}$, $-\text{SF}_5$, $-\text{R}^Y$, $-\text{OR}^Y$, $-\text{OC}(=\text{O})\text{R}^Y$, $-\text{NR}^Y\text{R}^Z$, $-\text{NR}^Y\text{C}(=\text{O})\text{R}^Z$, $-\text{SR}^Y$, $-\text{S}(=\text{O})\text{R}^Y$, $-\text{S}(=\text{O})_2\text{R}^Y$, $-\text{C}(=\text{O})\text{R}^Y$, $-\text{C}(=\text{O})\text{OR}^Y$, or $-\text{C}(=\text{O})\text{NR}^Y\text{R}^Z$.

Preferably, the 5-14-membered heteroaryl within the definition of V is selected from the group consisting of furane, thiophene, imidazole, pyrazole, oxazole, isoxazole, thiazole, triazole, pyridine, isoquinoline, benzothiazole, pyridazine, pyrimidine, imidazopyridine; in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{CN}$, $-\text{NO}$, $-\text{NO}_2$, $=\text{O}$, $=\text{S}$, $-\text{SF}_5$, $-\text{R}^Y$, $-\text{OR}^Y$, $-\text{OC}(=\text{O})\text{R}^Y$, $-\text{NR}^Y\text{R}^Z$, $-\text{NR}^Y\text{C}(=\text{O})\text{R}^Z$, $-\text{SR}^Y$, $-\text{S}(=\text{O})\text{R}^Y$, $-\text{S}(=\text{O})_2\text{R}^Y$, $-\text{C}(=\text{O})\text{R}^Y$, $-\text{C}(=\text{O})\text{OR}^Y$, or $-\text{C}(=\text{O})\text{NR}^Y\text{R}^Z$.

Preferably, the 5-14-membered heteroaryl within the definition of V is selected from the group consisting of furan-2-yl, furan-3-yl, thiophen-2-yl, thiophen-3-yl, pyrazol-5-yl, oxazol-5-yl, isoxazol-4-yl, thiazol-2-yl, thiazol-5-yl, 1,2,4-triazol-3-yl, 1,2,3-triazol-4-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, isoquinolin-1-yl, isoquinolin-5-yl, benzo[d]thiazol-2-yl, pyridazin-3-yl, pyrimidin-5-yl, and imidazo[1,2-a]pyridin-6-yl; in each case unsubstituted, mono- or polysubstituted with substituents independently of one another

8

(II) selected from $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{CN}$, $-\text{NO}$, $-\text{NO}_2$, $=\text{O}$, $=\text{S}$, $-\text{SF}_5$, $-\text{R}^Y$, $-\text{OR}^Y$, $-\text{OC}(=\text{O})\text{R}^Y$, $-\text{NR}^Y\text{R}^Z$, $-\text{NR}^Y\text{C}(=\text{O})\text{R}^Z$, $-\text{SR}^Y$, $-\text{S}(=\text{O})\text{R}^Y$, $-\text{S}(=\text{O})_2\text{R}^Y$, $-\text{C}(=\text{O})\text{R}^Y$, $-\text{C}(=\text{O})\text{OR}^Y$, or $-\text{C}(=\text{O})\text{NR}^Y\text{R}^Z$.

5 In another preferred embodiment of the benzofuran derivative according to the invention V represents 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted with substituents independently of one another selected from $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{CN}$, $-\text{NO}$, $-\text{NO}_2$, $=\text{O}$, $=\text{S}$, $-\text{SF}_5$, $-\text{R}^Y$, $-\text{OR}^Y$, $-\text{OC}(=\text{O})\text{R}^Y$, $-\text{NR}^Y\text{R}^Z$, $-\text{NR}^Y\text{C}(=\text{O})\text{R}^Z$, $-\text{SR}^Y$, $-\text{S}(=\text{O})\text{R}^Y$, $-\text{S}(=\text{O})_2\text{R}^Y$, $-\text{C}(=\text{O})\text{R}^Y$, $-\text{C}(=\text{O})\text{OR}^Y$, or $-\text{C}(=\text{O})\text{NR}^Y\text{R}^Z$.

15 Preferably, the 3-14-membered heterocycloalkyl within the definition of V is selected from azepane, 1,4-oxazepane, azetane, azetidine, aziridine, azocane, diazepane, dioxane, dioxolane, dithiane, dithiolane, imidazolidine, isothiazolidine, isoxalidine, morpholine, oxazolidine, oxepane, oxetane, oxirane, piperazine, piperidine, pyrazolidine, pyrrolidine, quinuclidine, tetrahydrofuran, tetrahydropyran, tetrahydrothiopyran, thiazolidine, thietane, thirane, thiolane, thiomorpholine, indoline, dihydrobenzofuran, dihydrobenzothiophene, 1,1-dioxothiacyclohexane, 2-azaspiro[3.3]heptane, 2-oxaspiro[3.3]heptane, 7-azaspiro[3.5]nonane, 8-azabicyclo[3.2.1]octane, 9-azabicyclo[3.3.1]nonane, hexahydro-1H-pyrrolizine, hexahydrocyclopenta[c]pyrrole, octahydrocyclopenta[c]pyrrole, and octahydropyrrolo[1,2-a]pyrazine; in each case unsubstituted, 20 mono- or polysubstituted with substituents independently of one another selected from $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{CN}$, $-\text{NO}$, $-\text{NO}_2$, $=\text{O}$, $=\text{S}$, $-\text{SF}_5$, $-\text{R}^Y$, $-\text{OR}^Y$, $-\text{OC}(=\text{O})\text{R}^Y$, $-\text{NR}^Y\text{R}^Z$, $-\text{NR}^Y\text{C}(=\text{O})\text{R}^Z$, $-\text{SR}^Y$, $-\text{S}(=\text{O})\text{R}^Y$, $-\text{S}(=\text{O})_2\text{R}^Y$, $-\text{C}(=\text{O})\text{R}^Y$, $-\text{C}(=\text{O})\text{OR}^Y$, or $-\text{C}(=\text{O})\text{NR}^Y\text{R}^Z$.

30 Preferably, the 3-14-membered heterocycloalkyl within the definition of V is tetrahydropyran or pyrrolidine; in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{CN}$, $-\text{NO}$, $-\text{NO}_2$, $=\text{O}$, $=\text{S}$, $-\text{SF}_5$, $-\text{R}^Y$, $-\text{OR}^Y$, $-\text{OC}(=\text{O})\text{R}^Y$, $-\text{NR}^Y\text{R}^Z$, $-\text{NR}^Y\text{C}(=\text{O})\text{R}^Z$, $-\text{SR}^Y$, $-\text{S}(=\text{O})\text{R}^Y$, $-\text{S}(=\text{O})_2\text{R}^Y$, $-\text{C}(=\text{O})\text{R}^Y$, $-\text{C}(=\text{O})\text{OR}^Y$, or $-\text{C}(=\text{O})\text{NR}^Y\text{R}^Z$.

40 Preferably, the 3-14-membered heterocycloalkyl within the definition of V is tetrahydropyran-4-yl or pyrrolidin-3-yl; in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{CN}$, $-\text{NO}$, $-\text{NO}_2$, $=\text{O}$, $=\text{S}$, $-\text{SF}_5$, $-\text{R}^Y$, $-\text{OR}^Y$, $-\text{OC}(=\text{O})\text{R}^Y$, $-\text{NR}^Y\text{R}^Z$, $-\text{NR}^Y\text{C}(=\text{O})\text{R}^Z$, $-\text{SR}^Y$, $-\text{S}(=\text{O})\text{R}^Y$, $-\text{S}(=\text{O})_2\text{R}^Y$, $-\text{C}(=\text{O})\text{R}^Y$, $-\text{C}(=\text{O})\text{OR}^Y$, or $-\text{C}(=\text{O})\text{NR}^Y\text{R}^Z$.

50 In a preferred embodiment of the benzofuran derivative according to the invention V is unsubstituted, mono- or polysubstituted with substituents independently of one another selected from

$-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{CN}$, $-\text{C}(=\text{O})\text{OH}$, $-\text{NH}_2$, $-\text{NO}_2$, $-\text{OH}$, $=\text{O}$, $-\text{SF}_5$;
 $-\text{C}_{1-6}$ -alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
 $-\text{C}(=\text{O})\text{O}-\text{C}_{1-6}$ -alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
 $-\text{NHC}_{1-6}$ -alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
 $-\text{N}(\text{C}_{1-6}\text{-alkyl})_2$, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
 $-\text{O}-\text{C}_{1-6}$ -alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;

9

—S(=O)₂—C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;

3-14-membered cycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered cycloalkyl is optionally connected through —C₁-C₆-alkylene- or —C₁-C₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted; or

3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered heterocycloalkyl is optionally connected through —C₁-C₆-alkylene- or —C₁-C₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted.

Preferably, V is unsubstituted, mono- or polysubstituted with substituents independently of one another selected from

—OH, —F, —Cl, —Br, —I, —SH, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —CN, —NO₂, —C(=O)OH, —NH₂, or —N(CH₃)₂;

—C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted with substituents independently of one another, selected from the group consisting of —F, —Cl, —Br, —I, —C₁₋₆-alkyl, C₂₋₆-alkenyl, —C₂₋₆-alkynyl, —OH, —O, —SH, —S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, C(=O)CHF₂, and —C(=O)NH₂;

—C₁₋₆-heteroalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted with substituents independently of one another, selected from the group consisting of —F, —Cl, —Br, —I, —C₁₋₆-alkyl, C₂₋₆-alkenyl, —C₂₋₆-alkynyl, —OH, —O, —SH, —S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, C(=O)CHF₂, and —C(=O)NH₂;

—OC₁₋₆-alkyl, unsubstituted, mono- or polysubstituted with substituents independently of one another, selected from the group consisting of —F, —Cl, —Br, —I, —C₁₋₆-alkyl, C₂₋₆-alkenyl, —C₂₋₆-alkynyl, —OH, —O, —SH, —S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, C(=O)CHF₂, and —C(=O)NH₂;

—O(C=O)C₁₋₆-alkyl, unsubstituted, mono- or polysubstituted with substituents independently of one another, selected from the group consisting of —F, —Cl, —Br, —I, —C₁₋₆-alkyl, C₂₋₆-alkenyl, —C₂₋₆-alkynyl, —OH, —O, —SH, —S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, C(=O)CHF₂, and —C(=O)NH₂;

—C(=O)OC₁₋₆-alkyl, unsubstituted, mono- or polysubstituted with substituents independently of one another, selected from the group consisting of —F, —Cl, —Br, —I, —C₁₋₆-alkyl, C₂₋₆-alkenyl, —C₂₋₆-alkynyl, —OH, —O, —SH, —S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, C(=O)CHF₂, and —C(=O)NH₂;

3-14-membered cycloalkyl selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl; in each case unsubstituted, mono- or polysubstituted with substituents independently of one another, selected from the group consisting of —F, —Cl, —Br, —I, —C₁₋₆-alkyl, C₂₋₆-alkenyl, —C₂₋₆-alkynyl, —OH, —O, —SH, —S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, C(=O)CHF₂, and —C(=O)NH₂;

10

3-14-membered heterocycloalkyl selected from the group consisting of azepane, 1,4-oxazepane, azetane, azetidine, aziridine, azocane, diazepane, dioxane, dioxolane, dithiane, dithiolane, imidazolidine, isothiazolidine, isoxalidine, morpholine, oxazolidine, oxepane, oxetane, oxirane, piperazine, piperidine, pyrazolidine, pyrrolidine, quinuclidine, tetrahydrofuran, tetrahydropyran, tetrahydrothiopyran, thiazolidine, thietane, thiirane, thiolane, thiomorpholine, indoline, dihydrobenzofuran, dihydrobenzothiophene, 1,1-dioxothiacyclohexane, 2-azaspiro[3.3]heptane, 2-oxaspiro[3.3]heptane, 7-azaspiro[3.5]nonane, 8-azabicyclo[3.2.1]octane, 9-azabicyclo[3.3.1]nonane, hexahydro-1H-pyrrolizine, hexahydrocyclopenta[c]pyrrole, octahydrocyclopenta[c]pyrrole, and octahydropyrrolo[1,2-a]pyrazin, in each case unsubstituted, mono- or polysubstituted with substituents independently of one another, selected from the group consisting of —F, —Cl, —Br, —I, —C₁₋₆-alkyl, C₂₋₆-alkenyl, —C₂₋₆-alkynyl, —OH, —O, —SH, —S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, C(=O)CHF₂, and —C(=O)NH₂.

Preferably, V is unsubstituted, mono- or polysubstituted with substituents independently of one another selected from —F, —Cl, —CN, —OH, —O, —C₁₋₆-alkyl, —CHF₂, —CF₃, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NHC(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-NHC(=O)—O—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkyl, —N(C₁₋₆-alkyl)₂, —OC₁₋₆-alkyl, —OCF₃, —O—C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —S(=O)₂—C₁₋₆-alkyl, -azetidine, —C₁₋₆-alkylene-O-tetrahydropyran, or -piperazine substituted with —C₁₋₆-alkyl.


In a preferred embodiments of the benzofuran derivative according to the invention V is

- (i) unsubstituted;
- (ii) monosubstituted;
- (iii) disubstituted;
- (iv) trisubstituted; or
- (v) tetrasubstituted.

In a preferred embodiments of the benzofuran derivative according to the invention V is








- (i) unsubstituted;
- (ii) monosubstituted; or
- (iii) disubstituted.

In preferred embodiments, V represents a 3-14-membered heterocycloalkyl (preferably 5-membered heterocycloalkyl), saturated or unsaturated; or 5-14-membered heteroaryl (preferably 5-membered heteroaryl); in each case unsubstituted, mono- or polysubstituted; preferably a residue selected from the group consisting of:




11

-continued




12

-continued



In a preferred embodiment, V represents -oxetanyl, unsubstituted, mono- or polysubstituted; preferably



In preferred embodiments, V represents a residue according to general formula (E)




wherein

Y^{E1} represents $-N=$, $-NR^{E2}-$, S, O, or $-CR^{E3}=$; Y^{E2} represents $-N=$, $-NR^{E3}-$, S, O, or $-CR^{E4}=$; and Y^{E3} represent $-N=$, $-NR^{E4}-$, S, O, or $-CR^{E5}=$; with the proviso that at least one of Y^{E1} , Y^{E2} and Y^{E3} is not $-CR^{E3}=$, $-CR^{E4}=$, and $-CR^{E5}=$, respectively. In another preferred embodiment, V represents a residue according to general formula (E) wherein Y^{E1} represents $-N=$, $-NR^{E2}-$, S, or $-CR^{E3}=$; Y^{E2} represents $-N=$, $-NR^{E3}-$, S, or $-CR^{E4}=$; and Y^{E3} represent $-N=$, $-NR^{E4}-$, S, or $-CR^{E5}=$; with the proviso that at least one of Y^{E1} , Y^{E2} , and Y^{E3} is not $-CR^{E3}=$, $-CR^{E4}=$, and $-CR^{E5}=$, respectively.


R^{E1} , R^{E2} , R^{E3} and R^{E4} independently of one another represent $-H$, $-CH_3$, $-CH_2$ -cyclopropyl, $-CH_2CF_3$, $-CH_2CHF_2$ or $-CF_3$; more in particular R^{E1} , R^{E2} , R^{E3} and R^{E4} independently of one another represent $-H$, $-CH_3$, or $-CF_3$; preferably with the proviso that only one of R^{E1} , R^{E2} , R^{E3} and R^{E4} represents a residue that is not $-H$.

In preferred embodiments, V represents 2-pyridine, unsubstituted, mono- or polysubstituted. In preferred embodiments, V represents a residue selected from the group consisting of:

13




In preferred embodiments, V represents 3-pyridine, unsubstituted, mono- or polysubstituted. In preferred embodiments, V represents a residue selected from the group consisting of:




14


-continued



In preferred embodiments, V represents 4-pyridine, unsubstituted, mono- or polysubstituted. In preferred embodiments, V represents a residue selected from the group consisting of:




In preferred embodiments, V represents a residue selected from the group consisting of:




In preferred embodiments, V represents a bicyclic heteroaryl, unsubstituted, mono- or polysubstituted, preferably selected from the group consisting of:

15




In preferred embodiments, V represents a residue according to general formula (F')



wherein

Y^{F1} represents $-N=$ or $-CR^{F4}=$; and Y^{F2} represents $-N=$ or $-CR^{F5}=$; and Y^{F3} represents $-N=$ or $-CR^{F3}=$; with the proviso that at least one of Y^{F1} and Y^{F2} is not $-CR^{F4}=$ and $-CR^{F5}=$, respectively; R^{F1} , R^{F2} , R^{F3} , R^{F4} , and R^{F5} independently of one another represent $-H$, $-CH_3$, $-CF_3$, $-OH$, $-OCH_3$, $-OCH_2CH_3$, $-Cl$, or $-azetidiny$; preferably with the proviso that only one of R^{F1} , R^{F2} , R^{F3} , R^{F4} , and R^{F5} represents a residue that is not $-H$.

In another preferred embodiment, V represents a residue according to general formula (F)




wherein

Y^{F1} represents $-N=$ or $-CR^{F4}=$; and Y^{F2} represents $-N=$ or $-CR^{F5}=$; with the proviso that at least one of Y^{F1} and Y^{F2} is not $-CR^{F4}=$ and $-CR^{F5}=$, respectively; R^{F1} , R^{F2} , R^{F3} , R^{F4} , and R^{F5} independently of one another represent $-H$, $-CH_3$, $-CF_3$, $-OH$, $-OCH_3$, $-OCH_2CH_3$, $-Cl$, or $-azetidiny$; preferably with the proviso that only one of R^{F1} , R^{F2} , R^{F3} , R^{F4} , and R^{F5} represents a residue that is not $-H$.


16

In preferred embodiments, V represents a residue according to general formula (G) or (H)

5 (G)




10 (H)




20 wherein R^{G1} and R^{H1} are selected from the group consisting of $-H$, $-CH_3$, $-CF_3$, $-OH$, $-OCH_3$, $-OCH_2CH_3$, $-Cl$, azetidiny, -cyclopropyl, -O-cyclopropyl, and $-CHF_2$; or wherein R^{G1} and R^{H1} are selected from the group consisting of $-H$, $-CH_3$, $-CF_3$, $-OH$, $-OCH_3$, $-OCH_2CH_3$, $-Cl$, and azetidiny.

In other preferred embodiments, V represents a residue according to general formula (G') or (H')

30 (G')



35 (H')



45 wherein R^{G1} and R^{H1} are selected from the group consisting of $-H$, $-CH_3$, $-CF_3$, $-OH$, $-OCH_3$, $-OCH_2CH_3$, $-Cl$, azetidiny, -cyclopropyl, -O-cyclopropyl, and $-CHF_2$; or wherein R^{G1} and R^{H1} are selected from the group consisting of $-H$, $-CH_3$, $-CF_3$, $-OH$, $-OCH_3$, $-OCH_2CH_3$, $-Cl$, and azetidiny;

In a preferred embodiment of the benzofuran derivative according to the invention R^1 represents

55 $-H$, $-F$, $-Cl$, $-Br$, $-I$, $-CN$;
 $-C_{1-6}$ -alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; $-O-C_{1-6}$ -alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
 $-C(=O)C_{1-6}$ -alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
 $-C(=O)OC_{1-6}$ -alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
 $-C(=O)NHC_{1-6}$ -alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
 $-C(=O)N(C_{1-6}\text{-alkyl})_2$, saturated or unsaturated, unsubstituted, mono- or polysubstituted;

17

—S(=O)C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
 —S(=O)₂—C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
 —C₁-C₆-heteroalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; or
 3-14-membered cycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered cycloalkyl is optionally connected through —C₁-C₆-alkylene- or —C₁-C₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted.

Preferably, R¹ represents —H, —F, —Cl, —Br, —I, —C₁₋₆-alkyl, —O—C₁₋₆-alkyl, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH(C₁₋₆-alkyl), —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —CF₃, —CF₂H, —CFH₂, —CF₂Cl, —CFCl₂, —C₁₋₆-alkylene-CF₃, —C₁₋₆-alkylene-CF₂H, —C₁₋₆-alkylene-CFH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)-C₁₋₆-alkylene-CF₃, —C(=O)C₁₋₆-alkyl, —C(=O)OC₁₋₆-alkyl, —C(=O)NH₂, —C(=O)NHC₁₋₆-alkyl, —C(=O)N(C₁₋₆-alkyl)₂, —S(=O)—C₁₋₆-alkyl, —S(=O)₂—C₁₋₆-alkyl, —O—C₁₋₆-alkyl, -cyclopropyl unsubstituted, cyclobutyl unsubstituted, cyclopentyl unsubstituted or cyclohexyl unsubstituted.

Preferably, R¹ represents —H, —C₁₋₆-alkyl, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —CH₂F, —CHF₂, —CF₃, -cyclopentyl, unsubstituted, or -cyclopropyl. Preferably, R¹ represents —H, —C₁₋₆-alkyl, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —CH₂F, —CHF₂, —CF₃, -cyclopentyl, or unsubstituted. Preferably, R¹ represents —CH₃.

Preferably, R¹ represents —CH₂F, —CHF₂, —CH₃, or -cyclopropyl. Preferably, R¹ represents —CH₂F, —CHF₂, or —CH₃. Preferably, R¹ represents —C(=O)NH₂, or —CHF₂.

Preferably, R¹ represents —H, —C₁₋₃-alkyl, —CF₃, —CF₂H, —CFH₂, —CF₂Cl, —CFCl₂, —C₁₋₃-alkylene-CF₃, —C₁₋₃-alkylene-CF₂H, —C₁₋₃-alkylene-CFH₂, or -cyclopropyl; preferably, R¹ represents —H, —C₁₋₃-alkyl, —CF₃, —CF₂H, —CFH₂, —CF₂Cl, —CFCl₂, —C₁₋₃-alkylene-CF₃, —C₁₋₃-alkylene-CF₂H, or —C₁₋₃-alkylene-CFH₂; more preferably —CH₃.

In a preferred embodiment of the benzofuran derivative according to the invention R² represents

—H;
 —C₁-C₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
 —C₁-C₆-heteroalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
 3-14-membered cycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered cycloalkyl is optionally connected through —C₁-C₆-alkylene- or —C₁-C₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted; or
 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered heterocycloalkyl is optionally connected through —C₁-C₆-alkylene- or —C₁-C₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted.

Preferably, R² represents —H, —C₁₋₆-alkyl, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH(C₁₋₆-alkyl), —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —CF₃, —CF₂H, —CFH₂, —CF₂Cl, —CFCl₂, —C₁₋₆-alkylene-CF₃, —C₁₋₆-alkylene-

18

CF₂H, —C₁₋₆-alkylene-CFH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, or —C₁₋₆-alkylene-N(C₁₋₆-alkyl)-C₁₋₆-alkylene-CF₃.

Preferably, R² represents —H or —C₁₋₆-alkyl.

In a preferred embodiment of the benzofuran derivative according to the invention R³ represents

—H;
 —OH;
 —C₁-C₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; or
 —C₁-C₆-heteroalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted.

Preferably, R³ represents —H, —OH, —C₁₋₆-alkyl, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NH(C₁₋₆-alkyl), —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —CF₃, —CF₂H, —CFH₂, —CF₂Cl, —CFCl₂, —C₁₋₆-alkylene-CF₃, —C₁₋₆-alkylene-CF₂H, —C₁₋₆-alkylene-CFH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, or —C₁₋₆-alkylene-N(C₁₋₆-alkyl)-C₁₋₆-alkylene-CF₃.

Preferably, R³ represents —H, —OH, or —C₁₋₆-alkyl, saturated, unsubstituted or monosubstituted with —OH. Preferably, R³ represents —H.

Preferably, R³ represents —H and R⁴ represents a residue other than —H.

In a preferred embodiment of the benzofuran derivative according to the invention R⁴ represents

—H;
 —S(=O)C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
 —S(=O)₂—C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
 —C₁-C₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
 —C₁-C₆-heteroalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
 3-14-membered cycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered cycloalkyl is optionally connected through —C₁-C₆-alkylene- or —C₁-C₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted;

3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered heterocycloalkyl is optionally connected through —C₁-C₆-alkylene- or —C₁-C₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted;

6-14-membered aryl, unsubstituted, mono- or polysubstituted; wherein said 6-14-membered aryl is optionally connected through —C₁-C₆-alkylene- or —C₁-C₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted; or

5-14-membered heteroaryl, unsubstituted, mono- or polysubstituted; wherein said 5-14-membered heteroaryl is optionally connected through —C₁-C₆-alkylene- or —C₁-C₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted.

Preferably, R⁴ represents

—S(=O)₂C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted with substituents independently of one another selected from the group consisting of —F, —Cl, —C₁₋₆-alkyl, —C₁₋₆-alkylene-CF₃, —OH, —O, —OC₁₋₆-alkyl, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —NH₂, —NHC₁₋₆-alkyl, —N(C₁₋₆-alkyl)₂, —NHC(=O)O—C₁₋₆-alkyl, —N(C₁₋₆-alkyl)C(=O)O—C₁₋₆-alkyl, —C₁₋₆-alk-

19

kylene-NHC(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkyl, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, —C(=O)—C₁₋₆-alkyl, —C(=O)OH, —C(=O)O—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkylene-CF₃, —C(=O)NH₂, —C(=O)NH(C₁₋₆-alkyl), —C(=O)N(C₁₋₆-alkyl)₂, —S(=O)₂C₁₋₆-alkyl, -phenyl, —C₁₋₆-alkylene-phenyl, 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted; and 5-14-membered heteroaryl, unsubstituted;

—S(=O)₂(3-14-membered cycloalkyl), wherein said 3-14-membered cycloalkyl is selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted with substituents independently of one another selected from the group consisting of —F, —Cl, —C₁₋₆-alkyl, —C₁₋₆-alkylene-CF₃, —OH, =O, —OC₁₋₆-alkyl, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —NH₂, —NHC₁₋₆-alkyl, —N(C₁₋₆-alkyl)₂, —NHC(=O)O—C₁₋₆-alkyl, —N(C₁₋₆-alkyl)C(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NHC(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkyl, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, —C(=O)—C₁₋₆-alkyl, —C(=O)OH, —C(=O)O—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkylene-CF₃, —C(=O)NH₂, —C(=O)NH(C₁₋₆-alkyl), —C(=O)N(C₁₋₆-alkyl)₂, —S(=O)₂C₁₋₆-alkyl, -phenyl, —C₁₋₆-alkylene-phenyl, 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted; and 5-14-membered heteroaryl, unsubstituted;

—C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted with substituents independently of one another selected from the group consisting of —F, —Cl, —C₁₋₆-alkyl, —C₁₋₆-alkylene-CF₃, —OH, =O, —OC₁₋₆-alkyl, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —NH₂, —NHC₁₋₆-alkyl, —N(C₁₋₆-alkyl)₂, —NHC(=O)O—C₁₋₆-alkyl, —N(C₁₋₆-alkyl)C(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NHC(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkyl, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, —C(=O)—C₁₋₆-alkyl, —C(=O)OH, —C(=O)O—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkylene-CF₃, —C(=O)NH₂, —C(=O)NH(C₁₋₆-alkyl), —C(=O)N(C₁₋₆-alkyl)₂, —S(=O)₂C₁₋₆-alkyl, -phenyl, —C₁₋₆-alkylene-phenyl, 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted; and 5-14-membered heteroaryl, unsubstituted;

3-14-membered cycloalkyl or —C₁₋₆-alkylene-(3-14-membered cycloalkyl), wherein —C₁₋₆-alkylene- is unsubstituted or monosubstituted with —OH, wherein said 3-14-membered cycloalkyl is selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl, in each case saturated or unsaturated, in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from the group consisting of —F, —Cl, —C₁₋₆-alkyl, —C₁₋₆-alkylene-CF₃, —OH, =O, —OC₁₋₆-alkyl, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —NH₂, —NHC₁₋₆-alkyl, —N(C₁₋₆-alkyl)₂, —NHC(=O)O—C₁₋₆-alkyl, —N(C₁₋₆-alkyl)C(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NHC(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkyl, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, —C(=O)—C₁₋₆-alkyl, —C(=O)OH, —C(=O)O—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkylene-CF₃, —C(=O)NH₂, —C(=O)NH(C₁₋₆-alkyl), —C(=O)N(C₁₋₆-alkyl)₂, —S(=O)₂C₁₋₆-alkyl, -phenyl, —C₁₋₆-alkylene-phenyl, 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted; and 5-14-membered heteroaryl, unsubstituted;

5-14-membered heteroaryl or —C₁₋₆-alkylene-(5-14-membered heteroaryl), wherein —C₁₋₆-alkylene- is unsubstituted or monosubstituted with —OH, wherein said 5-14-membered heteroaryl in each case is selected from the group consisting of benzimidazole, benzisoxazole, benzoazole, benzodioxole, benzofuran, benzothiadiazole, benzothiazole, benzothiophene, carbazole, cinoline, dibenzofuran, furane, furazane, imidazole,

20

C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkylene-CF₃, —C(=O)NH₂, —C(=O)NH(C₁₋₆-alkyl), —C(=O)N(C₁₋₆-alkyl)₂, —S(=O)₂C₁₋₆-alkyl, -phenyl, —C₁₋₆-alkylene-phenyl, 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted; and 5-14-membered heteroaryl, unsubstituted;

3-14-membered heterocycloalkyl or —C₁₋₆-alkylene-(3-14-membered heterocycloalkyl), wherein —C₁₋₆-alkylene- is unsubstituted or monosubstituted with —OH, wherein said 3-14-membered heterocycloalkyl in each case is selected from the group consisting of azepane, 1,4-oxazepane, azetane, azetidine, aziridine, azocane, diazepane, dioxane, dioxolane, dithiane, dithiolane, imidazolidine, isothiazolidine, isoxalidine, morpholine, oxazolidine, oxepane, oxetane, oxirane, piperazine, piperidine, pyrazolidine, pyrrolidine, quinuclidine, tetrahydrofuran, tetrahydropyran, tetrahydrothiopyran, thiazolidine, thietane, thiirane, thiolane, thiomorpholine, indoline, dihydrobenzofuran, dihydrobenzothiophene, 1,1-dioxothiacyclohexane, 2-azaspiro[3.3]heptane, 2-oxaspiro[3.3]heptane, 7-azaspiro[3.5]nonane, 8-azabicyclo[3.2.1]octane, 9-azabicyclo[3.3.1]nonane, hexahydro-1H-pyrrolizine, hexahydrocyclopenta[c]pyrrole, octahydrocyclopenta[c]pyrrole, and octahydropyrrolo[1,2-a]pyrazine; in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from the group consisting of —F, —Cl, —C₁₋₆-alkyl, —C₁₋₆-alkylene-CF₃, —OH, =O, —OC₁₋₆-alkyl, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —NH₂, —NHC₁₋₆-alkyl, —N(C₁₋₆-alkyl)₂, —NHC(=O)O—C₁₋₆-alkyl, —N(C₁₋₆-alkyl)C(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NHC(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkyl, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, —C(=O)—C₁₋₆-alkyl, —C(=O)OH, —C(=O)O—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkylene-CF₃, —C(=O)NH₂, —C(=O)NH(C₁₋₆-alkyl), —C(=O)N(C₁₋₆-alkyl)₂, —S(=O)₂C₁₋₆-alkyl, -phenyl, —C₁₋₆-alkylene-phenyl, 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted; and 5-14-membered heteroaryl, unsubstituted;

-phenyl unsubstituted, mono- or polysubstituted with substituents independently of one another selected from the group consisting of —F, —Cl, —CN, —C₁₋₆-alkyl, —C₁₋₆-alkylene-CF₃, —OH, =O, —OC₁₋₆-alkyl, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —NH₂, —NHC₁₋₆-alkyl, —N(C₁₋₆-alkyl)₂, —NHC(=O)O—C₁₋₆-alkyl, —N(C₁₋₆-alkyl)C(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NHC(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkyl, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, —C(=O)—C₁₋₆-alkyl, —C(=O)OH, —C(=O)O—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkylene-CF₃, —C(=O)NH₂, —C(=O)NH(C₁₋₆-alkyl), —C(=O)N(C₁₋₆-alkyl)₂, —S(=O)₂C₁₋₆-alkyl, -phenyl, —C₁₋₆-alkylene-phenyl, 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted; and 5-14-membered heteroaryl, unsubstituted;

5-14-membered heteroaryl or —C₁₋₆-alkylene-(5-14-membered heteroaryl), wherein —C₁₋₆-alkylene- is unsubstituted or monosubstituted with —OH, wherein said 5-14-membered heteroaryl in each case is selected from the group consisting of benzimidazole, benzisoxazole, benzoazole, benzodioxole, benzofuran, benzothiadiazole, benzothiazole, benzothiophene, carbazole, cinoline, dibenzofuran, furane, furazane, imidazole,

21

imidazopyridine, indazole, indole, indolizine, isobenzofuran, isoindole, isoquinoline, isothiazole, isoxazole, naphthyridine, oxadiazole, oxazole, oxindole, phthalazine, purine, pyrazine, pyrazole, pyridazine, pyridine, pyrimidine, pyrrole, quinazoline, quinoline, quinoxaline, tetrazole, thiadiazole, thiazole, thiophene, triazine, triazole, and [1,2,4]triazolo[4,3-a]pyrimidine; in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from the group consisting of —F, —Cl, —CN, —C₁₋₆-alkyl, —C₁₋₆-alkylene-CF₃, —OH, =O, —OC₁₋₆-alkyl, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —NH₂, —NHC₁₋₆-alkyl, —N(C₁₋₆-alkyl)₂, —NHC(=O)—C₁₋₆-alkyl, —N(C₁₋₆-alkyl)C(=O)—C₁₋₆-alkyl, —C₁₋₆-alkylene-NHC(=O)—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkyl, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, —C(=O)—C₁₋₆-alkyl, —C(=O)OH, —C(=O)O—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkylene-CF₃, —C(=O)NH₂, —C(=O)NH(C₁₋₆-alkyl), —C(=O)N(C₁₋₆-alkyl)₂, —S(=O)₂C₁₋₆-alkyl, —phenyl, —C₁₋₆-alkylene-phenyl, 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted; and 5-14-membered heteroaryl, unsubstituted.

Preferably, R⁴ represents

—H;
—S(=O)₂C₁₋₆-alkyl, saturated, unsubstituted, monosubstituted or polysubstituted with —F;
—S(=O)₂(3-14-membered cycloalkyl), saturated, unsubstituted;
—C₁₋₆-alkyl, saturated, unsubstituted, monosubstituted or disubstituted with substituents independently of one another selected from the group consisting of —OH, =O, —NH₂, —NHC₁₋₆-alkyl, —N(C₁₋₆-alkyl)₂, —OC₁₋₆-alkyl, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkyl, —C(=O)NH₂, —C(=O)—NH—C₁₋₃-alkyl, —C(=O)—N(C₁₋₃-alkyl)₂, —phenyl unsubstituted;

3-14-membered cycloalkyl or —C₁₋₆-alkylene-(3-14-membered cycloalkyl), wherein —C₁₋₆-alkylene- is unsubstituted or monosubstituted with —OH, wherein said 3-14-membered cycloalkyl is saturated, unsubstituted, monosubstituted or disubstituted with substituents independently of one another selected from the group consisting of —C₁₋₆-alkyl, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-NHC(=O)—C₁₋₆-alkyl, —OH, —OC₁₋₆-alkyl, —NH₂, —N(C₁₋₆-alkyl)₂, —NHC(=O)—C₁₋₆-alkyl;

3-14-membered heterocycloalkyl or —C₁₋₆-alkylene-(3-14-membered heterocycloalkyl), wherein —C₁₋₆-alkylene- is unsubstituted or monosubstituted with —OH, wherein said 3-14-membered heterocycloalkyl in each case is selected from azetane, 1,4-oxazepane, pyrrolidine, piperidine, azepane, diazepane, tetrahydrofuran, tetrahydropyran, oxetane, morpholine, piperazine, hexahydrocyclopenta[c]pyrrole, octahydrocyclopenta[c]pyrrole, octahydropyrrolo[1,2-a]pyrazin, 8-azabicyclo[3.2.1]octane, 9-azabicyclo[3.3.1]nonane, quinuclidine, hexahydro-1H-pyrrolizine, 2-oxaspiro[3.3]heptane, 2-azaspiro[3.3]heptane, 7-azaspiro[3.5]nonane, 1,1-dioxothiacyclohexane, in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from the group consisting of —F, —OH, =O, —C₁₋₆-alkyl, —C₁₋₆-alkylene-CF₃, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —NH₂, —N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —C(=O)—C₁₋₆-alkyl, —C(=O)OH, —C(=O)O—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkylene-CF₃, —C(=O)NH₂, —C(=O)NH(C₁₋₆-alkyl), —C(=O)N(C₁₋₆-alkyl)₂, —S(=O)₂C₁₋₆-alkyl, —phenyl, —C₁₋₆-alkylene-phenyl, 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted; and 5-14-membered heteroaryl, unsubstituted.

22

—C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —C(=O)—C₁₋₆-alkyl, —C(=O)OH, —C(=O)O—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkylene-CF₃, —C(=O)NH₂, —C(=O)NH(C₁₋₆-alkyl), —S(=O)₂C₁₋₆-alkyl, oxetanyl, pyrimidinyl, —C₁₋₆-alkylene-phenyl;

—phenyl unsubstituted;

5-14-membered heteroaryl or —C₁₋₆-alkylene-(5-14-membered heteroaryl), wherein —C₁₋₆-alkylene- is unsubstituted or monosubstituted with —OH, wherein said 5-14-membered heteroaryl in each case is selected from the group consisting of pyridine, pyridazine, pyrazine, pyrazole, isoxazole, triazole, and [1,2,4]triazolo[4,3-a]pyrimidine, in each case unsubstituted, monosubstituted or disubstituted with substituents independently of one another selected from the group consisting of —C₁₋₆-alkyl, —OH.

In a preferred embodiment of the benzofuran derivative according to the invention R³ and R⁴ together form a 5- or 6-membered heterocycle containing 1 or 2 heteroatoms selected from N, O and S, saturated or unsaturated, unsubstituted or mono- or polysubstituted.

Preferably, R³ and R⁴ together form a heterocycle selected from the group consisting of pyrrolidine, piperidine, morpholine, and piperazine, in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from the group consisting of —F, —C₁₋₆-alkyl, —NH₂, —NHCH₃, —N(CH₃)₂, —C(=O)NH—C₁₋₆-alkyl, —C(=O)N(C₁₋₆-alkyl)₂, —C(=O)O—C₁₋₆-alkyl, —NHC(=O)—C₁₋₆-alkyl, —pyridyl unsubstituted, and 1,2,4-oxadiazole unsubstituted or monosubstituted with —C₁₋₆-alkyl. In a preferred embodiment, R³ and R⁴ together do not form morpholine unsubstituted, mono- or polysubstituted.

Preferably, R³ and R⁴ together form a


pyrrolidine ring, unsubstituted or monosubstituted with —N(CH₃)₂;

piperidine ring, unsubstituted or monosubstituted with a substituent selected from the group consisting of —C₁₋₆-alkyl, —NH₂, —N(CH₃)₂, —C(=O)NH—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkyl, —NHC(=O)—C₁₋₆-alkyl, and 1,2,4-oxadiazole unsubstituted or monosubstituted with —C₁₋₆-alkyl;

morpholine ring, unsubstituted; or


piperazine ring, unsubstituted or N-substituted with a substituent selected from the group consisting of —C₁₋₆-alkyl and —pyridyl unsubstituted.

In a preferred embodiment, R³ and R⁴ both do not represent —H. In preferred embodiments, R³ and R⁴ together with the nitrogen atom to which they are attached form a residue selected from the group consisting of:




23

-continued




In other preferred embodiments R^3 represents $-H$ and R^4 does not represent $-H$.

In preferred embodiments, R^3 represents $-H$ and R^4 represent C_1 - C_6 -alkyl saturated or unsaturated, unsubstituted, mono- or polysubstituted. In preferred embodiments, R^3 represents $-H$ and R^4 represents a residue selected from the group consisting of:



24


-continued




In preferred embodiments, R^3 represents $-H$ and R^4 represents a residue $-CR'R''-(CH_2)_m-OH$, wherein m is an integer of from 1 to 6, preferably from 1 to 3; and wherein R' and R'' independently of one another represent $-H$, $-C_{1-3}$ -alkyl, $-CF_3$, $-CF_2H$, $-CFH_2$, $-C_{1-3}$ -alkylene- CF_3 , $-C_{1-3}$ -alkylene- CF_2H , $-C_{1-3}$ -alkylene- CFH_2 , $-C_{1-3}$ -alkylene- $O-C_{1-3}$ -alkyl, $-C_{1-3}$ -alkylene- OH , $-C(=O)-NH_2$, or $C(=O)-NH-C_{1-3}$ -alkyl; preferably $-H$, $-CH_3$, $-C_{1-3}$ -alkylene- OH , $-C(=O)-NH_2$, or $C(=O)-NH-C_{1-3}$ -alkyl. In a preferred embodiment, at least R' or R'' does not represent $-H$. In a preferred embodiment, neither R' nor R'' represents $-H$.

In other preferred embodiments, R^3 represents $-H$ and R^4 represents a 3-14-membered cycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; or a 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted, mono- or poly substituted.

In preferred embodiments, R^3 represents $-H$ and R^4 represents a 3-membered cycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; or a 3-membered heterocycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted. In preferred embodiments, R^3 represents $-H$ and R^4 represents a residue selected from the group consisting of:




In preferred embodiments, R^3 represents $-H$ and R^4 represents a 3-14-membered cycloalkyl (preferably a 4-membered cycloalkyl), saturated or unsaturated, unsubstituted, mono- or polysubstituted; or a 3-14-membered heterocycloalkyl (preferably a 4-membered heterocycloalkyl), saturated or unsaturated, unsubstituted, mono- or polysubstituted. In preferred embodiments, R^3 represents $-H$ and R^4 represents a residue selected from the group consisting of:




25

-continued



In preferred embodiments, R^3 represents $-H$ and R^4 represents a residue according to general formula (A),



wherein

m^4 is 0 or 1;

Y^4 is selected from $-O-$, $-NR^{A6}-$ and $-CR^{A7}R^{A8}-$; and

R^{A1} , R^{A2} , R^{A3} , R^{A4} , R^{A5} , R^{A6} , R^{A7} , and R^{A8} independently of one another represent $-H$, F , $-C_{1-3}$ -alkyl, $-C_{1-3}$ -alkylene-OH, $-C_{1-3}$ -alkylene-NH $_2$, $-C_{1-3}$ -alkylene-NH(C $_{1-3}$ -alkyl), $-C_{1-3}$ -alkylene-N(C $_{1-3}$ -alkyl) $_2$, $-C_{1-3}$ -alkylene-NH(C $_{1-3}$ -alkylene-CF $_3$), $-C_{1-3}$ -alkylene-C(=O)NH $_2$, $-C_{1-3}$ -alkylene-NH-C(=O)

26

OC $_{1-4}$ -alkyl, $-C(=O)NH_2$, $-C(=O)NH-C_{1-3}$ -alkyl, $-C(=O)N(C_{1-3}$ -alkyl) $_2$, -3 -oxetanyl, or $-CHF_2$; preferably, R^{A1} , R^{A2} , R^{A3} , R^{A4} , R^{A5} , R^{A6} , R^{A7} , and R^{A8} independently of one another represent $-H$, F , $-C_{1-3}$ -alkyl, $-C_{1-3}$ -alkylene-OH, $-C_{1-3}$ -alkylene-NH $_2$, $-C_{1-3}$ -alkylene-NH(C $_{1-3}$ -alkyl), $-C_{1-3}$ -alkylene-N(C $_{1-3}$ -alkyl) $_2$, $-C_{1-3}$ -alkylene-NH(C $_{1-3}$ -alkylene-CF $_3$), $-C_{1-3}$ -alkylene-C(=O)NH $_2$, $-C_{1-3}$ -alkylene-NH-C(=O)OC $_{1-4}$ -alkyl, $-C(=O)NH_2$, $-C(=O)NH-C_{1-3}$ -alkyl, $-C(=O)N(C_{1-3}$ -alkyl) $_2$, or -3 -oxetanyl; or R^{A7} and R^{A8} together with the carbon atom to which they are attached form a ring and represent $-CH_2OCH_2-$, $-CH_2OCH_2CH_2-$ or $-CH_2CH_2OCH_2CH_2-$, $-CH_2NHCH_2-$, $-CH_2NHCH_2CH_2-$ or $-CH_2CH_2NHCH_2CH_2-$.

In preferred embodiments, R^3 represents $-H$ and R^4 represents a residue according to general formula (A) as defined above, wherein

m^4 is 0 or 1;

Y^4 is selected from $-O-$ and $-CR^{A7}R^{A8}-$; and

R^{A1} , R^{A2} , R^{A3} , R^{A4} , R^{A5} , R^{A7} , and R^{A8} independently of one another represent $-H$, $-C_{1-3}$ -alkylene-OH, $-C_{1-3}$ -alkylene-N(C $_{1-3}$ -alkyl) $_2$, $-C(=O)NH_2$, or $-CHF_2$; preferably R^{A1} , R^{A2} , R^{A3} , R^{A4} , R^{A5} , R^{A7} , and R^{A8} independently of one another represent $-H$, $-C_{1-3}$ -alkylene-OH, $-C_{1-3}$ -alkylene-N(C $_{1-3}$ -alkyl) $_2$, or $-C(=O)NH_2$; preferably with the proviso that only one of R^{A1} , R^{A2} , R^{A3} , R^{A4} , R^{A5} , R^{A7} , and R^{A8} represents a residue that is not $-H$.

In preferred embodiments, R^3 represents $-H$ and R^4 represents a residue according to general formula (A) as defined above, wherein


m^4 is 0 or 1;

Y^4 is selected from $-O-$ and $-CR^{A7}R^{A8}-$; and

R^{A1} represents $-C_{1-3}$ -alkylene-OH, $-C_{1-3}$ -alkylene-N(C $_{1-3}$ -alkyl) $_2$, $-C(=O)NH_2$, or $-CHF_2$; preferably R^{A1} represents $-C_{1-3}$ -alkylene-OH, $-C_{1-3}$ -alkylene-N(C $_{1-3}$ -alkyl) $_2$, or $-C(=O)NH_2$; and


R^{A1} , R^{A2} , R^{A3} , R^{A4} , R^{A5} , R^{A7} , and R^{A8} represent $-H$.

In preferred embodiments, R^3 represents $-H$ and R^4 represents a 3-14-membered cycloalkyl (preferably a 5-membered cycloalkyl), saturated or unsaturated, unsubstituted, mono- or polysubstituted; or a 3-14-membered heterocycloalkyl (preferably a 5-membered heterocycloalkyl), saturated or unsaturated, unsubstituted, mono- or polysubstituted; or a 5-14-membered heteroaryl (preferably a 5-membered heteroaryl), unsubstituted, mono- or polysubstituted. In preferred embodiments, R^3 represents $-H$ and R^4 represents a residue selected from the group consisting of:




27

-continued




28

-continued



29

In preferred embodiments, R^3 represents $-H$ and R^4 represents a residue according to general formula (B),




wherein

Y^B is selected from $-O-$, $-NR^{B8}-$ and $-CR^{B9}R^{B10}-$; and R^{B1} , R^{B2} , R^{B3} , R^{B4} , R^{B5} , R^{B6} , R^{B7} , R^{B8} , R^{B9} and R^{B10} independently of one another represent $-H$, $-F$, $-OH$, $-C_{1-3}$ -alkyl, $-C_{1-3}$ -alkylene- OH , $-C_{1-3}$ -alkylene- $O-C_{1-3}$ -alkyl, $-C_{1-3}$ -alkylene- CF_3 , $-C_{1-3}$ -alkylene- CO_2H , $-C_{1-3}$ -alkylene- $C(=O)O-C_{1-3}$ -alkyl, $-C(=O)NH_2$, $-C(=O)NH-C_{1-3}$ -alkyl, or $-C(=O)N(C_{1-3}\text{-alkyl})_2$; or R^{B2} and R^{B3} together represent $=O$; or R^{B4} and R^{B5} together represent $=O$.

In preferred embodiments, R^3 represents $-H$ and R^4 represents a residue according to general formula (B) as defined above, wherein

Y^B is selected from $-O-$ and $-NR^{B8}-$; and R^{B1} , R^{B2} , R^{B3} , R^{B4} , R^{B5} , R^{B6} , R^{B7} , R^{B8} independently of one another represent $-H$, $-F$, $-C_{1-3}$ -alkyl, $-C_{1-3}$ -alkylene- OH , $-C_{1-3}$ -alkylene- CF_3 or $-C(=O)NH_2$; or R^{B2} and R^{B3} together represent $=O$; or R^{B4} and R^{B5} together represent $=O$; preferably with the proviso that only 1, 2 or 3 of R^{A1} , R^{A2} , R^{A3} , R^{A4} , R^{A5} , R^{A7} , and R^{A8} represent a residue that is not $-H$; preferably with the proviso that at least one of R^{A1} , R^{A2} , R^{A3} , R^{A4} , R^{A5} , R^{A7} , and R^{A8} represent a residue that is not $-H$.

In preferred embodiments, R^3 represents $-H$ and R^4 represents a 3-14-membered cycloalkyl (preferably a 6-membered cycloalkyl), saturated or unsaturated, unsubstituted, mono- or polysubstituted; or a 3-14-membered heterocycloalkyl (preferably a 6-membered heterocycloalkyl), saturated or unsaturated, unsubstituted, mono- or polysubstituted; or a 6-14-membered aryl (preferably a 6-membered aryl), unsubstituted, mono- or polysubstituted; or a 5-14-membered heteroaryl (preferably a 6-membered heteroaryl), unsubstituted, mono- or polysubstituted. In preferred embodiments, R^3 represents $-H$ and R^4 represents a residue selected from the group consisting of:



30

-continued

(B) 5

10

15

20

25

30

35

40


45

50

55


60

65




31

-continued




32

-continued



In preferred embodiments, R^3 represents $-H$ and R^4 represents a residue according to general formula (C),



wherein

Y^{C1} is selected from $-O-$, $-S(=O)_2-$, $-NR^{C8}-$ and $-CR^{C9}R^{C10}-$ and Y^{C2} represents $-CR^{C11}R^{C12}-$, or Y^{C1} represents $-CR^{C9}R^{C10}-$ and Y^{C2} is selected from $-O-$, $-S(=O)_2-$, and $-NR^{C8}-$;

R^{C1} , R^{C2} , R^{C3} , R^{C4} , R^{C5} , R^{C6} , R^{C7} , R^{C8} , R^{C9} , R^{C10} , R^{C11} and R^{C12} independently of one another represent $-H$, $-F$, $-OH$, $-C(=O)OC_{1-3}\text{-alkyl}$, $-NH_2$, $-NH(C_{1-3}\text{-alkyl})$, $-N(C_{1-3}\text{-alkyl})_2$, $-C_{1-3}\text{-alkyl}$, $-C_{1-3}\text{-alkylene-OH}$, $-C_{1-3}\text{-alkylene-}$, $-C(=O)NH_2$, $-C(=O)NH-C_{1-3}\text{-alkyl}$, or $-C(=O)N(C_{1-3}\text{-alkyl})_2$; or R^{C2} and R^{C3} together represent $=O$; or R^{C4} and R^{C5} together represent $=O$; or R^{C9} and R^{C10} together represent $=O$; or R^{C11} and R^{C12} together represent $=O$.

In preferred embodiments, R^3 represents $-H$ and R^4 represents a residue according to general formula (C) as defined above, wherein


Y^{C1} is selected from $-O-$ or $-NR^{C8}-$ and Y^{C2} represents $-CR^{C11}R^{C12}-$, or Y^{C1} represents $-CR^{C9}R^{C10}-$ and Y^{C2} is selected from $-O-$, and $-NR^{C8}-$;

R^{C1} , R^{C2} , R^{C3} , R^{C4} , R^{C5} , R^{C6} , R^{C7} , R^{C8} , R^{C9} , R^{C10} , R^{C11} and R^{C12} independently of one another represent $-H$,


33

—F, —C₁₋₃-alkyl, —C₁₋₃-alkylene-OH, or —C(=O)NH₂; preferably with the proviso that only 1, 2 or 3 of R^{C1}, R^{C2}, R^{C3}, R^{C4}, R^{C5}, R^{C6}, R^{C7}, R^{C8}, R^{C9}, R^{C10}, R^{C11} and R^{C12} represent a residue that is not —H; preferably with the proviso that at least one of R^{C1}, R^{C2}, R^{C3}, R^{C4}, R^{C5}, R^{C6}, R^{C7}, R^{C8}, R^{C9}, R^{C10}, R^{C11} and R^{C12} represent a residue that is not —H.

In preferred embodiments, R³ represents —H and R⁴ represents a 7-membered cycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; or a 7-membered heterocycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted. In preferred embodiments, R³ represents —H and R⁴ represents a residue:




In preferred embodiments, R³ represents —H and R⁴ represents a 3-14-membered cycloalkyl (preferably a 4, 5 or 6-membered cycloalkyl), saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered cycloalkyl is connected through —C₁-C₆-alkylene-, saturated or unsaturated, unsubstituted, mono- or polysubstituted; or a 3-14-membered heterocycloalkyl (preferably a 4, 5 or 6-membered heterocycloalkyl), saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered heterocycloalkyl is connected through —C₁-C₆-alkylene-, saturated or unsaturated, unsubstituted, mono- or polysubstituted; or a 6-14-membered aryl (preferably a 6-membered aryl), unsubstituted, mono- or polysubstituted; wherein said 36-14-membered aryl is connected through —C₁-C₆-alkylene-, saturated or unsaturated, unsubstituted, mono- or polysubstituted; or a 5-14-membered heteroaryl (preferably a 5 or 6-membered heteroaryl), unsubstituted, mono- or polysubstituted; wherein said 5-14-membered heteroaryl is connected through —C₁-C₆-alkylene-, saturated or unsaturated, unsubstituted, mono- or polysubstituted. In preferred embodiments, R³ represents —H and R⁴ represents a residue selected from the group consisting of:



34

-continued




In preferred embodiments, R³ represents —H and R⁴ represents a 5-membered heterocycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 5-membered heterocycloalkyl is connected through —C₁-C₆-alkylene-, saturated or unsaturated, unsubstituted, mono- or polysubstituted; or a 5-membered het-

35


eroaryl, unsubstituted, mono- or polysubstituted; wherein said 5-membered heteroaryl is connected through $-C_1-C_6-$ alkylene-, saturated or unsaturated, unsubstituted, mono- or poly substituted.

In preferred embodiments, R^3 represents $-H$ and R^4 represents a residue selected from the group consisting of:



In preferred embodiments, R^3 represents $-H$ and R^4 represents

- (i) a residue $-CR'R''-(CH_2)_m-OH$, wherein m is an integer of from 1 to 6, preferably from 1 to 3; and wherein R' and R'' independently of one another represent $-H$, $-C_{1-3}$ -alkyl, $-CF_3$, $-CF_2H$, $-CFH_2$, $-C_{1-3}$ -alkylene- CF_3 , $-C_{1-3}$ -alkylene- CF_2H , $-C_{1-3}$ -alkylene- CFH_2 , $-C_{1-3}$ -alkylene- $O-C_{1-3}$ -alkyl, or $-C_{1-3}$ -alkylene- OH ; preferably $-H$, $-CH_3$, or $-C_{1-3}$ -alkylene- OH . In a preferred embodiment, at least R' or R'' does not represent $-H$. In a preferred embodiment, neither R' nor R'' represents $-H$; or
- (ii) a residue according to general formula (D),



wherein

m^D and n^D independently of one another are 0, 1, 2, or 3; preferably with the proviso that $m^D+n^D \leq 3$;

Y^{D1} is selected from $-O-$, $-S(=O)_2-$, $-S(=O)(=NH)-$, $-NR^{D8}-$ and $-CR^{D9}R^{D10}-$ and Y^{D2} represents $-CR^{D11}R^{D12}-$;

or Y^{D1} is selected from $-O-$, $-S(=O)_2-$, $-NR^{D8}-$ and $-CR^{D9}R^{D10}-$ and Y^{D2} represents $-CR^{D11}R^{D12}-$; or Y^{D1} represents $-CR^{D9}R^{D10}-$ and Y^{D2} is selected from $-O-$, $-S(=O)_2-$, and $-NR^{D8}-$;

R^{D1} , R^{D2} , R^{D3} , R^{D4} , R^{D5} , R^{D6} , R^{D7} , R^{D8} , R^{D9} , R^{D10} , R^{D11} and R^{D12} independently of one another represent $-H$, $-F$, $-OH$, $-C_{1-3}$ -alkylene- OH , $-C(=O)NH_2$, $-C_{1-3}$ -alkylene- $C(=O)NH_2$, $-C(=O)O-C_{1-3}$ -alkyl, $-NH_2$, $-C_{1-3}$ -alkylene- NH_2 , $-NH(C_{1-3}$ -alkyl), $-N(C_{1-3}$ -alkyl) $_2$, $-NH(C_{1-3}$ -alkylene- CF_3), $-C_{1-3}$ -alkylene- OCH_3 , $-C_{1-3}$ -alkyl, $-C_{1-3}$ -alkylene- CF_3 ; or R^{D2} and R^{D3} together represent $=O$; or R^{D4} and R^{D5} together represent $=O$; or R^{D9} and R^{D10} together represent $=O$; or R^{D11} and R^{D12} together represent $=O$;

36

preferably wherein

m^D and n^D independently of one another are 0, 1, 2 or 3; preferably with the proviso that $m^D+n^D \leq 3$.

Y^{D1} is selected from $-O-$, $-NR^{D8}-$ and $-CR^{D9}R^{D10}-$ and Y^{D2} represents $-CR^{D11}R^{D12}-$; or Y^{D1} represents $-CR^{D9}R^{D10}-$ and Y^{D2} is selected from $-O-$ and $-NR^{D8}-$; R^{D1} , R^{D2} , R^{D3} , R^{D4} , R^{D5} , R^{D6} , R^{D7} , R^{D8} , R^{D9} , R^{D10} , R^{D11} and R^{D12} independently of one another represent $-H$, $-F$, $-OH$, $-C_{1-3}$ -alkylene- OH , $-C(=O)NH_2$, $-CH_2NH_2$, $-CH_2N(CH_3)_2$, $-NHCH_2CF_3$, $-CH_3$, or $-CH_2CF_3$; or R^{D2} and R^{D3} together represent $=O$; or R^{D4} and R^{D5} together represent $=O$; or R^{D9} and R^{D10} together represent $=O$; or R^{D11} and R^{D12} together represent $=O$; preferably with the proviso that only 1, 2 or 3 of R^{D1} , R^{D2} , R^{D3} , R^{D4} , R^{D5} , R^{D6} , R^{D7} , R^{D8} , R^{D9} , R^{D10} , R^{D11} and R^{D12} represent a residue that is not $-H$; preferably with the proviso that at least one of R^{D1} , R^{D2} , R^{D3} , R^{D4} , R^{D5} , R^{D6} , R^{D7} , R^{D8} , R^{D9} , R^{D10} , R^{D11} and R^{D12} represent a residue that is not $-H$.

In a preferred embodiment of the benzofuran derivative according to the invention R^5 and $R^{5'}$ independently of one another represent


- $-H$;
- $-C_1-C_6$ -alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
- $-C_1-C_6$ -heteroalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
- 3-14-membered cycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered cycloalkyl is optionally connected through $-C_1-C_6$ -alkylene- or $-C_1-C_6$ -heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or poly substituted.

Preferably, R^5 and $R^{5'}$ independently of one another represent $-H$, $-C_1-C_6$ -alkyl, or $-C_1-C_6$ -alkylene- $N(C_1-C_6$ -alkyl) $_2$.

In a preferred embodiment of the benzofuran derivative according to the invention, at least one of R^5 and $R^{5'}$ is not $-H$.


In a preferred embodiment of the benzofuran derivative according to the invention, R^5 and $R^{5'}$ are both $-H$.

In preferred embodiments, T represents $-O-$ and U represents $-CR^5R^{5'}$ and the resultant moiety $-O-CR^5R^{5'}$ represents a residue selected from the group consisting of:



37

In preferred embodiments, T represents $-\text{CR}^5\text{R}^{5*}-$ and U represents $-\text{O}-$ and the resultant moiety $-\text{CR}^5\text{R}^{5*}-\text{O}-$ represents a residue:



In preferred embodiments, R^5 represents $-\text{H}$ and R^{5*} represents a residue selected from the group consisting of $-\text{H}$, $-\text{C}_{1-3}\text{-alkyl}$, $-\text{CF}_3$, $-\text{CF}_2\text{H}$, $-\text{CFH}_2$, $-\text{C}_{1-3}\text{-alkylene-CF}_3$, $-\text{C}_{1-3}\text{-alkylene-CF}_2\text{H}$, $-\text{C}_{1-3}\text{-alkylene-CFH}_2$, and $-\text{C}_{1-3}\text{-alkylene-OH}$; preferably $-\text{H}$ or $\text{C}_{1-3}\text{-alkyl}$.

In a preferred embodiment of the benzofuran derivative according to the invention R^6 , R^7 and R^8 independently of one another represent

- $-\text{H}$;
- $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{SF}_5$, $-\text{CN}$, $-\text{NO}_2$, $-\text{C}(=\text{O})\text{OH}$, $-\text{NH}_2$;
- $-\text{C}_{1-6}\text{-alkyl}$, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
- $-\text{O}-\text{C}_{1-6}\text{-alkyl}$, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
- $-\text{NHC}_{1-6}\text{-alkyl}$, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
- $-\text{N}(\text{C}_{1-6}\text{-alkyl})_2$, saturated or unsaturated, unsubstituted, mono- or poly substituted;
- $-\text{C}(=\text{O})\text{OC}_{1-6}\text{-alkyl}$, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
- $-\text{OC}(=\text{O})\text{C}_{1-6}\text{-alkyl}$, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
- $-\text{C}_{1-6}\text{-heteroalkyl}$, saturated or unsaturated, unsubstituted, mono- or poly substituted.

Preferably, R^6 , R^7 and R^8 independently of one another represent

- $-\text{H}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{SH}$, $-\text{SF}_5$, $-\text{CN}$, $-\text{NO}_2$, $-\text{C}(=\text{O})\text{OH}$, $-\text{NH}_2$,
- $-\text{C}_{1-6}\text{-alkyl}$, $-\text{CF}_3$, $-\text{CHF}_2$, $-\text{CH}_2\text{F}$,
- $-\text{O}-\text{C}_{1-6}\text{-alkyl}$, $-\text{OCF}_3$, $-\text{OCHF}_2$, $-\text{OCH}_2\text{F}$,
- $-\text{NHC}_{1-6}\text{-alkyl}$ unsubstituted or substituted with one or more substituents independently of one another selected from $-\text{OH}$, $=\text{O}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{SH}$, $=\text{S}$, $-\text{CN}$, $-\text{CF}_3$, $-\text{CHF}_2$, $-\text{CH}_2\text{F}$, $-\text{OCF}_3$, $-\text{OCHF}_2$, $-\text{OCH}_2\text{F}$, SF_5 , $-\text{NO}_2$, $-\text{C}(=\text{O})\text{OH}$, $-\text{NH}_2$, and $-\text{C}(=\text{O})\text{NH}_2$;
- $-\text{N}(\text{C}_{1-6}\text{-alkyl})_2$ unsubstituted or substituted with one or more substituents independently of one another selected from $-\text{OH}$, $=\text{O}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{SH}$, $=\text{S}$, $-\text{CN}$, $-\text{CF}_3$, $-\text{CHF}_2$, $-\text{CH}_2\text{F}$, $-\text{OCF}_3$, $-\text{OCHF}_2$, $-\text{OCH}_2\text{F}$, SF_5 , $-\text{NO}_2$, $-\text{C}(=\text{O})\text{OH}$, $-\text{NH}_2$, and $-\text{C}(=\text{O})\text{NH}_2$;
- $-\text{C}(=\text{O})\text{OC}_{1-6}\text{-alkyl}$ unsubstituted or substituted with one or more substituents independently of one another selected from $-\text{OH}$, $=\text{O}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{SH}$, $=\text{S}$, $-\text{CN}$, $-\text{CF}_3$, $-\text{CHF}_2$, $-\text{CH}_2\text{F}$, $-\text{OCF}_3$, $-\text{OCHF}_2$, $-\text{OCH}_2\text{F}$, SF_5 , $-\text{NO}_2$, $-\text{C}(=\text{O})\text{OH}$, $-\text{NH}_2$, and $-\text{C}(=\text{O})\text{NH}_2$;
- $-\text{OC}(=\text{O})\text{C}_{1-6}\text{-alkyl}$ unsubstituted or substituted with one or more substituents independently of one another selected from $-\text{OH}$, $=\text{O}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{SH}$, $=\text{S}$, $-\text{CN}$, $-\text{CF}_3$, $-\text{CHF}_2$, $-\text{CH}_2\text{F}$, $-\text{OCF}_3$, $-\text{OCHF}_2$, $-\text{OCH}_2\text{F}$, SF_5 , $-\text{NO}_2$, $-\text{C}(=\text{O})\text{OH}$, $-\text{NH}_2$, and $-\text{C}(=\text{O})\text{NH}_2$; or

38

$-\text{C}_{1-6}\text{-heteroalkyl}$ unsubstituted or substituted with one or more substituents independently of one another selected from $-\text{OH}$, $=\text{O}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{SH}$, $=\text{S}$, $-\text{CN}$, $-\text{CF}_3$, $-\text{CHF}_2$, $-\text{CH}_2\text{F}$, $-\text{OCF}_3$, $-\text{OCHF}_2$, $-\text{OCH}_2\text{F}$, SF_5 , $-\text{NO}_2$, $-\text{C}(=\text{O})\text{OH}$, $-\text{NH}_2$, and $-\text{C}(=\text{O})\text{NH}_2$.

In preferred embodiments, R^6 , R^7 and R^8 independently of one another represents a residue selected from the group consisting of $-\text{H}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{CN}$, $\text{C}_{1-3}\text{-alkyl}$, $-\text{CF}_3$, $-\text{CF}_2\text{H}$, and $-\text{CFH}_2$; preferably $-\text{H}$ or $-\text{F}$.

In a preferred embodiment of the benzofuran derivative according to the invention R^6 represents $-\text{H}$, $-\text{F}$, $-\text{Cl}$, $-\text{CN}$, or $-\text{C}_1\text{-C}_6\text{-alkyl}$.

In a preferred embodiment of the benzofuran derivative according to the invention, R^6 does not represent $-\text{H}$.


In preferred embodiments, R^6 represents a residue selected from the group consisting of $-\text{H}$, $-\text{F}$, $-\text{Cl}$, $-\text{CN}$ or $-\text{CH}_3$; preferably $-\text{H}$, $-\text{F}$, $-\text{CN}$ or $-\text{CH}_3$.

In a preferred embodiment of the benzofuran derivative according to the invention R^7 represents $-\text{H}$, $-\text{F}$, $-\text{Cl}$, $-\text{CN}$, or $-\text{C}_1\text{-C}_6\text{-alkyl}$.

In a preferred embodiment of the benzofuran derivative according to the invention R^7 does not represent $-\text{H}$.

In preferred embodiments, especially when Q represents $-\text{NR}^3\text{R}^4$, R^7 represents a residue selected from the group consisting of $-\text{H}$, $-\text{F}$, $-\text{Cl}$, $-\text{CN}$ or CH_3 ; preferably $-\text{H}$, $-\text{F}$, $-\text{C}_1$ or $-\text{CH}_3$.

In preferred embodiments, especially when Q represents $-\text{OR}^2$, R^7 represents a residue selected from the group consisting of $-\text{H}$ or



In a preferred embodiment of the benzofuran derivative according to the invention R^8 represents $-\text{H}$, $-\text{F}$, $-\text{Cl}$, $-\text{CN}$, or $-\text{C}_1\text{-C}_6\text{-alkyl}$.

In a preferred embodiment of the benzofuran derivative according to the invention R^8 does not represent $-\text{H}$.

In preferred embodiments, R^8 represents a residue selected from the group consisting of $-\text{H}$, $-\text{F}$, $-\text{Cl}$, $-\text{CN}$ or CH_3 ; preferably $-\text{F}$.

In a preferred embodiment of the benzofuran derivative according to the invention

- (i) R^6 , R^7 and R^8 each represent $-\text{H}$; or
- (ii) two of R^6 , R^7 and R^8 represent $-\text{H}$ and the other of R^6 , R^7 and R^8 represents $-\text{F}$, $-\text{Cl}$, $-\text{CN}$, or $-\text{CH}_3$; or
- (iii) one of R^6 , R^7 and R^8 represents $-\text{H}$ and the other of R^6 , R^7 and R^8 independently of one another represent $-\text{F}$, $-\text{Cl}$, $-\text{CN}$, or $-\text{CH}_3$.

39

In a particularly preferred embodiment, the compound is according to general formula (I), wherein

R¹ represents —CH₃; and/or

R⁶, R⁷ and R⁵ each represent —H; and/or

T represents —O—; and/or


U represents —CH₂—; and/or

V represents thiazolyl, pyridyl, or pyrazolyl; wherein said thiazolyl, pyridyl, and pyrazolyl each independently from one another can be unsubstituted, monosubstituted or disubstituted with a substituent selected from the group consisting of —CH₃; —F; —CH₂CHF₂; and —CF₃; and/or

Q represents NR³R⁴; and/or


R³ represents H; and/or

R⁴ represents



40

-continued



In preferred embodiments of the invention, the benzofuran derivative is selected from the group consisting of:

cpd 001	5-(furan-3-ylmethoxy)-2-methylbenzofuran-3-carboxylic acid
cpd 002	2-methyl-5-(oxazol-5-ylmethoxy)benzofuran-3-carboxylic acid
cpd 003	2-methyl-5-(pyrrolidin-3-ylmethoxy)benzofuran-3-carboxylic acid
cpd 004	2-methyl-5-(pyridin-4-ylmethoxy)benzofuran-3-carboxylic acid
cpd 005	2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxylic acid
cpd 006	2-methyl-5-(pyridin-3-ylmethoxy)benzofuran-3-carboxylic acid
cpd 007	2-methyl-5-(pyridazin-3-ylmethoxy)benzofuran-3-carboxylic acid
cpd 008	2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxylic acid
cpd 009	2-methyl-5-((1-methyl-1H-pyrazol-3-yl)methoxy)benzofuran-3-carboxylic acid
cpd 010	2-methyl-5-((1-methyl-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxylic acid
cpd 011	2-methyl-5-(thiophen-3-ylmethoxy)benzofuran-3-carboxylic acid
cpd 012	2-methyl-5-(thiophen-2-ylmethoxy)benzofuran-3-carboxylic acid
cpd 013	2-methyl-5-(thiazol-2-ylmethoxy)benzofuran-3-carboxylic acid
cpd 014	2-methyl-5-(thiazol-5-ylmethoxy)benzofuran-3-carboxylic acid
cpd 015	2-methyl-5-((tetrahydro-2H-pyran-4-yl)methoxy)benzofuran-3-carboxylic acid
cpd 016	5-((3,5-dimethylisoxazol-4-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
cpd 017	5-(benzo[d]thiazol-2-ylmethoxy)-2-methylbenzofuran-3-carboxylic acid
cpd 018	2-methyl-5-((5-(trifluoromethyl)furan-2-yl)methoxy)benzofuran-3-carboxylic acid
cpd 019	5-((1-(tert-butoxycarbonyl)pyrrolidin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
cpd 020	(S)-2-methyl-5-(oxazol-5-ylmethoxy)-N-(pyrrolidin-3-yl)benzofuran-3-carboxamide
cpd 021	(S)-2-methyl-5-(pyridin-4-ylmethoxy)-N-(pyrrolidin-3-yl)benzofuran-3-carboxamide
cpd 022	(S)-2-methyl-5-(pyridin-3-ylmethoxy)-N-(pyrrolidin-3-yl)benzofuran-3-carboxamide
cpd 023	(S)-2-methyl-5-((1-methyl-1H-pyrazol-3-yl)methoxy)-N-(pyrrolidin-3-yl)benzofuran-3-carboxamide
cpd 024	(S)-2-methyl-N-(pyrrolidin-3-yl)-5-(thiazol-5-ylmethoxy)benzofuran-3-carboxamide
cpd 025	(S)-2-methyl-N-(pyrrolidin-3-yl)-5-(thiazol-2-ylmethoxy)benzofuran-3-carboxamide
cpd 026	2-methyl-N-(2-oxopyrrolidin-3-yl)-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
cpd 027	2-methyl-N-(2-oxopyrrolidin-3-yl)-5-(pyridin-3-ylmethoxy)benzofuran-3-carboxamide
cpd 028	(S)-2-methyl-5-((5-methylpyridin-3-yl)methoxy)-N-(pyrrolidin-3-yl)benzofuran-3-carboxamide
cpd 029	(S)-2-methyl-5-((6-methylpyridin-2-yl)methoxy)-N-(pyrrolidin-3-yl)benzofuran-3-carboxamide
cpd 030	2-methyl-N-(2-oxopyrrolidin-3-yl)-5-(pyridazin-3-ylmethoxy)benzofuran-3-carboxamide
cpd 031	2-methyl-N-(2-oxopyrrolidin-3-yl)-5-(pyrazin-2-ylmethoxy)benzofuran-3-carboxamide
cpd 032	2-methyl-N-(2-oxopyrrolidin-3-yl)-5-(pyrimidin-5-ylmethoxy)benzofuran-3-carboxamide
cpd 033	2-methyl-5-((6-methylpyridin-3-yl)methoxy)-N-(tetrahydrofuran-3-yl)benzofuran-3-carboxamide
cpd 034	2-methyl-5-((4-methylpyridin-3-yl)methoxy)-N-(tetrahydrofuran-3-yl)benzofuran-3-carboxamide
cpd 035	2-methyl-N-(2-oxopyrrolidin-3-yl)-5-(thiophen-3-ylmethoxy)benzofuran-3-carboxamide
cpd 036	2-methyl-N-(2-oxopyrrolidin-3-yl)-5-(thiazol-2-ylmethoxy)benzofuran-3-carboxamide
cpd 037	2-methyl-5-((4-methylpyridin-3-yl)methoxy)-N-(2-oxopyrrolidin-3-yl)benzofuran-3-carboxamide
cpd 038	2-methyl-5-((5-methylpyridin-3-yl)methoxy)-N-(2-oxopyrrolidin-3-yl)benzofuran-3-carboxamide
cpd 039	2-methyl-N-(1-methylpiperidin-4-yl)-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
cpd 040	5-((4-methoxypyridin-3-yl)methoxy)-2-methyl-N-(tetrahydrofuran-3-yl)benzofuran-3-carboxamide

-continued

cpd 041	2-methyl-N-(1-methylpiperidin-4-yl)-5-(thiophen-2-ylmethoxy)benzofuran-3-carboxamide
cpd 042	2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(2-oxopyrrolidin-3-yl)benzofuran-3-carboxamide
cpd 043	2-methyl-5-((4-methylthiazol-2-yl)methoxy)-N-(2-oxopyrrolidin-3-yl)benzofuran-3-carboxamide
cpd 044	5-((2,6-dimethylpyridin-3-yl)methoxy)-2-methyl-N-(2-oxopyrrolidin-3-yl)benzofuran-3-carboxamide
cpd 045	2-methyl-N-(1-methylpiperidin-4-yl)-5-(1-(pyridin-4-yl)ethoxy)benzofuran-3-carboxamide
cpd 046	5-((2-methoxypyridin-3-yl)methoxy)-2-methyl-N-(2-oxopyrrolidin-3-yl)benzofuran-3-carboxamide
cpd 047	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
cpd 048	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
cpd 049	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-(thiazol-2-ylmethoxy)benzofuran-3-carboxamide
cpd 050	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-(thiazol-2-ylmethoxy)benzofuran-3-carboxamide
cpd 051	5-((2-methoxypyridin-3-yl)methoxy)-2-methyl-N-(1-methylpiperidin-4-yl)benzofuran-3-carboxamide
cpd 052	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylpyridin-3-yl)methoxy)-benzofuran-3-carboxamide
cpd 053	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((3-methylpyridin-2-yl)methoxy)benzofuran-3-carboxamide
cpd 054	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)-benzofuran-3-carboxamide
cpd 055	(S)-2-methyl-N-(pyrrolidin-3-yl)-5-((5-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 056	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 057	2-methyl-N-(tetrahydrofuran-3-yl)-5-((4-(trifluoromethyl)thiazol-2-yl)methoxy)benzofuran-3-carboxamide
cpd 058	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methyl-benzofuran-3-carboxamide
cpd 059	5-(isoquinolin-1-ylmethoxy)-2-methyl-N-(1-methylpiperidin-4-yl)benzofuran-3-carboxamide
cpd 060	5-(isoquinolin-5-ylmethoxy)-2-methyl-N-(1-methylpiperidin-4-yl)benzofuran-3-carboxamide
cpd 061	5-(isoquinolin-6-ylmethoxy)-2-methyl-N-(1-methylpiperidin-4-yl)benzofuran-3-carboxamide
cpd 062	N-(3,3-difluoropiperidin-4-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
cpd 063	N-(3,3-difluoropiperidin-4-yl)-5-((6-methoxypyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide
cpd 064	2-methyl-N-(2-oxopyrrolidin-3-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 065	2-methyl-N-(2-oxopyrrolidin-3-yl)-5-((6-(trifluoromethyl)pyridin-2-yl)methoxy)benzofuran-3-carboxamide
cpd 066	2-methyl-N-(2-oxopyrrolidin-3-yl)-5-((6-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 067	5-((2-chloropyridin-3-yl)methoxy)-N-(3,3-difluoropiperidin-4-yl)-2-methylbenzofuran-3-carboxamide
cpd 068	5-((4-chloropyridin-3-yl)methoxy)-N-(3,3-difluoropiperidin-4-yl)-2-methylbenzofuran-3-carboxamide
cpd 069	2-methyl-N-(1-methylpiperidin-4-yl)-5-((5-(trifluoromethyl)furan-2-yl)methoxy)benzofuran-3-carboxamide
cpd 070	N-(3,3-difluoropiperidin-4-yl)-5-(imidazo[1,2-a]pyridin-6-ylmethoxy)-2-methylbenzofuran-3-carboxamide
cpd 071	N-(3,3-difluoropiperidin-4-yl)-5-((6-(dimethylamino)pyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide
cpd 072	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)-methoxy)benzofuran-3-carboxamide
cpd 073	tert-butyl (S)-3-(2-methyl-5-((5-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate
cpd 074	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 075	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((5-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 076	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((3-(trifluoromethyl)pyridin-2-yl)methoxy)benzofuran-3-carboxamide
cpd 077	tert-butyl (S)-3-(2-methyl-5-((5-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate
cpd 078	tert-butyl 4-(5-((6-(dimethylamino)pyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamido)-3,3-difluoropiperidine-1-carboxylate
cpd 079	tert-butyl 3-(((3-(ethoxycarbonyl)-2-methylbenzofuran-5-yl)oxy)methyl)pyrrolidine-1-carboxylate
cpd 080	tert-butyl (R)-3-(2-methyl-5-(oxazol-5-ylmethoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate
cpd 081	tert-butyl (R)-3-(2-methyl-5-(pyridin-4-ylmethoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate
cpd 082	tert-butyl (R)-3-(2-methyl-5-(pyridin-3-ylmethoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate
cpd 083	tert-butyl (R)-3-(2-methyl-5-((1-methyl-1H-pyrazol-3-yl)methoxy)benzofuran-3-carboxamido)-pyrrolidine-1-carboxylate
cpd 084	tert-butyl (R)-3-(2-methyl-5-(thiazol-5-ylmethoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate
cpd 085	tert-butyl (R)-3-(2-methyl-5-(thiazol-2-ylmethoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate
cpd 086	tert-butyl (R)-3-(2-methyl-5-((6-methylpyridin-2-yl)methoxy)benzofuran-3-carboxamido)-pyrrolidine-1-carboxylate
cpd 087	tert-butyl 3,3-difluoro-4-(2-methyl-5-(thiazol-2-ylmethoxy)benzofuran-3-carboxamido)piperidine-1-carboxylate
cpd 088	tert-butyl 3,3-difluoro-4-(2-methyl-5-((3-methylpyridin-2-yl)methoxy)benzofuran-3-carboxamido)piperidine-1-carboxylate

-continued

cpd 089	tert-butyl 3,3-difluoro-4-(5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamido)piperidine-1-carboxylate
cpd 090	tert-butyl 3,3-difluoro-4-(2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)piperidine-1-carboxylate
cpd 091	tert-butyl 3,3-difluoro-4-(5-((6-methoxypyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamido)piperidine-1-carboxylate
cpd 092	tert-butyl 4-(5-((2-chloropyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamido)-3,3-difluoropiperidine-1-carboxylate
cpd 093	tert-butyl 4-(5-((4-chloropyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamido)-3,3-difluoropiperidine-1-carboxylate
cpd 094	tert-butyl 3,3-difluoro-4-(5-(imidazo[1,2-a]pyridin-6-yl)methoxy)-2-methylbenzofuran-3-carboxamido)piperidine-1-carboxylate
cpd 095	tert-butyl 3,3-difluoro-4-(2-methyl-5-((5-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)piperidine-1-carboxylate
cpd 096	tert-butyl 3,3-difluoro-4-(2-methyl-5-((3-(trifluoromethyl)pyridin-2-yl)methoxy)benzofuran-3-carboxamido)piperidine-1-carboxylate
cpd 097	tert-butyl 3,3-difluoro-4-(2-methyl-5-(pyridin-2-yl)methoxy)benzofuran-3-carboxamido)piperidine-1-carboxylate
cpd 098	tert-butyl 3,3-difluoro-4-(2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)piperidine-1-carboxylate
cpd 099	2-methyl-5-((4-methylpyridin-3-yl)methoxy)benzofuran-3-carboxylic acid
cpd 100	5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
cpd 101	2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid
cpd 102	2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid
cpd 103	N-(4,4-difluoropiperidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 104	N-(1-hydroxy-2-methylpropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 105	N-(4,4-difluoropyrrolidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 106	N-(4,4-difluoropyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 107	N-(1-(hydroxymethyl)cyclobutyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 108	N-(1-(dimethylamino)methyl)cyclobutyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 109	N-(1-hydroxy-2-methylpropan-2-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 110	N-(3,3-difluoro-1-methylpiperidin-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 111	N-(4,4-difluoropiperidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 112	N-(3,3-difluoropiperidin-4-yl)-4-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 113	N-(3,3-difluoropiperidin-4-yl)-6-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 114	N-(1-(2-hydroxyethyl)cyclobutyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 115	N-(5,5-difluoropiperidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 116	2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(1-(2,2,2-trifluoroethyl)pyrrolidin-3-yl)benzofuran-3-carboxamide
cpd 117	N-(1-(hydroxymethyl)cyclobutyl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 118	N-(4,4-difluoro-1-methylpiperidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 119	N-(3-carbamoyloxetan-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 120	N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 121	N-(3-(2-hydroxyethyl)tetrahydrofuran-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 122	7-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 123	N-(4,4-difluoro-1-methylpyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 124	N-(4,4-difluoro-1-methylpyrrolidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 125	N-(5,5-difluoropiperidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 126	N-(3-(2-hydroxyethyl)tetrahydrofuran-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 127	N-(1-(2-hydroxyethyl)cyclobutyl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 128	N-(3-(2-hydroxyethyl)oxetan-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 129	N-(3-(2-hydroxyethyl)oxetan-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 130	N-(3,3-difluoro-1-methylpiperidin-4-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide

-continued

cpd 131	N-(4,4-difluoro-1-methylpiperidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
cpd 132	N-(3-(hydroxymethyl)tetrahydrofuran-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)-benzofuran-3-carboxamide
cpd 133	N-(3-(hydroxymethyl)oxetan-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
cpd 134	N-(4,4-difluoro-1-(2-hydroxyethyl)pyrrolidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 135	N-(3-(hydroxymethyl)tetrahydrofuran-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)-methoxy)benzofuran-3-carboxamide
cpd 136	N-(1-(aminomethyl)cyclobutyl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
cpd 137	N-(1-(aminomethyl)cyclobutyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 138	2-methyl-N-(1H-pyrazol-3-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 139	N-(1-(hydroxymethyl)cyclopropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 140	N-(1-(dimethylamino)methyl)cyclobutyl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
cpd 141	N-(5,5-difluoro-1-methylpiperidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
cpd 142	N-(3-(hydroxymethyl)oxetan-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 143	2-methyl-N-(1-methyl-1H-pyrazol-4-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 144	2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(1H-pyrazol-3-yl)benzofuran-3-carboxamide
cpd 145	2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(1-((2,2,2-trifluoroethyl)amino)methyl)cyclobutyl)benzofuran-3-carboxamide
cpd 146	2-methyl-N-(1-methyl-1H-pyrazol-3-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 147	N-(4-hydroxytetrahydrofuran-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
cpd 148	2-methyl-N-(tetrahydrofuran-3-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 149	2-methyl-N-(1-(2,2,2-trifluoroethyl)pyrrolidin-3-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)-methoxy)benzofuran-3-carboxamide
cpd 150	2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(1H-pyrazol-4-yl)benzofuran-3-carboxamide
cpd 151	N-(4,4-difluoro-1-(2-hydroxyethyl)pyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)-benzofuran-3-carboxamide
cpd 152	2-methyl-N-(1-methyl-1H-pyrazol-4-yl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 153	2-methyl-N-(1-((2,2,2-trifluoroethyl)amino)methyl)cyclobutyl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 154	2-methyl-N-(1-methyl-1H-pyrazol-5-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 155	N-(4,4-difluoro-1-(2,2,2-trifluoroethyl)pyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)-methoxy)benzofuran-3-carboxamide
cpd 156	2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(2-oxaspiro[3.3]heptan-6-yl)benzofuran-3-carboxamide
cpd 157	N-(4,4-difluoro-1-(2,2,2-trifluoroethyl)pyrrolidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 158	2-methyl-N-(1H-pyrazol-4-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 159	2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(tetrahydrofuran-3-yl)benzofuran-3-carboxamide
cpd 160	N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 161	2-methyl-N-(1-methyl-1H-pyrazol-3-yl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 162	N-(4-hydroxytetrahydrofuran-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 163	N-(5,5-difluoro-1-methylpiperidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 164	N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((2-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 165	6-chloro-N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
cpd 166	2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)benzofuran-3-carboxamide
cpd 167	2-methyl-N-(2-oxaspiro[3.3]heptan-6-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 168	2-methyl-N-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
cpd 169	N-(1-(2-methoxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 170	(3-aminotetrahydrofuran-3-yl)methyl 2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxylate
cpd 171	2-methyl-N-(1-methyl-1H-pyrazol-5-yl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 172	2,2,2-trifluoroethyl 3,3-difluoro-4-(2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate
cpd 173	2-methyl-N-(2-azaspiro[3.3]heptan-6-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 174	6-chloro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)-methoxy)benzofuran-3-carboxamide

-continued

cpd 175	2,2,2-trifluoroethyl (R)-3,3-difluoro-4-(2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamido)pyrrolidine-1-carboxylate
cpd 176	N-(3,3-difluoropiperidin-4-yl)-7-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
cpd 177	4-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)-methoxy)benzofuran-3-carboxamide
cpd 178	(3-amino-1H-pyrazol-1-yl)(2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-yl)methanone
cpd 179	N-(3,3-difluoropiperidin-4-yl)-2,6-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
cpd 180	N-(1-(2-hydroxyethyl)-5-oxopyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 181	(4-amino-1H-pyrazol-1-yl)(2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-yl)methanone
cpd 182	6-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)-methoxy)benzofuran-3-carboxamide
cpd 183	N-(3,3-difluoropiperidin-4-yl)-2,7-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
cpd 184	7-chloro-N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
cpd 185	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2,6-dimethyl-5-((4-methylthiazol-5-yl)methoxy)-benzofuran-3-carboxamide
cpd 186	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2,7-dimethyl-5-((4-methylthiazol-5-yl)methoxy)-benzofuran-3-carboxamide
cpd 187	7-chloro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)-methoxy)benzofuran-3-carboxamide
cpd 188	7-cyano-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)-methoxy)benzofuran-3-carboxamide
cpd 189	N-((3-aminooxetan-3-yl)methyl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
cpd 190	2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(2-azaspiro[3.3]heptan-6-yl)benzofuran-3-carboxamide
cpd 191	4-cyano-N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
cpd 192	(3-amino-1H-pyrazol-1-yl)(2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-yl)-methanone
cpd 193	4-cyano-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)-methoxy)benzofuran-3-carboxamide
cpd 194	6-cyano-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)-methoxy)benzofuran-3-carboxamide
cpd 195	6-cyano-N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
cpd 196	7-cyano-N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
cpd 197	2-(difluoromethyl)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((4-methylthiazol-5-yl)-methoxy)benzofuran-3-carboxamide
cpd 198	2-(difluoromethyl)-N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-5-((4-methylthiazol-5-yl)-methoxy)benzofuran-3-carboxamide
cpd 199	2-(difluoromethyl)-N-(3,3-difluoropiperidin-4-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
cpd 200	N-(1-(hydroxymethyl)cyclopropyl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
cpd 201	N-(3-(hydroxymethyl)azetidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 202	N-(3-(hydroxymethyl)azetidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
cpd 203	N-(4,4-difluorotetrahydrofuran-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
cpd 204	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)-methoxy)benzofuran-3-carboxamide
cpd 205	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide
cpd 206	5-((2-ethoxypyridin-3-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-benzofuran-3-carboxamide
cpd 207	N-(3,3-difluoropiperidin-4-yl)-5-((2-ethoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 208	2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 209	2-methyl-5-((2-(trifluoromethyl)thiazol-5-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 210	2-methyl-5-((2-methyl-4-(trifluoromethyl)thiazol-5-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 211	2-methyl-5-((2-methylloxazol-5-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 212	2-methyl-5-((2-methylthiazol-4-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 213	2-methyl-5-((3-(trifluoromethyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 214	2-methyl-5-((4-methylpyridin-3-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 215	2-methyl-5-((5-methylisoxazol-3-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 216	2-methyl-5-((5-methylthiazol-4-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 217	2-methyl-5-((6-methylpyridin-3-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 218	2-methyl-5-(oxetan-3-ylmethoxy)benzofuran-3-carboxylic acid
Cpd 219	2-methyl-5-(pyrimidin-4-ylmethoxy)benzofuran-3-carboxylic acid
Cpd 220	5-((1-(2,2-difluoroethyl)-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 221	5-((2-(difluoromethoxy)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid

-continued

Cpd 222	5-((2-(dimethylamino)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 223	5-((2,4-dimethylthiazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 224	5-((2,5-dimethylthiazol-4-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 225	5-((2-aminopyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 226	5-((3-(difluoromethyl)pyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 227	5-((6-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 228	5-((4-fluoro-1-methyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 229	2-(difluoromethyl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 230	2-(difluoromethyl)-5-((2-methoxypyridin-3-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 231	2-cyclopropyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 232	2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 233	4-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 234	4-fluoro-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 235	5-((1-(2,2-difluoroethyl)-1H-imidazol-2-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 236	5-((2-(azetidin-1-yl)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 237	5-((2-(difluoromethyl)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 238	5-((2-cyanopyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 239	5-((2-ethoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 240	5-((3-fluoropyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 241	5-((5-(2-hydroxyethoxy)pyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 242	5-((5-fluoropyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 243	6-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 244	6-fluoro-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 245	7-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 246	7-fluoro-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 247	2-methyl-5-((2-methylpyridin-3-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 248	2-methyl-5-(1-(2-(trifluoromethyl)thiazol-5-yl)ethoxy)benzofuran-3-carboxylic acid
Cpd 249	2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 250	2-methyl-5-((1-methyl-2-oxo-1,2-dihydropyridin-3-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 251	2,4-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 252	2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)oxy)methylbenzofuran-3-carboxylic acid
Cpd 253	2-methyl-5-((1-methyl-1H-1,2,3-triazol-4-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 254	2-methyl-5-((1-methyl-1H-1,2,3-triazol-5-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 255	2-methyl-5-((1-methyl-1H-1,2,4-triazol-3-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 256	2-methyl-5-((1-methyl-6-oxo-1,6-dihydropyridin-2-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 257	2-methyl-5-((2-methylpyrimidin-4-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 258	2-methyl-5-((3-methylloxetan-3-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 259	2-methyl-5-((4-methylloxazol-2-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 260	2-methyl-5-((4-methylthiazol-2-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 261	2-methyl-5-((5-methylpyrazin-2-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 262	2-methyl-5-((5-methylthiazol-2-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 263	2-methyl-5-((6-methylpyridazin-3-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 264	2-methyl-5-(pyrimidin-2-yl)methoxybenzofuran-3-carboxylic acid
Cpd 265	2-cyclopropyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 266	5-((6-methoxypyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 267	5-(imidazo[1,2-a]pyridin-6-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 268	2-methyl-5-(1-(4-methylthiazol-5-yl)ethoxy)benzofuran-3-carboxylic acid
Cpd 269	5-((2-methoxypyridin-4-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 270	2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylserine
Cpd 271	(R)-N-(1-amino-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 272	(R)-N-(1-amino-3-(methylsulfonyl)-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 273	(R)-N-(1-amino-3-methyl-1-oxobutan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 274	(R)-N-(1-hydroxy-3-(methylsulfonyl)propan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 275	(R)-N-(1-hydroxy-3-(methylsulfonyl)propan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 276	(R)-N-(3-hydroxy-1-(methylamino)-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 277	(S)-N-(1-(3,3-difluoropyrrolidin-1-yl)-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 278	(S)-N-(1-(dimethylamino)-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 279	(S)-N-(1-amino-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 280	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 281	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((2-(trifluoromethyl)thiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 282	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((2-methyl-4-(trifluoromethyl)thiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 283	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((2-methylloxazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 284	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((2-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 285	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-4-yl)methoxy)benzofuran-3-carboxamide

-continued

Cpd 286	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((3-(trifluoromethyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxamide
Cpd 287	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((4-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 288	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 289	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((5-methylisoxazol-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 290	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((5-methylthiazol-4-yl)methoxy)benzofuran-3-carboxamide
Cpd 291	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((6-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 292	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-(oxetan-3-ylmethoxy)benzofuran-3-carboxamide
Cpd 293	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-(pyridazin-3-ylmethoxy)benzofuran-3-carboxamide
Cpd 294	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-(pyrimidin-4-ylmethoxy)benzofuran-3-carboxamide
Cpd 295	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((1-(2,2-difluoroethyl)-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 296	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((2-(difluoromethoxy)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 297	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((2-(dimethylamino)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 298	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((2,4-dimethylthiazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 299	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((2,5-dimethylthiazol-4-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 300	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((2-aminopyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 301	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((3-(difluoromethyl)pyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 302	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((4-fluoro-1-methyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 303	(S)-N-(1-amino-3-methoxy-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 304	(S)-N-(1-amino-3-methyl-1-oxobutan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 305	(S)-N-(3-hydroxy-1-(methylamino)-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 306	(S)-N-(3-hydroxy-1-(methylamino)-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 307	(S)-N-(3-hydroxy-1-oxo-1-(pyrrolidin-1-yl)propan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 308	2-(difluoromethyl)-N-(1-(hydroxymethyl)cyclopropyl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 309	2-(difluoromethyl)-N-(3-(hydroxymethyl)oxetan-3-yl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 310	2-(difluoromethyl)-N-(3,3-difluoropiperidin-4-yl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 311	2-(difluoromethyl)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-hydroxypyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 312	2-(difluoromethyl)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-methoxypyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 313	2-cyclopropyl-N-(3,3-difluoropiperidin-4-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 314	2-cyclopropyl-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 315	2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(1-(oxazol-2-yl)ethyl)benzofuran-3-carboxamide
Cpd 316	2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(1,1,1-trifluoro-3-hydroxypropan-2-yl)benzofuran-3-carboxamide
Cpd 317	2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(5-oxo-1,4-oxazepan-6-yl)benzofuran-3-carboxamide
Cpd 318	2-methyl-N-(1-(methylcarbamoyl)cyclobutyl)-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 319	2-methyl-N-(1-(methylcarbamoyl)cyclobutyl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 320	2-methyl-N-(1-(methylcarbamoyl)cyclobutyl)-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 321	2-methyl-N-(2-(methylamino)-2-oxoethyl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 322	2-methyl-N-(2-(methylsulfonylamido)ethyl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 323	2-methyl-N-(2-(methylsulfonyl)ethyl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 324	2-methyl-N-(3-(methylcarbamoyl)oxetan-3-yl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 325	3-(((3-((4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)carbamoyl)-2-methylbenzofuran-5-yl)oxy)methyl)picolinamide

-continued

Cpd 326	4-fluoro-N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 327	4-fluoro-N-(1-(hydroxymethyl)cyclopropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 328	4-fluoro-N-(3-(hydroxymethyl)oxetan-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 329	4-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 330	4-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 331	5-((1-(2,2-difluoroethyl)-1H-imidazol-2-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
Cpd 332	5-((1-(2,2-difluoroethyl)-1H-pyrazol-5-yl)methoxy)-N-(3-(hydroxymethyl)-2-oxopiperidin-3-yl)-2-methylbenzofuran-3-carboxamide
Cpd 333	5-((1-(2,2-difluoroethyl)-1H-pyrazol-5-yl)methoxy)-N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methylbenzofuran-3-carboxamide
Cpd 334	5-((1-(2,2-difluoroethyl)-1H-pyrazol-5-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
Cpd 335	5-(2-(azetidin-1-yl)pyridin-3-yl)methoxy)-N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methylbenzofuran-3-carboxamide
Cpd 336	5-(2-(azetidin-1-yl)pyridin-3-yl)methoxy)-N-(3,3-difluoropiperidin-4-yl)-2-methylbenzofuran-3-carboxamide
Cpd 337	5-(2-(azetidin-1-yl)pyridin-3-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
Cpd 338	5-(2-(difluoromethoxy)pyridin-3-yl)methoxy)-N-(3,3-difluoropiperidin-4-yl)-2-methylbenzofuran-3-carboxamide
Cpd 339	5-(2-(difluoromethoxy)pyridin-3-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
Cpd 340	5-(2-(difluoromethyl)pyridin-3-yl)methoxy)-N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methylbenzofuran-3-carboxamide
Cpd 341	5-(2-(difluoromethyl)pyridin-3-yl)methoxy)-N-(3-(hydroxymethyl)oxetan-3-yl)-2-methylbenzofuran-3-carboxamide
Cpd 342	5-(2-(difluoromethyl)pyridin-3-yl)methoxy)-N-(3,3-difluoropiperidin-4-yl)-2-methylbenzofuran-3-carboxamide
Cpd 343	5-(2-(difluoromethyl)pyridin-3-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
Cpd 344	5-(2-(dimethylamino)pyridin-3-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
Cpd 345	5-(2,4-dimethylthiazol-5-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
Cpd 346	5-(2-aminopyridin-3-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
Cpd 347	5-(2-cyanopyridin-3-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
Cpd 348	5-(2-ethoxypyridin-3-yl)methoxy)-N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methylbenzofuran-3-carboxamide
Cpd 349	5-(2-hydroxypyridin-3-yl)methoxy)-N-(cis-4-hydroxytetrahydrofuran-3-yl)-2-methylbenzofuran-3-carboxamide
Cpd 350	5-(2-hydroxypyridin-3-yl)methoxy)-N-(trans-4-hydroxytetrahydrofuran-3-yl)-2-methylbenzofuran-3-carboxamide
Cpd 351	5-(3-(difluoromethyl)pyridin-2-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
Cpd 352	5-(3-fluoropyridin-2-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
Cpd 355	5-(4-fluoro-1-methyl-1H-pyrazol-5-yl)methoxy)-N-(3-(hydroxymethyl)-2-oxopiperidin-3-yl)-2-methylbenzofuran-3-carboxamide
Cpd 356	5-(4-fluoro-1-methyl-1H-pyrazol-5-yl)methoxy)-N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methylbenzofuran-3-carboxamide
Cpd 357	5-(4-fluoro-1-methyl-1H-pyrazol-5-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
Cpd 358	5-((5-(2-hydroxyethoxy)pyridin-2-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
Cpd 359	5-((5-fluoropyridin-2-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
Cpd 360	6-fluoro-N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 361	6-fluoro-N-(1-(hydroxymethyl)cyclopropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 362	6-fluoro-N-(3-(hydroxymethyl)oxetan-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 363	6-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 364	6-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 365	7-fluoro-N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 366	7-fluoro-N-(1-(hydroxymethyl)cyclopropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide

-continued

Cpd 367	7-fluoro-N-(3-(hydroxymethyl)oxetan-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 368	7-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 369	7-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 370	N-((1-cyanocyclopropyl)methyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 371	N-((1-cyanocyclopropyl)methyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 372	N-((1-fluorocyclopropyl)methyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 373	N-((1-fluorocyclopropyl)methyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 374	N-((1H-pyrazol-3-yl)methyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 375	N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-2-methyl-5-((2-methyloxazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 376	N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-2-methyl-5-((2-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 377	N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 378	N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 379	N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 380	N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-2-methyl-5-((6-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 381	N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-5-((1-(2,2-difluoroethyl)-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 382	N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-5-((2,4-dimethylthiazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 383	N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-5-((4-fluoro-1-methyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 384	N-((3-fluorooxetan-3-yl)methyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 385	N-((3-fluorooxetan-3-yl)methyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 386	N-(cis-4-hydroxytetrahydrofuran-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 387	N-(cis-4-hydroxytetrahydrofuran-3-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 388	N-(trans-4-hydroxytetrahydrofuran-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 389	N-(trans-4-hydroxytetrahydrofuran-3-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 390	N-((3S,4R)-4-hydroxy-2-oxopyrrolidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 391	N-((3S,4R)-4-hydroxy-2-oxopyrrolidin-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 392	N-((3S,4R)-4-hydroxy-2-oxopyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 393	N-((4-fluorotetrahydro-2H-pyran-4-yl)methyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 394	N-((4-fluorotetrahydro-2H-pyran-4-yl)methyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 395	N-((S)-1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-(1-(2-methylthiazol-5-yl)ethoxy)benzofuran-3-carboxamide
Cpd 396	N-(1-((2-hydroxyethyl)(methyl)amino)-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 397	N-(1-((2-hydroxyethyl)amino)-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 398	N-(1-((2-hydroxyethyl)amino)-2-methyl-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 399	N-(1-(difluoromethoxy)methyl)cyclopropyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 400	N-(1-(difluoromethoxy)methyl)cyclopropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 401	N-(1-(1H-benzo[d]imidazol-2-yl)-2-hydroxyethyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 402	N-(1-(2-(2-hydroxyethoxy)ethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 403	N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide
Cpd 404	N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 405	N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((1-methyl-2-oxo-1,2-dihydropyridin-3-yl)methoxy)benzofuran-3-carboxamide

-continued

Cpd 406	N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 407	N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-(thiazol-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 408	N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 409	N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 410	N-(1-(2-hydroxyethyl)-3-methyl-2-oxopyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 411	N-(1-(dimethylamino)-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 412	N-(1-(dimethylamino)-3-hydroxypropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 413	N-(1-(dimethylamino)-3-hydroxypropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 414	N-(1-(hydroxymethyl)cyclobutyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 415	N-(1-(hydroxymethyl)cyclobutyl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 416	N-(1-(hydroxymethyl)cyclobutyl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 417	N-(1-(hydroxymethyl)cyclobutyl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 418	N-(1-(hydroxymethyl)cyclopropyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 419	N-(1-(hydroxymethyl)cyclopropyl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 420	N-(1-(hydroxymethyl)cyclopropyl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 421	N-(1-(hydroxymethyl)cyclopropyl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 422	N-(1,3-dihydroxy-2-methylpropan-2-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide
Cpd 423	N-(1,3-dihydroxy-2-methylpropan-2-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 424	N-(1,3-dihydroxy-2-methylpropan-2-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 425	N-(1,3-dihydroxy-2-methylpropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 426	N-(1,3-dihydroxy-2-methylpropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 427	N-(1,3-dihydroxy-2-methylpropan-2-yl)-2-methyl-5-(pyridazin-3-ylmethoxy)benzofuran-3-carboxamide
Cpd 428	N-(1,3-dihydroxy-2-methylpropan-2-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 429	N-(1,3-dihydroxy-2-methylpropan-2-yl)-2-methyl-5-(pyrimidin-4-ylmethoxy)benzofuran-3-carboxamide
Cpd 430	N-(1,3-dihydroxy-2-methylpropan-2-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 431	N-(1,3-dihydroxy-2-methylpropan-2-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 432	N-(1-amino-1-oxo-3-(1H-1,2,4-triazol-1-yl)propan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 433	N-(1-amino-1-oxopropan-2-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 434	N-(1-amino-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 435	N-(1-amino-1-oxopropan-2-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 436	N-(1-amino-1-oxopropan-2-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 437	N-(1-amino-1-oxopropan-2-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 438	N-(1-amino-2-methyl-1-oxopropan-2-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 439	N-(1-amino-2-methyl-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 440	N-(1-amino-2-methyl-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 441	N-(1-amino-2-methyl-1-oxopropan-2-yl)-2-methyl-5-(pyridazin-3-ylmethoxy)benzofuran-3-carboxamide
Cpd 442	N-(1-amino-2-methyl-1-oxopropan-2-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 443	N-(1-amino-2-methyl-1-oxopropan-2-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 444	N-(1-amino-2-methyl-1-oxopropan-2-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 445	N-(1-amino-3-(cyclopropyl(methyl)amino)-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 446	N-(1-amino-3-(cyclopropyl(methyl)amino)-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide

-continued

Cpd 447	N-(1-amino-3-(difluoromethoxy)-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 448	N-(1-amino-3-(difluoromethoxy)-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 449	N-(1-amino-3-(dimethylamino)-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 450	N-(1-amino-3-(dimethylamino)-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 451	N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide
Cpd 452	N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 453	N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 454	N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 455	N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 456	N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((1-(2,2-difluoroethyl)-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 457	N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((2-(difluoromethyl)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 458	N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((2,4-dimethylthiazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 459	N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 460	N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((4-fluoro-1-methyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 461	N-(1-amino-3-hydroxy-2-methyl-1-oxopropan-2-yl)-2-methyl-5-((2-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 462	N-(1-amino-3-hydroxy-2-methyl-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 463	N-(1-amino-3-hydroxy-2-methyl-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 464	N-(1-amino-3-hydroxy-2-methyl-1-oxopropan-2-yl)-2-methyl-5-((6-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 465	N-(1-amino-3-hydroxy-2-methyl-1-oxopropan-2-yl)-5-((1-(2,2-difluoroethyl)-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 466	N-(1-amino-3-hydroxy-2-methyl-1-oxopropan-2-yl)-5-((4-fluoro-1-methyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 467	N-(1-amino-3-methoxy-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 468	N-(1-amino-3-methoxy-2-methyl-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 469	N-(1-amino-4-hydroxy-1-oxobutan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 470	N-(1-carbamoyl-3,3-difluorocyclobutyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 471	N-(1-carbamoylcyclobutyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 472	N-(1-carbamoylcyclobutyl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 473	N-(1-carbamoylcyclobutyl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 474	N-(1-carbamoylcyclobutyl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 475	N-(1-carbamoylcyclopropyl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 476	N-(1-carbamoylcyclopropyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 477	N-(1-carbamoylcyclopropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 478	N-(1-carbamoylcyclopropyl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 479	N-(1-carbamoylcyclopropyl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 480	N-(1-carbamoylcyclopropyl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 481	N-(1-hydroxy-2-methylpropan-2-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide
Cpd 482	N-(1-hydroxy-2-methylpropan-2-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 483	N-(1-hydroxy-2-methylpropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 484	N-(1-hydroxy-2-methylpropan-2-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 485	N-(1-hydroxy-2-methylpropan-2-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 486	N-(1-hydroxy-2-methylpropan-2-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 487	N-(1-hydroxy-3-morpholinopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide

-continued

Cpd 488	N-(1-hydroxy-3-morpholinopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 489	N-(1-hydroxypropan-2-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 490	N-(1-hydroxypropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 491	N-(1-hydroxypropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 492	(S)-N-(1-hydroxypropan-2-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 493	(R)-N-(1-hydroxypropan-2-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 494	(S)-N-(1-hydroxypropan-2-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 495	(R)-N-(1-hydroxypropan-2-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 496	(S)-N-(1-hydroxypropan-2-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 497	(R)-N-(1-hydroxypropan-2-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 498	N-(2,2-difluoroethyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 499	N-(2,2-difluoropropyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 500	N-(2,2-difluoropropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 501	N-(2-acetamidoethyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 502	N-(2-amino-1-cyano-2-oxoethyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 503	N-(2-amino-2-oxoethyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 504	N-(2-fluoro-2-methylpropyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 505	N-(2-fluoro-2-methylpropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 506	N-(2-hydroxy-2-methylpropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 507	N-(3-(aminomethyl)oxetan-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 508	N-(3-(dimethylamino)oxetane-3-carbonyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 509	N-(3-(dimethylcarbamoyl)oxetan-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 510	N-(3-(hydroxymethyl)-2-oxopiperidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 511	N-(3-(hydroxymethyl)-2-oxopiperidin-3-yl)-2-methyl-5-((2-methyloxazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 512	N-(3-(hydroxymethyl)-2-oxopiperidin-3-yl)-2-methyl-5-((2-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 513	N-(3-(hydroxymethyl)-2-oxopiperidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 514	N-(3-(hydroxymethyl)-2-oxopiperidin-3-yl)-2-methyl-5-((6-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 515	N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((2-methyloxazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 516	N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((2-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 517	N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 518	N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 519	N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((6-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 520	N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 521	N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 522	N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 523	N-(3-(hydroxymethyl)oxetan-3-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide
Cpd 524	N-(3-(hydroxymethyl)oxetan-3-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 525	N-(3-(hydroxymethyl)oxetan-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 526	N-(3-(hydroxymethyl)oxetan-3-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 527	N-(3-(hydroxymethyl)oxetan-3-yl)-2-methyl-5-(thiazol-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 528	N-(3-(hydroxymethyl)oxetan-3-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 529	N-(3-(hydroxymethyl)oxetan-3-yl)-N,2-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 530	N-(3-(hydroxymethyl)pyrrolidin-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 531	N-(3-(hydroxymethyl)tetrahydro-2H-pyran-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 532	N-(3-(hydroxymethyl)tetrahydro-2H-pyran-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide

-continued

Cpd 533	N-(3-(hydroxymethyl)tetrahydrofuran-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 534	N-(3-(hydroxymethyl)tetrahydrofuran-3-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 535	N-(3-(hydroxymethyl)tetrahydrofuran-3-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 536	N-(3-(hydroxymethyl)tetrahydrofuran-3-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 537	N-(3,3-difluoro-1-(hydroxymethyl)cyclobutyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 538	N-(3,3-difluoropiperidin-4-yl)-2,4-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 539	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-(((2-(trifluoromethyl)pyridin-3-yl)oxy)methyl)benzofuran-3-carboxamide
Cpd 540	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((1-methyl-1H-1,2,3-triazol-4-yl)methoxy)benzofuran-3-carboxamide
Cpd 541	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((1-methyl-1H-1,2,3-triazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 542	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((1-methyl-1H-1,2,4-triazol-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 543	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 544	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((1-methyl-2-oxo-1,2-dihydropyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 545	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((1-methyl-6-oxo-1,6-dihydropyridin-2-yl)methoxy)benzofuran-3-carboxamide
Cpd 546	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 547	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((2-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 548	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((2-methylpyrimidin-4-yl)methoxy)benzofuran-3-carboxamide
Cpd 549	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 550	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((3-methyloxetan-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 551	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((4-methyloxazol-2-yl)methoxy)benzofuran-3-carboxamide
Cpd 552	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((4-methylthiazol-2-yl)methoxy)benzofuran-3-carboxamide
Cpd 553	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((5-methylisoxazol-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 554	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((5-methylpyrazin-2-yl)methoxy)benzofuran-3-carboxamide
Cpd 555	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((5-methylthiazol-2-yl)methoxy)benzofuran-3-carboxamide
Cpd 556	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((6-methylpyridazin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 557	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-(pyridazin-3-ylmethoxy)benzofuran-3-carboxamide
Cpd 558	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-(pyridin-3-ylmethoxy)benzofuran-3-carboxamide
Cpd 559	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-(pyrimidin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 560	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-(thiazol-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 561	N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-(thiazol-5-ylmethoxy)benzofuran-3-carboxamide
Cpd 562	N-(3,3-difluoropiperidin-4-yl)-4-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 563	N-(3,3-difluoropiperidin-4-yl)-5-((2,4-dimethylthiazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 564	N-(3,3-difluoropiperidin-4-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 565	N-(3,3-difluoropiperidin-4-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 566	N-(3,3-difluoropiperidin-4-yl)-5-((3-fluoropyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 567	N-(3,3-difluoropiperidin-4-yl)-5-((5-(2-hydroxyethoxy)pyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 568	N-(3,3-difluoropiperidin-4-yl)-5-((5-fluoropyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 569	N-(3,3-difluoropiperidin-4-yl)-5-((6-hydroxypyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 570	N-(3,3-difluoropiperidin-4-yl)-5-((6-methoxypyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 571	N-(3,3-difluoropiperidin-4-yl)-5-(imidazo[1,2-a]pyridin-6-ylmethoxy)-2-methylbenzofuran-3-carboxamide
Cpd 572	N-(3,3-difluoropiperidin-4-yl)-6-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 573	N-(3,3-difluoropiperidin-4-yl)-7-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide

-continued

Cpd 574	N-(3,3-difluoropropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 575	N-(3-amino-2-(dimethylamino)-3-oxopropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 576	N-(3-amino-2,2-dimethyl-3-oxopropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 577	N-(3-carbamoyloxetan-3-yl)-2-(difluoromethyl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 578	N-(3-carbamoyloxetan-3-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide
Cpd 579	N-(3-carbamoyloxetan-3-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 580	N-(3-carbamoyloxetan-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 581	N-(3-carbamoyloxetan-3-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 582	N-(3-carbamoyloxetan-3-yl)-2-methyl-5-(thiazol-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 583	N-(3-carbamoyloxetan-3-yl)-4-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 584	N-(3-carbamoyloxetan-3-yl)-5-((2-(difluoromethyl)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 585	N-(3-carbamoyloxetan-3-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 586	N-(3-carbamoyloxetan-3-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 587	N-(3-carbamoyloxetan-3-yl)-6-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 588	N-(3-carbamoyloxetan-3-yl)-7-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 589	N-(3-carbamoyloxetan-3-yl)-N,2-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 590	N-(3-carbamoyltetrahydro-2H-pyran-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 591	N-(3-carbamoyltetrahydrofuran-3-yl)-2-methyl-5-((2-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 592	N-(3-carbamoyltetrahydrofuran-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 593	N-(3-carbamoyltetrahydrofuran-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 594	N-(3-carbamoyltetrahydrofuran-3-yl)-2-methyl-5-((6-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 595	N-(3-carbamoyltetrahydrofuran-3-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 596	N-(3-carbamoyltetrahydrofuran-3-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 597	N-(3-carbamoyltetrahydrofuran-3-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 598	N-(3-carbamoyltetrahydrofuran-3-yl)-5-((4-fluoro-1-methyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 599	N-(3-cyanooxetan-3-yl)-N,2-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 600	N-(3-hydroxy-1-(methylamino)-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 601	N-(3-hydroxy-1-(methylamino)-1-oxopropan-2-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 602	N-(3-hydroxy-2,2-dimethylpropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 603	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2,4-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 604	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((1-methyl-1H-1,2,3-triazol-4-yl)methoxy)benzofuran-3-carboxamide
Cpd 605	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((1-methyl-1H-1,2,3-triazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 606	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((1-methyl-1H-1,2,4-triazol-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 607	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((1-methyl-1H-imidazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 608	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 609	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((1-methyl-2-oxo-1,2-dihydropyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 610	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((1-methyl-6-oxo-1,6-dihydropyridin-2-yl)methoxy)benzofuran-3-carboxamide
Cpd 611	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((2-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 612	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((2-methylpyrimidin-4-yl)methoxy)benzofuran-3-carboxamide
Cpd 613	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 614	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((2-oxo-1,2-dihydropyridin-3-yl)methoxy)benzofuran-3-carboxamide

Cpd 615	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((3-(trifluoromethyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxamide
Cpd 616	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methyloxazol-2-yl)methoxy)benzofuran-3-carboxamide
Cpd 617	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-2-yl)methoxy)benzofuran-3-carboxamide
Cpd 618	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((5-methylisoxazol-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 619	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((5-methylpyrazin-2-yl)methoxy)benzofuran-3-carboxamide
Cpd 620	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((5-methylthiazol-2-yl)methoxy)benzofuran-3-carboxamide
Cpd 621	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((6-methylpyridazin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 622	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((6-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 623	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-(1-(4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 624	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-(pyridazin-3-ylmethoxy)benzofuran-3-carboxamide
Cpd 625	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-(pyridin-3-ylmethoxy)benzofuran-3-carboxamide
Cpd 626	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-(pyrimidin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 627	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-(thiazol-5-ylmethoxy)benzofuran-3-carboxamide
Cpd 628	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-hydroxypyridin-4-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 629	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-methoxypyridin-4-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 630	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((6-hydroxypyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 631	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((6-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 632	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((6-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 633	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-(imidazo[1,2-a]pyridin-6-ylmethoxy)-2-methylbenzofuran-3-carboxamide
Cpd 634	N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-N,2-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 635	N-(4,4-difluoropiperidin-3-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide
Cpd 636	N-(4,4-difluoropiperidin-3-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 637	N-(4,4-difluoropiperidin-3-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 638	N-(4,4-difluoropyrrolidin-3-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide
Cpd 639	N-(4,4-difluoropyrrolidin-3-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 640	N-(4,4-difluoropyrrolidin-3-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 641	N-(4,4-difluorotetrahydrofuran-3-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide
Cpd 642	N-(4,4-difluorotetrahydrofuran-3-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 643	N-(4,4-difluorotetrahydrofuran-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 646	N-(4,4-difluorotetrahydrofuran-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 647	N-(4,4-difluorotetrahydrofuran-3-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 648	N-(4,4-difluorotetrahydrofuran-3-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 649	N-(4-amino-2-methyl-4-oxobutan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 650	N-(4-carbamoyltetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 651	N-(4-hydroxy-2-methylbutan-2-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 652	N-(4-hydroxy-2-methylbutan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 653	N-(cis-4-hydroxytetrahydrofuran-3-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide
Cpd 654	N-(trans-4-hydroxytetrahydrofuran-3-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide
Cpd 655	N-(cis-4-hydroxytetrahydrofuran-3-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 656	N-(trans-4-hydroxytetrahydrofuran-3-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide

-continued

Cpd 657	N3-(3,3-difluoropiperidin-4-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-2,3-dicarboxamide
Cpd 658	N3-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-2,3-dicarboxamide
Cpd 659	N-cyclobutyl-N-(2-hydroxyethyl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 660	N-cyclobutyl-N-(2-hydroxyethyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 661	2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(2-oxazepan-3-yl)benzofuran-3-carboxamide
Cpd 662	N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-(pyridin-3-yl)methoxybenzofuran-3-carboxamide
Cpd 663	N-(4,4-difluoropyrrolidin-3-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 664	N-(4,4-difluorotetrahydrofuran-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 665	N-(4,4-difluorotetrahydrofuran-3-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 666	2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(2,2,2-trifluoroethyl)benzofuran-3-carboxamide
Cpd 667	5-((1,4-Dimethyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 668	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-5-((1,4-dimethyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 669	5-((4-Chloro-1-methyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 670	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-5-((4-chloro-1-methyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 671	5-((4-chloro-1-isopropyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 672	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((4-chloro-1-isopropyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 673	2-Methyl-5-((3-methylisoxazol-5-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 674	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((3-methylisoxazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 675	5-((1-isopropyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 676	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((1-isopropyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 677	5-((2,4-Dimethyloxazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 678	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-5-((2,4-dimethyloxazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 679	2-Methyl-5-((4-methyloxazol-5-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 680	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((4-methyloxazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 681	5-((2-Isopropylthiazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 682	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-5-((2-isopropylthiazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 683	N-(1-(Difluoromethyl)cyclobutyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 684	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((5-methylthiazol-2-yl)methoxy)benzofuran-3-carboxamide
Cpd 685	N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-2-methyl-5-((3-(trifluoromethyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxamide
Cpd 686	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((1-methyl-1H-pyrazol-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 687	2-Methyl-5-((1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 688	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 689	5-((1-(Cyclopropylmethyl)-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 690	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((1-(cyclopropylmethyl)-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 691	2-Methyl-5-((1-methyl-3-(trifluoromethyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 692	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((1-methyl-3-(trifluoromethyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxamide
Cpd 693	2-Methyl-5-((4-(trifluoromethyl)thiazol-5-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 694	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((4-(trifluoromethyl)thiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 695	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-(thiazol-2-yl)methoxybenzofuran-3-carboxamide
Cpd 696	N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-5-((5-fluoropyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 697	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-(thiazol-5-yl)methoxybenzofuran-3-carboxamide
Cpd 698	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((1-methyl-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxamide
Cpd 699	2-Methyl-5-((5-methylpyridin-2-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 700	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((5-methylpyridin-2-yl)methoxy)benzofuran-3-carboxamide
Cpd 701	5-((2-cyclopropylpyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 702	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-5-((2-cyclopropylpyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 703	5-((3-Cyclopropylpyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 704	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-5-((3-cyclopropylpyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide;

-continued


Cpd 705	5-((3-cyclopropylpyridin-2-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
Cpd 706	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((2-(azetidin-1-yl)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 707	2-Methyl-5-((2-methylpyrimidin-5-yl)methoxy)benzofuran-3-carboxylic acid
Cpd 708	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((2-methylpyrimidin-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 709	5-((2-Cyclopropoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
Cpd 710	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((2-cyclopropoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 711	5-((2-cyclopropoxypyridin-3-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
Cpd 712	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-(pyrimidin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 713	2-Methyl-5-(pyrazin-2-ylmethoxy)benzofuran-3-carboxylic acid
Cpd 714	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-(pyrazin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 715	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 716	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-(pyridin-4-ylmethoxy)benzofuran-3-carboxamide
Cpd 717	5-((3-(difluoromethyl)pyridin-2-yl)methoxy)-N-(3,3-difluoropiperidin-4-yl)-2-methylbenzofuran-3-carboxamide
Cpd 718	N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 719	N-(3-(hydroxymethyl)-2-oxopiperidin-3-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 720	N-(1-amino-3-hydroxy-2-methyl-1-oxopropan-2-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 721	N-(1-(Difluoromethyl)cyclobutyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 722	N-(1-Amino-3-hydroxy-2-methyl-1-oxopropan-2-yl)-2-methyl-5-((2-methylxazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 723	N-(3,3-difluorotetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 724	N-(3,3-Difluorotetrahydro-2H-pyran-4-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 725	N-(3-carbamoyltetrahydrofuran-3-yl)-2-methyl-5-((2-methylxazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 726	N-(1-Amino-3-(methylsulfonyl)-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 727	N-(1-Imino-1-oxidotetrahydro-1H-116-thiophen-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 728	N-(3-(hydroxymethyl)pyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 729	N-(1-Amino-1-oxo-3-(1H-1,2,4-triazol-1-yl)propan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 730	N-(3-(Hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((3-(trifluoromethyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxamide
Cpd 731	N-(3-(hydroxymethyl)-2-oxopiperidin-3-yl)-2-methyl-5-((3-(trifluoromethyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxamide
Cpd 732	N-(3-(Hydroxymethyl)-2-oxopiperidin-3-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 733	N-(3-(Hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 734	N-(3-(Hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-(pyridazin-3-ylmethoxy)benzofuran-3-carboxamide
Cpd 735	5-((5-Fluoropyridin-2-yl)methoxy)-N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methylbenzofuran-3-carboxamide
Cpd 736	5-((2-(Difluoromethyl)pyridin-3-yl)methoxy)-N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methylbenzofuran-3-carboxamide
Cpd 737	5-((2-(Difluoromethyl)pyridin-3-yl)methoxy)-N-(3-(hydroxymethyl)-2-oxopiperidin-3-yl)-2-methylbenzofuran-3-carboxamide
Cpd 738	N-(1-Amino-3-hydroxy-2-methyl-1-oxopropan-2-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 739	N-(1-Amino-3-hydroxy-2-methyl-1-oxopropan-2-yl)-5-((5-fluoropyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 740	N-(3-carbamoyltetrahydrofuran-3-yl)-5-((1-(2,2-difluoroethyl)-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 741	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-5-((5-fluoropyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide
Cpd 742	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-(difluoromethyl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 743	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-(difluoromethyl)-5-((6-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 744	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-(difluoromethyl)-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 745	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-(difluoromethyl)-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide

Cpd 746	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-(difluoromethyl)-5-((2-methyloxazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 747	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-cyclopropyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
Cpd 748	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-cyclopropyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 749	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-cyclopropyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
Cpd 750	(S)-N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-cyclopropyl-5-((6-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
Cpd 751	(S)-N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-cyclopropyl-5-((2-methyloxazol-5-yl)methoxy)benzofuran-3-carboxamide

and the physiologically acceptable salts thereof.


The benzofuran derivative according to the invention is for use in the treatment of pain which is preferably selected from nociceptive pain, inflammatory pain, and neuropathic pain. More preferably, the pain is post-operative pain.

Another aspect of the invention relates to a compound of formula (I),




a stereo-isomeric form, a physiologically acceptable salt, solvate and/or polymorph thereof, as defined above; preferably wherein

- (a-1) Q represents —OR^2 ; and R^1 represents $\text{—CH}_2\text{F}$, —CHF_2 , or —CF_3 ; and/or
- (a-2) Q represents —OR^2 ; and at least one of R^5 and $\text{R}^{5'}$ does not represent —H ; and/or
- (a-3) Q represents —OR^2 ; and R^6 does not represent —H ; and/or
- (a-3) Q represents —OR^2 ; and R^8 does not represent —H ; or
- (b-1) Q represents $\text{—NR}^3\text{R}^4$; with the proviso that the following compounds and their salts are excluded



-continued



and/or

- (b-2) Q represents $\text{—NR}^3\text{R}^4$; and R^1 represents $\text{—CH}_2\text{F}$, —CHF_2 , —CF_3 , or —cyclopropyl ; and/or
- (b-3) Q represents $\text{—NR}^3\text{R}^4$; and R^1 represents $\text{—CH}_2\text{F}$, —CHF_2 , or —CF_3 ; and/or
- (b-4) Q represents $\text{—NR}^3\text{R}^4$; and at least one of R^5 and $\text{R}^{5'}$ does not represent —H ; and/or
- (b-5) Q represents $\text{—NR}^3\text{R}^4$; and R^3 represents —H .

In preferred embodiments of the benzofuran derivatives according to the invention (a-1), (a-2), (a-3), (b-1), (b-2), (b-3), and (b-4) T represents —O— and U represents $\text{—CR}^5\text{R}^{5'}$ (i.e., the benzofuran derivatives is of formula (II)).

All definitions, preferred embodiments and preferred meanings of Q, T, U, V, R^1 , R^2 , R^3 , R^4 , R^5 , $\text{R}^{5'}$, R^6 , R^7 , and R^8 including the preferred substituents also analogously apply the benzofuran derivatives according to the invention, including but not limited to (a-1), (a-2), (a-3), (b-1), (b-2), (b-3), and (b-4), which are not necessarily restricted for use in the treatment of pain. Thus, this aspect of the invention relates to the benzofuran derivatives as such, compositions comprising the benzofuran derivatives, medicaments comprising the benzofuran derivatives, and the benzofuran derivatives for use in the prevention and/or treatment of TRPM3 mediated disorders such as pain and/or inflammatory hypersensitivity; and/or for counteracting pain and/or inflammatory hypersensitivity. Preferably, the pain is selected from nociceptive pain, inflammatory pain, and neuropathic pain. More preferably, the pain is post-operative pain.

In preferred embodiments of the invention, the benzo-furan derivative is selected from the group consisting of cpd 001 to cpd 207 as mentioned above and the physiologically acceptable salts thereof.

Another aspect of the invention relates to a pharmaceutical composition or a medicament comprising a compound according to the invention as described above.

Reference throughout this specification to "one embodiment" or "an embodiment" means that a particular feature, structure or characteristic described in connection with the embodiment is included in at least one embodiment of the invention. Thus, appearances of the phrases "in one embodiment" or "in an embodiment" in various places throughout this specification are not necessarily all referring to the same embodiment, but may. Furthermore, the particular features, structures or characteristics may be combined in any suitable manner, as would be apparent to one of ordinary skill in the art from this disclosure, in one or more embodiments. Also, embodiments described for an aspect of the invention may be used for another aspect of the invention and can be combined. Where an indefinite or definite article is used when referring to a singular noun e.g., "a" or "an", "the", this includes a plural of that noun unless something else is specifically stated.

Similarly, it should be appreciated that in the description of exemplary embodiments of the invention, various features of the invention are sometimes grouped together in a single embodiment, figure, or description thereof for the purpose of streamlining the disclosure and aiding in the understanding of one or more of the various inventive aspects.

In each of the following definitions, the number of carbon atoms represents the maximum number of carbon atoms generally optimally present in the substituent or linker; it is understood that where otherwise indicated in the present application, the number of carbon atoms represents the optimal maximum number of carbon atoms for that particular substituent or linker.

The term "leaving group" or "LG" as used herein means a chemical group which is susceptible to be displaced by a nucleophile or cleaved off or hydrolyzed in basic or acidic conditions. In a particular embodiment, a leaving group is selected from a halogen atom (e.g., Cl, Br, I) or a sulfonate (e.g., mesylate, tosylate, triflate).

The term "protecting group" refers to a moiety of a compound that masks or alters the properties of a functional group or the properties of the compound as a whole. The chemical substructure of a protecting group varies widely. One function of a protecting group is to serve as intermediates in the synthesis of the parental drug substance. Chemical protecting groups and strategies for protection/deprotection are well known in the art. See: "Protective Groups in Organic Chemistry", Theodora W. Greene (John Wiley & Sons, Inc., New York, 1991). Protecting groups are often utilized to mask the reactivity of certain functional groups, to assist in the efficiency of desired chemical reactions, e.g., making and breaking chemical bonds in an ordered and planned fashion. Protection of functional groups of a compound alters other physical properties besides the reactivity of the protected functional group, such as the polarity, lipophilicity (hydrophobicity), and other properties which can be measured by common analytical tools. Chemically protected intermediates may themselves be biologically active or inactive.

Protected compounds may also exhibit altered, and in some cases, optimized properties in vitro and in vivo, such as passage through cellular membranes and resistance to

enzymatic degradation or sequestration. In this role, protected compounds with intended therapeutic effects may be referred to as prodrugs. Another function of a protecting group is to convert the parental drug into a prodrug, whereby the parental drug is released upon conversion of the prodrug in vivo. Because active prodrugs may be absorbed more effectively than the parental drug, prodrugs may possess greater potency in vivo than the parental drug. Protecting groups are removed either in vitro, in the instance of chemical intermediates, or in vivo, in the case of prodrugs. With chemical intermediates, it is not particularly important that the resulting products after deprotection, e.g., alcohols, be physiologically acceptable, although in general it is more desirable if the products are pharmacologically innocuous.

The term "heteroatom(s)" as used herein means an atom selected from nitrogen, which can be quaternized; oxygen; and sulfur, including sulfoxide and sulfone.

The term "alkyl, saturated or unsaturated" as used herein encompasses saturated alkyl as well as unsaturated alkyl such as alkenyl, alkynyl, and the like. The term "alkyl" as used herein means normal, secondary, or tertiary, linear or branched hydrocarbon with no site of unsaturation. Examples are methyl, ethyl, 1-propyl (n-propyl), 2-propyl (iPr), 1-butyl, 2-methyl-1-propyl (i-Bu), 2-butyl (s-Bu), 2-dimethyl-2-propyl (t-Bu), 1-pentyl (n-pentyl), 2-pentyl, 3-pentyl, 2-methyl-2-butyl, 3-methyl-2-butyl, 3-methyl-1-butyl, 2-methyl-1-butyl, 1-hexyl, 2-hexyl, 3-hexyl, 2-methyl-2-pentyl, 3-methyl-2-pentyl, 4-methyl-2-pentyl, 3-methyl-3-pentyl, 2-methyl-3-pentyl, 2,3-dimethyl-2-butyl, and 3,3-dimethyl-2-butyl. The term "alkenyl" as used herein means normal, secondary or tertiary, linear or branched hydrocarbon with at least one site (usually 1 to 3, preferably 1) of unsaturation, namely a carbon-carbon, sp² double bond. Examples include, but are not limited to: ethylene or vinyl ($-\text{CH}=\text{CH}_2$), allyl ($-\text{CH}_2\text{CH}=\text{CH}_2$), and 5-hexenyl ($-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}=\text{CH}_2$). The double bond may be in the cis or trans configuration. The term "alkynyl" as used herein means normal, secondary, tertiary, linear or branched hydrocarbon with at least one site (usually 1 to 3, preferably 1) of unsaturation, namely a carbon-carbon, sp triple bond. Examples include, but are not limited to: ethynyl ($-\text{C}\equiv\text{CH}$), and 1-propynyl (propargyl, $-\text{CH}_2\text{C}\equiv\text{CH}$).

The term "alkylene, saturated or unsaturated" as used herein encompasses saturated alkylene as well as unsaturated alkylene such as alkenylene, alkynylene, alkenynylene and the like. The term "alkylene" as used herein means saturated, linear or branched chain hydrocarbon radical having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent alkane. Typical alkylene radicals include, but are not limited to: methylene ($-\text{CH}_2-$), 1,2-ethyl ($-\text{CH}_2\text{CH}_2-$), 1,3-propyl ($-\text{CH}_2\text{CH}_2\text{CH}_2-$), 1,4-butyl ($-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$), and the like. The term "alkenylene" as used herein means linear or branched chain hydrocarbon radical with at least one site (usually 1 to 3, preferably 1) of unsaturation, namely a carbon-carbon, sp² double bond, and having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent alkene. The term "alkynylene" as used herein means linear or branched chain hydrocarbon radical with at least one site (usually 1 to 3, preferably 1) of unsaturation, namely a carbon-carbon, sp triple bond, and having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent alkyne.

The term "heteroalkyl, saturated or unsaturated" as used herein encompasses saturated heteroalkyl as well as unsaturated heteroalkyl such as heteroalkenyl, heteroalkynyl, heteroalkenynyl and the like. The term "heteroalkyl" as used herein means linear or branched chain alkyl wherein one or more carbon atoms (usually 1, 2 or 3) are replaced by a heteroatom, i.e., an oxygen, nitrogen or sulfur atom, with the proviso that said chain may not contain two adjacent O atoms or two adjacent S atoms. This means that one or more —CH₃ of said alkyl can be replaced by —NH₂ and/or that one or more —CH₂— of said alkyl can be replaced by —NH—, —O— or —S—. The S atoms in said chains may be optionally oxidized with one or two oxygen atoms, to afford sulfoxides and sulfones, respectively. Furthermore, the heteroalkyl groups in the benzofuran derivatives of the invention can contain an oxo or thio group at any carbon or heteroatom that will result in a stable compound. Exemplary heteroalkyl groups include, but are not limited to, alcohols, alkyl ethers (such as for example -methoxy, -ethoxy, -butoxy . . .), primary, secondary, and tertiary alkyl amines, amides, ketones, esters, alkyl sulfides, and alkyl sulfones. The term "heteroalkenyl" means linear or branched chain alkenyl wherein one or more carbon atoms (usually 1, 2 or 3) are replaced by an oxygen, nitrogen or sulfur atom, with the proviso that said chain may not contain two adjacent O atoms or two adjacent S atoms. The term heteroalkenyl thus comprises imines, —O-alkenyl, —NH-alkenyl, —N(alkenyl)₂, —N(alkyl)(alkenyl), and —S-alkenyl. The term "heteroalkynyl" as used herein means linear or branched chain alkynyl wherein one or more carbon atoms (usually 1, 2 or 3) are replaced by an oxygen, nitrogen or sulfur atom, with the proviso that said chain may not contain two adjacent O atoms or two adjacent S atoms. The term heteroalkynyl thus comprises -cyano, —O-alkynyl, —NH-alkynyl, —N(alkynyl)₂, —N(alkyl)(alkynyl), —N(alkenyl)(alkynyl), and —S-alkynyl.

The term "heteroalkylene, saturated or unsaturated" as used herein encompasses saturated heteroalkylene as well as unsaturated heteroalkylene such as heteroalkenylene, heteroalkynylene, heteroalkenynylene and the like. The term "heteroalkylene" as used herein means linear or branched chain alkylene wherein one or more carbon atoms (usually 1, 2 or 3) are replaced by a heteroatom, i.e., an oxygen, nitrogen or sulfur atom, with the proviso that said chain may not contain two adjacent O atoms or two adjacent S atoms. The term "heteroalkynylene" as used herein means linear or branched chain alkynylene wherein one or more carbon atoms (usually 1, 2 or 3) are replaced by an oxygen, nitrogen or sulfur atom, with the proviso that said chain may not contain two adjacent O atoms or two adjacent S atoms.

The term "cycloalkyl, saturated or unsaturated" as used herein encompasses saturated cycloalkyl as well as unsaturated cycloalkyl such as cycloalkenyl, cycloalkynyl and the like. The term "cycloalkyl" as used herein and unless otherwise stated means a saturated cyclic hydrocarbon radical, such as for instance cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, norbornyl, fenchyl, decalinyl, adamantyl and the like. The term "cycloalkenyl" as used herein means a non-aromatic cyclic hydrocarbon radical with at least one site (usually 1 to 3, preferably 1) of unsaturation, namely a carbon-carbon, sp² double bond. Examples include, but are not limited to cyclopentenyl and

cyclohexenyl. The double bond may be in the cis or trans configuration. The term "cycloalkynyl" as used herein means a non-aromatic cyclic hydrocarbon radical with at least one site (usually 1 to 3, preferably 1) of unsaturation, namely a carbon-carbon, sp triple. An example is cyclohept-1-yne. Fused systems of a cycloalkyl ring with a heterocycloalkyl ring are considered as heterocycloalkyl irrespective of the ring that is bound to the core structure. Fused systems of a cycloalkyl ring with an aryl ring are considered as aryl irrespective of the ring that is bound to the core structure. Fused systems of a cycloalkyl ring with a heteroaryl ring are considered as heteroaryl irrespective of the ring that is bound to the core structure.

The term "heterocycloalkyl, saturated or unsaturated" as used herein encompasses saturated heterocycloalkyl as well as unsaturated non-aromatic heterocycloalkyl including at least one heteroatom, i.e., an N, O, or S as ring member. The term "heterocycloalkyl" as used herein and unless otherwise stated means "cycloalkyl" wherein one or more carbon atoms (usually 1, 2 or 3) are replaced by an oxygen, nitrogen or sulfur atom, with the proviso that said chain may not contain two adjacent O atoms or two adjacent S atoms. The term "heterocycloalkenyl" as used herein and unless otherwise stated means "cycloalkenyl" wherein one or more carbon atoms (usually 1, 2 or 3) are replaced by an oxygen, nitrogen or sulfur atom, with the proviso that said chain may not contain two adjacent O atoms or two adjacent S atoms. The term "heterocycloalkynyl" as used herein and unless otherwise stated means "cycloalkynyl" wherein one or more carbon atoms (usually 1, 2 or 3) are replaced by an oxygen, nitrogen or sulfur atom, with the proviso that said chain may not contain two adjacent O atoms or two adjacent S atoms. Examples of saturated and unsaturated heterocycloalkyl include but are not limited to azepane, 1,4-oxazepane, azetane, azetidine, aziridine, azocane, diazepane, dioxane, dioxolane, dithiane, dithiolane, imidazolidine, isothiazolidine, isoxalidine, morpholine, oxazolidine, oxepane, oxetane, oxirane, piperazine, piperidine, pyrazolidine, pyrrolidine, quinuclidine, tetrahydrofuran, tetrahydropyran, tetrahydrothiopyran, thiazolidine, thietane, thiirane, thiolane, thiomorpholine, indoline, dihydrobenzofuran, dihydrobenzothiophene, 1,1-dioxothiacyclohexane, 2-azaspiro[3.3]heptane, 2-oxaspiro[3.3]heptane, 7-azaspiro[3.5]nonane, 8-azabicyclo[3.2.1]octane, 9-azabicyclo[3.3.1]nonane, hexahydro-1H-pyrrolizine, hexahydrocyclopenta[c]pyrrole, octahydrocyclopenta[c]pyrrole, and octahydropyrrolo[1,2-a]pyrazin. Further heterocycloalkyls in the meaning of the invention are described in Paquette, Leo A. "Principles of Modern Heterocyclic Chemistry" (W. A. Benjamin, New York, 1968), particularly Chapters 1, 3, 4, 6, 7, and 9; "The Chemistry of Heterocyclic Compounds, A series of Monographs" (John Wiley & Sons, New York, 1950 to present), in particular Volumes 13, 14, 16, 19, and 28; Katritzky, Alan R., Rees, C. W. and Scriven, E. "Comprehensive Heterocyclic Chemistry" (Pergamon Press, 1996); and J. Am. Chem. Soc. (1960) 82:5566. When the heterocycloalkyl contains no nitrogen as ring member, it is typically bonded through carbon. When the heterocycloalkyl contains nitrogen as ring member, it may be bonded through nitrogen or carbon. Fused systems of heterocycloalkyl ring with a cycloalkyl ring are considered as heterocycloalkyl irrespective of the ring that is bound to the core structure. Fused systems of a heterocycloalkyl ring with an aryl ring are considered as heterocycloalkyl irrespective of the ring that is bound to the core structure. Fused systems of a

heterocycloalkyl ring with a heteroaryl ring are considered as heteroaryl irrespective of the ring that is bound to the core structure.

The term "aryl" as used herein means an aromatic hydrocarbon. Typical aryl groups include, but are not limited to 1 ring, or 2 or 3 rings fused together, radicals derived from benzene, naphthalene, anthracene, biphenyl, and the like. Fused systems of an aryl ring with a cycloalkyl ring are considered as aryl irrespective of the ring that is bound to the core structure. Fused systems of an aryl ring with a heterocycloalkyl ring are considered as heterocycloalkyl irrespective of the ring that is bound to the core structure. Thus, indoline, dihydrobenzofuran, dihydrobenzothiophene and the like are considered as heterocycloalkyl according to the invention. Fused systems of an aryl ring with a heteroaryl ring are considered as heteroaryl irrespective of the ring that is bound to the core structure.

The term "heteroaryl" as used herein means an aromatic ring system including at least one heteroatom, i.e., N, O, or S as ring member of the aromatic ring system. Examples of heteroaryl include but are not limited to benzimidazole, benzisoxazole, benzoazole, benzodioxole, benzofuran, benzothiadiazole, benzothiazole, benzothiophene, carbazole, cinnoline, dibenzofuran, furane, furazane, imidazole, imidazopyridine, indazole, indole, indolizine, isobenzofuran, isoindole, isoquinoline, isothiazole, isoxazole, naphthyridine, oxadiazole, oxazole, oxindole, phthalazine, purine, pyrazine, pyrazole, pyridazine, pyridine, pyrimidine, pyrrole, quinazoline, quinoxaline, tetrazole, thiadiazole, thiazole, thiophene, triazine, triazole, and [1,2,4] triazolo[4,3-a]pyrimidine.

By further way of example, carbon bonded heterocyclic rings are bonded at position 2, 3, 4, 5, or 6 of a pyridine, position 3, 4, 5, or 6 of a pyridazine, position 2, 4, 5, or 6 of a pyrimidine, position 2, 3, 5, or 6 of a pyrazine, position 2, 3, 4, or 5 of a furan, tetrahydrofuran, thiophene, pyrrole or tetrahydropyrrole, position 2, 4, or 5 of an oxazole, imidazole or thiazole, position 3, 4, or 5 of an isoxazole, pyrazole, or isothiazole, position 2 or 3 of an aziridine, position 2, 3, or 4 of an azetidine, position 2, 3, 4, 5, 6, 7, or 8 of a quinoline or position 1, 3, 4, 5, 6, 7, or 8 of an isoquinoline.

Preferred carbon bonded heterocycles include 2-pyridyl, 3-pyridyl, 4-pyridyl, 5-pyridyl, 6-pyridyl, 3-pyridazinyl, 4-pyridazinyl, 5-pyridazinyl, 6-pyridazinyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 6-pyrimidinyl, 2-pyrazinyl, 3-pyrazinyl, 5-pyrazinyl, 6-pyrazinyl, 2-thiazolyl, 4-thiazolyl, or 5-thiazolyl. By way of example, nitrogen bonded heterocyclic rings are bonded at position 1 of an aziridine, azetidine, pyrrole, pyrrolidine, 2-pyrroline, 3-pyrroline, imidazole, imidazolidine, 2-imidazoline, 3-imidazoline, pyrazole, pyrazoline, 2-pyrazoline, 3-pyrazoline, piperidine, piperazine, indole, indoline, 1H-indazole, position 2 of an isoindole, or isoindoline, position 4 of a morpholine, and position 9 of a carbazole, or β -carboline. Preferred nitrogen bonded heterocycles include 1-aziridyl, 1-azetetyl, 1-pyrrolyl, 1-imidazolyl, 1-pyrazolyl, and 1-piperidinyl. Further heteroaryls in the meaning of the invention are described in Paquette, Leo A. "Principles of Modern Heterocyclic Chemistry" (W. A. Benjamin, New York, 1968), particularly Chapters 1, 3, 4, 6, 7, and 9; "The Chemistry of Heterocyclic Compounds, A series of Monographs" (John Wiley & Sons, New York, 1950 to present), in particular Volumes 13, 14, 16, 19, and 28; Katritzky, Alan R., Rees, C. W. and Scriven, E. "Comprehensive Heterocyclic Chemistry" (Pergamon Press, 1996); and J. Am. Chem. Soc. (1960) 82:5566.

As used herein with respect to a substituting group, and unless otherwise stated, the terms "monosubstituted", "disubstituted", "trisubstituted", "polysubstituted" and the like means chemical structures defined herein, wherein the respective moiety is substituted with one or more substituents, meaning that one or more hydrogen atoms of said moiety are each independently replaced with a substituent. For example, $-C_{1-6}$ -alkyl that may be polysubstituted with $-F$ covers $-CH_2F$, $-CHF_2$, $-CF_3$, $-CH_2CF_3$, CF_2CF_3 , and the like. Likewise, $-C_{1-6}$ -alkyl that may be polysubstituted with substituents independently of one another selected from $-F$ and $-Cl$ covers $-CH_2F$, $-CHF_2$, $-CF_3$, $-CH_2CF_3$, CF_2CF_3 , $-CH_2Cl$, $-CHCl_2$, $-CCl_3$, $-CH_2CCl_3$, CCl_2CCl_3 , $-CHClF$, $-CClF_2$, $-CCl_2CF_3$, $-CF_2CCl_3$, $-CClFCCl_2F$, and the like. Any substituent designation that is found in more than one site in a compound of this invention shall be independently selected.

As used herein and unless otherwise stated, the term "solvate" includes any combination which may be formed by a derivative of this invention with a suitable inorganic solvent (e.g., hydrates) or organic solvent, such as but not limited to alcohols, ketones, esters, ethers, nitriles and the like.

The term "subject" as used herein, refers to an animal including humans, preferably a mammal, most preferably a human, who has been the object of treatment, observation or experiment.

The term "therapeutically effective amount" as used herein, means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes alleviation or partial alleviation of the symptoms of the disease or disorder being treated.

The term "composition" as used herein is intended to encompass a product comprising the specified ingredients in the therapeutically effective amounts, as well as any product which results, directly or indirectly, from combinations of the specified ingredients in the specified amounts.

The term "antagonist" or "inhibitor" as used herein refers to a compound capable of producing, depending on the circumstance, a functional antagonism of the TRPM3 ion channel, including competitive antagonists, non-competitive antagonists, desensitizing agonists, and partial agonists.

For purposes of the invention, the term "TRPM3-mediated" is used to refer to the condition of being affected by the modulation of the TRPM3 ion channel, including the state of being mediated by the TRPM3 ion channel.

The term "TRPM3 mediated disorder" as used herein refers to disorders or diseases for which the use of an antagonist of TRPM3 would prevent, treat, (partially) alleviate or improve the symptoms and consist of pain and inflammatory hypersensitivity condition. According to the International Association for the Study of Pain and for the purpose of the invention, pain is an unpleasant sensory and emotional experience associated with actual or potential tissue damage, or described in terms of such damage. Preferably, the TRPM3 mediated disorder is pain which is preferably selected from nociceptive pain, inflammatory pain, and neuropathic pain. More preferably, the pain is post-operative pain. For the purpose of the invention, the term "inflammatory hypersensitivity" is used to refer to a condition that is characterized by one or more hallmarks of inflammation, including edema, erythema, hyperthermia and pain, and/or by an exaggerated physiologic or pathophysiologic response to one or more than one type of stimulation, including thermal, mechanical and/or chemical stimulation.

The benzofuran derivatives of the invention have been shown to be antagonists of TRPM3 and the invention therefore provides the compounds as such, the compounds for use as a medicine, more specifically for use as a medicine in the prevention or treatment of TRPM3 mediated disorders in a subject with a therapeutically effective amount of a benzofuran derivative of the invention.

In a preferred embodiment of the invention, the benzofuran derivative of the invention is the sole pharmacologically active compound to be administered for therapy. In another preferred embodiment of the invention, the benzofuran derivative of the invention may be employed in combination with other therapeutic agents for the treatment or prophylaxis of TRPM3 mediated disorders. The invention therefore also relates to the use of a composition comprising:

one or more compounds of the formulae and embodiments herein, and

one or more further therapeutic or preventive agents that are used for the prevention or treatment of TRPM3 mediated disorders as biologically active agents in the form of a combined preparation for simultaneous, separate or sequential use.

The pharmaceutical composition or combined preparation according to this invention may contain benzofuran derivatives of the invention over a broad content range depending on the contemplated use and the expected effect of the preparation. Generally, the content of the benzofuran derivatives of the invention of the combined preparation is within the range of 0.1 to 99.9% by weight, preferably from 1 to 99% by weight, more preferably from 5 to 95% by weight.

In view of the fact that, when several active ingredients are used in combination, they do not necessarily bring out their joint therapeutic effect directly at the same time in the mammal to be treated, the corresponding composition may also be in the form of a medical kit or package containing the two ingredients in separate but adjacent repositories or compartments. In the latter context, each active ingredient may therefore be formulated in a way suitable for an administration route different from that of the other ingredient, e.g., one of them may be in the form of an oral or parenteral formulation whereas the other is in the form of an ampoule for intravenous injection or an aerosol.

Those of skill in the art will also recognize that the benzofuran derivatives of the invention may exist in many different protonation states, depending on, among other things, the pH of their environment. While the structural formulae provided herein depict the compounds in only one of several possible protonation states, it will be understood that these structures are illustrative only, and that the invention is not limited to any particular protonation state—any and all protonated forms of the compounds are intended to fall within the scope of the invention.

The term “pharmaceutically acceptable salts” as used herein means the therapeutically active non-toxic salt forms which the compounds of formulae herein are able to form. Therefore, the compounds of this invention optionally comprise salts of the compounds herein, especially pharmaceutically acceptable non-toxic salts containing, for example, Na^+ , Li^+ , K^+ , Ca^{2+} and Mg^{2+} . Such salts may include those derived by combination of appropriate cations such as alkali and alkaline earth metal ions or ammonium and quaternary amino ions with an acid anion moiety, typically a carboxylic acid. The benzofuran derivatives of the invention may bear multiple positive or negative charges. The net charge of the benzofuran derivatives of the invention may be either positive or negative. Any associated counter ions are typically dictated by the synthesis and/or isolation methods by which

the compounds are obtained. Typical counter ions include, but are not limited to ammonium, sodium, potassium, lithium, halides, acetate, trifluoroacetate, etc., and mixtures thereof. It will be understood that the identity of any associated counter ion is not a critical feature of the invention, and that the invention encompasses the compounds in association with any type of counter ion. Moreover, as the compounds can exist in a variety of different forms, the invention is intended to encompass not only forms of the compounds that are in association with counter ions (e.g., dry salts), but also forms that are not in association with counter ions (e.g., aqueous or organic solutions). Metal salts typically are prepared by reacting the metal hydroxide with a compound of this invention. Examples of metal salts which are prepared in this way are salts containing Li^+ , Na^+ , and K^+ . A less soluble metal salt can be precipitated from the solution of a more soluble salt by addition of the suitable metal compound. In addition, salts may be formed from acid addition of certain organic and inorganic acids to basic centers, typically amines, or to acidic groups. Examples of such appropriate acids include, for instance, inorganic acids such as hydrohalogen acids, e.g. hydrochloric or hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like; or organic acids such as, for example, acetic, propanoic, hydroxyacetic, 2-hydroxypropanoic, 2-oxopropanoic, lactic, pyruvic, oxalic (i.e. ethanedioic), malonic, succinic (i.e. butanedioic acid), maleic, fumaric, malic, tartaric, citric, methanesulfonic, ethanesulfonic, benzenesulfonic, p-toluenesulfonic, cyclohexanesulfamic, salicylic (i.e. 2-hydroxybenzoic), p-aminosalicylic and the like. Furthermore, this term also includes the solvates which the compounds of formulae herein as well as their salts are able to form, such as for example hydrates, alcoholates and the like. Finally, it is to be understood that the compositions herein comprise benzofuran derivatives of the invention in their unionized, as well as zwitterionic form, and combinations with stoichiometric amounts of water as in hydrates.

Also included within the scope of this invention are the salts of the parental compounds with one or more amino acids, especially the naturally-occurring amino acids found as protein components. The amino acid typically is one bearing a side chain with a basic or acidic group, e.g., lysine, arginine or glutamic acid, or a neutral group such as glycine, serine, threonine, alanine, isoleucine, or leucine.

The benzofuran derivatives of the invention also include physiologically acceptable salts thereof. Examples of physiologically acceptable salts of the benzofuran derivatives of the invention include salts derived from an appropriate base, such as an alkali metal (for example, sodium), an alkaline earth (for example, magnesium), ammonium and NX_4^+ (wherein X is $-\text{C}_{1-6}\text{-alkyl}$). Physiologically acceptable salts of a hydrogen atom or an amino group include salts of organic carboxylic acids such as acetic, benzoic, lactic, fumaric, tartaric, maleic, malonic, malic, isethionic, lactobionic and succinic acids; organic sulfonic acids, such as methanesulfonic, ethanesulfonic, benzenesulfonic and p-toluenesulfonic acids; and inorganic acids, such as hydrochloric, sulfuric, phosphoric and sulfamic acids. Physiologically acceptable salts of a compound containing a hydroxy group include the anion of said compound in combination with a suitable cation such as Na^+ and NX_4^+ (wherein X typically is independently selected from $-\text{H}$ or a $-\text{C}_{1-4}\text{-alkyl}$ group). However, salts of acids or bases which are not physiologically acceptable may also find use, for example, in the preparation or purification of a physiologically accept-

able compound. All salts, whether or not derived from a physiologically acceptable acid or base, are within the scope of the invention.

As used herein and unless otherwise stated, the term "enantiomer" means each individual optically active form of a benzofuran derivative of the invention, having an optical purity or enantiomeric excess (as determined by methods standard in the art) of at least 80% (i.e., at least 90% of one enantiomer and at most 10% of the other enantiomer), preferably at least 90% and more preferably at least 98%.

The term "isomers" as used herein means all possible isomeric forms, including tautomeric and stereochemical forms, which the compounds of formulae herein may possess, but not including position isomers. Typically, the structures shown herein exemplify only one tautomeric or resonance form of the compounds, but the corresponding alternative configurations are contemplated as well. Unless otherwise stated, the chemical designation of compounds denotes the mixture of all possible stereochemically isomeric forms, said mixtures containing all diastereomers and enantiomers (since the compounds of formulae herein may have at least one chiral center) of the basic molecular structure, as well as the stereochemically pure or enriched compounds. More particularly, stereogenic centers may have either the R- or S-configuration, and multiple bonds may have either cis- or trans-configuration.

Pure isomeric forms of the said compounds are defined as isomers substantially free of other enantiomeric or diastereomeric forms of the same basic molecular structure. In particular, the term "stereoisomerically pure" or "chirally pure" relates to compounds having a stereoisomeric excess of at least about 80% (i.e., at least 90% of one isomer and at most 10% of the other possible isomers), preferably at least 90%, more preferably at least 94% and most preferably at least 97%. The terms "enantiomerically pure" and "diastereomerically pure" should be understood in a similar way, having regard to the enantiomeric excess, respectively the diastereomeric excess, of the mixture in question.

Separation of stereoisomers is accomplished by standard methods known to those in the art. One enantiomer of a benzofuran derivative of the invention can be separated substantially free of its opposing enantiomer by a method such as formation of diastereomers using optically active resolving agents ("Stereochemistry of Carbon Compounds," (1962) by E. L. Eliel, McGraw Hill; Lochmuller, C. H., (1975) J. Chromatogr., 113(3) 283-302). Separation of isomers in a mixture can be accomplished by any suitable method, including: (1) formation of ionic, diastereomeric salts with chiral compounds and separation by fractional crystallization or other methods, (2) formation of diastereomeric compounds with chiral derivatizing reagents, separation of the diastereomers, and conversion to the pure enantiomers, or (3) enantiomers can be separated directly under chiral conditions. Under method (1), diastereomeric salts can be formed by reaction of enantiomerically pure chiral bases such as brucine, quinine, ephedrine, strychnine, a-methyl-b-phenylethylamine (amphetamine), and the like with asymmetric compounds bearing acidic functionality, such as carboxylic acid and sulfonic acid. The diastereomeric salts may be induced to separate by fractional crystallization or ionic chromatography. For separation of the optical isomers of amino compounds, addition of chiral carboxylic or sulfonic acids, such as camphorsulfonic acid, tartaric acid, mandelic acid, or lactic acid can result in formation of the diastereomeric salts. Alternatively, by method (2), the substrate to be resolved may be reacted with one enantiomer of a chiral compound to form a diastereo-

meric pair (Eliel, E. and Wilen, S. (1994) Stereochemistry of Organic Compounds, John Wiley & Sons, Inc., p. 322). Diastereomeric compounds can be formed by reacting asymmetric compounds with enantiomerically pure chiral derivatizing reagents, such as menthyl derivatives, followed by separation of the diastereomers and hydrolysis to yield the free, enantiomerically enriched compound. A method of determining optical purity involves making chiral esters, such as a menthyl ester or Mosher ester, a-methoxy-a-(trifluoromethyl)phenyl acetate (Jacob III. (1982) J. Org. Chem. 47:4165), of the racemic mixture, and analyzing the NMR spectrum for the presence of the two atropisomeric diastereomers. Stable diastereomers can be separated and isolated by normal- and reverse-phase chromatography following methods for separation of atropisomeric naphthyl-isoquinolines (Hoye, T., WO 96/15111). Under method (3), a racemic mixture of two asymmetric enantiomers is separated by chromatography using a chiral stationary phase. Suitable chiral stationary phases are, for example, polysaccharides, in particular cellulose or amylose derivatives. Commercially available polysaccharide based chiral stationary phases are ChiralCel® CA, OA, OB5, OC5, OD, OF, OG, OJ and OK, and Chiralpak® AD, AS, OP(+) and OT(+). Appropriate eluents or mobile phases for use in combination with said polysaccharide chiral stationary phases are hexane and the like, modified with an alcohol such as ethanol, isopropanol and the like. ("Chiral Liquid Chromatography" (1989) W. J. Lough, Ed. Chapman and Hall, New York; Okamoto, (1990) "Optical resolution of dihydropyridine enantiomers by High-performance liquid chromatography using phenylcarbamates of polysaccharides as a chiral stationary phase", J. of Chromatogr. 513:375-378).

The terms cis and trans are used herein in accordance with Chemical Abstracts nomenclature and include reference to the position of the substituents on a ring moiety. The absolute stereochemical configuration of the compounds of the formulae described herein may easily be determined by those skilled in the art while using well-known methods such as, for example, X-ray diffraction.

When a compound is crystallized from a solution or slurry, it can be crystallized in a different arrangement lattice of spaces (this property is called "polymorphism") to form crystals with different crystalline forms, each of which is known as "polymorphs". The term "Polymorph" as used herein, therefore, refers to a crystal form of a compound of Formula (I), where the molecules are localized in the three-dimensional lattice sites. Different polymorphs of the compound of Formula (I) may be different from each other in one or more physical properties, such as solubility and dissolution rate, true specific gravity, crystal form, accumulation mode, flowability and/or solid state stability. etc.

Benzofuran derivatives of the invention and their physiologically acceptable salts (hereafter collectively referred to as the active ingredients) may be administered by any route appropriate to the condition to be treated, suitable routes including oral, rectal, nasal, topical (including ocular, buccal and sublingual), vaginal and parenteral (including subcutaneous, intramuscular, intranasal, intravenous, intraarterial, intradermal, intrathecal and epidural). The preferred route of administration may vary with for example the condition of the recipient.

The therapeutically effective amount of the preparation of the compound(s), especially for the treatment of TRPM3 mediated disorders in humans and other mammals or in animals, preferably is a TRPM3 ion channel inhibiting amount of the compounds as defined herein and corresponds

to an amount which ensures a plasma level of between 1 µg/ml and 100 mg/ml, optionally of 10 mg/ml.

Suitable dosages of the compounds or compositions of the invention should be used to treat or prevent the TRPM3 mediated disorders in a subject. Depending upon the pathologic condition to be treated and the patient's condition, the said effective amount may be divided into several sub-units per day or may be administered at more than one day intervals.

The invention further provides (pharmaceutical) compositions comprising one or more benzofuran derivatives of the invention, more in particular of all the Formula (I) and other formulas and embodiments described herein and the more particular aspects or embodiments thereof. Furthermore, the invention provides the compounds or (pharmaceutical) compositions of the invention, more in particular of all the Formula (I) and other formulas and embodiments described herein and the more particular aspects or embodiments thereof, for use as a medicine, more in particular for use in the treatment of pain. The TRPM3 mediated disorders are selected from pain and an inflammatory hypersensitivity condition.

The benzofuran derivatives of the invention may be formulated with conventional carriers and excipients, which will be selected in accord with ordinary practice. Tablets will contain excipients, glidants, fillers, binders and the like. Aqueous formulations are prepared in sterile form, and when intended for delivery by other than oral administration generally will be isotonic. Formulations optionally contain excipients such as those set forth in the "Handbook of Pharmaceutical Excipients" (1986).

Subsequently, the term "pharmaceutically acceptable carrier" as used herein means any material or substance with which the active ingredient is formulated in order to facilitate its application or dissemination to the locus to be treated, for instance by dissolving, dispersing or diffusing the said composition, and/or to facilitate its storage, transport or handling without impairing its effectiveness. The pharmaceutically acceptable carrier may be a solid or a liquid or a gas which has been compressed to form a liquid, i.e., the compositions of this invention can suitably be used as concentrates, emulsions, solutions, granulates, dusts, sprays, aerosols, suspensions, ointments, creams, tablets, pellets or powders.

Suitable pharmaceutical carriers for use in the said pharmaceutical compositions and their formulation are well known to those skilled in the art, and there is no particular restriction to their selection within the invention. They may also include additives such as wetting agents, dispersing agents, stickers, adhesives, emulsifying agents, surface-active agents, solvents, coatings, antibacterial and antifungal agents, isotonic agents and the like, provided the same are consistent with pharmaceutical practice, i.e., carriers and additives which do not create permanent damage to mammals. The pharmaceutical compositions of the invention may be prepared in any known manner, for instance by homogeneously mixing, coating and/or grinding the active ingredients, in a one-step or multi-steps procedure, with the selected carrier material and, where appropriate, the other additives such as surface-active agents, may also be prepared by micronisation, for instance in view to obtain them in the form of microspheres usually having a diameter of about 1 to 10 µm, namely for the manufacture of microcapsules for controlled or sustained release of the active ingredients.

While it is possible for the benzofuran derivatives to be administered alone it is preferable to present them as phar-

maceutical formulations. The formulations, both for veterinary and for human use, of the invention comprise at least one active ingredient, as above described, together with one or more pharmaceutically acceptable carriers therefore and optionally other therapeutic ingredients. The carrier(s) optimally are "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof. The formulations include those suitable for oral, rectal, nasal, topical (including buccal and sublingual), vaginal or parenteral (including subcutaneous, intramuscular, intravenous, intradermal, intrathecal and epidural) administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. Such methods include the step of bringing into association the active ingredient with the carrier which constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

Formulations of the invention suitable for oral administration may be presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient; as a powder or granules; as solution or a suspension in an aqueous liquid or a non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. The active ingredient may also be presented as a bolus, electuary or paste.

A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, preservative, surface active or dispersing agent. Molded tablets may be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent. The tablets may optionally be coated or scored and may be formulated so as to provide slow or controlled release of the active ingredient therein. For infections of the eye or other external tissues e.g., mouth and skin, the formulations are optionally applied as a topical ointment or cream containing the active ingredient(s) in an amount of, for example, 0.075 to 20% w/w (including active ingredient(s) in a range between 0.1% and 20% in increments of 0.1% w/w such as 0.6% w/w, 0.7% w/w, etc.), preferably 0.2 to 15% w/w and most preferably 0.5 to 10% w/w. When formulated in an ointment, the active ingredients may be employed with either a paraffinic or a water-miscible ointment base. Alternatively, the active ingredients may be formulated in a cream with an oil-in-water cream base. If desired, the aqueous phase of the cream base may include, for example, at least 30% w/w of a polyhydric alcohol, i.e., an alcohol having two or more hydroxyl groups such as propylene glycol, butane 1,3-diol, mannitol, sorbitol, glycerol and polyethylene glycol (including PEG400) and mixtures thereof. The topical formulations may desirably include a compound which enhances absorption or penetration of the active ingredient through the skin or other affected areas. Examples of such dermal penetration enhancers include dimethylsulfoxide and related analogs.

The oily phase of the emulsions of this invention may be constituted from known ingredients in a known manner. While the phase may comprise merely an emulsifier (otherwise known as an emulgent), it desirably comprises a mixture of at least one emulsifier with a fat or an oil or with both a fat and an oil. Optionally, a hydrophilic emulsifier is

included together with a lipophilic emulsifier which acts as a stabilizer. It is also preferred to include both an oil and a fat. Together, the emulsifier(s) with or without stabilizer(s) make up the so-called emulsifying wax, and the wax together with the oil and fat make up the so-called emulsifying ointment base which forms the oily dispersed phase of the cream formulations.

The choice of suitable oils or fats for the formulation is based on achieving the desired cosmetic properties, since the solubility of the active compound in most oils likely to be used in pharmaceutical emulsion formulations is very low. Thus, the cream should optionally be a non-greasy, non-staining and washable product with suitable consistency to avoid leakage from tubes or other containers. Straight or branched chain, mono- or dibasic alkyl esters such as di-isoadipate, isocetyl stearate, propylene glycol diester of coconut fatty acids, isopropyl myristate, decyl oleate, isopropyl palmitate, butyl stearate, 2-ethylhexyl palmitate or a blend of branched chain esters known as Crodamol CAP may be used, the last three being preferred esters. These may be used alone or in combination depending on the properties required. Alternatively, high melting point lipids such as white soft paraffin and/or liquid paraffin or other mineral oils can be used.

Formulations suitable for topical administration to the eye also include eye drops wherein the active ingredient is dissolved or suspended in a suitable carrier, especially an aqueous solvent for the active ingredient. The active ingredient is optionally present in such formulations in a concentration of 0.5 to 20%, advantageously 0.5 to 10% particularly about 1.5% w/w. Formulations suitable for topical administration in the mouth include lozenges comprising the active ingredient in a flavored basis, usually sucrose and acacia or tragacanth; pastilles comprising the active ingredient in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouthwashes comprising the active ingredient in a suitable liquid carrier.

Formulations for rectal administration may be presented as a suppository with a suitable base comprising for example cocoa butter or a salicylate. Formulations suitable for nasal administration wherein the carrier is a solid include a coarse powder having a particle size for example in the range 20 to 500 microns (including particle sizes in a range between 20 and 500 microns in increments of 5 microns such as 30 microns, 35 microns, etc.), which is administered in the manner in which snuff is taken, i.e. by rapid inhalation through the nasal passage from a container of the powder held close up to the nose. Suitable formulations wherein the carrier is a liquid, for administration as for example a nasal spray or as nasal drops, include aqueous or oily solutions of the active ingredient. Formulations suitable for aerosol administration may be prepared according to conventional methods and may be delivered with other therapeutic agents.

Formulations suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or spray formulations containing in addition to the active ingredient such carriers as are known in the art to be appropriate.

Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze-dried

(lyophilized) condition requiring only the addition of the sterile liquid carrier, for example water for injections, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the kind previously described.

Preferred unit dosage formulations are those containing a daily dose or unit daily sub-dose, as herein above recited, or an appropriate fraction thereof, of an active ingredient.

It should be understood that in addition to the ingredients particularly mentioned above the formulations of this invention may include other agents conventional in the art having regard to the type of formulation in question, for example those suitable for oral administration may include flavoring agents.

Benzofuran derivatives of the invention can be used to provide controlled release pharmaceutical formulations containing as active ingredient one or more benzofuran derivatives of the invention ("controlled release formulations") in which the release of the active ingredient can be controlled and regulated to allow less frequency dosing or to improve the pharmacokinetic or toxicity profile of a given invention compound. Controlled release formulations adapted for oral administration in which discrete units comprising one or more benzofuran derivatives of the invention can be prepared according to conventional methods.

Another embodiment of this invention relates to various precursor or "prodrug" forms of the benzofuran derivatives of the invention. It may be desirable to formulate the benzofuran derivatives of the invention in the form of a chemical species which itself is not significantly biologically-active, but which when delivered to the animal, mammal or human will undergo a chemical reaction catalyzed by the normal function of the body, inter alia, enzymes present in the stomach or in blood serum, said chemical reaction having the effect of releasing a compound as defined herein. The term "prodrug" thus relates to these species which are converted in vivo into the active pharmaceutical ingredient.

The prodrugs of the benzofuran derivatives of the invention can have any form suitable to the formulator, for example, esters are non-limiting common pro-drug forms. In the present case, however, the pro-drug may necessarily exist in a form wherein a covalent bond is cleaved by the action of an enzyme present at the target locus. For example, a C—C covalent bond may be selectively cleaved by one or more enzymes at said target locus and, therefore, a pro-drug in a form other than an easily hydrolysable precursor, inter alia an ester, an amide, and the like, may be used. The counterpart of the active pharmaceutical ingredient in the pro-drug can have different structures such as an amino acid or peptide structure, alkyl chains, sugar moieties and others as known in the art.

For the purpose of the invention the term "therapeutically suitable pro-drug" is defined herein as "a compound modified in such a way as to be transformed in vivo to the therapeutically active form, whether by way of a single or by multiple biological transformations, when in contact with the tissues of the animal, mammal or human to which the pro-drug has been administered, and without undue toxicity, irritation, or allergic response, and achieving the intended therapeutic outcome".


More specifically the term "prodrug", as used herein, relates to an inactive or significantly less active derivative of a compound such as represented by the structural formulae herein described, which undergoes spontaneous or enzymatic transformation within the body in order to release the pharmacologically active form of the compound. For a comprehensive review, reference is made to Rautio J. et al.

("Prodrugs: design and clinical applications" Nature Reviews Drug Discovery, 2008, doi: 10.1038/nrd2468).

Representative benzofuran derivatives of the invention can be synthesized in accordance with the general synthetic methods described below and illustrated in the schemes that follow. Since the schemes are an illustration, the invention should not be construed as being limited by the specific chemical reaction and specific conditions described in the schemes and examples. The various starting material used in the schemes are commercially available or may be prepared by methods well within the skill persons versed in the art. The variables are as defined herein and within the skill of persons versed in the art.

Preferred embodiments of the invention are summarized as Clauses 1 to 51 hereinafter:

1. A compound of formula (I), a stereo-isomeric form, a physiologically acceptable salt, solvate and/or polymorph thereof.



preferably, the compound of formula (I), a stereo-isomeric form, a physiologically acceptable salt, solvate and/or polymorph thereof, for use in the treatment of pain;

wherein

R^1 represents $-F$, $-Cl$, $-Br$, $-I$, $-CN$, $-R^W$, $-OR^W$, $-OC(=O)R^W$, $-NR^WR^X$, $-NR^WC(=O)R^X$, $-SR^W$, $-S(=O)R^W$, $-S(=O)_2R^W$, $-C(=O)R^W$, $-C(=O)OR^W$, or $-C(=O)NR^WR^X$;

Q represents $-OR^2$ or $-NR^3R^4$;

R^2 represents $-R^Y$;

R^3 represents $-OH$ or $-R^Y$;

R^4 represents $-R^Y$ or $-S(=O)_2R^Y$;

or R^3 and R^4 together form a 4, 5, 6, 7 or 8 membered heterocycle containing 1 to 3 heteroatoms selected from N, O and S, saturated or unsaturated, unsubstituted or mono- or polysubstituted;

T represents $-O-$ and U represents $-CR^5R^{5i}-$; or T represents $-CR^5R^{5i}-$ and U represents $-O-$;

R^5 and R^{5i} independently of one another represent $-R^Y$;

R^6 , R^7 and R^8 independently of one another represent $-F$, $-Cl$, $-Br$, $-I$, $-CN$, $-NO_2$, $-SF_5$, $-R^W$, $-OR^W$, $-OC(=O)R^W$, $-NR^WR^X$, $-NR^WC(=O)R^X$, $-SR^W$, $-S(=O)R^W$, $-S(=O)_2R^W$, $-C(=O)R^W$, $-C(=O)OR^W$, or $-C(=O)NR^WR^X$;

V represents 3-14-membered heterocycloalkyl, saturated or unsaturated; or 5-14-membered heteroaryl; in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from $-F$, $-Cl$, $-Br$, $-I$, $-CN$, $-NO$, $-NO_2$, $=O$, $=S$, $-SF_5$, $-R^Y$, $-OR^Y$, $-OC(=O)R^Y$, $-NR^YR^Z$, $-NR^YC(=O)R^Z$, $-SR^Y$, $-S(=O)R^Y$, $-S(=O)_2R^Y$, $-C(=O)R^Y$, $-C(=O)OR^Y$, or $-C(=O)NR^YR^Z$;

wherein

R^W and R^X independently of one another in each case independently represent

$-H$;

$-C_1-C_6$ -alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;

$-C_1-C_6$ -heteroalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;

3-14-membered cycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered cycloalkyl is optionally connected through $-C_1-C_6$ -alkylene- or $-C_1-C_6$ -heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted; or

3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;

wherein said 3-14-membered heterocycloalkyl is optionally connected through $-C_1-C_6$ -alkylene- or $-C_1-C_6$ -heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted;

R^Y and R^Z independently of one another in each case independently represent

$-H$;

$-C_1-C_6$ -alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;

$-C_1-C_6$ -heteroalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;

3-14-membered cycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered cycloalkyl is optionally connected through $-C_1-C_6$ -alkylene- or $-C_1-C_6$ -heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted;

3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;

wherein said 3-14-membered heterocycloalkyl is optionally connected through $-C_1-C_6$ -alkylene- or $-C_1-C_6$ -heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted;

6-14-membered aryl, unsubstituted, mono- or polysubstituted; wherein said 6-14-membered aryl is optionally connected through $-C_1-C_6$ -alkylene- or $-C_1-C_6$ -heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted; or

5-14-membered heteroaryl, unsubstituted, mono- or polysubstituted; wherein said 5-14-membered heteroaryl is optionally connected through $-C_1-C_6$ -alkylene- or $-C_1-C_6$ -heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted;

or R^Y and R^Z together form a 4, 5, 6, 7 or 8 membered heterocycle containing 1 to 3 heteroatoms selected from N, O and S, saturated or unsaturated, unsubstituted or mono- or polysubstituted;

and wherein "mono- or polysubstituted" in each case independently means substituted with one or more substituents independently of one another selected from $-F$, $-Cl$, $-Br$, $-I$, $-CN$, $-C_1-C_6$ -alkyl, $-CF_3$, $-CF_2H$, $-CFH_2$, $-CF_2Cl$, $-CFCl_2$, $-C_1-C_6$ -alkylene- CF_3 , $-C_1-C_6$ -alkylene- CF_2H , $-C_1-C_6$ -alkylene- CFH_2 , $-C_1-C_6$ -alkylene- $O-CF_3$, $-C_1-C_6$ -alkylene- $O-CF_2H$, $-C_1-C_6$ -alkylene- $O-CFH_2$, $-C_1-C_6$ -alkylene- $NH-C_1-C_6$ -alkylene- CF_3 ,

91

$-C_{1-6}$ -alkylene- $N(C_{1-6}$ -alkyl)- C_{1-6} -alkylene-
 CF_3 , $-C(=O)-C_{1-6}$ -alkyl, $-C_{1-6}$ -alkylene- C
 $(=O)-C_{1-6}$ -alkyl, $-C(=O)OH$, $-C_{1-6}$ -al-
 $kylene-C(=O)-OH$, $-C(=O)-OC_{1-6}$ -alkyl,
 $-C_{1-6}$ -alkylene- $C(=O)-OC_{1-6}$ -alkyl, 5
 $-C(=O)O-C_{1-6}$ -alkylene- CF_3 , $-C(=O)-$
 NH_2 , $-C_{1-6}$ -alkylene- $C(=O)-NH_2$,
 $-C(=O)-NH(C_{1-6}$ -alkyl), $-C_{1-6}$ -alkylene- C
 $(=O)-NH(C_{1-6}$ -alkyl), $-C(=O)-N(C_{1-6}$ -al-
 $kyl)_2$, $-C_{1-6}$ -alkylene- $C(=O)-N(C_{1-6}$ -alkyl)₂, 10
 $-C(=O)-NH(OH)$, $-C_{1-6}$ -alkylene- $C(=O)-$
 $NH(OH)$, $-OH$, $-C_{1-6}$ -alkylene- OH , $=O$,
 $-OCF_3$, $-OCF_2H$, $-OCFH_2$, $-OCF_2Cl$,
 $-OCFCl_2$, $-O-C_{1-6}$ -alkyl, $-C_{1-6}$ -alkylene-
 $O-C_{1-6}$ -alkyl, $-O-C_{1-6}$ -alkylene- $O-C_{1-6}$ -al- 15
 kyl , $-O-C_{1-6}$ -alkylene- NH_2 , $-O-C_{1-6}$ -al-
 $kylene-NH-C_{1-6}$ -alkyl, $-O-C_{1-6}$ -alkylene- N
 $(C_{1-6}$ -alkyl)₂, $-O-C(=O)-C_{1-6}$ -alkyl, $-C_{1-6}$ -
 $alkylene-O-C(=O)-C_{1-6}$ -alkyl, $-O-C$
 $(=O)-O-C_{1-6}$ -alkyl, $-C_{1-6}$ -alkylene- $O-C$ 20
 $(=O)-O-C_{1-6}$ -alkyl, $-O-C(=O)-NH(C_{1-6}$ -
 $alkyl)$, $-C_{1-6}$ -alkylene- $O-C(=O)-NH(C_{1-6}$ -
 $alkyl)$, $-O-C(=O)-N(C_{1-6}$ -alkyl)₂, $-C_{1-6}$ -al-
 $kylene-O-C(=O)-N(C_{1-6}$ -alkyl)₂, $-O-S$
 $(=O)_2-NH_2$, $-C_{1-6}$ -alkylene- $O-S(=O)_2-$ 25
 NH_2 , $-O-S(=O)_2-NH(C_{1-6}$ -alkyl), $-C_{1-6}$ -
 $alkylene-O-S(=O)_2-NH(C_{1-6}$ -alkyl), $-O-S$
 $(=O)_2-N(C_{1-6}$ -alkyl)₂, $-C_{1-6}$ -alkylene- $O-S$
 $(=O)_2-N(C_{1-6}$ -alkyl)₂, $-NH_2$, $-NO$, $-NO_2$,
 $-C_{1-6}$ -alkylene- NH_2 , $-NH(C_{1-6}$ -alkyl), $-N(3-$ 30
 14 -membered cycloalkyl) $(C_{1-6}$ -alkyl), $-N(C_{1-6}$ -
 $alkyl)-C_{1-6}$ -alkylene- OH , $-N(H)-C_{1-6}$ -al-
 $kylene-OH$, $-C_{1-6}$ -alkylene- $NH(C_{1-6}$ -alkyl),
 $-N(C_{1-6}$ -alkyl)₂, $-C_{1-6}$ -alkylene- $N(C_{1-6}$ -
 $alkyl)_2$, $-NH-C(=O)-C_{1-6}$ -alkyl, $-C_{1-6}$ -al- 35
 $kylene-NH-C(=O)-C_{1-6}$ -alkyl, $-NH-C$
 $(=O)-O-C_{1-6}$ -alkyl, $-C_{1-6}$ -alkylene- $NH-C$
 $(=O)-O-C_{1-6}$ -alkyl, $-NH-C(=O)-NH_2$,
 $-C_{1-6}$ -alkylene- $NH-C(=O)-NH_2$, $-NH-C$
 $(=O)-NH(C_{1-6}$ -alkyl), $-C_{1-6}$ -alkylene- $NH-C$ 40
 $(=O)-NH(C_{1-6}$ -alkyl), $-NH-C(=O)-N$
 $(C_{1-6}$ -alkyl)₂, $-C_{1-6}$ -alkylene- $NH-C(=O)-N$
 $(C_{1-6}$ -alkyl)₂, $-N(C_{1-6}$ -alkyl)- $C(=O)-C_{1-6}$ -
 $alkyl$, $-C_{1-6}$ -alkylene- $N(C_{1-6}$ -alkyl)- $C(=O)-$ 45
 C_{1-6} -alkyl, $-N(C_{1-6}$ -alkyl)- $C(=O)-O-C_{1-6}$ -
 $alkyl$, $-C_{1-6}$ -alkylene- $N(C_{1-6}$ -alkyl)- $C(=O)-$
 $O-C_{1-6}$ -alkyl, $-N(C_{1-6}$ -alkyl)- $C(=O)-NH_2$,
 $-C_{1-6}$ -alkylene- $N(C_{1-6}$ -alkyl)- $C(=O)-NH_2$,
 $-N(C_{1-6}$ -alkyl)- $C(=O)-NH(C_{1-6}$ -alkyl),
 $-C_{1-6}$ -alkylene- $N(C_{1-6}$ -alkyl)- $C(=O)-NH$ 50
 $(C_{1-6}$ -alkyl), $-N(C_{1-6}$ -alkyl)- $C(=O)-N(C_{1-6}$ -
 $alkyl)_2$, $-C_{1-6}$ -alkylene- $N(C_{1-6}$ -alkyl)- $C(=O)-$
 $N(C_{1-6}$ -alkyl)₂, $-NH-S(=O)_2OH$, $-C_{1-6}$ -
 $alkylene-NH-S(=O)_2OH$, $-NH-S(=O)_2-$ 55
 C_{1-6} -alkyl, $-C_{1-6}$ -alkylene- $NH-S(=O)_2-C_{1-6}$ -
 $alkyl$, $-NH-S(=O)_2-O-C_{1-6}$ -alkyl, $-C_{1-6}$ -
 $alkylene-NH-S(=O)_2-O-C_{1-6}$ -alkyl,
 $-NH-S(=O)_2-NH_2$, $-NH-S(=O)_2-NH(C_{1-6}$ -al-
 $kyl)$, $-C_{1-6}$ -alkylene- $NH-S(=O)_2-NH(C_{1-6}$ - 60
 $alkyl)$, $-NH-S(=O)_2N(C_{1-6}$ -alkyl)₂, $-C_{1-6}$ -
 $alkylene-NH-S(=O)_2N(C_{1-6}$ -alkyl)₂, $-N(C_{1-6}$ -
 $alkyl)-S(=O)_2-OH$, $-C_{1-6}$ -alkylene- $N(C_{1-6}$ -
 $alkyl)-S(=O)_2-OH$, $-N(C_{1-6}$ -alkyl)- $S(=O)_2-$ 65
 C_{1-6} -alkyl, $-C_{1-6}$ -alkylene- $N(C_{1-6}$ -alkyl)- S
 $(=O)_2-C_{1-6}$ -alkyl, $-N(C_{1-6}$ -alkyl)- $S(=O)_2-$
 $O-C_{1-6}$ -alkyl, $-C_{1-6}$ -alkylene- $N(C_{1-6}$ -alkyl)- S

92

$(=O)_2-O-C_{1-6}$ -alkyl, $-N(C_{1-6}$ -alkyl)- S
 $(=O)_2-NH_2$, $-C_{1-6}$ -alkylene- $N(C_{1-6}$ -alkyl)- S
 $(=O)_2-NH_2$, $-N(C_{1-6}$ -alkyl)- $S(=O)_2-NH$
 $(C_{1-6}$ -alkyl), $-C_{1-6}$ -alkylene- $N(C_{1-6}$ -alkyl)- S
 $(=O)_2-NH(C_{1-6}$ -alkyl), $-N(C_{1-6}$ -alkyl)- S
 $(=O)_2-N(C_{1-6}$ -alkyl)₂, $-C_{1-6}$ -alkylene- $N(C_{1-6}$ -
 $alkyl)-S(=O)_2-N(C_{1-6}$ -alkyl)₂, $-SH$, $=S$,
 $-SF_5$, $-SCF_3$, $-SCF_2H$, $-SCFH_2$, $-S-C_{1-6}$ -
 $alkyl$, $-C_{1-6}$ -alkylene- $S-C_{1-6}$ -alkyl,
 $-S(=O)-C_{1-6}$ -alkyl, $-C_{1-6}$ -alkylene- S
 $(=O)-C_{1-6}$ -alkyl, $-S(=O)_2-C_{1-6}$ -alkyl,
 $-C_{1-6}$ -alkylene- $S(=O)_2-C_{1-6}$ -alkyl, $-S(=O)_2$
 $-OH$, $-C_{1-6}$ -alkylene- $S(=O)_2-OH$,
 $-S(=O)_2-O-C_{1-6}$ -alkyl, $-C_{1-6}$ -alkylene- S
 $(=O)_2-O-C_{1-6}$ -alkyl, $-S(=O)_2-NH_2$,
 $-C_{1-6}$ -alkylene- $S(=O)_2-NH_2$, $-S(=O)_2-$
 $NH(C_{1-6}$ -alkyl), $-C_{1-6}$ -alkylene- $S(=O)_2-NH$
 $(C_{1-6}$ -alkyl), $-S(=O)_2-N(C_{1-6}$ -alkyl)₂, $-C_{1-6}$ -
 $alkylene-S(=O)_2-N(C_{1-6}$ -alkyl)₂, 3-14-
 $membered$ cycloalkyl, $-C_{1-6}$ -alkylene-(3-14-
 $membered$ cycloalkyl), 3 to 14-membered
 $heterocycloalkyl$, $-C_{1-6}$ -alkylene-(3 to 14-membered
 $heterocycloalkyl)$, -phenyl, $-C_{1-6}$ -al-
 $kylene-phenyl$, 5 to 14-membered heteroaryl,
 $-C_{1-6}$ -alkylene-(5 to 14-membered heteroaryl),
 $-O-(3-14-membered$ cycloalkyl), $-O-(3$ to
 $14-membered$ heterocycloalkyl), $-O-phenyl$,
 $-O-(5$ to 14-membered heteroaryl), $-C(=O)-$
 $(3-14-membered$ cycloalkyl), $-C(=O)-(3$ to
 $14-membered$ heterocycloalkyl), $-C(=O)-phe-$
 nyl , $-C(=O)-(5$ to 14-membered heteroaryl),
 $-S(=O)_2-(3-14-membered$ cycloalkyl),
 $-S(=O)_2-(3$ to 14-membered heterocycloalkyl),
 $-S(=O)_2-phenyl$, $-S(=O)_2-(5$ to 14-mem-
 $bered$ heteroaryl).
2. The compound per se, or for use according to Clause 1,
wherein T represents $-O-$ and U represents
 $-CR^5R^{5'}$ —
3. The compound per se, or for use according to Clause 1,
wherein T represents $-CR^5R^{5'}$ — and U represents
 $-O-$.
4. The compound per se, or for use according to any one
of Clauses 1 to 3, wherein Q represents $-NR^3R^4$.
5. The compound per se, or for use according to any one
of Clauses 1 to 3, wherein Q represents $-OR^2$.
6. The compound per se, or for use according to any one
of Clauses 1 to 5, wherein V represents 5-14-membered
heteroaryl, unsubstituted, mono- or polysubstituted
with substituents independently of one another selected
from $-F$, $-Cl$, $-Br$, $-I$, $-CN$, $-NO$, $-NO_2$, $=O$,
 $=S$, $-SF_5$, $-R^Y$, $-OR^Y$, $-OC(=O)R^Y$, $-NR^YR^Z$,
 $-NR^YC(=O)R^Z$, $-SR^Y$, $-S(=O)R^Y$, $-S(=O)_2R^Y$,
 $-C(=O)R^Y$, $-C(=O)OR^Y$, or $-C(=O)NR^YR^Z$.
7. The compound per se, or for use according to Clause 6,
wherein the 5-14-membered heteroaryl is not benzo-
furan.
8. The compound per se, or for use according to Clause 6
or 7, wherein the 5-14-membered heteroaryl is selected
from benzimidazole, benzisoxazole, benzoxazole, ben-
zodioxole, benzofuran, benzothiadiazole, benzothiaz-
ole, benzothiophene, carbazole, cinnoline, dibenzo-
furan, furane, furazane, imidazole, imidazopyridine,
indazole, indole, indolizine, isobenzofuran, isoindole,
isoquinoline, isothiazole, isoxazole, naphthyrindine,
oxadiazole, oxazole, oxindole, phthalazine, purine,
pyrazine, pyrazole, pyridazine, pyridine, pyrimidine,
pyrrole, quinazoline, quinoline, quinoxaline, tetrazole,
thiadiazole, thiazole, thiophene, triazine, triazole, and

- [1,2,4]triazolo[4,3-a]pyrimidine; in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from —F, —Cl, —Br, —I, —CN, —NO, —NO₂, =O, =S, —SF₅, —R^Y, —OR^Y, —OC(=O)R^Y, —NR^YR^Z, —NR^YC(=O)R^Z, —SR^Y, —S(=O)R^Y, —S(=O)₂R^Y, —C(=O)R^Y, —C(=O)OR^Y, or —C(=O)NR^YR^Z. 5
9. The compound per se, or for use according to any one of Clauses 6 to 8, wherein the 5-14-membered heteroaryl is selected from the group consisting of furane, thiophene, imidazole, pyrazole, oxazole, isoxazole, thiazole, triazole, pyridine, isoquinoline, benzothiazole, pyridazine, pyrimidine, imidazopyridine; in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from —F, —Cl, —Br, —I, —CN, —NO, —NO₂, =O, =S, —SF₅, —R^Y, —OR^Y, —OC(=O)R^Y, —NR^YR^Z, —NR^YC(=O)R^Z, —SR^Y, —S(=O)R^Y, —S(=O)₂R^Y, —C(=O)R^Y, —C(=O)OR^Y, or —C(=O)NR^YR^Z. 10
10. The compound per se, or for use according to any one of Clauses 6 to 9, wherein the 5-14-membered heteroaryl is selected from the group consisting of furan-2-yl, furan-3-yl, thiophen-2-yl, thiophen-3-yl, pyrazol-5-yl, oxazol-5-yl, isoxazol-4-yl, thiazol-2-yl, thiazol-5-yl, 1,2,4-triazol-3-yl, 1,2,3-triazol-4-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, isoquinolin-1-yl, isoquinolin-5-yl, benzod[thiazol-2-yl, pyridazin-3-yl, pyrimidin-5-yl, and imidazo[1,2-a]pyridin-6-yl; in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from —F, —Cl, —Br, —I, —CN, —NO, —NO₂, =O, =S, —SF₅, —R^Y, —OR^Y, —OC(=O)R^Y, —NR^YR^Z, —NR^YC(=O)R^Z, —SR^Y, —S(=O)R^Y, —S(=O)₂R^Y, —C(=O)R^Y, —C(=O)OR^Y, or —C(=O)NR^YR^Z. 15
11. The compound per se, or for use according to any one of Clauses 1 to 5, wherein V represents 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted with substituents independently of one another selected from —F, —Cl, —Br, —I, —CN, —NO, —NO₂, =O, =S, —SF₅, —R^Y, —OR^Y, —OC(=O)R^Y, —NR^YR^Z, —NR^YC(=O)R^Z, —SR^Y, —S(=O)R^Y, —S(=O)₂R^Y, —C(=O)R^Y, —C(=O)OR^Y, or —C(=O)NR^YR^Z. 20
12. The compound per se, or for use according to Clause 11, wherein the 3-14-membered heterocycloalkyl is selected from azepane, 1,4-oxazepane, azetane, azetidine, aziridine, azocane, diazepane, dioxane, dioxolane, dithiane, dithiolane, imidazolidine, isothiazolidine, isoxalidine, morpholine, oxazolidine, oxepane, oxetane, oxirane, piperazine, piperidine, pyrazolidine, pyrrolidine, quinuclidine, tetrahydrofuran, tetrahydropyran, tetrahydrothiopyran, thiazolidine, thietane, thirane, thiolane, thiomorpholine, indoline, dihydrobenzofuran, dihydrobenzothiophene, 1,1-dioxothiacyclohexane, 2-azaspiro[3.3]heptane, 2-oxaspiro[3.3]heptane, 7-azaspiro[3.5]nonane, 8-azabicyclo[3.2.1]octane, 9-azabicyclo[3.3.1]nonane, hexahydro-1H-pyrrolizine, hexahydrocyclopenta[c]pyrrole, octahydrocyclopenta[c]pyrrole, and octahydropyrrolo[1,2-a]pyrazine; in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from —F, —Cl, —Br, —I, —CN, —NO, —NO₂, =O, =S, —SF₅, —R^Y, —OR^Y, —OC(=O)R^Y, —NR^YR^Z, —NR^YC(=O)R^Z, —SR^Y, —S(=O)R^Y, —S(=O)₂R^Y, —C(=O)R^Y, —C(=O)OR^Y, or —C(=O)NR^YR^Z. 25
13. The compound per se, or for use according to Clause 11 or 12, wherein the 3-14-membered heterocycloalkyl is tetrahydropyran or pyrrolidine; in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from —F, —Cl, —Br, —I, —CN, —NO, —NO₂, =O, =S, —SF₅, —R^Y, —OR^Y, —OC(=O)R^Y, —NR^YR^Z, —NR^YC(=O)R^Z, —SR^Y, —S(=O)R^Y, —S(=O)₂R^Y, —C(=O)R^Y, —C(=O)OR^Y, or —C(=O)NR^YR^Z. 30
14. The compound per se, or for use according to any one of Clauses 11 to 13, wherein the 3-14-membered heterocycloalkyl is tetrahydropyran-4-yl or pyrrolidin-3-yl; in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from —F, —Cl, —Br, —I, —CN, —NO, —NO₂, =O, =S, —SF₅, —R^Y, —OR^Y, —OC(=O)R^Y, —NR^YR^Z, —NR^YC(=O)R^Z, —SR^Y, —S(=O)R^Y, —S(=O)₂R^Y, —C(=O)R^Y, —C(=O)OR^Y, or —C(=O)NR^YR^Z. 35
15. The compound per se, or for use according to any one of the preceding Clauses, wherein V is unsubstituted, mono- or polysubstituted with substituents independently of one another selected from —F, —Cl, —Br, —I, —CN, —C(=O)OH, —NH₂, —NO₂, —OH, =O, —SF₅; —C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; —C(=O)O—C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; —NHC₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; —N(C₁₋₆-alkyl)₂, saturated or unsaturated, unsubstituted, mono- or polysubstituted; —O—C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; —S(=O)₂—C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; 3-14-membered cycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered cycloalkyl is optionally connected through —C₁-C₆-alkylene- or —C₁-C₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted; or 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered heterocycloalkyl is optionally connected through —C₁-C₆-alkylene- or —C₁-C₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted. 40
16. The compound per se, or for use according to any one of the preceding Clauses, wherein V is unsubstituted, mono- or polysubstituted with substituents independently of one another selected from —OH, —F, —Cl, —Br, —I, —SH, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, —CN, —NO₂, —C(=O)OH, —NH₂, —N(CH₃)₂, —cyclopropyl, or —O-cyclopropyl; preferably selected from —OH, —F, —Cl, —Br, —I, —SH, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, —SF₅, —CN, —NO₂, —C(=O)OH, —NH₂, or —N(CH₃)₂; —C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted with substituents independently of one another, selected from the group consisting of —F, —Cl, —Br, —I, —C₁₋₆-alkyl, C₂₋₆-alkenyl, —C₂₋₆-alkynyl, —OH, =O, —SH, =S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, 45

13. The compound per se, or for use according to Clause 11 or 12, wherein the 3-14-membered heterocycloalkyl is tetrahydropyran or pyrrolidine; in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from —F, —Cl, —Br, —I, —CN, —NO, —NO₂, =O, =S, —SF₅, —R^Y, —OR^Y, —OC(=O)R^Y, —NR^YR^Z, —NR^YC(=O)R^Z, —SR^Y, —S(=O)R^Y, —S(=O)₂R^Y, —C(=O)R^Y, —C(=O)OR^Y, or —C(=O)NR^YR^Z. 5
14. The compound per se, or for use according to any one of Clauses 11 to 13, wherein the 3-14-membered heterocycloalkyl is tetrahydropyran-4-yl or pyrrolidin-3-yl; in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from —F, —Cl, —Br, —I, —CN, —NO, —NO₂, =O, =S, —SF₅, —R^Y, —OR^Y, —OC(=O)R^Y, —NR^YR^Z, —NR^YC(=O)R^Z, —SR^Y, —S(=O)R^Y, —S(=O)₂R^Y, —C(=O)R^Y, —C(=O)OR^Y, or —C(=O)NR^YR^Z. 10
15. The compound per se, or for use according to any one of the preceding Clauses, wherein V is unsubstituted, mono- or polysubstituted with substituents independently of one another selected from —F, —Cl, —Br, —I, —CN, —C(=O)OH, —NH₂, —NO₂, —OH, =O, —SF₅; —C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; —C(=O)O—C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; —NHC₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; —N(C₁₋₆-alkyl)₂, saturated or unsaturated, unsubstituted, mono- or polysubstituted; —O—C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; —S(=O)₂—C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; 3-14-membered cycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered cycloalkyl is optionally connected through —C₁-C₆-alkylene- or —C₁-C₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted; or 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered heterocycloalkyl is optionally connected through —C₁-C₆-alkylene- or —C₁-C₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted. 15
16. The compound per se, or for use according to any one of the preceding Clauses, wherein V is unsubstituted, mono- or polysubstituted with substituents independently of one another selected from —OH, —F, —Cl, —Br, —I, —SH, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, —CN, —NO₂, —C(=O)OH, —NH₂, —N(CH₃)₂, —cyclopropyl, or —O-cyclopropyl; preferably selected from —OH, —F, —Cl, —Br, —I, —SH, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, —SF₅, —CN, —NO₂, —C(=O)OH, —NH₂, or —N(CH₃)₂; —C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted with substituents independently of one another, selected from the group consisting of —F, —Cl, —Br, —I, —C₁₋₆-alkyl, C₂₋₆-alkenyl, —C₂₋₆-alkynyl, —OH, =O, —SH, =S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, 20

95

—OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, C(=O)CHF₂, —C(=O)NH₂, and -cyclopropyl; preferably selected from the group consisting of —F, —Cl, —Br, —I, —C₁₋₆-alkyl, C₂₋₆-alkenyl, —C₂₋₆-alkynyl, —OH, —O, —SH, —S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, C(=O)CHF₂, and —C(=O)NH₂; 5

—C₁₋₆-heteroalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted with substituents independently of one another, selected from the group consisting of —F, —Cl, —Br, —I, —C₁₋₆-alkyl, C₂₋₆-alkenyl, —C₂₋₆-alkynyl, —OH, —O, —SH, —S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, C(=O)CHF₂, and —C(=O)NH₂; 10

—OC₁₋₆-alkyl, unsubstituted, mono- or polysubstituted with substituents independently of one another, selected from the group consisting of —F, —Cl, —Br, —I, —C₁₋₆-alkyl, C₂₋₆-alkenyl, —C₂₋₆-alkynyl, —OH, —O, —SH, —S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, C(=O)CHF₂, and —C(=O)NH₂; 15

—O(C=O)C₁₋₆-alkyl, unsubstituted, mono- or polysubstituted with substituents independently of one another, selected from the group consisting of —F, —Cl, —Br, —I, —C₁₋₆-alkyl, C₂₋₆-alkenyl, —C₂₋₆-alkynyl, —OH, —O, —SH, —S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, C(=O)CHF₂, and —C(=O)NH₂; 20

—C(=O)OC₁₋₆-alkyl, unsubstituted, mono- or polysubstituted with substituents independently of one another, selected from the group consisting of —F, —Cl, —Br, —I, —C₁₋₆-alkyl, C₂₋₆-alkenyl, —C₂₋₆-alkynyl, —OH, —O, —SH, —S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, C(=O)CHF₂, and —C(=O)NH₂; 25

3-14-membered cycloalkyl selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl; in each case unsubstituted, mono- or polysubstituted with substituents independently of one another, selected from the group consisting of —F, —Cl, —Br, —I, —C₁₋₆-alkyl, C₂₋₆-alkenyl, —C₂₋₆-alkynyl, —OH, —O, —SH, —S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, C(=O)CHF₂, and —C(=O)NH₂; 30

3-14-membered heterocycloalkyl selected from the group consisting of azepane, 1,4-oxazepane, azetane, azetidine, aziridine, azocane, diazepane, dioxane, dioxolane, dithiane, dithiolane, imidazolidine, isothiazolidine, isoxalidine, morpholine, oxazolidine, oxepane, oxetane, oxirane, piperazine, piperidine, pyrazolidine, pyrrolidine, quinuclidine, tetrahydrofuran, tetrahydropyran, tetrahydrothiopyran, thiazolidine, thietane, thiirane, thiolane, thiomorpholine, indoline, dihydrobenzofuran, dihydrobenzothio- 35

phene, 1,1-dioxothiacyclohexane, 2-azaspiro[3.3]heptane, 2-oxaspiro[3.3]heptane, 7-azaspiro[3.5]nonane, 8-azabicyclo[3.2.1]octane, 9-azabicyclo[3.3.1]nonane, hexahydro-1H-pyrrolizine, hexahydrocyclopenta[c]pyrrole, octahydrocyclopenta[c]pyrrole, and octahydropyrrolo[1,2-a]pyrazin, in each case unsubstituted, mono- or polysubstituted with substituents independently of one another, selected from the group consisting of —F, —Cl, —Br, —I, —C₁₋₆-alkyl, C₂₋₆-alkenyl, —C₂₋₆-alkynyl, —OH, —O, —SH, —S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, C(=O)CHF₂, and —C(=O)NH₂. 96

96

ta[c]pyrrole, and octahydropyrrolo[1,2-a]pyrazin, in each case unsubstituted, mono- or polysubstituted with substituents independently of one another, selected from the group consisting of —F, —Cl, —Br, —I, —C₁₋₆-alkyl, C₂₋₆-alkenyl, —C₂₋₆-alkynyl, —OH, —O, —SH, —S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, C(=O)CHF₂, and —C(=O)NH₂. 96

17. The compound per se, or for use according to any one of the preceding Clauses, wherein V is unsubstituted, mono- or polysubstituted with substituents independently of one another selected from —F, —Cl, —CN, —OH, —O, —C₁₋₆-alkyl, —CHF₂, —CF₃, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NHC(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-NHC(=O)O—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkyl, —N(C₁₋₆-alkyl)₂, —OC₁₋₆-alkyl, —OCF₃, —O—C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —S(=O)₂—C₁₋₆-alkyl, -azetidine, —C₁₋₆-alkylene-O-tetrahydropyran, or -piperazine substituted with —C₁₋₆-alkyl; or represents oxetanyl, unsubstituted, mono- or polysubstituted. 17

18. The compound per se, or for use according to any one of the preceding Clauses, wherein V is

- unsubstituted;
- monosubstituted;
- disubstituted;
- trisubstituted; or
- tetrasubstituted.

19. The compound per se, or for use according to any one of the preceding Clauses, wherein R¹ represents —H, —F, —Cl, —Br, —I; —C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; —O—C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; —C(=O)C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; —C(=O)OC₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; —C(=O)NHC₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; —C(=O)N(C₁₋₆-alkyl)₂, saturated or unsaturated, unsubstituted, mono- or polysubstituted; —S(=O)C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; —S(=O)₂—C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; —C₁₋₆-heteroalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; or 3-14-membered cycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered cycloalkyl is optionally connected through —C₁₋₆-alkylene- or —C₁₋₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted. 18

20. The compound per se, or for use according to any one of the preceding Clauses, wherein R¹ represents —H, —F, —Cl, —Br, —I, —C₁₋₆-alkyl, —O—C₁₋₆-alkyl, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH(C₁₋₆-alkyl), —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —CF₃, —CF₂H, —CFH₂, —CF₂Cl, —CFCl₂, —C₁₋₆-alkylene-CF₃, —C₁₋₆-alkylene-CF₂H, —C₁₋₆-alkylene-CFH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)—C₁₋₆-alkylene-CF₃, —C(=O)C₁₋₆-alkyl, —C(=O)OC₁₋₆-alkyl, —C(=O)NHC₁₋₆-alkyl, —C(=O)N(C₁₋₆-alkyl)₂, —S(=O)— 19

- C₁₋₆-alkyl, —S(=O)₂—C₁₋₆-alkyl, —O—C₁₋₆-alkyl, -cyclopropyl unsubstituted, cyclobutyl unsubstituted, cyclopentyl unsubstituted or cyclohexyl unsubstituted.
21. The compound per se, or for use according to any one of the preceding Clauses, wherein R¹ represents —H, —C₁₋₆-alkyl, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —CH₂F, —CHF₂, —CF₃, -cyclopentyl, unsubstituted, or -cyclopropyl, unsubstituted; preferably wherein R¹ represents —H, —C₁₋₆-alkyl, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —CH₂F, —CHF₂, —CF₃, or -cyclopentyl, unsubstituted.
22. The compound per se, or for use according to any one of the preceding Clauses, wherein R¹ represents —CH₂F, —CHF₂, —CH₃, or -cyclopropyl, unsubstituted; preferably wherein R¹ represents —CH₂F, —CHF₂, or —CH₃.
23. The compound per se, or for use according to any one of the preceding Clauses, wherein R² represents —H; —C₁-C₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; —C₁-C₆-heteroalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; 3-14-membered cycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered cycloalkyl is optionally connected through —C₁-C₆-alkylene- or —C₁-C₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted; or 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered heterocycloalkyl is optionally connected through —C₁-C₆-alkylene- or —C₁-C₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted.
24. The compound per se, or for use according to any one of the preceding Clauses, wherein R² represents —H, —C₁₋₆-alkyl, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH(C₁₋₆-alkyl), —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —CF₃, —CF₂H, —CFH₂, —CF₂Cl, —CFCl₂, —C₁₋₆-alkylene-CF₃, —C₁₋₆-alkylene-CF₂H, —C₁₋₆-alkylene-CFH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, or —C₁₋₆-alkylene-N(C₁₋₆-alkyl)-C₁₋₆-alkylene-CF₃.
25. The compound per se, or for use according to any one of the preceding Clauses, wherein R² represents —H or —C₁₋₆-alkyl.
26. The compound per se, or for use according to any one of the preceding Clauses, wherein R³ represents —H; —OH; —C₁-C₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; or —C₁-C₆-heteroalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted.
27. The compound per se, or for use according to any one of the preceding Clauses, wherein R³ represents —H, —OH, —C₁₋₆-alkyl, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NH(C₁₋₆-alkyl), —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —CF₃, —CF₂H, —CFH₂, —CF₂Cl, —CFCl₂, —C₁₋₆-alkylene-CF₃, —C₁₋₆-alkylene-CF₂H, —C₁₋₆-alkylene-CFH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, or —C₁₋₆-alkylene-N(C₁₋₆-alkyl)-C₁₋₆-alkylene-CF₃.

28. The compound per se, or for use according to any one of the preceding Clauses, wherein R³ represents —H, —OH, or —C₁₋₆-alkyl, saturated, unsubstituted or monosubstituted with —OH.
29. The compound per se, or for use according to any one of the preceding Clauses, wherein R⁴ represents —H; —S(=O)C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; —S(=O)₂—C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; —C₁-C₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; —C₁-C₆-heteroalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; 3-14-membered cycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered cycloalkyl is optionally connected through —C₁-C₆-alkylene- or —C₁-C₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted; 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered heterocycloalkyl is optionally connected through —C₁-C₆-alkylene- or —C₁-C₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted; 6-14-membered aryl, unsubstituted, mono- or polysubstituted; wherein said 6-14-membered aryl is optionally connected through —C₁-C₆-alkylene- or —C₁-C₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted; or 5-14-membered heteroaryl, unsubstituted, mono- or polysubstituted; wherein said 5-14-membered heteroaryl is optionally connected through —C₁-C₆-alkylene- or —C₁-C₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted.
30. The compound per se, or for use according to any one of the preceding Clauses, wherein R⁴ represents —S(=O)₂C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted with substituents independently of one another selected from the group consisting of —F, —Cl, —C₁₋₆-alkyl, —C₁₋₆-alkylene-CF₃, —OH, =O, —OC₁₋₆-alkyl, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —NH₂, —NHC₁₋₆-alkyl, —N(C₁₋₆-alkyl)₂, —NHC(=O)O—C₁₋₆-alkyl, —N(C₁₋₆-alkyl)C(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NHC(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkyl, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, —C(=O)—C₁₋₆-alkyl, —C(=O)OH, —C(=O)O—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkylene-CF₃, —C(=O)NH₂, —C(=O)NH(C₁₋₆-alkyl), —C(=O)N(C₁₋₆-alkyl)₂, —S(=O)₂C₁₋₆-alkyl, -phenyl, —C₁₋₆-alkylene-phenyl, 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted; and 5-14-membered heteroaryl, unsubstituted; —S(=O)₂(3-14-membered cycloalkyl), wherein said 3-14-membered cycloalkyl is selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted with substituents independently of

one another selected from the group consisting of —F, —Cl, —C₁₋₆-alkyl, —C₁₋₆-alkylene-CF₃, —OH, =O, —OC₁₋₆-alkyl, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —NH₂, —NHC(=O)-alkyl, —N(C₁₋₆-alkyl)₂, —NHC(=O)O—C₁₋₆-alkyl, —N(C₁₋₆-alkyl)C(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NHC(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkyl, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, —C(=O)—C₁₋₆-alkyl, —C(=O)OH, —C(=O)O—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkylene-CF₃, —C(=O)NH₂, —C(=O)NH(C₁₋₆-alkyl), —C(=O)N(C₁₋₆-alkyl)₂, —S(=O)₂C₁₋₆-alkyl, -phenyl, —C₁₋₆-alkylene-phenyl, 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted; and 5-14-membered heteroaryl, unsubstituted;

—C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted with substituents independently of one another selected from the group consisting of —F, —Cl, —C₁₋₆-alkyl, —C₁₋₆-alkylene-CF₃, —OH, =O, —OC₁₋₆-alkyl, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —NH₂, —NHC₁₋₆-alkyl, —N(C₁₋₆-alkyl)₂, —NHC(=O)O—C₁₋₆-alkyl, —N(C₁₋₆-alkyl)C(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NHC(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkyl, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, —C(=O)—C₁₋₆-alkyl, —C(=O)OH, —C(=O)O—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkylene-CF₃, —C(=O)NH₂, —C(=O)NH(C₁₋₆-alkyl), —C(=O)N(C₁₋₆-alkyl)₂, —S(=O)₂C₁₋₆-alkyl, -phenyl, —C₁₋₆-alkylene-phenyl, 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted; and 5-14-membered heteroaryl, unsubstituted;

3-14-membered cycloalkyl or —C₁₋₆-alkylene-(3-14-membered cycloalkyl), wherein —C₁₋₆-alkylene- is unsubstituted or monosubstituted with —OH, wherein said 3-14-membered cycloalkyl is selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl, in each case saturated or unsaturated, in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from the group consisting of —F, —Cl, —C₁₋₆-alkyl, —C₁₋₆-alkylene-CF₃, —OH, =O, —OC₁₋₆-alkyl, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —NH₂, —NHC₁₋₆-alkyl, —N(C₁₋₆-alkyl)₂, —NHC(=O)O—C₁₋₆-alkyl, —N(C₁₋₆-alkyl)C(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NHC(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkyl, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, —C(=O)—C₁₋₆-alkyl, —C(=O)OH, —C(=O)O—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkylene-CF₃, —C(=O)NH₂, —C(=O)NH(C₁₋₆-alkyl), —C(=O)N(C₁₋₆-alkyl)₂, —S(=O)₂C₁₋₆-alkyl, -phenyl, —C₁₋₆-alkylene-phenyl, 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted; and 5-14-membered heteroaryl, unsubstituted;

3-14-membered heterocycloalkyl or —C₁₋₆-alkylene-(3-14-membered heterocycloalkyl), wherein —C₁₋₆-alkylene- is unsubstituted or monosubstituted with —OH, wherein said 3-14-membered heterocycloalkyl in each case is selected from the group consisting of azepane, 1,4-oxazepane, azetane, azetidine, aziri-

dine, azocane, diazepane, dioxane, dioxolane, dithiane, dithiolane, imidazolidine, isothiazolidine, isoxalidine, morpholine, oxazolidine, oxepane, oxetane, oxirane, piperazine, piperidine, pyrazolidine, pyrrolidine, quinuclidine, tetrahydrofuran, tetrahydropyran, tetrahydrothiopyran, thiazolidine, thietane, thiirane, thiolane, thiomorpholine, indoline, dihydrobenzofuran, dihydrobenzothiophene, 1,1-dioxothiacyclohexane, 2-azaspiro[3.3]heptane, 2-oxaspiro[3.3]heptane, 7-azaspiro[3.5]nonane, 8-azabicyclo[3.2.1]octane, 9-azabicyclo[3.3.1]nonane, hexahydro-1H-pyrrolizine, hexahydrocyclopenta[c]pyrrole, octahydrocyclopenta[c]pyrrole, and octahydropyrrolo[1,2-a]pyrazine; in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from the group consisting of —F, —Cl, —C₁₋₆-alkyl, —C₁₋₆-alkylene-CF₃, —OH, =O, —OC₁₋₆-alkyl, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —NH₂, —NHC₁₋₆-alkyl, —N(C₁₋₆-alkyl)₂, —NHC(=O)O—C₁₋₆-alkyl, —N(C₁₋₆-alkyl)C(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NHC(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkyl, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, —C(=O)—C₁₋₆-alkyl, —C(=O)OH, —C(=O)O—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkylene-CF₃, —C(=O)NH₂, —C(=O)NH(C₁₋₆-alkyl), —C(=O)N(C₁₋₆-alkyl)₂, —S(=O)₂C₁₋₆-alkyl, -phenyl, —C₁₋₆-alkylene-phenyl, 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted; and 5-14-membered heteroaryl, unsubstituted;

—phenyl unsubstituted, mono- or polysubstituted with substituents independently of one another selected from the group consisting of —F, —Cl, —CN, —C₁₋₆-alkyl, —C₁₋₆-alkylene-CF₃, —OH, =O, —OC₁₋₆-alkyl, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —NH₂, —NHC₁₋₆-alkyl, —N(C₁₋₆-alkyl)₂, —NHC(=O)O—C₁₋₆-alkyl, —N(C₁₋₆-alkyl)C(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NHC(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkyl, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, —C(=O)—C₁₋₆-alkyl, —C(=O)OH, —C(=O)O—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkylene-CF₃, —C(=O)NH₂, —C(=O)NH(C₁₋₆-alkyl), —C(=O)N(C₁₋₆-alkyl)₂, —S(=O)₂C₁₋₆-alkyl, -phenyl, —C₁₋₆-alkylene-phenyl, 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted; and 5-14-membered heteroaryl, unsubstituted;

5-14-membered heteroaryl or —C₁₋₆-alkylene-(5-14-membered heteroaryl), wherein —C₁₋₆-alkylene- is unsubstituted or monosubstituted with —OH, wherein said 5-14-membered heteroaryl in each case is selected from the group consisting of benzimidazole, benzisoxazole, benzoazole, benzodioxole, benzofuran, benzothiadiazole, benzothiazole, benzothiophene, carbazole, cinnoline, dibenzofuran, furane, furazane, imidazole, imidazopyridine, indazole, indole, indolizine, isobenzofuran, isoindole, isoquinoline, isothiazole, isoxazole, naphthyridine, oxadiazole, oxazole, oxindole, phthalazine, purine, pyrazine, pyrazole, pyridazine, pyridine, pyrimidine, pyrrole, quinazoline, quinoxaline, tetrazole, thiadiazole, thiazole, thiophene, triazine, triazole, and [1,2,4]triazolo[4,3-a]pyrimidine; in each

101

- case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from the group consisting of —F, —Cl, —CN, —C₁₋₆-alkyl, —C₁₋₆-alkylene-CF₃, —OH, =O, —OC₁₋₆-alkyl, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —NH₂, —NHC₁₋₆-alkyl, —N(C₁₋₆-alkyl)₂, —NHC(=O)O—C₁₋₆-alkyl, —N(C₁₋₆-alkyl)C(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NHC(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkyl, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, —C(=O)—C₁₋₆-alkyl, —C(=O)OH, —C(=O)O—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkylene-CF₃, —C(=O)NH₂, —C(=O)NH(C₁₋₆-alkyl), —C(=O)N(C₁₋₆-alkyl)₂, —S(=O)₂C₁₋₆-alkyl, —phenyl, —C₁₋₆-alkylene-phenyl, 3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted; and 5-14-membered heteroaryl, unsubstituted.
31. The compound per se, or for use according to any one of the preceding Clauses, wherein R⁴ represents —H; —S(=O)₂C₁₋₆-alkyl, saturated, unsubstituted, monosubstituted or polysubstituted with —F; —S(=O)₂(3-14-membered cycloalkyl), saturated, unsubstituted; —C₁₋₆-alkyl, saturated, unsubstituted, monosubstituted or disubstituted with substituents independently of one another selected from the group consisting of —OH, —OC₁₋₆-alkyl, —N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkyl, —phenyl unsubstituted; 3-14-membered cycloalkyl or —C₁₋₆-alkylene-(3-14-membered cycloalkyl), wherein —C₁₋₆-alkylene- is unsubstituted or monosubstituted with —OH, wherein said 3-14-membered cycloalkyl is saturated, unsubstituted, monosubstituted or disubstituted with substituents independently of one another selected from the group consisting of —C₁₋₆-alkyl, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-NHC(=O)O—C₁₋₆-alkyl, —OH, —OC₁₋₆-alkyl, —NH₂, —N(C₁₋₆-alkyl)₂, —NHC(=O)O—C₁₋₆-alkyl; 3-14-membered heterocycloalkyl or —C₁₋₆-alkylene-(3-14-membered heterocycloalkyl), wherein —C₁₋₆-alkylene- is unsubstituted or monosubstituted with —OH, wherein said 3-14-membered heterocycloalkyl in each case is selected from azetane, 1,4-oxazepane, pyrrolidine, piperidine, azepane, diazepane, tetrahydrofuran, tetrahydropyran, oxetane, morpholine, piperazine, hexahydrocyclopenta[c]pyrrole, octahydrocyclopenta[c]pyrrole, octahydropyrrolo[1,2-a]pyrazine, 8-azabicyclo[3.2.1]octane, 9-azabicyclo[3.3.1]nonane, quinuclidine, hexahydro-1H-pyrrolizine, 2-oxaspiro[3.3]heptane, 2-azaspiro[3.3]heptane, 7-azaspiro[3.5]nonane, 1,1-dioxothiacyclohexane, in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from the group consisting of —F, —OH, =O, —C₁₋₆-alkyl, —C₁₋₆-alkylene-CF₃, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —NH₂, —N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —C(=O)—C₁₋₆-alkyl, —C(=O)OH, —C(=O)O—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alk-

102


- ylene-CF₃, —C(=O)NH₂, —C(=O)NH(C₁₋₆-alkyl), —S(=O)₂C₁₋₆-alkyl, oxetanyl, pyrimidinyl, —C₁₋₆-alkylene-phenyl; —phenyl unsubstituted; 5-14-membered heteroaryl or —C₁₋₆-alkylene-(5-14-membered heteroaryl), wherein —C₁₋₆-alkylene- is unsubstituted or monosubstituted with —OH, wherein said 5-14-membered heteroaryl in each case is selected from the group consisting of pyridine, pyridazine, pyrazine, pyrazole, isoxazole, triazole, and [1,2,4]triazolo[4,3-a]pyrimidine, in each case unsubstituted, monosubstituted or disubstituted with substituents independently of one another selected from the group consisting of —C₁₋₆-alkyl, —OH.
32. The compound per se, or for use according to any one of the preceding Clauses, wherein R³ and R⁴ together form a 5- or 6-membered heterocycle containing 1 or 2 heteroatoms selected from N, O and S, saturated or unsaturated, unsubstituted or mono- or poly substituted.
33. The compound per se, or for use according to any one of the preceding Clauses, wherein R³ and R⁴ together form a heterocycle selected from the group consisting of pyrrolidine, piperidine, morpholine, and piperazine, in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from the group consisting of —C₁₋₆-alkyl, —NH₂, —NHCH₃, —N(CH₃)₂, —C(=O)NH—C₁₋₆-alkyl, —C(=O)N(C₁₋₆-alkyl)₂, —C(=O)O—C₁₋₆-alkyl, —NHC(=O)O—C₁₋₆-alkyl, —pyridyl unsubstituted, and 1,2,4-oxadiazole unsubstituted or monosubstituted with —C₁₋₆-alkyl.
34. The compound per se, or for use according to any one of the preceding Clauses, wherein R³ and R⁴ together form a pyrrolidine ring, unsubstituted or monosubstituted with —N(CH₃)₂; piperidine ring, unsubstituted or monosubstituted with a substituent selected from the group consisting of —C₁₋₆-alkyl, —NH₂, —N(CH₃)₂, —C(=O)NH—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkyl, —NHC(=O)O—C₁₋₆-alkyl, and 1,2,4-oxadiazole unsubstituted or monosubstituted with —C₁₋₆-alkyl; morpholine ring, unsubstituted; or piperazine ring, unsubstituted or N-substituted with a substituent selected from the group consisting of —C₁₋₆-alkyl and —pyridyl unsubstituted.
35. The compound per se, or for use according to any one of the preceding Clauses, wherein R⁵ and R^{5'} independently of one another represent —H; —C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; —C₁₋₆-heteroalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; 3-14-membered cycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered cycloalkyl is optionally connected through —C₁-C₆-alkylene- or —C₁-C₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted.
36. The compound per se, or for use according to any one of the preceding Clauses, wherein R⁵ and R^{5'} independently of one another represent —H, —C₁₋₆-alkyl, or —C₁-C₆-alkylene-N(C₁-C₆-alkyl)₂.

103

37. The compound per se, or for use according to any one of the preceding Clauses, wherein at least one of R⁵ and R⁵ⁱ does not represent —H.
38. The compound per se, or for use according to any one of the preceding Clauses, wherein R⁶, R⁷ and R⁸ independently of one another represent
- H;
 - F, —Cl, —Br, —I, —OH, —SH, —SF₅, —CN, —NO₂, —C(=O)OH, —NH₂;
 - C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
 - O—C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
 - NHC₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
 - N(C₁₋₆-alkyl)₂, saturated or unsaturated, unsubstituted, mono- or poly substituted;
 - C(=O)OC₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
 - OC(=O)C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;
 - C₁₋₆-heteroalkyl, saturated or unsaturated, unsubstituted, mono- or poly substituted.
39. The compound per se, or for use according to any one of the preceding Clauses, wherein R⁶, R⁷ and R⁸ independently of one another represent
- H, —F, —Cl, —Br, —I, —OH, —SH, —SF₅, —CN, —NO₂, —C(=O)OH, —NH₂,
 - C₁₋₆-alkyl, —CF₃, —CHF₂, —CH₂F,
 - O—C₁₋₆-alkyl, —OCF₃, —OCHF₂, —OCH₂F,
 - NHC₁₋₆-alkyl unsubstituted or substituted with one or more substituents independently of one another selected from —OH, —O, —F, —Cl, —Br, —I, —SH, —S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, and —C(=O)NH₂;
 - N(C₁₋₆-alkyl)₂ unsubstituted or substituted with one or more substituents independently of one another selected from —OH, —O, —F, —Cl, —Br, —I, —SH, —S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, and —C(=O)NH₂;
 - C(=O)OC₁₋₆-alkyl unsubstituted or substituted with one or more substituents independently of one another selected from —OH, —O, —F, —Cl, —Br, —I, —SH, —S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, and —C(=O)NH₂;
 - OC(=O)C₁₋₆-alkyl unsubstituted or substituted with one or more substituents independently of one another selected from —OH, —O, —F, —Cl, —Br, —I, —SH, —S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, and —C(=O)NH₂; or
 - C₁₋₆-heteroalkyl unsubstituted or substituted with one or more substituents independently of one another selected from —OH, —O, —F, —Cl, —Br, —I, —SH, —S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, and —C(=O)NH₂.
40. The compound per se, or for use according to any one of the preceding Clauses, wherein R⁶ represents —H, —F, —Cl, —CN, or —C₁₋₆-alkyl.
41. The compound per se, or for use according to any one of the preceding Clauses, wherein R⁶ does not represent —H.


104

42. The compound per se, or for use according to any one of the preceding Clauses, wherein R⁷ represents —H, —F, —Cl, —CN, or —C₁₋₆-alkyl.
43. The compound per se, or for use according to any one of the preceding Clauses, wherein R⁷ does not represent —H.
44. The compound per se, or for use according to any one of the preceding Clauses, wherein R⁵ represents —H, —F, —Cl, —CN, or —C₁₋₆-alkyl.
45. The compound per se, or for use according to any one of the preceding Clauses, wherein R⁸ does not represent —H.
46. The compound per se, or for use according to any one of the preceding Clauses, wherein
- (i) R⁶, R⁷ and R⁸ each represent —H; or
 - (ii) two of R⁶, R⁷ and R⁸ represent —H and the other of R⁶, R⁷ and R⁸ represents —F, —Cl, —CN, or —CH₃; or
 - (iii) one of R⁶, R⁷ and R⁸ represents —H and the other of R⁶, R⁷ and R⁸ independently of one another represent —F, —Cl, —CN, or —CH₃.
47. The compound per se, or for use according to any one of the preceding Clauses, which is selected from the group consisting of cpd 001 to cpd 207 as mentioned above and the physiologically acceptable salts thereof.
48. The compound per se, or for use according to any one of the preceding Clauses, wherein the pain is selected from nociceptive pain, inflammatory pain, and neuropathic pain.
49. The compound per se, or for use according to any one of the preceding Clauses, wherein the pain is post-operative pain.
50. A compound of formula (I), a stereo-isomeric form, a physiologically acceptable salt, solvate and/or polymorph thereof, as defined in any one of the preceding Clauses, wherein
- (a) Q represents —OR²; and
 - (a-1) R¹ represents —CH₂F, —CHF₂, or —CF₃; and/or
 - (a-2) at least one of R⁵ and R⁵ⁱ does not represent —H; and/or
 - (a-3) R⁶ does not represent —H; and/or
 - (a-4) R⁵ does not represent —H;
 - or
 - (b) Q represents —NR³R⁴; and
 - (b-1) with the proviso that the following compounds and their salts are excluded



105

-continued



and/or

(b-2) R^1 represents $-\text{CH}_2\text{F}$, $-\text{CHF}_2$, or $-\text{CF}_3$, $-\text{CN}$, -propyl, or -cyclopropyl; and/or(b-3) at least one of R^5 and R^{51} does not represent $-\text{H}$; and/or(b-4) R^3 represents $-\text{H}$.

51. A pharmaceutical composition or a medicament comprising a compound according to any one of the preceding Clauses.

EXAMPLES

The following examples are provided for the purpose of illustrating the invention and by no means should be interpreted to limit the scope of the invention.

Representative compounds of the present invention can be synthesized in accordance with the general synthetic methods described below and illustrated in the schemes that follow. Since the schemes are an illustration, the invention should not be construed as being limited by the specific chemical reaction and specific conditions described in the schemes and examples. The various starting material used in the schemes are commercially available or may be prepared by methods well within the skill of persons versed in the art. The variables are as defined herein and within the skill of persons versed in the art.

Abbreviations used in the instant specification, particularly in the schemes and examples, are as follows: ACN—acetonitrile, AcOH—Acetic acid, ADDP—1,1'-(Azodicarbonyl)dipiperidine, aq.—Aqueous, AIBN—Azobisisobutyronitrile, CAN—Ceric ammonium nitrate, COMU—(1-Cyano-2-ethoxy-2-oxoethylidenaminooxy)dimethylamino-morpholino-carbenium, hexafluorophosphate, DABCO—1,4-diazabicyclo[2.2.2]octane, DAST—Diethylaminosulfur trifluoride, DBU—1,8-Diazabicyclo[5.4.0]undec-7-ene, DCC—N,N'-dicyclohexylcarbodiimide, DCM—Dichloromethane, DEAD—Diethyl azodicarboxylate, DIA—Diastereomer, DIAD—Diisopropyl azodicarboxylate, DEA—Diethylamine, DIPEA—Diisopropyl-ethylamine, DME—1,2-Dimethoxyethane, DMF—N,N-Dimethylformamide, DMSO—Dimethylsulfoxide, DTBAD—tert-Butylazodicarboxylate, EDCI or EDC—1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide, En—Enantiomer, Et₂O—


106

Diethyl ether, EtOH—Ethanol, EtOAc—Ethyl acetate, Eq.—Equivalent, FA—Formic acid, FCC—Flash column chromatography, h—Hour, HATU—O-(7-Azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate, HPLC—High performance liquid chromatography, IPA—
 5 isopropyl alcohol, LAH—Lithiumaluminumhydride, LG—Leaving group, MeOH—methanol, MgSO₄—Magnesium sulfate, min.—Minute, Na₂SO₄—Sodium sulfate, NBS—N-Bromosuccinimide, NMP—1-Methyl-2-pyrrolidinone, Pd(PPh₃)₄—Tetrakis-(triphenylphosphine)-palladium(0), Pd₂(dba)₃—Tris(dibenzylideneacetone)dipalladium, Pet
 10 ether—Petroleum ether, PPh₃—Triphenylphosphine, PS-DIEA—Diisopropyl-ethyl amine supported on Polystyrene, PS—PPh₃—Triphenylphosphine supported on Polystyrene, PyBop—Benzotriazol-1-yl-oxytripyrrolidinophosphonium hexafluorophosphate, PTSA—p-Toluenesulfonic acid, RF: ratio of frontiers, RM—Reaction mixture, RP—
 15 Reverse phase, RT—Room temperature, sat.—Saturated, SEM—[2-(Trimethylsilyl)ethoxy]methyl acetal, SFC—Supercritical fluid chromatography, SPE—Solid Phase Extraction, TBDMS—Tert-Butyldimethylsilyl ether, TBAF—Tetrabutylammonium fluoride hydrate, TBAI—Tetrabutylammonium iodide, TEA—Triethylamine, THF—
 20 Tetrahydrofuran, TFA—Trifluoroacetic acid, TLC—thin layer chromatography, TPP—triphenyl phosphine, IPA—
 25 isopropyl alcohol, TMS—trimethyl silyl, T3P—Propylphosphonic anhydride,


30 The compounds of interest having a structure according to the general formula (A) and all other formulas described herein and embodiments thereof can be prepared as outlined in the general chemical scheme 1.

35

Scheme 1: all R₁, R₂, R₃, R₄ and R₅ are as described for the compounds of the present invention.



107




para-Benzoquinone of formula 1, may be condensed with a ketoester of formula 2 (commercially available or synthesized by procedures known to those skilled in the art), wherein R_2 is an ester protecting group (e.g., methyl, ethyl, t-Bu and the like), in the presence of a Lewis acid (e.g., Titanium(IV) chloride, zinc(II) chloride and the like) in a polar solvent (e.g., DCM, MeOH, EtOH, and the like) at a temperature ranging from 0 to 100° C. to provide intermediates of formula 4. More detailed information can be found in the following references (*Bioorg. Med. Chem.* 2012, 20, 4237-4244 and FR 1319594). Alternatively, para-benzoquinone of formula 1 may be reacted with an enamine of formula 3 (commercially available or synthesized by procedures known to those skilled in the art), in the presence of

108

a protic acid (e.g., trifluoroacetic acid, para-toluenesulfonic acid, and the like) in a polar solvent (e.g., DCM, MeOH, EtOH, and the like) at a temperature ranging from 0 to 100° C. to provide intermediates of formula 4. More detailed information can be found in the following reference (*J. Heterocyclic Chem.* 2006, 43, 873). Intermediates of formula 4 may then be converted into the desired compounds of formula 7 via nucleophilic substitution using intermediates of formula 5 (commercially available or synthesized), wherein LG is a leaving group, in the presence of a base (e.g., DIPEA, DBU, triethylamine, Cs_2CO_3 , and the like) in a polar solvent (e.g., acetonitrile, DMF, NMP, and the like), with or without a chelating agent (e.g., 18-crown-6, cis-anti-cis-dicyclohexano-18-crown-6, and the like) at a temperature ranging from 0 to 100° C. Alternatively, intermediates of formula 4 may also be reacted with intermediates of formula 6 (commercially available or synthesized) in the presence of an azodicarboxylate reagent (e.g., DEAD, DIAD, ADDP, and the like) and a phosphine (e.g., tributylphosphine, triphenylphosphine and the like) in a solvent (e.g., THF, toluene, and the like) at a temperature ranging from 0 to 100° C., to provide the desired compounds of formula 7. Ester derivatives 7 may then be converted into the desired compounds of formula 8 via standard saponification reactions. The desired compounds of formula 10 may be obtained from acid derivatives of formula 8 by reaction with amine derivatives of formula 9 (commercially available or synthesized by procedures known in the art or as set forth in the examples below) under standard peptide coupling conditions (e.g., DCC, EDCI, HATU, PyBop and the like) in a polar aprotic solvent (e.g., DCM, DMF and the like). Alternatively, carboxylic acid derivatives of formula 8, may be converted into acid chloride derivatives by procedures known to those skilled in the art or as set forth in the examples below, and then reacted with amines of formula 9 to obtain the desired compounds of formula 10 by procedures known to those skilled in the art or as set forth in the examples below.

In a more particular embodiment, the compounds of the present invention may be synthesized as depicted in scheme 2.

Scheme 2: all R_1 , R_2 , R_3 , R_4 and R_5 are as described for the compounds of the present invention.






109

5-Hydroxy-benzofuran-3-carboxylic acid derivatives 11 (commercially available or synthesized by procedures known in the art or as set forth in the examples below) may be reacted with amine derivatives of formula 9 (commercially available or synthesized by procedures known in the art or as set forth in the examples below) under standard peptide coupling conditions (e.g. DCC, EDCI, HATU, PyBop and the like) in a polar aprotic solvent (e.g. DCM, DMF and the like) to provide intermediates of formula 12. Alternatively, compounds of formula 13 (synthesized as described in scheme 1) may be converted into intermediates of formula 12 via a hydrogenation reaction with a reducing agent (e.g., hydrogen gas, ammonium formate, cyclohexadiene and the like) using a catalyst (more preferably Pd or Pt) in a solvent (e.g., THF, EtOH, and the like). Intermediates of formula 12 may then be converted into the desired compounds of formula 10 via nucleophilic substitution using intermediates of formula 5 (commercially available or synthesized), wherein LG is a leaving group, in the presence of a base (e.g., DIPEA, DBU, triethylamine, Cs_2CO_3 , and the like) in a polar solvent (e.g., acetonitrile, DMF, NMP, and the like), with or without a chelating agent (e.g., 18-crown-6, cis-anti-cis-dicyclohexano-18-crown-6, and the like) at a temperature ranging from 0 to 100° C. Alternatively, compounds of formula 12 may also be reacted with intermediates of formula 6 (commercially available or synthesized) in the presence of an azodicarboxylate reagent (e.g., DEAD, ADDP, DIAD, tert-butylazodicarboxylate, and the like) and a phosphine (e.g., tributylphosphine, triphenylphosphine and the like) in a solvent (e.g., THF, toluene, and the like) at a temperature ranging from 0 to 100° C., to provide the desired compounds of formula 10.

In a more particular embodiment, the compounds of the present invention may be synthesized as depicted in scheme 3.







Scheme 3: all R1, R2, R3, R4, R5, R7 and R8 are as described for the compounds of the present invention.



16

110


-continued



22

111

-continued




A substituted para-benzoquinone derivatives of formula 14 may be condensed with a ketoester of formula 2 (commercially available or synthesized by procedures known to those skilled in the art), wherein R_2 is an ester protecting group (e.g. methyl, ethyl, t-Bu and the like), in the presence of a Lewis acid (e.g., Titanium(IV) chloride, zinc(II) chloride and the like) in a polar solvent (e.g., DCM, MeOH, EtOH, and the like) at a temperature ranging from 0 to 100° C. to provide a mixture of substituted intermediates of formula 15 and 16. More detailed information can be found in the following references (*Bioorg. Med. Chem.* 2012, 20,

112

4237-4244 and FR 1319594). Intermediates of formula 15 and/or 16 may then be converted into the desired compounds of formula 17 and/or 18 via nucleophilic substitution using intermediates of formula 5 (commercially available or synthesized), wherein LG is a leaving group, in the presence of a base (e.g., DIPEA, DBU, triethylamine, Cs_2CO_3 , and the like) in a polar solvent (e.g., acetonitrile, DMF, NMP, and the like), with or without a chelating agent (e.g., 18-crown-6, cis-anti-cis-dicyclohexano-18-crown-6, and the like) at a temperature ranging from 0 to 100° C. Alternatively, intermediates of formula 15 and/or 16 may also be reacted with intermediates of formula 6 (commercially available or synthesized) in the presence of an azodicarboxylate reagent (e.g., DEAD, DIAD, ADDP, and the like) and a phosphine (e.g., tributylphosphine, triphenylphosphine and the like) in a solvent (e.g., THF, toluene, and the like) at a temperature ranging from 0 to 100° C., to provide the desired compounds of formula 17 and/or 18. Ester derivatives 17 and/or 18 may then be converted into the desired carboxylic acid of formula 19 and/or 20 via standard saponification reactions. The desired compounds of formula 21 and/or 22 may be obtained from acid derivatives of formula 19 and/or 20 by reaction with amine derivatives of formula 9 (commercially available or synthesized by procedures known in the art or as set forth in the examples below) under standard peptide coupling conditions (e.g., DCC, EDCI, HATU, PyBop and the like) in a polar aprotic solvent (e.g., DCM, DMF and the like). A mixture of compounds 21 and 22 may be separated (e.g., silica gel, HPLC, SFC or preparative CFC) to provide the desired compounds of formula 21 or 22.


In a more particular embodiment, the compounds of the present invention may be synthesized as depicted in scheme 4.

Scheme 4: all R_1 , R_2 , R_3 , R_4 , R_5 , R_7 and R_8 are as described for the compounds of the present invention.



113


114



Intermediates of formula 4 may be halogenated with a suitable halogenating agent (e.g., Bromine, N-bromosuccinimide and the like) in a solvent (e.g., chloroform, water and the like) to provide the desired intermediates 23. The desired compounds of formula 26 may be obtained by an Ullmann type reaction with CuCN followed by a Mitsunobu type reaction with intermediates of formula 6 (commercially available or synthesized). Alternatively, the desired compounds of formula 26 may be obtained via a Mitsunobu type reaction with intermediates of formula 6 (commercially available or synthesized) followed by a Suzuki reaction. Ester derivatives 26 may then be converted into the desired compounds of formula 27 via standard saponification reactions. The desired compounds of formula 28 may be obtained from acid derivatives of formula 27 by reaction with amine derivatives of formula 9 (commercially available or synthesized by procedures known in the art or as set forth in the examples below) under standard peptide coupling conditions (e.g., DCC, EDCI, HATU, PyBop and the like) in a polar aprotic solvent (e.g., DCM, DMF and the like).


In a more particular embodiment, the compounds of the present invention may be synthesized as depicted in scheme 5.

Scheme 5: all R_1 , R_2 , R_3 and are as described for the compounds of the present invention.





115

-continued

**116**

-continued




Intermediates of formula 4 may be halogenated with a suitable halogenating agent (e.g. select fluor and the like) in a solvent (e.g. chloroform, acetonitrile and the like) to provide the desired intermediates 29 and 30. Intermediates of formula 29 and/or 30 may be reacted with intermediates of formula 6 (commercially available or synthesized) in the presence of an azodicarboxylate reagent (e.g., DEAD, DIAD, ADDP, and the like) and a phosphine (e.g., tributylphosphine, triphenylphosphine and the like) in a solvent (e.g., THF, toluene, and the like) at a temperature ranging from 0 to 100° C., to provide the desired compounds of formula 31 and/or 32. Ester derivatives 31 and/or 32 may then be converted into the corresponding carboxylic acid of formula 33 and/or 34 via standard saponification reactions. The desired compounds of formula 35 and/or 36 may be obtained from acid derivatives of formula 33 and/or 34 by reaction with amine derivatives of formula 9 (commercially available or synthesized by procedures known in the art or as set forth in the examples below) under standard peptide coupling conditions (e.g., DCC, EDCI, HATU, PyBop and the like) in a polar aprotic solvent (e.g., DCM, DMF and the like). A mixture of compounds 35 and 36 may be separated (e.g., silica gel, HPLC, SFC or preparative CFC) to provide the desired compounds of formula 35 or 36.


EXAMPLES

Structure/CODE


cpd 001




Structure/CODE




cpd 002




cpd 003




cpd 004




cpd 005




cpd 006



cpd 007




cpd 008




cpd 009

-continued


Structure/CODE




cpd 010




cpd 011




cpd 012




cpd 013




cpd 014



cpd 015



cpd 016




cpd 017

121


-continued

122


Structure/CODE




cpd 018




cpd 019




cpd 020



cpd 021




cpd 022




cpd 023


Structure/CODE




cpd 024




cpd 025




cpd 026



cpd 027




cpd 028




cpd 029


Structure/CODE




cpd 030




cpd 031




cpd 032



cpd 033




cpd 034




cpd 035


Structure/CODE




cpd 037




cpd 038



cpd 039




cpd 040




cpd 041


Structure/CODE




cpd 042




cpd 043



cpd 044








cpd 045



cpd 047

Structure/CODE



Structure/CODE








135


-continued

136


Structure/CODE




Structure/CODE	
----------------	--




cpd 064




cpd 065




cpd 066



cpd 067




cpd 068




cpd 069

-continued


Structure/CODE




cpd 070




cpd 071



cpd 072




cpd 073




cpd 074

Structure/CODE




cpd 075




cpd 076

cpd 077



cpd 078




cpd 079

143


-continued

144


Structure/CODE




cpd 080



cpd 081




cpd 082




cpd 083


Structure/CODE




cpd 084



cpd 085







cpd 086







cpd 087

-continued

Structure/CODE



Structure/CODE














151

-continued

152

Structure/CODE




Structure/CODE	
	cpd 101
	cpd 102
	cpd 103
	cpd 104
	cpd 105
	cpd 106

155

-continued


156

Structure/CODE




cpd 107


cpd 108




cpd 109




cpd 110




cpd 111




Structure/CODE




cpd 112




cpd 113



cpd 114




cpd 115




cpd 116

-continued


Structure/CODE




cpd 117




cpd 118




cpd 119



cpd 120




cpd 121




cpd 122

Structure/CODE


cpd 123




cpd 124




cpd 125



cpd 126









cpd 127



-continued

Structure/CODE









165

-continued

166

Structure/CODE




167


-continued

168


Structure/CODE




cpd 140




cpd 141




cpd 142



cpd 143



cpd 144




cpd 145


Structure/CODE




Structure/CODE




cpd 152




cpd 153



cpd 154









cpd 155



cpd 156

Structure/CODE




175


-continued

176


Structure/CODE




cpd 163



cpd 164



cpd 165




cpd 166

177


-continued

178


Structure/CODE




cpd 167




cpd 168



cpd 169




cpd 170




cpd 171

-continued


Structure/CODE




cpd 172




cpd 173



cpd 174



cpd 175




cpd 176

181

-continued

182

Structure/CODE



Structure/CODE




185


-continued

186


Structure/CODE




cpd 188




cpd 189



cpd 190




cpd 191




cpd 192

-continued


Structure/CODE




cpd 193




cpd 194



cpd 195



cpd 196




cpd 197

189

-continued


190

Structure/CODE




cpd 198


cpd 199




cpd 200



cpd 201



cpd 202




191


-continued

192


Structure/CODE




cpd 203




cpd 204




cpd 205



cpd 206











cpd 207




cpd 208

-continued


Structure/CODE




Structure/CODE




cpd 217




cpd 218




cpd 219




cpd 220




cpd 221



cpd 222




cpd 223




cpd 224

-continued


Structure/CODE




cpd 225




cpd 226




cpd 227




cpd 228




cpd 229



cpd 230




cpd 231




cpd 232

-continued


Structure/CODE	
----------------	--




cpd 233




cpd 234




cpd 235




cpd 236




cpd 237











cpd 238











cpd 239












cpd 240

Structure/CODE	
	cpd 241
	cpd 242
	cpd 243
	cpd 244
	cpd 245
	cpd 246
	cpd 247
	cpd 248

-continued

Structure/CODE	
	cpd 249
	cpd 250
	cpd 251
	cpd 252
	cpd 253
	cpd 254
	cpd 255
	cpd 256

-continued


Structure/CODE	
	cpd 257
	cpd 258
	cpd 259
	cpd 260
	cpd 261
	cpd 262
	cpd 263
	cpd 264
	cpd 265

Structure/CODE




-continued


Structure/CODE




cpd 273




cpd 274




cpd 275



cpd 276




cpd 277




cpd 278


Structure/CODE




cpd 279




cpd 280




cpd 281



cpd 282




cpd 283




cpd 284


Structure/CODE




cpd 285




cpd 286




cpd 287



cpd 288




cpd 289




cpd 290


Structure/CODE




cpd 291




cpd 292




cpd 293



cpd 294




cpd 295




cpd 296


Structure/CODE




cpd 297




cpd 298




cpd 299



cpd 300




cpd 301




cpd 302


Structure/CODE




cpd 303




cpd 304




cpd 305



cpd 306




cpd 307




cpd 308

Structure/CODE	
	cpd 309
	cpd 310
	cpd 311
	cpd 312
	cpd 313
	cpd 314


Structure/CODE




cpd 315




cpd 316




cpd 317



cpd 318



cpd 319




cpd 320

225


-continued

226


Structure/CODE




cpd 321




cpd 322



cpd 323









cpd 324




cpd 325


-continued

Structure/CODE	
	cpd 326
	cpd 327
	cpd 328
	cpd 329
	cpd 330
	cpd 331


Structure/CODE




cpd 332




cpd 333



cpd 334



cpd 335



cpd 336

Structure/CODE









Structure/CODE




-continued


Structure/CODE




Structure/CODE




cpd 357




cpd 358




cpd 359



cpd 360




cpd 361




cpd 362

-continued


Structure/CODE




cpd 363




cpd 364



cpd 365




cpd 366




cpd 367


Structure/CODE




cpd 368




cpd 369



cpd 370




cpd 371




cpd 372

-continued


Structure/CODE




cpd 373




cpd 374




cpd 375



cpd 376









cpd 377









cpd 378

-continued







Structure/CODE	
	cpd 379
	cpd 380
	cpd 381
	cpd 382
	cpd 383
	cpd 384

-continued

Structure/CODE




-continued


Structure/CODE	
	cpd 391
	cpd 392
	cpd 393
	cpd 394
	cpd 395
	cpd 396

-continued


Structure/CODE




cpd 397




cpd 398



cpd 399








cpd 400








cpd 401

Structure/CODE




-continued


Structure/CODE




Structure/CODE




cpd 412




cpd 413




cpd 414



cpd 415




cpd 416




cpd 417

-continued


Structure/CODE




cpd 418




cpd 419




cpd 420



cpd 421



cpd 422









cpd 423

261


-continued

262


Structure/CODE




Structure/CODE




Structure/CODE




cpd 436




cpd 437




cpd 438



cpd 439




cpd 440




cpd 441

-continued


Structure/CODE	
----------------	--




cpd 442




cpd 443




cpd 444



cpd 445









cpd 446




cpd 447


Structure/CODE




Structure/CODE




cpd 454




cpd 455




cpd 456



cpd 457




cpd 458




cpd 459


Structure/CODE




cpd 460




cpd 461




cpd 462



cpd 463









cpd 464




cpd 465

Structure/CODE




Structure/CODE


cpd 472




cpd 473




cpd 474




cpd 475




cpd 476




cpd 477




Structure/CODE




cpd 478




cpd 479




cpd 480



cpd 481



cpd 482




cpd 483

281


-continued

282


Structure/CODE




cpd 484




cpd 485



cpd 486









cpd 487









cpd 488


Structure/CODE




Structure/CODE




Structure/CODE




cpd 501




cpd 502




cpd 503



cpd 504









cpd 505




cpd 506


-continued

Structure/CODE	
	cpd 507
	cpd 508
	cpd 509
	cpd 510
	cpd 511
	cpd 512


Structure/CODE




cpd 513




cpd 514



cpd 515









cpd 516




cpd 517


Structure/CODE	
----------------	--




Structure/CODE




cpd 524




cpd 525




cpd 526



cpd 527



cpd 528




cpd 529

Structure/CODE




-continued


Structure/CODE




cpd 536




cpd 537



cpd 538




cpd 539




cpd 540


Structure/CODE




cpd 541




cpd 542



cpd 543








cpd 544








cpd 545






Structure/CODE




Structure/CODE




Structure/CODE




Structure/CODE




cpd 561




cpd 562



cpd 563




cpd 564




cpd 565


Structure/CODE




cpd 566




cpd 567



cpd 568








cpd 569



cpd 570

Structure/CODE




315


-continued

316


Structure/CODE




cpd 576




cpd 577




cpd 578



cpd 579



cpd 580




cpd 581

Structure/CODE




-continued


Structure/CODE




cpd 588




cpd 589




cpd 590



cpd 591




cpd 592




cpd 593


Structure/CODE




cpd 594




cpd 595




cpd 596



cpd 597









cpd 598




cpd 599

-continued


Structure/CODE




Structure/CODE




cpd 606




cpd 607




cpd 608



cpd 609









cpd 610









cpd 611


Structure/CODE




Structure/CODE




Structure/CODE




cpd 624




cpd 625




cpd 626



cpd 627









cpd 628



cpd 629

Structure/CODE




335


-continued

336


Structure/CODE




cpd 636




cpd 637




cpd 638



cpd 639




cpd 640



cpd 641

Structure/CODE



Structure/CODE




341

-continued


342

Structure/CODE




cpd 656


cpd 657




cpd 658




cpd 659










cpd 660



Structure/CODE



-continued


Structure/CODE	
	cpd 667
	cpd 668
	cpd 669
	cpd 670
	cpd 671
	cpd 672
	cpd 673

347


-continued

348


Structure/CODE




cpd 674




cpd 675




cpd 676



cpd 677




cpd 678




cpd 679

-continued


Structure/CODE




cpd 680




cpd 681




cpd 682



cpd 683









cpd 684









cpd 685

Structure/CODE




-continued

Structure/CODE




Structure/CODE




cpd 698


cpd 699




cpd 700




cpd 701




cpd 702




cpd 703




Structure/CODE




cpd 704




cpd 705




cpd 706



cpd 707




cpd 708




cpd 709

-continued


Structure/CODE




cpd 710




cpd 711




cpd 712



cpd 713



cpd 714









cpd 715

361

-continued


362

Structure/CODE




-continued


Structure/CODE




cpd 722




cpd 723



cpd 724



cpd 725




cpd 726

365


-continued

366


Structure/CODE




cpd 727




cpd 728



cpd 729



cpd 730




cpd 731

367


368

-continued


Structure/CODE




cpd 732




cpd 733



cpd 734




cpd 735




cpd 736

-continued


Structure/CODE




cpd 737




cpd 738




cpd 739



cpd 740




cpd 741




cpd 742


Structure/CODE




cpd 743




cpd 744




cpd 745



cpd 746




cpd 747




cpd 748


Structure/CODE



cpd 749



cpd 750



cpd 751

The following examples are provided for the purpose of illustrating the present invention and by no means should be interpreted to limit the scope of the present invention.

Part A represent the preparation of the compounds whereas Part B represents the pharmacological examples.

Part A

All starting materials which are not explicitly described were either commercially available (the details of suppliers such as for example Acros, Avocado, Aldrich, Fluka, FluoroChem, MatrixScientific, Maybridge, Merck, Sigma, etc. can be found in the SciFinder® Database for example) or the synthesis thereof has already been described precisely in the specialist literature (experimental guidelines can be found in the Reaxys® Database or the SciFinder® Database respectively, for example) or can be prepared using the conventional methods known to the person skilled in the art.

The reactions were, if necessary, carried out under an inert atmosphere (mostly argon and N₂). The number of equivalents of reagents and the amounts of solvents employed as well as the reaction temperatures and times can vary slightly between different reactions carried out by analogous methods. The work-up and purification methods were adapted according to the characteristic properties of each compound and can vary slightly for analogous methods. The yields of the compounds prepared are not optimized.

The indication "equivalents" ("eq." or "eq" or "equiv.") means molar equivalents, "RT" or "rt" means room temperature T (23±7° C.), "M" are indications of concentration in mol/l, "sol." means solution, "conc." means concentrated. The mixing ratios of solvents are usually stated in the volume/volume ratio.

Key analytical characterization was carried out by means of ¹H-NMR spectroscopy and/or mass spectrometry (MS, m/z for [M+H]⁺ and/or [M-H]⁻) for all the exemplary compounds and selected intermediate products. In certain cases, where e.g., regioisomers and/or diastereomers could be/were formed during the reaction, additional analytics, such as, e.g., ¹³C NMR and NOE (nuclear overhauser effect) NMR experiments were in some cases performed.

Analytical instruments employed were e.g., for NMR analysis a BRUKER 400 MHz or a BRUKER 500 MHz machine (Software Topspin), alternatively a BRUKER AVANCE 300 MHz and 400 Mhz was employed. For LC/MS analysis e.g., an Agilent 1290 infinity, Mass: 6150 SQD (ESI/APCI) or an Agilent 1200 SERIES, Mass: 6130 SQD (ESI/APCI) (Software Chemistation) was employed. Analytical HPLCs were measured e.g., on Waters (Software Empower), an Agilent-1200-ELSD (Software Chemistation) or an Agilent-1260 (Software OpenLAB). Analytical SFC were performed e.g., on a PIC solution (Software: SFC PICLAB ONLINE), a WATERS-X5 (Software MASSLYNX) or a WATERS-UPC2 (Empower).

Preparative HPLC were performed e.g., on a Waters 2998 (Software Empower) or a YMC (Software K-Prep). Preparative SFC were performed e.g., on a Waters, SFC-200 (Software Chromscope or Super chrome), a Waters, SFC-80 (Super chrome) or a PIC,PIC-175 (Software S10-100).

Structures of example compounds that contain stereocentres are drawn and named with absolute stereochemistry, if known. In case of unknown absolute stereochemistry, the

375

compounds can be either racemic, a mixture of diastereomers, a pure diastereomer of unknown stereochemistry, or a pure enantiomer of unknown stereochemistry. Dia 1 and Dia 2 means that diastereoisomers were separated but the stereochemistry is unknown. En 1 and En 2 means that both enantiomers were separated but the absolute configuration is unknown. No suffix given after the compound code means that a compound containing stereocentres was obtained as a racemic mixture or a mixture of diastereomers, respectively, unless the chemical name of the compound specifies the exact stereochemistry.

The LC/MS analysis mentioned in the experimental part were also performed on a Dionex Ultimate 3000 HPLC system (equipped with a PDA detector) connected to a mass spectrometer Brucker Esquire 6000 (equipped with a multimode source, ESI/APCI) (Method L in the table below). Or the LC/MS analysis mentioned in the experimental part were performed on a Waters system combining an Acquity UPLC H-Class equipped with a Acquity UPLC PDA Detector and an Acquity TQ Detector (ESI) (Method U in the table below).

376

Conditions used for the HPLC analysis in experimental part. The LC/MS analysis mentioned in the experimental part were performed on a Dionex Ultimate 3000 HPLC system (equipped with a PDA detector) connected to a mass spectrometer Brucker Esquire 6000 (equipped with a multimode source, ESI/APCI). The separations were performed with a SunFireC18, 3.5 μ m 3.0 $\times$ 100 mm, column equipped with a SunFire C18, 3.5 μ m, 3.0 $\times$ 20 mm Guard column or a X-Bridge C18 100 $\times$ 3.0 mm column equipped with a X-Bridge C18, 3.5 μ m, 3.0 $\times$ 20 mm Guard column thermostated to 30° C. and the DAD acquisition wavelength was set in the range of 190-420 nm (Method L in the table below). Elutions were carried out with the methods described in the following tables. Solvent A: Formic Acid LC-MS grade 0.1% in milliQ water Solvent B: NH₄OAc (LC-MS grade) 10 mMol in milliQ water, adjusted at pH10 with an aq. solution of NH₃, LC-MS grade. Solvent C: Acetonitrile LC-MS grade.

LC/MS		Time (min)	Solvents			Flow	
Method	System		A (%)	B (%)	C (%)	(mL/min)	Column
L1	Dionex Ultimate 3000 HPLC	0	80	—	20	1	SunFire C18
		0.2	80	—	20	1	
		7	40	—	60	1	
		8	10	—	90	1	
		10.8	10	—	90	1	
		11	80	—	20	1	
		14	80	—	20	1	
L2	Dionex Ultimate 3000 HPLC	0	50	—	50	1	SunFire C18
		0.2	50	—	50	1	
		6	10	—	90	1	
		10.8	10	—	90	1	
		11	50	—	50	1	
		14	50	—	50	1	
L3	Dionex Ultimate 3000 HPLC	0	—	80	20	1	X-Bridge C18
		0.2	—	80	20	1	
		7	—	40	60	1	
		8	—	10	90	1	
		10.8	—	10	90	1	
		11	—	80	20	1	
		14	—	80	20	1	
L4	Dionex Ultimate 3000 HPLC	0	—	50	50	1	X-Bridge C18
		0.2	—	50	50	1	
		6	—	10	90	1	
		10.8	—	10	90	1	
		11	—	50	50	1	
		14	—	50	50	1	
L5	Dionex Ultimate 3000 HPLC	0	95	—	5	1	SunFire C18
		1	95	—	5	1	
		7	50	—	50	1	
		8	10	—	90	1	
		10.8	10	—	90	1	
		11	95	—	5	1	
		14	95	—	5	1	
L6	Dionex Ultimate 3000 HPLC	0	—	95	5	1	X-Bridge C18
		1	—	95	5	1	
		7	—	50	50	1	
		8	—	10	90	1	
		10.8	—	10	90	1	
		11	—	95	5	1	
		14	—	95	5	1	

377

Conditions used for the UPLC analysis in experimental part. The LC/MS analysis mentioned in the experimental part were performed on a Waters system combining an Acquity UPLC H-Class equipped with a Acquity UPLC PDA Detector and an Acquity TQ DetectorS (ES). The separations were performed with an Acquity UPLC HSS C18, 2.1×50 mm, 1.8 μM column equipped with a prefilter and thermostated at 40° C. or an Acquity UPLC BEH C18, 2.1×50 mm, 1.7 μM column equipped with a prefilter and thermostated at 40° C. and the PAD acquisition wavelength was set in the range of 210-420 nm (Method U in the table below). Elutions were carried out with the methods described in the following table. Solvent A: Formic Acid LC-MS grade 0.1% in milliQ water Solvent B: NH₄OAc (LC-MS grade) 10 mMol in milliQ water, adjusted at pH10 with an aq. solution of NH₃, LC-MS grade. Solvent C: Acetonitrile LC-MS grade.

		Solvents					
LC/MS Method	System	Time (min)	A (%)	B (%)	C (%)	Flow (mL/min)	Column
U1	Waters UPLC	0	95	—	5	0.5	Acquity
		3.18	50	—	50	0.5	UPLC
		4	10	—	90	0.5	HSS C18
U2	Waters UPLC	5	10	—	90	0.5	
		0	80	—	20	0.5	Acquity
		3.4	40	—	60	0.5	UPLC
U3	Waters UPLC	4	10	—	90	0.5	HSS C18
		5	10	—	90	0.5	
		0	50	—	50	0.5	Acquity
U4	Waters UPLC	3.5	10	—	90	0.5	UPLC
		5	10	—	90	0.5	HSS C18
		0	—	95	5	0.5	Acquity
U5	Waters UPLC	3.18	—	50	50	0.5	UPLC
		4	—	10	90	0.5	BEH C18
		5	—	10	90	0.5	
U6	Waters UPLC	0	—	80	20	0.5	Acquity
		3.4	—	40	60	0.5	UPLC
		4	—	10	90	0.5	BEH C18
U6	Waters UPLC	5	—	10	90	0.5	
		0	—	50	50	0.5	Acquity
		3.5	—	10	90	0.5	UPLC
		5	—	10	90	0.5	BEH C18

Preparative HPLC purifications mentioned in this experimental part have been carried out with the following system: a Waters 2489 UV/Visible Detector, a Waters 2545 Binary Gradient Module, a Waters Fraction Collector III and a Waters Dual Flex Injector. The separations were performed with a X-Bridge Prep C18 column, 100×19 mm, 5 μm column equipped with a X-Bridge C18, 19×10 mm, 5 μm Guard column or with a SunFire Prep C18 ODB column (5 μm; 19×100 mm) equipped with a SunFire C18 guard column (5 μm; 19×10 mm). Elutions were carried out with the methods described in the following tables, and detection wavelengths were fixed at 210 and 254 nm. Solvent A: Formic Acid LC-MS grade 0.1% in milliQ water Solvent B: NH₄OAc (LC-MS grade) 10 mMol in milliQ water, adjusted at pH10 with an aq. solution of NH₃, LC-MS grade. Solvent C: Acetonitrile LC-MS grade.


HPLC	Time	Solvent		Flow	
Method	(min)	A (%)	B (%)	(mL/min)	Column
H1	0	80	20	20	X-Bridge
	2	80	20	20	Prep C18
	8	10	90	20	

378

-continued

HPLC Method	Time (min)	Solvent		Flow (mL/min)	Column
5	10.8	10	90	20	
	11	80	20	20	
	16	80	20	20	
H2	0	95	5	20	SunFire
	2	95	5	20	Prep C18
	8	50	50	20	ODB
10	9	10	90	20	
	13	10	90	20	
	14	95	5	20	
15	16	95	5	20	

Synthesis of 2-methyl-5-((5-(trifluoromethyl)furan-2-yl)methoxy)benzofuran-3-carboxylic acid (Cpd 018)



Step 1: To a solution of ethyl 5-hydroxy-2-methylbenzofuran-3-carboxylate (0.100 g; 0.45 mmol) in THF (10 mL) and NMP (1 mL) was added cesium carbonate (0.444 g; 1.36 mmol) and the RM was stirred at RT for 25 min. Then, 2-(bromomethyl)-5-(trifluoromethyl)furan (0.104 mg; 0.45 mmol) was added and the stirred solution was heated at 95° C. until the consumption of 5-hydroxy-2-methylbenzofuran-3-carboxylate. The mixture was cooled down to RT, poured in water and extracted twice with EtOAc. The combined organic layer was dried over sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by FC on silica gel using a gradient of EtOAc (0% to 80%) in heptane to afford 0.167 g (100%) of ethyl 2-methyl-5-((5-(trifluoromethyl)furan-2-yl)methoxy)benzofuran-3-carboxylate.


Step 2: To a solution of ethyl 2-methyl-5-((5-(trifluoromethyl)furan-2-yl)methoxy)benzofuran-3-carboxylate (0.167 g; 0.45 mmol) in a mixture of water-EtOH-MeOH-THF (6:3:3:1, 12 mL) was added sodium hydroxide (0.102 g; 1.8 mmol) and the RM was heated under reflux until the consumption of ethyl 5-(benzyloxy)-2-methylbenzofuran-3-carboxylate. After cooling, the volatiles were removed under reduced pressure and the remaining residue was dissolved in

379

water and washed with DCM. The mixture was acidified with an aq. solution of hydrochloric acid (6 N) until pH~2 and extracted with EtOAc. The resulting organic layer was washed with water and brine, dried over magnesium sulfate and filtered to afford 147.6 mg (89%) of 2-methyl-5-((5-(trifluoromethyl)furan-2-yl)methoxy)benzofuran-3-carboxylic acid (cpd 018).

Cpd 016 and cpd 017 were prepared in a manner similar (use of appropriate reagents and purification methods known to those skilled in the art) to cpd 018.

Synthesis of 2-methyl-5-((2-(trifluoromethyl)benzyl)oxy)benzofuran-3-carboxylic acid (Cpd 101)



Step 1: Tributylphosphine (1.7 mL; 6.5 mmol) was added dropwise to a stirred mixture of methyl 5-hydroxy-2-methylbenzofuran-3-carboxylate (1 g; 4.6 mmol), 2-(trifluoromethyl)pyridin-3-ylmethanol (1.28 g; 6.9 mmol) and ADDP (1.66 g; 6.5 mmol) in dry THF (5 mL) under argon. The mixture was stirred for 2 h and was concentrated under reduced pressure. The residue was purified by FCC on silica gel using a gradient of EtOAc (0-20%) in heptane to afford 1.4 g (83%) of methyl 2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate. M/Z(+): 366 (M+H). M/Z(-): 364 (M-H).

Step 2: To a solution of methyl 2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (1.40 g; 3.8 mmol) in a mixture of MeOH-THF (1:1, 40 mL) was added a solution of sodium hydroxide (2 N; 5 mL; 40 mmol) and the RM was heated at 75° C. overnight. After cooling, the volatiles were removed under reduced pressure and the remaining residue was dissolved in water. The mixture was acidified with a solution of HCl (6 N) until pH~5. The white precipitate was washed with water and dried under reduced pressure to afford 1.30 g (96%) of 2-methyl-5-((2-(trifluoromethyl)benzyl)oxy)benzofuran-3-carboxylic acid (cpd 101).


Cpd 099, cpd 100 and cpd 102 were prepared in a manner similar (use of appropriate reagents and purification methods known to those skilled in the art) to cpd 101.

The following compounds were prepared in a manner similar (use of appropriate reagents and purification methods known to the person skilled in the art) to cpd 101:

380

Cpd. Nr.	[M + H] ⁺ (m/z)	Cpd. Nr.	[M - H] ⁻ (m/z)
208	304.0	261	297.0
210	372.1	263	297.0
211	288.1	679	288.1
212	304.1	681	332.1
214	298.1	713	285.1
216	304.1		
217	298.1		
218	263.1		
219	285.0		
220	337.1		
223	318.0		
224	318.1		
227	314.2		
228	303.1		
235	337.0		
239	328.1		
240	302.1		
242	302.1		
248	318.1		
249	287.1		
257	299.1		
260	304.0		
262	304.0		
264	285.0		
268	318.1		
269	314.1		

Synthesis of 2-methyl-5-((1-methyl-1H-pyrazol-3-yl)methoxy)benzofuran-3-carboxylic acid (Cpd 008)



Step 1: Ethyl 5-hydroxy-2-methylbenzofuran-3-carboxylate (100 mg; 0.45 mmol), (1-methyl-1H-pyrazol-3-yl)methanol (0.075 g; 0.68 mmol) and PPh₃ (119 mg; 0.45 mmol) were dissolved in THF (5 mL) and cooled down to 0° C. Then DIAD (0.134 mL; 0.68 mmol) was added dropwise and the RM was stirred 36 h at RT. The volatiles were removed under reduced pressure and the residue was diluted with DCM and washed successively with an aq. solution of NaOH (1 N), water and brine. The organic layer was dried over magnesium sulfate, filtered and the solvent was removed under reduced pressure. The compound was purified by FCC on silica gel using a gradient of EtOAc (20 to 100%) in heptane to afford 0.071 g (50%) of the desired compound.

381


Step 2: An aq. solution of NaOH (2 N; 6 mL; 12 mmol) was added to a solution of ethyl 2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxylate (0.071 g; 0.25 mmol) in MeOH (4 mL) and ethanol (2 mL) and the mixture was stirred for 4 h at 80° C. After cooling, the mixture was diluted with water and washed with DCM. The aq. layer was acidified till pH=4 with an aq. solution of HCl (6 N). The solid formed was filtered and dried to afford 33 mg (51%) of 2-methyl-5-((1-methyl-1H-pyrazol-3-yl)methoxy)benzofuran-3-carboxylic acid (cpd 008).

The following compounds were prepared in a manner similar (use of appropriate reagents and purification methods known to the person skilled in the art) to cpd 008: Cpd 001, 002, 004, 005, 006, 007, 009, 010, 011, 012, 013 and 014.

The following compounds were prepared in a manner similar (use of appropriate reagents and purification methods known to the person skilled in the art) to cpd 008:

Cpd. Nr.	[M + H] ⁺ (m/z)
255	288.1
266	328.1
267	323.1

Synthesis of 2-methyl-5-((5-methylisoxazol-3-yl)methoxy)benzofuran-3-carboxylic acid (Cpd 215)



Step 1: To a stirred solution of ethyl 5-hydroxy-2-methylbenzofuran-3-carboxylate (0.4 g, 1.8 mmol) and (5-methylisoxazol-3-yl)methyl methanesulfonate (0.52 g, 2.7 mmol) in acetonitrile (25 mL) was added Cs₂CO₃ (1.18 g, 3.6 mmol) at RT. The reaction mixture was heated 70° C. and maintained at this temperature for 16 h. Reaction progress was monitored by TLC. The reaction mixture was filtered and concentrated under reduced pressure to give the crude compound. The crude was purified by 100-200 mesh silica-gel column chromatography using 5% ethyl acetate in pet ether as an eluent to afford ethyl 2-methyl-5-((5-methylisoxazol-3-yl)methoxy)benzofuran-3-carboxylate (0.4 g, 70%) as a pale yellow thick mass. TLC system: 30% Ethyl acetate in pet-ether; RF: 0.2.


382

Step 2: 2N aqueous NaOH (5 mL) was added to a stirred solution of ethyl 2-methyl-5-((5-methylisoxazol-3-yl)methoxy)benzofuran-3-carboxylate (0.4 g, 1.26 mmol) in methanol (5 mL) and THF (5 mL) at RT and the resulting reaction mixture was stirred at RT for 16 h. The reaction progress was monitored by TLC. The reaction mixture was diluted with water (10 mL) and acidified to pH ~2 with 1 N aq. HCl (10 mL), the precipitated solid was filtered and dried under vacuum to afford 2-methyl-5-((5-methylisoxazol-3-yl)methoxy)benzofuran-3-carboxylic acid (250 mg, 70%) as a white solid. TLC system: 50% Ethyl acetate in pet-ether; RF: 0.2. [M+H]⁺ (m/z): 288.1

The following compounds were prepared in a manner similar (use of appropriate reagents (including mesylates, chlorides and bromides) and purification methods known to the person skilled in the art) to cpd 215:

Cpd. Nr.	[M + H] ⁺ (m/z)	Cpd. Nr.	[M + H] ⁺ (m/z)
209	358.1	237	334.0
213	383.3	250	314.1
221	364.1	256	314.0
226	334.0	258	277.2
667	301.2	669	335.1
671	349.1	673	288.1
675	315.2	677	302.2
689	327.1	691	355.1
693	358.0	699	298.1
703	324.2	707	299.1

Synthesis of 5-((2-aminopyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid (Cpd 225)




Step 1: To a pre-stirred solution of ethyl 5-hydroxy-2-methylbenzofuran-3-carboxylate (0.900 g, 4.09 mmol) in THF (30 mL), was added NaH (60%) (0.313 g, 8.18 mmol) lot wise at 0° C. under argon atmosphere, followed by the addition of 3-(chloromethyl)pyridin-2-amine (2239-1) (0.871 g, 6.13 mmol) at 0° C. under argon atmosphere. The reaction mixture was heated up to 60° C. for 18 h. The reaction progress was monitored by TLC. The reaction mixture was diluted with water (30 mL) and extracted with ethyl acetate (2x50 mL). The combined organic layer was

383

washed with water (30 mL) and brine (30 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to give the crude material. The crude was purified by column chromatography over silica gel (100-200 mesh) using 0-30% ethyl acetate in pet-ether as an eluent to afford ethyl 5-((2-aminopyridin-3-yl) methoxy)-2-methylbenzofuran-3-carboxylate (0.30 g, 23%) as a white solid. TLC system: 50% EtOAc in Pet ether; Rf: 0.3.

Step 2: To a pre-stirred solution of ethyl 5-((2-aminopyridin-3-yl) methoxy)-2-methylbenzofuran-3-carboxylate (0.30 g, 0.92 mmol), in THF:MeOH:H₂O (1:1:1, 30 ml), was added NaOH (0.11 g, 2.76 mmol) at RT, then the reaction was heated to 50° C., and the reaction mixture was stirred for 6 h at 50° C. Reaction progress was monitored by TLC. The reaction mixture was concentrated to get the crude material, which was diluted with ice water and pH was adjusted to ~2 with 1 N aq. HCl solution to give a solid. The obtained solid was filtered and dried to afford 5-((2-aminopyridin-3-yl) methoxy)-2-methylbenzofuran-3-carboxylic acid (0.22 g, 80%) as an off-white solid. TLC system: 10% MeOH in CH_2Cl_2 ; Rf: 0.2. $[\text{M}+\text{H}]^+$ (m/z): 299.1

Synthesis of 2-methyl-5-((2-methylpyridin-3-yl) methoxy)benzofuran-3-carboxylic acid (Cpd 247)




Step 1: To a solution of ethyl 5-hydroxy-2-methylbenzofuran-3-carboxylate (800 g, 3.636 mmol) and (2-methylpyridin-3-yl)methyl methanesulfonate (1.4 g 7.272 mmol) in DMF (5 mL) at 0° C. was added K_2CO_3 (1.5 g, 10.909 mmol). The reaction mixture was allowed to come to RT and was stirred for 12 h at the same temperature. Reaction progress was monitored by TLC. The reaction mixture was quenched with ice water (100 mL) and stirred for 30 minutes. The solid material was collected by filtration and dried to afford 1.2 g (92%) of ethyl 2-methyl-5-((2-methylpyridin-3-yl)methoxy)benzofuran-3-carboxylate as an off white solid. TLC system: 20% Ethyl acetate in pet-ether; Rf: 0.20

Step 2: A solution of ethyl 2-methyl-5-((2-methylpyridin-3-yl)methoxy)benzofuran-3-carboxylate (1.2 g, 3.692 mmol), NaOH (295 mg, 7.384 mmol) in EtOH:THF:H₂O (2:2:1, 5.0 mL) was stirred at 70° C. for 16 h. The reaction progress was monitored by TLC. The solvent was distilled

384

off, then water was added (20 mL) to the crude residue and the aqueous phase was washed with diethyl ether (2×10 mL). The aqueous layer was acidified with 2N HCl, the resulting precipitated solid was collected by filtration, washed with water (10 mL) and dried under reduced pressure to afford 2-methyl-5-((2-methyl-4-(trifluoromethyl)thiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (1.0 g, 91%) as an off-white solid. TLC system: 10% MeOH in dichloromethane; Rf: 0.10. $[\text{M}+\text{H}]^+$ (m/z): 298.1

Synthesis of 5-((5-(2-hydroxyethoxy)pyridin-2-yl) methoxy)-2-methylbenzofuran-3-carboxylic acid (Cpd 241)



Step 1: To a stirred solution of ethyl 5-hydroxy-2-methylbenzofuran-3-carboxylate (5 g, 22.7 mmol), and (5-methoxypyridin-2-yl)methanol (3.79 g, 27.2 mmol) in THF (150 mL) was added ADDP (11.4 g, 46 mmol) followed by Bu_3P (9.2 g, 46 mmol) at 0° C. The reaction mixture was stirred at RT for 16 hours and was then concentrated under reduced pressure. The residue was partitioned between water and ethyl acetate. After separation, the aqueous layer was extracted with ethyl acetate (150 mL×2). The combined organic extracts were concentrated under reduced pressure to give a residue. The residue was purified by flash chromatography (silica gel, 100-200 mesh size) using a gradient of ethyl acetate (5-40%) in hexane to afford ethyl 5-((5-(2-methoxyethoxy)pyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxylate (3 g, 8.7 mmol, 38%) as a pale yellow solid. TLC system: 30% EtOAc in Pet ether; Rf: 0.35.


385

Step 2: To a stirred solution of ethyl 5-((5-methoxypyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxylate (500 mg, 1.4 mmol) in CH_2Cl_2 (20 mL) was added portion wise AlCl_3 (390 mg, 2.9 mmol) at 0°C ., the reaction mixture was then stirred at 50°C . for 2 hours. Reaction progress was monitored by LCMS. After completion of the reaction, the reaction mass was concentrated under reduced pressure to give a residue. The residue was treated with sat. NaHCO_3 solution (pH ~ 7) and was partitioned between water and dichloromethane. After separation, the aqueous layer was extracted with dichloromethane (50 mL $\times 2$). The combined organic extracts were concentrated under reduced pressure to afford ethyl 5-((5-hydroxypyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxylate (180 mg, 5 mmol) as a pale yellow solid. TLC system: 50% EtOAc in pet ether; RF: 0.3

Step 3: To a stirred solution of ethyl 5-((5-hydroxypyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxylate (600 mg, 1.8 mmol) in DMF (20 mL) was added (2-bromoethoxy)(tert-butyl)dimethylsilane (657 mg, 2.7 mmol) at 0°C ., the reaction mixture was then stirred at 60°C . for 6 hours. Reaction progress was monitored by LCMS. After completion of the reaction, the reaction mass was partitioned between water and ethyl acetate. After separation, the aqueous layer was extracted with ethyl acetate (50 mL $\times 2$). Combined organic extracts were concentrated under reduced pressure to afford ethyl 5-((5-(2-((tert-butyl)dimethylsilyl)oxy)ethoxy)pyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxylate (600 mg, 1.2 mmol) as a pale yellow solid. TLC system: 30% EtOAc in Pet ether; RF: 0.7.


Step 4: To a stirred solution of ethyl 5-((5-(2-((tert-butyl)dimethylsilyl)oxy)ethoxy)pyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxylate (600 mg, 1.2 mmol) in THF (20 mL) was added NaOH (148 mg, 3 mmol) in water (3 mL), and MeOH (3 mL) at 0°C ., the reaction mixture was then stirred at RT for 16 hours. Reaction progress was monitored by LCMS. After completion of the reaction, the reaction mass was concentrated under reduced pressure. The residue was stirred in 1 N HCl (pH ~ 2) for 30 min and was then partitioned between water and dichloromethane. After separation, the aqueous layer was extracted with dichloromethane (50 mL $\times 2$). Combined organic extracts were concentrated under reduced pressure to afford 5-((5-(2-hydroxyethoxy)pyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid (300 mg, 8 mmol) as an off-white solid. TLC system: 5% MeOH in dichloromethane; RF: 0.4. $[\text{M}+\text{H}]^+$ (m/z): 344.0

Synthesis of 5-((2-(dimethylamino)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid (Cpd 222)



386

-continued




Step 1: To a pre-stirred solution of ethyl 5-hydroxy-2-methylbenzofuran-3-carboxylate (2.5 g, 0.011 mmol) in acetonitrile (50 mL) was added Cs_2CO_3 (7.3 g, 0.022 mmol), 2-chloro-3-(chloromethyl) pyridine (2.1 g, 0.0132 mmol) under argon atmosphere at RT. The reaction mixture was heated to 80°C . for 16 h. The reaction progress was monitored by TLC. The reaction mixture was filtered and the filter cake was washed with acetonitrile (50 mL), and the filtrate was concentrated under reduced pressure to give the crude material. The crude material was purified by grace chromatography using 0-20% ethyl acetate in pet-ether as an eluent to afford ethyl 5-((2-chloropyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylate (2.0 g, 63%) as an off-white solid. TLC system: 30% EtOAc in Pet ether; RF: 0.6.

Step 2: A solution of ethyl 5-((2-chloropyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylate (500 mg, 1.44 mmol) and N,N-dimethyl amine solution in ethanol (10 mL, 5 M) was heated to 150°C . in a microwave for 2 h. The reaction progress was monitored by TLC. The reaction mixture was concentrated under reduced pressure to give the crude material. The crude material was purified by Grace flash chromatography using 0-5% ethyl acetate in pet-ether as an eluent to afford ethyl 5-((2-(dimethylamino)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylate (180 mg, 39%) as an off-white solid. TLC system: 30% EtOAc in Pet ether; RF: 0.5.

Step 3: To a stirred solution of ethyl 5-((2-(dimethylamino)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylate (300 mg, 0.847 mmol), in THF:MeOH:H₂O (1:1:0.5, 25 mL) was added NaOH (170 mg, 4.27 mmol) at RT and the mixture was heated to 50°C . The reaction mixture was stirred for 4 h at 50°C . and reaction progress was monitored by TLC. The reaction mixture was concentrated under vacuum and diluted with ice water, the pH was adjusted to ~ 2 with 1 N aq. HCl solution to give a solid which was filtered and dried to afford 5-((2-(dimethylamino)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid (170 mg, 61%) as a white solid. TLC system: 30% EtOAc in Pet ether; RF: 0.1. $[\text{M}+\text{H}]^+$ (m/z): 327.2.

387

Synthesis of 5-((2-cyclopropylpyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid (Cpd 701)



Step 1: To a pre-stirred solution of ethyl 5-hydroxy-2-methylbenzofuran-3-carboxylate (500 mg, 2.283 mmol) and (2-bromopyridin-3-yl)methanol (643.4 mg, 3.42 mmol) in THF (10 mL) was added ADDP (806.08 mg, 3.196 mmol) and *n*-Bu₃P (646.5 mg, 3.196 mmol) at RT. The RM was stirred for 16 h at RT. The reaction progress was monitored by TLC. The RM was diluted with water (100 mL) and extracted with ethyl acetate (2×50 mL). Combined organic layers were washed with brine (50 mL), dried over anhydrous Na₂SO₄ and evaporated under vacuum. The crude was purified by flash column chromatography using 0.1% formic acid in water and acetonitrile in water as an eluent to afford ethyl 5-((2-bromopyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylate (520 mg, crude) as yellow solid. Proceeded to next step without purification.


Step 2: Pd(PPh₃)₂Cl₂ (45.04 mg, 0.06 mmol) was added to a stirred solution of ethyl 5-((2-bromopyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylate (250 mg, 0.6426 mmol), cyclopropylboronic acid (83.38 mg, 0.9640 mmol), K₃PO₄ (409.15 mg, 1.9278 mmol) in 1,4-dioxane (10 mL) and water (4 mL) at RT. Reaction mixture was degassed with Ar gas for 15 minutes at RT. The reaction mixture was heated to 100° C. and stirred for 16 hours at 100° C. The reaction progress monitored by TLC. Reaction mass was filtered over celite bed, filtrate was diluted with water (20 mL) and extracted with ethyl acetate (2×50 mL). Combined organic layers were concentrated under reduced pressure to get crude. Crude was purified by Grace flash column chromatography using 0.1% formic acid in water

388

and acetonitrile as an eluent to afford ethyl 5-((2-cyclopropylpyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylate 150 mg, crude) as a brown solid. Proceeded to next step without purification. TLC system: 100% Ethyl acetate, RF: 0.4.

Step 3: A solution of NaOH (91.4 mg, 2.285 mmol) in water (8 mL) was added to a stirred solution of ethyl 5-((2-cyclopropylpyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylate (80 mg, 0.228 mmol) in methanol (2 mL) and THF (2 mL) at RT. The resulting reaction mixture was stirred for 16 h at RT. The reaction progress was monitored by TLC. The reaction mixture was concentrated under reduced pressure and diluted with water (15 mL), pH was adjusted to ~5 with 1 N HCl solution. The precipitated solid was extracted with ethyl acetate (2×50 mL). Combined organic layers were dried over Na₂SO₄ and concentrated under reduced pressure to afford 5-((2-cyclopropylpyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid (6) (100 mg, crude) as a brown gummy liquid. Proceeded to next step without purification. TLC system: 10% MeOH in dichloromethane; RF: 0.2.

Synthesis of 5-((2-aminopyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid (Cpd 230)




Step 1: To a stirred solution of ethyl 2-formyl-5-((2-methoxypyridin-3-yl)methoxy)benzofuran-3-carboxylate (1.5 g, 13.8 mmol) in CH₂Cl₂ (100 mL) was added DAST (4.47 g, 28 mmol) at 0° C. and the reaction mixture was stirred for 16 h at RT, the reaction progress was monitored by TLC. Ice cold water was added to the reaction mixture and aq. NaHCO₃ solution was added to get to pH ~7, followed by extraction with ethyl acetate (3×100 mL). The combined extracts were washed with brine (150 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to get the material crude (1.4 g). The crude was purified by column chromatography using silica gel (100-200 mesh) and 20% ethyl acetate in pet ether as eluent to afford ethyl 2-(difluoromethyl)-5-((2-methoxypyridin-3-yl)

389

methoxy)benzofuran-3-carboxylate (1 g) as a white solid. TLC system: 20% Ethyl acetate in pet-ether; Rf: 0.6.

Step 2: To a stirred solution of ethyl 2-(difluoromethyl)-5-((2-methoxypyridin-3-yl)methoxy)benzofuran-3-carboxylate (1 g, 1.4 mmol), was added LiOH·H₂O (110 mg, 2.8 mmol) in THF:H₂O:MeOH (3:1:1) (25 mL) at RT under an argon atmosphere. The reaction mixture was stirred for 2 h at RT. The reaction progress was monitored by TLC. The solvents were removed and the reaction mixture was treated with citric acid to adjust the pH to ~1 and the resulting precipitate was filtered. The solid was dried under vacuum to afford 2-(difluoromethyl)-5-((2-methoxypyridin-3-yl)methoxy)benzofuran-3-carboxylic acid (1 g, 68%) as an off-white solid. TLC system: 5% MeOH in CH₂Cl₂; Rf: 0.2. [M+H]⁺ (m/z): 350.0.

Synthesis of 5-((2-(azetidin-1-yl)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid (Cpd 236)




Step 1: To a stirred solution of ethyl 5-((2-chloropyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylate (4.0 g, 11.59 mmol, 1.0 eq) in DMSO (40 ml), at 0° C. was added K₂CO₃ (15.99 g, 115.90 mmol, 10 eq) at 0° C. and the mixture was stirred for 5 min. Azetidine (5.39 g, 57.97 mmol, 5 eq) was added to the reaction and the mixture was stirred for 18 h at 120° C. Reaction progress was monitored by TLC. The reaction mixture was diluted with ice water (50 ml), the product was extracted with EtOAc (300 ml). The ethyl acetate layer was washed with ice water (50 ml) and brine (25 ml) and was then dried over anhydrous Na₂SO₄, filtered and concentrated to get the crude product, which was purified by Grace flash column chromatography using 20% EtOAc-pet ether as eluent to afford ethyl 5-((2-(azetidin-1-yl)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylate (0.380 g, 9%, Off-white solid). TLC system: 30% Ethyl acetate in pet-ether; Rf: 0.3.

Step 2: To a stirred solution of ethyl 5-((2-(azetidin-1-yl)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylate (0.700 g, 1.91 mmol, 1.0 eq), in THF:MeOH:H₂O (1:1:1)

390

(60 ml), was added NaOH (0.29 g, 5.73 mmol, 3.0 eq) at RT, the mixture was then heated up to 50° C. The reaction mixture was stirred for 18 h at 50° C. and reaction progress was monitored by TLC. The reaction mixture was concentrated to get the crude material which was diluted with ice water and the pH was adjusted to 2 with 1 N HCl solution causing precipitation. The contents were further stirred for 30 min, the solids were then filtered off and dried to afford 5-((2-(azetidin-1-yl)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid (0.400 g, 61% white solid). TLC system: 100% Ethyl acetate; Rf: 0.2. [M+H]⁺ (m/z): 339.0.

Synthesis of 5-((2-cyanopyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid (Cpd 238)



Step 1: To a stirred solution of ethyl 5-hydroxy-2-methylbenzofuran-3-carboxylate (0.8 g, 3.63 mmol, 1.0 eq) in ACN (10 ml) at 0° C. was added Cs₂CO₃ (3.55 g, 10.89 mmol, 3 eq) and the mixture was stirred for 5 minutes followed by the addition of 3-(bromomethyl)picolinonitrile (0.85 g, 4.36 mmol, 1.2 eq) at 0° C. The reaction mixture was stirred for 18 h at RT, monitored by TLC. The reaction mixture was diluted with ice water (100 ml) and extracted with EtOAc (300 ml). The organic layer was washed with ice water (100 ml) and brine (50 ml), dried over anhydrous Na₂SO₄, filtered and concentrated to get the crude product, which was purified by Grace flash column chromatography using eluent 20% EtOAc in pet ether to afford ethyl 5-((2-cyanopyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylate (1.0 g, 81%, off-white solid). TLC system: 20% EtOAc-pet ether; 2:8; Rf: 0.3.

Step 2: To ethyl 5-((2-cyanopyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylate (1.0 g, 2.97 mmol, 1.0 eq), in THF:MeOH:H₂O (1:1:1) (120 ml) was added NaOH (0.35 g, 8.92 mmol, 3.0 eq) at RT, the reaction mixture was then heated to 50° C. for 18 h, the reaction progress was monitored by TLC. The reaction mixture was concentrated and diluted with ice water, the pH was adjusted to ~2 with 1 N HCl solution to give a precipitate while stirring for 30 min. The precipitate was filtered off and dried under vacuum to afford a mixture of 5-((2-cyanopyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid (cpd 238) and 5-((2-


391

carbamoylpyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid (0.75 g, crude). $[M+H]^+$ (m/z): 309.0, TLC system: 50% EtOAc-pet ether; Rf: 0.1.


The following compounds were prepared in a manner similar (use of appropriate reagents and purification methods known to the person skilled in the art) to cpd 238:

Cpd. Nr.	$[M+H]^+$ (m/z)
259	288.1
687	355.1
709	340.2

Synthesis of 2-methyl-5-(((2-(trifluoromethyl)pyridin-3-yl)oxy)methyl)benzofuran-3-carboxylic acid (Cpd 252)



392



cpd 252

Step 1. DABCO (2.78 g, 24.75 mmol) was added to a stirred mixture of 3-bromo-4-hydroxybenzaldehyde (5.0 g, 24.75 mmol) and ethyl but-2-ynoate (3.32 g, 29.70 mmol) in acetonitrile (100 mL) at room temperature and the resulting reaction mixture was then heated to reflux for 4 h. The solvent was distilled off under reduced pressure, the residue was diluted with ethyl acetate (300 mL) and the resulting solution was sequentially washed with 1 N HCl (100 mL), aq. 1 N NaOH (2x75 mL), water (100 mL), brine (150 mL) and was then dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The crude compound was purified by column chromatography over silica-gel (100-200 mesh) using 10% ethyl acetate in pet-ether as an eluent to afford ethyl 3-(2-bromo-4-formylphenoxy)but-2-enoate (1.5 g, 29%) as an off-white solid. TLC system: 20% Ethyl acetate in pet-ether; Rf: 0.7.

Step 2: A solution mixture of ethyl 3-(2-bromo-4-formylphenoxy)but-2-enoate (1.2 g, 3.821 mmol) and TEA (0.21 mL, 1.528 mmol) in acetonitrile (25.0 mL) was degassed with argon for 5 min and then was added bis(tri-tert-butylphosphine)palladium(0)] (195 mg, 0.382 mmol). The resulting reaction mixture was heated to reflux for 2 h. The reaction progress was monitored by TLC. (Note: Another batch on 250 mg scale using the same procedure was combined with this reaction mixture for work-up). The reaction mixture was directly concentrated under reduced pressure. The crude was partitioned between water (100 mL) and ethyl acetate (2x100 mL). The combined organic layers were washed with brine (150 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford ethyl 5-formyl-2-methylbenzofuran-3-carboxylate (1.0 g, 90%) as an off white crystalline solid. TLC system: 10% Ethyl acetate in pet-ether; Rf: 0.7.

Step 3: NaBH_4 (398 mg, 10.775 mmol) was added portionwise to a solution mixture of ethyl 5-formyl-2-methylbenzofuran-3-carboxylate (1.0 g, 4.310 mmol) in EtOH: THF (1:1) (40.0 mL) at 0°C . and the resulting reaction mixture was stirred for 30 min. The reaction mixture was then quenched with water (5.0 mL) and was concentrated under reduced pressure. The crude compound was diluted with ethyl acetate (50 mL), washed with water (2x50 mL), brine (100 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford ethyl 5-(hydroxymethyl)-2-methylbenzofuran-3-carboxylate (900 mg, crude) as a colorless oil. TLC system: 20% Ethyl acetate in pet-ether; Rf: 0.4.

Step 4: A mixture of ethyl 5-(hydroxymethyl)-2-methylbenzofuran-3-carboxylate (700 mg, 2.991 mmol), TPP (1.17 g, 4.487 mmol) and CBr_4 (1.48 g, 4.487 mmol) in THF (40.0 mL) was stirred at 0°C . was then warmed to ambient temperature for 2 h. The reaction progress was monitored by TLC. The reaction mixture was diluted with ethyl acetate (100 mL), washed with NaHCO_3 solution (50 mL), water (50 mL), brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford ethyl 5-(bromomethyl)-2-methylbenzofuran-3-carboxylate (1.25 g,


393

crude) as a pale yellow thick gummy mass. TLC system: 10% Ethyl acetate in pet-ether; Rf: 0.6.


Step 5: To a suspension of 2-(trifluoromethyl)pyridin-3-ol (600 mg, 3.680 mmol) and K_2CO_3 (1.26 g, 9.20 mmol) in acetonitrile (30 mL) was added ethyl 5-(bromomethyl)-2-methylbenzofuran-3-carboxylate (1.32 g, 4.41 mmol) and the resulting reaction mixture was maintained at 90° C. for 1 h. The reaction progress was monitored by TLC. The reaction mixture was diluted with ethyl acetate (100 mL) and was filtered. The filtrate was washed with water (100 mL), brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The crude compound was purified by column chromatography over silica-gel (100-200 mesh) using 20% ethyl acetate in pet-ether as an eluent to afford ethyl 2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)oxy)methyl)benzofuran-3-carboxylate (800 mg, 58%) as a red solid. TLC system: 20% Ethyl acetate in pet-ether; Rf: 0.5.

Step 6: To a stirred solution of ethyl 2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)oxy)methyl)benzofuran-3-carboxylate (800 mg, 2.110 mmol) in ethanol:THF (1:1) (40 mL) was added a solution of NaOH (253 mg, 6.332 mmol) dissolved in water (10 mL) and the resulting reaction mixture was maintained under stirring at 60° C. for 1 h. The reaction progress was monitored by TLC. The reaction mixture was cooled to room temperature and poured into ice cold water (100 mL), acidified with 1 N HCl to pH~5.0 and extracted with ethyl acetate (2x100 mL). The combined organic layer was washed with water (100 mL) followed by brine (100 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford 2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)oxy)methyl)benzofuran-3-carboxylic acid (800 mg, crude) as a red solid. TLC system: 10% MeOH in dichloromethane; Rf: 0.4. $[M+H]^+$ (m/z): 352.0.

Synthesis of 2,4-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid (Cpd 251)



394




Step 1: 2-(Trifluoromethyl)pyridin-3-yl)methanol (0.887 mg, 5.01 mmol), ADDP (0.946 mg, 4.67 mmol) and tri-N-butylphosphine (1.18 g, 4.67 mmol) were added sequentially to a pre-stirred solution of ethyl 4-bromo-5-hydroxy-2-methylbenzofuran-3-carboxylate (1.0 g, 3.34 mmol) in THF (50 mL) at 0° C. under an argon atmosphere. The reaction mixture was warmed to room temperature and the stirred for 2 h. The reaction progress was monitored by TLC. The reaction mixture was poured into water (80 mL) followed by extraction with ethyl acetate (2x60 mL). The combined organic layers were sequentially washed with water (30 mL), brine (30 mL), were then dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The crude product was purified by column chromatography over silica gel (100-200 mesh) using 0-20% ethyl acetate in pet-ether as an eluent to afford ethyl 4-bromo-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (0.600 g, crude) as a pale yellow liquid. TLC system: 20% Ethyl acetate in pet-ether; Rf: 0.6.

Step 2: A suspension of ethyl 4-bromo-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (0.6 g, 1.31 mmol), methylboronic acid (0.157 g, 2.62 mmol) and K_2CO_3 (0.542 g, 3.93 mmol) in 1,4-dioxane:water (9:1) (10 mL) in a sealed tube was degassed with argon for 10 min and then $Pd(PPh_3)_4$ (0.151 g, 0.13 mmol) was added. The resulting reaction mixture was again degassed for 10 minutes and was then heated to 120° C. for 16 h. The reaction progress was monitored by LC-MS. The reaction mixture was cooled to room temperature and filtered through a celite pad, the celite pad was washed with ethyl acetate (2x30 mL). The combined filtrate was dried over anhydrous Na_2SO_4 and evaporated in vacuo. The crude product was purified by flash column chromatography over silica gel (100-200 mesh) using 0-20% ethyl acetate in pet-ether as an eluent to afford ethyl 2,4-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (0.5 g, 71%) as a colorless liquid. TLC system: 20% Ethyl acetate in pet-ether; Rf: 0.7.

Step 3: A solution of NaOH (0.203 g, 5.08 mmol) in water (4 mL) was added to a pre-stirred solution ethyl 2,4-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (0.5 g, 1.27 mmol) in a mixture of methanol:THF (1:1) (14 mL) and the resulting reaction mixture was heated to 60° C. for 4 h. The reaction progress was monitored by TLC. The reaction mixture was cooled and poured into ice cold water (30 mL) and was acidified with 1 N HCl to pH~2. The crude product was extracted with ethyl acetate (2x50 mL). The combined organic layer was washed with water (30 mL), brine (30 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford 2,4-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid (0.350 g, 65%) as an off-white solid. This crude product was used for next step without any further purification. TLC system: 50% Ethyl acetate in pet-ether; Rf: 0.3. $[M+H]^+$ (m/z): 366.1

395

Synthesis of 2-cyclopropyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (Cpd 265)



Step 1: To a stirred solution of benzoquinone (1.5 g, 13.8 mmol) in IPA (50 ml) was added ZnCl_2 (9.4 g, 69.44 mmol) followed by the addition of methyl 3-cyclopropyl-3-oxopropanoate (7.8 g, 55.55 mmol) at RT. The reaction mixture was stirred for 6 h at reflux, the reaction progress was monitored by TLC. Ice cold water was added to the reaction mixture and the mixture was then extracted with ethyl acetate (2x300 mL). The combined extracts were washed with brine (100 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to get the crude material (3 g). The crude material was purified by column chromatography using silica gel (100-200 mesh) and 20% ethyl acetate in pet ether as an eluent to afford methyl 2-cyclopropyl-5-hydroxybenzofuran-3-carboxylate (800 mg, 74%) as a brown solid. TLC system: 50% ethyl acetate in pet-ether; RF: 0.63.

Step 2: To a stirred solution of methyl 2-cyclopropyl-5-hydroxybenzofuran-3-carboxylate (300 mg, 1.29 mmol) in CH_3CN (25 mL) was added Cs_2CO_3 (1.2 g, 3.87 mmol) followed by the addition of (4-methylthiazol-5-yl) methyl methanesulfonate (370 mg, 1.93 mmol) at 0°C . under an argon atmosphere. The reaction mixture was stirred for 16 h at 50°C . The reaction progress was monitored by TLC. Solvents were removed, ice cold water was added to the reaction mixture and the mixture was extracted with ethyl acetate (2x100 mL). The combined extracts were washed with brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to get the crude material (400 mg). The crude was purified by column chromatography using silica gel (100-200 mesh) and 25% ethyl acetate in pet ether as an eluent to afford methyl 2-cyclopropyl-5-((4-methylthiazol-5-yl) methoxy) benzofuran-3-carboxylate (200 mg, 45%) as a brown solid. TLC system: 50% ethyl acetate in pet-ether; RF: 0.42.

Step 3: To a stirred solution of methyl 2-cyclopropyl-5-((4-methylthiazol-5-yl) methoxy) benzofuran-3-carboxylate (200 mg, 0.583 mmol) in MeOH & THF (1:1, 10 ml) was added 2N NaOH (4 ml) at 0°C ., then the reaction mixture was stirred for 4 h at 50°C . Reaction progress was monitored by TLC. The reaction mixture was concentrated,


396

acidified to pH ~2 with 1 N aq HCl and extracted with ethyl acetate (3x50 mL). The combined extracts were washed with brine (20 ml), dried over anhydrous Na_2SO_4 , filtered, and concentrated under reduced pressure to get crude 2-cyclopropyl-5-((4-methylthiazol-5-yl) methoxy) benzofuran-3-carboxylic acid (120 mg, crude) as an off-white solid. $[\text{M}+\text{H}]^+$ (m/z): 330.1

The following compounds were prepared in a manner similar (use of appropriate reagents and purification methods known to the person skilled in the art) to cpd 265:

Cpd. Nr.	$[\text{M} + \text{H}]^+$ (m/z)
231	378.1

Synthesis of tert-butyl 3-(((3-(ethoxycarbonyl)-2-methylbenzofuran-5-yl)oxy)methyl)pyrrolidine-1-carboxylate (cpd 079), 5-((1-(tert-butoxycarbonyl)pyrrolidin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid (Cpd 019) and 2-methyl-5-(pyrrolidin-3-ylmethoxy)benzofuran-3-carboxylic acid (Cpd 003)



Step 1: To a cold solution of ethyl 5-hydroxy-2-methylbenzofuran-3-carboxylate (0.110 g; 0.5 mmol), tert-butyl

397


3-(hydroxymethyl)pyrrolidine-1-carboxylate (0.148 g; 0.75 mmol) and PS—PPh₃ (loading ~1.83 mmol/g; 0.546 g; 1 mmol) in THF (4 mL by mmol of ethyl 5-hydroxy-2-methylbenzofuran-3-carboxylate) was slowly added a solution of DIAD (0.148 mL; 0.75 mmol) in THF (0.4 mL). The RM was stirred at RT for 18 h and the volatiles were removed under reduced pressure. The crude material was purified by FCC on silica gel using a gradient of DCM (0% to 100%) in heptane to furnish 0.057 g (28%) of tert-butyl 3-(((3-(ethoxycarbonyl)-2-methylbenzofuran-5-yl)oxy)methyl)pyrrolidine-1-carboxylate (cpd 079).

Step 2: To a solution of tert-butyl 3-(((3-(ethoxycarbonyl)-2-methylbenzofuran-5-yl)oxy)methyl)pyrrolidine-1-carboxylate (cpd 079) (0.057 g; 0.141 mmol) in a mixture of MeOH:EtOH (2:1; 4.2 mL) was added an aq. solution of NaOH (1 N; 4.2 mL; 4.2 mmol) and the RM was heated under reflux for 18 h. After cooling, the volatiles were removed under reduced pressure and the remaining residue was solubilized in water. The mixture was acidified with an aq. solution of HCl (6 N) until pH~2. The white precipitate was washed with water and dried under high vacuum to furnish 0.052 g (98%) of 5-((1-(tert-butoxycarbonyl)pyrrolidin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid (cpd 019).

Step 3: To a solution of 5-((1-(tert-butoxycarbonyl)pyrrolidin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid (cpd 019) (0.050 g; 0.133 mmol) in THF (1 mL) was added a solution of HCl (4 N in dioxane; 1 mL) and the RM was stirred for 18 h. The volatiles were removed under reduced pressure and the remaining residue was dried under high vacuum to furnish 0.029 g (70%) of 3-(((3-carboxy-2-methylbenzofuran-5-yl)oxy)methyl)pyrrolidin-1-ium chloride (cpd 003).

Cpd 015 was prepared in a manner similar (use of appropriate reagents and purification methods known to the person skilled in the art) to cpd 019.

Synthesis of 2-methyl-N-(1-methylpiperidin-4-yl)-5-((5-(trifluoromethyl)furan-2-yl)methoxy)benzofuran-3-carboxamide (Cpd 069)



Step 1: HATU (235 mg; 0.62 mmol) was added to a solution of 2-methyl-5-((5-(trifluoromethyl)furan-2-yl)


398


methoxy)benzofuran-3-carboxylic acid (cpd 018) (140 mg; 0.41 mmol) and DIPEA (0.21 mL; 1.2 mmol) in DCM (5 mL). After 4 h at RT, 1-methylpiperidin-4-amine (I-7) (59 mg; 0.51 mmol) was added and the RM was stirred overnight. The RM was then diluted with DCM and washed successively with a sat. solution of sodium hydrogen carbonate and brine. The organic layer was dried over magnesium sulfate and filtered. The volatiles were removed under reduced pressure and the crude material was purified by FCC on silica gel using a gradient of MeOH (0% to 20%) in DCM to afford 128 mg (71%) of 2-methyl-N-(1-methylpiperidin-4-yl)-5-((5-(trifluoromethyl)furan-2-yl)methoxy)benzofuran-3-carboxamide (cpd 069).

The following compounds were prepared in a manner similar (use of appropriate reagents and purification methods (including chiral HPLC or chiral SFC) known to the person skilled in the art) to cpd 069:

Cpd 037, 039, 047, 049, 052, 054, 058, 059, 060, 061, 072, 078, 097, 098, 104, 107, 109, 117, 119, 128, 129, 132-En1, 132-En2, 133, 135-En1, 135-En 1, 139, 142, 143, 146, 148-En1, 148-En2, 150, 152, 154, 156, 158, 159-En1, 159-En2, 161, 167, racemic 170, 171, 178, 181, 192, 200, racemic 203, 204, 206, 270-En 1, 270-En 2, 274, 275, 308, 309, 312, 314, 315-En 1, 315-En 2, 316-En 1, 316-En 2, 317-En 1, 317-En 2, 322, 323, 331, 334, 337, 339, 341, 343, 344, 345, 346, 351, 352, 357, 358, 359, 370, 371, 372, 373, 374, 384, 385, 390, 391, 392, 393, 394, 399, 400, 401-En 1, 401-En 2, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 481, 482, 483, 484, 485, 486, 489-En 1, 489-En 2, 490-En 1, 490-En 2, 491-En 1, 491-En 2, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 504, 505, 506, 507, 523, 524, 525, 526, 527, 528, 529, 531-En 1, 531-En 2, 532-En 1, 532-En 2, 533-En 1, 533-En 2, 534-En 1, 534-En 2, 535-En 1, 535-En 2, 536-En 1, 536-En 2, 537, 574, 576, 602, 603, 606, 607, 608, 609, 610, 611, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, Cpd 623-En 1, Cpd 623-En 2, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 641-En1, 641-En2, 642-En1, 642-En2, 643-En1, 643-En2, 646-En1, 646-En 2, 647-En 1, 647-En 2, 648-En 1, 648-En 2, 649, 651, 652, 659, 660, 661-En 1, 661-En 2, 664, 665, 666, 668, 670, 672, 674, 676, 678, 680, 682, 683, 684, 685, 686, 688, 690, 692, 694, 695, 696, 697, 698, 700, 702, 704, 705, 706, 708, 710, 711, 712, 714, 715, 716, 718, 721, 722-En 1, 722-En 2, 723-En 1, 723-En 2, 724-En 1, 724-En 2, 726-En 1, 726-En 2, 727-En 1, 727-En 2, 727-En 3, 727-En 4, 738-En 1, 738-En 2, 739-En1, 739-En2, 741

Synthesis of 2-methyl-N-(2-oxopyrrolidin-3-yl)-5-(thiophen-3-ylmethoxy)benzofuran-3-carboxamide (Cpd 035)







Step 1: Oxalyl chloride (0.300 mL; 2.96 mmol) and DMF (2 drops) were added to a mixture of 2-methyl-5-(thiophen-3-ylmethoxy)benzofuran-3-carboxylic acid (Cpd 011) (100 mg; 0.35 mmol) in DCM (5 mL). The resulting mixture was stirred for 4 h. The RM was concentrated under reduced pressure to afford 2-methyl-5-(thiophen-3-ylmethoxy)benzofuran-3-carbonyl chloride.

Step 2: A solution of 2-methyl-5-(thiophen-3-ylmethoxy)benzofuran-3-carbonyl chloride dissolved in DCM (3 mL) was added to a mixture of 3-aminopyrrolidin-2-one (0.052 g; 0.520 mmol) and DIPEA (0.121 mL; 0.520 mmol). The RM was stirred at RT for 18 h. The solid formed was filtered off and purified by FCC on silica gel using a gradient of MeOH (0% to 20%) in DCM to afford 0.023 g (18%) 2-methyl-N-(2-oxopyrrolidin-3-yl)-5-(thiophen-3-ylmethoxy)benzofuran-3-carboxamide (cpd 035).

The following compounds were prepared in a manner similar (use of appropriate reagents and purification methods known to the person skilled in the art) to cpd 035:

Cpd 036, 080, 081, 082, 083, 084 and 085.

Synthesis of tert-butyl (S)-3-(2-methyl-5-((5-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate (Cpd 073)



Step 1: Tert-butyl (S)-3-aminopyrrolidine-1-carboxylate (1.1 g; 5.7 mmol) was added to a solution of 5-hydroxy-2-methylbenzofuran-3-carboxylic acid (1 g; 5.2 mmol), HATU (1.98 g; 5.2 mmol) and DIPEA (2.7 mL; 15.6 mmol) in DMF (10 mL). After 60 h, the reaction was concentrated under reduced pressure. The residue was partitioned between water and EtOAc. After separation, the aq. layer was extracted twice with EtOAc. Combined EtOAc extracts were dried over magnesium sulphate, filtered, and concentrated under reduced pressure. The residue was purified by FCC on silica gel using a gradient of MeOH (0 to 6%) in DCM to afford 0.94 g (50%) of tert-butyl-(S)-3-(5-hydroxy-2-methylbenzofuran-3-carboxamido)pyrrolidine-1-carboxylate. M/Z(+): 361 (M+H). M/Z(-): 359 (M-H). ¹H NMR (DMSO-d₆, 300 MHz) δ ppm: 9.26 (s, 1H), 8.17 (d, J=6.4 Hz, 1H), 7.32 (d, J=9.0 Hz, 1H), 6.99 (d, J=1.88 Hz, 1H), 6.70 (dd, J=8.9, 2.1 Hz, 1H), 4.35-4.50 (m, 1H), 3.51-3.3.65 (m, 1H), 3.35-3.49 (m, 1H), 3.15-3.31 (m, 1H), 2.53 (s, 3H), 2.02-2.19 (m, 1H), 1.84-1.99 (m, 1H), 1.41 (s, 9H).


Step 2: Tributylphosphine (0.155 mL; 0.583 mmol) was added dropwise to a stirred mixture of tert-butyl-(S)-3-(5-hydroxy-2-methylbenzofuran-3-carboxamido)pyrrolidine-1-carboxylate (0.150 g; 0.416 mmol), (5-methylpyridin-3-yl)methanol (0.081 g; 0.624 mmol) and ADDP (0.150 g; 0.583 mmol) in dry THF (5 mL) under argon. The mixture was stirred for 2 h and then the reaction was concentrated under reduced pressure. The residue was purified twice by FCC on silica gel using a gradient of MeOH (0-6%) in DCM for the first time and a gradient of MeOH (0-4%) in EtOAc for the second time to afford 0.113 g (58%) of tert-butyl (S)-3-(2-methyl-5-((5-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate (cpd 073).

The following compounds were prepared in a manner similar (use of appropriate reagents and purification methods known to the person skilled in the art) to cpd 073:

Cpd 026, 027, 030, 031, 032, 033, 034, 038, 040, 042, 043, 044, 046, 051, 057, 064, 065, 066, 077, 078; 086; 087; 088; 089; 090; 091; 092; 093; 094; 095 and 096.

401

(S)-2-methyl-5-((5-methylpyridin-3-yl)methoxy)-N-(pyrrolidin-3-yl)benzofuran-3-carboxamide (Cpd 028)




A solution of hydrogen chloride in 1,4-dioxane (4 M; 4 mL; 16 mmol) was added to a solution of tert-butyl (S)-3-((2-methyl-5-((5-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate (cpd 073) (0.105 g; 0.22 mmol) in DCM (3 mL). The mixture was stirred overnight at RT and was concentrated under reduced pressure. The residue was then purified by column of supported SiliaBond-PropylSulfonic Acid using a gradient of a solution of 3N ammonia in MeOH (0-100%) in DCM to afford 78 mg (95%) of (S)-2-methyl-5-((5-methylpyridin-3-yl)methoxy)-N-(pyrrolidin-3-yl)benzofuran-3-carboxamide (cpd 028).


The following compounds were prepared in a manner similar (use of appropriate reagents and purification methods known to the person skilled in the art) to cpd 028:

Cpd 020, 021, 022, 023, 034, 025, 029, 048, 050, 053, 055, 056, 062, 063, 067, 068, 070, 071, 074, 075 and 076.

Synthesis of 2-methyl-N-(1-methylpiperidin-4-yl)-5-(thiophen-2-ylmethoxy)benzofuran-3-carboxamide (Cpd 041)

**402**

-continued



Step 1: 5-(Benzyloxy)-2-methylbenzofuran-3-carboxylic acid (2.6 g; 9.2 mmol), HATU (5.2 g; 13.8 mmol; 1.5) and DIPEA (4.8 mL; 27.6 mmol) were dissolved in DCM (50 mL). After 4 h of stirring at RT, 1-methylpiperidin-4-amine (1.2 mL; 9.7 mmol) was added and the RM was stirred at RT overnight. The RM was diluted in 80 mL of DCM and washed with a sat. solution of sodium hydrogen carbonate and brine. The solvent was removed under reduced pressure. The residue was purified by FCC on silica gel using a gradient of MeOH (0-20%) in DCM to afford 1.4 g (48%) of 5-(benzyloxy)-2-methyl-N-(1-methylpiperidin-4-yl)benzofuran-3-carboxamide. ¹H NMR (DMSO d₆, 300 MHz) δ ppm 7.89 (d, 1H); 7.50-7.29 (m, 6H); 7.22 (d, 1H); 6.96 (dd, 1H); 5.13 (s, 2H); 3.85-3.65 (m, 1H); 2.75 (br d, 2H); 2.56 (s, 3H); 2.16 (s, 3H); 1.98 (ddd, 2H); 1.85-1.74 (m, 2H); 1.67-1.51 (m, 2H).



Step 2: A suspension of 5-(benzyloxy)-2-methyl-N-(1-methylpiperidin-4-yl)benzofuran-3-carboxamide (500 mg; 1.32 mmol) and Pd/C (70 mg, 0.065 mmol) in THF (13 mL) was stirred at RT under an atmospheric pressure of hydrogen until disappearance of the starting material. The RM was then filtered through a pad of Celite and the filtrate was evaporated under reduced pressure to afford 350 mg (92%) of 5-hydroxy-2-methyl-N-(1-methylpiperidin-4-yl)benzofuran-3-carboxamide, which was used without further purification. ¹H NMR (DMSO d₆, 300 MHz) δ (ppm): 9.25 (s, 1H); 7.84 (d, 1H); 7.31 (d, 1H); 6.99 (d, 1H); 6.69 (dd, 1H); 3.80-3.65 (m, 1H); 2.80-2.70 (m, 2H); 2.52 (s, 3H); 2.16 (s, 3H); 2.04-1.89 (m, 2H); 1.84-1.74 (m, 2H); 1.58 (ddd, 2H).

403

Step 3: To a cold solution of 5-hydroxy-2-methyl-N-(1-methylpiperidin-4-yl)benzofuran-3-carboxamide (75 mg, 0.26 mmol), thiophen-2-ylmethanol (45 mg; 0.39 mmol) and PS-PPH₃ (1.8 mmol/g; 289 mg; 0.52 mmol) in THF (3 mL) was slowly added a solution of DIAD (75 μ L; 0.39 mmol) in THF (0.8 mL). The RM was stirred at RT for 18 h and the volatiles were removed under reduced pressure. The crude material was purified by preparative HPLC (method H1) to furnish 8 mg (8%) of 2-methyl-N-(1-methylpiperidin-4-yl)-5-(thiophen-2-ylmethoxy)benzofuran-3-carboxamide (cpd 041).

Cpd 045 was prepared in a manner similar (use of appropriate reagents and purification methods known to the person skilled in the art) to cpd 041.

Synthesis of N-(4,4-difluoropyrrolidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide (Cpd 105)



Step 1: To a stirred solution cpd 101 (250 mg, 0.71 mmol) in DCM (50 mL), DIPEA (0.3 mL, 1.78 mmol), HATU (297 mg, 0.78 mmol) and tert-butyl 4-amino-3,3-difluoropyrrolidine-1-carboxylate (237 mg, 1.07 mmol) in DCM (1 mL) were added at RT. The RM was stirred for 18 h. Reaction progress was monitored by TLC. After completion of the reaction, the RM was diluted with DCM (300 mL), washed with sat. NaHCO₃ (50 mL), water (2x50 mL) and brine solution (50 mL), dried over anhydrous Na₂SO₄, filtered and concentrated. The residue was purified by Grace FCC using eluent 50% EtOAc-Pet ether to afford tert-butyl 3,3-difluoro-4-(2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate (350 mg, 88%).

Step 2: To a stirred solution of tert-butyl 3,3-difluoro-4-(2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)



404

benzofuran-3-carboxamido)pyrrolidine-1-carboxylate (350 mg, 0.63 mmol) in DCM (20 mL), TFA (1 mL) was added at 0° C. The RM was stirred for 18 h at RT. Reaction progress was monitored by TLC. The RM was concentrated and diluted with DCM (300 mL), washed with sat. NaHCO₃ (2x50 mL) and brine solution (50 mL), dried over anhydrous Na₂SO₄, filtered and concentrated. The residue was triturated with Et₂O (10 mL) to afford cpd 105 (250 mg, 77% over 2 steps). A preparative chiral SFC was performed to provide cpd 105-En 1 and cpd 105-En 2.

The following compounds were prepared in a manner similar (use of appropriate reagents and purification methods known to the person skilled in the art) to cpd 105:

Cpd 103-En1, 103-En2, 106-En1, 106-En2, 111-En1, 111-En2, 115-En1, 115-En2, 125-En1, 125-En2, 205-En1, 205-En2, 207-En1, 207-En2, 173, 189, 190, 310-En 1, 310-En 2, 313-En 1, 313-En 2, 336-En 1, 336-En 2, 338-En 1, 338-En 2, 342-En 1, 342-En 2, 538-En 1, 539-En 1, 539-En 2, 540-En 1, 540-En 2, 541-En 1, 541-En 2, 542-En 1, 542-En 2, 543-En 1, 543-En 2, 544-En 1, 544-En 2, 545-En 1, 545-En 2, 546-En 1, 547-En 1, 547-En 2, 548-En 1, 548-En 2, 549-En 1, 549-En 2, 550-En 1, 550-En 2, 551-En 1, 551-En 2, 552-En 1, 552-En 2, 553-En 1, 553-En 2, 554-En 1, 554-En 2, 555-En 1, 555-En 2, 556-En 1, 556-En 2, 557-En 1, 557-En 2, 558-En 1, 558-En 2, 559-En 1, 559-En 2, 560-En 1, 560-En 2, 561-En 1, 561-En 2, 562-En 1, 562-En 2, 563-En 1, 563-En 2, 564-En 1, 564-En 2, 565-En 1, 565-En 2, 566-En 1, 566-En 2, 567-En 1, 567-En 2, 568-En 1, 568-En 2, 569-En 1, 569-En 2, 570-En 1, 570-En 2, 571-En 1, 571-En 2, 572-En 1, 572-En 2, 573-En 1, 573-En 2, 635-En 1, 635-En 2, 636-En 1, 636-En 2, 637-En 1, 637-En 2, 638-En 1, 638-En 2, 639-En 1, 639-En 2, 663, 717.

Synthesis of N-(1-((dimethylamino)methyl)cyclobutyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 108)




Step 1: 37% Formaldehyde in water (0.21 mL, 2.5 mmol) was added to a pre-stirred solution of cpd 137 (100 mg, 0.26 mmol) in MeOH (10 mL) at 0° C. under argon atmosphere.

405

Then, glacial acetic acid (2 drops) and NaCNBH_3 (32.6 mg, 0.51 mmol) were added to the reaction. The RM was allowed to attain RT and stirred for 16 h. The reaction progress was monitored by TLC. The RM was poured into water (25 mL) and extracted with EtOAc (3×25 mL). The combined organic layer was washed with sat. NaHCO_3 (25 mL), water (25 mL) and brine (25 mL), dried over Na_2SO_4 and excess solvent was removed under reduced pressure. The crude product was purified by flash chromatography using 0-50% acetonitrile in 0.1% FA as an eluent followed by reverse phase prep-HPLC to afford cpd 108 (55 mg, 51%).

Cpd 140 was prepared in a manner similar (use of appropriate reagents and purification methods known to the person skilled in the art) to cpd 108:

Synthesis of N-(3,3-difluoropiperidin-4-yl)-4-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide (Cpd 112)



Step 1: 2-(Trifluoromethyl)pyridin-3-yl)methanol (2.23 g, 12.6 mmol), ADDP (2.96 g, 11.7 mmol) and tri-N-butylphosphine (2.37 g, 11.7 mmol) were added to a mixture of ethyl 4-fluoro-5-hydroxy-2-methylbenzofuran-3-carboxylate and ethyl 6-fluoro-5-hydroxy-2-methylbenzofuran-3-carboxylate (2.0 g, 8.4 mmol) in THF (50 mL) at 0° C. under argon atmosphere. The RM was allowed to attain RT and was stirred for 3 h. The reaction progress was monitored by TLC. The RM was poured into water (100 mL) and extracted with EtOAc (3×100 mL). The combined organic layer was washed with brine (2×100 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The crude product was purified by column chromatography over silica gel using 10-20% EtOAc in pet-ether as

406

an eluent followed by reverse phase prep-HPLC to afford a mixture of ethyl 4-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (300 mg, 8%) and ethyl 6-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (400 mg, 11%).

Step 2: 2N NaOH (10 mL) was added to a solution of ethyl 4-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (250 mg, 0.62 mmol) in a mixture of MeOH (15 mL) and THF (5 mL) at RT. The resulting RM was heated at 80° C. for 3 h. The reaction progress was monitored by TLC. The RM was cooled to RT, poured into ice cold water (50 mL) and acidified with 1 N HCl to pH~2. The crude product was extracted with EtOAc (3×50 mL), washed with water (100 mL), followed by brine (100 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford 4-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid (220 mg, 95%). The crude product was used for next step without purification.

Step 3: To a solution of a mixture of 4-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid (200 mg, 0.54 mmol) and tert-butyl 4-amino-3,3-difluoropiperidine-1-carboxylate (191 mg, 0.81 mmol) in DMF (15 mL) was added DIPEA (0.2 mL, 1.08 mmol) followed by HATU (412 mg, 1.08 mmol) at 0° C. under argon atmosphere. The RM was allowed to attain RT and stirred for 1 h. The reaction progress was monitored by TLC. The RM was diluted with ice cold water (50 mL) and filtered. The solid thus obtained was washed with water (50 mL) and dried under reduced pressure to afford tert-butyl 3,3-difluoro-4-(4-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)piperidine-1-carboxylate (250 mg, 78%).

407


408

Step 4: 4M HCl in dioxane (5 mL) was added dropwise to a solution of tert-butyl 3,3-difluoro-4-(4-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)piperidine-1-carboxylate (0.25 g, 0.42 mmol) in DCM (20 mL) at 0° C. under an argon atmosphere. The RM was warmed to RT and stirred for 5 h. The reaction progress was monitored by TLC. The excess solvents were evaporated in vacuo and the residue was cooled to 0° C., basified with sat. NaHCO₃ to pH ~9 and extracted with EtOAc (3x50 mL). The combined organic layer was washed with water (50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford cpd 112 (200 mg, 96%). A preparative chiral SFC was performed on the racemic mixture of cpd 112 to afford cpd 112-En 1 and cpd 112-En 2.

Synthesis of N-(3,3-difluoropiperidin-4-yl)-6-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide (Cpd 113) and N-(3,3-difluoropiperidin-4-yl)-7-fluoro-2-methyl-5-((2-(trifluoro-methyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide (Cpd 176)

409


410



411

412

-continued



413

Step 1: To a pre-stirred solution of CAN (27.41 g, 50.00 mmol) in water (60 mL) was added 2-fluorobenzene-1,4-diol (3.0 g, 23.80 mmol) at 0° C. The resulting RM was stirred at RT for 4 h. The reaction progress was monitored by TLC. The organic compound was extracted with Et₂O (3×50 mL). The combined organic layer was washed with brine (50 mL), dried over anhydrous Na₂SO₄ and filtered. The clear filtrate was passed through a silica gel column, eluted with diethyl ether and the collected fractions were evaporated in vacuo to afford 2-fluorocyclohexa-2,5-diene-1,4-dione (2.5 g, 84%).

Step 2: To a pre-stirred solution of 2-fluorocyclohexa-2,5-diene-1,4-dione (2.5 g, 20.16 mmol) in toluene (30 mL) was added ethyl 3-oxobutanoate (7.86 g, 60.48 mmol) followed by anhydrous ZnCl₂ (3.29 g, 24.19 mmol) at RT under argon atmosphere. The resulting RM was heated to reflux and maintained at this temperature for 16 h using a Dean-Stark apparatus. The reaction progress was monitored by TLC. The RM was cooled to RT, filtered through a celite pad and the celite pad was washed with EtOAc (70 mL). The combined clear filtrate was concentrated under reduced pressure. The crude product was purified by FCC over silica gel using 0-10% EtOAc in pet-ether as an eluent to afford a mixture of ethyl 7-fluoro-5-hydroxy-2-methylbenzofuran-3-carboxylate and ethyl 6-fluoro-5-hydroxy-2-methylbenzofuran-3-carboxylate (0.8 g, 21%).

Step 3: (2-(Trifluoromethyl)pyridin-3-yl)methanol (1.11 g, 6.30 mmol), ADDP (1.48 g, 5.88 mmol) and tri-N-butylphosphine (1.38 mL, 5.88 mmol) were added sequentially to a pre-stirred solution of mixture of ethyl 6-fluoro-5-hydroxy-2-methylbenzofuran-3-carboxylate and ethyl 7-fluoro-5-hydroxy-2-methylbenzofuran-3-carboxylate (1.0 g, 4.20 mmol) in THF (30 mL) at RT under argon atmosphere. The RM was stirred for 2 h at the same temperature. The reaction progress was monitored by TLC. The RM was poured into water (100 mL) and extracted with EtOAc (2×100 mL). The combined organic layer was sequentially washed with water (50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography over silica gel using 0-20% EtOAc in pet-ether as an eluent to afford a mixture of ethyl 6-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate and ethyl 7-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (1.1 g, 66%). The crude product was used for further step without purification.

Step 4: A solution of NaOH (0.44 g, 11.08 mmol) in water (8 mL) was added dropwise to a pre-stirred solution of a mixture of ethyl 6-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate and ethyl 7-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (1.1 g, 2.77 mmol) in a mixture of MeOH (15 mL) and THF (7 mL) at RT. The resulting RM was heated to 60° C. and maintained for 4 h. The reaction progress was monitored by TLC. The RM was cooled to RT, poured into ice cold water (50 mL) and acidified with 1 N HCl to a pH~2. The crude product was extracted with EtOAc (2×50 mL). The combined organic layer was washed with water (50 mL) followed by brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford a mixture of 6-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid and 7-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid (1.0 g, 99%). This crude product was used for next step without purification.


414

Step 5: To a pre-stirred solution of mixture of 6-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid and 7-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid (1.0 g, 2.71 mmol) and tert-butyl 4-amino-3,3-difluoropiperidine-1-carboxylate (0.74 g, 3.25 mmol) in DMF (20 mL) was added DIPEA (1.45 mL, 8.13 mmol) followed by HATU (2.05 g, 5.42 mmol) at 0° C. under an argon atmosphere. The RM was allowed to attain RT and stir for 16 h. The reaction progress was monitored by TLC. The RM was diluted with water (100 mL) and the organic compound was extracted with EtOAc (2×70 mL). The combined organic layer was washed with water (50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography over silica using 0-55% EtOAc and pet-ether as an eluent to afford a mixture of tert-butyl 3,3-difluoro-4-(6-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)piperidine-1-carboxylate and tert-butyl 3,3-difluoro-4-(7-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)piperidine-1-carboxylate (1.0 g, 62%).

Step 6: 4.0 M HCl in dioxane (13.62 mL, 13.62 mmol) was added dropwise to a pre-stirred solution of mixture of tert-butyl 4-(6-chloro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)-3,3-difluoropiperidine-1-carboxylate and tert-butyl 4-(7-chloro-2-methyl-5-((2-(trifluoro-methyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)-3,3-difluoropiperidine-1-carboxylate (1.0 g, 1.70 mmol) in DCM (15 mL) at 0° C. The RM was allowed to attain RT and stirred for 6 h. The reaction progress was monitored by TLC. The RM was concentrated under reduced pressure, the residue was basified with sat. NaHCO₃ (100 mL) and the organic compound was extracted with 10% MeOH in DCM (3×50 mL). The combined organic layer was washed with water (50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by GRACE flash chromatography using 0-60% acetonitrile and 0.1% FA in water as an eluent to afford a mixture of cpd 113 and cpd 176 (0.72 g, 87%). A preparative chiral SFC was performed on the mixture of racemic cpd 113 and racemic cpd 176 to provide cpd 113-En 1, cpd 113-En 2, cpd 176-En 1 and cpd 176-En 2.


The following compounds were prepared in a similar manner (use of appropriate reagents (chiral or racemic) and purification methods (including chiral HPLC or chiral SFC) known to the person skilled in the art) as described for the synthesis of cpd. 113 and cpd. 176: Cpd. 572-En 1, 572-En 2, 573-En 1, 573-En 2.

Synthesis of N-(1-(2-hydroxyethyl)cyclobutyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 114)





415

-continued




Step 1: To a pre-stirred solution of cpd 102 (0.3 g, 0.989 mmol) and 2-(1-aminocyclobutyl)ethan-1-ol (0.148 g, 1.285 mmol) in DMF (10 mL) was added DIPEA (0.675 mL, 3.959 mmol) followed by HATU (0.752 g, 1.978 mmol) at 0° C. under argon atmosphere. The RM was stirred for 1 h at same temperature. The reaction progress was monitored by LC-MS. The RM was diluted with sat. NaHCO₃ (80 mL) and then the crude product was extracted with EtOAc (2×50 mL). The combined organic layer was washed with water (30 mL) and brine (30 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by reverse phase prep-HPLC to afford cpd 114 (0.18 g, 45%)

Synthesis of 2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(1-(2,2,2-trifluoroethyl)pyrrolidin-3-yl)benzofuran-3-carboxamide (Cpd 116)



416

-continued




Step 1: To a solution of cpd 102 (600 mg, 1.98 mmol) in DCM (10 mL) at 0° C., DIPEA (1.77 mL, 9.90 mmol, 5 eq) was added dropwise at 0° C. and the RM stirred for 10 min. Tert-Butyl 3-aminopyrrolidine-1-carboxylate (552 mg, 2.97 mmol) and HATU (1.12 g, 2.97 mmol) were added to the RM. After completion of addition, the ice bath was removed and the RM was stirred at RT for 16 h. Reaction progress was monitored by TLC. The RM was extracted with DCM (2×100 mL), the combined organic layer was washed with water (100 mL) and brine (100 mL), dried over anhydrous Na₂SO₄, filtered, and concentrated to afford crude 2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (1.03 g).

Step 2: To a solution of 2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (1.0 g, 2.12 mmol) in DCM (15 mL) at 0° C., 4M aq. HCl (3 mL, 5.3 mmol) was added dropwise over 5 min at 0° C. and then the ice bath was removed. The RM was stirred at RT for 16 h. Reaction progress was monitored by TLC. The reaction was directly concentrated to get the crude product, which was washed with Et₂O (15 mL) to afford pure 2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(pyrrolidin-3-yl)benzofuran-3-carboxamide hydrochloride (700 mg, 94%).

Step 3: To a stirred solution of 2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(pyrrolidin-3-yl)benzofuran-3-carboxamide hydrochloride (700 mg, 1.71 mmol) in acetonitrile (20 mL), Cs₂CO₃ (328 mg, 5.13 mmol) and CF₃CH₂OTf (598 mg, 2.57 mmol) were added at RT. The resulting mixture was stirred for 3 h at RT. Reaction progress was monitored by TLC. After completion of the reaction, the RM was cooled to 0° C., quenched with ice-cold water (100 mL), and extracted with EtOAc (2×100 mL). The combined organic layer was washed with water (120 mL) and brine solution (120 mL), dried over anhydrous Na₂SO₄, and concentrated to afford a residue. This residue was purified by FCC using 80% EtOAc in pet ether as eluent to afford cpd 116 (280 mg, 74%). Cpd 116 was further purified by normal phase SFC-Prep to yield 190 mg (24%) of cpd 116. A preparative chiral SFC was performed on the racemic mixture of cpd 116 to afford cpd 116 En 1 and cpd 116 En 2.

417

Synthesis of N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide (Cpd 120)



Step 1: To a suspension of 60% NaH (314 g, 7.852 mmol) in DMF (10 mL) under argon atmosphere at 0° C., cpd 064 (850 mg, 1.963 mmol) in DMF (5 mL) and methyl 2-bromoacetate (450.4 mg, 2.944 mmol) were added at 0° C. The RM was stirred at RT for 3 h. Reaction progress was monitored by TLC. After completion of the reaction, the RM was diluted with ice-water (150 mL), extracted with EtOAc (3×300 mL), and the combined organic layer was washed with brine (100 mL), dried over anhydrous Na₂SO₄, filtered and concentrated. The residue was purified by FCC to afford the crude product (650 mg).


Step 2: To a solution of methyl 2-(3-(2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)pyrrolidin-1-yl)acetate (600 mg, 1.18 mmol) in MeOH (15 mL) under an argon atmosphere at 0° C., NaBH₄ (136.5 mg, 4.15 mmol) was added portionwise and then the solution was heated to 80° C. for 3 h. Reaction progress was monitored by TLC. After completion of the reaction, the RM was quenched with acetone (25 mL) and ice-water (150 mL), extracted with EtOAc (3×300 mL), and the combined organic layer was washed with brine solution (100 mL), dried over anhydrous Na₂SO₄, filtered and concentrated. The residue was purified by FCC to afford cpd 120 (280 mg, 30% over 2 steps). A preparative chiral SFC was performed on racemate cpd 120 to afford cpd 120-En 1 and cpd 120-En 2.

The following compounds were prepared in a manner similar (use of appropriate reagents and purification methods known to the person skilled in the art) to cpd 120-En 1 and cpd 120-En 2:

Cpd 134-En1, 134-En2, 151-En1 and 151-En2.

418

Synthesis of N-(3-(2-hydroxyethyl)tetrahydrofuran-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzo-furan-3-carboxamide (Cpd 121)



Step 1: To a stirred solution of dihydrofuran-3(2H)-one (5.0 g, 58.11 mmol) in toluene (60 mL), was added Ph₃P=CHC(O)OCH₂CH₃ (23.22 mL, 69.74 mmol), and then the RM was heated to 110° C. and stirred for 16 h. Reaction progress was monitored by TLC. After completion of the reaction, the RM was cooled to RT, and concentrated to yield the crude product, which was purified by Grace FCC using 20% EtOAc in Pet ether as eluent to afford ethyl (Z)-2-(dihydrofuran-3(2H)-ylidene)acetate (7.2 g, 80%).

Step 2: To a stirred solution of ethyl (Z)-2-(dihydrofuran-3(2H)-ylidene)acetate (5.0 g, 32.05 mmol) in toluene (60 mL), was added methanolic NH₃ (7M, 25 mL, 128.2 mmol), and then the RM was heated to 80° C. and stirred for 16 h in a steel bomb. The reaction progress was monitored by TLC. The RM was cooled to RT and concentrated to give the crude methyl 2-(3-aminotetrahydrofuran-3-yl)acetate (4.7 g, 92%).

Step 3: To a stirred solution of cpd 102 (800 mg, 2.64 mmol) in DCM (30 mL), were added DIPEA (1.89 mL, 10.56 mmol) and HATU (1.5 g, 3.96 mmol), followed by addition of methyl 2-(3-aminotetrahydrofuran-3-yl)acetate (629 mg, 3.96 mmol) at 0° C. and the RM was stirred for 2 h at RT. Reaction progress was monitored by TLC. After completion of the reaction, the RM was diluted with water (100 mL), and extracted with DCM (100 mL). The combined organic layer was washed with water (30 mL) and brine solution (25 mL), dried over anhydrous Na₂SO₄, filtered and concentrated to

419

afford the crude product which was purified by Grace FCC using 3% MeOH in DCM as eluent to afford methyl 2-(3-(2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)tetrahydrofuran-3-yl)acetate (650 mg, 55%).

Step 4: To a stirred solution of methyl 2-(3-(2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)tetrahydrofuran-3-yl)acetate (650 mg, 1.463 mmol) in THF (20 ml) at -30° C., was added LAH powder (61.19 mg, 1.61 mmol) portionwise. The RM was slowly warmed to RT and stirred for 2 h at RT. Reaction progress was monitored by TLC. After completion of the reaction, the RM was quenched with sat. Na₂SO₄ solution (100 mL), and extracted with EtOAc (100 ml). The combined organic layer was washed with water (30 ml) and brine solution (25 ml), dried over anhydrous Na₂SO₄, filtered and concentrated to


420

afford the crude product which was purified by Grace FCC using 6% MeOH in CH₂Cl₂ as eluent to afford cpd 121 (210 mg, 35%). A preparative chiral SFC was performed on racemic cpd 121 to provide cpd 121-En 1 and cpd 121-En 2.


Cpd 126-En 1 and cpd 126-En 2 was prepared in a manner similar (use of appropriate reagents and purification methods known to the person skilled in the art) to cpd 121-En 1 and cpd 121-En 2.

Synthesis of 7-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 122) and 6-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 182)

421



422



423

Step 1: (4-Methylthiazol-5-yl)methanol (0.65 g, 5.04 mmol), ADDP (1.18 g, 4.70 mmol) and tri-N-butylphosphine (1.15 mL, 4.70 mmol) were added sequentially to a pre-stirred solution of ethyl 7-fluoro-5-hydroxy-2-methylbenzofuran-3-carboxylate and ethyl 6-fluoro-5-hydroxy-2-methylbenzofuran-3-carboxylate (0.8 g, 3.36 mmol) in THF (30 mL) at 0° C. under an argon atmosphere. The RM was allowed to attain RT and stirred for 2 h. The reaction progress was monitored by TLC. The RM was poured into water (50 mL) and extracted with EtOAc (2×50 mL). The combined organic layer was washed with water (50 mL) and brine (50 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by FCC over silica gel using 0-20% EtOAc in pet-ether as an eluent to afford a mixture of ethyl 6-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate and ethyl 7-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate (0.7 g, 59%).

Step 2: A solution of NaOH (0.27 g, 6.87 mmol) in water (5 mL) was added to a pre-stirred solution of ethyl 6-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate compound and ethyl 7-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate (0.6 g, 1.71 mmol) in a mixture of MeOH (6 mL) and THF (3 mL) at RT. The resulting RM was heated to 60° C. and maintained at this temperature for 4 h. The reaction progress was monitored by TLC. The RM was cooled to RT, poured into ice cold water (50 mL), and acidified with 1 N HCl (pH~2). The crude product was extracted with EtOAc (2×50 mL). The combined organic layer was washed with water (2×50 mL), followed by brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford a mixture of 6-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (cpd 243) and 7-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (cpd 245) (0.6 g, 92%).

Step 3: To a pre-stirred solution of a mixture of 6-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid and 7-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (600 mg, 1.86 mmol), and methyl 4-aminotetrahydro-2H-pyran-4-carboxylate (0.356 g, 2.24 mmol) in DMF (10 mL) was added DIPEA (0.96 mL, 5.60 mmol) followed by HATU (1.41 g, 3.72 mmol) at 0° C. under an argon atmosphere. The RM was allowed to attain RT and stirred for 16 h. The reaction progress was monitored by TLC. The RM was diluted with water (50 mL) and then the organic compound was extracted with EtOAc (2×50 mL), and washed with water (50 mL) and brine (50 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography over silica gel using 0-55% EtOAc and pet ether as an eluent to afford a mixture of methyl 4-(6-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido) tetrahydro-2H-pyran-4-carboxylate and methyl 4-(7-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido) tetrahydro-2H-pyran-4-carboxylate (0.5 g, 57%).

Step 4: NaBH₄ (1.02 g, 27.05 mmol) was added portion-wise to a pre-stirred solution of a mixture of methyl 4-(6-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido) tetrahydro-2H-pyran-4-carboxylate and methyl 4-(7-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido) tetrahydro-2H-pyran-4-carboxylate (0.5 g, 1.08 mmol) in a mixture of MeOH (10 mL) and THF (5 mL) at 0° C. The RM was allowed to attain


424

RT and stirred for 16 h. The reaction progress was monitored by TLC. The RM was diluted with water (50 mL) and the organic compound was extracted with EtOAc (2×50 mL). The combined organic layer was washed with water (50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by GRACE flash chromatography using 0-60% acetonitrile and 0.1% FA in water as an eluent to afford a mixture of cpd 122 and cpd 182 (0.28 g, 59%). A preparative chiral SFC was performed on a mixture of cpd 111 and 182 to provide cpd 122 and cpd 182.

The following compounds were prepared in a similar manner (use of appropriate reagents (chiral or racemic) and purification methods (including chiral HPLC or chiral SFC) known to the person skilled in the art) as described in the synthesis of cpd. 112 and cpd. 182: Cpd. 360-En 1, 360-En 2, 361, 362, 364, 365-En 1, 365-En 2, 366, 367, 369, 587, 588.

The following compounds were prepared in a similar manner (use of appropriate reagents (chiral or racemic) and purification methods (including chiral HPLC or chiral SFC) known to the person skilled in the art) as described for step 3 in the synthesis of cpd. 243 and cpd. 245: Cpd. 244, 246.

Synthesis of N-(4,4-difluoro-1-methylpyrrolidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide (Cpd 124)



Step 1: To a stirred solution of cpd 105 (400 mg, 0.88 mmol) in MeOH (20 mL), 37% HCHO (0.5 mL) and acetic acid (0.1 mL) were added at 0° C. The RM was stirred for 30 min at 0° C. and then added NaCNBH₃ (109 mg, 1.76 mmol) at 0° C. The RM was stirred for 18 h at RT. Reaction progress was monitored by TLC. After completion of the reaction, the RM was quenched with water (50 mL). The product was extracted with EtOAc (3×100 mL), and the combined organic layer was washed with sat. NaHCO₃ (50 mL) and brine solution (50 mL), dried over anhydrous Na₂SO₄, filtered and concentrated. The residue was triturated with Et₂O (10 mL) to afford cpd 124 (370 mg, 90%).


425

A preparative chiral SFC was performed on the racemic mixture of cpd 124 to provide cpd 124-En 1 and cpd 124-En 2.

The following compounds were prepared in a manner similar (use of appropriate reagents and purification methods known to the person skilled in the art) to cpd 105:


Cpd 110-En1, 110-En2, 118-En1, 118-En2, 123-En1, 123-En2, 130-En1, 130-En2, 131-En1, 131-En2, 141-En1, 141-En2, 163-En1 and 163-En2.

Synthesis of N-(1-(2-hydroxyethyl)cyclobutyl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide (Cpd 127)




Step 1: To a pre-stirred solution of cpd 101 (0.52 g, 1.480 mmol) and 2-(1-aminocyclobutyl)ethan-1-ol (0.222 g, 1.924 mmol) in DMF (20 mL) was added DIPEA (1.01 mL, 5.920 mmol) followed by HATU (1.13 g, 2.96 mmol) at 0° C. under an argon atmosphere. The RM was stirred for 2 h at same temperature. The reaction progress was monitored by LC-MS. The RM was diluted with sat. NaHCO₃ (100 mL) and then the crude product was extracted with EtOAc (2x60 mL). The combined organic layer was washed with water (50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by reverse phase prep-HPLC to afford cpd 127 (0.110 g, 16%).

Synthesis of N-(1-(aminomethyl)cyclobutyl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide (Cpd 136)




426

-continued



A solution of N-(1-cyanocyclobutyl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide (450 mg, 1.048 mmol) in MeOH (10 mL) was added to a slurry of Raney-Ni (200 mg) in MeOH (10 mL), dropwise at RT. The RM was stirred under H₂ balloon pressure for 16 h. The RM was filtered through a celite pad and the celite pad was washed with MeOH (80 mL). The combined clear filtrate was concentrated under reduced pressure. The crude product was purified by reverse phase prep-HPLC to afford cpd 136 (0.3 g, 66%).

Synthesis of N-(1-(aminomethyl)cyclobutyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 137)




Step 1: To a solution of cpd 102 (2.0 g, 6.6 mmol) and 1-aminocyclobutane-1-carbonitrile (950 mg, 9.9 mmol) in DMF (20 mL) was added DIPEA (2.43 mL, 13.2 mmol) followed by HATU (5.0 g, 13.2 mmol) at 0° C. under argon atmosphere. The RM was allowed to attain RT and stirred for 1 h. The reaction progress was monitored by TLC. The RM was quenched with cold water (120 mL), the precipitated solid was filtered and dried under reduced pressure to

427

afford N-(1-cyanocyclobutyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (1.7 g, 68%).

Step 2: To a pre-stirred solution of N-(1-cyanocyclobutyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy) benzofuran-3-carboxamide (500 mg, 1.3 mmol) in a mixture of MeOH (10 mL) and THF (5 mL) was added $\text{COCl}_2 \cdot 6\text{H}_2\text{O}$ (468 mg, 1.96 mmol) followed by NaBH_4 (150 mg, 3.9 mmol) at 0°C . The RM was stirred for 30 min at same temperature. The reaction progress was monitored by LC-MS. The RM was poured into water (30 mL) and extracted with EtOAc (3 $\times$ 50 mL). The combined organic layer was washed with water (2 $\times$ 50 mL) and brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The crude product was purified by flash chromatography using 0-20% acetonitrile in 0.1% FA as an eluent, followed by reverse phase prep-HPLC to afford cpd 137 (80 mg, 16%).

Synthesis of 2-methyl-N-(1H-pyrazol-3-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide (Cpd 138)



Step 1: To a stirred solution of 1H-pyrazol-3-amine (5 g, 60.24 mmol) in DMF (50 mL), was added NaH (60% in oil) (4.819 g, 120.48 mmol) at 0°C . The RM was stirred for 30 min at 0°C and SEMCl (11.7 mL, 66.26 mmol) was added at 0°C . The RM was stirred for 1 h at RT. Reaction progress was monitored by TLC. After completion of the reaction, the RM was diluted with EtOAc (700 mL). The organic layer was washed with water (5 $\times$ 100 mL) and brine (100 mL), dried over anhydrous Na_2SO_4 , filtered and concentrated to give the crude product. The crude product was purified by Grace FCC using 100% EtOAc as eluent to afford 1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-3-amine (2.3 g, 18%).


Step 2: To a stirred solution of cpd 101 (200 mg, 0.56 mmol) in DCM (10 mL), was added DIPEA (0.3 mL, 1.71

428

mmol), HATU (325 mg, 0.85 mmol) and 1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-3-amine (182 mg, 0.85 mmol) at RT. The RM was stirred for 18 h at RT. Reaction progress was monitored by TLC. After completion of the reaction, the RM was diluted with DCM (200 mL), the organic layer was washed with water (2 $\times$ 50 mL), sat. NaHCO_3 (50 mL) and brine (50 mL), dried over anhydrous Na_2SO_4 , filtered and concentrated to give the crude product. The residue was purified by Grace FCC using 45% EtOAc in Pet ether as eluent to get 2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-N-(1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-3-yl)benzofuran-3-carboxamide (100 mg, 32%).

Step 3: To a stirred solution of 2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-N-(1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-3-yl)benzofuran-3-carboxamide (100 mg, 0.18 mmol) in DCM (10 mL), TFA (0.5 mL) was added at 0°C . The RM was stirred for 24 h at RT. Reaction progress was monitored by TLC. After completion of the reaction, the RM was poured into ice water (20 mL), basified to pH ~9 using sat. NaHCO_3 solution (20 mL), and the product was extracted with DCM (3 $\times$ 50 mL). The combined organic layer was washed with brine solution (50 mL), dried over anhydrous Na_2SO_4 , filtered and concentrated to give the crude product. The residue was purified by Grace FCC using 100% EtOAc as eluent to afford cpd 138. The product was purified further by HPLC.

Synthesis of 2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(1-(((2,2,2-trifluoroethyl)amino)methyl)cyclobutyl)-benzofuran-3-carboxamide (Cpd 145)




Step 1: 2,2,2-Trifluoroacetaldehyde monohydrate (77% in water, 0.85 mL, 5.1 mmol) was added to a solution of cpd 137 (200 mg, 0.51 mmol) in MeOH (15 mL) at 0°C under an argon atmosphere. Then, glacial acetic acid (3 drops) and NaCNBH_3 (65 mg, 1.0 mmol) were added to the reaction. The resulting RM was heated at 80°C for 16 h. The RM was cooled to RT, poured into water (50 mL) and extracted with EtOAc (2 $\times$ 50 mL). The combined organic layer was washed with sat. NaHCO_3 (25 mL), water (50 mL) and brine (50 mL), dried over Na_2SO_4 and excess solvent was removed

429


under reduced pressure. The crude product was purified by flash chromatography using 0-50% acetonitrile in 0.1% FA as an eluent followed by reverse phase prep-HPLC to afford cpd 145 (30 mg, 12%).

Cpd 153 was prepared in a manner similar (use of appropriate reagents and purification methods known to the person skilled in the art) to cpd 145.

Synthesis of N-(4-hydroxytetrahydrofuran-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide (Cpd 147)



Step 1: To a stirred solution of 3,6-dioxabicyclo[3.1.0]hexane (0.2 g, 2.27 mmol) in EtOH (10 ml) in a 100 mL steel bomb was added aq. NH_4OH (5 ml), and the steel bomb was closed tightly. The RM was heated up to 80° C. for 18 h in the steel bomb. Reaction progress was monitored by TLC. After completion of the reaction, the RM was concentrated to give the crude product, and the crude product was co-distilled with toluene three times to give 4-aminotetrahydrofuran-3-ol (0.16 g, 65%).




Step 2: To a solution of cpd 101 (0.35 g, 0.99 mmol) in DCM (20 ml), were added DIPEA (0.91 ml, 4.98 mmol) and HATU (0.568 g, 1.49 mmol), at 0° C., followed by a solution of 4-aminotetrahydrofuran-3-ol (0.151 g, 1.49 mmol) in DCM, dropwise at 0° C. Then the RM was stirred for 3 h at RT, and the reaction monitored by TLC. After completion of the reaction, the RM was diluted with water (20 ml), extracted with DCM (3×30 ml), and the combined organic layer was washed with water (50 ml) and brine solution (25 ml), dried over anhydrous Na_2SO_4 , filtered and concentrated to yield the crude product. The crude product was purified by FCC using 60% EtOAc-pet ether as eluent to afford cpd 147 (0.30 g, 69%). A preparative chiral SFC was performed on racemic cpd 147 to afford cpd 147 Dia 1-En 1 and cpd 147 Dia 1-En 2.

430

Cpd 162-Dia 1-En 1, 162-Dia 1-En 2, 386-En 1, 386-En 2, 387-En 1, 387-En 2, 388-En 1, 388-En 2, 389-En 1, 389-En 2, 653-En 1, 653-En 2, 654-En 1, 654-En 2, 655-En 1, 655-En 2, 656-En 1, 656-En 2 were prepared in a manner similar (use of appropriate reagents and purification methods known to the person skilled in the art) to cpd 147-Dia 1-En 1 and cpd 147-Dia 1-En 2.

Synthesis of 2-methyl-N-(1-(2,2,2-trifluoroethyl)pyrrolidin-3-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide (Cpd 149)



Step 1: To a stirred solution of cpd 101 (400 mg, 1.14 mmol) in DCM (15 mL), DIPEA (0.5 mL, 2.85 mmol) and HATU (476 mg, 1.25 mmol) were added at RT. The RM was stirred for 5 min at RT and tert-butyl 3-aminopyrrolidine-1-carboxylate (233 mg, 1.25 mmol) was added at RT. The RM was stirred for 18 h at RT. Reaction progress was monitored by TLC. After completion of the reaction, the RM was diluted with DCM (200 mL). The organic layer was washed with sat. NaHCO_3 solution (100 mL), water (100 mL) and brine (100 mL), dried over anhydrous Na_2SO_4 ,


431

filtered and concentrated to give a residue. The residue was purified by Grace FCC using 45% EtOAc in Pet ether as eluent to afford tert-butyl 3-(2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate (450 mg, 76%).

Step 2: To a stirred solution of tert-butyl 3-(2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate (450 mg, 0.86 mmol) in DCM (10 ml), 4N HCl in dioxane (5 mL) was added at 0° C. The RM was stirred for 18 h at RT. Reaction progress was monitored by TLC. After completion of the reaction, the RM was concentrated and dried to afford 2-methyl-N-(pyrrolidin-3-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide (520 mg, crude product).


Step 3: To a stirred solution of 2-methyl-N-(pyrrolidin-3-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide (500 mg, 1.02 mmol) in CH₃CN (50 ml), Cs₂CO₃ (1.33 g, 4.07 mmol) and CF₃CH₂OTf (472 mg, 2.03 mmol) were added at RT. The RM was heated to 50° C. and stirred for 1 h at 50° C. Reaction progress was monitored by LCMS. After completion of the reaction, the RM was cooled to RT, diluted with EtOAc (250 mL), washed with water (2×50 mL) and brine solution (50 mL), dried over anhydrous Na₂SO₄, filtered and concentrated to give a residue. The residue was purified by RP-Prep HPLC to afford cpd 149 (280 mg, 64% over 2 steps). A preparative chiral SFC was performed on racemic cpd 149 to provide cpd 149-En 1 and cpd 149-En 2.

Synthesis of N-(4,4-difluoro-1-(2,2,2-trifluoroethyl)pyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 155) and 2,2,2-trifluoroethyl 3,3-difluoro-4-(2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate (Cpd 172)



432


-continued



Step 1: To a solution of cpd 106 (400 mg, 0.982 mmol) in DMF (20 ml), Cs₂CO₃ (958 mg, 2.948 mmol) was added at RT. The RM was stirred for 5 min at RT and then CF₃CH₂OTf (342 mg, 1.474 mmol) was added at RT. The RM was stirred for 16 h at RT. Reaction progress was monitored by TLC. After completion of the reaction, the mixture was diluted with water (100 ml) and extracted with EtOAc (3×100 mL). The combined organic layer was washed with brine (50 mL), dried over anhydrous Na₂SO₄, filtered and concentrated. The residue was purified by RP-Prep HPLC to afford a mixture of cpd 155 and cpd 172 (320 mg, 45%). A preparative chiral SFC was performed on the mixture of racemic cpd 155 and racemic cpd 152 to afford cpd 155-En 1, cpd 155-En 2, cpd 172-En 1 and cpd 172-En 2.

Racemic Cpd 175 was prepared in a manner similar (use of appropriate reagents and purification methods known to the person skilled in the art) to cpd 155-En 1, cpd 155-En 2, cpd 172-En1 and cpd 172-En 2.


Synthesis of N-(4,4-difluoro-1-(2,2,2-trifluoroethyl)pyrrolidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide (Cpd 157)



433

Step 1: To a stirred solution of cpd 105 (600 mg, 1.318 mmol) in DMF (20 ml), NaH (56.9 mg, 3.954 mmol) was added at 0° C. The RM was stirred for 5 min at 0° C. and then CF₃CH₂OTf (609 mg, 2.637 mmol) was added at 0° C. The reaction was stirred for 18 h at RT. Reaction progress was monitored by TLC. After completion of the reaction, the mixture was diluted with ice water (50 mL) and extracted with EtOAc (3×100 mL). The combined organic layer was washed with brine (50 mL), dried over anhydrous Na₂SO₄, filtered and concentrated. The residue was purified by RP-Prep HPLC to afford cpd 157 (170 mg, 24%). A preparative chiral SFC was performed on racemic cpd 157 to afford cpd 157-En 1 and cpd 157-En 2.

Synthesis of N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzo-furan-3-carboxamide (Cpd 160)




434

Step 1: To a suspension of 60% NaH (240 mg, 6.21 mmol) in DMF (5.0 ml), cpd 042 (800 mg, 2.07 mmol) in DMF (10 ml) and (2-bromoethoxy)(tert-butyl)dimethylsilane (980 mg, 4.14 mmol) was added at 0° C. and the RM was allowed to rise to RT and was stirred for 16 h. Reaction progress was monitored by TLC. The RM was extracted with DCM (2×100 ml), the combined organic layer was washed with water (50 mL) and brine solution (50 mL), and then dried over anhydrous Na₂SO₄, filtered, and concentrated to afford N-(1-(2-((tert-butyldimethylsilyl)oxy)ethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzo-furan-3-carboxamide (850 mg, crude).

Step 2: To a solution of N-(1-(2-((tert-butyldimethylsilyl)oxy)ethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzo-furan-3-carboxamide (850 mg, 1.63 mmol) in THF (15 ml) at 0° C., TBAF (2.45 ml, 2.45 mmol), was added dropwise at 0° C. and the RM was stirred at RT for 2 h. Reaction progress was monitored by TLC. The RM was extracted with EtOAc (2×100 ml), the combined organic layer was washed with water (150 mL) and brine solution (150 mL); dried over anhydrous Na₂SO₄, filtered, and concentrated to afford the crude product, which was purified by FCC to afford the desired product cpd 160 (190 mg). A preparative chiral SFC was performed on the racemic mixture of cpd 160 to provide cpd 160-En 1 and cpd 160-En 2. Cpd 402 was isolated as a byproduct that occurred during the synthetic sequence.


The following compounds were prepared in a similar manner (use of appropriate reagents (chiral or racemic) and purification methods (including chiral HPLC or chiral SFC) known to the person skilled in the art) as described for cpd 160: Cpd. 403-En 1, 403-En 2, 404-En 1, 404-En 2, 405-En 1, 405-En 2, 409-En 1, 409-En 2, 410-En 1, 410-En 2,

Synthesis of N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((2-methylpyridin-3-yl)methoxy)benzo-furan-3-carboxamide (Cpd 164)



435

-continued



Step 1: (2-Methylpyridin-4-yl)methanol (1.8 g, 13.63 mmol), ADDP (3.2 g, 13.63 mmol) and tri-N-butyl phosphine (50% in EtOAc) (3.1 mL, 12.72 mmol) were added to a stirred solution of ethyl 5-hydroxy-2-methylbenzofuran-3-carboxylate (2.0 mg, 9.09 mmol) in THF (30 mL) at 0° C. under argon atmosphere. The resulting RM was stirred at rt for 24 h. The reaction progress was monitored by TLC. The RM was poured into water (100 mL) and extracted with EtOAc (2×100 mL). The combined organic layer was washed with brine (100 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by flash chromatography using 0.1% FA and acetonitrile as eluent to afford ethyl 2-methyl-5-((2-methylpyridin-4-yl)methoxy)benzofuran-3-carboxylate (1.2 g, 40%).

Step 2: A solution of 2-methyl-5-((2-methylpyridin-4-yl)methoxy)benzofuran-3-carboxylate (1.2 g, 5.08 mmol) and 4 M NaOH (10 mL) solution in MeOH:THF (1:1) (24 mL) was heated to 80° C. and maintained for 2 h. The reaction progress was monitored by TLC. The RM was cooled to RT, poured into ice cold water (100 mL), acidified with 1 N HCl and extracted with EtOAc (2×100 mL). The combined organic layer was washed with brine (100 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford 2-methyl-5-((2-methylpyridin-4-yl)methoxy)benzofuran-3-carboxylic acid (800 mg, 72%).

Step 3: To a stirred solution of 3-amino-1-(2-hydroxyethyl) pyrrolidin-2-one hydrochloride (0.362 g, 2.02 mmol) in DMF (10 mL) were added DIPEA (2.4 mL, 13.46 mmol) followed by 2-methyl-5-((2-methylpyridin-4-yl)methoxy)benzofuran-3-carboxylic acid (0.5 g, 1.68 mmol) and HATU (1.0 g, 3.36 mmol) sequentially at 0° C. under an argon atmosphere. The resulting RM was allowed warm to RT and stirred for 16 h. The reaction progress was monitored by TLC. The RM was diluted with ice cold water (100 mL) and extracted with 10% MeOH in DCM (2×50 mL). The combined organic layer was washed with brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by flash chromatography using 0.1% FA and acetonitrile as eluent to afford cpd 164. A preparative chiral SFC was performed on racemic cpd 164 to afford cpd 164-En 1 and cpd 164-En 2.


The following compounds were prepared in a manner similar (use of appropriate reagents and purification methods known to the person skilled in the art) to cpd 164-En 1 and cpd 164-En 2: cpd 169-En 1, 169-En 2, 180-En 1 and 180-En 2.

The following compounds were prepared in a manner similar (use of appropriate reagents and purification methods known to the person skilled in the art) to step 3 in the preparation of cpd 164-En 1 and cpd 164-En 2 req: Cpd.

436

335-En1, 335-En2, 340-En1, 340-En2, 348-En1, 348-En2, 406-En1, 406-En2, 407-En 1, 407-En 2, 408-En 1, 408-En 2.

Synthesis of 2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)benzofuran-3-carboxamide (Cpd 166)



Step 1: To a stirred solution of cpd 102 (500 mg, 1.65 mmol) in DCM (50 mL), were added DIPEA (1.4 mL, 8.25 mmol) and HATU (940 mg, 2.47 mmol), followed by tert-butyl 4-aminopiperidine-1-carboxylate (495 mg, 2.47 mmol) and the RM was stirred for 2 h at RT. Reaction progress was monitored by TLC. The RM was diluted with water (100 mL), extracted with EtOAc (3×50 mL), washed with water (50 mL) and brine (30 mL), dried over anhydrous Na₂SO₄, filtered and concentrated to give the crude product. The crude product was purified by Grace FCC using 20% EtOAc in Pet ether as eluent to afford tert-butyl 4-(2-methyl-

437

5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)piperidine-1-carboxylate (600 mg, 75%).


Step 2: To a solution of tert-butyl 4-(2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)piperidine-1-carboxylate (600 mg, 1.237 mmol) in DCM (80 ml) at 0° C., 4N HCl in dioxane (2.3 ml, 9.277 mmol) was added dropwise over 5 min at 0° C., and the ice bath was removed. The RM was stirred at RT for 16 h. Reaction progress was monitored by TLC. The RM was directly concentrated to afford the crude product, which was washed with Et₂O (15 ml) to afford 2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(piperidin-4-yl)benzofuran-3-carboxamide hydrochloride (700 mg crude).

438

(120 ml) and brine (120 ml), dried over anhydrous Na₂SO₄, and concentrated to afford a residue, which was purified by Grace FCC using 80% EtOAc in pet ether as eluent to afford cpd 166 (150 mg, 14%).

Cpd 168 was prepared in a manner similar (use of appropriate reagents and purification methods known to the person skilled in the art) to cpd 166.

Synthesis of 4-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)-methoxy)benzofuran-3-carboxamide (Cpd 177)



Step 3: To a stirred solution of 2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(piperidin-4-yl)benzofuran-3-carboxamide (700 mg, 1.71 mmol) in acetonitrile (40 ml), Cs₂CO₃ (167 mg, 5.13 mmol) and CF₃CH₂OTf (595 mg, 2.57 mmol) were added at RT. The resulting RM was stirred for 4 h at RT. Reaction progress was monitored by TLC. After completion of the reaction, the RM was quenched with ice-cold water (100 ml), and extracted with EtOAc (2×100 ml). The combined organic layer was washed with water

Step 1: To a solution of ethyl 5-hydroxy-2-methylbenzofuran-3-carboxylate (5.0 g, 22.7 mmol) in acetonitrile (300 mL) was added selectfluor (9.65 g, 27.2 mmol) at RT under argon atmosphere. The resulting RM was stirred for 16 h at same temperature. The reaction progress was monitored by TLC. The excess solvent was removed under reduced pressure and the crude compound was dissolved in EtOAc (500 mL). The above solution was washed with water (2×250 mL) and brine (250 mL), dried over anhydrous Na₂SO₄ and

439

concentrated under reduced pressure. The crude product was purified by FCC over silica using 0-20% EtOAc in pet-ether as an eluent followed by a flash chromatography using 0-47% of acetonitrile in 0.1% FA in water as an eluent to afford a mixture of ethyl 4-fluoro-5-hydroxy-2-methylbenzofuran-3-carboxylate and ethyl 6-fluoro-5-hydroxy-2-methylbenzofuran-3-carboxylate (1.0 g, 19%).

Step 2: (4-Methylthiazol-5-yl)methanol (1.23 g, 9.5 mmol), ADDP (2.81 g, 11.1 mmol) and tri-N-butylphosphine (2.75 mL, 11.1 mmol) were added sequentially to a solution of a mixture of ethyl 4-fluoro-5-hydroxy-2-methylbenzofuran-3-carboxylate and ethyl 6-fluoro-5-hydroxy-2-methylbenzofuran-3-carboxylate (1.9 g, 7.98 mmol) in THF (100 mL) at 0° C. under an argon atmosphere. The RM was allowed to attain RT and stir for 1 h. The reaction progress was monitored by TLC. The RM was poured into water (200 mL) and extracted with EtOAc (2×250 mL). The combined organic layer was washed with water (200 mL) and brine (200 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by FCC over silica using 0-30% EtOAc in pet-ether as an eluent to afford a mixture of ethyl 4-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate and ethyl 6-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate (1.5 g, 54%).

Step 3: 2N NaOH in water (20 mL) was added to a solution of a mixture of ethyl 4-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate and ethyl 6-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate (0.8 g, 2.29 mmol) in a mixture of MeOH (20 mL) and THF (5 mL) at RT. The resulting RM was heated to 80° C. for 3 h. The reaction progress was monitored by TLC. The RM was cooled to RT, poured into ice cold water (100 mL), and acidified with 1 N HCl (pH~2). The crude product was extracted with EtOAc (3×100 mL). The combined organic layer was washed with water (2×100 mL) followed by brine (100 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by flash chromatography using 0-50%


440

of acetonitrile in 0.1% FA in water as eluent to afford 4-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (cpd 233) (280 mg, 38%). The corresponding undesired acid from was separately collected.

Step 4: To a pre-stirred solution of 4-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (250 mg, 0.77 mmol) and methyl 4-aminotetrahydro-2H-pyran-4-carboxylate (185.7 mg, 1.10 mmol) in DMF (5 mL) was added DIPEA (0.28 mL, 1.55 mmol) followed by HATU (592 mg, 1.55 mmol) at 0° C. under argon atmosphere. The RM was allowed to attain RT and stirred for 1 h. The reaction progress was monitored by TLC. The RM was quenched with cold water (20 mL) and the solid was filtered and dried under reduced pressure to afford methyl 4-(4-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)tetrahydro-2H-pyran-4-carboxylate (260 mg, 74%).

Step 4: NaBH₄ (507 mg, 13.5 mmol) was added portionwise to a pre-stirred solution of methyl 4-(4-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)tetrahydro-2H-pyran-4-carboxylate (250 mg, 0.54 mmol) in a mixture of MeOH (10 mL) and THF (10 mL) at 0° C. The resulting RM was heated to 50° C. and maintained at this temperature for 16 h. The reaction progress was monitored by TLC. The RM was cooled to RT, diluted with water (50 mL) and the organic compound was extracted with EtOAc (2×50 mL). The combined organic layer was washed with water (50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by GRACE flash chromatography using 0-50% acetonitrile and 0.1% FA in water as an eluent to afford cpd 177 (50 mg, 21%).


Synthesis of N-(3,3-difluoropiperidin-4-yl)-2,6-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide (Cpd 179) and N-(3,3-difluoropiperidin-4-yl)-2,7-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide (Cpd 183)



441

442

-continued



Step 1: To a solution of 2-methylcyclohexa-2,5-diene-1,4-dione (20.0 g, 163.93 mmol) in toluene (500 mL) was added ethyl 3-oxobutanoate (63.9 g, 491.79 mmol) and anhydrous ZnCl_2 (26.8 g, 196.71 mmol) at RT under argon atmosphere. The resulting RM was refluxed for 16 h using a Dean-Stark apparatus. The reaction progress was monitored by TLC. The RM was cooled to RT, filtered through a celite pad and the celite pad was washed with EtOAc (300 mL). The filtrate was concentrated under reduced pressure. The crude product was purified by FCC over silica using 0-10% EtOAc in pet-ether as an eluent to afford a mixture of ethyl 5-hydroxy-2,7-dimethylbenzofuran-3-carboxylate and ethyl 5-hydroxy-2,6-dimethylbenzofuran-3-carboxylate (20.0 g, 52%).

Step 2: (2-(Trifluoromethyl)pyridin-3-yl)methanol (11.35 g, 64.10 mmol), ADDP (15.0 g, 59.82 mmol) and tri-N-butylphosphine (14.75 mL, 59.82 mmol) were added sequentially to a solution of mixture of ethyl 5-hydroxy-2,7-dimethylbenzofuran-3-carboxylate and ethyl 5-hydroxy-2,6-dimethylbenzofuran-3-carboxylate (10.0 g, 42.73 mmol) in THF (200 mL) at 0° C. under argon atmosphere. The RM was allowed to attain RT and was stirred for 3 h. The reaction progress was monitored by TLC. The RM was poured into water (200 mL) and extracted with EtOAc

(2×200 mL). The combined organic layer was washed with water (200 mL) and brine (200 mL). The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The crude product was purified by FCC over silica using 0-20% EtOAc in pet-ether as an eluent to afford a mixture of ethyl 2,6-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate and ethyl 2,7-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (8.0 g, 47%).

Step 3: A solution of NaOH (2.03 g, 50.89 mmol) in water (50 mL) was added to a solution of mixture of ethyl 2,6-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate and ethyl 2,7-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (5.0 g, 12.72 mmol) in a mixture of MeOH (50 mL) and THF (50 mL) at RT. The RM was heated to 60° C. for 3 h. The reaction progress was monitored by TLC. The RM was cooled to RT and poured into ice cold water (200 mL), then acidified with 1 N HCl (pH~2). The crude product was extracted with EtOAc (2×100 mL). The combined organic layer was washed with water (2×100 mL) followed by brine (100 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford a mixture of 2,6-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-

443

carboxylic acid and 2,7-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid (3.5 g, 76%). The crude product thus obtained was used for next step without further purification.

Step 4: To a solution of a mixture of 2,6-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxylic acid and 2,7-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid (2.0 g, 5.47 mmol) in DMF (50 mL) were added tert-butyl 4-amino-3,3-difluoropiperidine-1-carboxylate (1.5 g, 6.57 mmol), DIPEA (2.9 mL, 16.41 mmol) and HATU (4.1 g, 10.94 mmol) at 0° C. under argon atmosphere. The RM was allowed to attain RT for 3 h. The reaction progress was monitored by TLC. The RM was diluted with water (100 mL) and the crude product was extracted with EtOAc (2×100 mL). The combined organic layer was washed with water (100 mL) and brine (100 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography over silica using 0-55% EtOAc and pet- ether as an eluent to afford a mixture of tert-butyl 4-(2,6-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)-3,3-difluoropiperidine-1-carboxylate and tert-butyl 4-(2,7-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)-3,3-difluoropiperidine-1-carboxylate (1.5 g, 48%).


Step 5: TFA (1.5 mL) was added dropwise to a pre-stirred solution of a mixture of tert-butyl 4-(2,6-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)-3,3-difluoropiperidine-1-carboxylate and tert-

444


butyl 4-(2,7-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)-3,3-difluoropiperidine-1-carboxylate (1.0 g, 1.71 mmol) in DCM (10 mL) at 0° C. under an argon atmosphere. The RM was allowed to attain RT and stirred for 16 h. The reaction progress was monitored by TLC. The RM was concentrated under reduced pressure and basified with sat. NaHCO₃ solution (pH~8). The crude product was extracted with EtOAc (2×100 mL). The combined organic layer was washed with water (100 mL), followed by brine (100 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by GRACE flash chromatography using 0-50% acetonitrile and 0.1% FA in water as an eluent to afford a mixture of cpd 179 and cpd 183 (450 mg, 84%). A preparative chiral SFC was performed on the racemate cpd 179 and the racemate cpd 183 to provide cpd 179-En 1, cpd 179-En 2, cpd 183-En 1 and cpd 183-En 2.

The following compounds were prepared in a manner similar (use of appropriate reagents and purification methods known to the person skilled in the art) to cpd 179-En 1, cpd 179-En 2, cpd 183-En 1 and cpd 183-En 2. Cpd: 165-En1, 165-En2, 184-En1, 184-En2, 174 and 187.


Synthesis of N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2,6-dimethyl-5-((4-methylthiazol-5-yl)methoxy)-benzofuran-3-carboxamide (Cpd 185) and N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2,7-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 186)



445



446




Step 1: (4-Methylthiazol-5-yl)methanol (8.26 g, 64.10 mmol), ADDP (15.0 g, 59.82 mmol) and tri-N-butylphosphine (14.75 mL, 59.82 mmol) were added sequentially to a pre-stirred solution of a mixture of ethyl 5-hydroxy-2,7-dimethylbenzofuran-3-carboxylate and ethyl 5-hydroxy-2,6-dimethylbenzofuran-3-carboxylate (10.0 g, 42.73 mmol) in THF (200 mL) at 0° C. under an argon atmosphere. The RM was allowed to attain RT and was stirred for 3 h. The reaction progress was monitored by TLC. The RM was poured into water (200 mL) and extracted with EtOAc (2×200 mL). The combined organic layer was washed with water (200 mL) and brine (200 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by FCC over silica gel using 0-20% EtOAc in pet-ether as an eluent to afford a mixture of ethyl 2,7-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate and ethyl 2,6-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate (8.0 g, 57%).


Step 2: A solution of NaOH (0.927 g, 23.18 mmol) in water (14 mL) was added to a pre-stirred solution of a mixture of ethyl 2,7-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate and ethyl 2,6-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate (2.0 g, 5.79 mmol) in a mixture of MeOH (20 mL) and THF (20 mL) at RT. The resulting RM was heated to 60° C. and maintained for 3 h. The reaction progress was monitored by TLC. The RM was cooled to RT, poured into ice cold water (100 mL) and acidified with 1 N HCl (pH~2). The crude product was extracted with EtOAc (2×100 mL). The combined organic layer was washed with water (2×100 mL),

followed by brine (100 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford a mixture of 2,7-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid and 2,6-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (1.6 g, 89%). The crude product thus obtained was used for next step without further purification.

Step 3: To a pre-stirred solution of a mixture of 2,7-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3- and 2,6-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (1.0 g, 3.15 mmol) and (4-amino-tetrahydro-2H-pyran-4-yl)methanol (0.619 g, 4.73 mmol) in DMF (10 mL) was added DIPEA (1.69 mL, 9.45 mmol), followed by HATU (2.39 g, 6.30 mmol) at 0° C. under an argon atmosphere. The RM was allowed to attain RT and stirred for 3 h. The reaction progress was monitored by TLC. The RM was diluted with water (100 mL), and the crude product was extracted with EtOAc (2×100 mL). The combined organic layer was washed with water (100 mL) and brine (100 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography over silica gel using 0-55% EtOAc and pet-ether as an eluent to afford a mixture of cpd 186 and cpd 185 (0.5 g, 57%). A preparative chiral SFC was performed on a mixture of cpd 186 and 185 to provide cpd 186 and cpd 185.

Synthesis of 7-cyano-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 188)





Step 1: (4-Methylthiazol-5-yl)methanol (0.47 g, 3.67 mmol), ADDP (0.64 g, 3.42 mmol) and tri-N-butylphosphine (0.80 mL, 3.42 mmol) were added sequentially to a pre-stirred solution of ethyl 6-cyano-5-hydroxy-2-methylbenzofuran-3-carboxylate and ethyl 7-cyano-5-hydroxy-2-methylbenzofuran-3-carboxylate (0.6 g, 2.44 mmol) in THF (30 mL) at 0° C. under argon atmosphere. The RM was allowed to attain RT and stirred for 2 h. The reaction progress was monitored by TLC. The RM was poured into water (100 mL) and extracted with EtOAc (2×100 mL). The combined organic layer was washed with water (50 mL) and brine (50 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography over silica gel using 0-20% EtOAc in pet-ether as an eluent to afford a mixture of ethyl 6-cyano-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate and ethyl 7-cyano-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate (0.6 g, 68%).

Step 2: A solution of NaOH (0.26 g, 6.74 mmol) in water (8 mL) was added dropwise to a solution of ethyl 6-cyano-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate and ethyl 7-cyano-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate (0.6 g, 1.68 mmol) in a mixture of MeOH (10 mL) and THF (5 mL) at RT. The resulting RM was heated to 60° C. and maintained at this temperature for 4 h. The reaction progress was monitored by TLC. The RM was cooled to RT, poured into ice cold water (50 mL) and acidified with 1 N HCl (pH~2). The crude product was extracted with EtOAc (2×50 mL). The combined organic layer was washed with water (50 mL) followed by brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford a mixture of 6-cyano-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid and 7-cyano-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (0.55 g, 99%).


449

Step 3: To a solution of a mixture of 6-cyano-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid and 7-cyano-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (0.55 g, 1.67 mmol), and methyl 4-aminotetrahydro-2H-pyran-4-carboxylate (0.31 g, 2.01 mmol) in DMF (10 mL) was added DIPEA (0.86 mL, 5.03 mmol) followed by HATU (1.27 g, 3.35 mmol) at 0° C. under an argon atmosphere. The RM was allowed to attain RT and stirred for 16 h. The reaction progress was monitored by TLC. The RM was diluted with water (50 mL) and the organic compound was extracted with EtOAc (2×50 mL). The organic layer was washed with water (50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography over silica gel using 0-55% EtOAc in pet ether as an eluent to afford a mixture of methyl 4-(6-cyano-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)tetrahydro-2H-pyran-4-carboxylate and methyl 4-(7-cyano-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)tetrahydro-2H-pyran-4-carboxylate (0.5 g,

450

63%). Step 4: NaBH₄ (1.01 g, 26.65 mmol) was added portionwise to a pre-stirred solution of a mixture of methyl 4-(6-chloro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)tetrahydro-2H-pyran-4-carboxylate and methyl 4-(7-chloro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)tetrahydro-2H-pyran-4-carboxylate (0.5 g, 1.06 mmol) in a mixture of MeOH (10 mL) and THF (5 mL) at 0° C. The RM was allowed to attain RT and stirred for 16 h. The reaction progress was monitored by TLC. The RM was diluted with water (50 mL) and then the crude product was extracted with EtOAc (2×50 mL). The combined organic layer was washed with water (50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by GRACE flash chromatography using 0-60% acetonitrile and 0.1% FA in water as an eluent followed by RP prep-HPLC to afford cpd 188 (25 mg, 5%).

Synthesis of 4-cyano-N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide (Cpd 191)



451

Step 1: NBS (24.2 g, 136.22 mmol) was added to a solution of ethyl 5-hydroxy-2-methylbenzofuran-3-carboxylate (20.0 g, 90.81 mmol) in acetonitrile (600 mL) at RT under an argon atmosphere. The RM was stirred for 16 h at same temperature. The reaction progress was monitored by TLC. The RM was diluted with water (300 mL) and acidified with 1N-HCl to pH~2. The crude product was extracted with EtOAc (2×300 mL). The combined organic layer was washed with water (200 mL) followed by brine (200 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography over silica gel using 0-20% EtOAc and pet-ether as an eluent to afford ethyl 4-bromo-5-hydroxy-2-methylbenzofuran-3-carboxylate (5.0 g, 19%).

Step 2: CuCN (0.74 g, 8.36 mmol) was added to a solution of ethyl 4-bromo-5-hydroxy-2-methylbenzofuran-3-carboxylate (1.0 g, 3.44 mmol) in N,N-dimethylformamide (25 mL) at RT under an argon atmosphere. The resulting mixture was heated to 160° C. and maintained for 3 h. The reaction progress was monitored by TLC. The RM was cooled to RT, poured into water (100 mL) and extracted with EtOAc (2×100 mL). The combined organic layer was washed with water (2×50 mL) followed by brine (50 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography over silica gel using 0-20% EtOAc in pet-ether as an eluent to afford ethyl 4-cyano-5-hydroxy-2-methylbenzofuran-3-carboxylate (0.45 g, 55%).

Step 3: (2-(Trifluoromethyl)pyridin-3-yl)methanol (0.65 g, 3.67 mmol), ADDP (0.861 g, 3.41 mmol) and tri-N-butylphosphine (0.84 mL, 3.41 mmol) were added sequentially to a pre-stirred solution of ethyl 4-cyano-5-hydroxy-2-methylbenzofuran-3-carboxylate (0.6 g, 2.44 mmol) in THF (20 mL) at 0° C. under an argon atmosphere. The RM was allowed to attain RT and stirred for 3 h. The reaction progress was monitored by TLC. The RM was poured into water (50 mL) and extracted with EtOAc (2×50 mL). The combined organic layer was washed with water (50 mL) and brine (50 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by FCC over silica gel using 0-20% EtOAc in pet-ether as an eluent to afford ethyl 4-cyano-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (0.45 g, crude). The crude product thus obtained was used for next step without further purification.

Step 4: A solution of NaOH (0.178 g, 4.45 mmol) in water (5.0 mL) was added to a solution of ethyl 5-(benzyloxy)-4-cyano-2-methylbenzofuran-3-carboxylate (0.450 g, 1.11 mmol) in a mixture of MeOH (10 mL) and THF (10 mL) at RT. The resulting RM was heated to 60° C. and maintained for 3 h. The reaction progress was monitored by TLC. The RM was cooled to RT, poured into ice cold water (50 mL) and acidified with 1 N HCl (pH~2). The crude product was extracted with EtOAc (2×50 mL). The combined organic layer was washed with water (2×50 mL) followed by brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford 4-cyano-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid (0.4 g, crude). The crude product thus obtained was used for next step without further purification.


Step 5: To a solution of a mixture of 4-cyano-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid (0.4 g, 1.06 mmol) and tert-butyl 4-amino-3,3-difluoropiperidine-1-carboxylate (0.376 g, 1.59 mmol) in DMF (10 mL) was added DIPEA (0.50 mL, 3.18 mmol) followed by HATU (0.805 g, 2.12 mmol) at 0° C. under

452

argon atmosphere. The RM was allowed to attain RT and stirred for 3 h. The reaction progress was monitored by TLC. The RM was diluted with water (50 mL), and the crude product was extracted with EtOAc (2×50 mL). The combined organic layer was washed with water (50 mL), brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography over silica gel using 0-55% EtOAc and pet-ether as an eluent to afford tert-butyl 4-(4-cyano-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)-3,3-difluoropiperidine-1-carboxylate (0.3 g, crude).


Step 6: To a solution of tert-butyl 4-(4-cyano-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)-3,3-difluoropiperidine-1-carboxylate (0.3 g, 0.50 mmol) in DCM (2.5 mL) was added TFA (2.5 mL) dropwise at 0° C. under an argon atmosphere. The RM was allowed to attain RT and stirred for 16 h. The reaction progress was monitored by TLC. The RM was concentrated under reduced pressure, and the residue was basified with NaHCO₃ (pH~8). The crude product was extracted with EtOAc (2×50 mL). The combined organic layer was washed with water (50 mL) followed by brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by flash chromatography using 0-50% acetonitrile in 0.1% FA in water as an eluent to afford racemic cpd 191 (0.21 g, 89%). A preparative chiral SFC was performed on the racemic mixture of cpd 191 to afford cpd 191-En 1 and cpd 191-En 2.

Synthesis of 4-cyano-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 193)



453

-continued



Step 1: (4-Methylthiazol-5-yl)methanol (0.31 g, 2.44 mmol), ADDP (0.57 g, 2.28 mmol) and tri-N-butylphosphine (0.56 mL, 2.28 mmol) were added sequentially to a pre-stirred solution of ethyl 4-cyano-5-hydroxy-2-methylbenzofuran-3-carboxylate (0.4 g, 1.63 mmol) in THF (20 mL) at 0° C. under an argon atmosphere. The RM was allowed to warm to RT and stirred for 3 h. The reaction progress was monitored by TLC. The RM was poured into water (50 mL) and extracted with EtOAc (2×50 mL). The combined organic layer was washed with water (50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by FCC over silica using 0-20% EtOAc in pet-ether as an eluent to afford ethyl 4-cyano-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate (0.300 g, 52%). The crude product thus obtained was used for next step without further purification.


Step 2: A solution of NaOH (0.13 g, 3.38 mmol) in water (5.0 mL) was added to a pre-stirred solution of ethyl 5-(benzyloxy)-4-cyano-2-methylbenzofuran-3-carboxylate (0.30 g, 0.84 mmol) in MeOH (10 mL) and THF (10 mL) at RT. The resulting RM was heated to 60° C. and maintained at this temperature for 3 h. The reaction progress was monitored by TLC. The RM was cooled to RT, poured into ice cold water (50 mL), acidified with 1 N HCl (pH~2.0) and extracted with EtOAc (2×50 mL). The combined organic layer was washed with water (2×50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford 4-cyano-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (0.20 g, crude). The crude product thus obtained was used for next step without further purification.

Step 3: To a pre-stirred solution of 4-cyano-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (0.20 g, 0.61 mmol) and (4-aminotetrahydro-2H-pyran-4-yl)methanol (0.12 g, 0.92 mmol) in DMF (5 mL) were added DIPEA (0.29 mL, 1.83 mmol) followed by HATU (0.46 g, 1.22 mmol) at 0° C. under an argon atmosphere. The RM was stirred at room temperature for 3 h. The reaction progress was monitored by TLC. The RM was diluted with water (50 mL) and extracted with EtOAc (2×50 mL). The

454


combined organic layer was washed with water (50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by RP-HPLC to afford cpd 193 (0.90 g, 60%).

Synthesis of 6-cyano-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 194)




Step 1: To a solution of a mixture of 6-cyano-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (0.2 g, 0.60 mmol) and (4-aminotetrahydro-2H-pyran-4-yl)methanol (0.15 g, 1.21 mmol) in DMF (5 mL) was added DIPEA (0.32 mL, 1.82 mmol) followed by HATU (0.46 g, 1.21 mmol) at 0° C. under an argon atmosphere. The resulting RM was stirred at RT for 16 h. The reaction progress was monitored by TLC. The RM was diluted with water (30 mL) and extracted with EtOAc (2×30 mL). The combined organic layer was washed with water (30 mL) and brine (30 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by GRACE flash chromatography using 0.1% FA in water and acetonitrile as an eluent to afford cpd 194 (85 mg, 32%).

Synthesis of 6-cyano-N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide (Cpd 195) and 7-cyano-N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((2-(trifluoro-methyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide (Cpd 196)



455

456



457

Step 1: To a pre-stirred solution of CAN (199.18 g, 363.32 mmol) in water (500 mL) was added 2-chlorobenzene-1,4-diol (25.0 g, 173.01 mmol) at 0° C. The resulting RM was stirred at RT for 4 h. The reaction progress was monitored by TLC. The organic compound was extracted with Et₂O (3×200 mL). The combined organic layer was washed with brine (50 mL) and dried over anhydrous Na₂SO₄. The dried organic layer was passed through a silica gel column using diethyl ether as eluant and thus collected fractions were concentrated under reduced pressure to afford 2-chlorocyclohexa-2,5-diene-1,4-dione (20 g, 83%).

Step 2: To a pre-stirred solution of 2-chlorocyclohexa-2,5-diene-1,4-dione (20 g, 140.84 mmol) in toluene (300 mL) was added ethyl 3-oxobutanoate (54.92 g, 422.53 mmol) followed by anhydrous ZnCl₂ (23.0 g, 169.0 mmol) at RT under argon atmosphere. The resulting RM was heated to reflux and maintained for 16 h using Dean-Stark apparatus. The reaction progress was monitored by TLC. The RM was cooled to RT, filtered through a celite pad and the celite pad was washed with EtOAc (500 mL). The combined clear filtrate was concentrated under reduced pressure. The crude product was purified by FCC over silica gel using 0-10% EtOAc in pet-ether as an eluent to afford a mixture of ethyl 6-chloro-5-hydroxy-2-methylbenzofuran-3-carboxylate and ethyl 7-chloro-5-hydroxy-2-methylbenzofuran-3-carboxylate (7.0 g, 20%).

Step 3: A solution of a mixture of ethyl 6-chloro-5-hydroxy-2-methylbenzofuran-3-carboxylate and ethyl 7-chloro-5-hydroxy-2-methylbenzofuran-3-carboxylate (1.0 g, 3.93 mmol) and Zn(CN)₂ (2.01 g, 17.71 mmol) in dimethylacetamide (12 mL) was degassed with argon for 10 min. To the above RM, Pd(P(t-Bu)₃)₂ (0.60 g, 1.18 mmol) was added in one portion and it was then degassed for 5 min. The resulting mixture was heated to 160° C. and maintained at this temperature for 2 h under microwave heating. The reaction progress was monitored by TLC. The RM was poured into water (100 mL) and extracted with EtOAc (2×100 mL). The combined organic layer was washed with water (2×50 mL) and brine (50 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography over silica gel using 0-20% EtOAc in pet-ether as an eluent to afford a mixture of ethyl 6-cyano-5-hydroxy-2-methylbenzofuran-3-carboxylate and ethyl 7-cyano-5-hydroxy-2-methylbenzofuran-3-carboxylate (0.6 g, 62.5%).

Step 4: (2-(Trifluoromethyl)pyridin-3-yl)methanol (1.30 g, 7.34 mmol), ADDP (1.72 g, 6.85 mmol) and tri-N-butylphosphine (1.60 mL, 6.85 mmol) were added to a solution of a mixture of ethyl 6-cyano-5-hydroxy-2-methylbenzofuran-3-carboxylate and ethyl 7-cyano-5-hydroxy-2-methylbenzofuran-3-carboxylate (1.2 g, 4.89 mmol) in THF (30 mL) at RT under an argon atmosphere. The RM was stirred for 2 h at same temperature. The reaction progress was monitored by TLC. The RM was poured into water (100 mL) and extracted with EtOAc (2×100 mL). The combined organic layer was sequentially washed with water (50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography over silica gel using 0-20% EtOAc in pet-ether as an eluent to afford a regioisomeric mixture of ethyl 6-cyano-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate and ethyl 7-cyano-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (1.2 g, 60%).

458


Step 5: A solution of NaOH (0.47 g, 11.88 mmol) in water (8 mL) was added dropwise to a solution of a mixture of ethyl 6-cyano-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate and ethyl 7-cyano-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (1.2 g, 2.97 mmol) in a mixture of MeOH:THF (2:1) (23 mL) at RT. The resulting RM was heated to 60° C. for 4 h. The reaction progress was monitored by TLC. The RM was cooled to RT and poured into ice cold water (50 mL), and acidified with 1 N HCl to a pH~2. The crude product was extracted with EtOAc (2×50 mL). The combined organic layer was washed with water (50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford a regioisomeric mixture of 6-cyano-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid and 7-cyano-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxylic acid (1.0 g, 90%). This crude regioisomeric mixture was used in the next step without any separation.

Step 6: To a pre-stirred solution of a mixture of 6-cyano-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid and 7-cyano-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxylic acid (1.0 g, 2.65 mmol), and tert-butyl 4-amino-3,3-difluoropiperidine-1-carboxylate (0.90 g, 3.97 mmol) in DMF (20 mL) was added DIPEA (1.36 mL, 7.95 mmol) followed by HATU (2.02 g, 5.31 mmol) at 0° C. under an argon atmosphere. The RM was allowed to attain RT and stirred for 16 h. The reaction progress was monitored by TLC. The RM was diluted with water (100 mL) and the organic compound was extracted with EtOAc (2×70 mL). The combined organic layer was washed with water (50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography over silica using a gradient mixture of 0-55% EtOAc in pet-ether as an eluent to afford a regioisomeric mixture of tert-butyl 4-(6-cyano-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamido)-3,3-difluoropiperidine-1-carboxylate and tert-butyl 4-(7-cyano-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)-3,3-difluoropiperidine-1-carboxylate (1.0 g, 63%).

Step 7: 4.0 M HCl in dioxane (10.0 mL) was added dropwise to a solution of a mixture of tert-butyl 4-(6-cyano-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)-3,3-difluoropiperidine-1-carboxylate and tert-butyl 4-(7-cyano-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamido)-3,3-difluoropiperidine-1-carboxylate (1.0 g, 1.68 mmol) in DCM (15 mL) at 0° C. The RM was allowed to attain RT and was stirred for 5 h. The reaction progress was monitored by TLC. The RM was concentrated under reduced pressure. The residue was basified with sat. NaHCO₃ (100 mL) and the organic compound was extracted with 10% MeOH in DCM (3×50 mL). The combined organic layer was washed with water (50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by flash chromatography using 0.1% FA in water and acetonitrile as an eluent to afford an isomeric mixtures of racemic cpd 195 and racemic cpd 196 (0.60 g, 72%). A preparative chiral SFC was performed on the regioisomeric mixture of cpd 195 and cpd 196 to afford cpd 195-En 1, cpd 195-En 2, cpd 196-En 1 and cpd 196-En 2.

459

Synthesis of (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-cyclopropyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide (Cpd 747)



Step 1: To a stirred solution of methyl 2-cyclopropyl-5-hydroxybenzofuran-3-carboxylate (250 mg, 1.082 mmol) in THF (20 mL) was added ADDP (0.54 g, 2.164 mmol), $n\text{-Bu}_3\text{P}$ (2.1 mL, 2.164 mmol) and followed by addition of pyridin-2-ylmethanol (140 mg, 1.298 mmol) at 0°C . The RM was stirred for 16 h at RT and the reaction progress was monitored by TLC. The RM was diluted with water (40 mL) and extracted with ethyl acetate (3x50 mL). Combined

460


extracts were washed with water (20 mL), brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to get crude. Crude was purified by column chromatography over silica gel (100-200 mesh) using 5% ethyl acetate in pet-ether as an eluent to afford methyl-2-cyclopropyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxylate (3) (150 mg, 43%) as an off-white solid. TLC system: 20% ethyl acetate in pet ether; RF: 0.42.

Step 2: To a stirred solution of 2-cyclopropyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxylate (150 mg, 0.464 mmol) in MeOH:THF (1:1, 20 mL) was added 2N aqueous NaOH (4 mL) at RT. The RM was stirred for 16 h at RT and the reaction progress was monitored by TLC. The RM was concentrated, acidified to pH ~2 with 1 N HCl. The precipitated solid was filtered and dried under vacuum to afford 2-cyclopropyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxylic acid (120 mg) as an off-white solid. Crude used in the next step without purification. TLC system: 70% ethyl acetate in Pet ether; RF: 0.30.

Step 3: To a stirred solution of 2-cyclopropyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxylic acid (0.120 mg, 0.388 mmol) in DMF (5 mL) was added DIPEA (0.2 mL, 1.553 mmol), HATU (0.295 g, 0.776 mmol) at RT and followed by addition of (S)-2-amino-3-hydroxypropanamide-HCl (0.081 mg, 0.582 mmol). The RM was stirred for 16 h at RT. Reaction progress was monitored by TLC. The RM was diluted with water (50 mL), extracted with ethyl acetate (3x50 mL). The combined extracts were washed with water (20 mL), brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to get crude. The crude was purified by GRACE flash column chromatography using 0.1% aqueous formic acid and acetonitrile as an eluent to afford (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-cyclopropyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide (cpd 747) (35 mg, 19% over 2 steps).


The following compounds were prepared in a similar manner (use of appropriate reagents (chiral or racemic) and purification methods (including chiral HPLC or chiral SFC) known to the person skilled in the art) as described for cpd 747: Cpd: 748, 749, 750, 751.

Synthesis of 2-(difluoromethyl)-N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-5-((4-methylthiazol-5-yl)-methoxy)benzofuran-3-carboxamide (Cpd 198)



461

462



Step 1: To a solution of ethyl 5-hydroxy-2-methylbenzofuran-3-carboxylate (5.0 g, 22.7 mmol) in DCM (100 mL) was added imidazole (3.09 g, 45.4 mmol) followed by TBDMS-Cl (4.10 g, 27.2 mmol) at 0° C. under argon atmosphere. The RM was warmed to RT and stirred for 16 h. The reaction progress was monitored by TLC. The RM was poured into water (100 mL) and extracted with DCM (3×100 mL). The combined organic layer was washed with water (2×100 mL) followed by brine (200 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by FCC over silica-gel using 5% EtOAc in pet-ether as an eluent to obtain ethyl 5-((tert-butyldimethylsilyl)oxy)-2-methylbenzofuran-3-carboxylate (5.1 g, 68%).

Step 2: To a solution of ethyl 5-((tert-butyldimethylsilyl)oxy)-2-methylbenzofuran-3-carboxylate (6.0 g, 17.9 mmol) in 1,4-dioxane (250 mL) was added SeO₂ (39.8 g, 358 mmol) at RT under argon atmosphere. The resulting RM was heated to 120° C. for 48 h. The reaction progress was monitored by TLC. The RM was cooled to RT and filtered. The filtrate was concentrated under reduced pressure. The crude product was purified by FCC over silica gel using 2% EtOAc in pet-ether as an eluent to afford ethyl 5-((tert-butyldimethylsilyl)oxy)-2-formylbenzofuran-3-carboxylate (2.6 g, 42%).

Step 3: To a solution of ethyl 5-((tert-butyldimethylsilyl)oxy)-2-formylbenzofuran-3-carboxylate (2.6 g, 7.7 mmol) in DCM (100 mL) was added DAST (3.0 mL, 23.3 mmol) at 0° C. under argon atmosphere. The RM was allowed to warm to RT and stirred for 3 h. The reaction progress was monitored by TLC. The RM was diluted with DCM (100 mL), washed with sat. NaHCO₃ solution (100 mL), water (100 mL), and brine (100 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford ethyl 5-((tert-butyldimethylsilyl)oxy)-2-(difluoromethyl)benzofuran-3-carboxylate (2.4 g, crude product).

Step 4: To a solution of ethyl 5-((tert-butyldimethylsilyl)oxy)-2-(difluoromethyl)benzofuran-3-carboxylate (2.4 g, 6.48 mmol) in THF (50 mL) was added TBAF (1 M in THF) (12.9 mL, 12.9 mmol) at 0° C. under argon atmosphere. The RM was allowed to warm to RT and stirred for 1 h. The reaction progress was monitored by TLC. The RM was poured into water (100 mL) and extracted with EtOAc (3×100 mL). The combined organic layer was washed with water (100 mL) and brine (200 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford

ethyl 2-(difluoromethyl)-5-hydroxybenzofuran-3-carboxylate (1.8 g, crude). This crude product was used for next step without further purification.

Step 5: (4-Methylthiazol-5-yl)methanol (1.36 g, 10.5 mmol), ADDP (2.48 g, 9.8 mmol) and tri-N-butylphosphine (1.99 g, 9.8 mmol) were added sequentially to a solution of ethyl 2-(difluoromethyl)-5-hydroxybenzofuran-3-carboxylate (1.8 g, 7.0 mmol) in THF (150 mL) at 0° C. under an argon atmosphere. The resulting RM was stirred at RT for 3 h. The reaction progress was monitored by TLC. The RM was poured into water (200 mL) and extracted with EtOAc (3×100 mL). The combined organic layer was washed with water (100 mL) and brine (100 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography over silica gel using 0-20% EtOAc in pet-ether as an eluent to afford ethyl 2-(difluoromethyl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate (1.5 g, 55% for 3 steps).

Step 6: 2 M NaOH (20 mL) solution was added to a solution of ethyl 2-(difluoromethyl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate (800 mg, 2.17 mmol) in MeOH (25 mL) and THF (10 mL) at RT. The resulting RM was heated to 60° C. for 3 h. The reaction progress was monitored by TLC. The RM was cooled to RT and poured into ice cold water (100 mL) and acidified with 1 N HCl. The crude product was extracted with EtOAc (2×100 mL). The combined organic layer was washed with water (2×100 mL) and brine (100 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford 2-(difluoromethyl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (cpd 229) (400 mg, 55%).

Step 7: To a solution mixture of 2-(difluoromethyl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (0.4 g, 1.17 mmol) and 3-amino-1-(2-hydroxyethyl)pyrrolidin-2-one hydrochloride (255 mg, 1.41 mmol) in DMF (25 mL) were added DIPEA (0.43 mL, 2.35 mmol) followed by HATU (0.897 g, 2.35 mmol) at 0° C. under an argon atmosphere. The resulting RM was stirred at RT for 1 h. The reaction progress was monitored by TLC. The RM was diluted with water (100 mL) and extracted with EtOAc (2×100 mL). The combined organic layer was washed with water (100 mL) and brine (100 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by flash chromatography using 0.1% FA in water and acetonitrile as an eluent to afford cpd 198


463

(190 mg, 35%). A preparative chiral SFC was performed on the racemic mixture of cpd 198 to afford cpd 198-En 1 and cpd 198-En 2.

Cpd 197 was prepared in a manner similar (use of appropriate reagents and purification methods known to the person skilled in the art) to cpd 198-En 1 and cpd 198-En 2.

The following compounds were prepared in a similar manner (use of appropriate reagents (chiral or racemic) and purification methods (including chiral HPLC or chiral SFC) known to the person skilled in the art) as described for cpd 198: Cpd 742, 743, 744, 745, 746.

Synthesis of 2-(difluoromethyl)-N-(3,3-difluoropiperidin-4-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide (Cpd 199)



Step 1: To a pre-stirred solution of ethyl 2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (2.0 g, 5.27 mmol) in 1,4-dioxane (150 mL) was added SeO_2 (3.51 g, 31.6 mmol) at RT under argon atmosphere. The resulting RM was heated to 120° C. and maintained under stirring for 16 h. Additional amount of SeO_2 (5.8 g, 52.7 mmol) was added and the reaction was continued for another 24 h. The reaction progress was monitored by TLC. The RM was cooled to RT, filtered and the filtrate was concentrated under reduced pressure. The crude product was purified by FCC with silica gel using 10% EtOAc in pet-ether as an eluent to afford ethyl 2-formyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (900 mg, 43%).

Step 2: To a solution of ethyl 2-formyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (0.8 g, 2.03 mmol) in DCM (25 mL) was added DAST (0.53 mL,

4.07 mmol) at 0° C. under argon atmosphere. The RM was allowed to cool to RT and was stirred for 2 h. The reaction progress was monitored by TLC. The RM was diluted with DCM (100 mL), washed with sat. NaHCO_3 solution (100 mL), water (50 mL), brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford ethyl 2-(difluoromethyl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (800 mg). This crude product was used for next step without further purification.

Step 3: An aq. NaOH (2M solution) (25 mL) was added to a solution of ethyl 2-(difluoromethyl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (700 mg, 1.68 mmol) in MeOH (15 mL) and THF (5 mL) at RT. The resulting RM was heated to 60° C. for 3 h. The reaction progress was monitored by TLC. The RM was cooled to RT and poured into ice cold water (100 mL), acidified with 1 N HCl and extracted with EtOAc (2×100 mL). The combined


465

organic layer was washed with water (2×100 mL), brine (100 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford 2-(difluoromethyl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid (600 mg, crude product).

Step 4: To a pre-stirred solution of mixture of 2-(difluoromethyl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid (0.5 g, 1.29 mmol) and tert-butyl 4-amino-3,3-difluoropiperidine-1-carboxylate (0.365 g, 1.55 mmol) in DMF (25 mL) was added DIPEA (0.47 mL, 2.58 mmol) followed by HATU (0.982 g, 2.58 mmol) at 0° C. under argon atmosphere. The RM was allowed to attain RT and was stirred for 1 h. The reaction progress was monitored by TLC. The RM was diluted with water (100 mL), and extracted with EtOAc (2×100 mL). The combined organic layer was washed with water (100 mL) and brine (100 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography over silica gel using a gradient mixture of 0-25% EtOAc in pet-ether as an eluent to afford tert-butyl 4-(2-(difluoromethyl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)-3,3-difluoropiperidine-1-carboxylate (0.6 g, 55% over 3 steps).


Step 5: HCl (4M in dioxane) (15 mL) was added dropwise to a solution of 2-(difluoromethyl)-N-(3,3-difluoropiperidin-4-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide (0.6 g, 0.99 mmol) in DCM (50 mL) at 0° C. under an argon atmosphere. The RM was warmed to RT and stirred for 4 h. The reaction progress was monitored by TLC. The solvents were evaporated in vacuo. The resulting crude residue was partitioned between sat. NaHCO₃ (pH=9) and EtOAc (3×50 mL). The combined organic layer was washed with water (100 mL) and brine (100 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford 2-(difluoromethyl)-N-(3,3-difluoropiperidin-4-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide (0.4 g, 80%). A preparative chiral SFC was performed on the racemic mixture of cpd 199 to afford cpd 199-En 1 and cpd 199-En 2.

Synthesis of N-(3-(hydroxymethyl)azetidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 201)



466

-continued




Step 1: To a pre-stirred solution of 1-(tert-butyl) 3-ethyl 3-aminoazetidine-1,3-dicarboxylate (0.483 g, 1.98 mmol) in DMF (10 mL) was added DIPEA (3.5 mL, 19.80 mmol) followed by cpd 102 (0.6 g, 1.98 mmol) and HATU (1.5 g, 3.96 mmol) sequentially at 0° C. under an argon atmosphere. The RM was heated to 55° C. and stirred for 2 h. The reaction progress was monitored by TLC. The RM was diluted with ice cold water (100 mL) and extracted with EtOAc (150 mL). The combined organic layer was washed with brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by GRACE flash chromatography using 0.1% FA in water and acetonitrile as an eluent to afford 1-(tert-butyl) 3-ethyl 3-(2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)azetidine-1,3-dicarboxylate (0.45 g, 42%).

Step 2: NaBH₄ (0.113 g, 2.97 mmol) was added in small portions to a solution of 1-(tert-butyl) 3-ethyl 3-(2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)azetidine-1,3-dicarboxylate (0.45 g, 0.85 mmol) in MeOH (10 mL) at 0° C. and the resulting RM was stirred at RT for 2 h. The reaction progress was monitored by TLC. The RM was cooled to 0° C., diluted with water (30 mL) and extracted with 10% MeOH in DCM (2×50 mL). The combined organic layer was washed with brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford tert-butyl 3-(hydroxymethyl)-3-(2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)azetidine-1-carboxylate (0.4 g, crude). This crude product was used for next step without further purification.

Step 3: To a pre-stirred solution of tert-butyl 3-(hydroxymethyl)-3-(2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)azetidine-1-carboxylate (0.4 g, crude) in DCM (5.0 mL) was added TFA (1.0 mL) dropwise at 0° C. under an argon atmosphere. The RM was allowed to warm to room temperature and stirred for 5 h. The reaction progress was monitored by LC-MS. The RM was concentrated under reduced pressure. The crude product was purified by RP prep-HPLC to afford cpd 201 (75 mg, 25%).

467

Synthesis of N-(3-(hydroxymethyl)azetidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzo-furan-3-carboxamide (Cpd 202)



Step 1: To a solution of 1-(tert-butyl) 3-ethyl 3-aminoazetidine-1,3-dicarboxylate (0.417 g, 1.709 mmol) in DMF (10 mL) were added DIPEA (3.0 mL, 17.094 mmol), cpd 101 (0.6 g, 1.709 mmol) and HATU (1.2 g, 3.418 mmol) sequentially at 0° C. under an argon atmosphere. The resulting RM was heated to 55° C. for 2 h. The reaction progress was monitored by TLC. The RM was diluted with ice cold water (100 mL) and extracted with EtOAc (150 mL). The combined organic layer was washed with brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by flash chromatography using a gradient mixture of 30-50% of EtOAc in pet ether as an eluent to afford 1-(tert-butyl) 3-ethyl 3-(2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)azetidine-1,3-dicarboxylate (0.45 g, 42%).


Step 2: NaBH₄ (0.074 g, 1.94 mmol) was added in small portions to a solution of 1-(tert-butyl) 3-ethyl 3-(2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)azetidine-1,3-dicarboxylate (0.45 g, 0.779 mmol) in MeOH (10 mL) at 0° C. and the resulting RM was stirred at RT for 2 h. The reaction progress was monitored by TLC. The RM was cooled to 0° C., diluted with water (30 mL) and extracted with 10% MeOH in DCM (2x50 mL). The combined organic layer was washed with brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced

468

pressure to afford tert-butyl 3-(hydroxymethyl)-3-(2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)azetidine-1-carboxylate (0.4 g, crude). This crude product was used for next step without further purification.


Step 3: To a solution of tert-butyl 3-(hydroxymethyl)-3-(2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)azetidine-1-carboxylate (0.4 g, crude) in MeOH (15.0 mL) was added acetyl chloride (2.6 mL) dropwise at 0° C. under an argon atmosphere. The RM was allowed to attain RT and was stirred for 24 h. The reaction progress was monitored by LC-MS. The RM was concentrated under reduced pressure. The crude product was purified by reverse phase prep-HPLC to afford cpd 202 (100 mg, 30%).

Synthesis of 2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 232)




Step 1: To a pre-stirred solution mixture of 2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (0.2 g, 0.66 mmol), HATU (0.501 g, 1.32 mmol) and DIPEA (0.24 mL, 1.32 mmol) in DMF (10 mL) was added ammonium chloride (0.176 g, 3.3 mmol) at 0° C. and the resulting reaction mixture was stirred at room temperature for 2 h. The reaction progress was monitored by TLC. The reaction mixture was diluted with ice cold water (50 mL) and the precipitated solid was collected by filtration and dried under vacuum. The solid was washed with diethyl ether (20 mL) and dried to afford 2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (130 mg, 65%) as an off white solid. TLC system: 100% Ethyl acetate; Rf: 0.15.

Synthesis of (S)-N-(1-amino-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 279)



469

-continued




Step 1: To a stirred solution of 2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (150 mg, 0.495 mmol), HATU (376.2 mg, 0.99 mmol) and DIPEA (0.36 mL, 1.98 mmol) in DMF (10 mL) was added L-alaninamide hydrochloride (61.6 mg, 0.495 mmol) at 0° C. and the resulting reaction mixture was warmed to room temperature and stirred for 1 h. The reaction mass was diluted with water (20 mL) and extracted with ethyl acetate (2×20 mL). The combined organic layer was washed with water (2×20 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting compound was washed with n-pentane (2×10 mL) and was then filtered off to afford (S)-N-(1-amino-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (79 mg, 44%) as an off white solid. TLC: 10% Methanol in dichloromethane; R_f: 0.4.

The following compounds were prepared in a similar manner (use of appropriate reagents (chiral or racemic) and purification methods (including chiral HPLC or chiral SFC) known to the person skilled in the art) as described for cpd 279: Cpd. 271, 272, 273, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 318, 319, 320, 321, 324, 375, 376, 377, 378, 379, 380, 381, 382, 383, 395-Dia 1, 395-Dia 2, 396-En 1, 396-En 2, 397-En 1, 397-En 2, 398, 411-En 1, 411-En 2, 432-En 1, 432-En 2, 433-En 1, 433-En 2, 434-En 1, 434-En 2, 435-En 1, 435-En 2, 437-En 1, 437-En 2, 438, 439, 440, 441, 442, 443, 444, 445-En 1, 445-En 2, 446-En 1, 446-En 2, 447-En 1, 447-En 2, 448-En 1, 448-En 2, 449-En 1, 449-En 2, 450-En 1, 450-En 2, 451-En1, 451-En2, 452-En1, 452-En2, 453-En1, 453-En2, 454, 455-En1, 455-En2, 456-En1, 456-En 2, 457-En 1, 457-En 2, 458-En 1, 458-En 2, 459-En 1, 459-En 2, 460-En 1, 460-En 2, 461-En 1, 461-En 2, 462-En 1, 462-En 2, 463-En 1, 463-En 2, 464-En 1, 464-En 2, 465-En 1, 465-En 2, 466-En 1, 466-En 2, 467-En 1, 467-En 2, 468-En 1, 468-En 2, 469-En 1, 469-En 2, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 502, 503, 509, 577, 578, 579, 580, 581, 582, 584, 586, 590-En 1, 590-En2, 591-En1, 591-En2, 592-En1, 592-En2, 593-En1, 593-En2, 594-En1, 594-En2, 595-En1, 595-En 2, 596-En 1, 596-En 2, 597-En 1, 597-En 2, 598-En 1, 598-En 2, 600-En 1, 600-En 2, 601-En 1, 601-En 2, 650, 662.

470

Synthesis of 2-(difluoromethyl)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-methoxy-3-pyridin-3-yl)methoxy)benzofuran-3-carboxamide (Cpd 311)



Step 1: To a stirred solution of 2-(difluoromethyl)-5-((2-methoxypyridin-3-yl)methoxy)benzofuran-3-carboxylic acid (0.5 g, 9 mmol) in CH₂Cl₂ (30 mL) were added DIPEA (0.23 g, 1.8 mmol) and HATU (0.69 g, 1.8 mmol) at 0° C. followed by the addition of tert-butyl 4-amino-3,3-difluoropiperidine-1-carboxylate (273 mg, 1.1 mmol). The reaction mixture was stirred for 16 h at RT, the reaction progress was monitored by TLC. The reaction mixture was diluted with water (150 mL) and extracted with CH₂Cl₂ (3×100 mL). The combined extracts were washed with brine (150 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to get the crude material (0.7 g). The crude compound was purified by silica gel (100-200 mesh) column chromatography to afford 2-(difluoromethyl)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-methoxypyridin-3-yl)methoxy)benzofuran-3-carboxamide (0.35 g) as a white solid. TLC system: 5% MeOH in CH₂Cl₂; R_f: 0.3.


Step 2: To a stirred solution of tert-butyl 3,3-difluoro-4-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamido)piperidine-1-carboxylate (0.25 g, 1.5 mmol) in acetonitrile (15 mL), was added TMSCl (0.26 g, 2.4 mmol) followed by NaI (0.350 g, 2.4 mmol) at 0° C. and the reaction mixture was stirred for 3 h at 70° C. The reaction progress was monitored by TLC. The reaction mixture was diluted with water (150 mL) and extracted with EtOAc (2×20 mL). The combined extracts were washed with brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to get the crude material (500 mg, LCMS: 45%) as an off-white solid. The crude material was

471

purified by reverse phase prep-HPLC purification to afford 2-(difluoromethyl)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-oxo-1,2-dihydropyridin-3-yl)methoxy) benzofuran-3-carboxamide (130 mg) as an off-white solid. TLC system: 10% Methanol in Dichloromethane; Rf: 0.3.

The following compounds were prepared in a similar manner (use of appropriate reagents (chiral or racemic) and purification methods (including chiral HPLC or chiral SFC) known to the person skilled in the art) as described for cpd 311: Cpd. 349-En 1, 349-En 2, 350-En 1, 350-En 2, 408-En 1, 408-En 2, 420, 430, 436-En 1, 436-En 2, 443, 459-En 1, 459-En 2, 479, 485, 494, 495, 521-En 1, 521-En 2, 535-En 1, 535-En 2, 585, 596-En 1, 596-En 2, 614, 628, 630, 648-En 1, 648-En 2.

Synthesis of 2-methyl-N-(1-(methylcarbamoyl)cyclobutyl)-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 318)



Step 1: To a pre-stirred solution of 2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxylic acid (400 mg, 1.320 mmol) in CH_2Cl_2 (30 mL) were added DIPEA (0.9 mL, 5.280 mmol) and methyl 1-aminocyclobutane-1-carboxylate hydrochloride (304 mg, 1.848 mmol) followed by HATU (702 mg, 1.848 mmol) at 0°C . under an argon atmosphere. The reaction mixture was stirred at room temperature for 16 h. The reaction progress was monitored by TLC. The reaction mixture was diluted with water (30 mL) and extracted with EtOAc (3x50 mL). The combined organic layers were washed with water (1x20 mL) and brine (1x20 mL), dried over anhydrous Na_2SO_4 and concentrated under

472


reduced pressure to give the crude material, which was purified by flash chromatography with silica gel (100-200 mesh) using 30% ethyl acetate in pet ether as an eluent to afford methyl 1-(2-methyl-5-((2-methylthiazol-5-yl)methoxy) benzofuran-3-carboxamido) cyclobutane-1-carboxylate (300 mg, 55%) as an off-white solid. TLC system: 80% EtOAc in Pet ether; Rf: 0.66

Step 2: To a pre-stirred solution of methyl 1-(2-methyl-5-((2-methylthiazol-5-yl)methoxy) benzofuran-3-carboxamido) cyclobutane-1-carboxylate (300 mg, 0.724 mmol), in $\text{THF}:\text{MeOH}$ (1:1, 20 mL), was added 2N NaOH (3.5 mL) at RT, then the reaction mixture was heated to 60°C . The reaction mixture was stirred at this temperature for 3 h, and the reaction progress was monitored by TLC. The reaction mixture was concentrated, diluted with ice water, the pH was adjusted to ~ 2 with 1 N aq. HCl solution to give a solid. The solid was filtered off and dried under reduced pressure to afford 1-(2-methyl-5-((2-methylthiazol-5-yl)methoxy) benzofuran-3-carboxamido) cyclobutane-1-carboxylic acid (300 mg, crude) as an off-white solid. TLC system: 80% EtOAc in Pet ether; Rf: 0.24.


Step 3: To a pre-stirred solution of 1-(2-methyl-5-((2-methylthiazol-5-yl)methoxy) benzofuran-3-carboxamido) cyclobutane-1-carboxylic acid (300 mg, 0.750 mmol) in CH_2Cl_2 (30 mL) were added DIPEA (0.5 mL, 3.00 mmol) and methyl amine (2M in THF) (2.1 mL, 1.05 mmol) followed by HATU (399 mg, 1.05 mmol) at 0°C . under an argon atmosphere. The reaction mixture was stirred at room temperature for 2 h. The reaction progress was monitored by TLC. The reaction mixture was diluted with water (30 mL) and extracted with EtOAc (2x40 mL). The combined organic layers were washed with water (1x20 mL) and brine (1x30 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford the crude material, which was purified by flash chromatography with silica gel (100-200 mesh) using 20% ethyl acetate in pet ether as eluent to afford 2-methyl-N-(1-(methylcarbamoyl) cyclobutyl)-5-((2-methylthiazol-5-yl)methoxy) benzofuran-3-carboxamide (cpd 318) (85 mg, 28% over two steps) as a white solid. TLC system: 50% EtOAc in Pet ether; Rf: 0.54.

The following compounds were prepared in a similar manner (use of appropriate reagents (chiral or racemic) and purification methods known to the person skilled in the art) as described for cpd 318: cpds. 319, 320, 324, 725-En 1, 725-En 2, 740-En 1, 740-En 2.

Synthesis of 5-((2-cyanopyridin-3-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide (Cpd 325) and 3-(((3-((4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)carbamoyl)-2-methylbenzofuran-5-yl)oxy)methyl) picolinamide (Cpd 347)



473



Step 1: To a stirred solution of ethyl 5-hydroxy-2-methylbenzofuran-3-carboxylate (0.8 g, 3.63 mmol, 1.0 eq) in ACN (10 ml) at 0° C. was added Cs₂CO₃ (3.55 g, 10.89 mmol, 3 eq) and the mixture was stirred for 5 minutes, followed by the addition of 3-(bromomethyl)picolinonitrile (0.85 g, 4.36 mmol, 1.2 eq) at 0° C. The reaction mixture was stirred for 18 h at RT, monitored by TLC. The reaction mixture was diluted with ice water (100 ml) and extracted with EtOAc (300 ml). The organic layer was washed with ice water (100 ml) and brine (50 ml), dried over anhydrous Na₂SO₄, filtered and concentrated to get the crude product, which was purified by Grace flash column chromatography using eluent 20% EtOAc in pet ether to afford ethyl 5-((2-cyanopyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylate (1.0 g, 81%, off-white solid). TLC system: 20% EtOAc-pet ether; 2:8; Rf: 0.3.


Step 2: To ethyl 5-((2-cyanopyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylate (1.0 g, 2.97 mmol, 1.0 eq), in THF:MeOH:H₂O (1:1:1) (120 ml) was added NaOH (0.35 g, 8.92 mmol, 3.0 eq) at RT, the reaction mixture was then heated to 50° C. for 18 h, the reaction progress was monitored by TLC. The reaction mixture was concentrated and diluted with ice water, the pH was adjusted to ~2 with 1 N HCl solution to give a precipitate while stirring for 30 min. The precipitate was filtered off and was dried under vacuum to afford a mixture of 5-((2-cyanopyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid and 5-((2-carbam-

474

oylpyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid (0.75 g, crude). TLC system: 50% EtOAc-pet ether; Rf: 0.1.


Step 3: To a pre-stirred solution of a mixture of 5-((2-cyanopyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid, 5-((2-carbamoylpyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid (0.70 g, 2.27 mmol, 1 eq) and (4-aminotetrahydro-2H-pyran-4-yl)methanol (0.35 g, 2.72 mmol, 1.2 eq) in CH₂Cl₂ (100 mL) was added DIPEA (2.0 mL, 11.35 mmol, 5 eq) followed by HATU (1.29 g, 3.40 mmol, 1.5 eq) at 0° C. under an argon atmosphere. The reaction mixture was stirred at room temperature for 3 h and the reaction progress was monitored by TLC. The reaction mixture was diluted with water (100 mL) and extracted with CH₂Cl₂ (2×100 mL). The combined organic layers were washed with water (100 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford the crude material, which was purified by column chromatography/prep HPLC to afford 3-(((3-((4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)carbamoyl)-2-methylbenzofuran-5-yl)oxy)methyl)picolinamide (cpd 325) (0.110 g, 8% over 2 steps) and 5-((2-cyanopyridin-3-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide (cpd 347) (0.130 g, 10% over 2 steps) as white solids.

Synthesis of 4-fluoro-N-(3-(hydroxymethyl)oxetan-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 328) and 6-fluoro-N-(3-(hydroxymethyl)oxetan-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 362)



475

-continued



Step 1: To a pre-stirred solution of ethyl 5-hydroxy-2-methylbenzofuran-3-carboxylate (15 g, 68.18 mmol) in acetonitrile (500 mL) was added selectfluor (28.72 g, 81.13 mmol) at RT under an argon atmosphere. The reaction mixture was stirred at room temperature for 16 h. The reaction progress was monitored by TLC. The solvent was removed under reduced pressure to give a residue. The residue was dissolved in ethyl acetate (1000 mL), washed with water (500 mL) and brine (250 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to give the crude material. The crude product was purified by column chromatography over silica gel (100-200 mesh) using 0-20% ethyl acetate in pet-ether as an eluent to afford mixture of ethyl 4-fluoro-5-hydroxy-2-methylbenzofuran-3-carboxylate and ethyl 6-fluoro-5-hydroxy-2-methylbenzofuran-3-carboxylate (3.0 g, 18%) as a yellow solid. TLC system: 30% Ethyl acetate in pet-ether; Rf: 0.5.

Step 2: (4-Methylthiazol-5-yl)methanol (4.06 g, 31.51 mmol), ADDP (10.58 g 42.01 mmol) and tri-N-butyl phosphine (10.60 ml 42.01 mmol) were added sequentially to a pre-stirred solution of a mixture of ethyl 4-fluoro-5-hy-

476

droxy-2-methylbenzofuran-3-carboxylate and ethyl 6-fluoro-5-hydroxy-2-methylbenzofuran-3-carboxylate (5.0 g, 21.00 mmol) in THF (500 mL) at 0° C. under an argon atmosphere. The reaction mixture was stirred at room temperature for 1 h. The reaction progress was monitored by TLC. The reaction mixture was diluted with water (500 mL) and extracted with ethyl acetate (2×500 mL). The combined organic layers were washed with water (250 mL) and brine (250 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to give the crude material. The crude material was purified by column chromatography over silica gel (100-200 mesh) using 0-30% ethyl acetate in pet-ether as eluent to afford a mixture of ethyl 4-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate and ethyl 6-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate (6.0 g, 81%) as white solid. TLC system: 50% Ethyl acetate in pet-ether; Rf: 0.4.

Step 3: 2N NaOH in water (40 mL) was added to a pre-stirred solution of mixture of ethyl 4-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate and ethyl 6-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate (6.0 g, 17.19 mmol) in a mixture of methanol (40 mL) and THF (10 mL) at RT. The reaction mixture was heated to 80° C. for 3 h. The reaction progress was monitored by TLC. The reaction mixture was cooled to RT and then concentrated under reduced pressure to get the crude material. The crude material was poured into ice water (100 mL), acidified to pH~2 with 1 N HCl, the obtained white solid was filtered off and was washed with water (20 mL) to afford a mixture of 4-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (cpd 233) and 6-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (cpd 243) (4.5 g, 81%) as white solid. TLC system: 80% Ethyl acetate in pet-ether; Rf: 0.2.

Step 4: To a stirred solution of a mixture of 4-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid compound and 6-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (1:1) (550 mg, 1.71 mmol) in DMF (10 mL), were added DIPEA (1.2 mL, 6.84 mmol), HATU (970 mg, 2.57 mmol) and (3-(methylamino)oxetan-3-yl)methanol (260 mg, 2.57 mmol) at 0° C. and the reaction mixture was stirred for 2 h at RT. The reaction progress was monitored by TLC. After completion of the reaction, the reaction mixture was diluted with water (150 mL) and extracted with ethyl acetate (3×100 mL). The combined organic layers were washed with brine (150 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to get the crude compound (715 mg, LCMS: 37% & 23% ratio). The crude compound was purified by reverse phase prep-HPLC purification (column: INERTSIL-ODS 2 (250*19 mm), 5 μm, mobile phase: A: 10 mM Ammonium Bi Carbonate in H₂O, B: MeCN, Gradient: 0/30, 8/55, 9/55, 9.1/98, 11/98, 11.1/30, 14/30; flow rate: 18 mL/min; diluent: MeCN+H₂O+THF) to afford cpd. 328 (325 mg) and cpd. 362 (170.9 mg) as an off-white solid.





The following compounds were prepared in a similar manner (use of appropriate reagents and purification methods (including chiral HPLC or chiral SFC) known to the person skilled in the art) as described for cpds 328 and 362: Cpd. 326-En 1, 326-En 2, 327, 330, 360-En 1, 360-En 2, 361, 364, 572-En 1, 572-En 2, 583, 587.

The following compounds were prepared in a similar manner (use of appropriate reagents and purification meth-

477

ods (including chiral HPLC or chiral SFC) known to the person skilled in the art) as described for cpds 233: Cpd 234.

Synthesis of 4-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((2-methoxypyridin-3-yl)methoxy)benzofuran-3-carboxamide (Cpd 329) and 6-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((2-methoxypyridin-3-yl)methoxy)benzofuran-3-carboxamide (Cpd 363)




Step 1: To a pre-stirred solution of a mixture of 4-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide (cpd 330) and 6-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide (cpd 364)

478

(1.5 g, 3.37 mmol) in acetonitrile (100 mL) was added NaI (0.99 g, 6.75 mmol) at 0° C. followed by the dropwise addition of TMSCl (0.91 mL, 6.75 mmol) at 0° C. under an argon atmosphere. The reaction mixture was heated to 40° C. for 24 h. The reaction progress was monitored by TLC. The reaction mixture was diluted with water (100 mL) and extracted with 10% Methanol in CH₂Cl₂ (2×100 mL). The combined organic layers were washed with water (100 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford the crude product, which was purified by prep-HPLC purification to afford 4-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((2-methoxypyridin-3-yl)methoxy)benzofuran-3-carboxamide (cpd 329) (135 mg) as an off-white solid and 6-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((2-methoxypyridin-3-yl)methoxy)benzofuran-3-carboxamide (cpd 363) (30 mg) as a white solid.

The following compounds were prepared in a similar manner (use of appropriate reagents (chiral or racemic) and purification methods known to the person skilled in the art) as described for cpds 329 and 363: Cpd. 368.

Synthesis of 5-((1-(2,2-difluoroethyl)-1H-pyrazol-5-yl)methoxy)-N-(3-(hydroxymethyl)-2-oxopiperidin-3-yl)-2-methylbenzofuran-3-carboxamide (Cpd 332)



Step 1: To a stirred solution of 5-((1-(2,2-difluoroethyl)-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid (500 mg, 1.48 mmol) in DMF (20 mL) were added DIPEA (1.0 mL, 5.95 mmol), HATU (787 mg, 2.07 mmol) and ethyl 3-amino-2-oxopiperidine-3-carboxylate (385 mg, 2.07 mmol) at 0° C. The reaction mixture was stirred for 16

479


h at RT, and the reaction progress was monitored by TLC. The reaction mixture was diluted with water (50 mL) and extracted with ethyl acetate (2×100 mL). The combined organic layers were washed with water (40 mL) and brine (20 mL), dried over anhydrous Na₂SO₄, filtered and concentrated to get the crude material. The crude material was purified by Grace chromatography using 35% EtOAc in pet ether as an eluent to afford ethyl 3-(5-((1-(2,2-difluoroethyl)-1H-pyrazol-5-yl) methoxy)-2-methylbenzofuran-3-carboxamido)-2-oxopiperidine-3-carboxylate (350 mg, 46%) as a brown solid. TLC system: 60% Ethyl acetate in per ether; RF: 0.48.

Step 2: To a stirred solution of ethyl 3-(5-((1-(2,2-difluoroethyl)-1H-pyrazol-5-yl) methoxy)-2-methylbenzofuran-3-carboxamido)-2-oxopiperidine-3-carboxylate (350 mg, 0.69 mmol) in MeOH (40 mL) was added NaBH₄ (131 mg, 3.47 mmol) at 0° C. The reaction mixture was stirred for 16 h at RT. The reaction progress was monitored by TLC. The reaction mixture was concentrated under reduced pressure, diluted with water (100 mL), and extracted with ethyl acetate (2×100 mL). The combined extracts were washed with brine (50 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to get the crude material which was purified by GRACE flash chromatography using 0.1% aqueous formic acid in acetonitrile as an eluent to afford 5-((1-(2,2-difluoroethyl)-1H-pyrazol-5-yl) methoxy)-N-(3-(hydroxymethyl)-2-oxopiperidin-3-yl)-2-methylbenzofuran-3-carboxamide (250 mg, 78%) as an off-white solid. TLC system: 100% EtOAc; RF: 0.24.

A preparative chiral SFC was performed on the racemic mixture of cpd 332 to afford cpd 332-En 1 and cpd 332-En 2.


The following compounds were prepared in a similar manner (use of appropriate reagents and purification methods (including chiral HPLC or chiral SFC) known to the person skilled in the art) as described for cpd 332: Cpd. 333-En 1, 333-En 2, 355-En 1, 355-En 2, 356-En 1, 356-En 2, 510-En 1, 510-En 2, 511-En 1, 511-En 2, 512-En 1, 512-En 2, 513-En 1, 513-En 2, 514-En 1, 514-En 2, 515-En 1, 515-En 2, 516-En 1, 516-En 2, 517-En 1, 517-En 2, 518-En 1, 518-En 2, 519-En 1, 519-En 2, 520-En 1, 520-En 2, 521-En 1, 521-En 2, 522-En 1, 522-En 2, 719-En 1, 719-En 2, 732-En 1, 732-En 2, 733-En 1, 733-En 2, 734-En 1, 734-En 2, 735-En 1, 735-En 2, 736-En 1, 736-En 2, 737-En 1, 737-En 2.

Synthesis of N-(1-((difluoromethoxy)methyl)cyclopropyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 399)



480


-continued



Step 1: To a suspension of N-(1-(hydroxymethyl)cyclopropyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (cpd 418) (300 mg, 0.866 mmol) and CuI (30.6 mg, 0.168 mmol) in acetonitrile (20 mL) was added a solution of 2,2-difluoro-2-(fluorosulfonyl)acetic acid (215 mg, 1.209 mmol) in acetonitrile (3 mL) at room temperature and the resulting reaction mixture was heated to 50° C.-55° C. for 1 h. The reaction progress was monitored by TLC. The reaction mixture was cooled to room temperature and filtered. The filtrate was evaporated under reduced pressure to give the crude material which was purified by GRACE flash chromatography using 0.1% of formic acid in water and acetonitrile as an eluent to afford N-(1-((difluoromethoxy)methyl)cyclopropyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (cpd 399) (40 mg, 12%) as a white solid. TLC system: 50% Ethyl acetate in pet-ether; Rf: 0.8

The following compounds were prepared in a similar manner (use of appropriate reagents and purification methods known to the person skilled in the art) as described for cpd 399: cpd 400.

Synthesis of (R)-N-(1-(1H-benzo[d]imidazol-2-yl)-2-hydroxyethyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 401-En 1)




Step 1: EDC·HCl (0.378 g, 1.9801 mmol), HOBT (267 mg, 1.9801 mmol) and DIPEA (510 mg, 3.960 mmol) were added sequentially to a stirred solution of (S)-2-amino-2-(1H-benzo[d]imidazol-2-yl)ethan-1-ol hydrochloride (400 mg, 1.3201 mmol) in DMF (5.0 mL) at room temperature.

481

The resulting reaction mixture was stirred at room temperature for 2 h. The reaction progress was monitored by TLC. The reaction mixture was diluted with ice cold water (50 mL) and the precipitated solid was filtered and dried under vacuum to afford (S)—N-(1-(1H-benzo[d]imidazol-2-yl)-2-hydroxyethyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (cpd 401-En 1) (140 mg) as a white solid. TLC system: 100% Ethyl acetate; Rf: 0.2.

The following compounds were prepared in a similar manner (use of appropriate reagents and purification methods known to the person skilled in the art) as described for cpd 401-En 1: cpd 401-En 2.

Synthesis of N-(1-(dimethylamino)-3-hydroxypropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 413)



Step 1: To a stirred solution of 2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (1.0 g, 3.3 mmol) in DMF (10 mL) was added DIPEA (1.7 mL, 9.9 mmol), HATU (2.5 g, 6.6 mmol) at 0° C. followed by addition of methyl aziridine-2-carboxylate (0.45 g, 4.95 mmol). The reaction mixture was then warmed to room temperature and stirred for 16 h at the same temperature. The reaction progress was monitored by TLC. The reaction mixture was diluted with water (20 mL) and extracted with EtOAc (2×20 mL). The combined extracts were washed with water (3×20 mL) and brine (20 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford methyl 1-(2-methyl-5-((4-methylthiazol-5-yl)

482

methoxy)benzofuran-3-carbonyl)aziridine-2-carboxylate (1.0 g, crude) as a pale green gum. TLC system: 10% MeOH in dichloromethane; Rf: 0.6.


Step 2: To a stirred solution of methyl 1-(2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carbonyl)aziridine-2-carboxylate (1.0 g, 2.591 mmol) in dimethylamine (69.5 mmol, 5.8 g of dimethylamine HCl was therefore treated with TEA (10 mL) in CHCl₃ (10 mL)) at RT was added boron trifluoride diethyl etherate (0.3 mL, 2.59 mmol) at 0° C., then the reaction mixture was stirred for 16 h at RT. The reaction progress was monitored by TLC. The reaction mixture was diluted with water (30 mL) and extracted with dichloromethane (2×30 mL). The combined extracts were washed with brine (2×20 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to get the crude compound, which was purified by GRACE flash chromatography using 0.1% aqueous formic acid in acetonitrile to afford methyl 3-(dimethylamino)-2-(2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)propanoate (650 mg, 59%) as a pale yellow gum. TLC system: 80% EtOAc in pet-ether; Rf: 0.3.

Step 3: To a stirred solution of methyl 3-(dimethylamino)-2-(2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)propanoate (300 mg, 0.694 mmol) in MeOH (10 mL) was added NaBH₄ (131.8 mg, 3.47 mmol) at 0° C. The reaction mixture was stirred for 16 h at RT and the reaction progress was monitored by TLC. Then the reaction mixture was evaporated under reduced pressure, diluted with water (20 mL) and extracted with ethyl acetate (2×20 mL). The combined organic layers were dried over anhydrous Na₂SO₄ and concentrated to get the crude material which was purified by GRACE flash chromatography using 0.1% aqueous formic acid in acetonitrile as an eluent to afford N-(1-(dimethylamino)-3-hydroxypropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (cpd 413) (130 mg, 46%) as an off-white gum. TLC system: 10% Methanol in dichloromethane; Rf: 0.4


A preparative chiral SFC was performed on the racemic mixture of cpd 413 to afford cpd 413-En 1 and cpd 413-En 2.

The following compounds were prepared in a similar manner (use of appropriate reagents (chiral or racemic) and purification methods (including chiral HPLC or chiral SFC) known to the person skilled in the art) as described for cpd 413: 412-En 1, 412-En 2, 446-En 1, 446-En 2, 449-En 1, 449-En 2, 450-En 1, 450-En 2, 487-En 1, 487-En 2, 488-En 1, 488-En 2, 575-En 1, 575-En 2.

Synthesis of N-(1-amino-1-oxo-3-(1H-1,2,4-triazol-1-yl)propan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 432)



483



Step 1: To a stirred solution of methyl ((benzyloxy) carbonyl)serinate (1.0 g, 3.95 mmol) in chloroform (10 mL) was added PCl_5 (0.4 mL, 4.34 mmol) at 0°C . The reaction mixture was stirred for 16 h at RT. The reaction progress was monitored by TLC. The reaction mixture was diluted with water (50 mL) and extracted with dichloromethane (2x20 mL). The combined extracts were washed with sat. NaHCO_3 solution (2x20 mL) and brine (2x20 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to get the crude compound, which was purified by column chromatography using silica gel (100-200) and 10% ethyl acetate in pet ether as an eluent to afford methyl 2-(((benzyloxy)carbonyl)amino)-3-chloropropanoate (500 mg, 47%) as an off-white solid. TLC system: 30% EtOAc in Pet ether; RF: 0.6.

Step 2: To a stirred solution of 1H-1,2,4-triazole (200 mg, 2.89 mmol) in chloroform (10 mL) was added methyl 2-(((benzyloxy)carbonyl)amino)-3-chloropropanoate (946 mg, 3.47 mmol) at RT. To the reaction mixture was added DBU (878 mg, 5.78 mmol) at 0°C . then the reaction mixture was stirred for 16 h at 60°C . The reaction progress was monitored by TLC. The reaction mixture was diluted with water (50 mL) and extracted with dichloromethane (3x30 mL). The combined extracts were washed with brine (2x20 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to get the crude compound, which was purified by GRACE flash chromatography using 0.1% aque-

484

ous formic acid in acetonitrile as an eluent to afford methyl 2-(((benzyloxy)carbonyl)amino)-3-(1H-1,2,4-triazol-1-yl)propanoate (500 mg, 57%) as a pale yellow gum. TLC system: 50% EtOAc in Pet ether; RF: 0.6.

Step 3: To a stirred solution of methyl 2-(((benzyloxy) carbonyl)amino)-3-(1H-1,2,4-triazol-1-yl)propanoate (400 mg, 1.31 mmol) in MeOH (10 mL) was added 10% Pd/C (170 mg) at RT. The reaction mixture was stirred for 2 h at RT under H_2 (balloon atmosphere). The reaction progress was monitored by TLC. The reaction mixture was diluted with MeOH (10 mL), filtered through a celite bed and the celite bed was washed with MeOH (2x10 mL). The filtrate was concentrated under reduced pressure to afford methyl 2-amino-3-(1H-1,2,4-triazol-1-yl)propanoate (200 mg, 42%) as a pale yellow gum. TLC system: 5% MeOH in dichloromethane; RF: 0.4.


Step 4: To a stirred solution of 2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (300 mg, 0.99 mmol) and methyl 2-amino-3-(1H-1,2,4-triazol-1-yl)propanoate (175 mg, 1.48 mmol) in DMF (10 mL) were added DIPEA (0.5 mL, 2.97 mmol) and HATU (752 mg, 1.98 mmol) at 0°C . under an argon atmosphere. The reaction mixture was stirred at room temperature for 2 h. The reaction progress was monitored by TLC, the reaction mixture was diluted with cold water (25 mL) and extracted with ethyl acetate (2x40 mL). The combined organic layers were washed with water (15 mL) and brine (15 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford the crude product which was purified by GRACE flash chromatography using 0.1% aqueous formic acid in acetonitrile as an eluent to afford methyl 2-(2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)-3-(1H-1,2,4-triazol-1-yl)propanoate (200 mg, 44%) as an off-white gum. TLC system: 10% MeOH in dichloromethane; RF: 0.3

Step 5: A stirred solution of methyl 2-(2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)-3-(1H-1,2,4-triazol-1-yl)propanoate (200 mg, 0.439 mmol) in 7.0 M NH_3 in methanol (20 mL) was stirred for 16 h at RT and the reaction progress was monitored by TLC. Then the reaction mixture was evaporated under reduced pressure to get the crude compound, which was purified by GRACE flash chromatography using 0.1% aqueous formic acid in acetonitrile as an eluent to afford (120 mg, 62%) of N-(1-amino-1-oxo-3-(1H-1,2,4-triazol-1-yl)propan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (cpd 432) as an off-white gum. TLC system: 10% MeOH in dichloromethane; RF: 0.2

A preparative chiral SFC was performed on the racemic mixture of cpd 432 to afford cpd 432-En 1 and cpd 432-En 2.


The following compounds were prepared in a similar manner (use of appropriate reagents (chiral or racemic) and purification methods (including chiral HPLC or chiral SFC) known to the person skilled in the art) as described for cpd 432: 729-En 1, 729-En 2.

Synthesis of N-(1-amino-3-(difluoromethoxy)-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 447)



485

-continued



Step 1: 2-Amino-3-hydroxypropanamide·HCl (686 mg, 6.600 mmol) was added to a stirred solution of 2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (500 mg, 1.65 mmol), HATU (1.14 g, 3.00 mmol) and DIPEA (0.88 mL, 4.95 mmol) in DMF (10 mL) at room temperature. The resulting reaction mixture was stirred at room temperature and under argon atmosphere for 2 h. The reaction progress was monitored by TLC. (Note: Another batch was repeated on 500 mg scale and both the reaction mixtures were mixed together for work-up). The reaction mixture was diluted with water (100 mL) and extracted with ethyl acetate (2×100 mL). The combined organic layers were washed with water (100 mL) and brine (100 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (cpd 453) (600 mg, 46%) as an off-white solid. TLC system: 100% Ethyl acetate; R_f: 0.2.


Step 2: A solution of 2,2-difluoro-2-(fluorosulfonyl)acetic acid (68.5 mg, 0.38 mmol) in acetonitrile (1 mL) at 50° C. was added to a suspension of N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (150 mg, 0.38 mmol) and CuI (78.0 mg, 0.38 mmol) in acetonitrile (5 mL) and the mixture was heated to 50° C. for 16 h under an argon atmosphere. The reaction progress was monitored by TLC. (Note: Another batch was repeated on 150 mg scale, both the reaction mixtures were mixed together for work-up). The reaction mixture was cooled to room temperature and filtered. The filtrate was evaporated under reduced pressure. The crude compound was purified by GRACE flash chromatography using 0.1% of formic acid in water and acetonitrile as an eluent to afford N-(1-amino-3-(difluoromethoxy)-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (cpd 447) (120 mg, 35%) as a white solid. TLC system: 100% Ethyl acetate; R_f: 0.4.

A preparative chiral SFC was performed on the racemic mixture of cpd 447 to afford cpd 447-En 1 and cpd 447-En 2.

486

The following compounds were prepared in a similar manner (use of appropriate reagents (chiral or racemic) and purification methods (including chiral HPLC or chiral SFC) known to the person skilled in the art) as described for cpd 447: 448-En 1, 448-En 2.

Synthesis of N-(3-(dimethylamino)oxetane-3-carbonyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 508), N-(3-carbamoyloxetan-3-yl)-N,2-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (cpd 589) and N-(3-cyanooxetan-3-yl)-N,2-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 599)



Step 1: To a solution of 3-Oxetanone (3 g, 41.66 mmol) in CH₂Cl₂ (50 mL) was added 33% methyl amine in ethanol (39 mL, 416.6 mmol) and the solution was cooled to 0° C. TMSCN (8.3 mL, 66.66 mmol) was added slowly dropwise for 15 min to the reaction mixture and stirring was continued for 48 h at RT. The reaction mixture was diluted with dichloromethane (100 mL), washed with water (2×50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄, filtered and the filtrate was concentrated under reduced pressure to give the crude material, which was purified by grace column chromatography using 2% methanol in dichloromethane as

487


an eluent to afford 3-(methylamino)oxetane-3-carbonitrile (350 mg) as a light brown liquid. TLC system: 5% Methanol in dichloromethane; Rf: 0.4.

Step 2: To a stirred solution of 2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (500 mg, 1.65 mmol) in dichloromethane (30 mL), was added thionyl chloride (0.24 mL, 3.30 mmol) at 0° C. and the mixture was then heated to 40° C. for 2 h. The reaction progress was monitored by TLC. After completion of the reaction volatiles were removed under reduced pressure. To this was added an already prepared mixture of 3-(methylamino)oxetane-3-carbonitrile (369 mg, 3.30 mmol) in dichloromethane (20 mL), the mixture was cooled to 0° C. followed by the addition of triethyl amine (1.15 mL, 8.25 mmol). The reaction mixture was stirred for 16 h at RT, reaction progress was monitored by TLC. The reaction mixture was diluted with water (50 mL) and extracted with dichloromethane (2x50 mL). The combined extracts were washed with brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to get the crude material (0.6 g) which was purified by grace column chromatography to afford cpd 599 (300 mg) as an off-white solid. TLC system: 5% Methanol in dichloromethane; Rf: 0.6.

Step 3: To a solution of N-(3-cyanooxetan-3-yl)-N,2-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (1 g, 2.51 mmol) in THF (50 mL) was added NaOH (201 mg, 5.03 mmol) at 0° C. 30% H₂O₂ in water (0.2 mL, 2.51 mmol) was added slowly (dropwise) over 5 min to the reaction mixture and the mixture was then stirred for 1 h at RT. The reaction progress was monitored by TLC. The reaction mixture was diluted with dichloromethane (100 mL). The organic layer was washed with water (2x50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to give the crude material which was purified by grace chromatography using 3% Methanol in dichloromethane as an eluent to afford N-(3-carbamoyloxetan-3-yl)-N,2-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (cpd 589) (530 mg) as an off-white solid. TLC system: 5% Methanol in Dichloromethane; Rf: 0.2.


Step 4: To a solution of N-(3-carbamoyloxetan-3-yl)-N,2-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (500 mg, 1.20 mmol) in THF (50 mL) was added 60% NaH (72.2 mg, 1.80 mmol) at 0° C., and the mixture was stirred for 15 min. Mel (0.082 mL, 1.32 mmol) was added slowly (dropwise) over 5 min to the reaction mixture and the mixture was stirred for 2 h at 0° C. The reaction progress was monitored by TLC. The reaction mixture was diluted with dichloromethane (100 mL). The organic layer was washed with water (2x50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to give the crude material, which was purified by prep-HPLC to afford N-(3-(dimethylamino)oxetane-3-carbonyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (cpd 508) (60 mg) as an off-white solid. TLC system: 5% Methanol in Dichloromethane; Rf: 0.5.

Synthesis of N-(3-(hydroxymethyl)pyrrolidin-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 530)



488

-continued



Step 1: To a stirred solution of 3-amino-1-(tert-butoxycarbonyl)pyrrolidine-3-carboxylic acid (0.6 g, 2.606 mmol) in dichloromethane (10 mL) and MeOH (5 mL) was added a 0.6 M TMS-diazomethane solution in n-Hexane (8.6 mL, 5.212 mmol) at 0° C. The reaction mixture was stirred for 16 h at RT. Reaction progress was monitored by TLC. The reaction mixture was diluted with water (20 mL) and extracted with dichloromethane (2x20 mL). The combined extracts were washed with brine (2x20 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford the crude product of 1-(tert-butyl) 3-methyl 3-aminopyrrolidine-1,3-dicarboxylate (0.6 g) as a pale yellow gum. TLC system: 5% MeOH in dichloromethane; RF: 0.3.

Step 2: To a stirred solution of 2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid (300 mg, 0.99 mmol) in DMF (10 mL) were added DIPEA (0.53 mL, 2.97 mmol), HATU (564.3 mg, 1.485 mmol) and 1-(tert-butyl) 3-methyl 3-aminopyrrolidine-1,3-dicarboxylate (0.45 g, 4.95 mmol) at 0° C. The reaction mixture was stirred at room temperature for 16 h. The reaction progress was monitored by TLC. The reaction mixture was diluted with water (30 mL) and extracted with EtOAc (2x30 mL). The combined extracts were washed with water (3x20 mL) and brine (20 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to get the crude material, which was purified by column chromatography over silica gel (100-200 mesh) using 40% ethyl acetate in pet-ether as an eluent to afford 1-(tert-butyl) 3-methyl 3-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)pyrrolidine-1,3-dicarboxylate (330 mg, 62%) as a pale green gum. TLC system: 10% MeOH in dichloromethane; RF: 0.5.

489


Step 3: To a stirred solution of 1-(tert-butyl) 3-methyl 3-(2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)pyrrolidine-1,3-dicarboxylate (300 mg, 0.566 mmol) in MeOH (15 mL) was added NaBH₄ (107.8 mg, 2.834 mmol) at 0° C. The reaction mixture was stirred for 16 h at RT and the reaction progress was monitored by TLC. The reaction mixture was evaporated under reduced pressure, diluted with water (20 mL), and extracted with ethyl acetate (2×20 mL). The combined organic layers were dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford (250 mg, 85%) of tert-butyl 3-(hydroxymethyl)-3-(2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate as an off-white gum. TLC system: 10% Methanol in dichloromethane; RF: 0.2

Step 4: To a stirred solution of tert-butyl 3-(hydroxymethyl)-3-(2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate (250 mg, 0.498 mmol) in dichloromethane (10 mL) was added TFA (0.4 mL) at 0° C. The reaction mixture was stirred for 6 h at RT and the reaction progress was monitored by TLC. The reaction mixture was diluted with dichloromethane (50 mL). The organic layer was washed with saturated NaHCO₃ solution (2×20 mL) and brine (20 mL), dried over anhydrous Na₂SO₄ and concentrated to get the crude material which was purified by GRACE flash chromatography using 0.1% aqueous formic acid in acetonitrile as an eluent to afford N-(3-(hydroxymethyl)pyrrolidin-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide (100 mg, 50%) as an off-white gum. TLC system: 10% Methanol in dichloromethane; RF: 0.1.


A preparative chiral SFC was performed on the racemic mixture of cpd 530 to afford cpd 530-En 1 and cpd 530-En2.

The following compounds were prepared in a similar manner (use of appropriate reagents (chiral or racemic) and purification methods (including chiral HPLC or chiral SFC) known to the person skilled in the art) as described for cpd 530-En 1 & cpd 530-En 2: Cpd. 728-En 1, 728-En 2.

Synthesis of N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((3-(trifluoromethyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxamide (Cpd 730)



490



Step 1: To a stirred solution of ethyl 3-(trifluoromethyl)-1H-pyrazole-4-carboxylate (3.0 g, 14.42 mmol) in DMF (50 mL) was added K₂CO₃ (3.98 g, 28.84 mmol) at RT. The RM was cooled to 0° C., and added SEM-C₁ (3.0 mL, 17.30 mmol). The RM was stirred at 0° C. for 1 h and reaction progress was monitored by TLC. The reaction mixture was diluted with water (100 mL), and extracted with ethyl acetate (2×100 mL). Combined organic layers were washed with water (3×100 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to get a crude. Crude was purified by column chromatography using silica gel (100-200 mesh) and 10% ethyl acetate in pet ether as an eluent to afford ethyl 3-(trifluoromethyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazole-4-carboxylate (2.8 g, 58%) as a solid. TLC system: 10% ethyl acetate in pet ether; Rf: 0.6.

491

Step 2: To a stirred solution of ethyl 3-(trifluoromethyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazole-4-carboxylate (2.8 g, 8.28 mmol) in THF (50 mL) was added 1M LAH solution in THF (12.4 mL, 12.42 mmol) at 0° C. The RM was stirred at 0° C. for 2 h and the reaction progress was monitored by TLC. Reaction mixture was quenched with aq. sat. sodium sulphate solution at 0° C., diluted with ethyl acetate (50 mL), filtered through celite bed, and celite bed was washed with ethyl acetate (50 mL). Combined filtrate was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford (3-(trifluoromethyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-4-yl)methanol (2.5 g). TLC system: 10% ethyl acetate in pet ether; RF: 0.23.

Step 3: To a stirred solution of (3-(trifluoromethyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-4-yl)methanol (1.5 g, 5.06 mmol) in DCM (50 mL) was added triethylamine (1.4 mL, 10.13 mmol) and followed by addition of Mesyl Chloride (0.58 mL, 7.60 mmol) at 0° C. The RM was stirred for 16 h at RT and the reaction progress was monitored by TLC. The RM was diluted with water (40 mL) and extracted with DCM (3×50 mL). Combined extracts were washed with brine (20 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford (3-(trifluoromethyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-4-yl)methyl methanesulfonate (1.7 g) as a brown liquid. Proceeded to next step without further purification; TLC system: 10% ethyl acetate in pet ether; RF: 0.7.

Step 4: To a stirred solution of ethyl 5-hydroxy-2-methylbenzofuran-3-carboxylate (1.5 g, 6.81 mmol) in acetonitrile (100 mL) was added Cs₂CO₃ (6.6 g, 20.45 mmol) and followed by addition of (3-(trifluoromethyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-4-yl)methyl methanesulfonate (2.7 g, 7.49 mmol) at 0° C. The RM was stirred for 6 h at 80° C. and the reaction progress was monitored by TLC. The RM was filtered and washed the filter bed with ethyl acetate (100 mL). Filtrate was concentrated under reduced pressure to get crude. Crude was purified by column chromatography over silica gel (100-200 mesh) using 10% ethyl acetate in pet-ether as an eluent to afford ethyl 2-methyl-5-((3-(trifluoromethyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxylate (1.3 g, 59% over 3 steps) as an off-white solid. TLC system: 20% ethyl acetate in pet ether; RF: 0.7.

Step 5: To a stirred solution of ethyl 2-methyl-5-((3-(trifluoromethyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxylate (1.3 g, 2.61 mmol) in MeOH:THF:H₂O (1:1:0.5, 50 mL) was added NaOH (1.0 g, 26.10 mmol) at RT. The RM was stirred for 4 h at 50° C. and the reaction progress was monitored by TLC. The RM was concentrated, acidified to pH ~2 with aq. 1 N HCl solution, and stirred for 20 minutes. The precipitated solid was filtered, washed with water (10 mL) and dried under vacuum to afford 2-methyl-5-((3-(trifluoromethyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxylic acid (1.0 g) as an off-white solid. TLC system: 20% ethyl acetate in pet ether; RF: 0.1.

Step 6: To a stirred solution of 2-methyl-5-((3-(trifluoromethyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxylic acid (1.0 g, 2.12 mmol) in DMF (20 mL) was added HATU (1.21 g, 3.19 mmol), DIPEA (1.5 mL, 8.51 mmol) at 0° C. and followed by addition of ethyl 3-amino-2-oxopyrrolidine-3-carboxylate (548 mg, 3.19 mmol). The RM was stirred for 16 h at RT and reaction progress was monitored by TLC. The RM was diluted with water (100 mL), and extracted with ethyl acetate (2×100 mL). Combined organic layers were washed

492

with water (3×100 mL), brine (100 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford ethyl 3-(2-methyl-5-((3-(trifluoromethyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxamido)-2-oxopyrrolidine-3-carboxylate (1.0 g) as a brown residue. TLC system: 60% ethyl acetate in pet ether; RF: 0.2.


Step 7: To a stirred solution of ethyl 3-(2-methyl-5-((3-(trifluoromethyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxamido)-2-oxopyrrolidine-3-carboxylate (1.0 g, 1.60 mmol) in MeOH (20 mL) was added NaBH₄ (608 mg, 16.02 mmol) at 0° C. The RM was stirred for 16 h at RT and reaction progress was monitored by TLC. The RM was concentrated, diluted with water (50 mL), and extracted with ethyl acetate (2×100 mL). Combined organic layers were dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to get N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((3-(trifluoromethyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxamide (800 mg, crude) as a thick residue. TLC system: 60% ethyl acetate in pet-ether; RF: 0.2.

Step 8: To a stirred solution of N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((3-(trifluoromethyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxamide (800 mg, 1.37 mmol) in DCM (30 mL) was added TFA (5 mL) at 0° C. The RM was stirred for 6 h at RT and the reaction progress was monitored by TLC. The reaction mixture was concentrated, basified to pH ~8 with aq. sat NaHCO₃ solution and extracted with ethyl acetate (2×100 mL). Combined organic layers were dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to get crude. Crude was purified by Grace flash chromatography using 0.1% formic acid in acetonitrile as an eluent to afford N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((3-(trifluoromethyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxamide (Cpd 730) (500 mg, 34% for 4 steps).

A preparative chiral SFC was performed on the racemic mixture of cpd 730 to afford cpd 730-En 1 and cpd 730-En 2.


The following compounds were prepared in a similar manner (use of appropriate reagents (chiral or racemic) and purification methods (including chiral HPLC or chiral SFC) known to the person skilled in the art) as described for cpd 730-En 1 and cpd 730-En 2: Cpd. 731-En 1, 731-En 2.

Synthesis of N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((6-oxo-1,6-dihydropyridin-2-yl)methoxy)benzofuran-3-carboxamide (Cpd 569)



493

-continued



Step 1: To a pre-stirred solution mixture of 5-((6-methoxypyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid (1.0 g, 3.19 mmol), HATU (1.8 g, 4.79 mmol) and DIPEA (3.4 mL, 6.38 mmol) in DMF (20.0 mL) was added tert-butyl 4-amino-3,3-difluoropiperidine-1-carboxylate (0.904 g, 3.83 mmol) at 0° C. and the resulting reaction mixture was stirred at room temperature for 2 h. The reaction progress was monitored by TLC. (Note: Another 1.0 g of 5-((6-methoxypyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid was similarly reacted using the same procedure and was combined with the reaction described in detail for work-up). The combined reaction mixture was diluted with ice cold water (100 mL) and extracted with ethyl acetate (2×100 mL). The combined organic layers were washed with water (2×100 mL) and brine (100 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by GRACE flash chromatography using 0.1% formic acid in water and acetonitrile as an eluent to afford tert-butyl 3,3-difluoro-4-(5-((6-methoxypyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamido)piperidine-1-carboxylate (1.2 g, 36%) as an off-white solid. TLC system: 50% Ethyl acetate in pet ether; Rf: 0.4.

Step 2: HCl (4.0 M in dioxane) (10 mL) was added dropwise to a pre-stirred solution of tert-butyl 3,3-difluoro-4-(5-((6-methoxypyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamido)piperidine-1-carboxylate (600 mg, 1.12 mmol) in dichloromethane (5 mL) at 0° C. under an argon atmosphere. The resulting reaction mixture was allowed to attain room temperature and was stirred for 5 h. The reaction progress was monitored by TLC. (Note: 600

494

mg of tert-butyl 3,3-difluoro-4-(5-((6-methoxypyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamido)piperidine-1-carboxylate was similarly treated following the same procedure and was combined with the reaction mixture for work-up). The combined reaction mixture was concentrated under reduced pressure, the residue was basified with saturated NaHCO₃ (80 mL) solution and the organic compound was extracted with 10% methanol in dichloromethane (3×50 mL). The combined organic layers were washed with water (50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by GRACE flash chromatography using 0.1% formic acid in water and acetonitrile as an eluent to afford N-(3,3-difluoropiperidin-4-yl)-5-((6-methoxypyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide (900 mg, 92%) as an off-white solid. TLC system: 5% Methanol in dichloromethane; Rf: 0.2.



Step 3: TMS-C₁ (189 mg, 1.740 mmol) was slowly added to a pre-stirred mixture of N-(3,3-difluoropiperidin-4-yl)-5-((6-methoxypyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide (500 mg, 1.156 mmol) and NaI (347 mg, 2.32 mmol) in acetonitrile (10 mL) at room temperature and the resulting reaction mixture was heated to 60° C. for 24 h. The reaction progress was monitored by TLC. (Note: 400 mg of N-(3,3-difluoropiperidin-4-yl)-5-((6-methoxypyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide was similarly reacted following the same procedure and was combined with the reaction mixture for work-up). The combined reaction mixture was cooled to room temperature, quenched with brine solution (50 mL) and extracted with ethyl acetate (3×50 mL). The combined organic layers were dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by GRACE flash chromatography using 0.1% formic acid in water and acetonitrile as an eluent to afford N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((6-oxo-1,6-dihydropyridin-2-yl)methoxy)benzofuran-3-carboxamide (340 mg, 33%) as an off white solid. TLC system: 5% MeOH in dichloromethane; Rf: 0.3.

A preparative chiral SFC was performed on the racemic mixture of cpd 569 to afford cpd 569-En 1 and cpd 569-En 2.

The following compounds were prepared in a similar manner (use of appropriate reagents (chiral or racemic) and purification methods (including chiral HPLC or chiral SFC) known to the person skilled in the art) as described for cpd 569: Cpd. 564-En 1, 564-En 2, 569-En 1, 569-En 2, 637-En 1, 637-En 2, 640-En 1, 640-En 2.

Synthesis of 4-(((3-((4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)carbamoyl)-2-methylbenzofuran-5-yl)oxy)methyl)-1-methylene-1H-1,2,3-triazol-1-ium (Cpd 604) and N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((1-methyl-1H-1,2,3-triazol-5-yl)methoxy)benzofuran-3-carboxamide (Cpd 605)





Step 1: Methyl iodide (55.5 mL, 891.7 mmol) was added dropwise to a suspension of NaN_3 (27.8 g, 428.1 mmol) and prop-2-yn-1-ol (20.0 g, 356.7 mmol) in DMF (200 mL) at room temperature under an argon atmosphere. The reaction mixture was heated to 130° C. for 18 h. The reaction progress was monitored by TLC. The reaction mixture was cooled and treated with 200 mL of diethyl ether and the precipitate was removed by filtration. The filtrate was concentrated under reduced pressure. The crude compound was subjected to distillation at 130° C. to obtain a mixture of (1-methyl-1H-1,2,3-triazol-5-yl)methanol and (1-methyl-1H-1,2,3-triazol-4-yl)methanol (6.0 g, 15%) as a pale yellow liquid. TLC system: 5% Methanol in dichloromethane; Rf: 0.2.

Step 2: Thionyl chloride (4.7 mL 66.3 mmol) was added dropwise to a mixture of (1-methyl-1H-1,2,3-triazol-5-yl)methanol and (1-methyl-1H-1,2,3-triazol-4-yl)methanol (5.0 g, 44.2 mmol) in dichloromethane (50 mL) at 0° C. under an argon atmosphere. The reaction mixture was heated to 40° C. and was stirred for 4 h. Excess thionyl chloride and solvent were then evaporated in vacuo. The residue was cooled to 0° C., basified with saturated NaHCO_3 to pH~9 and extracted with dichloromethane (3×50 mL). The combined organic layer was washed with water (100 mL) and brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford a mixture of 5-(chloromethyl)-1-methyl-1H-1,2,3-triazole and 4-(chloromethyl)-1-methyl-1H-1,2,3-triazole (4.2 g, 71%) as a pale yellow liquid. TLC system: 50% Ethyl acetate in pet-ether; Rf: 0.5.

Step 3: NaH (60%) (1.9 g, 81.6 mmol) was added in small portions to a solution of ethyl 5-hydroxy-2-methylbenzofuran-3-carboxylate (6.0 g, 27.2 mmol) and a mixture of 5-(chloromethyl)-1-methyl-1H-1,2,3-triazole and 4-(chloromethyl)-1-methyl-1H-1,2,3-triazole (4.2 g, 32.7 mmol) in DMF (50 mL) at 0° C. under an argon atmosphere. The reaction mixture was warmed to room temperature and was stirred for 16 h. The reaction progress was monitored by TLC. The reaction mixture was diluted with ice water (100 mL) and extracted with ethyl acetate (2×100 mL). The combined organic layers were dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The crude product was purified by column chromatography over silica gel (100-200 mesh) using a gradient mixture of 0-40% ethyl acetate in pet-ether as an eluent to afford a mixture of ethyl 2-methyl-5-((1-methyl-1H-1,2,3-triazol-5-yl)methoxy)benzofuran-3-carboxylate (cpd 253) and ethyl 2-methyl-5-((1-methyl-1H-1,2,3-triazol-4-yl)methoxy)benzofuran-3-carboxylate (cpd 254) (2.2 g, 22%) as a pale yellow semi solid. TLC system: 70% Ethyl acetate in pet-ether; Rf: 0.2.


Step 4: A solution of NaOH (1.1 g, 27.9 mmol) in water (15 mL) was added to a pre-stirred solution of a mixture of ethyl 2-methyl-5-((1-methyl-1H-1,2,3-triazol-5-yl)methoxy)benzofuran-3-carboxylate (cpd 253) and ethyl 2-methyl-5-((1-methyl-1H-1,2,3-triazol-4-yl)methoxy)benzofuran-3-carboxylate (cpd 254) (2.2 g, 22%) as a pale yellow semi solid. TLC system: 70% Ethyl acetate in pet-ether; Rf: 0.2.

497

methoxy)benzofuran-3-carboxylate and ethyl 2-methyl-5-((1-methyl-1H-1,2,3-triazol-4-yl)methoxy)benzofuran-3-carboxylate (2.2 g, 6.98 mmol) in methanol:THF (1:1) (40 mL) at room temperature and the resulting reaction mixture was heated to 60° C. for 3 h. The reaction progress was monitored by TLC. The reaction mixture was cooled to room temperature and poured into ice cold water (100 mL), acidified with 1 N HCl to pH~2.0 and extracted with ethyl acetate (2x100 mL). The combined organic layers were washed with water (100 mL) followed by brine (100 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford a mixture of 2-methyl-5-((1-methyl-1H-1,2,3-triazol-5-yl)methoxy)benzofuran-3-carboxylic acid and 2-methyl-5-((1-methyl-1H-1,2,3-triazol-4-yl)methoxy)benzofuran-3-carboxylic acid (1.7 g, 85%) as a pale yellow solid. TLC system: 5% Methanol in dichloromethane; Rf: 0.2.


Step 5: (4-Aminotetrahydro-2H-pyran-4-yl)methanol (164.3 mg, 1.25 mmol) was added to a stirred solution of a mixture of 2-methyl-5-((1-methyl-1H-1,2,3-triazol-5-yl)methoxy)benzofuran-3-carboxylic acid and 2-methyl-5-((1-methyl-1H-1,2,3-triazol-4-yl)methoxy)benzofuran-3-carboxylic acid (300 mg, 1.04 mmol), HATU (790.8 mg, 2.08 mmol) and DIPEA (0.56 mL, 3.12 mmol) in DMF (10 mL). The resulting reaction mixture was stirred at room temperature for 2 h. The reaction progress was monitored by TLC. The reaction mixture was diluted with ice water (20 mL) and extracted with ethyl acetate (2x30 mL). The combined organic layers were dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by GRACE flash chromatography using 0.1% formic acid in water and acetonitrile as an eluent followed by chiral-SFC to afford N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((1-methyl-1H-1,2,3-triazol-5-yl)methoxy)benzofuran-3-carboxamide (cpd 605) (90 mg) and N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((1-methyl-1H-1,2,3-triazol-4-yl)methoxy)benzofuran-3-carboxamide (cpd 604) (50 mg) as off-white solids. TLC system: 70% Ethyl acetate in pet-ether; Rf: 0.4.

Synthesis of N3-(3,3-difluoropiperidin-4-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-2,3-dicarboxamide (Cpd 657)



498

-continued



Step 1: To a stirred solution of ethyl 2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (4 g, 10.5 mmol) in 1,4-dioxane (50 ml) was added SeO₂ (24 g, 21.1 mmol) at RT and the reaction mixture was stirred for 48 h at 120° C., the reaction progress was monitored by TLC. The reaction mixture was filtered through a celite pad and the filtrate was evaporated under reduced pressure to get the crude material (6 g), which was purified by silica gel (100-200 mesh) column chromatography to afford (1 g, 24%) as a pale yellow solid. TLC system: 30% Ethyl acetate in pet-ether; Rf: 0.4

Step 2: To a stirred solution of ethyl 2-formyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (1 g, 2.54 mmol) in ethanol (50 ml) was added NH₂OH·HCl (880 mg, 12.8 mmol) and pyridine (405 mg, 4.1 mmol) at RT and the resulting reaction mixture was then heated to 80° C. for 6 h. Ice cold water was added to the reaction mixture and the mixture was then extracted with

499

EtOAc (2×10 mL). The combined organic layers were washed with water (50 mL) and brine (20 mL), dried over anhydrous Na₂SO₄, filtered and evaporated under reduced pressure to get the crude material which was used in the next step without further purification. TLC system: 100% Ethyl acetate in pet-ether; Rf: 0.2

Step 3: To a stirred solution of (E)-N-(3-(ethoxycarbonyl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-2-yl)formimidic acid (1 g, 2.4 mmol) in THF (50 ml) were added DIPEA (0.722 g, 3 mmol) and T₃P (2.64 g, 3 mmol) at 0° C. and the resulting reaction mixture was stirred at RT for 16 h. Ice cold water was added to the reaction mixture and the mixture was extracted with EtOAc (2×10 mL). The combined organic layers were washed with water (50 mL) and brine (20 mL), dried over anhydrous Na₂SO₄, filtered and evaporated under reduced pressure to get the crude material which was purified by flash column chromatography over neutral alumina using 20% ethyl acetate in pet-ether as an eluent to afford ethyl 2-cyano-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (0.5 g, 52%) as an off-white solid. TLC system: 50% Ethyl acetate in pet-ether; Rf: 0.7

Step 4: To ethyl 2-cyano-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylate (550 mg, 1.4 mmol) was added LiOH·H₂O (110 mg, 2.8 mmol) in THF: H₂O (20 mL) at RT under an argon atmosphere. The reaction mixture was stirred for 2 h at RT. The reaction progress was monitored by TL C. The reaction mixture was acidified to pH ~1 with citric acid and the resulting precipitate was filtered off. The solid obtained was dried under vacuum to afford 2-cyano-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid (0.35 g, 68%) as an off-white solid. TLC system: 5% MeOH in CH₂Cl₂; Rf: 0.2.


Step 5: To a stirred solution of 2-cyano-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid (0.35 g, 9 mmol) in CH₂Cl₂ (30 ml) were added DIPEA (0.23 g, 1.8 mmol) and HATU (0.69 g, 1.8 mmol) at 0° C. followed by the addition of tert-butyl 4-amino-3,3-difluoropiperidine-1-carboxylate (273 mg, 1.1 mmol), then the reaction mixture was stirred for 6 h at RT, and the reaction progress was monitored by TLC. The reaction mixture was diluted with water (150 mL), and extracted with CH₂Cl₂ (3×100 mL). The combined organic extracts were washed with brine (150 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to get the crude compound (0.5 g) which was used without further purification in the next step.

Step 6: To a stirred solution of tert-butyl 4-(2-cyano-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)-3,3-difluoropiperidine-1-carboxylate (25 mg, 0.043 mmol) in 1,4-dioxane (10 ml), were added 4 M HCl in 1,4-dioxane (5 ml) at 0° C. and the mixture was stirred for 2 h at RT, the reaction progress was monitored by TLC. The reaction mixture was concentrated under reduced pressure to get the crude compound (30 mg, LCMS: 88%). The crude compound was purified by prep-SFC to afford of (12 mg) as an off-white solid. TLC system: 10% Methanol in Dichloromethane; Rf: 0.3.

Cpd 657 can be obtained in enantiomerically pure form as cpd 657-En 1 and cpd 657-En 2 using a chiral SFC separation of the product obtained after step 5. Both obtained enantiomers are then forwarded independently to obtain cpd 657-En 1 and cpd 657-En 2.


500

Synthesis of N3-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-2,3-dicarboxamide (Cpd 658)



501

-continued



Step 1: PTSA (0.051 g, 0.3 mmol) was added to a stirred solution of ethyl 5-(benzyloxy)-2-formylbenzofuran-3-carboxylate (1 g, 3 mmol) and ethylene glycol (0.925 g, 15.4 mmol) in toluene (20 mL) at RT. The resulting reaction mixture was stirred at 120° C. for 12 h using a Dean-stark apparatus. The reaction mixture was evaporated under reduced pressure to get a residue. The residue obtained was dissolved in ethyl acetate (20 mL) and washed with sat. NaHCO₃ solution (100 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered, and evaporated under reduced pressure to get the crude material which was purified by neutral alumina column chromatography using 20% ethyl acetate in pet ether to afford ethyl 5-(benzyloxy)-2-(1,3-dioxolan-2-yl)benzofuran-3-carboxylate (900 mg, 81%) as an off-white solid. TLC system: 20% EtOAc in Hexane; Rf: 0.4.

Step 2: To a stirred solution of ethyl 5-(benzyloxy)-2-(1,3-dioxolan-2-yl)benzofuran-3-carboxylate (0.9 g, 2.4 mmol) in ethanol (20 mL) was added 10% Pd/C (150 mg, 10 mol %), and then the reaction mixture was stirred for 6 h under H₂ atm pressure (60 psi) at RT, the reaction progress was monitored by TLC. The reaction mixture was filtered through a celite pad and the filtrate was evaporated under reduced pressure to get the crude material (1 g) as a colorless oil, which was used in the next step without further purification. TLC system: 40% Ethyl acetate in pet-ether; Rf: 0.5.

Step 3: (4-Methylthiazol-5-yl)methanol (2044-6) (0.229 g, 1.7 mmol), ADPP (0.604 g, 2.4 mmol) and tri-N-butylphosphine (0.484 g, 2.4 mmol) were added sequentially to a pre-stirred solution of ethyl 2-(1,3-dioxolan-2-yl)-5-hydroxybenzofuran-3-carboxylate (0.45 g, 1.6 mmol) in THF (15 mL) at RT under an argon atmosphere. The reaction mixture was stirred for 18 h at RT and the reaction progress was monitored by TLC. The reaction mixture was filtered and washed with diethyl ether (300 mL), the filtrate was concentrated under reduced pressure to give the crude material which was purified by flash column chromatography over neutral alumina using 10% ethyl acetate in pet-ether as an eluent to afford ethyl 2-(1,3-dioxolan-2-yl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate (0.2 g, 30%) as an off-white solid. TLC system: 30% Ethyl acetate in pet-ether; Rf: 0.35

Step 4: To a stirred solution of ethyl 2-(1,3-dioxolan-2-yl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate (100 mg, 2.57 mmol) in CH₂Cl₂ (10 mL) was added TFA (58 mg, 5.14 mmol) at 0° C. and the resulting reaction mixture was stirred at rt for 16 h. After completion of the reaction, ice-cold water was added to the reaction mixture and the mixture was extracted with CH₂Cl₂ (2×10 mL). The combined organic layers were washed with sat. NaHCO₃ solution (50 mL), dried over anhydrous Na₂SO₄, filtered and evaporated under reduced pressure to get the crude material,

502


which was used in the next step without further purification. TLC system: 50% Ethyl acetate in pet-ether; Rf: 0.3

Step 5: To a stirred solution of ethyl 2-formyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate (50 mg, 0.14 mmol) in ethanol (10 mL) were added NH₂OH·HCl (39 mg, 5 mmol) and pyridine (22 mg, 2.8 mmol) at RT and the resulting mixture was stirred at 90° C. for 6 h. After completion of the reaction ice-cold water was added to the reaction mixture and the mixture was extracted with CH₂Cl₂ (2×10 mL). The combined organic layers were washed with sat. NaHCO₃ solution (50 mL), dried over anhydrous Na₂SO₄, filtered and evaporated under reduced pressure to get the crude material, which was used in the next step without further purification. TLC system: 100% Ethyl acetate in pet-ether; Rf: 0.2.

Step 6: To a stirred solution of (E)-ethyl 2-((hydroxyimino)methyl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate (1 g, 2.8 mmol) in THF (50 mL) were added DIPEA (0.722 g, 3 mmol) and T₃P (2.64 g, 3 mmol) at 0° C. and the resulting reaction mixture was then stirred at RT for 16 h. Ice cold water was added to the reaction mixture and the mixture was extracted with EtOAc (2×10 mL). The combined organic layers were washed with brine (20 mL), dried over anhydrous Na₂SO₄, filtered and evaporated under reduced pressure to get the crude material which was purified by flash column chromatography over neutral alumina using 20% ethyl acetate in pet-ether as an eluent to afford ethyl 2-cyano-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate (0.3 g, 31%) as an off-white solid. TLC system: 50% Ethyl acetate in pet-ether; Rf: 0.6.

Step 7: To a stirred solution of ethyl 2-cyano-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylate (0.15 g, 4 mmol) in THF (50 mL) were added (4-aminotetrahydro-2H-pyran-4-yl)methanol (0.068 g, 5.2 mmol) and 2,3,4,6,7,8-hexahydro-1H-pyrimido[1,2-a]pyrimidine (0.108 g, 7.8 mmol) at RT and the resulting reaction mixture was heated to 80° C. for 40 min. Ice cold water was added to the reaction mixture and the mixture was extracted with EtOAc (2×10 mL). The combined organic layers were washed with brine (20 mL), dried over anhydrous Na₂SO₄, filtered and evaporated under reduced pressure to get the crude compound, which was purified by reverse phase prep-HPLC purification to afford 13 mg of cpf 658 as an off-white solid. TLC system: 50% Ethyl acetate in pet-ether; Rf: 0.2.


Synthesis of Small Synthetic Intermediates Used for the Synthesis of Cpd. 001-666



503


Step 1: A solution of methyl S-methyl-L-cysteinate (1) (2.0 g, 0.013 mmol) and 7 M NH_3 solution in methanol (20 mL) in a sealed tube was heated to 90°C . and stirred for 16 h at the same temperature. The reaction progress was monitored by TLC. The reaction mixture was concentrated under reduced pressure, and was then co-distilled with EtOAc (2x20 mL). To the reaction mixture was added EtOAc (30 mL), the reaction mixture was then stirred for 2 h, and the solid was filtered off and dried to afford (R)-2-amino-3-(methylthio)propanamide (1.6 g, 88%) as a brown solid. TLC system: 5% Methanol in dichloromethane; Rf: 0.1

Step 2: To a stirred solution of (R)-2-amino-3-(methylthio) propanamide (1.0 g, 0.007 mmol) in methanol and water (2:1), was added oxone (5.0 g, 0.008 mmol) at RT. The reaction mixture was stirred for 16 h at RT. The reaction mixture was concentrated under reduced pressure, and was then co-distilled with methanol (2x20 mL) to afford (R)-2-amino-3-(methylsulfonyl)propanamide (6.0 g crude) as a white solid which was used without any further purification.




Step 1: To a stirred solution of methyl S-methyl-L-cysteinate (1.0 g, 6.71 mmol) and THF (20 mL) was added 1 M LAH solution in THF (8 mL, 8.0 mmol) at 0°C . The reaction mixture was stirred for 1 h at 0°C . The reaction progress was monitored by TLC. Then the reaction mixture was quenched with saturated Na_2SO_4 solution and was filtered. The filtrate was concentrated under reduced pressure to afford (R)-2-amino-3-(methylthio)propan-1-ol (800 mg, crude) as a thick residue. TLC system: 10% Methanol in dichloromethane; Rf: 0.1

Step 2: To a stirred solution of (R)-2-amino-3-(methylthio)propan-1-ol (800 mg, 6.61 mmol) in methanol and water (2:1) was added oxone (4.4 g, 7.27 mmol) at RT. The reaction mixture was stirred for 16 h at RT. The reaction mixture was concentrated under reduced pressure, and was then co-distilled with methanol (2x20 mL) to afford (R)-2-amino-3-(methylsulfonyl)propan-1-ol (5.1 g crude) as a white solid, which was used without any further purification.




504



Step 1: EDC-HCl (1.3 g, 4.87 mmol), HOBT (0.98 g, 7.31 mmol), DIPEA (2.5 mL, 14.63 mmol) and 3,3-difluoropyrrolidine (0.837 g, 5.85 mmol) were added sequentially to a stirred solution of (tert-butoxycarbonyl)-L-serine (1.0 g, 4.87 mmol) at room temperature. The resulting reaction mixture was stirred at room temperature for 16 h. The reaction progress was monitored by TLC. The reaction mixture was diluted with ice cold water (100 mL) and the precipitated solid was filtered and dried under vacuum to afford t-butyl (S)-(1-(3,3-difluoropyrrolidin-1-yl)-3-hydroxy-1-oxopropan-2-yl)carbamate (650 mg, crude) as a yellow solid. TLC system: 30% Ethyl acetate in pet-ether; Rf: 0.5.

Step 2: 4M HCl in 1,4-dioxane (6.5 mL) was added to t-butyl (S)-(1-(3,3-difluoropyrrolidin-1-yl)-3-hydroxy-1-oxopropan-2-yl)carbamate (0.650 g, 2.210 mmol) at room temperature under an argon atmosphere. The resulting reaction mixture was stirred at room temperature for 30 min. The reaction progress was monitored by TLC. The reaction mixture was concentrated under reduced pressure to afford ((S)-2-amino-1-(3,3-difluoropyrrolidin-1-yl)-3-hydroxypropan-1-one (450 mg, crude) as an off-white solid. TLC system: 30% ethyl acetate in pet-ether; Rf: 0.3.




Step 1: To a stirred solution of 2-(benzyloxycarbonylamino)-3-hydroxypropanoic acid (2 g, 0.3 mmol) and methanamine hydrochloride (0.616 g, 9.2 mmol) in DCM (30 mL) was added DIPEA (0.077 g, 6 mmol) followed by HATU (0.228 g, 6 mmol) at 0°C . and the resulting mixture was stirred at RT for 18 h. Ice cold water was added to the reaction mixture and the mixture was extracted with DCM (2x50 mL). The combined organic layers were washed with water (50 mL) and brine (50 mL), dried over anhydrous Na_2SO_4 , filtered and evaporated under reduced pressure to get the crude material, which was purified by neutral alumina column chromatography using 5% methanol in CHCl_3 as eluent. All appropriate fractions were combined and evaporated under reduced pressure to get benzyl 3-hydroxy-

505

1-(methylamino)-1-oxopropan-2-ylcarbamate (1 g) as pale yellow solid. TLC system: 20% Methanol in CHCl_3 ; Rf: 0.3

Step 2: To a stirred solution of benzyl 3-hydroxy-1-(methylamino)-1-oxopropan-2-ylcarbamate (1 g, 3.9 mmol) in methanol (50 mL) was added Pd/C (200 mg, 10 mol %) under a nitrogen atmosphere and the reaction mixture was stirred under H_2 pressure of 50 psi for 16 h at RT. The reaction mixture was then filtered through a pad of celite and the celite bed was washed with methanol (40 mL). The filtrate was evaporated under reduced pressure to afford crude 2-amino-3-hydroxy-N-ethylpropanamide (0.35 g) as a colorless semi-solid residue. The crude product thus obtained was used in the next step without further purification. TLC system: 20% Ethyl acetate in pet-ether; Rf: 0.1.




Step 1: At 0°C ., ethyl 2-chloro-2-oxoacetate (3.6 mL, 30.985 mmol) was added drop-wise to a solution of cyclobutylamine (2.0 g, 28.169 mmol) and Et_3N (8.6 mL, 61.971 mmol) in diethyl ether (20 mL). The resulting reaction mixture was stirred at 0°C . to RT for 1 h. The reaction progress was monitored by TLC. The reaction mixture was filtered and the filtrate washed with diethyl ether (2x100 mL). The filtrate was concentrated under reduced pressure to afford ethyl 2-(cyclobutylamino)-2-oxoacetate (3.0 g, 62%) as a brown oil. This crude compound was used without further purification. TLC system: 10% MeOH in dichloromethane; Rf: 0.5.

Step 2: A solution of LAH (1 M in THF, 35 mL, 35.08 mmol) was added to a suspension of ethyl 2-(cyclobutylamino)-2-oxoacetate (3.0 g, 17.54 mmol) in THF at 0°C . and the resulting reaction mixture was stirred at 0°C .-RT for 3 h. The reaction was monitored by TLC. The reaction mixture was quenched with aq. 1M NaOH solution (5 mL) at 0°C . under stirring. The solids were filtered off, and the filter cake was washed with 10% MeOH in dichloromethane. The filtrate was concentrated under reduced pressure to afford N-cyclobutyl-2-hydroxyacetamide (1.8 g, 79%) as a pale yellow solid. This compound was used in the next step without any further purification. TLC system: 10% MeOH in dichloromethane; Rf: 0.3.

Step 3: LiAlH_4 (1M solution in THF) (16.7 mL, 16.74 mmol) was added to a suspension of N-cyclobutyl-2-hydroxyacetamide (1.8 g, 13.95 mmol) in THF at 0°C . and the resulting reaction mixture was heated to 70°C . for 16 h. The reaction was monitored by TLC. The reaction mixture was quenched with aq. 1M NaOH solution (5 mL) at 0°C . under stirring, the precipitated solids were filtered off, the filter cake was washed with 10% MeOH in dichloromethane. The combined filtrate was dried over anhyd. Na_2SO_4 and concentrated under reduced pressure to afford 2-(cyclobuty-

506

lamino)ethan-1-ol (1.0 g, 62%) as a brown gummy solid. TLC system: 10% Methanol in dichloromethane; Rf: 0.2



Step 1: To a pre-stirred mixture of benzene-1,2-diamine (1.0 g, 9.250 mmol), HATU (4.210 g, 11.10 mmol) and DIPEA (4.8 mL, 27.75 mmol) in DMF (10.0 mL) was added O-benzyl-N-(tert-butoxycarbonyl)-D-serine (3.0 g, 10.175 mmol) at 0°C . and the resulting reaction mixture was stirred at room temperature for 16 h. The reaction progress was monitored by TLC. The reaction mixture was diluted with water (75 mL) and extracted with ethyl acetate (2x80 mL). The combined extracts were washed with brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated to give the crude material which was purified by column chromatography over silica-gel (100-200 mesh) eluting with 25% ethyl acetate in pet-ether as an eluent to afford tert-butyl (S)-1-((2-aminophenyl)amino)-3-(benzyloxy)-1-oxopropan-2-ylcarbamate (1.1 g) as a yellow semi-solid. TLC system: 5% Methanol in dichloromethane; Rf: 0.1.


Step 2: A solution of tert-butyl (S)-1-((2-aminophenyl)amino)-3-(benzyloxy)-1-oxopropan-2-ylcarbamate (500 mg, 1.298 mmol) and acetic acid (5.0 mL) was stirred at 80°C . for 1 h. The reaction progress was monitored by TLC. The reaction mixture was concentrated under reduced pressure to afford tert-butyl (R)-1-(1H-benzo[d]imidazol-2-yl)-2-(benzyloxy)ethylcarbamate (400 mg, crude) as a colorless semi solid. This crude compound was directly used in the next step. TLC system: 5% methanol in dichloromethane; Rf: 0.7.

507

Step 3: A solution of tert-butyl (R)-1-(1H-benzo[d]imidazol-2-yl)-2-(benzyloxy)ethylcarbamate (2.3 g, 6.259 mmol) and HCl (4 M in dioxane) (23.0 mL) was stirred at room temperature for 1 h. The reaction progress was monitored by TLC. The reaction mixture was concentrated under reduced pressure to afford (R)-1-(1H-benzo[d]imidazol-2-yl)-2-(benzyloxy)ethan-1-amine hydrochloride (1.8 g, crude) as a pale yellow semi-solid. This crude was directly used in the next step. TLC system: 5% Methanol in dichloromethane; Rf: 0.3.

Step 4: 10% Pd—C (230 mg) was added to a stirred solution of (R)-1-(1H-benzo[d]imidazol-2-yl)-2-(benzyloxy)ethan-1-amine hydrochloride (0.500 g, 1.872 mmol) in ethyl acetate (30 mL) and concentrated HCl (0.5 mL) at room temperature. The resulting reaction mixture was stirred at room temperature under hydrogen gas balloon pressure for 16 h. The reaction progress was monitored by TLC. The reaction mixture was filtered through a celite bed. The filtrate was concentrated under reduced pressure to afford (R)-2-amino-2-(1H-benzo[d]imidazol-2-yl)ethan-1-ol hydrochloride (400 mg, crude) as an off-white solid. TLC system: 30% ethyl acetate in pet-ether; Rf: 0.2.

(S)-2-amino-2-(1H-benzo[d]imidazol-2-yl)ethan-1-ol hydrochloride can be prepared using the same synthetic sequence starting from O-benzyl-N-(tert-butoxycarbonyl)-L-serine.




Step 1: To a pre-stirred solution of methyl 1-methyl-1H-pyrazole-5-carboxylate (5 g, 35.71 mmol) in acetonitrile (200 mL) was added selectfluor (28.72 g, 42.85 mmol) and acetic acid (5 mL) at 0° C. under an argon atmosphere. The reaction mixture was heated up to 120° C. for 24 h. The reaction progress was monitored by TLC. The excess solvent was removed under reduced pressure and the crude product was dissolved in ethyl acetate (500 mL). The ethyl acetate layer was washed with water (250 mL) and brine (150 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to give the crude material, which was purified by silica gel (100-200 mesh) column chromatography using 0-20% ethyl acetate in pet-ether as an eluent to afford methyl 4-fluoro-1-methyl-1H-pyrazole-5-carboxylate (0.8 g, 14%) as an off-white solid. TLC system: 20% EtOAc in Pet ether; Rf: 0.4.


Step 2: To a stirred solution of methyl 4-fluoro-1-methyl-1H-pyrazole-5-carboxylate (0.8 g, 5.06 mmol 1 eq) in THF (30 mL) at 0° C., was added LAH (1M) (6 mL, 6.07 mmol, 1.2 eq) solution in THF drop wise at 0° C. under an argon atmosphere. The reaction mixture was stirred at room temperature for 2 h. The reaction progress was monitored by TLC. The reaction mixture was quenched with sat. Na₂SO₄ solution and extracted with 10% methanol in dichloromethane (2x50 mL). The combined organic layers were washed with water (20 mL) and brine (20 mL), dried over anhydrous

508

Na₂SO₄ and concentrated under reduced pressure to give the crude material, which was purified by column chromatography over silica gel (100-200 mesh) using 0-20% ethyl acetate in pet-ether as an eluent to afford (4-fluoro-1-methyl-1H-pyrazol-5-yl)methanol (0.45 g, 69%) as a colorless liquid. TLC system: 20% EtOAc in Pet ether; Rf: 0.3.



Step 1: To a stirred solution of (3-(trifluoromethyl)-1H-pyrazol-4-yl)methanol (600 mg, 3.61 mmol, 1.0 eq) and dichloromethane (50 mL) was added phosphorous tribromide (0.51 mL, 5.42 mmol, 1.5 eq) at 0° C. under an argon atmosphere. The reaction mixture was stirred for 2 h at 0° C. The reaction progress was monitored by TLC. The reaction mixture was diluted with ice water (100 mL) and extracted with dichloromethane (2x50 mL). The combined organic layers were washed with water (50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford 4-(bromomethyl)-3-(trifluoromethyl)-1H-pyrazole (430 mg, 52%) as a thick residue. TLC system: 5% methanol in dichloromethane; Rf: 0.7.




Step 1: To a stirred solution of 2-oxo-1,2-dihydropyridine-3-carbaldehyde (10 g, 81.3 mmol, 1.0 eq) in DMF (10 mL) was added Cs₂CO₃ (31.7 g, 97.56 mmol, 1.2 eq) followed by MeI (6.07 mL, 97.56 mmol, 1.2 eq) at 0° C. The reaction mixture was stirred for 3 h at RT and the reaction progress was monitored by TLC. After completion of the reaction, the mixture was diluted with water (30 mL) and extracted with EtOAc (3x50 mL). The combined organic layers were washed with water (20 mL) and brine (20 mL), dried over anhydrous Na₂SO₄, filtered and concentrated to afford 1-methyl-2-oxo-1,2-dihydropyridine-3-carbaldehyde (6.8 g, 61%, off-white solid). TLC system: 40% EtOAc/Pet ether; Rf: 0.4.

Step 2: To a stirred solution of 1-methyl-2-oxo-1,2-dihydropyridine-3-carbaldehyde (6.8 g, mmol, 1.0 eq) in methanol (10 mL) was added NaBH₄ (208.6 mg, 6.347 mmol, 1.2 eq) at 0° C. The reaction mixture was stirred for 3 h at RT and the reaction progress was monitored by TLC. After completion of the reaction, the reaction mixture was diluted with water (30 mL) and extracted with EtOAc (3x50 mL). The combined organic layers were washed with water (20 mL) and brine (20 mL), dried over anhydrous Na₂SO₄,

509

filtered and concentrated to get the crude material which was purified by Grace flash column chromatography to afford 3-(hydroxymethyl)-1-methylpyridin-2(1H)-one (3.5 g, 52%, off-white solid). TLC system: 40% EtOAc/Pet ether; Rf: 0.4.

Step 3: To a stirred solution of 3-(hydroxymethyl)-1-methylpyridin-2(1H)-one (600 g, 4.37 mmol, 1.0 eq) in CH_2Cl_2 (15 ml) was added Et_3N (1.5 ml, 10.86 mmol, 2.5 eq) followed by methanesulfonyl chloride (0.5 ml, 6.52 mmol, 1.5 eq) at 0°C . The reaction mixture was stirred for 3 h at RT and the reaction progress was monitored by TLC. After completion of the reaction, the reaction mixture was diluted with water (30 ml) and extracted with EtOAc (3×50 ml). The combined organic layers were washed with water (20 ml) and brine (20 ml), dried over anhydrous Na_2SO_4 , filtered and concentrated to get (1-methyl-2-oxo-1,2-dihydropyridin-3-yl)methyl methanesulfonate (365 mg crude, yellow color liquid). TLC system: 40% EtOAc/Pet ether; Rf: 0.75. Note: (1-Methyl-2-oxo-1,2-dihydropyridin-3-yl) methyl methanesulfonate was unstable even in work up and distillation at RT under vacuum. Crude was directly taken to next step.



Step 1: Thionyl chloride (5 mL) was added dropwise to a solution of 6-oxo-1,6-dihydropyridine-2-carboxylic acid (12.0 g, 86.26 mmol) in ethanol (300 mL) at 0°C . under an argon atmosphere. The reaction mixture was warmed to room temperature and stirred for 16 h. The excess of thionyl chloride and solvent was evaporated in vacuo and the residue was cooled to 0°C ., the remains were basified with saturated NaHCO_3 to pH ~9 and the resulting mixture was extracted with ethyl acetate (3×200 mL). The combined organic layer was washed with water (200 mL) and brine (200 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford ethyl 6-oxo-1,6-dihydro-


510

pyridine-2-carboxylate (8.0 g, 57%) as an off-white solid. TLC system: 50% Ethyl acetate in pet-ether; Rf: 0.5.


Step 2: Methyl iodide (8.1 g, 57.48 mmol) was added to a suspension of ethyl 6-oxo-1,6-dihydropyridine-2-carboxylate (8.0 g, 47.90 mmol) and Cs_2CO_3 (46.7 g, 143.71 mmol) in DMF (100 mL) at 0°C . under an argon atmosphere. The reaction mixture was warmed to room temperature and was stirred for 5 h. The reaction progress was monitored by TLC. The reaction mixture was diluted with ice cold water (200 mL) and extracted with ethyl acetate (3×100 mL). The combined organic layers were dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The crude product was purified by column chromatography over silica-gel (100-200 mesh) using a solvent gradient mixture of 0-30% ethyl acetate in pet-ether to obtain ethyl 1-methyl-6-oxo-1,6-dihydropyridine-2-carboxylate (2.0 g) & ethyl 6-methoxypicolinate (2.5 g, 52%) as pale yellow solid.

Step 3: NaBH_4 (1.25 g, 33.14 mmol) was added in small portions to a solution of ethyl 1-methyl-6-oxo-1,6-dihydropyridine-2-carboxylate (2.0 g, 11.04 mmol) in t-butanol (50 mL) at room temperature. The reaction mixture was heated to 60°C ., and was then stirred for 1 h before being rapidly cooled to room temperature. Methanol (2.5 mL) was added dropwise to the reaction mixture was then heated to 60°C . for 3 h. The reaction progress was monitored by TLC. The reaction mixture was cooled to 0°C . and was diluted with water (100 mL). The mixture was extracted with 10% methanol in dichloromethane (2×100 mL). The combined organic layers were washed with brine (100 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford 6-(hydroxymethyl)-1-methylpyridin-2(1H)-one (1.2 g 80%) as a pale yellow thick mass. TLC system: 100% Ethyl acetate; Rf: 0.2.

Step 4: Methanesulfonyl chloride (1.2 mL) was added dropwise to a solution mixture of 6-(hydroxymethyl)-1-methylpyridin-2(1H)-one (1.2 g, 8.63 mmol) and TEA (3.5 mL, 25.89 mmol) in dichloromethane (30 mL) at 0°C . under an argon atmosphere. The reaction mixture was warmed to room temperature and was stirred for 3 h. The reaction progress was monitored by TLC. The solvents were evaporated in vacuo and the residue was cooled to 0°C ., basified with saturated NaHCO_3 to a pH ~8 and extracted with ethyl acetate (3×50 mL). The combined organic layers were washed with water (50 mL) and brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford (1-methyl-6-oxo-1,6-dihydropyridin-2-yl)methyl methanesulfonate (2.4 g, crude) as a pale yellow thick mass. The resulting crude product was used for next step without further purification. TLC system: 30% Ethyl acetate in pet-ether; Rf: 0.5



511



Step 1: To a stirred solution of 2-hydroxynicotinic acid (3.0 g, 21.58 mmol) in ethanol (50 mL) was added SOCl_2 (21 mL) at room temperature, and reaction mixture was heated to 90°C . for 8 h. The reaction progress was monitored by TLC. The mixture was cooled to RT, the volatiles were evaporated under reduced pressure and ice-cold water (400 mL) was added to the reaction mixture. The pH was adjusted to $\sim 7-8$ with NaHCO_3 powder and the mixture was then extracted with chloroform (3×500 mL). The combined extracts were washed with brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to get ethyl 2-hydroxynicotinate (1.0 g, 27%, crude) as an off-white solid. TLC system: 100% Ethyl acetate; RF: 0.43. The material was used in the next step without further purification.


Step 2: To a stirred solution of ethyl 2-hydroxynicotinate (500 mg, 2.99 mmol) in acetonitrile (10 mL) was added 6M KOH (6 mL) followed by the addition of difluoromethyl trifluoromethanesulfonate (1.7 g, 8.98 mmol) at 0°C . under an argon atmosphere. The reaction mixture was stirred for 16 h at RT. The reaction progress was monitored by TLC. Solvents were removed under vacuum and ice cold water was added to the reaction mixture. The mixture was then extracted with ethyl acetate (2×100 mL). The combined extracts were washed with brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford crude 2-(difluoromethoxy) nicotinic (500 mg) as a yellow solid, which was used in the next step without further purification. TLC system: 10% MeOH in CH_2Cl_2 ; RF: 0.38.

Step 3: To a stirred solution of 2-(difluoromethoxy) nicotinic acidoxylate (500 mg, 2.64 mmol) in THF (15 mL) was added NaBH_4 (201 mg, 5.29 mmol) followed by the addition of BF_3 etherate (0.7 mL, 5.29 mmol) at 0°C . The reaction mixture was heated to 80°C . for 16 h. The reaction progress was monitored by TLC. The reaction mixture was concentrated, diluted with H_2O (50 mL) and extracted with ethyl acetate (3×50 mL). The combined extracts were washed with brine (20 mL), dried over anhydrous Na_2SO_4 , filtered and concentrated under vacuum to get the crude material, which was purified by flash chromatography to afford (2-(difluoromethoxy) pyridin-3-yl) methanol (100 mg, 19% over 2 steps) as a colorless liquid. TLC system: 80% Ethyl acetate in pet ether; RF: 0.56.

Step 4: To a stirred solution of (2-(difluoromethoxy) pyridin-3-yl) methanol (100 mg, 0.57 mmol) in dichloromethane (10 mL) was added Et_3N (0.1 mL, 0.85 mmol) followed by the addition of MsCl (98 mg, 0.85 mmol) at 0°C . The reaction mixture was stirred for 16 h at RT and the reaction progress was monitored by TLC. The reaction


512

mixture was diluted with water (50 mL) and extracted with ethyl acetate (3×50 mL). The combined extracts were washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford crude (2-(difluoromethoxy) pyridin-3-yl) methyl methanesulfonate (150 mg) as a brown solid. TLC system: 50% EtOAc in pet-ether; RF: 0.62. Crude was used in the next step without further purification and analytical data.




Step 1: To a stirred solution of 2-methylnicotinaldehyde (1.0 g, 8.26 mmol) in dichloromethane (50 mL) was added DAST (1.63 mL, 12.3 mmol) at 0°C . and reaction mixture was stirred for 2 h at RT. The reaction progress was monitored by TLC, reaction mass was diluted with dichloromethane (100 mL) and washed with water (2×50 mL) followed by brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to get the crude material, which was purified by Grace flash chromatography using 20% ethyl acetate in pet ether as an eluent to afford 3-(difluoromethyl)-2-methylpyridine (500 mg, 42%) as a thick colorless liquid. TLC system: 30% Ethyl acetate in pet-ether; RF: 0.3.

Step 2: To a stirred solution of 3-(difluoromethyl)-2-methylpyridine (500 mg, 3.49 mmol) in carbon tetrachloride (30 mL) were added NBS (742 mg, 4.19 mmol) and AIBN (58 mg, 0.349 mmol) at 0°C . under an argon atmosphere. The reaction mixture was stirred for 16 h at 80°C . The reaction progress was monitored by TLC. The reaction mass was diluted with dichloromethane (100 mL) and was washed with water (2×50 mL) followed by brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to get the crude material, which was purified by Grace flash chromatography using 15% ethyl acetate in pet ether as an eluent to afford 2-(bromomethyl)-3-(difluoromethyl)pyridine (150 mg, 20%). TLC system: 30% Ethyl acetate in pet-ether; RF: 0.6




513

-continued



Step 1: To a stirred solution of methyl 1H-pyrazole-5-carboxylate (1.5 g, 11.90 mmol) in CH₃CN (50 mL) was added Cs₂CO₃ (11.6 g, 35.712 mmol) followed by the addition of 2,2-difluoroethyl trifluoromethanesulfonate (3.8 g, 17.857 mmol) at RT and the reaction mixture was then heated to 70° C. for 2 h. The reaction progress was monitored by TLC. The reaction mixture was diluted with water (100 mL) and was extracted with ethyl acetate (3×100 mL). The combined extracts were washed with brine (30 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to get the crude material, which was purified by flash chromatography using 10% ethyl acetate in pet ether as eluent to afford methyl 1-(2,2-difluoroethyl)-1H-pyrazole-5-carboxylate (1 g) as a colorless liquid. (TLC: 40% Ethyl acetate in pet ether: R_f=0.68)

Step 2: To a stirred solution of methyl 1-(2,2-difluoroethyl)-1H-pyrazole-5-carboxylate (1 g, 5.263 mmol) in MeOH (30 mL), was added NaBH₄ (600 mg, 15.789 mmol) at 0° C. and the reaction mixture was stirred at reflux for 16 h. The reaction progress was monitored by TLC. The reaction mixture was concentrated, diluted with water (120 mL) and extracted with ethyl acetate (3×100 mL). The combined extracts were washed with brine (20 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford (1-(2,2-difluoroethyl)-1H-pyrazol-5-yl) methanol (900 mg, crude) as a colorless liquid. The obtained crude material was used in the next step without further purification. (TLC: 100% Ethyl acetate: R_f=0.2).




Step 1: To a stirred solution of methyl 1-methyl-1H-pyrazole-5-carboxylate (700 mg, 5.00 mmol) in Acetic acid (15 mL) was added NaOCl (2.2 g, 30.00 mmol) at 0° C. The RM was stirred for 2 h at RT and the reaction progress was monitored by TLC. The RM was concentrated under reduced pressure, diluted with water (50 mL) and extracted with ethyl acetate (3×50 mL). Combined extracts were washed with brine (20 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified by column chromatography by using 5% ethyl acetate in pet ether as eluents.

Step 2: To a stirred solution of methyl 4-fluoro-1-methyl-1H-pyrazole-5-carboxylate (0.8 g, 5.06 mmol 1 eq) in THF

514

(30 mL) at 0° C., was added LAH (1M) (6 mL, 6.07 mmol, 1.2 eq) solution in THF drop wise at 0° C. under an argon atmosphere. The RM was stirred at RT for 2 h. The reaction progress was monitored by TLC. The reaction mixture was quenched with sat. Na₂SO₄ solution and extracted with 10% methanol in dichloromethane (2×50 mL). The combined organic layers were washed with water (20 mL) and brine (20 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to give the crude material, which was purified by column chromatography over silica gel (100-200 mesh) using 0-20% ethyl acetate in pet-ether as an eluent to afford (4-fluoro-1-methyl-1H-pyrazol-5-yl)methanol (0.45 g, 69%) as a colorless liquid. TLC system: 20% EtOAc in Pet ether; R_f: 0.3.




Step 1: To a stirred solution of ethyl 1-isopropyl-1H-pyrazole-5-carboxylate (1.5 g, 8.287 mmol) in Acetic acid (15 mL) was added NaOCl (3.3 mL, 49.723 mmol) at 0° C. The RM was stirred for 6 h at RT, and the reaction progress was monitored by TLC. The RM was diluted with water (50 mL) and extracted with ethyl acetate (3×50 mL). Combined extracts were washed with water (20 mL), brine (20 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified by column chromatography over silica gel (100-200 mesh) using 15% ethyl acetate in pet-ether as an eluent to afford ethyl 4-chloro-1-isopropyl-1H-pyrazole-5-carboxylate (500 mg, 28%). TLC system: 30% ethyl acetate in pet ether; R_f: 0.68

Step 2: To a stirred solution of methyl 4-fluoro-1-methyl-1H-pyrazole-5-carboxylate (0.8 g, 5.06 mmol 1 eq) in THF (30 mL) at 0° C., was added LAH (1M) (6 mL, 6.07 mmol, 1.2 eq) solution in THF drop wise at 0° C. under an argon atmosphere. The RM was stirred at RT for 2 h. The reaction progress was monitored by TLC. The RM was quenched with sat. Na₂SO₄ solution and extracted with 10% methanol in dichloromethane (2×50 mL). The combined organic layers were washed with water (20 mL) and brine (20 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified by column chromatography over silica gel (100-200 mesh) using 0-20% ethyl acetate in pet-ether as an eluent to afford (4-fluoro-1-methyl-1H-pyrazol-5-yl)methanol (0.45 g, 69%) as a colorless liquid. TLC system: 20% EtOAc in Pet ether; R_f: 0.3.


Step 3: To a pre-stirred solution of (4-chloro-1-isopropyl-1H-pyrazol-5-yl) methanol (350 mg, 2.011 mmol) in dichloromethane (15 mL) was added triethylamine (0.5 mL, 4.02 mmol) and mesyl chloride (0.2 mL, 3.016 mmol) at 0° C. and the RM was stirred at RT for 16 h. The reaction progress was monitored by TLC. The RM was diluted with water (50 mL)

515


and extracted with dichloromethane (2×30 mL). Combined organic layers were dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford (4-chloro-1-isopropyl-1H-pyrazol-5-yl) methyl methanesulfonate (450 mg, crude) as a color less liquid. Crude was used in the next without further purification. TLC system: 20% ethyl acetate in pet ether; RF: 0.75.



Step 1: To a pre-stirred solution of (1-isopropyl-1H-pyrazol-5-yl) methanol (500 mg, 3.571 mmol) in dichloromethane (15 mL) was added triethylamine (1 mL, 7.142 mmol) and mesyl chloride (0.4 mL, 5.357 mmol) at 0° C. The RM was stirred at RT for 16 h. The reaction progress was monitored by TLC. The RM was diluted with water (50 mL) and extracted with dichloromethane (2×30 mL). Combined organic layers were dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford ((1-isopropyl-1H-pyrazol-5-yl) methyl methanesulfonate (550 mg, crude) as a color less liquid. Crude was used in next without further purification. TLC system: 50% ethyl acetate in pet ether; RF: 0.65.




Step 1: To a pre-stirred solution of (2,4-dimethyloxazol-5-yl)methanol (500 mg, 3.937 mmol) in DCM (50 mL) was added triethylamine (1.6 mL, 11.811 mmol) and Mesyl chloride (676.1 mg, 5.905 mmol) at 0° C. and the RM was stirred at RT for 3 h. The RM was diluted with water (25 mL) and extracted with dichloromethane (2×25 mL). The combined organic layer was dried over anhydrous Na_2SO_4 and evaporated under vacuum at 35° C. to afford (2,4-dimethylthiazol-5-yl) methyl methanesulfonate (560 mg, crude) as a brown liquid. TLC system: 30% Ethyl acetate in Pet ether; RF: 0.6. Proceeded to next step without purification and based on TLC.




Step 1: To a stirred solution of (1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)methanol (200 mg, 1.11 mmol) in DCM (10 mL), was added CBr_4 (551 mg, 1.66 mmol) and followed by addition of TPP (582 mL, 262 mmol) at 0° C. The RM was stirred at RT for 3 h. The reaction progress was monitored by TLC. The RM was diluted with water (30 mL) and extracted with dichloromethane (2×40 mL). Combined

516


organic layers were dried over anhydrous Na_2SO_4 and evaporated under reduced pressure to afford 5-(bromomethyl)-1-(2,2,2-trifluoroethyl)-1H-pyrazole (2355-3) (500 mg, crude) as an off-white solid. Proceeded to next step without further purification and based on TLC. TLC system: 30% ethyl acetate in pet ether; RF: 0.3.



Step 1: To a stirred solution of (1-(cyclopropylmethyl)-1H-pyrazol-5-yl)methanol (1.0 6.570 mmol) in dichloromethane (20 mL) was added Et_3N (2.7 mL, 19.71 mmol) and followed by addition of mesyl chloride (1.12 g, 9.855 mmol) at 0° C. The RM was stirred at RT for 2 h and the reaction progress was monitored by TLC. The RM was diluted with water (30 mL) and extracted with dichloromethane (2×30 mL). Combined organic layers were dried over anhydrous Na_2SO_4 and evaporated under reduced pressure to afford (1-(cyclopropylmethyl)-1H-pyrazol-5-yl)methyl methanesulfonate (1.4 g, crude) as a pale brown liquid. Crude was used in the next step without purification and based on TLC. TLC system: 40% Ethyl acetate in pet ether; RF: 0.5.




Step 1: To a stirred solution of (1-methyl-3-(trifluoromethyl)-1H-pyrazol-4-yl)methanol (400 mg, 2.222 mmol) in dichloromethane (20 mL) was added triethylamine (673 mg, 6.666 mmol) and followed by addition of methanesulfonyl chloride (379 mg, 3.333 mmol) at 0° C. The reaction mixture was stirred for 16 h at RT and the reaction progress was monitored by TLC. The reaction mixture was diluted with water (40 mL) and extracted with dichloromethane (3×50 mL). Combined extracts were washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford (1-methyl-3-(trifluoromethyl)-1H-pyrazol-4-yl)methyl methanesulfonate (3) (360 mg) as a pink solid. Crude was used in the next step without purification. TLC system: 20% Ethyl acetate in pet ether; RF: 0.54



Step 1: To a stirred solution of (3-cyclopropylpyridin-2-yl) methanol (500 mg, 3.355 mmol) in DCM (25 mL) was added triethylamine (1 mL, 6.711 mmol) and mesyl chloride

517

(0.4 mL, 5.032 mmol) at 0° C. and the resulting reaction mixture was stirred at room temperature for 16 h. The reaction progress was monitored by TLC. The reaction mixture was diluted with water (50 mL) and extracted with dichloromethane (2×50 mL). Combined organic layer were dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford (3-cyclopropylpyridin-2-yl) methyl methanesulfonate (650 mg, crude) as a brown liquid. Crude was used in the next without further purification. TLC system: 50% Ethyl acetate in pet ether; RF: 0.68.



Step 1: To a stirred solution of methyl 2-fluoronicotinate (2 g, 12.90 mmol) in DMF (50 mL) was added Cs₂CO₃ (6.2 g, 19.354 mmol) and followed by addition of cyclopropanol (1.1 g, 19.354 mmol) at RT. The reaction mixture was stirred for 16 h at 80° C., and the reaction progress was monitored by TLC. The reaction mixture was diluted with water (50

518

mL) and extracted with ethyl acetate (3×40 mL). Combined organic layers were washed with water (40 mL), brine (40 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to give crude. Crude was purified by column chromatography using silica gel (100-200 mesh size) and 20% ethyl acetate in pet ether as an eluent to afford methyl 2-cyclopropoxynicotinate (1.0 g, 40%) as a brown liquid. TLC system: 40% Ethyl acetate in Pet ether; RF: 0.62.

Step 2: To a stirred solution of methyl 2-cyclopropoxynicotinate (1.0 g, 5.181 mmol) in THF (15 mL) was added 1M LAH (5 mL, 5.181 mmol) at 0° C. The reaction mixture was stirred for 6 h at RT, and the reaction progress was monitored by TLC. The reaction mixture was quenched with Aq Na₂SO₄ and stirred for 10 min, solid was filtered and the filtrate was dried under vacuum to afford (2-cyclopropoxy-3-pyridyl) methanol (600 mg, crude) as a brown liquid. TLC system: 50% Ethyl acetate; RF: 0.20.

Step 3: To a stirred solution of (2-cyclopropoxypyridin-3-yl) methanol (500 mg, 3.030 mmol) in CCl₄ (20 mL) was added PPh₃ (1.58 g, 6.060 mmol) and followed by addition of CBr₄ (2.0 g, 6.060 mmol) at 0° C. The reaction mixture was stirred for 16 h at RT, and the reaction progress was monitored by TLC. The reaction mixture was diluted with water (50 mL), and extracted with ethyl acetate (3×40 mL). Combined organic layers were washed with water (20 mL), brine (20 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford crude. Crude was purified by column chromatography using silica gel (100-200 mesh size) and 15% ethyl acetate in pet ether as an eluent to afford 3-(bromomethyl)-2-cyclopropoxypyridine (800 mg) as a brown liquid. TLC system: 40% Ethyl acetate in Pet ether; RF: 0.71.

Analytical Data

Cpd	[M + H] ⁺ (m/z):	[M - H] ⁻ (m/z):	LC/MS Method	RT (min.)	¹ H NMR (δ ppm)
001	—	271	L1	7.6	(300 MHz, DMSO-d ₆): 12.94 (1H, s); 7.78 (1H, s); 7.68 (1H, s); 7.45-7.53 (2H, m); 6.96 (1H, dd); 6.60 (1H, br s); 4.99 (2H, s); 2.71 (3H, s)
002	274	272	L1	6.2	—
003	276	274	L5	5.1	—
004	284	282	L5	6.1	(300 MHz, DMSO-d ₆): 8.58 (d, 2H); 7.43-7.56 (4H, m); 7.02 (1H, dd); 5.22 (2H, s); 2.71 (3H, s)
005	284	282	L5	7.8	(300 MHz, DMSO-d ₆): 8.51-8.65 (1H, m); 7.96 (1H, s); 7.83 (1H, td); 7.72 (1H, d); 7.53 (1H, d); 7.19-7.39 (2H, m); 6.77-6.97 (1H, m); 5.16 (2H, s); 2.66 (3H, s)
006	284	282	L1	3.5	(300 MHz, DMSO-d ₆): 8.70 (1H, s); 8.55 (1H, d); 7.91 (1H, d); 7.33-7.61 (3H, m); 6.99-7.16 (1H, m); 5.19 (2H, s); 2.71 (3H, s)
007	285	283	L5	7.6	(300 MHz, DMSO-d ₆): 12.96 (1H, br s); 9.22 (1H, dd); 7.82-7.92 (1H, m); 7.67-7.80 (1H, m); 7.45-7.54 (2H, m); 7.05 (1H, dd); 5.27-5.63 (2H, m); 2.71 (3H, s)
008	287	285	L1	6.6	(300 MHz, DMSO-d ₆): 7.48-7.73 (2H, m); 7.37 (1H, d); 6.92-7.08 (1H, m); 6.37 (1H, d); 5.20 (2H, s); 3.85-3.98 (3H, m); 2.71 (3H, s)
009	287	285	L1	6.5	—
010	287	285	L1	6.0	—
011	—	287	L1	9.2	(300 MHz, DMSO-d ₆): 12.04 (1H, br s); 7.53-7.63 (2H, m); 7.42-7.51 (2H, m); 7.21 (1H, dd); 6.97 (1H, dd); 5.11 (2H, s); 2.71 (3H, s)
012	—	287	L1	8.9	—
013	290	288	L1	7.0	(300 MHz, DMSO-d ₆): 12.96 (1H, br s); 7.85 (1H, d); 7.78 (1H, d); 7.39-7.60 (2H, m); 7.04 (1H, dd); 5.48 (2H, s); 2.71 (3H, s)
014	290	288	L1	6.7	—
015	0	289	L1	7.3	—
016	302	300	L1	8.4	—
017	340	338	L2	4.8	—
018	—	339	L1	8.8	—

-continued

Cpd	[M + H] ⁺ (m/z):	[M - H] ⁻ (m/z):	LC/MS Method	RT (min.)	¹ H NMR (δ ppm)
019	—	374	L1	9.1	—
020	342	—	L6	6.80	—
021	352	350	L3	4.30	(DMSO d6, 300 MHz): 8.87 (2 H, d); 8.12 (2 H, d); 7.51 (1 H, d); 7.34 (1 H, br s); 7.08-7.03 (1 H, m); 5.51 (2 H, s); 4.60-4.50 (1 H, m); 3.01-3.55 (4 H, m); 2.59 (3 H, s); 2.30-2.19 (1 H, m); 1.92-2.10 (1 H, m)
022	352	350	L3	4.90	(DMSO d6-D2O, 300 MHz): 8.93 (1H, s); 8.79 (1H, d); 8.58 (1H, d); 7.01 (1H, dd); 7.53 (1H, d); 7.34 (1H, d); 7.08 (1H, dd); 5.36 (1H, s); 4.51-4.69 (1H, m); 3.22-3.62 (2H, m); 2.61 (1H, s); 2.20-2.43 (1H, m); 1.91-2.16 (1H, m)
023	355	—	L3	4.40	—
024	358	356	L6	7.20	—
025	358	—	L3	4.70	(DMSO d6-D2O, 300 MHz): 7.86 (1 H, d); 7.76 (1 H, d); 7.52 (1 H, d); 7.32 (1 H, d); 6.98-7.12 (1 H, m); 5.49 (2 H, s); 4.6-4.5 (1 H, m); 3.6-3.2 (4 H, m); 2.61 (3 H, s); 2.40-2.2 (1 H, m); 2.18-1.97 (1H, m)
026	366	364	U5	1.8	(300 MHz, DMSO-d6): 8.43-8.73 (1 H, m); 8.01-8.25 (1 H, m); 7.70-7.96 (2 H, m); 7.20-7.65 (4 H, m); 7.05-6.95 (1 H, m); 5.08-5.30 (2 H, s) 4.35-4.76 (1 H, m) 3.51-3.54 (1 H, m) 2.94-3.27 (4 H, m) 2.61 (3 H, s) 1.92-2.17 (1 H, m)
027	366	—	U5	1.6	—
028	366	364	U5	2.24	(DMSO d6, 300 MHz): 8.52 (br. s., 1H), 8.42 (br. s., 1H), 8.27 (d, 1H), 7.73 (br. s., 1H), 7.49 (d, 1H), 7.31 (br. s., 1H), 7.00 (d, , 1H), 5.18 (br. s., 2H), 4.48 (br. s., 1H), 3.06-3.30 (m, 2H), 2.87-3.03 (m, 2H), 2.61 (s., 3H), 2.34 (s, 3H), 2.00-2.21 (m, 1H), 1.74-1.96 (m, 1H).
029	366	—	U5	2.20	(CDCl3, 300 MHz): 7.61 (1 H, t); 7.29-7.42 (2 H, m); 7.22-7.43 (1 H, m); 7.27 (1 H, s); 7.16 (1 H, d); 7.03-7.13 (1 H, m); 6.95 (1 H, dd); 6.18-6.30 (1 H, m); 5.22 (2 H, s); 4.40-4.82 (1 H, m); 3.06-3.39 (2 H, m); 2.90-3.10 (2 H, m); 2.68 (3 H, s); 2.58 (3 H, s); 2.08-2.38 (1 H, m); 1.59-1.98 (1 H, m)
030	397	—	U4	2.4	—
031	366	364	U4	2.5	—
032	367	365	U5	1.3	—
033	367	365	U5	2.4	—
034	367	—	U5	2.3	—
035	371	—	U3	0.6	—
036	372	370	U2	2.01	(300 MHz, DMSO-d6): 8.15 (1H, d); 7.89 (1H, s); 7.85 (1H, d); 7.79 (1H, d); 7.50 (1H, d); 7.41 (1H, d); 7.03 (1H, dd); 5.47 (2H, s); 4.59 (1H, m); 3.25 (3H, dd); 2.62 (3H, s); 1.80-2.19 (m, 1H)
037	380	—	U5	2.05	—
038	380	—	U5	2.2	(300 MHz, CDCl3): 8.52 (1 H, s); 8.41 (1 H, s); 7.64 (1 H, s); 7.35-7.39 (1 H, m); 7.34 (1 H, s); 6.82-7.01 (1 H, m); 6.38-6.56 (1 H, m); 5.72-6.00 (1 H, m); 5.11 (2 H, s); 4.54 (1 H, d); 3.48 (2 H, dd); 2.83-3.08 (1 H, m); 2.58-2.77 (3 H, m); 2.37 (3 H, s); 1.86-2.23 (1 H, m)
039	380	378	L3	4.50	—
040	383	381	U5	2.0	(300 MHz, DMSO-d6): 8.42-8.53 (m, 2H), 8.23 (d, 1H), 7.46 (d, 1H), 7.23 (d, 1H), 7.12 (d, 1H), 6.96 (dd, 1H), 5.09 (s, 2H), 4.41-4.57 (m, 1H), 3.91 (s, 3H), 3.80-3.89 (m, 2H), 3.69-3.78 (m, 1H), 3.61 (dd, 1H), 2.57 (s, 3H), 2.08-2.23 (m, 1H), 1.85-2.02 (m, 1H).
041	385	—	L3	6.5	—
042	386	—	U5	1.8	—
043	386	384	U5	2.0	(400 MHz, DMSO-d6): 8.14 (1 H, d); 7.88 (1 H, s); 7.46-7.54 (1 H, m), 7.41 (1 H, d); 7.30 (1 H, s); 6.86-7.09 (1 H, m); 5.41 (2 H, s); 4.53-4.67 (1 H, m); 3.14-3.27 (2 H, m); 2.62 (3 H, s); 2.30-2.45 (4 H, m) 1.97-2.17 (1 H, m)
044	394	392	U5	2.1	—
045	394	—	L3	4.8	—
046	396	394	U5	2.4	(300 MHz, DMSO-d6): 8.12-8.22 (m, 2H), 7.89 (s, 1H), 7.84 (d, 1H), 7.47 (d, , 1H), 7.37 (d, 1H), 7.04 (dd, 1H), 6.97 (dd, , 1H), 5.08 (s, 2H), 4.51-4.65 (m, 1H), 3.93 (s, 3H), 3.20-3.31 (m, 2H), 2.62 (s, 3H), 2.30-2.44 (m, 1H), 1.93-2.11 (m, 1H).
047	397	395	U4	3.15	(DMSO d6, 400 MHz): 8.58 (1H, m); 7.84 (1H, td); 7.54 (1H, d); 7.48-7.42 (2H, m); 7.34 (1H, ddd); 7.26 (1H, d); 6.99 (1H, dd); 5.21 (2H, s); 4.92 (1H, t); 3.72-3.58 (6H, m); 2.60 (3H, s), 2.20 (1H, m); 2.16 (1H, m); 1.63 (2H, ddd).
048	402	—	U4	2.38	(DMSO d6, 300 MHz): 8.58 (1H, d); 8.22 (1H, d); 7.82 (1H, td); 7.54 (1H, d); 7.48 (1H, d); 7.00 (1H, dd); 5.19 (2H, s); 4.57-4.38 (1H, m); 3.11-3.07 (1H, m); 2.95-2.76 (2H, m); 2.66-2.57 (4H, m); 1.79-1.65 (2H, m).

-continued

Cpd	[M + H] ⁺ (m/z):	[M - H] ⁻ (m/z):	LC/MS Method	RT (min.)	¹ H NMR (δ ppm)
049	403	401	U5	1.95	—
050	408	—	U5	1.10	(CDCl ₃ , 300 MHz): 7.81 (1 H, d); 7.33-7.43 (2 H, m); 7.27-7.30 (1 H, m); 6.89-7.08 (1 H, m); 6.05 (1 H, d); 5.43 (2 H, s); 3.30-3.40 (1 H, m); 3.17 (1 H, d); 2.73-3.07 (2 H, m); 2.71 (3 H, s); 2.20-2.25 (1 H, m)
051	410	408	U5	2.9	(300MHz, CDCl ₃): 8.14 (1H, dd); 7.77 (1H, d); 7.34 (1H, d); 7.18 (1H, d); 6.97-6.90 (2H, m); 5.67 (1H, d); 5.10 (2H, m); 4.10-3.96 (4H, m); 2.83 (2H, d); 2.69 (3H, s); 2.32 (3H, s); 2.20 (2H, t); 2.09 (2H, m).
052	411	409	U4	3.18	—
053	416	414	U4	3.41	—
054	417	415	U5	2.24	(CDCl ₃ , 300 MHz): 8.72 (1 H, s); 7.38 (1 H, d); 7.12-7.31 (2 H, m); 6.95 (1 H, dd); 5.79 (1 H, s); 5.23 (2 H, s); 4.56 (1 H, br. s.); 3.68-3.98 (6 H, m); 2.71 (3 H, s); 2.49 (3 H, s); 1.85-2.12 (4 H, m)
055	366	364	U5	2.24	—
056	422	420	U4	2.76	(CDCl ₃ , 300 MHz): 8.72 (1 H, s); 7.37 (1 H, d); 6.83-7.00 (1 H, m); 6.02 (1 H, d); 5.23 (2 H, s); 3.28-3.46 (1 H, m); 3.13-3.20 (1 H, m); 2.75-3.04 (2 H, m); 2.71 (3 H, s); 2.50 (3 H, s); 2.15-2.25 (1 H, m); 1.61-1.79 (1 H, m) 1.25 (2 H, s)
057	427	425	U5	3.3	—
058	427	425	U5	2.56	—
059	430	428	L3	6.00	—
060	430	428	L6	7.70	—
061	430	428	L6	7.70	—
062	432	430	U5	2.98	(CDCl ₃ 300 MHz): 8.00-8.29 (1 H, m); 7.75 (1 H, d); 7.36 (1 H, d); 7.20-7.26 (1 H, m); 6.82-7.03 (2 H, m); 6.01 (1 H, d); 5.11 (2 H, s); 4.45-4.74 (1 H, m); 4.01 (3 H, s); 3.30-3.40 (1 H, m); 3.10-3.20 (1 H, m); 2.75-3.02 (2 H, m) 2.71 (3 H, s) 2.15-2.25 (1 H, m); 1.15-1.37 (1 H, m)
063	432	—	U5	3.17	(CDCl ₃ , 300 MHz): 7.58 (1 H, t); 7.35 (1 H, d); 7.27.21 (1 H, d); 7.08 (1 H, d); 6.98 (1 H, dd) 6.66 (1 H, d); 6.00 (1 H, d); 5.14 (2 H, s); 4.44-4.73 (1 H, m); 3.94 (3 H, s); 3.26-3.44 (1 H, m); 3.10-3.20 (1 H, d); 2.74-3.02 (2 H, m); 2.70 (3 H, s); 2.17 (1 H, d); 1.50-1.69 (1 H, m); 1.25 (1 H, s)
064	434	432	U5	2.5	(300 MHz, DMSO-d ₆): 8.74 (d, 1H), 8.28 (d, 1H), 8.19 (d, 1H), 7.90 (s, 1H), 7.79 (dd, , 1H), 7.50 (d, 1H), 7.38 (d, 1H), 6.99 (dd, 1H), 5.33 (s, 2H), 4.59 (q, 1H), 3.19-3.31 (m, 2H), 2.62 (s, 3H), 2.31-2.42 (m, 1H), 1.93-2.11 (m, 1H).
065	434	432	U5	2.9	—
066	434	432	U5	2.8	—
067	436	434	U5	2.87	(DMSO d ₆ , 300 MHz): 8.42 (1 H, d); 8.26 (1 H, d); 8.06 (1 H, d); 7.41-7.60 (2 H, m); 7.24 (1 H, d); 7.02 (1 H, dd); 5.19 (2 H, s); 4.34-4.69 (1 H, m); 3.04-3.20 (1 H, m); 2.69-3.01 (3 H, m); 2.55-2.68 (5 H, m); 1.59-1.91 (2 H, m)
068	436	—	U5	2.69	(CDCl ₃ , 300 MHz): 8.77 (1 H, s); 8.50 (1 H); 7.32-7.48 (2 H, m); 7.22-7.32 (1 H, m); 6.91-7.08 (1 H, m); 6.08 (1 H, d); 5.05-5.34 (2 H, m); 4.43-4.79 (1 H, m); 3.25-3.40 (1 H, m); 3.05-3.23 (1 H, m); 2.76-3.04 (2 H, m); 2.55-2.75 (3 H, m); 2.20 (1 H, d); 1.94 (1 H, br. s.); 1.38-1.76 (1 H, m); 0.62-0.96 (1 H, m)
069	437	435	L3	7.20	—
070	441	—	U5	2.33	—
071	445	—	U5	3.37	(DMSO d ₆ , 300 MHz): 8.19 (1 H, d); 7.38-7.56 (2 H, m); 7.15-7.26 (1 H, m); 6.88-7.05 (1 H, m); 6.68 (1 H, d); 6.46-6.60 (1 H, m); 4.88-5.09 (2 H, m); 4.36-4.62 (1 H, m); 3.05-3.20 (1 H, m); 3.02 (6 H, s); 2.75-2.96 (2 H, m); 2.54-2.70 (4 H, m); 1.61-1.86 (2 H, m)
072	465	463	U4	3.76	—
073	361	359	U5	3.5	—
074	470	468	U5	3.10	(CDCl ₃ , 300 MHz): 8.66 (1 H, d); 8.19 (1 H, d); 7.50-7.72 (1 H, m); 7.38 (1 H, d); 7.21 (1 H, d); 6.89-7.02 (1 H, m); 6.02 (1 H, d); 5.35 (2 H, s); 3.20-3.50 (2H, m); 2.74-3.11 (2 H, m); 2.70 (3 H, s) 2.05-2.28 (1 H, m); 1.25 (3 H, s); 0.87 (1 H, d)
075	470	468	U5	3.18	—
076	470	468	U4	3.78	—
077	520	518	U5	1.5	—

cpd	[M + H] ⁺ (m/z)	¹ H NMR (δ ppm)
103 - En 1	422.1	(DMSO-d ₆) δ = 1.85-1.96 (m, 1H), 2.04-2.13 (m, 1H), 2.41-2.45 (m, 4H), 2.57 (s, 3H), 2.62-2.69 (m, 2H), 2.89-2.99 (m, 2H), 4.30-4.41 (m, 1H), 5.30 (s, 2H), 6.97 (dd, 1H), 7.21 (d, 1H), 7.47 (d, 1H), 8.08 (d, 1H), 8.98 (s, 1H).
104	375.1	(DMSO-d ₆) δ = 1.34 (s, 6H), 2.42 (s, 3H), 2.57 (s, 3H), 3.48 (d, 2H), 5.04 (t, 1H), 5.31 (s, 2H), 6.95 (dd, 1H), 7.2 (s, 1H), 7.25 (s, 1H), 7.45 (d, 1H), 8.98 (s, 1H).
105 - En 1	456.3	(DMSO-d ₆) δ = 2.59 (s, 3H), 2.85-3.01 (m, 3H), 3.17-3.21 (m, 2H), 4.57-4.65 (m, 1H), 5.33 (s, 2H), 7.01 (dd, 1H), 7.25 (d, 1H), 7.52 (d, 1H), 7.77-7.80 (m, 1H), 8.20-8.33 (m, 2H), 8.73 (d, 1H).
106 - En 1	408.1	(DMSO-d ₆) δ = 2.32 (s, 3H), 2.59 (s, 3H), 2.83-3.06 (m, 3H), 3.21-3.27 (m, 2H), 4.57-4.66 (m, 1H), 5.32 (s, 2H), 6.98 (dd, 1H), 7.25 (d, 1H), 7.49 (d, 1H), 8.18 (br d, 1H), 8.98 (s, 1H).
107	386.8	(DMSO-d ₆): δ = 8.98 (s, 1H), 7.75 (s, 1H), 7.45 (d, 1H), 7.27 (d, 1H), 6.94 (dd, 1H), 5.31 (s, 2H), 4.90 (t, 1H), 3.66 (d, 2H), 2.58 (s, 3H), 2.42 (s, 3H), 2.38-2.29 (m, 2H), 2.15-2.09 (m, 2H), 1.90-1.81 (m, 1H), 1.80-1.70 (m, 1H).
108	414.1	(DMSO-d ₆): δ = 8.98 (s, 1H), 7.80 (s, 1H), 7.47 (d, 1H), 7.30 (d, 1H), 6.96 (dd, 1H), 5.31 (s, 2H), 2.70 (s, 2H), 2.65-2.55 (m, 5H), 2.42 (s, 3H), 2.24 (s, 6H), 2.10-2.00 (m, 2H), 1.99-1.86 (m, 1H), 1.79-1.70 (m, 1H).
109	423.1	(DMSO-d ₆) δ = 1.33 (s, 6H), 2.60 (s, 3H), 3.49-3.53 (m, 2H), 5.01-5.04 (m, 1H), 5.31 (s, 2H), 6.97 (d, 1H), 7.23 (d, 2H), 7.48 (d, 1H), 7.78 (s, 1H), 8.26 (d, 1H), 8.73 (s, 1H).
110 - En 1	436.1	(DMSO): δ = 1.86 (m, 2H), 2.15-2.21 (m, 1H), 2.25 (s, 3H), 2.34-2.44 (m, 5H), 2.569 (s, 3H), 2.77-2.80 (m, 1H), 3.04-3.11 (m, 1H), 4.31-4.41 (m, 1H), 5.30 (s, 2H), 6.95-6.98 (dd, 1H), 7.21 (s, 1H), 7.47 (d, 1H), 8.20 (d, 1H), 8.98 (s, 1H).
111 - En 1	470.2	(DMSO-d ₆) δ = 1.85-2.11 (m, 2H), 2.31-2.32 (m, 1H), 2.50-2.69 (m, 5H), 2.89-2.96 (m, 2H), 4.31-4.38 (m, 1H), 5.31 (s, 2H), 7.00 (dd, 1H), 7.22 (d, 1H), 7.50 (d, 1H), 7.77-7.80 (m, 1H), 8.10 (d, 1H), 8.25 (d, 1H), 8.73 (d, 1H).
112 - En 1	488.1	(DMSO-d ₆) δ = 8.74 (d, 1H), 8.42 (d, 1H), 8.25 (d, 1H), 7.82-7.79 (m, 1H), 7.39-7.36 (m, 1H), 7.24 (t, 1H), 5.36 (s, 2H), 4.50-4.37 (m, 1H), 3.12-3.04 (m, 1H), 2.95-2.72 (m, 2H), 2.70-2.52 (m, 1H), 2.52 (s, 3H), 2.45-2.35 (m, 1H), 1.82-1.75 (m, 1H), 1.68-1.55 (m, 1H).
113 - En 1	488.2	(DMSO-d ₆) δ = 8.75 (d, 1H), 8.30-8.24 (m, 2H), 7.82-7.79 (m, 1H), 7.67 (d, 1H), 7.37 (d, 1H), 5.37 (s, 2H), 4.52-4.42 (m, 1H), 3.14-3.07 (m, 1H), 2.94-2.72 (m, 2H), 2.60-2.56 (m, 4H), 2.49-2.39 (m, 1H), 1.82-1.68 (m, 2H).
114	401.1	(DMSO-d ₆): δ = 8.99 (s, 1H), 7.95 (s, 1H), 7.45 (d, 1H), 7.27 (d, 1H), 6.96-6.93 (m, 1H), 5.32 (s, 2H), 4.57 (t, 1H), 3.57-3.53 (m, 2H), 2.51 (s, 3H), 2.49-2.43 (m, 2H), 2.42 (s, 3H), 2.09-2.04 (m, 4H), 1.80-1.62 (m, 2H).
115 - En 1	422.1	(DMSO-d ₆) δ = 2.02-2.13 (m, 1H), 2.45-2.51 (m, 4H), 2.54 (s, 1H), 2.58-2.65 (m, 3H), 2.67-2.73 (m, 2H), 2.98 (d, 2H), 4.11-4.13 (m, 1H), 5.32 (s, 2H), 6.95-6.98 (m, 1H), 7.26 (s, 1H), 7.47 (d, 1H), 7.87 (d, 1H), 8.98 (s, 1H).
116	454.1	(DMSO) δ = 1.80-1.84 (m, 1H), 2.12-2.19 (m, 1H), 2.43 (s, 3H), 2.56 (s, 3H), 2.68-2.75 (m, 2H), 2.84-2.89 (m, 1H), 3.02-3.05 (m, 1H), 3.26-3.30 (m, 2H), 4.39-4.43 (m, 1H), 5.31 (s, 2H), 6.94-6.96 (dd, 1H), 7.23 (s, 1H), 7.45 (d, 1H), 8.12 (d, 1H), 8.98 (s, 1H).
117	435.1	(DMSO-d ₆) δ = 8.73 (d, 1H), 8.26 (d, 1H), 7.78 (t, 2H), 7.47 (d, 1H), 7.27 (d, 1H), 6.99-6.96 (m, 1H), 5.32 (s, 2H), 4.88 (t, 1H), 3.65 (d, 2H), 2.58 (s, 3H), 2.33-2.24 (m, 2H), 2.16-2.09 (m, 2H), 1.90-1.81 (m, 1H), 1.80-1.70 (m, 1H).
118 - En 1	436.1	(DMSO-d ₆) δ = 2.00-2.14 (m, 7H), 2.41 (s, 3H), 2.67-2.50 (m, 3H), 2.69-2.78 (m, 2H), 4.44-4.53 (m, 1H), 5.30 (s, 2H), 6.97 (dd, 1H), 7.25 (d, 1H), 7.47 (m, 1H), 8.15 (d, 1H), 8.98 (s, 1H).
119	402.1	(DMSO-d ₆) δ = 8.99 (s, 1H), 8.68 (s, 1H), 7.51-7.45 (m, 2H), 7.23-7.20 (m, 2H), 7.00-6.97 (m, 1H), 5.35 (s, 2H), 4.88-4.86 (m, 2H), 4.69-4.67 (m, 2H), 2.66 (s, 3H), 2.43 (s, 3H).
120 - En 1	478.2	(DMSO): δ = 1.96-2.01 (m, 1H), 2.32-2.38 (m, 1H), 2.61 (s, 3H), 3.13-3.19 (m, 1H), 3.36-3.47 (m, 3H), 3.52 (m, 2H), 4.65-4.72 (m, 2H), 5.32 (s, 2H), 6.97-7.005 (dd, 1H), 7.36 (s, 1H), 7.49 (d, 1H), 7.77-7.80 (m, 2H), 8.19-8.27 (dd, 1H), 8.735 (s, 1H).
121	417.1	(DMSO-d ₆) δ = 1.93-1.97 (m, 1H), 2.03-2.07 (m, 1H), 2.12-2.16 (m, 1H), 2.41-2.46 (s, 4H), 2.55 (s, 3H), 3.52-3.54 (m, 2H), 3.61 (d, 1H), 3.80-3.83 (m, 2H), 4.09 (d, 1H), 4.49 (s, 1H), 5.31 (s, 2H), 6.93-6.95 (dd, 1H), 7.2 (s, 1H), 7.45 (d, 1H), 7.86 (s, 1H), 8.98 (s, 1H).
122	435.1	(DMSO-d ₆): δ = 8.99 (s, 1H), 7.60 (d, 1H), 7.43-7.41 (m, 2H), 5.37 (s, 2H), 4.89 (br s, 1H), 3.73-3.56 (m, 6H), 2.59 (s, 3H), 2.41 (s, 3H), 2.16 (d, 2H), 1.65-1.59 (m, 2H).
123 - En 1	422.1	(DMSO-d ₆) δ = 2.29 (s, 3H), 2.42 (s, 3H), 2.51-2.58 (m, 1H), 2.61 (s, 3H), 2.68-2.71 (m, 1H), 3.17-3.25 (m, 2H), 4.68-4.76 (m, 1H), 5.31 (s, 2H), 6.98 (dd, 1H), 7.23 (d, 1H), 7.48 (d, 1H), 8.27 (br d, 1H), 8.98 (s, 1H).
124 - En 1	470.1	(DMSO-d ₆) δ = 2.29 (s, 3H), 2.45-2.49 (m, 1H), 2.56-2.62 (m, 3H), 2.66-2.69 (m, 1H), 3.17-3.25 (m, 2H), 4.67-4.75 (m, 1H), 5.33 (s, 2H), 6.99-7.02 (m, 1H), 7.24 (d, 1H), 7.51 (d, 1H), 7.77-7.80 (m, 1H), 8.24-8.30 (m, 2H), 8.73 (d, 1H).
125 - En 1	470.1	(DMSO-d ₆) δ = 2.04-2.11 (m, 1H), 2.32-2.35 (m, 1H), 2.50 (br s, 1H), 2.58 (s, 3H), 2.67-2.82 (m, 2H), 2.98 (d, 2H), 4.06-4.12 (m, 1H), 5.33 (s, 2H), 7.00 (dd, 1H), 7.26 (d, 1H), 7.50 (d, 1H), 7.77-7.80 (m, 1H), 7.90 (d, 1H), 8.26 (d, 1H), 8.73 (d, 1H).
126 - En 1	465.1	(DMSO-d ₆) δ = 1.93 (td, 1H), 2.03 (td, 1H), 2.12-2.18 (m, 1H), 2.39-2.46 (m, 1H), 2.56 (s, 3H), 3.47-3.56 (m, 2H), 3.60 (d, 1H), 3.80 (dd, 2H), 4.08 (d, 1H), 4.47 (d, 1H), 5.31 (s, 2H), 6.99 (dd, 1H), 7.21 (d, 1H), 7.49 (d, 1H), 7.79 (dd, 1H), 7.88 (s, 1H), 8.26 (d, 1H), 8.72 (d, 1H).
127	449.1	(DMSO-d ₆): δ = 8.37 (d, 1H), 8.26 (d, 1H), 7.97 (s, 1H), 7.80-7.77 (m, 1H), 7.48 (d, 1H), 7.27 (d, 1H), 6.99-6.97 (m, 1H), 5.33 (s, 2H), 4.53 (t, 1H), 3.56-3.51 (m, 2H), 2.58 (s, 3H), 2.46-2.40 (m, 2H), 2.10-2.02 (m, 4H), 1.99-1.72 (m, 2H).
128	403.1	(DMSO-d ₆) δ = 8.98 (s, 1H), 8.37 (s, 1H), 7.47 (m, 1H), 7.31 (d, 1H), 6.97 (dd, 1H), 5.32 (s, 2H), 4.75-4.74 (m, 2H), 4.53-4.51 (m, 3H), 3.59-3.56 (m, 2H), 2.60 (s, 3H), 2.42 (s, 3H), 2.25 (t, 2H).
129	451.1	(DMSO-d ₆) δ = 8.74 (d, 1H), 8.39 (s, 1H), 8.26 (d, 1H), 7.80-7.77 (m, 1H), 7.51-7.49 (m, 1H), 7.31 (d, 1H), 7.00 (dd, 1H), 5.32 (s, 2H), 4.73 (d, 2H), 4.53-4.48 (m, 3H), 3.59-3.54 (m, 2H), 2.61 (s, 3H), 2.25 (t, 2H).

-continued

cpd	[M + H] ⁺ (m/z)	¹ H NMR (δ ppm)
130 - En 1	484.1	(DMSO): δ = 1.8-1.813 (m, 2H), 2.14-2.17 (m, 1H), 2.21 (s, 3H), 2.32-2.41 (m, 1H), 2.54 (s, 3H), 2.77-2.79 (m, 1H), 3.03-3.09 (m, 1H), 4.33-4.40 (m, 1H), 5.30 (s, 2H), 6.98-7.01 (dd, 1H), 7.221 (s, 1H), 7.50 (d, 1H), 7.76-7.80 (m, 1H), 8.21-8.21-8.26 (m, 2H), 8.73 (d, 1H).
131 - En 1	484.2	(DMSO-d ₆) δ = 2.07-2.33 (m, 7H), 2.57 (s, 3H), 2.61-2.77 (m, 2H), 4.43-4.53 (m, 1H), 5.31 (s, 2H), 7.00 (dd, 1H), 7.22 (d, 1H), 7.51 (d, 1H), 7.77-7.80 (m, 1H), 8.17 (d, 1H), 8.26 (d, 1H), 8.73 (d, 1H).
132 - En 1	403.1	(DMSO-d ₆) δ = 2.04-2.07 (m, 1H), 2.23-2.25 (m, 1H), 2.42 (s, 3H), 2.57 (s, 3H), 3.66-3.71 (m, 2H), 3.78-3.82 (m, 3H), 3.92 (d, 1H), 5.07 (s, 1H), 5.30 (s, 2H), 6.93-6.95 (dd, 1H), 7.25 (s, 1H), 7.47 (d, 1H), 7.84 (s, 1H), 8.98 (s, 1H).
133	475.1	(DMSO-d ₆) δ = 8.73 (d, 1H), 8.37 (s, 1H), 8.26 (d, 1H), 7.80-7.77 (m, 1H), 7.49 (d, 1H), 7.34-7.33 (m, 1H), 7.00-6.98 (m, 1H), 5.32 (s, 2H), 5.18 (t, 1H), 4.66 (d, 2H), 4.55 (d, 2H), 3.76 (d, 2H), 2.61 (s, 3H).
134 - En 1	500.2	(DMSO-d ₆): δ = 2.49-2.58 (m, 6H), 2.69-2.80 (m, 1H), 3.24-3.31 (m, 2H), 3.51 (s, 2H), 4.54 (s, 1H), 4.65-4.73 (m, 1H), 5.32 (s, 2H), 7.01 (dd, 1H), 7.24 (d, 1H), 7.51 (d, 1H), 7.77-7.80 (m, 1H), 8.26 (t, 2H), 8.73 (d, 1H).
135 - En1	451.0	(DMSO-d ₆) δ = 2.06 (td, 1H), 2.22 (td, 1H), 2.58 (s, 3H), 3.63-3.73 (m, 2H), 3.75-3.82 (m, 3H), 3.92 (d, 1H), 5.05 (t, 1H), 5.32 (s, 2H), 6.98 (dd, 1H), 7.25 (d, 1H), 7.48 (d, 1H), 7.78 (dd, 1H), 7.86 (s, 1H), 8.25 (d, 1H), 8.72 (d, 1H).
136	434.1	(DMSO-d ₆) δ = 8.73 (d, 1H), 8.26 (d, 1H), 7.82-7.76 (m, 1H), 7.75 (s, 1H), 7.48 (d, 1H), 7.30 (d, 1H), 6.98 (dd, 1H), 5.32 (s, 2H), 2.87 (s, 2H), 2.60 (s, 3H), 2.42-2.35 (m, 2H), 2.09-2.00 (m, 2H), 1.90-1.80 (m, 1H), 1.79-1.52 (m, 3H).
137	468.1	(DMSO-d ₆) δ = 8.98 (s, 1H), 7.79 (s, 1H), 7.45 (d, 1H), 7.26 (d, 1H), 6.94 (dd, 1H), 5.31 (s, 2H), 3.29-3.26 (m, 2H), 3.05 (d, 2H), 2.58 (s, 3H), 2.41 (s, 3H), 2.40-2.32 (m, 3H), 2.15-2.08 (m, 2H), 1.90-1.81 (m, 1H), 1.80-1.73 (m, 1H).
138	383.1	(DMSO-d ₆) δ = 2.41 (s, 3H), 2.69 (s, 3H), 3.75 (s, 3H), 5.34 (s, 2H), 6.31 (d, 1H), 6.98 (dd, 1H), 7.39-7.40 (m, 2H), 7.53 (d, 1H), 8.98 (s, 1H), 10.0 (s, 1H).
139	373.1	(DMSO-d ₆) δ = 0.75-0.78 (m, 4H), 2.42 (s, 3H), 2.56 (s, 3H), 3.55 (d, 2H), 4.77 (t, 1H), 5.32 (s, 2H), 6.93 (dd, 1H), 7.26 (s, 1H), 7.44 (d, 1H), 8.14 (s, 1H), 8.98 (s, 1H).
140	462.2	(DMSO-d ₆) δ = 8.73 (d, 1H), 8.26 (d, 1H), 7.83 (s, 1H), 7.82-7.78 (m, 1H), 7.50 (d, 1H), 7.28 (d, 1H), 6.99 (dd, 1H), 5.31 (s, 2H), 2.68 (s, 2H), 2.59 (s, 3H), 2.58-2.51 (m, 2H), 2.21 (s, 6H), 2.10-2.01 (m, 2H), 1.95-1.85 (m, 1H), 1.80-1.70 (m, 1H).
141 - En 1	484.2	(DMSO-d ₆) δ = 1.95-2.10 (m, 1H), 2.12-2.15 (m, 1H), 2.29-2.49 (m, 5H), 2.67 (s, 3H), 2.82-2.93 (m, 2H), 4.19-4.21 (m, 1H), 5.33 (s, 2H), 7.01 (dd, 1H), 7.25 (d, 1H), 7.51 (d, 1H), 7.77-7.80 (m, 1H), 7.84 (br d, 1H), 8.26 (d, 1H), 8.73 (d, 1H).
142	388.8	(DMSO-d ₆) δ = 8.98 (s, 1H), 8.33 (s, 1H), 7.47 (d, 1H), 7.33 (d, 1H), 6.96 (dd, 1H), 5.32 (s, 2H), 5.20 (t, 1H), 4.67 (d, 2H), 4.54 (d, 2H), 3.77 (d, 2H), 2.60 (s, 3H), 2.43 (s, 3H).
143	431.1	(DMSO) δ = 2.63 (s, 3H), 3.82 (s, 3H), 5.34 (s, 2H), 7.02 (br dd, 1H), 7.31 (d, 1H), 7.51-7.54 (m, 2H), 7.77-7.80 (m, 1H), 8.01 (s, 1H), 8.27 (d, 1H), 8.73 (d, 1H), 10.07 (s, 1H).
144	369.1	(DMSO-d ₆) δ = 2.41 (s, 3H), 2.62 (s, 3H), 5.32 (s, 2H), 6.63 (s, 1H), 6.94-6.97 (dd, 1H), 7.3 (s, 1H), 7.47 (d, 1H), 7.67 (s, 1H), 8.97 (s, 1H), 10.45 (s, 1H), 12.43 (s, 1H).
145	468.1	(DMSO-d ₆) δ = 8.98 (s, 1H), 7.79 (s, 1H), 7.45 (d, 1H), 7.26 (d, 1H), 6.94 (dd, 1H), 5.31 (s, 2H), 3.29-3.26 (m, 2H), 3.05 (d, 2H), 2.58 (s, 3H), 2.41 (s, 3H), 2.40-2.32 (m, 3H), 2.15-2.08 (m, 2H), 1.90-1.81 (m, 1H), 1.80-1.73 (m, 1H).
146	431.1	(DMSO) δ = 2.63 (s, 3H), 3.78 (s, 3H), 5.32 (s, 2H), 6.57 (s, 1H), 7.00 (dd, 1H), 7.30 (s, 1H), 7.51 (d, 1H), 7.61 (d, 1H), 7.77-7.79 (m, 1H), 8.27 (d, 1H), 8.73 (d, 1H), 10.46 (s, 1H).
147 - Dia 1 - En 1	437.1	(DMSO-d ₆) δ = 2.56 (s, 3H), 3.56 (dd, 1H), 3.65 (dd, 1H), 3.88-3.92 (m, 1H), 3.99-4.02 (m, 1H), 4.23 (s, 2H), 5.30 (t, 3H), 6.99 (dd, 1H), 7.23 (d, 1H), 7.50 (d, 1H), 7.77-7.80 (m, 1H), 8.17 (dd, 1H), 8.25 (d, 1H), 8.73 (t, 1H).
148	421.1	(DMSO) δ = 1.89-1.92 (m, 1H), 2.14-2.18 (m, 1H), 2.57 (s, 3H), 3.58-3.61 (m, 1H), 3.71-3.75 (m, 1H), 3.81-3.89 (m, 2H), 4.42-4.47 (m, 1H), 5.32 (s, 2H), 7.00 (dd, 1H), 7.23 (s, 1H), 7.50 (d, 1H), 7.77-7.80 (m, 1H), 8.20 (d, 2H), 8.26 (d, 2H), 8.73 (d, 1H).
149	502.2	(DMSO-d ₆): δ = 1.78-1.82 (m, 1H), 2.12-2.16 (m, 1H), 2.57 (s, 3H), 2.66-2.77 (m, 2H), 2.82-2.89 (m, 1H), 3.02-3.07 (m, 1H), 3.25-3.32 (m, 2H), 4.37-4.45 (m, 1H), 5.31 (s, 2H), 6.98-7.00 (m, 1H), 7.22 (d, 1H), 7.49 (d, 1H), 7.79 (dd, 1H), 8.15 (d, 1H), 8.26 (d, 1H), 8.73 (d, 1H).
150	369.1	(DMSO-d ₆) δ = 2.41 (s, 3H), 2.63 (s, 3H), 5.34 (s, 2H), 6.97-6.99 (m, 1H), 7.32 (s, 1H), 7.50 (d, 1H), 7.67 (br s, 1H), 7.98 (br s, 1H), 8.98 (s, 1H), 10.0 (s, 1H), 12.62 (br s, 1H).
151 - En 1	452.2	(DMSO-d ₆) δ = 2.41 (s, 3H), 2.53-2.61 (m, 6H), 2.73-2.78 (m, 1H), 3.25-3.28 (m, 2H), 3.50-3.51 (m, 2H), 4.54 (m, 1H), 4.65-4.75 (m, 1H), 5.31 (s, 2H), 6.96-6.98 (dd, 1H), 7.23 (s, 1H), 7.48 (d, 1H), 8.25 (d, 1H), 8.98 (s, 1H).
152	383.1	(DMSO-d ₆) δ = 2.41 (s, 3H), 2.62 (s, 3H), 3.82 (s, 3H), 5.33 (s, 2H), 6.98 (dd, 1H), 7.30 (s, 1H), 7.49 (d, 1H), 7.56 (s, 1H), 8.01 (s, 1H), 8.98 (s, 1H), 10.04 (s, 1H).
153	516.2	(DMSO-d ₆) δ = 8.73 (d, 1H), 8.25 (d, 1H), 7.82-7.77 (m, 2H), 7.48 (d, 1H), 7.26 (d, 1H), 6.99 (dd, 1H), 5.32 (s, 2H), 3.29-3.25 (m, 2H), 3.04 (d, 2H), 2.58 (s, 3H), 2.38-2.28 (m, 3H), 2.13-2.08 (m, 2H), 1.91-1.72 (m, 2H).
154	431.1	(DMSO-d ₆): δ = 2.70 (s, 3H), 3.74 (s, 3H), 5.35 (s, 2H), 6.32 (s, 1H), 7.04 (dd, 1H), 7.37-7.41 (m, 2H), 7.56 (d, 1H), 7.76-7.79 (m, 1H), 8.27 (d, 1H), 8.74 (d, 1H), 10.04 (s, 1H).
155 - En 1	490.1	(DMSO-d ₆) δ = 2.42 (s, 3H), 2.58 (s, 3H), 2.88-2.92 (m, 1H), 3.04-3.07 (m, 1H), 3.38-3.44 (m, 4H), 4.74-4.82 (m, 1H), 5.31 (s, 2H), 6.96 (dd, 1H), 7.24 (d, 1H), 7.48 (d, 1H), 8.31 (d, 1H), 8.98 (s, 1H).
156	398.8	(DMSO-d ₆) δ = 8.99 (s, 1H), 8.14 (d, 1H), 7.46 (d, 1H), 7.24 (d, 1H), 6.96 (dd, 1H), 5.32 (s, 2H), 4.64 (s, 2H), 4.52 (s, 2H), 4.28-4.21 (m, 1H), 2.61-2.55 (m, 5H), 2.42 (s, 3H), 2.30-2.23 (m, 2H).

-continued

cpd	[M + H] ⁺ (m/z)	¹ H NMR (δ ppm)
157 - En 1	538.1	(DMSO-d ₆): δ = 2.60 (s, 3H), 2.85-2.90 (m, 1H), 3.02- 3.13 (m, 1H), 3.38-3.39 (m, 4H), 4.73-4.81 (m, 1H), 5.32 (s, 2H), 7.01 (dd, 1H), 7.24 (d, 1H), 7.52 (d, 1H), 7.77-7.80 (m, 1H), 8.26 (d, 1H), 8.35 (d, 1H), 8.73 (d, 1H).
158	417.1	(DMSO-d ₆): δ = 2.64 (s, 3H), 5.34 (s, 2H), 7.02 (dd, 1H), 7.33 (s, 1H), 7.53 (d, 1H), 7.65 (br s, 1H), 7.76-7.79 (m, 1H), 7.99 (br s, 1H), 8.27 (d, 1H), 8.73 (d, 1H), 10.07 (s, 1H), 12.64 (br s, 1H).
159	373.1	(DMSO-d ₆): δ = 1.89-1.94 (m, 1H), 2.15-2.19 (m, 1H), 2.49 (m, 3H), 2.56 (s, 3H), 3.59-3.62 (m, 1H), 3.71-3.75 (m, 1H), 3.83-3.89 (m, 2H), 4.47-4.48 (m, 1H), 5.32 (s, 2H), 6.94-6.96 (dd, 1H), 7.22-7.23 (d, 1H), 7.45-7.46 (d, 1H), 8.18-8.19 (d, 1H), 8.98 (s, 1H).
160	430.1	(DMSO-d ₆): δ = 1.94-2.00 (m, 1H), 2.32-2.37 (m, 1H), 2.42 (s, 3H), 2.61 (s, 3H), 3.14-3.19 (m, 1H), 3.31-3.45 (m, 3H), 3.50-3.54 (m, 2H), 4.65-4.73 (m, 2H), 5.32 (s, 2H), 6.95-6.98 (dd, 1H), 7.36 (s, 1H), 7.47 (d, 1H), 8.17 (d, 1H), 8.98 (s, 1H).
161	383.1	(DMSO-d ₆): δ = 2.41 (s, 3H), 2.61 (s, 3H), 3.76 (s, 3H), 5.32 (s, 2H), 6.58 (s, 1H), 6.96 (dd, 1H), 7.28 (s, 1H), 7.47 (d, 1H), 7.61 (s, 1H), 8.98 (s, 1H), 10.45 (s, 1H).
162 - Dial - En 1	389.1	(DMSO-d ₆): δ = 2.42 (s, 3H), 2.56 (s, 3H), 3.57 (dd, 1H), 3.67 (dd, 1H), 3.89-3.93 (m, 1H), 3.99-4.03 (m, 1H), 4.22-4.24 (m, 2H), 5.31 (s, 3H), 6.96 (dd, 1H), 7.22 (d, 1H), 7.47 (d, 1H), 8.14 (d, 1H), 8.98 (s, 1H).
163 - En 1	436.2	(DMSO-d ₆): δ = 1.9-2.17 (m, 1H), 2.30-2.33 (m, 1H), 2.34-2.37 (m, 3H), 2.39-2.42 (m, 2H), 2.49-2.51 (m, 3H), 2.58 (s, 3H), 2.84- 2.91 (m, 2H), 4.68-4.76 (m, 1H), 5.32 (s, 2H), 6.95-6.98 (m, 1H), 7.26 (s, 1H), 7.47 (d, 1H), 7.87 (d, 1H), 8.99 (s, 1H).
164 - En 1	424.2	(DMSO-d ₆): δ = 8.41 (d, 1H), 8.19 (d, 1H), 7.81 (d, 1H), 7.48 (d, 1H), 7.40 (d, 1H), 7.24 (t, 1H), 7.01 (dd, 1H), 5.16 (s, 2H), 4.73-4.65 (m, 2H), 3.52 (s, 2H), 3.48-3.37 (m, 3H), 3.21-3.14 (m, 1H), 2.63 (s, 3H), 2.53 (s, 3H), 2.40-2.30 (m, 1H), 2.01-1.90 (m, 1H).
165 - En 1	504.1	(DMSO-d ₆): δ = 8.75 (d, 1H), 8.31 (d, 2H), 7.84-7.81 (m, 2H), 7.37 (s, 1H), 5.39 (s, 2H), 4.49-4.42 (m, 1H), 3.14-3.07 (m, 1H), 2.93-2.72 (m, 2H), 2.60-2.56 (m, 4H), 2.49-2.39 (m, 1H), 1.82-1.68 (m, 2H).
166	468.1	(DMSO-d ₆): δ = 1.58-1.65 (m, 2H), 1.81-1.83 (m, 2H), 2.42-2.49 (m, 5H), 2.56 (s, 3H), 2.92-2.94 (m, 2H), 3.13-3.20 (m, 2H), 3.78-3.80 (m, 1H), 5.32 (s, 2H), 6.96 (dd, 1H), 7.23 (brs, 1H), 7.45 (d, 1H), 7.86 (brd, 1H), 8.98 (s, 1H).
167	447.1	(DMSO-d ₆): δ = 8.73 (d, 1H), 8.25 (d, 1H), 8.17 (d, 1H), 7.80-7.77 (m, 1H), 7.49 (d, 1H), 7.23-7.23 (d, 1H), 7.00-6.98 (m, 1H), 5.34 (s, 2H), 4.63 (s, 2H), 4.51 (s, 2H), 4.26-4.21 (m, 1H), 2.60-2.54 (m, 5H), 2.26-2.22 (m, 2H).
168	516.1	(DMSO-d ₆): δ = 1.55-1.60 (m, 2H), 1.79-1.84 (m, 2H), 2.41-2.48 (m, 2H), 2.57 (s, 3H), 2.92 (br d, 2H), 3.10-3.16 (m, 2H), 3.74-3.81 (m, 1H), 5.33 (s, 2H), 6.99 (dd, 1H), 7.23 (d, 1H), 7.49 (d, 1H), 7.76-7.80 (m, 1H), 7.89 (d, 1H), 8.26 (d, 1H), 8.73 (d, 1H).
169 - En 1	444.2	(DMSO-d ₆): δ = 8.99 (s, 1H), 8.36 (d, 1H), 7.47 (d, 1H), 7.27 (d, 1H), 6.96 (dd, 1H), 5.32 (s, 2H), 4.72 (t, 1H), 4.62-4.52 (m, 1H), 3.85-3.77 (m, 1H), 3.53-3.48 (m, 2H), 3.42-3.33 (m, 1H), 3.28-3.23 (m, 2H), 2.70-2.62 (m, 1H), 2.58 (s, 3H), 2.43-2.35 (m, 4H).
170	451.1	(DMSO-d ₆): δ = 2.08 (td, 1H), 2.26 (br s, 1H), 2.76 (s, 3H), 3.80-3.91 (m, 3H), 3.95-4.02 (m, 1H), 4.57 (br d, 2H), 5.35 (br s, 2H), 7.08 (br dd, 1H), 7.46 (br d, 1H), 7.60 (br d, 1H), 7.80 (br dd, 1H), 8.20-8.31 (m, 3H), 8.75 (br d, 1H).
171	383.1	(DMSO-d ₆): δ = 2.41 (s, 3H), 2.69 (s, 3H), 3.75 (s, 3H), 5.34 (s, 2H), 6.31 (d, 1H), 6.98 (dd, 1H), 7.39-7.40 (m, 2H), 7.53 (d, 1H), 8.98 (s, 1H), 10.0 (s, 1H).
172 - En 1	534.1	(DMSO-d ₆): δ = 2.41 (s, 3H), 2.59 (s, 3H), 3.50-3.58 (m, 1H), 3.86-3.96 (m, 3H), 4.71-4.77 (m, 2H), 5.08-5.12 (m, 1H), 5.31 (s, 2H), 6.96 (dd, 1H), 7.24 (d, 1H), 7.49 (d, 1H), 8.51 (t, 1H), 8.98 (s, 1H).
173	446.2	(CDCl ₃): δ = 8.66 (d, 1H), 8.17 (d, 1H), 7.56-7.53 (m, 1H), 7.36 (d, 1H), 7.13 (s, 1H), 6.94 (d, 1H), 5.78 (d, 1H), 5.36 (s, 2H), 4.48-4.42 (m, 1H), 3.72 (s, 2H), 3.61 (s, 2H), 2.74-2.68 (m, 6H), 2.04 (t, 2H).
174	451.1	(DMSO-d ₆): δ = 8.99 (s, 1H), 7.76 (s, 1H), 7.47-7.43 (m, 2H), 5.39 (s, 2H), 4.91 (t, 1H), 3.73-3.56 (m, 6H), 2.61 (s, 3H), 2.42 (s, 3H), 2.20-2.15 (m, 2H), 1.68-1.59 (m, 2H).
175	582.1	(DMSO-d ₆): δ = 2.59 (s, 3H), 3.50-3.56 (m, 1H), 3.82- 3.96 (m, 3H), 4.70-4.77 (m, 2H), 5.05-5.15 (m, 1H), 5.32 (s, 2H), 7.02 (dd, 1H), 7.24 (d, 1H), 7.53 (d, 1H), 7.77-7.80 (m, 1H), 8.25 (d, 1H), 8.54 (t, 1H), 8.73 (d, 1H).
176 - En 1	488.1	(DMSO-d ₆): δ = 8.74 (d, 1H), 8.35 (d, 1H), 8.26 (d, 1H), 7.80-7.77 (m, 1H), 7.06-7.03 (m, 2H), 5.32 (s, 2H), 4.51-4.40 (m, 1H), 3.13-3.07 (m, 1H), 2.93-2.78 (m, 2H), 2.65-2.59 (m, 4H), 2.49-2.44 (m, 1H), 1.80-1.65 (m, 2H).
177	435.2	(DMSO-d ₆): δ 8.98 (s, 1H), 7.72 (s, 1H), 7.34 (d, 1H), 7.23 (t, 1H), 5.37 (s, 2H), 4.81 (t, 1H), 3.71-3.55 (m, 6H), 2.52 (s, 3H), 2.39 (s, 3H), 2.10 (d, 2H), 1.66-1.58 (m, 2H).
178	417.1	(DMSO-d ₆): δ = 2.54 (s, 3H), 5.29 (s, 2H), 5.65 (s, 2H), 6.02 (d, 1H), 6.99-7.01 (m, 1H), 7.16 (d, 1H), 7.53 (d, 1H), 7.75-7.78 (m, 1H), 8.16 (d, 1H), 8.23 (d, 1H), 8.71-8.72 (m, 1H).
179 - En 1	484.2	(DMSO-d ₆): δ = 8.74 (d, 1H), 8.29 (d, 1H), 8.20 (d, 1H), 7.82-7.79 (m, 1H), 7.41 (s, 1H), 7.20 (s, 1H), 5.30 (s, 2H), 4.50-4.43 (m, 1H), 3.18-3.05 (m, 1H), 2.95-2.72 (m, 2H), 2.63-2.40 (m, 5H), 2.28 (s, 3H), 1.71-1.64 (m, 2H).
180 - En 1	430.1	(DMSO-d ₆): δ = 8.99 (s, 1H), 8.36 (d, 1H), 7.47 (d, 1H), 7.27 (d, 1H), 6.96 (dd, 1H), 5.32 (s, 2H), 4.72 (t, 1H), 4.62-4.52 (m, 1H), 3.85-3.77 (m, 1H), 3.53-3.48 (m, 2H), 3.42-3.33 (m, 1H), 3.28-3.23 (m, 2H), 2.70-2.62 (m, 1H), 2.58 (s, 3H), 2.43-2.35 (m, 4H).
181	417.1	(DMSO-d ₆): δ = 2.52 (s, 3H), 4.72 (s, 2H), 5.27 (s, 2H), 7.00-7.03 (m, 1H), 7.10 (d, 1H), 7.47 (s, 1H), 7.54 (d, 1H), 7.60 (s, 1H), 7.76-7.78 (m, 1H), 8.22 (d, 1H), 8.72 (d, 1H).
182	435.1	(DMSO-d ₆): δ = 8.99 (s, 1H), 7.51 (s, 1H), 7.07 (d, 1H), 7.00-6.97 (m, 1H), 5.33 (s, 2H), 4.89 (t, 1H), 3.73-3.57 (m, 6H), 2.62 (s, 3H), 2.42 (s, 3H), 2.16 (d, 2H), 1.65-1.59 (m, 2H).
183 - En 1	484.2	(DMSO-d ₆): δ = 8.73 (d, 1H), 8.24 (d, 1H), 8.18 (d, 1H), 7.79-7.76 (m, 1H), 7.04 (d, 1H), 6.86 (d, 1H), 5.28 (s, 2H), 4.49-4.40 (m, 1H), 3.13-3.06 (m, 1H), 2.92-2.77 (m, 2H), 2.63-2.56 (m, 4H), 2.49-2.29 (m, 4H), 1.79-1.70 (m, 2H).

-continued

cpd	[M + H] ⁺ (m/z)	¹ H NMR (δ ppm)
184 - En 1	504.1	(DMSO-d ₆) δ = 8.74 (d, 1H), 8.35 (d, 1H), 8.26 (d, 1H), 7.80-7.77 (m, 1H), 7.19 (dd, 2H), 5.33 (s, 2H), 4.51-4.40 (m, 1H), 3.14-3.07 (m, 1H), 2.93-2.27 (m, 2H), 2.65-2.59 (m, 4H), 2.50-2.44 (m, 1H), 1.80-1.67 (m, 2H).
185	431.1	(DMSO-d ₆) δ = 8.97 (s, 1H), 7.36 (s, 1H), 7.32 (s, 1H), 7.28 (s, 1H), 5.33 (s, 2H), 4.88 (t, 1H), 3.73-3.59 (m, 6H), 2.59 (s, 3H), 2.42 (s, 3H), 2.23 (s, 3H), 2.22-2.16 (m, 2H), 1.68-1.60 (m, 2H).
186	431.1	(DMSO-d ₆) δ = 8.97 (s, 1H), 7.32 (s, 1H), 7.08 (d, 1H), 6.81 (d, 1H), 5.28 (s, 2H), 4.88 (t, 1H), 3.72-3.58 (m, 6H), 2.60 (s, 3H), 2.41 (s, 6H), 2.18-2.16 (m, 2H), 1.66-1.60 (m, 2H).
187	451.1	(DMSO-d ₆) δ = 8.99 (s, 1H), 7.76 (s, 1H), 7.47-7.43 (m, 2H), 5.39 (s, 2H), 4.91 (t, 1H), 3.73-3.56 (m, 6H), 2.61 (s, 3H), 2.42 (s, 3H), 2.20-2.15 (m, 2H), 1.68-1.59 (m, 2H).
188	442.2	(DMSO-d ₆) δ = 9.01 (s, 1H), 7.63 (s, 1H), 7.58 (d, 1H), 7.56 (d, 1H), 5.39 (s, 2H), 4.91 (t, 1H), 3.73-3.56 (m, 6H), 2.67 (s, 3H), 2.43 (s, 3H), 2.17 (d, 2H), 1.65-1.58 (m, 2H).
189	436.2	(DMSO-d ₆) δ = 8.73 (d, 1H), 8.26 (d, 1H), 7.80-7.76 (m, 1H), 7.56 (d, 1H), 7.45 (d, 1H), 6.95 (dd, 1H), 6.52-6.13 (br s, 1H), 5.31 (s, 2H), 4.57 (br s, 2H), 3.49-3.38 (m, 6H), 2.64 (s, 3H).
190	398.2	(CDCl ₃) δ = 8.72 (d, 1H), 7.35 (d, 1H), 7.20 (d, 1H), 6.92 (dd, 1H), 5.81 (d, 1H), 5.23 (s, 2H), 4.50-4.43 (m, 1H), 3.73 (s, 2H), 3.61 (s, 2H), 2.78-2.71 (m, 2H), 2.68 (s, 3H), 2.50 (s, 3H), 2.11-2.05 (m, 2H).
191 - En 1	495.1	(DMSO-d ₆) δ = 8.78-8.74 (m, 2H), 8.30 (d, 1H), 7.92 (m, 1H), 7.86-7.83 (m, 1H), 7.30 (d, 1H), 5.49 (s, 2H), 4.52-4.40 (m, 1H), 3.18-3.07 (m, 1H), 2.93-2.77 (m, 2H), 2.63-2.51 (m, 2H), 2.52 (s, 3H), 1.90-1.84 (m, 1H), 1.70-1.61 (m, 1H).
192	369.1	(DMSO-d ₆) δ = 2.38 (s, 3H), 2.54 (s, 3H), 5.28 (s, 2H), 5.67 (s, 2H), 6.032 (s, 1H), 6.96 (dd, 1H), 7.16 (s, 1H), 7.5 (d, 1H), 8.17 (s, 1H), 8.97 (s, 1H).
193	442.1	(DMSO-d ₆) δ = 9.00 (s, 1H), 7.93 (s, 1H), 7.88 (d, 1H), 7.33 (d, 1H), 5.52 (s, 2H), 4.77 (t, 1H), 3.70-3.55 (m, 6H), 2.56 (s, 3H), 2.44 (s, 3H), 2.11-2.07 (m, 2H), 1.69-1.65 (m, 2H).
194	442.2	(DMSO-d ₆) δ = 9.01 (s, 1H), 8.09 (s, 1H), 7.56 (s, 1H), 7.48 (s, 1H), 5.49 (s, 2H), 4.91 (t, 1H), 3.73-3.56 (m, 6H), 2.65 (s, 3H), 2.45 (s, 3H), 2.19-2.15 (m, 2H), 1.67-1.58 (m, 2H).
195 - En 1	495.1	(DMSO-d ₆) δ = 8.77-8.75 (m, 1H), 8.43 (d, 1H), 8.30 (d, 1H), 8.17 (s, 1H), 7.86-7.83 (m, 1H), 7.39 (s, 1H), 5.46 (s, 2H), 4.52-4.41 (m, 1H), 3.15-3.07 (m, 1H), 2.93-2.80 (m, 2H), 2.63-2.48 (m, 5H), 1.85-1.75 (m, 1H), 1.73-1.65 (m, 1H).
196 - En 1	495.1	(DMSO-d ₆) δ = 8.74 (m, 1H), 8.44 (d, 1H), 8.27 (d, 1H), 7.82-7.77 (m, 1H), 7.65 (d, 1H), 7.53 (d, 1H), 5.37 (s, 2H), 4.53-4.40 (m, 1H), 3.15-3.06 (m, 1H), 2.93-2.75 (m, 2H), 2.65-2.55 (m, 4H), 2.49-1.82-1.78 (m, 1H), 1.72-1.65 (m, 2H).
197	453.1	(DMSO-d ₆) δ = 9.00 (s, 1H), 8.00 (s, 1H), 7.70 (d, 1H), 7.45-7.15 (m, 3H), 5.36 (s, 2H), 4.93 (t, 1H), 3.75-3.56 (m, 6H), 2.42 (s, 3H), 2.21-2.14 (m, 2H), 1.67-1.54 (m, 2H).
198 - En 1	466.1	(DMSO-d ₆) δ = 9.00 (s, 1H), 8.75 (d, 1H), 7.73 (d, 1H), 7.51 (d, 1H), 7.41 (t, 1H), 7.23 (dd, 1H), 5.37 (s, 2H), 4.75-4.68 (m, 2H), 3.58-3.50 (m, 2H), 3.48-3.38 (m, 3H), 3.22-3.14 (m, 1H), 2.43 (s, 3H), 2.42-2.32 (m, 1H), 2.08-1.95 (m, 1H).
199 - En 1	506.1	(DMSO-d ₆) δ = 8.86 (d, 1H), 8.74 (d, 1H), 8.26 (d, 1H), 7.81-7.73 (m, 2H), 7.41-7.14 (m, 3H), 5.35 (s, 2H), 4.58-4.39 (m, 1H), 3.16-3.05 (m, 1H), 2.95-2.78 (m, 2H), 2.68-2.55 (m, 1H), 2.50-2.40 (m, 1H), 1.85-1.76 (m, 1H), 1.75-1.60 (m, 1H).
200	421.1	(DMSO-d ₆) δ = 0.71-0.79 (m, 4H), 2.56 (s, 3H), 3.55 (d, 2H), 4.76 (t, 1H), 5.32 (s, 2H), 6.96 (dd, 1H), 7.26 (d, 1H), 7.47 (d, 1H), 7.77-7.80 (m, 1H), 8.18 (s, 1H), 8.26 (d, 1H), 8.73 (d, 1H).
201	388.1	(DMSO-d ₆) δ = 8.98 (s, 1H), 8.04 (s, 1H), 7.45 (d, 1H), 7.30 (d, 1H), 6.95 (dd, 1H), 5.31 (s, 2H), 4.97 (s, 1H), 3.78 (s, 2H), 3.67 (d, 2H), 3.43 (d, 2H), 2.59 (s, 3H), 2.42 (s, 3H).
202	436.1	(DMSO-d ₆) δ = 8.73 (d, 1H), 8.26 (d, 1H), 8.07 (s, 1H), 7.79 (dd, 1H), 7.47 (d, 1H), 7.31 (d, 1H), 6.97 (dd, 1H), 5.31 (s, 2H), 4.95 (s, 1H), 3.78 (s, 2H), 3.64 (d, 2H), 3.41 (d, 2H), 2.59 (s, 3H).
203	409.1	(DMSO) δ = 2.42 (s, 3H), 2.59 (s, 3H), 3.83-3.94 (m, 2H), 4.09-4.16 (m, 1H), 4.31-4.34 (m, 1H), 4.82-4.88 (m, 1H), 5.32 (s, 2H), 6.98 (dd, 1H), 7.25 (s, 1H), 7.49 (d, 1H), 8.39 (d, 1H), 8.98 (s, 1H).
204	400.1	(DMSO) δ = 1.59-1.67 (m, 2H), 2.25 (d, 2H), 2.59 (s, 3H), 3.64-3.71 (m, 9H), 4.95 (t, 1H), 5.14 (s, 2H), 6.86 (d, 1H), 7.00 (dd, 1H), 7.18 (d, 1H), 7.33 (d, 1H), 7.36 (s, 1H), 7.46 (d, 1H).
205 - En 1	405.1	(DMSO) δ = 1.74-1.87 (m, 2H), 2.41-2.49 (m, 1H), 2.57-2.59 (m, 4H), 2.78-2.98 (m, 2H), 3.11-3.19 (m, 1H), 3.68 (s, 3 H), 4.42-4.53 (m, 1H), 5.12 (s, 2H), 6.86 (d, 1H), 7.02 (dd, 1H), 7.18 (d, 1H), 7.26 (d, 1H), 7.48 (d, 1H), 8.19 (d, 1H).
206	441.2	(DMSO-d ₆) δ = 1.31-1.34 (m, 3H), 1.59-1.66 (m, 2H), 2.15 (d, 2H), 2.59 (s, 3H), 3.56-3.72 (m, 6H), 4.36-4.41 (m, 2H), 4.86 (t, 1H), 5.06 (s, 2H), 6.94-7.01 (m, 2H), 7.23 (d, 1H), 7.40 (s, 1H), 7.46 (d, 1H), 7.78 (m, 1H), 8.11-8.13 (m, 1H).
207 - En 1	446.1	(DMSO-d ₆) δ = 1.32 (t, 3H), 1.68-1.80 (m, 2H), 2.44-2.49 (m, 1H), 2.56-2.63 (m, 4H), 2.77-2.93 (m, 2H), 3.07-3.14 (m, 1H), 4.35-4.52 (m, 3H), 5.05 (s, 2H), 6.95-7.01 (m, 2H), 7.19-7.22 (m, 1H), 7.47 (d, 1H), 7.78-7.81 (m, 1H), 8.11-8.13 (m, 1H), 8.22 (d, 1H).

cpd Nr	NMR Data
232	¹ H NMR (400 MHz, DMSO) δ = 8.99 (s, 1H), 7.45 (d, 1H), 7.38 (d, 1H), 6.93 (dd, 1H), 5.33 (s, 2H), 2.62 (s, 3H), 2.43 (s, 3H).
270 - En 1	¹ H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 7.79 (d, 1H), 7.52-7.42 (m, 2H), 6.98 (dd, 1H), 5.32 (s, 2H), 4.30 (d, 1H), 3.80 (dd, 1H), 3.68 (dd, 1H), 2.64 (s, 3H), 2.42 (s, 3H).
271	¹ H NMR (500 MHz, DMSO) δ = 8.98 (s, 1H), 7.88 (d, 1H), 7.47 (s, 1H), 7.47 (d, 1H), 7.36 (d, 1H), 7.09 (s, 1H), 6.96 (dd, 1H), 5.32 (d, 2H), 4.48 (p, 1H), 2.62 (s, 3H), 2.42 (s, 3H), 1.36 (d, 3H).

-continued

cpd Nr	NMR Data
272	¹ H NMR (500 MHz, DMSO) δ = 8.99 (s, 1H), 7.59 (s, 1H), 7.52 (d, 1H), 7.48 (d, 1H), 7.35 (s, 1H), 6.97 (dd, 1H), 5.36-5.25 (m, 2H), 4.94 (td, 1H), 3.73 (dd, 1H), 3.66 (dd, 1H), 3.07 (s, 3H), 2.63 (s, 3H), 2.42 (s, 3H).
273	¹ H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 7.62 (d, 1H), 7.57-7.52 (m, 1H), 7.48 (d, 1H), 7.30 (d, 1H), 7.16 (s, 1H), 6.97 (dd, 1H), 5.32 (s, 2H), 4.37 (dd, 1H), 2.61 (s, 3H), 2.41 (s, 3H), 2.17-2.04 (m, 1H), 0.95 (dd, 6H).
274	¹ H NMR (500 MHz, DMSO) δ = 7.95 (d, 1H), 7.70 (s, 1H), 7.46 (d, 1H), 7.38 (d, 1H), 6.95 (dd, 1H), 5.31 (d, 2H), 5.16 (t, 1H), 4.57-4.51 (m, 1H), 3.60 (dt, 1H), 3.53-3.40 (m, 3H), 3.04 (s, 3H), 2.63 (s, 3H), 2.60 (s, 3H).
275	¹ H NMR (500 MHz, DMSO) δ = 8.99 (s, 1H), 7.96 (d, 1H), 7.47 (d, 1H), 7.41 (d, 1H), 6.96 (dd, 1H), 5.36-5.26 (m, 2H), 5.15 (s, 1H), 4.54 (d, 1H), 3.60 (dd, 1H), 3.53-3.43 (m, 3H), 3.04 (s, 3H), 2.60 (s, 3H), 2.42 (s, 3H).
276	¹ H NMR (500 MHz, DMSO) δ = 8.99 (s, 1H), 7.94 (d, 1H), 7.68 (d, 1H), 7.49 (d, 1H), 7.39 (d, 1H), 6.98 (dd, 1H), 5.32 (s, 2H), 5.02 (t, 1H), 4.49 (dt, 1H), 3.72 (t, 2H), 2.63 (d, 6H), 2.42 (s, 3H).
277	¹ H NMR (500 MHz, DMSO) δ = 8.01 (dd, 1H), 7.71 (s, 1H), 7.47 (d, 1H), 7.30 (d, 1H), 6.97 (dd, 1H), 5.32 (s, 2H), 5.08 (q, 1H), 4.79 (dq, 1H), 4.14 (dt, 1H), 3.93 (q, 1H), 3.89-3.66 (m, 3H), 3.72-3.51 (m, 1H), 2.62 (d, 6H), 2.42 (dq, 1H).
278	¹ H NMR (500 MHz, DMSO) δ = 7.86 (d, 1H), 7.72 (s, 1H), 7.47 (d, 1H), 7.33 (d, 1H), 6.96 (dd, 1H), 5.33 (s, 2H), 5.04 (dt, 1H), 4.99 (t, 1H), 3.71 (dt, 1H), 3.65 (dt, 1H), 3.13 (s, 3H), 2.89 (s, 3H), 2.62 (d, 6H).
279	¹ H NMR (500 MHz, DMSO) δ = 8.98 (s, 1H), 7.88 (d, 1H), 7.47 (s, 1H), 7.47 (d, 1H), 7.36 (d, 1H), 7.09 (s, 1H), 6.96 (dd, 1H), 5.32 (d, 2H), 4.48 (p, 1H), 2.62 (s, 3H), 2.42 (s, 3H), 1.36 (d, 3H).
280	¹ H NMR (500 MHz, DMSO) δ = 8.74 (dd, 1H), 8.27 (dd, 1H), 7.79 (dd, 1H), 7.64 (d, 1H), 7.54-7.45 (m, 2H), 7.43 (d, 1H), 7.18 (s, 1H), 7.00 (dd, 1H), 5.32 (s, 2H), 5.00 (t, 1H), 4.48 (dt, 1H), 3.74 (t, 2H), 2.65 (s, 3H).
281	¹ H NMR (400 MHz, DMSO) δ = 8.27-8.22 (m, 1H), 7.64 (d, 1H), 7.55-7.47 (m, 2H), 7.45 (d, 1H), 7.20 (s, 1H), 7.02 (dd, 1H), 5.53 (s, 2H), 5.02 (t, 1H), 4.49 (dt, 1H), 3.75 (t, 2H), 2.64 (s, 3H).
282	¹ H NMR (500 MHz, DMSO) δ = 7.65 (d, 1H), 7.51 (d, 1H), 7.47 (s, 1H), 7.40 (d, 1H), 7.18 (s, 1H), 6.99 (dd, 1H), 5.45 (d, 2H), 5.01 (t, 1H), 4.47 (dt, 1H), 3.74 (t, 2H), 2.66 (d, 6H).
283	¹ H NMR (400 MHz, DMSO) δ = 7.64 (d, 1H), 7.49 (d, 2H), 7.42 (d, 1H), 7.19 (d, 2H), 6.98 (dd, 1H), 5.13 (s, 2H), 5.07 (t, 1H), 4.49 (dt, 1H), 3.75 (q, 2H), 2.64 (s, 3H), 2.42 (s, 3H).
284	¹ H NMR (500 MHz, DMSO) δ = 8.41 (dd, 1H), 7.83 (dd, 1H), 7.65 (d, 1H), 7.50 (d, 2H), 7.46 (d, 1H), 7.24 (dd, 1H), 7.19 (s, 1H), 7.01 (dd, 1H), 5.16 (s, 2H), 5.05 (s, 1H), 4.48 (dt, 1H), 3.75 (d, 2H), 2.65 (s, 3H), 2.54 (s, 3H).
285	¹ H NMR (500 MHz, DMSO) δ = 7.61 (d, 1H), 7.58 (s, 1H), 7.48 (d, 2H), 7.43 (d, 1H), 7.20 (s, 1H), 6.99 (dd, 1H), 5.13 (s, 2H), 5.03 (t, 1H), 4.49 (dt, 1H), 3.81-3.70 (m, 2H), 2.65 (d, 6H).
286	¹ H NMR (500 MHz, DMSO) δ = 13.70 (s, 1H), 8.13 (s, 1H), 7.65 (d, 1H), 7.52-7.44 (m, 2H), 7.40 (d, 1H), 7.19 (s, 1H), 6.93 (dd, 1H), 5.06 (s, 2H), 5.04 (s, 1H), 4.48 (dt, 1H), 3.78-3.72 (m, 2H), 2.64 (s, 3H).
287	¹ H NMR (500 MHz, DMSO) δ = 8.57 (s, 1H), 8.42 (d, 1H), 7.64 (d, 1H), 7.53-7.44 (m, 3H), 7.28 (d, 1H), 7.19 (s, 1H), 7.01 (dd, 1H), 5.17 (s, 2H), 5.02 (t, 1H), 4.48 (dt, 1H), 3.78-3.71 (m, 2H), 2.65 (s, 3H), 2.38 (s, 3H).
288	¹ H NMR (500 MHz, DMSO) δ = 8.98 (s, 1H), 7.63 (d, 1H), 7.49 (d, 1H), 7.47 (s, 1H), 7.42 (d, 1H), 7.21-7.17 (m, 1H), 6.98 (dd, 1H), 5.32 (s, 2H), 5.01 (t, 1H), 4.48 (dt, 1H), 3.74 (tt, 2H), 2.64 (s, 3H), 2.42 (s, 3H).
289	¹ H NMR (400 MHz, DMSO) δ = 7.62 (d, 1H), 7.53-7.45 (m, 2H), 7.42 (d, 1H), 7.19 (s, 1H), 6.99 (dd, 1H), 6.36 (d, 1H), 5.16 (s, 2H), 5.01 (t, 1H), 4.48 (dt, 1H), 3.74 (tt, 2H), 2.64 (s, 3H), 2.41 (d, 3H).
290	¹ H NMR (500 MHz, DMSO) δ = 8.88 (s, 1H), 7.62 (d, 1H), 7.51-7.42 (m, 3H), 7.19 (s, 1H), 6.99 (dd, 1H), 5.15 (s, 2H), 5.02 (t, 1H), 4.48 (dt, 1H), 3.75 (tt, 2H), 2.64 (s, 3H), 2.53-2.49 (m, 3H).
291	¹ H NMR (500 MHz, DMSO) δ = 8.55 (d, 1H), 7.78 (dd, 1H), 7.62 (d, 1H), 7.48 (s, 1H), 7.48 (d, 1H), 7.42 (d, 1H), 7.28 (d, 1H), 7.19 (s, 1H), 6.98 (dd, 1H), 5.13 (s, 2H), 5.02 (t, 1H), 4.48 (dt, 1H), 3.81-3.69 (m, 2H), 2.64 (s, 3H), 2.47 (s, 3H).
292	¹ H NMR (600 MHz, DMSO) δ = 7.60 (d, 1H), 7.47 (d, 2H), 7.35 (d, 1H), 7.16 (s, 1H), 6.94 (dd, 1H), 5.07-4.99 (m, 1H), 4.73 (dd, 2H), 4.49 (d, 1H), 4.45 (t, 2H), 4.24 (d, 2H), 3.76 (h, 2H), 3.47-3.39 (m, 1H), 2.64 (s, 3H).
293	¹ H NMR (500 MHz, DMSO) δ = 9.22 (dd, 1H), 7.87 (dd, 1H), 7.76 (dd, 1H), 7.64 (d, 1H), 7.52-7.45 (m, 3H), 7.21-7.17 (m, 1H), 7.04 (dd, 1H), 5.43 (s, 2H), 5.02 (t, 1H), 4.48 (dt, 1H), 3.75 (td, 2H), 2.64 (s, 3H).
294	¹ H NMR (600 MHz, DMSO) δ = 9.18 (d, 1H), 8.83 (d, 1H), 7.67 (dd, 1H), 7.61 (d, 1H), 7.50 (d, 1H), 7.46 (s, 1H), 7.43 (d, 1H), 7.16 (s, 1H), 7.03 (dd, 1H), 5.24 (s, 2H), 4.48 (dt, 1H), 3.79-3.71 (m, 2H), 2.64 (s, 3H).
295	¹ H NMR (500 MHz, DMSO) δ = 7.63 (d, 1H), 7.55-7.47 (m, 3H), 7.45 (d, 1H), 7.20 (s, 1H), 7.00 (dd, 1H), 6.54-6.24 (m, 2H), 5.24 (s, 2H), 5.02 (t, 1H), 4.67 (td, 2H), 4.49 (dt, 1H), 3.75 (tt, 2H), 2.64 (s, 3H).
296	¹ H NMR (500 MHz, DMSO) δ = 8.26 (dd, 1H), 8.06 (dd, 1H), 7.98-7.57 (m, 2H), 7.48 (s, 1H), 7.57-7.24 (m, 3H), 7.24-7.09 (m, 1H), 7.00 (dd, 1H), 5.13 (s, 2H), 5.00 (t, 1H), 4.48 (dt, 1H), 3.87-3.62 (m, 2H), 2.65 (s, 3H).
297	¹ H NMR (500 MHz, DMSO) δ = 8.19 (dd, 1H), 7.77 (dd, 1H), 7.63 (d, 1H), 7.53-7.45 (m, 2H), 7.42 (d, 1H), 7.18 (s, 1H), 6.94 (ddd, 2H), 5.08 (s, 2H), 5.00 (t, 1H), 4.48 (dt, 1H), 3.75 (td, 2H), 2.86 (s, 6H), 2.64 (s, 3H).
298	¹ H NMR (500 MHz, DMSO) δ = 7.62 (d, 1H), 7.48 (d, 2H), 7.39 (d, 1H), 7.19 (s, 1H), 6.96 (dd, 1H), 5.23 (s, 2H), 5.01 (t, 1H), 4.48 (dt, 1H), 3.80-3.70 (m, 2H), 2.64 (s, 3H), 2.57 (s, 3H), 2.32 (s, 3H).
299	¹ H NMR (500 MHz, DMSO) δ = 7.62 (d, 1H), 7.51-7.45 (m, 2H), 7.42 (d, 1H), 7.18 (s, 1H), 6.97 (dd, 1H), 5.08-4.99 (m, 3H), 4.48 (dt, 1H), 3.81-3.69 (m, 2H), 2.64 (s, 3H), 2.58 (s, 3H), 2.42 (s, 3H).

-continued

cpd Nr	NMR Data
300	¹ H NMR (500 MHz, DMSO) δ = 7.92 (dd, 1H), 7.61 (d, 1H), 7.55 (dd, 1H), 7.51-7.46 (m, 2H), 7.44 (d, 1H), 7.21-7.17 (m, 1H), 6.99 (dd, 1H), 6.57 (dd, 1H), 5.86 (s, 2H), 5.05-4.96 (m, 3H), 4.49 (dt, 1H), 3.81-3.70 (m, 2H), 2.64 (s, 3H).
301	¹ H NMR (500 MHz, DMSO) δ = 8.85-8.64 (m, 1H), 8.22-8.01 (m, 1H), 7.74-7.16 (m, 6H), 7.29-7.09 (m, 1H), 6.99 (dd, 1H), 5.35 (s, 2H), 5.01 (t, 1H), 4.48 (dt, 1H), 3.75 (td, 2H), 2.64 (s, 3H).
302	¹ H NMR (500 MHz, DMSO) δ = 7.65 (d, 1H), 7.51 (d, 1H), 7.46 (dd, 3H), 7.21-7.16 (m, 1H), 7.01 (dd, 1H), 5.18 (s, 2H), 5.01 (t, 1H), 4.48 (dt, 1H), 3.83 (s, 3H), 3.75 (td, 2H), 2.64 (s, 3H).
303	¹ H NMR (500 MHz, DMSO) δ = 7.79 (d, 1H), 7.71 (s, 1H), 7.56 (s, 1H), 7.48 (d, 1H), 7.35 (d, 1H), 7.24 (s, 1H), 6.96 (dd, 1H), 5.33 (s, 2H), 4.66 (ddd, 1H), 3.73-3.61 (m, 2H), 3.30 (s, 3H), 2.62 (d, 6H).
304	¹ H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 7.62 (d, 1H), 7.58-7.52 (m, 1H), 7.48 (d, 1H), 7.30 (d, 1H), 7.16 (s, 1H), 6.97 (dd, 1H), 5.32 (s, 2H), 4.37 (dd, 1H), 2.61 (s, 3H), 2.41 (s, 3H), 2.11 (h, 1H), 0.95 (dd, 6H).
305	¹ H NMR (500 MHz, DMSO) δ = 7.99-7.93 (m, 1H), 7.72 (s, 1H), 7.67 (d, 1H), 7.48 (d, 1H), 7.38 (d, 1H), 6.98 (dd, 1H), 5.33 (s, 2H), 5.07 (s, 1H), 4.49 (dt, 1H), 3.77-3.67 (m, 2H), 2.67-2.57 (m, 9H).
306	¹ H NMR (500 MHz, DMSO) δ = 8.99 (s, 1H), 7.95 (q, 1H), 7.69 (d, 1H), 7.49 (d, 1H), 7.39 (d, 1H), 6.98 (dd, 1H), 5.32 (s, 2H), 5.05 (s, 1H), 4.49 (dt, 1H), 3.72 (d, 2H), 2.63 (d, 6H), 2.42 (s, 3H).
307	¹ H NMR (400 MHz, DMSO) δ = 7.88 (d, 1H), 7.72 (s, 1H), 7.47 (d, 1H), 7.33 (d, 1H), 6.96 (dd, 1H), 5.33 (s, 2H), 5.02 (s, 1H), 4.83 (q, 1H), 3.76-3.56 (m, 3H), 3.44-3.30 (m, 1H), 2.62 (d, 6H), 1.91 (p, 2H), 1.80 (p, 2H).
308	¹ H NMR (400 MHz, DMSO) δ = 9.01 (s, 1H), 8.71 (s, 1H), 7.69 (d, 1H), 7.58-7.06 (m, 3H), 5.37 (s, 2H), 4.82 (t, 1H), 3.57 (d, 2H), 2.44 (s, 3H), 0.80 (dt, 4H).
309	¹ H NMR (400 MHz, DMSO) δ = 9.01 (s, 1H), 8.95 (s, 1H), 7.72 (d, 1H), 7.78-7.09 (m, 3H), 5.38 (s, 2H), 5.28 (s, 1H), 4.72 (d, 2H), 4.54 (d, 2H), 3.81 (s, 2H), 2.44 (s, 3H).
310 - En 1	¹ H NMR (400 MHz, DMSO) δ = 9.00 (s, 1H), 8.84 (d, 1H), 7.73 (d, 1H), 7.52-7.02 (m, 3H), 5.36 (s, 2H), 4.62-4.37 (m, 1H), 3.13 (s, 1H), 3.03-2.73 (m, 2H), 2.60 (d, 1H), 2.42 (s, 4H), 1.97-1.55 (m, 2H).
311	¹ H NMR (500 MHz, DMSO) δ = 11.77 (s, 1H), 8.00 (s, 1H), 7.80-7.48 (m, 2H), 7.48-7.04 (m, 4H), 6.22 (t, 1H), 4.94 (s, 2H), 4.92 (s, 1H), 3.81-3.46 (m, 6H), 2.17 (d, 2H), 1.60 (ddd, 2H).
312	¹ H NMR (500 MHz, DMSO) δ = 8.17 (dd, 1H), 8.01 (s, 1H), 7.91-7.60 (m, 2H), 7.51-7.12 (m, 3H), 7.03 (dd, 1H), 5.12 (s, 2H), 4.91 (t, 1H), 3.93 (s, 3H), 3.81-3.54 (m, 4H), 3.56 (td, 2H), 2.16 (d, 2H), 1.72-1.47 (m, 2H).
313 - En 1	¹ H NMR (500 MHz, DMSO) δ = 8.73 (dd, 1H), 8.30-8.20 (m, 2H), 7.78 (dd, 1H), 7.44 (d, 1H), 7.19 (d, 1H), 6.96 (dd, 1H), 5.31 (s, 2H), 4.55-4.43 (m, 1H), 3.19-3.04 (m, 1H), 2.97-2.77 (m, 2H), 2.67-2.50 (m, 2H), 1.78 (s, 2H), 1.21-0.98 (m, 4H).
314	¹ H NMR (500 MHz, DMSO) δ = 8.98 (s, 1H), 7.43-7.37 (m, 2H), 7.25 (d, 1H), 6.92 (dd, 1H), 5.30 (s, 2H), 4.91 (t, 1H), 3.75-3.58 (m, 6H), 2.60-2.50 (m, 1H), 2.41 (s, 3H), 2.20 (d, 2H), 1.65 (ddd, 2H), 1.10 (dt, 2H), 1.11-1.02 (m, 2H).
315 - En 1	¹ H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 8.59 (d, 1H), 8.08 (d, 1H), 7.47 (d, 1H), 7.33 (d, 1H), 7.19 (d, 1H), 6.97 (dd, 1H), 5.40-5.28 (m, 1H), 5.32 (s, 2H), 2.61 (s, 3H), 2.42 (s, 3H), 1.58 (d, 3H).
316 - En 1	¹ H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 8.44 (d, 1H), 7.50 (d, 1H), 7.24 (d, 1H), 6.99 (dd, 1H), 5.31 (s, 2H), 5.24 (t, 1H), 4.81 (t, 1H), 3.77 (dp, 2H), 2.59 (s, 3H), 2.42 (s, 3H).
317 - En 1	¹ H NMR (500 MHz, DMSO) δ = 8.99 (s, 1H), 8.26 (dd, 1H), 7.69 (d, 1H), 7.53-7.45 (m, 2H), 6.99 (dd, 1H), 5.38-5.28 (m, 2H), 4.86 (ddd, 1H), 3.92 (td, 2H), 3.55 (ddd, 1H), 3.47 (dd, 1H), 3.38-3.32 (m, 1H), 3.10 (ddd, 1H), 2.66 (s, 3H), 2.42 (s, 3H).
318	¹ H NMR (500 MHz, DMSO) δ = 8.24 (s, 1H), 7.70 (s, 1H), 7.50-7.43 (m, 2H), 7.37 (d, 1H), 6.97 (dd, 1H), 5.34 (s, 2H), 2.63 (d, 6H), 2.62 (d, 3H), 2.55 (dd, 1H), 2.26 (ddd, 2H), 2.00-1.80 (m, 2H).
319	¹ H NMR (500 MHz, DMSO) δ = 8.99 (s, 1H), 8.24 (s, 1H), 7.50-7.42 (m, 2H), 7.37 (d, 1H), 6.97 (dd, 1H), 5.34 (s, 2H), 2.66-2.57 (m, 6H), 2.61-2.51 (m, 2H), 2.43 (s, 3H), 2.26 (tdt, 2H), 2.00-1.79 (m, 2H).
320	¹ H NMR (500 MHz, DMSO) δ = 8.58 (d, 1H), 8.24 (s, 1H), 7.84 (t, 1H), 7.56 (d, 1H), 7.46 (dd, 2H), 7.36 (dd, 2H), 7.01 (d, 1H), 5.23 (s, 2H), 2.65-2.59 (m, 5H), 2.24 (q, 2H), 1.93 (s, 1H), 1.86 (d, 1H).
321	¹ H NMR (500 MHz, DMSO) δ = 8.99 (s, 1H), 7.99 (t, 1H), 7.87 (d, 1H), 7.51-7.42 (m, 2H), 6.97 (dd, 1H), 5.33 (s, 2H), 3.88 (d, 2H), 2.67-2.60 (m, 6H), 2.43 (s, 3H).
322	¹ H NMR (600 MHz, DMSO) δ = 8.99 (s, 1H), 7.92 (s, 1H), 7.47 (d, 1H), 7.38 (d, 1H), 7.15 (t, 1H), 6.96 (dd, 1H), 5.33 (s, 2H), 3.42 (q, 1H), 3.17 (q, 2H), 2.93 (s, 3H), 2.62 (s, 3H), 2.43 (s, 3H).
323	¹ H NMR (600 MHz, DMSO) δ = 8.99 (s, 1H), 8.08 (t, 1H), 7.48 (d, 1H), 7.43 (d, 1H), 6.97 (dd, 1H), 5.33 (s, 2H), 3.74 (td, 2H), 3.44 (t, 2H), 3.07 (s, 3H), 2.63 (s, 3H), 2.43 (s, 3H).
324	¹ H NMR (400 MHz, DMSO) δ = 8.99 (s, 1H), 8.67 (s, 1H), 7.72 (d, 1H), 7.54-7.43 (m, 2H), 6.99 (dd, 1H), 5.35 (s, 2H), 4.86 (d, 2H), 4.68 (d, 2H), 2.70-2.61 (m, 6H), 2.43 (s, 3H).
325	¹ H NMR (400 MHz, DMSO) δ = 8.58 (dd, 1H), 8.15 (dd, 2H), 7.68-7.57 (m, 2H), 7.48 (d, 1H), 7.39 (s, 1H), 7.22 (d, 1H), 6.97 (dd, 1H), 5.59 (s, 2H), 4.86 (s, 1H), 3.74-3.50 (m, 6H), 2.61 (s, 3H), 2.14 (d, 2H), 1.62 (ddd, 2H).
326 - En 1	¹ H NMR (400 MHz, DMSO) δ = 8.99 (s, 1H), 8.43 (d, 1H), 7.35 (dd, 1H), 7.24 (dd, 1H), 5.36 (s, 2H), 4.72 (s, 1H), 4.64 (dt, 1H), 3.56-3.32 (m, 5H), 3.18 (dt, 1H), 2.55 (s, 3H), 2.39 (s, 3H), 2.44-2.31 (m, 1H), 1.89 (dq, 1H).
327	¹ H NMR (500 MHz, DMSO) δ = 8.99 (s, 1H), 8.42 (s, 1H), 7.33 (d, 1H), 7.22 (t, 1H), 5.36 (s, 2H), 4.73 (s, 1H), 3.55 (d, 2H), 2.39 (s, 3H), 0.83-0.75 (m, 2H), 0.71-0.65 (m, 2H).
328	¹ H NMR (400 MHz, DMSO) δ = 8.99 (s, 1H), 8.65 (s, 1H), 7.36 (dd, 1H), 7.25 (dd, 1H), 5.37 (s, 2H), 5.18 (t, 1H), 4.65-4.53 (m, 4H), 3.73 (d, 2H), 2.53 (s, 3H), 2.39 (s, 3H).
329	¹ H NMR (400 MHz, DMSO) δ = 11.75 (s, 1H), 7.72 (s, 1H), 7.62-7.54 (m, 1H), 7.39 (dd, 1H), 7.32 (dd, 1H), 7.25-7.15 (m, 1H), 6.23 (t, 1H), 4.95 (s, 2H), 4.81 (t, 1H), 3.72-3.55 (m, 6H), 2.52 (s, 3H), 2.15-2.05 (m, 2H), 1.70-1.57 (m, 1H).
330	¹ H NMR (500 MHz, DMSO) δ = 8.17 (dd, 1H), 7.81 (dd, 1H), 7.73 (s, 1H), 7.33 (d, 1H), 7.20 (t, 1H), 7.03 (dd, 1H), 5.13 (s, 2H), 4.81 (t, 1H), 3.91 (s, 3H), 3.72-3.56 (m, 6H), 2.52 (s, 3H), 2.14-2.06 (m, 2H), 1.63 (ddd, 2H).

-continued

cpd Nr	NMR Data
331	¹ H NMR (500 MHz, DMSO) δ = 7.47 (d, 1H), 7.37 (s, 1H), 7.32 (d, 1H), 7.26 (d, 1H), 7.10-6.85 (m, 2H), 6.55-6.13 (m, 1H), 5.18 (s, 2H), 4.91 (t, 1H), 4.58 (td, 2H), 3.82-3.52 (m, 6H), 2.60 (s, 3H), 2.20 (d, 2H), 1.76-1.51 (m, 2H).
332 - En 1	¹ H NMR (500 MHz, DMSO) δ = 7.69-7.30 (m, 4H), 7.31 (s, 1H), 6.99 (d, 1H), 6.58-6.16 (m, 2H), 5.23 (s, 3H), 4.67 (td, 2H), 3.74 (d, 1H), 3.64 (d, 1H), 3.38-3.20 (m, 1H), 3.20-3.02 (m, 1H), 2.61 (s, 3H), 2.20 (dt, 2H), 1.79 (q, 2H).
333 - En 1	¹ H NMR (500 MHz, DMSO) δ = 7.79 (s, 1H), 7.71-7.39 (m, 3H), 7.29 (d, 1H), 6.99 (dd, 1H), 6.59-6.16 (m, 2H), 5.23 (s, 3H), 4.68 (td, 2H), 3.61 (q, 2H), 3.41-3.08 (m, 2H), 2.60 (s, 3H), 2.63-2.28 (m, 2H).
334	¹ H NMR (400 MHz, DMSO) δ = 7.52 (d, 1H), 7.47 (d, 1H), 7.40 (s, 1H), 7.28 (d, 1H), 6.98 (dd, 1H), 6.65-6.11 (m, 2H), 5.23 (s, 2H), 4.89 (t, 1H), 4.67 (td, 2H), 3.84-3.47 (m, 6H), 2.61 (s, 3H), 2.18 (d, 2H), 1.67 (d, 0H), 1.64 (s, 1H).
335 - En 1	¹ H NMR (400 MHz, DMSO) δ = 8.18 (d, 1H), 8.09 (dd, 1H), 7.60 (dd, 1H), 7.47 (d, 1H), 7.36 (d, 1H), 6.97 (dd, 1H), 6.69 (dd, 1H), 4.95 (s, 2H), 4.70 (dt, 2H), 4.08 (t, 4H), 3.52 (d, 2H), 3.49-3.36 (m, 3H), 3.17 (dt, 1H), 2.62 (s, 3H), 2.23 (p, 2H), 2.05-1.89 (m, 1H).
336 - En 1	¹ H NMR (400 MHz, DMSO) δ = 8.22 (d, 1H), 8.09 (dd, 1H), 7.58 (dd, 1H), 7.47 (d, 1H), 7.21 (d, 1H), 6.97 (dd, 1H), 6.69 (dd, 1H), 4.93 (s, 2H), 4.46 (dd, 1H), 4.07 (t, 4H), 3.20-3.04 (m, 1H), 2.98-2.74 (m, 2H), 2.62 (d, 1H), 2.57 (s, 3H), 2.43 (s, 1H), 2.23 (p, 2H), 1.83-1.62 (m, 2H).
337	¹ H NMR (400 MHz, DMSO) δ = 8.08 (dd, 1H), 7.59 (dd, 1H), 7.46 (d, 1H), 7.41 (s, 1H), 7.25 (d, 1H), 6.96 (dd, 1H), 6.69 (dd, 1H), 4.94 (s, 2H), 4.89 (t, 1H), 4.08 (t, 4H), 3.75-3.66 (m, 2H), 3.71-3.56 (m, 4H), 2.60 (s, 3H), 2.29-2.13 (m, 4H), 1.70-1.57 (m, 2H).
338 - En 1	¹ H NMR (500 MHz, DMSO) δ = 8.36-8.14 (m, 2H), 8.04 (dd, 1H), 7.77 (t, 1H), 7.49 (d, 1H), 7.40-7.12 (m, 2H), 7.00 (dd, 1H), 5.24-5.00 (m, 2H), 4.47 (dt, 1H), 3.26-3.00 (m, 1H), 3.00-2.70 (m, 2H), 2.58 (d, 1H), 2.57 (s, 3H), 2.42 (s, 1H), 1.90-1.57 (m, 2H).
339	¹ H NMR (500 MHz, DMSO) δ = 8.25 (dd, 1H), 8.04 (dd, 1H), 7.76 (t, 1H), 7.55-7.16 (m, 4H), 6.98 (dd, 1H), 5.13 (s, 2H), 4.88 (s, 1H), 3.80-3.47 (m, 6H), 2.60 (s, 3H), 2.16 (d, 2H), 1.62 (ddd, 2H).
340 - En 1	¹ H NMR (500 MHz, DMSO) δ = 8.67 (dd, 1H), 8.20 (d, 1H), 8.13 (d, 1H), 7.75-6.87 (m, 5H), 5.36 (s, 2H), 4.69 (dt, 1H), 3.53 (t, 2H), 3.48-3.35 (m, 3H), 3.25-3.05 (m, 1H), 2.62 (s, 3H), 2.48-2.23 (m, 1H), 1.96 (dq, 1H).
341	¹ H NMR (400 MHz, DMSO) δ = 8.67 (dd, 1H), 8.36 (s, 1H), 8.13 (dd, 1H), 7.78-7.36 (m, 2H), 7.48-6.86 (m, 3H), 5.36 (s, 2H), 5.28-5.06 (m, 1H), 4.67 (d, 2H), 4.55 (d, 2H), 3.77 (d, 2H), 2.61 (s, 3H).
342 - En 1	¹ H NMR (400 MHz, DMSO) δ = 8.66 (dd, 1H), 8.22 (d, 1H), 8.12 (d, 1H), 7.77-7.37 (m, 2H), 7.37-6.86 (m, 3H), 5.34 (s, 2H), 4.73-4.20 (m, 1H), 3.11 (s, 1H), 3.02-2.65 (m, 2H), 2.57 (s, 3H), 2.42 (s, 1H), 1.93-1.53 (m, 2H).
343	¹ H NMR (400 MHz, DMSO) δ = 8.67 (dd, 1H), 8.26-8.00 (m, 1H), 7.78-7.29 (m, 3H), 7.44-6.85 (m, 3H), 5.35 (s, 2H), 4.88 (t, 1H), 3.84-3.46 (m, 6H), 2.61 (s, 3H), 2.16 (d, 2H), 1.87-1.47 (m, 2H).
344	¹ H NMR (500 MHz, DMSO) δ = 8.18 (dd, 1H), 7.76 (dd, 1H), 7.48-7.39 (m, 2H), 7.23 (d, 1H), 6.97-6.90 (m, 2H), 5.08 (s, 2H), 4.88 (s, 1H), 3.74-3.67 (m, 2H), 3.67-3.57 (m, 4H), 2.85 (s, 6H), 2.59 (s, 3H), 2.17 (d, 2H), 1.68-1.58 (m, 2H).
345	¹ H NMR (400 MHz, DMSO) δ = 7.45 (d, 1H), 7.40 (s, 1H), 7.23 (d, 1H), 6.93 (dd, 1H), 5.22 (s, 2H), 4.89 (t, 1H), 3.75-3.56 (m, 6H), 2.60 (s, 3H), 2.57 (s, 3H), 2.32 (s, 3H), 2.17 (d, 2H), 1.63 (ddd, 2H).
346	¹ H NMR (500 MHz, DMSO) δ = 7.92 (dd, 1H), 7.51 (dd, 1H), 7.46 (d, 1H), 7.37 (s, 1H), 7.27 (d, 1H), 6.97 (dd, 1H), 6.56 (dd, 1H), 5.85 (s, 2H), 4.97 (s, 2H), 4.90 (s, 1H), 3.70 (dt, 2H), 3.66 (s, 2H), 3.61 (td, 2H), 2.61 (s, 3H), 2.17 (d, 2H), 1.64 (ddd, 2H).
347	¹ H NMR (400 MHz, DMSO) δ = 8.75 (dd, 1H), 8.22 (dd, 1H), 7.80 (dd, 1H), 7.50 (d, 1H), 7.43 (s, 1H), 7.30 (d, 1H), 7.01 (dd, 1H), 5.33 (s, 2H), 4.88 (s, 1H), 3.74-3.55 (m, 6H), 2.61 (s, 3H), 2.17 (d, 2H), 1.70-1.57 (m, 2H).
348 - En 1	¹ H NMR (400 MHz, DMSO) δ = 8.20 (d, 1H), 8.13 (dd, 1H), 7.82 (dd, 1H), 7.47 (d, 1H), 7.35 (d, 1H), 6.99 (ddd, 2H), 5.07 (s, 2H), 4.77-4.61 (m, 1H), 4.39 (q, 2H), 3.52 (q, 2H), 3.49-3.36 (m, 3H), 3.17 (dt, 1H), 2.61 (s, 3H), 2.41-2.31 (m, 1H), 1.96 (p, 1H), 1.33 (t, 3H), 1.23 (s, 1H).
349 - En 1	¹ H NMR (400 MHz, DMSO) δ = 11.76 (s, 1H), 7.61-7.54 (m, 1H), 7.48 (dd, 2H), 7.42-7.31 (m, 2H), 6.95 (dd, 1H), 6.22 (t, 1H), 5.65 (s, 1H), 4.89 (s, 2H), 4.45-4.34 (m, 1H), 4.31 (ddd, 1H), 4.02-3.90 (m, 2H), 3.63 (dd, 1H), 3.58 (t, 1H), 2.63 (s, 3H).
350 - En 1	¹ H NMR (500 MHz, DMSO) δ = 8.15 (d, 1H), 7.54 (dd, 1H), 7.47-7.38 (m, 2H), 7.34 (s, 1H), 6.92 (dd, 1H), 6.20 (t, 1H), 4.96-4.86 (m, 2H), 4.33 (dt, 1H), 4.19 (tt, 1H), 4.05 (dd, 1H), 3.91 (dd, 1H), 3.63 (dd, 1H), 3.54 (dd, 1H), 2.57 (s, 3H).
351	¹ H NMR (500 MHz, DMSO) δ = 8.73 (d, 1H), 8.11 (d, 1H), 7.69-7.16 (m, 5H), 6.96 (dd, 1H), 5.36 (s, 2H), 4.88 (t, 1H), 3.71 (d, 2H), 3.75-3.52 (m, 4H), 2.59 (s, 3H), 2.17 (d, 2H), 1.64 (s, 2H).
352	¹ H NMR (500 MHz, DMSO) δ = 8.45 (dt, 1H), 7.79 (ddd, 1H), 7.53 (dt, 1H), 7.45 (d, 1H), 7.40 (s, 1H), 7.30 (d, 1H), 6.96 (dd, 1H), 5.25 (d, 2H), 4.88 (t, 1H), 3.74-3.58 (m, 6H), 2.59 (s, 3H), 2.18 (d, 2H), 1.63 (ddd, 2H).
355 - En 1	¹ H NMR (500 MHz, DMSO) δ = 7.58 (d, 1H), 7.55-7.45 (m, 3H), 7.31 (d, 1H), 7.00 (dd, 1H), 5.22 (t, 1H), 5.17 (s, 2H), 3.83 (s, 3H), 3.73 (dd, 1H), 3.64 (dd, 1H), 3.27 (dt, 1H), 3.16-3.09 (m, 1H), 2.60 (s, 3H), 2.27-2.14 (m, 2H), 1.85-1.73 (m, 1H).
356 - En 1	¹ H NMR (500 MHz, DMSO) δ = 7.78 (s, 1H), 7.54-7.45 (m, 3H), 7.29 (d, 1H), 6.99 (dd, 1H), 5.19 (d, 3H), 3.83 (s, 3H), 3.60 (qd, 2H), 3.20 (q, 1H), 2.60 (s, 3H), 2.49-2.35 (m, 2H).
357	¹ H NMR (400 MHz, DMSO) δ = 7.51-7.44 (m, 2H), 7.41 (s, 1H), 7.29 (d, 1H), 6.97 (dd, 1H), 5.17 (s, 2H), 4.89 (s, 1H), 3.83 (d, 3H), 3.76-3.56 (m, 6H), 2.60 (s, 3H), 2.18 (d, 2H), 1.64 (ddd, 2H).
358	¹ H NMR (500 MHz, DMSO) δ = 8.28 (dd, 1H), 7.50-7.40 (m, 3H), 7.39 (s, 1H), 7.24 (d, 1H), 6.96 (dd, 1H), 5.12 (s, 2H), 4.90 (s, 2H), 4.07 (dd, 2H), 3.74-3.57 (m, 8H), 2.59 (s, 3H), 2.21-2.13 (m, 2H), 1.63 (ddd, 2H).
359	¹ H NMR (400 MHz, DMSO) δ = 8.58 (d, 1H), 7.77 (td, 1H), 7.62 (dd, 1H), 7.46 (d, 1H), 7.39 (s, 1H), 7.24 (d, 1H), 6.98 (dd, 1H), 5.20 (s, 2H), 4.88 (t, 1H), 3.74-3.54 (m, 6H), 2.59 (s, 3H), 2.16 (d, 2H), 1.69-1.56 (m, 2H).

-continued

cpd Nr	NMR Data
360 - En 1	1H NMR (400 MHz, DMSO) δ = 9.00 (s, 1H), 8.24 (d, 1H), 7.62 (d, 1H), 7.56 (d, 1H), 5.40 (s, 2H), 4.69 (dt, 1H), 3.53 (s, 2H), 3.51-3.36 (m, 3H), 3.18 (dt, 1H), 2.61 (s, 3H), 2.42 (s, 3H), 2.38 (dd, 1H), 1.97 (dq, 1H).
361	1H NMR (400 MHz, DMSO) δ = 9.00 (s, 1H), 8.20 (s, 1H), 7.59 (d, 1H), 7.46 (d, 1H), 5.39 (s, 2H), 4.79 (s, 1H), 3.57 (s, 2H), 2.55 (s, 3H), 2.42 (s, 3H), 0.77 (dt, 4H).
362	1H NMR (400 MHz, DMSO) δ = 9.00 (s, 1H), 8.39 (s, 1H), 7.62 (d, 1H), 7.51 (d, 1H), 5.40 (s, 2H), 5.22 (s, 1H), 4.68 (d, 2H), 4.56 (d, 2H), 3.79 (d, 2H), 2.60 (s, 3H), 2.42 (s, 3H).
363	1H NMR (500 MHz, DMSO) δ = 7.70 (dd, 2H), 7.55 (d, 1H), 7.40 (d, 1H), 6.23 (t, 1H), 4.97 (s, 2H), 4.08 (s, 2H), 3.72 (t, 2H), 3.62 (d, 2H), 2.74 (s, 3H), 1.75-1.64 (m, 2H), 1.33 (d, 2H).
364	1H NMR (500 MHz, DMSO) δ = 8.17 (dd, 1H), 7.81 (dd, 1H), 7.62 (d, 1H), 7.44 (s, 1H), 7.39 (d, 1H), 7.04 (dd, 1H), 5.14 (s, 2H), 4.88 (t, 1H), 3.93 (s, 3H), 3.74-3.62 (m, 4H), 3.58 (td, 2H), 2.60 (s, 3H), 2.15 (d, 2H), 1.62 (ddd, 2H).
365 - En 1	1H NMR (500 MHz, DMSO) δ = 9.00 (s, 1H), 8.29 (d, 1H), 7.20 (d, 1H), 7.02 (dd, 1H), 5.34 (s, 2H), 4.68 (dt, 1H), 3.52 (t, 2H), 3.47-3.37 (m, 3H), 3.22-3.13 (m, 1H), 2.64 (s, 3H), 2.43 (s, 3H), 2.35 (s, 1H), 1.96 (dq, 1H).
366	1H NMR (500 MHz, DMSO) δ = 9.00 (s, 1H), 8.25 (s, 1H), 7.10 (d, 1H), 6.98 (dd, 1H), 5.34 (s, 2H), 4.78 (d, 1H), 3.55 (d, 2H), 2.58 (s, 3H), 2.43 (s, 3H), 0.76 (dt, 4H).
367	1H NMR (500 MHz, DMSO) δ = 9.00 (s, 1H), 8.44 (s, 1H), 7.16 (d, 1H), 7.02 (dd, 1H), 5.34 (s, 2H), 5.20 (t, 1H), 4.67 (d, 2H), 4.54 (d, 2H), 3.78 (d, 2H), 2.63 (s, 3H), 2.43 (s, 3H).
368	1H NMR (400 MHz, DMSO) δ = 11.77 (s, 1H), 7.58 (dd, 1H), 7.51 (s, 1H), 7.39 (d, 1H), 7.05 (d, 1H), 6.96 (dd, 1H), 6.22 (t, 1H), 4.89 (d, 3H), 3.74-3.53 (m, 6H), 2.62 (s, 3H), 2.16 (d, 2H), 1.68-1.55 (m, 2H).
369	1H NMR (400 MHz, DMSO) δ = 8.16 (dd, 1H), 7.82 (dd, 1H), 7.52 (s, 1H), 7.07 (d, 1H), 7.07-6.94 (m, 2H), 5.08 (s, 2H), 4.88 (t, 1H), 3.92 (s, 3H), 3.75-3.52 (m, 6H), 2.62 (s, 3H), 2.15 (d, 2H), 1.62 (ddd, 2H).
370	1H NMR (500 MHz, DMSO) δ = 8.43 (t, 1H), 7.69 (s, 1H), 7.48 (d, 1H), 7.34 (d, 1H), 6.97 (dd, 1H), 5.32 (s, 2H), 3.44 (d, 2H), 2.62 (d, 6H), 1.29-1.20 (m, 2H), 1.20-1.11 (m, 2H).
371	1H NMR (500 MHz, DMSO) δ = 8.99 (s, 1H), 8.45 (t, 1H), 7.49 (d, 1H), 7.36 (d, 1H), 6.98 (dd, 1H), 5.32 (s, 2H), 3.44 (d, 2H), 2.62 (s, 3H), 2.41 (s, 3H), 1.26 (q, 2H), 1.15 (q, 2H).
372	1H NMR (500 MHz, DMSO) δ = 8.29 (t, 1H), 7.69 (s, 1H), 7.47 (d, 1H), 7.28 (d, 1H), 6.96 (dd, 1H), 5.32 (s, 2H), 3.72 (dd, 2H), 2.61 (d, 6H), 1.10-0.92 (m, 2H), 0.92-0.76 (m, 1H).
373	1H NMR (500 MHz, DMSO) δ = 8.98 (s, 1H), 8.30 (t, 1H), 7.48 (d, 1H), 7.29 (d, 1H), 6.97 (dd, 1H), 5.32 (s, 2H), 3.72 (dd, 2H), 2.60 (s, 3H), 2.42 (s, 3H), 1.13-0.91 (m, 2H), 0.98-0.77 (m, 2H).
374	1H NMR (600 MHz, CDCl ₃) δ = 8.72 (s, 1H), 7.57 (d, 1H), 7.37 (d, 1H), 7.32 (d, 1H), 6.94 (dd, 1H), 6.57 (s, 1H), 6.33 (d, 1H), 5.25 (s, 2H), 4.73 (d, 2H), 2.74 (s, 3H), 2.50 (s, 3H).
375	1H NMR (500 MHz, DMSO) δ = 7.51 (d, 1H), 7.48-7.42 (m, 2H), 7.37 (d, 1H), 7.18 (d, 2H), 6.99 (dd, 1H), 5.12 (s, 2H), 5.07 (d, 1H), 4.39 (dd, 1H), 4.14 (td, 1H), 2.65 (s, 3H), 2.41 (s, 3H), 1.14 (d, 3H).
376	1H NMR (500 MHz, DMSO) δ = 8.41 (dd, 1H), 7.82 (dd, 1H), 7.54-7.47 (m, 2H), 7.46 (s, 1H), 7.40 (d, 1H), 7.24 (dd, 1H), 7.19-7.15 (m, 1H), 7.03 (dd, 1H), 5.16 (s, 2H), 5.08 (d, 1H), 4.37 (dd, 1H), 4.14 (td, 1H), 2.65 (s, 3H), 2.53 (s, 3H), 1.14 (d, 3H).
377	1H NMR (500 MHz, DMSO) δ = 7.72 (s, 1H), 7.50 (d, 1H), 7.48-7.41 (m, 2H), 7.36 (d, 1H), 7.18 (s, 1H), 6.98 (dd, 1H), 5.33 (s, 2H), 5.07 (d, 1H), 4.38 (dd, 1H), 4.14 (td, 1H), 2.65 (s, 3H), 2.63 (s, 3H), 1.14 (d, 3H).
378	1H NMR (500 MHz, DMSO) δ = 8.98 (s, 1H), 7.50 (d, 1H), 7.44 (d, 2H), 7.38 (d, 1H), 7.16 (s, 1H), 6.99 (dd, 1H), 5.32 (s, 2H), 5.06 (d, 1H), 4.37 (dd, 1H), 4.19-4.09 (m, 1H), 2.65 (s, 3H), 2.42 (s, 3H), 1.14 (d, 3H).
379	1H NMR (600 MHz, DMSO) δ = 8.98 (s, 1H), 7.51 (d, 1H), 7.48-7.40 (m, 2H), 7.37 (d, 1H), 7.14 (s, 1H), 7.00 (dd, 1H), 5.32 (s, 2H), 5.05 (d, 1H), 4.38 (dd, 1H), 4.15 (t, 1H), 2.66 (s, 3H), 2.42 (s, 3H), 1.15 (d, 3H).
380	1H NMR (500 MHz, DMSO) δ = 8.54 (d, 1H), 7.78 (dd, 1H), 7.50 (d, 1H), 7.46 (s, 1H), 7.45 (d, 1H), 7.38 (d, 1H), 7.27 (d, 1H), 7.19-7.15 (m, 1H), 6.99 (dd, 1H), 5.13 (s, 2H), 5.09 (s, 1H), 4.37 (dd, 1H), 4.18-4.10 (m, 1H), 2.65 (s, 3H), 2.47 (s, 3H), 1.14 (d, 3H).
381	1H NMR (500 MHz, DMSO) δ = 7.62-7.28 (m, 5H), 7.28-6.90 (m, 2H), 6.59-6.16 (m, 2H), 5.24 (s, 2H), 5.07 (s, 1H), 4.67 (td, 2H), 4.38 (dd, 1H), 4.14 (s, 1H), 2.66 (s, 3H), 1.14 (d, 3H).
382	1H NMR (500 MHz, DMSO) δ = 7.50 (d, 1H), 7.46-7.40 (m, 2H), 7.38 (d, 1H), 7.17 (s, 1H), 6.97 (dd, 1H), 5.23 (s, 2H), 5.06 (d, 1H), 4.37 (dd, 1H), 4.18-4.09 (m, 1H), 2.65 (s, 3H), 2.57 (s, 3H), 2.32 (s, 3H), 1.14 (d, 3H).
383	1H NMR (500 MHz, DMSO) δ = 7.52 (d, 1H), 7.50-7.43 (m, 3H), 7.41 (d, 1H), 7.17 (s, 1H), 7.02 (dd, 1H), 5.17 (s, 2H), 5.06 (d, 1H), 4.37 (dd, 1H), 4.14 (s, 1H), 3.83 (s, 3H), 2.66 (s, 3H), 1.14 (d, 3H).
384	1H NMR (500 MHz, DMSO) δ = 8.35 (t, 1H), 7.69 (s, 1H), 7.48 (d, 1H), 7.27 (d, 1H), 6.97 (dd, 1H), 5.32 (s, 2H), 4.79-4.60 (m, 4H), 3.82 (dd, 2H), 2.61 (d, 6H).
385	1H NMR (500 MHz, DMSO) δ = 8.99 (s, 1H), 8.36 (t, 1H), 7.48 (d, 1H), 7.29 (d, 1H), 6.97 (dd, 1H), 5.31 (s, 2H), 4.79-4.60 (m, 4H), 3.82 (dd, 2H), 2.59 (s, 3H), 2.42 (s, 3H).
386 - En 1	1H NMR (500 MHz, DMSO) δ = 7.70 (s, 1H), 7.48 (dd, 2H), 7.34 (d, 1H), 6.97 (dd, 1H), 5.61 (s, 1H), 5.31 (s, 2H), 4.45-4.36 (m, 1H), 4.31 (td, 1H), 4.00-3.91 (m, 2H), 3.64 (dd, 1H), 3.58 (t, 1H), 2.63 (d, 6H).
387 - En 1	1H NMR (400 MHz, DMSO) δ = 8.59 (ddd, 1H), 7.84 (td, 1H), 7.60-7.53 (m, 1H), 7.48 (dd, 2H), 7.40-7.31 (m, 2H), 7.00 (dd, 1H), 5.60 (s, 1H), 5.19 (s, 2H), 4.46-4.30 (m, 1H), 4.30 (s, 1H), 4.01-3.90 (m, 2H), 3.64 (dd, 1H), 3.58 (t, 1H), 2.63 (s, 3H).
388 - En 1	1H NMR (500 MHz, DMSO) δ = 8.13 (d, 1H), 7.69 (s, 1H), 7.46 (d, 1H), 7.21 (d, 1H), 6.95 (dd, 1H), 5.32 (s, 3H), 4.23 (ddq, 2H), 4.01 (dd, 1H), 3.92 (dd, 1H), 3.66 (dd, 1H), 3.56 (dd, 1H), 2.63 (s, 3H), 2.56 (s, 3H).
389 - En 1	1H NMR (400 MHz, DMSO) δ = 8.58 (ddd, 1H), 8.13 (d, 1H), 7.84 (td, 1H), 7.54 (d, 1H), 7.47 (d, 1H), 7.35 (ddd, 1H), 7.21 (d, 1H), 6.99 (dd, 1H), 5.31 (s, 1H), 5.20 (s, 2H), 4.23 (s, 2H), 4.01 (dd, 1H), 3.91 (dd, 1H), 3.65 (dd, 1H), 3.56 (dd, 1H), 2.56 (s, 3H).

-continued

cpd Nr	NMR Data
390	1H NMR (500 MHz, DMSO) δ = 8.74 (dd, 1H), 8.28 (d, 1H), 8.18 (d, 1H), 7.83-7.76 (m, 2H), 7.50 (d, 1H), 7.42 (d, 1H), 6.99 (dd, 1H), 5.57 (s, 1H), 5.32 (s, 2H), 4.45-4.35 (m, 2H), 3.48-3.40 (m, 1H), 3.02-2.94 (m, 1H), 2.63 (s, 3H).
391	1H NMR (500 MHz, DMSO) δ = 8.17-8.11 (m, 1H), 7.81 (d, 1H), 7.71 (s, 1H), 7.47 (d, 1H), 7.39 (d, 1H), 6.96 (dd, 1H), 5.58 (s, 1H), 5.32 (s, 2H), 4.40 (dd, 2H), 3.44 (td, 1H), 3.02-2.94 (m, 1H), 2.63 (d, 6H).
392	1H NMR (500 MHz, DMSO) δ = 8.99 (s, 1H), 8.15 (d, 1H), 7.80 (d, 1H), 7.48 (d, 1H), 7.41 (d, 1H), 6.97 (dd, 1H), 5.57 (s, 1H), 5.32 (s, 2H), 4.46-4.35 (m, 2H), 3.44 (ddd, 1H), 2.98 (dd, 1H), 2.63 (s, 3H), 2.43 (s, 3H).
393	1H NMR (400 MHz, DMSO) δ = 8.19 (t, 1H), 7.68 (s, 1H), 7.47 (d, 1H), 7.26 (d, 1H), 6.96 (dd, 1H), 5.32 (s, 2H), 3.75 (dt, 2H), 3.66-3.47 (m, 4H), 2.61 (d, 6H), 1.91-1.66 (m, 4H).
394	1H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 8.19 (t, 1H), 7.47 (d, 1H), 7.27 (d, 1H), 6.97 (dd, 1H), 5.31 (s, 2H), 3.75 (dt, 2H), 3.57 (dq, 4H), 2.59 (s, 3H), 2.41 (s, 3H), 1.92-1.66 (m, 4H).
395 - Dia 1	1H NMR (500 MHz, DMSO) δ = 7.64 (s, 1H), 7.57 (d, 1H), 7.50 (s, 1H), 7.44 (d, 1H), 7.37 (d, 1H), 7.21 (s, 1H), 6.94 (dd, 1H), 5.82 (q, 1H), 5.05 (t, 1H), 4.47 (dt, 1H), 3.76 (hept, 2H), 2.62 (s, 3H), 2.58 (s, 3H), 1.65 (d, 3H).
395 - Dia 2	1H NMR (500 MHz, DMSO) δ = 7.65 (s, 1H), 7.56 (d, 1H), 7.51 (s, 1H), 7.44 (d, 1H), 7.39 (d, 1H), 7.23 (s, 1H), 6.93 (dd, 1H), 5.82 (q, 1H), 5.02 (t, 1H), 4.49 (dt, 1H), 3.75 (hept, 2H), 2.62 (s, 3H), 2.57 (s, 3H), 1.65 (d, 3H).
396 - En 1	1H NMR (500 MHz, DMSO) δ = 8.99 (s, 1H), 8.04 (dd, 1H), 7.48 (dd, 1H), 7.33 (dd, 1H), 6.97 (dd, 1H), 5.33 (t, 2H), 4.99 (dp, 1H), 4.88 (t, 1H), 4.67 (t, 1H), 3.65-3.59 (m, 1H), 3.59-3.41 (m, 2H), 3.01 (d, 3H), 2.62 (d, 3H), 2.43 (d, 3H), 1.33 (t, 3H).
397 - En 1	1H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 8.04-7.92 (m, 2H), 7.47 (d, 1H), 7.34 (d, 1H), 6.96 (dd, 1H), 5.32 (s, 2H), 4.68 (t, 1H), 4.53 (p, 1H), 3.42 (q, 2H), 3.17 (q, 2H), 2.61 (s, 3H), 2.42 (s, 3H), 1.34 (d, 3H).
398	1H NMR (400 MHz, DMSO) δ = 8.99 (s, 1H), 7.89 (s, 1H), 7.66 (t, 1H), 7.48 (d, 1H), 7.35 (d, 1H), 6.98 (dd, 1H), 5.33 (s, 2H), 4.58 (t, 1H), 3.41 (q, 2H), 3.18 (q, 2H), 2.62 (s, 3H), 2.43 (s, 3H), 1.52 (s, 6H).
399	1H NMR (400 MHz, DMSO) δ = 8.34 (s, 1H), 7.69 (s, 1H), 7.57-7.10 (m, 2H), 6.94 (dd, 1H), 6.73 (t, 1H), 5.31 (s, 2H), 4.01 (s, 2H), 2.63 (s, 3H), 2.54 (s, 3H), 0.89 (d, 4H).
400	1H NMR (500 MHz, DMSO) δ = 8.99 (s, 1H), 8.34 (s, 1H), 7.45 (d, 1H), 7.23 (d, 1H), 6.94 (dd, 1H), 6.73 (t, 1H), 5.31 (s, 2H), 4.02 (s, 2H), 2.55 (s, 3H), 2.42 (s, 3H), 0.89 (dt, 4H).
401 - En 1	1H NMR (400 MHz, DMSO) δ = 12.35 (s, 1H), 8.20 (d, 1H), 7.71 (s, 1H), 7.55-7.46 (m, 3H), 7.50 (s, 1H), 7.18-7.11 (m, 2H), 6.97 (dd, 1H), 5.39 (d, 1H), 5.36 (s, 2H), 5.21 (s, 1H), 4.04-3.90 (m, 2H), 2.64 (d, 6H).
402 - En 1	1H NMR (400 MHz, DMSO) δ = 8.99 (s, 1H), 8.20 (d, 1H), 7.47 (d, 1H), 7.37 (d, 1H), 6.97 (dd, 1H), 5.32 (s, 2H), 4.72-4.62 (m, 1H), 4.59 (t, 1H), 3.59-3.43 (m, 4H), 3.47-3.38 (m, 4H), 2.61 (s, 3H), 2.43 (s, 3H), 2.41-2.31 (m, 1H), 1.98 (dq, 1H).
403 - En 1	1H NMR (400 MHz, DMSO) δ = 8.16 (d, 1H), 7.48 (d, 1H), 7.41 (d, 1H), 7.20 (d, 1H), 7.02 (dd, 1H), 6.88 (d, 1H), 5.15 (s, 2H), 4.76-4.64 (m, 2H), 3.70 (s, 3H), 3.52 (q, 2H), 3.48-3.35 (m, 3H), 3.23-3.12 (m, 1H), 2.62 (s, 3H), 2.36 (s, 1H), 2.07-1.91 (m, 1H).
404 - En 1	1H NMR (400 MHz, DMSO) δ = 8.18 (d, 1H), 7.48 (d, 1H), 7.38 (dd, 2H), 6.99 (dd, 1H), 6.39 (d, 1H), 5.20 (s, 2H), 4.73 (d, 1H), 4.73-4.63 (m, 1H), 3.85 (s, 3H), 3.53 (q, 2H), 3.49-3.37 (m, 3H), 3.18 (dt, 1H), 2.61 (s, 3H), 2.42-2.32 (m, 1H), 2.05-1.90 (m, 1H).
405	1H NMR (400 MHz, DMSO) δ = 8.18 (d, 1H), 7.72 (dd, 1H), 7.61-7.53 (m, 1H), 7.46 (d, 1H), 7.33 (d, 1H), 6.95 (dd, 1H), 6.26 (t, 1H), 4.93 (s, 2H), 4.68 (dt, 1H), 3.53 (t, 2H), 3.48 (s, 3H), 3.48-3.35 (m, 3H), 3.17 (dt, 1H), 2.61 (s, 3H), 2.42-2.32 (m, 1H), 1.98 (dq, 1H).
406 - En 1	1H NMR (400 MHz, DMSO) δ = 8.58 (ddd, 1H), 8.18 (d, 1H), 7.84 (td, 1H), 7.56 (dt, 1H), 7.47 (d, 1H), 7.40-7.31 (m, 2H), 7.00 (dd, 1H), 5.21 (s, 2H), 4.76-4.63 (m, 2H), 3.53 (q, 2H), 3.49-3.35 (m, 3H), 3.17 (dt, 1H), 2.61 (s, 3H), 2.36 (s, 1H), 2.05-1.89 (m, 1H).
407 - En 1	1H NMR (400 MHz, DMSO) δ = 8.18 (d, 1H), 7.85 (d, 1H), 7.78 (d, 1H), 7.50 (d, 1H), 7.40 (d, 1H), 7.03 (dd, 1H), 5.47 (s, 2H), 4.76-4.63 (m, 1H), 3.53 (q, 1H), 3.48-3.35 (m, 3H), 3.23-3.12 (m, 1H), 2.62 (s, 3H), 2.36 (s, 1H), 1.98 (dq, 1H).
408 - En 1	1H NMR (400 MHz, DMSO) δ = 8.19 (d, 1H), 7.57 (dd, 1H), 7.46 (d, 1H), 7.42-7.33 (m, 2H), 6.95 (dd, 1H), 6.22 (t, 1H), 4.90 (s, 2H), 4.75-4.63 (m, 1H), 3.53 (t, 2H), 3.42 (dt, 3H), 3.19 (dt, 1H), 2.61 (s, 3H), 2.42-2.32 (m, 1H), 2.07-1.92 (m, 1H).
409 - En 1	1H NMR (500 MHz, DMSO) δ = 8.22-8.13 (m, 2H), 7.83 (dd, 1H), 7.47 (d, 1H), 7.35 (d, 1H), 7.03 (dd, 1H), 6.97 (dd, 1H), 5.08 (s, 2H), 4.75-4.63 (m, 2H), 3.93 (s, 3H), 3.53 (q, 2H), 3.47-3.35 (m, 3H), 3.22-3.13 (m, 1H), 2.61 (s, 3H), 2.35 (q, 1H), 1.97 (dq, 1H).
410 - En 1	1H NMR (500 MHz, DMSO) δ = 8.98 (s, 1H), 7.92 (s, 1H), 7.47 (d, 1H), 7.22 (d, 1H), 6.96 (dd, 1H), 5.32 (s, 2H), 4.68 (s, 1H), 3.53 (t, 2H), 3.49-3.35 (m, 3H), 3.22-3.13 (m, 1H), 2.58 (s, 3H), 2.43 (s, 3H), 2.03 (ddd, 1H), 1.35 (s, 3H).
411 - En 1	1H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 7.89 (d, 1H), 7.48 (d, 1H), 7.34 (d, 1H), 6.97 (dd, 1H), 5.32 (s, 2H), 5.04 (dt, 1H), 4.99 (s, 1H), 3.68 (dd, 1H), 3.13 (s, 3H), 2.89 (s, 3H), 2.61 (s, 3H), 2.42 (s, 3H).
412 - En 1	1H NMR (400 MHz, DMSO) δ = 7.70 (d, 1H), 7.51 (d, 1H), 7.46 (d, 1H), 7.33 (d, 1H), 6.94 (dd, 1H), 5.31 (s, 2H), 4.79 (s, 1H), 4.10 (q, 1H), 3.55-3.48 (m, 2H), 2.61 (d, 6H), 2.50-2.42 (m, 1H), 2.39 (dd, 1H), 2.20 (s, 6H).
413 - En 1	1H NMR (500 MHz, DMSO) δ = 8.99 (s, 1H), 7.52 (d, 1H), 7.46 (d, 1H), 7.34 (d, 1H), 6.95 (dd, 1H), 5.31 (d, 2H), 4.78 (s, 1H), 4.10 (d, 0H), 3.52 (tt, 2H), 2.59 (s, 3H), 2.48-2.43 (m, 1H), 2.42 (s, 3H), 2.39 (dd, 1H), 2.20 (s, 6H).
414	1H NMR (400 MHz, DMSO) δ = 7.74 (s, 1H), 7.69 (s, 1H), 7.45 (d, 1H), 7.27 (d, 1H), 6.94 (dd, 1H), 5.31 (s, 2H), 4.91 (t, 1H), 3.66 (d, 2H), 2.63 (s, 3H), 2.58 (s, 3H), 2.33 (q, 1H), 2.18-2.05 (m, 3H), 1.86 (dd, 1H), 1.81-1.69 (m, 1H).

-continued

cpd Nr	NMR Data
415	¹ H NMR (400 MHz, DMSO) δ = 8.58 (ddd, 1H), 7.84 (td, 1H), 7.74 (s, 1H), 7.55 (dt, 1H), 7.45 (d, 1H), 7.35 (ddd, 1H), 7.28 (d, 1H), 6.98 (dd, 1H), 5.20 (s, 2H), 4.89 (d, 1H), 3.65 (d, 2H), 2.58 (s, 3H), 2.31 (q, 2H), 2.19-2.07 (m, 1H), 1.92-1.69 (m, 2H).
416	¹ H NMR (400 MHz, DMSO) δ = 11.76 (s, 1H), 7.76 (s, 1H), 7.61-7.54 (m, 1H), 7.44 (d, 1H), 7.38 (dd, 1H), 7.24 (d, 1H), 6.93 (dd, 1H), 6.22 (t, 1H), 4.90 (t, 1H), 4.89 (s, 2H), 3.66 (d, 2H), 2.58 (s, 3H), 2.38-2.25 (m, 2H), 2.13 (s, 2H), 1.94-1.66 (m, 2H).
417	¹ H NMR (400 MHz, DMSO) δ = 8.16 (dd, 1H), 7.82 (dd, 1H), 7.76 (s, 1H), 7.45 (d, 1H), 7.26 (d, 1H), 7.03 (dd, 1H), 6.95 (dd, 1H), 5.07 (s, 2H), 4.89 (t, 1H), 3.93 (s, 3H), 3.65 (d, 2H), 2.58 (s, 3H), 2.37-2.23 (m, 2H), 2.13 (ddd, 1H), 1.93-1.66 (m, 2H).
418	¹ H NMR (400 MHz, DMSO) δ = 8.14 (s, 1H), 7.69 (s, 1H), 7.44 (d, 1H), 7.26 (d, 1H), 6.93 (dd, 1H), 5.32 (s, 2H), 4.77 (t, 1H), 3.56 (d, 2H), 2.64 (s, 3H), 2.56 (s, 3H), 0.83 (s, 1H), 0.82-0.71 (m, 3H).
419	¹ H NMR (400 MHz, DMSO) δ = 8.58 (ddd, 1H), 8.14 (s, 1H), 7.84 (td, 1H), 7.55 (d, 1H), 7.44 (d, 1H), 7.35 (ddd, 1H), 7.26 (d, 1H), 6.97 (dd, 1H), 5.21 (s, 2H), 4.76 (t, 1H), 3.55 (d, 2H), 2.56 (s, 3H), 0.83-0.70 (m, 4H).
420	¹ H NMR (400 MHz, DMSO) δ = 11.76 (s, 1H), 8.16 (s, 1H), 7.57 (ddd, 1H), 7.43 (d, 1H), 7.38 (dd, 1H), 7.24 (d, 1H), 6.93 (dd, 1H), 6.23 (t, 1H), 4.90 (s, 2H), 4.76 (s, 1H), 3.56 (s, 2H), 2.56 (s, 3H), 0.76 (dt, 4H).
421	¹ H NMR (400 MHz, DMSO) δ = 8.20-8.12 (m, 2H), 7.83 (dd, 1H), 7.44 (d, 1H), 7.25 (d, 1H), 7.03 (dd, 1H), 6.95 (dd, 1H), 5.07 (s, 2H), 4.76 (t, 1H), 3.93 (s, 3H), 3.55 (d, 2H), 2.56 (s, 3H), 0.76 (dt, 4H).
422	¹ H NMR (400 MHz, DMSO) δ = 7.48 (d, 1H), 7.37 (d, 1H), 7.19 (d, 1H), 7.04-6.96 (m, 2H), 6.88 (d, 1H), 5.18-5.08 (m, 4H), 3.68 (d, 5H), 3.56 (dd, 2H), 2.60 (s, 3H), 1.33 (s, 3H).
423	¹ H NMR (400 MHz, DMSO) δ = 7.48 (d, 1H), 7.37 (d, 1H), 7.33 (d, 1H), 7.05 (s, 1H), 6.98 (dd, 1H), 6.39 (d, 1H), 5.18 (s, 2H), 4.98 (t, 2H), 3.85 (s, 3H), 3.65 (dd, 2H), 3.55 (dd, 2H), 2.60 (s, 3H), 1.32 (s, 3H).
424	¹ H NMR (400 MHz, DMSO) δ = 8.74 (d, 1H), 8.27 (d, 1H), 7.79 (dd, 1H), 7.49 (d, 1H), 7.31 (d, 1H), 7.07 (s, 1H), 6.98 (dd, 1H), 5.31 (s, 2H), 4.94 (t, 2H), 3.64 (dd, 2H), 3.55 (dd, 2H), 2.60 (s, 3H), 1.31 (s, 3H).
425	¹ H NMR (400 MHz, DMSO) δ = 7.70 (s, 1H), 7.47 (d, 1H), 7.29 (d, 1H), 7.03 (s, 1H), 6.95 (dd, 1H), 5.31 (s, 2H), 4.98 (t, 2H), 3.65 (dd, 2H), 3.55 (dd, 2H), 2.61 (d, 6H), 1.32 (s, 3H).
426	¹ H NMR (400 MHz, DMSO) δ = 8.99 (s, 1H), 7.47 (d, 1H), 7.30 (d, 1H), 7.05 (s, 1H), 6.96 (dd, 1H), 5.30 (s, 2H), 4.96 (t, 2H), 3.65 (dd, 2H), 3.55 (dd, 2H), 2.60 (s, 3H), 2.42 (s, 3H), 1.32 (s, 3H).
427	¹ H NMR (600 MHz, DMSO) δ = 9.22 (dd, 1H), 7.86 (dd, 1H), 7.76 (dd, 1H), 7.49 (d, 1H), 7.35 (d, 1H), 7.06-7.00 (m, 2H), 5.43 (s, 2H), 4.96 (tq, 2H), 3.66 (dd, 2H), 3.56 (dd, 2H), 2.60 (s, 3H), 1.32 (s, 3H).
428	¹ H NMR (500 MHz, DMSO) δ = 8.58 (ddd, 1H), 7.84 (td, 1H), 7.59-7.53 (m, 1H), 7.47 (d, 1H), 7.35 (ddd, 1H), 7.30 (d, 1H), 7.03 (s, 1H), 6.99 (dd, 1H), 5.19 (s, 2H), 4.96 (t, 2H), 3.64 (dd, 2H), 3.55 (dd, 2H), 2.59 (s, 3H), 1.31 (s, 3H).
429	¹ H NMR (600 MHz, DMSO) δ = 9.18 (d, 1H), 8.86-8.79 (m, 1H), 7.67 (dd, 1H), 7.49 (d, 1H), 7.30 (d, 1H), 7.05-7.00 (m, 2H), 5.24 (s, 2H), 4.98-4.92 (m, 2H), 3.64 (dd, 2H), 3.55 (dd, 2H), 2.60 (s, 3H), 1.31 (s, 3H).
430	¹ H NMR (500 MHz, DMSO) δ = 11.77 (s, 1H), 7.58 (ddd, 1H), 7.46 (d, 1H), 7.41-7.35 (m, 1H), 7.26 (d, 1H), 7.04 (s, 1H), 6.94 (dd, 1H), 6.22 (t, 1H), 4.95 (t, 2H), 4.88 (s, 2H), 3.64 (dd, 2H), 3.55 (dd, 2H), 2.59 (s, 3H), 1.31 (s, 3H).
431	¹ H NMR (500 MHz, DMSO) δ = 8.16 (dd, 1H), 7.83 (dd, 1H), 7.46 (d, 1H), 7.29 (d, 1H), 7.07-6.99 (m, 2H), 6.96 (dd, 1H), 5.06 (s, 2H), 4.94 (t, 2H), 3.93 (s, 3H), 3.64 (dd, 2H), 3.55 (dd, 2H), 2.60 (s, 3H), 1.31 (s, 3H).
432 - En 1	¹ H NMR (500 MHz, DMSO) δ = 8.46 (s, 1H), 8.03 (d, 1H), 7.98 (s, 1H), 7.72 (d, 2H), 7.46 (d, 1H), 7.37 (s, 1H), 7.29 (d, 1H), 6.95 (dd, 1H), 5.32 (s, 2H), 4.94 (td, 1H), 4.65 (dd, 1H), 4.60 (dd, 1H), 2.64 (s, 3H), 2.52 (s, 3H).
433 - En 1	¹ H NMR (400 MHz, DMSO) δ = 7.88 (d, 1H), 7.48 (d, 2H), 7.38 (dd, 2H), 7.11 (s, 1H), 6.98 (dd, 1H), 6.39 (d, 1H), 5.20 (s, 2H), 4.48 (p, 1H), 3.85 (s, 3H), 2.62 (s, 3H), 1.36 (d, 3H).
434 - En 1	¹ H NMR (400 MHz, DMSO) δ = 7.87 (d, 1H), 7.71 (s, 1H), 7.52-7.43 (m, 2H), 7.36 (d, 1H), 7.11 (s, 1H), 6.95 (dd, 1H), 5.33 (s, 2H), 4.48 (p, 1H), 2.62 (d, 6H), 1.36 (d, 3H).
435 - En 1	¹ H NMR (400 MHz, DMSO) δ = 8.58 (ddd, 1H), 7.91-7.79 (m, 2H), 7.56 (dt, 1H), 7.51-7.44 (m, 2H), 7.40-7.31 (m, 2H), 7.10 (s, 1H), 7.00 (dd, 1H), 5.21 (s, 2H), 4.47 (p, 1H), 2.62 (s, 3H), 1.36 (d, 3H).
436 - En 1	¹ H NMR (400 MHz, DMSO) δ = 11.74 (s, 1H), 7.89 (d, 1H), 7.58 (d, 1H), 7.52-7.42 (m, 2H), 7.36 (dd, 2H), 7.07 (s, 1H), 6.95 (dd, 1H), 6.22 (t, 1H), 4.90 (s, 2H), 4.46 (q, 1H), 2.62 (s, 3H), 1.36 (d, 3H).
437 - En 1	¹ H NMR (400 MHz, DMSO) δ = 8.16 (dd, 1H), 7.83 (dd, 1H), 7.47 (s, 1H), 7.47 (d, 1H), 7.35 (d, 1H), 7.08 (s, 1H), 7.03 (dd, 1H), 6.97 (dd, 1H), 5.08 (s, 2H), 4.47 (p, 1H), 3.93 (s, 3H), 2.62 (s, 3H), 1.35 (d, 3H).
438	¹ H NMR (400 MHz, DMSO) δ = 7.83 (s, 1H), 7.49 (d, 1H), 7.44 (d, 1H), 7.37 (d, 2H), 7.19 (s, 1H), 7.00 (dd, 1H), 6.39 (d, 1H), 5.21 (s, 2H), 3.86 (s, 3H), 2.64 (s, 3H), 1.57 (s, 6H).
439	¹ H NMR (400 MHz, DMSO) δ = 7.81 (s, 1H), 7.48 (d, 1H), 7.39 (d, 1H), 7.37 (s, 1H), 7.19 (s, 1H), 6.97 (dd, 1H), 5.34 (s, 2H), 2.63 (d, 6H), 1.56 (s, 6H).
440	¹ H NMR (400 MHz, DMSO) δ = 8.99 (s, 1H), 7.82 (s, 1H), 7.48 (d, 1H), 7.40 (d, 1H), 7.16 (s, 1H), 6.98 (dd, 1H), 5.33 (s, 2H), 2.64 (s, 3H), 2.42 (s, 3H), 1.56 (s, 6H).
441	¹ H NMR (600 MHz, DMSO) δ = 7.86 (d, 2H), 7.50 (d, 1H), 7.47-7.42 (m, 1H), 7.11 (s, 1H), 5.45 (s, 2H), 2.64 (s, 3H), 1.56 (s, 6H).
442	¹ H NMR (400 MHz, DMSO) δ = 8.58 (ddd, 1H), 7.89-7.79 (m, 2H), 7.57 (dt, 1H), 7.49 (d, 1H), 7.41 (d, 1H), 7.39-7.31 (m, 1H), 7.34 (s, 1H), 7.16 (s, 1H), 7.01 (dd, 1H), 5.21 (s, 2H), 2.63 (s, 3H), 1.55 (s, 6H).
443	¹ H NMR (400 MHz, DMSO) δ = 11.75 (s, 1H), 7.82 (s, 1H), 7.63-7.55 (m, 1H), 7.47 (d, 1H), 7.42-7.31 (m, 3H), 7.12 (s, 1H), 6.95 (dd, 1H), 6.22 (t, 1H), 4.90 (s, 2H), 2.63 (s, 3H), 1.55 (s, 6H).

-continued

cpd Nr	NMR Data
444	1H NMR (400 MHz, DMSO) δ = 8.16 (dd, 1H), 7.87-7.80 (m, 1H), 7.82 (s, 1H), 7.48 (d, 1H), 7.39 (d, 1H), 7.34 (s, 1H), 7.14 (s, 1H), 7.00 (ddd, 2H), 5.08 (s, 2H), 3.93 (s, 3H), 2.64 (s, 3H), 1.55 (s, 6H).
445 - En 1	1H NMR (500 MHz, DMSO) δ = 7.69 (d, 2H), 7.51-7.44 (m, 2H), 7.32 (d, 1H), 7.13-7.09 (m, 1H), 6.97 (dd, 1H), 5.31 (s, 2H), 4.56 (td, 1H), 2.88 (dd, 1H), 2.82 (dd, 1H), 2.62 (d, 6H), 2.34 (s, 3H), 1.75 (ddd, 1H), 0.48-0.37 (m, 3H), 0.32 (dt, 1H).
446 - En 1	1H NMR (500 MHz, DMSO) δ = 8.99 (s, 1H), 7.70 (d, 1H), 7.51-7.45 (m, 2H), 7.33 (d, 1H), 7.10 (s, 1H), 6.98 (dd, 1H), 5.31 (d, 2H), 4.55 (td, 1H), 2.88 (dd, 1H), 2.81 (dd, 1H), 2.60 (s, 3H), 2.42 (s, 3H), 2.33 (s, 3H), 1.75 (dq, 1H), 0.45-0.37 (m, 3H), 0.40-0.29 (m, 1H).
447 - En 1	1H NMR (500 MHz, DMSO) δ = 7.97 (d, 1H), 7.79-7.25 (m, 4H), 7.08-6.47 (m, 2H), 5.32 (s, 2H), 4.87-4.64 (m, 1H), 4.19 (qd, 2H), 3.33 (s, 2H), 2.63 (d, 5H).
448 - En 1	1H NMR (500 MHz, DMSO) δ = 8.98 (s, 1H), 7.98 (d, 1H), 7.66 (s, 1H), 7.58-7.26 (m, 2H), 7.37 (s, 1H), 6.98 (dd, 1H), 6.72 (t, 1H), 5.31 (s, 2H), 4.75 (td, 1H), 4.19 (qd, 2H), 2.62 (s, 3H), 2.42 (s, 3H).
449 - En 1	1H NMR (500 MHz, DMSO) δ = 7.77-7.69 (m, 2H), 7.59 (s, 1H), 7.47 (d, 1H), 7.41 (d, 1H), 7.14 (s, 1H), 6.95 (dd, 1H), 5.32 (s, 2H), 4.52 (q, 1H), 2.63 (d, 7H), 2.55 (dd, 1H), 2.24 (s, 6H).
450 - En 1	1H NMR (500 MHz, DMSO) δ = 8.99 (s, 1H), 7.76 (d, 1H), 7.58 (s, 1H), 7.48 (d, 1H), 7.15-7.11 (m, 1H), 6.97 (dd, 1H), 5.32 (d, 2H), 4.51 (td, 1H), 2.66 (dd, 1H), 2.62 (s, 3H), 2.55 (dd, 1H), 2.42 (s, 3H), 2.24 (s, 6H).
451 - En 1	1H NMR (400 MHz, DMSO) δ = 7.63 (d, 1H), 7.53-7.43 (m, 3H), 7.19 (t, 2H), 7.02 (dd, 1H), 6.87 (d, 1H), 5.22 (s, 1H), 5.16 (s, 2H), 4.48 (dt, 1H), 3.77 (q, 2H), 3.70 (s, 3H), 2.64 (s, 3H).
452 - En 1	1H NMR (400 MHz, DMSO) δ = 7.64 (d, 1H), 7.49 (s, 1H), 7.53-7.42 (m, 2H), 7.37 (d, 1H), 7.20 (s, 1H), 7.00 (dd, 1H), 6.40 (d, 1H), 5.20 (s, 2H), 5.07 (s, 1H), 4.53-4.44 (m, 1H), 3.85 (s, 3H), 3.81-3.69 (m, 2H), 2.64 (s, 3H).
453 - En 1	1H NMR (400 MHz, DMSO) δ = 7.72 (s, 1H), 7.62 (d, 1H), 7.52-7.45 (m, 2H), 7.41 (d, 1H), 7.20 (s, 1H), 6.97 (dd, 1H), 5.33 (s, 2H), 5.04 (s, 1H), 4.48 (dt, 1H), 3.78-3.72 (m, 2H), 2.63 (m, 6H).
454	1H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 7.63 (d, 1H), 7.49 (d, 2H), 7.42 (d, 1H), 7.19 (s, 1H), 6.98 (dd, 1H), 5.32 (s, 2H), 5.02 (s, 1H), 4.48 (dt, 1H), 3.75 (s, 2H), 2.64 (s, 3H), 2.42 (s, 3H).
455 - En 1	1H NMR (400 MHz, DMSO) δ = 8.58 (ddd, 1H), 7.84 (td, 1H), 7.62 (d, 1H), 7.57 (dt, 1H), 7.49 (d, 2H), 7.43 (d, 1H), 7.35 (ddd, 1H), 7.19 (s, 1H), 7.01 (dd, 1H), 5.20 (s, 2H), 5.02 (t, 1H), 4.48 (dt, 1H), 3.75 (td, 2H), 2.64 (s, 3H).
456 - En 1	1H NMR (500 MHz, DMSO) δ = 7.65 (d, 1H), 7.56-7.47 (m, 3H), 7.45 (d, 1H), 7.20 (s, 1H), 6.99 (dd, 1H), 6.55-6.23 (m, 2H), 5.24 (s, 2H), 5.05 (s, 1H), 4.67 (td, 2H), 4.49 (dt, 1H), 3.75 (dd, 2H), 2.64 (s, 3H).
457 - En 1	1H NMR (500 MHz, DMSO) δ = 8.67 (dd, 1H), 8.24-8.03 (m, 1H), 7.75-7.34 (m, 5H), 7.44-6.89 (m, 3H), 5.35 (s, 2H), 5.01 (s, 1H), 4.48 (dt, 1H), 3.74 (d, 2H), 2.65 (s, 3H).
458 - En 1	1H NMR (500 MHz, DMSO) δ = 7.62 (d, 1H), 7.48 (d, 2H), 7.39 (d, 1H), 7.18 (s, 1H), 6.96 (dd, 1H), 5.23 (s, 2H), 5.01 (t, 1H), 4.48 (dt, 1H), 3.74 (td, 2H), 2.64 (s, 3H), 2.57 (s, 3H), 2.32 (s, 3H).
459 - En 1	1H NMR (400 MHz, DMSO) δ = 11.75 (s, 1H), 7.66-7.56 (m, 2H), 7.51-7.44 (m, 2H), 7.38 (dd, 2H), 7.17 (s, 1H), 6.96 (dd, 1H), 6.22 (t, 1H), 5.04 (s, 1H), 4.90 (s, 2H), 4.47 (dt, 1H), 3.78-3.72 (m, 2H), 2.64 (s, 3H).
460 - En 1	1H NMR (500 MHz, DMSO) δ = 7.65 (d, 1H), 7.51 (d, 1H), 7.46 (dd, 3H), 7.21-7.16 (m, 1H), 7.01 (dd, 1H), 5.18 (s, 2H), 5.01 (t, 1H), 4.48 (dt, 1H), 3.83 (s, 3H), 3.75 (td, 2H), 2.64 (s, 3H).
461 - En 1	1H NMR (400 MHz, DMSO) δ = 8.41 (dd, 1H), 7.83 (dd, 1H), 7.65 (s, 1H), 7.54-7.47 (m, 2H), 7.45 (s, 1H), 7.30 (s, 1H), 7.24 (dd, 1H), 7.02 (dd, 1H), 5.17 (s, 2H), 3.97 (d, 1H), 3.77 (d, 1H), 2.66 (s, 3H), 2.54 (s, 3H), 1.53 (s, 3H).
462 - En 1	1H NMR (500 MHz, DMSO) δ = 7.72 (s, 1H), 7.63 (s, 1H), 7.52-7.44 (m, 3H), 7.31 (s, 1H), 6.97 (dd, 1H), 5.33 (s, 2H), 5.14 (s, 1H), 3.97 (d, 1H), 3.77 (d, 1H), 2.64 (d, 6H), 1.53 (s, 3H).
463 - En 1	1H NMR (400 MHz, DMSO) δ = 8.99 (s, 1H), 7.63 (s, 1H), 7.53-7.40 (m, 2H), 7.44 (s, 1H), 7.28 (s, 1H), 6.98 (dd, 1H), 5.32 (s, 2H), 5.16 (s, 1H), 3.96 (d, 1H), 3.76 (d, 1H), 2.65 (s, 3H), 2.42 (s, 3H), 1.53 (s, 3H).
464 - En 1	1H NMR (400 MHz, DMSO) δ = 8.55 (d, 1H), 7.78 (dd, 1H), 7.64 (s, 1H), 7.52-7.45 (m, 2H), 7.45 (s, 1H), 7.33-7.24 (m, 2H), 6.98 (dd, 1H), 5.13 (s, 2H), 3.97 (d, 1H), 3.77 (d, 1H), 2.65 (s, 3H), 2.47 (s, 3H), 1.53 (s, 3H).
465 - En 1	1H NMR (500 MHz, DMSO) δ = 7.74-7.38 (m, 5H), 7.30 (s, 1H), 7.00 (dd, 1H), 6.59-6.16 (m, 2H), 5.25 (s, 2H), 5.15 (s, 1H), 4.68 (td, 2H), 3.97 (d, 1H), 3.77 (d, 1H), 2.66 (s, 3H), 1.53 (s, 3H).
466 - En 1	1H NMR (500 MHz, DMSO) δ = 7.64 (s, 1H), 7.54-7.45 (m, 3H), 7.43 (s, 1H), 7.28 (s, 1H), 7.02 (dd, 1H), 5.18 (s, 2H), 5.12 (s, 1H), 3.96 (d, 1H), 3.83 (s, 3H), 3.77 (d, 1H), 2.66 (s, 3H), 1.53 (s, 3H).
467 - En 1	1H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 7.81 (d, 1H), 7.57-7.52 (m, 1H), 7.48 (d, 1H), 7.35 (d, 1H), 7.25-7.20 (m, 1H), 5.32 (s, 2H), 4.66 (ddd, 1H), 3.73-3.60 (m, 2H), 3.30 (s, 3H), 2.62 (s, 3H), 2.42 (s, 3H).
468 - En 1	1H NMR (400 MHz, DMSO) δ = 8.99 (s, 1H), 7.71 (s, 1H), 7.50 (d, 2H), 7.43 (d, 1H), 7.37 (s, 1H), 6.99 (dd, 1H), 5.33 (s, 2H), 4.00 (d, 1H), 3.75 (d, 1H), 3.26 (s, 3H), 2.65 (s, 3H), 2.42 (s, 3H), 1.55 (s, 3H).
469 - En 1	1H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 7.93 (d, 1H), 7.51-7.42 (m, 2H), 7.39 (d, 1H), 7.08 (s, 1H), 6.96 (dd, 1H), 5.31 (s, 2H), 4.75 (t, 1H), 4.51 (td, 1H), 3.56 (dt, 2H), 2.63 (s, 3H), 2.42 (s, 3H), 2.03-1.81 (m, 2H).
470	1H NMR (400 MHz, DMSO) δ = 8.99 (s, 1H), 8.62 (s, 1H), 7.49 (d, 1H), 7.38 (d, 1H), 7.27 (s, 1H), 7.20 (s, 1H), 6.99 (dd, 1H), 5.32 (s, 2H), 2.99-2.87 (m, 3H), 2.63 (s, 3H), 2.42 (s, 3H).
471	1H NMR (400 MHz, DMSO) δ = 8.24 (s, 1H), 7.70 (s, 1H), 7.47 (d, 1H), 7.36 (d, 1H), 7.02-6.93 (m, 2H), 6.88 (s, 1H), 5.34 (s, 2H), 2.63 (s, 6H), 2.59-2.53 (m, 2H), 2.26 (q, 2H), 2.00-1.80 (m, 2H).
472	1H NMR (400 MHz, DMSO) δ = 8.58 (d, 1H), 8.25 (s, 1H), 7.84 (t, 1H), 7.56 (d, 1H), 7.48 (d, 1H), 7.40-7.31 (m, 2H), 7.05-6.95 (m, 2H), 6.88 (s, 1H), 5.23 (s, 2H), 2.62 (s, 3H), 2.24 (q, 2H), 1.93 (s, 2H), 1.88 (dd, 2H).
473	1H NMR (500 MHz, DMSO-d ₆): δ 8.16 (dd, 1H), 7.82 (dd, 1H), 7.46 (d, 1H) 7.29-7.26 (m, 2H), 7.06-7.00 (m, 1H), 6.96 (dd, 1H), 5.06 (s, 2H), 5.04 (t, 1H), 3.92 (s, 3H), 3.50 (d, 2H), 3.58 (s, 3H), 1.33 (s, 6H).

-continued

cpd Nr	NMR Data
474	¹ H NMR (500 MHz, DMSO-d ₆): δ 8.98 (s, 1H), 7.58 (d, 1H), 7.46 (d, 1H), 7.33 (d, 1H), 6.97 (dd, 1H), 5.31 (s, 2H), 4.78 (s, 1H), 4.14-4.18 (m, 1H), 3.50-3.58 (m, 6H), 2.60 (s, 3H), 2.40-2.50 (m, 4H), 2.34-2.37 (m, 5H).
475	¹ H NMR (400 MHz, DMSO) δ = 8.78-8.70 (m, 1H), 8.27 (d, 1H), 7.79 (dd, 1H), 7.48 (d, 1H), 7.36 (d, 1H), 7.20 (s, 1H), 7.05 (s, 1H), 6.98 (dd, 1H), 5.33 (s, 2H), 2.61 (s, 3H), 1.35 (q, 2H), 1.01 (q, 2H).
476	¹ H NMR (400 MHz, DMSO) δ = 8.29 (s, 1H), 7.70 (s, 1H), 7.45 (d, 1H), 7.35 (d, 1H), 7.19 (s, 1H), 7.05 (s, 1H), 6.94 (dd, 1H), 5.33 (s, 2H), 2.62 (d, 6H), 1.35 (q, 2H), 1.03 (q, 2H).
477	¹ H NMR (400 MHz, DMSO) δ = 8.99 (s, 1H), 8.29 (s, 1H), 7.45 (d, 1H), 7.36 (d, 1H), 7.05 (s, 1H), 6.95 (dd, 1H), 5.33 (s, 2H), 2.60 (s, 3H), 2.42 (s, 3H), 1.35 (q, 2H), 1.04 (q, 2H).
478	¹ H NMR (400 MHz, DMSO) δ = 8.58 (ddd, 1H), 8.30 (s, 1H), 7.84 (td, 1H), 7.55 (d, 1H), 7.45 (d, 1H), 7.39-7.31 (m, 2H), 7.18 (s, 1H), 7.05 (s, 1H), 6.98 (dd, 1H), 5.23 (s, 2H), 2.60 (s, 3H), 1.35 (q, 2H), 1.02 (q, 2H).
479	¹ H NMR (400 MHz, DMSO) δ = 11.76 (s, 1H), 8.32 (s, 1H), 7.62-7.54 (m, 1H), 7.44 (d, 1H), 7.41-7.30 (m, 2H), 7.20 (s, 1H), 7.04 (s, 1H), 6.94 (dd, 1H), 6.23 (t, 1H), 4.92 (s, 2H), 2.60 (s, 3H), 1.34 (q, 2H), 1.03 (g, 2H).
480	¹ H NMR (400 MHz, DMSO) δ = 8.32 (s, 1H), 8.16 (dd, 1H), 7.83 (dd, 1H), 7.45 (d, 1H), 7.35 (d, 1H), 7.20 (s, 1H), 7.03 (dd, 2H), 6.96 (dd, 1H), 5.09 (s, 2H), 3.93 (s, 3H), 2.60 (s, 3H), 1.34 (q, 2H), 1.01 (g, 2H).
481	¹ H NMR (400 MHz, DMSO) δ = 7.46 (d, 1H), 7.34 (d, 1H), 7.19 (d, 1H), 7.15 (s, 1H), 6.99 (dd, 1H), 6.87 (d, 1H), 5.25 (t, 1H), 5.16 (s, 2H), 3.69 (s, 3H), 3.52 (d, 2H), 2.58 (s, 3H), 1.36 (s, 6H).
482	¹ H NMR (500 MHz, DMSO) δ = 7.46 (d, 1H), 7.37 (d, 1H), 7.29 (d, 1H), 7.20 (s, 1H), 6.97 (dd, 1H), 6.38 (d, 1H), 5.18 (s, 2H), 5.07 (t, 1H), 3.85 (s, 3H), 3.51 (d, 2H), 2.58 (s, 3H), 1.35 (s, 6H).
483	¹ H NMR (400 MHz, DMSO) δ = 7.70 (s, 1H), 7.46 (d, 1H), 7.25 (d, 1H), 7.19 (s, 1H), 6.95 (dd, 1H), 5.32 (s, 2H), 5.08 (t, 1H), 3.51 (d, 2H), 2.64 (s, 3H), 2.58 (s, 3H), 1.35 (s, 6H).
484	¹ H NMR (500 MHz, DMSO) δ = 8.58 (dt, 1H), 7.84 (td, 1H), 7.55 (d, 1H), 7.46 (d, 1H), 7.35 (ddd, 1H), 7.25 (d, 1H), 7.18 (s, 1H), 6.98 (dd, 1H), 5.20 (s, 2H), 5.05 (t, 1H), 3.50 (d, 2H), 2.57 (s, 3H), 1.34 (s, 6H).
485	¹ H NMR (400 MHz, DMSO) δ = 11.76 (s, 1H), 7.61-7.54 (m, 1H), 7.45 (d, 1H), 7.41-7.34 (m, 1H), 7.25-7.17 (m, 2H), 6.94 (dd, 1H), 6.22 (t, 1H), 5.03 (t, 1H), 4.89 (s, 2H), 3.50 (d, 2H), 2.57 (s, 3H), 1.34 (s, 6H).
486	¹ H NMR (400 MHz, DMSO) δ = 8.16 (dd, 1H), 7.82 (dd, 1H), 7.45 (d, 1H), 7.27-7.17 (m, 2H), 7.03 (dd, 1H), 6.95 (dd, 1H), 5.10-4.99 (m, 3H), 3.93 (s, 3H), 3.50 (d, 2H), 2.57 (s, 3H), 1.33 (s, 6H).
487 -	¹ H NMR (400 MHz, DMSO) δ = 7.69 (s, 1H), 7.57 (d, 1H), 7.46 (d, 1H), 7.32 (d, 1H), 6.97 (dd, 1H),
En 1	5.31 (s, 2H), 4.80 (s, 1H), 4.19-4.13 (m, 1H), 3.55 (t, 7H), 2.62 (d, 7H), 2.40 (s, 3H).
488 -	¹ H NMR (500 MHz, DMSO) δ = 8.98 (s, 1H), 7.58 (d, 1H), 7.47 (d, 1H), 7.34 (d, 1H), 6.98 (dd, 1H),
En 1	5.31 (s, 2H), 4.79 (s, 1H), 4.16 (s, 1H), 3.64-3.46 (m, 7H), 2.60 (s, 3H), 2.48 (d, 1H), 2.42 (s, 3H).
489 -	¹ H NMR (400 MHz, DMSO) δ = 8.73 (dd, 1H), 8.29-8.22 (m, 1H), 7.79 (dd, 1H), 7.67 (d, 1H), 7.48
En 1	(d, 1H), 7.27 (d, 1H), 6.98 (dd, 1H), 5.32 (d, 2H), 4.76 (t, 1H), 4.08-3.97 (m, 1H), 3.48 (dt, 1H), 3.43-3.33 (m, 1H), 2.58 (s, 3H), 1.14 (d, 3H).
490 -	¹ H NMR (400 MHz, DMSO) δ = 7.69 (s, 1H), 7.63 (d, 1H), 7.45 (d, 1H), 7.26 (d, 1H), 6.95 (dd, 1H),
En 1	5.32 (s, 2H), 4.80 (s, 1H), 4.04 (dt, 1H), 3.49 (dd, 1H), 3.39 (dd, 1H), 2.60 (d, 6H), 1.15 (d, 3H).
491 -	¹ H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 7.65 (d, 1H), 7.46 (d, 1H), 7.27 (d, 1H), 6.95 (dd, 1H),
En 1	5.32 (s, 2H), 4.78 (t, 1H), 4.11-3.98 (m, 1H), 3.49 (dt, 1H), 3.44-3.35 (m, 1H), 2.58 (s, 3H), 2.42 (s, 3H), 1.16 (d, 3H).
492	¹ H NMR (400 MHz, DMSO) δ = 8.58 (ddd, 1H), 7.84 (td, 1H), 7.63 (d, 1H), 7.55 (d, 1H), 7.46 (d, 1H), 7.35 (ddd, 1H), 7.26 (d, 1H), 6.99 (dd, 1H), 5.20 (s, 2H), 4.78 (t, 1H), 4.09-3.97 (m, 1H), 3.48 (dt, 1H), 3.43-3.34 (m, 1H), 2.57 (s, 3H), 1.15 (d, 3H).
493	¹ H NMR (400 MHz, DMSO) δ = 8.58 (ddd, 1H), 7.84 (td, 1H), 7.63 (d, 1H), 7.55 (d, 1H), 7.46 (d, 1H), 7.35 (ddd, 1H), 7.27 (d, 1H), 6.99 (dd, 1H), 5.20 (s, 2H), 4.78 (t, 1H), 4.03 (dt, 1H), 3.54-3.43 (m, 1H), 3.38 (dt, 1H), 2.57 (s, 3H), 1.15 (d, 3H).
494	¹ H NMR (400 MHz, DMSO) δ = 11.76 (s, 1H), 7.64 (d, 1H), 7.57 (ddd, 1H), 7.45 (d, 1H), 7.38 (dd, 1H), 7.25 (d, 1H), 6.94 (dd, 1H), 6.22 (t, 1H), 4.90 (s, 2H), 4.79 (s, 1H), 4.03 (p, 1H), 3.49 (dd, 1H), 3.39 (dd, 1H), 2.58 (s, 3H), 1.15 (d, 3H).
495	¹ H NMR (400 MHz, DMSO) δ = 11.76 (s, 1H), 7.64 (d, 1H), 7.61-7.53 (m, 1H), 7.45 (d, 1H), 7.38 (dd, 1H), 7.25 (d, 1H), 6.94 (dd, 1H), 6.22 (t, 1H), 4.90 (s, 2H), 4.79 (s, 1H), 4.03 (p, 1H), 3.49 (dd, 1H), 3.39 (dd, 1H), 2.58 (s, 3H), 1.15 (d, 3H).
496	¹ H NMR (400 MHz, DMSO) δ = 8.16 (dd, 1H), 7.82 (dd, 1H), 7.66 (d, 1H), 7.45 (d, 1H), 7.25 (d, 1H), 7.03 (dd, 1H), 6.96 (dd, 1H), 5.07 (s, 2H), 4.77 (s, 1H), 4.03 (p, 1H), 3.93 (s, 3H), 3.48 (t, 1H), 3.40 (s, 1H), 2.58 (s, 3H), 1.15 (d, 3H).
497	¹ H NMR (400 MHz, DMSO) δ = 8.16 (dd, 1H), 7.82 (dd, 1H), 7.65 (d, 1H), 7.45 (d, 1H), 7.25 (d, 1H), 7.03 (dd, 1H), 6.96 (dd, 1H), 5.07 (s, 2H), 4.77 (t, 1H), 4.03 (p, 1H), 3.93 (s, 3H), 3.48 (dt, 1H), 3.43-3.34 (m, 1H), 2.58 (s, 3H), 1.15 (d, 3H).
498	¹ H NMR (600 MHz, DMSO) δ = 8.98 (s, 1H), 8.25 (t, 1H), 7.49 (d, 1H), 7.34 (d, 1H), 6.99 (dd, 1H), 6.18 (tt, 1H), 5.33 (s, 2H), 3.72 (tdd, 2H), 2.62 (s, 3H), 2.43 (s, 3H).
499	¹ H NMR (500 MHz, DMSO) δ = 8.35 (t, 1H), 7.69 (s, 1H), 7.48 (d, 1H), 7.27 (d, 1H), 6.97 (dd, 1H), 5.32 (s, 2H), 3.75 (td, 2H), 2.61 (d, 6H), 1.66 (t, 3H).
500	¹ H NMR (500 MHz, DMSO) δ = 8.98 (s, 1H), 8.36 (t, 1H), 7.48 (d, 1H), 7.28 (d, 1H), 6.98 (dd, 1H), 5.32 (s, 2H), 3.75 (td, 2H), 2.60 (s, 3H), 2.42 (s, 3H), 1.66 (t, 3H).
501	¹ H NMR (600 MHz, CDCl ₃) δ = 8.73 (s, 1H), 7.45 (d, 1H), 7.36 (d, 1H), 6.94 (dd, 1H), 6.78 (s, 1H), 6.14 (s, 1H), 5.34 (s, 2H), 3.69-3.61 (m, 2H), 3.61-3.55 (m, 2H), 2.75 (s, 3H), 2.53 (s, 3H), 2.02 (s, 3H).
502	¹ H NMR (400 MHz, DMSO) δ = 8.99 (s, 1H), 8.87 (d, 1H), 7.85 (s, 1H), 7.50 (d, 1H), 7.45 (d, 1H), 7.00 (dd, 1H), 5.65 (d, 1H), 5.33 (s, 2H), 2.67 (s, 3H), 2.43 (s, 3H).
503	¹ H NMR (400 MHz, DMSO) δ = 8.99 (s, 1H), 7.94 (t, 1H), 7.52-7.41 (m, 2H), 7.42 (s, 1H), 7.07 (s, 1H), 6.97 (dd, 1H), 5.33 (s, 2H), 3.87 (d, 2H), 2.64 (s, 3H), 2.42 (s, 3H).

-continued

cpd Nr	NMR Data
504	¹ H NMR (500 MHz, DMSO) δ = 8.17 (t, 1H), 7.68 (s, 1H), 7.47 (d, 1H), 7.26 (d, 1H), 6.96 (dd, 1H), 5.32 (s, 2H), 3.50 (dd, 2H), 2.61 (d, 6H), 1.37 (d, 6H).
505	¹ H NMR (500 MHz, DMSO) δ = 8.98 (s, 1H), 8.17 (t, 1H), 7.47 (d, 1H), 7.27 (d, 1H), 6.97 (dd, 1H), 5.31 (s, 2H), 3.50 (dd, 2H), 2.60 (s, 3H), 2.41 (s, 3H), 1.37 (d, 6H).
506	¹ H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 7.74 (t, 1H), 7.47 (d, 1H), 7.32 (d, 1H), 6.97 (dd, 1H), 5.31 (s, 2H), 4.61 (s, 1H), 3.28 (d, 2H), 2.61 (s, 3H), 2.41 (s, 3H), 1.15 (s, 6H).
507	¹ H NMR (600 MHz, DMSO) δ = 8.97 (s, 1H), 7.52 (d, 1H), 7.47 (d, 1H), 6.98 (dd, 1H), 5.34 (s, 2H), 4.27 (q, 3H), 3.55 (d, 1H), 2.76 (d, 1H), 2.69 (d, 1H), 2.68 (s, 3H), 2.46 (s, 3H).
508	¹ H NMR (400 MHz, DMSO) δ = 10.65 (s, 1H), 9.00 (s, 1H), 7.54 (d, 1H), 7.37 (s, 1H), 7.02 (dd, 1H), 5.33 (s, 2H), 4.70 (t, 4H), 2.67 (s, 3H), 2.42 (s, 3H), 2.35 (s, 6H).
509	¹ H NMR (400 MHz, DMSO) δ = 9.14 (s, 1H), 8.99 (s, 1H), 7.49 (d, 1H), 7.25 (d, 1H), 6.99 (dd, 1H), 5.33 (s, 2H), 5.02 (d, 2H), 4.78 (d, 2H), 2.85 (s, 6H), 2.60 (s, 3H), 2.41 (s, 3H).
510 - En 1	¹ H NMR (400 MHz, DMSO) δ = 8.77-8.71 (m, 1H), 8.31-8.24 (m, 1H), 7.79 (dd, 1H), 7.57 (d, 1H), 7.55-7.47 (m, 2H), 7.29 (d, 1H), 6.99 (dd, 1H), 5.31 (s, 2H), 5.18 (s, 1H), 3.73 (d, 1H), 3.62 (d, 1H), 3.30-3.21 (m, 1H), 3.17-3.09 (m, 1H), 2.60 (s, 3H), 2.23 (dd, 0H), 2.18 (s, 2H), 1.79 (q, 2H).
511 - En 1	¹ H NMR (400 MHz, DMSO) δ = 7.59 (d, 1H), 7.53-7.45 (m, 2H), 7.27 (d, 1H), 7.17 (s, 1H), 6.97 (dd, 1H), 5.25 (s, 1H), 5.12 (s, 2H), 3.74 (d, 1H), 3.64 (d, 1H), 3.26 (dd, 1H), 3.16-3.10 (m, 1H), 2.60 (s, 3H), 2.42 (s, 3H), 2.28-2.14 (m, 2H), 1.85-1.72 (m, 2H).
512 - En 1	¹ H NMR (400 MHz, DMSO) δ = 8.41 (dd, 1H), 7.83 (dd, 1H), 7.62-7.56 (m, 1H), 7.54-7.46 (m, 2H), 7.32 (d, 1H), 7.25 (dd, 1H), 7.01 (dd, 1H), 5.24 (s, 1H), 5.15 (s, 2H), 3.74 (d, 1H), 3.63 (d, 1H), 3.26 (dd, 1H), 3.17-3.08 (m, 1H), 2.57 (d, 6H), 2.29-2.13 (m, 2H), 1.86-1.73 (m, 2H).
513 - En 1	¹ H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 7.58 (s, 1H), 7.53-7.45 (m, 2H), 7.27 (d, 1H), 6.97 (dd, 1H), 5.31 (s, 2H), 5.21 (t, 1H), 3.74 (dd, 1H), 3.63 (dd, 1H), 3.12 (d, 1H), 2.60 (s, 3H), 2.42 (s, 3H), 2.21 (d, 2H), 1.78 (s, 2H), 1.24 (s, 1H).
514 - En 1	¹ H NMR (500 MHz, DMSO) δ = 8.55 (d, 1H), 7.78 (dd, 1H), 7.58 (d, 1H), 7.51 (s, 1H), 7.48 (d, 1H), 7.31-7.24 (m, 2H), 6.98 (dd, 1H), 5.28 (s, 1H), 5.12 (s, 2H), 3.74 (d, 1H), 3.64 (d, 1H), 3.26 (dd, 1H), 3.13 (dd, 1H), 2.60 (s, 3H), 2.47 (s, 3H), 2.27-2.14 (m, 2H), 1.85-1.73 (m, 1H).
515 - En 1	¹ H NMR (500 MHz, DMSO) δ = 7.80 (s, 1H), 7.51-7.45 (m, 2H), 7.25 (d, 1H), 7.17 (s, 1H), 6.97 (dd, 1H), 5.22 (t, 1H), 5.12 (s, 2H), 3.67-3.55 (m, 2H), 3.29 (s, 0H), 3.21 (q, 1H), 2.59 (s, 3H), 2.49-2.36 (m, 2H), 2.42 (s, 3H).
516 - En 1	¹ H NMR (400 MHz, DMSO) δ = 8.41 (dd, 1H), 7.86-7.77 (m, 2H), 7.53-7.45 (m, 2H), 7.30 (d, 1H), 7.25 (dd, 1H), 7.01 (dd, 1H), 5.22 (s, 1H), 5.15 (s, 2H), 3.67-3.55 (m, 2H), 3.31-3.14 (m, 2H), 2.57 (d, 6H), 2.50-2.37 (m, 1H).
517 - En 1	¹ H NMR (500 MHz, DMSO) δ = 7.79 (s, 1H), 7.71 (s, 1H), 7.51 (s, 1H), 7.47 (d, 1H), 7.24 (d, 1H), 6.96 (dd, 1H), 5.32 (s, 2H), 3.63 (d, 1H), 3.58 (d, 1H), 3.30 (d, 1H), 3.21 (q, 1H), 2.61 (d, 6H), 2.48-2.35 (m, 1H).
518 - En 1	¹ H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 7.78 (s, 1H), 7.53-7.44 (m, 2H), 7.25 (d, 1H), 6.97 (dd, 1H), 5.31 (s, 2H), 5.21 (t, 1H), 3.62 (td, 2H), 3.20 (d, 1H), 2.59 (s, 3H), 2.49-2.34 (m, 1H), 2.42 (s, 3H).
519 - En 1	¹ H NMR (400 MHz, DMSO) δ = 8.55 (d, 1H), 7.81-7.74 (m, 2H), 7.51-7.44 (m, 2H), 7.31-7.22 (m, 2H), 6.97 (dd, 1H), 5.23 (s, 1H), 5.12 (s, 2H), 3.67-3.54 (m, 2H), 3.21 (q, 1H), 2.59 (s, 3H), 2.47 (s, 3H), 2.45 (d, 0H), 2.44-2.34 (m, 1H).
520 - En 1	¹ H NMR (500 MHz, DMSO) δ = 8.58 (d, 1H), 7.88-7.80 (m, 1H), 7.78 (s, 1H), 7.56 (d, 1H), 7.52-7.45 (m, 2H), 7.35 (dd, 1H), 7.27 (d, 1H), 7.00 (dd, 1H), 5.25 (t, 1H), 5.19 (s, 2H), 3.60 (qd, 2H), 3.20 (q, 1H), 2.59 (s, 3H), 2.49-2.37 (m, 3H).
521 - En 1	¹ H NMR (500 MHz, DMSO) δ = 11.77 (s, 1H), 7.78 (s, 1H), 7.59 (dd, 1H), 7.51 (s, 1H), 7.47 (d, 1H), 7.38 (dd, 1H), 7.26-7.22 (m, 1H), 6.95 (dd, 1H), 6.23 (t, 1H), 5.24 (s, 1H), 4.88 (s, 2H), 3.63 (d, 1H), 3.58 (d, 1H), 3.24-3.14 (m, 1H), 2.59 (s, 3H), 2.49-2.36 (m, 1H).
522 - En 1	¹ H NMR (500 MHz, DMSO) δ = 8.16 (dd, 1H), 7.84 (dd, 1H), 7.78 (s, 1H), 7.51 (s, 1H), 7.47 (d, 1H), 7.25 (d, 1H), 7.03 (dd, 1H), 6.97 (dd, 1H), 5.21 (s, 1H), 5.07 (s, 2H), 3.93 (s, 3H), 3.60 (q, 2H), 3.20 (q, 1H), 2.59 (s, 3H), 2.48-2.37 (m, 1H).
523	¹ H NMR (400 MHz, DMSO) δ = 8.31 (s, 1H), 7.47 (d, 1H), 7.41 (s, 1H), 7.19 (s, 1H), 7.00 (d, 1H), 6.87 (s, 1H), 5.26 (t, 1H), 5.16 (s, 2H), 4.70 (d, 2H), 4.55 (d, 2H), 3.79 (d, 2H), 3.69 (s, 3H), 2.60 (s, 3H).
524	¹ H NMR (400 MHz, DMSO) δ = 8.34 (s, 1H), 7.48 (d, 1H), 7.37 (dd, 2H), 6.99 (dd, 1H), 6.38 (d, 1H), 5.20 (d, 3H), 4.68 (d, 2H), 4.55 (d, 2H), 3.85 (s, 3H), 3.78 (s, 2H), 2.61 (s, 3H).
525	¹ H NMR (400 MHz, DMSO-d ₆) δ = 8.35 (s, 1H), 7.69 (s, 1H), 7.46 (d, 1H), 7.32 (s, 1H), 7.95 (d, 1H), 5.32 (s, 2H), 5.19 (t, 1H), 4.67 (d, 2H), 4.54 (d, 2H), 3.77 (d, 2H), 2.63 (s, 3H), 2.60 (s, 3H).
526	¹ H NMR (400 MHz, DMSO) δ = 8.62-8.55 (m, 1H), 8.33 (s, 1H), 7.84 (td, 1H), 7.56 (d, 1H), 7.47 (d, 1H), 7.39-7.31 (m, 2H), 7.00 (dd, 1H), 5.21 (s, 2H), 5.19 (d, 1H), 4.67 (d, 2H), 4.55 (d, 2H), 3.77 (d, 2H), 2.60 (s, 3H).
527	¹ H NMR (400 MHz, DMSO) δ = 8.33 (s, 1H), 7.85 (d, 1H), 7.79 (d, 1H), 7.49 (d, 1H), 7.38 (d, 1H), 7.03 (dd, 1H), 5.48 (s, 2H), 5.20 (t, 1H), 4.68 (d, 2H), 4.55 (d, 2H), 3.78 (d, 2H), 2.61 (s, 3H).
528	¹ H NMR (400 MHz, DMSO) δ = 8.35 (s, 1H), 8.16 (dd, 1H), 7.88-7.79 (m, 1H), 7.47 (d, 1H), 7.33 (d, 1H), 7.03 (dd, 1H), 6.97 (dd, 1H), 5.19 (t, 1H), 5.08 (s, 2H), 4.67 (d, 2H), 4.55 (d, 2H), 3.93 (s, 3H), 3.77 (d, 2H), 2.61 (s, 3H).
529	¹ H NMR (400 MHz, DMSO) δ = 8.99 (s, 1H), 7.48 (dd, 1H), 7.08 (d, 1H), 6.96 (dd, 1H), 5.35 (s, 2H), 5.35-5.24 (m, 1H), 4.77 (d, 2H), 4.47-4.39 (m, 2H), 4.00 (d, 2H), 2.87 (s, 3H), 2.45 (d, 6H).
530 - En 1	¹ H NMR (500 MHz, DMSO) δ = 7.70 (s, 1H), 7.54 (s, 1H), 7.45 (d, 1H), 7.25 (d, 1H), 6.93 (dd, 1H), 5.31 (s, 2H), 5.08 (s, 1H), 3.66 (dd, 2H), 3.00 (d, 1H), 2.95-2.85 (m, 1H), 2.79 (ddd, 1H), 2.63 (s, 3H), 2.56 (d, 4H), 2.06 (ddd, 1H), 1.79 (dt, 1H).
531 - En 1	¹ H NMR (400 MHz, DMSO) δ = 8.74 (dd, 1H), 8.27 (d, 1H), 7.79 (dd, 1H), 7.49 (d, 1H), 7.31 (d, 1H), 7.24 (s, 1H), 6.98 (dd, 1H), 5.31 (s, 2H), 4.88 (t, 1H), 4.00 (dd, 1H), 3.76-3.60 (m, 3H), 3.47 (dd, 2H), 2.60 (s, 3H), 2.29 (dd, 1H), 1.70 (dt, 2H), 1.51 (d, 1H).
532 - En 1	¹ H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 7.46 (d, 1H), 7.31 (d, 1H), 7.21 (s, 1H), 6.95 (dd, 1H), 5.30 (s, 2H), 4.90 (t, 1H), 4.05-3.96 (m, 1H), 3.76-3.61 (m, 3H), 3.51 (d, 1H), 3.46 (s, 1H), 2.60 (s, 3H), 2.42 (s, 3H), 2.31 (d, 1H), 1.75-1.65 (m, 2H), 1.52 (d, 1H).

-continued

cpd Nr	NMR Data
533 - En 1	1H NMR (500 MHz, DMSO) δ = 7.84 (s, 1H), 7.69 (s, 1H), 7.45 (d, 1H), 7.24 (d, 1H), 6.94 (dd, 1H), 5.31 (s, 2H), 5.11-5.05 (m, 1H), 3.92 (d, 1H), 3.86-3.75 (m, 3H), 3.72 (dd, 1H), 3.66 (dd, 1H), 2.63 (s, 3H), 2.57 (s, 3H), 2.25 (ddd, 1H), 2.06 (dt, 1H).
534 - En 1	1H NMR (400 MHz, DMSO) δ = 8.58 (ddd, 1H), 7.88-7.79 (m, 2H), 7.55 (d, 1H), 7.46 (d, 1H), 7.35 (ddd, 1H), 7.25 (d, 1H), 6.98 (dd, 1H), 5.20 (s, 2H), 5.08 (d, 1H), 3.92 (d, 1H), 3.87-3.74 (m, 3H), 3.71 (dd, 1H), 3.65 (dd, 1H), 2.57 (s, 3H), 2.23 (ddd, 1H), 2.06 (dt, 1H).
535 - En 1	1H NMR (400 MHz, DMSO) δ = 11.77 (s, 1H), 7.85 (s, 1H), 7.61-7.54 (m, 1H), 7.45 (d, 1H), 7.41-7.34 (m, 1H), 7.22 (d, 1H), 6.94 (dd, 1H), 6.22 (t, 1H), 5.08 (s, 1H), 4.88 (s, 2H), 3.92 (d, 1H), 3.80 (dt, 3H), 3.75-3.62 (m, 2H), 2.57 (s, 3H), 2.24 (dt, 1H), 2.06 (dt, 1H).
536 - En 1	1H NMR (400 MHz, DMSO) δ = 8.16 (dd, 1H), 7.88-7.79 (m, 2H), 7.45 (d, 1H), 7.24 (d, 1H), 7.03 (dd, 1H), 6.95 (dd, 1H), 5.06 (d, 3H), 3.92 (d, 4H), 3.80 (d, 2H), 3.78 (d, 1H), 3.68 (qd, 2H), 2.57 (s, 3H), 2.23 (dt, 1H), 2.05 (dt, 1H).
537 - En 1	1H NMR (500 MHz, DMSO) δ = 8.99 (s, 1H), 8.24 (s, 1H), 7.47 (d, 1H), 7.27 (d, 1H), 6.96 (dd, 1H), 5.31 (s, 2H), 5.22 (t, 1H), 3.65 (d, 2H), 2.87 (dd, 4H), 2.58 (s, 3H), 2.42 (s, 3H).
538 - En 1	1H NMR (400 MHz, DMSO) δ = 8.78-8.65 (m, 2H), 8.31-8.20 (m, 1H), 7.79 (dd, 1H), 7.33 (dd, 1H), 7.03 (d, 1H), 5.33-5.24 (m, 2H), 4.47 (d, 1H), 3.09 (s, 1H), 2.97-2.72 (m, 2H), 2.69-2.55 (m, 1H), 2.42 (s, 3H), 2.26 (s, 3H), 1.79 (d, 1H), 1.72-1.55 (m, 1H), 1.24 (s, 1H).
539 - En 1	1H NMR (400 MHz, DMSO) δ = 8.33-8.21 (m, 2H), 7.91 (d, 1H), 7.79-7.64 (m, 2H), 7.60 (dd, 1H), 7.39 (dd, 1H), 5.41 (s, 2H), 4.49 (s, 1H), 3.20-3.04 (m, 1H), 2.98-2.75 (m, 2H), 2.70-2.54 (m, 1H), 2.59 (s, 3H), 1.86-1.60 (m, 2H).
540 - En 1	1H NMR (500 MHz, DMSO) δ = 8.20 (d, 1H), 8.15 (s, 1H), 7.47 (d, 1H), 7.23 (d, 1H), 6.97 (dd, 1H), 5.14 (d, 2H), 4.54-4.42 (m, 1H), 4.05 (s, 2H), 3.30 (s, 1H), 3.11 (q, 1H), 2.92 (d, 1H), 2.84 (dd, 1H), 2.62 (d, 1H), 2.57 (s, 3H), 2.44 (s, 1H), 1.73 (ddd, 2H).
541 - En 1	1H NMR (400 MHz, DMSO) δ = 8.23 (d, 1H), 7.79 (s, 1H), 7.49 (d, 1H), 7.25 (d, 1H), 7.00 (dd, 1H), 5.30 (s, 2H), 4.55-4.40 (m, 1H), 4.06 (s, 3H), 3.13 (t, 1H), 2.99-2.74 (m, 2H), 2.63 (d, 1H), 2.57 (s, 3H), 1.83-1.63 (m, 2H).
542 - En 1	1H NMR (400 MHz, DMSO) δ = 8.46 (s, 1H), 8.18 (d, 1H), 7.46 (d, 1H), 7.23 (d, 1H), 6.97 (dd, 1H), 5.05 (d, 2H), 4.47 (s, 1H), 3.86 (s, 3H), 3.20-3.04 (m, 1H), 2.98-2.74 (m, 2H), 2.62 (d, 1H), 2.57 (s, 3H), 2.43 (s, 1H), 1.78 (s, 1H), 1.86-1.63 (m, 1H).
543 - En 1	1H NMR (400 MHz, DMSO) δ = 8.21 (d, 1H), 7.48 (d, 1H), 7.37 (d, 1H), 7.24 (d, 1H), 6.99 (dd, 1H), 6.36 (d, 1H), 5.18 (s, 2H), 4.47 (dd, 1H), 3.85 (s, 3H), 3.21-3.04 (m, 1H), 2.98-2.74 (m, 2H), 2.57 (s, 3H), 2.43 (s, 1H), 1.78 (s, 2H), 1.86-1.62 (m, 1H).
544 - En 1	1H NMR (500 MHz, DMSO) δ = 8.20 (d, 1H), 7.71 (d, 1H), 7.55 (d, 1H), 7.46 (d, 1H), 7.18 (s, 1H), 6.95 (d, 1H), 6.26 (t, 1H), 4.92 (t, 2H), 4.47 (s, 1H), 3.48 (s, 3H), 3.39 (s, 2H), 3.11 (s, 1H), 2.97-2.75 (m, 2H), 2.59 (d, 3H), 1.78 (s, 1H), 1.71 (d, 1H).
545 - En 1	1H NMR (400 MHz, DMSO) δ = 8.22 (d, 1H), 7.50 (d, 1H), 7.39 (dd, 1H), 7.24 (d, 1H), 7.03 (dd, 1H), 6.48-6.36 (m, 2H), 5.14 (s, 2H), 3.49 (s, 3H), 3.20-3.04 (m, 1H), 2.98-2.74 (m, 2H), 2.62 (d, 1H), 2.58 (s, 3H), 2.43 (s, 1H), 1.86-1.61 (m, 2H).
546 - En 1	1H NMR (DMSO-d6) δ = 8.73 (d, 1H), 8.25 (t, 2H), 7.79 (dd, 1H), 7.50 (d, 1H), 7.22 (d, 1H), 7.00 (dd, 1H), 5.31 (s, 2H), 4.37-4.58 (m, 1H), 3.04-3.17 (m, 1H), 2.74-2.96 (m, 2H), 2.59-2.72 (m, 1H), 2.58 (s, 3H), 1.63-1.85 (m, 2H).
547 - En 1	1H NMR (500 MHz, DMSO) δ = 8.41 (dd, 1H), 8.22 (d, 1H), 7.80 (dd, 1H), 7.48 (d, 1H), 7.29-7.19 (m, 2H), 7.00 (dd, 1H), 5.14 (s, 2H), 4.47 (dd, 1H), 3.12 (s, 0H), 2.92 (d, 1H), 2.84 (dd, 1H), 2.55 (d, 7H), 2.44 (s, 1H), 1.82-1.64 (m, 2H).
548 - En 1	1H NMR (400 MHz, DMSO) δ = 8.70 (d, 1H), 8.20 (d, 1H), 7.49 (d, 1H), 7.41 (d, 1H), 7.19 (d, 1H), 7.02 (dd, 1H), 5.18 (s, 2H), 4.46 (d, 1H), 3.19-3.03 (m, 1H), 2.98-2.73 (m, 2H), 2.60 (d, 7H), 1.76 (s, 1H), 1.80-1.62 (m, 1H).
549 - En 1	1H NMR (400 MHz, DMSO) δ = 8.20 (d, 1H), 7.68 (s, 1H), 7.47 (d, 1H), 7.20 (d, 1H), 6.96 (dd, 1H), 5.31 (s, 2H), 4.48 (s, 1H), 3.12 (s, 1H), 2.92 (d, 2H), 2.80 (d, 1H), 2.60 (s, 1H), 2.60 (d, 6H), 1.77 (s, 2H), 1.78-1.66 (m, 0H).
550 - En 1	1H NMR (400 MHz, DMSO) δ = 8.22 (d, 1H), 7.46 (dd, 1H), 7.17 (d, 1H), 6.94 (dt, 1H), 4.50 (d, 2H), 4.45 (s, 1H), 4.32 (d, 2H), 4.06 (d, 2H), 3.22-3.04 (m, 1H), 2.99-2.74 (m, 2H), 2.59 (d, 1H), 2.57 (s, 3H), 1.78 (s, 2H), 1.81-1.63 (m, 1H), 1.38 (s, 3H).
551 - En 1	1H NMR (400 MHz, DMSO) δ = 8.20 (d, 1H), 7.83 (q, 1H), 7.48 (d, 1H), 7.22 (d, 1H), 6.98 (dd, 1H), 5.16 (s, 2H), 4.47 (s, 1H), 3.20-3.04 (m, 1H), 2.98-2.74 (m, 2H), 2.63 (d, 1H), 2.57 (s, 3H), 2.43 (s, 1H), 2.09 (d, 3H), 1.74 (ddd, 2H).
552 - En 1	1H NMR (400 MHz, DMSO) δ = 8.21 (d, 1H), 7.49 (d, 1H), 7.29 (d, 1H), 7.24 (d, 1H), 7.01 (dd, 1H), 5.38 (s, 2H), 4.46 (dd, 1H), 3.22-3.04 (m, 0H), 3.12 (s, 1H), 2.99-2.74 (m, 2H), 2.61 (s, 1H), 2.57 (s, 3H), 2.37 (d, 4H), 1.74 (ddd, 2H).
553 - En 1	1H NMR (400 MHz, DMSO) δ = 8.20 (d, 1H), 7.48 (d, 1H), 7.21 (d, 1H), 6.97 (dd, 1H), 6.34 (d, 1H), 5.14 (s, 2H), 4.47 (s, 1H), 3.20-3.04 (m, 1H), 2.98-2.74 (m, 2H), 2.62 (d, 1H), 2.57 (s, 3H), 2.41 (d, 3H), 1.78 (s, 1H), 1.80-1.62 (m, 1H).
554 - En 1	1H NMR (400 MHz, DMSO) δ = 8.67 (d, 1H), 8.56 (d, 1H), 8.20 (d, 1H), 7.48 (d, 1H), 7.23 (d, 1H), 7.01 (dd, 1H), 5.22 (s, 2H), 4.53-4.40 (m, 1H), 3.13 (s, 1H), 2.93 (s, 2H), 2.57 (s, 3H), 1.80 (s, 1H), 1.71 (d, 1H).
555 - En 1	1H NMR (400 MHz, DMSO) δ = 8.20 (d, 1H), 7.54-7.43 (m, 2H), 7.23 (d, 1H), 7.00 (dd, 1H), 5.35 (s, 2H), 4.45 (s, 1H), 3.12 (s, 1H), 2.99-2.74 (m, 2H), 2.63 (d, 1H), 2.57 (s, 3H), 2.44 (d, 3H), 1.82 (d, 1H), 1.62 (m, 2H).
556 - En 1	1H NMR (400MHz, DMSO) δ = 8.19 (d, 1H), 7.71 (d, 1H), 7.60 (d, 1H), 7.47 (d, 1H), 7.23 (d, 1H), 6.99 (dd, 1H), 5.36 (s, 2H), 4.41-4.5 (m, 1H), 3.09-3.14 (t, 1H), 2.70-2.93 (m, 2H), 2.57-2.78 (m, 7H), 2.50 (bs, 1H), 1.72-1.78 (m, 2H).
557 - En 1	1H NMR (400 MHz, DMSO) δ = 9.22 (dd, 1H), 8.21 (d, 1H), 7.85 (dd, 1H), 7.76 (dd, 1H), 7.49 (d, 1H), 7.25 (d, 1H), 7.03 (dd, 1H), 5.42 (s, 2H), 4.47 (d, 1H), 3.11 (q, 1H), 2.98-2.73 (m, 2H), 2.62 (d, 1H), 2.57 (s, 3H), 1.85-1.63 (m, 2H).

-continued

cpd Nr	NMR Data
558 - En 1	1H NMR (400 MHz, DMSO) δ = 8.68 (d, 1H), 8.55 (dd, 1H), 8.21 (d, 1H), 7.88 (dt, 1H), 7.53-7.37 (m, 2H), 7.23 (d, 1H), 6.99 (dd, 1H), 5.17 (s, 2H), 4.55-4.40 (m, 1H), 3.19-3.04 (m, 1H), 2.98-2.74 (m, 2H), 2.62 (d, 1H), 2.57 (s, 3H), 1.72 (ddd, 2H).
559 - En 1	1H NMR (400 MHz, DMSO) δ = 8.83 (d, 2H), 8.16 (d, 1H), 7.52-7.39 (m, 2H), 7.15 (d, 1H), 6.95 (dd, 1H), 5.28 (s, 2H), 3.10 (s, 1H), 2.98-2.72 (m, 2H), 2.56 (s, 4H), 2.42 (s, 1H), 1.81-1.60 (m, 2H).
560 - En 1	1H NMR (400 MHz, DMSO) δ = 8.21 (d, 1H), 7.84 (d, 1H), 7.78 (d, 1H), 7.49 (d, 1H), 7.25 (d, 1H), 7.03 (dd, 1H), 5.45 (s, 2H), 3.20-3.04 (m, 1H), 2.99-2.74 (m, 2H), 2.63 (d, 1H), 2.57 (s, 3H), 2.44 (s, 1H), 1.78 (s, 1H), 1.86-1.62 (m, 1H).
561 - En 1	1H NMR (400 MHz, DMSO) δ = 9.11 (d, 1H), 8.21 (d, 1H), 7.99 (d, 1H), 7.48 (d, 1H), 7.22 (d, 1H), 6.98 (dd, 1H), 5.40 (s, 2H), 4.45 (s, 0H), 3.20-3.04 (m, 1H), 2.98-2.74 (m, 2H), 2.61 (t, 1H), 2.57 (s, 3H), 1.74 (dt, 2H).
562 - En 1	1H NMR (400 MHz, DMSO) δ = 8.99 (s, 1H), 8.40 (d, 1H), 7.36 (dd, 1H), 7.24 (dd, 1H), 5.37 (s, 2H), 3.08 (q, 1H), 2.96-2.72 (m, 2H), 2.59 (d, 2H), 2.51 (s, 3H), 2.39 (s, 3H), 1.79 (dt, 1H), 1.63 (qd, 1H).
563 - En 1	1H NMR (500 MHz, DMSO) δ = 8.20 (d, 1H), 7.46 (d, 1H), 7.20 (d, 1H), 6.94 (dd, 1H), 5.22 (s, 2H), 4.47 (s, 1H), 3.12 (s, 1H), 2.87 (dt, 2H), 2.62 (d, 1H), 2.57 (s, 6H), 2.32 (s, 3H), 1.84-1.64 (m, 2H).
564 - En 1	1H NMR (400 MHz, DMSO) δ = 11.75 (s, 1H), 8.20 (d, 1H), 7.56 (dd, 1H), 7.46 (d, 1H), 7.38 (dd, 1H), 7.18 (d, 1H), 6.95 (dd, 1H), 6.22 (t, 1H), 4.96-4.80 (m, 2H), 4.48 (d, 1H), 3.12 (t, 1H), 2.92 (d, 1H), 2.80 (s, 1H), 2.57 (s, 3H), 2.42 (s, 1H), 1.83-1.66 (m, 2H).
565 - En 1	1H NMR (400 MHz, DMSO) δ = 8.22 (d, 1H), 8.15 (dd, 1H), 7.81 (dd, 1H), 7.47 (d, 1H), 7.20 (d, 1H), 7.00 (ddd, 2H), 5.06 (s, 2H), 4.47 (dt, 1H), 3.92 (s, 3H), 3.11 (s, 1H), 2.98-2.73 (m, 2H), 2.60 (s, 1H), 2.57 (s, 3H), 2.43 (s, 1H), 1.85-1.61 (m, 2H).
566 - En 1	1H NMR (500 MHz, DMSO) δ = 8.45 (dt, 1H), 8.20 (d, 1H), 7.79 (ddd, 1H), 7.53 (dt, 1H), 7.47 (d, 1H), 7.24 (d, 1H), 6.98 (dd, 1H), 5.29-5.17 (m, 2H), 3.18-3.06 (m, 1H), 2.88 (s, 1H), 2.97-2.75 (m, 1H), 2.61 (s, 1H), 2.57 (s, 3H), 2.42 (s, 1H), 1.84-1.64 (m, 2H).
567 - En 1	1H NMR (400 MHz, DMSO) δ = 8.28 (d, 1H), 8.19 (d, 1H), 7.44 (qd, 3H), 7.21 (d, 1H), 6.97 (dd, 1H), 5.10 (s, 2H), 4.91 (t, 1H), 4.48 (d, 1H), 4.07 (dd, 2H), 3.73 (q, 2H), 3.11 (d, 1H), 2.98-2.74 (m, 2H), 2.69-2.53 (m, 1H), 2.57 (s, 3H), 1.78 (s, 2H), 1.24 (s, 1H).
568 - En 1	1H NMR (500 MHz, DMSO) δ = 8.58 (d, 1H), 8.19 (d, 1H), 7.77 (td, 1H), 7.63 (dd, 1H), 7.48 (d, 1H), 7.21 (d, 1H), 6.99 (dd, 1H), 5.19 (s, 2H), 4.55-4.41 (m, 1H), 3.12 (t, 1H), 2.86 (dt, 2H), 2.57 (s, 4H), 2.47-2.39 (m, 1H), 1.82-1.74 (m, 1H), 1.71 (qd, 1H).
569 - En 1	1H NMR (400 MHz, DMSO) δ = 11.68 (s, 1H), 8.21 (d, 1H), 7.45 (dd, 2H), 7.18 (d, 1H), 6.97 (dd, 1H), 6.35-6.24 (m, 2H), 4.91 (s, 2H), 4.45 (s, 1H), 3.10 (d, 1H), 2.98-2.74 (m, 2H), 2.61 (s, 1H), 2.57 (s, 3H), 1.78 (s, 1H), 1.79-1.62 (m, 1H).
570 - En 1	1H NMR (400 MHz, DMSO) δ = 8.20 (d, 1H), 7.72 (dd, 1H), 7.47 (d, 1H), 7.23 (d, 1H), 7.10 (dd, 1H), 6.99 (dd, 1H), 6.75 (dd, 1H), 5.11 (s, 2H), 3.86 (s, 3H), 3.19-3.03 (m, 1H), 2.98-2.73 (m, 2H), 2.62 (d, 1H), 2.57 (s, 3H), 2.43 (s, 1H), 1.76 (s, 1H), 1.84-1.61 (m, 1H).
571 - En 1	1H NMR (400 MHz, DMSO) δ = 8.72-8.63 (m, 1H), 8.19 (dd, 1H), 7.96 (dt, 1H), 7.64-7.53 (m, 2H), 7.48 (dd, 1H), 7.38-7.20 (m, 2H), 6.99 (ddd, 1H), 5.13 (s, 2H), 4.43 (s, 2H), 4.11 (d, 1H), 3.12 (s, 1H), 2.88 (dd, 1H), 2.57 (d, 3H), 1.82-1.61 (m, 2H).
572 - En 1	1H NMR (400 MHz, DMSO) δ = 8.99 (s, 1H), 8.26 (d, 1H), 7.62 (d, 1H), 7.40 (d, 1H), 5.38 (s, 2H), 4.48 (dt, 1H), 3.19-3.04 (m, 1H), 2.99-2.74 (m, 2H), 2.61 (s, 1H), 2.56 (s, 3H), 2.44 (s, 1H), 2.41 (s, 3H), 1.86-1.61 (m, 2H).
573 - En 1	1H NMR (500 MHz, DMSO) δ = 8.99 (s, 1H), 8.33 (d, 1H), 7.09-6.95 (m, 2H), 5.33 (s, 2H), 4.47 (d, 1H), 3.11 (s, 1H), 2.98-2.73 (m, 2H), 2.59 (s, 3H), 2.43 (d, 1H), 2.42 (s, 3H), 1.72 (ddd, 2H), 1.23 (s, 1H).
574 - En 1	1H NMR (500 MHz, DMSO) δ = 8.21 (dd, 2H), 8.06 (s, 1H), 7.95 (t, 1H), 7.57 (d, 1H), 4.24 (t, 1H), 3.74 (d, 1H), 3.21-3.11 (m, 3H), 2.55 (d, 3H), 2.17 (d, 1H), 2.11-1.96 (m, 1H), 1.99 (s, 1H), 1.74 (d, 1H), 1.56 (s, 3H), 1.43 (d, 1H), 1.01 (d, 6H), 0.86 (s, 0H).
575 - En 1	1H NMR (500 MHz, DMSO) δ = 8.99 (s, 1H), 7.78 (t, 1H), 7.46 (d, 2H), 7.40 (d, 1H), 7.16 (s, 1H), 6.95 (dd, 1H), 5.32 (s, 2H), 3.56 (dt, 1H), 3.46 (ddd, 1H), 3.23 (t, 1H), 2.60 (s, 3H), 2.42 (s, 3H), 2.30 (s, 6H).
576 - En 1	1H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 7.47 (d, 1H), 7.36 (d, 1H), 7.28 (s, 1H), 7.00 (s, 1H), 6.95 (dd, 1H), 5.33 (s, 2H), 3.40 (d, 2H), 2.61 (s, 3H), 2.42 (s, 3H), 1.16 (s, 6H).
577 - En 1	1H NMR (400 MHz, DMSO) δ = 9.30 (s, 1H), 9.02 (s, 1H), 7.75 (dd, 1H), 7.64-7.54 (m, 1H), 7.42 (d, 1H), 7.28 (s, 1H), 7.33-7.22 (m, 1H), 5.42 (s, 2H), 4.89 (d, 2H), 4.74 (d, 2H), 2.45 (s, 3H).
578 - En 1	1H NMR (500 MHz, DMSO) δ = 8.70 (s, 1H), 7.56 (d, 1H), 7.49 (d, 1H), 7.23 (d, 2H), 7.19 (d, 1H), 7.02 (dd, 1H), 6.87 (d, 1H), 5.19 (s, 2H), 4.88 (d, 2H), 4.71 (d, 2H), 3.70 (s, 3H), 2.65 (s, 3H).
579 - En 1	1H NMR (400 MHz, DMSO) δ = 8.69 (s, 1H), 7.54-7.46 (m, 2H), 7.38 (d, 1H), 7.22 (d, 2H), 7.01 (dd, 1H), 6.38 (d, 1H), 5.21 (s, 2H), 4.88 (d, 2H), 4.68 (d, 2H), 3.86 (s, 3H), 2.67 (s, 3H).
580 - En 1	1H-NMR (400 MHz, DMSO-d ₆) δ = 8.67 (s, 1H), 7.70 (s, 1H), 7.51-7.44 (m, 2H), 7.24-7.19 (m, 2H), 6.98 (dd, 1H), 5.34 (s, 2H), 4.87 (d, 2H), 4.68 (d, 2H), 2.65 (s, 3H), 2.63 (s, 3H).
581 - En 1	1H NMR (400 MHz, DMSO) δ = 8.70 (s, 1H), 8.59 (ddd, 1H), 7.85 (td, 1H), 7.57 (dt, 1H), 7.53-7.45 (m, 2H), 7.35 (ddd, 1H), 7.21 (d, 2H), 7.02 (dd, 1H), 5.24 (s, 2H), 4.88 (d, 2H), 4.68 (d, 2H), 2.65 (s, 3H).
582 - En 1	1H NMR (400 MHz, DMSO) δ = 8.70 (s, 1H), 7.85 (d, 1H), 7.79 (d, 1H), 7.55-7.47 (m, 2H), 7.22 (d, 2H), 7.05 (dd, 1H), 5.50 (s, 2H), 4.88 (d, 2H), 4.69 (d, 2H), 2.66 (s, 3H).
583 - En 1	1H NMR (400 MHz, DMSO) δ = 9.11-9.05 (m, 1H), 9.00 (s, 1H), 7.40 (dd, 1H), 7.34-7.24 (m, 2H), 7.03 (s, 1H), 5.39 (s, 2H), 4.87 (d, 2H), 4.60 (d, 2H), 2.57 (s, 3H), 2.40 (s, 3H).
584 - En 1	1H NMR (500 MHz, DMSO) δ = 8.74 (s, 1H), 8.67 (dd, 1H), 8.14 (d, 1H), 7.75-7.39 (m, 3H), 7.36-6.91 (m, 4H), 5.37 (s, 2H), 4.88 (d, 2H), 4.67 (d, 2H), 2.67 (s, 3H).
585 - En 1	1H NMR (500 MHz, DMSO) δ = 11.78 (s, 1H), 8.78 (s, 1H), 7.61 (dd, 1H), 7.51-7.43 (m, 2H), 7.39 (dd, 1H), 7.28 (s, 1H), 7.17 (s, 1H), 6.97 (dd, 1H), 6.23 (t, 1H), 4.93 (s, 2H), 4.87 (d, 2H), 4.68 (d, 2H), 2.65 (s, 3H).
586 - En 1	1H NMR (500 MHz, DMSO) δ = 8.70 (s, 1H), 8.16 (dd, 1H), 7.85 (dd, 1H), 7.52-7.43 (m, 2H), 7.23 (s, 1H), 7.20 (s, 1H), 7.02 (ddd, 2H), 5.10 (s, 2H), 4.87 (d, 2H), 4.67 (d, 2H), 3.92 (s, 3H), 2.67 (s, 3H).
587 - En 1	1H NMR (400 MHz, DMSO) δ = 9.00 (s, 1H), 8.72 (s, 1H), 7.68-7.60 (m, 2H), 7.25 (d, 2H), 5.42 (s, 2H), 4.88 (d, 2H), 4.69 (d, 2H), 2.67 (s, 3H), 2.42 (s, 3H).

-continued

cpd Nr	NMR Data
588	1H NMR (500 MHz, DMSO) δ = 9.00 (s, 1H), 8.78 (s, 1H), 7.29 (d, 1H), 7.22 (s, 1H), 7.05 (dd, 1H), 5.36 (s, 2H), 4.87 (d, 2H), 4.68 (d, 2H), 2.69 (s, 3H), 2.43 (s, 3H).
589	1H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 7.49 (d, 1H), 7.44 (s, 2H), 7.13 (d, 1H), 6.97 (dd, 1H), 5.35 (s, 2H), 4.84 (d, 2H), 4.76 (d, 2H), 2.90 (s, 3H), 2.47 (s, 3H), 2.43 (s, 3H).
590 - En 1	1H NMR (500 MHz, DMSO) δ = 7.69 (s, 1H), 7.59 (s, 1H), 7.49 (d, 1H), 7.34 (d, 1H), 7.08 (d, 2H), 6.98 (dd, 1H), 5.32 (s, 2H), 4.08 (dd, 1H), 3.83 (d, 1H), 3.65 (d, 1H), 3.43 (td, 1H), 2.63 (d, 6H), 2.34 (d, 1H), 1.95 (td, 1H), 1.76 (d, 1H).
591 - En 1	1H NMR (500 MHz, DMSO) δ = 8.41 (dd, 1H), 8.24 (s, 1H), 7.81 (dd, 1H), 7.49 (d, 1H), 7.25 (dd, 1H), 7.13 (s, 1H), 7.09 (s, 1H), 7.02 (dd, 1H), 5.16 (s, 2H), 4.18 (d, 1H), 3.94 (d, 1H), 3.90-3.79 (m, 2H), 2.62 (s, 3H), 2.53 (s, 3H), 2.41-2.28 (m, 2H).
592 - En 1	1H NMR (500 MHz, DMSO) δ = 8.22 (s, 1H), 7.70 (s, 1H), 7.47 (d, 1H), 7.31 (d, 1H), 7.12 (s, 1H), 7.08 (s, 1H), 5.33 (s, 2H), 4.18 (d, 1H), 3.94 (d, 1H), 3.86 (dd, 2H), 2.62 (d, 6H), 2.36 (hept, 2H).
593 - En 1	1H NMR (500 MHz, DMSO) δ = 8.99 (s, 1H), 7.48 (d, 1H), 7.33 (d, 1H), 7.14 (s, 1H), 7.08 (s, 1H), 6.98 (dd, 1H), 5.33 (s, 2H), 4.19 (d, 1H), 3.94 (d, 1H), 3.85 (dd, 2H), 2.62 (s, 3H), 2.42 (s, 3H), 2.41-2.28 (m, 2H).
594 - En 1	1H NMR (500 MHz, DMSO) δ = 8.54 (d, 1H), 8.24 (s, 1H), 7.77 (dd, 1H), 7.47 (d, 1H), 7.28 (d, 1H), 7.13 (s, 1H), 7.08 (s, 1H), 6.98 (dd, 1H), 5.13 (s, 2H), 4.18 (d, 1H), 3.94 (d, 1H), 3.85 (dd, 2H), 2.62 (s, 3H), 2.47 (s, 3H), 2.41-2.28 (m, 2H).
595 - En 1	1H NMR (400 MHz, DMSO) δ = 8.58 (ddd, 1H), 8.22 (s, 1H), 7.84 (td, 1H), 7.59-7.45 (m, 2H), 7.39-7.30 (m, 2H), 7.10 (d, 2H), 7.01 (dd, 1H), 5.22 (s, 2H), 4.19 (d, 1H), 3.93 (d, 1H), 3.89-3.79 (m, 2H), 2.61 (s, 3H), 2.42-2.26 (m, 2H), 1.25 (dd, 0H).
596 - En 1	1H NMR (500 MHz, DMSO) δ = 8.26 (s, 1H), 7.58 (dd, 1H), 7.46 (d, 1H), 7.38 (dd, 1H), 7.32 (d, 1H), 7.17 (s, 1H), 7.05 (s, 1H), 6.95 (dd, 1H), 6.22 (t, 1H), 4.91 (s, 2H), 4.20 (d, 1H), 3.93 (d, 1H), 3.84 (t, 2H), 2.61 (s, 3H), 2.37-2.31 (m, 1H).
597 - En 1	1H NMR (500 MHz, DMSO) δ = 8.25 (s, 1H), 8.16 (dd, 1H), 7.83 (dd, 1H), 7.48 (d, 1H), 7.13 (s, 1H), 7.08 (s, 1H), 7.03 (dd, 1H), 6.98 (dd, 1H), 5.08 (s, 2H), 4.18 (d, 1H), 3.93 (d, 1H), 3.93 (s, 3H), 3.90-3.78 (m, 2H), 2.62 (s, 3H), 2.38-2.27 (m, 2H).
598 - En 1	1H NMR (500 MHz, DMSO) δ = 8.27 (s, 1H), 7.53-7.45 (m, 2H), 7.36 (d, 1H), 7.13 (s, 1H), 7.08 (s, 1H), 7.00 (dd, 1H), 5.19 (s, 2H), 4.20 (d, 1H), 3.94 (d, 1H), 3.85 (dd, 2H), 3.83 (s, 3H), 2.62 (s, 3H), 2.41-2.29 (m, 2H).
599	1H NMR (500 MHz, DMSO) δ = 8.99 (s, 1H), 7.53 (d, 1H), 7.18 (d, 1H), 7.00 (dd, 1H), 5.37 (s, 2H), 4.98-4.92 (m, 2H), 4.91-4.85 (m, 2H), 2.93 (s, 3H), 2.54 (s, 3H), 2.43 (s, 3H).
600 - En 1	1H NMR (400 MHz, DMSO) δ = 8.99 (s, 1H), 7.95 (d, 1H), 7.68 (d, 1H), 7.49 (d, 1H), 7.39 (d, 1H), 6.98 (dd, 1H), 5.32 (s, 2H), 5.04 (s, 1H), 4.49 (dt, 1H), 3.72 (d, 2H), 2.63 (d, 6H), 2.42 (s, 3H).
601 - En 1	1H NMR (500 MHz, DMSO) δ = 8.58 (dt, 1H), 7.94 (d, 1H), 7.84 (td, 1H), 7.67 (d, 1H), 7.57 (d, 1H), 7.49 (d, 1H), 7.40 (d, 1H), 7.35 (ddd, 1H), 7.01 (dd, 1H), 5.20 (s, 2H), 5.03 (t, 1H), 4.48 (dt, 1H), 3.72 (t, 2H), 2.63 (d, 6H).
602	1H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 7.93 (t, 1H), 7.48 (d, 1H), 7.30 (d, 1H), 6.97 (dd, 1H), 5.31 (s, 2H), 4.75 (t, 1H), 3.21 (t, 4H), 2.61 (s, 3H), 2.41 (s, 3H), 0.88 (s, 6H).
603	1H NMR (400 MHz, DMSO) δ = 8.96 (s, 1H), 7.93 (s, 1H), 7.30 (d, 1H), 7.08 (d, 1H), 5.28 (s, 2H), 4.83 (t, 1H), 3.74-3.51 (m, 6H), 2.49 (s, 2H), 2.38 (s, 3H), 2.25 (s, 3H), 2.12 (d, 2H), 1.68-1.55 (m, 2H).
604	1H NMR (500 MHz, DMSO) δ = 8.16 (s, 1H), 7.45 (d, 1H), 7.39 (s, 1H), 7.28 (d, 1H), 6.95 (dd, 1H), 5.15 (s, 2H), 4.90 (s, 1H), 4.05 (s, 3H), 3.70 (dt, 2H), 3.67-3.58 (m, 4H), 2.59 (s, 3H), 2.19 (d, 2H), 1.63 (ddd, 2H).
605	1H NMR (500 MHz, DMSO) δ = 7.80 (s, 1H), 7.47 (d, 1H), 7.41 (s, 1H), 7.29 (d, 1H), 6.99 (dd, 1H), 5.30 (s, 2H), 4.93-4.87 (m, 1H), 4.06 (s, 3H), 3.66 (dddd, 6H), 2.61 (s, 3H), 2.18 (d, 2H), 1.63 (ddd, 2H).
606	1H NMR (400 MHz, DMSO) δ = 8.46 (s, 1H), 7.44 (d, 1H), 7.37 (s, 1H), 7.29 (d, 1H), 6.95 (dd, 1H), 5.07 (s, 2H), 4.88 (t, 1H), 3.86 (s, 3H), 3.74-3.63 (m, 4H), 3.67-3.58 (m, 2H), 2.59 (s, 3H), 2.19 (d, 2H), 1.63 (ddd, 2H).
607	1H NMR (500 MHz, DMSO) δ = 7.64 (s, 1H), 7.45 (d, 1H), 7.40 (s, 1H), 7.27 (d, 1H), 7.00 (d, 1H), 6.96 (dd, 1H), 5.10 (s, 2H), 4.90 (t, 1H), 3.74-3.68 (m, 2H), 3.68-3.58 (m, 7H), 2.60 (s, 3H), 2.18 (d, 2H), 1.68-1.58 (m, 2H).
608	1H NMR (400 MHz, DMSO) δ = 7.46 (d, 1H), 7.40 (s, 1H), 7.37 (d, 1H), 7.28 (d, 1H), 6.97 (dd, 1H), 6.37 (d, 1H), 5.18 (s, 2H), 4.90 (t, 1H), 3.85 (s, 3H), 3.76-3.56 (m, 6H), 2.60 (s, 3H), 2.18 (d, 2H), 1.70-1.57 (m, 2H).
609	1H NMR (400 MHz, DMSO) δ = 7.71 (dd, 1H), 7.53 (dd, 1H), 7.49-7.38 (m, 2H), 7.21 (d, 1H), 6.94 (dd, 1H), 6.25 (t, 1H), 4.92 (s, 2H), 4.87 (t, 1H), 3.73-3.52 (m, 6H), 3.48 (s, 3H), 2.60 (s, 3H), 2.16 (d, 2H), 1.68-1.55 (m, 2H).
610	1H NMR (400 MHz, DMSO) δ = 7.49 (d, 1H), 7.44-7.35 (m, 2H), 7.29 (d, 1H), 7.02 (dd, 1H), 6.47-6.38 (m, 2H), 5.14 (s, 2H), 4.90 (t, 1H), 3.75-3.55 (m, 6H), 3.49 (s, 3H), 2.61 (s, 3H), 2.18 (d, 2H), 1.63 (ddd, 2H).
611	1H NMR (500 MHz, DMSO) δ = 8.41 (dd, 1H), 7.81 (dd, 1H), 7.47 (d, 1H), 7.40 (s, 1H), 7.29 (d, 1H), 7.24 (dd, 1H), 6.99 (dd, 1H), 5.15 (s, 2H), 4.89 (s, 1H), 3.74-3.56 (m, 6H), 2.60 (s, 3H), 2.53 (s, 3H), 2.17 (d, 2H), 1.63 (ddd, 2H).
612	1H NMR (400 MHz, DMSO) δ = 8.70 (d, 1H), 7.48 (d, 1H), 7.43-7.37 (m, 2H), 7.21 (d, 1H), 7.00 (dd, 1H), 5.19 (s, 2H), 4.87 (t, 1H), 3.73-3.61 (m, 4H), 3.56 (td, 2H), 2.61 (d, 6H), 2.15 (d, 2H), 1.62 (ddd, 2H).
613	1H NMR (400 MHz, DMSO) δ = 7.68 (s, 1H), 7.45 (d, 1H), 7.40 (s, 1H), 7.24 (d, 1H), 6.94 (dd, 1H), 5.31 (s, 2H), 4.89 (t, 1H), 3.75-3.56 (m, 6H), 2.61 (d, 6H), 2.18 (d, 2H), 1.63 (s, 1H).
614	1H NMR (400 MHz, DMSO) δ = 11.75 (s, 1H), 7.56 (dd, 1H), 7.45 (d, 1H), 7.42-7.34 (m, 2H), 7.22 (d, 1H), 6.94 (dd, 1H), 6.22 (t, 1H), 4.88 (s, 3H), 3.75-3.54 (m, 6H), 2.60 (s, 3H), 2.17 (d, 2H), 1.62 (ddd, 2H).
615	1H NMR (500 MHz, DMSO) δ = 13.66 (s, 1H), 8.10 (s, 1H), 7.47-7.40 (m, 2H), 7.23 (d, 1H), 6.91 (dd, 1H), 5.04 (s, 2H), 4.89 (s, 1H), 3.70 (dt, 2H), 3.67-3.57 (m, 4H), 2.60 (s, 3H), 2.17 (d, 2H), 1.63 (ddd, 2H).

-continued

cpd Nr	NMR Data
616	¹ H NMR (400 MHz, DMSO-d ₆): δ 7.82-7.83 (m, 1H), 7.46 (d, 1H), 7.38 (s, 1H), 7.26 (d, 1H), 6.95 (dd, 1H), 5.17 (s, 2H), 4.88 (bs, 1H), 3.63-3.71 (m, 6H), 2.59 (s, 3H), 2.08 (d, 2H), 1.86 (s, 3H), 1.63-1.65 (m, 2H).
617	¹ H NMR (400 MHz, DMSO) δ = 7.48 (d, 1H), 7.40 (s, 1H), 7.32-7.25 (m, 2H), 6.99 (dd, 1H), 5.39 (s, 2H), 4.88 (t, 1H), 3.71 (dt, 2H), 3.68-3.56 (m, 4H), 2.60 (s, 3H), 2.37 (d, 3H), 2.17 (d, 2H), 1.63 (ddd, 2H).
618	¹ H NMR (400 MHz, DMSO) δ = 7.46 (d, 1H), 7.38 (s, 1H), 7.25 (d, 1H), 6.96 (dd, 1H), 6.34 (d, 1H), 5.15 (s, 2H), 4.88 (t, 1H), 3.75-3.56 (m, 6H), 2.60 (s, 3H), 2.40 (d, 3H), 2.18 (d, 2H), 1.69-1.57 (m, 2H).
619	¹ H NMR (400 MHz, DMSO) δ = 8.67 (d, 1H), 8.55 (d, 1H), 7.47 (d, 1H), 7.40 (s, 1H), 7.26 (d, 1H), 6.99 (dd, 1H), 5.23 (s, 2H), 4.89 (s, 1H), 3.74-3.54 (m, 7H), 2.59 (s, 3H), 2.17 (d, 2H), 1.63 (ddd, 2H).
620	¹ H NMR (400 MHz, DMSO) δ = 7.52-7.43 (m, 2H), 7.38 (s, 1H), 7.27 (d, 1H), 6.98 (dd, 1H), 5.36 (s, 2H), 4.88 (t, 1H), 3.75-3.68 (m, 1H), 3.64 (qd, 5H), 2.59 (s, 3H), 2.44 (d, 3H), 2.17 (d, 2H), 1.63 (ddd, 2H).
621	¹ H NMR (400 MHz, DMSO-d ₆): δ 7.74 (d, 1H), 7.62 (d, 1H), 7.45 (b, 1H), 7.38 (s, 1H), 7.27 (d, 1H), 6.98 (dd, 1H), 5.37 (s, 2H), 4.88 (t, 1H), 3.57-3.71 (m, 6H), 2.62 (s, 3H), 2.59 (s, 3H), 2.07 (d, 2H), 1.6-1.66 (t, 2H).
622	¹ H NMR (500 MHz, DMSO) δ = 8.53 (d, 1H), 7.76 (dd, 1H), 7.45 (d, 1H), 7.39 (s, 1H), 7.31-7.23 (m, 2H), 6.95 (dd, 1H), 5.11 (s, 2H), 4.89 (t, 1H), 3.70 (dt, 2H), 3.67-3.56 (m, 4H), 2.60 (s, 3H), 2.47 (s, 3H), 2.17 (d, 2H), 1.63 (ddd, 2H).
623 - En 1	¹ H NMR (400 MHz, DMSO) δ = 8.88 (s, 1H), 7.43-7.36 (m, 2H), 7.13 (d, 1H), 6.87 (dd, 1H), 5.81 (q, 1H), 4.88 (t, 1H), 3.71 (d, 2H), 3.65 (q, 1H), 3.62 (s, 1H), 3.58 (s, 1H), 2.57 (s, 3H), 2.37 (s, 3H), 2.14 (d, 2H), 1.63 (t, 5H).
624	¹ H NMR (400 MHz, DMSO) δ = 9.22 (dd, 1H), 7.85 (dd, 1H), 7.76 (dd, 1H), 7.48 (d, 1H), 7.39 (s, 1H), 7.29 (d, 1H), 7.01 (dd, 1H), 5.43 (s, 2H), 4.89 (s, 1H), 3.75-3.55 (m, 6H), 2.60 (s, 3H), 2.17 (d, 2H), 1.63 (s, 1H).
625	¹ H NMR (400 MHz, DMSO) δ = 8.68 (d, 1H), 8.58-8.51 (m, 1H), 7.89 (dt, 1H), 7.50-7.37 (m, 3H), 7.27 (d, 1H), 6.97 (dd, 1H), 5.17 (s, 2H), 4.90 (t, 1H), 3.75-3.55 (m, 6H), 2.60 (s, 3H), 2.17 (d, 2H), 1.63 (ddd, 2H).
626	¹ H NMR (400 MHz, DMSO) δ = 8.83 (d, 2H), 7.50-7.40 (m, 2H), 7.36 (s, 1H), 7.18 (d, 1H), 6.94 (dd, 1H), 5.29 (s, 2H), 4.86 (t, 1H), 3.73-3.51 (m, 6H), 2.57 (s, 3H), 2.14 (d, 2H), 1.62 (ddd, 2H).
627	¹ H NMR (400 MHz, DMSO) δ = 9.11 (d, 1H), 7.99 (d, 1H), 7.46 (d, 1H), 7.40 (s, 1H), 7.26 (d, 1H), 6.96 (dd, 1H), 5.40 (d, 2H), 4.89 (t, 1H), 3.75-3.68 (m, 2H), 3.68-3.56 (m, 4H), 2.60 (s, 3H), 2.18 (d, 2H), 1.69-1.57 (m, 2H).
628	¹ H NMR (500 MHz, DMSO) δ = 11.46 (s, 1H), 7.47 (d, 1H), 7.41-7.33 (m, 2H), 7.21 (d, 1H), 6.95 (dd, 1H), 6.36 (d, 1H), 6.21 (dd, 1H), 4.99 (d, 2H), 4.89 (s, 1H), 3.70 (dt, 2H), 3.65 (s, 2H), 3.60 (td, 2H), 2.60 (s, 3H), 2.16 (d, 2H), 1.63 (ddd, 2H).
629	¹ H NMR (400 MHz, DMSO) δ = 8.16 (dd, 1H), 7.46 (d, 1H), 7.38 (s, 1H), 7.22 (d, 1H), 7.05 (dd, 1H), 6.97 (dd, 1H), 6.89-6.83 (m, 1H), 5.16 (s, 2H), 4.90 (s, 1H), 3.84 (s, 3H), 3.75-3.63 (m, 4H), 3.59 (td, 2H), 2.59 (s, 3H), 2.16 (d, 2H), 1.62 (ddd, 2H).
630	¹ H NMR (400 MHz, DMSO) δ = 11.71 (s, 1H), 7.45 (dd, 2H), 7.40 (s, 1H), 7.22 (d, 1H), 6.95 (dd, 1H), 6.30 (d, 1H), 4.90 (d, 3H), 3.76-3.55 (m, 6H), 2.61 (s, 3H), 2.17 (d, 2H), 1.63 (ddd, 2H), 1.23 (s, 0H).
631	¹ H NMR (400 MHz, DMSO) δ = 11.59 (s, 1H), 7.57-7.48 (m, 2H), 7.48-7.37 (m, 2H), 7.22 (d, 1H), 6.92 (dd, 1H), 6.35 (d, 1H), 4.89 (t, 1H), 4.82 (s, 2H), 3.75-3.56 (m, 6H), 2.60 (s, 3H), 2.18 (d, 2H), 1.69-1.57 (m, 2H).
632	¹ H NMR (500 MHz, DMSO) δ = 8.26 (d, 1H), 7.80 (dd, 1H), 7.45 (d, 1H), 7.40 (s, 1H), 7.25 (d, 1H), 6.94 (dd, 1H), 6.85 (d, 1H), 5.06 (s, 2H), 4.89 (t, 1H), 3.85 (s, 3H), 3.74-3.57 (m, 6H), 2.60 (s, 3H), 2.17 (d, 2H), 1.63 (ddd, 2H).
633	¹ H NMR (400 MHz, DMSO) δ = 8.68 (dd, 1H), 7.95 (t, 1H), 7.62-7.55 (m, 2H), 7.47 (d, 1H), 7.41 (s, 1H), 7.36-7.26 (m, 2H), 6.98 (dd, 1H), 5.13 (s, 2H), 4.90 (t, 1H), 3.74-3.55 (m, 5H), 2.60 (s, 3H), 2.17 (d, 2H), 1.69-1.57 (m, 2H).
634	¹ H NMR (400 MHz, DMSO) δ = 8.99 (s, 1H), 7.58-7.50 (m, 2H), 7.01 (dd, 1H), 5.33 (s, 2H), 4.26 (s, 2H), 3.75-3.63 (m, 2H), 3.58 (dt, 2H), 2.75 (s, 3H), 2.42 (s, 3H), 2.27 (s, 3H), 1.65 (d, 3H), 1.53 (d, 2H).
635 - En 1	¹ H NMR (400 MHz, DMSO) δ = 8.06 (d, 1H), 7.48 (d, 1H), 7.27 (d, 1H), 7.18 (d, 1H), 7.02 (dd, 1H), 6.86 (d, 1H), 5.13 (s, 2H), 4.36 (ddd, 1H), 3.69 (s, 3H), 3.02-2.85 (m, 1H), 2.66 (td, 3H), 2.58 (s, 3H), 2.16-2.03 (m, 1H), 1.94 (s, 1H).
636 - En 1	¹ H NMR (500 MHz, DMSO) δ = 8.09 (d, 1H), 7.48 (d, 1H), 7.37 (d, 1H), 7.24 (d, 1H), 6.99 (dd, 1H), 6.36 (d, 1H), 5.18 (s, 2H), 3.85 (s, 3H), 2.93 (t, 2H), 2.66 (dt, 2H), 2.58 (s, 3H), 2.09 (q, 1H), 1.91 (d, 1H).
637 - En 1	¹ H NMR (400 MHz, DMSO) δ = 11.75 (s, 1H), 8.07 (d, 1H), 7.55 (dd, 1H), 7.47 (d, 1H), 7.37 (dd, 1H), 7.18 (d, 1H), 6.95 (dd, 1H), 6.22 (t, 1H), 4.96-4.80 (m, 2H), 4.35 (dd, 1H), 2.93 (s, 2H), 2.92 (d, 1H), 2.67 (q, 2H), 2.57 (s, 3H), 2.41 (s, 1H), 2.13-1.98 (m, 1H), 1.90 (d, 1H).
638 - En 1	¹ H NMR (400 MHz, DMSO) δ = 8.15 (d, 1H), 7.49 (d, 1H), 7.31 (d, 1H), 7.19 (d, 1H), 7.02 (dd, 1H), 6.86 (d, 1H), 5.14 (s, 2H), 4.62 (dt, 1H), 3.69 (s, 3H), 3.13-2.97 (m, 1H), 2.90 (t, 2H), 2.59 (s, 3H).
639 - En 1	¹ H NMR (400 MHz, DMSO) δ = 8.19 (d, 1H), 7.49 (d, 1H), 7.37 (d, 1H), 7.27 (d, 1H), 7.00 (dd, 1H), 6.36 (d, 1H), 5.19 (s, 2H), 4.63 (dt, 1H), 3.85 (s, 3H), 3.30-3.18 (m, 1H), 3.12-2.80 (m, 2H), 2.59 (s, 3H).
640 - En 1	¹ H NMR (500 MHz, DMSO) δ = 11.75 (s, 1H), 8.18 (d, 1H), 7.60-7.52 (m, 1H), 7.47 (d, 1H), 7.38 (d, 1H), 7.23 (s, 1H), 6.96 (d, 1H), 6.22 (t, 1H), 4.90 (s, 2H), 4.62 (s, 1H), 3.58 (s, 1H), 3.03 (s, 1H), 2.88 (s, 2H), 2.59 (s, 3H), 1.24 (s, 0H).
641 - En 1	¹ H NMR (500 MHz, DMSO) δ = 8.38 (d, 1H), 7.49 (d, 1H), 7.31 (d, 1H), 7.19 (s, 1H), 7.03 (dd, 1H), 6.86 (s, 1H), 5.15 (s, 2H), 4.86 (dt, 1H), 4.34 (t, 1H), 4.13 (q, 1H), 4.00-3.83 (m, 2H), 3.69 (s, 3H), 2.60 (s, 3H).

-continued

cpd Nr	NMR Data
642 - En 1	1H NMR (400 MHz, DMSO) δ = 8.41 (d, 1H), 7.50 (d, 1H), 7.37 (d, 1H), 7.27 (d, 1H), 7.01 (dd, 1H), 6.37 (d, 1H), 5.19 (s, 2H), 4.85 (dq, 1H), 4.33 (t, 1H), 4.13 (q, 1H), 3.90 (ddd, 2H), 3.85 (s, 3H), 2.60 (s, 3H).
643 - En 1	1H NMR (400 MHz, DMSO) δ = 8.39 (d, 1H), 7.69 (s, 1H), 7.49 (d, 1H), 7.24 (d, 1H), 6.98 (dd, 1H), 5.32 (s, 2H), 4.85 (dq, 1H), 4.33 (dd, 1H), 4.13 (q, 1H), 4.01-3.79 (m, 2H), 2.61 (d, 6H).
646 - En 1	1H NMR (500 MHz, DMSO) δ = 8.99 (s, 1H), 8.40 (d, 1H), 7.49 (d, 1H), 7.25 (d, 1H), 6.99 (dd, 1H), 5.32 (s, 2H), 4.85 (dp, 1H), 4.33 (dd, 1H), 4.13 (q, 1H), 3.98-3.80 (m, 2H), 2.60 (s, 3H), 2.42 (s, 3H).
647 - En 1	1H NMR (400 MHz, DMSO) δ = 8.58 (ddd, 1H), 8.39 (d, 1H), 7.84 (td, 1H), 7.60-7.44 (m, 2H), 7.35 (ddd, 1H), 7.25 (d, 1H), 7.02 (dd, 1H), 5.21 (s, 2H), 4.84 (dq, 1H), 4.39-4.26 (m, 1H), 4.13 (q, 1H), 4.00-3.78 (m, 2H), 2.59 (s, 3H).
648 - En 1	1H NMR (500 MHz, DMSO) δ = 11.76 (s, 1H), 8.39 (d, 1H), 7.56 (dd, 1H), 7.48 (d, 1H), 7.42-7.34 (m, 1H), 7.22 (d, 1H), 6.97 (dd, 1H), 6.22 (t, 1H), 4.90 (s, 2H), 4.96-4.77 (m, 1H), 4.33 (t, 1H), 4.12 (g, 1H), 3.98-3.80 (m, 2H), 2.59 (s, 3H).
649 - En 1	1H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 8.07 (s, 1H), 7.65-7.52 (m, 2H), 7.45 (d, 1H), 7.16-7.07 (m, 1H), 6.92 (dd, 1H), 5.35 (s, 2H), 2.63 (s, 3H), 2.46 (s, 2H), 2.42 (s, 3H), 1.50 (s, 6H).
650 - En 1	1H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 7.82 (s, 1H), 7.49 (d, 1H), 7.04 (s, 1H), 6.99 (dd, 1H), 6.96 (s, 1H), 5.32 (s, 2H), 3.80-3.70 (m, 2H), 3.64 (td, 2H), 2.64 (s, 3H), 2.41 (s, 3H), 2.13-1.94 (m, 4H).
651 - En 1	1H NMR (400 MHz, DMSO) δ = 8.73 (dd, 1H), 8.30-8.22 (m, 1H), 7.79 (dd, 1H), 7.70 (s, 1H), 7.48 (d, 1H), 7.28 (d, 1H), 6.98 (dd, 1H), 5.36-5.30 (m, 2H), 4.75 (s, 1H), 3.61 (t, 2H), 2.58 (s, 3H), 1.87 (t, 2H), 1.42 (s, 6H).
652 - En 1	1H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 7.69 (s, 1H), 7.45 (d, 1H), 7.28 (d, 1H), 6.93 (dd, 1H), 5.31 (s, 2H), 4.78 (s, 1H), 3.62 (d, 2H), 2.57 (s, 3H), 2.42 (s, 3H), 1.88 (t, 2H), 1.43 (s, 6H).
653 - En 1	1H NMR (400 MHz, DMSO) δ = 7.53-7.43 (m, 3H), 7.20 (d, 1H), 7.00 (dd, 1H), 6.89 (d, 1H), 5.97 (s, 1H), 5.22-5.10 (m, 2H), 4.42 (qd, 1H), 4.36-4.31 (m, 1H), 4.02-3.91 (m, 2H), 3.69 (s, 3H), 3.65 (dd, 1H), 3.59 (t, 1H), 2.63 (s, 3H).
654 - En 1	1H NMR (400 MHz, DMSO) δ = 8.12 (d, 1H), 7.46 (d, 1H), 7.30 (d, 1H), 7.18 (d, 1H), 7.00 (dd, 1H), 6.86 (d, 1H), 5.37-5.32 (m, 1H), 5.15 (s, 2H), 4.25 (ddt, 2H), 4.02 (dd, 1H), 3.94 (dd, 1H), 3.69 (s, 3H), 3.71-3.63 (m, 1H), 3.56 (dd, 1H), 2.56 (s, 3H).
655 - En 1	1H NMR (400 MHz, DMSO) δ = 7.49 (d, 2H), 7.38 (dd, 2H), 7.00 (dd, 1H), 6.39 (d, 1H), 5.61 (d, 1H), 5.18 (s, 2H), 4.41 (qd, 1H), 4.35-4.26 (m, 1H), 4.02-3.90 (m, 2H), 3.85 (s, 3H), 3.64 (dd, 1H), 3.59 (t, 1H), 2.64 (s, 3H).
656 - En 1	1H NMR (400 MHz, DMSO) δ = 8.14 (d, 1H), 7.47 (d, 1H), 7.37 (d, 1H), 7.25 (d, 1H), 6.98 (dd, 1H), 6.37 (d, 1H), 5.31 (d, 1H), 5.19 (s, 2H), 4.24 (s, 1H), 4.02 (dd, 1H), 3.92 (dd, 1H), 3.66 (dd, 1H), 3.56 (dd, 1H), 2.56 (s, 3H).
657 - En 1	1H NMR (400 MHz, DMSO) δ = 11.25 (d, 1H), 8.75 (d, 2H), 8.47 (s, 1H), 8.28 (d, 1H), 7.99 (d, 1H), 7.80 (dd, 1H), 7.63 (d, 1H), 7.27 (dd, 1H), 5.36 (s, 2H), 2.96 (d, 2H), 2.72 (d, 1H), 2.38-2.27 (m, 2H), 1.93 (s, 1H), 1.61 (d, 1H), 1.24 (s, 1H).
658 - En 1	1H NMR (400 MHz, DMSO) δ = 8.99 (s, 1H), 8.68 (s, 1H), 8.39 (s, 1H), 7.97 (d, 1H), 7.57 (d, 1H), 7.24-7.14 (m, 1H), 5.35 (s, 2H), 4.77 (s, 1H), 3.76-3.54 (m, 6H), 2.45 (s, 3H), 2.08 (d, 2H), 1.71-1.58 (m, 2H).
659 - En 1	1H NMR (400 MHz, DMSO) δ = 8.72 (dd, 1H), 8.23 (dd, 1H), 7.77 (dd, 1H), 7.50 (d, 1H), 6.99 (dd, 1H), 6.90 (d, 1H), 5.32 (s, 2H), 4.75 (s, 1H), 4.27-4.17 (m, 1H), 3.43 (s, 3H), 2.40 (s, 3H), 2.11 (s, 3H), 1.98 (s, 2H), 1.59 (s, 1H), 1.47 (s, 1H).
660 - En 1	1H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 7.50-7.42 (m, 1H), 6.99-6.90 (m, 2H), 5.31 (s, 2H), 4.75 (s, 1H), 4.24 (s, 1H), 3.46 (s, 4H), 2.40 (d, 6H), 2.14 (h, 2H), 2.00 (s, 2H), 1.60 (s, 1H), 1.50 (s, 1H).
661 - En 1	1H NMR (400 MHz, DMSO) δ = 8.98 (s, 1H), 8.01 (t, 1H), 7.82 (d, 1H), 7.50 (d, 1H), 7.44 (d, 1H), 6.99 (dd, 1H), 5.33 (d, 2H), 4.69-4.59 (m, 1H), 3.29-3.21 (m, 1H), 3.15-3.06 (m, 1H), 2.66 (s, 3H), 2.42 (s, 3H), 2.04-1.90 (m, 2H), 1.84-1.69 (m, 2H), 1.52 (q, 1H), 1.25 (q, 1H).
662 - En 1	1H NMR (500 MHz, DMSO) δ = 8.70 (s, 1H), 8.55 (d, 1H), 7.90 (d, 1H), 7.63 (d, 1H), 7.49 (d, 2H), 7.43 (dd, 2H), 7.19 (s, 1H), 7.00 (dd, 1H), 5.18 (s, 2H), 5.03 (t, 1H), 4.48 (dd, 1H), 3.74 (dt, 2H), 2.64 (s, 3H).
663 - En 1	1H NMR (500 MHz, DMSO) δ = 8.23-8.12 (m, 2H), 7.81 (dd, 1H), 7.48 (d, 1H), 7.23 (d, 1H), 7.00 (ddd, 2H), 5.08 (s, 2H), 4.61 (dt, 1H), 3.93 (s, 3H), 3.26 (dd, 1H), 3.00 (dt, 1H), 2.85 (dd, 1H), 2.59 (s, 3H).
664 - En 1	1H NMR (400 MHz, DMSO) δ = 8.73 (dd, 1H), 8.43 (d, 1H), 8.26 (d, 1H), 7.79 (dd, 1H), 7.52 (d, 1H), 7.25 (d, 1H), 7.02 (dd, 1H), 5.33 (s, 2H), 4.85 (dt, 1H), 4.32 (t, 1H), 4.13 (q, 1H), 3.98-3.77 (m, 2H), 2.60 (s, 3H).
665 - En 1	1H NMR (500 MHz, DMSO) δ = 8.41 (d, 1H), 8.16 (dd, 1H), 7.81 (dd, 1H), 7.49 (d, 1H), 7.24 (d, 1H), 7.01 (ddd, 2H), 5.08 (s, 2H), 4.85 (dp, 1H), 4.32 (dd, 1H), 4.13 (q, 1H), 3.93 (s, 3H), 3.97-3.79 (m, 2H), 2.60 (s, 3H).
666 - En 1	1H NMR (400 MHz, DMSO) δ = 8.99 (s, 1H), 8.57 (t, 1H), 7.50 (d, 1H), 7.29 (d, 1H), 7.00 (dd, 1H), 5.33 (s, 2H), 4.19-4.03 (m, 2H), 2.61 (s, 3H), 2.42 (s, 3H).
668 - En 1	1H NMR (400 MHz, DMSO) δ = 7.66 (d, 1H), 7.57 (bs, 1H), 7.51 (d, 1H), 7.45 (d, 2H), 7.18 (bs, 1H), 7.02 (dd, 1H), 5.16 (s, 2H), 5.01 (t, 1H), 4.48-4.46 (m, 1H), 3.88 (s, 3H), 3.76-3.73 (m, 2H), 2.64 (s, 3H).
670 - En 1	1H NMR (400 MHz, DMSO) δ = 7.65 (d, 1H), 7.62 (s, 1H), 7.55 (d, 2H), 7.18 (bs, 1H), 7.01-6.96 (m, 1H), 5.17 (s, 2H), 5.02 (t, 1H), 4.69-4.63 (m, 1H), 4.52-4.40 (m, 1H), 3.77-3.73 (m, 2H), 2.64 (s, 3H), 1.40 (d, 6H).
672 - En 1	1H NMR (400 MHz, DMSO) δ = 7.65 (d, 1H), 7.62 (s, 1H), 7.55 (d, 2H), 7.18 (bs, 1H), 7.01-6.96 (m, 1H), 5.17 (s, 2H), 5.02 (t, 1H), 4.69-4.63 (m, 1H), 4.52-4.40 (m, 1H), 3.77-3.73 (m, 2H), 2.64 (s, 3H), 1.40 (d, 6H).
674 - En 1	1H NMR (500 MHz, DMSO) δ = 7.63 (d, 1H), 7.52-7.48 (m, 2H), 7.42 (d, 1H), 7.19 (bs, 1H), 6.99 (dd, 1H), 6.51 (s, 1H), 5.25 (s, 2H), 5.03 (t, 1H), 4.50-4.45 (m, 1H), 3.76-3.73 (m, 2H), 2.63 (s, 3H), 2.23 (s, 3H).

-continued

cpd Nr	NMR Data
676	¹ H NMR (500 MHz, DMSO) δ = 7.61 (d, 1H), 7.52-7.40 (m, 4H), 7.19 (s, 1H), 7.00-6.94 (m, 1H), 6.37 (d, 1H), 5.20 (m, 2H), 5.02 (t, 1H), 4.65-4.58 (m, 1H), 4.52-4.44 (m, 2H), 3.80-3.70 (m, 2H), 2.64 (s, 3H), 1.40 (d, 6H).
678	¹ H NMR (500 MHz, DMSO): δ 7.62 (d, 1H), 7.50-7.46 (m, 2H), 7.40 (d, 1H), 7.18 (s, 1H), 6.96 (dd, 1H), 5.07 (s, 2H), 5.01 (t, 1H), 4.50-4.45 (m, 1H), 3.79-3.70 (m, 2H), 2.64 (s, 3H), 2.36 (s, 3H), 2.09 (s, 3H).
680	¹ H NMR (400 MHz, DMSO): δ 8.28 (s, 1H), 7.62 (d, 1H), 7.49-7.46 (m, 2H), 7.41 (d, 1H), 7.18 (s, 1H), 6.97 (dd, 1H), 5.15 (s, 2H), 5.01 (t, 1H), 4.49-4.45 (m, 1H), 3.76-3.72 (m, 2H), 2.64 (s, 3H), 2.16 (s, 3H).
682	¹ H NMR (500 MHz, DMSO): δ 7.76 (s, 1H), 7.61 (d, 1H), 7.50-7.47 (m, 2H), 7.42 (d, 1H), 7.19 (s, 1H), 6.98 (dd, 1H), 5.33 (s, 2H), 5.02 (t, 1H), 4.51-4.45 (m, 1H), 3.79-3.70 (m, 2H), 3.29-3.21 (m, 1H), 2.63 (s, 3H), 1.31 (d, 6H).
683	¹ H NMR (500 MHz, DMSO): δ 8.48 (s, 1H), 7.68 (s, 1H), 7.47 (d, 1H), 7.22 (d, 1H), 6.96 (dd, 1H), 6.34 (t, 1H), 5.32 (s, 2H), 2.63 (s, 3H), 2.58 (s, 3H), 2.49-2.39 (m, 2H), 2.37-2.28 (m, 2H), 2.05-1.97 (m, 1H), 1.90-1.80 (m, 1H).
684	¹ H NMR (500 MHz, DMSO): δ 7.62 (d, 1H), 7.54-7.46 (m, 3H), 7.42 (d, 1H), 7.19 (s, 1H), 7.01 (dd, 1H), 5.38 (s, 1H), 5.01 (t, 1H), 4.50-4.45 (m, 1H), 3.78-3.70 (m, 2H), 2.64 (s, 3H), 2.44 (s, 3H).
685	¹ H NMR (400 MHz, DMSO-d ₆): δ 7.88 (bs, 1H), 7.52-7.36 (m, 4H), 7.15 (bs, 1H), 6.94-6.90 (m, 1H), 5.44 (s, 1H), 5.07-5.01 (m, 2H), 4.40-4.35 (m, 1H), 4.17-4.12 (m, 1H), 2.65 (s, 3H), 1.15 (d, 3H).
686	¹ H NMR (500 MHz, DMSO): δ 7.65 (d, 1H), 7.47 (d, 1H), 7.40 (d, 1H), 6.96 (dd, 1H), 6.33 (d, 1H), 5.01 (s, 2H), 4.48 (t, 1H), 3.82 (s, 3H), 3.77-3.73 (m, 2H), 2.63 (s, 3H).
688	¹ H NMR (400 MHz, DMSO): δ 7.62 (d, 1H), 7.57 (d, 1H), 7.50-7.49 (m, 2H), 7.42 (d, 1H), 7.19 (bs, 1H), 6.99-6.96 (dd, 1H), 6.53 (d, 1H), 5.25 (s, 2H), 5.19-5.12 (m, 2H), 5.02 (bs, 1H), 4.50-4.46 (m, 1H), 3.74 (d, 2H), 2.64 (s, 2H).
690	¹ H NMR (500 MHz, DMSO): δ 7.63 (d, 1H), 7.51-7.43 (m, 3H), 7.40 (d, 1H), 7.20 (bs, 1H), 6.98 (dd, 1H), 6.41 (s, 1H), 5.19 (s, 2H), 5.02 (t, 1H), 4.51-4.45 (m, 1H), 4.01 (d, 2H), 3.77-3.72 (m, 2H), 2.64 (s, 3H), 1.30-1.25 (m, 1H), 0.50-0.45 (m, 2H), 0.38-0.33 (m, 2H).
692	¹ H NMR (500 MHz, DMSO): δ 8.10 (s, 1H), 7.62 (d, 1H), 7.50 (s, 1H), 7.47 (d, 1H), 7.39 (d, 1H), 7.21 (s, 1H), 6.92 (dd, 1H), 5.05-5.00 (m, 3H), 4.51-4.47 (m, 1H), 3.91 (s, 3H), 3.78-3.73 (m, 2H), 2.63 (s, 3H).
694	¹ H NMR (500 MHz, DMSO): δ 9.24 (s, 1H), 7.67 (d, 1H), 7.52-7.41 (m, 3H), 7.17 (s, 1H), 7.02 (dd, 1H), 5.53 (s, 2H), 5.01 (s, 1H), 4.49-4.45 (m, 1H), 3.77-3.72 (m, 2H), 2.64 (s, 3H).
695	¹ H NMR (500 MHz, DMSO): δ 7.85 (d, 1H), 7.78 (d, 1H), 7.64 (d, 1H), 7.53-7.43 (m, 3H), 7.19 (bs, 1H), 7.05-7.02 (dd, 1H), 5.46 (s, 2H), 5.01 (t, 1H), 4.50-4.45 (m, 1H), 3.76-3.73 (m, 2H), 2.64 (s, 3H).
696	¹ H-NMR (500 MHz, DMSO): δ 8.58 (d, 1H), 7.80-7.65 (m, 1H), 7.67-7.62 (m, 1H), 7.50-7.43 (m, 1H), 7.39 (d, 1H), 7.17 (s, 1H), 7.01 (dd, 1H), 5.19 (s, 2H), 5.09-5.06 (m, 1H), 4.39-4.34 (m, 1H), 4.16-4.11 (m, 1H), 2.64 (s, 3H), 1.13 (d, 3H).
697	¹ H NMR (500 MHz, DMSO): δ 9.11 (s, 1H), 8.02 (s, 1H), 7.62 (d, 1H), 7.51-7.47 (m, 2H), 7.43 (d, 1H), 7.20 (bs, 1H), 6.99 (dd, 1H), 5.42 (s, 2H), 5.02 (t, 1H), 4.50-4.46 (m, 1H), 3.77-3.72 (m, 2H), 2.63 (s, 3H).
698	¹ H NMR (400 MHz, DMSO): δ 7.81 (s, 1H), 7.59 (d, 1H), 7.54 (bs, 2H), 7.46 (d, 1H), 7.39 (d, 1H), 7.20 (bs, 1H), 6.92 (dd, 1H), 5.04 (bs, 1H), 4.97 (s, 2H), 4.52-4.45 (m, 1H), 3.81 (s, 3H), 3.77-3.73 (m, 2H), 2.63 (s, 3H).
700	¹ H NMR (500 MHz, DMSO): δ 8.41 (d, 1H), 7.65-7.58 (m, 2H), 7.49-7.40 (m, 4H), 7.18 (s, 1H), 6.98 (dd, 1H), 5.15 (s, 2H), 5.01 (t, 1H), 4.49-4.45 (m, 1H), 3.76-3.73 (m, 2H), 2.63 (s, 3H), 2.30 (s, 3H).
702	¹ H NMR (500 MHz, DMSO): δ 8.38 (dd, 1H), 7.79 (dd, 1H), 7.64 (d, 1H), 7.52-7.46 (d, 3H), 7.19-7.13 (m, 2H), 7.02 (dd, 1H), 5.28 (s, 2H), 5.03 (t, 1H), 4.50-4.61 (m, 1H), 3.76-3.73 (m, 2H), 2.64 (s, 3H), 2.28-2.22 (m, 1H), 1.02-0.95 (m, 4H).
704	¹ H NMR (500 MHz, DMSO): δ 8.37 (d, 1H), 7.65 (d, 1H), 7.48-7.41 (m, 4H), 7.31-7.28 (m, 1H), 7.18 (bs, 1H), 7.00 (dd, 1H), 5.32 (s, 1H), 5.07 (bs, 1H), 4.47 (d, 1H), 3.76 (bs, 2H), 2.66 (s, 3H), 2.15 (t, 1H), 0.96 (d, 2H), 0.74 (d, 2H).
705	¹ H NMR (500 MHz, DMSO): δ 8.36 (d, 1H), 7.47-7.38 (m, 3H), 7.32-7.25 (m, 2H), 7.0-6.90 (m, 1H), 5.32 (s, 2H), 4.92 (bs, 1H), 3.73-3.59 (m, 6H), 2.58 (s, 3H), 2.20-2.10 (m, 3H), 1.68-1.58 (m, 2H), 1.00-0.96 (m, 2H), 0.72-0.70 (m, 2H).
706	¹ H NMR (500 MHz, DMSO): δ 8.10-8.07 (m, 1H), 7.64-7.57 (m, 2H), 7.50-7.47 (m, 2H), 7.42 (d, 1H), 7.19 (s, 1H), 6.98 (dd, 1H), 6.69-6.67 (m, 1H), 5.01 (t, 1H), 4.94 (s, 2H), 4.50-4.45 (m, 1H), 4.07 (t, 4H), 3.77-3.73 (m, 2H), 2.64 (s, 3H), 2.26-2.18 (m, 2H).
708	¹ H NMR (500 MHz, DMSO): δ 8.80 (s, 2H), 7.64 (d, 1H), 7.51-7.47 (d, 2H), 7.43 (d, 1H), 7.19 (brs, 1H), 7.01 (dd, 1H), 5.20 (s, 2H), 5.03 (t, 1H), 4.49-4.46 (m, 1H), 3.78-3.71 (m, 2H), 2.64 (s, 3H), 2.62 (s, 3H).
710	¹ H NMR (400 MHz, DMSO): δ 8.18 (dd, 1H), 7.82 (dd, 1H), 7.61 (d, 1H), 7.47 (d, 2H), 7.38 (d, 1H), 7.16 (bs, 1H), 7.07-7.02 (m, 1H), 6.97-6.92 (m, 1H), 4.99 (t, 3H), 4.51-4.44 (m, 1H), 4.36-4.31 (m, 1H), 3.74 (t, 2H), 2.64 (s, 3H), 0.76-0.62 (m, 4H).
711	¹ H NMR (400 MHz, DMSO): δ 8.17 (dd, 1H), 7.80 (dd, 1H), 7.45 (d, 2H), 7.37 (bs, 1H), 7.20 (d, 1H), 7.07-7.02 (m, 1H), 6.94 (dd, 1H), 5.00 (s, 2H), 4.85 (s, 1H), 3.72-3.56 (m, 6H), 2.59 (s, 3H), 2.16 (d, 2H), 1.68-1.58 (m, 2H), 0.78-0.64 (m, 4H).
712	¹ H NMR (500 MHz, DMSO): δ 8.83 (d, 2H), 7.60 (d, 1H), 7.48-7.46 (m, 3H), 7.38 (d, 1H), 7.17 (brs, 1H), 6.96 (dd, 1H), 5.28 (s, 2H), 5.00 (t, 1H), 4.48-4.44 (m, 1H), 3.74-3.72 (m, 2H), 2.63 (s, 3H).
714	¹ H NMR (500 MHz, DMSO): δ 8.84 (d, 1H), 8.67 (d, 1H), 8.63 (d, 1H), 7.61 (d, 1H), 7.50 (d, 1H), 7.47-7.44 (m, 2H), 7.15 (bs, 1H), 7.04 (dd, 1H), 5.28 (s, 2H), 4.99 (t, 1H), 4.49-4.46 (m, 1H), 3.76-3.72 (m, 2H), 2.64 (s, 3H).
715	¹ H NMR (500 MHz, DMSO): δ 8.15 (dd, 1H), 7.86-7.81 (m, 1H), 7.63 (d, 1H), 7.50-7.45 (m, 2H), 7.40 (d, 1H), 7.18 (s, 1H), 7.05-6.95 (m, 2H), 5.07 (s, 2H), 5.00 (t, 1H), 4.50-4.46 (m, 1H), 3.92 (s, 3H), 3.76-3.71 (m, 2H), 2.64 (s, 3H).
716	¹ H NMR (500 MHz, DMSO): δ 8.58 (dd, 2H), 7.62 (d, 1H), 7.50-7.47 (m, 4H), 7.41 (d, 1H), 7.19 (s, 1H), 7.00 (dd, 1H), 5.21 (s, 2H), 5.02 (t, 1H), 4.49-4.45 (m, 1H), 3.75-3.74 (m, 2H), 2.63 (s, 3H).
717	¹ H NMR (500 MHz, DMSO): δ 9.87 (bs, 1H), 9.31 (bs, 1H), 8.74 (d, 1H), 8.47 (d, 1H), 8.12 (d, 1H), 7.61-7.57 (m, 1H), 7.50-7.22 (m, 3H), 7.02-6.97 (m, 1H), 5.35 (s, 2H), 4.88-4.80 (m, 1H), 3.87-3.79 (m, 2H), 3.36-3.30 (m, 1H), 3.26-3.15 (m, 1H), 2.63 (s, 3H), 2.10-2.00 (m, 2H).

-continued

cpd Nr	NMR Data
718	¹ H NMR (500 MHz, DMSO): δ 8.59-8.57 (m, 1H), 7.83 (td, 1H), 7.56 (d, 1H), 7.50 (d, 1H), 7.45-7.42 (m, 2H), 7.40-7.33 (m, 2H), 7.17 (s, 1H), 7.02 (dd, 1H), 5.19 (s, 2H), 5.06 (d, 1H), 4.36 (dd, 1H), 4.18-4.10 (m, 1H), 2.64 (s, 3H), 1.13 (d, 3H).
719 - En 1	¹ H NMR (400 MHz, DMSO): δ 8.57 (dd, 1H), 7.83 (t, 1H), 7.57-7.47 (m, 4H), 7.35 (t, 1H), 7.27 (d, 1H), 7.00 (dd, 1H), 5.24 (s, 1H), 5.19 (s, 1H), 3.74-3.62 (m, 2H), 3.31 (s, 1H), 3.11 (d, 1H), 2.59 (s, 3H), 2.21-2.15 (m, 2H), 1.83-1.74 (m, 2H).
720 - En 1	¹ H NMR (500 MHz, DMSO): δ 8.17 (dd, 1H), 7.86-7.82 (m, 1H), 7.64 (s, 1H), 7.50-7.42 (m, 3H), 7.28 (s, 1H), 7.05-7.01 (m, 1H), 6.97 (dd, 1H), 5.13-5.05 (m, 3H), 4.00-3.92 (m, 4H), 3.78-3.72 (m, 1H), 2.65 (s, 3H), 1.52 (s, 3H).
721	¹ H NMR (500 MHz, DMSO): δ 8.98 (s, 1H), 8.49 (s, 1H), 7.47 (d, 1H), 7.23 (d, 1H), 6.96 (dd, 1H), 6.34 (t, 1H), 5.32 (s, 2H), 2.58 (s, 3H), 2.47-2.41 (m, 5H), 2.39-2.30 (m, 2H), 2.05-1.95 (m, 1H), 1.90-1.82 (m, 1H).
722 - En 1	¹ H NMR (500 MHz, DMSO): δ 7.63 (s, 1H), 7.49 (d, 1H), 7.48-7.43 (m, 2H), 7.30 (s, 1H), 7.18 (s, 1H), 6.97 (dd, 1H), 5.14-5.10 (m, 3H), 4.00-3.93 (m, 1H), 3.80-3.73 (m, 1H), 2.65 (s, 3H), 2.49 (s, 3H), 1.53.
723 - En 1	¹ H NMR (500 MHz, DMSO): δ 8.98 (s, 1H), 8.29 (d, 1H), 7.48 (d, 1H), 7.22 (d, 1H), 6.97 (dd, 1H), 5.31 (s, 2H), 4.73-4.62 (m, 1H), 4.00-3.88 (m, 2H), 3.75-3.59 (m, 2H), 2.58 (s, 3H), 2.41 (s, 3H), 1.95-1.86 (m, 2H).
724 - En 1	¹ H NMR (500 MHz, DMSO): δ 8.28 (d, 1H), 7.68 (s, 1H), 7.47 (d, 1H), 7.21 (d, 1H), 6.97 (dd, 1H), 5.31 (s, 2H), 4.75-4.62 (m, 1H), 4.00-3.88 (m, 2H), 3.74-3.58 (m, 2H), 2.63 (s, 3H), 2.57 (s, 3H), 1.95-1.88 (m, 2H).
725 - En 1	¹ H NMR (500 MHz, DMSO): δ 8.23 (s, 1H), 7.47 (d, 1H), 7.32 (d, 1H), 7.16-7.06 (m, 3H), 6.97 (dd, 1H), 5.13 (s, 2H), 4.18 (d, 1H), 3.94 (d, 1H), 3.85 (t, 2H), 2.61 (s, 3H), 2.41 (s, 3H), 2.38-2.31 (m, 2H).
726 - En 1	¹ H NMR (500 MHz, DMSO): δ = 8.20 (d, 1H), 7.70 (s, 1H), 7.59 (bs, 1H), 7.49-7.46 (m, 2H), 7.35 (bs, 1H), 6.95 (dd, 1H), 5.34-5.28 (m, 2H), 4.97-4.93 (m, 1H), 3.74-3.63 (m, 2H), 3.07 (s, 3H), 2.63 (s, 3H), 2.62 (s, 3H).
727 - En 1	¹ H-NMR (500 MHz, DMSO): δ 8.26 (d, 1H), 7.70 (s, 1H), 7.47 (d, 1H), 7.35 (d, 1H), 6.95 (dd, 1H), 5.33 (s, 2H), 4.78-4.74 (m, 1H), 4.17 (s, 1H), 3.44-3.40 (m, 1H), 3.30-3.27 (m, 1H), 3.17-3.12 (m, 2H), 2.63 (s, 3H), 2.60 (s, 3H), 2.49-2.46 (s, 1H), 2.27-2.23 (m, 1H).
727 - En 3	¹ H-NMR (400 MHz, DMSO): δ 8.26 (d, 1H), 7.69 (s, 1H), 7.47 (d, 1H), 7.30 (d, 1H), 6.95 (dd, 1H), 5.32 (s, 2H), 4.70-4.64 (m, 1H), 4.07 (s, 1H), 3.45-3.40 (m, 1H), 3.30-3.27 (m, 1H), 3.18-3.10 (m, 2H), 2.63 (s, 3H), 2.59 (s, 3H), 2.46-2.41 (m, 1H), 2.33-2.26 (m, 1H).
727 - En 4	¹ H-NMR (400 MHz, DMSO): δ 8.26 (d, 1H), 7.69 (s, 1H), 7.47 (d, 1H), 7.30 (d, 1H), 6.95 (dd, 1H), 5.32 (s, 2H), 4.70-4.66 (m, 1H), 4.07 (s, 1H), 3.45-3.40 (m, 1H), 3.30-3.26 (m, 1H), 3.15-3.10 (m, 2H), 2.63 (s, 3H), 2.59 (s, 3H), 2.46-2.41 (m, 1H), 2.33-2.27 (m, 1H).
728 - En 1	¹ H NMR (500 MHz, DMSO): δ 8.98 (s, 1H), 7.55 (s, 1H), 7.44 (d, 1H), 7.26-7.20 (m, 1H), 6.93 (dd, 1H), 5.30 (s, 2H), 5.12-4.93 (m, 1H), 3.70-3.62 (m, 3H), 3.02-2.75 (m, 3H), 2.59-2.52 (m, 4H), 2.49 (s, 3H), 2.10-2.01 (m, 1H), 1.84-1.77 (m, 1H).
729 - En 1	¹ H NMR (500 MHz, DMSO): δ 8.99 (s, 1H), 8.47 (s, 1H), 8.06 (d, 1H), 7.97 (s, 1H), 7.73 (s, 1H), 7.46 (d, 1H), 7.36 (s, 1H), 7.27 (d, 1H), 6.97 (dd, 1H), 5.34-5.28 (s, 2H), 4.97-4.90 (m, 1H), 4.68-4.57 (m, 2H), 2.52 (s, 3H), 2.43 (s, 3H).
730 - En 1	¹ H NMR (400 MHz, DMSO): δ 8.08 (s, 1H), 7.78 (s, 1H), 7.51 (s, 1H), 7.45 (d, 1H), 7.23 (d, 1H), 6.91 (dd, 1H), 5.04 (s, 3H), 3.65-3.55 (m, 2H), 3.24-3.16 (m, 2H), 2.58 (s, 3H), 2.48-2.38 (m, 2H).
731 - En 1	¹ H NMR (400 MHz, DMSO): δ 8.12 (s, 1H), 7.58 (bs, 1H), 7.52 (s, 1H), 7.46 (d, 1H), 7.25 (d, 1H), 6.92 (dd, 1H), 5.05 (s, 2H), 3.73 (d, 1H), 3.63 (d, 1H), 3.29-3.24 (m, 1H), 3.15-3.10 (d, 1H), 2.59 (s, 3H), 2.27-2.14 (m, 2H), 1.87-1.75 (m, 2H).
732 - En 1	¹ H NMR (400 MHz, DMSO): δ 7.59 (brs, 1H), 7.51-7.47 (m, 2H), 7.37 (d, 1H), 7.31 (d, 1H), 7.00 (dd, 1H), 6.39 (d, 1H), 5.23 (brs, 1H), 5.18 (s, 2H), 3.85 (s, 3H), 3.75-3.72 (m, 1H), 3.65-3.62 (m, 1H), 3.28-3.25 (m, 1H), 3.15-3.12 (m, 1H), 2.60 (s, 3H), 2.23-2.2.17 (m, 2H), 1.85-1.73 (m, 2H).
733 - En 1	¹ H NMR (400 MHz, DMSO): δ 7.79 (brs, 1H), 7.51-7.47 (m, 2H), 7.37 (d, 1H), 7.29 (d, 1H), 7.00 (dd, 1H), 6.39 (d, 1H), 5.22 (brs, 1H), 5.18 (s, 2H), 3.85 (s, 3H), 3.62-3.60 (m, 2H), 3.21-3.19 (m, 2H), 2.59 (s, 3H), 2.49-2.39 (m, 2H).
734 - En 1	¹ H NMR (400 MHz, DMSO): δ 9.23-9.20 (m, 1H), 7.89-7.86 (m, 1H), 7.78-7.74 (m, 2H), 7.52-7.48 (m, 2H), 7.30 (d, 1H), 7.04 (dd, 1H), 5.42 (s, 2H), 5.25 (brs, 1H), 3.65-3.55 (m, 2H), 3.29-3.25 (m, 1H), 3.24-3.17 (m, 1H), 2.59 (s, 3H), 2.49-2.39 (m, 2H).
735 - En 1	¹ H NMR (400 MHz, DMSO): δ 8.60-7.57 (m, 1H), 7.79-7.72 (m, 2H), 7.67-7.62 (m, 1H), 7.50-7.44 (m, 2H), 7.28-7.23 (m, 1H), 7.00 (dd, 1H), 5.25-5.16 (m, 3H), 3.64-3.55 (m, 2H), 3.30-3.26 (m, 1H), 3.25-3.16 (m, 1H), 2.58 (s, 3H), 2.46-2.38 (m, 2H).
736 - En 1	¹ H NMR (400 MHz, DMSO): δ 8.67 (d, 1H), 8.13 (d, 1H), 7.78 (brs, 1H), 7.66-7.61 (m, 1H), 7.54-7.46 (m, 2H), 7.30-6.96 (m, 3H), 5.34 (s, 2H), 5.21 (brs, 1H), 3.64-3.55 (m, 2H), 3.21-3.15 (m, 2H), 2.62 (s, 3H), 2.45-2.36 (m, 2H).
737 - En 1	¹ H NMR (400 MHz, DMSO): δ 8.66 (d, 1H), 8.13 (d, 1H), 7.66-7.46 (m, 4H), 7.31-6.96 (m, 3H), 5.34 (s, 2H), 5.23 (brs, 1H), 3.75-3.70 (m, 1H), 3.65-3.60 (m, 1H), 3.25-3.20 (m, 1H), 3.18-3.10 (m, 1H), 2.60 (s, 3H), 2.25-2.12 (m, 2H), 1.90-1.70 (m, 2H).
738 - En 1	¹ H-NMR (400 MHz, DMSO): δ 8.58 (d, 1H), 7.84 (td, 1H), 7.63 (s, 1H), 7.58 (d, 1H), 7.51-7.46 (m, 2H), 7.43 (s, 1H), 7.36-7.33 (m, 1H), 7.29 (s, 1H), 7.02 (dd, 1H), 5.20 (s, 2H), 5.12 (t, 1H), 3.98-3.95 (m, 1H), 3.78-3.74 (m, 1H), 2.65 (s, 3H), 1.52 (s, 3H).
739 - En 1	¹ H-NMR (400 MHz, DMSO): δ 8.58 (d, 1H), 7.80-7.75 (m, 1H), 7.67-7.63 (m, 2H), 7.51-7.47 (m, 2H), 7.43 (s, 1H), 7.29 (s, 1H), 7.00 (dd, 1H), 5.19 (s, 2H), 5.12 (s, 1H), 3.96 (d, 1H), 3.76 (d, 1H), 2.65 (s, 3H), 1.52 (s, 3H).
740 - En 1	¹ H-NMR (500 MHz, DMSO): δ 8.25 (s, 1H), 7.53-7.45 (m, 2H), 7.34 (d, 1H), 7.13(bs, 1H), 7.08 (bs, 1H), 7.00 (dd, 1H), 6.50-6.26 (m, 2H), 5.24 (s, 2H), 4.72-4.63 (m, 2H), 4.18 (d, 1H), 3.94 (d, 1H), 3.89-3.83 (m, 2H), 2.62 (s, 3H), 2.40-2.35 (m, 2H).
741	¹ H NMR (500 MHz, DMSO): δ 8.59 (d, 1H), 7.80-7.76 (m, 1H), 7.67-7.62 (m, 2H), 7.50-7.47 (m, 2H), 7.43 (d, 1H), 7.20 (bs, 1H), 7.00 (dd, 1H), 5.20 (s, 2H), 5.03 (t, 1H), 4.49-4.45 (m, 1H), 3.76-3.73 (m, 2H), 2.63 (s, 3H).

-continued

cpd Nr	NMR Data
742	¹ H NMR (500 MHz, DMSO): δ 9.00 (s, 1H), 8.35 (d, 1H), 7.73 (d, 1H), 7.54-7.31 (m, 3H), 7.26-7.21 (m, 2H), 5.36 (s, 2H), 5.05 (t, 1H), 4.53-4.47 (m, 1H), 3.80-3.73 (m, 2H), 2.43 (s, 3H).
743	¹ H NMR (500 MHz, DMSO): δ 8.56 (d, 1H), 8.35 (d, 1H), 7.80 (d, 1H), 7.73 (d, 1H), 7.53-7.28 (m, 3H), 7.28 (d, 1H), 7.24-7.22 (m, 2H), 5.17 (s, 2H), 5.08 (bs, 1H), 3.77 (bs, 2H), 2.47 (s, 3H).
744	¹ H NMR (500 MHz, DMSO): δ 8.59 (d, 1H), 8.35 (bs, 1H), 7.86 (t, 1H), 7.73 (d, 1H), 7.68-7.03 (m, 7H), 5.24 (s, 2H), 5.10 (bs, 1H), 4.49 (bs, 1H), 3.81-3.75 (m, 2H).
745	¹ H NMR (500 MHz, DMSO): δ 8.32 (d, 1H), 7.75-7.70 (m, 2H), 7.55-7.32 (m, 3H), 7.24-7.21 (m, 2H), 5.37 (s, 2H), 5.07 (t, 2H), 4.53-4.49 (m, 1H), 3.80-3.74 (m, 2H), 2.63 (s, 3H).
746	¹ H NMR (500 MHz, DMSO): δ 8.31 (d, 1H), 7.73 (d, 1H), 7.56-7.30 (m, 3H), 7.21 (d, 3H), 5.18 (s, 2H), 5.07 (s, 1H), 4.55-4.45 (m, 1H), 3.77 (bs, 2H), 2.41 (s, 3H).
747	¹ H NMR (500 MHz, DMSO): δ 8.58 (s, 1H), 7.86-7.81 (m, 1H), 7.61-7.56 (m, 2H), 7.50 (s, 1H), 7.44-7.41 (m, 2H), 7.36-7.34 (m, 1H), 7.20 (s, 1H), 6.98 (dd, 1H), 5.19 (s, 2H), 5.04 (t, 1H), 4.51-4.47 (m, 1H), 3.80-3.71 (m, 2H), 2.77-2.70 (m, 1H), 1.16-1.07 (m, 4H).
748	¹ H NMR (500 MHz, DMSO): δ 8.98 (s, 1H), 7.60 (d, 1H), 7.49 (s, 1H), 7.44-7.40 (m, 2H), 7.19 (s, 1H), 6.95 (dd, 1H), 5.31 (s, 2H), 5.03 (t, 1H), 4.52-4.46 (m, 1H), 3.80-3.71 (m, 2H), 2.76-2.70 (m, 1H), 2.41 (s, 3H), 1.16-1.06 (m, 4H).
749	¹ H NMR (500 MHz, DMSO): δ 7.71 (s, 1H), 7.59 (d, 1H), 7.52 (s, 1H), 7.43-7.40 (m, 2H), 7.22 (s, 1H), 6.94 (dd, 1H), 5.32 (s, 2H), 5.05 (t, 1H), 4.51-4.46 (m, 1H), 3.82-3.70 (m, 2H), 2.78-2.69 (m, 1H), 2.62 (s, 3H), 1.16-1.05 (m, 4H).
750	¹ H NMR (500 MHz, DMSO): δ 8.54 (d, 1H), 7.78 (dd, 1H), 7.59 (d, 1H), 7.51 (s, 1H), 7.44-7.40 (m, 2H), 7.27 (d, 1H), 7.21 (s, 1H), 6.94 (dd, 1H), 5.12 (s, 2H), 5.04 (t, 1H), 4.51-4.47 (m, 1H), 3.80-3.72 (m, 2H), 2.76-2.71 (m, 1H), 2.46 (s, 3H), 1.15-1.07 (m, 4H).
751	¹ H NMR (500 MHz, DMSO): δ 7.59 (d, 1H), 7.51 (bs, 1H), 7.43-7.41 (m, 2H), 7.20-7.18 (m, 2H), 6.96-6.92 (dd, 1H), 5.12 (s, 1H), 5.04 (t, 1H), 4.52-4.48 (m, 2H), 3.81-3.72 (m, 2H), 2.77-2.73 (m, 1H), 2.41 (s, 3H), 1.14-1.09 (m, 4H).

In all above cases the analytical LCMS and ¹H NMR data collected for En2 matched the data for En1.
Chiral Analytical Data

Chiral SFC analysis						
Code	Column Name	co-solvent	co- solvent %	Flow [g/min]	RT [min]	Purity [%]
cpd 103-En 1	(1)	MeOH	40	4	3.23	99.0
cpd 103-En 2	(1)	MeOH	40	4	1.32	100.0
cpd 105-En 1	(2)	0.5% DEA in MeOH	25	3	3.76	99.5
cpd 105-En 2	(2)	0.5% DEA in MeOH	25	3	2.49	99.1
cpd 106-En 1	(3)	MeOH	35	3	7.81	99.2
cpd 106-En 2	(3)	MeOH	35	3	3.75	99.6
cpd 110-En 1	(4)	MeOH	40	3	2.74	99.9
cpd 110-En 2	(4)	MeOH	40	3	1.66	99.9
cpd 111-En 1	(5)	MeOH	40	3	2.85	99.8
cpd 111-En 2	(5)	MeOH	40	3	1.16	99.9
cpd 112-En 1	(5)	0.5% DEA in MeOH	40	3	1.35	95.2
cpd 112-En 2	(5)	0.5% DEA in MeOH	40	3	2.03	99.1
cpd 113-En 1	(6)	0.5% DEA in EtOH	20	3	4.59	99.9
cpd 113-En 2	(6)	0.5% DEA in EtOH	20	3	3.34	99.8
cpd 115-En 1	(6)	0.5% DEA in MeOH	40	3	6.14	99.6
cpd 115-En 2	(6)	0.5% DEA in MeOH	40	3	3.98	99.7
cpd 116-En 1	(4)	MeOH	20	3	3.58	99.1
cpd 116-En 2	(4)	MeOH	20	3	3.07	99.3
cpd 118-En 1	(7)	MeOH	15	3	3.90	99.9
cpd 118-En 2	(7)	MeOH	15	3	2.50	99.9
cpd 120-En 1	(6)	MeOH	40	3	6.10	99.5
cpd 120-En 2	(6)	MeOH	40	3	4.52	99.9
cpd 121-En 1	(5)	IPA	20	3	11.20	96.9
cpd 121-En 2	(5)	IPA	20	3	9.36	99.7
cpd 123-En 1	(4)	MeOH	30	3	1.61	99.9
cpd 123-En 2	(4)	MeOH	30	3	2.22	99.7
cpd 124-En 1	(8)	MeOH	25	3	5.77	99.9
cpd 124-En 2	(8)	MeOH	25	3	4.48	99.9
cpd 125-En 1	(6)	0.5% DEA in MeOH	30	3	3.48	99.5
cpd 125-En 2	(6)	0.5% DEA in MeOH	30	3	2.59	99.6
cpd 126-En 1	(9)	IPA	30	3	3.57	95.1
cpd 126-En 2	(9)	IPA	30	3	2.69	98.7
cpd 130-En 1	(4)	MeOH	30	3	1.85	99.9
cpd 130-En 2	(4)	MeOH	30	3	1.29	99.9
cpd 131-En 1	(7)	MeOH	10	3	3.04	99.0
cpd 131-En 2	(7)	MeOH	10	3	2.43	98.0
cpd 132-En 1	(1)	MeOH	30	3	4.50	99.3
cpd 132-En 2	(1)	MeOH	30	3	2.98	99.7

Chiral SFC analysis						
Code	Column Name	co-solvent	co-solvent %	Flow [g/min]	RT [min]	Purity [%]
cpd 134-En 1	(4)	MeOH	25	3	3.20	99.2
cpd 134-En 2	(4)	MeOH	25	3	1.48	99.5
cpd 135-En 1	(1)	MeOH	25	3	2.19	99.8
cpd 135-En 2	(1)	MeOH	25	3	3.33	99.9
cpd 141-En 1	(6)	MeOH	15	3	4.48	99.8
cpd 141-En 2	(6)	MeOH	15	3	5.61	99.9
cpd 147-Dia 1-En 1	(9)	IPA	25	3	3.24	99.6
cpd 147-Dia 1-En 2	(9)	IPA	25	3	6.21	98.3
cpd 148-En 1	(10)	MeOH	30	3	5.02	99.3
cpd 148-En 2	(10)	MeOH	30	3	4.22	99.8
cpd 151-En 1	(3)	MeOH	30	3	6.04	98.5
cpd 151-En 2	(3)	MeOH	30	3	9.85	97.9
cpd 155-En 1	(11)	MeOH	10	3	8.77	95.3
cpd 155-En 2	(11)	MeOH	10	3	9.41	100.0
cpd 157-En 1	(2)	MeOH	10	3	8.08	99.7
cpd 157-En 2	(2)	MeOH	10	3	5.55	98.9
cpd 159-En 1	(1)	MeOH	30	3	3.21	99.9
cpd 159-En 2	(1)	MeOH	30	3	4.08	99.8
cpd 160-En 1	(9)	MeOH	40	4	6.91	99.9
cpd 160-En 2	(9)	MeOH	40	4	9.94	99.6
cpd 162-Dia 1-En1	(5)	MeOH	40	3	2.01	99.7
cpd 162-Dia 1-En2	(5)	MeOH	40	3	2.53	99.5
cpd 163-En 1	(3)	MeOH	35	3	5.40	96.0
cpd 163-En 2	(3)	MeOH	35	3	3.30	98.3
cpd 164-En 1	(9)	MeOH	40	3	5.39	99.0
cpd 164-En 2	(9)	MeOH	40	3	8.72	97.6
cpd 165-En 1	(6)	0.5% DEA in MeOH	30	3	6.35	96.4
cpd 165-En 2	(6)	0.5% DEA in MeOH	30	3	4.51	97.4
cpd 169-En 1	(7)	MeOH	20	3	3.68	94.7
cpd 169-En 2	(7)	MeOH	20	3	3.10	99.6
cpd 172-En 1	(11)	MeOH	10	3	13.12	97.6
cpd 172-En 2	(11)	MeOH	10	3	16.13	99.9
cpd 176-En 1	(6)	0.5% DEA in EtOH	20	3	2.69	99.6
cpd 176-En 2	(6)	0.5% DEA in EtOH	20	3	3.87	99.5
cpd 179-En 1	(6)	0.5% DEA in EtOH	30	3	5.15	98.8
cpd 179-En 2	(6)	0.5% DEA in EtOH	30	3	3.42	99.3
cpd 180-En 1	(12)	MeOH	40	3	4.23	99.0
cpd 180-En 2	(12)	MeOH	40	3	6.03	97.7
cpd 183-En 1	(6)	0.5% DEA in EtOH	20	3	6.70	99.3
cpd 183-En 2	(6)	0.5% DEA in EtOH	20	3	3.73	97.1
cpd 184-En 1	(6)	0.5% DEA in MeOH	30	3	5.84	96.7
cpd 184-En 2	(6)	0.5% DEA in MeOH	30	3	3.89	99.9
cpd 191-En 1	(4)	MeOH	40	3	2.24	99.8
cpd 191-En 2	(4)	MeOH	40	3	0.83	99.9
cpd 195-En 1	(6)	0.5% DEA in MeOH	30	3	1.79	99.6
cpd 195-En 2	(6)	0.5% DEA in MeOH	30	3	1.54	99.8
cpd 196-En 1	(9)	0.5% DEA in MeOH	40	3	1.81	95.6
cpd 196-En 2	(9)	0.5% DEA in MeOH	40	3	1.47	99.9
cpd 198-En 1	(4)	0.5% DEA in MeOH	40	3	2.58	99.7
cpd 198-En 2	(4)	0.5% DEA in MeOH	40	3	2.00	99.9
cpd 199-En 1	(5)	MeOH	20	3	2.27	99.9
cpd 199-En 2	(5)	MeOH	20	3	1.80	99.4
cpd 205-En 1	(7)	MeOH	20	3	3.59	99.7
cpd 205-En 2	(7)	MeOH	20	3	2.17	99.7
cpd 207-En 1	(5)	0.5% DEA in MeOH	30	3	2.55	97.2
cpd 207-En 2	(5)	0.5% DEA in MeOH	30	3	1.92	98.1
Cpd 270-En 1	(8)	30 mM Ammonia in MeOH	25	3	10.80	91.0
Cpd 270-En 2	(8)	30 mM Ammonia in MeOH	25	3	7.55	99.5
Cpd 313-En 1	(6)	0.5% DEA in MeOH	30	3	7.02	99.3
Cpd 313-En 2	(6)	0.5% DEA in MeOH	30	3	3.69	99.2
Cpd 315-En 1	(9)	IPA	40	3	4.60	99.6
Cpd 315-En 2	(9)	IPA	40	3	3.62	99.9
Cpd 316-En 1	(6)	MeOH	40	3	1.22	97.6
Cpd 316-En 2	(6)	MeOH	40	3	1.48	95.6
Cpd 317-En 1	(13)	MeOH	40	3	5.18	99.8
Cpd 317-En 2	(13)	MeOH	40	3	3.11	99.9
Cpd 326-En 1	(3)	MeOH	40	3	7.76	100.0
Cpd 326-En 2	(3)	MeOH	40	3	9.84	99.6
Cpd 332-En 1	(5)	MeOH	30	3	2.82	100.0
Cpd 332-En 2	(5)	MeOH	30	3	3.66	99.6
Cpd 333-En 1	(11)	MeOH	40	3	2.32	100.0

-continued

Chiral SFC analysis						
Code	Column Name	co-solvent	co-solvent %	Flow [g/min]	RT [min]	Purity [%]
Cpd 333-En 2	(11)	MeOH	40	3	3.30	98.2
Cpd 336-En 1	(1)	MeOH	40	3	2.88	99.4
Cpd 336-En 2	(1)	MeOH	40	3	4.59	98.7
Cpd 342-En 1	(6)	MeOH	40	3	5.69	99.8
Cpd 342-En 2	(6)	MeOH	40	3	2.70	99.9
Cpd 348-En 1	(9)	MeOH	40	3	5.84	96.7
Cpd 348-En 2	(9)	MeOH	40	3	3.57	99.2
Cpd 349-En 1	(9)	MeOH	40	3	3.63	96.9
Cpd 349-En 2	(9)	MeOH	40	3	2.99	99.2
Cpd 350-En 1	(9)	MeOH	40	3	4.43	97.6
Cpd 350-En 2	(9)	MeOH	40	3	3.44	99.2
Cpd 355-En 1	(5)	MeOH	30	3	3.58	99.8
Cpd 355-En 2	(5)	MeOH	30	3	4.65	99.0
Cpd 356-En 1	(11)	MeOH	40	3	3.30	99.9
Cpd 356-En 2	(11)	MeOH	40	3	4.70	96.8
Cpd 360-En 1	(11)	MeOH	40	4	12.59	99.2
Cpd 360-En 2	(11)	MeOH	40	4	9.56	100.0
Cpd 365-En 1	(4)	MeOH	40	3	3.18	99.9
Cpd 365-En 2	(4)	MeOH	40	3	4.57	99.8
Cpd 386-En 1	(4)	0.5% DEA in MeOH	40	3	4.48	99.9
Cpd 386-En 2	(4)	0.5% DEA in MeOH	40	3	7.10	99.9
Cpd 387-En 1	(3)	MeOH	50	4	4.32	99.2
Cpd 387-En 2	(3)	MeOH	50	4	9.59	99.7
Cpd 388-En 1	(1)	MeOH	40	3	2.04	100.0
Cpd 388-En 2	(1)	MeOH	40	3	2.86	99.9
Cpd 389-En 1	(3)	MeOH	40	4	6.07	99.6
Cpd 389-En 2	(3)	MeOH	40	4	9.29	99.8
Cpd 395-Dia 1	(4)	IPA	20	3	5.94	96.5
Cpd 395-Dia 2	(4)	IPA	20	3	4.25	99.5
Cpd 396-En 1	(1)	0.5% DEA in MeOH	40	3	1.86	92.2
Cpd 396-En 2	(3)	MeOH	40	3	5.59	96.3
Cpd 397-En 1	(1)	0.5% DEA in MeOH	40	3	2.79	97.4
Cpd 397-En 2	(4)	0.5% DEA in MeOH	30	3	3.77	97.6
Cpd 401-En 1	(13)	0.5% DEA in MeOH	40	3	2.90	99.9
Cpd 401-En 2	(13)	0.5% DEA in MeOH	40	3	2.19	99.7
Cpd 402-En 1	(2)	MeOH	25	3	5.92	99.8
Cpd 403-En 1	(9)	0.5% DEA in MeOH	40	3	7.35	100.0
Cpd 403-En 2	(9)	0.5% DEA in MeOH	40	3	5.07	100.0
Cpd 404-En 1	(9)	MeOH	35	3	12.13	96.1
Cpd 404-En 2	(9)	MeOH	35	3	9.74	99.8
Cpd 406-En 1	(4)	EtOH	40	3	5.69	99.7
Cpd 406-En 2	(4)	EtOH	40	3	7.65	99.1
Cpd 407-En 1	(3)	MeOH	40	4	7.98	99.8
Cpd 407-En 2	(3)	MeOH	40	4	13.25	99.6
Cpd 408-En 1	(3)	MeOH	50	4	11.11	99.1
Cpd 408-En 2	(3)	MeOH	50	4	6.04	99.9
Cpd 409-En 1	(9)	MeOH	40	3	6.82	99.9
Cpd 409-En 2	(9)	MeOH	40	3	4.46	100.0
Cpd 410-En 1	(4)	MeOH	40	3	3.93	99.9
Cpd 410-En 2	(4)	MeOH	40	3	1.65	99.7
Cpd 411-En 1	(4)	MeOH	40	3	6.88	96.5
Cpd 411-En 2	(4)	MeOH	40	3	2.97	98.2
Cpd 412-En 1	(4)	IPA	25	3	3.30	98.8
Cpd 412-En 2	(4)	IPA	25	3	2.48	99.3
Cpd 413-En 1	(6)	0.5% DEA in MeOH	40	3	2.56	99.5
Cpd 413-En 2	(6)	0.5% DEA in MeOH	40	3	3.51	95.1
Cpd 432-En 1	(7)	MeOH	40	3	1.86	99.1
Cpd 432-En 2	(7)	MeOH	40	3	3.67	99.9
Cpd 433-En 1	(1)	MeOH	30	3	5.13	99.9
Cpd 433-En 2	(1)	MeOH	30	3	2.46	99.4
Cpd 434-En 1	(8)	MeOH	40	3	7.31	100.0
Cpd 434-En 2	(8)	MeOH	40	3	2.97	100.0
Cpd 435-En 1	(9)	0.5% DEA in MeOH	40	3	3.14	99.9
Cpd 435-En 2	(9)	0.5% DEA in MeOH	40	3	6.33	99.9
Cpd 436-En 1	(9)	MeOH	40	3	10.57	99.9
Cpd 436-En 2	(9)	MeOH	40	3	5.48	99.9
Cpd 437-En 1	(5)	0.5% DEA in MeOH	40	3	1.84	99.7
Cpd 437-En 2	(5)	0.5% DEA in MeOH	40	3	5.23	99.8
Cpd 445-En 1	(9)	0.5% DEA in MeOH	40	3	5.65	98.7
Cpd 445-En 2	(9)	0.5% DEA in MeOH	40	3	9.32	98.2
Cpd 446-En 1	(4)	MeOH	40	3	3.08	99.3
Cpd 446-En 2	(4)	MeOH	40	3	2.08	99.7
Cpd 447-En 1	(4)	MeOH	35	3	6.58	99.9

-continued

Chiral SFC analysis						
Code	Column Name	co-solvent	co-solvent %	Flow [g/min]	RT [min]	Purity [%]
Cpd 447-En 2	(4)	MeOH	35	3	2.35	99.7
Cpd 448-En 1	(9)	MeOH	30	3	3.26	98.9
Cpd 448-En 2	(9)	MeOH	30	3	4.80	99.1
Cpd 449-En 1	(9)	0.5% DEA in MeOH	40	3	4.71	99.9
Cpd 449-En 2	(9)	0.5% DEA in MeOH	40	3	8.69	99.5
Cpd 450-En 1	(5)	0.5% DEA in MeOH	30	3	4.32	99.9
Cpd 450-En 2	(5)	0.5% DEA in MeOH	30	3	6.95	99.6
Cpd 451-En 1	(9)	0.5% DEA in MeOH	40	3	3.49	99.9
Cpd 451-En 2	(9)	0.5% DEA in MeOH	40	3	5.14	99.9
Cpd 452-En 1	(4)	MeOH	40	3	4.12	97.3
Cpd 452-En 2	(4)	MeOH	40	3	3.42	99.2
Cpd 455-En 1	(9)	MeOH	40	3	3.10	99.9
Cpd 455-En 2	(9)	MeOH	40	3	4.74	99.8
Cpd 456-En 1	(4)	MeOH	35	3	3.92	99.5
Cpd 456-En 2	(4)	MeOH	35	3	2.54	99.9
Cpd 457-En 1	(5)	MeOH	40	3	2.05	99.9
Cpd 457-En 2	(5)	MeOH	40	3	3.79	99.0
Cpd 458-En 1	(9)	MeOH	40	3	2.84	99.5
Cpd 458-En 2	(9)	MeOH	40	3	4.36	97.3
Cpd 459-En 1	(9)	0.5% DEA in MeOH	40	4	9.06	98.8
Cpd 459-En 2	(9)	0.5% DEA in MeOH	40	4	6.34	94.1
Cpd 460-En 1	(4)	MeOH	35	3	4.26	100.0
Cpd 460-En 2	(4)	MeOH	35	3	2.18	100.0
Cpd 461-En 1	(4)	MeOH	40	3	2.58	99.8
Cpd 461-En 2	(4)	MeOH	40	3	4.69	98.4
Cpd 462-En 1	(1)	0.5% IP Amine in IPA	40	3	1.71	99.9
Cpd 462-En 2	(1)	0.5% IP Amine in IPA	40	3	2.23	98.8
Cpd 463-En 1	(14)	EtOH	30	3	5.45	99.4
Cpd 463-En 2	(14)	EtOH	30	3	4.01	96.6
Cpd 464-En 1	(4)	MeOH	30	3	6.03	99.1
Cpd 464-En 2	(4)	MeOH	30	3	5.62	95.4
Cpd 465-En 1	(4)	MeOH	30	3	2.84	99.5
Cpd 465-En 2	(4)	MeOH	30	3	4.27	99.2
Cpd 466-En 1	(15)	0.5% IP Amine in IPA	30	3	3.47	98.9
Cpd 466-En 2	(15)	0.5% IP Amine in IPA	30	3	2.58	99.1
Cpd 467-En 1	(3)	MeOH	40	4	3.40	99.9
Cpd 467-En 2	(3)	MeOH	40	4	5.24	99.7
Cpd 468-En 1	(8)	MeOH	40	4	13.80	99.8
Cpd 468-En 2	(8)	MeOH	40	4	17.00	97.3
Cpd 469-En 1	(4)	MeOH	40	3	3.03	99.7
Cpd 469-En 2	(4)	MeOH	40	3	2.04	97.5
Cpd 487-En 1	(6)	MeOH	40	3	7.18	99.9
Cpd 487-En 2	(6)	MeOH	40	3	12.10	99.6
Cpd 488-En 1	(4)	MeOH	20	3	5.99	99.9
Cpd 488-En 2	(4)	MeOH	20	3	8.52	97.9
Cpd 489-En 1	(6)	MeOH	30	3	1.82	99.3
Cpd 489-En 2	(6)	MeOH	30	3	2.58	100.0
Cpd 491-En 1	(4)	IPA	30	3	1.66	99.5
Cpd 491-En 2	(4)	IPA	30	3	2.73	99.7
Cpd 510-En 1	(3)	EtOH	40	4	6.45	100.0
Cpd 510-En 2	(3)	EtOH	40	4	13.34	99.9
Cpd 511-En 1	(5)	MeOH	35	3	3.15	99.9
Cpd 511-En 2	(5)	MeOH	35	3	4.04	99.8
Cpd 512-En 1	(5)	MeOH	30	3	7.00	99.9
Cpd 512-En 2	(5)	MeOH	30	3	9.01	99.8
Cpd 513-En 1	(13)	MeOH	30	3	2.86	99.8
Cpd 513-En 2	(13)	MeOH	30	3	3.54	99.3
Cpd 514-En 1	(13)	MeOH	35	3	5.70	99.7
Cpd 514-En 2	(13)	MeOH	35	3	7.10	99.3
Cpd 515-En 1	(11)	MeOH	35	3	6.52	99.8
Cpd 515-En 2	(11)	MeOH	35	3	10.07	97.9
Cpd 516-En 1	(13)	0.5% DEA in MeOH	35	3	2.25	99.8
Cpd 516-En 2	(13)	0.5% DEA in MeOH	35	3	2.62	98.3
Cpd 517-En 1	(3)	MeOH	50	4	18.69	99.9
Cpd 518-En 1	(11)	MeOH	50	4	5.46	99.9
Cpd 518-En 2	(11)	MeOH	50	4	8.69	97.9
Cpd 519-En 1	(11)	MeOH	40	3	8.75	99.1
Cpd 519-En 2	(11)	MeOH	40	3	14.26	97.8
Cpd 520-En 1	(11)	MeOH	50	4	6.38	99.1
Cpd 520-En 2	(11)	MeOH	50	4	10.10	95.3
Cpd 521-En 1	(9)	0.5% DEA in MeOH	40	3	5.78	97.7
Cpd 521-En 2	(9)	0.5% DEA in MeOH	40	3	6.56	96.7
Cpd 522-En 1	(13)	MeOH	30	3	2.53	99.7

Chiral SFC analysis						
Code	Column Name	co-solvent	co-solvent %	Flow [g/min]	RT [min]	Purity [%]
Cpd 522-En 2	(13)	MeOH	30	3	3.00	98.3
Cpd 530-En 1	(4)	0.5% DEA in MeOH	40	3	1.36	99.7
Cpd 530-En 2	(4)	0.5% DEA in MeOH	40	3	3.41	99.4
Cpd 531-En 1	(2)	MeOH	35	3	2.28	99.9
Cpd 531-En 2	(2)	MeOH	35	3	3.15	99.9
Cpd 532-En 1	(2)	MeOH	25	3	4.46	99.7
Cpd 532-En 2	(2)	MeOH	25	3	6.30	97.1
Cpd 533-En 1	(16)	MeOH	35	3	4.06	99.7
Cpd 533-En 2	(16)	MeOH	35	3	5.13	99.5
Cpd 534-En 1	(4)	MeOH	40	3	4.12	99.9
Cpd 534-En 2	(4)	MeOH	40	3	2.95	99.5
Cpd 535-En 1	(16)	0.5% DEA in MeOH	40	4	4.12	97.2
Cpd 535-En 2	(16)	0.5% DEA in MeOH	40	4	3.49	98.8
Cpd 536-En 1	(1)	MeOH	30	3	2.43	99.3
Cpd 536-En 2	(1)	MeOH	30	3	3.49	99.9
Cpd 538-En 1	(5)	MeOH	30	3	1.68	99.5
Cpd 539-En 1	(5)	0.5% DEA in MeOH	20	3	1.45	99.8
Cpd 539-En 2	(5)	0.5% DEA in MeOH	20	3	2.08	99.8
Cpd 540-En 1	(6)	0.5% DEA in EtOH	40	3	8.33	97.2
Cpd 540-En 2	(6)	0.5% DEA in EtOH	40	3	7.01	98.9
Cpd 541-En 1	(6)	0.5% DEA in EtOH	40	3	5.65	99.1
Cpd 541-En 2	(6)	0.5% DEA in EtOH	40	3	4.02	98.6
Cpd 542-En 1	(8)	0.5% DEA in MeOH	40	4	9.68	98.4
Cpd 542-En 2	(8)	0.5% DEA in MeOH	40	4	12.98	97.0
Cpd 543-En 1	(4)	EtOH	40	3	4.58	99.2
Cpd 543-En 2	(4)	EtOH	40	3	2.34	98.8
Cpd 544-En 1	(7)	0.5% DEA in MeOH	30	3	1.72	99.4
Cpd 544-En 2	(7)	0.5% DEA in MeOH	30	3	2.43	98.7
Cpd 545-En 1	(4)	0.5% DEA in MeOH	40	3	2.65	97.7
Cpd 545-En 2	(4)	0.5% DEA in MeOH	40	3	3.57	95.0
Cpd 547-En 1	(4)	EtOH	40	3	7.20	99.8
Cpd 547-En 2	(4)	EtOH	40	3	2.19	99.4
Cpd 548-En 1	(5)	MeOH	40	3	1.59	99.9
Cpd 548-En 2	(5)	MeOH	40	3	2.44	99.0
Cpd 549-En 1	(7)	MeOH	30	3	1.46	99.9
Cpd 549-En 2	(7)	MeOH	30	3	2.18	99.7
Cpd 550-En 1	(4)	0.5% DEA in MeOH	25	3	4.63	95.6
Cpd 550-En 2	(4)	0.5% DEA in MeOH	25	3	3.28	99.5
Cpd 551-En 1	(2)	MeOH	25	3	2.69	99.2
Cpd 551-En 2	(2)	MeOH	25	3	4.40	98.4
Cpd 552-En 1	(5)	MeOH	30	3	2.27	99.5
Cpd 552-En 2	(5)	MeOH	30	3	2.96	99.6
Cpd 553-En 1	(7)	MeOH	25	3	1.40	98.9
Cpd 553-En 2	(7)	MeOH	25	3	2.03	98.1
Cpd 554-En 1	(5)	MeOH	30	3	2.51	99.0
Cpd 554-En 2	(5)	MeOH	30	3	3.10	99.8
Cpd 555-En 1	(7)	IPA	40	3	1.13	99.9
Cpd 555-En 2	(7)	IPA	40	3	1.92	99.4
Cpd 556-En 1	(5)	0.5% DEA in MeOH	30	3	3.91	97.8
Cpd 556-En 2	(5)	0.5% DEA in MeOH	30	3	4.83	98.8
Cpd 557-En 1	(17)	MeOH	40	4	6.63	99.2
Cpd 557-En 2	(17)	MeOH	40	4	9.10	98.3
Cpd 558-En 1	(16)	MeOH	30	3	6.15	98.8
Cpd 558-En 2	(16)	MeOH	30	3	7.89	99.4
Cpd 559-En 1	(4)	0.5% DEA in MeOH	40	3	4.26	97.3
Cpd 559-En 2	(4)	0.5% DEA in MeOH	40	3	2.54	97.7
Cpd 561-En 1	(7)	MeOH	40	3	1.23	98.8
Cpd 561-En 2	(7)	MeOH	40	3	1.63	99.6
Cpd 562-En 1	(5)	MeOH	40	3	2.14	99.6
Cpd 562-En 2	(5)	MeOH	40	3	3.30	97.1
Cpd 563-En 1	(8)	MeOH	40	4	11.84	99.6
Cpd 563-En 2	(8)	MeOH	40	4	5.84	99.9
Cpd 564-En 1	(2)	MeOH	25	3	4.83	99.3
Cpd 564-En 2	(2)	MeOH	25	3	8.28	97.3
Cpd 566-En 1	(2)	MeOH	30	3	2.60	99.4
Cpd 566-En 2	(2)	MeOH	30	3	3.85	97.0
Cpd 567-En 1	(5)	0.5% DEA in MeOH	30	3	4.17	99.0
Cpd 567-En 2	(5)	0.5% DEA in MeOH	30	3	5.60	96.2
Cpd 568-En 1	(6)	0.5% DEA in MeOH	40	3	6.17	99.9
Cpd 568-En 2	(6)	0.5% DEA in MeOH	40	3	11.11	99.8
Cpd 569-En 1	(3)	0.5% DEA in MeOH	40	4	11.89	97.8
Cpd 569-En 2	(3)	0.5% DEA in MeOH	40	4	7.67	96.2
Cpd 570-En 1	(6)	0.5% DEA in EtOH	30	3	4.64	96.6

-continued

Chiral SFC analysis						
Code	Column Name	co-solvent	co-solvent %	Flow [g/min]	RT [min]	Purity [%]
Cpd 570-En 2	(6)	0.5% DEA in EtOH	30	3	3.96	97.2
Cpd 571-En 1	(4)	0.5% DEA in MeOH	40	3	7.15	99.8
Cpd 571-En 2	(4)	0.5% DEA in MeOH	40	3	3.90	99.2
Cpd 572-En 1	(6)	MeOH	40	3	4.63	99.4
Cpd 572-En 2	(6)	MeOH	40	3	2.94	100.0
Cpd 573-En 1	(6)	0.5% DEA in MeOH	40	3	3.87	93.9
Cpd 573-En 2	(6)	0.5% DEA in MeOH	40	3	2.76	97.1
Cpd 575-En 1	(13)	0.5% DEA in MeOH	20	3	6.59	99.5
Cpd 575-En 2	(13)	0.5% DEA in MeOH	20	3	7.57	99.6
Cpd 590-En 1	(4)	IPA	35	3	4.05	96.6
Cpd 590-En 2	(4)	IPA	35	3	2.96	99.4
Cpd 591-En 1	(17)	MeOH	40	3	4.20	99.9
Cpd 591-En 2	(17)	MeOH	40	3	6.10	99.6
Cpd 592-En 1	(3)	MeOH	50	4	9.41	98.7
Cpd 592-En 2	(3)	MeOH	50	4	4.05	99.2
Cpd 593-En 1	(3)	MeOH	30	3	13.39	99.6
Cpd 593-En 2	(3)	MeOH	30	3	10.89	99.9
Cpd 594-En 1	(4)	MeOH	40	3	12.23	99.8
Cpd 594-En 2	(4)	MeOH	40	3	5.43	99.4
Cpd 595-En 1	(4)	MeOH	40	3	4.72	100.0
Cpd 595-En 2	(4)	MeOH	40	3	7.39	99.8
Cpd 597-En 1	(17)	MeOH	40	4	8.05	99.8
Cpd 597-En 2	(17)	MeOH	40	4	4.95	99.9
Cpd 598-En 1	(3)	MeOH	30	3	7.79	99.7
Cpd 598-En 2	(3)	MeOH	30	3	10.31	98.2
Cpd 600-En 1	(4)	MeOH	40	3	4.15	96.5
Cpd 600-En 2	(4)	MeOH	40	3	2.49	98.8
Cpd 601-En 1	(9)	MeOH	40	3	3.22	99.7
Cpd 601-En 2	(9)	MeOH	40	3	4.69	99.3
Cpd 623-En 1	(12)	MeOH	20	3	5.96	99.7
Cpd 623-En 2	(12)	MeOH	20	3	9.51	99.2
Cpd 635-En 1	(5)	0.5% DEA in MeOH	20	3	3.51	98.9
Cpd 635-En 2	(5)	0.5% DEA in MeOH	20	3	2.24	99.0
Cpd 636-En 1	(10)	MeOH	40	4	2.78	98.4
Cpd 636-En 2	(10)	MeOH	40	4	4.44	97.7
Cpd 637-En 1	(1)	MeOH	40	3	2.71	98.8
Cpd 637-En 2	(1)	MeOH	40	3	3.93	99.9
Cpd 638-En 1	(9)	0.5% DEA in MeOH	40	3	2.17	99.9
Cpd 638-En 2	(9)	0.5% DEA in MeOH	40	3	1.70	99.8
Cpd 639-En 1	(6)	0.5% DEA in MeOH	40	3	4.09	97.8
Cpd 639-En 2	(6)	0.5% DEA in MeOH	40	3	2.82	99.7
Cpd 640-En 1	(4)	MeOH	40	3	8.41	98.4
Cpd 640-En 2	(4)	MeOH	40	3	3.13	99.4
Cpd 641-En 1	(4)	EtOH	40	3	2.13	99.3
Cpd 641-En 2	(4)	EtOH	40	3	3.53	99.4
Cpd 642-En 1	(9)	MeOH	15	3	3.43	99.9
Cpd 642-En 2	(9)	MeOH	15	3	4.41	98.9
Cpd 643-En 1	(9)	MeOH	30	3	3.59	99.9
Cpd 643-En 2	(9)	MeOH	30	3	2.52	100.0
Cpd 646-En 1	(8)	MeOH	45	3	5.57	99.9
Cpd 646-En 2	(8)	MeOH	45	3	7.23	99.4
Cpd 647-En 1	(6)	MeOH	40	3	3.26	99.4
Cpd 647-En 2	(6)	MeOH	40	3	4.85	100.0
Cpd 648-En 1	(6)	EtOH	40	3	2.04	98.6
Cpd 648-En 2	(6)	EtOH	40	3	3.16	97.1
Cpd 653-En 1	(9)	MeOH	4	3	3.26	99.1
Cpd 653-En 2	(9)	MeOH	4	3	2.60	99.3
Cpd 654-En 1	(9)	MeOH	4	3	3.00	99.4
Cpd 654-En 2	(9)	MeOH	4	3	2.47	99.4
Cpd 655-En 1	(6)	MeOH	40	3	4.72	98.4
Cpd 655-En 2	(6)	MeOH	40	3	7.67	97.1
Cpd 656-En 1	(9)	MeOH	35	3	3.1	99.6
Cpd 656-En 2	(9)	MeOH	35	3	2.48	99.9
Cpd 657-En 1	(9)	MeOH	25	3	15.35	99.3
Cpd 657-En 2	(9)	MeOH	25	3	11.72	99.1
Cpd 661-En 1	(3)	MeOH	40	4	9.70	98.6
Cpd 661-En 2	(3)	MeOH	40	4	6.77	99.7
Cpd 719-En 1	(15)	MeOH	40	3	4.31	99.91
Cpd 719-En 2	(15)	MeOH	40	3	5.49	99.79
Cpd 720-En 1	(3)	0.5% DEA in EtOH	40	3	2.19	98.40
Cpd 720-En 2	(3)	0.5% DEA in EtOH	40	3	2.70	96.48
Cpd 723-En 1	(7)	MeOH	20	3	1.94	99.52
Cpd 723-En 2	(7)	MeOH	20	3	3.21	98.87

Chiral SFC analysis						
Code	Column Name	co-solvent	co-solvent %	Flow [g/min]	RT [min]	Purity [%]
Cpd 724-En 1	(7)	MeOH	15	3	2.51	99.76
Cpd 724-En 2	(7)	MeOH	15	3	5.19	99.54
Cpd 725-En 1	(4)	MeOH	40	3	2.33	99.90
Cpd 725-En 2	(4)	MeOH	40	3	3.78	99.18
Cpd 726-En 1	(3)	MeOH	50	4	7.04	99.74
Cpd 726-En 2	(3)	MeOH	50	4	13.63	99.03
Cpd 727-En 1	(9)	0.5% DEA in MeOH	40	3	7.45	99.88
Cpd 727-En 2	(9)	0.5% DEA in MeOH	40	3	9.55	99.41
Cpd 727-En 3	(9)	MeOH	40	3	7.24	99.95
Cpd 727-En 4	(9)	MeOH	40	3	8.81	98.96
Cpd 728-En 1	(6)	0.5% DEA in MeOH	40	3	2.68	98.62
Cpd 728-En 2	(6)	0.5% DEA in MeOH	40	3	4.26	91.00
Cpd 730-En 1	(4)	MeOH	35	3	2.38	99.98
Cpd 730-En 2	(4)	MeOH	35	3	4.47	99.89
Cpd 731-En 1	(4)	MeOH	35	3	2.91	99.73
Cpd 731-En 2	(4)	MeOH	35	3	7.73	99.87
Cpd 732-En 1	(11)	MeOH	40	3	6.38	99.59
Cpd 732-En 2	(11)	MeOH	40	3	11.44	99.00
Cpd 733-En 1	(11)	MeOH	35	3	6.49	99.96
Cpd 733-En 2	(11)	MeOH	35	3	9.83	96.78
Cpd 734-En 1	(4)	0.5% IP Amine in IPA	40	3	1.80	99.73
Cpd 734-En 2	(4)	0.5% IP Amine in IPA	40	3	3.15	99.10
Cpd 735-En 1	(11)	MeOH	40	3	5.11	99.88
Cpd 735-En 2	(11)	MeOH	40	3	7.47	96.72
Cpd 736-En 1	(11)	MeOH	40	3	3.76	99.15
Cpd 736-En 2	(11)	MeOH	40	3	5.26	97.35
Cpd 737-En 1	(5)	MeOH	30	3	4.52	95.25
Cpd 737-En 2	(5)	MeOH	30	3	6.07	95.40
Cpd 738-En 1	(4)	EtOH	40	3	4.42	98.28
Cpd 738-En 2	(4)	EtOH	40	3	6.48	97.26
Cpd 739-En 1	(4)	MeOH	40	3	4.49	98.44
Cpd 739-En 2	(4)	MeOH	40	3	5.55	94.15
Cpd 740-En 1	(11)	MeOH	35	3	2.83	99.41
Cpd 740-En 2	(11)	MeOH	35	3	3.47	96.41

CHIRALCEL OZ-3 (4.6*150 mm) 3 μ m = (1); CHIRACEL OD-H (250 mm $\times$ 4.6), 5 μ m = (2); CHIRALPAK AD-H (4.6 $\times$ 250 mm), 5 μ m = (3); CHIRALPAK AD-3 (4.6*150 mm) 3 μ m = (4); CHIRALCEL OX-3 (4.6*150 mm) 3 μ m = (5); Chiralpak IG-3 (4.6 $\times$ 150 mm), 3 μ m = (6); CHIRALCEL OD-3 (4.6*150 mm) 3 μ m = (7); CHIRALPAK IG-3 (4.6*250 mm) 5 μ m = (8); CHIRALPAK IC-3 (4.6*150 mm) 3 μ m = (9); Lux Cellulose-2 (4.6 $\times$ 250 mm, 5 μ m = (10); CHIRALPAK IE-3 (4.6*150 mm) 3 μ m = (11); Lux-Amylose-2 (4.6 $\times$ 250 mm, 5 μ m = (12); (R,R) WHELK-01 (4.6*150 mm) 3.5 μ m = (13); CHIRALPAK AD-H (4.6*150 mm) 3 μ m = (14); Lux Cellulose-2 (4.6 $\times$ 150 mm) 3 μ m = (15); CHIRALCEL OX-H (4.6*250 mm, 5 μ m = (16); CHIRALPAK IF-3 (4.6*150 mm), 3 μ m = (17)

Chiral HPLC analysis						
Cpd	Column Name	Phase A/B	Ratio A/B	Flow [mL/min]	RT [min]	Purity [%]
Cpd 729-En 1	CHIRALCEL ODH (250 $\times$ 4.6 mm); 5 μ	n-Hex/ EtOH	3/7	1	5.44	96.50
Cpd 729-En 2	CHIRALCEL ODH (250 $\times$ 4.6 mm); 5 μ	n-Hex/ EtOH	3/7	1	9.58	99.89

Part B

Monitoring the TRPM3 Ion Channel Driven Ca²⁺ Uptake.

In order to monitor the inhibition of the mouse TRPM3 α 2 (mTRPM3) ion channel by the compounds of the invention, a cellular system making use of an mTRPM3 α 2 or hTRPM3 overexpressing cell line (flip-in HEK293) was used. The TRPM3 channel was stimulated/opened with Pregnenolone sulfate (PS) (50 μ M) which results in Ca²⁺ influx.

For mTRPM3, the intracellular Ca²⁺ was measured with a Calcium responsive dye, Fluor-4 AM ester (Invitrogen). Cells were cultured until a confluence of 80-90%, washed with Versene (Invitrogen) and detached from the surface by a short incubation with 0.05% Trypsin (Invitrogen). The

trypsination process was stopped by the addition of complete cell culture medium (DMEM, glutamax, 10% FCS, NEAA, Pen-Strep). Cells were collected and resuspended in Krebs buffer without Calcium at RT.

Prior the cell seeding (+2000 cells/well into a black, 384 well plate (Greiner)) the diluted compound was added in the assay plate, together with the PS dissolved in Krebs buffer containing Calcium. This resulted in a 2.4 mM Ca²⁺ assay solution. Directly after cell addition the plates were read on an Envision fluorescence reader (Perkin Elmer) by an Excitation of 485 nM and emission at 535 nM.

Channel inhibition was calculated compared to a non-PS stimulated control versus a condition stimulated with PS (50 μ M) with vehicle. The ability of the compounds of the

577

invention to inhibit this activity was determined as: Percentage inhibition=[1-((RFU determined for sample with test compound present-RFU determined for sample with positive control inhibitor) divided by (RFU determined in the presence of vehicle-RFU determined for sample with positive control inhibitor))]*100.

The activities of the Example compounds cpd 001-199 tested are depicted in the table below. The activity ranges A, B and C refer to IC₅₀ values in the Fluo-4 AM assay as follows: "A": IC₅₀<1 μM; "B": 1 μM≤IC₅₀≤20 μM and "C": IC₅₀>20 μM

cpd	IC ₅₀	
001	B	15
002	C	
003	C	
004	C	
005	B	20
006	B	
007	B	
008	C	
009	B	25
010	B	
011	B	
012	B	
013	B	30
014	B	
015	C	
016	C	
017	C	35
018	B	
019	C	
020	B	
021	B	40
022	A	
023	B	
024	A	
025	A	45
026	A	
027	A	
028	A	
029	A	50
030	A	
031	B	
032	B	
033	C	55
034	B	
035	A	
036	A	
037	B	60
038	A	
039	B	
040	C	
041	A	65
042	A	
043	A	
044	B	
045	B	
046	A	
047	A	
048	A	
049	A	
050	A	
051	A	
052	B	
053	B	
054	A	
055	C	
056	A	
057	C	
058	A	
059	B	
060	B	
061	C	
062	A	
063	A	
064	A	

578

-continued

cpd	IC ₅₀
065	B
066	B
067	A
068	A
069	B
070	A
071	A
072	A
073	B
074	A
075	C
076	B
077	C
078	B
079	B
080	B
081	B
082	B
083	B
084	B
085	B
086	B
087	B
088	B
089	B
090	B
091	B
092	B
093	B
094	B
095	B
096	B
097	B
098	B
099	B
100	B
101	B
102	B
103-En 1	A
103-En 2	A
104	A
105-En 1	A
105-En 2	A
106-En 1	A
106-En 2	A
107	A
108	A
109	A
110-En 1	A
110-En 2	A
111-En 1	A
111-En 2	A
112-En 1	A
112-En 2	A
114	A
115-En 1	A
115-En 2	A
116-En 1	A
116-En 2	A
117	A
118-En 1	A
118-En 2	A
119	A
120-En 1	A
120-En 2	A
121-En 1	A
121-En 2	A
122	B
123-En 1	A
123-En 2	A
124-En 1	A
124-En 2	A
125-En 1	A
125-En 2	A
126-En 1	A
126-En 2	A
127	A

579

-continued

cpd	IC ₅₀	
128	A	
129	A	5
130-En 1	A	
130-En 2	A	
131-En 1	A	
131-En 2	A	
132-En 1	A	
132-En 2	A	10
133	A	
134-En 1	B	
134-En 2	A	
135-En 1	A	
135-En 2	A	
136	A	15
137	A	
138	A	
139	A	
140	A	
141-En 1	A	
141-En 2	A	20
142	A	
143	A	
144	A	
145	A	
146	A	
147-Dia 1-En 1	A	25
147-Dia 1-En 2	A	
148-En 1	A	
148-En 2	A	
149-En 1	A	
149-En 2	A	
150	A	30
151-En 1	A	
151-En 2	A	
152	A	
153	A	
154	A	
155-En 1	A	
155-En 2	A	35
156	A	
157-En 1	A	
157-En 2	A	
158	A	
159-En 1	A	
159-En 2	A	40
160-En 1	A	
160-En 2	A	
161	A	
cpd 162-Dia 1-En2	A	
cpd 162-Dia 1-En1	A	
163-En 1	A	45
163-En 2	A	
164-En 1	A	
164-En 2	A	
165-En 1	A	
165-En 2	B	
166	A	
167	A	50
168	A	
169-En 1	A	
169-En 2	A	
170	A	
171	A	
172-En 1	A	55
172-En 2	A	
173	A	
174	A	
175	A	
176-En 1	B	
176-En 2	A	60
177	A	
178	B	
179-En 1	B	
179-En 2	C	
180-En 1	B	
180-En 2	B	65
181	B	

580

-continued

cpd	IC ₅₀
182	A
183-En 1	B
183-En 2	C
184-En 1	B
184-En 2	C
185	B
186	B
187	B
188	B
189	B
190	B
191-En 1	C
191-En 2	C
192	C
193	C
194	C
195-En 1	C
195-En 2	C
196-En 1	C
196-En 2	B
197	B
198-En 1	B
198-En 2	B
199-En 1	A
199-En 2	A

The activities of the Example compounds (cpds 200-299) tested are depicted in the table below. The activity ranges A, B and C refer to IC₅₀ values in the Fluo-4 AM assay as follows: “A”: IC₅₀<1 μM; “B”: 1 μM≤IC₅₀≤20 μM and “C”: IC₅₀>20 μM.

cpd	IC ₅₀
200	A
201	B
202	B
203	A
204	A
205 - En 1	B
205 - En 2	A
206	A
207 - En 1	A
207 - En 2	A
232	A
270 - En 1	C
270 - En 2	C
271	A
272	A
273	A
274	A
275	A
276	A
277	B
278	A
279	A
280	A
281	A
282	A
283	A
284	A
285	A
286	A
287	A
288	A
289	A
290	A
291	A
292	A
293	A
294	A
295	A
296	A
297	A

581

-continued

cpd	IC ₅₀
298	A
299	A

The activities of the Example compounds (cpds. 300-399) tested are depicted in the table below. The activity ranges A, B and C refer to IC₅₀ values in the Fluo-4 AM assay as follows: “A”: IC₅₀<1 μM; “B”: 1 μM≤IC₅₀≤20 μM and “C”: IC₅₀>20 μM

cpd	IC ₅₀	
300	A	15
301	A	
302	A	
303	A	
304	A	
305	A	20
306	A	
307	B	
308	B	
309	B	
310-En 1	A	25
310-En 2	A	
311	A	
312	B	
313-En 1	A	
313-En 2	B	30
314	A	
315-En 1	A	
315-En 2	A	
316-En 1	A	
316-En 2	A	35
317-En 1	A	
317-En 2	A	
318	A	
319	A	
320	A	40
321	A	
322	A	
323	A	
324	A	
325	B	45
326-En 1	A	
326-En 2	B	
327	B	
328	A	
329	B	50
330	A	
331	B	
332-En 1	A	
332-En 2	A	
333-En 1	A	55
333-En 2	A	
334	A	
335-En 1	A	
335-En 2	A	
336-En 1	A	60
336-En 2	A	
337	A	
338-En 1	A	
338-En 2	A	
339	A	65
340-En 1	A	
340-En 2	A	
341	A	
342-En 1	A	
342-En 2	A	
343	A	
344	A	
345	A	
346	A	
347	A	
348-En 1	A	
348-En 2	A	

582

-continued

cpd	IC ₅₀
349-En 1	A
349-En 2	A
350-En 1	A
350-En 2	A
351	B
352	A
355-En 1	A
355-En 2	A
356-En 1	A
356-En 2	A
357	A
358	C
359	A
360-En 1	A
360-En 2	A
361	A
362	A
363	A
364	A
365-En 1	C
365-En 2	C
366	B
367	A
368	B
369	B
370	A
371	A
372	A
373	A
374	A
375	A
376	A
377	A
378	A
379	A
380	A
381	A
382	A
383	A
384	A
385	A
386-En 1	A
386-En 2	B
387-En 1	A
387-En 2	A
388-En 1	A
388-En 2	B
389-En 1	A
389-En 2	A
390	A
391	A
392	A
393	A
394	A
395-Dia 1	A
395-Dia 2	A
396-En 1	A
396-En 2	A
397-En 1	A
397-En 2	A
398	A
399	A

The activities of the Example compounds (cpds. 400-499) tested are depicted in the table below. The activity ranges A, B and C refer to IC₅₀ values in the Fluo-4 AM assay as follows: “A”: IC₅₀<1 μM; “B”: 1 μM≤IC₅₀≤20 μM and “C”: IC₅₀>20 μM

cpd	IC ₅₀
400	A
401-En 1	A
401-En 2	B

583

-continued

cpd	IC ₅₀	
402-En 1	B	5
403-En 1	B	
403-En 2	C	
404-En 1	B	
404-En 2	B	
405	C	10
406-En 1	A	
406-En 2	A	
407-En 1	A	
407-En 2	B	
408-En 1	A	
408-En 2	A	
409-En 1	A	
409-En 2	A	
410-En 1	A	15
410-En 2	B	
411-En 1	A	
411-En 2	A	
412-En 1	A	
412-En 2	A	
413-En 1	A	
413-En 2	A	
414	A	
415	A	25
416	A	
417	A	
418	A	
419	A	
420	A	
421	A	
422	A	
423	A	
424	A	30
425	A	
426	A	
427	A	
428	A	
429	A	
429	A	
430	A	
431	A	
432-En 1	B	40
432-En 2	A	
433-En 1	A	
433-En 2	A	
434-En 1	A	
434-En 2	A	
435-En 1	A	
435-En 2	A	
436-En 1	A	
436-En 2	A	45
437-En 1	A	
437-En 2	A	
438	A	
439	A	
440	A	
441	A	
442	A	50
443	A	
444	A	
445-En 1	A	
445-En 2	B	
446-En 1	A	
446-En 2	B	
447-En 1	A	
447-En 2	A	
448-En 1	A	
448-En 1	A	55
449-En 1	A	
449-En 2	A	
450-En 1	A	
450-En 2	A	
451-En 1	A	
451-En 2	B	
452-En 1	A	
452-En 2	A	
453-En 1	A	

584

-continued

cpd	IC ₅₀
453-En 2	A
454	A
455-En 2	A
456-En 1	A
456-En 2	A
457-En 1	A
457-En 2	A
458-En 1	A
458-En 2	A
459-En 1	A
459-En 2	A
460-En 1	A
460-En 2	A
461-En 1	A
461-En 2	A
462-En 1	A
462-En 2	A
463-En 1	A
463-En 2	A
464-En 1	A
464-En 2	A
465-En 1	A
465-En 2	A
466-En 1	A
466-En 2	A
467-En 1	A
467-En 2	A
468-En 1	A
468-En 2	A
469-En 1	A
469-En 2	A
470	A
471	A
472	A
473	A
474	A
475	A
476	A
477	A
478	A
479	B
480	A
481	A
482	A
483	A
484	A
485	A
486	A
487-En 1	B
487-En 2	B
488-En 1	A
488-En 2	B
489-En 1	A
489-En 2	A
490-En 1	A
490-En 2	A
491-En 1	A
491-En 2	A
492	A
493	A
494	A
495	A
496	A
497	A
498	A
499	A

The activities of the Example compounds (cpds. 500-599) tested are depicted in the table below. The activity ranges A, B and C refer to IC₅₀ values in the Fluo-4 AM assay as follows: "A": IC₅₀<1 μM; "B": 1 μM≤IC₅₀≤20 μM and "C": IC₅₀>20 μM

585

cpd	IC ₅₀
500	A
501	A
502	A
503	A
504	A
505	A
506	A
507	B
508	A
509	A
510-En 1	A
510-En 2	A
511-En 1	A
511-En 2	B
512-En 1	A
512-En 2	A
513-En 1	A
513-En 2	A
514-En 1	A
514-En 2	A
515-En 1	A
515-En 1	B
516-En 1	A
516-En 2	A
517-En 1	A
517-En 2	A
518-En 1	A
518-En 2	A
519-En 1	A
519-En 2	A
520-En 1	A
520-En 2	A
521-En 1	A
521-En 2	B
522-En 1	A
522-En 2	A
523	B
524	A
525	A
526	A
527	A
528	A
529	A
530-En 1	B
530-En 2	A
531-En 1	A
531-En 2	A
532-En 1	A
532-En 2	A
533-En 1	A
533-En 2	A
534-En 1	A
534-En 2	A
535-En 1	A
535-En 2	A
536-En 1	A
536-En 2	A
537	A
538-En 1	B
539-En 1	B
539-En 2	A
540-En 1	B
540-En 2	C
541-En 1	B
541-En 2	C
542-En 1	B
542-En 2	C
543-En 1	A
543-En 2	A
544-En 1	B
544-En 2	B
545-En 1	B
545-En 2	B
546-En 1	A
547-En 1	A
547-En 2	A
548-En 1	B
548-En 2	C
549-En 1	A

586

-continued

cpd	IC ₅₀
549-En 2	B
550-En 1	B
550-En 2	B
551-En 1	A
551-En 2	B
552-En 1	A
552-En 2	B
553-En 1	A
553-En 2	B
554-En 1	A
554-En 2	C
555-En 1	A
555-En 2	B
556-En 1	B
556-En 2	C
557-En 1	A
557-En 2	A
558-En 1	A
558-En 2	A
559-En 1	B
559-En 2	C
560-En 1	A
560-En 2	A
561-En 1	A
561-En 2	A
562-En 1	A
562-En 2	A
563-En 1	A
563-En 2	A
564-En 1	A
564-En 2	A
565-En 1	A
565-En 2	A
566-En 1	A
566-En 2	B
567-En 1	C
567-En 2	C
568-En 1	A
568-En 2	B
569-En 1	B
569-En 2	C
570-En 1	A
570-En 2	B
571-En 1	B
571-En 2	B
572-En 1	A
572-En 2	A
573-En 1	A
573-En 2	B
574	A
575-En 1	A
575-En 1	A
576	A
577	A
578	B
579	A
580	A
581	A
582	A
583	A
584	A
585	A
586	A
587	A
588	A
589	B
590-En 1	A
590-En 2	A
591-En 1	A
591-En 2	A
592-En 1	A
592-En 2	A
593-En 1	A
593-En 2	A
594-En 1	A
594-En 2	A
595-En 1	A

587

-continued

cpd	IC ₅₀	
595-En 2	A	5
596-En 1	A	
596-En 2	A	
597-En 1	A	
597-En 2	A	
598-En 1	A	10
598-En 2	A	
599	B	

The activities of the Example compounds (cpds. 600-666) tested are depicted in the table below. The activity ranges A, B and C refer to IC₅₀ values in the Fluo-4 AM assay as follows: “A”: IC₅₀<1 μM; “B”: 1 μM≤IC₅₀≤20 μM and “C”: IC₅₀>20 μM

cpd	IC ₅₀	
600-En 1	A	15
600-En 2	A	
601-En 1	A	
601-En 2	A	
602	A	
603	B	20
604	C	
605	C	
606	B	
607	B	
608	A	25
609	B	
610	B	
611	A	
612	B	
613	A	30
614	A	
615	A	
616	B	
617	B	
618	A	35
619	B	
620	A	
621	C	
622	B	
623-En 1	A	40
623-En 2	A	
624	A	
625	A	
626	B	
627	A	45
628	B	
629	A	
630	C	
631	C	
632	B	50
633	B	
634	A	
635-En 1	A	
635-En 2	B	
636-En 1	A	55
636-En 2	A	
637-En 1	A	
637-En 2	A	
638-En 1	A	
638-En 2	B	60
639-En 1	A	
639-En 2	A	
640-En 1	A	
640-En 2	A	
641-En 1	B	65
641-En 2	B	
642-En 1	A	
642-En 2	A	
643-En 1	A	
643-En 2	A	
646-En 1	A	

588

-continued

cpd	IC ₅₀	
646-En 2	A	5
647-En 1	A	
647-En 2	A	
648-En 1	A	
648-En 2	A	
649	A	10
650	A	
651	A	
652	A	
653-En 1	B	
654-En 1	B	15
653-En 2	B	
654-En 2	B	
655-En 1	A	
655-En 2	A	
656-En 1	B	20
656-En 2	B	
657-En 1	C	
657-En 2	C	
658	C	
659	B	25
660	C	
661-En 1	A	
661-En 2	A	
662	A	
663	A	30
664	A	
665	A	
666	A	

The activities of the Example compounds (cpds. 667-675) tested are depicted in the table below. The activity ranges A, B and C refer to IC₅₀ values in the Fluo-4 AM assay as follows: “A”: IC₅₀<1 μM; “B”: 1 μM≤IC₅₀≤20 μM and “C”: IC₅₀>20 μM

cpd	IC ₅₀	
667		35
668	A	
669		
670	A	
671		
672	A	40
673		
674	A	
675		
676	A	45
677		
678	A	
679		
680	A	
681		50
682	A	
683	A	
684	A	
685	A	
686	A	55
687		
688	A	
689		
690	A	
691		60
692	A	
693		
694	A	
695	A	
696	A	65
697	A	
698	A	
699		
700	A	
701		70
702	A	

589

-continued


cpd	IC ₅₀	
703		
704	A	5
705	B	
706	A	
707		
708	A	
709		
710	A	10
711	A	
712	A	
713		
714	A	
715	A	
716	A	15
717	A	
718	A	
719 - En 1	A	
719 - En 2	A	
720 - En 1	A	
720 - En 2	A	20
721	A	
722 - En 1	A	
722 - En 2	A	
723 - En 1	A	
723 - En 2	A	
724 - En 1	A	
724 - En 2	B	25
725 - En 1	A	
725 - En 2	A	
726 - En 1	A	
726 - En 2	A	
727 - En 1	B	30
727 - En 2	B	
727 - En 3	B	
727 - En 4	B	
728 - En 1	B	
728 - En 2	A	
729 - En 1	A	
729 - En 2	A	35
730 - En 1	A	
730 - En 2	A	
731 - En 1	A	
731 - En 2	A	
732 - En 1	A	
732 - En 2	A	40
733 - En 1	A	
733 - En 2	A	
734 - En 1	A	
734 - En 2	B	
735 - En 1	A	
735 - En 2	A	45
736 - En 1	A	
736 - En 2	A	
737 - En 1	A	
737 - En 2	A	
738 - En 1	A	
738 - En 2	A	
739 - En 1	A	50
739 - En 2	A	
740 - En 1	A	
740 - En 2	A	
741	A	
742	A	
743	A	55
744	A	
745	A	
746	A	
747	A	
748	A	
749	A	60
750	A	
751	A	

The invention claimed is:

1. A compound of formula (I), a stereo-isomeric form, a physiologically acceptable salt, solvate and/or polymorph thereof

590

(I)



wherein

R¹ represents —F, —Cl, —Br, —I, —CN, —R^W, —OR^W, —OC(=O)R^W, —NR^WR^X, —NR^WC(=O)R^X, —SR^W, —S(=O)R^W, —S(=O)₂R^W, —C(=O)R^W, —C(=O)OR^W, or —C(=O)NR^WR^X;

Q represents —NR³R⁴;

R³ represents —OH or —R^Y;

R⁴ represents —R^Y or —S(=O)₂R^Y;

or R³ and R⁴ together form a 4, 5, 6, 7 or 8 membered heterocycle containing 1 to 3 heteroatoms selected from N, O and S, saturated or unsaturated, unsubstituted or mono- or polysubstituted;

T represents —O— and U represents —CR⁵R^{5t}—; or T represents —CR⁵R^{5t}— and U represents —O—;

R⁵ and R^{5t} independently of one another represent —R^Y;

R⁶, R⁷ and R⁸ independently of one another represent —F, —Cl, —Br, —I, —CN, —NO₂, —SF₅, —R^W, —OR^W, —OC(=O)R^W, —NR^WR^X, —NR^WC(=O)R^X, —SR^W, —S(=O)R^W, —S(=O)₂R^W, —C(=O)R^W, —C(=O)OR^W, or —C(=O)NR^WR^X;

V represents 3-14-membered heterocycloalkyl, saturated or unsaturated; or 5-14-membered heteroaryl; in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from —F, —Cl, —Br, —I, —CN, —NO, —NO₂, —O, —S, —SF₅, —R^Y, —OR^Y, —OC(=O)R^Y, —NR^YR^Z, —NR^YC(=O)R^Z, —SR^Y, —S(=O)R^Y, —S(=O)₂R^Y, —C(=O)R^Y, —C(=O)OR^Y, or —C(=O)NR^YR^Z;

wherein

R^W and R^X independently of one another in each case independently represent

—H;

—C₁-C₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;

—C₁-C₆-heteroalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;

3-14-membered cycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered cycloalkyl is optionally connected through —C₁-C₆-alkylene- or —C₁-C₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted; or

3-14-membered heterocycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;

wherein said 3-14-membered heterocycloalkyl is optionally connected through —C₁-C₆-alkylene- or —C₁-C₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted;

R^Y and R^Z independently of one another in each case independently represent


—H;

—C₁-C₆-alkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted;

kyl)(C₁₋₆-alkyl), —N(C₁₋₆-alkyl)-C₁₋₆-alkylene-OH, —N(H)—C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-NH(C₁₋₆-alkyl), —N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —NH—C(=O)—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH—C(=O)—C₁₋₆-alkyl, —NH—C(=O)—O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH—C(=O)—O—C₁₋₆-alkyl, —NH—C(=O)—NH₂, —C₁₋₆-alkylene-NH—C(=O)—NH₂, —NH—C(=O)—NH(C₁₋₆-alkyl), —C₁₋₆-alkylene-NH—C(=O)—NH(C₁₋₆-alkyl), —NH—C(=O)—N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-NH—C(=O)—N(C₁₋₆-alkyl)₂, —N(C₁₋₆-alkyl)-C(=O)—C₁₋₆-alkyl, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)-C(=O)—C₁₋₆-alkyl, —N(C₁₋₆-alkyl)-C(=O)—O—C₁₋₆-alkyl, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)-C(=O)—O—C₁₋₆-alkyl, —N(C₁₋₆-alkyl)-C(=O)—NH₂, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)-C(=O)—NH₂, —N(C₁₋₆-alkyl)-C(=O)—NH(C₁₋₆-alkyl), —C₁₋₆-alkylene-N(C₁₋₆-alkyl)-C(=O)—NH(C₁₋₆-alkyl), —N(C₁₋₆-alkyl)-C(=O)—NH(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)-C(=O)—NH(C₁₋₆-alkyl)₂, —NH—S(O)₂OH, —C₁₋₆-alkylene-NH—S(O)₂OH, —NH—S(O)₂—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH—S(O)₂—C₁₋₆-alkyl, —NH—S(O)₂—O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH—S(O)₂—O—C₁₋₆-alkyl, —NH—S(O)₂—NH₂, —C₁₋₆-alkylene-NH—S(O)₂—NH₂, —NH—S(O)₂—NH(C₁₋₆-alkyl), —C₁₋₆-alkylene-NH—S(O)₂—NH(C₁₋₆-alkyl), —NH—S(O)₂N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-NH—S(O)₂N(C₁₋₆-alkyl)₂, —N(C₁₋₆-alkyl)-S(=O)₂—OH, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)-S(=O)₂—OH, —N(C₁₋₆-alkyl)-S(=O)₂—C₁₋₆-alkyl, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)-S(=O)₂—C₁₋₆-alkyl, —N(C₁₋₆-alkyl)-S(=O)₂—O—C₁₋₆-alkyl, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)-S(=O)₂—O—C₁₋₆-alkyl, —N(C₁₋₆-alkyl)-S(=O)₂—NH₂, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)-S(=O)₂—NH₂, —N(C₁₋₆-alkyl)-S(=O)₂—NH(C₁₋₆-alkyl), —C₁₋₆-alkylene-N(C₁₋₆-alkyl)-S(=O)₂—NH(C₁₋₆-alkyl), —N(C₁₋₆-alkyl)-S(=O)₂—N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)-S(=O)₂—N(C₁₋₆-alkyl)₂, —SH, =S, —SF₅, —SCF₃, —SCF₂H, —SCFH₂, —S—C₁₋₆-alkyl, —C₁₋₆-alkylene-S—C₁₋₆-alkyl, —S(=O)—C₁₋₆-alkyl, —C₁₋₆-alkylene-S(=O)—C₁₋₆-alkyl, —S(=O)₂—C₁₋₆-alkyl, —C₁₋₆-alkylene-S(=O)₂—C₁₋₆-alkyl, —S(=O)₂—OH, —C₁₋₆-alkylene-S(=O)₂—OH, —S(=O)₂—O—C₁₋₆-alkyl, —C₁₋₆-alkylene-S(=O)₂—O—C₁₋₆-alkyl, —S(=O)₂—NH₂, —C₁₋₆-alkylene-S(=O)₂—NH₂, —S(=O)₂—NH(C₁₋₆-alkyl), —C₁₋₆-alkylene-S(=O)₂—NH(C₁₋₆-alkyl), —S(=O)₂—N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-S(=O)₂—N(C₁₋₆-alkyl)₂, 3-14-membered cycloalkyl, —C₁₋₆-alkylene-(3-14-membered cycloalkyl), 3 to 14-membered heterocycloalkyl, —C₁₋₆-alkylene-(3 to 14-membered heterocycloalkyl), —phenyl, —C₁₋₆-alkylene-phenyl, 5 to 14-membered heteroaryl, —C₁₋₆-alkylene-(5 to 14-membered heteroaryl), —O-(3-14-membered cycloalkyl), —O-(3 to 14-membered heterocycloalkyl), —O-phenyl, —O-(5 to 14-membered heteroaryl), —C(=O)-(3-14-membered cycloalkyl), —C(=O)-(3 to 14-membered hetero-cycloalkyl), —C(=O)-phenyl, —C(=O)-(5 to 14-membered heteroaryl), —S(=O)₂-(3-14-membered cycloalkyl), —S(=O)₂-(3 to 14-membered heterocycloalkyl), —S(=O)₂-phenyl, —S(=O)₂-(5 to 14-membered heteroaryl);

with the proviso that the following compounds and their salts are excluded

593



2. A compound of formula (I), a stereo-isomeric form, a physiologically acceptable salt, solvate and/or polymorph thereof, as defined in claim 1, wherein R^3 represents —H.

3. A compound of formula (I), a stereo-isomeric form, a physiologically acceptable salt, solvate and/or polymorph thereof, as defined in claim 1, wherein R^1 represents —CH₂F, —CHF₂, or —CF₃.

4. The compound according to claim 1, wherein T represents —O— and U represents —CR⁵R⁵—.

5. The compound according to claim 1, wherein V represents (i) 5-14-membered heteroaryl selected from benzimidazole, benzisoxazole, benzoazole, benzodioxole, benzofuran, benzothiadiazole, benzothiazole, benzothiophene, carbazole, cinnoline, dibenzofuran, furane, furazane, imidazole, imidazopyridine, indazole, indole, indolizine, isobenzofuran, isoindole, isoquinoline, isothiazole, isoxazole, naphthyridine, oxadiazole, oxazole, oxindole, phthalazine, purine, pyrazine, pyrazole, pyridazine, pyridine, pyrimidine, pyrrole, quinazoline, quinoline, quinoxaline, tetrazole, thiadiazole, thiazole, thiophene, triazine, triazole, and [1,2,4] triazole[4,3-a]pyrimidine; in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from —F, —Cl, —CN, —OH, —O, —C₁₋₆-alkyl, —CHF₂, —CF₃, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NHC(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-CHF₂, —C₁₋₆-alkylene-CF₃, —C₁₋₆-alkylene-cyclopropyl, —cyclopropyl, —O-cyclopropyl, —C₁₋₆-alkylene-NHC(=O)—O—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkyl, —N(C₁₋₆-alkyl)₂, —OCF₃, —O—C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —S(=O)₂—C₁₋₆-alkyl, —azetidine, —C₁₋₆-alkylene-O-tetrahydropyran, or —piperazine substituted with —C₁₋₆-alkyl; particularly in each

594

case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from —F, —Cl, —CN, —OH, —O, —C₁₋₆-alkyl, —CHF₂, —CF₃, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NHC(=O)O—C₁₋₆-alkyl, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-NHC(=O)O—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkyl, —N(C₁₋₆-alkyl)₂, —OCF₃, —O—C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —S(=O)₂—C₁₋₆-alkyl, —azetidine, —C₁₋₆-alkylene-O-tetrahydropyran, or —piperazine substituted with —C₁₋₆-alkyl; or represents (ii) —oxetanyl, unsubstituted, mono- or polysubstituted.

6. The compound according to claim 1, wherein R^1 represents —H, —F, —Cl, —Br, —I, —C₁₋₆-alkyl, —O—C₁₋₆-alkyl, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH(C₁₋₆-alkyl), —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —CF₃, —CF₂H, —CFH₂, —CF₂Cl, —CFCl₂, —C₁₋₆-alkylene-CF₃, —C₁₋₆-alkylene-CF₂H, —C₁₋₆-alkylene-CFH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)-C₁₋₆-alkylene-CF₃, —C(=O)C₁₋₆-alkyl, —C(=O)OC₁₋₆-alkyl, —C(=O)NH₂, —C(=O)NHC₁₋₆-alkyl, —C(=O)N(C₁₋₆-alkyl)₂, —S(=O)—C₁₋₆-alkyl, —S(=O)₂—C₁₋₆-alkyl, —O—C₁₋₆-alkyl, —cyclopropyl unsubstituted, cyclobutyl unsubstituted, cyclopentyl unsubstituted or cyclohexyl unsubstituted.

7. The compound according to claim 1, wherein R^3 represents —H, —OH, —C₁₋₆-alkyl, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NH(C₁₋₆-alkyl), —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —CF₃, —CF₂H, —CFH₂, —CF₂Cl, —CFCl₂, —C₁₋₆-alkylene-CF₃, —C₁₋₆-alkylene-CF₂H, —C₁₋₆-alkylene-CFH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, or —C₁₋₆-alkylene-N(C₁₋₆-alkyl)-C₁₋₆-alkylene-CF₃.

8. The compound according to claim 1, wherein R^4 represents

—H;
—S(=O)₂C₁₋₆-alkyl, saturated, unsubstituted, monosubstituted or polysubstituted with —F;
—S(=O)₂(3-14-membered cycloalkyl), saturated, unsubstituted;
—C₁₋₆-alkyl, saturated, unsubstituted, monosubstituted or disubstituted with substituents independently of one another selected from the group consisting of —OH, —O, —OC₁₋₆-alkyl, —NH₂, —NHC₁₋₆-alkyl, —N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkyl, —C(=O)NH₂, —C(=O)—NH—C₁₋₆-alkyl, —C(=O)N(C₁₋₃-alkyl)₂, —phenyl unsubstituted;

3-14-membered cycloalkyl or —C₁₋₆-alkylene-(3-14-membered cycloalkyl), wherein —C₁₋₆-alkylene- is unsubstituted or monosubstituted with —OH, wherein said 3-14-membered cycloalkyl is saturated, unsubstituted, monosubstituted or disubstituted with substituents independently of one another selected from the group consisting of —C₁₋₆-alkyl, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-NH—C₁₋₆-alkylene-CF₃, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-NHC(=O)O—C₁₋₆-alkyl, —OH, —OC₁₋₆-alkyl, —NH₂, —N(C₁₋₆-alkyl)₂, —NHC(=O)O—C₁₋₆-alkyl;

3-14-membered heterocycloalkyl or —C₁₋₆-alkylene-(3-14-membered heterocycloalkyl), wherein —C₁₋₆-alkylene- is unsubstituted or monosubstituted with —OH, wherein said 3-14-membered heterocycloalkyl in each case is selected from azetane, 1,4-oxazepane, pyrrolidine, piperidine, azepane, diazepane, tetrahydrofuran, tetrahydropyran, oxetane, morpholine, piperazine, hexahydrocyclopenta[c]pyrrole, octahydrocyclopenta[c]pyrrole, octahydropyrrolo[1,2-a]pyrazine,

595

8-azabicyclo[3.2.1]octane, 9-azabicyclo[3.3.1]nonane, quinuclidine, hexahydro-1H-pyrrolizine, 2-oxaspiro [3.3]heptane, 2-azaspiro[3.3]heptane, 7-azaspiro[3.5] nonane, 1,1-dioxothiacyclohexane, in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from the group consisting of —F, —OH, =O, —C₁₋₆-alkyl, —C₁₋₆-alkylene-CF₃, —C₁₋₆-alkylene-OH, —C₁₋₆-alkylene-O—C₁₋₆-alkyl, —NH₂, —N(C₁₋₆-alkyl)₂, —C₁₋₆-alkylene-NH₂, —C₁₋₆-alkylene-N(C₁₋₆-alkyl)₂, —C(=O)—C₁₋₆-alkyl, —C(=O)OH, —C(=O)O—C₁₋₆-alkyl, —C(=O)O—C₁₋₆-alkylene-CF₃, —C(=O)NH₂, —C(=O)NH(C₁₋₆-alkyl), —S(=O)₂ C₁₋₆-alkyl, oxetanyl, pyrimidinyl, —C₁₋₆-alkylene-phenyl; 15
-phenyl unsubstituted;
5-14-membered heteroaryl or —C₁₋₆-alkylene-(5-14-membered heteroaryl), wherein —C₁₋₆-alkylene- is unsubstituted or monosubstituted with —OH, wherein said 5-14-membered heteroaryl in each case is selected from the group consisting of pyridine, pyridazine, pyrazine, pyrazole, isoxazole, triazole, and [1,2,4]triazolo[4,3-a]pyrimidine, in each case unsubstituted, monosubstituted or disubstituted with substituents independently of one another selected from the group consisting of —C₁₋₆-alkyl, —OH.

9. The compound according to claim 1, wherein R³ and R⁴ together form a heterocycle selected from the group consisting of pyrrolidine, piperidine, morpholine, and piperazine, in each case unsubstituted, mono- or polysubstituted with substituents independently of one another selected from the group consisting of —F, —C₁₋₆-alkyl, —NH₂, —NHCH₃, —N(CH₃)₂, —C(=O)NH—C₁₋₆-alkyl, —C(=O)N(C₁₋₆-alkyl)₂, —C(=O)O—C₁₋₆-alkyl, —NHC(=O)O—C₁₋₆-alkyl, —pyridyl unsubstituted, and 1,2,4-oxadiazole unsubstituted or monosubstituted with —C₁₋₆-alkyl.

10. The compound according to claim 1, wherein R⁵ and R^{5'} independently of one another represent

—H;
—C₁₋₆-alkyl, saturated or unsaturated, unsubstituted, mono- or poly substituted;
—C₁₋₆-heteroalkyl, saturated or unsaturated, unsubstituted, mono- or poly substituted;

3-14-membered cycloalkyl, saturated or unsaturated, unsubstituted, mono- or polysubstituted; wherein said 3-14-membered cycloalkyl is optionally connected through —C₁₋₆-alkylene- or —C₁₋₆-heteroalkylene-, in each case saturated or unsaturated, unsubstituted, mono- or polysubstituted.

11. The compound according to claim 1, wherein R⁶, R⁷ and R⁸ independently of one another represent

—H, —F, —Cl, —Br, —I, —OH, —SH, —SF₅, —CN, —NO₂, —C(=O)OH, —NH₂,
—C₁₋₆-alkyl, —CF₃, —CHF₂, —CH₂F,
—O—C₁₋₆-alkyl, —OCF₃, —OCHF₂, —OCH₂F,
—NHC₁₋₆-alkyl unsubstituted or substituted with one or more substituents independently of one another selected from —OH, =O, —F, —Cl, —Br, —I, —SH, =S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, and —C(=O)NH₂;
—N(C₁₋₆-alkyl)₂ unsubstituted or substituted with one or more substituents independently of one another selected from —OH, =O, —F, —Cl, —Br, —I, —SH, =S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, and —C(=O)NH₂;

596

—C(=O)OC₁₋₆-alkyl unsubstituted or substituted with one or more substituents independently of one another selected from —OH, =O, —F, —Cl, —Br, —I, —SH, =S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, and —C(=O)NH₂;
—OC(=O)C₁₋₆-alkyl unsubstituted or substituted with one or more substituents independently of one another selected from —OH, =O, —F, —Cl, —Br, —I, —SH, =S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, and —C(=O)NH₂; or
—C₁₋₆-heteroalkyl unsubstituted or substituted with one or more substituents independently of one another selected from —OH, =O, —F, —Cl, —Br, —I, —SH, =S, —CN, —CF₃, —CHF₂, —CH₂F, —OCF₃, —OCHF₂, —OCH₂F, SF₅, —NO₂, —C(=O)OH, —NH₂, and —C(=O)NH₂.

12. The compound according to claim 1, which is selected from the group consisting of

cpd 001 5-(furan-3-ylmethoxy)-2-methylbenzofuran-3-carboxylic acid
cpd 002 2-methyl-5-(oxazol-5-ylmethoxy)benzofuran-3-carboxylic acid
cpd 003 2-methyl-5-(pyrrolidin-3-ylmethoxy)benzofuran-3-carboxylic acid
cpd 004 2-methyl-5-(pyridin-4-ylmethoxy)benzofuran-3-carboxylic acid
cpd 005 2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxylic acid
cpd 006 2-methyl-5-(pyridin-3-ylmethoxy)benzofuran-3-carboxylic acid
cpd 007 2-methyl-5-(pyridazin-3-ylmethoxy)benzofuran-3-carboxylic acid
cpd 008 2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxylic acid
cpd 009 2-methyl-5-((1-methyl-1H-pyrazol-3-yl)methoxy)benzofuran-3-carboxylic acid
cpd 010 2-methyl-5-((1-methyl-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxylic acid
cpd 011 2-methyl-5-(thiophen-3-ylmethoxy)benzofuran-3-carboxylic acid
cpd 012 2-methyl-5-(thiophen-2-ylmethoxy)benzofuran-3-carboxylic acid
cpd 013 2-methyl-5-(thiazol-2-ylmethoxy)benzofuran-3-carboxylic acid
cpd 014 2-methyl-5-(thiazol-5-ylmethoxy)benzofuran-3-carboxylic acid
cpd 015 2-methyl-5-((tetrahydro-2H-pyran-4-yl)methoxy)benzofuran-3-carboxylic acid
cpd 016 5-((3,5-dimethylisoxazol-4-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
cpd 017 5-(benzo[d]thiazol-2-ylmethoxy)-2-methylbenzofuran-3-carboxylic acid
cpd 018 2-methyl-5-((5-(trifluoromethyl)furan-2-yl)methoxy)benzofuran-3-carboxylic acid
cpd 019 5-((1-(tert-butoxycarbonyl)pyrrolidin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
cpd 020 (S)-2-methyl-5-(oxazol-5-ylmethoxy)-N-(pyrrolidin-3-yl)benzofuran-3-carboxamide
cpd 021 (S)-2-methyl-5-(pyridin-4-ylmethoxy)-N-(pyrrolidin-3-yl)benzofuran-3-carboxamide
cpd 022 (S)-2-methyl-5-(pyridin-3-ylmethoxy)-N-(pyrrolidin-3-yl)benzofuran-3-carboxamide
cpd 023 (S)-2-methyl-5-((1-methyl-1H-pyrazol-3-yl)methoxy)-N-(pyrrolidin-3-yl)benzofuran-3-carboxamide

597

cpd 024 (S)-2-methyl-N-(pyrrolidin-3-yl)-5-(thiazol-5-yl-methoxy)benzofuran-3-carboxamide
 cpd 025 (S)-2-methyl-N-(pyrrolidin-3-yl)-5-(thiazol-2-yl-methoxy)benzofuran-3-carboxamide
 cpd 026 2-methyl-N-(2-oxopyrrolidin-3-yl)-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide 5
 cpd 027 2-methyl-N-(2-oxopyrrolidin-3-yl)-5-(pyridin-3-ylmethoxy)benzofuran-3-carboxamide
 cpd 028 (S)-2-methyl-5-((5-methylpyridin-3-yl)methoxy)-N-(pyrrolidin-3-yl)benzofuran-3-carboxamide 10
 cpd 029 (S)-2-methyl-5-((6-methylpyridin-2-yl)methoxy)-N-(pyrrolidin-3-yl)benzofuran-3-carboxamide
 cpd 030 2-methyl-N-(2-oxopyrrolidin-3-yl)-5-(pyridazin-3-ylmethoxy)benzofuran-3-carboxamide 15
 cpd 031 2-methyl-N-(2-oxopyrrolidin-3-yl)-5-(pyrazin-2-ylmethoxy)benzofuran-3-carboxamide
 cpd 032 2-methyl-N-(2-oxopyrrolidin-3-yl)-5-(pyrimidin-5-ylmethoxy)benzofuran-3-carboxamide 20
 cpd 033 2-methyl-5-((6-methylpyridin-3-yl)methoxy)-N-(tetrahydrofuran-3-yl)benzofuran-3-carboxamide
 cpd 034 2-methyl-5-((4-methylpyridin-3-yl)methoxy)-N-(tetrahydrofuran-3-yl)benzofuran-3-carboxamide
 cpd 035 2-methyl-N-(2-oxopyrrolidin-3-yl)-5-(thiophen-3-ylmethoxy)benzofuran-3-carboxamide 25
 cpd 036 2-methyl-N-(2-oxopyrrolidin-3-yl)-5-(thiazol-2-ylmethoxy)benzofuran-3-carboxamide
 cpd 037 2-methyl-5-((4-methylpyridin-3-yl)methoxy)-N-(2-oxopyrrolidin-3-yl)benzofuran-3-carboxamide 30
 cpd 038 2-methyl-5-((5-methylpyridin-3-yl)methoxy)-N-(2-oxopyrrolidin-3-yl)benzofuran-3-carboxamide
 cpd 039 2-methyl-N-(1-methylpiperidin-4-yl)-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
 cpd 040 5-((4-methoxypyridin-3-yl)methoxy)-2-methyl-N-(tetrahydrofuran-3-yl)benzofuran-3-carboxamide 35
 cpd 041 2-methyl-N-(1-methylpiperidin-4-yl)-5-(thiophen-2-ylmethoxy)benzofuran-3-carboxamide
 cpd 042 2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(2-oxopyrrolidin-3-yl)benzofuran-3-carboxamide 40
 cpd 043 2-methyl-5-((4-methylthiazol-2-yl)methoxy)-N-(2-oxopyrrolidin-3-yl)benzofuran-3-carboxamide
 cpd 044 5-((2,6-dimethylpyridin-3-yl)methoxy)-2-methyl-N-(2-oxopyrrolidin-3-yl)benzofuran-3-carboxamide 45
 cpd 045 2-methyl-N-(1-methylpiperidin-4-yl)-5-(1-(pyridin-4-yl)ethoxy)benzofuran-3-carboxamide
 cpd 046 5-((2-methoxypyridin-3-yl)methoxy)-2-methyl-N-(2-oxopyrrolidin-3-yl)benzofuran-3-carboxamide
 cpd 047 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide 50
 cpd 048 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
 cpd 049 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-(thiazol-2-ylmethoxy)benzofuran-3-carboxamide 55
 cpd 050 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-(thiazol-2-ylmethoxy)benzofuran-3-carboxamide
 cpd 051 5-((2-methoxypyridin-3-yl)methoxy)-2-methyl-N-(1-methylpiperidin-4-yl)benzofuran-3-carboxamide 60
 cpd 052 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylpyridin-3-yl)methoxy)-benzofuran-3-carboxamide
 cpd 053 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((3-methylpyridin-2-yl)methoxy)benzofuran-3-carboxamide 65

598

cpd 054 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)-benzofuran-3-carboxamide
 cpd 055 (S)-2-methyl-N-(pyrrolidin-3-yl)-5-((5-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
 cpd 056 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
 cpd 057 2-methyl-N-(tetrahydrofuran-3-yl)-5-((4-(trifluoromethyl)thiazol-2-yl)methoxy)benzofuran-3-carboxamide
 cpd 058 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
 cpd 059 5-(isoquinolin-1-ylmethoxy)-2-methyl-N-(1-methylpiperidin-4-yl)benzofuran-3-carboxamide
 cpd 060 5-(isoquinolin-5-ylmethoxy)-2-methyl-N-(1-methylpiperidin-4-yl)benzofuran-3-carboxamide
 cpd 061 5-(isoquinolin-6-ylmethoxy)-2-methyl-N-(1-methylpiperidin-4-yl)benzofuran-3-carboxamide
 cpd 062 N-(3,3-difluoropiperidin-4-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
 cpd 063 N-(3,3-difluoropiperidin-4-yl)-5-((6-methoxypyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide
 cpd 064 2-methyl-N-(2-oxopyrrolidin-3-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
 cpd 065 2-methyl-N-(2-oxopyrrolidin-3-yl)-5-((6-(trifluoromethyl)pyridin-2-yl)methoxy)benzofuran-3-carboxamide
 cpd 066 2-methyl-N-(2-oxopyrrolidin-3-yl)-5-((6-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
 cpd 067 5-((2-chloropyridin-3-yl)methoxy)-N-(3,3-difluoropiperidin-4-yl)-2-methylbenzofuran-3-carboxamide
 cpd 068 5-((4-chloropyridin-3-yl)methoxy)-N-(3,3-difluoropiperidin-4-yl)-2-methylbenzofuran-3-carboxamide
 cpd 069 2-methyl-N-(1-methylpiperidin-4-yl)-5-((5-(trifluoromethyl)furan-2-yl)methoxy)benzofuran-3-carboxamide
 cpd 070 N-(3,3-difluoropiperidin-4-yl)-5-(imidazo[1,2-a]pyridin-6-ylmethoxy)-2-methylbenzofuran-3-carboxamide
 cpd 071 N-(3,3-difluoropiperidin-4-yl)-5-((6-(dimethylamino)pyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide
 cpd 072 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
 cpd 073 tert-butyl (S)-3-(2-methyl-5-((5-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamido)-pyrrolidine-1-carboxylate
 cpd 074 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
 cpd 075 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((5-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
 cpd 076 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((3-(trifluoromethyl)pyridin-2-yl)methoxy)benzofuran-3-carboxamide

599

- cpd 077 tert-butyl (S)-3-(2-methyl-5-((5-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate
- cpd 078 tert-butyl 4-(5-((6-(dimethylamino)pyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamido)-3,3-difluoropiperidine-1-carboxylate 5
- cpd 079 tert-butyl 3-(((3-(ethoxycarbonyl)-2-methylbenzofuran-5-yl)oxy)methyl)pyrrolidine-1-carboxylate
- cpd 080 tert-butyl (R)-3-(2-methyl-5-(oxazol-5-yl-methoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate 10
- cpd 081 tert-butyl (R)-3-(2-methyl-5-(pyridin-4-yl-methoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate
- cpd 082 tert-butyl (R)-3-(2-methyl-5-(pyridin-3-yl-methoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate 15
- cpd 083 tert-butyl (R)-3-(2-methyl-5-((1-methyl-1H-pyrazol-3-yl)methoxy)benzofuran-3-carboxamido)-pyrrolidine-1-carboxylate 20
- cpd 084 tert-butyl (R)-3-(2-methyl-5-(thiazol-5-yl-methoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate
- cpd 085 tert-butyl (R)-3-(2-methyl-5-(thiazol-2-yl-methoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate 25
- cpd 086 tert-butyl (R)-3-(2-methyl-5-((6-methylpyridin-2-yl)methoxy)benzofuran-3-carboxamido)-pyrrolidine-1-carboxylate
- cpd 087 tert-butyl 3,3-difluoro-4-(2-methyl-5-(thiazol-2-ylmethoxy)benzofuran-3-carboxamido)piperidine-1-carboxylate 30
- cpd 088 tert-butyl 3,3-difluoro-4-(2-methyl-5-((3-methylpyridin-2-yl)methoxy)benzofuran-3-carboxamido)piperidine-1-carboxylate 35
- cpd 089 tert-butyl 3,3-difluoro-4-(5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamido)piperidine-1-carboxylate
- cpd 090 tert-butyl 3,3-difluoro-4-(2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)piperidine-1-carboxylate 40
- cpd 091 tert-butyl 3,3-difluoro-4-(5-((6-methoxypyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamido)piperidine-1-carboxylate
- cpd 092 tert-butyl 4-(5-((2-chloropyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamido)-3,3-difluoropiperidine-1-carboxylate 45
- cpd 093 tert-butyl 4-(5-((4-chloropyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamido)-3,3-difluoropiperidine-1-carboxylate 50
- cpd 094 tert-butyl 3,3-difluoro-4-(5-(imidazo[1,2-a]pyridin-6-ylmethoxy)-2-methylbenzofuran-3-carboxamido)piperidine-1-carboxylate
- cpd 095 tert-butyl 3,3-difluoro-4-(2-methyl-5-((5-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)piperidine-1-carboxylate 55
- cpd 096 tert-butyl 3,3-difluoro-4-(2-methyl-5-((3-(trifluoromethyl)pyridin-2-yl)methoxy)benzofuran-3-carboxamido)piperidine-1-carboxylate
- cpd 097 tert-butyl 3,3-difluoro-4-(2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamido)piperidine-1-carboxylate 60
- cpd 098 tert-butyl 3,3-difluoro-4-(2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamido)piperidine-1-carboxylate 65
- cpd 099 2-methyl-5-((4-methylpyridin-3-yl)methoxy)benzofuran-3-carboxylic acid

600

- cpd 100 5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- cpd 101 2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid
- cpd 102 2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid
- cpd 103 N-(4,4-difluoropiperidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- cpd 104 N-(1-hydroxy-2-methylpropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- cpd 105 N-(4,4-difluoropyrrolidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
- cpd 106 N-(4,4-difluoropyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- cpd 107 N-(1-(hydroxymethyl)cyclobutyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- cpd 108 N-(1-((dimethylamino)methyl)cyclobutyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- cpd 109 N-(1-hydroxy-2-methylpropan-2-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
- cpd 110 N-(3,3-difluoro-1-methylpiperidin-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- cpd 111 N-(4,4-difluoropiperidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
- cpd 112 N-(3,3-difluoropiperidin-4-yl)-4-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
- cpd 113 N-(3,3-difluoropiperidin-4-yl)-6-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
- cpd 114 N-(1-(2-hydroxyethyl)cyclobutyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- cpd 115 N-(5,5-difluoropiperidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- cpd 116 2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(1-(2,2,2-trifluoroethyl)pyrrolidin-3-yl)benzofuran-3-carboxamide
- cpd 117 N-(1-(hydroxymethyl)cyclobutyl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
- cpd 118 N-(4,4-difluoro-1-methylpiperidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- cpd 119 N-(3-carbamoyloxetan-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- cpd 120 N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
- cpd 121 N-(3-(2-hydroxyethyl)tetrahydrofuran-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)-benzofuran-3-carboxamide
- cpd 122 7-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide

601

- cpd 123 N-(4,4-difluoro-1-methylpyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- cpd 124 N-(4,4-difluoro-1-methylpyrrolidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide 5
- cpd 125 N-(5,5-difluoropiperidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
- cpd 126 N-(3-(2-hydroxyethyl)tetrahydrofuran-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide 10
- cpd 127 N-(1-(2-hydroxyethyl)cyclobutyl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide 15
- cpd 128 N-(3-(2-hydroxyethyl)oxetan-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- cpd 129 N-(3-(2-hydroxyethyl)oxetan-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide 20
- cpd 130 N-(3,3-difluoro-1-methylpiperidin-4-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide 25
- cpd 131 N-(4,4-difluoro-1-methylpiperidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
- cpd 132 N-(3-(hydroxymethyl)tetrahydrofuran-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)-benzofuran-3-carboxamide 30
- cpd 133 N-(3-(hydroxymethyl)oxetan-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide 35
- cpd 134 N-(4,4-difluoro-1-(2-hydroxyethyl)pyrrolidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
- cpd 135 N-(3-(hydroxymethyl)tetrahydrofuran-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide 40
- cpd 136 N-(1-(aminomethyl)cyclobutyl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
- cpd 137 N-(1-(aminomethyl)cyclobutyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 45
- cpd 138 2-methyl-N-(1H-pyrazol-3-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide 50
- cpd 139 N-(1-(hydroxymethyl)cyclopropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- cpd 140 N-(1-((dimethylamino)methyl)cyclobutyl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide 55
- cpd 141 N-(5,5-difluoro-1-methylpiperidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
- cpd 142 N-(3-(hydroxymethyl)oxetan-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 60
- cpd 143 2-methyl-N-(1-methyl-1H-pyrazol-4-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide 65
- cpd 144 2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(1H-pyrazol-3-yl)benzofuran-3-carboxamide

602

- cpd 145 2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(1-(((2,2,2-trifluoroethyl)amino)methyl)-cyclobutyl)benzofuran-3-carboxamide
- cpd 146 2-methyl-N-(1-methyl-1H-pyrazol-3-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
- cpd 147 N-(4-hydroxytetrahydrofuran-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
- cpd 148 2-methyl-N-(tetrahydrofuran-3-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
- cpd 149 2-methyl-N-(1-(2,2,2-trifluoroethyl)pyrrolidin-3-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
- cpd 150 2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(1H-pyrazol-4-yl)benzofuran-3-carboxamide
- cpd 151 N-(4,4-difluoro-1-(2-hydroxyethyl)pyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)-benzofuran-3-carboxamide
- cpd 152 2-methyl-N-(1-methyl-1H-pyrazol-4-yl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- cpd 153 2-methyl-N-(1-(((2,2,2-trifluoroethyl)amino)methyl)cyclobutyl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
- cpd 154 2-methyl-N-(1-methyl-1H-pyrazol-5-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
- cpd 155 N-(4,4-difluoro-1-(2,2,2-trifluoroethyl)pyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- cpd 156 2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(2-oxaspiro[3.3]heptan-6-yl)benzofuran-3-carboxamide
- cpd 157 N-(4,4-difluoro-1-(2,2,2-trifluoroethyl)pyrrolidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
- cpd 158 2-methyl-N-(1H-pyrazol-4-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
- cpd 159 2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(tetrahydrofuran-3-yl)benzofuran-3-carboxamide
- cpd 160 N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)-benzofuran-3-carboxamide
- cpd 161 2-methyl-N-(1-methyl-1H-pyrazol-3-yl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- cpd 162 N-(4-hydroxytetrahydrofuran-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- cpd 163 N-(5,5-difluoro-1-methylpiperidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- cpd 164 N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((2-methylpyridin-3-yl)methoxy)-benzofuran-3-carboxamide
- cpd 165 6-chloro-N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
- cpd 166 2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)benzofuran-3-carboxamide
- cpd 167 2-methyl-N-(2-oxaspiro[3.3]heptan-6-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide

603

- cpd 168 2-methyl-N-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
- cpd 169 N-(1-(2-methoxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)-benzofuran-3-carboxamide 5
- cpd 170 (3-aminotetrahydrofuran-3-yl)methyl 2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxylate
- cpd 171 2-methyl-N-(1-methyl-1H-pyrazol-5-yl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 10
- cpd 172 2,2,2-trifluoroethyl 3,3-difluoro-4-(2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamido)pyrrolidine-1-carboxylate 15
- cpd 173 2-methyl-N-(2-azaspiro[3.3]heptan-6-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
- cpd 174 6-chloro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 20
- cpd 175 2,2,2-trifluoroethyl (R)-3,3-difluoro-4-(2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamido)pyrrolidine-1-carboxylate 25
- cpd 176 N-(3,3-difluoropiperidin-4-yl)-7-fluoro-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)-methoxy)benzofuran-3-carboxamide
- cpd 177 4-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)-methoxy)benzofuran-3-carboxamide 30
- cpd 178 (3-amino-1H-pyrazol-1-yl)(2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-yl)methanone
- cpd 179 N-(3,3-difluoropiperidin-4-yl)-2,6-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide 35
- cpd 180 N-(1-(2-hydroxyethyl)-5-oxopyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)-benzofuran-3-carboxamide 40
- cpd 181 (4-amino-1H-pyrazol-1-yl)(2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-yl)methanone
- cpd 182 6-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)-methoxy)benzofuran-3-carboxamide 45
- cpd 183 N-(3,3-difluoropiperidin-4-yl)-2,7-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
- cpd 184 7-chloro-N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide 50
- cpd 185 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2,6-dimethyl-5-((4-methylthiazol-5-yl)methoxy)-benzofuran-3-carboxamide 55
- cpd 186 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2,7-dimethyl-5-((4-methylthiazol-5-yl)-methoxy)benzofuran-3-carboxamide
- cpd 187 7-chloro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 60
- cpd 188 7-cyano-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- cpd 189 N-((3-aminooxetan-3-yl)methyl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide 65

604

- cpd 190 2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(2-azaspiro[3.3]heptan-6-yl)benzofuran-3-carboxamide
- cpd 191 4-cyano-N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)-methoxy)benzofuran-3-carboxamide
- cpd 192 (3-amino-1H-pyrazol-1-yl)(2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-yl)-methanone
- cpd 193 4-cyano-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- cpd 194 6-cyano-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- cpd 195 6-cyano-N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)-methoxy)benzofuran-3-carboxamide
- cpd 196 7-cyano-N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)-methoxy)benzofuran-3-carboxamide
- cpd 197 2-(difluoromethyl)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((4-methylthiazol-5-yl)-methoxy)benzofuran-3-carboxamide
- cpd 198 2-(difluoromethyl)-N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-5-((4-methylthiazol-5-yl)-methoxy)benzofuran-3-carboxamide
- cpd 199 2-(difluoromethyl)-N-(3,3-difluoropiperidin-4-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
- cpd 200 N-(1-(hydroxymethyl)cyclopropyl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
- cpd 201 N-(3-(hydroxymethyl)azetidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- cpd 202 N-(3-(hydroxymethyl)azetidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)-benzofuran-3-carboxamide
- cpd 203 N-(4,4-difluorotetrahydrofuran-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- cpd 204 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)-methoxy)benzofuran-3-carboxamide
- cpd 205 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide
- cpd 206 5-((2-ethoxypyridin-3-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
- cpd 207 N-(3,3-difluoropiperidin-4-yl)-5-((2-ethoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 208 2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 209 2-methyl-5-((2-(trifluoromethyl)thiazol-5-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 210 2-methyl-5-((2-methyl-4-(trifluoromethyl)thiazol-5-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 211 2-methyl-5-((2-methylloxazol-5-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 212 2-methyl-5-((2-methylthiazol-4-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 213 2-methyl-5-((3-(trifluoromethyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxylic acid

605

- Cpd 214 2-methyl-5-((4-methylpyridin-3-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 215 2-methyl-5-((5-methylisoxazol-3-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 216 2-methyl-5-((5-methylthiazol-4-yl)methoxy)benzofuran-3-carboxylic acid ⁵
- Cpd 217 2-methyl-5-((6-methylpyridin-3-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 218 2-methyl-5-(oxetan-3-ylmethoxy)benzofuran-3-carboxylic acid ¹⁰
- Cpd 219 2-methyl-5-(pyrimidin-4-ylmethoxy)benzofuran-3-carboxylic acid
- Cpd 220 5-((1-(2,2-difluoroethyl)-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid ¹⁵
- Cpd 221 5-((2-(difluoromethoxy)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 222 5-((2-(dimethylamino)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 223 5-((2,4-dimethylthiazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid ²⁰
- Cpd 224 5-((2,5-dimethylthiazol-4-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 225 5-((2-aminopyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid ²⁵
- Cpd 226 5-((3-(difluoromethyl)pyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 227 5-((6-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 228 5-((4-fluoro-1-methyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid ³⁰
- Cpd 229 2-(difluoromethyl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 230 2-(difluoromethyl)-5-((2-methoxypyridin-3-yl)methoxy)benzofuran-3-carboxylic acid ³⁵
- Cpd 231 2-cyclopropyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 232 2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 233 4-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid ⁴⁰
- Cpd 234 4-fluoro-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 235 5-((1-(2,2-difluoroethyl)-1H-imidazol-2-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid ⁴⁵
- Cpd 236 5-((2-(azetidin-1-yl)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 237 5-((2-(difluoromethyl)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 238 5-((2-cyanopyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid ⁵⁰
- Cpd 239 5-((2-ethoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 240 5-((3-fluoropyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid ⁵⁵
- Cpd 241 5-((5-(2-hydroxyethoxy)pyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 242 5-((5-fluoropyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 243 6-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid ⁶⁰
- Cpd 244 6-fluoro-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 245 7-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid ⁶⁵
- Cpd 246 7-fluoro-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid

606

- Cpd 247 2-methyl-5-((2-methylpyridin-3-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 248 2-methyl-5-(1-(2-(trifluoromethyl)thiazol-5-yl)ethoxy)benzofuran-3-carboxylic acid
- Cpd 249 2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 250 2-methyl-5-((1-methyl-2-oxo-1,2-dihydropyridin-3-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 251 2,4-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 252 2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)oxy)methylbenzofuran-3-carboxylic acid
- Cpd 253 2-methyl-5-((1-methyl-1H-1,2,3-triazol-4-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 254 2-methyl-5-((1-methyl-1H-1,2,3-triazol-5-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 255 2-methyl-5-((1-methyl-1H-1,2,4-triazol-3-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 256 2-methyl-5-((1-methyl-6-oxo-1,6-dihydropyridin-2-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 257 2-methyl-5-((2-methylpyrimidin-4-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 258 2-methyl-5-((3-methyloxetan-3-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 259 2-methyl-5-((4-methyloxazol-2-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 260 2-methyl-5-((4-methylthiazol-2-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 261 2-methyl-5-((5-methylpyrazin-2-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 262 2-methyl-5-((5-methylthiazol-2-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 263 2-methyl-5-((6-methylpyridazin-3-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 264 2-methyl-5-(pyrimidin-2-ylmethoxy)benzofuran-3-carboxylic acid
- Cpd 265 2-cyclopropyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 266 5-((6-methoxypyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 267 5-(imidazo[1,2-a]pyridin-6-ylmethoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 268 2-methyl-5-(1-(4-methylthiazol-5-yl)ethoxy)benzofuran-3-carboxylic acid
- Cpd 269 5-((2-methoxypyridin-4-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 270 (2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carbonyl)serine
- Cpd 271 (R)—N-(1-amino-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 272 (R)—N-(1-amino-3-(methylsulfonyl)-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 273 (R)—N-(1-amino-3-methyl-1-oxobutan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 274 (R)—N-(1-hydroxy-3-(methylsulfonyl)propan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 275 (R)—N-(1-hydroxy-3-(methylsulfonyl)propan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 276 (R)—N-(3-hydroxy-1-(methylamino)-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide

607

- Cpd 277 (S)—N-(1-(3,3-difluoropyrrolidin-1-yl)-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 278 (S)—N-(1-(dimethylamino)-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 5
- Cpd 279 (S)—N-(1-amino-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 280 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide 10
- Cpd 281 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((2-(trifluoromethyl)thiazol-5-yl)methoxy)benzofuran-3-carboxamide 15
- Cpd 282 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((2-methyl-4-(trifluoromethyl)thiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 283 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((2-methyl-4-(trifluoromethyl)thiazol-5-yl)methoxy)benzofuran-3-carboxamide 20
- Cpd 284 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((2-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide 25
- Cpd 285 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-4-yl)methoxy)benzofuran-3-carboxamide
- Cpd 286 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((3-(trifluoromethyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxamide 30
- Cpd 287 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((4-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 288 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 35
- Cpd 289 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((5-methylisoxazol-3-yl)methoxy)benzofuran-3-carboxamide 40
- Cpd 290 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((5-methylthiazol-4-yl)methoxy)benzofuran-3-carboxamide
- Cpd 291 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((6-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide 45
- Cpd 292 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-(oxetan-3-ylmethoxy)benzofuran-3-carboxamide
- Cpd 293 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-(pyridazin-3-ylmethoxy)benzofuran-3-carboxamide 50
- Cpd 294 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-(pyrimidin-4-ylmethoxy)benzofuran-3-carboxamide 55
- Cpd 295 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((1-(2,2-difluoroethyl)-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 296 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((2-(difluoromethoxy)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide 60
- Cpd 297 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((2-(dimethylamino)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 298 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((2,4-dimethylthiazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide 65

608

- Cpd 299 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((2,5-dimethylthiazol-4-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 300 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((2-aminopyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 301 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((3-(difluoromethyl)pyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 302 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((4-fluoro-1-methyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 303 (S)—N-(1-amino-3-methoxy-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 304 (S)—N-(1-amino-3-methyl-1-oxobutan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 305 (S)—N-(3-hydroxy-1-(methylamino)-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 306 (S)—N-(3-hydroxy-1-(methylamino)-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 307 (S)—N-(3-hydroxy-1-oxo-1-(pyrrolidin-1-yl)propan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 308 2-(difluoromethyl)-N-(1-(hydroxymethyl)cyclopropyl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 309 2-(difluoromethyl)-N-(3-(hydroxymethyl)oxetan-3-yl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 310 2-(difluoromethyl)-N-(3,3-difluoropiperidin-4-yl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 311 2-(difluoromethyl)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-hydroxypyridin-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 312 2-(difluoromethyl)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-methoxypyridin-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 313 2-cyclopropyl-N-(3,3-difluoropiperidin-4-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 314 2-cyclopropyl-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 315 2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(1-(oxazol-2-yl)ethyl)benzofuran-3-carboxamide
- Cpd 316 2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(1,1,1-trifluoro-3-hydroxypropan-2-yl)benzofuran-3-carboxamide
- Cpd 317 2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(5-oxo-1,4-oxazepan-6-yl)benzofuran-3-carboxamide
- Cpd 318 2-methyl-N-(1-(methylcarbamoyl)cyclobutyl)-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 319 2-methyl-N-(1-(methylcarbamoyl)cyclobutyl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 320 2-methyl-N-(1-(methylcarbamoyl)cyclobutyl)-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 321 2-methyl-N-(2-(methylamino)-2-oxoethyl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide

609

- Cpd 322 2-methyl-N-(2-(methylsulfonylamino)ethyl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 323 2-methyl-N-(2-(methylsulfonyl)ethyl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 5
- Cpd 324 2-methyl-N-(3-(methylcarbamoyl)oxetan-3-yl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 325 3-(((3-((4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)carbamoyl)-2-methylbenzofuran-5-yl)oxy)methyl)picolinamide 10
- Cpd 326 4-fluoro-N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 15
- Cpd 327 4-fluoro-N-(1-(hydroxymethyl)cyclopropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 328 4-fluoro-N-(3-(hydroxymethyl)oxetan-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 20
- Cpd 329 4-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide 25
- Cpd 330 4-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 331 5-((1-(2,2-difluoroethyl)-1H-imidazol-2-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide 30
- Cpd 332 5-((1-(2,2-difluoroethyl)-1H-pyrazol-5-yl)methoxy)-N-(3-(hydroxymethyl)-2-oxopiperidin-3-yl)-2-methylbenzofuran-3-carboxamide
- Cpd 333 5-((1-(2,2-difluoroethyl)-1H-pyrazol-5-yl)methoxy)-N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methylbenzofuran-3-carboxamide 35
- Cpd 334 5-((1-(2,2-difluoroethyl)-1H-pyrazol-5-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide 40
- Cpd 335 5-((2-(azetidin-1-yl)pyridin-3-yl)methoxy)-N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methylbenzofuran-3-carboxamide
- Cpd 336 5-((2-(azetidin-1-yl)pyridin-3-yl)methoxy)-N-(3,3-difluoropiperidin-4-yl)-2-methylbenzofuran-3-carboxamide 45
- Cpd 337 5-((2-(azetidin-1-yl)pyridin-3-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
- Cpd 338 5-((2-(difluoromethoxy)pyridin-3-yl)methoxy)-N-(3,3-difluoropiperidin-4-yl)-2-methylbenzofuran-3-carboxamide 50
- Cpd 339 5-((2-(difluoromethoxy)pyridin-3-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide 55
- Cpd 340 5-((2-(difluoromethyl)pyridin-3-yl)methoxy)-N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methylbenzofuran-3-carboxamide
- Cpd 341 5-((2-(difluoromethyl)pyridin-3-yl)methoxy)-N-(3-(hydroxymethyl)oxetan-3-yl)-2-methylbenzofuran-3-carboxamide 60
- Cpd 342 5-((2-(difluoromethyl)pyridin-3-yl)methoxy)-N-(3,3-difluoropiperidin-4-yl)-2-methylbenzofuran-3-carboxamide
- Cpd 343 5-((2-(difluoromethyl)pyridin-3-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide 65

610

- Cpd 344 5-((2-(dimethylamino)pyridin-3-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
- Cpd 345 5-((2,4-dimethylthiazol-5-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
- Cpd 346 5-((2-aminopyridin-3-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
- Cpd 347 5-((2-cyanopyridin-3-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
- Cpd 348 5-((2-ethoxypyridin-3-yl)methoxy)-N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methylbenzofuran-3-carboxamide
- Cpd 349 5-((2-hydroxypyridin-3-yl)methoxy)-N-(cis-4-hydroxytetrahydrofuran-3-yl)-2-methylbenzofuran-3-carboxamide
- Cpd 350 5-((2-hydroxypyridin-3-yl)methoxy)-N-(trans-4-hydroxytetrahydrofuran-3-yl)-2-methylbenzofuran-3-carboxamide
- Cpd 351 5-((3-(difluoromethyl)pyridin-2-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
- Cpd 352 5-((3-fluoropyridin-2-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
- Cpd 355 5-((4-fluoro-1-methyl-1H-pyrazol-5-yl)methoxy)-N-(3-(hydroxymethyl)-2-oxopiperidin-3-yl)-2-methylbenzofuran-3-carboxamide
- Cpd 356 5-((4-fluoro-1-methyl-1H-pyrazol-5-yl)methoxy)-N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methylbenzofuran-3-carboxamide
- Cpd 357 5-((4-fluoro-1-methyl-1H-pyrazol-5-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
- Cpd 358 5-((5-(2-hydroxyethoxy)pyridin-2-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
- Cpd 359 5-((5-fluoropyridin-2-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide
- Cpd 360 6-fluoro-N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 361 6-fluoro-N-(1-(hydroxymethyl)cyclopropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 362 6-fluoro-N-(3-(hydroxymethyl)oxetan-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 363 6-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 364 6-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 365 7-fluoro-N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 366 7-fluoro-N-(1-(hydroxymethyl)cyclopropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 367 7-fluoro-N-(3-(hydroxymethyl)oxetan-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide

611

- Cpd 368 7-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 369 7-fluoro-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide 5
- Cpd 370 N-((1-cyanocyclopropyl)methyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 371 N-((1-cyanocyclopropyl)methyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 10
- Cpd 372 N-((1-fluorocyclopropyl)methyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 15
- Cpd 373 N-((1-fluorocyclopropyl)methyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 374 N-((1H-pyrazol-3-yl)methyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 20
- Cpd 375 N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-2-methyl-5-((2-methyloxazol-5-yl)methoxy)benzofuran-3-carboxamide 25
- Cpd 376 N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-2-methyl-5-((2-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 377 N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 30
- Cpd 378 N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 379 N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 35
- Cpd 380 N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-2-methyl-5-((6-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide 40
- Cpd 381 N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-5-((1-(2,2-difluoroethyl)-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 382 N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-5-((2,4-dimethylthiazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide 45
- Cpd 383 N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-5-((4-fluoro-1-methyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 384 N-((3-fluorooxetan-3-yl)methyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 50
- Cpd 385 N-((3-fluorooxetan-3-yl)methyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 55
- Cpd 386 N-(cis-4-hydroxytetrahydrofuran-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 387 N-(cis-4-hydroxytetrahydrofuran-3-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide 60
- Cpd 388 N-(trans-4-hydroxytetrahydrofuran-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 389 N-(trans-4-hydroxytetrahydrofuran-3-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide 65

612

- Cpd 390 N-((3S,4R)-4-hydroxy-2-oxopyrrolidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 391 N-((3S,4R)-4-hydroxy-2-oxopyrrolidin-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 392 N-((3S,4R)-4-hydroxy-2-oxopyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 393 N-((4-fluorotetrahydro-2H-pyran-4-yl)methyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 394 N-((4-fluorotetrahydro-2H-pyran-4-yl)methyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 395 N-((S)-1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-(1-(2-methylthiazol-5-yl)ethoxy)benzofuran-3-carboxamide
- Cpd 396 N-(1-((2-hydroxyethyl)(methyl)amino)-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 397 N-(1-((2-hydroxyethyl)amino)-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 398 N-(1-((2-hydroxyethyl)amino)-2-methyl-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 399 N-(1-((difluoromethoxy)methyl)cyclopropyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 400 N-(1-((difluoromethoxy)methyl)cyclopropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 401 N-(1-(1H-benzo[d]imidazol-2-yl)-2-hydroxyethyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 402 N-(1-(2-(2-hydroxyethoxy)ethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 403 N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide
- Cpd 404 N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 405 N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((1-methyl-2-oxo-1,2-dihydropyridin-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 406—N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 407—N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-(thiazol-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 408 N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 409 N-(1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 410 N-(1-(2-hydroxyethyl)-3-methyl-2-oxopyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 411 N-(1-(dimethylamino)-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide

613

- Cpd 412 N-(1-(dimethylamino)-3-hydroxypropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 413 N-(1-(dimethylamino)-3-hydroxypropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 5
- Cpd 414 N-(1-(hydroxymethyl)cyclobutyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 415 N-(1-(hydroxymethyl)cyclobutyl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide 10
- Cpd 416 N-(1-(hydroxymethyl)cyclobutyl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 417 N-(1-(hydroxymethyl)cyclobutyl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide 15
- Cpd 418 N-(1-(hydroxymethyl)cyclopropyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 20
- Cpd 419 N-(1-(hydroxymethyl)cyclopropyl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 420 N-(1-(hydroxymethyl)cyclopropyl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide 25
- Cpd 421 N-(1-(hydroxymethyl)cyclopropyl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 422 N-(1,3-dihydroxy-2-methylpropan-2-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide 30
- Cpd 423 N-(1,3-dihydroxy-2-methylpropan-2-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 424 N-(1,3-dihydroxy-2-methylpropan-2-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide 35
- Cpd 425 N-(1,3-dihydroxy-2-methylpropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 40
- Cpd 426 N-(1,3-dihydroxy-2-methylpropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 427 N-(1,3-dihydroxy-2-methylpropan-2-yl)-2-methyl-5-(pyridazin-3-ylmethoxy)benzofuran-3-carboxamide 45
- Cpd 428 N-(1,3-dihydroxy-2-methylpropan-2-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 429 N-(1,3-dihydroxy-2-methylpropan-2-yl)-2-methyl-5-(pyrimidin-4-ylmethoxy)benzofuran-3-carboxamide 50
- Cpd 430 N-(1,3-dihydroxy-2-methylpropan-2-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide 55
- Cpd 431 N-(1,3-dihydroxy-2-methylpropan-2-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 432 N-(1-amino-1-oxo-3-(1H-1,2,4-triazol-1-yl)propan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 60
- Cpd 433 N-(1-amino-1-oxopropan-2-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 434 N-(1-amino-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 65

614

- Cpd 435 N-(1-amino-1-oxopropan-2-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 436 N-(1-amino-1-oxopropan-2-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 437 N-(1-amino-1-oxopropan-2-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 438 N-(1-amino-2-methyl-1-oxopropan-2-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 439 N-(1-amino-2-methyl-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 440 N-(1-amino-2-methyl-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 441 N-(1-amino-2-methyl-1-oxopropan-2-yl)-2-methyl-5-(pyridazin-3-ylmethoxy)benzofuran-3-carboxamide
- Cpd 442 N-(1-amino-2-methyl-1-oxopropan-2-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 443 N-(1-amino-2-methyl-1-oxopropan-2-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 444 N-(1-amino-2-methyl-1-oxopropan-2-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 445 N-(1-amino-3-(cyclopropyl(methyl)amino)-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 446 N-(1-amino-3-(cyclopropyl(methyl)amino)-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 447 N-(1-amino-3-(difluoromethoxy)-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 448 N-(1-amino-3-(difluoromethoxy)-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 449 N-(1-amino-3-(dimethylamino)-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 450 N-(1-amino-3-(dimethylamino)-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 451 N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide
- Cpd 452 N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 453 N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 454 N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 455 N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 456 N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((1-(2,2-difluoroethyl)-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide

615

- Cpd 457 N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((2-(difluoromethyl)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 458 N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((2,4-dimethylthiazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide 5
- Cpd 459 N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 460 N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((4-fluoro-1-methyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide 10
- Cpd 461 N-(1-amino-3-hydroxy-2-methyl-1-oxopropan-2-yl)-2-methyl-5-((2-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide 15
- Cpd 462 N-(1-amino-3-hydroxy-2-methyl-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 463 N-(1-amino-3-hydroxy-2-methyl-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 20
- Cpd 464 N-(1-amino-3-hydroxy-2-methyl-1-oxopropan-2-yl)-2-methyl-5-((6-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 465 N-(1-amino-3-hydroxy-2-methyl-1-oxopropan-2-yl)-5-((1-(2,2-difluoroethyl)-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide 25
- Cpd 466 N-(1-amino-3-hydroxy-2-methyl-1-oxopropan-2-yl)-5-((4-fluoro-1-methyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide 30
- Cpd 467 N-(1-amino-3-methoxy-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 468 N-(1-amino-3-methoxy-2-methyl-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 35
- Cpd 469 N-(1-amino-4-hydroxy-1-oxobutan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 470 N-(1-carbamoylcyclobutyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 40
- Cpd 471 N-(1-carbamoylcyclobutyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 45
- Cpd 472 N-(1-carbamoylcyclobutyl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 473 N-(1-carbamoylcyclobutyl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide 50
- Cpd 474 N-(1-carbamoylcyclobutyl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 475 N-(1-carbamoylcyclopropyl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide 55
- Cpd 476 N-(1-carbamoylcyclopropyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 477 N-(1-carbamoylcyclopropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 60
- Cpd 478 N-(1-carbamoylcyclopropyl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 479 N-(1-carbamoylcyclopropyl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide 65

616

- Cpd 480 N-(1-carbamoylcyclopropyl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 481 N-(1-hydroxy-2-methylpropan-2-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide
- Cpd 482 N-(1-hydroxy-2-methylpropan-2-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 483 N-(1-hydroxy-2-methylpropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 484 N-(1-hydroxy-2-methylpropan-2-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 485 N-(1-hydroxy-2-methylpropan-2-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 486 N-(1-hydroxy-2-methylpropan-2-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 487 N-(1-hydroxy-3-morpholinopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 488 N-(1-hydroxy-3-morpholinopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 489 N-(1-hydroxypropan-2-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 490 N-(1-hydroxypropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 491 N-(1-hydroxypropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 492 (S)—N-(1-hydroxypropan-2-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 493 (R)—N-(1-hydroxypropan-2-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 494 (S)—N-(1-hydroxypropan-2-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 495 (R)—N-(1-hydroxypropan-2-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 496 (S)—N-(1-hydroxypropan-2-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 497 (R)—N-(1-hydroxypropan-2-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 498 N-(2,2-difluoroethyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 499 N-(2,2-difluoropropyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 500 N-(2,2-difluoropropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 501 N-(2-acetamidoethyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 502 N-(2-amino-1-cyano-2-oxoethyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 503 N-(2-amino-2-oxoethyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 504 N-(2-fluoro-2-methylpropyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide

617

- Cpd 505 N-(2-fluoro-2-methylpropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 506 N-(2-hydroxy-2-methylpropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 5
- Cpd 507 N-(3-(aminomethyl)oxetan-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 508 N-(3-(dimethylamino)oxetane-3-carbonyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 10
- Cpd 509 N-(3-(dimethylcarbamoyl)oxetan-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 15
- Cpd 510 N-(3-(hydroxymethyl)-2-oxopiperidin-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 511 N-(3-(hydroxymethyl)-2-oxopiperidin-3-yl)-2-methyl-5-((2-methylloxazol-5-yl)methoxy)benzofuran-3-carboxamide 20
- Cpd 512 N-(3-(hydroxymethyl)-2-oxopiperidin-3-yl)-2-methyl-5-((2-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 513 N-(3-(hydroxymethyl)-2-oxopiperidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 25
- Cpd 514 N-(3-(hydroxymethyl)-2-oxopiperidin-3-yl)-2-methyl-5-((6-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide 30
- Cpd 515 N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((2-methylloxazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 516 N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((2-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide 35
- Cpd 517 N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 518 N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 40
- Cpd 519 N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((6-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide 45
- Cpd 520 N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 521 N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide 50
- Cpd 522 N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 523 N-(3-(hydroxymethyl)oxetan-3-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide 55
- Cpd 524 N-(3-(hydroxymethyl)oxetan-3-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide 60
- Cpd 525 N-(3-(hydroxymethyl)oxetan-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 526 N-(3-(hydroxymethyl)oxetan-3-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide 65
- Cpd 527 N-(3-(hydroxymethyl)oxetan-3-yl)-2-methyl-5-(thiazol-2-ylmethoxy)benzofuran-3-carboxamide

618

- Cpd 528 N-(3-(hydroxymethyl)oxetan-3-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 529 N-(3-(hydroxymethyl)oxetan-3-yl)-N,2-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 530 N-(3-(hydroxymethyl)pyrrolidin-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 531 N-(3-(hydroxymethyl)tetrahydro-2H-pyran-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 532 N-(3-(hydroxymethyl)tetrahydro-2H-pyran-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 533 N-(3-(hydroxymethyl)tetrahydrofuran-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 534 N-(3-(hydroxymethyl)tetrahydrofuran-3-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 535 N-(3-(hydroxymethyl)tetrahydrofuran-3-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 536 N-(3-(hydroxymethyl)tetrahydrofuran-3-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 537 N-(3,3-difluoro-1-(hydroxymethyl)cyclobutyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 538 N-(3,3-difluoropiperidin-4-yl)-2,4-dimethyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 539 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-(((2-(trifluoromethyl)pyridin-3-yl)oxy)methyl)benzofuran-3-carboxamide
- Cpd 540 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((1-methyl-1H-1,2,3-triazol-4-yl)methoxy)benzofuran-3-carboxamide
- Cpd 541 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((1-methyl-1H-1,2,3-triazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 542 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((1-methyl-1H-1,2,4-triazol-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 543 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 544 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((1-methyl-2-oxo-1,2-dihydropyridin-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 545 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((1-methyl-6-oxo-1,6-dihydropyridin-2-yl)methoxy)benzofuran-3-carboxamide
- Cpd 546 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 547 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((2-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 548 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((2-methylpyrimidin-4-yl)methoxy)benzofuran-3-carboxamide
- Cpd 549 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide

619

- Cpd 550 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((3-methyloxetan-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 551 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((4-methyloxazol-2-yl)methoxy)benzofuran-3-carboxamide 5
- Cpd 552 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((4-methylthiazol-2-yl)methoxy)benzofuran-3-carboxamide
- Cpd 553 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((5-methylisoxazol-3-yl)methoxy)benzofuran-3-carboxamide 10
- Cpd 554 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((5-methylpyrazin-2-yl)methoxy)benzofuran-3-carboxamide 15
- Cpd 555 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((5-methylthiazol-2-yl)methoxy)benzofuran-3-carboxamide
- Cpd 556 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-((6-methylpyridazin-3-yl)methoxy)benzofuran-3-carboxamide 20
- Cpd 557 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-(pyridazin-3-ylmethoxy)benzofuran-3-carboxamide
- Cpd 558 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-(pyridin-3-ylmethoxy)benzofuran-3-carboxamide 25
- Cpd 559 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-(pyrimidin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 560 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-(thiazol-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 561 N-(3,3-difluoropiperidin-4-yl)-2-methyl-5-(thiazol-5-ylmethoxy)benzofuran-3-carboxamide 30
- Cpd 562 N-(3,3-difluoropiperidin-4-yl)-4-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 563 N-(3,3-difluoropiperidin-4-yl)-5-((2,4-dimethylthiazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide 35
- Cpd 564 N-(3,3-difluoropiperidin-4-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide 40
- Cpd 565 N-(3,3-difluoropiperidin-4-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 566 N-(3,3-difluoropiperidin-4-yl)-5-((3-fluoropyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide 45
- Cpd 567 N-(3,3-difluoropiperidin-4-yl)-5-((5-(2-hydroxyethoxy)pyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 568 N-(3,3-difluoropiperidin-4-yl)-5-((5-fluoropyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide 50
- Cpd 569 N-(3,3-difluoropiperidin-4-yl)-5-((6-hydroxypyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide 55
- Cpd 570 N-(3,3-difluoropiperidin-4-yl)-5-((6-methoxypyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 571 N-(3,3-difluoropiperidin-4-yl)-5-(imidazo[1,2-a]pyridin-6-ylmethoxy)-2-methylbenzofuran-3-carboxamide 60
- Cpd 572 N-(3,3-difluoropiperidin-4-yl)-6-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 573 N-(3,3-difluoropiperidin-4-yl)-7-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 65

620

- Cpd 574 N-(3,3-difluoropropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 575 N-(3-amino-2-(dimethylamino)-3-oxopropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 576 N-(3-amino-2,2-dimethyl-3-oxopropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 577 N-(3-carbamoyloxetan-3-yl)-2-(difluoromethyl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 578 N-(3-carbamoyloxetan-3-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide
- Cpd 579 N-(3-carbamoyloxetan-3-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 580 N-(3-carbamoyloxetan-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 581 N-(3-carbamoyloxetan-3-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 582 N-(3-carbamoyloxetan-3-yl)-2-methyl-5-(thiazol-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 583 N-(3-carbamoyloxetan-3-yl)-4-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 584 N-(3-carbamoyloxetan-3-yl)-5-((2-(difluoromethyl)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 585 N-(3-carbamoyloxetan-3-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 586 N-(3-carbamoyloxetan-3-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 587 N-(3-carbamoyloxetan-3-yl)-6-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 588 N-(3-carbamoyloxetan-3-yl)-7-fluoro-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 589 N-(3-carbamoyloxetan-3-yl)-N,2-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 590 N-(3-carbamoyltetrahydro-2H-pyran-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 591 N-(3-carbamoyltetrahydrofuran-3-yl)-2-methyl-5-((2-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 592 N-(3-carbamoyltetrahydrofuran-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 593 N-(3-carbamoyltetrahydrofuran-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 594 N-(3-carbamoyltetrahydrofuran-3-yl)-2-methyl-5-((6-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 595 N-(3-carbamoyltetrahydrofuran-3-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 596 N-(3-carbamoyltetrahydrofuran-3-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide

621

- Cpd 597 N-(3-carbamoyltetrahydrofuran-3-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 598 N-(3-carbamoyltetrahydrofuran-3-yl)-5-((4-fluoro-1-methyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide 5
- Cpd 599 N-(3-cyanooxetan-3-yl)-N,2-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 600 N-(3-hydroxy-1-(methylamino)-1-oxopropan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 10
- Cpd 601 N-(3-hydroxy-1-(methylamino)-1-oxopropan-2-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide 15
- Cpd 602 N-(3-hydroxy-2,2-dimethylpropyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 603 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2,4-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 20
- Cpd 604 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((1-methyl-1H-1,2,3-triazol-4-yl)methoxy)benzofuran-3-carboxamide 25
- Cpd 605 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((1-methyl-1H-1,2,3-triazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 606 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((1-methyl-1H-1,2,4-triazol-3-yl)methoxy)benzofuran-3-carboxamide 30
- Cpd 607 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((1-methyl-1H-imidazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 608 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide 35
- Cpd 609 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((1-methyl-2-oxo-1,2-dihydropyridin-3-yl)methoxy)benzofuran-3-carboxamide 40
- Cpd 610 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((1-methyl-6-oxo-1,6-dihydropyridin-2-yl)methoxy)benzofuran-3-carboxamide
- Cpd 611 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((2-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide 45
- Cpd 612 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((2-methylpyrimidin-4-yl)methoxy)benzofuran-3-carboxamide
- Cpd 613 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 50
- Cpd 614 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((2-oxo-1,2-dihydropyridin-3-yl)methoxy)benzofuran-3-carboxamide 55
- Cpd 615 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((3-(trifluoromethyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxamide
- Cpd 616 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methyloxazol-2-yl)methoxy)benzofuran-3-carboxamide 60
- Cpd 617 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-2-yl)methoxy)benzofuran-3-carboxamide
- Cpd 618 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((5-methylisoxazol-3-yl)methoxy)benzofuran-3-carboxamide 65

622

- Cpd 619 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((5-methylpyrazin-2-yl)methoxy)benzofuran-3-carboxamide
- Cpd 620 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((5-methylthiazol-2-yl)methoxy)benzofuran-3-carboxamide
- Cpd 621 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((6-methylpyridazin-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 622 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-((6-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 623 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-(1-(4-methylthiazol-5-yl)ethoxy)benzofuran-3-carboxamide
- Cpd 624 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-(pyridazin-3-ylmethoxy)benzofuran-3-carboxamide
- Cpd 625 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-(pyridin-3-ylmethoxy)benzofuran-3-carboxamide
- Cpd 626 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-(pyrimidin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 627 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methyl-5-(thiazol-5-ylmethoxy)benzofuran-3-carboxamide
- Cpd 628 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-hydroxypyridin-4-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 629 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((2-methoxypyridin-4-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 630 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((6-hydroxypyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 631 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((6-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 632 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((6-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 633 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-(imidazo[1,2-a]pyridin-6-ylmethoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 634 N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-N,2-dimethyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 635 N-(4,4-difluoropiperidin-3-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide
- Cpd 636 N-(4,4-difluoropiperidin-3-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 637 N-(4,4-difluoropiperidin-3-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 638 N-(4,4-difluoropyrrolidin-3-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide
- Cpd 639 N-(4,4-difluoropyrrolidin-3-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 640 N-(4,4-difluoropyrrolidin-3-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide

623

- Cpd 641 N-(4,4-difluorotetrahydrofuran-3-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide
- Cpd 642 N-(4,4-difluorotetrahydrofuran-3-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide 5
- Cpd 643 N-(4,4-difluorotetrahydrofuran-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 646 N-(4,4-difluorotetrahydrofuran-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 10
- Cpd 647 N-(4,4-difluorotetrahydrofuran-3-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 648 N-(4,4-difluorotetrahydrofuran-3-yl)-5-((2-hydroxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide 15
- Cpd 649 N-(4-amino-2-methyl-4-oxobutan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 20
- Cpd 650 N-(4-carbamoyltetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 651 N-(4-hydroxy-2-methylbutan-2-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide 25
- Cpd 652 N-(4-hydroxy-2-methylbutan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 653 N-(cis-4-hydroxytetrahydrofuran-3-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide 30
- Cpd 654 N-(trans-4-hydroxytetrahydrofuran-3-yl)-2-methyl-5-((1-methyl-1H-imidazol-2-yl)methoxy)benzofuran-3-carboxamide 35
- Cpd 655 N-(cis-4-hydroxytetrahydrofuran-3-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 656 N-(trans-4-hydroxytetrahydrofuran-3-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide 40
- Cpd 657 N3-(3,3-difluoropiperidin-4-yl)-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-2,3-dicarboxamide
- Cpd 658 N3-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-2,3-dicarboxamide 45
- Cpd 659 N-cyclobutyl-N-(2-hydroxyethyl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide 50
- Cpd 660 N-cyclobutyl-N-(2-hydroxyethyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 661 2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(2-oxazepan-3-yl)benzofuran-3-carboxamide 55
- Cpd 662 N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-(pyridin-3-ylmethoxy)benzofuran-3-carboxamide
- Cpd 663 N-(4,4-difluoropyrrolidin-3-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide 60
- Cpd 664 N-(4,4-difluorotetrahydrofuran-3-yl)-2-methyl-5-((2-(trifluoromethyl)pyridin-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 665 N-(4,4-difluorotetrahydrofuran-3-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide 65

624

- Cpd 666 2-methyl-5-((4-methylthiazol-5-yl)methoxy)-N-(2,2,2-trifluoroethyl)benzofuran-3-carboxamide
- Cpd 667 5-((1,4-Dimethyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 668 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-5-((1,4-dimethyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 669 5-((4-Chloro-1-methyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 670 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-5-((4-chloro-1-methyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 671 5-((4-chloro-1-isopropyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 672 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((4-chloro-1-isopropyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 673 2-Methyl-5-((3-methylisoxazol-5-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 674 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((3-methylisoxazol-5-yl)methyl)benzofuran-3-carboxamide
- Cpd 675 5-((1-isopropyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 676 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((1-isopropyl-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 677 5-((2,4-Dimethyloxazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 678 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-5-((2,4-dimethyloxazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 679 2-Methyl-5-((4-methyloxazol-5-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 680 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((4-methyloxazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 681 5-((2-Isopropylthiazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 682 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-5-((2-isopropylthiazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 683 N-(1-(Difluoromethyl)cyclobutyl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 684 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((5-methylthiazol-2-yl)methoxy)benzofuran-3-carboxamide
- Cpd 685 N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-2-methyl-5-((3-(trifluoromethyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxamide
- Cpd 686 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((1-methyl-1H-pyrazol-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 687 2-Methyl-5-((1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 688 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((1-(2,2,2-trifluoroethyl)-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 689 5-((1-(Cyclopropylmethyl)-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 690 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((1-(cyclopropylmethyl)-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 691 2-Methyl-5-((1-methyl-3-(trifluoromethyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxylic acid

625

- Cpd 692 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((1-methyl-3-(trifluoromethyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxamide
- Cpd 693 2-Methyl-5-((4-(trifluoromethyl)thiazol-5-yl)methoxy)benzofuran-3-carboxylic acid 5
- Cpd 694 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((4-(trifluoromethyl)thiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 695 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-(thiazol-2-ylmethoxy)benzofuran-3-carboxamide 10
- Cpd 696 N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-5-(5-fluoropyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide 15
- Cpd 697 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-(thiazol-5-ylmethoxy)benzofuran-3-carboxamide
- Cpd 698 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((1-methyl-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxamide 20
- Cpd 699 2-Methyl-5-((5-methylpyridin-2-yl)methoxy)benzofuran-3-carboxylic acid
- Cpd 700 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-((5-methylpyridin-2-yl)methoxy)benzofuran-3-carboxamide 25
- Cpd 701 5-((2-cyclopropylpyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 702 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-5-(2-cyclopropylpyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide 30
- Cpd 703 5-((3-Cyclopropylpyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid
- Cpd 704 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-5-((3-cyclopropylpyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide; 35
- Cpd 705 5-((3-cyclopropylpyridin-2-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide 40
- Cpd 706 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((2-(azetidin-1-yl)pyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 707 2-Methyl-5-(2-methylpyrimidin-5-yl)methoxy)benzofuran-3-carboxylic acid 45
- Cpd 708 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-(2-methylpyrimidin-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 709 5-((2-Cyclopropoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxylic acid 50
- Cpd 710 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-(2-cyclopropoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 711 5-((2-cyclopropoxypyridin-3-yl)methoxy)-N-(4-(hydroxymethyl)tetrahydro-2H-pyran-4-yl)-2-methylbenzofuran-3-carboxamide 55
- Cpd 712 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-(pyrimidin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 713 2-Methyl-5-(pyrazin-2-ylmethoxy)benzofuran-3-carboxylic acid 60
- Cpd 714 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-(pyrazin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 715 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide 65

626

- Cpd 716 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-methyl-5-(pyridin-4-ylmethoxy)benzofuran-3-carboxamide
- Cpd 717 5-((3-(Difluoromethyl)pyridin-2-yl)methoxy)-N-(3,3-difluoropiperidin-4-yl)-2-methylbenzofuran-3-carboxamide
- Cpd 718 N-((2S,3R)-1-amino-3-hydroxy-1-oxobutan-2-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 719 N-(3-(hydroxymethyl)-2-oxopiperidin-3-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 720 N-(1-amino-3-hydroxy-2-methyl-1-oxopropan-2-yl)-5-((2-methoxypyridin-3-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 721 N-(1-(Difluoromethyl)cyclobutyl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 722 N-(1-Amino-3-hydroxy-2-methyl-1-oxopropan-2-yl)-2-methyl-5-((2-methylloxazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 723 N-(3,3-difluorotetrahydro-2H-pyran-4-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 724 N-(3,3-Difluorotetrahydro-2H-pyran-4-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 725 N-(3-carbamoyltetrahydrofuran-3-yl)-2-methyl-5-((2-methylloxazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 726 N-(1-Amino-3-(methylsulfonyl)-1-oxopropan-2-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 727 N-(1-Imino-1-oxidotetrahydro-1H-116-thiophen-3-yl)-2-methyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 728 N-(3-(hydroxymethyl)pyrrolidin-3-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 729 N-(1-Amino-1-oxo-3-(1H-1,2,4-triazol-1-yl)propan-2-yl)-2-methyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 730 N-(3-(Hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((3-(trifluoromethyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxamide
- Cpd 731 N-(3-(hydroxymethyl)-2-oxopiperidin-3-yl)-2-methyl-5-((3-(trifluoromethyl)-1H-pyrazol-4-yl)methoxy)benzofuran-3-carboxamide
- Cpd 732 N-(3-(Hydroxymethyl)-2-oxopiperidin-3-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 733 N-(3-(Hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-((1-methyl-1H-pyrazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 734 N-(3-(Hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methyl-5-(pyridazin-3-ylmethoxy)benzofuran-3-carboxamide
- Cpd 735 5-((5-Fluoropyridin-2-yl)methoxy)-N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methylbenzofuran-3-carboxamide
- Cpd 736 5-((2-(Difluoromethyl)pyridin-3-yl)methoxy)-N-(3-(hydroxymethyl)-2-oxopyrrolidin-3-yl)-2-methylbenzofuran-3-carboxamide
- Cpd 737 5-((2-(Difluoromethyl)pyridin-3-yl)methoxy)-N-(3-(hydroxymethyl)-2-oxopiperidin-3-yl)-2-methylbenzofuran-3-carboxamide

627

- Cpd 738 N-(1-Amino-3-hydroxy-2-methyl-1-oxopropan-2-yl)-2-methyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 739 N-(1-Amino-3-hydroxy-2-methyl-1-oxopropan-2-yl)-5-((5-fluoropyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide 5
- Cpd 740 N-(3-carbamoyltetrahydrofuran-3-yl)-5-((1-(2,2-difluoroethyl)-1H-pyrazol-5-yl)methoxy)-2-methylbenzofuran-3-carboxamide
- Cpd 741 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-5-((5-fluoropyridin-2-yl)methoxy)-2-methylbenzofuran-3-carboxamide 10
- Cpd 742 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-(difluoromethyl)-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide 15
- Cpd 743 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-(difluoromethyl)-5-((6-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 744 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-(difluoromethyl)-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide 20
- Cpd 745 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-(difluoromethyl)-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide

628

- Cpd 746 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-(difluoromethyl)-5-((2-methyloxazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 747 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-cyclopropyl-5-(pyridin-2-ylmethoxy)benzofuran-3-carboxamide
- Cpd 748 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-cyclopropyl-5-((4-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 749 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-cyclopropyl-5-((2-methylthiazol-5-yl)methoxy)benzofuran-3-carboxamide
- Cpd 750 (S)—N-(1-Amino-3-hydroxy-1-oxopropan-2-yl)-2-cyclopropyl-5-((6-methylpyridin-3-yl)methoxy)benzofuran-3-carboxamide
- Cpd 751 (S)—N-(1-amino-3-hydroxy-1-oxopropan-2-yl)-2-cyclopropyl-5-((2-methyloxazol-5-yl)methoxy)benzofuran-3-carboxamide
- and the physiologically acceptable salts thereof.
13. A pharmaceutical composition comprising a compound according to claim 1 and a pharmaceutically acceptable carrier.

* * * * *