

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent Application Publication

20250280828

Kind Code

A1

Publication Date

September 11, 2025

Inventor(s)

DUVARESCH; Dhiego et al.

FUNGICIDAL COMPOSITIONS

Abstract

A fungicidal composition comprising a mixture of components (A) and (B), wherein components (A) and (B) are as defined in claim 1, and use of the compositions in agriculture or horticulture for controlling or preventing infestation of plants by phytopathogenic microorganisms, preferably fungi.

Inventors: DUVARESCH; Dhiego (Sao Paulo, BR), QUARANTA; Laura (Basel, CH), HAAS; Ulrich Johannes (Stein, CH)

Applicant: SYNGENTA CROP PROTECTION AG (Basel, CH)

Family ID: 78820229

Assignee: SYNGENTA CROP PROTECTION AG (Basel, CH)

Appl. No.: 18/255490

Filed (or PCT Filed): December 01, 2021

PCT No.: PCT/EP2021/083778

Foreign Application Priority Data

EP	20211348.6	Dec. 02, 2020
EP	20213345.0	Dec. 11, 2020
EP	21196079.4	Sep. 10, 2021

Publication Classification

Int. Cl.: A01N43/40 (20060101); A01N25/04 (20060101); A01N25/08 (20060101); A01N43/82 (20060101); A01P3/00 (20060101)

U.S. Cl.:

CPC **A01N43/40** (20130101); **A01N25/04** (20130101); **A01N25/08** (20130101); **A01N43/82** (20130101); **A01P3/00** (20210801);

Background/Summary

CROSS-REFERENCE TO RELATED APPLICATIONS [0001] This application is a U.S. National Stage Application, filed under 35 U.S.C. § 371, of International Application No. PCT/EP2021/083778, filed Dec. 1, 2021, which claims priority to EP 21196079.4, filed Sep. 10, 2021, EP 20213345.0, filed Dec. 11, 2020, and EP 20211348.6, filed Dec. 2, 2020, the entire contents of which are incorporated by reference herein.

[0002] The present invention relates to novel fungicidal compositions, to their use in agriculture or horticulture for controlling diseases caused by phytopathogens, especially phytopathogenic fungi, and to methods of controlling diseases on useful plants.

[0003] Certain oxadiazole derivatives are known as insecticidal and acaricidal agents, e.g., from CN 1927860. WO 2013/064079, EP 0 276 432 and WO 2015/185485 describe the use of substituted oxadiazoles for combating phytopathogenic fungi.

[0004] Whilst many fungicidal compounds and compositions, belonging to various different chemical classes, have been/are being developed for use as fungicides in crops of useful plants, crop tolerance and activity against particular phytopathogenic fungi do not always satisfy the needs of agricultural practice in many respects. Therefore, there is a continuing need to find new compounds and compositions having superior biological properties for use in controlling or preventing infestation of plants by phytopathogenic fungi. For example, compounds possessing a greater biological activity, an advantageous spectrum of activity, an increased safety profile, improved physico-chemical properties, or increased biodegradability. Or else, compositions possessing a broader spectrum of activity, improved crop tolerance, synergistic interactions or potentiating properties, or compositions which display a more rapid onset of action or which have longer lasting residual activity or which enable a reduction in the number of applications and/or a reduction in the application rate of the compounds and compositions required for effective control of a phytopathogen, thereby enabling beneficial resistance-management practices, reduced environmental impact and reduced operator exposure.

[0005] The use of compositions comprising mixtures of different fungicidal compounds possessing different modes of action can address some of these needs (e.g., by combining fungicides with differing spectrums of activity).

[0006] According to the present invention, there is provided a fungicidal composition comprising a mixture of components (A) and (B) as active ingredients, wherein [0007] component (A) is a compound of formula (I):

##STR00001## [0008] wherein [0009] R^{sup.1} is hydrogen, —C(=O)CH_{sub.3} or —C(=O)CH_{sub.2}CH_{sub.3}, [0010] R^{sup.2} is methyl or isopropyl, and [0011] R^{sup.3} is hydrogen or fluorine, [0012] or a salt, enantiomer, tautomer or N-oxide thereof; and [0013] component (B) is a compound selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-

oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof.

[0014] In general, the weight ratio of component (A) to component (B) may be from 100:1 to 1:100, preferably from 50:1 to 1:50, more preferably from 20:1 to 1:40, even more preferably from 15:1 to 1:30, still more preferably from 12:1 to 1:25, or from 10:1 to 1:20, or from 10:1 to 1:10, or from 5:1 and 1:15, or from 5:1 to 1:5, or from 4:1 to 1:4, or from 3:1 to 1:10, or from 3:1 to 1:3, or from 2:1 to 1:5, or 1:1.

[0015] Further, according to the invention, there is provided a method of controlling or preventing phytopathogenic diseases, especially phytopathogenic fungi, on useful plants or on propagation material thereof, which comprises applying to the useful plants, the locus thereof or propagation material thereof a fungicidal composition according to the invention.

[0016] The benefits provided by certain fungicidal mixture compositions according to the invention may also include, inter alia, advantageous levels of biological activity for protecting plants against diseases that are caused by fungi or superior properties for use as agrochemical active ingredients (for example, greater biological activity, an advantageous spectrum of activity, an increased safety profile, improved physico-chemical properties, or increased biodegradability).

[0017] In one composition according to the invention, component (A) is a compound of formula (I-a):

##STR00002## [0018] wherein R^{sup.1} is hydrogen, —C(=O)CH_{sub.3} or —C(=O)CH_{sub.2}CH_{sub.3}, [0019] or a salt, enantiomer, tautomer or N-oxide thereof.

[0020] In another composition according to the invention, component (A) is a compound of formula (I-b):

##STR00003## [0021] wherein R^{sup.1} is hydrogen, —C(=O)CH_{sub.3} or —C(=O)CH_{sub.2}CH_{sub.3}, [0022] or a salt, enantiomer, tautomer or N-oxide thereof.

[0023] Preferably, component (A) is a compound selected from compounds A.01, A.02, A.03, A.04, A.05, A.06 and A.07 as defined in Table A below.

TABLE-US-00001 TABLE A Compound No. Compound structure IUPAC name A.01 [00004]

 [(1S,2S)-1-methyl-2-(o-tolyl)propyl] (2S)- 2-[(3-hydroxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate A.02 [00005]

 [(1S,2S)-1-methyl-2-(o-tolyl)propyl] (2S)- 2-[(3-acetoxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate A.03 [00006]

 [(1S,2S)-1-methyl-2-(o-tolyl)propyl] (2S)- 2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate A.04 [00007]

 [(1S,2S)-2-(4-fluoro-2-methyl-phenyl)- 1,3-dimethyl-butyl] (2S)-2-[(3-hydroxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate A.05 [00008]

 [(1S,2S)-2-(4-fluoro-2-methyl-phenyl)- 1,3-dimethyl-butyl] (2S)-2-[(3-acetoxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate A.06 [00009]

 [(1S,2S)-2-(4-fluoro-2-methyl-phenyl)- 1,3-dimethyl-butyl] (2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate A.07 [00010]

 [(1R,2R)-1-methyl-2-(o-tolyl)propyl] (2S)- 2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate

[0024] The presence of one or more possible asymmetric carbon atoms in component (B) means that the compounds may occur in optically isomeric forms, i.e., enantiomeric or diastereomeric forms. Also, atropisomers may occur as a result of restricted rotation about a single bond. The

present invention includes all those possible isomeric forms (e.g. geometric isomers) and mixtures thereof for component (B). The present invention includes all possible tautomeric forms for component (B), and also a racemic compound, i.e., a mixture of at least two enantiomers in a ratio of substantially 50:50.

[0025] In each case, the components (B) according to the invention are in free form, in oxidized form as a N-oxide or in salt form, e.g. an agronomically usable salt form.

[0026] N-oxides are oxidized forms of tertiary amines or oxidized forms of nitrogen containing heteroaromatic compounds. They are described for instance in the book "Heterocyclic N-oxides" by A. Albini and S. Pietra, CRC Press, Boca Raton 1991.

[0027] Component (B) is a compound selected from compounds B.01, B.02, B.03, B.04, B.05, B.06, B.07, B.08, B.09, B.10, B.11, B.12, B.13, B.14 and B.15 as defined in the Table B below.

TABLE-US-00002 TABLE B Compound number Compound structure IUPAC name B.01 [00011]

 N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-

yl]phenyl]methyl]cyclopropanecarboxamide B.02 [00012]  N,2-dimethoxy-N-

[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide B.03 [00013]

 N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-

yl]phenyl]methyl]propanamide B.04 [00014]  1-methoxy-3-methyl-1-[[4-[5-

(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea B.05 [00015]  1,3-

dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea B.06 [00016]

 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-

yl]phenyl]methyl]urea B.07 [00017]  N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-

3-yl]phenyl]methyl]propanamide B.08 [00018]  4,4-dimethyl-2-[[4-[5-

(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one B.09 [00019]

 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-

yl]phenyl]methyl]isoxazolidin-3-one B.10 [00020]  ethyl 1-[[4-[5-

(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate B.11 [00021]

 N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-

yl]phenyl]methyl]-1,2,4-triazol-3-amine B.12 [00022]  (3-methylisoxazol-5-yl)-

[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone B.13 [00023] 

(5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone B.14

[00024]  2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-

yl]phenyl]acetamide B.15 [00025]  ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-

oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate

[0028] The presence of an asymmetric carbon atom in any of the compounds B.01, B.02, B.03, B.04, B.05, B.06, B.07, B.08, B.09, B.10, B.11, B.12, B.13, B.14, or B.15 as defined in the Table B above means that these compounds may occur in chiral enantiomeric forms, i.e., (R)- and (S)-enantiomers.

[0029] Enantiomerically pure final compounds may be obtained from racemic starting materials as appropriate via standard physical separation techniques, such as reverse phase chiral chromatography, or through stereoselective synthetic techniques, e.g., by using chiral starting materials.

[0030] Preferred embodiments E.01 to E.105 as listed in Table C below relate to binary mixtures, wherein the weight ratio of component (A) to component (B) may be from 100:1 to 1:100, preferably from 50:1 to 1:50, more preferably from 20:1 to 1:40, even more preferably from 15:1 to 1:30, still more preferably from 12:1 to 1:25, or from 10:1 to 1:20, or from 10:1 to 1:10, or from 5:1 and 1:15, or from 5:1 to 1:5, or from 4:1 to 1:4, or from 3:1 to 1:10, or from 3:1 to 1:3, or from 2:1 to 1:5, or 1:1.

TABLE-US-00003 TABLE C Embodiment (A) (B) E.01 A.01 B.01 E.02 A.01 B.02 E.03 A.01 B.03

E.04 A.01 B.04 E.05 A.01 B.05 E.06 A.01 B.06 E.07 A.01 B.07 E.08 A.01 B.08 E.09 A.01 B.09

E.10 A.01 B.10 E.11 A.01 B.11 E.12 A.01 B.12 E.13 A.01 B.13 E.14 A.01 B.14 E.15 A.01 B.15

E.16 A.02 B.01 E.17 A.02 B.02 E.18 A.02 B.03 E.19 A.02 B.04 E.20 A.02 B.05 E.21 A.02 B.06 E.22 A.02 B.07 E.23 A.02 B.08 E.24 A.02 B.09 E.25 A.02 B.10 E.26 A.02 B.11 E.27 A.02 B.12 E.28 A.02 B.13 E.29 A.02 B.14 E.30 A.02 B.15 E.31 A.03 B.01 E.32 A.03 B.02 E.33 A.03 B.03 E.34 A.03 B.04 E.35 A.03 B.05 E.36 A.03 B.06 E.37 A.03 B.07 E.38 A.03 B.08 E.39 A.03 B.09 E.40 A.03 B.10 E.41 A.03 B.11 E.42 A.03 B.12 E.43 A.03 B.13 E.44 A.03 B.14 E.45 A.03 B.15 E.46 A.04 B.01 E.47 A.04 B.02 E.48 A.04 B.03 E.49 A.04 B.04 E.50 A.04 B.05 E.51 A.04 B.06 E.52 A.04 B.07 E.53 A.04 B.08 E.54 A.04 B.09 E.55 A.04 B.10 E.56 A.04 B.11 E.57 A.04 B.12 E.58 A.04 B.13 E.59 A.04 B.14 E.60 A.04 B.15 E.61 A.05 B.01 E.62 A.05 B.02 E.63 A.05 B.03 E.64 A.05 B.04 E.65 A.05 B.05 E.66 A.05 B.06 E.67 A.05 B.07 E.68 A.05 B.08 E.69 A.05 B.09 E.70 A.05 B.10 E.71 A.05 B.11 E.72 A.05 B.12 E.73 A.05 B.13 E.74 A.05 B.14 E.75 A.05 B.15 E.76 A.06 B.01 E.77 A.06 B.02 E.78 A.06 B.03 E.79 A.06 B.04 E.80 A.06 B.05 E.81 A.06 B.06 E.82 A.06 B.07 E.83 A.06 B.08 E.84 A.06 B.09 E.85 A.06 B.10 E.86 A.06 B.11 E.87 A.06 B.12 E.88 A.06 B.13 E.89 A.06 B.14 E.90 A.06 B.15 E.91 A.07 B.01 E.92 A.07 B.02 E.93 A.07 B.03 E.94 A.07 B.04 E.95 A.07 B.05 E.96 A.07 B.06 E.97 A.07 B.07 E.98 A.07 B.08 E.99 A.07 B.09 E.100 A.07 B.10 E.101 A.07 B.11 E.102 A.07 B.12 E.103 A.07 B.13 E.104 A.07 B.14 E.105 A.07 B.15

[0031] In a preferred composition according to the invention, component (A) is [(1 S,2S)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(3-hydroxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.01), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 15:1 to 1:30.

[0032] In a preferred composition according to the invention, component (A) is [(1S,2S)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(3-acetoxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.02), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-

oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 15:1 to 1:30.

[0033] In a preferred composition according to the invention, component (A) is [(1 S,2S)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate (A.03), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 15:1 to 1:30.

[0034] In a preferred composition according to the invention, component (A) is [(1 S,2S)-2-(4-fluoro-2-methyl-phenyl)-1,3-dimethyl-butyl](2S)-2-[(3-hydroxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.04), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 15:1 to 1:30.

[0035] In a preferred composition according to the invention, component (A) is [(1S,2S)-2-(4-fluoro-2-methyl-phenyl)-1,3-dimethyl-butyl](2S)-2-[(3-acetoxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.05), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-

(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 15:1 to 1:30.

[0036] In a preferred composition according to the invention, component (A) is [(1S,2S)-2-(4-fluoro-2-methyl-phenyl)-1,3-dimethyl-butyl](2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate (A.06), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 15:1 to 1:30.

[0037] In a preferred composition according to the invention, component (A) is [(1R,2R)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate (A.07), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-

oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 15:1 to 1:30.

[0038] In a preferred composition according to the invention, component (A) is [(1S,2S)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(3-hydroxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.01), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group

consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 10:1 to 1:10.

[0039] In a preferred composition according to the invention, component (A) is [(1 S,2S)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(3-acetoxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.02), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group

consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 10:1 to 1:10.

[0040] In a preferred composition according to the invention, component (A) is [(1S,2S)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate (A.03), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-

yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 10:1 to 1:10.

[0041] In a preferred composition according to the invention, component (A) is [(1 S,2S)-2-(4-fluoro-2-methyl-phenyl)-1,3-dimethyl-butyl](2S)-2-[(3-hydroxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.04), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 10:1 to 1:10.

[0042] In a preferred composition according to the invention, component (A) is [(1 S,2S)-2-(4-fluoro-2-methyl-phenyl)-1,3-dimethyl-butyl](2S)-2-[(3-acetoxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.05), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-

oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 10:1 to 1:10.

[0043] In a preferred composition according to the invention, component (A) is [(1S,2S)-2-(4-fluoro-2-methyl-phenyl)-1,3-dimethyl-butyl](2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate (A.06), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 10:1 to 1:10.

[0044] In a preferred composition according to the invention, component (A) is [(1R,2R)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate (A.07), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 10:1 to 1:10.

[0045] In a preferred composition according to the invention, component (A) is [(1S,2S)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(3-hydroxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.01), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-

yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 5:1 to 1:5.

[0046] In a preferred composition according to the invention, component (A) is [(1S,2S)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(3-acetoxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.02), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 5:1 to 1:5.

[0047] In a preferred composition according to the invention, component (A) is [(1S,2S)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate (A.03), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-

oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 5:1 to 1:5. [0048] In a preferred composition according to the invention, component (A) is [(1 S,2S)-2-(4-fluoro-2-methyl-phenyl)-1,3-dimethyl-butyl](2S)-2-[(3-hydroxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.04), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 5:1 to 1:5. [0049] In a preferred composition according to the invention, component (A) is [(1 S,2S)-2-(4-fluoro-2-methyl-phenyl)-1,3-dimethyl-butyl](2S)-2-[(3-acetoxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.05), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 5:1 to 1:5. [0050] In a preferred composition according to the invention, component (A) is [(1S,2S)-2-(4-fluoro-2-methyl-phenyl)-1,3-dimethyl-butyl](2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate (A.06), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-

(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 5:1 to 1:5.

[0051] In a preferred composition according to the invention, component (A) is [(1R,2R)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate (A.07), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 5:1 to 1:5.

[0052] In a preferred composition according to the invention, component (A) is [(1S,2S)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(3-hydroxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.01), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-

oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 4:1 to 1:4.

[0053] In a preferred composition according to the invention, component (A) is [(1S,2S)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(3-acetoxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.02), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 4:1 to 1:4.

[0054] In a preferred composition according to the invention, component (A) is [(1S,2S)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate (A.03), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 4:1 to 1:4.

[0055] In a preferred composition according to the invention, component (A) is [(1S,2S)-2-(4-fluoro-2-methyl-phenyl)-1,3-dimethyl-butyl](2S)-2-[(3-hydroxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.04), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-

(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 4:1 to 1:4.

[0056] In a preferred composition according to the invention, component (A) is [(1 S,2S)-2-(4-fluoro-2-methyl-phenyl)-1,3-dimethyl-butyl](2S)-2-[(3-acetoxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.05), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 4:1 to 1:4.

[0057] In a preferred composition according to the invention, component (A) is [(1 S,2S)-2-(4-fluoro-2-methyl-phenyl)-1,3-dimethyl-butyl](2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate (A.06), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-

oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 4:1 to 1:4.

[0058] In a preferred composition according to the invention, component (A) is [(1R,2R)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate (A.07), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 4:1 to 1:4.

[0059] In a preferred composition according to the invention, component (A) is [(1 S,2S)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(3-hydroxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.01), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 3:1 to 1:3.

[0060] In a preferred composition according to the invention, component (A) is [(1 S,2S)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(3-acetoxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.02), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-

yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 3:1 to 1:3.

[0061] In a preferred composition according to the invention, component (A) is [(1S,2S)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate (A.03), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 3:1 to 1:3.

[0062] In a preferred composition according to the invention, component (A) is [(1 S,2S)-2-(4-fluoro-2-methyl-phenyl)-1,3-dimethyl-butyl](2S)-2-[(3-hydroxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.04), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-

oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 3:1 to 1:3.

[0063] In a preferred composition according to the invention, component (A) is [(1S,2S)-2-(4-fluoro-2-methyl-phenyl)-1,3-dimethyl-butyl](2S)-2-[(3-acetoxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.05), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 3:1 to 1:3.

[0064] In a preferred composition according to the invention, component (A) is [(1 S,2S)-2-(4-fluoro-2-methyl-phenyl)-1,3-dimethyl-butyl](2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate (A.06), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 3:1 to 1:3.

[0065] In a preferred composition according to the invention, component (A) is [(1R,2R)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate (A.07), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-

yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 3:1 to 1:3.

[0066] In a preferred composition according to the invention, component (A) is [(1S,2S)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(3-hydroxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.01), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 2:1 to 1:2.

[0067] In a preferred composition according to the invention, component (A) is [(1 S,2S)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(3-acetoxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.02), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-

oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 2:1 to 1:2.

[0068] In a preferred composition according to the invention, component (A) is [(1 S,2S)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate (A.03), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 2:1 to 1:2.

[0069] In a preferred composition according to the invention, component (A) is [(1 S,2S)-2-(4-fluoro-2-methyl-phenyl)-1,3-dimethyl-butyl](2S)-2-[(3-hydroxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.04), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 2:1 to 1:2.

[0070] In a preferred composition according to the invention, component (A) is [(1 S,2S)-2-(4-fluoro-2-methyl-phenyl)-1,3-dimethyl-butyl](2S)-2-[(3-acetoxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.05), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-

(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 2:1 to 1:2.

[0071] In a preferred composition according to the invention, component (A) is [(1S,2S)-2-(4-fluoro-2-methyl-phenyl)-1,3-dimethyl-butyl](2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate (A.06), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 2:1 to 1:2.

[0072] In a preferred composition according to the invention, component (A) is [(1R,2R)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate (A.07), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-

oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is from 2:1 to 1:2.

[0073] In a preferred composition according to the invention, component (A) is [(1 S,2S)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(3-hydroxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.01), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is 1:1.

[0074] In a preferred composition according to the invention, component (A) is [(1 S,2S)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(3-acetoxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.02), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is 1:1.

[0075] In a preferred composition according to the invention, component (A) is [(1 S,2S)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate (A.03), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-

yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is 1:1.

[0076] In a preferred composition according to the invention, component (A) is [(1 S,2S)-2-(4-fluoro-2-methyl-phenyl)-1,3-dimethyl-butyl](2S)-2-[(3-hydroxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.04), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is 1:1.

[0077] In a preferred composition according to the invention, component (A) is [(1 S,2S)-2-(4-fluoro-2-methyl-phenyl)-1,3-dimethyl-butyl](2S)-2-[(3-acetoxy-4-methoxy-pyridine-2-carbonyl)amino]propanoate (A.05), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-

oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is 1:1.

[0078] In a preferred composition according to the invention, component (A) is [(1S,2S)-2-(4-fluoro-2-methyl-phenyl)-1,3-dimethyl-butyl](2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate (A.06), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is 1:1.

[0079] In a preferred composition according to the invention, component (A) is [(1R,2R)-1-methyl-2-(o-tolyl)propyl](2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate (A.07), or a salt, enantiomer, tautomer or N-oxide thereof, and component (B) is selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof, wherein the weight ratio of component (A) to component (B) is 1:1.

[0080] In any of the compositions according to the invention, the composition may comprise an additional active ingredient component (C), which is different to component (B), and is selected from the group consisting of metyltetrapole, azoxystrobin, trifloxystrobin, picoxystrobin, pyraclostrobin, metominostrobin, prothioconazole, cyproconazole, difenoconazole, tebuconazole,

propiconazole, epoxiconazole, hexaconazole, mefentrifluconazole, metconazole, benzovindiflupyr, pydiflumetofen, fluxapyroxad, sedaxane, bixafen, isopyrazam, fluopyram, fluindapyr, isoflucypram, inpyrfluxam, quinofumelin, ipflufenquin, aminopyrifin, fluazinam, fludioxonil, fenpicoxamid, florylpicoxamid, fenpropidin, fenpropimorph, mancozeb, chlorothalonil, cyclobutrifluram, Cu-oxychloride, Cu-hydroxide, N'-[5-bromo-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-ethyl-N-methyl-formamidine, N'-[5-bromo-2-methyl-6-[(1S)-1-methyl-2-propoxy-ethoxy]-3-pyridyl]-N-ethyl-N-methyl-formamidine, N'-[5-bromo-2-methyl-6-[(1R)-1-methyl-2-propoxy-ethoxy]-3-pyridyl]-N-ethyl-N-methyl-formamidine, N'-[5-bromo-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-isopropyl-N-methyl-formamidine, N'-[5-chloro-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-ethyl-N-methyl-formamidine, N'-(2-chloro-5-methyl-4-phenoxyphenyl)-N-ethyl-N-methyl-formamidine, N'-[2-chloro-4-(2-fluorophenoxy)-5-methylphenyl]-N-ethyl-N-methyl-formamidine, N'-[2-chloro-4-(3-fluorophenoxy)-5-methylphenyl]-N-ethyl-N-methyl-formamidine, N'-(2-bromo-5-methyl-4-phenoxyphenyl)-N-ethyl-N-methyl-formamidine, N'-[2-bromo-4-(2-fluorophenoxy)-5-methylphenyl]-N-ethyl-N-methyl-formamidine, N'-[2-chloro-4-(2-fluorophenoxy)-5-methylphenyl]-N-isopropyl-N-methyl-formamidine, N'-[4-(2-cyanophenoxy)-2-methyl-5-(trifluoromethyl)phenyl]-N-ethyl-N-methyl-formamidine, N'-[5-bromo-2-methyl-4-(2-methylphenoxy)phenyl]-N-ethyl-N-methyl-formamidine, N-ethyl-N'-[4-(2-fluorophenoxy)-2-methyl-5-(trifluoromethyl)phenyl]-N-methyl-formamidine, (Z,E)-5-[1-(2,4-dichlorophenyl)pyrazol-3-yl]oxy-2-methoxyimino-N,3-dimethyl-pent-3-enamide, (Z,E)-5-[1-(4-chlorophenyl)pyrazol-3-yl]oxy-2-methoxyimino-N,3-dimethyl-pent-3-enamide, 4-[[6-[2-(2,4-difluorophenyl)-1,1-difluoro-2-hydroxy-3-(5-thioxo-4H-1,2,4-triazol-1-yl)propyl]-3-pyridyl]oxy]benzonitrile, methyl 3-[(4-chlorophenyl)methyl]-2-hydroxy-1-methyl-2-(1,2,4-triazol-1-ylmethyl)cyclopentanecarboxylate, 2-[6-(4-bromophenoxy)-2-(trifluoromethyl)-3-pyridyl]-1-(1,2,4-triazol-1-yl)propan-2-ol, 2-[6-(4-chlorophenoxy)-2-(trifluoromethyl)-3-pyridyl]-1-(1,2,4-triazol-1-yl)propan-2-ol, N-methyl-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzenecarbothioamide, N-methyl-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzamide, 2,2-difluoro-N-methyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide, N-[N-methoxy-C-methyl-carbonimidoyl]-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzamide, N-[(E)-methoxyiminomethyl]-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzamide, N-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]cyclopropanecarboxamide, N-(2-fluorophenyl)-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzamide, 1-(5,6-dimethyl-3-pyridyl)-4,4-difluoro-3,3-dimethyl-isoquinoline and 1-[6-(difluoromethyl)-5-methyl-3-pyridyl]-4,4-difluoro-3,3-dimethyl-isoquinoline.

[0081] Preferably, component (C), which is different to component (B), is a compound selected from the group consisting of metyltetrapole, azoxystrobin, trifloxystrobin, picoxystrobin, metominostrobin, prothioconazole, cyproconazole, difenoconazole, tebuconazole, propiconazole, mefentrifluconazole, metconazole, benzovindiflupyr, pydiflumetofen, sedaxane, isopyrazam, inpyrfluxam, fluazinam, florylpicoxamid, fenpropidin, mancozeb, chlorothalonil, cyclobutrifluram, Cu-oxychloride, Cu-hydroxide, N'-[5-bromo-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-ethyl-N-methyl-formamidine, N'-[5-bromo-2-methyl-6-[(1S)-1-methyl-2-propoxy-ethoxy]-3-pyridyl]-N-ethyl-N-methyl-formamidine, N'-[5-bromo-2-methyl-6-[(1R)-1-methyl-2-propoxy-ethoxy]-3-pyridyl]-N-ethyl-N-methyl-formamidine and N'-[5-chloro-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-ethyl-N-methyl-formamidine.

[0082] The component (C) compounds are referred to herein and above by a so-called "ISO common name" or another "common name" being used in individual cases or a trademark name. The component (C) compounds are known and are commercially available and/or can be prepared using procedures known in the art and/or procedures reported in the literature such as, for instance, WO 2015/155075 and WO 2016/202742, WO 2017/005710, WO 2018/108977, WO 2018/153707, WO 2018/098216, WO 2019/093522, WO 2018/145921, WO 2015/185485, WO 2017/076742, WO 2017/081311, WO 2017/085100, WO 2017/093019, WO 2017/178245 and WO 2017/211649.

[0083] In embodiments of the invention where the composition comprise a component (A), a component (B) and a component (C), the weight ratio of component (A) to the sum of component (B) and component (C) may be from 100:1 to 1:100, preferably from 50:1 to 1:50, more preferably from 20:1 to 1:40, even more preferably from 15:1 to 1:30, still more preferably from 12:1 to 1:25, or from 10:1 to 1:20, or from 10:1 to 1:10, or from 5:1 and 1:15, or from 5:1 to 1:5, or from 4:1 to 1:4, or from 3:1 to 1:10, or from 3:1 to 1:3, or from 2:1 to 1:5, or 1:1.

[0084] The term “fungicide” as used herein means a compound that controls, modifies, or prevents the growth of fungi. The term “fungicidally effective amount” means the quantity of such a compound or combination of such compounds that is capable of producing an effect on the growth of fungi. Controlling or modifying effects include all deviation from natural development, such as killing, retardation and the like, and prevention includes barrier or other defensive formation in or on a plant to prevent fungal infection.

[0085] The term “plants” refers to all physical parts of a plant, including seeds, seedlings, saplings, roots, tubers, stems, stalks, foliage, and fruits.

[0086] The term “plant propagation material” denotes all generative parts of a plant, for example seeds or vegetative parts of plants such as cuttings and tubers. It includes seeds in the strict sense, as well as roots, fruits, tubers, bulbs, rhizomes, and parts of plants.

[0087] The term “locus” as used herein means fields in or on which plants are growing, or where seeds of cultivated plants are sown, or where seed will be placed into the soil. It includes soil, seeds, and seedlings, as well as established vegetation.

[0088] Throughout this document the expression “composition” stands for the various mixtures or combinations of components (A) and (B) (including the above-defined embodiments), for example in a single “ready-mix” form, in a combined spray mixture composed from separate formulations of the single active ingredient components, such as a “tank-mix”, and in a combined use of the single active ingredients when applied in a sequential manner, i.e. one after the other with a reasonably short period, such as a few hours or days. The order of applying the components (A) and (B) is not essential for working the present invention.

[0089] The composition according to the invention is effective against harmful microorganisms, such as microorganisms, that cause phytopathogenic diseases, in particular against phytopathogenic fungi and bacteria.

[0090] The composition of the invention may be used to control plant diseases caused by a broad spectrum of fungal plant pathogens in the Basidiomycete, Ascomycete, Oomycete and/or Deuteromycete, Blasocladiomycete, Chytridiomycete, Glomeromycete and/or Mucoromycete classes.

[0091] The composition is effective in controlling a broad spectrum of plant diseases, such as foliar pathogens of ornamental, turf, vegetable, field, cereal, and fruit crops.

[0092] These pathogens may include: Oomycetes, including *Phytophthora* diseases such as those caused by *Phytophthora capsici*, *Phytophthora infestans*, *Phytophthora sojae*, *Phytophthora fragariae*, *Phytophthora nicotianae*, *Phytophthora cinnamomi*, *Phytophthora citricola*, *Phytophthora citrophthora* and *Phytophthora erythroseptica*; *Pythium* diseases such as those caused by *Pythium aphanidermatum*, *Pythium arrhenomanes*, *Pythium graminicola*, *Pythium irregulare* and *Pythium ultimum*; diseases caused by Peronosporales such as *Peronospora destructor*, *Peronospora parasitica*, *Plasmopara viticola*, *Plasmopara halstedii*, *Pseudoperonospora cubensis*, *Albugo candida*, *Sclerophthora macrospora* and *Bremia lactucae*; and others such as *Aphanomyces cochlioides*, *Labyrinthula zosterae*, *Peronosclerospora sorghi* and *Sclerospora graminicola*; Ascomycetes, including blotch, spot, blast or blight diseases and/or rots for example those caused by Pleosporales such as *Stemphylium solani*, *Stagonospora tainanensis*, *Spilocaea oleaginea*, *Setosphaeria turcica*, *Pyrenochaeta lycopersici*, *Pleospora herbarum*, *Phoma destructiva*, *Phaeosphaeria herpotrichoides*, *Phaeocryptococcus gaeumannii*, *Ophiosphaerella graminicola*, *Ophiobolus graminis*, *Leptosphaeria maculans*, *Hendersonia creberrima*,

Helminthosporium tritici-repentis, *Setosphaeria turcica*, *Methylobacterium glycines*, *Didymella bryoniae*, *Cyloconium oleagineum*, *Corynespora cassiicola*, *Cochliobolus sativus*, *Bipolaris cactivora*, *Venturia inaequalis*, *Pyrenophora teres*, *Pyrenophora tritici-repentis*, *Alternaria alternata*, *Alternaria brassicicola*, *Alternaria solani* and *Alternaria tomatophila*, Capnodiales such as *Septoria tritici*, *Septoria nodorum*, *Septoria glycines*, *Cercospora arachidicola*, *Cercospora sojae*, *Cercospora zeae-maydis*, *Cercospora capsellae* and *Cercospora herpotrichoides*, *Cladosporium carpophilum*, *Cladosporium effusum*, *Passalora fulva*, *Cladosporium oxysporum*, *Dothistroma septosporum*, *Isariopsis clavispora*, *Mycosphaerella fijiensis*, *Mycosphaerella graminicola*, *Mycovellosiella koepkei*, *Phaeoisariopsis bataticola*, *Pseudocercospora vitis*, *Pseudocercospora herpotrichoides*, *Ramularia beticola*, *Ramularia collo-cygni*, Magnaporthales such as *Gaeumannomyces graminis*, *Magnaporthe grisea*, *Pyricularia oryzae*, Diaporthales such as *Anisogramma anomala*, *Apiognomonia errabunda*, *Cytospora platani*, *Diaporthe phaseolorum*, *Discula destructiva*, *Gnomonia fructicola*, *Greeneria uvicola*, *Melanconium juglandinum*, *Phomopsis viticola*, *Sirococcus clavigignenti-juglandacearum*, *Tubakia dryina*, *Dicarpella* spp., *Valsa ceratosperma*, and others such as *Actinothyrium graminis*, *Ascochyta pisi*, *Aspergillus flavus*, *Aspergillus fumigatus*, *Aspergillus nidulans*, *Asperisporium caricae*, *Blumeriella jaapii*, *Candida* spp., *Capnodium ramosum*, *Cephalosporium* spp., *Cephalosporium gramineum*, *Ceratocystis paradoxa*, *Chaetomium* spp., *Hymenoscyphus pseudoalbidus*, *Coccidioides* spp., *Cylindrosporium padi*, *Diplocarpon malae*, *Drepanopeziza campestris*, *Elsinoe ampelina*, *Epicoccum nigrum*, *Epidermophyton* spp., *Eutypa lata*, *Geotrichum candidum*, *Gibellina cerealis*, *Gloeocercospora sorghi*, *Gloeodes pomigena*, *Gloeosporium perennans*; *Gloeotinia temulenta*, *Griphosphaeria corticola*, *Kabatiella lini*, *Leptographium microsporum*, *Leptosphaerulina crassiasca*, *Lophodermium seditiosum*, *Marssonina graminicola*, *Microdochium nivale*, *Monilinia fructicola*, *Monographella albescens*, *Monosporascus cannonballus*, *Naemacyclus* spp., *Ophiostoma novoulmi*, *Paracoccidioides brasiliensis*, *Penicillium expansum*, *Pestalotia rhododendri*, *Petriellidium* spp., *Pezicula* spp., *Phialophora gregata*, *Phyllachora pomigena*, *Phymatotrichum omnivora*, *Physalospora abdita*, *Plectosporium tabacinum*, *Polyscytalum pustulans*, *Pseudopeziza medicaginis*, *Pyrenopeziza brassicae*, *Ramulispora sorghi*, *Rhabdocline pseudotsugae*, *Rhynchosporium secalis*, *Sacrocladium oryzae*, *Scedosporium* spp., *Schizothyrium pomi*, *Sclerotinia sclerotiorum*, *Sclerotinia minor*; *Sclerotium* spp., *Typhula ishikariensis*, *Seimatosporium mariae*, *Lepteutypa cupressi*, *Septocytia ruborum*, *Sphaceloma perseae*, *Sporonema phacidoides*, *Stigmata palmivora*, *Tapesia yellundae*, *Taphrina bullata*, *Thielaviopsis basicola*, *Trichoseptoria fructigena*, *Zygophiala jamaicensis*; powdery mildew diseases for example those caused by Erysiphales such as *Blumeria graminis*, *Erysiphe polygoni*, *Uncinula necator*, *Sphaerotheca fuliginea*, *Podosphaera leucotricha*, *Podosphaera macularis* *Golovinomyces cichoracearum*, *Leveillula taurica*, *Microsphaera diffusa*, *Oidiopsis gossypii*, *Phyllactinia guttata* and *Oidium arachidis*; molds for example those caused by Botryosphaeriales such as *Dothiorella aromatica*, *Diplodia seriata*, *Guignardia bidwellii*, *Botrytis cinerea*, *Botryotinia allii*, *Botryotinia fabae*, *Fusicoccum amygdali*, *Lasioidiplodia theobromae*, *Macrophoma theicola*, *Macrophoma phaseolina*, *Phyllosticta cucurbitacearum*; anthracnoses for example those caused by Glomerellales such as *Colletotrichum gloeosporioides*, *Colletotrichum lagenarium*, *Colletotrichum gossypii*, *Glomerella cingulata*, and *Colletotrichum graminicola*; and wilts or blights for example those caused by Hypocreales such as *Acremonium strictum*, *Claviceps purpurea*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium virguliforme*, *Fusarium oxysporum*, *Fusarium subglutinans*, *Fusarium oxysporum* f. sp. *cubense*, *Gerlachia nivale*, *Gibberella fujikuroi*, *Gibberella zeae*, *Gliocladium* spp., *Myrothecium verrucaria*, *Nectria ramulariae*, *Trichoderma viride*, *Trichothecium roseum*, and *Verticillium theobromae*; [0093] Basidiomycetes, including smuts for example those caused by Ustilaginales such as *Ustilago indica*, *Ustilago nuda*, *Ustilago tritici*, *Ustilago zeae*, rusts for example those caused by Pucciniales such as *Cerotelium fici*, *Chrysomyxa arctostaphyli*, *Coleosporium ipomoeae*, *Hemileia vastatrix*,

Puccinia arachidis, *Puccinia cacabata*, *Puccinia graminis*, *Puccinia recondita*, *Puccinia sorghi*, *Puccinia hordei*, *Puccinia striiformis* f. sp. *Hordei*, *Puccinia striiformis* f. sp. *Secalis*, *Pucciniastrum coryli*, or Uredinales such as *Cronartium ribicola*, *Gymnosporangium juniperi-viginianae*, *Melampsora medusae*, *Phakopsora pachyrhizi*, *Phragmidium mucronatum*, *Physopella ampelosis*, *Tranzschelia discolor* and *Uromyces viciae-fabae*; and other rots and diseases such as those caused by *Cryptococcus* spp., *Exobasidium vexans*, *Marasmiellus inoderma*, *Mycena* spp., *Sphacelotheca reiliana*, *Typhula ishikariensis*, *Urocystis agropyri*, *Itersonilia perplexans*, *Corticium invisum*, *Laetisaria fuciformis*, *Waitea circinata*, *Rhizoctonia solani*, *Thanetophorus cucurmeris*, *Entyloma dahliae*, *Entylomella microspora*, *Neovossia molinae* and *Tilletia caries*; *Blastocladiomycetes*, such as *Physoderma maydis*; [0094] *Mucoromycetes*, such as *Choanephora cucurbitarum*; *Mucor* spp.; *Rhizopus arrhizus*; as well as diseases caused by other species and genera closely related to those listed above.

[0095] In addition to their fungicidal activity, the compositions may also have activity against bacteria such as *Erwinia amylovora*, *Erwinia caratovora*, *Xanthomonas campestris*, *Pseudomonas syringae*, *Streptomyces scabies* and other related species as well as certain protozoa.

[0096] The composition according to the invention is particularly effective against phytopathogenic fungi belonging to the following classes: Ascomycetes (e.g. *Venturia*, *Podosphaera*, *Erysiphe*, *Monilinia*, *Mycosphaerella*, *Uncinula*); Basidiomycetes (e.g. the genus *Hemileia*, *Rhizoctonia*, *Phakopsora*, *Puccinia*, *Ustilago*, *Tilletia*); *Fungi imperfecti* (also known as Deuteromycetes; e.g. *Botrytis*, *Helminthosporium*, *Rhynchosporium*, *Fusarium*, *Septoria*, *Cercospora*, *Alternaria*, *Pyricularia* and *Pseudocercospora*); Oomycetes (e.g. *Phytophthora*, *Peronospora*, *Pseudoperonospora*, *Albugo*, *Bremia*, *Pythium*, *Pseudosclerospora*, *Plasmopara*).

[0097] Crops of useful plants in which the composition according to the invention can be used include perennial and annual crops, such as berry plants for example blackberries, blueberries, cranberries, raspberries and strawberries; cereals for example barley, maize (corn), millet, oats, rice, rye, sorghum triticate and wheat; fibre plants for example cotton, flax, hemp, jute and sisal; field crops for example sugar and fodder beet, coffee, hops, mustard, oilseed rape (canola), poppy, sugar cane, sunflower, tea and tobacco; fruit trees for example apple, apricot, avocado, banana, cherry, citrus, nectarine, peach, pear and plum; grasses for example Bermuda grass, bluegrass, bentgrass, centipede grass, fescue, ryegrass, St. Augustine grass and *Zoysia* grass; herbs such as basil, borage, chives, coriander, lavender, lovage, mint, oregano, parsley, rosemary, sage and thyme; legumes for example beans, lentils, peas and soya beans; nuts for example almond, cashew, ground nut, hazelnut, peanut, pecan, pistachio and walnut; palms for example oil palm; ornamentals for example flowers, shrubs and trees; other trees, for example cacao, coconut, olive and rubber; vegetables for example asparagus, aubergine, broccoli, cabbage, carrot, cucumber, garlic, lettuce, marrow, melon, okra, onion, pepper, potato, pumpkin, rhubarb, spinach and tomato; and vines for example grapes.

[0098] Crops are to be understood as being those which are naturally occurring, obtained by conventional methods of breeding, or obtained by genetic engineering. They include crops which contain so-called output traits (e.g. improved storage stability, higher nutritional value and improved flavour).

[0099] Crops are to be understood as also including those crops which have been rendered tolerant to herbicides like bromoxynil or classes of herbicides such as ALS-, EPSPS-, GS-, HPPD- and PPO-inhibitors. An example of a crop that has been rendered tolerant to imidazolinones, e.g. imazamox, by conventional methods of breeding is Clearfield® summer canola. Examples of crops that have been rendered tolerant to herbicides by genetic engineering methods include e.g. glyphosate- and glufosinate-resistant maize varieties commercially available under the trade names RoundupReady®, Herculex I® and LibertyLink®.

[0100] Crops are also to be understood as being those which naturally are or have been rendered resistant to harmful insects. This includes plants transformed by the use of recombinant DNA

techniques, for example, to be capable of synthesising one or more selectively acting toxins, such as are known, for example, from toxin-producing bacteria. Examples of toxins which can be expressed include δ -endotoxins, vegetative insecticidal proteins (Vip), insecticidal proteins of bacteria colonising nematodes, and toxins produced by scorpions, arachnids, wasps and fungi.

[0101] An example of a crop that has been modified to express the *Bacillus thuringiensis* toxin is the Bt maize KnockOut® (Syngenta Seeds). An example of a crop comprising more than one gene that codes for insecticidal resistance and thus expresses more than one toxin is VipCot® (Syngenta Seeds). Crops or seed material thereof can also be resistant to multiple types of pests (so-called stacked transgenic events when created by genetic modification). For example, a plant can have the ability to express an insecticidal protein while at the same time being herbicide tolerant, for example Herculex I® (Dow AgroSciences, Pioneer Hi-Bred International).

[0102] The fungicidal compositions according to the present invention may be used in controlling or preventing phytopathogenic diseases, especially phytopathogenic fungi (such as *Phakopsora pachyrhizi*) on soybean plants.

[0103] In particular, transgenic soybean plants expressing toxins, for example insecticidal proteins such as delta-endotoxins, e.g. Cry1Ac (Cry1Ac Bt protein). Accordingly, this may include transgenic soybean plants comprising event MON87701 (see U.S. Pat. No. 8,049,071 and related applications and patents, as well as WO 2014/170327 A1 (e.g., see paragraph [008] reference to Intacta RR2 PRO™ soybean)), event MON87751 (US. Patent Application Publication No. 2014/0373191) or event DAS-81419 (U.S. Pat. No. 8,632,978 and related applications and patents).

[0104] Other transgenic soybean plants may comprise event SYHTOH2-HPPD tolerance (U.S. Patent Application Publication No. 2014/0201860 and related applications and patents), event MON89788-glyphosate tolerance (U.S. Pat. No. 7,632,985 and related applications and patents), event MON87708-dicamba tolerance (U.S. Patent Application Publication No. US 2011/0067134 and related applications and patents), event DP-356043-5-glyphosate and ALS tolerance (U.S. Patent Application Publication No. US 2010/0184079 and related applications and patents), event A2704-12-glufosinate tolerance (U.S. Patent Application Publication No. US 2008/0320616 and related applications and patents), event DP-305423-1-ALS tolerance (U.S. Patent Application Publication No. US 2008/0312082 and related applications and patents), event A5547-127-glufosinate tolerance (U.S. Patent Application Publication No. US 2008/0196127 and related applications and patents), event DAS-40278-9-tolerance to 2,4-dichlorophenoxyacetic acid and aryloxyphenoxypropionate (see WO 2011/022469, WO 2011/022470, WO 2011/022471, and related applications and patents), event 127-ALS tolerance (WO 2010/080829 and related applications and patents), event GTS 40-3-2-glyphosate tolerance, event DAS-68416-4-2,4-dichlorophenoxyacetic acid and glufosinate tolerance, event FG72-glyphosate and isoxaflutole tolerance, event BPS-CV127-9-ALS tolerance and GU262-glufosinate tolerance or event SYHTO4R-HPPD tolerance.

[0105] The fungicidal compositions according to the present invention may be used in controlling or preventing phytopathogenic diseases, especially phytopathogenic fungi (such as *Phakopsora pachyrhizi*) on soybean plants. In particular, there are known in the scientific literature certain Elite soybean plant varieties where R-gene stacks, conferring a degree of immunity or resistance to specific *Phakopsora pachyrhizi*, have been introgressed in the plant genome, see for example: “*Fighting Asian Soybean Rust*”, Langenbach C, et al, *Front Plant Science* 7(797) 2016).

[0106] An elite plant is any plant from an elite line, such that an elite plant is a representative plant from an elite variety. Non-limiting examples of elite soybean varieties that are commercially available to farmers or soybean breeders include: AG00802, A0868, AG0902, A1923, AG2403, A2824, A3704, A4324, A5404, AG5903, AG6202 AG0934; AG1435; AG2031; AG2035; AG2433; AG2733; AG2933; AG3334; AG3832; AG4135; AG4632; AG4934; AG5831; AG6534; and AG7231 (Asgrow Seeds, Des Moines, Iowa, USA); BPR0144RR, BPR 4077NRR and BPR 4390NRR (Bio Plant Research, Camp Point, Ill., USA); DKB17-51 and DKB37-51 (DeKalb

Genetics, DeKalb, Ill., USA); DP 4546 RR, and DP 7870 RR (Delta & Pine Land Company, Lubbock, Tex., USA); JG 03R501, JG 32R606C ADD and JG 55R503C (JGL Inc., Greencastle, Ind., USA); NKS 13-K2 (NK Division of Syngenta Seeds, Golden Valley, Minnesota, USA); 90M01, 91M30, 92M33, 93M11, 94M30, 95M30, 97B52, P008T22R2; P16T17R2; P22T69R; P25T51R; P34T07R2; P35T58R; P39T67R; P47T36R; P46T21R; and P56T03R2 (Pioneer Hi-Bred International, Johnston, Iowa, USA); SG4771NRR and SG5161NRR/STS (Soygenetics, LLC, Lafayette, Ind., USA); SOO-K5, S11-L2, S28-Y2, S43-B1, S53-A1, S76-L9, S78-G6, S0009-M2; S007-Y4; S04-D3; S14-A6; S20-T6; S21-M7; S26-P3; S28-N6; S30-V6; S35-C3; S36-Y6; S39-C4; S47-K5; S48-D9; S52-Y2; S58-Z4; S67-R6; S73-S8; and S78-G6 (Syngenta Seeds, Henderson, Ky., USA); Richer (Northstar Seed Ltd. Alberta, CA); 14RD62 (Stine Seed Co. Ia., USA); or Armor 4744 (Armor Seed, LLC, Ar., USA).

[0107] Thus, in a further preferred embodiment, the fungicidal compositions according to the present invention are used to control *Phakopsora pachyrhizi*, (including fungicidally-resistant strains thereof, as outlined below) on Elite soybean plant varieties where R-gene stacks, conferring a degree of immunity or resistance to specific *Phakopsora pachyrhizi*, have been introgressed in the plant genome. Numerous benefits may be expected to ensue from said use, e.g. improved biological activity, an advantageous or broader spectrum of activity (inc. sensitive and resistant strains of *Phakopsora pachyrhizi*), an increased safety profile, improved crop tolerance, synergistic interactions or potentiating properties, improved onset of action or a longer lasting residual activity, a reduction in the number of applications and/or a reduction in the application rate of the compounds and compositions required for effective control of the phytopathogen (*Phakopsora pachyrhizi*), thereby enabling beneficial resistance-management practices, reduced environmental impact and reduced operator exposure.

[0108] Under certain circumstances, fungicidal compositions according to the present invention, when used in controlling or preventing phytopathogenic diseases, especially phytopathogenic fungi (such as *Phakopsora pachyrhizi*) on soybean plants (in particular any of the transgenic soybean plants as described above), may display a synergistic interaction between the active ingredients.

[0109] The fungicidal compositions according to the present invention may be used in controlling or preventing phytopathogenic diseases, especially phytopathogenic fungi (in particular, *Phakopsora pachyrhizi*) on soybean plants.

[0110] Additionally, to date, no cross-resistance has been observed between the component (B) (including any one of compounds B.01 to B.15) and the current fungicidal solutions used to control *Phakopsora pachyrhizi*.

[0111] Indeed, fungicidal-resistant strains of *Phakopsora pachyrhizi* have been reported in the scientific literature, with strains resistant to one or more fungicides from at least each of the following fungicidal mode of action classes being observed: sterol demethylation-inhibitors (DMI), quinone-outside-inhibitors (QoI) and succinate dehydrogenase inhibitors (SDHI). See for example: “Sensitivity of *Phakopsora pachyrhizi* towards quinone-outside-inhibitors and demethylation-inhibitors, and corresponding resistance mechanisms.” Schmitz H K et al, *Pest Manag Sci* (2014) 70: 378-388; “First detection of a SDH variant with reduced SDHI sensitivity in *Phakopsora pachyrhizi*” Simões K et al, *J Plant Dis Prot* (2018) 125: 21-2; “Competitive fitness of *Phakopsora pachyrhizi* isolates with mutations in the CYP51 and CYTB genes.” Klosowski A C et al, *Phytopathology* (2016) 106: 1278-1284; “Detection of the F129L mutation in the cytochrome b gene in *Phakopsora pachyrhizi*.” Klosowski A C et al, *Pest Manag Sci* (2016) 72: 1211-1215.

[0112] Thus, in a preferred embodiment, the fungicidal compositions according to the present invention are used to control *Phakopsora pachyrhizi* which are resistant to one or more fungicides from any of the following fungicidal MoA classes: sterol demethylation-inhibitors (DMI), quinone-outside-inhibitors (QoI) and succinate dehydrogenase inhibitors (SDHI).

[0113] The component (A) used in accordance with the present invention can be made according to any of the synthetic techniques as described in WO 2019/173665.

[0114] The component (B) used in accordance with the present invention can be made according to any of the synthetic techniques as described in WO 2017/055469, WO 2017/055473, WO 2017/093348, WO 2017/118689, WO 2017/220485, WO 2018/065414, WO 2018/158365, WO 2018/177894, WO 2018/177880, WO 2018/219773 and WO 2019/002151.

[0115] It is understood that, when in aqueous media, the component (B) according to the invention may be present in a reversible equilibrium with the corresponding covalently hydrated forms. This dynamic equilibrium may be important for the biological activity of the component (B).

[0116] Compositions of this invention, including all of the above disclosed embodiments and preferred examples thereof, can be mixed with one or more further pesticides including further fungicides, insecticides, nematocides, bactericides, acaricides, growth regulators, chemosterilants, semiochemicals, repellents, attractants, pheromones, feeding stimulants or other biologically active compounds to form a multi-component pesticide giving an even broader spectrum of agricultural protection.

[0117] Examples of such agricultural protectants with which the composition of this invention can be formulated are:

[0118] Fungicides such as etridiazole, fluazinam, benalaxyl, benalaxyl-M (kiralaxyl), furalaxyl, metalaxyl, metalaxyl-M (mefenoxam), dodiclin, N'-(2,5-dimethyl-4-phenoxy-phenyl)-N-ethyl-N-methyl-formamidine, N'-[4-(4,5-dichloro-thiazol-2-yloxy)-2,5-dimethyl-phenyl]-N-ethyl-N-methyl-formamidine, N'-[4-[[3-[(4-chlorophenyl)methyl]-1,2,4-thiadiazol-5-yl]oxy]-2,5-dimethyl-phenyl]-N-ethyl-N-methyl-formamidine, ethirimol, 3'-chloro-2-methoxy-N-[(3RS)-tetrahydro-2-oxofuran-3-yl]acet-2',6'-xylidide (clozylacon), cyprodinil, mepanipyrim, pyrimethanil, dithianon, aureofungin, blasticidin-S, biphenyl, chloroneb, dicloran, benzovindiflupyr, pydiflumetofen, hexachlorobenzene, quintozone, tecnazene, (TCNB), tolclofos-methyl, metrafenone, 2,6-dichloro-N-(4-trifluoromethylbenzyl)-benzamide, fluopicolide (flupicolide), tioxyimid, flusulfamide, benomyl, carbendazim, carbendazim chlorhydrate, chlorfenazole, fuberidazole, thiabendazole, thiophanate-methyl, benthiavalicarb, chlobenthiazole, probenazole, acibenzolar, bethoxazin, pyriofenone (IKF-309), acibenzolar-S-methyl, pyribencarb (KIF-7767), butylamine, 3-iodo-2-propinyl n-butylcarbamate (IPBC), iodocarb (isopropanyl butylcarbamate), isopropanyl butylcarbamate (iodocarb), picarbutrazox, polycarbamate, propamocarb, tolprocarb, 3-(difluoromethyl)-N-(7-fluoro-1,1,3,3-tetramethyl-indan-4-yl)-1-methyl-pyrazole-4-carboxamide diclocymet, N-[(5-chloro-2-isopropyl-phenyl)methyl]-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-pyrazole-4-carboxamide N-cyclopropyl-3-(difluoromethyl)-5-fluoro-N-[(2-isopropylphenyl)methyl]-1-methyl-pyrazole-4-carboxamide carpropamid, chlorothalonil, flumorph, oxine-copper, cymoxanil, phenamacril, cyazofamid, flutianil, thicyofen, chlozolate, iprodione, procymidone, vinclozolin, bupirimate, dinocron, dinopenton, dinobuton, dinocap, meptyldinocap, diphenylamine, phosdiphen, 2,6-dimethyl-[1,4]dithiino[2,3-c:5,6-c']dipyrrole-1,3,5,7(2H,6H)-tetraone, azithiram, etem, ferbam, mancozeb, maneb, metam, metiram (polyram), metiram-zinc, nabam, propineb, thiram, vapam (metam sodium), zineb, ziram, dithioether, isoprothiolane, ethaboxam, fosetyl, fosetyl-Al (fosetyl-al), methyl bromide, methyl iodide, methyl isothiocyanate, cyclafuramid, fenfuram, validamycin, streptomycin, (2RS)-2-bromo-2-(bromomethyl)glutaronitrile (bromothalonil), dodine, doguadine, guazatine, iminoctadine, iminoctadine triacetate, 2,4-D, 2,4-DB, kasugamycin, dimethirimol, fenhexamid, hymexazole, hydroxyisoxazole imazalil, imazalil sulphate, oxpoconazole, pefurazoate, prochloraz, triflumizole, fenamidone, Bordeaux mixture, calcium polysulfide, copper acetate, copper carbonate, copper hydroxide, copper naphthenate, copper oleate, copper oxychloride, copper oxyquinolate, copper silicate, copper sulphate, copper tallate, cuprous oxide, sulphur, carbaryl, phthalide (fthalide), dingjunezuo (Jun Si Qi), oxathiapiprolin, fluoroimide, mandipropamid, KSF-1002, benzamorf, dimethomorph, fenpropimorph, tridemorph, dodemorph, diethofencarb, fentin acetate, fentin hydroxide, carboxin, oxycarboxin, drazoxolon, famoxadone, m-phenylphenol, p-phenylphenol, tribromophenol (TBP), 2-[2-[(7,8-difluoro-2-methyl-3-quinolyl)oxy]-6-fluoro-phenyl]propan-2-ol,

2-[2-fluoro-6-[(8-fluoro-2-methyl-3-quinolyl)oxy]phenyl]propan-2-ol, cyflufenamid, ofurace, oxadixyl, flutolanil, mepronil, isofetamid, fenpiclonil, fludioxonil, pencycuron, edifenphos, iprobenfos, pyrazophos, phosphorus acids, tecloftalam, captafol, captan, ditalimfos, triforine, fenpropidin, piperalin, osthonol, 1-methylcyclopropene, 4-CPA, chlormequat, clofencet, dichlorprop, dimethipin, endothal, ethephon, flumetralin, forchlorfenuron, gibberellic acid, gibberellins, hymexazol, maleic hydrazide, mepiquat, naphthalene acetamide, paclobutrazol, prohexadione, prohexadione-calcium, thidiazuron, tribufos (tributyl phosphorotrithioate), trinexapac, uniconazole, α -naphthalene acetic acid, polyoxin D (polyoxim), BLAD, chitosan, fenoxanil, folpet, 3-(difluoromethyl)-N-methoxy-1-methyl-N-[1-methyl-2-(2,4,6-trichlorophenyl)ethyl]pyrazole-4-carboxamide, bixafen, fluxapyroxad, furametpyr, isopyrazam, penflufen, penthiopyrad, sedaxane, fenpyrazamine, diclomezine, pyrifenoxy, boscalid, fluopyram, diflumetorim, fenarimol, 5-fluoro-2-(p-tolylmethoxy)pyrimidin-4-amine ferimzone, dimetachlone (dimethaclone), pyroquilon, proquinazid, ethoxyquin, quinoxifen, 4,4,5-trifluoro-3,3-dimethyl-1-(3-quinolyl)isoquinoline, 4,4-difluoro-3,3-dimethyl-1-(3-quinolyl)isoquinoline 5-fluoro-3,3,4,4-tetramethyl-1-(3-quinolyl)isoquinoline 9-fluoro-2,2-dimethyl-5-(3-quinolyl)-3H-1,4-benzoxazepine, tebufloquin, oxolinic acid, chinomethionate (oxythioquinox, quinoxymethionate), spiroxamine, (E)-N-methyl-2-[2-(2, 5-dimethylphenoxyethyl) phenyl]-2-methoxy-iminoacetamide, (mandestrobin), azoxystrobin, coumoxystrobin, dimoxystrobin, enestroburin, enoxastrobin, fenamistobin, flufenoxystrobin, fluoxastrobin, kresoxim-methyl, mandestrobin, metaminostrobin, metaminostrobin, orysastrobin, picoxystrobin, pyraclostrobin, pyrametostrobin, pyraoxystrobin, triclopyricarb, trifloxystrobin, amisulbrom, dichlofluanid, tolylfluanid, but-3-ynyl N-[6-[(Z)-[(1-methyltetrazol-5-yl)-phenyl-methylene]amino]oxymethyl]-2-pyridyl]carbamate, dazomet, isotianil, tiadinil, thifluzamide, benthiazole (TCMTB), silthiofam, zoxamide, anilazine, tricyclazole, (+.-)-cis-1-(4-chlorophenyl)-2-(1H-1,2,4-triazol-1-yl)-cycloheptanol (huanjunzuo), 1-(5-bromo-2-pyridyl)-2-(2,4-difluorophenyl)-1,1-difluoro-3-(1,2,4-triazol-1-yl)propan-2-ol 2-(1-tert-butyl)-1-(2-chlorophenyl)-3-(1,2,4-triazol-1-yl)-propan-2-ol (TCDP), azaconazole, bitertanol (biloxazol), bromuconazole, climbazole, cyproconazole, difenoconazole, dimetconazole, diniconazole, diniconazole-M, epoxiconazole, etaconazole, fenbuconazole, fluquinconazole, flusilazole, flutriafol, hexaconazole, imibenconazole, ipconazole, ipfentrifluconazole, metconazole, myclobutanil, penconazole, propiconazole, prothioconazole, simeconazole, tebuconazole, tetraconazole, triadimefon, triadimenol, triazoxide, triticonazole, mefentrifluconazole, 2-[[[(1R,5S)-5-[(4-fluorophenyl)methyl]-1-hydroxy-2,2-dimethyl-cyclopentyl]methyl]-4H-1,2,4-triazole-3-thione, 2-[[[3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl]-4H-1,2,4-triazole-3-thione, ametoctradin (imidium), iprovalicarb, valifenalate, 2-benzyl-4-chlorophenol (Chlorophene), allyl alcohol, azafenidin, benzalkonium chloride, chloropicrin, cresol, daracide, dichlorophen (dichlorophene), difenzoquat, dipyrithione, N-(2-p-chlorobenzoyl)ethyl)-hexaminium chloride, NNF-0721, octhilinone, oxasulfuron, propamidin and propionic acid.

[0119] Insecticides such as abamectin, acephate, acetamiprid, amidoflumet (S-1955), avermectin, azadirachtin, azinphos-methyl, bifenthrin, bifenazate, buprofezin, carbofuran, cartap, chlorantraniliprole (DPX-E2Y45), chlorfenapyr, chlorfluazuron, chlorpyrifos, chlorpyrifos-methyl, chromafenozide, clothianidin, cyflumetofen, cyfluthrin, beta-cyfluthrin, cyhalothrin, lambda-cyhalothrin, cypermethrin, cyromazine, deltamethrin, diafenthiuron, diazinon, dieldrin, diflubenzuron, dimefluthrin, dimethoate, dinotefuran, diofenolan, emamectin, endosulfan, esfenvalerate, ethiprole, fenothiocarb, fenoxycarb, fenpropathrin, fenvalerate, fipronil, flonicamid, flubendiamide, flucythrinate, tau-fluvalinate, flufenerim (UR-50701), flufenoxuron, fonophos, halofenozide, hexaflumuron, hydramethylnon, imidacloprid, indoxacarb, isofenphos, lufenuron, malathion, metaflumizone, metaldehyde, methamidophos, methidathion, methomyl, methoprene, methoxychlor, metofluthrin, monocrotophos, methoxyfenozide, nitenpyram, nithiazine, novaluron, noviflumuron (XDE-007), oxamyl, parathion, parathion-methyl, permethrin, phorate, phosalone, phosmet, phosphamidon, pirimicarb, profenofos, profluthrin, pymetrozine, pyrafluprole, pyrethrin,

pyridalyl, pyrifluquinazon, pyriprole, pyriproxyfen, rotenone, ryanodine, spinetoram, spinosad, spirotetramat, sulprofos, tebufenozide, teflubenzuron, tefluthrin, terbufos, tetrachlorvinphos, thiachlorid, thiamethoxam, thiodicarb, thiosultap-sodium, tralomeethrin, triazamate, trichlorfon and triflumuron; [0120] Bactericides such as streptomycin; [0121] Acaricides such as amitraz, chinomethionat, chlorobenzilate, cyenopyrafen, cyhexatin, dicofol, dienochlor, etoxazole, fenazaquin, fenbutatin oxide, fenpropathrin, fenpyroximate, hexythiazox, propargite, pyridaben and tebufenpyrad; and [0122] Biological agents such as *Bacillus thuringiensis*, *Bacillus thuringiensis* delta endotoxin, baculovirus, and entomopathogenic bacteria, virus and fungi.

[0123] The compositions as described above can be used in a method for controlling pests, which comprises applying a composition comprising a mixture as described above to the pests or their environment.

[0124] The mixtures comprising a component (B) selected from Table B (above) and one or more active ingredients as described above can be applied, for example, in a single “ready-mix” form, in a combined spray mixture composed from separate formulations of the single active ingredient components, such as a “tank-mix”, and in a combined use of the single active ingredients when applied in a sequential manner, i.e. one after the other with a reasonably short period, such as a few hours or days. The order of applying the component (B) selected from Table B (above) and the active ingredients as described above is not essential for working the present invention.

[0125] The compositions of the present invention may also be used in crop enhancement.

According to the present invention, ‘crop enhancement’ means an improvement in plant vigour, an improvement in plant quality, improved tolerance to stress factors, and/or improved input use efficiency.

[0126] According to the present invention, an ‘improvement in plant vigour’ means that certain traits are improved qualitatively or quantitatively when compared with the same trait in a control plant which has been grown under the same conditions in the absence of the method of the invention. Such traits include, but are not limited to, early and/or improved germination, improved emergence, the ability to use less seeds, increased root growth, a more developed root system, increased root nodulation, increased shoot growth, increased tillering, stronger tillers, more productive tillers, increased or improved plant stand, less plant verse (lodging), an increase and/or improvement in plant height, an increase in plant weight (fresh or dry), bigger leaf blades, greener leaf colour, increased pigment content, increased photosynthetic activity, earlier flowering, longer panicles, early grain maturity, increased seed, fruit or pod size, increased pod or ear number, increased seed number per pod or ear, increased seed mass, enhanced seed filling, less dead basal leaves, delay of senescence, improved vitality of the plant, increased levels of amino acids in storage tissues and/or less inputs needed (e.g. less fertiliser, water and/or labour needed). A plant with improved vigour may have an increase in any of the aforementioned traits or any combination or two or more of the aforementioned traits.

[0127] According to the present invention, an ‘improvement in plant quality’ means that certain traits are improved qualitatively or quantitatively when compared with the same trait in a control plant which has been grown under the same conditions in the absence of the method of the invention. Such traits include, but are not limited to, improved visual appearance of the plant, reduced ethylene (reduced production and/or inhibition of reception), improved quality of harvested material, e.g. seeds, fruits, leaves, vegetables (such improved quality may manifest as improved visual appearance of the harvested material), improved carbohydrate content (e.g. increased quantities of sugar and/or starch, improved sugar acid ratio, reduction of reducing sugars, increased rate of development of sugar), improved protein content, improved oil content and composition, improved nutritional value, reduction in anti-nutritional compounds, improved organoleptic properties (e.g. improved taste) and/or improved consumer health benefits (e.g. increased levels of vitamins and anti-oxidants), improved post-harvest characteristics (e.g.

enhanced shelf-life and/or storage stability, easier processability, easier extraction of compounds), more homogenous crop development (e.g. synchronised germination, flowering and/or fruiting of plants), and/or improved seed quality (e.g. for use in following seasons). A plant with improved quality may have an increase in any of the aforementioned traits or any combination or two or more of the aforementioned traits.

[0128] According to the present invention, an 'improved tolerance to stress factors' means that certain traits are improved qualitatively or quantitatively when compared with the same trait in a control plant which has been grown under the same conditions in the absence of the method of the invention. Such traits include, but are not limited to, an increased tolerance and/or resistance to abiotic stress factors which cause sub-optimal growing conditions such as drought (e.g. any stress which leads to a lack of water content in plants, a lack of water uptake potential or a reduction in the water supply to plants), cold exposure, heat exposure, osmotic stress, UV stress, flooding, increased salinity (e.g. in the soil), increased mineral exposure, ozone exposure, high light exposure and/or limited availability of nutrients (e.g. nitrogen and/or phosphorus nutrients). A plant with improved tolerance to stress factors may have an increase in any of the aforementioned traits or any combination or two or more of the aforementioned traits. In the case of drought and nutrient stress, such improved tolerances may be due to, for example, more efficient uptake, use or retention of water and nutrients.

[0129] According to the present invention, an 'improved input use efficiency' means that the plants are able to grow more effectively using given levels of inputs compared to the grown of control plants which are grown under the same conditions in the absence of the method of the invention. In particular, the inputs include, but are not limited to fertiliser (such as nitrogen, phosphorous, potassium, micronutrients), light and water. A plant with improved input use efficiency may have an improved use of any of the aforementioned inputs or any combination of two or more of the aforementioned inputs.

[0130] Other crop enhancements of the present invention include a decrease in plant height, or reduction in tillering, which are beneficial features in crops or conditions where it is desirable to have less biomass and fewer tillers.

[0131] Any or all of the above crop enhancements may lead to an improved yield by improving e.g. plant physiology, plant growth and development and/or plant architecture. In the context of the present invention 'yield' includes, but is not limited to, (i) an increase in biomass production, grain yield, starch content, oil content and/or protein content, which may result from (a) an increase in the amount produced by the plant per se or (b) an improved ability to harvest plant matter, (ii) an improvement in the composition of the harvested material (e.g. improved sugar acid ratios, improved oil composition, increased nutritional value, reduction of anti-nutritional compounds, increased consumer health benefits) and/or (iii) an increased/facilitated ability to harvest the crop, improved processability of the crop and/or better storage stability/shelf life. Increased yield of an agricultural plant means that, where it is possible to take a quantitative measurement, the yield of a product of the respective plant is increased by a measurable amount over the yield of the same product of the plant produced under the same conditions, but without application of the present invention. According to the present invention, it is preferred that the yield be increased by at least 0.5%, more preferred at least 1%, even more preferred at least 2%, still more preferred at least 4%, preferably 5% or even more.

[0132] Any or all of the above crop enhancements may also lead to an improved utilisation of land, i.e. land which was previously unavailable or sub-optimal for cultivation may become available. For example, plants which show an increased ability to survive in drought conditions, may be able to be cultivated in areas of sub-optimal rainfall, e.g. perhaps on the fringe of a desert or even the desert itself.

[0133] In one aspect of the present invention, crop enhancements are made in the substantial absence of pressure from pests and/or diseases and/or abiotic stress. In a further aspect of the

present invention, improvements in plant vigour, stress tolerance, quality and/or yield are made in the substantial absence of pressure from pests and/or diseases. For example, pests and/or diseases may be controlled by a pesticidal treatment that is applied prior to, or at the same time as, the method of the present invention. In a still further aspect of the present invention, improvements in plant vigour, stress tolerance, quality and/or yield are made in the absence of pest and/or disease pressure. In a further embodiment, improvements in plant vigour, quality and/or yield are made in the absence, or substantial absence, of abiotic stress.

[0134] The compositions of the present invention may also be used in the field of protecting storage goods against attack of fungi. According to the present invention, the term “storage goods” is understood to denote natural substances of vegetable and/or animal origin and their processed forms, which have been taken from the natural life cycle and for which long-term protection is desired. Storage goods of vegetable origin, such as plants or parts thereof, for example stalks, leaves, tubers, seeds, fruits or grains, can be protected in the freshly harvested state or in processed form, such as pre-dried, moistened, comminuted, ground, pressed or roasted. Also falling under the definition of storage goods is timber, whether in the form of crude timber, such as construction timber, electricity pylons and barriers, or in the form of finished articles, such as furniture or objects made from wood. Storage goods of animal origin are hides, leather, furs, hairs and the like. The composition according to the present invention can prevent disadvantageous effects such as decay, discoloration or mold. Preferably “storage goods” is understood to denote natural substances of vegetable origin and/or their processed forms, more preferably fruits and their processed forms, such as pomes, stone fruits, soft fruits and citrus fruits and their processed forms. In another preferred embodiment of the invention “storage goods” is understood to denote wood.

[0135] Therefore, a further aspect of the present invention is a method of protecting storage goods, which comprises applying to the storage goods a composition according to the invention.

[0136] The composition of the present invention may also be used in the field of protecting technical material against attack of fungi. According to the present invention, the term “technical material” includes paper; carpets; constructions; cooling and heating systems; wall-boards; ventilation and air conditioning systems and the like; preferably “technical material” is understood to denote wall-boards. The composition according to the present invention can prevent disadvantageous effects such as decay, discoloration or mold.

[0137] The composition according to the invention is generally formulated in various ways using formulation adjuvants, such as carriers, solvents and surface-active substances. The formulations can be in various physical forms, e.g. in the form of dusting powders, gels, wettable powders, water-dispersible granules, water-dispersible tablets, effervescent pellets, emulsifiable concentrates, micro-emulsifiable concentrates, oil-in-water emulsions, oil-flowables, aqueous dispersions, oily dispersions, suspo-emulsions, capsule suspensions, emulsifiable granules, soluble liquids, water-soluble concentrates (with water or a water-miscible organic solvent as carrier), impregnated polymer films or in other forms known e.g. from the Manual on Development and Use of FAO and WHO Specifications for Pesticides, United Nations, First Edition, Second Revision (2010). Such formulations can either be used directly or diluted prior to use. The dilutions can be made, for example, with water, liquid fertilisers, micronutrients, biological organisms, oil or solvents.

[0138] The formulations can be prepared e.g. by mixing the active ingredient with the formulation adjuvants in order to obtain compositions in the form of finely divided solids, granules, solutions, dispersions or emulsions. The active ingredients can also be formulated with other adjuvants, such as finely divided solids, mineral oils, oils of vegetable or animal origin, modified oils of vegetable or animal origin, organic solvents, water, surface-active substances or combinations thereof.

[0139] The active ingredients can also be contained in microcapsules. Microcapsules contain the active ingredients in a porous carrier. This enables the active ingredients to be released into the environment in controlled amounts (e.g. slow-release). Microcapsules usually have a diameter of from 0.1 to 500 microns. They contain active ingredients in an amount of about from 25 to 95% by

weight of the capsule weight. The active ingredients can be in the form of a monolithic solid, in the form of fine particles in solid or liquid dispersion or in the form of a suitable solution. The encapsulating membranes can comprise, for example, natural or synthetic rubbers, cellulose, styrene/butadiene copolymers, polyacrylonitrile, polyacrylate, polyesters, polyamides, polyureas, polyurethane or chemically modified polymers and starch xanthates or other polymers that are known to the person skilled in the art. Alternatively, very fine microcapsules can be formed in which the active ingredient is contained in the form of finely divided particles in a solid matrix of base substance, but the microcapsules are not themselves encapsulated.

[0140] The formulation adjuvants that are suitable for the preparation of the formulations according to the invention are known per se. As liquid carriers there may be used: water, toluene, xylene, petroleum ether, vegetable oils, acetone, methyl ethyl ketone, cyclohexanone, acid anhydrides, acetonitrile, acetophenone, amyl acetate, 2-butanone, butylene carbonate, chlorobenzene, cyclohexane, cyclohexanol, alkyl esters of acetic acid, diacetone alcohol, 1,2-dichloropropane, diethanolamine, p-diethylbenzene, diethylene glycol, diethylene glycol abietate, diethylene glycol butyl ether, diethylene glycol ethyl ether, diethylene glycol methyl ether, N,N-dimethylformamide, dimethyl sulfoxide, 1,4-dioxane, dipropylene glycol, dipropylene glycol methyl ether, dipropylene glycol dibenzoate, diproxitol, alkylpyrrolidone, ethyl acetate, 2-ethylhexanol, ethylene carbonate, 1,1,1-trichloroethane, 2-heptanone, alpha-pinene, d-limonene, ethyl lactate, ethylene glycol, ethylene glycol butyl ether, ethylene glycol methyl ether, gamma-butyrolactone, glycerol, glycerol acetate, glycerol diacetate, glycerol triacetate, hexadecane, hexylene glycol, isoamyl acetate, isobornyl acetate, isooctane, isophorone, isopropylbenzene, isopropyl myristate, lactic acid, laurylamine, mesityl oxide, methoxypropanol, methyl isoamyl ketone, methyl isobutyl ketone, methyl laurate, methyl octanoate, methyl oleate, methylene chloride, m-xylene, n-hexane, n-octylamine, octadecanoic acid, octylamine acetate, oleic acid, oleylamine, o-xylene, phenol, polyethylene glycol, propionic acid, propyl lactate, propylene carbonate, propylene glycol, propylene glycol methyl ether, p-xylene, toluene, triethyl phosphate, triethylene glycol, xylenesulfonic acid, paraffin, mineral oil, trichloroethylene, perchloroethylene, ethyl acetate, amyl acetate, butyl acetate, propylene glycol methyl ether, diethylene glycol methyl ether, methanol, ethanol, isopropanol, and alcohols of higher molecular weight, such as amyl alcohol, tetrahydrofurfuryl alcohol, hexanol, octanol, ethylene glycol, propylene glycol, glycerol, N-methyl-2-pyrrolidone and the like.

[0141] Suitable solid carriers are, for example, talc, titanium dioxide, pyrophyllite clay, silica, attapulgite clay, kieselguhr, limestone, calcium carbonate, bentonite, calcium montmorillonite, cottonseed husks, wheat flour, soybean flour, pumice, wood flour, ground walnut shells, lignin and similar substances.

[0142] A large number of surface-active substances can advantageously be used in both solid and liquid formulations, especially in those formulations which can be diluted with a carrier prior to use. Surface-active substances may be anionic, cationic, non-ionic or polymeric and they can be used as emulsifiers, wetting agents or suspending agents or for other purposes. Typical surface-active substances include, for example, salts of alkyl sulfates, such as diethanolammonium lauryl sulfate; salts of alkylarylsulfonates, such as calcium dodecylbenzenesulfonate; alkylphenol/alkylene oxide addition products, such as nonylphenol ethoxylate; alcohol/alkylene oxide addition products, such as tridecylalcohol ethoxylate; soaps, such as sodium stearate; salts of alkyl naphthalenesulfonates, such as sodium dibutyl naphthalenesulfonate; dialkyl esters of sulfosuccinate salts, such as sodium di(2-ethylhexyl)sulfosuccinate; sorbitol esters, such as sorbitol oleate; quaternary amines, such as lauryltrimethylammonium chloride, polyethylene glycol esters of fatty acids, such as polyethylene glycol stearate; block copolymers of ethylene oxide and propylene oxide; and salts of mono- and di-alkylphosphate esters; and also further substances described e.g. in McCutcheon's Detergents and Emulsifiers Annual, MC Publishing Corp., Ridgewood New Jersey (1981).

[0143] Further adjuvants that can be used in pesticidal formulations include crystallisation inhibitors, viscosity modifiers, suspending agents, dyes, anti-oxidants, foaming agents, light absorbers, mixing auxiliaries, antifoams, complexing agents, neutralising or pH-modifying substances and buffers, corrosion inhibitors, fragrances, wetting agents, take-up enhancers, micronutrients, plasticisers, glidants, lubricants, dispersants, thickeners, antifreezes, microbicides, and liquid and solid fertilisers.

[0144] The formulations according to the invention can include an additive comprising an oil of vegetable or animal origin, a mineral oil, alkyl esters of such oils or mixtures of such oils and oil derivatives. The amount of oil additive in the formulation according to the invention is generally from 0.01 to 10%, based on the mixture to be applied. For example, the oil additive can be added to a spray tank in the desired concentration after a spray mixture has been prepared. Preferred oil additives comprise mineral oils or an oil of vegetable origin, for example rapeseed oil, olive oil or sunflower oil, emulsified vegetable oil, alkyl esters of oils of vegetable origin, for example the methyl derivatives, or an oil of animal origin, such as fish oil or beef tallow. Preferred oil additives comprise alkyl esters of C.sub.8-C.sub.22 fatty acids, especially the methyl derivatives of C.sub.12-C.sub.18 fatty acids, for example the methyl esters of lauric acid, palmitic acid and oleic acid (methyl laurate, methyl palmitate and methyl oleate, respectively). Many oil derivatives are known from the Compendium of Herbicide Adjuvants, 10.sup.th Edition, Southern Illinois University, 2010.

[0145] The formulations generally comprise from 0.1 to 99% by weight, especially from 0.1 to 95% by weight, of compounds of component (A) and component (B) and from 1 to 99.9% by weight of a formulation adjuvant which preferably includes from 0 to 25% by weight of a surface-active substance. Whereas commercial products may preferably be formulated as concentrates, the end user will normally employ dilute formulations.

[0146] The rates of application vary within wide limits and depend on the nature of the soil, the method of application, the crop plant, the pest to be controlled, the prevailing climatic conditions, and other factors governed by the method of application, the time of application and the target crop. As a general guideline compounds may be applied at a rate of from 1 to 2000 l/ha, especially from 10 to 1000 l/ha.

[0147] Certain compositions as described above may show a synergistic effect. This occurs whenever the action of an active ingredient combination is greater than the sum of the actions of the individual components. The action to be expected E for a given active ingredient combination obeys the so-called COLBY formula and can be calculated as follows (COLBY, S. R. "Calculating synergistic and antagonistic responses of herbicide combination". Weeds, Vol. 15, pages 20-22; 1967): [0148] ppm=milligrams of active ingredient (=a.i.) per liter of spray mixture, [0149] X=% action by active ingredient A) using p ppm of active ingredient, [0150] Y=% action by active ingredient B) using q ppm of active ingredient.

[0151] According to COLBY, the expected (additive) action of active ingredients A)+B) using p+q ppm of active ingredient is:

$$[00001] E = X + Y - \frac{X \cdot \text{Math. } Y}{100}.$$

[0152] If the action actually observed (O) is greater than the expected action (E), then the action of the combination is super-additive, i.e. there is a synergistic effect. In mathematical terms, synergism corresponds to a positive value for the difference of (O-E). In the case of purely complementary addition of activities (expected activity), said difference (O-E) is zero. A negative value of said difference (O-E) signals a loss of activity compared to the expected activity.

[0153] However, besides the actual synergistic action with respect to fungicidal activity, the composition according to the invention may also have further surprising advantageous properties. Examples of such advantageous properties that may be mentioned are: more advantageous degradability; improved toxicological and/or ecotoxicological behaviour; or improved characteristics of the useful plants including: emergence, crop yields, more developed root system,

tillering increase, increase in plant height, bigger leaf blade, less dead basal leaves, stronger tillers, greener leaf colour, less fertilizers needed, less seeds needed, more productive tillers, earlier flowering, early grain maturity, less plant verse (lodging), increased shoot growth, improved plant vigor, and early germination.

[0154] The composition according to the invention can be applied to the phytopathogenic microorganisms, the useful plants, the locus thereof, the propagation material thereof, storage goods or technical materials threatened by microorganism attack.

[0155] The composition according to the invention may be applied before or after infection of the useful plants, the propagation material thereof, storage goods or technical materials by the microorganisms.

[0156] The amount of a composition according to the invention to be applied, will depend on various factors, such as the compounds employed; the subject of the treatment, such as, for example plants, soil or seeds; the type of treatment, such as, for example spraying, dusting or seed dressing; the purpose of the treatment, such as, for example prophylactic or therapeutic; the type of fungi to be controlled or the application time.

[0157] When applied to the useful plants, component (A) is typically applied at a rate of from 5 to 2000 g a.i./ha, particularly from 10 to 1000 g a.i./ha, preferably from 5 to 200 g a.i./ha, e.g. 20, 50, 75, 100 or 200 g a.i./ha, typically in association with a rate of from 1 to 5000 g a.i./ha, particularly from 2 to 2000 g a.i./ha, preferably from 5 to 200 g a.i./ha, e.g. 30, 60, 100, 250, 500, 800, 1000, 1500 g a.i./ha of component (B).

[0158] In agricultural practice the application rates of the composition according to the invention depend on the type of effect desired, and typically range from 20 to 4000 g of total composition per hectare.

[0159] When the composition according to the invention is used for treating seed, rates of 0.001 to 50 g of a compound of component (A) per kg of seed, preferably from 0.01 to 10 g per kg of seed, and 0.001 to 50 g of a compound of component (B), per kg of seed, preferably from 0.01 to 10 g per kg of seed, are generally sufficient.

[0160] For the avoidance of doubt, where a literary reference, patent application, or patent, is cited within the text of this application, the entire text of said citation is herein incorporated by reference.

Description

EXAMPLES

[0161] The Examples which follow serve to illustrate the invention. The compounds (and compositions) of the invention may be distinguished from known compounds (and compositions) by virtue of greater efficacy at low application rates, which can be verified by the person skilled in the art using the experimental procedures outlined in the Examples, using lower application rates if necessary, for example 50 ppm, 12.5 ppm, 6 ppm, 3 ppm, 1.5 ppm or 0.2 ppm of active ingredient(s).

[0162] Throughout this description, temperatures are given in degrees Celsius (° C.) and “mp.” means melting point. LC/MS means Liquid Chromatography Mass Spectrometry and the description of the apparatus and the method (Methods A and B) is as follows:

The description of the LC/MS apparatus and the method A is: [0163] SQ Detector 2 from Waters

[0164] Ionisation method: Electrospray [0165] Polarity: positive and negative ions [0166] Capillary (kV) 3.0, Cone (V) 30.00, Extractor (V) 2.00, Source Temperature (° C.) 150, Desolvation [0167] Temperature (° C.) 350, Cone Gas Flow (L/Hr) 0, Desolvation Gas Flow (L/Hr) 650 [0168] Mass range: 100 to 900 Da [0169] DAD Wavelength range (nm): 210 to 500

[0170] Method Waters ACQUITY UPLC with the following HPLC gradient conditions: [0171] (Solvent A: Water/Methanol 20:1+0.05% formic acid and Solvent B: Acetonitrile+0.05% formic

acid)

TABLE-US-00004 Time (minutes) A (%) B (%) Flow rate (ml/min) 0 100 0 0.85 1.2 0 100 0.85 1.5 0 100 0.85

[0172] Type of column: Waters ACQUITY UPLC HSS T3; Column length: 30 mm; Internal diameter of column: 2.1 mm; Particle Size: 1.8 micron; Temperature: 60° C.

The description of the LC/MS apparatus and the method B is: [0173] SQ Detector 2 from Waters

[0174] Ionisation method: Electrospray [0175] Polarity: positive ions [0176] Capillary (kV) 3.5, Cone (V) 30.00, Extractor (V) 3.00, Source Temperature (° C.) 150, Desolvation [0177]

Temperature (° C.) 400, Cone Gas Flow (L/Hr) 60, Desolvation Gas Flow (L/Hr) 700 [0178] Mass range: 140 to 800 Da [0179] DAD Wavelength range (nm): 210 to 400

[0180] Method Waters ACQUITY UPLC with the following HPLC gradient conditions [0181]

(Solvent A: Water/Methanol 9:1+0.1% formic acid and Solvent B: Acetonitrile+0.1% formic acid)

TABLE-US-00005 Time (minutes) A (%) B (%) Flow rate (ml/min) 0 100 0 0.75 2.5 0 100 0.75 2.8 0 100 0.75 3.0 100 0 0.75

[0182] Type of column: Waters ACQUITY UPLC HSS T3; Column length: 30 mm; Internal diameter of column: 2.1 mm; Particle Size: 1.8 micron; Temperature: 60° C.

[0183] Where necessary, enantiomerically pure final compounds may be obtained from racemic materials as appropriate via standard physical separation techniques, such as reverse phase chiral chromatography, or through stereoselective synthetic techniques, e.g., by using chiral starting materials.

[0184] A representative description of a chiral analysis apparatus is: [0185] Supercritical Fluid

Chromatography: [0186] Waters Acquity UPC.sup.2/QDa [0187] PDA Detector Waters Acquity

UPC.sup.2 [0188] Column: Daicel SFC CHIRALPAK® IC, 3 µm, 0.3 cm×10 cm, 40° C. [0189]

Mobile phase: A: CO.sub.2 B: iPr gradient: 05% B in 3.8 min [0190] ABPR: 1800 psi [0191] Flow

rate: 2.0 mL/min [0192] Detection: 247 nm [0193] Sample concentration: 1 mg/mL in

Acetonitril/iPrôpanol 50/50 [0194] Injection: 1 µL

FORMULATION EXAMPLES

TABLE-US-00006 Wettable powders a) b) c) active ingredients [components (A) and (B)] 25%

50% 75% sodium lignosulfonate 5% 5% — sodium lauryl sulfate 3% — 5% sodium

diisobutyl naphthalenesulfonate — 6% 10% phenol polyethylene glycol ether — 2% — (7-8 mol of ethylene oxide) highly dispersed silicic acid 5% 10% 10% Kaolin 62% 27% —

[0195] The active ingredient is thoroughly mixed with the adjuvants and the mixture is thoroughly ground in a suitable mill, affording wettable powders that can be diluted with water to give suspensions of the desired concentration.

TABLE-US-00007 Powders for dry seed treatment a) b) c) active ingredients [components (A) and

(B)] 25% 50% 75% light mineral oil 5% 5% 5% highly dispersed silicic acid 5% 5% —

Kaolin 65% 40% — Talcum — — 20%

[0196] The active ingredient is thoroughly mixed with the adjuvants and the mixture is thoroughly ground in a suitable mill, affording powders that can be used directly for seed treatment.

TABLE-US-00008 Emulsifiable concentrate active ingredients [components (A) and (B)] 10%

octylphenol polyethylene glycol ether 3% (4-5 mol of ethylene oxide) calcium

dodecylbenzenesulfonate 3% castor oil polyglycol ether (35 mol of ethylene oxide) 4%

Cyclohexanone 30% xylene mixture 50%

[0197] Emulsions of any required dilution, which can be used in plant protection, can be obtained from this concentrate by dilution with water.

TABLE-US-00009 Dusts a) b) c) active ingredients [components (A) and (B)] 5% 6% 4%

talcum 95% — — Kaolin — 94% — mineral filler — — 96%

[0198] Ready-for-use dusts are obtained by mixing the active ingredient with the carrier and grinding the mixture in a suitable mill. Such powders can also be used for dry dressings for seed.

TABLE-US-00010 Extruder granules active ingredients [components (A) and (B)] 15% sodium

lignosulfonate 2% carboxymethylcellulose 1% Kaolin 82%

[0199] The active ingredient is mixed and ground with the adjuvants, and the mixture is moistened with water. The mixture is extruded and then dried in a stream of air.

TABLE-US-00011 Coated granules active ingredients [components (A) and (B)] 8% polyethylene glycol (mol. wt. 200) 3% Kaolin 89%

[0200] The finely ground active ingredient is uniformly applied, in a mixer, to the kaolin moistened with polyethylene glycol. Non-dusty coated granules are obtained in this manner.

TABLE-US-00012 Suspension concentrate active ingredients [components (A) and (B)] 40% propylene glycol 10% nonylphenol polyethylene glycol ether (15 mol of ethylene oxide) 6% Sodium lignosulfonate 10% carboxymethylcellulose 1% silicone oil (in the form of a 75% emulsion in water) 1% Water 32%

[0201] The finely ground active ingredient is intimately mixed with the adjuvants, giving a suspension concentrate from which suspensions of any desired dilution can be obtained by dilution with water. Using such dilutions, living plants as well as plant propagation material can be treated and protected against infestation by microorganisms, by spraying, pouring or immersion.

TABLE-US-00013 Flowable concentrate for seed treatment active ingredients [components (A) and (B)] 40% propylene glycol 5% copolymer butanol PO/EO 2% tristyrenepheno with 10-20 moles EO 2% 1,2-benzisothiazolin-3-one 0.5% (in the form of a 20% solution in water) monoazo-pigment calcium salt 5% Silicone oil (in the form of a 75% emulsion in water) 0.2% Water 45.3%

[0202] The finely ground active ingredient is intimately mixed with the adjuvants, giving a suspension concentrate from which suspensions of any desired dilution can be obtained by dilution with water. Using such dilutions, living plants as well as plant propagation material can be treated and protected against infestation by microorganisms, by spraying, pouring or immersion.

Slow Release Capsule Suspension

[0203] 28 parts of a combination of the active ingredients [components (A) and (B)] is mixed with 2 parts of an aromatic solvent and 7 parts of toluene diisocyanate/polymethylene-polyphenylisocyanate-mixture (8:1). This mixture is emulsified in a mixture of 1.2 parts of polyvinylalcohol, 0.05 parts of a defoamer and 51.6 parts of water until the desired particle size is achieved. To this emulsion a mixture of 2.8 parts 1,6-diaminohexane in 5.3 parts of water is added. The mixture is agitated until the polymerization reaction is completed. The obtained capsule suspension is stabilized by adding 0.25 parts of a thickener and 3 parts of a dispersing agent. The capsule suspension formulation contains 28% of the active ingredients. The medium capsule diameter is 8-15 microns. The resulting formulation is applied to seeds as an aqueous suspension in an apparatus suitable for that purpose.

PREPARATION EXAMPLES

[0204] The compounds B.01 to B.15 of Table Ti were prepared using any of the synthetic techniques as described in WO 2017/055469, WO 2017/055473, WO 2017/093348, WO 2017/118689, WO 2017/220485, WO 2018/065414, WO 2018/158365, WO 2018/177894, WO 2018/177880, WO 2018/219773 and WO 2019/002151.

[0205] Where necessary, enantiomerically pure final compounds may be obtained from racemic materials as appropriate via standard physical separation techniques, such as reverse phase chiral chromatography, or through stereoselective synthetic techniques, (e.g., by using chiral starting materials).

TABLE-US-00014 TABLE T1 Melting point (mp) data and/or retention times (R.sub.t) for compounds B.01 to B.15: Compound RT [M + H] Entry name Structure (min) (measured) Method mp ° C. B.01 N-methoxy-N-[[4- [5-(trifluoromethyl)- 1,2,4-oxadiazol-3- yl]phenyl]methyl] cyclopropanecarboxamide [00026]

 0.99 316.3 A 38-41 B.02 N,2-dimethoxy-N-[[4-[5- (trifluoromethyl)- 1,2,4-oxadiazol-3- yl]phenyl]methyl] propanamide [00027]

 1.62 360.2 B 62-65 B.03 N-ethyl-2-methyl- N-[[4-[5- (trifluoromethyl)- 1,2,4-

oxadiazol-3-yl]phenyl]methyl] propanamide [00028]  1.78 342.3 B B.04 1-methoxy-3- methyl-1-[[4-[5- (trifluoromethyl)- 1,2,4-oxadiazol-3- yl]phenyl]methyl] urea [00029]  75-78 B.05 1,3-dimethoxy-1- [[4-[5- (trifluoromethyl)- 1,2,4-oxadiazol-3- yl]phenyl]methyl]urea [00030]  88-91 B.06 3-ethyl-1-methoxy- 1-[[4-[5- (trifluoromethyl)- 1,2,4-oxadiazol-3- yl]phenyl]methyl]urea [00031]  58-63 B.07 N-[[4-[5- (trifluoromethyl)-1,2,4- oxadiazol-3- yl]phenyl]methyl] propanamide [00032]  0.93 300.4 A 138-143 B.08 4,4-dimethyl-2-[[4- [5-(trifluoromethyl)- 1,2,4-oxadiazol-3- yl]phenyl]methyl] isoxazolidin-3-one [00033]  1.13 342 A 53-54 B.09 5,5-dimethyl-2-[[4- [5-(trifluoromethyl)- 1,2,4-oxadiazol-3- yl]phenyl]methyl] isoxazolidin-3-one [00034]  1.04 342 A B.10 ethyl 1-[[4-[5- (trifluoromethyl)- 1,2,4-oxadiazol-3- yl]phenyl]methyl] pyrazole-4-carboxylate [00035]  120-130 B.11 N,N-dimethyl-1-[[4- [5-(trifluoromethyl)- 1,2,4-oxadiazol-3- yl]phenyl]methyl]- 1,2,4-triazol-3-amine [00036]  0.97 339 A 106.2-110.5 B.12 (3-methylisoxazol- 5-yl)-[4-[5- (trifluoromethyl)- 1,2,4-oxadiazol-3- yl]phenyl]methanone [00037]  112-115 B.13 (5-methyl-2- pyridyl)-[4-[5- (trifluoromethyl)- 1,2,4-oxadiazol-3- yl]phenyl]methanone [00038]  1.15 [M + H] not detected A 1.37 B.14 2-oxo-N-propyl-2- [4-[5- (trifluoromethyl)- 1,2,4-oxadiazol-3- yl]phenyl]acetamide [00039]  78-83 B.15 ethyl 1-[[5-[5- (trifluoromethyl)- 1,2,4-oxadiazol-3- yl]-2- thienyl]methyl]pyrazole- 4-carboxylate [00040]  1.08 373.2 B 115.7-116.5

BIOLOGICAL EXAMPLES

Example A1: Preventative Control Against *Phakopsora pachyrhizi* (Asian Soybean Rust)/Soybean/Leaf Disc

[0206] Whole soybean plants are treated with the recited compositions of active ingredients four weeks after planting. One day after spraying application leaf discs are cut from the first trifoliate leaf. Five repetitions at each rate are conducted. The leaf discs are inoculated with *Phakopsora pachyrhizi* one day after treatment. Evaluation of the leaf discs is conducted 11 to 14 days after inoculation and the biological activity is derived from the relation of the treated compared to untreated, infested check plants. The rates of the active ingredients used are given in Tables 1 to 6 as g active ingredient (a.i.)/ha.

[0207] Results from the tests outlined above are shown below in Tables 1 to 6. These data show that synergistic fungicidal activity is observed for the combination of a component (A) and a component (B) against *Phakopsora pachyrhizi*. According to COLBY, in mathematical terms the synergism factor SF corresponds to O/E. In the agricultural practice an SF of ≥ 1.0 indicates significant improvement over the purely complementary addition of activities (expected activity), while an SF of ≤ 0.9 in the practical application routine signals a loss of activity compared to the expected activity.

TABLE-US-00015 TABLE 1 Fungicidal activity of a composition of [(1S,2S)- 1-methyl-2-(o-tolyl)propyl] (2S)-2-[(4-methoxy- 3-propanoyloxy-pyridine-2-carbonyl)amino]propanoate (A.03) and N-methoxy-N-[[4-[5-(trifluoromethyl)- 1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide (B.01) against *Phakopsora pachyrhizi* as described in Example A1 above. Expected (additive) Combined activity inhibition A.03 B.01 (Colby) (observed) (g/ha) (g/ha) (%) (%) 0.4 10 0.2 0 0.8 65 0.4 0.8 69 77 0.2 0.8 65 73

TABLE-US-00016 TABLE 2 Fungicidal activity of a composition of [(1S,2S)-1-methyl-2-(o-tolyl)propyl] (2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2- carbonyl)amino]propanoate (A.03) and N,2-dimethoxy-N-[[4-[5- (trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide (B.02) against *Phakopsora pachyrhizi* as described in Example A1 above. Expected (additive) Combined activity inhibition A.03 B.02 (Colby) (observed) (g/ha) (g/ha) (%) (%) 0.8 42 0.4 10 0.4 64 0.8 0.4 79 83 0.4 0.4 68 77

TABLE-US-00017 TABLE 3 Fungicidal activity of a composition of [(1S,2S)-1-methyl-2-(o-tolyl)propyl] (2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2- carbonyl)amino]propanoate (A.03)

and N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide (B.03) against *Phakopsora pachyrhizi* as described in Example A1 above. Expected (additive) Combined activity inhibition A.03 B.03 (Colby) (observed) (g/ha) (g/ha) (%) (%) 0.4 10 0.8 55 0.4 21 0.4 0.8 60 76 0.4 0.4 29 51

TABLE-US-00018 TABLE 4 Fungicidal activity of a composition of [(1S,2S)-1-methyl-2-(o-tolyl)propyl] (2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2- carbonyl)amino]propanoate (A.03) and 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea (B.04) against *Phakopsora pachyrhizi* as described in Example A1 above. Expected (additive) Combined activity inhibition A.03 B.04 (Colby) (observed) (g/ha) (g/ha) (%) (%) 0.8 42 0.4 10 0.4 44 0.8 0.4 68 83 0.4 0.4 50 64

TABLE-US-00019 TABLE 5 Fungicidal activity of a composition of [(1S,2S)-1-methyl-2-(o-tolyl)propyl] (2S)-2-[(4-methoxy-3-propanoyloxy-pyridine- 2-carbonyl)amino]propanoate (A.03) and 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea (B.06) against *Phakopsora pachyrhizi* as described in Example A1 above. Expected (additive) Combined activity inhibition A.03 B.06 (Colby) (observed) (g/ha) (g/ha) (%) (%) 0.8 42 0.4 10 0.4 22 0.8 0.4 55 60 0.4 0.4 30 54

TABLE-US-00020 TABLE 6 Fungicidal activity of a composition of [(1S,2S)-1-methyl-2-(o-tolyl)propyl] (2S)-2-[(4-methoxy-3-propanoyloxy-pyridine-2- carbonyl)amino]propanoate (A.03) and ethyl 1-[[4-[5-(trifluoromethyl)- 1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate (B.10) against *Phakopsora pachyrhizi* as described in Example A1 above. Expected (additive) Combined activity inhibition A.03 B.10 (Colby) (observed) (g/ha) (g/ha) (%) (%) 0.8 42 0.4 10 0.4 55 0.8 0.4 74 81 0.4 0.4 60 79

Claims

1. A fungicidal composition comprising a mixture of components (A) and (B) as active ingredients, wherein component (A) is a compound of formula (I): ##STR00041## wherein R.sup.1 is hydrogen, —C(=O)CH.sub.3 or —C(=O)CH.sub.2CH.sub.3, R.sup.2 is methyl or isopropyl, and R.sup.3 is hydrogen or fluorine, or a salt, enantiomer, tautomer or N-oxide thereof; and component (B) is a compound selected from the group consisting of N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, ethyl 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrazole-4-carboxylate, N,N-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-1,2,4-triazol-3-amine, (3-methylisoxazol-5-yl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, (5-methyl-2-pyridyl)-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methanone, 2-oxo-N-propyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide and ethyl 1-[[5-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-2-thienyl]methyl]pyrazole-4-carboxylate, or a salt, enantiomer, tautomer or N-oxide thereof.

2. A fungicidal composition according to claim 1, which further comprises an additional active ingredient component (C), which is different to component (B), and is selected from the group consisting of metyltetrapole, azoxystrobin, trifloxystrobin, picoxystrobin, pyraclostrobin, metominostrobin, prothioconazole, cyproconazole, difenoconazole, tebuconazole, propiconazole,

epoxiconazole, hexaconazole, metconazole, benzovindiflupyr, pydiflumetofen, fluxapyroxad, sedaxane, bixafen, isopyrazam, fluopyram, fluindapyr, isoflucypram, inpyrfluxam, quinofumelin, ipflufenquin, aminopyrifin, fluazinam, fludioxonil, fenpicoxamid, florylpicoxamid, fenpropidin, fenpropimorph, mancozeb, chlorothalonil, cyclobutrifluram, Cu-oxychloride, Cu-hydroxide, N'-[5-bromo-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-ethyl-N-methyl-formamidine, N'-[5-bromo-2-methyl-6-[(1S)-1-methyl-2-propoxy-ethoxy]-3-pyridyl]-N-ethyl-N-methyl-formamidine, N'-[5-bromo-2-methyl-6-[(1R)-1-methyl-2-propoxy-ethoxy]-3-pyridyl]-N-ethyl-N-methyl-formamidine, N'-[5-bromo-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-isopropyl-N-methyl-formamidine, N'-[5-chloro-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-ethyl-N-methyl-formamidine, N'-(2-chloro-5-methyl-4-phenoxyphenyl)-N-ethyl-N-methyl-formamidine, N'-[2-chloro-4-(2-fluorophenoxy)-5-methylphenyl]-N-ethyl-N-methyl-formamidine, N'-[2-chloro-4-(3-fluorophenoxy)-5-methylphenyl]-N-ethyl-N-methyl-formamidine, N'-(2-bromo-5-methyl-4-phenoxyphenyl)-N-ethyl-N-methyl-formamidine, N'-[2-bromo-4-(2-fluorophenoxy)-5-methylphenyl]-N-ethyl-N-methyl-formamidine, N'-[2-chloro-4-(2-fluorophenoxy)-5-methylphenyl]-N-isopropyl-N-methyl-formamidine, N'-[4-(2-cyanophenoxy)-2-methyl-5-(trifluoromethyl)phenyl]-N-ethyl-N-methyl-formamidine, N'-[5-bromo-2-methyl-4-(2-methylphenoxy)phenyl]-N-ethyl-N-methyl-formamidine, N-ethyl-N'-[4-(2-fluorophenoxy)-2-methyl-5-(trifluoromethyl)phenyl]-N-methyl-formamidine, (Z,E)-5-[1-(2,4-dichlorophenyl)pyrazol-3-yl]oxy-2-methoxyimino-N,3-dimethyl-pent-3-enamide, (Z,E)-5-[1-(4-chlorophenyl)pyrazol-3-yl]oxy-2-methoxyimino-N,3-dimethyl-pent-3-enamide, 4-[[6-[2-(2,4-difluorophenyl)-1,1-difluoro-2-hydroxy-3-(5-thioxo-4H-1,2,4-triazol-1-yl)propyl]-3-pyridyl]oxy]benzonitrile, methyl 3-[(4-chlorophenyl)methyl]-2-hydroxy-1-methyl-2-(1,2,4-triazol-1-ylmethyl)cyclopentanecarboxylate, 2-[6-(4-bromophenoxy)-2-(trifluoromethyl)-3-pyridyl]-1-(1,2,4-triazol-1-yl)propan-2-ol, 2-[6-(4-chlorophenoxy)-2-(trifluoromethyl)-3-pyridyl]-1-(1,2,4-triazol-1-yl)propan-2-ol, N-methyl-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzenecarbothioamide, N-methyl-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzamide, 2,2-difluoro-N-methyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide, N-[N-methoxy-C-methyl-carbonimidoyl]-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzamide, N-[(E)-methoxyiminomethyl]-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzamide, N-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]cyclopropanecarboxamide, N-(2-fluorophenyl)-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzamide, 1-(5,6-dimethyl-3-pyridyl)-4,4-difluoro-3,3-dimethyl-isoquinoline and 1-[6-(difluoromethyl)-5-methyl-3-pyridyl]-4,4-difluoro-3,3-dimethyl-isoquinoline.

3. A fungicidal composition according to claim 2, wherein component (C), which is different to component (B), is a compound selected from the group consisting of metyltetrapole, azoxystrobin, trifloxystrobin, picoxystrobin, metominostrobin, prothioconazole, cyproconazole, difenoconazole, tebuconazole, propiconazole, mefentrifluconazole, metconazole, benzovindiflupyr, pydiflumetofen, sedaxane, isopyrazam, inpyrfluxam, fluazinam, florylpicoxamid, fenpropidin, mancozeb, chlorothalonil, cyclobutrifluram, Cu-oxychloride, Cu-hydroxide, N'-[5-bromo-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-ethyl-N-methyl-formamidine, N'-[5-bromo-2-methyl-6-[(1S)-1-methyl-2-propoxy-ethoxy]-3-pyridyl]-N-ethyl-N-methyl-formamidine, N'-[5-bromo-2-methyl-6-[(1R)-1-methyl-2-propoxy-ethoxy]-3-pyridyl]-N-ethyl-N-methyl-formamidine and N'-[5-chloro-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-ethyl-N-methyl-formamidine.

4. A fungicidal composition according to claim 1, wherein the weight ratio of component (A) to component (B) is from 20:1 to 1:40.

5. A fungicidal composition according to claim 1, wherein the weight ratio of component (A) to component (B) is from 15:1 to 1:30.

6. A fungicidal composition according to claim 1, wherein the weight ratio of component (A) to component (B) is from 10:1 to 1:10.

7. A fungicidal composition according to claim 1, wherein the weight ratio of component (A) to

component (B) is from 5:1 to 1:5.

8. A fungicidal composition according to claim 1, wherein the weight ratio of component (A) to component (B) is from 4:1 to 1:4.

9. A fungicidal composition according to claim 1, wherein the weight ratio of component (A) to component (B) is from 3:1 to 1:3.

10. A fungicidal composition according to claim 1, wherein the weight ratio of component (A) to component (B) is from 2:1 to 1:2.

11. A fungicidal composition according to claim 1, wherein the weight ratio of component (A) to component (B) is 1:1.

12. A fungicidal composition according to claim 1, wherein the composition further comprises an agriculturally acceptable carrier and, optionally, a surfactant and/or formulation adjuvants.

13. A method of controlling or preventing phytopathogenic diseases, especially phytopathogenic fungi, on useful plants or on propagation material thereof, which comprises applying to the useful plants, the locus thereof or propagation material thereof a fungicidal composition as defined in claim 1.

14. A method according to claim 13, wherein component (A) is applied to the useful plants at a rate of from 5 to 2000 g a.i./ha, particularly from 10 to 1000 g a.i./ha, preferably from 5 to 200 g a.i./ha in association with a rate of from 1 to 5000 g a.i./ha, particularly from 2 to 2000 g a.i./ha, preferably from 5 to 200 g a.i./ha of component (B).

15. A method according to claim 13, wherein the composition components (A) and (B) are applied in a sequential manner.
