



US 20250283101A1

(19) **United States**

(12) Patent Application Publication
Ben Khaled et al.

(10) **Pub. No.: US 2025/0283101 A1**

(43) **Pub. Date:** **Sep. 11, 2025**

(54) **METHOD OF MODULATING THE
ALKALOID CONTENT OF TOBACCO**

Publication Classification

(71) Applicant: **Nicoventures Trading Limited**,
London (GB)

(72) Inventors: **Sara Ben Khaled**, London (GB);
Javier Galdon-Armero, London (GB);
Francisco Anastacio De Abreu E Lima, London (GB)

(51) **Int. Cl.**
C12N 15/82 (2006.01)
A24B 3/12 (2006.01)
A24B 13/00 (2006.01)
A24B 15/16 (2020.01)
A24B 15/20 (2006.01)
A24B 15/24 (2006.01)
C07K 14/415 (2006.01)

(52) **U.S. Cl.**
 CPC *C12N 15/8242* (2013.01); *A24B 3/12*
 (2013.01); *A24B 13/00* (2013.01); *A24B 15/16*
 (2013.01); *A24B 15/20* (2013.01); *A24B 15/24*
 (2013.01); *C07K 14/415* (2013.01); *C12N*
15/8205 (2013.01)

(73) Assignee: **Nicoventures Trading Limited**,
London (GB)

(21) Appl. No.: **18/859,891**

(22) PCT Filed: **Apr. 26, 2023**

(86) PCT No.: **PCT/GB2023/051109**

§ 371 (c)(1),
(2) Date:

(30) **Foreign Application Priority Data**

Apr. 27, 2022 (GB) 2206107.1


(57) **ABSTRACT**

The present invention provides a method for modulating the alkaloid content of a plant (e.g. a tobacco plant), the method comprising modifying said plant by modulating the activity or expression of a YIPF-related protein. The present invention also provides for the use of a YIPF-related protein for modulating the alkaloid content of a plant, as well as tobacco cells, plants, plant propagation materials, harvested leaves, processed tobaccos, or tobacco products obtainable in accordance with the invention.

Specification includes a Sequence Listing.

Gene: Nitab4.5_0005388g0090.2 Genomic sequence (SEQ ID No. 1)

[illegible]

Figure 1**Figure 2**

Gene: Nitab4.5_0005388g0090.2 Genomic sequence (SEQ ID No. 1):

```
GGAGTTGGGACCCATAACCCAAAGCCCAAATATGACAAGATCCATCACCCTTACCGAAAAAACCAAAACCAA
AAAAAAGGAGAAAAAGAAATTTAAATAACTAAGGAAACAAATAGAAAGATTTGCATATACAGGAAACCACTCC
AGATCTGTGTCCGAAACTGAGCTGAAAGAACAAATTTGCATATACAGTTTATACGAGAAGAAACGAAGATGATG
TCCAGCAATTACACCGCAATTGATAATCAGAATGTTTCCGGATCTGTTCTGTAATTCTTTTCTCATTTTTTTTG
GTTTGATTGAAATTTTATTCTGGCCGTCGTTTCAGATCTAATTGTCTAATTTCTTATATATTATAGGCAGTCGCAG
ATCCACCTGGCCAAAGTTGCCGTCAAATTCACCTGGTAATGTCTTCAGATTTTCAACAAAATCATAACAGTAGTATT
TTGTTGGATCATTTTGAATTGGTTTGTGAAAAATTTGGGTTTATTTTGTGATTGCAGATTTCGAATTTGCAGACATT
TCCACCTTCCAGTTCCAGGGGAAAAATCTCTGGTGCTTCTGGACCTCCACGTGACGCCGACGGTATCTCCTCCTA
AGTTATTCATATATAGTTACGACGTAAAGACGTAATAGTGTGTTGCTATTAACCTAAGCTGTGGTTTTTCAGCTAAA
TTCAGCTAATTTTCTGCAAAAAAAGGGAAGGAAGGGGGGGGGGGAAGCAGGGTTATTTAGAATTTAGACA
TACAAGAAGTGGAGCTAGTGCAGCTTTTTAGTTTGTCTTTAGATCTTTGAAGTTCATGATACAACACTTGTCTCT
ATAATAGCGATATAAACACACCGATGATAGTGCTTAACATTGCATTATGAAGGGGCTGTCCAGCATGTTTCAGCA
TTTTTTCTTTCAACTAAGCTGATGTTAACATCTTTCACTCTGCTAATTATGCTATGACTTCTTTCAAATTGAAA
TTTTTACTTCAAGGCTACAAAGCCTGGATTCAAATATACATTCTTCTTCAATGATTTTGAATTCAAATCTGATA
CTATATGAGGCAGTAGGATCATAAGAAAGAAATTTGTTTACCAGTACAAAATTTGGAAAGACGAATGGAATTTTGT
GTAAACGTATAAGTCAAAATGAGGTGAGGAAGAAACCATTTTGCCAGGGTAAAAATAAATACAAGGAAGTCCCAGG
ATCGGTTTCGAGGGATTACAATAATGTGGTTATGTTTAGTCTAGGGTTATTTAGAATTTAGACATACAAGAAGTG
GAGCTAGTGCAGCTTTTGTCTTTAGATCTTTGAAGTTCATGATACAACACTTGTCTCTATAATAGCGAT
ATAAACACACCGATGATAGTGCTTAACATTGCATTATGAAGGGGCTGTCCAGCATGTTTCAGCATTTTTTCTTTC
AACTAAGCTGATGTTAACATCTTTCACTGCTAATTATGCTATGACTTCTTTCAAATTGAAATTTTTACTTCAA
GGCTACAAAGCCTGGATTCAAATATACATTCTTCTTCAATGATTTTGAATTCAAATCTGATACTATATGAGGCA
GTAGGATCATAAGAAAGAAATTTGTTTACCAGTACAAAATTTGGAAAGACGAATGGAATTTTGTGTAAACGTATAA
GTCAAAATGAGGTGAGGAAGAAACCATTTTGCCAGGGTAAAAATAAATACAAGGAAGTCCCAGGATCGGTTTCGAG
GGATTACAATAATGTGGTTATGTTTAGTCTAGGTATACCTAGTTCCGTACAGGTTGTCTCATCCGGTGCTTCTTT
TTGATAAGGTAAGTTTATTGATGCGGACAACAAAGTGCTGCTTAGTTGTATATTCAAAAAAGTATGTGCATCAC
AAGTCATGCAATGAACGGAGGAAGTCCAAAATGGCTTCGGGCTCTAACACATTTTCCACAGCCATGTTGCACCAA
AACTAAGCAAAGATATATAATTTGTGTTTGTAGTTTGTATCACAGTATCTTCTTCTCTTAAAAAATTCACGTTT
```

CATCTGGTGCCTTCTCATAGAGTTATTTTGGTTATCTCTAAAAAGAGTTAACTCTCAAAGCTTAACTGAAGGAAGT
TGTGAGTCATGCTATTGCGTCCTTTTCATTAAGTTATTAGCAAATCTATTTTCCCTCTCCACTTTCCACAGTTC
AGGTTGAATTTACTTCTCTTTTAGATAATAATTATGACTTGTTGTTGTAGATACATTCTCTAAACCAGTATCTG
GTTCTGATGAGCCCCAGCAGAGCGGTTGGTTTCGCGCTTTCACATATTGCTGCGTACAAGCCCTATTTTGATGTTG
ATACATCAGATGTTCTGGAGAGAATAAAAAGATTCACCTCTTCCCTTTCACGGGATCCTTTTCGGAGAAGACCGCCA
ATAACCCAGATTTGTATGCTCCCTGCTCTTTTATTTCTATGTTTATTGAATTAGCATTTATTCTCGTGTTTTAAT
CCTAAGTATAACTCTTATAAAAACTTATTATGAGTAGTTATTTCCCTGGAAATATAGGCCTTTGGGCTTCTACCGT
TATTAATGGCATTATATAAATTCTTATCCCAATTCACCTCTATCAATTCCTGCGCTGTTTTTCCCGTTTACCA
CGGATGAATAAAGTTGGCAGTTTATAGGAGACCAAGAATATATCTTGCTCCCAGTAACACTTTCTTCTTCTTGAA
TGATATCGTTCTAATTTCTTATTATGGTTTTGGATGAGATGACCGTTAATTAATATCCCTGTCTATCAGTTTCAT
CCTACAATATTCTCATTGATCAGCTATTTAGTCCTCGACTAAGAGGCATTTACCTGCTAGGAGGTTGCCAATGGG
CAACCACTATGCTTAACTTGCTCGAATTGAATAATGATTTATTGTGGAACTTCCATGCATTAAATGCAGGCTGT
AATGGGGTTGTGTAAATTCAGGCTAATGGATATGGAACAATTCTTACATTTAACTAGATGGAGTTTGTTCAAAT
CTCTGGCGGTATATTTTGAAAATTACCATTAGGCATACAGAAGAACCAATAAGAATAATTTGGGCTGGTATCTTT
GTCGTTTTCGAGTTGGAATTTGTCTGCGCATTGTACTAAAGCTAGTGAGTTTGTGCAAGTCATCAGATTTCCC
TTCTAGAATGTATGTGCTTTGCTTTGATTTCTTCTTCTGCTCCCTATATCTATCTTCTATTATATAGGATTTGTTT
AGACATGCCGTTTTGCACTTCAGAACCTGTTATTTCTTAAGTTTCTGTTTCCAAATTTGTATTACTTTTTTGGGG
GGGGGGGGGGGGGGTGGGTGTGAATCATTCTTCTGTTTCATCTTCTTATGCATGTTCTGTGTGTACGTATGAC
CGTGATATAATAGCACATGTTTGTATTATGGCAAGCGGCAGAGCTCATAATGGTGATTATTGGCATATTTGTTTC
ATGAAAAGTTATTTTCAGTTGTTGGGCTTATGATACATTATGACTGCTTTTAAAGTCCTTGCTCCTTTTAACTTG
ATTGACACTTGGTTTGTCTTGTGGAGTTGTAAGTGCAGTGTGATATTTCAACTGTACCTTCATGGACAAAAAG
TAATAACCATCAAACAACCTACTAAAAGAATTTAAAAAAGAAAAGAACAGTGATTAGCAAGTGAAGTTTGGCAC
TGCTGCCCTCTGTTCTGCTTTAGCCAGCTTTACATGCTCTCTGAGTAAGAAGACGCTTGTTTCAGAACTTGCCCT
GCAGGGAAGTTTGTCTAAAGGGTGCAGTGCAGGAACAGAAATGATGTCCGCAAACCTTTGCATCTAACTTAACT
ATTGCTAGATGCTTATGCAGCCATTATCTGTACTGGGTGGTGTCTAGACTTCTTTGCCCTTAAGATCTTGTGAG
TGTATGTTTCAATTATTTCCATATTAGCGAATCTCTACCGTAATTACTCAATTTTTTCTTTTCAGGTATGGCC
CGTTCTGGATATGCAGTACCTTGGTATTTGTTGCAGCTTCCATTGGCACGTTTGTACCTATATAGCACACAAGC
TACAGAAAAGGAGTGGAACATATGACATCAACCTTGTCACTTGGTCTGCGGGTTTATTCTATGGTTATTCCACTG
TAGTTCTCTTTGCTTGTCTTAGTCCTCAAGTACTTCTCTGCCCATCTGCGCTCGTTTCAGCTGCTGTGCCTTT
ACGGCTATTCCCTTTTGTCTTCTTCCGCTTGGTCAAGTTTATTTAATTACAGTATACTTTGTTGAACCTTTG
GGAAACCTGTTGCTTGTATGAATCCTTTTCAATTTTAAAGTTTGGCATTAGGATCCATACTAAGATCTATGATTAAC
GCTTACAGTGTCTGTCTCTTGTTCCTGGGAGATAGTGAGATGGGTGGTTGCTGGTGTAGCTGGCTTAATGTCTG
CCACATTTGTCGCTCTTAATCTGAAACACCACATAGTATCGGCTGGTGAAAGGTGGTTCTTCAATTGTGGCTGCTA
TCTTCTGTTGCAACTTGCACCTTGTCTTGGTACTGAGGGTTTACTTGTTCACGTCACAGTATAACTACAAAATT
ATTGGAATGCGACTAGGCTCGGTTGGAACATGTCTATATAAATAACAGAGAAAGGACACTGTTCTTGGTATCAACT
CGACTGTAAGTGTGCTTGGTCATTGAAACGGAAGTGTATTCATTCTTTTACGATATACAATAATATAGTTTC
ATCAAAAGGTTCTCAGCTTGTGGTCTTCTTATTGAGAATGTCACACTTTTGAGCTTTAATAGTGGTCTGGGAGTT
CGAGGGTCGATGATTTAGTCAAACATTTGTGTTTAGTGGGGATTTAATTGTAATGTCAAGACTGATTAGCAAATT
CCTACAAGTCCACTTACAAGATAATCAAACCATGCGGAAATACTTTTGCTTGCCATGATGACCTTGAATGACCT
AATTTTCAACATGGTATCTACCTATTCTTACTTTGCTTGTCTCCATTGACGGTCA

Figure 3

Nitab4.5_0005388g0090.2 coding sequence (SEQ ID No: 2):

ATGATGTCCAGCAATTACACCGCAATTGATAATCAGAATGTTTCCGGATCTGTTCTCTGCAGTCGCAGATCCACCT
GGCCAAGTTGCCGTCAAATTCAGTGATTCGAATTTGCAGACATTTCCACCTTCCAGTTCCCAGGGGAAAATCTCT
GGTGCTTCTGGACCTCCACGTGACGCCGACGATAACATTCTCTAAACCAGTATCTGGTTCTGATGAGCCCCAGCAG
AGCGGTTGGTTTCGCGCTTTCACATATTGCTGCGTACAAGCCCTATTTTGATGTTGATAACATCAGATGTTCTGGAG
AGAATAAAAAGATTCACTCTTTCTTTTACGGGATCCTTTTCGGAGAAGACCGCCAATAACCCAGATTTGTATGGC
CCGTTCTGGATATGCAGTACCTTGGTATTTGTTGCAGCTTCCATTGGCACGTTTGTACCTATATAGCACACAAG
CTACAGAAAAGGAGTGGAACATATGACATCAACCTTGTCACTTGGTCTGCGGGTTTATTCTATGGTTATTCCACT
GTAGTTCTCTTTGCTTGTCTTAGTCCTCAAGTACTTCTCTGCCCATCTGCGCTCGTTTCAGCTGCTGTGCCTT
TACGGCTATTCCCTTTTTGTCTTCAATTCGGGCATTGTGTCTGTCTCTTGTTCCTGGGAGATAGTGAGATGGGTG
GTTGCTGGTGTAGCTGGCTTAATGTCTGCCACATTTGTGCTCTTAACTCTGAAACACCACATAGTATCGGCTGGT
GAAAGGTGGTCTTCAATTGTGGCTGCTATCTTCTGTTGCAACTTGCACCTTGGTACTGAGGGTTTACTTG
TTCAACGTCACAGTATAA

Figure 4

Nitab4.5_0005388g0090.2 Amino acid sequence: (SEQ ID No: 3)

MMSSNYTAIDNQNVSGSVPAVADPPGQVAVKFTDSNLQTFPPSSSQGKISGASGPPRDADDTFSKPVSGSDEPQQ
SGWFRRAFTIAAYKPYFDVDTSDVLERIKDSLFPFTGSFSEKTANNPDLYGPFWICSTLVFVAASIGTEFVTYIAHK
LQKKEWNYDINLVTWSAGLFYGYSTVVPLCLFLVLKYFSAPSGLVQLLCLYGYSLFVFIPALCLSLVPWEIVRWV
VAGVAGLMSATFVALNLKHHIVSAGERWFFIVAAIFLLQLALALVLRVYLFNVTV

Figure 5

Identifier	Nic 1 ERF - Gene
Nitab4.5_0003090g0020.1	<i>ERF17L3ΔN</i>
Nitab4.5_0003090g0030.1	<i>ERF199</i>
Nitab4.5_0003665g0040.1	<i>JRE5L2</i>
Nitab4.5_0004620g0010.1	<i>ERF210</i>
Nitab4.5_0004620g0030.1	<i>ERF91</i>
Nitab4.5_0004620g0080.1	<i>ERF29</i>
Nitab4.5_0004620g0090.3	<i>ERF130</i>
Nitab4.5_0004620g0095.1	<i>ERF16</i>
Nitab4.5_0006382g0040.1	<i>ERF110</i>

Figure 6

Identifier	Nic 2 ERF - Gene
Nitab4.5_0002924g0010.1	ERF17LI
Nitab4.5_0002924g0020.2	ERF179
Nitab4.5_0002924g0040.2	ERF17
Nitab4.5_0002924g0045.1	ERF168
Nitab4.5_0002924g0050.2	ERF115
Nitab4.5_0006499g0010.1	ERF104
Nitab4.5_0006499g0020.2	ERF221
Nitab4.5_0012667g0020.2	ERF91L1
Nitab4.5_0015055g0010.2	ERF189

Figure 7

Nitab4.5_0003090g0030.1 (ERF199)

SEQ ID No. 7. >Nitab4.5_0003090g0030.1 720 residues [genomic]
 ATGGCAATGGAAATGAATCCAGCTGACGAAACCTTGTTTTCTCCGACTCTCATCTCCTTGAATCGATAAAGCAA
 CATCTTCTTGACGATTCAGATTTTTCTGAAATTTTTTCGTCGATGAATTCTAGCAACGAAATATTGCCTAACAGT
 CCTAGCTCAAGTTTTAGCAGCTTCGACTTCGACTGCAGCTTCCTTAATTGGGATGAAAACCTCTGAGGAAACATTA
 ATACCAACTGATCAGAATCCTTCACATGAATCCCATGAAAAGTACTCCGAGTCCGAGGAGAAAACCCAGGGCCCT
 GGGGTGGCGCGTGAGAAAAACGCGCCGCGAGATTGGACGCGGTACATAGGAGTGAAACGGCGACCGTGGGGGACG
 TTTTCGGCGGAGACAAGAGACCCAAGTAGGAAAGGTGAAGGTGCAAGGCTGTGGTTAGGAACTTACGAGACCGCA
 GAGGATGCAGCGTTAGCTTACGATCAAGCCGCTTTCAAATCCGCGGCTCGAGAGCTCGGCTCAATTTTCCTCAC
 TTAATCGGCTCAAACATGCCTAAGCCGGCTAGAGTTACAGCGAGGCGTAGTCGTACGCGCTCACCCGAGCCATCG
 TCTTCTTCATCCACCTCATCATCAGAAAATGTACCAAGGAAAAGGAATATAGATGTGATAAATTCATAGCCAAA
 GCCAAATTCCTTTGTCATAGCTTGAATTTACAGAGATTAGCTTAA

Figure 8

SEQ ID No. 8. >Nitab4.5_0003090g0030.1 720 residues [cdna]
 ATGGCAATGGAAATGAATCCAGCTGACGAAACCTTGTTTTCTCCGACTCTCATCTCCTTGAATCGATAAAGCAA
 CATCTTCTTGACGATTCAGATTTTTCTGAAATTTTTTCGTCGATGAATTCTAGCAACGAAATATTGCCTAACAGT
 CCTAGCTCAAGTTTTAGCAGCTTCGACTTCGACTGCAGCTTCCTTAATTGGGATGAAAACCTCTGAGGAAACATTA
 ATACCAACTGATCAGAATCCTTCACATGAATCCCATGAAAAGTACTCCGAGTCCGAGGAGAAAACCCAGGGCCCT
 GGGGTGGCGCGTGAGAAAAACGCGCCGCGAGATTGGACGCGGTACATAGGAGTGAAACGGCGACCGTGGGGGACG
 TTTTCGGCGGAGACAAGAGACCCAAGTAGGAAAGGTGAAGGTGCAAGGCTGTGGTTAGGAACTTACGAGACCGCA
 GAGGATGCAGCGTTAGCTTACGATCAAGCCGCTTTCAAATCCGCGGCTCGAGAGCTCGGCTCAATTTTCCTCAC
 TTAATCGGCTCAAACATGCCTAAGCCGGCTAGAGTTACAGCGAGGCGTAGTCGTACGCGCTCACCCGAGCCATCG
 TCTTCTTCATCCACCTCATCATCAGAAAATGTACCAAGGAAAAGGAATATAGATGTGATAAATTCATAGCCAAA
 GCCAAATTCCTTTGTCATAGCTTGAATTTACAGAGATTAGCTTAA

Figure 9

SEQ ID No. 9. >Nitab4.5_0003090g0030.1 720 residues [cds]
 ATGGCAATGGAAATGAATCCAGCTGACGAAACCTTGTTTTCTCCGACTCTCATCTCCTTGAATCGATAAAGCAA
 CATCTTCTTGACGATTCAGATTTTTCTGAAATTTTTTCGTCGATGAATTCTAGCAACGAAATATTGCCTAACAGT
 CCTAGCTCAAGTTTTAGCAGCTTCGACTTCGACTGCAGCTTCCTTAATTGGGATGAAAACCTCTGAGGAAACATTA
 ATACCAACTGATCAGAATCCTTCACATGAATCCCATGAAAAGTACTCCGAGTCCGAGGAGAAAACCCAGGGCCCT
 GGGGTGGCGCGTGAGAAAAACGCGCCGCGAGATTGGACGCGGTACATAGGAGTGAAACGGCGACCGTGGGGGACG
 TTTTCGGCGGAGACAAGAGACCCAAGTAGGAAAGGTGAAGGTGCAAGGCTGTGGTTAGGAACTTACGAGACCGCA
 GAGGATGCAGCGTTAGCTTACGATCAAGCCGCTTTCAAATCCGCGGCTCGAGAGCTCGGCTCAATTTTCCTCAC
 TTAATCGGCTCAAACATGCCTAAGCCGGCTAGAGTTACAGCGAGGCGTAGTCGTACGCGCTCACCCGAGCCATCG
 TCTTCTTCATCCACCTCATCATCAGAAAATGTACCAAGGAAAAGGAATATAGATGTGATAAATTCATAGCCAAA
 GCCAAATTCCTTTGTCATAGCTTGAATTTACAGAGATTAGCTTAA

Figure 10

SEQ ID No. 10. >Nitab4.5_0003090g0030.1 239 residues [peptide]
 MAMEMNPADETLFFSDSHLLESIKQHLLDDSDSEIFSSMNSSNEILPNSPSSSFSSFDSDCSFLNWDENSEETL
 IPTDQNPESHESHEKYSEEEKTQGGPVAREKNAPRDWTRYIGVKRRPWGTFSAETRDPSRKGEARLWLGTYETA
 EDAALAYDQAFAFKIRGSRARLNFPHLIGSNMPKPARVTARRSRTRSPEPSSSSSTSSSENVPRKRNIDVINSIAK
 AKFLCHSLNLQRLA

Figure 11

Nitab4.5_0015055g0010.2 (ERF189)

SEQ ID No. 11. >Nitab4.5_0015055g0010.2 699 residues [genomic]
 ATGGAAATGAATCTAGCTGACGAAACCTTGTTTTCTCTGAGTCTCATCTCCTTGAATCGATAAAGCAACATCTT
 CTTGATGATTTCAGATTTTTCTGAAATTTTTTCGCCGATGAGTTCAAGCAACGAAATATTGCCTAACAGTCTTAGC
 TCAAGTTTTAGCAGCTTCGACTGCAGCTTCCTCAATTGGGATGAAAACCTTTGAGGAAACATTAATACCAACTGAT
 CAAAATCCTTCACATGAGAAGTGCTCCGAGTCCGAGGAGCAAACCCAGGGCCCAGCGGTGGTGCGTGAGAAAAAC
 GCGCCGCGAGATTGGACGCGGTATATAGGAGTGAAACGGCGGCCGTGGGGGACGTTTTTCGGCGGAGACAAGAGAC
 CCAAGTAGGAAAGGTGAAGGTGCAAGGCTGTGGTTAGGAACTTACGAGACCGCAGAGGATGCAGCGTTGGCTTAC
 GATCAAGCCGCTTTCAAATCCGCGGCTCGAGAGCTCGGCTCAATTTTCCCACTTAATTGGCTCAAACATGCCT
 AAGCCGGCTAGAGTAACAGCGAGGCGTAGTCGTACGCGCTCACCCGAGCCATCCTCTTCTTCATCCACTTCATCA
 TCAGAAAATGTGCCAAGAAAAAGGAATATAGATGTGATAAATTCATAGCCAAAGCCAAATTCCTTTGTCATAGC
 T'

Figure 12

SEQ ID No. 12. >Nitab4.5_0015055g0010.2 699 residues [cdna]
 ATGGAAATGAATCTAGCTGACGAAACCTTGTTTTCTCTGAGTCTCATCTCCTTGAATCGATAAAGCAACATCTT
 CTTGATGATTCAGATTTTTCTGAAATTTTTTCGCCGATGAGTTCAAGCAACGAAATATTGCCTAACAGTCCTAGC
 TCAAGTTTTAGCAGCTTCGACTGCAGCTTCCTCAATTGGGATGAAAACCTTGAGGAAACATTAATACCAACTGAT
 CAAAATCCTTCACATGAGAAGTGCTCCGAGTCCGAGGAGCAAACCCAGGGCCCAGCGGTGGTGCGTGAGAAAAAC
 GCGCCGCGAGATTGGACGCGGTATATAGGAGTGAAACGGCGGCCGTGGGGGACGTTTTTCGGCGGAGACAAGAGAC
 CCAAGTAGGAAAGGTGAAGGTGCAAGGCTGTGGTTAGGAACCTTACGAGACCGCAGAGGATGCAGCGTTGGCTTAC
 GATCAAGCCGCTTTCAAATCCGCGGCTCGAGAGCTCGGCTCAATTTTCCCCACTTAATTGGCTCAAACATGCCT
 AAGCCGGCTAGAGTAACAGCGAGGCGTAGTCGTACGCGCTCACCCGAGCCATCCTCTTCTTCATCCACTTCATCA
 TCAGAAAATGTGCCAAGAAAAAGGAATATAGATGTGATAAATTCATAGCCAAAGCCAAATTCCTTTGTCATAGC
 TTAAATTTACAGAGATTAGCTTAA

Figure 13

SEQ ID No. 13. >Nitab4.5_0015055g0010.2 699 residues [cds]
 ATGGAAATGAATCTAGCTGACGAAACCTTGTTTTCTCTGAGTCTCATCTCCTTGAATCGATAAAGCAACATCTT
 CTTGATGATTCAGATTTTTCTGAAATTTTTTCGCCGATGAGTTCAAGCAACGAAATATTGCCTAACAGTCCTAGC
 TCAAGTTTTAGCAGCTTCGACTGCAGCTTCCTCAATTGGGATGAAAACCTTGAGGAAACATTAATACCAACTGAT
 CAAAATCCTTCACATGAGAAGTGCTCCGAGTCCGAGGAGCAAACCCAGGGCCCAGCGGTGGTGCGTGAGAAAAAC
 GCGCCGCGAGATTGGACGCGGTATATAGGAGTGAAACGGCGGCCGTGGGGGACGTTTTTCGGCGGAGACAAGAGAC
 CCAAGTAGGAAAGGTGAAGGTGCAAGGCTGTGGTTAGGAACCTTACGAGACCGCAGAGGATGCAGCGTTGGCTTAC
 GATCAAGCCGCTTTCAAATCCGCGGCTCGAGAGCTCGGCTCAATTTTCCCCACTTAATTGGCTCAAACATGCCT
 AAGCCGGCTAGAGTAACAGCGAGGCGTAGTCGTACGCGCTCACCCGAGCCATCCTCTTCTTCATCCACTTCATCA
 TCAGAAAATGTGCCAAGAAAAAGGAATATAGATGTGATAAATTCATAGCCAAAGCCAAATTCCTTTGTCATAGC
 TTAAATTTACAGAGATTAGCTTAA

Figure 14

SEQ ID No. 14.
 >Nitab4.5_0015055g0010.2 232 residues [peptide]
 MEMNLADETLFFSESHLLESIKQHLLDDSDSEIFSPMSSSNEILNPSPPSSSFSSFDCSFLNWDENFEETLIPTD
 QNPSHEKCSSEEEQTQGPVAVREKNAPRDWTRYIGVKRPPWGTFSAE TRDPSRKGE GARLWLGT YETAEDAALAY
 DQAAFKIRGSRARLNFPHLIGSNMPKPARVTARRSRTRSPEPSSSSSTSSSENVPKRKNIDVINSIAKAKFLCHS
 LNLQRL

Figure 15

TRV1: SEQ ID No. 15
 GATGGTAAGAAGAAAGGCGCGCAGTATGCGATAGCTCTTCACAGCCTGTATGACTTCAAGTTGAAAGACTTGATG
 GCTACTATGGTTGAGAAGAAAACCTAAAGTGGTTTCATGCTGCTATGCTTTTGCTCCTGAAAGTATGTTAGTGGAC
 GAAGGTCCATTACCTTCTGTTGACGGTTACTACATGAAGAAGAACGGGAAGATCTATTTCCGGTTTTGAGAAAGAT
 CCTTCCTTTTCTTACATTTCATGACTGGGAAGAGTACAAGAAGTATCTACTGGGGAAGCCAGTGAGTTACCAAGGG
 AATGTGTCTACTTCGAACCGTGGCAGGTGAGAGGAGACACAATGCTTTTTTCGATCTACAGGATAGCTGGAGTT
 CCGAGGAGGTCTCTATCATCGCAAGAGTACTACCGAAGAATATATATCAGTAGATGGGAAAGCATGGTTGTTGTC
 CCAATTTTTCGATCTGGTTCGAATCAACGCGAGAGTTGGTCAAGAAAGACCTGTTTGTAGAGAAACAATTCATGGAC
 AAGTGTGTTGGATTACATAGCTAGGTTATCTGACCAGAGCTGACCATAAGCAATGTTAAATCATACTTGAGTTCA
 AATAATTGGGTCTTATTCATAAACGGGGCGGCCGTGAAGAACAAGCAAAGTGTAGATTCTCGAGATTTACAGTTG
 TTGGCTCAAACCTTTGCTAGTGAAGGAACAAGTGGCGAGACCTGTCTGAGGGAGTTGCGTGAAGCAATTCGACT
 GAGACGAAACCTATCACGTCATTGACTGATGTGCTGGGTTTAAATATCAAGAAAACCTGTGGAAGCAGTTTGCTAAC
 AAGATCGCAGTCGGCGGATTTCGTTGGCATGGTTGGTACTCTAATTGGATTCTATCCAAAGAAGGTACTAACCTGG
 GCGAAGGACACACCAAATGGTCCAGAACTATGTTACGAGAAGCTCGCACAAAACCAAGGTGATAGTATTTCTGAGT
 GTTGTGTATGCCATTGGAGGAATCACGCTTATGCGTCGAGACATCCGAGATGGACTGGTGAAAAAAGTATGTGAT
 ATGTTTGATATCAAACGGGGGGGCCATGTCTTAGACGTTGAGAATCCGTGCCGCTATTATGAAATCAACGATTTT
 TTTAGCAGTCTGTATTCGGCATCTGAGTCCGGTGAGACCGTTTTTACCAGATTTATCCGAGGTAAGCAAGTCT
 GATAAGCTATTGCAGCAGAAGAAAGAAATCGCTGACGAGTTTCTAAGTGCAAAATTCCTCTAACTATTCTGGCAGT
 TCGGTGAGAACTTCTCCACCATCGGTGGTTCGTTTCATCTCGAAGCGGACTGGGTCTGTTGTTGGAAGACAGTAAC
 GTGCTGACCCAAGCTAGAGTTGGAGTTTCAAGAAAGGTAGACGATGAGGAGATCATGGAGCAGTTTCTGAGTGGT
 CTTATTGACACTGAAGCAGAAATTGACGAGGTTGTTTTCAGCTGAAATGTGAAAGAGGGGAAACAAGC
 GTACAAAAGGTGTTGTGTAACCTTTAACGCCACAGGATTTGAGAACGTGTTGCCAGCTTGCAAAACCTTTGGTC
 AGCAAAGGAAAAACGGTCAACCGTTCGATTACTTCAAGTGATGGGAGGTGAGAGATTACCAAAAAGCCGGTT
 GTCAGTGGAGACGATTCTGTGGACGCTAGAAGAGAGTTTCTGTACTACTTAGATGCGGAGAGAGTTCGCTCAAAT

GATGAAATTATGTCTCTGTATCGTGACTATTCGAGAGGAGTTATTCGAACTGGAGGTCAGAATTACCCGCACGGA
CTGGGAGTGTGGGATGTGGAGATGAAGAACTGGTGCATACGTCCAGTGGTCACTGAACATGCTTATGTGTTCCAA
CCAGACAAACGTATGGATGATTGGTCGGGATACTTAGAAGTGGCTGTTTGGGAACGAGGTATGTTGGTCAACGAC
TTCGCGGTGCGAAAGGATGAGTGATTATGTCATAGTTTGGCATCAGACGTATCTTTGCAATAACAGGTAATAATCC
TCTCTCTTGATATTTTTAAATTATAGAATTAATTAGTTTACTTTATCTTTACTATATGATTTAAATAGTTTAAAT
CTTGTTTTTGGAGTAACTATTCGATTTTGATATTTGTATTCGTCTTACAAAGTTGGAAATACATGATGATATTTTC
TTTTGAACGTGATACCTACCAATACTAATCTTACGGAATCTTTTAAATAGAGCACTAATCAACATGGAACTAAAGA
CCAATTCCTTAAGTGTCTCTGTGTACAGTTTCATTTTAGTAGTGGCTTAAAGTATTATTATCTCCCTTCATGCGGG
GCAATTATGTAGATTAAAATCGAAATTATATAAAATTTACATAAGTCTAAGTCTAGGGTCTCCAGCTAATTGTTA
TTTTTTTAAACGATGTTGACTAAAGCAATAACGACGTTGACTTGTGTAAACAGGTTGATCTTGGACAATTTAAGT
GCCCTGGATCTAGGACCAGTTAACTGTTCTTTTGAATTAGTTGACGGTGTACCTGGTTGTGGTAAGTCGACAATG
ATTGTCAACTCAGCTAATCCTTGTGTCGATGTGGTTCTCTCTACTGGGAGAGCAGCAACCGACGACTTGATCGAG
AGATTGCGCAGCAAAGGTTTTCCATGCAAATTGAAAAGGAGAGTGAAGACGGTTGATTCTTTTTTGTATGCAATTGT
GTCGATGGTTCTTTAACCGGAGACGTGTTGCATTTTCGACGAAGCTCTCATGGCCCATGCTGGTATGGTGTACTTT
TGCGCTCAGATAGCTGGTGCTAAACGATGTATCTGTCAAGGAGATCAGAATCAAATTTCTTTCAAGCCTAGGGTA
TCTCAAGTTGATTTGAGGTTTTCTAGTCTGGTCGGAAAGTTTGACATTGTTACAGAAAAAGAGAACTTACAGA
AGTCCAGCAGATGTGGCTGCCGATTTGAACAAGTACTTACTGAGATGTGAGAACACATAACCGGACTGCTAAT
TCGATGACGGTGAGGAAGATTGTGTCTAAAGAACAGGTTTCTTTGAAGCCTGGTGCTCAGTACATAACTTTCCCTT
CAGTCTGAGAAGAAGGAGTTGGTAAATTTGTTGGCATTGAGGAAAGTGGCAGCTAAAGTGAGTACAGTACACGAG
TCGCAAGGAGAGACATTCAAAGATGTAGTCTTAGTCAGGACGAAACCTACGGATGACTCAATCGCTAGAGGTCGG
GAGTACTTAATCGTGGCATTGTGCGCTCACACACAATCACTTGTGTATGAACTGTGAAAGAGGACGATGTAAGC
AAAGAGATCAGGGAAAGTGCCGCGCTTACGAAGGCGGCTTTGGCAAGATTTTTTGTACTGAGACCGTCTTATGA
CGGTTTTCGGTCTAGGTTTGTATGCTTTAGACATCATGAAGGGCTTGCCTGCTCCAGATTGAGTACGATTACG
GACTTGGAGATGTGGTACGACGCTTGTGTTCCGGGAAATTCGTTAAGAGACTCAAGCCTAGACGGGTATTTGGTG
GCAACGACTGATTGCAATTTGCGATTAGACAATGTTACGATCAAAAGTGGAACTGGAAAGACAAGTTTGCTGAA
AAAGAAACGTTTTCTGAAACCGGTTATTCGTACTGCTATGCCGACAAAAGGAAGACTACTCAGTTGGAGAGTTTG
TTAGCATTGACAGAAAAGGAACCAAGCGGCACCCGATCTACAAGAAAATGTGCACGCGACAGTTCTAATCGAAGAG
ACGATGAAGAAGCTGAAATCTGTTGTCTACGATGTGGGAAAAATTCGGGCTGATCCTATTGTCAATAGAGCTCAA
ATGGAGAGATGGTGGAGAAAATCAAAGCACAGCGGTACAGGCTAAGGTAGTAGCAGATGTGAGAGAGTTACATGAA
ATAGACTATTGCTCTTACATGTATATGATCAAATCTGACGTGAAACCTAAGACTGATTTAACACCGCAATTTGAA
TACTCAGCTCTACAGACTGTTGTGTATCACGAGAAGTTGATCAACTCGTTGTTGGTCCAATTTTCAAAGAAATTT
AATGAACGCAAGTTGGATGCTATGCAACCACATTTTGTGTTCAACACGAGAATGACATCGAGTGATTTAAACGAT
CGAGTGAAGTTCTTAAATACGGAAGCGGCTTACGACTTTGTTGAGATAGACATGTCTAAATTCGACAAGTCGGCA
AATCGCTTCCATTTACAACCTGCAGCTGGAGATTTACAGGTTATTTGGGCTGGATGAGTGGGCGGCCCTTCCTTTGG
GAGGTGTGCGCACTCAAACCTACTGTGAGAGATATTCAAATGGTATGATGGCGCATATTTGGTACCAACAAAAG
AGTGGAGATGCTGATACTTATAATGCAAATTCAGATAGAACACTGTGTGCACTCTTGTCTGAATTACCATTGGAG
AAAGCAGTCATGGTTACATATGGAGGAGATGACTCACTGATTGCGTTTCTTAGAGGAACGCAGTTTGTGATCCG
TGTCCAAAGTTGGCTACTAAGTGAATTTTCGAGTGCAAGATTTTAAAGTACGATGTCCCAATGTTTTGTGGGAAG
TTCTTGTCTTAAGACGTCATCGTGTACGAGTTCTGTCGGATCCGGTAAAGTTCTGACGAAGTTGGGGAAAAAG
AGTATAAAGGATGTGCAACATTTAGCCGAGATCTACATCTCGCTGAATGATTTCAATAGAGCTCTGGGACTAC
ATGGTGGTATCCAACTGTCCGAGTCTGTTTCAGACCGGTATTTGTACAAAGGTGATTCTGTTTCATGCGCTTTGT
GCGCTATGGAAGCATATTAAGAGTTTTACAGCTCTGTGTACATTATTCGAGACGAAAACGATAAGGAATGAAC
CCGGCTAAGGTTGATTGGAAGAAGGCACAGAGAGCTGTGTCAAACCTTTTACGACTGGTAATATGGAAGACAAGTC
ATTGGTCACCTTGAAGAAGAAGACTTTGAAAGTCTCAAATTTCTCAAATCTAGGGGCCATTGAATTGTTGTGGA
CGGTAGGAGGAAGAGACCGAAGTATTTTACAGAGAAGAGAACTGTCTAAATCATGTTGGTGGGAAGAAGAG
TGAACACAAGTTAGACGTTTTTGGACAAAGGGATTACAAAATGATTAAATCTTACGCGTTTTCTAAAGGTAGTAGG
TGTACAACTAGTTGTAACATCACATCTACCTGCAGATACGCTGGGTTCAATCAAATCGATCTGTTGGATTTCGAG
ACTTACTGAGAAAAGAAAGAGAGGAAAGACTATTTCAGAGATTCAAAGCTCGAGCTTGCGATAACTGTTTCAGTTGC
GCAGTACAAGGTTGAATACAGTATTTCCACACAGGAGAACCTACTTGTATGTCTGGAAGGTGGGTTGATTTCTGA
GGGCGTTCGGGTCTGTGACGGTACATACCTTTTCAGTATCGAAGTGTGCTGCTAATATGGGTTGCTACTGATTCGAC
TAGGCGCCTCAATGTGGAAGAAGCTGAACAGTTTCGGATTACATTGAAGGCGATTTTACCGATCAAGAGGTTTTCGG
TGAGTTTCATGTCTTTGAAACAAGTGGAGATGAAGACGATTGAGGCGAAGTACGATGGTCTTACAGACCAGCTAC
TACTAGACCTAAGTCATTATTGTCAAGTGAAGATGTTAAGAGAGCGTCTAATAAGAAAACTCGTCTTAATGCAT
AAAGAAATTTATTGTCAATATGACGTGTGACTCAAGGGTTGTGTGAATGAAGTCACTGTTCTTGGTCACGAGAC
GTGTAGTATCGGTTCATGCTAACAAATTCGGAAGCAAGTTGCTGACATGGTGGTGTACACGTAGGTGTGCGGA
AAATAATTGTGGATGGTTTTGTGTGTGTTATCAATGATTTTACTTTTGATGTGTATAATTGTTGTGCGCGTAG
TCACCTTGAAGAGTGTGCTAAACGTTGTTGAAACAAGAAATTCGAGAAATTTGGAAACAAATTCGACGAAATCAAGC
TGAAACATGTCTGCGACAGCTAAAAAGTCTCATAATTCGAAGACCTCTAAGAAGAAATTCAAAGAGGACAGAGA
ATTTGGGACACCAAAAAGATTTTAAAGAGATGATGTTCCTTTCGGGATTGATCGTTTGTGTTTGTGATTTTA
TTTTATATTGTTATCTGTTTCTGTGTATAGACTGTTTGAGATTGGCGCTTGGCCGACTCATTGCTTACCATAGG
GGAACGGACTTTGTTTGTGTTGTTATTTTATTTGATTTTATTTAAATTTCTCAATGATCTGAAAAGGCCCTCGAGG

CTAAGAGATTATTGGGGGGTGAGTAAGTACTTTTAAAGTGATGATGGTTACAAAGGCAAAAGGGGTAAAACCCCT
CGCCTACGTAAGCGTTATTACGCCCCGATCCCCCGGGGAGCTCGAATTCGCTGAAATCACCAGTCTCTCTCTACA
AATCTATCTCTCTCTATTTTTTCCATAAATAATGTGTGAGTAGTTTCCCGATAAGGGAAATTAGGGTTCTTATAG
GGTTTCGCTCATGTGTTGAGCATATAAGAAACCCCTTAGTATGTATTTGTATTTGTAAAATACTTCTATTATCAAT
AAAATTTCTAATTCCTAAAACCAAAATCCAGTACTAAAATCCAGATCTCCTAAAGTCCCTATAGATCTTTGTCTGT
GAATATAAACCAGACACGAGACGACTAAACCTGGAGCCCAGACGCCGTTTCAAGCTAGAAGTACCCTTAGGCAG
GAGGCCGTTAGGGAAAAGATGCTAAGGCAGGGTTGGTTACGTTGACTCCCCCGTAGGTTTGGTTTAAATATGATG
AAGTGGACGGAAGGAAGGAGACAAGGAAGGATAAGGTTGCAGGCCCTGTGCAAGGTAAGAAGATGGAAATTT
TGATAGAGGTACGCTACTATACTTATACTATAACGCTAAGGGAATGCTTGTATTTATACCCTATACCCCCTAATAA
CCCCCTTATCAATTTAAGAAATAATCCGCATAAGCCCCCGCTTAAAAAATTGGTATCAGAGCCATGAATAGGTCTAT
GACCAAAACTCAAGAGGATAAAACCTCACCAAAATACGAAAGAGTTCTTAACTCTAAAGATAAAAGATCTTTCAA
GATCAAAACTAGTTCCCTCACACCGGAGCATGCGATATCCTCGACCTGCAGGCATGCAAGCTTGGCGTAATCATG
GTCATAGCTGTTTCTGTGTGAAATTGTTATCCGCTCACAATTCACACAACATACGAGCCGGAAGCATAAAGTG
TAAAGCCTGGGGTGCCATATGAGTGAGCTAACTCACATTAATTGCGTTGCGCTCACTGCCCCGCTTCCAGTCGGG
AAACCTGTCTGTGCCAGCTGCATTAATGAATCGGCCAACGCGCGGGGAGAGGCGGTTTGCATTTGGGCCAAAGAC
AAAAGGGGCGACATTCAACCGATTGAGGGAGGGAAGGTAATATTGACGGAAATTATTATTAAAGGTGAATTATC
ACCGTCACCGACTTGAGCCATTGCGAATTAGAGCCAGCAAAATCACCAGTAGCACCATTACCATTAGCAAGGCC
GGAAACGTCACCAATGAAACCATCGATAGCAGCACCCTAATCAGTAGCAGAGAATCAAGTTTGCCTTTAGCGTC
AGACTGTAGCGCGTTTTTCATCGGCATTTTTCGGTCATAGCCCCCTTATTAGCGTTTGGCATCTTTTTCATAACAAA
ATCACCGGAACAGAGCCACCACCGGAACCGCCTCCCTCAGAGCCGCCACCCTCAGAACCGCCACCCTCAGAGCC
ACCACCCTCAGAGCCGCCACCAGAACCACCAGAGCCGCCGCCAGCATTGACAGGAGGCCCGATCTAGTAACA
TAGATGACACCGCGCGCGATAATTTATCCTAGTTTTCGCGCTATATTTTGTTCCTATCGCGTATTAAATGTATA
ATTGCGGGACTCTAATCATAAAAACCCATCTCATAAATAACGTCATGCATTACATGTTAATTATTACATGCTTAA
CGTAATTCAACAGAAATTATATGATAATCATCGCAAGACCGGCAACAGGATTCAATCTTAAGAACTTTATTGCC
AAATGTTTGAACGATCGGGGATCATCCGGGTCTGTGGCGGGAATCCACGAAAATATCCGAACGCAGCAAGATAT
CGCGGTGCATCTCGGTCTTGCTTGGCAGTCGCGCGCGACGCCGTTGATGTGGACGCCGGGCCGATCATATTGT
CGCTCAGGATCGTGGCGTTGTGCTTGTTCGGCCGTTGCTGTCTGAATGATATCGGCACCTTCGACCGCCTGTTCGG
CAGAGATCCCCGTGGCGAAGAACTCCAGCATGAGATCCCCGCGCTGGAGGATCATCCAGCCGCGCTCCCGAAAA
CGATTCCGAAGCCCAACCTTTCATAGAAGGCGCGGTGGAATCGAAATCTCGTGATGGCAGGTTGGGCGCTCGCTT
GGTCGGTCATTTTCAACCCAGAGTCCCGCTCAGAAGAACTCGTCAAGAAGGCGATAGAAGGCGATGCGCTGCGA
ATCGGGAGCGCGATACCGTAAAGCACGAGGAAGCGGTACGCCATTTCGCGCCAAGCTCTTCAGCAATATCACG
GGTAGCCAACGCTATGTCTGATAGCGGTCCGCCACACCCAGCCGCCACAGTCGATGAATCCAGAAAAGCGGCC
ATTTTCCACCATGATATTCGGAAGCAGGCATCGCCATGGGTACGACGAGATCATCGCCGTCGGGCATGCGCGC
CTTGAGCCTGGCGAACAGTTTCGGCTGGCGCGAGCCCTGATGCTCTTCGTCCAGATCATCCTGATCGACAAGACC
GGCTTCCATCCGAGTACGTGCTCGCTCGATGCGATGTTTCGCTTGGTGGTCAATGGGCAGGTAGCCGATCAAG
CGTATGCAGCCCGCGCATTGCATCAGCCATGATGGATACTTTCTCGGCAGGAGCAAGGTGAGATGACAGGAGATC
CTGCCCCGGCACTTCGCCCCAATAGCAGCCAGTCCCTTCCCGCTTCAGTGACAACGTCGAGCACAGCTGCGCAAGG
AACGCCCGTTCGTGGCCAGCCACGATAGCCGCGCTGCCTCGTCTGAGTTTATTTCAGGGCACCAGGACAGGTTCGGT
CTTGACAAAAGAACCAGCGCGCCCTGCGCTGACAGCCGGAACACGCGCGCATCAGAGCAGCCGATGTCTGTTG
TGCCAGTATAGCCGAATAGCCTTCCACCCAGCGCGGAGAACCTGCGTGCAATCCATCTTGTTCATCAT
GCGAAACGATCCAGATCCGGTGCAGATTATTTGGATTGAGAGTGAATATGAGACTCTAATTGGATACCGAGGGGA
ATTTATGGAACGTCAGTGGAGCATTTTTGACAAGAAATATTTGCTAGCTGATAGTGACCTTAGGCGACTTTTGAA
CGCGCAATAATGGTTTTCTGACGTATGTGCTTAGCTCATTAACCTCCAGAAACCCGCGGCTGAGTGGCTCCTTCAA
CGTTGCGGTTCTGTGCTTCCAAACGTAACCGGCTTGTCCCGCTCATCGCGGGGGTTCATAACGTGACTCCCT
TAATTCTCCGCTCATGATCAGATTGTCGTTTCCCGCCTTCAGTTTAAACTATCAGTGTTTGACAGGATATATTGG
CGGGTAAACCTAAGAGAAAAGAGCGTTTATTAGAATAATCGGATATTTAAAGGGCGTGAAAAGGTTTATCCGTT
CGTCCATTTGTATGTGCATGCCAACACAGGGTTCCCCAGATCTGGCGCCGCCAGCGAGACGAGCAAGATTGGC
CGCCGCCGAAACGATCCGACAGCGCGCCAGCACAGGTGCGCAGGCAAAATTGCACCAACGCATACAGCGCCAGC
AGAATGCCATAGTGGCGGTGACGTCTGAGTGAACACAGATCCGCGCAGGAGGCCCGGACACCGGCATAATC
AGGCCGATGCCGACAGCTCGAGCGGACAGTGCTCAGAATTACGATCAGGGGTATGTTGGTTTACGCTTGGC
CTCCGGACAGCCTCCGCTGGTCCGATTGAACGCGCGGATTCTTTATCACTGATAAGTTGGTGGACATATTATGT
TTATCAGTGATAAAGTGTCAAGCATGACAAAGTTGCAGCCGAATACAGTGATCCGTGCCGCCCTGGACCTGTTGA
ACGAGGTGCGCGTAGACGGTCTGACGACACGCAAACTGGCGGAACGGTTGGGGGTTTACGAGCCGGCGCTTTACT
GGCACTTCAGGAACAAGCGGGCGCTGCTCGACGCACTGGCCGAAGCCATGCTGGCGGAGAATCATACGCATTCCG
TGCCGAGAGCCGACGACGACTGGCGCTCATTTCTGATCGGGAATGCCCGCAGCTTCAGGCAGGCGCTGCTCGCCT
ACCGCGATGGCGCGCGCATCCATGCCGGCACGCGACCGGGCGCACCGCAGATGGAAACGGCCGACGCGCAGCTTC
GCTTCTCTGCGAGGCGGGTTTTTCGGCCGGGGACGCCGTCAATGCGCTGATGACAATCAGCTACTTCACTGTTG
GGGCGGTGCTTGAGGAGCAGGCCGGCGACAGCGATGCCGGCGAGCGCGGCGGCACCGTTGAACAGGCTCCGCTCT
CGCCGCTGTTGCGGGCCGCGATAGACGCCCTTCGACGAAGCCGGTCCGGACGACGCTTCGAGCAGGACTCGCGG
TGATTGTCGATGGATTGGCGAAAAGGAGGCTCGTTGTGAGGAACGTTGAAGGACCGAGAAAGGTTGACGATTGAT
CAGGACCGCTGCCGAGCGCAACCCACTCACTACAGCAGAGCCATGTAGACAACATCCCCCTCCCCCTTCCACCG

CGTCAGACGCCCGTAGCAGCCCGCTACGGGCTTTTTTCATGCCCTGCCCTAGCGTCCAAGCCTCACGGCCGCGCTC
GGCCTCTCTGGCGGCCCTTCTGGCGCTCTTCCGCTTCCTCGCTCACTGACTCGCTGCGCTCGGTGCTTCGGCTGCG
GCGAGCGGTATCAGCTCACTCAAAGGCGGTAATACGGTTATCCACAGAATCAGGGGATAACGCAGGAAGAACAT
GTGAGCAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCC
CCCTGACGAGCATCACAAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATACCAGGC
GTTTTCCCCCTGGAAGCTCCCTCGTGCGCTCTCTGTTCCGACCTGCGCTTACCGGATACCTGTCCGCCTTTCT
CCCTTCGGGAAGCGTGGCGCTTTTCCGCTGCATAACCCCTGCTTCGGGGTCATTATAGCGATTTTTTCGGTATATC
CATCCTTTTTTCGCACGATATACAGGATTTTGCCAAAGGGTTCGTGTAGACTTTCCTTGGTGTATCCAAACGGCGTC
AGCCGGGCAGGATAGGTGAAGTAGGCCACCCGCGAGCGGGTGTTCCTTCTTCACTGTCCCTTATTTCGCACCTGG
CGGTGCTCAACGGGAATCTGCTCTGCGAGGCTGGCCGGCTACCGCCGGCGTAACAGATGAGGGCAAGCGGATGG
CTGATGAAACCAAGCCAACCAGGAAGGGCAGCCACCTATCAAGGTGTACTGCCTTCCAGACGAACGAAGAGCGA
TTGAGGAAAAGGCGGCGGCGGCCGCGCATGAGCCTGTGCGCTACCTGCTGGCCGTGCGCCAGGGCTACAAAATCA
CGGGCGTCTGAGTATGAGCACGTCCGCGAGCTGGCCCGCATCAATGGCGACCTGGGCGCGCTGGGCGGCTGC
TGAAACTCTGGCTCACCGACGACCCGCGCACGGCGCGGTTGCGGTGATGCCACGATCCTCGCCCTGCTGGCGAAGA
TCGAAGAGAAGCAGGACGAGCTTGGCAAGGTGATGATGGGCGTGGTCCGCCCCAGGGCAGAGCCATGACTTTTTT
AGCCGCTAAAACGGCCGGGGGTGCGCGTGATTGCCAAGCACGTCCCATGCGCTCCATCAAGAAGAGCGACTTC
GCGGAGCTGGTGAAGTACATCACCGACGAGCAAGGCAAGACCGAGCGCCTTTGCGACGCTCACCGGGCTGGTATGC
CCTCGCCGTGGGCTGGCGGCGCTCTATGGCCCTGCAACCGCGCAGAAACGCGCTCGAAGCCGTGTGCGAGACA
CCGCGGGCGCGCGGCTTGTGGATACCTCGCGGAAAACCTTGGCCCTCACTGACAGATGAGGGGCGGACCTTGACAC
TTGAGGGGCGGACTCACCGGCGCGGCGTTGACAGATGAGGGGCGGCTCGATTTGCGCCGGCGACGTGGAGCTG
GCCAGCCTCGCAATCGGCGAAAACGCTGATTTTACGCGAGTPTCCACAGATGATGTGGACAAGCCTGGGGAT
AAGTGCCCTGCGGTATTGACACTTGAGGGGCGGACTACTGACAGATGAGGGGCGCGATCCTTGACACTTGAGGG
GCAGAGTGCTGACAGATGAGGGGCGCACCTATTGACATTTGAGGGGCTGTCCACAGGCAGAAAATCCAGCATTTG
CAAGGGTTTTCCGCCCCGTTTTTCGGCCACCGCTAACCTGTCTTTTAACTGCTTTTAAACCAATATTTATAAACCT
TGTTTTTAACAGGGCTGCGCCCTGTGCGCGTGACCGCGCACGCCGAAGGGGGGTGCCCCCCTTCTCGAACCCCT
CCCGGCCCGCTAACGCGGGCTCCCATCCCCCAGGGGTGCGCCCCCTCGGCCGCGAACGGCTCACCCCAAAAA
TGGCAGCGCTGGCAGTCTTGGCATTGCGGGGATCGGGGCAGTAACGGGATGGGCGATCAGCCGAGCGCGACGC
CCGGAAGCATTTGACGTGCGCGAGGTGCTGGCATCGACATTCAGCGACAGGTGCGCGGCGAGTGAGGGCGCGGCG
TGGGTGGCGGCTGCCCTTCACTTCGGCCGTGCGGGCATTCACGGACTTCATGGCGGGGCGCGCAATTTTACCT
TGGGCATTCTTGGCATAGTGGTTCGCGGGTGCCTGCTGCTGTTTCGGGGGTGCGATAAACCCAGCGAACCATTTGA
GGTGATAGGTAAGATTATACCGAGGTATGAAAACGAGAATTGGACCTTTACAGAATTACTCTATGAAGCGCCATA
TTTAAAAAGCTACCAAGACGAAGAGGATGAAGAGGATGAGGAGGCAGATTGCCTTGAATATATTGACAATACTGA
TAAGATAATATATCTTTTATATAGAAGATATCGCCGTATGTAAGGATTTTCAAGGGGCAAGGCATAGGCAGCGCGC
TTATCAATATATCTATAGAATGGGCAAAAGCATAAAAACTTGATGGAATAATGCTTGAACCCAGGACAATAACC
TTATAGCTTGTAAATCTATCATAATTGGGTAATGACTCCAACCTTATTGATAGTGTTTTATGTTACAGATAATGCC
CGATGACTTTGTCTATGCAGCTCCACCGATTTTGAGAACGACAGCGACTTCCGTCCCAGCGGTGCCAGGTGCTGCC
TCAGATTACAGTTATGCCGCTCAATTCGCTGCGTATATCGCTTGTGATTACGTGCAGCTTTCCCTTCAGGCGGG
ATTCATACAGCGGCCAGCCATCCGTCATCCATATCACCACGTCAAAGGGTGACAGCAGGCTCATAAGACGCCCCA
GCGTCGCCATAGTGCCTTACCGAATACGTGCGCAACACCGCTCTTCGGGAGACTGTACACGCTAAACAGCC
AGCGCTGGCGCATTTAGCCCGACATAGCCCCACTGTTCGTCCATTTCCGCGCAGACGATGAGCTGCTGCCG
GCTGTATGCGCGAGGTTACCGACTGCGGCCTGAGTTTTTAAAGTGACGTAAAATCGTGTGAGGCCAACGCCCAT
AATGCGGGCTGTTGCCCGCATCCAACGCCATTGATGGCCATATCAATGATTTTCTGGTGCGTACCGGGTTGAGA
AGCGGTGTAAGTGAAGTGCAGTTGCCATGTTTTACGGCAGTGAGAGCAGAGATAGCGCTGATGTCCGGCGGTGCT
TTTGCCGTTACGCACCACCCGTCAGTAGCTGAACAGGAGGACAGCTGATAGACACAGAAGCCACTGGAGCACC
TCAAAAACACCATCATACACTAAATCAGTAAGTTGGCAGCATCACCCATAATTGTGGTTTTCAAATCGGCTCCGT
CGATACTATGTTATACGCCAACTTTGAAAACAACCTTTGAAAAGCTGTTTTCTGGTATTTAAGGTTTTAGAATGC
AAGGAACAGTGAATTGGAGTTCGTCTTGTATAATTAGCTTCTTGGGGTATCTTTAAATACTGTAGAAAAGAGGA
AGGAAATAATAAATGGCTAAAATGAGAATATCACCGGAATTGAAAAAACTGATCGAAAAATACCGCTGCGTAAAA
GATACGGAAGGAATGTCTCCTGCTAAGGTATATAAGCTGCTGGGAGAAAATGAAAACCTATATTTAAAAATGACG
GACAGCCGGTATAAAGGACCACTATGATGTGGAACGGGAGACATGATGCTATGGCTGGAAGGAAAGCTG
CCTGTTCCAAAGGTCCTGCACTTTGAACGGCATGATGGCTGGAAGCAATCTGCTCATGAGTGAGGCGGATGGCGTC
CTTTGCTCGGAAGAGTATGAAGATGAACAAAGCCCTGAAAAGATTATCGAGCTGTATGCGGAGTGATCAGGCTC
TTTCACTCCATCGACATATCGGATTGTCCCTATACGAATAGCTTAGACAGCCGCTTAGCCGAATTGGATTACTTA
CTGAATAACGATCTGGCCGATGTGGATTGCGAAAACCTGGGAAGAAGACACTCCATTTAAAGATCCGCGCGAGCTG
TATGATTTTTTTAAAGACGGAAAAGCCCCGAAGAGGAACCTTGTCTTTTCCACGGCGACCTGGGAGACAGCAACATC
TTTGTGAAAGATGGCAAAGTAAGTGGCTTTATTGATCTTGGGAGAAGCGGCAGGGCGGACAAGTGGTATGACATT
GCCTTCTGCGTCCGGTCGATCAGGGAGGATATCGGGGAAGAACAGTATGTCGAGCTATTTTTTGACTTACTGGGG
ATCAAGCCTGATTGGGAGAAAATAAAATATTATATTTTACTGGATGAATTGTTTTAGTACCTAGATGTGGCGCAA
CGATGCCGGCGACAAGCAGGAGCGCACCGACTTCTCCGCATCAAGTGTTTTGGCTCTCAGGCCGAGGCCACGG
CAAGTATTTGGGCAAGGGTCCGTGGTATTCTGTCAGGGCAAGATTGGAATACCAAGTACGAGAAGGACGGCCA
GACGGTCTACGGGACCGACTTCATTGCCGATAAGGTGGATTATCTGGACACCAAGGCACCAGGCGGTCAAATCA

GGAATAAGGGGCACATTGCCCGGCGTGAGTCGGGGCAATCCCGCAAGGAGGGTGAATGAATCGGACGTTTGACCG
GAAGGCATACAGGCAAGAACTGATCGACGCGGGGTTTTCCGCCGAGGATGCCGAAACCATCGCAAGCCGACCGT
CATGCGTGCGCCCCGCGAAACCTTCCAGTCCGTCGGCTCGATGGTCCAGCAAGCTACGGCCAAGATCGAGCGCGA
CAGCGTGCAACTGGCTCCCCCTGCCCTGCCCGCGCCATCGGCCGCCGTGGAGCGTTTCGCGTCGCTCGAACAGGA
GGCGGCAGGTTTGGCGAAGTCGATGACCATCGACACGCGAGGAACTATGACGACCAAGAAGCGAAAAACCGCCGG
CGAGGACCTGGCAAAACAGGTCAGCGAGGCCAAGCAGGCCCGGTTGCTGAAACACACGAAGCAGCAGATCAAGGA
AATCGAGCTTTCCCTTGTTCGATATTGCGCCGTGGCCGGACACGATGCGAGCGATGCCAAACGACACGCCCCGCTC
TGCCCTGTTTACCACGCGCAACAAGAAAAATCCCGCGCGAGGCGCTGCAAAACAAGGTCATTTTCCACGTCACAA
GGACGTGAAGATCACCTACACCGGCGTCGAGCTGCGGGCCGACGATGACGAACTGGTGTGGCAGCAGGTGTTGGA
GTACGCGAAGCGCACCCCTATCGGCGAGCCGATCACCTTACGTTTACGAGCTTTGCCAGGACCTGGGCTGGTC
GATCAATGGCCGGTATTACACGAAGGCCGAGGAATGCTGTGCGCCCTACAGGCGACGGCGATGGGCTTACGTC
CGACCGCGTTGGGCACCTGGAATCGGTGTGCTGCTGCACCGCTTCCGCGTCTGGACCGTGGCAAGAAAACGTC
CCGTTGCCAGGTCTGATCGACGAGGAAATCGTCTGCTGTTTGTGCTGGCGACCACTACACGAAATTCATATGGGA
GAAGTACCGCAAGCTGTGCGCGACGGCCCCGACGGATGTTGACTATTTTTCAGCTCGCACCGGGAGCCGTACCCGCT
CAAGCTGGAAACCTTCCGCCCTCATGTGCGGATCGGATTCACCCGCGTGAAGAAGTGGCGCGAGCAGGTGCGCGA
AGCCTGCGAAGAGTTGCGAGGACGCGGCTGGTGGAAACGCGCTGGGTCAATGATGACCTGGTGCATTGCAACG
CTAGGGCCTTTGTGGGTGAGTTCCGCTGGGGGTTTCAGCAGCCAGCGCTTACTGGCATTTCAGGAACAAGCGGG
CACTGCTCGAGCACTTGTCTGCTCAGTATCGCTCGGGACGACGCGCGCTCTACGAACCTGCCGATAAACAGA
GGATTAAAATTGACAATTTGTGATTAAGGCTCAGATTTCAGCGGCTGGAGCGGCCGACGTGCAGGATTTCGCGAG
ATCCGATTGTGCGCCCTGAAGAAAGCTCCAGAGATGTTGCGGTCCGTTTACGAGCACGAGGAGAAAAAGCCCATG
GAGGCGTTGCTGAACGGTTGCGAGATGCCGTGGCATTGCGCGCCTACATCGACGGCGAGATCATTTGGGCTGTG
GTCTTCAAACAGGAGGACGGCCCAAGGACGCTCACAAGGCGCATCTGTCCGGCGTTTTCTGGAGCCCGAACAG
CGAGGCCGAGGGGTGCGCGGTATGCTGCTGCGGGCGTTGCCGGCGGTTTATTGCTCGTGATGATCGTCCGACAG
ATTCCAACGGGAATCTGGTGGATGCGCATCTTCATCTCGGCGCACTTAATATTTTCGCTATTTCTGGAGCTTGTTG
TTTATTTTCGGTCTACCGCCTGCCGGGCGGGGTGCGGGCGACGGTAGGCGCTGTGCAGCCGCTGATGGTCTGTTT
ATCTCTGCCGCTCTGCTAGGTAGCCCGATACGATTGATGGCGGTCTCGGGGGCTATTTGCGGAACGCGGGCGTG
GCGCTGTTGGTGTGACACCAAACGCGAGCGCTAGATCTGTGCGGCTCGCAGCGGGCCTGGCGGGGGCGGTTTCC
ATGGCGTTTCGGAACCGTGCTGACCCGCAAGTGGCAACCTCCCGTGCCCTCTGCTCACCTTTACCGCCTGGCAACTG
GCGGCGGAGGACTTCTGCTCGTTCCAGTAGCTTTAGTGTGTTGATCCGCCAATCCCGATGCCCTACAGGAACCAAT
GTTCTCGGCGCTGGCGTGGCTCGGCGCTGATCGGAGCGGGTTTAACTTACTTCTTTTGGTTCCGGGGGATCTCGCGA
CTCGAACCTACAGTTGTTTCTTACTGGGCTTTCTCAGCCCCAGATCTGGGGTTCGATCAGCCGGGGATGCATCAG
GCCGACAGTCGGAACCTTCGGGTCCCCGACCTGTACCATTCCGGTGAGCAATGGATAGGGGAGTTGATATCGTCAAC
GTTCACTTCTAAAGAAATAGCGCCACTCAGCTTCTCAGCGGCTTTATCCAGCGATTTCTTATTATGTGCGCATA
GTTCTCAAGATCGACAGCCTGTACGGTTAAGCGAGAAATGAATAAGAAGGCTGATAATTCGGATCTCTGCGAGG
GAGATGATATTTGATCACAGGCGCAACGCTCTGTCTATCGTTACAATCAACATGCTACCCTCCGCGAGATCATCC
GTGTTTCAAACCCGGCAGCTTAGTTGCCGTTCTTCCGAATAGCATCGGTAACATGAGCAAAGTCTGCCGCTTAC
AACGGCTCTCCCGCTGACGCCGTCCCGGACTGATGGGCTGCCGTATCGAGTGGTGAATTTGTGCCGAGCTGCCG
GTCGGGGAGCTGTTGGCTGGCTGGTGGCAGGATATATTGTGGTGTAACAAATTGACGCTTAGACAACTTAATAA
CACATTGCGGACGTTTTTAATGTACTGGGGTGGTTTTCTTTTACCAGTGAGACGGGCAACAGCTGATTGCCCT
TCACCGCTTGGCCCTGAGAGAGTTGCAGCAAGCGGTTCCACGTTGGTTTGGCCCCAGCAGGCGAAATCTGTGTTGA
TGGTGGTTCCGAAATCGGCAAAATCCCTTATAAATCAAAGAATAGCCCCGAGATAGGGTTGAGTGTGTTCCAGT
TTGGAACAAGAGTCCACTATTAAAGAACGTGGACTCCAACGTCAAAGGGCGAAAAACCGTCTATCAGGGCGATGG
CCCCTACGTGAACCATCACCCAAATCAAGTTTTTTGGGGTCGAGGTGCCGTAAAGCACTAAATCGGAACCCCTAA
AGGGAGCCCCGATTTAGAGCTTGACGGGGAAGCCGGCGAACGTGGCGAGAAAGGAAGGGAAGAAAGCGAAAGG
AGCGGGCGCCATTACGGCTGCGCAACTGTTGGGAAGGGCGATCGGTGCGGGCCTCTTCGCTATTACGCCAGCTGG
CGAAAGGGGGATGTGCTGCAAGCGGATTAAGTTGGGTAACGCCAGGGTTTTTCCAGTACAGACGTTGTAAAACGA
CGGCCAGTGAATTCGAGCTCGGTACCCCCCTACTCCAAAAATGTCAAAGATACAGTCTCAGAAGACCAAAGGGCT
ATTGAGACTTTTCAACAAAGGGTAATTTCCGGAAACCTCCTCGGATTCATTGCCCAGCTATCTGTCACTTCATC
GAAAGGACAGTAGAAAGGAAGGTGGCTCCTACAAATGCCATCATTTGCGATAAAGGAAAGGCTATCATTTCAAGAT
GCCCTGCGCAGTAGTGGTCCCAAAGATGGACCCCAACAGCAGGAGCATCGTGGAAGAAAGAACGCTCCCAACC
ACGTCTTCAAAGCAAGTGGATTGATGTGACATCTCCACTGACGTAAGGGATGACGCAACATCCCACTATCCTTCG
CAAGACCCCTTCTTCTATATAAGGAAGTTTCAATTTTCAATTTGGAGAGGACAGCCCCAAGCTTTCTAGAGGATCCATAAA
ACATTTCAATCCTTTGAACGCGGTAGAACGTGCTAATTGGATTTTGGTGAGAACGCGGTAGAACGTACTTATCAC
CTACAGTTTTATTTTGTGTTTTCTTTTTGGTTAATCTATCCAGCTTAGTACCGAGTGGGGGAAAGTGACTGGTGT
GCCTAAAACCTTTTTCTTTTGATACTTTGTAAAAATACATACAGATACAATGGCGAACGGTAACCTCAAGTTGTCTC
AATTGCTCAATGTGGACGAGATGTCTGCTGAGCAGAGGAGTCATTTCTTTGACTTGATGCTGACTAAACCTGATT
GTGAGATCGGGCAAATGATGCAAAGAGTTGTTGTTGATAAAGTCGATGACATGATTAGAGAAAGAAAGACTAAAG
ATCCAGTGATTGTTTCATGAAGTTCTTTCTCAGAAGGAACAGAACAGTTGATGGAAATTTATCCTGAATTCATA
TCGTGTTTAAAGACGACAAAAACATGGTTCATGGGTTTGGCGCTGCTGAGCGAAAACCTACAAGCTTTATTGCTTT
TAGATAGAGTTCTGCTCTGCAAGAGGTGGATGACATCGGTGGTCAATGGTCTGTTTTGGGTAACCTAGAGGTGAGA
AAAGGATTCAATCCTGTTGTCCAAATCTAGATATTCGGGATGATCAGAGAGAAATTTCTCGACAGATATTTCTTA

CTGCTATTGGTGATCAAGCTAGAAGTGGTAAGAGACAGATGTCGGAGAATGAGCTGTGGATGTATGACCAATTTT
GTGAAAATATTGCTGCGCCTAACGCGGTTAGGTGCAATAATACATATCAGGGTTGTACATGTAGGGGTTTT

METHOD OF MODULATING THE ALKALOID CONTENT OF TOBACCO

FIELD OF THE INVENTION

[0001] The present invention relates to methods of modulating the alkaloid content a plant or part thereof or cell or cell culture. The invention also extends to methods of modulating the expression and/or activity of polypeptides which modulate alkaloid content within plants. Alternatively, the invention provides methods of modulating the expression and/or activity of genes which encode polypeptides which modulate alkaloid content within plants. The invention also extends to constructs, which can be used to modulate the polypeptides. The invention further relates to plant cells and plants modified to achieve a modulation in alkaloid content. The invention also relates to a processed and harvested leaf from such modulated plants and use thereof in a tobacco industry product, including combustible smoking articles.

BACKGROUND

[0002] Alkaloids are a group of naturally occurring compounds which mostly contain basic nitrogen atoms and are produced by a large variety of organisms including bacteria, fungi, plants and animals.

[0003] Alkaloids may be classified according to the similarity of the carbon skeleton e.g. indole-, isoquinoline- and pyridine-like. Pyridine derivatives are one class of monomeric alkaloids; this class includes simple derivatives of pyridine, polycyclic condensed and noncondensing pyridine derivatives and sesquiterpene pyridine derivatives. Examples are nicotine, normicotine, pseudooxynicotine, anabasine, myosmine and anatabine.

[0004] Most of the known biological functions of alkaloids are related to protection. Neuroactive molecules, such as caffeine, cocaine, morphine, and nicotine, act as defence compounds against invading predators. The accumulation of these alkaloids is the result of signal transduction cascades that monitor gene expression, enzyme activities, and alkaloid concentrations. The fine-tuning of alkaloid content in the plant involves negative feedback loops and degradative pathways.

[0005] Nicotine occurs naturally in several varieties of plants but is found at the highest level in the tobacco plant. Cultivated tobacco produces 2-4% alkaloids of total dry weight. Nicotine is produced in wild and cultivated *Nicotiana* species and plays an important role in plant defence against herbivores and insects (Voelckel et al. (2001) *Oecologia* 127 (2): 274-280, incorporated herein by reference). It accounts for ~90% of the total alkaloid content. The remaining 10% of the alkaloid pool is mostly constituted by the structurally related compounds normicotine, anatabine, anabasine and pseudooxynicotine (PON).

[0006] The regulation of alkaloid content in tobacco is complex. Several factors including genotype, environment, fertilization and agronomic practices (e.g. topping) affect alkaloid levels in tobacco plants. Some key regulators of nicotine biosynthesis are well characterized, for example putrescine N-methyltransferase (PMT), which plays a pivotal role in this pathway, is activated by members of the ethylene responsive factor (ERF) superfamily, the largest

transcription factor family in the tobacco genome (Rushton et al. (2008) *Plant Physiol.* 147 (1): 280-295 incorporated herein by reference).

[0007] Tobacco pyridine alkaloids are precursors of tobacco-specific nitrosamines (TSNAs) that form during the post-harvest leaf curing. The four primary TSNAs found in cured tobacco leaves are N'-nitrosanornicotine (NNN), N'-nitrosoanatabine (NAT), N'-nitrosoanabasine (NAB) and 4-(methyl nitrosamino)-1-(3-pyridyl)-1-butanone (NNK). During the post-harvest leaf curing, reactions between pyridine alkaloids and nitrosating species leads to the formation of TSNAs. PON is likely to function as the direct precursor in the synthesis of the TSNA NNK (Bush et al., 2001 incorporated herein by reference). Reducing the production and accumulation of TSNAs is of high importance. The CYP82E family of nicotine demethylase genes is one of the primary regulators of nicotine to normicotine conversion, and altering their activity or accumulation may result in a decrease in NNN levels.

[0008] As described in the Examples, the inventors sought to investigate genes responsible for alkaloid and/or TSNA precursor synthesis, with the aim of modulating alkaloid content in plants, e.g. decreasing TSNA content in tobacco.

SUMMARY OF THE INVENTION

[0009] It has been surprisingly found that by modulating the activity or expression of a YIPF-related protein, the alkaloid content and/or TSNA content or precursor of TSNA content of plants can be modulated. The gene(s) as taught herein, for example Nitab4.5_0005388g0090.2, are regulators of alkaloid and TSNA precursor content in cultivated tobacco. In particular the gene(s) as taught herein, for example Nitab4.5_0005388g0090.2, is a regulator of alkaloid content in cultivated tobacco. Nitab4.5_0005388g0090.2 encodes a YIPF-related protein according to the present invention. Homologues of Nitab4.5_0005388g0090.2 are provided in Table 1.

[0010] According to the present invention, tobacco industry products with modulated alkaloid content and commercially desirable traits sought after by consumers of tobacco industry products can be produced. In some instances, consumers may desire a product with low levels of alkaloid content. In some instances, consumers may desire a product with low levels of TSNA precursors.

[0011] The present invention may be particularly useful in the field of plant molecular farming, where plants (such as tobacco and other *Nicotiana* spp.) are used for the production of proteins, peptides, and metabolites e.g. for the production of therapeutics and pharmaceuticals such as antibiotics, virus like particles, or neutraceuticals or small molecules. Tobacco has been used for the development of an HIV-neutralising antibody in an EU-funded project called PharmPlant and Medicago Inc., Canada have worked on a tobacco-based platform for the production of virus-like particles for flu vaccine manufacture.

[0012] Thus, a plant according to the present invention may be used for molecular farming to reduce or eliminate the presence of nicotinic alkaloids. The use of a low nicotine plant or rootstock is beneficial in molecular farming and would reduce downstream processing costs associated with purification.

[0013] The present inventors have surprisingly determined a method for modulating (e.g. decreasing) the alkaloid content, of a plant (e.g. a tobacco plant) by modulating (e.g.

decreasing) the activity or expression of a YIPF-related protein. The alkaloid content (e.g. the content of one or more of nicotine, nornicotine, PON, anabasine, anatabine or myosmine, suitably the content of one or more of nicotine, nornicotine, PON, anabasine or anatabine) of a plant (e.g. tobacco plant) may be decreased by decreasing the activity or expression of a YIPF-related protein or may be increased by increasing the activity or expression of a YIPF-related protein.

[0014] Prior to the present invention it had not been known that modulation of the activity or expression of a YIPF-related protein as described herein could be used to modulate alkaloid content or modulate TSNA precursor content in particular, nornicotine, PON, anabasine and/or anatabine content.

[0015] The present invention provides a method of modulating (e.g. decreasing) the alkaloid content of a tobacco plant or a part thereof or tobacco plant cell, the method comprising modifying said plant or plant cell by modulating (e.g. decreasing) the activity or expression of a YIPF-related protein. In another aspect, the present invention provides a method of modulating (e.g. decreasing) the content of a tobacco specific nitrosamine (TSNA) or a precursor of a TSNA in a tobacco plant or plant part thereof or tobacco plant cell, the method comprising modifying said plant or plant cell by modulating (e.g. decreasing) the activity or expression of a YIPF-related protein.

[0016] In another aspect, the present invention provides a method for producing a plant or part thereof, a cell or cell culture, a plant propagation material, a leaf, a cut harvested leaf, a processed leaf or a cut and processed leaf which has modulated (e.g. decreased) alkaloid content, the method comprising modifying said plant or cell culture to modulate the activity or expression of a YIPF-related protein.

[0017] In another aspect, the present invention provides the use of at least one gene encoding a YIPF-related protein for modulating alkaloid content of a tobacco cell or tobacco plant or part thereof. Suitably the YIPF-related protein may:

[0018] a) comprise an amino acid sequence as set out in SEQ ID No. 3; or a functional variant or functional fragment or orthologue of SEQ ID No. 3; or a sequence which has at least 80% identity to SEQ ID No. 3; or a homologue of SEQ ID No. 3; or

[0019] b) is encoded by a nucleotide sequence as set out in SEQ ID No. 1 or 2; or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2; or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2.

[0020] Suitably, the alkaloid content is modulated (e.g. decreased) in comparison to a plant or cell culture which has not been modified to modulate the activity or expression of a YIPF-related protein. In another aspect, the present invention provides a tobacco plant or part thereof or a tobacco cell or cell culture which has been modified to modulate (e.g. decrease) the activity or expression of a YIPF-related protein and the tobacco plant or part thereof or tobacco cell or cell culture has decreased alkaloid and or TSNA precursor content in comparison to an unmodified plant or unmodified cell or cell culture.

[0021] In a further aspect, the present invention provides a plant propagation material obtainable (e.g. obtained) from

a plant according to the present invention or from a plant or cell or cell culture produced by the method or use according to the present invention.

[0022] Suitably, the alkaloid content of the plant may be decreased in comparison to a plant or cell culture which has not been modified to modulate the activity or expression of a YIPF-related protein.

[0023] Suitably, the content of one or more alkaloids selected from nicotine, nornicotine, PON, anabasine, myosmine and anatabine may be modulated (e.g. decreased), preferably the content of nicotine, nornicotine and/or PON is modulated (e.g. decreased).

[0024] Suitably, the nicotine content may be decreased.

[0025] In one aspect, the present invention provides the use of a tobacco plant or part thereof or tobacco cell or cell culture according to the present invention, or of a plant produced according to the present invention to breed a plant.

[0026] In another aspect, the present invention provides the use of a tobacco plant or part thereof or a tobacco cell or cell culture according to the present invention, or of a plant produced by a method according to the present invention for production of a product.

[0027] In another aspect, the present invention provides the use of a tobacco plant or part thereof according to the present invention, or of a plant produced by a method according to the present invention to grow a crop.

[0028] In another aspect, the present invention provides the use of a tobacco plant or part thereof according to the present invention, or of a plant produced by a method according to the present invention to produce a leaf.

[0029] In one aspect, the present invention provides a harvested leaf of a plant according to the present invention, or obtainable from a plant propagated from a propagation material according to the present invention, or obtainable from a plant obtained by a use according to the present invention, or obtainable from a plant produced by a method according to the present invention. Suitably, the harvested leaf of a plant may be a cut harvested leaf.

[0030] In one aspect, the present invention provides a processed leaf, preferably a processed tobacco leaf, preferably a non-viable processed tobacco leaf:

[0031] obtainable (e.g. obtained) from a plant obtainable from a use according to the present invention;

[0032] obtainable (e.g. obtained) by processing a plant according to the present invention;

[0033] obtainable (e.g. obtained) from a plant propagated from a plant propagation material according to the present invention; or

[0034] obtainable (e.g. obtained) by processing a harvested leaf of a plant according to the present invention; or

[0035] obtainable (e.g. obtained) from a plant produced by a method according to the present invention.

[0036] Suitably, the processed leaf may be processed by curing, fermenting, pasteurising or a combination thereof.

[0037] Suitably, the processed leaf may be a cut processed leaf.

[0038] In one aspect, the present invention provides a cured tobacco material made from a plant or a part thereof according to the present invention, or a harvested leaf according to the present invention, or a processed leaf according to the present invention.

[0039] In another aspect, the present invention provides a tobacco blend comprising said cured tobacco material according to the present invention.

[0040] In one aspect, the present invention provides a tobacco industry product prepared from:

[0041] a tobacco plant or part thereof or tobacco cell or cell culture according to the present invention;

[0042] a tobacco plant or part thereof propagated from a tobacco plant propagation material according to the present invention;

[0043] a harvested leaf of a plant according to the present invention;

[0044] a processed leaf according to the present invention.

[0045] Suitably, the tobacco product may be a combustible smoking article.

[0046] Suitably, the tobacco product may be a smokeless tobacco product.

[0047] Suitably, the tobacco product may be a non-combustible aerosol provision system such as a tobacco heating device or an aerosol-generating device.

[0048] In another aspect, the present invention provides a combustible smoking article, non-combustible aerosol provisioning system, smokeless tobacco product or tobacco heating device comprising a plant or a part thereof according to the present invention or an extract (e.g. a tobacco extract) thereof or a tobacco cell culture according to the present invention; or a cured tobacco material according to the present invention; or a tobacco blend according to the present invention.

[0049] In one aspect, the present invention provides the use of a nucleotide sequence encoding a YIPF-related protein which:

[0050] a) encodes an amino acid sequence as set out in SEQ ID No. 3; or a functional variant or functional fragment or orthologue of SEQ ID No. 3; or a sequence which has at least 80% identity to SEQ ID No. 3; or a homologue of SEQ ID No. 3; or

[0051] b) comprises a sequence as set out in SEQ ID No. 1 or 2; or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2; or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2;

[0052] to select a plant having modulated (e.g. reduced) alkaloid content and/or modulated (e.g. reduced) content of TSNA or a precursor of a TSNA.

[0053] In another aspect, the present invention provides a mutant of a plant carrying a heritable mutation in a nucleotide sequence which:

[0054] a) encodes an amino acid sequence as set out in SEQ ID No. 3; or a functional variant or functional fragment or orthologue of SEQ ID No. 3; or a sequence which has at least 80% identity to SEQ ID No. 3; or a homologue of SEQ ID No. 3; or

[0055] b) comprises a sequence as set out in SEQ ID No. 1 or 2; or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2; or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2;

[0056] wherein said heritable mutation modulates (e.g. decreases) the activity or expression of YIPF-related protein and wherein the mutant plant has modulated (e.g. decreased) alkaloid content and/or modulated con-

tent of a TSNA or a precursor of a TSNA relative to a comparable plant which does not carry said heritable mutation.

[0057] In another aspect, the present invention provides progeny or seed of a mutant plant which carries the heritable mutation according to the present invention.

[0058] In one aspect, the present invention provides a harvested leaf, a processed leaf or cured tobacco material produced from a plant comprising a modification in a nucleotide sequence which:

[0059] a) encodes an amino acid sequence as set out in SEQ ID No. 3; or a functional variant or functional fragment or orthologue of SEQ ID No. 3; or a sequence which has at least 80% identity to SEQ ID No. 3; or a homologue of SEQ ID No. 3; or

[0060] b) comprises a sequence as set out in SEQ ID No. 1 or 2; or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2; or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2;

[0061] wherein said modification modulates (e.g. decreases) the activity or expression of YIPF-related protein and wherein said plant has modulated (e.g. decreased) alkaloid content and/or modulated content of a TSNA or a precursor of a TSNA relative to a comparable plant which does not carry said modification in said YIPF-related protein.

[0062] In one aspect, the activity or expression of at least one Nic1 ERF gene (such as any one or more of those in FIG. 5) and/or at least one Nic2 ERF gene (such as any one or more of those in FIG. 6) is modulated in addition to the at least one YIPF-related protein. Suitably, said at least one Nic1 ERF and/or Nic2 ERF gene may comprise a mutation which decreases its expression and/or activity. In one aspect, the activity of ERF199 (e.g. at least one of SEQ ID No. 7-10) is modulated (e.g. decreased) in addition to the modulation (e.g. decrease in activity and or expression) of the at least one YIPF-related protein. In one aspect, the activity of ERF189 (e.g. at least one of SEQ ID No. 11-14) is modulated (e.g. decreased) in addition to the modulation (e.g. decrease in activity and or expression) of the at least one YIPF-related protein. In one aspect, the activity of ERF199 (e.g. at least one of SEQ ID No. 7-10) and ERF189 (e.g. at least one of SEQ ID No. 11-14) is modulated (e.g. decreased) in addition to the modulation (e.g. decrease in activity and or expression) of the at least one YIPF-related protein. Exemplary sequences of ERF 199 and ERF189 are provided in FIGS. 7-14 (SEQ ID Nos 7-14). See also WO2018237107, which is incorporated herein in by reference in its entirety.

BRIEF DESCRIPTION OF THE DRAWINGS

[0063] Embodiments of the invention will now be described, by way of example only, with reference to the accompanying drawings, in which:

[0064] FIG. 1 shows the alkaloid content of plants targeted with VIGS construct for Nitab4.5_0005388g0090.2. Graph is representative of 2 biological replicates (N=12) analysed by one-way ANOVA and Tukey's multiple-comparison post-test. Values are shown as means $\pm$ SEM. Asterisks indicate statistical significance of P value $\leq$ 0.001. Pyridine alkaloids: nicotine, anabasine (ANAB), pseudoxyt nicotine (PON) and anatabine (ANAT).

[0065] FIG. 2 shows the genomic sequence of Nitab4.5_0005388g0090.2 (SEQ ID No. 1).

[0066] FIG. 3 shows the coding sequence of Nitab4.5_0005388g0090.2 (SEQ ID No. 2).

[0067] FIG. 4 shows the amino acid sequence of Nitab4.5_0005388g0090.2 (SEQ ID No. 3).

[0068] FIG. 5 provides a table of Nic1 ERFs.

[0069] FIG. 6 provides a table of Nic2 ERFs.

[0070] FIGS. 7-14 show the sequences of ERF199 (SEQ ID Nos 7-10) and ERF189 (SEQ ID Nos 11-14).

[0071] FIG. 15 shows the sequence of TRV1 (SEQ ID No 15) as used in Example 1.

[0072] Some sequences disclosed herein contain “X” or “N” in nucleotide sequences. “X” or “N” can be any nucleotide or a deletion or insertion of one or more nucleotides. For example, in some cases a string of “X”s or “N”s are shown. The number of “X”s or “N”s does not necessarily correlate with the actual number of nucleotides at that position. There may be more or fewer nucleotides than shown as “X” or “N” in the sequence.

DETAILED DESCRIPTION

[0073] For the first time the present inventors have shown that by modulating the activity or expression of at least one YIPF-related protein in a plant (e.g. a tobacco plant) or a cell (e.g. tobacco cell), the alkaloid and/or TSNA precursor content of the plant (or processed plant) or cell can be modulated.

[0074] Nitab4.5_0005388g0090.2 is identified as a member of the YIP (for YPT/RAB GTPase Interacting Protein) family which is a trans-Golgi (TGN)-localised complex involved in the secretory pathway, as a positive regulator of pyridine alkaloid biosynthesis. This provides a mechanistic insight into the cross-talk between the adaptive cell wall composition, hormone signalling, and nicotine biosynthesis, while offering a genetic solution to alter nicotine content in tobacco plants. The present invention provides a method of modulating (e.g. decreasing) the alkaloid content of a plant or a part thereof, the method comprising modifying said plant by modulating (e.g. decreasing) the activity or expression of at least one YIPF-related protein.

[0075] Also provided is a method of modulating (e.g. decreasing) the content of a tobacco specific nitrosamine (TSNA) precursor in a tobacco plant or plant part thereof, the method comprising modifying said plant by modulating (e.g. decreasing) the activity or expression of at least one YIPF-related protein.

[0076] The at least one YIPF-related protein may be selected from an amino acid sequence as set out in SEQ ID No. 3, or a functional variant or functional fragment or orthologue thereof, or a sequence which has at least 80% identity to SEQ ID No. 3, or a homologue of SEQ ID No. 3; or said at least one YIPF-related protein may be encoded by a which comprises a sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2, or a homologue of SEQ ID No. 1 or 2.

[0077] In one embodiment at least two genes encoding a YIPF-related protein are modified selected from the group of: genes which encode polypeptides comprising an amino acid sequence as set out in SEQ ID No. 3 or a functional variant or functional fragment or orthologue thereof, or a sequence which has at least 80% identity to SEQ ID No. 3,

or a homologue of SEQ ID No. 3; or genes encoding a YIPF-related protein comprising a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2, or a homologue of SEQ ID No. 1 or 2.

[0078] In one embodiment, at least three, such as at least four, such as at least five, such as at least six, such as at least seven, such as at least eight, such as at least nine, such as ten YIPF-related proteins are modulated, wherein the YIPF-related proteins comprise an amino acid sequence as set out in SEQ ID No. 3 or a functional variant or functional fragment or orthologue thereof, or a sequence which has at least 80% identity to SEQ ID No. 3, or a homologue of SEQ ID No. 3; or comprise a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2, or a homologue of SEQ ID No. 1 or 2.

[0079] In one aspect, the at least one YIPF-related protein comprises or consists of an amino acid sequence as set out in: SEQ ID No. 3 or a functional variant or functional fragment or orthologue thereof, or a sequence which has at least 80% identity to SEQ ID No. 3; or wherein the YIPF-related protein comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2 or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2; or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2.

[0080] In one aspect, the activity or expression of at least one further gene is modulated. Suitably, at least two (or at least three or at least four or at least five or at least six or at least seven or at least eight or at least nine) additional genes selected from Table 1 or a sequence having at least 80% sequence identity thereto may also be modulated.

[0081] The “expression” of a YIPF-related protein may refer to the level of transcription, translation i.e. protein expression.

[0082] Measurement of the level or amount of a gene product may be carried out by any suitable method, for example comparison of mRNA transcript levels, protein or peptide levels, and/or phenotype of a plant, between a modified plant and comparable plant which has not been modified according to the present invention.

[0083] The term “a comparable product” as defined herein would be one derived from a plant (e.g. a tobacco plant) which had not been modified according to the present invention, but in which all other relevant features were the same (e.g. plant species, growing conditions, method of processing the plant, e.g. tobacco, etc.). The comparable product according to the present invention may mean a plant (e.g. a tobacco plant) or a part thereof, such as a leaf (e.g. a tobacco leaf), a harvested leaf (e.g. a harvested tobacco leaf), a cut harvested leaf (e.g. a cut harvested tobacco leaf), a processed leaf (e.g. a processed tobacco leaf) or plant propagation material (e.g. tobacco plant propagation material), or a product comprising said plant or part therefore, e.g. a tobacco industry product or combinations thereof obtainable or obtained from a plant which has not been modified in accordance with the present invention, e.g. to modulate the activity or expression of gene encoding a YIPF-related protein. In one embodiment a comparable

product is one which does not comprise gene encoding a YIPF-related protein whose activity or expression has been modulated.

[0084] The term “modifying” or “modified” as used herein means a plant (e.g. a tobacco plant) or nucleic acid sequence that has been altered or changed. The present invention comprises the modification of plants using techniques for genetic modification of plants or non-genetic modification of plants. Such methods are well known in the art and examples of genetic modification techniques include transformation, transgenics, cisgenics, and gene editing methods. Examples of non-genetic modification techniques include fast-neutron mutagenesis, chemical mutagenesis e.g. ethyl methane-sulfonate (EMS) mutagenesis and modern population analysis approaches.

[0085] In one embodiment a natural variant which has a modified gene encoding a YIPF-related protein is selected and that trait or gene is bred into a second plant which may have commercially desirable traits.

[0086] In one embodiment the plant according to the present invention is a transgenic plant. In one embodiment the plant according to the invention is a non-transgenic plant.

[0087] The term “unmodified plant” as defined herein would be a plant (e.g. a tobacco plant) which had not been modified according to the present invention, e.g. to modulate the activity or expression of a YIPF-related protein or to modify the nucleic acid sequence of at least one gene encoding a YIPF-related protein; and in which all other relevant features were the same (e.g. plant species, growing conditions, method of processing tobacco, etc.). In one embodiment an unmodified plant is one which does not comprise a gene encoding YIPF-related protein whose activity or expression has been modulated. In one embodiment, an unmodified plant is one which does not comprise a modified nucleic acid sequence which encodes at least one gene encoding a YIPF-related protein.

YIPF-Related Protein

[0088] A “YIPF-related protein” as used herein refers to a protein which comprises a YPT/RAB GTPase Interacting Protein domain. Suitably, the YIPF-related protein comprises a YIP1 domain. An illustrative sequence of a YIPF-related protein from tobacco is shown in SEQ ID No. 3. Amino acids 112-269 of SEQ ID No. 3 relate to a YIP1 domain. Amino acids 123-148, 168-188, 200-218 and 224-242 of SEQ ID No. 3 relate to transmembrane alpha helices.

[0089] A YIPF-related protein may be identified by comparing the amino acid sequence of the protein in question to the amino acid sequence of SEQ ID No. 3. A YIPF-related protein may be identified by comparing the amino acid sequence of the protein in question to amino acids 112-269 of SEQ ID No. 3.

[0090] A YIPF-related protein as referred to herein comprises a YIP1 domain. In one aspect, a YIP1 domain in accordance with the present invention comprises or consists of an amino acid sequence set forth in amino acids 112-269 of SEQ ID No. 3 or a sequence with at least 80% identity thereto; or a functional fragment thereof which has YIP1 activity.

[0091] Suitably, a YIPF-related protein may refer to a sequence set forth in amino acids 112-269 of SEQ ID No. 3 or a sequence which has at least 80% identity thereto. Suitably, a YIPF-related protein as used herein may refer to

a sequence which corresponds to amino acids 112-269 of SEQ ID No. 3 when aligned with SEQ ID No. 3.

[0092] Domains within the amino acid sequence of a protein may be identified using domain prediction software known in the art. Domains are also described in protein databases such as UniprotKB.

[0093] Without wishing to be bound by theory, it is hypothesized that modulating content of a YIPF-related protein in a plant cell or modulating activity, of a YIPF-related protein in a plant would alter the metabolic pathways producing alkaloids and TSNA precursors, resulting in modulated alkaloid and/or TSNA precursor content.

[0094] In one embodiment, YIPF-related protein comprises an amino acid sequence shown as SEQ ID No. 3, or a sequence which has at least 80% identity thereto, or a homologue thereof. Suitably, a homologue of SEQ ID No. 3 may be selected from the group comprising the amino acid sequences provided in Table 1 or a sequence which has at least 80% identity thereto. In one embodiment, a YIPF-related protein comprises an amino acid sequence shown as SEQ ID No. 3, or a sequence which has at least 80% identity thereto (preferably at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identity thereto). In one embodiment a YIPF-related protein comprises an amino acid sequence shown in Table 1, or a sequence which has at least 80% identity thereto (preferably at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identity thereto).

[0095] In one embodiment, the YIPF-related protein according to the present invention comprises or consists of an amino acid shown as SEQ ID No. 3. In one embodiment, the YIPF-related protein according to the present invention comprises or consists of an amino acid shown in Table 1. Suitably, the protein may be from *Nicotiana tabacum*.

[0096] In one embodiment, a YIPF-related protein is encoded by a polynucleotide sequence wherein the gene (prior to mutation) comprises a polynucleotide sequence shown as SEQ ID No. 1, or a sequence which has at least 80% identity thereto, or a homologue thereof. Suitably, a homologue of SEQ ID No. 1 may be selected from the group comprising the polynucleotide sequences provided in Table 1 or a sequence which has at least 80% identity thereto.

[0097] In one embodiment, a YIPF-related protein is encoded by a polynucleotide sequence shown as SEQ ID No. 1, or a sequence which has at least 80% identity thereto (preferably at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identity thereto). In one embodiment a YIPF-related protein is encoded by a polynucleotide sequence shown in Table 1, or a sequence which has at least 80% identity thereto (preferably at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identity thereto).

[0098] In one embodiment, the YIPF-related protein is encoded by a polynucleotide sequence which comprises or consists of a polynucleotide sequence shown as SEQ ID No. 1. In one embodiment, the YIPF-related protein is encoded by a polynucleotide sequence which comprises or consists of a polynucleotide sequence shown in Table 1.

[0099] In one embodiment, a YIPF-related protein is encoded by a polynucleotide sequence wherein the gene (prior to mutation) comprises a polynucleotide sequence shown as SEQ ID No. 2, or a sequence which has at least 80% identity thereto, or a homologue thereof. Suitably, a homologue of SEQ ID No. 2 may be selected from the group

comprising the polynucleotide sequences provided in Table 1 or a sequence which has at least 80% identity thereto.

[0100] In one embodiment, a YIPF-related protein is encoded by a polynucleotide sequence shown as SEQ ID No. 2, or a sequence which has at least 80% identity thereto (preferably at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identity thereto). In one embodiment a YIPF-related protein is encoded by a polynucleotide sequence shown in Table 1, or a sequence which has at least 80% identity thereto (preferably at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identity thereto). In one embodiment, the YIPF-related protein is encoded by a polynucleotide sequence which comprises or consists of a polynucleotide sequence shown as SEQ ID No. 2. In one embodiment, the YIPF-related protein is encoded by a polynucleotide sequence which comprises or consists of a polynucleotide sequence shown in Table 1.

TABLE 1

Homologues of SEQ ID No. 1, 2 and 3
Homologue genomic sequence (SEQ ID NO: 4): GAAAAAACCAAAACCGAAGAAAAAAGGAGAAAAAGAAATTA AAATAACTAAGGAACAAATAGAAAGAGGAGGATCAGATCTGT GTCCGAAACTGAGCTGAAAGAAATAAATTTGCATATACAGTTTAT ACAGAGAAGAAACGAAGATGATGCCAGCAATTACACCCCAATTG ATAATCAGAAATGTTCTCGATCTGTTCTCTGTAATCTTTCTCAT TTTTTTTGGTTTGGATTGAATTTATTTCTGGCCGTCGTTTCAGAT CTAATTGTGTAATTTCTTATATATTATAGGCAGTCGCAGATCCAC CTGGCCAGTTGCGCTCAAATTCAGTGGTAATGCTCTCAGATTTT CGACAAATCATACAGTAGTATTTTGTGGATCATTTTGAATTGG TTTGTGAAATTTTGGGTTTATTTTGTGTTTGCAGATTCGAATTT GCAGACATTTCCACCTTCCAGTTCAGGGGAAAAATCTCTGGTGC TTCTGGACCTCCACGTGATGCTGACGGTATCTCTCTCAACTTAT TCATATATAGTTTACGACGTAAATAGTGTGTACTATTAACACG TGTGGTTTTAGCTAAATGGAGTTAGTGCAGCTTTTAAAGTTGT CTTTAGATCTTTGAAGTTTCATGATACAACTTGTCTCTGTAATA GTGATATAAACACACCAATGGTAGTGCTTAAATTTGCGTTATGAA GGGCTGTCCAGCATGTTTCAGCATTTTTTATTTTCACTAAGCTGA TGTATAAATCTTACACACAGCTAATATGCTATGGCTCTTTTCAA TTTGAAATTTTCTTCAAGGCTACAAACCTTAATCTTGAATTC AATATACATTTCTCTACAATGATTTTGAATTCAAATCTGATACT ATATGAGGAGTAGGATCATAAGAAAGAGATGTTTCTCCAGTAC AAAATTTGGAAGATGAATAAAATTTTGTGTAACATATAAGTCAA AATGAGGTGAGGAAGAACCATTTTGCAGGGTAAAATAAATACA AGGAAGTCCAGGATGGGTTTCGAGGGATTACAATAATGTGGTTA GTTTAGTCTAGGTATACCTAGTTCCGTACAGTTGTCTCATCCG GTGCGTCTTTTGATAGGGTAAAGTTTGTGTGATGCGAACAGCAA GTGCTGCTTAGTTGTATATTTAAAAAGCATGTGCATTACAAGTC ATGTAATGAACAGAGGAAGTATTTTCCACAGCCAGTTGCACCAA AAACCTAAGCAAGATATATAATTTGTGTTTGTGATTCACAGTA TCTTTCTTCTCTTAAAAAATTTGCACGTTTCACTCTGGTGCTTCT CATAGAGTTATTTTGTATCTCTAAAAAGAGTTAACTCTCAAAGC TTAACCTGAAGGAAGTTGTGAGTCTGCTATTGCGTCTTTTCATT AAGTTATTAGCAGATCTATTTTCCCTCTCCACTTTTCCACGCTCA GGTGAAATTTACTTCTCTTTTATGATAATAATTATGACTTCTGT GTAGATACATTTCTTAAACAGTATCTGGTCTGTATGAGCCCCA GCAGAGCGGTTGGTTTCGCGCTTCTACTATTGCTGCGTACAAGCC CTATTTGTAGTTTATGCTCGTGTCTTAATCCGAGGATAAATCTA TTCACTCTTTCTTTCACGGGATCCTTTTCGGAGAAGACTGCCAA TAAACAGATTTGTATGCTCCCTGCTCTTTTATTTCTATGTTTAT TGAATTAGCATTTATGCTCGTGTCTTAATCCGAGGATAAATCTA ATAAAACTTATTATGAGTAGTTATTTCTGGAAGTATAGGCCCTT TGGGCTTCTACTGTTATTAATGGCATTATATAAATCTTATCCCA TTTCACTTCTATCAATTTCTGCGCTGTTTTTCCCGTTTACACAA GATGAATAAAGTTGGCAGTTATAGGAGACCAGAAATATATCTTGC TCCCACTAACACTTTCTTCTTCTTGAATGTTATGTTCTAATTTT TTGTTATGATTTTGGATGAGATGACGTTAATTAATATCCCTGTC TATCAATTTTCATCTCTCAATATTCTCATGATCAGTATTTAGTCT CTCGACTAAGAGGCATTTACCTGCTGGGAGCTTGCCAAATGAGGCA ATGAGCTACCAATGCTTAACCTGTTTGAATTTGAATAGTGAATTT TGTGGAACCTTCCATGCATTAATGACGCTGTAATGGGTTGTG

TABLE 1-continued

Homologues of SEQ ID No. 1, 2 and 3
TAAATTGAGCTAATGGATATGGAACAATTTCTACATTTAACTAG ATGGAGTTTGTTCAAATCTCTGGCGGTATATTTTGAAATTAACC ATTAGGCATACAGAAGAACCAATACGAAAAATTTGGGTTGGTATC TTTGTCAATTTTCGAGTTGGAATTTGTCCTGCGCATTTGTAAG CTAGTGAGTTTGTGCAAGTCATCAGATTTCCCTTCTAGAAATTA TGTGCTTTCAATTTGATTTCTTCTCTGTCCTATATCTATCTGC TATTATATAGGATTTGTTAGACATTTCCGTTTACACTTTCAGAACCT GTTATTTCTTAAGTTTCTGTTCCAAATTTGTGATTAATTTTCTTT GGGGTGGGGTGGGGGTGTAATCATTACGTAATCATTCTCTCT TGTTTCATCTTCTTATGATGTTCTCTGTGTTACGTATGACCGTG ATATAAAGCACATGTTTGTATTGTCAGCGGCAGAGCTCATATA ATGGTGATTATTGGCATATTTGTTTCATGAAAAGTTATTTTCAGTT GTTGGGCTTATGATACATTATGACTGCTTTTAAAGTCTTGTCTCT TTTTAACTTGATTGACACTTGGTTTGGTTTGTGAGTTGGAACCT GGCATATGAAATTTCAACTGTACCTTCAAGGACAAAAAGTAAT AATCATCAACCACTACTAAAAAGAAATTTGAAAAAGAAAGAAC GTTAGTAGCAAGTGAAGTTTGGCCACTGCTGCCCCCTCTGTACATC TTTAGCCAGCTTTACATGCTCTCTGAGTAAGAAGAAGCTTGTTT AGAACTTGGCTGCAGGAAGTTTGTCTAAGGTTGCAGTCGACGA ACAGATATGATGTCGCAACGTTGCATCTAACTTAACTATTG CTAGATGCTTATGACGCCATTATCTGACTGGAATGGTGTCTAGAC TTCTTTGTCCCTTAAGATGTCATGAGTGTATGTTTCAATTTATTT CATATTAGCGAACTTCTCTACCGTAATTAATCAATTTTCTTTT CAGGTATGGCCCGTTCTGGATATGCAGTACCTTGGTATTTTGTG AGCTTCCATTGGCAGATTTGTTACGTATATAGCACACAAGCTACA GAAAAAGGAGTGGAACTACGACATCAACCTTGTCACTTGGTCTGC GGGTTTATTTTATGGTTATTCACATTTGTTCTCTTTTGTCTGTT TCTAGTCTCAAGTACTTCTCTGCACCATCTGGCCTCGTTTCACT GCTGTGCTTTACGGCTATTTCCCTTTTCTGCTTCACTCCGGCATT GGTAAGTTTATTTAATGTCAGTATACCTTGTGAACTTTTGGGAA AGCTGTTGCTTGTATGAATCTTTCACTATCATTTTAAAGTTTGC ATTAGGATCCATACTAAGATCTATGATTACCTGCTTACAGTGTCT GTCTCTGTTTCCCTGGGAGATAGTGAGATGGGTAGTTGTCTGTGT AGCTGGCTTCATGTCTGCCACATTTTGGGCTCTTAATCTGAAACA CCACATAGCATCGGCTGGTGAAGGTTGGTTCTTCAATTTGGGCTGC TATCTTCTGTTGCAACTTGCACCTTGTCTTGGTACTGAGGGTTTA CTGTGTTCAACGTCACAGTATACTACAAATTTATTTGAATGTGAC TAGGCACATGTCTATATAATCAGAGAAAGGACACTGTTCTTCGT ATCAACTTGAAGTGAAGTGTGCTTGGTCAATGAAACGGAAGTGA TTCATTCAATTTTGGACGATATACAATAATATAGTTTTCATCAAAAG GTTCTCAGCTTGTGCTCTTTTATGAATGTGCACTTTTGGAGC TTTAATAGTGGTCTGCGAGTTCCAGGGGTCGATGATTTACTCAAA CATTTGTGTTTGTAGTGGGATTTAATTTGTAATGTCAAGATTTGGTTA GCAAAATCTTACAAGTCCACTTAGAAGATAATCAAAACCATGCGGA AATACTTTTGTCTGCCATGATGACCTGAATGACCTAATTT
Homologue coding sequence (SEQ ID NO: 5): ATGATGTCCAGCAATTACACGCAATTGATAATCAGAATGTTTCT GGATCTGTTCTGTCAGTCGAGATCCACCTGGCCAGTTGCGGTC AAATTCAGTGATTGAAATTTGCAGACATTTCCACCTCCAGTTTCC CAGGGGAAAAATCTCTGGTGTCTTGGACCTCCACGTGATGCTGAC GATACATTTCTTAAACAGTATCTGGTCTGATGAGCCCCAGCAG AGCGGTTGGTTTCGCGCTTCTACTATTGCTGCGTACAAGCCCTAT TTTGATGTTGATACCACTGATGTTCTAGAGAGAATTAAGATTCA CTCCTTCCCTTTCACGGGATCCTTTTCGGAGAAGACTGCCAAATAA CCAGATTTGTATGGCCCGTTCTGGATATGCAGTACCTTGGTATTT GTTGCAGCTTCCATTGGCCACATTTGTTACGTATATAGCACACAAG CTCAGAAAAAGGAGTGGAACTACGACATCAACCTTGTCACTTGG CTCGCGGGTTTATTTATGGTTATTTCCACTATTTGTTCTCTTGG TTGTTTCTAGTCTCAAGTACTTCTCTGACCATCTGGCCCTGTT CAGCTGCTGTGCTTTACGGCTATTTCCCTTTTCTGCTTCAATCCG GCATTGTGCTGTCTCTTGTCTCCCTGGGAGATAGTGAGATGGGTA GTTGTGTTGTAGTGGCTTCATGTCTGCCACATTTTGGGCTCTT AATCTGAAACACCATAGCATCGGCTGGTGAAGGTTGGTCTTCTC ATTGTGGTGTCTATCTCTGTTGCAACTTGCACCTTGTCTTGGTA CTGAGGGTTTACTTGTTCACAGTCACAGTATAA

TABLE 1-continued

Homologues of SEQ ID No. 1, 2 and 3
Homologueaminoacidsequence (SEQIDNO:6): MMSNYTAIDNQNVSGSVPAVADPPGQVAVKFTDSNLQTFPPSSS QGKISGASGPPRADDTFSPKPVSGSDEPQSGWFRAFTIAAYKPY FDVDTTDVLERIKDSLPPFTGFSSEKTANKPDLYGPFWICSTLVF VAASIGTFVTYIAHKLQKKEWNYDINLVTSAGLFYGYSTIVPLC LFLVLKYFSAPSGLVQLLCLYGSLEVFPALCLSLVPWEIVRWV VAGVAGEMSATFVALNLKHHSASAGERWFFIVAAIFLLQLALALV LRVYLENTVT

[0101] Suitably, the protein for use according to the present invention may be encoded by a polynucleotide sequence from *Nicotiana tabacum*.

[0102] In one aspect the present invention provides a method of decreasing the alkaloid content of a plant or part thereof or cell (e.g. plant cell), the method comprising modifying said plant by decreasing or inhibiting the activity or expression of at least one YIPF-related protein.

[0103] In one aspect the present invention provides a method of decreasing the alkaloid content of a plant or part thereof or plant cell, the method comprising modifying said plant by decreasing or inhibiting the activity or expression of at least one YIPF-related protein comprising the amino acid sequence shown as SEQ ID No. 3, or a sequence which has at least 80% identity thereto or wherein the at least one gene encoding a YIPF-related protein comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2.

[0104] In one aspect the present invention provides a method of decreasing the content of a tobacco specific nitrosamine (TSNA) precursor in a plant or part thereof (e.g. leaf), the method comprising modifying said plant by decreasing or inhibiting the activity or expression of at least one YIPF-related protein. Suitably, the method comprising modifying said plant by decreasing or inhibiting the activity or expression of at least one YIPF-related protein comprising the amino acid sequence shown as SEQ ID No. 3, or a sequence which has at least 80% identity thereto or wherein the at least one gene encoding a YIPF-related protein comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2. In one aspect the present invention provides a method of decreasing the content of a tobacco specific nitrosamine (TSNA) precursor in a plant or part thereof (e.g. leaf), the method comprising modifying said plant by decreasing or inhibiting the activity or expression of at least one YIPF-related protein.

[0105] In one aspect the present invention provides a method of decreasing the content of a TSNA in a processed leaf, such as a cured leaf, the method comprising:

[0106] modifying a plant by decreasing or inhibiting the activity or expression of at least one YIPF-related protein;

[0107] harvesting a leaf from said plant;

[0108] and processing e.g. curing said harvested leaf.

[0109] Suitably, the method of decreasing the content of a TSNA in a processed leaf may comprise: modifying said plant by decreasing or inhibiting the activity or expression of

at least one YIPF-related protein comprising the amino acid sequence shown as SEQ ID No. 3, or a sequence which has at least 80% identity thereto or wherein the at least one gene encoding a YIPF-related protein comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2. The term “decreasing” or “inhibiting” (e.g. inhibiting the activity or expression of a YIPF-related protein) as used herein means that the activity or expression of the gene encoding the YIPF-related protein is lower or decreased compared with the activity or expression of the gene in a comparable product.

[0110] In one aspect, the present invention provides a method of increasing the alkaloid content of a plant or part thereof or cell (e.g. plant cell), the method comprising modifying said plant by increasing or enhancing the activity or expression of at least one gene encoding a YIPF-related protein.

[0111] In one aspect, the present invention provides a method of increasing the alkaloid content of a plant or part thereof or plant cell, the method comprising modifying said plant by increasing or enhancing the activity or expression of at least one YIPF-related protein comprising the amino acid sequence shown as SEQ ID No. 3, or a sequence which has at least 80% identity thereto or wherein the at least one gene encoding a YIPF-related protein comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2. In one aspect the present invention provides a method of increasing the content of a tobacco specific nitrosamine (TSNA) precursor in a plant or part thereof (e.g. leaf), the method comprising modifying said plant by increasing or enhancing the activity or expression of at least one YIPF-related protein.

[0112] In one aspect the present invention provides a method of increasing the content of a tobacco specific nitrosamine (TSNA) precursor in a plant or part thereof (e.g. leaf), the method comprising modifying said plant by increasing or enhancing the activity or expression of at least one YIPF-related protein comprising the amino acid sequence shown as SEQ ID No. 3, or a sequence which has at least 80% identity thereto, or wherein the at least one gene encoding a YIPF-related protein comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2.

[0113] The term “increasing” or “enhancing” (e.g. increasing the activity or expression of gene encoding YIPF-related protein) as used herein means that the activity or expression of the gene encoding the YIPF-related protein is higher or increased compared with the activity or expression of the gene in a comparable product.

[0114] According to the present invention, the activity or expression of a YIPF-related protein is modulated.

[0115] In one aspect the present invention provides a method of modulating (i.e. increasing or decreasing) the alkaloid content of a plant or part thereof or cell (e.g. plant cell), the method comprising modifying said plant by modulating (i.e. increasing or decreasing) the activity of at least one YIPF-related protein.

[0116] The term “activity” refers to any functionality of the YIPF-related protein.

[0117] For example the activity of the YIPF-related protein may refer to its ability to interact with Ypt family small GTPases. Methods for measuring protein: protein interactions are known in the art.

[0118] The activity of the YIPF-related protein may relate to its role in ER to Golgi and intra-Golgi transport. Thus modifying e.g. decreasing the activity of the YIPF-related protein may abrogate ER to Golgi and intra-Golgi transport. Without wishing to be bound by theory, immune process such as nicotine production in plants are tightly regulated by subcellular trafficking which may be modified by modulation of the YIPF-related protein according to the present invention.

[0119] Alternatively, the activity of a YIPF protein may be abrogated by disrupting or deleting at least part of a transmembrane domain. Disrupting or deleting at least part of a transmembrane domain may affect the cellular localisation of the YIPF-related protein.

[0120] Modulation of the activity of a YIPF-related protein may entail increasing or decreasing the activity of the YIPF-related protein.

[0121] Increasing the activity of a YIPF-related protein refers to enhancing or improving the ability of the YIPF-related protein to carry out a particular function in comparison to a YIPF-related protein in a plant that has not been modified in accordance with the invention. Decreasing the activity of a YIPF-related protein refers to reducing, inhibiting or disrupting the ability of the YIPF-related protein to carry out a particular function in comparison to a YIPF-related protein in a plant that has not been modified in accordance with the invention. The activity of a YIPF-related protein may be reduced to such an extent that the activity is prevented or eliminated. In some embodiments the activity of a YIPF-related protein be modulated (i.e. increased or decreased) by at least about 10% 20% 30%, or 40%, suitably at least about 50%, 60%, 70%, more suitably at least about 80%, 90%, 95% or 100% in comparison to the activity of a gene encoding a YIPF-related protein in a plant (e.g. a tobacco plant) which has not been modified in accordance with the present invention. Suitably, the activity may be decreased.

[0122] In some embodiments the modulated YIPF-related protein exhibits increased or decreased activity compared to an unmodified YIPF-related protein. The modulated YIPF-related protein may exhibit at least about 1%, at least about 3%, at least about 5%, at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80% or at least about 90% increased or decreased activity compared to an unmodified YIPF-related protein.

[0123] In one aspect the present invention provides a method of modulating (i.e. increasing or decreasing) the alkaloid content of a plant or part thereof or cell (e.g. plant cell), the method comprising modifying said plant by modulating (i.e. increasing or decreasing) the expression of at least one YIPF-related protein.

[0124] The “expression” of a gene refers to the degree to which the information encoded in the gene is converted to a functionality. The level of expression of a gene may be equated with the amount of the product of that gene present in a cell or organism. A modification that modulates (i.e. increases or decreases) the expression of a gene is one that

increases the amount of the product of that gene in a plant or cell in comparison to an unmodified plant or cell.

[0125] In some embodiments the expression of a YIPF-related protein gene is modulated (i.e. increased or decreased) in comparison to the expression of a gene encoding a YIPF-related protein in a plant (e.g. a tobacco plant) which has not been modified in accordance with the present invention.

[0126] In some embodiments the expression of a YIPF-related protein gene may be modulated (i.e. increased or decreased) by at least about 10% 20% 30%, or 40%, suitably at least about 50%, 60%, 70%, more suitably at least about 80%, 90%, 95% or 100% in comparison to the expression of a gene encoding a YIPF-related protein in a plant (e.g. a tobacco plant) which has not been modified in accordance with the present invention. Suitably, expression may be decreased.

[0127] In some embodiments, the modulated YIPF-related protein exhibits increased or decreased expression compared to an unmodified YIPF-related protein. The modulated YIPF-related protein may exhibit at least about 1%, at least about 3%, at least about 5%, at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80% or at least about 90% increased or decreased expression compared to an unmodified YIPF-related protein. Suitably, expression may be decreased. Typically, genes are transcribed to mRNA, which is translated to protein, the final gene product. Proteins may be sequestered in cellular stores and/or degraded. The expression of a gene may be modulated by modulating any or all of these steps. Accordingly, in some embodiments the modification modulates (e.g. decreases) expression of at least one gene encoding a YIPF-related protein in one of the following ways:

[0128] modulating (e.g. decreasing) transcription from the at least one gene encoding a YIPF-related protein;

[0129] modulating (e.g. decreasing) translation of the mRNA from the at least one gene encoding a YIPF-related protein;

[0130] modulating (e.g. decreasing) release of the YIPF-related protein from intracellular stores;

[0131] modulating (e.g. increasing) the rate of degradation of the YIPF-related protein and/or introducing a mutation which modifies the amino acid sequence of the YIPF-related protein to decrease or increase its activity e.g. to decrease YIPF-related protein activity.

[0132] The expression of specific genes encoding a YIPF-related protein can be measured by measuring transcription and/or translation of the gene. Methods for measuring transcription are well known in the art and include, amongst others, northern blot, RNA-Seq, in situ hybridization, DNA microarrays and RT-PCR. Alternatively, the expression of a gene may be measured indirectly by measuring the level of the gene product for example the protein encoded by said gene. For example, the expression of a YIPF-related protein may be determined by measuring the presence of the protein using an antibody specific for the YIPF-related protein by western blot.

Modifying

[0133] The plant or cell may be modified in any way that modulates activity or expression of at least one YIPF-related protein. Types of modifications to plants and cells that modulate activity or expression of genes, as well as tech-

niques to achieve those modifications, are known in the art. In some embodiments the present invention provides a method of decreasing the alkaloid content of a plant or part thereof or cell (e.g. plant cell), the method comprising modifying said plant by decreasing or inhibiting the activity or expression of at least one YIPF-related protein as described herein.

[0134] In some embodiments the present invention provides a method of decreasing the content of a tobacco specific nitrosamine (TSNA) or a precursor of a TSNA in a tobacco plant or plant part thereof, the method comprising modifying said plant or a cell culture by decreasing the activity or expression of at least one YIPF-related protein as described herein.

[0135] Any method known in the art for decreasing or inhibiting the activity or expression of a gene or protein may be used in the methods according to the present invention.

[0136] Suitably, the activity or expression of the gene encoding a YIPF-related protein may be reduced, partly inactivated, inhibited, eliminated, knocked out or lost such that the protein activity, expression or function of the gene encoding a YIPF-related protein may not be detectable.

[0137] In one aspect, the at least one gene encoding a YIPF-related protein is knocked out. In other words, the gene encoding a YIPF-related protein has been rendered completely inoperative. By way of example, the present method may comprise:

[0138] providing a mutation in a nucleic acid sequence which encodes a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto;

[0139] providing a mutation in a regulatory region (e.g. a promoter or an enhancer) which contributes to controlling the expression of a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto;

[0140] providing an antisense RNA, siRNA or miRNA which reduces the level of nucleic acid sequence encoding a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto.

[0141] Each of the above approaches results in the reduction or prevention of activity or expression of a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto or wherein the at least one gene encoding a YIPF-related protein comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2.

[0142] As used herein, the term “mutation” encompasses a natural genetic variant or an engineered variant. In particular, the term “mutation” refers to a variation in the nucleotide sequence encoding the amino acid sequence or in the amino acid sequence compared to the sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% (preferably at least 85%, preferably at least 90%, preferably at least 93%, preferably at least 95%, preferably at least 98%, preferably at least 99%) sequence identity thereto.

[0143] In one embodiment the mutation decreases the alkaloid content of a plant. In another embodiment, the mutation decreases the content of at least one TSNA precursor in a plant or part thereof, or leaf such as a harvested or processed leaf. In one embodiment the mutation decreases the content of one or more TSNA selected from NNN, NNK, NAT, NAB, preferably NNN and/or NNK content is decreased in a processed leaf. Suitably, the TSNA content is reduced in relation to a comparable product.

[0144] In one embodiment, a method according to the present invention may comprise providing a nucleic acid sequence to a plant or part thereof or plant cell, wherein said nucleic acid results in the reduction or elimination of the activity or expression of at least one YIPF-related protein.

[0145] In one embodiment, a method according to the present invention may comprise providing a nucleic acid sequence to a plant or part thereof or plant cell, wherein said nucleic acid results in the modification of the nucleic acid sequence of at least one YIPF-related protein.

[0146] Suitably said nucleic acid sequence may be introduced to the plant or part thereof or cell. Suitably an endogenous nucleic acid sequence in the plant or part thereof or cell may be modified to encode the polypeptide according to the present invention (e.g. by gene editing). For example, an endogenous nucleotide sequence may be modified to decrease the activity or expression of at least one YIPF-related protein.

[0147] In a preferred embodiment, each copy of a nucleic acid sequence encoding a protein comprising a sequence shown as SEQ ID No. 3, or a sequence which has at least 80% sequence identity thereto or wherein the at least one gene encoding a YIPF-related protein comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2, which is present in the plant is modified e.g. mutated as defined herein (e.g. each genomic copy of a gene encoding said protein in a plant is mutated). For example, each copy of the gene in the allotetraploid genome of *Nicotiana tabacum* may be mutated. In a preferred embodiment, some or all of the homologues of the YIPF-related protein as described herein are modified e.g. inhibited or mutated. Suitably, some or all of the homologues listed in Table 1, or corresponding sequences which have at least 80% sequence identity thereto are modified e.g. inhibited or mutated.

[0148] In some embodiments the plant or plant cell according to the present invention is homozygous. Suitably, the plant or plant cell may be homozygous for the modification e.g. inhibition or mutation. In some embodiments the plant or plant cell according to the present invention expresses only the modified e.g. mutated nucleic acid encoding at least one YIPF-related protein. In other words, in some embodiments no endogenous (or endogenous and functional protein) is present in the plant according to the present invention. In other words, if any endogenous protein is present it is preferably in an inactive form.

[0149] In one embodiment the present method may comprise providing a mutation in the nucleic acid sequence shown as SEQ ID No. 1, or 2, or a nucleic acid sequence which has at least 80% identity thereto, or a homologue of SEQ ID No. 1 or SEQ ID No. 2.

[0150] The mutation may alter the plant genome such that a nucleic acid sequence encoding a protein comprising the

amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto, or a homologue of SEQ ID No. 3, is completely or partially deleted or otherwise modified to inhibit or eliminate the activity of the YIPF-related protein. In some embodiments the mutation does not alter the level or expression of the protein but reduces inhibits or eliminates the activity of the YIPF-related protein.

[0151] Suitably, at least one mutation may be in a non-cytoplasmic domain, a cytoplasmic domain or a transmembrane domain of the YIPF-related protein.

[0152] The mutation may interrupt the nucleic acid sequence which encodes a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto, or a homologue of SEQ ID No. 3.

[0153] The interruption may cause the nucleic acid sequence to not be transcribed and/or translated. The nucleic acid sequence may be interrupted, for example, by deleting or otherwise modifying the ATG start codon of the nucleic acid sequence such that translation of the protein is reduced or prevented.

[0154] The nucleic acid sequence may comprise one or more nucleotide change(s) that reduce or prevent expression of the protein or affect protein trafficking. For example, expression of the protein may be reduced or prevented by introduction of one or more pre-mature stop codons, a frame shift, a splice mutation or a non-tolerated amino acid substitution in the open reading frame. A premature stop codon refers to a mutation which introduces a stop codon into the open reading frame and prevents translation of the entire amino acid sequence. The premature stop codon may be a TAG (“amber”), TAA (“ochre”), or TGA (“opal” or “umber”) codon.

[0155] A frame-shift mutation (also called a framing error or a reading frame shift) is a mutation caused by indels (insertions or deletions) of a number of nucleotides in a nucleic acid sequence that is not divisible by three. Due to the triplet nature of gene expression by codons, the insertion or deletion can change the reading frame, resulting in a completely different translation from the original. A frame-shift mutation will often cause the reading of the codons after the mutation to code for different amino acids. The frameshift mutation will commonly result in the introduction of a premature stop codon.

[0156] A splice mutation inserts, deletes or changes a number of nucleotides in the specific site at which splicing takes place during the processing of precursor messenger RNA into mature messenger RNA. The deletion of the splicing site results in one or more introns remaining in mature mRNA and may lead to the production of abnormal proteins.

[0157] A non-tolerated amino acid substitution refers to a mutation which causes a non-synonymous amino acid substitution in the protein which results in reduced or ablated function of the protein.

[0158] Any method known in the art for providing a mutation in a nucleic acid sequence may be used in the method according to the present invention. For example, homologous recombination may be used, in which a vector is created in which the relevant nucleic acid sequence(s) are mutated and used to transform plants or plant cells. Recombinant plants or plant cells expressing the mutated sequence may then be selected.

[0159] In one embodiment the mutation introduces a non-tolerated amino acid substitution in a protein comprising an amino acid sequence shown as SEQ ID No. 3, or a sequence which has at least 80% sequence identity thereto, or a homologue of SEQ ID No. 3.

[0160] In some embodiments, the YIPF-related protein may contain a mutation which decreases the expression or activity of the YIPF-related protein.

[0161] The mutation may be a deletion, a splice mutant or codon encoding a non-tolerated amino acid substitution.

[0162] In one embodiment, the nucleic acid sequence encoding the YIPF-related protein may be wholly or partially deleted. The deletion may be continuous, or may comprise a plurality of sections of sequence. The deletion preferably removes a sufficient amount of nucleotide sequence such that the nucleic acid sequence no longer encodes a functional YIPF-related protein. The deletion may be total, in which case 100% of the coding portion of the nucleic acid sequence is absent, when compared to the corresponding genome of a comparable unmodified plant. The deletion may, for example, remove at least 50, 60, 70, 80 or 90% of the coding portion of the nucleic acid sequence.

[0163] Suitably, at least part of the protein may be deleted. The deletion may, for example, remove at least 10, 20, 30, 40, 50, 60, 70, 80 or 90% of the coding portion of the protein.

[0164] The deletion may remove at least 10 amino acids (such as at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90 amino acids) from the YIPF-related protein. Suitably, the deletion may remove at least 10 amino acids (such as at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90 amino acids) of the YIPF-related protein wherein the sequence of the YIPF-related protein is aligned with SEQ ID No. 3. Suitably, the deletion may remove at least 10 amino acids (such as at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90 amino acids) from the YIPF-related protein wherein the YIPF-related protein prior to deletion comprises an amino acid sequence set forth in SEQ ID No. 3, or a sequence which has at least 80% sequence identity thereto, or a homologue of SEQ ID No. 3. Suitably, the YIPF-related protein prior to deletion comprises an amino acid sequence set forth in SEQ ID No. 3.

[0165] Suitably, the protein for use according to the present invention may comprise a truncated YIPF-related protein. Suitably, the truncated protein may be a truncated version of an amino acid sequence set forth in SEQ ID No. 3, or a sequence which has at least 80% sequence identity thereto, or a homologue of SEQ ID No. 3. Suitably, the truncated protein lacks at least 10 amino acids (such as at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90 amino acids, at least 100 amino acids, at least 110 amino acids) from the YIPF-related protein.

[0166] The deletion may remove at least part of a domain. The deletion may for example, remove at least 10, 20, 30, 40, 50, 60, 70, 80 or 90% of a domain. Suitably, the deletion may remove at least amino acids, at least 10 amino acids, at least 15, at least 20, at least 25, at least 30 amino acids, at least 40 amino acids, at least 50 amino acids, at least 60 amino acids, at least 70 amino acids, at least 80 amino acids of a domain. Suitably, the deletion may remove 5 amino acids, 10 amino acids, 15, 20 amino acids, 25 amino acids,

30 amino acids, 40 amino acids, 50 amino acids, 60 amino acids, 70 amino acids, 80 amino acids of a domain.

[0167] The deletion may remove at least part of a domain. The deletion may, for example, remove at least one or at least two amino acids from a domain. Suitably, a domain may be completely deleted.

[0168] Methods for deletion of nucleic acid sequences in plants are known in the art. For example, homologous recombination may be used, in which a vector is created in which the relevant nucleic acid sequence(s) are missing and used to transform plants or plant cells. Recombinant plants or plant cells expressing the new portion of sequence may then be selected.

[0169] Plant cells transformed with a vector as described herein may be grown and maintained in accordance with well-known tissue culturing methods such as by culturing the cells in a suitable culture medium supplied with the necessary growth factors such as amino acids, plant hormones, vitamins, etc.

[0170] Modification of the nucleic acid sequence may be performed using targeted mutagenesis methods (also referred to as targeted nucleotide exchange (TNE) or oligo-directed mutagenesis (ODM)). Targeted mutagenesis methods include, without limitation, those employing zinc finger nucleases, TALENs (see WO2011/072246 and WO2010/079430), Cas9-like, Cas9/crRNA/tracrRNA, Cas9/gRNA, or other CRISPR systems (see WO 2014/071006 and WO2014/093622), meganucleases (see WO2007/047859 and WO2009/059195), or targeted mutagenesis methods employing mutagenic oligonucleotides, possibly containing chemically modified nucleotides for enhancing mutagenesis with sequence complementarity to the gene, into plant protoplasts (e.g., KeyBase® or TALENs).

[0171] Alternatively, mutagenesis systems such as TILLING (Targeting Induced Local Lesions IN Genomics; McCallum et al. (2000) *Nat. Biotech.* 18:455, and McCallum et al. (2000) *Plant Physiol.* 123, 439-442, both incorporated herein by reference) may be used to generate plant lines which comprise a gene encoding a protein having a mutation. TILLING uses traditional chemical mutagenesis (e.g. ethyl methanesulfonate (EMS) mutagenesis, which produces random mutations) followed by high-throughput screening for mutations. Thus, plants, seeds, cells and tissues comprising a gene having the desired mutation may be obtained.

[0172] The method may comprise the steps of mutagenizing plant seeds (e.g. EMS mutagenesis), pooling of plant individuals or DNA, PCR amplification of a region of interest, heteroduplex formation and high-throughput detection, identification of the mutant plant, sequencing of the mutant PCR product. It is understood that other mutagenesis and selection methods may equally be used to generate such modified plants. Seeds may, for example, be radiated or chemically treated and the plants may be screened for a modified phenotype.

[0173] Fast neutron deletion mutagenesis may be used in a reverse genetics sense (i.e. with PCR) to identify plant lines carrying a deletion in the endogenous gene. See for example Ohshima et al. (1998) *Virology* 213:472-481; Okubara et al. (1994) *Genetics* 137:867-874; and Quesada et al. (2000) *Genetics* 154:421-4315 which are incorporated herein by reference.

[0174] In another approach, dominant mutants may be used to trigger RNA silencing due to gene inversion and

recombination of a duplicated gene locus. See for example Kusaba et al. (2003) *Plant Cell* 15:1455-1467 (incorporated herein by reference).

[0175] Modified plants may be distinguished from non-modified plants, i.e., wild type plants, by molecular methods, such as the mutation(s) present in the DNA, and by the modified phenotypic characteristics. The modified plants may be homozygous or heterozygous for the modification. Preferably modified plants are homozygous for the modification.

[0176] In one embodiment the method of reducing or preventing the activity or expression of a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto; or a homologue of SEQ ID No. 3 does not comprise treating the plant with a chemical (e.g. an agro-chemical).

[0177] Other ways of reducing or preventing the expression will be apparent to one skilled in the art and include the use of virus-induced gene silencing (VIGs), micro RNA silencing, RNAi, antisense, tDNA insertions, or dominant negative constructs (or antimorphic mutations).

[0178] In one embodiment the expression of a YIPF-related protein, such as a gene encoding a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto, may be reduced or eliminated by virus-induced gene silencing.

[0179] In one embodiment the expression of a YIPF-related protein, such as a gene encoding a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto, may be reduced or eliminated by microRNAs.

[0180] In one embodiment the expression of a YIPF-related protein, such as a gene encoding a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto, may be reduced or eliminated by RNAi.

[0181] In one embodiment the expression of a YIPF-related protein, such as a gene encoding a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto, may be reduced or eliminated by antisense suppression.

[0182] In one embodiment the expression of a YIPF-related protein, such as a gene encoding a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto, may be reduced or eliminated by sense suppression.

[0183] In one embodiment the expression of a YIPF-related protein, such as a gene encoding a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto, may be reduced or eliminated by tDNA insertions. In one embodiment the expression of a YIPF-related protein, such as a gene encoding a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto, may be reduced or eliminated by dominant negative constructs (or antimorphic mutations).

[0184] In one embodiment the expression of a YIPF-related protein, such as a gene encoding a protein compris-

ing the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto may be reduced or eliminated by a targeted mutagenesis based system.

[0185] In one embodiment the expression of a YIPF-related protein, such as a gene encoding a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto, may be reduced or eliminated by a gene editing e.g. CRISPR based system.

[0186] In one embodiment the expression of a gene encoding a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto may be reduced or eliminated by zinc finger nuclease, TALENs, meganucleases, mutagenic oligonucleotides or TILLING.

[0187] In some embodiments the present invention provides a method of increasing the alkaloid content of a plant or part thereof or cell (e.g. plant cell), the method comprising modifying said plant by increasing or enhancing the activity or expression of at least one YIPF-related protein. Any method known in the art for increasing or enhancing the activity or expression of a gene may be used in the methods according to the present invention.

[0188] In some embodiments the method may comprise overexpressing at least one gene encoding a YIPF-related protein. Suitably the method may comprise expressing one or more additional copies of the at least one gene encoding a YIPF-related protein in the plant or cell. Suitably the method may comprise modifying the endogenous copy of the at least one gene encoding a YIPF-related protein such that its expression is increased. The method may comprise mutating the coding sequence of the at least one gene encoding a YIPF-related protein. The method may comprise mutating a regulatory sequence that regulates expression of the at least one gene encoding a YIPF-related protein.

[0189] Suitably the method may comprise transforming a cell of a plant (e.g. a tobacco plant) with a genetic construct which encodes at least one YIPF-related protein comprising an amino acid sequence as set out in SEQ ID No. 3, or a functional variant or functional fragment or orthologue thereof, or a sequence which has at least 80% identity to SEQ ID No. 3; or a homologue of SEQ ID No. 3; or wherein the at least one gene encoding a YIPF-related protein comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2; or which comprises a nucleotide sequence which encodes a protein which is capable of promoting or augmenting at least one endogenous YIPF-related protein. It will be appreciated that each of these options would result in an increased activity and expression of the polypeptide encoded by the at least one YIPF-related protein. The method may comprise regenerating the plant from the transformed cell. There is provided use of genetic construct which is capable of increasing the activity and/or expression of a polypeptide encoded by at least one gene encoding a YIPF-related protein for increasing the alkaloid content (e.g. nicotine content) in a plant or part thereof or cell transformed with the construct.

[0190] The genetic construct may encode a polypeptide comprising the amino acid SEQ ID No. 3, or a functional variant or functional fragment or orthologue thereof, or a

sequence which has at least 80% identity to SEQ ID No. 3; or comprise a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2.

[0191] In another embodiment, the invention relates to a method of increasing the alkaloid content of a plant or part thereof or a cell, comprising modifying said plant or cell by increasing the activity of at least one YIPF-related protein.

[0192] In one embodiment the activity of at least one gene encoding a YIPF-related protein may be increased by introducing (or providing) a mutation to at least one gene encoding a YIPF-related protein.

[0193] Suitably, the activity of at least one gene encoding a YIPF-related protein may be increased by introducing a mutation to at least one gene encoding a YIPF-related protein which comprises an amino acid sequence as set out in SEQ ID No. 3; or a functional variant or functional fragment or orthologue thereof, or a sequence which has at least 80% identity to SEQ ID No. 3; or wherein the at least one gene encoding a YIPF-related protein comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2. In some embodiments a modification which increases the activity or expression of at least one YIPF-related protein and thereby increases alkaloid content by one of the following: modulating (e.g. increasing) transcription from the at least one gene encoding a YIPF-related protein;

[0194] modulating (e.g. increasing) translation of the mRNA from the at least one gene encoding a YIPF-related protein;

[0195] modulating (e.g. increasing) release of the YIPF-related protein from intracellular stores; and/or modulating (e.g. decreasing) the rate of degradation of the YIPF-related protein.

Alkaloid Content

[0196] In one embodiment the present invention provides a method of modulating the alkaloid content of a plant (e.g. a tobacco plant) or a part thereof, the method comprising modifying said plant by modulating the activity or expression of at least one YIPF-related protein.

[0197] The term “modulating” is used herein to mean either increasing or decreasing.

[0198] The term “increasing alkaloid content” is used herein to mean that the alkaloid content in the product of the present invention (e.g. plant, part thereof (e.g. leaf), processed leaf or a product made from the plant (e.g. a tobacco industry product)) is higher compared with a comparable product which has not been modified in accordance with the present invention.

[0199] The term “decreasing alkaloid content” is used herein to mean that alkaloid content in the product of the present invention (e.g. plant, part thereof (e.g. leaf), processed leaf or a product made from the plant (e.g. a tobacco industry product)) is lower compared with a comparable product which has not be modified in accordance with the present invention.

[0200] In some embodiments, the modulation of alkaloid content refers to an increase in alkaloid content wherein the activity or expression of at least one YIPF-related protein is increased (or example the protein is overexpressed).

[0201] In some embodiments, the modulation of alkaloid content refers to a decrease in alkaloid content wherein the expression of at least one YIPF-related protein is decreased or inhibited or eliminated. In a further aspect, the alkaloid content is measured from leaves. In one aspect the alkaloid content is measured from green leaves. In a further aspect, the alkaloid content is measured from cured leaves, e.g. air-cured, flue-cured, fire-cured or sun-cured leaves. In a further aspect, the alkaloid content is measured from flue-cured leaves. In a further aspect, the alkaloid content is measured from air-cured leaves.

[0202] The term “alkaloid content” is used herein to mean the concentration and/or total amount of the entire group of compounds classified as alkaloids or the concentration and/or total amount of one or more compounds classified as alkaloids. Alkaloids typically present in tobacco include normicotine, PON, anatabine, anabasine, nicotine, and myosmine. In some embodiments the content of one or more alkaloids, such as two or more alkaloids, such as three or more alkaloids, such as four or more alkaloids, such as five or more alkaloids, such as all six alkaloids, selected from nicotine, normicotine, PON, anatabine, anabasine and myosmine is modulated. In some embodiments the content of one or more alkaloids, such as two or more alkaloids, such as three or more alkaloids, such as four or more alkaloids, such as five or more alkaloids, such as all six alkaloids, selected from nicotine, normicotine, PON, anatabine, anabasine and myosmine is increased. In some embodiments the content of one or more alkaloids, such as two or more alkaloids, such as three or more alkaloids, such as four or more alkaloids, such as five or more alkaloids, such as all six alkaloids, selected from nicotine, normicotine, PON, anatabine, anabasine and myosmine is decreased. In some embodiments the total alkaloid content of the plant or cell is modulated. In some embodiments the total alkaloid content is increased. In some embodiments the total alkaloid content is increased.

[0203] Any method known in the art for determining the concentration and/or total content of alkaloids may be used. One preferred method for analysing alkaloid content involves the analysis by gas chromatography-flame ionization detection method (GC-FID) or by reversed phase high performance liquid chromatography with tandem mass spectrometry (LC-MS/MS).

[0204] In one embodiment there is provided a method for producing a plant (e.g. a tobacco plant) or part thereof, a plant propagation material (e.g. a tobacco plant propagation material), a cell (e.g. a tobacco cell), a leaf (e.g. a tobacco leaf), a harvested leaf (e.g. a harvested tobacco leaf), a cut harvested leaf (e.g. a cut harvested tobacco leaf), a processed leaf (e.g. a processed tobacco leaf), a cut and processed leaf (e.g. a cut and processed tobacco leaf), a product comprising said plant or part thereof (e.g. a tobacco industry product) or combinations thereof obtainable or obtained by a plant of the invention which has modulated alkaloid content, the method comprising modifying said plant to modulate the activity or expression of a YIPF-related protein. The modulated alkaloid content may be determined by comparing the alkaloid content in the plant (e.g. tobacco plant) or part thereof, plant propagation material (e.g. tobacco plant propagation material), a cell (e.g. a tobacco cell), leaf (e.g. tobacco leaf), harvested leaf (e.g. a harvested tobacco leaf), cut harvested leaf (e.g. a cut harvested tobacco leaf), processed leaf (e.g. processed tobacco leaf), cut and processed leaf (e.g. cut and

processed tobacco leaf), a product comprising a plant or part thereof of the present invention, e.g. a tobacco industry product, or combinations thereof with a comparable product.

[0205] Suitably the alkaloid content may be modulated in a plant, e.g. a tobacco plant e.g. modified tobacco plant. Suitably the alkaloid content may be modulated in a leaf (e.g. a tobacco leaf e.g. a tobacco leaf from a modified tobacco plant). Suitably the alkaloid content may be modulated in a harvested leaf (e.g. a harvested tobacco leaf from a modified tobacco plant). Suitably the alkaloid content may be modulated in a cut harvested leaf (e.g. a cut harvested tobacco leaf from a modified tobacco plant). Suitably the alkaloid content may be modulated in a processed leaf (e.g. a processed tobacco leaf e.g. a processed tobacco leaf from a modified tobacco plant). Suitably the alkaloid content may be modulated in a cut and processed leaf (e.g. a cut and processed tobacco leaf e.g. a cut and processed tobacco leaf from a modified tobacco plant). Suitably the alkaloid content may be modulated in a cured leaf (e.g. cured a tobacco leaf from a modified tobacco plant). Suitably the alkaloid content may be modulated in an extract of a green leaf (e.g. a green tobacco leaf from a modified tobacco plant). Suitably the alkaloid content may be modulated in a product comprising the plant of the present invention or part thereof (e.g. a tobacco industry product, for example a tobacco industry product produced from a modified tobacco plant or part thereof). Suitably the alkaloid content may be modulated in any one of the above products or combinations thereof. Suitably the modulation of alkaloid content described above may be an increase in alkaloid content. Suitably the modulation of alkaloid content described above may be a decrease in alkaloid content (e.g. a decrease in normicotine and/or PON content). In one embodiment the content of one or more alkaloids selected from normicotine, PON, anatabine and anabasine is decreased. In one embodiment the content of normicotine is decreased. In one embodiment the content of PON is decreased. In one embodiment the content of anatabine is decreased. In one embodiment the content of anabasine is decreased.

[0206] In one embodiment the nicotine content of a modified plant (e.g. tobacco plant), plant propagation material (e.g. tobacco plant propagation material), leaf (e.g. tobacco leaf), harvested leaf (e.g. harvested tobacco leaf), cut harvested leaf (e.g. cut harvested tobacco leaf), processed leaf (e.g. processed tobacco leaf), cut and processed leaf (e.g. cut and processed tobacco leaf) or tobacco industry product from a modified tobacco plant is not substantially decreased. Suitably, the nicotine content is at least 85% (such as at least 90%, such as at least 95%, such as at least 98%, such as at least 99%) of the nicotine content of a comparable product. In one embodiment the alkaloid content of a plant (e.g. tobacco plant) or part thereof may be modulated by at least 0.5, 1.5, 2, 3 or 4 fold when compared to the alkaloid content of a plant (e.g. tobacco plant) or part thereof, respectively, which has not been modified to modulate the activity or expression of at least one gene encoding a YIPF-related protein and which has been grown under similar growth conditions. Suitably the alkaloid content may be modulated by about 0.5 fold to about 4 fold. Suitably the alkaloid content may be modulated by about 4 fold. Suitably the modification may be an increase or a decrease in alkaloid content. Suitably the modulation may be of one or more alkaloids selected from nicotine, normicotine, PON, anatabine, anabasine and myosmine. Suitably the modulation

may be of one or more alkaloids selected from nicotine, nornicotine, PON, anatabine and anabasine. Suitably, the nornicotine content may be reduced. Suitably, the PON content may be reduced. Suitably, the anatabine content may be reduced. Suitably, the anabasine content may be reduced.

[0207] In one embodiment of the invention the alkaloid content of a plant (e.g. a tobacco plant) or part thereof may be modulated by at least 1%, 2%, 5%, 8%, 10%, 12%, 15%, 20%, 25%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 100% in comparison to a plant (e.g. a tobacco plant) or part thereof which has not been modified according to the present invention. In one embodiment the alkaloid content may be modulated by at least 30% in comparison to an unmodified plant or part thereof. In one embodiment the alkaloid content may be modulated by at least 40% in comparison to an unmodified plant or part thereof. In one embodiment the alkaloid content may be modulated by at least 50% in comparison to an unmodified plant or part thereof. In one embodiment the alkaloid content may be modulated by at least 60% in comparison to an unmodified plant or part thereof.

[0208] The modulation may be an increase or a decrease in alkaloid content when compared to an unmodified plant (e.g. a tobacco plant) or part thereof.

[0209] Suitably the modulation may be of total alkaloid content. Suitably the modulation may be of one or more alkaloids selected from nicotine, nornicotine, PON, anatabine, anabasine and myosmine. Suitably the modulation may be of one or more alkaloids selected from nornicotine, nicotine, PON, anatabine and anabasine. Suitably the modulation may be of nornicotine content, such as decrease in nornicotine content. Suitably the modulation may be of anabasine content, such as decrease in anabasine content. Suitably the modulation may be of PON content, such as decrease in PON content. Suitably the modulation may be of anatabine content, such as decrease in anatabine content.

[0210] Suitably the modulation may be of more than one alkaloid, such as two or more alkaloids, such as three or more alkaloids, such as four or more alkaloids, such as five or more alkaloids, such as all six alkaloids, selected from nicotine, nornicotine, PON, anatabine, anabasine and myosmine.

[0211] In some embodiments the alkaloid content of the plant may be modulated by between about 5% and about 100%, by between about 10% and about 90%, by between about 20% and about 80%, by between about 30% and about 70%, by between about 40% and 60%, by between about 40% and 50%, or by between about 50% and 60%.

Tobacco-Specific Nitrosamine (TSNA) Content

[0212] In one embodiment the present invention provides a method of decreasing the content of at least one tobacco-specific nitrosamine (TSNA) precursor in a plant (e.g. a tobacco plant) or a part thereof or in a tobacco cell. Suitably, the method may comprise modifying said plant by modulating the activity or expression of at least one YIPF-related protein. In one embodiment, the present invention provides a method of producing a processed leaf with decreased TSNA content (e.g. relative to a comparable product). The method of producing a processed leaf with decreased TSNA content may comprise:

[0213] modifying a plant by decreasing or inhibiting the activity or expression of at least one YIPF-related protein;

[0214] harvesting a leaf from said plant;

[0215] and processing e.g. curing said harvested leaf.

[0216] The TSNA may be measured in a processed tobacco, e.g. cured tobacco or reconstituted tobacco. In one embodiment the TSNA content is measured and/or modified (e.g. reduced) in a cured tobacco plant or part thereof (e.g. in cured tobacco leaf).

[0217] The term “tobacco-specific nitrosamine” or “TSNA” as used herein has its usual meaning in the art, namely a nitrosamine which is found only in tobacco industry products or other nicotine-containing products. Suitably the at least one tobacco-specific nitrosamine may be N'-nitrosonornicotine (NNN), 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK), N'-nitrosoanatabine (NAT) or N-nitrosoanabasine (NAB).

[0218] The term “precursor thereto” when used in relation to at least one tobacco-specific nitrosamine refers to one or more chemicals or compounds of a tobacco plant that give rise to the formation of a tobacco-specific nitrosamine or are involved in the nitrosation reaction leading to tobacco-specific nitrosamine production.

[0219] In one embodiment the TSNA may be one or more of group selected from: N'-nitrosonornicotine (NNN), 4-(methyl nitrosamino)-1-(3-pyridyl)-1-butanone (NNK), N'-nitrosoanatabine (NAT) and N'-nitrosoanabasine (NAB). Suitably the at least one tobacco-specific nitrosamine may be NNK or NNN. In one embodiment the tobacco-specific nitrosamine is NNN. In another embodiment the tobacco-specific nitrosamine is NNK.

[0220] In one embodiment the precursor of the TSNA is one or more of the group selected from nornicotine, anabasine, anatabine, and an oxidised derivative of nicotine such as pseudooxynicotine (PON).

[0221] In one embodiment the TSNA is N'-nitrosonornicotine (NNN) and/or the precursor is nornicotine. In one embodiment the content of NNN is decreased. In one embodiment the content of nornicotine is decreased. In one embodiment the content of NNN and nornicotine is decreased.

[0222] In one embodiment the TSNA is 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) and/or the precursor is PON. In one embodiment the content of NNK is decreased. In one embodiment the content of PON is decreased. In one embodiment the content of NNK and PON is decreased. In one embodiment the TSNA is N'-nitrosoanatabine (NAT) and/or the precursor is anatabine. In one embodiment the content of NAT is decreased. In one embodiment the content of anatabine is decreased. In one embodiment the content of NAT and anatabine is decreased.

[0223] In one embodiment the TSNA is N'-nitrosoanabasine (NAB) and/or the precursor is anabasine. In one embodiment the content of NAB is decreased. In one embodiment the content of nornicotine is decreased. In one embodiment the content of NAB and anabasine is decreased. The precursor of the TSNA (e.g. NNK, NNN, NAB and/or NAT) may be measured in green tobacco leaf, e.g. prior to processing, e.g. prior to curing. In one embodiment the precursor of the TSNA (e.g. NNK, NNN, NAB and/or NAT) is measured and/or modified (e.g. reduced) in a green tobacco leaf, e.g. prior to processing, e.g. prior to curing.

[0224] In one embodiment carrying out a method and/or use of the invention results in a reduction of at least one TSNA or a precursor thereto in the modified tobacco plant

(or part thereof) when compared to a tobacco plant (or part thereof) which has not been modified in accordance with the present invention.

[0225] The terms “reducing at least one TSNA or precursor thereto” or “reduction of at least one TSNA or precursor thereto” are used herein to mean that the concentration and/or total content of the at least one TSNA or precursor thereto in the product, method or use of the invention is lower in relation to a comparable product, method or use. For example, a comparable tobacco industry product would be derived from a tobacco plant which had not been modified according to the present invention, but in which all other relevant features were the same (e.g. plant species, growing conditions, method of processing tobacco, etc.).

[0226] Any method known in the art for determining the concentration and/or levels of at least one TSNA or precursor thereto may be used. In particular a method such may comprise the addition of deuterium labelled internal standard, an aqueous extraction and filtration, followed by analysis using reversed phase high performance liquid chromatography with tandem mass spectrometry (LC-MS/MS) may be used. Other examples for determining the concentration and/or level of a precursor to a tobacco-specific nitrosamine include a method such as the one detailed in CORESTA recommended method CRM-72: Determination of Tobacco Specific Nitrosamines in Tobacco and Tobacco Products by LC-MS/MS; CRM being developed into ISO/DIS 21766 or Wagner et al. (2005) Analytical Chemistry 77 (4), 1001-1006 all of which are incorporated herein by reference.

[0227] Suitably the concentration and/or total content of the at least one tobacco-specific nitrosamine or precursor thereto may be reduced by carrying out a method and/or use of the present invention. Suitably the concentration and/or level of the at least one tobacco-specific nitrosamine or precursor thereto may be reduced in a tobacco plant of the invention (e.g. obtainable or obtained by a method and/or use of the invention) when compared to the concentration and/or level of the at least one tobacco-specific nitrosamine (s) or precursor thereto in a tobacco plant which has not been modified in accordance with present invention.

[0228] The concentration and/or total content of the at least one tobacco-specific nitrosamine(s) or precursor thereto may be reduced in a tobacco leaf, harvested leaf, processed tobacco leaf, tobacco industry product or combinations thereof obtainable or obtained from a tobacco plant (or part of a tobacco plant or a tobacco cell culture) of the invention when compared with a tobacco leaf, harvested leaf, processed tobacco leaf, tobacco industry product or combinations thereof obtainable or obtained from a tobacco plant (or part of a tobacco plant or a tobacco cell culture) which has not been modified in accordance with the present invention.

[0229] Suitably the concentration and/or total content of the at least one tobacco-specific nitrosamine or precursor thereto may be reduced in a processed tobacco leaf.

[0230] Suitably the concentration and/or level of the at least one tobacco-specific nitrosamine or precursor thereto may be reduced in a tobacco industry product.

[0231] In one embodiment the at least one tobacco-specific nitrosamine or precursor thereto may be reduced by at least about 1%, at least about 3%, at least about 5%, at least about 10%, at least about 20%, at least about 30%, at least about 40% or at least about 50%. In some embodiments the

at least one tobacco-specific nitrosamine or precursor thereto may be reduced by between about 5% and about 50%, by between about 10% and about 50%, by between about 20% and about 50%, by between about 30% and about 50%, or by between about 40% and 50%.

[0232] In relation to processed (e.g. cured) tobacco leaf (e.g. cured or reconstituted), the at least one tobacco-specific nitrosamine or precursor thereto may be reduced by between about 5000 ng/g and about 50 ng/g, by between about 4000 ng/g and about 100 ng/g, by between about 3000 ng/g and 500 ng/g or by between 2000 ng/g and 1000 ng/g. In some embodiments the at least one tobacco-specific nitrosamine or precursor thereto may be reduced by at least about 5000 ng/g, at least about 4000 ng/g, at least about 3000 ng/g, at least about 2000 ng/g, at least about 1000 ng/g, at least about 500 ng/g, at least about 100 ng/g or at least about 50 ng/g.

Biomass Production

[0233] In some instances, it may be desirable to produce plants or biomass with high alkaloid levels e.g. high levels of nicotine content so that nicotine may be purified to produce a pure nicotine product for example for use in devices which utilize liquid containing nicotine (e.g. e-cigarettes) or within tobacco heating devices. For example, the production of nicotine in this way could reduce costs of nicotine extraction for the production of e-liquids for e-cigarettes.

[0234] In one aspect, the present invention provides a method of producing a biomass comprising: growing a cell which has been engineered to modulate (e.g. increase) the activity or expression of a gene encoding a YIPF-related protein under conditions to produce a biomass. Suitably, the activity or expression of a YIPF-related protein may be increased in order to increase the concentration and/or total nicotine content.

[0235] In one embodiment, the present invention provides a method of producing a biomass having modified (e.g. increased) concentration and/or total content of nicotine, comprising growing a cell which has been engineered to increase the activity or expression of at least one YIPF-related protein comprising an amino acid sequence as set out in SEQ ID No. 3, or a functional variant or functional fragment or orthologue thereof, or a sequence which has at least 80% identity to SEQ ID No. 3; or wherein the at least one YIPF-related protein comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2.

[0236] The cell may be engineered by any method known in the art to modify the activity or expression of at least one YIPF-related protein. Suitably, the cell may be engineered to express an exogenous gene encoding a YIPF-related protein. Suitably, the cell may be engineered to overexpress a gene encoding a YIPF-related protein.

[0237] Suitably, the biomass may contain a higher concentration and/or total content of nicotine compared with the biomass produced by a comparable cell which has not been modified in accordance with the present invention.

[0238] Suitably the cell for use in biomass production may be a plant cell, such as a tobacco cell.

[0239] Suitably the cell for use in biomass production may be a yeast cell.

[0240] In one embodiment the cell (e.g. yeast cell) may be further modified to comprise one or more sequences that increases nicotinic alkaloid biosynthesis. Suitably these one or more sequences may be incorporated into a nucleic acid construct that is suitable for cell (e.g. yeast cell) transformation. The one or more sequences may be overexpressed in the cell (e.g. yeast cell).

[0241] The sequences may be selected from one or more of the following genes: MPO (or Methylputrescine Oxidase or MPO1 or MPO2); A622 (or Isoflavone reductase-like protein or Isoflavone reductase homolog or Isoflavone reductase-like protein); BBL (or Berberine bridge enzyme or Berberine bridge enzyme-like or BBE or NBB1); PMT (or Putrescine N-Methyltransferase or putrescine methyltransferase or S-adenosyl-L-methionine: putrescine N-methyltransferase or PMT1 or PMT2 or PMT3 or PMT4) and QPT (or quinolinate phosphoribosyltransferase). In one embodiment, the sequences may be selected from one or more of the following genes: BBL, A622, PMT and MPO (MPO1 or MPO2). Genes suitable for modification in this way may be taught in US2016032299 for example, which is incorporated herein by reference.

Commercially Desirable Traits

[0242] In one embodiment the plants of the present invention have modified (i.e. increased or decreased) total alkaloid content and/or modified (i.e. increased or decreased) content of one or more alkaloids, whilst the flavour characteristics and/or other commercially desirable traits are at least maintained. Suitably, the plants of the present invention may have decreased total alkaloid content and/or decreased content of one or more alkaloids, whilst the flavour characteristics and/or other commercially desirable traits are at least maintained.

[0243] In one embodiment, the plants of the present invention produce leaves of a similar grade and/or quality to plants which have not been modified according to the invention.

[0244] In one embodiment, the plants of the present invention have reduced nornicotine and/or PON and/or anabasine and/or anatabine content without a significant change in the flavour characteristics of the plant (e.g. compared with the same plant which has not been modified in accordance with the present invention).

[0245] In one embodiment the plants of the present invention have decreased TSNA precursor content without a significant change (e.g. decrease) in other commercially desirable traits of the plant (e.g. compared with the same plant which has not been modified in accordance with the present invention). In particular, the yield of the modified plant is preferably not reduced compared with the same plant which has not been modified in accordance with the present invention.

[0246] Therefore in one embodiment the methods and uses of the present invention relate to decreasing TSNA precursor content whilst maintaining the flavour characteristics and/or other commercially desirable traits (e.g. yield).

[0247] The term “commercially desirable traits” as used herein will include traits such as yield, mature plant height, harvestable leaf number, average node length, cutter leaf length, cutter leaf width, quality (e.g. leaf quality, suitably cured leaf quality), abiotic (for instance drought) stress tolerance, herbicide tolerance and/or biotic (for instance insect, bacteria or fungus) stress tolerance.

[0248] Leaf quality may be measured based on colour, texture and aroma of the cured leaf, for example according to United States Department of Agriculture (USDA) grades and standards.

[0249] Tobacco grades are evaluated based on factors including, but not limited to, the leaf stalk position, leaf size, leaf colour, leaf uniformity and integrity, ripeness, texture, elasticity, sheen (related with the intensity and the depth of coloration of the leaf as well as the shine), hygroscopicity (the faculty of the tobacco leaves to absorb and to retain the ambient moisture), and green nuance or cast. Leaf grade can be determined using standard methods known in the art, for example, using an Official Standard Grade published by the Agricultural Marketing Service of the US Department of Agriculture (7 U.S.C. § 511). See, e.g., Official Standard Grades for Burley Tobacco (U.S. Type 31 and Foreign Type 93), effective Nov. 5, 1990 (55 F.R. 40645); Official Standard Grades for Flue-Cured Tobacco (U.S. Types 11, 12, 13, 14 and Foreign Type 92), effective Mar. 27, 1989 (54 F.R. 7925); Official Standard Grades for Pennsylvania Seedleaf Tobacco (U.S. Type 41), effective Jan. 8, 1965 (29 F.R. 16854); Official Standard Grades for Ohio Cigar-Leaf Tobacco (U.S. Types 42, 43, and 44), effective Dec. 8, 1963 (28 F.R. 11719 and 28 F.R. 11926); Official Standard Grades for Wisconsin Cigar-Binder Tobacco (U.S. Types 54 and 55), effective Nov. 20, 1969 (34 F.R. 17061); Official Standard Grades for Wisconsin Cigar-Binder Tobacco (U.S. Types 54 and 55), effective Nov. 20, 1969 (34 F.R. 17061); Official Standard Grades for Georgia and Florida Shade-Grown Cigar-Wrapper Tobacco (U.S. Type 62), Effective April 1971. A USDA grade index value can be determined according to an industry accepted grade index. See e.g. Bowman et al. (1988) Tobacco Science, 32:39-40; Legacy Tobacco Document Library (Bates Document #523267826-523267833, Jul. 1, 1988, Memorandum on the Proposed Burley Tobacco Grade Index); and Miller et al. (1990) Tobacco Intern., 192:55-57 (all foregoing references are incorporated herein in their entirety). In one aspect, a USDA grade index is a 0-100 numerical representation of federal grade received and is a weighted average of all stalk positions. A higher grade index indicates higher quality. Alternatively, leaf grade may be determined via hyperspectral imaging. See e.g. WO 2011/027315 (which is incorporated herein by reference).

[0250] In one embodiment, a tobacco plant of the present invention provides tobacco of commercially acceptable grade.

[0251] Suitably, the tobacco plant of the present invention provides cured tobacco of commercially acceptable grade.

[0252] In one embodiment, a tobacco plant of the present invention is capable of producing leaves having a USDA grade index value of at least about 70% of the USDA grade index value of leaves of a comparable plant when grown in similar growth conditions. Suitably, tobacco plants disclosed herein may be capable of producing leaves having a USDA grade index value of at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 98% of the USDA grade index value of a control plant when grown in similar growth conditions. Suitably, tobacco plants disclosed herein may be capable of producing leaves having a USDA grade index value of between 65% and 130%, between 70% and 130%, between 75% and 130%, between 80% and 130%, between 85% and 130%, between 90% and

130%, between 95% and 130%, between 100% and 130%, between 105% and 130%, between 110% and 130%, between 115% and 130%, or between 120% and 130% of the USDA grade index value of a comparable plant.

[0253] In one aspect, the tobacco plant of the present invention is capable of producing leaves having a USDA grade index value of at least 50. Suitably, tobacco plants disclosed herein may be capable of producing leaves having a USDA grade index value of 55 or more, 60 or more, 65 or more, 70 or more, 75 or more, 80 or more, 85 or more, 90 or more, and 95 or more.

[0254] Unless specified otherwise, used herein, tobacco yield refers to cured leaf yield which is calculated based on the weight of cured tobacco leaves per acre under standard field conditions following standard agronomic and curing practice.

[0255] In one aspect, a plant (e.g. a tobacco plant) of the present invention has a yield between 50% and 150%, between 55% and 145%, between 60% and 140%, between 65% and 135%, between 70% and 130%, between 75% and 125%, between 80% and 120%, between 85% and 115%, between 90% and 110%, between 95% and 105%, between 50% and 100%, between 55% and 100%, between 60% and 100%, between 65% and 100%, between 70% and 100%, between 75% and 100%, between 80% and 100%, between 85% and 100%, between 90% and 100%, between 95% and 100%, between 100% and 150%, between 105% and 150%, between 110% and 150%, between 115% and 150%, between 120% and 150%, between 125% and 150%, between 130% and 150%, between 135% and 150%, between 140% and 150%, or between 145% and 150% of the yield of a comparable plant when grown in similar field conditions.

[0256] In another aspect, the plant (e.g. a tobacco plant) yield of the present invention is approximately 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, or 3.0 times of the yield of a comparable plant when grown in similar field conditions.

[0257] In another aspect, the yield of a tobacco plant of the present invention is comparable to the yield of the flue cured comparable plant when grown in similar field conditions.

[0258] In one aspect, a tobacco plant of the present invention provides a yield selected from the group consisting of about between 1200 and 3500, between 1300 and 3400, between 1400 and 3300, between 1500 and 3200, between 1600 and 3100, between 1700 and 3000, between 1800 and 2900, between 1900 and 2800, between 2000 and 2700, between 2100 and 2600, between 2200 and 2500, and between 2300 and 2400 lbs/acre.

[0259] In another aspect, a tobacco plant of the present invention provides a yield selected from the group consisting of about between 1200 and 3500, between 1300 and 3500, between 1400 and 3500, between 1500 and 3500, between 1600 and 3500, between 1700 and 3500, between 1800 and 3500, between 1900 and 3500, between 2000 and 3500, between 2100 and 3500, between 2200 and 3500, between 2300 and 3500, between 2400 and 3500, between 2500 and 3500, between 2600 and 3500, between 2700 and 3500, between 2800 and 3500, between 2900 and 3500, between 3000 and 3500, and between 3100 and 3500 lbs/acre.

[0260] In a further aspect, a tobacco plant of the present invention provides a yield selected from the group consisting of about between 1200 and 3500, between 1200 and

3400, between 1200 and 3300, between 1200 and 3200, between 1200 and 3100, between 1200 and 3000, between 1200 and 2900, between 1200 and 2800, between 1200 and 2700, between 1200 and 2600, between 1200 and 2500, between 1200 and 2400, between 1200 and 2300, between 1200 and 2200, between 1200 and 2100, between 1200 and 2000, between 1200 and 1900, between 1200 and 1800, between 1200 and 1700, between 1200 and 1600, between 1200 and 1500, and between 1200 and 1400 lbs/acre.

Plant Breeding

[0261] In one embodiment the present invention provides a method of producing a plant having a modified alkaloid content and/or modified content of a tobacco specific nitrosamine (TSNA) precursor comprising:

[0262] a. crossing a donor plant having modified (e.g. decreased) nicotine content and/or modified (e.g. decreased) content of a tobacco specific nitrosamine (TSNA) precursor and wherein the activity or expression of at least one YIPF-related protein according to the present invention has been modulated (e.g. decreased) in the donor plant in accordance with the present invention with a recipient tobacco plant that does not have modified nicotine content or modified content of a tobacco specific nitrosamine (TSNA) precursor and possesses commercially desirable traits;

[0263] b. isolating genetic material from a progeny of said donor plant crossed with said recipient plant; and

[0264] c. performing molecular marker-assisted selection with a molecular marker comprising:

[0265] i. identifying an introgressed region comprising a mutation in a polynucleotide sequence encoding a protein defined in a.

[0266] Suitably, the activity or expression of a protein comprising an amino acid sequence as set out in SEQ ID No. 3, or a functional variant or functional fragment or orthologue thereof, or a sequence which has at least 80% identity to SEQ ID No. 3; or a homologue of SEQ ID No. 3; or a protein encoded by a nucleotide sequence as set out in SEQ ID No. 1 or 2 or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2 or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2; is modulated in the donor plant when compared to a comparable plant. Suitably, the alkaloid content and/or TSNA precursor content is decreased by said method. Suitably, the alkaloid content and/or TSNA precursor content is decreased and the activity or expression of said YIPF-related protein is decreased or inhibited.

[0267] The molecular marker assisted selection may comprise performing PCR to identify an introgressed nucleic acid sequence comprising a mutation which modulates the activity or expression of a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% identity thereto.

Tobacco Plants

[0268] The present invention provides methods, uses directed to plants (e.g. tobacco plants) as well as a cell (e.g. a tobacco cell), a cell culture, a plant (e.g. a tobacco plant) and a plant propagation material.

[0269] The term “tobacco plant” as used herein refers to a plant in the genus *Nicotiana* that is used in the production of

tobacco industry products. Non-limiting examples of suitable “tobacco” plants include *N. tabacum* and *N. rustica* (for example, *N. tabacum* L., LA B21, LN KY171, TI 1406, Basma, *Galpao*, Perique, Beinhart 1000-1, and Petico).

[0270] The tobacco material can be derived or obtained from varieties of *Nicotiana tabacum* types, commonly known as Burley varieties, flue or bright varieties and dark varieties. In some embodiments, the tobacco material is derived from a Burley, Virginia or a dark tobacco plant. The tobacco plant may be selected from Burley tobacco, rare tobacco, speciality tobacco, expanded tobacco or the like.

[0271] The use of tobacco cultivars and elite tobacco cultivars is also contemplated herein. The tobacco plant for use herein may therefore be a tobacco variety or elite tobacco cultivar. Particularly useful *Nicotiana tabacum* varieties include Flue-cured Virginia type, Burley type, and Oriental type. In some embodiments, the tobacco plant may be, for example, selected from one or more of the following varieties: L. cultivar T.I. 1068, AA 37-1, B 13P, Xanthi (Mitchell-Mor), KT D #3 Hybrid 107, Bel-W3, 79-615, Samsun Holmes NN, F4 from cross BU21×Hoja Parado, line 97, KTRDC #2 Hybrid 49, KTRDC #4 Hybrid 1 10, Burley 21, PM016, KTRDC #5 KY 160 SI, KTRDC #7 FCA, KTRDC #6 TN 86 SI, PM021, K 149, K 326, K 346, K 358, K 394, K 399, K 730, KY 10, KY 14, KY 160, KY 17, KY 8959, KY 9, KY 907, MD 609, McNair 373, NC 2000, PG 01, PG 04, P01, P02, P03, RG 11, RG 17, RG 8, Speight G-28, TN 86, TN 90, VA 509, AS44, Banket A1, Basma Drama B84/31, Basma I Zichna ZP4/B, Basma Xanthi BX 2A, Batek, Besuki Jember, C104, Coker 319, Coker 347, Criollo Misionero, PM092, Delcrest, Djebel 81, DVH 405, *Galpao* Comum, HB04P, Hicks Broadleaf, Kabakulak Ellassona, PM102, Kutsage E1, KY 14×L8, KY 171, LA BU 21, McNair 944, NC 2326, NC 71, NC 297, NC 3, PVH 03, PVH 09, PVH 19, PVH 21 10, Red Russian, Samsun, Saplak, Simmaba, Talgar 28, PM132, Wislica, Yayaldag, NC 4, TR Madole, Prilep HC-72, Prilep P23, Prilep PB 156/1, Prilep P12-2/1, Yaka JK-48, Yaka JB 125/3, TI-1068, KDH-960, TI-1070, TW136, PM204, PM205, Basma, TKF 4028, L8, TKF 2002, TN 90, GR141, Basma xanthi, GR149, GR153, and Petit Havana.

[0272] Non-limiting examples of varieties or cultivars are: BD 64, CC 101, CC 200, CC 27, CC 301, CC 400, CC 500, CC 600, CC 700, CC 800, CC 900, Coker 176, Coker 319, Coker 371 Gold, Coker 48, CD 263, DF91 1, DT 538 LC, *Galpao* tobacco, GL 26H, GL 350, GL 600, GL 737, GL 939, GL 973, HB 04P, HB 04P LC, HB3307PLC, Hybrid 403LC, Hybrid 404LC, Hybrid 501 LC, K 149, K 326, K 346, K 358, K394, K 399, K 730, KDH 959, KT 200, KT204LC, KY10, KY14, KY 160, KY 17, KY 171, KY 907, KY907LC, KTY14×L8 LC, Little Crittenden, McNair 373, McNair 944, msKY 14×L8, Narrow Leaf Madole, Narrow Leaf Madole LC, NBH 98, N-126, N-777LC, N-7371 LC, NC 100, NC 102, NC 2000, NC 291, NC 297, NC 299, NC 3, NC 4, NC 5, NC 6, NC7, NC 606, NC 71, NC 72, NC 810, NC BH 129, NC 2002, Neal Smith Madole, OXFORD 207, PD 7302 LC, PD 7309 LC, PD 7312 LC ‘Periq’e’ tobacco, PVH03, PVH09, PVH19, PVH50, PVH51, R 610, R 630, R 7-1 1, R 7-12, RG 17, RG 81, RG H51, RG H 4, RG H 51, RS 1410, Speight 168, Speight 172, Speight 179, Speight 210, Speight 220, Speight 225, Speight 227, Speight 234, Speight G-28, Speight G-70, Speight H-6, Speight H20, Speight NF3, TI 1406, TI 1269, TN 86, TN86LC, TN 90, TN 97, TN97LC, TN D94, TN D950, TR (Tom Rosson) Madole, VA 309,

VA359, AA 37-1, B 13P, Xanthi (Mitchell-Mor), Bel-W3, 79-615, Samsun Holmes NN, KTRDC number 2 Hybrid 49, Burley 21, KY 8959, KY 9, MD 609, PG 01, PG 04, P01, P02, P03, RG 1 1, RG 8, VA 509, AS44, Banket A1, Basma Drama B84/31, Basma I Zichna ZP4/B, Basma Xanthi BX 2A, Batek, Besuki Jember, C104, Coker 347, Criollo Misionero, Delcrest, Djebel 81, DVH 405, *Galpao* Comum, HB04P, Hicks Broadleaf, Kabakulak Ellassona, Kutsage E1, LA BU 21, NC 2326, NC 297, PVH 21 10, Red Russian, Samsun, Saplak, Simmaba, Talgar 28, Wislica, Yayaldag, Prilep HC-72, Prilep P23, Prilep PB 156/1, Prilep P12-2/1, Yaka JK-48, Yaka JB 125/3, TI-1068, KDH-960, TI-1070, TW136, Basma, TKF 4028, L8, TKF 2002, GR141, Basma xanthi, GR149, GR153, Petit Havana. Low converter sub-varieties of the above, even if not specifically identified herein, are also contemplated.

[0273] The tobacco plant may be a Burley, Flue-cured Virginia, or Oriental.

[0274] In one embodiment the plant propagation material may be obtainable from a plant (e.g. a tobacco plant) of the invention.

[0275] A “plant propagation material” as used herein refers to any plant matter taken from a plant from which further plants may be produced. Suitably, a plant propagation material may be selected from a seed, plant calli and plant clumps. Suitably the plant propagation material may be a seed. Suitably, the plant propagation material may be plant calli. Suitably the plant propagation material may be plant clumps.

[0276] In one embodiment the cell (e.g. tobacco cell), cell culture, tobacco plant and/or plant propagation material may be obtainable (e.g. obtained) by a method according to the invention. Suitably a tobacco plant according to the present invention may have modulated (e.g. decreased) nicotine content when compared to an unmodified tobacco plant, wherein the tobacco plant has been modified to modulate (e.g. decrease) the activity or expression of at least one YIPF-related protein. Suitably a tobacco plant according to the present invention may have decreased nicotine content when compared to an unmodified tobacco plant, wherein the tobacco plant has been modified to decrease or inhibit the activity or expression of at least one YIPF-related protein.

[0277] Suitably a tobacco plant according to the present invention may have modulated (e.g. reduced) content of a tobacco specific nitrosamine (TSNA) precursor when compared to an unmodified tobacco plant, wherein the tobacco plant has been modified to modulate (e.g. decrease) the activity or expression of at least one YIPF-related protein.

[0278] Suitably a tobacco plant according to the present invention may have decreased tobacco specific nitrosamine (TSNA) precursor content when compared to an unmodified tobacco plant, wherein the tobacco plant has been modified to decrease or inhibit the activity or expression of at least one YIPF-related protein.

[0279] In one embodiment the tobacco plant in accordance with the present invention comprises a tobacco cell of the invention.

[0280] In another embodiment the plant propagation material may be obtainable (e.g. obtained) from a tobacco plant of the invention.

[0281] In one embodiment there is provided the use of a tobacco plant as described herein to breed a tobacco plant.

[0282] The present invention also provides in another embodiment the use of a tobacco plant of the foregoing embodiments for the production of a tobacco industry product.

[0283] In another embodiment there is provided the use of a tobacco plant of the invention to grow a crop.

[0284] In one embodiment there is provided the use of a cell as provided for in the foregoing embodiments for production of a tobacco industry product.

[0285] In one embodiment the present invention provides a cell culture (e.g. in vitro culture).

[0286] The tobacco cell culture may be a cell suspension culture. These cells cultured in vitro may be incorporated into a tobacco industry product, e.g. as a substitute for conventional tobacco particles, shreds, fine cut or long cut tobacco lamina, as an additive ingredient or as both a substitute and an additive. Suitably, the cell culture may produce nicotine.

[0287] In one embodiment there is provided the use of a cell culture, e.g. a harvested and/or processed cell culture according to the present invention for the production of a tobacco industry product.

[0288] The tobacco cells harvested from an in vitro culture may be dried, e.g. freeze-dried, for example to produce a powder.

[0289] In one embodiment, the cell culture is a tobacco cell culture. The skilled person will be aware of known methods for establishing in vitro cultures of tobacco cells. By way of example only, the following method may be used: collecting seeds from a tobacco plant of interest and sterilising their exterior to eliminate unwanted organisms, planting said seeds to grow a tobacco plant of interest, removing tissue from the tobacco plant (for example, from the tobacco stem) for use as an explant, establishing a callus culture from the tobacco explant, establishing a cell suspension culture from the callus culture, and harvesting culture material (e.g. including tobacco cells) to produce a tobacco cell culture.

[0290] The tobacco cells can be harvested by various methods, including filtration, e.g. vacuum filtration. The sample may be washed in the filter by adding water and the remaining liquid removed with the filtration, e.g. vacuum filtration.

[0291] The harvested tobacco cell culture may be further processed, e.g. dried, such as air-dried and/or freeze-dried. The harvested tobacco cell culture or dried harvested tobacco cell culture or an extract therefrom may be incorporated into tobacco industry products according to the present invention.

[0292] In one embodiment, the present invention provides a plant (e.g. tobacco plant) or part thereof for use in molecular farming. Suitably, a plant or part thereof modified in accordance with the present invention may be used in the manufacture of proteins such as therapeutics e.g. antibiotics, virus like particles, neutraceuticals or small molecules.

[0293] In one embodiment, the present invention provides a method for the production of proteins (e.g. therapeutic proteins), the method comprising modifying a plant or part thereof capable of producing said protein (e.g. therapeutic protein) by modulating (e.g. decreasing) the activity or expression of at least one YIPF-related protein having an amino acid sequence as set out in SEQ ID No.3, or a functional variant or functional fragment or orthologue thereof, or a sequence which has at least 80% identity to SEQ ID No. 3; or a homologue of SEQ ID No. 3; or wherein

the at least one YIPF-related protein comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2; and culturing the plant under conditions sufficient to allow the production of said protein (e.g. therapeutic protein).

Products

[0294] The present invention also provides for products obtainable or obtained from plants according to the present invention. Products are provided which are obtainable or obtained from a plant in which the activity or expression of a YIPF-related protein has been modulated.

[0295] In one embodiment, the product may comprise a construct of the invention which modulates the activity or expression of at least one YIPF-related protein as defined herein. In one embodiment, the product may comprise a construct of the invention which modifies the nucleic acid sequence of at least one YIPF-related protein as defined herein.

[0296] The present invention also provides for products obtainable or obtained from tobacco according to the present invention.

[0297] In one embodiment there is provided the use of a tobacco plant of the invention to produce a tobacco leaf.

[0298] Suitably the tobacco leaf may be subjected to downstream applications such as processing.

[0299] Thus in one embodiment the use of the foregoing embodiment may provide a processed tobacco leaf. Suitably the tobacco leaf may be subjected to curing, fermenting, pasteurising or combinations thereof. In another embodiment the tobacco leaf may be cut. In some embodiments the tobacco leaf may be cut before or after being subjected to curing, fermenting, pasteurising or combinations thereof.

[0300] In one embodiment the present invention provides a harvested leaf of a tobacco plant of the invention.

[0301] In a further embodiment the harvested leaf may be obtainable (e.g. obtained) from a tobacco plant propagated from a propagation material of the present invention.

[0302] In another embodiment there is provided a harvest leaf obtainable from a method or use of the present invention.

[0303] Suitably the harvested leaf may be a cut harvested leaf.

[0304] In some embodiments the harvested leaf may comprise viable tobacco cells. In other embodiments the harvested leaf may be subjected to further processing.

[0305] There is also provided a processed tobacco leaf.

[0306] The processed tobacco leaf may be obtainable from a tobacco plant of the invention. Suitably the processed tobacco leaf may be obtainable from a tobacco plant obtained in accordance with any of the methods and/or uses of the present invention.

[0307] Suitably, the processed leaf may comprise reduced content of one or more TSNA's selected from NNN, NNK, NAT and NAB. Suitably, the content of NNN may be reduced. Suitably, the content of NNK may be reduced. Suitably, the content of NAT may be reduced. Suitably, the content of NAB may be reduced. Suitably, the reduction in TSNA content is in relation to a comparable product which has not been modified according to the present invention.

[0308] In another embodiment the processed tobacco leaf may be obtainable (e.g. obtained) from a tobacco plant propagated from a tobacco plant propagation material according to the present invention.

[0309] The processed tobacco leaf of the present invention may be obtainable (e.g. obtained) by processing a harvested leaf of the invention.

[0310] The term “processed tobacco leaf” as used herein refers to a tobacco leaf that has undergone one or more processing steps to which tobacco is subjected to in the art. A “processed tobacco leaf” comprises no or substantially no viable cells.

[0311] The term “viable cells” refers to cells which are able to grow and/or are metabolically active. Thus, if a cell is said to not be viable, also referred to as “non-viable” then a cell does not display the characteristics of a viable cell.

[0312] The term “substantially no viable cells” means that less than about 5% of the total cells are viable.

[0313] Preferably, less than about 3%, more preferably less than about 1%, even more preferably less than about 0.1% of the total cells are viable.

[0314] In one embodiment the processed tobacco leaf may be processed by one or more of: curing, fermenting and/or pasteurising.

[0315] Suitably the processed tobacco leaf may be processed by curing.

[0316] Tobacco leaf may be cured by any method known in the art. In one embodiment tobacco leaf may be cured by one or more of the curing methods selected from the group consisting of: air curing, fire curing, flue curing and sun curing.

[0317] Suitably the tobacco leaf may be air cured.

[0318] Typically air curing is achieved by hanging tobacco leaf in well-ventilated barns and allowing to dry. This is usually carried out over a period of four to eight weeks. Air curing is especially suitable for burley tobacco.

[0319] Suitably the tobacco leaf may be fire cured. Fire curing is typically achieved by hanging tobacco leaf in large barns where fires of hardwoods are kept on continuous or intermittent low smoulder and usually takes between three days and ten weeks, depending on the process and the tobacco.

[0320] In another embodiment the tobacco leaf may be flue cured. Flue curing may comprise stringing tobacco leaves onto tobacco sticks and hanging them from tier-poles in curing barns. The barns usually have a flue which runs from externally fed fire boxes. Typically this results in tobacco that has been heat-cured without being exposed to smoke. Usually the temperature will be raised slowly over the course of the curing with the whole process taking approximately 1 week.

[0321] Suitably the tobacco leaf may be sun cured. This method typically involves exposure of uncovered tobacco to the sun.

[0322] Suitably the processed tobacco leaf may be processed by fermenting.

[0323] Fermentation can be carried out in any manner known in the art. Typically during fermentation, the tobacco leaves are piled into stacks (a bulk) of cured tobacco covered in e.g. burlap to retain moisture. The combination of the remaining water inside the leaf and the weight of the tobacco generates a natural heat which ripens the tobacco. The temperature in the centre of the bulk is monitored daily. In some methods every week, the entire bulk is opened. The

leaves are then removed to be shaken and moistened and the bulk is rotated so that the inside leaves go outside and the bottom leaves are placed on the top of the bulk. This ensures even fermentation throughout the bulk. The additional moisture on the leaves, plus the actual rotation of the leaves themselves, generates heat, releasing the tobacco's natural ammonia and reducing nicotine, while also deepening the colour and improving the tobacco's aroma. Typically the fermentation process continues for up to 6 months, depending on the variety of tobacco, stalk position on the leaf, thickness and intended use of leaf.

[0324] Suitably the processed tobacco leaf may be processed by pasteurising. Pasteurising may be particularly preferred when the tobacco leaf will be used to make a smokeless tobacco industry product, most preferably snus.

[0325] Tobacco leaf pasteurisation may be carried out by any method known in the art. For example pasteurisation may be carried out as detailed in J Foulds, L Ramstrom, M Burke, K Fagerstrom. Effect of smokeless tobacco (snus) on smoking and public health in Sweden Tobacco Control (2003) 12:349-359, the teaching of which is incorporated herein by reference.

[0326] During the production of snus, pasteurisation is typically carried out by a process in which the tobacco is heat treated with steam for 24-36 hours (reaching temperatures of approximately 100° C.). This results in an almost sterile product and without wishing to be bound by theory one of the consequences of this is believed to be a limitation of further TSNA formation. In one embodiment the pasteurisation may be steam pasteurisation.

[0327] In some embodiments the processed tobacco leaf may be cut. The processed tobacco leaf may be cut before or after processing. Suitably, the processed tobacco leaf may be cut after processing.

[0328] In one embodiment, the use of the foregoing embodiment may provide reconstituted tobacco. In one embodiment, there is provided reconstituted tobacco.

[0329] “Reconstituted” as used herein may also be referred to as recon, recycled or homogenized sheet tobacco and refers to tobacco material generated from remnants of tobacco leaf after processing.

[0330] Reconstituted tobacco allows the production of a consistent, high quality blend and allows the adjustment of the ratio of individual components.

[0331] Reconstituted tobacco may be nano fibre recon (nanofibers can be extracted in solid or liquid form), paper making recon (which uses stems, scraps, and midribs, etc. as the raw material) or slurry type recon (which uses a mixture of fines and tobacco stems, ground to power, mixed with water and vegetable binding agent; the soluble residue is formed to sheets by extracting the water).

[0332] Any method known in the art may be used for making reconstituted tobacco, for example see CORESTA Congress, Sapporo, 2012, Smoke Science/Product Technology Groups, SSPT 12 (incorporated herein by reference).

[0333] In some embodiments the tobacco plant, harvested leaf of a tobacco plant and/or processed tobacco leaf may be used to extract nicotine. The extraction of nicotine can be achieved using any method known in the art. For example a method for extracting nicotine from tobacco is taught in U.S. Pat. No. 2,162,738 which is incorporated herein by reference.

[0334] In one aspect, the present invention provides cured tobacco material made from a tobacco plant or part thereof according to the invention.

[0335] Suitably, the cured tobacco may comprise a reduced content of one or more TSNA's selected from NNK, NNN, NAT and NAB. Suitably, the content of NNN may be reduced. Suitably, the content of NNK may be reduced. Suitably, the content of NAT may be reduced. Suitably, the content of NAB may be reduced. Suitably, the reduction in TSNA content is in relation to a comparable product which has not been modified according to the present invention.

[0336] In another aspect, the present invention provides a tobacco blend comprising tobacco material made from a tobacco plant or part thereof according to the present invention, or from a tobacco cell culture according to the present invention. In one aspect, the present invention provides a tobacco blend comprising cured tobacco material according to the present invention.

[0337] Suitably, the tobacco blend according to the present invention may comprise approximately 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80% or 90% tobacco from a tobacco plant or part thereof according to the present invention, or from a tobacco cell culture according to the present invention. Suitably, the tobacco blend may comprise approximately 10% tobacco from a tobacco plant or part thereof according to the present invention, or from a tobacco cell culture according to the present invention. Suitably, the tobacco blend may comprise approximately 20% tobacco from a tobacco plant or part thereof according to the present invention, or from a tobacco cell culture according to the present invention. Suitably, the tobacco blend may comprise approximately 30% tobacco from a tobacco plant or part thereof according to the present invention, or from a tobacco cell culture according to the present invention. Suitably, the tobacco blend may comprise approximately 40% tobacco from a tobacco plant or part thereof according to the present invention, or from a tobacco cell culture according to the present invention. Suitably, the tobacco blend may comprise approximately 50% tobacco from a tobacco plant or part thereof according to the present invention, or from a tobacco cell culture according to the present invention. Suitably, the tobacco blend may comprise approximately 60% tobacco from a tobacco plant or part thereof according to the present invention, or from a tobacco cell culture according to the present invention. Suitably, the tobacco blend may comprise approximately 70% tobacco from a tobacco plant or part thereof according to the present invention, or from a tobacco cell culture according to the present invention. Suitably, the tobacco blend may comprise approximately 80% tobacco from a tobacco plant or part thereof according to the present invention, or from a tobacco cell culture according to the present invention. Suitably, the tobacco blend may comprise approximately 90% tobacco from a tobacco plant or part thereof according to the present invention, or from a tobacco cell culture according to the present invention.

[0338] In one aspect, a tobacco blend product of the present invention comprises at least about 5, 10, 20, 30, 40, 50, 60, 70, 80, 90, or 95 percent by dry weight of tobacco cured from a tobacco plant or part thereof according to the present invention, or a tobacco cell culture according to the present invention.

[0339] Suitably, the cured tobacco material may be air cured. Suitably, the cured tobacco material may be flue

cured. Suitably, the cured tobacco material may be sun cured. Suitably, the cured tobacco material may be fire cured.

[0340] A tobacco industry product or smoking article according to the present invention may comprise the tobacco material (e.g. cured tobacco material or reconstituted tobacco material) according to the present invention.

[0341] In another aspect the present invention provides a tobacco industry product.

[0342] In one embodiment the tobacco industry product according to the present invention may be a blended tobacco industry product. Suitably, the tobacco blend may comprise cured tobacco material according to the present invention.

[0343] In one embodiment the tobacco industry product may be prepared from a tobacco plant of the invention or a part thereof.

[0344] Suitably the tobacco plant or part thereof may be propagated from a tobacco plant propagation material according to the present invention.

[0345] The term "part thereof" as used herein in the context of a tobacco plant refers to a portion of the tobacco plant. Suitably, the "part thereof" may be a leaf, root or stem of a tobacco plant or the flowers. Suitably, the "part thereof" may be a leaf, root or stem of a tobacco plant.

Tobacco Industry Product

[0346] As used herein, the term "tobacco industry product" is intended to include combustible smoking articles such as cigarettes, cigarillos, cigars, tobacco for pipes or for roll-your-own cigarettes, (whether based on tobacco, tobacco derivatives, expanded tobacco, reconstituted tobacco, tobacco substitutes or other smokable material), non-combustible aerosol provision systems such as heating products that release compounds from substrate materials without burning such as electronic cigarettes, tobacco heating products, and hybrid systems to generate aerosol from a combination of substrate materials, for example hybrid systems containing a liquid or gel or solid substrate, as well as aerosolizable substrate materials used within these aerosol provision systems; and aerosol-free delivery articles such as lozenges, gums, patches, articles comprising breathable powders and smokeless tobacco industry products such as snus and snuff, which aerosol-free delivery articles may or may not deliver nicotine.

[0347] In one embodiment the tobacco industry product may be prepared from (e.g. may comprise) a tobacco plant of the invention or a part thereof.

[0348] Suitably the tobacco plant or part thereof may be propagated from a tobacco plant propagation material according to the present invention.

[0349] The term "part thereof" as used herein in the context of a tobacco plant refers to a portion of the tobacco plant. Preferably the "part thereof" is a leaf of a tobacco plant.

[0350] In another embodiment the tobacco industry product may be prepared from a harvested leaf of the invention.

[0351] In a further embodiment the tobacco industry product may be prepared from a processed tobacco leaf of the invention.

[0352] Suitably the tobacco industry product may be prepared from a tobacco leaf processed by one or more of: curing, fermenting and/or pasteurising.

[0353] Suitably the tobacco industry product may comprise a cut tobacco leaf, optionally processed as per the foregoing embodiment.

[0354] In another embodiment, the tobacco industry product may be prepared from a tobacco cell culture according to the present invention.

[0355] In another embodiment, the tobacco industry product may be prepared from (e.g. may comprise) a cured tobacco material according to the present invention.

[0356] In another embodiment, the tobacco industry product may be prepared from (e.g. may comprise) a tobacco blend according to the present invention.

[0357] In one embodiment the tobacco industry product may be a smoking article.

[0358] As used herein, the term “smoking article” can include smokeable products, such as rolling tobacco, cigarettes, cigars and cigarillos whether based on tobacco, tobacco derivatives, expanded tobacco, reconstituted tobacco or tobacco substitutes.

[0359] In another embodiment the tobacco industry product may be a smokeless tobacco industry product.

[0360] The term “smokeless tobacco industry product” as used herein refers to a tobacco industry product that is not intended to be smoked and/or subjected to combustion.

[0361] Smokeless tobacco industry products (including heat-not-burn materials) may contain tobacco in any form, including dried particles, shreds, granules, powders, or slurry, deposited on, mixed in, surrounded by, or combined with other ingredients in any format, such as flakes, films, tabs, foams, or beads.

[0362] In one embodiment a smokeless tobacco industry product may include snus, snuff, chewing tobacco or the like.

[0363] In one embodiment, the tobacco industry product is a combustible smoking article, selected from the group consisting of a cigarette, a cigarillo and a cigar.

[0364] In one embodiment, the tobacco industry product comprises one or more components of a combustible smoking article, such as a filter, a filter rod, a filter rod segments, tobacco, a tobacco rod, a tobacco rod segment, a spill, an additive release component such as a capsule, a thread, beads, a paper such as a plug wrap, a tipping paper or a cigarette paper.

[0365] In one embodiment, the tobacco industry product is a non-combustible aerosol provision system. In one embodiment, the tobacco industry product comprises one or more components of a non-combustible aerosol provision system, such as a heater and an aerosolizable substrate. In one embodiment, the aerosol provision system is an electronic cigarette also known as a vaping device.

[0366] In one embodiment the electronic cigarette comprises a heater, a power supply capable of supplying power to the heater, an aerosolizable substrate such as a liquid or gel, a housing and optionally a mouthpiece.

[0367] In one embodiment the aerosolizable substrate is contained in a substrate container. In one embodiment the substrate container is combined with or comprises the heater.

[0368] In one embodiment, the tobacco industry product is a heating product which releases one or more compounds by heating, but not burning, a substrate material. The substrate material is an aerosolizable material which may be for example tobacco or other non-tobacco products, which may

or may not contain nicotine. In one embodiment, the heating product is a tobacco heating product.

[0369] In one embodiment, the heating product is an electronic device.

[0370] In one embodiment, the tobacco heating product comprises a heater, a power supply capable of supplying power to the heater, an aerosolizable substrate such as a solid or gel material. In one embodiment the heating product is a non-electronic article.

[0371] In one embodiment the heating product comprises an aerosolizable substrate such as a solid or gel material and a heat source which is capable of supplying heat energy to the aerosolizable substrate without any electronic means, such as by burning a combustion material, such as charcoal.

[0372] In one embodiment the heating product also comprises a filter capable of filtering the aerosol generated by heating the aerosolizable substrate.

[0373] In some embodiments the aerosolizable substrate material may comprise a vapour or aerosol generating agent or a humectant, such as glycerol, propylene glycol, triacetin or diethylene glycol.

[0374] In one embodiment, the tobacco industry product is a hybrid system to generate aerosol by heating, but not burning, a combination of substrate materials. The substrate materials may comprise for example solid, liquid or gel which may or may not contain nicotine. In one embodiment, the hybrid system comprises a liquid or gel substrate and a solid substrate. The solid substrate may be for example tobacco or other non-tobacco products, which may or may not contain nicotine. In one embodiment, the hybrid system comprises a liquid or gel substrate and tobacco.

[0375] In a further embodiment the tobacco industry product may be a tobacco heating device or hybrid device or e-cigarette or the like.

[0376] Typically in tobacco heating devices or hybrid devices, an aerosol is generated by the transfer of heat from a heat source to a physically separate aerosol-forming substrate or material, which may be located within, around or downstream of the heat source. During smoking, volatile compounds are released from the aerosol-forming substrate by heat transfer from the heat source and entrained in air drawn through the smoking article. As the released compounds cool, they condense to form an aerosol that is inhaled by the user.

[0377] Aerosol-generating articles and devices for consuming or smoking tobacco heating devices are known in the art. They can include, for example, electrically heated aerosol-generating devices in which an aerosol is generated by the transfer of heat from one or more electrical heating elements of the aerosol-generating device to the aerosol-forming substrate of a tobacco heating device. Suitably the tobacco heating device may be an aerosol-generating device.

[0378] Preferably the tobacco heating device may be a heat-not-burn device. Heat-not-burn devices are known in the art and release compounds by heating, but not burning, tobacco.

[0379] An example of a suitable, heat-not-burn device may be one taught in WO2013/034459 or GB2515502 which are incorporated herein by reference.

[0380] In one embodiment the aerosol-forming substrate of a tobacco heating device may be a tobacco industry product in accordance with the present invention.

[0381] In one embodiment the tobacco heating device may be a hybrid device.

Polynucleotides/Polypeptides/Constructs

[0382] In certain embodiments of the present invention, constructs which modulate activity or expression of at least one YIPF-related protein as described herein may be transformed into plant cells, suitably under the direction of a promoter.

[0383] In certain embodiments of the present invention, constructs which decrease (i.e. inhibit) the activity or expression of at least one YIPF-related protein as described herein may be transformed into plant cells under the direction of a promoter. For example, the genetic construct may be a gene editing construct or may comprise an RNAi molecule, which may comprise a small interfering RNA (siRNA) molecule, or a short hairpin loop (shRNA) molecule.

[0384] In certain embodiments of the present invention, constructs which increase activity or expression of gene encoding a YIPF-related protein as described herein may be transformed into plant cells, suitably under the direction of a promoter e.g. constructs which encode a gene encoding a YIPF-related protein such as an endogenous YIPF-related protein.

[0385] Constructs may be introduced into plants according to the present invention by means of suitable vector, e.g. plant transformation vectors. A plant transformation vector may comprise an expression cassette comprising 5'-3' in the direction of transcription, a promoter sequence, a construct sequence targeting gene encoding a YIPF-related protein as described herein and, optionally a 3' untranslated, terminator sequence including a stop signal for RNA polymerase and a polyadenylation signal for polyadenylase. The promoter sequence may be present in one or more copies, and such copies may be identical or variants of a promoter sequence as described above. The terminator sequence may be obtained from plant, bacterial or viral genes. Suitable terminator sequences are the pea *rbcs* E9 terminator sequence, the *nos* terminator sequence derived from the nopaline synthase gene of *Agrobacterium tumefaciens* and the 35S terminator sequence from cauliflower mosaic virus, for example. A person skilled in the art will be readily aware of other suitable terminator sequences.

[0386] The construct of the present invention may also comprise a gene expression enhancing mechanism to increase the strength of the promoter. An example of such an enhancer element is one derived from a portion of the promoter of the pea plastocyanin gene, and which is the subject of International Patent Application No. WO 97/20056 which is incorporated herein by reference. Suitable enhancer elements may be the *nos* enhancer element derived from the nopaline synthase gene of *Agrobacterium tumefaciens* and the 35S enhancer element from cauliflower mosaic virus, for example.

[0387] These regulatory regions may be derived from the same gene as the promoter DNA sequence or may be derived from different genes, from *Nicotiana tabacum* or other organisms, for example from a plant of the family Solanaceae, or from the subfamily Cestroideae. All of the regulatory regions should be capable of operating in cells of the tissue to be transformed.

[0388] The promoter DNA sequence may be derived from the same gene as the gene of interest, e.g. the gene the promoter is going to direct, for instance a gene encoding a YIPF-related protein according to the invention, a coding sequence used in the present invention or may be derived

from a different gene, from *Nicotiana tabacum*, or another organism, for example from a plant of the family Solanaceae, or from the subfamily Cestroideae.

[0389] The expression cassette may be incorporated into a basic plant transformation vector, such as pBIN 19 Plus, pBI 101, pKYLX71: 35S2, pCambia2300 or other suitable plant transformation vectors known in the art. In addition to the expression cassette, the plant transformation vector will contain such sequences as are necessary for the transformation process. These may include the *Agrobacterium* vir genes, one or more T-DNA border sequences, and a selectable marker or other means of identifying transgenic plant cells.

[0390] The term “expression vector or plant transformation vector” means a construct capable of in vivo or in vitro expression. Preferably, the expression vector is incorporated in the genome of the organism. In one embodiment the vector of the present invention expresses a protein e.g. a YIPF-related protein as described herein. The term “incorporated” preferably relates to stable incorporation into the genome.

[0391] Techniques for transforming plants are well known within the art and include *Agrobacterium*-mediated transformation, for example. The basic principle in the construction of genetically modified plants is to insert genetic information in the plant genome so as to obtain a stable maintenance of the inserted genetic material. A review of the general techniques may be found in articles by Potrykus (*Annu Rev Plant Physiol Plant Mol Biol* 42:205-225) and Christon (*AgroFood-Industry Hi-Tech March/April 1994* 17-27), which are incorporated herein by reference.

[0392] Typically, in *Agrobacterium*-mediated transformation a binary vector carrying a foreign DNA of interest, i.e. a construct according to the present invention, is transferred from an appropriate *Agrobacterium* strain to a target plant by the co-cultivation of the *Agrobacterium* with explants from the target plant. Transformed plant tissue is then regenerated on selection media, which selection media comprises a selectable marker and plant growth hormones. An alternative is the floral dip method (Clough & Bent, 1998 *Plant J.* 1998 December; 16 (6): 735-43, which is incorporated herein by reference) whereby floral buds of an intact plant are brought into contact with a suspension of the *Agrobacterium* strain containing the chimeric gene, and following seed set, transformed individuals are germinated and identified by growth on selective media. Direct infection of plant tissues by *Agrobacterium* is a simple technique which has been widely employed and which is described in Butcher et al. (1980) *Tissue Culture Methods for Plant Pathologists*, eds.: D. S. Ingrams and J. P. Helgeson, 203-208 which is incorporated herein by reference. Further suitable transformation methods include direct gene transfer into protoplasts using polyethylene glycol or electroporation techniques, particle bombardment, micro-injection and the use of silicon carbide fibres for example. Transforming plants using ballistic transformation and production of fertile transgenic maize plants by silicon carbide whisker-mediated transformation is taught in Frame et al. (1994) *The Plant Journal* 6 (6): 941-948, which is incorporated herein by reference, and viral transformation techniques is taught in, for example, Meyer et al. (1992) *Mol. Gen. Genet.* 231 (3): 345-352, which is incorporated herein by reference. The use of cassava mosaic virus as a vector system for plants is taught in Meyer et al. (1992) *Gene* 110:213-217, which is incor-

porated herein by reference. Further teachings on plant transformation may be found in EP-A-0449375, incorporated herein by reference.

[0393] In a further aspect, the present invention relates to a vector system which carries a construct and introducing it into the genome of an organism, such as a plant, suitably a tobacco plant. The vector system may comprise one vector, but it may comprise two vectors. In the case of two vectors, the vector system is normally referred to as a binary vector system. Binary vector systems are described in further detail in Gynheung et al. (1980) Binary Vectors, Plant Molecular Biology Manual A3, 1-19, which is incorporated herein by reference.

[0394] One extensively employed system for transformation of plant cells uses the Ti plasmid from *Agrobacterium tumefaciens* or a Ri plasmid from *Agrobacterium rhizogenes* described by An et al. (1986) Plant Physiol. 81, 301-305 and Butcher et al. (1980) Tissue Culture Methods for Plant Pathologists eds.: D. S. Ingrams and J. P. Helgeson, 203-208 which are incorporated herein by reference. After each introduction method of the desired exogenous gene according to the present invention in the plants, the presence and/or insertion of further DNA sequences may be necessary. The use of T-DNA for the transformation of plant cells has been intensively studied and is described in EP-A-120516; Hoekema (1985) The Binary Plant Vector System, Offset-drukkerij Kanters B. B., Amsterdam Chapter V; Fraley et al. Crit. Rev. Plant Sci. 4:1-46; and An et al. (1985) EMBO J 4:277-284, all incorporated herein by reference.

[0395] Plant cells transformed with construct(s) which modulate the activity or expression of at least one YIPF-related protein may be grown and maintained in accordance with well-known tissue culturing methods such as by culturing the cells in a suitable culture medium supplied with the necessary growth factors such as amino acids, plant hormones, vitamins, etc.

[0396] The term “transgenic plant” in relation to the present invention includes any plant that comprises a construct which modulates the activity or expression of at least one YIPF-related protein according to the invention. Accordingly a transgenic plant is a plant which has been transformed with a construct according to the invention. Preferably the transgenic plant exhibits modulated (e.g. reduced) alkaloid content and/or modulated (e.g. reduced) TSNA precursor content according to the present invention. The term “transgenic plant” does not cover native nucleotide coding sequences in their natural environment when they are under the control of their native promoter which is also in its natural environment.

[0397] In one aspect, a gene encoding a YIPF-related protein, a construct, a plant transformation vector or a plant cell according to the present invention is in an isolated form. The term “isolated” means that the sequence is at least substantially free from at least one other component with which the sequence is naturally associated in nature and as found in nature.

[0398] In one aspect, a gene encoding a YIPF-related protein, a construct, plant transformation vector or a plant cell according to the invention is in a purified form. The term “purified” means in a relatively pure state, e.g. at least about 90% pure, or at least about 95% pure or at least about 98% pure.

[0399] The term “nucleotide sequence” as used herein refers to an oligonucleotide sequence or polynucleotide

sequence, and variant, homologues, fragments and derivatives thereof (such as portions thereof). The nucleotide sequence may be of genomic or synthetic or recombinant origin, which may be double-stranded or single-stranded whether representing the sense or anti-sense strand.

[0400] The term “nucleotide sequence” in relation to the present invention includes genomic DNA, cDNA, synthetic DNA, and RNA. Preferably it means DNA, more preferably cDNA sequence coding for the present invention.

[0401] In a preferred embodiment, the nucleotide sequence when relating to and when encompassed by the per se scope of the present invention, i.e. the gene encoding a YIPF-related protein according to the present invention, includes the native nucleotide sequence when in its natural environment and when it is linked to its naturally associated sequence(s) that is/are also in its/their natural environment. For ease of reference, we shall call this preferred embodiment the “native nucleotide sequence”. In this regard, the term “native nucleotide sequence” means an entire nucleotide sequence that is in its native environment and when operatively linked to an entire promoter with which it is naturally associated, which promoter is also in its native environment.

[0402] The nucleotide sequence for use in the present invention may be present in a vector in which the nucleotide sequence is operably linked to regulatory sequences capable of providing for the expression of the nucleotide sequence by a suitable host organism. The constructs for use in the present invention may be transformed into a suitable host cell as described herein to provide for expression of a polypeptide of the present invention. The choice of vector e.g. a plasmid, cosmid, or phage vector will often depend on the host cell into which it is to be introduced. Vectors may be used in vitro, for example for the production of RNA or used to transfect, transform, transduce or infect a host cell.

[0403] In some applications, the nucleotide sequence for use in the present invention is operably linked to a regulatory sequence which is capable of providing for the expression of the nucleotide sequence, such as by the chosen host cell. By way of example, the present invention covers a vector comprising the nucleotide sequence of gene encoding a YIPF-related protein as described herein operably linked to such a regulatory sequence, i.e. the vector is an expression vector. The term “operably linked” refers to a juxtaposition wherein the components described are in a relationship permitting them to function in their intended manner. A regulatory sequence “operably linked” to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences.

[0404] The term “regulatory sequences” includes promoters and enhancers and other expression regulation signals. The term “promoter” is used in the normal sense of the art, e.g. an RNA polymerase binding site. The nucleotide sequence within a construct which encodes a YIPF-related protein may be operably linked to at least a promoter.

[0405] The term “construct”—which is synonymous with terms such as “cassette” or “vector”—includes a nucleotide sequence for use according to the present invention directly or indirectly attached to a promoter.

[0406] An example of an indirect attachment is the provision of a suitable spacer group such as an intron sequence, such as the Sh1-intron or the ADH intron, intermediate the promoter and the nucleotide sequence of the present inven-

tion. The same is true for the term “fused” in relation to the present invention which includes direct or indirect attachment. In some cases, the terms do not cover the natural combination of the nucleotide sequence coding for the protein ordinarily associated with the wild type gene promoter and when they are both in their natural environment. The construct may even contain or express a marker, which allows for the selection of the genetic construct. In some embodiments, a promoter may be operably linked to nucleotide sequence in a construct or vector which is used to modulate the concentration and/or total content of nicotine in a cell or cell culture or tobacco plant or part thereof.

[0407] In some embodiments the promoter may be selected from the group consisting of: a constitutive promoter, a tissue-specific promoter, a developmentally-regulated promoter and an inducible promoter.

[0408] In one embodiment the promoter may be a constitutive promoter.

[0409] A constitutive promoter directs the expression of a gene throughout the various parts of a plant continuously during plant development, although the gene may not be expressed at the same level in all cell types. Examples of known constitutive promoters include those associated with the cauliflower mosaic virus 35S transcript (Odell J T, Nagy F, Chua N H. (1985). Identification of DNA sequences required for activity of the cauliflower mosaic virus 35S promoter. *Nature*. 313 810-2), the rice actin 1 gene (Zhang W, McElroy D, Wu R. (1991). Analysis of rice Act1 5' region activity in transgenic rice plants. *Plant Cell* 3 1155-65) and the maize ubiquitin 1 gene (Comejo M J, Luth D, Blankenship K M, Anderson O D, Blechl A E. (1993). Activity of a maize ubiquitin promoter in transgenic rice. *Plant Molec. Biol.* 23 567-81). Constitutive promoters such as the Carnation Etched Ring Virus (CERV) promoter (Hull R, Sadler J, Longstaff M (1986) (CaMV/35S), figwort mosaic virus 35S promoter. The sequence of carnation etched ring virus DNA: comparison with cauliflower mosaic virus and retroviruses. *EMBO Journal*, 5 (2): 3083-3090).

[0410] The constitutive promoter may be selected from a: a carnation etched ring virus (CERV) promoter, a cauliflower mosaic virus (CaMV 35S promoter), a promoter from the rice actin 1 gene or the maize ubiquitin 1 gene.

[0411] The promoter may be a tissue specific promoter. A tissue-specific promoter is one which directs the expression of a gene in one (or a few) parts of a plant, usually throughout the lifetime of those plant parts. The category of tissue-specific promoter commonly also includes promoters whose specificity is not absolute, i.e. they may also direct expression at a lower level in tissues other than the preferred tissue. Tissue specific promoters include the phaseolin-promoter, legumin b4-promoter, usp-promoter, sbp-promoter, ST-LS1 promoter, B33 (patatin class I promoter). In another embodiment the promoter may be a developmentally-regulated promoter.

[0412] A developmentally-regulated promoter directs a change in the expression of a gene in one or more parts of a plant at a specific time during plant development. The gene may be expressed in that plant part at other times at a different (usually lower) level, and may also be expressed in other plant parts.

[0413] In one embodiment the promoter may be an inducible promoter.

[0414] An inducible promoter is capable of directing the expression of a gene in response to an inducer. In the

absence of the inducer the gene will not be expressed. The inducer may act directly upon the promoter sequence, or may act by counteracting the effect of a repressor molecule. The inducer may be a chemical agent such as a metabolite, a protein, a growth regulator (such as auxin and salicylic acid which activate the OCS promoter), or a toxic element, a physiological stress such as heat, light (such as the soybean SSU promoter), wounding (e.g. the nos, nopaline synthase promoter), or osmotic pressure, or an indirect consequence of the action of a pathogen or pest. A developmentally-regulated promoter might be described as a specific type of inducible promoter responding to an endogenous inducer produced by the plant or to an environmental stimulus at a particular point in the life cycle of the plant. Examples of known inducible promoters include those associated with wound response, such as described by Warner SA, Scott R, Draper J. ((1993) *Plant J.* 3 191-201), temperature response as disclosed by Benfey & Chua (1989) (Benfey, P. N., and Chua, N H. ((1989) *Science* 244 174-181), and chemically induced, as described by Gatz ((1995) *Methods in Cell Biol.* 50 411-424).

[0415] A nucleotide sequence encoding either a protein which has the specific properties as gene encoding YIPF-related protein as defined herein or a protein which is suitable for modification may be identified and/or isolated and/or purified from any cell or organism producing said protein. Various methods are well known within the art for the identification and/or isolation and/or purification of nucleotide sequences. By way of example, PCR amplification techniques to prepare more of a sequence may be used once a suitable sequence has been identified and/or isolated and/or purified.

[0416] In a yet further alternative, the nucleotide sequence encoding the YIPF-related protein may be prepared synthetically by established standard methods, e.g. the phosphoramidite method described by Beaucage et al. (1981) *Tetrahedron Letters* 22, 1859-1869 which is incorporated herein by reference, or the method described by Matthes et al. (1984) *EMBO J.* 3, 801-805 which is incorporated herein by reference. In the phosphoramidite method, oligonucleotides are synthesised, e.g. in an automatic DNA synthesiser, purified, annealed, ligated and cloned in appropriate vectors.

[0417] As used herein, the term “amino acid sequence” is synonymous with the term “polypeptide” and/or the term “protein”. In some instances, the term “amino acid sequence” is synonymous with the term “peptide”. In some instances, the term “amino acid sequence” is synonymous with the term “enzyme”.

[0418] The present invention also encompasses the use of sequences having a degree of sequence identity or sequence homology with amino acid sequence(s) of a polypeptide having the specific properties defined herein or of any nucleotide sequence i.e. a gene encoding a YIPF-related protein (hereinafter referred to as a “homologous sequence (s)”). Here, the term “homologue” means an entity having a certain homology with the subject amino acid sequences and the subject nucleotide sequences. Here, the term “homology” can be equated with “identity”.

[0419] The homologous amino acid sequence and/or nucleotide sequence and/or fragments should provide and/or encode a polypeptide which retains the functional activity and/or enhances the activity of the YIPF-related protein. Typically, the homologous sequences will comprise the same catalytic sites etc. as the subject amino acid sequence

for instance or will encode the same catalytic sites. Although homology can also be considered in terms of similarity (i.e. amino acid residues having similar chemical properties/functions), in the context of the present invention it is preferred to express homology in terms of sequence identity. Homologous sequences typically retain functional domains or motifs.

[0420] In one embodiment, a homologous sequence is taken to include an amino acid sequence or nucleotide sequence which has one, two or several additions, deletions and/or substitutions compared with the subject sequence.

[0421] In one embodiment, homologues of YIPF-related protein SEQ ID No. 3 are provided in Table 1. The present invention extends to the utilisation of homologues listed in Table 1 and to sequences having at least 80% sequence identity thereto.

Sequence Identity

[0422] Sequence identity comparisons can be conducted by eye, or more usually, with the aid of readily available sequence comparison programs. These commercially available computer programs can calculate % homology between two or more sequences. % homology or % identity may be calculated over contiguous sequences, i.e. one sequence is aligned with the other sequence and each amino acid in one sequence is directly compared with the corresponding amino acid in the other sequence, one residue at a time. This is called an “ungapped” alignment. Typically, such ungapped alignments are performed only over a relatively short number of residues.

[0423] Although this is a very simple and consistent method, it fails to take into consideration that, for example, in an otherwise identical pair of sequences, one insertion or deletion will cause the following amino acid residues to be put out of alignment, thus potentially resulting in a large reduction in % homology when a global alignment is performed. Consequently, most sequence comparison methods are designed to produce optimal alignments that take into consideration possible insertions and deletions without penalising unduly the overall homology score. This is achieved by inserting “gaps” in the sequence alignment to try to maximise local homology.

[0424] However, these more complex methods assign “gap penalties” to each gap that occurs in the alignment so that, for the same number of identical amino acids, a sequence alignment with as few gaps as possible-reflecting higher relatedness between the two compared sequences-will achieve a higher score than one with many gaps. “Affine gap costs” are typically used that charge a relatively high cost for the existence of a gap and a smaller penalty for each subsequent residue in the gap. This is the most commonly used gap scoring system. High gap penalties will of course produce optimised alignments with fewer gaps. Most alignment programs allow the gap penalties to be modified. However, it is preferred to use the default values when using such software for sequence comparisons.

[0425] Calculation of maximum % homology therefore firstly requires the production of an optimal alignment, taking into consideration gap penalties. A suitable computer program for carrying out such an alignment is the Vector NTI (Invitrogen Corp.). Examples of software that can perform sequence comparisons include, but are not limited to, the BLAST package (see Ausubel et al. (1999) Short Protocols in Molecular Biology, 4th Ed-Chapter 18),

BLAST 2 (see FEMS Microbiol Lett 1999 174 (2): 247-50; FEMS Microbiol Lett 1999 177 (1): 187-8 and tatiana@ncbi.nlm.nih.gov), FASTA (Altschul et al. 1990 J. Mol. Biol. 403-410) and AlignX for example. At least BLAST, BLAST 2 and FASTA are available for offline and online searching (see Ausubel et al. 1999, pages 7-58 to 7-60).

[0426] Although the final % homology can be measured in terms of identity, the alignment process itself is typically not based on an all-or-nothing pair comparison. Instead, a scaled similarity score matrix is generally used that assigns scores to each pairwise comparison based on chemical similarity or evolutionary distance. An example of such a matrix commonly used is the BLOSUM62 matrix—the default matrix for the BLAST suite of programs. Vector NTI programs generally use either the public default values or a custom symbol comparison table if supplied (see user manual for further details). For some applications, it is preferred to use the default values for the Vector NTI package.

[0427] Alternatively, percentage homologies may be calculated using the multiple alignment feature in Vector NTI (Invitrogen Corp.), based on an algorithm, analogous to CLUSTAL (Higgins DG & Sharp PM (1988), Gene 73 (1), 237-244). Once the software has produced an optimal alignment, it is possible to calculate % homology, preferably % sequence identity. The software typically does this as part of the sequence comparison and generates a numerical result.

[0428] Should gap penalties be used when determining sequence identity, then preferably the following parameters are used for pairwise alignment:

FOR BLAST			
GAP OPEN		0	
GAP EXTENSION		0	

FOR CLUSTAL			
	DNA	PROTEIN	
WORD SIZE	2	1	K triple
GAP PENALTY	15	10	
GAP EXTENSION	6.66	0.1	

[0429] In one embodiment, CLUSTAL may be used with the gap penalty and gap extension set as defined above. In some embodiments the gap penalties used for BLAST or CLUSTAL alignment may be different to those detailed above. The skilled person will appreciate that the standard parameters for performing BLAST and CLUSTAL alignments may change periodically and will be able to select appropriate parameters based on the standard parameters detailed for BLAST or CLUSTAL alignment algorithms at the time.

[0430] Suitably, the degree of identity with regard to a nucleotide sequence is determined over at least 50 contiguous nucleotides, preferably over at least 60 contiguous nucleotides, preferably over at least 70 contiguous nucleotides, preferably over at least 80 contiguous nucleotides, preferably over at least 90 contiguous nucleotides, preferably over at least 100 contiguous nucleotides, preferably over at least 150 contiguous nucleotides, preferably over at least 200 contiguous nucleotides, preferably over at least 250 contiguous nucleotides, preferably over at least 300 contiguous nucleotides, preferably over at least 350 con-

tiguous nucleotides, preferably over at least 400 contiguous nucleotides, preferably over at least 450 contiguous nucleotides, preferably over at least 500 contiguous nucleotides, preferably over at least 550 contiguous nucleotides, preferably over at least 600 contiguous nucleotides, preferably over at least 650 contiguous nucleotides, or preferably over at least 700 contiguous nucleotides.

[0431] Suitably, the degree of identity with regard to a nucleotide, cDNA, cds or amino acid sequence may be determined over the whole sequence.

[0432] The sequences may also have deletions, insertions or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent substance. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues as long as the secondary binding activity of the substance is retained. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values include leucine, isoleucine, valine, glycine, alanine, asparagine, glutamine, serine, threonine, phenylalanine, and tyrosine.

[0433] Conservative substitutions may be made, for example according to the Table below. Amino acids in the same block in the second column and preferably in the same line in the third column may be substituted for each other:

ALIPHATIC	Non-polar	GAP ILV
	Polar-uncharged	CSTM NQ
	Polar-charged	DE KR
AROMATIC		HFYW

[0434] The present invention also encompasses homologous substitution (substitution and replacement are both used herein to mean the interchange of an existing amino acid residue, with an alternative residue) that may occur i.e. like-for-like substitution such as basic for basic, acidic for acidic, polar for polar etc. Non-homologous substitution may also occur i.e. from one class of residue to another or alternatively involving the inclusion of unnatural amino acids such as ornithine (hereinafter referred to as Z), diaminobutyric acid ornithine (hereinafter referred to as B), norleucine ornithine (hereinafter referred to as O), pyrroline, thienylalanine, naphthylalanine and phenylglycine.

[0435] Replacements may also be made by unnatural amino acids include; alpha* and alpha-disubstituted* amino acids, N-alkyl amino acids*, lactic acid*, halide derivatives of natural amino acids such as trifluorotyrosine*, p-Cl-phenylalanine*, p-Br-phenylalanine*, p-I-phenylalanine*, L-allyl-glycine*, beta-alanine*, L-alpha-amino butyric acid*, L-gamma-amino butyric acid*, L-alpha-amino isobutyric acid*, L-epsilon-amino caproic acid #, 7-amino heptanoic acid*, L-methionine sulfone **, L-norleucine*, L-norvaline*, p-nitro-L-phenylalanine*, L-hydroxyproline #, L-thioproline*, methyl derivatives of phenylalanine (Phe) such as 4-methyl-Phe*, pentamethyl-Phe*, L-Phe (4-amino) #, L-Tyr (methyl)*, L-Phe (4-isopropyl)*, L-Tic (1,2,3,4-tetrahydroisoquinoline-3-carboxyl acid)*, L-diaminopropionic acid # and L-Phe (4-benzyl)*. The notation * has been utilised for the

purpose of the discussion above (relating to homologous or non-homologous substitution), to indicate the hydrophobic nature of the derivative whereas #has been utilised to indicate the hydrophilic nature of the derivative, ** indicates amphipathic characteristics.

[0436] Variant amino acid sequences may include suitable spacer groups that may be inserted between any two amino acid residues of the sequence including alkyl groups such as methyl, ethyl or propyl groups in addition to amino acid spacers such as glycine or beta-alanine residues. A further form of variation, involves the presence of one or more amino acid residues in peptoid form, which will be well understood by those skilled in the art. For the avoidance of doubt, "the peptoid form" is used to refer to variant amino acid residues wherein the alpha-carbon substituent group is on the residue's nitrogen atom rather than the alpha-carbon. Processes for preparing peptides in the peptoid form are known in the art, for example Simon et al. (1992) PNAS 89 (20), 9367-9371 and Horwell (1995) Trends Biotechnol. 13 (4), 132-134.

[0437] The nucleotide sequences for use in the present invention may include within them synthetic or modified nucleotides. A number of different types of modification to oligonucleotides are known in the art. These include methylphosphonate and phosphorothioate backbones and/or the addition of acridine or polylysine chains at the 3' and/or 5' ends of the molecule. For the purposes of the present invention, it is to be understood that the nucleotide sequences described herein may be modified by any method available in the art. Such modifications may be carried out in order to enhance the in vivo activity or life span of nucleotide sequences of the present invention. The present invention also encompasses sequences that are complementary to the nucleic acid sequences of the present invention or sequences that are capable of hybridising either to the sequences of the present invention or to sequences that are complementary thereto. The term "hybridisation" as used herein shall include "the process by which a strand of nucleic acid joins with a complementary strand through base pairing" as well as the process of amplification as carried out in polymerase chain reaction (PCR) technologies.

[0438] The present invention also relates to nucleotide sequences that can hybridise to the nucleotide sequences of the present invention (including complementary sequences of those presented herein). Preferably, hybridisation is determined under stringency conditions (e.g. 50° C. and 0.2xSSC {1xSSC=0.15 M NaCl, 0.015 M Na3citrate pH 7.0}). More preferably, hybridisation is determined under high stringency conditions (e.g. 65° C. and 0.1xSSC {1xSSC=0.15 M NaCl, 0.015 M Nascitrate pH 7.0}).

[0439] A review of the general techniques used for transforming plants may be found in articles such as Potrykus et al. (1991) Annu Rev Plant Physiol. Plant Mol. Biol. 42:205-225 and Christou et al. (1994) Agro-Food-Industry Hi-Tech March/April 17-27, which are incorporated herein by reference. Further teachings on plant transformation may be found in EP-A-0449375, incorporated herein by reference.

[0440] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs.

[0441] Singleton, et al., DICTIONARY OF MICROBIOLOGY AND MOLECULAR BIOLOGY, 20 ED., John Wiley and Sons, New York (1994), and Hale & Marham, THE HARPER COLLINS DICTIONARY OF BIOLOGY,

Harper Perennial, NY (1991) provide one of skill with a general dictionary of many of the terms used in this disclosure.

[0442] This disclosure is not limited by the exemplary methods and materials disclosed herein, and any methods and materials similar or equivalent to those described herein can be used in the practice or testing of embodiments of this disclosure. Numeric ranges are inclusive of the numbers defining the range. Unless otherwise indicated, any nucleic acid sequences are written left to right in 5' to 3' orientation; amino acid sequences are written left to right in amino to carboxy orientation, respectively.

[0443] The headings provided herein are not limitations of the various aspects or embodiments of this disclosure which can be had by reference to the specification as a whole. Accordingly, the terms defined immediately below are more fully defined by reference to the specification as a whole. Amino acids are referred to herein using the name of the amino acid, the three letter abbreviation or the single letter abbreviation. The term “protein”, as used herein, includes proteins, polypeptides, and peptides. As used herein, the term “amino acid sequence” is synonymous with the term “polypeptide” and/or the term “protein”. In some instances, the term “amino acid sequence” is synonymous with the term “peptide”. In some instances, the term “amino acid sequence” is synonymous with the term “enzyme”.

[0444] In the present disclosure and claims, the conventional one-letter and three-letter codes for amino acid residues may be used. The 3-letter code for amino acids as defined in conformity with the IUPACIUB Joint Commission on Biochemical Nomenclature (JCBN). It is also understood that a polypeptide may be coded for by more than one nucleotide sequence due to the degeneracy of the genetic code.

[0445] Other definitions of terms may appear throughout the specification. Before the exemplary embodiments are described in more detail, it is to understand that this disclosure is not limited to particular embodiments described, as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present disclosure will be limited only by the appended claims.

[0446] Where a range of values is provided, it is understood that each intervening value, to the tenth of the unit of the lower limit unless the context clearly dictates otherwise, between the upper and lower limits of that range is also specifically disclosed. Each smaller range between any stated value or intervening value in a stated range and any other stated or intervening value in that stated range is encompassed within this disclosure. The upper and lower limits of these smaller ranges may independently be included or excluded in the range, and each range where either, neither or both limits are included in the smaller ranges is also encompassed within this disclosure, subject to any specifically excluded limit in the stated range. Where the stated range includes one or both of the limits, ranges excluding either or both of those included limits are also included in this disclosure.

[0447] It must be noted that as used herein and in the appended claims, the singular forms “a”, “an”, and “the” include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to “an enzyme” or

“YIPF-related protein” includes a plurality of such candidate agents and equivalents thereof known to those skilled in the art, and so forth.

Advantages

[0448] It has been surprisingly found that by modulating the activity or expression of a YIPF-related protein as taught herein which acts as a positive regulator of alkaloid content in tobacco, the alkaloid and/or TSNA precursor content of plants can be modulated. Thereby tobacco industry products with modulated (e.g. reduced) alkaloid content) and/or reduced TSNA precursor content and commercially desirable traits sought after by consumers of tobacco industry products can be produced.

[0449] The present inventors have surprisingly determined a method for modulating the alkaloid content and/or TSNA precursor content of a plant (e.g. tobacco plant) by modulating the activity or expression of a YIPF-related protein. Alkaloid or TSNA precursor content of a plant (e.g. tobacco plant) may be decreased by decreasing or inhibiting the activity or expression of a YIPF-related protein as described herein. Prior to the present invention it had not been known that modulation of the activity or expression of a YIPF-related protein as described herein could be used to modulate alkaloid (and/or TSNA precursor content of a plant (e.g. a tobacco plant).

[0450] The publications discussed herein are provided solely for their disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that such publications constitute prior art to the claims appended hereto.

EXAMPLES

Example 1—Virus-Induced Gene Silencing (VIGS)
of Nitab4.5_0005388g0090.2 Decreases Alkaloid
Content in Leaves

Virus-Induced Gene Silencing (VIGS)

[0451] For virus induced gene silencing, a 300-nucleotide cDNA fragment of Nitab4.5_0005388g0090.2:

(TRV2: SEQ ID No. 16)
ATGATGTCCAGCAATTACACCGCAATTGATAATCAGAATGTTTCC
GGATCTGTTCCTCGAGTCGCAGATCCACCTGGCCAAGTTGCCGTC
AAATTCAGTATCGAATTTCAGACATTTCCACCTTCCAGTTCC
CAGGGGAAATCTCTGGTGCTTCTGGACCTCCACGTGACGCCGAC
GATACATTCTCTAAACAGTATCTGGTTCTGATGAGCCCCAGCAG
AGCGGTTGGTTTCGCGCTTCTCACTATTGCTGCGTACAAGCCCTAT
TTTGATGTTGATACATCAGATGTTCTGGAG

was synthesized and cloned with In-Fusion cloning kit into pTV00 (between EcoRI and XhoI sites). The plasmid was then transformed into *A. tumefaciens* GV3101.

[0452] The TRV vector comprising both (TRV RNA1 SEQ ID No. 15) and (TRV RNA2) comprising the targeted nucleotide sequence were separately propagated in *A. tumefaciens*. These cultures were mixed (1:1) and syringe-infiltration into 2-week-old TN90 plants. The silencing effect

was assessed five weeks post-virus infection by assessing the expression level of the target gene.

Silencing

[0453] VIGS assays were performed as previously described (Ratcliff et al., 2001) Ratcliff, F et al., (2001), The Plant Journal, 25:237-245 (incorporated herein by reference). Briefly, independent cultures of *A. tumefaciens* GV3101 carrying TRV2 and TRV1 plasmids were propagated overnight in LB medium supplemented with appropriate antibiotics. Cultures were resuspended in VIGS buffer (10 mM morpholineethanesulfonic acid pH 5.6, 10 mM MgCL₂, and 100 μM acetocyringone) adjusting optical density to OD₆₀₀=1, and incubated overnight at room temperature in the dark.

[0454] These cultures were mixed (1:1) and syringe-infiltrated into 2-week-old TN90 plants. The silencing effect was assessed two weeks post-virus infection by assessing the expression level of the target gene. TRV-Luciferase was used as a negative control and TRV-PDS (reduced chlorophyll content of the silenced leaves) was used as a phenotypic silencing control.

Alkaloid Measurement

[0455] Relative content of pyridine alkaloids was determined by reversed phase high performance liquid chromatography with tandem mass spectrometry (LC-MS/MS). Chromatographic separation was achieved using a Gemini-NX column (100 mm×3.0 mm, particle size 3 μm, Phenomenex) and gradient chromatographic separation using 6.5 mM ammonium acetate buffer (aq) (pH10) and Methanol. [0456] Mass Spectrometer operates in electrospray (ESI) positive mode using scheduled MRM data acquisition. Two MRM transitions were monitored for each analyte and one for the isotope labelled internal standard.

Analyte	Precursor Ion	Daughter Ion (quant/confirm)
Nicotine	163.1	130/106
Nicotine d4	167.1	134.1
Anabasine	163.1	80/120
Anatabine	161.1	144/80
Nornicotine	149.1	80/130
Nornicotine d4	153.1	84.1
PON	176.1	106.0/148
PON d4	183.1	110.0

Results

[0457] Alkaloid content of plants targeted with VIGS construct for Nitab4.5_0005388g0090.2. Graph is representative of 2 biological replicates (N=12) analysed by one-way ANOVA and Tukey’s multiple-comparison post-test. Values are shown as means±SEM. Asterisks indicate statistical significance of P value≤0.001.

[0458] VIGS of Nitab4.5_0005388g0090.2 leads to a decrease in alkaloid content in leaves, in particular a decrease in nicotine, anabasine, PON and anatabine content.

CONCLUSIONS

[0459] Nitab4.5_0005388g0090.2 is a positive regulator of alkaloid content, in particular alkaloid content in leaves and is a regulator of pyridine alkaloids in tobacco.

Example 2—Homologue Testing

[0460] The effects of the homologues of SEQ ID No. 3, namely those listed in Table 1, are tested in assays as described in the above examples.

SEQUENCE LISTING						
Sequence total quantity: 16						
SEQ ID NO: 1	moltype = DNA length = 5004					
FEATURE	Location/Qualifiers					
source	1..5004					
	mol_type = genomic DNA					
	organism = Nicotiana tabacum					
SEQUENCE: 1						
ggagttggga	cccataaccc	aaagcccaaa	tatgacaaga	tccatcaccc	cttaccgaaa	60
aaaacccaaa	acccaaaaaa	aaaggagaaa	agaaaattaa	aataactaag	gaaacaaaata	120
gaaagatttg	catatacagg	aaaccactcc	agatctgtgt	ccgaaactga	gctgaaagaa	180
caaatttgca	tatacagttt	tatacagaaa	gaaacgaaga	tgatgtccag	caattacacc	240
gcaattgata	atcagaatgt	ttccggatct	gttctctgtaa	ttctttttctc	attttttttg	300
gtttgattga	attttttattc	tggccgtcgt	tcagatctaa	ttgtctaat	tcttatatat	360
tataggcagt	cgcagatcca	cctggccaag	ttgccgtcaa	attcactggt	aatgtcttca	420
gattttcaac	aaaatcatac	agtagtattt	tttgtggatca	ttttgaattg	gtttgtgaaa	480
attttggggt	tatttttgta	ttgcagattc	gaatttgcag	acatttccac	cttccagttc	540
ccaggggaaa	atctctggtg	cttctggacc	tccacgtgac	gccgacggta	tctctccta	600
agttattcat	atatagttac	gacgtaaaga	cgtaatagtg	tgttgctatt	aactaacgtg	660
tggttttcag	ctaatttcag	ctaatttttc	tgcaaaaaaa	agggaaggaa	gggggggggg	720
ggggaagcag	gggtattttg	aatttagaca	tacaagaagt	ggagctagtg	cagcttttta	780
gtttgtcttt	agatctttga	agttcatgat	acaacacttg	tctctataat	agcgatataa	840
acacaccgat	gatagtgtct	aacattgcat	tatgaagggg	ctgtccagca	tgtttcagca	900
tttttttctt	tcaactaagc	tgatgttaac	atctttcact	ctgctaatta	tgctatgact	960
tctttcaaat	tgaatttttt	acttcaaggc	tacaaagcct	ggattcaaat	atacattctt	1020
cctacaatga	ttttgaattc	aaatctgata	ctatatgagg	cagtaggata	ataagaaaga	1080
aattgtttca	ccagtacaaa	attggaaa	cgaatggaat	tttgtgtaaa	cgtataagtc	1140
aaaatgaggt	gaggaagaaa	ccatttttgc	agggtaaaat	aaatacaagg	aagtcacag	1200
atcgggttcg	agggattaca	ataatgtggt	tatgttttagt	ctagggttat	ttagaattta	1260
gacatacaag	aagtgagact	agtgcagctt	tttagtttgt	ctttagatct	ttgaagttca	1320

-continued

tgatacaaca	cttgtctcta	taatagcgat	ataaacacac	cgatgatagt	gcttaacatt	1380
gcattatgaa	ggggctgtcc	agcatgtttc	agcatttttt	ctttcaacta	agctgatgtt	1440
aacatcttct	acactgtctaa	ttatgtctatg	acttctttca	aattgaaatt	tttacttcaa	1500
ggctacaaa	cctggattca	aataacatt	cttctctaca	tgattttgaa	ttcaaatctg	1560
atactataatg	aggcagtagg	atcataagaa	agaaattgtt	tcaccagtac	aaaattggaa	1620
agacgaatgg	aatttttgtg	aaacgtataa	gtcaaaatga	ggtaggaag	aaaccatttt	1680
gccagggtaa	aataaatata	aggaagtccc	aggatcggtt	tcgagggatt	acaataatgt	1740
ggttatgttt	agctaggta	tacctagttc	cgtacaggtt	gtctcatccg	gtgcttcttt	1800
ttgataaggt	aaagtttatt	gatgctggaca	acaaagtgtc	gcttagttgt	atattcaaaa	1860
aagtatgtgc	atcacaaagt	atgcaatgaa	cggaggaagt	ccaaaatggc	ttcgggctct	1920
aacacatttt	ccacagccat	gttgacccaa	aaactaagca	aagatatata	attgtgtttg	1980
agtttgatca	cagtatcttt	cttctcctaa	aaaaattgca	cgtttcatct	gggtgcttct	2040
catagagtta	ttttgtttatc	cttaaaaaga	gttaactctc	aaagcttaac	tgaaggaagt	2100
tgtgagtcac	gctatttgcg	ccttttcatt	aagttattag	caaactctat	tttccctctc	2160
cactttccac	agttcaggtt	gaatttactt	ctctttttag	ataataatta	tgacttggtg	2220
ttgtagatca	attctctaaa	ccagtatctg	gttctgatga	gccccagcag	agcggttggt	2280
ttcgcgcttt	cactattgct	gcgtacaagc	cctattttga	tgtagatata	tcagatgttc	2340
tgagagaaat	aaaagattca	ctctttccct	tcacgggac	cttttcggag	aagaccgcca	2400
ataacccaga	tttgtatgct	cctgctcttt	ttatttctat	gtttattgaa	ttagcattta	2460
ttctcggtgt	ttaatctctaa	gtataactct	tataaaaaact	tattatgagt	agttattttcc	2520
tggaatatata	ggcctttggg	cttctaccgt	tattaaatgg	catttatata	attcttatcc	2580
cattcacctc	tatcaattct	gtccctgtt	ttttcccggt	taccacggat	gaataaagtt	2640
ggcagtttat	aggagaccaa	gaatatatct	tgctcccagt	aacactttct	tcttcttgaa	2700
tgatatcggt	ctaatttctt	attatgggtt	tgtaggagat	gaccgttaat	taatatccct	2760
gtctatcagt	ttcatctctac	aatttctca	ttgatcagct	atttagctct	cgactaagag	2820
gcatttacct	gctaggaggt	tgccaatggg	caaccactat	gcttaacttg	ctcgaattga	2880
ataatgattt	attgtggaat	cttccatgca	ttaaatgcag	gctgtaattg	ggttgtgtga	2940
attcaggcta	atggatagg	aacaattctt	acatttaact	agatggagtt	tgtttcaaat	3000
ctctggcggt	atattttgaa	aattaccatt	aggcatacag	aagaaccaat	aagaataaatt	3060
tggtgtggt	ttcttctgt	ttcgagttg	gaatttgtcc	tgccgattgt	actaaagcta	3120
gtgagttttg	tgcaagtctat	cagatttccc	ttctagaatg	tatgtgcttt	cgttttgatt	3180
ttcttctctg	ttccctatct	tatcttctat	tatataggat	tgtttagaca	tgccgttttg	3240
cacttcagaa	cctgttattt	cttaagtttc	tgttccaaat	ttgtattact	ttttttgggg	3300
gggggggggg	gggggtgggt	tgaaatcttc	cttctgttcc	atcttcttta	tgcatgttcc	3360
tggtttacgt	atgacctgga	tataatagca	catgtttgtt	tattggcaag	cggcagagct	3420
cataatgggt	attattggca	tatttgtttc	atgaaaagtt	atttcagttg	ttgggcttat	3480
gatacattat	gactgtcttt	taagtccttg	ctccttttta	acttgattga	cacttggttt	3540
gctttgtgga	gttgtaactg	cgcagtgtat	atttcaactg	taccttcatg	gacaaaaaag	3600
taataaccat	caaacacact	actaaaagaa	tttaaaaaag	aaaagaacag	tgattagcaa	3660
gtgaagtttt	gccactgctt	cccctctgtt	cgctctttag	ccagctttac	atgctctctg	3720
agtaagaaga	gcctgtttca	gaacttgcc	gcagggaagt	tttgctaaag	gggtgcagtgc	3780
aggaacagaa	atgatgtccg	caaactttgc	atctaaactt	aaactattgc	tagatgctta	3840
tgacgccatt	atctgtactg	ggtggtgtct	agacttcttt	gccccttaag	atcttgtgag	3900
tgatgtttc	aattattttc	atattagcga	acttctctac	cgtaattact	caattttttc	3960
ttttcaggt	tgcccgcttc	tggtatgca	gtaccttggt	atttggtgca	gcttccattg	4020
gcacgtttgt	tacctattta	gcacacaagc	tacagaaaaa	ggagtggaa	tatgacatca	4080
accttgtcac	ttggtctgct	gggtttattct	atggtttatc	cactgtagtt	cctctttgct	4140
tggttctagt	cctcaagtac	ttctctgccc	catctggcct	cgctcagctg	ctgtgccttt	4200
acggctattc	cctttttgtc	ttcattccgg	cattggtaag	tttattta	tacagtatac	4260
tttgttgaa	ttttgggaaa	cctgttgctt	gatgaatcct	ttcattttta	agtttgccat	4320
taggatccat	actaagatct	atgattaa	cgttacagtg	tctgtctctt	gttccctggg	4380
agatagtgag	atgggtgggt	gctggtgtag	ctggctta	gtctgcacac	tttgtcgctc	4440
ttaatctgaa	acaccacata	gtatcggtg	gtgaaaggtg	gtttcttcat	gtggctgcta	4500
cttctcgtgt	gcaacttgca	cttgctttgg	tactgaggtt	ttactgttcc	aacgtcacag	4560
tataactaca	aaattatttg	aatgcgacta	ggctcgggtg	gaacatgtca	tataaataac	4620
agagaaagga	cactgttctt	gggtatcaact	cgactgtaag	tgtgcttggt	cattgaaacg	4680
gaactgtatt	cattcatttt	tgacgatata	caataatata	gtttcatcaa	aagggttcca	4740
gcttgtggtc	ttcttattga	gaatgtcaca	cttttgagct	ttaatagtgg	ctcgggagtt	4800
cgaggggtcga	tgatttagtc	aaacatttgt	gttttagtggg	gatttaattg	taattgtcaag	4860
actgattagc	aaattcctac	aagtccactt	acaagataat	caaaccatgc	ggaataactt	4920
ttgcttgcca	tgatgacctt	tgaatgacct	aattttcaac	atggtatcta	cctattctta	4980
ctttgcttgc	tccattgacg	gtca				5004

SEQ ID NO: 2 moltype = DNA length = 843
FEATURE Location/Qualifiers
source 1..843
 mol_type = genomic DNA
 organism = Nicotiana tabacum

SEQUENCE: 2

atgatgtcca	gcaattacac	cgcaattgat	aatcagaatg	tttccggatc	tggtcctgca	60
gtcgagatc	cactgtgcca	agttgccgtg	aaattcactg	attcgaattt	gcagacattt	120
ccaccttcca	gttcccgagg	gaaaatctct	ggtgcttctg	gacctccacg	tgacgccgac	180
gatacattct	ctaaaccagt	atctggttct	gatgagcccc	agcagagcgg	ttgggttccg	240
gctttcacta	ttgctgcgta	caagccctat	tttgatgttg	atacatcaga	tggttctggag	300
agaataaaag	attcactctt	tcctttccag	ggatcccttt	cggagaagac	cgccaataac	360
ccagatttgt	atggcccgtt	ctggatatgc	agtaccttgg	tatttgttgc	agcttccatt	420

-continued

```

ggcacgtttg ttacctatat agcacacaag ctacagaaaa aggagtggaa ctatgacatc 480
aaccttgtea cttggtctgc ggggtttattc tatggttatt ccaactgtagt tcctctttgc 540
ttggtttctag tctccaagta cttctctgcc ccatctggcc tcgttcagct gctgtgacct 600
tacggctatt cctttttgt cttcattccg gcattgtgtc tgtctctgt tcctgggag 660
atagtgagat ggggtgggttc tgggttagct ggcttaagt ctgccacatt tgtcgtctct 720
aatctgaaac accacatagt atcggctggt gaaagggtgt tcttcattgt ggctgctatc 780
ttcctgttgc aacttgcaact tgccttggtg ctgagggttt acttgttcaa cgtcacagta 840
taa
843

```

```

SEQ ID NO: 3      moltype = AA length = 280
FEATURE          Location/Qualifiers
source           1..280
                 mol_type = protein
                 organism = Nicotiana tabacum

```

```

SEQUENCE: 3
MMSSNYTAID NQNVSGSVPA VADPPGQVAV KFTDSNLQTF PPSQQGKIS GASGPPRDAD 60
DTFSKPVSGS DEQQSGWFR AFTIAAYKPY FDVDTSDVLE RIKDSLFPFT GSFSEKTANN 120
PDLYGPFWIC STLVFVAASI GTFVTYIAHK LQKKEWNYDI NLVTSAGLGF YGYSTVVPLC 180
LFLVLKYFSA PSGLVQLLCL YGYSLFVFIPL ALCLSLVPWE IWRVWVAGVA GLMSATFVAL 240
NLKHHIVSAG ERWFFIVAAI FLLQLALALV LRVYLFNVTV
280

```

```

SEQ ID NO: 4      moltype = DNA length = 4271
FEATURE          Location/Qualifiers
source           1..4271
                 mol_type = genomic DNA
                 organism = unidentified

```

```

SEQUENCE: 4
gaaaaaacca aaaaaccgaag aaaaaaaaag gagaaaagaa aatataaaata actaaggaaa 60
caaatagaaa gagaggagatc acagatctgt gtccgaaact gagctgaaag aataaatttg 120
catatagagt tttatagaga gaagaaacga agatgatgtc cagcaattac accgcaattg 180
ataatcagaa tgtttctgga tctgttctgt taattctttt ctcatttttt tttggtttga 240
ttgaattatt attctggccg tctgttcagat ctaattgtgt aatttcttat atattatagg 300
cagtcgcaga tccacctggc caagtgccg tcaaatcac tggtaagtgc ttcagatttt 360
cgacaaaatc atacagtagt atttttggg atcattttga attggtttgt gaaaattttg 420
ggtttatttt gtgtttgcag attcgaattt gcagacattt ccaccttcca gttcccaggg 480
gaaaatctct ggtgcttctg gacctccacg tgatgctgac ggtatctcct cctaacttat 540
tcataatag ttacgacgta aatagtgtgt tactattaac taacgtgttg tttcagcta 600
aattggagtt agtcagctt ttaagtgtgt ctttagatct ttgaagttca tgatacaaca 660
cttgtctctg taatagtgat ataacacac caatggtagt gcttaaaatt gcgttatgaa 720
gggctgtcca gcatgttcca gcatttttta tttcatctaa gctgatgtta aaatcttaca 780
cacagctaat tatgctatgg cttctttcaa tttgaaattt ttacttcaag gctacaaacc 840
ctaactctga attcaaatat acattcttcc tacaatgatt ttgaattcaa atctgatact 900
atagaggcca gtagagtcac aagaaagaga ttgtttctcc agtacaaaat tggaaagatg 960
aataaaaatt tgtgtaaaata taaagtcaa aatgaggtga ggaagaaacc attttgccag 1020
ggtaaaataa atacaaggaa gtcccaggat ggggttcgag gggattacaat aatgtgggta 1080
tggttagtct aggtataact agttccgtac aagttgtctc atccgggtgcg tctttttgat 1140
agggtaaaag ttgttgatgc gaacagcaaa gtgctgctta gttgtatatt caaaaaagca 1200
tgtgcattac aagtcagtga atgaacagag gaagtatttt ccacagccac gttgcaccaa 1260
aaactaagca aagatagata attgtgtttg agtttgatca cagtatcttt cttctctcaa 1320
aaaatttgca cgtttcatct ggtgcttctt catagagtta tttgtttatc tctaaaaaga 1380
gttaactctc aaagcttaac tgaaggaaat tgtgagtcgt gctattgcgt ccttttcaat 1440
aagttattag cagatctatt ttccctctc cactttccac gctcagggtg aatttacttc 1500
tctttttaga taataattat gacttcttgt tgtatagata ttctctaaac cagtatctgg 1560
ttctgatgag cccagcaga gcgggtgggt tcgcgcttcc actattgctg cgtacaagcc 1620
ctattttgat gttgatacca ctgatgttct agagagaatt aaagattcac tctttccttt 1680
cacgggatcc ttttcggaga agactgccaa taaaccagat ttgtatgctc cctgctcttt 1740
tatttctatg tttattgaat tagcatttat gctcgtgtct taatccgaag tataactcta 1800
ataaaaaact attatgagta gttatttctt ggaagtatag gcctttgggc ttctactggt 1860
attaatggca tttatataat tcttatccca ttcacttcta tcaattctct gccctgtttt 1920
ttcccggtta ccacagatga ataaagtggc cagttatagg agaccaagaa tatatcttgc 1980
tcccactaac actttcttct tcttgaatgt tattgttcta attttttgtt atgatttttg 2040
atgagatgac cgtaaatata tatccctgtc tatcaatttc atcctgcaat attctcatcg 2100
atcagctatt tagtctctga ctaagaggca tttactgtct gggagcttgc caatgaggca 2160
atgagctacc aatgcttaac ttgttcgaat tgaatagtg tttattgttg aaacttccat 2220
gcattaaatg caggctgtgaa tgggggttggt taaattcagg ctaatggata tggacaat 2280
cttacattta actagatgga gtttgtttca aatctctggc ggtatatttt gaaaattacc 2340
attaggcata cagaagaacc aatacgaaaa atttgggttg gtatctttgt cattttcgag 2400
ttggaatttg tctgcgcgat tgtactgaag ctagtgaagt ttgtgcaagt catcagattt 2460
cccttctaga atttatgtgc tttcattttg atttcttctt ctgtccctat atctatctgc 2520
tattatagat gattgtttag acattccgtt tacacttcag aacctgttat ttcttaagt 2580
tctgttccaa atttgtatca cttttttttt ggggtggggg tgggggggtg gaatcattca 2640
gctaatactt cctctgtgtt catcttcttt atgcatgttc ctgtgttacg tatgaccgtg 2700
atataaaagc acatgtttgt ttattggcaa gcggcagagc tcataatggt gattattggc 2760
atatttgttt catgaaaagt tatctcagtt gttgggctta tgatacatta tgactgcttt 2820
taagtccttt gctctttttt aacttgattg tggtttgttg agttggaact 2880
gcgcataatg aatttcaact gtaccttcaa ggacaaaaaa gtaataatca tcaacaacc 2940

```

-continued

tactaaaaga	atttgaaaaa	gaaaagaaca	gttagtagca	agtgaagttt	tgccactgct	3000
gcccctctgt	acatctttag	cccagcttta	catgctctct	gagtaagaag	aagcttggtc	3060
agaacttggc	tgccaggaag	ttttgcttaa	ggttgcagtg	caggaacaga	tatgatgtct	3120
gcaaacgttg	catctaaact	taaaactattg	ctagatgctt	atgcagccat	tatctgtact	3180
ggatgggtgc	tagacttctt	tgtcccttaa	gatgtcatga	gtgtatgttt	caattatttc	3240
catattagcg	aacttctcta	cogtaattac	tcaatttttt	cttttcaggt	atggcccgtt	3300
ctggatatgc	agtaccttgg	tatttgttgc	agcttccatt	ggcacatttg	ttacgtatat	3360
agcacacaag	ctacagaaaa	aggagtggaa	ctacgacatc	aaccttgtca	cttgggtctgc	3420
gggtttatatt	tatggttatt	ccactattgt	tccctcttgc	ttgtttctag	tcctcaagta	3480
cttctctgca	ccatctggcc	tcgttcagct	gctgtgcctt	tacggctatt	cccttttcgt	3540
cttcattccg	gcatttggtta	gtttatttaa	ttgcagtata	ctttgttgaa	cttttgggaa	3600
agctgttgct	tgatgaatcc	tttcactatc	atttttaagt	ttgccattag	gatccatact	3660
aagatctatg	attacctgct	tacagtgtct	gtctcttgtt	ccctggggaga	tagtgagatg	3720
ggtagttgtc	ggtagtagct	gcttcattgc	tgccacattt	gtggctctta	atctgaaaca	3780
ccacatagca	tcggctgggtg	aaaggtgggt	cttcattgtg	gctgctatct	tcctgttgca	3840
acttgcaact	gctttggtag	tgagggttta	cttgttcaac	gtcacagtat	aactacaaaa	3900
ttattggaat	gtgactaggc	acatgtcata	taatcacaga	gaaaggacac	tgttcttcgt	3960
atcaacttga	ctgtaagtgt	gcttggtcat	tgaaacggaa	ctgtattcat	tcatttttga	4020
cgatatacaa	taataatagt	tcatacaaaag	gttctcagct	tgtgtcttct	tttttagaat	4080
gtcacacttt	tgagctttaa	tagtggtctg	cgagttccag	gggtcgatga	tttactcaaa	4140
catttgtgtt	tagtggggat	ttaattgtaa	tgtcaagatt	gggttagcaa	ttcttacaag	4200
tcacttaga	agataatcaa	accatgcgga	aatacttttg	cttgccatga	tgacctgaa	4260
tgacctaat	t					4271

SEQ ID NO: 5 moltype = DNA length = 843
 FEATURE Location/Qualifiers
 source 1..843
 mol_type = genomic DNA
 organism = unidentified

SEQUENCE: 5

atgatgtcca	gcaattacac	cgcaattgat	aatcagaatg	tttctggatc	tgttcctgca	60
gtcgcagatc	cacctggcca	agttgccgtc	aaattcactg	attcgaattt	gcagacattt	120
ccaccttcca	gttcccagg	gaaatctctt	ggtgcttctg	gacctccacg	tgatgctgac	180
gatacattct	ctaaaccagt	atctggttct	gatgagcccc	agcagagcgg	ttgggttcgc	240
gctttcacta	ttgctgcgta	caagccctat	tttgatgttg	ataccactga	tgttctagag	300
agaattaaag	attcactctt	tcctttccacg	ggatcctttt	cggagaagac	tgccaataaa	360
ccagatttgt	atggcccgtt	ctggatatgc	agtagccttg	tatttgttgc	agcttccatt	420
ggcacatttg	ttacgtatat	agcacacaag	ctacagaaaa	aggagtggaa	ctacgacatc	480
aaccttgtca	cttgggtctgc	gggtttatatt	tatggttatt	ccactattgt	tcctctttgc	540
ttgtttctag	tcctcaagta	cttctctgca	ccatctggcc	tcgttcagct	gctgtgcctt	600
tacggctatt	cccttttcgt	cttcattccg	gcattgtgtc	tgtctcttgt	tcctggggag	660
atagttagat	gggtagttgc	tggtgttagct	ggcttcattg	ctgcccacatt	tggtgctctt	720
aatctgaaac	accacatcag	atcggctggt	gaaaggtggt	tcctcattgt	ggctgctatc	780
ttcctgttgc	aacttgcact	tgctttggta	ctgagggttt	acttgttcaa	cgtcacagta	840
taa						843

SEQ ID NO: 6 moltype = AA length = 280
 FEATURE Location/Qualifiers
 source 1..280
 mol_type = protein
 organism = unidentified

SEQUENCE: 6

MMSSNYTAID	NQNVSGSVPA	VADPPGQVAV	KFTDSNLQTF	PPSSSQKIS	GASGPPRDAD	60
DTFSKPVS	DEPQSGWFR	AFTIAAYKPY	FDVDTDVLE	RIKDSLFPFT	GSFSEKTANK	120
PDLYGFWIC	STLVFVAASI	GTFVTYIAHK	LQKEWNYDI	NLVTWSAGLF	YGSTIVPLC	180
LFLVLKYFSA	PSGLVQLLCL	YGYSLFVFI	ALCLSLVPWE	IVRWVAVGA	GFMSATFVAL	240
NLKHIIASAG	ERWFFIVAAI	FLLQLALALV	LRVYLFNVTV			280

SEQ ID NO: 7 moltype = DNA length = 720
 FEATURE Location/Qualifiers
 source 1..720
 mol_type = genomic DNA
 organism = Nicotiana tabacum

SEQUENCE: 7

atggcaatgg	aaatgaatcc	agctgacgaa	accttgtttt	tctccgactc	tcattctcctt	60
gaatcgataa	agcaacatct	tcttgacgat	tcagattttt	ctgaaatttt	ttcgtcgatg	120
aattctagca	acgaaatatt	gcctaacagt	cctagctcaa	gttttagcag	cttcgacttc	180
gactgcagct	tccttaattg	ggatgaaaac	tctgaggaaa	cattaatacc	aactgatcag	240
aatccttcac	atgaatccca	tgaagagtac	tccgagtcgg	aggagaaaac	ccaggggcct	300
gggggtggcg	gtgagaaaaa	cgcgcgcgca	gattggagcg	ggatcatagg	agtgaacagg	360
cgaccgtggg	ggacgttctt	ggcggagaca	agagacccaa	gtaggaaaag	tgaaggtgca	420
aggctgtggt	taggaactta	cgagacgcga	gaggatgcag	cgtagactta	cgatcaagcc	480
gctttcaaaa	tccgcgctc	gagagctcgg	ctcaattttc	ctcacttaat	cggtccaaac	540
atgcctaagc	cgcttagagt	tacagcgagg	cgtagtctga	cgcgctcacc	cgagccatcg	600
tcctcttcac	ccactcatc	atcagaaaaa	gtaccaagga	aaaggaatat	agatgtgata	660
aattccatag	ccaaagccaa	attcctttgt	catagcttga	atttacagag	attagcttaa	720

-continued

```

SEQ ID NO: 8           moltype = DNA length = 720
FEATURE               Location/Qualifiers
source                1..720
                     mol_type = other DNA
                     organism = synthetic construct

SEQUENCE: 8
atggcaatgg aaatgaatcc agctgacgaa accttgtttt tctccgactc tcatctcctt 60
gaatcgataa agcaacatct tcttgacgat tcagattttt ctgaaatttt ttcgtcgatg 120
aattctagca acgaaatatt gcctaacagt cctagctcaa gttttagcag cttcgacttc 180
gactgcagct tccttaattg ggatgaaaac tctgaggaaa cattaatacc aactgatcag 240
aatccttcac atgaatccca tgaaaagtac tccgagtccg aggagaaaac ccagggccct 300
ggggtggcgc gtgagaaaaa cgcgcgcgca gattggacgc ggtacatagg agtgaaacgg 360
cgaccgtggg ggacgttttc gccggagaca agagacccaa gtaggaaagg tgaaggtgca 420
aggctgtggt taggaactta cgagaccgca gaggatgcag cgttagctta cgatcaagcc 480
gctttcaaaa tccgcggctc gagagctcgg ctcaattttc ctcaactaat cggtcctaac 540
atgcctaagc cggctagagt tacagcgagg cgtagtcgta cgcgctcacc cgagccatcg 600
tcttcttcac ccacctcatc atcagaaaat gtaccaagga aaaggaatat agatgtgata 660
aattccatag ccaaagccaa attcctttgt catagcttga atttacagag attagcttaa 720

```

```

SEQ ID NO: 9           moltype = DNA length = 720
FEATURE               Location/Qualifiers
source                1..720
                     mol_type = genomic DNA
                     organism = Nicotiana tabacum

SEQUENCE: 9
atggcaatgg aaatgaatcc agctgacgaa accttgtttt tctccgactc tcatctcctt 60
gaatcgataa agcaacatct tcttgacgat tcagattttt ctgaaatttt ttcgtcgatg 120
aattctagca acgaaatatt gcctaacagt cctagctcaa gttttagcag cttcgacttc 180
gactgcagct tccttaattg ggatgaaaac tctgaggaaa cattaatacc aactgatcag 240
aatccttcac atgaatccca tgaaaagtac tccgagtccg aggagaaaac ccagggccct 300
ggggtggcgc gtgagaaaaa cgcgcgcgca gattggacgc ggtacatagg agtgaaacgg 360
cgaccgtggg ggacgttttc gccggagaca agagacccaa gtaggaaagg tgaaggtgca 420
aggctgtggt taggaactta cgagaccgca gaggatgcag cgttagctta cgatcaagcc 480
gctttcaaaa tccgcggctc gagagctcgg ctcaattttc ctcaactaat cggtcctaac 540
atgcctaagc cggctagagt tacagcgagg cgtagtcgta cgcgctcacc cgagccatcg 600
tcttcttcac ccacctcatc atcagaaaat gtaccaagga aaaggaatat agatgtgata 660
aattccatag ccaaagccaa attcctttgt catagcttga atttacagag attagcttaa 720

```

```

SEQ ID NO: 10          moltype = AA length = 239
FEATURE               Location/Qualifiers
source                1..239
                     mol_type = protein
                     organism = Nicotiana tabacum

SEQUENCE: 10
MAMEMNPAD ETLFFSDSHLL ESIIQHLLDD SDFSEIFSSM NSSNEILPNS PSSSFSSFDF 60
DCSFLNWDEN SEETLIPTDQ NPSHESHEKY SESEKTKQGP GVAREKNAPR DWTRYIGVKR 120
RPWGTFS AET RDP SRKGE GA RLWLGT YETA EDAA LAYDQA AFKIRGSRAR LNFPHLIGSN 180
MPKPARV TAR RSRTRSPEPS SSSSTSSSEN VPRKRNIDVI NSIAKAKFLC HSLNLQRLA 239

```

```

SEQ ID NO: 11          moltype = DNA length = 699
FEATURE               Location/Qualifiers
source                1..699
                     mol_type = genomic DNA
                     organism = Nicotiana tabacum

SEQUENCE: 11
atggaaatga atctagctga cgaacacctg tttttctctg agtctcatct ccttgaatcg 60
ataaagcaac atcttcttga tgattcagat ttttctgaaa ttttttcgcc gatgagttca 120
agcaacgaaa tattgcctaa cagtcctagc tcaagtttta gcagcttcga ctgcagcttc 180
ctcaattggg atgaaaactt tgaggaaaca ttaataccaa ctgatcaaaa tccttcacat 240
gagaagtgc tccagctccga ggagcaaaacc cagggccccag cgggtggcg tgagaaaaac 300
gcccgcgcgag attggacgcg gtatatagga gtgaaacggc ggccgtgggg gacgttttcg 360
gcggagacaa gagacccaag taggaaaggt gaaggtgcaa ggctgtggtt aggaacttac 420
gagaccgcag aggatgcagc gttggcttac gatcaagccg ctttcaaaa cgcgggctcg 480
agagctcgcc tcaatttttc ccacttaatt ggctcaaaaca tgccctaaagg ggctagagta 540
acagcgaggc gtagtctgtac gcgctcacc gagccatcct cttcttcac cacttcatca 600
tcagaaaatg tgccaagaaa aaggaatata gatgtgataa attccatagc caaagccaaa 660
ttcctttgtc atagcttaaa tttacagaga ttagcttaa 699

```

```

SEQ ID NO: 12          moltype = DNA length = 699
FEATURE               Location/Qualifiers
source                1..699
                     mol_type = other DNA
                     organism = synthetic construct

SEQUENCE: 12
atggaaatga atctagctga cgaacacctg tttttctctg agtctcatct ccttgaatcg 60

```

-continued

ataaagcaac	atcttcttga	tgattcagat	ttttctgaaa	tttttctgcc	gatgagttca	120
agcaacgaaa	tattgcctaa	cagtcctagc	tcaagtttta	gcagcttcga	ctgcagcttc	180
ctcaattggg	atgaaaactt	tgaggaaaca	ttaataccaa	ctgatcaaaa	tccttcacat	240
gagaagtgc	ccgagtcgga	ggagcaaac	cagggcccag	cggtggtgcg	tgagaaaaac	300
gcgcgcgag	attggacgcg	gtatatagga	gtgaaacggc	ggccgtgggg	gacgttttcg	360
gcggagacaa	gagaccaag	taggaaaggt	gaaggtgcaa	ggctgtggtt	aggaacttac	420
gagaccgcag	aggatgcagc	gttggtttac	gatcaagccg	ctttcaaaat	ccgcggctcg	480
agagctcggc	tcaattttcc	ccacttaatt	ggctcaaaac	tgccaaagcc	ggctagagta	540
acagcgaggc	gtagtcgtac	gcgctcacc	gagccatcct	cttcttcac	cacttcacat	600
tcagaaaatg	tgccaagaaa	aaggaatata	gatgtgataa	attccatagc	caaagccaaa	660
ttcctttgtc	atagcttaaa	tttacagaga	ttagcttaa			699

SEQ ID NO: 13 moltype = DNA length = 699
 FEATURE Location/Qualifiers
 source 1..699
 mol_type = genomic DNA
 organism = Nicotiana tabacum

SEQUENCE: 13

atggaatga	atctagctga	cgaaaccttg	ttttctctg	agtctcatct	ccttgaatcg	60
ataaagcaac	atcttcttga	tgattcagat	ttttctgaaa	tttttctgcc	gatgagttca	120
agcaacgaaa	tattgcctaa	cagtcctagc	tcaagtttta	gcagcttcga	ctgcagcttc	180
ctcaattggg	atgaaaactt	tgaggaaaca	ttaataccaa	ctgatcaaaa	tccttcacat	240
gagaagtgc	ccgagtcgga	ggagcaaac	cagggcccag	cggtggtgcg	tgagaaaaac	300
gcgcgcgag	attggacgcg	gtatatagga	gtgaaacggc	ggccgtgggg	gacgttttcg	360
gcggagacaa	gagaccaag	taggaaaggt	gaaggtgcaa	ggctgtggtt	aggaacttac	420
gagaccgcag	aggatgcagc	gttggtttac	gatcaagccg	ctttcaaaat	ccgcggctcg	480
agagctcggc	tcaattttcc	ccacttaatt	ggctcaaaac	tgccaaagcc	ggctagagta	540
acagcgaggc	gtagtcgtac	gcgctcacc	gagccatcct	cttcttcac	cacttcacat	600
tcagaaaatg	tgccaagaaa	aaggaatata	gatgtgataa	attccatagc	caaagccaaa	660
ttcctttgtc	atagcttaaa	tttacagaga	ttagcttaa			699

SEQ ID NO: 14 moltype = AA length = 231
 FEATURE Location/Qualifiers
 source 1..231
 mol_type = protein
 organism = Nicotiana tabacum

SEQUENCE: 14

MEMNLADETL	FFSESHLLES	IKQHLDDSD	FSEIFSPMSS	SNEILPNSPS	SSFSSFDCSF	60
LNWDENFEET	LIPDQNSH	EKSESEEEQT	QGPVAVREKN	APRDWTRYIG	VKRRPWGTF	120
AETRDPSRKG	EGARLWLGT	ETAEDALALY	DQAAFKIRGS	RARLNFPHLI	GSNMPKPARV	180
TARRSRTRSP	EPSSSSSTSS	SENVPRKRNI	DVINSIAKAK	FLCHSLNLQR	L	231

SEQ ID NO: 15 moltype = DNA length = 20171
 FEATURE Location/Qualifiers
 source 1..20171
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 15

gatggtaaga	agaaaggcgc	gcagtatgcy	atagctcttc	acagcctgta	tgacttcaag	60
ttgaaagact	tgatggctac	tatgggttag	aagaaaacta	aagtgggttc	tgctgctatg	120
ctttttgtct	ctgaaagtat	gttagtgagc	gaaggtccat	taccttctgt	tgacggttac	180
tacatgaaga	agaacgggaa	gactatcttc	ggttttgaga	aagatccctc	cttttcttac	240
attcatgact	gggaagagta	caagaagtat	ctactgggga	agccagttag	ttaccaaggg	300
aatgtgttct	acttcgaacc	gtggcagggt	agaggagaca	caatgctttt	ttcgatctac	360
aggatagctg	gagttccgag	gaggtctcta	tcactcgcaag	agtactaccg	aagaatatat	420
atcagtagat	gggaaagcat	gggtgtgttc	ccaattttcg	atctgggtcg	atcaacgcga	480
gagttggtca	agaaagacct	gtttgttaga	aaacaattca	tggaacaagt	ttgggattac	540
atagctaggt	tatctgacca	gcagctgacc	ataagcaatg	ttaaatcata	cttgagttca	600
aataattggg	tcttattcat	aaacggggcg	gccgtgaaga	acaagcaaa	tgtagattct	660
cgagatttac	agttgttggt	tcaaaacttg	ctagtgaagg	aacaagtggt	gagacctgtc	720
atgagggagt	tcggtggaag	aattctgact	gagacgaaac	ctatcacgtc	attgactgat	780
gtgctggggt	taatatcaag	aaaactgtgg	aagcagtttg	ctaacaagat	cgagctcggc	840
ggattcggtg	gcattggttg	tactctaatt	ggattctatc	caaagaaggt	actaacctgg	900
gcgaaggaca	caccaaatgg	tccagaacta	tgttacgaga	actcgcaaca	aaccaaggtg	960
atagtatttc	tgagtgttgt	gtagtccatt	ggaggaatca	cgcttatgcy	tcgagacatc	1020
cgagatggac	tggtgaaaaa	actatgtgat	atgtttgata	tcaaacgggg	ggcccatgct	1080
ttagacgttg	agaatccgtg	ccgctattat	gaaatcaacg	atctctttag	cagctctgat	1140
tcggcatctg	agtcocgtga	gaccgtttta	ccagatttat	ccgaggtaaa	agccaagtct	1200
gataagctat	tcgacgagaa	gaaagaaatc	gctgacgagt	ttctaagtcg	aaaattctct	1260
aactattctg	gcagttcggt	gagaacttct	ccaccatcgg	tggtcggttc	atctcgaaagc	1320
ggactgggtc	tgtgtgtgga	agacagtaac	gtgctgacct	aagctagagt	tgaggtttca	1380
agaaaggtag	tcgatgagga	gatcatggag	cagtttctga	gtgttcttat	tgacactgaa	1440
gcagaaattg	acgaggttgt	ttcagccttt	tcagctgaat	gtgaaagagg	ggaacaagc	1500
ggtacaaaag	tggtgtgtaa	acctttaacg	ccaccaggat	ttgagaacgt	ggtgccagct	1560
gtcaaacctt	tggtcagcaa	aggaaaaaac	gtcaaacgtg	tcgattactt	ccaagtgtat	1620
ggaggtgaga	gattacaaaa	aaggccggtt	gtcagtgagg	acgattctgt	ggacgctaga	1680

-continued

agagagtttc	tgtactactt	agatgcggag	agagtcgctc	aaaatgatga	aattatgtct	1740
ctgtatcgtg	actattcgag	aggagttatt	cgaactggag	gtcagaatta	cccgacgga	1800
ctgggagtg	gggatgtgga	gatgaagaac	tgggtgcatac	gtccagtggt	cactgaacat	1860
gcttatgtgt	tccaaccaga	caaacgtatg	gatgattggt	cgggatactt	agaagtggct	1920
gtttgggaac	gaggtatggt	ggtcaacgac	tccgcggtcg	aaaggatgag	tgattatgtc	1980
atagtttgcg	atcagacgta	tctttgcaat	aacaggtaat	aatcctctct	cttgatattt	2040
ttaaattata	gaattaatta	gtttacttta	ttctttacta	tatgatttaa	atagtttaat	2100
cttgtttttg	agtaaacat	tcgattttga	tatttgtatt	cgtcctacaa	agttggaaat	2160
actgatgata	ttttcttttg	aacgtgatac	ctaccaatac	taatcttacg	gaatctttta	2220
atagagcact	aatcaacatg	gaactaaaga	ccaattctta	agtgtctctg	ttgtacagtt	2280
cattttagta	gtgcgtttaa	gtattattat	ctcccttcac	gcggggcaat	tatgtagatt	2340
aaaatcgaaa	ttatataaaa	tttacataag	tctaagtcta	gggtctccag	ctaattgtta	2400
tttttttaac	gatgttgact	aaagcaataa	cgacgttgac	ttgtgtttaa	caggttgatc	2460
ttggacaatt	taagtgcctc	ggatctagga	ccagtttaact	gttcttttga	attagttgac	2520
ggtgtacctg	gttgtgttaa	gtcgacaatg	attgtcaact	cagctaatcc	ttgtgtcgat	2580
gtggttctct	ctactgggag	agcagcaacc	gacgacttga	tcgagagatt	cgcgagcaaa	2640
ggttttccat	gcaaatgaa	aaggagagtg	aagacggttg	attctttttt	gatgcattgt	2700
gtcgatggtt	ctttaaccgg	agacgtgttg	catctcgacg	aagctctcat	ggcccatgct	2760
ggtaggtgtg	acttttgcgc	tcagatagct	ggtgctaaac	gatgtatctg	tcaaggagat	2820
cagaatcaaa	tttctttcaa	gcttagggta	tctcaagttg	atttgaggtt	ttctagtctg	2880
gtcggaaagt	ttgacattgt	tacagaaaaa	agagaaactt	acagaagtcc	agcagatgtg	2940
gctgcccgtat	tgaacaagta	ctatactgga	gatgtcagaa	cacataacgc	gactgtcaat	3000
tcgatgacgg	tgaggagaat	tgtgtctaaa	gaacaggttt	ctttgaagcc	tggtgctcag	3060
tacataaact	tccttcagtc	tgagaagaag	gagttggtaa	atttggtggc	attgagggaa	3120
gtggcagcta	aagttagtat	agtacacgag	tcgcaaggag	agacattcaa	agatgtagtc	3180
ctagtcagga	cgaaacctac	ggatgactca	atcgctagag	gtcgggagta	cttaactcgt	3240
gcattgtcgc	gtcacacaca	atcacttgtg	tatgaaactg	tgaagaggga	cgatgtaagc	3300
aaagagatca	gggaaagtgc	cgcgcttacg	aaggcggctt	tggcaagatt	tttgttact	3360
gagaccgtct	tatgacgggt	tcggcttagg	tttgatgtct	ttagacatca	tgaaggccct	3420
tgccgctgtc	cagattcagg	tacgattacg	gacttgagga	tgtggtacga	cgcttgtttt	3480
ccgggaaatt	cgtaagaga	ctcaagccta	gacgggtatt	tggtggcaac	gactgattgc	3540
aatttgcgat	tgacaatgt	tacgatcaaa	agtggaact	ggaaagacaa	gtttgcgtga	3600
aaagaaactg	ttctgaaacc	ggttattcgt	actgctatgc	ctgacaaaag	gaagactact	3660
cagttggaga	gtttgttagc	attgcagaaa	aggaaccaag	cggcacccga	tctacaagaa	3720
aatgtgcacg	cgacagttct	aatcgaaag	acgatgaaga	agctgaaatc	tgttgtctac	3780
gatgtgggaa	aaattcgggc	tgatcctatt	gtcaatagag	ctcaaatgga	gagatgggtg	3840
agaaatcaaa	gcacagcgg	acaggctaag	gtagtgcag	atgtgagaga	gttacctgaa	3900
atagactatt	cgtcttacat	gtatatgac	aaatctgacg	tgaaccttaa	gactgattta	3960
acaccgcaat	ttgaatactc	agctctacag	actgttgtgt	atcacgagaa	gttgatcaac	4020
tcgttgttgc	gtccaatttt	caagaaat	aatgaacgca	agttggatgc	tatgcaacca	4080
caatttgtgt	tcaacacgag	aatgacatcg	agtgatttaa	acgatcgagt	gaagtcttta	4140
aatacggaag	cggcttacga	ctttgttgag	atagacatgt	ctaaattcga	caagtccgca	4200
aatcgcttcc	atttacaact	gcagctggag	atttacaggt	tatttgggct	ggatgagttg	4260
gcgcccttcc	tttgggaggt	gtcgcacact	caaactactg	tgagagatat	tcaaaatggt	4320
atgatggcgc	atatttggta	ccaacaaaag	agtgagatg	ctgatactta	taatgcaaat	4380
tcagatagaa	cactgtgtgc	actcttgtct	gaattacat	tgagaaaagc	agtcactggt	4440
acatatggag	gagatgactc	actgattgct	tttcttagag	gaacgcagtt	tgttgatccg	4500
tgtccaaagt	tggtacttaa	gtggaatttc	gagtgcaaga	tttttaagta	cgatgtccca	4560
atgttttgtg	ggaagtctct	gcttaagacg	tcatcgtgtt	acgagttcgt	gccagatccg	4620
gtaaaagtgc	tgacgaagtt	ggggaaaaag	agtataaagg	atgtgcaaca	tttagccgag	4680
atctacatct	cgctgaatga	ttccaataga	gctcttggga	actacatggt	ggatcccaaa	4740
ctgtccgagt	ctgtttcaga	ccgggtattg	tacaaaagggt	attctgttca	tgccgtttgt	4800
gcgctatgga	agcatattaa	gagttttaca	gctctgtgta	cattattccg	agacgaaaac	4860
gataaggaat	tgaaccggcg	taaggttgat	tggaagaagg	cacagagagc	tgtgtcaaac	4920
ttttacgact	ggtaatatgg	aagacaagtc	attggtcacc	tgaagaaga	agactttcga	4980
agtctcaaaa	ttctcaaatc	taggggcat	tgaattgttt	gtggacggta	ggagggaagag	5040
accgaagtat	tttcacagaa	gaagagaaac	tgtcctaact	catgttggtg	ggaagaagag	5100
tgaacacaag	ttagacgttt	ttgaccaaag	ggattacaaa	atgattaaat	cttacgcgtt	5160
tctaaaggta	gtagggtgac	aactagttgt	aacatcacat	ctacctgcag	atacgccctg	5220
gttcattcaa	atcgatctgt	tggtatcgag	acttactgag	aaaagaaaga	gagggaagac	5280
tattcagaga	ttcaaaagtc	gagcttgcca	taactgttca	gttgccgagt	acaaggttga	5340
atacagattt	tccacacagg	agaacgtact	tgatgtctg	aaggtgggtt	gtatttctga	5400
gggcgttccg	gtctgtgacg	gtacataccc	tttcagtatc	gaagtgtcgc	taatatgggt	5460
tgctactgat	tcgactaggc	gcctcaatgt	ggaagaactg	aacagttcgg	attacattga	5520
agcgattttt	accgatcaag	aggttttcgg	tgagttcatg	tccttgaaac	aagttggagat	5580
gaagacgatt	gaggcgaagt	acgatggtcc	ttacagacca	gctactacta	gacctaaagt	5640
attattgtca	agtgaagatg	ttaagagagc	gtctaataag	aaaaactcgt	cttaatgcat	5700
aaagaaattt	attgtcaata	tgacgtgtgt	actcaagggt	tgtgtgaatg	aagtcactgt	5760
tcttgggtcac	gagacgtgta	gtatcgggtc	tgctaacaaa	ttgcgaaagc	aagttgtctga	5820
catggttggt	gtcacacgta	ggtgtgcgga	aaataattgt	ggatgggttg	tctgtgttgt	5880
tatcaatgat	tttacttttg	atgtgtataa	ttgtgtggcg	cgtagtccac	ttgaaaagt	5940
tcgtaaacgt	gttgaaacaa	gaaatcgaga	aatttggaaa	caaattcgac	gaaatcaagc	6000
tgaaaacatg	tctgcgacag	ctaaaaagtc	tcataattcg	aagacctcta	agaagaaatt	6060
caaagaggac	agagaatttg	ggacacccaa	aagattttta	agagatgatg	ttcctttcgg	6120
gattgatcgt	ttgtttgcctt	tttgattttta	ttttatattg	ttatctgttt	ctgtgtatag	6180
actgttttag	attggcgctt	ggccgactca	ttgtcttacc	ataggggaac	ggacttttgt	6240

-continued

tgtgttgta	ttttattt	atttattaa	aattctcaat	gatctgaaa	ggcctcgagg	6300
ctaagagatt	attggggggt	gagtaagtag	ttttaaagtg	atgatgggta	caaaggcaaa	6360
aggggtaaaa	cccctcgcc	acgtaagcgt	tattacgccc	ggatccccg	gggagctcga	6420
attcgctgaa	atcacccagtc	tctctctaca	aatctatctc	tctctatttt	ttccataaat	6480
aatgtgtgag	tagtttccc	ataaggga	ttaggggtct	tataggggtt	cgctcatgtg	6540
tfagagcatat	aagaaaccc	tagtatgtat	ttgtatttgt	aaaatacttc	tattatcaat	6600
aaaatttcta	attcctaaaa	caaaaatcca	gtactaaaat	ccagatctcc	taaagtcctc	6660
atagatcttt	gtcgtgaata	ttaaccagac	acgagacgac	taaaacctga	gcccagacgc	6720
cgttcgaagc	tagaagtagc	gcttaggcag	gaggccgtta	gggaaaagat	gctaaggcag	6780
ggttgggtac	gttgactccc	ccgtagggtt	ggtttaata	tgatgaagtg	gacggaagga	6840
aggaggaaga	caagggaaga	taagggttga	ggccctgtgc	aaggtaaaga	gatggaaatt	6900
tgatagaggt	acgctactat	acttatacta	tacgctaagg	gaatgcttgt	atttataccc	6960
tataccccct	aataacccct	tatcaattta	agaaataatc	cgcataagcc	cccgttataa	7020
aattgggtatc	agagccatga	ataggcttat	gacaaaaact	caagaggata	aaacctcacc	7080
aaaatacga	agagttctta	actcctaaaga	ttaaagatct	ttcaagatca	aaactagttc	7140
cctcacaccg	gagcatcgca	tatcctcgac	ctgcaggcat	gcaagcttgg	cgtaatcatg	7200
gtcatagctg	tttctgtgtg	gaaattgtta	tccgctcaca	attccacaca	acatacgagc	7260
cggaagcata	aagtgtaaag	cctgggggtg	ctaagttagtg	agctaaactca	cattaattgc	7320
gttgcgctca	ctgcccgtt	tccagtcggg	aaacctgtcg	tgcagctgc	attaatgaat	7380
cgcccaacgc	gcggggagag	gcgggttgcg	tattgggcca	aagacaaaag	ggcgacattc	7440
aaccgattga	gggaggaag	gtaaatattg	acggaaatta	ttcattaaag	gtgaattatc	7500
accgtcacgc	acttgagcca	tttgggaatt	agagccagca	aaatcaccag	tagcaccatt	7560
accattagca	aggccgga	cgtcaccaat	gaaaccatcg	atagcagcac	cgtaatcagt	7620
agcgacagaa	tcaagtttgc	ctttagcgtc	agactgtagc	gcgttttcat	cggtattttc	7680
ggctatagcc	cccttattag	cgtttgccat	cttttcaata	tcaaaatcac	cggaaccaga	7740
gccaccaccg	gaaccgcctc	ctccagagcc	gccaccctca	gaaccgccc	cctcagagcc	7800
accaccctca	gagccgccc	cagaaccacc	accagagccg	ccgcccagcat	tgacaggagg	7860
cccgatctag	taacatagat	gacaccgcgc	gcgataattt	atcctagtgt	gcgcgctata	7920
ttttgttttc	tatcgctgat	ttaattgtata	attgcgggac	tctaatcata	aaaacccatc	7980
tcataaataa	cgcatgcat	tacatgttaa	ttattacatg	cttaacgtaa	ttcaacagaa	8040
atttatatgat	aatcatcgca	agaccggcaa	caggattcaa	tcttaagaaa	ctttattgcc	8100
aaatgtttga	acgatcgggg	atcatccggg	tctgtggcgg	gaactccacg	aaaatatccg	8160
aacgcagcaa	gatatcgccg	tgcatctcgg	tcttgccctg	gcagtcgccc	ccgacgccgt	8220
tgatgtggac	gccgggccc	atcatattgt	cgctcaggat	cggtggcgtg	tgcttgtcgg	8280
ccgttgctgt	cgtaattgata	tccggcacctt	cgaccgcctg	ttccgcagag	atcccgtggg	8340
cgaaagaactc	cagcatgaga	tccccgcgct	ggaggatcat	ccagccggcg	tcccggaaaa	8400
cgattccgaa	gcccaacgtt	tcatagaagg	cgccgggtga	atcgaaatct	cgtagtgga	8460
ggttgggctg	cgcttggtgc	gtcatttcga	acccagagtg	cccgcctcaga	agaactcgtc	8520
aagaaggcga	tagaaggcga	tgcgctgcga	atcgggagcg	gcgataccgt	aaagcacgag	8580
gaagcgggtca	gcccattcgc	gcacaagctc	ttcagcaata	tcacgggtag	ccaacgctat	8640
gtcctgatag	cggtcccgca	gccccagccg	gccacagtcg	atgaatccag	aaaagcggcc	8700
attttccacc	atgatattcg	gcaagcaggc	atcgccatgg	gtcacagcga	gatcatcgcc	8760
gtcgggcatg	atgggcccga	gacctggcga	cagttcgccg	ggcgcgagcc	cctgatgctc	8820
ttcgctccaga	tcatcctgat	cgacaagacc	ggcttccatc	cgagtacgtg	ctcgctcgat	8880
gcgatgtttc	gcttggtgtg	cgaatgggca	ggtagccgga	tcaagcgtat	gcagccgccc	8940
caattgcatca	gcccattcgc	atactttctc	ggcaggagca	aggtgagatg	acaggagatc	9000
ctgccccggc	acttcgccc	atagcagcca	gtcccttccc	gcttcagtga	caacgtcgag	9060
cacagctgcg	caagggaacgc	ccgtcggtgc	cagccacgat	agccgcgctg	cctcgctcgt	9120
cagttcattc	agggcaccgc	acagggtcgt	cttgacaaaa	agaacccggc	gcccctgcgc	9180
tgacagccgg	aacacggcgg	catcagagca	gccgattgtc	tggtgtgccc	agtcatagcc	9240
gaatagcctc	tccaccaagc	cgccgggaga	acctgcgtgc	aatccatctt	gttccaatcat	9300
gcgaaacgat	ccagatccgg	tgagattatg	agtgaaatag	agactcta	agactcta	9360
tggtatccga	ggggaattta	tggaaacgtc	gtggagcatt	tttgacaaga	aatatttgct	9420
agctgatagt	gaccttaggc	gacttttgaa	cgcgcaataa	tggtttctga	cgtagtgctg	9480
tagctcatta	aaactccagaa	accccgccgt	gagtggtctc	ttcaacgttg	cggttctgtc	9540
agttccaaac	gtaaaacggc	ttgtcccgcg	tcatcgccgg	gggtcataac	gtgactccct	9600
taattctccg	ctcatgatca	gatttgtcgt	tcccgccctc	agttttaaact	atcagtggtt	9660
gacaggatata	attggcgggt	aaacctaaaga	gaaaagagcg	tttattagaa	taatcgggata	9720
tttaaaaggg	cgtgaaaagg	tttatccgtt	cgtccatttg	tatgtgcagt	ccaaccacag	9780
ggttccccag	atctggcgcc	ggccagcgag	acgagcaaga	ttggccgccc	cccgaacaga	9840
tccgacagcg	cgcccagcac	aggtgcgcag	gcaaaattgca	ccaacgcata	cagcgccagc	9900
agaatgccat	agtgggcggt	gacgtcgctc	gagtgaaaca	gatcgcgcac	gaggcccgcc	9960
agcaccggca	taatcaggcc	gatgccgaca	gcgtcgagcg	cgacagtgct	cagaattacg	10020
atcaggggta	tgttgggttt	cacgtctggc	ctccggacca	gcctccgctg	gtccgattga	10080
acgcgcggat	tctttatcac	tgataagttg	gtggacatat	tatgtttatc	agtgataaaag	10140
tgtaagcat	gacaaagtgt	cagccgaata	cagtgatccg	tgccgcctcg	gacctgttga	10200
acgaggtcgg	cgtagacggg	ctgacgacac	gcaaaactgg	ggaaacgggt	gggggttcagc	10260
agccggcgct	ttaactggcac	ttcaggaaaca	agccggcgct	gctcgacgca	ctggccgaag	10320
ccatgctggc	ggagaatcat	acgcattcgg	tgccgagagc	cgacgacgac	tgccgctcat	10380
ttctgatcgg	gaatgcccg	agcttcaggc	agggcgctgt	cgctaccgc	gatggcgccg	10440
gcatccatgc	cgccacgcga	ccgggcccac	cgcagatgga	aacggccgac	gcgcagcttc	10500
gcttcctctg	cgaggccggg	ttttcgcccg	gggacgcgct	caatgcgctg	atgacaatca	10560
gctacttcac	tggtggggcc	gtgcttgagg	agcagggccg	cgacagcgat	gcccggcagc	10620
gcccggccac	cgttgaacag	gctccgctct	cgccgctgtt	gcccggcccg	atagacgcct	10680
tcgacgaagc	cggtccggag	agcaggttcg	agcagggact	cgcggtgatt	gtcgatggat	10740
tgccgaaaag	gaggtcgtt	gtcaggaacg	ttgaaggacc	gagaaagggg	gacgattgat	10800

-continued

caggaccgct	gccggagcgc	aaccactca	ctacagcaga	gccatgtaga	caacatcccc	10860
tcccccttct	caccgcgtca	gacgcccgt	gcagcccgt	acgggctttt	tcatgcoctg	10920
ccctagcgtc	caagcctcac	ggcgcgctc	ggcctctctg	gcggccttct	ggcgtcttct	10980
cgctctctcg	ctcactgact	cgctgcgctc	ggtcgttcgg	ctgcggcgag	cggtatcagc	11040
tcactcaaa	gcggtataac	ggttatccac	agaatcaggg	gataacgcag	gaaagaacat	11100
gtgagcaaaa	ggccagcaaa	aggccaggaa	ccgtaaaaag	cccgcgttgc	tggcggtttt	11160
ccataggctc	cgccccctg	acgagcatca	caaaaatcga	cgctcaagtc	agagggtggcg	11220
aaaccgcaca	ggactataaa	gataccaggc	gtttccccct	ggaagctccc	tcgtgcgctc	11280
tcctgttccg	accctgcgcg	ttaccggata	cctgtccgcc	tttctccctt	cggaagcgt	11340
ggcgcttttc	cgctgcataa	ccctgcttcg	gggtcattat	agcgattttt	tcggtatctc	11400
catccttttt	cgacagatat	acaggatttt	gcccagggtt	tcgtgtagac	tttcttgggt	11460
gtatccaacg	cgctcagcgc	ggcaggatag	gtgaagttag	cccaccgcgc	agcgggtggt	11520
ccttcttcac	tgtcccttat	tcgcacctgg	cggtgctcaa	cggaatctct	gctctgcgag	11580
gctggccggc	taccgcggcg	gtaacagatg	agggcaagcg	gatggctgat	gaaaccaagc	11640
caaccaggaa	gggcagccca	cctatcaagg	tgtactgcct	tcagagcga	cgaagagcga	11700
ttgaggaaaa	ggcggcgggc	gcccgcctga	gcctgtcggc	ctacctgctg	gcccgcggcc	11760
agggctacaa	aatcacgggc	gtcgtggact	atgagcacgt	ccgcgagctg	gcccgcacac	11820
atggcgacct	gggcgcctcg	ggcggcctgc	tgaactctcg	gctcaccgac	gacccgcgca	11880
cggcgcggtt	cggtgatgac	acgatcctcg	ccctgctggc	gaagatcgaa	gagaagcagg	11940
acgagcttgg	caaggctcatg	atgggcgtgg	tcgcgccgag	ggcagagcca	tgactttttt	12000
agccgctaaa	acggccgggg	ggtgcgcgtg	attgccaaag	acgtcccatc	gcgctccatc	12060
aagaagagcg	acttcgcgga	gctggtgaag	tacatcacgc	acgagcaagg	caagaccgag	12120
cgcccttgcg	acgctcacgc	ggctggttgc	cctgcgcgct	gggctggcgg	ccgtctatgg	12180
ccctgcacac	gcgcagaaaa	cgccgtcgaa	gcctgtgctg	agacaccgcg	gcccgcggcg	12240
ttgtggatag	ctcgcgaaa	acttggccct	cactgacaga	tgagggcgcg	acgttgacac	12300
ttgagggggc	gactcacccg	gcgcggcgtt	gacagatgag	gggcaggctc	gatttcggcc	12360
ggcgacgtgg	agctggccag	cctcgcaaat	cggcgaaaa	gcctgatttt	acgcgagttt	12420
cccacagatg	atgtggacaa	gcctggggat	aagtgcctcg	cggtattgac	acttgagggg	12480
cgcgactact	gcagatgag	gggcgcgcat	cttgacactt	gaggggcaga	gtgctgacag	12540
atgagggggc	cacctattga	catttgaggg	gctgtccaca	ggcagaaaa	ccagcatttg	12600
caagggtttc	gcgccgtttt	tcggccaccg	ctaacctgtc	ttttaacctg	cttttaaac	12660
aatatttata	aaccttgggt	ttaacagggg	ctgcgccctg	tgccgctgac	cgcgccgcgc	12720
gaaggggggt	gccccccctt	ctcgaaacct	cccggcccg	taacgcgggc	ctcccatccc	12780
cccaggggct	gcgccccctg	gcgcggaacg	gcctcacccc	aaaaatggca	gcgctggcag	12840
tccttgccat	tgccgggatc	ggggcgagta	cggggatggc	gatcagcccg	agcgcgacgc	12900
ccggaagcat	tgacgtgcgc	cagggtgctg	catcgacatt	cagcgaccag	gtgccgggca	12960
gtgagggcgg	cgccctgggt	ggcggcctgc	ccttcaactt	ggcgcgcggg	gcattcacgg	13020
acttcatggc	ggggcccgga	atttttacct	tgggcattct	tgccatagtg	gtgcgggggt	13080
cgctgctcgt	ggttcggggg	gcgataaaac	cagcgaacca	tttgagggtg	taggttaaat	13140
tataccgagg	tatgaaaacg	agaattggac	ctttacagaa	ttactctatg	aagcgccata	13200
tttaaaaagc	taccaagacg	aagaggatga	agaggatgag	gaggcagatt	gccttgaata	13260
tattgacaat	actgataaga	taatatatct	tttatataga	agatatcgcc	gtatgtaagg	13320
atttcagggg	gcaaggcaca	ggcagcgccg	ttatcaatat	atctatagaa	tgggcgaagg	13380
ataaaaactt	gcatggacta	atgcttgaaa	cccaggacaa	taaccttata	gcttgttaaa	13440
tcctatcata	ttggggaatg	actccaactt	attgatagtg	ttttatgttc	agataatgcc	13500
cgatgaactt	gtcatgcagc	tccaccgatt	ttgagaacga	cagcgacttc	cgcccgacc	13560
gtgccagggt	gtgcctcaga	ttcaggttat	gcgcctcaat	tcgctgcgta	tatcgcttgc	13620
tgattacgtg	cagctttccc	ttcaggcggg	attcatacag	cgcccgacca	tcctgcatcc	13680
atatcaccac	gtcaaaaggt	gacagcaggc	tcataagacg	cccagcgctc	gccatagtgc	13740
gttcaccgaa	tacgtgcgca	acaacgcgtc	tcgggagact	gtcatacgcg	taaaacagcc	13800
agcgtggcgc	cgattttagc	ccgacatagc	cccactgttc	gtccatttcc	gcgcgagcga	13860
tgacgtcact	cccgcgctgt	ttaccgactg	cgccctgagt	tttttaagtg		13920
acgtaaaaac	gtgttgaggg	caacgcacct	aatgcgggct	gttgcccgcc	atccaacgcc	13980
attcatggcc	atatcaatga	ttttctgggt	cgtaccgggt	tgagaagcgg	tgtaagtga	14040
ctgcagttgc	catgtttttac	ggcagtgaag	gcagagatag	cgctgatgtc	cgccggtgct	14100
tttgccgtta	cgccaccacc	cgctcagtgc	tgaacaggag	ggacagctga	tagacacaga	14160
agccactgga	gcacctcaaa	aacaccatca	tacactaaat	cagtaagttg	gcagcatcac	14220
ccataaattg	ggtttcaaaa	tcggctccgt	cgatactatg	ttatacgcca	actttgaaaa	14280
caactttgaa	aaagctgttt	tctgggtatt	aaggttttag	aatgcaagga	acagtgaatt	14340
ggagttcgct	ttgttataat	tagcttcttg	gggtatcttt	aaatactgta	gaaaagagga	14400
aggaataaat	aaatggctaa	tacccggaat	tgaaaaaact	gatcgaaaaa		14460
taccgctgcg	taaaagatag	ggaagggaat	tctcctgcta	aggtatataa	gctggtggga	14520
gaaaatgaaa	acctatattt	aaaaatgacg	gacagccggt	ataaagggac	cacctatgat	14580
gtggaacggg	aaaaggacat	gatgctatgg	ctggaaggaa	agctgcctgt	tccaaaggtc	14640
ctgcactttg	aacggcctga	tggtctggagc	aatctgctca	tgagtggggc	cgatggcgct	14700
ctttgctcgg	aagagtatga	agatgaacaa	agccctgaaa	agatttatcg	gctgatgcgc	14760
gagtgcatca	ggctctttca	ctccatcgac	atatcggatt	gtccctatag	gaatagctta	14820
gacagccgct	tagccgaatt	ggattactta	ctgaataacg	atctggccga	tgtggattgc	14880
gaaaactggg	aagaagacac	tccatttaaa	gatccgcgcg	agctgtatga	ttttttaaag	14940
acggaaaagc	ccgaagagga	acttgtcttt	tcccacggcg	acctgggaga	cagcaacatc	15000
tttgtgaaag	atggcgaagt	aagtggcctt	attgatcttg	ggagaagcgg	cagggcgagc	15060
aagtggatg	acattgcctt	ctgcgtcccg	tcgatcaggg	aggtatcggg	ggaagaacag	15120
tatgtcgagc	tattttttga	cttactgggg	atcaagcctg	attggggaga	aataaaaat	15180
tatatattac	tggatgaatt	gttttagtac	ctagatgtgg	cgcaacgatg	ccggcgacaa	15240
gcaggagcgc	accgacttct	tcgcgatcaa	gtgttttggc	tctcaggcgc	aggccacagg	15300
caagtatttg	ggcaaggggt	cgctggtatt	cgtgcaggcg	aagattcgga	ataccaagta	15360

-continued

cgagaaggac	ggccagacgg	tctacgggac	cgacttcatt	gccgataagg	tggattatct	15420
ggacaccaag	gcaccaggcg	ggtaaaatca	ggaataaagg	cacattgccc	cggcgtgagt	15480
cggggcaatc	ccgcaaggag	ggatgaatgaa	tccggacgttt	gaccggaagg	catcacaggca	15540
agaactgatc	gacgcgggggt	tttccgcgca	ggatgccgaa	accatcgcaa	gccgcaccgt	15600
catgcgtcgc	cccgcgcaaa	ccttcagtc	cgtcggctcg	atgggtccagc	aagctacggc	15660
caagatcgag	cgcgacagcg	tgaactggc	tccccctgcc	ctgccccgcg	catcgccgcg	15720
cgtggagcgt	tccgctcgct	tccaacagga	ggcggcaggt	ttggcggaagt	cgatgaccat	15780
cgacacgcga	ggaactatga	cgaccaagaa	gcgaaaaacc	gccggcgagg	acctggcaaa	15840
acaggtcagc	gaggccaagc	aggccgcgtt	gctgaaacac	acgaagcagc	agatcaagga	15900
aatgcagctt	tccctgttgc	atattgcgcc	gtggccggac	acgatgcgag	cgatgccaaa	15960
cgacacggcc	cgctctgcgc	tgttaccac	gcgcaacaag	aaaatccgcg	gcgaggcgct	16020
gcaaaacaag	gtcattttcc	acgtcaacaa	ggacgtgaag	atcacctaca	ccggcgctcga	16080
gctgcccggc	gacgatgacg	aactggtgtg	gcagcaggtg	ttggagtacg	cgaagcgcac	16140
ccctatcgcc	gagcgcgatc	ccttcacgtt	ctacgagctt	tgcaggacc	tgggctgggtc	16200
gatcaatggc	cggtattaca	cgaaggccga	ggaatgcctg	tccgcgctac	aggcgacggc	16260
gatgggcttc	acgtccgacc	gcgttgggca	cctggaatcg	gtgtcgctgc	tgcaccgctt	16320
ccgcgtcctg	gacggtggca	agaaaacgtc	ccgttgccag	gtcctgatcg	acgaggaaat	16380
cgtcgtgctg	tttgcgtggc	accactacac	gaaattcata	tgggagaagt	accgcaagct	16440
gtcgcgcagc	gcccgcagga	tgttcgacta	tttcagctcg	caccgggagc	cgtaccgctt	16500
caagctggaa	accttcgcgc	tcatgtcggc	atcggattcc	accgcgctga	agaagtggcg	16560
cgagcaggtc	ggcgaaagcct	gcgaagagtt	gcgaggcagc	ggcctggtgg	aacacgcctg	16620
ggtaaatgat	gacctggtgc	attgcaaacg	ctagggcctt	gtggggtcag	ttccggctgg	16680
gggttcagca	gcccagcgtt	tactggcatt	tcaggaaaca	gcgggcactg	ctcgacgcac	16740
ttgcttcgct	cagtatcgct	cgggacgcac	ggcgcgctct	acgaactgcc	gataaacaga	16800
ggattaaaat	tgacaattgt	gattaaaggct	cagattcgac	ggcctggagc	ggccgacgtg	16860
caggatttcc	cgagatccg	attgtcggcc	ctgaagaaag	ctccagagat	gttcgggttc	16920
gtttacgagc	acgaggagaa	aaagcccatg	gaggcggttc	ctgaacgggt	gcgagatgcc	16980
gtggcattcg	gcgcctacat	cgacggcgag	atcattgggc	tgtcggctct	caaacaggag	17040
gacggcccga	aggacgcctc	caaggcgcat	ctgtccggcg	ttttcgtgga	gcccgaacag	17100
cgaggccgag	gggtccgcgg	tatgctgctg	cgggcggtgc	cggcggtgtt	attgctcgtg	17160
atgatcgtcc	gacagattcc	aacgggaaatc	tgggtggatgc	gcattctcat	cctccggcga	17220
cttaatat	gcctattctg	gagcttgttg	tttatttcgg	tctaccgctt	gcccggcggg	17280
gtcgcggcga	cggtaggcgc	tgtgcagccg	ctgatggctg	tgttcatctc	tgcgcgtctg	17340
ctaggtagcc	gcatacgatt	gatggcggtc	ctgggggcta	tttgccgaac	tgcgggcgtg	17400
gcgctgttgc	ttgtgacacc	aaacgcagcg	ctagatcctg	tcggcgctgc	agcgggcctg	17460
gcggggcgcg	tttccatggc	gttcggaacc	gtgctgaccc	gcaagtggca	acctcccgtg	17520
cctctgctca	cctttaccgc	ctggcaactg	gcggccggag	gacttctgct	cgttcagta	17580
gctttagtgt	ttgatccgc	aatcccgatg	cctacaggaa	ccaatgttct	cggcctggcg	17640
tggctcggcc	tgatcggagc	gggtttaacc	tacttccctt	gggtccgggg	gatctcgcga	17700
ctcgaaacct	cagttgtttc	cttactgggc	tttctcagcc	ccagatctgg	ggtcgatcag	17760
ccggggatgc	atcaggccga	cagtcggaa	ttcgggtccc	cgacctgtac	cattcgggtg	17820
gcaatggata	ggggagttga	tatcgtcaac	gttcacttct	aaagaaatag	cgccactcag	17880
cttctcagc	ggctttatcc	agcgatttcc	tattatgtcg	gcatagttct	caagatcgac	17940
agcctgtcac	gggttaagcga	gaaatgaata	agaaggctga	taattcggat	ctctgcgagg	18000
gagatgatat	ttgatccagc	gcagcaacgc	tctgtcatcg	ttacaatcaa	catgctaccc	18060
tcgcgcgagat	catcgtgtgt	tcaaaccccg	cagcttagtt	gccgttcttc	cgaatagcat	18120
cggttaacatg	agcaaaagtct	gccgccttac	aacggctctc	ccgctgacgc	cgtcccggac	18180
tgatgggctg	cctgtatcga	gtgggtgatt	tgtccgcagc	tgccggtcgg	ggagctgttg	18240
gctggctggg	ggcaggtatg	attgtgggtg	aaacaaattg	acgcttagac	aacttaataa	18300
cacattgcgg	acgtttttta	tgtactgggg	tggtttttct	tttcaccagt	gagacgggca	18360
acagctgatt	gcccttcaac	gcctggccct	gagagagttg	cagcaagcgg	tccacgctgg	18420
tttgccccag	caggcgaaaa	tctcttttga	tgggtggttc	gaaatcggca	aaatccotta	18480
taaatcaaaa	gaatagcccg	agataggggt	gagtggttgt	ccagtttggg	acaagagtc	18540
actattaaag	aacgtggagc	ccaacgtcaa	agggcgaaaa	accgtctatc	agggcgatgg	18600
ccactacgt	gaacctcac	ccaaatcaag	tttttggggg	tcgaggtgcc	gtaaaagcact	18660
aaatcggaac	cctaagggga	gcccccgatt	tagagcttga	cgggggaaagc	cgccgaacgt	18720
ggcgagaaag	gaaggggaag	aagcgaaagg	agcgggcgcc	attcaggctg	cgcaactggt	18780
gggaaggcg	atcgggtcgg	gcctcttcgc	tattacgcca	gctggcgaaa	gggggatgtg	18840
ctgcaaggcg	attaagttgg	gtaacgccag	ggttttccca	gtcacgacgt	tgtaaaaacga	18900
cggcagtgga	attcgagctc	ggtaaccccc	tactccaaaa	atgtcaaaag	tacagtctca	18960
gaagaccaaa	gggctattga	gacttttcaa	caaagggtaa	tttcgggaaa	cctcctcgga	19020
ttccattgcc	cagctatctg	tcacttcac	gaaaggacag	tagaaaagga	aggtggctcc	19080
tacaaatgcc	atcattgcga	taaaggaaag	gctatcattc	aagatgcctc	tgcgacaggt	19140
ggtcccaaa	atggaccccc	accacagagg	agcatcgtgg	aaaaagaaga	cgtcccaacc	19200
acgtcttcaa	agcaagtggg	ttgatgtgac	atctccactg	acgtaaaggga	tgacgcacaa	19260
tcccactatc	cttcgcaaga	cccttcttct	atataaggaa	gttcatttca	tttgagaggg	19320
acagcccaag	ctttctagag	gatccataaa	acatttcaat	cctttgaacg	cggtagaacg	19380
tgctaattgg	attttgtgga	gaacgcggta	gaacgtactt	atcacctaca	gttttatttt	19440
gtttttcttt	ttggtttaat	ctatccagct	tagtaccgag	tggggggaaag	tgactgggtg	19500
gcctaaaacc	ttttctttga	tactttgtga	aaatacatc	agatacaatg	gcgaacggta	19560
acttcaagtt	gttccaagt	ctcaatgtgg	acgagatgtc	tgtgagcag	aggagtcaat	19620
tctttgactt	gatgctgact	aaacctgatt	gtgagatcgg	gcaaatgatg	caaagagttg	19680
ttgttgataa	agtcgatgac	atgattagag	aaagaaagac	taagatccca	gtgattgttc	19740
atgaagttct	ttctcagaag	gaacagaaca	agttgatgga	aatttatcct	gaattcaata	19800
tcgtgtttaa	agacgacaaa	aacatgggtc	atgggtttgc	ggctgctgag	cgaaaactac	19860
aagctttatt	gcttttagat	agagttcctg	ctctgcaaga	gggtgatgac	atcgggtggtc	19920

-continued

```

aatggctgctt ttgggtaact agaggtgaga aaaggattca ttctgtgtgt ccaaatctag 19980
atattcggga tgatcagaga gaaatttctc gacagatatt tcttactgct attgggtgatc 20040
aagctagaag tggtaagaga cagatgtcgg agaattgagct gtggatgtat gaccaatttc 20100
gtgaaaatat tgctgcgcct aacgcgggta ggtgcaataa tacatatcag ggtgtacat 20160
gtaggggttt t 20171

```

```

SEQ ID NO: 16      moltype = DNA length = 300
FEATURE           Location/Qualifiers
source            1..300
                  mol_type = other DNA
                  organism = synthetic construct

```

```

SEQUENCE: 16
atgatgtcca gcaattacac cgcaattgat aatcagaatg ttccggatc tgttctctgca 60
gtcgcagatc cacttgccca agttgcgctc aaattcactg attcgaattt gcagacattt 120
ccaccttcca gttcccgagg gaaaatctct ggtgcttctg gacctccacg tgacgccgac 180
gatacattct ctaaacagct atctggttct gatgagcccc agcagagcgg ttggtttcgc 240
gctttcacta ttgctgcgta caagccctat tttgatgttg atacatcaga tgttctggag 300

```

1. A method of modulating (e.g. decreasing) the alkaloid content of a tobacco plant or a part thereof or tobacco plant cell, the method comprising modifying said plant or plant cell by modulating (e.g. decreasing) the activity or expression of a YIPF-related protein.

2. A method of modulating (e.g. decreasing) the content of a tobacco specific nitrosamine (TSNA) or a precursor of a TSNA in a tobacco plant or plant part thereof or tobacco plant cell, the method comprising modifying said plant or plant cell by modulating (e.g. decreasing) the activity or expression of a YIPF-related protein.

3. A method for producing a plant or part thereof, a cell or cell culture, a plant propagation material, a leaf, a cut harvested leaf, a processed leaf or a cut and processed leaf which has modulated (e.g. decreased) alkaloid content, the method comprising modifying said plant or cell culture to modulate the activity or expression of a YIPF-related protein.

4. Use of at least one gene encoding a YIPF-related protein for modulating alkaloid content of a tobacco cell or tobacco plant or part thereof.

5. A method or use according to any preceding claim, wherein the YIPF-related protein:

- a) comprises an amino acid sequence as set out in SEQ ID No. 3; or a functional variant or functional fragment or orthologue of SEQ ID No. 3; or a sequence which has at least 80% identity to SEQ ID No. 3; or a homologue of SEQ ID No. 3; or
- b) is encoded by a nucleotide sequence as set out in SEQ ID No. 1 or 2; or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2; or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2.

6. A method or use according to any one of the preceding claims, wherein the alkaloid content is modulated (e.g. decreased) in comparison to a plant or cell culture which has not been modified to modulate the activity or expression of a YIPF-related protein.

7. A tobacco plant or part thereof or a tobacco cell or cell culture which has been modified to modulate (e.g. decrease) the activity or expression of a YIPF-related protein and the tobacco plant or part thereof or tobacco cell or cell culture has decreased alkaloid and or TSNA precursor content in comparison to an unmodified plant or unmodified cell or cell culture.

8. A plant propagation material obtainable (e.g. obtained) from a plant according to claim 7 or from a plant or cell or cell culture produced by the method or use of any one of claims 1 to 6.

9. A method or use according to any one of claims 1-6, or a plant or part thereof or cell or cell culture according to claim 7, or a plant propagation material according to claim 8, wherein the alkaloid content of the plant is decreased in comparison to a plant or cell culture which has not been modified to modulate the activity or expression of a YIPF-related protein.

10. A method or use according to any one of claim 1-6 or 9, a tobacco plant or part thereof or tobacco cell or cell culture according to claim 7 or 9, or a plant propagation material according to claim 8, wherein the content of one or more alkaloids selected from nicotine, nornicotine, PON, anabesine, myosmine and anatabine is modulated (e.g. decreased), preferably the content of nicotine, nornicotine and/or PON is modulated (e.g. decreased).

11. A method or use according to claim 10, a tobacco plant or part thereof or tobacco cell or tobacco cell culture according to claim 10, or a plant propagation material according to claim 10 wherein the nicotine content is decreased.

12. Use of a tobacco plant or part thereof or tobacco cell or cell culture according to any one of claim 7 or 9-11, or of a plant produced by the method of any one of claims 1 to 6, or 9 to 11 to breed a plant.

13. Use of a tobacco plant or part thereof or a tobacco cell or cell culture according to any one of claim 7 or 9-11, or of a plant produced by the method of any one of claims 1 to 6, or 9 to 11 for production of a product.

14. Use of a tobacco plant or part thereof according to any one of claim 7 or 9-11, or of a plant produced by the method of any one of claims 1 to 6, or 9 to 11 to grow a crop.

15. Use of a tobacco plant or part thereof according to any one of claim 7 or 9-11, or of a plant produced by the method of any one of claims 1 to 6, or 9 to 11 to produce a leaf.

16. A harvested leaf of a plant according to any one of claim 7 or 9-11, or obtainable from a plant propagated from a propagation material according to any one of claims 8-11, or obtainable from a plant obtained by a use according to any

one of claim 4-6 or 9-11, or obtainable from a plant produced by the method of any one of claim 1-6, or 9-11.

17. A harvested leaf of a plant according to claim 16, wherein the harvested leaf of a plant is a cut harvested leaf.

18. A processed leaf, preferably a processed tobacco leaf, preferably a non-viable processed tobacco leaf:

obtainable (e.g. obtained) from a plant obtainable from a use according to any one of claim 4-6 or 9-11;

obtainable (e.g. obtained) by processing a plant according to any one of claim 7 or 9-11;

obtainable (e.g. obtained) from a plant propagated from a plant propagation material according to any one of claims 8-11; or

obtainable (e.g. obtained) by processing a harvested leaf of a plant according to claim 16 or 17; or

obtainable (e.g. obtained) from a plant produced by the method of any one of claims 1 to 6, or 9 to 11.

19. A processed leaf according to claim 18, wherein the leaf is processed by curing, fermenting, pasteurising or a combination thereof.

20. A processed leaf according to claim 18 or 19, wherein the processed leaf is a cut processed leaf.

21. Cured tobacco material made from a plant or a part thereof according to any one of claim 7 or 9-11, or a harvested leaf according to claim 16 or 17, or a processed leaf according to any of claims 18-20.

22. A tobacco blend comprising said cured tobacco material of claim 21.

23. A tobacco industry product prepared from:

a tobacco plant or part thereof or tobacco cell or cell culture according to any one of claim 7 or 9-11;

a tobacco plant or part thereof propagated from a tobacco plant propagation material according to claim 8;

a harvested leaf of a plant according to claim 16 or 17;

a processed leaf according to any one of claims 18-20.

24. A tobacco industry product according to claim 23, wherein the tobacco product is a combustible smoking article.

25. A tobacco industry product according to claim 23, wherein the tobacco product is a smokeless tobacco product.

26. A tobacco product according to claim 23, wherein the tobacco product is a non-combustible aerosol provision system such as a tobacco heating device or an aerosol-generating device.

27. A combustible smoking article, non-combustible aerosol provisioning system, smokeless tobacco product or tobacco heating device comprising a plant or a part thereof according to any one of claim 7 or 9-11 or an extract (e.g. a tobacco extract) thereof or a tobacco cell culture according to any one of claim 7 or 9-11; or a cured tobacco material according to claim 21; or a tobacco blend according to claim 22.

28. Use of a nucleotide sequence encoding a YIPF-related protein which:

a) encodes an amino acid sequence as set out in SEQ ID No. 3; or a functional variant or functional fragment or orthologue of SEQ ID No. 3; or a sequence which has at least 80% identity to SEQ ID No. 3; or a homologue of SEQ ID No. 3; or

b) comprises a sequence as set out in SEQ ID No. 1 or 2; or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2; or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2;

to select a plant having modulated (e.g. reduced) alkaloid content and/or modulated (e.g. reduced) content of TSNA or a precursor of a TSNA.

29. A mutant of a plant carrying a heritable mutation in a nucleotide sequence which:

a) encodes an amino acid sequence as set out in SEQ ID No. 3; or a functional variant or functional fragment or orthologue of SEQ ID No. 3; or a sequence which has at least 80% identity to SEQ ID No. 3; or a homologue of SEQ ID No. 3; or

b) comprises a sequence as set out in SEQ ID No. 1 or 2; or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2; or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2;

wherein said heritable mutation modulates (e.g. decreases) the activity or expression of YIPF-related protein and wherein the mutant plant has modulated (e.g. decreased) alkaloid content and/or modulated content of a tobacco specific nitrosamine (TSNA) or a precursor of a TSNA relative to a comparable plant which does not carry said heritable mutation.

30. Progeny or seed of a mutant plant which carries the heritable mutation according to claim 29.

31. A harvested leaf, a processed leaf or cured tobacco material produced from a plant comprising a modification in a nucleotide sequence which:

a) encodes an amino acid sequence as set out in SEQ ID No. 3; or a functional variant or functional fragment or orthologue of SEQ ID No. 3; or a sequence which has at least 80% identity to SEQ ID No. 3; or a homologue of SEQ ID No. 3; or

b) comprises a sequence as set out in SEQ ID No. 1 or 2; or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2; or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2;

wherein said modification modulates (e.g. decreases) the activity or expression of YIPF-related protein and wherein said plant has modulated (e.g. decreased) alkaloid content and/or modulated content of a tobacco specific nitrosamine (TSNA) or a precursor of a TSNA relative to a comparable plant which does not carry said modification in said YIPF-related protein.

* * * * *