

Figure 1




Figure 2




Figure 3

Figure 3A (SEQ ID NOs:18-25)




Figure 3B




Figure 3C




Figure 4

Figure 4A




Figure 4B (SEQ ID NOs:26 and 19-21)

ATGTTATCGAGCGACC

ATAGCTCGCTGGTTAG

AATCGCATCGGCATAT

CGTAGCCGTATATATC

O_1

O_1

0

O_2

0

O_3

0

O_4

0

O_2

0

O_3

1

O_4

0

O_3

0

O_4

1

O_4

0

O_4

0

O_4

0

O_4

0

$w = 1$

Figure 4 (continued)

Figure 4C (SEQ ID NOs: 18-21)

ATAGTATCGAGCGACC

ATAGCTCGCTGGTTTAG

AATCGCATCGGCATAT

CGTAGCCGTATATATC

$M =$

	O_1	O_2	O_3	O_4
O_1	0	0	0	1
O_2	0	0	1	0
O_3	0	1	0	0
O_4	1	0	0	0

$w = 0$

Figure 5

Figure 5A




Figure 5 (continued)

Figure 5B




Figure 6




Figure 7 (continued)

Figure 7A

[illegible][illegible][illegible][illegible]

Figure 8

```
>Seq_01 (SEQ ID NO:1)
AGCCATAAGACGTACTGCACCTGTTGGGTTACAATCATATTCTTTCATGAGAGTCACGCATCCC
AATCGCCCCGTAACCTCCATCCTTTGACCCGTCAAATTGCTTCCCCGTGTTAACAGCAACCTCC
AAAA
```

```
>Oligo a+ (SEQ ID NO:2) AGCCATAAGACGTACT
>Oligo a- (SEQ ID NO:3) GTGCAGTACGTCTTAT
>Oligo b+ (SEQ ID NO:4) GCACCTGTTGGGTTAC
>Oligo b- (SEQ ID NO:5) GATTGTAACCCAACAG
>Oligo c+ (SEQ ID NO:6) AATCATATTCTTTCAT
>Oligo c- (SEQ ID NO:7) TCTCATGAAAGAATAT
>Oligo d+ (SEQ ID NO:8) GAGAGTCACGCATCCC
>Oligo d- (SEQ ID NO:9) GATTGGGATGCGTGAC
>Oligo e+ (SEQ ID NO:10) AATCGCCCCGTAACCT
>Oligo e- (SEQ ID NO:11) ATGGAAGTTACGGGGC
>Oligo f+ (SEQ ID NO:12) CCATCCTTTGACCCGT
>Oligo f- (SEQ ID NO:13) TTTGACGGGTCAAAGG
>Oligo g+ (SEQ ID NO:14) CAAATTGCTTCCCCGT
>Oligo g- (SEQ ID NO:15) TAACACGGGGAAGCAA
>Oligo h+ (SEQ ID NO:16) GTTAACAGCAACCTCC
>Oligo h- (SEQ ID NO:17) TTTTGGAGGTTGCTGT
>Oligo a+ (SEQ ID NO:18) ATAGTATCGAGCGACC
>Oligo a- (SEQ ID NO:19) GATTGGTCGCTCGATA
>Oligo b+ (SEQ ID NO:20) AATCGCATCGGCATAT
>Oligo b- (SEQ ID NO:21) CTATATATGCCGATGC
>Oligo c+ (SEQ ID NO:22) ATAGTATGGCCATATG
>Oligo c- (SEQ ID NO:23) TGAGCATATGGCCATA
>Oligo d+ (SEQ ID NO:24) CTCATTGAGAGTACTG
>Oligo d- (SEQ ID NO:25) GTCACAGTACTCTCAA
>Oligo a+ (SEQ ID NO:26) ATGTTATCGAGCGACC
```

```
>Seq_02 (SEQ ID NO:27)
GCACCAAACGAAAATACTATGGGCCAACGATACCGAGTGTACGGACACAGACGATTACGTCTGA
ATCTCATGCCCAGTAGTCGGAGGCGCGGCCAGAGATATTGAGGCGATCAGCCACCCGAATGGAG
AACAAAGCCACAACCCGTATTGCGTCTCTATTTGTTTTGATTGCCGCATCATCTAGTGATGAGG
GCACCTGAACACCTTAATACCTAATTCGTCACCACATTGTTGTGAGCCATAGGGGGACATCCTG
TTTA
```

Figure 8 (continued)

```

>Oligo a+ (SEQ ID NO:28) GCACCAAACGAAAATA
>Oligo a- (SEQ ID NO:29) ATAGTATTTTCGTTTG
>Oligo b+ (SEQ ID NO:30) CTATGGGCCAACGATA
>Oligo b- (SEQ ID NO:31) TCGGTATCGTTGGCCC
>Oligo c+ (SEQ ID NO:32) CCGAGTGTACGGACAC
>Oligo c- (SEQ ID NO:33) GTCTGTGTCCGTACAC
>Oligo d+ (SEQ ID NO:34) AGACGATTACGTCTGA
>Oligo d- (SEQ ID NO:35) AGATTCAGACGTAATC
>Oligo e+ (SEQ ID NO:36) ATCTCATGCCCAGTAG
>Oligo e- (SEQ ID NO:37) CCGACTACTGGGCATG
>Oligo f+ (SEQ ID NO:38) TCGGAGGCGCGGCCAG
>Oligo f- (SEQ ID NO:39) ATCTCTGGCCGCGCCT
>Oligo g+ (SEQ ID NO:40) AGATATTGAGGCGATC
>Oligo g- (SEQ ID NO:41) GGCTGATCGCCTCAAT
>Oligo h+ (SEQ ID NO:42) AGCCACCCGAATGGAG
>Oligo h- (SEQ ID NO:43) TGTTCTCCATTTCGGGT
>Oligo i+ (SEQ ID NO:44) AACAAAGCCACAACCC
>Oligo i- (SEQ ID NO:45) ATACGGGTTGTGGCTT
>Oligo j+ (SEQ ID NO:46) GTATTGCGTCTCTATT
>Oligo j- (SEQ ID NO:47) AACAAATAGAGACGCA
>Oligo k+ (SEQ ID NO:48) TGTTTTGATTGCCGCA
>Oligo k- (SEQ ID NO:49) ATGATGCGGCAATCAA
>Oligo l+ (SEQ ID NO:50) TCATCTAGTGATGAGG
>Oligo l- (SEQ ID NO:51) GTGCCCTCATCACTAG
>Oligo m+ (SEQ ID NO:52) GCACCTGAACACCTTA
>Oligo m- (SEQ ID NO:53) GTATTAAGGTGTTTCAG
>Oligo n+ (SEQ ID NO:54) ATACCTAATTCGTCAC
>Oligo n- (SEQ ID NO:55) TGTGGTGACGAATTAG
>Oligo o+ (SEQ ID NO:56) CACATTGTTGTGAGCC
>Oligo o- (SEQ ID NO:57) CTATGGCTCACAACAA
>Oligo p+ (SEQ ID NO:58) ATAGGGGGACATCCTG
>Oligo p- (SEQ ID NO:59) TAAACAGGATGTCCCC

```

Figure 8 (continued)

>Seq_03 (SEQ ID NO:60)

CACTTAGTGCTCGTCCCGTTGATCGCTGGTGCATCAGGGCCTTGAGCTGGCCTGTTGGTGGGCC
TGCTAACGCCGCCCTCCTGTGGACGAAGTGGAATGAGAGACGTGCGCCATGTTATCTCCGCCAG
CCGTCCATGCGTGACCGGGTCGGGTACGACCGGGCCGGAACGGACTCACACCCATGGTGAGCAA
TACGTATCCACCGGGAGCTTAGTTTGGCCCCCTTGGGCCGACCCGGGGAGTCCGAACGCTCGGG
ACTACTACTTCACGAGCAAGCATCTCCGGCGCCGCTGCCCCGCGATCTGGCACGCCCCCTTCCAGA
CTCCTGGGCAGCGACAGCGCCGAGAGGGCGGTGCCCCAGGGGAACCGCCCTACAGCCCCCGAGA
TCGTTTCTCCCCCGCTTCCACCCCTGGGCGAGGTGCATACCTTTCAACCCGTCAAATGCCACT
TAGAACGTCCTGCTCGTCCGTAGAAAACAGGCAGACGCCGACGCCCATATGGGAGGCGCTCGT
TGGCCCCGCTGGGCGACTGGGCGTGACTGGCGACCGCCAGGTATCCCCGGGGGGAGCGCGCCACG
CTGCAGGCGGACGCGAGGGAGATGCAGGGACGGGGTGGTCAGCTTGACCCGGGCCTGGGAGGCT
CGTCCGGCCCCCTCGCGGCTAGACTGGGATAGCGGTATACCAGCGGGAGGAGTCTCTCCGGCGTC
CGGCCCTGCAGGCAAAGGCGTGAGCGTTTCCCGCTGTCCGGGCGGTAGCGGCTGGGGCCATGTC
CGACGGCGCGCGATGTAGTCGGGGCCCATTTCCACGGTGCAACGTCCCCCTACAGTAGGTCAATC
GGCAGGCACAGGCGCGCGCGCCCGCCGCGCTAGTTCCGTGGCCGTCGTACAGAGCAGAGGAATC
CGAAGTCGGGTTGCCCGGGCTCCTGTCCGGCCACCGCCCCGAGCTCCCTACGGGACAGCCCGGG
AAAGCTGGGAGGGACAGGGTAGTCGGCCCCCCCCGCTCTGTTCGTTCCGCATCCGCCCTTGCGGGCC
CCGC

Oligo (SEQ ID NO:61) TCGAGGAACGC
Oligo (SEQ ID NO:62) ATATGCGTTCCTCGA
Oligo (SEQ ID NO:63) ATATCGGTAAA
Oligo (SEQ ID NO:64) TTTAGCG
Oligo (SEQ ID NO:65) TCGAGGAACG
Oligo (SEQ ID NO:66) TATGCGTTCCTCGA
Oligo (SEQ ID NO:67) CATATCGGTAAA
Oligo (SEQ ID NO:68) TTTAGCGA
Oligo (SEQ ID NO:69) TCGAGGAACGCA
Oligo (SEQ ID NO:70) TATCGGTAAA
Oligo (SEQ ID NO:71) TCGAGGAACG
Oligo (SEQ ID NO:72) CATATCGGTAAA

Figure 10 (continued)

Figure 10B




Figure 11




Figure 12




Figure 13




Figure 14

Leading Strands:

CACTTAGTGCTCGTCC, SEQ ID NO:73
 CGTTGATCGCTGGTGC, SEQ ID NO:74
 ATCAGGGCCTTGAGCT, SEQ ID NO:75
 GGCCTGTTGGTGGGCC, SEQ ID NO:76
 TGCTAACGCCGCCCTC, SEQ ID NO:77
 CTGTGGACGAAGTGGA, SEQ ID NO:78
 ATGAGAGACGTGCGCC, SEQ ID NO:79
 ATGTTATCTCCGCCAG, SEQ ID NO:80
 CCGTCCATGCGTGACC, SEQ ID NO:81
 GGGTCGGGTACGACCG, SEQ ID NO:82
 GGCCGGAACGGACTCA, SEQ ID NO:83
 CACCCATGGTGAGCAA, SEQ ID NO:84
 TACGTATCCACCGGA, SEQ ID NO:85
 GCTTAGTTTGGCCCCC, SEQ ID NO:86
 TTGGGCCGACCCGGGG, SEQ ID NO:87
 AGTCCGAACGCTCGGG, SEQ ID NO:88
 ACTACTACTTCACGAG, SEQ ID NO:89
 CAAGCATCTCCGGCGC, SEQ ID NO:90
 CGCTGCCCCGCGATCTG, SEQ ID NO:91
 GCACGCCCTTCCAGA, SEQ ID NO:92
 CTCCTGGGCAGCGACA, SEQ ID NO:93
 GCGCCGAGAGGGCGGT, SEQ ID NO:94
 CGCCCAGGGGAACCGC, SEQ ID NO:95
 CCTACAGCCCCCGAGA, SEQ ID NO:96
 TCGTTTCTCCCCCGC, SEQ ID NO:97
 TTCCACCCCTGGGCGA, SEQ ID NO:98
 GGTGCATACCTTTCAA, SEQ ID NO:99
 CCCGTCAAATGCCACT, SEQ ID NO:100
 TAGAACGTCCTGCTCG, SEQ ID NO:101
 TCCGTAGAAAACAGGC, SEQ ID NO:102
 AGACGCCGGACGCCCA, SEQ ID NO:103
 TATGGGAGGCGCTCGT, SEQ ID NO:104
 TGGCCCGCTGGGCGAC, SEQ ID NO:105
 TGGGCGTGACTGGCGA, SEQ ID NO:106
 CCGCCAGGTATCCCCG, SEQ ID NO:107
 GGGGGAGCGCGCCACG, SEQ ID NO:108
 CTGCAGGCGGACGCGA, SEQ ID NO:109
 GGGAGATGCAGGGACG, SEQ ID NO:110
 GGGTGGTCAGCTTGAC, SEQ ID NO:111
 CCGGGCCTGGGAGGCT, SEQ ID NO:112
 CGTCCGGCCCCCTCGCG, SEQ ID NO:113
 GCTAGACTGGGATAGC, SEQ ID NO:114
 GGTATAACCAGCGGGAG, SEQ ID NO:115
 GAGTCTCTCCGGCGTC, SEQ ID NO:116
 CGGCCCTGCAGGCAAA, SEQ ID NO:117
 GGCGTGAGCGTTTCCC, SEQ ID NO:118

Figure 14 (continued)

GCTGTCCGGGCGGTAG, SEQ ID NO:119
 CGGCTGGGGCCATGTC, SEQ ID NO:120
 CGACGGCGCGCGATGT, SEQ ID NO:121
 AGTCGGGGCCCATTCC, SEQ ID NO:122
 ACGGTGCAACGTCCCC, SEQ ID NO:123
 CTACAGTAGGTCAATC, SEQ ID NO:124
 GGCAGGCACAGGCGCG, SEQ ID NO:125
 CGCGCCCGCCGCGCTA, SEQ ID NO:126
 GTTCCGTGGCCGTCGT, SEQ ID NO:127
 ACAGAGCAGAGGAATC, SEQ ID NO:128
 CGAAGTCGGGTTGCCC, SEQ ID NO:129
 GGGCTCCTGTCCGGCC, SEQ ID NO:130
 ACCGCCCCGAGCTCCC, SEQ ID NO:131
 TACGGGACAGCCCGGG, SEQ ID NO:132
 AAAGCTGGGAGGGACA, SEQ ID NO:133
 GGGTAGTCGGCCCCCC, SEQ ID NO:134
 CGCTCTGTCTGTTCCGC, SEQ ID NO:135
 ATCCGCCTTGCGGGCC, SEQ ID NO:136

Lagging Strands:

AACGGGACGAGCACTA, SEQ ID NO:137
 TGATGCACCAGCGATC, SEQ ID NO:138
 GGCCAGCTCAAGGCCC, SEQ ID NO:139
 AGCAGGCCCACCAACA, SEQ ID NO:140
 ACAGGAGGGCGGCGTT, SEQ ID NO:141
 TCATTCCACTTCGTCC, SEQ ID NO:142
 ACATGGCGCACGTCTC, SEQ ID NO:143
 ACGGCTGGCGGAGATA, SEQ ID NO:144
 ACCCGGTCACGCATGG, SEQ ID NO:145
 GGCCCGGTCTGTACCCG, SEQ ID NO:146
 GGTGTGAGTCCGTTCC, SEQ ID NO:147
 CGTATTGCTCACCATG, SEQ ID NO:148
 AAGCTCCCGGTGGATA, SEQ ID NO:149
 CCAAGGGGGCCAAACT, SEQ ID NO:150
 GACTCCCCGGGTCGGC, SEQ ID NO:151
 TAGTCCCGAGCGTTCG, SEQ ID NO:152
 CTTGCTCGTGAAGTAG, SEQ ID NO:153
 AGCGGCGCCGGAGATG, SEQ ID NO:154
 GTGCCAGATCGCGGGC, SEQ ID NO:155
 GGAGTCTGGAAGGGGC, SEQ ID NO:156
 GCGCTGTCTGCTGCCCA, SEQ ID NO:157
 GGCGACCGCCCTCTCG, SEQ ID NO:158
 TAGGGCGGTTCCCCTG, SEQ ID NO:159
 ACGATCTCGGGGGCTG, SEQ ID NO:160
 GGAAGCGGGGGGAGAA, SEQ ID NO:161
 CACCTCGCCCAGGGGT, SEQ ID NO:162
 CGGGTTGAAAGGTATG, SEQ ID NO:163

Figure 14 (continued)

TCTAAGTGGCATTGTA, SEQ ID NO:164
 CGGACGAGCAGGACGT, SEQ ID NO:165
 GTCTGCCTGTTTTCTA, SEQ ID NO:166
 CATATGGGCGTCCGGC, SEQ ID NO:167
 GCCAACGAGCGCCTCC, SEQ ID NO:168
 CCCAGTCGCCCAGCGG, SEQ ID NO:169
 GCGGTCGCCAGTCACG, SEQ ID NO:170
 CCCCCGGGGATACCTG, SEQ ID NO:171
 GCAGCGTGGCGCGCTC, SEQ ID NO:172
 TCCCTCGCGTCCGCCT, SEQ ID NO:173
 ACCCCGTCCCTGCATC, SEQ ID NO:174
 CCGGGTCAAGCTGACC, SEQ ID NO:175
 GACGAGCCTCCCAGGC, SEQ ID NO:176
 TAGCCGCGAGGGGCCG, SEQ ID NO:177
 TACCGCTATCCCAGTC, SEQ ID NO:178
 ACTCCTCCCGCTGGTA, SEQ ID NO:179
 GCCGGACGCCGGAGAG, SEQ ID NO:180
 CGCCTTTGCCTGCAGG, SEQ ID NO:181
 CAGCGGGAAACGCTCA, SEQ ID NO:182
 GCCGCTACCGCCCGGA, SEQ ID NO:183
 GTCGGACATGGCCCCA, SEQ ID NO:184
 GACTACATCGCGCGCC, SEQ ID NO:185
 CCGTGGAATGGGCCCC, SEQ ID NO:186
 GTAGGGGGACGTTGCA, SEQ ID NO:187
 TGCCGATTGACCTACT, SEQ ID NO:188
 CGCGCGCGCCTGTGCC, SEQ ID NO:189
 GAACTAGCGCGGCGGG, SEQ ID NO:190
 CTGTACGACGGCCACG, SEQ ID NO:191
 TTCGGATTCCCTCTGCT, SEQ ID NO:192
 GCCCGGGCAACCCGAC, SEQ ID NO:193
 CCGTGGCCGGACAGGA, SEQ ID NO:194
 CGTAGGGAGCTCGGGG, SEQ ID NO:195
 CTTTCCCGGGCTGTCC, SEQ ID NO:196
 ACCCTGTCCCTCCCAG, SEQ ID NO:197
 AGCGGGGGGGCCGACT, SEQ ID NO:198
 GGATGCGGAACGACAG, SEQ ID NO:199
 GCGGGGCCCGCAAGGC, SEQ ID NO:200

Figure 15

Leading Strands:

CACTTAGTGCTCGTC, SEQ ID NO:201
CCGTTGATCGCTGGT, SEQ ID NO:202
GCATCAGGGCCTTGAGC, SEQ ID NO:203
TGGCCTGTTGGTGGGC, SEQ ID NO:204
CTGCTAACGCCGCCCT, SEQ ID NO:205
CCTGTGGACGAAGTGGA, SEQ ID NO:206
ATGAGAGACGTGCGCCA, SEQ ID NO:207
TGTTATCTCCGCCAGCC, SEQ ID NO:208
GTCCATGCGTGACCGGG, SEQ ID NO:209
TCGGGTACGACCGGGCC, SEQ ID NO:210
GGAACGGACTCACAC, SEQ ID NO:211
CCATGGTGAGCAATACG, SEQ ID NO:212
TATCCACCGGGAGCT, SEQ ID NO:213
TAGTTTGGCCCCCTT, SEQ ID NO:214
GGGCCGACCCGGGGAG, SEQ ID NO:215
TCCGAACGCTCGGGACT, SEQ ID NO:216
ACTACTTCACGAGCAAG, SEQ ID NO:217
CATCTCCGGCGCCGCTG, SEQ ID NO:218
CCCGCATCTGGCACGC, SEQ ID NO:219
CCCTTCCAGACTCCT, SEQ ID NO:220
GGGCAGCGACAGCGC, SEQ ID NO:221
CGAGAGGGCGGTCGC, SEQ ID NO:222
CCAGGGGAACCGCCC, SEQ ID NO:223
TACAGCCCCCGAGATCG, SEQ ID NO:224
TTTCTCCCCCGCTTCC, SEQ ID NO:225
ACCCCTGGGCGAGGTGC, SEQ ID NO:226
ATACCTTTCAACCGTC, SEQ ID NO:227
AAATGCCACTTAGAACG, SEQ ID NO:228
TCCTGCTCGTCCGTAGA, SEQ ID NO:229
AAACAGGCAGACGCCGG, SEQ ID NO:230
ACGCCCATATGGGAGG, SEQ ID NO:231
CGCTCGTTGGCCCGC, SEQ ID NO:232
TGGGCGACTGGGCGTGA, SEQ ID NO:233
CTGGCGACCGCCAGGTA, SEQ ID NO:234
TCCCCGGGGGAGCGCG, SEQ ID NO:235
CCACGCTGCAGGCGGAC, SEQ ID NO:236
GCGAGGGAGATGCAGG, SEQ ID NO:237
GACGGGGTGGTCAGCTT, SEQ ID NO:238
GACCCGGGCCTGGGAGG, SEQ ID NO:239
CTCGTCCGGCCCCTCGC, SEQ ID NO:240
GGCTAGACTGGGATAGC, SEQ ID NO:241
GGTATACCAGCGGGAGG, SEQ ID NO:242
GCCCTGCAGGCAAAGG, SEQ ID NO:243
CGTGAGCGTTTCCCG, SEQ ID NO:244
CTGTCCGGGCGGTAGCG, SEQ ID NO:245
GCTGGGGCCATGTCCGA, SEQ ID NO:246

Figure 15 (continued)

CGGCGCGCGATGTAGTC, SEQ ID NO:247
 GGGGCCCATTCACGGT, SEQ ID NO:248
 GCAACGTCCCCCTAC, SEQ ID NO:249
 AGTAGGTCAATCGGCAG, SEQ ID NO:250
 GCACAGGCGCGCGCGCC, SEQ ID NO:251
 CGCCGCGCTAGTTCC, SEQ ID NO:252
 GTGGCCGTCGTACAG, SEQ ID NO:253
 AGCAGAGGAATCCGAA, SEQ ID NO:254
 GTCGGGTTGCCCCGGG, SEQ ID NO:255
 CTCCTGTCCGGCCACCG, SEQ ID NO:256
 CCCCAGAGCTCCCTAC, SEQ ID NO:257
 GGGACAGCCCCGGGAAA, SEQ ID NO:258
 GCTGGGAGGGACAGGG, SEQ ID NO:259
 TAGTCGGCCCCCCCCG, SEQ ID NO:260
 CTCTGTCGTTCCGCA, SEQ ID NO:261
 TCCGCCTTGCGGGCC, SEQ ID NO:262

Lagging Strands:


ACGGGACGAGCACTA, SEQ ID NO:263
 ATGCACCAGCGATCA, SEQ ID NO:264
 GCCAGCTCAAGGCCCTG, SEQ ID NO:265
 GCAGGCCCACCAACAG, SEQ ID NO:266
 CAGGAGGGCGGCGTTA, SEQ ID NO:267
 TCATTCCACTTCGTCCA, SEQ ID NO:268
 AACATGGCGCACGTCTC, SEQ ID NO:269
 GGACGGCTGGCGGAGAT, SEQ ID NO:270
 CCGACCCGGTCACGCAT, SEQ ID NO:271
 TTCCGGCCCGGTCGTAC, SEQ ID NO:272
 ATGGGTGTGAGTCCG, SEQ ID NO:273
 GATACGTATTGCTCACC, SEQ ID NO:274
 ACTAAGCTCCCGGTG, SEQ ID NO:275
 GCCCAAGGGGGCCAA, SEQ ID NO:276
 CGGACTCCCCGGGTCG, SEQ ID NO:277
 TAGTAGTCCCGAGCGTT, SEQ ID NO:278
 GATGCTTGCTCGTGAAG, SEQ ID NO:279
 CGGGCAGCGGCGCCGGA, SEQ ID NO:280
 AGGGGCGTGCCAGATCG, SEQ ID NO:281
 GCCCAGGAGTCTGGA, SEQ ID NO:282
 CTCGGCGCTGTCGCT, SEQ ID NO:283
 CTGGGCGACCGCCCT, SEQ ID NO:284
 TGTAGGGCGGTTCCC, SEQ ID NO:285
 GAAACGATCTCGGGGGC, SEQ ID NO:286
 GGGTGGAAGCGGGGGGA, SEQ ID NO:287
 GTATGCACCTCGCCCAG, SEQ ID NO:288
 ATTTGACGGGTTGAAAG, SEQ ID NO:289
 AGGACGTTCTAAGTGGC, SEQ ID NO:290
 GTTTTCTACGGACGAGC, SEQ ID NO:291

Figure 15 (continued)


GGGTCCGGCGTCTGCCT, SEQ ID NO:292
AGCGCCTCCCATATGG, SEQ ID NO:293
CCCAGCGGGCCAACG, SEQ ID NO:294
CCAGTCACGCCCAGTCG, SEQ ID NO:295
GGGATACCTGGCGGTCG, SEQ ID NO:296
GTGGCGCGCTCCCCCG, SEQ ID NO:297
TCGCGTCCGCCTGCAGC, SEQ ID NO:298
CGTCCCTGCATCTCCC, SEQ ID NO:299
GGTCAAGCTGACCACCC, SEQ ID NO:300
CGAGCCTCCCAGGCCCG, SEQ ID NO:301
AGCCGCGAGGGGCCGGA, SEQ ID NO:302
TACCGCTATCCCAGTCT, SEQ ID NO:303
GACTCCTCCCGCTGGTA, SEQ ID NO:304
GGGCCGGACGCCGAGA, SEQ ID NO:305
CACGCCTTTGCCTGCA, SEQ ID NO:306
ACAGCGGGAAACGCT, SEQ ID NO:307
CAGCCGCTACCGCCCGG, SEQ ID NO:308
GCCGTGCGACATGGCCC, SEQ ID NO:309
CCCCGACTACATCGCGC, SEQ ID NO:310
TTGCACCGTGGAATGGG, SEQ ID NO:311
TACTGTAGGGGGACG, SEQ ID NO:312
GTGCCTGCCGATTGACC, SEQ ID NO:313
GGCGGGCGCGCGCCT, SEQ ID NO:314
CCACGGAAGTAGCGC, SEQ ID NO:315
TGCTCTGTACGACGG, SEQ ID NO:316
CGACTTCGGATTCTC, SEQ ID NO:317
GGAGCCCCGGGCAACC, SEQ ID NO:318
GGGGCGGTGGCCGACA, SEQ ID NO:319
TCCCGTAGGGAGCTC, SEQ ID NO:320
CAGCTTTCCCGGGCTG, SEQ ID NO:321
ACTACCCTGTCCCTCC, SEQ ID NO:322
AGAGCGGGGGGGCCG, SEQ ID NO:323
CGGATGCGGAACGAC, SEQ ID NO:324
GCGGGGCCCGCAAGG, SEQ ID NO:325

Fig. 16

A)



B)



ASYMMETRIC ASSEMBLY OF POLYNUCLEOTIDES

FIELD OF THE INVENTION

[0001] The invention relates to a novel method for synthesizing a double stranded (ds) polynucleotide having a predefined sequence using a workflow of multiple assembly tiers including at least one asymmetric assembly of nucleic acid (NA) building blocks.

BACKGROUND OF THE INVENTION

[0002] Some methods for the synthesis and assembly of nucleic acids implement a hierarchical assembly approach where multiple pairs of DNA, RNA, TNA or XNA oligonucleotides, single or double stranded, are ligated in parallel and independently in order to obtain a first set of intermediate products that are longer.

[0003] WO2019073072A1 discloses a hierarchical assembly approach, wherein, after completing these first steps, the first intermediate products are again combined in pairs, thereby obtaining even longer polynucleotides. Further steps proceed in the same fashion until a long target polynucleotide is obtained (FIG. 1).

[0004] This and other methods rely on matching overhangs between any two starting oligonucleotides (oligos) or intermediate products and are thus subject to ambiguities either because overhangs other than the intended ones can ligate, resulting in a wrong intermediate product or “mis-assembly”. These assemblies can be represented as graphs where the vertices indicate transfer/pooling steps, and the nodes represent the DNA oligos or intermediate polynucleotides. When the specific workflow is disclosed, the graph is not attributed to specific experimental steps or reaction or to any particular method but is used only to indicate a visual aid to imply the hierarchical nature. Prior art hierarchical assemblies use a symmetrical workflow, a sequential addition of oligos or pooled reactions, and the notion of “hierarchical” is used to denote a tiered, but generally sequence-independent approach. This is understood as a workflow where all oligos react to obtain reaction products that are longer than the components of the reaction products, which is repeated in several steps.

[0005] Although a symmetric hierarchical assembly method is improved over methods using pooled or sequential workflows, it does not in its own prevent mis-assemblies, if there is a variety of constituting building blocks (oligo or polynucleotides) which have a region of sufficient sequence overlap, that could result in an erroneous assembly (see e.g., FIGS. 2,3).

[0006] Although the last few years have seen considerable progress in the techniques for synthesizing DNA, there are still severe restrictions in terms of volume, throughput and, specially, length of DNA.

[0007] WO2021055962A1 discloses compositions and methods for template-free double stranded geometric enzymatic nucleic acid synthesis of arbitrarily programmed nucleic acid sequences.

[0008] WO2018203056A1 discloses a nucleic acid for use in DNA assembly, wherein the nucleic acid comprises at least one methylation-protectable restriction element, the methylation-protectable restriction element comprising: a restriction enzyme recognition sequence that is recognized by a restriction enzyme that cleaves outside of the recogni-

tion sequence; and a DNA methylase recognition sequence, wherein the restriction enzyme recognition sequence and the DNA methylase recognition sequence overlap such that the base modified by the DNA methylase lies within the restriction enzyme recognition sequence, wherein the DNA methylase recognition sequence is not identical to or enclosed by the restriction enzyme recognition sequence, and wherein the DNA methylase recognition sequence does not overlap with the sequence that would form the overhang end sequence generated by the restriction enzyme.

SUMMARY OF THE INVENTION

[0009] It is an objective of the present invention to provide an improved method for synthesizing double stranded (ds) polynucleotides, aiming to reduce the number of mis-assemblies when synthesizing a target ds polynucleotide. It is a further objective of the present invention to provide an improved method for synthesizing ds polynucleotides where a variety of building blocks (oligo or polynucleotides) comprising the same matching region of sequence overlap are used.

[0010] The object is solved by the subject of the present invention.

[0011] According to the invention, there is provided a method of assembling nucleic acid (NA) building blocks to produce a double stranded (ds) target polynucleotide, comprising assembling NA building blocks according to a workflow comprising multiple assembly tiers, wherein at least one asymmetric pair of NA building blocks which are of different length, preferably at least one asymmetric pair of ds NA building blocks, is connected to produce an intermediate polynucleotide in at least one of said multiple assembly tiers, wherein said intermediate polynucleotide is used as a building block to connect further NA building blocks at both ends of said intermediate polynucleotide in one or more further assembly tiers.

[0012] According to a specific aspect, there is provided a method of assembling nucleic acid (NA) building blocks to produce a double stranded (ds) target polynucleotide, comprising assembling NA building blocks according to a workflow comprising multiple assembly tiers, wherein, in a hierarchical assembly, at least one asymmetric pair of NA building blocks of different length that originate from different assembly tiers is connected to produce an intermediate polynucleotide in at least one of said multiple assembly tiers, wherein said intermediate polynucleotide is used as a building block to connect further NA building blocks at both ends of said intermediate polynucleotide in one or more further assembly tiers.

[0013] Specifically, the workflow is an asymmetric assembly workflow. Specifically, the assembly comprises at least one asymmetric assembly of NA building blocks.

[0014] Specifically, the workflow is a hierarchical asymmetric workflow. Specifically, the hierarchical assembly comprises at least one asymmetric assembly of NA building blocks.

[0015] Specifically said workflow comprises at least one tier, which comprises at least one asymmetric assembly of NA building blocks.

[0016] Specifically said workflow comprises at least one tier, in which at least one asymmetric pair of NA building blocks of different length is assembled. Specifically, said at least one asymmetric pair of NA building blocks of different length is assembled by an asymmetric assembly.

[0017] Specifically, the asymmetric assembly is non-geometric.

[0018] Specifically, said hierarchical assembly comprises at least one asymmetric assembly of said at least one asymmetric pair of NA building blocks.

[0019] Specifically, a pair of NA building blocks consists of two NA building blocks with matching sequences to allow an assembly of the two NA building blocks.

[0020] Specifically, a pair of ds NA building blocks, such that the pair of ds NA building blocks is assembled by connecting their ends such as by hybridizing the 5'-end of one NA building block of said pair to the 3'-end of the other one of said pair, to produce an assembled ds polynucleotide or nucleic acid molecule comprising a site of assembly, which is where the matching sequences have hybridized.

[0021] Specifically, said asymmetric pair of NA building blocks is a pair of one ds NA building block and one ss NA building block, or a pair of two ds NA building blocks.

[0022] Specifically, a pair of ds NA building blocks which is an asymmetric pair of ds NA building blocks consists of two ds NA building blocks of different length. In contrast, in a symmetric pair of ds NA building blocks, both NA building blocks have the same length.

[0023] Specifically, the length of an asymmetric pair of NA building blocks may differ such that at least one of the NA building blocks has a shorter length than the other NA building block of said asymmetric pair of ds NA building blocks e.g., at least any one of 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 95%, up to 99% shorter.

[0024] Specifically, the asymmetric pair of ds NA building blocks is a pair of ds NA building blocks, and the difference in length is determined comparing the double stranded part of the ds NA building blocks.

[0025] Specifically, where a pair of two of NA building blocks of different length is assembled, the present disclosure provides for the following embodiments:

[0026] a) both NA building blocks are de novo synthesized and used as a starting NA building block in a first assembly tier; or

[0027] b) one of the NA building blocks is produced in a previous assembly tier, and the other one is de novo synthesized as a starting NA building block; or

[0028] c) both of the NA building blocks are produced by previous assembly tiers, optionally wherein the number of assembly tiers to produce one of the NA building blocks differs from the number of assembly tiers to produce the other one of the NA building blocks.

[0029] According to a specific aspect, one NA building block of said at least one asymmetric pair is a starting NA building block, and said asymmetric pair is assembled in a second or further tier of assembly.

[0030] Specifically, one NA building block of said at least one asymmetric pair is a starting NA building block, in particular a ss or ds NA building block, such as a ss or ds oligo, and the other one of the asymmetric pair is an NA building block which is a reaction product obtained by assembly of NA building blocks in a first, second or further assembly tier.

[0031] By assembling an asymmetric pair of ds NA building blocks, a site of assembly can be chosen that is not in the center of the assembly product, but within a different region that avoids mis-assemblies.

[0032] Specifically, one of the NA building blocks of the asymmetric pair is a reaction product obtained by assembly of NA building blocks in a previous assembly tier, and the other one of the asymmetric pair is a starting NA building block, or a reaction product obtained by assembly of NA building blocks in an assembly tier that is different from said previous assembly tier.

[0033] Specifically, in an assembly tier, one or more pairs of NA building blocks are assembled. The number of assembly tiers, reaction steps and corresponding reaction products is basically determined by the length of the target polynucleotide. In order to synthesize large target polynucleotides, a series of assembly tiers may be needed for assembly into the target polynucleotide e.g., at least 5, 10, 20, 50, 100, 500, 1,000, 5,000 or more may be necessary. Each tier comprises a series of parallel assembly reactions. Typically, the number of assembly reactions decreases with increasing length of the NA building blocks in a tier.

[0034] Specifically, an intermediate polynucleotide as used herein is a ds NA building block which is a product of assembling a pair of NA building blocks. Specifically, the intermediate polynucleotide is used as a building block to connect further NA building blocks at both ends of said intermediate polynucleotide, such as to assembly at least one NA building block in both directions, in one or more further assembly tiers. Assembling of NA building blocks in both directions results in an extension of the 3'-end and an extension of the 5'-end of said intermediate polynucleotide. Such extensions can be done in parallel and/or in the same assembly tier, or consecutively.

[0035] Specifically, the intermediate polynucleotide is assembled with further NA building blocks at its 5'-end and its 3'-end (i.e., at both of its ends) in the same assembly tier, optionally wherein assembly at both ends is carried out simultaneously or consecutively.

[0036] Specifically, the intermediate polynucleotide is assembled with further NA building blocks at its 5'-end and its 3'-end (i.e., at both of its ends) in the different assembly tiers, optionally wherein assembly of one of the ends of the intermediate polynucleotide with a NA building block is carried out in one assembly tier, and assembly of the other end of the intermediate polynucleotide with another NA building block is carried out in another assembly tier.

[0037] In a specific embodiment, the intermediate polynucleotide is assembled with a NA building block at its 5'-end in a further assembly tier, thereby obtaining a reaction product, which reaction product is assembled with another NA building block at its 3'-end in the same or another further assembly tier.

[0038] In any case, assembly of the intermediate polynucleotide with further NA building blocks at both ends can be carried out simultaneously or consecutively.

[0039] Where the intermediate polynucleotide is assembled with further NA building blocks at both ends, such polynucleotide is considered as an intermediate that is different from a prefinal polynucleotide.

[0040] Typically, in a final step of assembly, a prefinal polynucleotide is used that has been produced in a prefinal assembly step, which is then assembled with a final NA building block in the final step of assembly. Optionally, a final NA building block is used which comprises a blunt end, or an end that is designed to match a site of a nucleotide construct (such as a vector or plasmid) to incorporate the target ds polynucleotide in said nucleotide construct. In such

final step of assembly, the prefinal polynucleotide is assembled with another NA building block only at one of its ends.

[0041] Specifically, the NA building blocks are oligonucleotides or polynucleotides e.g., fragments of the ds target polynucleotide.

[0042] Specifically, NA building blocks are either double stranded such as ds oligos or ds polynucleotides, in particular ds intermediate polynucleotides or prefinal polynucleotides, or single stranded such as ss oligos or ss polynucleotides.

[0043] In a specific aspect of the method described herein, pairs of matching ss oligos are assembled to produce ds NA building blocks. Preferably, said pairs of matching ss oligos are transferred into separate reaction containments to produce the respective ds NA building blocks.

[0044] Specifically, the NA building blocks have a length ranging from 8 nucleotides up to 20% of the length of the target polynucleotide.

[0045] In a specific embodiment, ss NA building blocks are ss oligos or ss polynucleotides with a length of at least 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, or even more nucleotides. Specifically, ss NA building blocks can have a length of at least 30, 40, 50, 60, 70, 80, 90, or 100 nucleotides. In specific cases, the length of ss NA building blocks is limited, and may have a length of less than 900, 800, 700, 600, 500, 400, 300, 200 or 100 nucleotides. Specifically, ss NA building blocks may have a length of up to 26, 25, 24, 23, 22, 21, 20, 19, 18, 17 or 16 nucleotides.

[0046] In a specific embodiment, ds NA building blocks are ds oligos or ds intermediate or prefinal polynucleotides, which comprise a double stranded part that has a length of at least 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, or even more base pairs. Specifically, the double stranded part of the ds NA building blocks can have a length of at least 30, 40, 50, 60, 70, 80, 90, or 100 base pairs. In specific cases, the length of the ds NA building blocks is limited, such as ds NA building blocks with a length of the double stranded part of less than 900, 800, 700, 600, 500, 400, 300, 200 or 100 base pairs. Specifically, the double stranded part of said ds NA building blocks have a length of up to 26, 25, 24, 23, 22, 21, 20, 19, 18, 17 or 16 base pairs.

[0047] Specifically, the NA building blocks can have a length of up to any one of 20%, 15%, or 10% of the target nucleotide. Specifically, any building blocks that have a length of at least any one of 5%, 6%, 7%, 8%, 9%, or 10% of the target nucleotide are ds.

[0048] Specifically, a ds NA building block has one or two overhangs, which is at one or both of its ends, in particular at the 5'-end and/or at the 3'-end of the ds NA building block.

[0049] Specifically, a ds NA building block comprising overhangs on both ends and no blunt end may be used as an intermediate polynucleotide.

[0050] Specifically, NA building blocks are assembled according to a defined workflow.

[0051] Specifically, the workflow may provide for the production of ds NA building blocks from respective single stranded (ss) NA building blocks, such as by assembly of ss oligos. Specifically, ds NA building blocks can be produced by assembly of ss NA building blocks before their use in a method to assemble larger polynucleotides.

[0052] Specifically, the workflow may provide for the production of ds NA building blocks from respective ds NA building blocks, such as by assembly of ds oligos. Specifically, ds NA building blocks can be produced by assembly of ds NA building blocks before their use in a method to assemble larger polynucleotides.

[0053] According to a specific aspect described herein, an asymmetric hierarchical workflow is employed.

[0054] Specifically, said workflow is determined using an algorithm.

[0055] Specifically, said algorithm selects pairs of matching NA building blocks and optionally ss NA building block linkers (such as ss oligo linkers), if necessary, and determines the assembly workflow, not by a mere sequence partitioning, but by determining an optimal or near-optimal way to assemble the target ds polynucleotide, avoiding mismatches or undesired reaction products as far as possible.

[0056] NA building blocks and assembly workflow are specifically selected to avoid undesired (incorrect) reactions or reaction products, such as palindromic sequences, runaway reactions and unambiguous assembly.

[0057] Specifically, the assembly workflow is automated.

[0058] Specifically, the automated workflow employs microfluidic handlers that are capable of transferring serially or in parallel the full or partial contents of one or several compartments into other pre-specified compartments that may or may not be empty.

[0059] In certain embodiments of the method described herein, purification of the reaction products may be necessary at one more assembly step. If there are incorrect reaction products besides the correct reaction products, such incorrect reaction products may be suitably separated from the correct ones e.g., as follows: using gel electrophoresis to detect oligonucleotides or polynucleotides of a certain size and excising and purifying bands of the gel corresponding to the size of the desired reaction product. Specifically, correct reaction products can be detected by incorporation of tags or labels into the sequence. Specifically, oligos or polynucleotides may be captured using biotinylated oligonucleotide adapters capable of hybridizing with an overhang of a NA building block wherein, said adapters are fixed to the substrate and coated with streptavidin. Non-captured incorrect products are eliminated by washing and subsequently, the correct products are released from the adapters by increasing the temperature. Specifically, further separation methods well-known in the art may be applied. Specifically, such methods may involve chromatographic or affinity separation methods.

[0060] According to the method provided herein, NA building blocks are assembled to intermediate polynucleotides and ultimately the target polynucleotide.

[0061] Specifically, said assembly is by connecting matching sequences.

[0062] Specifically, said assembly is by connecting a matching ss NA building block to another ss NA building block and/or to an overhang of a ds NA building block; and/or by connecting matching overhangs of a pair of ds NA building blocks.

[0063] Specifically, matching NA building blocks or respective parts of the NA building blocks such as overhangs comprise or consist of ss nucleotide sequences which are sufficiently complementary to each other, to allow hybridizing said nucleotide sequences.

[00664] Specifically, assembly is by hybridizing complementary nucleotide sequences (or sufficiently complementary sequences).

[00665] According to a specific aspect, assembly may also include connecting NA building blocks by a ligation reaction which comprises enzymatic, chemical, or an adaptor ligation.

[00666] Specifically, said ligation reaction is an enzymatic ligation reaction using a ligase, preferably a ligase, such as any one of a T3, T4 or T7 DNA ligase, or a RNA ligase, a polymerase or ribozymes. Preferably T4 DNA ligase, T7 DNA Ligase, T3 DNA Ligase, Taq DNA Ligase, DNA polymerase, or engineered enzymes are used in the ligation reaction. Preferably, the following ligation reaction is used: T4 DNA Ligase, at a concentration of 10 cohesive end units per L supplemented with 1 mM ATP (Sambrook and Russell, 2014, Chapter 1, Protocol 17).

[00667] Specifically, said assembly is directly by hybridizing matching overhangs of a ds NA building block, or indirectly by hybridizing a suitable ss NA building block linker (i.e., a ss NA linker), which ss NA linker can be a ss oligo or ss polynucleotide, optionally contained in and obtained from a respective library.

[00668] In a specific embodiment of the method described herein, a solid carrier is used to immobilize one or more of said NA building blocks, the target polynucleotide, or one or more intermediate(s) of assembly. Specifically, said nucleic acids may be immobilized on a solid phase using a tag, for example a biotin tag.

[00669] Specifically, the NA building blocks are provided in separate containments e.g., a library of NA building blocks. A library of NA building blocks is conveniently used as a source of NA building blocks, which can be selected from the library on demand, such as to allow assembling the selected NA building blocks according to a pre-defined workflow.

[0070] Specifically, ss and ds NA building blocks can be provided in separate containments of a library, which can be readily used as a source of NA building blocks for the purpose described herein. In a specific embodiment, ds NA building blocks are prepared in and sourced from the same containments.

[0071] In a specific embodiment of the method described herein, said NA building blocks are obtained from a library of NA building blocks, which comprises a diversity of NA building blocks, each contained in a separate library containment.

[0072] Specifically, said library comprises ss NA building blocks, preferably wherein said ss NA building blocks have been obtained by de novo synthesis, such as by chemical or enzymatic means.

[0073] Specifically, said library comprises ds NA building blocks, preferably produced from respective ss NA building blocks from the library.

[0074] The library described herein specifically comprises or consists of library members which are ss and/or ds NA building blocks, preferably the library comprises or consists of both of ss and ds NA building blocks. Preferably, library members are pre-built, optionally provided in a storage stable form, and located at defined positions within an array device. Library members can be synthesized and stored in the array device until needed.

[0075] Specifically, the NA building blocks can be produced by any suitable prior art method, such as chemical

polynucleotide (or oligonucleotide) synthesis methods, including the H-phosphonate, phosphodiester, phosphotriester or phosphite triester synthesis methods, or any of the massively parallel oligonucleotide synthesis methods e.g., microarray or microfluidics-based oligonucleotide synthesis e.g., as described in Gao et al. 2001, LeProust et al. 2010, or Bonde et al. 2014.

[0076] Specifically, the NA building blocks can be produced by any of the enzymatic polynucleotide (or oligonucleotide) synthesis methods e.g., ssDNA synthesis by DNA polymerase proteins or by reverse transcriptase proteins, which produce hybrid RNA-ssDNA molecules. Specifically, the enzymatic polynucleotide synthesis reaction is performed in vitro.

[0077] Specific embodiments refer to NA building blocks which are modified by any one or more of phosphorylation, methylation, biotinylation, or linkage to a fluorophore or quencher. Therefore, the library described herein may comprise library members which are NA building blocks that can be any or all of the following: unmodified ss; phosphorylated ss; methylated ss; biotinylated ss; phosphorylated, biotinylated and methylated ss; unmodified ds; phosphorylated ds; methylated ds; biotinylated and phosphorylated ds; biotinylated and methylated ds. Preferably, library members comprise a 5'-phosphorylation. Specifically, the library described herein comprises ss NA building blocks comprising fluorophores or quenchers and ds NA building blocks comprising fluorophores or quenchers.

[0078] Specifically, said NA building blocks are provided in a storage-stable form, preferably a form which is storage-stable for at least 6 months at room-temperature.

[0079] Specifically, the library of NA building blocks described herein is storage stable, comprising the NA building blocks in storage containments.

[0080] Specifically, the library may comprise dried NA building blocks. NA building blocks can be stored in storage containments in a dry state. Dry-state is, for example, achieved by lyophilization, freeze drying, evaporation, crystallization or the like. The enzymes which catalyze the degradation of nucleic acids are typically active at room temperature in a fluid biomolecule preparation. Dry-state storage inhibits such enzymatic activity because such enzymes are generally inactive upon de-hydration and because the degradative chemical reactions which they catalyze typically entail the addition of water (i.e., hydrolysis) of a protein or nucleic acid molecule, thus producing protein or nucleic acid backbone cleavage. In the dry state, there is little or no water (e.g., less than 5%, 4%, 3%, 2% or 1% (w/w) water) as a chemical reactant to support such enzyme catalysis. Additionally, any non-enzymatic hydrolysis of protein or nucleic acid is similarly inhibited, since water is generally unavailable for such reactions.

[0081] In a specific embodiment, the ds target polynucleotide produced according to the method described herein, is provided and/or stored in a dry state.

[0082] Specifically, a reaction containment or a storage containment is a compartment unit, such as a well, of any one of a microtiter plate, a microfluidic microplate, a set of capillaries, a microarray or a biochip, preferably a DNA and/or RNA biochip.

[0083] Specifically, for hybridization of matching nucleic acid molecules, said matching nucleic acid molecules are placed in a reaction containment that features an environment in which one nucleic acid strand bonds to a second

nucleic acid strand by complementary strand interactions and hydrogen bonding to produce a double stranded oligonucleotide or polynucleotide. Such conditions include a mixture of the matching nucleic acid molecules as reaction components in an aqueous or organic solution, further components (e.g., salts, chelating agents, formamide) and their concentrations in said solution, and the temperature of the mixture. Other well-known factors, such as the length of incubation time or reaction chamber dimensions may contribute to the environment.

[0084] Specifically, the NA building blocks described herein are produced by synthesizing the oligonucleotide or polynucleotide sequence from nucleotide building blocks by any of the polynucleotide synthesis methods, wherein the building blocks are comprised of “A” denoting deoxyadenosine, “T” denoting deoxythymidine, “G” denoting deoxyguanosine, or “C” denoting deoxycytidine or other natural nucleosides (e.g. adenosine, thymidine, guanosine, cytidine, uridine), nucleotide-analogs e.g., inosine and 2'-deoxyinosine and their derivatives (e.g. 7'-deaza-2'-deoxyinosine, 2'-deaza-2'-deoxyinosine),azole- (e.g. benzimidazole, indole, 5-fluoroindole) or nitroazole analogues (e.g. 3-nitropyrrol, 5-nitroindol, 5-nitroimidazole, 4-nitropyrazole, 4-nitrobenzimidazole) and their derivatives, acyclic sugar analogues (e.g. those derived from hypoxanthine- or indazole derivatives, 3-nitroimidazole, or imidazole-4,5-dicarboxamide), 5'-triphosphates of universal base analogues (e.g. derived from indole derivatives), isocarbostyryl and its derivatives (e.g. methylisocarbostyryl, 7-propynylisocarbostyryl), hydrogen bonding universal base analogues (e.g. pyrrolopyrimidine), or any of the other chemically modified bases (such as diaminopurine, 5-methylcytosine, isoguanine, 5-methyl-isocytosine, K-2'-deoxyribose, P-2'-deoxyribose). Likewise, the NA building blocks may comprise “U” denoting uracil, a deoxyuridine, or deoxyuridine site. An enzyme Uracil DNA glycosylase can be used to transform the deoxyuridine site into an abasic site, which subsequently can be cut using glycosylate-lyase endonuclease VIII, thereby releasing the DNA.

[0085] The building blocks are linked by phosphodiester linkage or peptidyl linkages or by phosphorothioate linkages or by any of the other types of nucleotide linkages.

[0086] Specifically, the target polynucleotide has a length of at least 132 base pairs (bps). Specifically, said target polynucleotide has a length of at least 150, 200, 250, 300, 400, 500, 600, 700, 800, 900, 1,000 bps.

[0087] Specifically, by the method described herein target polynucleotides with a length up to 1,000, 5,000, 10,000 or 100,000 bps or even longer can be produced, at a low price and at a high speed.

[0088] Specifically, the nucleotide sequence of a target ds polynucleotide or respective template can be of one of natural or artificial origin.

[0089] Specifically, the target ds polynucleotide has blunt ends on both ends.

[0090] According to a specific aspect, a blunt end of the target ds polynucleotide can be produced by using a pair of NA building blocks wherein one of the NA building blocks comprises a blunt end, thereby obtaining a terminal NA building block with a blunt end. According to another specific aspect, a ss NA building block and a ds NA building block are assembled, wherein the ss NA building block is complementary to an overhang of the ds NA building block,

such as to hybridize the complementary sequences without generating any further overhang, thereby producing a blunt end.

[0091] Specifically, the method provided herein comprises a finalization step.

[0092] Specifically, said finalization step serves to add one or more nucleotide(s) which correspond to those previously removed from the 3'-end and 5'-end, respectively, to prepare a template of such target ds polynucleotide for the purpose of assembly of the target ds polynucleotide according to a template sequence, such as e.g., to generate blunt ends. Specifically, one or more NA building blocks may be selected for producing blunt ends, which are complementary to any overhang of a prefinal intermediate polynucleotide i.e., complementary to the sticky ends of the polynucleotide. Specifically, respective oligos can be used as primers in a PCR reaction to amplify the final product and to add the remaining NA building blocks to each strand to synthesize the complete target polynucleotide with blunt ends.

[0093] Specifically, said finalization step comprises a purification step of a PCR product that has been produced employing standard kits, such as the Monarch PCR & DNA clean up kit from New England Biolabs (product no. T1030), to eliminate remaining NA building blocks, oligos, enzymes and reagents, thereby obtaining the target ds polynucleotide as a purified DNA product, ready for further use.

[0094] According to a specific aspect, the method described herein further comprises a step of enriching the target polynucleotide or one or more intermediates of assembly, by polymerase chain reaction (PCR).

[0095] Specifically, the target polynucleotide or one or more intermediates of assembly may be purified by immobilization on a solid phase using a tag, for example a biotin tag, and enrichment using, e.g., PCR amplification. According to a preferred embodiment, two sets of primers are used for target specific enrichment and simultaneous elimination of the tag. Specifically, by using a set of primers specific to the 5' end of the leading strand and a set of primers specific to the 5' end of the lagging strand of the polynucleotide that is to be enriched, each comprising a primer that is complementary to at least the overhang and a primer that is complementary to the core sequence of the polynucleotide, the target polynucleotide is amplified without the tag sequence. This has the profound advantage that no additional step is required to remove the tag sequence, e.g., by enzymatic digestion.

[0096] The degree of purification is understood as the amount of NA building block copies per volume or per total (poly) nucleotide mass. Various methods to determine the degree of purity of a preparation of nucleic acid molecules are known to a person skilled in the art. Specifically, the degree of purity may be determined using gel electrophoresis, next generation sequencing or qPCR.

[0097] In a specific embodiment of the invention, said target ds polynucleotide is sequenced to verify the degree of identity with the sequence of a template or a sequence of interest (SOI). Any suitable sequencing method may be used, for example any one of SNP genotyping methods, including hybridization-based methods (e.g. molecular beacons, SNP microarrays, restriction fragment length polymorphism, PCR-based methods, including Allele-specific PCR, primer extension-, 5'-nuclease or Oligonucleotide Ligation Assay, Single strand conformation polymorphism, Temperature gradient gel electrophoresis, Denaturing high perfor-

mance liquid chromatography, High-resolution Melting of the entire amplicon (HRM), SNPlex and surveyor nuclease assay; Sequencing based mutation analysis, including capillary sequencing or high-throughput sequencing of an entire PCR amplicon of the PTR (amplicon sequencing). Such high-throughput (HT) amplicon sequencing methods include, but are not restricted to polony sequencing, pyro-sequencing, Illumina (Solexa) sequencing, SOLID sequencing, semiconductor sequencing, DNA nanoball sequencing, Heliscope single molecule sequencing, Single molecule real time (SMRT) sequencing, Nanopore DNA sequencing, tunnelling currents DNA sequencing, sequencing by hybridization, sequencing with mass spectrometry, Microfluidic Sanger sequencing, Microscopy-based sequencing, RNAP sequencing.

[0098] In a specific embodiment, the method described herein further comprises a step of processing the target polynucleotide or intermediates of assembly, by means of enzymatic modification employing restriction enzymes or kinases, or by chemical means, to facilitate further ligation, or to facilitate cloning the target polynucleotide into a vector or plasmid.

[0099] In a further specific embodiment of the invention, the target ds polynucleotide can be further modified to produce a derivative thereof, which is any of a ds DNA, ss DNA or RNA molecule, e.g. comprised in a vector, such as a plasmid.

[0100] Specifically, said target ds polynucleotide can be modified by enzymatic modification, employing any one or more of methyltransferases, kinases, CRISPR/Cas9, multiplex automated genome engineering (MAGE) using A-red recombination, conjugative assembly genome engineering (CAGE), the Argonaute protein family (Ago) or a derivative thereof, zinc-finger nucleases (ZFNs), transcription activator-like effector nucleases (TALENs), meganucleases, tyrosine/serine site-specific recombinases (Tyr/Ser SSRs), hybridizing molecules, sulfurylases, recombinases, nucleases, DNA polymerases, RNA polymerases or TNases.

FIGURES

[0101] FIG. 1. An example of a symmetric assembly tree with the corresponding tiers indicated.

[0102] FIG. 2. An example of an asymmetric assembly tree where two components (c and d) would fail to assemble, causing all the connected nodes to fail as well.

[0103] FIG. 3. (a) Double stranded oligonucleotides with sticky ends used as assembly reagents (Sequence IDs 18-25). (b-c) Alternative assembly trees for the four reagents.

[0104] FIG. 4. (a) Method for calculating the Adjacency Matrix for two ds oligonucleotides with given overhangs. (b-c) Examples of the 4x4 adjacency matrices with minor modifications to the overhangs.

[0105] FIG. 5. (a) An example of a symmetric assembly tree with the corresponding scores indicated for every node. (b) The optimised version of the assembly tree where the asymmetric structure resolves the problematic nodes and gives an overall acceptable score. (c) The two Ligation Matrices used to calculate the scores. The non-zero elements are the ones corresponding to overhangs with matching sequences, and the gray areas are the ones that will cause misligations.

[0106] FIG. 6. The introduction of “zeromers” in the automated assembly setup allows the use of repetitive,

symmetric movements of a liquid handler arm to assemble a molecule following an asymmetric tree.

[0107] FIG. 7. (a) The layout of reagents in a plate used to assemble a molecule following a symmetric tree. (b) The layout of reagents in a plate used to assemble a molecule following an asymmetric tree.

[0108] FIG. 8. Nucleotide sequences referred to herein.

[0109] FIG. 9. Misligation matrices for symmetric and asymmetric assembly trees, used to calculate the scores.

[0110] FIG. 10. (a) The symmetric assembly of a 256 bp molecule shows two problematic nodes that cause the assembly to fail. (b) Optimized asymmetric assembly solves the issues caused by the misligations and enables the assembly.

[0111] FIG. 11. Electropherograms of the final assembly products of symmetric (continuous line) and asymmetric (dashed line) assemblies. The correct size molecule (256 bp) is only visible in the asymmetric case.

[0112] FIG. 12. Three examples of assembly trees/partitions for the same SOI, showing how the score is maximized by many folds when the adaptive partitioning method is combined with the optimization of the asymmetric assembly tree.

[0113] FIG. 13 shows a graphical representation of the product of the ligation and misligation matrix (Example 5).

[0114] FIG. 14 shows a list of oligo sequences resulting from partitioning a SOI (Example 5).

[0115] FIG. 15 shows a list of oligo sequences resulting from adaptive partitioning a SOI (Example 5).

[0116] FIG. 16: graphical representation of assembly trees: A) Fully symmetric tree. At each tier, reaction products from the same previous tier are used to have further reaction products. The “leaves” of the tree are the starting oligos. T=Tiers. R.P.=Reaction products. Note that at every step the number of reaction product decreases, in particular, they halve at every step; B) Asymmetric tree. The reactions may involve products (double dashed arrows) or starting oligos (double arrows) that are from any other previous tier (not all instances indicated). The “leaves” of the tree are the starting oligos. T=Tiers. R.P.=Reaction products. Note that at any step the number of reaction product may increase and do not halve at every step.

DETAILED DESCRIPTION

[0117] Specific terms as used throughout the specification have the following meaning.

[0118] As used herein, the terms “a”, “an” and “the” are used herein to refer to one or more than one i.e., to at least one. The terms “comprise”, “contain”, “have” and “include” as used herein can be used synonymously and shall be understood as an open definition, allowing further members or parts or elements. “Consisting” is considered as a closest definition without further elements of the consisting definition feature. Thus “comprising” is broader and contains the “consisting” definition.

[0119] The term “5'-end” or “3'-end” of a NA building block is herein understood to refer to the side or orientation of an end of the NA building block, and specifically understood in the following way.

[0120] The 5'-end of a ss NA building block is understood to comprise or consist of the 5'-terminal region e.g., a part of the ss oligo or polynucleotide, such as e.g., comprising or consisting of at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, nucleotides or more e.g., at least 10%, 20%, 30%, 40%, 50%, 60%, 70%,

80%, 90%, 95%, 96%, 97%, 98%, 99%, up to or close to 100% of the full length ss molecule, which 5'-terminal region is including the 5'-terminus, i.e. the 5'-terminal nucleotide.

[0121] The 3'-end of a ss NA building block is understood to comprise or consist of the 3'-terminal region e.g., a part of the ss oligo or polynucleotide, such as e.g., comprising or consisting of at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, nucleotides or more e.g., at least 10%, 20%, 30%; 40%, 50%, 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, 99%, up to or close to 100% of the full length ss molecule, which 3'-terminal region is including the 3'-terminus, i.e. the 3'-terminal nucleotide.

[0122] The 5'-end of a ds NA building block is understood to comprise or consist of the 5'-terminal region of the leading strand e.g., a part of the leading strand of the ds oligo or polynucleotide, such as e.g., comprising or consisting of at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, nucleotides or more e.g., at least 10%, 20%, 30%; 40%, 50%, 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, 99%, up to or close to 100% of the full length ds molecule, which 5'-terminal region is including the 5'-terminus, i.e. the 5'-terminal nucleotide.

[0123] The 3'-end of a ds NA building block is understood to comprise or consist of the 3'-terminal region e.g., a part of the leading strand of the ds oligo or polynucleotide, such as e.g., comprising or consisting of at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, nucleotides or more e.g., at least 10%, 20%, 30%; 40%, 50%, 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, 99%, up to or close to 100% of the full length ds molecule, which 3'-terminal region is including the 3'-terminus, i.e. the 3'-terminal nucleotide.

[0124] Typically, assembly of the NA building blocks as described herein is by end-to-end ligating or connecting (e.g., annealing, hybridizing or linking), wherein a 5'-end of a NA building block is connected to the 3'-end of another NA building block. NA building blocks can be ss or ds. A ss NA building block can be connected to another ss NA building block, or a ds NA building block, and vice versa. A ds NA building block can be connected to another ds building block. The assembly products will comprise both assembled NA building blocks.

[0125] Single-strand annealing of complementary sequences may lead to ds NA building blocks with overhangs or blunt ends. Assembly of a pair of ss NA building blocks typically results in a ds NA building block, wherein one ss NA building block is the leading strand and the other one is the lagging strand. In such case, assembly is not an end-to-end assembly, but a side-to-side assembly to a ds molecule, wherein the 5'-end and 3'-end of the ss NA building block which is comprised as a leading strand defines the 5'-end and 3'-end of the ds molecule, respectively.

[0126] The term “algorithm” as used herein refers to a self-contained sequence of actions to be performed. An algorithm is an effective method that can be expressed within a finite amount of space and time and in a well-defined formal language for calculating a function. Starting from an initial state and initial input the instructions describe a computation that, when executed, proceeds through a finite number of well-defined successive states, eventually producing “output” and terminating at a final ending state. The transition from one state to the next is necessarily deterministic.

[0127] The term “assembly” or “assemble” refers to the formation of an oligonucleotide or polynucleotide by a ligation reaction e.g., linking and/or hybridizing single stranded and/or double stranded NA building blocks. Specifically, said assembly is by any method of annealing or hybridizing nucleotide sequences which are complementary to each other, or comprise regions that are complementary or at least partially complementary to each other.

[0128] Assembly of the target ds polynucleotide can either be directly by hybridizing matching ss NA building blocks, overhangs of ds NA building blocks, or indirectly by hybridizing one or more suitable ss NA building block linkers, wherein e.g., a ss oligo linker is used, to allow assembling a pair of NA building blocks (e.g., both ss or ds, or one being ss and the other one being ds) that would not hybridize to each other without using the linker(s).

[0129] For direct assembly, NA building blocks are joined together by their single stranded parts or overlaps (i.e., the overlapping parts or overhangs), such that the overlaps are included in the continuous sequence only once. Upon aligning two NA building blocks with an overlap, a continuous sequence is formed which has a length that is the length of both individual NA building blocks taken together, minus the length of the overlap. Consequently, a continuous sequence is obtained which comprises a segment of each of the aligned NA building blocks.

[0130] For indirect assembly, the target ds polynucleotide or any intermediates can be formed upon assembly of ss NA building blocks and joining them through one or more ss linker(s). For example, two ss NA building blocks, each of e.g., 10 bases length, may be joined by a ss oligo linker of e.g., 6 bases length, such that 3 bases of the 3' terminal end of the first ss NA building block align with the 3 bases of the 5' end of the ss linker and that 3 bases of the 5' end of the second ss NA building block align with the 3 bases of the 3' end of the ss linker.

[0131] Single stranded oligo linkers having a sequence complementary to the combined overhangs may connect adjacent NA building blocks in the target polynucleotide. Linkers may e.g., consist of 6 bases to connect two adjacent NA building blocks, each with a 3 base long overhang, one on the 3' end and the other on the 5' end, respectively.

[0132] Each assembly will result in an extension of one or both ends of an NA building block, such as an extension of the 5'-end and/or the 3'-end of an NA building block.

[0133] Multiple parallel reactions can be carried out. Specifically, an assembly tier is characterized by at least two assembly reactions in parallel, in particular wherein the respective reaction products will be further reacted in one or more further steps. Specifically, one or more parallel assembly reactions can be performed in separation reaction containments during one assembly period (“tier” or “round”), before one or more further assembly reactions are performed in any of the following tiers, using assembly product(s) of any of the previous tiers.

[0134] The term “single-tier” refers to a single compartment assembly method, such as to allow pooled assembly of NA building blocks. For example, all starting oligos can be assembled in a pooled way i.e., at the same time in one reaction containment (“one pot”).

[0135] The term “multiple-tier” refers to a sequential multi compartment assembly method, where assembly reactions are performed in parallel during one tier, and reaction products obtained by such parallel assembly reactions are

used as intermediates to perform further assembly steps in one or more further tiers of assembly.

[0136] Specifically, the method described herein is a multiple-tier assembly method to produce the ds target polynucleotide in more than one tiers of assembly reactions e.g., at least two, three, four, five, six, seven, eight, nine, or ten tiers, or even more than 10 tiers such as at least 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20, up to a number of tiers which is conveniently used in the synthesis of long target polynucleotides e.g., up to 30, or up to 20, or up to 10 tiers.

[0137] Assembly tiers are typically sequential. Specifically, an assembly tier starting with assembly of only starting NA building blocks is termed “tier 1”. Assembly of reaction products obtained by such tier 1 assembly, are typically used in one or more further tiers of assembly, which can be consecutive or non-consecutive.

[0138] Multiple-tier assembly is herein specifically understood as a multi-part, modular assembly method. Contrary to single-tier or “one-pot” type of assembly, it particularly refers to a hierarchical or sequential assembly, where the component DNA pieces are assembled in.

[0139] Multi-tier DNA assembly methods can be performed with methods that combine or include e.g., Golden Gate assembly, BASIC, BioBrick assembly (and variants such as BglBrick) and Gateway cloning. Bespoke multi-tier DNA assembly methods can also include Gibson Assembly, AQUA cloning, Twin Primer Assembly, ligase cycling reaction, SLIC, SLICE, overlap extension PCR and CPEC.

[0140] To compare, single-tier assembly is understood to assembly a polynucleotide in one step, e.g., using pool reactions or “one-pot” assembly reactions.

[0141] The term “assembly workflow” or simply “workflow” as used herein refers to the pre-defined sequence of assembly of NA building blocks to the target ds polynucleotide. The workflow is typically optimized to allow efficient synthesis of the target polynucleotide, aiming to avoid mismatches of NA building blocks which would lead to a polynucleotide comprising an incorrect sequence.

[0142] The workflow is specifically designed to avoid mismatches or reaction products which cannot be used for assembly to produce the target ds polynucleotide. If there are partial constructs that can anneal in alternative ways, a runaway i.e., an uncontrolled polymerization reaction, can occur. To avoid combinations of pairs of matching NA building blocks that would result in unwanted constructs or runaway reactions, the pairs of matching NA building blocks are assembled in a predetermined sequence of assembly steps i.e., a specific workflow. Preferably, said specific workflow is not linear but hierarchical i.e., following an algorithm that provides for intermediate reaction products which are defined non-consecutive parts of the target ds polynucleotide conveniently produced avoiding undesired reaction products to the extent possible, before such intermediate reaction products are further assembled into further intermediate reaction products or into the target ds polynucleotide sequence.

[0143] In a linear workflow, the polynucleotide is assembled in a linear fashion starting at the 3' end of the leading strand, and adding the next oligo to link the 3' end of the leading strand with the 5' end of the next oligo. For example, oligo B is ligated to oligo A, oligo C is ligated to oligo B, oligo D is ligated to oligo C and so forth. This assembly may be achieved simultaneously by adding all oligos to the reaction containment at the same time (such as

in a pool assembly), or the polynucleotide is extended progressively by successively adding oligos A, B, C, D and so forth to the reaction containment.

[0144] Specifically, a linear assembly workflow employs only a starting oligo at a time, reacting it with an intermediate product (a reaction product of the previous tier) of the growing ds polynucleotide. In this case there are as many tiers as starting oligos, always using a starting oligo at each “tier”.

[0145] In contrast, a hierarchical assembly of NA building blocks, is a tiered molecular assembly where at least two reaction products of a previous tier are used in further tiers.

[0146] In a hierarchical workflow, the polynucleotide is typically assembled in at least two tiers. To generate a target polynucleotide A-B-C-D, in the first tier, oligo B is ligated to oligo A, oligo D is ligated to oligo C, producing the reaction products A-B and C-D. In the second tier, the reaction product A-B is ligated to C-D, thereby producing the target polynucleotide A-B-C-D.

[0147] However, undesired reaction products may be obtained where oligos A and B are capable of hybridizing not only in the orientation A-B but also B-A due to complementary sequences or overhangs. A linear hierarchical workflow as described above would result in the unwanted polynucleotide A-B-A-B-C-D, and fragments like A-B-A or B-A, in addition to the desired polynucleotide A-B-C-D. Such unwanted side reactions drastically reduce the yield and purity of the target polynucleotide.

[0148] According to a specific aspect described herein, an asymmetric hierarchical workflow is employed. Accordingly, in the above example, only oligos B and C are ligated in a first tier, producing the ds NA building block B-C. In a second tier, the ds NA building block B-C is reacted with the ss NA building block D to produce the intermediate polynucleotide B-C-D. The intermediate polynucleotide B-C-D is then connected with the ss building block A, wherein upon ligation the desired polynucleotide A-B-C-D is formed.

[0149] Geometric (also called “symmetric”) assembly is a type of hierarchical assembly where at every step, pairs of assembly products of a same previous tier are used, thereby halving (or nearly halving, in case of odd number of reactions) the number of NA building blocks to be assembled, at every tier, see specific Examples A and B below.

[0150] Example A: 16 reaction products of previous tier are used, to obtain 8 intermediates, and then these 8 intermediates are used to get further 4 intermediates, in turn used to get 2 intermediates and a final reaction of 1 molecule (the target one). In this case, 8 starting oligos were used in the first tier, to result in 4 intermediates which are assembled in the second tier, to result in 2 intermediates which are assembled in the third tier, to result in one target polynucleotide.

[0151] Example B: Two reaction products of a tier 1 and two reaction products of tier 2 are assembled in parallel, to get two intermediates (of different length). In a following tier, the two intermediates are assembled to obtain a target molecule. In each step, the assembly was symmetric, because the number of NA building blocks used in each previous tier was reduced by half.

[0152] According to a specific aspect as described herein, a specific type of hierarchical assembly is performed which is an asymmetric (also called “non-geometric”) assembly. In such asymmetric assembly method, NA building blocks are

assembled in multiple assembly tiers, wherein in at least one tier, an asymmetric pair of NA building blocks is assembled; see specific Example C below

[0153] Example C: in a first reaction, a reaction product of tier 3 is assembled with a starting oligo, and in parallel, in a second reaction, a reaction product of tier 2 is assembled with a reaction product of tier 3, followed by another tier where in a third reaction, the reaction products of the first and second reactions are combined (assembled), and in yet another tier, the reaction product of the third reaction is reacted with another NA building block (from any tier).

[0154] Specifically, at least one NA building block of the asymmetric pair of NA building blocks is a starting oligo (e.g., a ss oligo), or a reaction product from an earlier tier where at any assembly tier reaction product of any previous tier (not the immediate previous tier), is used.

[0155] According to a specific aspect, as described herein, an asymmetric, hierarchical assembly workflow is employed, which comprises the assembly of at least one asymmetric pair of NA building blocks. Such asymmetric pair of NA building blocks particularly consists of two NS building blocks of different length, which originate from different assembly tiers.

[0156] The asymmetric, hierarchical assembly method can involve also reacting two products of the same previous tier, but not every reaction at every tier is composed from reaction products of the previous tier. In this case, the number of reaction products is not halved at each tier.

[0157] Specifically, NA building blocks can be assembled in one particular tier, wherein at least one NA building block originates from an assembly tier other than the preceding tier. For example, at least one NA building block is assembled with another NA building block, wherein said NA building blocks originate from tiers that are at least one or more e.g., at least two, three, four, five, six, seven, eight, nine, or ten tiers different.

[0158] According to a specific example of a multi-tier assembly method, one NA building block of the asymmetric pair of NA building blocks is a starting oligo e.g., an ss oligo, and the other NA building block of the asymmetric pair of NA building blocks, can be a reaction product (e.g., a ds NA building block) of any preceding tier.

[0159] According to a specific example, one of the NA building blocks is a starting NA building block, and the other NA building block that is assembled with the starting NA building block, has been obtained as a reaction product of the tier 1 assembly, or of any of the further tiers of assembly, up to the immediate preceding assembly tier which results in the ds target polynucleotide.

[0160] According to another specific example of a multi-tier assembly method, the two NA building blocks can be ds NA building blocks which are each reaction products from assembly reactions of different tiers

[0161] Specifically, unwanted side products of assembly can be avoided by changing the length of the respective NA building blocks thereby changing the overhang sequence. However, changing the length or the sequence of oligos is often not possible or desired, e.g., when oligos are derived from restriction digest or from a library comprising oligos of a specific length only, making assembly of the desired polynucleotide difficult or impossible.

[0162] In one specific embodiment described herein, the process of determining the assembly workflow is carried out by an algorithm. Candidate divisions of the sequence of the

template are systematically examined to find the optimal number, length of the NA building blocks of pairs of NA building blocks (including e.g., asymmetric pairs of different length), and assembly sequence of subsets to divide it into for synthesis in accordance with the method provided herein. Initially, the entire target sequence can be taken as a single subset, after which smaller and smaller subsets are formed with increasing numbers of candidate NA building blocks in decreasing size until a partitioning is found that fulfills the subset criteria listed above.

[0163] In the method provided herein, a SOI which is the sequence of a target polynucleotide or the sequence of a template thereof, may be divided into sub-sequences, corresponding to subsets of NA building blocks, avoiding particular nucleotide synthesis problems, such as palindromic sequences, runaway reactions and unambiguous assembly. In particular, such division into shorter NA building blocks may be very efficient to shorten the assembly process and to avoid the need of separating unwanted reaction products. Specifically, ligation of subsets of NA building blocks yields intermediate reaction products, also called intermediates, and assembly of intermediate reaction products ultimately yields the target ds polynucleotide. Preferably, additional criteria to those listed above may be used for selecting subsets of NA building blocks. Such additional criteria include, but are not limited to, minimization of the size of the subset of NA building blocks employed in any single ligation reaction (for example to avoid mismatch ligations), minimizing the difference in annealing temperature of members of a subset of NA building block precursors, minimizing the difference in annealing temperatures of the overhangs of different ds subunits, whether to employ frame-shifting adaptors or ss oligo linkers and whether to minimize the degree of cross-hybridization among the hybrid forming portions of different NA building blocks that make up a subset.

[0164] Specifically, the method described herein employs an “asymmetric assembly workflow”, comprising multiple assembly tiers, wherein in at least one assembly tier at least one asymmetric pair of NA building blocks is assembled. To avoid combinations of pairs of matching NA building blocks that would result in unwanted constructs or runaway reactions, pairs of matching NA building blocks are assembled in a predetermined sequence of asymmetric assembly steps. Preferably, said specific workflow follows an algorithm that provides for intermediate reaction products which are defined non-consecutive parts of the target ds polynucleotide conveniently produced avoiding undesired reaction products to the extent possible, before such intermediate reaction products are further assembled into further intermediate reaction products or into the target ds polynucleotide sequence.

[0165] The term “asymmetric pair of NA building blocks” as used herein shall refer to two NA building blocks, which differ in length. Specifically, said asymmetric pair of NA building blocks is assembled in one assembly tier, optionally followed by assembly of a further asymmetric pair of NA building blocks in a further assembly tier, wherein one of said NA building blocks of said further asymmetric pair is a product of assembly of said asymmetric pair in the previous assembly tier. In an asymmetric workflow, at least one pair of asymmetric NA building blocks is assembled in at least one tier, preferably in at least two or more tiers asymmetric pairs of NA building blocks are assembled.

[0166] In a symmetric workflow, NA building blocks of equal length are assembled e.g., in each assembly tier.

[0167] A nucleic acid (NA) building block is herein particularly understood as an oligonucleotide or polynucleotide which is used in the assembly of the target polynucleotide described herein. NA building blocks may be single stranded or double stranded oligonucleotides or polynucleotides as described herein. Upon assembly in multiple tiers, they form the target polynucleotide.

[0168] “Pairs of matching NA building blocks” are herein understood as two (in some particular cases more than two, e.g., three) NA building blocks which comprise matching nucleotide sequences that are complimentary to each other, or at least partially complementary to each other, such as to allow ligation or a hybridization reaction between the matching nucleotide sequences. Sufficiently complementary nucleotide sequences can be at least 50%, 60%, 70%, 80%, more 90%, 95%, 96%, 97%, 98%, 99%, or 100% complementary to each other.

[0169] Specifically, matching nucleotide sequences are of regions of two single stranded nucleic acids or overhang parts of ds nucleic acids, and have a nucleotide base composition that allow the respective single-stranded regions to anneal to a stable, double-stranded hydrogen-bonded region, in particular applying stringent annealing conditions, such annealing is also referred to as “hybridization”. When a contiguous sequence of nucleotides of one single-stranded region is able to form a series of “canonical” hydrogen-bonded base pairs with an analogous sequence of nucleotides of the other single-stranded region, such that A is paired with U or T and C is paired with G, the nucleotide sequences are 100% complementary. Besides conventional bases (A, G, C, T), analogs e.g., inosine and 2'-deoxyinosine and their derivatives (e.g. 7'-deaza-2'-deoxyinosine, 2'-deaza-2'-deoxyinosine), azole- (e.g. benzimidazole, indole, 5-fluoroindole) or nitroazole analogues (e.g. 3-nitropyrrol, 5-nitroindol, 5-nitroimidazole, 4-nitropyrazole, 4-nitrobenzimidazole) and their derivatives, acyclic sugar analogues (e.g. those derived from hypoxanthine- or indazole derivatives, 3-nitroimidazole, or imidazole-4,5-dicarboxamide), 5'-triphosphates of universal base analogues (e.g. derived from indole derivatives), isocarbostyryl and other hydrophobic analogues, and any of its derivatives (e.g. methylisocarbostyryl, 7-propynylisocarbostyryl), hydrogen bonding universal base analogues (e.g. pyrrolopyrimidin), and other chemically modified bases (such as diaminopurine, 5-methylcytosine, isoguanine, 5-methyl-isocytosine, K-2'-deoxyribose, P-2'-deoxyribose) can have different base-pairing preferences and can pair with more than one natural nucleobase with similar stringency/probability. In certain cases, the monomers are linked by phosphodiester or by peptidyl linkages or by phosphorothioate linkages.

[0170] The term “single stranded (ss) NA building block”, shall refer to a ss oligonucleotide or polynucleotide of a DNA (herein also referred to as “ssDNA”), which is a linear polymer of nucleotide monomers. Monomers making up oligonucleotides or polynucleotide are capable of specifically binding to a natural polynucleotide by way of a regular pattern of monomer-to-monomer interactions, such as Watson-Crick type of base pairing, base stacking, Hoogsteen or reverse Hoogsteen types of base pairing, wobble base pairing, or the like.

[0171] Ss NA building blocks described herein, which are oligonucleotides, typically range in size between 6 and 26,

but may be longer e.g., between 6 and 220 nucleotides, such as at least 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, or 27, up to 220, 210, 200, 190, 180, 170, 160, 150, 140, 130, 120, 110, 100, 90, 80, 70, 60, 50, 40, 30, 29, 28, 27, or 26 nucleotides. Whenever an oligonucleotide is represented by a sequence of letters (upper or lower case), such as “ATGC,” it will be understood that the nucleotides are in 5'→3' order from left to right and that “A” denotes deoxyadenosine, “T” denotes deoxythymidine, “G” denotes deoxyguanosine, and “C” denotes deoxycytidine. Besides conventional nucleotides (A, G, C, T), modified nucleotides e.g. K-2'-deoxyribose, P-2'-deoxyribose, 2'-deoxyinosine, 2'-deoxyxanthosine or nucleotides with nucleobase analogs may be used e.g., inosine, or 5-methylisocytosine, or 3-nitropyrrole, 5-nitroindole, pyrrolidine, 4-nitroimidazole, 4-nitropyrazole, 4-nitrobenzimidazole, 4-aminobenzimidazole, 5-nitroindazole, 3-nitroimidazole, 5-aminoindole, benzimidazole, 5-fluoroindole, indole, methylisocarbostyryl, pyrrolopyrimidine 7-propynylisocarbostyryl. The terminology and atom numbering conventions follow those disclosed in Strachan and Read, *Human Molecular Genetics 2* (Wiley-Liss, New York, 1999). Usually, an oligonucleotide or polynucleotide comprises the four natural nucleosides (e.g., deoxyadenosine, deoxycytidine, deoxyguanosine, deoxythymidine for DNA or their ribose counterparts for RNA) linked by phosphodiester or by peptidyl linkages or by phosphorothioate linkages; however, they may also comprise non-natural nucleotide analogs e.g., including modified bases, sugars, or internucleosidic linkages.

[0172] Ss NA building blocks, such as oligonucleotides, can be produced using chemical synthesis methods e.g., by synthesizing the oligonucleotide sequence from monomer-phosphoramidites, dimer-phosphoramidites (Neuner, Cortese, and Monaci 1998) or trimer-phosphoramidites (Sondek and Shortle 1992), mixture of monomer-phosphoramidites, mixture of dimer-phosphoramidites, mixture of trimer-phosphoramidites or their combination thereof.

[0173] In some embodiments, NA building blocks are produced and purified from naturally-occurring sources, or synthesized in vivo, within the cell undergoing in vivo mutagenesis using any of a variety of well-known enzymatic methods e.g., as described in Farzadfard et al. (2014). Specifically, enzymes that synthesize soft-randomized NA building blocks include, but are not limited to low fidelity DNA polymerase proteins or low fidelity reverse transcriptase proteins which incorporate mismatching nucleotides during synthesis with high frequency. Alternatively, mismatching nucleotides are incorporated into the NA building blocks with a higher frequency by the DNA polymerases or reverse transcriptases due to the presence of chemical substances, which are well-known to those skilled in the art.

[0174] The term “double stranded (ds) NA building block” as used herein, shall refer to a NA building block which is a linear polymer of nucleotide dimers. Dimers making up oligonucleotides or polynucleotides comprise complementary nucleotides bound by way of a regular pattern of monomer-to-monomer interactions, such as Watson-Crick type of base pairing, base stacking, Hoogsteen or reverse Hoogsteen types of base pairing, wobble base pairing, or the like.

[0175] Double stranded NA building blocks described herein, which are oligonucleotides, typically range in size between 6 and 26 base pairs (bp), but may be longer. dsDNA

oligonucleotides described herein may range in size between 6 and 200 base pairs, e.g., between 6 and 220 base pairs, such as at least 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, or 27, up to 220, 210, 200, 190, 180, 170, 160, 150, 140, 130, 120, 110, 100, 90, 80, 70, 60, 50, 40, 30, 29, 28, 27, or 26 base pairs. Whenever an oligonucleotide or polynucleotide is represented by a sequence of letters (upper or lower case), such as "ATGC," it will be understood that the nucleotides are in 5'→3' order from left to right and that "A" denotes deoxyadenosine, "T" denotes deoxythymine, "G" denotes deoxyguanosine, and "C" denotes deoxycytidine. Besides conventional nucleotides (A, G, C, T), modified nucleotides e.g. K-2'-deoxyribose, P-2'-deoxyribose, 2'-deoxyinosine, 2'-deoxyxanthosine or nucleotides with nucleobase analogs may be used e.g., inosine, or 5-methylisocytosine, or 3-nitropyrrole, 5-nitroindole, pyrrolidine, 4-nitroimidazole, 4-nitropyrazole, 4-nitrobenzimidazole, 4-aminobenzimidazole, 5-nitroindazole, 3-nitroimidazole, 5-aminoindole, benzimidazole, 5-fluoroindole, indole, methylisocarbostryl, pyrrolopyrimidine 7-propynylisocarbostryl. The terminology and atom numbering conventions follow those disclosed in Strachan and Read, *Human Molecular Genetics 2* (Wiley-Liss, New York, 1999). Usually, an oligonucleotide or polynucleotide comprises the four natural nucleosides (e.g. deoxyadenosine, deoxycytidine, deoxyguanosine, deoxythymine for DNA or their ribose counterparts for RNA) linked by phosphodiester or by peptidyl linkages or by phosphorothioate linkages; however, they may also comprise non-natural nucleotide analogs, e.g. including modified bases, sugars, or internucleosidic linkages.

[0176] The simplest end of a double stranded molecule is called a blunt end. In a blunt-ended molecule, both strands terminate in a base pair. Non-blunt ends are created by various overhangs. The term "overhang" as used herein refers to a stretch of unpaired nucleotides at an end of a ds NA building block molecule. These unpaired nucleotides can be in either strand, creating either 3' or 5' overhangs.

[0177] An overhang is specifically characterized by a ss terminal stretch of one or more nucleotides that is reactive to hybridize with a matching ss sequence of a reaction partner (such as another ss or ds NA building block), which ss terminal stretch is extending the double stranded part of the ds NA building block. In particular, the overhang is reactive insofar that it is capable of hybridizing with another ss oligo or overhang that comprises a matching sequence. A reactive overhang is herein also understood to result in a "sticky end" of the molecule.

[0178] The overhang can be part of the leading strand or the lagging strand.

[0179] Where the overhang is at the 5'-end of the ds NA building block and is part of the leading strand, the overhang comprises the 5'-terminal region including the 5'-terminus of the leading strand.

[0180] Where the overhang is at the 3'-end of the ds NA building block and is part of the leading strand, the overhang comprises the 3'-terminal region including the 3'-terminus of the leading strand.

[0181] Where the overhang is at the 5'-end of the ds NA building block and is part of the lagging strand, the overhang comprises the 3'-terminal region including the 3'-terminus of the lagging strand.

[0182] Where the overhang is at the 3'-end of the ds NA building block and is part of the lagging strand, the overhang comprises the 5'-terminal region including the 5'-terminus of the lagging strand.

[0183] Specifically, a ds NA building block may be used which comprises one overhang at one end, and a blunt end at the other end. A blunt end is specifically characterized by a ds terminal stretch of one or more base pairs which includes the terminus of both strands without an overhang at such end of the ds NA building block.

[0184] An overhang may consist of one nucleotide, or can be longer. An overhang may comprise or consist of any one of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 nucleotides, or at least any one of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 nucleotides. An overhang is typically not more than half of a ds NA building block length. For example, if said ds NA building block is a ds oligo that is 6 nucleotides long (in the ds part of the oligo), the overhang is typically not more than 3 nucleotides long, meaning the overhang can also be 1 or 2 nucleotides long. According to a specific example, if said ds oligo is 24 nucleotides long (in the ds part of the oligo), the overhang is typically not more than 12 nucleotides long, meaning it can also be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or 11 nucleotides long.

[0185] The term "derivative" refers to an oligonucleotide or a polynucleotide differing from the original oligonucleotide or polynucleotide, but retaining essential properties thereof. Derivatives may e.g., be produced using a ds polynucleotide (e.g. DNA) as a starting material to engineer single stranded DNA, or complementary RNA molecule, to introduce one or more point mutations, or to bind heterologous moieties or tags by chemical and/or enzymatic means.

[0186] Generally, derivatives are overall closely similar, and, in many regions, identical to the original oligonucleotide or polynucleotide. As a practical matter, whether any particular nucleic acid molecule or polypeptide is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to a nucleotide sequence of the present invention can be determined conventionally using known computer programs. A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. (1990) 6:237-245). In a sequence alignment the query and subject sequences are both DNA sequences. An RNA sequence can be compared by converting U's to T's. The result of said global sequence alignment is in percent identity. If the subject sequence is shorter than the query sequence because of 5' or 3' deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for 5' and 3' truncations of the subject sequence when calculating percent identity. For example, a 90 base subject sequence is aligned to a 100 base query sequence to determine percent identity. The deletions occur at the 5' end of the subject sequence and therefore, the FASTDB alignment does not show a matched/alignment of the first 10 bases at 5' end. The 10 impaired bases represent 10% of the sequence (number of bases at the 5' and 3' ends not matched/total number of bases in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 bases were perfectly matched the final percent identity would be 90%.

In another example, a 90 base subject sequence is compared with a 100 base query sequence. This time the deletions are internal deletions so that there are no bases on the 5' or 3' of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only bases 5' and 3' of the subject sequence which are not matched/aligned with the query sequence are manually corrected for.

[0187] The terms “hybridize,” “hybridization,” “hybridizing,” “anneal,” and “annealing,” as used herein, generally refer to a reaction in which one or more polynucleotides react to form a complex that is stabilized via hydrogen bonding between the bases of the nucleotide residues. The hydrogen bonding may occur by Watson Crick base pairing, Hoogsteen binding, or in any other sequence specific manner. The complex may comprise two strands forming a duplex structure, three or more strands forming a multi stranded complex, a single self-hybridizing strand, or any combination of these. A hybridization reaction may constitute a step in a more extensive process, such as the initiation of a PCR, or the enzymatic cleavage of a polynucleotide by a ribozyme.

[0188] The term “intermediate polynucleotide” as used herein, refers to a product of the assembly of two NA building blocks. For example, a pair of matching NA building blocks is transferred into a reaction containment and the matching NA building blocks are assembled thereby forming a new NA building block that is double stranded and considered as an intermediate because used in a further assembly step. Specifically, said ds NA building block may comprise at least one overhang. Such overhang allows further assembly with another matching NA building block in the direction of the overhang. Where an intermediate comprises two overhangs i.e., on both of its ends, such intermediate is designed to assemble with further NA building blocks in both directions.

[0189] As used herein, the term “ligation” is intended to mean the process during which two nucleic acid sequences anneal to one another and/or assemble with intermolecular chemical bonds (e.g., hydrogen bonds) so as to form a double strand under appropriate conditions.

[0190] Ligation products, herein also referred to as reaction products, can be formed from both double stranded nucleic acids and single stranded nucleic acids. Double-stranded nucleic acids can be ligated by “sticky end” ligation or “blunt end” ligation. In sticky end ligation, staggered ends comprising terminal overhangs can hybridize to a ligation partner. In blunt end ligation, terminal overhangs are not present and successful ligation depends on transient associations of 5'-ends and 3'-ends. Blunt end ligations in general are less efficient than sticky end ligations, and various optimizations, such as adjusting concentrations, incubation times, and temperatures, can be applied to improve efficiencies. Single-stranded polynucleotides can also be ligated.

[0191] The ligation efficiency between two complementary sequences or sufficiently complementary sequences depends on the operating conditions that are used, and in particular the stringency. The stringency may be understood to denote the degree of homology; the higher the stringency, the higher percent homology between the sequences. The stringency may be defined in particular by the base composition of the two nucleic sequences, and/or by the degree of mismatching between these two nucleic sequences. By varying the conditions e.g., salt concentration and tempera-

ture, a given nucleic acid sequence may be allowed to ligate only with its exact complement (high stringency) or with any somewhat related sequences (low stringency). Increasing the temperature or decreasing the salt concentration may tend to increase the selectivity of a ligation reaction.

[0192] The ligation reaction can be performed by hybridizing nucleic acid molecules which are sufficiently complementary to each other.

[0193] The ligation reaction can be performed by an enzyme, specifically a DNA ligase enzyme. A DNA ligase catalyzes the formation of covalent phosphodiester linkages, which permanently join the nucleotides together. In addition, T4 DNA ligase can also ligate ssDNA if no dsDNA templates are present, although this is generally a slow reaction. Non-limiting examples of enzymes that can be used for ligation reactions are ATP-dependent double-stranded polynucleotide ligases, NAD⁺ dependent DNA or RNA ligases, and single-strand polynucleotide ligases. Non-limiting examples of ligases are *Escherichia coli* DNA ligase, *Thermus filiformis* DNA ligase, *Thermus thermophilus* DNA ligase, *Thermus scotoductus* DNA ligase (I and II), CircLi-gase™ (Epicentre; Madison, WI), T3 DNA ligase, T4 DNA ligase, T4 RNA ligase, T7 DNA ligase, Taq ligase, Ampli-gase (Epicentre® Technologies Corp.), VanC-type ligase, 9° N DNA Ligase, Tsp DNA ligase, DNA ligase I, DNA ligase III, DNA ligase IV, Sso7-T3 DNA ligase, Sso7-T4 DNA ligase, Sso7-T7 DNA ligase, Sso7-Taq DNA ligase, Sso7-*E. coli* DNA ligase, Sso7-Amp ligase DNA ligase, and thermostable ligases. Ligase enzymes may be wild-type, mutant isoforms, and genetically engineered variants. Ligation reactions can contain a buffer component, small molecule ligation enhancers, and other reaction components.

[0194] Specifically, a T4 DNA ligase can be used in a ligation reaction. In the method provided herein, the ligation reaction can be performed under high-fidelity conditions that block side reactions and minimize mismatches.

[0195] Assembly of NA building blocks into intermediate polynucleotides or into the target polynucleotide may be carried out using suitable ligation buffer solutions. The ligation buffer solution is e.g., an aqueous solution, typically in a nuclease-free environment, at a pH that ensures the selected ligase will be active; typically, this is a pH of between about 7-9. Preferably, the pH is maintained by Tris-HCl at a concentration of between about 5 mM to 50 mM. The ligation buffer solution may include one or more nuclease inhibitors, usually calcium ion chelators, such as EDTA. Typically, EDTA is included at a concentration of between about 0.1 to 10 mM. The ligation buffer solution includes whatever cofactors are required for the selected ligase to be active. Usually, this is a divalent magnesium ion at a concentration of between about 0.2 mM to 20 mM, typically provided as a chloride salt. For T4 DNA ligase ATP is required as a cofactor. The ligase buffer solution may also include a reducing agent, such as dithiothreitol (DTT) or dithioerythritol (DTE), typically at a concentration of between about 0.1 mM to about 10 mM. Optionally, the ligase buffer may contain agents to reduce nonspecific binding of the oligonucleotides and polynucleotides. Exemplary agents include salmon sperm DNA, herring sperm DNA, serum albumin, Denhardt's solution, and the like. Preferably, ligation conditions are adjusted so that ligation will occur if matching NA building blocks form perfectly matched duplexes with the bases of a contiguous complementary region. However, it is understood that it may be

advantageous to permit non-pairing nucleotides on the 5'-end of one NA building block of a matching pair of NA building blocks and the 3'-end of the other one of the matching pair of NA building blocks to aid in detection or to reduce blunt-end ligation. Important parameters in the ligation reaction include temperature, salt concentration, presence or absence and concentration of denaturants such as formamide, concentration of the pair of NA building blocks and type of ligase employed. Methods of selecting hybridization conditions for the reaction are known to those skilled in the art.

[0196] Preferably, ligation is performed under stringent hybridization conditions to ensure that only matching NA building blocks hybridize. Typically, stringency is controlled by adjusting the temperature at which hybridization occurs while holding salt concentration at some constant value e.g., at about (+/-20%) 100 mM NaCl, or the equivalent salt concentration of other salts. Other factors can be relevant, such as the particular sequence or length of the matching pair of NA building blocks, and/or the heat lability of the selected ligase. Preferably, the ligation reaction is carried out at a temperature close to the melting temperature of hybridized NA building blocks in the ligation buffer. More preferably, the ligation reaction is carried out at a temperature within 10° C. below the melting temperature of hybridized NA building blocks in the ligation buffer solution. Most preferably, the ligation reaction is carried out at a temperature in the range of 0 to 5° C. below the melting temperature of hybridized NA building blocks in the ligation buffer solution.

[0197] Ligation may be followed by one or more amplification reactions. In some embodiments, the ligation products, or target polynucleotides are isolated or enriched prior to amplification.

[0198] Isolation can be achieved by various suitable purification methods including affinity purification and gel electrophoresis. For example, ligation products, or target polynucleotides can be isolated by binding of a selective binding agent immobilized on a support to a tag attached to the capture probe. The support can then be used to separate or isolate the capture probe and any polynucleotide hybridized to the capture probe from the other contents of the sample reaction volume. The isolated polynucleotides can then be used for amplification and further sample preparation steps. In some embodiments, the capture probe is degraded or selectively removed prior to amplification of the circular target polynucleotides. Amplification of reaction products, or target polynucleotides can be achieved by various suitable amplification methods known to those skilled in the art.

[0199] Reaction and storage compartments may be conveniently provided within one or more parts of a device, which together are provided as "array device". Such array device may be any one or more of a microtiter plate, microfluidic microplate, set of capillaries, microarray or a biochip, preferably a DNA or RNA biochip.

[0200] A microarray is herein understood as a supporting material (such as a glass or plastic slide) onto which numerous molecules or fragments such as DNA or NA building blocks are attached in a regular pattern. More specifically, it refers to microscope slides that are printed with thousands of tiny spots in defined positions, wherein said spots are capable of binding DNA. Such slides are often also referred to as biochips, DNA chips, or gene chips. Microarrays can bind DNA in a covalent or non-covalent manner and can

thus serve as array devices in which NA building blocks are stored in pre-defined locations, i.e., spots.

[0201] In a specific example, said separate library containments are micro-well plates, arranged as stacked plates, optionally barcode labelled, and accessible by an automated microdroplet handler. Library members may be conveniently stored in said stacked micro-well plates, wherein the order and stacking is according to decreasing frequency of use. One example is to store into micro-well plates where the first plate comprises the most common pair combinations of NA building blocks, in decreasing order until the last micro-well plate which contains the least-frequently used library members.

[0202] According to a specific aspect, a microfluidics chip can be used in order to pipe droplets containing NA building blocks in such a way as to implement the assembly workflow. The design of the chip can be that of a symmetric workflow, but by leaving certain channels free of the NA building blocks, an asymmetric workflow is enabled. Similarly, digital microfluidics e.g., platforms of electro-wetting on dielectric (EWOD) can also be used to have accurate control on the movement of individual droplets, thereby facilitating the implementation of complex hierarchical workflows.

[0203] According to a specific aspect, a microtiter plate can be used, which term is understood to refer to well plates, multi-well plates or micro-well plates. These plates are commonly manufactured in a 2:3 rectangular mix with 96, 384, or 1536 wells, although other cavity configurations are available. Some of the other sizes, far less common, available are 6, 24, 3456, and 9600 wells. The wells of the microplate typically hold between tens of nanoliters to several milliliters of liquid.

[0204] Capillaries can be any glass capillaries, microfluidic capillaries and autonomous microfluidic capillary systems. Capillary microfluidics are important tools in many different fields. Due to their axisymmetric flow and ability to withstand organic solvents, when compared with their lithographically fabricated polydimethylsiloxane (PDMS) counterparts, glass capillary devices possess advantages for microfluidic applications. In particular, a circular tube is inserted into a square outer flow channel, which greatly simplifies alignment and centering of these devices. These devices can produce small and large droplets, ranging from 10 to multiple hundreds in μm size.

[0205] NA building blocks may be conveniently transferred from an array device to reaction containments by automated means e.g., either robotically or via dedicated fluids using, for example, an automated liquid handler, from such compartments into other compartments herein referred to as reaction compartments, i.e. from one vessel to another. In order to facilitate time efficient assembly of polynucleotides, hierarchies of reactions and respective vessels may be employed corresponding to frequency of use of NA building block library members. The transfer to a new vessel involves the physical movement of a device that picks one or more molecules of a NA building block from the respective location, or the pneumatic/hydraulic deposition through microfluidics. Due to the large number of NA building blocks required to theoretically build any given sequence, most spatial distributions would incur into wasted time and resources due to scanning and lengthy travel times of the liquid handler. However, by using a specific distribution of

the library members or reaction compartments, it can be ensured that there is minimal movement according to a target sequence.

[0206] According to a specific aspect, transfer of NA building blocks described herein, such as to transfer from one containment to another, e.g., transfer from a library containment to a reaction containment, is done using an automated system e.g., using a liquid handler such as further described herein.

[0207] As used herein the terms “liquid handler”, “automated handler” or “microdroplet handler” refer to any device used in a method of liquid handling, preferably, automated liquid handling, preferably a device as used in sensor-integrated robotic systems. As low-volume dispensing becomes increasingly common in life science, microsyringes have emerged which have a high level of precision with hermetic seals. Some manual or electronic holders are designed to precisely control the piston displacement to ensure the accuracy of the dispensed volume. Besides the syringe, a pipette is another popular tool for liquid handling. The dispensed volume can be at the micro- or sub-microliter level. Multichannel pipettes are recommended for multi-outlet pipetting at one time. There are both fixed- and adjustable-volume pipettes on the market. The former is more accurate and precise, whereas the latter has a larger scope of applications because the operator can choose different volumes according to need. Besides, high throughput has become important in life science research. One of the representative applications is microarray printing. This technology creates an array of biosample spots each at the nanoliter scale to enable the analysis of large numbers of experiments in parallel with only tiny quantities of samples. The process of spotting thousands of biosamples is almost an impossible task with a handheld dispensing tool, making robotic liquid handling an important aspect.

[0208] Robotic workstations have multiple advantages over manual liquid handling since robots can work without fatigue, increase the throughput, perform consistently, and ensure accuracy and precision. According to the requirements for the platform with integration and multifunction, there are still more complex systems in which the liquid-handling task is only one part of the function. The generic architecture of liquid-handling may be built up as follows. First, the control center controls a robot that moves between the dispensing part and the washing station of the robotic workstation. The washing station is used to clean the dispensing head for lengthening its life and for ensuring the safety of the sample. Liquid samples are expelled from the dispensing head and deposited on the substrates for further processing. Sensors are incorporated to monitor the status of the dispensing part such that feedback control can be performed by the control center. Sensors are not always installed on all the workstations but are more and more used to construct the feedback loop for delivering a better performance.

[0209] Specifically, a liquid handler can be used in a method described herein, in particular a microdroplet handler. Specifically, the liquid handler is automated. Using a liquid handler, a suitable volume of at least any one of 10, 20, 30, 40, 50, 60, 70, 80, 90 100, 200 or 500 nL can be transferred. Typically, at least any one of 10^9 , 10^{10} , 10^{11} or 10^{12} copies of a library member e.g., NA building blocks sourced from one library containment, or matching pairs of NA building blocks sourced from separate library contain-

ments, are placed into one reaction containment. For example, at least about 10^{11} copies (e.g., 6.06×10^{11} copies) of specific NA building blocks are placed into one reaction containment to react with each other. Typically, the volume in which NA building blocks are transferred by a liquid handler is between 10 and 1000 nL. Preferably it is between 10 and 500 nL, and more preferably it is between 50 and 250 nL.

[0210] “Microfluidic devices” typically enable the manipulation of discrete fluid packets in the form of microdroplets that provide numerous benefits for conducting biological and chemical assays. Among these benefits are a large reduction in the volume of reagent required for assays, the size of sample required, and the size of the equipment itself. Such technology also enhances the speed of biological and chemical assays by reducing the volumes over which processes such as heating, diffusion, and convective mixing occur. Once the droplets are generated, carefully designed droplet operations allow for the multiplexing of a large number of droplets to enable large-scale complex biological and chemical assays.

[0211] The term “microfluidic microplate” as used herein, refers to a combination of microfluidic technology with standard SBS-configured 96-well microplate architecture, in the form of microfluidic microplate technology. A microfluidic microplate allows for the improvement of essential workflows, conservation of samples and reagents, improved reaction kinetics, and the ability to improve the sensitivity of the assay by multiple analyte loading.

[0212] The term “library” as used herein shall refer to a collection of library members which are NA building blocks (e.g., an oligonucleotide or polynucleotide library). Library members can be ss NA building blocks or ds NA building blocks. A library typically contains library members which are diverse. The library described herein comprises library members suitably composed of NA building blocks of varying lengths and different sequences.

[0213] According to a specific example, the library is a focused library which is designed to assemble a pre-defined target polynucleotide e.g., according to a template or comprising a SOI, wherein the library members are NA building blocks which correspond to a certain region of the target polynucleotide, template or SOI. According to a specific further example, a library may be larger to comprise a diversity that allows assembly of the library members to produce a variety of ds target polynucleotides. Specifically, the diversity may even cover the entire genetic space of a species or organism. For example, a library may comprise a diversity of NA building blocks as necessary to potentially synthesize any and all naturally occurring polynucleotides of the human chromosomal genome or mitochondrial genome by assembly of the respective NA building blocks. In a further example, said diversity may cover any and all naturally occurring polynucleotides of eukaryotic species other than human, such as e.g., mouse, rat, rabbit, pig, sheep, plants, fungi or yeast. In yet another example, said diversity may cover any and all naturally occurring polynucleotides of prokaryotes, such as e.g., archaea or bacteria.

[0214] The library provided herein, specifically comprises pairs of matching NA building blocks, each consisting of two ss NA building blocks, comprising partially or fully complementary sequences. Said pair(s) of matching NA building blocks may be present in the library, wherein each of said two ss NA building blocks are comprised in separate

containments, or wherein both of said two ss NA building blocks are comprised in the same containments where they may anneal and form a ds NA building block. The nucleotide sequences of a pair of matching ss NA building blocks may be complementary in at least 1, 2 or 3 nucleotides, preferably at least 4 or more nucleotides, such that a matching pair can form a new ds polynucleotide molecule by hybridization of the ss NA building block sequences, preferably wherein the ss NA building blocks hybridize in part, thereby obtaining a ds polynucleotide with an overhang.

[0215] The library preferably comprises NA building blocks which are artificially or chemically synthesized, or chemically modified (e.g., including peptidyl nucleic acids or phosphorothioate bond) NA building blocks which are synthesized by suitable methods well-known in the art. The NA building blocks comprised in the library can also be generated by enzymatic digestion of naturally occurring DNAs.

[0216] The library described herein may comprise thousands of NA building blocks as library members. Specifically, the library described herein comprises a diversity of library members, wherein each of the library members has a different nucleotide sequence.

[0217] The library described herein may specifically comprise a diversity of ds NA building block library members, wherein each of the ds NA building block library members has a different nucleotide sequence. Specifically, said diversity covers at least 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 150, 200, 250, 300, 350, 400, 450, 500, 750, 1,000, 2,000, 3,000, 4,000, 5,000, 10,000, 20,000, 40,000, 60,000, 80,000, 100,000, 120,000, 140,000, 160,000, 180,000 or 200,000 different ds NA building blocks or library members.

[0218] The library described herein may specifically comprise a diversity of ss NA building block library members, wherein each of the ss NA building block library members has a different nucleotide sequence. Specifically, said diversity covers at least 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 150, 200, 250, 300, 350, 400, 450, 500, 750, 1,000, 2,000, 3,000, 4,000, 5,000, 10,000, 20,000, 40,000, 60,000, 80,000, 100,000, 120,000, 140,000, 160,000, 180,000 or 200,000 different ss NA building blocks. Specifically, ss NA building blocks may be used as linkers, such as suitably used in the assembly of two ds NA building blocks to facilitate linkage.

[0219] Specifically, said diversity means, that different library members differ in at least one base or base pair. One library member may actually encompass multiple copies of the respective ss or ds NA building blocks, which copies consist of the same sequence. Such multiple copies of a library member are specifically contained in only one library containment.

[0220] In a specific embodiment described herein, said diversity covers NA building blocks which are modified, preferably phosphorylated.

[0221] The library described herein may comprise library members which are phosphorylated, methylated, biotinylated or which are linked to fluorophores or quenchers. As

described herein, library members may comprise one or more additional phosphoryl groups.

[0222] Methylation of library members comprises the addition of a methyl group to a DNA molecule, preferably to cysteine or adenine, and is performed according to suitable DNA methylation methods well-known in the art.

[0223] As used herein, biotinylation refers to a method of covalently attaching one or more biotin molecules to a nucleic acid, such as ss or ds NA building blocks. The library members described herein may be biotinylated by suitable methods well-known in the art; preferably it is a method of chemical biotinylation. NA building blocks can be readily biotinylated in the course of synthesis by phosphoramidite methods well-known in the art, which use biotin phosphoramidite.

[0224] Library members described herein may be conjugated to a fluorophore by suitable chemical and enzymatic methods well-known in the art. Exemplary methods used for the fluorescent labeling of nucleic acids may employ a method for enzymatic labeling of DNA with fluorescent dyes e.g., using a Thermo Fisher's ARES DNA labeling kit, which employ a two-step method for enzymatic labeling of DNA with fluorescent dyes. Further exemplary methods may employ a chemical method for labeling nucleic acids without enzymatic incorporation of labeled nucleotides e.g., using a ULYSIS Nucleic Acid Labeling Kit. Further exemplary methods may employ chemical labeling of amine-terminated oligonucleotides to prepare singly labeled fluorescent oligonucleotide conjugates e.g., using an Alexa Fluor Oligonucleotide Amine Labeling Kit. Further exemplary methods may employ DNA arrays/microarrays and other hybridization techniques.

[0225] Library members may be linked to one or more quenchers, e.g., substances that absorb excitation energy from a fluorophore, by suitable methods well-known in the art. Examples of quenchers include but are not limited to Dabsyl (dimethylaminoazobenzenesulfonic acid), Black Hole Quenchers, Qxl quenchers, Iowa black FQ, Iowa black RQ and IRDye QC-1.

[0226] The library described herein specifically comprises or consists of NA building blocks which are preferably purified, may comprise modifications and are preferably kept at a standard concentration and volume in an appropriate buffer, solvent and/or excipient, e.g., ready-to-use for assembling, or be provided in a storage-stable form, such as in the dry state.

[0227] Specifically, the library members are contained in the library in a storage stable form, such as in solution or in a dry state.

[0228] Specifically, any of the following buffer and/or excipients may be used to keep the NA building blocks in solution: Tris Buffer, T.E. Buffer (Tris-EDTA Buffer) or Nuclease Free Water. Specifically, library members may be kept in Tris Buffer, wherein said Tris Buffer is provided at a concentration of about 10 mM (+/-1 mM or 2 mM). Specifically, library members may be kept in T.E. Buffer. Specifically, said T.E. Buffer is at least composed of Tris, at a concentration of about 10 mM (+/-1 mM or 2 mM), and EDTA, at a concentration of any one of 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9 or 1.0 mM. Specifically, Nuclease Free Water, is water which has been de-ionized, filtered and autoclaved and is essentially free of contaminating non-specific endonuclease, exonuclease and RNase activity.

[0229] Specifically, any of the following solvents may be used to provide a solution of NA building blocks: ethanol, methanol, propanol, formamide, pyridine, or dimethyl sulfoxide, acetonitrile, dimethylformamide, formamide, tetrahydrofuran, MDSO, DMF, glycerol or ionic solvents, or mixtures of any of the foregoing e.g., in varying concentrations.

[0230] Specifically, all library members are kept in a compartmented array device.

[0231] In a specific embodiment, a library of NA building blocks is provided within an array device, wherein library members are contained in separate library containments.

[0232] Specifically, said array device is any of a microtiter plate, a microfluidic microplate, a set of capillaries, a microarray or a biochip, preferably a DNA and/or RNA biochip. Said array device may comprise only one, all or any type of the aforementioned containments.

[0233] Specifically, more than one different library members may be contained in only one library containment.

[0234] Specifically, said different library members contained in one library containment can be a mixture of ss NA building blocks of such sequences that are not capable of annealing to each other, thus, the ss NA building blocks are contained in the mixture as ss molecules. Specifically, said different library members contained in one library containment can be a mixture of ds and ss NA building blocks of such sequences that are not capable of assembly with or ligating to any of the other ds and ss NA building blocks contained in the mixture. Specifically, said different library members contained in one library containment can be ss and ds NA building blocks of such sequences that are not capable of assembly or annealing to one another.

[0235] The term “library diversity” or “diversity” as used herein, refers to a degree of versatility characterizing the library provided herein. Specifically, said diversity comprises aa and/or ds NA building blocks of different lengths and different sequences.

[0236] Specifically, the diversity of a NA building block library means that a variety of library members differ in at least one base or base pair. One library member may actually encompass multiple copies of ss or ds NA building blocks of the same sequence. Such multiple copies of a library member are specifically contained in only one library containment.

[0237] For example, the library may comprise all possible sequence variations of 8 nucleobase long ss oligos (herein referred to as octamers), which are 65,536 different ss oligos of 8 nucleobases length, and in addition, further ss oligos or ds oligos of different lengths, which are commonly comprised in target sequences and are, thus, used more often to build any pre-defined sequence. Including commonly used ss or ds NA building blocks into a library’s diversity decreases synthesis cost and increases time efficiency.

[0238] Specifically, said diversity may cover an entire genome, for example the human genome. Specifically, said diversity may cover the entire genetic space. Specifically, said diversity may cover a genome or the entire genetic space multiple times in multiple ways. For example, the diversity may encompass all possible hexamer, heptamer and/or octamer sequence combinations, or even encompass all or selected 9-mers, 10-mers and/or up to 26-mers.

[0239] Each library member may be individually characterized and marked by a selectable marker or a DNA sequence tag or barcode, to facilitate the selection of a

library member in the library or the identification of a library member in the library. Alternatively, the genetic mutation may be determined directly by a suitable determination method e.g., high-throughput sequencing, capillary sequencing or employing specific probes hybridizing with a pre-defined sequence, to select the corresponding oligonucleotide.

[0240] It may be desirable to place each of the diverse library members in separate containers, to obtain a library of NA building blocks in separate containers. According to a specific embodiment, the library is provided in an array e.g., a DNA biochip, wherein the array comprises a series of containments or spots on a solid carrier.

[0241] The term “sequence of interest” or “SOI” as used herein refers to the desired nucleotide or base pair sequence of the ds target polynucleotide which is to be produced by the method provided herein.

[0242] The term “solid support” as used herein is understood as a solid material often used in oligo or polynucleotide synthesis because it allows easy washing and cleanup of a NA reaction mixture from other chemicals, enzymes, unreacted oligos, etc. By keeping a NA assembly attached to a substrate, there is more control on the assembly. However, the yield tends to be lower than in solutions posing other challenges. Nevertheless, use of solid-support attachment for enzymatic and/or chemical assembly may be useful e.g., for purification or amplification steps. In general, the purpose for using a solid support-attached oligo or polynucleotide is to capture something, hold on to it, enrich it, and identify and/or purify it.

[0243] Examples of common solid supports include microarrays or beads. The most generally accessible approach for producing NA building block microarrays is to synthesize the individual NA building blocks separately prior to immobilization on the solid surface. In this case, the NA building block can be modified with a functional group that allows attachment to a reactive group on a solid surface. Another important class is the micro-bead-attached NA building block. Some modifications include: amine-NA building blocks covalently linked to an activated carboxylate group or succinimidyl ester, SH-NA building blocks covalently linked via an alkylating reagent such as an iodoacetamide or maleimide, acrydite-NA building blocks covalently linked through a thioether, and biotin-NA building blocks captured by immobilized Streptavidin.

[0244] Specific methods include photo-cleavable chemical moieties that allow detachment from a solid support by irradiation. WO2019140353A1 discloses a method where oligos are attached through an anchor oligonucleotide that contains an uracil base. By treating with uracil-N-deglycosylase (UNG) abasic nucleotide is left in place of the uracil. Then, treating the sample with either Endo IV or APE-I, which are endonucleases capable of cutting at a single stranded abasic site, the fragments are released from the surface.

[0245] The term “target double stranded (ds) polynucleotide” refers to a polynucleotide having a predefined sequence, which is produced by the method provided herein. Specifically, said target ds polynucleotide is characterized by a sequence which is identical and/or corresponding to a SOI. If the target ds polynucleotide sequence has a sequence which is less than 100% identical to a SOI, the target ds polynucleotide is understood as a proxy ds polynucleotide

that can be further modified to produce a ds polynucleotide that has a sequence which is identical and/or corresponding to the SOI.

[0246] The foregoing description will be more fully understood with reference to the following examples. Such examples are, however, merely representative of methods of practicing one or more embodiments of the present invention and should not be read as limiting the scope of invention.

EXAMPLES

[0247] The methods disclosed herein refer to a hierarchical assembly, albeit not a symmetric one. Instead of reacting all oligos or reaction products at every tier of the assembly, only subsets of these are assembled at any given tier, with the remaining ones assembled at later tiers, achieving an asymmetric assembly. In such a way, ambiguous assemblies are avoided, thereby successfully completing the desired product which cannot be otherwise completed with a symmetric assembly (FIG. 3).

[0248] For example, to assemble oligos a,b,c and d into the construct abcd, a symmetrical assembly workflow would be: in a first tier oligos a+b and c+d are reacted in two independent compartments to give first reaction products ab and cd. In a second reaction these two products are reacted in a compartment to yield the desired abcd polynucleotide. This assembly workflow can be represented as a binary tree in bracket notation $\{\{a,b\},\{c,d\}\}$.

[0249] According to a specific example, a and b are assembled by ligation with sticky-ends and the sticky ends permit not only ab as a reaction product but also ba as a second (undesired) reaction product (FIG. 3). Necessarily, in this particular case, other reaction products such as $(ab)_n$, $(ba)_n$, would also form, impeding a successful assembly at the second tier.

[0250] This problem can be circumvented using two convoluted principles:

- [0251] (i) a method of partitioning the sequence into sub-sequences, and
- [0252] (ii) a method for determining an assembly workflow.

[0253] Partitioning is the process of determining a set of sub-sequences of oligonucleotides or polynucleotides in such a way that when processed with an appropriate assembly workflow they result in an assembled polynucleotide with the sequence of interest.

[0254] The assembly workflow is the series of hierarchical steps that are carried out to achieve that assembly. For instance, the assembly workflow above $\{\{a,b\},\{c,d\}\}$ is the simplest hierarchical workflow possible. But others can be more efficient, depending on the sequence.

[0255] According to this example, the method that is disclosed herein comprises three aspects:

- [0256] (i) a partitioning of the sequence, preferably using an algorithm, in a way that it eliminates as many non-specific assembly steps as possible;
- [0257] (ii) a sequence-specific assembly workflow that ensures that no ambiguities remain and
- [0258] (iii) the actual enzymatic or chemical assembly of the oligo/polynucleotides starting from the sequences defined in (i) following the steps defined in (ii) into the target sequence.

[0259] More specifically, the partitioning of a target sequence into smaller polynucleotides or oligonucleotides

can be done in a way where these building blocks have different lengths, but equal overhang length, or where they have different overhang lengths, and then, when assembled in non-trivial assembly workflows that are sequence-dependent and non-symmetric (or asymmetric), as explained below, a successful synthesis of a target polynucleotide is achieved in a way that cannot be done with naïve partitions that do not consider the mis-assembly risk and/or hierarchical but sequence-independent symmetric workflows.

[0260] In the following subsections the partitioning and the assembly workflows are described in more detail.

A. Asymmetric Assembly Workflows.

[0261] In some cases, for instance, if the components (oligos) are derived from restriction digests, there may not be ways to redesign the oligos as indicated above. Thus, one cannot rely on oligo design alone. To overcome this problem, the way in which the oligos are assembled and an asymmetric assembly strategy was developed.

[0262] In the strategy of asymmetric assembly trees, in a first reaction b and c can be assembled into bc, then this reaction product can be reacted with, d to give bcd and only after, in a third tier react the last product with a to obtain abcd (FIG. 3b). This assembly can be represented by a binary tree $\{a,\{\{b,c\},d\}\}$. Note that also other alternative assemblies could potentially work, such as $\{\{a,\{b,c\}\},d\}$ or $\{\{a,\{b,\{c,d\}\}\}$ (FIG. 3c). However, the number of possible assembly trees for a given number of starting oligos N is combinatorically big, thus, the problem arises of deciding which of these is sufficient in order to successfully complete an assembly. For longer target sequences it is thus beneficial to use a computational method as a first step.

[0263] First, a measure for the uniqueness (or not uniqueness) of the assembly of two oligos is assigned, in order to verify their validity in the assembly (examples given below). This measure can be assigned to every node in the tree and, then, all nodes' measures convolved by, for example, multiplying the weights at each node, to give a single measure to the complete workflow in a way that correlates with the effectiveness of the actual assembly and/or to the success probability of the assembly (FIG. 4).

[0264] By using a measure with these properties, it is then possible to identify an optimal, or nearly so, assembly workflow by performing some search in the space of all potential trees with an optimization method or selection criteria.

[0265] Once an optimal or nearly optimal workflow has been identified, this workflow can be employed in order to complete the assembly e.g., by means of liquid handlers or microfluidic devices. Considering that there are many possible assembly workflows, the measures in question can consider also as a second criterion such as minimizing the number of tiers or to have the most symmetric assembly possible that does not lead to failure.

B. Modification of Oligo Length with Equal Oligo Size

[0266] Where there are oligos that have self-complementary overhangs or that are able to self-ligate (e.g., with overhangs like ATAT, TATA, CGCG, GCGC or variations that allow complementarity with wobble pairs, Hoogsteen pairs or by any other criterion that indicates that the oligos can self-ligate), then the partition is deemed invalid. In this case, the problematic oligonucleotide can be re-designed together with the immediate pair so that they are slightly

larger or shorter, until the ambiguity is resolved. For example, consider the two oligo dimers:

Oligo a	Oligo b
...TCGAGGAACGC	<u>ATATCGGTAA</u> ...
...AGCTCCTTGCGT <u>TATA</u>	GCGATTT...

[0267] Oligo sequences referred to above, regions that are complementary to each other are underlined:

Oligo a, leading strand:
(SEQ ID NO: 61)
TCGAGGAACGC

Oligo a, lagging strand:
(SEQ ID NO: 62)
ATATGCGTTCCTCGA

Oligo b, leading strand:
(SEQ ID NO: 63)
ATATCGGTAA

Oligo b, lagging strand:
(SEQ ID NO: 64)
TTTAGCG

[0268] Both, Oligo a and Oligo B will be self-ligating because the overhang is ambiguous. The oligo design can be modified by transferring the 3'C and 5'A of Oligo a to the 5' and 3' ends, respectively, of Oligo b.

Oligo a'	Oligo b'
...TCGAGGAACG	<u>CATATCGGTAA</u> ...
...AGCTCCTTGCGT <u>TAT</u>	<u>AGCGATTT</u> ...

[0269] Oligo sequences referred to above, regions that are complementary to each other are underlined, transferred nucleosides are bold and underlined:

Oligo a', leading strand:
(SEQ ID NO: 65)
TCGAGGAACG

Oligo a', lagging strand:
(SEQ ID NO: 66)
TATGCGTTCCTCGA

Oligo b', leading strand:
(SEQ ID NO: 67)
CATATCGGTAA

Oligo b', lagging strand:
(SEQ ID NO: 68)
TTTAGCGA

[0270] Now, Oligo a' is one base shorter than Oligo a and Oligo b' is one base longer than Oligo b, which is an asymmetric pair of oligos, and, importantly, they do not self-ligate anymore.

[0271] This strategy can be employed to find more efficient partitions even if the oligos are not self-ligating.

C. Modification of Overhang Length

[0272] The problem of self-ligation or any other ambiguity in the ligation can be solved also by a modification of the

length of the overhang. Considering the same example as above, namely the two self-ligating oligos that appear contiguously in the sequence:

Oligo a	Oligo b
...TCGAGGAACGC	<u>ATATCGGTAA</u> ...
...AGCTCCTTGCGT <u>TATA</u>	GCGATTT...

[0273] Oligo sequences referred to above, regions that are complementary to each other are underlined:

Oligo a, leading strand:
(SEQ ID NO: 61)
TCGAGGAACGC

Oligo a, lagging strand:
(SEQ ID NO: 62)
ATATGCGTTCCTCGA

Oligo b, leading strand:
(SEQ ID NO: 63)
ATATCGGTAA

Oligo b, lagging strand:
(SEQ ID NO: 64)
TTTAGCG

[0274] can be partitioned in a different manner so that one base of one oligo sequence in the naïve partition is instead included in the other oligo, as for example:

Oligo a''	Oligo b''
...TCGAGGAACG <u>A</u>	<u>TATCGGTAA</u> ...
...AGCTCCTTGCGT <u>TATA</u>	GCGATTT...

[0275] Oligo sequences referred to above, regions that are complementary to each other are underlined, transferred nucleosides are bold and underlined:

Oligo a'', leading strand:
(SEQ ID NO: 69)
TCGAGGAACG**A**

Oligo a'', lagging strand:
(SEQ ID NO: 62)
ATATGCGTTCCTCGA

Oligo b'', leading strand:
(SEQ ID NO: 70)
TATCGGTAA

Oligo b'', lagging strand:
(SEQ ID NO: 64)
TTTAGCG

[0276] or, even in a manner that increases the oligo overhang length, i.e.

Oligo a'''	Oligo b'''
...TCGAGGAACG	<u>CATATCGGTAA</u> ...
...AGCTCCTTGCGT <u>TATA</u>	GCGATTT...

[0277] Oligo sequences referred to above, regions that are complementary to each other are underlined, transferred nucleosides are bold and underlined:

```
Oligo a''', leading strand:
                               (SEQ ID NO: 71)
TCGAGGAACG
Oligo a''', lagging strand:
                               (SEQ ID NO: 62)
ATATGCGTTCCTCGA
Oligo b''', leading strand:
                               (SEQ ID NO: 72)
CATATCGGTAAA
Oligo b''', lagging strand:
                               (SEQ ID NO: 64)
TTTAGCG
```

D. Computation of the Assembly Workflow.

[0278] A given target Sequence of Interest (SOI) is processed by the algorithm to provide (a) a list of ss oligonucleotide sequences (oligos) and (b) the workflow *W* to assemble those sequences. Note that oligo sequences and *W* may be different depending on the assembly method (see below). Also, depending on the nature of the algorithm that is used a different output can be produced although of similar efficiency of assembly. Different criteria can be chosen to partition the sequence, as for example, computational parameters that consider Watson-Crick and wobble and/or Hoogsteen base pairs, a percentage of mismatch, enzyme kinetic parameters, annealing thermodynamics, indices for ligation/cloning efficiency, physico-chemical, statistical or phenomenological formulas, or algorithmic processes. Note that there are sequences that cannot be partitioned (or assembled), such as $(AT)_n$ $(TA)_n$ $(CG)_n$ $(GC)_n$ and other extreme cases of high regularity.

1. Provision of the Oligonucleotides or Polynucleotide Fragments.

[0279] Oligonucleotides may be provided as a library in array format in micro-well plate, procured from any provider or made using phosphoramidite synthesis, TdT enzymatic or any other method. They could also be directly synthesized in the microfluidic chip ready for assembly. However, it may be needed that these are annealed, if a ligation-based assembly will be used. If a PCR-based assembly is used instead, they may be provided and used directly as single stranded oligos. The oligos or polynucleotides may have any length.

2. Assembly of the Polynucleotides.

[0280] The method can be used to assemble short oligos (e.g., 8 nt) into polynucleotides by ligation with sticky ends, by Gibson Assembly (GA), considering that the different oligos have sufficient overlap, and where at any tier a ligation or GA reaction is incubated by providing the necessary enzymes, or gene fragments assembled by golden gate. Oligos may be attached to solid substrate (e.g., avidin attachment through biotin modifications or any other method) in magnetic beads or coated surfaces, thereby increasing accuracy and facilitating purification steps in between assembly tiers. Implementation of the asymmetric assembly steps can be done by using liquid transfers (by automated handlers or microfluidic devices).

3. Postprocessing of the Polynucleotides.

[0281] This can include purification, amplification by means of high-quality PCR, bacterial cloning, or any other process required to finalize and enrich the assembled molecule that has the SOI.

Example 1. Algorithm for Partitioning the Sequences and Computing the Tree

Choosing the Objective Function

[0282] An objective function (score) was chosen for evaluating a partition by defining a weight to every node *n* according to whether the oligos at each node can react in a unique manner (*w*=1) or not (*w*=0). More specifically, any two oligos have a total of four overhangs (*O*₁, . . . , *O*₄) (see FIG. 4a for numbering of the overhangs) where we want that

[0283] *O*₂ and *O*₃ match according to Watson-Crick pairing

[0284] any other pair of overhangs does not match, also excluding “wobble” or Hoogsteen pair variations.

[0285] Mathematically, this can be represented by a symmetric adjacency matrix *M* (*M*_{*ij*}=*M*_{*ji*}) where a 1 indicates a matching pair and a 0 indicates the pair does not match. The score *w_n* at node *n* is then defined as

$$w = \begin{cases} 0 & \text{if } O_{ij} = 1 \text{ for } i, j \neq 2, 3 \\ 1 & \text{otherwise} \end{cases}$$

[0286] For example, FIG. 4b shows a pair of oligos that match in a unique manner, so only the element *M*_{2,3} whereas FIG. 4c, aside from having the desired matching *O*₁ and *O*₄ also match which would lead to a ligation in an undesired manner. Therefore, the elements

[0287] *M*_{1,4}=1 and therefore *w*=0.

[0288] As indicated above, other metrics can be incorporated based, for example, in relation to ligation activity, on frequency of recovered clones from a controlled experiment with proper design of overhang pairs. Metrics need not to be binary, but were chosen to be in this example for simplicity and clarity (see Example 5).

Partitioning the Sequence

[0289] SEQ ID NO:1 was chosen as SOI. In this example, it was first partitioned in a naïve manner, that is in equal lengths of 16 nt and with overhangs of length=4. This results in 16 single stranded oligonucleotide sequences that, once the target double stranded polynucleotide is assembled, will be identical in sequence to the SOI.

Symmetric and Asymmetric Assembly Trees

[0290] From the resulting selection of oligos we compare two partitions: a symmetric and an asymmetric one. The former is, as disclosed in the prior art, the simplest binary tree (FIG. 1). To choose an asymmetric tree the following steps were applied:

[0291] i. Start with a symmetric tree

[0292] ii. Evaluate the score at each node

[0293] iii. If there are nodes that have score *w*=0, the branches are modified by re-splitting at one deeper node in such a way that the conflicting oligos are not in the same immediate node

[0294] iv. Repeat steps ii-iii 1 million times until there is a solution

[0295] v. If there is no solution after 1 million trials, the sequence is re-partitioned and the assembly tree computed by starting at step i.

[0296] Other approaches can be used, especially when sequences are longer and not all assembly trees can be computed due to combinatorial complexity; for example, by Monte Carlo sampling a set of assembly trees that are valid as measured by the metric above followed by picking a valid tree. This can result in asymmetric assembly workflows represented by trees that are “unbalanced” (in the technical graph jargon) and that lead to successful assemblies.

[0297] FIG. 5 shows the result of a flat, naïve partition (symmetric) that presents one problematic node in oligos c and d and an asymmetric assembly tree that resolves the conflict. FIG. 5c shows the adjacency matrices of the problematic node (oligos matching in an undesired manner) revealing that under the symmetric assembly (left matrix) workflow there are anticipated assembly problems, whereas under the asymmetric one (right matrix), there are none. The adjacency matrices consider not only the overhang matching but also the local adjacency as determined by the assembly workflow.

[0298] In bracket notation, the result of the partition of the asymmetric assembly tree is represented as $\{\{\{a, b\}, c\}, \{d, e\}, \{f, g\}\}, h\}$, which in graph representation is the tree of FIG. 5b.

Example 2. Assembly of a Target Molecule of 128 bp with an Asymmetric Workflow and Comparison to a Symmetric Workflow by Ligation of 16-Mers

[0299] After partitioning the target sequence (SEQ ID NO:1) and defining the assembly trees as described in Example 1, the oligos are assembled by following both a symmetric assembly (FIG. 5a), and by following the asymmetric tree described above (FIG. 5b), namely $\{\{\{a, b\}, c\}, \{d, e\}, \{f, g\}\}, h\}$.

[0300] All oligos a-h were procured by Integrated DNA Technologies (IDT) with standard desalting purification and were provided normalized at a concentration of 50 μ M on IDTE Buffer (pH 7.5). The oligos used in the assemblies below were single-stranded and pure.

Preparing the Annealing Solutions and Annealing

[0301] Some commercial buffers are ready to mix in H_2O such as New England Biolabs’ (NEB) T4 Ligase Reaction Buffer, product nr B0202S, and readily contain the ATP necessary for the ligase activity. In a microcentrifuge tube, 216 μ L of dd H_2O with NEB T4 ligase buffer were prepared and the solution was mixed well by vortexing. 24 μ L of this solution mix were dispensed into to 8 reaction tubes labelled a-h. 3 μ L of each oligo was transferred to 8 predefined tubes and mixed well by pipetting (sequences see FIG. 8, SEQ ID NOs: 2 to 17):

[0302] Oligos a- and a+ into tube a

[0303] Oligos b- and b+ into tube b

[0304] Oligos c- and c+ into tube c

[0305] Oligos d- and d+ into tube d

[0306] Oligos e- and e+ into tube e

[0307] Oligos f- and f+ into tube f

[0308] Oligos g- and g+ into tube h

[0309] Oligos h- and h+ into tube h

[0310] The tubes were sealed and incubated in a thermocycler for 30 sec at 98° C. The temperature was then decreased from 95° C. to 24° C. with a ramp function that diminished the temperature by 1° C. per minute allowing the matching pairs of ss oligos to anneal. Once finished the double stranded oligos were kept at 4° C.

Preparing the Ligation Solution.

[0311] The ligation solution was prepared on ice by mixing, in the following order, 32.5 μ L of nuclease free dd H_2O , 7.5 μ L of ligase buffer and 5 μ L of ATP for a final concentration of 10 mM. The ligation solution was mixed well by vortexing and spun down. 2.5 μ L of T4 Ligase (NEB, product nr. M0202) & 2.5 μ L of T4 polynucleotide kinase (PNK) (NEB, product nr. M0201B) were added for a total of 10 and 0.25 units per μ L of final solution, respectively, and mixed well by gently pipetting. The solution was kept on ice until needed. 10 μ L of the ligation solution were transferred to each of 4 tubes (b, d, f, h) containing 5 μ L the corresponding ds oligos and mixed by pipetting. Afterwards the tubes were sealed again.

Symmetric Assembly

[0312] The oligos are arrayed in rows on a 96-micro-well plate and pairs of oligos or reaction products are transferred in tiers (see FIG. 1):

[0313] i. First assembly tier. Transfer the contents of oligo mixtures a into mixture of oligo b (plus ligation solution), of oligo mixtures c into mixture of oligo d (plus ligation solution), of oligo mixtures e into mixture of oligo f (plus ligation solution), and of oligo mixtures g into mixture of oligo h (plus ligation solution).

[0314] ii. After gently mixing by pipetting, these two reactions were incubated for 30 min at 24° C.

[0315] iii. Second assembly tier. Transfer the contents of oligo mixtures ab (tube b) into mixture of oligo cd (tube d) and of oligo mixtures ef (tube f) into mixture of oligo gh (tube h).

[0316] iv. After gently mixing by pipetting, these two reactions were incubated for 30 min at 24° C.

[0317] v. Third assembly tier. Transfer the contents of oligo mixtures a.d (tube d) into mixture of oligo e.h (tube h).

[0318] vi. After gently mixing by pipetting, these two reactions were incubated for 30 min at 24° C.

[0319] vii. The final volume containing the 128 bp product was 80 μ L.

[0320] viii. Heat inactivating the ligation reaction by incubating for 10 min at 65° C.

Asymmetric Assembly by Step-by-Step Hand-Pipetting

[0321] In this case, it suffices to keep track of the order in which the oligos and their reaction products are combined, by proceeding according to the tree in FIG. 5b. The reaction conditions are the same as in the symmetric assembly. Although the asymmetric assembly will follow the steps in a similar manner as with the symmetric assembly, it is important to notice that the order of the transfers is crucial.

[0322] i. First assembly tier. Transfer the contents of oligo mixtures a (tube a) into mixture b (tube b), oligo mixtures d (tube d) into mixture e (tube e) and oligo mixtures f (tube f) into mixture g (tube g).

- [0323] ii. After gently mixing by pipetting these two reactions were incubated for 30 min at 24° C.
- [0324] iii. Second assembly tier. Transfer the contents of oligo mixtures c (tube c) into mixture a.b (tube b), the contents of oligo mixture d.e (tube e) into mixture f.g (tube g).
- [0325] iv. After gently mixing by pipetting these two reactions were incubated for 30 min at 24° C.
- [0326] v. Third assembly tier. Combine the reaction product a.g (tube g) with the mixture h (tube h).
- [0327] vi. Gently mix by pipetting and incubate the reaction for 30 min at room temperature.
- [0328] vii. The final volume containing the 128 bp product was 70 μ L.
- [0329] viii. Heat inactivating the ligation reaction by incubating for 10 min at 65° C.
- [0330] ix. Optionally, perform a bead purification by using MagMax beads of the final product following the provider's protocol. Alternatively, separation on agarose gel can be performed and the DNA extracted using purification kits, such as the Monarch PCR & DNA clean up kit from New England Biolabs (product nr. T1030).

Example 3. Assembly of a Target Molecule of 128 bp (SEQ ID NO:1) with an Asymmetric Workflow by Ligation of 16-Mers Using Liquid Handlers

[0331] In order to adapt the method to the use of automated liquid handlers, the order of the oligos is re-arranged and then liquid-handling steps are performed. Although this seems unnecessarily complicated for a set of 8 oligo dimers, longer sequences will require hundreds of dimers arrayed in a complex manner making the process highly prone to error (see Example 4). The detailed step-by-step instructions as described above may be confusing, in particular because the order of steps will vary according to the target sequence. For this reason, an alternative was developed, which is based on a scheme where the oligos are initially arrayed in a specific manner (also sequence dependent).

Arraying the Oligos

[0332] To facilitate the process of assembly by means of liquid handlers (multichannel pipettes or preferably automatic liquid handlers), a particular arraying of the oligos can be done that directly maps the results of the assembly tree to non-trivial placement of oligonucleotides on a microwell plate.

[0333] The asymmetric assembly tree can be casted into a larger tree of the same depth (nr. of tiers) that is symmetric but has empty leaves, or "Zeromers", denoted by $\emptyset$ (FIG. 6). Note that the zeromers are only an abstraction; in an experimental set-up they will correspond to empty wells or solution without oligos. In this example, the zeromers are always introduced to the left of any oligo (this has to do with mechanical transfer reasons, namely, to avoid moving liquids unnecessarily, however the same result can be achieved by inverting the direction of the transfers and the position of the zeromers) or sub-tree. In the case of the 8 oligos, the smallest symmetric tree is one having a total of 16 leaves containing 8 zeromers. FIG. 7a shows how the 8 oligos would be arrayed to perform regular transfers in a symmetric assembly workflow. FIG. 9b shows how the tree with zeromers is arrayed in a non-trivial manner, which will

facilitate transfer by a skilled person or by a liquid handler without worrying about the contents or specific movements of the wells. This is particularly important for longer sequences. For clarity and didactic reasons, however, it is described herein using a shorter sequence.

Procuring the Oligos

[0334] The arraying is a 1-to-1 mapping from the assembly tree to a 96 (or 384) micro-well plate. Thus, these can be directly procured or provided with the necessary array format.

[0335] In the case of the asymmetric assembly tree $\{\{\{a, b\}, c\}, \{\{d, e\}, \{f, g\}\}, h\}$ the oligos are arrayed as in the Starting Plate of FIG. 7, which is according to the following table 1:

TABLE 1		
Oligo Array		
Oligo	Micro-well	
	Symmetric Assembly	Asymmetric assembly
a	A1	A1
b	A7	A7
c	A2	A8
d	A8	A3
e	B1	A9
f	B7	A4
g	B2	A10
h	B8	B10

Assembling the Oligos

[0336] Arrayed in this manner, both the symmetric and asymmetric assembly trees can be implemented by regular movements. In the case of asymmetric trees, as indicated above, there are always one or more tiers of assembly. Yet the movements of the handler can be regular, even if they sometimes include some actions that do not carry liquids (e.g. wells A5-A6 or well B1-B9).

Example 4. Assembly of a Target Molecule of 256 bp with an Asymmetric Workflow by Ligation of Oligonucleotides of 16 bp and Comparing to Symmetric Assembly

[0337] In this example, a 256 bp sequence was used (SEQ ID NO:27). After partitioning the target sequence and defining the assembly trees as described in Example 1. The adjacency matrices (FIG. 9) show two problematic nodes in the symmetric assembly. The oligos are assembled by following both a symmetric assembly (FIG. 12a), and by following the asymmetric tree described above (FIG. 12b), namely $\{\{\{\{a, b\}, c\}, \{d, e\}\}, \{\{f, g\}, h\}\}, \{\{\{i, j, k\}, \{l, m\}, \{n, o\}\}\}, p\}$.

Asymmetric Assembly by Step-by-Step Hand-Pipetting

[0338] The assembly procedure is analogous to the one described in Example 2, but this time the number of required steps is larger to account for a larger assembly tree. Specifically, steps i-viii. follow a similar pattern as in the aforementioned example (with a different combination of mixing steps according to the different assembly tree), for

the two main branches of the tree (FIG. 10b). At this point we have tube h containing mixture a.h and tube p containing mixture i.p. We proceed as follows:

- [0339] ix. A bead purification of 128 bp products was performed by using MagMax bead technology and following manufacturer's instructions with the following modifications to the protocol: binding solution buffer to sample ratio used was 1.66. After beads purification, DNA was eluted in 17 μ l of ligation mix (T7 ligase (75 U/ μ l), T4 PNK (0.25 U/ μ l), 10 mM ATP, 10 $\times$ T4 NEB Buffer and H₂O)
- [0340] x. Fourth assembly tier. Combine the reaction product a.h (tube h) with the mixture i.p (tube p).
- [0341] xi. Gently mix by pipetting and incubate the reaction for 60 min at 24 C.
- [0342] xii. A bead purification of final 256 bp products was performed by using MagMax beads following the manufacturer's protocol with the following modifications to the protocol: binding solution buffer to sample ratio used was 1, elution in 17 μ l ddH₂O.

Visualization and Comparison of the Results

[0343] Following ligations, final products were subjected to quality control using capillary electrophoresis (Fragment Analyzer). Prior to the loading of the samples on the FA, samples were diluted 2 $\times$ in 1 $\times$ TE Buffer. Samples were run on the FA using Small Fragment Kit (Agilent, DNF-476-0500) and using corresponding SS Small Fragment method (Agilent, DNF-476-33) following manufacturer's instructions.

[0344] FIG. 11 shows the electropherograms of the two assemblies as obtained after the purification procedure. The symmetric assembly fails, since there is no clear peak at the target length and the target molecule could not be recovered even with purification. In contrast, the asymmetric assembly leads to results where the target molecule can be recovered after purification as seen in a clear peak.

[0345] Optionally, the successful construct can be isolated from the gel by using standard kits (e.g., Zymoclean), amplified by PCR (e.g., Sambrook and Russell, 2014; Chapter 8) and sequenced.

TABLE 2

Experimental yield and purity of polynucleotides with pooled, symmetric and asymmetric assembly methods.		
Assembly	Purity (% Conc)	Yield (ng/ μ L)
Tiered	<1.0	<0.1
Pooled	7.6	0.1
Asymmetric	40.0	0.37

Example 5. Assembly Tree Optimisation of a GC-Rich Target Molecule of 1028 bp with an Asymmetric Workflow by Ligation of Oligonucleotides of Different Lengths Using Liquid Handlers

[0346] In this example, the advantage of using an asymmetric workflow and optimization algorithms to partition molecules with challenging sequences into building blocks (oligos) that are of unequal length. The SOI (Seq_03, SEQ ID NO:60) is of 1028 bp and is partitioned in two ways: into fixed-length 16 mers and with a size adaptive between 15

and 17 nt. The scores of the balanced (symmetric) assembly tree for the naïve partition are compared to the optimised assembly trees of both the naïve and the adaptive partitions.

Choosing the Objective Function

[0347] In order to score an assembly tree, a "Misligation Matrix" ($M_{i,j}$) is generated. Each element of this matrix is 1 if the corresponding overhang pair is exposed during the ligation process, as a consequence of the chosen assembly tree, but it is not supposed to ligate; 0 otherwise. The ligation matrix ($L_{i,j}$) assigns a score (between 0 and 4000) directly related to the chance of ligation of the corresponding pair, irrespective of whether they are exposed during the assembly process. High GC content pairs have a larger score. Specifically, if the two overhangs have at least 3 out of 4 matching nucleotides according to Watson-Crick pairing, the matrix element is increased by a factor 10 for every A/T pair and by a factor 1000 for every G/C pair. To calculate the score, the two matrices are multiplied, element-wise, and then the total sum of each element is taken.

[0348] Overall, the score is:

$$S = -\left(\sum_{i,j} L_{i,j} \times M_{i,j}\right)/n^2$$

[0349] Where n is the total number of overhangs and the minus sign is necessary to make sure that assembly trees with more misligation propensity have lower scores.

Partitioning the Sequence and Selecting the Assembly Tree

[0350] By using the score defined in the previous section an optimal partition and assembly tree is computed under three different conditions.

[0351] First, a uniform, "naïve" partition is done by segmenting the SOI and its reverse complement into 16 mers (with 4 nt offset to allow for 4 nt 5' overhangs). The list of oligo sequences is provided in FIG. 14. A symmetric tree is used ad hoc as an assembly workflow. The score is parsed on this partition and topology. The naïve partition resulted in 128 oligonucleotide sequences.

[0352] Second, the same naïve partition above is used, but an asymmetric tree is computed in order to minimise mis-assemblies by means of maximising the score.

[0353] Third, an adaptive partition is done by locally maximising the score. Oligonucleotide sequences range between 15 and 17 nt and each dimer is constrained to have to 4 nt 5' overhangs. This partition resulted in 126 oligonucleotide sequences. The list of oligo sequences is in FIG. 15. After a locally-optimal partition has been completed, an optimal tree is computed by further maximising the score.

[0354] FIG. 12 shows three scored assembly trees, namely a naïve partition with a symmetric tree, a naïve partition with the asymmetric tree and an adaptive partition with a symmetric tree. FIG. 13 shows a graphical representation of the product of the ligation and misligation matrix.

[0355] As it appears evident from the scoring, the best result is obtained when the adaptive partitioning method meets the asymmetric assembly tree. In this case the score is increased by more than a factor two. The choice of an asymmetric tree implies additional steps are required for the assembly, but it is largely compensated by the extremely

increased chances of a successful assembly, even more so for a sequence like the SOI that has a GC content larger than 60%.

Provision of the Oligonucleotides

[0356] The synthesis of the oligos was outsourced to an established genomic company. The oligo providers were contacted to ensure that arraying included empty wells as required.

[0357] For the oligos from the naïve partition with symmetric tree, 1 plate of 364 micro wells was procured, reflecting the systematic order of the oligos (see Example 2) for the symmetric assembly.

[0358] The oligos of the naïve partition with an asymmetric tree were procured in 1 plate of 364 micro wells. Even though these are the same oligos as in the symmetric tree, we chose to acquire the oligos readily arrayed and avoid potential mistakes. Similar as in Example 4, the arraying was done by re-casting the oligos into a larger symmetric tree and adding zeroimers to the rightmost unassigned terminal branches of the tree. The same logic and procedure were followed to procure the oligos of the adaptive partition and asymmetric tree.

Assembly of the Sequences

[0359] Once oligonucleotides have been provided, these were annealed as in Examples 2-4 to form dimers with 5' overhangs.

[0360] The assembly steps were implemented through symmetric movement of and MCA arm on a Tecan Fluent under the same conditions as in Examples 2-4.

[0361] Comparison of the results by means of capillary electrophoresis indicated that the naïve assembly with the symmetric tree failed (no detectable peak). The naïve and adaptive partitions with asymmetric trees both resulted in detectable peaks in the electropherogram. The yield and purity proved to be a higher for the assembly based on the adaptive partition with asymmetric tree than on the naïve partition with asymmetric tree.

[0362] A PCR amplification of the target product was performed to enrich the sample. After amplification and clean-up with a Zymoclean kit, sequencing was performed on an Oxford Nanopore minION. The analysis of sequencing results indicate that the assembly is correct.

Assembly on Acoustic Dispensers

[0363] In a different embodiment, the transfer of oligonucleotides can be achieved by means of acoustic transfer (Echo Liquid Handler 525, Beckman-Coulter). In this embodiment, the arraying with zeroimers is unnecessary and it is possible to maximise use of the micro-well plates. Instead of mapping a tree into an array that is to be handled with symmetric movements of a multi-channel pipette, the mapping is performed as "worklist", i.e. a set of instructions (for example in XML format) that indicate the precise order of transfers to be done. Whilst this is not strictly parallel, the accelerated speed of acoustic dispensing makes it effectively parallel, e.g., as compared to the required incubation time of the reactions.

Example 6. Exemplary Materials and Features that May be Included in an Asymmetric Assembly Using Solid-Support Attachment

[0364] As a means to improve the yield of the assemblies, it may be convenient to use purification methods after some or every ligation or assembly reaction. Embodiments may use attachment of the oligonucleotides to a solid support. The oligos may have amino modifications in the 3' or 5' ends allowing a covalent or non-covalent bond to an adaptor or to a surface moiety. Solid support may comprise different treatments allowing stable or covalent links through carbonyl amide, carboxamide, sulfonamide or thiourea.

[0365] In some embodiments, nucleic acid polynucleotides of 50 or more nt or bp may be attached to solid supports by covalent modification of terminal phosphate groups to cellulose or polysaccharides. Other embodiments for attaching shorter oligonucleotides (20-50 nt) may use using phosphoramidate or amide linkages.

[0366] Some embodiments may use proteins or glycoproteins such as avidin, streptavidin or similar variants as part of the solid support to facilitate a non-covalent attachment to a biotin terminal modification. The biotin may be directly attached to the oligo/polynucleotides, thereby establishing a stable, non-covalent interaction. This can be reversed by incubating at high temperatures. Some embodiments use a light-sensitive release chemical moiety in the biotin, allowing cleaving of the covalent bond by exposure to the appropriate wave-length.

Partitioning the Sequence and Selecting the Assembly Tree

[0367] Partitioning of a target sequence is done as in Example 1 and this process is not influenced by the choice of any strategy for solid support and required modifications.

Provision of the Oligonucleotides on a Solid Support

[0368] Some or all oligonucleotides may be provided attached to a solid support by directly deriving products from column chemical synthesis by means of phosphoramidite chemistry, or by means of enzymatic synthesis using native or modified TdTs, amongst other enzymatic methods of DNA synthesis. In some embodiments oligonucleotides may be provided on a solid-support, where each oligo is in a separate compartment in a microwell plate with a treated surface allowing stable or covalent links. Some embodiments may attach the oligonucleotides to magnetic beads provided on separate compartments. Other embodiments may attach oligonucleotides to treated glass surfaces forming microarrays.

Preparation of the Assembly Through Annealing or Polymerase Extension

[0369] In some embodiments, the provided oligonucleotide building blocks may be converted into double stranded oligonucleotides by hybridisation of the attached oligos with other single stranded oligos that are in solution and may have no modifications. The hybridisation may be with oligos of equal or similar sizes so that the hybridised oligo results in blunt ends or may leave an overhang. The annealing may be done by controlling the annealing temperature with a thermocycler.

[0370] In other embodiments the annealing may be done with DNA primers in order to create a double stranded by

elongation using a polymerase such as TAQ or Klenow, thereby forming double stranded DNA or DNA-RNA hybrid.

[0371] After hybridisation or elongation, the sample may be subject to washing steps to eliminate unreacted oligonucleotides, cofactors and enzymes.

Detachment and Assembly of Oligonucleotides and Polynucleotides

[0372] Detachment methods depend on the chemical nature of the attachment and on the modification used on the oligonucleotide building blocks. In some embodiments, the detachment may use sequence-dependent enzymatic methods such as restriction enzymes to cleavage the DNA in specific sequences included in an attachment adaptor.

[0373] Another embodiment may use detaching a biotin-modified oligo or polynucleotide by means of thermal methods.

[0374] Since the presence of biotin may interfere in the assembly, or the use of thermal detachment may disrupt the secondary structure of the ds polynucleotide, some embodiments may use photo-cleavable linkages e.g., amino C6 (5' end) modifications covalently bonded either to the solid support or to biotin (Olejnik et al 1998). In this case, cleavage is achieved by exposure of the sample to long-wave UV light in the wave length range of 300-350, thereby releasing the oligo with a 5'-phosphate group which can be readily used in further chemical or enzymatic assembly steps.

[0375] Other detachment methods of DNA molecules are achieved by including a deoxyuridine site (U) at the 3' terminus as part of the linker to the solid support. The enzyme Uracil DNA glycosylase can then be used to transform the U into an abasic site, which subsequently can be cut using glycosylate-lyase endonuclease VIII, thereby releasing the DNA.

[0376] In any of these or other embodiments, the detachment at any step or tier is done for less than about half of the reaction components and only and exclusively (a) after pooling and reacting specific pairs or intermediate reaction products of oligonucleotides in separate reaction containers indicated by the asymmetric assembly tree, (b) performing a washing step to eliminate unwanted unreacted reagents from said reaction containers, and (c) before the need to transfer the reaction products to another reaction container which may contain another attached oligo or polynucleotide. Any other reaction compartment whose contents are not to be transferred at a specific step—as indicated by the assembly tree—, may be left momentarily attached. This is in comparison to symmetric trees where about half of the contents need to be detached and transferred.

Example 7. Use of Electro-Wetting on Dielectric Microfluidics Platform to Assemble a Target Polynucleotide Using an Asymmetric Workflow

[0377] An equivalent but different embodiment of an asymmetric assembly can be implemented in microfluidics platforms like those of Electro Wetting on Dielectric (EWOD), also known as “digital microfluidics”.

[0378] Although the way of handling the liquids is different in different microfluidics platforms, they may allow an even more efficient implementation of complex assembly workflows. Microfluidics have several advantages over stan-

dard liquid handling which include but are not limited to low consumable amounts thanks to the small droplet sizes and sparing a multitude of plastic tips that are costly, pollutant, imprecise and potentially pose bottlenecks to the workflow.

[0379] In some embodiments, a digital microfluidic platform is used in order to mix and react micro-droplets in a controlled manner. Droplets may be provided in micro-well plates and prepared with the enzymatic milieu as in FIGS. 2-5 plus necessary surfactants that are required to manipulate the droplets by electro-mechanical means. Droplets may be transferred to the EWOD platform by a liquid handler in small amounts, less than 1 μ L, preferably 0.1 μ L. The assembly tree is translated into digital movements that merge and mix droplets in accordance to each assembly tree. Once droplets are larger than the characteristic size of the EWOD's platform array, these can be combinatorically parallelized to have, effectively, replicas of the assemblies.

[0380] In addition, by using appropriate photosensitive probes it is feasible to discard microdroplets that have resulted on misassemblies and to gather data to improve assembly scores.

REFERENCES

- [0381]** Bonde, M. T., Kosuri, S., Genee, H. J., Sarup-Lytzen, K., Church, G. M., Sommer, M. O. A. and Wang H. H. (2014) Direct Mutagenesis of Thousands of Genomic Targets Using Microarray-Derived Oligonucleotides. *ACS Synthetic Biology* 4(1):17-22.
- [0382]** Farzadfard F, Lu TK. Synthetic biology. Genomically encoded analog memory with precise in vivo DNA writing in living cell populations. *Science*. 2014 Nov. 14; 346(6211):1256272. doi: 10.1126/science.1256272. PMID: 25395541; PMCID: PMC4266475.
- [0383]** Gao, X., LeProust, E., Zhang, H., Srivannavit, O., Gulari, E., Yu, P., . . . & Zhou, X. (2001). A flexible light-directed DNA chip synthesis gated by deprotection using solution photogenerated acids. *Nucleic Acids Research*, 29(22), 4744-4750.
- [0384]** Kai, J., Puntambekar A., Santiago N., Lee S. H., Sehly D. W., Moore V., Han J. and Ahn C. H. (2012) A novel microfluidic microplate as the next generation assay platform for enzyme linked immunoassays (ELISA). *Lab Chip*, 12(21): 4257-62.
- [0385]** LeProust, E. M., Peck, B. J., Spirin, K., McCuen, H. B., Moore, B., Namsaraev, E., and Caruthers, M. H. (2010) Synthesis of high-quality libraries of long (150 mer) oligonucleotides by a novel depurination controlled process. *Nucleic Acids Research*, 38(8), 2522-2540.
- [0386]** Neuner, P., Cortese, R., & Monaci, P. (1998). Codon-based mutagenesis using dimer-phosphoramidites. *Nucleic acids research*, 26(5), 1223-1227.
- [0387]** Sondek, J., & Shortle, D. (1992). A general strategy for random insertion and substitution substoichiometric mutagenesis: coupling of trinucleotide phosphoramidites. *Proceedings of the National Academy of Sciences*, 89(8), 3581-3585.
- [0388]** Olejnik, J., Krzymanska-Olejnik, E., Rothschild, K. J. (1998) Photocleavable aminotag phosphoramidites for 5'-termini DNA/RNA labeling. *Nucleic Acids Res.* 26:3572-3576.

 SEQUENCE LISTING

Sequence total quantity: 325

SEQ ID NO: 1 moltype = DNA length = 132

FEATURE Location/Qualifiers

misc_feature 1..132

note = Sequence of Interest

source 1..132

mol_type = other DNA

organism = synthetic construct

SEQUENCE: 1

agccataaga cgtactgcac ctgttgggtt acaatcatat tctttcatga gagtcacgca 60

tcccaatcgc cccgtaactt ccatacctttg acccgtaaaa ttgcttcccc gtgtaaacag 120

caacctccaa aa 132

SEQ ID NO: 2 moltype = DNA length = 16

FEATURE Location/Qualifiers

misc_feature 1..16

note = Oligo

source 1..16

mol_type = other DNA

organism = synthetic construct

SEQUENCE: 2

agccataaga cgtact 16

SEQ ID NO: 3 moltype = DNA length = 16

FEATURE Location/Qualifiers

misc_feature 1..16

note = Oligo

source 1..16

mol_type = other DNA

organism = synthetic construct

SEQUENCE: 3

gtgcagtacg tcttat 16

SEQ ID NO: 4 moltype = DNA length = 16

FEATURE Location/Qualifiers

misc_feature 1..16

note = Oligo

source 1..16

mol_type = other DNA

organism = synthetic construct

SEQUENCE: 4

gcacctgttg gggttac 16

SEQ ID NO: 5 moltype = DNA length = 16

FEATURE Location/Qualifiers

misc_feature 1..16

note = Oligo

source 1..16

mol_type = other DNA

organism = synthetic construct

SEQUENCE: 5

gattgtaacc caacag 16

SEQ ID NO: 6 moltype = DNA length = 16

FEATURE Location/Qualifiers

misc_feature 1..16

note = Oligo

source 1..16

mol_type = other DNA

organism = synthetic construct

SEQUENCE: 6

aatcatattc tttcat 16

SEQ ID NO: 7 moltype = DNA length = 16

FEATURE Location/Qualifiers

misc_feature 1..16

note = Oligo

source 1..16

mol_type = other DNA

organism = synthetic construct

SEQUENCE: 7

tctcatgaaa gaatat 16

SEQ ID NO: 8 moltype = DNA length = 16

FEATURE Location/Qualifiers

-continued

misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 8		
gagagtcacg catccc		16
SEQ ID NO: 9	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 9		
gattgggatg cgtgac		16
SEQ ID NO: 10	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 10		
aatcgccccg taactt		16
SEQ ID NO: 11	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 11		
atggaagtta cggggc		16
SEQ ID NO: 12	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 12		
ccatcctttg acccgt		16
SEQ ID NO: 13	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 13		
tttgacgggt caaagg		16
SEQ ID NO: 14	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 14		
caaattgctt ccccg		16
SEQ ID NO: 15	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 15		

-continued

taacacgggg aagcaa	16
SEQ ID NO: 16	moltype = DNA length = 16
FEATURE	Location/Qualifiers
misc_feature	1..16
	note = Oligo
source	1..16
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 16	
gttaacagca acctcc	16
SEQ ID NO: 17	moltype = DNA length = 16
FEATURE	Location/Qualifiers
misc_feature	1..16
	note = Oligo
source	1..16
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 17	
ttttggaggt tgctgt	16
SEQ ID NO: 18	moltype = DNA length = 16
FEATURE	Location/Qualifiers
misc_feature	1..16
	note = Oligo
source	1..16
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 18	
atagtatcga gcgacc	16
SEQ ID NO: 19	moltype = DNA length = 16
FEATURE	Location/Qualifiers
misc_feature	1..16
	note = Oligo
source	1..16
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 19	
gattggtcgc tcgata	16
SEQ ID NO: 20	moltype = DNA length = 16
FEATURE	Location/Qualifiers
misc_feature	1..16
	note = Oligo
source	1..16
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 20	
aatcgcatcg gcatat	16
SEQ ID NO: 21	moltype = DNA length = 16
FEATURE	Location/Qualifiers
misc_feature	1..16
	note = Oligo
source	1..16
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 21	
ctatatatgc cgatgc	16
SEQ ID NO: 22	moltype = DNA length = 16
FEATURE	Location/Qualifiers
misc_feature	1..16
	note = Oligo
source	1..16
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 22	
atagtatggc catatg	16
SEQ ID NO: 23	moltype = DNA length = 16
FEATURE	Location/Qualifiers
misc_feature	1..16
	note = Oligo

-continued

```

source                1..16
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 23
tgagcatatg gccata                                16

SEQ ID NO: 24      moltype = DNA length = 16
FEATURE            Location/Qualifiers
misc_feature       1..16
                      note = Oligo
source             1..16
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 24
ctcattgaga gtactg                                16

SEQ ID NO: 25      moltype = DNA length = 16
FEATURE            Location/Qualifiers
misc_feature       1..16
                      note = Oligo
source             1..16
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 25
gtcacagtac tctcaa                                16

SEQ ID NO: 26      moltype = DNA length = 16
FEATURE            Location/Qualifiers
misc_feature       1..16
                      note = Oligo
source             1..16
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 26
atgttatcga gcgacc                                16

SEQ ID NO: 27      moltype = DNA length = 260
FEATURE            Location/Qualifiers
misc_feature       1..260
                      note = Sequence of Interest
source             1..260
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 27
gcaccaaacy aaaatactat gggccaacga taccgagtgt acggacacag acgattacgt 60
ctgaatctca tgcccagtag tcggaggcgc ggccagagat attgaggcga tcagccaccc 120
gaatggagaa caaagccaca acccgatttg cgtctctatt tgttttgatt gccgcatcat 180
ctagtgatga gggcacctga acaccttaat acctaattcg tcaccacatt gttgtgagcc 240
atagggggac atcctgttta                                260

SEQ ID NO: 28      moltype = DNA length = 16
FEATURE            Location/Qualifiers
misc_feature       1..16
                      note = Oligo
source             1..16
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 28
gcaccaaacy aaaata                                16

SEQ ID NO: 29      moltype = DNA length = 16
FEATURE            Location/Qualifiers
misc_feature       1..16
                      note = Oligo
source             1..16
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 29
atagtatttt cgtttg                                16

SEQ ID NO: 30      moltype = DNA length = 16
FEATURE            Location/Qualifiers
misc_feature       1..16
                      note = Oligo
source             1..16
                      mol_type = other DNA

```

-continued

SEQUENCE: 30	organism = synthetic construct	
ctatgggcca acgata		16
SEQ ID NO: 31	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 31		
tcggtatcgt tggccc		16
SEQ ID NO: 32	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 32		
cggagtgtac ggacac		16
SEQ ID NO: 33	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 33		
gtctgtgtcc gtacac		16
SEQ ID NO: 34	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 34		
agacgattac gtctga		16
SEQ ID NO: 35	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 35		
agattcagac gtaatc		16
SEQ ID NO: 36	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 36		
atctcatgcc cagtag		16
SEQ ID NO: 37	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 37		
ccgactactg ggcata		16
SEQ ID NO: 38	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	

-continued

misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 38		
tcggaggcgc gccag		16
SEQ ID NO: 39	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 39		
atctctggcc gcgcct		16
SEQ ID NO: 40	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 40		
agatattgag gcgac		16
SEQ ID NO: 41	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 41		
ggctgatcgc ctcaat		16
SEQ ID NO: 42	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 42		
agccacccga atggag		16
SEQ ID NO: 43	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 43		
tggtctccat tcgggt		16
SEQ ID NO: 44	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 44		
aacaaagcca caaccc		16
SEQ ID NO: 45	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 45		

-continued

atacgggttg tggett	16
SEQ ID NO: 46	moltype = DNA length = 16
FEATURE	Location/Qualifiers
misc_feature	1..16
	note = Oligo
source	1..16
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 46	
gtattgcgtc tctatt	16
SEQ ID NO: 47	moltype = DNA length = 16
FEATURE	Location/Qualifiers
misc_feature	1..16
	note = Oligo
source	1..16
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 47	
aacaaataga gacgca	16
SEQ ID NO: 48	moltype = DNA length = 16
FEATURE	Location/Qualifiers
misc_feature	1..16
	note = Oligo
source	1..16
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 48	
tgttttgatt gccgca	16
SEQ ID NO: 49	moltype = DNA length = 16
FEATURE	Location/Qualifiers
misc_feature	1..16
	note = Oligo
source	1..16
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 49	
atgatgcggc aatcaa	16
SEQ ID NO: 50	moltype = DNA length = 16
FEATURE	Location/Qualifiers
misc_feature	1..16
	note = Oligo
source	1..16
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 50	
tcatctagtg atgagg	16
SEQ ID NO: 51	moltype = DNA length = 16
FEATURE	Location/Qualifiers
misc_feature	1..16
	note = Oligo
source	1..16
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 51	
gtgccctcat cactag	16
SEQ ID NO: 52	moltype = DNA length = 16
FEATURE	Location/Qualifiers
misc_feature	1..16
	note = Oligo
source	1..16
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 52	
gcacctgaac acctta	16
SEQ ID NO: 53	moltype = DNA length = 16
FEATURE	Location/Qualifiers
misc_feature	1..16
	note = Oligo

-continued

source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 53		
gtattaaggt gttcag		16
SEQ ID NO: 54	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 54		
atacctaatt cgtcac		16
SEQ ID NO: 55	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 55		
tgtggtgacg aattag		16
SEQ ID NO: 56	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 56		
cacattgttg tgagcc		16
SEQ ID NO: 57	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 57		
ctatggctca caacaa		16
SEQ ID NO: 58	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 58		
atagggggac atcctg		16
SEQ ID NO: 59	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 59		
taaacaggat gtcccc		16
SEQ ID NO: 60	moltype = DNA length = 1028	
FEATURE	Location/Qualifiers	
misc_feature	1..1028	
	note = Sequence of Interest	
source	1..1028 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 60		
cacttagtgc tcgtcccggt gatcgctggt gcatcagggc cttgagctgg cctgttggtg		60
ggcctgctaa cgccgccctc ctgtggacga agtgaatga gagacgtgcg ccatgttatc		120

```
SEQ ID NO: 67          moltype = DNA   length = 12  
FEATURE               Location/Qualifiers  
misc_feature           1..12  
                        note = Oligo  
source                 1..12
```

-continued

SEQUENCE: 67 catatcggta aa	mol_type = other DNA organism = synthetic construct	12
SEQ ID NO: 68 SEQUENCE: 68 000	moltype = length =	
SEQ ID NO: 69 FEATURE misc_feature source	moltype = DNA length = 12 Location/Qualifiers 1..12 note = Oligo 1..12 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 69 tcgaggaacg ca		12
SEQ ID NO: 70 FEATURE misc_feature source	moltype = DNA length = 10 Location/Qualifiers 1..10 note = Oligo 1..10 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 70 tatcggtaaa		10
SEQ ID NO: 71 FEATURE misc_feature source	moltype = DNA length = 10 Location/Qualifiers 1..10 note = Oligo 1..10 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 71 tcgaggaacg		10
SEQ ID NO: 72 FEATURE misc_feature source	moltype = DNA length = 12 Location/Qualifiers 1..12 note = Oligo 1..12 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 72 catatcggta aa		12
SEQ ID NO: 73 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 73 cacttagtgc tcgtcc		16
SEQ ID NO: 74 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 74 cgttgatcgc tgggtgc		16
SEQ ID NO: 75 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct	

-continued

SEQUENCE: 75 atcagggcct tgagct		16
SEQ ID NO: 76 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 76 ggcctgttgg tgggcc		16
SEQ ID NO: 77 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 77 tgctaacgcc gccctc		16
SEQ ID NO: 78 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 78 ctgtggacga agtggga		16
SEQ ID NO: 79 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 79 atgagagacg tgcgcc		16
SEQ ID NO: 80 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 80 atgttatctc cgccag		16
SEQ ID NO: 81 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 81 ccgtccatgc gtgacc		16
SEQ ID NO: 82 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 82 gggtcgggta cgaccg		16
SEQ ID NO: 83 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16	

-continued

source	note = Oligo 1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 83 ggccggaacg gactca		16
SEQ ID NO: 84 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 84 cacccatggt gagcaa		16
SEQ ID NO: 85 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 85 tacgtatcca ccggga		16
SEQ ID NO: 86 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 86 gcttagtttg gcccc		16
SEQ ID NO: 87 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 87 ttgggccgac ccgggg		16
SEQ ID NO: 88 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 88 agtccgaacg ctcggg		16
SEQ ID NO: 89 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 89 actactactt cacgag		16
SEQ ID NO: 90 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 90 caagcatctc cggcgc		16

-continued

SEQ ID NO: 91	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 91		
cgctgcccgc gatctg		16
SEQ ID NO: 92	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 92		
gcacgcccct tccaga		16
SEQ ID NO: 93	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 93		
ctcctgggca gcgaca		16
SEQ ID NO: 94	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 94		
gcgccgagag ggcggt		16
SEQ ID NO: 95	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 95		
cgccagggg aaccgc		16
SEQ ID NO: 96	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 96		
cctacagccc ccgaga		16
SEQ ID NO: 97	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 97		
tcgtttctcc cccgc		16
SEQ ID NO: 98	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	

-continued

SEQUENCE: 98	mol_type = other DNA	
ttccaccctt gggcga	organism = synthetic construct	16
SEQ ID NO: 99	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
source	note = Oligo	
	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 99		
ggtgcataacc tttcaa		16
SEQ ID NO: 100	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
source	note = Oligo	
	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 100		
ccgtcaaat gccact		16
SEQ ID NO: 101	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
source	note = Oligo	
	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 101		
tagaacgtcc tgctcg		16
SEQ ID NO: 102	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
source	note = Oligo	
	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 102		
tccgtagaaa acaggc		16
SEQ ID NO: 103	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
source	note = Oligo	
	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 103		
agacgccgga cgccca		16
SEQ ID NO: 104	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
source	note = Oligo	
	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 104		
tatgggaggc gctcgt		16
SEQ ID NO: 105	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
source	note = Oligo	
	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 105		
tggcccgtg ggcgac		16
SEQ ID NO: 106	moltype = DNA length = 16	

-continued

FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 106		
tgggcgtgac tggcga		16
SEQ ID NO: 107	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 107		
ccgccaggta tccccg		16
SEQ ID NO: 108	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 108		
gggggagcgc gccacg		16
SEQ ID NO: 109	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 109		
ctgcaggcgg acgcga		16
SEQ ID NO: 110	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 110		
gggagatgca gggacg		16
SEQ ID NO: 111	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 111		
gggtggtcag cttgac		16
SEQ ID NO: 112	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 112		
ccgggcctgg gaggct		16
SEQ ID NO: 113	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	

-continued

SEQUENCE: 113 cgcccgcccc ctcgcg	16
SEQ ID NO: 114 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct
SEQUENCE: 114 gctagactgg gatagc	16
SEQ ID NO: 115 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct
SEQUENCE: 115 ggtataccag cgggag	16
SEQ ID NO: 116 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct
SEQUENCE: 116 gagtcctctcc ggcgtc	16
SEQ ID NO: 117 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct
SEQUENCE: 117 cggccctgca ggcaaa	16
SEQ ID NO: 118 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct
SEQUENCE: 118 ggcgtgagcg tttccc	16
SEQ ID NO: 119 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct
SEQUENCE: 119 gctgtccggg cggtag	16
SEQ ID NO: 120 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct
SEQUENCE: 120 cggctggggc catgtc	16
SEQ ID NO: 121 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16

-continued

source	note = Oligo 1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 121 cgacggcgcg cgatgt		16
SEQ ID NO: 122 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 122 agtcggggcc cattcc		16
SEQ ID NO: 123 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 123 acggtgcaac gtcccc		16
SEQ ID NO: 124 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 124 ctacagtagg tcaatc		16
SEQ ID NO: 125 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 125 ggcaggcaca ggcgcg		16
SEQ ID NO: 126 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 126 cgcgcccgcc gcgcta		16
SEQ ID NO: 127 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 127 gttccgtggc cgtcgt		16
SEQ ID NO: 128 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 128 acagagcaga ggaatc		16

-continued

SEQ ID NO: 129	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 129		
cgaagtcggg ttgccc		16
SEQ ID NO: 130	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 130		
gggtctctgt ccggcc		16
SEQ ID NO: 131	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 131		
accgccccga gctccc		16
SEQ ID NO: 132	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 132		
tacgggacag cccggg		16
SEQ ID NO: 133	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 133		
aaagctggga gggaca		16
SEQ ID NO: 134	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 134		
gggtagtcgg cccccc		16
SEQ ID NO: 135	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 135		
cgctctgtcg ttccgc		16
SEQ ID NO: 136	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	

-continued

SEQUENCE: 136 atccgccttg cgggcc	mol_type = other DNA organism = synthetic construct	16
SEQ ID NO: 137 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 137 aacgggacga gcacta		16
SEQ ID NO: 138 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 138 tgatgcacca gcgata		16
SEQ ID NO: 139 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 139 ggccagctca aggccc		16
SEQ ID NO: 140 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 140 agcaggccca ccaaca		16
SEQ ID NO: 141 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 141 acaggagggc ggcgtt		16
SEQ ID NO: 142 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 142 tcattccact tcgtcc		16
SEQ ID NO: 143 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 143 acatggcgca cgtctc		16
SEQ ID NO: 144	moltype = DNA length = 16	

-continued

FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 144		
acggctggcg gagata		16
SEQ ID NO: 145	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 145		
acccggtcac gcatgg		16
SEQ ID NO: 146	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 146		
ggcccggtcg taccgg		16
SEQ ID NO: 147	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 147		
ggtgtgagtc cgttcc		16
SEQ ID NO: 148	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 148		
cgtattgctc accatg		16
SEQ ID NO: 149	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 149		
aagctcccgg tggata		16
SEQ ID NO: 150	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 150		
ccaagggggc caaact		16
SEQ ID NO: 151	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	

-continued

SEQUENCE: 151 gactccccgg gtcggc	16
SEQ ID NO: 152 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct
SEQUENCE: 152 tagtccccgag cgttcg	16
SEQ ID NO: 153 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct
SEQUENCE: 153 cttgctcgtg aagtag	16
SEQ ID NO: 154 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct
SEQUENCE: 154 agcggcgccg gagatg	16
SEQ ID NO: 155 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct
SEQUENCE: 155 gtgccagatc gcgggc	16
SEQ ID NO: 156 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct
SEQUENCE: 156 ggagtctgga aggggc	16
SEQ ID NO: 157 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct
SEQUENCE: 157 gcgctgtcgc tgccca	16
SEQ ID NO: 158 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct
SEQUENCE: 158 ggcgaccgcc ctctcg	16
SEQ ID NO: 159 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16

-continued

source	note = Oligo 1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 159 tagggcggtt cccctg		16
SEQ ID NO: 160 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 160 acgatctcgg gggctg		16
SEQ ID NO: 161 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 161 ggaagcgggg ggagaa		16
SEQ ID NO: 162 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 162 cacctcgccc aggggt		16
SEQ ID NO: 163 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 163 cgggttgaaa ggtatg		16
SEQ ID NO: 164 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 164 tctaagtggc atttga		16
SEQ ID NO: 165 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 165 cggacgagca ggacgt		16
SEQ ID NO: 166 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 166 gtctgctgt tttcta		16

-continued

SEQ ID NO: 167	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 167		
catatgggcg tccggc		16
SEQ ID NO: 168	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 168		
gccaacgagc gcctcc		16
SEQ ID NO: 169	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 169		
cccagtcgcc cagcgg		16
SEQ ID NO: 170	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 170		
gcggtcgcca gtcacg		16
SEQ ID NO: 171	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 171		
ccccgggga tacctg		16
SEQ ID NO: 172	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 172		
gcagcgtggc gcgctc		16
SEQ ID NO: 173	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 173		
tccctcgcgt ccgcct		16
SEQ ID NO: 174	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	

-continued

SEQUENCE: 174 accccgctccc tgcatac	mol_type = other DNA organism = synthetic construct	16
SEQ ID NO: 175 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 175 cgggtcaag ctgacc		16
SEQ ID NO: 176 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 176 gacgagcctc ccaggc		16
SEQ ID NO: 177 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 177 tagccgcgag gggccg		16
SEQ ID NO: 178 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 178 taccgctatc ccagtc		16
SEQ ID NO: 179 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 179 actcctcccg ctggta		16
SEQ ID NO: 180 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 180 gccggacgcc ggagag		16
SEQ ID NO: 181 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 181 cgcccttgcc tgcagg		16
SEQ ID NO: 182	moltype = DNA length = 16	

-continued

FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 182		
cagcgggaaa cgctca		16
SEQ ID NO: 183	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 183		
gccgctaccg cccgga		16
SEQ ID NO: 184	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 184		
gtcggacatg gcccca		16
SEQ ID NO: 185	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 185		
gactacatcg cgcgcc		16
SEQ ID NO: 186	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 186		
ccgtggaatg ggcccc		16
SEQ ID NO: 187	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 187		
gtagggggac gttgca		16
SEQ ID NO: 188	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 188		
tgccgattga cctact		16
SEQ ID NO: 189	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	

-continued

SEQUENCE: 189 cgcgcgcgcc tgtgcc	16
SEQ ID NO: 190 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct
SEQUENCE: 190 gaactagcgc ggcggg	16
SEQ ID NO: 191 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct
SEQUENCE: 191 ctgtacgacg gccacg	16
SEQ ID NO: 192 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct
SEQUENCE: 192 ttcggattcc tctgct	16
SEQ ID NO: 193 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct
SEQUENCE: 193 gcccgggcaa cccgac	16
SEQ ID NO: 194 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct
SEQUENCE: 194 cgttgccggg acagga	16
SEQ ID NO: 195 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct
SEQUENCE: 195 cgtagggagc tcgggg	16
SEQ ID NO: 196 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct
SEQUENCE: 196 ctttcccggg ctgtcc	16
SEQ ID NO: 197 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16

-continued

source	note = Oligo 1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 197 accctgtccc tcccag		16
SEQ ID NO: 198 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 198 agcggggggg ccgact		16
SEQ ID NO: 199 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 199 ggatgcgga cgcacag		16
SEQ ID NO: 200 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 200 gcggggcccg caaggc		16
SEQ ID NO: 201 FEATURE misc_feature	moltype = DNA length = 15 Location/Qualifiers 1..15 note = Oligo	
source	1..15 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 201 cacttagtgc tcgtc		15
SEQ ID NO: 202 FEATURE misc_feature	moltype = DNA length = 15 Location/Qualifiers 1..15 note = Oligo	
source	1..15 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 202 ccgttgatcg ctggt		15
SEQ ID NO: 203 FEATURE misc_feature	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo	
source	1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 203 gcacagggc cttgagc		17
SEQ ID NO: 204 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 204 tggcctgttg gtgggc		16

-continued

SEQ ID NO: 205	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 205		
ctgctaacgc cgccct		16
SEQ ID NO: 206	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 206		
cctgtggacg aagtgga		17
SEQ ID NO: 207	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 207		
atgagagacg tgcgcca		17
SEQ ID NO: 208	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 208		
tgttatctcc gccagcc		17
SEQ ID NO: 209	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 209		
gtccatgcgt gaccggg		17
SEQ ID NO: 210	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 210		
tcgggtacga ccggggc		17
SEQ ID NO: 211	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
	note = Oligo	
source	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 211		
ggaacggact cacac		15
SEQ ID NO: 212	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	

-continued

SEQUENCE: 212	mol_type = other DNA	
ccatggtgag caatacg	organism = synthetic construct	17
SEQ ID NO: 213	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
source	note = Oligo	
	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 213		
tatccaccgg gagct		15
SEQ ID NO: 214	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
source	note = Oligo	
	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 214		
tagtttgcc ccctt		15
SEQ ID NO: 215	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
source	note = Oligo	
	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 215		
gggccgaccc ggggag		16
SEQ ID NO: 216	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
source	note = Oligo	
	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 216		
tccgaacgct cgggact		17
SEQ ID NO: 217	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
source	note = Oligo	
	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 217		
actacttcac gagcaag		17
SEQ ID NO: 218	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
source	note = Oligo	
	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 218		
catctccggc gccgctg		17
SEQ ID NO: 219	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
source	note = Oligo	
	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 219		
cccgcgatct ggcacgc		17
SEQ ID NO: 220	moltype = DNA length = 15	

-continued

FEATURE	Location/Qualifiers	
misc_feature	1..15	
	note = Oligo	
source	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 220		
cccttcagaga ctcct		15
SEQ ID NO: 221	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
	note = Oligo	
source	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 221		
gggcagcgac agcgc		15
SEQ ID NO: 222	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
	note = Oligo	
source	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 222		
cgagagggcg gtcgc		15
SEQ ID NO: 223	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
	note = Oligo	
source	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 223		
ccaggggaac cgccc		15
SEQ ID NO: 224	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 224		
tacagccccc gagatcg		17
SEQ ID NO: 225	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 225		
tttctccccc cgcttcc		17
SEQ ID NO: 226	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 226		
accctgggc gaggtgc		17
SEQ ID NO: 227	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	

-continued

SEQUENCE: 227 atacctttca acccgtc		17
SEQ ID NO: 228 FEATURE misc_feature source	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo 1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 228 aaatgccact tagaacg		17
SEQ ID NO: 229 FEATURE misc_feature source	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo 1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 229 tcctgctcgt ccgtaga		17
SEQ ID NO: 230 FEATURE misc_feature source	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo 1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 230 aaacaggcag acgccgg		17
SEQ ID NO: 231 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 231 acgcccataat gggagg		16
SEQ ID NO: 232 FEATURE misc_feature source	moltype = DNA length = 15 Location/Qualifiers 1..15 note = Oligo 1..15 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 232 cgctcgttgg cccgc		15
SEQ ID NO: 233 FEATURE misc_feature source	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo 1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 233 tgggcgactg ggcgtga		17
SEQ ID NO: 234 FEATURE misc_feature source	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo 1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 234 ctggcgaccg ccaggta		17
SEQ ID NO: 235 FEATURE misc_feature	moltype = DNA length = 17 Location/Qualifiers 1..17	

-continued

source	note = Oligo 1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 235 tccccggggg gagcgcg		17
SEQ ID NO: 236 FEATURE misc_feature	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo	
source	1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 236 ccacgctgca ggcggac		17
SEQ ID NO: 237 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 237 gcgagggaga tgcagg		16
SEQ ID NO: 238 FEATURE misc_feature	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo	
source	1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 238 gacggggtgg tcagctt		17
SEQ ID NO: 239 FEATURE misc_feature	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo	
source	1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 239 gaccggggcc tgggagg		17
SEQ ID NO: 240 FEATURE misc_feature	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo	
source	1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 240 ctcgtccggc ccctcgc		17
SEQ ID NO: 241 FEATURE misc_feature	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo	
source	1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 241 ggctagactg ggatagc		17
SEQ ID NO: 242 FEATURE misc_feature	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo	
source	1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 242 ggtataccag cgggagg		17

-continued

SEQ ID NO: 243	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 243		
gccctgcagg caaagg		16
SEQ ID NO: 244	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
	note = Oligo	
source	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 244		
cgtgagcggtt tcccg		15
SEQ ID NO: 245	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 245		
ctgtccgggc ggtagcg		17
SEQ ID NO: 246	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 246		
gctggggcca tgtccga		17
SEQ ID NO: 247	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 247		
cggcgcgcca tgtagtc		17
SEQ ID NO: 248	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 248		
ggggccatt ccacggt		17
SEQ ID NO: 249	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
	note = Oligo	
source	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 249		
gcaacgtccc cctac		15
SEQ ID NO: 250	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	

-continued

SEQUENCE: 250	mol_type = other DNA	
agtaggtcaa tcggcag	organism = synthetic construct	17
SEQ ID NO: 251	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
source	note = Oligo	
	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 251		
gcacaggcgc gcgcgcc		17
SEQ ID NO: 252	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
source	note = Oligo	
	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 252		
cgccgcgcta gttcc		15
SEQ ID NO: 253	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
source	note = Oligo	
	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 253		
gtggccgtcg tacag		15
SEQ ID NO: 254	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
source	note = Oligo	
	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 254		
agcagaggaa tccgaa		16
SEQ ID NO: 255	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
source	note = Oligo	
	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 255		
gtcgggttgc ccggg		15
SEQ ID NO: 256	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
source	note = Oligo	
	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 256		
ctcctgtccg gccaccg		17
SEQ ID NO: 257	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
source	note = Oligo	
	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 257		
ccccgagctc cctac		15
SEQ ID NO: 258	moltype = DNA length = 16	

-continued

FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 258		
gggacagccc gggaaa		16
SEQ ID NO: 259	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 259		
gctgggaggg acaggg		16
SEQ ID NO: 260	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
	note = Oligo	
source	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 260		
tagtcggccc ccccg		15
SEQ ID NO: 261	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
	note = Oligo	
source	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 261		
ctctgtcggt cgcga		15
SEQ ID NO: 262	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
	note = Oligo	
source	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 262		
tccgccttgc gggcc		15
SEQ ID NO: 263	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
	note = Oligo	
source	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 263		
acgggacgag cacta		15
SEQ ID NO: 264	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
	note = Oligo	
source	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 264		
atgcaccagc gatca		15
SEQ ID NO: 265	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	

-continued

SEQUENCE: 265 gccagctcaa ggccctg		17
SEQ ID NO: 266 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 266 gcaggcccac caacag		16
SEQ ID NO: 267 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 267 caggagggcg gcgtta		16
SEQ ID NO: 268 FEATURE misc_feature source	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo 1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 268 tcattccact tcgtcca		17
SEQ ID NO: 269 FEATURE misc_feature source	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo 1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 269 aacatggcgc acgtctc		17
SEQ ID NO: 270 FEATURE misc_feature source	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo 1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 270 ggacggctgg cggagat		17
SEQ ID NO: 271 FEATURE misc_feature source	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo 1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 271 ccgaccgggt cacgcat		17
SEQ ID NO: 272 FEATURE misc_feature source	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo 1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 272 ttccggcccg gtcgtac		17
SEQ ID NO: 273 FEATURE misc_feature	moltype = DNA length = 15 Location/Qualifiers 1..15	

-continued

source	note = Oligo 1..15 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 273 atgggtgtga gtccg		15
SEQ ID NO: 274 FEATURE misc_feature	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo	
source	1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 274 gatacgtatt gctcacc		17
SEQ ID NO: 275 FEATURE misc_feature	moltype = DNA length = 15 Location/Qualifiers 1..15 note = Oligo	
source	1..15 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 275 actaagctcc cgggtg		15
SEQ ID NO: 276 FEATURE misc_feature	moltype = DNA length = 15 Location/Qualifiers 1..15 note = Oligo	
source	1..15 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 276 gccaagggg gccaa		15
SEQ ID NO: 277 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 277 cggactcccc gggtcg		16
SEQ ID NO: 278 FEATURE misc_feature	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo	
source	1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 278 tagtagtccc gagcggt		17
SEQ ID NO: 279 FEATURE misc_feature	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo	
source	1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 279 gatgcttgct cgtgaag		17
SEQ ID NO: 280 FEATURE misc_feature	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo	
source	1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 280 cgggcagcgg cgccgga		17

-continued

SEQ ID NO: 281	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 281		
agggcggtgc cagatcg		17
SEQ ID NO: 282	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
	note = Oligo	
source	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 282		
gcccaggagt ctgga		15
SEQ ID NO: 283	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
	note = Oligo	
source	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 283		
ctcgcgctg tcgct		15
SEQ ID NO: 284	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
	note = Oligo	
source	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 284		
ctggcgacc gccct		15
SEQ ID NO: 285	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
	note = Oligo	
source	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 285		
tgtaggcggt ttccc		15
SEQ ID NO: 286	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 286		
gaaacgatct cgggggc		17
SEQ ID NO: 287	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 287		
gggtggaagc gggggga		17
SEQ ID NO: 288	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	

-continued

SEQUENCE: 288	mol_type = other DNA	
gtatgcacct cgcccag	organism = synthetic construct	17
SEQ ID NO: 289	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
source	note = Oligo	
	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 289		
atttgacggg ttgaaag		17
SEQ ID NO: 290	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
source	note = Oligo	
	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 290		
aggacgttct aagtggc		17
SEQ ID NO: 291	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
source	note = Oligo	
	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 291		
gttttctacg gacgagc		17
SEQ ID NO: 292	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
source	note = Oligo	
	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 292		
gcgtccggcg tctgctt		17
SEQ ID NO: 293	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
source	note = Oligo	
	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 293		
agcgctccc atatgg		16
SEQ ID NO: 294	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
source	note = Oligo	
	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 294		
cccagcgggc caacg		15
SEQ ID NO: 295	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
source	note = Oligo	
	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 295		
ccagtcacgc ccagtcg		17
SEQ ID NO: 296	moltype = DNA length = 17	

-continued

FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 296		
gggataacctg gcggtcg		17
SEQ ID NO: 297	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 297		
gtggcgcgct ccccccg		17
SEQ ID NO: 298	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 298		
tcgcgctccgc ctgcagc		17
SEQ ID NO: 299	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 299		
cgtccctgca tctccc		16
SEQ ID NO: 300	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 300		
ggccaagctg accaccc		17
SEQ ID NO: 301	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 301		
cgagcctccc aggcccg		17
SEQ ID NO: 302	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 302		
agccgcgagg ggccgga		17
SEQ ID NO: 303	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	

-continued

SEQUENCE: 303 taccgctatc ccagtct		17
SEQ ID NO: 304 FEATURE misc_feature source	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo 1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 304 gactcctccc gctggta		17
SEQ ID NO: 305 FEATURE misc_feature source	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo 1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 305 gggccggacg ccggaga		17
SEQ ID NO: 306 FEATURE misc_feature source	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo 1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 306 cacgcctttg cctgca		16
SEQ ID NO: 307 FEATURE misc_feature source	moltype = DNA length = 15 Location/Qualifiers 1..15 note = Oligo 1..15 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 307 acagcgggaa acgct		15
SEQ ID NO: 308 FEATURE misc_feature source	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo 1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 308 cagccgctac cgcccg		17
SEQ ID NO: 309 FEATURE misc_feature source	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo 1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 309 gccgtcggac atggccc		17
SEQ ID NO: 310 FEATURE misc_feature source	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo 1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 310 ccccgactac atcgcg		17
SEQ ID NO: 311 FEATURE misc_feature	moltype = DNA length = 17 Location/Qualifiers 1..17	

-continued

source	note = Oligo 1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 311 ttgcaccgtg gaatggg		17
SEQ ID NO: 312 FEATURE misc_feature	moltype = DNA length = 15 Location/Qualifiers 1..15 note = Oligo	
source	1..15 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 312 tactgtaggg ggacg		15
SEQ ID NO: 313 FEATURE misc_feature	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo	
source	1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 313 gtgcctgccc attgacc		17
SEQ ID NO: 314 FEATURE misc_feature	moltype = DNA length = 17 Location/Qualifiers 1..17 note = Oligo	
source	1..17 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 314 ggcgggcccgcg cgcgcct		17
SEQ ID NO: 315 FEATURE misc_feature	moltype = DNA length = 15 Location/Qualifiers 1..15 note = Oligo	
source	1..15 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 315 ccacggaact agcgc		15
SEQ ID NO: 316 FEATURE misc_feature	moltype = DNA length = 15 Location/Qualifiers 1..15 note = Oligo	
source	1..15 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 316 tgctctgtac gacgg		15
SEQ ID NO: 317 FEATURE misc_feature	moltype = DNA length = 16 Location/Qualifiers 1..16 note = Oligo	
source	1..16 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 317 cgacttcgga ttcctc		16
SEQ ID NO: 318 FEATURE misc_feature	moltype = DNA length = 15 Location/Qualifiers 1..15 note = Oligo	
source	1..15 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 318 ggagcccggg caacc		15

-continued

SEQ ID NO: 319	moltype = DNA length = 17	
FEATURE	Location/Qualifiers	
misc_feature	1..17	
	note = Oligo	
source	1..17	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 319		
ggggcggtgg ccggaca		17
SEQ ID NO: 320	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
	note = Oligo	
source	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 320		
tcccgtaggg agctc		15
SEQ ID NO: 321	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 321		
cagctttccc gggctg		16
SEQ ID NO: 322	moltype = DNA length = 16	
FEATURE	Location/Qualifiers	
misc_feature	1..16	
	note = Oligo	
source	1..16	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 322		
actaccctgt ccctcc		16
SEQ ID NO: 323	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
	note = Oligo	
source	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 323		
agagcggggg ggccg		15
SEQ ID NO: 324	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
	note = Oligo	
source	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 324		
cggatgcgga acgac		15
SEQ ID NO: 325	moltype = DNA length = 15	
FEATURE	Location/Qualifiers	
misc_feature	1..15	
	note = Oligo	
source	1..15	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 325		
gcggggcccg caagg		15

1. A method of assembling nucleic acid (NA) building blocks to produce a double stranded (ds) target polynucleotide, comprising assembling NA building blocks according to a workflow comprising multiple assembly tiers, wherein

in a hierarchical assembly, at least one asymmetric pair of NA building blocks of different length that originate from different assembly tiers is connected to produce an intermediate polynucleotide in at least one of said multiple assembly

tiers, wherein said intermediate polynucleotide is used as a building block to connect further NA building blocks at both ends of said intermediate polynucleotide in one or more further assembly tiers.

2. The method of claim 1, wherein one of the NA building blocks of the asymmetric pair is a reaction product obtained by assembly of NA building blocks in a previous assembly tier, and the other NA building block of the asymmetric pair is a starting NA building block or a reaction product obtained by assembly of NA building blocks in an assembly tier that is different from said previous assembly tier.

3. The method of claim 1, wherein the NA building blocks are oligonucleotides or polynucleotides.

4. The method of claim 1, wherein the NA building blocks are double stranded or single stranded (ss).

5. The method of claim 1, wherein the workflow provides for the production of ds NA building blocks from respective ss NA building blocks.

6. The method of claim 1, wherein the NA building blocks have a length ranging from 8 nucleotides up to 20% of the length of the target polynucleotide.

7. The method of claim 1, wherein assembly is accomplished by connecting matching ss NA building blocks and/or connecting matching overhangs of a pair of ds NA building blocks.

8. The method of claim 1, wherein assembly is by hybridizing complementary nucleotide sequences, or by a ligation reaction which comprises enzymatic, chemical, or an adaptor ligation.

9. The method of claim 8, wherein said ligation reaction is an enzymatic ligation reaction using a DNA ligase an RNA ligase, a polymerase or ribozymes.

10. The method of claim 1, wherein a solid carrier is used to immobilize one or more of said NA building blocks, the target polynucleotide, or one or more intermediate(s) of assembly.

11. The method of claim 1, wherein said NA building blocks are obtained from a library of NA building blocks, which comprises a diversity of NA building blocks, each contained in a separate library containment.

12. The method of claim 11, wherein said library comprises

a) ss NA building blocks, and/or

b) ds NA building blocks.

13. The method of claim 1, wherein said target polynucleotide has a length of at least 132 bps.

14. The method of claim 1, further comprising a step of enriching the target polynucleotide or one or more intermediates of assembly, by polymerase chain reaction (PCR).

15. The method of claim 1, further comprising a step of processing the target polynucleotide or intermediates of assembly by means of enzymatic modification employing restriction enzymes or kinases, or by chemical means, to facilitate further ligation or to facilitate cloning the target polynucleotide into a vector or plasmid.

16. The method of claim 1, wherein one NA building block of said at least one asymmetric pair is a starting NA building block, and said asymmetric pair is assembled in a second or further tier of assembly.

17. The method of claim 1, wherein said hierarchical assembly comprises at least one asymmetric assembly of said at least one asymmetric pair of NA building blocks.

18. The method of claim 9, wherein the DNA ligase is a T3, T4 or T7 DNA ligase.

19. The method of claim 12, wherein said ss NA building blocks have been obtained by de novo synthesis.

20. The method of claim 12, wherein said ds NA building blocks are produced by assembly of respective ss NA building blocks.

* * * * *