

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent	12397006
Kind Code	B2
Date of Patent	August 26, 2025
Inventor(s)	Krill; Steven L. et al.

Fospropofol formulations

Abstract

The present disclosure provides pharmaceutical compositions for oral administration of fospropofol, or pharmaceutically acceptable salts of fospropofol, as well as methods of oral administration of fospropofol.

Inventors:	Krill; Steven L. (San Clemente, CA), Chen; Feng-Jing (Irvine, CA)
Applicant:	Epalex Corporation (Mountain View, CA)
Family ID:	1000008781168
Assignee:	EPALEX CORPORATION (Palo Alto, CA)
Appl. No.:	17/882711
Filed:	August 08, 2022

Prior Publication Data

Document Identifier	Publication Date
US 20220378808 A1	Dec. 01, 2022

Related U.S. Application Data

continuation parent-doc US 17217656 20210330 US 11207334 20211228 child-doc US 17466016
division parent-doc US 17466016 20210903 US 11439653 child-doc US 17882711

Publication Classification

Int. Cl.: **A61K31/661** (20060101); **A61K9/00** (20060101); **A61K9/20** (20060101); **A61K31/683** (20060101); **A61K47/12** (20060101)

U.S. Cl.:

CPC **A61K31/661** (20130101); **A61K9/0053** (20130101); **A61K9/2013** (20130101);
A61K31/683 (20130101); **A61K47/12** (20130101);

Field of Classification Search

CPC: A61K (31/661)

References Cited

U.S. PATENT DOCUMENTS

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
6383471	12/2001	Chen et al.	N/A	N/A
7230003	12/2006	Gallop et al.	N/A	N/A
7241807	12/2006	Xu et al.	N/A	N/A
7538092	12/2008	Orlando et al.	N/A	N/A
7550155	12/2008	Zhang et al.	N/A	N/A
7645792	12/2009	Xu et al.	N/A	N/A
8354454	12/2012	Mills et al.	N/A	N/A
8383687	12/2012	Harris et al.	N/A	N/A
8470861	12/2012	Anders et al.	N/A	N/A
8541400	12/2012	Johnsson et al.	N/A	N/A
8962696	12/2014	Harris et al.	N/A	N/A
9023813	12/2014	Shull	N/A	N/A
9272978	12/2015	Zhang et al.	N/A	N/A
9339553	12/2015	Zhang et al.	N/A	N/A
9556156	12/2016	Dugar et al.	N/A	N/A
9643917	12/2016	Li et al.	N/A	N/A
9757334	12/2016	Lovell et al.	N/A	N/A
10239851	12/2018	Li et al.	N/A	N/A
10568834	12/2019	Garti et al.	N/A	N/A
11207334	12/2020	Krill	N/A	A61K 9/2013
11331271	12/2021	Slusher et al.	N/A	N/A
11439653	12/2021	Krill et al.	N/A	N/A
11478490	12/2021	Krill	N/A	A61K 31/661
11547714	12/2022	Murphy et al.	N/A	N/A
11628178	12/2022	Krill	514/130	A61K 31/661
2005/0004381	12/2004	Gallop et al.	N/A	N/A
2006/0205969	12/2005	Xu et al.	N/A	N/A
2007/0135390	12/2006	West et al.	N/A	N/A
2007/0202158	12/2006	Slusher et al.	N/A	N/A
2007/0259933	12/2006	Virsik et al.	N/A	N/A
2008/0161400	12/2007	Virsik et al.	N/A	N/A
2008/0214508	12/2007	Slusher et al.	N/A	N/A
2008/0306285	12/2007	Hale et al.	N/A	N/A
2009/0076141	12/2008	Virsik	N/A	N/A
2009/0156562	12/2008	Winch	N/A	N/A
2009/0221532	12/2008	Gibiansky et al.	N/A	N/A
2009/0286763	12/2008	Xu et al.	N/A	N/A
2010/0311698	12/2009	Patel et al.	N/A	N/A

2011/0269844	12/2010	LeDonne	N/A	N/A
2012/0289470	12/2011	Heit et al.	N/A	N/A
2012/0295866	12/2011	Shull et al.	N/A	N/A
2012/0316247	12/2011	Xie et al.	N/A	N/A
2019/0151458	12/2018	Ciufolini et al.	N/A	N/A
2019/0224123	12/2018	Theisinger et al.	N/A	N/A
2020/0289404	12/2019	Slusher et al.	N/A	N/A

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
1698588	12/2004	CN	N/A
101675917	12/2009	CN	N/A
101675918	12/2009	CN	N/A
102351895	12/2011	CN	N/A
102382005	12/2011	CN	N/A
103172658	12/2012	CN	N/A
103816542	12/2013	CN	N/A
106138015	12/2015	CN	N/A
WO 2000/008033	12/1999	WO	N/A
WO 2000/048572	12/1999	WO	N/A
WO 2002/013810	12/2001	WO	N/A
WO 2003/057153	12/2002	WO	N/A
WO 2003/086413	12/2002	WO	N/A
WO 2004/030658	12/2003	WO	N/A
WO 2004/032971	12/2003	WO	N/A
WO 2005/044201	12/2004	WO	N/A
WO 2006/017351	12/2005	WO	N/A
WO 2006/033911	12/2005	WO	N/A
WO 2008/157627	12/2007	WO	N/A
WO 2009/016269	12/2008	WO	N/A
WO 2011/160267	12/2010	WO	N/A
WO 2011/160268	12/2010	WO	N/A
WO 2013/093931	12/2012	WO	N/A
WO 2017/205632	12/2016	WO	N/A

OTHER PUBLICATIONS

International Patent Application No. PCT/US2021/065225; Int'l Search Report and the Written Opinion; dated Mar. 15, 2022; 15 pages. cited by applicant

Abdelmalak et al.; "Fospropofol, A New Sedative Anesthetic, and Its Utility in the Perioperative Period"; Current Pharmaceutical Design; vol. 18; 2012; p. 6241-6252. cited by applicant

Meek et al.; "Comparing propofol with placebo for early resolution of acute migraine in adult emergency department patients: A double-blind randomised controlled trial"; Emergency Medicine Australasia; 2020; 8 pages. cited by applicant

Mitra et al.; "Propofol for migraine in the emergency department: A pilot randomised controlled trial"; Emergency Medicine Australasia; vol. 32; 2020; p. 542-547. cited by applicant

Piatka et al.; "Propofol for Treatment of Acute Migraine in the Emergency Department: A Systematic Review"; Academic Emergency Medicine; 2019; 13 pages. cited by applicant

Wei et al.; "Oral Delivery of Propofol with Methoxymethylphosphonic Acid as the Delivery Vehicle"; Journal of Medicinal Chemistry; vol. 60; 2017; p. 8580-8590. cited by applicant

Supplemental Tables S1 and S2 from Wozniak et al. ; "Gastrointestinal delivery of propofol from

fospropofol: its bioavailability and activity in rodents and human volunteers”; *Journal of Translational Medicine*; vol. 13; 2015; 3 pages. cited by applicant

Ahmad et al.; “Interactions between opioid drugs and propofol in laboratory models of seizures”; *British Journal of Anaesthesia*; vol. 74; 1995; p. 311-314. cited by applicant

Alessandri et al.; “Seizures and Sepsis: A Narrative Review”; *Journal of Clinical Medicine*; vol. 10; 2021; 10 pages. cited by applicant

Al-Hader et al.; “The Comparative Effects of Propofol, Thiopental, and Diazepam, Administered Intravenously, on Pentylentetrazol Seizure Threshold in The Rabbit”; *Life Sciences*; vol. 51; 1992; p. 779-786. cited by applicant

“The American Headache Society Position Statement on Integrating New Migraine Treatments Into Clinical Practice”; *Headache*; vol. 59; 2019; 18 pages. cited by applicant

Bauman et al.; “Seizure Clusters: Morbidity and Mortality”; *Frontier in Neurology*; vol. 12; Feb. 2021; 5 pages. cited by applicant

Baumgartner et al.; “A survey of the European Reference Network EpiCARE on clinical practice for selected rare epilepsies”; *Epilepsia Open*; vol. 6; 2021; p. 160-470. cited by applicant

Beghi; “Addressing the burden of epilepsy: Many unmet needs”; *Pharmacological Research*; vol. 107; 2016; p. 79-84. cited by applicant

Beghi; “The Epidemiology of Epilepsy”; *Neuroepidemiology*; vol. 54; 2020; p. 185-191. cited by applicant

Begley et al.; “The direct cost of epilepsy in the United States: A systematic review of estimates”; *Epilepsia*; vol. 56; 2015; p. 1376-1387. cited by applicant

Bengalorkar et al.; “Fospropofol: Clinical Pharmacology”; *Journal of Anaesthesiology Clinical Pharmacology*; vol. 27; 2011; p. 79-83. cited by applicant

Bialer et al.; “Progress report on new antiepileptic drugs: A summary of the Fifteenth Eilat Conference on New Antiepileptic Drugs and Devices (EILAT XV). I. Drugs in preclinical and early clinical development”; *Epilepsia*; vol. 61; 2020; p. 2340-2364. cited by applicant

Binnie et al.; “Acute effects of lamotrigine (BW430C) in persons with epilepsy”; *Epilepsia*; vol. 27; 1986; p. 248-254 (abstract only). cited by applicant

Binnie et al.; “Photosensitivity as a model for acute antiepileptic drug studies”; *Electroencephalogr Clin Neurophysiol.*; vol. 63; Jan. 1986; p. 35-41 (abstract only). cited by applicant

Binnie; “Preliminary evaluation of potential anti-epileptic drugs by single dose electrophysiological and pharmacological studies in patients”; *J. Neural Transm.*; vol. 72; 1988; p. 259-266 (abstract only). cited by applicant

Bonafede et al.; “Direct and Indirect Healthcare Resource Utilization and Costs Among Migraine Patients in the United States”; *Headache*; May 2018; p. 700-714. cited by applicant

Bond et al.; “The use of analogue scales in rating subjective feelings”; *Br. J. Med. Psychol.*; vol. 47; 1974; p. 211-218. cited by applicant

Borgdorff; “Arguments against the role of cortical spreading depression in migraine”; *Neurological Research*; vol. 40 No. 3; 2018; p. 173-181. cited by applicant

Brodie et al.; “Patterns of treatment response in newly diagnosed epilepsy”; *Neurology*; vol. 78; 2012; p. 1548-1554. cited by applicant

Brophy et al.; “Guidelines for the Evaluation and Management of Status Epilepticus”; *Neurocrit Care*; vol. 17; 2012; p. 3-23. cited by applicant

Burch et al.; “The Prevalence and Impact of Migraine and Severe Headache in the United States: Figures and Trends From Government Health Studies”; *Headache*; vol. 58; 2018; p. 496-505. cited by applicant

Cameron; “Opisthotonos again”; *Anaesthesia*; vol. 42; 1987; p. 1124. cited by applicant

“Summary Health Statistics: National Health Interview Survey”; https://ftp.cdc.gov/pub/Health_Statistics/NCHS/NHIS/SHS/2018_SHS_Table_A-5.pdf; U.S. Department of Health and Human Services; 2018; 9 pages. cited by applicant

Chernik et al.; “Validity and reliability of the Observer's Assessment of Alertness/Sedation Scale: study with Intravenous Midazolam”; *Journal of Clinical Psychopharmacology*; vol. 10 No. 4; 1990; p. 244-251. cited by applicant

“Clinical Brief—Examining the Economic Impact and Implications of Epilepsy”; *The American Journal of Managed Care*; Feb. 2020; 8 pages. cited by applicant

Cohen; “Clinical trial: a dose-response study of fospropofol disodium for moderate sedation during colonoscopy”; *Alimentary Pharmacology & Therapeutics*; vol. 27; 2008; p. 597-608. cited by applicant

Contreras et al.; “Bioavailability of Oral Propofol in Humans”; *Int'l Society for Anaesthetic Pharmacology*; 2011; p. 21-23 (abstracts). cited by applicant

Dalic et al.; “Managing drug-resistant epilepsy: challenges and solutions”; *Neuropsychiatric Disease and Treatment*; vol. 12; 2016; p. 2605-2616. cited by applicant

De Riu et al.; “Propofol Anticonvulsant Activity in Experimental Epileptic Status”; *British Journal of Anaesthesia*; vol. 69; 1992; p. 177-181. cited by applicant

Dinis-Oliveria; “Metabolic Profiles of Propofol and Fospropofol: Clinical and Forensic Interpretative Aspects”; *BioMed Research Int'l*; vol. 2018 Article 6852857; 2018; 16 pages. cited by applicant

Falco-Water; “Epilepsy—Definition, Classification, Pathophysiology, and Epidemiology”; *Seminars in Neurology*; vol. 40; 2020; p. 617-623. cited by applicant

Farzana et al.; “Parosmia and Dysgeusia after Intravenous Propofol-Based General Anesthesia: A Case Report”; *Annals of Cardiac Anaesthesia*; vol. 25; 2022; p. 112-115. cited by applicant

Feist et al.; “Prevalence and incidence of epilepsy”; *Neurology*; vol. 88; 2017; p. 296-303. cited by applicant

Fisher et al.; “Epileptic Seizures and Epilepsy: Definitions Proposed by the International League Against Epilepsy (ILAE) and the International Bureau for Epilepsy (IBE)”; *Epilepsia*; vol. 46 No. 4; 2005; p. 470-472. cited by applicant

Fisher et al.; “A practical clinical definition of epilepsy”; *Epilepsia*; vol. 55; 2014; p. 475-482. cited by applicant

Fodale et al.; “Propofol Infusion Syndrome an Overview of a Perplexing Disease”; *Drug Safety*; vol. 31; 2008; p. 293-303. cited by applicant

French et al.; “Efficacy and Tolerability of the New Antiepileptic Drugs, II: Treatment of Refractory Epilepsy: Report of the TTA and QSS Subcommittees of the American Academy of Neurology and the American Epilepsy Society”; *Epilepsia*; vol. 45; 2004; p. 410-423. cited by applicant

French et al.; “Inhaled alprazolam rapidly suppresses epileptic activity in photosensitive participants”; *Epilepsia*; vol. 60; 2019; p. 1602-1609. cited by applicant

French et al.; “Time to Start Calling Things by Their Own Names? The Case for Antiseizure Medicines”; *Epilepsia Current*; vol. 20; 2020; p. 69-72. cited by applicant

Gan et al.; “Determination of plasma concentrations of propofol associated with 50% reduction in postoperative nausea”; *Anesthesiology*; vol. 87; 1997; p. 779-784. cited by applicant

Garcia et al.; “General Anesthetic Actions on GABAA Receptors”; *Current Neuropharmacology*; vol. 8, 2010; p. 2-9. cited by applicant

“Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990-2015: a systematic analysis for the Global Burden of Disease Study 2015”; *GBD 2015 Disease and Injury Incidence and Prevalence Collaborators*; *Lancet*; vol. 388; Oct. 2016; p. 1545-1602. cited by applicant

“Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016”; *GBD 2016 Disease and Injury Incidence and Prevalence Collaborators*; *Lancet*; vol. 390; Sep. 2017; p. 1211-1259. cited by applicant

“Global, regional, and national burden of epilepsy, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016”; GBD 2016 Epilepsy Collaborators; Lancet; vol. 18; Apr. 2019; p. 357-375. cited by applicant

“Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017”; GBD 2017 Disease and Injury Incidence and Prevalence Collaborators; Lancet; vol. 392; Nov. 2018; p. 1789-1858. cited by applicant

Hasan; “Evaluation of the Anticonvulsant Effects of Propofol in Pentylenetetrazole kindled Rats”; The FASEB Journal; vol. 31; 2017; p. 813.6 (abstract only). cited by applicant

Hasan et al.; “Comparison between the effects of propofol and midazolam on pentylenetetrazole kindled convulsions in rats”; The FASEB Journal; vol. 33; 2019; p. 665.4 (abstract only). cited by applicant

“Headache Classification Committee of the International Headache Society (IHS)”; The International Classification of Headache Disorders, 3rd edition; Cephalalgia; vol. 38; 2018; 211 pages. cited by applicant

Hiraoka et al.; “Changes in drug plasma concentrations of an extensively bound and highly extracted drug, propofol, in response to altered plasma binding”; Clinical Pharmacology & Therapeutics; vol. 75; 2004; p. 324-330. cited by applicant

Hiraoka et al.; “Kidneys contribute to the extrahepatic clearance of propofol in humans, but not lungs and brain”; British Journal of Clinical Pharmacology; vol. 60; 2005; p. 176-182. cited by applicant

Hoymork et al.; “Why do women wake up faster than men from propofol anaesthesia?”; British Journal of Anaesthesia; vol. 95; 2005; p. 627-633. cited by applicant

Johannessen et al.; “Therapeutic drug monitoring of antiepileptic drugs: current status and future prospects”; Expert Opinion on Drug Metabolism & Toxicology; vol. 16; 2020; p. 227-238. cited by applicant

Kasteleijn-Nolst et al.; “Photosensitive epilepsy: a model to study the effects of antiepileptic drugs. Evaluation of the piracetam analogue, levetiracetam”; Epilepsy Research; vol. 25; 1996; p. 225-230. cited by applicant

Kasteleijn-Nolst et al.; “Evaluation of brivaracetam, a novel SV2A ligand, in the photosensitivity model”; Neurology; vol. 69; 2007; p. 1027-1034. cited by applicant

Katsarava et al.; “Defining the Differences Between Episodic Migraine and Chronic Migraine”; Curr Pain Headache Rep; vol. 16; 2012; p. 86-92. cited by applicant

Kumar et al.; “Intraoperative refractory status epilepticus caused by propofol—a case report—”; Korean Journal of Anesthesiology; vol. 74; 2021; p. 70-72. cited by applicant

Lee et al.; “Diagnosis and Treatment of Status Epilepticus”; Journal of Epilepsy Research; vol. 10; 2020; p. 45-54. cited by applicant

Lingamaneni et al.; “Anesthetic Properties of 4-Iodopropofol”; Anesthesiology; vol. 94; Jun. 2001; p. 1050-1057. cited by applicant

Lipton et al.; “Migraine prevalence, disease burden, and the need for preventive therapy”; Neurology; vol. 68; 2007; p. 343-349. cited by applicant

Lipton et al.; “Predicting Inadequate Response to Acute Migraine Medication: Results From the American Migraine Prevalence and Prevention (AMPP) Study”; Headache; vol. 56; 2016; p. 1635-1648. cited by applicant

Lowson et al.; “Anticonvulsant Properties of Propofol and Thiopentone: Comparison Using Two Tests in Laboratory Mice”; British Journal of Anaesthesia; vol. 64; 1990; p. 59-63. cited by applicant

Lu et al.; “Propofol-induced refractory status epilepticus at remission age in benign epilepsy with centrotemporal spikes—A case report and literature review”; Medicine; vol. 98; 2019; 5 pages. cited by applicant

Mahmoud et al.; “Migraine and the risk of cardiovascular and cerebrovascular events: a metaanalysis of 16 cohort studies including 1 152 407 subjects”; *BMJ Open*; 2018; 10 pages. cited by applicant

Marmura et al.; “The Acute Treatment of Migraine in Adults: The American Headache Society Evidence Assessment of Migraine Pharmacotherapies”; *Headache*; vol. 55; 2015; p. 3-20. cited by applicant

Mathew et al.; “Intravenous Valproate Sodium (Depacon) Aborts Migraine Rapidly: A Preliminary Report”; *Headache*; vol. 40; 2000; p. 720-723. cited by applicant

Meyer et al.; “Propofol: Pro- or Anticonvulsant Drug?”; *Int'l Anesthesia Research Society*; vol. 108; Jun. 2009; p. 1993-1994. cited by applicant

Munakata et al.; “Economic Burden of Transformed Migraine: Results From the American Migraine Prevalence and Prevention (AMPP) Study”; *Headache*; vol. 49; 2009; p. 498-508. cited by applicant

Ngugi et al.; “Estimation of the burden of active and life-time epilepsy: A meta-analytic approach”; *Epilepsia*; vol. 51(5); 2010; p. 883-890. cited by applicant

Nicolodi et al.; “Exploration of NMDA Receptors in Migraine: Therapeutic and Theoretic Implications”; *Int'l J. Clin. Pharm. Res*; vol. 15; 1995; p. 181-189. cited by applicant

Nishikawa et al.; “Inhibitory Influence of GABA on Central Serotonergic Transmission. Involvement of the Habenulo-Raph6 Pathways in the GABAergic Inhibition of Ascending Cerebral Serotonergic Neurons”; *Brain Research*; vol. 331; 1985; p. 81-90. cited by applicant

Oei-Lim et al.; “Pharmacokinetics of propofol during conscious sedation using target-controlled infusion in anxious patients undergoing dental treatment”; *British Journal of Anaesthesia*; vol. 80; 1998; p. 324-331. cited by applicant

Ohmori et al.; “The Anticonvulsant Action of Propofol on Epileptiform Activity in Rat Hippocampal Slices”; *Anesth Analg*; vol. 99; 2004; p. 1095-1101. cited by applicant

Pack et al.; “Epilepsy Overview and Revised Classification of Seizures and Epilepsies”; *Continuum Journal*; vol. 25; Apr. 2019; p. 306-321. cited by applicant

Penovich; “Acute Repetitive Seizures (ARS) or Cluster Seizures”; <https://www.epilepsyfoundationmn.org/2020/01/14/acute-repetitive-seizures-ars-or-cluster-seizures/>; Jan. 2021; 4 pages. cited by applicant

Perucca et al.; “30 years of second-generation antiseizure medications: impact and future perspectives”; *Lancet Neurology*; vol. 19; 2020; 12 pages. cited by applicant

Puledda et al.; “Non-Pharmacological Approaches for Migraine”; *Neurotherapeutics*; vol. 15; 2018; p. 336-345. cited by applicant

Puri G.D; “Target controlled infusion total intravenous anaesthesia and Indian patients: Do we need our own data?”; *Indian Journal of Anaesthesia*; vol. 62; 2018; p. 245-248. cited by applicant

Rampil; “A primer for EEG signal processing in anesthesia”; *Anesthesiology*; vol. 89; 1998; p. 980-1002. cited by applicant

Raoof et al.; “In vivo assessment of intestinal, hepatic, and pulmonary first pass metabolism of propofol in the rat”; *Pharmaceutical Research*; vol. 13; 1996; p. 891-895. cited by applicant

Rogawski, MD; “Common Pathophysiologic Mechanisms in Migraine and Epilepsy”; *Arch Neurol*; vol. 65 No. 6; Jun. 2008; p. 709-714. cited by applicant

Rohmann et al.; “Migraine, headache, and mortality in women: a cohort study”; *The Journal of Headache and Pain*; vol. 21:27; 2020; 8 pages. cited by applicant

Rui et al.; “National Hospital Ambulatory Medical Care Survey: 2017 Emergency Department Summary Tables”; *CDC National Center for Health Statistics*; https://www.cdc.gov/nchs/data/nhamcs/web_tables/2017_ed_web_tables-508.pdf; 2017; accessed Apr. 2020; 37 pages. cited by applicant

Sahinovic et al.; “Clinical Pharmacokinetics and Pharmacodynamics of Propofol”; *Clin Pharmacokinet*; vol. 57; 2018; p. 1539-1558. cited by applicant

Samra et al.; “Effects of propofol sedation on seizures and intracranially recorded epileptiform activity in patients with partial epilepsy”; *Anesthesiology*; vol. 82; 1995; p. 843-851. cited by applicant

Schwedt et al.; “Acute treatment of migraine in adults”; Wolters Kluwer; 2021; 36 pages. cited by applicant

Silberstein; “Practice parameter: evidence-based guidelines for migraine headache (an evidence-based review): report of the Quality Standards Subcommittee of the American Academy of Neurology”; *Neurology*; vol. 55; 2000; p. 754-762. cited by applicant

Silvestri et al.; “Fospropofol Disodium for Sedation in Elderly Patients Undergoing Flexible Bronchoscopy”; *J Bronchology Interv Pulmonol.*; vol. 18(1); Jan. 2011; p. 15-22. cited by applicant

Simon et al.; “Disposition and pharmacology of propofol glucuronide administered intravenously to animals”; *Xenobiotica*; vol. 22 No. 11; 1992; p. 1267-1273. cited by applicant

J. R. Sneyd; “Excitatory events associated with propofol anaesthesia: a review”; *Journal of the Royal Society of Medicine*; vol. 85; May 1992; p. 288-291. cited by applicant

Straube et al.; “Primary headaches during lifespan”; *The Journal of Headache and Pain*; vol. 20; 2019; 14 pages. cited by applicant

Strzelczyk et al.; “Expanding the Treatment Landscape for Lennox-Gastaut Syndrome: Current and Future Strategies”; *CNS Drugs*; vol. 35; 2021; p. 61-83. cited by applicant

“Summary Health Statistics: National Health Interview Survey, 2018”; U.S. Dept. of Health and Human Services; 2018; 9 pages. cited by applicant

Trapani et al.; “Propofol in Anesthesia. Mechanism of Action, Structure-Activity Relationships, and Drug Delivery”; *Current Medicinal Chemistry*; vol. 7; 2000; p. 249-271. cited by applicant

Vanhaerents et al.; “Epilepsy Emergencies: Status Epilepticus, Acute Repetitive Seizures, and Autoimmune Encephalitis”; *Continuum: Lifelong Learning in Neurology*; vol. 25; 2019; p. 454-476. cited by applicant

Vasileiou et al.; “Propofol: A review of its non-anaesthetic effects”; *European Journal of Pharmacology*; vol. 605; 2009; 8 pages. cited by applicant

Vasquez et al.; “Pediatric refractory and super-refractory status epilepticus”; *Seizure*; vol. 68; 2019; p. 62-71. cited by applicant

Veselis et al.; “Low-dose Propofol—induced Amnesia is Not due to a Failure of Encoding”; *Anesthesiology*; vol. 109; Aug. 2008; p. 213-224. cited by applicant

Wood et al.; “Propofol Infusion for the Treatment of Status Epilepticus”; *The Lancet*; Feb. 1988; p. 480-481. cited by applicant

“Epilepsy—A public health imperative”; World Health Organization; 2019; 171 pages. cited by applicant

H.F. Yanny; “Propofol infusions for status epilepticus”; *Anaesthesia*; vol. 43; 1988; p. 514. cited by applicant

Zack et al.; “National and State Estimates of the Numbers of Adults and Children with Active Epilepsy—United States, 2015”; *Morbidity and Mortality Weekly Report*; vol. 66 No. 31; Aug. 2017; p. 821-825. cited by applicant

Zhang et al.; “Systematic review and meta-analysis of propofol versus barbiturates for controlling refractory status epilepticus”; *BMC Neurology*; vol. 19; 2019; 11 pages. cited by applicant

Brodie; “Road to refractory epilepsy: The Glasgow story”; *Epilepsia*; vol. 54 Supplemental 2; 2013; p. 5-8. cited by applicant

Kasteleijn-Nolst Trenite; “Photosensitivity in epilepsy. Electrophysiological and clinical correlates”; *Acta Neurol Scan Suppl.*; vol. 125; 1989; p. 3-147. cited by applicant

Kasteleijn-Nolst Trenite et al.; “Preliminary assessment of the efficacy of Org 6370 in photosensitive epileptic patients: paradoxical enhancement of photosensitivity and provocation of myoclonic seizures”; *Epilpsia*; vol. 33(1); 1992; p. 135-141. cited by applicant

Kasteleijn-Nolst Trenite et al.; “Evaluation of carisbamate, a novel antiepileptic drug, in

photosensitive patients. An exploratory, placebo-controlled study”; Epilepsy Research; vol. 74; 2007; p. 193-200. cited by applicant

Nishikawa et al.; “Inhibitory Influence of GABA on Central Serotonergic Transmission. Raph6 Nuclei as the Neuroanatomical Site of the GABAergic Inhibition of Cerebral Serotonergic Neurons”; Brain Research; vol. 331; 1985; p. 91-103. cited by applicant

“Initial Investigational New Drug Application—(Fospropofol Disodium) For Oral Administration”; Investigator's Brochure; Version 1.0; Jul. 2020; 69 pages. cited by applicant

“US 21 CFR Part 58. Good Laboratory Practice for Nonclinical Laboratory Studies”; Available at https://www.ecfr.gov/cgi-bin/text-idx?SID=3be49f31878defa85f39ed3b84fcbe1&mc=true&node=se21.1.58_11&rgn=div8; 16 pages. cited by applicant

“Diprivan® (propofol) injectable emulsion, USP”; https://www.accessdata.fda.gov/drugsatfda_docs/label/2017/019627s066lbl.pdf; Fresenius-Kabi; Apr. 2017; 54 pages. cited by applicant

Sheehan et al.; “Comparative Validation of the S-STS, the ISSTPlus, and the C-SSRS for Assessing the Suicidal Thinking and Behavior FDA 2012 Suicidality Categories”; Innov Clin Neurosci.; vol. 11; 2014; p. 32-46. cited by applicant

Sheehan et al.; “Status Update on the Sheehan-Suicidality Tracking Scale (S-STS) 2014”; Innov Clin Neurosci.; vol. 11; 2014; p. 93-140. cited by applicant

“Bioanalytical Method Validation Guidance for Industry”; U.S. Dept. of Health and Human Services; May 2018; Biopharmaceutics; 41 pages. cited by applicant

“E6(R2) Good Clinical Practice: Integrated Addendum to ICH E6(R1) Guidance for Industry”; U.S. Dept. of Health and Human Services; Mar. 2018; 69 pages. cited by applicant

“Lusedra™ (fospropofol disodium) Injection, for intravenous use”; FDA printed label. Revised Oct. 2009. Available at https://www.accessdata.fda.gov/drugsatfda_docs/label/2010/022244s006lbl.pdf; 3 pages. cited by applicant

Aeschbacher et al.; “Propofol in rabbits. 2. Long-term anesthesia.”; Laboratory Animal Science; vol. 43(4); Aug. 1993; p. 328-335 (abstract only). cited by applicant

Glen et al.; “Interaction Studies and other investigations of the pharmacology of propofol (‘Diprivan’)”; Postgrad Med J.; vol. 61 Suppl. 3; 1985; p. 7-14 (abstract only). cited by applicant

Muir et al.; “Respiratory depression and apnea induced by propofol in dogs”; Am J Vet Res.; vol. 59(2); Feb. 1998; p. 157-161 (abstract only). cited by applicant

Ebbing; “General Chemistry—Second Edition”; Houghton Mifflin Company; 1987; p. 412. cited by applicant

European Patent Application No. 21925083.4; Extended Search Report; dated Jan. 21, 2025; 10 pages. cited by applicant

“IHS Classification ICHD-3”; <https://ichd-3.org/evolution-of-ihs-classification-1-3/>; International Headache Society; © 2021; accessed Feb. 9, 2021; 3 pages. cited by applicant

International Patent Application No. PCT/US2021/065225; Int'l Preliminary Report on Patentability; dated Aug. 17, 2023; 9 pages. cited by applicant

U.S. Appl. No. 17/066,957, filed Oct. 9, 2020, Rogawski et al. cited by applicant

U.S. Appl. No. 17/168,365, filed Feb. 5, 2021, Murphy et al. cited by applicant

U.S. Appl. No. 17/387,059, filed Jul. 28, 2021, Krill et al. cited by applicant

U.S. Appl. No. 17/465,966, filed Sep. 3, 2021, Krill et al. cited by applicant

U.S. Appl. No. 17/562,605, filed Dec. 27, 2021, Krill et al. cited by applicant

Feng et al., “Novel propofol derivatives and implications for anesthesia practice”, J. Anaesthesiol Clin Pharmacol., Jan.-Mar. 2017; vol. 33(1), p. 9-15. cited by applicant

Dhir; “Propofol in the treatment of refractory migraine headaches”; Expert Review of Neurotherapeutics; vol. 16 No. 9; 2016; p. 1007-1011. cited by applicant

Harris et al.; "Monitored anesthesia care (MAC) sedation: clinical utility of fospropofol"; Therapeutics and Clinical Risk Management; vol. 5; 2009; p. 949-959. cited by applicant

Ovesen et al.; "Intraluminal pH in the Stomach, Duodenum, and Proximal Jejunum in Normal Subjects and Patients With Exocrine Pancreatic Insufficiency"; Gastroenterology; vol. 90; 1986; p. 958-962. cited by applicant

Wozniak et al.; "Gastrointestinal delivery of propofol from fospropofol: its bioavailability and activity in rodents and human volunteers"; Journal of Translational Medicine; vol. 13; 2015; 13 pages. cited by applicant

Wilson et al.; "The abuse potential of propofol"; Clinical Toxicology; vol. 48; 2010; p. 165-170. cited by applicant

Ward et al.; "Use of intravenous propofol in the treatment of migraine headache"; EMA; 2013; p. 619. cited by applicant

Soleimanpour et al.; "Improvement of refractory migraine headache by propofol: case series"; Int'l Journal of Emergency Medicine; vol. 5; 2012; 4 pages. cited by applicant

Soleimanpour et al.; "Effectiveness of intravenous Dexamethasone versus Propofol for pain relief in the migraine headache: A prospective double blind randomized clinical trial"; BMC Neurology; vol. 12; 2012; 7 pages. cited by applicant

Simmonds et al.; "The Effect of Single-Dose Propofol Injection on Pain and Quality of Life in Chronic Daily Headache: A Randomized, Double-Blind, Controlled Trial"; Int'l Anesthesia Research Society; vol. 109 No. 6; Dec. 2009; p. 1972-1980. cited by applicant

Sheridan et al.; "Low-Dose Propofol for the Abortive Treatment of Pediatric Migraine in the Emergency Department"; Pediatric Emergency Care; vol. 28 No. 12; Dec. 2012; p. 1293-1296. cited by applicant

Sheridan et al.; "Low-Dose Propofol for Pediatric Migraine: a Prospective, Randomized Controlled Trial"; The Journal of Emergency Medicine; vol. 54 No. 5; 2018; p. 600-606. cited by applicant

Schneider et al.; "Propofol dependency after treatment of tension headache"; Addiction Biology; vol. 6; 2001; p. 263-265. cited by applicant

Sato et al.; "Low-dose intravenous propofol as a possible therapeutic option for acute confusional migraine"; American Journal of Emergency Medicine; vol. 35; 2017; 2 pages. cited by applicant

Reinsel et al.; "The P300 event-related potential during propofol sedation: a possible marker for amnesia?"; British Journal of Anesthesia; vol. 74; 1995; 674-680. cited by applicant

Razavi et al.; "Propofol and Alfentanil in Treatment of a Patient with Episodic Cluster Headache"; Anesth Pain Medicine; vol. 4(2); May 2014; 3 pages. cited by applicant

Mosier et al.; "Sedative Dosing of Propofol For Treatment of Migraine Headache in the Emergency Department: A Case Series"; Western Journal of Emergency Medicine; vol. 14 No. 6; Nov. 2013; p. 646-649. cited by applicant

Moshtaghion et al.; "The Efficacy of Propofol vs. Subcutaneous Sumatriptan for Treatment of Acute Migraine Headaches in the Emergency Department: A Double-Blinded Clinical Trial"; World Institute of Pain; 2014; 5 pages. cited by applicant

Mohseni et al.; "Propofol Alleviates Intractable Migraine Headache: A Case Report"; Anesthesiology and Pain Medicine; vol. 2(2); 2012; p. 94-96. cited by applicant

Mendes et al.; "Intravenous Propofol in the Treatment of Refractory Headache"; Headache; vol. 42; 2002; p. 638-641. cited by applicant

Long et al.; "Benign Headache Management in the Emergency Department"; The Journal of Emergency Medicine; vol. 54 No. 4; 2018; p. 458-468. cited by applicant

Krusz et al.; "Intravenous Propofol: Unique Effectiveness in Treating Intractable Migraine"; Headache; vol. 40; Mar. 2000; p. 224-230. cited by applicant

Ferrari et al.; "Oral triptans (serotonin 5-HT_{1B/1D} agonists) in acute migraine treatment: a meta-analysis of 53 trials"; The Lancet; vol. 358; Nov. 2001; p. 1668-1675. cited by applicant

Drummond-Lewis et al.; "Propofol: A New Treatment Strategy for Refractory Migraine

Headache”; Pain Medicine; vol. 3 No. 4; 2002; p. 366-369. cited by applicant

Dhir et al.; “Propofol hemisuccinate suppresses cortical spreading depression”; Neuroscience Letters; vol. 514; 2012; p. 67-70. cited by applicant

Dhir et al.; “Seizure Protection by Intrapulmonary Delivery of Propofol Hemisuccinate”; The Journal of Pharmacology and Experimental Therapeutics; vol. 336 No. 1; 2011; p. 215-222. cited by applicant

Bloomstone; “Propofol: A Novel Treatment for Breaking Migraine Headache”; Anesthesiology; vol. 106; 2007; p. 405-406. cited by applicant

Baker; “The Anticonvulsant Effects of Propofol and a Propofol Analog, 2,6-Diisopropyl-4-(1-Hydroxy-2,2,2-Trifluoroethyl)Phenol, in a 6 Hz Partial Seizure Model”; Int'l Anesthesia Research Society; vol. 112 No. 2; Feb. 2011; p. 340-344. cited by applicant

Kurt et al.; “Anxiolytic-Like Profile of Propofol, a General Anesthetic, in the Plus-Maze Test in Mice”; Polish Journal of Pharmacology; vol. 55; 2003; p. 973- 977. cited by applicant

Zacny et al.; “Propofol at Conscious Sedation Doses Produces Mild Analgesia to Cold Pressor-Induced Pain in Healthy Volunteers”; Journal of Clinical Anesthesia; vol. 8; 1996; p. 469-474. cited by applicant

Nishiyama et al.; “Intrathecal propofol has analgesic effects on inflammation-induced pain in rats”; Canadian Journal of Anesthesia; vol. 51(9); 2004; p. 899-904. cited by applicant

Bennett et al.; “Postoperative Infections Traced to Contamination of an Intravenous Anesthetic, Propofol”; The New England Journal of Medicine; vol. 333; 1995; p. 147-154. cited by applicant

Pytliak et al.; “Serotonin Receptors—From Molecular Biology to Clinical Applications”; Physiological Research; 2011; 19 pages. cited by applicant

Fechner et al.; “Pharmacokinetics and Clinical Pharmacodynamics of the New Propofol Prodrug GPI 15715 in Volunteers”; Anesthesiology; vol. 99 No. 2; Aug. 2003; p. 303-313. cited by applicant

Mahajan et al.; “Fospropofol”; Journal of Pharmacology and Pharmacotherapeutics; vol. 3 No. 3; Jul.-Sep. 2012; p. 293-296. cited by applicant

Borgeat et al.; “Subhypnotic Doses of Propofol Relieve Pruritus Associated with Liver Disease”; Gastroenterology; vol. 104; Jan. 1993; p. 244-247. cited by applicant

Kam et al.; “Pruritus—itching for a cause and relief?”; Anaesthesia; vol. 51; 1996; p. 1133-1138. cited by applicant

Pain et al.; “Effect of Nonsedative Doses of Propofol on an Innate Anxiogenic Situation in Rats”; Anesthesiology; vol. 90; 1999; p. 191-196. cited by applicant

Dwivedi et al., Evergreening: A deceptive device in patent rights, Technology in Society, 32: 324-330, 2010. doi: 10.1016/j.tecyhsoc.2010.10.009. cited by applicant

Feldman, Robin, Understanding, ‘Evergreening’: Making Minor Modifications of Existing Medications to Extend Protections, Health Affairs, 41(6): 801-804, 2022. cited by applicant

Primary Examiner: Shiao; Rei Tsang

Attorney, Agent or Firm: BakerHostetler

Background/Summary

CROSS-REFERENCE TO RELATED APPLICATION (1) This application is a divisional of U.S. application Ser. No. 17/466,016, filed Sep. 3, 2021, which is a continuation of U.S. application Ser. No. 17/217,656, filed Mar. 30, 2021, the entirety of which is incorporated by reference herein.

TECHNICAL FIELD

(1) The present disclosure pertains to pharmaceutical compositions for oral administration of fospropofol, or pharmaceutically acceptable salts of fospropofol, as well as methods for oral administration of fospropofol.

BACKGROUND

(2) Propofol (2,6-diisopropylphenol) is approved for use as an intravenous, short-acting anesthetic agent for inducing and maintaining anesthesia. Fospropofol disodium, a water-soluble, phosphono-O-methyl prodrug of propofol, was approved for use as an intravenous sedative, but its marketing has been discontinued.

(3) While intravenous administration is useful for anesthesia applications, oral administration of fospropofol would be desirable for other uses. Oral dosing, however, requires a dosage form having adequate bioavailability with minimal subject-to-subject variability. There is a need for dosage forms of fospropofol that are orally bioavailable with minimal inter-subject variability.

SUMMARY

(4) The present disclosure provides pharmaceutical dosage forms for oral administration comprising fospropofol or a pharmaceutically acceptable salt of fospropofol, and a pharmaceutically acceptable acid.

(5) The disclosure also provides methods of orally administering fospropofol, or a pharmaceutically acceptable salt thereof, to a subject in need thereof, the method comprising orally co-administering fospropofol, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable acid, to the subject.

Description

BRIEF DESCRIPTION OF THE DRAWINGS

(1) FIG. 1 depicts the dissolution profiles of 600 mg fospropofol disodium salt acidified tablets in 300 mL of 0.1 N HCl at 37° C.

(2) FIG. 2 depicts the propofol plasma profile comparison of 600 mg fospropofol disodium tablet with ascorbic acid acidifier (Composition A2) without pentagastrin pretreatment; 600 mg fospropofol disodium tablet with tartaric acid acidifier (Composition A3) without pentagastrin pretreatment; and a 1300 mg fospropofol disodium aqueous solution (30 mg/mL) without pentagastrin pretreatment which propofol plasma concentration is normalized to 600 mg fospropofol disodium.

(3) FIG. 3 depicts the propofol plasma profile comparison of 600 mg fospropofol disodium tablet with ascorbic acid acidifier (Composition A2) with pentagastrin pretreatment; and 600 mg fospropofol disodium tablet control (i.e., no acidifier) with pentagastrin pretreatment.

(4) FIG. 4 depicts the propofol plasma profile comparison of 600 mg fospropofol disodium tablet with tartaric acid acidifier (Composition A3) without pentagastrin pretreatment; and 600 mg fospropofol disodium tablet control (i.e., no acidifier) with pentagastrin pretreatment.

DETAILED DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

(5) The present disclosure may be understood more readily by reference to the following detailed description of desired embodiments and the examples included therein. In the following specification and the claims that follow, reference will be made to a number of terms which have the following meanings.

(6) Unless indicated to the contrary, the numerical values should be understood to include numerical values which are the same when reduced to the same number of significant figures and numerical values which differ from the stated value by less than the experimental error of conventional measurement technique of the type described in the present application to determine the value.

- (7) All ranges disclosed herein are inclusive of the recited endpoint and independently combinable (for example, the range of “from 2 to 10” is inclusive of the endpoints, 2 and 10, and all the intermediate values). The endpoints of the ranges and any values disclosed herein are not limited to the precise range or value; they are sufficiently imprecise to include values approximating these ranges and/or values.
- (8) As used herein, approximating language may be applied to modify any quantitative representation that may vary without resulting in a change in the basic function to which it is related. Accordingly, a value modified by a term or terms, such as “about” and “substantially,” may not be limited to the precise value specified, in some cases. In at least some instances, the approximating language may correspond to the precision of an instrument for measuring the value. The modifier “about” should also be considered as disclosing the range defined by the absolute values of the two endpoints. For example, the expression “from about 2 to about 4” also discloses the range “from 2 to 4.” The term “about” may refer to plus or minus 10% of the indicated number. For example, “about 10%” may indicate a range of 9% to 11%, and “about 1” may mean from 0.9-1.1. Other meanings of “about” may be apparent from the context, such as rounding off, so, for example “about 1” may also mean from 0.5 to 1.4.
- (9) In the present disclosure the singular forms “a,” “an,” and “the” include the plural reference, and reference to a particular numerical value includes at least that particular value, unless the context clearly indicates otherwise. Thus, for example, a reference to “a compound” is a reference to one or more of such compounds and equivalents thereof known to those skilled in the art, and so forth. The term “plurality”, as used herein, means more than one.
- (10) The term “subject,” is used herein to refer to an animal, for example a human, to whom treatment, including prophylactic treatment, with the pharmaceutical compositions or methods according to the present invention, is provided. The term “subject” as used herein refers to human and non-human animals.
- (11) As used herein, the term “treating” means reducing or eliminating the signs or symptoms of the condition for which fospropofol is being administered.
- (12) In some aspects, the disclosure is directed to pharmaceutical dosage forms for oral administration comprising fospropofol or a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable acid.
- (13) In some embodiments, the pharmaceutical dosage forms of the disclosure comprise fospropofol:
- (14) ##STR00001##
- (15) In other embodiments, the pharmaceutical dosage forms of the disclosure comprise a pharmaceutically acceptable salt of fospropofol. As used herein, “pharmaceutically acceptable salt of fospropofol” refers to a salt of fospropofol that is pharmaceutically acceptable and that possesses the desired pharmacologic activity. Such salts are generally non-toxic, and may be inorganic or organic base addition salts. Specifically, such salts include: salts formed when at least one acidic proton present in fospropofol either is replaced by at least one metal ion, e.g., an alkali metal ion, an alkaline earth ion, or an aluminum ion; or by an organic base such as aliphatic, alicyclic, or aromatic organic amines, such as ammonia, methylamine, dimethylamine, diethylamine, picoline, ethanolamine, diethanolamine, triethanolamine, ethylenediamine, lysine, arginine, ornithine, choline, N,N'-dibenzylethylene-diamine, chloroprocaine, diethanolamine, procaine, N-benzylphenethylamine, N-methylglucamine piperazine, tris(hydroxymethyl)-aminomethane, tetramethylammonium hydroxide, and the like. Examples of pharmaceutically acceptable salts of fospropofol include monosodium, monopotassium, disodium, dipotassium salts, diethylamine, t-butyl amine, ethylene diamine, benzathine, piperazine, ethanolamine, diethanolamine, ammonium, tromethamine, benethamine, histidine, calcium, magnesium, and zinc salts.
- (16) In some embodiments, the pharmaceutical dosage forms of the disclosure comprise fospropofol disodium:

(17) ##STR00002##

(18) The pharmaceutical dosage forms of the disclosure including a pharmaceutically acceptable salt of fospropofol comprise a pharmaceutically acceptable acid.

(19) Pharmaceutically acceptable acids are known in the art. Exemplary pharmaceutically acceptable acids include all applicable stereoisomers including diastereomers, enantiomers and mixtures thereof, for example, 1-hydroxy-2-naphthoic acid, 2,2-dichloroacetic acid, 2-hydroxethanesulfonic acid, 4-acetamidobenzoic acid, 4-aminosalicylic acid, acetic acid, aceturic acid, Acid hydrolyzed proteins, Acid Modified Starch, Aconitic Acid, adipic acid, alginic acid, α -oxo-glutaric acid, benzenesulfonic acid, benzoic acid, butyric acid, camphor-10-sulfonic acid, camphoric acid, capric acid, caproic acid, caprylic acid, carbonic acid, Cholic acid, cinnamic acid, citric acid, cyclamic acid, D(-)-Lactic acid, Desoxycholic acid, D-glucaric acid, D-glucoheptonic acid, D-glucuronic acid, Di(tert-butyl)naphthalenedisulfonic acid, Di(tert-butyl)naphthalenesulfonic acid, DL-lactic acid, DL-mandelic acid, DL-tartaric acid, tartaric acid, dodecylsulfuric acid, Erythorbic acid (D-isoascorbic acid), ethane-1,2-disulfonic acid, ethanesulfonic acid, formic acid, fumaric acid, galactaric acid, gentisic acid, glutaric acid, glycerophosphoric acid, Glycocholic acid, glycolic acid, hexanedioic acid, hippuric acid, hydrobromic acid, Hydrochloric acid, Iron naphthenate, iron salts, iron salts, isobutyric acid, L(+)-lactic acid, L(+)-potassium acid tartrate, tartaric acid, L(+)-tartaric acid, Lactic acid, lactobionic acid, L-ascorbic acid, ascorbic acid, L-aspartic acid, lauric acid, L-glutamic acid, L-Glutamic acid hydrochloride, Linoleic acid, L-Malic acid, L-pyroglutamic acid, L-tartaric acid, maleic acid, malic acid, malonic acid, methanesulfonic acid, monobasic potassium phosphate, monobasic sodium phosphate, naphthalene-1,5-disulfonic acid, naphthalene-2-sulfonic acid, naphthenic acids, Niacin (nicotinic acid), nicotinic acid, nitric acid, octanoic acid, oleic acid, orotic acid, oxalic acid, palmitic acid, pamoic acid, Pectin low acid, Pectinic acid, phosphoric acid, propanoic acid, Propionic acid, p-toluenesulfonic acid, pyruvic acid, saccharin, salicylic acid, sebacic acid, Sodium acid pyrophosphate, Sodium aluminum phosphate, sodium metabisulfite, Sorbic acid, stearic acid, succinic acid, sulfuric acid, tall oil fatty acids, Tannic acid (hydrolyzable gallotannins), Taurocholic acid, thiocyanic acid, Thiodipropionic acid, trifluoroacetic acid, undec-10-enoic acid, orange juice, apple juice, grapefruit juice, as well as combinations thereof. As used herein, a “pharmaceutically acceptable acid” preferably refers to an acid that is on the FDA published inactive ingredient database (IID) for approved drug products or the generally recognized as safe (GRAS) database for use in food. In some embodiments, the pharmaceutically acceptable acid is a polyacid such as hyaluronic acid, polyacrylic acid (e.g., Carbomers), polyaspartic acid, polyglutamic acid or mixtures thereof.

(20) In some embodiments, the pharmaceutically acceptable acid is ascorbic acid, citric acid, malic acid, tartaric acid, succinic acid, fumaric acid, maleic acid, lactic acid, monobasic sodium phosphate, monobasic potassium phosphate, or sodium metabisulfite.

(21) In other embodiments, the pharmaceutically acceptable acid is ascorbic acid, citric acid, malic acid, or tartaric acid, or all applicable stereoisomers including diastereomers, enantiomers and mixtures thereof.

(22) In some embodiments, the pharmaceutically acceptable acid is ascorbic acid. As used herein, the term “ascorbic acid” refers to DL-ascorbic acid, L-ascorbic acid, D-ascorbic acid, or mixtures thereof. In some embodiments, the ascorbic acid is DL-ascorbic acid. In other embodiments, the ascorbic acid is L-ascorbic acid. In other embodiments, the ascorbic acid is D-ascorbic acid.

(23) In some embodiments, the pharmaceutically acceptable acid is tartaric acid. As used herein, the term “tartaric acid” refers to DL- tartaric acid, L- tartaric acid, D- tartaric acid, meso-tartaric acid, or mixtures thereof. In some embodiments, the tartaric acid is DL-tartaric acid. In other embodiments, the tartaric acid is L- tartaric acid. In other embodiments, the tartaric acid is D-tartaric acid.

(24) In some embodiments, the pharmaceutically acceptable acid is malic acid. As used herein, the

term “malic acid” refers to DL- malic acid, L- malic acid, D- malic acid, or mixtures thereof. In some embodiments, the malic acid is DL- malic acid. In other embodiments, the malic acid is L- malic acid. In other embodiments, the malic acid is D-malic acid.

(25) In some embodiments, the pharmaceutically acceptable acid is citric acid.

(26) In some embodiments, the pharmaceutically acceptable acid is fumaric acid, i.e., the trans isomer of butenedioic acid. In some embodiments, the pharmaceutically acceptable acid is maleic acid, i.e., the cis isomer of butenedioic acid. In some embodiments, the pharmaceutically acceptable acid is a mixture of the cis isomer of butenedioic acid and the trans isomer of butenedioic acid. An exemplary composition comprising fumaric acid comprises 384.2 mg fospropofol disodium hydrate (equivalent to 300 mg fospropofol disodium), 28.2 mg of polyplasdone XL, 2.8 mg of magnesium stearate, and 150.0 mg of fumaric acid. Another exemplary composition comprising fumaric acid comprises 127.8 mg fospropofol disodium hydrate (equivalent to 100 mg fospropofol disodium), 14.7 mg of polyplasdone XL, 1.46 mg of magnesium stearate, and 150.0 mg of fumaric acid.

(27) In some embodiments in which the pharmaceutical dosage forms of the disclosure comprise fospropofol in acid form, i.e.,

(28) ##STR00003##

(29) the dosage forms do not include an additional pharmaceutically acceptable acid because in such embodiments, the fospropofol acid itself provides the acidity to achieve oral bioavailability and/or reduced inter-subject variability.

(30) In some aspects, the pharmaceutical dosage forms of the disclosure further comprise a pharmaceutically acceptable excipient. In such aspects, the pharmaceutically acceptable excipient is present in addition to the pharmaceutically acceptable acid.

(31) A “pharmaceutically acceptable excipient” refers to a substance that is non-toxic, biologically tolerable, and otherwise biologically suitable for administration to a subject, such as an inert substance, added to a pharmacological composition or otherwise used as a vehicle, carrier, diluent, or release modifier to facilitate administration of an agent and that is compatible therewith.

(32) In some embodiments, the pharmaceutically acceptable excipient may be water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents, starches, sugars, micro-crystalline cellulose, surfactants, polymers, diluents, granulating agents, lubricants, binders, fillers, and disintegrants.

(33) Binders suitable for use in pharmaceutical compositions and dosage forms include, but are not limited to, corn starch, potato starch, or other starches, gelatin, natural and synthetic gums such as acacia, sodium alginate, alginic acid, other alginates, powdered tragacanth, guar gum, cellulose and its derivatives (e.g., ethyl cellulose, cellulose acetate, carboxymethyl cellulose calcium, sodium carboxymethyl cellulose), polyvinyl pyrrolidone, methyl cellulose, pre-gelatinized starch, dicalcium phosphate, hydroxypropyl methyl cellulose, microcrystalline cellulose, and mixtures thereof.

(34) Fillers for use in the pharmaceutical compositions and dosage forms disclosed herein include, but are not limited to, talc, calcium carbonate (e.g., granules or powder), microcrystalline cellulose, powdered cellulose, dextrates, kaolin, mannitol, silicic acid, sorbitol, starch, dicalcium phosphate, pre-gelatinized starch, and mixtures thereof.

(35) Disintegrants that can be used to form pharmaceutical compositions and dosage forms of the invention include, but are not limited to, agar-agar, alginic acid, calcium carbonate, microcrystalline cellulose, croscarmellose sodium, crospovidone, polacrillin potassium, sodium starch glycolate, potato or tapioca starch, other starches, pre-gelatinized starch, other starches, clays, other algin, other celluloses, gums or mixtures thereof.

(36) Lubricants which can be used to form pharmaceutical compositions and dosage forms of the invention include, but are not limited to, calcium stearate, magnesium stearate, mineral oil, light mineral oil, glycerin, sorbitol, mannitol, polyethylene glycol, other glycols, stearic acid, sodium

lauryl sulfate, talc, hydrogenated vegetable oil (e.g., peanut oil, cottonseed oil, sunflower oil, sesame oil, olive oil, corn oil, and soybean oil), zinc stearate, ethyl oleate, ethyl laurate, agar, or mixtures thereof.

(37) In some embodiments, the solid pharmaceutical dosage form is uncoated or coated to delay disintegration and absorption in the gastrointestinal tract. For example, a time delay material such as glyceryl monostearate or glyceryl distearate can be employed.

(38) Surfactants which can be used to form pharmaceutical compositions and dosage forms of the invention include, but are not limited to, hydrophilic surfactants, lipophilic surfactants, ionic surfactants, and mixtures thereof.

(39) Suitable ionic surfactants include, but are not limited to, alkylammonium salts; fusidic acid salts; fatty acid derivatives of amino acids, oligopeptides, and polypeptides; glyceride derivatives of amino acids, oligopeptides, and polypeptides; lecithins and hydrogenated lecithins; lysolecithins and hydrogenated lysolecithins; phospholipids and derivatives thereof; lysophospholipids and derivatives thereof; carnitine fatty acid ester salts; salts of alkylsulfates; fatty acid salts; sodium docusate; acyl lactylates; mono- and di-acetylated tartaric acid esters of mono- and di-glycerides; succinylated mono- and di-glycerides; citric acid esters of mono- and di-glycerides; and mixtures thereof, lecithins, lysolecithin, phospholipids, lysophospholipids and derivatives thereof; carnitine fatty acid ester salts; salts of alkylsulfates; fatty acid salts; sodium docusate; acylactylates; mono- and di-acetylated tartaric acid esters of mono- and di-glycerides; succinylated mono- and di-glycerides; citric acid esters of mono- and di-glycerides; and mixtures thereof.

(40) Ionic surfactants may be the ionized forms of lecithin, lysolecithin, phosphatidylcholine, phosphatidylethanolamine, phosphatidylglycerol, phosphatidic acid, phosphatidylserine, lysophosphatidylcholine, lysophosphatidylethanolamine, lysophosphatidylglycerol, lysophosphatidic acid, lysophosphatidylserine, PEG-phosphatidylethanolamine, PVP-phosphatidylethanolamine, lactic esters of fatty acids, stearyl-2-lactylate, stearyl lactylate, succinylated monoglycerides, mono/diacetylated tartaric acid esters of mono/diglycerides, citric acid esters of mono/diglycerides, choly sarcosine, caproate, caprylate, caprate, laurate, myristate, palmitate, oleate, ricinoleate, linoleate, linolenate, stearate, lauryl sulfate, teracecyl sulfate, docusate, lauroyl carnitines, palmitoyl carnitines, myristoyl carnitines, and salts and mixtures thereof.

(41) Hydrophilic non-ionic surfactants may include, but are not limited to, alkylglucosides; alkylmaltosides; alkylthiogluconates; lauryl macroglycerides; polyoxyalkylene alkyl ethers such as polyethylene glycol alkyl ethers; polyoxyalkylene alkylphenols such as polyethylene glycol alkyl phenols; polyoxyalkylene alkyl phenol fatty acid esters such as polyethylene glycol fatty acids monoesters and polyethylene glycol fatty acids diesters; polyethylene glycol glycerol fatty acid esters; polyglycerol fatty acid esters; polyoxyalkylene sorbitan fatty acid esters such as polyethylene glycol sorbitan fatty acid esters; hydrophilic transesterification products of a polyol with at least one member of the group consisting of glycerides, vegetable oils, hydrogenated vegetable oils, fatty acids, and sterols; polyoxyethylene sterols, derivatives, and analogues thereof; polyoxyethylated vitamins and derivatives thereof; polyoxyethylene-polyoxypropylene block copolymers; and mixtures thereof, polyethylene glycol sorbitan fatty acid esters and hydrophilic transesterification products of a polyol with at least one member of the group consisting of triglycerides, vegetable oils, and hydrogenated vegetable oils. The polyol may be glycerol, ethylene glycol, polyethylene glycol, sorbitol, propylene glycol, pentaerythritol, or a saccharide.

(42) Other hydrophilic-non-ionic surfactants include, without limitation, PEG- 10 laurate, PEG- 12 laurate, PEG-20 laurate, PEG-32 laurate, PEG-32 dilaurate, PEG- 12 oleate, PEG- 15 oleate, PEG-20 oleate, PEG-20 dioleate, PEG-32 oleate, PEG-200 oleate, PEG-400 oleate, PEG- 15 stearate, PEG-32 distearate, PEG-40 stearate, PEG- 100 stearate, PEG-20 dilaurate, PEG-25 glyceryl trioleate, PEG-32 dioleate, PEG-20 glyceryl laurate, PEG-30 glyceryl laurate, PEG-20 glyceryl stearate, PEG-20 glyceryl oleate, PEG-30 glyceryl oleate, PEG-30 glyceryl laurate, PEG-40

glyceryl laurate, PEG-40 palm kernel oil, PEG-50 hydrogenated castor oil, PEG-40 castor oil, PEG-35 castor oil, PEG-60 castor oil, PEG-40 hydrogenated castor oil, PEG-60 hydrogenated castor oil, PEG-60 corn oil, PEG-6 caprate/caprylate glycerides, PEG-8 caprate/caprylate glycerides, polyglyceryl-10 laurate, PEG-30 cholesterol, PEG-25 phyto sterol, PEG-30 soya sterol, PEG-20 trioleate, PEG-40 sorbitan oleate, PEG-80 sorbitan laurate, polysorbate 20, polysorbate 80, POE-9 lauryl ether, POE-23 lauryl ether, POE-10 oleyl ether, POE-20 oleyl ether, POE-20 stearyl ether, tocopheryl PEG- 100 succinate, PEG-24 cholesterol, polyglyceryl-10-oleate, Tween 40, Tween 60, sucrose monostearate, sucrose mono laurate, sucrose monopalmitate, PEG 10-100 nonyl phenol series, PEG 15-100 octyl phenol series, and poloxamers.

(43) Suitable lipophilic surfactants include, by way of example only: fatty alcohols; glycerol fatty acid esters; acetylated glycerol fatty acid esters; lower alcohol fatty acids esters; propylene glycol fatty acid esters; sorbitan fatty acid esters; polyethylene glycol sorbitan fatty acid esters; sterols and sterol derivatives; polyoxyethylated sterols and sterol derivatives; polyethylene glycol alkyl ethers; sugar esters; sugar ethers; lactic acid derivatives of mono- and di-glycerides; hydrophobic transesterification products of a polyol with at least one member of the group consisting of glycerides, vegetable oils, hydrogenated vegetable oils, fatty acids and sterols; oil-soluble vitamins/vitamin derivatives; and mixtures thereof.

(44) Solubilizers include, but are not limited to, the following: alcohols and polyols, such as ethanol, isopropanol, butanol, benzyl alcohol, ethylene glycol, propylene glycol, butanediols and isomers thereof, glycerol, pentaerythritol, sorbitol, mannitol, transcitol, dimethyl isosorbide, polyethylene glycol, polypropylene glycol, polyvinylalcohol, hydroxypropyl methylcellulose and other cellulose derivatives, cyclodextrins and cyclodextrin derivatives; ethers of polyethylene glycols having an average molecular weight of about 200 to about 6000, such as tetrahydrofurfuryl alcohol PEG ether (glycofuro) or methoxy PEG; amides and other nitrogen-containing compounds such as 2-pyrrolidone, 2-piperidone, ϵ -caprolactam, N-alkylpyrrolidone, N-hydroxyalkylpyrrolidone, N-alkylpiperidone, N-alkylcaprolactam, dimethylacetamide and polyvinylpyrrolidone; esters such as ethyl propionate, tributylcitrate, acetyl triethylcitrate, acetyl tributyl citrate, triethylcitrate, ethyl oleate, ethyl caprylate, ethyl butyrate, triacetin, propylene glycol monoacetate, propylene glycol diacetate, ϵ -caprolactone and isomers thereof, δ -valerolactone and isomers thereof, β -butyrolactone and isomers thereof and other solubilizers known in the art, such as dimethyl acetamide, dimethyl isosorbide, N-methyl pyrrolidones, mono-octanoin, diethylene glycol monoethyl ether, and water.

(45) Release modifiers may include coatings or matrix materials.

(46) Release modifying coatings include but are not limited to polymer coating materials, such as cellulose acetate phthalate, cellulose acetate trimaletate, hydroxy propyl methylcellulose phthalate, polyvinyl acetate phthalate, ammonio methacrylate copolymers such as those sold under the Trade Mark Eudragit® RS and RL, poly acrylic acid and poly acrylate and methacrylate copolymers such as those sold under the Trade Mark Eudragite S and L, polyvinyl acetaldiethylamino acetate, hydroxypropyl methylcellulose acetate succinate, shellac; hydrogels and gel-forming materials, such as carboxyvinyl polymers, sodium alginate, sodium carmellose, calcium carmellose, sodium carboxymethyl starch, poly vinyl alcohol, hydroxyethyl cellulose, methyl cellulose, gelatin, starch, and cellulose based cross-linked polymers—in which the degree of crosslinking is low so as to facilitate adsorption of water and expansion of the polymer matrix, hydroxypropyl cellulose, hydroxypropyl methylcellulose, polyvinylpyrrolidone, crosslinked starch, microcrystalline cellulose, chitin, aminoacryl-methacrylate copolymer (Eudragit® RS-PM, Rohm & Haas), pullulan, collagen, casein, agar, gum arabic, sodium carboxymethyl cellulose, (swellable hydrophilic polymers) poly(hydroxyalkyl methacrylate) (m. wt. .sup.~5 k-5,000 k), polyvinylpyrrolidone (m. wt. .sup.~10 k-360 k), anionic and cationic hydrogels, polyvinyl alcohol having a low acetate residual, a swellable mixture of agar and carboxymethyl cellulose, copolymers of maleic anhydride and styrene, ethylene, propylene or isobutylene, pectin (m. wt. .sup.~30 k-300

k), polysaccharides such as agar, acacia, karaya, tragacanth, algin and guar, polyacrylamides, Polyox® polyethylene oxides (m. wt. .sup.~100 k-5,000 k), AquaKeep® acrylate polymers, diesters of polyglucan, crosslinked polyvinyl alcohol and poly N-vinyl-2-pyrrolidone, sodium starch glucolate (e.g. Explotab®; Edward Mandell C. Ltd.); hydrophilic polymers such as polysaccharides, methyl cellulose, sodium or calcium carboxymethyl cellulose, hydroxypropyl methyl cellulose, hydroxypropyl cellulose, hydroxyethyl cellulose, nitro cellulose, carboxymethyl cellulose, cellulose ethers, polyethylene oxides (e.g. Polyox®, Union Carbide), methyl ethyl cellulose, ethylhydroxy ethylcellulose, cellulose acetate, cellulose butyrate, cellulose propionate, gelatin, collagen, starch, maltodextrin, pullulan, polyvinyl pyrrolidone, polyvinyl alcohol, polyvinyl acetate, glycerol fatty acid esters, polyacrylamide, polyacrylic acid, copolymers of methacrylic acid or methacrylic acid (e.g. Eudragit®, Rohm and Haas), other acrylic acid derivatives, sorbitan esters, natural gums, lecithins, pectin, alginates, ammonia alginate, sodium, calcium, potassium alginates, propylene glycol alginate, agar, and gums such as arabic, karaya, locust bean, tragacanth, carrageens, guar, xanthan, scleroglucan and mixtures and blends thereof. As will be appreciated by the person skilled in the art, excipients such as plasticisers, lubricants, solvents and the like may be added to the coating. Suitable plasticisers include for example acetylated monoglycerides; butyl phthalyl butyl glycolate; dibutyl tartrate; diethyl phthalate; dimethyl phthalate; ethyl phthalyl ethyl glycolate; glycerin; propylene glycol; triacetin; citrate; tripropioin; diacetin; dibutyl phthalate; acetyl monoglyceride; polyethylene glycols; castor oil; triethyl citrate; polyhydric alcohols, glycerol, acetate esters, glycerol triacetate, acetyl triethyl citrate, dibenzyl phthalate, dihexyl phthalate, butyl octyl phthalate, diisononyl phthalate, butyl octyl phthalate, dioctyl azelate, epoxidised tallate, triisooctyl trimellitate, diethylhexyl phthalate, di-n-octyl phthalate, di-i-octyl phthalate, di-i-decyl phthalate, di-n-undecyl phthalate, di-n-tridecyl phthalate, tri-2-ethylhexyl trimellitate, di-2-ethylhexyl adipate, di-2-ethylhexyl sebacate, di-2-ethylhexyl azelate, dibutyl sebacate.

(47) Release-modifying matrix materials include hydrophilic polymers, hydrophobic polymers and mixtures thereof, dicalcium phosphate, microcrystalline cellulose, sodium carboxymethylcellulose, hydroxyalkylcelluloses such as hydroxypropylmethylcellulose and hydroxypropylcellulose, polyethylene oxide, alkylcelluloses such as methylcellulose and ethylcellulose, polyethylene glycol, polyvinylpyrrolidone, cellulose acetate, cellulose acetate butyrate, cellulose acetate phthalate, cellulose acetate trimellitate, polyvinylacetate phthalate, polyalkylmethacrylates, polyvinyl acetate, Poly(-hydroxy ethyl methacrylate), Poly(N-vinyl pyrrolidone), Poly(methyl methacrylate), Poly(vinyl alcohol), Poly(acrylic acid), Polyacrylamide, Poly(ethylene-co-vinyl acetate), Poly(ethylene glycol), Poly(methacrylic acid), Polylactides (PLA), Polyglycolides (PGA), Poly(lactide-co-glycolides) (PLGA), Polyanhydrides, and Polyorthoesters, and mixture thereof.

(48) In some embodiments, the fospropofol (or pharmaceutically acceptable salt thereof) and the pharmaceutically acceptable acid can be present in the same unit dosage form. In other embodiments, all or a portion of the pharmaceutically acceptable acid can be administered separately from the unit dosage form comprising the fospropofol (or pharmaceutically acceptable salt thereof).

(49) In some aspects, the pharmaceutical dosage forms of the disclosure are those that when dissolved in water, contain an amount of pharmaceutically acceptable acid necessary to drive the ionic equilibrium of the aqueous solution towards a pH lower than 7 and preferably <4.5.

(50) In some aspects, the pharmaceutical dosage forms of the disclosure are those wherein dissolving an amount of the dosage form containing a therapeutically effective amount of fospropofol, or a pharmaceutically acceptable salt thereof, in 300 mL of water at 25° C., results in a solution having a pH of less than or equal to 6.3, such as, for example, a pH of less than or equal to 6.3, 6.2, 6.1, 6.0, 5.9, 5.8, 5.7, 5.6, 5.5, 5.4, 5.3, 5.2, 5.1, 5.0, 4.9, 4.8, 4.7, 4.6, 4.5, 4.4, 4.3, 4.2, 4.1, 4.0, 3.9, 3.8, 3.7, 3.6, 3.5, 3.4, 3.3, 3.2, 3.1, 3.0, 2.9, 2.8, 2.7, 2.6, 2.5, 2.4, 2.3, 2.2, 2.1, 2.0, 1.9, 1.8, 1.7, 1.6, 1.5, 1.4, 1.3, 1.2, 1.1, or 1.0.

(51) The term “therapeutically effective amount”, as used herein, refers to an amount sufficient to reduce or eliminate the signs or symptoms of the condition for which fospropofol is being administered. The therapeutically effective amount may be contained in a single dosage form, or may be the cumulative amount contained in multiple dosage forms.

(52) In some embodiments, a therapeutically effective amount of fospropofol, or a pharmaceutically acceptable salt of fospropofol, is 10-4800 mg (on a fospropofol basis), for example, an amount that is about (i.e., the specified number $\pm$ 10%) any one of 10 mg, 20 mg, 30 mg, 40 mg, 50 mg, 60 mg, 70 mg, 80 mg, 90 mg, 100 mg, 150 mg, 200 mg, 250 mg, 300 mg, 350 mg, 400 mg, 450 mg, 500 mg, 550 mg, 600 mg, 650 mg, 700 mg, 750 mg, 800 mg, 850 mg, 900 mg, 950 mg, 1000 mg, 1050 mg, 1100 mg, 1150 mg, 1200 mg, 1250 mg, 1300 mg, 1350 mg, 1400 mg, 1450 mg, 1500 mg, 1550 mg, 1600 mg, 1650 mg, 1700 mg, 1750 mg, 1800 mg, 1850 mg, 1900 mg, 1950 mg, 2000 mg, 2050 mg, 2100 mg, 2150 mg, 2200 mg, 2250 mg, 2300 mg, 2350 mg, 2400 mg, 2450 mg, 2500 mg, 2550 mg, 2600 mg, 2650 mg, 2700 mg, 2750 mg, 2800 mg, 2850 mg, 2900 mg, 2950 mg, 3000 mg, 3050 mg, 3100 mg, 3150 mg, 3200 mg, 3250 mg, 3300 mg, 3350 mg, 3400 mg, 3450 mg, 3500 mg, 3550 mg, 3600 mg, 3650 mg, 3700 mg, 3750 mg, 3800 mg, 3850 mg, 3900 mg, 3950 mg, 4000 mg, 4050 mg, 4100 mg, 4150 mg, 4200 mg, 4250 mg, 4300 mg, 4350 mg, 4400 mg, 4450 mg, 4500 mg, 4550 mg, 4600 mg, 4650 mg, 4700 mg, 4750 mg, or 4800 mg.

(53) In other embodiments, a therapeutically effective amount of fospropofol, or a pharmaceutically acceptable salt of fospropofol, is about 1 mg/kg to about 80 mg/kg (on a fospropofol basis), for example, an amount that is about (i.e., the specified number $\pm$ 10%) any one of 1 mg/kg, 2 mg/kg, 3 mg/kg, 4 mg/kg, 5 mg/kg, 6 mg/kg, 7 mg/kg, 8 mg/kg, 9 mg/kg, 10 mg/kg, 11 mg/kg, 12 mg/kg, 13 mg/kg, 14 mg/kg, 15 mg/kg, 16 mg/kg, 17 mg/kg, 18 mg/kg, 19 mg/kg, 20 mg/kg, 21 mg/kg, 22 mg/kg, 23 mg/kg, 24 mg/kg, 25 mg/kg, 26 mg/kg, 27 mg/kg, 28 mg/kg, 29 mg/kg, 30 mg/kg, 31 mg/kg, 32 mg/kg, 33 mg/kg, 34 mg/kg, 35 mg/kg, 36 mg/kg, 37 mg/kg, 38 mg/kg, 39 mg/kg, 40 mg/kg, 41 mg/kg, 42 mg/kg, 43 mg/kg, 44 mg/kg, 45 mg/kg, 46 mg/kg, 47 mg/kg, 48 mg/kg, 49 mg/kg, 50 mg/kg, 51 mg/kg, 52 mg/kg, 53 mg/kg, 54 mg/kg, 55 mg/kg, 56 mg/kg, 57 mg/kg, 58 mg/kg, 59 mg/kg, 60 mg/kg, 61 mg/kg, 62 mg/kg, 63 mg/kg, 64 mg/kg, 65 mg/kg, 66 mg/kg, 67 mg/kg, 68 mg/kg, 69 mg/kg, 70 mg/kg, 71 mg/kg, 72 mg/kg, 73 mg/kg, 74 mg/kg, 75 mg/kg, 76 mg/kg, 77 mg/kg, 78 mg/kg, 79 mg/kg, or 80 mg/kg.

(54) In some embodiments, the pharmaceutical dosage forms of the disclosure are those wherein dissolving an amount of the dosage form containing a therapeutically effective amount of fospropofol, or a pharmaceutically acceptable salt thereof, in 300 mL of water at 25° C., results in a solution having a pH of less than or equal to 4.5.

(55) In other embodiments, the pharmaceutical dosage forms of the disclosure are those wherein dissolving an amount of the dosage form containing a therapeutically effective amount of fospropofol, or a pharmaceutically acceptable salt thereof, in 300 mL of water at 25° C., results in a solution having a pH of less than or equal to 4.2.

(56) In other embodiments, the pharmaceutical dosage forms of the disclosure are those wherein dissolving an amount of the dosage form containing a therapeutically effective amount of fospropofol, or a pharmaceutically acceptable salt thereof, in 300 mL of water at 25° C., results in a solution having a pH of less than or equal to 4.0.

(57) In some aspects, the pharmaceutical dosage forms of the disclosure are those releasing the active ingredient (e.g., the fospropofol or the fospropofol salt) immediately or in modified or extended-release manner.

(58) In some aspects, the pharmaceutical dosage forms of the disclosure are those wherein dissolution of an amount of the dosage form containing a therapeutically effective amount of fospropofol, or a pharmaceutically acceptable salt thereof, in 300 mL of 0.1 N HCl at 37° C., results in at least 30% (e.g., 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, or 99% or greater than 99%) of the fospropofol or pharmaceutically acceptable

salt thereof (on a fospropofol basis) in the dosage form being released from the dosage form within 60 minutes or less (e.g., within 60 minutes, 55 minutes, 50 minutes, 45 minutes, 40 minutes, 35 minutes, 30 minutes, 25 minutes, 20 minutes, 15 minutes, 10 minutes, or within 5 minutes).

(59) In some embodiments, the pharmaceutical dosage forms of the disclosure are those wherein dissolution of an amount of the dosage form containing a therapeutically effective amount of fospropofol, or a pharmaceutically acceptable salt thereof, in 300 mL of 0.1 N HCl at 37° C. results in at least 30% of the fospropofol or pharmaceutically acceptable salt thereof (on a fospropofol basis) in the dosage form being released from the dosage form within 30 minutes.

(60) In some embodiments, the pharmaceutical dosage forms of the disclosure are those wherein dissolution of an amount of the dosage form containing a therapeutically effective amount of fospropofol, or a pharmaceutically acceptable salt thereof, in 300 mL of 0.1 N HCl at 37° C. results in at least 50% of the fospropofol or pharmaceutically acceptable salt thereof (on a fospropofol basis) in the dosage form being released from the dosage form within 60 minutes.

(61) In other embodiments, the pharmaceutical dosage forms of the disclosure are those wherein dissolution of an amount of the dosage form containing a therapeutically effective amount of fospropofol, or a pharmaceutically acceptable salt thereof, in 300 mL of 0.1 N HCl at 37° C. results in at least 90% of the fospropofol or pharmaceutically acceptable salt thereof (on a fospropofol basis) in the dosage form being released from the dosage form within 30 minutes.

(62) In some aspects, the pharmaceutical dosage forms of the disclosure are those wherein the mole ratio of fospropofol, or a pharmaceutically acceptable salt thereof, to pharmaceutically acceptable acid is 3:1 or less, such as, for example, 0.2:1, 0.25:1, 0.3:1, 0.35:1, 0.4:1, 0.45:1, 0.5:1, 0.55:1, 0.6:1, 0.65:1, 0.7:1, 0.75:1, 0.8:1, 0.85:1, 0.9:1, 0.95:1, 1:1, 1.1:1, 1.2:1, 1.3:1, 1.4:1, 1.5:1, 1.6:1, 1.7:1, 1.8:1, 1.9:1, 2:1, 2.1:1, 2.2:1, 2.3:1, 2.4:1, 2.5:1, 2.6:1, 2.7:1, 2.8:1, 2.9:1, or 3:1.

(63) The pharmaceutical dosage forms of the disclosure may be tablets, capsules, caplets, softgels, sterile aqueous or organic solutions, reconstitutable powders, elixirs, liquids, colloidal or other types of suspensions, emulsions, beads, beadlets, granules, microparticles, nanoparticles, and combinations thereof.

(64) In some embodiments, the pharmaceutical dosage form of the disclosure is a tablet, capsule, or softgel.

(65) In some aspects, the disclosure is directed to methods of administering fospropofol or a pharmaceutically acceptable salt thereof to a subject in need thereof, the method comprising orally administering fospropofol, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable acid, to the subject.

(66) In some embodiments, the disclosure is directed to methods of administering fospropofol or a pharmaceutically acceptable salt thereof to a subject in need thereof, comprising orally administering to the subject a pharmaceutical dosage form of this disclosure.

(67) In other embodiments of the disclosed methods, the subject is orally co-administered fospropofol (or a pharmaceutically acceptable salt thereof) and a pharmaceutically acceptable acid.

(68) As used herein, the term “orally co-administering” refers to simultaneous administration, or sequential administration in such a manner that the fospropofol, or a pharmaceutically acceptable salt thereof, and the pharmaceutically acceptable acid are present in the subject's stomach at the same time.

(69) In some embodiments of the disclosed methods, the fospropofol (or pharmaceutically acceptable salt thereof) and the pharmaceutically acceptable acid are administered in the same unit dosage form (i.e., the fospropofol or pharmaceutically acceptable salt thereof are present in the same dosage form together with the pharmaceutically acceptable acid).

(70) In other embodiments, all or a portion of the pharmaceutically acceptable acid is administered separately from the dosage form comprising the fospropofol (or pharmaceutically acceptable salt thereof).

(71) In some embodiments, the methods are for orally administering fospropofol.

(72) In other embodiments, the methods are for orally administering a pharmaceutically acceptable salt of fospropofol.

(73) In other embodiments, the methods are for orally administering fospropofol disodium.

(74) In some embodiments of the disclosed methods, the pharmaceutically acceptable acid used in the methods of the disclosure is 1-hydroxy-2-naphthoic acid, 2,2-dichloroacetic acid, 2-hydroxethanesulfonic acid, 4-acetamidobenzoic acid, 4-aminosalicylic acid, acetic acid, aceturic acid, Acid hydrolyzed proteins, Acid Modified Starch, Aconitic Acid, adipic acid, alginic acid, a-oxo-glutaric acid, benzenesulfonic acid, benzoic acid, butyric acid, camphor-10-sulfonic acid, camphoric acid, capric acid, caproic acid, caprylic acid, carbonic acid, Cholic acid, cinnamic acid, citric acid, cyclamic acid, D(-)-Lactic acid, Desoxycholic acid, D-glucaric acid, D-glucoheptonic acid, D-glucuronic acid, Di(tert-butyl)naphthalenedisulfonic acid, Di(tert-butyl)naphthalenesulfonic acid, DL-lactic acid, DL-mandelic acid, DL-tartaric acid, tartaric acid, dodecylsulfuric acid, Erythorbic acid (D-isoascorbic acid), ethane-1,2-disulfonic acid, ethanesulfonic acid, formic acid, fumaric acid, galactaric acid, gentisic acid, glutaric acid, glycerophosphoric acid, Glycocholic acid, glycolic acid, hexanedioic acid, hippuric acid, hydrobromic acid, Hydrochloric acid, Iron naphthenate, iron salts, iron salts, isobutyric acid, L(+)-lactic acid, L(+)-potassium acid tartrate, L(+)-tartaric acid, Lactic acid, lactobionic acid, L-ascorbic acid, ascorbic acid, L-aspartic acid, lauric acid, L-glutamic acid, L-Glutamic acid hydrochloride, Linoleic acid, L-Malic acid, L-pyroglutamic acid, L-tartaric acid, maleic acid, malic acid, malonic acid, methanesulfonic acid, monobasic potassium phosphate, monobasic sodium phosphate, naphthalene-1,5-disulfonic acid, naphthalene-2-sulfonic acid, naphthenic acids, Niacin (nicotinic acid), nicotinic acid, nitric acid, octanoic acid, oleic acid, orotic acid, oxalic acid, palmitic acid, pamoic acid, Pectin low acid, Pectinic acid, phosphoric acid, propanoic acid, Propionic acid, p-toluenesulfonic acid, pyruvic acid, saccharin, salicylic acid, sebacic acid, Sodium acid pyrophosphate, Sodium aluminum phosphate, sodium metabisulfite, Sorbic acid, stearic acid, succinic acid, sulfuric acid, tall oil fatty acids, Tannic acid (hydrolyzable gallotannins), Taurocholic acid, thiocyanic acid, Thiodipropionic acid, trifluoroacetic acid, undec-10-enoic acid, orange juice, apple juice, grapefruit juice, or a combination thereof.

(75) In some embodiments, the pharmaceutically acceptable acid used in the methods of the disclosure is ascorbic acid. In some embodiments, the ascorbic acid is DL-ascorbic acid. In other embodiments, the ascorbic acid is L-ascorbic acid. In other embodiments, the ascorbic acid is D-ascorbic acid.

(76) In other embodiments, the pharmaceutically acceptable acid used in the methods of the disclosure is tartaric acid. In some embodiments, the tartaric acid is DL- tartaric acid. In other embodiments, the tartaric acid is L- tartaric acid. In other embodiments, the tartaric acid is D-tartaric acid.

(77) In some embodiments, the pharmaceutically acceptable acid used in the methods of the disclosure is citric acid.

(78) In some embodiments, the pharmaceutically acceptable acid used in the methods of the disclosure is malic acid. In some embodiments, the malic acid is DL- malic acid. In other embodiments, the malic acid is L- malic acid. In other embodiments, the malic acid is D- malic acid.

(79) In some methods of the disclosure, the administration of fospropofol (or a pharmaceutically acceptable salt thereof) and a pharmaceutically acceptable acid results in a propofol C_{max} that is greater than the C_{max} resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without co-administration of the pharmaceutically acceptable acid. The term "C_{max}", as used herein, refers to the peak concentration of propofol observed in the subject's plasma following administration of fospropofol or a pharmaceutically acceptable salt thereof. The concentration of propofol in the subject's plasma samples can be determined using standard analytical methods. In some embodiments of such methods, the administration of fospropofol (or a

pharmaceutically acceptable salt thereof) and a pharmaceutically acceptable acid comprises administering a pharmaceutical dosage form as described herein.

(80) In some embodiments of the methods of the disclosure, the administration of fospropofol (or a pharmaceutically acceptable salt thereof) and a pharmaceutically acceptable acid results in a propofol C_{max} that is at least 1.2 times greater than a propofol C_{max} resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of the pharmaceutically acceptable acid, such as for example, at least 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3.0, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9, or 4 times greater.

(81) In some embodiments of the methods of the disclosure, the propofol C_{max} is 1.2 times greater than a propofol C_{max} resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof without administration of the pharmaceutically acceptable acid.

(82) In some embodiments of the methods of the disclosure, the propofol C_{max} is at least 1.5 times greater than a propofol C_{max} resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of the pharmaceutically acceptable acid.

(83) In some embodiments of the methods of the disclosure, the propofol C_{max} is at least 2 times greater than a propofol C_{max} resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of the pharmaceutically acceptable acid.

(84) In some embodiments of the methods of the disclosure, the propofol C_{max} is at least 2.5 times greater than a propofol C_{max} resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of the pharmaceutically acceptable acid.

(85) In some embodiments of the methods of the disclosure, the propofol C_{max} is 3 times greater than a propofol C_{max} resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of the pharmaceutically acceptable acid.

(86) In some methods of the disclosure, the administration of fospropofol (or a pharmaceutically acceptable salt thereof) and a pharmaceutically acceptable acid results in a propofol AUC that is greater than a propofol AUC resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of the pharmaceutically acceptable acid. In some embodiments, the AUC is AUC.sub.t. The term "AUC.sub.t", as used herein, refers to the area under the plasma propofol concentration-time curve from time zero to time t. In other embodiments, the AUC is AUC.sub.∞. The term "AUC.sub.∞", as used herein, refers to the AUC obtained by extrapolation of AUC.sub.0 to ∞. In some embodiments of such methods, the administration of fospropofol (or a pharmaceutically acceptable salt thereof) and a pharmaceutically acceptable acid comprises administering a pharmaceutical dosage form as described herein.

(87) In some embodiments of the methods of the disclosure, the propofol AUC.sub.∞ is at least 1.5 times greater than a propofol AUC.sub.∞ resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of the pharmaceutically acceptable acid, such as for example, at least 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3.0, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9, or 4 times greater.

(88) In some embodiments of the methods of the disclosure, the propofol AUC.sub.∞ is 1.5 times greater than a propofol AUC.sub.∞ resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of the pharmaceutically acceptable acid.

(89) In some embodiments of the methods of the disclosure, the propofol AUC.sub.∞ is at least 1.6 times greater than a propofol AUC.sub.∞ resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of the pharmaceutically acceptable acid.

(90) In some embodiments of the methods of the disclosure, the propofol AUC.sub.∞ is at least 2 times greater than a propofol AUC.sub.∞ resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of the pharmaceutically acceptable

acid.

(91) In some embodiments of the methods of the disclosure, the propofol AUC.sub.∞ is at least 2.5 times greater than a propofol AUC.sub.∞ resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of the pharmaceutically acceptable acid.

(92) In some embodiments of the methods of the disclosure, the propofol AUC.sub.∞ is at least 3 times greater than a propofol AUC.sub.∞ resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of the pharmaceutically acceptable acid.

(93) In some embodiments of the methods disclosed herein, the subject is a human.

(94) In some embodiments of the methods disclosed herein, the subject is experiencing hypochlorhydria or achlorhydria prior to the administration of a dosage form as described herein. In some embodiments, the subject is diagnosed with hypochlorhydria or achlorhydria prior to the administration. In other embodiments of the methods disclosed herein, the subject is suspected of having hypochlorhydria or achlorhydria prior to the administration.

(95) In some embodiments of the methods disclosed herein, the subject has been administered a proton pump inhibitor (PPI) prior to the administration of a dosage form as described herein. Exemplary proton pump inhibitors include Omeprazole (Prilosec), Esomeprazole (Nexium), Lansoprazole (Prevacid), Rabeprazole (AcipHex), Pantoprazole (Protonix), Dexlansoprazole (Dexilant), and Zegerid (omeprazole with sodium bicarbonate).

(96) In some aspects, the methods described herein are directed to treating a disease or disorder in a subject in need thereof.

(97) In some embodiments, the methods described herein are directed to treating a disease or disorder comprising orally administering to a subject a pharmaceutical dosage form described herein.

(98) In other embodiments, the methods described herein are directed to treating a disease or disorder comprising orally administering fospropofol, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable acid, to a subject.

(99) In some embodiments of the disclosed methods of treating a disease or disorder, the disease or disorder is migraine, acute repetitive seizures, seizure clusters, neuropathic pain, postherpetic neuralgia, traumatic brain injury, Alzheimer's disease, epilepsy, anxiety, fragile X syndrome, post-traumatic stress disorder, lysosomal storage disorders (Niemann-Pick type C disease), depression (including post-partum depression), premenstrual dysphoric disorder, alcohol craving, or smoking cessation.

(100) In some embodiments of the disclosed methods of treating a disease or disorder, the disease or disorder is migraine. The term "migraine," as used herein, refers to a chronic neurovascular disorder characterized by recurrent attacks of often severe headache ("migraine attacks"), typically accompanied by nausea and sensitivity to light and/or sound. Migraine is a clinical diagnosis, criteria for which would be known and understood by those practicing in the treatment of migraine, and would include, for example, the criteria proposed by the International Headache Society (IHS). See <http://ihs-classification.org/en/>.

(101) In some embodiments, the subject's migraine is migraine with aura. Migraine with aura (also referred to as classic migraine) is characterized by focal neurological symptoms that typically precede, or sometimes accompany, the headache.

(102) In other embodiments, the subject's migraine is migraine without aura. Migraine without aura (also referred to as common migraine) is characterized by the absence of focal neurological symptoms that typically precede, or sometimes accompany, the headache.

(103) In other embodiments, the subject's migraine is cluster headache.

(104) In other embodiments, the subject's migraine is intractable migraine.

(105) In some embodiments, the patient's migraine is refractory migraine. Refractory migraine, as

used herein, refers to migraine that fails to respond to pharmacologic treatment. Failure to respond in this regard includes, for example, failure of a pharmacological treatment to eliminate migraine pain, as well as failure of a pharmacological treatment to reduce severe or moderate migraine pain to mild migraine pain. Refractory migraine may fail to respond one or more types of pharmacologic treatment. Examples of pharmacologic treatment to which refractory migraine may fail to respond include CGRP inhibitors (e.g., gepants, anti-CGRP antibodies), and triptans (e.g., sumatriptan (Imitrex), rizatriptan (Maxalt), almotriptan (Axert), naratriptan (Amerge), zolmitriptan (Zomig), frovatriptan (Frova) and eletriptan (Relpax)).

(106) In some embodiments of the disclosed methods, the subject's refractory migraine may fail to respond to CGRP inhibitors, and is referred to as CGRP inhibitor-refractory migraine.

(107) In some embodiments, the subject's CGRP-inhibitor refractory migraine fails to respond to gepant treatment, and is referred to as gepant-refractory migraine. In other embodiments, the subject's CGRP-inhibitor refractory migraine fails to respond to anti-CGRP antibodies, and is referred to as anti-CGRP antibody-refractory migraine.

(108) In other embodiments, the subject's refractory migraine may fail to respond to triptans, and is referred to as triptan-refractory migraine.

(109) In other embodiments, the subject's refractory migraine may fail to respond to NSAIDs and is referred to as NSAID-refractory migraine.

(110) In other embodiments, the subject's refractory migraine may fail to respond to dihydroergotamine (DHE) and is referred to as DHE-refractory migraine.

(111) In some embodiments of the disclosed methods of treating a disease or disorder, the disease or disorder is epilepsy.

EXAMPLES

Example 1: Dissolution and pH Studies

(112) Fospropofol Disodium tablets, 600 mg (label claim; 1.c.), were manufactured for dissolution testing and use in a pharmacokinetics study. A total of three tablets for each lot were weighed prior to testing.

(113) TABLE-US-00001 TABLE 1 Acidified Tablets-Compositions

	A1	A2	A3	A4	%	%	%	%
Composition mg w/w	mg w/w	mg w/w	mg w/w	Fospropofol	666.9*	42.96	666.9*	42.96
56.43	666.9*	65.18	Disodium hydrate*	Polyplasdone	77.6	5.00	77.6	5.00
59.1	5.00	51.2	5.00	Magnesium	7.8	0.50	7.8	0.50
5.9	0.50	5.1	0.50	Stearate	Citric Acid, 800.0	51.54	monohydrate	Ascorbic 800.0
51.54	Acid	Tartaric Acid	450.0	38.07	Malic Acid	300.0	29.32	Total
1552.3	100.00	1181.9	100.00	1023.2	100.00	*equivalent to 600 mg fospropofol disodium adjusted for water content.		

(114) Dissolution studies were performed under the conditions shown in Table 2 (n=3 vessels, 300-mL media volume and a single tablet per vessel). Analyses were performed by HPLC under the conditions shown in Table 3.

(115) TABLE-US-00002 TABLE 2 Conditions for Dissolution Studies

Conditions	Setting
Dissolution Medium	0.1 NHCl de-aerated
Apparatus	USP apparatus II (rotating paddles)
Vessel Volume	300 mL
Temperature	37.0° C. ± 0.5° C.
Rotation Speed	50 RPM (200 RPM from 60-75 or 60-90 minutes)
Sample Volume	1.5 mL
Sample Times	5, 10, 15, 20, 25, 30, 40, 45, and 60 minutes
with an infinity pull at	75 or 90 minutes
In-line Filter	QLA 10 µm Porous (Full Flow)
Filters	

(116) TABLE-US-00003 TABLE 3 HPLC Instrumental Conditions

Instrument	A	suitable gradient	
HPLC system equipped with an inline degasser.	Column	Phenomenex Luna C18(2), 50 × 4.6 mm, 3 µm/PIN 00B-4251-E0	
Detection	UV, 220 nm	Column Temperature	25° C.
Sample Temperature		Ambient	
Flow Rate	2.0 mL/minute	Injection Volume	10 µL
Run Time	8 minutes	(See gradient table below)	
Mobile Phase A	0.1% TFA in HPW	Mobile Phase B	0.1% TFA in ACN
Column/Needle Wash	50:50 ACN:HPW	Attenuation	1000 mAUN (when applicable)
Gradient Time (min), % A, % B	0.0, 95, 5	6.0, 5, 95	6.2, 95, 5
	8.0, 95, 5		

(117) TABLE-US-00004 TABLE 4 A1 (citric acid monohydrate) Dissolution Results for

Fospropofol Disodium Tablets, 600 mg, (% of label claim recovery) Pull Point (minutes) Vessel 5
10 15 20 25 30 40 45 60 1 10 22 31 42 48 53 61 63 71 2 10 21 30 38 44 49 55 62 67 3 12 26 36 44
49 54 62 65 71 Average 11 23 32 41 47 52 59 63 70 SD 1.2 2.6 3.2 3.1 2.6 2.6 3.8 1.5 2.3 % RSD
10.8 11.5 9.9 7.4 5.6 5.1 6.4 2.4 3.3

(118) TABLE-US-00005 TABLE 5 A2 (ascorbic acid) Dissolution Results for Fospropofol
Disodium Tablets, 600 mg, (% of label claim recovery) Pull Point (minutes) Vessel 5 10 15 20 25
30 40 45 60 1 26 58 76 86 89 90 93 91 92 2 31 62 78 86 90 91 93 93 93 3 29 63 78 88 89 91 89 88
91 Average 29 61 77 87 89 91 92 91 92 SD 2.5 2.6 1.2 1.2 0.6 0.6 2.3 2.5 1.0 % RSD 8.8 4.3 1.5
1.3 0.6 0.6 2.5 2.8 1.1

(119) TABLE-US-00006 TABLE 6 A3 (tartaric acid) Dissolution Results for Fospropofol
Disodium Tablets, 600 mg, (% of label claim recovery) Pull Point (minutes) Vessel 5 10 15 20 25
30 40 45 60 1 9 20 28 33 37 41 47 50 54 2 8 18 27 33 37 40 46 49 56 3 7 16 24 30 33 37 43 46 52
Average 8 18 26 32 36 39 45 48 54 SD 1.0 2.0 2.1 1.7 2.3 2.1 2.1 2.1 2.0 % RSD 12.5 11.1 7.9 5.4
6.5 5.3 4.6 4.3 3.7

(120) TABLE-US-00007 TABLE 7 A4 (malic acid) Dissolution Results for Fospropofol Disodium
Tablets, 600 mg, (% of label claim recovery) Pull Point (minutes) Vessel 5 10 15 20 25 30 40 45 60
1 21 48 63 70 76 77 82 82 85 2 23 48 59 66 74 78 79 81 84 3 23 49 61 67 70 75 77 79 83 Average
22 48 61 68 73 77 79 81 84 SD 1.2 0.6 2.0 2.1 3.1 1.5 2.5 1.5 1.0 % RSD 5.2 1.2 3.3 3.1 4.2 2.0 3.2
1.9 1.2

(121) As used herein, the term “label claim” refers the total weight of the fospropofol disodium
within the dosage unit.

(122) FIG. 1 shows the dissolution profiles of the acidified tablets.

(123) The solution pH values after the acidified tablets were dissolved in 300 mL of water are
shown in Table 8.

(124) TABLE-US-00008 TABLE 8 pH After Acidified Tablet Dissolved in 300 mL of Water
Acidifier Excipient pH Citric Acid 3.11 Tartaric Acid 3.13 Malic Acid 3.94 Ascorbic Acid 4.10

Example 2: Bioavailability Studies

(125) The bioavailability of propofol from the fospropofol-containing dosage forms is assessed in
dogs using the following general protocol.

(126) The dogs used in the studies have the following characteristics: Strain: Beagle Condition:
Purpose-bred, non-naïve Source: Marshall Farms. North Rose, NY Number of Males: 6 (plus 1
alternate) Target Age at the Initiation of Dosing: At least 8 months. Target Weight at the Initiation
of Dosing: 6 to 13 kg

(127) All animals used in the studies have documentation of immunization for parvovirus,
distemper, adenovirus type 2, parainfluenza, *Bordetella*, papilloma, and rabies.

(128) Animals are identified with a tattoo or a subcutaneously implanted electronic identification
chip.

(129) Each animal is inspected by a clinical veterinarian upon receipt. Animals judged to be in
good health are placed immediately in acclimation for at least 10 days.

(130) Animals judged to be suitable for testing are assigned to groups randomly based on body
weight stratification into a block design using computer program. Animals are arbitrarily
reassigned to a different group at the discretion of the study director based on acclimation data. The
animals are dosed as follows: Dose Route: Tablet; Acidified Tablet Frequency: Once daily; single
administration Method: The first day of dosing is designated as Day 1 (exception: alternate animals
used for replacement after Day 1 assume the day of the animal being replaced). Tablet
Administration: A single dose of the test article is administered orally via tablet. Each subject
receives 1 (600 mg) tablet. Doses are irrespective of body weight. In studies wherein the dogs are
pretreated with pentagastin, the pentagastin is administered intravenously before the dogs are
administered the fospropofol-containing tablet.

(131) In a separate study, the dogs are orally administered a solution containing 30 mg/mL

fospropofol at a dose of 160 mg/kg in water rather than a tablet.

(132) Bioanalytical Methods:

(133) Venipuncture from a jugular vein (saphenous or cephalic vein is used, if necessary). Target Volume (mL): Approximately 1 mL/time point collected without anesthesia. Anticoagulant: Sodium Heparin Prior to blood collection, approximately 0.05 mL of 200 mg/mL of sodium orthovanadate (SOV) solution is added to the heparinized blood collection tubes to prevent ex vivo conversion via alkaline phosphatase. Special Requirements: After collection, blood collection tubes are kept on wet ice until centrifugation within 30 minutes of collection. Processing: Plasma Samples are mixed gently and centrifuged within 30 minutes of collection. The samples are centrifuged at 2-8° C. and the resultant plasma is separated, transferred to duplicate uniquely labeled polypropylene tubes, and kept on wet ice until transferred to storage. Samples are stored in a freezer set to maintain a target of -70° C. Bioanalytical samples are analyzed for concentration of test article (fospropofol) and specified metabolites (propofol) using a validated analytical procedure.

(134) FIG. 2 shows the propofol plasma profile comparisons of (1) 600 mg fospropofol disodium tablet with ascorbic acid acidifier (Composition A2) with no pentagastrin pretreatment; (2) 600 mg fospropofol disodium tablet with tartaric acid acidifier (Composition A3) with no pentagastrin pretreatment; and (3) 1300 mg fospropofol disodium aqueous solution (30 mg/mL) with no pentagastrin pretreatment.

(135) FIG. 3 shows the propofol plasma profile comparisons of (1) 600 mg fospropofol disodium tablet with ascorbic acid acidifier (Composition A2) with pentagastrin pretreatment; and (2) 600 mg fospropofol disodium tablet with no acidifier (Control) with pentagastrin pretreatment.

(136) FIG. 4 shows the propofol plasma profile comparisons of (1) 600 mg fospropofol disodium tablet with tartaric acid acidifier (Composition A3) with no pentagastrin pretreatment; and (2) 600 mg fospropofol disodium tablet with no acidifier (Control) with pentagastrin pretreatment.

(137) The pharmacokinetic (PK) analysis results are shown in Table 9.

(138) TABLE-US-00009 TABLE 9 PK Results C_{max} AUC_{sub.∞} Half-life (ng/mL) (ng * hr/mL) (h)
1300 mg Fospropofol disodium aqueous solution (30 mg/mL) with no pentagastrin pretreatment
Mean 399.43 711.06 1.45 SD 281.70 450.13 0.58 CV 71% 63% 40%
600 mg Fospropofol disodium tablet with ascorbic acid acidifier (Composition A2) with no pentagastrin pretreatment
Mean 714.40 1191.91 1.48 SD 295.70 506.15 0.2 CV 41% 42% 14%
600 mg Fospropofol disodium tablet with tartaric acid acidifier (Composition A3) with no pentagastrin pretreatment
Mean 1107.50 1656.87 2.93 SD 685.43 998.42 3.10 CV 62% 60% 106%
600 mg Fospropofol disodium tablet with control (i.e., no acidifier) with pentagastrin pretreatment
Mean 732.5 1002.22 1.34 SD 312.47 340.86 0.34 CV 43% 34% 26%
600 mg Fospropofol disodium tablet with ascorbic acid acidifier (Composition A2) with pentagastrin pretreatment
Mean 928.33 1623.93 3.84 SD 167.12 378.36 3.22 CV 18% 23% 84%

(139) The pharmacokinetic results demonstrate that administering fospropofol together with an acid, such as in the acidified tablets, results in increased propofol C_{max} and AUC_{sub.∞} when compared to the dose adjusted exposure seen with the oral solution. See FIG. 2; Table 9. This effect is seen using both ascorbic acid and tartaric acid as the tablet acidifiers.

(140) In addition, the results demonstrate that administration of the acidified tablets with pentagastrin pretreatment results in increased plasma propofol C_{max} and AUC_{sub.∞} compared to the plasma propofol C_{max} and AUC_{sub.∞} observed upon administration of a control tablet (i.e., with no acidifier) with pentagastrin pretreatment. See FIG. 3; Table 9. Pentagastrin pretreatment acidifies the subject's stomach contents to an acidity comparable to that of normal human stomach contents.

(141) In addition, these results demonstrate that co-administration of fospropofol disodium with tartaric acid without pentagastrin pretreatment results in increased plasma propofol C_{max} and AUC_{sub.∞} compared to the plasma propofol C_{max} and AUC_{sub.∞} observed upon administration

of a control tablet (i.e., with no acidifier) with pentagastrin pretreatment. See FIG. 4; Table 9.

(142) The experiments described above surprisingly demonstrate that oral administration of fospropofol together with a pharmaceutically acceptable acid increases the plasma propofol C_{max} and AUC_{sub.∞} relative to controls in which fospropofol is orally administered without an acidifier. Moreover, this effect is seen with multiple acids (e.g., ascorbic acid, tartaric acid) and is seen under conditions expected in human subjects. Thus, the disclosed invention solves the problem of providing an oral dosage form that can provide useful propofol pharmacokinetics.

(143) These results also demonstrate that co-administration of fospropofol disodium with a pharmaceutically acceptable acid can result in decreased variability in C_{max} and AUC_∞ relative to administration without acid.

(144) In some embodiments, the disclosure is directed to the following aspects: Aspect 1. A pharmaceutical dosage form for oral administration comprising fospropofol or a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable acid. Aspect 2. The pharmaceutical dosage form according to aspect 1 comprising fospropofol. Aspect 3. The pharmaceutical dosage form according to aspect 1 comprising a pharmaceutically acceptable salt of fospropofol. Aspect 4. The pharmaceutical dosage form according to aspect 3, wherein the pharmaceutically acceptable salt of fospropofol is fospropofol disodium. Aspect 5. The pharmaceutical dosage form according to any one of the preceding aspects, wherein the pharmaceutically acceptable acid is on the FDA inactive ingredient database (IID) for approved drug products or generally recognized as safe (GRAS).

Aspect 6. The pharmaceutical dosage form according to any one of the preceding aspects, wherein the pharmaceutically acceptable acid is 1-hydroxy-2-naphthoic acid, 2,2-dichloroacetic acid, 2-hydroxethanesulfonic acid, 4-acetamidobenzoic acid, 4-aminosalicylic acid, acetic acid, aceturic acid, Acid hydrolyzed proteins, Acid Modified Starch, Aconitic Acid, adipic acid, alginic acid, α-oxo-glutaric acid, benzenesulfonic acid, benzoic acid, butyric acid, camphor-10-sulfonic acid, camphoric acid, capric acid, caproic acid, caprylic acid, carbonic acid, Cholic acid, cinnamic acid, citric acid, cyclamic acid, D(-)-Lactic acid, Desoxycholic acid, D-glucaric acid, D-glucoheptonic acid, D-glucuronic acid, Di(tert-butyl)naphthalenedisulfonic acid, Di(tert-butyl)naphthalenesulfonic acid, DL-lactic acid, DL-mandelic acid, DL-tartaric acid, tartaric acid, dodecylsulfuric acid, Erythorbic acid (D-isoascorbic acid), ethane-1,2-disulfonic acid, ethanesulfonic acid, formic acid, fumaric acid, galactaric acid, gentisic acid, glutaric acid, glycerophosphoric acid, Glycocholic acid, glycolic acid, hexanedioic acid, hippuric acid, hydrobromic acid, Hydrochloric acid, Iron naphthenate, iron salts, iron salts, isobutyric acid, L(+)-lactic acid, L(+)-potassium acid tartrate, L(+)-tartaric acid, Lactic acid, lactobionic acid, L-ascorbic acid, ascorbic acid, L-aspartic acid, lauric acid, L-glutamic acid, L-Glutamic acid hydrochloride, Linoleic acid, L-Malic acid, L-pyroglutamic acid, L-tartaric acid, maleic acid, malic acid, malonic acid, methanesulfonic acid, monobasic potassium phosphate, monobasic sodium phosphate, naphthalene-1,5-disulfonic acid, naphthalene-2-sulfonic acid, naphthenic acids, Niacin (nicotinic acid), nicotinic acid, nitric acid, octanoic acid, oleic acid, orotic acid, oxalic acid, palmitic acid, pamoic acid, Pectin low acid, Pectinic acid, phosphoric acid, propanoic acid, Propionic acid, p-toluenesulfonic acid, pyruvic acid, saccharin, salicylic acid, sebacic acid, Sodium acid pyrophosphate, Sodium aluminum phosphate, sodium metabisulfite, Sorbic acid, stearic acid, succinic acid, sulfuric acid, tall oil fatty acids, Tannic acid (hydrolyzable gallotannins), Taurocholic acid, thiocyanic acid, Thiodipropionic acid, trifluoroacetic acid, undec-10-enoic acid, orange juice, apple juice, grapefruit juice, or a combination thereof or a combination thereof. Aspect 7. The pharmaceutical dosage form according to any one of the preceding aspects, wherein the pharmaceutically acceptable acid is ascorbic acid, citric acid, malic acid, or tartaric acid. Aspect 8. The pharmaceutical dosage form according to any one of the preceding aspects, wherein the pharmaceutically acceptable acid is ascorbic acid. Aspect 9. The pharmaceutical dosage form according to any one of the preceding aspects, wherein the pharmaceutically acceptable acid is tartaric acid. Aspect 10. The pharmaceutical dosage form according to any one of the preceding

aspects, wherein the pharmaceutically acceptable acid is citric acid. Aspect 11. The pharmaceutical dosage form according to any one of the preceding aspects, wherein the pharmaceutically acceptable acid is malic acid. Aspect 12. The pharmaceutical dosage form according to any one of the preceding aspects, wherein dissolution of an amount of the dosage form containing a therapeutically effective amount of fospropofol, or a pharmaceutically acceptable salt thereof, in 300 mL of water at 25° C. results in a solution having a pH of 6.3 or less. Aspect 13. The pharmaceutical dosage form according to any one of the preceding aspects, wherein dissolution of an amount of the dosage form containing a therapeutically effective amount of fospropofol, or a pharmaceutically acceptable salt thereof, in 300 mL of water at 25° C. results in a solution having a pH of 4.5 or less. Aspect 14. The pharmaceutical dosage form according to any one of the preceding aspects, wherein dissolution of an amount of the dosage form containing a therapeutically effective amount of fospropofol, or a pharmaceutically acceptable salt thereof, in 300 mL of water at 25° C. results in a solution having a pH of 4.2 or less. Aspect 15. The pharmaceutical dosage form according to any one of the preceding aspects, wherein dissolution of an amount of the dosage form containing a therapeutically effective amount of fospropofol, or a pharmaceutically acceptable salt thereof, in 300 mL of water at 25° C. results in a solution having a pH of 4.0 or less. Aspect 16. The pharmaceutical dosage form according to any one of the preceding aspects, wherein dissolution of an amount of the dosage form containing a therapeutically effective amount of fospropofol, or a pharmaceutically acceptable salt thereof, in 300 mL of 0.1 N HCl at 37° C., results in at least 30% of the fospropofol or pharmaceutically acceptable salt thereof (on a fospropofol basis) in the dosage form being released from the dosage form within 60 minutes. Aspect 17. The pharmaceutical dosage form according to aspect 16, wherein dissolution of an amount of the dosage form containing a therapeutically effective amount of fospropofol, or a pharmaceutically acceptable salt thereof, in 300 mL of 0.1 N HCl at 37° C. results in at least 30% of the fospropofol or pharmaceutically acceptable salt thereof (on a fospropofol basis) in the dosage form being released from the dosage form within 30 minutes. Aspect 18. The pharmaceutical dosage form according to any one of the preceding aspects, wherein dissolution of an amount of the dosage form containing a therapeutically effective amount of fospropofol, or a pharmaceutically acceptable salt thereof, in 300 mL of 0.1 N HCl at 37° C. results in at least 50% of the fospropofol or pharmaceutically acceptable salt thereof (on a fospropofol basis) in the dosage form being released from the dosage form within 60 minutes. Aspect 19. The pharmaceutical dosage form according to any one of the preceding aspects, wherein dissolution of an amount of the dosage form containing a therapeutically effective amount of fospropofol, or a pharmaceutically acceptable salt thereof, in 300 mL of 0.1 N HCl at 37° C. results in at least 50% of the fospropofol or pharmaceutically acceptable salt thereof (on a fospropofol basis) in the dosage form being released from the dosage form within 30 minutes. Aspect 20. The pharmaceutical dosage form according to any one of the preceding aspects, wherein dissolution of an amount of the dosage form containing a therapeutically effective amount of fospropofol, or a pharmaceutically acceptable salt thereof, in 300 mL of 0.1 N HCl at 37° C. results in at least 90% of the fospropofol or pharmaceutically acceptable salt thereof (on a fospropofol basis) in the dosage form being released from the dosage form within 30 minutes. Aspect 21. The pharmaceutical dosage form according to any one of the preceding aspects, wherein dissolution of an amount of the dosage form containing a therapeutically effective amount of fospropofol, or a pharmaceutically acceptable salt thereof, in 300 mL of 0.1 N HCl at 37° C. results in at least 90% of the fospropofol or pharmaceutically acceptable salt thereof (on a fospropofol basis) in the dosage form being released from the dosage form within 60 minutes. Aspect 22. The pharmaceutical dosage form according to any one of the preceding aspects, wherein dissolution of an amount of the dosage form containing a therapeutically effective amount of fospropofol, or a pharmaceutically acceptable salt thereof, in 300 mL of 0.1 N HCl at 37° C. results in release of at least 90% of the fospropofol from the dosage form within 30 minutes. Aspect 23. The pharmaceutical dosage form according to any one of the

preceding aspects, wherein the mole ratio of fospropofol or a pharmaceutically acceptable salt thereof (on a fospropofol basis) to pharmaceutically acceptable acid is 3:1 or less. Aspect 24. A method of administering fospropofol or a pharmaceutically acceptable salt thereof to a subject in need thereof, comprising orally administering to said subject a pharmaceutical dosage form according to any one of the preceding aspects. Aspect 25. A method of administering fospropofol or a pharmaceutically acceptable salt thereof to a subject in need thereof, said method comprising orally administering fospropofol, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable acid, to the subject. Aspect 26. The method of aspect 24 or aspect 25, wherein said fospropofol, or a pharmaceutically acceptable salt thereof, and said pharmaceutically acceptable acid are present in the same unit dosage form. Aspect 27. The method of aspect 25, wherein all or a portion of said pharmaceutically acceptable acid is administered separately from the unit dosage form comprising said fospropofol, or a pharmaceutically acceptable salt thereof. Aspect 28. The method according to any one of aspects 24 to 27, wherein said pharmaceutically acceptable acid is ascorbic acid, citric acid, malic acid, tartaric acid, succinic acid, fumaric acid, maleic acid, lactic acid, monobasic sodium phosphate, monobasic potassium phosphate, sodium metabisulfite, apple juice, orange juice, or grapefruit juice. Aspect 29. The method according to aspect 28, wherein said pharmaceutically acceptable acid is ascorbic acid, citric acid, malic acid, or tartaric acid. Aspect 30. The method according to aspect 29, wherein said pharmaceutically acceptable acid is ascorbic acid. Aspect 31. The method according to aspect 29, wherein said pharmaceutically acceptable acid is citric acid. Aspect 32. The method according to aspect 29, wherein said pharmaceutically acceptable acid is malic acid. Aspect 33. The method according to aspect 29, wherein said pharmaceutically acceptable acid is tartaric acid. Aspect 34. The method of any one of aspects 24 to 33, wherein said subject is experiencing hypochlorhydria or achlorhydria prior to the administration. Aspect 35. The method according to aspect 34, wherein said subject has been administered a proton pump inhibitor (PPI) prior to the administration. Aspect 36. The method of any one of aspects 24 to 35, wherein said administration results in a propofol C_{max} that is greater than a propofol C_{max} resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of the pharmaceutically acceptable acid. Aspect 37. The method of aspect 36, wherein the propofol C_{max} is at least 1.5 times greater than a propofol C_{max} resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of the pharmaceutically acceptable acid. Aspect 38. The method of aspect 36, wherein the propofol C_{max} is at least 2 times greater than a propofol C_{max} resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of the pharmaceutically acceptable acid. Aspect 39. The method of aspect 36, wherein the propofol C_{max} is at least 2.5 times greater than a propofol C_{max} resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of the pharmaceutically acceptable acid. Aspect 40. The method of aspect 36, wherein the propofol C_{max} is at least 3 times greater than a propofol C_{max} resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof without administration of the pharmaceutically acceptable acid. Aspect 41. The method of any one of aspects 24 to 40, wherein said administration results in a propofol AUC_{sub.∞} that is greater than a propofol AUC_{sub.∞} resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of the pharmaceutically acceptable acid. Aspect 42. The method of aspect 41, wherein the propofol AUC_{sub.∞} is at least 1.5 times greater than a propofol AUC_{sub.∞} resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of the pharmaceutically acceptable acid. Aspect 43. The method of aspect 41, wherein the propofol AUC_{sub.∞} is at least 2 times greater than a propofol AUC_{sub.∞} resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of the pharmaceutically acceptable acid. Aspect 44. The method of aspect 41, wherein the propofol AUC_{sub.∞} is at least 2.5 times greater than a propofol AUC_{sub.∞} resulting from administering fospropofol, or a pharmaceutically acceptable

salt thereof, without administration of the pharmaceutically acceptable acid. Aspect 45. The method of aspect 41, wherein the propofol AUC.sub.∞ is at least 3 times greater than a propofol AUC.sub.∞ resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without co-administration of the pharmaceutically acceptable acid. Aspect 46. The method of any one of aspects 24 to 45, wherein said method is directed to treating a disease or disorder in a subject in need thereof. Aspect 47. The method of aspect 46, wherein said method comprises orally administering to a subject the pharmaceutical dosage form of any one of aspects 1 to 23. Aspect 48. The method of aspect 46, comprising orally administering fospropofol, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable acid, to a subject. Aspect 49. The method of any one of aspects 46 to 48, wherein the disease or disorder is migraine, acute repetitive seizures, seizure clusters, neuropathic pain, postherpetic neuralgia, traumatic brain injury, Alzheimer's disease, epilepsy, anxiety, fragile X syndrome, post-traumatic stress disorder, lysosomal storage disorders (Niemann-Pick type C disease), depression (including post-partum depression), premenstrual dysphoric disorder, alcohol craving, or smoking cessation. Aspect 50. The method of aspect 49, wherein the disease or disorder is migraine.

Claims

1. A method of treating migraine or epilepsy in a subject in need thereof, said method comprising orally administering to said subject a pharmaceutical dosage form comprising fospropofol, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable acid, wherein the mole ratio of fospropofol, or a pharmaceutically acceptable salt thereof, to the pharmaceutically acceptable acid is 3:1 or less.
2. The method according to claim 1, wherein said subject is a human.
3. The method of claim 1, wherein the method comprises treating epilepsy in the subject.
4. The method of claim 1, wherein the method comprises treating migraine in the subject.
5. The method of claim 1, wherein said subject has or is suspected of having hypochlorhydria or achlorhydria prior to the administration of the pharmaceutical dosage form.
6. The method according to claim 5, wherein said subject has been administered a proton pump inhibitor (PPI) prior to the administration of the pharmaceutical dosage form.
7. The method of claim 1, wherein said administration of the pharmaceutical dosage form results in a propofol C_{max} that is greater than a propofol C_{max} resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of a pharmaceutically acceptable acid.
8. The method of claim 7, wherein the propofol C_{max} is at least 1.2 times greater than a propofol C_{max} resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of the pharmaceutically acceptable acid.
9. The method of claim 8, wherein the propofol C_{max} is at least 3 times greater than a propofol C_{max} resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of the pharmaceutically acceptable acid.
10. The method of claim 1, wherein said administration of the pharmaceutical dosage form results in a propofol AUC.sub.∞ that is greater than a propofol AUC.sub.∞ resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of a pharmaceutically acceptable acid.
11. The method of claim 10, wherein the propofol AUC.sub.∞ is at least 1.5 times greater than a propofol AUC.sub.∞ resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of a pharmaceutically acceptable acid.
12. The method of claim 11, wherein the propofol AUC.sub.∞ is at least 3 times greater than a propofol AUC.sub.∞ resulting from administering fospropofol, or a pharmaceutically acceptable salt thereof, without administration of a pharmaceutically acceptable acid.

13. The method of claim 1, wherein said pharmaceutical dosage form comprises fospropofol.
 14. The method of claim 1, wherein said pharmaceutical dosage form comprises a pharmaceutically acceptable salt of fospropofol.
 15. The method of claim 14, wherein said pharmaceutically acceptable salt of fospropofol is fospropofol disodium.
 16. The method of claim 1, wherein said pharmaceutical dosage form is in the form of a tablet, capsule, or softgel.
 17. The method of claim 1, wherein the mole ratio of fospropofol, or a pharmaceutically acceptable salt thereof, to the pharmaceutically acceptable acid is 2:1 or less.
 18. The method of claim 1, wherein the mole ratio of fospropofol, or a pharmaceutically acceptable salt thereof, to the pharmaceutically acceptable acid is 1.5:1 or less.
-