



US012398434B2

(12) **United States Patent**
Monpoeho et al.

(10) **Patent No.:** **US 12,398,434 B2**

(45) **Date of Patent:** ***Aug. 26, 2025**

(54) **COMPOSITIONS AND METHODS FOR
DETECTING A BIOLOGICAL
CONTAMINANT**

(71) Applicant: **Regeneron Pharmaceuticals, Inc.**,
Tarrytown, NY (US)

(72) Inventors: **Serge Monpoeho**, East Greenbush, NY
(US); **Sheldon Mink**, Rensselaer, NY
(US); **Paul Vescio**, Malta, NY (US)

(73) Assignee: **Regeneron Pharmaceuticals, Inc.**,
Tarrytown, NY (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 239 days.

This patent is subject to a terminal dis-
claimer.

(21) Appl. No.: **18/061,712**

(22) Filed: **Dec. 5, 2022**

(65) **Prior Publication Data**

US 2023/0295747 A1 Sep. 21, 2023

Related U.S. Application Data

(60) Continuation of application No. 16/855,217, filed on
Apr. 22, 2020, now Pat. No. 11,549,154, which is a
division of application No. 15/080,859, filed on Mar.
25, 2016, now Pat. No. 10,669,594.

(60) Provisional application No. 62/139,321, filed on Mar.
27, 2015.

(51) **Int. Cl.**

C12Q 1/70 (2006.01)

C12Q 1/68 (2018.01)

C12Q 1/6876 (2018.01)

(52) **U.S. Cl.**

CPC **C12Q 1/701** (2013.01); **C12Q 1/6876**
(2013.01); **C12Q 2600/158** (2013.01); **C12Q**
2600/166 (2013.01)

(58) **Field of Classification Search**

CPC **C12Q 1/701**; **C12Q 1/6876**
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,683,202 A 7/1987 Mullis
5,583,236 A 12/1996 Brush
5,585,254 A 12/1996 Maxwell et al.
6,927,004 B2 8/2005 Eurlings et al.
7,087,411 B2 8/2006 Daly et al.
7,279,159 B2 10/2007 Daly et al.
7,771,997 B2 8/2010 Chen et al.
9,834,815 B2 12/2017 Brewer et al.
10,669,594 B2 * 6/2020 Monpoeho C12Q 1/701
11,549,154 B2 * 1/2023 Monpoeho C12Q 1/701
2002/0177222 A1 11/2002 Li
2003/0170612 A1 9/2003 Pichuanes et al.
2006/0166232 A1 7/2006 Vickery et al.
2007/0031850 A1 2/2007 Mounts et al.

2010/0331527 A1 12/2010 Davis et al.
2013/0109742 A1 * 5/2013 Hewitt C12N 15/8645
435/235.1

2015/0093749 A1 4/2015 Ling
2020/0248278 A1 8/2020 Monpoeho et al.

FOREIGN PATENT DOCUMENTS

CN 102206713 A 10/2011
DE 102006034844 B3 12/2007
EP 1077260 A1 2/2001
EP 2098599 A1 9/2009
EP 2397542 A1 12/2011
GB 2525024 A * 10/2015 C12Q 1/6806
JP 2014513921 A 6/2014
KR 20140006963 A 1/2014
WO WO-0146463 A2 6/2001
WO WO-03002753 A2 1/2003
WO WO-2004044247 A2 5/2004
WO WO-2012052158 A2 4/2012
WO WO-2012114312 A2 8/2012
WO WO-2013012708 A1 1/2013
WO WO-2014006432 A2 1/2014
WO WO-2014055515 A1 * 4/2014 A61K 38/162

OTHER PUBLICATIONS

Atkinson et al., Development of a real-time RT-PCR assay for the
detection of Crimean-Congo hemorrhagic fever virus. Vector Borne
Zoonotic Dis. Sep. 2012; 12(9):786-93. Epub Jan. 4, 2012. (Year:
2012).*

Genbank Accession No. CP032919—Agrobacterium tumefaciens
strain 15955 plasmid pAt15955, complete, submitted Oct. 9, 2018,
retrieved on Sep. 22, 2024 from <http://www.ncbi.nlm.nih.gov/nucleotide/CP032919>. (Year: 2018).*

Gerriets et al., Implementation of a T4 extraction control for
molecular assays of cerebrospinal fluid and stool specimens. J Mol
Diagn. Jan. 2008; 10(1):28-32. Epub Dec. 28, 2007. (Year: 2008).*

Ninove et al., 2011. RNA and DNA bacteriophages as molecular
diagnosis controls in clinical virology: a comprehensive study of
more than 45,000 routine PCR tests. PloS one, 6(2), e16142, pp. 1-7.
(Year: 2011).*

Altamirano, C., et al., "Analysis of CHO cells metabolic redistri-
bution in a glutamate-based defined medium in continuous culture",
Biotechnology Progress (2001); 17(6): 1032-1041.

Ashkenazi, A., et al., "Protection against endotoxic shock by a
tumor necrosis factor receptor immunoadhesin", Proceedings of the
National Academy of Sciences (1991); 88(23): 10535-10539.

(Continued)

Primary Examiner — Gary Benzion

Assistant Examiner — Olayinka A Oyeyemi

(74) *Attorney, Agent, or Firm* — Fenwick & West LLP

(57)

ABSTRACT

Provided are compositions and methods useful to the deter-
mination of whether a microbial contaminant is present in a
biological therapeutic production process. Specifically, an
artificial positive amplification control plasmid and unique
quantitative PCR detection probe are provided, which
enables the rapid and real-time detection of a false positive
result.

16 Claims, No Drawings

Specification includes a Sequence Listing.

(56)

References Cited

OTHER PUBLICATIONS

- Byrn, R. A., et al., "Biological properties of a CD4 immunoadhesin", *Nature* (Apr. 1990); 344: 667-670.
- Hollenbaugh, D., et al., "Construction of immunoglobulin fusion proteins", *Current Protocols in Immunology* (2002); 48(1): 10.19A.1-10.19A.11.
- Holliger, P., et al., "Diabodies": Small bivalent and bispecific antibody fragments, *Proceedings of the National Academy of Sciences USA* (1993); 90: 6444-6448.
- Huang, Y., et al., "Maximizing productivity of CHO cell-based fed-batch culture using chemically defined media conditions and typical manufacturing equipment", *Biotechnology progress* (2010); 26(5): 1400-1410.
- Innis, M. A., et al., "DNA sequencing with *Thermus aquaticus* DNA polymerase and direct sequencing of polymerase chain reaction-amplified DNA", *Proceedings of the National Academy of Sciences* (1988); 85(24): 9436-9440.
- Kaufman, R. J., "Use of recombinant DNA technology for engineering mammalian cells to produce proteins", *Large-Scale Mammalian Cell Culture*, 1st Edition (1990); 10: 15-69.
- Kipriyanov, S. M., et al., "Single-chain antibody streptavidin fusions: tetrameric bifunctional scFv-complexes with biotin binding activity and enhanced affinity to antigen", *Human Antibodies and Hybridomas* (1995); 6(3): 93-101.
- Kipriyanov, S. M., et al., "Recombinant single-chain Fv fragments carrying C-terminal cysteine residues: Production of bivalent and biotinylated miniantibodies", *Molecular Immunology* (1994); 31(14): 1047-1058.
- Poljak, R. J., et al., "Production and structure of diabodies," *Structure* (1994); 2(12): 1121-1123.
- Southwick, P. L., et al., "Cyanine dye labeling reagents-carboxymethylindocyanine succinimidyl esters", *Cytometry: The Journal of the International Society for Analytical Cytology* (1990); 11(3): 418-430.
- Sylvain, K., et al., "Rapid screening for HLA-B27 by a TaqMan-PCR assay using sequence-specific primers and a minor groove binder probe, a novel type of TaqMan™ probe", *Journal of Immunological Methods* (2004); 287(1-2): 179-186.
- Taggart, E. W., et al., "Use of heat labile UNG in an RT-PCR assay for enterovirus detection", *Journal of Virological Methods* (2002); 105(1): 57-65.
- Taylor, L. D., et al., "A transgenic mouse that expresses a diversity of human sequence heavy and light chain immunoglobulins", *Nucleic Acids Research* (1992); 20(23): 6287-6295.
- Ward, E. S., et al., "Binding activities of a repertoire of single immunoglobulin variable domains secreted from *Escherichia coli*", *Nature* (1989); 341: 544-546.
- [Author Unknown] "QIASymphony DNA Handbook", QIAGEN (Sep. 2010); 32 pages.
- Atkinson, B. et al., "Development of a real-time RT-PCR assay for the detection of Crimean-Congo hemorrhagic fever virus", *Vector Borne Zoonotic Dis.* (Sep. 2012); 12(9): 786-93.
- Berting, A., et al., "Virus susceptibility of Chinese hamster ovary (CHO) cells and detection of viral contaminations by adventitious agent testing", *Biotechnology and Bioengineering* (2010); 106(4): 598-607.
- Besselsen, D. G., et al., "Identification of novel murine parvovirus strains by epidemiological analysis of naturally infected mice", *Journal of General Virology* (2006); 87(6): 1543-1556.
- Cotmore, S. F., and Tattersall, P., "The autonomously replicating parvoviruses of vertebrates", *Advances in Virus Research* (1987); 33: 91-174.
- Cotmore, S. F., et al., "Replication initiator protein NS1 of the parvovirus minute virus of mice binds to modular divergent sites distributed throughout duplex viral DNA", *Journal of Virology* (2007); 81(23): 13015-13027.
- Database EMBL [Online] "Sequence 3 from Patent EP1077260" Retrieved from EBI Accession No. EM_PAT: AX137738, Database Accession No. AX137738, The Whole Document & EP 1 077 260 A1 (Deutsches Krebsforsch [DE] (Feb. 21, 2001); 1 page.
- Decaro, N., et al., "A real-time PCR assay for rapid detection and quantitation of canine parvovirus type 2 in the feces of dogs", *Veterinary Microbiology* (2005); 105(1): 19-28.
- Diez-Valcarce, M., et al., "Construction and analytical application of internal amplification controls (IAC) for detection of food supply chain-relevant viruses by real-time PCR-based assays", *Food Analytical Methods* (2011); 4(3): 437-445.
- Drosten, C., et al., "Evaluation of a new PCR assay with competitive internal control sequence for blood donor screening", *Transfusion* (2000); 40(6): 718-724.
- European Office Action for European Application No. 16734060.3, mailed Nov. 12, 2018, 7 pages.
- Gerriets, J. E., et al., "Implementation of a T4 extraction control for molecular assays of cerebrospinal fluid and stool specimens", *The Journal of Molecular Diagnostics* (2008); 10(1): 28-32.
- Gonçalves-De-Albuquerque, S. C., et al., "Tracking false-negative results in molecular diagnosis: proposal of a triplex-PCR based method for leishmaniasis diagnosis", *Journal of Venomous Animals and Toxins including Tropical Diseases* (2014); 20: 1-6.
- International Search Report and Written Opinion for PCT/US2016/024216, mailed Aug. 12, 2016, 14 pages.
- Jacoby, R. O. et al., "Rodent Parvovirus Infections", *Laboratory Animal Science* (Aug. 1996); 46(4): 370-380.
- Jones, M. D., et al., "Single-stranded M13 DNA: use as a cloning vector", *Nucleic Acids Research* (1986); 14(24): 10116.
- Kerr, A., and Nims, R., "Adventitious viruses detected in biopharmaceutical bulk harvest samples over a 10 Year Period", *PDA Journal of Pharmaceutical Science and Technology* (2010); 64(5): 481-485.
- Kothapalli, R., et al., "Problems associated with product enhancement reverse transcriptase assay using bacteriophage MS2 RNA as a template", *Journal of Virological Methods* (2003); 109(2): 203-207.
- Lee, A. V., et al., "Comparative evaluation of the QIAGEN QIASymphony® SP system and bioMérieux NucliSens easyMAG automated extraction platforms in a clinical virology laboratory", *Journal of Clinical Virology* (2011); 52(4): 339-343.
- Liu, H. F., et al., "Recovery and purification process development for monoclonal antibody production", *mAbs* (2010); 2(5): 480-499.
- Mackay, I. M., et al., "Real-time PCR in virology", *Nucleic Acids Research* (2002); 30(6): 1292-1305.
- Marras, S. A. E., "Selection of fluorophore and quencher pairs for fluorescent nucleic acid hybridization probes", *Methods in Molecular Biology: Fluorescent Energy Transfer Nucleic Acid Probes* (2006): 3-16.
- Marras, S. A. E., et al., "Efficiencies of fluorescence resonance energy transfer and contact-mediated quenching in oligonucleotide probes", *Nucleic Acids Research* (2002); 30(21): e122; 8 pages.
- Marras, S. A. E., "Interactive fluorophore and quencher pairs for labeling fluorescent nucleic acid hybridization probes", *Molecular Biotechnology* (2008); 38(3): 247-255.
- Merten, O.-W., "Virus contaminations of cell cultures—a biotechnological view", *Cytotechnology* (2002); 39(2): 91-116.
- Miesegaes, G., et al., "Analysis of viral clearance unit operations for monoclonal antibodies", *Biotechnology and Bioengineering* (2010); 106(2): 238-246.
- Müller, J., et al., "Development and validation of a real-time PCR assay for routine testing of blood donations for parvovirus B19 DNA", *Infusion Therapy and Transfusion Medicine* (2002); 29(5): 254-258.
- Moniwa, M., et al., "Performance of a foot-and-mouth disease virus reverse transcription-polymerase chain reaction with amplification controls between three real-time instruments", *Journal of Veterinary Diagnostic Investigation* (2007); 19(1): 9-20.
- Moody, M., et al., "Mouse minute virus (MMV) contamination—a case study: detection, root cause determination, and corrective actions", *PDA Journal of Pharmaceutical Science and Technology* (2011); 65(6): 580-588.
- Ninove, L., et al., "RNA and DNA bacteriophages as molecular diagnosis controls in clinical virology: a comprehensive study of more than 45,000 routine PCR tests", *PloS One* (2011); 6(2): e16142; 7 pages.

(56)

References Cited

OTHER PUBLICATIONS

- Redig, A. J., and Besselsen, D. G., "Detection of rodent parvoviruses by use of fluorogenic nuclease polymerase chain reaction assays", *Comparative Medicine* (2001); 51(4): 326-331.
- Sachsenröder, J., et al., "Simultaneous identification of DNA and RNA viruses present in pig faeces using process-controlled deep sequencing", *PloS One* (2012); 7(4): e34631; 11 pages.
- Sambrook, J., and Russell, D. W., "Cloning into Bacteriophage M13 Vectors", *Cold Spring Harbor Protocols* (2006); 3 pages.
- Shen, X. Z., et al., "A preliminary analysis of antineoplastic activity of parvovirus MVMP NS-1 proteins", *Cell Research* (1997); 7(2): 217-227.
- Shien, J-H., et al., "Identification of sequence changes in live attenuated goose parvovirus vaccine strains developed in Asia and Europe", *Avian Pathology* (2008); 37(5): 499-505.
- Skiaopoulos, M. H., et al., "Characterization of linker insertion and point mutations in the NS-1 gene of minute virus of mice: effects on DNA replication and transcriptional activation functions of NS-1", *Virology* (1992); 188(1): 122-134.
- Van Wezenbeek, P. M.G. F., et al., "Nucleotide sequence of the filamentous bacteriophage M13 DNA genome: comparison with phage fd", *Gene* (1980); 11(1-2): 129-148.
- Vieira, J. and Messing, J., "The pUC plasmids, an M13mp7-derived system for insertion mutagenesis and sequencing with synthetic universal primers", *Gene* (1982); 19(3): 259-268.
- Yu, Q., "Cloning into M13 bacteriophage vectors", *Methods in Molecular Biology* (1996); 58(Chapter 41): 343-348.
- Zhan, D., et al., "Detection of minute virus of mice using real time quantitative PCR in assessment of virus clearance during the purification of mammalian cell substrate derived biotherapeutics", *Biologicals* (2002); 30(4): 259-270.

* cited by examiner

1

COMPOSITIONS AND METHODS FOR DETECTING A BIOLOGICAL CONTAMINANT

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation of U.S. patent application Ser. No. 16/855,217, filed on Apr. 22, 2020, now U.S. Pat. No. 11,549,154, which is a divisional of U.S. patent application Ser. No. 15/080,859, filed Mar. 25, 2016, now U.S. Pat. No. 10,669,594, which claims the benefit of priority under 35 USC § 119(e) of U.S. Provisional Patent Application No. 62/139,321, filed 27 Mar. 2015. The contents of each of these applications is herein incorporated by reference in their entirety.

INCORPORATION BY REFERENCE OF SEQUENCE LISTING

The contents of the electronic sequence listing are incorporated herein by reference in their entirety: a computer readable format copy of the Sequence Listing (filename: REGE-008CO1US_SeqList_ST26.xml, date recorded: Apr. 30, 2023, file size: 97,011 bytes).

FIELD OF THE INVENTION

The present invention relates generally to the process of manufacturing biological molecules via cell culture. The present invention relates more specifically, but not exclusively, to compositions and methods for detecting biological contaminants in cell culture.

BACKGROUND OF THE INVENTION

Biopharmaceutical drugs, especially therapeutic antibodies, are produced by mammalian cell culture. Chinese hamster ovary (CHO) cells are the most commonly used host cell. These production systems are prone to adventitious and endogenous virus infection, which presents a potential safety problem for the biopharmaceutical drug. Viral clearance procedures and viral load measurements are therefore used to promote the safety of the drug. Steps employed to reduce viral load include nanofiltration, virus inactivation by heat or pH hold, and chromatography. Viral load and the effectiveness of virus removal can be monitored by time consuming infectivity assays or by fast quantitative assays such as real-time PCR or quantitative polymerase chain reaction (Q-PCR).

Q-PCR requires the proper negative and positive controls to be reliable. Nucleic acid extraction controls are added to test samples to control for proper nucleic acid extraction. If the nucleic acid extraction control is negative or outside of the expected recovery range during Q-PCR, then the sample is rejected. Conversely, if the nucleic acid extraction control is positive or within the expected recovery range during Q-PCR, nucleic acid extraction from the test sample is deemed reliable. A negative control is usually included in the Q-PCR assay, such as a sample-free buffer. The presence of a positive signal in the negative control might signify the contamination of the Q-PCR or nucleic acid extraction reagents with virus material.

A positive amplification control may also be included in a Q-PCR viral load assay. Such a positive control may include conserved viral nucleic acid sequences that are amplified using primers that amplify genuine viral contami-

2

nants. The failure of the Q-PCR to detect the positive control may indicate that the amplification procedure would have failed to detect a viral contaminant if one were present in the test sample.

The use of positive amplification controls, which emulate the target contaminant, creates its own problems. If the test sample is contaminated with even a slight amount of positive control, given the exquisite sensitivity of Q-PCR, the test sample may show a false positive. False positives can be mitigated to some extent by using a low level of positive amplification control, using segregated rooms for positive control work, using the UNG/dUTP system to selectively degrade PCR products containing dUTP, using single-use containers and displacement pipettes, and by thoroughly cleaning work areas and equipment. Regardless of the fastidious use of those mitigators, false positive results still occur during PCR testing.

In biopharmaceutical manufacturing, the risk of getting a false positive for a biological contaminant is non-negligible and may result in costly corrective actions. There is a great need for systems and methods to determine whether a given positive PCR result is a true positive or a false positive resulting from cross-contamination of the positive control. Applicants have developed and now disclose positive control compositions, systems and methods that permit the real-time determination of false positive Q-PCR signals.

SUMMARY OF THE INVENTION

Applicant has solved the problem of identifying in real-time whether a positive Q-PCR signal for a target contaminant in a test sample is a true positive or a false positive due to cross-contamination. Applicant has created a positive amplification control (PAC) plasmid that includes a biological contaminant target sequence (i.e., the positive control sequence) and a unique artificial plasmid-specific sequence. This unique artificial plasmid-specific sequence enables the assay technician to specifically identify the plasmid when it is in a sample. Thus, when a positive contaminant signal is detected and the artificial plasmid-specific sequence is determined not to be present, the technician can be confident that the result is a true positive result. Conversely, in the event of a false positive, the presence of the unique artificial plasmid-sequence allows the technician to quickly rule out the ostensible positive result as a false positive.

In some embodiments, the positive control unique artificial plasmid-specific sequence ("unique sequence"; a.k.a. PACP or positive amplification control polynucleotide) is detected using a fluorescently labeled artificial oligonucleotide detection probe ("unique detection probe" or "UDP") included in the Q-PCR reaction mix. The unique detection probe comprises a nucleic acid polymer covalently bound to a fluorophore and a quencher. The "unique" nucleic acid polymer is designed to specifically anneal to the unique sequence and incorporate into amplicon copies of the unique sequence during PCR. The "unique" nucleic acid polymer is designed to not recognize or anneal to any naturally occurring parvovirus under hybridization conditions employed in the operation of the subject assay. In one embodiment, the "unique" nucleic acid polymer comprises from 17 to 20 nucleotides, wherein no more than seven (7) to 10 internal consecutive nucleotides and no more than six (6) consecutive three-prime nucleotides are identical to any parvovirus sequence. In one embodiment, the "unique" nucleotide polymer comprises from 17 to 20 nucleotides, wherein no more than 7 to 10 consecutive nucleotides and no more than six (6) consecutive three-prime nucleotides are identical to

any parvovirus sequence set forth in SEQ ID NOs: 9 and 12-37. When the nucleic acid polymer of the unique detection probe is not incorporated into amplicon copies, the quencher remains in close proximity to the fluorophore. If the fluorophore is excited, then the quencher absorbs the emitted light and prevents that light from being detected (by FRET or contact quenching). When the nucleic acid polymer is incorporated into the amplicon copies (i.e., when the unique sequence is present in the sample), the fluorophore and the quencher are released from the unique detection probe and are therefore spatially separated. In that case, when the fluorophore is excited, the quencher is sufficiently far away that it cannot efficiently quench the emitted light. The emitted light can therefore be detected. Thus, when the unique sequence is present in the sample, the detectable emission wavelength increases in intensity as the PCR proceeds. When the unique sequence is absent, the quencher does its work and no emission wavelength is detected as the PCR advances. In one embodiment, the fluorophore is attached at or near the 5-prime end of the nucleic acid polymer, and the quencher at or near the 3-prime end. In an alternative embodiment, the fluorophore is attached at or near the 3-prime end of the nucleic acid polymer, and the quencher at or near the 5-prime end.

Any fluorophore-quencher pair now known or later discovered may be used in the practice of this invention. (See for example S. A. Marras, "Selection of fluorophore and quencher pairs for fluorescent nucleic acid hybridization probes," *Methods Mol. Biol.* 2006; 335:3-16.) In some embodiments, the fluorophore has an excitation wavelength of 495 nm, 538 nm, or 646 nm and an emission wavelength of 520 nm, 554 nm, or 669 nm, respectively. In some embodiments, the quencher is a dye with an absorbance peak of 430 nm to 672 nm. In some embodiments, the quencher is selected from the group consisting of DDQ-I, Dabcyl, Eclipse, Iowa Black FQ, BHQ-1, QSY-7, BHQ-2, DDDQ-II, Iowa Black RQ, QSY-21, and DHQ-3. In one embodiment, the fluorophore has an excitation wavelength of 495 nm and an emission wavelength of 520 nm, for example FAM, and the quencher is BHQ-1. In one embodiment, the nucleic acid polymer comprises the nucleic acid sequence of SEQ ID NO:3 (5'-TGTCGATGGCGAATGGCTA-3').

One aspect of the invention is the unique detection probe itself, as described above, containing the nucleic acid polymer and the linked fluorophore and quencher. Other aspects of the invention include the positive amplification control (PAC) plasmid itself, which contains a biological contaminant target sequence and the unique artificial plasmid-specific sequence, and the use of that plasmid as a positive control to assess the presence of a target biological contaminant in a cell culture. The PAC plasmid is used to control for the successful PCR amplification reaction designed to amplify target biological contaminant sequences. For example, a separate and parallel PCR reaction, which contains the identical components and is run under identical parameters as the test sample, but containing the positive control plasmid in lieu of a test sample, is run. If the positive control plasmid-containing sample yields a positive "contaminant" signal, but the test sample does not, then one might conclude that the test sample is devoid of the target biological contaminant. In some embodiments, the test sample is obtained from a mammalian cell culture, such as a bioreactor culture containing CHO cells engineered to produce a therapeutic protein of interest.

In one embodiment, the PAC plasmid contains (a) a target amplification polynucleotide (TAP) sequence, such as for example a parvovirus nucleic acid sequence or sequence of

another target contaminant, and (b) a plasmid amplification control polynucleotide (PACP) sequence. The PACP sequence (sense or Crick strand) is complementary to the nucleic acid polymer of the unique sequence probe (antisense or Watson strand).

In one embodiment, the PAC plasmid is deployed in a separate Q-PCR reaction in parallel with a Q-PCR reaction containing a test sample. The TAP sequence is designed to be representative of the target biological contaminant. For example, if the test sample Q-PCR reaction fails to produce TAP amplicons, and the positive control (i.e., PAC plasmid-containing sample) Q-PCR fails to produce TAP amplicons, then one may assume that the Q-PCR reaction failed. In one embodiment, the target biological contaminant is a rodent parvovirus and the TAP sequence comprises a rodent parvovirus sequence. In one embodiment, the TAP sequence comprises all or part of the parvovirus NS1 sequence, which is conserved across several rodent parvovirus strains. See Cotmore, et al., "Replication Initiator Protein NS1 of the parvovirus Minute Virus of Mice Binds to Modular Divergent Sites Distributed throughout Duplex Viral DNA," *J. Virol.* 2007 December; 81(23):13015-13027. In some embodiments, the several rodent parvovirus strains include minute virus of mice prototype strain (MVMp), minute virus of mice immunosuppressive strain (MVMi), minute virus of mice Cutter strain (MVMc), mouse parvovirus 1b (MPV-1b), mouse parvovirus 1a (MPV-1a), mouse parvovirus 1c (MVP-1c), hamster parvovirus (HaPV), Toolan's parvovirus (H-1), Kilham rat virus (KRV), Rat parvovirus 1a (RPV-1a), rat minute virus (RMV), and University of Massachusetts strain of rat virus L (RV-Umass). See O.-W. Merten, "Virus Contaminations of Cell Cultures—A Biotechnological View," *Cytotechnology.* 2002 July; 39(2):91-116. In one embodiment, the TAP sequence comprises the nucleic acid sequence of SEQ ID NO:1, SEQ ID NO:2, and the complement of SEQ ID NO:4.

In other aspects, the invention is directed to a PCR cocktail composition and method of using the PCR cocktail to detecting a target contaminant in a test sample and ruling out a false positive due to contamination of the test sample with the PAC.

In one embodiment, the PCR cocktail contains inter alia a target contaminant specific forward oligonucleotide primer, target contaminant specific oligonucleotide detection probe, an artificial oligonucleotide detection probe, such as the unique detection probe (UDP) described above, and a target contaminant specific reverse oligonucleotide primer. In an embodiment in which the target contaminant is a rodent parvovirus, the PCR cocktail contains inter alia a rodent parvovirus specific forward oligonucleotide primer, a rodent parvovirus specific oligonucleotide detection probe, an artificial oligonucleotide detection probe, such as the unique detection probe (UDP) described above, and a rodent parvovirus specific reverse oligonucleotide primer. Each detection probe (i.e., target contaminant specific oligonucleotide detection probe, such as rodent parvovirus specific oligonucleotide detection probe, and artificial oligonucleotide detection probe) contains a nucleic acid sequence linked to a fluorophore at one end (either 5-prime or 3-prime) and a quencher at the other end (either 3-prime or 5-prime, respectively).

Here, the rodent parvovirus specific forward oligonucleotide primer and the nucleic acid sequence of the rodent parvovirus specific oligonucleotide detection probe hybridize to the antisense strand of the parvovirus. The rodent parvovirus specific reverse oligonucleotide primer hybridizes to the sense strand of the parvovirus. In one embodi-

ment, the parvovirus sequence to which the primers and parvovirus specific oligonucleotide probe hybridize is a conserved rodent parvovirus sequence. In one case, the conserved parvovirus sequence is a parvovirus NS1 sequence, such as for example the nucleic acid sequence described in SEQ ID NO:9. By using a conserved sequence, a single probe will be effective to detect multiple strains of rodent parvovirus.

In one embodiment, the artificial oligonucleotide detection probe does not hybridize to a parvovirus nucleic acid or any biological contaminant sequence. The nucleic acid of the artificial oligonucleotide detection probe is synthetic and will not hybridize with any stringency to any biological contaminants nucleic acid sequence. In one embodiment, the nucleic acid of the artificial oligonucleotide detection probe (a.k.a. "unique" nucleic acid polymer or unique sequence) is designed to not recognize or anneal to any naturally occurring parvovirus under hybridization conditions employed in the operation of the subject assay. In one embodiment, the "unique" nucleic acid polymer comprises from 17 to 20 nucleotides, wherein no more than seven (7) to 10 internal consecutive nucleotides and no more than six (6) consecutive three-prime nucleotides are identical to any parvovirus sequence. In one embodiment, the "unique" nucleotide polymer comprises from 17 to 20 nucleotides, wherein no more than 7 to 10 consecutive nucleotides and no more than six (6) consecutive three-prime nucleotides are identical to any parvovirus sequence set forth in SEQ ID NOs: 9 and 12-37. However, the nucleic acid of the artificial oligonucleotide detection probe (i.e., the unique sequence) hybridizes to the PACP sequence of the PAC plasmid. Thus, the artificial oligonucleotide detection probe detects the PAC plasmid, but not a parvovirus or other biological contaminant sequence.

In one embodiment, the PCR cocktail can be used to determine whether the PAC plasmid is present in the test biological sample, rendering a false positive result. Here, if the test sample shows a positive Q-PCR signal for the parvovirus specific oligonucleotide probe, and a negative Q-PCR signal for the artificial oligonucleotide detection probe, then the test sample is presumed to be free of PAC contamination (i.e., true positive).

In one embodiment, the target contaminant specific forward oligonucleotide primer comprises the sequence of SEQ ID NO:1; the target contaminant specific oligonucleotide detection probe nucleic acid comprises the sequence of SEQ ID NO:2; the artificial oligonucleotide detection probe nucleic acid (i.e., the UDP) comprises the sequence of SEQ ID NO:3; and the target contaminant specific reverse oligonucleotide primer comprises the sequence of SEQ ID NO:4.

In other aspects, the invention provides a system and method for detecting a biological contaminant in a test sample. Here, the test sample is a cell culture, such as for example an industrial scale mammalian cell culture for the production of a therapeutic protein. Mammalian cells useful in the practice of this invention include, but are not limited to CHO cells, CHO-K1 cells, and EESYR cells (see U.S. Pat. No. 7,771,997). The system and method include—in addition to the primers, probes, cocktails and PAC plasmid used as described above—the use of a nucleic acid extraction control (NEC). In one embodiment, the NEC is an M13K07 phage, which is included in the test sample prior to nucleic acid extraction. If the nucleic acid extraction is adequate for the purposes of detecting contaminant DNA or RNA, then the NEC nucleic acid (e.g., M13K07 nucleic acid) is detected via Q-PCR in the "spiked" test sample. In a particular embodiment, the Q-PCR reaction admixture

contains a target contaminant specific forward oligonucleotide primer, target contaminant specific oligonucleotide detection probe, an artificial oligonucleotide detection probe, such as the unique detection probe (UDP) described above, a target contaminant specific reverse oligonucleotide primer, an NEC specific forward oligonucleotide primer, an NEC specific oligonucleotide detection probe, and an NEC specific reverse oligonucleotide primer. In one embodiment, the test sample is taken from a therapeutic protein production cell culture, which is spiked with an NEC (e.g., an M13K07 phage) prior to nucleic acid extraction and subsequent Q-PCR analysis.

In a specific embodiment, the target contaminant is a rodent parvovirus, in which case (1) the target contaminant specific forward oligonucleotide primer is a rodent parvovirus specific forward oligonucleotide primer, more specifically comprising the sequence of SEQ ID NO:1; (2) the target contaminant specific oligonucleotide detection probe is a rodent parvovirus specific oligonucleotide detection probe, more specifically comprising the nucleic acid sequence of SEQ ID NO:2; (3) the artificial oligonucleotide detection probe is the unique detection probe (UDP), more specifically comprising the nucleic acid sequence of SEQ ID NO:3; (4) the target contaminant specific reverse oligonucleotide primer is a rodent parvovirus specific reverse oligonucleotide primer, more specifically comprising the nucleic acid sequence of SEQ ID NO:4; (5) the NEC specific forward oligonucleotide primer is an M13 forward oligonucleotide primer, more specifically comprising the nucleic acid sequence of SEQ ID NO:5; (6) the NEC specific oligonucleotide detection probe is an M13 detection probe, more specifically comprising the nucleic acid sequence of SEQ ID NO:6; and (7) the NEC specific reverse oligonucleotide primer is an M13 reverse oligonucleotide primer, more specifically comprising the nucleic acid of SEQ ID NO:8. In one embodiment of the method, if an NEC Q-PCR signal is detected (i.e., the signal is within the expected recovery range), then one can assume that the nucleic acid extraction of the test sample was successful. On the other hand, if no NEC signal is detected (i.e., the signal is outside the expected recovery range), then one can assume that the nucleic acid extraction of the test sample failed, and any negative Q-PCR target contaminant signal is considered invalid (i.e., false negative).

DETAILED DESCRIPTION OF THE INVENTION

Before the present invention is described, it is to be understood that this invention is not limited to particular methods and experimental conditions described, as such methods and conditions may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only by the appended claims.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. As used herein, the term "about," when used in reference to a particular recited numerical value, means that the value may vary from the recited value by no more than 1%. For example, as used herein, the expression "about 100" includes 99 and 101 and all values in between (e.g., 99.1, 99.2, 99.3, 99.4, etc.).

Although any methods and materials similar or equivalent to those described herein can be used in the practice of the

present invention, the preferred methods and materials are now described. All publications mentioned herein are incorporated by reference in their entirety. Other embodiments will become apparent from a review of the ensuing detailed description.

In order that the invention herein described may be fully understood, the following detailed description is set forth.

The invention relates to improved materials and methods for detecting any biological contaminant in any cell culture that produces a therapeutic protein. Specifically, the invention relates to quantitative polymerase chain reaction (Q-PCR) materials and methods that incorporate a positive amplification control element or step that is easily detected to eliminate false positives.

PCR and Quantitative PCR

As used herein, the phrase “polymerase chain reaction” (“PCR”) means a method for making copies of a nucleic acid (e.g., DNA) by employing multiple cycles of denaturation (the separation of template DNA strands), annealing (the hybridization of single stranded oligonucleotides to the single stranded template DNA strands), and DNA synthesis (DNA polymerase catalyzes the synthesis of a new DNA strand primed from the 3-prime end of the hybridized oligonucleotide, using the template DNA strand as a template). For amplification to occur, at least two different oligonucleotide primers (known as “primer” for short) are used in the PCR reaction. One primer, which is generally called the forward primer, hybridizes to the antisense strand of the template DNA and forms the 5-prime end of the newly synthesized sense strand. The other primer, which is generally called the reverse primer, hybridizes to the sense strand of the template DNA and forms the 5-prime end of the newly synthesized antisense strand. At each cycle, each template strand is copied to form a new double stranded DNA molecule, which is also known as an “amplicon”. Thus, with non-limiting amounts of oligonucleotide primers, DNA polymerase (i.e., Taq polymerase or other thermostable DNA polymerase; see Innis et al., *DNA sequencing with Thermus aquaticus DNA polymerase and direct sequencing of polymerase chain reaction-amplified DNA*, 85(24) Proc Natl Acad Sci USA. 9436-40 (1988)), and nucleotide triphosphates, the number of DNA molecules (templates and amplicons) doubles at each cycle. PCR is described in U.S. Pat. No. 4,683,202 (issued Jul. 28, 1987). See also PCR Primer: A Laboratory Manual (Carl W. Dieffenbach & Gabriela S. Dveksler eds., 1995).

As used herein, the term “cycle” means a single round of (1) DNA strand melting called “denaturation”, followed by (2) hybridizing of oligonucleotide primers to the resultant single-stranded DNA by the rules of base-pairing, a process called “annealing”, and (3) polymerization of a new strand of DNA starting at the 3-prime end of the oligonucleotide primer and moving in a 5-prime to 3-primer direction, a process called “amplification” or “extension”. Generally, polymerization uses a DNA polymerase enzyme, such as Taq polymerase, to catalyze the formation of phosphodiester bonds between adjacent deoxynucleotide triphosphates (“dNTPs”), which are laid down along the exposed single-stranded template DNA by hydrogen bonding according to the rules of base pairing. Denaturation, annealing, and amplification are performed at certain temperatures based in part upon the GC content of the DNA template and the oligonucleotide primers, and the length of the DNA strand to be copied. The temperature of denaturation and annealing, as well as the ionic strength of the reaction buffer controls the stringency of hybridization and the fidelity of DNA copying.

“Quantitative PCR” or “qPCR” or “Q-PCR” (also known as “real time PCR”) is a type of PCR that enables the monitoring of amplicon formation during the PCR cycling process. Q-PCR can be used to quantify the amount of a specific template DNA in a sample. Q-PCR incorporates at least one oligonucleotide detection probe in the reaction mix in addition to the forward oligonucleotide primer and the reverse oligonucleotide primer. The detection probe is a single-stranded oligonucleotide that hybridizes to the sense or antisense strand of the target template DNA somewhere between the forward primer binding site and the reverse primer binding site. During the annealing step, the oligonucleotide detection probe anneals to the single-stranded template. As polymerization occurs, the probe is cleaved and degraded by the 5-prime nuclease activity of the DNA polymerase. Thus, as amplification of the specific template sequence occurs, detection probes are degraded at an exponential rate.

Q-PCR oligonucleotide detection probes are generally constructed with an attached fluorophore (also known as a reporter fluorescent dye, or simply “reporter”) and an attached quencher. In most cases, the fluorophore is attached at or near the 5-prime end of the oligonucleotide and the quencher is attached at or near the 3-prime end of the oligonucleotide. However, any workable architecture may be used in the practice of this invention. When the oligonucleotide detection probe is intact, the fluorophore and the quencher are proximal such that the quencher absorbs light emitted by the excited fluorophore, thereby significantly reducing detectable fluorophore emission. When the oligonucleotide detection probe is cleaved or degraded, the fluorophore and quencher are released and consequently separated in space. The quencher is no longer close enough to quench the fluorophore emission. As more specific amplicons are formed, more oligonucleotide detection probes are cleaved, more fluorophores and quenchers are released such that more fluorophore/quencher pairs are separated, and the fluorescence emission amplitude increases. In other words, an increasing fluorophore emission signal correlates to the amount of specific target DNA in the sample. For a review of Q-PCR, see Ian M. Mackay et al., *Survey and Summary: Real-Time PCR in Virology*, 30(6) Nucleic Acids Research 1292-1305 (2002).

Fluorescence quenching can occur by direct contact between reporter and quencher (also known as static quenching) or by fluorescence resonance energy transfer (FRET) between the reporter and quencher when the reporter and quencher are both within the Forster radius of each other. A specific fluorophore can be excited by one or more specific wavelengths of light, or a range of wavelengths with a maximum. This is called the excitation wavelength. After a fluorophore is excited, it returns to a ground state and emits light at a longer wavelength than the excitation wavelength. This is called the emission wavelength. During FRET, a second fluorophore, dye, lanthanide series molecule, or the like, which has an absorbance spectrum that matches or overlaps the emission spectrum of the fluorophore, absorbs the light emitted by the excited fluorophore within the Forster radius, thereby quenching or reducing the wavelength of the fluorophore emission. Contact or static quenching results when the reporter and quencher form a ternary complex at the ground state of the fluorophore. This ternary complex is non-fluorescent, i.e., essentially non-excitable and therefore does not emit light at an expected emission wavelength.

For a review on static quenching and FRET, see Salvatore A. E. Marras et al., *Efficiencies of Fluorescence Resonance*

Energy Transfer and Contact-Mediated Quenching in Oligonucleotide Probes, 30(21) Nucleic Acids Research e122, pp. 1-8 (2002). Marras et al. also discuss the selection of reporter/quencher pairs for use in the application of Q-PCR. Nucleic Acids

The terms “polynucleotide”, “oligonucleotide”, “probe”, “primer” or “nucleotide primer” or “oligonucleotide primer”, “template” or “template nucleic acid” or “template DNA” are used herein according to their ordinary meaning to one of ordinary skill in the molecular biological arts. For a detailed explanation of each, see for example PCR Primer: A Laboratory Manual (Carl W. Dieffenbach & Gabriela S. Dveksler eds., 1995)

As used herein, “amplicon” refers to a DNA product resulting from the amplification of a template nucleic acid sequence through PCR. As PCR proceeds and template is amplified, newly formed DNA amplicons serve as templates for subsequent rounds of DNA synthesis.

Cell Cultures

The invention is directed to an improved Q-PCR method for detecting biological contaminants in a cell culture. Cell cultures are often used to produce complex biological molecules for therapeutic use, such as antibodies, trap molecules and Fc fusion proteins. These cultures should remain free of biological contaminants. The detection of contaminants is important to determine whether a particular batch is to be discarded or subject to remediation.

Cell cultures include culture media and cells usually derived from a single cell line. Here, the cell line comprises cells capable of producing a biotherapeutic protein. Examples of cell lines that are routinely used to produce protein biotherapeutics include inter alia primary cells, BSC cells, HeLa cells, HepG2 cells, LLC-MK cells, CV-1 cells, COS cells, VERO cells, MDBK cells, MDCK cells, CRFK cells, RAF cells, RK cells, TCMK-1 cells, LLCPK cells, PK15 cells, LLC-RK cells, MDOK cells, BHK cells, BHK-21 cells, CHO cells, CHO-K1 cells, NS-1 cells, MRC-5 cells, WI-38 cells, 3T3 cells, 293 cells, Per.C6 cells and chicken embryo cells. A Chinese hamster ovary (CHO) cell line or one or more of several specific CHO cell variants, such as the CHO-K1 cell line are optimized for large-scale protein production. The EESYR® cell line is a specialized CHO cell line optimized for enhanced production of proteins of interest. For a detailed description of EESYR® cells, see U.S. Pat. No. 7,771,997 (issued Aug. 10, 2010).

“Cell culture” or “culture” means the growth and propagation of cells outside of a multicellular organism or tissue. Suitable culture conditions for mammalian cells are known in the art. See for example *Animal cell culture: A Practical Approach* (D. Rickwood, ed., 1992). Mammalian cells may be cultured in suspension or attached to a solid substrate. Fluidized bed bioreactors, hollow fiber bioreactors, roller bottles, shake flasks, or stirred tank bioreactors, with or without microcarriers, and operated in a batch, fed batch, continuous, semi-continuous, or perfusion mode are available for mammalian cell culture. Cell culture media or concentrated feed media may be added to the culture continuously or at intervals during the culture (i.e., batch fed). For example, a culture may be fed once per day, every other day, every three days, or may be fed when the concentration of a specific medium component, which is being directly or indirectly monitored, falls outside a desired range.

Animal cells, such as CHO cells or EESYR® cells, may be cultured in small scale cultures, such as in 125 ml containers having about 25 ml of media, 250 ml containers having about 50 to 100 ml of media, 500 ml containers having about 100 to 200 ml of media. Alternatively, cultures

may be large scale such as for example 1000 ml containers having about 300 to 1000 ml of media, 3000 ml containers having about 500 ml to 3000 ml of media, 8000 ml containers having about 2000 ml to 8000 ml of media, and 15000 ml containers having about 4000 ml to 15000 ml of media. Cultures for manufacturing can contain 10,000 L of media or more. Large scale cell cultures, such as for clinical manufacturing of protein therapeutics, are typically maintained for days or weeks while the cells produce the desired protein(s). During this time, samples of the culture may be removed and tested for the presence of biological contaminants.

Production of Therapeutic Proteins

The cell culture which is monitored for biological contamination may be used to produce a protein or other biological molecule of interest, such as a therapeutically effective antibody or other biopharmaceutical drug substance. The protein product (protein of interest) can be inter alia an antibody, a human antibody, a humanized antibody, a chimeric antibody, a monoclonal antibody, a multispecific antibody, a bispecific antibody, an antigen binding antibody fragment, a single chain antibody, a diabody, triabody or tetrabody, a Fab fragment or a F(ab')₂ fragment, an IgA antibody, an IgD antibody, an IgE antibody, an IgM antibody, an IgG antibody, an IgG1 antibody, an IgG2 antibody, an IgG3 antibody, or an IgG4 antibody. In one embodiment, the antibody is an IgG1 antibody. In one embodiment, the antibody is an IgG2 antibody. In one embodiment, the antibody is an IgG4 antibody.

The protein of interest can be a recombinant protein that contains an Fc moiety and another domain, (e.g., an Fc-fusion protein). An Fc-fusion protein can be a receptor Fc-fusion protein, which contains one or more of one or more extracellular domain(s) of a receptor coupled to an Fc moiety. In some case, the Fc moiety comprises a hinge region followed by a CH2 and CH3 domain of an IgG. In some cases, the receptor Fc-fusion protein contains two or more distinct receptor chains that bind to either a single ligand or multiple ligands. For example, an Fc-fusion protein is a trap, such as for example an IL-1 trap (e.g., rilonacept, which contains the IL-1RACp ligand-binding region fused to the IL-1R1 extracellular region fused to an Fc domain of hIgG1; see U.S. Pat. No. 6,927,004), or a VEGF trap (e.g., aflibercept, which contains the Ig domain 2 of the VEGF receptor Flt1 fused to the Ig domain 3 of the VEGF receptor Flk1 fused to an Fc domain of hIgG1; see U.S. Pat. Nos. 7,087,411 and 7,279,159).

The present invention is not limited to any particular type of cell or cell line for protein production. Examples of cell types suitable for protein production include mammalian cells, such as a CHO-derived cell like EESYR®, insect cells, avian cells, bacterial cells, and yeast cells. The cells may be stem cells or recombinant cells transformed with a vector for recombinant gene expression, or cells transfected with a virus for producing viral products. The cells may contain a recombinant heterologous polynucleotide construct that encodes a protein of interest. That construct may be an episomal (such as an extrachromosomal plasmid or fragment) or it may be physically integrated into the genome of the cell. The cells may also produce a protein of interest without having that protein encoded on a heterologous polypeptide construct. In other words, the cell may naturally encode the protein of interest, such as a B-cell producing an antibody. Methods and vectors for genetically engineering cells or cell lines to express a protein of interest are well known to those of ordinary skill in the art. For example, various techniques are illustrated in *Current Protocols in*

Molecular Biology, Ausubel et al., eds. (Wiley & Sons, New York, 1988, and quarterly updates); Sambrook et al., *Molecular Cloning: A Laboratory Manual* (Cold Spring Laboratory Press, 1989); Kaufman, R. J., *Large Scale Mammalian Cell Culture*, 1990, pp. 15-69. A wide variety of cell lines suitable for growth in culture are available from the American Type Culture Collection (Manassas, Va.) and commercial vendors.

The cells may also be primary cells, such as chicken embryo cells, or primary cell lines. Examples of useful cells include BSC cells, LLC-MK cells, CV-1 cells, COS cells, VERO cells, MDBK cells, MDCK cells, CRFK cells, RAF cells, RK cells, TCMK-1 cells, LLC-PK cells, PK15 cells, LLC-RK cells, MDOK cells, BHK-21 cells, chicken embryo cells, NS-1 cells, MRC-5 cells, WI-38 cells, BHK cells, 293 cells, Per.C6 cells and CHO cells. In various embodiments, the cell line is a CHO cell derivative, such as CHO-K1, CHO DUX B-11, CHO DG-44, Veggie-CHO, GS-CHO, S-CHO, CHO lec mutant lines, or an EESYR® cell line.

In one particular scenario, the cell is a CHO cell derivative such as a EESYR® cell that ectopically (heterologously) expresses a protein. That protein comprises an immunoglobulin heavy chain region, such as a CH1, CH2, or CH3 region. In one embodiment, the protein comprises a human or rodent immunoglobulin CH2 and CH3 region. In one embodiment, the protein comprises a human or rodent immunoglobulin CH1, CH2, and CH3 region. In one embodiment, the protein comprises a hinge region and a CH1, CH2, and CH3 region. In a specific embodiment, the protein comprises an immunoglobulin heavy chain variable domain. In a specific embodiment, the protein comprises an immunoglobulin light chain variable domain. In a specific embodiment, the protein comprises an immunoglobulin heavy chain variable domain and an immunoglobulin light chain variable domain. In a specific embodiment, the protein is an antibody, such as a human antibody, a rodent antibody, or a chimeric human/rodent antibody (e.g., human/mouse, human/rat, or human hamster).

A protein production phase of a cell culture can be conducted at any scale of culture, from individual flasks and shaker flasks or wave bags, to one-liter bioreactors, and to large scale industrial bioreactors. A large scale process can be conducted in a volume of about 100 liters to 20,000 liters or more. One or more of several means may be used to control protein production, such as temperature shift or chemical induction. The growth phase of the cell may occur at a higher temperature than the production phase during which protein is expressed and/or secreted. For example, the growth phase may occur at a first temperature of about 35° C. to 38° C., and the production phase may occur at a second temperature of about 29° C. to 37° C., optionally from about 30° C. to 36° C. or from about 30° C. to 34° C. In addition, chemical inducers of protein production, such as caffeine, butyrate, tamoxifen, estrogen, tetracycline, doxycycline, and hexamethylene bisacetamide (HMB A), may be added concurrently, before, or after a temperature shift. If inducers are added after a temperature shift, they can be added from one hour to five days after the temperature shift, such as from one to two days after the temperature shift. Production cell cultures may be run as continuous feed culture system, as in a chemostat (see C. Altamirano et al., *Biotechnol Prog.* 2001 November-December; 17(6):1032-41), or according to a fed-batch (batch-fed) process (Huang, 2010).

Therapeutic Protein Products

As used herein "peptide," "polypeptide" and "protein" are used interchangeably throughout and refer to a molecule comprising two or more amino acid residues joined to each

other by a peptide bond. Peptides, polypeptides and proteins may also include modifications such as glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, alkylation, hydroxylation and ADP-ribosylation. Peptides, polypeptides, and proteins can be of scientific or commercial interest, including protein-based drugs. Peptides, polypeptides, and proteins include, among other things, antibodies and chimeric or fusion proteins. Peptides, polypeptides, and proteins are produced by recombinant animal cell lines using cell culture methods.

"Antibody" refers to an immunoglobulin molecule consisting of four polypeptide chains, two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds. Each heavy chain has a heavy chain variable region (HCVR or VH) and a heavy chain constant region. The heavy chain constant region contains three domains, CH1, CH2 and CH3. Each light chain has a light chain variable region and a light chain constant region. The light chain constant region consists of one domain (CL). The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR). Each VH and VL is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. The term "antibody" includes reference to both glycosylated and non-glycosylated immunoglobulins of any isotype or subclass. The term "antibody" includes antibody molecules prepared, expressed, created or isolated by recombinant means, such as antibodies isolated from a host cell transfected to express the antibody. The term antibody also includes bispecific antibody, which includes a heterotetrameric immunoglobulin that can bind to more than one epitope. Bispecific antibodies are generally described in US Patent Application Publication No. 2010/0331527, which is incorporated herein by reference.

The term "antigen-binding portion" of an antibody (or "antibody fragment"), refers to one or more fragments of an antibody that retain the ability to specifically bind to an antigen. Examples of binding fragments encompassed within the term "antigen-binding portion" of an antibody include (i) a Fab fragment, a monovalent fragment consisting of the VL, VH, CL and CH1 domains; (ii) a F(ab')₂ fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) a Fd fragment consisting of the VH and CH1 domains; (iv) a Fv fragment consisting of the VL and VH domains of a single arm of an antibody, (v) a dAb fragment (Ward et al. (1989) *Nature* 241:544-546), which consists of a VH domain, (vi) an isolated CDR, and (vii) an scFv, which consists of the two domains of the Fv fragment, VL and VH, joined by a synthetic linker to form a single protein chain in which the VL and VH regions pair to form monovalent molecules. Other forms of single chain antibodies, such as diabodies are also encompassed under the term "antibody" (see e.g., Holliger et al. (1993) *PNAS USA* 90:6444-6448; Poljak et al. (1994) *Structure* 2:1121-1123).

An antibody or antigen-binding portion thereof may be part of a larger immunoadhesion molecule, formed by covalent or noncovalent association of the antibody or an antibody portion with one or more other proteins or peptides. Examples of such immunoadhesion molecules include use of the streptavidin core region to make a tetrameric scFv molecule (Kipriyanov et al. (1995) *Human Antibodies and Hybridomas* 6:93-101) and use of a cysteine residue, a marker peptide and a C-terminal polyhistidine tag to make bivalent and biotinylated scFv molecules (Kipriyanov et al.

(1994) Mol. Immunol. 31:1047-1058). Antibody portions, such as Fab and F(ab')₂ fragments, can be prepared from whole antibodies using conventional techniques, such as by papain or pepsin digestion of whole antibodies. Moreover, antibodies, antibody portions and immunoadhesion molecules can be obtained using standard recombinant DNA techniques (see Sambrook et al., 1989).

The term "human antibody" is intended to include antibodies having variable and constant regions derived from human germline immunoglobulin sequences. The human antibodies of the invention may include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or site-specific mutagenesis in vitro or by somatic mutation in vivo), for example in the CDRs and in particular CDR3. However, the term "human antibody", as used herein, does not include antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences.

The term "recombinant human antibody", as used herein, is intended to include all human antibodies that are prepared, expressed, created or isolated by recombinant means, such as antibodies expressed using a recombinant expression vector transfected into a host cell, antibodies isolated from a recombinant, combinatorial human antibody library, antibodies isolated from an animal (e.g., a mouse) that is transgenic for human immunoglobulin genes (see e.g., Taylor et al. (1992) Nucl. Acids Res. 20:6287-6295) or antibodies prepared, expressed, created or isolated by any other means that involves splicing of human immunoglobulin gene sequences to other DNA sequences. Such recombinant human antibodies have variable and constant regions derived from human germline immunoglobulin sequences. In certain embodiments, however, such recombinant human antibodies are subjected to in vitro mutagenesis (or, when an animal transgenic for human Ig sequences is used, in vivo somatic mutagenesis) and thus the amino acid sequences of the VH and VL regions of the recombinant antibodies are sequences that, while derived from and related to human germline VH and VL sequences, may not naturally exist within the human antibody germline repertoire in vivo.

"Fc fusion proteins" comprise part or all of two or more proteins, one of which is an Fc portion of an immunoglobulin molecule, which are not otherwise found together in nature. Preparation of fusion proteins comprising certain heterologous polypeptides fused to various portions of antibody-derived polypeptides (including the Fc domain) has been described, e.g., by Ashkenazi et al., Proc. Natl. Acad. Sci. USA 88: 10535, 1991; Byrn et al., Nature 344:677, 1990; and Hollenbaugh et al., "Construction of Immunoglobulin Fusion Proteins", in Current Protocols in Immunology, Suppl. 4, pages 10.19.1-10.19.11, 1992. "Receptor Fc-fusion proteins" comprise one or more extracellular domain(s) of a receptor coupled to an Fc moiety, which in some embodiments comprises a hinge region followed by a CH2 and CH3 domain of an immunoglobulin. In some embodiments, the Fc-fusion protein contains two or more distinct receptor chains that bind to a one or more ligand(s). For example, an Fc-fusion protein is a trap, such as for example an IL-1 trap (e.g., rilonacept, which contains the IL-1RAcP ligand binding region fused to the IL-1R1 extracellular region fused to Fc of hIgG1; see U.S. Pat. No. 6,927,004), or a VEGF trap (e.g., aflibercept, which contains the Ig domain 2 of the VEGF receptor Flt1 fused to the Ig domain 3 of the VEGF receptor Flk1 fused to Fc of hIgG1; see U.S. Pat. Nos. 7,087,411 and 7,279,159).

Biological Contaminants

As used herein, the term "biological contaminant" means any unwanted, undesirable, harmful or potentially harmful biological entity. Those entities include inter alia prions (the etiological cause of bovine/transmissible spongiform encephalopathy), virions, viruses, mycoplasma, other bacteria, contaminating metazoan cells, DNAs, RNAs, transposons, other transposable elements, yeast, other fungi, algae, protists, and other adventitious and endogenous agents. Of particular concern to biotherapeutic production processes that use rodent cells (like CHO cells and derivatives of CHO cells) are adventitious viruses associated with cells or media or manufacturing raw materials. Contamination of cell culture bulk process materials and the resultant drug product poses a direct risk to the patient as well as the indirect risk of interrupting the supply of medicine.

A non-exhaustive list of adventitious and/or endogenous agents that can infect CHO cell cultures includes: single stranded (–) RNA viruses such as cache valley virus, influenza A/B virus, parainfluenza 1/2/3, simian virus 5, mumps virus, bovine respiratory syncytial virus, and vesicular stomatitis virus; single stranded (+) RNA viruses such as bovine coronavirus, vesivirus 2117, encephalomyocarditis virus, coxsackie virus B-3, semliki forest virus, and sindbis virus; double stranded RNA viruses such as bluetongue virus, epizootic hemorrhagic disease virus, and reovirus 1/2/3; single stranded DNA viruses, such as porcine circovirus 1, and the particularly problematic parvoviruses that include mice minute virus (also known as minute virus of mice); and double stranded DNA viruses such as adenovirus and pseudorabies virus. Of these potential adventitious agents, four viruses dominate bulk harvest samples from CHO cell cultures across various manufacturers. Those viruses are reovirus type 2, cache valley virus, epizootic disease hemorrhagic virus, and the rodent parvovirus mouse minute virus. For a detailed review of adventitious viral contaminations of CHO cell cultures, see Andreas Berting et al., *Virus Susceptibility of Chinese Hamster Ovary (CHO) Cells and Detection of Viral Contaminations by Adventitious Agent Testing*, 106(4) Biotechnology and Bioengineering 598-607 (2010), and Andrew Kerr & Raymond Nims, *Adventitious Viruses Detected in Biopharmaceutical Bulk Harvest Samples over a 10 Year Period*, 64(5) PDA Journal of Pharmaceutical Science & Technology 481-485 (2010).

Adventitious agent testing falls into two general categories. The first is a classical virology approach using an in vitro virus assay. Here, the test sample is applied to an indicator cell line, the cells are incubated and passaged for 14 to 28 days, and then end-points, such as cytopathic effect or hemagglutination, are measured (Berting, 2010). The second is a PCR-based assay, which measures in real-time the presence of nucleic acids associated with adventitious or endogenous agents. For example, see Zhan et al., *Detection of Minute Virus of Mice Using Real Time Quantitative PCR in Assessment of Virus Clearance during the Purification of Mammalian Cell Substrate Derived Biotherapeutics*, 30(4) Biologicals 259-270 (2002).

Mouse minute virus (a.k.a. MMV, minute virus of mouse, or MVM) poses a special problem for biotherapeutic manufacturing. Both the U.S. Food and Drug Administration (FDA) and the European Medicines require testing specifically for MVM. This virus is a member of the parvoviridae family (parvovirus) and common in mice. It is excreted in urine and feces, is robust, and is persistent in the environment. It easily can be introduced into biotherapeutic manufacturing processes. See Moody et al., *Mouse Minute Virus (MAMV) Contamination—A Case Study: Detection, Root*

15

Cause Determination, and Corrective Actions, 65(6) PDA Journal of Pharmaceutical Science and Technology 580-288 (2011). Other rodent parvoviruses may negatively impact cell culture-based biopharmaceutical production. In addition to the prototype MVM strain, those rodent parvoviruses include inter alia the MVM immunosuppressive strain and cutter strain, mouse parvovirus 1a (MPV-1a), MPV-1b, MPV-1c, hamster parvovirus, toolan's parvovirus (parvovirus H-1), kilham rat virus, rat parvovirus 1a, rat minute virus, and the Umass strain of rat virus L. See S. F. Cotmore & P. Tattersal, *The Autonomously Replicating Parvoviruses of Vertebrates*, 33 *Advances in Virus Research* 91-174 (1987), and Jacoby et al., *Rodent Parvovirus Infections*, 46(4) *Lab Anim Sci.* 370-80 (1996).

These parvoviruses share a conserved nucleic acid sequence called NS-1 (NS1), which encodes a large non-structural protein involved in the amplification of the viral genome. The conserved NS1 nucleotide sequence from MVM is depicted in SEQ ID NO:9. Nucleotides 875-956 of that sequence are at least 97% conserved across a wide array of rodent parvovirus NS1 sequences, and therefore serve as good target sequences for PCR-based adventitious and endogenous agent testing. PCR-based testing for rodent parvoviruses (and other adventitious and endogenous agents) can be performed on raw materials, pre-harvested culture media, at various points along the bulk process purification of the biotherapeutic molecule, and at the formulation and packaging stages. The detection of a contaminant may require remediation steps such as disposal of contaminated material, replacement of raw material, and decontamination of the facility.

In addition to testing for adventitious agents, endogenous agents, and other biological contaminants, which enables corrective and preventative actions (CAPAs) during manufacturing, special manufacturing and bulk process steps (i.e., unit operations) may be employed to eliminate, reduce, or inactivate viral contaminants. Chemical inactivation, virus retentive filtration, and chromatography have been shown to be effective in reducing herpesvirus, retrovirus, and parvovirus in harvested or partially purified cell culture fluid. The chemical inactivation step most frequently used is a low pH treatment, which some of ordinary skill in the art believe is due to the denaturation of virus envelope proteins. Protein A, hydroxyapatite, cation exchange and anion exchange chromatography steps have all been shown to remove virus to some extent. See for example Miesegaes et al., *Analysis of Viral Clearance Unit Operations for Monoclonal Antibodies*, 106(2) *Biotechnology and Bioengineering* 238-246 (2010), and Liu et al., *Recovery and Purification Process Development for Monoclonal Antibody production*, 2(5) *mAbs* 480-499 (2010).

Q-PCR Testing: Positive and Negative Controls

Tests for biological contaminants should be properly controlled to ensure accuracy, reliability, and believability. As used herein, a "negative control" incorporates most or all experimental reagents and conditions, but without the test sample. The test sample may be replaced with a buffer or sham culture media known not to contain the biological contaminant-of-interest. Further as used herein, the "negative control" should generate a negative result for the biological contaminant. If the negative control generates a positive biological contaminant result, then the skilled technician or scientist can assume that a positive result from the parallel test sample might not accurately reflect whether the test sample contains the biological contaminant.

As used herein, one or more "positive controls" are employed assess whether the experimental conditions are

16

adequately operable to detect the biological contaminant. Positive controls can be used at any step along the experimental process to ensure that each step is operating, and to determine at which step the process breaks down.

In some embodiments, a positive control is employed at the nucleic acid extraction step to assess whether the intended extraction of any biological contaminant nucleic acid was efficient enough to detect the biological contaminant. Such a positive control is called a "nucleic acid extraction control" or "NEC". In some cases, the NEC is selected to mimic the target biological contaminant in terms of protein-nucleic acid structure. If the NEC is extracted in a manner sufficient to be detected, then the technician may assume that the target biological nucleic acid was also extracted in a manner sufficient to be detected. In one embodiment, the NEC is a single stranded DNA phage, not altogether unlike parvoviruses.

In a specific embodiment, the NEC is an M13 bacteriophage, which is composed of a circular single stranded DNA of about 6407 nucleotides. In a more specific embodiment, the NEC is the M13K07 strain, which is a generally available molecular biology reagent used for cloning and other laboratory purposes. See van Wezenbeek et al., *Nucleotide Sequence of the Filamentous Bacteriophage M13 DNA Genome: Comparison with Phage fd*, 11(1-2) *Gene* 129-148 (1980). A nucleotide sequence of the M13K07 phage is depicted in SEQ ID NO:8. While the M13 bacteriophage, or the M13K07 bacteriophage strain can be used as the NEC in specific embodiments, the invention is in no way limited to using that particular reagent as an NEC. One of ordinary skill in the art may substitute another reagent as an NEC in the practice of this invention without departing from the scope of the invention (e.g., MS2 phage for Reverse Transcriptase-PCR assays; see Kothapalli et al., *Problems associated with product enhancement reverse transcriptase assay using bacteriophage MS2 RNA as a template*, 109(2) *J. Virol. Methods* 203-207 (2003)).

In some embodiments of the invention, a positive control is employed at the PCR amplification step to assess whether the PCR reagents (including primers) and PCR steps are/were sufficient to detect the biological contaminant template nucleic acids. Such a positive control is called a "plasmid amplification control" or "PAC". In some cases, the PAC is selected or designed to match the target biological contaminant nucleic acid sequence, and exactly match the test sample forward and reverse oligonucleotide primers. In a specific embodiment, the PAC additionally contains a "unique sequence" that is not found in the target biological contaminant nucleic acid. In a more specific embodiment, the unique sequence is not found in nature. A "unique" nucleic acid polymer is designed to specifically anneal to this unique sequence and incorporate into amplicon copies of the unique sequence during PCR. The "unique" nucleic acid polymer is designed to not recognize or anneal to any naturally occurring parvovirus under hybridization conditions employed in the operation of the subject assay. In one embodiment, the "unique" nucleic acid polymer comprises from 17 to 20 nucleotides, wherein no more than seven (7) to 10 internal consecutive nucleotides and no more than six (6) consecutive three-prime nucleotides are identical to any parvovirus sequence. In one embodiment, the "unique" nucleotide polymer comprises from 17 to 20 nucleotides, wherein no more than 7 to 10 consecutive nucleotides and no more than six (6) consecutive three-prime nucleotides are identical to any parvovirus sequence set forth in SEQ ID NOs: 9 and 12-37.

17

This unique sequence enables the skilled technician to distinguish between a bona fide target biological contaminant sequence, and cross-contamination of the test Q-PCR reaction with the PAC plasmid.

In one embodiment, the PAC (as a plasmid, a.k.a. PAC plasmid) is included in an identical but separate Q-PCR reaction to control for PCR amplification. The PAC reaction use the exact same oligonucleotide primers and probes as is used in the test sample Q-PCR reaction to accurately reflect amplification of an actual target biological contaminant nucleic acid. The test sample Q-PCR reaction and the separate PAC plasmid Q-PCR reaction include a target detection probe and a unique sequence probe. The PAC reaction, if properly functioning, is expected to generate a positive target signal and a positive unique sequence signal. Obtaining a positive target signal and a positive unique sequence signal in the PAC plasmid-containing reaction indicates that the test sample Q-PCR reagents and conditions are adequately working to produce a positive target signal in the test sample whenever a target biological contaminant nucleic acid is present.

Therefore, the PAC acts as a control for proper PCR amplification. If a negative target signal is obtained in the test sample, but the PAC shows a positive target signal, then the skilled technician can assume that no detectable target contaminant DNA is present in the test sample. Conversely, if a positive target sequence signal and a positive unique sequence signal are obtained in the test sample, then the technician can assume that the PAC plasmid cross-contaminated the test sample, and therefore the positive target sequence signal may be a false positive.

DETAILED DESCRIPTION OF SEVERAL EMBODIMENTS

In some embodiments, an improved positive control system (i.e., compositions and methods) is provided for detecting biological contaminants by the use of polymerase chain reaction, more specifically, Q-PCR.

In one aspect, the invention provides a unique sequence probe (USP) to detect the positive amplification control plasmid (PAC plasmid). The USP contains an artificial nucleotide sequence capable of hybridizing to the unique artificial plasmid-specific sequence (a.k.a. unique sequence, or PACP unique sequence, or UAPS), a fluorophore, and a quencher. In a specific embodiment, the artificial nucleotide sequence capable of hybridizing to the UAPS comprises the

18

nucleic acid sequence of SEQ ID NO:3 (5'-TGTC-GATGGCGAATGGCTA-3'), which is the anti-sense to the UAPS sense strand sequence (e.g., i.e., SEQ ID NO:10-5'-TAGCCATTCGCC ATCGACA-3').

As described above, Q-PCR detection probes contain a fluorophore and a quencher. Quenching may be by contact quenching or by FRET, depending on the fluorophore/quencher pair. Here, a fluorophore and a quencher are covalently attached the USP oligonucleotide. In one embodiment, the fluorophore has an excitation wavelength of 495 nm to 680 nm and an emission wavelength of 515 nm to 710 nm. In some embodiments, the fluorophore has an excitation wavelength of 495 nm, 538 nm, or 646 nm and an emission wavelength of 520 nm, 554 nm, or 669 nm, respectively. In some embodiments, the quencher is a dye with an absorbance peak of 430 nm to 672 nm. Examples of useful quenchers include DDQ®-I, Dabcyl, Eclipse®, Iowa Black FQ®, BHQ®-1, QSY®-7, BHQ®-2, DDQ®-II, Iowa Black RQ®, QSY®-21, and DHQ®-3. In a specific embodiment, the fluorophore is fluorescein amidite (FAM; described in U.S. Pat. No. 5,583,236 (issued Dec. 10, 1996), which has an absorbance maximum of about 495 nm and an emission maximum of about 520 nm, and the quencher is Black Hole Quencher®-1 (BHQ®-1, Biosearch Technologies, Inc., Petaluma, CA), which absorbs at 480 nm to 580 nm. BHQ®-1 quenches via FRET and contact quenching. Generally, but not always, the quencher is attached via an ether bond to the 3-prime hydroxyl group of the oligonucleotide, and the fluorophore is attached via an ester bond to the 5-prime phosphate group of the oligonucleotide.

In another aspect, the invention provides a mixture of Q-PCR reagents comprising multiple oligonucleotides and probes, useful for the detection of a biological contaminant in cell cultures, raw materials, partially purified and purified biological molecules, and the like. In one embodiment, the biological contaminant is a DNA virus, more specifically a parvovirus, and more specifically still, a rodent parvovirus, such as MVM. The parvovirus contains an NS1 gene having a nucleic acid sequence that is at least 88% identical to any sequence listed in Table 1. In another embodiment, the parvovirus contains an NS1 gene having a nucleic acid sequence that is at least 97% identical to the sequence set forth in SEQ ID NO:9 (i.e., Mouse Minute Virus (MVM) NS1 gene). In yet another embodiment, the parvovirus contains an NS1 gene comprising the consensus sequence of SEQ ID NO:37.

TABLE 1

Parvoviridae NS1 Sequences					
Gene Name	SEQ ID NO:	Identity to SEQ ID NO: 9	Gene Name	SEQ ID NO:	Identity to SEQ ID NO: 9
MVM lymphotropic variant	12	97%	MVM, strain M	25	95%
Mouse parvovirus 4b	13	96%	Parvovirus LulII	26	90%
Mouse parvovirus 4a	14	96%	Rat parvovirus UT	27	89%
Mouse parvovirus 1b	15	96%	Kilham rat virus	28	89%
MVM immune-suppressive variant	16	96%	Rat minute virus 1c	29	89%
Mouse parvovirus 1	17	96%	Rat minute virus 1b	30	89%
Mouse parvovirus	18	96%	Rat minute virus 1a	31	89%

TABLE 1-continued

Parvoviridae NS1 Sequences					
Gene Name	SEQ ID NO:	Identity to SEQ ID NO: 9	Gene Name	SEQ ID NO:	Identity to SEQ ID NO: 9
Mouse parvovirus UT	19	96%	H-1 parvovirus	32	89%
Mouse parvovirus 1e	20	96%	Rat minute virus isolate NTU1	33	89%
Mouse parvovirus 1c	21	95%	Parvovirus h-1	34	89%
Hamster parvovirus	22	95%	Rat minute virus isolate NTU2	35	88%
Mouse parvovirus 3	23	95%	Rat minute virus 2a	36	88%
Mouse minute virus	24	95%	Consensus NS1	37	100%

In a specific embodiment, the mixture contains (1) a rodent parvovirus specific forward oligonucleotide primer; (2) a rodent parvovirus specific oligonucleotide detection probe; (3) an artificial oligonucleotide detection probe, such as the USP; and (4) a rodent parvovirus specific reverse oligonucleotide primer. Such a mixture may be used in both the test sample and the positive control sample. In a more specific embodiment, the oligonucleotide primers and parvovirus specific oligonucleotide detection probe hybridize to an NS1 sequence, such as for example the NS1 sequence set forth in SEQ ID NO:9. In a more specific embodiment, the (1) forward primer comprises the sequence of SEQ ID NO:1; (2) the parvovirus specific oligonucleotide detection probe comprises the sequence of SEQ ID NO:2; (3) the artificial probe is a USP and comprises the sequence of SEQ ID NO:3; and (4) the reverse primer comprises the sequence of SEQ ID NO:4.

As described above, the USP contains a fluorophore and a quencher, so that amplicons containing that sequence, i.e., PAC DNA, can be detected in real time as Q-PCR proceeds. In a similar manner, the parvovirus specific detection probe contains an oligonucleotide with a fluorophore and quencher covalently attached. In practice, the fluorophore emission wavelength of the parvovirus detection probe should be different than the emission wavelength of the USP to differentiate between a true parvovirus positive signal and a false positive due to contamination of the test sample with a PAC plasmid or other PACP (e.g., a PACP contaminating amplicon). Thus, in those embodiments in which the USP fluorophore is FAM, the fluorophore attached to the parvovirus detection probe oligonucleotide should have an emission wavelength other than about 520 nm. In a specific embodiment, the fluorophore attached to the rodent parvovirus specific oligonucleotide detection probe is VIC® Dye (Life Technologies, Inc., Carlsbad, CA), which has an absorbance maximum at 538 nm and an emission maximum of about 554 nm. Here, the quencher can be a FRET quencher or a contact quencher. In one embodiment the quencher is a minor groove binding non-fluorescent quencher (MGBNFQ) (see Sylvain et al., *Rapid Screening for HLA-B27 by a TaqMan-PCR Assay Using Sequence-Specific Primers and a Minor Groove Binder Probe, a Novel Type of TaqMan™ Probe*, 287(1-2) Journal of Immunological Methods 179-186 (2004)).

The mixture of primers and probes described above is used to test and distinguish a positive amplification control construct from a bonafide parvovirus contaminant. In some embodiments, the mixture of primers also includes a set of primers and a probe to detect a nucleic acid extraction control (NEC). In a particular embodiment, the NEC is an

M13 bacteriophage, for example the M13K07 strain (SEQ ID NO:8). Thus, in some embodiments, in addition to the parvovirus primers and probe and the USP probe, the mixture of primers and probes contains (5) an M13 specific forward oligonucleotide primer; (6) an M13 specific oligonucleotide detection probe; and (7) an M13 specific reverse oligonucleotide primer.

In some embodiments, the M13 primers and probe hybridize to the sequence of SEQ ID NO:8. In a specific embodiment, the M13 specific forward oligonucleotide primer comprises the nucleic acid sequence of SEQ ID NO:5; the M13 specific oligonucleotide detection probe comprises the nucleic acid sequence of SEQ ID NO:6; and the M13 specific reverse oligonucleotide primer comprises the nucleic acid sequence of SEQ ID NO:7. As in the case of the non-overlapping fluorophore emission spectra for the parvovirus probe and the USP, the NEC probe contains a fluorophore that emits light at a non-overlapping spectrum. In a specific embodiment, the fluorophore attached to the M13 specific oligonucleotide detection probe is Cy5, a cyanine dye having an absorbance maximum of about 650 nm and an emission maximum of about 670 nm (see Southwick et al., *Cyanine Dye Labeling Reagents: Carboxymethylindocyanine Succinimidyl Esters*, 11 Cytometry, 418-430 (1990)). Here, the quencher may operate via FRET or contact quenching. In one embodiment, the quencher is Black Hole Quencher®-2 (BHQ®-2, Biosearch Technologies, Inc., Petaluma, CA), which absorbs at a maximum of about 579 nm, and quenches in the range of about 550 nm to about 650 nm.

In another aspect, the invention provides a method of detecting a biological contaminant in a biological molecule production process. The biological contaminant more specifically comprises a parvovirus, more specifically a rodent parvovirus, and most specifically those parvoviruses sharing at least 97% identity to the MVM NS1 gene. In some embodiments, the NS1 gene comprises the sequence of SEQ ID NO:9. In one embodiment, the biological molecule production process is a mammalian cell culture process for manufacturing an antibody, a trap molecule, or other therapeutic antibody. A test sample is taken from the cell culture (or bulk ingredients) and extracted for nucleic acids. In some cases, the test sample is spiked with an NEC, such as M13 (e.g., SEQ ID NO:8), to serve as a control for proper extraction of nucleic acids prior to Q-PCR. The method comprises the steps of (1) combining (a) the nucleic acid sample extracted from the test sample, (b) oligonucleotide primers and probes (as described above), and (c) a DNA polymerase, preferably a thermostable DNA polymerase with 5' exonuclease activity, such as Taq polymerase; (2)

subjecting that combination to polymerase chain reaction (PCR); and (3) monitoring the production of various amplicons via fluorescence emission amplitude.

Formation of specific amplicons is correlated to the presence and amount of various template nucleic acids in the test sample. The specific amplicons include (1) target amplification polynucleotides (TAPs), such as for example rodent parvovirus sequences (e.g., those biological contaminants containing NS1 sequences), (2) nucleic acid extraction control (NEC), such as for example an M13 (e.g., M13K07) polynucleotides (NECPs), and (3) plasmid amplification control polynucleotides (PACPs), such as the unique artificial plasmid-specific sequence (UAPS). If TAPs and NECPs are produced, and PACPs are not produced, then it can be concluded that the test sample contains a biological contaminant and does not contain a cross-contaminating positive amplification control plasmid. However, if both TAPs and PACPs (i.e., UAPs) are produced in the test sample Q-PCR reaction, then it may be concluded that the test sample was cross-contaminated with the PAC plasmid and the TAP result may be a false positive.

In a specific embodiment—in which the primers and probes comprise (1) a rodent parvovirus specific forward oligonucleotide primer comprising the sequence of SEQ ID NO:1, (2) a rodent parvovirus specific oligonucleotide detection probe comprising the sequence of SEQ ID NO:2, a VIC fluorophore, and an MGBNFQ quencher, (3) an artificial oligonucleotide detection probe, such as the USP comprising the sequence of SEQ ID NO:3 and labeled with 6-FAM and BHQ®-1, (4) a rodent parvovirus specific reverse oligonucleotide primer comprising the sequence of SEQ ID NO:4, (5) an M13 specific forward oligonucleotide primer comprising the sequence of SEQ ID NO:5, (6) an M13 specific oligonucleotide detection probe comprising the sequence of SEQ ID NO:6 and labeled with Cy5 and BHQ®-2, and (7) an M13 specific reverse oligonucleotide primer comprising the sequence of SEQ ID NO:7—the production of TAP is monitored at about 533 nm to about 580 nm, the production of PACP is monitored at about 465 nm to about 510 nm, and the production of NECP is monitored at about 618 nm to about 660 nm.

In one embodiment, the test sample is taken from a production CHO cell culture, e.g., a production culture of EESYR® cells transformed with nucleic acids encoding a protein-of-interest, 96 to 72 hours before the time of harvesting the culture. If a TAP and an NECP is detected in the test sample, but no PACP is detected in the test sample, then a confirmation test may be performed on a second test sample obtained from the same production cell culture 48 hours before the time of harvest. If a TAP and an NECP again are detected in the second test sample, but no PACP is detected in the test sample the cell culture is deemed to be contaminated and may not be processed further. Alternatively, a second test may not be performed, yet the cell culture may be deemed to be contaminated and the culture not processed further.

In one embodiment, nucleic acid is extracted from the test sample taken from the production cell culture. Here, one milliliter of the test sample is subjected to cell lysis, proteolysis, and heat denaturation, followed by combining the sample with a nucleic acid extraction control (NEC) sample and then extracting the nucleic acids from the sample. In one embodiment, the nucleic acids are extracted from the test sample (or NEC-spiked test sample) using an automated nucleic acid extraction system, such as a QIAAsymphony® instrument (Qiagen, Inc., Valencia, CA) (see Lee et al., *Comparative evaluation of the QIAGEN QIAAsymphony® SP*

system and bioMérieux NucliSens easyMAG automated extraction platforms in a clinical virology laboratory, 52(4) J. Clin. Virol. 339-43 (2011)).

In one embodiment, the enzyme uracil-N-glycosylase (UNG) is added to the Q-PCR reaction mixture prior to performing the PCR to selectively degrade contaminating amplicons (see Taggart et al., *Use of heat labile UNG in an RT-PCR assay for enterovirus detection*, 105(1) J. Virol. Methods. 57-65 (2002)). The reaction mixture is incubated at 50° C. for at least 2 minutes, more specifically in some cases for 2 minutes or 5 minutes.

In a specific embodiment, after the optional UNG treatment the reaction mixture is incubated at 95° C. for 2 minutes, followed by eight cycles of (1) denaturation at 95° C. for 10 seconds, followed by (2) annealing and extension for 30 seconds, such that the first annealing temperature is 70° C., then the annealing temperature is decreased by 1° C. at each cycle, such that the eighth annealing is 62° C. After the initial eight cycles, 40 cycles of DNA amplification is performed comprising the steps of (1) denaturation at 95° C. for 10 seconds, followed by (2) annealing and extension at 62° C. for 30 seconds. In a particular embodiment, the temperature change rate from the denaturation temperature to the annealing temperature is about 4.4° C. per second, and from the annealing temperature to the denaturation temperature is about 2.2° C. per second.

In some embodiments, the method of detecting a biological contaminant in a production cell culture medium or product thereof includes performing an external positive amplification control (PAC) assay run separately from the test sample assay, and performing an external negative control assay run separately from the test sample assay and PAC assay. If either the negative control or the positive control fails, then the result obtained from the test sample assay is rejected.

In one embodiment, the external positive control comprises the steps of (1) combining inter alia and without a test sample (a) a positive amplification control (PAC) plasmid, which in a specific embodiment comprises the sequence of SEQ ID NO:11, (b) a rodent parvovirus specific forward oligonucleotide primer comprising the sequence of SEQ ID NO:1, (c) a rodent parvovirus specific oligonucleotide detection probe comprising the sequence of SEQ ID NO:2 labeled with VIC and MGBNFQ, (d) an artificial oligonucleotide detection probe, such as the USP comprising the sequence of SEQ ID NO:3 labeled with 6-FAM and BHQ®-1, (e) a rodent parvovirus specific reverse oligonucleotide primer comprising the sequence of SEQ ID NO:4, and (f) a DNA polymerase, preferably a thermostable DNA polymerase with 5' exonuclease activity, such as Taq polymerase; (2) subjecting the positive control mixture to positive control polymerase chain reaction (PCR); and (3) monitoring the production of (a) target amplification polynucleotides (TAPs), (b) nucleic acid extraction control amplification polynucleotides (NECPs), and (c) plasmid amplification control polynucleotides (PACPs) during the PCR. The production of TAP is monitored at about 533 nm to about 580 nm, the production of PACPs is monitored at about 465 nm to about 510 nm, and the production of NECP is monitored at about 618 nm to about 660 nm.

The positive amplification control PCR reaction is run identically to the test sample PCR reaction (as described above). If TAPs and PACPs are produced in the positive control reaction, then it can be concluded that the PCR amplification procedure is operating properly. If TAPs are not produced in the positive amplification control reaction, then any negative TAP in the test sample may be discounted

23

as a failed PCR reaction. In one embodiment, the NEC (i.e., e.g., M13K07) is included in the positive amplification control. A properly functioning control should also reveal a positive NECP signal (see Table 1).

In one embodiment, the external negative control comprises the steps of (1) combining inter alia and without a test sample and without a PAC plasmid (a) a blank, which can be a buffer that mimics the test sample buffer system, or simply water, (b) a rodent parvovirus specific forward oligonucleotide primer comprising the sequence of SEQ ID NO:1, (c) a rodent parvovirus specific oligonucleotide detection probe comprising the sequence of SEQ ID NO:2 labeled with VIC and MGBNFQ, (d) an artificial oligonucleotide detection probe, such as the USP comprising the sequence of SEQ ID NO:3 labeled with 6-FAM and BHQ®-1, (e) a rodent parvovirus specific reverse oligonucleotide primer comprising the sequence of SEQ ID NO:4, and (f) a DNA polymerase, preferably a thermostable DNA polymerase with 5' exonuclease activity, such as Taq polymerase; (2) subjecting the positive control mixture to positive control polymerase chain reaction (PCR); and (3) monitoring the production of (a) target amplification polynucleotides (TAP), (b) nucleic acid extraction control amplification polynucleotides (NECP), and (c) plasmid amplification control polynucleotides (PACP) during the PCR. The production of TAP is monitored at about 533 nm to about 580 nm, and the production of PACP is monitored at about 465 nm to about 510 nm.

The negative control PCR reaction is run identically to the test sample PCR reaction (as described above). If TAPs and PACPs are produced in the negative control reaction, then it can be concluded that the PCR reagents are contaminated with the PAC plasmid. If TAPs, but no PACPs are produced in the negative control reaction, then it can be concluded that the PCR reagents are contaminated with a parvovirus. In both cases, the test sample results are discarded. However, if both TAP and PACP production is negative in the negative control reaction, TAP and PACP production is positive in the positive amplification control, and TAP (and optionally NECP) production is positive, and PACP production is negative in the test sample reaction, then the technician may conclude that the test sample is contaminated (see Table 2 and Table 3).

In other aspects, the invention provides a positive amplification control plasmid (PAC plasmid), and a mixture of positive control reagents including the positive control plasmid. In one embodiment, the PAC plasmid comprises (1) a parvovirus nucleic acid sequence, (2) an M13K07 nucleic acid sequence, and (3) an artificial nucleic acid sequence unique to the plasmid (a.k.a. UAPS or "unique" sequence). In a specific embodiment, the parvovirus nucleic acid sequence comprises the sequences of SEQ ID NO:1, SEQ ID NO:2, and SEQ ID NO:4; the M13K07 nucleic acid sequence comprises the sequences of SEQ ID NO:5, SEQ ID NO:6, and SEQ ID NO:7, and the unique sequence comprises the antisense sequence of SEQ ID NO:3. In a more specific embodiment, the nucleotide sequence of the PAC plasmid consists of the sequence set forth in SEQ ID NO:11.

In some embodiments, the mixture of positive control reagents includes inter alia the PAC plasmid described above, a rodent parvovirus specific forward oligonucleotide primer, a rodent parvovirus specific oligonucleotide detection probe, an artificial oligonucleotide detection probe (i.e., the USP), a rodent parvovirus specific reverse oligonucleotide primer, an M13 specific forward oligonucleotide primer, an M13 specific oligonucleotide detection probe,

24

and an M13 specific reverse oligonucleotide primer, and a buffer. The mixture optionally contains dNTPs and Taq polymerase.

TABLE 2

Assay Controls and Test Session Status			
Assay Controls	Parvovirus Signal (TAP)	UAP Signal PACP)	M13K07 Signal (NECP)
NEC	0	0	+
PAC	+	+	+
Negative control (buffer or water)	0	0	0
Test session status	☐ VALID		☐ INVALID

TABLE 3

Test Session Status				
Sample Description	Parvovirus	UAP Signal	M13K07	Is test sample suitable?
Negative test sample	0	0	+	Yes
False negative test sample	0	0	0	No
Positive test sample	+	0	+	Yes
False positive test sample	+	+	+	Yes

In a specific embodiment, the rodent parvovirus specific forward oligonucleotide primer comprises the nucleic acid sequence of SEQ ID NO:1, the rodent parvovirus specific oligonucleotide detection probe comprises a VIC fluorophore, a minor groove binding quencher (MGBNFQ), and the nucleic acid sequence of SEQ ID NO:2, the USP comprises a VIC fluorophore, a non-fluorescent quencher BHQ, and the nucleic acid sequence of SEQ ID NO:3, the rodent parvovirus specific reverse oligonucleotide primer comprises the nucleic acid sequence of SEQ ID NO:4, the M13 specific forward oligonucleotide primer comprises the nucleic acid sequence of SEQ ID NO:5, the M13 specific oligonucleotide detection probe comprises a Cy5 fluorophore, a BHQ-2 quencher, and the nucleic acid sequence of SEQ ID NO:6, and the M13 specific reverse oligonucleotide primer comprises the nucleic acid sequence of SEQ ID NO:7.

Example 1: Oligonucleotides and Nucleic Acid Reagents

Rodent parvovirus, M13K07, and artificial unique oligonucleotides (Oligos) were obtained from various vendors at various scales and in various formats, which are described in Table 4. All oligos were assigned a three-year expiration date upon receipt from the vendor.

The oligonucleotides were reconstituted in water to a concentration of 100 μ M to produce master stocks. The master stocks were further diluted to produce 10 $\times$ stock solutions prior to setting up the (PCR reactions. Table 5 depicts the 10 $\times$ and 1 $\times$ concentrations of the qPCR oligonucleotides (primers and probes).

TABLE 4

Parvovirus, M13, and unique artificial sequence primers and probes			
Type of Oligo	Name	Sequence (5'-3')	Vendor/ Scale/Purification/Format
Forward Oligonucleotide Primer	MVM- FWD	5'/TGC ATA AAA GAG TAA CCT CAC CAG/3' (SEQ ID NO: 1)	IDT/1 μ M/HPLC/lyophilized or Lab Ready Life Technologies/80K pMol/ HPLC/dry or liquid
Sequence Detection Probe	MVM- MGB1	5'/VIC/ ACT GGA TGA TGA TGC AGC /MGBNFQ/3' (SEQ ID NO: 2)	Life Technologies/ 50K/HPLC/liquid; 100 pmol/ μ L
Sequence Detection Probe	parvovirus AntiSense Flag	5'/6-FAM/TGT CGA TGG CGA ATG GCT A /BHQ [®] - 1/3' (SEQ ID NO: 3)	IDT/1 μ M/HPLC/lyophilized or Lab Ready
Reverse Oligonucleotide Primer	MVM- REV	5'/CCA CCT GGT TGA GCC ATC/3' (SEQ ID NO: 4)	IDT/1 μ M/HPLC/lyophilized or Lab Ready Life Technologies/80K pMol/ HPLC/dry or liquid
Forward Oligonucleotide Primer	M13- FWD	5'/AAG CCT CAG CGA CCG AAT AT/3' (SEQ ID NO: 5)	IDT/1 μ M/HPLC/lyophilized or Lab Ready
Sequence Detection Probe	M13- probe	/5'/CY5/ TAT GCG TGG GCG ATG GTT GTT GTC A /BHQ2/3' (SEQ ID NO: 6)	IDT/1 μ M/HPLC/lyophilized or Lab Ready
Reverse Oligonucleotide Primer	M13-REV	5'/TCA GCT TGC TTT CGA GGT GAA T/3' (SEQ ID NO: 7)	IDT/1 μ M/HPLC/lyophilized or Lab Ready

TABLE 5

Oligonucleotide Concentrations			
Oligo name	1X oligo concentration per PCR reaction	10X concentration	Volume of Oligo Master Stock for 1 mL 10X concentration
MVM-fwd	0.15 μ M	1.5 μ M	15 μ L
MVM-MGB1	0.15 μ M	1.5 μ M	15 μ L
parvovirus AntiSense Flag	0.15 μ M	1.5 μ M	15 μ L
MVM-rev	0.15 μ M	1.5 μ M	15 μ L
M13-Fwd	0.1 μ M	1.0 μ M	10 μ L
M13-Probe	0.1 μ M	1.0 μ M	10 μ L
M13-Rev	0.1 μ M	1.0 μ M	10 μ L

Example 2: Positive Amplification Control Plasmid

The parvo-M13 positive amplification control (PAC) plasmid was prepared in pUC57-Kan plasmid (GeneWiz, Inc., South Plainfield, NJ). This PAC plasmid consists of the sequence set forth in SEQ ID NO:11. The parvovirus-M13 PAC plasmid contains 2.1×10^8 copies/ng calculated using the following formula: $[\text{number} = (\text{amount} * \text{number}/\text{mole}) / (\text{bp} * \text{ng}/\text{g} * \text{g}/\text{mole} \text{ of } \text{bp})]$; where amount=ng, number/mole= 6.022×10^{23} , bp=4372, ng/g= 1×10^9 , and g/mole of bp=650. The concentration of the plasmid was calculated and expressed as ng/ μ L and diluted serially to a 10 $\times$ concentration of 10^2 copies/ μ L. From the 10^2 copies/ μ L dilution, 50 μ L aliquots were prepared and stored in 2 mL sterile screw-cap tubes. All aliquots were stored at <minus 60° C.

Example 3: Preparation of M13K07 Phage

Ten-fold serial dilutions of M13K07 phage were performed using MEM as a diluent to obtain M13K07 at a titer of 100 pfu/ μ L. 200 μ L aliquots were prepared and stored at <minus 60° C. and assigned a three-year expiration date.

Example 4: Q-PCR Reaction Reagent Preparation

The Rodent Parvovirus Real-Time PCR Detection assay is a fully automated TaqMan[®] PCR process consisting of an automated DNA purification using QIA Symphony[®] (Qia-gen) followed by nucleic acid amplification and real-time PCR product detection on the LightCycler[®] 480 instrument (Roche Diagnostic). Each test article was automatically spiked with the phage M13K07 as an internal control (IC) in order to assess for presence of PCR inhibitors. The rodent parvovirus primer oligonucleotides were designed to hybridize within a highly conserved region of the rodent parvovirus genome (NS-1 region) in order to ensure broad range detection. The assay was run in a duplex (i.e. two targets) format and the rodent parvovirus and the M13K07 primers respectively generated PCR products of approximately 110 and 97 bp. The PCR product was detected in real-time through cleavage of two probes labeled with different reporter dyes: VIC fluorophore for rodent parvovirus and Cy5 fluorophore for M13K07 phage. A positive amplification control (PAC) plasmid was used at a concentration of 100 copies/ μ L. This plasmid contains a unique (Flag) sequence (USP) differentiating it from the wild type parvovirus using a specific probe labeled with the fluorophore FAM.

Master Mix was assembled in a 2 mL tube according to Table 6 in amounts sufficient for at least three times the

number of test articles including the controls. The tubes were stored at 2-8° C. The negative amplification control (NAC) vial was prepared by adding 50 µL of water to a 2 mL tube.

TABLE 6

Reaction Mixtures			
	Reagent Stock Concentration	Reagent Concentration for One Reaction	Reagent Stock Concentration for One Reaction
qPCR 2X Master Mix (TaqMan or VeriQuest)	2X	1X	12.5 µL
Parvo-qPCR-Oligo Mix	10X	1X	2.5 µL
Mix Volume per each well			15 µL
Sample volume per well (extracted DNA)			10 µL

Example 5: Test Sample Preparation

Since all samples have the potential to contain or be contaminated with adventitious agent, aseptic practices were adhered to for all sample pre-processing steps. The following steps were performed in a clean biosafety cabinet. Test articles (samples) were obtained from antibody or trap molecule producing EESYR® cell cultures and frozen or used directly.

1000 µL of PBS was aliquoted into appropriate 2 mL-tubes to serve as negative extraction controls (NEC). When the PCR is used as an end-point to a cell culture step, the positive and negative controls of the cell culture were used as positive extraction control and negative extraction control, respectively.

Each test article was thawed at room temperature. Those samples in 60 mL bags were transferred to 50 mL-Falcon tubes and then aliquoted. 1000 µL of each sample was pipetted into 2 mL-tubes for pre-processing. 400 µL of lysis buffer (ChargeSwitch® Lysis Buffer L13, (Invitrogen, Cat #CS 11202 or equivalent, Carlsbad, CA) was added into each tube and vortexed for at least 10 seconds. 20 µL of Proteinase K (>10 mg/mL, >800 units/mL in 40% glycerol (v/v) containing 10 mM Tris-HCl, pH 7.5, with 1 mM calcium acetate; SAFC, Cat #P4850-5ML, Sigma Aldrich, St. Louis, MO) was then added into each sample and vortexed for at least 10 seconds. Samples were then incubated at 65° C. for 30 minutes. After incubation, the samples were vortexed and centrifuged at 17,000×g for 10 minutes.

Concurrently, the positive amplification control (PAC; minimum 50 µL of 10² copies/µL plasmid in 2 mL tube) and M13K07 (10² PFU/mL) Internal Control (IC; minimum 150 µL/12 samples) were thawed.

Example 6: Nucleic Acid Extraction

Internal Control (IC) was required for the programmed extraction protocol on the QIASymphony® Instrument (Qiagen, Valencia, CA). The instrument automatically added 120 µL of reconstituted IC to each sample. For every 12 samples, 1.8 mL of IC (i.e., 1 vial) was required by the instrument. Vials of IC were prepared by adding 1650 µL AVE buffer (RNase-free water containing 0.04% sodium azide) to 150 µL of M13K07 IC.

Each prepared sample was loaded into the Sample Carrier of the instrument within the biosafety cabinet (BSC). The vial(s) of IC were loaded into a separated (dedicated) Sample Carrier within the BSC at a ratio of one vial of IC per 12 samples. All drawers were closed and the instrument was run according to manufacturer's recommended protocol (see QIASymphony DNA Handbook, 09/2010, available at <http://www.algimed.by/download/EN-QIASymphony-DNA-Handbook.pdf>).

The Reagent Prep Cartridge was prepared with the QIASymphony® DSP Virus/Pathogen reagent kit, which contained all reagents required for performing an extraction (see QIASymphony® DSP Virus/Pathogen Kit Instructions for Use (Handbook, April 2013, available at <https://www.qiagen.com/us/resources/download.aspx?id=f8bc0b3c-0aff-46ee-8807-5ed145f9e969&lang=en>). The reagent prep cartridge contained proteinase K and had a shelf-life of about two weeks.

The nucleic acid extraction was performed in a 96-well format reaction plate to facilitate integration with the Q-PCR. After the automated nucleic acid extraction procedure was completed and passed the status check, the reaction plate was cooled and sealed with LightCycler® 480 Sealing Foil (Roche, Branchburg, NJ). The 96-well plate was placed in a plate spinner balanced with a suitable counterweight (e.g., another 96-well plate) and spun for 30 to 60 seconds. The wells were checked for bubbles and the spin was repeated if necessary.

Example 7: Q-PCR

Q-PCR was performed on the LightCycler® 480 Instrument (Roche, Branchburg, NJ) using any one of two programs—TaqMan Triplex, and/or Veriquest Triplex. The TaqMan Triplex protocol used the TaqMan Fast Advance Custom Master Mix without the reference dye (ROX). Three fluorescence channels (FAM, VIC, and CY5) were selected and a UNG step time of 2 minutes. The reaction parameters that were used are outlined in Table 7.

When using VeriQuest® Master Mix, also without ROX, a UNG step time of 5 minutes was used. The reaction parameters that were used are outlined in Table 8.

TABLE 7

Rodent Parvovirus PCR TaqMan Triplex Program Steps						
Step Name	# of Cycles	Analysis mode	Temperature (° C.)	Time	Block ramp rate (° C./Sec)	Step Size
UNG	1	None	50	2 min	4.4	
Pre-Incubation	1	None	95	2 min	4.4	
TD	8	None	95	10 sec	4.4	1° C./step
			70 --> 62	30 sec	2.2	

TABLE 7-continued

Rodent Parvovirus PCR TaqMan Triplex Program Steps						
Step Name	# of Cycles	Analysis mode	Temperature (° C.)	Time	Block ramp rate (° C./Sec)	Step Size
Amplification	40	Quantification	95	10 sec	4.4	
			62	30 sec	2.2	
Cooling	1	None	40	30 sec	2.2	

The crossing point (Cp) fluorescence signal for each of the M13 internal control, rodent parvovirus, and positive amplification control plasmid was determined by one or both of two algorithms. The first algorithm is the Automated Second Derivative method. This method does not require user input and generally resulted in greater consistency, and therefore was considered to be the preferred method. The second algorithm is the Fit Points method. This method allows the user to set the threshold line, in cases of divergent background. The point where the log-linear curve crosses that threshold line becomes the crossing point. The following filter combs were used to monitor fluorescent signals: 533-580 nm (VIC signal) for Rodent parvovirus (subset "Sample-Parvo"); 618-660 nm (Cy5 signal) for the internal control M13K07 (subset "Sample-M13"); and 465-510 nm (FAM signal) for the positive amplification control plasmid (subset "Sample-FAM"). The fluorescence background level was determined using the Fit Point analysis in case of ambiguous fluorescence signal. Acceptable background fluorescence signal was considered to be <1 unit on the amplification plot scale. Any fluorescence signal ≤1 unit was considered within acceptable background, and thus negative.

TABLE 8

Rodent Parvovirus PCR VeriQuest Triplex Program Steps						
Step Name	# of Cycles	Analysis mode	Temperature (° C.)	Time	Block ramp rate (° C./Sec)	Step Size
UNG	1	None	50	5 min	4.4	
Pre-Incubation	1	None	95	2 min	4.4	
TD	8	None	95	10 sec	4.4	1° C./step
			70 --> 62	30 sec	2.2	
Amplification	40	Quantification	95	10 sec	4.4	
			62	30 sec	2.2	
Cooling	1	None	40	30 sec	2.2	

Example 8: Conditions for a Valid Test Session

For the test session to be considered valid, the following conditions must be met. The Negative Amplification Control (NAC, i.e. water) must be negative for fluorescence signal in all three channels. The Negative Extraction Control (NEC, i.e. PBS) or the cell Culture Negative Control flask must be negative for fluorescence signal in the [533-580] channel (i.e. rodent parvovirus probe VIC signal), negative for fluorescence signal in the [465-510] channel (i.e. positive amplification control [PAC] Antisense Flag probe-FAM signal), and positive for fluorescence signal in the [618-660] channel (i.e. M13K07 probe-CY5 signal). The PAC should be positive for fluorescence signals all three channels. The M13K07 Cp value in the NEC was used as a reference for assessing the presence of inhibitory substance in the samples.

If the PCR was used as an end-point for a test article from a cell culture step with a rodent parvovirus positive control, the Positive Virus Control must be positive for fluorescence signal in the [533-580] channel (i.e. rodent parvovirus probe VIC signal), positive for fluorescence signal in the [618-660] channel (i.e., M13K07 probe-CY5 signal), and negative for fluorescence signal in the [465-510] channel (i.e., PAC Antisense Flag probe-FAM signal). The M13K07 Cp value in the cell culture positive control sample was expected to be within ±4 cycles range of the NEC-M13 Cp value.

For all PAC plasmid-containing control reactions, the fluorescent signals must be positive in all three channels.

Example 9: Conditions for an Invalid Test Session

An assay was considered invalid when any one or more of the following conditions was met: (1) a very low amplification curve (<1 unit on the fluorescence scale) for the PAC, (2) a determinant error (machine, software, or human error) was confirmed, (3) the NAC was positive for amplification in any of the three channels, (4) the NEC was positive for amplification in the [533-580] VIC channel, positive for

amplification in the [465-510] FAM channel, or negative for amplification in the [618-660] Cy5 channel, and (5) the PAC was negative for amplification signal in any of the three channels. Investigation and retesting of the test sample was performed whenever the assay was determined to be invalid.

Example 10: Conditions for Negative Sample Result in Valid Test

All of the following conditions must be met for a valid negative parvovirus test result. The sample must be positive for M13K07 DNA amplification signal in the M13 [618-660] channel within an expected Cp value range of NEC-M13 Cp±4 cycles, suggesting the absence of PCR inhibitors. The sample must be negative for parvovirus DNA amplification signal in the parvovirus [533-580] channel. A fluorescence signal below 1 unit on the fluorescence scale,

31

regardless the Cp value was considered within the acceptable fluorescence background level, and was reported as negative for parvovirus DNA amplification. The sample must be negative for PAC amplification signal in the Antisense Flag [465-510] channel. Note that a fluorescence signal below 1 unit on the fluorescence scale, regardless the Cp value (automatically generated by the instrument), was considered within the acceptable fluorescence background level, and was reported as negative for PAC-plasmid DNA amplification.

Example 11: Conditions for “No Sample” Result in a Valid Test Session

A “No Sample” result occurred when any one or more of the following conditions was met. Whenever the sample was negative for M13K07 DNA amplification signal in the Cy5 channel [618-660], negative for parvovirus DNA amplification signal in the parvovirus channel [533-580], and negative for PAC amplification in the antisense Flag FAM channel [465-510], the sample or PCR reagents were investigated per standard operating procedure. This condition suggests the presence of PCR inhibitors or a failure of proper nucleic acid extraction. Samples could be diluted (1:2, 1:5, 1:10) to overcome inhibition. A Cp value for the fluorescence signal in the M13 channel [618-660] that was out of range (NEC M13 Cp $\pm$ 4 cycles), indicated partial inhibition of the PCR or error in phage spiking and warranted a repeat test. Sample dilution (1:2, 1:5, 1:10) could be considered to overcome any inhibition of PCR. Any evidence of determinant error or unexpected very low fluorescence signal (<1 unit on the fluorescence scale) observed in the M13 channel [618-660] that did not allow a conclusive evaluation of the sample suitability was considered to be a “no sample result”, and suggested failure during amplification or DNA extraction. This warranted a repeat test.

Example 12: Condition for Initial Out of Specification (iOOS) Sample Result in a Valid Test Session

A sample was considered to be iOOS when it was (i) positive for parvovirus DNA amplification signal in the parvovirus channel [533-580] (i.e., rodent parvovirus probe VIC signal) with a fluorescence signal above 1 unit on the fluorescence scale (for at least one of the two replicate wells), and (ii) negative for PAC amplification in the antisense Flag FAM channel [465-510], indicating that there was no cross-contamination from the PAC. As a result, the technician must (i) initiate a GLIF (General Laboratory Investigation Form), (ii) notify the department that submitted the sample to QC Virology for testing, (iii) initiated an NOE, (iv) preserve the amplification tube (freeze at -20° C.) for further investigation (e.g. Flag sequence screening or sequencing), and (v) repeat the assay and retest the sample.

Example 13: Repeat and Retest Plan

A repeat test was initiated whenever the assay was invalid or a sample result was considered a “no result.” A retest was performed for confirmation of an iOOS event (i.e. the first time a sample result was positive for DNA amplification signal in the parvovirus channel [533-580]). Retests or repeat tests must be executed using fresh reagent aliquots (i.e., master mix reagents, extraction kit).

If NAC was positive for fluorescence in any channel; the entire test session would be repeated starting from the

32

amplification (PCR) step using the already purified DNA samples with freshly made master mix. If NEC was negative for fluorescence in the M13 channel [618-660], the entire test session would be repeated starting from the DNA extraction step. If NEC was positive for fluorescence in the parvovirus channel [533-580] or the Antisense Flag channel [465-510], the entire test session would be repeated starting from the DNA extraction step. If PAC was negative for fluorescence in any channel, the test session would be repeated from the amplification (PCR step), using the same purified DNA samples with freshly made master mix.

For those samples with a “no sample” result, the test session would be repeated for the impacted sample starting from the DNA extraction step. Samples could be diluted (1:2, 1:5, or 1:10) (in addition to the non-diluted sample) before DNA extraction as part of the investigation and issue resolution process in order to demonstrate the presence of inhibitors.

The retest to confirm an initial positive result (iOOS) was performed using four separate aliquots as follows: two aliquots of test article from the original sampling event (e.g., one day after final feed), and two aliquots of test article from a different sampling event (e.g., two days after final feed, if possible) or different sample bag. If any of the additional testing of the 4 aliquots resulted in a positive signal with no evidence of determinate error (demonstrated via the investigation), the lot was considered as failing the requirement of absence of rodent parvovirus virus genomic material. An infectivity assay was warranted for final disposition for such a positive Q-PCR result. CHO-K1 cells are used as the indicator cell line to determine the infection state of the detected nucleic acid.

Whenever the retest results are negative, a confirmatory test is required at a different sampling event (e.g., three days after final feed) to confirm absence of rodent parvovirus genomic material.

Example 14: Retest Plan for Other Sample Types with iOOS

Other sample types including, e.g., unprocessed bulk material, end-of-production cells, cells at the limit of in vitro age, and soy material, which had an initial positive result (iOOS), were retested as follows using four separate aliquots:

Two aliquots of test article from the original sample container (i.e., e.g., bag) and two aliquots from a different sample container were retested according to standard operating procedures. The CHO-K1 indicator cell was inoculated with one sample aliquot of the test article from the original sample container. The inoculated indicator cell CHO-K1 was harvested after 1-3 days in culture per standard operating procedure to determine the infectious status of the detected nucleic acid. A standard curve for quantification of the re-extracted nucleic acid may be used.

Whenever any of the additional PCR testing of the four aliquots was positive with no evidence of determinate error (demonstrated via the investigation), the infectivity assay determines the final disposition of the article. Upon confirmation of the initial OOS, sequencing of the nucleic acid and transmission electron microscopy (TEM) could be considered in order to identify the microorganism and rule out any laboratory error.

The infectivity assay using CHO-K1 as an indicator cell for the suspected contaminated material is required to determine the infectious state of the nucleic acid detected. Whenever the investigation fails to support the possibility of

33

parvovirus virus contamination and the additional PCR tests of the four aliquots of test article and the CHOK1 culture are all negative, the lot was considered as meeting the requirements for absence of infectious rodent parvovirus virus.

Example 15: Parvovirus Testing During Recombinant Protein Production

The Q-PCR procedure described above was performed as a critical in-process control for the testing of cell culture fluid, i.e., the unprocessed bulk material from the production bioreactor containing a CHO cell derivative containing heterologous antibody heavy chain and light chain constructs. Each heterologous monoclonal antibody (mAb) binds to a different target or epitope. Good manufacturing process (GMP) tests were performed by QC virology scientists at a large-scale bioprocess production facility. Sixteen (16) of those tests are listed in Table 9. In each case, the test sessions were valid, the assay system suitability criteria had been met, and false positive detection was not observed. In test sessions #13 and #14, false negative detection did occur, which suggested the presence of PCR inhibitors or a failure of nucleic acid extraction.

34

TABLE 9

Rodent Parvovirus PCR Tests					
Test #	Type of Sample	Product	Lot	Testing	
1	UPB EA	mAb1	'51	Rodent Parvo PCR	
2	UPB	mAb1	'52	Rodent Parvo PCR	
3	UPB	mAb3	'07	Rodent Parvo PCR	
4	UPB	mAb1	'51	Rodent Parvo PCR	
5	UPB EA	mAb2	'35	Rodent Parvo PCR	
6	UPB EA	mAb4	'05	Rodent Parvo PCR	
7	UPB	mAb2	'36	Rodent Parvo PCR	
8	UPB	mAb1	'53	Rodent Parvo PCR	
9	UPB	mAb1	'53	Rodent Parvo PCR	
10	UPB EA	mAb3	'08	Rodent Parvo PCR	
11	UPB EA	mAb5	'01	Rodent Parvo PCR	
12	UPB EA	mAb1	'54	Rodent Parvo PCR	
13	UPB	mAb6	'01	Rodent Parvo PCR	
14	UPB	mAb4	'05	Rodent Parvo PCR Qualification	
15	UPB	mAb2	'35	Rodent Parvo PCR Qualification	
16	UPB	mAb2	'36	Rodent Parvo PCR Qualification	

UPB = Unprocessed Bulk Material;
EA = Early Alert;
Rodent Parvo PCR = Q-PCR Procedure described above.

SEQUENCE LISTING

```

Sequence total quantity: 37
SEQ ID NO: 1          moltype = DNA  length = 24
FEATURE              Location/Qualifiers
source               1..24
                    mol_type = genomic DNA
                    organism = Rodent protoparvovirus 1

SEQUENCE: 1
tgcataaaag agtaacctca ccag                                24

SEQ ID NO: 2          moltype = DNA  length = 18
FEATURE              Location/Qualifiers
source               1..18
                    mol_type = genomic DNA
                    organism = Rodent protoparvovirus 1

SEQUENCE: 2
actggatgat gatgcagc                                      18

SEQ ID NO: 3          moltype = DNA  length = 19
FEATURE              Location/Qualifiers
source               1..19
                    mol_type = other DNA
                    organism = synthetic construct

SEQUENCE: 3
tgtcgatggc gaatggcta                                    19

SEQ ID NO: 4          moltype = DNA  length = 18
FEATURE              Location/Qualifiers
source               1..18
                    mol_type = genomic DNA
                    organism = Rodent protoparvovirus 1

SEQUENCE: 4
ccacctgggt gagccatc                                     18

SEQ ID NO: 5          moltype = DNA  length = 20
FEATURE              Location/Qualifiers
source               1..20
                    mol_type = genomic DNA
                    organism = Inovirus M13

SEQUENCE: 5
aagcctcagc gaccgaatat                                   20

SEQ ID NO: 6          moltype = DNA  length = 24
FEATURE              Location/Qualifiers
source               1..24
                    mol_type = genomic DNA
                    organism = Inovirus M13

SEQUENCE: 6
tatgcgtggg cgatggttgt tgtc                                24

```

-continued

```

SEQ ID NO: 7          moltype = DNA length = 22
FEATURE              Location/Qualifiers
source                1..22
                     mol_type = genomic DNA
                     organism = Inovirus M13

SEQUENCE: 7
tcagcttgct ttcgaggtga at                                     22

SEQ ID NO: 8          moltype = DNA length = 6407
FEATURE              Location/Qualifiers
source                1..6407
                     mol_type = genomic DNA
                     organism = Inovirus M13

SEQUENCE: 8
aacgtacta ctattagtag aatgatgcc accttttcag ctgcgcccc aaatgaaaat 60
atagctaaac aggttattga ccatttgcca aatgtatcta atgggtcaaac taaatctact 120
cgttcgcaga attgggaatc aactgttaca tgggaatgaaa ctccagaca ccgactactta 180
gttgcataatt taaaacatgt tgagctacag caccagattc agcaattaag ctctaagcca 240
tccgcaaaaa tgacotctta tcaaaaggag caattaaagg tactctctaa tcttgacctg 300
tgggagtttg ctcccggtct ggttcgcttt gaagctcgaa ttaaaacgcg atatttgaag 360
tcttcgggc ttcctcttaa tctttttgat gcaatccgct ttgcttctga ctataatagt 420
cagggtaaaag acctgatttt tgatttatgg tcattctcgt tttctgaact gtttaagca 480
tttgaggggg attcaatgaa tatttatgac gattccgcag tattggacgc tatccagtct 540
aaacatttta ctattacccc ctctggcaaa acttcttttg caaaagcctc tcgctatttt 600
ggtttttatc gtcgtctggt aaacgaggt tatgatagtg ttgctcttac tatgcctcgt 660
aattcctttt ggcgttatgt atctgcatta gttgaatgtg gtattcctaa atctcaactg 720
atgaatcttt ctacctgtaa taatgttgtt ccggtagttc gttttattaa cgtagatttt 780
tcttcccaac gtctgtactg gtataatgag ccagttctta aaatcgcata aggttaattca 840
caatgattaa agttgaaatt aaacctctc aagcccaatt tactactcgt tctgggtgtt 900
ctcgtcaggg caagccttat tcactgaatg agcagcttgg ttacgttgat ttgggtaagt 960
aatatccggg tctgtcaag attactcttg atgaaggcca gccagcctat gcgcctgggc 1020
tgtacaccgt tcactctgct tcttccaaag ttggtcagtt cgggtccctt atgattgacc 1080
gtctgcgcct cgtccggct agtaaacatg gagcaggtcg cggatttcga cacaatttat 1140
caggcgatga tcaaaactc cgttgtactt tgttcgcgc ttggtataat cgtgggggt 1200
caaaagatgag tgttttagtg tatctttcgc cctcttctgt tttaggttgg tgccttcgta 1260
gtggcattac gtattttacc cgtttaatgg aaacttcctc atgaaaaagt ctttagtcct 1320
caaagcctct gttagcgttg ctaccctcgt tccgatgctg tctttcgtcg ctgaggggtga 1380
cgatcccgca aaagcggcct ttaactccct gcaagcctca gcgaccgaat atatcggtta 1440
tcgctggggc atggttgttg tcattgtcgg cgcaactatc ggtatcaagc tgtttaagaa 1500
attcacctcg aaagcaagct gataaaacga tacaattaaa ggctcctttt ggagcctttt 1560
tttttgagat tttcaacat gaaaaaatta ttatcgcaa ttctcttagt tgttctttc 1620
tattctact ccgtgaaacg tgtgaaagt tgtttagcaa aaccccatc agaaaattca 1680
tttactaacg tctggaaga cgacaaaact ttagatcggt acgctaacta tgaggggtgt 1740
ctgtggaatg ctacaggcgt tgtagtttgt actggtgacg aaactcagtg ttacggtaaa 1800
tgggttccta ttgggttgac tatccctgaa aatgaggggtg gtggtctgga ggggtggcgt 1860
tctgaggggtg gcggttctga ggggtggcgt actaaacctc ctgagtagcg tgataacact 1920
attccgggct atacttatat caacctctc gacggcactt atccgcctgg tactgagcaa 1980
aaacccgcta atccctaacc ttctcttgag gagtctcagc ctcttaatac tttcatgttt 2040
cagaataata ggttcgaaa taggcagggg gcattaaactg tttatacggg cactgttact 2100
caaggcactg acccgttaa aacttattac cagtacactc ctgtatcatc aaaagccatg 2160
tatgacgctt actggaacgg taaattcaga gactgcgctt tccattctgg ctttaatgag 2220
gatccattcg tttgtgaata tcaaggccaa tctgtctgacc tgcctcaacc tctgtcaat 2280
gctggcgccg gctctggttg tggttctggt ggcggtctcg aggggtggtg ctctgaggg 2340
ggcgggtctg aggggtggcg cctctgagga ggcggttccg gtggtggctc tggttccgg 2400
gattttgatt atgaaaagat ggcacaaactc aataaggggg ctatgaccca aaatgccgat 2460
gaaaacgcgc tacagctcga cgctaaaggg aaacttgatt ctgtcgctac tgattacggt 2520
gctgctatcg atggtttcat tggtagcgtt tccggccttg ctaatggtaa tgggtctact 2580
ggtagtttg ctggctctaa ttcccaatg gctcaagtcg gtgacggtag taattcacct 2640
ttaatgaata attccgtca atattacct tccctccctc aatcggttga atgtccgct 2700
tttgtcttta gcgctggtaa accatatgaa ttttctattg attgtgacaa aataaactta 2760
ttccgtgggt tctttgcgtt tcttttatat gttgccacct ttatgtatgt attttctacg 2820
tttgtaaca tactgcgtaa taaggagtct taatcatgcc agttcttttg ggtattccgt 2880
tattattcgc tttcctcggt ttctctcgtg taactttgtt cggctatctg cttacttttc 2940
ttaaaaaggg cttcggttaag atagctattg ctatttcatt gtttcttctg cttattattg 3000
ggcttaactc aattctctgt ggttatctct ctgatattag cgctcaatta ccctctgact 3060
ttgttcaggg tggtcagtta attctccgct ctaatgcgct tccctgtttt tatgttatc 3120
tctctgtaaa ggcgtgctatt ttcattttg acgttaaaaa aaaaactggt tcttatttgg 3180
attgggataa ataatatggc tgtttatttt gtaactggca aattaggctc tggaaagacg 3240
ctcgttagcg ttggtaagat tcaggataaa attgtagctg ggtgcaaaa agcaactaat 3300
cttgatttaa ggcctcaaaa cctcccgcaa gtccggagggt tgcgtaaaaa gcctcgcggt 3360
cttagaatac cggataagcc ttctatatct gatttgcttg ctattgggag cggtaatgat 3420
tccacgatg aaaaataaaa cggcttgctt gttctcgatg agtcgggtac ttgggttaat 3480
accggtctct ggaatgataa ggaagacag ccgattattg attgggttct acatgctcgt 3540
aaattaggat gggatattat ttttcttgt caggacttat ctattgttga taaacaggcg 3600
cgttctgcat tagctgaaca tgttgtttat tgcgtcgtc tggacagaat tactttacct 3660
tttgtcggta ctttatattc tcttattact ggctcgaaaa tgcctctgcc taaattacat 3720
gttggcggtg ttgaatatgg cgattctcaa ctggttagcg ttgggtttat 3780
actggtgaaga attgtataa cgcatatgat actaaacagg ctttttctag taattatgat 3840

```

-continued

```

tccggtgttt attcttattt aacgccttat ttatcacacg gtcggtattt caaaccatta 3900
aatttaggtc agaagatgaa attaaactaaa atatatattga aaaagttttc tcgcgttctt 3960
tgtcttgccga ttggatttgc atcagcattht acatatagtt atataaccca acctaaagccg 4020
gagggttaaaa aggtagtcctc tcagacctat gattttgata aattcactat tgactcttct 4080
cagcgctcta atctaagcta tcgctatgtt ttcaaggatt ctaagggaaa attaattaat 4140
agcgacgatt tacagaagca aggttatttca ctcacatata ttgatttatg tactgtttcc 4200
attaaaaaag gtaattcaaa tgaattgtt aaatgtaatt aattttgttt tcttgatgtt 4260
tgtttcatca tcttcttttg ctcaggtaat tgaaatgaat aattgcctc tcgcgattt 4320
tgtaacttgg tattcaaaagc aatcaggcga atccgttatt gtttctcccg atgtaaaagg 4380
tactgttact gtatattcat ttgacgttaa acctgaaaat ctacgcaatt tctttatttc 4440
tgttttactg gctaataatt ttgatattgt tggttcaatt ccttccataa ttcagaagta 4500
taatccaaac aatcaggatt atattgatga attgccatca totgataatc aggaatatga 4560
tgataattcc gctccttctt gtggtttctt tgttccgcaa atgataatg ttactcaaac 4620
ttttaaaatt aataacgttc gggcaaaagg ttttaacga gttgtcgaat tgtttgtaaa 4680
gtctaatact tctaaatcct caaatgtatt atctattgac ggctctaatc tattagtgtt 4740
tagtgacctc aaagatattt tagataacct tctcacttc cttctactg ttgatttggc 4800
aactgaccag atattgttg aggggttgat atttgaggtt cagcaagggt atgctttaga 4860
tttttcattht gctgctggct ctcagcggtg cactgttgca ggcggtgtta atactgaccg 4920
ctcacctctt gttttcttct ctgctgggtg ttggtcgtt atttttaag gcgatgtttt 4980
agggctatca gttcgcgcct taagactaa tagccattca aaaatattgt ctgtgccacg 5040
tattcttacc ctttcagggtc agaagggttc tatctctgtt ggccagaatg tcccttttat 5100
tactggtcgt gtgactgggt aatctgccaa tgtaataaat ccatttcaga cgattgagcg 5160
tcaaaaatga ggtatttcca tgagcggttt tctgttgca atggctggcg gtaatatgtt 5220
tctggatatt accagcaagg ccgatagttt gagttcttct actcaggcaa gtgatgttat 5280
tactaatcaa agaagtattg ctaccaagggt taatttgcgt gatggacaga ctcttttact 5340
cgggtggcctc actgattata aaaacacttc tcaagattct ggcgtaacct tctgtcttaa 5400
aatcccttta atcgccctcc tgtttagctc ccgctctgat tccaacgagg aaagcacgtt 5460
atcgtgctct gtcaaaagca ccatagtacg cgccctgtag cggcgacata agcgcggcg 5520
gtgtgggtgt tacgcgcagc gtgacgcgta cacttgccag cgccctagcg cccgctcctt 5580
tcgctttctt ccttctcttt ctgcgcacgt tcgcgcgctt tcccgctcaa gctctaaatc 5640
gggggctccc ttagggttct cgatttagtg ctttacggca cctcgacccc aaaaaacttg 5700
atttgggtga tgggtcacgt agtgggacct cgccctgata gacgggtttt cgccctttga 5760
cgttggagtc cacgttcttt aatagtggac tcttgttcca aactggaaca acactcaacc 5820
ctatctcggt ctattctttt gatttataag ggattttgcc gatttcggcc tatttggttaa 5880
aaaatgagct gatttaacaa aaattttaa cgaattttaa caaaatatta acgtttacaa 5940
tttaaatatt tgcttataca atcttctgt ttttggggtt tttctgatta tcaaccgggg 6000
tacatatgat tgacatgcta gttttacgat taccgttcat cgattctctt gtttgctcca 6060
gactctcagg caatgacctg atagcctttg tagacctctc aaaaatagct acctctccg 6120
gcatgaattt atcagctaga acggttgaat atcatattga tgggtgattg actgtctccg 6180
gcctttctca ccttttgtaa tctttaccta cacattactc aggcattgca tttaaaatat 6240
atgaggggtc taaaaatttt tacccttgcg ttgaaataaa ggcttctccc gcaaaagtat 6300
tacaggggtc taatgttttt ggtacaaccg atttagcttt atgctctgag gctttattgc 6360
ttaattttgc taattctttg ccttgccgtg atgatttatt ggatgtt 6407

```

```

SEQ ID NO: 9      moltype = DNA length = 2019
FEATURE          Location/Qualifiers
source            1..2019
                  mol_type = genomic DNA
                  organism = Rodent protoparvovirus 1

```

```

SEQUENCE: 9
atggctggaa atgcttactc tgatgaagtt ttgggagcaa ccaactgggt aaaggaaaaa 60
agtaaccagg aagtgttctc atttgttttt aaaaatgaaa atgttcaact gaattgaaaa 120
gatatcggat ggaatagtta caaaaaagag ctgcaggagg acgagctgaa atctttacaa 180
cgaggagcgg aaactacttg ggaccaaaagc gaggacatgg aatgggaaac cacagtggat 240
gaaatgacca aaaagcaagt attcattttt gattcttttg ttaaaaaatg tttatttgaa 300
gtgcttaaca caaagaatat atttctgtgt gatgttaatt ggtttgtgca acatgaatgg 360
ggaaaagacc aaggctggca ctgccatgta ctaattggag gaaaggactt tagtcaagct 420
caagggaagt ggtggagaag gcaactaaat gtttactgga gcagatgggt ggtaacagcc 480
tgtaatgtgc aactaacacc agctgaaaga attaaactaa gagaatagc agaagacaat 540
gagtgggtta ctctacttac ttataagcat aagcaaacca aaaaagacta taccagtggt 600
gttctttttg gaaacatgat tgcttactat tttttaacta aaaagaaaat aagcactagt 660
ccaccaagag acggaggcta ttttcttagc agtgactctg gctggaaaac taacttttta 720
aaagaaggcg agcgccatct agtgagcaaaa ctatacactg atgacatgcg gccagaaacg 780
gttgaaacca cagtaaccac tgcgcaggaa actaagcgcg gcagaattca aactaaaaaa 840
gaagtttcta ttaaaactac acttaaagag ctggtgcata aaagagtaac ctaccagag 900
gactggatga tgatgcagcc agacagttac attgaaatga tggctcaacc aggtggagaa 960
aacctgtcga aaaatacgct agagatttgt acactaactc tagccagaac caaaacagca 1020
tttgacttaa ttttagaaaa agctgaaacc ccaacttttc actgcctgac 1080
acaagaacct gcagaatttt tgcttttcat ggctgggaact atgttaaagt ttgccatgct 1140
atthtctgtg ttttaaacag acaaggaggc aaaaagaaata ctgttttatt tcatggacca 1200
gccagcacag gcaaatctat tattgcacaa gccatagcac aagcagttgg caatgttggt 1260
tgctataatg cagccaatgt aaactttcca tttaatgact gtaccaacaa gaacttgatt 1320
tgggtagaag aagctggtta ctttgacag caagtaaac agtttaaagc catttgctct 1380
ggtcaaaacta ttcgcaattg tcaaaaaagg aaaggcagca aacagattga accaacacca 1440
gtcatcatga ccaacaaatg gaacattaca gtggtcagaa taggctgcga agaaagacca 1500
gaacacactc aaccaatcag agacagaatg cttaacatct atctaacaca taccctgcct 1560
ggtgactttg gtttggttga caaaaatgaa tggcccatga tttgtgcttg gttggtaaa 1620
aatggttacc aatctaccat ggcaagctac tgtgctaaat ggggcaaaag tctgtattgg 1680
tcagaaaact gggcgagacc aaagggtcca actcctataa atttactagg ttcggcacgc 1740

```

-continued

```

tcaccattca cgacaccgaa aagtacgcct ctcagccaga actatgcact aactccactt 1800
gcatcggatc tcgaggacct ggcttttagag ccttgaggca caccaaatac tctgtttgcy 1860
ggcactgcag aaaccacgaa cactggggaa gctggttcca aagcctgcca agatgggtcaa 1920
ctgagcccaa cttggtcaga gatcgaggag gatttgagag cgtgcttcgg tgcggaaccg 1980
ttgaagaaag acttcagcga gccgctgaac ttggactaa 2019

```

```

SEQ ID NO: 10      moltype = DNA length = 19
FEATURE            Location/Qualifiers
misc_feature       1..19
                    note = synthetic
source             1..19
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 10
tagccattcg ccacgcaga 19

```

```

SEQ ID NO: 11      moltype = DNA length = 1707
FEATURE            Location/Qualifiers
misc_feature       1..1707
                    note = synthetic
source             1..1707
                    mol_type = other DNA
                    organism = Inovirus M13

```

```

SEQUENCE: 11
aagcctcagc gacgaatat atcgggttatg cgtggggcat ggtgtgtgtc attgtcggcg 60
caactatcgg tatcaagctg ttttaagaaat tcacctcgaa agcaagctga cagcctacc 120
gcgatgctga atgacccgga ctgagatggc gaaatttatg gcgtgtgacc cgttatgtct 180
catttcggtc agtgggtcat tgctagtagt cgattgcatt gccattctcc gagtgtatta 240
gcgtgacagc cgcaggggaa ccataaaatg caatcgtagt ccacctgacg gtacttagaa 300
atgaggggtcc ccttttggcc acgcacctgt tcgctcgtcg tttgctttta agaaccgcac 360
gaaccacaga gcataaagag aacctctagc tcctttacaa ggtactgggt ccttttcag 420
cgggatgcct tatctaaacg caatgacaga cgtattctcc aggccacatc gcttctact 480
tcgctgggga tccatcattg gcggccgaag ccgccattcc atagttagtc cttcgtctgt 540
gtctttctgt gccagatcgt ctagcaaat cccgatccag tttatctcac gaaactatag 600
tcgtacagac cgaaatctta agtcaaatca cgcgactagg ctcagctcta ttttagtggt 660
catgggtttt ggtccgcgcg agcgggtgcaa ccgattagga ccatgtaaaa catttggtac 720
aagtcttctt ttaaacacaa tcttcctgct cagtggcgca tgattatcgt tgttgctagc 780
cagcgtggta agtaacagca ccactgcgag cctaagtgtc cctttccacg aacacaggcg 840
tgtccgatcc tatattagga ctccgcaatg ggggttagca gtgcacacct aaacgatgtt 900
gaagactcgc gatgtacatg ctctgttaca atacatacgt gttccggctg ttatcctgca 960
tcggaacctc aatcatgcat cgcaccagcg tattcgtgtc atctaggagg ggcgcgtagg 1020
ataaataatt caattaagat gtcgttatgc tagtatacgc ctaccgctca ccggccatct 1080
gtgtgcagat ggggcgacga gttactggcc ctgatttctc cgcttctaata accacacact 1140
gggcaatacg agtccaagcc agtctcgcag taacgctcat cagctaacga aagaggttaga 1200
ggctcgctaa cggagacgag ttaaaagacac gagttcccaa aaccaggcgg gctcgccacg 1260
acggctaata cgtgtagttt acgtgaacaa tgttctgaag aaaatttctg aaagaaggac 1320
ccgtcaccgc ctacaattac ctacaacggt cggccgcacc ttcgattgtc gtggccaccc 1380
tcggattaca cggcagaggt ggttgtgtcc cgacaggcca gcataattac ctgaggcggt 1440
acccaatcgc ttctccgtcg gatttgctac agcccctgag cgctacatgc acgaaaccaa 1500
gttatgtatg cactgggccc gtaaataggac gttagcctgt agttagcacg tagcccgccc 1560
gcattagtac agtagagcct ccgcccgcac cctgtttatt aagtatttcc tgcataaaag 1620
agtaacctca ccagaggact ggaatgatga gcagccagac agttagccat tcgccatcga 1680
cattgaaatg atggctcaac caggtggg 1707

```

```

SEQ ID NO: 12      moltype = DNA length = 2019
FEATURE            Location/Qualifiers
misc_feature       1..2019
                    note = lymphotropic variant sequence
source             1..2019
                    mol_type = genomic DNA
                    organism = Rodent protoparvovirus 1

```

```

SEQUENCE: 12
atggctggaa atgcttactc tgatgaagtt ttgggaacaa ccaactgggt aaaggaaaaa 60
agtaaccagg aagtgtttctc atttgttttt aaaactgagg atgttcaact aaatggaaaa 120
gatatcggat ggaataatta caaaaaggag ctgcaggagg acgagctgaa atctttacaa 180
cgaggagcgg aaactacctg ggaccaaagc gaggacatgg aatgggaatc tacagtggat 240
gaaatgacca aaaagcaagt attcatttat gactctttag ttaaaaaatg tttgtttgaa 300
gtgcttagca caaaaaatat agctcctgct gatgttactt ggtttgtgca gcatgaatgg 360
gggaaagacc aaggctggca ctgccatgta ctaattggag gcaaggactt tagtcaagct 420
caaggaaaaa ggtggagaag gcagctaaat gtttactgga gcagatgggt ggtaacagcc 480
tgtaatgtgc agctaacacc agctgaaaga attaaactaa gagaatagc agaagacagt 540
gagtgggtta ctttactcac ttataaacat aagcaaacca aaaaggacta tactaaatgt 600
gttctttttg gaaatatgat tgcttactac tttttaacca aaaagaaaaa aagcaccagt 660
ccgccaaggg ccggaggcta ttttctaagc agtgactctg gctggaaaac taacttttta 720
aaagaggggc aacgcatctc agtgagcaaa ttatacactg atgacatgcy gccagaaacg 780
gttgaaccca cagtaaccac tgcgcaggaa actaagcgcy gcagaattca aactaaaaaa 840
gaggtttcta ttaaaaccac acttaagag ctatgcata aaagagtaac ctcaccagaa 900
gactggatga tgatgcagc agacagttac attgaaatga tggctcaacc aggtggagaa 960
aacctgctga aaaatacgtc agagatttgt acgctaactc tagccagaac aaaaacagca 1020

```

-continued

```

tttgacttga ttttagaaaa agctgaacc agcaaaactaa ccaacttttc actgacctgac 1080
acaagaaacct gcaagatttt tgcttttcat ggctggaact atgtttaaagt ttgccatgct 1140
at ttgctgtg ttctaaacag acaaggaggc aaaagaaata ctgttttatt tcacggacca 1200
gccagtacag gcaaatctat tattgcacaa gccatagcac aggcagttgg taatgttgg 1260
tgctataatg cagctaattg gaactttcca tttaatgact gtaccaacaa gaacttgatt 1320
tgggtagaag aagctggtaa ctttggacag caagtaaacc agtttaaagc catttgctct 1380
ggtcaaaacta ttgcatttga tcaaaaagga aaaggcagca acaaaattga accaacacca 1440
gtcatcatga ccacaaatga gaacattaca gtggtcagaa taggctgcga agagagacca 1500
gaacacactc aaccaattag agacagaatg ctcaacattc atctaacaca tacattgcct 1560
ggtgactttg gtttgggtga caagaatgaa tggccatga tttgtgcttg gttggtaaag 1620
aatggttacc aatctaccat ggcaagctac tgcgctaaat ggggcaaaagt tctgatttg 1680
tcagaaaact gggcgagacc aaaggtgccc actcctataa attcactagg ttccggcacgc 1740
tcaccattca cgacaccgaa aagtacgcct ctcagccaga actatgcaat aactccactt 1800
gcatcggtac tcagggacct ggctttagag ccttggagca caccaaaatac tctgtttgct 1860
ggcactgcag aaacccagaa cactggggaa gctggttcca aagcctgcc aagatggtcaa 1920
ctgagcccaa cttggtcaga gatcgaggag gatttgagag cgtgcttcgg tcggaaccg 1980
ttgaagagag acttcagcga gccgctgaac ttggactaa 2019

```

```

SEQ ID NO: 13      moltype = DNA length = 2019
FEATURE
source             Location/Qualifiers
                   1..2019
                   mol_type = genomic DNA
                   organism = Rodent protoparvovirus 1

```

```

SEQUENCE: 13
atggctggaa atgcttactc tgatgaagtt ttgggagcaa ccaactgggt aaaggaaaaa 60
agtaaccagg aagtattctc atttgtttt aaaaatgagg atgttcaact gaatggaaaa 120
gatattggat ggaatagcta caaaaagag ctacaggagg acgagctgaa atctttacaa 180
cgaggagcgg aaactacatg ggacaaaagc gaggacatgg aatgggaatc tacagtggat 240
gaagtaacca aaagcaagt attcatttt gactctttag ttaaaaaatg ttgtttgaa 300
gtgcttaaca caaagaacat agctcctagt gatgttaatt ggtttgtaca gcatgaatgg 360
ggaaaagacc aaggctggca ttgccatgta ctaattggag gcaaggactt tagtcaagct 420
caaggaaagt ggtggagaag gcagctaaat gtttactgga gcagatgggt ggttaacagcc 480
tgtaatgtgc agctatcacc agctgaaaga attaaactaa gagaatagc agaagacaat 540
gagtggttga gcttgcctac ttataagcat aagcaaacca aaaaggacta tactaagtgt 600
gttctttttg gcaacatgat tgcttactac tttttaacca aaaagaaaat aagcactagt 660
ccaccaaggg acggaggcta ttttctaagc agtgactctg gctggaaaac taacttttta 720
aaagaaggcg aacgccatct agtgagcaaa ctatacactg atgacatgcg gccagaaacg 780
gttgaacca cagtaaccac tgcacaggaa actaagcgcg gcagaattca aactaaaaaa 840
gaggtttcta ttaaaaccac acttaagag ctggtgcata aaagagtaac ctaccagaa 900
gactggatga tgatgcagcc agacagttat attgaaatga tggctcaacc aggtggagaa 960
aacctgctga aaaatacgct agagatttgt acactaactc tagctagaac caaaacagca 1020
tttgacttga ttttagaaaa agctgaaacc agcaaaactaa ctaacttttc actgccggac 1080
acaagaaacct gcaagatttt tgcttttcat ggctggaact acattaaagt ttgcatgct 1140
at ttgctgtg ttttaaacag acaaggaggc aaaagaaata ctgttttatt tcatggacca 1200
gccagtacag gcaaatccat tattgcacaa gccatagcac aggcagttgg taatgttgg 1260
tgctataatg cagcaaatgt aaactttcca ttcaatgact gtaccaacaa gaacttgatt 1320
tgggtggaag aagctggtaa ctttggacag caagtaaacc agttcaaagc catttgctct 1380
ggtcaaaacta ttgcatttga tcaaaaagga aaaggcagca aacagattga accaacacca 1440
gtcatcatga ccacaaatga aaacattaca gtggtcagaa taggctgtga agaaagacca 1500
gagcacactc aaccaatcag agacagaatg cttaacattc atctaacaca tacattgcct 1560
ggtgactttg gtttgggtga caaaaatgag tggcccatga tttgtgcttg gttggtaaag 1620
aatggttacc aatctaccat ggcaagctac tgtgctaaat ggggcaaaagt tctgatttg 1680
acggaaaact gggcgagacc gaaggtgccc actcctataa attcactagg ttccggcacgc 1740
tcaccattca cgacaccgaa aagtacgcct ctcagccaga actatgcact aactccactt 1800
gcatcggtac tcagggacct ggctttagag ccttggagca caccaaaatac tctgtttgt 1860
ggcactgcag aaacccagaa cactggggaa gctggttcca aagcctgcc aagatggtcaa 1920
ctgagcccaa cttggtcaga gatcgaggag gatttgagag cgtgcttcgg tcggaaccg 1980
ttgaagaaag acttcagcga gccgctgaac ttggactaa 2019

```

```

SEQ ID NO: 14      moltype = DNA length = 2042
FEATURE
source             Location/Qualifiers
                   1..2042
                   mol_type = genomic DNA
                   organism = Rodent protoparvovirus 1

```

```

SEQUENCE: 14
atggctggaa atgcttactc tgatgaagtt ttgggagcaa ccaactgggt aaaggaaaaa 60
aataatcagg aagtgttctc atttgtttt aaaaatgagg atgttcaact gaatggaaaa 120
gatatcggtat ggaatagtta caaaaagag ctgacaggagg acgagctgaa atctttacaa 180
cgaggagcgg aaactacatg ggacaaaagc gaggacatgg aatgggaatc tgcaagtggat 240
gaagtgacca aaaagcaagt attcattttt gactctttag ttaaaaaatg ttgtttgaa 300
gtgcttaaca caaagaacat agctcctgct gatgttaatt ggtttgtgca gcatgaatgg 360
ggaaaagacc aaggctggca ctgccatgta ctaattggag gcaaggactt tagtcaagct 420
caaggaaagt ggtggagaag gcagctaaat gtttactgga gcagatgggt agtaacagcc 480
tgtaatgtac agctatcacc agctgaaaga attaaactaa gagaatagc agaagacaat 540
gagtggttga ccttgcctac ttataagcat aagcaaacca aaaaggacta tactaagtgt 600
gttctttttg gcaacatgat tgcttactac tttttaacca aaaagaaaat aagcactagt 660
ccaccaaggg acggaggcta ttttctgagc agtgactctg gctggaaaac taacttttta 720
aaagaggcg aacgccatct agtgagcaaa ctatatactg atgacatgcg gccagaaacg 780
gttgaacca cagtaaccac tgcgcaggaa actaagcgcg gcagaattca aactaaaaaa 840

```


-continued

```

gagggtttcaa ttaaaaccac acttaaagag ctggtgcata aaagagtaac ctcaccagaa 900
gactggatga tgaatgcagcc agacagttac attgaaatga tggctcaacc aggtggagaa 960
aacctgctga aaaatacgcct agagatttgt acactaactc tagccagaac caaaacagca 1020
tttgacttga ttttagaaaa agctgaaacc agcaaaactaa caaacttttc actgctgat 1080
acaagaacct gcaagatttt tgcttttcat ggctggaact acattaaagt ttgcatgct 1140
atttgcctgt ttttaaacag acaaggaggc aaaagaaata ctgttttatt tcatggacca 1200
gccagtacag gcaaatctat cattgcacaa gccatagcac aggcagttgg taatgttgg 1260
tgctataatg cagcaaatgt gaactttcca ttcaatgact gtaccaacaa gaacttaatt 1320
tgggtggaag aagctggtta ctttgacaa caagtaaaacc agtttaaagc ctttgcct 1380
ggtcaaaacta ttcgcattga tcaaaaagga aaaggcagca agcagattga accaacacca 1440
gtcatcatga ccacaaatga aaacatcaca gtggtcagaa taggctgcga agagagacca 1500
gaacacactc aaccaatcag agacagaatg cttaacatc atctaacaca tacattgcct 1560
ggtgactttg gtttggttga caaaaatgag tggcccatga tttgtgcttg gttggtaaag 1620
aatgggtacc aatctaccat ggcaagctac tgtgctaaat ggggcaaaagt tctgatttg 1680
acagaaaaact gggcgagcc gaaggtgccg actcctataa attcactagg ttcggcacgc 1740
tcaccattca cgcaccgaa aagtacgcct ctacgccaga actatgcact aactccactt 1800
gcatcgatc tcgaggacct ggctttagag ccttgaggca caccaaatac tctgtttgcg 1860
ggcactgcag aaacccagaa cactggggaa gctggttcca aagcctgcc aagatggtcaa 1920
ctgagcccaa cttggtcaga gatcgaggag gatttgagag cgtgcttcgg tgcggaaccg 1980
tgaagaaag acttcaacga gccgctgaac ttggactaag gtacgatggc gcctccagct 2040
aa 2042

```

```

SEQ ID NO: 15      moltype = DNA length = 2019
FEATURE            Location/Qualifiers
source             1..2019
                   mol_type = genomic DNA
                   organism = Rodent protoparvovirus 1

```

```

SEQUENCE: 15
atggctggaa atgcttactc tgatgaagtt ttgggaacaa ccaactgggt aaaggaaaaa 60
agtaaccagg aagtgtcttc atttgttttt aaaactgagg atgttcaact aaatggaaaa 120
gatatcggat ggaataatta caaaaaggag ctgcaggagg acgagctgaa atctttacaa 180
cgaggagcgg aaactacctg ggaccaaagc gaggacatgg aatgggaatc tacagtggat 240
gaaatgacca aaaagcaagt attcatttat gactctttag ttaaaaaatg tttgtttgaa 300
gtgcttagca caaagaatat agctcctgct gatgttactt gggtttgtgca gcatgaatgg 360
gggaaagacc aaggctggca ctgccatgta ctaattggag gcaaggactt tagtcaagct 420
caaggaaaat ggtggagaag gcagctaaat gtttactgga gcagatgggt ggtaacagcc 480
tgtaatgtgc agctaacacc agctgaaaga attaaactaa gagaatagc agaagacagt 540
gagtggttga ctttgcctac ttataagcat aagcaaacca aaaaggacta taccagtggt 600
gttctttttg gaaacatgat tgcttactac tttttaacca agaagaaaat aagcactagt 660
ccgccaaggg ccgagggtca ttttctgagc agtgactctg gctggaaaac taacttttta 720
aaagaggggc aagccatctc agtgagcaaa ctatacactg atgacatgag gccagaaacg 780
gttgaaccca cagtaaccac tgcgcaggaa actaagcgcg gcagaattca aactaaaaaa 840
gagggtttct ttaaaaccac acttaaagag ctggtgcata aaagagtaac ctcaccagaa 900
gactggatga tgaatgcagc agacagttac attgaaatga tggctcaacc aggtggagaa 960
aacctgctga aaaatacgcct agagatctgt acactaactc tagctagaac caaaacagca 1020
tttgacttga ttttagaaaa agctgaaacc agcaaaactaa ccaacttttc actgctgac 1080
acaagaacct gcaagatctt tgcttttcat ggctggaact acattaaagt ttgcatgct 1140
atttgcctgt ttctaaacag acaaggaggc aaaagaaata ctgttttatt tcatggacca 1200
gccagtacag gcaaatccat tattgcacaa gccatagcac aggcagttgg taatgttgg 1260
tgctataatg cagcaaatgt gaactttcca ttcaatgact gtaccaacaa gaacttgatt 1320
tgggtggaag aagctggtta ctttgacaa caagtaaaacc agtttaaagc ctttgcct 1380
ggtcaaacaa ttcgcattga tcaaaaagga aaaggcagca agcagattga accaacacca 1440
gtcatcatga ccacaaatga gaacattaca gtggtcacaa taggctgcga ggagagacca 1500
gaacacactc aaccaatcag agacagaatg cttaacatc atctaacaca tacattgcct 1560
ggtgactttg gtttggttga caaaagttag tggcccatga tctgtgcttg gttggtaaag 1620
aatgggtacc aatctaccat ggcaagctac tgtgctaaat ggggcaaaagt tctgatttg 1680
acagaaaaact gggcgagcc gaaggtgccg actcctataa attcactagg ttcggcacgc 1740
tcaccattca cgcaccgaa aagtacgcct ctacgccaga actatgcact aactccactt 1800
gcatcgatc tcgaggacct ggctttagag ccttgaggca caccaaatac tctgtttgcg 1860
ggcactgcag aaacccagaa cactggggaa gctggttcca aagcctgcc aagatggtcaa 1920
ctgagcccaa cttggtcaga gatcgaggag gatttgagag cgtgcttcgg tgcggaaccg 1980
tgaagagag acttcagcga gccgctgaac ttggactaa 2019

```

```

SEQ ID NO: 16      moltype = DNA length = 2019
FEATURE            Location/Qualifiers
misc_feature       1..2019
                   note = immunosuppressive variant sequence
source             1..2019
                   mol_type = genomic DNA
                   organism = Rodent protoparvovirus 1

```

```

SEQUENCE: 16
atggctggaa atgcttactc tgatgaagtt ttgggaacaa ccaactgggt aaaggaaaaa 60
agtaaccagg aagtgtcttc atttgttttt aaaactgagg atgttcaact aaatggaaaa 120
gatatcggat ggaataatta caaaaaggag ctgcaggagg acgagctgaa atctttacaa 180
cgaggagcgg aaactacctg ggaccaaagc gaggacatgg aatgggaatc tacagtggat 240
gaaatgacca aaaagcaagt attcatttat gactctttag ttaaaaaatg tttgtttgaa 300
gtgcttagca caaaaaatat agctcctgct gatgttactt gggtttgtgca gcatgaatgg 360
gggaaagacc aaggctggca ctgccatgta ctaattggag gcaaggactt tagtcaagct 420
caaggaaaat ggtggagaag gcagctaaat gtttactgga gcagatgggt ggtaacagcc 480

```

-continued

```

tgtaatgtgc agctaaccac agctgaaaga attaaactaa gagaaatagc agaagacagt 540
gagtggggtta ctttactcac ttataaacat aagcaaacca aaaaggacta tactaaatgt 600
gttcttttttg gaaatatgat tgcttactac tttttaacca aaaagaaaat aagcaccagt 660
ccgccaaggcg acggagggcta ttttctaagc agtgactctg gctggaaaac taacttttta 720
aaagagggcg aacgccatct agtgagcaaa ttatacactg atgacatgcg gccagaaacg 780
gttgaaacca cagtaaccac tgcgcaggaa actaagcgcg gcagaattca aactaaaaaa 840
gagggtttcta ttaaaaccac acttaaagag ctagtgcata aaagagtaac ctcaccagaa 900
gactggatga tgatgcagcc agacagttac attgaaatga tggctcaacc aggtggagaa 960
aacctgctga aaaatacgtc agagatttgt acgctaactc tagccagaac aaaaacagca 1020
tttgacttga ttttagaaaa agctgaaacc agcaaaacta ccaacttttc actgctgac 1080
acaagaacct gcaagatttt tgcttttcat ggctggaact atgttaaagt ttgccatgct 1140
atthtctgtg ttctaaccag acaaggagcg aaaagaaata ctgttttatt tcacggacca 1200
gccagtacag gcaaatctat tattgcacaa gccatagcac aggcagttgg taatgttgg 1260
tgctataatg cagctaatgt gaactttcca tttaatgact gtacgaacaa aaacttgatt 1320
tgggtagaag aagctggtta ctttgacag caagtaaaac agtttaaagc catttgctct 1380
ggtcaaaacta ttcgcattga tcaaaaagga aaaggcagca acaaaatga accaacacca 1440
gtcatcatga ccacaaatga gaacattaca gtggtcagaa taggctgcga agagagacca 1500
gaacacactc aaccaattag agacagaatg ctcaacattc atctaacaca tacattgcct 1560
ggtgactttg gtttggttga caagaatgaa tggcccatga tttgtgcttg gttggtaaag 1620
aatgggtacc aatctaccat ggcaagctac tgcgctaaat ggggcaaaagt tctgatttgg 1680
tcagaaaaact gggcgagcc gaaggtgccc actcctataa attcactagg ttccggcacgc 1740
tcaccattca cgacaccgaa aagtacgcct ctacgccaga actatgcact aactccactt 1800
gcatcgatc tcgaggaact ggctttagag ccttgagaca caccaaaatac tctgttgcg 1860
ggcactgcag aaacccagaa cactggggaa gctggttcca aagcctgcc aagatggtcaa 1920
ctgagcccaa cttggtcaga gatcgaggag gatttgagag cgtgcttcgg tgcggaaccg 1980
ttgaagagag acttcagcga gccgctgaac ttggactaa 2019

```

```

SEQ ID NO: 17      moltype = DNA length = 2019
FEATURE            Location/Qualifiers
source              1..2019
                    mol_type = genomic DNA
                    organism = Rodent protoparvovirus 1

```

```

SEQUENCE: 17
atggctggaa atgcttactc tgatgaagtt ttgggaacaa ccaactgggt aaaggaaaaa 60
agtaaccagg aagtggtctc atttgttttt aaaactgagg atgttcaact aaatggaaaa 120
gatatcggtt ggaataatta caaaaaggag ctgcaggagg acgagctgaa atctttacaa 180
cgaggagcgg aaactacatg ggaccaaagc gaggacatgg aatgggaatc tacagtggat 240
gaaatgacca aaaagcaagt attcatttat gactctttag ttaaaaaatg tttggttgaa 300
gtgcttagca caaagaatat agctcctgct gatgttactt ggtttgtgca gcatgaatgg 360
gggaaagacc aaggtctggca ctgccatgta ctaattggag gcaaggactt tagtcaagct 420
caaggaaaaa ggtggagaag gcagctaaat gtttactgga gcagatgggt ggtaacagcc 480
tgtaatgtgc agctaaccac agctgaaaga attaaactaa gagaaatagc agaagacagt 540
gagtggttta ctttacttac ttataaacat aagcaaacca aaaaggacta tactaaatgt 600
gttctttttg gaaatatgat tgcttactac tttttaacca aaaaaaaaat aagcaccagt 660
ccgccaagag acggaggcta ttttctaagc agtgactctg gctggaaaac taacttttta 720
aaagaaggcg acgcccattc agtgagcaaa ctatacactg atgacatgcg accagaaacg 780
gttgaaacca cagtaaccac tgcgcaggaa actaagcgcg gcagaattca aactaaaaaa 840
gagggtttcta ttaaaaccac acttaaagag ctggtgcata aaagagtaac ctcaccagaa 900
gactggatga tgatgcagcc agacagttac attgaaatga tggctcaacc aggtggagaa 960
aacctgctga aaaatacgtc agagatctgt acactaactc tagctagaac caaaacagca 1020
tttgacttga ttttagaaaa agctgaaacc agcaaaacta ccaacttttc actgctgac 1080
acaagaacct gcaagattct tgcttttcat ggctggaact acattaaagt ttgccatgct 1140
atthtctgtg ttctaaccag acaaggagcg aaaagaaata ctgttttatt tcacggacca 1200
gccagtacag gcaaatccat tattgcacaa gccatagcac aggcagttgg taatgttgg 1260
tgctataatg cagcaaatgt gaactttcca ttcaatgact gtaccaacaa gaacttaatt 1320
tgggtggaag aagctggtta ctttgacaa caagtaaaac agtttaaagc catttgctct 1380
ggtcaaacaa ttcgcattga tcaaaaagga aaaggcagca agcagattga accaacacca 1440
gtcatcatga ccacaaatga aaacattaca gtggtcaaaa taggctgcga ggaagacca 1500
gaacacactc aaccaatcag agacagaatg cttaacattc atctaacaca tacattgcct 1560
ggtgactttg gtttggttga caaaaatgag tggcccatga tttgtgcttg gttggtaaag 1620
aatgggtacc aatctaccat ggcaagctac tgtgctaaat ggggcaaaagt tctgatttgg 1680
acggaaaaact gggcgagcc gaaggtgccc actcctataa attcactagg ttccggcacgc 1740
tcaccattca cgacaccgaa aagtacgcct ctacgccaga actatgcact aactccactt 1800
gcatcgatc tcgaggaact ggctttagag ccttgagaca caccaaaatac tctgttgcg 1860
ggcactgcag aaacccagaa cactggggaa gctggttcca aagcctgcc aagatggtcaa 1920
ctgagcccaa cttggtcaga gatcgaggag gatttgagag cgtgcttcgg tgcggaaccg 1980
ttgaagaaag acttcagcga gccgctgaac ttggactaa 2019

```

```

SEQ ID NO: 18      moltype = DNA length = 2019
FEATURE            Location/Qualifiers
source              1..2019
                    mol_type = genomic DNA
                    organism = Rodent protoparvovirus 1

```

```

SEQUENCE: 18
atggctggaa atgcttactc tgatgaagtt ttgggaacaa ccaactgggt aaaggaaaaa 60
agtaaccagg aagtggtctc atttgttttt aaaactgaag atgttcaact gaatggaaaa 120
gatattggtt ggaataatta cgaaaaggag ctgcaaggag acgagctaaa atctttacaa 180
cgaggagcgg aaactacatg ggaccaaagc gaggacatgg aatgggaatc tacagtggat 240
gaagtgacca aaaagcaagt attcatttat gactctttag ttaaaaaatg tttggttgaa 300

```

-continued

```

gtacttagca caaagaacat agctcctagt gatgttaatt gggttgtgca gcatgaatgg 360
ggaagagacc aagcgtggca ttgccatgta ctaattggag gcaaagactt tagtcaagct 420
caaggaaagt ggtggagaag gcagctaaagt gtttactgga gcagatgggt ggtaacagct 480
tgtaatgtac agctaaccac agctgaaaga attaaactaa gagaatagc agaagacagt 540
gaatggggtta ccttgctcac ttataagcat aagcaaacca aaaaggacta tacciaagtgt 600
gttctttttg gaaacataat tgcttactac tttttaacta aaaagaaat aagcaccagt 660
ccgccaagag acggaggcta tttcttagc agtgactctg gctggaaaac taacttttta 720
aaagagggcg aacgccatct agtgagcaaa ctatacactg atgacatgcg gccagaaacg 780
gttgaacca cagtaaccac tgcgcaggaa actaagcgcg gcagaattca aactaaaaaa 840
gagggttcaa ttaaaaccac acttaaagag ctggtgcata agagagtaac ctcaccagaa 900
gactggatga tgaatgagcc agacagttat attgaaatga tggctcaacc aggtggagaa 960
aacctgctaa aaaatacgct agagatttgt acactaactc tagctagaac caaacacgca 1020
tttgacttga ttttagaaaa agctgaaacc agcaaaactaa ccaacttttc actgccggac 1080
acaagaacct gcaaaatfff tgcttttcat ggctggaact atgttaaagt ttgcatgct 1140
atcttctgtg ttctaaacag acaaggaggc aagagaaata ctgttttatt tcatggacca 1200
gccagcacag gcaaatccat tattgcacaa gccatagcac aggcagttgg taatgttgg 1260
tgctataatg cagcaaatgt gaactttcca ttcaatgact gtaccaacaa gaacttaatt 1320
tgggtggaag aagctggtta ctttgacaaa caagtaaac accgttaaagc ctttctct 1380
ggtcaacaaa ttcgcattga tcaaaaagga aaaggcagca agcagattga accaacacca 1440
gtcatcatga ccacaaatga aaacattaca gtgggtcaaaa taggctgcga ggaaagacca 1500
gaacacactc aaccaatcag agacagaatg cttaacattc atctaacaca tacattgcct 1560
ggtgactttg atttggttga caaaaatgag tggcccatga tttgtgcttg gttggtaaag 1620
aatggttacc aatctaccat ggcgaagctac tgtgctaaat ggggcaaaagt tctgattgg 1680
acggaaaact gggcgagacc gaaggtgccc actcctataa attcactagg ttccggcacgc 1740
tcaccattca cgacaccgaa aagtagccct ctacgccaga actatgcact aactccactt 1800
gcatcgatc tcgaggacct ggctttagag ccttgagaca caccaaaatac tctgttgcg 1860
ggcactgcag aaacccagaa cactggggaa gctggttcca aagcctgccca agatggtcaa 1920
ctgagcccaa cttggtcaga gatcgaggag gatttgagag cgtgcttcgg tgcggaaccg 1980
ttgaagaaag acttcagcga gccgctgaac ttggactaa 2019

```

```

SEQ ID NO: 19      moltype = DNA length = 2019
FEATURE            Location/Qualifiers
source              1..2019
                    mol_type = genomic DNA
                    organism = Rodent protoparvovirus 1

```

```

SEQUENCE: 19
atggctggaa acgcttactc tgatgaagtt ttgggaacaa ccaactgggt aaaggaaaaa 60
agtaaccagg aagtggtctc atttgttttt aaaactgagg atgttcaact aaatggaaaa 120
gatatcggat ggaataatta cagaaaggag ctgcaggagg acgagctgaa atctttacaa 180
cgaggagcag aaactacatg ggaccaaagc gaggacatgg aatgggaatc tacatgggat 240
gaagtgaaca aaaagcaagt attcatttat gactctttag ttaaaaaatg ttgtttgaa 300
gtgcttaaca caaagaacat atctcctggt gatgttaatt gggttgtgca gcatgaatgg 360
ggaaaagacc aagcgtggca ttgccatgta ctaattggag gcaaagactt tagccaagct 420
caaggaaaat ggtggagaag gcagctaagt gtttactgga gcagatgggt agtaacagcc 480
tgtaatgtgc agctatcacc agctgaaaga attaaactaa gagaatagc agaagacagt 540
gagtggtgta ccttgctcac ttataagcat aagcaaacca aaaagacta tactaagtgt 600
gttctttttg gcaacataat tgcttactac tttttaacca agaagaaaaa aagcactagt 660
ccgccaaggg acggaggcta tttctgagc agtgactctg gctggaaaac taacttttta 720
aaagagggcg aacgccatct agtgagcaaa ctatacactg atgacatgcg gccagaaacg 780
gttgaacca cagtaaccac tgcgcaggaa actaagcgcg gcagaattca aactaaaaaa 840
gaagtttcta ttaaaaccac acttaaagaa ctggtgcata aaagagtaac ctcaccagaa 900
gactggatga tgatgcagct agacagttac attgaaatga tggctcaacc aggtggagaa 960
aacctgctaa aaaatacgct agagatttgt acgctaactc tagctagaac caaacacgca 1020
tttgacttga ttttagaaaa agctgaaacc agcaaaactaa ctaacttttc actgcctgac 1080
acaagaacct gcaagatttt tgcttttcat ggctggaact acattaaagt ttgcatgct 1140
atcttctgtg ttctaaacag acaaggaggc aaaagaaata ctgttttatt tcatggacca 1200
gccagtacag gcaaatccat tattgcacaa gccatagcac aggcagttgg taatgttgg 1260
tgctataatg cagcaaatgt gaactttcca ttcaatgact gtaccaacaa gaacttgatt 1320
tgggtggaag aagctggtta ctttgacaaa caagtaaac accgttaaagc ctttctct 1380
ggtcaacaaa ttcgcattga tcaaaaagga aaaggcagca agcagattga accaacacca 1440
gtcatcatga ccacaaatga gaacattaca gtgggtcaaaa taggctgcga ggagagacca 1500
gaacacactc aaccaataag agacagaatg cttaacattc atctaacaca tacattgcct 1560
ggtgactttg gtttggttga caaaaagtga tggcccatga tctgtgcttg gttggtaaag 1620
aatggttacc aatctaccat ggcgaagctac tgtgctaaat ggggcaaaagt tctgattgg 1680
tcagaaaact gggcgagacc gaaggtgccc actcctataa attcactagg ttccggcacgc 1740
tcaccattca cgacaccgaa aagtagccct ctacgccaga actatgcact aactccactt 1800
gcatcgatc tcgaggacct ggctttagag ccttgagaca caccaaaatac tctgttgcg 1860
ggcactgcag aaacccagaa cactggggaa gctggttcca aagcctgccca agatggtcaa 1920
ctgagcccaa cttggtcaga gatcgaggag gatttgagag cgtgcttcgg tgcggaaccg 1980
ttgaagaaag acttcagcga gccgctgaac ttggactaa 2019

```

```

SEQ ID NO: 20      moltype = DNA length = 2009
FEATURE            Location/Qualifiers
source              1..2009
                    mol_type = genomic DNA
                    organism = Rodent protoparvovirus 1

```

```

SEQUENCE: 20
atgcttactc tgatgaagtt ttgggaacaa ccaactgggt aaaggaaaaa agtaaccagg 60
aagtggtctc atttgttttt aaaactgaag atattcaact gaatggaaaa gatattggat 120

```

-continued

```

ggaataatta cagaagaggag ctgcaagagg acgagctaaa atctttacaa cgaggagcgg 180
aaactacctg ggacccaaagc gagggacatgg aatgggaatc tacagtggat gaagtgacca 240
aaaagcaagt attcatttat gactcttttag ttaaaaaatg tttgtttgaa gtgcttaaca 300
caaagaacat atctcctgggt gatgttaatt ggtttgtgca gcatgaatgg ggaagagacc 360
aaggctggca ttgcoattgta ctaattggag gcaaagactt tagtcaagct caaggaaagt 420
ggtggagaag gcagctaaagt gtttactgga gcagatggtt ggtaacagcc tghtaatgtac 480
agctaaccac agctgaaaga attaaactaa gagaatatgc agaagacagt gaatgggtta 540
ccttgctcac ttataagcat aagcaaacca aaaaggacta taccaagtgt gttctttttg 600
gcaacataat tgcttactac tttttaacca agaagaaaaa aagcaccagt ccgccaaagg 660
acggaggcta ttttcttagt agtgactctg gctggaaaac taacttttta aaagagggcg 720
aacgccatct agtgagcaaa ctatacactg atgacatgcg gccagaaacg gttgaaacca 780
cagtaaacac tgcgcaggaa actaagcgcg gcagaattca aactaaaaaa gaggtttcta 840
ttaaaaccac acttaaagag ctagtgcata agagagtaac ctcaccagaa gactggatga 900
tgatgcagcc agacagttaa attgaaatga tggctcaacc aggtggagaa aacctgctga 960
aaaaacgct agagatctgt acactaactc tagctagaac caaacagca tttgacttga 1020
tttagaaaa agctgaaac agcaaaactaa ccaacttttc actgcctgac acaagaacct 1080
gcaagatctt tgcttttcat ggctggaact acattaaagt ttgccatgct atttgctgtg 1140
ttctaaccag acaaggaggc aaaagaaata ctgttttatt tcatggacca gccagtacag 1200
gcaaatccat tatgcaaaa gccatagcac aggcagttgg taatgttggg tgctataatg 1260
cagcaaatgt gaactttcca ttcaatgact gtaccaacaa gaacttaatt tgggtggaag 1320
aagctggtaa ctttgacaaa caagttaacc agtttaagc catttgctct ggtcaacaaa 1380
ttgcattga tcaaaaagga aaaggcagca agcagattga accaacacca gtcacatga 1440
ccacaaatga aaacattaca gtggctcaaaa taggctgcga ggaagacca gaacacactc 1500
aaccaatcag agacagaatg ctttaacattc atctaacaca tacattgcct ggtgactttg 1560
gtttggttga caaaaatgag tggcccatga tttgtgctg gttggtaaa aatgggtacc 1620
aatctaccat ggaagctcac tgtgctaaat ggggcaaaagt tctgatttg acggaaaaact 1680
gggcgaggcc gaaggtgccc actcctataa attcactagg ttccggcacgc tcaccattca 1740
cgacaccgaa aagtagccct ctacgcccaga actatgcact aactccactt gcatcggtac 1800
tcgaggacct ggtcttagag ccttggagca caccaaaatac tctgtgtgcg ggcactgcag 1860
aaaccagaa cactggggaa gctggttcca aagcctgcc aagatggtcaa ctgagcccaa 1920
cttggtcaga gctcgaggag gatttgagag cgtgcttcgg tgcggaaccg ttgaagagag 1980
acttcagcga gccgctgaac ttggactaa 2009

```

```

SEQ ID NO: 21      moltype = DNA length = 2019
FEATURE
variation          Location/Qualifiers
                    609
                    replace =
variation          1590
                    replace =
source             1..2019
                    mol_type = genomic DNA
                    organism = Rodent protoparvovirus 1

```

```

SEQUENCE: 21
atggctggaa atgcttactc tgatgaagtt ttgggaacaa ccaactgggt aaaggaaaaa 60
agtaaccagg aagtgttctc atttgttttt aaaactgaag atgttcaact aaatggaaaa 120
gatattggat ggaataatta cagaagaggag ctgcaagagg acgagctaaa atctttacaa 180
cgaggagcgg aaactacctg ggacccaaagc gagggacatgg aatgggaatc tacagtggat 240
gaagtgacca aaaagcaagt attcatttat gactcttttag ttaaaaaatg tttgtttgaa 300
gtgcttaaca caaagaacat atctcctgggt gatgttaatt ggtttgtgca gcatgaatgg 360
ggaagagacc aaggctggca ttgccatgta ctaattggag gcaaagactt tagtcaagct 420
caaggaaagt ggtggagaag gcagctaaagt gtttactgga gcagatggtt ggtaacagcc 480
tgtaatgtac agctaaacac agctgaaaga attaaactaa gagaatatgc agaagacagt 540
gaatgggtta ccttgctcac ttataagcat aagcaaacca aaaaggacta taccaagtgt 600
gttcttttng gcaacataat tgcttactac tttttaacca agaagaaaaa aagcaccagt 660
ccgccaaagg agcgaggcta ttttcttagt agtgactctg gctggaaaac taacttttta 720
aaagagggcg aacgccatct agtgagcaaa ctatacactg atgacatgcg gccagaaacg 780
gttgaacca cagtaaccac tgcgcaggaa actaagcgcg gcagaattca aactaaaaaa 840
gagggtttct ttaaaaccac acttaaagag ctagtgcata agagagtaac ctcaccagaa 900
gactggatga tgatgcagcc agacagttaa attgaaatga tggctcaacc aggtggagaa 960
aacctgctga aaaatacgtg agagatctgt acactaactc tagctagaac caaacagca 1020
tttgacttga ttttagaaaa agctgaaacc agcaaaactaa ccaacttttc actgcctgac 1080
acaagaacct gcaagatctt tgcttttcat ggctggaact acattaaagt ttgccatgct 1140
atthtctgtg ttctaaccag acaaggaggc aaaagaaata ctgttttatt tcatggacca 1200
gccagtacag gcaaatccat tattgcacaa gccatagcac aggcagttgg taatgttggg 1260
tggtataatg cagcaaatgt gaactttcca ttcaatgact gtaccaacaa gaacttaatt 1320
tgggtggaag aagctggtta ctttgacaaa caagttaacc agtttaagc catttgctct 1380
ggtcaacaaa ttgcgattga tcaaaaagga aaaggcagca agcagattga accaacacca 1440
gtcatctatg ccacaaatga aaacattaca gtggctcaaaa taggctgcga ggaagacca 1500
gaacacactc aaccaatcag agacagaatg ctttaacattc atctaacaca tacattgcct 1560
ggtgactttg gtttgggtga caaacatgan tggcccatga tttgtgcttg gttggtaaa 1620
aatgggtacc aatctaccat ggcaagctac tgtgctaaat ggggcaaaagt tctgatttg 1680
acggaaaact gggcgaggcc gaaggtgccc actcctataa attcactagg ttccggcacgc 1740
tcaccattca cgacaccgaa aagtacgcct ctcagccaga actatgcact aactccactt 1800
gcatcgatc tcgaggacct ggtcttagag ccttggagca caccaaaatac tctgtgtgcg 1860
ggcactgcag aaaccocgaa cactggggaa gctggttcca aagcctgcc aagatggtcaa 1920
ctgagcccaa cttggtcaga gatcgaggag gatttgaaag cgtgcttcgg tgcggaaccg 1980
ttgaagagag acttcagcga gccgctgaac ttggactaa 2019

```

```

SEQ ID NO: 22      moltype = DNA length = 2019

```

-continued

FEATURE	Location/Qualifiers					
source	1..2019					
	mol_type = genomic DNA					
	organism = unidentified					
SEQUENCE: 22						
atggctggaa	atgcttactc	tgatgaagtt	ttgggaacaa	ccaactgggt	aaaagagaaa	60
agtaaccagg	aagtgttctc	atttgttttt	aaaaatgaag	atgttcagct	caatggaaaa	120
gatatcggat	ggaatagtta	caaaaaggag	ctgcaagagg	aagagctgaa	atctttacaa	180
cgaggagcgg	aaactacctg	ggaccagagc	gaggacatgg	aatgggaatc	ttcagtggtg	240
gaactaacca	aaaagcaagt	attcattttt	gactctttag	ttaaaaaatg	tttgtttgaa	300
gtgctgagta	caaagaacat	agcacctagt	gatgttactt	ggtttgtaca	gcacgaatgg	360
ggaaaagacc	aaggtctgca	ctgtcatgtg	ctaattggag	gcaaggactt	tagccaagct	420
caaggaaaat	gggtggagaag	gcagttaa	gtttactgga	gcagatgggt	ggtaacagcc	480
tgtagtgtgc	agctattacc	agctgaaaga	attaaagctg	gagagatagc	ggaagaccaa	540
gaatgggtca	ctttgcttac	ttataagcat	aagcaaacca	aaaagacta	taccaagtgt	600
gtttgctttg	gaaatatagt	tgcttactac	tttttatcca	agaagaaaat	atgcaccagt	660
ccaccaaggg	acggaggtga	ttttcttagc	agtgaactcg	gctggaaaac	taacttttta	720
aaagaaggcg	aacgccatct	agtgaagcaa	ctatacactg	atgacatgcg	gccagaaacg	780
gttgaiaacc	cagtaaccac	tgccgaggaa	actaagcgcg	gcagaattca	aactaaaaaa	840
gaggtctcta	ttaaaaccac	acttaagag	ctgggtgcata	agagagtaac	ctcaccagaa	900
gactggatga	tgtatgcagc	agacagttac	attgaaatga	tggctcaacc	aggtggagaa	960
aaactgctga	aaaatacgtg	agagatttgt	acactaactc	tagccagaac	aaaaacagca	1020
tttgacttaa	ttttgaaaaa	agctgaaacc	agcaaaactaa	ccaacttttc	actgccggac	1080
acaagaacct	gcaagatctt	tgcttttcat	ggctggaaac	atattaaagt	ttgccatgct	1140
atttgctgtg	ttctaaacag	acaaggagcg	aaaagaaata	ctgttttatt	tcacggacca	1200
gccagtacag	gcaaatccat	tattgcacaa	gccatagcac	aggcagttgg	taatgttggt	1260
tgtataatg	cagcaaatgt	gaactttcca	tttaagtact	gcaccaacaa	aaacctgatt	1320
tggttggaag	aagctggtta	ctttggacag	caagtaaac	agtttaaacg	catttgctct	1380
gggtcaaaact	tttcgattga	tcaaaaagga	aaaggcagca	aacagattga	accaaacacca	1440
gtcatcatga	ccacaatga	aaacattaca	gtgggtcaaaa	taggctgtga	agaaagacca	1500
gaacacactc	aaccaatcag	agacagaatg	cttaacattc	atctaacaca	tacattgcct	1560
ggtagctttg	gtttggttga	caaacatgaa	tggtccatga	tttgtgcttg	gttggttaag	1620
aatggttacc	aattaccat	ggcaagctac	tgtgctaaat	ggggcaaaag	tcctgattgg	1680
acggaaaact	ggggcgagcc	gaaggtgccg	actcctataa	attcactagg	ttcggcacgc	1740
tcaccattca	cgacacagaa	aagtacgcct	ctcagccaga	actatgcact	aactccactt	1800
gcacggatc	tcgaggacct	ggcttttagag	ccttgagcga	caccaaaatac	tcctgttgcg	1860
ggcactcgag	aaacccagaa	cactggggaa	gctgggtcca	aagcctgccca	agatgggtcaa	1920
ctgagcccaa	cttggtcaga	gatcgaggag	gatttgagag	cgtgcttcgg	tgcggaaccg	1980
ttgaagagag	acttcagcga	gccgctgaac	ttggactaa		2019	

SEQ ID NO: 23	moltype = DNA	length = 2019
FEATURE	Location/Qualifiers	
source	1..2019	
	mol_type = genomic DNA	
	organism = Rodent protoparvovirus 1	

SEQUENCE: 23						
atggctggaa	acgcttactc	tgatgaagtt	ttaggaacaa	ccaactgggt	aaaggaaaaa	60
agtaaccagg	aagtgttctc	atttgttttt	aaaaatgaag	atgttcaact	gaatggaaaa	120
gatatcggat	ggaatagtta	cagaaggag	ctgcaggagg	acgagctgaa	atctttacaa	180
cgaggagcgg	aaactacctg	ggaccaaagc	gaggacatgg	aatgggaatc	tcagtggtg	240
gaactaacca	aaaagcaagt	attcattttt	gactctttag	ttaaaaaatg	tttgtttgaa	300
gtgctgagta	caaagaacat	agctcctagt	gatgttactt	ggtttgtacg	gcacgaatgg	360
ggaaaagacc	aaggtctgca	ctgtcatgtg	ctcattggag	gcaaggactt	tagccaagct	420
caaggaaaat	gggtggagaag	gcagttaa	gtttactgga	gcagatgggt	agtaacagcc	480
tgtaatgtgc	agttatcacc	agctgaaaga	attaaagctg	gagagatagc	ggaagaccaa	540
gaatgggtca	ctttgcttac	ttataagcat	aagcaaacca	aaaaggacta	taccaagtgt	600
gttctttttg	gaaatatagt	tgcttactac	tttttaacca	agaagaaaat	aagcaccagt	660
ccaccaaggg	acggagacta	ttttctgagc	agtgaactcg	gctggaaaac	taacttttta	720
aaagaggggc	aacgccatct	agtgaacaaa	ctatacactg	atgacatgcg	accagaaacg	780
gttgaiaacc	cagtaaccac	tgccgaggaa	actaagcgcg	gcagaattca	aactaagaaa	840
gaggtctcta	ttaaaaccac	acttaagag	ctgggtgcata	aaagagtaac	ctcaccagaa	900
gactggatga	tgtatgcagc	agacagttac	attgaaatga	tggctcaacc	aggtggagaa	960
aaactgctga	aaaatacgtg	agagatttgt	acactaactc	tagccagaac	aaaaacagca	1020
tttgacttga	ttttgaaaaa	agctgaaacc	agcaaaactaa	ccaacttttc	actgccggac	1080
acaagaacct	gcaagatctt	tgcttttcat	ggctggaaac	atattaaagt	ttgccatgct	1140
atttgctgtg	ttctaaacag	acaaggagcg	aaaagaaata	ctgttttatt	tcacggacca	1200
gccagtacag	gcaaatccat	tattgcacaa	gccatagcac	aggcagttgg	taatgttggt	1260
tgtataatg	cagcaaatgt	gaactttcca	ttcaatgact	gcaccaacaa	aaacctgatt	1320
tggttagaag	aagctggtta	ctttggacag	caagtaaac	aatttaaacg	catttgctct	1380
ggtcaaaact	tttcgattga	tcaaaaagga	aaaggcagca	aacagattga	accaaacacca	1440
gtcatcatga	ccacaatga	aaacattaca	gtgggtcaaaa	taggctgtga	agaaagacca	1500
gaacacactc	aaccaatcag	agatagaatg	cttaacattc	atctaacaca	tacattgcct	1560
ggtagctttg	gtttggttga	caaacatgaa	tggtccatga	tttgtgcttg	gttggttaag	1620
aatggttacc	aattaccat	ggcaagctac	tgtgctaaat	ggggcaaaag	tcctgattgg	1680
acggaaaact	ggggcgagcc	gaaggtgccg	actcctataa	attcactagg	ttcggcacgc	1740
tcaccattca	cgacacagaa	aagtacgcct	ctcagccaga	actatgcact	aactccactt	1800
gcacggatc	tcaggagacct	ggcttttagag	ccttgagcga	caccaaaatac	tcctgttgcg	1860
ggcactcgag	aaacccagaa	cactggggaa	gctgggtcca	aagcctgccca	agatgggtcaa	1920
ctgagcccaa	cttggtcaga	gatcgaggag	gatttgagag	cgtgcttcgg	tgcggaaccg	1980

-continued

ttgaagagag acttcagcga gccgctgaac ttggactaa 2019

SEQ ID NO: 24 moltype = DNA length = 2019
 FEATURE Location/Qualifiers
 source 1..2019
 mol_type = genomic DNA
 organism = Rodent protoparvovirus 1

SEQUENCE: 24

atggctggaa	atgcttactc	tgatgaggtt	ttgggagcaa	ccaactgggt	aaaggaaaaa	60
agtaaccagt	tagtattctc	atttgttttt	aaaaatgaag	atgttcaatt	gaatggaaaa	120
gatatcggat	ggaatagtta	cagaaaggag	ctgcaagagg	acgagctaaa	atctttacaa	180
cgaggagcgg	aaactacctg	ggaccagagc	gaggacatgg	aatgggaatc	ttcagtggat	240
gaactaacca	caaagcaagt	attcattttt	gactcttttg	ttaaaaagtg	tttatttgaa	300
gtgctaagta	caaagaacat	agctcctagt	gatgttaatt	ggtagtgca	gcataaatgg	360
ggaaaagacc	aaggctggca	ttgccatgta	ctaattggag	gcaaagactt	tagccaagct	420
caaggaaagt	ggtaggagaag	gcagctaaat	gtttactgga	gcagatgggt	ggtaacagcc	480
tcagtggtgc	agctatcacc	agccgaaaga	attaaagctg	gagaaatagc	ggaagaccaa	540
gaatgggtca	ctttgcttac	ttataagcat	aagcaaacca	aaaaagacta	taccaaatgt	600
gtttgctttg	gaaatatgat	tgtttactac	tttttaacca	agaagaaaat	atgcactagt	660
ccaccaaggg	acggaggcta	ttttcttagc	agtgaactcg	gctggaaaaa	taacttttta	720
aaagaaggcg	aacgccatct	agtgaagcaa	ctatacactg	atgacatgcg	gccagaaacg	780
gttgaaacca	cagtaaccac	tgcgcaggaa	actaagcgcg	gcagaattca	aactaaaaaa	840
gagggtttcta	ttaaaaccac	acttaagag	ctgggtgcata	agagagtaac	ctcaccagaa	900
gactggatga	tgatgcagcc	agacagttac	attgaaatga	tggctcagcc	aggtggagaa	960
aaactgctta	aaaatacgtg	agagatctgt	acgctaactc	tagctagaac	caaaaacagcc	1020
tttgacttga	ttttagaaaa	agctgaaacc	agcaaaactaa	ccaacttttc	actgccggac	1080
acaagaacct	gtaagatttt	tgtctttcat	ggctggaact	acattaaagt	ttgccatgct	1140
atttgctgtg	ttctaaccag	acaaggagcg	aaaagaaata	ctgttttatt	tcattggacca	1200
gccagtacag	gcaaatccat	cattgcacaa	gccatagcac	aggcagttgg	taatgttggt	1260
tgctataatg	cagcaaatgt	gaactttcca	ttcaatgact	gcaccaacaa	aaacctgatt	1320
tgggtggaag	aagctggtta	ctttggacag	caagtaaaacc	aatttaaagc	catttgctct	1380
ggtaaaacta	ttcgctttga	tcaaaaagga	aaaggcagca	agcagattga	accaaacacca	1440
gtcatcatga	ccacaaatga	gaacattaca	gtggctcagaa	taggctgcga	agagagacca	1500
gaacacactc	aaccaattag	agacagaatg	ctcaacattc	atctaacaca	tacattgcct	1560
ggtgactttg	gtttggttga	caaaaatgaa	tggcccatga	tttgtgcttg	gttggtaaag	1620
aatggttacc	aatctacat	ggcaagctac	tgtgctaaat	ggggcaaaag	tcctgatttg	1680
tcagaaaact	gggcggagcg	gaaggtgccc	actcctataa	attcaatagg	ttcgggacgc	1740
tcaccattca	cgacacagaa	aagtaacgct	ctcagccaga	actatgcact	aactccactt	1800
gcacggatc	tcgaggacct	ggcttttagag	ccttgagagca	caccaaaatac	tcctgttgcg	1860
ggcactcgag	aaacccagaa	cactggggaa	gctggttcca	aagcctgccca	agatggtcaa	1920
ctgagcccaa	cttggtcaga	gatcgaggag	gatttgagag	cgtgcttcgg	tgcggaaaccg	1980
ttgaagaaag	acttcagcga	gccgctgaac	ttggactaa			2019

SEQ ID NO: 25 moltype = DNA length = 2019
 FEATURE Location/Qualifiers
 misc_feature 1..2019
 note = strain M sequence
 source 1..2019
 mol_type = genomic DNA
 organism = Rodent protoparvovirus 1

SEQUENCE: 25

atggctggaa	atgcttactc	tgatgaagtt	ttgggaacaa	ccaactgggt	aaaagagaaa	60
agtaaccagg	aagtggtctc	atttgttttt	aaaaatgaag	atgttcagct	caatggaaaa	120
gatatcggat	ggaatagtta	caaaaaggag	ctgcaagagg	aagagctgaa	atctttacaa	180
cgaggagcgg	aaactacctg	ggaccagagc	gaggacatgg	aatgggaatc	ttcagtggat	240
gaactaacca	aaaagcaagt	attcattttt	gactcttttg	ttaaaaaatg	tttggttgaa	300
gtgctgagta	caaagaacat	agctcctagt	gatgttactt	ggtttgtaca	gcataaatgg	360
ggaaaagacc	aaagctggca	ctgtcatgta	ctaattggag	gcaaggaact	tagccaagct	420
caaggaaaaa	ggtaggagaag	gcagttaaat	gtttactgga	gcagatgggt	ggtaacagcc	480
tgtagtgtgc	agctatcacc	agctgaaaga	attaaagctg	gagagatagc	ggaagaccaa	540
gaatgggtca	ctttgcttac	ttataagcat	aagcaaacca	aaaaagacta	taccaagtgt	600
gtttgctttg	gaaatatagt	tgtttactac	tttttatcca	agaagaaaaa	atgcaccagt	660
ccaccaaggg	acggaggcta	ttttcttagc	agtgaactcg	gctggaaaaa	taacttttta	720
aaagaaggcg	aacgccatct	agtgaagcaa	ctatacactg	atgacatgcg	gccagaaacg	780
gttgaaacca	caataaccac	tgcgcaggaa	actaagcgcg	gcagaattca	aactaaaaaa	840
gagggtctcta	ttaaaaccac	acttaagag	ctgggtacata	agagagtaac	ctcaccagaa	900
gactggatga	tgatgcagcc	agacagttac	attgaaatga	tggctcaacc	aggtggagaa	960
aaactgctta	aaaatacgtg	agaaatctgt	acgctaactc	tagctagaac	caaaaacagca	1020
tttgacttga	ttttagaaaa	agctgaaacc	agcaaaactaa	ccaacttttc	actgccggac	1080
acaagaacct	gcaagatctt	tgtctttcat	ggctggaact	atgttaaagt	ttgccatgct	1140
atttgctgtg	ttctaaccag	acaaggagcg	aaaagaaata	ctgttttatt	tcattggacca	1200
gccagtacag	gcaaatccat	tattgcacaa	gccatagcac	aggcagttgg	taatgttggt	1260
tgctataatg	cagcaaatgt	gaactttcca	tttaatgact	gcaccaacaa	aaacctgatt	1320
tgggtggaag	aagctggtta	ctttggacag	caagtaaaacc	agtttaaagc	catttgctct	1380
ggtaaaacta	ttcgctttga	tcaaaaagga	aaaggcagca	agcagattga	accaaacacca	1440
gtcatcatga	ccacaaatga	aaacattaca	gtggctcagaa	taggctgcga	ggagagacca	1500
gagcacactc	aaccaatcag	agacagaatg	ctcaacattc	atctgacaca	tacattgcct	1560
ggtgactttg	gtttggttga	caagaatgaa	tggcccatga	tttgtgcttg	gttggtaaag	1620
aatggttacc	aatctacat	ggcaagctac	tgtgctaaat	ggggcaaaag	acctgatttg	1680

-continued

tcagaaaact	gggcagagcc	gaaggtagcc	actcctataa	attcactagg	ttcagcacgc	1740
tcaccattca	cgacaccgaa	aagtacgcct	ctcagccaga	actatgcact	aactccactt	1800
gcacgagatc	tcgaggacct	ggctttagag	ccttgagaga	caccaaatac	tcctgttgcc	1860
ggcactgcag	aaacccagaa	cactggggaa	gctgggtcca	aagcctgcca	agatggtcaa	1920
ctgagcccaa	cttggtcaga	gacgagggag	gatttgagag	cggtcctcgg	tgcggaaccg	1980
ttgaagaaag	acttcagcga	gccgctgaac	ttggactaa			2019

SEQ ID NO: 26 moltype = DNA length = 1966
 FEATURE Location/Qualifiers
 source 1..1966
 mol_type = genomic DNA
 organism = Rodent protoparvovirus 1

SEQUENCE: 26

atggctggaa	acgcgtactc	tgatgaagtt	ttgggaacaa	ctaactgggt	gaaggataag	60
agcaaccagg	aagtattctc	atttgttttt	aaaaatgagg	atgttcagct	caatggaaaa	120
aatatcggtg	ggaacagtta	cagaaaggag	ctgcaagagg	aggagctgaa	atctttacaa	180
cgaggagctg	aaactacctg	ggaccagagc	gaggacatgg	aatgggaatc	ttcagtggat	240
gaactgacca	aaaagcaagt	attcattttt	gactctttag	ttaaaaagtg	tcctcttgaa	300
gtactgagca	caaagaacat	agctcctagt	gatgttactt	ggtttgtaca	gcatgaatgg	360
ggaaaagacc	aaggctggca	ctgtcatgtg	ctcattggag	gcaagaactt	tagccagcct	420
caaggaaaat	gggtggaggag	acaattaaat	gtttactgga	gtagatgggt	ggtaacagcc	480
tgtagcgtgc	agctatcacc	agctgaaaga	attaaactaa	gagaaatagc	agaagaccaa	540
gaatgggtta	ctctgcttac	ttataagcat	aagcaaacca	aaaaagacta	tactaagtgt	600
gtttgtcttg	gaaatatggt	tgcttactac	tttttaacca	aaaagaaaaa	atgtaccagt	660
ccaccaaggg	acggaggcta	ttttctcagt	agtgaactcg	gctggaaaaa	taactttttg	720
aaagaaggcg	aacgccatct	agtgaacaaa	ctataactcg	atgacatgag	gccagaaacg	780
gttgagacca	cagtaaccac	agcgcaggaa	actaagcgcg	gcagaattca	aactaagaag	840
gaagtctcta	tttaagactac	acttaagag	ctgggtacata	agagagtaac	ctcaccagaa	900
gactggatga	tgatgcagcc	agacagttac	attgaaatga	tggctcaacc	agggggagaa	960
aacctactta	agaatacgct	agagatctgt	acgctgactc	tagccagaac	caaaacagcc	1020
tttgacttga	ttttagaaaa	agctgaaacc	agcaaaactaa	ccaacttttt	actggctgat	1080
acaagaacct	gtagaatctt	tgcttttcat	ggctggaact	acatcaaaag	ctgtcatgct	1140
atttgtgtgt	tcttgaacag	acagggaggc	aaaagaaata	ctgttctggt	tcattggacca	1200
gccagtcacg	gcaaatcagt	tattgcacag	gccatagcac	aggcagttgg	taatgttggt	1260
tggtataacg	cagccaatgt	gaactttcca	tttaagtact	gtaccaacaa	gaacttaatc	1320
tgggtggaag	aagctggtta	ctttggacag	caagtaaaac	agtttaaaag	catttgttct	1380
ggtcagacca	gtaacatgta	caaaaaggga	aaaggcagca	aacagattga	accaacacca	1440
gtgatcctga	ccacaaatga	aaacatcaca	gtgggtcaaaa	taggggtgtga	agagagacca	1500
gaacacactc	aaccaatcag	agacagaatg	ttaaacattc	atctgacaca	tacattgcct	1560
ggtgactttg	gtttggttga	tataaacgaa	tggcctatga	tatgtgcttg	gttggttaag	1620
aacggttacc	aatcgaccat	ggcaagttac	tgtgctaaat	ggggcaaaag	tcctgattgg	1680
acagaaaact	gggaggagcc	aaaagtaacg	actgaaataa	attcggtagg	ttcaaccaac	1740
tcaccatctc	cgaaaagtac	gcctctcagc	cagaactacg	cactaaactc	gtcggatctc	1800
gaggacctgg	ctctggagcc	ttggagcaca	ccaagtactc	ctgttggtgg	cactgtcaaa	1860
accccgaaac	ctggggaaac	tggttcaaca	gcctgtcaag	aagctcaacg	gagcccaact	1920
tggtccgaga	tcgaggagga	tttgagagcg	tgcttcagtt	cggaaac		1966

SEQ ID NO: 27 moltype = DNA length = 2019
 FEATURE Location/Qualifiers
 source 1..2019
 mol_type = genomic DNA
 organism = unidentified

SEQUENCE: 27

atggctggaa	acgcttactc	cgatgaggtt	ttgggagcaa	ccaactgggt	aaaggacaaa	60
agtagccaaag	aggtgttctc	atttgttttt	aaaaatgaga	acgtccagct	aaatgggaag	120
gacatcggtt	gggaatagta	cagaaaggag	ctacaagatg	acgagctgaa	gtctctacaa	180
cgaggagcgg	aaaccacttg	ggaccaaaagc	gaggacatgg	aatgggagag	cgcagtggtg	240
gacatgacca	aaaagcaagt	attcattttt	gattcttttg	ttaagaaatg	tttgtttgaa	300
gtgctcagca	caaagaacat	aactcctagt	gatgttactt	gggtcgtgca	gcatgaatgg	360
ggaaaggacc	aaggctggca	ctgtcatgtg	ctaattggag	gcaaaagact	tagtcaagct	420
caaggaaaaat	ggtggagaag	gcagctaaat	gtgtactgga	gtagatgggt	ggtgactgcc	480
tgtaatgttc	aactaacacc	agctgaaaga	ataaaactga	gagaaatagc	agaggacagt	540
gaatgggtca	ctttgtctac	ctataagcat	aagcacacca	agaaggacta	taccaagtgt	600
gttctttttg	gaaacatgat	tgcttattac	tttttaagca	aaaagaaaaa	atgtaccagt	660
ccaccaaggg	acggaggcta	ttttcttagc	agtgaactcg	gctggaaaaa	taactttttg	720
aaagaaggcg	agcgccatct	agtgaacaa	ctgtatactg	atgagatgaa	accagaaacg	780
gttgagacca	cagtgaaccac	agcacaggaa	gctaagcgcg	gcagaattca	aactagaaag	840
gaaggtctcta	ttataaacac	acttaagag	ttgggtacata	aaagagtaac	ctcaccagaa	900
gactggatga	tgatgcagcc	agacagttac	attgaaatga	tggctcaacc	aggtggagaa	960
aacttgctta	aaaatacact	agagatctgt	acactgactc	tagcaagaac	caaaacagca	1020
tttgacttga	ttctggaaaa	agctgaaacc	agcaaaactag	ccaacttttc	catggctaac	1080
accagaacct	gtagaatctt	tgctgaacat	ggctggaact	atattaaagt	ctgtcatgcc	1140
atctgtgtgt	tgctaaatag	acaaggaggc	aaaaggaaac	ctgtgctctt	tcattggacca	1200
gccagcagag	gcaaatctat	tattgcacaa	gccatagcac	aagcagttgg	taatgttggt	1260
tggtataatg	tgtccaatgt	gaactttcca	tttaagtact	gcaccaacaa	aaacttgatt	1320
tgggtggaag	aagctggtta	ctttggccag	caagtaaaac	aattcaaaag	tatttgttct	1380
ggccaaacca	tacgcattga	tcaaaaagga	aaaggcagca	aacagattga	accaacacca	1440
gttatcctga	ccaccaacga	gaacattacc	gtgggtcagaa	taggctgtga	ggaaagacca	1500
gaacacactc	aaccaatcag	agacagaatg	ctcaacattc	acctgacacg	tacactgcct	1560

-continued

```

ggtgactttg gtctggttga taagcacgaa tggcctctga tctgtgcttg gttggtgaag 1620
aatggttacc aatctaccat ggcttgtttc tgtgctaaat ggggcaaagt tctgatttgg 1680
tcagaagact gggcgagacc gaagctagag actcctataa attcactagg ttcaatgcgc 1740
tcaccatctc tgactccgag aagtacgcct ctcagccaga actacgctct tactccactt 1800
gcatcggacc ttgcggaact agctctagag ccttggagca caccaaatac tctgtttgtg 1860
ggcactgcag caagccagaa cactggggag gctggtttca cagcctgcc aagtgctcaa 1920
cggagcccaa cctggtccga gatcgaggcg gatctgagag cgtgcttcag ccaggaacag 1980
ctggagaaag acttcagcga ttcactgacc ttggactaa 2019

```

```

SEQ ID NO: 28      moltype = DNA length = 2019
FEATURE           Location/Qualifiers
source            1..2019
                  mol_type = genomic DNA
                  organism = Rodent protoparvovirus 1

```

```

SEQUENCE: 28
atggctggaa acgcttactc cgatgaggtt ttgggagcaa ccaactggct aaaggacaaa 60
agtagccagg aggtgtcttc atttgttttt aaaaatgaga acgtccaact aaatgggaag 120
gacatcgggt ggaatagtta cagaaaaggag ctacaagatg acgagctgaa gtctctacaa 180
cgagggcgcg agaccacttg ggaccaaagc gaggacatgg aatgggagag cgcagtggat 240
gacatgacca aaaagaagt attcattttt gattcttttg ttaagaagtg tttgtttgaa 300
gtgctcagca caaagaacat agctcctagt gatgttactt ggttcgtgca gcatgaatgg 360
ggaaaggacc aaggctggca ctgtcatgtg ctgattggag gcaaggactt tagtcaagct 420
caaggaaaat ggtggagaag gcagctaaat gtgtactgga gtagatgggt ggtgactgcc 480
tgtaatgttc aactaacacc agctgaaaga attaaactga gagaatagc agaagacagt 540
gaatgggtca ctttgcttac ctataagcat aagcacacca agaaggacta taccagtggt 600
gttctttttg gaaacatgat tgcttattac tttctaagca aaaagaaaat atgtaccagt 660
ccaccaaggg acggaggcta tttcttagc agtgactctg gctggaaaac taactttttg 720
aaagagggcg agcgccatct agtgagcaaa ctataactg atgagatgaa accagaaacg 780
gtcagagacca cagtgaaccac tgcgcaggaa gctaagcgcg gcagaattca aactagaaaag 840
gaggtctcga ttaaaaccac actcaaagag ttggtgcata aaagagtaac ctaccagaaa 900
gactggatga tgatgcagcc agacagttac attgaaatga tggctcaacc aggtggagaa 960
aacttgctta aaaatacact agagatctgt acactgactc tagcaagaac caaaacagcc 1020
tttgacttga ttctggaaaa agctgaaacc agcaaaactag ccaacttttc catggctagc 1080
accagaacct gtagaactct tgcctgagcat ggctggaact atattaaagt ctgccatgcc 1140
atctgttgtg tgctaaatag acaaggagcg aaaaaggaaca ctgtgctctt tcacggacca 1200
gccagcacag gcaaatctat cattgcacaa gccatagcac aaggagttgg taatgttgg 1260
tgttataatg ctgccaatgt gaactttcca tttaatgact gtaccaacaa aaacttgatt 1320
tgggtggaag aagctggtta ctttggccag caagtaaaacc aattcaaagc tatttgttct 1380
ggccaaacca tacgcattga tcaaaaagga aaaggcagca aacagattga accaacacca 1440
gttatcatga ccaccaacga gaacattacc gtggtcagaa taggctgtga ggaaagacca 1500
gaacacactc aaccaatcag agacagaatg ctcaacattc acctgacacg tacactgcct 1560
ggtgactttg gtctggttga taagcacgaa tggcctctga tctgtgcttg gttggtgaag 1620
aatggttacc aatctaccat ggcttgttac tgtgctaaat ggggcaaagt tctgatttgg 1680
tcagaggact gggcgagacc gaagctagag actcctataa tttcgtagg ttcaatgcgc 1740
tcaccatctc tgactccgag aagtacgcct ctcagccaga actacgctct tactccactt 1800
gcatcggacc ttgcggaact agctctagag ccttggagca caccaaatac tctgtttgcg 1860
ggcactgcag caagccagaa cactggggag gctggtttca cagcctgcc aagtgctcaa 1920
cggagcccaa cctggtccga gatcgaggcg gatctgagag cgtgcttcag ccaggaacag 1980
ctggagaaag acttcagcga ttcactgaca ttggactaa 2019

```

```

SEQ ID NO: 29      moltype = DNA length = 2024
FEATURE           Location/Qualifiers
source            1..2024
                  mol_type = genomic DNA
                  organism = unidentified

```

```

SEQUENCE: 29
atggctggaa acgcttactc cgatgaagtt ttgggagcaa ccaactggct aaaggacaaa 60
agtagccagg aagtgtcttc atttgttttt aaaaatgaga acgtccaact aaatgggaag 120
gacatcgggt ggaatagtta cagaaaaggag ctacaagatg acgagctgaa gtctctacaa 180
cgagggcgcg agaccacttg ggaccaaagc gaggacatgg aatgggagag cgcagtggat 240
gacatgacca aaaagaagt attcattttt gattcttttg ttaagaagtg tctgtttgaa 300
gtgctcagca caaagaacat agctcctagt gatgttactt ggttcgtgca gcatgaatgg 360
ggaaaggacc aaggctggca ctgtcatgtg ctgattggag gcaaggactt tagtcaagct 420
caaggaaaat ggtggagaag gcagctaaat gtgtactgga gtagatgggt agtgactgcc 480
tgtaatgttc aactaacacc agctgaaaga attaaactga gagaatagc agaggacagt 540
gaatgggtca ctttgcttac ctataagcat aagcacacca agaaggacta taccagtggt 600
gttctttttg gaaacatgat tgcttattac tttctaagca aaaagaaaat atgtaccagt 660
ccaccaaggg acggaggcta tttcttagc agtgactctg gctggaaaac taactttttg 720
aaagagggcg agcgccatct agtgagcaaa ctgtatactg atgagatgaa accagaaacg 780
gtcagagacca cagtgaaccac tgcgcaggaa gctaagcgcg gcagaattca aactagaaaag 840
gaggtctcga ttaaaaccac actcaaagag ttggtacata aaagagtaac ctaccagaaa 900
gactggatga tgatgcagcc agacagttac attgaaatga tggctcaacc aggtggagaa 960
aacttgctta aaaatacact agagatttgt acactgactc tagcaagaac caaaacagcc 1020
tttgacttga ttctggaaaa agctgaaacc agcaaaactag ccaacttttc catggctagc 1080
accagaacct gtagaactct tgcctgagcat ggctggaact atattaaagt ctgccatgcc 1140
atctgttgtg tactaaatag acaaggagcg aaaaaggaaca ctgtgctctt tcacggacca 1200
gccagcacag gcaaatctat cattgcacaa gccatagcac aaggagttgg taatgttgg 1260
tgttacaatg ctgccaatgt gaactttcca tttaatgact gtaccaacaa aaacttgatt 1320
tgggtggaag aagctggtta ctttggccag caagtaaaacc aattcaaagc tatttgttct 1380

```


-continued

ggccaaacca	tacgcatgga	tcaaaaagga	aaaggcagca	aacagattga	accaacacca	1440
gttatcatga	ccaccaacga	gaacattaca	gtggtcagaa	taggctgtga	ggaaagacca	1500
gaacacactc	aaccaatcag	agacagaatg	ctcaacattc	acctgacacg	tacactgcct	1560
ggtgactttg	gtctggtgga	taagcacgaa	tggcctctaa	tctgtgcttg	gttgggtgaag	1620
aatgggttacc	aatctaccat	ggcttggttac	tgtgctaaat	ggggcacaag	tctgtattgg	1680
tcagaggact	gggaggagcc	gaagctagag	actcctataa	attcgctagg	ttcaatgcgc	1740
tcaccatctc	tgactccgag	aagtacgcct	ctcagccaga	actacgctct	tactccactt	1800
gcatcggacc	ttgaggacct	agccctagag	ccttggagca	caccaaatac	tctgtttgcy	1860
ggcactgcag	caagccagaa	cactggggag	gctggtttca	cagcctgtca	aggtgctcaa	1920
cggagcccaa	cctgggtccga	gatcgaggcg	gatctgagag	cgtgcttcag	ccaggaacag	1980
ctggagaaaag	acttcagcga	ttcactgacc	ttggactaag	gtac		2024

SEQ ID NO: 30 moltype = DNA length = 2019
 FEATURE Location/Qualifiers
 source 1..2019
 mol_type = genomic DNA
 organism = unidentified

SEQUENCE: 30

atggctggaa	acgcttactc	cgatgaggtt	ttgggagcaa	ccaactggct	aaaggacaaa	60
agtagccagg	agggtgtctc	atttgttttt	aaaaatgaga	acgtccaact	aaatgggaag	120
gacatcggtt	ggaatagtta	cagaaaggag	ctacaagatg	acgagctgaa	gtctctgcaa	180
cgaggggcgg	agaccacttg	ggaccaaaagc	gaggacatgg	aatgggagag	cgcagtggat	240
gacatgacca	aaaagcaagt	attcattttt	gattcttttg	ttaagaagtg	tctgtttgaa	300
gtgctcagca	caaagaacat	agctcctagt	gatgttactt	ggttttgtgca	gcatgaatgg	360
ggaaaaagacc	aaggctggca	ctgtcatgtg	ctgattggag	gcaaggactt	tagtcaagct	420
caaggaaaaat	gggtggagaag	gcagctaaat	gtgtactgga	gtagatgggt	ggtgactgcc	480
tgtaatgttc	aactaacacc	agctgaaaga	attaaactga	gagaataagc	agaggacagt	540
gaatgggtga	ctttgcttct	ctataagcat	aagcacacca	agaaggacta	taccaagtgt	600
gttctttttg	gaaacatgat	tgcttattac	tttctaagca	aaaagaaaaa	atgtaccagt	660
ccaccaagggg	acggaggcta	ttttcttagc	agtgaactctg	gctggaaaac	taactttttg	720
aaagagggcg	agcgccatct	agtgaacaaa	ctgtatactg	atgagatgaa	accagaaacg	780
gtcgagacca	cagtgcaccac	tgcgagggaa	gctaagcgcg	gcagaattca	aactagaaaag	840
gaggtctcga	ttaaaaccac	actcaaagag	ttggtgcata	aaagagtaac	ctcaccagaa	900
gactggatga	tgatgcagcc	agacagttac	attgaaatga	tggctcaacc	aggtggagaa	960
aacttgctta	aaaatacact	agagatctgt	acactgactc	tagcaagaac	caaaaacagcc	1020
tttgactttg	ttctggaaaa	agctgaaacc	agcaaaactag	ccaacttttt	catggctaac	1080
accagaacct	gtagaattct	tgtcgagcat	ggctggaact	atattaaagt	ctgtcatgcc	1140
atctgttgtg	tgctaaatag	acaaggaggg	aaaagggaaca	ctgtgctctt	tcacggacca	1200
gccagcacag	gcaaatctat	cattgcacaa	gccatagcac	aaggagttgg	taatgttgtt	1260
tggtataatg	ctgccaatgt	gaaactttcca	tttaatgact	gtaccaacaa	aaacttgatt	1320
tgggtggaag	aagctggtaa	ctttggccag	caagtaaaac	aattcaaaag	tattttgtct	1380
ggccaaacca	tacgcatgga	tcaaaaagga	aaaggcagca	aacagattga	accaacacca	1440
gttatcatga	ccaccaacga	gaacattacc	gtggtcaaaa	taggctgtga	ggaaagacca	1500
gaacacactc	aaccaatcag	agacagaatg	ctcaacattc	acctgacacg	tacactgcct	1560
ggtgactttg	gtctggtgga	taagcacgaa	tggcctctga	tctgtgcttg	gttggtaaaag	1620
aatgggttacc	aatctaccat	ggcttggttac	tgtgctaaat	ggggcacaag	tctgtattgg	1680
tcagaggact	gggaggagcc	gaagctagag	actcctataa	attcgctagg	ttcaatgcgc	1740
tcaccatctc	tgactccgag	aagtacgcct	ctcagccaga	actacgctct	tactccactt	1800
gcatcggatc	tcgaggacct	ggcactggaa	ccttggagca	caccaaatac	tctgtttgtg	1860
gacactgtac	aaaccccgaa	cactggggag	gctggtttca	cagcctgtca	aggtgctcaa	1920
cggagcccaa	cctgggtccga	gatcgaggcg	gatctgagag	cgtgcttcag	ccaggaacag	1980
ctggagaaaag	acttcagcga	ttcactgacc	ttggactaa			2019

SEQ ID NO: 31 moltype = DNA length = 1997
 FEATURE Location/Qualifiers
 source 1..1997
 mol_type = genomic DNA
 organism = unidentified

SEQUENCE: 31

atgaggtttt	gggagcaacc	aactggctaa	aggacaaaag	tagccaggag	gtgttctcat	60
ttgtttttta	aaatgagaac	gtccaaacta	atgggaagga	catcggttgg	aatagtataa	120
gaaaggagct	acaagatgac	gagctgaagt	ctttgcaacg	agggggcggag	accacttggg	180
accaaaagca	ggacatggaa	tgggagagcg	cagtggaatg	catgaccaa	aagcaagtat	240
tcatttttga	ttctttgggt	aaaaagtgtc	tgtttgaagt	gctcagcaca	aagaacatag	300
ctcctagtga	tgttacttgg	ttcgtgcagc	atgaatgggg	aaaggaccaa	ggctggcact	360
gtcatgtgct	gatggaggc	aaggacttta	gtcaacctca	aggaaagtgg	tgagaaggcc	420
agctaaatgt	gtactggagt	agatggttgg	tgactgcctg	taatgttcaa	ctaacaccag	480
ctgaaagaat	taactgtaga	aggacagtga	atgggtcact	ttgcttacct		540
ataagcataa	acacaccaag	aaggactata	ccaagtgtgt	tttttttgga	aacatgattg	600
cttattactt	tctaagcaaa	aagaaaatat	gtaccagtcc	accaagggac	ggaggctatt	660
ttcttagcag	tgactctggc	tggaaaacta	actttttgaa	agaggggcag	cgccatctag	720
tgagcaaaact	gtatactgat	gagatgaaac	cagaaacggt	cgagaccaca	gtgaccactg	780
cgcagggaagc	taagcgcggc	agaattcaaa	ctagaaaagga	ggctctgatt	aaaaccacac	840
tcaaaagatt	ggtgcataaa	agagtaacct	caccagaaga	ctggatgatg	atgcagccag	900
acagttacat	ggaatgatg	gctcaaccag	gtggagaaaa	cttgcttaaa	aatacactag	960
agatctgtac	actgactcta	gcaagaacca	aaacagcctt	tgacttgatt	ctagaaaaag	1020
ctgaaaccag	caaactagcc	aactttttcca	tggctaacac	cagaacctgt	agaatctttg	1080
ctgagcatgg	ctggaaactat	attaaagtct	gccatgccat	ctgttgtgtg	ctgaatagac	1140
aaggaggcaa	aaggaaact	gtgctctttc	acggaccagc	cagcacaggc	aaatctatca	1200

-continued

ttgcacaagc	catagcacaa	ggagttggta	atgttgggtg	ttacaatgct	gccaatgtga	1260
actttccatt	taatgactgt	accaacaaaa	acttgatttg	ggtggaagaa	gctggtaact	1320
ttggccagca	agtaaaccaa	ttcaaaagcta	tttgttcttg	ccaaaccata	cgcattgatc	1380
aaaaaggaaa	aggcagcaaa	cagattgaac	caacaccagt	tatcatgacc	accaacgaga	1440
acattaccgt	ggtcagaata	ggctgtgagg	aaaggccaga	acacactcaa	ccaatcagag	1500
acagaatgct	caacattcac	ctgacacgta	cactgcctgg	tgactttggt	ctgggtggata	1560
agcacgaatg	gcctctgatc	tgtgcttggg	tggggaagaa	tggttaccaa	tctaccatgg	1620
cttgttactg	tgctaaaagg	ggcaaagttc	ctgattggtc	agaggactgg	gcggagccga	1680
agctagagac	tcctataaat	tcgctagggt	caatgcgctc	accatctctg	actccgagaa	1740
gtacgcctct	cagccagaac	tacgctctta	ctccacttgc	atcggacctt	gcggacctag	1800
ctctagagcc	ttggagcaca	ccaaatactc	ctggtgcggg	cactgcagca	agccagaaca	1860
ctggggaggc	tggtttcaca	gcctcccaag	gtgctcaacg	gagcccaacc	tggtccgaga	1920
tcgagggcga	ctgagagcgc	tgtctcagcc	aggaacagct	ggagaaagac	ttcagcgatt	1980
cactgacctt	gctactaa					1997

SEQ ID NO: 32 moltype = DNA length = 2019
 FEATURE Location/Qualifiers
 source 1..2019
 mol_type = genomic DNA
 organism = Rodent protoparvovirus 1

SEQUENCE: 32

atggctggaa	acgcttactc	cgatgaggtt	ttgggagtaa	ccaactggct	gaaggacaaa	60
agtagccagg	agggtgtctc	atttgttttt	aaaaatgaaa	acgtcccaact	aaatggaaaag	120
gacatcggtt	ggaatagtta	cagaaaaggag	ctacaagatg	acgagctgaa	gtctctacaa	180
cgaggggagg	agaccacttg	ggaccaaaagc	gaggacatgg	aatgggagag	cgcagtggat	240
gacatgacca	aaaagcaagt	atttattttt	gattcttttg	ttaagaagtg	tttgtttgaa	300
gtgctcagca	caaagaacat	agctcctagt	aatgttactt	ggttcgtgca	gcatgaatgg	360
ggaaaggacc	aaggctggca	ctgtcatgtg	ctgattggag	gcaaggactt	tagtcaacct	420
caaggaaaaat	gggtggagaag	gcagctaaat	gtgtactgga	gtagatgggt	ggtgactgcc	480
tgtaatgttc	aactaacacc	agctgaaaaga	attaaactga	gagaaatagc	agaggacagt	540
gaatgggtca	ctttgctttc	ctataagcat	aagcacacca	agaaggacta	taccaagtgt	600
gttctttttg	gaaacatgat	tgcttattac	tttttaagca	aaaagaaaaat	atgtaccagt	660
ccaccaaggg	acggaggcta	ttttcttagc	agtgaactctg	gctggaaaac	taactttttg	720
aaagaggggc	agcgccatct	agtgaagcaaa	ctgtatactg	atgagatgaa	accagaaaacg	780
gtcgagacca	cagtgaccac	tgccacaggaa	gctaagcgcg	gcagaattca	aactagaaaag	840
gaggtctcga	ttaaaaccac	actcaaagag	ttggtacata	aaagagtaac	ctcaccagaa	900
gactggatga	ttagtcagcg	agacagttac	attgaaatga	tggctcaacc	aggtggagaa	960
aaactgtcta	aaaatacact	agagatctgt	acactgactc	tagcaagaac	caaaaacagcc	1020
tttgacttga	ttctggaaaa	agctgaaaacc	agcaaaactag	ccaacttttc	catggctagc	1080
accagaacct	gtagaattct	gcttgagcat	ggctggaact	atattaaagt	ctgccatgcc	1140
atctgttgtg	tgctgaatag	acaaggaggc	aaaagggaaca	ctgtgctctt	tcacggacca	1200
gccagcacag	gcaaatctat	tattgcacaa	gccatagcac	aagcagttgg	taatgttggg	1260
tggtacaatg	ctgccaatgt	gaaactttcca	tttaatgact	gtaccaacaa	aaacttgatt	1320
tgggtggaag	aagctggtta	ctttggccag	caagtaaaac	aattcaaaagc	tatttgtttct	1380
ggccaaacca	tacgcattga	tcaaaaagga	aaaggcagca	aacagattga	accaaacacca	1440
gttattatga	ccaccaacga	gaacattacc	gtggttagaa	taggtgtgta	ggaaagacca	1500
gaacacactc	aaccaatcag	agacagaatg	ctcaacattc	acctgacacg	tacactacct	1560
ggtgactttg	gtttggtgga	taagcacgaa	tggcctctga	tctgtgcttg	gttggtgaa	1620
aatgggtacc	aatctaccat	ggcttgttac	tgtgctaaat	ggggcaaaagt	tctgtattgg	1680
tcagaggact	ggggcgagcc	gaagctagac	actcctataa	attcgctagg	ttcaatgcgc	1740
tcaccatctc	tgactccgag	aagtacgcct	ctcagccaaa	actacgctct	tactccactt	1800
gcactcggacc	ttgcggagct	agctctagag	ccttgagaca	caccaaaatc	tctgtgtgcg	1860
ggcactgcag	caaggcaaaa	actgggggag	gctggttcca	cagcctgcca	aggtgtctca	1920
cggagcccaa	cctgggtccga	gatcgaggcg	gatttgagag	cttgcttcag	tcaagaacag	1980
ttggagagcg	acttcaacga	ggagctgacc	ttggactaa			2019

SEQ ID NO: 33 moltype = DNA length = 2019
 FEATURE Location/Qualifiers
 source 1..2019
 mol_type = genomic DNA
 organism = unidentified

SEQUENCE: 33

atggctggaa	acgcttactc	cgatgaagtt	ttgggagcaa	ccaactggct	aaaggacaaa	60
agtagccagg	agggtgtctc	atttgttttt	aaaaatgaga	acgtcccaact	aaatgggaag	120
gacatcggtt	ggaatagtta	cagaaaaggag	ctacaagatg	acgagctgaa	gtctctacaa	180
cgaggggagg	agaccacttg	ggaccaaaagc	gaggacatgg	aatgggagag	cgcagtggat	240
gacatgacca	aaaagcaagt	attcattttt	gattcttttg	ttaagaagtg	tttgtttgaa	300
gtgctcagca	caaagaacat	agctcctagt	gatgttactt	ggttcgtgca	gcatgaatgg	360
ggaaaggacc	aaggctggca	ctgtcatgtg	ttgattggag	gcaaggactt	tagtcaagct	420
caaggaaaaat	gggtggagaag	gcagctaaat	gtgtactgga	gtagatgggt	ggtgactgcc	480
tgtaatgttc	aactaacacc	agctgaaaaga	attaaactga	gagaaatagc	agaggacagt	540
gaatgggtca	ctttgctttc	ctataagcat	aagcacacca	agaaggacta	taccaagtgt	600
gttctttttg	gaaacatgat	tgcttattac	tttctaagca	aaaagaaaaat	atgtaccagt	660
ccaccaaggg	acggaggcta	ttttcttagc	agtgaactctg	gctggaaaac	taactttttg	720
aaagaggggc	agcgccatct	agtgaagcaaa	ctgtatactg	atgagatgaa	accagaaaacg	780
gtcgagacca	cagtgaccac	tgccacaggaa	gctaagcgcg	gcagaattca	aactagaaaag	840
gaggtctcga	ttaaaaccac	actcaaagag	ttggtacata	aaagagtaac	ctcaccagaa	900
gactggatga	tgatgcagcc	agacagttat	attgaaatga	tggctcaacc	aggtggagaa	960
aaactgtcta	aaaatacact	agagatatgt	acactgactc	tagcaagaac	caaaaacagcc	1020

-continued

tttgacttga	ttctggaaaa	agctgaaacc	agcaaaactag	ccaacttttc	tatggctaac	1080
accagaacct	gtagaatctt	tgctgagcat	ggctggaact	atattaaagt	ctgccatgcc	1140
atctgttgtg	tgctaaatag	acaaggaggc	aaaaggaaca	ctgtgctctt	tcacggacca	1200
gccagcacag	gcaaatctat	cattgcacaa	gccatagcac	aagcagttgg	taatgttgg	1260
tgttacaatg	ctgccaatgt	gaactttcca	tttaatgact	gtaccaacaa	aaacttgatt	1320
tgggtggaag	aagctggtaa	ctttggccag	caagtaaaac	aattcaaagc	tattttgtct	1380
ggccaaacca	tacgcattga	tcaaaaagga	aaaggcagca	aacagattga	accaacacca	1440
gttatcatga	ccaccaacga	gaacattacc	gtggtcagaa	taggctgtga	ggaaagacca	1500
gaacacactc	aaccaatcag	agacagaatg	ctcaacattc	acctgacacg	tacactgcct	1560
ggtgactttg	gtctgggtga	taagcacgaa	tggcctctga	tctgtgcttg	gttgggtgaag	1620
aatggttacc	aatctacaat	ggcttggtac	tgtgctaagt	ggggcaaaag	tctgtattgg	1680
tcagaagact	gggaggagcc	gaagctagag	actcctataa	attcgctagg	ttcaatgcgc	1740
tcaccatctc	tgaactcgag	aagtacgcct	ctcagccaga	actacgctct	tactccactt	1800
gcacgggacc	ttgcggacct	agctctggag	ccttgaggca	caccaaatat	tctgtttgag	1860
ggcactgcag	caagccagaa	cactggggag	gctggtttcg	cagcctgcca	aggtgtctca	1920
cggagcccaa	cctgggtccga	gatagaagca	gaacttgagag	cttgcctcag	tcaagaacag	1980
ttggagagcg	acttcaacga	ggaactgacc	ttggactaa			2019

SEQ ID NO: 34 moltype = DNA length = 2019
 FEATURE Location/Qualifiers
 source 1..2019
 mol_type = genomic DNA
 organism = Rodent protoparvovirus 1

SEQUENCE: 34

atggctggaa	acgcttactc	cgatgaggtt	ttgggagtaa	caaactggct	gaaggacaaa	60
agtagccagg	aggtgtcttc	atttgttttt	aaaaatgaaa	acgtccaact	aaatggaaaag	120
gacatcggtt	ggaatagtta	cagaaaggag	ctacaagatg	acgagctgaa	gtctctacaa	180
cgaggggcgg	agaccacttg	ggaccaaaag	gaggacatgg	aatgggagag	cgcagtggtg	240
gacatgacca	aaaagcaagt	atttattttt	gattcttttg	ttaagaagtg	tttgtttgaa	300
gtgctcagca	caaagaacat	agctcctagt	aatgttactt	ggttcgtgca	gcatgaatgg	360
ggaaaggacc	caggtctggca	ctgtcatgtg	ctgattggag	gcaaggactt	tagtcaacct	420
caaggaaaat	gggtggagaag	gcagctaaat	gtgtactgga	gtagatgggt	gggtgactgc	480
tgtaatgttc	aactaacacc	agctgaaaag	attaaactga	gagaaatagc	agaggacagt	540
gaatgggttc	cttgtctttc	ctataagcat	aagcacacca	agaaggacta	taccaagtgt	600
gttctttttg	gaaacatgat	tgtctattac	tttttaagca	aaaagaaaat	atgtaccagt	660
ccaccaaggg	acggaggcta	ttttcttagc	agtgaactct	gctggaaaac	taactttttg	720
aaagaggggc	agcggccatc	attgagcaaa	ctgtatactg	atgagatgaa	accagaaacg	780
gtcgagacca	cagtgaccac	tgcacagga	gctaagcgcg	gcagaattca	aactagagag	840
gaggtctcga	ttaaaaccac	actcaaagag	ttggtacata	aaagagtaac	ctcaccagaa	900
gactggatga	ttagtcagcc	agacagttac	attgaaatga	tggctcaacc	aggtggagaa	960
aacttgctta	aaaatacact	agagatctgt	acactgactc	tagcaagaac	caaaaacagc	1020
tttgacttga	ttctggaaaa	agctgaaaac	agcaaaactag	ccaacttttc	catggctagc	1080
accagaacct	gtagaatctt	gctgagcat	ggctggaact	atattaaagt	ctgccatgcc	1140
atctgttgtg	tgtcgaatag	acaaggaggc	aaaaggaaca	ctgtgctctt	tcacggacca	1200
gccagcacag	gcaaatctat	tattgcacaa	gccatagcac	aagcagttgg	taatgttgg	1260
tgttacaatg	ctgccaatgt	gaactttcca	tttaatgact	gtaccaacaa	aaacttgatt	1320
tgggtggaag	aagctggtaa	ctttggccag	caagtaaaac	aattcaaagc	tattttgtct	1380
ggccaaacca	tacgcattga	tcaaaaagga	aaaggcagca	aacagattga	accaacacca	1440
gttattatga	ccaccaacga	gaacattacc	gtgggttagaa	taggctgtga	ggaaagacca	1500
gaacacactc	aaccaatcag	agacagaatg	ctcaacattc	acctgacacg	tacactacct	1560
ggtgactttg	gtttgggtga	taagcacgaa	tggcctctga	tctgtgcttg	gttgggtgaag	1620
aatgggtacc	aatctacaat	ggcttggtac	tgtgctaagt	ggggcaaaag	tctgtattgg	1680
tcagaggact	gggaggagcc	gaagctagac	actcctataa	attcgctagg	ttcaatgcgc	1740
tcaccatctc	tgaactcgag	aagtacgcct	ctcagccaaa	actacgctct	tactccactt	1800
gcacgggacc	ttgcggacct	agctctagag	ccttgaggca	caccaaatat	tctgtttgag	1860
ggcactgcag	caagccaaaa	cactggggag	gctggtttca	cagcctgcca	aggtgtctca	1920
cggagcccaa	cctgggtccga	gatcgaggcg	gatttgagag	cttgcctcag	tcaagaacag	1980
ttggagagcg	acttcaacga	ggagctgacc	ttggactaa			2019

SEQ ID NO: 35 moltype = DNA length = 1997
 FEATURE Location/Qualifiers
 source 1..1997
 mol_type = genomic DNA
 organism = unidentified

SEQUENCE: 35

atgagggtttt	gggagcaacc	aactggctaa	aggacaaaag	tagccaggag	gtattctcat	60
ttgttttttaa	aaatgagaac	gtccaactaa	atgggaagga	catcggttgg	aatagttaca	120
gaaaggagct	acaagatgac	gagctgaagt	ctctacaacg	aggggaggag	accacttggg	180
accaaagcga	ggacatggaa	tgggagagcg	cagtggtatga	catgaccaaa	aagcaagtat	240
tcatttttga	ttcttttggt	aagaagtgtc	tgtttgaaat	gctcagcaca	aagaacatag	300
ctcctagtga	gtgtacttgg	ttcgtgcagc	atgaatgggg	aaaggaccaa	ggctggcact	360
gtcatgtgct	gattggaggc	aaggacttta	gtcaagctca	aggaaaatgg	tggagaaggc	420
agctaaatgt	tgaactggag	agatggttgg	tgaactgcctg	taatgttcaa	ctaaccacag	480
ctgaaagaat	gtaactaaga	gaaatagcag	aggagcagtg	atgggtcaact	ttgcttacct	540
ataagcataa	gcacaccaag	aaggactata	ccaagtgtgt	ttcttttggg	aacatgattg	600
cttattactt	tctaagcaaa	aagaaaatat	gtaccagtc	accaagggac	ggaggctatt	660
ttcttagcag	tgactctggc	tggaaaacta	actttttgaa	agagggcgag	cgccatctag	720
tgagcaaac	gtactactgt	gagatgaaac	cagaaacgg	cgagaccaca	gtgaccactg	780
cacaggaagc	taagcgcgcc	agaattcaaa	ctagaaagga	ggtctcgatt	aaaaccacac	840

-continued

tcaaagaggtt	ggtagataaa	agagtaacct	caccagaaga	ctggatgatg	atgcagccag	900
acagttatat	tgaatgatg	gctcaaccag	gtggagaaaa	cttgcttaaa	aatacactag	960
agatatgtac	actgactcta	gcaagaacca	aaacagcctt	tgacttgatt	ctggaaaaag	1020
ctgaaaccag	caaactagcc	aacttttcta	tggctagcac	cagaacctgt	agaatctttg	1080
ctgagcatgg	ctggaactat	attaaagtct	gccatgccat	ctgttggtg	ctaaatagac	1140
aaggaggcaa	aaggaaact	gtgctctttc	acggaccagc	cagcacaggc	aaatctatca	1200
ttgcacaagc	catagcacia	gcagttggta	atgttggttg	ttacaatgct	gccaatgtga	1260
actttccatt	taatgactgt	accaacaaaa	acttgatttg	ggtggaagaa	gctggtaact	1320
ttggccagca	agtaaaccaa	ttcaaaagcta	ttgttctctg	ccaaaccata	cgcatgatc	1380
aaaaaggaaa	aggcagcaaa	cagattgaac	caacaccagt	tatcatgacc	accaacgaga	1440
acattaccgt	ggtcagaata	ggctgtgagg	aaagaccaga	acacactcaa	ccaatcagag	1500
acagaatgct	caacattcac	ctgacacgta	caactgcctg	tgactttggt	ctggtggata	1560
agcagcaatg	gcctctgatc	tgtgcttggt	tgggtgaagaa	tggttaccaa	tctaccatgg	1620
ctgtttactg	tgctaaatgg	ggcaaaagttc	ctgatttggtc	agaagactgg	gcggagccga	1680
agctagagac	tcctataaat	tcgctagggt	caatgcgctc	accatctctg	actccgagaa	1740
gtacgcctct	cagccagAAC	tacgctctta	ctccacttgt	atcggaacct	gcggagcctag	1800
ctctggagcc	ttggagcaca	ccaaatactc	ctgttgccggg	cactgcagca	agccagaaaca	1860
ctggggaggc	tggtttcgca	gcctgtcaag	gtgctcaacg	gagcccaacc	tggtccgaga	1920
tagaagcaga	cttgagagct	tgcttcagtc	aagaacagtt	ggagagcgac	ttcaacgagg	1980
agctgacctt	ggactaa					1997

SEQ ID NO: 36 moltype = DNA length = 2019
 FEATURE Location/Qualifiers
 source 1..2019
 mol_type = genomic DNA
 organism = unidentified

SEQUENCE: 36

atggctgaa	acgcttactc	cgatgaggtt	ttgggagcaa	ccaactggct	aaaggacaaa	60
agtagccagg	agggtgtctc	atttgttttt	aaaaatgaga	acgtccaact	aatggggaag	120
gacatcggtt	ggaatagtta	cagaaaggag	ctacaagatg	acgagctgaa	gtctctacaa	180
cgaggggcgg	agaccacttg	ggaccaaagc	gaggacatgg	aatgggagag	cgagtggtat	240
gacatgacca	aaaagcaagt	attcattttt	gattcttttg	ttaagaagtg	tctgtttgaa	300
gtgctcagca	caaagaacat	agctcctagt	gatgttactt	ggctcgtgca	gcatgaatgg	360
ggaaaggacc	aaggctggca	ctgtcatgtg	ctgattggag	gcaaggactt	tagtcaagct	420
caaggaaaaat	ggtaggagaag	gcagctaaat	gtgtactgga	gtagatgggt	ggtagactgc	480
tgtaatgttc	aactaaacac	agctgaaaga	attaaactga	gagaaatagc	agaggacagt	540
gaatgggtca	ctttgtctac	ctataagcat	aagcacacca	agaaggacta	taccaagtgt	600
gttctttttg	gaaacatgat	tgctttattac	tttctaagca	aaaagaaaat	atgtaccagt	660
ccaccaaggg	acggaggcta	ttttcttagc	agtgactctg	gctggaaaac	taacttcttg	720
aaagagggcg	agcgccatct	agttagcaaa	ctgtatactg	atgagatgaa	accagaaacg	780
gtcgagacca	cagtgaccac	tgcgcaggaa	gctaagcgcg	gcagaattca	aactagaaag	840
gaggtctcga	ttaaaaccac	actcaaaag	ttggtgcata	aaagagtaac	ctcaccagaa	900
gactggatga	tgatgcagcc	agacagttac	attgaaatga	tggctcaacc	aggtggagaa	960
aacttgctta	aaaatacact	agagatatgt	acactgactc	tagcaagaac	caaaacagcc	1020
tttgacttga	ttctggaaaa	agctgaaacc	agcaaaactag	ccaacttttc	catggctagc	1080
accagaacct	gtagaatctt	tgttgagcat	ggctggaact	atattaaagt	ctgccatgcc	1140
atctgtgtgt	tactaaatag	acaaggaggc	aaaagggaaca	ctgtgctctt	tcacggacca	1200
gccagcacag	gcaaatctat	cattgcacaa	gccatagcac	aagcagttgg	taatgttggt	1260
tgttacaatg	ctgccaatgt	gaactttcca	tttaatgact	gtaccaacaa	aaacttgatt	1320
tgggtggaag	aagctgtgtaa	ctttggccag	caagtaaaac	aattcaaaag	tatttgttct	1380
ggccaacca	tacgcattga	tcaaaaagga	aaaggcagca	aacagattga	accaaacacca	1440
gttatcatga	ccaccaacga	aaacattacc	gtggtcagaa	taggctgtga	ggaaagacca	1500
gaacacactc	aaccaatcag	agacagaatg	ctcaacattc	acctgacacg	tacactgcct	1560
ggtgactttg	gtctgggtga	taagcacgaa	tggcctctga	tctgtgcttg	gttggtgaag	1620
aatggttacc	aatctacaat	ggcttggtac	tgtgctaaat	ggggcaaaag	tcttgatttg	1680
tcagaagact	ggggcgagcc	gaagctagag	actcctataa	attcgctagg	ttcaatgcgc	1740
tcaccatctc	tgactccgag	aagtacgcct	ctcagccaga	actacgctct	tactccactt	1800
gcacggagcc	ttcgggacct	agctctggag	ccttgaggca	caccaaatac	tcctgttgcg	1860
ggcactgtag	caagccagaa	cactggggag	gctggtttcg	cagcctgcca	aggtgctcaa	1920
cggagcccaa	cctggtccga	gatagaagca	gacttgagag	cttgcttcag	tcaagaacag	1980
ttggagagcg	acttcaacga	ggaactgacc	ttggactaa			2019

SEQ ID NO: 37 moltype = DNA length = 192
 FEATURE Location/Qualifiers
 misc_feature 1..192
 note = consensus sequence
 source 1..192
 mol_type = other DNA
 organism = Rodent protoparvovirus 1

SEQUENCE: 37

ctarrrrarga	rgtytcdatt	aaaacyacac	tyaaagaryt	rgtrcataar	agagtaacct	60
caccagarga	ctggatgatg	atgcagccag	acagttayac	tgaatgatg	gctcarccag	120
gkggagaaaa	cytrctdaar	aatacrctag	aratygtac	retractcta	gchagaacma	180
aaacagcmtt	tg					192

67

The invention claimed is:

1. A positive amplification control (PAC) plasmid comprising:

- a unique artificial plasmid-specific sequence (UAPS) comprising SEQ ID NO: 10, and
- a target amplification polynucleotide (TAP) sequence, wherein said TAP comprises all or part of a parvovirus NS-1 sequence.

2. The PAC plasmid of claim 1, wherein said UAPS does not recognize or anneal to any natural parvovirus.

3. The PAC plasmid of claim 1, wherein the parvovirus NS-1 sequence is at least 88% identical to any one of SEQ ID NOs: 12-37 or 97% identical to SEQ ID NO: 9.

4. The PAC plasmid of claim 1, wherein the parvovirus NS-1 sequence comprises SEQ ID NO: 9 or any one of SEQ ID NOs: 12-37.

5. The PAC plasmid of claim 1, wherein the parvovirus NS-1 sequence comprises SEQ ID NO: 37.

6. The PAC plasmid of claim 1, comprising a nucleic acid extraction control (NEC) nucleotide sequence, wherein said NEC comprises an M13 bacteriophage nucleotide sequence.

7. The PAC plasmid of claim 6, wherein said M13 bacteriophage nucleotide sequence is a M13K07 nucleotide sequence.

8. The PAC plasmid of claim 7, wherein said M13K07 nucleotide sequence comprises SEQ ID NO. 8.

9. The PAC plasmid of claim 1, further comprising binding sequences for one or more oligonucleotide primers.

10. The PAC plasmid of claim 9, wherein said one or more oligonucleotide primers are selected from the group con-

68

sisting of a forward oligonucleotide primer, a reverse oligonucleotide primer, and a combination thereof.

11. The PAC plasmid of claim 9, wherein said one or more oligonucleotide primers comprise rodent parvovirus oligonucleotide primers.

12. The PAC plasmid of claim 11, wherein said rodent parvovirus oligonucleotide primers are selected from the group consisting of a forward oligonucleotide primer, a reverse oligonucleotide primer, and a combination thereof.

13. The PAC plasmid of claim 1, further comprising binding sequences for one or more M13 oligonucleotide primers.

14. The PAC plasmid of claim 13, wherein said one or more M13 oligonucleotide primers are selected from the group consisting of a forward oligonucleotide primer, a reverse oligonucleotide primer, and a combination thereof.

15. The PAC plasmid of claim 1, wherein said NS-1 sequence is selected from the group consisting of minute virus of mice prototype strain (MVMp), minute virus of mouse immunosuppressive strain (MVMi), minute virus of mouse Cutter strain (MVMc); mouse parvovirus 1b (MPV-1b), mouse parvovirus 1a (MPV-1a), mouse parvovirus 1c (MPV-1c), hamster parvovirus (HaPV), Toolan's parvovirus (H-1), Kilham rat virus (KRV), rat parvovirus 1a, rat minute virus, and Umass strain of Rat virus L (RV-Umass).

16. The PAC plasmid of claim 1, comprising the sequence of SEQ ID NO: 11.

* * * * *