



US 20250282863A1

(19) **United States**

(12) **Patent Application Publication**
Raum et al.

(10) **Pub. No.: US 2025/0282863 A1**

(43) **Pub. Date: Sep. 11, 2025**

(54) **ANTIBODY CONSTRUCTS FOR MSLN AND CD3**

(71) Applicants: **AMGEN RESEARCH (MUNICH) GMBH**, Munich (DE); **AMGEN INC.**, Thousand Oaks, CA (US)

(72) Inventors: **Tobias Raum**, Munich (DE); **Jonas Anlahr**, Munich (DE); **Julie Bailis**, Thousand Oaks, CA (US); **Claudia Blümel**, Munich (DE); **Patrick Hoffmann**, Munich (DE); **Peter Kufer**, Munich (DE); **Elisabeth Nahrwold**, Munich (DE); **Doris Rau**, Munich (DE)

(21) Appl. No.: **19/022,048**

(22) Filed: **Jan. 15, 2025**

Related U.S. Application Data

(60) Continuation of application No. 18/538,279, filed on Dec. 13, 2023, which is a division of application No. 15/225,423, filed on Aug. 1, 2016, now Pat. No. 11,884,720.

(60) Provisional application No. 62/199,939, filed on Jul. 31, 2015, provisional application No. 62/290,861, filed on Feb. 3, 2016.

Publication Classification

(51) **Int. Cl.**

C07K 16/28 (2006.01)

C07K 16/30 (2006.01)

(52) **U.S. Cl.**

CPC **C07K 16/28** (2013.01); **C07K 16/2809** (2013.01); **C07K 16/30** (2013.01); **C07K 2317/31** (2013.01); **C07K 2317/33** (2013.01); **C07K 2317/56** (2013.01); **C07K 2317/565** (2013.01); **C07K 2317/622** (2013.01); **C07K 2317/73** (2013.01); **C07K 2317/92** (2013.01); **C07K 2317/94** (2013.01)

(57)

ABSTRACT

The present invention relates to a bispecific antibody construct comprising a first binding domain which binds to human MSLN on the surface of a target cell and a second binding domain which binds to human CD3 on the surface of a T cell. Moreover, the invention provides a polynucleotide encoding the antibody construct, a vector comprising said polynucleotide and a host cell transformed or transfected with said polynucleotide or vector. Furthermore, the invention provides a process for the production of the antibody construct of the invention, a medical use of said antibody construct and a kit comprising said antibody construct.

Specification includes a Sequence Listing.

Figure 1




Figure 2A




Figure 2B




Figure 3A




Figure 3B




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




FIGURE 12A




FIGURE 12B




FIGURE 13A




FIGURE 13B




FIGURE 14A




FIGURE 14B




Figure 15A




Figure 15B




Figure 16A




Figure 16B




Figure 16C




Figure 17




Figure 18





Figure 19



ANTIBODY CONSTRUCTS FOR MSLN AND CD3

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application is a Continuation of U.S. application Ser. No. 18/538,279, filed Dec. 13, 2023, which is a Divisional of U.S. application Ser. No. 15/225,423, filed Aug. 1, 2016, now U.S. Pat. No. 11,884,720, issued Jan. 30, 2024, which claims the priority benefit under 35 U.S.C. § 119(e) of U.S. Provisional Application No. 62/290,861, filed Feb. 3, 2016 and U.S. Provisional Application No. 62/199,939, filed Jul. 31, 2015, which are incorporated herein by reference in their entirety.

INCORPORATION BY REFERENCE OF MATERIAL SUBMITTED ELECTRONICALLY

[0002] This application contains, as a separate part of disclosure, a Sequence Listing in computer-readable form (Filename: 49897C_SeqUsting.xml; Size: 482,662 bytes; Created: Jan. 8, 2025) which is incorporated by reference herein in its entirety.

FIELD

[0003] The present invention relates to a bispecific antibody construct comprising a first binding domain which binds to human MSLN on the surface of a target cell and a second binding domain which binds to human CD3 on the surface of a T cell. Moreover, the invention provides a polynucleotide encoding the antibody construct, a vector comprising said polynucleotide and a host cell transformed or transfected with said polynucleotide or vector. Furthermore, the invention provides a process for the production of the antibody construct of the invention, a medical use of said antibody construct and a kit comprising said antibody construct.

INTRODUCTION

[0004] Mesothelin is a cell surface protein that was initially discovered as an antigen recognized by K1, a monoclonal antibody derived by immunization of mice with the OVCAR-3 human ovarian cancer cell line (Chang et al., Int. J. Cancer 1992). Expression cloning was used to identify the mesothelin cDNA from a HeLa library (Chang et al., PNAS 1996). Molecular analysis demonstrated that the mesothelin gene is encoded within a 69 kD precursor protein, which is cleaved by furin to give two distinct proteins: megakaryocyte potentiating factor (MPF), a 31 kD protein that is shed, and mesothelin, a 40 kD protein that associates with the cell membrane through a glycosylphosphatidylinositol linkage (GPI anchor) (Chang et al., PNAS 1996). The mesothelin protein is organized into superhelical domains, with ARM-type repeats.

[0005] Mesothelin is highly expressed in ovarian cancer as well as in other tumor types, including pancreatic cancer, mesothelioma, lung cancer, gastric cancer and triple negative breast cancer. In normal tissues, mesothelin is mainly expressed in the mesothelial cell layer of the pleural, pericardial, and peritoneal cavities. Mesothelin is also expressed on the surface epithelia of normal ovary, Fallopian tube and tonsil.

[0006] In addition to its expression on the cell surface, mesothelin is also shed into serum through the action of

ADAM17/TACE. Serum levels of shed mesothelin are elevated in patients with ovarian cancer and other cancer. MESOMARK®, an ELISA test for shed serum mesothelin, is approved by the FDA for humanitarian use and may help in the diagnosis or monitoring of mesothelioma. Shed mesothelin has also been used, alone or together with other markers, to aid in diagnosis or prognosis of other cancer types.

[0007] The correlation of serum levels of shed mesothelin with disease suggested a potential role for the mesothelin protein in cancer progression. While the biological function of mesothelin is not well understood—knockout mice appear normal—mesothelin has been shown to bind the mucin MUC16/CA-125. The mesothelin-CA-125 interaction has been proposed to function in cell adhesion, invasion and metastasis.

[0008] The first antibodies generated against mesothelin for therapeutic intervention were designed to interfere with the interaction between mesothelin and CA-125. Phage display identified the Fv SS, which was affinity optimized and used to generate a recombinant immunotoxin targeting mesothelin, SS1P. The MORAb-009 antibody amatuximab, which also uses SS1, recognizes a non-linear epitope in the amino terminal 64 amino acids of mesothelin. The SS1 Fv was also used to generate chimeric antigen receptor-engineered T cells. Anti-mesothelin antibodies have also been used to generate drug conjugates, such as Anetumab ravtansine, containing the MF-T antibody coupled to DM4, and the 7D9 antibody conjugated to monomethyl auristatin E. These anti-mesothelin antibodies also recognize the amino terminal region of the protein (amino acids 296-390) although they do not compete with. These anti-mesothelin targeted therapeutics that are currently in clinical trials have shown limited efficacy as single agents.

[0009] Very recently, new anti-mesothelin antibodies have been reported that recognize other regions of the mesothelin protein. There still remains a need to understand whether antibodies against different epitopes of mesothelin than SS1/MORAb-009 (for example, antibodies that are not competed by shed mesothelin or shed CA-125) would have improved efficacy in patients.

[0010] As there is still a need for having available further options for the treatment of solid tumor diseases related to the overexpression of MSLN, such as ovarian cancer, pancreatic cancer, mesothelioma, lung cancer, gastric cancer and triple negative breast cancer, there are provided herewith means and methods for the solution of this problem in the form of a bispecific antibody construct having a binding domain directed to MSLN on the surface of tumor target cells and a second binding domain directed to CD3 on the surface of T cells.

[0011] Thus, in a first aspect, the present invention provides a bispecific antibody construct comprising a first binding domain which binds to human MSLN on the surface of a target cell and a second binding domain which binds to human CD3 on the surface of a T cell, wherein the first binding domain binds to an epitope of MSLN which is comprised within the MSLN variants as depicted in SEQ ID NOs: 231, 232, and 233 and further binds to *Macaca fascicularis* MSLN as depicted in SEQ ID NOs: 234.

[0012] It must be noted that as used herein, the singular forms “a”, “an”, and “the” include plural references unless the context clearly indicates otherwise. Thus, for example, reference to “a reagent” includes one or more of such

different reagents and reference to “the method” includes reference to equivalent steps and methods known to those of ordinary skill in the art that could be modified or substituted for the methods described herein.

[0013] Unless otherwise indicated, the term “at least” preceding a series of elements is to be understood to refer to every element in the series. Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the present invention.

[0014] The term “and/or” wherever used herein includes the meaning of “and”, “or” and “all or any other combination of the elements connected by said term”.

[0015] The term “about” or “approximately” as used herein means within $\pm 20\%$, preferably within $\pm 15\%$, more preferably within $\pm 10\%$, and most preferably within $\pm 5\%$ of a given value or range.

[0016] Throughout this specification and the claims which follow, unless the context requires otherwise, the word “comprise”, and variations such as “comprises” and “comprising”, will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integer or step. When used herein the term “comprising” can be substituted with the term “containing” or “including” or sometimes when used herein with the term “having”.

[0017] When used herein “consisting of” excludes any element, step, or ingredient not specified in the claim element. When used herein, “consisting essentially of” does not exclude materials or steps that do not materially affect the basic and novel characteristics of the claim.

[0018] In each instance herein any of the terms “comprising”, “consisting essentially of” and “consisting of” may be replaced with either of the other two terms.

[0019] The term “antibody construct” refers to a molecule in which the structure and/or function is/are based on the structure and/or function of an antibody, e.g., of a full-length or whole immunoglobulin molecule. An antibody construct is hence capable of binding to its specific target or antigen. Furthermore, an antibody construct according to the invention comprises the minimum structural requirements of an antibody which allow for the target binding. This minimum requirement may e.g. be defined by the presence of at least the three light chain CDRs (i.e. CDR1, CDR2 and CDR3 of the VL region) and/or the three heavy chain CDRs (i.e. CDR1, CDR2 and CDR3 of the VH region), preferably of all six CDRs. The antibodies on which the constructs according to the invention are based include for example monoclonal, recombinant, chimeric, deimmunized, humanized and human antibodies.

[0020] Within the definition of “antibody constructs” according to the invention are full-length or whole antibodies also including camelid antibodies and other immunoglobulin antibodies generated by biotechnological or protein engineering methods or processes. These full-length antibodies may be for example monoclonal, recombinant, chimeric, deimmunized, humanized and human antibodies. Also within the definition of “antibody constructs” are fragments of full-length antibodies, such as VH, VHH, VL, (s)dAb, Fv, Fd, Fab, Fab', F(ab')₂ or “r IgG” (“half antibody”). Antibody constructs according to the invention may also be modified fragments of antibodies, also called antibody variants, such as scFv, di-scFv or bi(s)-scFv, scFv-Fc,

scFv-zipper, scFab, Fab2, Fab3, diabodies, single chain diabodies, tandem diabodies (Tandab's), tandem di-scFv, tandem tri-scFv, “minibodies” exemplified by a structure which is as follows: (VH-VL-CH3)₂, (scFv-CH3)₂, ((scFv) 2-CH3+CH3), ((scFv)₂-CH3) or (scFv-CH3-scFv)₂, multi-bodies such as triabodies or tetrabodies, and single domain antibodies such as nanobodies or single variable domain antibodies comprising merely one variable domain, which might be VHH, VH or VL, that specifically bind an antigen or epitope independently of other V regions or domains.

[0021] A binding domain may typically comprise an antibody light chain variable region (VL) and an antibody heavy chain variable region (VH); however, it does not have to comprise both. Fd fragments, for example, have two VH regions and often retain some antigen-binding function of the intact antigen-binding domain. Additional examples for the format of antibody fragments, antibody variants or binding domains include (1) a Fab fragment, a monovalent fragment having the VL, VH, CL and CH1 domains; (2) a F(ab')₂ fragment, a bivalent fragment having two Fab fragments linked by a disulfide bridge at the hinge region; (3) an Fd fragment having the two VH and CH1 domains; (4) an Fv fragment having the VL and VH domains of a single arm of an antibody, (5) a dAb fragment (Ward et al., (1989) Nature 341:544-546), which has a VH domain; (6) an isolated complementarity determining region (CDR), and (7) a single chain Fv (scFv), the latter being preferred (for example, derived from an scFV-library). Examples for embodiments of antibody constructs according to the invention are e.g. described in WO 00/006605, WO 2005/040220, WO 2008/119567, WO 2010/037838, WO 2013/026837, WO 2013/026833, US 2014/0308285, US 2014/0302037, WO 2014/144722, WO 2014/151910, and WO 2015/048272.

[0022] Furthermore, the definition of the term “antibody construct” includes monovalent, bivalent and polyvalent/multivalent constructs and, thus, monospecific constructs, specifically binding to only one antigenic structure, as well as bispecific and polyspecific/multispecific constructs, which specifically bind more than one antigenic structure, e.g. two, three or more, through distinct binding domains. Moreover, the definition of the term “antibody construct” includes molecules consisting of only one polypeptide chain as well as molecules consisting of more than one polypeptide chain, which chains can be either identical (homodimers, homotrimers or homo oligomers) or different (heterodimer, heterotrimer or heterooligomer). Examples for the above identified antibodies and variants or derivatives thereof are described inter alia in Harlow and Lane, Antibodies a laboratory manual, CSHL Press (1988) and Using Antibodies: a laboratory manual, CSHL Press (1999), Kontermann and Dübel, Antibody Engineering, Springer, 2nd ed. 2010 and Little, Recombinant Antibodies for Immunotherapy, Cambridge University Press 2009.

[0023] The antibody constructs of the present invention are preferably “in vitro generated antibody constructs”. This term refers to an antibody construct according to the above definition where all or part of the variable region (e.g., at least one CDR) is generated in a non-immune cell selection, e.g., an in vitro phage display, protein chip or any other method in which candidate sequences can be tested for their ability to bind to an antigen. This term thus preferably excludes sequences generated solely by genomic rearrangement in an immune cell in an animal. A “recombinant

antibody” is an antibody made through the use of recombinant DNA technology or genetic engineering.

[0024] The term “monoclonal antibody” (mAb) or monoclonal antibody construct as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations and/or post-translation modifications (e.g., isomerizations, amidations) that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site or determinant on the antigen, in contrast to conventional (polyclonal) antibody preparations which typically include different antibodies directed against different determinants (or epitopes). In addition to their specificity, the monoclonal antibodies are advantageous in that they are synthesized by the hybridoma culture, hence uncontaminated by other immunoglobulins. The modifier “monoclonal” indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method.

[0025] For the preparation of monoclonal antibodies, any technique providing antibodies produced by continuous cell line cultures can be used. For example, monoclonal antibodies to be used may be made by the hybridoma method first described by Koehler et al., *Nature*, 256: 495 (1975), or may be made by recombinant DNA methods (see, e.g., U.S. Pat. No. 4,816,567). Examples for further techniques to produce human monoclonal antibodies include the trioma technique, the human B-cell hybridoma technique (Kozbor, *Immunology Today* 4 (1983), 72) and the EBV-hybridoma technique (Cole et al., *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc. (1985), 77-96).

[0026] Hybridomas can then be screened using standard methods, such as enzyme-linked immunosorbent assay (ELISA) and surface plasmon resonance (BIAcore™) analysis, to identify one or more hybridomas that produce an antibody that specifically binds with a specified antigen. Any form of the relevant antigen may be used as the immunogen, e.g., recombinant antigen, naturally occurring forms, any variants or fragments thereof, as well as an antigenic peptide thereof. Surface plasmon resonance as employed in the BIAcore system can be used to increase the efficiency of phage antibodies which bind to an epitope of a target antigen, such as MSLN or CD3 epsilon (Schier, *Human Antibodies Hybridomas* 7 (1996), 97-105; Malmborg, *J. Immunol. Methods* 183 (1995), 7-13).

[0027] Another exemplary method of making monoclonal antibodies includes screening protein expression libraries, e.g., phage display or ribosome display libraries. Phage display is described, for example, in Ladner et al., U.S. Pat. No. 5,223,409; Smith (1985) *Science* 228:1315-1317; Clackson et al., *Nature*, 352: 624-628 (1991) and Marks et al., *J. Mol. Biol.*, 222: 581-597 (1991).

[0028] In addition to the use of display libraries, the relevant antigen can be used to immunize a non-human animal, e.g., a rodent (such as a mouse, hamster, rabbit or rat). In one embodiment, the non-human animal includes at least a part of a human immunoglobulin gene. For example, it is possible to engineer mouse strains deficient in mouse antibody production with large fragments of the human Ig (immunoglobulin) loci. Using the hybridoma technology, antigen-specific monoclonal antibodies derived from the

genes with the desired specificity may be produced and selected. See, e.g., XENOMOUSE™, Green et al. (1994) *Nature Genetics* 7:13-21, US 2003-0070185, WO 96/34096, and WO 96/33735.

[0029] A monoclonal antibody can also be obtained from a non-human animal, and then modified, e.g., humanized, deimmunized, rendered chimeric etc., using recombinant DNA techniques known in the art. Examples of modified antibody constructs include humanized variants of non-human antibodies, “affinity matured” antibodies (see, e.g. Hawkins et al. *J. Mol. Biol.* 254, 889-896 (1992) and Lowman et al., *Biochemistry* 30, 10832-10837 (1991)) and antibody mutants with altered effector function(s) (see, e.g., U.S. Pat. No. 5,648,260, Kontermann and Dübel (2010), loc. cit. and Little (2009), loc. cit.).

[0030] In immunology, affinity maturation is the process by which B cells produce antibodies with increased affinity for antigen during the course of an immune response. With repeated exposures to the same antigen, a host will produce antibodies of successively greater affinities. Like the natural prototype, the in vitro affinity maturation is based on the principles of mutation and selection. The in vitro affinity maturation has successfully been used to optimize antibodies, antibody constructs, and antibody fragments. Random mutations inside the CDRs are introduced using radiation, chemical mutagens or error-prone PCR. In addition, the genetical diversity can be increased by chain shuffling. Two or three rounds of mutation and selection using display methods like phage display usually results in antibody fragments with affinities in the low nanomolar range.

[0031] A preferred type of an amino acid substitutional variation of the antibody constructs involves substituting one or more hypervariable region residues of a parent antibody (e. g. a humanized or human antibody). Generally, the resulting variant(s) selected for further development will have improved biological properties relative to the parent antibody from which they are generated. A convenient way for generating such substitutional variants involves affinity maturation using phage display. Briefly, several hypervariable region sites (e. g. 6-7 sites) are mutated to generate all possible amino acid substitutions at each site. The antibody variants thus generated are displayed in a monovalent fashion from filamentous phage particles as fusions to the gene III product of M13 packaged within each particle. The phage-displayed variants are then screened for their biological activity (e. g. binding affinity) as herein disclosed. In order to identify candidate hypervariable region sites for modification, alanine scanning mutagenesis can be performed to identify hypervariable region residues contributing significantly to antigen binding. Alternatively, or additionally, it may be beneficial to analyze a crystal structure of the antigen-antibody complex to identify contact points between the binding domain and, e.g., human MSLN. Such contact residues and neighbouring residues are candidates for substitution according to the techniques elaborated herein. Once such variants are generated, the panel of variants is subjected to screening as described herein and antibodies with superior properties in one or more relevant assays may be selected for further development.

[0032] The monoclonal antibodies and antibody constructs of the present invention specifically include “chimeric” antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a par-

ticular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is/are identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Pat. No. 4,816,567; Morrison et al., Proc. Natl. Acad. Sci. USA, 81: 6851-6855 (1984)). Chimeric antibodies of interest herein include "primitized" antibodies comprising variable domain antigen-binding sequences derived from a non-human primate (e.g., Old World Monkey, Ape etc.) and human constant region sequences. A variety of approaches for making chimeric antibodies have been described. See e.g., Morrison et al., Proc. Natl. Acad. Sci. U.S.A. 81:6851, 1985; Takeda et al., Nature 314:452, 1985; Cabilly et al., U.S. Pat. No. 4,816,567; Boss et al., U.S. Pat. No. 4,816,397; Tanaguchi et al., EP 0171496; EP 0173494; and GB 2177096.

[0033] An antibody, antibody construct, antibody fragment or antibody variant may also be modified by specific deletion of human T cell epitopes (a method called "deimmunization") by the methods disclosed for example in WO 98/52976 or WO 00/34317. Briefly, the heavy and light chain variable domains of an antibody can be analyzed for peptides that bind to MHC class II; these peptides represent potential T cell epitopes (as defined in WO 98/52976 and WO 00/34317). For detection of potential T cell epitopes, a computer modeling approach termed "peptide threading" can be applied, and in addition a database of human MHC class II binding peptides can be searched for motifs present in the VH and VL sequences, as described in WO 98/52976 and WO 00/34317. These motifs bind to any of the 18 major MHC class II DR allotypes, and thus constitute potential T cell epitopes. Potential T cell epitopes detected can be eliminated by substituting small numbers of amino acid residues in the variable domains, or preferably, by single amino acid substitutions. Typically, conservative substitutions are made. Often, but not exclusively, an amino acid common to a position in human germline antibody sequences may be used. Human germline sequences are disclosed e.g. in Tomlinson, et al. (1992) J. Mol. Biol. 227:776-798; Cook, G. P. et al. (1995) Immunol. Today Vol. 16 (5): 237-242; and Tomlinson et al. (1995) EMBO J. 14: 14:4628-4638. The V BASE directory provides a comprehensive directory of human immunoglobulin variable region sequences (compiled by Tomlinson, L.A. et al. MRC Centre for Protein Engineering, Cambridge, UK). These sequences can be used as a source of human sequence, e.g., for framework regions and CDRs. Consensus human framework regions can also be used, for example as described in U.S. Pat. No. 6,300,064.

[0034] "Humanized" antibodies, antibody constructs, variants or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) are antibodies or immunoglobulins of mostly human sequences, which contain (a) minimal sequence(s) derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a hypervariable region (also CDR) of the recipient are replaced by residues from a hypervariable region of a non-human (e.g., rodent) species (donor antibody) such as mouse, rat, hamster or rabbit having the desired specificity, affinity, and capacity. In some instances, Fv framework region (FR) residues of the human immuno-

globulin are replaced by corresponding non-human residues. Furthermore, "humanized antibodies" as used herein may also comprise residues which are found neither in the recipient antibody nor the donor antibody. These modifications are made to further refine and optimize antibody performance. The humanized antibody may also comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones et al., Nature, 321: 522-525 (1986); Reichmann et al., Nature, 332: 323-329 (1988); and Presta, Curr. Op. Struct. Biol., 2: 593-596 (1992).

[0035] Humanized antibodies or fragments thereof can be generated by replacing sequences of the Fv variable domain that are not directly involved in antigen binding with equivalent sequences from human Fv variable domains. Exemplary methods for generating humanized antibodies or fragments thereof are provided by Morrison (1985) Science 229:1202-1207; by Oi et al. (1986) BioTechniques 4:214; and by U.S. Pat. Nos. 5,585,089; 5,693,761; 5,693,762; 5,859,205; and 6,407,213. Those methods include isolating, manipulating, and expressing the nucleic acid sequences that encode all or part of immunoglobulin Fv variable domains from at least one of a heavy or light chain. Such nucleic acids may be obtained from a hybridoma producing an antibody against a predetermined target, as described above, as well as from other sources. The recombinant DNA encoding the humanized antibody molecule can then be cloned into an appropriate expression vector.

[0036] Humanized antibodies may also be produced using transgenic animals such as mice that express human heavy and light chain genes, but are incapable of expressing the endogenous mouse immunoglobulin heavy and light chain genes. Winter describes an exemplary CDR grafting method that may be used to prepare the humanized antibodies described herein (U.S. Pat. No. 5,225,539). All of the CDRs of a particular human antibody may be replaced with at least a portion of a non-human CDR, or only some of the CDRs may be replaced with non-human CDRs. It is only necessary to replace the number of CDRs required for binding of the humanized antibody to a predetermined antigen.

[0037] A humanized antibody can be optimized by the introduction of conservative substitutions, consensus sequence substitutions, germline substitutions and/or back mutations. Such altered immunoglobulin molecules can be made by any of several techniques known in the art, (e.g., Teng et al., Proc. Natl. Acad. Sci. U.S.A., 80: 7308-7312, 1983; Kozbor et al., Immunology Today, 4: 7279, 1983; Olsson et al., Meth. Enzymol., 92: 3-16, 1982, and EP 239 400).

[0038] The term "human antibody", "human antibody construct" and "human binding domain" includes antibodies, antibody constructs and binding domains having antibody regions such as variable and constant regions or domains which correspond substantially to human germline immunoglobulin sequences known in the art, including, for example, those described by Kabat et al. (1991) (loc. cit.). The human antibodies, antibody constructs or binding domains of the invention may include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or site-specific mutagenesis in vitro or by somatic mutation in vivo), for example in the CDRs, and in particular, in CDR3. The human antibodies, antibody constructs or binding domains can have at least one, two, three, four, five, or more positions replaced

with an amino acid residue that is not encoded by the human germline immunoglobulin sequence. The definition of human antibodies, antibody constructs and binding domains as used herein also contemplates fully human antibodies, which include only non-artificially and/or genetically altered human sequences of antibodies as those can be derived by using technologies or systems such as the Xenomouse.

[0039] In some embodiments, the antibody constructs of the invention are “isolated” or “substantially pure” antibody constructs. “Isolated” or “substantially pure”, when used to describe the antibody constructs disclosed herein, means an antibody construct that has been identified, separated and/or recovered from a component of its production environment. Preferably, the antibody construct is free or substantially free of association with all other components from its production environment. Contaminant components of its production environment, such as that resulting from recombinant transfected cells, are materials that would typically interfere with diagnostic or therapeutic uses for the polypeptide, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. The antibody constructs may e.g. constitute at least about 5%, or at least about 50% by weight of the total protein in a given sample. It is understood that the isolated protein may constitute from 5% to 99.9% by weight of the total protein content, depending on the circumstances. The polypeptide may be made at a significantly higher concentration through the use of an inducible promoter or high expression promoter, such that it is made at increased concentration levels. The definition includes the production of an antibody construct in a wide variety of organisms and/or host cells that are known in the art. In preferred embodiments, the antibody construct will be purified (1) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (2) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Ordinarily, however, an isolated antibody construct will be prepared by at least one purification step.

[0040] The term “binding domain” characterizes in connection with the present invention a domain which (specifically) binds to/interacts with/recognizes a given target epitope or a given target site on the target molecules (antigens), here: MSLN and CD3, respectively. The structure and function of the first binding domain (recognizing MSLN), and preferably also the structure and/or function of the second binding domain (recognizing CD3), is/are based on the structure and/or function of an antibody, e.g. of a full-length or whole immunoglobulin molecule. According to the invention, the first binding domain is characterized by the presence of three light chain CDRs (i.e. CDR1, CDR2 and CDR3 of the VL region) and/or three heavy chain CDRs (i.e. CDR1, CDR2 and CDR3 of the VH region). The second binding domain preferably also comprises the minimum structural requirements of an antibody which allow for the target binding. More preferably, the second binding domain comprises at least three light chain CDRs (i.e. CDR1, CDR2 and CDR3 of the VL region) and/or three heavy chain CDRs (i.e. CDR1, CDR2 and CDR3 of the VH region). It is envisaged that the first and/or second binding domain is produced by or obtainable by phage-display or library screening methods rather than by grafting CDR sequences from a pre-existing (monoclonal) antibody into a scaffold.

[0041] According to the present invention, binding domains are in the form of one or more polypeptides. Such polypeptides may include proteinaceous parts and non-proteinaceous parts (e.g. chemical linkers or chemical cross-linking agents such as glutaraldehyde). Proteins (including fragments thereof, preferably biologically active fragments, and peptides, usually having less than 30 amino acids) comprise two or more amino acids coupled to each other via a covalent peptide bond (resulting in a chain of amino acids). The term “polypeptide” as used herein describes a group of molecules, which usually consist of more than 30 amino acids. Polypeptides may further form multimers such as dimers, trimers and higher oligomers, i.e., consisting of more than one polypeptide molecule. Polypeptide molecules forming such dimers, trimers etc. may be identical or non-identical. The corresponding higher order structures of such multimers are, consequently, termed homo- or heterodimers, homo- or heterotrimers etc. An example for a heteromultimer is an antibody molecule, which, in its naturally occurring form, consists of two identical light polypeptide chains and two identical heavy polypeptide chains. The terms “peptide”, “polypeptide” and “protein” also refer to naturally modified peptides/polypeptides/proteins wherein the modification is effected e.g. by post-translational modifications like glycosylation, acetylation, phosphorylation and the like. A “peptide”, “polypeptide” or “protein” when referred to herein may also be chemically modified such as pegylated. Such modifications are well known in the art and described herein below.

[0042] Preferably the binding domain which binds to MSLN and/or the binding domain which binds to CD3 is/are human binding domains. Antibodies and antibody constructs comprising at least one human binding domain avoid some of the problems associated with antibodies or antibody constructs that possess non-human such as rodent (e.g. murine, rat, hamster or rabbit) variable and/or constant regions. The presence of such rodent derived proteins can lead to the rapid clearance of the antibodies or antibody constructs or can lead to the generation of an immune response against the antibody or antibody construct by a patient. In order to avoid the use of rodent derived antibodies or antibody constructs, human or fully human antibodies/antibody constructs can be generated through the introduction of human antibody function into a rodent so that the rodent produces fully human antibodies.

[0043] The ability to clone and reconstruct megabase-sized human loci in YACs and to introduce them into the mouse germline provides a powerful approach to elucidating the functional components of very large or crudely mapped loci as well as generating useful models of human disease. Furthermore, the use of such technology for substitution of mouse loci with their human equivalents could provide unique insights into the expression and regulation of human gene products during development, their communication with other systems, and their involvement in disease induction and progression.

[0044] An important practical application of such a strategy is the “humanization” of the mouse humoral immune system. Introduction of human immunoglobulin (Ig) loci into mice in which the endogenous Ig genes have been inactivated offers the opportunity to study the mechanisms underlying programmed expression and assembly of antibodies as well as their role in B-cell development. Furthermore, such a strategy could provide an ideal source for

production of fully human monoclonal antibodies (mAbs)—an important milestone towards fulfilling the promise of antibody therapy in human disease. Fully human antibodies or antibody constructs are expected to minimize the immunogenic and allergic responses intrinsic to mouse or mouse-derivatized mAbs and thus to increase the efficacy and safety of the administered antibodies/antibody constructs. The use of fully human antibodies or antibody constructs can be expected to provide a substantial advantage in the treatment of chronic and recurring human diseases, such as inflammation, autoimmunity, and cancer, which require repeated compound administrations.

[0045] One approach towards this goal was to engineer mouse strains deficient in mouse antibody production with large fragments of the human Ig loci in anticipation that such mice would produce a large repertoire of human antibodies in the absence of mouse antibodies. Large human Ig fragments would preserve the large variable gene diversity as well as the proper regulation of antibody production and expression. By exploiting the mouse machinery for antibody diversification and selection and the lack of immunological tolerance to human proteins, the reproduced human antibody repertoire in these mouse strains should yield high affinity antibodies against any antigen of interest, including human antigens. Using the hybridoma technology, antigen-specific human mAbs with the desired specificity could be readily produced and selected. This general strategy was demonstrated in connection with the generation of the first XenoMouse mouse strains (see Green et al. *Nature Genetics* 7:13-21 (1994)). The XenoMouse strains were engineered with yeast artificial chromosomes (YACs) containing 245 kb and 190 kb-sized germline configuration fragments of the human heavy chain locus and kappa light chain locus, respectively, which contained core variable and constant region sequences. The human Ig containing YACs proved to be compatible with the mouse system for both rearrangement and expression of antibodies and were capable of substituting for the inactivated mouse Ig genes. This was demonstrated by their ability to induce B cell development, to produce an adult-like human repertoire of fully human antibodies, and to generate antigen-specific human mAbs. These results also suggested that introduction of larger portions of the human Ig loci containing greater numbers of V genes, additional regulatory elements, and human Ig constant regions might recapitulate substantially the full repertoire that is characteristic of the human humoral response to infection and immunization. The work of Green et al. was recently extended to the introduction of greater than approximately 80% of the human antibody repertoire through introduction of megabase sized, germline configuration YAC fragments of the human heavy chain loci and kappa light chain loci, respectively. See Mendez et al. *Nature Genetics* 15:146-156 (1997) and U.S. patent application Ser. No. 08/759,620.

[0046] The production of the XenoMouse mice is further discussed and delineated in U.S. patent application Ser. No. 07/466,008, Ser. No. 07/610,515, Ser. No. 07/919,297, Ser. No. 07/922,649, Ser. No. 08/031,801, Ser. No. 08/112,848, Ser. No. 08/234,145, Ser. No. 08/376,279, Ser. No. 08/430,938, Ser. No. 08/464,584, Ser. No. 08/464,582, Ser. No. 08/463,191, Ser. No. 08/462,837, Ser. No. 08/486,853, Ser. No. 08/486,857, Ser. No. 08/486,859, Ser. No. 08/462,513, Ser. No. 08/724,752, and Ser. No. 08/759,620; and U.S. Pat. Nos. 6,162,963; 6,150,584; 6,114,598; 6,075,181, and

5,939,598 and Japanese Patent Nos. 3 068 180 B2, 3 068 506 B2, and 3 068 507 B2. See also Mendez et al. *Nature Genetics* 15:146-156 (1997) and Green and Jakobovits J. *Exp. Med.* 188:483-495 (1998), EP 0 463 151 B1, WO 94/02602, WO 96/34096, WO 98/24893, WO 00/76310, and WO 03/47336.

[0047] In an alternative approach, others, including GenPharm International, Inc., have utilized a “minilocus” approach. In the minilocus approach, an exogenous Ig locus is mimicked through the inclusion of pieces (individual genes) from the Ig locus. Thus, one or more VH genes, one or more DH genes, one or more JH genes, a mu constant region, and a second constant region (preferably a gamma constant region) are formed into a construct for insertion into an animal. This approach is described in U.S. Pat. No. 5,545,807 to Surani et al. and U.S. Pat. Nos. 5,545,806; 5,625,825; 5,625,126; 5,633,425; 5,661,016; 5,770,429; 5,789,650; 5,814,318; 5,877,397; 5,874,299; and 6,255,458 each to Lonberg and Kay, U.S. Pat. Nos. 5,591,669 and 6,023,010 to Krimpenfort and Berns, U.S. Pat. Nos. 5,612,205; 5,721,367; and U.S. Pat. No. 5,789,215 to Berns et al., and U.S. Pat. No. 5,643,763 to Choi and Dunn, and GenPharm International U.S. patent application Ser. No. 07/574,748, Ser. No. 07/575,962, Ser. No. 07/810,279, Ser. No. 07/853,408, Ser. No. 07/904,068, Ser. No. 07/990,860, Ser. No. 08/053,131, Ser. No. 08/096,762, Ser. No. 08/155,301, Ser. No. 08/161,739, Ser. No. 08/165,699, Ser. No. 08/209,741. See also EP 0 546 073 B1, WO 92/03918, WO 92/22645, WO 92/22647, WO 92/22670, WO 93/12227, WO 94/00569, WO 94/25585, WO 96/14436, WO 97/13852, and WO 98/24884 and U.S. Pat. No. 5,981,175. See further Taylor et al. (1992), Chen et al. (1993), Tuailon et al. (1993), Choi et al. (1993), Lonberg et al. (1994), Taylor et al. (1994), and Tuailon et al. (1995), Fishwild et al. (1996).

[0048] Kirin has also demonstrated the generation of human antibodies from mice in which, through microcell fusion, large pieces of chromosomes, or entire chromosomes, have been introduced. See European Patent Application Nos. 773 288 and 843 961. Xenerex Biosciences is developing a technology for the potential generation of human antibodies. In this technology, SCID mice are reconstituted with human lymphatic cells, e.g., B and/or T cells. Mice are then immunized with an antigen and can generate an immune response against the antigen. See U.S. Pat. Nos. 5,476,996; 5,698,767; and 5,958,765.

[0049] Human anti-mouse antibody (HAMA) responses have led the industry to prepare chimeric or otherwise humanized antibodies. It is however expected that certain human anti-chimeric antibody (HACA) responses will be observed, particularly in chronic or multi-dose utilizations of the antibody. Thus, it would be desirable to provide antibody constructs comprising a human binding domain against MSLN and a human binding domain against CD3 in order to vitiate concerns and/or effects of HAMA or HACA response.

[0050] The terms “(specifically) binds to”, “(specifically) recognizes”, “is (specifically) directed to”, and “(specifically) reacts with” mean in accordance with this invention that a binding domain interacts or specifically interacts with a given epitope or a given target site on the target molecules (antigens), here: MSLN and CD3, respectively.

[0051] The term “epitope” refers to a site on an antigen to which a binding domain, such as an antibody or immuno-

globulin, or a derivative, fragment or variant of an antibody or an immunoglobulin, specifically binds. An “epitope” is antigenic and thus the term epitope is sometimes also referred to herein as “antigenic structure” or “antigenic determinant”. Thus, the binding domain is an “antigen interaction site”. Said binding/interaction is also understood to define a “specific recognition”.

[0052] “Epitopes” can be formed both by contiguous amino acids or non-contiguous amino acids juxtaposed by tertiary folding of a protein. A “linear epitope” is an epitope where an amino acid primary sequence comprises the recognized epitope. A linear epitope typically includes at least 3 or at least 4, and more usually, at least 5 or at least 6 or at least 7, for example, about 8 to about 10 amino acids in a unique sequence.

[0053] A “conformational epitope”, in contrast to a linear epitope, is an epitope wherein the primary sequence of the amino acids comprising the epitope is not the sole defining component of the epitope recognized (e.g., an epitope wherein the primary sequence of amino acids is not necessarily recognized by the binding domain). Typically a conformational epitope comprises an increased number of amino acids relative to a linear epitope. With regard to recognition of conformational epitopes, the binding domain recognizes a three-dimensional structure of the antigen, preferably a peptide or protein or fragment thereof (in the context of the present invention, the antigenic structure for one of the binding domains is comprised within the MSLN protein). For example, when a protein molecule folds to form a three-dimensional structure, certain amino acids and/or the polypeptide backbone forming the conformational epitope become juxtaposed enabling the antibody to recognize the epitope. Methods of determining the conformation of epitopes include, but are not limited to, x-ray crystallography, two-dimensional nuclear magnetic resonance (2D-NMR) spectroscopy and site-directed spin labeling and electron paramagnetic resonance (EPR) spectroscopy.

[0054] A method for epitope mapping is described in the following: When a region (a contiguous amino acid stretch) in the human MSLN protein is exchanged/replaced with its corresponding region of a non-human and non-primate MSLN antigen (e.g., mouse MSLN, but others like chicken, rat, hamster, rabbit etc. might also be conceivable), a decrease in the binding of the binding domain is expected to occur, unless the binding domain is cross-reactive for the non-human, non-primate MSLN used. Said decrease is preferably at least 10%, 20%, 30%, 40%, or 50%; more preferably at least 60%, 70%, or 80%, and most preferably 90%, 95% or even 100% in comparison to the binding to the respective region in the human MSLN protein, whereby binding to the respective region in the human MSLN protein is set to be 100%. It is envisaged that the aforementioned human MSLN/non-human MSLN chimeras are expressed in CHO cells. The human MSLN/non-human MSLN chimeras may also be fused with a transmembrane domain and/or cytoplasmic domain of a different membrane-bound protein such as EpCAM, although such technique was not necessary for the method described in Examples 1 and 2.

[0055] In an alternative or additional method for epitope mapping, several truncated versions of the human MSLN extracellular domain can be generated in order to determine a specific region that is recognized by a binding domain. In these truncated versions, the different extracellular MSLN

domains/sub-domains or regions are stepwise deleted, starting from the N-terminus. The truncated MSLN versions that may be expressed in CHO cells. It is also envisaged that the truncated MSLN versions may be fused with a transmembrane domain and/or cytoplasmic domain of a different membrane-bound protein such as EpCAM. It is also envisaged that the truncated MSLN versions may encompass a signal peptide domain at their N-terminus, for example a signal peptide derived from mouse IgG heavy chain signal peptide. It is furthermore envisaged that the truncated MSLN versions may encompass a v5 domain at their N-terminus (following the signal peptide) which allows verifying their correct expression on the cell surface. A decrease or a loss of binding is expected to occur with those truncated MSLN versions which do not encompass any more the MSLN region that is recognized by the binding domain. The decrease of binding is preferably at least 10%, 20%, 30%, 40%, 50%; more preferably at least 60%, 70%, 80%, and most preferably 90%, 95% or even 100%, whereby binding to the entire human MSLN protein (or its extracellular region or domain) is set to be 100%.

[0056] A further method to determine the contribution of a specific residue of a target antigen to the recognition by a antibody construct or binding domain is alanine scanning (see e.g. Morrison K L & Weiss G A. *Cur Opin Chem Biol.* 2001 June; 5(3):302-7), where each residue to be analyzed is replaced by alanine, e.g. via site-directed mutagenesis. Alanine is used because of its non-bulky, chemically inert, methyl functional group that nevertheless mimics the secondary structure references that many of the other amino acids possess. Sometimes bulky amino acids such as valine or leucine can be used in cases where conservation of the size of mutated residues is desired. Alanine scanning is a mature technology which has been used for a long period of time.

[0057] The interaction between the binding domain and the epitope or the region comprising the epitope implies that a binding domain exhibits appreciable affinity for the epitope/the region comprising the epitope on a particular protein or antigen (here: MSLN and CD3, respectively) and, generally, does not exhibit significant reactivity with proteins or antigens other than MSLN or CD3. “Appreciable affinity” includes binding with an affinity of about 10^{-6} M (KD) or stronger. Preferably, binding is considered specific when the binding affinity is about 10^{-12} to 10^{-8} M, 10^{-12} to 10^{-9} M, 10^{-12} to 10^{-10} M, 10^{-11} to 10^{-8} M, preferably of about 10^{-11} to 10^{-9} M. Whether a binding domain specifically reacts with or binds to a target can be tested readily by, inter alia, comparing the reaction of said binding domain with a target protein or antigen with the reaction of said binding domain with proteins or antigens other than MSLN or CD3. Preferably, a binding domain of the invention does not essentially or substantially bind to proteins or antigens other than MSLN or CD3 (i.e., the first binding domain is not capable of binding to proteins other than MSLN and the second binding domain is not capable of binding to proteins other than CD3).

[0058] The term “does not essentially/substantially bind” or “is not capable of binding” means that a binding domain of the present invention does not bind a protein or antigen other than MSLN or CD3, i.e., does not show reactivity of more than 30%, preferably not more than 20%, more preferably not more than 10%, particularly preferably not more than 9%, 8%, 7%, 6% or 5% with proteins or antigens other

than MSLN or CD3, whereby binding to MSLN or CD3, respectively, is set to be 100%.

[0059] Specific binding is believed to be effected by specific motifs in the amino acid sequence of the binding domain and the antigen. Thus, binding is achieved as a result of their primary, secondary and/or tertiary structure as well as the result of secondary modifications of said structures. The specific interaction of the antigen-interaction-site with its specific antigen may result in a simple binding of said site to the antigen. Moreover, the specific interaction of the antigen-interaction-site with its specific antigen may alternatively or additionally result in the initiation of a signal, e.g. due to the induction of a change of the conformation of the antigen, an oligomerization of the antigen, etc.

[0060] It is also preferred in one embodiment of the invention that the second binding domain binds to human CD3 epsilon and to *Callithrix jacchus*, *Saguinus Oedipus* or *Saimir sciureus* CD3 epsilon.

[0061] In another aspect, the present invention provides a bispecific antibody construct comprising a first binding domain which binds to human MSLN on the surface of a target cell and a second binding domain which binds to human CD3 on the surface of a T cell, wherein the first binding domain binds to an epitope of MSLN which is comprised within the region of the human MSLN having a sequence as depicted in SEQ ID NO selected from the group consisting of SEQ ID NO: 244 (cluster 1+2), SEQ ID NO: 245 (cluster 2+3) and SEQ ID NO: 241 (cluster 4).

[0062] Preferably, the first binding domain of the bispecific antibody construct of the invention comprises a VH region comprising CDR-H1, CDR-H2 and CDR-H3 and a VL region comprising CDR-L1, CDR-L2 and CDR-L3 selected from the group consisting of:

[0063] a) CDR-H1 as depicted in SEQ ID NO: 151, CDR-H2 as depicted in SEQ ID NO: 152, CDR-H3 as depicted in SEQ ID NO: 153, CDR-L1 as depicted in SEQ ID NO: 154, CDR-L2 as depicted in SEQ ID NO: 155 and CDR-L3 as depicted in SEQ ID NO: 156;

[0064] b) CDR-H1 as depicted in SEQ ID NO: 161, CDR-H2 as depicted in SEQ ID NO: 162, CDR-H3 as depicted in SEQ ID NO: 163, CDR-L1 as depicted in SEQ ID NO: 164, CDR-L2 as depicted in SEQ ID NO: 165 and CDR-L3 as depicted in SEQ ID NO: 166;

[0065] c) CDR-H1 as depicted in SEQ ID NO: 171, CDR-H2 as depicted in SEQ ID NO: 172, CDR-H3 as depicted in SEQ ID NO: 173, CDR-L1 as depicted in SEQ ID NO: 174, CDR-L2 as depicted in SEQ ID NO: 175 and CDR-L3 as depicted in SEQ ID NO: 176;

[0066] d) CDR-H1 as depicted in SEQ ID NO: 181, CDR-H2 as depicted in SEQ ID NO: 182, CDR-H3 as depicted in SEQ ID NO: 183, CDR-L1 as depicted in SEQ ID NO: 184, CDR-L2 as depicted in SEQ ID NO: 185 and CDR-L3 as depicted in SEQ ID NO: 186;

[0067] e) CDR-H1 as depicted in SEQ ID NO: 191, CDR-H2 as depicted in SEQ ID NO: 192, CDR-H3 as depicted in SEQ ID NO: 193, CDR-L1 as depicted in SEQ ID NO: 194, CDR-L2 as depicted in SEQ ID NO: 195 and CDR-L3 as depicted in SEQ ID NO: 196;

[0068] f) CDR-H1 as depicted in SEQ ID NO: 201, CDR-H2 as depicted in SEQ ID NO: 202, CDR-H3 as depicted in SEQ ID NO: 203, CDR-L1 as depicted in SEQ ID NO: 204, CDR-L2 as depicted in SEQ ID NO: 205 and CDR-L3 as depicted in SEQ ID NO: 206;

[0069] g) CDR-H1 as depicted in SEQ ID NO: 211, CDR-H2 as depicted in SEQ ID NO: 212, CDR-H3 as depicted in SEQ ID NO: 213, CDR-L1 as depicted in SEQ ID NO: 214, CDR-L2 as depicted in SEQ ID NO: 215 and CDR-L3 as depicted in SEQ ID NO: 216; and

[0070] h) CDR-H1 as depicted in SEQ ID NO: 221, CDR-H2 as depicted in SEQ ID NO: 222, CDR-H3 as depicted in SEQ ID NO: 223, CDR-L1 as depicted in SEQ ID NO: 224, CDR-L2 as depicted in SEQ ID NO: 225 and CDR-L3 as depicted in SEQ ID NO: 226.

[0071] The term “variable” refers to the portions of the antibody or immunoglobulin domains that exhibit variability in their sequence and that are involved in determining the specificity and binding affinity of a particular antibody (i.e., the “variable domain(s)”). The pairing of a variable heavy chain (VH) and a variable light chain (VL) together forms a single antigen-binding site.

[0072] Variability is not evenly distributed throughout the variable domains of antibodies; it is concentrated in sub-domains of each of the heavy and light chain variable regions. These sub-domains are called “hypervariable regions” or “complementarity determining regions” (CDRs). The more conserved (i.e., non-hypervariable) portions of the variable domains are called the “framework” regions (FRM or FR) and provide a scaffold for the six CDRs in three dimensional space to form an antigen-binding surface. The variable domains of naturally occurring heavy and light chains each comprise four FRM regions (FR1, FR2, FR3, and FR4), largely adopting a β -sheet configuration, connected by three hypervariable regions, which form loops connecting, and in some cases forming part of, the β -sheet structure. The hypervariable regions in each chain are held together in close proximity by the FRM and, with the hypervariable regions from the other chain, contribute to the formation of the antigen-binding site (see Kabat et al., loc. cit.).

[0073] The terms “CDR”, and its plural “CDRs”, refer to the complementarity determining region of which three make up the binding character of a light chain variable region (CDR-L1, CDR-L2 and CDR-L3) and three make up the binding character of a heavy chain variable region (CDR-H1, CDR-H2 and CDR-H3). CDRs contain most of the residues responsible for specific interactions of the antibody with the antigen and hence contribute to the functional activity of an antibody molecule: they are the main determinants of antigen specificity.

[0074] The exact definitional CDR boundaries and lengths are subject to different classification and numbering systems. CDRs may therefore be referred to by Kabat, Chothia, contact or any other boundary definitions, including the numbering system described herein. Despite differing boundaries, each of these systems has some degree of overlap in what constitutes the so called “hypervariable regions” within the variable sequences. CDR definitions according to these systems may therefore differ in length and boundary areas with respect to the adjacent framework region. See for example Kabat (an approach based on cross-species sequence variability), Chothia (an approach based on crystallographic studies of antigen-antibody complexes), and/or MacCallum (Kabat et al., loc. cit.; Chothia et al., J. Mol. Biol, 1987, 196: 901-917; and MacCallum et al., J. Mol. Biol, 1996, 262: 732). Still another standard for characterizing the antigen binding site is the AbM definition used by Oxford Molecular’s AbM antibody modeling soft-

ware. See, e.g., Protein Sequence and Structure Analysis of Antibody Variable Domains. In: Antibody Engineering Lab Manual (Ed.: Duebel, S. and Kontermann, R., Springer-Verlag, Heidelberg). To the extent that two residue identification techniques define regions of overlapping, but not identical regions, they can be combined to define a hybrid CDR. However, the numbering in accordance with the so-called Kabat system is preferred.

[0075] Typically, CDRs form a loop structure that can be classified as a canonical structure. The term “canonical structure” refers to the main chain conformation that is adopted by the antigen binding (CDR) loops. From comparative structural studies, it has been found that five of the six antigen binding loops have only a limited repertoire of available conformations. Each canonical structure can be characterized by the torsion angles of the polypeptide backbone. Correspondent loops between antibodies may, therefore, have very similar three dimensional structures, despite high amino acid sequence variability in most parts of the loops (Chothia and Lesk, J. Mol. Biol., 1987,196: 901; Chothia et al., Nature, 1989, 342: 877; Martin and Thornton, J. Mol. Biol, 1996, 263: 800). Furthermore, there is a relationship between the adopted loop structure and the amino acid sequences surrounding it. The conformation of a particular canonical class is determined by the length of the loop and the amino acid residues residing at key positions within the loop, as well as within the conserved framework (i.e., outside of the loop). Assignment to a particular canonical class can therefore be made based on the presence of these key amino acid residues.

[0076] The term “canonical structure” may also include considerations as to the linear sequence of the antibody, for example, as catalogued by Kabat (Kabat et al., loc. cit.). The Kabat numbering scheme (system) is a widely adopted standard for numbering the amino acid residues of an antibody variable domain in a consistent manner and is the preferred scheme applied in the present invention as also mentioned elsewhere herein. Additional structural considerations can also be used to determine the canonical structure of an antibody. For example, those differences not fully reflected by Kabat numbering can be described by the numbering system of Chothia et al. and/or revealed by other techniques, for example, crystallography and two- or three-dimensional computational modeling. Accordingly, a given antibody sequence may be placed into a canonical class which allows for, among other things, identifying appropriate chassis sequences (e.g., based on a desire to include a variety of canonical structures in a library). Kabat numbering of antibody amino acid sequences and structural considerations as described by Chothia et al., loc. cit. and their implications for construing canonical aspects of antibody structure, are described in the literature. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known in the art. For a review of the antibody structure, see Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, eds. Harlow et al., 1988.

[0077] The CDR3 of the light chain and, particularly, the CDR3 of the heavy chain may constitute the most important determinants in antigen binding within the light and heavy chain variable regions. In some antibody constructs, the heavy chain CDR3 appears to constitute the major area of contact between the antigen and the antibody. In vitro selection schemes in which CDR3 alone is varied can be

used to vary the binding properties of an antibody or determine which residues contribute to the binding of an antigen. Hence, CDR3 is typically the greatest source of molecular diversity within the antibody-binding site. H3, for example, can be as short as two amino acid residues or greater than 26 amino acids.

[0078] In a classical full-length antibody or immunoglobulin, each light (L) chain is linked to a heavy (H) chain by one covalent disulfide bond, while the two H chains are linked to each other by one or more disulfide bonds depending on the H chain isotype. The CH domain most proximal to VH is usually designated as CH1. The constant (“C”) domains are not directly involved in antigen binding, but exhibit various effector functions, such as antibody-dependent, cell-mediated cytotoxicity and complement activation. The Fc region of an antibody is comprised within the heavy chain constant domains and is for example able to interact with cell surface located Fc receptors.

[0079] The sequence of antibody genes after assembly and somatic mutation is highly varied, and these varied genes are estimated to encode 10^{10} different antibody molecules (Immunoglobulin Genes, 2nd ed., eds. Jonio et al., Academic Press, San Diego, CA, 1995). Accordingly, the immune system provides a repertoire of immunoglobulins. The term “repertoire” refers to at least one nucleotide sequence derived wholly or partially from at least one sequence encoding at least one immunoglobulin. The sequence(s) may be generated by rearrangement in vivo of the V, D, and J segments of heavy chains, and the V and J segments of light chains. Alternatively, the sequence(s) can be generated from a cell in response to which rearrangement occurs, e.g., in vitro stimulation. Alternatively, part or all of the sequence(s) may be obtained by DNA splicing, nucleotide synthesis, mutagenesis, and other methods, see, e.g., U.S. Pat. No. 5,565,332. A repertoire may include only one sequence or may include a plurality of sequences, including ones in a genetically diverse collection.

[0080] A preferred antibody construct according to the invention can also be defined as a bispecific antibody construct comprising a first (preferably human) binding domain which binds to human MSLN on the surface of a target cell and a second binding domain which binds to human CD3 on the surface of a T cell, wherein the first binding domain binds to the same epitope of MSLN as an antibody selected from the group consisting of MS_1 to MS_8, i.e., an antibody comprising a VH region comprising CDR-H1, CDR-H2 and CDR-H3 and a VL region comprising CDR-L1, CDR-L2 and CDR-L3 selected from the group consisting of:

[0081] a) CDR-H1 as depicted in SEQ ID NO: 151, CDR-H2 as depicted in SEQ ID NO: 152, CDR-H3 as depicted in SEQ ID NO: 153, CDR-L1 as depicted in SEQ ID NO: 154, CDR-L2 as depicted in SEQ ID NO: 155 and CDR-L3 as depicted in SEQ ID NO: 156;

[0082] b) CDR-H1 as depicted in SEQ ID NO: 161, CDR-H2 as depicted in SEQ ID NO: 162, CDR-H3 as depicted in SEQ ID NO: 163, CDR-L1 as depicted in SEQ ID NO: 164, CDR-L2 as depicted in SEQ ID NO: 165 and CDR-L3 as depicted in SEQ ID NO: 166;

[0083] c) CDR-H1 as depicted in SEQ ID NO: 171, CDR-H2 as depicted in SEQ ID NO: 172, CDR-H3 as depicted in SEQ ID NO: 173, CDR-L1 as depicted in SEQ ID NO: 174, CDR-L2 as depicted in SEQ ID NO: 175 and CDR-L3 as depicted in SEQ ID NO: 176;

- [0084] d) CDR-H1 as depicted in SEQ ID NO: 181, CDR-H2 as depicted in SEQ ID NO: 182, CDR-H3 as depicted in SEQ ID NO: 183, CDR-L1 as depicted in SEQ ID NO: 184, CDR-L2 as depicted in SEQ ID NO: 185 and CDR-L3 as depicted in SEQ ID NO: 186;
- [0085] e) CDR-H1 as depicted in SEQ ID NO: 191, CDR-H2 as depicted in SEQ ID NO: 192, CDR-H3 as depicted in SEQ ID NO: 193, CDR-L1 as depicted in SEQ ID NO: 194, CDR-L2 as depicted in SEQ ID NO: 195 and CDR-L3 as depicted in SEQ ID NO: 196;
- [0086] f) CDR-H1 as depicted in SEQ ID NO: 201, CDR-H2 as depicted in SEQ ID NO: 202, CDR-H3 as depicted in SEQ ID NO: 203, CDR-L1 as depicted in SEQ ID NO: 204, CDR-L2 as depicted in SEQ ID NO: 205 and CDR-L3 as depicted in SEQ ID NO: 206;
- [0087] g) CDR-H1 as depicted in SEQ ID NO: 211, CDR-H2 as depicted in SEQ ID NO: 212, CDR-H3 as depicted in SEQ ID NO: 213, CDR-L1 as depicted in SEQ ID NO: 214, CDR-L2 as depicted in SEQ ID NO: 215 and CDR-L3 as depicted in SEQ ID NO: 216; and
- [0088] h) CDR-H1 as depicted in SEQ ID NO: 221, CDR-H2 as depicted in SEQ ID NO: 222, CDR-H3 as depicted in SEQ ID NO: 223, CDR-L1 as depicted in SEQ ID NO: 224, CDR-L2 as depicted in SEQ ID NO: 225 and CDR-L3 as depicted in SEQ ID NO: 226.

[0089] Whether or not an antibody construct binds to the same epitope of MSLN as another given antibody construct can be measured e.g. by epitope mapping with chimeric or truncated target molecules, e.g. as described herein above and in the appended Examples 1 and 2.

[0090] A preferred antibody construct according to the invention can also be defined as a bispecific antibody construct comprising a first (preferably human) binding domain which binds to human MSLN on the surface of a target cell and a second binding domain which binds to human CD3 on the surface of a T cell, wherein the first binding domain competes for binding with an antibody selected from the group consisting of MS_1 to MS_8, i.e., an antibody comprising a VH region comprising CDR-H1, CDR-H2 and CDR-H3 and a VL region comprising CDR-L1, CDR-L2 and CDR-L3 selected from the group consisting of those described above.

[0091] Whether or not an antibody construct competes for binding with another given antibody construct can be measured in a competition assay such as a competitive ELISA or a cell-based competition assay. Avidin-coupled microparticles (beads) can also be used. Similar to an avidin-coated ELISA plate, when reacted with a biotinylated protein, each of these beads can be used as a substrate on which an assay can be performed. Antigen is coated onto a bead and then precoated with the first antibody. The second antibody is added and any additional binding is determined. Possible means for the read-out includes flow cytometry.

[0092] In one embodiment, the first binding domain of the antibody construct of the invention comprises a VH region selected from the group consisting of those depicted in SEQ ID NO: 157, SEQ ID NO: 167, SEQ ID NO: 177, SEQ ID NO: 187, SEQ ID NO: 197, SEQ ID NO: 207, SEQ ID NO: 217, and SEQ ID NO: 227.

[0093] In a further embodiment, the first binding domain of the antibody construct of the invention comprises a VL region selected from the group consisting of those depicted in SEQ ID NO: 158, SEQ ID NO: 168, SEQ ID NO: 178,

SEQ ID NO: 188, SEQ ID NO: 198, SEQ ID NO: 208, SEQ ID NO: 218, and SEQ ID NO: 228.

[0094] In another embodiment, the first binding domain of the antibody construct of the invention comprises a VH region and a VL region selected from the group consisting of pairs of a VH region and a VL region as depicted in SEQ ID NO: 157+158, SEQ ID NO: 167+168, SEQ ID NO: 177+178, SEQ ID NO: 187+188, SEQ ID NO: 197+198, SEQ ID NO: 207+208, SEQ ID NO: 217+218, and SEQ ID NO: 227+228.

[0095] In yet a further embodiment, the first binding domain of the antibody construct of the invention comprises a polypeptide selected from the group consisting of those depicted in SEQ ID NO: 159, SEQ ID NO: 169, SEQ ID NO: 179, SEQ ID NO: 189, SEQ ID NO: 199, SEQ ID NO: 209, SEQ ID NO: 219, and SEQ ID NO: 229.

[0096] The above first binding domains (which are specified by their CDRs, VH region and VL region and combinations thereof) characterize as binding domains which bind to an epitope of MSLN as depicted in SEQ ID NOs: 231, 232, and 233.

[0097] The term “bispecific” as used herein refers to an antibody construct which is “at least bispecific”, i.e., it comprises at least a first binding domain and a second binding domain, wherein the first binding domain binds to one antigen or target (here: MSLN), and the second binding domain binds to another antigen or target (here: CD3). Accordingly, antibody constructs according to the invention comprise specificities for at least two different antigens or targets.

[0098] The term “bispecific antibody construct” of the invention also encompasses multispecific antibody constructs such as trispecific antibody constructs, the latter ones including three binding domains, or constructs having more than three (e.g. four, five . . .) specificities.

[0099] Given that the antibody constructs according to the invention are (at least) bispecific, they do not occur naturally and they are markedly different from naturally occurring products. A “bispecific” antibody construct or immunoglobulin is hence an artificial hybrid antibody or immunoglobulin having at least two distinct binding sites with different specificities. Bispecific antibody constructs can be produced by a variety of methods including fusion of hybridomas or linking of Fab' fragments. See, e.g., Song-sivilai & Lachmann, Clin. Exp. Immunol. 79:315-321 (1990).

[0100] The at least two binding domains and the variable domains of the antibody construct of the present invention may or may not comprise peptide linkers (spacer peptides). The term “peptide linker” comprises in accordance with the present invention an amino acid sequence by which the amino acid sequences of one (variable and/or binding) domain and another (variable and/or binding) domain of the antibody construct of the invention are linked with each other. An essential technical feature of such peptide linker is that it does not comprise any polymerization activity. Among the suitable peptide linkers are those described in U.S. Pat. Nos. 4,751,180 and 4,935,233 or WO 88/09344. The peptide linkers can also be used to attach other domains or modules or regions (such as half-life extending domains) to the antibody construct of the invention.

[0101] In the event that a linker is used, this linker is preferably of a length and sequence sufficient to ensure that each of the first and second domains can, independently

from one another, retain their differential binding specificities. For peptide linkers which connect the at least two binding domains (or two variable domains) in the antibody construct of the invention, those peptide linkers are preferred which comprise only a few number of amino acid residues, e.g. 12 amino acid residues or less. Thus, peptide linkers of 12, 11, 10, 9, 8, 7, 6 or 5 amino acid residues are preferred. An envisaged peptide linker with less than 5 amino acids comprises 4, 3, 2 or one amino acid(s), wherein Gly-rich linkers are preferred. A particularly preferred “single” amino acid in the context of said “peptide linker” is Gly. Accordingly, said peptide linker may consist of the single amino acid Gly. Another preferred embodiment of a peptide linker is characterized by the amino acid sequence Gly-Gly-Gly-Gly-Ser, i.e. Gly₄Ser (SEQ ID NO: 1), or polymers thereof, i.e. (Gly₄Ser)_x, where x is an integer of 1 or greater (e.g. 2 or 3). Preferred linkers are depicted in SEQ ID NOs: 1-9. The characteristics of said peptide linker, which comprise the absence of the promotion of secondary structures, are known in the art and are described e.g. in Dall’Acqua et al. (Biochem. (1998) 37, 9266-9273), Cheadle et al. (Mol Immunol (1992) 29, 21-30) and Raag and Whitlow (FASEB (1995) 9(1), 73-80). Peptide linkers which furthermore do not promote any secondary structures are preferred. The linkage of said domains to each other can be provided, e.g., by genetic engineering, as described in the examples. Methods for preparing fused and operatively linked bispecific single chain constructs and expressing them in mammalian cells or bacteria are well-known in the art (e.g. WO 99/54440 or Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 2001).

[0102] As described herein above, the invention provides a preferred embodiment wherein the antibody construct is in a format selected from the group consisting of (scFv)₂, scFv-single domain mAb, diabodies and oligomers of any of the those formats.

[0103] According to a particularly preferred embodiment, and as documented in the appended examples, the antibody construct of the invention is a “bispecific single chain antibody construct”, more preferably a bispecific “single chain Fv” (scFv). Although the two domains of the Fv fragment, VL and VH, are coded for by separate genes, they can be joined, using recombinant methods, by a synthetic linker—as described hereinbefore—that enables them to be made as a single protein chain in which the VL and VH regions pair to form a monovalent molecule; see e.g., Huston et al. (1988) Proc. Natl. Acad. Sci USA 85:5879-5883). These antibody fragments are obtained using conventional techniques known to those with skill in the art, and the fragments are evaluated for function in the same manner as are whole or full-length antibodies. A single-chain variable fragment (scFv) is hence a fusion protein of the variable region of the heavy chain (VH) and of the light chain (VL) of immunoglobulins, usually connected with a short linker peptide of about ten to about 25 amino acids, preferably about 15 to 20 amino acids. The linker is usually rich in glycine for flexibility, as well as serine or threonine for solubility, and can either connect the N-terminus of the VH with the C-terminus of the VL, or vice versa. This protein retains the specificity of the original immunoglobulin, despite removal of the constant regions and introduction of the linker.

[0104] Bispecific single chain molecules are known in the art and are described in WO 99/54440, Mack, J. Immunol. (1997), 158, 3965-3970, Mack, PNAS, (1995), 92, 7021-7025, Kufer, Cancer Immunol. Immunother., (1997), 45, 193-197, Löffler, Blood, (2000), 95, 6, 2098-2103, Brühl, Immunol., (2001), 166, 2420-2426, Kipriyanov, J. Mol. Biol., (1999), 293, 41-56. Techniques described for the production of single chain antibodies (see, inter alia, U.S. Pat. No. 4,946,778, Kontermann and Dübel (2010), loc. cit. and Little (2009), loc. cit.) can be adapted to produce single chain antibody constructs specifically recognizing (an) elected target(s).

[0105] Bivalent (also called divalent) or bispecific single-chain variable fragments (bi-scFvs or di-scFvs having the format (scFv)₂ can be engineered by linking two scFv molecules (e.g. with linkers as described hereinbefore). If these two scFv molecules have the same binding specificity, the resulting (scFv)₂ molecule will preferably be called bivalent (i.e. it has two valences for the same target epitope). If the two scFv molecules have different binding specificities, the resulting (scFv)₂ molecule will preferably be called bispecific. The linking can be done by producing a single peptide chain with two VH regions and two VL regions, yielding tandem scFvs (see e.g. Kufer P. et al., (2004) Trends in Biotechnology 22(5):238-244). Another possibility is the creation of scFv molecules with linker peptides that are too short for the two variable regions to fold together (e.g. about five amino acids), forcing the scFvs to dimerize. This type is known as diabodies (see e.g. Hollinger, Philipp et al., (July 1993) Proceedings of the National Academy of Sciences of the United States of America 90 (14): 6444-8.).

[0106] According to an also preferred embodiment of the antibody construct of the invention the heavy chain (VH) and of the light chain (VL) of a binding domain binding either to the target antigen MSLN or CD3 are not directly connected via an above described peptide linker but the binding domain is formed due to the formation of a bispecific molecule as described for the diabody. Thus, the VH chain of the CD3 binding domain may be fused to the VL of the MSLN binding domain via such peptide linker, while the VH chain of the MSLN binding domain is fused to the VL of the CD3 binding domain via such peptide linker.

[0107] Single domain antibodies comprise merely one (monomeric) antibody variable domain which is able to bind selectively to a specific antigen, independently of other V regions or domains. The first single domain antibodies were engineered from heavy chain antibodies found in camelids, and these are called V_HH fragments. Cartilaginous fishes also have heavy chain antibodies (IgNAR) from which single domain antibodies called V_{NAR} fragments can be obtained. An alternative approach is to split the dimeric variable domains from common immunoglobulins e.g. from humans or rodents into monomers, hence obtaining VH or VL as a single domain Ab. Although most research into single domain antibodies is currently based on heavy chain variable domains, nanobodies derived from light chains have also been shown to bind specifically to target epitopes. Examples of single domain antibodies are called sdAb, nanobodies or single variable domain antibodies.

[0108] A (single domain mAb)₂ is hence a monoclonal antibody construct composed of (at least) two single domain monoclonal antibodies, which are individually selected from the group comprising VH, VL, V_HH and V_{NAR}. The linker is preferably in the form of a peptide linker. Similarly, an

“scFv-single domain mAb” is a monoclonal antibody construct composed of at least one single domain antibody as described above and one scFv molecule as described above. Again, the linker is preferably in the form of a peptide linker.

[0109] It is furthermore envisaged that the present invention provides a bispecific antibody construct comprising a first binding domain which binds to human MSLN on the surface of a target cell and a second binding domain which binds to human CD3 on the surface of a T cell, wherein the first binding domain binds to an epitope of MSLN which is comprised within the region as depicted in SEQ ID NO: 245 (cluster 2+3).

[0110] Accordingly, in a further aspect of the invention, the first binding domain of the bispecific antibody construct comprises a VH region comprising CDR-H1, CDR-H2 and CDR-H3 and a VL region comprising CDR-L1, CDR-L2 and CDR-L3 selected from the group consisting of:

[0111] a) CDR-H1 as depicted in SEQ ID NO: 161, CDR-H2 as depicted in SEQ ID NO: 162, CDR-H3 as depicted in SEQ ID NO: 163, CDR-L1 as depicted in SEQ ID NO: 164, CDR-L2 as depicted in SEQ ID NO: 165 and CDR-L3 as depicted in SEQ ID NO: 166;

[0112] b) CDR-H1 as depicted in SEQ ID NO: 171, CDR-H2 as depicted in SEQ ID NO: 172, CDR-H3 as depicted in SEQ ID NO: 173, CDR-L1 as depicted in SEQ ID NO: 174, CDR-L2 as depicted in SEQ ID NO: 175 and CDR-L3 as depicted in SEQ ID NO: 176;

[0113] c) CDR-H1 as depicted in SEQ ID NO: 181, CDR-H2 as depicted in SEQ ID NO: 182, CDR-H3 as depicted in SEQ ID NO: 183, CDR-L1 as depicted in SEQ ID NO: 184, CDR-L2 as depicted in SEQ ID NO: 185 and CDR-L3 as depicted in SEQ ID NO: 186;

[0114] d) CDR-H1 as depicted in SEQ ID NO: 191, CDR-H2 as depicted in SEQ ID NO: 192, CDR-H3 as depicted in SEQ ID NO: 193, CDR-L1 as depicted in SEQ ID NO: 194, CDR-L2 as depicted in SEQ ID NO: 195 and CDR-L3 as depicted in SEQ ID NO: 196;

[0115] e) CDR-H1 as depicted in SEQ ID NO: 201, CDR-H2 as depicted in SEQ ID NO: 202, CDR-H3 as depicted in SEQ ID NO: 203, CDR-L1 as depicted in SEQ ID NO: 204, CDR-L2 as depicted in SEQ ID NO: 205 and CDR-L3 as depicted in SEQ ID NO: 206.

[0116] In one embodiment, the first binding domain of the antibody construct of the invention comprises a VH region selected from the group consisting of those depicted in SEQ ID NO: 167, SEQ ID NO: 177, SEQ ID NO: 187, SEQ ID NO: 197, and SEQ ID NO: 207.

[0117] In a further embodiment, the first binding domain of the antibody construct of the invention comprises a VL region selected from the group consisting of those depicted in SEQ ID NO: 168, SEQ ID NO: 178, SEQ ID NO: 188, SEQ ID NO: 198, and SEQ ID NO: 208.

[0118] In another embodiment, the first binding domain of the antibody construct of the invention comprises a VH region and a VL region selected from the group consisting of pairs of a VH region and a VL region as depicted in SEQ ID NO: 167+168, SEQ ID NO: 177+178, SEQ ID NO: 187+188, SEQ ID NO: 197+198, and SEQ ID NO: 207+208.

[0119] In a further embodiment, the first binding domain of the antibody construct of the invention comprises a polypeptide selected from the group consisting of those depicted in SEQ ID NO: 169, SEQ ID NO: 179, SEQ ID NO: 189, SEQ ID NO: 199, and SEQ ID NO: 209.

[0120] It is also envisaged that the present invention provides a bispecific antibody construct comprising a first binding domain which binds to human MSLN on the surface of a target cell and a second binding domain which binds to human CD3 on the surface of a T cell, wherein the first binding domain binds to an epitope of MSLN which is comprised within the region as depicted in SEQ ID NO: 244 (cluster 1+2).

[0121] Accordingly, in a further aspect of the invention, the first binding domain of the bispecific antibody construct comprises a VH region comprising CDR-H1, CDR-H2 and CDR-H3 and a VL region comprising CDR-L1, CDR-L2 and CDR-L3 as follows:

[0122] (a) CDR-H1 as depicted in SEQ ID NO: 151, CDR-H2 as depicted in SEQ ID NO: 152, CDR-H3 as depicted in SEQ ID NO: 153, CDR-L1 as depicted in SEQ ID NO: 154, CDR-L2 as depicted in SEQ ID NO: 155 and CDR-L3 as depicted in SEQ ID NO: 156; or

[0123] (b) CDR-H1 as depicted in SEQ ID NO: 221, CDR-H2 as depicted in SEQ ID NO: 222, CDR-H3 as depicted in SEQ ID NO: 223, CDR-L1 as depicted in SEQ ID NO: 224, CDR-L2 as depicted in SEQ ID NO: 225 and CDR-L3 as depicted in SEQ ID NO: 226.

[0124] In one embodiment, the first binding domain of the antibody construct of the invention comprises a VH region depicted in SEQ ID NO: 157 or SEQ ID NO: 227.

[0125] In a further embodiment, the first binding domain of the antibody construct of the invention comprises a VL region depicted in SEQ ID NO: 158, and SEQ ID NO: 228.

[0126] In another embodiment, the first binding domain of the antibody construct of the invention comprises a VH region and a VL region selected from the group consisting of pairs of a VH region and a VL region as depicted in SEQ ID NO: 157+158, and SEQ ID NO: 227+228.

[0127] In a further embodiment, the first binding domain of the antibody construct of the invention comprises a polypeptide selected from the group consisting of those depicted in SEQ ID NO: 159, and SEQ ID NO: 229.

[0128] It is also envisaged that the present invention provides a bispecific antibody construct comprising a first binding domain which binds to human MSLN on the surface of a target cell and a second binding domain which binds to human CD3 on the surface of a T cell, wherein the first binding domain binds to an epitope of MSLN which is comprised within the region as depicted in SEQ ID NO: 241 (cluster 4).

[0129] Accordingly, in a further aspect of the invention, the first binding domain of the bispecific antibody construct comprises a VH region comprising CDR-H1 as depicted in SEQ ID NO: 211, CDR-H2 as depicted in SEQ ID NO: 212, CDR-H3 as depicted in SEQ ID NO: 213, CDR-L1 as depicted in SEQ ID NO: 214, CDR-L2 as depicted in SEQ ID NO: 215 and CDR-L3 as depicted in SEQ ID NO: 216.

[0130] In one embodiment, the first binding domain of the antibody construct of the invention comprises a VH region depicted in SEQ ID NO: 217.

[0131] In a further embodiment, the first binding domain of the antibody construct of the invention comprises a VL region depicted in SEQ ID NO: 218.

[0132] In another embodiment, the first binding domain of the antibody construct of the invention comprises a VH region and a VL region as depicted in SEQ ID NO: 217+218.

[0133] In a further embodiment, the first binding domain of the antibody construct of the invention comprises a polypeptide as depicted in SEQ ID NO: 219.

[0134] Another preferred antibody construct according to the invention can also be defined as a bispecific antibody construct comprising a first (preferably human) binding domain which binds to human MSLN on the surface of a target cell and a second binding domain which binds to human CD3 on the surface of a T cell, wherein the first binding domain competes for binding with an antibody selected from the group consisting of MS-3, MS-4 and MS-5, i.e., an antibody comprising a VH region comprising CDR-H1, CDR-H2 and CDR-H3 and a VL region comprising CDR-L1, CDR-L2 and CDR-L3 selected from the group consisting of those described above.

[0135] T cells or T lymphocytes are a type of lymphocyte (itself a type of white blood cell) that play a central role in cell-mediated immunity. There are several subsets of T cells, each with a distinct function. T cells can be distinguished from other lymphocytes, such as B cells and NK cells, by the presence of a T cell receptor (TCR) on the cell surface. The TCR is responsible for recognizing antigens bound to major histocompatibility complex (MHC) molecules and is composed of two different protein chains. In 95% of the T cells, the TCR consists of an alpha (α) and beta (β) chain. When the TCR engages with antigenic peptide and MHC (peptide/MHC complex), the T lymphocyte is activated through a series of biochemical events mediated by associated enzymes, co-receptors, specialized adaptor molecules, and activated or released transcription factors

[0136] The CD3 receptor complex is a protein complex and is composed of four chains. In mammals, the complex contains a CD3 γ (gamma) chain, a CD3 δ (delta) chain, and two CD3 ϵ (epsilon) chains. These chains associate with the T cell receptor (TCR) and the so-called ζ (zeta) chain to form the T cell receptor CD3 complex and to generate an activation signal in T lymphocytes. The CD3 γ (gamma), CD3 δ (delta), and CD3 ϵ (epsilon) chains are highly related cell-surface proteins of the immunoglobulin superfamily containing a single extracellular immunoglobulin domain. The intracellular tails of the CD3 molecules contain a single conserved motif known as an immunoreceptor tyrosine-based activation motif or ITAM for short, which is essential for the signaling capacity of the TCR. The CD3 epsilon molecule is a polypeptide which in humans is encoded by the CD3 ϵ gene which resides on chromosome 11. The most preferred epitope of CD3 epsilon is comprised within amino acid residues 1-27 of the human CD3 epsilon extracellular domain.

[0137] The redirected lysis of target cells via the recruitment of T cells by a multispecific, at least bispecific, antibody construct involves cytolytic synapse formation and delivery of perforin and granzymes. The engaged T cells are capable of serial target cell lysis, and are not affected by immune escape mechanisms interfering with peptide antigen processing and presentation, or clonal T cell differentiation; see, for example, WO 2007/042261.

[0138] Cytotoxicity mediated by MSLNxCD3 bispecific antibody constructs can be measured in various ways. See Examples 5. Effector cells can be e.g. stimulated enriched (human) CD8 positive T cells or unstimulated (human) peripheral blood mononuclear cells (PBMC). If the target cells are of macaque origin or express or are transfected with macaque MSLN, the effector cells should also be of

macaque origin such as a macaque T cell line, e.g. 4119LnPx. The target cells should express (at least the extracellular domain of) MSLN, e.g. human or macaque MSLN. Target cells can be a cell line (such as CHO) which is stably or transiently transfected with MSLN, e.g. human or macaque MSLN. Alternatively, the target cells can be a MSLN positive natural expresser cell line, such as the human cell line OVCAR-8. Usually EC₅₀ values are expected to be lower with target cell lines expressing higher levels of MSLN on the cell surface. The effector to target cell (E:T) ratio is usually about 10:1, but can also vary. Cytotoxic activity of MSLNxCD3 bispecific antibody constructs can be measured in a 51-chromium release assay (incubation time of about 18 hours) or in a FACS-based cytotoxicity assay (incubation time of about 48 hours). Modifications of the assay incubation time (cytotoxic reaction) are also possible. Other methods of measuring cytotoxicity are well-known to the skilled person and comprise MTT or MTS assays, ATP-based assays including bioluminescent assays, the sulforhodamine B (SRB) assay, WST assay, clonogenic assay and the ECIS technology.

[0139] The cytotoxic activity mediated by MSLNxCD3 bispecific antibody constructs of the present invention is preferably measured in a cell-based cytotoxicity assay. It may also be measured in a 51-chromium release assay. It is represented by the EC₅₀ value, which corresponds to the half maximal effective concentration (concentration of the antibody construct which induces a cytotoxic response halfway between the baseline and maximum). Preferably, the EC₅₀ value of the MSLNxCD3 bispecific antibody constructs is ≤ 5000 pM or ≤ 4000 pM, more preferably ≤ 3000 pM or ≤ 2000 pM, even more preferably ≤ 1000 pM or ≤ 500 pM, even more preferably ≤ 400 pM or ≤ 300 pM, even more preferably ≤ 200 pM, even more preferably ≤ 100 pM, even more preferably ≤ 50 pM, even more preferably ≤ 20 pM or ≤ 10 pM, and most preferably ≤ 5 pM.

[0140] The above given EC₅₀ values can be measured in different assays. The skilled person is aware that an EC₅₀ value can be expected to be lower when stimulated/enriched CDB+ T cells are used as effector cells, compared with unstimulated PBMC. It can furthermore be expected that the EC₅₀ values are lower when the target cells express a high number of the target antigen compared with a low target expression rate. For example, when stimulated/enriched human CD8+ T cells are used as effector cells (and either MSLN transfected cells such as CHO cells or a MSLN positive human cell line OVCAR-8 are used as target cells), the EC₅₀ value of the MSLN xCD3 bispecific antibody construct is preferably ≤ 1000 pM, more preferably ≤ 500 pM, even more preferably ≤ 250 pM, even more preferably ≤ 100 pM, even more preferably ≤ 50 pM, even more preferably ≤ 10 pM, and most preferably ≤ 5 pM. When human PBMCs are used as effector cells, the EC₅₀ value of the MSLNxCD3 bispecific antibody construct is preferably ≤ 5000 pM or ≤ 4000 pM (in particular when the target cells are a MSLN positive human cell line OVCAR-8), more preferably ≤ 2000 pM (in particular when the target cells are MSLN transfected cells such as CHO cells), more preferably ≤ 1000 pM or ≤ 500 pM, even more preferably ≤ 200 pM, even more preferably ≤ 150 pM, even more preferably ≤ 100 pM, and most preferably ≤ 50 pM, or lower. When a macaque T cell line such as LnPx4119 is used as effector cells, and a macaque MSLN transfected cell line such as CHO cells is used as target cell line, the EC₅₀ value of the MSLN xCD3

bispecific antibody construct is preferably ≤ 2000 pM or ≤ 1500 pM, more preferably ≤ 1000 pM or ≤ 500 pM, even more preferably ≤ 300 pM or ≤ 250 pM, even more preferably ≤ 100 pM, and most preferably ≤ 50 pM.

[0141] Preferably, the MSLNxCD3 bispecific antibody constructs of the present invention do not induce/mediate lysis or do not essentially induce/mediate lysis of MSLN negative cells such as CHO cells. The term “do not induce lysis”, “do not essentially induce lysis”, “do not mediate lysis” or “do not essentially mediate lysis” means that an antibody construct of the present invention does not induce or mediate lysis of more than 30%, preferably not more than 20%, more preferably not more than 10%, particularly preferably not more than 9%, 8%, 7%, 6% or 5% of MSLN negative cells, whereby lysis of a MSLN positive human cell line OVCAR-8 (see above) is set to be 100%. This usually applies for concentrations of the antibody construct of up to 500 nM. The skilled person knows how to measure cell lysis without further ado. Moreover, the present specification teaches specific instructions how to measure cell lysis.

[0142] The difference in cytotoxic activity between the monomeric and the dimeric isoform of individual MSLN xCD3 bispecific antibody constructs is referred to as “potency gap”. This potency gap can e.g. be calculated as ratio between EC_{50} values of the molecule’s monomeric and dimeric form, see Example 5.8. Potency gaps of the MSLNxCD3 bispecific antibody constructs of the present invention are preferably ≤ 5 , more preferably ≤ 4 , even more preferably ≤ 3 , even more preferably ≤ 2 , furthermore preferably ≤ 1 , and most preferably ≤ 0.3 .

[0143] The first and/or the second (or any further) binding domain(s) of the antibody construct of the invention is/are preferably cross-species specific for members of the mammalian order of primates. Cross-species specific CD3 binding domains are, for example, described in WO 2008/119567. According to one embodiment, the first and/or second binding domain, in addition to binding to human MSLN and human CD3, respectively, will also bind to MSLN/CD3 of primates including (but not limited to) new world primates (such as *Callithrix jacchus*, *Saguinus Oedipus* or *Saimir sciureus*), old world primates (such as baboons and macaques), gibbons, orangutans, and non-human homininae. It is envisaged that the first binding domain of the antibody construct of the invention which binds to human MSLN on the surface of a target cell also binds at least to macaque MSLN, and/or the second binding domain which binds to human CD3 on the surface of a T cell also binds at least to macaque CD3. A preferred macaque is *Macaca fascicularis*. *Macaca mulatta* (Rhesus) is also envisaged.

[0144] In one aspect of the invention, the first binding domain binds to human MSLN and further binds to macaque MSLN, such as MSLN of *Macaca fascicularis*, and more preferably, to macaque MSLN expressed on the surface macaque cells. A preferred *Macaca fascicularis* MSLN is depicted in SEQ ID NO: 234. The affinity of the first binding domain for macaque MSLN is preferably ≤ 15 nM, more preferably ≤ 10 nM, even more preferably ≤ 5 nM, even more preferably ≤ 1 nM, even more preferably ≤ 0.5 nM, even more preferably ≤ 0.1 nM, and most preferably ≤ 0.05 nM or even ≤ 0.01 nM.

[0145] Preferably the affinity gap of the antibody constructs according to the invention for binding macaque MSLN versus human MSLN [ma MSLN:hu MSLN] (as determined e.g. by BiaCore or by Scatchard analysis) is

<100 , preferably <20 , more preferably <15 , further preferably <10 , even more preferably <8 , more preferably <6 and most preferably <2 . Preferred ranges for the affinity gap of the antibody constructs according to the invention for binding macaque MSLN versus human MSLN are between 0.1 and 20, more preferably between 0.2 and 10, even more preferably between 0.3 and 6, even more preferably between 0.5 and 3 or between 0.5 and 2.5, and most preferably between 0.5 and 2 or between 0.6 and 2. See Examples 3.

[0146] Accordingly, in line with this invention antibody construct comprising a MSLN binder for the epitope cluster 2+3 are preferred, which all are shown to have an affinity gap ≤ 15 . Further preferred are antibody construct comprising a MSLN binder for the epitope cluster 2+3 having an affinity gap ≤ 6 , such as MS-3, MS-4 or MS-5 based bispecific antibody constructs.

[0147] In one embodiment of the antibody construct of the invention, the second binding domain binds to human CD3 epsilon and to *Callithrix jacchus*, *Saguinus Oedipus* or *Saimir sciureus* CD3 epsilon. Preferably, the second binding domain binds to an extracellular epitope of these CD3 epsilon chains. It is also envisaged that the second binding domain binds to an extracellular epitope of the human and the *Macaca* CD3 epsilon chain. The most preferred epitope of CD3 epsilon is comprised within amino acid residues 1-27 of the human CD3 epsilon extracellular domain. Even more specifically, the epitope comprises at least the amino acid sequence Gln-Asp-Gly-Asn-Glu. *Callithrix jacchus* and *Saguinus oedipus* are both new world primate belonging to the family of Callitrichidae, while *Saimir sciureus* is a new world primate belonging to the family of Cebidae.

[0148] It is particularly preferred for the antibody construct of the present invention that the second binding domain which binds to human CD3 on the surface of a T cell comprises a VL region comprising CDR-L1, CDR-L2 and CDR-L3 selected from:

[0149] (a) CDR-L1 as depicted in SEQ ID NO: 27 of WO 2008/119567, CDR-L2 as depicted in SEQ ID NO: 28 of WO 2008/119567 and CDR-L3 as depicted in SEQ ID NO: 29 of WO 2008/119567;

[0150] (b) CDR-L1 as depicted in SEQ ID NO: 117 of WO 2008/119567, CDR-L2 as depicted in SEQ ID NO: 118 of WO 2008/119567 and CDR-L3 as depicted in SEQ ID NO: 119 of WO 2008/119567; and

[0151] (c) CDR-L1 as depicted in SEQ ID NO: 153 of WO 2008/119567, CDR-L2 as depicted in SEQ ID NO: 154 of WO 2008/119567 and CDR-L3 as depicted in SEQ ID NO: 155 of WO 2008/119567.

[0152] In an alternatively preferred embodiment of the antibody construct of the present invention, the second binding domain which binds to human CD3 on the surface of a T cell comprises a VH region comprising CDR-H 1, CDR-H2 and CDR-H3 selected from:

[0153] (a) CDR-H1 as depicted in SEQ ID NO: 12 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 13 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 14 of WO 2008/119567;

[0154] (b) CDR-H1 as depicted in SEQ ID NO: 30 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 31 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 32 of WO 2008/119567;

[0155] (c) CDR-H1 as depicted in SEQ ID NO: 48 of WO 2008/119567, CDR-H2 as depicted in SEQ ID

- NO: 49 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 50 of WO 2008/119567;
- [0156] (d) CDR-H1 as depicted in SEQ ID NO: 66 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 67 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 68 of WO 2008/119567;
- [0157] (e) CDR-H1 as depicted in SEQ ID NO: 84 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 85 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 86 of WO 2008/119567;
- [0158] (f) CDR-H1 as depicted in SEQ ID NO: 102 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 103 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 104 of WO 2008/119567;
- [0159] (g) CDR-H1 as depicted in SEQ ID NO: 120 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 121 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 122 of WO 2008/119567;
- [0160] (h) CDR-H1 as depicted in SEQ ID NO: 138 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 139 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 140 of WO 2008/119567;
- [0161] (i) CDR-H1 as depicted in SEQ ID NO: 156 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 157 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 158 of WO 2008/119567; and
- [0162] (j) CDR-H1 as depicted in SEQ ID NO: 174 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 175 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 176 of WO 2008/119567.
- [0163] It is further preferred for the antibody construct of the present invention that the second binding domain which binds to human CD3 on the surface of a T cell comprises a VL region selected from the group consisting of a VL region as depicted in SEQ ID NO: 18, SEQ ID NO: 27, SEQ ID NO: 36, SEQ ID NO: 45, SEQ ID NO: 54, SEQ ID NO: 63, SEQ ID NO: 72, SEQ ID NO: 81, SEQ ID NO: 90, SEQ ID NO: 99, and SEQ ID NO: 102 (see also SEQ ID NO: 35, 39, 125, 129, 161 or 165 of WO 2008/119567).
- [0164] It is alternatively preferred that the second binding domain which binds to human CD3 on the surface of a T cell comprises a VH region selected from the group consisting of a VH region as depicted in SEQ ID NO: 17, SEQ ID NO: 26, SEQ ID NO: 35, SEQ ID NO: 44, SEQ ID NO: 53, SEQ ID NO: 62, SEQ ID NO: 71, SEQ ID NO: 80, SEQ ID NO: 89, SEQ ID NO: 98, and SEQ ID NO: 101 (see also SEQ ID NO: 15, 19, 33, 37, 51, 55, 69, 73, 87, 91, 105, 109, 123, 127, 141, 145, 159, 163, 177 or 181 of WO 2008/119567).
- [0165] More preferably, the antibody construct of the present invention is characterized by the second binding domain which binds to human CD3 on the surface of a T cell comprising a VL region and a VH region selected from the group consisting of:
- [0166] (a) a VL region as depicted in SEQ ID NO: 17 or 21 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 15 or 19 of WO 2008/119567;
- [0167] (b) a VL region as depicted in SEQ ID NO: 35 or 39 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 33 or 37 of WO 2008/119567;
- [0168] (c) a VL region as depicted in SEQ ID NO: 53 or 57 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 51 or 55 of WO 2008/119567;
- [0169] (d) a VL region as depicted in SEQ ID NO: 71 or 75 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 69 or 73 of WO 2008/119567;
- [0170] (e) a VL region as depicted in SEQ ID NO: 89 or 93 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 87 or 91 of WO 2008/119567;
- [0171] (f) a VL region as depicted in SEQ ID NO: 107 or 111 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 105 or 109 of WO 2008/119567;
- [0172] (g) a VL region as depicted in SEQ ID NO: 125 or 129 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 123 or 127 of WO 2008/119567;
- [0173] (h) a VL region as depicted in SEQ ID NO: 143 or 147 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 141 or 145 of WO 2008/119567;
- [0174] (i) a VL region as depicted in SEQ ID NO: 161 or 165 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 159 or 163 of WO 2008/119567; and
- [0175] (j) a VL region as depicted in SEQ ID NO: 179 or 183 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 177 or 181 of WO 2008/119567.
- [0176] Also preferred in connection with the antibody construct of the present invention is a second binding domain which binds to human CD3 on the surface of a T cell comprising a VL region as depicted in SEQ ID NO: 102 and a VH region as depicted in SEQ ID NO: 101.
- [0177] According to a preferred embodiment of the antibody construct of the present invention, the binding domains and in particular the second binding domain (which binds to human CD3 on the surface of a T cell) have the following format: The pairs of VH regions and VL regions are in the format of a single chain antibody (scFv). The VH and VL regions are arranged in the order VH-VL or VL-VH. It is preferred that the VH-region is positioned N-terminally of a linker sequence, and the VL-region is positioned C-terminally of the linker sequence.
- [0178] A preferred embodiment of the above described antibody construct of the present invention is characterized by the second binding domain which binds to human CD3 on the surface of a T cell comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 19, SEQ ID NO: 28, SEQ ID NO: 37, SEQ ID NO: 46, SEQ ID NO: 55, SEQ ID NO: 64, SEQ ID NO: 73, SEQ ID NO: 82, SEQ ID NO: 91, SEQ ID NO: 100, and SEQ ID NO: 103 (see also SEQ ID NOs: 23, 25, 41, 43, 59, 61, 77, 79, 95, 97, 113, 115, 131, 133, 149, 151, 167, 169, 185 or 187 of WO 2008/119567).
- [0179] It is also envisaged that the antibody construct of the invention has, in addition to its function to bind to the target molecules MSLN and CD3, a further function. In this format, the antibody construct is a trifunctional or multifunctional antibody construct by targeting target cells through binding to MSLN, mediating cytotoxic T cell activity through CD3 binding and providing a further function such as a fully functional Fc constant domain mediating antibody-dependent cellular cytotoxicity through recruitment of effector cells like NK cells, a label (fluorescent etc.), a therapeutic agent such as a toxin or radionuclide, and/or means to enhance serum half-life, etc.

[0180] Examples for means to extend serum half-life of the antibody constructs of the invention include peptides, proteins or domains of proteins, which are fused or otherwise attached to the antibody constructs. The group of peptides, proteins or protein domains includes peptides binding to other proteins with preferred pharmacokinetic profile in the human body such as serum albumin (see WO 2009/127691). An alternative concept of such half-life extending peptides includes peptides binding to the neonatal Fc receptor (FcRn, see WO 2007/098420), which can also be used in the constructs of the present invention. The concept of attaching larger domains of proteins or complete proteins includes e.g. the fusion of human serum albumin, variants or mutants of human serum albumin (see WO 2011/051489, WO 2012/059486, WO 2012/150319, WO 2013/135896, WO 2014/072481, WO 2013/075066) or domains thereof as well as the fusion of constant region of immunoglobulins (Fc domains) and variants thereof. Such variants of Fc domains may be optimized/modified in order to allow the desired pairing of dimers or multimers, to abolish Fc receptor binding (e.g. the Fc γ receptor) or for other reasons. A further concept known in the art to extend the half-life of small protein compounds in the human body is the pegylation of those compounds such as the antibody construct of the present invention.

[0181] In a preferred embodiment, the antibody construct of the invention is described as follows:

[0182] (a) a polypeptide comprising in the following order starting from the N-terminus:

[0183] a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 159, SEQ ID NO: 169, SEQ ID NO: 179, SEQ ID NO: 189, SEQ ID NO: 199, SEQ ID NO: 209, SEQ ID NO: 219, and SEQ ID NO: 229

[0184] a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 1-9; and

[0185] a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 19, SEQ ID NO: 28, SEQ ID NO: 37, SEQ ID NO: 46, SEQ ID NO: 55, SEQ ID NO: 64, SEQ ID NO: 73, SEQ ID NO: 82, SEQ ID NO: 91, SEQ ID NO: 100, and SEQ ID NO: 103; and

[0186] optionally a His-tag, such as the one depicted in SEQ ID NO 10;

[0187] (b) a polypeptide comprising in following order starting from the N-terminus:

[0188] a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 159, SEQ ID NO: 169, SEQ ID NO: 179, SEQ ID NO: 189, SEQ ID NO: 199, SEQ ID NO: 209, SEQ ID NO: 219, and SEQ ID NO: 229

[0189] a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 1-9;

[0190] a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 19, SEQ ID NO: 28, SEQ ID NO: 37, SEQ ID NO: 46, SEQ ID NO: 55, SEQ ID NO: 64, SEQ ID NO: 73, SEQ ID NO: 82, SEQ ID NO: 91, SEQ ID NO: 100, and SEQ ID NO: 103;

[0191] optionally a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 1-9;

[0192] a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NOs: 104-134; and

[0193] optionally a His-tag, such as the one depicted in SEQ ID NO 10;

[0194] (c) a polypeptide comprising in the following order starting from the N-terminus:

[0195] a polypeptide having the amino acid sequence QRFVTGHFGGLX₁PANG (SEQ ID NO: 135) whereas X₁ is Y or H; and

[0196] a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 159, SEQ ID NO: 169, SEQ ID NO: 179, SEQ ID NO: 189, SEQ ID NO: 199, SEQ ID NO: 209, SEQ ID NO: 219, and SEQ ID NO: 229;

[0197] a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 1-9;

[0198] a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 19, SEQ ID NO: 28, SEQ ID NO: 37, SEQ ID NO: 46, SEQ ID NO: 55, SEQ ID NO: 64, SEQ ID NO: 73, SEQ ID NO: 82, SEQ ID NO: 91, SEQ ID NO: 100, and SEQ ID NO: 103;

[0199] a polypeptide having the amino acid sequence QRFVTGHFGGLHPANG (SEQ ID NO: 137) or QRFCTGHFGGLHPCNG (SEQ ID NO: 139); and

[0200] optionally a His-tag, such as the one depicted in SEQ ID NO 10;

[0201] (d) a polypeptide comprising in the following order starting from the N-terminus

[0202] a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 17, SEQ ID NO: 26, SEQ ID NO: 35, SEQ ID NO: 44, SEQ ID NO: 53, SEQ ID NO: 62, SEQ ID NO: 71, SEQ ID NO: 80, SEQ ID NO: 89, SEQ ID NO: 98, and SEQ ID NO: 101;

[0203] a peptide linker having the amino acid sequence depicted in SEQ ID NO: 8;

[0204] a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 158, SEQ ID NO: 168, SEQ ID NO: 178, SEQ ID NO: 188, SEQ ID NO: 198, SEQ ID NO: 208, SEQ ID NO: 218, and SEQ ID NO: 228 and a serine residue at the C-terminus;

[0205] a polypeptide having the amino acid sequence depicted in SEQ ID NO: 140; and

[0206] a polypeptide comprising in the following order starting from the N-terminus:

[0207] a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 157, SEQ ID NO: 167, SEQ ID NO: 177, SEQ ID NO: 187, SEQ ID NO: 197, SEQ ID NO: 207, SEQ ID NO: 217, and SEQ ID NO: 227;

[0208] a peptide linker having the amino acid sequence depicted in SEQ ID NO: 8;

[0209] a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 18, SEQ ID NO: 27, SEQ ID NO: 36, SEQ ID NO: 45, SEQ ID NO: 54, SEQ ID NO: 63, SEQ ID NO: 72, SEQ ID NO: 81, SEQ ID NO: 90, SEQ ID NO: 99, and SEQ ID NO: 102 and a serine residue at the C-terminus;

- [0210] a polypeptide having the amino acid sequence depicted in SEQ ID NO: 141;
- [0211] (e) a polypeptide comprising in the following order starting from the N-terminus:
- [0212] a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 17, SEQ ID NO: 26, SEQ ID NO: 35, SEQ ID NO: 44, SEQ ID NO: 53, SEQ ID NO: 62, SEQ ID NO: 71, SEQ ID NO: 80, SEQ ID NO: 89, SEQ ID NO: 98, and SEQ ID NO: 101;
- [0213] a peptide linker having the amino acid sequence depicted in SEQ ID NO: 8;
- [0214] a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 158, SEQ ID NO: 168, SEQ ID NO: 178, SEQ ID NO: 188, SEQ ID NO: 198, SEQ ID NO: 208, SEQ ID NO: 218, and SEQ ID NO: 228;
- [0215] a polypeptide having the amino acid sequence depicted in SEQ ID NO: 142; and
- [0216] a polypeptide comprising in the following order starting from the N-terminus:
- [0217] a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 157, SEQ ID NO: 167, SEQ ID NO: 177, SEQ ID NO: 187, SEQ ID NO: 197, SEQ ID NO: 207, SEQ ID NO: 217, and SEQ ID NO: 227;
- [0218] a peptide linker having an amino acid sequence depicted in SEQ ID NO: 8;
- [0219] a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 18, SEQ ID NO: 27, SEQ ID NO: 36, SEQ ID NO: 45, SEQ ID NO: 54, SEQ ID NO: 63, SEQ ID NO: 72, SEQ ID NO: 81, SEQ ID NO: 90, SEQ ID NO: 99, and SEQ ID NO: 102 and a serine residue at the C-terminus;
- [0220] a polypeptide having the amino acid sequence depicted in SEQ ID NO: 143;
- [0221] (f) a polypeptide comprising in the following order starting from the N-terminus:
- [0222] a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 159, SEQ ID NO: 169, SEQ ID NO: 179, SEQ ID NO: 189, SEQ ID NO: 199, SEQ ID NO: 209, SEQ ID NO: 219, and SEQ ID NO: 229;
- [0223] a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 1-9; and
- [0224] a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 19, SEQ ID NO: 28, SEQ ID NO: 37, SEQ ID NO: 46, SEQ ID NO: 55, SEQ ID NO: 64, SEQ ID NO: 73, SEQ ID NO: 82, SEQ ID NO: 91, SEQ ID NO: 100, and SEQ ID NO: 103; and
- [0225] a polypeptide having the amino acid sequence depicted in SEQ ID NO: 144; and
- [0226] a polypeptide having the amino acid sequence depicted in SEQ ID NO: 145;
- [0227] (g) a polypeptide comprising in the following order starting from the N-terminus:
- [0228] a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 159, SEQ ID NO: 169, SEQ ID NO: 179, SEQ ID NO: 189, SEQ ID NO: 199, SEQ ID NO: 209, SEQ ID NO: 219, and SEQ ID NO: 229; and
- [0229] a polypeptide having the amino acid sequence depicted in SEQ ID NO: 146; and
- [0230] a polypeptide comprising in the following order starting from the N-terminus:
- [0231] a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 19, SEQ ID NO: 28, SEQ ID NO: 37, SEQ ID NO: 46, SEQ ID NO: 55, SEQ ID NO: 64, SEQ ID NO: 73, SEQ ID NO: 82, SEQ ID NO: 91, SEQ ID NO: 100, and SEQ ID NO: 103; and
- [0232] a polypeptide having the amino acid sequence depicted in SEQ ID NO: 147;
- [0233] (h) a polypeptide comprising in the following order starting from the N-terminus:
- [0234] a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 159, SEQ ID NO: 169, SEQ ID NO: 179, SEQ ID NO: 189, SEQ ID NO: 199, SEQ ID NO: 209, SEQ ID NO: 219, and SEQ ID NO: 229; and
- [0235] a polypeptide having the amino acid sequence depicted in SEQ ID NO: 148; and
- [0236] a polypeptide comprising in the following order starting from the N-terminus:
- [0237] a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 19, SEQ ID NO: 28, SEQ ID NO: 37, SEQ ID NO: 46, SEQ ID NO: 55, SEQ ID NO: 64, SEQ ID NO: 73, SEQ ID NO: 82, SEQ ID NO: 91, SEQ ID NO: 100, and SEQ ID NO: 103; and
- [0238] a polypeptide having the amino acid sequence depicted in SEQ ID NO: 149; or
- [0239] (i) a polypeptide comprising in the following order starting from the N-terminus:
- [0240] a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 159, SEQ ID NO: 169, SEQ ID NO: 179, SEQ ID NO: 189, SEQ ID NO: 199, SEQ ID NO: 209, SEQ ID NO: 219, and SEQ ID NO: 229;
- [0241] a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 1-9; and
- [0242] a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 19, SEQ ID NO: 28, SEQ ID NO: 37, SEQ ID NO: 46, SEQ ID NO: 55, SEQ ID NO: 64, SEQ ID NO: 73, SEQ ID NO: 82, SEQ ID NO: 91, SEQ ID NO: 100, and SEQ ID NO: 103; and
- [0243] a polypeptide having the amino acid sequence depicted in SEQ ID NO: 150.
- [0244] As described above, several preferred antibody constructs of the invention are modified by fusion with another moiety such as albumin or albumin variants. If these fusion constructs are characterized for their properties, in particular target affinity or cytotoxic activity, the skilled person will be aware that similar fusion constructs or unmodified bispecific antibody constructs can be expected to have similar (or even better) properties. For example, if a bispecific antibody construct fused with albumin has an appreciable or desirable cytotoxic activity or target affinity, it can be expected that the same/similar or even a higher cytotoxic activity/target affinity will be observed for the construct w/o albumin.
- [0245] According to another preferred embodiment, the bispecific antibody construct of the invention comprises (in

addition to the two binding domains) a third domain which comprises two polypeptide monomers, each comprising a hinge, a CH2 and a CH3 domain, wherein said two polypeptides (or polypeptide monomers) are fused to each other via a peptide linker. Preferably, said third domain comprises in an N- to C-terminal order: hinge-CH2-CH3-linker-hinge-CH2-CH3. Preferred amino acid sequences for said third domain are depicted in SEQ ID NOs: 260-267. Each of said two polypeptide monomers preferably has an amino acid sequence that is selected from the group consisting of SEQ ID NOs: 252-259, or that is at least 90% identical to those sequences. In another preferred embodiment, the first and second binding domains of the bispecific antibody construct of the invention are fused to the third domain via a peptide linker which is for example selected from the group consisting of SEQ ID NOs: 1, 2, 3, 4, 5, 6, 7, 8 and 9.

[0246] In line with the present invention, a “hinge” is an IgG hinge region. This region can be identified by analogy using the Kabat numbering, see Kabat positions 223-243. In line with the above, the minimal requirement for a “hinge” are the amino acid residues corresponding to the IgG₁ sequence stretch of D231 to P243 according to the Kabat numbering. The terms CH2 and CH3 refer to the immunoglobulin heavy chain constant regions 2 and 3. These regions can as well be identified by analogy using the Kabat numbering, see Kabat positions 244-360 for CH2 and Kabat positions 361-478 for CH3. It is understood that there is some variation between the immunoglobulins in terms of their IgG₁ Fc region, IgG₂ Fc region, IgG₃ Fc region, IgG₄ Fc region, IgM Fc region, IgA Fc region, IgD Fc region and IgE Fc region (see, e.g., Padlan, Molecular Immunology, 31(3), 169-217 (1993)). The term Fc monomer refers to the last two heavy chain constant regions of IgA, IgD, and IgG, and the last three heavy chain constant regions of IgE and IgM. The Fc monomer can also include the flexible hinge N-terminal to these domains. For IgA and IgM, the Fc monomer may include the J chain. For IgG, the Fc portion comprises immunoglobulin domains CH2 and CH3 and the hinge between the first two domains and CH2. Although the boundaries of the Fc portion of an immunoglobulin may vary, an example for a human IgG heavy chain Fc portion comprising a functional hinge, CH2 and CH3 domain can be defined e.g. to comprise residues D231 (of the hinge domain) to P476 (of the C-terminus of the CH3 domain), or D231 to L476, respectively, for IgG₄, wherein the numbering is according to Kabat.

[0247] The antibody construct of the invention may hence comprise in an N- to C-terminal order:

- [0248]** (a) the first binding domain;
- [0249]** (b) a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 2, 8 and 9;
- [0250]** (c) the second binding domain;
- [0251]** (d) a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 1, 2, 4, 5, 6, 8 and 9;
- [0252]** (e) the first polypeptide monomer of the third domain (comprising a hinge, a CH2 and a CH3 domain);
- [0253]** (f) a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 301, 302, 303 and 304; and

[0254] (g) the second polypeptide monomer of the third domain (comprising a hinge, a CH2 and a CH3 domain).

[0255] It is also preferred that the antibody construct of the invention comprises in an N- to C-terminal order:

[0256] the first binding domain having an amino acid sequence selected from the group consisting of SEQ ID NO: 159, SEQ ID NO: 169, SEQ ID NO: 179, SEQ ID NO: 189, SEQ ID NO: 199, SEQ ID NO: 209, SEQ ID NO: 219, and SEQ ID NO: 229;

[0257] a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 2, 8 and 9;

[0258] the second binding domain having an amino acid sequence selected from the group consisting of SEQ ID NO: 19, SEQ ID NO: 28, SEQ ID NO: 37, SEQ ID NO: 46, SEQ ID NO: 55, SEQ ID NO: 64, SEQ ID NO: 73, SEQ ID NO: 82, SEQ ID NO: 91, SEQ ID NO: 100, and SEQ ID NO: 103 (see also SEQ ID NOs: 23, 25, 41, 43, 59, 61, 77, 79, 95, 97, 113, 115, 131, 133, 149, 151, 167, 169, 185 or 187 of WO 2008/119567);

[0259] a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 1, 2, 4, 5, 6, 8 and 9; and

[0260] the third domain having an amino acid sequence selected from the group consisting of SEQ ID NOs: 260-267.

[0261] Hence, in a preferred embodiment, the antibody construct of the present invention comprises or consists of a polypeptide selected from the group consisting of those depicted in SEQ ID NOs: 268 to 299.

[0262] It is also preferably in one embodiment of the invention that the antibody construct of the present invention comprises or consists of a polypeptide selected from the group consisting of those depicted in SEQ ID NOs: 276 to 287.

[0263] Also in a preferred embodiment of the invention that the antibody construct of the present invention comprises or consists of a polypeptide selected from the group consisting of those depicted in SEQ ID NOs: 276, 280 and 284, more preferably comprises or consists of a polypeptide as depicted in SEQ ID NO: 280.

[0264] Covalent modifications of the antibody constructs are also included within the scope of this invention, and are generally, but not always, done post-translationally. For example, several types of covalent modifications of the antibody construct are introduced into the molecule by reacting specific amino acid residues of the antibody construct with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C-terminal residues.

[0265] Cysteiny residues most commonly are reacted with α -haloacetates (and corresponding amines), such as chloroacetic acid or chloroacetamide, to give carboxymethyl or carboxyamidomethyl derivatives. Cysteiny residues also are derivatized by reaction with bromotrifluoroacetone, α -bromo- β -(5-imidazolyl)propionic acid, chloroacetyl phosphate, N-alkylmaleimides, 3-nitro-2-pyridyl disulfide, methyl 2-pyridyl disulfide, p-chloromercuribenzoate, 2-chloromercuri-4-nitrophenol, or chloro-7-nitrobenzo-2-oxa-1,3-diazole.

[0266] Histidyl residues are derivatized by reaction with diethylpyrocarbonate at pH 5.5-7.0 because this agent is relatively specific for the histidyl side chain. Para-brom-

phenacyl bromide also is useful; the reaction is preferably performed in 0.1 M sodium cacodylate at pH 6.0. Lysiny and amino terminal residues are reacted with succinic or other carboxylic acid anhydrides. Derivatization with these agents has the effect of reversing the charge of the lysiny residues. Other suitable reagents for derivatizing alpha-amino-containing residues include imidoesters such as methyl picolinimidate; pyridoxal phosphate; pyridoxal; chloroborohydride; trinitrobenzenesulfonic acid; O-methylisourea; 2,4-pentanedione; and transaminase-catalyzed reaction with glyoxylate.

[0267] Arginy residues are modified by reaction with one or several conventional reagents, among them phenylglyoxal, 2,3-butanedione, 1,2-cyclohexanedione, and ninhydrin. Derivatization of arginine residues requires that the reaction be performed in alkaline conditions because of the high pKa of the guanidine functional group. Furthermore, these reagents may react with the groups of lysine as well as the arginine epsilon-amino group.

[0268] The specific modification of tyrosyl residues may be made, with particular interest in introducing spectral labels into tyrosyl residues by reaction with aromatic diazonium compounds or tetranitromethane. Most commonly, N-acetylimidazole and tetranitromethane are used to form O-acetyl tyrosyl species and 3-nitro derivatives, respectively. Tyrosyl residues are iodinated using ^{125}I or ^{131}I to prepare labeled proteins for use in radioimmunoassay, the chloramine T method described above being suitable.

[0269] Carboxyl side groups (aspartyl or glutamyl) are selectively modified by reaction with carbodiimides ($\text{R}'\text{—N}=\text{C}=\text{N—R}'$), where R and R' are optionally different alkyl groups, such as 1-cyclohexyl-3-(2-morpholinyl-4-ethyl) carbodiimide or 1-ethyl-3-(4-azonia-4,4-dimethylpentyl) carbodiimide. Furthermore, aspartyl and glutamyl residues are converted to asparaginy and glutaminyl residues by reaction with ammonium ions.

[0270] Derivatization with bifunctional agents is useful for crosslinking the antibody constructs of the present invention to a water-insoluble support matrix or surface for use in a variety of methods. Commonly used crosslinking agents include, e.g., 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidylpropionate), and bifunctional maleimides such as bis-N-maleimido-1,8-octane. Derivatizing agents such as methyl-3-[(p-azidophenyl)dithio]propioimidate yield photo-activatable intermediates that are capable of forming crosslinks in the presence of light. Alternatively, reactive water-insoluble matrices such as cyanogen bromide-activated carbohydrates and the reactive substrates as described in U.S. Pat. Nos. 3,969,287; 3,691,016; 4,195,128; 4,247,642; 4,229,537; and 4,330,440 are employed for protein immobilization.

[0271] Glutaminyl and asparaginy residues are frequently deamidated to the corresponding glutamyl and aspartyl residues, respectively. Alternatively, these residues are deamidated under mildly acidic conditions. Either form of these residues falls within the scope of this invention.

[0272] Other modifications include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the α -amino groups of lysine, arginine, and histidine side chains (T. E. Creighton, *Proteins: Structure and Molecular Properties*, W. H. Free-

man & Co., San Francisco, 1983, pp. 79-86), acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

[0273] Another type of covalent modification of the antibody constructs included within the scope of this invention comprises altering the glycosylation pattern of the protein. As is known in the art, glycosylation patterns can depend on both the sequence of the protein (e.g., the presence or absence of particular glycosylation amino acid residues, discussed below), or the host cell or organism in which the protein is produced. Particular expression systems are discussed below.

[0274] Glycosylation of polypeptides is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tri-peptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tri-peptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-acetylgalactosamine, galactose, or xylose, to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

[0275] Addition of glycosylation sites to the antibody construct is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tri-peptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the starting sequence (for O-linked glycosylation sites). For ease, the amino acid sequence of an antibody construct is preferably altered through changes at the DNA level, particularly by mutating the DNA encoding the polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids.

[0276] Another means of increasing the number of carbohydrate moieties on the antibody construct is by chemical or enzymatic coupling of glycosides to the protein. These procedures are advantageous in that they do not require production of the protein in a host cell that has glycosylation capabilities for N- and O-linked glycosylation. Depending on the coupling mode used, the sugar(s) may be attached to (a) arginine and histidine, (b) free carboxyl groups, (c) free sulfhydryl groups such as those of cysteine, (d) free hydroxyl groups such as those of serine, threonine, or hydroxyproline, (e) aromatic residues such as those of phenylalanine, tyrosine, or tryptophan, or (f) the amide group of glutamine. These methods are described in WO 87/05330, and in Aplin and Wriston, 1981, *CRC Crit. Rev. Biochem.*, pp. 259-306.

[0277] Removal of carbohydrate moieties present on the starting antibody construct may be accomplished chemically or enzymatically. Chemical deglycosylation requires exposure of the protein to the compound trifluoromethanesulfonic acid, or an equivalent compound. This treatment results in the cleavage of most or all sugars except the linking sugar (N-acetylglucosamine or N-acetylgalactosamine), while leaving the polypeptide intact. Chemical deglycosylation is described by Hakimuddin et al., 1987, *Arch. Biochem. Biophys.* 259:52 and by Edge et al., 1981, *Anal. Biochem.* 118:131. Enzymatic cleavage of carbohydrate moieties on

polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura et al., 1987, *Meth. Enzymol.* 138:350. Glycosylation at potential glycosylation sites may be prevented by the use of the compound tunicamycin as described by Duskin et al., 1982, *J. Biol. Chem.* 257:3105. Tunicamycin blocks the formation of protein-N-glycoside linkages.

[0278] Other modifications of the antibody construct are also contemplated herein. For example, another type of covalent modification of the antibody construct comprises linking the antibody construct to various non-proteinaceous polymers, including, but not limited to, various polyols such as polyethylene glycol, polypropylene glycol, polyoxalkylenes, or copolymers of polyethylene glycol and polypropylene glycol, in the manner set forth in U.S. Pat. Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337. In addition, as is known in the art, amino acid substitutions may be made in various positions within the antibody construct, e.g. in order to facilitate the addition of polymers such as PEG.

[0279] In some embodiments, the covalent modification of the antibody constructs of the invention comprises the addition of one or more labels. The labelling group may be coupled to the antibody construct via spacer arms of various lengths to reduce potential steric hindrance.

[0280] Various methods for labelling proteins are known in the art and can be used in performing the present invention. The term "label" or "labelling group" refers to any detectable label. In general, labels fall into a variety of classes, depending on the assay in which they are to be detected—the following examples include, but are not limited to:

[0281] a) isotopic labels, which may be radioactive or heavy isotopes, such as radioisotopes or radionuclides (e.g., ^3H , ^{14}C , ^{15}N , ^{35}S , ^{89}Zr , ^{90}Y , ^{99}Tc , ^{111}In , ^{125}I , ^{131}I)

[0282] b) magnetic labels (e.g., magnetic particles)

[0283] c) redox active moieties

[0284] d) optical dyes (including, but not limited to, chromophores, phosphors and fluorophores) such as fluorescent groups (e.g., FITC, rhodamine, lanthanide phosphors), chemiluminescent groups, and fluorophores which can be either "small molecule" fluors or proteinaceous fluors

[0285] e) enzymatic groups (e.g. horseradish peroxidase, β -galactosidase, luciferase, alkaline phosphatase)

[0286] f) biotinylated groups

[0287] g) predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags, etc.)

[0288] By "fluorescent label" is meant any molecule that may be detected via its inherent fluorescent properties. Suitable fluorescent labels include, but are not limited to, fluorescein, rhodamine, tetramethylrhodamine, eosin, erythrosin, coumarin, methyl-coumarins, pyrene, Malachite green, stilbene, Lucifer Yellow, Cascade BlueJ, Texas Red, IAE-DANS, EDANS, BODIPY FL, LC Red 640, Cy 5, Cy 5.5, LC Red 705, Oregon green, the Alexa-Fluor dyes (Alexa Fluor 350, Alexa Fluor 430, Alexa Fluor 488, Alexa Fluor 546, Alexa Fluor 568, Alexa Fluor 594, Alexa Fluor 633, Alexa Fluor 660, Alexa Fluor 680), Cascade Blue, Cascade Yellow and R-phycoerythrin (PE) (Molecular Probes, Eugene, OR), FITC, Rhodamine, and Texas Red (Pierce, Rockford, IL), Cy5, Cy5.5, Cy7 (Amersham Life Science,

Pittsburgh, PA). Suitable optical dyes, including fluorophores, are described in *Molecular Probes Handbook* by Richard P. Haugland.

[0289] Suitable proteinaceous fluorescent labels also include, but are not limited to, green fluorescent protein, including a *Renilla*, *Ptilosarcus*, or *Aequorea* species of GFP (Chalfie et al., 1994, *Science* 263:802-805), EGFP (Clontech Laboratories, Inc., Genbank Accession Number U55762), blue fluorescent protein (BFP, Quantum Biotechnologies, Inc. 1801 de Maisonneuve Blvd. West, 8th Floor, Montreal, Quebec, Canada H3H 1J9; Stauber, 1998, *Biotechniques* 24:462-471; Heim et al., 1996, *Curr. Biol.* 6:178-182), enhanced yellow fluorescent protein (EYFP, Clontech Laboratories, Inc.), luciferase (Ichiki et al., 1993, *J. Immunol.* 150:5408-5417), P galactosidase (Nolan et al., 1988, *Proc. Natl. Acad. Sci. U.S.A.* 85:2603-2607) and *Renilla* (WO92/15673, WO95/07463, WO98/14605, WO98/26277, WO99/49019, U.S. Pat. Nos. 5,292,658; 5,418,155; 5,683,888; 5,741,668; 5,777,079; 5,804,387; 5,874,304; 5,876,995; 5,925,558).

[0290] Leucine zipper domains are peptides that promote oligomerization of the proteins in which they are found. Leucine zippers were originally identified in several DNA-binding proteins (Landschulz et al., 1988, *Science* 240:1759), and have since been found in a variety of different proteins. Among the known leucine zippers are naturally occurring peptides and derivatives thereof that dimerize or trimerize. Examples of leucine zipper domains suitable for producing soluble oligomeric proteins are described in PCT application WO 94/10308, and the leucine zipper derived from lung surfactant protein D (SPD) described in Hoppe et al., 1994, *FEBS Letters* 344:191. The use of a modified leucine zipper that allows for stable trimerization of a heterologous protein fused thereto is described in Fanslow et al., 1994, *Semin. Immunol.* 6:267-78. In one approach, recombinant fusion proteins comprising a MSLN antibody fragment or derivative fused to a leucine zipper peptide are expressed in suitable host cells, and the soluble oligomeric MSLN antibody fragments or derivatives that form are recovered from the culture supernatant.

[0291] The antibody construct of the invention may also comprise additional domains, which are e.g. helpful in the isolation of the molecule or relate to an adapted pharmacokinetic profile of the molecule. Domains helpful for the isolation of an antibody construct may be selected from peptide motives or secondarily introduced moieties, which can be captured in an isolation method, e.g. an isolation column. Non-limiting embodiments of such additional domains comprise peptide motives known as Myc-tag, HAT-tag, HA-tag, TAP-tag, GST-tag, chitin binding domain (CBD-tag), maltose binding protein (MBP-tag), Flag-tag, Strep-tag and variants thereof (e.g. StrepII-tag) and His-tag. All herein disclosed antibody constructs characterized by the identified CDRs are preferred to comprise a His-tag domain, which is generally known as a repeat of consecutive His residues in the amino acid sequence of a molecule, preferably of five, and more preferably of six His residues (hexahistidine). The His-tag may be located e.g. at the N- or C-terminus of the antibody construct, preferably it is located at the C-terminus. Most preferably, a hexa-histidine tag (HHHHHH) (SEQ ID NO:10) is linked via peptide bond to the C-terminus of the antibody construct according to the invention.

[0292] The first binding domain of the antibody construct of the present invention binds to human MSLN on the surface of a target cell. The preferred amino acid sequence of human MSLN is represented by NOs: 231, 232, and 233. The first binding domain according to the invention hence preferably binds to MSLN when it is expressed by naturally expressing cells or cell lines, and/or by cells or cell lines transformed or (stably/transiently) transfected with MSLN. In a preferred embodiment the first binding domain also binds to MSLN when MSLN is used as a “target” or “ligand” molecule in an in vitro binding assay such as BIAcore or Scatchard. The “target cell” can be any prokaryotic or eukaryotic cell expressing MSLN on its surface; preferably the target cell is a cell that is part of the human or animal body, such as an ovarian cancer cell, pancreatic cancer cell, mesothelioma cell, lung cancer cell, gastric cancer cell and triple negative breast cancer cell.

[0293] The affinity of the first binding domain for human MSLN is preferably ≤ 20 nM, more preferably ≤ 10 nM, even more preferably ≤ 5 nM, even more preferably ≤ 2 nM, even more preferably ≤ 1 nM, even more preferably ≤ 0.6 nM, even more preferably ≤ 0.5 nM, and most preferably ≤ 0.4 nM. The affinity can be measured for example in a BIAcore assay or in a Scatchard assay, e.g. as described in the Examples. Other methods of determining the affinity are also well-known to the skilled person; see e.g. appended Example 3.

[0294] Amino acid sequence modifications of the antibody constructs described herein are also contemplated. For example, it may be desirable to improve the binding affinity and/or other biological properties of the antibody construct. Amino acid sequence variants of the antibody constructs are prepared by introducing appropriate nucleotide changes into the antibody constructs nucleic acid, or by peptide synthesis. All of the below described amino acid sequence modifications should result in an antibody construct which still retains the desired biological activity (binding to MSLN and to CD3) of the unmodified parental molecule.

[0295] The term “amino acid” or “amino acid residue” typically refers to an amino acid having its art recognized definition such as an amino acid selected from the group consisting of: alanine (Ala or A); arginine (Arg or R); asparagine (Asn or N); aspartic acid (Asp or D); cysteine (Cys or C); glutamine (Gln or Q); glutamic acid (Glu or E); glycine (Gly or G); histidine (His or H); isoleucine (Ile or I); leucine (Leu or L); lysine (Lys or K); methionine (Met or M); phenylalanine (Phe or F); proline (Pro or P); serine (Ser or S); threonine (Thr or T); tryptophan (Trp or W); tyrosine (Tyr or Y); and valine (Val or V), although modified, synthetic, or rare amino acids may be used as desired. Generally, amino acids can be grouped as having a nonpolar side chain (e.g., Ala, Cys, Ile, Leu, Met, Phe, Pro, Val); a negatively charged side chain (e.g., Asp, Glu); a positively charged sidechain (e.g., Arg, His, Lys); or an uncharged polar side chain (e.g., Asn, Cys, Gln, Gly, His, Met, Phe, Ser, Thr, Trp, and Tyr).

[0296] Amino acid modifications include, for example, deletions from, and/or insertions into, and/or substitutions of, residues within the amino acid sequences of the antibody constructs. Any combination of deletion, insertion, and substitution is made to arrive at the final construct, provided that the final construct possesses the desired characteristics. The amino acid changes also may alter post-translational

processes of the antibody constructs, such as changing the number or position of glycosylation sites.

[0297] For example, 1, 2, 3, 4, 5, or 6 amino acids may be inserted or deleted in each of the CDRs (of course, dependent on their length), while 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or 25 amino acids may be inserted or deleted in each of the FRs. Preferably, amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 residues to polypeptides containing a hundred or more residues, as well as intra-sequence insertions of single or multiple amino acid residues. An insertional variant of the antibody construct of the invention includes the fusion to the N-terminus or to the C-terminus of the antibody construct of an enzyme or the fusion to a polypeptide which increases the serum half-life of the antibody construct.

[0298] The sites of greatest interest for substitutional mutagenesis include the CDRs of the heavy and/or light chain, in particular the hypervariable regions, but FR alterations in the heavy and/or light chain are also contemplated. The substitutions are preferably conservative substitutions as described herein. Preferably, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 amino acids may be substituted in a CDR, while 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or 25 amino acids may be substituted in the framework regions (FRs), depending on the length of the CDR or FR. For example, if a CDR sequence encompasses 6 amino acids, it is envisaged that one, two or three of these amino acids are substituted. Similarly, if a CDR sequence encompasses 15 amino acids it is envisaged that one, two, three, four, five or six of these amino acids are substituted.

[0299] A useful method for identification of certain residues or regions of the antibody constructs that are preferred locations for mutagenesis is called “alanine scanning mutagenesis” as described by Cunningham and Wells in *Science*, 244: 1081-1085 (1989). Here, a residue or group of target residues within the antibody construct is/are identified (e.g. charged residues such as arg, asp, his, lys, and glu) and replaced by a neutral or negatively charged amino acid (most preferably alanine or polyalanine) to affect the interaction of the amino acids with the epitope.

[0300] Those amino acid locations demonstrating functional sensitivity to the substitutions are then refined by introducing further or other variants at, or for, the sites of substitution. Thus, while the site or region for introducing an amino acid sequence variation is predetermined, the nature of the mutation per se needs not to be predetermined. For example, to analyze or optimize the performance of a mutation at a given site, alanine scanning or random mutagenesis may be conducted at a target codon or region, and the expressed antibody construct variants are screened for the optimal combination of desired activity. Techniques for making substitution mutations at predetermined sites in the DNA having a known sequence are well known, for example, M13 primer mutagenesis and PCR mutagenesis. Screening of the mutants is done using assays of antigen binding activities, such as MSLN or CD3 binding.

[0301] Generally, if amino acids are substituted in one or more or all of the CDRs of the heavy and/or light chain, it is preferred that the then-obtained “substituted” sequence is at least 60% or 65%, more preferably 70% or 75%, even more preferably 80% or 85%, and particularly preferably 90% or 95% identical to the “original” CDR sequence. This

means that it is dependent of the length of the CDR to which degree it is identical to the “substituted” sequence.

[0302] For example, a CDR having 5 amino acids is preferably 80% identical to its substituted sequence in order to have at least one amino acid substituted. Accordingly, the CDRs of the antibody construct may have different degrees of identity to their substituted sequences, e.g., CDRL1 may have 80%, while CDRL3 may have 90%.

[0303] Preferred substitutions (or replacements) are conservative substitutions. However, any substitution (including non-conservative substitution or one or more from the “exemplary substitutions” listed in Table 1, below) is envisaged as long as the antibody construct retains its capability to bind to MSLN via the first binding domain and to CD3 or CD3 epsilon via the second binding domain and/or its CDRs have an identity to the then substituted sequence (at least 60% or 65%, more preferably 70% or 75%, even more preferably 80% or 85%, and particularly preferably 90% or 95% identical to the “original” CDR sequence).

[0304] Conservative substitutions are shown in Table 1 under the heading of “preferred substitutions”. If such substitutions result in a change in biological activity, then more substantial changes, denominated “exemplary substitutions” in Table 1, or as further described below in reference to amino acid classes, may be introduced and the products screened for a desired characteristic.

TABLE 1

Amino acid substitutions		
Original	Exemplary Substitutions	Preferred Substitutions
Ala (A)	val, leu, ile	val
Arg (R)	lys, gln, asn	lys
Asn (N)	gln, his, asp, lys, arg	gln
Asp (D)	glu, asn	glu
Cys (C)	ser, ala	ser
Gln (Q)	asn, glu	asn
Glu (E)	asp, gln	asp
Gly (G)	Ala	ala
His (H)	asn, gln, lys, arg	arg
Ile (I)	leu, val, met, ala, phe	leu
Leu (L)	norleucine, ile, val, met, ala	ile
Lys (K)	arg, gln, asn	arg
Met (M)	leu, phe, ile	leu
Phe (F)	leu, val, ile, ala, tyr	tyr
Pro (P)	Ala	ala
Ser (S)	Thr	thr
Thr (T)	Ser	ser
Trp (W)	tyr, phe	tyr
Tyr (Y)	trp, phe, thr, ser	phe
Val (V)	ile, leu, met, phe, ala	leu

[0305] Substantial modifications in the biological properties of the antibody construct of the present invention are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties: (1) hydrophobic: norleucine, met, ala, val, leu, ile; (2) neutral hydrophilic: cys, ser, thr; (3) acidic: asp, glu; (4) basic: asn, gln, his, lys, arg; (5) residues that influence chain orientation: gly, pro; and (6) aromatic: trp, tyr, phe.

[0306] Non-conservative substitutions will entail exchanging a member of one of these classes for another

class. Any cysteine residue not involved in maintaining the proper conformation of the antibody construct may be substituted, generally with serine, to improve the oxidative stability of the molecule and prevent aberrant crosslinking. Conversely, cysteine bond(s) may be added to the antibody to improve its stability (particularly where the antibody is an antibody fragment such as an Fv fragment).

[0307] For amino acid sequences, sequence identity and/or similarity is determined by using standard techniques known in the art, including, but not limited to, the local sequence identity algorithm of Smith and Waterman, 1981, *Adv. Appl. Math.* 2:482, the sequence identity alignment algorithm of Needleman and Wunsch, 1970, *J. Mol. Biol.* 48:443, the search for similarity method of Pearson and Lipman, 1988, *Proc. Nat. Acad. Sci. U.S.A.* 85:2444, computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Drive, Madison, Wis.), the Best Fit sequence program described by Devereux et al., 1984, *Nucl. Acid Res.* 12:387-395, preferably using the default settings, or by inspection. Preferably, percent identity is calculated by FastDB based upon the following parameters: mismatch penalty of 1; gap penalty of 1; gap size penalty of 0.33; and joining penalty of 30, “Current Methods in Sequence Comparison and Analysis,” *Macromolecule Sequencing and Synthesis, Selected Methods and Applications*, pp 127-149 (1988), Alan R. Liss, Inc.

[0308] An example of a useful algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments. It can also plot a tree showing the clustering relationships used to create the alignment. PILEUP uses a simplification of the progressive alignment method of Feng & Doolittle, 1987, *J. Mol. Evol.* 35:351-360; the method is similar to that described by Higgins and Sharp, 1989, *CABIOS* 5:151-153. Useful PILEUP parameters including a default gap weight of 3.00, a default gap length weight of 0.10, and weighted end gaps.

[0309] Another example of a useful algorithm is the BLAST algorithm, described in: Altschul et al., 1990, *J. Mol. Biol.* 215:403-410; Altschul et al., 1997, *Nucleic Acids Res.* 25:3389-3402; and Karin et al., 1993, *Proc. Natl. Acad. Sci. U.S.A.* 90:5873-5787. A particularly useful BLAST program is the WU-BLAST-2 program which was obtained from Altschul et al., 1996, *Methods in Enzymology* 266: 460-480. WU-BLAST-2 uses several search parameters, most of which are set to the default values. The adjustable parameters are set with the following values: overlap span=1, overlap fraction=0.125, word threshold (T)=II. The HSP S and HSP S2 parameters are dynamic values and are established by the program itself depending upon the composition of the particular sequence and composition of the particular database against which the sequence of interest is being searched; however, the values may be adjusted to increase sensitivity.

[0310] An additional useful algorithm is gapped BLAST as reported by Altschul et al., 1993, *Nucl. Acids Res.* 25:3389-3402. Gapped BLAST uses BLOSUM-62 substitution scores; threshold T parameter set to 9; the two-hit method to trigger ungapped extensions, charges gap lengths of k a cost of 10+k; Xu set to 16, and Xg set to 40 for database search stage and to 67 for the output stage of the algorithms. Gapped alignments are triggered by a score corresponding to about 22 bits.

[0311] Generally, the amino acid homology, similarity, or identity between individual variant CDRs are at least 60% to the sequences depicted herein, and more typically with preferably increasing homologies or identities of at least 65% or 70%, more preferably at least 75% or 80%, even more preferably at least 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, and almost 100%. In a similar manner, “percent (%) nucleic acid sequence identity” with respect to the nucleic acid sequence of the binding proteins identified herein is defined as the percentage of nucleotide residues in a candidate sequence that are identical with the nucleotide residues in the coding sequence of the antibody construct. A specific method utilizes the BLASTN module of WU-BLAST-2 set to the default parameters, with overlap span and overlap fraction set to 1 and 0.125, respectively.

[0312] Generally, the nucleic acid sequence homology, similarity, or identity between the nucleotide sequences encoding individual variant CDRs and the nucleotide sequences depicted herein are at least 60%, and more typically with preferably increasing homologies or identities of at least 65%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%, and almost 100%. Thus, a “variant CDR” is one with the specified homology, similarity, or identity to the parent CDR of the invention, and shares biological function, including, but not limited to, at least 60%, 65%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% of the specificity and/or activity of the parent CDR.

[0313] In one embodiment, the percentage of identity to human germline of the antibody constructs according to the invention is $\geq 70\%$ or $\geq 75\%$, more preferably $\geq 80\%$ or $\geq 85\%$, even more preferably $\geq 90\%$, and most preferably $\geq 91\%$, $\geq 92\%$, $\geq 93\%$, $\geq 94\%$, $\geq 95\%$ or even $\geq 96\%$. Identity to human antibody germline gene products is thought to be an important feature to reduce the risk of therapeutic proteins to elicit an immune response against the drug in the patient during treatment. Hwang & Foote (“Immunogenicity of engineered antibodies”; Methods 36 (2005) 3-10) demonstrate that the reduction of non-human portions of drug antibody constructs leads to a decrease of risk to induce anti-drug antibodies in the patients during treatment. By comparing an exhaustive number of clinically evaluated antibody drugs and the respective immunogenicity data, the trend is shown that humanization of the V-regions of antibodies makes the protein less immunogenic (average 5.1% of patients) than antibodies carrying unaltered non-human V regions (average 23.59% of patients). A higher degree of identity to human sequences is hence desirable for V-region based protein therapeutics in the form of antibody constructs. For this purpose of determining the germline identity, the V-regions of VL can be aligned with the amino acid sequences of human germline V segments and J segments (<http://vbase.mrc-cpe.cam.ac.uk/>) using Vector NTI software and the amino acid sequence calculated by dividing the identical amino acid residues by the total number of amino acid residues of the VL in percent. The same can be for the VH segments (<http://vbase.mrc-cpe.cam.ac.uk/>) with the exception that the VH CDR3 may be excluded due to its high diversity and a lack of existing human germline VH CDR3 alignment partners. Recombinant techniques can then be used to increase sequence identity to human antibody germline genes.

[0314] In a further embodiment, the bispecific antibody constructs of the present invention exhibit high monomer yields under standard research scale conditions, e.g., in a standard two-step purification process. Preferably the monomer yield of the antibody constructs according to the invention is ≥ 0.25 mg/L supernatant, more preferably ≥ 0.5 mg/L, even more preferably ≥ 1 mg/L, and most preferably ≥ 3 mg/L supernatant.

[0315] Likewise, the yield of the dimeric antibody construct isoforms and hence the monomer percentage (i.e., monomer:(monomer+dimer)) of the antibody constructs can be determined.

[0316] The productivity of monomeric and dimeric antibody constructs and the calculated monomer percentage can e.g. be obtained in the SEC purification step of culture supernatant from standardized research-scale production in roller bottles. In one embodiment, the monomer percentage of the antibody constructs is $\geq 80\%$, more preferably $\geq 85\%$, even more preferably $\geq 90\%$, and most preferably $\geq 95\%$.

[0317] In one embodiment, the antibody constructs have a preferred plasma stability (ratio of EC50 with plasma to EC50 w/o plasma) of ≤ 5 or ≤ 4 , more preferably ≤ 3.5 or ≤ 3 , even more preferably ≤ 2.5 or ≤ 2 , and most preferably ≤ 1.5 or ≤ 1 . The plasma stability of an antibody construct can be tested by incubation of the construct in human plasma at 37° C. for 24 hours followed by EC50 determination in a 51-chromium release cytotoxicity assay. The effector cells in the cytotoxicity assay can be stimulated enriched human CD8 positive T cells. Target cells can e.g. be CHO cells transfected with human MSLN. The effector to target cell (E:T) ratio can be chosen as 10:1. The human plasma pool used for this purpose is derived from the blood of healthy donors collected by EDTA coated syringes. Cellular components are removed by centrifugation and the upper plasma phase is collected and subsequently pooled. As control, antibody constructs are diluted immediately prior to the cytotoxicity assay in RPMI-1640 medium. The plasma stability is calculated as ratio of EC50 (after plasma incubation) to EC50 (control). See Example 6.

[0318] It is furthermore preferred that the monomer to dimer conversion of antibody constructs of the invention is low. The conversion can be measured under different conditions and analyzed by high performance size exclusion chromatography. See Example 9. For example, incubation of the monomeric isoforms of the antibody constructs can be carried out for 7 days at 37° C. and concentrations of e.g. 100 $\mu\text{g/ml}$ or 250 $\mu\text{g/ml}$ in an incubator. Under these conditions, it is preferred that the antibody constructs of the invention show a dimer percentage that is $\leq 5\%$, more preferably $\leq 4\%$, even more preferably $\leq 3\%$, even more preferably $\leq 2.5\%$, even more preferably $\leq 2\%$, even more preferably $\leq 1.5\%$, and most preferably $\leq 1\%$ or $\leq 0.5\%$ or even 0%.

[0319] It is also preferred that the bispecific antibody constructs of the present invention present with very low dimer conversion after a number of freeze/thaw cycles. For example, the antibody construct monomer is adjusted to a concentration of 250 $\mu\text{g/ml}$ e.g. in generic formulation buffer and subjected to three freeze/thaw cycles (freezing at -80°C . for 30 min followed by thawing for 30 min at room temperature), followed by high performance SEC to determine the percentage of initially monomeric antibody construct, which had been converted into dimeric antibody construct. Preferably the dimer percentages of the bispecific

antibody constructs are $\leq 5\%$, more preferably $\leq 4\%$, even more preferably $\leq 3\%$, even more preferably $\leq 2.5\%$, even more preferably $\leq 2\%$, even more preferably $\leq 1.5\%$, and most preferably $\leq 1\%$ or even $\leq 0.5\%$, for example after three freeze/thaw cycles.

[0320] The bispecific antibody constructs of the present invention preferably show a favorable thermostability with aggregation temperatures $\geq 45^\circ\text{C}$. or $\geq 50^\circ\text{C}$., more preferably $\geq 51^\circ\text{C}$., $\geq 52^\circ\text{C}$., $\geq 53^\circ\text{C}$. or $\geq 54^\circ\text{C}$., even more preferably $\geq 56^\circ\text{C}$. or $\geq 57^\circ\text{C}$., and most preferably $\geq 58^\circ\text{C}$. or $\geq 59^\circ\text{C}$. The thermostability parameter can be determined in terms of antibody aggregation temperature as follows: Antibody solution at a concentration 250 $\mu\text{g/ml}$ is transferred into a single use cuvette and placed in a Dynamic Light Scattering (DLS) device. The sample is heated from 40°C . to 70°C . at a heating rate of $0.5^\circ\text{C}/\text{min}$ with constant acquisition of the measured radius. Increase of radius indicating melting of the protein and aggregation is used to calculate the aggregation temperature of the antibody. See Example 10.

[0321] Alternatively, temperature melting curves can be determined by Differential Scanning Calorimetry (DSC) to determine intrinsic biophysical protein stabilities of the antibody constructs. These experiments are performed using a MicroCal LLC (Northampton, MA, U.S.A) VP-DSC device. The energy uptake of a sample containing an antibody construct is recorded from 20°C . to 90°C . compared to a sample containing only the formulation buffer. The antibody constructs are adjusted to a final concentration of 250 $\mu\text{g/ml}$ e.g. in SEC running buffer. For recording of the respective melting curve, the overall sample temperature is increased stepwise. At each temperature T energy uptake of the sample and the formulation buffer reference is recorded. The difference in energy uptake C_p (kcal/mole/ $^\circ\text{C}$.) of the sample minus the reference is plotted against the respective temperature. The melting temperature is defined as the temperature at the first maximum of energy uptake.

[0322] The MSLN xCD3 bispecific antibody constructs of the invention are also envisaged to have a turbidity (as measured by OD340 after concentration of purified monomeric antibody construct to 2.5 mg/ml and over night incubation) of ≤ 0.2 , preferably of ≤ 0.15 , more preferably of ≤ 0.12 , even more preferably of ≤ 0.1 or even ≤ 0.09 , and most preferably of ≤ 0.08 or ≤ 0.07 . See Example 11.

[0323] In a further embodiment the antibody construct according to the invention is stable at acidic pH. The more tolerant the antibody construct behaves at unphysiologic pH such as pH 5.5 (a pH which is required to run e.g. a cation exchange chromatography), the higher is the recovery of the antibody construct eluted from an ion exchange column relative to the total amount of loaded protein. Recovery of the antibody construct from an ion (e.g., cation) exchange column at pH 5.5 is preferably $\geq 30\%$, more preferably $\geq 40\%$, more preferably $\geq 50\%$, even more preferably $\geq 60\%$, even more preferably $\geq 70\%$, even more preferably $\geq 80\%$, even more preferably $\geq 90\%$, even more preferably $\geq 95\%$, and most preferably $\geq 99\%$. See Example 7.

[0324] It is furthermore envisaged that the bispecific antibody constructs of the present invention exhibit therapeutic efficacy or anti-tumor activity. This can e.g. be assessed in a study as disclosed in the following example of an advanced stage human tumor xenograft model:

[0325] On day 1 of the study, 5×10^6 cells of a human MSLN positive cancer cell line (e.g. OVCAR-8) are subcutaneously injected in the right dorsal flank of female NOD/SCID mice. When the mean tumor volume reaches about 100 mm^3 , in vitro expanded human CD3 positive T cells are transplanted into the mice by injection of about 2×10^7 cells into the peritoneal cavity of the animals. Mice of vehicle control group 1 do not receive effector cells and are used as an untransplanted control for comparison with vehicle control group 2 (receiving effector cells) to monitor the impact of T cells alone on tumor growth. The antibody treatment starts when the mean tumor volume reaches about 200 mm^3 . The mean tumor size of each treatment group on the day of treatment start should not be statistically different from any other group (analysis of variance). Mice are treated with 0.5 mg/kg/day of a MSLN xCD3 bispecific antibody construct by intravenous bolus injection for about 15 to 20 days. Tumors are measured by caliper during the study and progress evaluated by intergroup comparison of tumor volumes (TV). The tumor growth inhibition T/C [%] is determined by calculating TV as $\text{T/C \%} = 100 \times (\text{median TV of analyzed group}) / (\text{median TV of control group 2})$.

[0326] The skilled person knows how to modify or adapt certain parameters of this study, such as the number of injected tumor cells, the site of injection, the number of transplanted human T cells, the amount of bispecific antibody constructs to be administered, and the timelines, while still arriving at a meaningful and reproducible result. Preferably, the tumor growth inhibition T/C [%] is ≤ 70 or ≤ 60 , more preferably ≤ 50 or ≤ 40 , even more preferably ≤ 30 or ≤ 20 and most preferably ≤ 10 or ≤ 5 or even ≤ 2.5 .

[0327] The invention further provides a polynucleotide/nucleic acid molecule encoding an antibody construct of the invention.

[0328] A polynucleotide is a biopolymer composed of 13 or more nucleotide monomers covalently bonded in a chain. DNA (such as cDNA) and RNA (such as mRNA) are examples of polynucleotides with distinct biological function. Nucleotides are organic molecules that serve as the monomers or subunits of nucleic acid molecules like DNA or RNA. The nucleic acid molecule or polynucleotide can be double stranded and single stranded, linear and circular. It is preferably comprised in a vector which is preferably comprised in a host cell. Said host cell is, e.g. after transformation or transfection with the vector or the polynucleotide of the invention, capable of expressing the antibody construct. For that purpose the polynucleotide or nucleic acid molecule is operatively linked with control sequences.

[0329] The genetic code is the set of rules by which information encoded within genetic material (nucleic acids) is translated into proteins. Biological decoding in living cells is accomplished by the ribosome which links amino acids in an order specified by mRNA, using tRNA molecules to carry amino acids and to read the mRNA three nucleotides at a time. The code defines how sequences of these nucleotide triplets, called codons, specify which amino acid will be added next during protein synthesis. With some exceptions, a three-nucleotide codon in a nucleic acid sequence specifies a single amino acid. Because the vast majority of genes are

encoded with exactly the same code, this particular code is often referred to as the canonical or standard genetic code. While the genetic code determines the protein sequence for a given coding region, other genomic regions can influence when and where these proteins are produced.

[0330] Furthermore, the invention provides a vector comprising a polynucleotide/nucleic acid molecule of the invention.

[0331] A vector is a nucleic acid molecule used as a vehicle to transfer (foreign) genetic material into a cell. The term “vector” encompasses—but is not restricted to—plasmids, viruses, cosmids and artificial chromosomes. In general, engineered vectors comprise an origin of replication, a multicloning site and a selectable marker. The vector itself is generally a nucleotide sequence, commonly a DNA sequence, that comprises an insert (transgene) and a larger sequence that serves as the “backbone” of the vector. Modern vectors may encompass additional features besides the transgene insert and a backbone: promoter, genetic marker, antibiotic resistance, reporter gene, targeting sequence, protein purification tag. Vectors called expression vectors (expression constructs) specifically are for the expression of the transgene in the target cell, and generally have control sequences.

[0332] The term “control sequences” refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

[0333] A nucleic acid is “operably linked” when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, “operably linked” means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

[0334] “Transfection” is the process of deliberately introducing nucleic acid molecules or polynucleotides (including vectors) into target cells. The term is mostly used for non-viral methods in eukaryotic cells. Transduction is often used to describe virus-mediated transfer of nucleic acid molecules or polynucleotides. Transfection of animal cells typically involves opening transient pores or “holes” in the cell membrane, to allow the uptake of material. Transfection can be carried out using calcium phosphate, by electroporation, by cell squeezing or by mixing a cationic lipid with the material to produce liposomes, which fuse with the cell membrane and deposit their cargo inside.

[0335] The term “transformation” is used to describe non-viral transfer of nucleic acid molecules or polynucleotides (including vectors) into bacteria, and also into non-animal eukaryotic cells, including plant cells. Transforma-

tion is hence the genetic alteration of a bacterial or non-animal eukaryotic cell resulting from the direct uptake through the cell membrane(s) from its surroundings and subsequent incorporation of exogenous genetic material (nucleic acid molecules). Transformation can be effected by artificial means. For transformation to happen, cells or bacteria must be in a state of competence, which might occur as a time-limited response to environmental conditions such as starvation and cell density.

[0336] Moreover, the invention provides a host cell transformed or transfected with the polynucleotide/nucleic acid molecule or with the vector of the invention.

[0337] As used herein, the terms “host cell” or “recipient cell” are intended to include any individual cell or cell culture that can be or has/have been recipients of vectors, exogenous nucleic acid molecules, and polynucleotides encoding the antibody construct of the present invention; and/or recipients of the antibody construct itself. The introduction of the respective material into the cell is carried out by way of transformation, transfection and the like. The term “host cell” is also intended to include progeny or potential progeny of a single cell. Because certain modifications may occur in succeeding generations due to either natural, accidental, or deliberate mutation or due to environmental influences, such progeny may not, in fact, be completely identical (in morphology or in genomic or total DNA complement) to the parent cell, but is still included within the scope of the term as used herein. Suitable host cells include prokaryotic or eukaryotic cells, and also include but are not limited to bacteria, yeast cells, fungi cells, plant cells, and animal cells such as insect cells and mammalian cells, e.g., murine, rat, macaque or human.

[0338] The antibody construct of the invention can be produced in bacteria. After expression, the antibody construct of the invention is isolated from the *E. coli* cell paste in a soluble fraction and can be purified through, e.g., affinity chromatography and/or size exclusion. Final purification can be carried out similar to the process for purifying antibody expressed e.g., in CHO cells.

[0339] In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for the antibody construct of the invention. *Saccharomyces cerevisiae*, or common baker’s yeast, is the most commonly used among lower eukaryotic host microorganisms. However, a number of other genera, species, and strains are commonly available and useful herein, such as *Schizosaccharomyces pombe*, *Kluyveromyces* hosts such as *K. lactis*, *K. fragilis* (ATCC 12424), *K. bulgaricus* (ATCC 16045), *K. wickerhamii* (ATCC 24178), *K. waltii* (ATCC 56500), *K. drosophilae* (ATCC 36906), *K. thermotolerans*, and *K. marxianus*; *Yarrowia* (EP 402 226); *Pichia pastoris* (EP 183 070); *Candida*; *Trichoderma reesei* (EP 244 234); *Neurospora crassa*; *Schwanniomyces* such as *Schwanniomyces occidentalis*; and filamentous fungi such as *Neurospora*, *Penicillium*, *Tolypocladium*, and *Aspergillus* hosts such as *A. nidulans* and *A. niger*.

[0340] Suitable host cells for the expression of glycosylated antibody construct of the invention are derived from multicellular organisms. Examples of invertebrate cells include plant and insect cells. Numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts such as *Spodoptera frugiperda* (caterpillar), *Aedes aegypti* (mosquito), *Aedes albopictus* (mosquito), *Drosophila melanogaster* (fruit fly), and *Bombyx mori* have

been identified. A variety of viral strains for transfection are publicly available, e.g., the L-1 variant of *Autographa californica* NPV and the Bm-5 strain of *Bombyx mori* NPV, and such viruses may be used as the virus herein according to the present invention, particularly for transfection of *Spodoptera frugiperda* cells.

[0341] Plant cell cultures of cotton, corn, potato, soybean, *petunia*, tomato, *Arabidopsis* and tobacco can also be used as hosts. Cloning and expression vectors useful in the production of proteins in plant cell culture are known to those of skill in the art. See e.g. Hiatt et al., *Nature* (1989) 342: 76-78, Owen et al. (1992) *Bio/Technology* 10: 790-794, Artsenko et al. (1995) *The Plant J* 8: 745-750, and Fecker et al. (1996) *Plant Mol Biol* 32: 979-986.

[0342] However, interest has been greatest in vertebrate cells, and propagation of vertebrate cells in culture (tissue culture) has become a routine procedure. Examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., *J. Gen Virol.* 36: 59 (1977)); baby hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells/-DHFR (CHO, Urlaub et al., *Proc. Natl. Acad. Sci. USA* 77:4216 (1980)); mouse sertoli cells (TM4, Mather, *Biol. Reprod.* 23: 243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, 1413 8065); mouse mammary tumor (MMT 060562, ATCC CCL5 1); TRI cells (Mather et al., *Annals N. Y Acad. Sci.* (1982) 383: 44-68); MRC 5 cells; FS4 cells; and a human hepatoma line (Hep G2).

[0343] In a further embodiment the invention provides a process for the production of an antibody construct of the invention, said process comprising culturing a host cell of the invention under conditions allowing the expression of the antibody construct of the invention and recovering the produced antibody construct from the culture.

[0344] As used herein, the term "culturing" refers to the in vitro maintenance, differentiation, growth, proliferation and/or propagation of cells under suitable conditions in a medium. The term "expression" includes any step involved in the production of an antibody construct of the invention including, but not limited to, transcription, post-transcriptional modification, translation, post-translational modification, and secretion.

[0345] When using recombinant techniques, the antibody construct can be produced intracellularly, in the periplasmic space, or directly secreted into the medium. If the antibody construct is produced intracellularly, as a first step, the particulate debris, either host cells or lysed fragments, are removed, for example, by centrifugation or ultrafiltration. Carter et al., *Bio/Technology* 10: 163-167 (1992) describe a procedure for isolating antibodies which are secreted to the periplasmic space of *E. coli*. Briefly, cell paste is thawed in the presence of sodium acetate (pH 3.5), EDTA, and phenylmethylsulfonyl fluoride (PMSF) over about 30 min. Cell debris can be removed by centrifugation. Where the antibody is secreted into the medium, supernatants from such expression systems are generally first concentrated using a commercially available protein concentration filter, for

example, an Amicon or Millipore Pellicon ultrafiltration unit. A protease inhibitor such as PMSF may be included in any of the foregoing steps to inhibit proteolysis and antibiotics may be included to prevent the growth of adventitious contaminants.

[0346] The antibody construct of the invention prepared from the host cells can be recovered or purified using, for example, hydroxylapatite chromatography, gel electrophoresis, dialysis, and affinity chromatography. Other techniques for protein purification such as fractionation on an ion-exchange column, ethanol precipitation, Reverse Phase HPLC, chromatography on silica, chromatography on heparin SEPHAROSE™, chromatography on an anion or cation exchange resin (such as a polyaspartic acid column), chromatofocusing, SDS-PAGE, and ammonium sulfate precipitation are also available depending on the antibody to be recovered. Where the antibody construct of the invention comprises a CH3 domain, the Bakerbond ABX resin (J.T. Baker, Phillipsburg, NJ) is useful for purification.

[0347] Affinity chromatography is a preferred purification technique. The matrix to which the affinity ligand is attached is most often agarose, but other matrices are available. Mechanically stable matrices such as controlled pore glass or poly (styrenedivinyl) benzene allow for faster flow rates and shorter processing times than can be achieved with agarose.

[0348] Moreover, the invention provides a pharmaceutical composition comprising an antibody construct of the invention or an antibody construct produced according to the process of the invention. It is preferred for the pharmaceutical composition of the invention that the homogeneity of the antibody construct is $\geq 80\%$, more preferably $\geq 81\%$, $\geq 82\%$, $\geq 83\%$, $\geq 84\%$, or $\geq 85\%$, further preferably $\geq 86\%$, $\geq 87\%$, $\geq 88\%$, $\geq 89\%$, or $\geq 90\%$, still further preferably, $\geq 91\%$, $\geq 92\%$, $\geq 93\%$, $\geq 94\%$, or $\geq 95\%$ and most preferably $\geq 96\%$, $\geq 97\%$, $\geq 98\%$ or $\geq 99\%$.

[0349] As used herein, the term "pharmaceutical composition" relates to a composition which is suitable for administration to a patient, preferably a human patient. The particularly preferred pharmaceutical composition of this invention comprises one or a plurality of the antibody construct(s) of the invention, preferably in a therapeutically effective amount. Preferably, the pharmaceutical composition further comprises suitable formulations of one or more (pharmaceutically effective) carriers, stabilizers, excipients, diluents, solubilizers, surfactants, emulsifiers, preservatives and/or adjuvants. Acceptable constituents of the composition are preferably nontoxic to recipients at the dosages and concentrations employed. Pharmaceutical compositions of the invention include, but are not limited to, liquid, frozen, and lyophilized compositions.

[0350] The inventive compositions may comprise a pharmaceutically acceptable carrier. In general, as used herein, "pharmaceutically acceptable carrier" means any and all aqueous and non-aqueous solutions, sterile solutions, solvents, buffers, e.g. phosphate buffered saline (PBS) solutions, water, suspensions, emulsions, such as oil/water emulsions, various types of wetting agents, liposomes, dispersion media and coatings, which are compatible with pharmaceutical administration, in particular with parenteral administration. The use of such media and agents in pharmaceutical compositions is well known in the art, and the compositions comprising such carriers can be formulated by well-known conventional methods.

[0351] Certain embodiments provide pharmaceutical compositions comprising the antibody construct of the invention and further one or more excipients such as those illustratively described in this section and elsewhere herein. Excipients can be used in the invention in this regard for a wide variety of purposes, such as adjusting physical, chemical, or biological properties of formulations, such as adjustment of viscosity, and or processes of the invention to improve effectiveness and or to stabilize such formulations and processes against degradation and spoilage due to, for instance, stresses that occur during manufacturing, shipping, storage, pre-use preparation, administration, and thereafter.

[0352] In certain embodiments, the pharmaceutical composition may contain formulation materials for the purpose of modifying, maintaining or preserving, e.g., the pH, osmolarity, viscosity, clarity, color, isotonicity, odor, sterility, stability, rate of dissolution or release, adsorption or penetration of the composition (see, REMINGTON'S PHARMACEUTICAL SCIENCES, 18th Edition, (A.R. Genmo, ed.), 1990, Mack Publishing Company). In such embodiments, suitable formulation materials may include, but are not limited to:

[0353] amino acids such as glycine, alanine, glutamine, asparagine, threonine, proline, 2-phenylalanine, including charged amino acids, preferably lysine, lysine acetate, arginine, glutamate and/or histidine

[0354] antimicrobials such as antibacterial and antifungal agents

[0355] antioxidants such as ascorbic acid, methionine, sodium sulfite or sodium hydrogen-sulfite;

[0356] buffers, buffer systems and buffering agents which are used to maintain the composition at physiological pH or at a slightly lower pH, typically within a pH range of from about 5 to about 8 or 9; examples of buffers are borate, bicarbonate, Tris-HCl, citrates, phosphates or other organic acids, succinate, phosphate, histidine and acetate; for example Tris buffer of about pH 7.0-8.5, or acetate buffer of about pH 4.0-5.5;

[0357] non-aqueous solvents such as propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate;

[0358] aqueous carriers including water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media;

[0359] biodegradable polymers such as polyesters;

[0360] bulking agents such as mannitol or glycine;

[0361] chelating agents such as ethylenediamine tetraacetic acid (EDTA);

[0362] isotonic and absorption delaying agents;

[0363] complexing agents such as caffeine, polyvinylpyrrolidone, beta-cyclodextrin or hydroxypropyl-beta-cyclodextrin)

[0364] fillers;

[0365] monosaccharides; disaccharides; and other carbohydrates (such as glucose, mannose or dextrans); carbohydrates may be non-reducing sugars, preferably trehalose, sucrose, octasulfate, sorbitol or xylitol;

[0366] (low molecular weight) proteins, polypeptides or proteinaceous carriers such as human or bovine serum albumin, gelatin or immunoglobulins, preferably of human origin;

[0367] coloring and flavouring agents;

[0368] sulfur containing reducing agents, such as glutathione, thioctic acid, sodium thioglycolate, thioglycerol, [alpha]-monothioglycerol, and sodium thio sulfate

[0369] diluting agents;

[0370] emulsifying agents;

[0371] hydrophilic polymers such as polyvinylpyrrolidone)

[0372] salt-forming counter-ions such as sodium;

[0373] preservatives such as antimicrobials, anti-oxidants, chelating agents, inert gases and the like; examples are: benzalkonium chloride, benzoic acid, salicylic acid, thimerosal, phenethyl alcohol, methylparaben, propylparaben, chlorhexidine, sorbic acid or hydrogen peroxide);

[0374] metal complexes such as Zn-protein complexes;

[0375] solvents and co-solvents (such as glycerin, propylene glycol or polyethylene glycol);

[0376] sugars and sugar alcohols, such as trehalose, sucrose, octasulfate, mannitol, sorbitol or xylitol stachyose, mannose, sorbose, xylose, ribose, myoinositol, galactose, lactitol, ribitol, myoinositol, galactitol, glycerol, cyclitols (e.g., inositol), polyethylene glycol; and polyhydric sugar alcohols;

[0377] suspending agents;

[0378] surfactants or wetting agents such as pluronics, PEG, sorbitan esters, polysorbates such as polysorbate 20, polysorbate, triton, tromethamine, lecithin, cholesterol, tyloxapal; surfactants may be detergents, preferably with a molecular weight of >1.2 KD and/or a polyether, preferably with a molecular weight of >3 KD; non-limiting examples for preferred detergents are Tween 20, Tween 40, Tween 60, Tween 80 and Tween 85; non-limiting examples for preferred polyethers are PEG 3000, PEG 3350, PEG 4000 and PEG 5000;

[0379] stability enhancing agents such as sucrose or sorbitol;

[0380] tonicity enhancing agents such as alkali metal halides, preferably sodium or potassium chloride, mannitol sorbitol;

[0381] parenteral delivery vehicles including sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's, or fixed oils;

[0382] intravenous delivery vehicles including fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose).

[0383] It is evident to those skilled in the art that the different constituents of the pharmaceutical composition (e.g., those listed above) can have different effects, for example, and amino acid can act as a buffer, a stabilizer and/or an antioxidant; mannitol can act as a bulking agent and/or a tonicity enhancing agent; sodium chloride can act as delivery vehicle and/or tonicity enhancing agent; etc.

[0384] It is envisaged that the composition of the invention might comprise, in addition to the polypeptide of the invention defined herein, further biologically active agents, depending on the intended use of the composition. Such agents might be drugs acting on the gastro-intestinal system, drugs acting as cytostatics, drugs preventing hyperurikemia, drugs inhibiting immunoreactions (e.g. corticosteroids), drugs modulating the inflammatory response, drugs acting on the circulatory system and/or agents such as cytokines known in the art. It is also envisaged that the antibody construct of the present invention is applied in a co-therapy, i.e., in combination with another anti-cancer medicament.

[0385] In certain embodiments, the optimal pharmaceutical composition will be determined by one skilled in the art depending upon, for example, the intended route of administration, delivery format and desired dosage. See, for example, REMINGTON'S PHARMACEUTICAL SCIENCES, supra. In certain embodiments, such compositions may influence the physical state, stability, rate of in vivo release and rate of in vivo clearance of the antibody construct of the invention. In certain embodiments, the primary vehicle or carrier in a pharmaceutical composition may be either aqueous or non-aqueous in nature. For example, a suitable vehicle or carrier may be water for injection, physiological saline solution or artificial cerebrospinal fluid, possibly supplemented with other materials common in compositions for parenteral administration. Neutral buffered saline or saline mixed with serum albumin are further exemplary vehicles. In certain embodiments, the antibody construct of the invention compositions may be prepared for storage by mixing the selected composition having the desired degree of purity with optional formulation agents (REMINGTON'S PHARMACEUTICAL SCIENCES, supra) in the form of a lyophilized cake or an aqueous solution. Further, in certain embodiments, the antibody construct of the invention may be formulated as a lyophilizate using appropriate excipients such as sucrose.

[0386] When parenteral administration is contemplated, the therapeutic compositions for use in this invention may be provided in the form of a pyrogen-free, parenterally acceptable aqueous solution comprising the desired antibody construct of the invention in a pharmaceutically acceptable vehicle. A particularly suitable vehicle for parenteral injection is sterile distilled water in which the antibody construct of the invention is formulated as a sterile, isotonic solution, properly preserved. In certain embodiments, the preparation can involve the formulation of the desired molecule with an agent, such as injectable microspheres, bio-erodible particles, polymeric compounds (such as polylactic acid or polyglycolic acid), beads or liposomes, that may provide controlled or sustained release of the product which can be delivered via depot injection. In certain embodiments, hyaluronic acid may also be used, having the effect of promoting sustained duration in the circulation. In certain embodiments, implantable drug delivery devices may be used to introduce the desired antibody construct.

[0387] Additional pharmaceutical compositions will be evident to those skilled in the art, including formulations involving the antibody construct of the invention in sustained- or controlled-delivery/release formulations. Techniques for formulating a variety of other sustained- or controlled-delivery means, such as liposome carriers, bio-erodible microparticles or porous beads and depot injections, are also known to those skilled in the art. See, for example, International Patent Application No. PCT/US93/00829, which describes controlled release of porous polymeric microparticles for delivery of pharmaceutical compositions.

[0388] Sustained-release preparations may include semi-permeable polymer matrices in the form of shaped articles, e.g., films, or microcapsules. Sustained release matrices may include polyesters, hydrogels, polylactides (as disclosed in U.S. Pat. No. 3,773,919 and European Patent Application Publication No. EP 058481), copolymers of L-glutamic acid and gamma ethyl-L-glutamate (Sidman et al., 1983, Biopolymers 2:547-556), poly (2-hydroxyethyl-methacrylate)

(Langer et al., 1981, J. Biomed. Mater. Res. 15:167-277 and Langer, 1982, Chem. Tech. 12:98-105), ethylene vinyl acetate (Langer et al., 1981, supra) or poly-D(-)-3-hydroxybutyric acid (European Patent Application Publication No. EP 133,988). Sustained release compositions may also include liposomes that can be prepared by any of several methods known in the art. See, e.g., Eppstein et al., 1985, Proc. Natl. Acad. Sci. U.S.A. 82:3688-3692; European Patent Application Publication Nos. EP 036,676; EP 088,046 and EP 143,949.

[0389] The antibody construct may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization (for example, hydroxymethylcellulose or gelatine-microcapsules and poly (methylmethacrylate) microcapsules, respectively), in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nanoparticles and nanocapsules), or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences, 16th edition, Oslo, A., Ed., (1980).

[0390] Pharmaceutical compositions used for in vivo administration are typically provided as sterile preparations. Sterilization can be accomplished by filtration through sterile filtration membranes. When the composition is lyophilized, sterilization using this method may be conducted either prior to or following lyophilization and reconstitution. Compositions for parenteral administration can be stored in lyophilized form or in a solution. Parenteral compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

[0391] Another aspect of the invention includes self-buffering antibody construct of the invention formulations, which can be used as pharmaceutical compositions, as described in international patent application WO 06138181A2 (PCT/US2006/022599). A variety of expositions are available on protein stabilization and formulation materials and methods useful in this regard, such as Arakawa et al., "Solvent interactions in pharmaceutical formulations," Pharm Res. 8(3): 285-91 (1991); Kendrick et al., "Physical stabilization of proteins in aqueous solution" in: RATIO-NAL DESIGN OF STABLE PROTEIN FORMULATIONS: THEORY AND PRACTICE, Carpenter and Manning, eds. Pharmaceutical Biotechnology. 13: 61-84 (2002), and Randolph et al., "Surfactant-protein interactions", Pharm Biotechnol. 13: 159-75 (2002), see particularly the parts pertinent to excipients and processes of the same for self-buffering protein formulations in accordance with the current invention, especially as to protein pharmaceutical products and processes for veterinary and/or human medical uses.

[0392] Salts may be used in accordance with certain embodiments of the invention to, for example, adjust the ionic strength and/or the isotonicity of a formulation and/or to improve the solubility and/or physical stability of a protein or other ingredient of a composition in accordance with the invention. As is well known, ions can stabilize the native state of proteins by binding to charged residues on the protein's surface and by shielding charged and polar groups in the protein and reducing the strength of their electrostatic interactions, attractive, and repulsive interactions. Ions also can stabilize the denatured state of a protein by binding to, in particular, the denatured peptide linkages ($-\text{CONH}$) of the protein. Furthermore, ionic interaction with charged and

polar groups in a protein also can reduce intermolecular electrostatic interactions and, thereby, prevent or reduce protein aggregation and insolubility.

[0393] Ionic species differ significantly in their effects on proteins. A number of categorical rankings of ions and their effects on proteins have been developed that can be used in formulating pharmaceutical compositions in accordance with the invention. One example is the Hofmeister series, which ranks ionic and polar non-ionic solutes by their effect on the conformational stability of proteins in solution. Stabilizing solutes are referred to as “kosmotropic”. Destabilizing solutes are referred to as “chaotropic”. Kosmotropes commonly are used at high concentrations (e.g., >1 molar ammonium sulfate) to precipitate proteins from solution (“salting-out”). Chaotropes commonly are used to denature and/or to solubilize proteins (“salting-in”). The relative effectiveness of ions to “salt-in” and “salt-out” defines their position in the Hofmeister series.

[0394] Free amino acids can be used in the antibody construct of the invention formulations in accordance with various embodiments of the invention as bulking agents, stabilizers, and antioxidants, as well as other standard uses. Lysine, proline, serine, and alanine can be used for stabilizing proteins in a formulation. Glycine is useful in lyophilization to ensure correct cake structure and properties. Arginine may be useful to inhibit protein aggregation, in both liquid and lyophilized formulations. Methionine is useful as an antioxidant.

[0395] Polyols include sugars, e.g., mannitol, sucrose, and sorbitol and polyhydric alcohols such as, for instance, glycerol and propylene glycol, and, for purposes of discussion herein, polyethylene glycol (PEG) and related substances. Polyols are kosmotropic. They are useful stabilizing agents in both liquid and lyophilized formulations to protect proteins from physical and chemical degradation processes. Polyols also are useful for adjusting the tonicity of formulations. Among polyols useful in select embodiments of the invention is mannitol, commonly used to ensure structural stability of the cake in lyophilized formulations. It ensures structural stability to the cake. It is generally used with a lyoprotectant, e.g., sucrose. Sorbitol and sucrose are among preferred agents for adjusting tonicity and as stabilizers to protect against freeze-thaw stresses during transport or the preparation of bulks during the manufacturing process. Reducing sugars (which contain free aldehyde or ketone groups), such as glucose and lactose, can glycate surface lysine and arginine residues. Therefore, they generally are not among preferred polyols for use in accordance with the invention. In addition, sugars that form such reactive species, such as sucrose, which is hydrolyzed to fructose and glucose under acidic conditions, and consequently engenders glycation, also is not among preferred polyols of the invention in this regard. PEG is useful to stabilize proteins and as a cryoprotectant and can be used in the invention in this regard.

[0396] Embodiments of the antibody construct of the invention formulations further comprise surfactants. Protein molecules may be susceptible to adsorption on surfaces and to denaturation and consequent aggregation at air-liquid, solid-liquid, and liquid-liquid interfaces. These effects generally scale inversely with protein concentration. These deleterious interactions generally scale inversely with protein concentration and typically are exacerbated by physical agitation, such as that generated during the shipping and

handling of a product. Surfactants routinely are used to prevent, minimize, or reduce surface adsorption. Useful surfactants in the invention in this regard include polysorbate 20, polysorbate 80, other fatty acid esters of sorbitan polyethoxylates, and poloxamer 188. Surfactants also are commonly used to control protein conformational stability. The use of surfactants in this regard is protein-specific since, any given surfactant typically will stabilize some proteins and destabilize others.

[0397] Polysorbates are susceptible to oxidative degradation and often, as supplied, contain sufficient quantities of peroxides to cause oxidation of protein residue side-chains, especially methionine. Consequently, polysorbates should be used carefully, and when used, should be employed at their lowest effective concentration. In this regard, polysorbates exemplify the general rule that excipients should be used in their lowest effective concentrations.

[0398] Embodiments of the antibody construct of the invention formulations further comprise one or more antioxidants. To some extent deleterious oxidation of proteins can be prevented in pharmaceutical formulations by maintaining proper levels of ambient oxygen and temperature and by avoiding exposure to light. Antioxidant excipients can be used as well to prevent oxidative degradation of proteins. Among useful antioxidants in this regard are reducing agents, oxygen/free-radical scavengers, and chelating agents. Antioxidants for use in therapeutic protein formulations in accordance with the invention preferably are water-soluble and maintain their activity throughout the shelf life of a product. EDTA is a preferred antioxidant in accordance with the invention in this regard. Antioxidants can damage proteins. For instance, reducing agents, such as glutathione in particular, can disrupt intramolecular disulfide linkages. Thus, antioxidants for use in the invention are selected to, among other things, eliminate or sufficiently reduce the possibility of themselves damaging proteins in the formulation.

[0399] Formulations in accordance with the invention may include metal ions that are protein co-factors and that are necessary to form protein coordination complexes, such as zinc necessary to form certain insulin suspensions. Metal ions also can inhibit some processes that degrade proteins. However, metal ions also catalyze physical and chemical processes that degrade proteins. Magnesium ions (10-120 mM) can be used to inhibit isomerization of aspartic acid to isoaspartic acid. Ca^{+2} ions (up to 100 mM) can increase the stability of human deoxyribonuclease. Mg^{+2} , Mn^{+2} , and Zn^{+2} , however, can destabilize rhDNase. Similarly, Ca^{+2} and Sr^{+2} can stabilize Factor VIII, it can be destabilized by Mg^{+2} , Mn^{+2} and Zn^{+2} , Cu^{+2} and Fe^{+2} , and its aggregation can be increased by Al^{+3} ions.

[0400] Embodiments of the antibody construct of the invention formulations further comprise one or more preservatives. Preservatives are necessary when developing multi-dose parenteral formulations that involve more than one extraction from the same container. Their primary function is to inhibit microbial growth and ensure product sterility throughout the shelf-life or term of use of the drug product. Commonly used preservatives include benzyl alcohol, phenol and m-cresol. Although preservatives have a long history of use with small-molecule parenterals, the development of protein formulations that includes preservatives can be challenging. Preservatives almost always have a destabilizing effect (aggregation) on proteins, and this

has become a major factor in limiting their use in multi-dose protein formulations. To date, most protein drugs have been formulated for single-use only. However, when multi-dose formulations are possible, they have the added advantage of enabling patient convenience, and increased marketability. A good example is that of human growth hormone (hGH) where the development of preserved formulations has led to commercialization of more convenient, multi-use injection pen presentations. At least four such pen devices containing preserved formulations of hGH are currently available on the market. Norditropin (liquid, Novo Nordisk), Nutropin AQ (liquid, Genentech) & Genotropin (lyophilized—dual chamber cartridge, Pharmacia & Upjohn) contain phenol while Somatropo (Eli Lilly) is formulated with m-cresol. Several aspects need to be considered during the formulation and development of preserved dosage forms. The effective preservative concentration in the drug product must be optimized. This requires testing a given preservative in the dosage form with concentration ranges that confer antimicrobial effectiveness without compromising protein stability.

[0401] As might be expected, development of liquid formulations containing preservatives are more challenging than lyophilized formulations. Freeze-dried products can be lyophilized without the preservative and reconstituted with a preservative containing diluent at the time of use. This shortens the time for which a preservative is in contact with the protein, significantly minimizing the associated stability risks. With liquid formulations, preservative effectiveness and stability should be maintained over the entire product shelf-life (about 18 to 24 months). An important point to note is that preservative effectiveness should be demonstrated in the final formulation containing the active drug and all excipient components.

[0402] The antibody constructs disclosed herein may also be formulated as immuno-liposomes. A “liposome” is a small vesicle composed of various types of lipids, phospholipids and/or surfactant which is useful for delivery of a drug to a mammal. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes. Liposomes containing the antibody construct are prepared by methods known in the art, such as described in Epstein et al., Proc. Natl. Acad. Sci. USA, 82: 3688 (1985); Hwang et al., Proc. Natl. Acad. Sci. USA, 77: 4030 (1980); U.S. Pat. Nos. 4,485,045 and 4,544,545; and WO 97/38731. Uposomes with enhanced circulation time are disclosed in U.S. Pat. No. 5,013,556. Particularly useful liposomes can be generated by the reverse phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol and PEG-derivatized phosphatidylethanolamine (PEG-PE). Uposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody construct of the present invention can be conjugated to the liposomes as described in Martin et al. J. Biol. Chem. 257: 286-288 (1982) via a disulfide interchange reaction. A chemotherapeutic agent is optionally contained within the liposome. See Gabizon et al. J. National Cancer Inst. 81 (19) 1484 (1989).

[0403] Once the pharmaceutical composition has been formulated, it may be stored in sterile vials as a solution, suspension, gel, emulsion, solid, crystal, or as a dehydrated or lyophilized powder. Such formulations may be stored

either in a ready-to-use form or in a form (e.g., lyophilized) that is reconstituted prior to administration.

[0404] The biological activity of the pharmaceutical composition defined herein can be determined for instance by cytotoxicity assays, as described in the following examples, in WO 99/54440 or by Schlereth et al. (Cancer Immunol. Immunother. 20 (2005), 1-12). “Efficacy” or “in vivo efficacy” as used herein refers to the response to therapy by the pharmaceutical composition of the invention, using e.g. standardized NCI response criteria. The success or in vivo efficacy of the therapy using a pharmaceutical composition of the invention refers to the effectiveness of the composition for its intended purpose, i.e. the ability of the composition to cause its desired effect, i.e. depletion of pathologic cells, e.g. tumor cells. The in vivo efficacy may be monitored by established standard methods for the respective disease entities including, but not limited to white blood cell counts, differentials, Fluorescence Activated Cell Sorting, bone marrow aspiration. In addition, various disease specific clinical chemistry parameters and other established standard methods may be used. Furthermore, computer-aided tomography, X-ray, nuclear magnetic resonance tomography (e.g. for National Cancer Institute-criteria based response assessment [Cheson B D, Horning S J, Coiffier B, Shipp M A, Fisher R I, Connors J M, Lister T A, Vose J, Grillo-Lopez A, Hagenebeck A, Cabanillas F, Klippensten D, Hiddemann W, Castellino R, Harris N L, Armitage J O, Carter W, Hoppe R, Canellos G P. Report of an international workshop to standardize response criteria for non-Hodgkin's lymphomas. NCI Sponsored International Working Group. J Clin Oncol. 1999 April; 17(4):1244]), positron-emission tomography scanning, white blood cell counts, differentials, Fluorescence Activated Cell Sorting, bone marrow aspiration, lymph node biopsies/histologies, and various lymphoma specific clinical chemistry parameters (e.g. lactate dehydrogenase) and other established standard methods may be used.

[0405] Another major challenge in the development of drugs such as the pharmaceutical composition of the invention is the predictable modulation of pharmacokinetic properties. To this end, a pharmacokinetic profile of the drug candidate, i.e. a profile of the pharmacokinetic parameters that affect the ability of a particular drug to treat a given condition, can be established. Pharmacokinetic parameters of the drug influencing the ability of a drug to treating a certain disease entity include, but are not limited to: half-life, volume of distribution, hepatic first-pass metabolism and the degree of blood serum binding. The efficacy of a given drug agent can be influenced by each of the parameters mentioned above.

[0406] “Half-life” means the time where 50% of an administered drug are eliminated through biological processes, e.g. metabolism, excretion, etc. By “hepatic first-pass metabolism” is meant the propensity of a drug to be metabolized upon first contact with the liver, i.e. during its first pass through the liver. “Volume of distribution” means the degree of retention of a drug throughout the various compartments of the body, like e.g. intracellular and extracellular spaces, tissues and organs, etc. and the distribution of the drug within these compartments. “Degree of blood serum binding” means the propensity of a drug to interact with and bind to blood serum proteins, such as albumin, leading to a reduction or loss of biological activity of the drug.

[0407] Pharmacokinetic parameters also include bioavailability, lag time (Tlag), Tmax, absorption rates, more onset and/or Cmax for a given amount of drug administered. “Bioavailability” means the amount of a drug in the blood compartment. “Lag time” means the time delay between the administration of the drug and its detection and measurability in blood or plasma. “Tmax” is the time after which maximal blood concentration of the drug is reached, and “Cmax” is the blood concentration maximally obtained with a given drug. The time to reach a blood or tissue concentration of the drug which is required for its biological effect is influenced by all parameters. Pharmacokinetic parameters of bispecific antibody constructs exhibiting cross-species specificity, which may be determined in preclinical animal testing in non-chimpanzee primates as outlined above, are also set forth e.g. in the publication by Schlereth et al. (Cancer Immunol. Immunother. 20 (2005), 1-12).

[0408] One embodiment provides the antibody construct of the invention or the antibody construct produced according to the process of the invention for use in the prevention, treatment or amelioration of a tumor or cancer disease or of a metastatic cancer disease.

[0409] According to a preferred embodiment of the invention said tumor or cancer disease is a solid tumor disease.

[0410] The formulations described herein are useful as pharmaceutical compositions in the treatment, amelioration and/or prevention of the pathological medical condition as described herein in a patient in need thereof. The term “treatment” refers to both therapeutic treatment and prophylactic or preventative measures. Treatment includes the application or administration of the formulation to the body, an isolated tissue, or cell from a patient who has a disease/disorder, a symptom of a disease/disorder, or a predisposition toward a disease/disorder, with the purpose to cure, heal, alleviate, relieve, after, remedy, ameliorate, improve, or affect the disease, the symptom of the disease, or the predisposition toward the disease.

[0411] The term “amelioration” as used herein refers to any improvement of the disease state of a patient having a tumor or cancer or a metastatic cancer as specified herein below, by the administration of an antibody construct according to the invention to a subject in need thereof. Such an improvement may also be seen as a slowing or stopping of the progression of the tumor or cancer or metastatic cancer of the patient. The term “prevention” as used herein means the avoidance of the occurrence or re-occurrence of a patient having a tumor or cancer or a metastatic cancer as specified herein below, by the administration of an antibody construct according to the invention to a subject in need thereof.

[0412] The term “disease” refers to any condition that would benefit from treatment with the antibody construct or the pharmaceutical composition described herein. This includes chronic and acute disorders or diseases including those pathological conditions that predispose the mammal to the disease in question.

[0413] A “neoplasm” is an abnormal growth of tissue, usually but not always forming a mass. When also forming a mass, it is commonly referred to as a “tumor”. Neoplasms or tumors or can be benign, potentially malignant (pre-cancerous), or malignant. Malignant neoplasms are commonly called cancer. They usually invade and destroy the surrounding tissue and may form metastases, i.e., they spread to other parts, tissues or organs of the body. Hence,

the term “metastatic cancer” encompasses metastases to other tissues or organs than the one of the original tumor. Lymphomas and leukemias are lymphoid neoplasms. For the purposes of the present invention, they are also encompassed by the terms “tumor” or “cancer”.

[0414] In a preferred embodiment of the invention, the tumor or cancer disease is a solid tumor disease and the metastatic cancer disease can be derived from any of the foregoing.

[0415] Preferred tumor or cancer diseases in connection with this invention are selected from a group consisting of breast cancer, Carcinoid, cervical cancer, colorectal cancer, endometrial cancer, gastric cancer, head and neck cancer, mesothelioma, liver cancer, lung cancer, ovarian cancer, pancreatic cancer, prostate cancer, skin cancer, renal cancer and stomach cancer. More preferably, the tumor or cancer disease, which is preferably a solid tumor disease, can be selected from the group consisting of ovarian cancer, pancreatic cancer, mesothelioma, lung cancer, gastric cancer and triple negative breast cancer. The metastatic cancer disease can be derived from any of the foregoing.

[0416] The invention also provides a method for the treatment or amelioration of tumor or cancer disease or a metastatic cancer disease, comprising the step of administering to a subject in need thereof the antibody construct of the invention or the antibody construct produced according to the process of the invention.

[0417] The terms “subject in need” or those “in need of treatment” includes those already with the disorder, as well as those in which the disorder is to be prevented. The subject in need or “patient” includes human and other mammalian subjects that receive either prophylactic or therapeutic treatment.

[0418] The antibody construct of the invention will generally be designed for specific routes and methods of administration, for specific dosages and frequencies of administration, for specific treatments of specific diseases, with ranges of bio-availability and persistence, among other things. The materials of the composition are preferably formulated in concentrations that are acceptable for the site of administration.

[0419] Formulations and compositions thus may be designed in accordance with the invention for delivery by any suitable route of administration. In the context of the present invention, the routes of administration include, but are not limited to

[0420] topical routes (such as epicutaneous, inhalational, nasal, ophthalmic, auricular/aural, vaginal, mucosal);

[0421] enteral routes (such as oral, gastrointestinal, sublingual, sublabial, buccal, rectal); and

[0422] parenteral routes (such as intravenous, intraarterial, intraosseous, intramuscular, intracerebral, intracerebroventricular, epidural, intrathecal, subcutaneous, intraperitoneal, extra-amniotic, intraarticular, intracardiac, intradermal, intralesional, intrauterine, intravesical, intravitreal, transdermal, intranasal, transmucosal, intrasynovial, intraluminal).

[0423] The pharmaceutical compositions and the antibody construct of this invention are particularly useful for parenteral administration, e.g., subcutaneous or intravenous delivery, for example by injection such as bolus injection, or by infusion such as continuous infusion. Pharmaceutical compositions may be administered using a medical device.

Examples of medical devices for administering pharmaceutical compositions are described in U.S. Pat. Nos. 4,475,196; 4,439,196; 4,447,224; 4,447, 233; 4,486,194; 4,487,603; 4,596,556; 4,790,824; 4,941,880; 5,064,413; 5,312,335; 5,312,335; 5,383,851; and 5,399,163.

[0424] In particular, the present invention provides for an uninterrupted administration of the suitable composition. As a non-limiting example, uninterrupted or substantially uninterrupted, i.e. continuous administration may be realized by a small pump system worn by the patient for metering the influx of therapeutic agent into the body of the patient. The pharmaceutical composition comprising the antibody construct of the invention can be administered by using said pump systems. Such pump systems are generally known in the art, and commonly rely on periodic exchange of cartridges containing the therapeutic agent to be infused. When exchanging the cartridge in such a pump system, a temporary interruption of the otherwise uninterrupted flow of therapeutic agent into the body of the patient may ensue. In such a case, the phase of administration prior to cartridge replacement and the phase of administration following cartridge replacement would still be considered within the meaning of the pharmaceutical means and methods of the invention together make up one "uninterrupted administration" of such therapeutic agent.

[0425] The continuous or uninterrupted administration of the antibody constructs of the invention may be intravenous or subcutaneous by way of a fluid delivery device or small pump system including a fluid driving mechanism for driving fluid out of a reservoir and an actuating mechanism for actuating the driving mechanism. Pump systems for subcutaneous administration may include a needle or a cannula for penetrating the skin of a patient and delivering the suitable composition into the patient's body. Said pump systems may be directly fixed or attached to the skin of the patient independently of a vein, artery or blood vessel, thereby allowing a direct contact between the pump system and the skin of the patient. The pump system can be attached to the skin of the patient for 24 hours up to several days. The pump system may be of small size with a reservoir for small volumes. As a non-limiting example, the volume of the reservoir for the suitable pharmaceutical composition to be administered can be between 0.1 and 50 ml.

[0426] The continuous administration may also be transdermal by way of a patch worn on the skin and replaced at intervals. One of skill in the art is aware of patch systems for drug delivery suitable for this purpose. It is of note that transdermal administration is especially amenable to uninterrupted administration, as exchange of a first exhausted patch can advantageously be accomplished simultaneously with the placement of a new, second patch, for example on the surface of the skin immediately adjacent to the first exhausted patch and immediately prior to removal of the first exhausted patch. Issues of flow interruption or power cell failure do not arise.

[0427] If the pharmaceutical composition has been lyophilized, the lyophilized material is first reconstituted in an appropriate liquid prior to administration. The lyophilized material may be reconstituted in, e.g., bacteriostatic water for injection (BWI), physiological saline, phosphate buffered saline (PBS), or the same formulation the protein had been in prior to lyophilization.

[0428] The compositions of the present invention can be administered to the subject at a suitable dose which can be

determined e.g. by dose escalating studies by administration of increasing doses of the antibody construct of the invention exhibiting cross-species specificity described herein to non-chimpanzee primates, for instance macaques. As set forth above, the antibody construct of the invention exhibiting cross-species specificity described herein can be advantageously used in identical form in preclinical testing in non-chimpanzee primates and as drug in humans. The dosage regimen will be determined by the attending physician and clinical factors. As is well known in the medical arts, dosages for any one patient depend upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently.

[0429] The term "effective dose" or "effective dosage" is defined as an amount sufficient to achieve or at least partially achieve the desired effect. The term "therapeutically effective dose" is defined as an amount sufficient to cure or at least partially arrest the disease and its complications in a patient already suffering from the disease. Amounts or doses effective for this use will depend on the condition to be treated (the indication), the delivered antibody construct, the therapeutic context and objectives, the severity of the disease, prior therapy, the patient's clinical history and response to the therapeutic agent, the route of administration, the size (body weight, body surface or organ size) and/or condition (the age and general health) of the patient, and the general state of the patient's own immune system. The proper dose can be adjusted according to the judgment of the attending physician such that it can be administered to the patient once or over a series of administrations, and in order to obtain the optimal therapeutic effect.

[0430] A typical dosage may range from about 0.1 pg/kg to up to about 30 mg/kg or more, depending on the factors mentioned above. In specific embodiments, the dosage may range from 1.0 pg/kg up to about 20 mg/kg, optionally from 10 µg/kg up to about 10 mg/kg or from 100 µg/kg up to about 5 mg/kg.

[0431] A therapeutic effective amount of an antibody construct of the invention preferably results in a decrease in severity of disease symptoms, an increase in frequency or duration of disease symptom-free periods or a prevention of impairment or disability due to the disease affliction. For treating MSLN-expressing tumors, a therapeutically effective amount of the antibody construct of the invention, e.g. an anti-MSLN/anti-CD3 antibody construct, preferably inhibits cell growth or tumor growth by at least about 20%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, or at least about 90% relative to untreated patients. The ability of a compound to inhibit tumor growth may be evaluated in an animal model predictive of efficacy in human tumors.

[0432] The pharmaceutical composition can be administered as a sole therapeutic or in combination with additional therapies such as anti-cancer therapies as needed, e.g. other proteinaceous and non-proteinaceous drugs. These drugs may be administered simultaneously with the composition comprising the antibody construct of the invention as defined herein or separately before or after administration of said antibody construct in timely defined intervals and doses.

[0433] The term "effective and non-toxic dose" as used herein refers to a tolerable dose of an inventive antibody

construct which is high enough to cause depletion of pathologic cells, tumor elimination, tumor shrinkage or stabilization of disease without or essentially without major toxic effects. Such effective and non-toxic doses may be determined e.g. by dose escalation studies described in the art and should be below the dose inducing severe adverse side events (dose limiting toxicity, DLT).

[0434] The term “toxicity” as used herein refers to the toxic effects of a drug manifested in adverse events or severe adverse events. These side events might refer to a lack of tolerability of the drug in general and/or a lack of local tolerance after administration. Toxicity could also include teratogenic or carcinogenic effects caused by the drug.

[0435] The term “safety”, “in vivo safety” or “tolerability” as used herein defines the administration of a drug without inducing severe adverse events directly after administration (local tolerance) and during a longer period of application of the drug. “Safety”, “in vivo safety” or “tolerability” can be evaluated e.g. at regular intervals during the treatment and follow-up period. Measurements include clinical evaluation, e.g. organ manifestations, and screening of laboratory abnormalities. Clinical evaluation may be carried out and deviations to normal findings recorded/coded according to NCI-CTC and/or MedDRA standards. Organ manifestations may include criteria such as allergy/immunology, blood/bone marrow, cardiac arrhythmia, coagulation and the like, as set forth e.g. in the Common Terminology Criteria for adverse events v3.0 (CTCAE). Laboratory parameters which may be tested include for instance hematology, clinical chemistry, coagulation profile and urine analysis and examination of other body fluids such as serum, plasma, lymphoid or spinal fluid, liquor and the like. Safety can thus be assessed e.g. by physical examination, imaging techniques (i.e. ultrasound, x-ray, CT scans, Magnetic Resonance Imaging (MRI), other measures with technical devices (i.e. electrocardiogram), vital signs, by measuring laboratory parameters and recording adverse events. For example, adverse events in non-chimpanzee primates in the uses and methods according to the invention may be examined by histopathological and/or histochemical methods.

[0436] The above terms are also referred to e.g. in the Preclinical safety evaluation of biotechnology-derived pharmaceuticals S6; ICH Harmonised Tripartite Guideline; ICH Steering Committee meeting on Jul. 16, 1997.

[0437] In a further embodiment, the invention provides a kit comprising an antibody construct of the invention, an antibody construct produced according to the process of the invention, a polynucleotide of the invention, a vector of the invention, and/or a host cell of the invention.

[0438] In the context of the present invention, the term “kit” means two or more components—one of which corresponding to the antibody construct, the pharmaceutical composition, the vector or the host cell of the invention—packaged together in a container, recipient or otherwise. A kit can hence be described as a set of products and/or utensils that are sufficient to achieve a certain goal, which can be marketed as a single unit.

[0439] The kit may comprise one or more recipients (such as vials, ampoules, containers, syringes, bottles, bags) of any appropriate shape, size and material (preferably waterproof, e.g. plastic or glass) containing the antibody construct or the pharmaceutical composition of the present invention in an appropriate dosage for administration (see above). The kit may additionally contain directions for use (e.g. in the form

of a leaflet or instruction manual), means for administering the antibody construct of the present invention such as a syringe, pump, infuser or the like, means for reconstituting the antibody construct of the invention and/or means for diluting the antibody construct of the invention.

[0440] The invention also provides kits for a single-dose administration unit. The kit of the invention may also contain a first recipient comprising a dried/lyophilized antibody construct and a second recipient comprising an aqueous formulation. In certain embodiments of this invention, kits containing single-chambered and multi-chambered pre-filled syringes (e.g., liquid syringes and lysyringes) are provided.

[0441] The Figures show:

[0442] FIG. 1:

[0443] Schematic representation of the human-mouse MSLN chimera. The MSLN chimera were generated with 6 distinct sequence stretch exchanges from human to mouse MSLN. Respective variants were expressed on the surface of CHO cell clones (E1-E6). See Example 1.

[0444] FIGS. 2A-2B

[0445] Cross-Reactivity of anti-MSLN antibody constructs as detected by flow cytometry: binding to human and macaque MSLN and CD3. See Example 4.

[0446] FIGS. 3A-3B

[0447] Analysis of anti-MSLN antibody constructs by flow cytometry: binding to human variants v2 and v6. See Example 4.

[0448] FIG. 4

[0449] The potency of MSLNxCD3 bispecific antibody constructs in redirecting stimulated human CD8+ effector T cells against human MSLN-transfected CHO cells was measured in an 18 hour ⁵¹Cr release assay (Effector target ration 10:1).

[0450] FIG. 5

[0451] The potency of MSLNxCD3 bispecific antibody constructs in redirecting stimulated human CD8+ effector T cells against the MSLN positive OVCAR-8 was measured in an 18 hour ⁵¹Cr release assay (Effector target ration 10:1).

[0452] FIG. 6

[0453] The potency of MSLNxCD3 bispecific antibody constructs in redirecting the T cells in unstimulated human PBMC (CD14⁺/CD56⁺) against human MSLN-transfected CHO cells in the absence and presence of soluble MSLN was measured in a 48 hour FACS-based cytotoxicity assay (Effector target ration 10:1).

[0454] FIG. 7

[0455] The potency of MSLNxCD3 bispecific antibody constructs in redirecting the T cells in unstimulated human PBMC (CD14⁺/CD56⁺) against the MSLN-positive human cell line OVCAR-8 was measured in a 48 hour FACS-based cytotoxicity assay.

[0456] FIG. 8

[0457] Confirmation of the cross-reactive MSLNxCD3 bispecific antibody constructs are capable of redirecting macaque T cells against macaque MSLN-transfected CHO cells, a 48 hour FACS-based cytotoxicity assay was performed with a macaque T cell line LnPx4119 as effector T cells (Effector target ration 10:1).

[0458] FIG. 9

[0459] Potency gap between the monomeric and the dimeric isoforms of MSLNxCD3 bispecific antibody constructs in redirecting the T cells in stimulated human CD8+

effector T cells against human MSLN-transfected CHO cells was measured in an 18 hour ^{51}Cr release assay (Effector target ration 10:1).

[0460] FIG. 10

[0461] Stability of MSLNxCD3 bispecific antibody constructs after incubation for 96 hours in human plasma. 18-hour ^{51}Cr based assay. Effector cells: stimulated enriched human CD8 T cells. Target cells: hu MSLN transfected CHO cells. Effector to target cell (E:T) ratio: 10:1. BiTE protein as indicated.

[0462] FIG. 11

[0463] The protein homogeneity of the antibody constructs of the invention analyzed High Resolution Cation Exchange Chromatography CIEC

[0464] FIGS. 12A-12B

[0465] The surface hydrophobicity of bispecific antibody constructs of the invention tested in Hydrophobic Interaction Chromatography HIC in flow-through mode

[0466] FIGS. 13A-13B

[0467] Monomer to dimer conversion after three freeze/thaw cycles.

[0468] FIGS. 14A-14B

[0469] Monomer to dimer conversion after 7 days of incubation at 250 $\mu\text{g}/\text{ml}$

[0470] FIGS. 15A-15B

[0471] Evaluation of Target-independent T Cell Activation by Mesothelin (MS) HLE BiTE® antibody constructs. 2(a) antibody construct of the invention in 48 h activation assay with human PBMC (3 $\times$); HLE BiTE® serial dilutions (start 20 nM; 1:5, 7 $\times$ +blank); w/o or with FcR blocking [10 mg/mL huIgG (Kiovog, Baxter)]; FACS measurement of CD69 and CD25 [not shown] expression on CD4 $^{+}$, CD8 $^{+}$ T cells. 2(b) Hetero-Fc antibody construct in 48 h activation assay with human PBMC and CD14 $^{+}$ /CD33 $^{+}$ cell depleted PBMC (3 $\times$); HLE BiTE® serial dilutions (start 20 nM; 1:5, 7 $\times$ +blank); FACS measurement of CD69 and CD25 [not shown] expression on CD4 $^{+}$, CD8 $^{+}$ T cells.

[0472] FIGS. 16A-16C

[0473] Evaluation of Target-independent T Cell Activation by CDH19 HLE BiTE® antibody constructs. 3(a) antibody construct of the invention in 48 h activation assay with human PBMC (3 $\times$); HLE BiTE® serial dilutions (start 20 nM; 1:5, 7 $\times$ +blank); w/o or with FcR blocking [10 mg/mL huIgG (Kiovog, Baxter)]; FACS measurement of CD69 and CD25 [not shown] expression on CD4 $^{+}$, CD8 $^{+}$ T cells. 3(b) Hetero-Fc antibody construct in 48 h activation assay with human PBMC and CD14 $^{+}$ /CD33 $^{+}$ cell depleted PBMC (3 $\times$); HLE BiTE® serial dilutions (start 20 nM; 1:5, 7 $\times$ +blank); FACS measurement of CD69 and CD25 [not shown] expression on CD4 $^{+}$, CD8 $^{+}$ T cells. 3(c) X-body construct in 48 h activation assay with human PBMC and CD14 $^{+}$ /CD33 $^{+}$ cell depleted PBMC (3 $\times$); HLE BiTE® serial dilutions (start 20 nM; 1:5, 7 $\times$ +blank); FACS measurement of CD69 and CD25 [not shown] expression on CD4 $^{+}$, CD8 $^{+}$ T cells.

[0474] FIG. 17

[0475] Complement C1q Binding of BiTE® Fc fusion antibody constructs. BiTE® Fc fusion antibody constructs (BiTE® single chain Fc (triangle), BiTE® hetero Fc (squares), canonical BiTE® (circle)) were coated on a Maxisorp plate (in dilution series), prior to incubation with pooled human serum and incubation with polyclonal anti human CC1q murine antibody, visualized by goat anti-mouse Fc-AP conjugate.

[0476] FIG. 18

[0477] Mean PK profiles of four pairs of BiTE®-HLE fusion proteins after single dose administration in cynomolgus monkeys. For reasons of comparability, serum concentrations were dose-normalized to 15 $\mu\text{g}/\text{kg}$ and indicated in nmol.

[0478] FIG. 19

[0479] Mean PK profiles of five different BiTE® antibody constructs, each fused to a scFc half-life extending moiety. For reasons of comparability, serum concentrations were dose-normalized to 15 $\mu\text{g}/\text{kg}$ and indicated in nmol.

EXAMPLES

[0480] The following examples illustrate the invention. These examples should not be construed as to limit the scope of this invention. The present invention is limited only by the claims.

Example 1

Generation of CHO Cells Expressing Wild Type and Chimeric MSLN

[0481] The MSLN antigen can be subdivided into six different sub-domains or regions that are defined, for the purposes of Examples 1 and 2. The aa sequence of those six sub-domains is depicted in SEQ ID NOs: 238-243.

[0482] The following molecules were generated; see also FIG. 1:

hu orl MSLN-E1 mu	SEQ ID NO: 246
hu orl MSLN-E2 mu	SEQ ID NO: 247
hu orl MSLN-E3 mu	SEQ ID NO: 248
hu orl MSLN-E4 mu	SEQ ID NO: 249
hu orl MSLN-E5 mu	SEQ ID NO: 250
hu orl MSLN-E6 mu	SEQ ID NO: 251
hu orl MSLN-full human	SEQ ID NO: 231

[0483] For the generation of CHO cells expressing human, cynomolgus macaque ("cyno") and truncated human N-terminal V5 tagged MSLN, the respective coding sequences for human MSLN (SEQ ID NO: 231; see also GeneBank accession number NM_005823), cyno MSLN (SEQ ID NO: 234, see LMR C52457) and the six chimeric human/murine MSLN versions (see above) were cloned into a plasmid designated pEF-DHFR (pEF-DHFR is described in Raum et al. Cancer Immunol Immunother 50 (2001) 141-150). For cell surface expression of human and cyno MSLN the original signal peptide was used. All cloning procedures were carried out according to standard protocols (Sambrook, Molecular Cloning; A Laboratory Manual, 3rd edition, Cold Spring Harbour Laboratory Press, Cold Spring Harbour, New York (2001)). For each construct, a corresponding plasmid was transfected into DHFR deficient CHO cells for eukaryotic expression, as described by Kaufman R.J. (1990) Methods Enzymol. 185, 537-566.

[0484] The expression of human, chimeric and cyno MSLN on CHO cells was verified in a FACS assay using a monoclonal mouse IgG2b anti-human MSLN antibody. As negative control, cells were incubated with isotype control antibody instead of the first antibody. The samples were measured by flow cytometry.

Example 2

Epitope Mapping of Anti-MSLN Antibody Constructs

[0485] Cells transfected with human MSLN and with the chimeric human MSLN molecules (see Example 1) were stained with crude, undiluted periplasmic extract containing bispecific MSLNxCD3 antibody constructs (with the CD3 binding domain being denominated 12C) fused to a human albumin (variant 1), in PBS/1.5% FCS. Bound molecules were detected with an in-house mouse monoclonal anti-CD3 binding domain antibody (50 µl) followed by an anti-mouse IgG Fc-gamma-PE (1:100, 50 µl; Jackson ImmunoResearch #115-116-071) All antibodies were diluted in PBS/1.5% FCS. As negative control, cells were incubated with PBS/2% FCS instead of the periplasmic extract. The samples were measured by flow cytometry.

[0486] The regions that were recognized by the respective MSLN binding domains are indicated in the sequence table (Table 2). Loss of the FACS signal in the respective chimeric MSLN constructs comprising the murine epitope cluster was the read out for the relevance of the respective cluster for the binding. In case the FACS signal was negatively impacted for more than one, respectively two, chimeric MSLN clone, it was concluded that both cluster are relevant.

TABLE 2

Mapping of the epitope clusters for MS-1 to MS-8	
Epitope cluster	Binder
E 1 + 2	MS-1
E 2 + 3	MS-2
E 2 + 3	MS-3
E 2 + 3	MS-4
E 2 + 3	MS-5
E 2 + 3	MS-6
E 4	MS-7
E 1 + 2	MS-8

Example 3

Scatchard-Based Analysis of MSLNxCD3 Bispecific Antibody Construct Affinity to Human and Macaque MSLN on Target Antigen Positive Cells and Determination of the Interspecies Affinity Gap

[0487] The affinities of MSLNxCD3 bispecific antibody constructs to CHO cells transfected with human or macaque MSLN were also determined by Scatchard analysis as the most reliable method for measuring potential affinity gaps between human and macaque MSLN. For the Scatchard analysis, saturation binding experiments are performed using a monovalent detection system to precisely determine monovalent binding of the MSLNxCD3 bispecific antibody constructs to the respective cell line.

[0488] 2×10^4 cells of the respective cell line (recombinantly human MSLN-expressing CHO cell line, recombinantly macaque MSLN-expressing CHO cell line) were incubated each with 50 µl of a triplet dilution series (twelve dilutions at 1:2) of the respective MSLNxCD3 bispecific antibody construct (until saturation is reached) starting at 10-20 nM followed by 16 h incubation at 4° C. under agitation and one residual washing step. Then, the cells were incubated for another hour with 30 µl of a CD3xALEXA488 conjugate solution. After one washing step, the cells were

resuspended in 150 µl FACS buffer containing 3.5% formaldehyde, incubated for further 15 min, centrifuged, resuspended in FACS buffer and analyzed using a FACS CantoII machine and FACS Diva software. Data were generated from two independent sets of experiments, each using triplicates. Respective Scatchard analysis was calculated to extrapolate maximal binding (Bmax). The concentrations of MSLNxCD3 bispecific antibody constructs at half-maximal binding were determined reflecting the respective KDs. Values of triplicate measurements were plotted as hyperbolic curves and as S-shaped curves to demonstrate proper concentration ranges from minimal to optimal binding.

[0489] Values depicted in Table 3 were derived from two independent experiments per MSLNxCD3 bispecific antibody construct. Cell based Scatchard analysis confirmed that the MSLNxCD3 bispecific antibody constructs of the invention are subnanomolar in affinity to human MSLN and to mac MSLN and present with a small cyno/human interspecies affinity gap of around 1.

TABLE 3

Affinities (KD) of MSLN x CD3 bispecific antibody constructs as determined in cell based Scatchard analysis with the calculated affinity gap KD macaque MSLN/KD human MSLN. Antibody constructs were measured in two independent experiments, each using triplicates.			
BiTE antibody MS x 12C-HALB	Cell based affinity hu MSLN* [nM]	Cell based affinity mac MSLN* [nM]	Affinity gap KD ^{mac} /KD ^{hu} MSLN
MS-1	1.25 ± 0.8	13.86 ± 2.9	11.08
MS-8	4.68 ± 0.4	85.86 ± 16.7	18.35
MS-2	1.40 ± 0.9	10.16 ± 3.5	7.28
MS-3	0.74 ± 0.8	0.82 ± 0.5	1.12
MS-4	0.91 ± 1.0	0.77 ± 0.7	0.85
MS-5	0.39 ± 0.3	2.31 ± 3.7	5.91
MS-6	0.60 ± 0.3	9.03 ± 3.7	15.04
MS-7	1.06 ± 0.5	4.07 ± 1.0	3.86

Example 4

Bispecific Binding and Interspecies Cross-Reactivity

[0490] For confirmation of binding to human MSLN and CD3 and to cyno MSLN and CD3, bispecific antibody constructs of the invention were tested by flow cytometry using

[0491] CHO cells transfected with human MSLN, with human MSLN isoform (NM_013404=SEQ ID NO:232 and AY743922=SEQ ID NO:233), and with macaque MSLN, respectively,

[0492] the human MSLN positive human cell line OVCAR-8,

[0493] CD3-expressing human T cell leukemia cell line HPB-all (DSMZ, Braunschweig, ACC483), and

[0494] the cynomolgus CD3-expressing T cell line LnPx 4119

[0495] For flow cytometry 200,000 cells of the respective cell lines were incubated for 60 min at 4° C. with 50 µl of purified bispecific antibody construct at a concentration of 5 µg/ml. The cells were washed twice in PBS/2% FCS and then incubated with an in-house mouse antibody (2 µg/ml) specific for the CD3 binding part of the bispecific antibody constructs for 30 min at 4° C. After washing, bound mouse antibodies were detected with a goat anti-mouse Fcy-PE

(1:100) for 30 min at 4° C. Samples were measured by flow cytometry. Non-transfected CHO cells were used as negative control.

[0496] The results are shown in FIGS. 2 and 3. The MSLNxCD3 bispecific antibody constructs of the invention stained CHO cells transfected with human MSLN, the artificial MSLN isoform and with cyno MSLN, and they also stained the human MSLN positive human cell line OVCAR-8 (natural expresser). Human and cyno T cell lines expressing CD3 were also recognized by the bispecific antibody constructs. Moreover, there was no staining of the negative control cells (non-transfected CHO).

Example 5

Cytotoxic Activity

[0497] The potency of MSLNxCD3 bispecific antibody constructs of the invention in redirecting effector T cells against MSLN-expressing target cells was analyzed in five in vitro cytotoxicity assays:

[0498] The potency of MSLNxCD3 bispecific antibody constructs in redirecting stimulated human CD8+ effector T cells against human MSLN-transfected CHO cells was measured in an 18 hour ⁵¹Cr release assay (Effector target ration 10:1). FIG. 4 and Table 4

[0499] The potency of MSLNxCD3 bispecific antibody constructs in redirecting stimulated human CD8+ effector T cells against the MSLN positive cell line OVCAR-8 was measured in an 18 hour ⁵¹Cr release assay (Effector target ration 10:1). FIG. 5 Table 5

[0500] The potency of MSLNxCD3 bispecific antibody constructs in redirecting the T cells in unstimulated human PBMC (CD14⁻/CD56⁻) against human MSLN-transfected CHO cells in the absence and presence of soluble MSLN was measured in a 48 hour FACS-based cytotoxicity assay (Effector target ration 10:1). FIG. 6 and Table 6

[0501] The potency of MSLNxCD3 bispecific antibody constructs in redirecting the T cells in unstimulated human PBMC (CD14⁻/CD56⁻) against the MSLN-positive human cell line OVCAR-8 was measured in a 48 hour FACS-based cytotoxicity assay. FIG. 7

[0502] For confirmation that the cross-reactive MSLNxCD3 bispecific antibody constructs are capable of redirecting macaque T cells against macaque MSLN-transfected CHO cells, a 48 hour FACS-based cytotoxicity assay was performed with a macaque T cell line LnPx4119 as effector T cells (Effector target ration 10:1). FIG. 8

[0503] Potency gap between the monomeric and the dimeric isoforms of MSLNxCD3 bispecific antibody constructs in redirecting the T cells in stimulated human CD8+effector T cells against human MSLN-transfected CHO cells was measured in an 18 hour ⁵¹Cr release assay (Effector target ration 10:1). FIG. 9

Example 5.1

Chromium Release Assay with Stimulated Human T Cells

[0504] Stimulated T cells enriched for CD8+ T cells were obtained as described in the following. A petri dish (145 mm diameter, Greiner bio-one GmbH, Kremsmünster) was coated with a commercially available anti-CD3 specific antibody (OKT3, Orthoclone) in a final concentration of 1

µg/ml for 1 hour at 37° C. Unbound protein was removed by one washing step with PBS. 3-5×10⁷ human PBMC were added to the precoated petri dish in 120 ml of RPMI 1640 with stabilized glutamine/10% FCS/IL-2 20 U/ml (Pro-leukin®, Chiron) and stimulated for 2 days. On the third day, the cells were collected and washed once with RPMI 1640. IL-2 was added to a final concentration of 20 U/ml and the cells were cultured again for one day in the same cell culture medium as above. CD8⁺ cytotoxic T lymphocytes (CTLs) were enriched by depletion of CD4⁺ T cells and CD56+NK cells using Dynal-Beads according to the manufacturer's protocol.

[0505] Cyno MSLN- or human MSLN-transfected CHO target cells were washed twice with PBS and labeled with 11.1 MBq ⁵¹Cr in a final volume of 100 µl RPMI with 50% FCS for 60 minutes at 37° C. Subsequently, the labeled target cells were washed 3 times with 5 ml RPMI and then used in the cytotoxicity assay. The assay was performed in a 96-well plate in a total volume of 200 µl supplemented RPMI with an E:T ratio of 10:1. A starting concentration of 0.01-1 µg/ml of purified bispecific antibody construct and threefold dilutions thereof were used.

[0506] Incubation time for the assay was 18 hours. Cytotoxicity was determined as relative values of released chromium in the supernatant relative to the difference of maximum lysis (addition of Triton-X) and spontaneous lysis (without effector cells). All measurements were carried out in quadruplicates. Measurement of chromium activity in the supernatants was performed in a Wizard 3" gamma counter (Perkin Elmer Life Sciences GmbH, Köln, Germany). Analysis of the results was carried out with Prism 5 for Windows (version 5.0, GraphPad Software Inc., San Diego, California, USA). EC50 values calculated by the analysis program from the sigmoidal dose response curves were used for comparison of cytotoxic activity.

Example 5.2

Potency of Redirecting Stimulated Human Effector T Cells Against Human MSLN-Transfected CHO Cells

[0507] The cytotoxic activity of MSLNxCD3 bispecific antibody constructs according to the invention was analyzed in a 51-chromium (⁵¹Cr) release cytotoxicity assay using CHO cells transfected with human MSLN as target cells, and stimulated human CDB+ T cells as effector cells. The experiment was carried out as described in Example 8.1.

[0508] The results are shown in Table 4. The MSLNxCD3 bispecific antibody constructs showed very potent cytotoxic activity against human MSLN transfected CHO cells in the 1-digit picomolar range.

TABLE 4

EC50 values [pM] of MSLN × CD3 bispecific antibody constructs analyzed in a 51-chromium (⁵¹ Cr) release cytotoxicity assay using CHO cells transfected with human MSLN as target cells, and stimulated human CD8 T cells as effector cells.	
BiTE × I2C-HALB	EC50 [pM]
MS 1	1.5
MS 8	6.5
MS 2	2.0
MS 3	0.2
MS 4	0.1

TABLE 4-continued

EC50 values [pM] of MSLN × CD3 bispecific antibody constructs analyzed in a 51-chromium (⁵¹ Cr) release cytotoxicity assay using CHO cells transfected with human MSLN as target cells, and stimulated human CD8 T cells as effector cells.	
BiTE × I2C-HALB	EC50 [pM]
MS 5	0.2
MS 6	0.5
MS 7	6.0

Example 5.3

Potency of Redirecting Stimulated Human Effector T Cells Against the MSLN Positive Human Cell Line OVCAR-8

[0509] The cytotoxic activity of MSLN×CD3 bispecific antibody constructs was analyzed in a 51-chromium (⁵¹Cr) release cytotoxicity assay using the MSLN-positive human cell line OVCAR-8 as source of target cells, and stimulated human CD8+ T cells as effector cells. The assay was carried out as described in Example 8.1.

[0510] In accordance with the results of the 51-chromium release assays with stimulated enriched human CD8+ T lymphocytes as effector cells and human MSLN-transfected CHO cells as target cells, MSLN×CD3 bispecific antibody constructs of the present invention are also potent in cytotoxic activity against natural expresser target cells (see Table 5).

TABLE 5

EC50 values [pM] of MSLN × CD3 bispecific antibody constructs analyzed in an 18-hour 51-chromium (⁵¹ Cr) release cytotoxicity assay with the MSLN-positive human cell line OVCAR-8 as source of target cells, and stimulated enriched human CD8 T cells as effector cells.	
BiTE × I2C-HALB	EC50 [pM]
MS 1	0.2
MS 8	1.3
MS 2	0.8
MS 3	0.1
MS 4	0.1
MS 5	0.3
MS 6	0.2
MS 7	0.5

Example 5.4

FACS-Based Cytotoxicity Assay with Unstimulated Human PBMC Isolation of Effector Cells

[0511] Human peripheral blood mononuclear cells (PBMC) were prepared by Ficoll density gradient centrifugation from enriched lymphocyte preparations (buffy coats), a side product of blood banks collecting blood for transfusions. Buffy coats were supplied by a local blood bank and PBMC were prepared on the same day of blood collection. After Ficoll density centrifugation and extensive washes with Dulbecco's PBS (Gibco), remaining erythrocytes were removed from PBMC via incubation with erythrocyte lysis buffer (155 mM NH₄Cl, 10 mM KHCO₃, 100 pM EDTA). Platelets were removed via the supernatant upon centrifugation of PBMC at 100×g. Remaining lymphocytes mainly

encompass B and T lymphocytes, NK cells and monocytes. PBMC were kept in culture at 37° C./5% CO₂ in RPMI medium (Gibco) with 10% FCS (Gibco).

Depletion of CD14⁺ and CD56⁺ Cells

[0512] For depletion of CD14⁺ cells, human CD14 MicroBeads (Milteny Biotec, MACS, #130-050-201) were used, for depletion of NK cells human CD56 MicroBeads (MACS, #130-050-401). PBMC were counted and centrifuged for 10 min at room temperature with 300 ×g. The supernatant was discarded and the cell pellet resuspended in MACS isolation buffer [80 μL/10⁷ cells; PBS (Invitrogen, #20012-043), 0.5% (v/v) FBS (Gibco, #10270-106), 2 mM EDTA (Sigma-Aldrich, #E-6511)]. CD14 MicroBeads and CD56 MicroBeads (20 μL/10⁷ cells) were added and incubated for 15 min at 4-8° C. The cells were washed with MACS isolation buffer (1-2 mL/10⁷ cells). After centrifugation (see above), supernatant was discarded and cells resuspended in MACS isolation buffer (500 μL/10⁸ cells). CD14/CD56 negative cells were then isolated using LS Columns (Milteny Biotec, #130-042-401). PBMC w/o CD14+/CD56+cells were cultured in RPMI complete medium i.e. RPMI1640 (Biochrom AG, #FG1215) supplemented with 10% FBS (Biochrom AG, #S0115), 1× non-essential amino acids (Biochrom AG, #K0293), 10 mM Hepes buffer (Biochrom AG, #L1613), 1 mM sodium pyruvate (Biochrom AG, #L0473) and 100 U/mL penicillin/streptomycin (Biochrom AG, #A2213) at 37° C. in an incubator until needed.

Target Cell Labeling

[0513] For the analysis of cell lysis in flow cytometry assays, the fluorescent membrane dye DiOC₁₈ (DiO) (Molecular Probes, #V22886) was used to label human MSLN- or macaque MSLN-transfected CHO cells as target cells and distinguish them from effector cells. Briefly, cells were harvested, washed once with PBS and adjusted to 10⁶ cell/mL in PBS containing 2% (v/v) FBS and the membrane dye DiO (5 μL/10⁶ cells). After incubation for 3 min at 37° C., cells were washed twice in complete RPMI medium and the cell number adjusted to 1.25×10⁵ cells/mL. The vitality of cells was determined using 0.5% (v/v) isotonic EosinG solution (Roth, #45380).

Flow Cytometry Based Analysis

[0514] This assay was designed to quantify the lysis of cyno or human MSLN-transfected CHO cells in the presence of serial dilutions of MSLN bispecific antibody constructs. Equal volumes of DiO-labeled target cells and effector cells (i.e., PBMC w/o CD14⁺ cells) were mixed, resulting in an E:T cell ratio of 10:1. 160 μL of this suspension were transferred to each well of a 96-well plate. 40 μL of serial dilutions of the MSLN×CD3 bispecific antibody constructs and a negative control bispecific (a CD3-based bispecific antibody construct recognizing an irrelevant target antigen) or RPMI complete medium as an additional negative control were added. The bispecific antibody-mediated cytotoxic reaction proceeded for 48 hours in a 7% CO₂ humidified incubator. Then cells were transferred to a new 96-well plate and loss of target cell membrane integrity was monitored by adding propidium iodide (PI) at a final concentration of 1 μg/mL. PI is a membrane impermeable dye

that normally is excluded from viable cells, whereas dead cells take it up and become identifiable by fluorescent emission.

[0515] Samples were measured by flow cytometry on a FACSCanto II instrument and analyzed by FACSDiva software (both from Becton Dickinson). Target cells were identified as DiO-positive cells. PI-negative target cells were classified as living target cells. Percentage of cytotoxicity was calculated according to the following formula:

$$\text{Cytotoxicity [\%]} = \frac{n_{\text{dead target cells}}}{n_{\text{target cells}}} \times 100$$

n = number of events

[0516] Using GraphPad Prism 5 software (Graph Pad Software, San Diego), the percentage of cytotoxicity was plotted against the corresponding bispecific antibody construct concentrations. Dose response curves were analyzed with the four parametric logistic regression models for evaluation of sigmoid dose response curves with fixed hill slope and EC50 values were calculated.

Example 5.5

Potency of Redirecting Unstimulated Human PBMC Against Human MSLN-Transfected CHO Cells in Absence and Presence of Soluble MSLN

[0517] The cytotoxic activity of MSLNxCD3 bispecific antibody constructs was analyzed in a FACS-based cytotoxicity assay using CHO cells transfected with human MSLN as target cells, and unstimulated human PBMC as effector cells. The assay was carried out as described in Example 8.4 above.

[0518] The results of the FACS-based cytotoxicity assays with unstimulated human PBMC as effector cells and human MSLN-transfected CHO cells as targets are shown in Table 7.

TABLE 6

Cytotoxic Activity of Unstimulated Human PBMC Against Human MSLN-transfected CHO Cells in absence and presence of soluble MSLN			
BiTE × I2C-HALB	w/o sMSLN EC50 [pM]	with 50 ng/ml sMSLN EC50 [pM]	with 400 ng/ml sMSLN EC50 [pM]
MS-1	12	53	215
MS-8	3.4	8.3	39
MS-2	2.4	2.9	6.3
MS-3	1.2	3.0	16
MS-4	1.5	3.9	11
MS-5	1.0	2.2	11
MS-6	1.1	2.6	12
MS-7	8.1	28	166

[0519] Expectedly, EC50 values were generally higher in cytotoxicity assays with unstimulated PBMC as effector cells compared with cytotoxicity assays using stimulated human CD8+ T cells (see Example 8.2).

Example 5.6

Potency of Redirecting Unstimulated Human PBMC Against the MSLN-Positive Cell Line OVCAR-8 Cells

[0520] The cytotoxic activity of MSLNxCD3 bispecific antibody constructs was furthermore analyzed in a FACS-based cytotoxicity assay using the MSLN-positive human cell line OVCAR-8 as a source of target cells and unstimulated human PBMC as effector cells. The assay was carried out as described in Example 8.4 above. The results are shown in Table 7.

TABLE 7

EC50 values [pM] of MSLN × CD3 bispecific antibody constructs as measured in a 48-hour FACS-based cytotoxicity assay with unstimulated human PBMC as effector cells and the human cell line OVCAR-8 as source of target cells.	
BiTE × I2C-HALB	EC50 [pM]
MS 1	5.9
MS 8	13
MS 2	18
MS 3	1.5
MS 4	2.0
MS 5	1.9
MS 6	2.3
MS 7	9.7

Example 5.7

Potency of Redirecting Macaque T Cells Against Macaque MSLN-Expressing CHO Cells

[0521] The cytotoxic activity of MSLNxCD3 bispecific antibody constructs was analyzed in a FACS-based cytotoxicity assay using CHO cells transfected with macaque (cyno) MSLN as target cells, and the macaque T cell line 4119LnPx (Knappe et al. Blood 95:3256-61 (2000)) as source of effector cells. Target cell labeling of macaque MSLN-transfected CHO cells and flow cytometry based analysis of cytotoxic activity was performed as described above.

[0522] Results are shown in Table 8. Macaque T cells from cell line 4119LnPx were induced to efficiently kill macaque MSLN-transfected CHO cells by MSLNxCD3 bispecific antibody constructs of the invention.

TABLE 8

EC50 values [pM] of MSLN × CD3 bispecific as measured in a 48-hour FACS-based cytotoxicity assay with macaque T cell line 4119LnPx as effector cells and CHO cells transfected with macaque MSLN as target cells.	
BiTE × I2C-HALB	EC50 [pM]
MS 1	434
MS 8	1589
MS 2	583
MS 3	56
MS 4	67
MS 5	85
MS 6	100
MS 7	1791

Example 5.8

Potency Gap Between the Monomeric and the Dimeric Isoform of Bispecific Antibody Constructs

[0523] In order to determine the difference in cytotoxic activity between the monomeric and the dimeric isoform of individual MSLN $\times$ CD3 bispecific antibody constructs (referred to as potency gap), an 18 hour 51-chromium release cytotoxicity assay was carried out as described hereinabove (Example 8.1) with purified bispecific antibody construct monomer and dimer. Effector cells were stimulated enriched human CD8 $^{+}$ T cells. Target cells were hu MSLN transfected CHO cells. Effector to target cell (E:T) ratio was 10:1. The potency gap was calculated as ratio between EC50 values.

[0524] The results of the assays with stimulated enriched human CD $^{+}$ T cells as effector cells and human MSLN-transfected CHO cells as targets are shown in Table 9.

TABLE 9

Cytotoxic Activity of Unstimulated Human PBMC Against Human MSLN-transfected CHO Cells using monomeric and dimeric MSLN $\times$ CD3 bispecific antibody constructs			
BiTE $\times$ I2C-HALB	EC ₅₀ [pM]		Monomer to dimer ratio (EC ₅₀ Monomer/ EC ₅₀ Dimer)
	Monomer	Dimer	
MS 1	1.5	1.7	0.9
MS 8	6.5	9.5	0.7
MS 2	2.0	2.0	1.0
MS 3	0.2	0.7	0.3
MS 4	0.1	0.9	0.1
MS 5	0.2	1.1	0.2
MS 6	0.5	3.4	0.1
MS 7	6.0	21	0.3

Example 6

Stability after Incubation for 24 Hours in Human Plasma

[0525] Purified bispecific antibody constructs were incubated at a ratio of 1:5 in a human plasma pool at 37° C. for 96 hours at a final concentration of 2-20 pg/ml. After plasma incubation the antibody constructs were compared in a 51-chromium release assay with stimulated enriched human CD8 $^{+}$ T cells and human MSLN-transfected CHO cells at a starting concentration of 0.01-0.1 μ g/ml and with an effector to target cell (E:T) ratio of 10:1 (assay as described in Example 8.1). Non-incubated, freshly thawed bispecific antibody constructs were included as controls.

[0526] The results are shown in Table 10 below; All tested antibody constructs had a very favourable plasma stability (EC₅₀ plasma/EC₅₀ control) of ≤ 4 .

TABLE 10

EC50 values of the antibody constructs with and without plasma incubation and calculated plasma/control value			
BiTE $\times$ I2C-HALB	EC ₅₀ [pM]		Plasma to Control Ratio (EC ₅₀ Plasma/ EC ₅₀ Control)
	w/ Plasma	w/o Plasma	
MS 1	3.0	1.5	2.0
MS 8	4.8	6.5	0.7

TABLE 10-continued

EC50 values of the antibody constructs with and without plasma incubation and calculated plasma/control value			
BiTE $\times$ I2C-HALB	EC ₅₀ [pM]		Plasma to Control Ratio (EC ₅₀ Plasma/ EC ₅₀ Control)
	w/ Plasma	w/o Plasma	
MS 2	6.7	2.0	3.4
MS 3	0.3	0.2	1.5
MS 4	0.4	0.1	4
MS 5	0.3	0.2	1.5
MS 6	0.6	0.5	1.2
MS 7	11.4	6.0	1.9

Example 7

Protein Homogeneity by High Resolution Cation Exchange Chromatography

[0527] The protein homogeneity the antibody constructs of the invention was analyzed by high resolution cation exchange chromatography CIEC.

[0528] 50 μ g of antibody construct monomer were diluted with 50 ml binding buffer A (20 mM sodium dihydrogen phosphate, 30 mM NaCl, 0.01% sodium octanate, pH 5.5), and 40 ml of this solution were applied to a 1 ml BioPro SP-F column (YMC, Germany) connected to an Äkta Micro FPLC device (GE Healthcare, Germany). After sample binding, a wash step with further binding buffer was carried out. For protein elution, a linear increasing salt gradient using buffer B (20 mM sodium dihydrogen phosphate, 1000 mM NaCl, 0.01% sodium octanate, pH 5.5) up to 50% percent buffer B was applied over 10 column volumes. The whole run was monitored at 280, 254 and 210 nm optical absorbance. Analysis was done by peak integration of the 280 nm signal recorded in the Äkta Unicomp software run evaluation sheet.

[0529] The results are shown in Table 11 below. Almost all tested antibody constructs have a very favourable homogeneity of $\geq 80\%$ (area under the curve (=AUC) of the main peak). The only exception is MS-2 $\times$ CD3-HALB bispecific construct with merely 67% homogeneity.

TABLE 11

Protein homogeneity of the antibody constructs (% AUC of main peak)	
MSLN HALB BiTE	AUC of Main Peak [%]
MS 1	100
MS 8	93
MS 2	67
MS 3	87
MS 4	80
MS 5	89
MS 6	81
MS 7	100

Example 8

Surface Hydrophobicity as Measured by HIC Butyl

[0530] The surface hydrophobicity of bispecific antibody constructs of the invention was tested in Hydrophobic Interaction Chromatography HIC in flow-through mode.

[0531] 50 µg of antibody construct monomer were diluted with generic formulation buffer to a final volume of 500 µl (10 mM citric acid, 75 mM lysine HCl, 4% trehalose, pH 7.0) and applied to a 1 ml Butyl Sepharose FF column (GE Healthcare, Germany) connected to a Äkta Purifier FPLC system (GE Healthcare, Germany). The whole run was monitored at 280, 254 and 210 nm optical absorbance. Analysis was done by peak integration of the 280 nm signal recorded in the Äkta Unicom software run evaluation sheet. Elution behavior was evaluated by comparing area and velocity of rise and decline of protein signal thereby indicating the strength of interaction of the BiTE albumin fusion with the matrix.

[0532] The antibody constructs had a good elution behaviour, which was mostly rapid and complete; see Table 12.

TABLE 12

The surface hydrophobicity of bispecific antibody constructs	
MSLN HALB BiTE	HIC Butyl Elution behavior
MS 1	1
MS 8	1
MS 2	1
MS 3	1
MS 4	1
MS 5	1
MS 6	1
MS 7	1

Example 9

Monomer to Dimer Conversion after (I) Three Freeze/Thaw Cycles and (II) 7 Days of Incubation at 250 µg/ml

[0533] Bispecific MSLNxCD3 antibody monomeric construct were subjected to different stress conditions followed by high performance SEC to determine the percentage of initially monomeric antibody construct, which had been converted into dimeric antibody construct.

[0534] (i) 25 µg of monomeric antibody construct were adjusted to a concentration of 250 µg/ml with generic formulation buffer and then frozen at -80° C. for 30 min followed by thawing for 30 min at room temperature. After three freeze/thaw cycles the dimer content was determined by HP-SEC.

[0535] (ii) 25 µg of monomeric antibody construct were adjusted to a concentration of 250 µg/ml with generic formulation buffer followed by incubation at 37° C. for 7 days. The dimer content was determined by HP-SEC.

[0536] A high resolution SEC Column TSK Gel G3000 SWXL (Tosoh, Tokyo-Japan) was connected to an Äkta Purifier 10 FPLC (GE Lifesciences) equipped with an A905 Autosampler. Column equilibration and running buffer consisted of 100 mM KH₂PO₄—200 mM Na₂SO₄ adjusted to pH 6.6. The antibody solution (25 µg protein) was applied to the equilibrated column and elution was carried out at a flow rate of 0.75 ml/min at a maximum pressure of 7 MPa. The whole run was monitored at 280, 254 and 210 nm optical absorbance. Analysis was done by peak integration of the 210 nm signal recorded in the Äkta Unicom software run evaluation sheet. Dimer content was calculated by dividing the area of the dimer peak by the total area of monomer plus dimer peak.

[0537] The results are shown in Table 13 below. The MSLNxCD3 bispecific antibody constructs of the invention presented with dimer percentages of 0.0% after three freeze/thaw cycles, and with dimer percentages of ≤2.2% after 7 days of incubation at 37° C.

TABLE 13

Percentage of monomeric versus dimeric MSLN x CD3 bispecific antibody constructs as determined by High Performance Size Exclusion Chromatography (HP-SEC).	
MSLN HALB BiTE	BiTE Dimer Conversion [%]
MS 1	0.2
MS 8	1.6
MS 2	0.0
MS 3	1.7
MS 4	0.3
MS 5	1.2
MS 6	0.4
MS 7	2.2

Example 10

Thermostability

[0538] Antibody aggregation temperature was determined as follows: 40 µl of antibody construct solution at 250 µg/ml were transferred into a single use cuvette and placed in a Wyatt Dynamic Light Scattering device DynaPro Nanostar (Wyatt). The sample was heated from 40° C. to 70° C. at a heating rate of 0.5° C./min with constant acquisition of the measured radius. Increase of radius indicating melting of the protein and aggregation was used by the software package delivered with the DLS device to calculate the aggregation temperature of the antibody construct.

[0539] All tested MSLNxCD3 bispecific antibody constructs of the invention showed thermal stability with aggregation temperatures 249° C., as shown in Table 14 below. The group of antibody constructs binding to epitope cluster 2+3 even had a thermal stability of 251° and up to more than 56° C.

TABLE 14

Thermostability of the bispecific antibody constructs as determined by DLS (dynamic light scattering)	
MSLN HALB BiTE	Thermal Stability DLS T _d [° C.]
MS 1	49.3
MS 8	51.0
MS 2	53.1
MS 3	51.6
MS 4	56.5
MS 5	54.4
MS 6	53.9
MS 7	51.4

Example 11

Turbidity at 2500 µg/ml Antibody Concentration

[0540] 1 ml of purified antibody construct solution of a concentration of 250 µg/ml was concentrated by spin concentration units to 2500 µg/ml. After 16h storage at 5° C. the

turbidity of the antibody solution was determined by OD340 nm optical absorption measurement against the generic formulation buffer.

[0541] The results are shown in Table 15 below. All tested antibody constructs have a very favourable turbidity of ≤ 0.09 .

TABLE 15

Turbidity of the antibody constructs after concentration to 2.5 mg/ml over night	
MSLN HALB BiTE	Turbidity after 16 h @ 2500 μ g/ml [OD340]
MS 1	0.080
MS 8	0.079
MS 2	0.070
MS 3	0.087
MS 4	0.076
MS 5	0.071
MS 6	0.070
MS 7	0.066

Example 12: BiTE® Induced CD69 Expression on T Cells in Absence of Target Cells

[0542] Isolated PBMC from healthy human donors were cultured with increasing CDH19/CD3 or MSLN/CD3 HLE bispecific antibody constructs for 48 h. The expression of the activation marker CD69 on T cells was determined by immunostaining and flow cytometry and antigen specific conjugates mAb.

[0543] Target-independent T cell activation in terms of CD69 upregulation was observed for all anti-CDH 19 constructs but was most pronounced for heteroFc and crossbody molecules. Upregulation of CD69 by antiCDH19-scFc occurred at higher concentrations and the amplitude was in part lower compared to the other two Fc-based constructs.

[0544] For the anti-MSLN almost no target-independent T cell activation was observed for the scFc-containing molecule, while the heteroFc construct induced a strong upregulation of CD69 on the cell surface T cells in the absence of target cells.

[0545] Target-independent T cell activation induced by BiTE® constructs containing a single chain-Fc, or hetero-Fc fusion at the C-terminus was evaluated for the following constructs:

[0546] BiTE® constructs (serial dilutions: 0.1 pM-2 μ M)

[0547] a. MSLN scFc; 1.14 mg/mL;

[0548] b. MSLN Hetero Fc; 1.02 mg/

[0549] Human PBMC effector cells (3 donors; #065, #823, #836 (scFc) #401, #415, #433 (heteroFc); #590, #595, 598, #605 (X-body)).

[0550] 48 h incubation time.

[0551] Determination of CD69 expression on CD4+ and CD8+ T cells with flow cytometer and antigen-specific conjugates mAb. Results see FIG. 15.

[0552] Target-independent T cell activation induced by BiTE® antibody constructs containing a single chain-Fc, hetero-Fc or crossbody fusion at the C-terminus was evaluated for the following constructs:

[0553] BiTE® constructs (serial dilutions: 0.1 pM-2 μ M)

[0554] a. CDH19 scFc; 245.3 μ g/mL (

[0555] b. CDH-19 Hetero Fc; 1 mg/mL

[0556] c. CDH19 Xbody; 6.3 mg/mL

[0557] Human PBMC effector cells (3 to 4 donors; #386, #392, #401 (scFc) #282, #284, #287 (heteroFc)).

[0558] 48 h incubation time.

[0559] Determination of CD69 expression on CD4+ and CD8+ T cells with flow cytometer and antigen-specific conjugates mAb. Results see FIG. 16.

Example 13

[0560] Purified BiTE® antibody constructs were coated on a Maxisorb Plate in decreasing concentration (100 nM, 1:4 dilutions). After 3 $\times$ washing with PBS-T and blocking with PBS/3% (w/v) BSA (60 min, 37° C.), pooled human plasma was incubated for 60 min, 80 rpm at room temperature. After 3 $\times$ washing a mouse monoclonal antibody specific for human C1q subunit A (CC1q) was added (Thermo MA1-83963, 1:500) for 60 min, 80 rpm, room temperature, after described washing steps a goat anti mouse Fc-POX mAb (1:5,000) was incubated for 60 min, 80 rpm, room temperature. After additional washing, TMB substrate was incubated and stopped after colorimetric reaction by addition of H₂SO₄. The absorption was determined at 450 nm.

[0561] Result: As shown in FIG. 17 at high concentrations, the BiTE® hetero Fc construct (squares) showed higher binding signals for human CC1q compared to a BiTE® single chain Fc construct (triangle). As a negative control a canonical BiTE® (circle) was used, which showed no significant CC1q binding.

Example 14: Pharmacokinetics of BiTE® Antibody Constructs Fused to Single-Chain Fc(scFc) and Hetero-Fc (hetFc) Proteins

[0562] Various target binding BiTE® antibodies were fused to two different half-life extending moieties. The two different HLE-variants per BiTE® antibody, subsequently named as BiTE®-scFc and BiTE®-hetFc, were tested in the cynomolgus monkey in the context of pharmacokinetic (PK) studies. The corresponding nomenclature of these molecules is briefly summarized in Table 16 below.

TABLE 16

Compound nomenclature of nine single dosed BiTE® HLE antibody constructs	
compound synonym	test compound name
Compound 1a	CD33cc-scFc
Compound 1b	CD33cc-hetFc
Compound 2a	MSLN-scFc
Compound 2b	MSLN-hetFc
Compound 3a	CDH19-scFc
Compound 3b	CH19-hetFc
Compound 4	CD20-scFc
Compound 5a	DLL3-scFc
Compound 5b	DLL3-hetFc

[0563] The BiTE® HLE antibody construct were administered as intravenous bolus (compounds 1b-5b) and intravenous infusion (compound 1a, 30 min) at 6 µg/kg (compound 2b), 12 µg/kg (compounds 2a and 3a-5b) and 15 µg/kg (compounds 1a and 1b), respectively. For each of the above named compounds a group of at least two to three animals was used. Blood samples were collected and serum was prepared for determination of serum concentrations. Serum BiTE® antibody construct levels were measured using an immunoassay. The assay is performed by capturing the BiTE® via its target moiety, while an antibody directed against the CD3-binding part of the construct was used for detection. The serum concentration-time profiles were used to determine PK parameters. Blood sampling time points are listed in Table 17 below.

TABLE 17

Blood sampling time points during the PK study		
blood sampling time points cmpd. 1a [h]	blood sampling time points cmpd. 1b [h]	blood sampling time points cmpd. 2a-5b [h]
0.085	0.085	0.05
4.00	1	0.25
24.00	4	0.5
48.00	12	1
72.00	24	4
96.00	48	8
120.00	72	24
144.00	96	48
168	144	72
	168	168
		240
		336

[0564] The pharmacokinetics of four BiTE®-HLE antibody construct pairs are shown exemplarily in FIG. 18. Each pair stands for the same BiTE® protein fused to either a scFc- or a hetFc extension. For all proteins serum levels were quantifiable for all time points in all animals after BiTE®-HLE administration. The PK profiles describe a biphasic, exponential decline after each of the single test item administrations.

[0565] The pharmacokinetic parameters were determined using standard non-compartmental analysis (NCA) methods. Using non compartmental analysis, the following PK parameters were estimated: AUC_{inf} (Area under the serum concentration—time curve), V_{ss} (volume of distribution at steady state), CL (systemic clearance) and terminal t_{1/2} (terminal half-life).

[0566] The PK parameter for each tested compound are summarized as mean of n=2 and n=3, respectively in Table 18 below.

TABLE 18

Pharmacokinetic parameter of scFc- and heteroFc-variants from different BiTE®-target binders in cynomolgus monkeys.				
test item	terminal t _{1/2} [h]	AUC _{inf} [h*ng/mL]	CI [mL/h/kg]	V _{ss} [mL/kg]
Compound 1a	167	6645	1.4	256
Compound 1b	95	4955	2.6	261
Compound 2a	213	12072	0.73	179
Compound 2b	116	6971	0.8	78
Compound 3a	61	3293	3.6	129
Compound 3b	59	3633	3.3	79

TABLE 18-continued

Pharmacokinetic parameter of scFc- and heteroFc-variants from different BiTE®-target binders in cynomolgus monkeys.				
test item	terminal t _{1/2} [h]	AUC _{inf} [h*ng/mL]	CI [mL/h/kg]	V _{ss} [mL/kg]
Compound 4	97	6266	1.9	180
Compound 5a	234	24769	0.48	144
Compound 5b	173	14639	0.82	166

[0567] Overall, the AUC_{inf} for the different BiTE®-HLE pairs of target binders fused to either -scFc and -hetFc moiety, respectively, ranged between 3293 h*ng/mL and 24769 h*ng/mL, depending on the BiTE® target context. All analyzed HLE fusions achieved systemic clearance values of 0.48 to 3.6 mL/h/kg. The corresponding volumes of distribution ranged between 78 and 261 mL/kg.

Example 15

[0568] Preformulated drug substances containing purified MSLN-hALB, MSLN-hFc, and MSLN-scFc respectively were buffer exchanged via ultrafiltration/diafiltration using membranes with a molecular weight cut-off (MWCO) of 10 kDa. Final formulation was achieved by adding concentrated stock solutions. Resulting formulations for each construct are listed in Table 19 and comprise K60RTrT composed of 20 mM potassium phosphate, 150 mM L-arginine hydrochloride, 6% (w/V) trehalose dihydrate, 0.01% (w/V) polysorbate 80 at pH 6.0 and G40MSuT composed of 10 mM glutamate, 4% (w/V) mannitol, 2% (w/V) sucrose, 0.01% (w/V) polysorbate 80 at pH 4.0. The target protein concentration was 1.0 mg/mL. Formulated MSLN constructs were filled to 1 mL in type I glass vials which were stoppered with butyl rubber stoppers and crimped with aluminum seals. Filled vials were incubated at -20, 5, 25 and 3700. One vial of each version was subjected to five freeze and thaw (FIT) cycles. Target freezing temperature was -29° C. Target thawing temperature was 2° C. The ramp rate was approximately 0.3 K/min.

[0569] Visual particles were assessed in accordance to the method described by Ph Eur 2.9.20 by trained operators. Visual particle counts per vial are depicted in Table 7. The number of visual (larger than 125 µm) proteinaceous particles was higher for MSLN-hFc if compared to both MSLN-hALB and MSLN-scFc.

TABLE 19

Number of visual proteinaceous particles per vial for stressed and unstressed (T0) samples containing different half-life extended anti-Mesothelin (MSLN) BiTE® constructs. K60RTrT denotes for a formulation containing 20 mM potassium phosphate, 150 mM L-arginine hydrochloride, 6% (w/V) trehalose dihydrate, 0.01% (w/V) polysorbate 80, pH 6.0. G40MSuT denotes for a formulation containing 10 mM glutamate, 4% (w/V) mannitol, 2% (w/V) sucrose, 0.01% (w/V) polysorbate 80, pH 4.0					
Construct	hALB	hFc		scFc	
Formulation	K60RTrT	K60RTrT	G40MSuT	K60RTrT	G40MSuT
Number of visible (>125 µm) proteinaceous particles per vial					
T0	0	0	1	0	0
5 F/T cycles	0	2	2	0	1
2 w 5° C.	0	2	2	0	0
2 w 25° C.	0	2	1	0	0
2 w 37° C.	0	2	2	0	0
4 w -20° C.	0	2	1	0	0

TABLE 19-continued

Number of visual proteinaceous particles per vial for stressed and unstressed (T0) samples containing different half-life extended anti-Mesothelin (MSLN) BiTE® constructs. K60RTTrT denotes for a formulation containing 20 mM potassium phosphate, 150 mM L-arginine hydrochloride, 6% (w/v) trehalose dihydrate, 0.01% (w/v) polysorbate 80, pH 6.0. G40MSuT denotes for a formulation containing 10 mM glutamate, 4% (w/v) mannitol, 2% (w/v) sucrose, 0.01% (w/v) polysorbate 80, pH 4.0					
Construct	hALB	hFc		scFc	
Formulation	K60RTTrT	K60RTTrT	G40MSuT	K60RTTrT	G40MSuT
Number of visible (>125 µm) proteinaceous particles per vial					
4 w 5° C.	0	1	2	0	0
4 w 25° C.	0	2	2	0	0
4 w 37° C.	0	2	2	0	0

[0570] The samples described above were also analyzed by size exclusion ultra-high performance chromatography (SE-UPLC) in order to quantify the percentaged content of high molecular weight species (HMWS). SE-UPLC was performed on an AcquityH-Class UPLC system (Waters) using an Acquity UPLC BEH200 SEC 150 mm column (Waters). Column temperature was set to 25° C. Separation of size variants was achieved by applying an isocratic method with a flow rate of 0.4 mL/min. The mobile phase was composed of 100 mM sodium phosphate, 250 mM NaCl at pH 6.8. The run time totals 6.0 minutes. Samples were held at 8° C. within the autosampler until analysis. A total amount of 3 pg protein was injected. In order to avoid carry over an intermediate injection with 40% acetonitrile was performed after each sample. Detection was based on fluorescence emission (excitation at 280 nm, emission at 325 nm). Peak integration was performed using Empower® software. Relative area under the curve of HMWS was reported (Table 20).

[0571] Fc based constructs exhibited lower HMWS contents in the formulation variant G40MSuT than in K60RTTrT independent on the stress condition. It became evident that MSLN-scFc contained less HMWS than MSLN-hFc in both G40MSuT as well as K60RTTrT preparations. MSLN-scFc in its preferred formulation (G40MSuT) was less prone to HMWS formation than MSLN-hALB formulated in K60RTTrT. In previous experiments this buffer showed improved stabilizing potential for hALB based constructs if compared to formulations with more acidic pH values.

TABLE 20

Overview on HMWS contents in stressed and unstressed (T0) MSLN-hALB, -hFc, and -scFc preparations determined via SE-UPLC					
Construct	hALB	hFc		scFc	
Formulation	K60RTTrT	K60RTTrT	G40MSuT	K60RTTrT	G40MSuT
% HMWS					
T0	1.8	6.7	3.3	2.5	1.3
5 F/T cycles	2.0	7.2	4.1	3.0	1.5
2 w 5° C.	n.t.	n.t.	n.t.	2.9	1.1
2 w 25° C.	n.t.	6.6	2.7	2.4	0.5
2 w 37° C.	2.6	6.3	2.1	2.7	0.3

TABLE 20-continued

Overview on HMWS contents in stressed and unstressed (T0) MSLN-hALB, -hFc, and -scFc preparations determined via SE-UPLC					
Construct	hALB	hFc		scFc	
Formulation	K60RTTrT	K60RTTrT	G40MSuT	K60RTTrT	G40MSuT
% HMWS					
4 w -20° C.	5.9	8.7	1.6	6.6	0.3
4 w 5° C.	2.0	8.2	2.8	3.6	0.6
4 w 25° C.	2.2	6.8	2.6	2.7	0.4
4 w 37° C.	3.5	7.6	1.9	4.3	0.3

n.t. = not tested

[0572] The abundance of chemical modifications upon heat stress (incubation at 37° C.) was monitored by peptide mapping. Protein samples were enzymatically digested and the resulting peptides were separated using reversed phase chromatography. The column eluate was directly injected into the ion source of a mass spectrometer for identification and quantitation of the peptides.

[0573] In order to achieve maximum coverage, two separate enzyme digests were performed: once with trypsin and once with chymotrypsin. In each case, the proteins were denatured with guanidinium chloride and then reduced with dithiothreitol (DTT). After incubation in DTT, free cysteine residues were alkylated by the addition of iodoacetic acid. Samples were then buffer exchanged into 50 mM Tris pH 7.8 for digestion. Trypsin and chymotrypsin were added to separate reaction tubes at a ratio of 1:10 (sample:enzyme) each. Samples were digested for 30 min at 37° C. and the reaction was quenched by adding trifluoroacetic acid.

[0574] A load of 5 µg of each digest was separately injected onto a Zorbax SB-C18 (Agilent #859700-902) reversed phase column equilibrated in 0.1% (v/v) formic acid (FA). A 156 minutes gradient of up to 90% acetonitrile containing 0.1% FA was used to elute the peptides directly into the electrospray ion source of a Q-Exactive Plus mass spectrometer (Thermo Scientific). Data was collected in data dependent mode using a top 12 method in which a full scan (resolution 70 000; scan range 200-2000 m/z) was followed by high energy collision dissociation (HCD) of the 12 most abundant ions (resolution 17 500).

[0575] Peptides were identified based on accurate mass and tandem mass spectrum using in-house software. Identifications were manually verified. Relative quantities of modified and unmodified peptides were calculated based on ion abundance using Pinpoint software (Thermo Scientific).

[0576] Percentages of chemical modifications of the complement determining regions (CDRs) and of the half-life extending portion (either hALB or Fc) detected in MSLN-hALB, -hFc, and -scFc preparations are given by Table 21. When comparing similar formulation conditions, it became obvious that overall, chemical modifications were least abundant in scFc constructs.

TABLE 21

Overview on chemical modifications in stressed and unstressed (T0) MSLN-hALB, -hFc, and -scFc preparations determined via peptide mapping					
Construct	hALB	hFc		scFc	
Formulation	K60RTrT	K60RTrT	G40MSuT	K60RTrT	G40MSuT
% N101 deamidation (CDR)					
T0	0.1	0.2	0.2	0.2	0.2
2 w 37° C.	0.7	0.8	3.0	0.7	3.2
4 w 37° C.	1.3	n.t.	8.5	n.t.	6.4
% N162 deamidation (CDR)					
T0	3.0	1.7	1.9	2.3	2.5
2 w 37° C.	15.9	11.6	2.7	15.0	3.3
4 w 37° C.	26.8	n.t.	3.7	n.t.	4.1
% M279 oxidation (CDR)					
T0	0.6	1.4	1.6	0.6	1.0
2 w 37° C.	1.2	0.8	0.8	0.6	1.0
4 w 37° C.	0.9	n.t.	0.8	n.t.	0.6
% N348 deamidation (CDR)					
T0	0.5	3.2	3.3	0.5	0.9
2 w 37° C.	20.5	21.6	1.9	9.4	1.3
4 w 37° C.	22.8	n.t.	2.0	n.t.	2.9
% N351 deamidation (CDR)					
T0	0.2	2.9	2.6	0.5	1.0
2 w 37° C.	6.6	12.7	0.9	3.8	0.4
4 w 37° C.	8.7	n.t.	0.8	n.t.	0.8
% M530 oxidation (Fc)					
T0	n.a.	3.9	4.1	2.6	3.2
2 w 37° C.	n.a.	9.0	3.1	4.0	4.3
4 w 37° C.	n.a.	n.t.	3.4	n.t.	3.5
% N603 deamidation (Fc)					
T0	n.a.	1.3	1.9	1.3	1.4
2 w 37° C.	n.a.	7.9	4.6	7.0	5.6
4 w 37° C.	n.a.	n.t.	6.9	n.t.	8.1
% M706 oxidation (Fc)					
T0	n.a.	3.2	3.6	1.5	2.1
2 w 37° C.	n.a.	6.0	2.8	2.1	2.5
4 w 37° C.	n.a.	n.t.	2.6	n.t.	2.0
% M587 oxidation (hALB)					
T0	1.0	n.a.	n.a.	n.a.	n.a.
2 w 37° C.	2.2	n.a.	n.a.	n.a.	n.a.
4 w 37° C.	2.3	n.a.	n.a.	n.a.	n.a.
% M623 oxidation (hALB)					
T0	1.9	n.a.	n.a.	n.a.	n.a.
2 w 37° C.	2.4	n.a.	n.a.	n.a.	n.a.
4 w 37° C.	3.0	n.a.	n.a.	n.a.	n.a.
% M798 oxidation (hALB)					
T0	1.4	n.a.	n.a.	n.a.	n.a.
2 w 37° C.	3.3	n.a.	n.a.	n.a.	n.a.
4 w 37° C.	3.5	n.a.	n.a.	n.a.	n.a.
% M829 oxidation (hALB)					
T0	8.9	n.a.	n.a.	n.a.	n.a.
2 w 37° C.	42.9	n.a.	n.a.	n.a.	n.a.
4 w 37° C.	44.1	n.a.	n.a.	n.a.	n.a.

n.a. = not applicable;
n.t. = not tested

Example 16

[0577] MSLN-hALB, -hFc, -scFc formulated as described under Example 15 were subjected to a pH jump experiment.

The concentration of the starting materials was 1.0 mg/mL. A volume of 0.38 mL of each starting material was filled in a glass vial. After preconditioning at 37° C. the solutions were spiked with 20 fold phosphate buffered saline (PBS) which was composed of 0.090 M potassium phosphate, 0.480 M sodium phosphate (both dibasic), 0.052 M potassium chloride and 2.76 M NaCl. The spiked samples were incubated at 37° C. for two weeks. After incubation they were analyzed by SE-UPLC using the method described under Example 15 and the percentaged content of HMWS was reported (Table 22). When comparing all constructs formulated in K60RTrT the HMWS content increased in the following order: hALB<scFc<hFc. MSLN-scFc also showed a lower HMWS content than MSLN-hFc when formulated in G40MSuT.

TABLE 22

Overview on HMWS contents in stressed (pH jump + 2 w 37° C.) MSLN-hALB, -hFc, and -scFc preparations determined via SE-UPLC					
Construct	hALB	hFc		scFc	
Formulation	K60RTrT	K60RTrT	G40MSuT	K60RTrT	G40MSuT
% HMWS					
2 w 37° C.	1.5	8.3	7.1	5.4	5.1

Example 17

[0578] MSLN-hALB, -hFc, and -scFc formulated as described under Example 15 were subjected to agitation stress. The concentration of the starting materials was 1.0 mg/mL. A volume of 0.5 mL of each solution was filtered through an appropriate 0.22 µm filter and filled into 3 cc glass vials. The vials were placed in a plastic box ensuring that the vials were not displaced within the box during agitation. The box was placed onto an orbital shaker. The samples were agitated at 500 rpm for 65 hours. Visual particles were assessed in accordance to the method described by Ph Eur 2.9.20. The method was conducted by trained operators. Visual particle counts per vial are depicted in Table 23. Visible proteinaceous particles were only observed in MSLN-hFc preparations.

TABLE 23

Number of visual proteinaceous particles per vial in agitated samples					
Construct	hALB	hFc		scFc	
Formulation	K60RTrT	K60RTrT	G40MSuT	K60RTrT	G40MSuT
Number of visible (>125 µm) proteinaceous particles per vial					
65 h, 500 rpm	0	1	1	0	0

[0579] Above samples were also analyzed by size exclusion ultra-high performance chromatography (SE-UPLC) in order to quantify the percentaged content of high molecular weight species (HMWS). The same method as described in Example 15 was applied. The HMWS contents of agitated samples are outlined by Table 24. The formation of HMWS was most pronounced in MSLN-hFc when comparing K60RTrT preparations. For Fc based constructs the HMWS contents could be reduced by lowering the formulation pH

(G40MSuT). But again HMWS were more abundant in MSLN-hFc than in MSLN-scFc.

TABLE 24

Overview on HMWS contents in stressed (pH jump + 2 w 37° C.) MSLN-hALB, -hFc, and -scFc preparations determined via SE-UPLC					
Construct	hALB	hFc		scFc	
Formulation	K60RTTrT	K60RTTrT	G40MSuT	K60RTTrT	G40MSuT
% HMWS					
65 h, 500 rpm	1.8	5.8	2.4	1.8	0.3

Example 17

[0580] MSLN-hALB, -hFc, and -scFc formulated as described under Example 15 were exposed to visible and UVA light (photo stress). Protein concentration totaled 1 mg/mL in all preparations. Protein solutions were filtered through a filter with 0.22 µm pore size and filled to 0.5 mL in type I glass vials. MSLN-hALB and -scFc were subjected to two different tests including 0.2 MLux visible light/25 W*/h/m² UVA light and 1.2MLux visible light/173 W*/h/m² respectively. MSLN-hFc was subjected to two different tests including 0.2 MLux visible light without UVA light and 1.2 MLux visible light/30 W*/h/m² UVA light respectively. Chamber temperatures were adjusted to 25° C. After light exposure samples were analyzed by visible inspection (Table 25), SE-UPLC (Table 26) and peptide map (Table 27). Aforementioned methods were performed according to the procedures described under Example 15. Although MSLN-hALB, and -scFc were exposed to higher doses of UVA light, no visible proteinaceous particles was observed whereas MSLN-hFc samples exhibited one visible proteinaceous particle per vial for both tests irrespective of the formulation.

TABLE 25

Overview on the number of visible proteinaceous particles per vial in MSLN-hALB, -hFc, and -scFc preparations determined after light exposure					
Construct	hALB	hFc		scFc	
Formulation	K60RTTrT	K60RTTrT	G40MSuT	K60RTTrT	G40MSuT
Number of visible (>125 µm) proteinaceous particles per vial					
T0	0	0	1	0	0
Test 1	0 ¹⁾	1 ²⁾	1 ²⁾	0 ¹⁾	0 ¹⁾
Test 2	0 ³⁾	1 ⁴⁾	1 ⁴⁾	0 ³⁾	0 ³⁾

¹⁾0.2 MLux visible light/25 W*/h/m² UVA light,

²⁾0.2 MLux visible light without UVA light,

³⁾1.2 MLux visible light/173 W*/h/m²,

⁴⁾1.2 MLux visible light/30 W*/h/m²

[0581] HMWS increased in the following order MSLN-hALB <-scFc <-hFc when the protein was formulated in K60RTTrT. HMWS could be reduced for Fc based constructs when formulated in G40MSuT. However HMWS were again less pronounced for MSLN-scFc. MSLN-hFc revealed to be especially sensitive towards UVA light exposure.

TABLE 26

Overview on HMWS contents in MSLN-hALB, -hFc, and -scFc preparations determined after light exposure via SE-UPLC					
Construct	hALB	hFc		scFc	
Formulation	K60RTTrT	K60RTTrT	G40MSuT	K60RTTrT	G40MSuT
% HMWS					
T0	1.8	6.7	3.3	2.5	1.3
Test 1	1.8 ¹⁾	6.3 ²⁾	2.5 ²⁾	2.1 ¹⁾	0.4 ¹⁾
Test 2	2.0 ³⁾	11.0 ⁴⁾	2.1 ⁴⁾	2.4 ³⁾	0.3 ³⁾

¹⁾0.2 MLux visible light/25 W*/h/m² UVA light,

²⁾0.2 MLux visible light without UVA light,

³⁾1.2 MLux visible light/173 W*/h/m²,

⁴⁾1.2 MLux visible light/30 W*/h/m²

[0582] Percentages of chemical modifications of the complement determining regions (CDRs) and of the half-life extending portion (either hALB or Fc) detected in MSLN-hALB, -hFc, and -scFc preparations are given by Table 27. When comparing similar formulation conditions, it became obvious that overall, chemical modifications were least abundant in scFc constructs.

TABLE 27

Overview on chemical modifications in MSLN-hALB, -hFc, and -scFc preparations determined after light exposure via peptide mapping					
Construct	hALB	hFc		scFc	
Formulation	K60RTTrT	K60RTTrT	G40MSuT	K60RTTrT	G40MSuT
% N101 deamidation (CDR)					
T0	0.1	0.2	0.2	0.2	0.2
Test 1	0.2 ¹⁾	n.t.	0.3 ²⁾	n.t.	0.5 ¹⁾
Test 2	0.2 ³⁾	n.t.	0.6 ⁴⁾	n.t.	0.7 ³⁾
% N162 deamidation (CDR)					
T0	3.0	1.7	1.9	2.3	2.5
Test 1	3.0 ¹⁾	n.t.	2.1 ²⁾	n.t.	2.7 ¹⁾
Test 2	3.6 ³⁾	n.t.	3.1 ⁴⁾	n.t.	2.8 ³⁾
% M279 oxidation (CDR)					
T0	0.6	1.4	1.6	0.6	1.0
Test 1	0.8 ¹⁾	n.t.	2.6 ²⁾	n.t.	0.6 ¹⁾
Test 2	1.0 ³⁾	n.t.	6.3 ⁴⁾	n.t.	0.7 ³⁾
% N348 deamidation (CDR)					
T0	0.5	3.2	3.3	0.5	0.9
Test 1	0.4 ¹⁾	n.t.	2.7 ²⁾	n.t.	0.2 ¹⁾
Test 2	0.9 ³⁾	n.t.	3.9 ⁴⁾	n.t.	0.2 ³⁾
% N351 deamidation (CDR)					
T0	0.2	2.9	2.6	0.5	1.0
Test 1	0.4 ¹⁾	n.t.	2.0 ²⁾	n.t.	0.3 ¹⁾
Test 2	0.5 ³⁾	n.t.	2.6 ⁴⁾	n.t.	0.3 ³⁾
% M530 oxidation (Fc)					
T0	n.a.	3.9	4.1	2.6	3.2
Test 1	n.a.	n.t.	7.6 ²⁾	n.t.	3.1 ¹⁾
Test 2	n.a.	n.t.	21.8 ⁴⁾	n.t.	4.1 ³⁾
% M706 oxidation (Fc)					
T0	n.a.	3.2	3.6	1.5	2.1
Test 1	n.a.	n.t.	6.5 ²⁾	n.t.	1.8 ¹⁾
Test 2	n.a.	n.t.	17.8 ⁴⁾	n.t.	2.7 ³⁾
% M587 oxidation (hALB)					
T0	1.0	n.a.	n.a.	n.a.	n.a.
Test 1	1.5	n.a.	n.a.	n.a.	n.a.
Test 2	2.4	n.a.	n.a.	n.a.	n.a.

TABLE 27-continued

Overview on chemical modifications in MSLN-hALB, -hFc, and -scFc preparations determined after light exposure via peptide mapping					
Construct	hALB	hFc		scFc	
Formulation	K60RTrT	K60RTrT	G40MSuT	K60RTrT	G40MSuT
% M623 oxidation (hALB)					
T0	1.9	n.a.	n.a.	n.a.	n.a.
Test 1	4.0	n.a.	n.a.	n.a.	n.a.
Test 2	4.1	n.a.	n.a.	n.a.	n.a.
% M798 oxidation (hALB)					
T0	1.4	n.a.	n.a.	n.a.	n.a.
Test 1	2.1	n.a.	n.a.	n.a.	n.a.
Test 2	3.1	n.a.	n.a.	n.a.	n.a.
% M829 oxidation (hALB)					
T0	8.9	n.a.	n.a.	n.a.	n.a.
Test 1	31.0	n.a.	n.a.	n.a.	n.a.
Test 2	25.2	n.a.	n.a.	n.a.	n.a.

n.a. = not applicable;
n.t. = not tested

Example 18

[0583] MSLN-hALB was formulated in K60RTrT and MSLN-scFc was formulated in G40MSuT according to the procedure described in Example 15. Protein concentrations totaled 0.05 mg/mL. Glass (borosilicate, type I, 13 mm 3 cc vial from West, Art. No. 68000375) and polypropylene test containers (2 mL with O-ring, e.g. from Sarstedt, Art No. 72.694.005) are filled with 500 μ L of the test solution. The test solution was left for five minutes in the first test container. Then a 150 μ L aliquot was sampled for analysis. The remaining test solution (350 μ L) was transferred sequentially from one test container to the next (five containers in total). In each vial, the solution was left for five minutes before the next transfer. The same pipette tip was used for each transfer step. The same test was performed using 30 mL polycarbonate bottles (Nalgene, PCS-000295 with closure, PP/20-415/ZTPE). For this container type the first container was filled with 5 mL. After a 150 μ L aliquot was sampled, the residual volume was transferred from one test container to the next (according to the procedure described above). Samples pulled from container #1 and #5 were analyzed by SE-UPLC (method as described under Example 15). In addition protein detection was carried out with a PDA detector (280 nm) in order to determine protein concentrations. Percentaged protein recovery from each test

container is given by Table 28. It was shown that protein recovery was more pronounced for MSLN-scFc than for MSLN-hALB irrespective of the container type.

TABLE 28

Protein recovery from different container types for MSLN-hALB, and -scFc determined by SE-UPLC		
Construct	hALB	scFc
Formulation	K60RTrT	G40MSuT
% Protein recovery (from nominal)		
Type I glass	80.0	92.0
Polypropylene	87.0	97.3
Polycarbonate	87.0	96.0

Example 19

[0584] MSLN-hALB was formulated in K60RTrT and MSLN-scFc was formulated in K60RTrT and G40MSuT according to the procedure described in Example 15. The protein concentration totaled 1.0 mg/mL. 1950 μ L of each test solution was spiked with 50 μ L of a 1000 ppm silicon standard solution (Specpure from AlfaAesar, Art. No. 38717) resulting in a 25 ppm spike. An unspiked test solution served as control sample. The spiked test solution as well as the control sample were filled into 3 cc type I glass vials and were incubated at 37° C. for 24 hours. All samples were analyzed by SE-UPLC according to the method described in Example 15 in order to quantify the amount of HMWS (Table 29). When formulated in K60RTrT, MSLN-hALB and -scFc showed similar increases in HMWS upon silicon spiking. For the scFc construct it could be shown that this increase could be reduced by lowering the formulation pH to 4.0. According to preliminary experiments, this approach was not feasible for MSLN-hALB since it revealed to undergo fragmentation at formulation pH values of 5.0 and below.

TABLE 29

Overview on HMWS contents in MSLN-hALB, and -scFc preparations determined via SE-UPLC after spiking with 25 ppm silicon			
Construct	hALB	scFc	
Formulation	K60RTrT	K60RTrT	G40MSuT
Δ % HMWS (compared to unspiked control)			
25 ppm spike	1.0	1.0	0.2

TABLE 30

Sequence Listing			
SEQ ID NO:	Description	Source	Sequence
1.	Peptide linker	artificial	GGGG
2.	Peptide linker	artificial	GGGGS
3.	Peptide linker	artificial	GGGGQ

TABLE 30-continued

Sequence Listing			
SEQ ID NO:	Description	Source	Sequence
4.	Peptide linker	artificial	PGGGGS
5.	Peptide linker	artificial	PGGDGS
6.	Peptide linker	artificial	SGGGGS
7.	Peptide linker	artificial	GGGSGGGGS
8.	Peptide linker	artificial	GGGSGGGGS
9.	Peptide linker	artificial	GGGSGGGSGGGGS
10.	Hexa-histidine	artificial	HHHHHH
11.	CDR-L1 of F6A	artificial	GSSTGAVTSGYYPN
12.	CDR-L2 of F6A	artificial	GTKFLAP
13.	CDR-L3 of F6A	artificial	ALWYSNRWV
14.	CDR-H1 of F6A	artificial	IYAMN
15.	CDR-H2 of F6A	artificial	RIRSKYNNYATYYADSVKS
16.	CDR-H3 of F6A	artificial	HGNFGNSYVSFFAY
17.	VH of F6A	artificial	EVQLVESGGGLVQPGGSLKLSCAASGFTFN IYAMNWVRQAPGKLEWVARIRSKYNNYATYYADSVKSRFTISRDDSKN TAYLQMNNLKTEDTAVYYCVRHGNFGNSYVSFFAYWGQGLTVTVSS
18.	VL of F6A	artificial	QTVVTQEPSLTVSPGGTVTLTCGSSTGAVTSGYYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLGGAALTLSG VQPEDEAEYYCALWYSNRWVFGGGTKLTVL
19.	VH-VL of F6A	artificial	EVQLVESGGGLVQPGGSLKLSCAASGFTFN IYAMNWVRQAPGKLEWVARIRSKYNNYATYYADSVKSRFTISRDDSKN TAYLQMNNLKTEDTAVYYCVRHGNFGNSYVSFFAYWGQGLTVTVSSGGGSGGGSGGGSGTQVVTQEPSLTVSPGGTV TLTCGSSTGAVTSGYYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLGGAALTLSGVQPEDEAEYYCALWYSNR WVFGGGTKLTVL
20.	CDR-L1 of H2C	artificial	GSSTGAVTSGYYPN
21.	CDR-L2 of H2C	artificial	GTKFLAP
22.	CDR-L3 of H2C	artificial	ALWYSNRWV
23.	CDR-H1 of H2C	artificial	KYAMN
24.	CDR-H2 of H2C	artificial	RIRSKYNNYATYYADSVKD
25.	CDR-H3 of H2C	artificial	HGNFGNSYISYWAY
26.	VH of H2C	artificial	EVQLVESGGGLVQPGGSLKLSCAASGFTFN KYAMNWVRQAPGKLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKN TAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSS

TABLE 30-continued

Sequence Listing			
SEQ ID NO:	Description	Source	Sequence
27.	VL of H2C	artificial	QTVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGYYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYYCALWYSNRWVFGGGTKLTVL
28.	VH-VL of H2C	artificial	EVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYLSYWAYWGQGTTLVTVSSGGGGSGGGSGGGGSQTVVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGYYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYYCALWYSNRWVFGGGTKLTVL
29.	CDR-L1 of H1E	artificial	GSSTGAVTSGYYPN
30.	CDR-L2 of H1E	artificial	GTKFLAP
31.	CDR-L3 of H1E	artificial	ALWYSNRWV
32.	CDR-H1 of H1E	artificial	SYAMN
33.	CDR-H2 of H1E	artificial	RIRSKYNNYATYYADSVKG
34.	CDR-H3 of H1E	artificial	HGNFGNSYLSFWAY
35.	VH of H1E	artificial	EVQLVESGGGLEQPGGSLKLSCAASGFTFNSYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKGRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYLSFWAYWGQGTTLVTVSS
36.	VL of H1E	artificial	QTVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGYYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYYCALWYSNRWVFGGGTKLTVL
37.	VH-VL of H1E	artificial	EVQLVESGGGLEQPGGSLKLSCAASGFTFNSYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKGRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYLSFWAYWGQGTTLVTVSSGGGGSGGGSGGGGSQTVVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGYYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYYCALWYSNRWVFGGGTKLTVL
38.	CDR-L1 of G4H	artificial	GSSTGAVTSGYYPN
39.	CDR-L2 of G4H	artificial	GTKFLAP
40.	CDR-L3 of G4H	artificial	ALWYSNRWV
41.	CDR-H1 of G4H	artificial	RYAMN
42.	CDR-H2 of G4H	artificial	RIRSKYNNYATYYADSVKG
43.	CDR-H3 of G4H	artificial	HGNFGNSYLSYPAY
44.	VH of G4H	artificial	EVQLVESGGGLVQPGGSLKLSCAASGFTFNRYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKGRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYLSYFAYWGQGTTLVTVSS
45.	VL of G4H	artificial	QTVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGYYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYYCALWYSNRWVFGGGTKLTVL
46.	VH-VL of G4H	artificial	EVQLVESGGGLVQPGGSLKLSCAASGFTFNRYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKGRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYLSYFAYWGQGTTLVTVSSGGGGSGGGSGGGGSQTVVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGYYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYYCALWYSNRWVFGGGTKLTVL
47.	CDR-L1 of A2J	artificial	RSSTGAVTSGYYPN
48.	CDR-L2 of A2J	artificial	ATDMRPS

TABLE 30-continued

Sequence Listing			
SEQ ID NO:	Descrip- tion	Source	Sequence
49.	CDR-L3 of A2J	artifi- cial	ALWYSNRWV
50.	CDR-H1 of A2J	artifi- cial	VYAMN
51.	CDR-H2 of A2J	artifi- cial	RIRSKYNNYATYYADSVKK
52.	CDR-H3 of A2J	artifi- cial	HGNFGNSYLSWWAY
53.	VH of A2J	artifi- cial	EVQLVESGGGLVQPGGSLKLSCAASGFTFNVIYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKKRFTISRDDSKN TAYLQMNNLKTEDTAVYYCVRHGNFGNSYLSWWAYWGQGTLLTVSS
54.	VL of A2J	artifi- cial	QTVVTQEPSLTVSPGGTVTLTCSRSTGAVTSGYYPNWVQKPGQAPRGLIGATDMRPSGTPARFSGSLGGKAALTLSG VQPEDEAEYYCALWYSNRWVFGGGTKLTVL
55.	VH-VL of A2J	artifi- cial	EVQLVESGGGLVQPGGSLKLSCAASGFTFNVIYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKKRFTISRDDSKN TAYLQMNNLKTEDTAVYYCVRHGNFGNSYLSWWAYWGQGTLLTVSSGGGGSGGGSGGGSGTQVVTQEPSLTVSPGGTV TLTCSRSTGAVTSGYYPNWVQKPGQAPRGLIGATDMRPSGTPARFSGSLGGKAALTLSGVQPEDEAEYYCALWYSNR WVFGGGTKLTVL
56.	CDR-L1 of E1L	artifi- cial	GSSTGAVTSGYYPN
57.	CDR-L2 of E1L	artifi- cial	GTKFLAP
58.	CDR-L3 of E1L	artifi- cial	ALWYSNRWV
59.	CDR-H1 of E1L	artifi- cial	KYAMN
60.	CDR-H2 of E1L	artifi- cial	RIRSKYNNYATYYADSVKS
61.	CDR-H3 of E1L	artifi- cial	HGNFGNSYTSYYAY
62.	VH of E1L	artifi- cial	EVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKSRFTISRDDSKN TAYLQMNNLKTEDTAVYYCVRHGNFGNSYTSYYAYWGQGTLLTVSS
63.	VL of E1L	artifi- cial	QTVVTQEPSLTVSPGGTVTLTCGSSTGAVTSGYYPNWVQKPGQAPRGLIGGTFKFLAPGTPARFSGSLGGKAALTLSG VQPEDEAEYYCALWYSNRWVFGGGTKLTVL
64.	VH-VL of E1L	artifi- cial	EVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKSRFTISRDDSKN TAYLQMNNLKTEDTAVYYCVRHGNFGNSYTSYYAYWGQGTLLTVSSGGGGSGGGSGGGSGTQVVTQEPSLTVSPGGTV TLTCGSSTGAVTSGYYPNWVQKPGQAPRGLIGGTFKFLAPGTPARFSGSLGGKAALTLSGVQPEDEAEYYCALWYSNR WVFGGGTKLTVL
65.	CDR-L1 of E2M	artifi- cial	RSSTGAVTSGYYPN
66.	CDR-L2 of E2M	artifi- cial	ATDMRPS
67.	CDR-L3 of E2M	artifi- cial	ALWYSNRWV
68.	CDR-H1 of E2M	artifi- cial	GYAMN
69.	CDR-H2 of E2M	artifi- cial	RIRSKYNNYATYYADSVKE
70.	CDR-H3 of E2M	artifi- cial	HRNFGNSYLSWFAY

TABLE 30-continued

Sequence Listing			
SEQ ID NO:	Description	Source	Sequence
71.	VH of E2M	artificial	EVQLVESGGGLVQPGGSLKLSAASGFTFNGYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKERFTISRDDSKN TAYLQMNNLKTEDTAVYYCVRHRNFGNSYLSWFAYWGQGTLLTVSS
72.	VL of E2M	artificial	QTVVTQEPSTLVSPGGTVTLTCSRSTGAVTSGYYPNWVQKPGQAPRGLIGATDMRPSGTPARFSGSLGGKAALTLSG VQPEDEAEYYCALWYSNRWVFGGGTKLTVL
73.	VH-VL of E2M	artificial	EVQLVESGGGLVQPGGSLKLSAASGFTFNGYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKERFTISRDDSKN TAYLQMNNLKTEDTAVYYCVRHRNFGNSYLSWFAYWGQGTLLTVSSGGGGSGGGSGGGGSQTVVTQEPSTLVSPGGTV TLTCSRSTGAVTSGYYPNWVQKPGQAPRGLIGATDMRPSGTPARFSGSLGGKAALTLSGVQPEDEAEYYCALWYSNR WVFGGGTKLTVL
74.	CDR-L1 of F70	artificial	GSSTGAVTSGYYPN
75.	CDR-L2 of F70	artificial	GTKFLAP
76.	CDR-L3 of F70	artificial	ALWYSNRWV
77.	CDR-H1 of F70	artificial	VYAMN
78.	CDR-H2 of F70	artificial	RIRSKYNNYATYYADSVKK
79.	CDR-H3 of F70	artificial	HGNFGNSYISWWAY
80.	VH of F70	artificial	EVQLVESGGGLVQPGGSLKLSAASGFTFNVYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKKRFTISRDDSKN TAYLQMNNLKTEDTAVYYCVRHGNFGNSYISWWAYWGQGTLLTVSS
81.	VL of F70	artificial	QTVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGYYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLGGKAALTLSG VQPEDEAEYYCALWYSNRWVFGGGTKLTVL
82.	VH-VL of F70	artificial	EVQLVESGGGLVQPGGSLKLSAASGFTFNVYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKKRFTISRDDSKN TAYLQMNNLKTEDTAVYYCVRHGNFGNSYISWWAYWGQGTLLTVSSGGGGSGGGSGGGGSQTVVTQEPSTLVSPGGTV TLTCGSSTGAVTSGYYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLGGKAALTLSGVQPEDEAEYYCALWYSNR WVFGGGTKLTVL
83.	CDR-L1 of F12Q	artificial	GSSTGAVTSGNYPN
84.	CDR-L2 of F12Q	artificial	GTKFLAP
85.	CDR-L3 of F12Q	artificial	VLWYSNRWV
86.	CDR-H1 of F12Q	artificial	SYAMN
87.	CDR-H2 of F12Q	artificial	RIRSKYNNYATYYADSVKG
88.	CDR-H3 of F12Q	artificial	HGNFGNSYVSWWAY
89.	VH of F12Q	artificial	EVQLVESGGGLVQPGGSLKLSAASGFTFNSYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKGRFTISRDDSKN TAYLQMNNLKTEDTAVYYCVRHGNFGNSYVSWWAYWGQGTLLTVSS
90.	VL of F12Q	artificial	QTVVTQEPSTLVSPGGTVTLTCGSSTGAVTSGNYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLGGKAALTLSG VQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
91.	VH-VL of F12Q	artificial	EVQLVESGGGLVQPGGSLKLSAASGFTFNSYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKGRFTISRDDSKN TAYLQMNNLKTEDTAVYYCVRHGNFGNSYVSWWAYWGQGTLLTVSSGGGGSGGGSGGGGSQTVVTQEPSTLVSPGGTV TLTCGSSTGAVTSGNYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLGGKAALTLSGVQPEDEAEYYCVLWYSNR WVFGGGTKLTVL
92.	CDR-L1 of I2C	artificial	GSSTGAVTSGNYPN

TABLE 30-continued

Sequence Listing			
SEQ ID NO:	Description	Source	Sequence
93.	CDR-L2 of I2C	artificial	GTKFLAP
94.	CDR-L3 of I2C	artificial	VLWYSNRWV
95.	CDR-H1 of I2C	artificial	KYAMN
96.	CDR-H2 of I2C	artificial	RIRSKYNNYATYYADSVKD
97.	CDR-H3 of I2C	artificial	HGNFGNSYISYWAY
98.	VH of I2C	artificial	EVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKN TAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSS
99.	VL of I2C	artificial	QTVVTQEPSLTVSPGGTVTLTCGSSGTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLGGKAALTLSG VQPEDEAEYYCVLWYSNRWVFGGKLTVL
100.	VH-VL of I2C	artificial	EVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKN TAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGSGGGSGGSGTQVVTQEPSLTVSPGGTV TLTCGSSGTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLGGKAALTLSGVQPEDEAEYYCVLWYSNR WVFGGKLTVL
101.	VH of F12q	artificial	EVQLVESGGGLVQPGGSLRLSCAASGFTFNSYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKGRFTISRDDSKN TAYLQMNSLKTEDTAVYYCVRHGNFGNSYVSWWAYWGQGLTVTVSS
102.	VL of F12q	artificial	QTVVTQEPSLTVSPGGTVTLTCGSSGTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLGGKAALTLSG VQPEDEAEYYCVLWYSNRWVFGGKLTVL
103.	F12q	scFv	EVQLVESGGGLVQPGGSLRLSCAASGFTFNSYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKGRFTISRDDSKN TAYLQMNSLKTEDTAVYYCVRHGNFGNSYVSWWAYWGQGLTVTVSSGGGGSGGGSGGGSGGSGTQVVTQEPSLTVSPGGTV TLTCGSSGTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLGGKAALTLSGVQPEDEAEYYCVLWYSNR WVFGGKLTVL
104.	HALB	human	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQCPFDHVKLVNEVTEFAKTCVADESAENCDSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFQHKDDNPPLRLVRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCSLQKFGERAFKAWAVARLSQRFPKAEFAEVSKLVT LTKVHTECCGDLLECADRADLAKYICENQDSISSKLEKCEKPLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIKQNCSELF EQLGEYKFQNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALEVDETYVPKEFNAETFTFHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAVMDDFAA FVEKCKADDKETCFABEGKKLVAASQAALGL
105.	HALB7	artificial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQCPFDHVKLVNEVTEFAKTCVADESAENCDSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFQHKDDNPPLRLVRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCSLQKFGERAFKAWAVARLSQRFPKAEFAEVSKLVT LTKVHTECCGDLLECADRADLAKYICENQDSISSKLEKCEKPLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIKQNCSELF EQLGEYKFQNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALEVDETYVPKEFNAETFTFHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAVMDDFAA FVEKCKADDKETCFABEGKKLVAASQAALGL
106.	HALB098	artificial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQCPFDHVKLVNEVTEFAKTCVADESAENCDSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFQHKDDNPPLRLVRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCSLQKFGERAFKAWAVARLSQRFPKAEFAEVSKLVT LTKVHTECCGDLLECADRADLAKYICENQDSISSKLEKCEKPLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIKQNCSELF EQLGEYKFQNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALEVDETYVPKEFNAETFTFHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAVMDDFAA FVEKCKADDKETCFABEGKKLVAASQAALGL

TABLE 30-continued

Sequence Listing			
SEQ ID NO:	Description	Source	Sequence
107.	HALB114	artificial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQCPFEDHVKLVNEVTEFAKTCVADESAENCCKSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLOHKDDNPPLRLVRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCASLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLEKCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIKQNCSELF EQLGEYKFQNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALDVEDYVPKEFNAETFTPHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAVMDDFAA FVEKCKKADDKETCFAEEGPHLVAASKAALGL
108.	HALB254	artificial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQCPFEDHVKLVNEVTEFAKTCVADESAENCCKSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLOHKDDNPPLRLVRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCASLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLEKCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIKQNCSELF EQLGEYKFQNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALDVEDYVPKEFNAETFTPHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAVMDKFAA FVEKCKKADDKETCFAEEGPKLVAASQAALGL
109.	HALB253	artificial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQCPFEDHVKLVNEVTEFAKTCVADESAENCCKSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLOHKDDNPPLRLVRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCASLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLEKCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIKQNCSELF EQLGEYKFQNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALDVEDYVPKEFNAETFTPHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAVMDKFAA FVEKCKKADDKETCFAEEGPKLVAASQAALGL
110.	HALB131	artificial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQCPFEDHVKLVNEVTEFAKTCVADESAENCCKSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLOHKDDNPPLRLVRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCASLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLEKCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIKQNCSELF EQLGEYKFQNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALDVEDYVPKEFNAETFTPHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAVMDKFAA FVEKCKKADDKETCFAEEGPHLVAASQAALGL
111.	HALB135	artificial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQCPFEDHVKLVNEVTEFAKTCVADESAENCCKSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLOHKDDNPPLRLVRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCASLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLEKCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIKQNCSELF EQLGEYKFQNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALDVEDYVPKEFNAETFTPHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAVMDDFAA FVEKCKKADDKETCFAEEGPHLVAASKAALGL
112.	HALB133	artificial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQCPFEDHVKLVNEVTEFAKTCVADESAENCCKSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLOHKDDNPPLRLVRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCASLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLEKCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIKQNCSELF EQLGEYKFQNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALDVEDYVPKEFNAETFTPHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAVMDKFAA FVEKCKKADDKETCFAEEGPKLVAASKAALGL
113.	HALB234	artificial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQCPFEDHVKLVNEVTEFAKTCVADESAENCCKSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLOHKDDNPPLRLVRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCASLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLEKCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIKQNCSELF EQLGEYKFQNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALDVEDYVPKEFNAETFTPHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAVMDDFAA FVEKCKKADDKETCFAEEGPKLVAASKAALGL

TABLE 30-continued

Sequence Listing			
SEQ ID NO:	Description	Source	Sequence
114.	HALB C34S	artificial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQSPFEDHVKLVNEVTEFAKTCVADESAENCCKSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLQHKDDNPNLPRLVPRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCASLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIQNCSELF EQLGEYKFNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALEVDETYVPKEFNAETFTPHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAVMDDFAA FVEKCKADDDKETCFAEEGKCLVAASQAALGL
115.	HALB7 C34S	artificial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQSPFEDHVKLVNEVTEFAKTCVADESAENCCKSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLQHKDDNPNLPRLVPRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCASLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIQNCSELF EQLGEYKFNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALEVDETYVPKEFNAETFTPHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAVMDDFAA FVEKCKADDDKETCFAEEGKCLVAASQAALGL
116.	HALB098 C34S	artificial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQSPFEDHVKLVNEVTEFAKTCVADESAENCCKSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLQHKDDNPNLPRLVPRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCASLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIQNCSELF EQLGEYKFNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALEVDETYVPKEFNAETFTPHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAVMDDFAA FVEKCKADDDKETCFAEEGPKLVAASQAALGL
117.	HALB114 C34S	artificial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQSPFEDHVKLVNEVTEFAKTCVADESAENCCKSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLQHKDDNPNLPRLVPRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCASLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIQNCSELF EQLGEYKFNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALDVEDETYVPKEFNAETFTPHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAVMDDFAA FVEKCKADDDKETCFAEEGPHLVAASQAALGL
118.	HALB254 C34S	artificial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQSPFEDHVKLVNEVTEFAKTCVADESAENCCKSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLQHKDDNPNLPRLVPRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCASLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIQNCSELF EQLGEYKFNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALGVEDETYVPKEFNAETFTPHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAVMDKFAA FVEKCKADDDKETCFAEEGPKLVAASQAALGL
119.	HALB253 C34S	artificial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQSPFEDHVKLVNEVTEFAKTCVADESAENCCKSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLQHKDDNPNLPRLVPRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCASLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIQNCSELF EQLGEYKFNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALDVEDETYVPKEFNAETFTPHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAVMDKFAA FVEKCKADDDKETCFAEEGPKLVAASQAALGL
120.	HALB131 C34S	artificial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQSPFEDHVKLVNEVTEFAKTCVADESAENCCKSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLQHKDDNPNLPRLVPRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCASLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIQNCSELF EQLGEYKFNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALEVDETYVPKEFNAETFTPHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAVMDKFAA FVEKCKADDDKETCFAEEGPHLVAASQAALGL

TABLE 30-continued

Sequence Listing			
SEQ ID NO:	Description	Source	Sequence
121.	HALB135 C34S	artificial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQSPFEDHVKLVNEVTEFAKTCVADESAENCCKSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLQHKDDNPMLPRLVRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCCASLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIKQNCSELF EQLGEYKFQNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALEVDETYVPKEFNAETFTPHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAVMDDFAA FVEKCKKADDKETCFAEEGPHLVAASKAALGL
122.	HALB133 C34S	artificial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQSPFEDHVKLVNEVTEFAKTCVADESAENCCKSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLQHKDDNPMLPRLVRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCCASLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIKQNCSELF EQLGEYKFQNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALEVDETYVPKEFNAETFTPHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAVMDKFAA FVEKCKKADDKETCFAEEGPKLVAASKAALGL
123.	HALB234 C34S	artificial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQSPFEDHVKLVNEVTEFAKTCVADESAENCCKSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLQHKDDNPMLPRLVRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCCASLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIKQNCSELF EQLGEYKFQNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALEVDETYVPKEFNAETFTPHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAVMDDFAA FVEKCKKADDKETCFAEEGPKLVAASKAALGL
124.	HALB C34A	artificial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQAPFEDHVKLVNEVTEFAKTCVADESAENCCKSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLQHKDDNPMLPRLVRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCCASLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIKQNCSELF EQLGEYKFQNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALEVDETYVPKEFNAETFTPHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAVMDDFAA FVEKCKKADDKETCFAEEGKLVAAQAALGL
125.	HALB7 C34A	artificial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQAPFEDHVKLVNEVTEFAKTCVADESAENCCKSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLQHKDDNPMLPRLVRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCCASLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIKQNCSELF EQLGEYKFQNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALEVDETYVPKEFNAGTFTPHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAAMDDFAA FVEKCKKADDKETCFAEEGKLVAAQAALGL
126.	HALB098 C34A	artificial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQAPFEDHVKLVNEVTEFAKTCVADESAENCCKSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLQHKDDNPMLPRLVRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCCASLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIKQNCSELF EQLGEYKFQNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALEVDETYVPKEFNAETFTPHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAVMDDFAA FVEKCKKADDKETCFAEEGPKLVAQAALGL
127.	HALB114 C34A	artificial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQAPFEDHVKLVNEVTEFAKTCVADESAENCCKSLHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLQHKDDNPMLPRLVRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELLFFFA KRYKAAPTECCQAADKAACLLPKLDELDRDEGKASSAKQRLKCCASLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYFYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLIKQNCSELF EQLGEYKFQNALLVRYTKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALEVDETYVPKEFNAETFTPHADICTLSEKERQIKQTALVELVKHKPKATKEQLKAVMDDFAA FVEKCKKADDKETCFAEEGPHLVAASKAALGL

TABLE 30-continued

Sequence Listing			
SEQ ID No:	Descrip- tion	Source	Sequence
128.	HALB254 C34A	artifi- cial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQAPPFEDHVKLVNEVTEFAKTCVADESAENCDKS LHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLOHKDDNPNLPRLVRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELFFFA KRYKAAPTECCQAADKAACLLPKLDELRLDEGKASSAKQRLKCSLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDS ISSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYEYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLI KQNCSELF EQLGEYKFQNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALGDVDETYPKEFNAETFTFHADICTLSEKERQIKKQTALVELVKHKPKATKEQLKAVMDKFAA FVEKCKADDKETCFAEEGPKLVAASQAALGL
129.	HALB253 C34A	artifi- cial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQAPPFEDHVKLVNEVTEFAKTCVADESAENCDKS LHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLOHKDDNPNLPRLVRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELFFFA KRYKAAPTECCQAADKAACLLPKLDELRLDEGKASSAKQRLKCSLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDS ISSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYEYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLI KQNCSELF EQLGEYKFQNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALGDVDETYPKEFNAETFTFHADICTLSEKERQIKKQTALVELVKHKPKATKEQLKAVMDKFAA FVEKCKADDKETCFAEEGPKLVAASQAALGL
130.	HALB131 C34A	artifi- cial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQAPPFEDHVKLVNEVTEFAKTCVADESAENCDKS LHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLOHKDDNPNLPRLVRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELFFFA KRYKAAPTECCQAADKAACLLPKLDELRLDEGKASSAKQRLKCSLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDS ISSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYEYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLI KQNCSELF EQLGEYKFQNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALGDVDETYPKEFNAETFTFHADICTLSEKERQIKKQTALVELVKHKPKATKEQLKAVMDKFAA FVEKCKADDKETCFAEEGPKLVAASQAALGL
131.	HALB135 C34A	artifi- cial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQAPPFEDHVKLVNEVTEFAKTCVADESAENCDKS LHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLOHKDDNPNLPRLVRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELFFFA KRYKAAPTECCQAADKAACLLPKLDELRLDEGKASSAKQRLKCSLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDS ISSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYEYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLI KQNCSELF EQLGEYKFQNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALGDVDETYPKEFNAETFTFHADICTLSEKERQIKKQTALVELVKHKPKATKEQLKAVMDFAA FVEKCKADDKETCFAEEGPKLVAASKAALGL
132.	HALB133 C34A	artifi- cial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQAPPFEDHVKLVNEVTEFAKTCVADESAENCDKS LHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLOHKDDNPNLPRLVRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELFFFA KRYKAAPTECCQAADKAACLLPKLDELRLDEGKASSAKQRLKCSLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDS ISSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYEYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLI KQNCSELF EQLGEYKFQNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALGDVDETYPKEFNAETFTFHADICTLSEKERQIKKQTALVELVKHKPKATKEQLKAVMDKFAA FVEKCKADDKETCFAEEGPKLVAASKAALGL
133.	HALB234 C34A	artifi- cial	DAHKSEVAHRFKDLGEENFKALVLIIFAQYLQQAPPFEDHVKLVNEVTEFAKTCVADESAENCDKS LHTLFGDKLCTVAT LRETYGEMADCCAKQEPERNECFLOHKDDNPNLPRLVRPEVDVMCTAFHDNEETFLKKYLYEIAARRHPYFYAPELFFFA KRYKAAPTECCQAADKAACLLPKLDELRLDEGKASSAKQRLKCSLQKFGERAFAKAWAVARLSQRFPKAEFAEVS KLVT LTKVHTECCHGDLLECADDRADLAKYICENQDS ISSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVC KNYAEAKDVFLGMFLYEYARRHPDYSVVLRLAKTYETTLKCCAAADPHECYAKVFDEFKPLVEEPQNLI KQNCSELF EQLGEYKFQNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVT KCCTESLVNRRPCFSALGDVDETYPKEFNAETFTFHADICTLSEKERQIKKQTALVELVKHKPKATKEQLKAVMDFAA FVEKCKADDKETCFAEEGPKLVAASKAALGL
134.	Ab156	artifi- cial	RDWDFDFVGGGTPVGG
135.	linear FcRn binding peptide	artifi- cial	QRFVTGHFGGLXPANG

TABLE 30-continued

Sequence Listing			
SEQ ID NO:	Description	Source	Sequence
136.	linear FcRn binding peptide Y	artificial	QRFVTGHFGGLYPANG
137.	linear FcRn binding peptide H	artificial	QRFVTGHFGGLHPANG
138.	core FcRn binding peptide H	artificial	TGHFGGLHP
139.	cyclic FcRn binding peptide H	artificial	QRFCTGHFGGLHPCNG
140.	Cross body 1 HC		ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQ TYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKF NWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYDTTPPVLDSDGSFFLYSDLTVDKSRWQQGNVFCSCVMHEALH NHYTQKSLSLSPGK
141.	Cross body 1 LC		GQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKADSSPVKAGVETTTPSKQSNNKYAASSYLSLTPEQW KSHRSYSQCQVTHEGSTVEKTVAPTECSEPKSSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPE VKFNWYVDGVEVHNAKTKPCEEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTL PPSRKEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLKSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHE ALHNHNTQKSLSLSPGK
142.	Cross body 2 HC		ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSNFGTQ TYTCNVDHKPSNTKVDKTVEPKSSDKTHTCPPCPAEEAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKF NWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYDTTPPVLDSDGSFFLYSDLTVDKSRWQQGNVFCSCVMHEALH NHYTQKSLSLSPGK
143.	Cross body 2 LC		GQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKADSSPVKAGVETTTPSKQSNNKYAASSYLSLTPEQW KSHRSYSQCQVTHEGSTVEKTVAPTECSEPKSSDKTHTCPPCPAEEAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDV HEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREP QVYTLPPSRKEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLKSDGSFFLYSKLTVDKSRWQQGNVFC SVMHEALHNHNTQKSLSLSPGK
144.	Hetero-Fc binder Fc		DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGST YRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRKEMTKNQVSLTCLVKGFYPSDIA VEWESNGQPENNYKTTPPVLKSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNHNTQKSLSLSPGK
145.	Hetero-Fc partner Fc		DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGST YRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRKEMTKNQVSLTCLVKGFYPSDIA VEWESNGQPENNYDTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNHNTQKSLSLSPGK
146.	Maxibody 1 target Fc		EPKSSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEE QYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRKEMTKNQVSLTCLVKGFY PSDIAVEWESNGQPENNYKTTPPVLKSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNHNTQKSLSLSPGK

TABLE 30-continued

Sequence Listing			
SEQ ID NO:	Description	Source	Sequence
147.	Maxibody 1 CD3 Fc		EPKSSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEE QYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFY PSDIAVEWESNGQPENNYDTTPPVLDSDGSFFLYSDLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK
148.	Maxibody 2 target Fc		EPKSSDKTHTCPPCPAEEAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREE QYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRKEMTKNQVSLTCLVKGFY PSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK
149.	Maxibody 2 CD3 Fc		EPKSSDKTHTCPPCPAEEAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREE QYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFY PSDIAVEWESNGQPENNYDTTPPVLDSDGSFFLYSDLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK
150.	Mono	Fc	APELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYDTTPPVLDSDGSFFLYSDLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK
151.	MS_1	VH CDR1	SSSYW
152.	MS_1	VH CDR2	SIYYSGITNYPNLSKS
153.	MS_1	VH CDR3	PSNYDAFDI
154.	MS_1	VL CDR1	TGSSSNIGAGYDVH
155.	MS_1	VL CDR2	GNSKRPS
156.	MS_1	VL CDR3	QSYDSSLGGWV
157.	MS_1	VH	QVQLQESGPGLVKPSSETLSLTCTVSGGSISSSSYYWGWIQQPPGKLEWIGSIYYSGITNYPNLSKSRVTISVDTSKNQ FSLKLSSTAAADTAVYYCARPSNYDAFDIWIQQGTMTVSSSGGGSGGGSGGGSGSVLTQPPSVSGAPGQRVTISCTG
158.	MS_1	VL	QSVLTQPPSVSGAPGQRVTISCTGSSSNIGAGYDVHWYQQVPGTAPKLLIYGNSKRPSGVPDRFSGSKSGTSASLAITG LQAEDEADYYCQSYDSSLGGWVFGGKTLTVL
159.	MS_1	scFv	QVQLQESGPGLVKPSSETLSLTCTVSGGSISSSSYYWGWIQQPPGKLEWIGSIYYSGITNYPNLSKSRVTISVDTSKNQ FSLKLSSTAAADTAVYYCARPSNYDAFDIWIQQGTMTVSSSGGGSGGGSGGGSGSVLTQPPSVSGAPGQRVTISCTG SSSNIGAGYDVHWYQQVPGTAPKLLIYGNSKRPSGVPDRFSGSKSGTSASLAITGLQAEDEADYYCQSYDSSLGGWVFG GGKTLTVL
160.	MS_1	bi- speci- fic mole- cule	QVQLQESGPGLVKPSSETLSLTCTVSGGSISSSSYYWGWIQQPPGKLEWIGSIYYSGITNYPNLSKSRVTISVDTSKNQ FSLKLSSTAAADTAVYYCARPSNYDAFDIWIQQGTMTVSSSGGGSGGGSGGGSGSVLTQPPSVSGAPGQRVTISCTG SSSNIGAGYDVHWYQQVPGTAPKLLIYGNSKRPSGVPDRFSGSKSGTSASLAITGLQAEDEADYYCQSYDSSLGGWVFG GGKTLTVLSSGGGSEVLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKLEWVARIRSKYNNYATYYADS VKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYI SYWAYWGQTLTVTVSSGGGGSGGGSGGGGQTIV TQEPSTLVSPGGTVTLTCSGSTGAVTSGNYPNWNVQKPGQAPRGLIGGTFKFLAPGTAPRFGSLLGGKAALTLSGVQPE DEAEYYCVLWYSNRWVFGGKTLTVL
161.	MS_2	VH CDR1	DHYMS
162.	MS_2	VH CDR2	YISSSGSTIYADSVKG
163.	MS_2	VH CDR3	DLGPSFDY
164.	MS_2	VL CDR1	RASQGISSWLA
165.	MS_2	VL CDR2	AASRLQS
166.	MS_2	VL CDR3	QQANSFPRT
167.	MS_2	VH	QVQLVESGGGVVQPGRLRLSCLASGFTFSDHYMSWIRQTPGKLEWVSYISSSGSTIYADSVKGRFTISRDNAKNSL YLQMNSLRADTAVYYCARDLGPSFDYWGQSLTVTVSS
168.	MS_2	VL	DIQMTQSPSSVSASVGRVITITCRASQGISSWLAWYQQKPKGAPKLLIDAASRLQSGVPSRFSGSGSGTDFTLTISSLQ PEDFATYYCQQANSFPRTFGQGTKVEIK
169.	MS_2	scFv	QVQLVESGGGVVQPGRLRLSCLASGFTFSDHYMSWIRQTPGKLEWVSYISSSGSTIYADSVKGRFTISRDNAKNSL YLQMNSLRADTAVYYCARDLGPSFDYWGQSLTVTVSSGGGGSGGGSGGGSDIQMTQSPSSVSASVGRVITITCRAS QGIISSWLAWYQQKPKGAPKLLIDAASRLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQANSFPRTFGQGTKVE IK

TABLE 30-continued

Sequence Listing			
SEQ ID NO:	Description	Source	Sequence
170.	MS_2	bi-speci- fic mole- cule	QVQLVESGGGVQPGRSLRLSCAASGFTFSDHYMSWIRQTPGKGLEWVSYYSSSGSTIYYADSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDLGPSFDYWGQGLVTVSSGGGGSGGGSGGGGSDIQMTQSPSSVSASVGDRTITCRASQGIISSWLAWYQQKPGKAPKLLIDAASRLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQANSFPRTFGQGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGGSGGGSGGGGSQTVVTQEPSSLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
171.	MS_3	VH CDR1	DYYMT
172.	MS_3	VH CDR2	YISSSGSTIYYADSVKG
173.	MS_3	VH CDR3	DRNSHFDY
174.	MS_3	VL CDR1	RASQGITRWLA
175.	MS_3	VL CDR2	AASVLQS
176.	MS_3	VL CDR3	QQSNSFPRT
177.	MS_3	VH	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMTWIRQAPGKGLEWISYISSSGSTIYYADSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDRNHFDYWGQGLVTVSS
178.	MS_3	VL	DIQMTQSPSSVSASVGDRTITCRASQGITRWLAWYQQKPGKAPKLLIYAASVLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQSNSFPRTFGQGTKVEIK
179.	MS_3	scFv	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMTWIRQAPGKGLEWISYISSSGSTIYYADSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDRNHFDYWGQGLVTVSSGGGGSGGGSGGGGSDIQMTQSPSSVSASVGDRTITCRASQGITRWLAWYQQKPGKAPKLLIYAASVLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQSNSFPRTFGQGTKVEIK
180.	MS_3	bi-speci- fic mole- cule	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMTWIRQAPGKGLEWISYISSSGSTIYYADSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDRNHFDYWGQGLVTVSSGGGGSGGGSGGGGSDIQMTQSPSSVSASVGDRTITCRASQGITRWLAWYQQKPGKAPKLLIYAASVLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQSNSFPRTFGQGTKVEIKSRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGGSGGGSGGGGSQTVVTQEPSSLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
181.	MS_4	VH CDR1	DYYMT
182.	MS_4	VH CDR2	YISSSGSTIYYADSVKG
183.	MS_4	VH CDR3	DRNSHFDY
184.	MS_4	VL CDR1	RASQGINTWLA
185.	MS_4	VL CDR2	GASGLQS
186.	MS_4	VL CDR3	QQAQSFPRPT
187.	MS_4	VH	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMTWIRQAPGKGLEWLSYISSSGSTIYYADSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDRNHFDYWGQGLVTVSS
188.	MS_4	VL	DIQMTQSPSSVSASVGDRTITCRASQGINTWLAWYQQKPGKAPKLLIYGASGLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQAQSFPRPTFGQGTKVEIK
189.	MS_4	scFv	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMTWIRQAPGKGLEWLSYISSSGSTIYYADSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDRNHFDYWGQGLVTVSSGGGGSGGGSGGGGSDIQMTQSPSSVSASVGDRTITCRASQGINTWLAWYQQKPGKAPKLLIYGASGLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQAQSFPRPTFGQGTKVEIK
190.	MS_4	bi-speci- fic mole- cule	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMTWIRQAPGKGLEWLSYISSSGSTIYYADSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDRNHFDYWGQGLVTVSSGGGGSGGGSGGGGSDIQMTQSPSSVSASVGDRTITCRASQGINTWLAWYQQKPGKAPKLLIYGASGLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQAQSFPRPTFGQGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGGSGGGSGGGGSQTVVTQEPSSLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL

TABLE 30-continued

Sequence Listing			
SEQ ID NO:	Description	Source	Sequence
191. MS_5		VH CDR1	DHYMS
192. MS_5		VH CDR2	YISSSGGIYYADSVKG
193. MS_5		VH CDR3	DVGSHFDY
194. MS_5		VL CDR1	RASQDISRWLA
195. MS_5		VL CDR2	AASRLQS
196. MS_5		VL CDR3	QQAQSFPR
197. MS_5		VH	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDHYMSWIRQAPGKGLEWFSYISSSGGIYYADSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDVGSHFDYWGQGLTVTVSS
198. MS_5		VL	DIQMTQSPSSVSASVGDRTITCRASQDISRWLAHWYQKPKGAPKLLISAASRLQSGVPSRFSGSGSGTDFTLTISLQPEDFAIYYCQQAQSFPRTFGQGTKEIK
199. MS_5		scFv	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDHYMSWIRQAPGKGLEWFSYISSSGGIYYADSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDVGSHFDYWGQGLTVTVSSGGGGSGGGSGGGSDIQMTQSPSSVSASVGDRTITCRASQDISRWLAHWYQKPKGAPKLLISAASRLQSGVPSRFSGSGSGTDFTLTISLQPEDFAIYYCQQAQSFPRTFGQGTKEIK
200. MS_5	bi-specific molecule		QVQLVESGGGLVKPGGSLRLSCAASGFTFSDHYMSWIRQAPGKGLEWFSYISSSGGIYYADSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDVGSHFDYWGQGLTVTVSSGGGGSGGGSGGGSDIQMTQSPSSVSASVGDRTITCRASQDISRWLAHWYQKPKGAPKLLISAASRLQSGVPSRFSGSGSGTDFTLTISLQPEDFAIYYCQQAQSFPRTFGQGTKEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGSGGGGQTVVTQEPSLTVSPGGTVTLTCSGSTGAVTSGNYPNWWVQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYYCVLWYSNRWVFGGGLKLTVL
201. MS_6		VH CDR1	DHYMS
202. MS_6		VH CDR2	YISNSGSIYYVDSVKG
203. MS_6		VH CDR3	DVRTAFDY
204. MS_6		VL CDR1	RASQSIGSWLA
205. MS_6		VL CDR2	AASSLQS
206. MS_6		VL CDR3	QQANSFPR
207. MS_6		VH	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDHYMSWIRQAPGKGLEWISYISNSGSIYYVDSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDVRTAFDYWGQGLTVTVSS
208. MS_6		VL	DIQMTQSPSSVSASVGDRTITCRASQSIGSWLAHWYQKPKGAPNLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQAQSFPRTFGQGTKEIK
209. MS_6		scFv	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDHYMSWIRQAPGKGLEWISYISNSGSIYYVDSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDVRTAFDYWGQGLTVTVSSGGGGSGGGSGGGSDIQMTQSPSSVSASVGDRTITCRASQSIGSWLAHWYQKPKGAPNLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQAQSFPRTFGQGTKEIK
210. MS_6	bi-specific molecule		QVQLVESGGGLVKPGGSLRLSCAASGFTFSDHYMSWIRQAPGKGLEWISYISNSGSIYYVDSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDVRTAFDYWGQGLTVTVSSGGGGSGGGSGGGSDIQMTQSPSSVSASVGDRTITCRASQSIGSWLAHWYQKPKGAPNLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQAQSFPRTFGQGTKEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGSGGGGQTVVTQEPSLTVSPGGTVTLTCSGSTGAVTSGNYPNWWVQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYYCVLWYSNRWVFGGGLKLTVL
211. MS_7		VH CDR1	SKFMT
212. MS_7		VH CDR2	VIYSGGKTYADSVKG
213. MS_7		VH CDR3	DSGGWGYFDY
214. MS_7		VL CDR1	KSSQSVLYSSNNKNYLA
215. MS_7		VL CDR2	WASTRES

TABLE 30-continued

Sequence Listing			
SEQ ID NO:	Descrip- tion	Source	Sequence
216. MS_7		VL CDR3	QQYYSTPPT
217. MS_7		VH	EVQLVESGGGLIQPGSLRLSCAVSGFTVSSKFMWVRQAPGKGLEWVSVIYSGGKTYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCARDSSGGWGYFDYWGQGLTVTVSS
218. MS_7		VL	DIVMTQSPDSLAVSLGERATINCKSSQSVLYSSNNKNYLAWYQQKPGQPPKLLIYWASTRESGVPDRFSGSGSDTFTL TISSLQAEDVAVYYCQQYYSTPPTFGQGTKVEIK
219. MS_7		scFv	EVQLVESGGGLIQPGSLRLSCAVSGFTVSSKFMWVRQAPGKGLEWVSVIYSGGKTYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCARDSSGGWGYFDYWGQGLTVTVSSGGGSGGGGSGGGSDIVMTQSPDSLAVSLGERATINCKS SQSVLYSSNNKNYLAWYQQKPGQPPKLLIYWASTRESGVPDRFSGSGSDTFTLTISSLQAEDVAVYYCQQYYSTPPTF GQGTKVEIK
220. MS_7	bi- speci- fic mole- cule		EVQLVESGGGLIQPGSLRLSCAVSGFTVSSKFMWVRQAPGKGLEWVSVIYSGGKTYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCARDSSGGWGYFDYWGQGLTVTVSSGGGSGGGGSGGGSDIVMTQSPDSLAVSLGERATINCKS SQSVLYSSNNKNYLAWYQQKPGQPPKLLIYWASTRESGVPDRFSGSGSDTFTLTISSLQAEDVAVYYCQQYYSTPPTF GQGTKVEIKSGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYAD SVKDRFTISRDDS KNTAYLQMNNLKTEDTAVYYCVRHGNFNSYISYWAYWGQGLTVTVSSGGGSGGGGSGGGGSGQTV VTQEPSTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLGGKAALTL SGVQP EDEAEYYCVLWYSNRWVFGGKTLTVL
221. MS_8		VH CDR1	SYYN
222. MS_8		VH CDR2	RIYNGNTYINPSLKS
223. MS_8		VH CDR3	PKLGIDAFDI
224. MS_8		VL CDR1	TGSSSNIGAGYDVH
225. MS_8		VL CDR2	GNSNRPS
226. MS_8		VL CDR3	QSYDSSLSGWV
227. MS_8		VH	QVQLQESGPGLVKPSSETLSLTCTVSGGSISSYYWNWIRQPPGKGLEWIGRIYNGNTYINPSLKS RVTTISGDT SKNQFS LKLSSVTAADTAVYYCARPKLGIDAFDIWGQGTMTVTVSS
228. MS_8		VL	QSVLTQPPSVSGAPGQRVTISCTGSSSNIGAGYDVHWYQQLPGTAPKLLIYGNSNRPSGVPDRFSGSKSGTSASLAITG LQAED EADYYCQSYDSSLSGWVFGGKTLTVL
229. MS_8		scFv	QVQLQESGPGLVKPSSETLSLTCTVSGGSISSYYWNWIRQPPGKGLEWIGRIYNGNTYINPSLKS RVTTISGDT SKNQFS LKLSSVTAADTAVYYCARPKLGIDAFDIWGQGTMTVTVSSGGGSGGGGSGGGGQSVLTQPPSVSGAPGQRVTISCTGS SSNIGAGYDVHWYQQLPGTAPKLLIYGNSNRPSGVPDRFSGSKSGTSASLAITGLQAED EADYYCQSYDSSLSGWVFGG GTKLTVL
230. MS_8	bi- speci- fic mole- cule		QVQLQESGPGLVKPSSETLSLTCTVSGGSISSYYWNWIRQPPGKGLEWIGRIYNGNTYINPSLKS RVTTISGDT SKNQFS LKLSSVTAADTAVYYCARPKLGIDAFDIWGQGTMTVTVSSGGGSGGGGSGGGGQSVLTQPPSVSGAPGQRVTISCTGS SSNIGAGYDVHWYQQLPGTAPKLLIYGNSNRPSGVPDRFSGSKSGTSASLAITGLQAED EADYYCQSYDSSLSGWVFGG GTKLTVLSSGGGSEVQLVESGGGLVQPGGSLKLSAASGFTENKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSV KDRFTISRDDS KNTAYLQMNNLKTEDTAVYYCVRHGNFNSYISYWAYWGQGLTVTVSSGGGSGGGGSGGGGSGTVVT QEPSTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLGGKAALTL SGVQPED EAEYYCVLWYSNRWVFGGKTLTVL
231. Human MSLN v1 NM_005823		human	MALPTARPLLGSCGTPALGSLFLFLSLGWVQPSRTLAGETGQEAAPLDGVLANPPN ISSLSPRQLLGFP CAEVSGLSLST ERVRELAVALAQKNV KLS TEQLRCLAHRLSEPPEDLDALPLDLLLFLNPDAFSGPQACTRFFSRI TKANVDLLPRGAPE RQRLLPAALACWGVRSLLSEADVRLAGGLACDLPGRFVAESA EVLLPRLVSCPGPLDQDQQAARAALQGGGPPYGGP STWSVSTMDALRGLLPVLGQPI IRSIQGIVA AWRQRSSRDP SWRPQERTILRPRFRREVEKTACPSGKKAREIDESLI FYKWELEACVDAALLATQMDRVNAIPFTYEQLDVLKHKLDELYPQGYPESV IQHLGYLFLKMSPED IRKWNVT SLETL KALLEVNKGHEMSPQVATLIDRFVKGRGQLDKD TLDLTAFYPGYLCSLSPEELSVPSSIWAVRPQDLDCDPRQLD VLYPKARLAFQNMNGSEYFVKIQSFLGGAPTEDLKALSQQNVSMDLATFMKLRD DAVLPLTVAEVQKLLGPHVEGLKAE ERHRPVRDWILRQRQDDLDTLGLGLQGQIPNGYLVLDLSMQEALSGTPCLLGGPVLTVLALLLASTLA
232. Human MSLN v2 NM_013404		human	MALPTARPLLGSCGTPALGSLFLFLSLGWVQPSRTLAGETGQEAAPLDGVLANPPN ISSLSPRQLLGFP CAEVSGLSLST ERVRELAVALAQKNV KLS TEQLRCLAHRLSEPPEDLDALPLDLLLFLNPDAFSGPQACTRFFSRI TKANVDLLPRGAPE RQRLLPAALACWGVRSLLSEADVRLAGGLACDLPGRFVAESA EVLLPRLVSCPGPLDQDQQAARAALQGGGPPYGGP STWSVSTMDALRGLLPVLGQPI IRSIQGIVA AWRQRSSRDP SWRPQERTILRPRFRREVEKTACPSGKKAREIDESLI FYKWELEACVDAALLATQMDRVNAIPFTYEQLDVLKHKLDELYPQGYPESV IQHLGYLFLKMSPED IRKWNVT SLETL KALLEVNKGHEMSPQAPRRPLPQVATLIDRFVKGRGQLDKD TLDLTAFYPGYLCSLSPEELSVPSSIWAVRPQDLDCDPRQLD VLYPKARLAFQNMNGSEYFVKIQSFLGGAPTEDLKALSQQNVSMDLATFMKLRD DAVLPLTVAEVQKLLGPHVEGLKAE HVEGLKAEERHRPVRDWILRQRQDDLDTLGLGLQGQIPNGYLVLDLSMQEALSGTPCLLGGPVLTVLALLLASTLA

TABLE 30-continued

Sequence Listing			
SEQ ID NO:	Description	Source	Sequence
233.	Human MSLN v6 AY743922	human	MALPTARPLLGSCGTPALGSLFLFLSLGWVQPSRTLAGETGQEAAPLDGVLANPPNIISSLSRQLLGFP CAEVSGSLST ERVRELAVALAQKNVKLSTEQLRCLAHRLSEPPEDLDALPLDLLLFLNPDADFSGPQACTHFFSRI TKANVDLLPRGAPE RQRLPAAALACWGVGSGSLLEADVRALGGLACDLPGRFVAESA EVLLPRLVSCPGPLDQDQQAARAALQGGGPPYGGP STWSVTMDALRGLLPVLGQPI IRSIPQGIVA AWRQRSSRDPSWRQP ERTILRPRFRREVEKTACPSGKKAREIDESLI FYKKWELEACVDAALLATQMDRVNAIPFTYEQLDVLKHKLDELYPQGYPESVIQHLGYLFLKMSPEDIRKWNVT SLETL KALLEVNKGHEMSPQVATLIDRFVKGRGQLDKDTLDTLTAFYPGYLCSLSPEELSSVPSSIWAVRPQDLDTCDPRQLD VLYPKARLAFQNMNGSEYFVKIQSFLGGAPTE DLKALSQQNVSMDLATFMKLRTDAVLPLTVAEVQKLLGPHVEGLKAE ERHRPVRDWILRQRQDDLDTLGLGLQGGINPGYLVLDLSVQEAALSGTPCLLGGPVPVTLVALLLASTLA
234.	cyno MSLN v1 LMR C52457	macaque	MALPMARPLSGSCGTPALGSLFLFLSLGWVQPSRVLAGETRQEAAPLDGILTNPADIASLSRQLLGFT CVEVSGSLST ELVQELAVALGQKNVKLSAEQLRCLAHRLSEPPEDLDALPLDLLLFLNPDADFSGPQACTHFFSRVAKANVDLLPRGAPE RQRLPAAALTCWGVGSGSLLEADVRALGGLACDLPGRFVAESA EVVLPRLVRCLGPLDQDQQAARAALQGGGPPYGGP STWSISTLDDLQSLLPVLGQPI IHSIPQGILAAWRQRSSRDPSWQQPEQTVLRPRFRRDVERTTCPEKEVHEIDENLI FYKKRELEACVDAALLAAQMDRVDAIPFTYEQLDVLKHKLDELYPQGYPESVIRHLGHLFLKMSPEDIRKWNVT SLETL KALLKVS KGHEMSAQVATLIDRVVVG RGQLDKDTADTLTAFCPGCLCSLSPELSSVPSSIVIGAVRPQDLDTCDGRQLD VLYPKARLAFQNMNGSEYFVKIRPFLGGAPTE DLKALSQQNVSMDLATFMKLRTDAVLPLTVAEVQKLLGPHVEGLKVE EQHSPVRDWILKQRQDDLDTLGLGLQGGINPGYLI L DLSVREALSGTPCLLGGPVPVTLI LALLLASTLA
235.	Mega-karyocyte-potenti-ating factor NM_005823	human	MALPTARPLLGSCGTPALGSLFLFLSLGWVQPSRTLAGETGQEAAPLDGVLANPPNIISSLSRQLLGFP CAEVSGSLST ERVRELAVALAQKNVKLSTEQLRCLAHRLSEPPEDLDALPLDLLLFLNPDADFSGPQACTRFFSRI TKANVDLLPRGAPE RQRLPAAALACWGVGSGSLLEADVRALGGLACDLPGRFVAESA EVLLPRLVSCPGPLDQDQQAARAALQGGGPPYGGP STWSVTMDALRGLLPVLGQPI IRSIPQGIVA AWRQRSSRDPSWRQPER
236.	MSLN, cleaved form NM_005823	human	EVEKTACPSGKKAREIDESLIFYKKWELEACVDAALLATQMDRVNAIPFTYEQLDVLKHKLDELYPQGYPESVIQHLGY LFLKMSPEDIRKWNVT SLETLKALLEVNKGHEMSPQVATLIDRFVKGRGQLDKDTLDTLTAFYPGYLCSLSPEELSSVP PSSIWAVRPQDLDTCDPRQLDVLYPKARLAFQNMNGSEYFVKIQSFLGGAPTE DLKALSQQNVSMDLATFMKLRTDAVL PLTVAEVQKLLGPHVEGLKAEERHRPVRDWILRQRQDDLDTLGLGLQGGINPGYLVLDLSMQEALS
237.	C-terminal cleavage NM_005823	human	GTPCLLGGPVPVTLVALLLASTLA
238.	MSLN-E 1	human	EVEKTACPSGKKAREIDESLIFYKKWELEACVDAALLATQMDRVNAIPFTY
239.	MSLN-E 2	human	EQLDVLKHKLDELYPQGYPESVIQHLGYLFLKMSPEI
240.	MSLN-E 3	human	RKWNVT SLETLKALLEVNKGHEMSPQVATLIDRFVKGRGQLDKDTLDTLTAFYPGYLCSLSPEELSSVP
241.	MSLN-E 4	human	PSSIWAVRPQDLDTCDPRQLDVLYPKARLAFQNMNGSEYFVKIQSFLG
242.	MSLN-E 5	human	GAPTE DLKALSQQNVSMDLATFMKLRTDAVLPLTVAEVQKLLGP
243.	MSLN-E 6	human	HVEGLKAEERHRPVRDWILRQRQDDLDTLGLGLQGGINPGYLVLDLSMQEAL
244.	MSLN-E 1 + 2	human	EVEKTACPSGKKAREIDESLIFYKKWELEACVDAALLATQMDRVNAIPFTYEQLDVLKHKLDELYPQGYPESVIQHLGY LFLKMSPEI
245.	MSLN-E 2 + 3	human	EQLDVLKHKLDELYPQGYPESVIQHLGYLFLKMSPEDIRKWNVT SLETLKALLEVNKGHEMSPQVATLIDRFVKGRGQL DKDTLDTLTAFYPGYLCSLSPEELSSVP
246.	hu orl MSLN-E1 mu	artifi- cial	MALPTARPLLGSCGTPALGSLFLFLSLGWVQPSRTLAGETGQEAAPLDGVLANPPNIISSLSRQLLGFP CAEVSGSLST ERVRELAVALAQKNVKLSTEQLRCLAHRLSEPPEDLDALPLDLLLFLNPDADFSGPQACTRFFSRI TKANVDLLPRGAPE RQRLPAAALACWGVGSGSLLEADVRALGGLACDLPGRFVAESA EVLLPRLVSCPGPLDQDQQAARAALQGGGPPYGGP STWSVTMDALRGLLPVLGQPI IRSIPQGIVA AWRQRSSRDPSWRQP ERTILRPRFRREVEKTACPSGKKAREIDESLI FYQNWELEACVDGTM LARQMDLVNEIPFTYEQLDVLKHKLDELYPQGYPESVIQHLGYLFLKMSPEDIRKWNVT SLETL KALLEVNKGHEMSPQVATLIDRFVKGRGQLDKDTLDTLTAFYPGYLCSLSPEELSSVPSSIWAVRPQDLDTCDPRQLD VLYPKARLAFQNMNGSEYFVKIQSFLGGAPTE DLKALSQQNVSMDLATFMKLRTDAVLPLTVAEVQKLLGPHVEGLKAE ERHRPVRDWILRQRQDDLDTLGLGLQGGINPGYLVLDLSMQEALSGTPCLLGGPVPVTLVALLLASTLA
247.	hu orl MSLN-E2 mu	artifi- cial	MALPTARPLLGSCGTPALGSLFLFLSLGWVQPSRTLAGETGQEAAPLDGVLANPPNIISSLSRQLLGFP CAEVSGSLST ERVRELAVALAQKNVKLSTEQLRCLAHRLSEPPEDLDALPLDLLLFLNPDADFSGPQACTRFFSRI TKANVDLLPRGAPE RQRLPAAALACWGVGSGSLLEADVRALGGLACDLPGRFVAESA EVLLPRLVSCPGPLDQDQQAARAALQGGGPPYGGP STWSVTMDALRGLLPVLGQPI IRSIPQGIVA AWRQRSSRDPSWRQP ERTILRPRFRREVEKTACPSGKKAREIDESLI FYKKWELEACVDAALLATQMDRVNAIPFTYEQLSIFKHKLDKTYPQGYPESLIQQLGHFRYVSPEDIRKWNVT SLETL KALLEVNKGHEMSPQVATLIDRFVKGRGQLDKDTLDTLTAFYPGYLCSLSPEELSSVPSSIWAVRPQDLDTCDPRQLD VLYPKARLAFQNMNGSEYFVKIQSFLGGAPTE DLKALSQQNVSMDLATFMKLRTDAVLPLTVAEVQKLLGPHVEGLKAE ERHRPVRDWILRQRQDDLDTLGLGLQGGINPGYLVLDLSMQEALSGTPCLLGGPVPVTLVALLLASTLA

TABLE 30-continued

Sequence Listing			
SEQ ID NO:	Description	Source	Sequence
248.	hu orl MSLN- E3 mu	artifi- cial	MALPTARPLLGSCGTPALGSLFLFSLGWQPSRTLAGETGQEAAPLDGVLANPPNISSLSRQLLGFPFCAEVSGSLST ERVRELAVALAQKNVKLSTEQLRCLAHRLSEPPEDLDALPLDLLLFLNPDADFSGPQACTRFFFSRI TKANVDLLPRGAPE RQRLPAALACWGVRSLLSEADVRLGGLACDLPGRFVAESAIEVLLPRLVSCPGPLDQDQQAARAALQGGGPPYGGP STWSVTMDALRGLLPVLGQPI IRSIPQGIVAARQRSSRDPSWRQPETILRPRFRREVEKTACPSGKKAREIDESLI FYKKWELEACVDAALLATQMDRVNAIPFTYEQLDVLKHKLDELYPQGYPESVIQHLGYLFLKMSPEDIRKWNVTSPDTV KTLKVS KGQKMNAQAIALVACYL RGGGQLDEDMVKALGDI PLSYLCDFSPQDLHSVPPSSIWAVRPQDLDTCDPRQLD VLYPKARLAFQNMNGSEYFVKIQSFLGGAPTEDLKALSQQNVSMDLATFMKLRTDAVLP LPTVAEVQKLLGPHVEGLKAE ERHRPVRDWILRQRQDDLDLTGLGLQGGINPGYLVLDLSMQEALSGTPCLLGGPGPVLTVLALLLASTLA
249.	hu orl MSLN- E4 mu	artifi- cial	MALPTARPLLGSCGTPALGSLFLFSLGWQPSRTLAGETGQEAAPLDGVLANPPNISSLSRQLLGFPFCAEVSGSLST ERVRELAVALAQKNVKLSTEQLRCLAHRLSEPPEDLDALPLDLLLFLNPDADFSGPQACTRFFFSRI TKANVDLLPRGAPE RQRLPAALACWGVRSLLSEADVRLGGLACDLPGRFVAESAIEVLLPRLVSCPGPLDQDQQAARAALQGGGPPYGGP STWSVTMDALRGLLPVLGQPI IRSIPQGIVAARQRSSRDPSWRQPETILRPRFRREVEKTACPSGKKAREIDESLI FYKKWELEACVDAALLATQMDRVNAIPFTYEQLDVLKHKLDELYPQGYPESVIQHLGYLFLKMSPEDIRKWNVTSLLETL KALLEVNKGHEMSPQVATLIDRFVKGRGQLDKDTLDTLTAFYPGYLCSLSPEELSSVPPSSIWAVRPQDLDTCDPRQLD LLYQKACSAFQNVSGLEYFEKI KTFLLGGAPTEDLKALSQQNVSMDLATFMKLRTDAVLP LPTVAEVQKLLGPHVEGLKAE ERHRPVRDWILRQRQDDLDLTGLGLQGGINPGYLVLDLSMQEALSGTPCLLGGPGPVLTVLALLLASTLA
250.	hu orl MSLN- E5 mu	artifi- cial	MALPTARPLLGSCGTPALGSLFLFSLGWQPSRTLAGETGQEAAPLDGVLANPPNISSLSRQLLGFPFCAEVSGSLST ERVRELAVALAQKNVKLSTEQLRCLAHRLSEPPEDLDALPLDLLLFLNPDADFSGPQACTRFFFSRI TKANVDLLPRGAPE RQRLPAALACWGVRSLLSEADVRLGGLACDLPGRFVAESAIEVLLPRLVSCPGPLDQDQQAARAALQGGGPPYGGP STWSVTMDALRGLLPVLGQPI IRSIPQGIVAARQRSSRDPSWRQPETILRPRFRREVEKTACPSGKKAREIDESLI FYKKWELEACVDAALLATQMDRVNAIPFTYEQLDVLKHKLDELYPQGYPESVIQHLGYLFLKMSPEDIRKWNVTSLLETL KALLEVNKGHEMSPQVATLIDRFVKGRGQLDKDTLDTLTAFYPGYLCSLSPEELSSVPPSSIWAVRPQDLDTCDPRQLD VLYPKARLAFQNMNGSEYFVKIQSFLGGASVKDLRALSQHNVSMDIATFKRLQVDSLVLGSLVAEVQKLLGPHVEGLKAE ERHRPVRDWILRQRQDDLDLTGLGLQGGINPGYLVLDLSMQEALSGTPCLLGGPGPVLTVLALLLASTLA
251.	hu orl MSLN- E6 mu	artifi- cial	MALPTARPLLGSCGTPALGSLFLFSLGWQPSRTLAGETGQEAAPLDGVLANPPNISSLSRQLLGFPFCAEVSGSLST ERVRELAVALAQKNVKLSTEQLRCLAHRLSEPPEDLDALPLDLLLFLNPDADFSGPQACTRFFFSRI TKANVDLLPRGAPE RQRLPAALACWGVRSLLSEADVRLGGLACDLPGRFVAESAIEVLLPRLVSCPGPLDQDQQAARAALQGGGPPYGGP STWSVTMDALRGLLPVLGQPI IRSIPQGIVAARQRSSRDPSWRQPETILRPRFRREVEKTACPSGKKAREIDESLI FYKKWELEACVDAALLATQMDRVNAIPFTYEQLDVLKHKLDELYPQGYPESVIQHLGYLFLKMSPEDIRKWNVTSLLETL KALLEVNKGHEMSPQVATLIDRFVKGRGQLDKDTLDTLTAFYPGYLCSLSPEELSSVPPSSIWAVRPQDLDTCDPRQLD VLYPKARLAFQNMNGSEYFVKIQSFLGGAPTEDLKALSQQNVSMDLATFMKLRTDAVLP LPTVAEVQKLLGPNIVDLKTE EDKSPVRDWLFRQHQLDRLGLGLQGGINPGYLVLDLNVREAFSGTPCLLGGPGPVLTVLALLLASTLA
252.	Fc monomer-1 +c/-g		DKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNNAKTKPCEEQYGST YRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIA VEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNNHYTKQKSLSPGK
253.	Fc monomer-2 +c/-g/delGK		DKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNNAKTKPCEEQYGST YRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIA VEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNNHYTKQKSLSP
254.	Fc monomer-3 -c/+g		DKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIA VEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNNHYTKQKSLSPGK
255.	Fc monomer-4 -c/+g/delGK		DKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIA VEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNNHYTKQKSLSP
256.	Fc monomer-5 -c/-g		DKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNNAKTKPREEQYGST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIA VEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNNHYTKQKSLSPGK
257.	Fc monomer-6 -c/-g/delGK		DKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNNAKTKPREEQYGST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIA VEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNNHYTKQKSLSP
258.	Fc monomer-7 +c/+g		DKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNNAKTKPCEEQYNST YRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIA VEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNNHYTKQKSLSPGK

TABLE 30-continued

Sequence Listing			
SEQ ID NO:	Description	Source	Sequence
268.	MS_1xCD3-scFc	Bi-specific HLE molecule	QVQLQESGPGLVKPSSETLSLTCTVSGGSISSSSYYWGWIQQPPGKLEWIGSIYYSGITNYPNPSLKSRTVISVDTSKNQ FSLKLSSTVTAADTAVYYCARPSNYDAFDIWGQGTMTVSSGGGGSGGGGSGGGGSSQSVLTQPPSVSGAPGQRVTISCTG SSSNIAGYDVHWYQQVPGTAPKLLIYGN SKRPSGVPDRFSGSKSGTSASLAITGLQAEDEADYYCQSYDSSLGGWVFG GGTKLTVLSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKLEWVARIRSKYNNYATYYADS VKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTTLTVSSGGGGSGGGGSGGGGSSQTVV TQEPSTLTVSPGGTVTLTCSGSTGAVTSGNYPNWWQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTL SGVQPE DEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELLGGPSVFLFPPPKPDTLMI SRTPEVTCVVVDVSHE DPEVKFNWYVDGVEVHNAKTKPCEEQYGYSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPI EKTISKAKGQPREPQV YTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLY SKLTVDKSRWQQGNVFCSCV MHEALHNHYTQKSLSLSPGKGGGSGGGGSGGGGSGGGGSGGGGSGGGGSDKTHTCPPCPAPELLGGPSVFLFPPPKPDTL MI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGYSTYRCVSVLTVLHQDWLNGKEYKCKVSNKAL PAPI EKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLY SKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK
269.	MS_1xCD3-scFc_delGK	Bi-specific HLE molecule	QVQLQESGPGLVKPSSETLSLTCTVSGGSISSSSYYWGWIQQPPGKLEWIGSIYYSGITNYPNPSLKSRTVISVDTSKNQ FSLKLSSTVTAADTAVYYCARPSNYDAFDIWGQGTMTVSSGGGGSGGGGSGGGGSSQSVLTQPPSVSGAPGQRVTISCTG SSSNIAGYDVHWYQQVPGTAPKLLIYGN SKRPSGVPDRFSGSKSGTSASLAITGLQAEDEADYYCQSYDSSLGGWVFG GGTKLTVLSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKLEWVARIRSKYNNYATYYADS VKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTTLTVSSGGGGSGGGGSGGGGSSQTVV TQEPSTLTVSPGGTVTLTCSGSTGAVTSGNYPNWWQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTL SGVQPE DEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELLGGPSVFLFPPPKPDTLMI SRTPEVTCVVVDVSHE DPEVKFNWYVDGVEVHNAKTKPCEEQYGYSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPI EKTISKAKGQPREPQV YTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLY SKLTVDKSRWQQGNVFCSCV MHEALHNHYTQKSLSLSPGPGGGGGGGGGSGGGGSGGGGSGGGGSGGGGSDKTHTCPPCPAPELLGGPSVFLFPPPKPDTL MI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGYSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPA PI EKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLY SK LTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK
270.	MS_1_CCxC D3-scFc	Bi-specific HLE molecule	QVQLQESGPGLVKPSSETLSLTCTVSGGSISSSSYYWGWIQQPPGKLEWIGSIYYSGITNYPNPSLKSRTVISVDTSKNQ FSLKLSSTVTAADTAVYYCARPSNYDAFDIWGQGTMTVSSGGGGSGGGGSGGGGSSQSVLTQPPSVSGAPGQRVTISCTG SSSNIAGYDVHWYQQVPGTAPKLLIYGN SKRPSGVPDRFSGSKSGTSASLAITGLQAEDEADYYCQSYDSSLGGWVFG CGTKLTVLSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKLEWVARIRSKYNNYATYYADS VKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTTLTVSSGGGGSGGGGSGGGGSSQTVV TQEPSTLTVSPGGTVTLTCSGSTGAVTSGNYPNWWQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTL SGVQPE DEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELLGGPSVFLFPPPKPDTLMI SRTPEVTCVVVDVSHE DPEVKFNWYVDGVEVHNAKTKPCEEQYGYSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPI EKTISKAKGQPREPQV YTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLY SKLTVDKSRWQQGNVFCSCV MHEALHNHYTQKSLSLSPGKGGGSGGGGSGGGGSGGGGSGGGGSGGGGSDKTHTCPPCPAPELLGGPSVFLFPPPKPDTL MI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGYSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPA PI EKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLY SKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK
271.	MS_1_CCxC D3-scFc_delGK	Bi-specific HLE molecule	QVQLQESGPGLVKPSSETLSLTCTVSGGSISSSSYYWGWIQQPPGKLEWIGSIYYSGITNYPNPSLKSRTVISVDTSKNQ FSLKLSSTVTAADTAVYYCARPSNYDAFDIWGQGTMTVSSGGGGSGGGGSGGGGSSQSVLTQPPSVSGAPGQRVTISCTG SSSNIAGYDVHWYQQVPGTAPKLLIYGN SKRPSGVPDRFSGSKSGTSASLAITGLQAEDEADYYCQSYDSSLGGWVFG CGTKLTVLSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKLEWVARIRSKYNNYATYYADS VKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTTLTVSSGGGGSGGGGSGGGGSSQTVV TQEPSTLTVSPGGTVTLTCSGSTGAVTSGNYPNWWQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTL SGVQPE DEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELLGGPSVFLFPPPKPDTLMI SRTPEVTCVVVDVSHE DPEVKFNWYVDGVEVHNAKTKPCEEQYGYSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPI EKTISKAKGQPREPQV YTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLY SKLTVDKSRWQQGNVFCSCV MHEALHNHYTQKSLSLSPGPGGGGGGGGGSGGGGSGGGGSGGGGSGGGGSDKTHTCPPCPAPELLGGPSVFLFPPPKPDTL MI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGYSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPA PI EKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLY SK LTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK
272.	MS_2xCD3-scFc	Bi-specific HLE molecule	QVQLVESGGGVQPGRLSLRSCAASGFTFSDHYMSWIRQTPGKLEWVSYISSSGSTIYYADSVKGRFTISRDNAKNSL YLQMNLSRAEDTAVYYCARDLGPSPDYWGQGLVTVSSGGGGSGGGGSGGGGSDIQMTQSPSSVSASVGDRTVITCRAS QGIISSWLAWYQKPKGAPKLLIDAA SRLQSGVPSRFRSGSGSDTFLTITSSLPEDFATYYCQQA NSFPRTFGQGTKE IKSGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKLEWVARIRSKYNNYATYYADSVKDRFT ISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTTLTVSSGGGGSGGGGSGGGGSSQTVV TQEPSTLTVSPGGTVTLTCSGSTGAVTSGNYPNWWQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTL SGVQPE DEAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKTHTCPPCPAPELLGGPSVFLFPPPKPDTLMI SRTPEVTCVVVDVSHE DPEVKFNWYVDGVEVHNAKTKPCEEQYGYSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPI EKTISKAKGQPREPQV YTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLY SKLTVDKSRWQQGNVFCSCV MHEALHNHYTQKSLSLSPGKGGGSGGGGSGGGGSGGGGSGGGGSGGGGSDKTHTCPPCPAPELLGGPSVFLFPPPKPDTL MI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGYSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPA PI EKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLY SK LTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK

Sequence Listing				
SEQ ID NO:	Description	Source	Sequence	
273.	MS_2xCD3-scFc_delGK	Bi-specific HLE molecule	QVQLVSESGGGVVQPGRSRLRLSCAASGFTFS	DHYMSWIRQTPGKGLEWVS
274.	MS_2_CcxC-D3-scFc	Bi-specific HLE molecule	QVQLVSESGGGVVQPGRSRLRLSCAASGFTFS	DHYMSWIRQTPGKCLEWVS
275.	MS_2_CcxC-D3-scFc_delGK	Bi-specific HLE molecule	QVQLVSESGGGVVQPGRSRLRLSCAASGFTFS	DHYMSWIRQTPGKCLEWVS
276.	MS_3xCD3-scFc	Bi-specific HLE molecule	QVQLVSESGGLVKPGGSLRLSCAASGFTFS	DYMTWIRQAPGKGLEWIS
277.	MS_3xCD3-scFc_delGK	Bi-specific HLE molecule	QVQLVSESGGLVKPGGSLRLSCAASGFTFS	DYMTWIRQAPGKGLEWIS

TABLE 30-continued

Sequence Listing			
SEQ ID NO:	Description	Source	Sequence
			NWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFFLYSKLTVDKSRWQQGNVSCSVMHEALH NHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSDKTHHTCPPCPAPELLGGPSVFLFPPPKPDTLMISRTP EVTCTVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFFLYSKLTVDKSRWQQGNVSCSVMHEALH NHYTQKSLSLSPGK
278.	MS_3_CCxC D3-scFc	Bi-speci- fic HLE mole- cule	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDYIMTWRQAPGKCLEWISYISSSGSTIYYADSVKGRFTISRDNAKNSL YLQMNSLRRAEDTAVYYCARDNRSHFDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMTQSPSSVSASVGDRTITCRAS QGITRWLAWYQQKPGKAPKLLIYAASVLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQSNSFPRTFGCGTKVE IKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKLEWVARIRSKYNNYATYYADSVKDRFT ISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGGSGGGGSGTQVVTQEP SLTVSPGGTVTLTSGSSTGAVTSGNYPNWVQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYY CVLWYSNRWVFGGKTLTVLGGGDKTHTCPPCPAPELLGGPSVFLFPPPKPDTLMISRTPETVTCVVDVSHEDPEVKF NWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFFLYSKLTVDKSRWQQGNVSCSVMHEALH NHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSDKTHHTCPPCPAPELLGGPSVFLFPPPKPDTLMISR TPEVTCTVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKT ISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFFLYSKLTVD KSRWQQGNVSCSVMHEALH NHYTQKSLSLSPGK
279.	MS_3_CCxC D3-scFc_delGK	Bi-speci- fic HLE mole- cule	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDYIMTWRQAPGKCLEWISYISSSGSTIYYADSVKGRFTISRDNAKNSL YLQMNSLRRAEDTAVYYCARDNRSHFDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMTQSPSSVSASVGDRTITCRAS QGITRWLAWYQQKPGKAPKLLIYAASVLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQSNSFPRTFGCGTKVE IKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKLEWVARIRSKYNNYATYYADSVKDRFT ISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGGSGGGGSGTQVVTQEP SLTVSPGGTVTLTSGSSTGAVTSGNYPNWVQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYY CVLWYSNRWVFGGKTLTVLGGGDKTHTCPPCPAPELLGGPSVFLFPPPKPDTLMISRTPETVTCVVDVSHEDPEVKF NWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFFLYSKLTVDKSRWQQGNVSCSVMHEALH NHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSDKTHHTCPPCPAPELLGGPSVFLFPPPKPDTLMISRTP EVTCTVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFFLYSKLTVDKSRWQQGNVSCSVMHEALH NHYTQKSLSLSPGK
280.	MS_4xCD3-scFc	Bi-speci- fic HLE mole- cule	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDYIMTWRQAPGKLEWLSYISSSGSTIYYADSVKGRFTISRDNAKNSL FLQMNSLRRAEDTAVYYCARDNRSHFDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMTQSPSSVSASVGDRTITCRAS QGINTWLAWYQQKPGKAPKLLIYGASGLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQAISFPRTFGCGTKVE IKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKLEWVARIRSKYNNYATYYADSVKDRFT ISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGGSGGGGSGTQVVTQEP SLTVSPGGTVTLTSGSSTGAVTSGNYPNWVQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYY CVLWYSNRWVFGGKTLTVLGGGDKTHTCPPCPAPELLGGPSVFLFPPPKPDTLMISRTPETVTCVVDVSHEDPEVKF NWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFFLYSKLTVDKSRWQQGNVSCSVMHEALH NHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSDKTHHTCPPCPAPELLGGPSVFLFPPPKPDTLMISRTP EVTCTVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFFLYSKLTVDKSRWQQGNVSCSVMHEALH NHYTQKSLSLSPGK
281.	MS_4xCD3-scFc_delGK	Bi-speci- fic HLE mole- cule	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDYIMTWRQAPGKLEWLSYISSSGSTIYYADSVKGRFTISRDNAKNSL FLQMNSLRRAEDTAVYYCARDNRSHFDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMTQSPSSVSASVGDRTITCRAS QGINTWLAWYQQKPGKAPKLLIYGASGLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQAISFPRTFGCGTKVE IKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKLEWVARIRSKYNNYATYYADSVKDRFT ISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGGSGGGGSGTQVVTQEP SLTVSPGGTVTLTSGSSTGAVTSGNYPNWVQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYY CVLWYSNRWVFGGKTLTVLGGGDKTHTCPPCPAPELLGGPSVFLFPPPKPDTLMISRTPETVTCVVDVSHEDPEVKF NWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFFLYSKLTVDKSRWQQGNVSCSVMHEALH NHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSDKTHHTCPPCPAPELLGGPSVFLFPPPKPDTLMISRTP EVTCTVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFFLYSKLTVDKSRWQQGNVSCSVMHEALH NHYTQKSLSLSPGK
282.	MS_4_CCxC D3-scFc	Bi-speci- fic HLE mole- cule	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDYIMTWRQAPGKCLEWLSYISSSGSTIYYADSVKGRFTISRDNAKNSL FLQMNSLRRAEDTAVYYCARDNRSHFDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMTQSPSSVSASVGDRTITCRAS QGINTWLAWYQQKPGKAPKLLIYGASGLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQAISFPRTFGCGTKVE IKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKLEWVARIRSKYNNYATYYADSVKDRFT ISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGGSGGGGSGTQVVTQEP SLTVSPGGTVTLTSGSSTGAVTSGNYPNWVQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYY CVLWYSNRWVFGGKTLTVLGGGDKTHTCPPCPAPELLGGPSVFLFPPPKPDTLMISRTPETVTCVVDVSHEDPEVKF NWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFFLYSKLTVDKSRWQQGNVSCSVMHEALH NHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSDKTHHTCPPCPAPELLGGPSVFLFPPPKPDTLMISRTP EVTCTVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFFLYSKLTVDKSRWQQGNVSCSVMHEALH NHYTQKSLSLSPGK

TABLE 30-continued

Sequence Listing			
SEQ ID NO:	Description	Source	Sequence
			REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALH NHYTQKLSLSLSPGKGGGSGGGGSGGGGSGGGGSGGGGSGGGGSDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISR TPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEK TISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVD KSRWQQGNVFCSCVMHEALHNNHTYQKLSLSLSPGK
283.	MS_4_CCxC D3- scFc_delGK	Bi- speci- fic HLE mole- cule	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDYMTWIRQAPGKCLEWFSYISSSGSTIYYADSVKGRFTISRDNAKNSL YLQMNSLRRAEDTAVYYCARDVGSHPFDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMTQSPSSVSASVGDRTITCRAS QGINTWLAWYQQKPGKAPKLLIYGASGLQSGVPSRFSGSGSGTDFTLTISLQPEDFAIYYCQQAISFPRTFGCGTKVE IKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKLEWVARIRSKYNNYATYYADSVKDRFT ISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGGSGGGGSGTQVVTQEP TVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQPEDAEAY CVLWYSNRWVFGGGLKTLTVLGGGDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPETVTCVVVDVSHEDPEVKF NWYVDGVEVHNAKTKPCEEQYGSYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALH NHYTQKLSLSLPGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTP EVTCTVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNNHTYQKLSLSLSPGK
284.	MS_5xCD3- scFc	Bi- speci- fic HLE mole- cule	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDHYMSWIRQAPGKLEWFSYISSSGGIYYADSVKGRFTISRDNAKNSL YLQMNSLRRAEDTAVYYCARDVGSHPFDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMTQSPSSVSASVGDRTITCRAS QDISRWLAWYQQKPGKAPKLLISAASRLQSGVPSRFSGSGSGTDFTLTISLQPEDFAIYYCQQAISFPRTFGCGTKVE IKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKLEWVARIRSKYNNYATYYADSVKDRFT ISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGGSGGGGSGTQVVTQEP TVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQPEDAEAY CVLWYSNRWVFGGGLKTLTVLGGGDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPETVTCVVVDVSHEDPEVKF NWYVDGVEVHNAKTKPCEEQYGSYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALH NHYTQKLSLSLPGKGGGSGGGGSGGGGSGGGGSGGGGSGGGGSDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISR TPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEK TISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVD KSRWQQGNVFCSCVMHEALHNNHTYQKLSLSLSPGK
285.	MS_5xCD3- scFc_delGK	Bi- speci- fic HLE mole- cule	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDHYMSWIRQAPGKLEWFSYISSSGGIYYADSVKGRFTISRDNAKNSL YLQMNSLRRAEDTAVYYCARDVGSHPFDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMTQSPSSVSASVGDRTITCRAS QDISRWLAWYQQKPGKAPKLLISAASRLQSGVPSRFSGSGSGTDFTLTISLQPEDFAIYYCQQAISFPRTFGCGTKVE IKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKLEWVARIRSKYNNYATYYADSVKDRFT ISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGGSGGGGSGTQVVTQEP TVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQPEDAEAY CVLWYSNRWVFGGGLKTLTVLGGGDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPETVTCVVVDVSHEDPEVKF NWYVDGVEVHNAKTKPCEEQYGSYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALH NHYTQKLSLSLPGKGGGSGGGGSGGGGSGGGGSGGGGSGGGGSDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISR TPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEK TISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVD KSRWQQGNVFCSCVMHEALHNNHTYQKLSLSLSPGK
286.	MS_5_CCxC D3-scFc	Bi- speci- fic HLE mole- cule	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDHYMSWIRQAPGKLEWFSYISSSGGIYYADSVKGRFTISRDNAKNSL YLQMNSLRRAEDTAVYYCARDVGSHPFDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMTQSPSSVSASVGDRTITCRAS QDISRWLAWYQQKPGKAPKLLISAASRLQSGVPSRFSGSGSGTDFTLTISLQPEDFAIYYCQQAISFPRTFGCGTKVE IKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKLEWVARIRSKYNNYATYYADSVKDRFT ISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGGSGGGGSGTQVVTQEP TVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQPEDAEAY CVLWYSNRWVFGGGLKTLTVLGGGDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPETVTCVVVDVSHEDPEVKF NWYVDGVEVHNAKTKPCEEQYGSYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALH NHYTQKLSLSLPGKGGGSGGGGSGGGGSGGGGSGGGGSGGGGSDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISR TPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEK TISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVD KSRWQQGNVFCSCVMHEALHNNHTYQKLSLSLSPGK
287.	MS_5_CCxC D3- scFc_delGK	Bi- speci- fic HLE mole- cule	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDHYMSWIRQAPGKLEWFSYISSSGGIYYADSVKGRFTISRDNAKNSL YLQMNSLRRAEDTAVYYCARDVGSHPFDYWGQGLTVTVSSGGGGSGGGGSGGGGSDIQMTQSPSSVSASVGDRTITCRAS QDISRWLAWYQQKPGKAPKLLISAASRLQSGVPSRFSGSGSGTDFTLTISLQPEDFAIYYCQQAISFPRTFGCGTKVE IKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKLEWVARIRSKYNNYATYYADSVKDRFT ISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGGSGGGGSGTQVVTQEP TVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQPEDAEAY CVLWYSNRWVFGGGLKTLTVLGGGDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPETVTCVVVDVSHEDPEVKF NWYVDGVEVHNAKTKPCEEQYGSYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALH

TABLE 30-continued

Sequence Listing			
SEQ ID NO:	Description	Source	Sequence
			NHYTQKSLSLSPGGGGSGGGSGGGSGGGSGGGSGGGSGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTP EVTCTVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSLKTVDKS RWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
288.	MS_6xCD3-scFc	Bi-specific HLE molecule	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDHYMSWIRQAPGKGLEWISYISNSGSIYYVDSVKGRFTISRDNAKNSL YLQMNSLRAEDTAVYYCARDVRTAFDYWGQGLTVTVSSGGGGSGGGSGGGSGGGSDIQMTQSPSSVSASVGDRTITCRAS QSIGSWLAWYQQKPGKAPNLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQANSFPRTFGCGTKVD IKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFT ISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGSGGGSGGGQTVVTQEP TVSPGGTVTLTTCGSSGTGAVTSGNYPNWVQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLGSGVQPEDEAEY CVLWYSNRWVFGGKTTLVLGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPTEVTCVVVDVSHEDPEVKF NWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSLKTVDKS RWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
289.	MS_6xCD3-scFc_delGK	Bi-specific HLE molecule	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDHYMSWIRQAPGKGLEWISYISNSGSIYYVDSVKGRFTISRDNAKNSL YLQMNSLRAEDTAVYYCARDVRTAFDYWGQGLTVTVSSGGGGSGGGSGGGSGGGSDIQMTQSPSSVSASVGDRTITCRAS QSIGSWLAWYQQKPGKAPNLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQANSFPRTFGCGTKVD IKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFT ISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGSGGGSGGGQTVVTQEP TVSPGGTVTLTTCGSSGTGAVTSGNYPNWVQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLGSGVQPEDEAEY CVLWYSNRWVFGGKTTLVLGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPTEVTCVVVDVSHEDPEVKF NWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSLKTVDKS RWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
290.	MS_6_CCxC D3-scFc	Bi-specific HLE molecule	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDHYMSWIRQAPGKCLEWISYISNSGSIYYVDSVKGRFTISRDNAKNSL YLQMNSLRAEDTAVYYCARDVRTAFDYWGQGLTVTVSSGGGGSGGGSGGGSGGGSDIQMTQSPSSVSASVGDRTITCRAS QSIGSWLAWYQQKPGKAPNLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQANSFPRTFGCGTKVD IKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFT ISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGSGGGSGGGQTVVTQEP TVSPGGTVTLTTCGSSGTGAVTSGNYPNWVQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLGSGVQPEDEAEY CVLWYSNRWVFGGKTTLVLGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPTEVTCVVVDVSHEDPEVKF NWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSLKTVDKS RWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
291.	MS_6_CCxC D3-sc_Fc_delGK	Bi-specific HLE molecule	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDHYMSWIRQAPGKCLEWISYISNSGSIYYVDSVKGRFTISRDNAKNSL YLQMNSLRAEDTAVYYCARDVRTAFDYWGQGLTVTVSSGGGGSGGGSGGGSGGGSDIQMTQSPSSVSASVGDRTITCRAS QSIGSWLAWYQQKPGKAPNLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQANSFPRTFGCGTKVD IKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFT ISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGSGGGSGGGQTVVTQEP TVSPGGTVTLTTCGSSGTGAVTSGNYPNWVQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLGSGVQPEDEAEY CVLWYSNRWVFGGKTTLVLGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPTEVTCVVVDVSHEDPEVKF NWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSLKTVDKS RWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
292.	MS_7xCD3-scFc	Bi-specific HLE molecule	EVQLVESGGGLIQPGGSLRLSCAVSGFTVSSKFTWVRQAPGKGLEWVSVIYSGGKTYADSVKGRFTISRDNKNTLY LQMNSLRAEDTAVYYCARDSSGGWGYFDYWGQGLTVTVSSGGGGSGGGSGGGSGGGSDIVMTQSPDSLAVSLGERATINCK SQSVLYSNNKNYLAWYQQKPGQPKLLIYASTRESGVPRDFSGSGSGTDFTLTISLQPEDVAVYYCQQYVYTPPTF GQGTKEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYAD SVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGSGGGSGGGQTV VTQEPSTVSPGGTVTLTTCGSSGTGAVTSGNYPNWVQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLGSGVQ EDEAEYCVLWYSNRWVFGGKTTLVLGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPTEVTCVVVDVSH EDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSLKTVDKS RWQQGNVFSCSVMHEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPK

TABLE 30-continued

Sequence Listing			
SEQ ID NO:	Description	Source	Sequence
			DTLMSIRTPTEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAAKTKPCEEQYGSYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNNHTQKSLSLSPGK
293.	MS_7xCD3-scFc_delGK	Bi-specific HLE molecule	EVQLVESGGGLIQPGGSLRLSCAASGFTVSSKFMTWVRQAPGKGLEWVSVIYSGGKITYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARDSSGGWGYFDYWGQGTTLVTVSSGGGGSGGGSGGGGSDIVMTQSPDSLAVSLGERATINCKSSQSVLYSSNNKNYLAWYQQKPGQPPKLLIYWASTRESGVDRFSGSGSGTDFTLTISSLQAEDVAVYYCQQYYSTPPTFGQGTKEIKSGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTTLVTVSSGGGGSGGGSGGGGSGTQTVVTQEPSTLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLGGAALTLGSGVQPEDEAEYYCVLWYSNRWVFGGKTKLTVLGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAAKTKPCEEQYGSYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNNHTQKSLSLSPGK
294.	MS_7_CCxC D3-scFc	Bi-specific HLE molecule	EVQLVESGGGLIQPGGSLRLSCAASGFTVSSKFMTWVRQAPGKCLEWVSVIYSGGKITYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARDSSGGWGYFDYWGQGTTLVTVSSGGGGSGGGSGGGGSDIVMTQSPDSLAVSLGERATINCKSSQSVLYSSNNKNYLAWYQQKPGQPPKLLIYWASTRESGVDRFSGSGSGTDFTLTISSLQAEDVAVYYCQQYYSTPPTFGCGTKVEIKSGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTTLVTVSSGGGGSGGGSGGGGSGTQTVVTQEPSTLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLGGAALTLGSGVQPEDEAEYYCVLWYSNRWVFGGKTKLTVLGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAAKTKPCEEQYGSYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNNHTQKSLSLSPGK
295.	MS_7_CCxC D3-scFc_delGK	Bi-specific HLE molecule	EVQLVESGGGLIQPGGSLRLSCAASGFTVSSKFMTWVRQAPGKCLEWVSVIYSGGKITYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARDSSGGWGYFDYWGQGTTLVTVSSGGGGSGGGSGGGGSDIVMTQSPDSLAVSLGERATINCKSSQSVLYSSNNKNYLAWYQQKPGQPPKLLIYWASTRESGVDRFSGSGSGTDFTLTISSLQAEDVAVYYCQQYYSTPPTFGCGTKVEIKSGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTTLVTVSSGGGGSGGGSGGGGSGTQTVVTQEPSTLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLGGAALTLGSGVQPEDEAEYYCVLWYSNRWVFGGKTKLTVLGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAAKTKPCEEQYGSYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNNHTQKSLSLSPGK
296.	MS_8xCD3-scFc	Bi-specific HLE molecule	QVQLQESGPGLVKPSETLSLTCTVSGGSISSYYNWIRQPPGKLEWIGRIYYNGNTYYNP SLKSRVTISGDTSKNQFS LKLSSVTAADTAVYYCARPKLGIDAFDIWGQGTMTVTVSSGGGGSGGGSGGGGSGVSLTQPPSVSGAPGQRVTISCTGSSNIGAGYDVHWYQQLPGTAPKLLIYGNNSNRPSGVDRFSGSKSGTSASLAI TGLQAED EADYYCQSYDSLSGWFVGGGTLKLTVLSSGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTTLVTVSSGGGGSGGGSGGGGSGTQTVVTQEPSTLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLGGAALTLGSGVQPEDEAEYYCVLWYSNRWVFGGKTKLTVLGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAAKTKPCEEQYGSYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNNHTQKSLSLSPGK
297.	MS_8xCD3-scFc_delGK	Bi-specific HLE molecule	QVQLQESGPGLVKPSETLSLTCTVSGGSISSYYNWIRQPPGKLEWIGRIYYNGNTYYNP SLKSRVTISGDTSKNQFS LKLSSVTAADTAVYYCARPKLGIDAFDIWGQGTMTVTVSSGGGGSGGGSGGGGSGVSLTQPPSVSGAPGQRVTISCTGSSNIGAGYDVHWYQQLPGTAPKLLIYGNNSNRPSGVDRFSGSKSGTSASLAI TGLQAED EADYYCQSYDSLSGWFVGGGTLKLTVLSSGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTTLVTVSSGGGGSGGGSGGGGSGTQTVVTQEPSTLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLGGAALTLGSGVQPEDEAEYYCVLWYSNRWVFGGKTKLTVLGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHED

TABLE 30-continued

Sequence Listing			
SEQ ID NO:	Description	Source	Sequence
			PEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSSVMHEALHNNHYTQKLSLSLSPGK
298.	MS_8_CCxC D3-scFc	Bi-specific HLE molecule	QVQLQESGPGLVKPSSETLSLTCTVSGGSISSYYWNWIRQPPGKCLEWIGRIYYNGNTYYNP SLKSRVTISGDTSKNQFS LKLSSTVAADTAVYYCARPKLGIDAFDIWGQGTMTVTVSSGGGSGGGSGGGSGGGSGQSVLTQPPSVSGAPGQRVTISCTGSSSNIGAGYDVHVVYQQLPGTAPKLLIYGNISNRPSGVDRFSGSGSGTASLAITGLQAEDEADYYCQSYDSSLGQVFGCGTKLTVLSSGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGSGGGSGGGSGQSVTVTQEPSTLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKLAPGTPARFSGSLGGKAALTLGSGVQPEDAEYYCVLWYSNRWVFGGTKLTVLGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSSVMHEALHNNHYTQKLSLSLSPGK
299.	MS_8_CCxC D3-scFc_delGK	Bi-specific HLE molecule	QVQLQESGPGLVKPSSETLSLTCTVSGGSISSYYWNWIRQPPGKCLEWIGRIYYNGNTYYNP SLKSRVTISGDTSKNQFS LKLSSTVAADTAVYYCARPKLGIDAFDIWGQGTMTVTVSSGGGSGGGSGGGSGGGSGQSVLTQPPSVSGAPGQRVTISCTGSSSNIGAGYDVHVVYQQLPGTAPKLLIYGNISNRPSGVDRFSGSGSGTASLAITGLQAEDEADYYCQSYDSSLGQVFGCGTKLTVLSSGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGSGGGSGGGSGQSVTVTQEPSTLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKLAPGTPARFSGSLGGKAALTLGSGVQPEDAEYYCVLWYSNRWVFGGTKLTVLGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSSVMHEALHNNHYTQKLSLSLSPGK
300.	Peptide linker	(G ₄ S) ₄ linker	GGGGSGGGSGGGSGGGSGGGGS
301.	Peptide linker	(G ₄ S) ₅ linker	GGGGSGGGSGGGSGGGSGGGSGGGGS
302.	Peptide linker	(G ₄ S) ₆ linker	GGGGSGGGSGGGSGGGSGGGSGGGSGGGGS
303.	Peptide linker	(G ₄ S) ₇ linker	GGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGGS
304.	Peptide linker	(G ₄ S) ₈ linker	GGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGGS

SEQUENCE LISTING

```
Sequence total quantity: 327
SEQ ID NO: 1          moltype = AA length = 4
FEATURE               Location/Qualifiers
REGION               1..4
                     note = Synthetic peptide
source               1..4
                     mol_type = protein
                     organism = synthetic construct
```

4

```
SEQ ID NO: 2          moltype = AA  length = 5
FEATURE              Location/Qualifiers
REGION               1..5
```

-continued

source	note = Synthetic peptide 1..5 mol_type = protein organism = synthetic construct	
SEQUENCE: 2 GGGGS		5
SEQ ID NO: 3 FEATURE REGION	moltype = AA length = 5 Location/Qualifiers 1..5 note = Synthetic peptide	
source	1..5 mol_type = protein organism = synthetic construct	
SEQUENCE: 3 GGGGQ		5
SEQ ID NO: 4 FEATURE REGION	moltype = AA length = 6 Location/Qualifiers 1..6 note = Synthetic peptide	
source	1..6 mol_type = protein organism = synthetic construct	
SEQUENCE: 4 PGGGGS		6
SEQ ID NO: 5 FEATURE REGION	moltype = AA length = 6 Location/Qualifiers 1..6 note = Synthetic peptide	
source	1..6 mol_type = protein organism = synthetic construct	
SEQUENCE: 5 PGGDGS		6
SEQ ID NO: 6 FEATURE REGION	moltype = AA length = 6 Location/Qualifiers 1..6 note = Synthetic peptide	
source	1..6 mol_type = protein organism = synthetic construct	
SEQUENCE: 6 SGGGGS		6
SEQ ID NO: 7 FEATURE REGION	moltype = AA length = 9 Location/Qualifiers 1..9 note = Synthetic peptide	
source	1..9 mol_type = protein organism = synthetic construct	
SEQUENCE: 7 GGGSGGGGS		9
SEQ ID NO: 8 FEATURE REGION	moltype = AA length = 10 Location/Qualifiers 1..10 note = Synthetic peptide	
source	1..10 mol_type = protein organism = synthetic construct	
SEQUENCE: 8 GGGSGGGGS		10
SEQ ID NO: 9 FEATURE REGION	moltype = AA length = 15 Location/Qualifiers 1..15 note = Synthetic peptide	
source	1..15 mol_type = protein organism = synthetic construct	
SEQUENCE: 9 GGGSGGGGS GGGGS		15

-continued

SEQ ID NO: 10	moltype = AA length = 6	
FEATURE	Location/Qualifiers	
REGION	1..6	
source	note = Synthetic peptide	
	1..6	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 10		
HHHHHH		6
SEQ ID NO: 11	moltype = AA length = 14	
FEATURE	Location/Qualifiers	
REGION	1..14	
source	note = Synthetic peptide	
	1..14	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 11		
GSSTGAVTSG YYPN		14
SEQ ID NO: 12	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
source	note = Synthetic peptide	
	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 12		
GTKFLAP		7
SEQ ID NO: 13	moltype = AA length = 9	
FEATURE	Location/Qualifiers	
REGION	1..9	
source	note = Synthetic peptide	
	1..9	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 13		
ALWYSNRWV		9
SEQ ID NO: 14	moltype = AA length = 5	
FEATURE	Location/Qualifiers	
REGION	1..5	
source	note = Synthetic peptide	
	1..5	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 14		
IYAMN		5
SEQ ID NO: 15	moltype = AA length = 19	
FEATURE	Location/Qualifiers	
REGION	1..19	
source	note = Synthetic peptide	
	1..19	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 15		
RIRSKYNNYA TYYADSVKS		19
SEQ ID NO: 16	moltype = AA length = 14	
FEATURE	Location/Qualifiers	
REGION	1..14	
source	note = Synthetic peptide	
	1..14	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 16		
HGNFGNSYVS FFAY		14
SEQ ID NO: 17	moltype = AA length = 125	
FEATURE	Location/Qualifiers	
REGION	1..125	
source	note = Synthetic peptide	
	1..125	

-continued

mol_type = protein
organism = synthetic construct

SEQUENCE: 17
EVQLVESGGG LVQPGGSLKL SCAASGFTFN IYAMNWRQA PGKGLEWVAR IRSKYNNYAT 60
YYADSVKSRF TISRDDSKNT AYLQMNLLKT EDTAVYYCVR HGNFGNSYVS FFAYWGQGTL 120
VTVSS 125

SEQ ID NO: 18 moltype = AA length = 109
FEATURE Location/Qualifiers
REGION 1..109
note = Synthetic peptide
source 1..109
mol_type = protein
organism = synthetic construct

SEQUENCE: 18
QTVVTQEPSL TVSPGGTVTL TCGSSTGAVT SGYYPNWVQQ KPGQAPRGLI GGTKFLAPGT 60
PARFSGSLLG GKAALTLGSG QPEDEAEYYC ALWYSNRWVF GGGTKLTVL 109

SEQ ID NO: 19 moltype = AA length = 249
FEATURE Location/Qualifiers
REGION 1..249
note = Synthetic peptide
source 1..249
mol_type = protein
organism = synthetic construct

SEQUENCE: 19
EVQLVESGGG LVQPGGSLKL SCAASGFTFN IYAMNWRQA PGKGLEWVAR IRSKYNNYAT 60
YYADSVKSRF TISRDDSKNT AYLQMNLLKT EDTAVYYCVR HGNFGNSYVS FFAYWGQGTL 120
VTVSSGGGGS GGGGSGGGGS QTVVTQEPSL TVSPGGTVTL TCGSSTGAVT SGYYPNWVQQ 180
KPGQAPRGLI GGTKFLAPGT PARFSGSLLG GKAALTLGSG QPEDEAEYYC ALWYSNRWVF 240
GGGTKLTVL 249

SEQ ID NO: 20 moltype = AA length = 14
FEATURE Location/Qualifiers
REGION 1..14
note = Synthetic peptide
source 1..14
mol_type = protein
organism = synthetic construct

SEQUENCE: 20
GSSTGAVTSG YYPN 14

SEQ ID NO: 21 moltype = AA length = 7
FEATURE Location/Qualifiers
REGION 1..7
note = Synthetic peptide
source 1..7
mol_type = protein
organism = synthetic construct

SEQUENCE: 21
GTKFLAP 7

SEQ ID NO: 22 moltype = AA length = 9
FEATURE Location/Qualifiers
REGION 1..9
note = Synthetic peptide
source 1..9
mol_type = protein
organism = synthetic construct

SEQUENCE: 22
ALWYSNRWV 9

SEQ ID NO: 23 moltype = AA length = 5
FEATURE Location/Qualifiers
REGION 1..5
note = Synthetic peptide
source 1..5
mol_type = protein
organism = synthetic construct

SEQUENCE: 23
KYAMN 5

SEQ ID NO: 24 moltype = AA length = 19
FEATURE Location/Qualifiers
REGION 1..19
note = Synthetic peptide

-continued

source 1..19
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 24
 RIRSKYNNYA TYYADSVKD 19

SEQ ID NO: 25 moltype = AA length = 14
 FEATURE Location/Qualifiers
 REGION 1..14
 note = Synthetic peptide

source 1..14
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 25
 HGNFGNSYIS YWAY 14

SEQ ID NO: 26 moltype = AA length = 125
 FEATURE Location/Qualifiers
 REGION 1..125
 note = Synthetic peptide

source 1..125
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 26
 EVQLVESGGG LVQPGGSLKL SCAASGFTFN KYAMNWVRQA PGKGLEWVAR IRSKYNNYAT 60
 YYADSVKDRF TISRDDSKNT AYLQMNLLKT EDTAVYYCVR HGNFGNSYIS YWAYWGQGT 120
 VTSS 125

SEQ ID NO: 27 moltype = AA length = 109
 FEATURE Location/Qualifiers
 REGION 1..109
 note = Synthetic peptide

source 1..109
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 27
 QTVVTQEPSL TVSPGGTVTL TCGSSTGAVT SGYYPNWVQQ KPGQAPRGLI GGTKFLAPGT 60
 PARFSGSLLG GKAALTLSGV QPEDEAEYYC ALWYSNRWVF GGGTKLTVL 109

SEQ ID NO: 28 moltype = AA length = 249
 FEATURE Location/Qualifiers
 REGION 1..249
 note = Synthetic peptide

source 1..249
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 28
 EVQLVESGGG LVQPGGSLKL SCAASGFTFN KYAMNWVRQA PGKGLEWVAR IRSKYNNYAT 60
 YYADSVKDRF TISRDDSKNT AYLQMNLLKT EDTAVYYCVR HGNFGNSYIS YWAYWGQGT 120
 VTSSGGGGS GGGGSGGGS QTVVTQEPSL TVSPGGTVTL TCGSSTGAVT SGYYPNWVQQ 180
 KPGQAPRGLI GGTKFLAPGT PARFSGSLLG GKAALTLSGV QPEDEAEYYC ALWYSNRWVF 240
 GGGTKLTVL 249

SEQ ID NO: 29 moltype = AA length = 14
 FEATURE Location/Qualifiers
 REGION 1..14
 note = Synthetic peptide

source 1..14
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 29
 GSSTGAVTSG YYPN 14

SEQ ID NO: 30 moltype = AA length = 7
 FEATURE Location/Qualifiers
 REGION 1..7
 note = Synthetic peptide

source 1..7
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 30
 GTKFLAP 7

SEQ ID NO: 31 moltype = AA length = 9
 FEATURE Location/Qualifiers
 REGION 1..9

-continued

source	note = Synthetic peptide 1..9 mol_type = protein organism = synthetic construct	
SEQUENCE: 31 ALWYSNRWV		9
SEQ ID NO: 32 FEATURE REGION	moltype = AA length = 5 Location/Qualifiers 1..5 note = Synthetic peptide	
source	1..5 mol_type = protein organism = synthetic construct	
SEQUENCE: 32 SYAMN		5
SEQ ID NO: 33 FEATURE REGION	moltype = AA length = 19 Location/Qualifiers 1..19 note = Synthetic peptide	
source	1..19 mol_type = protein organism = synthetic construct	
SEQUENCE: 33 RIRSKYNNYA TYYADSVKG		19
SEQ ID NO: 34 FEATURE REGION	moltype = AA length = 14 Location/Qualifiers 1..14 note = Synthetic peptide	
source	1..14 mol_type = protein organism = synthetic construct	
SEQUENCE: 34 HGNFGNSYLS FWAY		14
SEQ ID NO: 35 FEATURE REGION	moltype = AA length = 125 Location/Qualifiers 1..125 note = Synthetic peptide	
source	1..125 mol_type = protein organism = synthetic construct	
SEQUENCE: 35 EVQLVESGGG LEQPGGSLKL SCAASGFTFN SYAMNWVRQA PGKGLEWVAR IRSKYNNYAT YYADSVKGRF TISRDDSKNT AYLQMNLLKT EDTAVYYCVR HGNFGNSYLS FWAYWGQGTL VTSS		60 120 125
SEQ ID NO: 36 FEATURE REGION	moltype = AA length = 109 Location/Qualifiers 1..109 note = Synthetic peptide	
source	1..109 mol_type = protein organism = synthetic construct	
SEQUENCE: 36 QTVVTQEPSL TVSPGGTVTL TCGSSTGAVT SGYYPNWVQQ KPGQAPRGLI GGTKFLAPGT PARFSGSLLG GKAALTLGTV QPEDEAEYYC ALWYSNRWVF GGGTKLTVL		60 109
SEQ ID NO: 37 FEATURE REGION	moltype = AA length = 249 Location/Qualifiers 1..249 note = Synthetic peptide	
source	1..249 mol_type = protein organism = synthetic construct	
SEQUENCE: 37 EVQLVESGGG LEQPGGSLKL SCAASGFTFN SYAMNWVRQA PGKGLEWVAR IRSKYNNYAT YYADSVKGRF TISRDDSKNT AYLQMNLLKT EDTAVYYCVR HGNFGNSYLS FWAYWGQGTL VTSSGGGGS GGGGSGGGGS QTVVTQEPSL TVSPGGTVTL TCGSSTGAVT SGYYPNWVQQ KPGQAPRGLI GGTKFLAPGT PARFSGSLLG GKAALTLGTV QPEDEAEYYC ALWYSNRWVF GGGKLTVL		60 120 180 240 249
SEQ ID NO: 38 FEATURE	moltype = AA length = 14 Location/Qualifiers	

-continued

REGION	1..14	
source	note = Synthetic peptide	
	1..14	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 38		
GSSTGAVTSG YYPN		14
SEQ ID NO: 39	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
	note = Synthetic peptide	
source	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 39		
GTKFLAP		7
SEQ ID NO: 40	moltype = AA length = 9	
FEATURE	Location/Qualifiers	
REGION	1..9	
	note = Synthetic peptide	
source	1..9	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 40		
ALWYSNRWV		9
SEQ ID NO: 41	moltype = AA length = 5	
FEATURE	Location/Qualifiers	
REGION	1..5	
	note = Synthetic peptide	
source	1..5	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 41		
RYAMN		5
SEQ ID NO: 42	moltype = AA length = 19	
FEATURE	Location/Qualifiers	
REGION	1..19	
	note = Synthetic peptide	
source	1..19	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 42		
RIRSKYNNYA TYYADSVKG		19
SEQ ID NO: 43	moltype = AA length = 14	
FEATURE	Location/Qualifiers	
REGION	1..14	
	note = Synthetic peptide	
source	1..14	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 43		
HGNFGNSYLS YFAY		14
SEQ ID NO: 44	moltype = AA length = 125	
FEATURE	Location/Qualifiers	
REGION	1..125	
	note = Synthetic peptide	
source	1..125	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 44		
EVQLVESGGG LVQPGGSLKL SCAASGFTFN RYAMNWRQA PGKGLEWVAR IRSKYNNYAT		60
YYADSVKGRF TISRDDSKNT AYLQMNNLKT EDTAVYYCVR HGNFGNSYLS YFAYWGQGTL		120
VTSS		125
SEQ ID NO: 45	moltype = AA length = 109	
FEATURE	Location/Qualifiers	
REGION	1..109	
	note = Synthetic peptide	
source	1..109	
	mol_type = protein	

-continued

```

                                organism = synthetic construct
SEQUENCE: 45
QTVVTQEPSL TVSPGGTVTL TCGSSTGAVT SGYYPNWVQQ KPGQAPRGLI GGTKFLAPGT 60
PARFSGSLLG GKAALTLGV QPEDEAEYYC ALWYSNRWVF GGGTKLTVL 109

SEQ ID NO: 46      moltype = AA length = 249
FEATURE           Location/Qualifiers
REGION            1..249
                  note = Synthetic peptide
source            1..249
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 46
EVQLVESGGG LVQPGGSLKL SCAASGFTFN RYAMNWRQA PGKGLEWVAR IRSKYNNYAT 60
YYADSVKGRF TISRDDSKNT AYLQMNLLKT EDTAVYYCVR HGNFGNSYLS YFAYWGQGTL 120
VTVSSGGGGS GGGGSGGGGS QTVVTQEPSL TVSPGGTVTL TCGSSTGAVT SGYYPNWVQQ 180
KPGQAPRGLI GGTKFLAPGT PARFSGSLLG GKAALTLGV QPEDEAEYYC ALWYSNRWVF 240
GGGTKLTVL 249

SEQ ID NO: 47      moltype = AA length = 14
FEATURE           Location/Qualifiers
REGION            1..14
                  note = Synthetic peptide
source            1..14
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 47
RSTGAVTSG YYPN 14

SEQ ID NO: 48      moltype = AA length = 7
FEATURE           Location/Qualifiers
REGION            1..7
                  note = Synthetic peptide
source            1..7
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 48
ATDMRPS 7

SEQ ID NO: 49      moltype = AA length = 9
FEATURE           Location/Qualifiers
REGION            1..9
                  note = Synthetic peptide
source            1..9
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 49
ALWYSNRWV 9

SEQ ID NO: 50      moltype = AA length = 5
FEATURE           Location/Qualifiers
REGION            1..5
                  note = Synthetic peptide
source            1..5
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 50
VYAMN 5

SEQ ID NO: 51      moltype = AA length = 19
FEATURE           Location/Qualifiers
REGION            1..19
                  note = Synthetic peptide
source            1..19
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 51
RIRSKYNNYA TYYADSVKK 19

SEQ ID NO: 52      moltype = AA length = 14
FEATURE           Location/Qualifiers
REGION            1..14
                  note = Synthetic peptide
source            1..14
                  mol_type = protein
                  organism = synthetic construct

```

-continued

SEQUENCE: 52
HGNFGNSYLS WWAY 14

SEQ ID NO: 53 moltype = AA length = 125
FEATURE Location/Qualifiers
REGION 1..125
note = Synthetic peptide
source 1..125
mol_type = protein
organism = synthetic construct

SEQUENCE: 53
EVQLVESGGG LVQPGGSLKL SCAASGFTFN VYAMNWRQA PGKGLEWVAR IRSKYNNYAT 60
YYADSVKKRF TISRDDSKNT AYLQMNLLKT EDTAVYYCVR HGNFGNSYLS WWAYWGQGTL 120
VTSS 125

SEQ ID NO: 54 moltype = AA length = 109
FEATURE Location/Qualifiers
REGION 1..109
note = Synthetic peptide
source 1..109
mol_type = protein
organism = synthetic construct

SEQUENCE: 54
QTVVTQEPSL TVSPGGTVTL TCRSSTGAVT SGYYPNWVQQ KPGQAPRGLI GATDMRPSGT 60
PARFSGSLLG GKAALTLGGV QPEDEAEYYC ALWYSNRWVF GGTGKLTVL 109

SEQ ID NO: 55 moltype = AA length = 249
FEATURE Location/Qualifiers
REGION 1..249
note = Synthetic peptide
source 1..249
mol_type = protein
organism = synthetic construct

SEQUENCE: 55
EVQLVESGGG LVQPGGSLKL SCAASGFTFN VYAMNWRQA PGKGLEWVAR IRSKYNNYAT 60
YYADSVKKRF TISRDDSKNT AYLQMNLLKT EDTAVYYCVR HGNFGNSYLS WWAYWGQGTL 120
VTVSSGGGGS GGGGSGGGGS QTVVTQEPSL TVSPGGTVTL TCRSSTGAVT SGYYPNWVQQ 180
KPGQAPRGLI GATDMRPSGT PARFSGSLLG GKAALTLGGV QPEDEAEYYC ALWYSNRWVF 240
GGTGKLTVL 249

SEQ ID NO: 56 moltype = AA length = 14
FEATURE Location/Qualifiers
REGION 1..14
note = Synthetic peptide
source 1..14
mol_type = protein
organism = synthetic construct

SEQUENCE: 56
GSTGAVTSG YYPN 14

SEQ ID NO: 57 moltype = AA length = 7
FEATURE Location/Qualifiers
REGION 1..7
note = Synthetic peptide
source 1..7
mol_type = protein
organism = synthetic construct

SEQUENCE: 57
GTKFLAP 7

SEQ ID NO: 58 moltype = AA length = 9
FEATURE Location/Qualifiers
REGION 1..9
note = Synthetic peptide
source 1..9
mol_type = protein
organism = synthetic construct

SEQUENCE: 58
ALWYSNRWV 9

SEQ ID NO: 59 moltype = AA length = 5
FEATURE Location/Qualifiers
REGION 1..5
note = Synthetic peptide
source 1..5
mol_type = protein

-continued

organism = synthetic construct
 SEQUENCE: 59
 KYAMN 5

SEQ ID NO: 60
 FEATURE Location/Qualifiers
 REGION 1..19
 note = Synthetic peptide
 source 1..19
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 60
 RIRSKYNNYA TYYADSVKS 19

SEQ ID NO: 61
 FEATURE Location/Qualifiers
 REGION 1..14
 note = Synthetic peptide
 source 1..14
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 61
 HGNFGNSYTS YYAY 14

SEQ ID NO: 62
 FEATURE Location/Qualifiers
 REGION 1..125
 note = Synthetic peptide
 source 1..125
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 62
 EVQLVESGGG LVQPGGSLKL SCAASGFTFN KYAMNWRQA PGKGLEWVAR IRSKYNNYAT 60
 YYADSVKSRF TISRDDSKNT AYLMNNLKT EDTAVYYCVR HGNFGNSYTS YYAYWGQGTL 120
 VTVSS 125

SEQ ID NO: 63
 FEATURE Location/Qualifiers
 REGION 1..109
 note = Synthetic peptide
 source 1..109
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 63
 QTVVTQEPSL TVSPGGTVTL TCGSSTGAVT SGYYPNWVQQ KPGQAPRGLI GGTKFLAPGT 60
 PARFSGSLLG GKAALTLSGV QPEDEAEYYC ALWYSNRWVF GGGTKLTVL 109

SEQ ID NO: 64
 FEATURE Location/Qualifiers
 REGION 1..249
 note = Synthetic peptide
 source 1..249
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 64
 EVQLVESGGG LVQPGGSLKL SCAASGFTFN KYAMNWRQA PGKGLEWVAR IRSKYNNYAT 60
 YYADSVKSRF TISRDDSKNT AYLMNNLKT EDTAVYYCVR HGNFGNSYTS YYAYWGQGTL 120
 VTVSSGGGGS GGGGSGGGGS QTVVTQEPSL TVSPGGTVTL TCGSSTGAVT SGYYPNWVQQ 180
 KPGQAPRGLI GGTKFLAPGT PARFSGSLLG GKAALTLSGV QPEDEAEYYC ALWYSNRWVF 240
 GGGTKLTVL 249

SEQ ID NO: 65
 FEATURE Location/Qualifiers
 REGION 1..14
 note = Synthetic peptide
 source 1..14
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 65
 RSSTGAVTSG YYPN 14

SEQ ID NO: 66
 FEATURE Location/Qualifiers
 REGION 1..7
 note = Synthetic peptide
 source 1..7

-continued

SEQUENCE: 66	mol_type = protein	
ATDMRPS	organism = synthetic construct	7
SEQ ID NO: 67	moltype = AA length = 9	
FEATURE	Location/Qualifiers	
REGION	1..9	
source	note = Synthetic peptide	
	1..9	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 67		
ALWYSNRWV		9
SEQ ID NO: 68	moltype = AA length = 5	
FEATURE	Location/Qualifiers	
REGION	1..5	
source	note = Synthetic peptide	
	1..5	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 68		
GYAMN		5
SEQ ID NO: 69	moltype = AA length = 19	
FEATURE	Location/Qualifiers	
REGION	1..19	
source	note = Synthetic peptide	
	1..19	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 69		
RIRSKYNNYA TYYADSVKE		19
SEQ ID NO: 70	moltype = AA length = 14	
FEATURE	Location/Qualifiers	
REGION	1..14	
source	note = Synthetic peptide	
	1..14	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 70		
HRNFGNSYLS WFAY		14
SEQ ID NO: 71	moltype = AA length = 125	
FEATURE	Location/Qualifiers	
REGION	1..125	
source	note = Synthetic peptide	
	1..125	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 71		
EVQLVESGGG LVQPGGSLKL SCAASGFTFN GYAMNWRQA PGKGLEWVAR IRSKYNNYAT		60
YYADSVKERF TISRDDSKNT AYLQMNNLKT EDTAVYYCVR HRNFGNSYLS WFAYWGQGTL		120
VTVSS		125
SEQ ID NO: 72	moltype = AA length = 109	
FEATURE	Location/Qualifiers	
REGION	1..109	
source	note = Synthetic peptide	
	1..109	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 72		
QTVVTQEPSTL TVSPGGTVTL TCRSSTGAVT SGYYPNWVQQ KPGQAPRGLI GATDMRPSGT		60
PARFSGSLLG GKAALTLSGV QPEDEAEYYC ALWYSNRWVF GGGTKLTVL		109
SEQ ID NO: 73	moltype = AA length = 249	
FEATURE	Location/Qualifiers	
REGION	1..249	
source	note = Synthetic peptide	
	1..249	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 73		

-continued

EVQLVESGGG	LVQPGGSLKL	SCAASGFTFN	GYAMNWVRQA	PGKGLEWVAR	IRSKYNNYAT	60
YYADSVKERF	TISRDDSKNT	AYLQMNNLKT	EDTAVYYCVR	HRNFGNSYLS	WFAYWGQGTL	120
VTVSSGGGGS	GGGGSGGGGS	QTVVTQEPSL	TVSPGGTVTL	TCRSSTGAVT	SGYYPNWVQQ	180
KPGQAPRGLI	GATDMRPSGT	PARFSGSLLG	GKAALTLSGV	QPEDEAEYYC	ALWYSNRWVF	240
GGGTKLTVL						249
SEQ ID NO: 74	moltype = AA length = 14					
FEATURE	Location/Qualifiers					
REGION	1..14					
	note = Synthetic peptide					
source	1..14					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 74						
GSSTGAVTSG YYPN						14
SEQ ID NO: 75	moltype = AA length = 7					
FEATURE	Location/Qualifiers					
REGION	1..7					
	note = Synthetic peptide					
source	1..7					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 75						
GTKFLAP						7
SEQ ID NO: 76	moltype = AA length = 9					
FEATURE	Location/Qualifiers					
REGION	1..9					
	note = Synthetic peptide					
source	1..9					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 76						
ALWYSNRWV						9
SEQ ID NO: 77	moltype = AA length = 5					
FEATURE	Location/Qualifiers					
REGION	1..5					
	note = Synthetic peptide					
source	1..5					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 77						
VYAMN						5
SEQ ID NO: 78	moltype = AA length = 19					
FEATURE	Location/Qualifiers					
REGION	1..19					
	note = Synthetic peptide					
source	1..19					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 78						
RIRSKYNNYA TYYADSVKK						19
SEQ ID NO: 79	moltype = AA length = 14					
FEATURE	Location/Qualifiers					
REGION	1..14					
	note = Synthetic peptide					
source	1..14					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 79						
HGNFGNSYIS WWAY						14
SEQ ID NO: 80	moltype = AA length = 125					
FEATURE	Location/Qualifiers					
REGION	1..125					
	note = Synthetic peptide					
source	1..125					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 80						
EVQLVESGGG	LVQPGGSLKL	SCAASGFTFN	VYAMNWVRQA	PGKGLEWVAR	IRSKYNNYAT	60
YYADSVKKRF	TISRDDSKNT	AYLQMNNLKT	EDTAVYYCVR	HGNFGNSYIS	WWAYWGQGTL	120

-continued

VTVSS	125
SEQ ID NO: 81	moltype = AA length = 109
FEATURE	Location/Qualifiers
REGION	1..109
source	note = Synthetic peptide
	1..109
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 81	
QTVVTQEPSL TVSPGGTVTL TCGSSTGAVT SGYYPNWVQQ KPGQAPRGLI GGTKFLAPGT	60
PARFSGSLLG GKAALTLGSG QPEDEAEYYC ALWYSNRWVF GGGTKLTVL	109
SEQ ID NO: 82	moltype = AA length = 249
FEATURE	Location/Qualifiers
REGION	1..249
source	note = Synthetic peptide
	1..249
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 82	
EVQLVESGGG LVQPGGSLKL SCAASGFTFN VYAMNWVRQA PGKGLEWVAR IRSKYNNYAT	60
YYADSVKKRF TISRDDSKNT AYLQMNLLKT EDTAVYVCVR HGNGNSYIS WWAYWGQGTL	120
VTVSSGGGGS GGGGSGGGS QTVVTQEPSL TVSPGGTVTL TCGSSTGAVT SGYYPNWVQQ	180
KPGQAPRGLI GGTKFLAPGT PARFSGSLLG GKAALTLGSG QPEDEAEYYC ALWYSNRWVF	240
GGGTKLTVL	249
SEQ ID NO: 83	moltype = AA length = 14
FEATURE	Location/Qualifiers
REGION	1..14
source	note = Synthetic peptide
	1..14
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 83	
GSSTGAVTSG NYPN	14
SEQ ID NO: 84	moltype = AA length = 7
FEATURE	Location/Qualifiers
REGION	1..7
source	note = Synthetic peptide
	1..7
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 84	
GTKFLAP	7
SEQ ID NO: 85	moltype = AA length = 9
FEATURE	Location/Qualifiers
REGION	1..9
source	note = Synthetic peptide
	1..9
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 85	
VLWYSNRWV	9
SEQ ID NO: 86	moltype = AA length = 5
FEATURE	Location/Qualifiers
REGION	1..5
source	note = Synthetic peptide
	1..5
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 86	
SYAMN	5
SEQ ID NO: 87	moltype = AA length = 19
FEATURE	Location/Qualifiers
REGION	1..19
source	note = Synthetic peptide
	1..19
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 87	
RIRSKYNNYA TYYADSVKG	19

SEQ ID NO: 88	moltype = AA length = 14		
FEATURE	Location/Qualifiers		
REGION	1..14		
	note = Synthetic peptide		
source	1..14		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 88			
HGNFGNSYVS WWAY			14
SEQ ID NO: 89	moltype = AA length = 125		
FEATURE	Location/Qualifiers		
REGION	1..125		
	note = Synthetic peptide		
source	1..125		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 89			
EVQLVESGGG LVQPGGSLKL	SCAASGFTFN SYAMNWVRQA	PGKGLEWVAR IRSKYNNYAT	60
YYADSVKGRF TISRDDSKNT	AYLQMNHLKT EDTAVYYCVR	HGNFGNSYVS WWAYWQGQTL	120
VTSS			125
SEQ ID NO: 90	moltype = AA length = 109		
FEATURE	Location/Qualifiers		
REGION	1..109		
	note = Synthetic peptide		
source	1..109		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 90			
QTVVTQEPSL TVSPGGTVTL	TCGSSTGAVT SGNYPNWVQQ	KPGQAPRGLI GGTKFLAPGT	60
PARFSGSLLG GKAAALTLSGV	QPEDEAEYYC VLWYSNRWVF	GGGTKLTVL	109
SEQ ID NO: 91	moltype = AA length = 249		
FEATURE	Location/Qualifiers		
REGION	1..249		
	note = Synthetic peptide		
source	1..249		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 91			
EVQLVESGGG LVQPGGSLKL	SCAASGFTFN SYAMNWVRQA	PGKGLEWVAR IRSKYNNYAT	60
YYADSVKGRF TISRDDSKNT	AYLQMNHLKT EDTAVYYCVR	HGNFGNSYVS WWAYWQGQTL	120
VTSSGGGGG GGGSGGGGS	QTVVTQEPSL TVSPGGTVTL	TCGSSTGAVT SGNYPNWVQQ	180
KPGQAPRGLI GGTKFLAPGT	PARFSGSLLG GKAAALTLSGV	QPEDEAEYYC VLWYSNRWVF	240
GGGTKLTVL			249
SEQ ID NO: 92	moltype = AA length = 14		
FEATURE	Location/Qualifiers		
REGION	1..14		
	note = Synthetic peptide		
source	1..14		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 92			
GSSTGAVTSG NYPN			14
SEQ ID NO: 93	moltype = AA length = 7		
FEATURE	Location/Qualifiers		
REGION	1..7		
	note = Synthetic peptide		
source	1..7		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 93			
GTKFLAP			7
SEQ ID NO: 94	moltype = AA length = 9		
FEATURE	Location/Qualifiers		
REGION	1..9		
	note = Synthetic peptide		
source	1..9		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 94			

-continued

VLWYSNRWV	9
SEQ ID NO: 95	moltype = AA length = 5
FEATURE	Location/Qualifiers
REGION	1..5
source	note = Synthetic peptide
	1..5
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 95	
KYAMN	5
SEQ ID NO: 96	moltype = AA length = 19
FEATURE	Location/Qualifiers
REGION	1..19
source	note = Synthetic peptide
	1..19
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 96	
RIRSKYNNYA TYYADSVKD	19
SEQ ID NO: 97	moltype = AA length = 14
FEATURE	Location/Qualifiers
REGION	1..14
source	note = Synthetic peptide
	1..14
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 97	
HGNFGNSYIS YWAY	14
SEQ ID NO: 98	moltype = AA length = 125
FEATURE	Location/Qualifiers
REGION	1..125
source	note = Synthetic peptide
	1..125
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 98	
EVQLVESGGG LVQPGGSLKL SCAASGFTFN KYAMNWVRQA PGKGLEWVAR IRSKYNNYAT	60
YYADSVKDRF TISRDDSKNT AYLQMNLLKT EDTAVYYCVR HGNFGNSYIS YWAYWGQGTL	120
VTVSS	125
SEQ ID NO: 99	moltype = AA length = 109
FEATURE	Location/Qualifiers
REGION	1..109
source	note = Synthetic peptide
	1..109
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 99	
QTVVTQEPSL TVSPGGTVTL TCGSSTGAVT SGNYPNWVQQ KPGQAPRGLI GGTKFLAPGT	60
PARFSGSLLG GKAALTLGTV QPEDEAEYYC VLWYSNRWVF GGGTKLTVL	109
SEQ ID NO: 100	moltype = AA length = 249
FEATURE	Location/Qualifiers
REGION	1..249
source	note = Synthetic peptide
	1..249
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 100	
EVQLVESGGG LVQPGGSLKL SCAASGFTFN KYAMNWVRQA PGKGLEWVAR IRSKYNNYAT	60
YYADSVKDRF TISRDDSKNT AYLQMNLLKT EDTAVYYCVR HGNFGNSYIS YWAYWGQGTL	120
VTVSSGGGGS GGGSGGGGS QTVVTQEPSL TVSPGGTVTL TCGSSTGAVT SGNYPNWVQQ	180
KPGQAPRGLI GGTKFLAPGT PARFSGSLLG GKAALTLGTV QPEDEAEYYC VLWYSNRWVF	240
GGGTKLTVL	249
SEQ ID NO: 101	moltype = AA length = 125
FEATURE	Location/Qualifiers
REGION	1..125
source	note = Synthetic peptide
	1..125
	mol_type = protein
	organism = synthetic construct

-continued

SEQUENCE: 101
EVQLVESGGG LVQPGGSLRL SCAASGFTFN SYAMNWRQA PGKGLEWVAR IRSKYNNYAT 60
YYADSVKGRF TISRDDSNT AYLMNSLKT EDTAVYVCVR HGNFGNSYVS WWAYWGQGT 120
VTSS 125

SEQ ID NO: 102 moltype = AA length = 109
FEATURE Location/Qualifiers
REGION 1..109
 note = Synthetic peptide
source 1..109
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 102
QTVVTQEPSL TVSPGGTVTL TCGSSTGAVT SGNYPNWVQQ KPGQAPRGLI GGTKFLAPGT 60
PARFSGSLLG GKAALTLSGV QPEDEAEYYC VLWYSNRWVF GGGTKLTVL 109

SEQ ID NO: 103 moltype = AA length = 249
FEATURE Location/Qualifiers
REGION 1..249
 note = Synthetic peptide
source 1..249
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 103
EVQLVESGGG LVQPGGSLRL SCAASGFTFN SYAMNWRQA PGKGLEWVAR IRSKYNNYAT 60
YYADSVKGRF TISRDDSNT AYLMNSLKT EDTAVYVCVR HGNFGNSYVS WWAYWGQGT 120
VTSSGGGGS GGGSGGGGS QTVVTQEPSL TVSPGGTVTL TCGSSTGAVT SGNYPNWVQQ 180
KPGQAPRGLI GGTKFLAPGT PARFSGSLLG GKAALTLSGV QPEDEAEYYC VLWYSNRWVF 240
GGGTKLTVL 249

SEQ ID NO: 104 moltype = AA length = 585
FEATURE Location/Qualifiers
source 1..585
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 104
DAHKSEVAHR FKDLGEENFK ALVLIIFAQY LQQCPFEDHV KLVNEVTEFA KTCVADESAAE 60
NCDKSLHTLF GDKLCTVATL RETYGEADAC CAKQEPERNE CFLQHKDDNP NLPRLVRPEV 120
DVMCTAFHDN EETFLKKYLY EIARRHPYFY APELLFFAKR YKAAFTCECC AADKAACLLP 180
KLDELRLDEG ASSAKQRLKC ASLQKFGERA FKAWAVARLS QRFPAEFPAE VSKLVTDLTK 240
VHTECCHGDL LECADDRADL AKYICENQDS ISSKLKECCE KPLLEKSHCI AEVENDEMPA 300
DLPSLAADFV ESKDVCKNYA EAKDVFLGMF LYEYARRHPD YSVVLLRLA KTYETTLEKC 360
CAAADPHECY AKVFDEFKPL VEEPQNLIKQ NCELFEQLGE YKFNALLVR YTKKVPQVST 420
PTLVEVSRNL GKVGSKCKKH PEAKRMPCAE DYLSVVLNQL CVLHEKTPVS DRVTKCCTES 480
LVNRRPCFSA LEVDETYVPK EFNATFTFH ADICTLSEKE RQIKKQALV ELVKHKPKAT 540
KEQLKAVMDD FAAFVEKCKC ADDKETCFPAE EGKKLVAASQ AALGL 585

SEQ ID NO: 105 moltype = AA length = 585
FEATURE Location/Qualifiers
REGION 1..585
 note = Synthetic peptide
source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 105
DAHKSEVAHR FKDLGEENFK ALVLIIFAQY LQQCPFEDHV KLVNEVTEFA KTCVADESAAE 60
NCDKSLHTLF GDKLCTVATL RETYGEADAC CAKQEPERNE CFLQHKDDNP NLPRLVRPEV 120
DVMCTAFHDN EETFLKKYLY EIARRHPYFY APELLFFAKR YKAAFTCECC AADKAACLLP 180
KLDELRLDEG ASSAKQRLKC ASLQKFGERA FKAWAVARLS QRFPAEFPAE VSKLVTDLTK 240
VHTECCHGDL LECADDRADL AKYICENQDS ISSKLKECCE KPLLEKSHCI AEVENDEMPA 300
DLPSLAADFV ESKDVCKNYA EAKDVFLGMF LYEYARRHPD YSVVLLRLA KTYETTLEKC 360
CAAADPHECY AKVFDEFKPL VEEPQNLIKQ NCELFEQLGE YKFNALLVR YTKKVPQVST 420
PTLVEVSRNL GKVGSKCKKH PEAKRMPCAE DYLSVVLNQL CVLHEKTPVS DRVTKCCTES 480
LVNRRPCFSA LEVDETYVPK EFNATFTFH ADICTLSEKE RQIKKQALV ELVKHKPKAT 540
KEQLKAAMD FAAFVEKCKC ADDKETCFPAE EGKKLVAASQ AALGL 585

SEQ ID NO: 106 moltype = AA length = 585
FEATURE Location/Qualifiers
REGION 1..585
 note = Synthetic peptide
source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 106
DAHKSEVAHR FKDLGEENFK ALVLIIFAQY LQQCPFEDHV KLVNEVTEFA KTCVADESAAE 60
NCDKSLHTLF GDKLCTVATL RETYGEADAC CAKQEPERNE CFLQHKDDNP NLPRLVRPEV 120

-continued

DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LDVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMD	FAAFVEKCKC	ADDKETCFAE	EGPKLVAAASQ	AALGL		585

SEQ ID NO: 107 moltype = AA length = 585
 FEATURE Location/Qualifiers
 REGION 1..585
 note = Synthetic peptide
 source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 107

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQCPFEDHV	KLVNEVTEFA	KTCVADES	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120
DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LDVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMD	FAAFVEKCKC	ADDKETCFAE	EGPHLVAAASQ	AALGL		585

SEQ ID NO: 108 moltype = AA length = 585
 FEATURE Location/Qualifiers
 REGION 1..585
 note = Synthetic peptide
 source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 108

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQCPFEDHV	KLVNEVTEFA	KTCVADES	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120
DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LDVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMD	FAAFVEKCKC	ADDKETCFAE	EGPKLVAAASQ	AALGL		585

SEQ ID NO: 109 moltype = AA length = 585
 FEATURE Location/Qualifiers
 REGION 1..585
 note = Synthetic peptide
 source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 109

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQCPFEDHV	KLVNEVTEFA	KTCVADES	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120
DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LDVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMD	FAAFVEKCKC	ADDKETCFAE	EGPKLVAAASQ	AALGL		585

SEQ ID NO: 110 moltype = AA length = 585
 FEATURE Location/Qualifiers
 REGION 1..585
 note = Synthetic peptide
 source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 110

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQCPFEDHV	KLVNEVTEFA	KTCVADES	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120

-continued

DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAKQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LEVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMDK	FAAFVEKCKK	ADDKETCFAE	EGPHLVAASQ	AALGL		585

SEQ ID NO: 111 moltype = AA length = 585
 FEATURE Location/Qualifiers
 REGION 1..585
 note = Synthetic peptide
 source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 111

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQCPFEDHV	KLVNEVTEFA	KTCVADESAE	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120
DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAKQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LEVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMDK	FAAFVEKCKK	ADDKETCFAE	EGPHLVAASK	AALGL		585

SEQ ID NO: 112 moltype = AA length = 585
 FEATURE Location/Qualifiers
 REGION 1..585
 note = Synthetic peptide
 source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 112

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQCPFEDHV	KLVNEVTEFA	KTCVADESAE	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120
DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAKQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LEVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMDK	FAAFVEKCKK	ADDKETCFAE	EGPKLVAASK	AALGL		585

SEQ ID NO: 113 moltype = AA length = 585
 FEATURE Location/Qualifiers
 REGION 1..585
 note = Synthetic peptide
 source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 113

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQCPFEDHV	KLVNEVTEFA	KTCVADESAE	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120
DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAKQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LDVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMDK	FAAFVEKCKK	ADDKETCFAE	EGPKLVAASK	AALGL		585

SEQ ID NO: 114 moltype = AA length = 585
 FEATURE Location/Qualifiers
 REGION 1..585
 note = Synthetic peptide
 source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 114

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQSPFEDHV	KLVNEVTEFA	KTCVADESAE	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120

-continued

DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAKQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LEVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMDD	FAAFVEKCKK	ADDKETCFAE	EGKKLVAAASQ	AALGL		585

SEQ ID NO: 115 moltype = AA length = 585
 FEATURE Location/Qualifiers
 REGION 1..585
 note = Synthetic peptide
 source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 115

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQSPFEDHV	KLVNEVTEFA	KTCVADESAE	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120
DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAKQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LEVDETYVPK	EFNAGTFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAAMD	FAAFVEKCKK	ADDKETCFAE	EGKKLVAAASQ	AALGL		585

SEQ ID NO: 116 moltype = AA length = 585
 FEATURE Location/Qualifiers
 REGION 1..585
 note = Synthetic peptide
 source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 116

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQSPFEDHV	KLVNEVTEFA	KTCVADESAE	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120
DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAKQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LEVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMDD	FAAFVEKCKK	ADDKETCFAE	EGPKLVAAASQ	AALGL		585

SEQ ID NO: 117 moltype = AA length = 585
 FEATURE Location/Qualifiers
 REGION 1..585
 note = Synthetic peptide
 source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 117

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQSPFEDHV	KLVNEVTEFA	KTCVADESAE	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120
DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAKQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LDVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMDD	FAAFVEKCKK	ADDKETCFAE	EGPHLVAAASK	AALGL		585

SEQ ID NO: 118 moltype = AA length = 585
 FEATURE Location/Qualifiers
 REGION 1..585
 note = Synthetic peptide
 source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 118

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQSPFEDHV	KLVNEVTEFA	KTCVADESAE	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120

-continued

DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTCECC	AADKAACLLP	180
KLDELRLDEGK	ASSAQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LGVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMDK	FAAFVEKCKK	ADDKETCFAE	EGPKLVAAASQ	AALGL		585

SEQ ID NO: 119 moltype = AA length = 585

FEATURE Location/Qualifiers

REGION 1..585

source note = Synthetic peptide

1..585

mol_type = protein

organism = synthetic construct

SEQUENCE: 119

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQSPFEDHV	KLVNEVTEFA	KTCVADESAE	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120
DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTCECC	AADKAACLLP	180
KLDELRLDEGK	ASSAQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LDVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMDK	FAAFVEKCKK	ADDKETCFAE	EGPKLVAAASQ	AALGL		585

SEQ ID NO: 120 moltype = AA length = 585

FEATURE Location/Qualifiers

REGION 1..585

source note = Synthetic peptide

1..585

mol_type = protein

organism = synthetic construct

SEQUENCE: 120

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQSPFEDHV	KLVNEVTEFA	KTCVADESAE	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120
DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTCECC	AADKAACLLP	180
KLDELRLDEGK	ASSAQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LEVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMDK	FAAFVEKCKK	ADDKETCFAE	EGPHLVAAASK	AALGL		585

SEQ ID NO: 121 moltype = AA length = 585

FEATURE Location/Qualifiers

REGION 1..585

source note = Synthetic peptide

1..585

mol_type = protein

organism = synthetic construct

SEQUENCE: 121

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQSPFEDHV	KLVNEVTEFA	KTCVADESAE	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120
DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTCECC	AADKAACLLP	180
KLDELRLDEGK	ASSAQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LEVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMDK	FAAFVEKCKK	ADDKETCFAE	EGPHLVAAASK	AALGL		585

SEQ ID NO: 122 moltype = AA length = 585

FEATURE Location/Qualifiers

REGION 1..585

source note = Synthetic peptide

1..585

mol_type = protein

organism = synthetic construct

SEQUENCE: 122

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQSPFEDHV	KLVNEVTEFA	KTCVADESAE	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120

-continued

DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAKQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LEVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMDK	FAAFVEKCKK	ADDKETCFAE	EGPKLVAASK	AALGL		585

SEQ ID NO: 123 moltype = AA length = 585
 FEATURE Location/Qualifiers
 REGION 1..585
 note = Synthetic peptide
 source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 123

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQSPFEDHV	KLVNEVTEFA	KTCVADESAE	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120
DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAKQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LEVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMDK	FAAFVEKCKK	ADDKETCFAE	EGPKLVAASK	AALGL		585

SEQ ID NO: 124 moltype = AA length = 585
 FEATURE Location/Qualifiers
 REGION 1..585
 note = Synthetic peptide
 source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 124

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQAPFEDHV	KLVNEVTEFA	KTCVADESAE	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120
DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAKQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LEVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMDK	FAAFVEKCKK	ADDKETCFAE	EGKKLVAASQ	AALGL		585

SEQ ID NO: 125 moltype = AA length = 585
 FEATURE Location/Qualifiers
 REGION 1..585
 note = Synthetic peptide
 source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 125

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQAPFEDHV	KLVNEVTEFA	KTCVADESAE	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120
DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAKQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LEVDETYVPK	EFNAGTFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAAMD	FAAFVEKCKK	ADDKETCFAE	EGKKLVAASQ	AALGL		585

SEQ ID NO: 126 moltype = AA length = 585
 FEATURE Location/Qualifiers
 REGION 1..585
 note = Synthetic peptide
 source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 126

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQAPFEDHV	KLVNEVTEFA	KTCVADESAE	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120

-continued

DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTCECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LDVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMDD	FAAFVEKCKK	ADDKETCFAE	EGPKLVAAASQ	AALGL		585

SEQ ID NO: 127 moltype = AA length = 585
 FEATURE Location/Qualifiers
 REGION 1..585
 note = Synthetic peptide
 source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 127

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQAPFEDHV	KLVNEVTEFA	KTCVADESAE	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120
DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTCECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LDVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMDD	FAAFVEKCKK	ADDKETCFAE	EGPHLVAAASK	AALGL		585

SEQ ID NO: 128 moltype = AA length = 585
 FEATURE Location/Qualifiers
 REGION 1..585
 note = Synthetic peptide
 source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 128

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQAPFEDHV	KLVNEVTEFA	KTCVADESAE	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120
DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTCECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LDVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMDK	FAAFVEKCKK	ADDKETCFAE	EGPKLVAAASQ	AALGL		585

SEQ ID NO: 129 moltype = AA length = 585
 FEATURE Location/Qualifiers
 REGION 1..585
 note = Synthetic peptide
 source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 129

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQAPFEDHV	KLVNEVTEFA	KTCVADESAE	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120
DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTCECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LDVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMDK	FAAFVEKCKK	ADDKETCFAE	EGPKLVAAASQ	AALGL		585

SEQ ID NO: 130 moltype = AA length = 585
 FEATURE Location/Qualifiers
 REGION 1..585
 note = Synthetic peptide
 source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 130

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQAPFEDHV	KLVNEVTEFA	KTCVADESAE	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120

-continued

DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAKQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LEVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMDK	FAAFVEKCKK	ADDKETCFAE	EGPHLVAASQ	AALGL		585

SEQ ID NO: 131 moltype = AA length = 585
 FEATURE Location/Qualifiers
 REGION 1..585
 note = Synthetic peptide
 source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 131

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQAPFEDHV	KLVNEVTEFA	KTCVADESAE	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120
DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAKQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LEVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMDK	FAAFVEKCKK	ADDKETCFAE	EGPHLVAASK	AALGL		585

SEQ ID NO: 132 moltype = AA length = 585
 FEATURE Location/Qualifiers
 REGION 1..585
 note = Synthetic peptide
 source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 132

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQAPFEDHV	KLVNEVTEFA	KTCVADESAE	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120
DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAKQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LEVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMDK	FAAFVEKCKK	ADDKETCFAE	EGPKLVAASK	AALGL		585

SEQ ID NO: 133 moltype = AA length = 585
 FEATURE Location/Qualifiers
 REGION 1..585
 note = Synthetic peptide
 source 1..585
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 133

DAHKSEVAHR	FKDLGEENFK	ALVLIIFAQY	LQQAPFEDHV	KLVNEVTEFA	KTCVADESAE	60
NCDKSLHTLF	GDKLCTVATL	RETYGEMADC	CAKQEPERNE	CFLQHKDDNP	NLPRLVRPEV	120
DVMCTAFHDN	EETFLKKYLY	EIARRHPYFY	APELLFFAKR	YKAAFTECCQ	AADKAACLLP	180
KLDELRLDEGK	ASSAKQRLKC	ASLQKFGERA	FKAWAVARLS	QRFPKAEFAE	VSKLVTDLTK	240
VHTECCHGDL	LECADDRADL	AKYICENQDS	ISSKLKECCE	KPLLEKSHCI	AEVENDEMPA	300
DLPSLAADFV	ESKDVCKNYA	EAKDVFLGMF	LYEYARRHPD	YSVVLRLRLA	KTYETTLEKC	360
CAAADPHECY	AKVFDEFKPL	VEEPQNLIKQ	NCELFEQLGE	YKFQNALLV	YTKKVPQVST	420
PTLVEVSRNL	GKVGSKCKH	PEAKRMPCAE	DYLSVVLNQL	CVLHEKTPVS	DRVTKCCTES	480
LVNRRPCFSA	LDVDETYVPK	EFNAETFTFH	ADICTLSEKE	RQIKKQTALV	ELVKHKPKAT	540
KEQLKAVMDK	FAAFVEKCKK	ADDKETCFAE	EGPKLVAASK	AALGL		585

SEQ ID NO: 134 moltype = AA length = 16
 FEATURE Location/Qualifiers
 REGION 1..16
 note = Synthetic peptide
 source 1..16
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 134
 RDWDFDVFGG GTPVGG

16

-continued

SEQ ID NO: 135	moltype = AA length = 16	
FEATURE	Location/Qualifiers	
REGION	1..16	
	note = Synthetic peptide	
SITE	12	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
source	1..16	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 135		
QRFVTGHFVG LXPANG		16
SEQ ID NO: 136	moltype = AA length = 16	
FEATURE	Location/Qualifiers	
REGION	1..16	
	note = Synthetic peptide	
source	1..16	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 136		
QRFVTGHFVG LYPANG		16
SEQ ID NO: 137	moltype = AA length = 16	
FEATURE	Location/Qualifiers	
REGION	1..16	
	note = Synthetic peptide	
source	1..16	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 137		
QRFVTGHFVG LHPANG		16
SEQ ID NO: 138	moltype = AA length = 9	
FEATURE	Location/Qualifiers	
REGION	1..9	
	note = Synthetic peptide	
source	1..9	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 138		
TGHFGGLHP		9
SEQ ID NO: 139	moltype = AA length = 16	
FEATURE	Location/Qualifiers	
REGION	1..16	
	note = Synthetic peptide	
source	1..16	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 139		
QRFCTGHFVG LHPCNG		16
SEQ ID NO: 140	moltype = AA length = 330	
FEATURE	Location/Qualifiers	
REGION	1..330	
	note = Synthetic peptide	
source	1..330	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 140		
ASTKGPSVFP LAPSSKSTSG GTAALGCLVK DYFPEPVTVS WNSGALTSGV HTPFAVLQSS		60
GLYSLSSVVT VPSSSLGTQT YICNVNHNKPS NTKVDKKVEP KSCDKTHTCP PCPAPELLGG		120
PSVFLFPPKP KDTLMISRT EVTCVVVDVS HEDPEVKFNW YVDGVEVHNA KTKPCEEQYG		180
STYRCVSVLT VLHQDWLNGK EYCKKVSNKA LPAPIEKTIS KAKGQPREPQ VYTLPPSREE		240
MTKNQVSLTC LVKGFYPSDI AVEWESNGQP ENNYDTTPPV LDSDGSFFLY SDLTVDKSRW		300
QQGNVFSCSV MHEALHNHYT QKSLSLSPGK		330
SEQ ID NO: 141	moltype = AA length = 333	
FEATURE	Location/Qualifiers	
REGION	1..333	
	note = Synthetic peptide	
source	1..333	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 141		

-continued

GQPKAAPSVT	LPPPSSEELQ	ANKATLVCLI	SDFYPGAVTV	AWKADSSPVK	AGVETTPSK	60
QSNKYAASS	YLSLTPEQWK	SHRSYSCQVT	HEGSTVEKTV	APTECSDKTH	TCPPCPAPEL	120
LGGPSVFLFP	PKPKDTLMIS	RTPEVTCVVV	DVSHEDPEVK	FNWYVDGVEV	HNAKTKPCEE	180
QYGSTYRCVS	VLTVLHQDWL	NGKEYKCKVS	NKALPAPIEK	TISKAKGQPR	EPQVYTLPPS	240
RKEMTKNQVS	LTCLVKGFYP	SDIAVEWESN	GQPENNYKTT	PPVLKSDGSF	FLYKLTVDK	300
SRWQQGNVFS	CSVMHEALHN	HYTQKSLSL	PGK			333

SEQ ID NO: 142 moltype = AA length = 330
 FEATURE Location/Qualifiers
 REGION 1..330
 note = Synthetic peptide
 source 1..330
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 142

ASTKGPSVFP	LAPCSRSTSE	STAALGCLVK	DYFPEPVTVS	WNSGALTSGV	HTFPAVLQSS	60
GLYSLSSVVT	VPSSNFGTQT	YTCNVDHKPS	NTKVDKTVEP	KSSDKTHTCP	PCPAPEAAGG	120
PSVFLFPPKP	KDTLMISRTPE	EVTCVVVDVS	HEDPEVKFNW	YVDGVEVHNA	KTKPREEQYN	180
STYRVVSVLT	VLHQDWLNGK	EYKCKVSNKA	LPAPIEKTIS	KAKGQPREPQ	VYTLPPSREE	240
MTKNQVSLTC	LVKGFYPSDI	AVEWESNGQP	ENNYDTTPPV	LDSDGSFFLY	SDLTVDKSRW	300
QQGNVFSCSV	MHEALHNHYT	QKSLSLSPGK				330

SEQ ID NO: 143 moltype = AA length = 338
 FEATURE Location/Qualifiers
 REGION 1..338
 note = Synthetic peptide
 source 1..338
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 143

GQPKAAPSVT	LPPPSSEELQ	ANKATLVCLI	SDFYPGAVTV	AWKADSSPVK	AGVETTPSK	60
QSNKYAASS	YLSLTPEQWK	SHRSYSCQVT	HEGSTVEKTV	APTECSEPKS	SDKTHTCPPC	120
PAPEAAGGPS	VFLFPPPKFD	TLMISRTPEV	TCVVVDVSHE	DPEVKFNWYV	DGVEVHNAKT	180
KPREEQYNST	YRVVSVLTVL	HQDWLNGKEY	KCKVSNKALP	APIEKTISKA	KGQPREPQVY	240
TLPPSRKEMT	KNQVSLTCLV	KGFYPSDIAV	EWESNGQPEN	NYKTTTPVLK	SDGSFFLYSK	300
LTVDKSRWQQ	GNVFSCSVMH	EALHNHYTQK	SLSLSPGK			338

SEQ ID NO: 144 moltype = AA length = 227
 FEATURE Location/Qualifiers
 REGION 1..227
 note = Synthetic peptide
 source 1..227
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 144

DKTHTCPPCP	APELLGGPSV	FLFPPKPKDT	LMISRTPEVT	CVVVDVSHED	PEVKFNWYVD	60
GVEVHNAKTK	PCREEQYGSTY	RCVSVLTVLH	QDWLNGKEYK	CKVSNKALPA	PIEKTISKAK	120
GQPREPQVYT	LPPSRKEMTK	NQVSLTCLVK	GFYPSDIAVE	WESNGQPENN	YKTTTPVLKS	180
DGSFFLYSKL	TVDKSRWQQG	NVFSCSVMHE	ALHNHYTQKS	LSLSPGK		227

SEQ ID NO: 145 moltype = AA length = 227
 FEATURE Location/Qualifiers
 REGION 1..227
 note = Synthetic peptide
 source 1..227
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 145

DKTHTCPPCP	APELLGGPSV	FLFPPKPKDT	LMISRTPEVT	CVVVDVSHED	PEVKFNWYVD	60
GVEVHNAKTK	PCREEQYGSTY	RCVSVLTVLH	QDWLNGKEYK	CKVSNKALPA	PIEKTISKAK	120
GQPREPQVYT	LPPSRKEMTK	NQVSLTCLVK	GFYPSDIAVE	WESNGQPENN	YDTTPPVLD	180
DGSFFLYSDL	TVDKSRWQQG	NVFSCSVMHE	ALHNHYTQKS	LSLSPGK		227

SEQ ID NO: 146 moltype = AA length = 232
 FEATURE Location/Qualifiers
 REGION 1..232
 note = Synthetic peptide
 source 1..232
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 146

EPKSSDKTHT	CPPCPAPELL	GGPSVFLFPP	KPKDTLMISR	TPEVTCVVVD	VSHEDPEVKF	60
NWYVDGVEVH	NAKTKPCEEQ	YGSTYRCVSV	LTVLHQDWLN	GKEYKCKVSN	KALPAPIEKT	120
ISKAKGQPRE	PQVYTLPPSR	KEMTKNQVSL	TCLVKGFYPS	DIAVEWESNG	QPENNYKTTT	180
PVLKSDGSFF	LYSKLTVDKS	RWQQGNVFSC	SVMHEALHNH	YTQKSLSLSP	GK	232

SEQ ID NO: 147	moltype = AA length = 232					
FEATURE	Location/Qualifiers					
REGION	1..232					
	note = Synthetic peptide					
source	1..232					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 147						
EPKSSDKTHT	CPPCPAPELL	GGPSVFLFPP	KPKDTLMISR	TPEVTCVVVD	VSHEDPEVKF	60
NWYVDGVEVH	NAKTKPCEEQ	YGSTYRCVSV	LTVLHQDWLN	GKEYKCKVSN	KALPAPIEKT	120
ISKAKGQPRE	PQVYTLPPSR	EEMTKNQVSL	TCLVKGFYPS	DIAVEWESNG	QPENNYDTTP	180
PVLDSGDGFF	LYSDLTVDKS	RWQQGNVFSC	SVMHEALHNN	YTQKSLSLSP	GK	232
SEQ ID NO: 148	moltype = AA length = 232					
FEATURE	Location/Qualifiers					
REGION	1..232					
	note = Synthetic peptide					
source	1..232					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 148						
EPKSSDKTHT	CPPCPAPEAA	GGPSVFLFPP	KPKDTLMISR	TPEVTCVVVD	VSHEDPEVKF	60
NWYVDGVEVH	NAKTKPREEQ	YNSTYRVVSV	LTVLHQDWLN	GKEYKCKVSN	KALPAPIEKT	120
ISKAKGQPRE	PQVYTLPPSR	KEMTKNQVSL	TCLVKGFYPS	DIAVEWESNG	QPENNYKTPP	180
PVLKSDGSFF	LYSKLTVDKS	RWQQGNVFSC	SVMHEALHNN	YTQKSLSLSP	GK	232
SEQ ID NO: 149	moltype = AA length = 232					
FEATURE	Location/Qualifiers					
REGION	1..232					
	note = Synthetic peptide					
source	1..232					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 149						
EPKSSDKTHT	CPPCPAPEAA	GGPSVFLFPP	KPKDTLMISR	TPEVTCVVVD	VSHEDPEVKF	60
NWYVDGVEVH	NAKTKPREEQ	YNSTYRVVSV	LTVLHQDWLN	GKEYKCKVSN	KALPAPIEKT	120
ISKAKGQPRE	PQVYTLPPSR	EEMTKNQVSL	TCLVKGFYPS	DIAVEWESNG	QPENNYDTTP	180
PVLDSGDGFF	LYSDLTVDKS	RWQQGNVFSC	SVMHEALHNN	YTQKSLSLSP	GK	232
SEQ ID NO: 150	moltype = AA length = 217					
FEATURE	Location/Qualifiers					
REGION	1..217					
	note = Synthetic peptide					
source	1..217					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 150						
APELLGGPSV	FLFPPPKPKDT	LMISRTPEVT	CVVVDVSHED	PEVKFNWYVD	GVEVHNAKTK	60
PREEQYNSTY	RVVSVLTVLH	QDWLNGKEYK	CKVSNKALPA	PIEKTISKAK	GQPREPQVTT	120
LPSPREEMTK	NQVSLTCLVK	GFYPSDIAVE	WESNGQPENN	YDTPPVLDL	DGSFFLYSDL	180
TVDKSRWQQG	NVFSFSVMHE	ALHNNHYTQK	LSLSPGK			217
SEQ ID NO: 151	moltype = AA length = 7					
FEATURE	Location/Qualifiers					
REGION	1..7					
	note = Synthetic peptide					
source	1..7					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 151						
SSSYWG						7
SEQ ID NO: 152	moltype = AA length = 16					
FEATURE	Location/Qualifiers					
REGION	1..16					
	note = Synthetic peptide					
source	1..16					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 152						
SIYSGITNY	NPSLKS					16
SEQ ID NO: 153	moltype = AA length = 9					
FEATURE	Location/Qualifiers					
REGION	1..9					
	note = Synthetic peptide					

-continued

source	1..9	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 153		
PSNYDAFDI		9
SEQ ID NO: 154	moltype = AA length = 14	
FEATURE	Location/Qualifiers	
REGION	1..14	
	note = Synthetic peptide	
source	1..14	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 154		
TGSSSNIGAG YDVH		14
SEQ ID NO: 155	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
	note = Synthetic peptide	
source	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 155		
GNSKRPS		7
SEQ ID NO: 156	moltype = AA length = 11	
FEATURE	Location/Qualifiers	
REGION	1..11	
	note = Synthetic peptide	
source	1..11	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 156		
QSYDSSLGGW V		11
SEQ ID NO: 157	moltype = AA length = 119	
FEATURE	Location/Qualifiers	
REGION	1..119	
	note = Synthetic peptide	
source	1..119	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 157		
QVQLQESGPG LVKPSSETLSL TCTVSGGSIS SSSYYWGWR QPPGKGLEWI GSIYYSGITN		60
YNPSLKSRVT ISVDTSKNQF SLKLSSVTAA DTAVYYCARP SNYDAFDIWG QGTMTVSS		119
SEQ ID NO: 158	moltype = AA length = 111	
FEATURE	Location/Qualifiers	
REGION	1..111	
	note = Synthetic peptide	
source	1..111	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 158		
QSVLTQPPSV SGAPGQRTVI SCTGSSSNIG AGYDVHWYQQ VPGTAPKLLI YGNSKRPSGV		60
PDRFSGSKSG TSASLAITGL QAEDEADYYC QSYDSSLGGW VFGGGTKLTV L		111
SEQ ID NO: 159	moltype = AA length = 245	
FEATURE	Location/Qualifiers	
REGION	1..245	
	note = Synthetic peptide	
source	1..245	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 159		
QVQLQESGPG LVKPSSETLSL TCTVSGGSIS SSSYYWGWR QPPGKGLEWI GSIYYSGITN		60
YNPSLKSRVT ISVDTSKNQF SLKLSSVTAA DTAVYYCARP SNYDAFDIWG QGTMTVSSG		120
GGGSGGGSG GGSQSVLTQ PPSVSGAPGQ RVTISCTGSS SNIGAGYDVH WYQQVPGTAP		180
KLLIYGNSKR PSGVPDRFSG SKSGTSASLA ITGLQAEDEA DYYCQSYDSS LGGWVFGGGT		240
KLTVL		245
SEQ ID NO: 160	moltype = AA length = 500	
FEATURE	Location/Qualifiers	
REGION	1..500	
	note = Synthetic peptide	

-continued

```

source                1..500
                      mol_type = protein
                      organism = synthetic construct

SEQUENCE: 160
QVQLQESGPG LVKPSSETLSL TCTVSGGSIS SSSYYGWIR QPPGKGLEWI GSIYYSGITN 60
YNPSLKSRVT ISVDTSKNQF SLKLSSVTAA DTAVYYCARP SNYDAFDIWG QGTMTVTVSSG 120
GGSGGGGGSG GGSQSQSLTQ PPSVSGAPGQ RVTISCTGSS SNIGAGYDVH WYQQVPGTAP 180
KLLIYGNSKR PSGVPDRFSG SKSGTSASLA ITGLQAEDEA DYYCQSYDSS LGGWVFGGTT 240
KLTVLSSGGG SEVQLVESGG GLVQPGGSLK LSCAASGFTF NKYAMNWVRQ APGKGLEWVA 300
RIRSKYNNYA TYYADSVKDR FTISRDDSKN TAYLQMNLLK TEDTAVYYCV RHGNFGNSYI 360
SYWAYWGQGT LVTVSSGGGG SGGGGSGGGG SQTIVTQEPS LTVSPGGTGT LTCGSSTGAV 420
TSGNYPNWVQ QKPGQAPRGL IGGTKFLAPG TPARFSGSLG GGKAALTLSG VQPEDEAEYY 480
CVLWYSNRWV PGGGTKLTVL                                     500

SEQ ID NO: 161        moltype = AA length = 5
FEATURE              Location/Qualifiers
REGION               1..5
                     note = Synthetic peptide
source               1..5
                     mol_type = protein
                     organism = synthetic construct

SEQUENCE: 161
DHYS                                     5

SEQ ID NO: 162        moltype = AA length = 17
FEATURE              Location/Qualifiers
REGION               1..17
                     note = Synthetic peptide
source               1..17
                     mol_type = protein
                     organism = synthetic construct

SEQUENCE: 162
YISSSGSTIY YADSVKG                                     17

SEQ ID NO: 163        moltype = AA length = 8
FEATURE              Location/Qualifiers
REGION               1..8
                     note = Synthetic peptide
source               1..8
                     mol_type = protein
                     organism = synthetic construct

SEQUENCE: 163
DLGPSFDY                                     8

SEQ ID NO: 164        moltype = AA length = 11
FEATURE              Location/Qualifiers
REGION               1..11
                     note = Synthetic peptide
source               1..11
                     mol_type = protein
                     organism = synthetic construct

SEQUENCE: 164
RASQGISSWL A                                     11

SEQ ID NO: 165        moltype = AA length = 7
FEATURE              Location/Qualifiers
REGION               1..7
                     note = Synthetic peptide
source               1..7
                     mol_type = protein
                     organism = synthetic construct

SEQUENCE: 165
AASRLQS                                     7

SEQ ID NO: 166        moltype = AA length = 9
FEATURE              Location/Qualifiers
REGION               1..9
                     note = Synthetic peptide
source               1..9
                     mol_type = protein
                     organism = synthetic construct

SEQUENCE: 166
QQANSFPRT                                     9

SEQ ID NO: 167        moltype = AA length = 117
FEATURE              Location/Qualifiers

```

-continued

REGION 1..117
 note = Synthetic peptide
 source 1..117
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 167
 QVQLVESGGG VVQPGRLRL SCAASGFTFS DHYMSWIRQT PGKGLEWVSY ISSSGSTIYY 60
 ADSVKGRFTI SRDPAKNSLY LQMNSLRAED TAVYYCARDL GPSFDYWGG SLVTVSS 117

SEQ ID NO: 168 moltype = AA length = 107
 FEATURE Location/Qualifiers
 REGION 1..107
 note = Synthetic peptide
 source 1..107
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 168
 DIQMTQSPSS VSASVGDRTV ITCRASQGIS SWLAWYQQK GKAPKLLIDA ASRLQSGVPS 60
 RFGSGSGGTD FTLTISSLQP EDFATYYCQQ ANSPRFTFGQ GTKVEIK 107

SEQ ID NO: 169 moltype = AA length = 239
 FEATURE Location/Qualifiers
 REGION 1..239
 note = Synthetic peptide
 source 1..239
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 169
 QVQLVESGGG VVQPGRLRL SCAASGFTFS DHYMSWIRQT PGKGLEWVSY ISSSGSTIYY 60
 ADSVKGRFTI SRDPAKNSLY LQMNSLRAED TAVYYCARDL GPSFDYWGG SLVTVSSGGG 120
 GSGGGSGGG GSDIQMTQSP SSVASVGDRTV ITCRASQG ISSWLAWYQQ KPGKAPKLLI 180
 DAASRLQSGV PSRFGSGSGG TDFTLTISSL QPEDFATYYC QQANSPRFTF GQGTKVEIK 239

SEQ ID NO: 170 moltype = AA length = 494
 FEATURE Location/Qualifiers
 REGION 1..494
 note = Synthetic peptide
 source 1..494
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 170
 QVQLVESGGG VVQPGRLRL SCAASGFTFS DHYMSWIRQT PGKGLEWVSY ISSSGSTIYY 60
 ADSVKGRFTI SRDPAKNSLY LQMNSLRAED TAVYYCARDL GPSFDYWGG SLVTVSSGGG 120
 GSGGGSGGG GSDIQMTQSP SSVASVGDRTV ITCRASQG ISSWLAWYQQ KPGKAPKLLI 180
 DAASRLQSGV PSRFGSGSGG TDFTLTISSL QPEDFATYYC QQANSPRFTF GQGTKVEIKS 240
 GGGGSEVQLV ESGGGLVQPG GSLKLSAAS GFTFNKYAMN WVRQAPGKGL EWVARIRSKY 300
 NNYATYYADS VKDRFTISR DSKNTAYLQM NNLKTEDTAV YYCVRHGNFG NSYISYWAYW 360
 GQGTLLVTSS GGGGSGGGGS GGGGSGTVVT QEPSTLVSPG GTVTLTCGSS TGAVTSGNYP 420
 NWVQKPGQA PRGLIGTKF LAPGTPARFS GSLLGGKAAL TSGVQPEDE AEYVCVLWYS 480
 NRWVFGGGTK LTVL 494

SEQ ID NO: 171 moltype = AA length = 5
 FEATURE Location/Qualifiers
 REGION 1..5
 note = Synthetic peptide
 source 1..5
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 171
 DYYMT 5

SEQ ID NO: 172 moltype = AA length = 17
 FEATURE Location/Qualifiers
 REGION 1..17
 note = Synthetic peptide
 source 1..17
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 172
 YISSSGSTIY YADSVKG 17

SEQ ID NO: 173 moltype = AA length = 8
 FEATURE Location/Qualifiers
 REGION 1..8
 note = Synthetic peptide
 source 1..8

-continued

		mol_type = protein organism = synthetic construct	
SEQUENCE: 173			
DRNSHPDY			8
SEQ ID NO: 174		moltype = AA length = 11	
FEATURE		Location/Qualifiers	
REGION		1..11	
		note = Synthetic peptide	
source		1..11	
		mol_type = protein	
		organism = synthetic construct	
SEQUENCE: 174			
RASQGITRWL A			11
SEQ ID NO: 175		moltype = AA length = 7	
FEATURE		Location/Qualifiers	
REGION		1..7	
		note = Synthetic peptide	
source		1..7	
		mol_type = protein	
		organism = synthetic construct	
SEQUENCE: 175			
AASVLQS			7
SEQ ID NO: 176		moltype = AA length = 9	
FEATURE		Location/Qualifiers	
REGION		1..9	
		note = Synthetic peptide	
source		1..9	
		mol_type = protein	
		organism = synthetic construct	
SEQUENCE: 176			
QQSNSFPRT			9
SEQ ID NO: 177		moltype = AA length = 117	
FEATURE		Location/Qualifiers	
REGION		1..117	
		note = Synthetic peptide	
source		1..117	
		mol_type = protein	
		organism = synthetic construct	
SEQUENCE: 177			
QVQLVESGGG LVKPGGSLRL SCAASGFTFS DYYMTWIRQA PGKGLEWISY ISSSGSTIYY	60		
ADSVKGRFTI SRDIAKNSLY LQMNSLRAED TAVYYCARDR NSHPDYWGQG TLVTVSS	117		
SEQ ID NO: 178		moltype = AA length = 107	
FEATURE		Location/Qualifiers	
REGION		1..107	
		note = Synthetic peptide	
source		1..107	
		mol_type = protein	
		organism = synthetic construct	
SEQUENCE: 178			
DIQMTQSPSS VSASVGDRVT ITCRASQGIT RWLAWYQQKPK GKAPKLLIYA ASVLQSGVPS	60		
RFGSGSGGTD FTLTISSLQP EDFATYYCQQ SNSFPRTFGQ GTKVEIK	107		
SEQ ID NO: 179		moltype = AA length = 239	
FEATURE		Location/Qualifiers	
REGION		1..239	
		note = Synthetic peptide	
source		1..239	
		mol_type = protein	
		organism = synthetic construct	
SEQUENCE: 179			
QVQLVESGGG LVKPGGSLRL SCAASGFTFS DYYMTWIRQA PGKGLEWISY ISSSGSTIYY	60		
ADSVKGRFTI SRDIAKNSLY LQMNSLRAED TAVYYCARDR NSHPDYWGQG TLVTVSSGGG	120		
GSGGGSGGG GSDIQMTQSP SSVASVGDR VTITCRASQG ITRWLAWYQQ KPGKAPKLLI	180		
YAASVLQSGV PSRFGSGSG TDFTLTISSL QPEDFATYYC QQSNSFPRTF GQGTKVEIK	239		
SEQ ID NO: 180		moltype = AA length = 494	
FEATURE		Location/Qualifiers	
REGION		1..494	
		note = Synthetic peptide	
source		1..494	
		mol_type = protein	

-continued

```

                                organism = synthetic construct
SEQUENCE: 180
QVQLVESGGG LVKPGGSLRL SCAASGFTFS DYYMTWIRQA PGKGLEWISY ISSSGSTIYY 60
ADSVKGRFTI SRDNAKNSLY LQMNSLRAED TAVYYCARDR NSHFDYWGQG TLVTVSSGGG 120
GSGGGGSGGG GSDIQMTQSP SSVSASVGDR VTITCRASQG ITRWLAWYQQ KPGKAPKLLI 180
YAASVLQSGV PSRFGSGSGG TDFTLTISSL QPEDFATYYC QQSNSFPRTF GQGKVEIKS 240
GGGGSEVQLV ESGGGLVQPG GSLKLSAAS GFTFNKYAMN WVRQAPGKGL EWVARIRSKY 300
NNYATYYADS VKDRFTISRD DSKNTAYLQM NNLKTEDTAV YYCVRHGNFG NSYISYWAYW 360
GGQTLVTVSS GGGGSGGGGS GGGGSQTVVT QEPSLTVSPG GTVTLTCGSS TGAVTSGNYP 420
NWVQKPGQA PRGLIGGTKF LAPGTPARFS GSLLGGAAL TSLGVQPEDE AEYYCVLWYS 480
NRWVFGGGTK LTVL 494

SEQ ID NO: 181      moltype = AA  length = 5
FEATURE            Location/Qualifiers
REGION            1..5
                  note = Synthetic peptide
source            1..5
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 181
DYYMT 5

SEQ ID NO: 182      moltype = AA  length = 17
FEATURE            Location/Qualifiers
REGION            1..17
                  note = Synthetic peptide
source            1..17
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 182
YISSSGSTIY YADSVKG 17

SEQ ID NO: 183      moltype = AA  length = 8
FEATURE            Location/Qualifiers
REGION            1..8
                  note = Synthetic peptide
source            1..8
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 183
DRNSHFDY 8

SEQ ID NO: 184      moltype = AA  length = 11
FEATURE            Location/Qualifiers
REGION            1..11
                  note = Synthetic peptide
source            1..11
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 184
RASQGINTWL A 11

SEQ ID NO: 185      moltype = AA  length = 7
FEATURE            Location/Qualifiers
REGION            1..7
                  note = Synthetic peptide
source            1..7
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 185
GASGLQS 7

SEQ ID NO: 186      moltype = AA  length = 9
FEATURE            Location/Qualifiers
REGION            1..9
                  note = Synthetic peptide
source            1..9
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 186
QQAKSPRT 9

SEQ ID NO: 187      moltype = AA  length = 117
FEATURE            Location/Qualifiers
REGION            1..117
                  note = Synthetic peptide

```

-continued

```

source                1..117
                      mol_type = protein
                      organism = synthetic construct

SEQUENCE: 187
QVQLVESGGG LVKPGGSLRL SCAASGFTFS DYYMTWIRQA PGKGLEWLSY ISSSGSTIYY 60
ADSVKGRFTI SRDIAKNSLF LQMNSLRAED TAVYYCARDR NSHFDYWGGG TLVTVSS 117

SEQ ID NO: 188        moltype = AA length = 107
FEATURE              Location/Qualifiers
REGION              1..107
                      note = Synthetic peptide
source              1..107
                      mol_type = protein
                      organism = synthetic construct

SEQUENCE: 188
DIQMTQSPSS VSASVGDRTV ITCRASQGIN TWLAWYQQKPKAPKLLIYG ASGLQSGVPS 60
RFSGSGSGTD FTLTISSLQP EDFATYYCQQ AKSPFRTEGQ GTKVEIK 107

SEQ ID NO: 189        moltype = AA length = 239
FEATURE              Location/Qualifiers
REGION              1..239
                      note = Synthetic peptide
source              1..239
                      mol_type = protein
                      organism = synthetic construct

SEQUENCE: 189
QVQLVESGGG LVKPGGSLRL SCAASGFTFS DYYMTWIRQA PGKGLEWLSY ISSSGSTIYY 60
ADSVKGRFTI SRDIAKNSLF LQMNSLRAED TAVYYCARDR NSHFDYWGGG TLVTVSSGGG 120
GSGGGGSGGG GSDIQMTQSP SSVASVGDRTV ITTCRASQG INTWLAWYQQ KPGKAPKLLI 180
YGASGLQSGV PSRFSGSGSG TDFTLTISSL QPEDFATYYC QQAQSPRTEF GQGTKVEIK 239

SEQ ID NO: 190        moltype = AA length = 494
FEATURE              Location/Qualifiers
REGION              1..494
                      note = Synthetic peptide
source              1..494
                      mol_type = protein
                      organism = synthetic construct

SEQUENCE: 190
QVQLVESGGG LVKPGGSLRL SCAASGFTFS DYYMTWIRQA PGKGLEWLSY ISSSGSTIYY 60
ADSVKGRFTI SRDIAKNSLF LQMNSLRAED TAVYYCARDR NSHFDYWGGG TLVTVSSGGG 120
GSGGGGSGGG GSDIQMTQSP SSVASVGDRTV ITTCRASQG INTWLAWYQQ KPGKAPKLLI 180
YGASGLQSGV PSRFSGSGSG TDFTLTISSL QPEDFATYYC QQAQSPRTEF GQGTKVEIKS 240
GGGGSEVQLV ESGGGLVQPG GSKLSCAAS GFTFNKYAMN WVRQAPGKGL EWARIRISKY 300
NNYATYYADS VKDRFTISR DSKNTAYLQM NNLKTEDTAV YVCVRHGNFG NSYISYWAYW 360
GQSTLVTVSS GGGGSGGGGS GGGGSGTQVT QEPSTLVSPG GTVTLTCGSS TGAVTSGNYP 420
NWVQKPGQA PRGLIGGTFK LAPGTPARFS GSLGGAAL TLGVPQPEDE AEYYCWLWYS 480
NRWFPGGTFK LTVL 494

SEQ ID NO: 191        moltype = AA length = 5
FEATURE              Location/Qualifiers
REGION              1..5
                      note = Synthetic peptide
source              1..5
                      mol_type = protein
                      organism = synthetic construct

SEQUENCE: 191
DHYS 5

SEQ ID NO: 192        moltype = AA length = 17
FEATURE              Location/Qualifiers
REGION              1..17
                      note = Synthetic peptide
source              1..17
                      mol_type = protein
                      organism = synthetic construct

SEQUENCE: 192
YISSGGIIY YADSVKG 17

SEQ ID NO: 193        moltype = AA length = 8
FEATURE              Location/Qualifiers
REGION              1..8
                      note = Synthetic peptide
source              1..8
                      mol_type = protein
                      organism = synthetic construct

```

-continued

SEQUENCE: 193
DVGSHFDY 8

SEQ ID NO: 194 moltype = AA length = 11
FEATURE Location/Qualifiers
REGION 1..11
note = Synthetic peptide
source 1..11
mol_type = protein
organism = synthetic construct

SEQUENCE: 194
RASQDISRWL A 11

SEQ ID NO: 195 moltype = AA length = 7
FEATURE Location/Qualifiers
REGION 1..7
note = Synthetic peptide
source 1..7
mol_type = protein
organism = synthetic construct

SEQUENCE: 195
AASRLQS 7

SEQ ID NO: 196 moltype = AA length = 9
FEATURE Location/Qualifiers
REGION 1..9
note = Synthetic peptide
source 1..9
mol_type = protein
organism = synthetic construct

SEQUENCE: 196
QQAKSPFRT 9

SEQ ID NO: 197 moltype = AA length = 117
FEATURE Location/Qualifiers
REGION 1..117
note = Synthetic peptide
source 1..117
mol_type = protein
organism = synthetic construct

SEQUENCE: 197
QQQLVESGGG LVKPGGSLRL SCAASGFTFS DHYMSWIRQA PGKGLEWFSY ISSSGGIYY 60
ADSVKGRFTI SRDIAKNSLY LQMNSLRAED TAVYYCARDV GSHFDYWQGQ TLVTVSS 117

SEQ ID NO: 198 moltype = AA length = 107
FEATURE Location/Qualifiers
REGION 1..107
note = Synthetic peptide
source 1..107
mol_type = protein
organism = synthetic construct

SEQUENCE: 198
DIQMTQSPSS VSASVGDRTV ITCRASQDIS RWLAWYQQKPKAPKLLISA ASRLQSGVPS 60
RFGSGSGGTD FTLTISSLQP EDFAIYYCQQ AKSPRTFGQ GTKVEIK 107

SEQ ID NO: 199 moltype = AA length = 239
FEATURE Location/Qualifiers
REGION 1..239
note = Synthetic peptide
source 1..239
mol_type = protein
organism = synthetic construct

SEQUENCE: 199
QQQLVESGGG LVKPGGSLRL SCAASGFTFS DHYMSWIRQA PGKGLEWFSY ISSSGGIYY 60
ADSVKGRFTI SRDIAKNSLY LQMNSLRAED TAVYYCARDV GSHFDYWQGQ TLVTVSSGGG 120
GGGGGGSGGG GSIDIQMTQSP SSVASVGDRTV ITCRASQDIS RWLAWYQQKAPKAPKLLI 180
SAASRLQSGV PSRFGSGSGG TDFTLTISSL QPEDFAIYYC QQAKSPRTFG QGTKVEIK 239

SEQ ID NO: 200 moltype = AA length = 494
FEATURE Location/Qualifiers
REGION 1..494
note = Synthetic peptide
source 1..494
mol_type = protein
organism = synthetic construct

SEQUENCE: 200

QVQLVESGGG	LVKPGGSLRL	SCAASGFTFS	DHYMSWIRQA	PGKGLEWFSY	ISSSGGIYY	60
ADSVKGRFTI	SRDNAKNSLY	LQMNSLRAED	TAVYYCARDV	GSHFDYWGQG	TLVTYSSGGG	120
GSGGGSGGG	GSDIQMTQSP	SSVSASVGD	VTITCRASQD	ISRWLAWYQQ	KPGKAPKLLI	180
SAASRLQSGV	PSRFSGSGSG	TDFTLTISSL	QPEDFAIYYC	QQAKSPFRTF	GQGTKVEIKS	240
GGGGSEVQLV	ESGGGLVQPG	GSLLKSCAAS	GFTFNKYAMN	WVRQAPGKGL	EWVARIRSKY	300
NNYATYYADS	VKDRFTISR	DSKNTAYLQM	NNLKTEDTAV	YYCVRHGNFG	NSYISYWAYW	360
GQGTILTVSS	GGGGSGGGGS	GGGGSQTVVT	QEPSLTVSPG	GTVTLTCGSS	TGAVTSGNYP	420
NWVQKPGQA	PRGLIGGTKF	LAPGTPARFS	GSLLGGKAAL	TLSGVQPEDE	AEYYCVLWYS	480
NRWVFGGGTK	LTVL					494
SEQ ID NO: 201	moltype = AA length = 5					
FEATURE	Location/Qualifiers					
REGION	1..5					
	note = Synthetic peptide					
source	1..5					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 201						
DHYMS						5
SEQ ID NO: 202	moltype = AA length = 17					
FEATURE	Location/Qualifiers					
REGION	1..17					
	note = Synthetic peptide					
source	1..17					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 202						
YISNSGSIYY YDVSVKG						17
SEQ ID NO: 203	moltype = AA length = 8					
FEATURE	Location/Qualifiers					
REGION	1..8					
	note = Synthetic peptide					
source	1..8					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 203						
DVRTAFDY						8
SEQ ID NO: 204	moltype = AA length = 11					
FEATURE	Location/Qualifiers					
REGION	1..11					
	note = Synthetic peptide					
source	1..11					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 204						
RASQSIGSWL A						11
SEQ ID NO: 205	moltype = AA length = 7					
FEATURE	Location/Qualifiers					
REGION	1..7					
	note = Synthetic peptide					
source	1..7					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 205						
AASSLQS						7
SEQ ID NO: 206	moltype = AA length = 9					
FEATURE	Location/Qualifiers					
REGION	1..9					
	note = Synthetic peptide					
source	1..9					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 206						
QQANSFPRT						9
SEQ ID NO: 207	moltype = AA length = 117					
FEATURE	Location/Qualifiers					
REGION	1..117					
	note = Synthetic peptide					
source	1..117					
	mol_type = protein					

-continued

```

                                organism = synthetic construct
SEQUENCE: 207
QVQLVESGGG LVKPGGSLRL SCAASGFTFS DHYMSWIRQA PGKGLEWISY ISNSGSIYY 60
VDSVKGRFTI SRDPAKNSLY LQMNSLRAED TAVYYCARDV RTAFDYWGQG TLVTVSS 117

SEQ ID NO: 208      moltype = AA length = 107
FEATURE            Location/Qualifiers
REGION             1..107
                   note = Synthetic peptide
source             1..107
                   mol_type = protein
                   organism = synthetic construct

SEQUENCE: 208
DIQMTQSPSS VSASVGDVRT ITCRASQSIG SWLAWYQQKP GKAPNLLIYA ASSLQSGVPS 60
RFSGSGSGTD FTLTISSLQP EDFATYICQQ ANSPRTFGQ GTKVDIK 107

SEQ ID NO: 209      moltype = AA length = 239
FEATURE            Location/Qualifiers
REGION             1..239
                   note = Synthetic peptide
source             1..239
                   mol_type = protein
                   organism = synthetic construct

SEQUENCE: 209
QVQLVESGGG LVKPGGSLRL SCAASGFTFS DHYMSWIRQA PGKGLEWISY ISNSGSIYY 60
VDSVKGRFTI SRDPAKNSLY LQMNSLRAED TAVYYCARDV RTAFDYWGQG TLVTVSSGGG 120
GSGGGSGGGG GSDIQMTQSP SSVASVGDRT VTITCRASQS IGSWLAWYQQ KPGKAPNLLI 180
YAASSLQSGV PSRFSGSGSG TDFTLTISSL QPEDFATYYC QQANSPRTF GQGTKVDIK 239

SEQ ID NO: 210      moltype = AA length = 494
FEATURE            Location/Qualifiers
REGION             1..494
                   note = Synthetic peptide
source             1..494
                   mol_type = protein
                   organism = synthetic construct

SEQUENCE: 210
QVQLVESGGG LVKPGGSLRL SCAASGFTFS DHYMSWIRQA PGKGLEWISY ISNSGSIYY 60
VDSVKGRFTI SRDPAKNSLY LQMNSLRAED TAVYYCARDV RTAFDYWGQG TLVTVSSGGG 120
GSGGGSGGGG GSDIQMTQSP SSVASVGDRT VTITCRASQS IGSWLAWYQQ KPGKAPNLLI 180
YAASSLQSGV PSRFSGSGSG TDFTLTISSL QPEDFATYYC QQANSPRTF GQGTKVDIKS 240
GGGSEVQLV ESDDLQVPGG GSKLSCAAS GFTFNKYAMN WVRQAPGKGL EWVARIRSKY 300
NNYATYYADS VKDRFTISR DSKNTAYLQM NNLKTEDTAV YYCVRHGNFG NSVISYWAYW 360
GQGTTLVTSS GGGGSGGGGS GGGGSGTQVT QEPSTLVSPG GTVTLTCGSS TGAVTSGNYP 420
NWVQKPGQA PRGLIGTKF LAPGTPARFS GSSLGKKAAL TSGVQPEDE AEYVCVLWYS 480
NRWVFGGK LTVL 494

SEQ ID NO: 211      moltype = AA length = 5
FEATURE            Location/Qualifiers
REGION             1..5
                   note = Synthetic peptide
source             1..5
                   mol_type = protein
                   organism = synthetic construct

SEQUENCE: 211
SKFMT 5

SEQ ID NO: 212      moltype = AA length = 16
FEATURE            Location/Qualifiers
REGION             1..16
                   note = Synthetic peptide
source             1..16
                   mol_type = protein
                   organism = synthetic construct

SEQUENCE: 212
VIYSGGKTY ADSVKG 16

SEQ ID NO: 213      moltype = AA length = 10
FEATURE            Location/Qualifiers
REGION             1..10
                   note = Synthetic peptide
source             1..10
                   mol_type = protein
                   organism = synthetic construct

SEQUENCE: 213
DSGGWGYFDY 10

```

-continued

SEQ ID NO: 214	moltype = AA	length = 17	
FEATURE	Location/Qualifiers		
REGION	1..17		
	note = Synthetic peptide		
source	1..17		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 214			
KSSQSVLYSS NNKNYLA			17
SEQ ID NO: 215	moltype = AA	length = 7	
FEATURE	Location/Qualifiers		
REGION	1..7		
	note = Synthetic peptide		
source	1..7		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 215			
WASTRES			7
SEQ ID NO: 216	moltype = AA	length = 9	
FEATURE	Location/Qualifiers		
REGION	1..9		
	note = Synthetic peptide		
source	1..9		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 216			
QQYYSTPPT			9
SEQ ID NO: 217	moltype = AA	length = 118	
FEATURE	Location/Qualifiers		
REGION	1..118		
	note = Synthetic peptide		
source	1..118		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 217			
EVQLVESGGG LIQPGGSLRL	SCAVSGFTVS	SKFMTWVRQA	PGKGLEWVSV IYSGGKTYYA 60
DSVKGRFTIS RDNSKNTLYL	QMNSLRAEDT	AVYYCARDSG	GWGYFDYWGG GTLVTVSS 118
SEQ ID NO: 218	moltype = AA	length = 113	
FEATURE	Location/Qualifiers		
REGION	1..113		
	note = Synthetic peptide		
source	1..113		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 218			
DIVMTQSPDS LAVSLGERAT	INCKSSQSVL	YSSNNKNYLA	WYQQKPGQP KLLIYWASTR 60
ESGVPDRFSG SGSGTDFTLT	ISSLQAEDEA	VYYCQQYYST	PPTFGQGTKV EIK 113
SEQ ID NO: 219	moltype = AA	length = 246	
FEATURE	Location/Qualifiers		
REGION	1..246		
	note = Synthetic peptide		
source	1..246		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 219			
EVQLVESGGG LIQPGGSLRL	SCAVSGFTVS	SKFMTWVRQA	PGKGLEWVSV IYSGGKTYYA 60
DSVKGRFTIS RDNSKNTLYL	QMNSLRAEDT	AVYYCARDSG	GWGYFDYWGG GTLVTVSSGG 120
GGSGGGGSGG GGSDIVMTQS	PDSLAVSLGE	RATINCKSSQ	SVLYSSNNKN YLAWYQQKPG 180
QPPKLLIYWA STRESGVDPDR	FSGSGSGTDF	TLTISSLQAE	DVAVVYCCQY YSTPTPTFGQG 240
TKVEIK			246
SEQ ID NO: 220	moltype = AA	length = 501	
FEATURE	Location/Qualifiers		
REGION	1..501		
	note = Synthetic peptide		
source	1..501		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 220			
EVQLVESGGG LIQPGGSLRL	SCAVSGFTVS	SKFMTWVRQA	PGKGLEWVSV IYSGGKTYYA 60

DSVKGRFTIS	RDNSKNTLYL	QMNSLRAEDT	AVYYCARDSDG	GWGYFDYWGQ	CTLVTVSSGG	120
GGSGGGGSGG	GGSDIVMTQS	PDSLAVSLGE	RATINCKSSQ	SVLYSSNNKN	YLAWYQQKPG	180
QPKLLIYWA	STRESGVDR	FSGSGSGTDF	TLTISSLQAE	DVAVYYCQY	YSTPPTFGQG	300
TKVEIKSGGG	GSEVQLVESG	GGLVQPGGSL	KLSCAASGFT	FNKYAMNWVR	QAPGKGLEWV	340
ARIRSKYNNY	ATYYADSVKD	RFTISRDDSK	NTAYLQMNNL	KTEDTAVYYC	VRHGNFGNSY	360
ISYWAYWGQG	TLVTVSSGGG	SGSGGGSGGG	GSQTVVTQEP	SLTVSPGGTV	TLTCGSSTGA	420
VTSGNYPNWV	QQKPGQAPRG	LIGGTGFLAP	GTPARFSGSL	LGGKAALTLS	GVQPEDEAEY	480
YCVLWYSNRW	VFGGGTKLTV	L				501
SEQ ID NO: 221	moltype = AA length = 5					
FEATURE	Location/Qualifiers					
REGION	1..5					
	note = Synthetic peptide					
source	1..5					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 221						
SYIWN						5
SEQ ID NO: 222	moltype = AA length = 16					
FEATURE	Location/Qualifiers					
REGION	1..16					
	note = Synthetic peptide					
source	1..16					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 222						
RIIYNGNTYY	NPSLKS					16
SEQ ID NO: 223	moltype = AA length = 10					
FEATURE	Location/Qualifiers					
REGION	1..10					
	note = Synthetic peptide					
source	1..10					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 223						
PKLGIDAFDI						10
SEQ ID NO: 224	moltype = AA length = 14					
FEATURE	Location/Qualifiers					
REGION	1..14					
	note = Synthetic peptide					
source	1..14					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 224						
TGSSSNIGAG	YDVH					14
SEQ ID NO: 225	moltype = AA length = 7					
FEATURE	Location/Qualifiers					
REGION	1..7					
	note = Synthetic peptide					
source	1..7					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 225						
GNSNRPS						7
SEQ ID NO: 226	moltype = AA length = 11					
FEATURE	Location/Qualifiers					
REGION	1..11					
	note = Synthetic peptide					
source	1..11					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 226						
QSYDSSLSGW	V					11
SEQ ID NO: 227	moltype = AA length = 118					
FEATURE	Location/Qualifiers					
REGION	1..118					
	note = Synthetic peptide					
source	1..118					
	mol_type = protein					
	organism = synthetic construct					

-continued

SEQUENCE: 227
 QVQLQESGPG LVKPSSETLSL TCTVSGGSIS SYYNWIRQP PGKGLEWIGR IYYNGNTYYN 60
 PSLKSRVTIS GDTSKNQFSL KLSSVTAADT AVYYCARPKL GIDAFDIWGQ GTMVTVSS 118

SEQ ID NO: 228 moltype = AA length = 111
 FEATURE Location/Qualifiers
 REGION 1..111
 note = Synthetic peptide
 source 1..111
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 228
 QSVLTQPPSV SGAPGQRTVI SCTGSSSNIG AGYDVHWYQQ LPGTAPKLLI YGNSNRPSGV 60
 PDRFSGSKSG TSASLAITGL QAEDEADYYC QSYDSSLGWW VFGGGTKLTV L 111

SEQ ID NO: 229 moltype = AA length = 244
 FEATURE Location/Qualifiers
 REGION 1..244
 note = Synthetic peptide
 source 1..244
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 229
 QVQLQESGPG LVKPSSETLSL TCTVSGGSIS SYYNWIRQP PGKGLEWIGR IYYNGNTYYN 60
 PSLKSRVTIS GDTSKNQFSL KLSSVTAADT AVYYCARPKL GIDAFDIWGQ GTMVTVSSGG 120
 GSGGGGSGG GGSQSVLTQP PSVSGAPGQR VTISCTGSSS NIGAGYDVHW YQQLPGTAPK 180
 LLIYGNSNRP SGVPDRFSGS KSGTSASLAI TGLQAEDEAD YYCQSYDSSL SGWVFGGGTK 240
 LTVL 244

SEQ ID NO: 230 moltype = AA length = 499
 FEATURE Location/Qualifiers
 REGION 1..499
 note = Synthetic peptide
 source 1..499
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 230
 QVQLQESGPG LVKPSSETLSL TCTVSGGSIS SYYNWIRQP PGKGLEWIGR IYYNGNTYYN 60
 PSLKSRVTIS GDTSKNQFSL KLSSVTAADT AVYYCARPKL GIDAFDIWGQ GTMVTVSSGG 120
 GSGGGGSGG GGSQSVLTQP PSVSGAPGQR VTISCTGSSS NIGAGYDVHW YQQLPGTAPK 180
 LLIYGNSNRP SGVPDRFSGS KSGTSASLAI TGLQAEDEAD YYCQSYDSSL SGWVFGGGTK 240
 LTVLSGGGGS EVQLVESGGG LVQPGGSLKL SCAASGFTFN KYAMNWVRQA PGKGLEWVAR 300
 IRSKYNRYAT YYADSVKDRF TISRDDSNT AYLMNNLKT EDTAVYYCVR HGNTFGNSYIS 360
 YWAYWGQGTI VTVSSGGGGS GGGGGGGGGS QTVVTQEPSL TVSPGGTVTL TCGSSTGAVT 420
 SGNYPNWVQQ KPGQAPRGLI GGTKFLAPGT PARFSGSLLG GKAAALTLGV QPEDEAEYYC 480
 VLWYSNRWVF GGGTKLTVL 499

SEQ ID NO: 231 moltype = AA length = 622
 FEATURE Location/Qualifiers
 source 1..622
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 231
 MALPTARPLL GSCGTPALGS LLFLFLSLGW VQPSRTLAGE TGQEAAPLDG VLANPPNISS 60
 LSPRQLLGFP CAEVSGLSTE RVRELAVALA QKNVKLSTEQ LRCLAHRLSE PPEDLDALPL 120
 DLLLFLNPDA FSGPQACTRF FSRITKANVD LLPRGAPERQ RLLPAALACW GVRGSLSEA 180
 DVALGGLAC DLPGRFVAES AEVLLPRLVS CPGPLDQDQQ EAARAALQGG GPPYGPSTW 240
 SVSTMALRG LLPVLGQPII RSIPQGIVAA WRQRSSRDPS WRQPRTILR PRFRREVEKT 300
 ACPGSGKKARE IDESLIFYKK WELEACVDAA LLATQMDRVN AIPFTYEQLD VLKHKLDELY 360
 PQGYPESVIQ HLGYLFLKMS PEDIRKWNVT SLETLKALLE VNKGHEMSPQ VATLIDRFVK 420
 GRGQLDKDTL DTLTAFYPGY LCSSLPEELS SVPPSSIWAV RPQDLDTCDP RQLDVLVYKA 480
 RLAFQNMNGS EYFVKIQSFL GGAPTEDLKA LSQQNVSMDL ATFMKLRTDA VLPLTVAEVQ 540
 KLLGPHVEGL KAERHRFVR DWILRQRQDD LDTLGLGLQG GIPNGYLVLD LSMQEALSGT 600
 PCLLGPGPVL TVLALLLAST LA 622

SEQ ID NO: 232 moltype = AA length = 630
 FEATURE Location/Qualifiers
 source 1..630
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 232
 MALPTARPLL GSCGTPALGS LLFLFLSLGW VQPSRTLAGE TGQEAAPLDG VLANPPNISS 60
 LSPRQLLGFP CAEVSGLSTE RVRELAVALA QKNVKLSTEQ LRCLAHRLSE PPEDLDALPL 120
 DLLLFLNPDA FSGPQACTRF FSRITKANVD LLPRGAPERQ RLLPAALACW GVRGSLSEA 180
 DVALGGLAC DLPGRFVAES AEVLLPRLVS CPGPLDQDQQ EAARAALQGG GPPYGPSTW 240
 SVSTMALRG LLPVLGQPII RSIPQGIVAA WRQRSSRDPS WRQPRTILR PRFRREVEKT 300

-continued

ACPSGKKARE	IDESLIFYKK	WELEACVDAA	LLATQMDRVN	AIPPTYEQLD	VLKHKLDLY	360
PQGYPESVIQ	HLGYLFLKMS	PEDIRKWNVT	SLETLKALLE	VNKGHEMSPQ	APRRPLPQVA	420
TLIDRFVKGR	GQLDKDTLDT	LTAFYPGYLC	SLSPHEELSSV	PPSSIWAVRP	QDLDTCDPRQ	480
LDVLYPKARL	AFQNMNGSEY	FVKIQSFLGG	APTEDLKALS	QQNVSMDLAT	FMKLRTDAVL	540
PLTVAEVQKL	LGPVHEGLKA	EERHRPVRDW	ILRQRQDDLD	TLGLGLQGGI	PNGYLVLDLS	600
MQEALSGTFC	LLGPGPVLTV	LALLLASTLA				630

SEQ ID NO: 233 moltype = AA length = 622
 FEATURE Location/Qualifiers
 source 1..622
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 233

MALPTARPLL	GSCGTPALGS	LLFLFLSLGW	VQPSRTLAGE	TGQEAAPLDG	VLANPPNISS	60
LSPRQLLGFP	CAEVSGSLSTE	RVRELAVALA	QKNVKLSTEQ	LRCLAHRLSE	PPEDLDALPL	120
DLLLFLNPDA	FSGPQACTHF	FSRITKANVD	LLPRGAPERQ	RLLPAALACW	GVRGSLSEA	180
DVRALGGLAC	DLPGRFVAES	AEVLLPRLVS	CPGPLDQDQQ	EAARAALQGG	GPPYGPSTW	240
SVSTMDALRG	LLPVLGQPII	RSIPQGIVAA	WRQRSSRDPS	WRQPRTLILR	PRFRREVEKT	300
ACPSGKKARE	IDESLIFYKK	WELEACVDAA	LLATQMDRVN	AIPPTYEQLD	VLKHKLDLY	360
PQGYPESVIQ	HLGYLFLKMS	PEDIRKWNVT	SLETLKALLE	VNKGHEMSPQ	VATLIDRFVK	420
GRGQLDKDTL	DTLTAIFYPGY	LCSLSPEELS	SVPPSSIWAV	RPQDLDTCDP	RQLDVLYPKA	480
RLAFQNMNGS	EYFVKIQSFL	GGAPTEDLKA	LSQQNVSMDL	ATFMKLRTDA	VLPLTVAEVQ	540
KLLGPHVEGL	KAEERHRPVR	DWILRQRQDD	LDTLGLGLQG	GIPNGYLVLD	LSVQEALSGT	600
PCLLGGPVL	TVLALLLAST	LA				622

SEQ ID NO: 234 moltype = AA length = 622
 FEATURE Location/Qualifiers
 source 1..622
 mol_type = protein
 organism = Macaca fascicularis

SEQUENCE: 234

MALPMARPLS	GSCGTPALGS	LLFLFLSLGW	VQPSRVLAGL	TRQEAAPLDG	ILTNAPDIAS	60
LSPRQLLGFT	CVEVSGSLSTE	LVQELAVALL	QKNVKLSAEQ	LRCLAHRLSE	PPEDLDALPL	120
DLLLFLNPDA	FSGPQACTHF	FSRVAKANVD	LLPRGAPERQ	RLLPAALTCW	GVRGSLSEA	180
DVRALGGLAC	DLPGRFVAES	AEVVLPRLVR	CLGPLDQDQQ	EAARAALQRG	GPPYGPSTW	240
SISTLDDLQS	LLPVLGQPII	HSIPQGILAA	WRQRSSRDPS	WQQPEQTVLR	PRFRRDVERT	300
TCPPEKEVHE	IDENLIFYKK	RELEACVDAA	LLAAQMDRVD	AIPPTYEQLD	VLKHKLDLY	360
PQGYPESVIR	HLGHLFLKMS	PEDIRKWNVT	SLETLKALLK	VSKGHEMSAQ	VATLIDRVVV	420
GRGQLDKDTA	DTLTAFCPGC	LCSLSPERLS	SVPPSVIGAV	RPQDLDTCPG	RQLDVLYPKA	480
RLAFQNMNGS	EYFVKIRPFL	GGAPTEDLKA	LSQQNVSMDL	ATFMKLRRRA	VLPLTVAEVQ	540
KLLGPHVEGL	KVEEQHPVR	DWILKQRQDD	LDTLGLGLQG	GIPNGYLILD	LSVREALSGT	600
PCLLGGPVL	TILALLLAST	LA				622

SEQ ID NO: 235 moltype = AA length = 286
 FEATURE Location/Qualifiers
 source 1..286
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 235

MALPTARPLL	GSCGTPALGS	LLFLFLSLGW	VQPSRTLAGE	TGQEAAPLDG	VLANPPNISS	60
LSPRQLLGFP	CAEVSGSLSTE	RVRELAVALA	QKNVKLSTEQ	LRCLAHRLSE	PPEDLDALPL	120
DLLLFLNPDA	FSGPQACTRF	FSRITKANVD	LLPRGAPERQ	RLLPAALACW	GVRGSLSEA	180
DVRALGGLAC	DLPGRFVAES	AEVLLPRLVS	CPGPLDQDQQ	EAARAALQGG	GPPYGPSTW	240
SVSTMDALRG	LLPVLGQPII	RSIPQGIVAA	WRQRSSRDPS	WRQPER		286

SEQ ID NO: 236 moltype = AA length = 303
 FEATURE Location/Qualifiers
 source 1..303
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 236

EVEKTACPSG	KKAREIDESL	IFYKKWELEA	CVDAALLATQ	MDRVNAIPFT	YEQLDVLKHK	60
LDELYPQGYP	ESVIQHLGYL	FLKMSPEDIR	KWNVTSLETL	KALLEVNKGH	EMSPQVATLI	120
DRFVKGRGQL	DKDTLDTLTA	FYPGYLCSLS	PEELSSVPPS	SIWAVRPQDL	DTCDPRQLDV	180
LYPKARLAFQ	NMNGSEYFVK	IQSFLGGAPT	EDLKALSQQN	VSMDLATFMK	LRTDAVLPLT	240
VAEVQKLLGP	HVEGLKAEER	HRPVRDWILR	QRQDDLTGL	LGLQGGIPNG	YLVLDLSMQE	300
ALS						303

SEQ ID NO: 237 moltype = AA length = 24
 FEATURE Location/Qualifiers
 source 1..24
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 237

GTPCLLGGP	VLTVLALLLA	STLA				24
-----------	------------	------	--	--	--	----

-continued

SEQ ID NO: 238	moltype = AA	length = 51	
FEATURE	Location/Qualifiers		
source	1..51		
	mol_type = protein		
	organism = Homo sapiens		
SEQUENCE: 238			
EVEKTACPSG KKAREIDESL IFYKKWELEA CVDAALLATQ MDRVNAIPFT Y			51
SEQ ID NO: 239	moltype = AA	length = 38	
FEATURE	Location/Qualifiers		
source	1..38		
	mol_type = protein		
	organism = Homo sapiens		
SEQUENCE: 239			
EQLDVLKHKL DELYPQGYE SVIQHLGYLF LKMSPEDI			38
SEQ ID NO: 240	moltype = AA	length = 69	
FEATURE	Location/Qualifiers		
source	1..69		
	mol_type = protein		
	organism = Homo sapiens		
SEQUENCE: 240			
RKWNVTSLT LKALLEVNKG HEMSPQVATL IDRFVKGRGQ LDKDTLDTLT AFYPGYLCSL			60
SPEELSSVP			69
SEQ ID NO: 241	moltype = AA	length = 48	
FEATURE	Location/Qualifiers		
source	1..48		
	mol_type = protein		
	organism = Homo sapiens		
SEQUENCE: 241			
PSSIWAVRPQ DLDTC DPRQL DVLYPKARLA FQNMNGSEYF VKIQSFLG			48
SEQ ID NO: 242	moltype = AA	length = 44	
FEATURE	Location/Qualifiers		
source	1..44		
	mol_type = protein		
	organism = Homo sapiens		
SEQUENCE: 242			
GAPTEDLKAL SQQNVSMDLA TFMKLRTDAV LPLTVAEVQK LLGP			44
SEQ ID NO: 243	moltype = AA	length = 52	
FEATURE	Location/Qualifiers		
source	1..52		
	mol_type = protein		
	organism = Homo sapiens		
SEQUENCE: 243			
HVEGLKAEER HRPVRDWILR QRQDDLDTLG LGLQGGIPNG YLVLDLSMQE AL			52
SEQ ID NO: 244	moltype = AA	length = 89	
FEATURE	Location/Qualifiers		
source	1..89		
	mol_type = protein		
	organism = Homo sapiens		
SEQUENCE: 244			
EVEKTACPSG KKAREIDESL IFYKKWELEA CVDAALLATQ MDRVNAIPFT YEQLDVLKHK			60
LDELYPQGYE SVIQHLGYL FLKMSPEDI			89
SEQ ID NO: 245	moltype = AA	length = 107	
FEATURE	Location/Qualifiers		
source	1..107		
	mol_type = protein		
	organism = Homo sapiens		
SEQUENCE: 245			
EQLDVLKHKL DELYPQGYE SVIQHLGYLF LKMSPEDIRK WNVTSLET LK ALLEVNKGHE			60
MSPQVATLID RFVKGRGQLD KDTLDTLTAF YPGYLCSLSP EELSSVP			107
SEQ ID NO: 246	moltype = AA	length = 622	
FEATURE	Location/Qualifiers		
REGION	1..622		
	note = Synthetic peptide		
source	1..622		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 246			
MALPTARPLL GSCGTPALGS LLFLFLSLGW VQPSRTLAGE TGQEAAPLDG VLANPPNISS			60

-continued

```

LSPRQLLGFP CAEVSGSLSTE RVRELAVALA QKNVKLSTEQ LRCLAHRLSE PPEDLDALPL 120
DLLLLFLNPDA FSGPQACTRF FSRITKANVD LLPRGAPERQ RLLPAALACW GVRGSLSEA 180
DVRALGGLAC DLPGRFVAES AEVLLPRLVS CPGPLDQDQQ EAARAALQGG GPPYGGPSTW 240
SVSTMALRG LLPVLGQPII RSIPQGIVAA WRQSSRDPS WRQPRTLILR PRFRDAEQK 300
ACPPGKEPYK VDEDLIFYQN WELEACVDGT MLARQMDLVN EIPFTYEQLD VLKHKLDLEY 360
PQGYPESVIQ HLGYLFLKMS PEDIRKWNVT SLETLKALLE VNKGHEMSPQ VATLIDRFVK 420
GRGQLDKDTL DTLTAFYPGY LCSLSPEELS SVPPSSIWAV RPQDLDTCDP RQLDVLYPKA 480
RLAFQNMNGS EYFVKIQSFL GGAPTEDLKA LSQQNVSM DL ATFMKLRTDA VLPLTVAEVQ 540
KLLGPHVEGL KAEERHRPVR DWILRQRQDD LDTLGLGLQG GIPNGYLVLD LSMQEALSGT 600
PCLLGGPVL TVLALLLAST LA 622

```

```

SEQ ID NO: 247      moltype = AA length = 622
FEATURE            Location/Qualifiers
REGION             1..622
                   note = Synthetic peptide
source             1..622
                   mol_type = protein
                   organism = synthetic construct

```

```

SEQUENCE: 247
MALPTARPLL GSCGTPALGS LLFLFLSLGW VQPSRTLAGE TGQEAAPLDG VLANPPNISS 60
LSPRQLLGFP CAEVSGSLSTE RVRELAVALA QKNVKLSTEQ LRCLAHRLSE PPEDLDALPL 120
DLLLLFLNPDA FSGPQACTRF FSRITKANVD LLPRGAPERQ RLLPAALACW GVRGSLSEA 180
DVRALGGLAC DLPGRFVAES AEVLLPRLVS CPGPLDQDQQ EAARAALQGG GPPYGGPSTW 240
SVSTMALRG LLPVLGQPII RSIPQGIVAA WRQSSRDPS WRQPRTLILR PRFRREVEKT 300
ACPSGKKARE IDESLIFYKK WELEACVDAA LLATQMDRVN AIPFTYEQLD IFKHKLDKTY 360
PQGYPESLIQ QLGHHFRYVS PEDIRKWNVT SLETLKALLE VNKGHEMSPQ VATLIDRFVK 420
GRGQLDKDTL DTLTAFYPGY LCSLSPEELS SVPPSSIWAV RPQDLDTCDP RQLDVLYPKA 480
RLAFQNMNGS EYFVKIQSFL GGAPTEDLKA LSQQNVSM DL ATFMKLRTDA VLPLTVAEVQ 540
KLLGPHVEGL KAEERHRPVR DWILRQRQDD LDTLGLGLQG GIPNGYLVLD LSMQEALSGT 600
PCLLGGPVL TVLALLLAST LA 622

```

```

SEQ ID NO: 248      moltype = AA length = 622
FEATURE            Location/Qualifiers
REGION             1..622
                   note = Synthetic peptide
source             1..622
                   mol_type = protein
                   organism = synthetic construct

```

```

SEQUENCE: 248
MALPTARPLL GSCGTPALGS LLFLFLSLGW VQPSRTLAGE TGQEAAPLDG VLANPPNISS 60
LSPRQLLGFP CAEVSGSLSTE RVRELAVALA QKNVKLSTEQ LRCLAHRLSE PPEDLDALPL 120
DLLLLFLNPDA FSGPQACTRF FSRITKANVD LLPRGAPERQ RLLPAALACW GVRGSLSEA 180
DVRALGGLAC DLPGRFVAES AEVLLPRLVS CPGPLDQDQQ EAARAALQGG GPPYGGPSTW 240
SVSTMALRG LLPVLGQPII RSIPQGIVAA WRQSSRDPS WRQPRTLILR PRFRREVEKT 300
ACPSGKKARE IDESLIFYKK WELEACVDAA LLATQMDRVN AIPFTYEQLD VLKHKLDLEY 360
PQGYPESVIQ HLGYLFLKMS PEDIHQNVNT SPDTVKTLK VSKGQKMNAQ AIALVACYLR 420
GGGQLDEDMV KALGDIPLSY LCDFSQDLH SVPPSSIWAV RPQDLDTCDP RQLDVLYPKA 480
RLAFQNMNGS EYFVKIQSFL GGAPTEDLKA LSQQNVSM DL ATFMKLRTDA VLPLTVAEVQ 540
KLLGPHVEGL KAEERHRPVR DWILRQRQDD LDTLGLGLQG GIPNGYLVLD LSMQEALSGT 600
PCLLGGPVL TVLALLLAST LA 622

```

```

SEQ ID NO: 249      moltype = AA length = 622
FEATURE            Location/Qualifiers
REGION             1..622
                   note = Synthetic peptide
source             1..622
                   mol_type = protein
                   organism = synthetic construct

```

```

SEQUENCE: 249
MALPTARPLL GSCGTPALGS LLFLFLSLGW VQPSRTLAGE TGQEAAPLDG VLANPPNISS 60
LSPRQLLGFP CAEVSGSLSTE RVRELAVALA QKNVKLSTEQ LRCLAHRLSE PPEDLDALPL 120
DLLLLFLNPDA FSGPQACTRF FSRITKANVD LLPRGAPERQ RLLPAALACW GVRGSLSEA 180
DVRALGGLAC DLPGRFVAES AEVLLPRLVS CPGPLDQDQQ EAARAALQGG GPPYGGPSTW 240
SVSTMALRG LLPVLGQPII RSIPQGIVAA WRQSSRDPS WRQPRTLILR PRFRREVEKT 300
ACPSGKKARE IDESLIFYKK WELEACVDAA LLATQMDRVN AIPFTYEQLD VLKHKLDLEY 360
PQGYPESVIQ HLGYLFLKMS PEDIRKWNVT SLETLKALLE VNKGHEMSPQ VATLIDRFVK 420
GRGQLDKDTL DTLTAFYPGY LCSLSPEELS SVPPSSMWLV GPQDLDKCSQ RHLGLLYQKA 480
CSAFQNVSGL EYFEKIKTFL GGAPTEDLKA LSQQNVSM DL ATFMKLRTDA VLPLTVAEVQ 540
KLLGPHVEGL KAEERHRPVR DWILRQRQDD LDTLGLGLQG GIPNGYLVLD LSMQEALSGT 600
PCLLGGPVL TVLALLLAST LA 622

```

```

SEQ ID NO: 250      moltype = AA length = 622
FEATURE            Location/Qualifiers
REGION             1..622
                   note = Synthetic peptide
source             1..622

```

-continued

mol_type = protein
organism = synthetic construct

SEQUENCE: 250

MALPTARPLL	GSCGTPALGS	LLFLLFSLGW	VQPSRTLAGE	TGQEAAPLDG	VLANPPNISS	60
LSPRQLLGFP	CAEVSGLSTE	RVRELAVALA	QKNVKLSTEQ	LRCLAHRLSE	PPEDLDALPL	120
DLLLFLNPDA	FSGPQACTRF	FSRITKANVD	LLPRGAPERQ	RLLPAALACW	GVRGSLSEA	180
DVRALGGLAC	DLPGRFVAES	AEVLLPRLVS	CPGPLDQDQQ	EAARAALQGG	GPPYGPPSTW	240
SVSTMDALRG	LLPVLGQPII	RSIPQGIVAA	WRQSSRDPS	WRQPRTLILR	PRFRREVEKT	300
ACPSGKKARE	IDESLIFYKK	WELEACVDAA	LLATQMDRVN	AIPFTYEQLD	VLKHKLDLEY	360
PQGYPESVIQ	HLGYLFLKMS	PEDIRKWNVT	SLETLKALLE	VNKGHEMSQP	VATLIDRFVK	420
GRGQLDKDTL	DTLTAFYPGY	LCSLSPEELS	SVPPSSIWAV	RPQDLDTCDP	RQLDVLYPKA	480
RLAFQNMNGS	EYFVKIQSFL	GGASVKDLRA	LSQHNVSMDI	ATFKRLQVDS	LVGLSVAEVQ	540
KLLGPHVEGL	KAERHRPVR	DWILRQRQDD	LDTLGLGLQG	GIPNGYLVLD	LSMQEALSGT	600
PCLLGPGPVL	TVLALLLAST	LA				622

SEQ ID NO: 251 moltype = AA length = 622
FEATURE Location/Qualifiers
REGION 1..622
 note = Synthetic peptide
source 1..622
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 251

MALPTARPLL	GSCGTPALGS	LLFLLFSLGW	VQPSRTLAGE	TGQEAAPLDG	VLANPPNISS	60
LSPRQLLGFP	CAEVSGLSTE	RVRELAVALA	QKNVKLSTEQ	LRCLAHRLSE	PPEDLDALPL	120
DLLLFLNPDA	FSGPQACTRF	FSRITKANVD	LLPRGAPERQ	RLLPAALACW	GVRGSLSEA	180
DVRALGGLAC	DLPGRFVAES	AEVLLPRLVS	CPGPLDQDQQ	EAARAALQGG	GPPYGPPSTW	240
SVSTMDALRG	LLPVLGQPII	RSIPQGIVAA	WRQSSRDPS	WRQPRTLILR	PRFRREVEKT	300
ACPSGKKARE	IDESLIFYKK	WELEACVDAA	LLATQMDRVN	AIPFTYEQLD	VLKHKLDLEY	360
PQGYPESVIQ	HLGYLFLKMS	PEDIRKWNVT	SLETLKALLE	VNKGHEMSQP	VATLIDRFVK	420
GRGQLDKDTL	DTLTAFYPGY	LCSLSPEELS	SVPPSSIWAV	RPQDLDTCDP	RQLDVLYPKA	480
RLAFQNMNGS	EYFVKIQSFL	GGAPTEDLKA	LSQQNVSMDL	ATFMKLRDA	VLPLTVAEVQ	540
KLLGPNIVDL	KTEEDKSPVR	DWLFQRQHKD	LDRLGLGLQG	GIPNGYLVLD	FNVREAFSGT	600
PCLLGPGPVL	TVLALLLAST	LA				622

SEQ ID NO: 252 moltype = AA length = 227
FEATURE Location/Qualifiers
REGION 1..227
 note = Fc monomer-1
source 1..227
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 252

DKTHTCPPCP	APELLGGPSV	FLFPPKPKDT	LMISRTPEVT	CVVVDVSHED	PEVKFNWYVD	60
GVEVHNAKTK	PCEEQYGSTY	RCVSVLTVLH	QDWLNGKEYK	CKVSNKALPA	PIEKTISKAK	120
GQPREPQVYT	LPPSREEMTK	NQVSLTCLVK	GFYPDSIAVE	WESNGQPENN	YKTTTPVLDS	180
DGSFFLYSKL	TVDKSRWQQG	NVFSQSVSMHE	ALHNHYTQKS	LSLSPGK		227

SEQ ID NO: 253 moltype = AA length = 225
FEATURE Location/Qualifiers
REGION 1..225
 note = Fc monomer-2
source 1..225
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 253

DKTHTCPPCP	APELLGGPSV	FLFPPKPKDT	LMISRTPEVT	CVVVDVSHED	PEVKFNWYVD	60
GVEVHNAKTK	PCEEQYGSTY	RCVSVLTVLH	QDWLNGKEYK	CKVSNKALPA	PIEKTISKAK	120
GQPREPQVYT	LPPSREEMTK	NQVSLTCLVK	GFYPDSIAVE	WESNGQPENN	YKTTTPVLDS	180
DGSFFLYSKL	TVDKSRWQQG	NVFSQSVSMHE	ALHNHYTQKS	LSLSPGK		225

SEQ ID NO: 254 moltype = AA length = 227
FEATURE Location/Qualifiers
REGION 1..227
 note = Fc monomer-3
source 1..227
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 254

DKTHTCPPCP	APELLGGPSV	FLFPPKPKDT	LMISRTPEVT	CVVVDVSHED	PEVKFNWYVD	60
GVEVHNAKTK	PREEQYNSTY	RVVSVLTVLH	QDWLNGKEYK	CKVSNKALPA	PIEKTISKAK	120
GQPREPQVYT	LPPSREEMTK	NQVSLTCLVK	GFYPDSIAVE	WESNGQPENN	YKTTTPVLDS	180
DGSFFLYSKL	TVDKSRWQQG	NVFSQSVSMHE	ALHNHYTQKS	LSLSPGK		227

SEQ ID NO: 255 moltype = AA length = 225
FEATURE Location/Qualifiers

-continued

REGION 1..225
 note = Fc monomer-4
 source 1..225
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 255
 DKTHTCPPCP APELLGGPSV FLFPPKPKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD 60
 GVEVHNAKTK PREEQYNSTY RVVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK 120
 GQPREPQVYT LPPSREEMTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTPPVLDL 180
 DGSFFLYSKL TVDKSRWQGG NVFSCSV MHE ALHNHYTQKS LSLSP 225

SEQ ID NO: 256 moltype = AA length = 227
 FEATURE Location/Qualifiers
 REGION 1..227
 note = Fc monomer-5
 source 1..227
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 256
 DKTHTCPPCP APELLGGPSV FLFPPKPKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD 60
 GVEVHNAKTK PREEQYGSTY RVVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK 120
 GQPREPQVYT LPPSREEMTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTPPVLDL 180
 DGSFFLYSKL TVDKSRWQGG NVFSCSV MHE ALHNHYTQKS LSLSPGK 227

SEQ ID NO: 257 moltype = AA length = 225
 FEATURE Location/Qualifiers
 REGION 1..225
 note = Fc monomer-6
 source 1..225
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 257
 DKTHTCPPCP APELLGGPSV FLFPPKPKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD 60
 GVEVHNAKTK PREEQYGSTY RVVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK 120
 GQPREPQVYT LPPSREEMTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTPPVLDL 180
 DGSFFLYSKL TVDKSRWQGG NVFSCSV MHE ALHNHYTQKS LSLSP 225

SEQ ID NO: 258 moltype = AA length = 227
 FEATURE Location/Qualifiers
 REGION 1..227
 note = Fc monomer-7
 source 1..227
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 258
 DKTHTCPPCP APELLGGPSV FLFPPKPKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD 60
 GVEVHNAKTK PCEEQYNSTY RCVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK 120
 GQPREPQVYT LPPSREEMTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTPPVLDL 180
 DGSFFLYSKL TVDKSRWQGG NVFSCSV MHE ALHNHYTQKS LSLSPGK 227

SEQ ID NO: 259 moltype = AA length = 225
 FEATURE Location/Qualifiers
 REGION 1..225
 note = Fc monomer-8
 source 1..225
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 259
 DKTHTCPPCP APELLGGPSV FLFPPKPKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD 60
 GVEVHNAKTK PCEEQYNSTY RCVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK 120
 GQPREPQVYT LPPSREEMTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTPPVLDL 180
 DGSFFLYSKL TVDKSRWQGG NVFSCSV MHE ALHNHYTQKS LSLSP 225

SEQ ID NO: 260 moltype = AA length = 484
 FEATURE Location/Qualifiers
 REGION 1..484
 note = scFc-1
 source 1..484
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 260
 DKTHTCPPCP APELLGGPSV FLFPPKPKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD 60
 GVEVHNAKTK PCEEQYGSTY RCVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK 120
 GQPREPQVYT LPPSREEMTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTPPVLDL 180
 DGSFFLYSKL TVDKSRWQGG NVFSCSV MHE ALHNHYTQKS LSLSPGKGGG GSGGGGSGGG 240
 GSGGGGSGGG GSGGGGSDKT HTCPCPAPE LLGGPSVFLF PPKKDTLMI SRTPEVTCVV 300

-continued

VDVSHEDPEV	KFNWYVDGVE	VHNAKTKPCE	EQYGSTYRCV	SVLTVLHQDW	LNGKEYKCKV	360
SNKALPAPIE	KTISKAKGQP	REPQVYTLPP	SREEMTKNQV	SLTCLVKGFY	PSDIAVEWES	420
NGQPENNYKT	TPPVLDSGDS	FFLYSKLTVD	KSRWQQGNVF	SCSVMHEALH	NHYTQKSLSL	480
SPGK						484

```

SEQ ID NO: 261      moltype = AA  length = 480
FEATURE            Location/Qualifiers
REGION             1..480
                   note = scFc-2
source             1..480
                   mol_type = protein
                   organism = synthetic construct

```

```

SEQUENCE: 261
DKTHTCPPCP APELLGGPSV FLFPPKPKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD 60
GVEVHNAKTK PCEEQYGSTY RCVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK 120
GQPREPQVYT LPPSREEMTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTPPVLD 180
DGSFFLYSKL TVDKSRWQQG NVFSCSVMHE ALHNHYTQKS LSLSPGGGGS GGGGSGGGGS 240
GGGGSGGGGS GGGGSDKTHT CPPCPAPELL GGPSVFLFPP KPKDTLMISR TPEVTCVVVD 300
VSHEDPEVKF NWWYVDGVEVH NAKTKPCEEQ YGSTRYCVSV LTVLHQDWLN GKEYKCKVSN 360
KALPAPIEKT ISKAKGQPRE PQVYTLPPSR EEMTKNQVSL TCLVKGFYPS DIAVEWESNG 420
QPENNYKTTT PVLDSGDSFF LYSKLTVDKS RWQQGNVFSC SVMHEALHNNH YTQKSLSLSP 480

```

```

SEQ ID NO: 262      moltype = AA  length = 484
FEATURE            Location/Qualifiers
REGION             1..484
                   note = scFc-3
source             1..484
                   mol_type = protein
                   organism = synthetic construct

```

```

SEQUENCE: 262
DKTHTCPPCP APELLGGPSV FLFPPKPKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD 60
GVEVHNAKTK PREEQYNSTY RVVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK 120
GQPREPQVYT LPPSREEMTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTPPVLD 180
DGSFFLYSKL TVDKSRWQQG NVFSCSVMHE ALHNHYTQKS LSLSPGKGGG GSGGGSGGGG 240
GSGGGSGGGG GSGGGGSDKT HTCPPCPAPE LLGGPSVFLF PPKPKDTLMI SRTPEVTCVV 300
VDVSHEDPEV KFNWYVDGVE VHNAKTKPRE EQYNSTYRVV SVLTVLHQDW LNGKEYKCKV 360
SNKALPAPIE KTISKAKGQP REPQVYTLPP SREEMTKNQV SLTCLVKGFY PSDIAVEWES 420
NGQPENNYKT TPPVLDSGDS FFLYSKLTVD KSRWQQGNVF SCSVMHEALH NHYTQKSLSL 480
SPGK

```

```

SEQ ID NO: 263      moltype = AA  length = 480
FEATURE            Location/Qualifiers
REGION             1..480
                   note = scFc-4
source             1..480
                   mol_type = protein
                   organism = synthetic construct

```

```

SEQUENCE: 263
DKTHTCPPCP APELLGGPSV FLFPPKPKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD 60
GVEVHNAKTK PREEQYNSTY RVVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK 120
GQPREPQVYT LPPSREEMTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTPPVLD 180
DGSFFLYSKL TVDKSRWQQG NVFSCSVMHE ALHNHYTQKS LSLSPGGGGS GGGGSGGGGS 240
GGGGSGGGGS GGGGSDKTHT CPPCPAPELL GGPSVFLFPP KPKDTLMISR TPEVTCVVVD 300
VSHEDPEVKF NWWYVDGVEVH NAKTKPREEQ YNSTYRVVSV LTVLHQDWLN GKEYKCKVSN 360
KALPAPIEKT ISKAKGQPRE PQVYTLPPSR EEMTKNQVSL TCLVKGFYPS DIAVEWESNG 420
QPENNYKTTT PVLDSGDSFF LYSKLTVDKS RWQQGNVFSC SVMHEALHNNH YTQKSLSLSP 480

```

```

SEQ ID NO: 264      moltype = AA  length = 484
FEATURE            Location/Qualifiers
REGION             1..484
                   note = scFc-5
source             1..484
                   mol_type = protein
                   organism = synthetic construct

```

```

SEQUENCE: 264
DKTHTCPPCP APELLGGPSV FLFPPKPKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD 60
GVEVHNAKTK PREEQYGSTY RVVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK 120
GQPREPQVYT LPPSREEMTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTPPVLD 180
DGSFFLYSKL TVDKSRWQQG NVFSCSVMHE ALHNHYTQKS LSLSPGKGGG GSGGGSGGGG 240
GSGGGSGGGG GSGGGGSDKT HTCPPCPAPE LLGGPSVFLF PPKPKDTLMI SRTPEVTCVV 300
VDVSHEDPEV KFNWYVDGVE VHNAKTKPRE EQYGSTYRVV SVLTVLHQDW LNGKEYKCKV 360
SNKALPAPIE KTISKAKGQP REPQVYTLPP SREEMTKNQV SLTCLVKGFY PSDIAVEWES 420
NGQPENNYKT TPPVLDSGDS FFLYSKLTVD KSRWQQGNVF SCSVMHEALH NHYTQKSLSL 480
SPGK

```

```

SEQ ID NO: 265      moltype = AA  length = 480

```

-continued

FEATURE Location/Qualifiers
 REGION 1..480
 note = scFc-6
 source 1..480
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 265

DKTHTCPPCP	APELLGGPSV	FLFPPKPKDT	LMISRTPEVT	CVVVDVSHED	PEVKFNWYVD	60
GVEVHNATK	PREEQYGSTY	RVVSVLTVLH	QDWLNGKEYK	CKVSNKALPA	PIEKTISKAK	120
GQPREPQVYT	LPPSREEMTK	NQVSLTCLVK	GFYPSTIAVE	WESNGQPENN	YKTPPVLD	180
DGSFFLYSKL	TVDKSRWQQG	NVFSCSVME	ALHNHYTQKS	LSLSPGGGGS	GGGGSGGGGS	240
GGGGSGGGGS	GGGGSDKTH	CPPCPAPELL	GGPSVFLFPP	KPKDTLMISR	TPEVTCVVVD	300
VSHEDPEVKF	NWYVDGVEVH	NAKTKPREEQ	YGSTYRVVSV	LTVLHQDWLN	GKEYKCKVSN	360
KALPAPIEKT	ISKAKGQPRE	PQVYTLPPSR	EEMTKNQVSL	TCLVKGFYPS	DIAVEWESNG	420
QPENNYKTTT	PVLDSGGSFF	LYSKLTVDKS	RWQQGNVFSC	SVMHEALHNNH	YTQKSLSLSP	480

SEQ ID NO: 266 moltype = AA length = 484
 FEATURE Location/Qualifiers
 REGION 1..484
 note = scFc-7
 source 1..484
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 266

DKTHTCPPCP	APELLGGPSV	FLFPPKPKDT	LMISRTPEVT	CVVVDVSHED	PEVKFNWYVD	60
GVEVHNATK	PCEEQYNSTY	RCVSVLTVLH	QDWLNGKEYK	CKVSNKALPA	PIEKTISKAK	120
GQPREPQVYT	LPPSREEMTK	NQVSLTCLVK	GFYPSTIAVE	WESNGQPENN	YKTPPVLD	180
DGSFFLYSKL	TVDKSRWQQG	NVFSCSVME	ALHNHYTQKS	LSLSPGKGGG	GSGGGSGGGG	240
GSGGGSGGGG	GSGGGGSDKT	HTCPPCPAPE	LLGGPSVFLF	PPKPKDTLMI	SRTPEVTCVV	300
VDVSHEDPEV	KFNWYVDGVE	VHNAKTKPCE	EQYNSTYRCV	SVLTVLHQDW	LNGKEYKCKV	360
SNKALPAPIE	KTISKAKGQP	REPQVYTLPP	SREEMTKNQV	SLTCLVKGFY	PSDIAVEWES	420
NGQPENNYKT	TPPVLDSDGS	FFLYSLKTV	KSRWQQGNVF	SCSVMEALH	NHYTQKSLSL	480
SPGK						484

SEQ ID NO: 267 moltype = AA length = 480
 FEATURE Location/Qualifiers
 REGION 1..480
 note = scFc-8
 source 1..480
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 267

DKTHTCPPCP	APELLGGPSV	FLFPPKPKDT	LMISRTPEVT	CVVVDVSHED	PEVKFNWYVD	60
GVEVHNATK	PCEEQYNSTY	RCVSVLTVLH	QDWLNGKEYK	CKVSNKALPA	PIEKTISKAK	120
GQPREPQVYT	LPPSREEMTK	NQVSLTCLVK	GFYPSTIAVE	WESNGQPENN	YKTPPVLD	180
DGSFFLYSKL	TVDKSRWQQG	NVFSCSVME	ALHNHYTQKS	LSLSPGGGGS	GGGGSGGGGS	240
GGGGSGGGGS	GGGGSDKTH	CPPCPAPELL	GGPSVFLFPP	KPKDTLMISR	TPEVTCVVVD	300
VSHEDPEVKF	NWYVDGVEVH	NAKTKPCEEQ	YNSTYRCVSV	LTVLHQDWLN	GKEYKCKVSN	360
KALPAPIEKT	ISKAKGQPRE	PQVYTLPPSR	EEMTKNQVSL	TCLVKGFYPS	DIAVEWESNG	420
QPENNYKTTT	PVLDSGGSFF	LYSKLTVDKS	RWQQGNVFSC	SVMHEALHNNH	YTQKSLSLSP	480

SEQ ID NO: 268 moltype = AA length = 988
 FEATURE Location/Qualifiers
 REGION 1..988
 note = MS_1xCD3-scFc
 source 1..988
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 268

QVQLQESGPG	LVKPSSETLSL	TCTVSGGSIS	SSSYWGWIR	QPPGKGLEWI	GSIIYSGITN	60
YNPSLKSRTV	ISVDTSKNQF	SLKLSVTA	DTAVYYCARP	SNYDAFDI	WQGMVTVSSG	120
GGGGSGGGGS	GGGQSGLTQ	PPSVSGAPGQ	RVTISCTGSS	SNIGAGYDVH	WYQQVPGTAP	180
KLLIYGNYSR	PSGVPDRFSG	SKSGTSASLA	ITGLQAEDEA	DYYCQSYDSS	LGWVFGGGT	240
KLTVLSGGGG	TEYQLVESGG	GLVQPGGSLK	LSCAASGFTF	NKYAMNWVRQ	APGKGLEWVA	300
RIRSKYNRYA	SVYADSVKDR	FTISRDDSKN	TAYLQMNHLK	TEDTAVYYCV	RHGNFGNSYI	360
SYWAYWGQGT	LVTYSSGGGG	SGGGSGGGGG	SQTIVTQEPS	LTVSPGGTIV	LTCGSSGTGAV	420
TSGNYPNWVQ	QKPGQAPRGL	IGGTFKFLAPG	TPARFSGSL	GGKAALTL	SGVQPEDEAEYY	480
CVLWYSNRWV	FGGGTKLTVL	GGGGDKTH	TPCPAPPELLG	GPSVFLFPPK	PKDTLMISR	540
PEVTCVVVDV	SHEDPEVKFN	WYVDGVEVHN	AKTKPCEEQY	GSTYRCVSVL	TVLHQDWLNG	600
KEYKCKVSNK	ALPAPIEKTI	SKAKGQPREP	QVYTLPPSRE	EMTKNQVSLT	CLVKGFYPSD	660
IAVEWESNGQ	PENNYKTTTP	VLDSDGSFFL	YSKLTVDKSR	WQQGNVFSCS	VMHEALHNNHY	720
TQKSLSLSPG	KGGGGSGGGG	SGGGSGGGGG	SGGGSGGGGG	SDKTHTCPPC	PAPPELLGGPS	780
VFLFPPKPKD	TLMSRTPEV	TCVVVDVSH	DPEVKFNWYV	DGVEVHNATK	KPCEEQYGST	840
YRCVSVLTVL	QDWLNGKEY	KCKVSNKALP	APIEKTISKA	KGQPREPQVY	TLPPSREEMT	900
KNQVSLTCLV	KGFYPSDIAV	EWESNGQPEN	NYKTPPVLD	SDGSFFLYSK	LTVDKSRWQQ	960
GNVFSCSVMH	EALHNHYTQK	SLSLSPGK				988

-continued

SEQ ID NO: 269 moltype = AA length = 986
FEATURE Location/Qualifiers
REGION 1..986
 note = MS_1xCD3-scFc_delGK
source 1..986
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 269

QVQLQESGPG	LVKPSSETLSL	TCTVSGGSIS	SSSYWGWIR	QPPGKGLEWI	GSIIYSGITN	60
YNPSLKSRVT	ISVDTSKNQF	SLKLSSVTAA	DTAVYYCARP	SNYDAFDIWG	QGTMTVTVSSG	120
GGSGGGGGSG	GGGSQSVLTQ	PPSVSGAPGQ	RVTISCTGSS	SNIGAGYDVH	WYQQVPGTAP	180
KLLIYGNRKR	PSGVPDRFSG	SKSGTSASLA	ITGLQAEDEA	DYYCQSYDSS	LGGWVFGGGT	240
KLTVLSSGGG	SEVQLVESGG	GLVQPGGSLK	LSCAASGFTF	NKYAMNWVRQ	APGKGLEWVA	300
RIRSKYNRYA	TYYADSVKDR	FTISRDDSKN	TAYLQMNNLK	TEDTAVYYCV	RHGNFGNSYI	360
SYWAYWQQGT	LVTYSSGGGG	SGGGSGGGGG	SQTVVTQEPS	LTVSPGGTVT	LTCGSSTGAV	420
TSGNYPNWVQ	QKPGQAPRGL	IGGTKFLAPG	TPARFSGSLL	GGKAALTLSG	VQPEDEAEYY	480
CVLWYSNRWV	FGGGTKLTVL	GGGGDKHTHC	PPCPAPELLG	GPSVFLFPPK	PKDTLMISRT	540
PEVTCVVVDV	SHEDPEVKFN	WYVDGVEVHN	AKTKPCEEQY	GSTYRCVSVL	TVLHQDWLNG	600
KEYKCKVSNK	ALPAPIEKTI	SKAKGQPREP	QVYTLPPSRE	EMTKNQVSLT	CLVKGFYPSD	660
IAVEWESNGQ	PENNYKTPPP	VLDSDGSFFL	YSKLTVDKSR	WQQGNVFS	VMHEALHNHY	720
TQKSLSLSPG	GGSGGGGGSG	GGSGGGGGSG	GGSGGGGGSD	KTHTCPPCPA	PELLGGPSVF	780
LFPKPKPDKL	MISRTPEVTC	VVDVSHEDP	EVKFNWYVDG	VEVHNAKTKP	CEEQYGSTYR	840
CVSVLTVLHQ	DWLNGKEYKC	KVSNKALPAP	IEKTISKAKG	QPREPQVYTL	PPSREEMTKN	900
QVSLTCLVKG	FYPSPDIAVEW	ESNGQPENNY	KTPPVVLDSD	GSFPLYSKLT	VDKSRWQQGN	960
VFSCSVMHEA	LHNHYTQKSL	SLSPGK				986

SEQ ID NO: 270 moltype = AA length = 988
FEATURE Location/Qualifiers
REGION 1..988
 note = MS_1_CxCD3-scFc
source 1..988
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 270

QVQLQESGPG	LVKPSSETLSL	TCTVSGGSIS	SSSYWGWIR	QPPGKCLEWI	GSIIYSGITN	60
YNPSLKSRVT	ISVDTSKNQF	SLKLSSVTAA	DTAVYYCARP	SNYDAFDIWG	QGTMTVTVSSG	120
GGSGGGGGSG	GGGSQSVLTQ	PPSVSGAPGQ	RVTISCTGSS	SNIGAGYDVH	WYQQVPGTAP	180
KLLIYGNRKR	PSGVPDRFSG	SKSGTSASLA	ITGLQAEDEA	DYYCQSYDSS	LGGWVFGCGT	240
KLTVLSSGGG	SEVQLVESGG	GLVQPGGSLK	LSCAASGFTF	NKYAMNWVRQ	APGKGLEWVA	300
RIRSKYNRYA	TYYADSVKDR	FTISRDDSKN	TAYLQMNNLK	TEDTAVYYCV	RHGNFGNSYI	360
SYWAYWQQGT	LVTYSSGGGG	SGGGSGGGGG	SQTVVTQEPS	LTVSPGGTVT	LTCGSSTGAV	420
TSGNYPNWVQ	QKPGQAPRGL	IGGTKFLAPG	TPARFSGSLL	GGKAALTLSG	VQPEDEAEYY	480
CVLWYSNRWV	FGGGTKLTVL	GGGGDKHTHC	PPCPAPELLG	GPSVFLFPPK	PKDTLMISRT	540
PEVTCVVVDV	SHEDPEVKFN	WYVDGVEVHN	AKTKPCEEQY	GSTYRCVSVL	TVLHQDWLNG	600
KEYKCKVSNK	ALPAPIEKTI	SKAKGQPREP	QVYTLPPSRE	EMTKNQVSLT	CLVKGFYPSD	660
IAVEWESNGQ	PENNYKTPPP	VLDSDGSFFL	YSKLTVDKSR	WQQGNVFS	VMHEALHNHY	720
TQKSLSLSPG	KGGSGGGGGG	SGGGSGGGGG	SGGGSGGGGG	SDKTHTCPPC	PAPELLGGPS	780
VFLFPPKPKD	LTMISRTPEV	TCVVVDVSHE	DPEVKFNWYV	DGVEVHNAKT	KPCEEQYGST	840
YRCVSVLTVL	HQDWLNGKEY	KCKVSNKALP	APIEKTISKA	KGQPREPQVY	TLPPSREEMT	900
KNQVSLTCLV	KGFYPSPDIAV	EWESNGQPEN	NYKTPPVLD	SDGSFFLYSK	LTVDKSRWQQ	960
GNVFS	CSVMH	EALHNHYTQK	SLSLSPGK			988

SEQ ID NO: 271 moltype = AA length = 986
FEATURE Location/Qualifiers
REGION 1..986
 note = MS_1_CxCD3-scFc_delGK
source 1..986
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 271

QVQLQESGPG	LVKPSSETLSL	TCTVSGGSIS	SSSYWGWIR	QPPGKCLEWI	GSIIYSGITN	60
YNPSLKSRVT	ISVDTSKNQF	SLKLSSVTAA	DTAVYYCARP	SNYDAFDIWG	QGTMTVTVSSG	120
GGSGGGGGSG	GGGSQSVLTQ	PPSVSGAPGQ	RVTISCTGSS	SNIGAGYDVH	WYQQVPGTAP	180
KLLIYGNRKR	PSGVPDRFSG	SKSGTSASLA	ITGLQAEDEA	DYYCQSYDSS	LGGWVFGCGT	240
KLTVLSSGGG	SEVQLVESGG	GLVQPGGSLK	LSCAASGFTF	NKYAMNWVRQ	APGKGLEWVA	300
RIRSKYNRYA	TYYADSVKDR	FTISRDDSKN	TAYLQMNNLK	TEDTAVYYCV	RHGNFGNSYI	360
SYWAYWQQGT	LVTYSSGGGG	SGGGSGGGGG	SQTVVTQEPS	LTVSPGGTVT	LTCGSSTGAV	420
TSGNYPNWVQ	QKPGQAPRGL	IGGTKFLAPG	TPARFSGSLL	GGKAALTLSG	VQPEDEAEYY	480
CVLWYSNRWV	FGGGTKLTVL	GGGGDKHTHC	PPCPAPELLG	GPSVFLFPPK	PKDTLMISRT	540
PEVTCVVVDV	SHEDPEVKFN	WYVDGVEVHN	AKTKPCEEQY	GSTYRCVSVL	TVLHQDWLNG	600
KEYKCKVSNK	ALPAPIEKTI	SKAKGQPREP	QVYTLPPSRE	EMTKNQVSLT	CLVKGFYPSD	660
IAVEWESNGQ	PENNYKTPPP	VLDSDGSFFL	YSKLTVDKSR	WQQGNVFS	VMHEALHNHY	720
TQKSLSLSPG	GGSGGGGGSG	GGSGGGGGSG	GGSGGGGGSD	KTHTCPPCPA	PELLGGPSVF	780
LFPKPKPDKL	MISRTPEVTC	VVDVSHEDP	EVKFNWYVDG	VEVHNAKTKP	CEEQYGSTYR	840
CVSVLTVLHQ	DWLNGKEYKC	KVSNKALPAP	IEKTISKAKG	QPREPQVYTL	PPSREEMTKN	900

-continued

QVSLTCLVKG	FYPSDIAVEW	ESNGQPENNY	KTPPVLDSD	GSFPLYSKLT	VDKSRWQQGN	960
VFSCSVMHEA	LHNHYTQKSL	SLSPGK				986

SEQ ID NO: 272 moltype = AA length = 982
FEATURE Location/Qualifiers
REGION 1..982
 note = MS_2xCD3-scFc
source 1..982
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 272
QVQLVESGGG VVQPGRLRL SCAASGFTFS DHYMSWIRQT PGKGLEWVSY ISSSGSTIYY 60
ADSVKGRFTI SRDNAKNSLY LQMNSLRAED TAVYYCARDL GPSFDYWQGG SLVTVSSGGG 120
GSGGGSGGGG GSDIQMTQSP SSVASVGDRL VTITCRASQG ISSWLAWYQQ KPGKAPKLLI 180
DAASRLQSGV PSRFGSGSGG TDFTLTISSL QPEDFATYYC QQANSFPRTF GQGTKVEIKS 240
GGGSEVQLV ESGGGLVQPG GSKLSCAAS GFTFNKYAMN WVRQAPGKGL EWVARIRSKY 300
NNYATYYADS VKDRFTISR DSKNTAYLQM NNLKTEDTAV YVCVRHGNFG NSYISYWAYW 360
GQGTLLTVSS GGGGSGGGGS GGGGSQTVVT QEPSLTVSPG GTVTLTCGSS TGAVTSGNYP 420
NWVQKPGQA PRGLIGTKF LAPGTPARFS GSLLGKKAAL TSGVQPEDE AEYYCVLWYS 480
NRWVFGGGTK LTVLGGGDK THTCPAP PAP ELLGGPSVFL FPPKPKDTLM ISRTPEVTCV 540
VVDVSHEDPE VKFNWYVDG EVHNAKTKPC EEQYGSTYRC VSVLTVLHGD WLNKEYKCK 600
VSNKALPAPI EKTISKAKGQ PREPQVYTL PPREMTKNQ VSLTCLVKGF YPSDIAVEWE 660
SNGQPENNYK TTPPVLDSDG SFFLYSKLTV DKSRLWQQGNV FSCSVMHEAL HNHYTQKSL 720
LSPGKGGGGS GGGGSGGGGS GGGGSGGGGS GGGGSDKTHT CPPCPAPELL GGPVFLFPP 780
KPKDTLMISR TPEVTCVVVD VSHEDPEVK NWYVDGVEVH NAKTKPCEEQ YGSTRYCVSV 840
LTVLHGDWLN GKEYKCKVSN KALPAPIEKT ISKAKGQPRE PQVYTLPPSR EEMTKNQVSL 900
TCLVKGFPYS DIAVEWESNG QPENNYKTP PVLDSDGSFF LYSKLTVDKS RWQQGNVFS 960
SVMHEALHNH YTQKSLSLSP GK 982

SEQ ID NO: 273 moltype = AA length = 980
FEATURE Location/Qualifiers
REGION 1..980
 note = MS_2xCD3-scFc_delGK
source 1..980
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 273
QVQLVESGGG VVQPGRLRL SCAASGFTFS DHYMSWIRQT PGKGLEWVSY ISSSGSTIYY 60
ADSVKGRFTI SRDNAKNSLY LQMNSLRAED TAVYYCARDL GPSFDYWQGG SLVTVSSGGG 120
GSGGGSGGGG GSDIQMTQSP SSVASVGDRL VTITCRASQG ISSWLAWYQQ KPGKAPKLLI 180
DAASRLQSGV PSRFGSGSGG TDFTLTISSL QPEDFATYYC QQANSFPRTF GQGTKVEIKS 240
GGGSEVQLV ESGGGLVQPG GSKLSCAAS GFTFNKYAMN WVRQAPGKGL EWVARIRSKY 300
NNYATYYADS VKDRFTISR DSKNTAYLQM NNLKTEDTAV YVCVRHGNFG NSYISYWAYW 360
GQGTLLTVSS GGGGSGGGGS GGGGSQTVVT QEPSLTVSPG GTVTLTCGSS TGAVTSGNYP 420
NWVQKPGQA PRGLIGTKF LAPGTPARFS GSLLGKKAAL TSGVQPEDE AEYYCVLWYS 480
NRWVFGGGTK LTVLGGGDK THTCPAP PAP ELLGGPSVFL FPPKPKDTLM ISRTPEVTCV 540
VVDVSHEDPE VKFNWYVDG EVHNAKTKPC EEQYGSTYRC VSVLTVLHGD WLNKEYKCK 600
VSNKALPAPI EKTISKAKGQ PREPQVYTL PPREMTKNQ VSLTCLVKGF YPSDIAVEWE 660
SNGQPENNYK TTPPVLDSDG SFFLYSKLTV DKSRLWQQGNV FSCSVMHEAL HNHYTQKSL 720
LSPGKGGGSG GGGGSGGGSG GGGGSGGGSG GGGSDKTHTCP PCPAPELLGG PSVFLFPPKP 780
KDTLMISRTPEVTCVVVDVSHEDPEVKFNW YVDGVEVHNA KTKPCEEQYQ STRYCVSVLT 840
VLHGDWLNKG EYKCKVSNKA LPAPIEKTIS KAKGQPREPQ VYTLPPSREE MTKNQVSLTC 900
LVKGFYPSDI AVEWESNGQP ENNYKTPPV LDSDGSFFLY SKLTVDKSRW QQGNVFS 960
MHEALHNHYT QKSLSLSPGK 980

SEQ ID NO: 274 moltype = AA length = 982
FEATURE Location/Qualifiers
REGION 1..982
 note = MS_2_CCD3-scFc
source 1..982
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 274
QVQLVESGGG VVQPGRLRL SCAASGFTFS DHYMSWIRQT PGKCLEWVSY ISSSGSTIYY 60
ADSVKGRFTI SRDNAKNSLY LQMNSLRAED TAVYYCARDL GPSFDYWQGG SLVTVSSGGG 120
GSGGGSGGGG GSDIQMTQSP SSVASVGDRL VTITCRASQG ISSWLAWYQQ KPGKAPKLLI 180
DAASRLQSGV PSRFGSGSGG TDFTLTISSL QPEDFATYYC QQANSFPRTF GCGTKVEIKS 240
GGGSEVQLV ESGGGLVQPG GSKLSCAAS GFTFNKYAMN WVRQAPGKGL EWVARIRSKY 300
NNYATYYADS VKDRFTISR DSKNTAYLQM NNLKTEDTAV YVCVRHGNFG NSYISYWAYW 360
GQGTLLTVSS GGGGSGGGGS GGGGSQTVVT QEPSLTVSPG GTVTLTCGSS TGAVTSGNYP 420
NWVQKPGQA PRGLIGTKF LAPGTPARFS GSLLGKKAAL TSGVQPEDE AEYYCVLWYS 480
NRWVFGGGTK LTVLGGGDK THTCPAP PAP ELLGGPSVFL FPPKPKDTLM ISRTPEVTCV 540
VVDVSHEDPE VKFNWYVDG EVHNAKTKPC EEQYGSTYRC VSVLTVLHGD WLNKEYKCK 600
VSNKALPAPI EKTISKAKGQ PREPQVYTL PPREMTKNQ VSLTCLVKGF YPSDIAVEWE 660
SNGQPENNYK TTPPVLDSDG SFFLYSKLTV DKSRLWQQGNV FSCSVMHEAL HNHYTQKSL 720
LSPGKGGGGS GGGGSGGGGS GGGGSGGGGS GGGGSDKTHT CPPCPAPELL GGPVFLFPP 780

-continued

```
KPKDTLMISR TPEVTCVVVD VSHEDPEVKF NWYVDGVEVH NAKTKPCEEQ YGSTYRCVSV 840
LTVLHQDWLN GKEYKCKVSN KALPAPIEKT ISKAKGQPRE PQVYTLPPSR EEMTKNQVSL 900
TCLVKGFYPS DIAVEWESNG QPENNYKTTT PVLDSGDSFF LYSKLTVDKS RWQQGNVFSF 960
SVMHEALHNH YTQKSLSLSP GK 982
```

```
SEQ ID NO: 275      moltype = AA length = 980
FEATURE            Location/Qualifiers
REGION             1..980
                   note = MS_2_CxCD3-scFc_delGK
source             1..980
                   mol_type = protein
                   organism = synthetic construct
```

```
SEQUENCE: 275
QVQLVESGGG VVQPGSRSLRL SCAASGFTFS DHYMSWIRQT PGKCLEWVSY ISSSGSTIYY 60
ADSVKGRFTI SRDNAKNSLY LQMNSLRAED TAVYYCARDL GPSFDYWGQG SLVTVSSGGG 120
GGGGGGSGGG GSDIQMTQSP SSVASVGDRT VTITCRASQG ISSWLAWYQQ KPGKAPKLLI 180
DAASRLQSGV PSRFGSGSGG TDFTLTISSL QPEDFATYYC QQANSFPRTF GCGTKVEIKS 240
GGGGSEVQLV ESGGGLVQPG GSKLSCAAS GFTFNKYAMN WVRQAPGKGL EWVARIRSKY 300
NNYATYYADS VKDRFTISR DSKNTAYLQM NNLKTEDTAV YYCVRHGNFG NSYISYWAYW 360
GQGTSLTVSS GGGGGSGGGG GGGGSQTVVT QEPSTLVSPG GTVTLTCGSS TGAVTSGNYP 420
NWVQKPGQA PRGLIGGTFK LAPGTPARFS GSLGGKAAL TSGVQPEDE AEYCVLWYS 480
NRWVFGGGTK LTVLGGGDK THTCPPCPAP ELLGGPSVFL FPPKPKDTLM ISRTPEVTCV 540
VVDVSHEDPE VKFNWYVDG EVHNAKTKPC EEQYGSTYRC VSVLTVLHQD WLNKEYKCK 600
VSNKALPAPI EKTISKAKGQ PREPQVYTL PSREEMTKNQ VSLTCLVKGF YPSDIAVEWE 660
SNGQPENNYK TTPVLDSDG SFPLYSKLT DKSRRQQGNV FSCSVMHEAL HNHYTQKSL 720
LSPGGGGSGG GGGGGGGSGG GGGGGGGSGG GGSDDKTHCP PCPAPELLGG PSVFLFPPKP 780
KDTLMISRTPEVTCVVVDVSHEDPEVKF NWYVDGVEVH NAKTKPCEEQ YGSTYRCVSV 840
VHLHQDWLNK GKEYKCKVSN KALPAPIEKT ISKAKGQPRE PQVYTLPPSR EEMTKNQVSL 900
LVKGFYPSDI AVEWESNGQP ENNYKTTTPV LDDSDGSFFLY SKLTVDKSRW QQGNVFSFCSV 960
MHEALHNHYT QKSLSLSPGK 982
```

```
SEQ ID NO: 276      moltype = AA length = 982
FEATURE            Location/Qualifiers
REGION             1..982
                   note = MS_3xCD3-scFc
source             1..982
                   mol_type = protein
                   organism = synthetic construct
```

```
SEQUENCE: 276
QVQLVESGGG LVKPGGSLRL SCAASGFTFS DYYMTWIRQA PGKGLEWISY ISSSGSTIYY 60
ADSVKGRFTI SRDNAKNSLY LQMNSLRAED TAVYYCARDR NSHFDYWGQG TLVTVSSGGG 120
GGGGGGSGGG GSDIQMTQSP SSVASVGDRT VTITCRASQG ITRWLAWYQQ KPGKAPKLLI 180
YAASVLQSGV PSRFGSGSGG TDFTLTISSL QPEDFATYYC QQNSFPRTF GQGTKVEIKS 240
GGGGSEVQLV ESGGGLVQPG GSKLSCAAS GFTFNKYAMN WVRQAPGKGL EWVARIRSKY 300
NNYATYYADS VKDRFTISR DSKNTAYLQM NNLKTEDTAV YYCVRHGNFG NSYISYWAYW 360
GQGTSLTVSS GGGGGSGGGG GGGGSQTVVT QEPSTLVSPG GTVTLTCGSS TGAVTSGNYP 420
NWVQKPGQA PRGLIGGTFK LAPGTPARFS GSLGGKAAL TSGVQPEDE AEYCVLWYS 480
NRWVFGGGTK LTVLGGGDK THTCPPCPAP ELLGGPSVFL FPPKPKDTLM ISRTPEVTCV 540
VVDVSHEDPE VKFNWYVDG EVHNAKTKPC EEQYGSTYRC VSVLTVLHQD WLNKEYKCK 600
VSNKALPAPI EKTISKAKGQ PREPQVYTL PSREEMTKNQ VSLTCLVKGF YPSDIAVEWE 660
SNGQPENNYK TTPVLDSDG SFPLYSKLT DKSRRQQGNV FSCSVMHEAL HNHYTQKSL 720
LSPGGGGSGG GGGGGGGSGG GGGGGGGSGG GGSDDKTHCP PCPAPELLGG PSVFLFPPKP 780
KPKDTLMISR TPEVTCVVVD VSHEDPEVKF NWYVDGVEVH NAKTKPCEEQ YGSTYRCVSV 840
LTVLHQDWLN GKEYKCKVSN KALPAPIEKT ISKAKGQPRE PQVYTLPPSR EEMTKNQVSL 900
TCLVKGFYPS DIAVEWESNG QPENNYKTTT PVLDSGDSFF LYSKLTVDKS RWQQGNVFSF 960
SVMHEALHNH YTQKSLSLSP GK 982
```

```
SEQ ID NO: 277      moltype = AA length = 980
FEATURE            Location/Qualifiers
REGION             1..980
                   note = MS_3xCD3-scFc_delGK
source             1..980
                   mol_type = protein
                   organism = synthetic construct
```

```
SEQUENCE: 277
QVQLVESGGG LVKPGGSLRL SCAASGFTFS DYYMTWIRQA PGKGLEWISY ISSSGSTIYY 60
ADSVKGRFTI SRDNAKNSLY LQMNSLRAED TAVYYCARDR NSHFDYWGQG TLVTVSSGGG 120
GGGGGGSGGG GSDIQMTQSP SSVASVGDRT VTITCRASQG ITRWLAWYQQ KPGKAPKLLI 180
YAASVLQSGV PSRFGSGSGG TDFTLTISSL QPEDFATYYC QQNSFPRTF GQGTKVEIKS 240
GGGGSEVQLV ESGGGLVQPG GSKLSCAAS GFTFNKYAMN WVRQAPGKGL EWVARIRSKY 300
NNYATYYADS VKDRFTISR DSKNTAYLQM NNLKTEDTAV YYCVRHGNFG NSYISYWAYW 360
GQGTSLTVSS GGGGGSGGGG GGGGSQTVVT QEPSTLVSPG GTVTLTCGSS TGAVTSGNYP 420
NWVQKPGQA PRGLIGGTFK LAPGTPARFS GSLGGKAAL TSGVQPEDE AEYCVLWYS 480
NRWVFGGGTK LTVLGGGDK THTCPPCPAP ELLGGPSVFL FPPKPKDTLM ISRTPEVTCV 540
VVDVSHEDPE VKFNWYVDG EVHNAKTKPC EEQYGSTYRC VSVLTVLHQD WLNKEYKCK 600
VSNKALPAPI EKTISKAKGQ PREPQVYTL PSREEMTKNQ VSLTCLVKGF YPSDIAVEWE 660
```

-continued

SNGQPENNYK	TTPPVLDSDG	SFFLYSKLTV	DKSRWQQGNV	FSCSVMHEAL	HNHYTQKSL	720
LSPGGGGSGG	GGSGGGSGG	GGSGGGSGG	GGSDKTHTCP	PCPAPELLGG	PSVFLFPPKP	780
KDTLMISRTP	EVTCTVVVDVS	HEDPEVKFNW	YVDGVEVHNA	KTKPCEEQYG	STYRCVSVLT	840
VLHQDWLNGK	EYCKVSNKA	LPAPIEKTIS	KAKGQPREPQ	VYTLPPSREE	MTKNQVSLTC	900
LVKGFYPSDI	AVEWESNGQP	ENNYKTTPPV	LDSDGSFFLY	SKLTVDKSRW	QQGNVFSCSV	960
MHEALHNHYT	QKSLSLSPGK					980

SEQ ID NO: 278 moltype = AA length = 982
 FEATURE Location/Qualifiers
 REGION 1..982
 note = MS_3_CCxCD3-scFc
 source 1..982
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 278

QVQLVESGGG	LVKPGGSLRL	SCAASGFTFS	DYYMTWIRQA	PGKCLEWISY	ISSSGSTIYY	60
ADSVKGRFTI	SRDNAKNSLY	LQMNSLRAED	TAVYYCARDR	NSHFDYWQQG	TLVTVSSGGG	120
GGSGGGSGGG	GSDIQMTQSP	SSVSASVGD	VTITCRASQG	ITRWLAHYQQ	KPGKAPKLLI	180
YAASVLQSGV	PSRFSGSGSG	TDFTLTISL	QPEDFATYYC	QQSNSFPRTF	GCGTKVEIKS	240
GGGGSEVQLV	ESGGGLVQPG	GSLKLSCAAS	GFTFNKYAMN	WVRQAPGKGL	EWVARIRSKY	300
NNYATYYADS	VKDRFTISR	DSKNTAYLQ	NNLKTEDTAV	YYCVRHGNFG	NSYISYWAYW	360
GQGTLLVTSS	GGGGSGGGG	GGGGSQT	QEPSTVSPG	GTVTLTCGSS	TGAVTSGNYP	420
NWVQKPGQA	PRGLIGGTFK	LAPGTPARFS	GSLGGKAAAL	TLSGVQPEDE	AEYYCVLWYS	480
NRWVFGGGTK	LTVLGGGDK	THTCPPCPAP	ELLGGPSVFL	PPPKPKDTLM	ISRTPEVTCV	540
VVDVSHEDPE	VKFNWYVDG	EVHNAKTKPC	EEQYGYSTYRC	VSVLTVLHQD	WLNKGEYKCK	600
VSNKALPAPI	EKTISKAKGQ	PREPQVYTL	PSREEMTKNQ	VSLTCLVKGF	YPSDIAVEWE	660
SNGQPENNYK	TTPPVLDSDG	SFFLYSKLTV	DKSRWQQGNV	FSCSVMHEAL	HNHYTQKSL	720
LSPGGGGSGG	GGSGGGSGG	GGSGGGSGG	GGSDKTHTCP	PCPAPELLGG	GGPSVFLFPP	780
KPKDTLMISR	TPEVTCVVVD	VSHEDPEVKF	NWYVDGVEVH	NAKTKPCEEQ	YGSTYRCVSV	840
LTVLHQDWLN	GKEYKCKVSN	KALPAPIEKT	ISKAKGQPRE	PQVYTLPPSR	EEMTKNQVSL	900
TCLVKGFYPS	DIAVEWESNG	QPENNYKTPP	PVLDSDGSFF	LYSKLTVDKS	RWQQGNVFSC	960
SVMHEALHNN	YTQKSLSLSP	GK				982

SEQ ID NO: 279 moltype = AA length = 980
 FEATURE Location/Qualifiers
 REGION 1..980
 note = MS_3_CCxCD3-scFc_delGK
 source 1..980
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 279

QVQLVESGGG	LVKPGGSLRL	SCAASGFTFS	DYYMTWIRQA	PGKCLEWISY	ISSSGSTIYY	60
ADSVKGRFTI	SRDNAKNSLY	LQMNSLRAED	TAVYYCARDR	NSHFDYWQQG	TLVTVSSGGG	120
GGSGGGSGGG	GSDIQMTQSP	SSVSASVGD	VTITCRASQG	ITRWLAHYQQ	KPGKAPKLLI	180
YAASVLQSGV	PSRFSGSGSG	TDFTLTISL	QPEDFATYYC	QQSNSFPRTF	GCGTKVEIKS	240
GGGGSEVQLV	ESGGGLVQPG	GSLKLSCAAS	GFTFNKYAMN	WVRQAPGKGL	EWVARIRSKY	300
NNYATYYADS	VKDRFTISR	DSKNTAYLQ	NNLKTEDTAV	YYCVRHGNFG	NSYISYWAYW	360
GQGTLLVTSS	GGGGSGGGG	GGGGSQT	QEPSTVSPG	GTVTLTCGSS	TGAVTSGNYP	420
NWVQKPGQA	PRGLIGGTFK	LAPGTPARFS	GSLGGKAAAL	TLSGVQPEDE	AEYYCVLWYS	480
NRWVFGGGTK	LTVLGGGDK	THTCPPCPAP	ELLGGPSVFL	PPPKPKDTLM	ISRTPEVTCV	540
VVDVSHEDPE	VKFNWYVDG	EVHNAKTKPC	EEQYGYSTYRC	VSVLTVLHQD	WLNKGEYKCK	600
VSNKALPAPI	EKTISKAKGQ	PREPQVYTL	PSREEMTKNQ	VSLTCLVKGF	YPSDIAVEWE	660
SNGQPENNYK	TTPPVLDSDG	SFFLYSKLTV	DKSRWQQGNV	FSCSVMHEAL	HNHYTQKSL	720
LSPGGGGSGG	GGSGGGSGG	GGSGGGSGG	GGSDKTHTCP	PCPAPELLGG	PSVFLFPPKP	780
KDTLMISRTP	EVTCTVVVDVS	HEDPEVKFNW	YVDGVEVHNA	KTKPCEEQYG	STYRCVSVLT	840
VLHQDWLNGK	EYCKVSNKA	LPAPIEKTIS	KAKGQPREPQ	VYTLPPSREE	MTKNQVSLTC	900
LVKGFYPSDI	AVEWESNGQP	ENNYKTTPPV	LDSDGSFFLY	SKLTVDKSRW	QQGNVFSCSV	960
MHEALHNHYT	QKSLSLSPGK					980

SEQ ID NO: 280 moltype = AA length = 982
 FEATURE Location/Qualifiers
 REGION 1..982
 note = MS_4xCD3-scFc
 source 1..982
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 280

QVQLVESGGG	LVKPGGSLRL	SCAASGFTFS	DYYMTWIRQA	PGKGLEWLSY	ISSSGSTIYY	60
ADSVKGRFTI	SRDNAKNSLF	LQMNSLRAED	TAVYYCARDR	NSHFDYWQQG	TLVTVSSGGG	120
GGSGGGSGGG	GSDIQMTQSP	SSVSASVGD	VTITCRASQG	INTWLAWYQQ	KPGKAPKLLI	180
YGASGLQSGV	PSRFSGSGSG	TDFTLTISL	QPEDFATYYC	QQAQSFRTF	GQGTKVEIKS	240
GGGGSEVQLV	ESGGGLVQPG	GSLKLSCAAS	GFTFNKYAMN	WVRQAPGKGL	EWVARIRSKY	300
NNYATYYADS	VKDRFTISR	DSKNTAYLQ	NNLKTEDTAV	YYCVRHGNFG	NSYISYWAYW	360
GQGTLLVTSS	GGGGSGGGG	GGGGSQT	QEPSTVSPG	GTVTLTCGSS	TGAVTSGNYP	420
NWVQKPGQA	PRGLIGGTFK	LAPGTPARFS	GSLGGKAAAL	TLSGVQPEDE	AEYYCVLWYS	480
NRWVFGGGTK	LTVLGGGDK	THTCPPCPAP	ELLGGPSVFL	PPPKPKDTLM	ISRTPEVTCV	540

-continued

```
VVDVSHEDPE VKFNWYVDGV EVHNAKTKPC EEQYGYSTYRC VSVLTVLHQD WLNKEYKCK 600
VSNKALPAPI EKTISKAKGQ PREPQVYTL PPREMTKNQ VSLTCLVKGF YPSDIAVEWE 660
SNGQPENNYK TTPPVLDSDG SFFLYSKLTV DKSRWQQGNV FSCSVMHEAL HNHYTQKSL 720
LSPGKGGGGS GGGGSGGGGS GGGGSGGGGS GGGGSDKTHT CPPCPAPELL GGPSVFLFPP 780
KPKDTLMISR TPEVTCVVVD VSHEDPEVKF NWYVDGVEVH NAKTKPCEEQ YGSTRYCVSV 840
LTVLHQDWLNL GKEYKCKVSN KALPAPIEKT ISKAKGQPRE PQVYTLPPSR EEMTKNQVSL 900
TCLVKGFPYPS DIAVEWESNG QPENNYKTTT PVLDSGDSFF LYSKLTVDKS RWQQGNVFSC 960
SVMHEALHNH YTQKSLSLSP GK 982
```

```
SEQ ID NO: 281      moltype = AA length = 980
FEATURE            Location/Qualifiers
REGION             1..980
                   note = MS_4xCD3-scFc_delGK
source             1..980
                   mol_type = protein
                   organism = synthetic construct
```

```
SEQUENCE: 281
QVQLVESGGG LVKPGGSLRL SCAASGFTFS DYYMTWIRQA PGKCLEWLSY ISSSGSTIYY 60
ADSVKGRFTI SRDIAKNSLF LQMNSLRAED TAVYYCARDR NSHFDYWGQG TLVTVSSGGG 120
GGGGGGGGGG GSDIQMTQSP SSVASVGD RVTITCRASQG INTWLAWYQQ KPGKAPKLLI 180
YGASGLQSGV PSRFGSGSGG TDFTLTISL QPEDFATYYC QQAQSFPRTF GCGTKVEIKS 240
GGGSEVQLV ESGGGLVQPG GSKLSCAAS GFTFNKYAMN WVRQAPGKGL EWVARIRSKY 300
NNYATYYADS VKDRFTISR DSKNTAYLQM NNLKTEDTAV YVCVRHGNFG NSYISYWAYW 360
GGGTLVTVSS GGGGSGGGGS GGGGSGTQVT QEPSTLVSPG GTVTLTCGSS TGAVTSGNYP 420
NWVQQKPGQA PRGLIGTKF LAPGTARFS GLLGGKAAL TSGVQPEDE AEYCVLWYS 480
NRWVFGGGTK LTVLGGGDK THTCPPCPAP ELLGGPSVFL FPPKPKDTLM ISRTPEVTCV 540
VVDVSHEDPE VKFNWYVDGV EVHNAKTKPC EEQYGYSTYRC VSVLTVLHQD WLNKEYKCK 600
VSNKALPAPI EKTISKAKGQ PREPQVYTL PPREMTKNQ VSLTCLVKGF YPSDIAVEWE 660
SNGQPENNYK TTPPVLDSDG SFFLYSKLTV DKSRWQQGNV FSCSVMHEAL HNHYTQKSL 720
LSPGKGGGGS GGGGSGGGGS GGGGSGGGGS GGGSDKTHTCP PCPAPPELLG PSVFLFPPKP 780
KDTLMISRTP EVTCVVVDVS HEDPEVKFNW YVDGVEVHNA KTKPCEEQYQ STRYCVSVLT 840
VLHQDWLNGK EYKCKVSNKA LPAPIEKTIS KAKGQPREPQ VYTLPPSR EMTKNQVSLTC 900
LVKGFYPSDI AVEWESNGQP ENNYKTTTPV LDSGDSFFLY SKLTVDKSRW QQGNVFSCSV 960
MHEALHNHYT QKSLSLSPGK 982
```

```
SEQ ID NO: 282      moltype = AA length = 982
FEATURE            Location/Qualifiers
REGION             1..982
                   note = MS_4_CCD3-scFc
source             1..982
                   mol_type = protein
                   organism = synthetic construct
```

```
SEQUENCE: 282
QVQLVESGGG LVKPGGSLRL SCAASGFTFS DYYMTWIRQA PGKCLEWLSY ISSSGSTIYY 60
ADSVKGRFTI SRDIAKNSLF LQMNSLRAED TAVYYCARDR NSHFDYWGQG TLVTVSSGGG 120
GGGGGGGGGG GSDIQMTQSP SSVASVGD RVTITCRASQG INTWLAWYQQ KPGKAPKLLI 180
YGASGLQSGV PSRFGSGSGG TDFTLTISL QPEDFATYYC QQAQSFPRTF GCGTKVEIKS 240
GGGSEVQLV ESGGGLVQPG GSKLSCAAS GFTFNKYAMN WVRQAPGKGL EWVARIRSKY 300
NNYATYYADS VKDRFTISR DSKNTAYLQM NNLKTEDTAV YVCVRHGNFG NSYISYWAYW 360
GGGTLVTVSS GGGGSGGGGS GGGGSGTQVT QEPSTLVSPG GTVTLTCGSS TGAVTSGNYP 420
NWVQQKPGQA PRGLIGTKF LAPGTARFS GLLGGKAAL TSGVQPEDE AEYCVLWYS 480
NRWVFGGGTK LTVLGGGDK THTCPPCPAP ELLGGPSVFL FPPKPKDTLM ISRTPEVTCV 540
VVDVSHEDPE VKFNWYVDGV EVHNAKTKPC EEQYGYSTYRC VSVLTVLHQD WLNKEYKCK 600
VSNKALPAPI EKTISKAKGQ PREPQVYTL PPREMTKNQ VSLTCLVKGF YPSDIAVEWE 660
SNGQPENNYK TTPPVLDSDG SFFLYSKLTV DKSRWQQGNV FSCSVMHEAL HNHYTQKSL 720
LSPGKGGGGS GGGGSGGGGS GGGGSGGGGS GGGSDKTHT CPPCPAPELL GGPSVFLFPP 780
KPKDTLMISR TPEVTCVVVD VSHEDPEVKF NWYVDGVEVH NAKTKPCEEQ YGSTRYCVSV 840
LTVLHQDWLNL GKEYKCKVSN KALPAPIEKT ISKAKGQPRE PQVYTLPPSR EEMTKNQVSL 900
TCLVKGFPYPS DIAVEWESNG QPENNYKTTT PVLDSGDSFF LYSKLTVDKS RWQQGNVFSC 960
SVMHEALHNH YTQKSLSLSP GK 982
```

```
SEQ ID NO: 283      moltype = AA length = 980
FEATURE            Location/Qualifiers
REGION             1..980
                   note = MS_4_CCD3-scFc_delGK
source             1..980
                   mol_type = protein
                   organism = synthetic construct
```

```
SEQUENCE: 283
QVQLVESGGG LVKPGGSLRL SCAASGFTFS DYYMTWIRQA PGKCLEWLSY ISSSGSTIYY 60
ADSVKGRFTI SRDIAKNSLF LQMNSLRAED TAVYYCARDR NSHFDYWGQG TLVTVSSGGG 120
GGGGGGGGGG GSDIQMTQSP SSVASVGD RVTITCRASQG INTWLAWYQQ KPGKAPKLLI 180
YGASGLQSGV PSRFGSGSGG TDFTLTISL QPEDFATYYC QQAQSFPRTF GCGTKVEIKS 240
GGGSEVQLV ESGGGLVQPG GSKLSCAAS GFTFNKYAMN WVRQAPGKGL EWVARIRSKY 300
NNYATYYADS VKDRFTISR DSKNTAYLQM NNLKTEDTAV YVCVRHGNFG NSYISYWAYW 360
GGGTLVTVSS GGGGSGGGGS GGGGSGTQVT QEPSTLVSPG GTVTLTCGSS TGAVTSGNYP 420
```


-continued

NWVQKPGQA	PRGLIGGTKF	LAPGTPARFS	GSLLGKKAAL	TLSGVQPEDE	AEYYCVLWYS	480
NRWVFGGK	LTVLGGGDK	THTCPPCPAP	ELLGGPSVFL	FPPKPKDTLM	ISRTPEVTCV	540
VVDVSHEDPE	VKFNWYVDG	EVHNAKTKPC	EEQYGSTYRC	VSVLTVLHQD	WLNKEYKCK	600
VSNKALPAPI	EKTISKAKGQ	PREPQVYTL	PSREEMTKNQ	VSLTCLVKGF	YPSDIAVEWE	660
SNGQPENNYK	TTPPVLDSDG	SFFLYSKLTV	DKSRWQQGNV	FSCSVMHEAL	HNHYTQKSL	720
LSPGGGGSGG	GGSGGGSGG	GGSGGGSGG	GGSDKTHTCP	PCPAPELLGG	PSVFLFPPKP	780
KDTLMISRTP	EVTCTVVDVS	HEDPEVKFNW	YVDGVEVHNA	KTKPCEEQYG	STYRCVSVLT	840
VLHQDWLNGK	EYKCKVSNKA	LPAPIEKTIS	KAKGQPREPQ	VYTLPPSREE	MTKNQVSLTC	900
LVKGFYPSDI	AVEWESNGQP	ENNYKTTTPV	LDSGGSFFLY	SKLTVDKSRW	QQGNVFS	960
MHEALHNHYT	QKSLSLSPGK					980

SEQ ID NO: 284 moltype = AA length = 982
 FEATURE Location/Qualifiers
 REGION 1..982
 note = MS_5xCD3-scFc
 source 1..982
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 284

QVQLVESGGG	LVKPGGSLRL	SCAASGFTFS	DHYMSWIRQA	PGKGLEWFSY	ISSSGGIIYY	60
ADSVKGRFTI	SRDNAKNSLY	LQMNSLRAED	TAVYYCARDV	GSHFDYWGQ	TLVTVSSGGG	120
GGGGGGSGGG	GSDIQMTQSP	SSVSASVGDR	VTITCRASQD	ISRWLAWYQQ	KPGKAPKLLI	180
SAASRLQSGV	PSRFGSGSGG	TDFTLTISSL	QPEDFAIYYC	QQAQSPRTF	GQGTKEIKS	240
GGGGSEVQLV	ESGGGLVQPG	GSLKLSKAAS	GFTFNKYAMN	WVRQAPGKGL	EWVARIRSKY	300
NNYATYYADS	VKDRFTISR	DSKNTAYLQM	NNLKTEDTAV	YVCVRHGNFG	NSYISYWAYW	360
GQGTLVTVSS	GGGGSGGGG	GGGGSGTVVT	QEPSTVSPG	GTVTLTCGSS	TGAVTSGNYP	420
NWVQKPGQA	PRGLIGGTKF	LAPGTPARFS	GSLLGKKAAL	TLSGVQPEDE	AEYYCVLWYS	480
NRWVFGGK	LTVLGGGDK	THTCPPCPAP	ELLGGPSVFL	FPPKPKDTLM	ISRTPEVTCV	540
VVDVSHEDPE	VKFNWYVDG	EVHNAKTKPC	EEQYGSTYRC	VSVLTVLHQD	WLNKEYKCK	600
VSNKALPAPI	EKTISKAKGQ	PREPQVYTL	PSREEMTKNQ	VSLTCLVKGF	YPSDIAVEWE	660
SNGQPENNYK	TTPPVLDSDG	SFFLYSKLTV	DKSRWQQGNV	FSCSVMHEAL	HNHYTQKSL	720
LSPGGGGSGG	GGGGSGGGG	GGGGSGGGG	GGGSDKTH	CPPCPAPELL	GGPSVFLFPP	780
KPKDTLMISR	TPEVTCVVVD	VSHEDPEVKF	NWYVDGVEVH	NAKTKPCEEQ	YGSTYRCVSV	840
LTVLHQDWLN	GKEYCKVSN	KALPAIEKT	ISKAKGQPRE	PQVYTLPPSR	EEMTKNQVSL	900
TCLVKGFYPS	DAVEWESNG	QPENNYKTT	PVLDSGGSFF	LYSKLTVDKS	RWQQGNVFS	960
SVMHEALHNH	YTQKSLSLSP	GK				982

SEQ ID NO: 285 moltype = AA length = 980
 FEATURE Location/Qualifiers
 REGION 1..980
 note = MS_5xCD3-scFc_delGK
 source 1..980
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 285

QVQLVESGGG	LVKPGGSLRL	SCAASGFTFS	DHYMSWIRQA	PGKGLEWFSY	ISSSGGIIYY	60
ADSVKGRFTI	SRDNAKNSLY	LQMNSLRAED	TAVYYCARDV	GSHFDYWGQ	TLVTVSSGGG	120
GGGGGGSGGG	GSDIQMTQSP	SSVSASVGDR	VTITCRASQD	ISRWLAWYQQ	KPGKAPKLLI	180
SAASRLQSGV	PSRFGSGSGG	TDFTLTISSL	QPEDFAIYYC	QQAQSPRTF	GQGTKEIKS	240
GGGGSEVQLV	ESGGGLVQPG	GSLKLSKAAS	GFTFNKYAMN	WVRQAPGKGL	EWVARIRSKY	300
NNYATYYADS	VKDRFTISR	DSKNTAYLQM	NNLKTEDTAV	YVCVRHGNFG	NSYISYWAYW	360
GQGTLVTVSS	GGGGSGGGG	GGGGSGTVVT	QEPSTVSPG	GTVTLTCGSS	TGAVTSGNYP	420
NWVQKPGQA	PRGLIGGTKF	LAPGTPARFS	GSLLGKKAAL	TLSGVQPEDE	AEYYCVLWYS	480
NRWVFGGK	LTVLGGGDK	THTCPPCPAP	ELLGGPSVFL	FPPKPKDTLM	ISRTPEVTCV	540
VVDVSHEDPE	VKFNWYVDG	EVHNAKTKPC	EEQYGSTYRC	VSVLTVLHQD	WLNKEYKCK	600
VSNKALPAPI	EKTISKAKGQ	PREPQVYTL	PSREEMTKNQ	VSLTCLVKGF	YPSDIAVEWE	660
SNGQPENNYK	TTPPVLDSDG	SFFLYSKLTV	DKSRWQQGNV	FSCSVMHEAL	HNHYTQKSL	720
LSPGGGGSGG	GGSGGGSGG	GGSGGGSGG	GGSDKTHTCP	PCPAPELLGG	PSVFLFPPKP	780
KDTLMISRTP	EVTCTVVDVS	HEDPEVKFNW	YVDGVEVHNA	KTKPCEEQYG	STYRCVSVLT	840
VLHQDWLNGK	EYKCKVSNKA	LPAPIEKTIS	KAKGQPREPQ	VYTLPPSREE	MTKNQVSLTC	900
LVKGFYPSDI	AVEWESNGQP	ENNYKTTTPV	LDSGGSFFLY	SKLTVDKSRW	QQGNVFS	960
MHEALHNHYT	QKSLSLSPGK					980

SEQ ID NO: 286 moltype = AA length = 982
 FEATURE Location/Qualifiers
 REGION 1..982
 note = MS_5_CCD3-scFc
 source 1..982
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 286

QVQLVESGGG	LVKPGGSLRL	SCAASGFTFS	DHYMSWIRQA	PGKCLEWFSY	ISSSGGIIYY	60
ADSVKGRFTI	SRDNAKNSLY	LQMNSLRAED	TAVYYCARDV	GSHFDYWGQ	TLVTVSSGGG	120
GGGGGGSGGG	GSDIQMTQSP	SSVSASVGDR	VTITCRASQD	ISRWLAWYQQ	KPGKAPKLLI	180
SAASRLQSGV	PSRFGSGSGG	TDFTLTISSL	QPEDFAIYYC	QQAQSPRTF	GQGTKEIKS	240
GGGGSEVQLV	ESGGGLVQPG	GSLKLSKAAS	GFTFNKYAMN	WVRQAPGKGL	EWVARIRSKY	300

-continued

```

NNYATYYADS VKDRFTISR DSKNTAYLQM NNLKTEDTAV YYCVRHGNFG NSYISYWAYW 360
GGQTLVTVSS GGGGSGGGGS GGGGSGTVVT QEPSLTVSPG GTVTLTCGSS TGAVTSGNYP 420
NWVQKPGQA PRGLIGGTKF LAPGTPARFS GSLGGKAAL TSGVQPEDE AEYCVLWYS 480
NRWVFGGK LTVLGGGDK THTCPAP ELLGGPSVFL FPPKPKDTLM ISRTPEVTCV 540
VVDVSHEDPE VKFNWYVDG EVHNAKTKPC EEQYGYSTYRC VSVLTVLHQD WLNKEYKCK 600
VSNKALPAPI EKTISKAKGQ PREPQVYTL PSREEMTKNQ VSLTCLVKGF YPSDIAVEWE 660
SNGQPENNYK TTPPVLDSDG SFPLYSKLTV DKSRWQQGNV FSCSVMHEAL HNHYTQKSL 720
LSPGKGGGS GGGGSGGGGS GGGGSGGGGS GGGGSDKTHT CPPCPAPELL GGPSVFLFPP 780
KPKDTLMISR TPEVTCVVVD VSHEDPEVKF NWYVDGVEVH NAKTKPCEEQ YGSTRYCVSV 840
LTVLHQDWLW GKEYKCKVSN KALPAPIEKT ISKAKGQPRE PQVYTLPPSR EEMTKNQVSL 900
TCLVKGFPYS DIAVEWESNG QPENNYKTP PVLDSDGSFF LYSKLTVDKS RWQQGNVFSC 960
SVMHEALHNH YTQKSLSLSP GK 982

```

```

SEQ ID NO: 287      moltype = AA length = 980
FEATURE            Location/Qualifiers
REGION             1..980
                   note = MS_5_CcCD3-scFc_delGK
source             1..980
                   mol_type = protein
                   organism = synthetic construct

```

```

SEQUENCE: 287
QVQLVESGGG LVKPGGSLRL SCAASGFTFS DHYMSWIRQA PGKCLEWFSY ISSSGGIYY 60
ADSVKGRFTI SRDNAKNSLY LQMNSLRAED TAVYYCARDV GSHFDYWGQG TLTVSSGGG 120
GGGGGGGGGS GSDIQTQSP SSVASVGD RVTITCRASQD ISRWLAWYQQ KPGKAPKLLI 180
SAASRLQSGV PSRFGSGSGG TDFTLTISSL QPEDFAIYYC QQAKSPRTF GCGTKVEIKS 240
GGGGSEVQLV ESGGGLVQPG GSKLSCAAS GFTFNKYAMN WVRQAPGKGL EWVARIRSKY 300
NNYATYYADS VKDRFTISR DSKNTAYLQM NNLKTEDTAV YYCVRHGNFG NSYISYWAYW 360
GGQTLVTVSS GGGGSGGGGS GGGGSGTVVT QEPSLTVSPG GTVTLTCGSS TGAVTSGNYP 420
NWVQKPGQA PRGLIGGTKF LAPGTPARFS GSLGGKAAL TSGVQPEDE AEYCVLWYS 480
NRWVFGGK LTVLGGGDK THTCPAP ELLGGPSVFL FPPKPKDTLM ISRTPEVTCV 540
VVDVSHEDPE VKFNWYVDG EVHNAKTKPC EEQYGYSTYRC VSVLTVLHQD WLNKEYKCK 600
VSNKALPAPI EKTISKAKGQ PREPQVYTL PSREEMTKNQ VSLTCLVKGF YPSDIAVEWE 660
SNGQPENNYK TTPPVLDSDG SFPLYSKLTV DKSRWQQGNV FSCSVMHEAL HNHYTQKSL 720
LSPGKGGGS GGGGSGGGGS GGGGSGGGGS GGGGSDKTHT CPPCPAPELL GGPSVFLFPP 780
KDTLMISRTP EVTCVVVDV HEDPEVKFNW YVDGVEVHNA KTKPCEEQYG STRYCVSVLT 840
VLHQDWLWNG EYKCKVSNKA LPAPIEKTIS KAKGQPREPQ VYTLPPSREE MTKNQVSLTC 900
LVKGFYPSDI AVEWESNGQP ENNYKTPPV LDSDGSFFLY SKLTVDKSRW QQGNVFSCSV 960
MHEALHNHYT QKSLSLSPGK 982

```

```

SEQ ID NO: 288      moltype = AA length = 982
FEATURE            Location/Qualifiers
REGION             1..982
                   note = MS_6xCD3-scFc
source             1..982
                   mol_type = protein
                   organism = synthetic construct

```

```

SEQUENCE: 288
QVQLVESGGG LVKPGGSLRL SCAASGFTFS DHYMSWIRQA PGKLEWISY ISNSGSIYY 60
VDSVKGRFTI SRDNAKNSLY LQMNSLRAED TAVYYCARDV RTAFDYWGQG TLTVSSGGG 120
GGGGGGGGGS GSDIQTQSP SSVASVGD RVTITCRASQS IGSWLAWYQQ KPGKAPNLLI 180
YAASRLQSGV PSRFGSGSGG TDFTLTISSL QPEDFATYYC QQANSFPRTF GQGTQVDIKS 240
GGGGSEVQLV ESGGGLVQPG GSKLSCAAS GFTFNKYAMN WVRQAPGKGL EWVARIRSKY 300
NNYATYYADS VKDRFTISR DSKNTAYLQM NNLKTEDTAV YYCVRHGNFG NSYISYWAYW 360
GGQTLVTVSS GGGGSGGGGS GGGGSGTVVT QEPSLTVSPG GTVTLTCGSS TGAVTSGNYP 420
NWVQKPGQA PRGLIGGTKF LAPGTPARFS GSLGGKAAL TSGVQPEDE AEYCVLWYS 480
NRWVFGGK LTVLGGGDK THTCPAP ELLGGPSVFL FPPKPKDTLM ISRTPEVTCV 540
VVDVSHEDPE VKFNWYVDG EVHNAKTKPC EEQYGYSTYRC VSVLTVLHQD WLNKEYKCK 600
VSNKALPAPI EKTISKAKGQ PREPQVYTL PSREEMTKNQ VSLTCLVKGF YPSDIAVEWE 660
SNGQPENNYK TTPPVLDSDG SFPLYSKLTV DKSRWQQGNV FSCSVMHEAL HNHYTQKSL 720
LSPGKGGGS GGGGSGGGGS GGGGSGGGGS GGGGSDKTHT CPPCPAPELL GGPSVFLFPP 780
KPKDTLMISR TPEVTCVVVD VSHEDPEVKF NWYVDGVEVH NAKTKPCEEQ YGSTRYCVSV 840
LTVLHQDWLW GKEYKCKVSN KALPAPIEKT ISKAKGQPRE PQVYTLPPSR EEMTKNQVSL 900
TCLVKGFPYS DIAVEWESNG QPENNYKTP PVLDSDGSFF LYSKLTVDKS RWQQGNVFSC 960
SVMHEALHNH YTQKSLSLSP GK 982

```

```

SEQ ID NO: 289      moltype = AA length = 980
FEATURE            Location/Qualifiers
REGION             1..980
                   note = MS_6xCD3-scFc_delGK
source             1..980
                   mol_type = protein
                   organism = synthetic construct

```

```

SEQUENCE: 289
QVQLVESGGG LVKPGGSLRL SCAASGFTFS DHYMSWIRQA PGKLEWISY ISNSGSIYY 60
VDSVKGRFTI SRDNAKNSLY LQMNSLRAED TAVYYCARDV RTAFDYWGQG TLTVSSGGG 120
GGGGGGGGGS GSDIQTQSP SSVASVGD RVTITCRASQS IGSWLAWYQQ KPGKAPNLLI 180

```

-continued

YAASSLQSGV	PSRFGSGSGG	TDFTLTISL	QPEDFATYYC	QQANSFPRTF	GQGTKVDIKS	240
GGGGSEVQLV	ESGGGLVQPG	GSLKLSCAAS	GFTFNKYAMN	WVRQAPGKGL	EWVARIRSKY	300
NNYATYYADS	VKDRFTISR	DSKNTAYLQM	NNLKTEDTAV	YVCVRHGNFG	NSYISYWAYW	360
GQGTLVTVSS	GGGGSGGGGS	GGGGSQTVVT	QEPSLTVSPG	GTVTLTCGSS	TGAVTSGNYP	420
NWVQKPGQA	PRGLIGGTFK	LAPGTPARFS	GSLLGKKAAL	TLSGVQPEDE	AEYYCVLWYS	480
NRWVFGGGTK	LTVLGGGGDK	THTCPPCPAP	ELLGGPSVFL	FPKPKDTLM	ISRTPEVTCV	540
VVDVSHEDPE	VKFNWYVDGV	EVHNAKTKPC	EEQYGSTYRC	VSVLTVLHQD	WLNKEYKCK	600
VSNKALPAPI	EKTISKAKGQ	PREPQVYTL	PSREEMTKNQ	VSLTCLVKGF	YPSDIAVEWE	660
SNGQPENNYK	TTPPVLDSDG	SFFLYSKLTV	DKSRWQQGNV	FSCSVMHEAL	HNHYTQKSLS	720
LSPGGGGSGG	GGSGGGSGG	GGSGGGSGG	GGSDKTHTCP	PCPAPELLGG	PSVFLFPPKP	780
KDTLMISRTP	EVTCTVVDVS	HEDPEVKFNW	YVDGVEVHNA	KTKPCEEQYG	STYRCVSVLT	840
VLHQDWLNGK	EYKCKVSNKA	LPAPIEKTIS	KAKGQPREPQ	VYTLPPSREE	MTKNQVSLTC	900
LVKGFYPSDI	AVEWESNGQP	ENNYKTPPV	LDSGGSFFLY	SKLTVDKSRW	QQGNVFSCSV	960
MHEALHNHYT	QKSLSLSPGK					980

SEQ ID NO: 290 moltype = AA length = 982
FEATURE Location/Qualifiers
REGION 1..982
 note = MS_6_CCxCD3-scFc
source 1..982
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 290

QVQLVESGGG	LVKPGGSLRL	SCAASGFTFS	DHYMSWIRQA	PGKCLEWISY	ISNSGSIYY	60
VDSVKGRFTI	SRDNAKNSLY	LQMNSLRAED	TAVYYCARDV	RTAFDYWGQG	TLVTVSSGGG	120
GSGGGSGGGG	GSDIQMTQSP	SSVSASVGDR	VTITCRASQS	IGSWLAWYQQ	KPGKAPNLLI	180
YAASSLQSGV	PSRFGSGSGG	TDFTLTISL	QPEDFATYYC	QQANSFPRTF	GCGTKVDIKS	240
GGGGSEVQLV	ESGGGLVQPG	GSLKLSCAAS	GFTFNKYAMN	WVRQAPGKGL	EWVARIRSKY	300
NNYATYYADS	VKDRFTISR	DSKNTAYLQM	NNLKTEDTAV	YVCVRHGNFG	NSYISYWAYW	360
GQGTLVTVSS	GGGGSGGGGS	GGGGSQTVVT	QEPSLTVSPG	GTVTLTCGSS	TGAVTSGNYP	420
NWVQKPGQA	PRGLIGGTFK	LAPGTPARFS	GSLLGKKAAL	TLSGVQPEDE	AEYYCVLWYS	480
NRWVFGGGTK	LTVLGGGGDK	THTCPPCPAP	ELLGGPSVFL	FPKPKDTLM	ISRTPEVTCV	540
VVDVSHEDPE	VKFNWYVDGV	EVHNAKTKPC	EEQYGSTYRC	VSVLTVLHQD	WLNKEYKCK	600
VSNKALPAPI	EKTISKAKGQ	PREPQVYTL	PSREEMTKNQ	VSLTCLVKGF	YPSDIAVEWE	660
SNGQPENNYK	TTPPVLDSDG	SFFLYSKLTV	DKSRWQQGNV	FSCSVMHEAL	HNHYTQKSLS	720
LSPGGGGSGG	GGSGGGSGG	GGSGGGSGG	GGGSDKTHT	CPPCPAPELL	GGPSVFLFPP	780
KPKDTLMISR	TPEVTCVVD	VSHEDPEVKF	NWYVDGVEVH	NAKTKPCEEQ	YGSTYRCVSV	840
LTVLHQDWLN	GKEYKCKVSN	KALPAPIEKT	ISKAKGQPRE	PQVYTLPPSR	EEMTKNQVSL	900
TCLVKGFPYS	DAVEWESNG	QPENNYKTP	PVLDSDGSFF	LYSKLTVDKS	RWQQGNVFSC	960
SVMHEALHNH	YTQKSLSLSP	GK				982

SEQ ID NO: 291 moltype = AA length = 980
FEATURE Location/Qualifiers
REGION 1..980
 note = MS_6_CCxCD3-scFc_delGK
source 1..980
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 291

QVQLVESGGG	LVKPGGSLRL	SCAASGFTFS	DHYMSWIRQA	PGKCLEWISY	ISNSGSIYY	60
VDSVKGRFTI	SRDNAKNSLY	LQMNSLRAED	TAVYYCARDV	RTAFDYWGQG	TLVTVSSGGG	120
GSGGGSGGGG	GSDIQMTQSP	SSVSASVGDR	VTITCRASQS	IGSWLAWYQQ	KPGKAPNLLI	180
YAASSLQSGV	PSRFGSGSGG	TDFTLTISL	QPEDFATYYC	QQANSFPRTF	GCGTKVDIKS	240
GGGGSEVQLV	ESGGGLVQPG	GSLKLSCAAS	GFTFNKYAMN	WVRQAPGKGL	EWVARIRSKY	300
NNYATYYADS	VKDRFTISR	DSKNTAYLQM	NNLKTEDTAV	YVCVRHGNFG	NSYISYWAYW	360
GQGTLVTVSS	GGGGSGGGGS	GGGGSQTVVT	QEPSLTVSPG	GTVTLTCGSS	TGAVTSGNYP	420
NWVQKPGQA	PRGLIGGTFK	LAPGTPARFS	GSLLGKKAAL	TLSGVQPEDE	AEYYCVLWYS	480
NRWVFGGGTK	LTVLGGGGDK	THTCPPCPAP	ELLGGPSVFL	FPKPKDTLM	ISRTPEVTCV	540
VVDVSHEDPE	VKFNWYVDGV	EVHNAKTKPC	EEQYGSTYRC	VSVLTVLHQD	WLNKEYKCK	600
VSNKALPAPI	EKTISKAKGQ	PREPQVYTL	PSREEMTKNQ	VSLTCLVKGF	YPSDIAVEWE	660
SNGQPENNYK	TTPPVLDSDG	SFFLYSKLTV	DKSRWQQGNV	FSCSVMHEAL	HNHYTQKSLS	720
LSPGGGGSGG	GGSGGGSGG	GGSGGGSGG	GGSDKTHTCP	PCPAPELLGG	PSVFLFPPKP	780
KDTLMISRTP	EVTCTVVDVS	HEDPEVKFNW	YVDGVEVHNA	KTKPCEEQYG	STYRCVSVLT	840
VLHQDWLNGK	EYKCKVSNKA	LPAPIEKTIS	KAKGQPREPQ	VYTLPPSREE	MTKNQVSLTC	900
LVKGFYPSDI	AVEWESNGQP	ENNYKTPPV	LDSGGSFFLY	SKLTVDKSRW	QQGNVFSCSV	960
MHEALHNHYT	QKSLSLSPGK					980

SEQ ID NO: 292 moltype = AA length = 989
FEATURE Location/Qualifiers
REGION 1..989
 note = MS_7xCD3-scFc
source 1..989
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 292

EVQLVESGGG	LIQPGGSLRL	SCAVSGFTVS	SKFMTWVRQA	PGKGLEWVSV	IYSGGKTYYA	60
------------	------------	------------	------------	------------	------------	----

-continued

```
DSVKGRFTIS RDNSKNTLYL QMNSLRAEDT AVYYCARD SG GWGYFDYWGQ GTLVTVSSGG 120
GGSGGGGGSGG GGS DIVMTQS PDSLAVSLGE RATINCKSSQ SVLYSSNNKN YLAWYQQKPG 180
QPPKLLIYWA STRESGVPDR FSGSGSGTDF TLTISSLQAE DVAVYYCQQY YSTPPTFGQG 240
TKVEIKSGGG GSEVQLVESG GGLVQPGGSL KLS CAASGFT FNKYAMNWVR QAPGKGLEWV 300
ARIRSKYNNY ATYYADSVKD RFTISRDDSK NTAYLQMNNL KTEDTAVYYC VRHGNFGNSY 360
ISYWAYWGQG TLVTVSSGGG GSGGGGSGGG GSQT VVTQEP SLTVSPGGTV TLTGSGSTGA 420
VTS GNYPNWV QQKPGQAPRG LIGGT KFLAP GTPARFSGSL LGGKAAL TLS GVQPEDEAEY 480
YCVLWYSNRW VFGGGTKLTV LGGGGDKTHT CPPCPAPELL GGPSVFLFPP KPKDTLMISR 540
TPEVTCVVVD VSHEDPEVKF NWYVDGVEVH NAKTKPCEEQ YGSTRYCVSV LTVLHQDWLN 600
GKEYKCKVSN KALPAPIEKT ISKAKGQPRE PQVYTLPPSR EEMTKNQVSL TCLVKGFYPS 660
DIAVEWESNG QPENNYK TTP PVLDSDGSFF LYSKLTVDKS RWQQGNVFSC SVMHEALHNNH 720
YTQKSLSLSP GKG GGGSGGGG GSGGGGSGGG GSGGGGSGGG GSKTHTCPP CPAPPELLGGP 780
SVFLFPPKPK DTL MISRTP VTCVVVDVSH EDPEVKFNWY VDGVEVHNAK TKPCEEQYGS 840
TYRCVSVLTV LHQDWLNGKE YKCKVSNKAL PAPIEKTISK AKGQPREPQV YTLPPSREEM 900
TKNQVSLTCL VKGFYPSDIA VEWESNGQPE NNYKTPPVLD DSDGSFFLYS KLTVDKSRWQ 960
QGNVFSCSVM HEALHNNHTQ KSLSLSPGK 989
```

```
SEQ ID NO: 293      moltype = AA length = 987
FEATURE            Location/Qualifiers
REGION              1..987
                    note = MS_7xCD3-scFc_delGK
source              1..987
                    mol_type = protein
                    organism = synthetic construct
```

```
SEQUENCE: 293
EVQLVESGGG LIQPGGSLRL SCAVSGFTVS SKFMTWVRQA PGKGLEWVSV IYSGGKTYYA 60
DSVKGRFTIS RDNSKNTLYL QMNSLRAEDT AVYYCARD SG GWGYFDYWGQ GTLVTVSSGG 120
GGSGGGGGSGG GGS DIVMTQS PDSLAVSLGE RATINCKSSQ SVLYSSNNKN YLAWYQQKPG 180
QPPKLLIYWA STRESGVPDR FSGSGSGTDF TLTISSLQAE DVAVYYCQQY YSTPPTFGQG 240
TKVEIKSGGG GSEVQLVESG GGLVQPGGSL KLS CAASGFT FNKYAMNWVR QAPGKGLEWV 300
ARIRSKYNNY ATYYADSVKD RFTISRDDSK NTAYLQMNNL KTEDTAVYYC VRHGNFGNSY 360
ISYWAYWGQG TLVTVSSGGG GSGGGGSGGG GSQT VVTQEP SLTVSPGGTV TLTGSGSTGA 420
VTS GNYPNWV QQKPGQAPRG LIGGT KFLAP GTPARFSGSL LGGKAAL TLS GVQPEDEAEY 480
YCVLWYSNRW VFGGGTKLTV LGGGGDKTHT CPPCPAPELL GGPSVFLFPP KPKDTLMISR 540
TPEVTCVVVD VSHEDPEVKF NWYVDGVEVH NAKTKPCEEQ YGSTRYCVSV LTVLHQDWLN 600
GKEYKCKVSN KALPAPIEKT ISKAKGQPRE PQVYTLPPSR EEMTKNQVSL TCLVKGFYPS 660
DIAVEWESNG QPENNYK TTP PVLDSDGSFF LYSKLTVDKS RWQQGNVFSC SVMHEALHNNH 720
YTQKSLSLSP GGGGSGGGGS GGGGSGGGGS GGGGSGGGGS GSKTHTCPPCP APPELLGGPSV 780
FLFPPKPKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD GVEVHNAKTK PCEEQYGSY 840
RCVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK GQPREPQVYT LPPSREEMTK 900
NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTPPVLDSD GSGFFLYSKL TVDKSRWQQG 960
NVFSCSVMHE ALHNNHTQKS LSLSPGK 987
```

```
SEQ ID NO: 294      moltype = AA length = 989
FEATURE            Location/Qualifiers
REGION              1..989
                    note = MS_7_CCD3-scFc
source              1..989
                    mol_type = protein
                    organism = synthetic construct
```

```
SEQUENCE: 294
EVQLVESGGG LIQPGGSLRL SCAVSGFTVS SKFMTWVRQA PGKCLEWVSV IYSGGKTYYA 60
DSVKGRFTIS RDNSKNTLYL QMNSLRAEDT AVYYCARD SG GWGYFDYWGQ GTLVTVSSGG 120
GGSGGGGGSGG GGS DIVMTQS PDSLAVSLGE RATINCKSSQ SVLYSSNNKN YLAWYQQKPG 180
QPPKLLIYWA STRESGVPDR FSGSGSGTDF TLTISSLQAE DVAVYYCQQY YSTPPTFGCG 240
TKVEIKSGGG GSEVQLVESG GGLVQPGGSL KLS CAASGFT FNKYAMNWVR QAPGKGLEWV 300
ARIRSKYNNY ATYYADSVKD RFTISRDDSK NTAYLQMNNL KTEDTAVYYC VRHGNFGNSY 360
ISYWAYWGQG TLVTVSSGGG GSGGGGSGGG GSQT VVTQEP SLTVSPGGTV TLTGSGSTGA 420
VTS GNYPNWV QQKPGQAPRG LIGGT KFLAP GTPARFSGSL LGGKAAL TLS GVQPEDEAEY 480
YCVLWYSNRW VFGGGTKLTV LGGGGDKTHT CPPCPAPELL GGPSVFLFPP KPKDTLMISR 540
TPEVTCVVVD VSHEDPEVKF NWYVDGVEVH NAKTKPCEEQ YGSTRYCVSV LTVLHQDWLN 600
GKEYKCKVSN KALPAPIEKT ISKAKGQPRE PQVYTLPPSR EEMTKNQVSL TCLVKGFYPS 660
DIAVEWESNG QPENNYK TTP PVLDSDGSFF LYSKLTVDKS RWQQGNVFSC SVMHEALHNNH 720
YTQKSLSLSP GKG GGGSGGGG GSGGGGSGGG GSGGGGSGGG GSKTHTCPP CPAPPELLGGP 780
SVFLFPPKPK DTL MISRTP VTCVVVDVSH EDPEVKFNWY VDGVEVHNAK TKPCEEQYGS 840
TYRCVSVLTV LHQDWLNGKE YKCKVSNKAL PAPIEKTISK AKGQPREPQV YTLPPSREEM 900
TKNQVSLTCL VKGFYPSDIA VEWESNGQPE NNYKTPPVLD DSDGSFFLYS KLTVDKSRWQ 960
QGNVFSCSVM HEALHNNHTQ KSLSLSPGK 989
```

```
SEQ ID NO: 295      moltype = AA length = 987
FEATURE            Location/Qualifiers
REGION              1..987
                    note = MS_7_CCD3-scFc_delGK
source              1..987
                    mol_type = protein
                    organism = synthetic construct
```

-continued

SEQUENCE: 295

```

EVQLVESGGG LIQPGGSLRL SCAVSGFTVS SKFMTWVRQA PGKCLEWVSV IYSGGKTYYA 60
DSVKGRFTIS RDNSKNTLYL QMNSLRAEDT AVYYCARDSDG GWGYFDYWGQ GTLVTVSSGG 120
GGSGGGGSGG GGSDIVMTQS PDSLAVSLGE RATINCKSSQ SVLYSSNNKN YLAWYQQKPG 180
QPPKLLIYWA STRESGVPDR FSGSGSGTDF TLTISLQAE DVAVYYCQQY YSTPPTFGCG 240
TKVEIKSGGG GSEVQLVESG GGLVQPGGSL KLSCAASGFT FNKYAMNWVR QAPGKGLEWV 300
ARIRSKYNNY ATYYADSVKD RFTISRDDSK NTAYLQMNNL KTEDTAVYYC VRHGNFGNSY 360
ISYWAYWGQG TLVTVSSGGG GSGGGGSGGG GSQTVVTQEP SLTVSPGGTV TLTGSSSTGA 420
VTSGNYPNWV QQKPGQAPRG LIGGTKFLAP GTPARFSGSL LGGKAALTLS GVQPEDEAEY 480
YCVLWYSNRW VFGGGTKLTV LGGGDKTHT CPPCPAPELL GGPSVFLEFP KPKDTLMISR 540
TPEVTCVVVD VSHEDPEVKF NWYVDGVEVH NAKTKPCEEQ YGSTRCVSV LTVLHQDWLN 600
GKEYKCKVSN KALPAPIEKT ISKAKGQPRE PQVYTLPPSR EEMTKNQVSL TCLVKGFYPS 660
DIAVEWESNG QPENNYKTTT PVLDSGGSFF LYSKLTVDKS RWQGNVFSC SVMHEALHNH 720
YTQKSLSLSP GGGGSGGGGS GGGGSGGGGS GGGGSGGGGS DKTHTCPPCP APELLGGPSV 780
FLFPPKPKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD GVEVHNATK PCIEQYGSTY 840
RCVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK GQPREPQVYT LPPSREEMTK 900
NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTTTPVLDS DGSFFLYSKL TVDKSRWQQG 960
NVFSCSVMHE ALHNHYTQKS LSLSPGK 987

```

SEQ ID NO: 296 moltype = AA length = 987

FEATURE Location/Qualifiers

REGION 1..987

note = MS_8xCD3-scFc

source 1..987

mol_type = protein

organism = synthetic construct

SEQUENCE: 296

```

QVQLQESGPG LVKPSSETLSL TCTVSGGSIS SYWNWIRQP PGKGLEWIGR IYNGNTYYN 60
PSLKSRTVIS GDTSKNQFSL KLSSTAAADT AVYYCARPKL GIDAFDIWQ GTMVTVSSGG 120
GGSGGGGSGG GGSQSVLTQP PSVSGAPGQR VTISCTGSSS NIGAGYDVHW YQQLPGTAPK 180
LLIYGNSNRP SGVPDRFSGS KSGTSASLAI TGLQAEDEAD YQCQSYDSSL SGWVFGGGTK 240
LTVLSGGGGS EVQLVESGGG LVQPGGSLKL SCAASGFTFN KYAMNWVRQA PGKGLEWVAR 300
IRSKYNNYAT YYADSVKDRF TISRDDSKNT AYLMNNLKT EDTAVYYCVR HGNFGNSYIS 360
YWAYWGQGT LTVTVSSGGGS GGGGSGGGGS QTVVTQEPSL TVSPGGTVTL TCGSSTGAVT 420
SGNYPNWVQQ KPGQAPRGLI GGTKFLAPGT PARFSGSLLG GKAALTLGTV QPEDEAEYYC 480
VLWYSNRWVF GGGTKLTVLG GGGDKTHTCP PCPAPELLGG PSVFLFPPKP KDTLMISRT 540
EVTCTVVVDV HEDPEVKFNW YVDGVEVHNA KTKPCEEQYG STYRCVSVLT VLHQDWLNGK 600
EYKCKVSNKA LPAPIEKTIS KAKGQPREPQ VYTLPPSREE MTKNQVSLTC LVKGFYPSDI 660
AVEWESNGQP ENNYKTTTPV LDSDGSFFLY SKLTVDKSRW QQGNVFSCSV MHEALHNHYT 720
QKSLSLSPGK GGGGSGGGGS GGGGSGGGGS DKTHTCPPCP APELLGGPSV 780
FLFPPKPKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD GVEVHNATK PCIEQYGSTY 840
RCVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK GQPREPQVYT LPPSREEMTK 900
NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTTTPVLDS DGSFFLYSKL TVDKSRWQQG 960
NVFSCSVMHE ALHNHYTQKS LSLSPGK 987

```

SEQ ID NO: 297 moltype = AA length = 985

FEATURE Location/Qualifiers

REGION 1..985

note = MS_8xCD3-scFc_delGK

source 1..985

mol_type = protein

organism = synthetic construct

SEQUENCE: 297

```

QVQLQESGPG LVKPSSETLSL TCTVSGGSIS SYWNWIRQP PGKGLEWIGR IYNGNTYYN 60
PSLKSRTVIS GDTSKNQFSL KLSSTAAADT AVYYCARPKL GIDAFDIWQ GTMVTVSSGG 120
GGSGGGGSGG GGSQSVLTQP PSVSGAPGQR VTISCTGSSS NIGAGYDVHW YQQLPGTAPK 180
LLIYGNSNRP SGVPDRFSGS KSGTSASLAI TGLQAEDEAD YQCQSYDSSL SGWVFGGGTK 240
LTVLSGGGGS EVQLVESGGG LVQPGGSLKL SCAASGFTFN KYAMNWVRQA PGKGLEWVAR 300
IRSKYNNYAT YYADSVKDRF TISRDDSKNT AYLMNNLKT EDTAVYYCVR HGNFGNSYIS 360
YWAYWGQGT LTVTVSSGGGS GGGGSGGGGS QTVVTQEPSL TVSPGGTVTL TCGSSTGAVT 420
SGNYPNWVQQ KPGQAPRGLI GGTKFLAPGT PARFSGSLLG GKAALTLGTV QPEDEAEYYC 480
VLWYSNRWVF GGGTKLTVLG GGGDKTHTCP PCPAPELLGG PSVFLFPPKP KDTLMISRT 540
EVTCTVVVDV HEDPEVKFNW YVDGVEVHNA KTKPCEEQYG STYRCVSVLT VLHQDWLNGK 600
EYKCKVSNKA LPAPIEKTIS KAKGQPREPQ VYTLPPSREE MTKNQVSLTC LVKGFYPSDI 660
AVEWESNGQP ENNYKTTTPV LDSDGSFFLY SKLTVDKSRW QQGNVFSCSV MHEALHNHYT 720
QKSLSLSPGK GGGGSGGGGS GGGGSGGGGS GGGGSGGGSDK THTCPPCPAP ELLGGPSVFL 780
FPPKPKDTLM ISRTPEVTCV VVDVSHEDPE VKFNWYVDGV EVHNATKPC EEQYGSTYRC 840
VSVLTVLHQD WLNKEYKCK VSNKALPAPI EKTISKAKGQ PREPQVYTL PPSREEMTKNQ 900
VSLTCLVKGF YPSDIAVEWE SNGQPENNYK TTPPVLDSDG SFFLYSKLTV DKSRWQQGNV 960
FSCSVMHEAL HNHYTQKSLS LSPGK 985

```

SEQ ID NO: 298 moltype = AA length = 987

FEATURE Location/Qualifiers

REGION 1..987

note = MS_8_CCxCD3-scFc

source 1..987

-continued

```

mol_type = protein
organism = synthetic construct

SEQUENCE: 298
QVQLQESGPG LVKPSSETLSL TCTVSGGSIS SYWNWIRQP PGKCLEWIGR IYNGNTYYN 60
PSLKSRTVIS GDTSKNQFSL KLSSTAAADT AVYYCARPKL GIDAFDIWQ GTMTVSSGG 120
GGSGGGGGSG GGSQSVLTQP PSVSGAPGQR VTISCTGSSS NIGAGYDVHW YQQLPGTAPK 180
LLIYGNSNRP SGVPDRFSGS KSGTSASLAI TGLQAEDEAD YQCQSYDSSL SGWVFGCGTK 240
LTVLSGGGGS EVQLVESGGG LVQPGGSLKL SCAASGFTFN KYAMNWVRQA PGKLEWVAR 300
IRSKYNNYAT YYADSVKDRF TISRDDSKNT AYLMNNLKT EDTAVYYCVR HGNFGNSYIS 360
YWAYWQGGTL VTVSSGGGGS GGGSGGGGGS QTVVTQEP SL TVSPGGTVTL TCGSSTGAVT 420
SGNYPNWVQQ KPGQAPRGLI GGTKFLAPGT PARFSGSLG GKAALTL SGV QPEDEAEYYC 480
VLWYSNRWVF GGGTKLTVLG GGGDKTHTCP PCPAPELLGG PSVFLFPPKP KDTLMISRT 540
EVTCTVVVDV HEDPEVKFNW YVDGVEVHNA KTKPCEEQYG STYRCVSVLT VLHQDWLNGK 600
EYKCKVSNKA LPAPIEKTIS KAKGQPREPQ VYTLPPSREE MTKNQVSLTC LVKGFYPSDI 660
AVEWESNGQP ENNYKTTTPPV LDSDGSEFFLY SKLTVDKSRW QQGNVFSCSV MHEALHNHYT 720
QKSLSLSPGK GGSQSVLTQP PSVSGAPGQR VTISCTGSSS NIGAGYDVHW YQQLPGTAPK 780
FLFPPKPKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD GVEVHNAKTK PCCEEQYGSTY 840
RCVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK GQPREPQVYT LPPSREEMTK 900
NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTTTPPVLD SPSFLYSKLT TVDKSRWQQG 960
NVFSCSVMEH ALHNHYTQKS LSLSPGK 987

SEQ ID NO: 299      moltype = AA length = 985
FEATURE             Location/Qualifiers
REGION              1..985
                    note = MS_8_CCxCD3-scFc_delGK
source              1..985
                    mol_type = protein
                    organism = synthetic construct

SEQUENCE: 299
QVQLQESGPG LVKPSSETLSL TCTVSGGSIS SYWNWIRQP PGKCLEWIGR IYNGNTYYN 60
PSLKSRTVIS GDTSKNQFSL KLSSTAAADT AVYYCARPKL GIDAFDIWQ GTMTVSSGG 120
GGSGGGGGSG GGSQSVLTQP PSVSGAPGQR VTISCTGSSS NIGAGYDVHW YQQLPGTAPK 180
LLIYGNSNRP SGVPDRFSGS KSGTSASLAI TGLQAEDEAD YQCQSYDSSL SGWVFGCGTK 240
LTVLSGGGGS EVQLVESGGG LVQPGGSLKL SCAASGFTFN KYAMNWVRQA PGKLEWVAR 300
IRSKYNNYAT YYADSVKDRF TISRDDSKNT AYLMNNLKT EDTAVYYCVR HGNFGNSYIS 360
YWAYWQGGTL VTVSSGGGGS GGGSGGGGGS QTVVTQEP SL TVSPGGTVTL TCGSSTGAVT 420
SGNYPNWVQQ KPGQAPRGLI GGTKFLAPGT PARFSGSLG GKAALTL SGV QPEDEAEYYC 480
VLWYSNRWVF GGGTKLTVLG GGGDKTHTCP PCPAPELLGG PSVFLFPPKP KDTLMISRT 540
EVTCTVVVDV HEDPEVKFNW YVDGVEVHNA KTKPCEEQYG STYRCVSVLT VLHQDWLNGK 600
EYKCKVSNKA LPAPIEKTIS KAKGQPREPQ VYTLPPSREE MTKNQVSLTC LVKGFYPSDI 660
AVEWESNGQP ENNYKTTTPPV LDSDGSEFFLY SKLTVDKSRW QQGNVFSCSV MHEALHNHYT 720
QKSLSLSPGK GGSQSVLTQP PSVSGAPGQR VTISCTGSSS NIGAGYDVHW YQQLPGTAPK 780
FPPKPKDTLM ISRTPEVTCV VVDVSHEDPE VKFNWYVDG EVHNAKTKPC EEQYGSTYRC 840
VSVLTVLHQA WLNKEYKCK VSNKALPAPI EKTISKAKGQ PREPQVYTL PSREEMTKNQ 900
VSLTCLVKGF YPSDIAVEWE SNGQPENNYK TTPPVLDSDG SFFLYSKLTV DKSRWQQGNV 960
FSCSVMEHAL HNHYTQKSLS LSPGK 985

SEQ ID NO: 300      moltype = AA length = 20
FEATURE             Location/Qualifiers
REGION              1..20
                    note = (G4S)4 linker
source              1..20
                    mol_type = protein
                    organism = synthetic construct

SEQUENCE: 300
GGGSGGGGGS GGGSGGGGGS 20

SEQ ID NO: 301      moltype = AA length = 25
FEATURE             Location/Qualifiers
REGION              1..25
                    note = (G4S)5 linker
source              1..25
                    mol_type = protein
                    organism = synthetic construct

SEQUENCE: 301
GGGSGGGGGS GGGSGGGGGS GGGGS 25

SEQ ID NO: 302      moltype = AA length = 30
FEATURE             Location/Qualifiers
REGION              1..30
                    note = (G4S)6 linker
source              1..30
                    mol_type = protein
                    organism = synthetic construct

SEQUENCE: 302
GGGSGGGGGS GGGSGGGGGS GGGSGGGGGS 30

```

SEQ ID NO: 303	multype = AA length = 35		
FEATURE	Location/Qualifiers		
REGION	1..35		
	note = (G4S)7 linker		
source	1..35		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 303			
GGGSGGGGS GGGSGGGGS GGGSGGGGS GGGGS			35
SEQ ID NO: 304	multype = AA length = 40		
FEATURE	Location/Qualifiers		
REGION	1..40		
	note = (G4S)8 linker		
source	1..40		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 304			
GGGSGGGGS GGGSGGGGS GGGSGGGGS GGGSGGGGS			40
SEQ ID NO: 305	multype = AA length = 109		
FEATURE	Location/Qualifiers		
REGION	1..109		
	note = Synthetic Peptide		
source	1..109		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 305			
ELVVTQEPSL TVSPGGTVTL TCGSSTGAVT SGYYPNWVQQ KPGQAPRGLI GGTKFLAPGT			60
PARFSGSLLG GKAALTLSGV QPEDEAEYYC ALWYSNRWVF GGGTKLTVL			109
SEQ ID NO: 306	multype = AA length = 109		
FEATURE	Location/Qualifiers		
REGION	1..109		
	note = Synthetic Peptide		
source	1..109		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 306			
ELVVTQEPSL TVSPGGTVTL TCRSSTGAVT SGYYPNWVQQ KPGQAPRGLI GATDMRPSGT			60
PARFSGSLLG GKAALTLSGV QPEDEAEYYC ALWYSNRWVF GGGTKLTVL			109
SEQ ID NO: 307	multype = AA length = 109		
FEATURE	Location/Qualifiers		
REGION	1..109		
	note = Synthetic Peptide		
source	1..109		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 307			
ELVVTQEPSL TVSPGGTVTL TCGSSTGAVT SGNYPNWVQQ KPGQAPRGLI GGTKFLAPGT			60
PARFSGSLLG GKAALTLSGV QPEDEAEYYC VLWYSNRWVF GGGTKLTVL			109
SEQ ID NO: 308	multype = AA length = 125		
FEATURE	Location/Qualifiers		
REGION	1..125		
	note = Synthetic Peptide		
source	1..125		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 308			
EVQLESGGG LVQPGGSLKL SCAASGFTFN IYAMNWVRQA PGKGLEWVAR IRSKYNNYAT			60
YYADSVKSRLF TISRDDSKNT AYLQMNNLKT EDTAVYYCVR HGNFGNSYVS FFAYWGQGTL			120
VTVSS			125
SEQ ID NO: 309	multype = AA length = 125		
FEATURE	Location/Qualifiers		
REGION	1..125		
	note = Synthetic Peptide		
source	1..125		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 309			
EVQLESGGG LVQPGGSLKL SCAASGFTFN KYAMNWVRQA PGKGLEWVAR IRSKYNNYAT			60
YYADSVKDRF TISRDDSKNT AYLQMNNLKT EDTAVYYCVR HGNFGNSYIS YWAYWGQGTL			120
			125

-continued

VTVSS		125
SEQ ID NO: 310		
FEATURE	moltype = AA length = 125	
REGION	Location/Qualifiers	
	1..125	
	note = Synthetic peptide	
source	1..125	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 310		
EVQLLESGGG	LEQPGGSLKL	60
YYADSVKGRF	TISRDDSKNT	120
VTVSS		125
SEQ ID NO: 311		
FEATURE	moltype = AA length = 125	
REGION	Location/Qualifiers	
	1..125	
	note = Synthetic Peptide	
source	1..125	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 311		
EVQLLESGGG	LVQPGGSLKL	60
YYADSVKGRF	TISRDDSKNT	120
VTVSS		125
SEQ ID NO: 312		
FEATURE	moltype = AA length = 125	
REGION	Location/Qualifiers	
	1..125	
	note = Synthetic Peptide	
source	1..125	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 312		
EVQLLESGGG	LVQPGGSLKL	60
YYADSVKGRF	TISRDDSKNT	120
VTVSS		125
SEQ ID NO: 313		
FEATURE	moltype = AA length = 125	
REGION	Location/Qualifiers	
	1..125	
	note = Synthetic Peptide	
source	1..125	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 313		
EVQLLESGGG	LVQPGGSLKL	60
YYADSVKSRF	TISRDDSKNT	120
VTVSS		125
SEQ ID NO: 314		
FEATURE	moltype = AA length = 125	
REGION	Location/Qualifiers	
	1..125	
	note = Synthetic Peptide	
source	1..125	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 314		
EVQLLESGGG	LVQPGGSLKL	60
YYADSVKGRF	TISRDDSKNT	120
VTVSS		125
SEQ ID NO: 315		
FEATURE	moltype = AA length = 125	
REGION	Location/Qualifiers	
	1..125	
	note = Synthetic Peptide	
source	1..125	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 315		
EVQLLESGGG	LVQPGGSLKL	60
YYADSVKKRF	TISRDDSKNT	120
VTVSS		125
SEQ ID NO: 316		
FEATURE	moltype = AA length = 125	
	Location/Qualifiers	

-continued

REGION 1..125
 note = Synthetic peptide
 source 1..125
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 316
 EVQLLES GGG LVQPGGSLKL SCAASGFTFN SYAMNWVRQA PGKGLEWVAR IRSKYNNYAT 60
 YYADSVKGRF TISRDDSKNT AYLQMNLLKT EDTAVYYCVR HGNFGNSYVS WWAYWGQGTL 120
 VTVSS 125
 SEQ ID NO: 317 moltype = AA length = 125
 FEATURE Location/Qualifiers
 REGION 1..125
 note = Synthetic Peptide
 source 1..125
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 317
 EVQLLES GGG LVQPGGSLKL SCAASGFTFN KYAMNWVRQA PGKGLEWVAR IRSKYNNYAT 60
 YYADSVKDRF TISRDDSKNT AYLQMNLLKT EDTAVYYCVR HGNFGNSYIS YWAYWGQGTL 120
 VTVSS 125
 SEQ ID NO: 318 moltype = AA length = 249
 FEATURE Location/Qualifiers
 REGION 1..249
 note = Synthetic Peptide
 source 1..249
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 318
 EVQLLES GGG LVQPGGSLKL SCAASGFTFN IYAMNWVRQA PGKGLEWVAR IRSKYNNYAT 60
 YYADSVKSRF TISRDDSKNT AYLQMNLLKT EDTAVYYCVR HGNFGNSYVS FFAYWGQGTL 120
 VTVSSGGGGS GGGGSGGGGS ELVVTQEPSL TVSPGGTVTL TCGSSTGAVT SGYYPNWVQQ 180
 KPGQAPRGLI GGTKFLAPGT PARFSGSLLG GKAAITLSGV QPEDEAEYYC ALWYSNRWVF 240
 GGGTKLTVL 249
 SEQ ID NO: 319 moltype = AA length = 249
 FEATURE Location/Qualifiers
 REGION 1..249
 note = Synthetic Peptide
 source 1..249
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 319
 EVQLLES GGG LVQPGGSLKL SCAASGFTFN KYAMNWVRQA PGKGLEWVAR IRSKYNNYAT 60
 YYADSVKDRF TISRDDSKNT AYLQMNLLKT EDTAVYYCVR HGNFGNSYIS YWAYWGQGTL 120
 VTVSSGGGGS GGGGSGGGGS ELVVTQEPSL TVSPGGTVTL TCGSSTGAVT SGYYPNWVQQ 180
 KPGQAPRGLI GGTKFLAPGT PARFSGSLLG GKAAITLSGV QPEDEAEYYC ALWYSNRWVF 240
 GGGTKLTVL 249
 SEQ ID NO: 320 moltype = AA length = 249
 FEATURE Location/Qualifiers
 REGION 1..249
 note = Synthetic Peptide
 source 1..249
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 320
 EVQLLES GGG LEQPGGSLKL SCAASGFTFN SYAMNWVRQA PGKGLEWVAR IRSKYNNYAT 60
 YYADSVKGRF TISRDDSKNT AYLQMNLLKT EDTAVYYCVR HGNFGNSYLS FWAYWGQGTL 120
 VTVSSGGGGS GGGGSGGGGS ELVVTQEPSL TVSPGGTVTL TCGSSTGAVT SGYYPNWVQQ 180
 KPGQAPRGLI GGTKFLAPGT PARFSGSLLG GKAAITLSGV QPEDEAEYYC ALWYSNRWVF 240
 GGGTKLTVL 249
 SEQ ID NO: 321 moltype = AA length = 249
 FEATURE Location/Qualifiers
 REGION 1..249
 note = Synthetic Peptide
 source 1..249
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 321
 EVQLLES GGG LVQPGGSLKL SCAASGFTFN RYAMNWVRQA PGKGLEWVAR IRSKYNNYAT 60
 YYADSVKGRF TISRDDSKNT AYLQMNLLKT EDTAVYYCVR HGNFGNSYLS YFAYWGQGTL 120
 VTVSSGGGGS GGGGSGGGGS ELVVTQEPSL TVSPGGTVTL TCGSSTGAVT SGYYPNWVQQ 180
 KPGQAPRGLI GGTKFLAPGT PARFSGSLLG GKAAITLSGV QPEDEAEYYC ALWYSNRWVF 240

-continued

GGGTKLTVL 249

SEQ ID NO: 322 moltype = AA length = 249
 FEATURE Location/Qualifiers
 REGION 1..249
 note = Synthetic Peptide
 source 1..249
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 322

EVQLLES GGG LVQPGGSLKL	SCAASGFTFN	VYAMNWRQA	PGKGLEWVAR	IRSKYNNYAT	60
YYADSVKKRF	TISRDDSKNT	AYLQMNNLKT	EDTAVYYCVR	HGNFGNSYLS	120
VTVSSGGGGS	GGGGSGGGGS	ELVVTQEPSL	TVSPGGTVTL	TCRSSTGAVT	180
KPGQAPRGLI	GATDMRPSGT	PARFSGSLLG	GKAALTLSGV	QPEDEAEYYC	240
GGGTKLTVL					249

SEQ ID NO: 323 moltype = AA length = 249
 FEATURE Location/Qualifiers
 REGION 1..249
 note = Synthetic Peptide
 source 1..249
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 323

EVQLLES GGG LVQPGGSLKL	SCAASGFTFN	KYAMNWRQA	PGKGLEWVAR	IRSKYNNYAT	60
YYADSVKSRF	TISRDDSKNT	AYLQMNNLKT	EDTAVYYCVR	HGNFGNSYLS	120
VTVSSGGGGS	GGGGSGGGGS	ELVVTQEPSL	TVSPGGTVTL	TCGSSTGAVT	180
KPGQAPRGLI	GGTKFLAPGT	PARFSGSLLG	GKAALTLSGV	QPEDEAEYYC	240
GGGTKLTVL					249

SEQ ID NO: 324 moltype = AA length = 249
 FEATURE Location/Qualifiers
 REGION 1..249
 note = Synthetic Peptide
 source 1..249
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 324

EVQLLES GGG LVQPGGSLKL	SCAASGFTFN	GYAMNWRQA	PGKGLEWVAR	IRSKYNNYAT	60
YYADSVKKRF	TISRDDSKNT	AYLQMNNLKT	EDTAVYYCVR	HRNFGNSYLS	120
VTVSSGGGGS	GGGGSGGGGS	ELVVTQEPSL	TVSPGGTVTL	TCRSSTGAVT	180
KPGQAPRGLI	GATDMRPSGT	PARFSGSLLG	GKAALTLSGV	QPEDEAEYYC	240
GGGTKLTVL					249

SEQ ID NO: 325 moltype = AA length = 249
 FEATURE Location/Qualifiers
 REGION 1..249
 note = Synthetic Peptide
 source 1..249
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 325

EVQLLES GGG LVQPGGSLKL	SCAASGFTFN	VYAMNWRQA	PGKGLEWVAR	IRSKYNNYAT	60
YYADSVKKRF	TISRDDSKNT	AYLQMNNLKT	EDTAVYYCVR	HGNFGNSYIS	120
VTVSSGGGGS	GGGGSGGGGS	ELVVTQEPSL	TVSPGGTVTL	TCGSSTGAVT	180
KPGQAPRGLI	GGTKFLAPGT	PARFSGSLLG	GKAALTLSGV	QPEDEAEYYC	240
GGGTKLTVL					249

SEQ ID NO: 326 moltype = AA length = 249
 FEATURE Location/Qualifiers
 REGION 1..249
 note = Synthetic Peptide
 source 1..249
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 326

EVQLLES GGG LVQPGGSLKL	SCAASGFTFN	SYAMNWRQA	PGKGLEWVAR	IRSKYNNYAT	60
YYADSVKGRF	TISRDDSKNT	AYLQMNNLKT	EDTAVYYCVR	HGNFGNSYVS	120
VTVSSGGGGS	GGGGSGGGGS	ELVVTQEPSL	TVSPGGTVTL	TCGSSTGAVT	180
KPGQAPRGLI	GGTKFLAPGT	PARFSGSLLG	GKAALTLSGV	QPEDEAEYYC	240
GGGTKLTVL					249

SEQ ID NO: 327 moltype = AA length = 249
 FEATURE Location/Qualifiers
 REGION 1..249
 note = Synthetic Peptide

-continued

```

source          1..249
                mol_type = protein
                organism = synthetic construct

SEQUENCE: 327
EVQLLESGGG LVQPGGSLKL SCAASGFTFN SYAMNWRQA PGKGLEWVAR IRSKYNNYAT 60
YYADSVKGRF TISRDDSKNT AYLQMNLLKT EDTAVYYCVR HGNFGNSYVS WWAYWGQGTL 120
VTVSSGGGGS GGGGSGGGGS ELVVTQEPSL TVSPGGTVTL TCGSSTGAVT SGNYPNWVQQ 180
KPGQAPRGLI GGTKFLAPGT PARFSGSLLG GKAAITLSGV QPEDEAEYYC VLWYSNRWVF 240
GGGTKLTVL                                     249

```

1-22. (canceled)

23. A bispecific antibody construct comprising a first binding domain which binds to human and macaque MSLN on the surface of a target cell and a second binding domain which binds to human CD3 on the surface of a T cell, wherein the first binding domain binds to an epitope of MSLN which is comprised within the region of the human MSLN having a sequence having the sequence of SEQ ID NO: 241, and wherein the first binding domain binds to the same epitope of MSLN as an antibody comprising a VH region comprising CDR-H1 having the sequence of SEQ ID NO: 211, CDR-H2 having the sequence of SEQ ID NO: 212, CDR-H3 having the sequence of SEQ ID NO: 213 and a VL region comprising CDR-L1 having the sequence of SEQ ID NO: 214, CDR-L2 having the sequence of SEQ ID NO: 215 and CDR-L3 having the sequence of SEQ ID NO: 216.

24. The antibody construct of claim **23**, wherein the first binding domain comprises a VH region comprising CDR-H1 having the sequence of SEQ ID NO: 211, CDR-H2 having the sequence of SEQ ID NO: 212, CDR-H3 having the sequence of SEQ ID NO: 213 and a VL region comprising CDR-L1 having the sequence of SEQ ID NO: 214, CDR-L2 having the sequence of SEQ ID NO: 215 and CDR-L3 having the sequence of SEQ ID NO: 216.

25. The antibody construct of claim **23**, wherein the second binding domain binds to human CD3 epsilon and to *Callithrix jacchus*, *Saguinus oedipus* or *Saimiri sciureus* CD3 epsilon.

26. The antibody construct of claim **23**, wherein the antibody construct is in a format of a (scFv)₂, a scFv-single domain mAb, or a diabody, or an oligomer of any of the (scFv)₂, scFv-single domain mAb, or diabody formats.

27. The antibody construct of claim **23**, wherein the first binding domain comprises a VH region comprising the sequence of SEQ ID NO: 217, and/or wherein the first binding domain comprises a VL region comprising the sequence of SEQ ID NO: 218, or wherein the first binding domain comprises a VH region and a VL region comprising the sequence of SEQ ID NO: 217 and 218, or wherein the first binding domain comprises a polypeptide comprising the sequence of SEQ ID NO: 219.

28. The antibody construct of claim **23**, comprising:

(a) a polypeptide comprising in the following order starting from the N-terminus:

a polypeptide having an amino acid sequence of SEQ ID NO: 219;

a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 1-9; and

a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 19, SEQ ID NO: 28, SEQ ID NO: 37, SEQ ID NO: 46, SEQ ID NO: 55, SEQ ID NO: 64, SEQ ID NO: 73, SEQ ID NO: 82, SEQ ID NO: 91, SEQ ID NO: 100, and SEQ ID NO: 103; and

optionally a His-tag, such as the one depicted in SEQ ID NO: 10;

(b) a polypeptide comprising in the following order starting from the N-terminus:

a polypeptide having an amino acid sequence of SEQ ID NO: 219;

a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 1-9;

a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 19, SEQ ID NO: 28, SEQ ID NO: 37, SEQ ID NO: 46, SEQ ID NO: 55, SEQ ID NO: 64, SEQ ID NO: 73, SEQ ID NO: 82, SEQ ID NO: 91, SEQ ID NO: 100, and SEQ ID NO: 103;

optionally a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 1-9;

a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NOs: 104-134; and

optionally a His-tag, such as the one depicted in SEQ ID NO: 10;

(c) a polypeptide comprising in the following order starting from the N-terminus:

a polypeptide having the amino acid sequence QRFVTGHFGGLX1PANG (SEQ ID NO: 135) whereas X1 is Y or H; and

a polypeptide having an amino acid sequence of SEQ ID NO: 219;

a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 1-9;

a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 19, SEQ ID NO: 28, SEQ ID NO: 37, SEQ ID NO: 46, SEQ ID NO: 55, SEQ ID NO: 64, SEQ ID NO: 73, SEQ ID NO: 82, SEQ ID NO: 91, SEQ ID NO: 100, and SEQ ID NO: 103;

a polypeptide having the amino acid sequence QRFVTGHFGGLHPANG (SEQ ID NO: 137) or QRFCTGHFGGLHPCNG (SEQ ID NO: 139); and optionally a His-tag, such as the one depicted in SEQ ID NO: 10;

- (d) a polypeptide comprising in the following order starting from the N-terminus:
- a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 17, SEQ ID NO: 26, SEQ ID NO: 35, SEQ ID NO: 44, SEQ ID NO: 53, SEQ ID NO: 62, SEQ ID NO: 71, SEQ ID NO: 80, SEQ ID NO: 89, SEQ ID NO: 98, and SEQ ID NO: 101;
 - a peptide linker having the amino acid sequence depicted in SEQ ID NO: 8;
 - a polypeptide having an amino acid sequence of SEQ ID NO: 218 and a serine residue at the C-terminus;
 - a polypeptide having the amino acid sequence depicted in SEQ ID NO: 140; and
 - a polypeptide comprising in the following order starting from the N terminus:
 - a polypeptide having an amino acid sequence of SEQ ID NO: 217;
 - a peptide linker having the amino acid sequence depicted in SEQ ID NO: 8;
 - a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 18, SEQ ID NO: 27, SEQ ID NO: 36, SEQ ID NO: 45, SEQ ID NO: 54, SEQ ID NO: 63, SEQ ID NO: 72, SEQ ID NO: 81, SEQ ID NO: 90, SEQ ID NO: 99, and SEQ ID NO: 102 and a serine residue at the C-terminus;
 - a polypeptide having the amino acid sequence depicted in SEQ ID NO: 141;
- (e) a polypeptide comprising in the following order starting from the N-terminus:
- a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 17, SEQ ID NO: 26, SEQ ID NO: 35, SEQ ID NO: 44, SEQ ID NO: 53, SEQ ID NO: 62, SEQ ID NO: 71, SEQ ID NO: 80, SEQ ID NO: 89, SEQ ID NO: 98, and SEQ ID NO: 101;
 - a peptide linker having the amino acid sequence depicted in SEQ ID NO: 8;
 - a polypeptide having an amino acid sequence of SEQ ID NO: 218;
 - a polypeptide having the amino acid sequence depicted in SEQ ID NO: 142; and
 - a polypeptide comprising in the following order starting from the N-terminus:
 - a polypeptide having an amino acid sequence of SEQ ID NO: 217;
 - a peptide linker having an amino acid sequence depicted in SEQ ID NO: 8;
 - a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 18, SEQ ID NO: 27, SEQ ID NO: 36, SEQ ID NO: 45, SEQ ID NO: 54, SEQ ID NO: 63, SEQ ID NO: 72, SEQ ID NO: 81, SEQ ID NO: 90, SEQ ID NO: 99, and SEQ ID NO: 102 and a serine residue at the C-terminus;
 - a polypeptide having the amino acid sequence depicted in SEQ ID NO: 143;
- (f) a polypeptide comprising in the following order starting from the N-terminus:
- a polypeptide having an amino acid sequence of SEQ ID NO: 219;
 - a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 1-9; and
 - a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 19, SEQ ID NO: 28, SEQ ID NO: 37, SEQ ID NO: 46, SEQ ID NO: 55, SEQ ID NO: 64, SEQ ID NO: 73, SEQ ID NO: 82, SEQ ID NO: 91, SEQ ID NO: 100, and SEQ ID NO: 103; and
 - a polypeptide having the amino acid sequence depicted in SEQ ID NO: 144; and
 - a polypeptide having the amino acid sequence depicted in SEQ ID NO: 145;
- (g) a polypeptide comprising in the following order starting from the N-terminus:
- a polypeptide having an amino acid sequence of SEQ ID NO: 219; and
 - a polypeptide having the amino acid sequence depicted in SEQ ID NO: 146; and
 - a polypeptide comprising in the following order starting from the N-terminus:
 - a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 19, SEQ ID NO: 28, SEQ ID NO: 37, SEQ ID NO: 46, SEQ ID NO: 55, SEQ ID NO: 64, SEQ ID NO: 73, SEQ ID NO: 82, SEQ ID NO: 91, SEQ ID NO: 100, and SEQ ID NO: 103; and
 - a polypeptide having the amino acid sequence depicted in SEQ ID NO: 147;
- (h) a polypeptide comprising in the following order starting from the N-terminus:
- a polypeptide having an amino acid sequence of SEQ ID NO: 219; and
 - a polypeptide having the amino acid sequence depicted in SEQ ID NO: 148; and
 - a polypeptide comprising in the following order starting from the N-terminus:
 - a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 19, SEQ ID NO: 28, SEQ ID NO: 37, SEQ ID NO: 46, SEQ ID NO: 55, SEQ ID NO: 64, SEQ ID NO: 73, SEQ ID NO: 82, SEQ ID NO: 91, SEQ ID NO: 100, and SEQ ID NO: 103; and
 - a polypeptide having the amino acid sequence depicted in SEQ ID NO: 149; or
- (i) a polypeptide comprising in the following order starting from the N-terminus:
- a polypeptide having an amino acid sequence of SEQ ID NO: 219;
 - a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 1-9; and
 - a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 19, SEQ ID NO: 28, SEQ ID NO: 37, SEQ ID NO: 46, SEQ ID NO: 55, SEQ ID NO: 64, SEQ ID NO: 73, SEQ ID NO: 82, SEQ ID NO: 91, SEQ ID NO: 100, and SEQ ID NO: 103; and
 - a polypeptide having the amino acid sequence depicted in SEQ ID NO: 150.

29. The antibody construct of claim **23** comprising in a N- to C-terminal order:

the first binding domain having an amino acid sequence of SEQ ID NO: 219;

a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 2, 8 and 9;

the second binding domain having an amino acid sequence selected from the group consisting of SEQ ID NO: 19, SEQ ID NO: 28, SEQ ID NO: 37, SEQ ID NO: 46, SEQ ID NO: 55, SEQ ID NO: 64, SEQ ID NO: 73, SEQ ID NO: 82, SEQ ID NO: 91, SEQ ID NO: 100, and SEQ ID NO: 103 (see also SEQ ID NOs: 23, 25, 41, 43, 59, 61, 77, 79, 95, 97, 113, 115, 131, 133, 149, 151, 167, 169, 185 or 187 of WO 2008/119567);

a peptide linker having an amino acid sequence selected from the group consisting of SEQ ID NOs: 1, 2, 4, 5, 6, 8 and 9; and

the third domain having an amino acid sequence selected from the group consisting of SEQ ID NOs: 260-267.

30. The antibody construct of claim **29**, comprising a sequence selected from the group consisting of SEQ ID NOs: 292-295.

31. A polynucleotide encoding the antibody construct of claim **23**.

32. A vector comprising the polynucleotide of claim **31**.

33. A host cell transformed or transfected with the polynucleotide of claim **31**.

34. A process for producing a polypeptide, comprising culturing a host cell comprising a polynucleotide encoding the polypeptide under conditions allowing the expression of the polypeptide, wherein the polypeptide comprises a first binding domain which binds to human

and macaque MSLN on the surface of a target cell and a second binding domain which binds to human CD3 on the surface of a T cell, wherein the first binding domain binds to an epitope of MSLN which is comprised within the region of the human MSLN having a sequence having the sequence of SEQ ID NO: 241, and wherein the first binding domain binds to the same epitope of MSLN as an antibody comprising a VH region comprising CDR-H1 having the sequence of SEQ ID NO: 211, CDR-H2 having the sequence of SEQ ID NO: 212, CDR-H3 having the sequence of SEQ ID NO: 213 and a VL region comprising CDR-L1 having the sequence of SEQ ID NO: 214, CDR-L2 having the sequence of SEQ ID NO: 215 and CDR-L3 having the sequence of SEQ ID NO: 216; and

recovering the polypeptide from the culture.

35. A pharmaceutical composition comprising the antibody construct of claim **23** and a pharmaceutically acceptable carrier.

36. A method of preventing, treating, or ameliorating a solid tumor disease or a metastatic cancer disease in a subject, the method comprising administering to the subject an effective amount of the antibody construct of claim **23**.

37. The method of claim **35**, wherein the solid tumor disease or metastatic cancer disease is ovarian cancer, pancreatic cancer, mesothelioma, lung cancer, gastric cancer, or breast cancer.

38. The method of claim **36**, wherein the breast cancer is triple negative breast cancer.

39. A kit comprising the antibody construct of claim **23** and instructions for use.

* * * * *