



US 20250283098A1

(19) **United States**

(12) **Patent Application Publication**
Cheng et al.

(10) **Pub. No.: US 2025/0283098 A1**

(43) **Pub. Date: Sep. 11, 2025**

(54) **MORPHOGENIC REGULATORS AND
METHODS OF USING THE SAME**

(60) Provisional application No. 62/880,131, filed on Jul.
30, 2019.

(71) Applicant: **Pairwise Plants Services, Inc.**,
Durham, NC (US)

Publication Classification

(72) Inventors: **Ming Cheng**, Cary, NC (US);
Qingchun Shi, Cary, NC (US);
Thomas J. Poorten, Durham, NC (US);
Brian Charles Wilding Crawford,
Cary, NC (US); **Sathya Sheela Jali**,
Cary, NC (US); **Doug Heckart**, Wake
Forest, NC (US); **Yongjoo Kim**,
Durham, NC (US)

(51) **Int. Cl.**
C12N 15/82 (2006.01)
(52) **U.S. Cl.**
CPC **C12N 15/821** (2013.01); **C12N 15/8216**
(2013.01)

(21) Appl. No.: **19/218,804**

(22) Filed: **May 27, 2025**

Related U.S. Application Data

(62) Division of application No. 17/630,973, filed on Jan.
28, 2022, filed as application No. PCT/US2020/
044268 on Jul. 30, 2020, now Pat. No. 12,344,850.

(57) **ABSTRACT**

Methods and compositions for generating transgene-free
plants are described. In particular, the transgene-free plants
are regenerated by transient expression of morphogenic
regulators in a plant cell that is precisely edited using a
gene-editing methodology. Also, provided are transformed
plants, plant cell, tissues, and seeds in which a target gene is
edited, but non-transgenic.

Specification includes a Sequence Listing.

Fig. 1

1	2	3	4	5	6	7	8	9	10
11	12	13	14	15	16	17	18	19	20
21	22	23	24	25	26	27	28	29	30
31	32	33	34	35	36	37	38	39	40
41	42	43	44	45	46				
47	48	49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64	65	66
67	68	69	70	71	72	73	74	75	76
77	78	79	80	81	82	83	84	85	86
87	88	89	90	91	92	93	94	95	96
97	98	99	100	101	102	103	104	105	106

MORPHOGENIC REGULATORS AND METHODS OF USING THE SAME

STATEMENT REGARDING ELECTRONIC FILING OF A SEQUENCE LISTING

[0001] A Sequence Listing in XML format, entitled 1499-39DV_ST26.xml, 283,095 bytes in size, generated on May 25, 2025, and filed herewith, is hereby incorporated by reference in its entirety for its disclosures.

FIELD

[0002] The present disclosure relates to morphogenic regulators and methods of using the same. The present disclosure further relates to next generation plant breeding, including breeding for vegetative propagated crops, along with methods and compositions for producing transgene-free plants, optionally in which a target gene is edited.

BACKGROUND

[0003] Recombinant DNA technology has emerged as a powerful tool for editing genes and genetic elements in vectors to produce novel recombinant DNA products. Due to the continuous improvement of DNA technology and the application of *Agrobacterium* for transgene delivery, transgenic plants have been successfully generated over two decades and genetically modified (GM) plant products have been commercialized to meet high demands of modern society. Not surprisingly, genetic transformation in plants has become a routine practice so that there are more than one hundred agricultural crops to have been genetically modified around the world.

[0004] The International Service for the Acquisition of Agri-Biotech Applications (ISAAA) reported that about 160 million hectares were used for cultivating transgenic crops in 2011, which was a 94-fold increase within five years from 1996. There is no doubt that GM technology has become an important plant breeding tool as an answer to a threat of food security in developing countries.

[0005] Along with rapid growth of transgenic crop cultivation and production, GM plants have raised concerns from regulatory agencies and the public in some countries, including Europe. One of main concerns is that a piece of foreign DNA and/or a selectable marker gene (usually an antibiotic- or herbicide-resistant gene) remains in the genomes of GM plants. There have been a few approaches to producing transgene-free transgenic plants. One approach is to segregate away integration of a transgene and/or a piece of foreign DNA by conventional breeding methods, for example, by backcrossing the transgenic plants with its parent and achieving offspring with a genetic identity which is closer to that of the parent. In this way, a desired trait in the transgenic plants can be transferred to the favored genetic background of another variety, while undesired foreign DNA can be eliminated.

[0006] The aforementioned segregating-away approach has been applied to row crops for removal of unwanted foreign DNA and/or transgene(s); however, they are not particularly adopted to vegetative propagated crops due to their heterozygotic nature and/or multiploidy. The recurrent crossing method to segregate away the transgene(s) is practically and economically undesirable for heterozygotic and/or multiploid plants because the crossings would be technically challenging and/or selection of progeny with only the

desired traits and/or gene of interest (GOI) would require multiple rounds of crossing to generate pure and/or near-isogenic lines, which could result in genetic drift, such as loss of beneficial alleles. Also, conventional genetic engineering methods still have not given solutions to the technical challenges of identifying random mutations caused by integration of foreign DNAs in a host genome and removing them in a selective, efficient, and economic manner. Therefore, there is a great need in the art for a method of creating transgene-free plants with high heterozygosity and/or multiploidy, along with introduction of targeted genetic modification by employing new targeted genome engineering technologies.

SUMMARY OF THE DISCLOSURE

[0007] One aspect of the present disclosure is directed to an isolated polynucleotide comprising a nucleotide sequence having at least 90%, 95%, or 98% sequence identity to the nucleotide sequence of any one of SEQ ID NOs: 1-18 or SEQ ID NOs: 28-49. In some embodiments, the isolated polynucleotide comprises a nucleotide sequence having at least 90%, 95%, or 98% sequence identity to the nucleotide sequence of any one of SEQ ID NOs: 1-18. In some embodiments, the isolated polynucleotide comprises the nucleotide sequence of any one of SEQ ID NOs: 1-18 or SEQ ID NOs: 28-49. In some embodiments, the isolated polynucleotide comprises the nucleotide sequence of any one of SEQ ID NOs: 1-18.

[0008] Another aspect of the present disclosure is directed to an isolated polynucleotide encoding an amino acid sequence having at least 90%, 95%, or 98% sequence identity to the polypeptide sequence of any one of SEQ ID NOs: 19-27 or SEQ ID NOs: 50-60. In some embodiments, the isolated polynucleotide encodes an amino acid sequence having at least 90%, 95%, or 98% sequence identity to the polypeptide sequence of any one of SEQ ID NOs: 19-27. In some embodiments, the isolated polynucleotide encodes the polypeptide sequence of any one of SEQ ID NOs: 19-27 or SEQ ID NOs: 50-60. In some embodiments, the isolated polynucleotide encodes the polypeptide sequence of any one of SEQ ID NOs: 19-27.

[0009] A further aspect of the present disclosure is directed to an isolated polypeptide comprising an amino acid sequence having at least 90%, 95%, or 98% sequence identity to the amino acid sequence of any one of SEQ ID NOs: 19-27 or SEQ ID NOs: 50-60. In some embodiments, the isolated polypeptide comprises an amino acid sequence having at least 90%, 95%, or 98% sequence identity to the amino acid sequence of any one of SEQ ID NOs: 19-27. In some embodiments, the isolated polypeptide comprises the amino acid sequence of any one of SEQ ID NOs: 19-27 or SEQ ID NOs: 50-60. In some embodiments, the isolated polypeptide comprises the amino acid sequence of any one of SEQ ID NOs: 19-27.

[0010] Another aspect of the present disclosure is directed to a method for transforming a plant cell, the method comprising: introducing a polynucleotide or a polypeptide comprising a heterologous morphogenic regulator into the plant cell, thereby transforming the plant cell.

[0011] A further aspect of the present disclosure is directed to a method of improving or increasing regeneration frequency in a group of plant cells, the method comprising: introducing a polynucleotide comprising a heterologous morphogenic regulator into a plant cell in the group of plant

cells, thereby improving or increasing regeneration frequency in the group of plant cells.

[0012] An additional aspect of the present disclosure is directed to a plant cell, plant part, and/or plant comprising: an edited polynucleotide, wherein the plant cell, plant part, and/or plant is transgene-free.

[0013] A further aspect of the present disclosure is directed to a plant cell comprising: a polynucleotide comprising a heterologous morphogenic regulator.

[0014] Another aspect of the present disclosure is directed to a group of plant cells comprising: a first plant cell comprising a polynucleotide comprising a heterologous morphogenic regulator; and a second plant cell comprising a polynucleotide encoding gene editing machinery or a gene editing complex (e.g., a ribonucleoprotein) that is capable of modifying at least one target nucleic acid.

[0015] The present disclosure can provide a solution to the aforementioned problem of conventional breeding and/or genetic engineering methods to avoid integrating undesired transgenes and/or foreign DNA pieces from stably integrated transgenic plants. In some embodiments, the present disclosure provides a method of modulating and/or modifying a target nucleic acid using a gene-editing system. Furthermore, the disclosure teaches a method of stimulating regeneration of a gene-edited plant or plant part from a plant cell, optionally in which a target nucleic acid is stably and precisely edited. The regeneration of a gene-edited plant or plant part can be induced, as elaborated upon herein, by at least one transiently expressed morphogenic regulator taught in the present disclosure.

[0016] The present disclosure further provides expression cassettes and/or vectors comprising a polynucleotide and/or polypeptide of the present invention, and cells comprising a polynucleotide and/or polypeptide of the present invention.

[0017] It is noted that aspects of the invention described with respect to one embodiment, may be incorporated in a different embodiment although not specifically described relative thereto. That is, all embodiments and/or features of any embodiment can be combined in any way and/or combination. Applicant reserves the right to change any originally filed claim and/or file any new claim accordingly, including the right to be able to amend any originally filed claim to depend from and/or incorporate any feature of any other claim or claims although not originally claimed in that manner. These and other objects and/or aspects of the present invention are explained in detail in the specification set forth below. Further features, advantages and details of the present invention will be appreciated by those of ordinary skill in the art from a reading of the figures and the detailed description of the preferred embodiments that follow, such description being merely illustrative of the present invention.

BRIEF DESCRIPTION OF THE FIGURES

[0018] FIG. 1 illustrates GUS assay results of one hundred and six shoots induced by the ipt gene expression. From one hundred six ipt-induced shoots, seven samples were GUS-positive, but the other ninety-nine samples were GUS-negative.

DETAILED DESCRIPTION

[0019] The present disclosure now will be described hereinafter with reference to the accompanying drawings and

examples, in which embodiments of the invention are shown. This description is not intended to be a detailed catalog of all the different ways in which the invention may be implemented, or all the features that may be added to the instant invention. For example, features illustrated with respect to one embodiment may be incorporated into other embodiments, and features illustrated with respect to a particular embodiment may be deleted from that embodiment. Thus, the invention contemplates that in some embodiments of the invention, any feature or combination of features set forth herein can be excluded or omitted. To illustrate, if the specification states that a composition comprises components A, B and C, it is specifically intended that any of A, B or C, or a combination thereof, can be omitted and disclaimed singularly or in any combination. In addition, numerous variations and additions to the various embodiments suggested herein will be apparent to those skilled in the art in light of the instant disclosure, which do not depart from the instant invention. Hence, the following descriptions are intended to illustrate some particular embodiments of the invention, and not to exhaustively specify all permutations, combinations and variations thereof.

[0020] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. The terminology used in the description of the invention herein is for the purpose of describing particular embodiments only and is not intended to be limiting of the invention.

[0021] All publications, patent applications, patents and other references cited herein are incorporated by reference in their entireties for the teachings relevant to the sentence and/or paragraph in which the reference is presented.

[0022] As used in the description of the invention and the appended claims, the singular forms “a,” “an” and “the” are intended to include the plural forms as well, unless the context clearly indicates otherwise.

[0023] Also as used herein, “and/or” refers to and encompasses any and all possible combinations of one or more of the associated listed items, as well as the lack of combinations when interpreted in the alternative (“or”).

[0024] The term “about,” as used herein when referring to a measurable value such as an amount or concentration and the like, is meant to encompass variations of $\pm 10\%$, $\pm 5\%$, $\pm 1\%$, $\pm 0.5\%$, or even $\pm 0.1\%$ of the specified value as well as the specified value. For example, “about X” where X is the measurable value, is meant to include X as well as variations of $\pm 10\%$, $\pm 5\%$, $\pm 1\%$, $\pm 0.5\%$, or even $\pm 0.1\%$ of X. A range provided herein for a measureable value may include any other range and/or individual value therein.

[0025] As used herein, phrases such as “between X and Y” and “between about X and Y” should be interpreted to include X and Y. As used herein, phrases such as “between about X and Y” mean “between about X and about Y” and phrases such as “from about X to Y” mean “from about X to about Y.”

[0026] Recitation of ranges of values herein are merely intended to serve as a shorthand method of referring individually to each separate value falling within the range, unless otherwise indicated herein, and each separate value is incorporated into the specification as if it were individually recited herein. For example, if the range 10 to 15 is disclosed, then 11, 12, 13, and 14 are also disclosed.

[0027] The term “comprise,” “comprises” and “comprising” as used herein, specify the presence of the stated features, integers, steps, operations, elements, and/or components, but do not preclude the presence or addition of one or more other features, integers, steps, operations, elements, components, and/or groups thereof.

[0028] As used herein, the transitional phrase “consisting essentially of” means that the scope of a claim is to be interpreted to encompass the specified materials or steps recited in the claim and those that do not materially affect the basic and novel characteristic(s) of the claimed invention. Thus, the term “consisting essentially of” when used in a claim of this invention is not intended to be interpreted to be equivalent to “comprising.”

[0029] As used herein, the terms “increase,” “increasing,” “enhance,” “enhancing,” “improve” and “improving” (and grammatical variations thereof) describe an elevation in a specified parameter or value of at least about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 100%, 150%, 200%, 300%, 400%, 500% or more as compared to a control.

[0030] As used herein, the terms “reduce,” “reduced,” “reducing,” “reduction,” “diminish,” and “decrease” (and grammatical variations thereof), describe, for example, a decrease in a specified parameter or value of at least about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99%, or 100% as compared to a control. In particular embodiments, the reduction can result in no or essentially no (i.e., an insignificant amount, e.g., less than about 10% or even 5%) detectable activity or amount.

[0031] A “heterologous” or a “recombinant” nucleotide sequence is a nucleotide sequence not naturally associated with a host cell into which it is introduced, including non-naturally occurring multiple copies of a naturally occurring nucleotide sequence.

[0032] A “native” or “wild type” nucleic acid, nucleotide sequence, polypeptide or amino acid sequence refer to a naturally occurring or endogenous nucleic acid, nucleotide sequence, polypeptide or amino acid sequence. Thus, for example, a “wild type mRNA” is an mRNA that is naturally occurring in or endogenous to the reference organism. A “homologous” nucleic acid sequence is a nucleotide sequence naturally associated with a host cell into which it is introduced.

[0033] As used herein, the terms “nucleic acid,” “nucleic acid molecule,” “nucleotide sequence” and “polynucleotide” refer to RNA or DNA that is linear or branched, single or double stranded, or a hybrid thereof. The term also encompasses RNA/DNA hybrids. When dsRNA is produced synthetically, less common bases, such as inosine, 5-methylcytosine, 6-methyladenine, hypoxanthine and others can also be used for antisense, dsRNA, and ribozyme pairing. For example, polynucleotides that contain C-5 propyne analogues of uridine and cytidine have been shown to bind RNA with high affinity and to be potent antisense inhibitors of gene expression. Other modifications, such as modification to the phosphodiester backbone, or the 2'-hydroxy in the ribose sugar group of the RNA can also be made.

[0034] As used herein, the term “nucleotide sequence” refers to a heteropolymer of nucleotides or the sequence of these nucleotides from the 5' to 3' end of a nucleic acid molecule and includes DNA or RNA molecules, including cDNA, a DNA fragment or portion, genomic DNA, syn-

thetic (e.g., chemically synthesized) DNA, plasmid DNA, mRNA, and anti-sense RNA, any of which can be single stranded or double stranded. The terms “nucleotide sequence” “nucleic acid,” “nucleic acid molecule,” “nucleic acid construct,” “recombinant nucleic acid,” “oligonucleotide” and “polynucleotide” are also used interchangeably herein to refer to a heteropolymer of nucleotides. Nucleic acid molecules and/or nucleotide sequences provided herein are presented herein in the 5' to 3' direction, from left to right and are represented using the standard code for representing the nucleotide characters as set forth in the U.S. sequence rules, 37 CFR §§ 1.821-1.825 and the World Intellectual Property Organization (WIPO) Standard ST.25. A “5' region” as used herein can mean the region of a polynucleotide that is nearest the 5' end of the polynucleotide. Thus, for example, an element in the 5' region of a polynucleotide can be located anywhere from the first nucleotide located at the 5' end of the polynucleotide to the nucleotide located halfway through the polynucleotide. A “3' region” as used herein can mean the region of a polynucleotide that is nearest the 3' end of the polynucleotide. Thus, for example, an element in the 3' region of a polynucleotide can be located anywhere from the first nucleotide located at the 3' end of the polynucleotide to the nucleotide located halfway through the polynucleotide.

[0035] As used herein, the term “gene” refers to a nucleic acid molecule capable of being used to produce mRNA, antisense RNA, miRNA, anti-microRNA antisense oligodeoxyribonucleotide (AMO) and the like. Genes may or may not be capable of being used to produce a functional protein or gene product. Genes can include both coding and non-coding regions (e.g., introns, regulatory elements, promoters, enhancers, termination sequences and/or 5' and 3' untranslated regions).

[0036] A polynucleotide, gene, or polypeptide may be “isolated” by which is meant a nucleic acid or polypeptide that is substantially or essentially free from components normally found in association with the nucleic acid or polypeptide, respectively, in its natural state. In some embodiments, such components include other cellular material, culture medium from recombinant production, and/or various chemicals used in chemically synthesizing the nucleic acid or polypeptide.

[0037] The term “mutation” refers to point mutations (e.g., missense, or nonsense, or insertions or deletions of single base pairs that result in frame shifts), insertions, deletions, and/or truncations. When the mutation is a substitution of a residue within an amino acid sequence with another residue, or a deletion or insertion of one or more residues within a sequence, the mutations are typically described by identifying the original residue followed by the position of the residue within the sequence and by the identity of the newly substituted residue.

[0038] The terms “complementary” or “complementarity,” as used herein, refer to the natural binding of polynucleotides under permissive salt and temperature conditions by base-pairing. For example, the sequence “A-G-T” (5' to 3') binds to the complementary sequence “T-C-A” (3' to 5'). Complementarity between two single-stranded molecules may be “partial,” in which only some of the nucleotides bind, or it may be complete when total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has

significant effects on the efficiency and strength of hybridization between nucleic acid strands.

[0039] “Complement” as used herein can mean 100% complementarity with the comparator nucleotide sequence or it can mean less than 100% complementarity (e.g., “substantially complementary,” e.g., about 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, and the like, complementarity).

[0040] A “portion” or “fragment” of a nucleotide sequence or polypeptide sequence will be understood to mean a nucleotide or polypeptide sequence of reduced length (e.g., reduced by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more residue(s) (e.g., nucleotide(s) or peptide(s)) relative to a reference nucleotide or polypeptide sequence, respectively, and comprising, consisting essentially of and/or consisting of a nucleotide or polypeptide sequence of contiguous residues, respectively, identical or almost identical (e.g., 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% identical) to the reference nucleotide or polypeptide sequence. Such a nucleic acid fragment or portion according to the invention may be, where appropriate, included in a larger polynucleotide of which it is a constituent. As an example, a repeat sequence of guide nucleic acid of this invention may comprise a portion of a wild type CRISPR-Cas repeat sequence (e.g., a wild type Type V CRISPR Cas repeat, e.g., a repeat from the CRISPR Cas system that includes, but is not limited to, Cas12a (Cpf1), Cas12b, Cas12c (C2c3), Cas12d (CasY), Cas12e (CasX), Cas12g, Cas12h, Cas12i, C2c1, C2c4, C2c5, C2c8, C2c9, C2c10, Cas14a, Cas14b, and/or Cas14c, and the like).

[0041] Different nucleic acids or proteins having homology are referred to herein as “homologues.” The term homologue includes homologous sequences from the same and other species and orthologous sequences from the same and other species. “Homology” refers to the level of similarity between two or more nucleic acid and/or amino acid sequences in terms of percent of positional identity (i.e., sequence similarity or identity). Homology also refers to the concept of similar functional properties among different nucleic acids or proteins. Thus, the compositions and methods of the invention further comprise homologues to the nucleotide sequences and polypeptide sequences of this invention. “Orthologous” and “orthologs” as used herein, refers to homologous nucleotide sequences and/or amino acid sequences in different species that arose from a common ancestral gene during speciation. A homologue or ortholog of a nucleotide sequence of this invention has a substantial sequence identity (e.g., at least about 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or 100%) to said nucleotide sequence of the invention.

[0042] As used herein “sequence identity” refers to the extent to which two optimally aligned polynucleotide or polypeptide sequences are invariant throughout a window of alignment of components, e.g., nucleotides or amino acids. “Identity” can be readily calculated by known methods including, but not limited to, those described in: *Computational Molecular Biology* (Lesk, A. M., ed.) Oxford University Press, New York (1988); *Biocomputing: Informatics*

and Genome Projects (Smith, D. W., ed.) Academic Press, New York (1993); *Computer Analysis of Sequence Data, Part I* (Griffin, A. M., and Griffin, H. G., eds.) Humana Press, New Jersey (1994); *Sequence Analysis in Molecular Biology* (von Heinje, G., ed.) Academic Press (1987); and *Sequence Analysis Primer* (Gribskov, M. and Devereux, J., eds.) Stockton Press, New York (1991).

[0043] As used herein, the term “percent sequence identity” or “percent identity” refers to the percentage of identical nucleotides in a linear polynucleotide sequence of a reference (“query”) polynucleotide molecule (or its complementary strand) as compared to a test (“subject”) polynucleotide molecule (or its complementary strand) when the two sequences are optimally aligned. In some embodiments, “percent identity” can refer to the percentage of identical amino acids in an amino acid sequence as compared to a reference polypeptide.

[0044] As used herein, the phrase “substantially identical,” or “substantial identity” in the context of two nucleic acid molecules, nucleotide sequences or protein sequences, refers to two or more sequences or subsequences that have at least about 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or 100% nucleotide or amino acid residue identity, when compared and aligned for maximum correspondence, as measured using one of the following sequence comparison algorithms or by visual inspection. In some embodiments of the invention, the substantial identity exists over a region of consecutive nucleotides of a nucleotide sequence of the invention that is about 10 nucleotides to about 20 nucleotides, about 10 nucleotides to about 25 nucleotides, about 10 nucleotides to about 30 nucleotides, about 15 nucleotides to about 25 nucleotides, about 30 nucleotides to about 40 nucleotides, about 50 nucleotides to about 60 nucleotides, about 70 nucleotides to about 80 nucleotides, about 90 nucleotides to about 100 nucleotides, or more nucleotides in length, and any range therein, up to the full length of the sequence. In some embodiments, the nucleotide sequences can be substantially identical over at least about 20 nucleotides (e.g., about 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40 nucleotides). In some embodiments, a substantially identical nucleotide or protein sequence performs substantially the same function as the nucleotide (or encoded protein sequence) to which it is substantially identical.

[0045] For sequence comparison, typically one sequence acts as a reference sequence to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are entered into a computer, subsequence coordinates are designated if necessary, and sequence algorithm program parameters are designated. The sequence comparison algorithm then calculates the percent sequence identity for the test sequence(s) relative to the reference sequence, based on the designated program parameters.

[0046] Optimal alignment of sequences for aligning a comparison window are well known to those skilled in the art and may be conducted by tools such as the local homology algorithm of Smith and Waterman, the homology alignment algorithm of Needleman and Wunsch, the search for similarity method of Pearson and Lipman, and optionally by computerized implementations of these algorithms such as GAP, BESTFIT, FASTA, and TFASTA available as part

of the GCG WISCONSIN PACKAGE® (Accelrys Inc., San Diego, CA) software suite for sequence analysis. An “identity fraction” for aligned segments of a test sequence and a reference sequence is the number of identical components which are shared by the two aligned sequences divided by the total number of components in the reference sequence segment, e.g., the entire reference sequence or a smaller defined part of the reference sequence. Percent sequence identity is represented as the identity fraction multiplied by 100. The comparison of one or more polynucleotide sequences may be to a full-length polynucleotide sequence or a portion thereof, or to a longer polynucleotide sequence. For purposes of this invention “percent identity” may also be determined using BLASTX version 2.0 for translated nucleotide sequences and BLASTN version 2.0 for polynucleotide sequences.

[0047] Two nucleotide sequences may also be considered substantially complementary when the two sequences hybridize to each other under stringent conditions. In some representative embodiments, two nucleotide sequences considered to be substantially complementary hybridize to each other under highly stringent conditions.

[0048] “Stringent hybridization conditions” and “stringent hybridization wash conditions” in the context of nucleic acid hybridization experiments such as Southern and Northern hybridizations are sequence dependent, and are different under different environmental parameters. An extensive guide to the hybridization of nucleic acids is found in Tijssen *Laboratory Techniques in Biochemistry and Molecular Biology-Hybridization with Nucleic Acid Probes* part I chapter 2 “Overview of principles of hybridization and the strategy of nucleic acid probe assays” Elsevier, New York (1993). Generally, highly stringent hybridization and wash conditions are selected to be about 5° C. lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH.

[0049] The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Very stringent conditions are selected to be equal to the T_m for a particular probe. An example of stringent hybridization conditions for hybridization of complementary nucleotide sequences which have more than 100 complementary residues on a filter in a Southern or northern blot is 50% formamide with 1 mg of heparin at 42° C., with the hybridization being carried out overnight. An example of highly stringent wash conditions is 0.1 M NaCl at 72° C. for about 15 minutes. An example of stringent wash conditions is a 0.2×SSC wash at 65° C. for 15 minutes (see, Sambrook, *infra*, for a description of SSC buffer). Often, a high stringency wash is preceded by a low stringency wash to remove background probe signal. An example of a medium stringency wash for a duplex of, e.g., more than 100 nucleotides, is 1×SSC at 45° C. for 15 minutes. An example of a low stringency wash for a duplex of, e.g., more than 100 nucleotides, is 4-6×SSC at 40° C. for 15 minutes. For short probes (e.g., about 10 to 50 nucleotides), stringent conditions typically involve salt concentrations of less than about 1.0 M Na ion, typically about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3, and the temperature is typically at least about 30° C. Stringent conditions can also be achieved with the addition of destabilizing agents such as formamide. In general, a signal to noise ratio of 2× (or higher) than that observed for an unrelated probe in the particular hybridization assay

indicates detection of a specific hybridization. Nucleotide sequences that do not hybridize to each other under stringent conditions are still substantially identical if the proteins that they encode are substantially identical. This can occur, for example, when a copy of a nucleotide sequence is created using the maximum codon degeneracy permitted by the genetic code.

[0050] A polynucleotide and/or recombinant nucleic acid construct of this invention can be codon optimized for expression. In some embodiments, a polynucleotide, nucleic acid construct, expression cassette, and/or vector of the present invention (e.g., that comprises/encodes a nucleic acid binding domain (e.g., a DNA binding domain such as a sequence-specific DNA binding domain from a polynucleotide-guided endonuclease, a zinc finger nuclease, a transcription activator-like effector nuclease (TALEN), an Argonaute protein, and/or a CRISPR-Cas effector protein), a guide nucleic acid, a cytosine deaminase, adenine deaminase, and/or a heterologous morphogenic regulator) may be codon optimized for expression in an organism (e.g., an animal, a plant, a fungus, an archaon, or a bacterium). In some embodiments, the codon optimized nucleic acid constructs, polynucleotides, expression cassettes, and/or vectors of the invention have about 70% to about 99.9% (e.g., 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5%, 99.9% or 100%) identity or more to the reference nucleic acid constructs, polynucleotides, expression cassettes, and/or vectors but which have not been codon optimized.

[0051] In any of the embodiments described herein, a polynucleotide or nucleic acid construct of the invention may be operatively associated with a variety of promoters and/or other regulatory elements for expression in an organism or cell thereof (e.g., a plant and/or a cell of a plant). Thus, in some embodiments, a polynucleotide or nucleic acid construct of this invention may further comprise one or more promoters, introns, enhancers, and/or terminators operably linked to one or more nucleotide sequences. In some embodiments, a promoter may be operably associated with an intron (e.g., Ubi1 promoter and intron). In some embodiments, a promoter associated with an intron may be referred to as a “promoter region” (e.g., Ubi1 promoter and intron).

[0052] By “operably linked” or “operably associated” as used herein in reference to polynucleotides, it is meant that the indicated elements are functionally related to each other, and are also generally physically related. Thus, the term “operably linked” or “operably associated” as used herein, refers to nucleotide sequences on a single nucleic acid molecule that are functionally associated. Thus, a first nucleotide sequence that is operably linked to a second nucleotide sequence means a situation when the first nucleotide sequence is placed in a functional relationship with the second nucleotide sequence. For instance, a promoter is operably associated with a nucleotide sequence if the promoter effects the transcription or expression of said nucleotide sequence. Those skilled in the art will appreciate that the control sequences (e.g., promoter) need not be contiguous with the nucleotide sequence to which it is operably associated, as long as the control sequences function to direct the expression thereof. Thus, for example, intervening untranslated, yet transcribed, nucleic acid sequences can be present

between a promoter and the nucleotide sequence, and the promoter can still be considered “operably linked” to the nucleotide sequence.

[0053] As used herein, the term “linked,” or “fused” in reference to polypeptides, refers to the attachment of one polypeptide to another. A polypeptide may be linked or fused to another polypeptide (at the N-terminus or the C-terminus) directly (e.g., via a peptide bond) or through a linker (e.g., a peptide linker).

[0054] The term “linker” in reference to polypeptides is art-recognized and refers to a chemical group, or a molecule linking two molecules or moieties, e.g., two domains of a fusion protein, such as, for example, a CRISPR-Cas effector protein and a peptide tag and/or a polypeptide of interest. A linker may be comprised of a single linking molecule (e.g., a single amino acid) or may comprise more than one linking molecule. In some embodiments, the linker can be an organic molecule, group, polymer, or chemical moiety such as a bivalent organic moiety. In some embodiments, the linker may be an amino acid or it may be a peptide. In some embodiments, the linker is a peptide. In some embodiments, a peptide linker useful with this invention may be about 2 to about 100 or more amino acids in length. In some embodiments, a peptide linker may be a GS linker.

[0055] In some embodiments, two or more polynucleotide molecules may be linked by a linker that can be an organic molecule, group, polymer, or chemical moiety such as a bivalent organic moiety. A polynucleotide may be linked or fused to another polynucleotide (at the 5' end or the 3' end) via a covalent or non-covalent linkage or binding, including e.g., Watson-Crick base-pairing, or through one or more linking nucleotides. In some embodiments, a polynucleotide motif of a certain structure may be inserted within another polynucleotide sequence (e.g. extension of the hairpin structure in guide RNA). In some embodiments, the linking nucleotides may be naturally occurring nucleotides. In some embodiments, the linking nucleotides may be non-naturally occurring nucleotides.

[0056] A “promoter” is a nucleotide sequence that controls or regulates the transcription of a nucleotide sequence (e.g., a coding sequence) that is operably associated with the promoter. The coding sequence controlled or regulated by a promoter may encode a polypeptide and/or a functional RNA. Typically, a “promoter” refers to a nucleotide sequence that contains a binding site for RNA polymerase II and directs the initiation of transcription. In general, promoters are found 5', or upstream, relative to the start of the coding region of the corresponding coding sequence. A promoter may comprise other elements that act as regulators of gene expression; e.g., a promoter region. These include a TATA box consensus sequence, and often a CAAT box consensus sequence (Breathnach and Chambon, (1981) *Annu. Rev. Biochem.* 50:349). In plants, the CAAT box may be substituted by the AGGA box (Messing et al., (1983) in *Genetic Engineering of Plants*, T. Kosuge, C. Meredith and A. Hollaender (eds.), Plenum Press, pp. 211-227). In some embodiments, a promoter region may comprise at least one intron (e.g., SEQ ID NO:61 or SEQ ID NO:62).

[0057] Promoters useful with this invention can include, for example, constitutive, inducible, temporally regulated, developmentally regulated, chemically regulated, tissue-preferred and/or tissue-specific promoters for use in the preparation of recombinant nucleic acid molecules, e.g.,

“synthetic nucleic acid constructs” or “protein-RNA complex.” These various types of promoters are known in the art.

[0058] The choice of promoter may vary depending on the temporal and spatial requirements for expression, and also may vary based on the host cell to be transformed. Promoters for many different organisms are well known in the art. Based on the extensive knowledge present in the art, the appropriate promoter can be selected for the particular host organism of interest. Thus, for example, much is known about promoters upstream of highly constitutively expressed genes in model organisms and such knowledge can be readily accessed and implemented in other systems as appropriate.

[0059] In some embodiments, a promoter functional in a plant may be used with the constructs of this invention. Non-limiting examples of a promoter useful for driving expression in a plant include the promoter of the RubisCo small subunit gene 1 (PrbcS1), the promoter of the actin gene (Pactin), the promoter of the nitrate reductase gene (Pnr) and the promoter of duplicated carbonic anhydrase gene 1 (Pdca1) (See, Walker et al. *Plant Cell Rep.* 23:727-735 (2005); Li et al. *Gene* 403:132-142 (2007); Li et al. *Mol Biol. Rep.* 37:1143-1154 (2010)). PrbcS1 and Pactin are constitutive promoters and Pnr and Pdca1 are inducible promoters. Pnr is induced by nitrate and repressed by ammonium (Li et al. *Gene* 403:132-142 (2007)) and Pdca1 is induced by salt (Li et al. *Mol Biol. Rep.* 37:1143-1154 (2010)).

[0060] Examples of constitutive promoters useful for plants include, but are not limited to, cestrum virus promoter (cmp) (U.S. Pat. No. 7,166,770), the rice actin 1 promoter (Wang et al. (1992) *Mol. Cell. Biol.* 12:3399-3406; as well as U.S. Pat. No. 5,641,876), CaMV 35S promoter (Odell et al. (1985) *Nature* 313:810-812), CaMV 19S promoter (Lawton et al. (1987) *Plant Mol. Biol.* 9:315-324), nos promoter (Ebert et al. (1987) *Proc. Natl. Acad. Sci USA* 84:5745-5749), Adh promoter (Walker et al. (1987) *Proc. Natl. Acad. Sci. USA* 84:6624-6629), sucrose synthase promoter (Yang & Russell (1990) *Proc. Natl. Acad. Sci. USA* 87:4144-4148), and the ubiquitin promoter. The constitutive promoter derived from ubiquitin accumulates in many cell types. Ubiquitin promoters have been cloned from several plant species for use in transgenic plants, for example, sunflower (Binet et al., 1991. *Plant Science* 79:87-94), maize (Christensen et al., 1989. *Plant Molec. Biol.* 12:619-632), and *arabidopsis* (Norris et al. 1993. *Plant Molec. Biol.* 21:895-906). The maize ubiquitin promoter (UbiP) has been developed in transgenic monocot systems and its sequence and vectors constructed for monocot transformation are disclosed in the European patent publication EP0342926. The ubiquitin promoter is suitable for the expression of the nucleotide sequences of the invention in transgenic plants, especially monocotyledons. Further, the promoter expression cassettes described by McElroy et al. (*Mol. Gen. Genet.* 231:150-160 (1991)) can be easily modified for the expression of the nucleotide sequences of the invention and are particularly suitable for use in monocotyledonous hosts.

[0061] In some embodiments, tissue specific/tissue preferred promoters can be used for expression of a heterologous polynucleotide in a plant cell. Tissue specific or preferred expression patterns include, but are not limited to, green tissue specific or preferred, root specific or preferred, stem specific or preferred, flower specific or preferred or pollen specific or preferred. Promoters suitable for expres-

sion in green tissue include many that regulate genes involved in photosynthesis and many of these have been cloned from both monocotyledons and dicotyledons. In one embodiment, a promoter useful with the invention is the maize PEPC promoter from the phosphoenol carboxylase gene (Hudspeth & Grula, *Plant Molec. Biol.* 12:579-589 (1989)). Non-limiting examples of tissue-specific promoters include those associated with genes encoding the seed storage proteins (such as β -conglycinin, cruciferin, napin and phaseolin), zein or oil body proteins (such as oleosin), or proteins involved in fatty acid biosynthesis (including acyl carrier protein, stearoyl-ACP desaturase and fatty acid desaturases (fad 2-1)), and other nucleic acids expressed during embryo development (such as Bcc4, see, e.g., Kridl et al. (1991) *Seed Sci. Res.* 1:209-219; as well as EP U.S. Pat. No. 255,378). Tissue-specific or tissue-preferential promoters useful for the expression of the nucleotide sequences of the invention in plants, particularly maize, include but are not limited to those that direct expression in root, pith, leaf or pollen. Such promoters are disclosed, for example, in WO 93/07278, incorporated by reference herein for its disclosure of promoters. Other non-limiting examples of tissue specific or tissue preferred promoters useful with the invention the cotton rubisco promoter disclosed in U.S. Pat. No. 6,040,504; the rice sucrose synthase promoter disclosed in U.S. Pat. No. 5,604,121; the root specific promoter described by de Framond (FEBS 290:103-106 (1991); European patent EP 0452269 to Ciba-Geigy); the stem specific promoter described in U.S. Pat. No. 5,625,136 (to Ciba-Geigy) and which drives expression of the maize trpA gene; the cestrum yellow leaf curling virus promoter disclosed in WO 01/73087; and pollen specific or preferred promoters including, but not limited to, ProOsLPS10 and ProOsLPS11 from rice (Nguyen et al. *Plant Biotechnol. Reports* 9(5):297-306 (2015)), ZmSTK2_USP from maize (Wang et al. *Genome* 60(6):485-495 (2017)), LAT52 and LAT59 from tomato (Twell et al. *Development* 109(3):705-713 (1990)), Zm13 (U.S. Pat. No. 10,421,972), PL_{A2}- δ promoter from *arabidopsis* (U.S. Pat. No. 7,141,424), and/or the ZmC5 promoter from maize (International PCT Publication No. WO1999/042587).

[0062] Additional examples of plant tissue-specific/tissue preferred promoters include, but are not limited to, the root hair-specific cis-elements (RHEs) (Kim et al. *The Plant Cell* 18:2958-2970 (2006)), the root-specific promoters RCc3 (Jeong et al. *Plant Physiol.* 153:185-197 (2010)) and RB7 (U.S. Pat. No. 5,459,252), the lectin promoter (Lindstrom et al. (1990) *Dev. Genet.* 11:160-167; and Vodkin (1983) *Prog. Clin. Biol. Res.* 138:87-98), corn alcohol dehydrogenase 1 promoter (Dennis et al. (1984) *Nucleic Acids Res.* 12:3983-4000), S-adenosyl-L-methionine synthetase (SAMS) (Vander Mijnsbrugge et al. (1996) *Plant and Cell Physiology*, 37(8):1108-1115), corn light harvesting complex promoter (Bansal et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:3654-3658), corn heat shock protein promoter (O'Dell et al. (1985) *EMBO J.* 5:451-458; and Rochester et al. (1986) *EMBO J.* 5:451-458), pea small subunit RuBP carboxylase promoter (Cashmore, "Nuclear genes encoding the small subunit of ribulose-1,5-bisphosphate carboxylase" pp. 29-39 In: Genetic Engineering of Plants (Hollaender ed., Plenum Press 1983; and Poulsen et al. (1986) *Mol. Gen. Genet.* 205:193-200), Ti plasmid mannopine synthase promoter (Langridge et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:3219-3223), Ti plasmid nopaline synthase promoter (Langridge et

al. (1989), supra), *petunia* chalcone isomerase promoter (van Tunen et al. (1988) *EMBO J.* 7:1257-1263), bean glycine rich protein 1 promoter (Keller et al. (1989) *Genes Dev.* 3:1639-1646), truncated CaMV 35S promoter (O'Dell et al. (1985) *Nature* 313:810-812), potato patatin promoter (Wenzler et al. (1989) *Plant Mol. Biol.* 13:347-354), root cell promoter (Yamamoto et al. (1990) *Nucleic Acids Res.* 18:7449), maize zein promoter (Kriz et al. (1987) *Mol. Gen. Genet.* 207:90-98; Langridge et al. (1983) *Cell* 34:1015-1022; Reina et al. (1990) *Nucleic Acids Res.* 18:6425; Reina et al. (1990) *Nucleic Acids Res.* 18:7449; and Wandelt et al. (1989) *Nucleic Acids Res.* 17:2354), globulin-1 promoter (Belanger et al. (1991) *Genetics* 129:863-872), α -tubulin cab promoter (Sullivan et al. (1989) *Mol. Gen. Genet.* 215:431-440), PEPCase promoter (Hudspeth & Grula (1989) *Plant Mol. Biol.* 12:579-589), R gene complex-associated promoters (Chandler et al. (1989) *Plant Cell* 1:1175-1183), and chalcone synthase promoters (Franken et al. (1991) *EMBO J.* 10:2605-2612).

[0063] Useful for seed-specific expression is the pea vicilin promoter (Czako et al. (1992) *Mol. Gen. Genet.* 235:33-40; as well as the seed-specific promoters disclosed in U.S. Pat. No. 5,625,136. Useful promoters for expression in mature leaves are those that are switched at the onset of senescence, such as the SAG promoter from *Arabidopsis* (Gan et al. (1995) *Science* 270:1986-1988).

[0064] In addition, promoters functional in chloroplasts can be used. Non-limiting examples of such promoters include the bacteriophage T3 gene 9 5' UTR and other promoters disclosed in U.S. Pat. No. 7,579,516. Other promoters useful with the invention include but are not limited to the S-E9 small subunit RuBP carboxylase promoter and the Kunitz trypsin inhibitor gene promoter (Kti3).

[0065] Additional regulatory elements useful with this invention include, but are not limited to, introns, enhancers, termination sequences and/or 5' and 3' untranslated regions.

[0066] An intron useful with this invention can be an intron identified in and isolated from a plant and then inserted into an expression cassette to be used in transformation of a plant. As would be understood by those of skill in the art, introns can comprise the sequences required for self-excision and are incorporated into nucleic acid constructs/expression cassettes in frame. An intron can be used either as a spacer to separate multiple protein-coding sequences in one nucleic acid construct, or an intron can be used inside one protein-coding sequence to, for example, stabilize the mRNA. If they are used within a protein-coding sequence, they are inserted "in-frame" with the excision sites included. Introns may also be associated with promoters to improve or modify expression. As an example, a promoter/intron combination useful with this invention includes but is not limited to that of the maize Ubi1 promoter and intron.

[0067] Non-limiting examples of introns useful with the present invention include introns from the ADHI gene (e.g., Adh1-S introns 1, 2 and 6), the ubiquitin gene (Ubi1), the RuBisCO small subunit (rbcS) gene, the RuBisCO large subunit (rbcL) gene, the actin gene (e.g., actin-1 intron), the pyruvate dehydrogenase kinase gene (pdh), the nitrate reductase gene (nr), the duplicated carbonic anhydrase gene 1 (Tdca1), the psbA gene, the atpA gene, or any combination thereof.

[0068] "Gene editing machinery" and a "gene editing complex" as used herein can each be referred to as an editing

system. “Gene editing machinery” and a “gene editing complex” useful with this invention can be any site-specific (sequence-specific) genome editing system now known or later developed, which system can introduce mutations in a target specific manner. For example, gene editing machinery or a gene editing complex can include, but are not limited to, a CRISPR-Cas editing system, a meganuclease editing system, a zinc finger nuclease (ZFN) editing system, a transcription activator-like effector nuclease (TALEN) editing system, a base editing system and/or a prime editing system, each of which can comprise one or more polypeptides and/or one or more polynucleotides that when present and/or expressed as a system in a cell can modify (mutate) a target nucleic acid in a sequence specific manner. In some embodiments, an editing system (e.g., site- or sequence-specific editing system) can comprise one or more polynucleotides and/or one or more polypeptides, including but not limited to a nucleic acid binding domain (DNA binding domain), a nuclease, and/or other polypeptide, and/or a polynucleotide.

[0069] In some embodiments, an editing system can comprise one or more sequence-specific nucleic acid binding domains (e.g., DNA binding domains) that can be from, for example, a polynucleotide-guided endonuclease, a CRISPR-Cas endonuclease (e.g., CRISPR-Cas effector protein), a zinc finger nuclease, a transcription activator-like effector nuclease (TALEN) and/or an Argonaute protein. In some embodiments, an editing system can comprise one or more cleavage domains (e.g., nucleases) including, but not limited to, an endonuclease (e.g., FokI), a polynucleotide-guided endonuclease, a CRISPR-Cas endonuclease (e.g., CRISPR-Cas effector protein), a zinc finger nuclease, and/or a transcription activator-like effector nuclease (TALEN).

[0070] A “nucleic acid binding domain” as used herein includes a DNA binding domain and may be a site- or sequence-specific nucleic acid binding domain. In some embodiments, a nucleic acid binding domain may be a sequence-specific nucleic acid binding domain such as, but not limited to, a sequence-specific binding domain from, for example, a polynucleotide-guided endonuclease, a CRISPR-Cas effector protein (e.g., CRISPR-Cas endonuclease), a zinc finger nuclease, a transcription activator-like effector nuclease (TALEN) and/or an Argonaute protein. In some embodiments, a nucleic acid binding domain comprises a cleavage domain (e.g., a nuclease domain) such as, but not limited to, an endonuclease (e.g., FokI), a polynucleotide-guided endonuclease, a CRISPR-Cas endonuclease, a zinc finger nuclease, and/or a transcription activator-like effector nuclease (TALEN). In some embodiments, the nucleic acid binding domain is a polypeptide that can associate (e.g., form a complex) with one or more nucleic acid molecules (e.g., form a complex with a guide nucleic acid as described herein) that can direct or guide the nucleic acid binding domain to a specific target nucleotide sequence (e.g., a gene locus of a genome) that is complementary to the one or more nucleic acid molecules (or a portion or region thereof), thereby causing the nucleic acid binding domain to bind to the nucleotide sequence at the specific target site. In some embodiments, the nucleic acid binding domain is a CRISPR-Cas effector protein as described herein. In some embodiments, reference is made to specifically to a CRISPR-Cas effector protein for simplicity, but a nucleic acid binding domain as described herein may be used.

[0071] In some embodiments, a gene editing complex comprises a ribonucleoprotein such as an assembled ribo-

nucleoprotein complex (e.g., that comprises a CRISPR-Cas effector protein and a guide nucleic acid). A gene editing complex, as used herein, may be assembled when introduced into a plant cell or may assemble into a complex (e.g., a covalently and/or non-covalently bound complex) after and/or during introduction into a plant cell. Exemplary ribonucleoproteins and methods of use thereof include, but are not limited to, those described in Malnoid et al., (2016) *Front. Plant Sci.* 7:1904; Subburaj et al., (2016) *Plant Cell Rep.* 35:1535; Woo et al., (2015) *Nat. Biotechnol.* 33:1162; Liang et al., (2017) *Nat. Commun.* 8:14261; Svitashhev et al., *Nat. Commun.* 7, 13274 (2016); Zhang et al., (2016) *Nat. Commun.* 7:12617; Kim et al., (2017) *Nat. Commun.* 8:14406.

[0072] In some embodiments, a polynucleotide and/or a nucleic acid construct of the invention can be an “expression cassette” or can be comprised within an expression cassette. As used herein, “expression cassette” means a recombinant nucleic acid molecule comprising, for example, a nucleic acid construct of the invention (e.g., a polynucleotide comprising a heterologous morphogenic regulator, a polynucleotide encoding a CRISPR-Cas effector protein, a polynucleotide encoding a CRISPR-Cas fusion protein, a polynucleotide encoding a cytosine deaminase, a polynucleotide encoding an adenine deaminase, and/or a polynucleotide comprising a guide nucleic acid), wherein the nucleic acid construct is operably associated with at least a control sequence (e.g., a promoter). Thus, some embodiments of the invention provide expression cassettes designed to express, for example, a nucleic acid construct of the invention. When an expression cassette comprises more than one polynucleotide, the polynucleotides may be operably linked to a single promoter that drives expression of all of the polynucleotides or the polynucleotides may be operably linked to one or more different promoters (e.g., three polynucleotides may be driven by one, two or three promoters in any combination). Thus, for example, a polynucleotide comprising a heterologous morphogenic regulator, a polynucleotide encoding a CRISPR-Cas effector protein, a polynucleotide encoding a deaminase, and a polynucleotide comprising a guide nucleic acid comprised in an expression cassette may each be operably associated with a single promoter or one or more of the polynucleotide(s) may be operably associated with separate promoters (e.g., two or three promoters) in any combination. In some embodiments, two or more different expression cassettes can be provided that one or more polynucleotides. For example, a first expression cassette may be provided that comprises a polynucleotide comprising a heterologous morphogenic regulator and a second expression cassette may be provided that comprises a polynucleotide encoding a CRISPR-Cas effector protein, a polynucleotide encoding a deaminase, and a polynucleotide comprising a guide nucleic acid that may each be operably associated with a single promoter or one or more of the polynucleotide(s) may be operably associated with separate promoters (e.g., two or three promoters) in any combination. The first expression cassette may comprise one or more polynucleotides that are the same as or different than one or more polynucleotides in the second expression cassette and vice versa.

[0073] In some embodiments, an expression cassette comprising the polynucleotides/nucleic acid constructs of the invention may be optimized for expression in an organism (e.g., an animal, a plant, a bacterium and the like).

[0074] An expression cassette comprising a nucleic acid construct of the invention may be chimeric, meaning that at least one of its components is heterologous with respect to at least one of its other components (e.g., a promoter from the host organism operably linked to a polynucleotide of interest to be expressed in the host organism, wherein the polynucleotide of interest is from a different organism than the host or is not normally found in association with that promoter). An expression cassette may also be one that is naturally occurring but has been obtained in a recombinant form useful for heterologous expression.

[0075] An expression cassette can optionally include a transcriptional and/or translational termination region (i.e., termination region) and/or an enhancer region that is functional in the selected host cell. A variety of transcriptional terminators and enhancers are known in the art and are available for use in expression cassettes. Transcriptional terminators are responsible for the termination of transcription and correct mRNA polyadenylation. A termination region and/or the enhancer region may be native to the transcriptional initiation region, may be native to a gene encoding a CRISPR-Cas effector protein or a gene encoding a deaminase, may be native to a host cell, or may be native to another source (e.g., foreign or heterologous to the promoter, to a gene encoding the CRISPR-Cas effector protein or a gene encoding the deaminase, to a host cell, or any combination thereof).

[0076] An expression cassette of the invention also can include a polynucleotide encoding a selectable marker, which can be used to select a transformed host cell. As used herein, “selectable marker” means a polynucleotide sequence that when expressed imparts a distinct phenotype to the host cell expressing the marker and thus allows such transformed cells to be distinguished from those that do not have the marker. Such a polynucleotide sequence may encode either a selectable or screenable marker, depending on whether the marker confers a trait that can be selected for by chemical means, such as by using a selective agent (e.g., an antibiotic and the like), or on whether the marker is simply a trait that one can identify through observation or testing, such as by screening (e.g., fluorescence). Many examples of suitable selectable markers are known in the art and can be used in the expression cassettes described herein.

[0077] The expression cassettes, the nucleic acid molecules/constructs and polynucleotide sequences described herein can be used in connection with vectors. The term “vector” refers to a composition for transferring, delivering or introducing a nucleic acid (or nucleic acids) into a cell. A vector comprises a nucleic acid construct comprising the nucleotide sequence(s) to be transferred, delivered or introduced. Vectors for use in transformation of host organisms are well known in the art. Non-limiting examples of general classes of vectors include viral vectors, plasmid vectors, phage vectors, phagemid vectors, cosmid vectors, fosmid vectors, bacteriophages, artificial chromosomes, minicircles, or *Agrobacterium* binary vectors in double or single stranded linear or circular form which may or may not be self transmissible or mobilizable. In some embodiments, a viral vector can include, but is not limited, to a retroviral, lentiviral, adenoviral, adeno-associated, or herpes simplex viral vector. A vector as defined herein can transform a prokaryotic or eukaryotic host either by integration into the cellular genome or exist extrachromosomally (e.g. autonomous replicating plasmid with an origin of replication).

Additionally included are shuttle vectors by which is meant a DNA vehicle capable, naturally or by design, of replication in two different host organisms, which may be selected from actinomycetes and related species, bacteria and eukaryotic (e.g. higher plant, mammalian, yeast or fungal cells). In some embodiments, the nucleic acid in the vector is under the control of, and operably linked to, an appropriate promoter or other regulatory elements for transcription in a host cell. The vector may be a bi-functional expression vector which functions in multiple hosts. In the case of genomic DNA, this may contain its own promoter and/or other regulatory elements and in the case of cDNA this may be under the control of an appropriate promoter and/or other regulatory elements for expression in the host cell. Accordingly, a nucleic acid construct of this invention and/or expression cassettes comprising the same may be comprised in vectors as described herein and as known in the art.

[0078] As used herein, “contact,” “contacting,” “contacted,” and grammatical variations thereof, refer to placing the components of a desired reaction together under conditions suitable for carrying out the desired reaction (e.g., transformation, transcriptional control, genome editing, nicking, and/or cleavage). Thus, for example, a target nucleic acid may be contacted with a nucleic acid construct of the invention encoding, for example, a nucleic acid binding domain (e.g., a DNA binding domain such as a sequence-specific DNA binding protein (e.g., polynucleotide-guided endonuclease, a CRISPR-Cas effector protein (e.g., CRISPR-Cas endonuclease), a zinc finger nuclease, a transcription activator-like effector nuclease (TALEN) and/or an Argonaute protein)), a guide nucleic acid, a deaminase (e.g., a cytosine and/or adenine deaminase), and a heterologous morphogenic regulator, under conditions whereby the CRISPR-Cas effector protein and heterologous morphogenic regulator are expressed, and the CRISPR-Cas effector protein forms a complex with the guide nucleic acid, the complex hybridizes to the target nucleic acid, and optionally the deaminase is recruited to the CRISPR-Cas effector protein (and thus, to the target nucleic acid) or the deaminase is fused to the CRISPR-Cas effector protein, thereby modifying the target nucleic acid. In some embodiments, the deaminase and the CRISPR-Cas effector protein localize at the target nucleic acid, optionally through covalent and/or non-covalent interactions.

[0079] As used herein, “modifying” or “modification” in reference to a target nucleic acid includes editing (e.g., mutating), covalent modification, exchanging/substituting nucleic acids/nucleotide bases, deleting, cleaving, nicking, and/or transcriptional control of a target nucleic acid.

[0080] “Recruit,” “recruiting” or “recruitment” as used herein refer to attracting one or more polypeptide(s) or polynucleotide(s) to another polypeptide or polynucleotide (e.g., to a particular location in a genome) using protein-protein interactions, RNA-protein interactions, and/or chemical interactions. Protein-protein interactions can include, but are not limited to, peptide tags (epitopes, multimerized epitopes) and corresponding affinity polypeptides, RNA recruiting motifs and corresponding affinity polypeptides, and/or chemical interactions. Example chemical interactions that may be useful with polypeptides and polynucleotides for the purpose of recruitment can include, but are not limited to, rapamycin-inducible dimerization of FRB-FKBP; Biotin-streptavidin interaction; SNAP tag (Hussain et al. *Curr Pharm Des.* 19(30):5437-42 (2013));

Halo tag (Los et al. *ACS Chem Biol.* 3(6):373-82 (2008)); CLIP tag (Gautier et al. *Chemistry & Biology* 15:128-136 (2008)); DmrA-DmrC heterodimer induced by a compound (Tak et al. *Nat Methods* 14(12):1163-1166 (2017)); Bifunctional ligand approaches (fuse two protein-binding chemicals together) (Voß et al. *Curr Opin Chemical Biology* 28:194-201 (2015)) (e.g. dihydrofolate reductase (DHFR) (Kopyteck et al. *Cell Chem Biol* 7(5):313-321 (2000)).

[0081] “Introducing,” “introduce,” “introduced” (and grammatical variations thereof) in the context of a polynucleotide of interest, polypeptide of interest, and/or editing system means presenting a nucleotide sequence of interest (e.g., polynucleotide, a nucleic acid construct, and/or a guide nucleic acid), a polypeptide of interest, and/or an editing system (e.g., a polynucleotide, polypeptide, and/or ribonucleoprotein) to a host organism or cell of said organism (e.g., host cell; e.g., a plant cell) in such a manner that the nucleotide sequence of interest, polypeptide of interest, and/or editing system gains access to the interior of a cell. Thus, for example, a polynucleotide comprising a heterologous morphogenic regulator and/or a nucleic acid construct encoding a CRISPR-Cas effector protein, a guide nucleic acid, and a deaminase may be introduced into a cell of an organism, thereby transforming the cell with the polynucleotide comprising a heterologous morphogenic regulator and/or the CRISPR-Cas effector protein, guide nucleic acid, and deaminase. Another example is that a polynucleotide comprising a heterologous morphogenic regulator and/or an editing system comprising a ribonucleoprotein may be introduced into a cell of an organism, thereby transforming the cell with the polynucleotide comprising a heterologous morphogenic regulator and/or the ribonucleoprotein. The ribonucleoprotein may comprise a CRISPR-Cas effector domain, guide nucleic acid, and optionally a deaminase domain. In some embodiments, a polypeptide comprising a heterologous morphogenic regulator and/or an editing system comprising a ribonucleoprotein may be introduced into a cell of an organism, thereby transforming the cell with the polypeptide comprising a heterologous morphogenic regulator and/or the ribonucleoprotein.

[0082] The term “transformation” as used herein refers to the introduction of a heterologous nucleic acid, polypeptide, and/or ribonucleoprotein into a cell. Transformation of a cell may be stable or transient. Thus, in some embodiments, a host cell or host organism may be stably transformed with a polynucleotide/nucleic acid molecule of the invention. In some embodiments, a host cell or host organism may be transiently transformed with a nucleic acid construct, a polypeptide and/or a ribonucleoprotein of the invention.

[0083] “Transient transformation” or “transiently transformed” in the context of a polynucleotide, polypeptide, and/or ribonucleoprotein means that a polynucleotide, polypeptide, and/or ribonucleoprotein is introduced into the cell and does not integrate into the genome of the cell.

[0084] By “stably introducing” or “stably introduced” in the context of a polynucleotide that is introduced into a cell is intended that the introduced polynucleotide is stably incorporated into the genome of the cell, and thus the cell is stably transformed with the polynucleotide.

[0085] “Stable transformation” or “stably transformed” as used herein means that a nucleic acid molecule is introduced into a cell and integrates into the genome of the cell. As such, the integrated nucleic acid molecule is capable of being inherited by the progeny thereof, more particularly, by the

progeny of multiple successive generations. “Genome” as used herein includes the nuclear and the plastid genome, and therefore includes integration of the nucleic acid into, for example, the chloroplast or mitochondrial genome. Stable transformation as used herein can also refer to a transgene that is maintained extrachromosomally, for example, as a minichromosome or a plasmid.

[0086] Transient transformation may be detected by, for example, an enzyme-linked immunosorbent assay (ELISA) or Western blot, which can detect the presence of a peptide or polypeptide encoded by one or more transgene introduced into an organism. Stable transformation of a cell can be detected by, for example, a Southern blot hybridization assay of genomic DNA of the cell with nucleic acid sequences which specifically hybridize with a nucleotide sequence of a transgene introduced into an organism (e.g., a plant). Stable transformation of a cell can be detected by, for example, a Northern blot hybridization assay of RNA of the cell with nucleic acid sequences which specifically hybridize with a nucleotide sequence of a transgene introduced into a host organism. Stable transformation of a cell can also be detected by, e.g., a polymerase chain reaction (PCR) or other amplification reactions as are well known in the art, employing specific primer sequences that hybridize with target sequence(s) of a transgene, resulting in amplification of the transgene sequence, which can be detected according to standard methods. Transformation can also be detected by direct sequencing and/or hybridization protocols well known in the art.

[0087] Accordingly, in some embodiments, nucleotide sequences, polynucleotides, nucleic acid constructs, and/or expression cassettes of the invention may be expressed transiently and/or they can be stably incorporated into the genome of the host organism. Thus, in some embodiments, a nucleic acid construct of the invention may be transiently introduced into a cell with a guide nucleic acid and as such, no DNA maintained in the cell.

[0088] A nucleic acid construct, polypeptide, and/or ribonucleoprotein of the invention can be introduced into a cell by any method known to those of skill in the art. In some embodiments of the invention, transformation of a cell comprises nuclear transformation. In other embodiments, transformation of a cell comprises plastid transformation (e.g., chloroplast transformation). In some embodiments of the invention, transformation of a cell comprises particle bombardment. In still further embodiments, the recombinant nucleic acid construct of the invention can be introduced into a cell via conventional breeding techniques.

[0089] Procedures for transforming both eukaryotic and prokaryotic organisms are well known and routine in the art and are described throughout the literature (See, for example, Jiang et al. 2013. *Nat. Biotechnol.* 31:233-239; Ran et al. *Nature Protocols* 8:2281-2308 (2013)).

[0090] A nucleotide sequence, polypeptide, and/or ribonucleoprotein therefore can be introduced into a host organism or its cell in any number of ways that are well known in the art. The methods of the invention do not depend on a particular method for introducing one or more nucleotide sequence(s), polypeptide(s), and/or ribonucleoprotein(s) into the organism, only that they gain access to the interior of at least one cell of the organism. Where more than one nucleotide sequence, polypeptide, and/or ribonucleoprotein is to be introduced, they can be assembled as part of a single construct (e.g., a single nucleic acid construct,) or as sepa-

rate constructs. Accordingly, a nucleotide sequence, polypeptide, and/or ribonucleoprotein can be introduced into the cell of interest in a single transformation event, and/or in separate transformation events, or, alternatively, where relevant, a nucleotide sequence can be incorporated into a plant, for example, as part of a breeding protocol.

[0091] The terms “genetically engineered host cell,” “recombinant host cell,” and “recombinant strain” are used interchangeably herein and refer to host cells that have been genetically engineered by a method of the present disclosure. Thus, the terms include a host cell (e.g., bacteria, yeast cell, fungal cell, CHO, human cell, plant cell, protoplast derived from plant, callus, etc.) that has been genetically altered, modified, or engineered, such that it exhibits an altered, modified, or different genotype and/or phenotype (e.g., when the genetic modification affects coding nucleic acid sequences), as compared to the naturally-occurring host cell from which it was derived. It is understood that the terms refer not only to the particular recombinant host cell in question, but also to the progeny or potential progeny of such a host cell.

[0092] The term “genetically engineered” may refer to any manipulation of a host cell’s genome (e.g., by insertion or deletion of nucleic acids).

[0093] The term “next generation plant breeding” refers to a host of plant breeding tools and methodologies that are available to today’s breeder. A key distinguishing feature of next generation plant breeding is that the breeder is no longer confined to relying upon observed phenotypic variation, in order to infer underlying genetic causes for a given trait. Rather, next generation plant breeding may include the utilization of molecular markers and marker assisted selection (MAS), such that the breeder can directly observe movement of alleles and genetic elements of interest from one plant in the breeding population to another, and is not confined to merely observing phenotype. Further, next generation plant breeding methods are not confined to utilizing natural genetic variation found within a plant population. Rather, the breeder utilizing next generation plant breeding methodology can access a host of modern genetic engineering tools that directly alter/change/edit the plant’s underlying genetic architecture in a targeted manner, in order to bring about a phenotypic trait of interest. In aspects, the plants bred with a next generation plant breeding methodology are indistinguishable from a plant that was bred in a traditional manner, as the resulting end product plant could theoretically be developed by either method. In particular aspects, a next generation plant breeding methodology may result in a plant that comprises: a genetic modification that is a deletion of any size; a genetic modification that is a single base pair substitution; a genetic modification that is an introduction of nucleic acid sequences from within the plant’s natural gene pool (e.g. any plant that could be crossed or bred with a plant of interest) or from editing of nucleic acid sequences in a plant to correspond to a sequence known to occur in the plant’s natural gene pool; and offspring of said plants.

[0094] The term “traditional plant breeding” refers to the utilization of natural variation found within a plant population as a source for alleles and genetic variants that impart a trait of interest to a given plant. Traditional breeding methods make use of crossing procedures that rely largely upon observed phenotypic variation to infer causative allele association. That is, traditional plant breeding relies upon

observations of expressed phenotype of a given plant to infer underlying genetic cause. These observations are utilized to inform the breeding procedure in order to move allelic variation into germplasm of interest. Further, traditional plant breeding has also been characterized as comprising random mutagenesis techniques, which can be used to introduce genetic variation into a given germplasm. These random mutagenesis techniques may include chemical and/or radiation-based mutagenesis procedures. Consequently, one key feature of traditional plant breeding, is that the breeder does not utilize a genetic engineering tool that directly alters/changes/edits the plant’s underlying genetic architecture in a targeted manner, in order to introduce genetic diversity and bring about a phenotypic trait of interest.

[0095] As used herein, the term “endogenous gene,” refers to a naturally occurring gene, in the location in which it is naturally found within the host cell genome. An endogenous gene as described herein can include alleles of naturally occurring genes that have been mutated according to any of a method of the present disclosure, i.e. an endogenous gene could have been modified at some point by traditional plant breeding methods and/or next generation plant breeding methods.

[0096] As used herein, the term “exogenous” refers to a substance coming from some source other than its native source. For example, the terms “exogenous protein,” or “exogenous gene” refer to a protein or gene from a non-native source, and that has been artificially supplied to a biological system. As used herein, the term “exogenous” is used interchangeably with the term “heterologous,” and refers to a substance coming from some source other than its native source.

[0097] As used herein, the term “heterologous” refers to a substance coming from some source or location other than its native source or location. In some embodiments, the term “heterologous nucleic acid” refers to a nucleic acid sequence that is not naturally found in the particular organism. For example, the term “heterologous promoter” may refer to a promoter that has been taken from one source organism and utilized in another organism, in which the promoter is not naturally found. However, the term “heterologous promoter” may also refer to a promoter that is from within the same source organism, but has merely been moved to a novel location, in which said promoter is not normally located.

[0098] As used herein, the term “nucleotide change” refers to, e.g., nucleotide substitution, deletion, and/or insertion, as is well understood in the art. For example, mutations contain alterations that produce single nucleotide substitutions, silent substitutions, additions, or deletions, but do not alter the properties or activities of the encoded protein or how the proteins are made.

[0099] As used herein, the term “protein modification” refers to, e.g., amino acid substitution, amino acid modification, deletion, and/or insertion, as is well understood in the art.

[0100] As used herein, the term “codon optimization” implies that the codon usage of a DNA or RNA is adapted to that of a cell or organism of interest to improve the transcription rate of said recombinant nucleic acid in the cell or organism of interest. The skilled person is well aware of the fact that a target nucleic acid can be modified at one position due to the codon degeneracy, whereas this modification will still lead to the same amino acid sequence at that

position after translation, which is achieved by codon optimization to take into consideration the species-specific codon usage of a target cell or organism.

[0101] As used herein, the term “naturally occurring” as applied to a nucleic acid, a polypeptide, a cell, or an organism, refers to a nucleic acid, polypeptide, cell, or organism that is found in nature. The term “naturally occurring” may refer to a gene or sequence derived from a naturally occurring source. Thus, for the purposes of this disclosure, a “non-naturally occurring” sequence is a sequence that has been synthesized, mutated, engineered, edited, or otherwise modified to have a different sequence from known natural sequences. In some embodiments, the modification may be at the protein level (e.g., amino acid substitutions, deletions, or insertions). In other embodiments, the modification may be at the DNA level (e.g., nucleotide substitutions, deletions, or insertions).

[0102] As used herein, the phrases “recombinant construct”, “expression construct”, “chimeric construct”, and “recombinant DNA construct” are used interchangeably herein. A recombinant construct comprises an artificial combination of nucleic acid fragments, e.g., regulatory and coding sequences that are not found together in nature. For example, a chimeric construct may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that found in nature. Such construct may be used by itself or may be used in conjunction with a vector. If a vector is used then the choice of vector is dependent upon the method that will be used to transform host cells as is well known to those skilled in the art. For example, a plasmid vector can be used. The skilled artisan is well aware of the genetic elements that must be present on the vector in order to successfully transform, select and propagate host cells comprising any of the isolated nucleic acid fragments of the disclosure. The skilled artisan will also recognize that different independent transformation events will result in different levels and patterns of expression (Jones et al., (1985) *EMBO J.* 4:2411-2418; De Almeida et al., (1989) *Mol. Gen. Genetics* 218:78-86), and thus that multiple events must be screened in order to obtain lines displaying the desired expression level and pattern. Such screening may be accomplished by Southern analysis of DNA, Northern analysis of mRNA expression, immunoblotting analysis of protein expression, or phenotypic analysis, among others. Vectors can be plasmids, viruses, bacteriophages, pro-viruses, phagemids, transposons, artificial chromosomes, and the like, that replicate autonomously or can integrate into a chromosome of a host cell. A vector can also be a naked RNA polynucleotide, a naked DNA polynucleotide, a polynucleotide composed of both DNA and RNA within the same strand, a poly-lysine-conjugated DNA or RNA, a peptide-conjugated DNA or RNA, a liposome-conjugated DNA, or the like, that is not autonomously replicating. As used herein, the term “expression” refers to the production of a functional end-product e.g., an mRNA or a protein (precursor or mature).

[0103] In some embodiments, an editing system of the present invention comprises a CRISPR-Cas effector protein. As used herein, a “CRISPR-Cas effector protein” is a protein or polypeptide or domain thereof that cleaves, cuts, or nicks a nucleic acid, binds a nucleic acid (e.g., a target nucleic acid and/or a guide nucleic acid), and/or that identifies, recog-

nizes, or binds a guide nucleic acid as defined herein. In some embodiments, a CRISPR-Cas effector protein may be an enzyme (e.g., a nuclease, endonuclease, nickase, etc.) or portion thereof and/or may function as an enzyme. In some embodiments, a CRISPR-Cas effector protein refers to a CRISPR-Cas nuclease polypeptide or domain thereof that comprises nuclease activity or in which the nuclease activity has been reduced or eliminated, and/or comprises nickase activity or in which the nickase has been reduced or eliminated, and/or comprises single stranded DNA cleavage activity (ss DNase activity) or in which the ss DNase activity has been reduced or eliminated, and/or comprises self-processing RNase activity or in which the self-processing RNase activity has been reduced or eliminated. A CRISPR-Cas effector protein may bind to a target nucleic acid. A CRISPR-Cas effector protein may be a Type I, II, III, IV, V, or VI CRISPR-Cas effector protein. In some embodiments, a CRISPR-Cas effector protein may be from a Type I CRISPR-Cas system, a Type II CRISPR-Cas system, a Type III CRISPR-Cas system, a Type IV CRISPR-Cas system, Type V CRISPR-Cas system, or a Type VI CRISPR-Cas system. In some embodiments, a CRISPR-Cas effector protein of the invention may be from a Type II CRISPR-Cas system or a Type V CRISPR-Cas system. In some embodiments, a CRISPR-Cas effector protein may be a Type II CRISPR-Cas effector protein, for example, a Cas9 effector protein. In some embodiments, a CRISPR-Cas effector protein may be Type V CRISPR-Cas effector protein, for example, a Cas12 effector protein. In some embodiments, a CRISPR-Cas effector protein may be a Cas12a polypeptide or domain thereof and optionally may have an amino acid sequence of any one of SEQ ID NOs:115-131 and/or a nucleotide sequence of any one of SEQ ID NOs:132-134.

[0104] In some embodiments, a CRISPR-Cas effector protein may be or include, but is not limited to, a Cas9, C2c1, C2c3, Cas12a (also referred to as Cpf1), Cas12b, Cas12c, Cas12d, Cas12e, Cas13a, Cas13b, Cas13c, Cas13d, Cas1, Cas1B, Cas2, Cas3, Cas3', Cas3'', Cas4, Cas5, Cas6, Cas7, Cas8, Cas9 (also known as Csn1 and Csx12), Cas10, Csy2, Csy3, Cse1, Cse2, Csc1, Csc2, Csa5, Csn2, Csm2, Csm3, Csm4, Csm5, Csm6, Cmr1, Cmr3, Cmr4, Cmr5, Cmr6, Csb1, Csb2, Csb3, Csx17, Csx14, Csx10, Csx16, CsaX, Csx3, Csx1, Csx15, Csf1, Csf2, Csf3, Csf4 (dinG), and/or Csf5 nuclease, optionally wherein the CRISPR-Cas effector protein may be a Cas9, Cas12a (Cpf1), Cas12b, Cas12c (C2c3), Cas12d (CasY), Cas12e (CasX), Cas12g, Cas12h, Cas12i, C2c4, C2c5, C2c8, C2c9, C2c10, Cas14a, Cas14b, and/or Cas14c effector protein.

[0105] In some embodiments, a CRISPR-Cas effector protein useful with the invention may comprise a mutation in its nuclease active site (e.g., RuvC, HNH, e.g., RuvC site of a Cas12a nuclease domain; e.g., RuvC site and/or HNH site of a Cas9 nuclease domain). A CRISPR-Cas effector protein having a mutation in its nuclease active site, and therefore, no longer comprising nuclease activity, is commonly referred to as “dead,” e.g., dCas9. In some embodiments, a CRISPR-Cas effector protein domain or polypeptide having a mutation in its nuclease active site may have impaired activity or reduced activity as compared to the same CRISPR-Cas effector protein without the mutation, e.g., a nickase, e.g., Cas9 nickase, Cas12a nickase.

[0106] A CRISPR Cas9 effector protein or CRISPR Cas9 effector domain useful with this invention may be any known or later identified Cas9 nuclease. In some embodi-

ments, a CRISPR Cas9 polypeptide can be a Cas9 polypeptide from, for example, *Streptococcus* spp. (e.g., *S. pyogenes*, *S. thermophilus*), *Lactobacillus* spp., *Bifidobacterium* spp., *Kandleria* spp., *Leuconostoc* spp., *Oenococcus* spp., *Pediococcus* spp., *Weissella* spp., and/or *Olsenella* spp. In some embodiments, a CRISPR-Cas effector protein may be a Cas9 polypeptide or domain thereof and optionally may have a nucleotide sequence of any one of SEQ ID NOs:63-73 and/or an amino acid sequence of any one of SEQ ID NOs:74-75.

[0107] In some embodiments, the CRISPR-Cas effector protein may be a Cas9 polypeptide derived from *Streptococcus pyogenes* and recognizes the PAM sequence motif NGG, NAG, NGA (Mali et al, Science 2013; 339(6121): 823-826). In some embodiments, the CRISPR-Cas effector protein may be a Cas9 polypeptide derived from *Streptococcus thermophilus* and recognizes the PAM sequence motif NGGNG and/or NNAGAAW (W=A or T) (See, e.g., Horvath et al, Science, 2010; 327(5962): 167-170, and Deveau et al, J Bacteriol 2008; 190(4): 1390-1400). In some embodiments, the CRISPR-Cas effector protein may be a Cas9 polypeptide derived from *Streptococcus mutans* and recognizes the PAM sequence motif NGG and/or NAAR (R=A or G) (See, e.g., Deveau et al, J BACTERIOL 2008; 190(4): 1390-1400). In some embodiments, the CRISPR-Cas effector protein may be a Cas9 polypeptide derived from *Streptococcus aureus* and recognizes the PAM sequence motif NNGRR (R=A or G). In some embodiments, the CRISPR-Cas effector protein may be a Cas9 protein derived from *S. aureus*, which recognizes the PAM sequence motif N GRRT (R=A or G). In some embodiments, the CRISPR-Cas effector protein may be a Cas9 polypeptide derived from *S. aureus*, which recognizes the PAM sequence motif N GRRV (R=A or G). In some embodiments, the CRISPR-Cas effector protein may be a Cas9 polypeptide that is derived from *Neisseria meningitidis* and recognizes the PAM sequence motif N GATT or N GCTT (R=A or G, V=A, G or C) (See, e.g., Hou et al, PNAS 2013, 1-6). In the aforementioned embodiments, N can be any nucleotide residue, e.g., any of A, G, C or T. In some embodiments, the CRISPR-Cas effector protein may be a Cas13a protein derived from *Leptotrichia shahii*, which recognizes a protospacer flanking sequence (PFS) (or RNA PAM (rPAM)) sequence motif of a single 3' A, U, or C, which may be located within the target nucleic acid.

[0108] A Type V CRISPR-Cas effector protein useful with embodiments of the invention may be any Type V CRISPR-Cas nuclease. A Type V CRISPR-Cas nuclease useful with this invention as an effector protein can include, but is not limited, to Cas12a (Cpf1), Cas12b, Cas12c (C2c3), Cas12d (CasY), Cas12e (CasX), Cas12g, Cas12h, Cas12i, C2c1, C2c4, C2c5, C2c8, C2c9, C2c10, Cas14a, Cas14b, and/or Cas14c nuclease. In some embodiments, a Type V CRISPR-Cas nuclease polypeptide or domain useful with embodiments of the invention may be a Cas12a polypeptide or domain. In some embodiments, a Type V CRISPR-Cas effector protein or domain useful with embodiments of the invention may be a nickase, optionally, a Cas12a nickase.

[0109] In some embodiments, the CRISPR-Cas effector protein may be derived from Cas12a, which is a Type V Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR)-Cas nuclease. Cas12a differs in several respects from the more well-known Type II CRISPR Cas9 nuclease. For example, Cas9 recognizes a G-rich protospacer-adjacent

motif (PAM) that is 3' to its guide RNA (gRNA, sgRNA, crRNA, crDNA, CRISPR array) binding site (protospacer, target nucleic acid, target DNA) (3'-NGG), while Cas12a recognizes a T-rich PAM that is located 5' to the target nucleic acid (5'-TTN, 5'-TTTN. In fact, the orientations in which Cas9 and Cas12a bind their guide RNAs are very nearly reversed in relation to their N and C termini. Furthermore, Cas12a enzymes use a single guide RNA (gRNA, CRISPR array, crRNA) rather than the dual guide RNA (sgRNA (e.g., crRNA and tracrRNA)) found in natural Cas9 systems, and Cas12a processes its own gRNAs. Additionally, Cas12a nuclease activity produces staggered DNA double stranded breaks instead of blunt ends produced by Cas9 nuclease activity, and Cas12a relies on a single RuvC domain to cleave both DNA strands, whereas Cas9 utilizes an HNH domain and a RuvC domain for cleavage.

[0110] A CRISPR Cas12a effector protein/domain useful with this invention may be any known or later identified Cas12a polypeptide (previously known as Cpf1) (see, e.g., U.S. Pat. No. 9,790,490, which is incorporated by reference for its disclosures of Cpf1 (Cas12a) sequences). The term "Cas12a", "Cas12a polypeptide" or "Cas12a domain" refers to an RNA-guided nuclease comprising a Cas12a polypeptide, or a fragment thereof, which comprises the guide nucleic acid binding domain of Cas12a and/or an active, inactive, or partially active DNA cleavage domain of Cas12a. In some embodiments, a Cas12a useful with the invention may comprise a mutation in the nuclease active site (e.g., RuvC site of the Cas12a domain). A Cas12a domain or Cas12a polypeptide having a mutation in its nuclease active site, and therefore, no longer comprising nuclease activity, is commonly referred to as deadCas12a (e.g., dCas12a). In some embodiments, a Cas12a domain or Cas12a polypeptide having a mutation in its nuclease active site may have impaired activity, e.g., may have nickase activity.

[0111] In some embodiments, a CRISPR-Cas effector protein may be optimized for expression in an organism, for example, in an animal, a plant, a fungus, an archaeon, or a bacterium. In some embodiments, a CRISPR-Cas effector protein (e.g., Cas12a polypeptide/domain or a Cas9 polypeptide/domain) may be optimized for expression in a plant.

[0112] Any deaminase domain/polypeptide useful for base editing may be used with this invention. A "cytosine deaminase" and "cytidine deaminase" as used herein refer to a polypeptide or domain thereof that catalyzes or is capable of catalyzing cytosine deamination in that the polypeptide or domain catalyzes or is capable of catalyzing the removal of an amine group from a cytosine base. Thus, a cytosine deaminase may result in conversion of cytosine to a thymidine (through a uracil intermediate), causing a C to T conversion, or a G to A conversion in the complementary strand in the genome. Thus, in some embodiments, the cytosine deaminase encoded by the polynucleotide of the invention generates a C→T conversion in the sense (e.g., "+"; template) strand of the target nucleic acid or a G→A conversion in antisense (e.g., "-", complementary) strand of the target nucleic acid. In some embodiments, a cytosine deaminase encoded by a polynucleotide of the invention generates a C to T, G, or A conversion in the complementary strand in the genome.

[0113] A cytosine deaminase useful with this invention may be any known or later identified cytosine deaminase from any organism (see, e.g., U.S. Pat. No. 10,167,457 and

Thuronyi et al. *Nat. Biotechnol.* 37:1070-1079 (2019), each of which is incorporated by reference herein for its disclosure of cytosine deaminases). Cytosine deaminases can catalyze the hydrolytic deamination of cytidine or deoxycytidine to uridine or deoxyuridine, respectively. Thus, in some embodiments, a deaminase or deaminase domain useful with this invention may be a cytidine deaminase domain, catalyzing the hydrolytic deamination of cytosine to uracil. In some embodiments, a cytosine deaminase may be a variant of a naturally-occurring cytosine deaminase, including, but not limited to, a primate (e.g., a human, monkey, chimpanzee, gorilla), a dog, a cow, a rat or a mouse. Thus, in some embodiments, an cytosine deaminase useful with the invention may be about 70% to about 100% identical to a wild type cytosine deaminase (e.g., about 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical, and any range or value therein, to a naturally occurring cytosine deaminase).

[0114] In some embodiments, a cytosine deaminase useful with the invention may be an apolipoprotein B mRNA-editing complex (APOBEC) family deaminase. In some embodiments, the cytosine deaminase may be an APOBEC1 deaminase, an APOBEC2 deaminase, an APOBEC3A deaminase, an APOBEC3B deaminase, an APOBEC3C deaminase, an APOBEC3D deaminase, an APOBEC3F deaminase, an APOBEC3G deaminase, an APOBEC3H deaminase, an APOBEC4 deaminase, a human activation induced deaminase (hAID), an rAPOBEC1, FERNY, and/or a CDA1, optionally a pmCDA1, an atCDA1 (e.g., At2g19570), and evolved versions of the same. In some embodiments, the cytosine deaminase may be an APOBEC1 deaminase having the amino acid sequence of SEQ ID NO:76. In some embodiments, the cytosine deaminase may be an APOBEC3A deaminase having the amino acid sequence of SEQ ID NO:77. In some embodiments, the cytosine deaminase may be an CDA1 deaminase, optionally a CDA1 having the amino acid sequence of SEQ ID NO:78. In some embodiments, the cytosine deaminase may be a FERNY deaminase, optionally a FERNY having the amino acid sequence of SEQ ID NO:79. In some embodiments, the cytosine deaminase may be a rAPOBEC1 deaminase, optionally a rAPOBEC1 deaminase having the amino acid sequence of SEQ ID NO:80. In some embodiments, the cytosine deaminase may be a hAID deaminase, optionally a hAID having the amino acid sequence of SEQ ID NO:81 or SEQ ID NO:82. In some embodiments, a cytosine deaminase useful with the invention may be about 70% to about 100% identical (e.g., 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or 100% identical) to the amino acid sequence of a naturally occurring cytosine deaminase (e.g., “evolved deaminases”) (see, e.g., SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85). In some embodiments, a cytosine deaminase useful with the invention may be about 70% to about 99.5% identical (e.g., about 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.5% identical) to the amino acid sequence of any one of SEQ ID NOs:76-85 (e.g., at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%,

at least 98%, at least 99%, or at least 99.5% identical to the amino acid sequence of any one of SEQ ID NOs:76-85). In some embodiments, a polynucleotide encoding a cytosine deaminase may be codon optimized for expression in a plant and the codon optimized polypeptide may be about 70% to 99.5% identical to the reference polynucleotide.

[0115] An “adenine deaminase” and “adenosine deaminase” as used herein refer to a polypeptide or domain thereof that catalyzes or is capable of catalyzing the hydrolytic deamination (e.g., removal of an amine group from adenine) of adenine or adenosine. In some embodiments, an adenine deaminase may catalyze the hydrolytic deamination of adenosine or deoxyadenosine to inosine or deoxyinosine, respectively. In some embodiments, the adenosine deaminase may catalyze the hydrolytic deamination of adenine or adenosine in DNA. In some embodiments, an adenine deaminase encoded by a nucleic acid construct of the invention may generate an A→G conversion in the sense (e.g., “+”; template) strand of the target nucleic acid or a T→C conversion in the antisense (e.g., “-”, complementary) strand of the target nucleic acid. An adenine deaminase useful with this invention may be any known or later identified adenine deaminase from any organism (see, e.g., U.S. Pat. No. 10,113,163, which is incorporated by reference herein for its disclosure of adenine deaminases).

[0116] In some embodiments, an adenosine deaminase may be a variant of a naturally-occurring adenine deaminase. Thus, in some embodiments, an adenosine deaminase may be about 70% to 100% identical to a wild type adenine deaminase (e.g., about 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical, and any range or value therein, to a naturally occurring adenine deaminase). In some embodiments, the deaminase or deaminase does not occur in nature and may be referred to as an engineered, mutated or evolved adenosine deaminase. Thus, for example, an engineered, mutated or evolved adenine deaminase polypeptide or an adenine deaminase domain may be about 70% to 99.9% identical to a naturally occurring adenine deaminase polypeptide/domain (e.g., about 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8% or 99.9% identical, and any range or value therein, to a naturally occurring adenine deaminase polypeptide or adenine deaminase domain). In some embodiments, the adenosine deaminase may be from a bacterium, (e.g., *Escherichia coli*, *Staphylococcus aureus*, *Haemophilus influenzae*, *Caulobacter crescentus*, and the like). In some embodiments, a polynucleotide encoding an adenine deaminase polypeptide/domain may be codon optimized for expression in a plant.

[0117] In some embodiments, an adenine deaminase domain may be a wild type tRNA-specific adenosine deaminase domain, e.g., a tRNA-specific adenosine deaminase (TadA) and/or a mutated/evolved adenosine deaminase domain, e.g., mutated/evolved tRNA-specific adenosine deaminase domain (TadA*). In some embodiments, a TadA domain may be from *E. coli*. In some embodiments, the TadA may be modified, e.g., truncated, missing one or more N-terminal and/or C-terminal amino acids relative to a full-length TadA (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13,

14, 15, 6, 17, 18, 19, or 20 N-terminal and/or C terminal amino acid residues may be missing relative to a full length TadA. In some embodiments, a TadA polypeptide or TadA domain does not comprise an N-terminal methionine. In some embodiments, a wild type *E. coli* TadA comprises the amino acid sequence of SEQ ID NO:86. In some embodiments, a mutated/evolved *E. coli* TadA* comprises the amino acid sequence of SEQ ID NOs:87-90 (e.g., SEQ ID NOs: 87, 88, 89, or 90). In some embodiments, a polynucleotide encoding a TadA/TadA* may be codon optimized for expression in a plant. In some embodiments, an adenine deaminase may comprise all or a portion of an amino acid sequence of any one of SEQ ID NOs:91-96. In some embodiments, an adenine deaminase may comprise all or a portion of an amino acid sequence of any one of SEQ ID NOs:86-96.

[0118] In some embodiments, a nucleic acid construct of this invention may further encode a glycosylase inhibitor (e.g., a uracil glycosylase inhibitor (UGI) such as uracil-DNA glycosylase inhibitor). Thus, in some embodiments, a nucleic acid construct encoding a CRISPR-Cas effector protein and a cytosine deaminase and/or adenine deaminase may further encode a glycosylase inhibitor, optionally wherein the glycosylase inhibitor may be codon optimized for expression in a plant. In some embodiments, the invention provides fusion proteins comprising a CRISPR-Cas effector polypeptide and a UGI and/or one or more polynucleotides encoding the same, optionally wherein the one or more polynucleotides may be codon optimized for expression in a plant. In some embodiments, the invention provides fusion proteins comprising a CRISPR-Cas effector polypeptide, a deaminase domain (e.g., an adenine deaminase domain and/or a cytosine deaminase domain) and optionally a UGI and/or one or more polynucleotides encoding the same, optionally wherein the one or more polynucleotides may be codon optimized for expression in a plant. In some embodiments, the invention provides fusion proteins, wherein a CRISPR-Cas effector polypeptide, a deaminase domain, and/or a UGI may be fused to any combination of peptide tags and affinity polypeptides as described herein, which may thereby recruit the deaminase domain and/or UGI to the CRISPR-Cas effector polypeptide and to a target nucleic acid. In some embodiments, a guide nucleic acid may be linked to a recruiting RNA motif and one or more of the deaminase domain and/or UGI may be fused to an affinity polypeptide that is capable of interacting with the recruiting RNA motif, thereby recruiting the deaminase domain and UGI to a target nucleic acid.

[0119] A “uracil glycosylase inhibitor” useful with the invention may be any protein or polypeptide that is capable of inhibiting a uracil-DNA glycosylase base-excision repair enzyme. In some embodiments, a UGI domain comprises a wild type UGI or a fragment thereof. In some embodiments, a UGI domain useful with the invention may be about 70% to about 100% identical (e.g., 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or 100% identical and any range or value therein) to the amino acid sequence of a naturally occurring UGI domain. In some embodiments, a UGI domain may comprise the amino acid sequence of SEQ ID NO:97 or a polypeptide having about 70% to about 99.5% identity to the amino acid sequence of SEQ ID NO:97 (e.g., at least 80%, at least 85%, at least 90%, at least 92%,

at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to the amino acid sequence of SEQ ID NO:97). For example, in some embodiments, a UGI domain may comprise a fragment of the amino acid sequence of SEQ ID NO:97 that is 100% identical to a portion of consecutive nucleotides (e.g., 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80 consecutive nucleotides; e.g., about 10, 15, 20, 25, 30, 35, 40, 45, to about 50, 55, 60, 65, 70, 75, 80 consecutive nucleotides) of the amino acid sequence of SEQ ID NO:97. In some embodiments, a UGI domain may be a variant of a known UGI (e.g., SEQ ID NO:97) having about 70% to about 99.5% identity (e.g., 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% identity, and any range or value therein) to the known UGI. In some embodiments, a polynucleotide encoding a UGI may be codon optimized for expression in a plant (e.g., a plant) and the codon optimized polypeptide may be about 70% to about 99.5% identical to the reference polynucleotide.

[0120] The nucleic acid constructs of the invention comprising a CRISPR-Cas effector protein or a fusion protein thereof may be used in combination with a guide nucleic acid (e.g., guide RNA (gRNA), CRISPR array, CRISPR RNA, crRNA), designed to function with the encoded CRISPR-Cas effector protein or domain, to modify a target nucleic acid. A guide nucleic acid useful with this invention may comprise at least one spacer sequence and at least one repeat sequence. The guide nucleic acid is capable of forming a complex with the CRISPR-Cas nuclease domain encoded and expressed by a nucleic acid construct of the invention and the spacer sequence is capable of hybridizing to a target nucleic acid, thereby guiding the complex to the target nucleic acid, wherein the target nucleic acid may be modified (e.g., cleaved or edited) and/or modulated (e.g., modulating transcription) by a deaminase (e.g., a cytosine deaminase and/or adenine deaminase, optionally present in and/or recruited to the complex).

[0121] As an example, a nucleic acid construct encoding a Cas9 domain linked to a cytosine deaminase domain (e.g., a fusion protein) may be used in combination with a Cas9 guide nucleic acid to modify a target nucleic acid, wherein the cytosine deaminase domain of the fusion protein deaminates a cytosine base in the target nucleic acid, thereby editing the target nucleic acid. In a further example, a nucleic acid construct encoding a Cas9 domain linked to an adenine deaminase domain (e.g., a fusion protein) may be used in combination with a Cas9 guide nucleic acid to modify a target nucleic acid, wherein the adenine deaminase domain of the fusion protein deaminates an adenosine base in the target nucleic acid, thereby editing the target nucleic acid. In some embodiments, a CRISPR-Cas effector protein (e.g., Cas9) is not fused to a cytosine deaminase and/or adenine deaminase.

[0122] Likewise, a nucleic acid construct encoding a Cas12a domain (or other selected CRISPR-Cas nuclease, e.g., C2c1, C2c3, Cas12b, Cas12c, Cas12d, Cas12e, Cas13a, Cas13b, Cas13c, Cas13d, Cas1, Cas1B, Cas2, Cas3, Cas3', Cas3'', Cas4, Cas5, Cas6, Cas7, Cas8, Cas9 (also known as Csn1 and Csx12), Cas10, Csy1, Csy2, Csy3, Cse1, Cse2, Csc1, Csc2, Csa5, Csn2, Csm2, Csm3, Csm4, Csm5, Csm6, Cmr1, Cmr3, Cmr4, Cmr5, Cmr6, Csb1, Csb2, Csb3, Csx17, Csx14, Csx10, Csx16, CsaX, Csx3, Csx1, Csx15, Csf1,

Csf2, Csf3, Csf4 (dinG), and/or Csf5) may be linked to a cytosine deaminase domain or adenine deaminase domain (e.g., fusion protein) and may be used in combination with a Cas12a guide nucleic acid (or the guide nucleic acid for the other selected CRISPR-Cas nuclease) to modify a target nucleic acid, wherein the cytosine deaminase domain or adenine deaminase domain of the fusion protein deaminates a cytosine base or adenosine base, respectively, in the target nucleic acid, thereby editing the target nucleic acid.

[0123] A “guide nucleic acid,” “guide RNA,” “gRNA,” “CRISPR RNA/DNA” “crRNA” or “crDNA” as used herein means a nucleic acid that comprises at least one spacer sequence, which is complementary to (and hybridizes to) a target DNA (e.g., protospacer), and at least one repeat sequence (e.g., a repeat of a Type V Cas12a CRISPR-Cas system, or a fragment or portion thereof; a repeat of a Type II Cas9 CRISPR-Cas system, or fragment thereof; a repeat of a Type V C2c 1 CRISPR Cas system, or a fragment thereof; a repeat of a CRISPR-Cas system of, for example, C2c3, Cas12a (also referred to as Cpf1), Cas12b, Cas12c, Cas12d, Cas12e, Cas13a, Cas13b, Cas13c, Cas13d, Cas1, Cas1B, Cas2, Cas3, Cas3', Cas4, Cas5, Cas6, Cas7, Cas8, Cas9 (also known as Csn1 and Csx12), Cas10, Csy1, Csy2, Csy3, Cse1, Cse2, Csc1, Csc2, Csa5, Csn2, Csm2, Csm3, Csm4, Csm5, Csm6, Cmr1, Cmr3, Cmr4, Cmr5, Cmr6, Csb1, Csb2, Csb3, Csx17, Csx14, Csx10, Csx16, CsaX, Csx3, Csx1, Csx15, Csf1, Csf2, Csf3, Csf4 (dinG), and/or Csf5, or a fragment thereof), wherein the repeat sequence may be linked to the 5' end and/or the 3' end of the spacer sequence. In some embodiments, the guide nucleic acid comprises DNA. In some embodiments, the guide nucleic acid comprises RNA (e.g., is a guide RNA). The design of a gRNA of this invention may be based on a Type I, Type II, Type III, Type IV, Type V, or Type VI CRISPR-Cas system.

[0124] In some embodiments, a Cas12a gRNA may comprise, from 5' to 3', a repeat sequence (full length or portion thereof (“handle”); e.g., pseudoknot-like structure) and a spacer sequence.

[0125] In some embodiments, a guide nucleic acid may comprise more than one repeat sequence-spacer sequence (e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, or more repeat-spacer sequences) (e.g., repeat-spacer-repeat, e.g., repeat-spacer-repeat-spacer-repeat-spacer-repeat-spacer-repeat-spacer, and the like). The guide nucleic acids of this invention are synthetic, human-made and not found in nature. A gRNA can be quite long and may be used as an aptamer (like in the MS2 recruitment strategy) or other RNA structures hanging off the spacer.

[0126] A “repeat sequence” as used herein, refers to, for example, any repeat sequence of a wild-type CRISPR Cas locus (e.g., a Cas9 locus, a Cas12a locus, a C2c1 locus, etc.) or a repeat sequence of a synthetic crRNA that is functional with the CRISPR-Cas effector protein encoded by the nucleic acid constructs of the invention. A repeat sequence useful with this invention can be any known or later identified repeat sequence of a CRISPR-Cas locus (e.g., Type I, Type II, Type III, Type IV, Type V or Type VI) or it can be a synthetic repeat designed to function in a Type I, II, III, IV, V or VI CRISPR-Cas system. A repeat sequence may comprise a hairpin structure and/or a stem loop structure. In some embodiments, a repeat sequence may form a pseudoknot-like structure at its 5' end (i.e., “handle”). Thus, in some embodiments, a repeat sequence can be identical to or

substantially identical to a repeat sequence from wild-type Type I CRISPR-Cas loci, Type II, CRISPR-Cas loci, Type III, CRISPR-Cas loci, Type IV CRISPR-Cas loci, Type V CRISPR-Cas loci and/or Type VI CRISPR-Cas loci. A repeat sequence from a wild-type CRISPR-Cas locus may be determined through established algorithms, such as using the CRISPRfinder offered through CRISPRdb (see, Grissa et al. *Nucleic Acids Res.* 35(Web Server issue):W52-7). In some embodiments, a repeat sequence or portion thereof is linked at its 3' end to the 5' end of a spacer sequence, thereby forming a repeat-spacer sequence (e.g., guide nucleic acid, guide RNA/DNA, crRNA, crDNA).

[0127] In some embodiments, a repeat sequence comprises, consists essentially of, or consists of at least 10 nucleotides depending on the particular repeat and whether the guide nucleic acid comprising the repeat is processed or unprocessed (e.g., about 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50 to 100 or more nucleotides, or any range or value therein; e.g., about). In some embodiments, a repeat sequence comprises, consists essentially of, or consists of about 10 to about 20, about 10 to about 30, about 10 to about 45, about 10 to about 50, about 15 to about 30, about 15 to about 40, about 15 to about 45, about 15 to about 50, about 20 to about 30, about 20 to about 40, about 20 to about 50, about 30 to about 40, about 40 to about 80, about 50 to about 100 or more nucleotides.

[0128] A repeat sequence linked to the 5' end of a spacer sequence can comprise a portion of a repeat sequence (e.g., 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 or more contiguous nucleotides of a wild type repeat sequence). In some embodiments, a portion of a repeat sequence linked to the 5' end of a spacer sequence can be about five to about ten consecutive nucleotides in length (e.g., about 5, 6, 7, 8, 9, 10 nucleotides) and have at least 90% sequence identity (e.g., at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more) to the same region (e.g., 5' end) of a wild type CRISPR Cas repeat nucleotide sequence. In some embodiments, a portion of a repeat sequence may comprise a pseudoknot-like structure at its 5' end (e.g., “handle”).

[0129] A “spacer sequence” as used herein is a nucleotide sequence that is complementary to a target nucleic acid (e.g., target DNA) (e.g., protospacer). The spacer sequence can be fully complementary or substantially complementary (e.g., at least about 70% complementary (e.g., about 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more)) to a target nucleic acid. Thus, in some embodiments, the spacer sequence can have one, two, three, four, or five mismatches as compared to the target nucleic acid, which mismatches can be contiguous or noncontiguous. In some embodiments, the spacer sequence can have 70% complementarity to a target nucleic acid. In other embodiments, the spacer nucleotide sequence can have 80% complementarity to a target nucleic acid. In still other embodiments, the spacer nucleotide sequence can have 85%, 90%, 95%, 96%, 97%, 98%, 99% or 99.5% complementarity, and the like, to the target nucleic acid (protospacer). In some embodiments, the spacer sequence is 100% complementary to the target nucleic acid. A spacer sequence may have a length from about 15 nucleotides to about 30 nucleotides (e.g., 15, 16, 17, 18, 19, 20, 21,

22, 23, 24, 25, 26, 27, 28, 29, or 30 nucleotides, or any range or value therein). Thus, in some embodiments, a spacer sequence may have complete complementarity or substantial complementarity over a region of a target nucleic acid (e.g., protospacer) that is at least about 15 nucleotides to about 30 nucleotides in length. In some embodiments, the spacer is about 20 nucleotides in length. In some embodiments, the spacer is about 21, 22, or 23 nucleotides in length.

[0130] In some embodiments, the 5' region of a spacer sequence of a guide nucleic acid may be identical to a target DNA, while the 3' region of the spacer may be substantially complementary to the target DNA (e.g., Type V CRISPR-Cas), or the 3' region of a spacer sequence of a guide nucleic acid may be identical to a target DNA, while the 5' region of the spacer may be substantially complementary to the target DNA (e.g., Type II CRISPR-Cas), and therefore, the overall complementarity of the spacer sequence to the target DNA may be less than 100%. Thus, for example, in a guide for a Type V CRISPR-Cas system, the first 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 nucleotides in the 5' region (i.e., seed region) of, for example, a 20 nucleotide spacer sequence may be 100% complementary to the target DNA, while the remaining nucleotides in the 3' region of the spacer sequence are substantially complementary (e.g., at least about 70% complementary) to the target DNA. In some embodiments, the first 1 to 8 nucleotides (e.g., the first 1, 2, 3, 4, 5, 6, 7, 8, nucleotides, and any range therein) of the 5' end of the spacer sequence may be 100% complementary to the target DNA, while the remaining nucleotides in the 3' region of the spacer sequence are substantially complementary (e.g., at least about 50% complementary (e.g., 50%, 55%, 60%, 65%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more)) to the target DNA.

[0131] As a further example, in a guide for a Type II CRISPR-Cas system, the first 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 nucleotides in the 3' region (i.e., seed region) of, for

or it may be inserted into the recruiting guide nucleic acid (e.g., within the hairpin loop).

[0132] In some embodiments, a seed region of a spacer may be about 8 to about 10 nucleotides in length, about 5 to about 6 nucleotides in length, or about 6 nucleotides in length.

[0133] A “target nucleic acid,” “target DNA,” “target nucleotide sequence,” “target region,” and a “target region in the genome” are used interchangeably herein and refer to a region of a plant’s genome that is targeted by an editing system (or a component thereof) as described herein. In some embodiments, a target nucleic acid is fully complementary (100% complementary) or substantially complementary (e.g., at least 70% complementary (e.g., 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more)) to a spacer sequence in a guide nucleic acid of this invention. A target region useful for a CRISPR-Cas system may be located immediately 3' (e.g., Type V CRISPR-Cas system) or immediately 5' (e.g., Type II CRISPR-Cas system) to a PAM sequence in the genome of the organism (e.g., a plant genome). A target region may be selected from any region of at least 15 consecutive nucleotides (e.g., 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 nucleotides, and the like) located immediately adjacent to a PAM sequence.

[0134] A “protospacer sequence” refers to the target double stranded DNA and specifically to the portion of the target DNA (e.g., or target region in the genome) that is fully or substantially complementary (and hybridizes) to the spacer sequence of the CRISPR repeat-spacer sequences (e.g., guide nucleic acids, CRISPR arrays, crRNAs).

[0135] In the case of Type V CRISPR-Cas (e.g., Cas12a) systems and Type II CRISPR-Cas (Cas9) systems, the protospacer sequence is flanked by (e.g., immediately adjacent to) a protospacer adjacent motif (PAM). For Type IV CRISPR-Cas systems, the PAM is located at the 5' end on the non-target strand and at the 3' end of the target strand (see below, as an example).

```

5'-NNNNNNNNNNNNNNNNNNNN-3'   RNA Spacer           (SEQ ID NO: 98)
   |||||
3'-AAANNNNNNNNNNNNNNNNNNNN-5' Target strand       (SEQ ID NO: 99)
   ||||
5'-TTNNNNNNNNNNNNNNNNNNNN-3' Non-target strand (SEQ ID NO: 100)

```

example, a 20 nucleotide spacer sequence may be 100% complementary to the target DNA, while the remaining nucleotides in the 5' region of the spacer sequence are substantially complementary (e.g., at least about 70% complementary) to the target DNA. In some embodiments, the first 1 to 10 nucleotides (e.g., the first 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 nucleotides, and any range therein) of the 3' end of the spacer sequence may be 100% complementary to the target DNA, while the remaining nucleotides in the 5' region of the spacer sequence are substantially complementary (e.g., at least about 50% complementary (e.g., at least about 50%, 55%, 60%, 65%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more or any range or value therein)) to the target DNA. A recruiting guide RNA further comprises one or more recruiting motifs as described herein, which may be linked to the 5' end of the guide or the 3' end

[0136] In the case of Type II CRISPR-Cas (e.g., Cas9) systems, the PAM is located immediately 3' of the target region. The PAM for Type I CRISPR-Cas systems is located 5' of the target strand. There is no known PAM for Type III CRISPR-Cas systems. Makarova et al. describes the nomenclature for all the classes, types and subtypes of CRISPR systems (*Nature Reviews Microbiology* 13:722-736 (2015)). Guide structures and PAMs are described in by R. Barrangou (*Genome Biol.* 16:247 (2015)).

[0137] Canonical Cas12a PAMs are T rich. In some embodiments, a canonical Cas12a PAM sequence may be 5'-TTN, 5'-TTTN, or 5'-TTTV. In some embodiments, canonical Cas9 (e.g., *S. pyogenes*) PAMs may be 5'-NGG-3'. In some embodiments, non-canonical PAMs may be used but may be less efficient.

[0138] Additional PAM sequences may be determined by those skilled in the art through established experimental and computational approaches. Thus, for example, experimental approaches include targeting a sequence flanked by all

possible nucleotide sequences and identifying sequence members that do not undergo targeting, such as through the transformation of target plasmid DNA (Esvelt et al. 2013. *Nat. Methods* 10:1116-1121; Jiang et al. 2013. *Nat. Biotechnol.* 31:233-239). In some aspects, a computational approach can include performing BLAST searches of natural spacers to identify the original target DNA sequences in bacteriophages or plasmids and aligning these sequences to determine conserved sequences adjacent to the target sequence (Briner and Barrangou. 2014. *Appl. Environ. Microbiol.* 80:994-1001; Mojica et al. 2009. *Microbiology* 155:733-740).

[0139] In some embodiments, the present invention provides expression cassettes and/or vectors comprising the nucleic acid constructs of the invention (e.g., one or more components of an editing system of the invention). In some embodiments, expression cassettes and/or vectors comprising the nucleic acid constructs of the invention and/or one or more guide nucleic acids may be provided. In some embodiments, a nucleic acid construct of the invention comprising a heterologous morphogenic regulator and encoding a base editor (e.g., a construct comprising a CRISPR-Cas effector domain and a deaminase domain (e.g., a fusion protein)) or the components for base editing (e.g., a CRISPR-Cas effector domain fused to a peptide tag or an affinity polypeptide, a deaminase domain fused to a peptide tag or an affinity polypeptide, and/or a UGI fused to a peptide tag or an affinity polypeptide), may be comprised on the same or on a separate expression cassette or vector from that comprising the one or more guide nucleic acids. When the nucleic acid construct encoding a base editor or the components for base editing is/are comprised on separate expression cassette(s) or vector(s) from that comprising the guide nucleic acid, a target nucleic acid may be contacted with (e.g., provided with) the expression cassette(s) or vector(s) encoding the base editor or components for base editing in any order from one another and the guide nucleic acid, e.g., prior to, concurrently with, or after the expression cassette comprising the guide nucleic acid is provided (e.g., contacted with the target nucleic acid).

[0140] Fusion proteins of the invention may comprise a sequence-specific DNA binding domain, a CRISPR-Cas effector protein, and/or a deaminase fused to a peptide tag or an affinity polypeptide that interact with the peptide tag, as known in the art, for use in recruiting the deaminase to the target nucleic acid. Methods of recruiting may also comprise a guide nucleic acids linked to an RNA recruiting motif and a deaminase fused to an affinity polypeptide capable of interacting with the RNA recruiting motif, thereby recruiting the deaminase to the target nucleic acid. Alternatively, chemical interactions may be used to recruit a polypeptide (e.g., a deaminase) to a target nucleic acid.

[0141] A peptide tag (e.g., epitope) useful with this invention may include, but is not limited to, a GCN4 peptide tag (e.g., Sun-Tag), a c-Myc affinity tag, an HA affinity tag, a His affinity tag, an S affinity tag, a methionine-His affinity tag, an RGD-His affinity tag, a FLAG octapeptide, a strep tag or strep tag II, a V5 tag, and/or a VSV-G epitope. Any epitope that may be linked to a polypeptide and for which there is a corresponding affinity polypeptide that may be linked to another polypeptide may be used with this invention as a peptide tag. In some embodiments, a peptide tag may comprise 1 or 2 or more copies of a peptide tag (e.g., repeat unit, multimerized epitope (e.g., tandem repeats)) (e.g., 1, 2,

3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 or more repeat units. In some embodiments, an affinity polypeptide that interacts with/binds to a peptide tag may be an antibody. In some embodiments, the antibody may be a scFv antibody. In some embodiments, an affinity polypeptide that binds to a peptide tag may be synthetic (e.g., evolved for affinity interaction) including, but not limited to, an affibody, an anticalin, a monobody and/or a DARPIn (see, e.g., Sha et al., *Protein Sci.* 26(5): 910-924 (2017)); Gilbreth (*Curr Opin Struc Biol* 22(4):413-420 (2013)), U.S. Pat. No. 9,982,053, each of which are incorporated by reference in their entireties for the teachings relevant to affibodies, anticalins, monobodies and/or DARPins.

[0142] In some embodiments, a guide nucleic acid may be linked to an RNA recruiting motif, and a polypeptide to be recruited (e.g., a deaminase) may be fused to an affinity polypeptide that binds to the RNA recruiting motif, wherein the guide binds to the target nucleic acid and the RNA recruiting motif binds to the affinity polypeptide, thereby recruiting the polypeptide to the guide and contacting the target nucleic acid with the polypeptide (e.g., deaminase). In some embodiments, two or more polypeptides may be recruited to a guide nucleic acid, thereby contacting the target nucleic acid with two or more polypeptides (e.g., deaminases).

[0143] In some embodiments of the invention, a guide RNA may be linked to one or two or more RNA recruiting motifs (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more motifs; e.g., at least 10 to about 25 motifs), optionally wherein the two or more RNA recruiting motifs may be the same RNA recruiting motif or different RNA recruiting motifs. In some embodiments, an RNA recruiting motif and corresponding affinity polypeptide may include, but is not limited to, a telomerase Ku binding motif (e.g., Ku binding hairpin) and the corresponding affinity polypeptide Ku (e.g., Ku heterodimer), a telomerase Sm7 binding motif and the corresponding affinity polypeptide Sm7, an MS2 phage operator stem-loop and the corresponding affinity polypeptide MS2 Coat Protein (MCP), a PP7 phage operator stem-loop and the corresponding affinity polypeptide PP7 Coat Protein (PCP), an SFMu phage Com stem-loop and the corresponding affinity polypeptide Com RNA binding protein, a PUF binding site (PBS) and the affinity polypeptide *Pumilio/fem-3* mRNA binding factor (PUF), and/or a synthetic RNA-aptamer and the aptamer ligand as the corresponding affinity polypeptide. In some embodiments, the RNA recruiting motif and corresponding affinity polypeptide may be an MS2 phage operator stem-loop and the affinity polypeptide MS2 Coat Protein (MCP). In some embodiments, the RNA recruiting motif and corresponding affinity polypeptide may be a PUF binding site (PBS) and the affinity polypeptide *Pumilio/fem-3* mRNA binding factor (PUF). Exemplary RNA recruiting motifs and corresponding affinity polypeptides that may be useful with this invention can include, but are not limited to, SEQ ID NOs:101-111.

[0144] In some embodiments, the components for recruiting polypeptides and nucleic acids may include those that function through chemical interactions that may include, but are not limited to, rapamycin-inducible dimerization of FRB-FKBP; Biotin-streptavidin; SNAP tag; Halo tag; CLIP tag; DmrA-DmrC heterodimer induced by a compound; bifunctional ligand (e.g., fusion of two protein-binding chemicals together; e.g. dihydrofolate reductase (DHFR).

[0145] In some embodiments, the nucleic acid constructs, expression cassettes or vectors of the invention that are optimized for expression in a plant may be about 70% to 100% identical (e.g., about 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or 100%) to the nucleic acid constructs, expression cassettes or vectors comprising the same polynucleotide(s) but which have not been codon optimized for expression in a plant.

[0146] As described herein, a “peptide tag” may be employed to recruit one or more polypeptides. A peptide tag may be any polypeptide that is capable of being bound by a corresponding affinity polypeptide. A peptide tag may also be referred to as an “epitope” and when provided in multiple copies, a “multimerized epitope.” Example peptide tags can include, but are not limited to, a GCN4 peptide tag (e.g., Sun-Tag), a c-Myc affinity tag, an HA affinity tag, a His affinity tag, an S affinity tag, a methionine-His affinity tag, an RGD-His affinity tag, a FLAG octapeptide, a strep tag or strep tag II, a V5 tag, and/or a VSV-G epitope. In some embodiments, a peptide tag may also include phosphorylated tyrosines in specific sequence contexts recognized by SH2 domains, characteristic consensus sequences containing phosphoserines recognized by 14-3-3 proteins, proline rich peptide motifs recognized by SH3 domains, PDZ protein interaction domains or the PDZ signal sequences, and an AGO hook motif from plants. Peptide tags are disclosed in WO2018/136783 and U.S. Patent Application Publication No. 2017/0219596, which are incorporated by reference for their disclosures of peptide tags. Peptide tags that may be useful with this invention can include, but are not limited to, SEQ ID NO:112 and SEQ ID NO:113. An affinity polypeptide useful with peptide tags includes, but is not limited to, SEQ ID NO:114.

[0147] A peptide tag may comprise or be present in one copy or in 2 or more copies of the peptide tag (e.g., multimerized peptide tag or multimerized epitope) (e.g., about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 or more peptide tags). When multimerized, the peptide tags may be fused directly to one another or they may be linked to one another via one or more amino acids (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more amino acids, optionally about 3 to about 10, about 4 to about 10, about 5 to about 10, about 5 to about 15, or about 5 to about 20 amino acids, and the like, and any value or range therein). Thus, in some embodiments, a CRISPR-Cas effector protein of the invention may comprise a CRISPR-Cas effector protein domain fused to one peptide tag or to two or more peptide tags, optionally wherein the two or more peptide tags are fused to one another via one or more amino acid residues. In some embodiments, a peptide tag useful with the invention may be a single copy of a GCN4 peptide tag or epitope or may be a multimerized GCN4 epitope comprising about 2 to about 25 or more copies of the peptide tag (e.g., about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 or more copies of a GCN4 epitope or any range therein).

[0148] In some embodiments, a peptide tag may be fused to a CRISPR-Cas polypeptide or domain. In some embodiments, a peptide tag may be fused or linked to the C-terminus of a CRISPR-Cas effector protein to form a CRISPR-Cas fusion protein. In some embodiments, a peptide tag may

be fused or linked to the N-terminus of a CRISPR-Cas effector protein to form a CRISPR-Cas fusion protein. In some embodiments, a peptide tag may be fused within a CRISPR-Cas effector protein (e.g., a peptide tag may be in a loop region of a CRISPR-Cas effector protein). In some embodiments, peptide tag may be fused to a cytosine deaminase and/or to an adenine deaminase.

[0149] An “affinity polypeptide” (e.g., “recruiting polypeptide”) refers to any polypeptide that is capable of binding to its corresponding peptide tag, peptide tag, or RNA recruiting motif. An affinity polypeptide for a peptide tag may be, for example, an antibody and/or a single chain antibody that specifically binds the peptide tag, respectively. In some embodiments, an antibody for a peptide tag may be, but is not limited to, an scFv antibody. In some embodiments, an affinity polypeptide may be fused or linked to the N-terminus of a deaminase (e.g., a cytosine deaminase or an adenine deaminase). In some embodiments, the affinity polypeptide is stable under the reducing conditions of a cell or cellular extract.

[0150] The nucleic acid constructs of the invention and/or guide nucleic acids may be comprised in one or more expression cassettes as described herein. In some embodiments, a nucleic acid construct of the invention may be comprised in the same or in a separate expression cassette or vector from that comprising a guide nucleic acid and/or a recruiting guide nucleic acid.

[0151] When used in combination with guide nucleic acids and recruiting guide nucleic acids, the nucleic acid constructs of the invention (and expression cassettes and vectors comprising the same) may be used to modify a target nucleic acid and/or its expression. A target nucleic acid may be contacted with a nucleic acid construct of the invention and/or expression cassettes and/or vectors comprising the same prior to, concurrently with or after contacting the target nucleic acid with the guide nucleic acid/recruiting guide nucleic acid (and/or expression cassettes and vectors comprising the same).

[0152] The term “seed region” refers to the critical portion of a crRNA’s or guide RNA’s guide sequence that is most susceptible to mismatches with their targets. In some embodiments, a single mismatch in the seed region of a crRNA/gRNA can render a CRISPR complex inactive at that binding site. In some embodiments, the seed regions for Cas9 endonucleases are located along the last ~12 nts of the 3’ portion of the guide sequence, which correspond (hybridize) to the portion of the protospacer target sequence that is adjacent to the PAM. In some embodiments, the seed regions for Cpf1 endonucleases are located along the first ~5 nts of the 5’ portion of the guide sequence, which correspond (hybridize) to the portion of the protospacer target sequence adjacent to the PAM.

[0153] The term “modified” refers to a substance or compound (e.g., a cell, a polynucleotide sequence, and/or a polypeptide sequence) that has been altered or changed as compared to the corresponding unmodified substance or compound. For example, a modified polynucleotide can be a nucleic acid that has been edited (e.g., mutated such as by substituting and/or deleting a nucleic acid) and/or changed compared to its unmodified or native sequence and/or structure.

[0154] An “edited polynucleotide” as used herein refers to a polynucleotide that has been edited (e.g., mutated such as by substituting and/or deleting a nucleic acid) and/or

changed compared to its unmodified or native sequence and/or structure. The term “gene edited plant, part or cell” as used herein refers to a plant, part or cell that comprises one or more endogenous genes that are edited by an editing system. An editing system of the present disclosure may comprise a targeting element and/or an editing element. The targeting element may be capable of recognizing and/or may be configured to recognize a target genomic sequence. The editing element may be capable of modifying and/or may be configured to modify a target nucleic acid, e.g., by substitution or insertion of one or more nucleotides, deletion of one or more nucleotides, alteration of a nucleotide sequence to include a regulatory sequence, insertion of a transgene at a safe harbor genomic site or other specific location in the genome, or any combination thereof. The targeting element and the editing element can be on the same nucleic acid molecule or different nucleic acid molecules. In some embodiments, the editing element is capable of precise genome editing by insertion and/or deletion of at least one nucleotide using a CRISPR/Cas system. In some embodiments, the editing element is capable of precise genome editing by substitution of a single nucleotide using a base editor, such as cytosine base editor (CBE) and/or adenine base editor (ABE), which is directly or indirectly fused to a CRISPR-associated effector protein.

[0155] The terms “transgene” or “transgenic” as used herein refer to at least one nucleic acid sequence that is taken from the genome of one organism, or produced synthetically, and which is then introduced into a host cell or organism or tissue of interest and which is subsequently integrated into the host’s genome by means of “stable” transformation or transfection approaches. In contrast, the term “transient” transformation or transfection or introduction refers to a way of introducing molecular tools including at least one nucleic acid (DNA, RNA, single-stranded or double-stranded or a mixture thereof) and/or at least one amino acid sequence, optionally comprising suitable chemical or biological agents, to achieve a transfer into at least one compartment of interest of a cell, including, but not restricted to, the cytoplasm, an organelle, including the nucleus, a mitochondrion, a vacuole, a chloroplast, or into a membrane, resulting in transcription and/or translation and/or association and/or activity of the at least one molecule introduced without achieving a stable integration or incorporation and thus inheritance of the respective at least one molecule introduced into the genome of a cell. The terms “transgene-free” refers to a condition in which a transgene is not present or found in the genome of a host cell or tissue or organism of interest.

[0156] As used herein, the term “tissue culture” indicates a composition comprising isolated cells of the same or a different type or a collection of such cells organized into parts of a plant. Exemplary types of tissue cultures are protoplasts, calli, plant clumps, and plant cells that can generate tissue culture that are intact in plants or parts of plants, such as embryos, pollen, flowers, seeds, leaves, stems, roots, root tips, anthers, pistils, meristematic cells, axillary buds, ovaries, seed coat, endosperm, hypocotyls, cotyledons and the like. The term “plant organ” refers to plant tissue or a group of tissues that constitute a morphologically and functionally distinct part of a plant. “Progeny” comprises any subsequent generation of a plant.

[0157] General methods in molecular and cellular biochemistry can be found in such standard textbooks as

Molecular Cloning: A Laboratory Manual, 3rd Ed. (Sambrook et al., HarBor Laboratory Press 2001); Short Protocols in Molecular Biology, 4th Ed. (Ausubel et al. eds., John Wiley & Sons 1999); Protein Methods (Bollag et al., John Wiley & Sons 1996); Nonviral Vectors for Gene Therapy (Wagner et al. eds., Academic Press 1999); Viral Vectors (Kaplift & Loewy eds., Academic Press 1995); Immunology Methods Manual (I. Lefkovits ed., Academic Press 1997); and Cell and Tissue Culture: Laboratory Procedures in Biotechnology (Doyle & Griffiths, John Wiley & Sons 1998), the disclosures of which are incorporated herein by reference.

[0158] As mentioned above, segregating to eliminate transgenes in vegetative propagated crops such as caneberries and cherries is not a preferred practice after a target gene is edited. Particularly, the heterozygotic nature of these crops requires several years or decades in the case of woody crops such as cherries, to breed newly edited traits to a desired elite cultivar for commercial production using traditional breeding methods. The present disclosure provides a solution to obtain transgene-free edited plants such as by transiently expressing the gene editing machinery (optionally, a Cas9 endonuclease and gRNA) in the plant cells and also by co-expressing at least one morphogenic regulator to regenerate edited cells without the aid of selection. The present disclosure also teaches methods for culturing plant tissues for transformation and regeneration as well as identification of the gene-edited event in the transformed plant, plant tissue, and plant cell.

[0159] A method of the present invention may comprise modifying a target nucleic acid using a nucleic acid construct of the invention, and/or an expression cassette and/or vector comprising the same. A method of the present invention may be carried out in an in vivo system (e.g., in a cell or in an organism) or in an in vitro system (e.g., cell free). A method, composition, and/or system of the present invention may generate and/or provide allelic diversity, optionally in a semi-random way. In some embodiments, a method of the present invention comprises determining a desired or preferred phenotype using and/or based on the modified target nucleic acid. A method of the present invention may provide one or more modified target nucleic acid(s), and the one or more modified target nucleic acid(s) may be analyzed for a desired or preferred phenotype.

[0160] According to some embodiments, a morphogenic regulator is provided. A morphogenic regulator may be present in and/or used in a composition, system, and/or method of the present invention. A “morphogenic regulator” as used herein refers to a nucleic acid sequence, gene, or polypeptide that can cause and/or stimulate morphogenesis in a cell in which it is present and expressed. A morphogenic regulator of the present invention may be used to regenerate a plant from a transformed plant cell. In some embodiments, ectopic overexpression of a morphogenic regulator involved in either embryo and/or meristem development stimulates growth of transgenic plants. Due to pleiotropic deleterious effect of some morphogenic regulators under a condition of constitutive expression, the present disclosure teaches the initial stimulating of morphogenic regulators while later restricting or eliminating their expression in the plant. In some embodiments, methods of controlling ectopic overexpression include, but are not limited to the use of transient expression, inducible promoters, tissue-specific promoters, and excision of the morphogenic regulators.

[0161] In some embodiments, ectopic overexpression of morphogenic regulators induce morphogenic growth response to enhance and/or improve transformation efficiencies, which may allow for transformation of numerous recalcitrant crops and/or successful regeneration of transformed plants. In some embodiments, transient expression of morphogenic regulators enhances and/or improves transformation efficiencies. In some embodiments, transient expression of a morphogenic regulator as described herein provides an increase and/or improvement in plant transformation efficiency and/or rate. In other embodiments, transient expression of morphogenic regulators provides an increase and/or improvement in a plant regeneration process and/or rate. In some embodiments, introduction of a polypeptide comprising a morphogenic regulator onto and/or into a plant cell provides an increase and/or improvement in plant transformation efficiency and/or rate and/or a plant regeneration process and/or rate. In some embodiments, the use of morphogenic regulators can solve current issues with plant genome modification and regeneration of genetically-modified plants and/or gene-edited plants. In some embodiments, plant transformation and regeneration improvements can be distinguished by their impact on either improving transformation efficiencies, or improving the regeneration process to recover transgenic plants.

[0162] Exemplary morphogenic regulators include, but are not limited to, a Wuschel (WUS) gene (e.g., from blackberry and/or cherry), Baby boom (BBM, e.g., from blackberry and/or cherry), Knotted1 (KN1) gene (e.g., from maize, blackberry, and/or cherry), isopentenyl transferase (ipt) gene (e.g., from *Agrobacterium tumefaciens*, blackberry, and/or cherry), CLAVATA3 (CLV3) mutant, miR156 (e.g., from citrus), AtGRF5 (e.g., from *Arabidopsis thaliana*), NTT (e.g., from *Arabidopsis thaliana*, blackberry, and/or cherry), HDZipII (e.g., from *Arabidopsis thaliana*, blackberry, and/or cherry), and orthologs or polypeptides thereof. Further exemplary morphogenic regulators include, but are not limited to those described in U.S. Pat. No. 7,256,322; *Plant Cell* 14:1737-1749 (2002); *Plant Cell Physiol.* 41(5) 583-590 (2000); *Plant Physiol.* 161(3):1076-1085 (2013); *J Exp Bot.* 69(12):2979-2993 (2018); *Plant Cell* 27(2):349-360 (2015); and *Plant Physiol.* 167(3):817-832 (2015).

[0163] In some embodiments, homologs and orthologs of WUS, BBM, KN1, IPT, CLV3, miR156, GRF5, NTT, and HDZipII genes are identified and used for improving plant transformation and/regeneration in a plant of interest.

[0164] Further exemplary morphogenic regulators include, but are not limited to, those having a nucleotide sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the nucleotide sequence of any one of SEQ ID NOs:1-18 or SEQ ID NOs:28-49. In some embodiments, a nucleotide sequence of the present invention may encode an amino acid sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the polypeptide sequence of any one of SEQ ID NOs:19-27 or SEQ ID NOs:50-60. Exemplary morphogenic regulators may have an amino acid sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the polypeptide sequence of any one of SEQ ID NOs:19-27 or SEQ ID NOs:50-60. A polynucleotide comprising a morphogenic regulator may be operably associated with a heterologous promoter. Compositions, systems, and/

or methods of the present invention may provide a cell, plant part, plant, expression cassette, and/or vector comprising a polynucleotide comprising a morphogenic regulator as described herein and/or a polypeptide comprising a morphogenic regulator as described herein. In some embodiments, the cell, plant part, plant, expression cassette, and/or vector also includes an editing system (e.g., a polynucleotide encoding gene editing machinery or a gene editing complex (e.g., a ribonucleoprotein)), and/or a polynucleotide encoding a selectable marker.

[0165] The catabolic enzyme isopentenyl transferase (IPT) directs the synthesis of cytokinins and plays a major role in controlling cytokinin levels in plant tissues. Multiple routes have been proposed for cytokinin biosynthesis. Transfer RNA degradation has been suggested to be a source of cytokinin, because some tRNA molecules contain an isopentenyladenosine (iPA) residue at the site adjacent to the anticodon (Swaminathan, et al., (1977) *Biochemistry* 16:1355-1360). The modification is catalyzed by tRNA isopentenyl transferase (tRNA IPT; EC 2.5.1.8), which has been identified in various organisms such as *Escherichia coli*, *Saccharomyces cerevisiae*, *Lactobacillus acidophilus*, *Homo sapiens*, and *Zea mays* (Bartz, et al., (1972) *Biochemie* 54:31-39; Kline, et al., (1969) *Biochemistry* 8:4361-4371; Holtz, et al., (1975) *Hoppe-Seyler's Z. Physiol. Chem.* 356:1459-1464; Golovko, et al., (2000) *Gene* 258:85-93; and, Holtz, et al., (1979) *Hoppe-Seyler's Z. Physiol. Chem.* 359:89-101). However, this pathway is not considered to be the main route for cytokinin synthesis (Chen, et al., (1997) *Physiol. Plant* 101:665-673 and McGraw, et al., (1995) *Plant Hormones, Physiology, Biochemistry and Molecular Biology*. Ed. Davies, 98-117, Kluwer Academic Publishers, Dordrecht).

[0166] The current knowledge of cytokinin biosynthesis in plants is largely deduced from studies on a possible analogous system in *Agrobacterium tumefaciens*. Cells of *A. tumefaciens* are able to infect certain plant species by inducing tumor formation in host plant tissues (Van Montagu, et al., (1982) *Curr Top Microbiol Immunol* 96:237-254; Hansen, et al., (1999). *Curr Top Microbiol Immunol* 240:21-57). To do so, the *A. tumefaciens* cells synthesize and secrete cytokinins which mediate the transformation of normal host plant tissues into tumors or calli. This process is facilitated by the *A. tumefaciens* tumor-inducing plasmid which contains genes encoding the necessary enzyme and regulators for cytokinin biosynthesis. Biochemical and genetic studies revealed that Gene 4 of the tumor-inducing plasmid encodes an isopentenyl transferase (IPT), which converts AMP and DMAPP into isopentenyladenosine-5'-monophosphate (iPMP), the active form of cytokinins (Akaiyoshi, et al., (1984) *Proc. Natl. Acad. Sci. USA* 81:5994-5998). Overexpression of the *Agrobacterium* ipt gene in a variety of transgenic plants has been shown to cause an increased level of cytokinins and elicit typical cytokinin responses in the host plant (Hansen, et al., (1999) *Curr Top Microbiol Immunol* 240:21-57). Therefore, it has been postulated that plant cells use machinery similar to that of *A. tumefaciens* cells for cytokinin biosynthesis. *Arabidopsis* IPT homologs have recently been identified in *Arabidopsis* and *Petunia* (Takei, et al., (2001) *J. Biol. Chem.* 276:26405-26410 and Kakimoto, (2001) *Plant Cell Physiol.* 42:677-685). Overexpression of the *Arabidopsis* IPT homologs in plants elevated cytokinin levels and elicited typical cyto-

nin responses in planta and under tissue culture conditions (Kakimoto, (2001) *Plant Cell Physiol.* 42:677-685).

[0167] In view of the influence of cytokinins on a wide variety of plant developmental processes, including root architecture, shoot and leaf development, and seed set, the ability to manipulate cytokinin levels in higher plant cells, and thereby drastically affect plant growth and productivity, offers significant commercial value (Mok, et al., (1994) *Cytokinins. Chemistry, Action and Function*. CRC Press, Boca Raton, Fla., pp. 155-166).

[0168] In some embodiments, the morphogenic regulator is an *ipt* gene encoding isopentenyl transferase. Isopentenyl transferase (*ipt*) gene is a cytokinin biosynthetic gene isolated from *Agrobacterium tumefaciens*.

[0169] In some embodiments, overexpression of a morphogenic regulator (e.g., an *ipt* gene) in transgenic plant cells induces shoot formation in media without additional plant growth regulators. In some embodiments, non-transgenic plants may be regenerated from the hormone-free media and they may be morphologically normal. In some embodiments, these non-transgenic plants are derived from cells where a morphogenic regulator was transiently expressed. In some embodiments, these non-transgenic plants are from the cells adjacent to the transformed cells which redistributed the cytokinin to the adjacent wild type cells.

[0170] Compositions, systems, and/or methods of the present invention may produce non-transgenic plants that are derived from cells under a condition of in which a morphogenic regulator is transiently expressed. In some embodiments, an expression cassette comprising gene editing machinery (Cas variants, Cpf1, gRNA targeting PDS or other genes) is transformed or co-transformed with another expression cassette comprising a morphogenic regulator. In some embodiments, the resultant transgenic (with an edited polynucleotide, but without expression of the morphogenic regulator) and non-transgenic (with an unedited polynucleotide, and with expression of the morphogenic regulator) plants can be analyzed for transgene integration and gene editing efficiency.

[0171] A method of the present invention may comprise introducing a polynucleotide or a polypeptide comprising a heterologous morphogenic regulator into a plant cell, thereby transforming the plant cell. A “heterologous morphogenic regulator” as used herein refers to a morphogenic regulator as described herein that is heterologous to the plant cell in which the morphogenic regulator is being or has been introduced, or to a morphogenic regulator as described herein that is present at a different protein concentration than it would normally be present in the plant cell to which the morphogenic regulator is being or has been introduced. Compared to a control, the method may improve or increase regeneration frequency in a group of plant cells in which at least one of the plant cells is transformed with the polynucleotide or polypeptide comprising a heterologous morphogenic regulator. A “group of plant cells” as used herein refers to two more plant cells. In some embodiments, regeneration frequency may be improved or increased in a group of plant cells that are not transformed with a polynucleotide or polypeptide comprising a heterologous morphogenic regulator, but are in proximity to a cell comprising the polynucleotide or polypeptide comprising a heterologous morphogenic regulator. The method may include introducing an editing system into the same plant cell or a

different plant cell as the heterologous morphogenic regulator. The step of introducing the polynucleotide or polypeptide comprising the heterologous morphogenic regulator and the step of introducing the editing system may be performed concurrently or sequentially in any order. A plant cell may be stably transformed with a polynucleotide comprising a heterologous morphogenic regulator and/or may be stably transformed with an editing system (e.g., a polynucleotide encoding a gene editing machinery or a gene editing complex such as a ribonucleoprotein). A plant cell may be transiently transformed with a polynucleotide or polypeptide comprising a heterologous morphogenic regulator and/or may be stably transformed with an editing system (e.g., a polynucleotide encoding a gene editing machinery or a gene editing complex).

[0172] A method of the present invention may not introduce (i.e., may be devoid of introducing) a polynucleotide encoding a selectable marker. Accordingly, a plant cell comprising the polynucleotide or polypeptide comprising a heterologous morphogenic regulator and/or editing system may not comprise a polynucleotide encoding a selectable marker. In some embodiments, a method of the present invention comprises introducing a polynucleotide encoding a selectable marker into the same plant cell or a different plant cell as the heterologous morphogenic regulator and/or editing system. A selectable marker may be stably or transiently transformed in a cell and/or method of the present invention.

[0173] A single polynucleotide may comprise a polynucleotide comprising a heterologous morphogenic regulator, a polynucleotide encoding an editing system, and/or a polynucleotide encoding a selectable marker, or two or more polynucleotides may include one or more of a polynucleotide comprising a heterologous morphogenic regulator, a polynucleotide encoding an editing system, and/or a polynucleotide encoding a selectable marker. In some embodiments, a first polynucleotide comprises a polynucleotide comprising a heterologous morphogenic regulator and a second polynucleotide comprises a polynucleotide encoding editing system.

[0174] A method of the present invention may stimulate embryogenesis and/or organogenesis in a plant cell. In some embodiments, a method of the present invention stimulates embryogenesis and/or organogenesis in a plant cell comprising a polynucleotide or polypeptide comprising a heterologous morphogenic regulator and optionally an editing system and/or selectable marker. In some embodiments, a plant cell and/or method of the present invention stimulates embryogenesis and/or organogenesis in a plant cell that does not comprise (i.e., is devoid of) a polynucleotide or polypeptide comprising a heterologous morphogenic regulator or does not have the heterologous morphogenic regulator at a same concentration (e.g., protein concentration) in the plant cell. For example, a method of the present invention may introduce a polynucleotide comprising a heterologous morphogenic regulator into a plant cell and the plant cell comprising the polynucleotide comprising a heterologous morphogenic regulator may stably or transiently express the polynucleotide comprising a heterologous morphogenic regulator. In stably or transiently expressing the polynucleotide comprising a heterologous morphogenic regulator, the plant cell may stimulate embryogenesis and/or organogenesis in a plant cell that does not comprise (i.e., is devoid of) a polynucleotide comprising a heterologous morphogenic

regulator. The plant cell devoid of the polynucleotide comprising a heterologous morphogenic regulator may be a neighboring plant cell.

[0175] A “neighboring plant cell” as used herein refers to the location of a plant cell relative to another plant cell (i.e., the reference plant cell), and a neighboring plant cell is one that is in physical contact with, adjacent to, and/or in chemical communication with (e.g., receives cytokines from) the reference plant cell. The neighboring plant cell may be a plant cell that is from a different plant species or genotype than that of the transformed cell comprising the heterologous morphogenic regulator as described herein (e.g., the reference plant cell). In some embodiments, a neighboring plant cell is a plant cell in which a plasmodesmata (e.g., a primary and/or secondary plasmodesmata) connects the plant cell to the reference plant cell and/or in which a plasmodesmata traverses or passes through the cell wall of both the reference plant cell and neighboring plant cell. In some embodiments, a neighboring plant cell is the plant cell that is immediately adjacent to the reference plant cell and middle lamella contacts or joins the cell walls of the two plant cells. In some embodiments, a neighboring plant cell is a plant cell in which an apoplast is formed by a portion of the cell wall of the reference plant cell and by a portion of the cell wall of the neighboring plant cell. In some embodiments, a neighboring plant cell is a plant cell in which 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more plant cells are between the cell wall of the neighboring plant cell and the cell wall of the reference plant cell. In some embodiments, a neighboring plant cell is one that is in the same explant as the reference plant cell and/or all cells in an explant are considered to be a neighboring plant cell. A neighboring plant cell may be devoid of a polynucleotide or a polypeptide comprising a heterologous morphogenic regulator, devoid of an editing system, and/or devoid of a selectable marker. In some embodiments, a first plant cell is transiently transformed with a polynucleotide or a polypeptide comprising a heterologous morphogenic regulator and the first plant cell stimulates a neighboring plant cell that is transiently transformed with an editing system to multiply and/or regenerate.

[0176] According to some embodiments, a composition, system, and/or method of the present invention causes or induces one or more plant cell(s) to multiply to provide one or more transgene-free plant cell(s), optionally using a plant cell in which a polynucleotide or a polypeptide comprising a heterologous morphogenic regulator has been introduced. The one or more transgene-free plant cell(s) may be undifferentiated plant cells or tissue or may be a callus. The one or more transgene-free plant cell(s) may be cultured explants. The one or more transgene-free cell(s) may form and/or may be grown to form one or more organ(s) and/or plant part(s). In some embodiments, a composition, system, and/or method of the present invention regenerates a transgene-free plant part or transgene-free plant, optionally using a plant cell in which a polynucleotide or a polypeptide comprising a heterologous morphogenic regulator has been introduced. In some embodiments, a transgene-free plant cell, a transgene-free plant part, and/or transgene-free plant is produced (e.g., regenerated) from the plant cell in which a polynucleotide or a polypeptide comprising a heterologous morphogenic regulator has been introduced. In some embodiments, the plant cell in which a polynucleotide or polypeptide comprising a heterologous morphogenic regu-

lator has been introduced stimulates and/or initiates one or more neighboring plant cell(s) to multiply and/or regenerate to provide a transgene-free plant cell, a transgene-free plant part, or transgene-free plant. A method of the present invention may include, responsive to a step of introducing a polynucleotide or a polypeptide comprising a heterologous morphogenic regulator into a plant cell and optionally responsive to introducing an editing system into the same plant cell or a different plant cell (e.g., a neighboring plant cell), producing a plant cell, plant part and/or plant that is non-transgenic. A transgene-free plant cell, plant part and/or plant of the present invention may comprise an edited polynucleotide. The edited polynucleotide may be responsive to introducing an editing system into the same plant cell or a different plant cell (e.g., a neighboring plant cell) as the plant cell in which a polynucleotide or a polypeptide comprising a heterologous morphogenic regulator is introduced.

[0177] A composition, system, and/or method of the present invention may induce shoot formation, optionally using a plant cell in which a polynucleotide or a polypeptide comprising a heterologous morphogenic regulator and/or editing system has been introduced. In some embodiments, the method comprises expressing the heterologous morphogenic regulator in the plant cell and inducing shoot formation from the plant cell or a neighboring plant cell. The shoot formation may comprise apical dominance and/or roots.

[0178] A method of the present invention may include growing (e.g., culturing) a plant cell in the presence of and/or in contact with (e.g., on) media devoid of plant growth regulators and/or hormones (e.g., hormone-free media). The plant cell may comprise a polynucleotide or a polypeptide comprising a heterologous morphogenic regulator, editing system, and/or a selectable marker. In some embodiments, growing the plant cell comprises growing the plant cell in the presence of, in contact with, and/or on media devoid of plant growth regulators and/or hormones and producing a plant part and/or plant from the plant cell.

[0179] In some embodiments, a method of the present invention comprises transiently expressing an editing system in a plant cell and regenerating plant cells comprising an edited polynucleotide, optionally without the aid of selection (e.g., without using or introducing a selection marker). A polynucleotide encoding the editing system may be introduced into the plant cell or a neighboring plant cell, and/or the editing system may be introduced into the plant cell or a neighboring plant cell as an assembled ribonucleoprotein complex. In some embodiments, the editing system comprises a CRISPR-Cas effector protein (e.g., Cas9/Cas12a) and a guide nucleic acid.

[0180] In some embodiments, a method of the present invention comprises introducing (e.g., delivering) a polynucleotide or a polypeptide comprising a heterologous morphogenic regulator and a polynucleotide encoding a selectable marker to a plant cell, and regenerating a stably transformed plant under herbicide and/or antibiotic selection. In some embodiments, the method comprises introducing an editing system. In some embodiments, an editing system is not introduced.

[0181] In some embodiments, a method of the present invention comprises introducing (e.g., delivering) a polynucleotide or a polypeptide comprising a heterologous morphogenic regulator to a plant cell and does not include introducing a selectable marker to the plant cell. The method further includes regenerating a stably transformed plant

without selection. In some embodiments, the method comprises introducing an editing system. In some embodiments, an editing system is not introduced.

[0182] In some embodiments, a composition, system and/or method of the present invention provides a genetic trigger for embryogenesis, organogenesis, and/or regeneration. In some embodiments, a composition, system and/or method of the present invention stimulates or causes transformed cells to multiply and/or regenerate faster and/or more efficiently than non-transformed cells. This may allow for transformed shoots to be more efficiently found among the regenerated shoots even without a selection system to inhibit the regeneration of untransformed cells.

[0183] In some embodiments, a method of the present invention comprises introducing (e.g., delivering) a polynucleotide or a polypeptide comprising a heterologous morphogenic regulator to a plant cell, and inducing a neighboring plant cell to multiply and/or regenerate. Certain types of morphogenic regulators (e.g., wuschel or WUS) organize embryo development on a concentration gradient from high to low. For this reason, cells that are transiently or stably transformed can induce their neighboring plant cells to organize an embryo and regenerate. Accordingly, a polynucleotide or a polypeptide comprising a heterologous morphogenic regulator may be introduced separately from another transgene and/or separately from an editing system. In some embodiments, the cells that receive the polynucleotide or polypeptide comprising a heterologous morphogenic regulator cause or induce neighboring plant cells, some of which have received the separate transgene and/or editing system, to multiply and/or regenerate. This can be with or without editing tools (e.g., an editing system) and/or with or without a selectable marker in the neighboring plant cells.

[0184] In some embodiments, a method of the present invention directly expresses transgenes encoding polynucleotides that initiate and/or define embryo identity. A composition, system, and/or method of the present invention may reduce or eliminate the dependence of a plant cell on media growth regulators and/or hormone supplements for regeneration. Media hormone supplementation enables many cells in a tissue, whether transformed or not, the potential to regenerate. However, an advantage of a composition, system, and/or method of the present invention can be that by introducing a polynucleotide or a polypeptide comprising a heterologous morphogenic regulator into a plant cell, the potential to regenerate is now tied to the transformation identity itself, representing a powerful approach to selectively encourage transformed cells to regenerate preferentially over untransformed cells. A polypeptide, polynucleotide, composition, system, and/or method of the present invention may increase or improve regeneration efficiency by about 2-, 4-, 6-, 8-, 10-, 20-, 30-, 40-, 50-, 60-, 70-, 80-, 90-100-, 200-, 300-, 400-, 500-, 600-, 700-, 800-, 900-, or 1000-fold compared to a control (e.g., a composition, system, cell (e.g., plant cell), and/or method that is devoid of a polypeptide and/or polynucleotide of the present invention).

[0185] A composition, system and/or method of the present invention may provide and/or produce a plant part and/or plant such as responsive to a step of introducing a polynucleotide or a polypeptide comprising a heterologous morphogenic regulator into a plant cell and optionally responsive to introducing an editing system into the same plant cell or a different plant cell (e.g., a neighboring plant cell) and

responsive to growing plant cells. The plant part and/or plant may be morphologically normal compared to a control plant part and/or plant (e.g., compared to a non-transformed plant part or plant). Methods for determining morphological structures are known to those of skill in the art.

[0186] A composition, system and/or method of the present invention may comprise and/or introduce a morphogenic regulator as described herein. In some embodiments, the heterologous morphogenic regulator is a Knotted 1 (KN1), Baby Boom (BBM), Wuschel (WUS) or Wuschel-related homeobox (WOX) gene, or an ortholog thereof. In some embodiments, the morphogenic regulator is ipt or an ortholog thereof. In some embodiments, the morphogenic regulator is kn1 or an orthology thereof, optionally kn1 homeobox gene of maize. The maize transcription factor KN1 functions in plant meristems, self-renewing structures consisting of stem cells and their immediate daughters, which is associated with cytokinins accumulation through the positive regulation of cytokinin synthesis genes. In some embodiments, kn1 gene can be assessed to see if its transient expression is able to enhance plant regeneration and consequently increase gene editing occurrences. In some embodiments, the morphogenic regulator is WUS. In some embodiments, the morphogenic regulator is BBM. In some embodiments, the morphogenic regulator is CLV3. In some embodiments, the morphogenic regulator is miR156. In some embodiments, the morphogenic regulator is GRF5. In some embodiments, the morphogenic regulator is NTT. In some embodiments, the morphogenic regulator is HDZipII.

[0187] In some embodiments, the polynucleotide comprising the heterologous morphogenic regulator has at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the nucleotide sequence of any one of SEQ ID NOs:1-18 or SEQ ID NOs:28-49, optionally wherein the polynucleotide comprising the heterologous morphogenic regulator comprises the nucleotide sequence of any one of SEQ ID NOs:1-18 or SEQ ID NOs:28-49. In some embodiments, the polynucleotide comprising the heterologous morphogenic regulator has at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the nucleotide sequence of any one of SEQ ID NOs:1-18, optionally wherein the polynucleotide comprising the heterologous morphogenic regulator comprises the nucleotide sequence of any one of SEQ ID NOs:1-18. A polynucleotide comprising the heterologous morphogenic regulator may be operably associated with a promoter (e.g., a heterologous promoter). In some embodiments, the polynucleotide comprising the heterologous morphogenic regulator encodes an amino acid sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the polypeptide sequence of any one of SEQ ID NOs:19-27 or SEQ ID NOs:50-60, optionally wherein the polynucleotide comprising the heterologous morphogenic regulator encodes the polypeptide sequence of any one of SEQ ID NOs:19-27 or SEQ ID NOs:50-60. In some embodiments, the polynucleotide comprising the heterologous morphogenic regulator encodes an amino acid sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the polypeptide sequence of any one of SEQ ID NOs:19-27, optionally wherein the polynucleotide comprising the heterologous morphogenic regulator encodes the polypeptide sequence of any one of SEQ ID NOs:19-27. In some embodiments, the polypeptide comprising the hetero-

ologous morphogenic regulator has an amino acid sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the polypeptide sequence of any one of SEQ ID NOs:19-27 or SEQ ID NOs:50-60, optionally wherein the polypeptide comprising the heterologous morphogenic regulator has the polypeptide sequence of any one of SEQ ID NOs:19-27 or SEQ ID NOs:50-60. In some embodiments, the polypeptide comprising the heterologous morphogenic regulator has an amino acid sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the polypeptide sequence of any one of SEQ ID NOs:19-27, optionally wherein the polypeptide comprising the heterologous morphogenic regulator has the polypeptide sequence of any one of SEQ ID NOs:19-27. In some embodiments, the present disclosure provides a delivery of one or more components of an editing system directly to the plant cell. This is of interest, inter alia, for the generation of non-transgenic plants. In some embodiments, one or more of the components of the editing system is prepared outside the plant or plant cell and delivered to the cell. In some embodiments, the editing system is prepared in vitro prior to introduction to the plant cell. A base-editing fusion protein may be prepared by various methods known by one of skill in the art and include recombinant production. After expression, the CRISPR-Cas effector protein may be isolated, refolded if needed, purified and optionally treated to remove any purification tags if a tag such as a His-tag is present in the fusion protein. Once crude, partially purified, or more completely purified CRISPR-Cas effector protein is obtained, the protein may be introduced to the plant cell.

[0188] In some embodiments, the CRISPR-Cas effector protein is mixed with a guide nucleic acid that targets a target nucleic acid to form a pre-assembled ribonucleoprotein.

[0189] The individual components or pre-assembled ribonucleoprotein can be introduced into the plant cell via electroporation, by bombardment with targeted gene-editing system coated particles, by chemical transfection and/or by some other means of transport across a cell membrane. For instance, transfection of a plant protoplast with a pre-assembled CRISPR-Cas9 ribonucleoprotein has been demonstrated to ensure targeted modification of the plant genome (as described by Woo et al. *Nature Biotechnology*, 2015; 33(11):1162-1164).

[0190] In some embodiments, an editing system and/or one or more component(s) thereof may be introduced into a plant cell using nanoparticles. The editing system and/or components thereof, either as protein or nucleic acid or in a combination thereof, can be uploaded onto and/or packaged in nanoparticles and applied to the plants (such as for instance described in WO2008/042156 and US2013/0185823). In particular, the disclosure teaches nanoparticles uploaded with or packed with DNA molecule(s) encoding the CRISPR-Cas effector protein, DNA molecule(s) encoding cytosine deaminase (which may be fused to the CRISPR-Cas protein or a linker), and DNA molecules encoding the guide RNA and/or isolated guide RNA as described in WO2015/089419.

[0191] Further means of introducing one or more components of an editing system to a plant cell is by using cell penetrating peptides (CPP). Accordingly, some embodiments of the disclosure comprise compositions comprising a cell penetrating peptide linked to the gene-editing fusion

protein. In some embodiments, a sequence-specific nucleic acid binding domain (e.g., a CRISPR Cas effector protein) and/or guide nucleic acid is coupled to one or more CPPs to effectively transport them inside plant protoplasts. Ramakrishna (*Genome Res.* 2014 June; 24(6): 1020-7 for Cas9 in human cells). In some embodiments, a sequence-specific nucleic acid binding domain (e.g., a CRISPR Cas effector protein) and/or guide nucleic acid are encoded by one or more circular or non-circular DNA molecule(s) which are coupled to one or more CPPs for plant protoplast delivery. The plant protoplasts are then regenerated to plant cells and further to plants. CPPs are generally described as short peptides of fewer than 35 amino acids either derived from proteins or from chimeric sequences which are capable of transporting biomolecules across cell membrane in a receptor independent manner. CPP can be cationic peptides, peptides having hydrophobic sequences, amphipathic peptides, peptides having proline-rich and anti-microbial sequence, and chimeric or bipartite peptides (Pooga and Langel 2005). CPPs are able to penetrate biological membranes and as such trigger the movement of various biomolecules across cell membranes into the cytoplasm and to improve their intracellular routing, and hence facilitate interaction of the biomolecule with the target.

[0192] In some embodiments, the methods described herein are used to modify a endogenous nucleic acid and/or gene and/or to modify their expression without the permanent introduction of any foreign gene including those encoding the editing system components into the genome of the plant, so as to avoid the presence of foreign DNA in the genome of the plant. In some embodiments, this may be provided by transient expression of the editing system components. In some embodiments, one or more of the components are expressed on one or more viral vectors which produce sufficient a sequence-specific nucleic acid binding domain (e.g., a CRISPR Cas effector protein), and guide nucleic acid to consistently steadily ensure modification of a nucleic acid of interest according to a method described herein. In some embodiments, transient expression of the editing system components is ensured in plant protoplasts and thus not integrated into the genome. The limited window of expression can be sufficient to allow the editing system to ensure modification of a target nucleic acid as described herein.

[0193] In some embodiments, the different components of the editing system are introduced in the plant cell, protoplast or plant tissue either separately or in mixture, with the aid of delivering molecules such as nanoparticles or CPP molecules as described herein above.

[0194] The expression of the editing system components can induce targeted modification of the genome. The different strategies described herein can allow CRISPR-mediated targeted genome editing without requiring the introduction of the editing system components into the plant genome. Components which are transiently introduced into the plant cell are used for editing a target nucleic acid and become non-functional naturally as they are stably integrated into a host genome.

[0195] In some embodiments, plant cells which have an edited polynucleotide and that are produced or obtained by any of the methods described herein, can be cultured to regenerate a whole plant that possesses the transformed or modified genotype and thus the desired phenotype. Conventional regeneration techniques are well known to those

skilled in the art. Particular examples of such regeneration techniques rely on manipulation of certain phytohormones in a tissue culture growth medium, and typically relying on a biocide and/or herbicide marker which has been introduced together with the desired nucleotide sequences. In some embodiments, plant regeneration is obtained from cultured protoplasts, plant callus, explants, organs, pollens, embryos or parts thereof (see e.g. Evans et al. (1983), Handbook of Plant Cell Culture, Klee et al (1987) Ann. Rev. of Plant Phys.). However, the regeneration efficiency and ratio is very low.

[0196] In some embodiments, transformed or improved plants as described herein can be self-pollinated to provide seed for homozygous trait-improved plants of the disclosure, which have a desired trait such as seedlessness, reduced seed size, reduced endocarp tissue, or less lignified endocarp (homozygous for the DNA modification) or crossed with non-transgenic plants or different trait-improved plants to provide seed for heterozygous plants. Where a recombinant DNA was introduced into the plant cell, the resulting plant of such a crossing is a plant which is heterozygous for the recombinant DNA molecule. Both such homozygous and heterozygous plants obtained by crossing from the trait-improved plants and comprising the genetic modification (which can be a recombinant DNA) are referred to herein as “progeny”. Alternatively, gene-edited plants can be obtained by one of the methods described herein using an editing system whereby no foreign DNA is incorporated into the genome using transient expression/delivery or whereby foreign DNA is incorporated into the genome using stable transformation but removed/segregated away upon crossing. Progeny of such plants, obtained by further breeding may also contain the genetic modification such as nucleotide substitutions. Breedings may be performed by any breeding methods that are commonly used for different crops (e.g., Allard, Principles of Plant Breeding, John Wiley & Sons, NY, U. of CA, Davis, CA, 50-98 (1960). However, this approach cannot be applied to plants and/or crops with high heterozygosity and/or multiploidy.

[0197] Developing a method to generate CRISPR-mediated non-transgenic events is highly desirable for many applications of genome editing, particularly for asexually propagated, heterozygous, perennial crop plants. It has been reported that pre-assembled CRISPR/Cas9 ribonucleoproteins can be delivered into protoplasts to induce mutations, without the need for stable integration of CRISPR/Cas9 genes into the host-plant genome (Malnoy et al. 2016, Subburaj et al. 2016, Woo et al. 2015). Particle bombardment has also been used to deliver CRISPR/Cas9 ribonucleoproteins to wheat and maize cells, producing non-transgenic mutants (Liang et al. 2017, Svitashv et al 2016, Zhang et al. 2016, Kim 2017).

[0198] A method for production and expedient screening of CRISPR/Cas9-mediated non-transgenic mutant plants via transient T-DNA expression has been reported using tobacco as a model species (Che et al 2018). The mutant rate of non-transgenic plant from total regenerated plants was <0.5% when no selection was applied. The present disclosure teaches that transient expression of a morphogenic regulator in dicot species is demonstrated to enhance regeneration and may subsequently increase gene editing efficiency.

[0199] As mentioned above, transient expression of an editing system via *Agrobacterium*-mediated T-DNA deliv-

ery was reported to generate edited events without transgene integration, the overall efficiency was low due to the lack of selection (Chen et al. 2018). The present disclosure teaches use of transient expression of morphogenic regulator(s) to enhance the regeneration of edited cells without transgene integration. In some embodiments, morphogenic regulators are transiently transformed using *Agrobacterium*-mediated T-DNA delivery along with gene editing machinery that generates edited events.

[0200] In some embodiments, a gene-edited plant cell produced according to a method of the present is grown, multiplied, and/or regenerated in a medium without growth hormones, in which case, for example, at most 2%, at most 1%, at most 0.9%, at most 0.8%, at most 0.7%, at most 0.6%, at most 0.5%, at most 0.4%, at most 0.3%, at most 0.2%, at most 0.1%, at most 0.05%, at most 0.01%, at most 0.005%, at most 0.001%, at most 0.0005%, or at most 0.0001% or lower of the transformed cells are regenerated into a whole plants. In some embodiments, a gene-edited plant cell produced according to a method of the present invention is transformed with an expression cassette comprising one or more (e.g., 1, 2, 3, 4, 5, or more) morphogenic regulator(s) (such as WUS, BBN, KN1, IPT, CLV3, miR156, GRF5, NTT, and/or HDZipII) and regenerated in a medium without growth hormones, in which case, for example, at least 1%, at least 2%, at least 3%, at least 4%, at least 5%, at least 6%, at least 7%, at least 8%, at least 9%, at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95% or higher, of the transformed cells are regenerated into a whole plants.

[0201] In some embodiments of the present invention, transient expression of at least one morphogenic regulator in a transformed cell is capable of regenerating a plant without regeneration-inducing hormones added.

[0202] In some embodiments, transient expression of at least one morphogenic regulator in a transformed cell is capable of regenerating a plant in combination with addition of regeneration-inducing hormones in an improved, enhanced and/or synergistic manner.

[0203] In some embodiments, a plant cell of interest can be co-transformed with an expression cassette comprising one or more gene editing components/machinery and another expression cassette comprising at least one morphogenic regulator. In embodiments, the transient expression of at least one morphogenic regulator increases, enhances, improves transformation efficiency of the gene-editing system. In further embodiments, the transient expression of at least one morphogenic regulator increases, enhances, improves regeneration of the gene-edited plant cell. In further embodiments, the transient expression of at least one morphogenic regulator increases, enhances, improves regeneration of the gene-edited plant cell even in a culture medium without hormones such as regeneration-inducing hormones.

[0204] In other embodiments, the morphogenic regulators taught in the present disclosure can be used as a positive selection marker that gives rise to plant regeneration in a hormone-less culture medium. This allows for avoidance of use of a selectable marker gene (usually an antibiotic- or herbicide-resistant gene), which can make the transformed cells and/or regenerated plants transgenic due to undesired DNA pieces that is stably integrated and inherited.

[0205] In some embodiments, transient expression of morphogenic regulators in a plant cell transformed with the gene-editing machinery taught in the present disclosure increases, improves, and/or enhance plant transformation efficiency at least 0.5%, at least 1%, at least 2%, at least 3%, at least 4%, at least 5%, at least 6%, at least 7%, at least 8%, at least 9%, at least 10%, at least 12%, at least 15%, at least 18%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 100%, at least 150%, at least 200%, at least 250%, at least 300%, at least 400%, at least 500% or higher than a plant cell transformed with the gene-editing machinery, but without morphogenic regulator expression.

[0206] In some embodiments, transient expression of morphogenic regulators in a plant cell transformed with the gene-editing machinery taught in the present disclosure increases, improves, and/or enhance plant regeneration at least 0.5%, at least 1%, at least 2%, at least 3%, at least 4%, at least 5%, at least 6%, at least 7%, at least 8%, at least 9%, at least 10%, at least 12%, at least 15%, at least 18%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 100%, at least 150%, at least 200%, at least 250%, at least 300%, at least 400%, at least 500% or higher than a plant cell transformed with the gene-editing machinery, but without morphogenic regulator expression.

[0207] In some embodiments, the modified plant cells described herein demonstrate an altered expression and/or function of one or more endogenous target genes.

[0208] In some embodiments, the expression of an endogenous target gene in a particular pathway is reduced in the modified plant cells. In some embodiments, the expression of a plurality (e.g., two or more) of endogenous target genes in a particular pathway are reduced in the modified plant cells. For example, the expression of 2, 3, 4, 5, 6, 7, 8, 9, 10, or more endogenous target genes in a particular pathway may be reduced. In some embodiments, the expression of an endogenous target gene in one pathway and the expression of an endogenous target genes in another pathway is reduced in the modified plant cells. In some embodiments, the expression of a plurality of endogenous target genes in one pathway and the expression of a plurality of endogenous target genes in another pathway are reduced in the modified plant cells. For example, the expression of 2, 3, 4, 5, 6, 7, 8, 9, 10, or more endogenous target genes in one pathway may be reduced and the expression of 2, 3, 4, 5, 6, 7, 8, 9, 10, or more endogenous target genes in another particular pathway may be reduced. In some embodiments, the expression of a plurality of endogenous target genes in a plurality of pathways is reduced. For example, the expression of one endogenous gene from each of a plurality of pathways (e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, or more pathways) may be reduced. In additional aspects, the expression of a plurality of endogenous genes (e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, or more genes) from each of a plurality of pathways (e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, or more pathways) may be reduced.

[0209] In some embodiments, the function of a protein encoded by an endogenous target gene in a particular pathway is altered in the modified plant cells. In some embodiments, the functions of proteins encoded by a plurality (e.g., two or more) of endogenous target genes in a particular pathway are altered in the modified plant cells. For example, the function of proteins encoded by 2, 3, 4, 5, 6, 7, 8, 9, 10, or more endogenous target genes in a particular

pathway may be altered. In some embodiments, the function of a protein encoded by an endogenous target gene in one pathway and the function of an endogenous target genes in another pathway is altered in the modified plant cells. In some embodiments, the functions of proteins encoded by a plurality of endogenous target genes in one pathway and the function of proteins encoded by a plurality of endogenous target genes in another pathway are altered in the modified plant cells. For example, the function of proteins encoded by 2, 3, 4, 5, 6, 7, 8, 9, 10, or more endogenous target genes in one pathway may be altered and the function of proteins encoded by 2, 3, 4, 5, 6, 7, 8, 9, 10, or more endogenous target genes in another particular pathway may be altered. In some embodiments, the functions of proteins encoded by a plurality of endogenous target genes in a plurality of pathways are altered. For example, the function of a protein encoded by one endogenous gene from each of a plurality of pathways (e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, or more pathways) may be altered. In additional aspects, the function of proteins encoded by a plurality of endogenous genes (e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, or more genes) from each of a plurality of pathways (e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, or more pathways) may be altered.

[0210] Gene editing generally refers to the process of modifying the nucleotide sequence of a genome, preferably in a precise or pre-determined manner. Examples of methods of gene editing described herein include methods of using site-directed nucleases to cut deoxyribonucleic acid (DNA) at precise target locations in the genome, thereby creating single-strand or double-strand DNA breaks at particular locations within the genome. Such breaks can be and regularly are repaired by natural, endogenous cellular processes, such as homology-directed repair (HDR) and non-homologous end joining (NHEJ). These two main DNA repair processes consist of a family of alternative pathways. NHEJ directly joins the DNA ends resulting from a double-strand break, sometimes with the loss or addition of nucleotide sequence, which may disrupt or enhance gene expression. HDR utilizes a homologous sequence, or donor sequence, as a template for inserting a defined DNA sequence at the break point. The homologous sequence can be in the endogenous genome, such as a sister chromatid. Alternatively, the donor can be an exogenous nucleic acid, such as a plasmid, a single-strand oligonucleotide, a double-stranded oligonucleotide, a duplex oligonucleotide or a virus, that has regions of high homology with the nuclease-cleaved locus, but which can also contain additional sequence or sequence changes including deletions that can be incorporated into the cleaved target locus. A third repair mechanism can be microhomology-mediated end joining (MMEJ), also referred to as "Alternative NHEJ," in which the genetic outcome is similar to NHEJ in that small deletions and insertions can occur at the cleavage site. MMEJ can make use of homologous sequences of a few basepairs flanking the DNA break site to drive a more favored DNA end joining repair outcome, and recent reports have further elucidated the molecular mechanism of this process; see, e.g., Cho and Greenberg, *Nature* 518, 174-76 (2015); Kent et al., *Nature Structural and Molecular Biology*, Adv. Online doi:10.1038/nsmb.2961 (2015); Mateos-Gomez et al., *Nature* 518, 254-57 (2015); Ceccaldi et al., *Nature* 528, 258-62 (2015). In some instances, it may be possible to predict likely repair outcomes based on analysis of potential microhomologies at the site of the DNA break.

[0211] “Recombination” refers to a process of exchange of genetic information between two polynucleotides, including but not limited to, donor capture by non-homologous end joining (NHEJ) and homologous recombination. For the purposes of this disclosure, “homologous recombination (HR)” refers to the specialized form of such exchange that takes place, for example, during repair of double-strand breaks in cells via homology-directed repair (HDR) mechanisms. This process requires nucleotide sequence homology, uses a “donor” molecule as a template to repair a “target” molecule (i.e., the one that experienced the double-strand break), and is variously known as “non-crossover gene conversion” or “short tract gene conversion,” because it leads to the transfer of genetic information from the donor to the target. Without wishing to be bound by any particular theory, such transfer can involve mismatch correction of heteroduplex DNA that forms between the broken target and the donor, and/or synthesis-dependent strand annealing, in which the donor is used to resynthesize genetic information that will become part of the target, and/or related processes. Such specialized HR often results in an alteration of the sequence of the target molecule such that part or all of the sequence of the donor polynucleotide is incorporated into the target polynucleotide.

[0212] Gene editing methods contemplated in various embodiments comprise engineered nucleases, designed to bind and cleave a target DNA sequence in a gene of interest. The engineered nucleases contemplated in particular embodiments, can be used to introduce a double-strand break in a target polynucleotide sequence, which may be repaired (i) by non-homologous end joining (NHEJ) in the absence of a polynucleotide template, e.g., a donor repair template, or (ii) by homology directed repair (HDR), i.e., homologous recombination, in the presence of a donor repair template. Engineered nucleases contemplated in certain embodiments, can also be engineered as nickases, which generate single-stranded DNA breaks that can be repaired using the cell’s base-excision-repair (BER) machinery or homologous recombination in the presence of a donor repair template.

[0213] Gene editing via sequence-specific nucleases is known in the art. See references (1) Carroll, D. (2011) Genome engineering with zinc-finger nucleases. *Genetics*, 188, 773-82; (2) Wood, A. J. et al. (2011) Targeted gene editing across species using ZFNs and TALENs. *Science* (New York, N.Y.), 333, 307; (3) Perez-Pinera, P. et al. (2012) Advances in targeted gene editing. *Current opinion in chemical biology*, 16, 268-77, each of which is hereby incorporated by reference in their entireties.

[0214] A nuclease-mediated double-stranded DNA (dsDNA) break in the genome can be repaired by two main mechanisms: Non-Homologous End Joining (NHEJ), which frequently results in the introduction of non-specific insertions and deletions (indels), or homology directed repair (HDR), which incorporates a homologous strand as a repair template. See Symington, L. S. and Gautier, J. (2011) Double-strand break end resection and repair pathway choice. *Annual review of genetics*, 45, 247-71, which is hereby incorporated by reference in its entirety.

[0215] When a sequence-specific nuclease is delivered along with a homologous donor DNA construct containing the desired mutations, gene targeting efficiencies are increased by 1000-fold compared to just the donor construct alone. See Urnov et al. (2005) Highly efficient endogenous

human gene correction using designed zinc-finger nucleases. *Nature*, 435, 646-51, which is hereby incorporated by reference in its entirety.

[0216] In some embodiments, the gene editing techniques of the present disclosure are used for plants that are modified using any gene editing tool, including, but not limited to: ZFNs, TALENs, CRISPR, and Mega nuclease technologies. In some embodiments, the gene editing tools of the present disclosure comprise proteins or polynucleotides which have been custom designed to target and cut at specific deoxyribonucleic acid (DNA) sequences. In some embodiments, gene editing proteins are capable of directly recognizing and binding to selected DNA sequences. In other embodiments, the gene editing tools of the present disclosure form complexes, wherein nuclease components rely on nucleic acid molecules for binding and recruiting the complex to the target DNA sequence.

[0217] In some embodiments, the single component gene editing tools comprise a binding domain capable of recognizing specific DNA sequences in the genome of the plant and a nuclease that cuts double-stranded DNA. The rationale of gene editing technology taught in the present disclosure is the use of a tool that allows the introduction of site-specific mutations in the plant genome or the site-specific integration of genes.

[0218] Many methods are available for delivering genes into plant cells, e.g. transfection, electroporation, viral vectors and *Agrobacterium* mediated transfer. Genes can be expressed transiently from a plasmid vector. Once expressed, the genes generate the targeted mutation that will be stably inherited, even after the degradation of the plasmid containing the gene.

[0219] Customizable nucleases can be used to make targeted double-stranded breaks (DSB) in living cells, the repair of which can be exploited to induce desired sequence changes. Two competing pathways effect repairs in most cells, including plant cells. Repair of a nuclease-induced DSB by non-homologous end joining (NHEJ) leads to the introduction of insertion/deletion mutations (indels) with high frequencies. By contrast, DSB repair by homology directed repair (HDR) with a user-supplied “donor template” DNA can lead to the introduction of specific alterations (e.g., point mutations and insertions) or the correction of mutant sequences back to wild-type.

[0220] In some embodiments, a plant cell of interest is generated by gene editing accomplished with engineered nucleases targeting one or more loci that contributes to a target gene of interest. Without wishing to be bound to any particular theory, it is contemplated that engineered nucleases are designed to precisely disrupt one or more target genes of interest through gene editing and, once nuclease activity and specificity are validated, lead to predictable disruption of target gene expression and/or function.

[0221] The engineered nucleases described herein generate single-stranded DNA nicks or double-stranded DNA breaks (DSB) in a target sequence. Furthermore, a DSB can be achieved in the target DNA by the use of two nucleases generating single-stranded nicks (nickases). Each nickase cleaves one strand of the DNA and the use of two or more nickases can create a double strand break (e.g., a staggered double-stranded break) in a target DNA sequence. In other embodiments, the nucleases are used in combination with a

donor repair template, which is introduced into the target sequence at the DNA break-site via homologous recombination at a DSB.

[0222] Engineered nucleases described herein that are suitable for gene editing comprise one or more DNA binding domains and one or more DNA cleavage domains (e.g., one or more endonuclease and/or exonuclease domains), and optionally, one or more linkers contemplated herein. An “engineered nuclease” refers to a nuclease comprising one or more DNA binding domains and one or more DNA cleavage domains, wherein the nuclease has been designed and/or modified to bind a DNA binding target sequence adjacent to a DNA cleavage target sequence. The engineered nuclease may be designed and/or modified from a naturally occurring nuclease or from a previously engineered nuclease.

[0223] Illustrative examples of nucleases that may be engineered to bind and cleave a target sequence include, but are not limited to homing endonucleases (meganucleases), megaTALs, transcription activator-like effector nucleases (TALENs), zinc finger nucleases (ZFNs), and clustered regularly-interspaced short palindromic repeats (CRISPR)/Cas nuclease systems.

[0224] In some embodiments, the nucleases contemplated herein comprise one or more heterologous DNA-binding and cleavage domains (e.g., ZFNs, TALENs, megaTALs), (Boissel et al., 2014; Christian et al., 2010). In other embodiments, the DNA-binding domain of a naturally-occurring nuclease may be altered to bind to a selected target site (e.g., a meganuclease that has been engineered to bind to site different than the cognate binding site). For example, meganucleases have been designed to bind target sites different from their cognate binding sites (Boissel et al., 2014). In particular embodiments, a nuclease requires a nucleic acid sequence to target the nuclease to a target site (e.g., CRISPR/Cas).

(i) TALEN

[0225] Transcription activator-like effector nucleases (TALENs) comprise a nonspecific DNA-cleaving nuclease (e.g., a Fok I cleavage domain) fused to a DNA-binding domain that can be easily engineered so that TALENs can target essentially any sequence (See, e.g., Joung and Sander, *Nature Reviews Molecular Cell Biology* 14:49-55 (2013)). Methods for generating engineered TALENs are known in the art, see, e.g., the fast ligation-based automatable solid-phase high-throughput (FLASH) system described in U.S. Ser. No. 61/610,212, and Reyon et al., *Nature Biotechnology* 30,460-465 (2012); as well as the methods described in Bogdanove & Voytas, *Science* 333, 1843-1846 (2011); Bogdanove et al., *Curr Opin Plant Biol* 13, 394-401 (2010); Scholze & Boch, *J. Curr Opin Microbiol* (2011); Boch et al., *Science* 326, 1509-1512 (2009); Moscou & Bogdanove, *Science* 326, 1501 (2009); Miller et al., *Nat Biotechnol* 29, 143-148 (2011); Morbitzer et al., *T. Proc Natl Acad Sci USA* 107, 21617-21622 (2010); Morbitzer et al., *Nucleic Acids Res* 39, 5790-5799 (2011); Zhang et al., *Nat Biotechnol* 29, 149-153 (2011); Geissler et al., *PLoS ONE* 6, e19509 (2011); Weber et al., *PLoS ONE* 6, e19722 (2011); Christian et al., *Genetics* 186, 757-761 (2010); Li et al., *Nucleic Acids Res* 39, 359-372 (2011); Mahfouz et al., *Proc Natl Acad Sci USA* 108, 2623-2628 (2011); Mussolino et al., *Nucleic Acids Res* (2011); Li et al., *Nucleic Acids Res* 39, 6315-6325 (2011); Cermak et al., *Nucleic Acids Res* 39, e82

(2011); Wood et al., *Science* 333, 307 (2011); Hockemeyer et al. *Nat Biotechnol* 29, 731-734 (2011); Tesson et al., *Nat Biotechnol* 29, 695-696 (2011); Sander et al., *Nat Biotechnol* 29, 697-698 (2011); Huang et al., *Nat Biotechnol* 29, 699-700 (2011); and Zhang et al., *Nat Biotechnol* 29, 149-153 (2011); all of which are incorporated herein by reference in their entirety.

[0226] In some embodiments, a TALEN that binds to and cleaves a target region of a locus that contributes to a target gene of interest. A “TALEN” refers to an engineered nuclease comprising an engineered TALE DNA binding domain contemplated elsewhere herein and an endonuclease domain (or endonuclease half-domain thereof), and optionally comprise one or more linkers and/or additional functional domains, e.g., an end-processing enzymatic domain of an end-processing enzyme that exhibits 5-3' exonuclease, 5-3' alkaline exonuclease, 3-5'exonuclease (e.g., Trex2), 5' flap endonuclease, helicase or template-independent DNA polymerases activity.

[0227] In some embodiments, plants of interest are modified through Transcription activator-like (TAL) effector nucleases (TALENs). TALENs are polypeptides with repeat polypeptide arms capable of recognizing and binding to specific nucleic acid regions. By engineering the polypeptide arms to recognize selected target sequences, the TAL nucleases can be used to direct double stranded DNA breaks to specific genomic regions. These breaks can then be repaired via recombination to edit, delete, insert, or otherwise modify the DNA of a host organism. In some embodiments, TALENs are used alone for gene editing (e.g., for the deletion or disruption of a gene). In other embodiments, TALs are used in conjunction with donor sequences and/or other recombination factor proteins that will assist in the Non-homologous end joining (NHEJ) process to replace the targeted DNA region. For more information on the TAL-mediated gene editing compositions and methods of the present disclosure, see U.S. Pat. Nos. 8,440,432; 8,440,432; 8,450,471; 8,586,526; 8,586,363; 8,592,645; 8,697,853; 8,704,041; 8,921,112; and 8,912,138, each of which is hereby incorporated in its entirety for all purposes.

(ii) MegaTALs

[0228] Various illustrative embodiments contemplate a megaTAL nuclease that binds to and cleaves a target region of a locus that contributes to a target gene of interest. A “megaTAL” refers to an engineered nuclease comprising an engineered TALE DNA binding domain and an engineered meganuclease, and optionally comprise one or more linkers and/or additional functional domains, e.g., an end-processing enzymatic domain of an end-processing enzyme that exhibits 5-3' exonuclease, 5-Y alkaline exonuclease, 3-5'exonuclease, 5' flap endonuclease, helicase or template-independent DNA polymerases activity.

[0229] A “TALE DNA binding domain” is the DNA binding portion of transcription activator-like effectors (TALE or TAL-effectors), which mimics plant transcriptional activators to manipulate the plant transcriptome (see e.g., Kay et al., 2007. *Science* 318:648-651). TALE DNA binding domains contemplated in particular embodiments are engineered de novo or from naturally occurring TALEs. Illustrative examples of TALE proteins for deriving and designing DNA binding domains are disclosed in U.S. Pat. No. 9,017,967, and references cited therein, all of which are incorporated herein by reference in their entirety.

[0230] In some embodiments, plants of interest are modified through megaTALs. In some embodiments, megaTALs are engineered endonucleases capable of targeting selected DNA sequences and inducing DNA breaks.

(iii) Meganucleases/Homing Endonucleases (HE)

[0231] Meganucleases are sequence-specific endonucleases originating from a variety of organisms such as bacteria, yeast, algae and plant organelles. A number of Meganucleases are known in the art, see, e.g., WO 2012010976 (Meganuclease variants cleaving DNA target sequences of the TERT gene); U.S. Pat. Nos. 8,021,867; 8,119,361 and 8,119,381 (I-CreI meganucleases); U.S. Pat. No. 7,897,372 (I-CreI Meganuclease Variants with Modified Specificity).

[0232] In some embodiments, a homing endonuclease or meganuclease is engineered to bind to, and to introduce single-stranded nicks or double-strand breaks (DSBs) in, one or more loci that contribute to a target gene of interest. "Homing endonuclease" and "meganuclease" are used interchangeably and refer to naturally-occurring nucleases or engineered meganucleases that recognize 12-45 base-pair cleavage sites and are commonly grouped into five families based on sequence and structure motifs: LAGLIDADG (SED ID NO:135), GIY-YIG, HNH, His-Cys box, and PD-(D/E)XK.

[0233] Engineered HEs do not exist in nature and can be obtained by recombinant DNA technology or by random mutagenesis. Engineered HEs may be obtained by making one or more amino acid alterations, e.g., mutating, substituting, adding, or deleting one or more amino acids, in a naturally occurring HE or previously engineered HE. In particular embodiments, an engineered HE comprises one or more amino acid alterations to the DNA recognition interface. Engineered HEs contemplated in particular embodiments may further comprise one or more linkers and/or additional functional domains, e.g., an end-processing enzymatic domain of an end-processing enzyme that exhibits 5-3' exonuclease, 5-3' alkaline exonuclease, 3-5' exonuclease, 5' flap endonuclease, helicase or template-independent DNA polymerases activity.

[0234] In some embodiments, plants of interest are modified through meganucleases. In some embodiments, meganucleases are engineered endonucleases capable of targeting selected DNA sequences and inducing DNA breaks. In some embodiments, new meganucleases targeting specific regions are developed through recombinant techniques which combine the DNA binding motifs from various other identified nucleases. In other embodiments, new meganucleases are created through semi-rational mutational analysis, which attempts to modify the structure of existing binding domains to obtain specificity for additional sequences. For more information on the use of meganucleases for genome editing, see Silva et al., 2011 *Current Gene Therapy* 11 pg 11-27; and Stoddard et al., 2014 *Mobile DNA* 5 pg 7, each of which is hereby incorporated in its entirety for all purposes.

(iv) ZFN

[0235] Zinc-finger nucleases (ZFNs) are composed of programmable, sequence-specific zinc finger DNA-binding modules (see above) linked to a nonspecific DNA cleavage domain, e.g., a Fok I cleavage domain. Methods for making and using ZFNs are known in the art, see, e.g., (Maeder et al., 2008, *Mol. Cell*, 31:294-301; Joung et al., 2010, *Nat. Methods*, 7:91-92; Isalan et al., 2001, *Nat. Biotechnol.*,

19:656-660; Sander et al., *Nat. Methods*, 8(1):67-9, 2011; Bhakta et al., *Genome Res.* 23(3):530-8, 2013). In some embodiments, the ZFNs are described in, or are generated as described in, WO 2011/017293 or WO 2004/099366. Additional suitable ZFNs are described in U.S. Pat. Nos. 6,511, 808, 6,013,453, 6,007,988, and 6,503,717 and U.S. patent application 2002/0160940.

[0236] In some embodiments, a zinc finger nuclease (ZFN) that binds to and cleaves a target region of a locus that contributes to a target gene of interest. A "ZFN" refers to an engineered nuclease comprising one or more zinc finger DNA binding domains and an endonuclease domain (or endonuclease half-domain thereof), and optionally comprise one or more linkers and/or additional functional domains, e.g., an end-processing enzymatic domain of an end-processing enzyme that exhibits 5-3' exonuclease, 5-3' alkaline exonuclease, 3-5' exonuclease, 5' flap endonuclease, helicase or template-independent DNA polymerases activity.

[0237] In one embodiment, targeted double-stranded cleavage is achieved using two ZFNs, each comprising an endonuclease half-domain can be used to reconstitute a catalytically active cleavage domain. In another embodiment, targeted double-stranded cleavage is achieved with a single polypeptide comprising one or more zinc finger DNA binding domains and two endonuclease half-domains.

[0238] In one embodiment, a ZFN comprises a TALE DNA binding domain contemplated elsewhere herein, a zinc finger DNA binding domain, and an endonuclease domain (or endonuclease half-domain) contemplated elsewhere herein.

[0239] In one embodiment, a ZFN comprises a zinc finger DNA binding domain, and a meganuclease contemplated elsewhere herein.

[0240] In particular embodiments, the ZFN comprises a zinc finger DNA binding domain that has one, two, three, four, five, six, seven, or eight or more zinc finger motifs and an endonuclease domain (or endonuclease half-domain). Typically, a single zinc finger motif is about 30 amino acids in length. Zinc fingers motifs include both canonical C₂H₂ zinc fingers, and non-canonical zinc fingers such as, for example, C₃H zinc fingers and C₄ zinc fingers.

[0241] Zinc finger binding domains can be engineered to bind any DNA sequence. Candidate zinc finger DNA binding domains for a given 3 bp DNA target sequence have been identified and modular assembly strategies have been devised for linking a plurality of the domains into a multi-finger peptide targeted to the corresponding composite DNA target sequence. Other suitable methods known in the art can also be used to design and construct nucleic acids encoding zinc finger DNA binding domains, e.g., phage display, random mutagenesis, combinatorial libraries, computer/rational design, affinity selection, PCR, cloning from cDNA or genomic libraries, synthetic construction and the like. (See, e.g., U.S. Pat. No. 5,786,538; Wu et al., *PNAS* 92:344-348 (1995); Jamieson et al., *Biochemistry* 33:5689-5695 (1994); Rebar & Pabo, *Science* 263:671-673 (1994); Choo & Klug, *PNAS* 91:11163-11167 (1994); Choo & Klug, *PNAS* 91:11168-11172 (1994); Desjarlais & Berg, *PNAS* 90:2256-2260 (1993); Desjarlais & Berg, *PNAS* 89:7345-7349 (1992); Pomerantz et al., *Science* 267:93-96 (1995); Pomerantz et al., *PNAS* 92:9752-9756 (1995); Liu et al., *PNAS* 94:5525-5530 (1997); Griesman & Pabo, *Science* 275:657-661 (1997); Desjarlais & Berg, *PNAS* 91:11-99-11103 (1994)).

[0242] Individual zinc finger motifs bind to a three or four nucleotide sequence. The length of a sequence to which a zinc finger binding domain is engineered to bind (e.g., a target sequence) will determine the number of zinc finger motifs in an engineered zinc finger binding domain. For example, for ZFNs in which the zinc finger motifs do not bind to overlapping subsites, a six-nucleotide target sequence is bound by a two-finger binding domain; a nine-nucleotide target sequence is bound by a three-finger binding domain, etc. In particular embodiments, DNA binding sites for individual zinc fingers motifs in a target site need not be contiguous, but can be separated by one or several nucleotides, depending on the length and nature of the linker sequences between the zinc finger motifs in a multi-finger binding domain.

[0243] In some embodiments, plants of interest are modified through Zinc Finger Nucleases. Three variants of the ZFN technology are recognized in plant genome engineering (with applications ranging from producing single mutations or short deletions/insertions in the case of ZFN-1 and -2 techniques up to targeted introduction of new genes in the case of the ZFN-3 technique):

[0244] ZFN-1: Genes encoding ZFNs are delivered to plant cells without a repair template. The ZFNs bind to the plant DNA and generate site specific double-strand breaks (DSBs). The natural DNA-repair process (which occurs through nonhomologous end-joining, NHEJ) leads to site specific mutations, in one or only a few base pairs, or to short deletions or insertions.

[0245] ZFN-2: Genes encoding ZFNs are delivered to plant cells along with a repair template homologous to the targeted area, spanning a few kilo base pairs. The ZFNs bind to the plant DNA and generate site-specific DSBs. Natural gene repair mechanisms generate site-specific point mutations e.g. changes to one or a few base pairs through homologous recombination and the copying of the repair template.

[0246] ZFN-3: Genes encoding ZFNs are delivered to plant cells along with a stretch of DNA which can be several kilo base pairs long and the ends of which are homologous to the DNA sequences flanking the cleavage site. As a result, the DNA stretch is inserted into the plant genome in a site specific manner.

(v) FokI

[0247] FokI is a type II restriction endonuclease that includes a DNA recognition domain and a catalytic (endonuclease) domain. The fusion proteins described herein can include all of FokI or just the catalytic endonuclease domain, e.g., amino acids 388-583 or 408-583 of GenBank Acc. No. AAA24927.1, e.g., as described in WO95/09233, Li et al., *Nucleic Acids Res.* 39(1): 359-372 (2011); Cathomen and Joung, *Mol. Ther.* 16: 1200-1207 (2008), or a mutated form of FokI as described in Miller et al. *Nat Biotechnol* 25: 778-785 (2007); Szczepek et al., *Nat Biotechnol* 25: 786-793 (2007); or Bitinaite et al., *Proc. Natl. Acad. Sci. USA*. 95:10570-10575 (1998). See also Tsai et al., *Nat Biotechnol*. 2014 June; 32(6):569-76

[0248] In some embodiments, plants of interest are modified through FokI endonucleases.

[0249] Herein, the term “targeted gene-editing system” refers to a protein, nucleic acid, or combination thereof that is capable of substituting, inserting, or deleting one or more nucleotides at a target site and modifying an endogenous

target DNA sequence when introduced into a cell, thereby causing one or more amino acid substitutions, insertions, or deletions. In some embodiments, the editing system taught in the present disclosure refers to the targeted gene editing system.

[0250] More specifically, the term “targeted base-editing system” refers to a protein, nucleic acid, or combination thereof that is capable of substituting one or more nucleotides at a target site and modifying an endogenous target DNA sequence when introduced into a cell, thereby causing one or more amino acid substitutions.

[0251] Numerous gene editing systems suitable for use in the methods of the present disclosure, include, but are not limited to, zinc-finger nuclease systems, TALEN systems, and CRISPR/Cas systems.

[0252] In some embodiments, a nuclease-inactivated CRISPR/Cas system having a base deaminase activity is utilized for a targeted base-editing. In other aspects, a nickase is used.

[0253] In some embodiments, the targeted base-editing system can mediate a change in the sequence of the endogenous target gene, for example, by introducing one or more point mutations into the endogenous target sequence, such as by substituting C with T (or G with A) or A with G (or T with C) in the endogenous target sequence.

[0254] In some embodiments, the targeted gene-editing system may mediate a change in the expression of the protein encoded by the endogenous target gene. In such embodiments, the targeted gene-editing system may regulate the expression of the encoded protein by modifications of the endogenous target DNA sequence, or by acting on the mRNA product encoded by the DNA sequence. In some embodiments, the targeted gene-editing system may result in the expression of a modified endogenous protein. In some embodiments, the modifications to the endogenous DNA sequence mediated by the targeted gene-editing system result in an altered function of the modified endogenous protein as compared to the corresponding endogenous protein in an unmodified plant cell. In such embodiments, the expression level of the modified endogenous protein may be increased, decreased or may be the same, or substantially similar to, the expression level of the corresponding endogenous protein in an unmodified plant cell.

[0255] The present disclosure provides a targeted gene-editing system to edit a target nucleotide sequence in the genome of a plant, comprising at least one of the followings; i) a gene editing fusion protein, and a guide RNA; ii) an expression construct comprising a nucleotide sequence encoding a gene editing fusion protein, and a guide RNA; iii) a gene editing fusion protein, and an expression construct comprising a nucleotide sequence encoding a guide RNA; iv) an expression construct comprising a nucleotide sequence encoding a gene editing fusion protein, and an expression construct comprising a nucleotide sequence encoding a guide RNA; v) an expression construct comprising a nucleotide sequence encoding gene editing fusion protein and a nucleotide sequence encoding guide RNA; wherein said gene editing fusion protein contains a CRISPR-associated effector domain and optionally a deaminase domain, said guide RNA can target said gene editing fusion protein to the target sequence in the plant genome.

[0256] In some embodiments, the present disclosure provides a polynucleotide encoding a gRNA. In some embodiments, a gRNA-encoding nucleic acid is comprised in an

expression vector, e.g., a recombinant expression vector. In some embodiments, the present disclosure provides a polynucleotide encoding a site-directed modifying polypeptide. In some embodiments, the polynucleotide encoding a site-directed modifying polypeptide is comprised in an expression vector, e.g., a recombinant expression vector.

[0257] In some embodiments, the targeted gene-editing system comprises a Cas protein as a CRISPR-associated effector domain. Cas molecules of a variety of species can be used in the methods and compositions described herein, including Cas molecules derived from *S. pyogenes*, *S. aureus*, *N. meningitidis*, *S. thermophiles*, *Acidovorax avenae*, *Actinobacillus pleuropneumoniae*, *Actinobacillus succinogenes*, *Actinobacillus suis*, *Actinomyces* sp., *Cycliphilus denitrificans*, *Aminomonas paucivorans*, *Bacillus cereus*, *Bacillus smithii*, *Bacillus thuringiensis*, *Bacteroides* sp., *Blastopirellula marina*, *Bradyrhizobium* sp., *Brevibacillus laterosporus*, *Campylobacter coli*, *Campylobacter jejuni*, *Campylobacter lari*, *Candidatus puniceispirillum*, *Clostridium cellulolyticum*, *Clostridium perfringens*, *Corynebacterium accolens*, *Corynebacterium diphtheriae*, *Corynebacterium matruchotii*, *Dinoroseobacter shibae*, *Eubacterium dolichum*, *Gammaproteobacterium*, *Gluconacetobacter diazotrophicus*, *Haemophilus parainfluenzae*, *Haemophilus sputorum*, *Helicobacter canadensis*, *Helicobacter cinaedi*, *Helicobacter mustelae*, *Ilyobacter polytropus*, *Kingella kingae*, *Lactobacillus crispatus*, *Listeria ivanovii*, *Listeria monocytogenes*, *Listeriaceae bacterium*, *Methylocystis* sp., *Methylosinus trichosporium*, *Mobiluncus mulieris*, *Neisseria bacilliformis*, *Neisseria cinerea*, *Neisseria flavescens*, *Neisseria lactamica*, *Neisseria meningitidis*, *Neisseria* sp., *Neisseria wadsworthii*, *Nitrosomonas* sp., *Parvibaculum lavamentivorans*, *Pasteurella multocida*, *Phascolarctobacterium succinatutens*, *Ralstonia syzygii*, *Rhodopseudomonas palustris*, *Rhodovulum* sp., *Simonsiella muelleri*, *Sphingomonas* sp., *Sporolactobacillus vineae*, *Staphylococcus aureus*, *Staphylococcus lugdunensis*, *Streptococcus* sp., *Subdoligranulum* sp., *Tistrella mobilis*, *Treponema* sp., or *Verminephrobacter eiseniae*.

[0258] In some embodiments, the Cas protein is a naturally-occurring Cas protein. In other embodiments, the Cas protein is an engineered Cas protein. In some embodiments, the Cas endonuclease is selected from the group consisting of C2C1, C2C3, Cpf1 (also referred to as Cas12a), Cas12b, Cas12c, Cas12d, Cas12e, Cas13a, Cas13b, Cas13c, Cas13d, Cas1, Cas1B, Cas2, Cas3, Cas4, Cas5, Cas6, Cas7, Cas8, Cas9 (also known as Csn1 and Csx12), Cas10, Csy1, Csy2, Csy3, Cse1, Cse2, Csc1, Csc2, Csa5, Csn2, Csm2, Csm3, Csm4, Csm5, Csm6, Cmr1, Cmr3, Cmr4, Cmr5, Cmr6, Csb1, Csb2, Csb3, Csx17, Csx14, Csx10, Csx16, CsaX, Csx3, Csx1, Csx15, Csf1, Csf2, Csf3, and Csf4.

[0259] In some embodiments, the Cas protein is an endonuclease such as a Cas13 protein. In some embodiments, the Cas13 protein is a Cas13a (Abudayyeh et al., Nature 550 (2017), 280-284), Cas13b (Cox et al., Science (2017) 358: 6336, 1019-1027), Cas13c (Cox et al., Science (2017) 358:6336, 1019-1027), or Cas13d (Zhang et al., Cell 175 (2018), 212-223) protein.

[0260] In some embodiments, the Cas protein is a wild type or naturally occurring Cas9 protein or a Cas9 ortholog. Wild type Cas9 is a multi-domain enzyme that uses an HNH nuclease domain to cleave the target strand of DNA and a RuvC-like domain to cleave the non-target strand. Binding of WT Cas9 to DNA based on gRNA specificity results in

double-stranded DNA breaks that can be repaired by non-homologous end joining (NHEJ) or homology-directed repair (HDR).

[0261] In some embodiments, the naturally occurring Cas9 polypeptide is selected from the group consisting of SpCas9, SpCas9-HF1, SpCas9-HF2, SpCas9-HF3, SpCas9-HF4, SaCas9, FnCpf, FnCas9, eSpCas9, and NmeCas9. In some embodiments, the Cas9 protein comprises an amino acid sequence having at least 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to a Cas9 amino acid sequence described in Chylinski et al., RNA Biology 2013 10:5, 727-737; Hou et al., PNAS Early Edition 2013, 1-6).

[0262] In some embodiments, the Cas polypeptide comprises one or more of the following activities: (a) a nickase activity, i.e., the ability to cleave a single strand, e.g., the non-complementary strand or the complementary strand, of a nucleic acid molecule; (b) a double stranded nuclease activity, i.e., the ability to cleave both strands of a double stranded nucleic acid and create a double stranded break, which in an embodiment is the presence of two nickase activities; (c) an endonuclease activity; (d) an exonuclease activity; and/or (e) a helicase activity, i.e., the ability to unwind the helical structure of a double stranded nucleic acid. In other embodiments, the Cas protein may be dead or inactive (e.g. dCas).

[0263] In some embodiments, the Cas polypeptide is fused to heterologous polypeptide/proteins that has base deaminase activity.

[0264] In some embodiments, different Cas proteins (i.e., Cas9 proteins from various species) may be advantageous to use in the various provided methods in order to capitalize on various enzymatic characteristics of the different Cas proteins (e.g., for different PAM sequence preferences; for increased or decreased enzymatic activity; for an increased or decreased level of cellular toxicity; to change the balance between NHEJ, homology-directed repair, single strand breaks, double strand breaks, etc.).

[0265] In some embodiments, the Cas protein is a Cas9 protein derived from *S. pyogenes* and recognizes the PAM sequence motif NGG, NAG, NGA (Mali et al, Science 2013; 339(6121): 823-826). In some embodiments, the Cas protein is a Cas9 protein derived from *S. thermophiles* and recognizes the PAM sequence motif NGGNG and/or NNAGAAW (W=A or T) (See, e.g., Horvath et al, Science, 2010; 327 (5962): 167-170, and Deveau et al, J Bacteriol 2008; 190(4): 1390-1400). In some embodiments, the Cas protein is a Cas9 protein derived from *S. mutans* and recognizes the PAM sequence motif NGG and/or NAAR (R=A or G) (See, e.g., Deveau et al, J BACTERIOL 2008; 190(4): 1390-1400). In some embodiments, the Cas protein is a Cas9 protein derived from *S. aureus* and recognizes the PAM sequence motif NNGRR (R=A or G). In some embodiments, the Cas protein is a Cas9 protein derived from *S. aureus* and recognizes the PAM sequence motif N GRRT (R=A or G). In some embodiments, the Cas protein is a Cas9 protein derived from *S. aureus* and recognizes the PAM sequence motif N GRRV (R=A or G). In some embodiments, the Cas protein is a Cas9 protein derived from *N. meningitidis* and recognizes the PAM sequence motif N GATT or N GCTT (R=A or G, V=A, G or C) (See, e.g., Hou et al, PNAS 2013, 1-6). In the aforementioned embodiments, N can be any nucleotide residue, e.g., any of A, G, C or T. In some embodiments, the

Cas protein is a Cas13a protein derived from *Leptotrichia shahii* and recognizes the PFS sequence motif of a single 3' A, U, or C.

[0266] In embodiments, a Cas protein as a CRISPR-associated effector domain is codon-optimized based on plant genomes of the present disclosure.

[0267] In some embodiments, the at least one CRISPR-associated effector is selected from at least one of a nuclease, comprising a CRISPR nuclease, including Cas or Cpf1 nucleases.

[0268] In some embodiments, the at least one CRISPR-associated effector is a Cas polypeptide. In some embodiments, the at least one CRISPR-associated effector is a Cas polypeptide, wherein the Cas polypeptide comprises a site-specific DNA binding domain linked to at least one base editor. The CRISPR-associated effector or the nucleic acid sequence encoding the same, is selected from the group comprising (i) Cas9, including SpCas9, SaCas9, SaKKH-Cas9, VQR-Cas9, StlCas9, (ii) Cpf1, including AsCpf1, LbCpf1, FnCpf1, (iii) CasX, or (iv) CasY, or any variant or derivative of the aforementioned CRISPR-associated effector, preferably wherein the at least one CRISPR-associated effector comprises a mutation in comparison to the respective wild type sequence so that the resulting CRISPR-associated effector is converted to a single-strand specific DNA nickase, or to a DNA binding effector lacking all DNA cleavage ability, as described below.

[0269] Therefore, according to the present disclosure, artificially modified CRISPR nucleases are envisaged, which might indeed not be any “nucleases” in the sense of double-strand cleaving enzymes, but which are nickases or nuclease-dead variants, which still have inherent DNA recognition and thus binding ability.

[0270] In some embodiments, the Cas protein described above can be engineered to alter one or more properties of the Cas polypeptide. For example, in some embodiments, the Cas polypeptide comprises altered enzymatic properties, e.g., altered nuclease activity, (as compared with a naturally occurring or other reference Cas molecule) or altered helicase activity. In some embodiments, an engineered Cas polypeptide can have an alteration that alters its size, e.g., a deletion of amino acid sequence that reduces its size without significant effect on another property of the Cas polypeptide. In some embodiments, an engineered Cas polypeptide comprises an alteration that affects PAM recognition. For example, an engineered Cas polypeptide can be altered to recognize a PAM sequence other than the PAM sequence recognized by the corresponding wild type Cas protein. In some embodiments, the targeted gene-editing system comprises a Cas protein as a CRISPR-associated effector domain.

[0271] Cas polypeptides with desired properties can be made in a number of ways, including alteration of a naturally occurring Cas polypeptide or parental Cas polypeptide, to provide a mutant or altered Cas polypeptide having a desired property. For example, one or more mutations can be introduced into the sequence of a parental Cas polypeptide (e.g., a naturally occurring or engineered Cas polypeptide). Such mutations and differences may comprise substitutions (e.g., conservative substitutions or substitutions of non-essential amino acids); insertions; or deletions. In some embodiments, a mutant Cas polypeptide comprises one or more mutations (e.g., at least 1, 2, 3, 4, 5, 10, 15, 20, 30, 40 or 50 mutations) relative to a parental Cas polypeptide.

[0272] In an embodiment, a mutant Cas polypeptide comprises a cleavage property that differs from a naturally occurring Cas polypeptide. In some embodiments, the Cas is a deactivated Cas (dCas) mutant, which is catalytically dead. In such embodiments, the Cas polypeptide does not comprise any intrinsic enzymatic activity and is unable to mediate target nucleic acid cleavage. In such embodiments, the dCas is fused with a heterologous protein that is capable of modifying the target nucleic acid in a non-cleavage based manner. In some embodiments, the targeted gene-editing system comprises a Cas protein as a CRISPR-associated effector domain.

[0273] In some embodiments, the dCas is a dCas9 mutant. In some embodiments, a dCas protein is fused to base deaminase domains (e.g., cytidine deaminase, or adenosine deaminase). In some such cases, the dCas fusion protein is targeted by the gRNA to a specific location (i.e., sequence) in the target nucleic acid and exerts locus-specific modification such as replacing C with T (or G with A) if the fusion protein has cytidine deaminase activity) or replacing A with G (or T with C) if the fusion protein has adenosine deaminase activity.

[0274] In some embodiments, the dCas is a dCas13 mutant (Koner mann et al., Cell 173 (2018), 665-676). These dCas13 mutants can then be fused to enzymes that modify RNA, including adenosine deaminases (e.g., ADAR1 and ADAR2). Adenosine deaminases convert adenine to inosine, which the translational machinery treats like guanine, thereby creating a functional A to G change in the target sequence.

[0275] In some embodiments, the CRISPR-associated effector protein is Cas9 endonuclease. In some embodiments, the CRISPR-associated effector protein is a CRISPR-Cas variant, which is dCas9 mutant or nCas9 nickase mutant. The Cas9 endonuclease has a DNA cleavage domain containing two subdomains: i) the RuvC subdomain cleaving the non-complementary single-stranded chain and ii) the HNH nuclease subdomain cleaving the chain that is complementary to gRNA. Mutations in these subdomains can inactivate Cas9 endonuclease to form deactivated Cas9 (dCas9), which is interchangeably used with “catalytically dead Cas9”. The nuclease-inactivated Cas9 retains DNA binding capacity directed by gRNA. Thus, in principle, when fused with an additional protein, the dCas9 can simply target said additional protein to almost any DNA sequence through co-expression with appropriate guide RNA. For example, catalytically dead Cas9 (dCas9), which contains Asp10Ala (D10A) and His840Ala (H840A) mutations that inactivate its nuclease activity, retains its ability to bind DNA in a guide RNA-programmed manner, but does not cleave the DNA backbone (Komor et al., nature (2016), Vol 533:420-424). In some embodiments, conjugation of dCas9 with an enzymatic or chemical catalyst that mediates the direct conversion of one base to another could enable RNA-programmed DNA base editing.

[0276] In some embodiments, the mutant Cas9 is a Cas9 nickase (nCas9) mutant. Cas9 nickase mutants comprise only one catalytically active domain (either the RuvC domain (D10A) or the HNH domain (H840A)). The Cas9 nickase mutants retain DNA binding based on gRNA specificity, but are capable of cutting only one strand of DNA resulting in a single-strand break (e.g. a “nick”). In some embodiments, two complementary Cas9 nickase mutants (e.g., one Cas9 nickase mutant with an inactivated RuvC

domain, and one Cas9 nickase mutant with an inactivated HNH domain) are expressed in the same cell with two gRNAs corresponding to two respective target sequences; one target sequence on the sense DNA strand, and one on the antisense DNA strand. This dual-nickase system results in staggered double stranded breaks and can increase target specificity, as it is unlikely that two off-target nicks will be generated close enough to generate a double stranded break. In some embodiments, a Cas9 nickase mutant is co-expressed with a nucleic acid repair template to facilitate the incorporation of an exogenous nucleic acid sequence by homology-directed repair.

[0277] The dCas9 of the present disclosure can be derived from Cas9 of different species, for example, derived from *S. pyogenes* Cas9 (SpCas9). Mutations in both the RuvC subdomain and the HNH nuclease subdomain of the SpCas9 (includes, for example, D10A and H840A mutations) inactivate *S. pyogenes* Cas9 nuclease, resulting in a nuclease-dead/catalytically dead Cas9 (dCas9). In some embodiments of the present disclosure, the nuclease-inactivated Cas9 comprises the dCas9. In some preferred embodiments, the nuclease-inactivated Cas9 comprises.

[0278] Inactivation of one of the subdomains by mutation allows Cas9 to gain nickase activity, i.e., resulting in a Cas9 nickase (nCas9), for example, nCas9 with a D10A mutation only.

[0279] In some embodiments, the nuclease-inactivated Cas9 comprises amino acid substitutions D10A and/or H840A relative to wild type Cas9. In some embodiments, the nuclease-inactivated Cas9 of the present disclosure loses nuclease activity completely, which is catalytically dead. In such embodiments, the nuclease-inactivated Cas9 is the dCas9 with D10A and H840A. Therefore, the term “nuclease-inactivated Cas9” refers to dCas9 and/or nCas9.

[0280] In other embodiments, the nuclease-inactivated Cas9 of the present disclosure has nickase activity. In some embodiments, the nuclease-inactivated Cas9 is a Cas9 nickase that retains the cleavage activity of the HNH subdomain of Cas9, whereas the cleavage activity of the RuvC subdomain is inactivated. For example, the nuclease-inactivated Cas9 contains an amino acid substitution D10A relative to wild type Cas9. In such embodiments, the nuclease-inactivated Cas9 is the nCas9 with D10A only. In some embodiments of the present disclosure, the nuclease-inactivated Cas9 comprises the nCas9.

[0281] In some embodiments, the Cas polypeptides described herein can be engineered to alter the PAM/PFS specificity of the Cas polypeptide. In some embodiments, a mutant Cas polypeptide has a PAM/PFS specificity that is different from the PAM/PFS specificity of the parental Cas polypeptide. For example, a naturally occurring Cas protein can be modified to alter the PAM/PFS sequence that the mutant Cas polypeptide recognizes to decrease off target sites, improve specificity, or eliminate a PAM/PFS recognition requirement. In some embodiments, a Cas protein can be modified to increase the length of the PAM/PFS recognition sequence. In some embodiments, the length of the PAM recognition sequence is at least 4, 5, 6, 7, 8, 9, 10 or 15 amino acids in length. Cas polypeptides that recognize different PAM/PFS sequences and/or have reduced off-target activity can be generated using directed evolution. Exemplary methods and systems that can be used for directed evolution of Cas polypeptides are described, e.g., in Esvelt et al. Nature 2011, 472(7344): 499-503.

[0282] Exemplary Cas mutants are described in International PCT Publication No. WO 2015/161276 and Konermann et al., Cell 173 (2018), 665-676, which are incorporated herein by reference in their entireties.

[0283] In some embodiments, the deaminase domain is fused to the N-terminus of the nuclease-inactivated Cas9 domain. In some embodiments, the deaminase domain is fused to the C-terminus of the nuclease-inactivated Cas9 domain. In some embodiments, the deaminase domain and the nuclease inactivated Cas9 domain are fused through a linker. The linker can be a non-functional amino acid sequence having no secondary or higher structure, N-terminus and one or more NLSs at the C-terminus. Where there are more than one NLS, each NLS may be selected as independent from other NLSs. In some embodiments, the targeted base-editing fusion protein comprises two NLSs, for example, the two NLSs are located at the N-terminus and the C-terminus, respectively.

[0284] In some embodiment, a targeted base modification is a conversion of any nucleotide C, A, T, or G, to any other nucleotide. Any one of a C, A, T or G nucleotide can be exchanged in a site-directed way as mediated by a base editor, or a catalytically active fragment thereof, to another nucleotide. A base editing complex can comprise any base editor, or a base editor domain or catalytically active fragment thereof, which can convert a nucleotide of interest into any other nucleotide of interest in a targeted way.

[0285] A base editing domain according to the present disclosure can comprise at least one cytidine deaminase, or a catalytically active fragment thereof. The at least one base editing complex can comprise the cytidine deaminase, or a domain thereof in the form of a catalytically active fragment, as base editor.

[0286] According to the present disclosure, cytidine deaminases that can be used in connection with the present disclosure include, but are not limited to, members of the enzyme family known as apolipoprotein B mRNA-editing complex (APOBEC) family deaminase, an activation-induced deaminase (AID), or a cytidine deaminase 1 (CDA1). In particular embodiments, the deaminase in an APOBEC1 deaminase, an APOBEC2 deaminase, an APOBEC3A deaminase, an APOBEC3B deaminase, an APOBEC3C deaminase, and APOBEC3D deaminase, an APOBEC3E deaminase, an APOBEC3F deaminase an APOBEC3G deaminase, an APOBEC3H deaminase, or an APOBEC4 deaminase.

[0287] In the methods and systems of the present disclosure, the cytidine deaminase is capable of targeting Cytosine in a DNA single strand. In certain example embodiments the cytidine deaminase may edit on a single strand present outside of the binding component e.g. bound Cas9 and/or Cas13. In other example embodiments, the cytidine deaminase may edit at a localized bubble, such as a localized bubble formed by a mismatch at the target edit site but the guide sequence.

[0288] In some embodiments, the cytidine deaminase protein recognizes and converts one or more target cytosine residue(s) in a single-stranded bubble of a DNA-RNA heteroduplex into uracil residues (s). In some embodiments, the cytidine deaminase protein recognizes a binding window on the single-stranded bubble of a DNA-RNA heteroduplex. In some embodiments, the binding window contains at least one target cytosine residue(s). In some embodiments, the binding window is in the range of about 3 bp to about 100

bp. In some embodiments, the binding window is in the range of about 5 bp to about 50 bp. In some embodiments, the binding window is in the range of about 10 bp to about 30 bp. In some embodiments, the binding window is about 1 bp, 2 bp, 3 bp, 5 bp, 7 bp, 10 bp, 15 bp, 20 bp, 25 bp, 30 bp, 40 bp, 45 bp, 50 bp, 55 bp, 60 bp, 65 bp, 70 bp, 75 bp, 80 bp, 85 bp, 90 bp, 95 bp, or 100 bp.

[0289] In some embodiments, the cytidine deaminase protein comprises one or more deaminase domains. Not intended to be bound by theory, it is contemplated that the deaminase domain functions to recognize and convert one or more target cytosine (C) residue(s) contained in a single-stranded bubble of a DNA-RNA heteroduplex into (an) uracil (U) residue (s). In some embodiments, the deaminase domain comprises an active center. In some embodiments, the active center comprises a zinc ion. In some embodiments, amino acid residues in or near the active center interact with one or more nucleotide(s) 5' to a target cytosine residue. In some embodiments, amino acid residues in or near the active center interact with one or more nucleotide(s) 3' to a target cytosine residue.

[0290] In some embodiments, the cytidine deaminase comprises human APOBEC1 full protein (hAPOBEC1) or the deaminase domain thereof (hAPOBEC1-D) or a C-terminally truncated version thereof (hAPOBEC-T). In some embodiments, the cytidine deaminase is an APOBEC family member that is homologous to hAPOBEC1, hAPOBEC-D or hAPOBEC-T. In some embodiments, the cytidine deaminase comprises human AID1 full protein (hAID) or the deaminase domain thereof (hAID-D) or a C-terminally truncated version thereof (hAID-T). In some embodiments, the cytidine deaminase is an AID family member that is homologous to hAID, hAID-D or hAID-T. In some embodiments, the hAIDT is a hAID which is C-terminally truncated by about 20 amino acids.

[0291] In some embodiments, the cytidine deaminase comprises the wild type amino acid sequence of a cytosine deaminase. In some embodiments, the cytidine deaminase comprises one or more mutations in the cytosine deaminase sequence, such that the editing efficiency, and/or substrate editing preference of the cytosine deaminase is changed according to specific needs.

[0292] Certain mutations of APOBEC 1 and APOBEC3 proteins have been described in Kim et al., *Nature Biotechnology* (2017) 35(4):371-377 and Harris et al. *Mol. Cell* (2002) 10:1247-1253, each of which is incorporated herein by reference in its entirety.

[0293] Additional embodiments of the cytidine deaminase are disclosed in WO2017/070632 and WO2018/213726, each of which is incorporated herein by reference in its entirety.

[0294] In some embodiments, at least one CRISPR-associated effector is temporarily or permanently linked to at least one base editor to form a targeted base editing complex, which is a base editing fusion protein, wherein the base editing complex mediates the at least one targeted base modification. The at least one CRISPR-associated effector can be non-covalently (temporarily) or covalently (permanently) be attached to at least one base editor. Any component of the at least one base editor can be temporarily or permanently linked to the at least one CRISPR-associated effector.

[0295] The adenosine deaminases (e.g. engineered adenosine deaminases, evolved adenosine deaminases) pro-

vided herein may be from any organism, such as a bacterium. In some embodiments, the deaminase or deaminase domain is a variant of a naturally-occurring deaminase from an organism. In some embodiments, the deaminase or deaminase domain does not occur in nature. For example, in some embodiments, the deaminase or deaminase domain is at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to a naturally-occurring deaminase. In some embodiments, the adenosine deaminase is from a bacterium, such as, *E. coli*, *S. aureus*, *S. typhi*, *S. putrefaciens*, *H. influenzae*, or *C. crescentus*. In some embodiments, the adenosine deaminase is a TadA deaminase. In some embodiments, the TadA deaminase is an *E. coli* TadA deaminase (ecTadA). In some embodiments, the TadA deaminase is a truncated *E. coli* TadA deaminase. For example, the truncated ecTadA may be missing one or more N-terminal amino acids relative to a full-length ecTadA. In some embodiments, the truncated ecTadA may be missing 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 6, 17, 18, 19, or 20 N-terminal amino acid residues relative to the full length ecTadA. In some embodiments, the truncated ecTadA may be missing 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 6, 17, 18, 19, or 20 C-terminal amino acid residues relative to the full length ecTadA. In some embodiments, the ecTadA deaminase does not comprise an N-terminal methionine.

[0296] Some aspects of the disclosure provide adenosine deaminases. In some embodiments, the adenosine deaminases provided herein are capable of deaminating adenine. In some embodiments, the adenosine deaminases provided herein are capable of deaminating adenine in a deoxyadenosine residue of DNA. The adenosine deaminase may be derived from any suitable organism (e.g., *E. coli*). In some embodiments, the adenine deaminase is a naturally-occurring adenosine deaminase that includes one or more mutations corresponding to any of the mutations provided herein (e.g., mutations in ecTadA). One of skill in the art will be able to identify the corresponding residue in any homologous protein and in the respective encoding nucleic acid by methods well known in the art, e.g., by sequence alignment and determination of homologous residues. Accordingly, one of skill in the art would be able to generate mutations in any naturally-occurring adenosine deaminase (e.g., having homology to ecTadA) that corresponds to any of the mutations described herein, e.g., any of the mutations identified in ecTadA. In some embodiments, the adenosine deaminase is from a prokaryote. In some embodiments, the adenosine deaminase is from a bacterium. In some embodiments, the adenosine deaminase is from *Escherichia coli*, *Staphylococcus aureus*, *Salmonella typhi*, *Shewanella putrefaciens*, *Haemophilus influenzae*, *Caulobacter crescentus*, or *Bacillus subtilis*. In some embodiments, the adenosine deaminase is from *E. coli*.

[0297] In other embodiments, adenosine deaminases that can be used that include, but are not limited to, members of the enzyme family known as adenosine deaminases that act on RNA (ADARs), members of the enzyme family known as adenosine deaminases that act on tRNA (ADATs), and other adenosine deaminase domain-containing (ADAD) family members. According to the present disclosure, the adenosine deaminase is capable of targeting adenine in a RNA/DNA heteroduplex. Indeed, Zheng et al. (*Nucleic Acids Res.* 2017, 45(6): 3369-3377) has demonstrated that

ADARs can carry out adenosine to inosine editing reactions on RNA/DNA heteroduplexes. In particular embodiments, the adenosine deaminase has been modified to increase its ability to edit DNA in a RNA/DNA heteroduplex as detailed herein below.

[0298] In some embodiments, the adenosine deaminase is derived from one or more metazoa species, including but not limited to, mammals, birds, frogs, squids, fish, flies and worms. In some embodiments, the adenosine deaminase is a human, squid or *Drosophila* adenosine deaminase.

[0299] In some embodiments, the adenosine deaminase is a human ADAR, including hADAR1, hADAR2, hADAR3. In some embodiments, the adenosine deaminase is a *Caenorhabditis elegans* ADAR protein, including ADR-1 and ADR-2. In some embodiments, the adenosine deaminase is a *Drosophila* ADAR protein, including dAdar. In some embodiments, the adenosine deaminase is a squid *Loligo pealeii* ADAR protein, including sqADAR2a and sqADAR2b. In some embodiments, the adenosine deaminase is a human ADAT protein. In some embodiments, the adenosine deaminase is a *Drosophila* ADAT protein. In some embodiments, the adenosine deaminase is a human ADAD protein, including TE R (hADAD1) and TE RL (hADAD2).

[0300] In some embodiments, the adenosine deaminase is a TadA protein such as *E. coli* TadA. See Kim et al., Biochemistry 45:6407-6416 (2006); Wolf et al., EMBO J. 21:3841-3851 (2002), each of which is incorporated herein by reference in its entirety. In some embodiments, the adenosine deaminase is mouse ADA (See Grunebaum et al., Curr. Opin. Allergy Clin. Immunol. 13:630-638 (2013)) or human ADAT2 (See Fukui et al., J. Nucleic Acids 2010: 260512 (2010)), each of which is incorporated herein by reference in its entirety.

[0301] Additional embodiments of the adenosine deaminase are disclosed in U.S. Pat. No. 10,113,163 and WO2018/213708, each of which is incorporated herein by reference in its entirety.

[0302] In some embodiments, at least one CRISPR-associated effector is temporarily or permanently linked to at least one base editor to form a targeted base editing complex, which is a base editing fusion protein, wherein the base editing complex mediates the at least one targeted base modification. The at least one CRISPR-associated effector can be non-covalently (temporarily) or covalently (permanently) be attached to at least one base editor. Any component of the at least one base editor can be temporarily or permanently linked to the at least one CRISPR-associated effector.

[0303] In one aspect the present disclosure provides methods for targeted deamination of adenine in a DNA, more particularly in a locus of interest. The disclosure teaches the adenosine deaminase (AD) protein is recruited specifically to the relevant Adenine in the locus of interest by a CRISPR-Cas complex which can specifically bind to a target sequence. In order to achieve this, the adenosine deaminase protein can either be covalently linked to the CRISPR-Cas enzyme or be provided as a separate protein, but adapted so as to ensure recruitment thereof to the CRISPR-Cas complex.

[0304] In particular embodiments, recruitment of the adenosine deaminase to the target locus is ensured by fusing the adenosine deaminase or catalytic domain thereof to the CRISPR-Cas protein, which is a Cas or Cpf1 protein.

Methods of generating a fusion protein from two separate proteins are known in the art and typically involve the use of spacers or linkers. The CRISPR-Cas protein can be fused to the adenosine deaminase protein or catalytic domain thereof on either the N- or C-terminal end thereof. In particular embodiments, the CRISPR-Cas protein is a Cas or Cpf1 protein and is linked to the N-terminus of the deaminase protein or its catalytic domain.

[0305] In the present disclosure, linker refers to a molecule which joins two entities such as two domains to form a fusion protein. Generally, such molecules have no specific biological activity other than to join or to preserve some minimum distance or other spatial relationship between the proteins. However, in certain embodiments, the linker may be selected to influence some property of the linker and/or the fusion protein such as the folding, net charge, or hydrophobicity of the linker.

[0306] Suitable linkers for use in the methods of the present disclosure are well known to those of skill in the art and include, but are not limited to, straight or branched-chain carbon linkers, heterocyclic carbon linkers, or peptide linkers. However, as used herein the linker may also be a covalent bond (carbon-carbon bond or carbon-heteroatom bond). In particular embodiments, the linker is used to separate the CRISPR-Cas protein and the cytidine deaminase by a distance sufficient to ensure that each protein retains its required functional property. Preferred peptide linker sequences adopt a flexible extended conformation and do not exhibit a propensity for developing an ordered secondary structure.

[0307] In some embodiments, the linker can be a chemical moiety which can be monomeric, dimeric, multimeric or polymeric. Exemplary linkers are disclosed in Maratea et al. (1985), Gene 40: 39-46; Murphy et al. (1986) Proc. Nat'l. Acad. Sci. USA 83: 8258-62; U.S. Pat. Nos. 4,935,233; and 4,751,180. For example, GlySer linkers GGS, GGGS or GSG can be used. GGS, GSG, GGGS or GGGGS linkers can be used in repeats of 3 such as (GGS)₃. In other embodiments, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12 or more of (GGGS), to provide suitable lengths. In some embodiments, linkers such as (GGGS)₁, (GGGS)₂, (GGGS)₃, (GGGS)₄, (GGGS)₅, (GGGS)₆, (GGGS)₇, (GGGS)₈, (GGGS)₉, (GGGS)₁₀, (GGGS)₁₁, or (GGGS)₁₂ may be used. In other embodiments, the linker is XTEN linker. In particular embodiments, the nuclease-inactivated CRISPR-associated effector protein such as dCas9 or nCas9 is linked to the deaminase protein or its catalytic domain by means of an XTEN linker. In some embodiments, the nuclease-inactivated Cas mutant is linked C-terminally to the N-terminus of a deaminase protein or its catalytic domain by means of an XTEN linker. In addition, N- and C-terminal NLSs can also function as linker.

[0308] The present disclosure provides guide RNAs (gRNAs) that direct a site-directed modifying polypeptide to a specific target nucleic acid sequence. A gRNA comprises a nucleic acid-targeting segment and protein-binding segment. The nucleic acid-targeting segment of a gRNA comprises a nucleotide sequence that is complementary to a sequence in the target nucleic acid sequence. As such, the nucleic acid-targeting segment of a gRNA interacts with a target nucleic acid in a sequence-specific manner via hybridization (i.e., base pairing), and the nucleotide sequence of the nucleic acid-targeting segment determines the location within the target nucleic acid that the gRNA will bind. The nucleic

acid-targeting segment of a gRNA can be modified (e.g., by genetic engineering) to hybridize to any desired sequence within a target nucleic acid sequence.

[0309] The protein-binding segment of a guide RNA interacts with a site-directed modifying polypeptide (e.g. a Cas protein) to form a complex. The guide RNA guides the bound polypeptide to a specific nucleotide sequence within target nucleic acid via the above-described nucleic acid-targeting segment. The protein-binding segment of a guide RNA comprises two stretches of nucleotides that are complementary to one another and which form a double stranded RNA duplex.

[0310] In some embodiments, a gRNA comprises two separate RNA molecules. In such embodiments, each of the two RNA molecules comprises a stretch of nucleotides that are complementary to one another such that the complementary nucleotides of the two RNA molecules hybridize to form the double-stranded RNA duplex of the protein-binding segment. In some embodiments, a gRNA comprises a single guide RNA molecule (sgRNA).

[0311] The specificity of a gRNA for a target loci is mediated by the sequence of the nucleic acid-binding segment, which comprises about 20 nucleotides that are complementary to a target nucleic acid sequence within the target locus. In some embodiments, the corresponding target nucleic acid sequence is approximately 20 nucleotides in length. In some embodiments, the nucleic acid-binding segments of the gRNA sequences of the present disclosure are at least 90% complementary to a target nucleic acid sequence within a target locus. In some embodiments, the nucleic acid-binding segments of the gRNA sequences of the present disclosure are at least 95%, 96%, 97%, 98%, or 99% complementary to a target nucleic acid sequence within a target locus. In some embodiments, the nucleic acid-binding segments of the gRNA sequences of the present disclosure are 100% complementary to a target nucleic acid sequence within a target locus.

[0312] In some embodiments, the target nucleic acid sequence is an RNA target sequence. In some embodiments, the target nucleic acid sequence is a DNA target sequence.

[0313] In some embodiments, the targeted gene-editing system comprises two or more gRNA molecules each comprising a DNA-binding segment, wherein at least one of the nucleic acid-binding segments binds to a target DNA sequence of a target gene.

[0314] In some embodiments, the guide RNA is a single guide RNA (sgRNA). Methods of constructing suitable sgRNAs according to a given target sequence are known in the art. See e.g., Wang, Y. et al. Simultaneous editing of three homoeoalleles in hexaploid bread wheat confers heritable resistance to powdery mildew. *Nat. Biotechnol.* 32, 947-951 (2014); Shan, Q. et al. Targeted genome modification of crop plants using a CRISPR-Cas system. *Nat. Biotechnol.* 31, 686-688 (2013); Liang, Z. et al. Targeted mutagenesis in *Zea mays* using TALENs and the CRISPR/Cas system. *J Genet Genomics.* 41, 63-68 (2014).

[0315] The addition of a uracil DNA glycosylase (UGI) domain further increased the base-editing efficiency. In some embodiments, the targeted base-editing system further comprises a base excision repair (BER) inhibitor. Cellular DNA-repair response to the presence of a U:G pairing in DNA may be responsible for a decrease in nucleobase editing efficiency in plant cells. Uracil DNA glycosylase catalyzes removal of uracil from DNA in plant cells, which may

initiate base excision repair, such that the U:G pair is reversed to C:G. In some embodiments, the BER inhibitor is an uracil glycosylase inhibitor or an active domain thereof.

[0316] In some embodiments, the BER inhibitor is an inhibitor of uracil DNA glycosylase (UDG). In some embodiments, the BER inhibitor is an inhibitor of UDG. In some embodiments, the BER inhibitor is a polypeptide inhibitor. In some embodiments, the BER inhibitor is a protein that binds single-stranded DNA. For example, the BER inhibitor may be a *Erwinia tasmaniensis* single-stranded binding protein. In some embodiments, the BER inhibitor is a protein that binds uracil. In some embodiments, the BER inhibitor is a protein that binds uracil in DNA. In some embodiments, the BER inhibitor is a catalytically inactive UDG or binding domain thereof. In some embodiments, the BER inhibitor is a catalytically inactive UDG or binding domain thereof that does not excise uracil from the DNA. Other proteins that are capable of inhibiting (e.g., sterically blocking) UDG are within the scope of this disclosure. Additionally, any proteins that block or inhibit base-excision repair as also within the scope of this disclosure.

[0317] Base excision repair may be inhibited by molecules that bind the edited strand, block the edited base, inhibit uracil DNA glycosylase, inhibit base excision repair, protect the edited base, and/or promote fixing of the non-targeted strand. Accordingly, the use of the BER inhibitor described herein can increase the editing efficiency of a cytidine deaminase that is capable of catalyzing a C to U change.

[0318] In particular embodiments, the uracil glycosylase inhibitor (UGI) is the uracil DNA glycosylase inhibitor of *Bacillus subtilis* bacteriophage PBS1 or an active fragment thereof, such as an 83 residue protein of *Bacillus subtilis* bacteriophage PBS1.

[0319] Suitable UGI protein and nucleotide sequences are provided herein and additional suitable UGI sequences are known to those in the art, and include, for example, those published in Wang et al., Uracil-DNA glycosylase inhibitor gene of bacteriophage PBS2 encodes a binding protein specific for uracil-DNA glycosylase. *J. Biol. Chem.* 264: 1 163-1 171(1989); Lundquist et al., Site-directed mutagenesis and characterization of uracil-DNA glycosylase inhibitor protein. Role of specific carboxylic amino acids in complex formation with *Escherichia coli* uracil-DNA glycosylase. *J. Biol. Chem.* 272:21408-21419(1997); Ravishankar et al., X-ray analysis of a complex of *Escherichia coli* uracil DNA glycosylase (EcUDG) with a proteinaceous inhibitor. The structure elucidation of a prokaryotic UDG. *Nucleic Acids Res.* 26:4880-4887(1998); and Putnam et al., Protein mimicry of DNA from crystal structures of the uracil-DNA glycosylase inhibitor protein and its complex with *Escherichia coli* uracil-DNA glycosylase. *J. Mol. Biol.* 287:331-346(1999), each of which incorporated herein by reference. Additional embodiments of the uracil glycosylase inhibitor (UGI) are disclosed in WO2018/086623, WO2018/205995, WO2017/70632, and WO2018/213726, each of which is incorporated herein by reference in its entirety.

[0320] In some embodiments, the UGI domain comprises a wild type UGI or a UGI. In some embodiments, the UGI proteins provided herein include fragments of UGI and proteins homologous to a UGI or a UGI fragment.

[0321] Additional proteins may be uracil glycosylase inhibitors. For example, other proteins that are capable of inhibiting (e.g., sterically blocking) a uracil-DNA glycosylase

lase base-excision repair enzyme are within the scope of this disclosure. Additionally, any proteins that block or inhibit base-excision repair as also within the scope of this disclosure. In some embodiments, a protein that binds DNA is used. In another embodiment, a substitute for UGI is used. In some embodiments, a uracil glycosylase inhibitor is a protein that binds single-stranded DNA. For example, a uracil glycosylase inhibitor may be a *Erwinia tasmaniensis* single-stranded binding protein. In some embodiments, a uracil glycosylase inhibitor is a protein that binds uracil. In some embodiments, a uracil glycosylase inhibitor is a protein that binds uracil in DNA. In some embodiments, a uracil glycosylase inhibitor is a catalytically inactive uracil DNA-glycosylase protein. In some embodiments, a uracil glycosylase inhibitor is a catalytically inactive uracil DNA-glycosylase protein that does not excise uracil from the DNA. As another example, a uracil glycosylase inhibitor is a catalytically inactive UDG.

[0322] In some embodiments, the base editing system comprises the following domains; i) the CRISPR-Cas protein (dCas9 or nCas9) and ii) the cytidine deaminase, which can be fused to or linked to a BER inhibitor (e.g., an inhibitor of uracil DNA glycosylase).

[0323] Uracil-DNA glycosylase (UDG) is an enzyme that reverts mutations in DNA. The most common mutation is the deamination of cytosine to uracil. UDG repairs these mutations and UDG is crucial in DNA repair. Various uracil-DNA glycosylases and related DNA glycosylases (EC) are present such as uracil-DNA glycosylase, thermophilic uracil-DNA glycosylase, G:T/U mismatch-specific DNA glycosylase (Mug), and single-strand selective monofunctional uracil-DNA glycosylase (SMUG1).

[0324] Uracil DNA glycosylases remove uracil from DNA, which can arise either by spontaneous deamination of cytosine or by the misincorporation of dU opposite dA during DNA replication. The prototypical member of this family is *E. coli* UDG, which was among the first glycosylases discovered. Four different uracil-DNA glycosylase activities have been identified in mammalian cells, including UNG, SMUG1, TDG, and MBD4, which vary in substrate specificity and subcellular localization. SMUG1 prefers single-stranded DNA as substrate, but also removes U from double-stranded DNA. In addition to unmodified uracil, SMUG1 can excise 5-hydroxyuracil, 5-hydroxymethyluracil and 5-formyluracil bearing an oxidized group at ring C5. [13] TDG and MBD4 are strictly specific for double-stranded DNA. TDG can remove thymine glycol when present opposite guanine, as well as derivatives of U with modifications at carbon 5.

[0325] TDG and SMUG1 are the major enzymes responsible for the repair of the U:G mispairs caused by spontaneous cytosine deamination, whereas uracil arising in DNA through dU misincorporation is mainly dealt with by UNG. MBD4 is thought to correct T:G mismatches that arise from deamination of 5-methylcytosine to thymine in CpG sites.

[0326] Uracil arising in DNA either from misincorporation of dUMP or from deamination of cytosine is actively removed through the multistep base excision repair (BER) pathway. BER of uracil is initiated by a uracil DNA glycosylase (UDG) activity that cleaves the N-glycosidic bond and excises uracil as a free base, generating an abasic (apurinic/apyrimidinic, AP) site in the DNA. Repair is completed through subsequent steps that include incision at the AP site, gap tailoring, repair synthesis, and ligation.

[0327] In some embodiments, the addition of a Uracil-DNA glycosylase (UDG) such as uracil N-glycosylase (UNG) can induce various mutations at targeted base. In some embodiments, the targeted base-editing system further comprises a Uracil-DNA glycosylase (UDG). Cellular DNA-repair response to the presence of a U:G pairing in DNA may be responsible for a decrease in nucleobase editing efficiency in plant cells. Uracil DNA glycosylase catalyzes removal of uracil from DNA in plant cells, which may initiate base excision repair, such that the U:G pair is reversed to C:G. In other embodiments, removal of uracil from DNA in plant cells are not always reversed to C for C:G pairing, but randomized to other bases such as T, A, and G.

[0328] In some embodiments, a Uracil-DNA glycosylase (UDG) is fused to the targeted base editing system taught in the present disclosure to introduce a stable and targeted, but randomized single nucleotide substitution in a target gene of interest.

[0329] The use of the UDG described herein can increase the base randomization in a targeted single nucleotide of a target gene.

[0330] In some embodiments, a UDG is provided in cis. In some embodiments, a UDG is provided in trans. In some embodiments, a UDG is fused to a base editor (or a base editing system) described in the present disclosure. In other embodiments, a UDG trigger a stall DNA replication for base randomization. In other embodiments, a UDG triggers a DNA repair through DNA replication, thereby including base randomization. In further embodiments, naturally occurring UDG variants can be used as a UDG domain. In further embodiments, non-naturally occurring UDG variants can be used as a UDG domain. In further embodiments, a UDG can be genetically engineered to enhance a functional UDG activity.

[0331] A nuclear localization signal (NLS), or any other organelle targeting signal, can be further required to ensure proper targeting of the complex. The present disclosure relate to modifying an cytosine in a target locus of interest, whereby the target locus is within a plant cell. In order to improve targeting of the CRISPR-Cas protein and/or the cytidine deaminase protein or catalytic domain thereof used in the methods of the present disclosure to the nucleus, it may be advantageous to provide one or both of these components with one or more nuclear localization sequences (NLSs).

[0332] In some embodiments, the NLSs used in the context are heterologous to the proteins. In general, NLS consists of one or more short sequences of positively charged lysine or arginine exposed on the surface of a protein, but other types of NLS are also known in the art.

[0333] In some embodiments, the N-terminus of the gene editing fusion protein comprises an NLS with an amino acid sequence. In some embodiments, the C-terminus of the gene-editing fusion protein comprises an NLS.

[0334] In addition, the gene editing fusion protein may also include other localization sequences, such as cytoplasmic localization sequences, chloroplast localization sequences, mitochondrial localization sequences, and the like, depending on the location of the DNA to be edited. In order to obtain efficient expression in plants, in some embodiments, the nucleotide sequence encoding the gene editing fusion protein is codon optimized for the plant to be gene edited.

[0335] Codon optimization refers to a process of modifying a nucleic acid sequence for enhanced expression in the host cells of interest by replacing at least one codon (e.g. about or more than about 1, 2, 3, 4, 5, 10, 15, 20, 25, 50, or more codons) of the native sequence with codons that are more frequently or most frequently used in the genes of that host cell while maintaining the native amino acid sequence. Various species exhibit particular bias for certain codons of a particular amino acid. Codon bias (differences in codon usage between organisms) often correlates with the efficiency of translation of messenger RNA (mRNA), which is in turn believed to be dependent on, among other things, the properties of the codons being translated and the availability of particular transfer RNA (tRNA) molecules. The predominance of selected tRNAs in a cell is generally a reflection of the codons used most frequently in peptide synthesis. Accordingly, genes can be tailored for optimal gene expression in a given organism based on codon optimization. Codon usage tables are readily available, for example, at the “Codon Usage Database” available at www.kazusa.or.jp/codon/ and these tables can be adapted in a number of ways. See Nakamura, Y., et al. “Codon usage tabulated from the international DNA sequence databases: status for the year 2000” *Nucl. Acids Res.* 28:292 (2000).

[0336] In some embodiments, the codon-optimized nucleotide sequence encoding the gene editing fusion protein is provided herein. In some embodiments, the guide RNA is a single guide RNA (sgRNA). Methods of constructing suitable sgRNAs according to a given target sequence are known in the art. See e.g., Wang, Y. et al. Simultaneous editing of three homeoalleles in hexaploid bread wheat confers heritable resistance to powdery mildew. *Nat. Biotechnol.* 32, 947-951 (2014); Shan, Q. et al. Targeted genome modification of crop plants using a CRISPR-Cas system. *Nat. Biotechnol.* 31, 686-688 (2013); Liang, Z. et al. Targeted mutagenesis in *Zea mays* using TALENs and the CRISPR/Cas system. *J Genet Genomics.* 41, 63-68 (2014).

[0337] In order to ensure appropriate expression in a plant cell, the components of the targeted gene-editing system described herein are typically placed under control of a plant promoter, i.e. a promoter operable in plant cells. The use of different types of promoters is envisaged.

[0338] A constitutive plant promoter is a promoter that is able to express the open reading frame (ORF) that it controls in all or nearly all of the plant tissues during all or nearly all developmental stages of the plant (referred to as “constitutive expression”). One non-limiting example of a constitutive promoter is the cauliflower mosaic virus 35S promoter. “Regulated promoter” refers to promoters that direct gene expression not constitutively, but in a temporally- and/or spatially-regulated manner, and includes tissue-specific, tissue-preferred and inducible promoters. Different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental conditions. In particular embodiments, one or more of the components of the targeted gene-editing system described herein are expressed under the control of a constitutive promoter, such as the cauliflower mosaic virus 35S promoter. Tissue-preferred promoters can be utilized to target enhanced expression in certain cell types within a particular plant tissue, for instance vascular cells in leaves or roots or in specific cells of the seed. Examples of particular promoters for use of the present disclosure can be found in Kawamata et al., (1997) *Plant*

Cell Physiol 38:792-803; Yamamoto et al., (1997) *Plant J* 12:255-65; Hire et al, (1992) *Plant Mol Biol* 20:207-18, Kuster et al, (1995) *Plant Mol Biol* 29:759-72, and Capana et al., (1994) *Plant Mol Biol* 25:681-91.

[0339] Inducible promoters can be of interest to express one or more of the components of the targeted gene-editing system described herein under limited circumstances to avoid non-specific activity of the deaminase. In particular embodiments, one or more elements of the targeted gene-editing system described herein are expressed under control of an inducible promoter.

[0340] Examples of promoters that are inducible and that allow for spatiotemporal control of gene editing or gene expression may use a form of energy. The form of energy may include but is not limited to sound energy, electromagnetic radiation, chemical energy and/or thermal energy. Examples of inducible systems include tetracycline inducible promoters (Tet-On or Tet-Off), small molecule two-hybrid transcription activations systems (FKBP, ABA, etc), or light inducible systems (Phytochrome, LOV domains, or cryptochrome), such as a Light Inducible Transcriptional Effector (LITE) that direct changes in transcriptional activity in a sequence-specific manner. The components of a light inducible system may include a fusion protein of the targeted gene-editing system and a light-responsive cytochrome heterodimer (e.g. from *Arabidopsis thaliana*). Further examples of inducible DNA binding proteins and methods for their use are provided in U.S. 61/736,465 and U.S. 61/721,283, which is hereby incorporated by reference in its entirety.

[0341] In particular embodiments, transient or inducible expression can be achieved by using, for example, chemical-regulated promoters, i.e. whereby the application of an exogenous chemical induces gene expression. Modulating of gene expression can also be obtained by a chemical-repressible promoter, where application of the chemical represses gene expression. Chemical-inducible promoters include, but are not limited to, the maize In2-2 promoter, activated by benzene sulfonamide herbicide safeners (De Veylder et al., (1997) *Plant Cell Physiol* 38:568-77), the maize GST promoter (GST-11-27, WO93/01294), activated by hydrophobic electrophilic compounds used as pre-emergent herbicides, and the tobacco PR-1 promoter (Ono et al., (2004) *Biosci Biotechnol Biochem* 68:803-7) activated by salicylic acid. Promoters which are regulated by antibiotics, such as tetracycline-inducible and tetracycline-repressible promoters (Gatz et al., (1991) *Mol Gen Genet* 227: 229-37; U.S. Pat. Nos. 5,814,618 and 5,789,156) can also be used herein.

[0342] In some embodiments, the nucleotide sequence encoding the gene-edited fusion protein and/or the nucleotide sequence encoding the guide RNA is operably linked to a plant expression regulatory element, such as a promoter. Examples of promoters that can be used in the present disclosure include, but are not limited to the cauliflower mosaic virus 35S promoter (Odell et al. (1985) *Nature* 313: 810-812), a maize Ubi-1 promoter, a wheat U6 promoter, a rice U3 promoter, a maize U3 promoter, a rice actin promoter, a TrpPro5 promoter (U.S. patent application Ser. No. 10/377,318; filed on Mar. 16, 2005), a pEMU promoter (Last et al. *Theor. Appl. Genet.* 8 1: 581-588), a MAS promoter (Velten et al. (1984) *EMBO J.* 3:2723-2730), a maize H3 histone promoter (Lepetit et al. *Mol. Gen. Genet.* 231:276-285 and Atanassova et al. (1992) *Plant J.* 2 (3): 291-300), and a *Brassica napus* ALS3 (PCT Application WO

97/41228) promoters. Promoters that can be used in the present disclosure also include the commonly used tissue specific promoters as reviewed in Moore et al. (2006) *Plant J.* 45 (4): 651-683.

[0343] In the present disclosure, the plants are intended to comprise without limitation angiosperm and gymnosperm plants such as acacia, alfalfa, amaranth, apple, apricot, artichoke, ash tree, asparagus, avocado, banana, barley, beans, beet, birch, beech, blackberry, black raspberry, blueberry, broccoli, Brussel's sprouts, cabbage, cane berry, canola, cantaloupe, carrot, cassava, cauliflower, cedar, a cereal, celery, chestnut, cherry, Chinese cabbage, citrus, Clementine, clover, coffee, corn, cotton, cowpea, cucumber, cypress, eggplant, elm, endive, *eucalyptus*, fennel, figs, fir, geranium, grape, grapefruit, groundnuts, ground cherry, gum hemlock, hickory, kale, kiwifruit, kohlrabi, larch, lettuce, leek, lemon, lime, locust, pine, maidenhair, maize, mango, maple, melon, millet, mushroom, mustard, nuts, oak, oats, oil palm, okra, onion, orange, an ornamental plant or flower or tree, *papaya*, palm, parsley, parsnip, pea, peach, peanut, pear, peat, pepper, persimmon, pigeon pea, peach, pine, pineapple, plantain, plum, pomegranate, potato, pumpkin, radicchio, radish, rapeseed, raspberry, rice, rye, sorghum, safflower, sallow, soybean, spinach, spruce, squash, strawberry, sugar beet, sugarcane, sunflower, sweet potato, sweet corn, tangerine, tea, tobacco, tomato, trees, triticale, turf grasses, turnips, vine, walnut, watercress, watermelon, wheat, wild strawberry, yams, yew, and zucchini.

[0344] The methods for targeted gene-editing system as described herein can be used to confer desired traits on essentially any plant. A wide variety of plants and plant cell systems may be engineered for the desired physiological and agronomic characteristics described herein using the nucleic acid constructs of the present disclosure and the various transformation methods. In preferred embodiments, target plants and plant cells for engineering include, but are not limited to, those monocotyledonous and dicotyledonous plants, such as crops including grain crops (e.g., wheat, maize, rice, millet, barley), fruit crops (e.g., tomato, apple, grape, peach, pear, plum, raspberry, black raspberry, blackberry, cane berry, cherry, avocado, strawberry, wild strawberry, orange), forage crops (e.g., alfalfa), root vegetable crops (e.g., carrot, potato, sugar beets, yam), leafy vegetable crops (e.g., lettuce, spinach); flowering plants (e.g., *petunia*, rose, *chrysanthemum*), conifers and pine trees (e.g., pine fir, spruce); plants used in phytoremediation (e.g., heavy metal accumulating plants); oil crops (e.g., sunflower, rape seed) and plants used for experimental purposes (e.g., *Arabidopsis*). In some embodiments, fruit crops such as tomato, apple, peach, pear, plum, raspberry, black raspberry, blackberry, cane berry, cherry, avocado, strawberry, wild strawberry, grape and orange.

[0345] The present disclosure provides methods for targeted editing in a plant cell, tissue, organ or plant. In one aspect, the present disclosure provides methods for producing a gene-edited plant by transient expression of a gene editing system. In another aspect, the present disclosure provides methods for a gene-edited and non-transgenic plant by transient expression of a gene editing system along with transient expression of at least one morphogenic regulator taught in the present disclosure. Therefore, the present disclosure teaches a gene-edited plant that is also transgene-free without any integration of foreign DNAs.

[0346] In some embodiments, an editing system as described herein is used to introduce targeted mutations, such as insertions, deletions, or substitutions, thereby causing a nonsense mutation (e.g., premature stop codon) or a missense mutation (e.g., encoding different amino acid residue). This is of interest where the single nucleotide mutations in certain endogenous genes can confer or contribute to a desired trait.

[0347] The methods described herein may result in the generation of gene-edited plants that have one or more desirable traits compared to the wild type plant.

[0348] In some embodiments, non-transgenic but gene-edited plants, plant parts, or cells are obtained, in that no exogenous DNA sequence is incorporated into the genome of any of the plant cells of the plant. In such embodiments, the gene-edited plants are non-transgenic. Where only the modification of an endogenous gene is ensured and no foreign genes are introduced or maintained in the plant genome, the resulting genetically modified crops contain no foreign genes and can thus basically be considered non-transgenic.

[0349] In some embodiments, modification of the target sequence can be accomplished simply by introducing or producing the gene editing protein and guide RNA in plant cells, and the modification can be stably inherited without the need of stable transformation of plants with an editing system. This avoids potential off-target effects of a stable editing system, and also avoids the integration of exogenous nucleotide sequences into the plant genome, and thereby resulting in higher biosafety. In some embodiments, the editing system is the targeted gene editing system. In further embodiments, the plant cells in which the genetic modification is stably inherited without stable transformation is edited by the gene editing system (i.e. the targeted gene editing system) are transformed and regenerated into a whole plant by the transient expression of one or more morphogenic regulators.

[0350] In other embodiments, the polynucleotides are delivered into the cell by a DNA virus (e.g., a geminivirus) or an RNA virus (e.g., a tobnavirus). In other embodiments, the introducing steps include delivering to the plant cell a T-DNA containing one or more polynucleotide sequences encoding an encoding system (e.g., a CRISPR-Cas effector protein and a guide nucleic acid), where the delivering is via *Agrobacterium*. The polynucleotide sequence encoding the components of CRISPR/Cas system can be operably linked to a promoter, such as a constitutive promoter (e.g., a cauliflower mosaic virus 35S promoter), or a cell specific or inducible promoter in a plant cell of interest described herein. In other embodiments, the polynucleotide is introduced by microprojectile bombardment.

[0351] In some embodiments, the gene editing system can be introduced into plants by various methods well known to people skilled in the art. Methods that can be used to introduce the gene editing system of the present disclosure into plants include but not limited to particle bombardment, PEG-mediated protoplast transformation, *Agrobacterium*-mediated transformation, plant virus-mediated transformation, pollen tube approach, and ovary injection approach.

[0352] In some embodiments, the introduction is performed in the absence of a selective pressure, thereby avoiding the integration of exogenous nucleotide sequences in the plant genome. In some embodiments, the introduction comprises transforming the gene editing system into isolated

plant cells or tissues, and then regenerating the transformed plant cells or tissues into an intact plant by the help of morphogenic regulator expression taught in the present disclosure. Preferably, the regeneration is performed in the absence of a selective pressure, i.e., no selective agent against the selective gene carried on the expression vector is used during the tissue culture. Without the use of a selective agent, the regeneration efficiency of the plant can be increased to obtain a modified plant that does not contain exogenous nucleotide sequences. To increase, enhance, improve the transformation and regeneration of a non-transgenic plant with a desired modification inherited, the morphogenic regulator(s) is transiently expressed in the plant cell.

[0353] In other embodiments, the editing system of the present disclosure can be transformed to a particular site on an intact plant, such as leaf, shoot tip, pollen tube, young ear, or hypocotyl. This is particularly suitable for the transformation of plants that are difficult to regenerate by tissue culture. In some embodiments, proteins expressed in vitro and/or RNA molecules transcribed in vitro are directly transformed into the plant. The proteins and/or RNA molecules are capable of achieving gene-editing in plant cells, and are subsequently degraded by the cells to avoid the integration of exogenous nucleotide sequences into the plant genome. Plants that can be gene-edited by the methods includes monocotyledon and dicotyledon. For example, the plant may be a crop plant such as wheat, rice, maize, soybean, sunflower, sorghum, rape, alfalfa, cotton, barley, millet, sugar cane, tomato, tobacco, cassava, or potato. For another example, the plant may be a fruit crops such as tomato, apple, peach, pear, plum, raspberry, black raspberry, blackberry, cane berry, cherry, avocado, strawberry, wild strawberry, grape and orange.

[0354] In some embodiments, the target sequence is associated with plant traits such as desired traits, and thereby the gene editing results in the plant having altered traits relative to a wild type plant. In the present disclosure, the target sequence to be modified (e.g., a target nucleic acid) may be located anywhere in the genome, for example, within a functional gene such as a protein-coding gene or, for example, may be located in a gene expression regulatory region such as a promoter region or an enhancer region, and thereby accomplish the functional modification of said gene or accomplish the modification of a gene expression.

[0355] In some embodiments, the method further comprises obtaining progeny of the gene-edited plant, which is also non-transgenic. In a further aspect, the disclosure also provides a gene-edited plant or progeny thereof or parts thereof, wherein the plant is obtained by the method described above.

[0356] In another aspect, the present disclosure also provides a plant breeding method comprising crossing a first gene-edited plant obtained by the above-mentioned method of the present disclosure with a second plant not containing said genetic modification, thereby introducing said genetic modification into said second plant.

[0357] In particular embodiments, the method further includes screening the plant cell after the introducing steps to determine whether the expression of the gene of interest has been modified. In particular embodiments, the methods include the step of regenerating a plant from the plant cell. In further embodiments, the methods include cross breeding the plant to obtain a genetically desired plant lineage.

[0358] In some embodiments, significant impact of transient expression of morphogenic regulators on plant transformation and regeneration can be applied for generation of gene-edited plant without foreign DNA integration. The regenerated plant from a gene-edited plant cell is transgene-free and/or DNA integration-free. Transient expression of at least one morphogenic regulators gives rise to successful regeneration of a plant cell of interest.

[0359] In other embodiments, significant impact of transient expression of morphogenic regulators on plant transformation and regeneration can be applied for plant regeneration of dicot species, especially those with heterozygotic nature and/or polyploidy.

[0360] In some embodiments, transient expression of morphogenic regulators can be used to regenerate plants from protoplast cells. In other embodiments, transient expression of morphogenic regulator can be used to regenerate plants from edited callus by the gene-editing system taught in the present disclosure.

[0361] As described herein, the nucleic acids of the invention and/or expression cassettes and/or vectors comprising the same may be codon optimized for expression in an organism. An organism useful with this invention may be any organism or cell thereof for which a morphogenic regulator and/or nucleic acid modification may be useful. An organism can include, but is not limited to, any animal, any plant, any fungus, any archaeon, or any bacterium. In some embodiments, the organism may be a plant or cell thereof.

[0362] A target nucleic acid of any cell, plant, or plant part may be modified using a composition, system, and/or method of the invention. Any plant (or groupings of plants, for example, into a genus or higher order classification) may be modified using a nucleic acid construct, composition, system, and/or method of this invention including an angiosperm, a gymnosperm, a monocot, a dicot, a C3, C4, CAM plant, a bryophyte, a fern and/or fern ally, a microalgae, and/or a macroalgae. A plant and/or plant part useful with this invention may be a plant and/or plant part of any plant species/variety/cultivar. The term "plant part," as used herein, includes but is not limited to, embryos, pollen, ovules, seeds, leaves, stems, shoots, flowers, branches, fruit, kernels, ears, cobs, husks, stalks, roots, root tips, anthers, plant cells including plant cells that are intact in plants and/or parts of plants, plant protoplasts, plant tissues, plant cell tissue cultures, plant calli, plant clumps, and the like. As used herein, "shoot" refers to the above ground parts including the leaves and stems. Further, as used herein, "plant cell" refers to a structural and physiological unit of the plant, which comprises a cell wall and also may refer to a protoplast. A plant cell can be in the form of an isolated single cell or can be a cultured cell or can be a part of a higher-organized unit such as, for example, a plant tissue or a plant organ.

[0363] Non-limiting examples of plants useful with the present invention include turf grasses (e.g., bluegrass, bentgrass, ryegrass, fescue), feather reed grass, tufted hair grass, *miscanthus*, *arundo*, switchgrass, vegetable crops, including artichokes, kohlrabi, arugula, leeks, asparagus, lettuce (e.g., head, leaf, romaine), malanga, melons (e.g., muskmelon, watermelon, crenshaw, honeydew, cantaloupe), cole crops (e.g., brussels sprouts, cabbage, cauliflower, broccoli, collards, kale, chinese cabbage, bok choy), cardoni, carrots, napa, okra, onions, celery, parsley, chick peas, parsnips, chicory, peppers, potatoes, cucurbits (e.g., marrow, cucum-

ber, zucchini, squash, pumpkin, honeydew melon, watermelon, cantaloupe), radishes, dry bulb onions, rutabaga, eggplant, salsify, escarole, shallots, endive, garlic, spinach, green onions, squash, greens, beet (sugar beet and fodder beet), sweet potatoes, chard, horseradish, tomatoes, turnips, and spices; a fruit crop such as apples, apricots, cherries, nectarines, peaches, pears, plums, prunes, cherry, quince, fig, nuts (e.g., chestnuts, pecans, pistachios, hazelnuts, pistachios, peanuts, walnuts, macadamia nuts, almonds, and the like), citrus (e.g., clementine, kumquat, orange, grapefruit, tangerine, mandarin, lemon, lime, and the like), blueberries, black raspberries, boysenberries, cranberries, currants, gooseberries, loganberries, raspberries, strawberries, blackberries, grapes (wine and table), avocados, bananas, kiwi, persimmons, pomegranate, pineapple, tropical fruits, pomes, melon, mango, *papaya*, and lychee, a field crop plant such as clover, alfalfa, timothy, evening primrose, meadow foam, corn/maize (field, sweet, popcorn), hops, jojoba, buckwheat, safflower, *quinoa*, wheat, rice, barley, rye, millet, sorghum, oats, triticale, sorghum, tobacco, kapok, a leguminous plant (beans (e.g., green and dried), lentils, peas, soybeans), an oil plant (rape, canola, mustard, poppy, olive, sunflower, coconut, castor oil plant, cocoa bean, groundnut, oil palm), duckweed, *Arabidopsis*, a fiber plant (cotton, flax, hemp, jute), *Cannabis* (e.g., *Cannabis sativa*, *Cannabis indica*, and *Cannabis ruderalis*), lauraceae (cinnamon, camphor), or a plant such as coffee, sugar cane, tea, and natural rubber plants; and/or a bedding plant such as a flowering plant, a cactus, a succulent and/or an ornamental plant (e.g., roses, tulips, violets), as well as trees such as forest trees (broad-leaved trees and evergreens, such as conifers; e.g., elm, ash, oak, maple, fir, spruce, cedar, pine, birch, cypress, *eucalyptus*, willow), as well as shrubs and other nursery stock. In some embodiments, the nucleic acid constructs of the invention and/or expression cassettes and/or vectors encoding the same may be used to modify maize, soybean, wheat, canola, rice, tomato, pepper, sunflower, raspberry, blackberry, black raspberry and/or cherry.

[0364] In some embodiments, a plant cell of the present invention and/or that is used in a composition, system and/or method of the present invention is a plant cell is from an asexually propagated, heterozygous, perennial plant. In some embodiments, a plant cell of the present invention and/or that is used in a composition, system and/or method of the present invention is a dicot plant cell, optionally wherein the plant cell is a blackberry, raspberry, or cherry plant cell.

[0365] In some embodiments, the invention provides cells (e.g., plant cells, animal cells, bacterial cells, archaeon cells, and the like) comprising a polypeptide, polynucleotide, nucleic acid construct, expression cassette and/or vector of the invention.

[0366] The present invention further comprises a kit or kits to carry out the methods of this invention. A kit of this invention can comprise reagents, buffers, and apparatus for mixing, measuring, sorting, labeling, etc, as well as instructions and the like as would be appropriate for modifying a target nucleic acid.

[0367] In some embodiments, the invention provides a kit for comprising one or more nucleic acid construct(s) of the invention, expression cassette(s), vector(s), and/or cells comprising the same as described herein, with optional instructions for the use thereof. In some embodiments, a kit may further comprise an editing system (e.g., a CRISPR-Cas

guide nucleic acid (corresponding to a CRISPR-Cas effector protein encoded by a polynucleotide of the invention) and/or an expression cassette, vector, and/or cell comprising the same. Accordingly, in some embodiments, kits are provided comprising a nucleic acid construct comprising (a) a polynucleotide(s) as provided herein and (b) a promoter that drives expression of the polynucleotide(s) of (a). In some embodiments, the kit may further comprise a nucleic acid construct encoding a guide nucleic acid, wherein the construct comprises a cloning site for cloning of a nucleic acid sequence identical or complementary to a target nucleic acid sequence into backbone of the guide nucleic acid.

[0368] In some embodiments, the nucleic acid construct of the invention may be an mRNA that may encode one or more introns within the encoded polynucleotide(s). In some embodiments, the nucleic acid constructs of the invention, and/or an expression cassettes and/or vectors comprising the same, may further encode one or more selectable markers useful for identifying transformants (e.g., a nucleic acid encoding an antibiotic resistance gene, herbicide resistance gene, and the like).

[0369] The invention will now be described with reference to the following examples. It should be appreciated that these examples are not intended to limit the scope of the claims to the invention, but are rather intended to be exemplary of certain embodiments. Any variations in the exemplified methods that occur to the skilled artisan are intended to fall within the scope of the invention.

EXAMPLES

[0370] The following examples are given for the purpose of illustrating various embodiments of the disclosure and are not meant to limit the present disclosure in any fashion. Changes therein and other uses which are encompassed within the spirit of the disclosure, as defined by the scope of the claims, will occur to those skilled in the art.

Example 1. Transformation of Tobacco Leaves

[0371] Transformation can be accomplished by various methods known to be effective in plants, including particle-mediated delivery, *Agrobacterium*-mediated transformation, PEG-mediated delivery, and electroporation.

[0372] The tobacco leaves are prepared for transformation with plasmids containing the gene-editing system and/or plasmids containing at least one morphogenic regulator. The expression vector containing the morphogenic regulators such as WUS, BBN, KN1, IPT, CLV3, miR156, GRF5, NTT, and HDZipII is co-bombarded or co-transformed with another expression vector containing the targeted gene editing system, for example, CRISPR/Cas 9 endonuclease with gRNA.

[0373] The tobacco leaf transformation can be achieved by the simultaneous expression of the morphogenic regulator and the gene editing machinery after the vacuum infiltration of tobacco leaves with two populations of recombinant *Agrobacterium*.

Example 2. Transient Expression of Morphogenic Regulators

[0374] Parameters of the transformation protocol can be modified to ensure that the activity of morphogenic regulators (including, but are not limited to, WUS, BBN, KN1, IPT, CLV3, miR156, GRF5, NTT, and HDZipII) is transient.

[0375] One method for transient expression can involve precipitating the expression vector in a manner that allows for transcription and expression, but precludes subsequent release of the DNA, for example, by using the chemical polyethylenimine.

[0376] In this example, three expression vectors were constructed to test integration efficiency of a morphogenic regulator; 1) GUS marker gene driven by a constitutive promoter, 2) ipt gene fused with GUS marker gene under control of a native promoter, 3) ipt gene fused with GUS marker gene under control of a constitutive promoter. Each vector was transformed into tobacco leaves (*Nicotiana benthamiana*).

[0377] The effect of an isopentenyl transferase (ipt) gene expression under a constitutive promoter could clearly be seen. For example, ipt gene expression induced multiple shoot formation in a hormone-free media when compared to two controls (i) without ipt gene expressed as well as (ii) with ipt gene expressed under a native promoter. The ipt transformed leaf explants showed darker green color and enlargement at 2-4 wks and numerous shoots emerged as early as 4 weeks after inoculation on hormone-free and nonselective medium.

[0378] The ipt gene under a constitutive promoter produced significantly more plants that were regenerated in a hormone-free media than the control where a construct without ipt gene was used, indicating that ipt gene enhanced plant regeneration in tobacco.

[0379] Over 90% of the plants regenerated were GUS negative in the first experiment. As shown in FIG. 1, GUS assay results of one hundred six shoots induced by the ipt gene expression indicate that about 93.4% (99 out of 106 ipt-induced shoots) were GUS negative, suggesting either that the GUS-negative shoots would not be stably integrated but transiently expressed. However, transgenic nature has not been determined. Thus, molecular analyses are being performed to determine the transgenic nature of these GUS negative plants is in progress.

[0380] On the other hand, seven samples were GUS-positive, which may indicate stable integration of ipt-GUS fusion cassette. Furthermore, the GUS-positive ipt-induced shoots generated too many shoots that could be explained by too high level of cytokinin accumulation due to the constitutive ipt gene expression. Thus, the GUS positive samples are considered to be stably integrated into a tobacco genome.

[0381] Importantly, molecular analyses are being performed to determine the transgenic nature of these GUS negative plants is in progress.

Example 3. Constructions of Expression Vectors for Gene-Editing System and/or Morphogenic Regulators

[0382] In this example, the construction of expression vector contains a marker-free T-DNA. Especially, three constructs are being generated and tested for efficiency of gene-editing and/or plant regeneration.

[0383] 1) Locate both ipt (Morphogenic regulator; Morgen) and gene-editing machinery outside of T-DNA Left Border

[0384] 2) Locate editing machinery inside of T-DNA, but ipt/Morgen outside of T-DNA Left Border

[0385] 3) Locate both ipt/Morgen and editing machinery inside of T-DNA

[0386] An expression vector comprising 1) expression cassette for ipt gene and 2) expression cassette for gene editing system (CRISPR/Cas9 and gRNA) will be transformed into a plant cell of interest in order to generate mutations at the corresponding targeted sites. DNA delivery (such as recombinant DNA, plasmid DNA) for targeted mutagenesis and gene editing in plants are presented herewith.

[0387] The gene-editing expression vectors will be transformed into plants of interest such as: black raspberry, blackberry, cane berry, cherry, peach, avocado, strawberry, wild strawberry, apple, tomato, grape, and peach as well as model plants such as *Arabidopsis* and tobacco. The disclosure teaches all types of transformation methods, including using *agrobacterium*-mediated protocols that are known in the art and/or developed by the inventors, as well as biolistic transformation methods. Tissue culture and regeneration of transformed plants will be performed accordingly.

Example 4: Molecular Analysis for Transformed Plants

[0388] In this example, the activity of the gene-editing machinery will be examined with efficiency of plant transformation and regeneration by control of transient ipt gene expression on the transformed plants of interest including *Arabidopsis*, black raspberry, cane berry, blackberry, cherry, avocado, strawberry, wild strawberry, peach, grape, apple, tomato and/or tobacco.

Example 5: Inducible Expression of Morphogenic Regulators

[0389] In this example, expression vectors were constructed to test inducible ipt expression (GR-LHG4-pOP4::ipt) in tobacco transformation. The vector contained two expression cassettes. The ipt was driven by a pOP4 promoter, whose transcription activity is controlled by the transcription factor LHG4. LHG4 was fused with glucocorticoid receptor (GR), constitutively expressed and located in cytoplasm. GR-LHG4 protein translocates to a cell nucleus upon the application of inducing chemical dexamethasone (Dex). Each vector was transformed into tobacco leaves (*Nicotiana benthamiana*). Leaf segments at about 1 square centimetre were prepared from tobacco plants maintained in vitro. An *Agrobacterium* suspension containing the plasmid was prepared at OD 0.2 as the inoculum. Leaf segments were inoculated by the *Agrobacterium* by submerging in the suspension for 10 minutes and then were transferred to sterile filter paper (1 mL of MS liquid added) for a 2-day co-culture. The leaf explants were then transferred to MS-based agar medium containing no plant growth regulators or selection agent for regeneration. Medium with and without inducing chemical Dex (μM) were compared for the effect of induced ipt expression.

[0390] The explants showed darker green color and enlargement at 2-4 wks than the control; and numerous shoots emerged as early as 4 weeks after inoculation on hormone-free and nonselective medium. In addition, numerous shoots emerged as early as 4 weeks on hormone-free nonselective medium supplemented with inducing chemical dexamethasone (Dex). As the control, no shoots were regenerated from the same medium without Dex.

Example 6: Non-Transgenic Editing Using Morphogenic Regulators and a Gene-Editing System

[0391] The efficacy of non-transgenic editing using ipt in the same T-DNA with PDS editing machinery was examined. The vectors contained a sequence encoding LbCpf1 nuclease and gRNA targeting tobacco PDS genes to create deletion mutations. In the same vector, an ipt constitutive expression cassette was built in the same T-DNA with the PDS editing machinery. *Agrobacterium* containing the vector was used for tobacco transformation. Leaf segments at about 1 square centimetre were prepared from tobacco plants maintained in vitro. An *Agrobacterium* suspension containing the plasmid was prepared at OD 0.2 as the inoculum. Leaf segments were inoculated by the *Agrobacterium* by submerging in the suspension for 10 minutes and then were transferred to sterile filter paper (1 mL of MS liquid added) for a 2-day co-culture. The leaf explants were then transferred to MS-based agar medium containing no plant growth regulators or selection agent for regeneration. **[0392]** The plants that were regenerated with ipt included many shoots when regenerated on the hormone-free nonselective medium. A population of 200 shoots was subjected to molecular determination of transgene integration and gene editing. A total of 7 samples had transgene integration among the 200 (providing 96.5% nontransgenic, which is similar with Example 2). Four samples showed deletion mutations targeted by gRNA, including 3 with transgene integration and 1 without the integration. As the control, 20 shoots regenerated from medium with selection and BAP were used for the same molecular determinations, and photobleaching phenotype was observed on some regenerated shoots.

Example 7: Strong Constitutive Expression

[0393] RoKN1 and PavKN1 (strong constitutive expression) in tobacco transformation was tested. Leaf segments at about 1 square centimeter were prepared from tobacco plants maintained in vitro. An *Agrobacterium* suspension containing the plasmid (RoKN1 or PavKN1 expression) was prepared at OD 0.2 as the inoculum. Leaf segments were inoculated by the *Agrobacterium* by submerging in the suspension for 10 minute and transferred to sterile filter paper (1 mL of MS liquid added) for a 2-day co-culture. Then the leaf explants were transferred to MS-based agar medium containing 1 mg/L of BAP and 100 mg/L of spectinomycin as the selective agent. In contrast to expression with ipt, no different phenotype was observed on hormone-free nonselective medium but strong a shooty phenotype was observed on regeneration medium (BAP+ selection).

Example 8: Expression of Morphogenic Regulators with Various Promoters

[0394] Various morphogenic regulators were tested together with different promoters in tobacco transformation. RoWUS, SorbiWUS, PavWUS, AtBBM, TaBBM-1, TaBBM-2, SobiBBM, RoBBM, and PavBBM were tested. Each gene was driven by a double viral strong constitutive promoter or a weaker promoter (pNOS) for a comparative test. Leaf segments at about 1 square centimetre were prepared from tobacco plants maintained in vitro. An *Agrobacterium* suspension containing each of the plasmids was

prepared at OD 0.2 as the inoculum. Leaf segments were inoculated by the *Agrobacterium* by submerging in the suspension for 10 minutes and were then transferred to sterile filter paper (1 mL of MS liquid added) for a 2-day co-culture. Then the leaf explants were transferred to four types of MS-based agar medium for regeneration: 1) hormone-free and selection-free; 2) BAP 1 mg/L and selection free; 3) BAP 1 mg/L and selection; 4) 2, 4D 0.2 mg/L and selection.

[0395] The medium conditions post inoculation include 1) hormone-free nonselective 2) BAP w/o selection 3) BAP+ selection 4) 2,4 D+selection. There was no significant morphogenic changes for all tested genes on medium conditions 1), 2) or 3). On the 2,4 D containing medium, somatic embryogenesis was observed from RoWUS, PavWUS and SobiWUS transformed explants, as showed by early globular callus (both promoters), and bipolar structures (strong promoters) and proliferation of shoots from regenerated structure (strong promoter). The level of WUS expression appeared to affect the normal development as informed by visual reporter ZsGreen expression.

Example 9

[0396] WOX2-GR and WOX8-GR in tobacco transformation were tested. Results: shoot regeneration was observed from WOX2-GR inoculated explants at the presence of Dex on medium with BAP and selection (image below). Strong shoot proliferations were observed from WOX8-GR inoculated explants at the presence of Dex on medium with BAP and selection.

Example 10

[0397] RoWUS was tested in blackberry transformation. Results: higher efficiency of regeneration was observed in RoWUS transformed leaf petiole explants, as shown by higher number of shoots regenerated per petiole. Both promoter types (constitutive strong and constitutive weak) showed similar gene effect. Two *Agrobacterium* strains were mixed at different ratios and co-infiltrated to blackberry explants. One strain delivered a T-DNA containing the RoWUS expression cassette and the other strain delivered a T-DNA containing a Cas12a expression cassette for genome editing. Many regenerated shoots were obtained due to RoWUS regeneration enhancing capacity. When delivered at a 1:1 ratio of RoWUS T-DNA to Cas12a T-DNA, 36 shoots were recovered and 2 were positive in a qPCR test for presence of a hygromycin selectable marker gene (5.6% efficiency). When delivered at a 1:9 ratio of RoWUS T-DNA to Cas12a T-DNA, 50 shoots were recovered and 5 were positive in a qPCR test for presence of a hygromycin selectable marker gene (10% efficiency). In contrast to these results, a control sample group transformed only with the T-DNA containing the Cas12a expression cassette yielded 41 shoots and only 1 was positive in a qPCR test for presence of a hygromycin selectable marker gene (2.4% efficiency). This indicates that transgenic shoots can be recovered at a higher efficiency when transforming with RoWUS than without it.

REFERENCES

[0398] All references, articles, publications, patents, patent publications, and patent applications cited herein are incorporated by reference in their entireties for all purposes.

However, mention of any reference, article, publication, patent, patent publication, and patent application cited herein is not, and should not, be taken as an acknowledgment or any form of suggestion that they constitute valid prior art or form part of the common general knowledge in any country in the world.

[0399] U.S. Pat. No. 7,256,322 B2

[0400] US20180273960 A1

[0401] Gordon-Kamm B., Sardesai N., Arling M., Lowe K., Hoerster G., Betts S., and Jones T. Using morphogenic genes to improve recovery and regeneration of transgenic plants. *Plants* 8(2). Pii: E38 (2019)

[0402] Chen et al. A method for the production and expedient screening of CRISPR/Cas9-mediated non-transgenic mutant plants. *Hort Research* 5:13 (2018)

[0403] Carimi F., De Pasquale F., Crescimanno F. G. Somatic embryogenesis and plant regeneration from pistil thin cell layers of citrus. *Plant Cell Rep.* 18, 935-940 (1999).

[0404] Paul H., Belaizi M., Sangwan-Norreel B.S. Somatic embryogenesis in apple. *J. Plant Physiol.* 143, 78-86 (1994).

[0405] Fischer R., Emans, N. Molecular farming of pharmaceutical proteins. *Transgenic Res.* 9, 279-299 (2000).

[0406] Pogue G. P. et al. Production of pharmaceutical-grade recombinant aprotinin and a monoclonal antibody product using plant-based transient expression systems. *Plant Biotechnol. J.* 8, 638-654 (2010).

[0407] Richael C. M., Kalyaeva M., Chretien R. C., Yan H., Adimulam S., Stivison A., Weeks J. T., Rommens C. M., Cytokinin vectors mediate marker-free and backbone-free plant transformation. *Transgenic Res.* 17:905-917 (2008).

[0408] Boutilier K., Offringa R., Sharma V K., Kieft H, Ouellet T, Zhang L, Hattori J, Liu C M, van Lammeren A A, Miki B L, Custers J B, van Lookeren Campagne MMEctopic expression of BABY BOOM triggers a conversion from vegetative to embryonic growth. *Plant Cell* 14:1737-1749 (2002)

[0409] Nishimura A., Tamaoki M., Sakamoto T., Matsuo M., Over-Expression of Tobacco knotted1-Type ClassI Homeobox Genes Alters Various Leaf Morphology. *Plant Cell Physiol.* 41(5) 583-590 (2000)

[0410] Song X F, Guo P., Ren S.-C, Xu T.-T., and Liu C.-M. Antagonistic Peptide Technology for Functional Dissection of CLV3/ESR Genes in *Arabidopsis*. *Plant Physiol.* 161(3):1076-1085 (2013)

[0411] Long J. M., Liu C. Y., Feng M. Q., Liu Y., Wu X. M., Guo W. W., miR156-SPLs Module Regulates Somatic Embryogenesis Induction in Citrus Callus. *J Exp Bot.* 69(12):2979-2993 (2018)

[0412] Zhang T. Q., Lian H., Tang H., Dolezal K., Zhou C. M., Yu S., Chen J. H., Chen Q., Liu H., Ljung K., Wang J. W. An intrinsic microRNA timer regulates progressive decline in shoot regenerative capacity in plant. *Plant Cell* 27(2):349-360 (2015)

[0413] Vercruyssen L, Tognetti V. B., Gonzalez N., Van Dingenen J., De Milde L., Bielach A., De Rycke R., Van Breusegem F., Inzé D. GROWTH REGULATING FACTOR5 Stimulates *Arabidopsis* Chloroplast Division, Photosynthesis, and Leaf Longevity. *Plant Physiol.* 167(3): 817-832 (2015)

[0414] Malnoy, M. et al. DNA-free genetically edited grapevine and apple protoplast using CRISPR/Cas9 ribonucleoproteins. *Front. Plant Sci.* 7, 1904 (2016).

[0415] Subburaj, S. et al. Site-directed mutagenesis in *Petunia* x hybrid protoplast system using direct delivery of purified recombinant Cas9 ribonucleoproteins. *Plant Cell Rep.* 35, 1535-1544 (2016).

[0416] Woo, J. W. et al. DNA-free genome editing in plants with preassembled CRISPR-Cas9 ribonucleoproteins. *Nat. Biotechnol.* 33, 1162-1164 (2015).

[0417] Liang, Z. et al. Efficient DNA-free genome editing of bread wheat using CRISPR/Cas9 ribonucleoprotein complexes. *Nat. Commun.* 8, 14261 (2017).

[0418] Svitashchev, S., Schwartz, C., Lenderts, B., Young, J. K. & Cigan, A. M. Genome editing in maize directed by CRISPR-Cas9 ribonucleoprotein complexes. *Nat. Commun.* 7, 13274 (2016).

[0419] Zhang, Y. et al. Efficient and transgene-free genome editing in wheat through transient expression of CRISPR/Cas9 DNA or RNA. *Nat. Commun.* 7, 12617 (2016).

[0420] Kim, H. et al. CRISPR/Cpf1-mediated DNA-free plant genome editing. *Nat. Commun.* 8, 14406 (2017).

SEQUENCE LISTING

Sequence total quantity: 135

SEQ ID NO: 1 moltype = DNA length = 972
FEATURE Location/Qualifiers
source 1..972
 mol_type = other DNA
 organism = *Rubus occidentalis*

SEQUENCE: 1
atggactgcg gcgcccaaac aacaacaag gacaagctcc tcgtcattat gggagccacc 60
ggcgccggca agtcgcggtc atccctcgac ctgcgcgtt gtttccctcc ggcaaccttc 120
gaagtcatca acgcggataa aatgcaggtc tacaaggagc tcgacatcac caccaacaag 180
ctccccgctc cggagcgact cggcgctccc caccacttgc tcggcgagtt cgactcgctc 240
gacggcgagg ttactccgcg caagtcttcg gagctcgagg ctgcgggtgt ttccggcggt 300
agtgtctgga ggaagtgc tgtgcttggt ggtgggtcca actcattcat ccacgcgctg 360
gtcgtcgacc ggttcgaacc aggttttaat gtctttgacg gcgactctgt gacagtatcg 420
atctcatcgg agctccggta caattggtgt tttctctggg ttgatgtgtc gtcggtggtg 480
ttgacggagt acttgcgaa gcgagtagac gaaatgctag actcggggat gttactcgag 540
ctggccgagt ttacaacct ggccgaggaa gactcggtgg ttcgaaacgg gataagaaag 600
tccatcggag tagcggagtt cagccggtat ttttaggaag acccgccgcc gggaggtagc 660
gaggaggtgg gggacgatga cccggtacgg agagaagcat ataaggacgc cgtgagggaa 720
attaagctga acacgtgtca gctggcgaag aggcagattg gaaaaatctc acggtcgagg 780

-continued

```

ggggcggggt ggaacctacg gaggttagat gcgacggagg cgtttagggc ggcggggaag 840
tcagatgaaa aagacgggaa taggtgggctg gagatttggg aaagagagggt ggttggacca 900
agcgtgaaga ttgtgaatca tttcttgagg gaggagattc aatcaacaaa agtacaaaat 960
caaaagattt ag                                     972

```

```

SEQ ID NO: 2      moltype = DNA length = 972
FEATURE          Location/Qualifiers
source           1..972
                 mol_type = other DNA
                 organism = Rubus occidentalis

```

```

SEQUENCE: 2
atggattgcg gcgacacctac taccaataag gataagctgt tggttattat gggcgctacc 60
ggcgccggaa agtctagatg gtcattggat ctgctgcctt gtttccacc agctacattc 120
gaagttatta atgctgataa gatgcagggt tacaaaggac tggatcacac aactaacaaa 180
ctgctgcac cagagcgctt ggggtgttcc catcaccttc tgggcgaatt tgattctctt 240
gatggagagg tgaccccgag taagtgcaga gaactggcag ctgcggtggt ttctggcggt 300
tcagccagac gcaaaagtgc tgttctgggt ggtggctcta attcttttat tcatgccttg 360
gtggttgata gattcgaaac aggcctttaa gtgttcgatg gcgattctgt tactgtgtca 420
atctcttcag agcttcgcta taattgctgt tttctgtggg ttgatgtgtc ttcagtgggt 480
cttaccgaat acctgtctaa gaggggtgat gagatgttgg attcaggcat gttgcttgaa 540
cttgcctgag tctacaactc tgcgagggaa gattcagtg ttcgcacagg cattaggaag 600
tctatcggag ttgtgtaatt ttcagggtat ttcagaaaac atccaccacc aggcggctct 660
gaggaagttg gagatgatga tctgtgagg agagaagcct acaaggatgc agtgcccgag 720
atcaaatgta atacatgcca gcttgccaag aggcagattg gtaaaatctc aagattgcgc 780
ggtgcaggct ggaacctgcg caggttggat gcaactgagg cttttagagc cgctgttaag 840
tctgatgaaa aagatggtaa caggtgggta gagatttggg aaagagagggt ggttggccca 900
tctgtgaaga ttgtgaatca cttcctggag gaagagatcc agtcaactaa ggtgcagaa 960
cagaaatct ga                                     972

```

```

SEQ ID NO: 3      moltype = DNA length = 1074
FEATURE          Location/Qualifiers
source           1..1074
                 mol_type = other DNA
                 organism = Rubus occidentalis

```

```

SEQUENCE: 3
atgacacttg attccttccc aactcttcat tattcttatt ataattattac cccccattct 60
ctctcctcgc cggcggtgac cctcactccc ccaccatctc tcccccaaaa gccgcgctgg 120
gccccaatgg actccggccc agcaacaact acaaccgtcc cactccggca caaggacaag 180
cttgtgtcta tcattgggagc caccggagcc ggaaggtccc gtctctcaat cgacctcgcc 240
actcaacttc cttcttgctt gaaatcatc aactcggaca aaatgcagggt ctacaaggga 300
ctcgacatca ccaccaacaa gattccagtt cccgaccgta tcggagtttc ccaccacctc 360
ctcggtcagt tcgaccgctc cgcaggggag ctaccccccg ccgagtttcg acaactcgcc 420
gggtcaagccg tttccgcat tacttctcgg cggaaagtgc ccatgcttgt ggggtgggtcc 480
aactctctaa tacacgcctt cctcgttgac cgggtcgaac cgggtgtaga tgtcttcgaa 540
ggcgcgccgc ccatatcgtc ggagctcagg tacaattgct gctttctctg ggttgatgtc 600
tccttgacgg ttttagcgga ctacttgtct aagcagtag acgaaatgct cgactcggga 660
atgttggagg agttggccga gttttgcgac ccggaagatg acgacttgcc gattcgaacc 720
gggttgagaa aggtatttgg agtaccggag ttcagccggt ttttcaaaaa ataccgcgcg 780
atttgtcaga gtcagaaggc gagggccgat gatccggtgc ggagaggagc atataaagaa 840
gcggttaggg ccattaagga caaacctgtt cagctggcca agacgcagat tggaaagatt 900
ctacggctta ggtggcgagg aggggtggac ctacagaggc tagatgcgac cgaggcggtc 960
agggcggttg tgacgtcaga agacgatgcc ggaaagagggt ggtcagagat atggcagaaa 1020
cagggtgggtg aagcaagcgt gaaaattgtg aagcgattct tggagcagga gtag 1074

```

```

SEQ ID NO: 4      moltype = DNA length = 1074
FEATURE          Location/Qualifiers
source           1..1074
                 mol_type = other DNA
                 organism = Rubus occidentalis

```

```

SEQUENCE: 4
atgacattgg attcttttcc tactcttcat tactcatact acaacatcac tctcactca 60
ctgtcttcac cagcagtgac tctgactcca ccaccatctc tgctcagaa gccaaagatgg 120
gtctgcattg attctggacc tgccactacc acaactgttc cacttagaca taaggataaa 180
ctgggtggta ttatgggtgc aacaggcgct ggaaaaatctc gcctgtcaat cgatttggcc 240
actcaacttc cttcttgctt cgagattatc aattcagata agatgcagggt ttacaaaggc 300
ctggatatta ccacaaacaa gatccctgtg ccagatagaa ttggagttcc acatcacctg 360
ttgggtcaat ttgatcctct tgatggcgag ctgaccccgag ctgaattccg ccagttggcc 420
ggtcaggcag tgtctgctat tacatcaaga cgaaagtgc ctatgctgggt tggcgatct 480
aattcattga tccatgccct tctggttgat aggtttgagc caggcggtgga tgttttcgaa 540
gggtggcgctg ccattctctt agagcttaga tataactgct gtttctctgt ggtggatgtt 600
tctctgactg tcttgccgca ttacttgtca aagagagttg atgaaatgct tgattctgga 660
atgcttgagg aactggcaga gttttgtgat cctgaagatg atgatttggc aattaggacc 720
ggtcttagaa aggtatctcg cgtgccagag ttctcaagat ttttcaagaa gtatcctcca 780
atctgccagt ctcaaaaagc ccgcccgtgat gatccagtga ggagaggcgc atacaaggaa 840
gccgttaggg caatcaaaaga taacacctgt cagcttgcta agacacagat tggaaaaaatc 900
ttgcgcctta ggggtgtctg cggctgggat ttgcagcgcc ttgatgtctac tgaggccttt 960

```

-continued

```

agggcagtggtg ttacctcaga agatgatgca ggcaagcgct ggtctgagat ttggcagaaa 1020
cagggtggttg aagcttctgt gaagatcggt aaaagggtct tggagcagga atga 1074

```

```

SEQ ID NO: 5      moltype = DNA length = 1125
FEATURE          Location/Qualifiers
source           1..1125
                 mol_type = other DNA
                 organism = Prunus avium

```

```

SEQUENCE: 5
atgacacttc atcccttccc aactcactcc caccaccatt attctaattc caatttttat 60
tcttatatct ctcgtttcca cttgttcccg ccggtcaacc acctctgccc caccactcct 120
tcccatcatc atcctctccc ccacatcaag ccccgcccgcg gggcccatat atcctccggc 180
cccaccaccc caccgcaaca acaacagcgc aaggacaagc ttctcattat cttgggcggc 240
accggcgccg gcaagtccecg cctctccctc gacctcgcca cccgcttccc tttcttcgaa 300
atcgtgaact ccgacaaaat gcaactctac gccggcctcg acatcaccac caacaagctt 360
cccatccccc accgctcggg gtccccccac catctcctcg gcgagttcga ccctcgccac 420
ggcgatttta cgcccgccca attccgcgcc gttgcgggtc aagccatttc cagcattacg 480
aatcgaagga aggtgcccga gctcgtccgc ggctccaact ccttcattca tgcgctcctc 540
gtggaccggg tcgaaccggg ctccaatgtc ttcgaaccgg gttccaatgg ctctgtctct 600
tccgagctca ggtataattg ctgttttctg tgggtggacg tgtcgttgac ggtcttgacg 660
gagtaacttg gcaagcagag cgacgaaatg ctcgactcgg ggatgttgga cgagttggcc 720
gagttctgca acccgagtag ccaggacgaa gacgaattga cggcgagtag gacagcgttg 780
agaaaggcga ttggcgtgcc cgagttcact cggatatttg aaaaatatcc accacagggg 840
aggggcccga aggggtgatg tcgggagcgg agggaaagcat acgaagaggg ggtgagggcg 900
atcaaggaca acacgtgtca gctggcgaag aggcagatag ggaagatcct acggctgaag 960
ggaggagggt ggagacctac gaggttagat gcaacgggac cgtttagggg ggtggttgcg 1020
acgacgtcgt cggaataagg cgacggaaaag aggtgggtcag agatatggga gaggcagggt 1080
gtcaaaccaa gcgtgaaggt tgtgaagcgt ttcttgagac agtag 1125

```

```

SEQ ID NO: 6      moltype = DNA length = 1125
FEATURE          Location/Qualifiers
source           1..1125
                 mol_type = other DNA
                 organism = Prunus avium

```

```

SEQUENCE: 6
atgactcttc atccatttcc taccactctc catcaccatt actcaaatct gaactttctac 60
tcttacattt caccgtttca cctgtttcca ccagttaatc acttggtccc aactaccctc 120
tctcaccatc acccattgcc tcatattaag ccaagacgcg gagcccatat ctcttcagggt 180
ccaacaactc caagacagca gcagcagaga aaggataaac tgttgattat ccttggtgca 240
acaggagctg gcaagtcctc cctgtcattg gatcttgcaa ctagggttcc atttttcgag 300
atcgtgaact cagataagat gcagctttac gctggcctgg atattaccac aaacaagttg 360
ccaatccctg ataggcttg cgttcctcat cacttctcgg gagaattcga tccaagacat 420
ggcgatttta cccctgcccc gttccgcgct gtggctggac aggcattctc ttcaatcaca 480
aacaggagaa aagtcctat gctgggttgc ggatctaact cttttattca cgccttgctt 540
gtggatcgct tcgagccagg ttctaattgt ttgaacctg gcttcaacgg atcagtgctc 600
tcagagctta ggtataattg ctgttttctg tgggtggatg tttctcttac agttctgact 660
gagtaacctg gcaagagagt ggtgaaatg ttggattcag gcatgttgga tgagcttgca 720
gaattctgtg atccagatcg ccaggatgag gatgaattga ccgctctcga gacagcactt 780
aggaaagcta ttggagttcc taggtttact agatatctcg aaaagtaacc acctcagggt 840
cgcggtggcg aaggcagatg tagggagcgc aggggaagctt atgaggaagc cgtgagagca 900
atcaaaagata acacttgtca gctggccaag agacagattg gcaaaatcct gagactgaag 960
ggcggcgctg gggatctgag aagactggat gctaccgatg ctttagggc agtggttgct 1020
actacctctt cagataagga tgatggaaaa aggtgggtctg agatctggga aagacagggt 1080
gttaagccat cagttaaggt ggttaaaaga ttcttgatg agtga 1125

```

```

SEQ ID NO: 7      moltype = DNA length = 1173
FEATURE          Location/Qualifiers
source           1..1173
                 mol_type = other DNA
                 organism = Rubus occidentalis

```

```

SEQUENCE: 7
atggaggaaat acaataacag taatgaaatg gatcatgaga gttcaggtag tcccagagga 60
aatctcctgt acgcatctcc aaatctcgga ggtaactatg gtagaactgc cgcagggtgat 120
cataatcagc agatcaaacac ctttcatctt caatcaagcg gtggcgagtg ttttcagttc 180
ggaggaaacag cacagcccac cgtgaagacc gaagcagcag gtagcacttc ccaacatgct 240
ccccacaaat ttcatcatca gtttctcctc cctctattaa tattgtcaag aggtcagaat 300
gatcagctac cagttcagca acaacagcac atagaaaaag agaatgaagt ggaggccatc 360
aaggctaaaa tcactgctca cctcactac tctaactctc tgcaagctta catggattgc 420
caaagggttg gagctccgtc tgaagtcgtg gctcgtctcg cagctgcgcg ccaggagttt 480
gaagcagcac agcgatcttt agtcagttcg agagatgctt cttcgaaaga cccagaactc 540
gatcagttca tggagcttta ctatgatag ttggttaaat atcgggagga actaacaagg 600
cccatacaag agcaaatgga tttcatgagg aagattgaaa ctacagctaa catgctcggc 660
aatgacagta gcaactactc tcttcgcac tctcggcctc cggcgcaaga ggacaaatgt 720
gacggaaaatg gttcatctga agaggaccag gacaacaata gtggtggaga aactgaagt 780
ccgagatcgt atccgagagc cgaagacaga gaactcaaga atcacctctt gagaaagtac 840
agtggttact taagttagct caagcaagaa ctctccaaga aaaagaagaa aggcacaaatg 900

```

-continued

```

cccaaagatg ccaggcagaa gcttctgagt tgggtgggagt tgcactacaa gtggccatat 960
ccttcggagt cggagaaggt ggcattggcg gagtcgacag gtttggatca gaagcagata 1020
aacaattggt tcataaatca aaggaagcgg cactggaagc catcggagga catgcagttc 1080
atggtgatgg atgggttgca ccacagaat gcagcaactc tctacatgga tggacactac 1140
attggtgatg gtcattaccg ttcgggtccc taa 1173

```

```

SEQ ID NO: 8      moltype = DNA length = 1362
FEATURE          Location/Qualifiers
source           1..1362
                 mol_type = other DNA
                 organism = Rubus occidentalis

```

```

SEQUENCE: 8
atggaggaat acaataacag taatgaatg gatcatgaga gttcaggtag tcccaggtaa 60
gtttctgctt ctacctttga tatatatata ataattatca ttaattagta gtaatatataat 120
atttcaaata tttttttcaa aataaaagaa tgtagtatat agcaattgct tttctgtagt 180
ttataagtgt gtatatttta atttataact ttctaatat atgacccaaa tttgttgatg 240
tgcaggggaa atttctgtga cgcatctcca aatctcggag gtaactatgg tagaactgcc 300
gcaggtgatc ataactcagc gatcaacacc ttcatcttc aatcaagcgg tggcgagtgt 360
tttcagtcgg gaggaacagc acagcccacc gtgaagaccg aagcagcagg tagcacttcc 420
caacatgctc ccacaaaatt tcatcatcag ttctctctc cctcattaat attgtcaaga 480
ggtcagaatg atcagctacc agttcagcaa caacagcaca tagaaaaaga gaataagtg 540
gagggccatca aggcataaat catcgctcac cctcactact ctaatctctt gcaagcttac 600
atggattgcc aaaggggtggg agctccgtct gaagtcgtgg ctctctctgc agctgcgcgc 660
caggagtttg aagcacgaca gcgatcttta gtcagttcga gagatgcttc ttcgaaagac 720
ccagaactcg atcagttcat ggaagcttac tatgatagt tggtaaata tcgggaggaa 780
ctaaacaaggc atacaaga agcaatggat ttcatgagga agattgaaac tcagcttaac 840
atgctcggca atgcagtag cactactcct ctctgcctct tctcggcctc ggcgcaagag 900
gacaaatgtg acggaatggt ttcatctgaa gaggaccagg acaacaatag tggtagagaa 960
actgaagtgg ccgagatcga tccgagagcc gaagacagag aactcaagaa tcacctcttg 1020
agaaagtaca gtggttactt aagtagcctc aagcaagaac tctccaagaa aaagaagaaa 1080
ggcaaatgca ccaaaagtgc caggcagaag cttctgagtt ggtgggagtt gcaactacaag 1140
tggccatata cttcggagtc ggagaagggt gcatctggcg agtcgacagg ttgggatcag 1200
aagcagataa acaattggtt cataaatcaa aggaagcggc actggaagcc atcggaggac 1260
atgcagttca tgggtgatga tgggttgcc cccacagaatg cagcaactct ctacatggat 1320
ggacactaca ttggtgatgg gtcattaccg ctcgggtccct aa 1362

```

```

SEQ ID NO: 9      moltype = DNA length = 1068
FEATURE          Location/Qualifiers
source           1..1068
                 mol_type = other DNA
                 organism = Prunus avium

```

```

SEQUENCE: 9
atggaagagt acaacccaat ggatcatgag agctcagggt gtagaggaaa ttctctgtac 60
gcctcaccaa atcttgagg caactatggg agggcaagcc atgatcatca gatgggcatc 120
aacgccttct atcttcaatc gagcgggtgt ggtgatcttc actctccagg aggagcacac 180
ccaattaatg tcaagactga agccagcaca tcccatcatg gccatcacia atttcagtac 240
aacaacaaca acaataataa taataataat cctcttctgt catcaagagg gcaccaacca 300
gtagttcaac aacagcataa tctggaccgt caaaacgatg atcacacctc gagctccaac 360
gaagtggaa g ccatcaaaagc caagatcatc gccaccctc agcagcgggt ttcagtggct 420
tcaagagagg cttcaaaaga tccagaactc gatcagttca tgggaagttta ctatgatag 480
ttggtgaaat atcgagagga gctaaacaag ccatacaag aagccatgga ttcatgctgg 540
aggattgaaa ctcagctcaa catgcttgcc acaataatg ctctctctct tgcactcttc 600
tcacctctgt aggcacaaat tgaaggaatt ggttcactct aagaggagca ggagaatagt 660
ggttgagaaa cagaagtgcc tgagattgat ccaagagctg aagacagaga gctcaagaac 720
cacctgttga aaaagtatag tggttactta agtagcctga agcaagagct ttccaagaaa 780
aagaagaaag ggaattgtcc caaagatgcc aggcagaagc ttctcagttg gtgggagttg 840
cattacaagt ggccatatcc ttcagaatcg gagaagggtg cattggcgga gtcaaccggt 900
ttggatcaga acaaaataaa caattgggtc ataaatcaga ggaagaggca ctggaaaccc 960
tcagaggaca tgcagttcat ggtgatggat ggccctacac ctcagaatgc ggccctgtat 1020
atggatggac actacatagg cgatggccac taccgtctgg ggccatga 1068

```

```

SEQ ID NO: 10     moltype = DNA length = 1068
FEATURE          Location/Qualifiers
source           1..1068
                 mol_type = other DNA
                 organism = Prunus avium

```

```

SEQUENCE: 10
atggaagagt acaacccaat ggatcatgag agctcagggt gtagaggaaa ttctctgtac 60
gcctcaccaa atcttgagg caactatggg agggcaagcc atgatcatca gatgggcatc 120
aacgccttct atcttcaatc gagcgggtgt ggtgatcttc actctccagg aggagcacac 180
ccaattaatg tcaagactga agccagcaca tcccatcatg gccatcacia atttcagtac 240
aacaacaaca acaataataa taataataat cctcttctgt catcaagagg gcaccaacca 300
gtagttcaac aacagcataa tctggaccgt caaaacgatg atcacacctc gagctccaac 360
gaagtggaa g ccatcaaaagc caagatcatc gccaccctc agcagcgggt ttcagtggct 420
tcaagagagg cttcaaaaga tccagaactc gatcagttca tgggaagttta ctatgatag 480
ttggtgaaat atcgagagga gctaaacaag ccatacaag aagccatgga ttcatgctgg 540

```

-continued

```

aggattgaaa ctcagctcaa catgcttggc aacaataatg ctctctctct tcgcatcttc 600
tcacctctcg aggacaaatg tgaaggaatt ggttcacatg aagaggagca ggagaatagt 660
ggtggagaaa cagaagtgcc tgagattgat ccaagagctg aagacagaga gctcaagaac 720
cacctgttga aaaagttagt tggttactta agtagcctga agcaagagct ttccaagaaa 780
aagaagaaa ggaaattgcc caaagatgcc aggcagaagc ttctcagttg gtgggagttg 840
cattacaagt ggccatatcc ttcagaatcg gagaaggtgg cattggcgga gtcaaccggg 900
ttggatcaga acaaaaataa caattgggtc ataaatcaga ggaagaggca ctggaaaccc 960
tcagaggaca tgcagttcat ggtgatggat ggccacacc ctcagaatgc ggccctgtat 1020
atggatggac actacatagg cgtatggcac taccgtctcg ggccatga 1068

```

```

SEQ ID NO: 11      moltype = DNA length = 2103
FEATURE           Location/Qualifiers
source            1..2103
                  mol_type = other DNA
                  organism = Rubus occidentalis

```

```

SEQUENCE: 11
atgaacatga actggttggg ttctctctct tccctcaag aagatgatgc tccaatatca 60
caccagctcg ctgaccaaga gacgttagcc tctcgccctg gcttcaactc caatgatcaa 120
atccaaggtg ccggcgggtg ggcagattgg aatatgaagg gtgtagacat gaactcatca 180
gatggcaata actacaaaag gacctcatca tcatccgata tgtcgccat gctaatggga 240
agcagtacta ctactagtac ttcatgcagc attatcagcc agccaaacct tgaatatcat 300
catcaacaac aacaactacc aaagcttgaa aacttcttg gccgacattc ttttgcggat 360
catgacaaaa gtagtactag tcatgattac atgttcgata tgaataatgg accgggtgca 420
attagcagta atgttatgaa tacttctaata actaatagta acaatatgg gctgtccatg 480
atcaagacct ggctgaggaa tcaacctcct cagccgcgcg ataacaacca ccagctggag 540
caggagagca agactactaa ggagagcaga aatcaaccac agagtagtct atcgctctca 600
atgagtactg gatcgcttca tacagtaaca actgcaactg caggtggtgc cagtggccagt 660
gccgtgtccg gggaaactac ctcatcatca aataataata gaaagcagtc gccagttttg 720
acagccacaa caactgatcc ccaaaactca actactgggt gtgccattga ggctgtgccc 780
agaaaggcca ttgatacatt tggccagagg acttctatat accgtgtgtg aaccaggcac 840
agatggacag gtagatatga agctcatcta tgggataata gctgccgaag agagggacaa 900
actcgcaagg gaaggcaagg tggttatgac aaagaagaca aggcagctag agcttatgac 960
ttagcagcat tgaatatatt gggcaccaca accactacca actttccgat cagtacctat 1020
gaaaaagaga tagatgaaat gaagccgatg acaagacagg agtatgttgc atctctgaga 1080
aggaagagta gccgggtttc tcgaggtgca tccatttacc gaggggtaac caggcaccac 1140
cagcatggga gatggcaagg aagaattggt agagttcgag ggaacaaaga tctgtacctt 1200
ggaaactttc gtacccaaga ggaagcagct gaggcgatg acattgctgc cataaagttc 1260
cgaggactca atgcggtaac aaactttgac atgagcagat atgatgtcaa gtccattctg 1320
gagagcagcg catgtcccat tactactggt gccctcgcaa agcgattgaa ggatgttcag 1380
cagcagcctc ctctctcggc agatcatcat catcagatca tgtgtcgtc tgtgctggat 1440
catcacggac agatgatcag atcttcatcg tcaacagagc atgatatcat gggcaatatc 1500
tttcatcagg atgatgatca tctccgtcaa caacttcaga tgggaggcac ccacaatttc 1560
ttccatgata atgatcacca gcagcgtagc ggcccttatg gattaatgga ttcttcggca 1620
gcagcagctt ctatggaaca cagctcaggc tctaattctg tcatatatag tgttggggat 1680
caccatggta ataataatgg atatggaagg aatgctagag ggactggtag tggctatata 1740
atgcccatgg ctatgagtac agttgtggct aatcatggcc aaaatcaagc tgatggaaat 1800
aatatcaatg gtttcggaga tgggtgatgt gatgatcagg aagtgaatat taaggcgagc 1860
cagcttgggt atgatcatca tcaaaatgtg tttctgggtt caagtaatac tacagatccc 1920
gcttatcagc atcagcgaag caacaggaaac ttgtattatc aacttcagtc acatgatgac 1980
cagcatgaat cgtcatcaga agcagtagct actagtagta ctacatgtaa catgaattgg 2040
gttccaacag ctgttccaac tcatcttgct cattctactt ttacagctcg gaacgcacaa 2100
tag 2103

```

```

SEQ ID NO: 12      moltype = DNA length = 2103
FEATURE           Location/Qualifiers
source            1..2103
                  mol_type = other DNA
                  organism = Rubus occidentalis

```

```

SEQUENCE: 12
atgaatatga actggttggg cttttctctt tctccacagg aagatgatgc tctattttct 60
catcagctgg ccgatacaga aactctggca tctagattgg gcttcaattc aaacgatcag 120
atccaggagg ctggcggaga ggcgagttgg aatatgaagg gtgtggatat gaactcttca 180
gatggcaata actataagcg cactcttca tcttcagatc tttctccaat gctgatgggt 240
tcttcaacta ccacatctac atcatgctct attatctcac agccaaattt ggagaacct 300
caccagcagc agcagttgcc taagcttgaa aattttcttg gcagacattc tttcgccgat 360
cacgatcagt cttcaacctc acacgattac atgtttgata tgaataacgg acctggtgca 420
atctcttcaa agcttatgaa cacttctaac accaactcaa ataacatcgg actgtctatg 480
atcaaaacat ggttgaggaa tcagccacct cagccaagag ataataacca tcagctggag 540
caggaatcta agactaccaa agagtcaagg aaccagcctc agtcttctact gtcattgtct 600
atgtcaactg gttctttgca caccgtgaca actgctacag ccggtggcgc atctgcttca 660
gctggcgtcg gcgagactac atcttcatct aataacaata gaaagcagtc accagtttg 720
accgctacta ccacagatcc tcagacacag actacgggag gtgccattga agcagtgcca 780
aggaagacca tcgatacttt cggacagaga acctctatct atcgcggtgt tacacgccat 840
aggtggagctg gcagatacga ggctcacctt tgggataatt catgtagacg cgaaggacag 900
actagaaaag gtgcgccagg ccgatatgat aaggaagata aagctgctag agcttatgat 960
ctggcagctt tgaataactg gggcacaact accacaacta acttcccaat ctctacttat 1020

```

-continued

```

gagaaggaaa tcgatgagat gaaacctatg accagacagg aatacgtggc ttcactgagg 1080
agaaagtcac ctggattctc tcgcggtgac tcaatctatc gcggagtgac taggcacac 1140
cagcatggta ggtggcaggc taggatcggc agagttgccc gaaataaggc tctttatctg 1200
ggcacatttt ctactcagga agaggctgca gaggcatacg atattgctgc catcaagttt 1260
cgcggaacta atgctgtgac caacttcgat atgtctagat acgatgttaa atcaatcttg 1320
gaatcatctg cccctccaat caccacaggc gcttctgcca agcgccgtga agatgtgcag 1380
cagcagccac caccaccagc tgatcatcac catcagatta tgttgtcacc tgttcttgat 1440
caccatggac agatgatcag gtcactttca tctactgagc atgatattat gggtaatatc 1500
ttccatcagg atgatgatca cctgagacag cagttgcaga tgggtggcac ccacaacttt 1560
ttccacgata atgatcacca gcagcgctct ggtcttatgg gcctgatgga ttcacttgca 1620
gctgcgcgat ctatggaaca ttcactctga tctaattctg tgatctattc agttggcgat 1680
catcatggca acaataacgg ttatggctct aacgcccgcg gaacaggttc aggctacac 1740
atgctatagg caatgtctac tgtggttgct aatcacggcc agaaccaggc agatggaaat 1800
aacattaatg cctttggcga tggcgatggc gatgatcagg aagtgaacat caaggctcag 1860
cagcttggtt atgatcacca tcagaatgtt ttccctgggt catctaaccac taccgatcca 1920
gcataccagc accatgcttc taataggaac ctttattacc agctgcctgt gcgatgatg 1980
cagcacgagt catcttcaga agcagttgct acatcttcaa caacttgcaa tatgaactgg 2040
gtgccaaccg cgttctctac acatttgga cactcaacct ttacagtttg gaatgatact 2100
tga 2103

```

```

SEQ ID NO: 13      moltype = DNA length = 1599
FEATURE           Location/Qualifiers
source            1..1599
                  mol_type = other DNA
                  organism = Prunus avium

```

```

SEQUENCE: 13
atgggttcca tgaactgggt ggggtttctc ttttccctc aagaagtcgt tctgtctgat 60
ccatccatat ctgaggggtga tcatcatcat caacatcatg acggccatga gttatctgat 120
gatcagaccg ccacactagt ctctcgctt ggcttcaagt actccgatga catctccggt 180
accgatgtca ccggaggctc ctttgatctc acttctcatg atgactcatc cgctactact 240
actgctgctg ctgttctctaa taatcattct cacaaccacc tcatccctcc accttttggc 300
atgcttcac  aagctgtgac attcaacacc accaatattc actcccaagg tgcagattgg 360
agtaatatga actcatcaga tagcagcacc tacaagacga cttcagacca cctgtctatg 420
ctgatgggaa gcaacagttc gtcagaagc caaaaccttg aaaataatca tcaacagcca 480
aagcttgaaa acttcccttg ccgacactcc ttcgttgccg atgagcacca catcggcggc 540
cagagtagta gtggcacgca tgcatacaat aataatagca ccagcaatac aatacagact 600
ctttcgcttt cgtatgagta tggatctcac actcatcata atactactag tactgctgct 660
gtgggagaaa cttgttcac  agatcgagat aatacttgta ataataataa caagctggca 720
gttgtaggga ccacgcctc aacgcctcct cccctgggga tccgatagca gactactagt 780
actactgcta ctgctattga ggcgtgccc agaaagtcgg ttgatacttt tggacagagg 840
acttctatat accgtggcac agatggacag gtagatatga agctcatctc tgggataata 900
gttgccggag agaaggacaa actcgcaagg gaaggcaaga tcagcaacta tgagaagag 960
atagacgaaa tgaagcacat gacgagacag gagtacgttg catctttaag aagggaagag 1020
agtggctttt ctctgtgtgc atccatttat cgaggggtta caagacacca ccagcatggg 1080
agatggcaag caaggattgg tccagtcgca gggacaacag atctctacct gggaaacttt 1140
agctccaact ctgtcatgta tagcactgct actactacta atggaggagg aggaggagga 1200
ggaggagatc atggtggcta tataatccca atgggtacag ttatatctaa tgatcacaa 1260
gatgggtttg gagatggcac taatcatgat catcaggtag aggcgggtgg gcttggattt 1320
gaaaatgtgt ttggtctaac ttctgcttct gttaatgcta caaatgttac tagggatgct 1380
tataataatc atgcaaggaa cttgtattat cttccgcagc agctgccgca atcgtcatca 1440
gtgtcatcag tatcatcagc aggtgtggcc aagggcgcca gcgcataatg tattcatgaa 1500
gcagggcaat gtaacaattg gtagcccaac gcagttccaa cgtcgactaa cagtaaacac 1560
aacatgctcc tacttttcc  tgtctggaat gacacataa 1599

```

```

SEQ ID NO: 14      moltype = DNA length = 1599
FEATURE           Location/Qualifiers
source            1..1599
                  mol_type = other DNA
                  organism = Prunus avium

```

```

SEQUENCE: 14
atgggctcta tgaattggct tggattttct ctgtctccac aggaagtggt gccagccgat 60
ccttctattt cagaaggcga tcatcaccat cagcaccatg atggacatga gctgtctgat 120
gatcagactg caacctgggt gtcacgcctt ggttttaaat actctgatga tatctcaggc 180
actgatgtta ccggcgggat cttcgatttg acctctcacg atgattcttc agcaactacc 240
acagctgcgc cagttctcaa taaccactca cataatcacc ttattccacc tccatttggt 300
atgctgcac  aggtcgccgc attcaatact accaacatcc actctcaggg cgctgattgg 360
tcaaacatga actcttcaga ttcttctact tataagacaa cttctgatca tttgtcaatg 420
cttatgggct ctaattcttc atacagatca cagaacttgg agaataacca tcagcagcca 480
aagctggaaa actttttggg acgccactct ttcgtggcag atgaacacca tattggtggc 540
cagcttctat cttggactca tgcattacaat aacaattcta catcaaacac tatccagacc 600
ctgtctttgt caatgtctac cggatctcac acacaccata acaccacac aacagctgct 660
gttgccgaga cttgtctac  gtagcggat aatacatgta acaataacaa taagctggct 720
gtgggttaga ctacgcctc aactgcacct ccacctgcca ttgattctca gacaacttca 780
accacageta ccgccatcga agctgtgcca agaaagctgt ttgatacttt cggacagaga 840
acctcaatct atcgcggaac agatggtcag gtggatatga agttgatctc tggaattatc 900
gttgccgggtg agaaggataa acttgcaaga gaaggcaaga tttcaacta cgagaaggaa 960

```


-continued

atcgatgaga	tgaagcatat	gaccagacag	gaatacgttg	cttctctgag	aagaaagtct	1020
tctggctttt	ctcggcgagc	ctcaatctat	cgcggagtga	caaggcacca	tcagcacggt	1080
aggtggcagg	ctaggatcgg	tagagtggcc	ggcaataaag	atctttatct	gggcaccttc	1140
tcactaact	ctgtgatgta	ctcaacagcc	actaccacta	atggcggcgg	cggcggcgcc	1200
ggcggcgatc	atggcggaata	cattatccca	atgggcactg	ttatttctaa	tgatcacaaac	1260
gatggatttg	gcgatggcac	taacctgat	caccaggtga	aagcagttgc	tcttggtttt	1320
gagaatgtgt	tcggctcaac	ttctgcatca	gtgaatgcta	caaacgttac	tagggatgcc	1380
tataacaatc	atgcaagaaa	cctgattac	ttgccacagc	agctgcctca	gtcatcttca	1440
gtgtcttcag	tttcttcagc	tggcgtggcc	aagggtggct	ctgcatacga	tattcacgaa	1500
gctggacagt	gtaacaattg	gatgccaaca	gctgttccta	catctactaa	ctcaaatatc	1560
aacatgccac	ctaccttcac	agtgtggaat	gatacttga			1599

SEQ ID NO: 15 moltype = DNA length = 894
 FEATURE Location/Qualifiers
 source 1..894
 mol_type = other DNA
 organism = Rubus occidentalis

SEQUENCE: 15

atggaaccac	aatcccagca	accaaactgag	gatggaggaa	gcaacaaagc	agctgggagt	60
gctaacaatgc	tttgcaggca	gagcagtagc	aggtggactc	ccactactga	tcagataaga	120
atcctcaagg	agctttacta	caacaaggga	gttaggtccc	caagtgcaga	gcagattcag	180
aggatctgtc	tccagctgaa	acggtacgga	aagatcgagg	gaaagaatgt	cttttactgg	240
ttccagaacc	acaaggctcg	agagaggcag	aagaagaggt	tcacttcgga	tgttcatgtg	300
cccatgcaaa	gatcagggtc	tgttgggaac	agtaacgtag	ctaattggaa	acctgaggat	360
cagtatctca	acatgaacaa	caagtagctc	aacatttgtc	aaggggtttc	ttcttctgca	420
tctgcttctg	cttcttcagc	tgggtgtggt	tcttttaatg	ggcagatggg	gaactatggg	480
gtttatggat	ctatgaacat	ggagaagagt	tttagggact	gttcaatctc	agctggagga	540
ggtaccagta	ctactggtgt	tggaattggt	ggatccatta	gtcacaatta	cgggtcatgg	600
gtcgggtgta	gtgttgaccc	gtattcctcc	tcactcgta	ccactactag	tactactttc	660
tttgacaacc	aaaaacaat	tttcatggaa	gaagaagaag	aggatcacca	ggagattgaa	720
acccttcccc	tgttccccat	gcattggtgag	gacatctttg	gcaacatcaa	gacaacttcc	780
gaaggcggcg	gcggcgacta	cggctactac	tccagcggct	ggggcggtag	ctctgcact	840
tcccttgagc	tcagcctcaa	ttcctacggc	tacaacaacc	ccaactatta	ctga	894

SEQ ID NO: 16 moltype = DNA length = 894
 FEATURE Location/Qualifiers
 source 1..894
 mol_type = other DNA
 organism = Rubus occidentalis

SEQUENCE: 16

atggaaccac	agtctcagca	gcctactgag	gatggcggat	caaataaggc	tgccggatct	60
gctaacaatgc	tgtgcaggca	gtcttcaaca	agatggactc	caactaccga	tcagatcagg	120
atcttgaagg	aactttacta	caataagggc	gtgagatcac	cttctgccga	gcagattcag	180
cgcactctgtc	tgcagttgaa	gagatatggc	aagattgaag	gcaagaatgt	gttttatttg	240
ttccagaacc	ataaggcaag	agagcgccag	aagaagagat	tcacttctga	tgtgcacgtt	300
ccaatgcagc	gctcaggtct	ggtgggcaat	tctaacgttg	ctaattggaa	gcctgaagat	360
cagtatttga	acatgaataa	caagtactca	aacatttgcc	aggggtgtgc	ttcatctgca	420
tcagcttctg	cctcatctgc	cggcgtgggt	tcttttaatg	gccagatggg	taactatggc	480
gtttacggat	ctatgaatat	ggagaatatc	ttccgcgatt	gttcaatctc	tgcaagggtg	540
ggaacttcaa	caactggcgt	gggaattggt	ggctcaatct	ctcataatta	tggctcatgg	600
gtgggaggtt	ctgttgatcc	atattcatct	tcattctaca	ccacaacttc	taccacattt	660
ttcgataacc	agaagcagat	tttcatggag	gaagaggaag	aggatcatca	ggaaatcgag	720
actcttccac	tgtttcttat	gcacggagaa	gatatcttgc	gtaatatcaa	gactacctca	780
gagggcggag	gtggcgatta	cggctattac	tcactctggat	ggggaggttc	atctagaaca	840
tctttggaac	tttcaactgaa	ttcttatggc	tacaataacc	ctaactatta	ctga	894

SEQ ID NO: 17 moltype = DNA length = 918
 FEATURE Location/Qualifiers
 source 1..918
 mol_type = other DNA
 organism = Prunus avium

SEQUENCE: 17

atggaacctc	aacaaaccca	acaacaacca	aatgaagatg	gcagcaacaa	agggactagt	60
ggtactaatt	ttctgtgcag	gcaaagcagt	acaagggtga	caccacaag	tgatcagatt	120
agaatattga	aggaccttta	ctacaacaat	ggtgtgaggt	ccccaaagtc	agagcagatt	180
cagaggatct	ctgctcggct	gagacagtag	ggcaagattg	aaggcaagaa	cgttttctat	240
tggtttcaga	accacaaggc	tcgtgagagg	cagaagaaaa	ggttcattga	tgtaccagcc	300
ccaccatca	tgcaaaagatc	agggcttggg	attaataata	atgctcctac	tgcttatgaa	360
cccattaatc	acagcaacaa	gtaccccaac	atttctgctt	ctgctggggg	ttcttctgca	420
tctgcttctt	cagctggtgt	gattgtctgt	gggcaaatgg	ggagctatgg	ctatggatct	480
atgaccatgg	agaagagttt	tagggactgc	tcaatatcag	ctgcaggagg	tactagttagt	540
ggtcatgttg	gtggatctaa	taataatatt	ggtcacaacg	ttggatcatg	ggttggtgtt	600
gatccatatt	cttcacccta	cactctcttt	gacaagagat	catcatcaag	acaagtgttt	660
ggtgaccaag	aaaatatgat	ggatgaagaa	gatcatgaat	accaagaaaa	cctgcaagat	720
tcccagaga	ttgagacctc	ccctctcttc	cccatgcatg	gtgaagacat	ccatggcttt	780
ggcaacatca	agtcacacctc	cgacagctac	tactccggct	ggtaccgcctc	cgacgagggc	840

-continued

aacaatggcc	gcacttcct	tgagetcagc	ctcaattcct	acggccacat	gacccctgat	900
tatttcagct	catattaa					918

SEQ ID NO: 18 moltype = DNA length = 1107
 FEATURE Location/Qualifiers
 source 1..1107
 mol_type = other DNA
 organism = Prunus avium

SEQUENCE: 18

atggaacctc	aacaaaccca	acaacaacca	aatgaagatg	gcagcaacaa	aggtaagttt	60
ctgcttttac	ctttgatata	tatataataa	ttatcattaa	ttagtagtaa	tataatattt	120
caaatatttt	tttcaaaata	aaagaatgta	gtatatagca	attgcttttc	tgtagtttat	180
aagtgtgtat	attttaattt	ataacttttc	taatatatga	ccaaaatttg	ttgatgtgca	240
gggactagtg	gtactaat	tctgtgcagg	caaagcagta	caagggtggac	acccacaagt	300
gatcagatta	gaattattgaa	ggacctttac	tacaacaatg	gtgtgaggtc	cccaagtgtca	360
gagcagattc	agaggatctc	tgctcggctg	agacagtacg	gcaagattga	aggcaagaac	420
gttttctatt	ggtttcagaa	ccacaaggct	cgtgagaggc	agaagaaaag	gttcattgat	480
gtaccagccc	cacctcat	gcaaaatca	gggcttggga	ttaataataa	tgctcctact	540
gcttatgaac	ccattaatca	cagcaacaag	tacccaaca	tttctgcttc	tgctggggtt	600
tcttctgcat	ctgctttctc	agctggtgtg	attgctgttg	ggcaaatggg	gagctatggc	660
tatggatcta	tgaccatgga	gaagagtttt	agggactgct	caatatcagc	tgaggaggt	720
actagtatgtg	gtcatgttgg	tggatcta	aataatattg	gtcacacgt	tggatcatgg	780
gttggtgttg	atccatattc	ttcaccttac	actctctttg	acaagagatc	atcatcaaga	840
caagtgtttg	gtgaccaaga	aaatatgatg	gatgaagaag	atcatgaata	ccaagaaaac	900
ctgcaagatt	cccagagat	tgagaccctc	cctctcttcc	ccatgcatgg	tgaagacatc	960
catggctttg	gcaacatcaa	gtccacctcc	gacagctact	actcggctg	gtatcgctcc	1020
gacgaggcca	acaatggcgc	cacttccctt	gagctcagcc	tcaattccta	cgccacatg	1080
acccctgatt	atttcagctc	atattaa				1107

SEQ ID NO: 19 moltype = AA length = 323
 FEATURE Location/Qualifiers
 source 1..323
 mol_type = protein
 organism = Rubus occidentalis

SEQUENCE: 19

MDCGGPTTNK	DKLLVIMGAT	GAGKSRLSLD	LAACFPPTATF	EVINADKMQV	YKGLDITTNK	60
LPAPERLGV	PHLLGEFDSL	DGEVTPAKFR	ELAAARVVSGV	SARRKVPVLV	GGNSFIHAL	120
VVDRFEPGPN	VFDGDSVTVS	ISSELRYNCC	FLWVDVSSV	LTEYLSKRVD	EMLDSGMLLE	180
LAEFYNSAE	DSVVRTGIRK	SIGVAEFSRY	FRKHPPPGGS	EEVGDDDPVR	REAYKDAVRE	240
IKLNTQLAK	RQIGKISRLR	GAGWNLRLRD	ATEAFRAAVK	SDEKDGNRWS	EIWEREVVGP	300
SVKIVNHFLE	EEIQSTKVQN	QKI				323

SEQ ID NO: 20 moltype = AA length = 357
 FEATURE Location/Qualifiers
 source 1..357
 mol_type = protein
 organism = Rubus occidentalis

SEQUENCE: 20

MTLDSFPPTLH	YSYINITHPS	LSSPAVLTLP	PSPSLPQKPRW	ARMDSGPATT	TTVPLRHKDK	60
LVVIMGATGA	GKSRLSIDLA	THFPSCFEII	NSDKMQVYKG	LDITTNKIPV	PDRIGVPHHL	120
LQGFDPDLGE	LTPAEFRQLA	GQAVSAITSR	RKVPMLVGGG	NSLIHALLVD	RFEPGVDFVE	180
GGAAISSSEL	YNCCFLWVDV	SLTVLADYLS	KRVDEMLDSG	MLEELAEFCD	PEDDLAIRT	240
GLRKAIGVPE	FSRFFKKYPP	ICQSQKARPD	DPVRRGAYKE	AVRAIKDNCT	QLAKTQIGKI	300
LRLRVAGGWD	LQRLDATEAF	RAVVTSEDDA	GKRWSEIWQK	QVVEASVKIV	KRFLEQE	357

SEQ ID NO: 21 moltype = AA length = 374
 FEATURE Location/Qualifiers
 source 1..374
 mol_type = protein
 organism = Prunus avium

SEQUENCE: 21

MTLHPPPTHS	HHHYSNLNFI	SYISRFHLFP	PVNHLCPPTP	SHHHPLPHIK	PRRGAIHSSG	60
PPTPRQQQQR	KDKLLIILGA	TGAGKSRLSL	DLATRPFPEF	IVNSDKMQLY	AGLDITTNKL	120
PIPDRLGVPH	HLLGEFDRH	GDFTPAQFRA	VAGQAISIT	NRKVPMLVVG	GSNSFIHALL	180
VDRFEPGSNV	FEPGFNGSVS	SELRYNCCFL	WVDVSLTVLT	EYLCKRVDEM	LDGMLDELA	240
EPCDPRQDE	DELTAQTAL	RKAIGVPEFT	RYFEKYPPQG	RGGEGRDER	REAYEAVRA	300
IKDNCTQLAK	RQIGKILRLK	GGGWLRLRLD	ATDAFRAVVA	TTSSDKDDGK	RWSEIWERQV	360
VKPSVKVVKR	FLDE					374

SEQ ID NO: 22 moltype = AA length = 446
 FEATURE Location/Qualifiers
 source 1..446
 mol_type = protein
 organism = Rubus occidentalis

SEQUENCE: 22

MEBYNNSNEM	DHESSGTPRV	SASTFDIYII	IIINYINISNI	FFKIKECSIQ	LLFCSLVCIF	60
------------	------------	------------	-------------	------------	------------	----

-continued

FITFLIYDQN	LLMCRGNFLY	ASPNLGGNYG	RTAAGDHNQO	INTPHLQSSG	GEFQSGGTA	120
QPTVKTEAAG	STSQHAPHKF	HHQFPFPPSLI	LSRGQNDQLP	VQQQQHIENE	NEVEAIKAKI	180
IAHPHYSNLL	QAYMDCQVRG	APSEVVARLA	AARQEFEARQ	RSLVSSRDAS	SKDPELDQFM	240
EAYYDMLVKY	REELTRPIQE	AMDFMRKIET	QLNMLGNDSS	TTPLRIFSAS	AQEDKCDGNG	300
SSEEDQDNNS	GGETEVAEID	PRAEDRELKN	HLLRKYSGYL	SSLKQELSKK	KKKGKLPKDA	360
RQKLLSWWEL	HYKWPYPSES	EKVALAESTG	LDQKQINNWF	INQRKRHWKP	SEDMQFMVMD	420
GLHPQNAATL	YMDGHYIGDG	HYRLGP				446

SEQ ID NO: 23 moltype = AA length = 355
 FEATURE Location/Qualifiers
 source 1..355
 mol_type = protein
 organism = Prunus avium

SEQUENCE: 23

MEEYNQMDHE	SSGGRGNFLY	ASPNLGGNYG	RASHDHQMG	NAFHLQSSGG	GDFHSPGGAH	60
PINVKTEAST	SHHGHHKQY	NNNNNNNNNN	PLSSSRGHQP	VVQQQHNLDR	QNDDHTLSSN	120
EVEAIKAKII	AHPQQRSSVA	SREASKDPEL	DQFMEAYYDM	LVKYREELTR	PIQEAMDFMR	180
RIETQLNMLG	NNNAPPLRIF	SPSEDKCEGI	GSSEEEQENS	GGETEVPEID	PRAEDRELKN	240
HLLKKYSGYL	SSLKQELSKK	KKKGKLPKDA	RQKLLSWWEL	HYKWPYPSES	EKVALAESTG	300
LDQKQINNWF	INQRKRHWKP	SEDMQFMVMD	GLHPQNAALY	MDGHYIGDGH	YRLGP	355

SEQ ID NO: 24 moltype = AA length = 700
 FEATURE Location/Qualifiers
 source 1..700
 mol_type = protein
 organism = Rubus occidentalis

SEQUENCE: 24

MNMNWLGFSL	SPQEDDAPIS	HQLADQETLA	SRLGFNSNDQ	IQGAGGEADW	NMKGVDMNSS	60
DGNNYKRTSS	SSDLSPMLMG	SSTTTSTSCS	IISQPNLENH	HQQQLPKLE	NFLGRHSFAD	120
HDQSTSHDY	MFDMMNGPGA	ISSNVMTSN	TNSNNIGLSM	IKTWLRNQP	QPRDNNHQLE	180
QESKTTKESE	NQPQSSLSLS	MSTGSLHTVT	TATAGGASAS	AGAGETTSSS	NNNRKQSPVL	240
TATTTDPQTQ	TTGGAIEAVP	RKAIDTFGQR	TSIYRGVTRH	RWTGRYEHL	WDNSCRREGQ	300
TRKGRQGGYD	KEDKAARAYD	LAALKYWGTT	TTTNFPISTY	EKEIDEMKPM	TRQEYVASLR	360
RKSSGFSRGA	SIYRGVTRHH	QHGRWQARIG	RVAGNKDLYL	GTFSTQEAAA	EAYDIAAIKF	420
RGLNAVTFND	MSRYDVKSIL	ESSALPITTG	ASAKRLKDVQ	QQPPPPADHH	HQIMLSSVLD	480
HHQMIRSSS	STEHDIMGNI	FHQDDHLRQ	QLQMGGTHNF	FHDHHDQORS	GLMGLMDSSA	540
AAASMEHSSG	SNSVIYSVGD	HGNNNGYGS	NARGTSGSYI	MPMAMSTVVA	NHGQNQADGN	600
NINGFGDGDG	DDQEVNIKAQ	QLGYDHHQNV	FLGSSNTTDP	AYQHHASNRN	LYYQLPVHDD	660
QHSSSEAVA	TSSTTCNMNW	VPTAVPHTLA	HSTFTVWNT			700

SEQ ID NO: 25 moltype = AA length = 532
 FEATURE Location/Qualifiers
 source 1..532
 mol_type = protein
 organism = Prunus avium

SEQUENCE: 25

MGSMNWLGFPS	LSPQEVVPAD	PSISEGDHHH	QHHDGHELS	DQTATLVSR	GFKYSDDISG	60
TDVTGGCPDL	TSHDDSSATT	TAAAVPNNHS	HNHLIPPPFG	MLHQAFAFNT	TNIHSQGADW	120
SNMSSDSST	YKTTSDHLSM	LMGSSSYRS	QNLNNHQOP	KLENFLGRHS	FVADEHHIGG	180
QSSSGTHAYN	NNSTNTIQT	LSLSMSTGSH	THHNTTSTAA	AGETCSSDRD	NTCNNNNKLA	240
VVRTTASTAP	PPGIDSQTTS	TTATAIEAVP	RKSVDTFGQR	TSIYRGTDGQ	VDMLISGII	300
VAGEKDKLAR	BGKISNYEKE	IDEMKHMTRQ	EYVASLRRKS	SGFSRGASII	RGVTRHHQHG	360
RWQARIGRVA	GKNDLYLGT	SSNSVMYSTA	TTTNGGGGGG	GGDHGGYIIP	MGTVISNDHN	420
DGFGDGTNHD	HQVKAVALGF	ENVFGSTSAS	VNATNVTRDA	YNNHARNLYY	LPQQLPQSSS	480
VSSVSSAGVA	KGGSAYDIHE	AGQCNNWMP	AVPTSTNSNI	NMPPTFTVWN	DT	532

SEQ ID NO: 26 moltype = AA length = 297
 FEATURE Location/Qualifiers
 source 1..297
 mol_type = protein
 organism = Rubus occidentalis

SEQUENCE: 26

MEPQSQQPTE	DGGSNKAAGS	ANMLCRQSST	RWTPTTDQIR	ILKELYNNKG	VRSPSAEQIQ	60
RICLQLKRYG	KIEGKNVFYW	FQNHKARERQ	KKRFTSDVHV	PMQRSGLVGN	SNVANWKPED	120
QYLNMMNKYS	NICQGVSSSA	SASASSAGVV	SFNGQMGNYG	VYGSMMNEKS	FRDCSISAGG	180
GTSTTGVGIG	GSISHNYGSW	VGGSVDPYSS	SSYTTTSTTF	FDNQKQIFME	EEEEHDQIE	240
TLPLPFMHGE	DIFGNIKTTS	EGGGGDYGY	SSGWSGSRT	SLSLSLNSYG	YNNPNYY	297

SEQ ID NO: 27 moltype = AA length = 305
 FEATURE Location/Qualifiers
 source 1..305
 mol_type = protein
 organism = Prunus avium

SEQUENCE: 27

MEPQQTQQQP	NEDGSNKGTS	GTNFLCRQSS	TRWTPTSDQI	RILKDLYYNN	GVRSPSAEQI	60
QRISARLRQY	GKIEGKNVYF	WFQNHKARER	QKKRFIDVPA	PPIMQRSGLG	INNNAPTAYE	120

-continued

PINHNSNKYPN	ISASAGVSSA	SASSAGVIAV	QOMGSYGYGS	MTMEKSPRDC	SISAAGGTSS	180
GHVGGSNMNI	GHNVGSWGV	DPYSSPYTLF	DKRSSSRQVF	GDQENMMDEE	DHEYQENLQD	240
SPEIETLPLF	PMHGEDIHGF	GNIKSTSDSY	YSGWYRSDEG	NNGRSTLELS	LNSYGHMTPD	300
YFSSY						305

SEQ ID NO: 28 moltype = DNA length = 723
 FEATURE Location/Qualifiers
 source 1..723
 mol_type = other DNA
 organism = Agrobacterium tumefaciens

SEQUENCE: 28

atggatctgc	gtctaatttt	cgggtccaact	tgcacaggaa	agacgtcgac	cgcggtagct	60
ctgtcccagc	agactgggct	tccagtcctt	tcgctcgatc	gggtccaatg	ttgtcctcag	120
ctgtcaaccg	gaagcggacg	accaacagtg	gaagaactga	aaggaacgag	cgtctctatac	180
cttgatgac	ggcctctggt	gaaggggtatc	atcgacagcca	agcaagctca	tgaagggtcg	240
atgggggagg	tgtataatta	tgaggccccc	ggcgggctta	ttcttgaggg	aggatctatc	300
tcgttgctca	agtgcatggc	gcaaaagcagt	tattggagtg	cggatttttcg	ttggcatatt	360
attcgccacg	agtttagcaga	cgaagagacc	ttcatgaacg	tggccaaggg	cagagttaag	420
cagatgttac	gccctgctgc	aggcctttct	attatccaag	agttgggtga	tctttggaaa	480
gagcctcgcc	tgaggcccat	actgaaagag	atcgatggat	atcgatatgc	catgtgtgtt	540
gctagccaga	accagatcac	atccgatatg	ctattgcagc	ttgacgcaga	tatggaggat	600
aagttgatcc	atgggatcgc	tcaggagtat	ctcatccatg	cacgccgaca	agaacagaaa	660
ttccctcgag	ttaacgcagc	cgcttacgac	ggattcgaag	gtcatccatt	cggaatgtat	720
tag						723

SEQ ID NO: 29 moltype = DNA length = 723
 FEATURE Location/Qualifiers
 source 1..723
 mol_type = other DNA
 organism = Agrobacterium tumefaciens

SEQUENCE: 29

atggatctgc	gtctaatttt	cgggtccaact	tgcacaggaa	agacgtcgac	cgcggtagct	60
ctgtcccagc	agactgggct	tccagtcctt	tcgctcgatc	gggtccaatg	ttgtcctcag	120
ctgtcaaccg	gaagcggacg	accaacagtg	gaagaactga	aaggaacgag	cgtctctatac	180
cttgatgac	ggcctctggt	gaaggggtatc	atcgacagcca	agcaagctca	tgaagggtcg	240
atgggggagg	tgtataatta	tgaggccccc	ggcgggctta	ttcttgaggg	aggatctatc	300
tcgttgctca	agtgcatggc	gcaaaagcagt	tattggagtg	cggatttttcg	ttggcatatt	360
attcgccacg	agtttagcaga	cgaagagacc	ttcatgaacg	tggccaaggg	cagagttaag	420
cagatgttac	gccctgctgc	aggcctttct	attatccaag	agttgggtga	tctttggaaa	480
gagcctcgcc	tgaggcccat	actgaaagag	atcgatggat	atcgatatgc	catgtgtgtt	540
gctagccaga	accagatcac	atccgatatg	ctattgcagc	ttgacgcaga	tatggaggat	600
aagttgatcc	atgggatcgc	tcaggagtat	ctcatccatg	cacgccgaca	agaacagaaa	660
ttccctcgag	ttaacgcagc	cgcttacgac	ggattcgaag	gtcatccatt	cggaatgtat	720
tag						723

SEQ ID NO: 30 moltype = DNA length = 957
 FEATURE Location/Qualifiers
 source 1..957
 mol_type = other DNA
 organism = Arabidopsis thaliana

SEQUENCE: 30

atgaagtgtg	atgacaaaat	gggtgtgac	atgggtgcaa	cgggttctgg	caagtcatac	60
ctctctgttg	atctcgtttt	acatttttaa	gccgagatca	tcaactctga	caaaatgcag	120
ttctacgatg	gcttgaagat	caccacgaat	caatcgacca	ttgaagatcg	acgtggagtg	180
ccacatcacc	ttctcgttga	actaaaaccg	gaggctggag	aagtcacagc	ggcagaattt	240
cgcgttatag	cggttgaagc	catctccgag	attactcaac	gtaaaaagct	cccaatcctt	300
gccgttggtg	ccaactcata	cattcatgct	ctccttgcaa	aatcttatga	ccctgaaaac	360
tatccgtttt	ctgatcacaa	gggctcaatc	tgctccgagt	tgaatatatga	ttgtttgttc	420
atgttgatag	atgtggatca	gtctgtgtta	ttcgagatc	tttctttacg	tttgatctt	480
atgatgaagt	caggtatgtt	cgaggagatc	gctgagttcc	accgctctaa	gaaggccccc	540
aaagagccat	tgggggatatg	gaaggctata	ggagtgcaag	agtttgatga	ctacctcaaa	600
atgtacaagt	gggacaatga	catggataaa	tgaggacccta	tgagaaagga	ggcttatgag	660
aaggcggtga	gagccatcaa	agaaaacaca	tttcagctca	caaaggatca	aatcacggaag	720
atcaacaagc	tgagaaatgc	cgggtggggc	ataaagaagg	tgatgctac	agcatcgttt	780
cgagaggcaa	ttagggcagc	caagggaagg	gaaggtgtag	cagagatgca	gagaaagata	840
tggaacaagg	aagtgttgga	accgtgtgtg	aagattgtca	ggagccactt	ggaccaaccg	900
atcaactatt	attattatta	cttttattta	ctaaaaagat	ttttaagtct	taactag	957

SEQ ID NO: 31 moltype = DNA length = 956
 FEATURE Location/Qualifiers
 source 1..956
 mol_type = other DNA
 organism = Arabidopsis thaliana

SEQUENCE: 31

atgaagtgtg	atgataaaat	gggtgttatt	atgggtgcaa	cagggttctgg	caagtcttca	60
ctgtcagttg	atctggcttt	gcattttcaag	gccgaatata	tcaattctga	taagatgcag	120

-continued

ttctacgatg	gcctgaagat	tactaccaac	cagtcaacta	tcgaggatag	acgcgggtgtg	180
ccacatcacc	tggtgggcga	attgaaccct	gaggctggag	aagtgaacgc	tgccgagttt	240
agagttatgg	cagctgaggc	tatttctgaa	atcacacagc	gcaagaaact	tccaattctg	300
gcccggcggt	ctaattcata	tatccatgca	cttctggcta	agtcttacga	tcagaaaaac	360
tatcctttct	cagatcacaa	aggatctatt	tggtcagagt	tgaagtaaga	ttgctgtttt	420
atttggtatg	atgtggatca	gtctgttctt	ttcagatata	tttcaactgc	cttggatctt	480
atgatgaaat	ctggaatgtt	tgaggaaatc	gcagagtcc	ataggtcaaa	gaaagctcca	540
aaggaaacct	ttggcatttg	gaaagcaatc	ggagtgcagg	aattcgatga	ttacctgaag	600
atgtataagt	gggataatga	tatggataag	tgggataccta	tgccgaagga	agcttacgaa	660
aaagccgtta	gggcaattaa	agagaacaca	ttccagttag	ctaaggatca	gatcactaag	720
atttaataagc	ttagaaaagc	cggatgggat	attaagaaag	tggatgctac	cgctcattc	780
agggaaagcca	tcagagccgc	aaaggaaggc	gaggggtgtg	cagagatgca	gaggaagatc	840
tggataaagg	aagtgtctga	acctgcgtg	aaaattgtta	gatctcactt	ggatcagcct	900
atcaattact	actactacta	ctctattttg	cttaagcgct	tcctgtcatt	gaactg	956

SEQ ID NO: 32 moltype = DNA length = 7908

FEATURE Location/Qualifiers

source 1..7908

mol_type = genomic DNA

organism = Zea mays

SEQUENCE: 32

agtgcactgt	ctctctccaa	aggcagctgc	attggcctcc	agcctttttc	ctactgtgcc	60
gcgcgccctc	cctctctctc	aatgatagca	tagggagaga	aggcatactc	cgaggcatcc	120
ttctcctttc	cctctccttc	cccaaacctt	ttctcctttt	ccctcgcccc	aagaacttca	180
ttctatctcc	agcgccctcc	tttgcgttg	cgcaggagga	gctcacgggg	acagtgggcg	240
gagagctcga	tcgctgcacc	actacttcac	tggagggtcc	cccactccca	tggaggagat	300
caccacaacac	tttggagtgt	gcgcaagcag	ccacggccat	ggccacggcc	agcaccacca	360
tcataccacac	caccaccacc	cgtgggcctc	ctccctcagc	gccgtcgtag	gcgcgtgccc	420
gcccgaaccg	ccaagcgcag	gcctgcgcgt	gacctgaac	acgggtggcg	ccactgggaa	480
cagcggcggt	agcggcaacc	cgggtgctga	gcttgccaac	gggtggcgcc	tcctcgacgc	540
atgctgaag	gcgaaggagc	cctcgtcgtc	gtctccctac	gcaggcgagc	tcgaggccat	600
caaggccaag	atcatctcgc	acccacacta	ctactcgtc	ctcactgcct	acctcgagtg	660
caacaaggca	agcaagacta	gtagccgcag	tctaacaata	gctctcctgc	ttccggttca	720
gctactgctg	ctagcgcaagt	ttggccacat	ctgatgtcaa	gagcatggcg	catgcacata	780
ctttctgctt	gcttttatcat	agcatactat	agtttatgca	tgctgctcat	gtagacatgt	840
gcatctggtg	ttcggaaactt	tctcaaacgg	ccgtgggtgt	ttgcagggtg	gggcaccacc	900
ggagggtgtg	gcgaggctga	gcagagatag	gcaggaggtg	gaggcgcgcc	agcgcacggc	960
gctcggcgcc	ctcggcgctg	cgaacggagc	ggagctggac	cagttcatgt	tgaggcatct	1020
ttaggtttat	ttcacgcgcg	cgcagcgcac	cattattagt	cgttttcagc	tcgggctcgg	1080
atttaatttg	cgtgtatatata	tgttaggagg	cgtaccacga	gatgctgggt	aagttcaggg	1140
aggagctgac	gaggccgctg	caggaggcga	tggagttcat	gcgaagggtg	gagtcgcagc	1200
tgaactcgct	ttccatctcc	ggaaggtcgc	tgcgcaacat	cctttcatct	ggtaactgaag	1260
ttgctgcggc	ccatgcacct	cttattttat	attagatgat	ttcttggatc	ggttgtcgcc	1320
tcatgcagcg	atgcattgcc	tcctctctgt	taaaatcttc	ctgtcctctc	cttcgtcttg	1380
ctgtcctctt	tcctgggttg	gcataatgtg	atgggttttt	ttttttgcat	ttcaaaaggg	1440
ttaaattttt	ccttgatctg	gtttctcagc	aattaaattt	ggcatctgtg	tagttccctt	1500
tttaagagca	tcttgctatg	catgcataat	ttgagtatag	atagctgcgg	aacgcagatg	1560
aatgtcttgt	ctgcgcttca	tgttttttaa	tgcgcggcag	cattctcaat	cagggggttc	1620
gaaagagaaat	cttctccctg	cttgcctttc	ctggccatgc	atctgtgaga	ttcttctctc	1680
tggcttcctg	cgatctttgc	ttctgctgga	aaggaaaatg	tcctcagtc	cacatgtgca	1740
ggagcgcagc	ttcaatctag	catggctgaa	agctctctgg	ttctttccat	acactccatc	1800
ctgcacattc	cttgcatata	ttctgctgtt	aaactgccta	gcgacgagga	tgaatgaaaa	1860
cacctgtctt	ttttactagt	gggagtaact	atttggctgc	tggtgcccag	gatgcgttgt	1920
tcactgctct	ccatttatatt	gatcgattcc	atttggaatg	gaacatgtgc	atgtgttctg	1980
cgagttagtt	ctggctcagc	gcagccatct	tttcccatga	tgggaacccc	agggagttcc	2040
cttgttctag	gggtttggaa	agggccacag	tgtactgggt	gctgtggcat	tgtgcaagca	2100
cagtcctatg	ccgaggagtc	tcactggcgt	atgtcagtag	aacatagtag	tatacacgtt	2160
gtagtgtggg	gagatcagag	agagagagag	atggggagtc	atccccatc	tgtcacgcaa	2220
gcttgtggcc	agaacattct	tgtcatcttt	ctccccctca	agagatgtag	ctacctgttg	2280
tcctagaatc	tatagtgttg	tgtgtgtgtc	ttccagggtc	gcacacagat	ctggccactg	2340
ctggccttag	tatttgctaa	ttttaaaatg	aacacatcag	gtagatcgag	acgatccatg	2400
gcaagcgtct	ttctaattgac	ctatcacaa	gctatagtaa	caactcgtac	caccacagtt	2460
acgcacagcc	cagagttttt	cacttctcgg	ttctgttctt	cttagagcac	tggctatggc	2520
tagtatatat	gcattggcagc	aacacacaca	cacacacaca	atcacttcat	tactggagca	2580
agttagcgaa	gaagctgcga	tctgatgtcg	tagaatgctg	caaaaatgaa	aggttcagag	2640
gcaggcctac	gggttgaaat	ggagcaccgt	atgcggcgcc	ccagtttttt	gtgttctgac	2700
cggcgagctg	gacaaaatgg	cctgtgtgct	tcgagaacca	tagagacagt	tgccatagcg	2760
atgagcgctg	agcggccctt	attatttgat	gtgagatctc	tctatcaatg	gatgtgtgat	2820
ctgctagggt	tatattatat	atgcgtctac	tatatcccta	ccagtccctg	tattttctga	2880
cagatagact	tctccccgcg	gcgttctctc	ataaataaat	gacggctcaag	gaacatttga	2940
tgccttccga	ggcgagctag	tacaagtttc	atctgcgttt	gcgtttgtat	ttttcagctg	3000
tcaactcaga	ttttctctca	gctccaaactg	ttctttgacc	ttcttggtag	tgttttcttg	3060
gaagaaacta	tttgttgacc	atttttgggt	ataataaaggc	cacagacaaa	ctgttgaaat	3120
taatcacaaa	acaacactac	tgcgtttttt	cttttctttt	ctgactgatg	ttgcacatgt	3180
actccaagat	tcttgggttg	atgcactctt	tcaaggctcat	gcaaaagcag	ggacacggcg	3240
tgtcacgagt	ttccgtttgt	ctcgaggtaa	gacacaaaaa	gataccggaa	gcaaacacaa	3300

-continued

taaaacaaa	ggtaggatgg	aaaagagctg	atggatttta	caatgtatcg	ccagagtggg	3360
aaagacaa	cgactagagg	aaaggaacga	tctggcactt	ccaactgtgt	gaattcttcc	3420
atgtgatgtg	cgtctcgtgc	atgtttat	tcttggttac	tggaaatcgg	gtgcacccat	3480
ttttattcag	gattcaatta	ttggttactg	gagccaggtc	cgttcgtttt	tattttcctt	3540
ttgagattga	ctggatcgat	tctctgggtc	tagtcactta	tgtctgttgt	tttatcggtt	3600
tcatggattt	ctattctatc	tctcccaata	ttcagtatca	cataaaatcc	tgtgactaga	3660
tttctttatc	acatacaatc	ttgttttttt	attttcttta	atttaacact	ggatttgggt	3720
gataatatgt	tctgaaaaac	gttaataaat	ttaagcaca	cagcctttta	aaaaatgta	3780
taattattta	ccttgaaacc	cagcatcggc	cgggaacaa	gaaagaact	caaggtttta	3840
aatgcattga	aatgaaacat	taatggagtg	tttggtttga	ataatgatgt	agtcacatcat	3900
cttctcactc	ctcacttttt	tggttgggtt	atggaatgga	gtgagttaat	ccatcaacac	3960
ctcattccctc	atagtttagtt	gttttagtact	aatatgtgga	atgaagtcat	cccaccaa	4020
ttgaagaatg	gactcatgat	gcaccacttc	attttagata	gagtgtattca	tcaaaccaaa	4080
caacctcata	gggcatctag	ggatcctagg	agctaaaaa	aaagtgtact	aagttttagtc	4140
acttttaggag	ctaaagatct	actaaatagg	aaactaaaa	tgactagaat	agtaaaaggt	4200
atcttttttag	tcattttttt	ctcctaagaa	ggagctaaat	tttagttagt	ttggttttagc	4260
tcttggatcc	aaacaggccc	taagaaaatg	tttcgacaa	atttgggaca	ataaaacagt	4320
tctaaggat	ttctttggca	agattaggcc	ttctttggaa	caaaagaaaa	tgaaggaa	4380
ttgaaggatt	gaaatccat	cgaagctttt	catatgcaa	gaattgtgtt	cctaggatga	4440
tttctaacaa	gaggctcatc	cccttgaaaa	ttgttctttg	tgtctatctc	tctcctctaa	4500
ttcatgttgt	cttatgttgc	attgaaacac	tattagaaaa	ttttcatgtg	ttttaattta	4560
tgtatgatgt	taagtgtcaa	gcagcactat	tctacatttt	tttctattcc	tgtgttttat	4620
cgatactgca	tccaatgaa	ggcctaagcg	tatcatgcaa	tcttttccag	aagtttttta	4680
gattgctgta	taatgacaag	gtgcaatgct	cttcgttttc	tttttgggtt	ttcctcattt	4740
tcagcaaggc	caccaaattt	ttccatgatg	ttcttgtctt	attcctgtag	tatccctcga	4800
aaaaatttca	tagcttttcta	catccatggt	atctaatgat	gttcttcatt	tcaaaagacg	4860
acacatttgg	cataagttct	gattgtgtac	taaaactttg	tggtacggtc	atttactttt	4920
actatacaca	tgtactgtat	tcatgcgtac	attgtacttt	gtccattggg	gcctttcttc	4980
tttttaatac	aacgggcatc	gtttcaaaaa	aaatgtgtca	gtcgtaaact	cacacttctt	5040
tagttttccc	cagttaaacg	catgacggaa	tagagttaca	gagtcctcat	tcaagtcaca	5100
tatatctcta	actcgtttct	aaaattaaag	taatattttc	attgttgctt	ccgtaagaat	5160
gcaccacaaa	cacaaaatat	catttctctt	atgcaaatat	atataatatga	atgtttgttt	5220
ctggcagtg	tgagaagctt	ctcactgagc	caaagggttt	agggatgcca	tgtgtctctt	5280
ttctttttta	agtgtagatt	aagcaaacct	ttataatttt	taaaactcca	ccagatacct	5340
tttctaatt	cataaatgct	ctggactatc	ttcacgaaaa	aggtcttggg	aatatgttca	5400
caagcctcct	ttatctgtta	tcttgggcta	gacataggat	gtgttggaag	aatacaagcg	5460
gagttgtttc	gtttaaagcaa	aaaaaaaact	gtaactttat	attggagaat	atacatcgtt	5520
ctcccatatt	tgttacagtg	ctcataaata	gacaaagttt	tttttttgaa	atcttaagcc	5580
ctgtgtttct	ttgtttatct	tatctctata	ttgtctccaa	atatgttgtc	atcaagatgg	5640
atatagataa	ttgtcaaaac	gaaaagtgtg	catggttcat	ttatgccaat	tctgaaaagc	5700
ataagttaaa	tattaagata	caataaaaga	aaaacatatg	cagtgttgac	tgttgggaaa	5760
agaacaaggt	ctgcacatac	acttgcaata	taaatctttt	taccagcaca	taaaaaaacg	5820
atcaaatatc	acgcaaacaa	agtaatacgt	ccagaaacac	ccatactttt	gaaattcgtc	5880
tttgaatg	agtaggttca	caaaaataaa	tggtgcaact	gcacatgttc	tttatatctg	5940
tactcaatgg	gttttctgga	gatgttagat	tgtattgggtg	gacaatatcc	taatgcaact	6000
cgcaaatcc	caaggccgaa	ctcaagtggg	agtttggatg	attttttttg	taaaagaagc	6060
gatggctg	ctagtttaca	tagaccatca	cacttcacaa	ttcacattca	tgaagtcata	6120
actttgttca	cttctgtgtt	aactatggca	tataaaaaa	tactgtgtgt	tgttttgcct	6180
gtatggaaca	tgtattttag	ttttttagta	aataaagtcca	gtaaatgtct	ttgccggaca	6240
aatttcacca	acttggctac	gaatgatgct	tgtacacttc	ttttgtttta	tctaaaagtt	6300
ccatgtctta	tgtctgaagg	ctcttctgag	gaggatcaag	aaggtagcgg	aggagagacc	6360
gagctccctg	aagttgattg	acatggtgtg	gaccaagagc	tgaagcacca	tctcctgaag	6420
aaatacagtg	gctatctaag	ctcgtccaag	caagaactgt	caaagaagaa	gaagaagagg	6480
aagctcccca	aggaggtctg	ccagcagctc	cttagctggg	gggacagca	ctacaaatgg	6540
cttaaccctc	cagtacgtct	tctttttatt	cttccatttt	aactattggt	ggtgacacat	6600
gatttagacg	atgccaatc	ttcatgaact	tttcatagcc	agctacccaa	tgttagtact	6660
gactgcacat	tgtaatcaa	gggttaagat	atatatacat	aaatccattt	tggcaaatct	6720
aagctacata	tgggtctttg	atcttccatg	acggtctgtt	gatctctgat	ttgcataatg	6780
gcataataaa	agtgaagcaa	aatatgtcag	agtcataata	tattgatcag	ggagtggcag	6840
gtgattattg	gtatttaattt	aaacttattt	aaggtatttt	gaaacttctg	tagcgtttctt	6900
actaaatacc	attgattttta	atttaagcaa	ctatatattt	atctgggtgaa	aaatgaagcc	6960
ttttctgata	tacaaattga	agagtctaca	atggtttcac	ttacatggct	gaaacagaaa	7020
atcatagtgc	cctgaattgt	gtgttgatac	tcaaaagcgc	agattcaaat	ttgtaatttt	7080
caagtttagg	gttctaagtg	aaaaaaaac	attgagttca	ggagcataca	ctgaactttt	7140
ttttatcat	atcttcattt	tgttgatgtg	ttgtatacgc	gcataatagc	tgtgcttccc	7200
tactggatgt	gaattaaaca	actcttccca	tccgtgagca	ggagactcag	aaggtggcac	7260
tggctgagtc	tacgggctt	gacctgaagc	agatcaacaa	ctggttcac	aaccagcgga	7320
agcggcactg	gaagccatcc	gaggagatgc	accacctgat	gatggacggg	taccacacca	7380
ccaatgcctt	ctacatggac	ggccacttca	tcaacgacgc	gggctgtac	cggtcggct	7440
agccaccggt	atctcgtctc	catcttcacac	cccacggcct	agctataaag	actaatgggt	7500
ccaggtgtct	gaagtactga	agacaggggg	gctagctatc	taattgttgt	gccgacgcga	7560
tgagctgtaa	ggaggccatg	cttaattatt	ctgttgccgt	tgtactctta	tctatatcgc	7620
cctatgcctc	cgtcatgaa	ctatgcttta	ggtgtgtgct	gctccacact	gtggtgtgtg	7680
gcttttgctt	ttgtgtggtc	gtattgtatg	cgtaacctga	cagatggatc	cctgattgct	7740
acatgtttga	ataatttgca	tgtactagct	agtttctgcc	taatctaag	gtacggctca	7800
tgtcttgtca	gggtgatttg	ttatgtgttt	tttttttcaa	attgaggtct	gtttgttttg	7860

-continued

ctctcaatcc atccatgtga attgggtgga attaaatgag tttaaatt 7908

SEQ ID NO: 33 moltype = DNA length = 1077
FEATURE Location/Qualifiers
source 1..1077
 mol_type = other DNA
 organism = Zea mays

SEQUENCE: 33

atggaggaga	tcacccaaca	ctttggagtt	ggcgcaagca	gccacggcca	tggccacggc	60
cagcaccacc	atcatcacca	ccaccaccac	ccgtgggcat	cctccctcag	cgccgtcgta	120
gcgcgcgtgc	cgccgcaacc	gccaaagcga	ggcctgcccgc	tgaccctgaa	cacgggtggcg	180
gccactggga	acagcggcgg	tagcggcaac	ccggtgctgc	agcttgccaa	cggtggcggc	240
ctcctcgacg	catgcgtcaa	ggcgaaggag	ccctcgctcg	cgtctcccta	cgcaggcgac	300
gtcgaggcca	tcaaggccaa	gatcatctcg	caccacacac	actactcgct	cctcactgcc	360
tacctcgagt	gcaacaaggt	ggggggacca	ccggagggtg	cggcgaggct	gacggagata	420
gcgcaggagg	tggaggcgcg	gcagcgacg	gcgctcgccg	gcctggccgc	tgcgacggag	480
ccggagctgg	accagttcat	ggaggcgtag	cacgagatgc	tggtagaagt	cagggaggag	540
ctgacgaggc	cgctgcaggga	ggcgatggag	ttcatgcgaa	gggtggagtc	gcagctgaac	600
tcgctttcca	tctccggaag	gtcgctgcgc	aacatccttt	catctggctc	ttctgaggag	660
gatcaagaag	gtagcggagg	agagaccgag	ctccctgaag	ttgatgcaca	tgggtgtggac	720
caagagctga	agcaccatct	cctgaagaaa	tacagtggct	atctaagctc	gctcaagcaa	780
gaactgtcaa	agaagaagaa	gaaagggaag	ctcccacaag	aggctcgcca	gcagctcctt	840
agctgggtgg	atcagcacta	caaattggct	taccctcag	agactcagaa	ggtggcactg	900
gctgagtcta	ccgggcttga	cctgaagcag	atcaacaact	ggttcatcaa	ccagcggaa	960
cggcactgga	agccatccga	ggagatgcac	cacctgatga	tggacggata	ccacaccacc	1020
aatgccttct	ataggacgg	ccacttcac	aacgacggcg	ggctgtaccg	gctcggc	1077

SEQ ID NO: 34 moltype = DNA length = 1755
FEATURE Location/Qualifiers
source 1..1755
 mol_type = other DNA
 organism = Arabidopsis thaliana

SEQUENCE: 34

atgaactcga	tgaataactg	gttaggcttc	tctctctctc	ctcatgatca	aaatcatcac	60
cgtacggatg	ttgactcctc	caccaccaga	accgcgtag	atgttgccgg	aggggtactgt	120
tttgatctgg	ccgctccctc	cgatgaatct	tctgcgcttc	aaacatcttt	tctttctcct	180
ttcgtgtgca	ccctcgaaagc	tttcaccaga	gacaataata	gtcactcccc	agattgggag	240
atcaatgggt	gtgcattgca	taacattaac	aataacgaac	aaaatggacc	aaagcttgag	300
aatttctctg	gccgcaccac	cacgatttac	aataccaacg	agaccgttgt	agatggaaat	360
ggcgattgtg	gaggaggaga	cggtgggtgt	ggcggtcac	taggcctttc	gatgataaaa	420
acatggctga	gtaatcattc	ggttgtctat	gctaatactc	aagacaatgg	taacggtgca	480
cgaggcttgt	ccctctctat	gaattcatct	actagtgtga	gcaacaacta	caacaacaat	540
gatgatgtcg	tccaagagaa	gactattgtt	gatgtcgtag	aaactacacc	gaagaaaact	600
attgagagtt	ttggacaaag	gacgtctata	taccgcggtg	ttacaaggca	tcggtgggaca	660
ggtagatcac	aggcacattt	atgggacaat	agttgcaaaa	gagaaggcca	gactcgcaaa	720
ggaagacaa	tttatctggg	aggttatgac	aaagaagaaa	aagcagctag	ggcttacgat	780
ttagccgcac	taaagtattg	gggaaccacc	actactacta	acttcccctt	gagtgaatat	840
gagaaagagg	tagaagagat	gaagcacatg	acgaggcca	agtatgttgc	ctctctgcgc	900
aggaaaagta	gtggtttctc	tcgtgggtgca	tcgatttatc	gaggagtaac	aaggcatcac	960
caacatggaa	ggtggcaagc	taggatcgga	agagtcgccc	gtaacaaaga	cctctacttg	1020
ggaactttcg	gcacacagga	agaggctgct	gaggcttatg	acattgcagc	catataaattc	1080
agaggattaa	gcgagtgac	taacttcgac	atgaacagat	acaatgttaa	agcaatcctc	1140
gagagcccca	gtctacctat	tggtagttct	gcgaaacgct	tcaaggacgt	taataatccg	1200
gttccagcta	tgatgattag	taataacgtt	tcagagagtg	caaataatgt	tagcggttgg	1260
caaaaacactg	cgtttcagca	tcacaggga	atggatttga	gcttattgca	gcaacagcag	1320
gagaggtacg	ttggttatta	caatggagga	aacttgtcta	ccgagagtag	taggggttgt	1380
ttcaaacaa	aggaggaaca	acaacacttc	ttgagaaact	cgccgagtag	catgactaat	1440
gttgatcatc	atagctcgac	ctctgatgat	tctgttaccg	tttgtggaaa	tgttgttagt	1500
tatggtggtt	atcaaggatt	cgcaatccct	gttgaacat	cggttaatta	cgatcccttt	1560
actgctgctg	agattgctta	caacgcaaga	aatcattatt	actatgctca	gcatcagcaa	1620
caacagcaga	ttcagcagtc	gccgggagga	gattttccgg	tggcgatttc	gaataaccat	1680
agctctaaca	tgtactttca	cggggaaggt	ggtggagaga	gggtcccaac	gttttcagtt	1740
tggaaacgaca	cttag					1755

SEQ ID NO: 35 moltype = DNA length = 1755
FEATURE Location/Qualifiers
source 1..1755
 mol_type = other DNA
 organism = Arabidopsis thaliana

SEQUENCE: 35

atgaattcta	tgaataactg	gttgggcttt	tctctttcac	cacatgatca	gaaccatcac	60
aggactgatg	tggattcttc	aactaccaga	accgcgtag	atgttgccag	cggtactctg	120
tttgatttgg	ctgcccacac	agatgaatct	tcagcagtag	agacttcttt	tctgtcacct	180
ttcggcggtta	ctcttgagcg	tttcaccaga	gataataact	ctcactcagc	cgattgggat	240
attaatgggt	gcgcctgcaa	taacatcaat	aacaatgagc	agaatggccc	taagctggaa	300
aacttcttgg	gaagaacaac	taccatctat	aataactaac	aaacggtggt	tgatggtaat	360

-continued

```

ggcgattgtg gaggtggcga tggaggtggc ggaggttctc ttggtctgtc aatgatcaaa 420
acctggctgt ctaatcattc agtggctaata gccaaaccacc aggataatgg caacggtgct 480
cgcggtctgt ctctttctat gaattcttca acatctgatt caaacaatta caacaataac 540
gatgatgtgg ttcaggagaa gactattgtg gatgtggttg aaacaactcc aaagaaaaca 600
attgagtctt ttggtcagcg cacttcaatc tacaggggcg ttacaaggca tagatggact 660
ggacgctatg aagctcactt gtgggataat tcttgaaga gggagggcca gactcgcaaa 720
ggaaggcagg tttacttggg cggatatgat aaggaagaga aagcagctag ggcctacgat 780
cttgccgcac tgaatatatt gggcaccaca actaccacaa attttctctt ttctgaatac 840
gagaaggaa gaggagagat gaagcatatg acccgccagg agtatgttgc ttctctgaga 900
agaaagtctt caggtttctc taggggcgcc tcaatctata ggggtgtgac aagacatcac 960
cagcacggca gatggcaggg tagaatcgga cgcgttgccg gtaataagga tctgtacttg 1020
ggcacatttg gaactcagga agaggctgcc gaagcatatg atattgcagc tatcaagttt 1080
agaggcttgt ctgctgtgac taacttcgat atgaatcgct ataactgtta agccattctt 1140
gagttctccat cactgcttat cggatcttca gccaaagcgc ttaagatgt gaataaccca 1200
gttctctgaa tgatgatctc taataacgtg tctgaatcag caaataacgt ttccaggatgg 1260
cagaataccg cttccagca taccagggt atggatcttt cactgttgca gcagcagcag 1320
gaaagatatg tgggatatga caatgtgttc aacctgtcta ccgagtcac aagggtttgc 1380
ttaagcagg aaggaggagc gcagcatttc ttgagaaatt ctccatcaca catgacaaac 1440
gttgatcctc actcttcaac ctctgatgat tcagtgcagc ttgtggcaa tgtggtttct 1500
tacggaggtt atcagggttt tgcatttcca gtgggcacct cagtttaacta cgatcctttc 1560
acagccgagc agatcgcata caacgctaga aaccattact actacgcccc gcaccagcag 1620
cagcagcaga ttcagcagtc ccaggcgga gatttccctg tggcaatctc aaataacccat 1680
tcttcaaaaa tgtacttcca cggagagggt ggcgagagaag gtgctcctac ttctctgtt 1740
tggaacgata cctga 1755

```

```

SEQ ID NO: 36      moltype = DNA length = 1986
FEATURE            Location/Qualifiers
source              1..1986
                    mol_type = other DNA
                    organism = Triticum aestivum

```

```

SEQUENCE: 36
atggcttctg gaaacgactg gttgggggttc tctctctcgg gccaaagaaa cccacagccg 60
catcaggata gcccgccgccc cgcgcgcatc gacatctccc gcgccagtga cttctatggc 120
gtaccacgcg cccagctccc cgacgctcaa ctcgcggtgc caggtcatca gccttcatac 180
gggatacagg aggacttcaa taggggagct catgaaaccc acgattggaa catgaggggg 240
gatctggact ataacggcgg tgcgtccgag ctctcgatgc tctcgggttc gaggcagctc 300
ggcgcaaga gggcgcccat ggaggccgag accgagccca agcttgagga cttctcgggt 360
tgcaactcgt tctctccga gcaagatcag gccggcggtt tctgttctc cggagtcctg 420
atgaccacca gcagcaacag caacacgggg agcaaaccca tggagctctc tatgatcaag 480
agctggctcc gtgacaacca gctaccocaa gcgcgcgacc cgcagtcgca gaccggggcg 540
cccgcgagc agcgcaacc ccacgaggag atgggcaccc acgcagcagc cttcgatccg 600
cgggggagaa acggcgcggt ggtggtggcg gggagctcgc agagcctggc gctctcgatg 660
agcacgggtt caggatcgca cttgccaatg gctgtggtca gcgcagtgcc cagtggggga 720
gtctcagaga gcaactctc ggagaacaag cgagccagcg gcgccatgga ttcccgggg 780
ggtgcggtag aagctgtcgc gagggaagtc atcgacacat tcgggcaaa gacgtcgata 840
tatcgagggt taacaagaca tagatggaca ggcggttatg aggtctatct ctgggacaat 900
agctgcagaa gagaagggca gactcgcaag ggccgacaag tttatcttgg tggctatgac 960
aaggaggaca aggcagcaag ggttatgat ttggcagcac tcaagtatg ggttacaaca 1020
acaactcaaa atttcccaat taatacctac gaaaagagg tggatgaaat gaaacatag 1080
actaggcagg aatatatcgc atacttaaga aggaatagca gcggatttct tctgtgtgag 1140
tcaaaatata gtgtgttaac taggcacccat cagcatggga gatggcaagc aagaatttgg 1200
cgggttgtag gaaacaaaaga tctctactta ggcaccttca gcaccgagga ggaggcgggc 1260
gaggcctacg acatcgcgcc gatcaagttc cgcggggtca acgcgctcac caacttcgac 1320
atgagtcgct acgacgtcaa gaccatcctc gagagcagca ccctgcccgt ggggtggcgcg 1380
gctaggcgcc tcaaggaggc cgtgatcat ccggaggccg gcgccacgat ctggcgagcc 1440
ggcatggagc gcgccatcat ctcccagctg accgacgtcg ggtggggcgc gtacgcgtcg 1500
taccaccagc gctggccgag catcgattc cagcagccgt cgccgctctc ggtgcactac 1560
ccatacgtcg ccagccgccc ccgcggtggg tgcaagcccg agcaggacgc caccgtcgcc 1620
acgcaaaagc tgcaggacct ccagcagctg cacctcggga ccgcccgtca caactatttc 1680
caggcatcgt cgagctcaac ggtctacaac ggccggcgcg tcggcgggta ccagcagggc 1740
accggtggca acgctctctt gatgccggct agcacggtt gggatgaaca ggggcacagc 1800
agcactgccg ccaaccaggg aagcacctgc agctacgggg acgaagaggg gaagctcaat 1860
cttgggtacg atgccatggc gatggcgagc accggcgccg accatacgc ggccggcgta 1920
actgtgagca tcgcgagggc gaacgggtac tccaacaact ggagctcgcc gttcagtggc 1980
atgtga 1986

```

```

SEQ ID NO: 37      moltype = DNA length = 1986
FEATURE            Location/Qualifiers
source              1..1986
                    mol_type = other DNA
                    organism = Triticum aestivum

```

```

SEQUENCE: 37
atggcctcag gaaatgattg gcttgggttt tctctgtcag gccaggagaa cctcagcca 60
catcaggatt ctccaccttc agcaggaatt gatattctag gtgcttctga ttctatggc 120
gtgcctactc agctgcccc agatgcacag ttgggagttc ctggtcacca gccatttac 180
ggtattactg aggattttta tagaggcgct catgaaaccc acgattggaa tatgcgcgga 240

```


-continued

```

gatcttgatt ataacggcgg agcctcagag ctttctatgc tgggtgggttc ttcagccggt 300
ggtggcaagc gcgctgccaat ggaagcagag actgaacctt aattggagga tttctctggc 360
tgcaactcat tcggtgtctga acaggatcag gccggagggt ttctgttctc aggcgttcca 420
atgactacct ctgcaaaactc aaacacagga tctaactata tgggaattgtc aatgatcaag 480
tcttgggtga gagataacca gttgcctcag gctcctcctc cacagtctca gaccggtgct 540
ccagcacagc agcctcagcc acacgaggaa atgggcacag atgcatcttc attcgatcct 600
agaggccgca atggcgccct tgtggttgca ggctcttcac agtcactggc cttgtctatg 660
tcaacaggct ctggatcaca tctgccaatg gccgtgggtt ctggctcagc atctggcgga 720
gtgtcagagt ctacttcttc agaaaacaaa cgcgcttcag gagccatgga ttctcctggt 780
ggcgctgtgg aggcagttgc taggaagtca attgatacct ttggacagcg cacatctatc 840
tataggggtg tgactaggca tagatggacc ggcagatacg aggccactt gtgggataat 900
tcatgtagac gcgaaggcca gtctcgcaaa ggaaggcagg tttatcttgg aggttacgat 960
aaggaagata aagcagctcg cgcttatgat ttggccgcac ttaagtactg gggcacaact 1020
accacaacta acttcccaat taacacttac gagaagggaag ttgatgagat gaaacatatg 1080
acccgccagg aatatactgc ttacctgagg agaaattctt caggatttcc aaggggtgcc 1140
tctaaataca ggggagtgac cagacatcac cagcacggta gatggcagcg tagaattggc 1200
cgcgttgccg gaaacaagga tctttatctg ggcacttttt ctacagagga agagcgtgcc 1260
gaggcatagc atattgcagc tatcaagttt aggggtctga atgctgtgac aaacttcgat 1320
atgtctagat acgatgttaa acaaatcctt gaatcttcaa ctctgcctgt gggaggagca 1380
gcaaggagggt tgaaggagcg tgcgatcat ccagaagccg gcgcaactat ctggagggct 1440
ggatgggatg gcgccattat ctcacagctt accgatgttg gtatgggcgc ttatgcctct 1500
taccatcacg catggcctac tatcgctttt cagcagcctt caccactgtc tgtgcactat 1560
ccttacgttg cacagcctcc aagaggatgg tgcaaacccag agcaggatgc aactgtggct 1620
acccagcttc tgcaggattt gcagcagttg catcttggtc cagcagctca caattacttt 1680
caggcctctt catcttcaac gtgtataaac ggtggcggag ttggtggcta ccagcagggc 1740
accggtggta atgcattctc gatgccagct tctacagtgg ttgatgaaca gggccactct 1800
tcaaccgcgc caaatcaggg atcaacatgt tcttatggcg atgaagaggg caagctgaac 1860
ttgggttacg atgcaatggc tatggcctct acaggcgctg atccttatgc tgccgcacat 1920
actgtttcta ttgcaagagc taatggctac tcaataaact ggtcttcacc attctctggt 1980
atgtga

```

```

SEQ ID NO: 38      moltype = DNA length = 1980
FEATURE            Location/Qualifiers
source              1..1980
                    mol_type = other DNA
                    organism = Triticum aestivum

```

```

SEQUENCE: 38
atggcttctg ggaacgactg gttgggggttc tcgctctccg gccaaagaaa cccacagccg 60
catcaggata gctcgccgcc cgcgcacatc gacatcgctg gcgccagatgg cttctatggc 120
ctaccaccgc agccagctcc cgaagctcaa ctccgtgtgc caggctcatca tcagccttca 180
tacgggatca cggaggcctt caatagggga gctcatgaaa tccacgattg gaacatgagg 240
ggtgatctcg actacaacgg cgggtcgctc gagctctcga tgctcgtcgg gtcgagcgca 300
gttggcgcca agagggcgcc catggaaggc gagaccgagc ccaagcttga ggacttctc 360
gttggcaact cgttcgtctc cgagcaagat caggccgggt gcttcttgtt ctccggtgtc 420
ccgatggcga ccagcaccac cagcaacagc aggaagcaaca ccattggagct ctctatgatc 480
aagagctggc tccgtgacaa ccaggtaacc caagcgcgcg acccgagtc gcagaccggg 540
gcgcccgcgc agcagccgca accccacgag gagatgggca ccgacgcgag cagcttcgat 600
ccgctgggga gaaacggcgc gttggtggtg gcggggagct cgcagagcct gccgctctcg 660
atgagcacgg gctcaggatc gcacttgccg atggctgtgc tcggcgccag tcccagtggt 720
ggagtctcag agagcacctc ctccggagaac aagcgggcca gcggcgccat ggattcgccg 780
ggcggtgcgg tagaagctgt cgcgaggaag tccatcgaca cattcgggca aaggacgtcg 840
atatatcgag gtgtaacaag acatagatgg acagggcgtt atgaggctca tctctgggac 900
aatagctgca gaagagaagg gcagagtcgc aagggtaggc aagtttatct ggggtggctat 960
gacaaggagg acaaggcagc aagggcttat gatitggcag cactcaagta ttgggtaca 1020
acaacaacaa caaatttccc aatttaatac tacgaaaaag aggtggatga aatgaaacat 1080
atgactagcc aggaatacat cgcataccta agaaggaata gcagcggatt ttcccggtgt 1140
gcgtcaaaat atcgtggtgt aactaggcac catcagcatg ggagatggca agcaagaatt 1200
gggaggggtg caggaaacaa ggaatctctc ttaggcacct tcagcaccga ggaggaggcg 1260
gcggaggcct acgacatcgc ggcgatcaag ttccgtgggc tgaacgccgt caccacactc 1320
gacatgagtc gctacgagct caagaccatc ctccagagca gcacctgccc ggtgggtggc 1380
gcggcgaggg cccccaagga ggtcgctgat catccagagg ctggcgccac gatctggcgg 1440
gcgggcatgg acggcgccgt catctcccag ctgaccgacg tcgggatggg cactgacggc 1500
tcgtaccacc acggctggcc aaccatcgca ttccagcagc cctcgccgct gtcggtgcac 1560
taccatacag tcgcccagcc gtctcgccgg ttggtgcaag ccgagcagga cgcacgcgtc 1620
gccacgcaaa gccctgcagg cctccagcag ctgcacctcg ggacgcgcgc tcacaactat 1680
ttccaggcat cgtcgagctc caccgtctac aacggcgccg ggtaccagca aggcaccagt 1740
ggcaacgcct tcttgatgcc ggctagcacc gtggtgggat aacaggggca cagcagcact 1800
gccaccaacc agggagacac atgcagctac ggggacgaag aggggaagct caatctcggt 1860
tacgatgcca tggcgatggc gagcaccggc gccgacccat acggcgccgc gccaaactgtg 1920
agcattgcga gggcgaaagg gtactccaac aactggagct cgccgttcaa tggcatgtga 1980

```

```

SEQ ID NO: 39      moltype = DNA length = 1980
FEATURE            Location/Qualifiers
source              1..1980
                    mol_type = other DNA
                    organism = Triticum aestivum

```

-continued

SEQUENCE: 39

```

atggcttctg gaaatgattg gcttggtttt tctctgtcag gccaggagaa cccacagcct 60
catcaggatt cttcaccacc tgcgatatt gatatcgag gagcttcagg tttctatggc 120
cttccaactc agccagctcc tgatgcccag ctgggagtg caggtcatca ccagccttct 180
tacggatta ccgaggcttt taatcgcggc gctcatgaaa tccacgattg gaatatgagg 240
ggcgatctgg attataacgg cggagcctca gagctttcta tgctgggtggg ctcttcagca 300
gttggtggca agcgcgctgc catggaagct gagactgaac caaaattgga ggattttctt 360
gtgggaaact cattcgtttc tgaacaggat caggctggag gttttctgtt ctccaggtgtg 420
cctatggcca catctactaa ttctaactca aggtctaata ccatggaact gtcaatgatt 480
aagctttggt tgagagataa ccaggttcca caggccccc atcctcagtc acagaccgga 540
gcacctgcac agcagccaca gcctcacgag gaaatgggaa cagatgcata ttcatttgat 600
ccattgggca gaaatggcgc tcttggtgtt gccggtctct cacagtctct gccattgtca 660
atgtctacag gctcaggatc tcacctgcca atggctgtgt tgggcggatc acctctggtt 720
ggcgtttcag agtctacttc ttcagaaaac aaacgcgat caggcgctat ggattctcct 780
ggaggtgctg tggaggctgt tgcagggaag tcaattgata cattcggaca gaggacttct 840
atctatagag gtgtgaccag acatcgctgg acaggcagat acgagggcca cttgtgggat 900
aattcatgca gacgcgaagg acagtctagg aaaggtagac aggtttatct tggcgataac 960
gataaggaa gataaagcag tcgcgcttat gatttggccg cacttaagta ctggggcact 1020
accacaacta ccaacttccc aattaacact tacgagaagg aagtggatga gatgaaacat 1080
atgacccgcc aggaatatat cgctactctg aggagaaatt cttcaggatt ctcaagggtt 1140
gcatctaat acagaggagt gacacgcat caccagcac gtagatggca ggctcgcat 1200
ggcagggttg ccggaaccaa ggaactttat ctgggcactt ttcaaccga ggaagaggct 1260
gccagggttg agcatattgc agctatcaag tttagaggac tgaatgcgtt gactaacttc 1320
gatattgtct gctacgatgt taaaactatc cttgaatctt caacctgcc tgtgggtgga 1380
gcagcacgca ggcaaaagga ggtgcagat catcctgaag caggcgctac catttgagg 1440
gctggcatgg atggaggtgt gatctcacag ctgacagatg ttggtatggg cacttatgcc 1500
tcttaccatc acggtgggcc aacaattgca ttccagcagc catcaccttt gtctgtgcac 1560
tatccatacg ttgccagccc ttctagaggt tggtgcaaac ctgagcagga tgctcagtg 1620
gcaacacagt cctcaggaga ttgacagcag ttgcatcttg gaactgctgc ccacaattac 1680
tttcaggctt cttcatcttc aactgtttat aatggcggag gttaccagca gggcacttct 1740
ggcaacgcct tcctgatgcc ctgactaca gtggttgatg aacagggcca ttcttaacc 1800
gcaacaaatc agggatcaac ctgttcttat ggcatgaag agggcaagct gaacttgggt 1860
tacgatgcca tggcaatggc ttctaccggc gccgatccat atgcagctgc ccctacagt 1920
tcaatcgcca gagcaaatgg ctactctaact aactggctctt cacttttcaa cggtatgtga 1980

```

SEQ ID NO: 40

moltype = DNA length = 2112

FEATURE

Location/Qualifiers

source

1..2112

mol_type = other DNA

organism = Sorghum bicolor

SEQUENCE: 40

```

atggctactg tgaacaactg gctcgctttc tccctctccc cgcaggagct gccgcccacc 60
cagacggact ccaccctcat cctcgccgcc accacgcagc atgtctccgg cgatgtctgc 120
ttcaacatcc cccaagattg gagcatgagg ggaatccgagc tttcggcgct cgtcgccgag 180
ccgaagctgg aggaacttct cggcggaatc tccctctccg agcagcacca caaggccaac 240
tgcaacatga tcccacgac tacgacgaca gcttgctacg cgaactcggg tgctaccgcc 300
ggctaccatc accagctgta ccaccagccc accagctccg cgctccactc cgtgacttcc 360
gtcatggttg cctcctcggc cggcgcgctc cagcagcagg gtgcatctgt cagcgcgggc 420
agcgctaagt ttagcgctgg cgtggcgctt gccagtgcca atggcagcgg cagcatcggg 480
ctgtccatga tcaagaactg gctgcggagc caaccagctc ccatgcagcc gaggggtggc 540
gcggtgaga gctgacaggg gctctctttg tccatgaaca tggcgggggc gacgcaaggc 600
gocgctggca tgcacttct tgcctggagag cgcggccggg cgcggagag tgctcagac 660
tcggcacagg gtggagccgt cgtcacggct ccaaggagg atagcggtag cagcgggtgt 720
gccgccaccg gcgcctagt agcctgagc acggacacgg gtggcagcgg cgcgtcggct 780
gacaacacgg caaggaagac ggtggacacg ttcgggcagc gcagctcgat ttaccgtggc 840
gtgacaaaggc atagatggac tgggagatat gaagcacatc tgtgggacaa cagtgtcaga 900
agggaaaggac aaactcgcaa gggctgtcaa gtctatttag gtggctatga taaagaggag 960
aaagctgcta gggcttatga tctggctgct ctttaagtact ggggtcccac gacaacaaca 1020
aattttccag tgaataacta cgaaaaggag ctggaggata tgaagcacat gacaaggcag 1080
gagttttaga cgtctctgag aaggaagagc agtggtttct ccagaggtgc atccatttac 1140
aggggagtgat taggcatca ccagcatgga agatggcaag cacggatttg acgagttgca 1200
gggaacaagg atctctactt gggcaccttc agcacgcagg aggagggcagc ggaggcatac 1260
gacattgcgg cgatcaagtt ccgcggcctc aacgcgctca caaacttca catgagccgc 1320
tacgacgtca agagcatcct ggacagcagt gcgctcccca tcggcagcgc cgccaagcgt 1380
ctcaaggagg ccgagggcgc cgcgtcgca cagcaccatg ccggcggtgt gagctacgac 1440
gtcggccgca tagcctcaaca gctcggcgac ggccggcgccc tggcgggcgc gtacggcgcg 1500
cactaccatg gcgcctggcc gaccatcgcg ttccagccga gcgcggccac gggcctgtac 1560
caccgtagc gcagccgat gcgcgggttg tgcaagcagg agcaggacca cgcgggtgatc 1620
gcccggcgcg acagcctgca ggagctccac cacctgaacc tgggtgctgc cgcggcgcg 1680
cacgacttct tctcggcggg gcagcaggcg gcgatgcacg gcctgggtag catggacaat 1740
gaatcactcg agcacgcac cggtccaac tccgtcgtgt acaacgggtt tggatgatac 1800
aacggcagca ccgctcgtcg cagtgtgtgc tacatgatgc ctatgagcgc tgccacggcg 1860
acggctacca cggcaatggt gagccacgag caggtgcata cacgggcaca gggatgatc 1920
cacgacgaag ccaagcaggc tgcctcagatg gggtagcaga gctacctggt gaacgcagag 1980
aactatggcg gcgggaggat gctcgggcc tctcagcgcc accggcgcca 2040
agcagcaacg ataacatggc ggacgtcggc catggcgcg cagactctt cagtgtctgg 2100

```

-continued

aacgatactt aa 2112

SEQ ID NO: 41 moltype = DNA length = 2112
 FEATURE Location/Qualifiers
 source 1..2112
 mol_type = other DNA
 organism = Sorghum bicolor

SEQUENCE: 41

atggccacag	tgaataactg	gcttgcat	tctctgtcac	cacaggaa	tccacctaca	60
cagaatgatt	caactctgat	ttctgtctgc	actaccgatg	atgtgtcagg	cgatgtttgc	120
ttcaatatcc	cacaggattg	gtctatgcgc	ggctcagaac	tttctgcact	ggttgctgaa	180
cctaagtttg	aggattttct	tggcggaa	tctttctcag	agcagcatca	caaagccaat	240
tgcaacatga	tcccatcaac	atcttcaact	gcctgttatg	catcttcagg	tgcaacagct	300
ggctatcatc	accagttgta	ccatcagcct	acttcttcag	cccttcactt	tgcaattctt	360
gtgatgtgtg	cttcttcagc	cggctggcgt	catgatggag	gtgctatgct	ttcagcagct	420
tctgcaaatg	gctcagcagg	tgcaaggagca	gcactctgcaa	acggatctgg	ttcaattggt	480
ttgtcaatga	tcaagaattg	gcttcgctct	cagccagctc	ctatgcagcc	aagggtggct	540
gocgcagaa	ctgttcaggg	tctttctctg	tctatgaata	tggtcggcgc	cacacaggga	600
gctgcccgtg	tgccactgtt	ggcaggagaa	aggggaagg	caactgagtc	tgtgtcaacc	660
tctgtcagg	ggcaggcgtg	gggttacagca	cctaaggagg	attcagggtg	ctctggagtt	720
gcagctactg	gtgcattggt	ggctgtttct	accgatacag	gaggttcagg	agcatctgct	780
gataataccg	cccgcataac	cgtggataca	ttcgccagca	ggacatcaat	ctatagagga	840
gttacaagac	atcgctggac	tggaagatac	gaagctcacc	tttgggataa	ctcttgacga	900
cgccaggggc	agaccaggaa	gggaagacag	gtgtatctgc	gcggtacaga	taaggaggaa	960
aaagccgcaa	ggccttatga	ttggcgtgcc	cttaataact	ggggcccaac	aactaccaca	1020
aattttctcg	tgataaacta	tgagaaggaa	ctggaggata	tgaaacacat	gaccaggcag	1080
gaatttgttg	cttcattgag	gagaaagtct	tcaggattct	caaggagtgc	ctctatctat	1140
agaggcgtga	cacgccatca	ccagcatgga	aggtggcagg	ctaggatcgg	aagggttgca	1200
ggaaacaaag	atttgtatct	tgccactttc	tctaccagg	aggaagcagc	tgaggcatac	1260
gatattgccg	caatcaagtt	tagaggattg	aatgctgtga	ctaacttcga	tatgtcacgc	1320
tacgatgtta	aatctatcct	ggattcttca	gccttgccaa	tcggctcagc	tgccaagagg	1380
cttaagaag	ctgaagcagc	tgccctctgt	cagcatcacg	ccggcgtggt	ttcatacagat	1440
gtgggaagaa	ttgcctctca	gcttgccgat	ggtggcgcac	tgccagctgc	ctatggtgcc	1500
cattaccacg	ccgcatggcc	aactatcgct	tttcagcctt	cagcagctac	cggactgtat	1560
caccatacgc	ccagccttat	gagagggttg	tgtaagcagg	aacaggatca	tgctgttatc	1620
gocgcagctc	actctctgca	ggagttgcat	cactcgaatt	tggttgccgc	agctggcgca	1680
catgatattc	tttctgtctg	tcagcaggcc	gcaatgcacg	gcctgggagc	tatggataat	1740
gcctcatttg	agcattctac	tggtcctaat	tctgtggttt	ataacggtgt	gggctgattca	1800
aacggatcta	ccgtggtttg	ttcaggaggt	tacatgatgc	ctatgtctgc	tgccactgcc	1860
accgcaacta	ccgctatggt	gtccatgaa	cagggttcacg	ctagagctca	gggcgatcat	1920
cacgatgagg	ccaacacagg	agctcagatg	ggatatgaat	cttacctggt	gaatgctgag	1980
aactacggag	gtggttaggat	gtcagcagca	tgggcaactg	tttctgtctc	acctgctgcc	2040
tcttcaaatg	ataacatggc	agatgtgggc	catggcggag	ctcagttggt	ctctgttttg	2100
aatgatacct	ga					2112

SEQ ID NO: 42 moltype = DNA length = 927
 FEATURE Location/Qualifiers
 source 1..927
 mol_type = other DNA
 organism = Triticum aestivum

SEQUENCE: 42

atggacaagc	agagcgtcat	gtggagacag	ctgcagcacc	agcaccacca	ggccagcggc	60
ggggcggtg	gcaatgccgt	cgtgtctgcc	ggtgtgtgtg	cgacgacgac	gaccgcgccc	120
gtccggcgga	gcggcgcgcg	gtggacgccc	acggcggagc	aggtcaagat	cctcaaggac	180
ctctactacg	actgcggcat	ccggctcgcc	acggcggagc	agatccagcg	gatcgcgccc	240
cggtccggc	agtaacggcg	gatcgagggc	aagaacgtct	tctactggtt	ccagaaccac	300
aaggcccgcg	agcgccagaa	gaagcgccct	ggcgctcgac	tcaacggctc	ccccctcgcc	360
accgccaccg	ccgcccagct	cctcgccctc	tccccctccg	gtgcggcggc	cggcttgta	420
ggcgccggca	gctgcgcggg	tagaggcgcg	gctgtccatc	cagatgcgag	cgccactact	480
actacttgct	ggggagacag	caccctgcag	gactacatgg	gcgcgaggag	cacggcgggg	540
gcgggcaacc	acggcgcgcg	cgccgcagca	gcacctacgc	cgtggccagc	tgcgagcttc	600
cctttctcca	ccaaccagac	gcccgcgaat	cccccgccgc	gggagctccc	gctcttcccc	660
accggcgagg	gcccggcaaga	aagcgccgac	gacttcaacg	gcagcagcta	ccatctccag	720
cccaacagct	cgcagtggtg	ggaagctgcc	gcccgcagca	acgccaacgc	catggcagtc	780
gtccatcatc	agctgctgca	agaacagcac	gagcagcagc	acagctttta	caacagcagc	840
ggcaaccagc	agcagatgat	gatgatgatg	cccatccagg	acggcggcac	ctccccggag	900
ctcactctcc	gcgcccctta	catgtga				927

SEQ ID NO: 43 moltype = DNA length = 927
 FEATURE Location/Qualifiers
 source 1..927
 mol_type = other DNA
 organism = Triticum aestivum

SEQUENCE: 43

atggataagc	agtcagttat	gtggaggcag	cttcagcatc	agcaccacca	ggcatctgga	60
gctgcccggc	gaaatgcagt	ggcagctgcc	ggagccggtg	caactaccac	aactgcacca	120

-continued

```

gttcgccctt caggagctag gtggacacct actccagagc aggttaagat tcttaaagat 180
ctgtactacg attgoggtat cagatctcca actgctgaac agattcagcg catcgagct 240
agactgcgco agtatggcgg cattgaggga aagaatgtgt tttactggtt ccagaaccac 300
aaagcaaggg aaagacagaa gaaaagggtg ggagtggatg ttaacggttc accacttgct 360
accgccacag ccgagatgt tcttgccctg gtgctgcgcg aggcctgtat 420
ggcgctggat cttgtgctgg tagaggcgct gctgtgcac ctagatgcac tgctaccaca 480
actacctgtt ggggcgattc aaccttgacg gattacatgg gagctcgctc tacagccgga 540
gcaggtaatc atggtggcgc agctgcgcga gctcctactc catggcctgc cgcactcttt 600
ccattctcaa ctaaccagac cccacctatg ccaccacca gagagttgcc actttttcct 660
actggcgggc gcagacagga atcagccgat gatttcaatg gctcttcata tcatttgacg 720
cctaactctt cacagtgggt ggaggctgcc gcagcttcaa atgctaagcg catggcagtg 780
gttcatcacc agctgttgca ggagcagcat gaacagcagc actcttttta taattcttca 840
ggtaaccagc agcagatgat gatgatgat ccaatccagg atgctggcac atctccagaa 900
ctgactttga gagccctta catgtga 927

```

```

SEQ ID NO: 44      moltype = DNA length = 984
FEATURE           Location/Qualifiers
source            1..984
                  mol_type = other DNA
                  organism = Sorghum bicolor

```

```

SEQUENCE: 44
atggcggcca atgtgggcgc gggcaggagt gccgtcggcg gcggcggtgg tggtagagcg 60
ggtaggactg gcaactgtgc tgccagcgcg agcgtggcga cgacagcggt gtgccgcccc 120
atcggtctcg ggtggacgac gacgcccagc cagatcagga tactcaagga gttctactac 180
ggctgcggca tccggtcgcc caactcggag cagatccagc gcataccgc catgctgcgc 240
cagcacggca agatcgaggg caagaacgtc tttactggt tccagaacca caaggcccg 300
gaacgccaga agcgcgcct caccacacct gacgtcaacg tgcccaccgc cgcagccg 360
ggcgcgggcg acgcccagcc ccaactaggc gtcctctcgc tgcgtcgcc ttcaggcgcg 420
gcgctccct cgcgccacct cggtcttctac gcgggcaatg gcggcgcgcg ctgcagcggt 480
ctgctggaca caggttccga ctgcgtgcc atggccactg agacatgctt cctgcaggac 540
tacatgggag tgatgggcac gggcagcgcg gccgcgcgt cgcgtgggc atgcttctcg 600
tcgtcgaaac catgggcgcg ggcgcggcga cgggcaccga cggtagcgcg ggcgcccag 660
acgctccctc tcttcccgac gggcgcgac gacagccagc cccgcgcgcc gcggcacgga 720
gtcccagttc cagttgcagc aggcgagggc atccgcggcg gcagcagcag cagcaggtac 780
ctgcggttct ggggtgcgcg gccacaact gccagtgcga cttccattgg gatccagcag 840
caacaccagc tgctgcagct gcaagagcag tacagctttt acagcaacgc catgcccggc 900
accgcgagcc aagatgcac agcagcatcc ctggagctca gcctcagctc ctggtgctcc 960
ccttaccctg cagggaacct gtga 984

```

```

SEQ ID NO: 45      moltype = DNA length = 984
FEATURE           Location/Qualifiers
source            1..984
                  mol_type = other DNA
                  organism = Sorghum bicolor

```

```

SEQUENCE: 45
atggcggcga acgtgggggc aggtcgggtc gccgtcgggg gagggggagg ggttagagcg 60
ggtaggactg gcacagcgcg agcgtcagcg tcagttgcca ctaccgcagt gtgcagacca 120
attggtcttc gctggacccc aacacctgag cagattagga tctgaagga attttattac 180
ggctgtggaa ttatgacacc taattctgag cagattcaga ggatcactgc tatgttgaga 240
cagcatggca agatcgaggg aaaaaatgtg ttttattggt tccagaacca caaggcccg 300
gaaagggcga aaagacgcct tacaatctg gatgtgaatg ttccaacagc tgccgcagct 360
ggcgccgca atgcatcaac tcatttggga gtgtgtgcac tttcttcacc atctggtgct 420
gccccacct cacctacct tggtttttat gctggcaatg gcggagccgg atctaccgtt 480
ctgttgata catcttcaga ttgcgcagct atggcaactg aaacctgctt tctgcaggat 540
tacatgggag tgatgggcac tggatcagcc gcagctgcct ctcttgggc ttgcttctct 600
tcattcaaca caatggcagc tgccgcagct cgcgcacca cagttactag ggtcctgag 660
actctgccat tgttccctac cgttgcgcat gattctcagc caagaagacc aagacatgga 720
gtgccagttc ctgtggctgc tggcgaggct atcagaggcg gctcatcttc atctcgctac 780
cttccatttt ggggtgctgc cctacaact gcatcagcta catctattgg catccagcag 840
cagcaccagc ttctgcagct gcaggagcag tattcattct actctaagc catgcccagg 900
actggttcac aggatgcac tcgcgcactc cttgaattgt ccttgcctcc ctggtgctct 960
ccctatcccg ctggaactat gtga 984

```

```

SEQ ID NO: 46      moltype = DNA length = 783
FEATURE           Location/Qualifiers
source            1..783
                  mol_type = other DNA
                  organism = Arabidopsis thaliana

```

```

SEQUENCE: 46
atggaaaacg aagtaaacgc aggaacagca agcagttcaa gatggaaccc aacgaaagat 60
cagatcacgc tactggaaaa tctttacaag gaaggaatac gaactccgag gcgcgatcag 120
attcagcaga tcacggtag gcttcgtgcy tacggccata tcgaaggtaa aaacgtcttt 180
tactggttcc agaaccataa ggctaggcaa cgccaaaagc agaaacagga gcgcattggct 240
tacttcaatc gctcctccca caaaacctcc cgtttcttct accccctccc ttgctcaaac 300
gtgggttggt tcagtcgcta ctatttacg caagcaagtg atcatcatat gaatcaacat 360
ggaagtgtat acacaaacga tcttcttcac agaaacaatg tgatgattcc aagtgtggc 420

```

-continued

```

tacgagaaac ggacagtcac acaacatcag aaacaacttt cagacataag aacaacagca 480
gccacaagaa tgccaatttc tccgagttca ctcagatttg acagatttgc cctccgtgat 540
aactgttatg ccggtgagga cattaacgtc aattccagtg gacggaaaac actccctctt 600
tttctcttcc agcctttgaa tgcaagtaat gctgatggta tgggaagttc cagttttgce 660
cttggtatgt attctccggt ggattgttct agcgatggag ccggccgaga gcagccggtt 720
attgatttct tttctggtgg ttctacttct actcgtttcc atagtaatgg taatgggttg 780
taa 783

```

```

SEQ ID NO: 47      moltype = DNA length = 783
FEATURE            Location/Qualifiers
source              1..783
                    mol_type = other DNA
                    organism = Arabidopsis thaliana

```

```

SEQUENCE: 47
atggaaaacg aagtaaacgc aggaacagca agcagttcaa gatggaaccc aacgaaagat 60
cagatcacgc tactgaaaaa tctttacaag gaaggaatac gaactccgag cgccgatcag 120
attcagcaga taccoggtag gcttcgtgcy tacggccata tcgaaggtaa aaacgtcttt 180
tactggttcc agaaccataa ggctaggcaa cgccaaaagc agaaacagga gcgcattggt 240
tacttcaatc gctctctcca caaacctccc cgtttcttct acccccctcc ttgctcaaac 300
gtgggttggt tccagtcgta ctatttacag caagcaagtg atcatcatat gaatcaacat 360
ggaagtgtat acacaaaacg tcttcttcac agaaacaatg tgatgatgcc aagtgggtggc 420
tacgagaaac ggacagtcac acaacatcag aaacaacttt cagacataag aacaacagca 480
gccacaagaa tgccaatttc tccgagttca ctcagatttg acagatttgc cctccgtgat 540
aactgttatg ccggtgagga cattaacgtc aattccagtg gacggaaaac actccctctt 600
tttctcttcc agcctttgaa tgcaagtaat gctgatggta tgggaagttc cagttttgce 660
cttggtatgt attctccggt ggattgttct agcgatggag ccggccgaga gcagccggtt 720
attgatttct tttctggtgg ttctacttct actcgtttcc atagtaatgg taatgggttg 780
taa 783

```

```

SEQ ID NO: 48      moltype = DNA length = 978
FEATURE            Location/Qualifiers
source              1..978
                    mol_type = other DNA
                    organism = Arabidopsis thaliana

```

```

SEQUENCE: 48
atgtcctcct caaacaacaa ttggccaagc atgttcaaat ccaaaccctg caacaataat 60
catcatcatc aacatgaaat cgatactcca tcttacatgc actactctaa ttgcaacctc 120
tcattcttct tttctcaga tcggatacca gatcctaacc cgagatggaa tcctaaaccg 180
gagcagatta ggatactcga atcaatcttc aattccggtg ctattaaacc acctagagag 240
gagattcaaa gaatccggat ccggcttcaa gaatatggtc aaatccggtg cgcaaacgtg 300
ttttactggt ttcaaaaaccg gaaatctcga gcaaaaacac agcttcgtgt tcatcacaaa 360
agccctaaaa tgtcaaagaa ggacaagacg gttattccta gtactgacgc tgatcattgt 420
tttggttttg ttaaccaaga aaccggatta tatccgggtc aaaaacaatg gttggtggta 480
accgaaccgg ccggttttct atttccggtt cataatgacg cgagcgctgc tcaatcagcg 540
tttggttttg gcgattttgt tgtaccggtg gtaacggaag aagggatggc attctctacc 600
gttaataacg gcgttaattt ggagactaac gaaaattttg ataaaattcc ggcgatcaat 660
ttatacggcg gagatggaaa tggcgggtga aattgttttc ctctttgac tgttccatta 720
accatcaatc aatctcaaga aaaacgagat gtaggattat ccggtggtga agacgtcgga 780
gataatgttt atccggtgag aatgacgggt tttattaacg agatgcctat cgaagtagtg 840
tctggattat tcaacgttaa ggcagctttc ggaaacgatg ccgttttgat caactcgttt 900
ggccagccta ttcttacaga tgaatttggt gttacttata aacctctcca aaatggcgca 960
atctattatc ttatttag 978

```

```

SEQ ID NO: 49      moltype = DNA length = 978
FEATURE            Location/Qualifiers
source              1..978
                    mol_type = other DNA
                    organism = Arabidopsis thaliana

```

```

SEQUENCE: 49
atgtcctcct caaacaacaa ttggccaagc atgttcaaat ccaaaccctg caacaataat 60
catcatcatc aacatgaaat cgatactcca tcttacatgc actactctaa ttgcaacctc 120
tcattcttct tttctcaga tcggatacca gatcctaacc cgagatggaa tcctaaaccg 180
gagcagatta ggatactcga atcaatcttc aattccggtg ctattaaacc acctagagag 240
gagattcaaa gaatccggat ccggcttcaa gaatatggtc aaatccggtg cgcaaacgtg 300
ttttactggt ttcaaaaaccg gaaatctcga gcaaaaacac agcttcgtgt tcatcacaaa 360
agccctaaaa tgtcaaagaa ggacaagacg gttattccta gtactgacgc tgatcattgt 420
tttggttttg ttaaccaaga aaccggatta tatccgggtc aaaaacaatg gttggtggta 480
accgaaccgg ccggttttct atttccggtt cataatgacg cgagcgctgc tcaatcagcg 540
tttggttttg gcgattttgt tgtaccggtg gtaacggaag aagggatggc attctctacc 600
gttaataacg gcgttaattt ggagactaac gaaaattttg ataaaattcc ggcgatcaat 660
ttatacggcg gagatggaaa tggcgggtga aattgttttc ctctttgac tgttccatta 720
accatcaatc aatctcaaga aaaacgagat gtaggattat ccggtggtga agacgtcgga 780
gataatgttt atccggtgag aatgacgggt tttattaacg agatgcctat cgaagtagtg 840
tctggattat tcaacgttaa ggcagctttc ggaaacgatg ccgttttgat caactcgttt 900
ggccagccta ttcttacaga tgaatttggt gttacttata aacctctcca aaatggcgca 960
atctattatc ttatttag 978

```

-continued

SEQ ID NO: 50 moltype = AA length = 240
 FEATURE Location/Qualifiers
 source 1..240
 mol_type = protein
 organism = *Agrobacterium tumefaciens*

SEQUENCE: 50

MDLRLLIFGPT	CTGKTSTAVA	LAQQTGLPVL	SLDRVQCCPQ	LSTGSGRPTV	EELKGTSLRY	60
LDDRPLVKGI	IAAKQAHERL	MGEVYNYEAH	GGLILEGGS	SLLKCMQSS	YWSADFRWHI	120
IRHELADEET	PMNVAKARVK	QMLRPAAGLS	IIQELVDLWK	EPRLRPILKE	IDGYRYAMLF	180
ASQNQITSDM	LLQLDADMED	KLIHGIAQEY	LIHARRQEQK	FPRVNAAYD	GFEGHPFGMY	240

SEQ ID NO: 51 moltype = AA length = 318
 FEATURE Location/Qualifiers
 source 1..318
 mol_type = protein
 organism = *Arabidopsis thaliana*

SEQUENCE: 51

MKCNCKMVVI	MGATGSGKSS	LSVDLALHFK	AEIINSDKMQ	FYDGLKITTN	QSTIEDRRGV	60
PHLLLGELNP	EAGEVTADEF	RVMAAEAEISE	ITQRKKLPIL	AGGSNSYIHA	LLAKSYDPEN	120
YFSDHKGSI	CSELKYDCCF	IWIDVDQSVL	FEYLSLRDL	MMKSGMFEEI	AEFHRSKKAP	180
KEPLGIWKAI	GVQEFDDYLK	MYKWDNDMDK	WDPMRKEAYE	KAVRAIKENT	FQLTKDQITK	240
INKLRNAGWD	IKKVDATASF	REAIRAAKEG	EGVAEMQRKI	WNKEVLEPCV	KIVRSHLDQP	300
INYYYYFYFL	LKRFLSLN					318

SEQ ID NO: 52 moltype = AA length = 359
 FEATURE Location/Qualifiers
 source 1..359
 mol_type = protein
 organism = *Zea mays*

SEQUENCE: 52

MEEITQHFGV	GASSHGHGHG	QHHHHHHHHH	PWASSLSAVV	APLPPQPPSA	GLPLTLNTVA	60
ATGNSGGSGN	PVLQLANGGG	LLDACVKAKE	PSSSPYAGD	VEAIKAKIIS	HPHYSSLTA	120
YLECNKVGAP	PEVSARLTEI	AQEVEARQRT	ALGGLAAATE	PELDQFMEAY	HEMLVKFREE	180
LTRPLQEAME	PMRRVESQLN	SLSISGRSLR	NILSSGSSEE	DQEGSGGETE	LPEVDAHGVD	240
QELKHHLLKK	YSGYLSLQK	ELSKKKKKGK	LPKEARQQLL	SWWDQHYKWP	YPSETQKVAL	300
AESTGLDLKQ	INNWFINQRK	RHWKPSSEMH	HLMDGYHTT	NAFYMDGHFI	NDGGLYRLG	359

SEQ ID NO: 53 moltype = AA length = 584
 FEATURE Location/Qualifiers
 source 1..584
 mol_type = protein
 organism = *Arabidopsis thaliana*

SEQUENCE: 53

MNSMNNWLGF	SLSPHDQNH	RTDVSSTTR	TAVDVAGGYC	FDLAAPSDDES	SAVQTSFLSP	60
FGVTLEAFTR	DNNSHSRDWD	INGGACNNIN	NNEQNGPKLE	NFLGRRTTIY	NTNETVVDGN	120
GDCGGDGGG	GGSLGLSMIK	TWLSNHSVAN	ANHQDNGNGA	RGLSLSMNSS	TSDSNNNYNN	180
DDVVQEKTI	DVVTTPPKT	IESFGQRTSI	YRGVTRHRWT	GRYEAHLWDN	SKREGQTRK	240
GRQVYLGVD	KEEKAARAYD	LAALKYWGTT	TTTNFPLSEY	EKEVEEMKHM	TRQEVASLR	300
RKSSGFSRGA	SIYRGVTRHH	QHGRWQARIG	RVAGNKDLYL	GTFGTQEEAA	EAYDIAAIKF	360
RGLSAVTNFD	MNRYNVKAIL	ESPSLPIGSS	AKRLKDVNNP	VPAMMISNNV	SESANNVSGW	420
QNTAFQHHQG	MDLSLLQQQQ	ERYVGYNGG	NLSTESTRVC	FKQEEEQQHF	LRNSPSHMTN	480
VDHSSSTSD	SVTVCGNVVS	YGGYQGFAIP	VGTSVNYDPF	TAAEIAYNAR	NHYYYAQHQQ	540
QQQIQQSPGG	DPPVAISNNH	SSNMYFHGEG	GGEGAPTFSV	WNDT		584

SEQ ID NO: 54 moltype = AA length = 661
 FEATURE Location/Qualifiers
 source 1..661
 mol_type = protein
 organism = *Triticum aestivum*

SEQUENCE: 54

MASGNDWLGF	SLSGQENPQP	HQDSPPPAGI	DISGASDFYG	VPTQPAPDAQ	LGVPGHQPSY	60
GITEDFNRGA	HETHDWNMRG	DLDYNGGASE	LSMLVGSSAV	GGKRAAMEAE	TEPKLEDFLG	120
CNSFVSEQDQ	AGGFLFSGVP	MTTSSNSNTG	SNTMELSMIK	SWLRDNQLPQ	APHPQSQTGA	180
PAQQQPQHEE	MGTDASSFDP	RGRNGALVVA	GSSQSLALSM	STGSGSHLPM	AVVSGSASGG	240
VSESTSSSENK	RASGAMDSPG	GAVEAVARKS	IDTFGQRTSI	YRGVTRHRWT	GRYEAHLWDN	300
SCRREGQSRK	GRQVYLGVD	KEDKAARAYD	LAALKYWGTT	TTTNFPINTY	EKEVDEMCKH	360
TRQYIAYLR	RNSSGFSRGA	SKYRGVTRHH	QHGRWQARIG	RVAGNKDLYL	GTFSTEEEA	420
EAYDIAAIKF	RGLNAVTFND	MSRYDVKTIL	ESSTLPVGG	ARRLKEAADH	PEAGATIWRA	480
GMDGAIISQL	TDVGMGAYAS	YHHGWPTIAF	QQSPSLSVHY	PYVAQPPRGW	CKPEQDATVA	540
TQSLQDLQQL	HLGTAAHNYF	QASSSSTVYN	GGGVGGYQQG	TGGNAFLMPA	STVDEQGH	600
STAAQGSTC	SYGDEEGKLN	LGADAMAMAS	TGADPYAAS	TVSIARANGY	SNNWSSPFG	660
M						661

SEQ ID NO: 55 moltype = AA length = 659
 FEATURE Location/Qualifiers

-continued

source 1..659
mol_type = protein
organism = *Triticum aestivum*

SEQUENCE: 55

MASGNDWLGF	SLSGQENPQP	HQDSSPPADI	DIAGASGFYG	LPTQPAPDAQ	LGVPGHHQPS	60
YGITEAFNRG	AHEIHDWNMR	GDLDYNGGAS	ELSMVLVGSSA	VGGKRAAMEA	ETEPKLEDFL	120
VGNSTFVSED	QAGGFLFSGV	PMATSTNSNS	RSNTMELSMI	KSWLRDNQVP	QAPHPQSQTG	180
APAQQPQPHE	EMGTDASSFD	PLGRNGALVV	AGSSQSLSLS	MSTGSGSHLP	MAVLGGSPSG	240
GVSESTSSEN	KRASGAMDSP	GGAVEAVARK	SIDTFGQRTS	IYRGVTRHRW	TGRYEHLWD	300
NSCRREGQSR	KGRQVYLGGY	DKEDKAARAY	DLAALKYWG	TTTTNFPINT	YEKEVDEMKH	360
MTRQEYIAYL	RRNSSGFSRG	ASKYRGVTRH	HQHGRWQARI	GRVAGNKDLY	LGTFSSTEEA	420
AEAYDIAAIK	FRGLNAVTFN	DMSRYDVKTI	LESSTLPVGG	AARRPKEVAD	HPEAGATIWR	480
AGMDGGVISQ	LTDVGMGTYA	SYHHGWPTIA	FQQPSPLSVH	YPYVAQPSRG	WCKPEQDASV	540
ATQSLQDLQQ	LHLGTAAHNY	FQASSSSTVY	NGGGYQQGTS	GNAFLMPAST	VVDEQGHSSST	600
ATNQGSTCSY	GDEEGKLNLG	YDAMAMASTG	ADPYAAAPT	SIARANGYSN	NWSSPFNGM	659

SEQ ID NO: 56 moltype = AA length = 703
FEATURE Location/Qualifiers
source 1..703
mol_type = protein
organism = *Sorghum bicolor*

SEQUENCE: 56

MATVNNWLAF	SLSPQELPPT	QTDSTLISAA	TTDDVSGDVC	FNIPQDWSMR	GSLSALVAE	60
PKLEDFLGGI	SFSEQHKKAN	CNMIPSTSST	ACYASSGATA	GYHHQLYHQP	TSSALHFADS	120
VMVASSAGGV	HDGGAMLSSA	SANGSAGAGA	ASANGSGSIG	LSMIKNWLRS	QPAPMQPRVA	180
AAESVQGLSL	SMNMAGATQG	AAGMPLLAG	RGRAPESVST	SAQGGAVVTA	PKEDSGSGSV	240
AATGALVAVS	TDTGGSGASA	DNTARKTVDT	FGQRTSIYRG	VTRHRWTGRY	EAHLWDNSCR	300
REGQTRKGRQ	VYLGGYDKKE	KAARAYDLAA	LKYWGPTTTT	NFPVNNYEKE	LEDMKHMTRO	360
EFVASLRRKS	SGFSRGASIV	RGVTRHHQHG	RWQARIGRVA	GNKDLYLGT	STQEEAAEAY	420
DIAAIKFRGL	NAVTFNFMDS	YDVKSILDSS	ALPIGSAKR	LKEAEEAASA	QHHAGVVSVD	480
VGRIASQLGD	GGALAAAYGA	HYHGAWPTIA	FQPSAATGLY	HPYAPQMRGW	CKEQDHAVID	540
AAAHSLQELH	HLNLGAAAGA	HDFFSAGQQA	AMHGLGSMND	ASLEHSTGSN	SVVYNGVGDS	600
NGSTVVGSGG	YMPMPSAATA	TATTAMVSHE	QVHARAQGDH	HDEAKQAAQM	GYESYLVNAE	660
NYGGGRMSAA	WATVSAPPAA	SSNDNMADVG	HGGAQLFSVW	NDT		703

SEQ ID NO: 57 moltype = AA length = 308
FEATURE Location/Qualifiers
source 1..308
mol_type = protein
organism = *Triticum aestivum*

SEQUENCE: 57

MDKQSVMMWRQ	LQHQQHQASG	AAGGNAVAAA	GAGATTTTAP	VRPSGARWTP	TPEQVKILKD	60
LYYDCGIRSP	TAEQIRIAA	RLRQYGRIEG	KNVFYWFQNH	KARERQKKRL	GVDVNGSPLA	120
TATAADVLLAL	SPSGAAAGLY	GAGSCAGRGA	AVHPDASATT	TTCWGDSTLQ	DYMGARSTAG	180
AGNHGGAAAA	APTPWPAASF	PFTNQTPPM	PPPRELPLFP	TGGGRQESAD	DFNGSSYHLQ	240
PNSSQWEEAA	AASNANAMAV	VHHQLLQEQH	EQQHSFYNS	GNQQQMMMMM	PIQDAGTSPE	300
LTLRAPYM						308

SEQ ID NO: 58 moltype = AA length = 327
FEATURE Location/Qualifiers
source 1..327
mol_type = protein
organism = *Sorghum bicolor*

SEQUENCE: 58

MAANVGAGRS	AVGGGGGGGA	GGTGTAASG	SVATTAVCRP	IGSRWTPTPE	QIRILKEFY	60
GCGIRSPNSE	QIQRTITAMLR	QHGKIEGKNV	FYWFQNHKAR	ERQKRRLTNL	DVNVPTAAAA	120
GAADASTHLG	VLSLSSPSGA	APPSPTLGFY	AGNGGAGSTV	LLDTSDDCAA	MATETCFLQD	180
YMGVMTGSGA	AAASPWACFS	SSNTMAAAAA	RAPTIVTRAPE	TLPLFPTGGD	DSQPRRPRHG	240
VPVPVAGEA	IRGSSSSSRY	LPFWGAAPT	ASATSIGIQQ	QHQLLQLQEQ	YSFYSNAMP	300
TGSQDASAAS	LELSLSSWCS	PYPAGTM				327

SEQ ID NO: 59 moltype = AA length = 260
FEATURE Location/Qualifiers
source 1..260
mol_type = protein
organism = *Arabidopsis thaliana*

SEQUENCE: 59

MENEVNAGTA	SSSRWNPTKD	QITLLENLYK	EGIRTPSADQ	IQQITGRLRA	YGHIEGKNVF	60
YWFQNHKARQ	RQKQKQERMA	YFNRLHKT	RFYPPPCSN	VGCVSPYYLQ	QASDHMMNQH	120
GSVYTNDLLH	RNNVIMPSGG	YEKRTVTQHQ	QQLSDIRTTA	ATRPIMPSS	LRFDRFALRD	180
NCYAGEDINV	NSSGRKTLPL	PPLQPLNASN	ADGMGSSSFA	LGSDSPVDCS	SDGAGREQPF	240
IDFFSGGSTS	TRFDSNGNGL					260

SEQ ID NO: 60 moltype = AA length = 325
FEATURE Location/Qualifiers
source 1..325

-continued

```

mol_type = protein
organism = Arabidopsis thaliana

SEQUENCE: 60
MSSSNKNWPS MFKSKPCNNN HHHQHEIDTP SYMHYSNCNL SSSFSSDRIP DPKPRWNPKP 60
EQIRILESIF NSGTINPPRE EIQRIRIRLQ EYQIGDANV FYWFQNRKSR AKHKLRVHHK 120
SPKMSKKDKT VIPSTDADHC FGFVNQETGL YPVQNNELVV TEPAGFLFPV HNDPSAAQSA 180
FGFGDFVVPV VTEEGMAFST VNNGVNLETN ENFDKIPAIN LYGGDNGGGG NCFPPLTVPL 240
TINQSQEKRD VGLSGGEDVG DNVYPVRMTV FINEMPIEVV SGLFNVKAAF GNDAVLINSF 300
GPILTDEFG VTYQLQNGA IYYLI 325

SEQ ID NO: 61      moltype = DNA length = 1592
FEATURE           Location/Qualifiers
source            1..1592
                  mol_type = genomic DNA
                  organism = Medicago truncatula

SEQUENCE: 61
actgttaata atttttaaac gtcagcgcac taaaaaaacg aaaagacgga cactgaaaa 60
taaaaaaacac acactagttt atgacgcaat actattttac ttagtatttg ggtacattag 120
acaaaaccggt gaaagagatg tatcagctat gaaacctgta tacttcaata cagagactta 180
ctcatatcgg atacgtacgc acgaagtatc atattaatta ttttaatttt taataaatat 240
ttatcgggat acttatgtga tactctacat atacacaagg atatttctaa gatactttat 300
agatacgtat cctagaaaaa catgaagagt aaaaaagtgga gacaatgttg taaaaattca 360
ttataaaatgt atatgattca atttttagata tgcacagta taattgattc tcgatgaaac 420
acttaaaatt atatttcttg tggagaacg tagcgagaga ggtgattcag ttagacaaca 480
ttaataaaaa ttaagtgtta gttcttttaa tgatgtttct ctcaatatca catcatatga 540
aaatgtaata tgattttata gaaaaatttt aaaaaattta ttttaataat cacatgtact 600
attttttaaa aattgtatct ttataataaa tacaataata aagagtaatc agtggttaatt 660
tttcttcaaa tataagtttt attataaatc attgttaacg tatcataagt cattaccgta 720
tcgtatctta tttttttttt aaaaacggct aattcacgta cccgtattgt attgtaccg 780
cacctgtatc acaatcgatc ttagttagaa gaattgtctc gaggcgggtc aagacagcat 840
ataatagacg tggactctct tataccaaac gttgtcgtat cacaagggt taggtaacaa 900
gtcacagttt tccacgtgtg cacgttttaa ttggaagagg tgcggtggc gtaataaac 960
agccaatcga ttttctgat aaaagcaaat caggtaaatc aaactcttct attcttttct 1020
tcccatcgcg tacaaaacgg gttccttttg aaaagagatt cattcaaacc tagcaccocaa 1080
ttccggtttca aggtataatc tactttctat tcttcgatta ttttattatt attagctact 1140
atcgtttaat cgatcttttc ttttgatccg tcaaatttaa attcaattag ggttttggtc 1200
ttttctttca tctgattgaa atccttctga attgaaccgt ttacttgatt ttactgttta 1260
ttgtatgatt taatcctttg tttttcaaag acagtcctta gattgtgatt aggggttcat 1320
ataaattttt agatttggat ttttgtattg tatgattcaa aaaatacgtc ctttaattag 1380
attagtacat ggatgttttt taaccgattt attgattgtc agggagaatt tgatgagcaa 1440
gtttttttga tgcctgttgt aaattgaatt gattataatt gctgatctgc tgcctccagt 1500
tttcataacc catattcttt taacctgtt gtacacacaa tgaaaaattg gtgattgatt 1560
catttgtttt tctttgtttt ggattataca gg 1592

SEQ ID NO: 62      moltype = DNA length = 2000
FEATURE           Location/Qualifiers
source            1..2000
                  mol_type = genomic DNA
                  organism = Zea mays

SEQUENCE: 62
gtcgtgcccc tctctagaga taaagagcat tgcattgtcta aagtataaaa aattaccaca 60
tatttttttg tcaacattat ttgaagtgtg gtttatctat ctctatacat atatttaaac 120
ttcactctac aaataatata gtctataata ctaaaataat attagtgttt tagaggatca 180
tataaataaa ctgctagaca tgggtctaaag gataattgaa tattttgaca atctacagt 240
ttatcttttt agtgtgcatg tgatctctct gttttttttg caaatagctt gacctatata 300
atacttcacg cacttttatta gtacatccat ttaggattta ggggttgatg tttctataga 360
ctaattttta gtacatccat tttattcttt ttagtctcta aattttttta aactaaaact 420
ctattttagt tttttattta ataatttaga tataaaatga aataaaataa attgactaca 480
aataaaacaa atacccttta agaaaataaaa aaactaagca aacatttttc ttgtttcgag 540
tagataaatga caggctgttc aacgccgtcg acgagtctaa cggacaccaa ccagcgaaac 600
agcagcgtcg cgtcgggccca agcgaagcag acggcacggc atctctgtag ctgcctctgg 660
acccctctcg agagtccgcg tccaccgttg gacttgcctc gctgtcggca tccagaaaatt 720
gctgtggcga gcggcagacg tagggcggca cggcaggcgg cctcttctct ctctcacggc 780
accggcagct acgggggatt cctttccccc cgtctctctg ctttcccttc ctgcgccgc 840
gtaataaata gacacccctc ccacacccct tttccccaac ctgctgttctg ttcggagcgc 900
acacacacgc aaccagatct cccccaatc cagccgtcgg cacctccgct tcaaggtaacg 960
cgcgtcatcc tccccccccc cctctctcta cctctcttag atcggcgatc cgggtccatgg 1020
ttagggcccg gtagtctcac tctgttcat gtttgtgta gagcaaacat gttcatgttc 1080
atgtttgtga tgaatgtgtc tgggtgggag gtcgttctag atcggagtag gatactgttt 1140
caagctacct ggtggattta ttaattttgt atctgtatgt gtgtgccata catcttcata 1200
gttacgagtt taagatgatg catggaaata tgcgtcagg ataggatata atgttgatgc 1260
gggttttact gatgcataa cagagatgct ttttttctcg cttgggtgtg atgataggt 1320
ctggttgggc ggtcgttcta gatcggagta gaatactgtt tcaaaactac tgggtgattt 1380
attaaaggat aaagggtcgt tctagatcgg agtagaatac tgtttcaaac tacctgggtg 1440
atttatataa ggatctgtat gtatgtgcct acatcttcat agttacgagt ttaagatgat 1500
ggatggaaat atcgatctag gataggtata catgttgatg cgggttttac tgatgcata 1560

```


-continued

acagagatgc	tttttttcgc	tgggtgtgga	tgatgtgggc	tggttggggc	gtcgttctag	1620
atcgagtag	aatactgttt	caaacacac	ggtggattta	ttaattttgt	atctttatgt	1680
gtgtgccata	catcttcata	gttacgagtt	taagatgatg	gatggaaata	ttgatctagg	1740
ataggtatag	atgttgatgt	gggttttact	gatgcatata	catgatggca	tatgcggcat	1800
ctattcatat	gctctaacct	tgagtacctt	tctattataa	taaacacagta	tgttttataa	1860
ttattttgat	cttgatatac	ttggatgatg	gcataatgac	cagctatatg	tggtattttt	1920
agccctgcct	tcatacgcta	tttatttgct	tggtactggt	tcttttgctc	gatgctcacc	1980
ctgtgtgttg	gtgatacttc					2000

SEQ ID NO: 63 moltype = DNA length = 4101
 FEATURE Location/Qualifiers
 misc_feature 1..4101
 note = Cas9
 source 1..4101
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 63

gacaagaagt	acagcatcgg	gctggcgatc	gggaccaact	ccgtcgctg	ggctgtgatt	60
accgacgagt	acaaggtgcc	atccaagaag	ttcaaggtcc	tcggcaacac	tgaccggcac	120
agcattaaaga	agaacctgat	tggggcgctg	ctgttcgatt	cgggggagac	tgccggaggcg	180
accaggctga	agcggactgc	gcgcccggagg	tacaccagga	ggaagaatcg	gatctgctac	240
ctccaggaga	ttttctcgaa	tgagatggcc	aaggtggacg	attccttctt	ccatcgccctg	300
gaggagtcgt	tcctcggtga	ggaggacaag	aagcatgaga	ggcatcccat	tttcgggaat	360
atcgttgacg	aggtggctta	ccatgagaag	tacccgacca	tctaccatct	gcggaagaag	420
ctcgtcgatt	cgaccgataa	ggccgacctg	cggctgatct	acctggccct	cgcgcacatg	480
attnaagtcc	ggggccattt	cctcatcgag	ggcgacctca	accgggacaa	ctcggacgtg	540
gataagctct	tcatttcagct	cgtgcagaca	tacaaccagc	tcttcgagga	gaatcccat	600
aacgcctcgg	gggtcgacgc	taaggctatt	ctctcggctc	ggctgtcgaa	gtcgcgcccg	660
ctggagaatc	tcatttgccc	gctcccaggc	gagaagaaga	acggcctctt	cggaacacct	720
attgcccctg	cgctggggct	cacaccgaat	ttcaagtcca	acttcgacct	cgcgaggac	780
gctaagctcc	agctcagcaa	ggatacttac	gatgatgacc	tcgataacct	gctcgccacg	840
attggggatc	agtacgcgga	tctgttcttc	gcgcccaaga	atctcagcga	tgctattctc	900
ctgtcggaca	ttctcccgct	caacacagag	attactaagg	ccccactgtc	ggcgagcatg	960
attnaagagt	acgatgagca	tcatacagg	ctgacactgc	tcaaggcgct	ggctcggcag	1020
cagctccccg	agaagtacaa	ggagattttc	ttcgatcagt	caaagaatgg	gtacgcgggc	1080
tacattgatg	gcgcccgcgt	ccaggaggag	ttctacaagt	tcatttaagc	catccctggag	1140
aagatggacg	ggaccgagga	gctgctggtg	aagctcaatc	gggaggacct	gctccggaag	1200
cagcgcacat	tcgacaatgg	ctcgatttct	caccagattc	acctggggcg	gctgcacgcc	1260
atttctccga	ggcaggagga	cttctaccgc	ttctccaagg	acaaccgcga	gaagatcgag	1320
aagatctcga	cttctccgat	tcataactac	gtggggccgc	tcgcgcgggg	gaactcccg	1380
ttcgcgtgga	tgactcgcga	gtccgaagaa	acgattacac	cgtggaattt	cgaggaggct	1440
gtcgacaagg	gcgctagtgc	gcagtcattc	attgagagga	tgaccaatct	cgataagaac	1500
ctgcctaacc	agcaggtgct	gccaagcat	ctcgtcgtct	acgagtactt	caccgtttac	1560
aatgagctga	ccaaggtgaa	gtatgtgact	gagggcatga	ggaagccagc	gttccctgagc	1620
ggcgagcaga	agaaggtctt	cgtggacctg	ctctccaaga	ctaaccggaa	gggtgactgtg	1680
aagcagctca	aggagactca	cttccaagaag	attgagtgtc	tcgattccgt	tgagattagc	1740
gggttgagg	atcgggtcaa	tgtctcgtc	gggacatacc	acgatctcct	gaagatcatt	1800
aaggataaag	acttctctga	caacgaggag	aacgaggaca	ttctcgaaga	tattgtcctg	1860
accctcaacc	ttctcagaga	tcgggagatg	atcgaggaga	ggctcaagac	atcgcctcat	1920
ctgttcgatg	ataaggtcat	gaagcagctg	aagcgcaggg	gggtacacagg	gtggggggcg	1980
ctgagccgga	agctgatcaa	cgggattcgg	gataaagcgt	ccgggaagac	aattctcgac	2040
ttctccaagt	ccgacggggt	gcctaaccgc	aacttcatgc	agctcaattc	tgatgactcg	2100
ctgacattca	aggaggtat	tcagaaggcg	cagggttctg	ggcagggcga	ctcgtctcac	2160
gagcatattg	cgaatctggc	gggctcccc	gcgattaaga	agggcattct	gcaaacctgc	2220
aaggtggttg	atgagctggt	caaggctcat	ggcgggcata	agccagagaa	tattgtcatc	2280
gagatggcgc	gggagaatca	gaccacacag	aaggggcaga	agaactcacg	ggagcggatg	2340
aagcgcacat	aggaggccat	caaggagctg	gggtcgcaga	tcctgaagga	gcacccctgc	2400
gagaacactc	agctgcaaaa	tgagaagctg	tacctctact	acctccagaa	cgggaggggac	2460
atgtatgtgg	atcaggagct	ggatattaat	aggctgagcg	attacgatgt	cgaccacatt	2520
gtccacacag	cgttctcgaa	ggacgacagc	attgacaaca	aggtgctgac	ccgctcggat	2580
aagaacaggg	gcaagagcga	taatgttcca	agcgaggagg	ttgtgaagaa	gatgaagaac	2640
tactggcggc	agctcctgaa	cgcgaagctc	atcacacagc	ggaagttcga	caacctcacc	2700
aaggctgagc	gcgggggcct	gagcgagctg	gacaaggcgg	gggttcattaa	gaggcagctg	2760
gtcgagacac	ggcagattac	aaagcatggt	gcgcagattc	tcgattcccc	gatgaacacc	2820
aagtacgatg	agaacgataa	gctgattcgg	gaggtcaagg	taattaccct	gaagtcacaa	2880
ctggtgtccg	acttcaggaa	ggacttccag	ttctacaagg	ttcgggagat	caacaactac	2940
caccacgcgc	atgatgccta	cctcaacgcg	gtcgtgggga	ccgctctcat	caagaagtac	3000
ccaaagctgg	agtcagagtt	cgtctacggg	gattacaagg	tttacgacgt	gcggaagatg	3060
atcgctaaga	gcgagcagga	gattggcaag	gctaccgcta	agtaacttct	ctactccaac	3120
atcatgaact	tcttcaagac	agagattacc	ctcgcgaatg	gcgagatccg	gaagaggccc	3180
ctcatcgaga	caaatgggga	gacaggggag	attgtctggg	ataagggggc	ggatttcgcg	3240
accgtccgga	aggctcctgc	gatgccccag	gttaaatattg	tcaagaagac	tgaggtccag	3300
actggcgctc	tctcaaggga	gtcgattctc	ccaaagagga	actccgataa	gctcattgct	3360
cggagaagag	attgggaccc	caagaagtac	gggggatttc	actccccacc	tgttgcttac	3420
tctgttctcg	ttgttgctaa	gggtggagaag	gggaagtcga	agaagctgaa	gagcgtgaag	3480
gagctgctcg	ggattacaat	tatggagagg	tcactcttcg	agaagaatcc	catcgacttc	3540

-continued

```

ctggaggcca agggctacaa ggagggtgaag aaggacctga ttattaagct gccaagtag 3600
tcgctcttcg agctggagaa tgggcggaag cggatgctgg cgcccgggg ggagctgcaa 3660
aaggggaacg agctggcgct cccctccaag tatgtgaact tcctctacct ggctcgcac 3720
tacgagaagc tgaaggggtc cccagaggat aatgagcaga agcagctctt cgtcgagcag 3780
cataagcact acctggacga gattatcgag cagattagcg agttctcgaa gcgggtcatc 3840
ctcgcgatg cgaacctgga taagggtgctc agcgccctaca ataagcaccg ggacaagccg 3900
attcgggagc aggcggagaa tattattcac ctcttcacac tcaccaacct cggggcacca 3960
gctgcgttca agtacttcca cactactatc gaccggaagc ggtacacctc gacgaaggag 4020
gtgctcgacg ccacctctat tcaccagtcg atcacaggcc tgtacgagac acggattgac 4080
ctgtccacgc tcgggggcga c 4101

```

```

SEQ ID NO: 64      moltype = DNA length = 4101
FEATURE
misc_feature       1..4101
                    note = Cas9
source              1..4101
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 64
gacaagaagt actccattgg cctggcgatt gggacaaaact cgggtggggtg ggccgtgatt 60
acggatgagt acaaggttcc aagcaagaag ttcaaggtcc tcgggaacac agatcgccat 120
tcgattaaga agaattctcat tggggcgctc ctcttcgact cgggggagac agcggaggct 180
accaggctca agcggacagc caggcgcgcg tacacaaggc ggaagaatcg catctgctac 240
ctccaggaga tttctcgaa tgagatggcg aaggtggacg acagcttctt ccacggctg 300
gaggagtctc tcctggtgga ggaggataag aagcacgaga ggcattccat ttccgggaac 360
acgctggagc aggttgcgta ccatgagaag taccctacaa tctaccatct gcggaagaag 420
ctggttgact ccacagacaa ggcggacctg aggtgatctt acctcgctct ggcacacatg 480
attaagttcc cggggcattt cctgatcgag ggggacctga atcccgacaa ttcggatgtg 540
gacaagctct tcatccagct ggtgcagacc tacaaccagc tgttcgagga gaatcccatc 600
aatgcgtcgg gcgttgacgc taaggccatt ctgtccgcta ggctgtcgaa gagcaggagg 660
ctggagaacc tgatcgccca gctgccaggc gagaagaaga atgggctctt cgggaatctg 720
attgcgtctt cctggggctt gacaccgaac ttcaagagca atttcgatct ggctgaggac 780
gcgaagctcc agctctcgaa ggacacttac gacgatgacc tcgataacct cctcgcgag 840
atcgggggac agtacgctga tctcttctc gccgctaaga acctctcgga tgcatactg 900
ctctccgaca tttccgggtt taataccgag attacaaaagg cccactgtc ggctgcatg 960
atcaagcggg acgatgagca tcatcaggat ctacacctgc tcaaggccct cgtcgcgag 1020
acgtgcctga agaagatcaa ggagattttc ttcgaccaga gcaagaatgg gtacgtggtc 1080
tacattgacg gcggggcctc acaggaggag ttctacaagt tcatcaagcc aatcctggag 1140
aagatggatg ggcagagaga gctgctggtg aagctcaacc gggaggatct gctcaggaa 1200
cagcggagct tgacacacgg ttctgatccc catcagatcc acctggggga gctgcacgcg 1260
atcctcgccc ggcaggagga ttctaccctt ttctgaagg ataactcgga gaagatcgag 1320
aagattctca cctccggat tccctactac gtccggccac tcgccccggg caatagcagg 1380
ttcgctgga tcgacacgga gagcagaggag acaatcaccc cctggaactt cgaggagggt 1440
gtcgacaagg gggcgctccg ccagtcattc ccaagagcat agcctcctct acgagtactt cactgtttac 1500
ctgccaatg agaaggttct cccaaagcat agcctcctct acgagtactt cactgtttac 1560
aacgagctga ccaaggtgaa gtatgtgacc gagggcatgc ggaagccgcg gttctgtctc 1620
ggcgagcaga agaagggcct ttgggacctc ctgttcaaga ccaatcgcaa ggtcacagtc 1680
aagcagctca aggaggatta ctccaagaag atcgagtgtc tcgactcggg tgagattagc 1740
aggggtgagg atcggttcaa cgcgagcctc ggcacttacc acgacctct gaagatcatc 1800
aaggataaag acttctcga caacgaggag aacgaggata ttctggagga catcgtgctc 1860
accctgacgc tgttcgagga tcgggagatg atcgaggagc gcctgaagac ctacgctcat 1920
ctcttcgatg ataaggctat gaagcagctg aagaggaggc ggtacaccgg gtggggccgc 1980
ctgagcagga agctcattaa ggggatcagg gacaagcaga gcggcaagac catcctggag 2040
ttctccaaga gcgatggctt cgccaaccgg aatttcatgc agctcatcca cgacgactcc 2100
ctcaccttca aggagggcat tcagaaggct caggtcagcg gccaggcgca ctcgctgcat 2160
gagcacatcg ctaacctggc gggcagccca gccatcaaga agggcatcct ccagacagt 2220
aaggctcgtg atgagctggt gaaggctcat ggcggcgata agcccgagaa tattgtgatt 2280
gagatggcgc gggagaatca gacctctcag aagggccaga agaactcgcg ggagcgcagt 2340
aagaggatcg aggggggat taaggagctg ggcagccaga ttctcaagga gcaccccggt 2400
gagaataccc agctccagaa cgagaagctg tacctctact acctccagaa tgggcgggac 2460
atgtatgttg atcaggagct ggaacatcaat cgctctcggg attacgacgt ggaccacatc 2520
gtgccccaga gcttctgaa ggatgatagc atcgacaata aggtcctgac ccgctccgac 2580
aagaatcgcg gcaagagcga caacgtgccg agcaggaggc tcgtgaagaa gatgaagaac 2640
tactggcgcg agctgtgaa cgcgaagctc attacacagc ggaagttcga taacctgagc 2700
aaggcgagga gggcgcgctt ctccgagctg gacaaggcgg gcttcattaa gaggcagctc 2760
gtggagactc gccagatcac caagcagctg gctcagatcc tcgatagccg gatgaatacg 2820
aagtacgatg agaatgacaa gctcatccgg gaggtgaagg taatcacctt gaagtcaaac 2880
ctcgttagcg atttccgaa ggaactccag ttctacaagg tcggggagat taaccaactc 2940
catcatgcgc acgatgcgta cctcaatgcg gtggtgggca cagccctgat taagaagtag 3000
cccaagctgg agagcgagtt cgtctacggg gactacaagg tgtacgatgt tcggaagatg 3060
atcgccaaga tctgacagga gattgggaag gccacgccta agtacttctt ctactcgaat 3120
attatgaatt tctcaagac cgagatcaca ctgcctaagc gggagattcg gaagcgccc 3180
ctcatcgaga ctaacgggga gactggcgag attgtgtggg acaaggggag cgacttcgct 3240
accgtgcgca aggtcctctc gatgccccag gttaaatatt ttaagaagac agagggtcag 3300
acggcggggt tctccaagga gcttatcctg ccgaagcgga actcgacaa gctgatcgcc 3360
cgcaagaagg attgggacct caagaagtag gggggattcg atagcccaac cgtggcttac 3420

```

-continued

```

agcgtcctcg tggtcgcca ggttgagaag gggaagtcga agaagctcaa gagcgtaag 3480
gagctgctcg gcatcaccat catggagcgg tccagcttcg agaagaatcc tatcgacttc 3540
ctggaggcta aggggtacaa ggaggccaag aaggacctga tcattaagct gcccaagtac 3600
tctctgttcg agctggagaa cgggaggaag cggatgctgg cgtctgctgg cgagctacag 3660
aagggaatg agctggcgct cccctcgaag tatgtcaact tctctacctt ggcttcccat 3720
tacgagaagc tgaagggctc gcccgaggat aatgagcaga agcagctctt cgtggagcag 3780
cacaagcact acctcgacga gatcattgag cagatttcgg agttctcgaa gcgggtcatt 3840
ctcgcgagcg cgaacctcga caaggtctct tggcggtaca acaagcaccg ggacaagccc 3900
atccgggagc aggcgagaa cattatccac ctcttcacac tgaccaacct cggcgctccc 3960
gccggttca agtacttcga caccaccatt gaccgcaaga gatacacatc caccaaggag 4020
gtgctggagc cgacctcat ccaccagagc atcacaggcc tctacgagac acggatcgac 4080
ctctcgagc tggggggcga t 4101

```

```

SEQ ID NO: 65      moltype = DNA length = 4092
FEATURE            Location/Qualifiers
misc_feature        1..4092
                    note = Cas9
source              1..4092
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 65
gacaagaagt actcgatcgg cctggcgatt ggcacaaaca gcgtgggggt ggctgtgac 60
actgatgagt acaaggtgcc atcgaagaag ttcaaggtgc tggggaatac agaccggcat 120
tcgatcaaga agaattctat tggcgctctc ctcttcgatt ccggcgagac tgcgtaggcg 180
accgcctga agcgcaccgc ccggcggcgc tacactcggc ggaagaatag gatttgctac 240
ctccaggaga tttctcgaa tgagatggcc aaggtggatg acagcttctt ccaccgctg 300
gaggagtcgt tctcgttcga ggaggacaag aagcatgagc ggcaccttat ctccgggaat 360
atcgttgatg aggtcgcccta ccacgagaag taccocacta tctaccatct ccgcaagaag 420
ctcgtggaca gcacagataa ggcgcacctc cgcctgatct acctcgccct cgcgcacatg 480
attaagttcc gggggcactt cctcattgag ggggatctga atcccgataa ctccgacgtg 540
gacaagctgt tcattccagc ggtgcagaca tacaaccagc tgttcgagga gaatcccatc 600
aacgcgagcg cgtgtggcgc taagggccatt ctgtcggcta ggctctcgaa gtcgagggcg 660
ctggagaacc tgattgcgca gctcccgccg gagaagaaga acgggctgtt cggaatctc 720
atcgccctct ccctcggcct caccacaaac ttcaagagca attcgacct ggctgaggac 780
gctaagctgc aactctcaaa ggatacatac gatgacgacc tggacaatct cctggctcag 840
atcggcgacc agtacgctga cctgttctct cgggccaaga atctgtcgga cgcgattctc 900
ctcagcgaca tctgcgcgt caataccgag attacgaagg ctccactgtc tgcgtcaatg 960
attaagcggg acgatgagca tcaccaggat ctgacctctc tgaaggcgct cgtgcggcag 1020
cagctgcccg agaagtacaa ggagattttc ttcgatcaga gcaagaatgg ctacgcggcg 1080
tacatcgacg gggcgcgagc ccaggaggag ttctacaagt tcataaagcc catcctggag 1140
aagatggacg gcacgcagga gctactcgtg aagctcaatc gggaggatct cctccggaag 1200
cagcggacat tcgataacgg gtctatccca caccagatcc acctcggcga gctgcatcgc 1260
attctgcggc ggcaggagga ttctaccctt ttctgaagg acaaccggga gaagatcgag 1320
aagatcctca cattccggat tccatactac gtcggccccc tggcgagggg caatagccgg 1380
tccgctgga tgacaaggaa gtcgagggag actattaccc cgtggaattt cgaggagggtg 1440
gttgacaagg gcgcttcgcg gcagagcttc attgagcgga tgacaaactc cgacaagaat 1500
ctccccaacg agaaggtcct gcggaagcat agcctcctgt acgagtactt caccgtctac 1560
aatgagctaa ctaaggtcaa gtatgtgaca gaggcgatga ggaagccagc ctctctctca 1620
ggcgagcaga aagaaggccat tttggacctc ctgttcaaga caaacccgaa ggtgacagt 1680
aagcagctga aggaggatta cttcaagaag attgagtgtc tcgactcagt ggagatttca 1740
ggcgtggagg atcgggtcaa cgcgagcctg gggacttacc acgacctgct gaagattatt 1800
aaggacaagg acttctcgga taacgaggag aatgaggaca tctggagga tattgtgctc 1860
acctcacccc tgttcgagga cagggagatg attgaggaga ggctcaagac ctacgcgcac 1920
ctgttcgatg acaaggtcat gaagcagctg aagaggcgcc gctacactgg gtggggcgcg 1980
ctgtcgcgga agctcgatcaa cggcattcgg gataagcagt ccgggaagac cattctgga 2040
ttctgaagt cggacggctt cgcacaacagg aatttcatgc agctgatcca cgacgactcc 2100
ctcaccttca aggaggacat tcagaaggcc cagggttagcg gccaggggga ctcactccac 2160
gagcatattg ccaatctggc cggctctcca gctatcaaga agggcatcct gcaaacagtt 2220
aaggttgttg acgagctggt taaggtcatg gggcggcata agcccgagaa cattgtcatc 2280
gagatggctc gggagaacca gacaactcag aagggccaga agaactccag ggagcgcatg 2340
aagcggattg aggaggccat taaggagctg ggggtccaga tctcaaggga gcacctgtc 2400
gagaacactc agctgcaaaa cgagaagctc tacctgtact acctccagaa cgggcgggat 2460
atgtatgtgg atcaggagct ggacatcaac aggtctctcg actacagct ggatcacatt 2520
gtccacagct ctttctcaa ggatgattcc atcgacaaca aggtgtgtgac gcgcagcgac 2580
aagaataggg ggaagtcgga caacgttccg agcgaggagg tcgtgaagaa gatgaagaat 2640
tactggaggc agctcctgaa tgcgaagctg atcactcaga ggaagttcga caatctgaca 2700
aaggcggaga gggcggggct ctcggagctg gataaggcgg gcttcatcaa gcggcagctc 2760
gttgaaaccc ggagatcac caagcatgtc gccagatcc tcgatagccg catgaacacc 2820
aagtacgatg agaacgacaa gctcattcgg gaggttaagg tcattacgct gaagtccaag 2880
ctcgtcagcg actcaggaa ggaattccag ttctacaagg ttcgggagat taacaactac 2940
caccacgcgc atgatcgcta cctgaacgct gttgtcgga ctgctctcat caagaagtac 3000
ccaaagctgg agtccgagtt cgtctacggg gactacaagg tctacgatgt ccggaagatg 3060
atcgccaagt cggagcagga gatcgggaag gctactcgga agtactctt ctacagcaac 3120
attatgaatt tcttcaagac ggagattacg ctggcgcaac gggagattag gaagaggccc 3180
ctcattgaga ctaatgggga gacaggcgag attgtttggg acaaggcccg cgacttcgcg 3240
actgtgcgga aggtcctgtc catgccacag gtgaattatt ttaagaagac agaggtgcag 3300

```

-continued

actgggggct	tctcgaagga	gagcattctc	ccaaagcgga	acagcgataa	gctcatcgcg	3360
cgcaagaag	attgggaccc	taagaagtac	ggcggttcg	attctcccac	tgtggcctac	3420
tcogttctcg	tggttgccaa	ggttgagaag	gggaagtcga	agaagctgaa	gtcggccaag	3480
gagctgctcg	ggattacaat	catggagcgg	agcagcttcg	agaagaaccc	tattgatttc	3540
ctggaggcca	agggtacaa	ggagggttaag	aaggatctca	ttatcaagct	ccctaagtac	3600
tctctgttcg	agctggagaa	tggccggaag	aggatgctgg	cctcggtcgg	cgagctacag	3660
aaggggaaatg	agctggccct	cccgtcgaag	tatgtgaatt	tcctgtacct	cgcgtcgac	3720
tacgagaagc	tcaagggcag	cccggaggat	aatgagcaga	agcagctctt	cgtggagcag	3780
cataagcact	acctggacga	gatcattgag	cagatcagcg	agttctcgaa	gcggttatt	3840
ctggctgatg	ctaaccctgga	caagggtctg	agcgcctaca	ataagcatcg	cgacaagccg	3900
attcgcgagc	aggcggagaa	tattatccac	ctgttcaccc	tcactaacct	cggggctccc	3960
gcgcccttca	agctacttga	taccacaata	gataggaagc	ggtacacctc	gacgaaggag	4020
gtcctcgacg	ccacactcat	ccatcagtcg	attacagggc	tgtacgagac	acggattgac	4080
ctctcgacg	tg					4092

SEQ ID NO: 66 moltype = DNA length = 4101
 FEATURE Location/Qualifiers
 misc_feature 1..4101
 note = Cas9
 source 1..4101
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 66

gacaagaagt	attccatagg	cctggctatc	ggcaccaca	gcgtgggctg	ggcgcgtcatc	60
accgacgagt	acaaagtggc	gagtaaaaa	ttcaaagtgc	tcggcaacac	cgaccgccac	120
tcataaaga	aaaacctgat	cggggcgctc	ctgttcgaca	gcggcgagac	ggcggaggcc	180
accgccttga	aacgcacggc	ccgacggcgc	tacacggcgc	gcaagaacccg	gatctgttac	240
ctacaggaga	ttttctctaa	cgagatggcg	aagggtggacg	actcgttctt	tcaccgcctc	300
gaagagtctc	tcctcgtgga	ggaggacaag	aaacacgagc	gccaccgat	cttcggcaac	360
atcgtggacg	aggtggccta	ccacgagaag	tacccgacca	tctaccacct	ccggaagaaa	420
ctcgtggaca	gcacggacaa	ggcgcacctg	aggctcatct	acctgcacct	ggcgcacatg	480
attaagtctc	ggggccactt	ctcgatcgag	ggcgacctga	accgggacaa	cagcgacgtg	540
gacaagctgt	tcattccagct	agtcacagac	tacaaccagc	ttttcgagga	aaaccccatc	600
aacgccagcg	gggtggacgc	gaaggcgatc	ctgtccgccc	ggctgagcaa	gtcccgccg	660
ctggagaacc	tcatcgcgca	gttgcccggc	gagaagaaga	acgggctgtt	cggaacccg	720
atcgccctct	ccctgggggt	caccccgaac	ttcaagtcca	acttcgacct	cgccgaggac	780
gccaaactac	agctgagcaa	ggacacctac	gacgacgacc	tcgacaacct	gctggccacg	840
atcggggacc	agtagcgaga	ctgtgtcttc	gccgccaa	acctctccga	cgcctcctg	900
ctgtcgga	ctctcgccgt	gaacacggag	atcacgaagg	ccccgctctc	ggcctcgatg	960
atataacgct	tcgacgagca	ccaccaggac	ttgacctctc	tcaaggcgct	ggctccgccc	1020
cagcttcccg	agaagtacaa	ggaaatcttt	ttcgatcaga	gcaagaacgg	gtacgcggg	1080
tacatcgacg	gcggggcgct	ccaggaggag	ttctacaagt	tcattcaagc	catcctggag	1140
aaaatggacg	ggacggagga	gctgctcgtg	aagctcaacc	gcgaagattt	gctccgcaag	1200
cagcgcacgt	tcgacaacgg	gtcgatcccg	caccagatcc	acctggggcg	gctgcacgg	1260
atcctcaggc	gtcaggga	cttctacccc	ttcctcaagg	acaaccgcga	gaagatagag	1320
aagattctga	ccttcagaa	tccttattac	gtgggcccgc	tggctcgggg	caactcgccg	1380
ttcgccctga	tgacgcgcaa	gtccgaggag	acctcacccc	cgtgggaact	cgaggagggtg	1440
gtggataaag	gtgcctcggc	ccagctcctt	atcgagcgga	tgaccaacct	cgacaagaac	1500
ctgccgaacg	gtgaaggtgct	ccccaaagc	agcctgctct	acgaatattt	cacggtgtac	1560
aacgagctga	cgaaggtcaa	gtactgtgac	gaggggaatga	ggaaacctgc	attcctctcc	1620
ggggagcaga	agaagccat	agtcgacctc	ctgttcaaga	ccaaccggaa	ggtcaccgtc	1680
aagcagctca	aggaggacta	cttcaagaag	atcgagtgtc	tcgattcagt	ggagatcagc	1740
ggcgtcgagg	accggttcaa	cgccagccctg	ggcactctacc	acgacctgct	caagatcatc	1800
aaggacaagg	acttctctga	caacgaggag	aacgaggaca	tcctggaggga	catcgtgctg	1860
accctgacgc	tcctcgagga	cgcgagatg	atcgaggagc	gcctcaagac	ctacgccac	1920
ctgttcgacg	acaaggtgat	gaagcagctc	aagcggcgga	gataactg	gtggggccgc	1980
ctctcccgga	agctcattaa	cgttatcagg	gataaagcgt	ccgggaagac	gatcctcgac	2040
ttctcaagt	cggaagggtt	cgccaaccgc	aacttcatgc	agctcatcca	cgacgactcc	2100
ctgacgttca	aggaggacat	ccagaaggcc	caagtgtctg	gtcaagggtga	ctcgctccac	2160
gagcacatcg	ccaacctcgc	gggcagcccc	gccatcaaga	agggaaact	ccagaccgtc	2220
aaggtggtgg	acgagctggt	gaaggtcatg	ggccgccaca	agccggagaa	catcgtcatc	2280
gagatggcgc	gggagaacca	gaccacgcag	aaggggcaga	aaaatagccg	tgagcgcag	2340
aagcgcacg	aggaggggat	taaggagtgt	ggcagccaga	tcctcaagga	gcacctgtg	2400
gagaacacgc	agttgcaaaa	cgagaagctc	tacctgtact	acctccagaa	cgggagggat	2460
atgtacgtgg	accaagaact	ggacatcaac	cgcctgtccg	actacgacgt	ggaccacatc	2520
gtgccgcaga	gcttctctaa	ggacgacagc	atcgacaaca	aggtgctcac	ccggtccgac	2580
aagaatcggg	gcaagtccga	caacgtgccc	agcgaggagg	tcgtcaaaaa	gatgaaaaac	2640
tactggcgac	aactactgaa	cgccaagctc	atcacccagc	gcaagttcga	caacctcaca	2700
aaagccgagc	gcggcggtt	gagcagctg	gacaaggccg	gggtctatcaa	gcgccagctc	2760
gtcgagacgc	gccagatcac	gaagcagctc	gcgcagatac	tcgacagccg	gatgaacacc	2820
aagtacgacg	agaacgacaa	gctcatccgg	gaggtgaagg	tcataccctc	caagtcgaa	2880
ctcgtgagcg	acttccgcaa	ggaactccag	ttctacaagg	tcggggagat	caacaactac	2940
caccacgccc	acgatgctta	ttttaacgccc	gtggtgggga	cgccctcat	taagaaatac	3000
cgaagctgg	agtccgagtt	cgtgtacggc	gactacaagg	tgtacgacgt	caggaagatg	3060
atcgccaagt	ccgaacagga	gatcgggaag	gccacggcga	aatacttctt	ctacagcaac	3120
atcatgaact	ttctcaagac	cgagatcacc	ctcgccaacg	gcgagatccg	caagcgcgcc	3180

-continued

```

ctcatcgaga cgaacgggga gacgggcgag atcgtctggg acaagggggc cgaattcgcc 3240
actgtgcgga aggtgctgtc gatgcccag gtcaacatcg tcaagaagac ggaggtcag 3300
acgggcgggt tcagcaagga gagcatcctg ccgaagcgca acagcgacaa gctgatcgcc 3360
cgcaaaaagg actgggatcc aaaaaagtac ggcggtctcg acagcccccac cgtcgccatc 3420
agcgtctctg tcgtogctaa agtcgagaag ggcaagtcca aaaagctcaa gacggtcaa 3480
gagctgctcg ggatcaccat catggagcgg tccagcttcg agaagaacct aattgatttc 3540
ctggaggcga aggggtacaa ggagggtcaag aaagacctca tcataaagct gccgaagtac 3600
tcactcttcg agctggagaa cgggcgcaag cggatgctgg cgtcgcccg agagctccaa 3660
aagggaacgc agctggcgct gccgagcaag tacgtgaact tccctacct ggctgccac 3720
tacgagaagc tcaaggcgag tccagaggat aacgagcaga agcagctatt cgtggagcag 3780
cacaagcact acctggacga gatcatcgag cagatcagcg agttctccaa gcgcgtcatc 3840
ctggcgagcg ccaacctgga caaggtgctg tccgcgtaca acaagcaccg cgacaagccg 3900
atccgcgagc aagccgagaa catcatccac ctgttcaccc tcacgaacct cggggcaccc 3960
gcgccttcca aatatttcga caccgacctc gaccgcaagc gctacaccag cagcaaggcg 4020
gtgctcgacg ccacctgat ccaccagagc atcacccggc tgtacgagac ccgcatcgac 4080
ctctcgacgc tcggcgggga c 4101

```

```

SEQ ID NO: 67      moltype = DNA length = 4101
FEATURE
misc_feature       1..4101
                    note = Cas9
source              1..4101
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 67
gacaagaagt acagtattgg attggccatc gggacgaaca gcgtgggctg ggccgctatc 60
accgacgagt acaagggtgcc atccaagaag ttttaaggttc tgggggaatac cgaccgccac 120
tcgatcaaga aaaatctcat cggggcgctg cttttcgaca gcggcgagac ggccgaagcg 180
acggcgctca agcgagcgcc tcgtcgccgt ttaccccggc gtaagaacct catctgttac 240
ctccaggaga tatcagcaa cgagatggcg aaggtggacg actcctttt ccaccgtctt 300
gaggagtcct tcctggtcga ggaggacaag aagcacgagc gccaccgat cttcgggaac 360
atcgtggagc aggtggcctc ccacgagaag taccaccaga tctaccacct ccgcaaaaaa 420
ctcgtggact caactgacaa ggccgatttg aggttatctt acctgcacct cgccacatg 480
attaagttcc gtgggcctca cctaactcga ggtgacctca acccgacaa cttgacgtg 540
gacaagctgt tcattccagct tgtgcagacc tacaatcagc tctttgagga gaatccgatc 600
aacgcatctg tgtgtgagcg aaaggccatc ctcagcgcg gcgtgagcaa gtctaggcgg 660
ttggagaacc tgatcgccca actgcccggc gagaagaaaa atggcctctt cgcaacctg 720
atcgccctgt cgtgggggct caccgcgaac ttcaagagta actttgacct ggccgaggac 780
gctaagctcc agctatctaa ggacacatac gacgacgacc tggacaacct gctggcccag 840
atcgcgacc tctcgccga cctcttcta gccgccaaga acctgtccga cgccatctc 900
ctcagcgaca tctgcgcgt gaacacggag atcacgaagg ctcgctcag cgccctcatg 960
attaagcgtg acgacgagc ccaccaagac ctaactttac tcaagccct cgtgcggcag 1020
cagcttcccg agaatcaaa agagatatct tttgatcagt ccaagaacgg ttatgcgggc 1080
tacatcgacg gcggcgcgag ccaggaggag ttctacaagt tcatcaagcc catcctggag 1140
aagatggacg gcacggagga gctgctcgtg aagctcaacc gtgaagacct cctgcgaaag 1200
cagcgaacct tcgacaacgg ttcgatcccg caccagatcc acctggggga gctgcacgcc 1260
atcctgaggg gacaggagga cttctacctt ttcctaaagg acaaccgca gaagattgaa 1320
aaaatcctga cgtttcgcat acctactac gtccgcccgc tggcgcgcg caactcccgg 1380
ttcgcttggg tgaccctgaa gagcgaggag acgatcacc cgtggaaact cgagagggtc 1440
gtggacaagg gcgcgagcgc gcagagcttc atcgagcgca tgaccaact cgacaagaac 1500
ctccgcaacg agaaggtgct cccaaagcac tccctcctgt acgagtatt caccgtgtac 1560
aacaggttga caaaggtgaa gtacgtgacg gagggaaatg ggaagcctgc gttcctctc 1620
ggcgagcaga agaaggcaat cgtggacctg ctcttcaaga ccaaccggaa ggtgacggtg 1680
aagcagctca aggaggacta cttcaaaaaa atcgagtgtc tcgactccgt ggagataaag 1740
ggcgtggagg accgattcaa gcctccctc ggcaacctacc acgacctct taagatcatc 1800
aaggacaagg acttcttggg caacgaggag aacgaggaca tcctggagga catcgtgctc 1860
acctgacccc tcttcgagga ccgggagatg atcgaggagc gcctcaagac gtacgccac 1920
ttgttcgacg acaaggtgat gaagcagctc aagcggcgcg gatcacccg gtggggcccg 1980
ctatcccgca aacttatcaa cggcatccgc gacaagcagt ccggcaagac gatcctggat 2040
ttcctcaagt cggacgggtt cgccaaccgg aactcatgc agctcatcca cgacgacagc 2100
ctcacgttca accaggtcat ccagaaggcc caagtgagcg gtcaagggga cagcctccac 2160
gagcacattg cgaaccttgc tgggagccct gcgatcaaga aggggatatt gcaaacctg 2220
aaggtcgtgg acgagtgtg gaaggtcatg gggcgacaca agcccgagaa catcgtgatc 2280
gagatggcca gggaaaatca gaccacgcag aagggccaaa aaaacagccg cgagcggatg 2340
aagcggatcg aggaggctc caaggagctg gggtcgcaga tctcaagga gcaccgggtg 2400
gagaacacgc agctccagaa cgagaagctg tacctctatt acctacagaa cggcggggat 2460
atgtacgtgg accaggagct agacatcaac cgcctgtccg actacgacgt ggaccatctc 2520
gtcccgcagt cgttcttgaa ggacgacagc atcgacaaca aggtgctcac aagatcggat 2580
aagaatcgag gcaagtccga caacgtgccc tcggaggagg tggtaagaa aatgaaaaac 2640
tactggcgcg agtgctgtaa cgccaagctc attacgcagc ggaagttcga caacctgacg 2700
aaggctgaac ggtgtgggct cagcgagcta gacaaggcgg ggttcatcaa gcggcagctc 2760
gtcgagaccc ggcagatcac caagcagtg gcgcagatcc tggactcgcg catgaacacc 2820
aagtacgacg agaacgacaa gctcatccgt gagggtgaag tcatcaccct taagtctaag 2880
ctggtcagtg acttcgcaa ggacttccag ttctacaagg tccgggagat caacaactac 2940
caccacgcgc acgacgcta cctcaacgcg gtgggtggga cggcgcttat taagaatat 3000
cccaagctgg aaagcgagtt cgtttacggc gactacaagg tgtacgacgt ccgcaagatg 3060

```

-continued

```

atcgcaaaagt cggaacagga aatcggaag gcgacggcca aatatttctt ttactccaac 3120
atcatgaatt tttttaagac ggagatcacc ctggcgaaacg gggagatccg caagcggccc 3180
ctcatcgaga ccaacgggga gacgggagcgc atcgtctggg acaagggccg ggacttcgcc 3240
accgtgcgga aggtgctttc tatgctctca gtcaatatcg tcaaaaagac agagggtgcag 3300
accggcgggg tcagcaagga gtctatcctg ccgaagcgca actcggacaa gctcatcgcg 3360
cgcaagaaag actgggaccc caaaaaatat ggcggggttcg actcgccgac cgtcgctac 3420
agcgtctctg tgggtggctaa ggtcgagaag ggcaagagca aaaagctaaa gtcgggtgaag 3480
gagctgctgg gcatcaccat catggagcgc tcgtcttttc agaagaatcc aatcgacttc 3540
ctagaggcga aggggtacaa ggagggtcaaa aaggatctta tcatcaaaact gccgaagtac 3600
agtctgttcg agctggagaa cgggcggaag cggatgctgg ctagtgcggg cgagtgcag 3660
aaggggcaacg agttggcact gccctccaag tacgtgaact tcctgtacct ggctccccc 3720
tacgagaagc tcaaggggag ccccgaggac aacgagcaga agcagctatt cgtcgagcag 3780
cacaagcact acctggagca gatcatcgag cagatcagtg agttctccaa gcgggtcatc 3840
ctcgcgagcg caaacctgga caaggtgctg agcgcgtaca acaagcacag ggacaagcca 3900
atcagggaaac aggcgagaa catcatccac ctgttcaccc tgaccaacct ggggtgcaccg 3960
gctgccttca agtactttga caccagcacc gaccggaagc gctacacctc cacgaaggag 4020
gtgctggacg ccacgctgat ccaccagagc atcacgggcg tctacgagac acggatcgac 4080
ctgagccagc ttggcgggga c 4101

```

```

SEQ ID NO: 68      moltype = DNA length = 4092
FEATURE           Location/Qualifiers
misc_feature       1..4092
                  note = Cas9
source            1..4092
                  mol_type = other DNA
                  organism = synthetic construct

```

```

SEQUENCE: 68
gacaaaaagt attccattgg actcgtatc ggcacgaaca gcgtcgggtg ggcgggtcatc 60
actgacgagt acaaggtgcc gagcaagaag tttaaggtgc tgggaaacac cgacaggcac 120
tcgatcaaga aaaatcttat cggggcccta ctcttcgact ccggagaaac cgcgagggcc 180
accgggttga agcgacgggc ccgctgtcgc tacaccaggc gcaagaaccg gatctgtac 240
ctccaggaga tatcagcaa tgagatggcg aaggtggacg actcgttttt tcacaggcta 300
gaggagtctt tcctcgtgga ggaggacaag aaacacgagc gccaccccat cttcggcaac 360
atcgtggatg aggtggcata tcacgagaag taccacaacca tctaccacct ccgcaaaaag 420
ctcgtggact ctaccgacaa ggcgacacct cgtctgatct acctcgcgct ggcccacatg 480
attaagttcc gaggacactt tctgatcgag ggcgacctga acccagacaa cagcgacgtg 540
gacaagctgt tcatccaact ttccagacc tacaatcagc tcttcgagga gaacctatc 600
aacgcctcgg cgtgggacgc gaaggccatc ctgtccgccc gcctgagcaa gtcgcgggcg 660
ctggagaacc tgatcgccca gctccccggc gaaaaaaga acggcctctt cggcaacctc 720
atcgcggtgt cctcggggct caccgccaa tcaagtcca acttcgacct ggcgagggac 780
gctaaaactc agctctcgaa ggatacctac gacgacgacc tcgacaacct gctggcccg 840
atcggcgacc agtacgggga ccttttctcg cgggccaaag acctgagcga cgcgatcctc 900
cttagcgaca tactcgtgt taacaccgag atcacgaagg ccccgctctc cgcgtccatg 960
attaagcgct acgacgagca ccaccaagac cttaccctgc ttaaggcgct ggtcaggcag 1020
cagttaccgg agaagtacaa ggagatcttt ttgatcaat ctaagaacgg ttacgcccgg 1080
tacatcgacg cgcgcgcgtc caaggaggag ttctacaagt tcatcaagcc gatctggag 1140
aaaatggacg ggaccgagga gctcgtcgtg aagctcaacc gcgaagacct cctccgcaag 1200
cagcgcacct tcgacaacgg gtcgcatccc caccagatcc acctgggaga gctgcacgcg 1260
atcctgcgga gacaagagga ctctacccc ttctcaagg acaaccggga gaagattgaa 1320
aaaatactta cttttcgtat ccggtactac gtcgggcccc ttgcgagggg caactccaga 1380
tccgctgga tgacccgcaa gtcgagggag acctacccc cgtggaaact cgaggagggtg 1440
gtggacaagg gcgcgtcggc ccagtcgttc atcgagcgca tgaccaactc cgacaagaac 1500
cttccgaacg agaaggtgct ccggaagcac agcctgctct acgaatatct tactgtgtac 1560
aacgagctga cgaaggtcaa gtacgttacg gaggggatga ggaagcccg cttcctctcc 1620
ggcgagcaga agaaagccat tgtggatctc ctgttcaaga ccaaccgcaa ggtgacgggtg 1680
aaacagctca aagaggacta cttcaagaag atcgagtgtc tcgactccgt agagatcagc 1740
ggggtcgagg accgcttcaa cgcctcgtg ggcacgtacc acgacctgct aaagattatc 1800
aaggacaaaag acttctaga caatgaggag aacgaggaca ttctggagga catcgtgctg 1860
actctgacgc tgttcgaaga ccgcgagatg atcgaggagc ggcttaagac gtacgcccac 1920
ctgttcgacg acaaggtgat gaagcagttg aaacggcggc gctacacccg gtggggcgcg 1980
ctctcccga gaactcctaa cgcctcccgc gacaagcagt cggggaagac gatcctggac 2040
ttctcaaga gcgacggctt cgcacaaccg aacttcatgc agctaatacc cgacgacagc 2100
ctgacgttca aggaggacat ccagaaggcc caagtgagcg gccagggaga ctcgctacac 2160
gagcatatcg ccaacctggc tggcagcccc gcgattaaga aaggaatcct ccaaaccgtc 2220
aaagtgggtg acgagctggt gaaggtgatg ggccgccaca agcccgagaa cattgtgatc 2280
gagatggcgc gggagaacca gacgacgcag aagggccaaa aaaatagcag ggaagggatg 2340
aagcgaatag aggaggggat caaggagctg gggagccaga ttctcaaaag gcacccggtc 2400
gagaacacac agctccagaa cgagaagctg tacctctact acctccaaaa cggcccgcat 2460
atgtactgtg accaggaact agacatcaac cggctgagcg actatgacgt ggaccacatc 2520
gtgcgcgagt ccttctctaa ggacgactcg attgacaaca aagtgtcac tagatccgac 2580
aagaacagag gcaagagcga taacgtcccg tcggaggagg tcgtcaagaa aatgaaaaac 2640
tactggcggc acctcctaaa cgccaagctc atcacgcagc gtaagttcga caacctgacg 2700
aaggcgagcg gggcggggct gagcgagctg gacaagcgcg ggttcacaa gcggcgagctc 2760
gttgagacgc ggcagatcac aaagcacgtc gcgcaaatcc tcgactcccc catgaacacc 2820
aagtacgacg gaacgacaa gctcatcccg gaggtgaagg tcattacctc taaatcgaag 2880
ctcgtcagcg actttcgtaa ggacttccag ttctacaagg tcagagagat caacaactac 2940

```

-continued

caccacgccc	acgacgccta	tctgaacgcc	gtggtgggca	ccgcgcttat	taagaagtac	3000
cccaagctgg	agtcogagtt	cgtgtacggc	gactacaagg	tttatgacgt	caggaagatg	3060
atcgccaagt	cggaacagga	gatcggaaaa	gctaccgcca	aatatttctt	ctatagcaac	3120
atcatgaact	tcttcaaaac	cgagatcacc	ctcgccaacg	gcgagatccg	gaagcgccc	3180
ctcatcgaga	ccaacgggga	gacgggggag	atcgtctggg	acaaggggcg	ggacttcgct	3240
actgtccgaa	agggtgcttc	catgccacaa	gtgaatatcg	tcaagaaaac	agagggtcag	3300
accggagggt	tcaagtaagga	gtccatcctg	ccaagcgga	actccgacaa	gctaattgct	3360
cgcaaaaagg	attgggatcc	taaaaaatat	ggcggtctcg	actcgcccac	ggtcgcctac	3420
tctgtgctgg	tctgtggcgaa	gggtggagaag	ggcaagtcca	agaagctcaa	gagcgtcaag	3480
gagctgctgg	ggatcacgat	catggagcgt	agttcgtttg	agaagaatcc	catcgacttc	3540
ctggaggcta	aggggtacaa	ggagggtcaaa	aaggacctca	tcattaaagt	gccgaagtac	3600
agcctcttcg	agctggagaa	cgggcgggaag	cgtatgctcg	cctccgctgg	ggagttacaa	3660
aaggggaacg	agctggcgct	gccgtctaag	tacgtcaact	tccgtgaact	ggcctccac	3720
tacgagaagc	tcaaggagtc	gccggaggac	aacgagcaga	agcagctctt	cgtagagcag	3780
cacaagcact	acctggacga	gatcatcgag	cagatttcag	agttctcaaa	gcgggtcatc	3840
ctcgccgacg	ccaacctgga	caaggtgctc	tgggcctaca	acaagcaccg	ggacaagccg	3900
atccgcgaac	agccgaaaaa	catcatccac	ctgttcacgc	tcaccaactc	cggtgcctcg	3960
gcggccttca	agctacttga	cacgaccatc	gaccggaagc	gctatacctc	gacgaaggag	4020
gtgctggacg	ccaccctgat	ccaccagtc	atcacggggc	tttacgagac	ccggaatcgac	4080
ctctcgacgc	ta					4092

SEQ ID NO: 69 moltype = DNA length = 4101
 FEATURE Location/Qualifiers
 misc_feature 1..4101
 note = Cas9
 source 1..4101
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 69

gacaagaagt	atagatttgg	actcgccatc	ggaaccaact	ctgtgggggtg	ggctgttatt	60
acagatgaat	ataaggtgac	atccaaaaag	tttaagttc	tgggcaatac	tgatagacac	120
tcaatcaaga	agaatctgat	agggtgcactt	ctgtttgata	gtggagagac	tgccgaggca	180
accagactta	aaaggactgc	aagaagaaga	tataccagaa	gaaagaatat	gattttgctat	240
ttgcaggaaa	tcttcagcaa	cgaaatggcc	aaggttgatg	actcattttt	ccataggttg	300
gaggagagtt	ttcttgtgga	ggaagataag	aagcacgaaa	gacacccaat	tttcggggaat	360
atagtgagac	aggtggctta	tcatgagaag	tatccacta	tctaccacct	gagaaagaaa	420
cttgtggact	caactgataa	ggctgatctt	aggcttatat	acttggccct	tgacatatg	480
atcaaatcca	ggggccattt	tcttatcgaa	ggcgatctta	atcccgataa	ctcagatgtg	540
gacaagctgt	ttatacaact	tgtgcaaac	tacaatcaac	tcttcgagga	gaatcccat	600
aacgcctccg	cgctggatgc	aaaagccata	ctgtcagcca	gactgagcaa	aagtaggaga	660
ctggagaaac	tatatagcca	actgcccgtg	gaaaagaaga	atggggtctt	cggaatctg	720
atcgctcttt	cattgggggt	gacacccaac	tttaagagta	actttgactt	ggcagaagat	780
gcaagtttgc	aggtcagtaa	agacacatat	gacgatgacc	ttgacaatct	cttggcacia	840
ataggggac	aatacgttga	ccttttcttc	gctgccaaga	acctcagcga	cgtatatact	900
ttgtccgaca	ttcttagggg	taataccgaa	attacaagg	ccccctctag	tgcaagtatg	960
atcaaaaggt	atgatgagca	tcaccaagac	cttacactgc	tgaaggctct	ggttagacag	1020
caactccctg	aaaagtataa	ggaatatatt	ttcgaccaaa	gtaagaacgg	gtacgcccgt	1080
tatatgtgat	ggggcgcaag	tcaagaagaa	ttttacaaat	tcatcaagcc	aattcttgaa	1140
aagatggacg	ggactgagga	attgctgggt	aaactgaata	gagaggacct	tcttagaaaa	1200
cagaggacat	ttgacaatgg	gtccatccca	caccagattc	atctggggga	actccacgca	1260
atattgagga	gacaagaaga	cttttaccca	ttccttaagg	ataatagaga	gaaaatcgaa	1320
aaaaatctga	ctttcaggat	tcttactact	gttggggcac	tggccagggt	gaactcaaga	1380
ttcgcttgga	tgacaaggaa	gtcagaagaa	accataaccc	cttgggaatt	tgaagagggt	1440
gttgataaag	gggcatcagc	ccagctcttt	atagagagga	tgaccaactt	tgataaaaat	1500
cttccaaatg	agaaggtttt	gccaaaacat	agtcttttgt	acgagtactt	tactgtttat	1560
aacgaattga	ccaaggtgaa	gtatgtgacc	gaggggaatg	ggaagccagc	atttttgttc	1620
ggggagcaaa	agaagcaat	cgttgatctt	ctctcaaga	ccaacagaaa	agtgaccgtg	1680
aaacaactga	aggaagacta	cttcaaaaag	atagaatgtt	tcgattcagt	ggaatttagc	1740
ggtgttgaag	acaggttcaa	tgttctattg	ggtacttacc	acgacctgtt	gaagataatc	1800
aaagacaagg	actttctcga	taatgaggag	aacgaagaca	tcttggaaga	catttgtgctt	1860
acactcactt	tggttgagga	cagggaatgt	attgaggaaa	gactcaaaac	ttacgtctcat	1920
ttgtttgatg	ataaggttat	gaaacaacta	aaaagaagaa	ggtacacccg	ctggggaaga	1980
ttgagtagga	aactgatcaa	cgttattaga	gataaacaat	ccggaaagac	tatcctcgat	2040
ttccttaaga	gtgatggctt	tgcaaatagg	aattttatgc	agctgattca	tgacgactca	2100
cttaccttca	aagaagacat	ccaaaagct	caggtgtctg	ggcaaggcga	cagtctgcat	2160
gaacatatag	ctaacttggc	tgggagtccc	gccatcaaga	aggggatact	tcaaacagtt	2220
aaagtgtgtg	acgaattggg	gaaggtaatg	ggaaggcaca	agcctgaaaa	tatagtgata	2280
gaaatggcaa	gggaaatcca	aacaaccag	aaggagacga	agaacagtag	ggaaggatg	2340
aaaaggatag	aagaggggat	caaagagctt	ggtagccaga	tctcaagga	acatccagtg	2400
gagaataccc	aacttcaaaa	cgagaaactc	tatttgtact	acttgacaga	cggaagagat	2460
ttgtatgtgg	accaagagct	tgatattaac	aggctgagcg	attatgacgt	tgaccacata	2520
gtgccccaat	cattctctcaa	ggatgactct	attgataata	aggtgctgac	aaggagtgtg	2580
aagaatagag	ggaaatccga	caacgttcca	tccgagggaag	tgtggaagaa	gatgaagaac	2640
tactggaggc	agttgtgtga	cgctaagctc	attaccacga	ggaaattcga	taacctgacc	2700
aaagcagaga	gaggcgggct	gagcgaaact	gataaagcag	gttctatcaa	gagacaaact	2760
gtggagacta	ggcaaatatt	taagcacgtg	gctcaaatac	tcgacagcag	gatgaacaca	2820

-continued

```

aagtagcagc agaacgacaa gctcattaga gaggttaagg ttattactct gaaaagtaaa 2880
ttgggttagc atttcagaaa ggatttccaa ttctataagg ttagagagat caacaattat 2940
catcatgcac atgatgccta tctgaatgct gtggttggtg cagcccttat caagaagtac 3000
cctaagctag agagcaggtt tgtgtacgga gattataagg tgtatgatgt gaggaaaatg 3060
ctcgctaaaa gtgagcaaga gattgaaaag gctacgcgca aatacttctt ttattccaat 3120
attatgaatt tcttcaagac agaataccac ctggctaacg gcgagataag gaagaggccg 3180
cttatcgaaa ctaatgggga gacaggcgaa atagtgtggg acaaaggag ggatttcgca 3240
actgtgagga aggttttgag catgcctcag gtgaatatcg ttaagaaaac cgaagttcaa 3300
actggagggt tctctaagga aagcattctc cccaagagga actccgacaa gctgattgct 3360
agaaagaaag actgggaccc caagaagtat ggccgattcg actcaccac tgtggcata 3420
agcgctctcg tgggggcaaa ggttgaaaag ggtaaatcca aaaaactcaa atccgtgaag 3480
gaactccttg gcataactat tatggaaagg agtagctttg aaaagaatcc catcgacttt 3540
ctcgaagcta agggctataa ggaagttaag aaggacctta taatcaaact tccaaaatac 3600
tccctttttg agttgaaaaa cggcagaaaag agaattgttg ccagtgcagg ggagcttcaa 3660
aagggcaacg aactggctct gccatagcaaa tatgtgaact ttttgatct ggcatcacac 3720
tacgagaaac ttaaaggctc tctgaggac aacgagcaaa aacagctctt tgtgaacag 3780
cataagcact acctcgacga gattattgag cagatcagcg agttctcaa gagagttatt 3840
ctggctgacg ctaatcttga caaggttttg tccgcttaca acaaacacag ggataagcca 3900
atcagggagc aggcagaaaa cataatccat ctctttacc tgacaaaact cggtgcccc 3960
gctgctttca agtattttga tactaccatt gacaggaaga gatatacttc cactaaggaa 4020
gtgctcgacg caaccctcat acaccaaagt atcacaggcc tctatgaaac taggtagat 4080
ttgtctcaac ttgggggcga t 4101

```

```

SEQ ID NO: 70      moltype = DNA length = 4101
FEATURE
misc_feature       1..4101
                    note = Cas9
source              1..4101
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 70
gacaaaaagt attccatcgg gcttgcctac ggaaccaact ctgtgggggtg ggcagttatt 60
accgacgaat acaagggtgcc cagcaagaag ttttaaggttc tggggaaacac agatagacat 120
agcataaaga aaaacctgat aggcgcactg ttgttcgact ccggggaacac agccgaagct 180
accaggctga agagaactgc aagaagaagg tacaccagaa gaaaaaacag aatatgttat 240
ctccaagaga tttctcttaa cgagatggcc aaggtggacg actcattctt tcacagactg 300
gaagaatctt tcttgttgga cgaagataag aaacacgaga ggcaccttat ttttggcaat 360
atcggtgatg aggtggctta ccacgaaaaa tacctacaa tataccacct caggaaaaaa 420
ttggttgata gtacagacaa ggcgcacctc aggtcctatc atttgccctt ggcccatatg 480
attaaattca gggggcactt tctcatcgag ggagatttga acccgacaa cagtgatgtt 540
gataagctct ttattcagct cgtgcagact tacaatcagt tgtttgagga aaacccctt 600
aatgcttcg ggggtggacg caaggcaatc ctttctgcaa gactctcaa gtcaaggaga 660
ctcgaataat tgatagcaaa gcttcaggga gagaagaaga acgggctctt tggaaacctg 720
atcgctctgt cactcggact cacaccaat ttcaaaagca attttgattt ggccagaggc 780
gctaagctgc aactcagtaa ggatacctac gacgatgact tggataatct gctcgacaa 840
attggggacc agtatgcaga cctgtttctc gcagctaaga acttgagtga cgccatattg 900
ctcagtgaca tctcagggtt taataccgag attacaaaag ctccactctc tgcaagcatg 960
atcaagaggt atgacagaca ccatcaagac ctgacactcc ttaaggcgtt ggtagggcag 1020
caacttcctg aaaagtctaa ggaatctctt ttcgatcaaa gcaaaaacgg ctacgcggcg 1080
tatatagacg ggggagcatc ccaagaagaa ttttataagt tcataaaacc tatattggag 1140
aagatggacg ggacagagga attgctcgtg aaactgaaca gggaggatct cctcaggaaag 1200
caaaggacct tcgacaatgg tccatccca catcagattc acctcggcga actgcagca 1260
atactgagaa gacaagagga cttttatcct ttcctgaagg acaacaggga gaaaatcgag 1320
aaaaatctga catcagaat ccatactac gttgggctc tggccagagg taacagtagg 1380
ttcgcttgga tgactaggaa actcagagg actattacac cctggaaact tgaagaagtt 1440
gttgataaag gagcttcagc acaatcattc atcgaaaaga tgacaaaact tgacaaaaat 1500
ctgcctaatg agaagtgct cccaaaacat tccctgctgt atgagtattt taccgtttat 1560
aacgagctta ccaaggtgaa atacgttact gaaggtatga gaaagccagc ttttctttca 1620
ggggagcaaa agaaggctat cgtggatctt ctctttaaga ccaacagaaa ggtagccgtg 1680
aagcagctta aggaagacta ctttaaaaag atcgagtgtt ttgactcagt ggaataaagc 1740
gggtttgaag atagattcaa cgcactcctt ggaacttctc atgatctct taagataatc 1800
aaggataaag actttctcga caacgaggaa aacgaagata tactggagga catagttctg 1860
acacttactt tgttcgagga tagggagatg atcgaggaaa gactgaaaac atatgctcac 1920
cttttcgacg acaaaagttt gaaacaactc aagagaagga gatatacagg gtgggggaga 1980
ttgagcagga aactgattaa tggatatcaga gacaaacagt caggaaaaac aatactcgac 2040
tttttgaaat cagacgggtt cgcaaatagg aatttcatgc agcttataca cgacgattca 2100
cttactttta aagaggacat tcaaaaggct caagttagtg gacaagggtg ctccctccac 2160
gaacacatcg caaatctcgc tggcagccct gcaattaaga aggtatact ccagacagtt 2220
aaggttggtg acgagctggt taaagtgatg ggaagacaca aacccgagaa catagtata 2280
gagatggcca gggaaaacca aaccactcaa aaagggcaga aaaaatccag agagaggatg 2340
aaaaggattg aagaaggtat caaggagctg ggtagccaaa ttctgaaaga acatcctgtg 2400
gaaaacactc aactocagaa tgagaaactc tatctgtact atctgcaaaa tgggagagat 2460
atgtatgttg accaggaact ggacataaac aggtctctcag attacgatgt ggatcatatc 2520
gtgccacagt cttttcttaa ggtatagatg atcgacaata aggtgcttac caggtccgac 2580
aagaacaggg gaaagtcaga taactgcctc tctgaagaag ttgttaaaaa gatgaagaa 2640
tactggagac agctgcttaa cgctaagctc ataacacaga ggaagtttga caacttgacc 2700

```


-continued

```

aaggccgaga gaggccgact ctccagaattg gataaggcag ggttcataaa aaggcagctg 2760
gtggaaacaa ggcagataac taacatgtg gctcagatcc tcgatagtag gatgaataca 2820
aaatacagatg agaacgacaa gctcataagg gaggttaaaag tgataactct gaaatccaaa 2880
ctgggttagcg attttaggaa ggatttccag ttttacaagg ttagggagat caacaattat 2940
catcacgccc acgatgccta cttgaacgca gttgtgggta ctgcacttat caaaaagtac 3000
cctaagctgg aatccgagtt tgtttatgga gactataagg tgtacgacgt tagaaaaatg 3060
attgcaaagt cagagcagga gatagggaaa gccactgcaa aatatttctt ttatagcaat 3120
atcatgaatt tctttaagac agaaatcaca ctggccaatg gggaaataag gaagaggccc 3180
ctgatcgaaa ctaatggcga gacaggggag attgtgtggg ataaaggtag ggactttgca 3240
acagtggagg aagtgtcgag catgccccaa gttaatatcg ttaaaaagac cgagggtcaa 3300
acagggggct ttagtaagga aagcattttt cccaagagga atagtacaa attgattgct 3360
aggaaaaaag attgggaccc caaaaagtat ggcggatttg atagcccccac tgttgcttac 3420
tccgtgctcg tgggtgcaaa ggtgggagaag ggaagagca agaaactgaa gtcagttaag 3480
gaactccttg gtatcactat catggaaaaga agctcctttg agaagaaccc tattgacttc 3540
ctggaggcta aagggtacaa agagggttaag aaagacctta tcattaaatt gcccaaatat 3600
agtcttttcg agcttgaaaa cggaagaaaag aggatgcttg catccgctgg cgaattgcaa 3660
aagggcaatg agcttgctct cctttccaaag tatgtgaact tcttttatct tgctcacac 3720
tatgaaaaac tcaaggttcc acccggaagac aacgaacaaa agcaactatt tgtggaacaa 3780
cacaagcact acctggacga aatcattgag caaatctctg agttttcaaa aagggttaac 3840
ttggctgacg caaatctcga caaagttttg tcagcttaca acaaacatag agataagcca 3900
attagagagc aagctgagaa tatcatccat ctgtttaccc tgactaacct tggagcgct 3960
gctgctttta aatatttcga caccacaatc gacaggaaga ggtacactag cactaaggaa 4020
gttctcgacg caacctcat ccaccagagt attacaggcc tgtacgagac aagaattgat 4080
ctttctcaac ttggtggtga c 4101

```

```

SEQ ID NO: 71      moltype = DNA length = 4101
FEATURE            Location/Qualifiers
misc_feature        1..4101
                    note = Cas9
source              1..4101
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 71
gataagaagt actcaatcgg tctggcaatc ggaaccaact ctgtgggttg ggcagtgatt 60
acagatgagt ataaggtgcc aagcaaaaaa ttcaaggtgc tgggtaatac cgacagacac 120
agcattaaga agaatttgat tggagcactc ctctttgact caggggaaac agcagaggca 180
acaaggctga agaggacagc aaggcggagg tacacaaggc ggaaaaaacag gatatgctac 240
ctccagaaaa tctttagcaa cgagatggct aaagtggatg atagcttttt ccatagactc 300
gaagaatcct ttcttgttga agaggacaaa aagcatgaaa ggcataccat ctccggcaat 360
atagttgatg aggttgcata ccatgagaag taccaccaaa tctaccacct cagaagaaaa 420
cttgtggact ccacagataa agcagacctg aggtcctat acctgcactc cgcacacatg 480
atcaagttca gagggcactt tctcatcgaa ggtgacctga atccagataa ttcatagtg 540
gataaactgt ttatcacagc ggtgcaaaac tacaaccaac ttttcgagga aaaccaatc 600
aatgcctcgc gtgttgatgc aaaggccatc ctgtcagcaa gactcagcaa aagcaggcgg 660
ctcgaaaaac tcctcgcccc gcttcccggt gaaaagaaga acgggctctt tggtaatctc 720
atcgatttga gccttggtct tactccaac ttcaagagca attttgatct ggcagagatt 780
gctaaactgc aactctcaaa ggacacatat gacgatgacc ttgacaatct gttggcccag 840
atcggggacc aatatgcaga cctcttctcg gccgcaaaag atctgtcaga tgcaatcctc 900
ttgtccgaca tacttgagat taacactgag atcacaaggc cacctctgtc cgctccatg 960
attaagagat acgatgagca tcaccaggat ctgactttgc tcaaaagcct cgttagacag 1020
cagttgcccag aaaagtacaa agaaatattc tttgatcaat caaaaaacgg atatgcaggg 1080
tacatcgacg gtggggcaag ccaggaaagag ttctacaaat tcatacaaac tatcctggaa 1140
aagatggatg ggacagaaga gctgctgggt aagctgaata ggggaagacct cctcagaaaag 1200
cagaggacat ttgataacgg gagcatccct catcaaatcc acctcggtga actccatgct 1260
atcctgagaa ggcaggaaga cttttatcca tttttgagg acaataggga gaaaatcgaa 1320
aaaaacctga catcagaat cccatactac gttggtctct tggcaagagg taacagtagg 1380
ttcgcatgga tgacaaggaa aagcgaggag acaatcacac cctggaattt tgaggaaagt 1440
gttgacaagg gtgccagcgc acaatccttt atcgaaagaa tgacaatttt cgacaagaat 1500
ctgcctaacg aaaaggttct cccaaagcat tcaactcctg acgaatattt tacagtttat 1560
aacgaactga ctaaagttaa atacgttacc gagggataga ggaagccagc attcctttcc 1620
ggggaacaga agaaagctat tgtggacctc ctgttcaaga caaatagaaa agtgacagtt 1680
aagcaactca aagaggatta cttcaaaaag atcgaatgtt ttgactctgt ggagatcagc 1740
ggggtggagg atagattcaa cgcagcctg ggtacatata atgactcctt gaaaatcatt 1800
aaagacaagg acttcttgga caacgaggag aacgaggaca ttctggaaga cattgttctg 1860
accctcacac tctttgagga tagggagatg attgagaaa gactgaagac ctacgcccac 1920
ctctttgacg ataaagtgat gaacagctc aagagaagaa ggtatacagg ttgggggaga 1980
ctgagcagga agttgatcaa tgggattagg gacaacaagt ccgggaaaaa aatcctcgat 2040
ttctgaagt cagacggttt cgcaaacaga aattttatgc agtcoattca cgatgacagc 2100
ttgacattca aggaagacat ccaaaaggct caagtgagcg gccaaaggga tagcctccac 2160
gagcatattg caaatctggc aggttcacca gccatcaaaa agggcatact tcagacagtt 2220
aagggttggtg cgaatttggt taaagttatg ggcaggcata agcagagaa tatcgttatc 2280
gaaatggcaa gggagaacca aacaactcaa aaagggcaga aaaaatagcag agagaggatg 2340
aaaaaatcg aggaagggat caaggaaactt ggggtccaaa tctcagaaga gcacccagtt 2400
gaaaatactc aactgcaaaa cgagaagctc tatctctact atctccaaaa cgggagggat 2460
atgtatgttg accaggagct ggatattaac agactgtcag attatgatgt tgatcatatc 2520
gtgccccagt catctctgaa ggacgattcc atcgacaaca aagttctcac aaggtccgat 2580

```

-continued

aaaaacaggg	gcaagtccga	taacgttcca	agcgaagaag	tgggtgaaaa	gatgaaaaac	2640
tattggagac	aacttctgaa	tgcaaagtgt	attactcaga	gaaagtgttg	caacctcaca	2700
aaagcagaaa	gaggcgggct	tagcgaactc	gataaggcag	ggtttatcaa	aagacagctg	2760
gttgagacaa	ggcagatcac	aaaacatgtg	gcacagatcc	ttgactcaag	gatgaatacc	2820
aagtatgatg	agaatgataa	gttgatcagg	gagggttaag	ttatcacact	caaatccaaa	2880
ctgggtgtcag	acttcaggaa	agactttcaa	ttttataaag	tgagggagat	caataactac	2940
caccatgcac	atgacgccta	cctgaacgca	gtggtgggta	cagcattgat	taaaaaatac	3000
cctaagctgg	agtctgagtt	tgtgtacggg	gactacaagg	tgtacgacgt	gaggaaaatg	3060
atagccaagt	ccgagcagga	gatcgggaaa	gcaacagcta	agtatctctt	ttacagtaat	3120
atcatgaatt	tctttaaacc	tgagattact	ctggcaaacg	gggagatcag	gaaaagaccc	3180
ctcatcgaga	ctaattggtga	aacaggtgag	atcgtttggg	acaaggggag	ggatttttgt	3240
actgttagaa	aagttctgag	tatgccacaa	gtgaatatgt	tgaaaaagac	agaagttcag	3300
acaggtgggt	tctccaaga	atccatcctg	ccaagagaa	attcagacaa	gctcatcgca	3360
agaaagaagg	actgggaccc	taagaagtac	ggaggatttg	acagccccac	cgtggcctat	3420
tcctgtcttg	ttgtggcaaa	ggtggagaaa	gggaagagca	aaaaactgaa	atccgtgaaa	3480
gaactgctgg	gaattaccat	ctggaaga	agctcctttg	agaagaaccc	aatcgacttc	3540
ctggaagcaa	aaggatataa	ggaagtga	aaggacctca	ttatcaagct	cccaaaaatac	3600
tcacttttctg	agtgtggagaa	cggtagaaag	aggatgctgg	caagcgcagg	ggaacttcag	3660
aaaggcaatg	agctggcatt	gccatcaaa	tatgtgaact	tcctctactt	ggccagccat	3720
tacgagaaac	ttaaaggtag	cccagaagat	aacgagcaaa	aacagctctt	tgtggaaacag	3780
cataagcatt	atctggatga	gatcatagaa	caaatctcag	agttttccaa	gagagtatatc	3840
ctcgcagatg	caaacttgga	taaggttctc	tcagcctata	ataagcatag	agacaagcca	3900
attagagagc	aagcagagaa	cattatccac	ttgttccact	ttacaaaacct	ggggggcacca	3960
gccgccttca	aatatctcga	tacaacaata	gacagaaaga	ggtataccag	caccaagaaa	4020
gttctcgacg	ccacactgat	ccatcaatca	atccacagcc	tttacgaaac	taggatcgac	4080
ttgtcacaaac	tgggtgggga	t				4101

SEQ ID NO: 72 moltype = DNA length = 3307
 FEATURE Location/Qualifiers
 misc_feature 1..3307
 note = Cas9
 source 1..3307
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 72

gagcaaggac	acctacgacg	acgacttgga	caacctattg	gcccagatag	gtgaccagta	60
tgacagacctc	ttccttgccg	ccaagaactt	gagtgacgct	atactgctca	gtgacatcct	120
gaggggtgaac	actgagatca	ctaaggcccc	tctctctgcc	tcaatgatta	agcgttacga	180
cgagcatcac	caggatctca	cctctgctta	ggcccttgtt	cggcagcagc	tcctcgagaa	240
gtacaaggag	atattttttg	accagtctaa	gaacgggtac	gccgggtaca	ttgacgggtg	300
ggcaagccag	gaggagtctt	acaagttcat	caagccgac	cttgagaaga	tggacggcac	360
cgaggagcta	cttgtcaagt	tgaaccggga	agacctgtct	cggaaacacg	gtacattcga	420
caacggcgac	atccctcaac	agatccacct	ggcggaacta	cacgccatcc	tcgcagctca	480
ggaggacttc	tatccattct	tgaagataa	cagggaaaaa	atcgaaaaaa	tacttacgtt	540
tgaataacct	tactacgtgg	ggccccctgc	tcgggggaaa	tcagatctcg	catggatgac	600
caggaaagtc	gaggagacca	tcacaccctg	gaactttgag	gaggtgggtg	acaaagggtg	660
ttctgccccag	tccctcatgt	agcggatgac	taacttcgac	aagaacctgc	ccaacgagaa	720
ggtgctgcc	aagcacagcc	tgtcttacga	atactttact	gtgtacaatg	agctgacgaa	780
ggtgaagtac	gtgacagagg	ggatgcggaa	gcccgcttct	ctgagcggcg	agcaaaaaaa	840
agcaatcgtg	gacctactgt	tcaagaccaa	ccgaaagggtg	acagtgaagc	agctcaagg	900
ggactacttc	aaaaaaatcg	agtgtctcga	ctctgttgag	ataagcggcg	tggaggaccg	960
attcaacgct	tcattgggaa	cctatcacga	cctgctcaag	atcattaaag	acaaggactt	1020
cctggataat	gaggagaatg	aggacatcct	ggaggatatt	gtgctgaccc	ttactctatt	1080
cgaggacagg	gagatgatcg	aggagcgact	caagacctac	gctcacctgt	tcgacgacaa	1140
ggttatgaag	caattgaagc	gtaggcgata	cacgggggtg	ggaagactct	cccgaaaact	1200
gataaacggc	atcagggaca	agcagtcagg	gaagacgac	ttggacttcc	tgaatccga	1260
cgggttcgcc	aaccgcaact	tcattgcagc	cattcacgac	gactcactaa	cgttcaaaga	1320
ggacattcag	aaggctcaag	tcagtggaca	aggcgactcc	ctgcacgagc	acattgcaaa	1380
ccttgccgggc	tccccggcga	ttaaaaagg	cattctccaa	acgggttaagg	tgggtggacga	1440
gctggtgaag	gtgatggggc	gacacaagcc	tgagaacatc	gtgatcgaga	tggccaggga	1500
gaaccagact	accagaagg	gtcagaagaa	ctctcgggaa	cgtatgaagc	gtattgagga	1560
ggggattaa	gagttgggct	ctcaaatcct	caaggagcac	cctgtggaga	acactcagct	1620
ccaaaacgag	aagctgtacc	tgtactacct	gcaaaacggg	cgcgatatgt	acgtggatca	1680
ggagttggac	atcaacaggg	ttagcgatta	cgacgtggac	cacatcgtgc	cacagtcatt	1740
cttaaaaggac	gacagatcgc	acaacaaggt	ttgacgagg	agcgacaaga	atcgagggaa	1800
aagtgaacaat	gttccatccg	aggaggtggt	caagaaaatg	aagaactatt	ggcgtcagct	1860
tctgaacgcc	aagctcatca	cccagcggaa	attcgacaac	ctgactaagg	ctgagcaggg	1920
cggactctcc	gagcttgaca	aggctggcct	catcaagcgg	cagttggctg	aaacccgaca	1980
gataacgaag	cacgttgccc	agatacttga	ctcccgtatg	aacaccaagt	acgacgagaa	2040
gcacaagctc	atcaggagg	tgaaggtcat	tacccttaag	tccaaactcg	tcagcgactt	2100
tcgtaaggac	ttccagttct	acaagggtgc	cgagatcaat	aactaccacc	acgcacacga	2160
cgcctacctg	aacgcagtg	ttggaacccg	gttgattaaa	aagtacccca	agttggagtc	2220
ggagttcgtt	tacggggact	acaaggtgta	cgacgttcgg	aagatgatcg	ccaagtctga	2280
acaggagatc	gggaaagcaa	ccgccaagta	tttctcttat	agcaacatca	tgaacttctt	2340
taaaaccgag	atcacacttg	ccaatggcga	gatccgtga	aggccgctga	tcgagacaaa	2400
tggggagact	ggcgagatcg	tgtgggacaa	gggcccgcgac	ttcgcaaccc	ttcgaaagt	2460

-continued

cttgtccatg	cctcaagtca	acatcgtcaa	gaagactgag	gtgcaaacag	gcgggttctc	2520
gaaggagtc	atactgcccc	agaggaaactc	agacaagctc	atagcacgca	aaaaagactg	2580
ggatccaaag	aaatacggcg	gggttcgactc	gccgacagtc	gcataactccg	tgttagtggt	2640
ggctaaagt	gaaaagggga	agtccaagaa	gctcaagtc	gtcaaggagt	tgctcgggat	2700
caccattatg	gaacgggtcc	cattcgagaa	gaatccatt	gacttcctag	aggcgaagg	2760
ctacaaagag	gtcaaaaagg	acctaattat	taagctcccc	aagtattcac	tcttcgaact	2820
tgaatgggt	cgtaagcgga	tgttggaag	cgctggagag	cttcagaagg	ggaacgagct	2880
tgactgcct	tccaagtacg	tgaacttcct	gtacctcgcc	tctcattacg	agaagttgaa	2940
gggtccaccg	gaggacaacg	agcagaagca	gttgttcgtg	gagcagcaca	agcactacct	3000
cgacgagatc	attgagcaga	taagtgaagt	cagcaaacgg	gtgatccttg	ccgacgctaa	3060
cctggacaag	gtgctgagcg	cctacaacaa	gcacagagac	aagccgatcc	gagagcaagc	3120
ggagaacatc	atacacctgt	tcaccctcac	gaacctcggg	gctcccgag	ccttcaaata	3180
ttttgacacg	accatcgacc	gtaaacgcta	cactagcacg	aaggaggtgc	tggacgctac	3240
ccttatccac	cagtcocatca	ccggcctgta	cgagacgaga	atcgacttgt	cgcagctcgg	3300
tggtgac						3307

SEQ ID NO: 73 moltype = DNA length = 4101
 FEATURE Location/Qualifiers
 misc_feature 1..4101
 note = Cas9
 source 1..4101
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 73

gacaaaaaat	actcaattgg	tctggcaatt	gggaccaaca	gtgtcggatg	ggccgtgatt	60
accgacgagt	acaaggtgcc	gtccaaaaaa	ttcaaggtgc	ttgggaacac	cgaccgccac	120
tcgatcaaga	aaaacctaat	cgggtcggtt	cttttcgaca	gtggggagac	cgccgaggca	180
acacgcttaa	aacgcacagc	taggaggaga	tatacacggc	gcaagaaccg	aatatgctac	240
ttacaggaga	tattctccaa	tgagatggcg	aaggtggacg	actctttctt	ccatcggtt	300
gaggaatcct	tcctgggtcg	ggaggacaag	aagcagcagc	gacacccgat	attcgggaac	360
atcggtgatg	agggtggcgt	caacgagaag	taccaacga	tataccactt	acgcaagaag	420
ctcgtggact	ctacggacaa	ggccgacttg	cgccttatct	acttggcact	ggcccacatg	480
attaagttcc	gaggccactt	ccttatcgag	ggtgacctga	accccgataa	ctccgacgtg	540
gacaagctct	tcattccaact	gtccagaca	tacaaccagc	tattcgagga	gaatcctatc	600
aacgcctctg	gggtggacgc	taaagctatc	ctctcagccc	gcctgtcaaa	gtcgaggagg	660
ttggagaacc	taatcgcccc	gcttcagggc	gagaagaaaa	atgggctgtt	cggaaacctt	720
atcgcaactc	cactgggcct	aaccccgaac	ttcaagtcca	acttcgacct	ggcagaggac	780
gcgaatattg	agttgtcgaa	agacacctat	gacgatgacc	tggaacaacct	gttggcccag	840
ataggggacc	agtacgcgga	cctgttccct	gcggccaaga	acctgtccga	cgccatcttg	900
ctgtcggata	tactgctggg	gaacaccgag	atcactaaag	cacctctctc	cgccagcatg	960
attaaagcgtt	acgacgagca	ccaccaagat	ttgacctgc	taaaggcact	tgtacggcag	1020
cagcttcctcg	agaagtacaa	ggagatcttt	ttcgacaaa	gcaagaaccg	ctacgcccgg	1080
tacatcgacg	gaggtgccc	ccaggaggag	ttctacaagt	tcattaagcc	catccttag	1140
aagatggacg	ggactgagga	actacttggt	aagctgaacc	gggaagactt	actacgggaag	1200
cagcgtaact	tcgacaacgg	ttctatccca	catcagatcc	atcttgggga	gttgacgcg	1260
atcctgagac	gcccagggga	cttttaccct	ttcctgaaa	acaaccgga	gaaaatcgag	1320
aagatactga	ccttcagaat	accttactac	gtcggacccc	ttgcgcgagg	caactcaaga	1380
ttcgcgtgga	tgaccaggaa	atcagaggag	accatcacac	cctggaattt	cgaggagggtg	1440
gttgacaagg	gtgcctccgc	ccagtccttt	atcgaaacga	tgaccaactt	cgacaagaac	1500
ttgcccacag	agaaggtgct	ccccaaacac	agcctcctct	acgaatatct	cacagtgtac	1560
aacgagctta	ctaaagttaa	gtatgttact	gagggcatga	ggaaaccgcg	cttctgtgca	1620
ggcagagcga	gaaaagctat	tgtggacctc	cttttcaaga	ccaaccggaa	ggtagacagt	1680
aagcagctca	aggaggacta	cttcaagaag	atagagtgtc	tcgacagcgt	ggagatcagc	1740
ggggtggagg	acagattcaa	tgctctcttc	ggaacatacc	acgacttgct	taagatcatc	1800
aaggacaagg	acttctctga	caacgaggaa	aacgaggata	ttctggagga	tattgttctg	1860
actcttacc	tgttcgagga	ccgggagatg	atcgaggagc	gtctcaagac	ctacgcccac	1920
ctgttcgacg	acaaagttaa	gaagcagctc	aagcgtcgga	gatataccgg	atggggccgt	1980
ctgtctcgga	agctcatcaa	cgggatcagg	gacaagcagt	cagggaagac	gatcttagac	2040
ttccttaagt	ctgacggctt	cgccaacagg	aacttcatgc	agttgatcca	cgacgacagc	2100
cttaccttca	aggaggacat	ccagaaggcc	caagtgaagt	gccagggtga	cagcctccac	2160
gagcatattg	ctaactctgc	gggttcccc	gcgattaaaa	aggcataact	tcaaacctgt	2220
aaggtggtgg	acgagcttgt	caaggtgatg	gggcgacaca	agcccagaga	catcgtgatc	2280
gagatggcca	gggagaacca	gaccacccag	aaggggcaga	agaatagccg	agaacgcatg	2340
aagcgcacat	aggaggggat	taaggagcta	gggagccaga	tcctcaagga	acatcccgtc	2400
gagaacacc	agctccagaa	cgagaagcta	tacctctact	acttgcaaaa	cgggagggat	2460
atgtacgtgg	atcaggagtt	ggacattaac	cgccaaagcg	actacgacgt	agatcacatc	2520
gtgcctcagt	cattcctcaa	agacgacagc	attgacaaca	aagtcctgac	ccgatccgac	2580
aagaaccgag	gaaaatccga	caatgtgccc	tcagaggagg	tcgtcaagaa	aatgaagaac	2640
tattggaggc	agctacttaa	cgccaaactc	ataaccacgc	ggaagttcga	caacctgaca	2700
aaggctgagc	ggggtggggt	cagcgagctt	gacaaggctg	gcttcaccaa	gcggcagttg	2760
gtggagacaa	gacagataac	gaagcagctg	gctcagatcc	tggactctcg	catgaacacg	2820
aagtacgacg	agaacgacaa	attgatcccg	gaggtcaagg	ttattacgct	caagagcaaaa	2880
cttgtcagcg	atttcoccaa	ggacttcacg	ttctacaagg	tgaggggagt	taacaactac	2940
caccatgcac	attgatgcta	cttgaacgca	gtggtgggga	ccgcgcttat	taaaaagtac	3000
cctaagttgg	agtcagagtt	cgtttatggg	gactacaagg	tgtacgacgt	ccggaagattg	3060
attgcaaaagt	ctgaacagga	aatcggaag	gccaccgcca	aatatttctt	ctacagttaac	3120

-continued

```

attatgaatt tttttaagac tgaattact ctgcgaaacg gcgagatcag gaagcgctcc 3180
ctcatcgaga caaacgggga gaccggggag atagtctggg acaaggggag ggacttcgct 3240
acgggtgagga aggtgctctc gatgccacaa gtgaacatcg tcaaaaagac agagggtcag 3300
accggtggct tctcaaagga gtcaatcctg ccaaaacgta acagcgacaa gctcatcgcc 3360
cgcaagaaaag actgggaccc taagaagatg ggtgggttcg actcaccgac ggtcgcatatc 3420
tcctgtcttg tctgggcaaa ggtggaaaag ggcaagtcca aaaaactgaa atccgtgaag 3480
gagttgcttg gcattaccat catggaacgc agcagcttcg agaagaaccc cattgacttc 3540
ctggaggcta aagggtacaa ggagggtcaa aaagatttaa ttattaagct acctaatgac 3600
agcttgcttg agctggagaa cggccgaaaa cgaatgctcg catccgcccgg ggaacttcaa 3660
aagggcaacg agcttgcgct gccctccaag tacgtgaact tcctgtactt ggcatccac 3720
tacgagaaac tcaagggtag cccagaggac aacgagcaga agcagctatt cgtggagcag 3780
cacaagcact acctcgacga gataatcgag cagatcagtg agttcagtaa gcgggtgata 3840
ctcgcggacg ccaacttgga caaggtgctt agtgcctaca acaagcaccg tgacaagccc 3900
atccgagaaac aggtgtgagaa catcatccac cttttcactc tgacaaaacct cggtgctccc 3960
gccgccttca aatacttcga cactaccatc gacaggaagc gctacacatc tacgaagaaa 4020
gttcttgacg ctacgcttat tcatcagctt atcacagggc tgtacgagac aaggatcgac 4080
cttagccaaac tcgcggggga t 4101

```

```

SEQ ID NO: 74      moltype = AA length = 1367
FEATURE           Location/Qualifiers
source            1..1367
                  mol_type = protein
                  organism = synthetic construct

```

```

SEQUENCE: 74
DKKYSIGLAI GTNSVGWAVI TDEYKVPSKK FKVLGNTDRH SIKKNLIGAL LFDSGETAEA 60
TRLKRTARRR YTRRKNRICY LQEIFSHEMA KVDDSFHRL EESFLVEEDK KHERHPFIGN 120
IVDEVAYHEK YPTIYHLRKK LVDSTDKADL RLIYLALAHM IKFRGHFLIE GDLNPDNSDV 180
DKLFIQLVQT YNQLFEENPI NASGVDAAAI LSARLSKSR LLENLIAQLPG EKKNGLFIGNL 240
IALSLGLTPN FKSDFDLAED AKLQLSKDLY DDDLDNLLAQ IGDQYADLFL AAKNLSAIL 300
LSDILRVNTE ITKAPLSASM IKRYDEHHQD LTLKALVRQ QLPEKYKEIF FDQSKNGYAG 360
YIDGGASQEE FYKFIKPILE KMDGTEELLV KLNREDLLRK QRTFDNGSIP HQIHLGELHA 420
ILRRQEDFYP FLKDNREKIE KILTFRIPTY VGPLARGNSR FAWMTRKSEE TITPWNFEEV 480
VDKGASQSF IERMTNFDKN LPNEKVLPHK SLLYEYFTVY NELTKVKYVT EGMKRPAPLS 540
GEQKKAIVDL LFKTNRKVTY KQLKEDYFKK IECFDSVEIS GVEDRFNASL GTYHDLKII 600
KDKDFLDNEE NEDILEDIVL TLTFLFEDRE IERLKYTAH LFDDKVMKQL KRRRYTGWGR 660
LSRKLINGIR DKQSGKTILD FLKSDGFANR NFMQLIHDDS LTFKEDIQKA QVSGQGDLSH 720
EHIANLAGSP AIKKGILQTV KVVDELVKVM GRHKPENIVI EMARENQTTQ KGQKNSRERM 780
KRIEIEGKEL GSQILKEHPV ENTQLQNEKL YLYYLQNGRD MYVDQELDIN RLSYDVIDHI 840
VPQSFLKDDI IDNKVLRSD KNRGKSDNVP SEEVVKMKMN YWRQLLNAKL ITQRKFDNLT 900
KAERGGLSEL DKAGFIKRLQ VETRQITKHV AQILDSRMNT KYDENDKLIR EVKVI TLKSK 960
LVSDPRKDFQ FYKVEINNY HHAHDAYLNA VGTALIKKY PKLESEFVYG DYKVYDVRKM 1020
IAKSEIEGK ATAKYFFYSN IMNFFKTEIT LANGEIRKP LIETNGETGE IVWDKGRDFA 1080
TVRKVLSMPQ VNIKKTEVQ TGGFSKESIL PKRNSDKLIA RKDWDPKKY GGFDSPTVAY 1140
SVLVVAKVEK GSKKLKSVK ELLGITIMER SSFEKNPIDF LEAKGYKEVK KDIIKLPKY 1200
SLFELENGRK RMLASAGELQ KGNELALPSK YVNFYLLASH YEKLKGSPE NEQQLFVEQ 1260
HKHYLDEIIE QISEFSKRV LADANLDKVL SAYNKHDKP IREQAENIIH LFTLTNLGAP 1320
AAFKYFDTTI DRKRYTSTKE VLDATLIHQ S ITGLYETRID LSQLGGD 1367

```

```

SEQ ID NO: 75      moltype = AA length = 1367
FEATURE           Location/Qualifiers
source            1..1367
                  mol_type = protein
                  organism = synthetic construct

```

```

SEQUENCE: 75
DKKYSIGLAI GTNSVGWAVI TDEYKVPSKK FKVLGNTDRH SIKKNLIGAL LFDSGETAEA 60
TRLKRTARRR YTRRKNRICY LQEIFSHEMA KVDDSFHRL EESFLVEEDK KHERHPFIGN 120
IVDEVAYHEK YPTIYHLRKK LVDSTDKADL RLIYLALAHM IKFRGHFLIE GDLNPDNSDV 180
DKLFIQLVQT YNQLFEENPI NASGVDAAAI LSARLSKSR LLENLIAQLPG EKKNGLFIGNL 240
IALSLGLTPN FKSDFDLAED AKLQLSKDLY DDDLDNLLAQ IGDQYADLFL AAKNLSAIL 300
LSDILRVNTE ITKAPLSASM IKRYDEHHQD LTLKALVRQ QLPEKYKEIF FDQSKNGYAG 360
YIDGGASQEE FYKFIKPILE KMDGTEELLV KLNREDLLRK QRTFDNGSIP HQIHLGELHA 420
ILRRQEDFYP FLKDNREKIE KILTFRIPTY VGPLARGNSR FAWMTRKSEE TITPWNFEEV 480
VDKGASQSF IERMTNFDKN LPNEKVLPHK SLLYEYFTVY NELTKVKYVT EGMKRPAPLS 540
GEQKKAIVDL LFKTNRKVTY KQLKEDYFKK IECFDSVEIS GVEDRFNASL GTYHDLKII 600
KDKDFLDNEE NEDILEDIVL TLTFLFEDRE IERLKYTAH LFDDKVMKQL KRRRYTGWGR 660
LSRKLINGIR DKQSGKTILD FLKSDGFANR NFMQLIHDDS LTFKEDIQKA QVSGQGDLSH 720
EHIANLAGSP AIKKGILQTV KVVDELVKVM GRHKPENIVI EMARENQTTQ KGQKNSRERM 780
KRIEIEGKEL GSQILKEHPV ENTQLQNEKL YLYYLQNGRD MYVDQELDIN RLSYDVIDHI 840
VPQSFLADDS IDNKVLRSD KNRGKSDNVP SEEVVKMKMN YWRQLLNAKL ITQRKFDNLT 900
KAERGGLSEL DKAGFIKRLQ VETRQITKHV AQILDSRMNT KYDENDKLIR EVKVI TLKSK 960
LVSDPRKDFQ FYKVEINNY HHAHDAYLNA VGTALIKKY PALESEFVYG DYKVYDVRKM 1020
IAKSEIEGK ATAKYFFYSN IMNFFKTEIT LANGEIRKP LIETNGETGE IVWDKGRDFA 1080
TVRKVLSMPQ VNIKKTEVQ TGGFSKESIL PKRNSDKLIA RKDWDPKKY GGFDSPTVAY 1140
SVLVVAKVEK GSKKLKSVK ELLGITIMER SSFEKNPIDF LEAKGYKEVK KDIIKLPKY 1200
SLFELENGRK RMLASAGELQ KGNELALPSK YVNFYLLASH YEKLKGSPE NEQQLFVEQ 1260
HKHYLDEIIE QISEFSKRV LADANLDKVL SAYNKHDKP IREQAENIIH LFTLTNLGAP 1320

```

-continued

AAFKYFDTTI DRKRYTSTKE VLDATLIHQ5 ITGLYETRID LSQLGGD 1367

SEQ ID NO: 76 moltype = AA length = 228
 FEATURE Location/Qualifiers
 source 1..228
 mol_type = protein
 organism = Rattus norvegicus

SEQUENCE: 76
 SSETGPVAVD PTLRRRIEPH EFEVFFDPRE LRKETCLLYE INWGGRHSIW RHTSQNTNKH 60
 VEVNFIKFT TERYFCPNTR CSITWFLSWS PCGECSRAIT EFLSRYPHVT LFIYIARLYH 120
 HADPRNRQGL RDLISSGVTI QIMTEQESGY CWRNFFVNYSP SNEAHWPYPY HLWVRLYVLE 180
 LYCIILGLLP CLNILRRKQP QLTFFTIALQ SCHYQLRPPH ILWATGLK 228

SEQ ID NO: 77 moltype = AA length = 199
 FEATURE Location/Qualifiers
 source 1..199
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 77
 MEASPASGPR HLMDFPHITS NFNNIGIRHK TYLCYEVERL DNGTSVKMDQ HRGFLHNQAK 60
 NLLCGFYGRH AELRFLDLVP SLQLDPAQIY RVTWFISWSP CFSWGCAGEV RAFLQENTHV 120
 RLRIFAARIY DYDPLYKEAL QMLRDAGAQV SIMTYDEFKH CWDTFVDHQG CPFQPWDGLD 180
 EHSQALSGRL RAILQNQGN 199

SEQ ID NO: 78 moltype = DNA length = 621
 FEATURE Location/Qualifiers
 source 1..621
 mol_type = genomic DNA
 organism = Petromyzon marinus

SEQUENCE: 78
 acagatgcag agtatgtgag aattcacgaa aagctggaca tctatacctt caagaagcag 60
 ttctttaaca ataagaagtc tgtgagccat aggtgctacg tgctgttcga gctgaagaga 120
 aggggtgaaa gaagggcatg tttttggggg tatgctgtga acaagcccca gtctggaact 180
 gagagaggca ttcacgccga aattttcagc atcagaaaagg tggaggaata cctgagggat 240
 aacctggac agtttacaat taattggtat tctagctggt ctcctatgcg tgactgtgcc 300
 gagaagatcc tggaaatgta caaccaggag ctgagaggaa atggccatac cctgaagatt 360
 tgggcctgca agctgtacta tgaaaagaac gcaagaaatc agatcggact gtggaacctg 420
 agggataatg gtgtggggct gaacgtgatg gtgtccgagc actatcagtg ctgtagaaag 480
 attttcattc agtcctcaca taatcagctg aacgagaata gatggctgga aaagactctg 540
 aagagggctg agaagagaag gtccgaactg tcaattatga tccaggtgaa gatcctgcac 600
 accactaagt cacctgccgt g 621

SEQ ID NO: 79 moltype = AA length = 160
 FEATURE Location/Qualifiers
 source 1..160
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 79
 FERNYDPREL RKETYLLYEI KWGKSGKLWR HWCQNNRTQH AEVYFLENIF NARRFNPSTH 60
 CSITWFLSWS PCAECSQKIV DFLKEHPNVL EIYVARLYYH EDERNRQGLR DLVNSGVTIR 120
 IMDLPDYNIC WKTFFVSDQG DEDYWPQHFA PWIKQYSLKL 160

SEQ ID NO: 80 moltype = AA length = 229
 FEATURE Location/Qualifiers
 source 1..229
 mol_type = protein
 organism = Rattus norvegicus

SEQUENCE: 80
 MSSETGPVAV DPTLRRRIEP HEFEVFFDPR ELRKETCLLY EINWGGRHSI WRHTSQNTNK 60
 HVEVNFIEKF TTERYFCPNT RCSITWFLSW SPCGECSRAI TEFLSRYPHV TLFYIARLY 120
 HHADPRNRQG LRDLISSGVT IQIMTEQESG YCWRNFFVNY PSNEAHWPY PHLWVRLYVL 180
 ELYCIILGLP PCLNILRRKQ PQLTFFTIAL QSCHYQLRPP HILWATGLK 229

SEQ ID NO: 81 moltype = AA length = 198
 FEATURE Location/Qualifiers
 source 1..198
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 81
 MDLLMNRRK FLYQFKNVRW AKGRRETYLC YVVKRDSAT SFSLDFGYLR NKNCHVELL 60
 FLRYISDWDL DPGRCYRVTW FTSWSPCYDC ARHVADFLRG NPNLSLRIFT ARLYFCEDRK 120
 AEPEGLRRLH RAGVQIAIMT FKDYFYCWNT FVENHERTFK AWEGLHENS VRLSRQLRRIL 180
 LPLYEVDDL R DAFRTLGL 198

SEQ ID NO: 82 moltype = AA length = 197
 FEATURE Location/Qualifiers

-continued

source 1..197
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 82
 MDSLLMNRRE FLYQFKNVRW AKGRRETYLC YVVKRRDSAT SFSLDFGYLR NKNCHVELL 60
 FLRYISDWDL DPGRCYRVTW FLSWSPCYDC ARHVADFLRG NPNLSLRIFT ARLYPCEDRK 120
 AEPEGLRLRH RAGVQIAIMT FKDYFYCWNT FVENHGRTPFK AWEGLHENS V RLSRQLRRIL 180
 LPLYEVDDL R DAFRTCT 197

SEQ ID NO: 83 moltype = AA length = 207
 FEATURE Location/Qualifiers
 source 1..207
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 83
 TDAEVVRIHE KLDIYTFKKQ FSNKKSVSH RCVLFLFKR RGERRACFWG YAVNKPQSGT 60
 ERGIHAEIFS IRKVEEYLRD NPGQFTINWY SSWSPCADCA EKILEWYNQE LRGNGHTLKI 120
 WVCKLYYEKN ARNQIGLWNL RDNGVGLNVM VSEHYQCCRK IFIQSSHNQL NENRWLEKTL 180
 KRAEKRRSEL SIMFQVKILH TTKSPAV 207

SEQ ID NO: 84 moltype = AA length = 228
 FEATURE Location/Qualifiers
 source 1..228
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 84
 SSKTGPVAVD PTLRRRIEPH EFVFFDPRE LRKETCLLYE INWGGRHISW RHTSQNTNKH 60
 VEVNPIEKFT TERYFCPNTR CSITWFLSWS PCGECRAIT EFLSRYPNVT LFIYIARLYH 120
 LANPRNRQGL RDLISSGVTI QIMTEQESGY CWHNFVNYSN SNESHWPRYP HLWVRLYVLE 180
 LYCIILGLPP CLNILRRKQS QLTSTIALQ SCHYQLRPPH ILWATGLK 228

SEQ ID NO: 85 moltype = AA length = 162
 FEATURE Location/Qualifiers
 source 1..162
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 85
 SPERNYDPRE LRKETYLLE IKWGKSGKLW RHWQNNRTQ HAEVYFLENI FNARRFPNST 60
 HCSITWYLSW SPCAECQKI VDFLKEHPNV NLEIYVARLY YPENERNRQG LRDLVNSGV T 120
 IRIMDLDPYN YCWKTFVSDQ GGDYVWPGH FAPWIKQYSL KL 162

SEQ ID NO: 86 moltype = AA length = 166
 FEATURE Location/Qualifiers
 source 1..166
 mol_type = protein
 organism = Escherichia coli

SEQUENCE: 86
 SEVEFSHEYW MRHALTLAKR AWDEREVPVG AVLVHNNRVI GEGWNRPIGR HDPTAHAEIM 60
 ALRQGGGLVMQ NYRLIDATLY VTLEPCVMCA GAMIHSRIGR VVFGARDAKT GAAGSLMDVL 120
 HHPGMNHRVE ITEGILADEC AALLSDFFRM RQEIKAQKK AQSSTD 166

SEQ ID NO: 87 moltype = AA length = 166
 FEATURE Location/Qualifiers
 source 1..166
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 87
 SEVEFSHEYW MRHALTLAKR ARDEREVPVG AVLVLNNRVI GEGWNRAIGL HDPTAHAEIM 60
 ALRQGGGLVMQ NYRLIDATLY VTLEPCVMCA GAMIHSRIGR VVFGVRNAKT GAAGSLMDVL 120
 HYPGMNHRVE ITEGILADEC AALLCYFFRM PRQVFNAQKK AQSSTD 166

SEQ ID NO: 88 moltype = AA length = 166
 FEATURE Location/Qualifiers
 source 1..166
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 88
 SEVEFSHEYW MRHALTLAKR AWDEREVPVG AVLVLNNRVI GEGWNRSIGL HDPTAHAEIM 60
 ALRQGGGLVMQ NYRLIDATLY VTLEPCVMCA GAMIHSRIGR VVFGVRNAKT GAAGSLMDVL 120
 HYPGMNHRVE ITEGILADEC AALLCYFFRM RQVFNAQKK AQSSTD 166

SEQ ID NO: 89 moltype = AA length = 166
 FEATURE Location/Qualifiers
 source 1..166
 mol_type = protein
 organism = synthetic construct

-continued

SEQUENCE: 89

SEVEFSHEYW	MRHALTLAKR	ALDEREVPVG	AVLVLMNRVI	GEGWNRAIGL	HDPTAHAEIM	60
ALRQGGLVMQ	NYRLIDATLY	VTPEPCVMCA	GAMIHSRIGR	VVFGVRNAKT	GAAGSLMDVL	120
HYPGMNHRVE	ITEGILADEC	NALLCYFFRM	RRQVFNAQKK	AQSSTD		166

SEQ ID NO: 90 moltype = AA length = 166
FEATURE Location/Qualifiers
source 1..166
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 90

SEVEFSHEYW	MRHALTLAKR	ALDEREVPVG	AVLVLMNRVI	GEGWNRAIGL	HDPTAHAEIM	60
ALRQGGLVMQ	NYRLIDATLY	VTPEPCVMCA	GAMIHSRIGR	VVFGVRNAKT	GAAGSLMDVL	120
HYPGMNHRVE	ITEGILADEC	NALLCYFFRM	PRQVFNAQKK	AQSSTD		166

SEQ ID NO: 91 moltype = AA length = 1763
FEATURE Location/Qualifiers
source 1..1763
 mol_type = protein
 organism = unidentified

SEQUENCE: 91

SEVEFSHEYW	MRHALTLAKR	AWDEREVPVG	AVLVHNNRVI	GEGWNRPIGR	HDPTAHAEIM	60
ALRQGGLVMQ	NYRLIDATLY	VTLEPCVMCA	GAMIHSRIGR	VVFGARDAKT	GAAGSLMDVL	120
HHPGMNHRVE	ITEGILADEC	AALLSDFFRM	RRQEIKAQKK	AQSSTDSSGS	SGGSSGSETP	180
GTSESATPES	SGGSSGSSSE	VEFSHEYWMR	HALTLAKRAR	DEREVPVGAV	LVLNNRVIGE	240
GWNRIGLHD	PTAHAEIMAL	RQGGLVMQNY	RLIDATLYVT	PEPCVMCAGA	MIHSRIGRVV	300
FGVRNAKTGA	AGSLMDVLHY	PGMNNRVEIT	EGILADECAA	LLCYFFRMPR	QVFNAQKKAQ	360
SSTDSSGSSG	GSSGSETPGT	SESATPESSG	GSSGSDKKY	SIGLAIGTNS	VGWAVITDEY	420
KVPSKKFKVL	GNTDRHSIKK	NLIGALLFDS	GETAEATRLK	RTARRRYTRR	KNRICYLQEI	480
FSNEMAKVDD	SFFHRLEESF	LVEEDKKHER	HPIFGNIVDE	VAYHEKYPTI	YHLRKKLVDS	540
TKADLRLLIY	LALAHMIKFR	GHFLIEGDLN	PDNSDVKLF	IQLVQTYNQL	FEENPINASG	600
VDAILKLSAR	LSKSRRLENL	IAQLPGEKKN	GLFGNLIALS	LGLTPNPKSN	FDLAEDAKLQ	660
LSKDTYDDDL	DNLLAQIGDQ	YADLFLLAAN	LSDAILLSDI	LRVNTETKA	PLSASMIKRY	720
DEHHQDLTLL	KALVRQQLPE	KYKEIFFDQS	KNGYAGYIDG	GASQEEFYKF	IKPILEKMDG	780
TEELLVKLNR	EDLLRKQRTF	DNGSIPHQIH	LGELHAILRR	QEDFYFPLKD	NREKIEKILT	840
FRIPYVVGPL	ARGNSRFAMW	TRKSEETITP	WNFEVVVDKG	ASAQSFIERM	TNFDKNLPNE	900
KVLPKHSLLY	EYFTVYNELT	KVKYVTEGMR	KPAFLSGEQK	KAIVDLLFKT	NRKVTVKQLK	960
EDYFKKIECF	DSVEISGVED	RFNASLGTYH	DLLKIICKDK	FLDNEENEDI	LEDIVLTLTL	1020
FEDREMIER	LKTYAHLFDD	KVMKQLKRRR	YTGWGRLSRK	LINGIRDKQS	GKTILDFLKS	1080
DGFANRNFMY	LIHDDSLTFK	EDIQKAQVSG	QGDSLHEHIA	NLAGSPAICK	GILQTVKVV	1140
ELVVKVMGRHK	PENIVIEMAR	ENQTTQKGQK	NSRERMKRIE	EGIKELGSQI	LKEHPVENTQ	1200
LQNEKLYLYY	LQNGRDMYVD	QELDINRLSD	YDVVDHIVQS	FLKDDSIDNK	VLTRSDKNRG	1260
KSDNVPSEEV	MKKMKNYWRQ	LLNAKLITQR	KFDNLTKAER	GGLSELDKAG	FIKRLQVETR	1320
QITKHVAQIL	DSRMNTKYDE	NDKLIREVKV	ITLKSCLVSD	FRKDFQFYKV	REINNYHHAH	1380
DAYLNAVVG	ALIKKYPKLE	SEFVYGDYKV	YDVVKMIAS	BQETGKATAK	YFFYSNIMNF	1440
FKETITLANG	BIRKRPLIET	NETGEIVWD	KGRDFATVRK	VLSMPQVNIV	KKTEVQTGGF	1500
SKESILPKRN	SDKLIARKKD	WDPKKYGGFD	SPTVAYSVLV	VAKVEKGKSK	KLKSVKELLG	1560
ITIMERSSEF	KNPIDFLBAK	GYKEVKKDLI	IKLPKYSLFE	LENGRKRMLA	SAGELQKGNE	1620
LALPSKYVMF	LYLASHYVNF	KGSPEDNEQK	QLFVEQHKHY	LDEIEQISE	FSKRVLILADA	1680
NLDKVL SAYN	KHRDKPIREQ	AENIIHLFTL	TNLGAPAAFK	YFDTTIDRKR	YTSTKEVLDA	1740
TLIHQSITSL	YETRIDLSQL	GGD				1763

SEQ ID NO: 92 moltype = AA length = 1565
FEATURE Location/Qualifiers
source 1..1565
 mol_type = protein
 organism = unidentified

SEQUENCE: 92

SEVEFSHEYW	MRHALTLAKR	ARDEREVPVG	AVLVLMNRVI	GEGWNRAIGL	HDPTAHAEIM	60
ALRQGGLVMQ	NYRLIDATLY	VTPEPCVMCA	GAMIHSRIGR	VVFGVRNSKR	GAAGSLMNVL	120
NYPGMNHRVE	ITEGILADEC	AALLCDFYRM	PRQVFNAQKK	AQSSINSGGS	SGGSSGSETP	180
GTSESATPES	SGGSSGSSDK	KYSIGLAIGT	NSVGWAVITD	EYKVPSSKKFK	VLGNTDRHSI	240
KKNLIGALLF	DSGETAEATR	LKRTARRRYT	RRKNRICYLQ	EIFSNEMAKV	DDSFPHRLEE	300
SFLVEEDKKH	ERHPHFNIV	DEVAYHEKYP	TIYHLRKKLV	DSTDADLRL	IYLALAHMIK	360
FRGHFLIEGD	LRPNDSVDK	LFIQLVQTYN	QLFEENPINA	SGVDKAKILS	ARLSKSRRLE	420
NLIAQLPGEK	KNGLFGNLIA	LSLGLTPNFK	SNFDLAEDAK	LQLSKDTYDD	DLNLLAQIG	480
DQYADLFLLA	KNLSDAILLS	DILRVNTEIT	KAPLSASMIK	RYDEHHQDLT	LLKALVRQQL	540
PEKYKEIFPD	QSKNGYAGYI	DGGASQEEFY	KFIKPILEKM	DGTEELLVKL	NREDLLRKQR	600
TFDNGSIPHQ	IHLGELHAIL	RRQEDFYFPL	KDNREKIEKI	LTFRIPYVVG	PLARGNSRFA	660
WMTRKSEETI	TPWNFEFVVD	KGASQAQSFIE	RMNTNFDKNLP	NEKVLPHKSL	LYEYFTVYNE	720
LTQVKYVTEG	MRKPAFLSGE	QKKAIVDLLF	KTNRKVTVKQ	LKEDYFKKIE	CFDSVEISGV	780
EDRFNASLGT	YHDLKKIICK	KDFLDNEENE	DILEDIVLTL	TLFEDREMIE	ERLKYTAHLF	840
DDKVMKQLKR	RRYTGWRGLS	RKLINGIRDK	QSGKTILDFL	KSDGFANRNF	MQLIHDDSLT	900
FKEDIQAQV	SGQGDSLHEH	IANLAGSPAI	KKGILQTVKV	VDELVKVMGR	HKPENIVIE	960
ARENQTTQKG	QKNREREMKR	IEBGIKELGS	QILKEHPVEN	TQLQNEKLYL	YYLQNGRDMY	1020
VDQELDINRL	SDYVDHIVP	QSFLKDDSID	NKVLTRSDKN	RGKSDNVPSE	EVVKMKKNY	1080

-continued

RQLLNAKLIT	QRKFDNLTKA	ERGGSELDDK	AGFIKRQLVE	TRQITKHVAQ	ILDSRMNTKY	1140
DENDKLIREV	KVITLKSCLV	SDFRKDFQFY	KVREINNYHH	AHDAYLNAV	GTALIKKYPK	1200
LESEFVYGDY	KVYDVRKMIA	KSEQEIGKAT	AKYFFYSNIM	NFFKTEITLA	NGEIRKRPLI	1260
ETNGETGEIV	WDKGRDFATV	RKVLSPMPQVN	IVKKTVEVQTG	GFSKESILPK	RNSDKLIARK	1320
KDWDPKKYGG	FDSPTVAYS	LVVAKVEKGK	SKKLKSVKEL	LGITIMERSS	FEKNPIDFLE	1380
AKGYKEVKKD	LIILPKYSL	FELNGRKRK	LASAGELQKG	NELALPSKYV	NFLYLASHYE	1440
KLKGSPEPNE	QKQLFVEQHK	HYLDEIIEQI	SEFSKRVLIA	DANLDKVLSA	YNKHRDKPIR	1500
EQAENIIHLF	TLTNLGAFAA	FKYFDTTIDR	KRYTSTKEVL	DATLIHQSI	GLYETRIDL	1560
QLGGD						1565

SEQ ID NO: 93 moltype = AA length = 1565
 FEATURE Location/Qualifiers
 source 1..1565
 mol_type = protein
 organism = unidentified

SEQUENCE: 93

SEVEFSHEYW	MRHALTLAKR	ARDEREVPVG	AVLVLMNRVI	GEGWNRAIGL	HDPTAHAEIM	60
ALRQGGGLVMQ	NYRLYDATLY	STFPCVMCA	GAMIHSRIGR	VVFGVRNAKT	GAAGSLMDVL	120
HHPGMNRHVE	ITEGILADE	AALLCRFFRM	PRRVFNAQKK	AQSSTDGSGS	SGGSSGSETP	180
GTSESATPES	SGGSSGSGDK	KYSIGLAIGT	NSVGWAVITD	EYKVPSSKKFK	VLGNTDRHSI	240
KKNLIGALLF	DSGETAEATR	LKRTARRRYT	RRKNRICYLQ	EIFSNEMAKV	DDSFPHRLEE	300
SFLVEEDKKH	ERHPIFGNIV	DEVAYHEKYP	TIYHLRKKLV	DSTDKADLRL	IYLALAHMIK	360
FRGHFLIEGD	LNPDSNDVDK	LFIQLVQTYN	QLFEENPINA	SGVDAKAILS	ARLSKSRRL	420
NLIAQLPGEK	KNLFGNLIA	LSLGLTPNFK	SNFDLAEDAK	LQLSKDITYDD	DLNLLAQIG	480
DQYADLFLAA	KNLSDAILLS	DILRVNTEIT	KAPLSASMIK	RYDEHHQDLT	LLKALVRQQL	540
PEKYKEIFPD	QSKNGYAGYI	DGGASQEEFY	KFIKPILEKM	DGTEELLVKL	NREDLLRKQR	600
TFDNGSIPHQ	IHLGELHAIL	RQEDFPYFPL	KDNREKIEKI	LTFRIPYVVG	PLARGNSRFA	660
WMTRKSEETI	TPWNFEVVVD	KGASAQSFIE	RMTNFDKNLP	NEKVLPHKSL	LYEYFTVYNE	720
LTKVKYVTEG	MRKPAFLSGE	QKKAIVDLLF	KTNRKVTVKQ	LKEDYFKKIE	CFDSVEISGV	780
EDRFNASLGT	YHDLKKIKD	KDFLDNEENE	DILEDIVLTL	TLFEDREMIE	ERLKYAHLEF	840
DDKVMKQLKR	RRYTGWGRLS	RKLINGIRDK	QSGKTILDFL	KSDGFANRNF	MQLIHDDSLT	900
FKEDIQKAQV	SGQGDSLHEH	IANLAGSPA	KKGILQTVKV	VDELVKVMGR	HKPENIVIE	960
ARENQTTQKG	QKNSRERMKR	IEEGIKELGS	QILKEHPVEN	TQLQNEKLYL	YYLQNGRDMY	1020
VDQELDLNRL	SDYVDHIVP	QSFLKDDSID	NKVLTRSDKN	RGKSDNVPSE	EVVKMKMNYW	1080
RQLLNAKLIT	QRKFDNLTKA	ERGGSELDDK	AGFIKRQLVE	TRQITKHVAQ	ILDSRMNTKY	1140
DENDKLIREV	KVITLKSCLV	SDFRKDFQFY	KVREINNYHH	AHDAYLNAV	GTALIKKYPK	1200
LESEFVYGDY	KVYDVRKMIA	KSEQEIGKAT	AKYFFYSNIM	NFFKTEITLA	NGEIRKRPLI	1260
ETNGETGEIV	WDKGRDFATV	RKVLSPMPQVN	IVKKTVEVQTG	GFSKESILPK	RNSDKLIARK	1320
KDWDPKKYGG	FDSPTVAYS	LVVAKVEKGK	SKKLKSVKEL	LGITIMERSS	FEKNPIDFLE	1380
AKGYKEVKKD	LIILPKYSL	FELNGRKRK	LASAGELQKG	NELALPSKYV	NFLYLASHYE	1440
KLKGSPEPNE	QKQLFVEQHK	HYLDEIIEQI	SEFSKRVLIA	DANLDKVLSA	YNKHRDKPIR	1500
EQAENIIHLF	TLTNLGAFAA	FKYFDTTIDR	KRYTSTKEVL	DATLIHQSI	GLYETRIDL	1560
QLGGD						1565

SEQ ID NO: 94 moltype = AA length = 364
 FEATURE Location/Qualifiers
 source 1..364
 mol_type = protein
 organism = unidentified

SEQUENCE: 94

SEVEFSHEYW	MRHALTLAKR	AWDEREVPVG	AVLVHNNRVI	GEGWNRPIGR	HDPTAHAEIM	60
ALRQGGGLVMQ	NYRLIDATLY	VTLEPCVMCA	GAMIHSRIGR	VVFGARDAKT	GAAGSLMDVL	120
HHPGMNRHVE	ITEGILADE	AALLSDFFRM	RQEIKAQKK	AQSSTDGSGS	SGGSSGSETP	180
GTSESATPES	SGGSSGSGSE	VEFSHEYWMR	HALTLAKRAR	DEREVPVGAV	LVLNNRIVGE	240
GNRAIGLHD	PTAHAEIMAL	RQGGGLVMQNY	RLIDATLYVT	FEPVCVMCAGA	MIHSRIGRVV	300
FGVRNAKTGA	AGSLMDVLHY	PGMNRHVEIT	EGILADECAA	LLCYFFRMPR	QVFNAQKKAQ	360
SSTD						364

SEQ ID NO: 95 moltype = AA length = 167
 FEATURE Location/Qualifiers
 source 1..167
 mol_type = protein
 organism = unidentified

SEQUENCE: 95

MSEVEFSHEY	WMRHALTLAK	RARDEREVPV	GAVLVNNRV	IGEGWNRAIG	LHDPTAHAEI	60
MALRQGGGLVM	QNYRLIDATL	YSTFPCVMC	AGAMIHSRIG	RVVFGVRNAK	TGAAGSLMDV	120
LHHPGMNHRV	EITEGILADE	CAALLCRFFR	MPRRVFNAQK	KAQSSTD		167

SEQ ID NO: 96 moltype = AA length = 167
 FEATURE Location/Qualifiers
 source 1..167
 mol_type = protein
 organism = unidentified

SEQUENCE: 96

MSEVEFSHEY	WMRHALTLAK	RARDEREVPV	GAVLVNNRV	IGEGWNRAIG	LHDPTAHAEI	60
MALRQGGGLVM	QNYRLIDATL	YVTFPCVMC	AGAMIHSRIG	RVVFGVRNSK	RGAAGSLMNV	120

-continued

LNYPGMNHRV EITEGILADE CAALLCDFYR MPRQVFNAQK KQSSIN 167

SEQ ID NO: 97 moltype = AA length = 83
 FEATURE Location/Qualifiers
 source 1..83
 mol_type = protein
 note = Bacillus phage AR9
 organism = unidentified

SEQUENCE: 97
 TNLSDIIEKE TKGQLVIOES ILMLPEEVVEE VIGNKPESDI LVHTAYDEST DENVMLLTSD 60
 APEYKPWALV IQDSNGENKI KML 83

SEQ ID NO: 98 moltype = length =
 SEQUENCE: 98
 000

SEQ ID NO: 99 moltype = length =
 SEQUENCE: 99
 000

SEQ ID NO: 100 moltype = length =
 SEQUENCE: 100
 000

SEQ ID NO: 101 moltype = DNA length = 66
 FEATURE Location/Qualifiers
 source 1..66
 mol_type = other DNA
 organism = unidentified

SEQUENCE: 101
 ttcttgctgt acttatagat cgctacgtta tttcaatddd gaaaatctga gtcctgggag 60
 tgcgga 66

SEQ ID NO: 102 moltype = AA length = 605
 FEATURE Location/Qualifiers
 source 1..605
 mol_type = protein
 organism = unidentified

SEQUENCE: 102
 MSGWESYYKT EGDEEAEEEQ EENLEASGDY KYSGRDSLIF LVDASKAMFE SQSEDELTPF 60
 DMSIQCIQSV YISKIISDR DLLAWFYGTE KDKNSVNFKI YVLQELDNPG AKRILELDQF 120
 KGQQGQKRFQ DMMGHGSDYS LSEVLWVCAN LFSVDVQFKMS HKRIMLFTNE DNPNGNDSAK 180
 ASRARTKAGD LRDTGIFLDL HLKKPGGFDI SLFYRDIISI AEDEDLRVHF EESSKLEDLL 240
 RKVRKETRK RALSRLKLLK NKDIVISVGI YNLVQKALKP PPIKLYRETN EPVTKTRTF 300
 NTSTGGLLLP SDTKRSQIYG SRQIILEKEE TEELKRFDPP GLMLMGFKPL VLLKKHHYLR 360
 PSLFVYPEES LVIGSSTLFS ALLIKCLEKE VAALCRYTPR RNIPPYFVAL VPQEEELDDQ 420
 KIQVTPPGFQ LVFLPFADDK RKMPPTEKIM ATPQVQGMK AIVEKLRFY RSDSFENPVL 480
 QQHFRNLEAL ALDMEPEQA VDLTLPKVEA MNKRLGSLVD EFKELVYPPD YNPEGKVTKR 540
 KHDNEGSGSK RPKVEYSEEE LKTHISKGTG GKFTVPLKEA CRAYGLKSGL KQELLEALT 600
 KHFQD 605

SEQ ID NO: 103 moltype = AA length = 482
 FEATURE Location/Qualifiers
 source 1..482
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 103
 MVRSGNKA AW LCMDVGFTMS NSIPGIESPF EQAKKVITMF VQRQVFAENK DEIALVLFGT 60
 DGTDNPLSGG DQYQNTVHR HLMLPDFDLL EDIESKIQPG SQQADFLDAL IVSMDVIQHE 120
 TIGKKFEKRH IEIFTDLSSR FSKSQLDIII HSLKKCDISE RSHIHWPCRL TIGSNLSIRI 180
 AAYKSILQER VKKTTWDAKT LKKEDIQKET VYCLNDDDET EVLKEDIIQG FRYGSDIVPF 240
 SKVDEEQMKY KSEGKCFSVL GFCKSSQVQR RFFMGNQVLK VFAARDDEAA AVALSSLIHA 300
 LDDLDIWAIV RYAYDKRANP QVGVAFFPHK HNYECLVYVQ LPFMEDLRQY MFSSLKNSKK 360
 YAPTEAQLNA VDALDSMSL AKKDEKDTL EDLPPTTKIP NPRQRLFQC LLHRLALPRE 420
 PLPPIQQHIW NMLNPPAEVT TKSQIPLSKI KTLFPLIEAK KKDQVTAQEI FQPNHEDGPT 480
 AK 482

SEQ ID NO: 104 moltype = DNA length = 10
 FEATURE Location/Qualifiers
 source 1..10
 mol_type = other DNA
 organism = unidentified

SEQUENCE: 104
 aatttttgga 10

SEQ ID NO: 105 moltype = AA length = 83

-continued

FEATURE	Location/Qualifiers	
source	1..83	
	mol_type = protein	
	organism = unidentified	
SEQUENCE: 105		
GSVIDVSSQR VNVQRPLDAL GNSLNSPVII KLKGDREFRG VLKSFDLHMN LVLNDAAEELE		60
DGEVTRRLGT VLIRGDNIVY ISP		83
SEQ ID NO: 106	moltype = DNA length = 25	
FEATURE	Location/Qualifiers	
source	1..25	
	mol_type = other DNA	
	note = bacteriophage MS2	
	organism = unidentified	
SEQUENCE: 106		
gcgcacatga ggatcaccca tgtgc		25
SEQ ID NO: 107	moltype = AA length = 116	
FEATURE	Location/Qualifiers	
source	1..116	
	mol_type = protein	
	note = bacteriophage MS2	
	organism = unidentified	
SEQUENCE: 107		
MASNFTQFVL VDNNGTGDVT VAPSNFANGI AEISSNSRSQ AYKVTCSVRQ SSAQNRKYTI		60
KVEVPKGAWR SYLNMELTIP IFATNSDCEL IVKAMQGLLK DGNPIPSAIA ANSGIY		116
SEQ ID NO: 108	moltype = DNA length = 26	
FEATURE	Location/Qualifiers	
source	1..26	
	mol_type = other DNA	
	note = bacteriophage PP7	
	organism = unidentified	
SEQUENCE: 108		
ataaggagtt tatatggaaa ccctta		26
SEQ ID NO: 109	moltype = AA length = 127	
FEATURE	Location/Qualifiers	
source	1..127	
	mol_type = protein	
	note = bacteriophage PP7	
	organism = unidentified	
SEQUENCE: 109		
MSKTIVLSVG EATRTLTEIQ STADRQIFEE KVGPLVGLRL LTASLRQNGA KTAYRVNLKL		60
DQADWDCSTS VCGELPKVRY TQVWSDVTI VANSTEASRK SLYDLTKSLV ATSQVEDLVV		120
NLVPLGR		127
SEQ ID NO: 110	moltype = DNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = genomic DNA	
	organism = Zea mays	
SEQUENCE: 110		
ctgaatgcct gcgagcatc		19
SEQ ID NO: 111	moltype = AA length = 62	
FEATURE	Location/Qualifiers	
source	1..62	
	mol_type = protein	
	organism = Zea mays	
SEQUENCE: 111		
MKSIRCKNCN KLLFKADSFH HIEIRCPRCK RHIIMLNACE HPTEKHCGKR EKITHSDETV		60
RY		62
SEQ ID NO: 112	moltype = AA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 112		
EELLSKNYHL ENEVARLKKG SGSG		24
SEQ ID NO: 113	moltype = AA length = 241	
FEATURE	Location/Qualifiers	
source	1..241	
	mol_type = protein	

-continued

organism = synthetic construct

SEQUENCE: 113

EEELLSKNYH	LENEVARLKK	GSGSGEELLS	KNYHLENEVA	RLKKGSGSGE	ELLSKNYHLE	60
NEVARLKKGS	GSGEELLSKN	YHLENEVARL	KKGSGSGEEL	LSKNYHLENE	VARLKKGSGS	120
GEELLSKNYH	LENEVARLKK	GSGSGEELLS	KNYHLENEVA	RLKKGSGSGE	ELLSKNYHLE	180
NEVARLKKGS	GSGEELLSKN	YHLENEVARL	KKGSGSGEEL	LSKNYHLENE	VARLKKGSGS	240
G						241

SEQ ID NO: 114 moltype = AA length = 277

FEATURE Location/Qualifiers

source 1..277

 mol_type = protein

 organism = synthetic construct

SEQUENCE: 114

MGPDIIVMTQS	PSSLSASVGD	RVTITCRSST	GAVTTSNYAS	WVQEKPGKLF	KGLIGGTNNR	60
APGVPSRFRSG	SLIGDKATLT	ISSLQPEDFA	TYFCALWYSN	HWVFGQGTKV	ELKRGGGGSG	120
GGGSGGGGSS	GGGSEVKLLE	SGGGLVQPGG	SLKLSCAVSG	FSLTDYGVNW	VRQAPGRGLE	180
WIGVIWGDGI	TDYNSALKDR	FIISKDNGKN	TVYLQMSKVR	SDDTALYYCV	TGLFDYWGQG	240
TLVTVSSYPY	DVPDYAGGGG	GSGGGSGGGG	GSGGGGS			277

SEQ ID NO: 115 moltype = AA length = 1227

FEATURE Location/Qualifiers

source 1..1227

 mol_type = protein

 organism = synthetic construct

SEQUENCE: 115

SKLEKFTNCY	SLSKTLRFKA	IPVGKTQENI	DNKRLLEVEDE	KRAEDYKGVK	KLLDRYYLSF	60
INDVLHSIKL	KNLNYYISLF	RKKTRTEKEN	KELENLEINL	RKEIAKAFKG	NEGYKSLFKK	120
DIETILPEF	LDDKDEIALV	NSFNGFTTAF	TGFFDNRENM	FSEEAKSTSI	AFRCINENLT	180
RYISNMDIFE	KVDAIFDKHE	VQEIKEKILN	SDYDVEDFFE	GEFFNFVLTQ	EGIDVYNAII	240
GGFVTESGEG	IKGLNEVINL	YNQKTKQKLP	KFKPLYKQVL	SDRESLSFYG	EGYTSDEEVL	300
EVFRNTLNKN	SEIFSSIKKL	EKLKKNFDEY	SSAGIFVKNG	PAISTISKDI	FGEWNVIRDK	360
WNAEYDDIHL	KKKAVVTEKY	EDDRKSFKK	IGSFSLEQLQ	EYADADLSV	EKLKEIIQK	420
VDEIYKVYGS	SEKLFADDFV	LEKSLKNDK	VVAIMKDLLD	SVKSFENYIK	APFGEGKETN	480
RDESFYGDVF	LAYDILLKVD	HIYDAIRNYV	TQKPYSKDKF	KLYFQNPQFM	GGWDKDKETD	540
YRATILRYGS	KYYLAIMDKK	YAKCLQKIDK	DDVNGNYEKI	NYKLLPGPNK	MLPKVFFSKK	600
WMAYYNPSID	IQKIYKNGTF	KKGDMFNLND	CHKLIDFFKD	SISRYPKWSN	AYDFNFSETE	660
KYKDIAGFYR	EVEEQGYKVS	FESASKKEVD	KLVEEGKLYM	FQIYNKDFSD	KSHGTPNLHT	720
MYFKLLFDEN	NHGQIRLSGG	AELFMRRASL	KKEELVVHPA	NSPIANKNPD	NPKKTTTLSY	780
DVYKDKRFE	QKYLHIPIA	INKCPKNIFK	INTEVRVLLK	HDDNPYVIGI	ARGERNLLYI	840
VVVDGKGNIV	EQYSLNEIIN	NFNGIRIKTD	YHSLLDKKEK	ERFEARQNW	SIENIKELKA	900
GYISQVVKHI	CELVKDYDAV	IALEDLNSGF	KNSRVKVEKQ	VYQKFEKMLI	DKLNYMVDKK	960
SNPCATGGAL	KGQITTNKFE	SFKSMSTQNG	FIFYIPAWLT	SKIDPSTGFV	NLLKTKYTSI	1020
ADSKKFISFF	DRIMYVPEED	LFEPALDYKN	FSRTDADYIK	KWKLYSYGNN	IRIFRNPCKN	1080
NVFDWEEVCL	TSAYKELFNK	YGINYQQGDI	RALLCEQSDK	AFYSSFMALM	SLMLQMRNSI	1140
TGRTDVFLLI	SPVKNSDGIF	YDSRNVEAQE	NAILPKNADA	NGAYNIARKV	LWAIGQFKKA	1200
EDEKLDKVKI	AISNKEWLEY	AQTSVKKH				1227

SEQ ID NO: 116 moltype = AA length = 1307

FEATURE Location/Qualifiers

source 1..1307

 mol_type = protein

 note = Acidaminococcus sp.

 organism = unidentified

SEQUENCE: 116

MTQFEGFTNL	YQVSKTLRFE	LIPQGKTLKH	IQEQGFIEED	KARNDHYKEL	KPIIDRIYKT	60
YADQCLQLVQ	LDWENLSAAI	DSYRKEKTEE	TRNALIEBQA	TYRNAIHDFY	IGRTDNLTD	120
INKRHAEIYK	GLFKAELFNG	KVLKQLGTVT	TTEHENALLR	SFDKFTTYFS	GFYENRKNVF	180
SAEDISTAIP	HRIVQDNFPK	FKENCHIFTR	LITAVPSLRE	HFENVKKAIG	IFVSTSIEEV	240
FSFPFYNQLL	TQTQIDLYNQ	LLGGISREAG	TEKIKGLNEV	LNLAIQKND	TAHIIASLPH	300
RFIPLFKQIL	SDRNTLSFIL	EEFKSDEEVI	QSFCYKTL	RNENVLETA	ALFNELNSID	360
LTHIFISHKK	LETISSALCD	HWDTLRNALY	ERRISELTGK	ITKSAKEKVQ	RLKHEDINL	420
QEIISAAGKE	LSEAFKQKTS	EILSHAHAA	DQPLPTTLKK	QEEKEILKSQ	LDSLLGLYHL	480
LDWFAVDEN	EVDPEFSARL	TGKLEMEPS	LSFYNKARNY	ATKKPYSVEK	FKLNFQMPTL	540
ASGWDVNKEK	NGAILLVKFN	GLYYLGIMPK	QKGRYKALSF	EPTKTESEGF	DKMYDYFPD	600
AAKMIPKCS	QLKAVTAHFQ	THTTPIILLSN	NFIEPLEITK	EIYDLNNEPK	EPKKFQTAYA	660
KKTGDQKGYR	EALCKWIDFT	RDFLSKYTKT	TSIDLSSLRP	SSQYKDLGEY	YAEINPLLYH	720
ISFQRIAEKE	IMDAVETGKL	YLFQIYNKDF	AKGHHGKPNL	HTLYWTGLFS	PENLAKTSIK	780
LNGQAELFYR	PKSRMKRMAH	RLGEKMLNKK	LKDQKTPIPD	TLYQELYDYV	NHRLSHDLSD	840
EARALLPNVI	TRKVSHEIHK	DRRFTSDKFF	FHVPIITLNYQ	AANSPSKFNQ	RVNAYLKEHP	900
ETPIIGIDRG	ENLIIYITVI	DSTGKILEQR	SLNTIQQFDY	QKKLDNREKE	RVAARQAWSV	960
VGTIKDLKQG	YLSQVIHEIV	DLMIHYQAVV	VLENLNFQFK	SKRTGIAEKA	VYQQFEKMLI	1020
DKLNLCLVLK	YPAEKVGVGL	NPYQLTDQFT	SFAKMGTSQG	FLFYVPAPYT	SKIDPLTGTV	1080
DPFVWKTIAK	HESRKHFLG	FDLHYDVKT	GDFILHFKMN	RNLSFQRLP	GFMPAWDIVF	1140
EKNETQFPAK	ETPIIAGKRI	VPVIENHRFT	GRYRDLYPAN	ELIALLEEKG	IVFRDGSNIL	1200
PKLLENDSSH	AIDTMVALIR	SVLQMRNSNA	ATGEDYINSP	VRDLNGVCFD	SRFQNPFWPM	1260

-continued

DADANGAYHI ALKGQLLLNH LKESKDLKLQ NGISNQDWLA YIQELRN 1307

SEQ ID NO: 117 moltype = AA length = 1241
FEATURE Location/Qualifiers
source 1..1241
 mol_type = protein
 organism = Butyrivibrio proteoclasticus

SEQUENCE: 117
MLLYENYTKR NQITKSLRLE LRPQGKTLRN IKELNLLQEQ KAIYALLERL KPVIDEGIKD 60
IARDTLKNCE LSFEEKLYEHF LSGDKKAYAK ESERLKKEIV KTLIKNLPEG IGKISEINSA 120
KYLNGVLVDF IDKTHKDSSE KQNILSDILE TKGYLALFSK FLTSRITTLE QSMPCRVIEN 180
FEIYAANIPK MQDALERGAV SFAIEYESIC SVDYNNQILS QEDIDSYNRL ISGIMDEDGA 240
KEKGINQTIS EKNIKIKSEH LEEKPFRILK QLHKQILEER EKAPTIDHID SDEEVQVTK 300
EAFEQTKEQW ENIKKINGFY AKDPGDITLF IVVGNQTHV LSQLIYGEHD RIRLLLEEYE 360
KNTLEVLPRR TKSEDARYDK FVNAVKKVA KESHTFDGLQ KMTGDDRLFI LYRDELARNY 420
MRIKAYGTF ERDILKSRRG IKGNRDVQES LVSFYDELTK FRSLARIINS GNDEKADPIF 480
YNTFDGIFEK ANRTYKAENL CRNYVTKSPA DDARIMASCL GTPARLRTHW WNGEENFAIN 540
DVAMIRRGDE YYYFVLTPDV KPVDLKTDE TDAQIFVQRK GAKSFLGLPK ALFKCILEPY 600
FESPEHKNDK NCVIEEYVSK PLTIDRRAYD IPKNGTFKKT NIGIDGLTEE KFKDDCRYLI 660
DVYKEFIADV TRYSCFNMSG LKRADEYNDI GEFFSDVDTR LCTMEWIPVS FERINDMVDK 720
KEGLLFLVRS MFLYNRPKP YERTFIQLFS DSNMEHTSML LNSRAMIQYR AASLPRRVTH 780
KKGSLVALR DSNGEHIPMH IREAIYKMK NFDISSEDFI MAKAYLAHD VAIKANEDI 840
IRNRRYTEDK PFLSLSYTKN ADISARTLDY INDKVEEDTQ DSRMAVIVTR NLKDLTYVAV 900
VDEKNNVLEE KSLNEIDGVN YRELLKERTK IKYHDKTRLW QYDVSSKGLK EAYVELAVTQ 960
ISKLATKYNA VVVVESMSST FKDKFSFLDE QIFKAFEARL CARMSDLSFN TIKEGEAGSI 1020
SNPIQVSNMN GNSYQDGVYI FLNNAYTRTL CPDTGFVDVF DKTRLITMQS KRQFFAKMKD 1080
IRIDDGEMLF TFNLEEYPTK RLDRKEWTV KIAAGDSYFD KDKGEYVYN DIVREQIIPA 1140
LLEDKAVFDG NMAEKFLDKT AISGKSVELI YKWFANALYG IITKKDGEKI YRSPITGTEI 1200
DVSKNTTYNF GKFKMFQKEY RGDGDFLDAF LNYMQAQDIA V 1241

SEQ ID NO: 118 moltype = AA length = 1238
FEATURE Location/Qualifiers
source 1..1238
 mol_type = protein
 organism = Methanoplasma termitum

SEQUENCE: 118
MNNYDEFTKL YPIQKTIRFE LKPQGRTEH LETFNFFEEED RDRAEKYKIL KEAIDEYHKK 60
FIDEHLTNMS LDWNSLKQIS EKYKSREEK DKKVFLSEQK RMRQEVSEF KKDDRFDLDF 120
SKKLFSELK EEIYKGNHQ EIDALKSFDK FSGYFGLHE NRKNMYSDDG EITAINSRIV 180
NNFPKFLDN LQKYQEARLK YPEWIIKAES ALVAHNIKMD IVFSLEYFNK VLNQEGIQRY 240
NLALGGYVTK SGEKMMGLND ALNLAHQSEK SSKGRIHMTF LFKQILSEKE SFSYIPDVFT 300
EDSQLPSIG GFFAQIENDK DGNIFDRALE LISSYAEYDT ERIYIRQADI NRVSNIIFGE 360
WGTGLGLMRE YKADSNIDIN LERTCKVVDK WLDKKEFALS DVLEAIDRTG NNDAFNEYIS 420
KMRTAREKID AARKEMKFI EKSIGDEESI HIIKTLDSV QQFLHFFNLF KARQDIPLDG 480
AFYAEFDEVH SKLPAIVPLY NKVRNYLTKN NLNTKKIKLN FKNPTLANGW DQNKVYDYAS 540
LIFLRDGNYY LGIINPKRKK NIKFEQSGSN GPFYRKMVYK QIPGPNKNLR PVFLTSTKKG 600
KEYKPSKEII EGYEADKHIR GDKFDLDFCH KLIDFFKESI EKHKDWSKFN FYFSPTESYG 660
DISEFYLDVE KQGYRMHFEN ISAEITIDEYV EKGDLFLFOI YNKDFVKAAT GKDKMHTIYW 720
NAAFSPENLQ KVVVKLNGEA ELFYRDKSDI KEIVHREGEI LVNRTYNGRT PVPDKHKKL 780
TDYHNGRTKD LGEAKEYLDK VRYFKAHYDI TKDRRYLNDK IYFHVPLTLN FKANGKKNLN 840
KMVIEKFLSD EKAHIIIGDR GERNLYYSI IDRSKGIIDQ QSLNVIDGFD YREKLNQREI 900
EMKDARQSNV AIGKIKDLKE GYLSKAVHEI TKMAIQYNAI VVMEELNYGF KRGRFKVEKQ 960
IYQKFENMLI DKMNYLVFKD APDESPGGVL NAYQLTNPLE SFAKLKGQTG ILFYVPAAYT 1020
SKIDPTTGFV NLFNTSKTN AQERKEFLQK FESISYSAKD GGIFAFAPDY RKFGTSKTDH 1080
KNVWTAYTNG ERMRYIKEKK RNELFDPSKE IKEALTSSGI KYDGGQNILP DILRSNNNGL 1140
IYTMYSFIA AIQMRVYDGK EDYIISPIKN SKGEFFRTDP KRRELPIAD ANGAYNIALR 1200
GELTMRAIAE KFDPDSEKMA KLELKHKDW EFMQTRGD 1238

SEQ ID NO: 119 moltype = AA length = 1281
FEATURE Location/Qualifiers
source 1..1281
 mol_type = protein
 organism = Eubacterium eligens

SEQUENCE: 119
MNGNRSIVYR EFGVGVPAK TLRNELRPVG HTQEHIIQNG LIQEDELQRE KSTELKNIMD 60
DYRYEYIDKS LSGVTDLDFT LLFELMNLVQ SSPSKDNKKA LEKEQSKMRE QICTHQSDS 120
NYKNIFNAKL LKEILPDFIK NYNQYDVVKD AGKLETALF NGFSTYFTDF FEKRKNVFTK 180
EAVSTSIAYR IVHENSILFL ANMYSYKKIS EKALDEIEVI EKNQDKMGD WELNQIFNPD 240
FYNMVLISQS IDFYNEICGV VNAHMNLYCQ QTKNNYNLFK MRKLHKQILA YTSTSFVVPK 300
MPEDDMSVYN AVNAFIDE TE KGNIIIGLKD IVNKYDELDE KRIYISKDFY ETLSCFMSGN 360
WNLITGCVEN ERMRYIHAKG KSKEEKVKKA VKEDKYK SIN DVNDLVEKYI DEKERNFKN 420
SNAKYIREI SNIITDTETA HLEYDDHISL IESEEKADEM KKRLDMMYMN YHWAKAFIVD 480
EVLDRDEMPY SDIDDIYNIL ENIVPLYNRV RNYVTQKPYN SKKIKLNFQS PTLANGWSQS 540
KEFDNNAIL IRDNKYLA I FNAKNKPKK IIQGNSDKKN DNDYKMMVYN LLPGANKMLP 600
KVFLSKKGIE TRPKSDYIIS GYNAKHKT SENFDISFCR DLIDYFKNSI EKHAERWKYE 660
FKFSATDSYS DISEFYREVE MQGYRIDWTY ISEADINKLD BEGKIYLFQI YNKDFAENST 720

-continued

GKENLHTMYF	KNIFSEENLD	KIIKLNGQAE	LPYRRASVKN	PVKHKKDSVL	VNKTYKNQLD	780
NGDVVRIPIP	DDIYNEIYKM	YNGYIKESDL	SEAAKEYLDK	VEVRTAQKDI	VKDYRYTVDK	840
YFIHTPITIN	YKVTARNNVN	DMVVKYIAQN	DDIHVIGIDR	GERNLIYISV	IDSHGNIVKQ	900
KSYNILNNYD	YKKKLVEKEK	TREYARKNWK	SIGNIKELKE	GYISGVVHEI	AMLIVEYNAI	960
IAMEDLNYGF	KRGRFKVERQ	VYQKFESMLI	NKLNYFASKE	KSVDEPGGLL	KGYQLTYVPD	1020
NIKNLGKQCG	VIFVVPAAFT	SKIDPSTGFI	SAFNFKSIST	NASRKQFFMQ	FDEIRYCAEK	1080
DMFSFGFDYN	NFDTYNTIMG	KTQWTVYTNQ	ERLQSEFNNA	RRTGKTKSIN	LTETIKLLLE	1140
DNEINYADGH	DIRIDMEKMD	EDKKSEFFAQ	LLSLYKLTVO	MRNSYTEAE	QENGISYDKI	1200
ISPVINDEGE	FFDSDNKYES	DDKECKMPKD	ADANGAYCIA	LKGLYEVLKI	KSEWTEGDFD	1260
RNCLKLPHAE	WLDFTQNKRY	E				1281

SEQ ID NO: 120 moltype = AA length = 1300
 FEATURE Location/Qualifiers
 source 1..1300
 mol_type = protein
 organism = Francisella tularensis

SEQUENCE: 120

MSIYQEFVVK	YLSKTLRFE	LIPQGKLTEN	IKARGLILDD	EKRAKDYKKA	KQIIDKYHQF	60
FIEEILSSVC	ISEDLLQNY	DVYFKLKKSD	DDNLQKDFKS	AKDTIKKQIS	EYIKDSEKFK	120
NLFNQNLIDA	KKGQESDLIL	WLKQSKDNGI	ELFKANSDIT	DIDEALEIIK	SFKGWTTYFK	180
GFHENRKVNY	SSNDIPTSI	YRIVDDNLPK	FLENKAKYES	LKDKAPEAIN	YEQIKKDLAE	240
ELTFDIDYKT	SEVNQRVFSL	DEVFEIANFN	NVLNQSGITK	FNTIIGGKFV	NGENTKRRKI	300
NEYINLYSQQ	INDKTLKKYK	MSVLFKQILS	DTESKSFVID	KLEDDSDVVT	TMQSFYEQIA	360
AFKTVEEKSI	KETLSLLFDD	LKAQKLDLSK	IYFKNDKSLT	DLSQQVDDY	SVIGTAVLEY	420
ITQOIAPKML	DNPSKKEQEL	IAKTEKAKY	LSLETIKLAL	EEFNKHRDID	KQCRFEIILA	480
NFAAIPMIFD	EIAQNKDNL	QISIKYQNGG	KKDLLQASAE	DDVKAIKDLL	DQTNLLHLKL	540
KTFHISQSED	KANILDKDEH	FYLVFEECYF	ELANIVPLYN	KIRNYITQKP	YSDEKPKLNF	600
ENSTLANGWD	KNKEPDNTAI	LFIKDDKYVL	GVMNKKNNKI	FDDKAIKENK	GEYKKIVYK	660
LLPGANKMLP	KVFFSAKSIK	FYNPSEDILR	IRNHSTHTKN	GSPQKGYEKF	EFNIEDCRKF	720
IDFYKQSISK	HPWEKDFGFR	FSDTQRYNSI	DEFYREVENQ	GYKLTFFENIS	ESYIDSVVNG	780
GKLYLFIQYN	KDFSAYSGFR	PNLHTLYWKA	LPDERNLQDV	VYKLNGEAE	FYRKQSIKK	840
ITHPAKEAIA	NKNKDNPKKE	SVFEYDLIKD	KRFTEDKFFF	HCPITINFPS	SGANKFNDEI	900
NLLLKEKAND	WHILSIDRGE	RHLAYYTLVD	GKGNIIKQDT	FNIIIGNDRMK	TNYHDKLAAI	960
EKDRDSARKD	WKKLNNIKEM	KEGYLSQVHV	ETAKLVIEYN	AIVVFEDLNF	GFKRGRFKVE	1020
KQVYQKLEKM	LIEKLNLYLVF	KDNEFDKTGG	VLRAYQLTAP	FETPKKMGKQ	TGIIYYVPAG	1080
FTSKICPVTV	FPVNLQPKYE	SVSKSQEFFS	KFDKICYNLD	KGYFEFSFDY	KNFGDKAAKG	1140
KWTIASFGSR	LINFNRNSDKN	HNWDTREVYP	TKELEKLLKD	YSIEYGHGEC	IKAAICGESD	1200
KKFFAKLTSV	LNTILQMRNS	KTGTEDLYLI	SPVADVNGNF	PDSRQAPKQM	PQDADANGAY	1260
HIGLKGLMLL	GRIKNNQEGK	KLNLVIKNEE	YFEFVQNRNN			1300

SEQ ID NO: 121 moltype = AA length = 1206
 FEATURE Location/Qualifiers
 source 1..1206
 mol_type = protein
 note = Lachnospiraceae bacterium
 organism = unidentified

SEQUENCE: 121

MYYESLTQKY	PVSKTIRNEL	IPIGKTLTDL	RQNNILESVD	KRKQNYEHVK	GILDEYHKQL	60
INEALDNCTL	PSLKIAAEIY	LKNQKEVSDR	EDFNKTQDLL	RKEVVEKLKA	HENFTKIGKK	120
DILDLEKLKP	SISEDYNYAL	ESFRNFYTYF	TSYNKVRENL	YSDKEKSSTV	AYRLINENFP	180
KPLDNVKSYP	FVKTAGILAD	GLGEEBQDSL	FIVETFNKTL	TQDGIDTYNS	QVGKINSIN	240
LYNQKNQKAN	GFRKIPKMKM	LYKQILSDRE	ESFIDEFQSD	EVLIDNVESY	GSVLIESLKS	300
SKVSAPFDAL	RESKGNVYV	KNDLAKTAMS	VIVFENWRTF	DDLLNQEYDL	ANENKKKDDK	360
YFEKRQKELK	KNKSYSLEHL	CNLSEDSCNL	IENYIHQISD	DIENIIINNE	TFLRIVINEH	420
DRSRKLAKNR	KAVKAIKDFL	DSIKVLEREL	KLINSSGQEL	EKDLIVYSAH	EELLVELKQV	480
DSLYNMTRNY	LTKKPFSTEK	VKLNFNRSTL	LNGWDRNKET	DNLGVLLKLD	GKYYLGIMNT	540
SANKAFVNP	VAKTEKVPFK	VDYKLLPVPN	QMLPKVFFAK	SNIDFYNPSS	EIYSNYKKG	600
HKKGNMFSLE	DCHNLIDFFK	ESISKHEDWS	KFGPKEDTQA	SYNDISEFYR	EVEKQGYKLT	660
YTDIDETIYN	DLIERNELYL	FQIYNKDFSM	YSKGLNLNLT	LYFMMLFDQR	NIDDVVYKLN	720
GEAEVFPYRPA	SISEDELIH	KAGEEIKNKN	PNRARTKETS	TFSYDIDVKD	RYSKDKFTLH	780
IPITMNFQVD	EVKRFENDAVN	SAIRIDENVN	VIGIDRGERN	LLYVVVIDSK	GNILEQISLN	840
SIINKEYDIE	TDYHALDLER	EGGRDKARKD	WNTVENIRDL	KAGLYLQVVN	VVAKLVLYKN	900
AIICLEDLNF	GFKRGRQKVE	KQVYQKFEKM	LIDKLNLYVI	DKSREQTSPK	ELGGALNALQ	960
LTSKPKSPKE	LGKQSGVIYY	VPAYLTSKID	PTTGAFANLFY	MKCNENVEKSK	RFPDGFDFIR	1020
FNALENVPEF	GFDYRSFTQR	ACGINSKQTV	CTNGERIIKY	RNPDKNNMFD	EKVVVVTDEM	1080
KNLFEQYKIP	YEDGRNVKDM	IISNEEAIFY	RRLYRLLQQT	LQMRNSTSDG	TRDYIISPVK	1140
NKREAYFNSE	LSDGSPVKDA	DANGAYNIAR	KGLWVLEQIR	QKSEGEKINL	AMTNAEWLEY	1200
AQTHLL						1206

SEQ ID NO: 122 moltype = AA length = 1233
 FEATURE Location/Qualifiers
 source 1..1233
 mol_type = protein
 note = Lachnospiraceae bacterium
 organism = unidentified

SEQUENCE: 122

-continued

MDYGNQGFER	RAPLTKTITL	RLKPIGETRE	TIREQKLLBQ	DAAPRKLVT	VTPIVDDCIR	60
KIADNALCHF	GTEYDFSCLG	NAISKNDKA	IKKETEKVEK	LLAKVLTEENL	PDGLRKVNDI	120
NSAAFIQDTL	TSPVQDDADK	RVLIQELKGG	TVLMQRFLLT	RITALTVWLP	DRVFENFNIF	180
IEENAEKMRIL	LDSPLNEKIM	KFDPDAEQYA	SLEFYQGCLS	QKDIDSYNLI	ISGIYADDEV	240
KNPGINEIVK	EYNQOIRGDK	DESPLPKLKK	LHKQILMPVE	KAFFVRVLSN	DSDARSILEK	300
ILKDTEMPLS	KIIEAMKEAD	AGDIAVYGSF	LHELSHVIYG	DHGKLSQIIY	DKESKRISL	360
MTLSPKERRK	ESKRLLEGLE	EHIRKSTYTF	DELNRVYAEK	VMAAYIAAVE	ESCAEIMRKE	420
KDLRTLSSKE	DVKIRGNRHN	TLIVKNYFNA	WTVFRNLIRI	LRRKSEAEID	SDFYDVLDD	480
VEVLSLTYKG	ENLCRSYITK	KIGSDLKPEI	ATYGSALRPN	SRWWSPEKEF	NVKFHTIVRR	540
DGRLYYFILP	KGAKPVELED	MDGDIECLQM	RKIPNPTIFL	PKLVFKDPEA	FFRDNPDAE	600
FVFLSGMKAP	VTITRETYEA	YRYKLYTVGK	LRDGEVSEEE	YKRALLQVLT	AYKEFLENRM	660
IYADLNFGFK	DLEEKSDSE	FIKQVETHNT	FMCWAKVSSS	QLDDLKSGN	GLLFEIWSER	720
LESYYKYGNE	KVLRGYEGVL	LSILKDENLV	SMRTLLNSRP	MLVYRPKESS	KPMVVHRDGS	780
RVVDRFPDKG	KYIPPEVHDE	LYRFFNNLLI	KEKLGEKARK	ILDNKKVKVK	VLESERVKWS	840
KFYDEQFAVT	FSVKNADCL	DTTKDLNAEV	MEQYSESNRL	ILIRNTTDLI	YYLVLDKNGK	900
VLKQSRSLNI	NDGARDVDWK	ERFRQVTKDR	NEGYNEWDYS	RTSNDLKVEY	LNVALKEIAE	960
AVIEYNAILI	IEKMSNAFKD	KYSFLDDVTF	KGFETKKLAK	LSDLHFRGIK	DGEPCSFSTNP	1020
LQLCQMSDNK	ILQDGVIFMV	PNSMTRSLDP	DTGFIPAIND	HNIRTKKAKL	NFLSKFDQLK	1080
VSEGGCLINM	YSGDSLVEPH	TGNRVWNCCC	NHPITNYDRE	TKKVEFIEEP	VEELSRVLEE	1140
NGIETDTLNL	KLNERENVPG	KVVDIAISLV	LNLYLRGTSG	VAGQRAVYYS	PVTGKKYDIS	1200
FIQAMNLRNK	CDYYRIGSKE	RGEWTFVQAQ	LIN			1233

SEQ ID NO: 123 moltype = AA length = 1227
 FEATURE Location/Qualifiers
 source 1..1227
 mol_type = protein
 note = Lachnospiraceae bacterium
 organism = unidentified

SEQUENCE: 123

MSKLEKFTNC	YLSKTLRFK	AIPVGKTQEN	IDNKRLLED	EKRAEDYKGV	KKLLDRYYLS	60
FINDVLHSIK	LKNLNNYISL	FRKKTREKE	NKELENLEIN	LRKEIAKAFK	GNEGYKSLFK	120
KDIIETILPE	FLDDKDEIAL	VNSFNGFTTA	FTGFFDNREN	MFSEEAKEST	IAFRGINENL	180
TRYISNMDF	EKVDAIFDKH	EVQEIKEKIL	NSDYDVEDFF	EGEFPNFVLT	QEGIDVYNAI	240
IGGFVTESE	KIKGLNEYIN	LYNQKTKQKL	PKPKPLYKQV	LSDRSLSFY	GEGYTSDEEV	300
LEVFRNTLNK	NSEIFSSIKK	LEKLKFNDE	YSSAGIFVKV	GPAISTISKD	IFGEWNVIRD	360
KWNAEYDDIH	LKKKAVVTEK	YEDDRKSPK	KIGSFSLEQL	QEYADADLSV	VEKLKEIIQ	420
KVDEIYKVG	SSEKLFDAF	VLEKSLKND	AVVAIMKDLL	DSVKSFENYI	KAFFGEGKET	480
NRDESFGDF	VLAYDILLKV	DHIYDAIRNY	VTQKPYSKDK	FKLYFQNPQF	MGGWDKDKET	540
DYRATILRYG	SKYYLAIMDK	KYAKCLQKID	KDDVNGNYEK	INYKLLPGPN	KMLPKVFFSK	600
KWMAYNPSSE	DIQKIYKNGT	FKKGDMMFNL	DCHKLIDFFK	DSISRYPKWS	NAYDFNFSST	660
EKYKDIAGPY	REVEEQGYKV	SPESASKKEV	DKLVEEGKLY	MFQIYNKDFS	DKSHGTPNLH	720
TMFYKLLPDE	NNHGQIRLSG	GAEFLMRRAS	LKKEELVVHP	ANSPIANKNP	DNPKKTTTSL	780
YDVYKDKRFS	BDQYELHPI	ANINKCPKNI	FKINTEVRVL	LKHDDNPYVI	GIDRGERNLL	840
YIVVVDGKGN	IVEQYSLNEI	INNFMGIRIK	TDYHSLDDK	EKERFEARQN	WTSIENIKEL	900
KAGYISQVHV	KICELVEKYD	AVIALEDLNS	GFKNRSRVKE	KQVYQKFEKM	LIDKLNYMVD	960
KKSNPCATGG	ALKGYQITNK	FESFKSMSTQ	NGFIFYIPAW	LTSKIDPSTG	FVNLLKTKYT	1020
SIADKKFISS	FDRIMYVPEE	DLFEFALDYK	NFSRTDADYI	KKWKLYSYGN	RIRIFRNPCK	1080
NNVFDWEVC	LTSAYKELFN	KYGINYQQGD	IRALLCEQSD	KAFYSSPMAL	MSLMLQMRNS	1140
ITGRTDVFDF	ISPVKNSDGI	FYDSRNYEAQ	ENAILPKNAD	ANGAYNIARK	VLWAIQGFKK	1200
AEDEKLKDKV	IASNKEWLEY	AQTSVKKH				1227

SEQ ID NO: 124 moltype = AA length = 1264
 FEATURE Location/Qualifiers
 source 1..1264
 mol_type = protein
 note = Lachnospiraceae bacterium
 organism = unidentified

SEQUENCE: 124

MEDYSGFVNI	YSIQKTLRFE	LKPVGKTLEH	IEKKGFLKDD	KIRAEDYKAV	KKIIDKYHRA	60
YIEEVFDSVL	HQKKKKDKTR	FSTQFIKEIK	EFSELYYKTE	KNIPDKERLE	ALSEKLRKML	120
VGAFKGEFSE	EVAEKYNKNL	FSKELIRNEI	EKFCETDEER	KQVSNFKSFT	TYFTGFHSNR	180
QNIYSDEKKS	TAIGYRIIHQ	NLPKFLDNLK	IIESIQRREF	DFFWSDLKKN	LKKIDKNIKL	240
TEYFSIDGFV	NVLNQKGIDA	YNTILGGKSE	ESGEKIQGLN	EYINLYRQKN	NIDRKNPLNV	300
KILFKQILGD	RETSKFIPEA	FPDDQSVLNS	ITEFAKYLLK	DKKKKSIIAE	LKKFLSSFNR	360
YELDGILYLN	DNLSASISTF	LFDDWSFIKK	SVSFKYDESV	GDPKKKIKSP	LKYEKEKEKW	420
LKQKYTTISF	LNDAESYSK	SQDEKRVKIR	LEAYFAEFKS	KDDAKKQFDL	LERIEEAYAI	480
VEPLLGAEPY	RDRNLKADKK	EVGKIKDFLD	SIKSLQFFLK	PLLSAEIFDE	KDLGFYNQLE	540
GYYEEDISG	HLYNKVRNYI	TGKIYSKEKF	KLNFENSTLL	KGWDENREVA	NLCVIFREDQ	600
KYYLGVMDEK	NNTILSDIPK	VKPNELFYEK	MVYKLIPTPH	MQLPRIIFSS	DNLISIYNPSK	660
SILKIREAKS	FKEGKNFKLK	DCHKFIDFYK	ESISKNEWDS	RDFPKFSKTS	SYENISEFYR	720
EVERQQYNLD	FKKYSKFYID	SLVEDGKLYL	FQIYNKDFS	FSKKGKPNLHT	IYFRSLFSKE	780
NLKDVCCLKN	GEAEMFFRKK	SINYDEKKKR	EGHHPPELFEK	LKYPILKDKR	YSEDKQPFHL	840
PISLNFKSKE	RLNPNLKVNE	FLKRNKDINI	IGIDRGERNL	LYLVMINQKG	EILKQTLDDS	900
MQSGKGRPEI	NYKEKLQKEK	IERDKARKSW	GTVENIKELK	EGYLSIVIHQ	ISKLMVENNA	960
IVLEDLNIG	FKRGRQKVER	QVYQKFEKML	IDKLNFLVFK	ENKPTPEGGV	LKAYQLTDEF	1020
QSFEKLKSKT	GFLFYVPSWN	TSKIDPRTGF	IDFLHPAYEN	IEKAKQWINK	FDSIRFNSKM	1080

-continued

DWFEPTADTR	KFSENMLGK	NRVWVICTTN	VERYFTSKTA	NSSIQYNSIQ	ITEKLKELFV	1140
DIPFSNGQDL	KPEILRKND	VFFKSLLFYI	KTTLRLQNN	GKKGEEKDF	ILSPVVDKSG	1200
RFFNSLEASD	DEPKDADANG	AYHIALKGLM	NLLVLNETKE	ENLSRPKWKI	KNKDWLEFVW	1260
ERNR						1264

SEQ ID NO: 125 moltype = AA length = 1373
 FEATURE Location/Qualifiers
 source 1..1373
 mol_type = protein
 organism = Moraxella bovoculi

SEQUENCE: 125

MLFQDFTHL	PLSKTVRFEL	FIDRTLEHII	AKNFLSQDET	MADMHQKV	ILDDYHRDFI	60
ADMMGEVKLT	KLAEFYDVYL	KFRKNPKDDE	LQKAQLKDLQ	AVLRKEIVKP	IGNGGKYKAG	120
YDRLFGAKLF	KDGKELGDLA	KFVIAQEGES	SPKLAHLAHF	EKFSTYFTGF	HDNRKNMYS	180
EDKHTAIAYR	LIHENLPRFI	DNLQILTTIK	QKHSALYDQI	INELTASGLD	VSLASHLDGY	240
HKLLTQEGIT	AYNTLLGGIS	GEAGSPKIQG	INELINSHHN	QHCHKSERIA	KLRPLHKQIL	300
SDGMSVSFLP	SKFADDSEMC	QAVNEFYRHY	ADVFAKVQSL	FDGFDHQQD	GIYVEHKNLN	360
ELSKQAFGDF	ALLGRVLDGY	YVDVNVPEFN	ERFAKAKTDN	AKAKLTKEKD	KFIKGVHSLA	420
SLEQAIEHYT	ARHDDSEVQA	GKLGQYFKHG	LAGVDNPIQK	IHNHSTIKG	FLERERPAGE	480
RALPKIKSGK	NPEMTQLRQL	KELLDNALNV	AHFAKLLTTK	TTLDNQDGNF	YGEFGVLYDE	540
LAKIPTLYNK	VRDYSQKPF	STEKYKLNFG	NPTLLNGWDL	NKEKDNFGVI	LQKDGCCYLA	600
LLDKAHKKVF	DNAPTNGKSI	YQKMIYKYLE	VRKQFPKVFF	SKEAIAINYH	PSKELVEIKD	660
KGRQRSDDER	LKLYRFILEC	LKIHPKYDKK	FEGAIGDIQL	FKKDKKGREV	PISEKDLFKD	720
INGIFSSKPK	LEMEDFFIGE	FKRYNPSQDL	VDQYNIYKKI	DSNDNRKKEN	FYNNHPKFKK	780
DLVRYYESM	CKHEEWESF	EFSKKLQDIG	CYVDVNELEF	EIETRRRLNYK	ISFCINADY	840
IDELVEQQQL	YLPQIYNKDF	SPKAHGKPNL	HTLYPKALFS	EDNLADPIYK	LNGEAQIFYR	900
KASLDMNETT	IHRAGEVLEN	KNPDNPKKRQ	FVYDIIDKDR	YTQKDFMLHV	PITMNFQVQG	960
MTIKEFNKKV	NQSIQQYDEV	NVIGIDRGER	HLLYLTVINS	KGEILEQCSL	NDITTASANG	1020
TQMTTPYHKI	LDKREIERLN	ARVGWGEIET	IKELKSGYLS	HVVHQISQLM	LKYNAIVVLE	1080
DLNFGFKRGR	FKVEKQIYQN	FENALIKKLN	HLVLKDKADD	EIGSYKNALQ	LNNFTDLKS	1140
IGKQTGLFVY	VPAWNTGKID	PETGFVDLLK	PRYENIQASQ	APFGKFDKIC	YNADKYFEF	1200
HIDYAKFTDK	AKNSRQIWTI	CSHGDKRYVY	DKTANQNKA	AKGINVNDIL	KSLFARHHIN	1260
EKQPNLVMID	CQNNDKEFHK	SLMYLLKTL	ALRYSNASSD	EDFILSPVAN	DEGVFPNSAL	1320
ADDTQPNAD	ANGAYHIALK	GLWLLNELKN	SDDLNVKCLA	IDNQTWLNFA	QNR	1373

SEQ ID NO: 126 moltype = AA length = 1352
 FEATURE Location/Qualifiers
 source 1..1352
 mol_type = protein
 note = Paracubacteria bacterium
 organism = unidentified

SEQUENCE: 126

MENIFDQFIG	KYSLSKTLRF	ELKPVGKTED	FLKINKVFEK	DQTIDDSYNQ	AKFYFDSLHQ	60
KFIDAAALASD	KTSLSFQNF	ADVLEKQNKI	ILDKKREMG	LKRKDKNAV	IDRLQKEIND	120
AEDITQKEKE	KIYKDVRTL	DNEAESWKTY	YQREVDGKK	ITESKADLKQ	KGADFLTAAG	180
ILKVLKYEF	EEKEKEFOAK	NQPSLFVEEK	ENPGQKRYIF	DSFDKFAGYL	TKFQQTKKNL	240
YAADGTSTAV	ATRIADNFII	FHONTKVFRD	KYKNNHTDLG	FDEENIFEIE	RYKNCLLQRE	300
IEHIKNENSY	NKIIGRINKK	IKERYDQKAK	DTKLTKSDFP	FFKNLDKQIL	GEVEKEKQLI	360
EKTREKTED	VLIERFKEFI	ENNEERFTAA	KKLMNAFCNG	EFSEYEGIY	LKNKAINTIS	420
RRWFVSDRDF	ELKLPQQKSK	NKSEKNPKV	KKFISIAEIK	NAVEELDGI	FKAVFYDKKI	480
IAQGGSKLEQ	FLVWKYEF	YLPFDIEREN	GEKLLGYDSC	LKIAKQLGIF	PQKEAREKA	540
TAVIKNYADA	GLGIFQMMKY	FSLDDKDRKN	TPGQLSTNFI	AEYDGYKDF	EFIKYNEFR	600
NFITKKPFDE	DKIKLNFENG	ALLKGWDENK	EYDFMGVILK	KEGRLYLGIM	HKNHRKLFQS	660
MGNAGKDNAN	RYQKMIYKQI	ADASKDVPR	LLTSKKAMEK	FKPSQEILRI	KKEKTFKRES	720
KNFSLRLDHA	LIEYRNCIP	QYSNWSFYDF	QFQDTGKYQN	IKEPTDDVQK	YGYKISFRDI	780
DDEYINQALN	EGKMYLFEV	NKDIYNTKNG	SKNLHTLYFE	HILSAENLND	PVFKLSGMAE	840
IFQRQPSVNE	REKITTQKNQ	CILDKGDRAY	KYRRYTEKKI	MPHMSLVLNT	GKGEIKQVQF	900
NKLIINQRISS	SDNEMRVNVI	GIDRGEKNLL	YYSVVKQNGE	IIEQASLNEI	NGVNYRDKLI	960
EREKERLKNR	QSWKPVVKIK	DLKKGYISHV	IHKICQLIEK	YSAIVVLEDL	NMRFKQIRGG	1020
IERSVYQQFE	KALIDKLGYL	VFKDNDRDLA	PGGVNLNGYQ	SAPFVSPEKM	RKQGTGILFYT	1080
QAEYTSKTD	ITGFRKNVYI	SNSASLDKIK	EAVKKFDAIG	WDGKEQSYFF	KYNPNYLADE	1140
KYKNSTVSKE	WAIFASAPRI	RRQKGEDGYW	KYDRVKVNEE	FEKLLKVWNF	VNPKATDIKQ	1200
EIIKKIKAGD	LQGEKELDGR	LRNFWHSFIY	LFNLVLELFR	SFSLQIKIKA	GEVIAVDEGV	1260
DFIASPVKPF	FTTPNPYIPS	NLCWLAVENA	DANGAYNIAR	KGVMILKKIR	EHAKKDFEKF	1320
KLPNLFISNA	EWDEAARDWG	KYAGTTALNL	DH			1352

SEQ ID NO: 127 moltype = AA length = 1260
 FEATURE Location/Qualifiers
 source 1..1260
 mol_type = protein
 organism = Porphyromonas crevioricanis

SEQUENCE: 127

MDSLKDFTNL	YPVSKTLRFE	LKPVGKTLEN	IEKAGILKED	EHRAESYRRV	KKIIDTYHKV	60
FIDSSLENMA	KMGIENEIKA	MLQSFCELYK	KDHRTEGEDK	ALDKIRAVLR	GLIVGAFTGV	120
CGRRENTVQN	KYESLPEKEK	LIKEILPDFV	LSTEAESELPF	SVEEATRSLE	EPDSFTSYFA	180
GFYENRKNYI	STKPQSTAI	YRLIHENLPK	FIDNILVFQK	IKEPIAKELE	HIRADFSAGG	240

-continued

YIKKDERLED	IFSLNYYIHV	LSQAGIEKYN	ALIGKIVTEG	DGEMKGLNEH	INLYNQQRGR	300
EDRLPLFRPL	YKQILSDREQ	LSYLPESFEK	DEELLRALKE	FYDHI AEDIL	GRTQQLMTSI	360
SEYDLSRIYV	RNDSQLTDIS	KKMLGDWNAI	YMARERAYDH	EQAPKRITAK	YERDRKALK	420
GEESISLANL	NSCIAFLDNV	RDCRVDTYLS	TLGQKEGPHG	LSNLVENVFA	SYHEAEQLLS	480
FPYPEENLI	QDKDNVVLK	NLLDNISDLQ	RFLKPLWGMG	DEPKDERFY	GEYNYIRGAL	540
DQVIFLYNKV	RNYLTRPKYS	TRKVKLNFGN	SQLLSGWDNR	KEKDNSCVIL	RKGQNFYLAI	600
MNNRHKRSFE	NKMLPEYKEG	EPYFEKMDYK	FLPDPNKMLP	KVFLSKKGIE	IYKPSPKLLE	660
QYGHGTHKKG	DTFSMDLHE	LIDFFKHSIE	AHEDWKQGF	KFSDTATYEN	VSSFYREVED	720
QGYKLSFRKV	SESYVYSLID	QGKLYLFQIY	NKDFSPCSKG	TPNLHTLYWR	MLFDERNLAD	780
VIYKLDGKAE	IPFREKSLKN	DHPHTPAGKP	IKKKSQKKKG	EESLFEYDLV	KDRRYTMDKF	840
QPHVPI TMNF	KCSAGSKVND	MVNAHIREAK	DMHVIGIDRG	ERNLLYICVI	DSRGITLDQI	900
SLNTINDIDY	HDLLERSDKD	RQQEHRNWQT	IEGIKELKQG	YLSQAVHRIA	ELMVAYKAVV	960
ALEDLNMGGF	RGRQKVESV	YQFEKQLID	KLNYLVDKKK	RPEDIGLLLR	AYQFTAPFKS	1020
FKEMGKQNGF	LFYIPAWNTS	NIDPTTGFEV	LFHVQYENVD	KAKSFFQKFD	SISYNPKKDW	1080
FEFAPDYKNF	TKKAEGSRSM	WILCTHGSRI	KNFRNSQKNG	QWDSEEFALT	EAFKSLFVRY	1140
EIDYTADLKT	AIVDEKQKDF	FVDLLKLFLK	TVQMRNSWKE	KDLDYILSPV	AGADGRFFDT	1200
REGNKS LPKD	ADANGAYNIA	LKGLWALRQI	RQTSEGGKLG	LAISNKEWLQ	FVQERSYEKD	1260

SEQ ID NO: 128 moltype = AA length = 1324
 FEATURE Location/Qualifiers
 source 1..1324
 mol_type = protein
 organism = Prevotella disiens

SEQUENCE: 128

MENYQEFNTL	FQLNKT LRFE	LKPIGKTCEL	LEEGKIFASG	SFLEKDKVRA	DNVSYVKKEI	60
DKKHKIFIEE	TLSSFSISND	LLKQYFDCYN	ELKAPKKDCK	SDEEEVKKTA	LRNKCTSIQR	120
AMREAI SQAF	LKSPQKLLA	IKNLIENVFK	ADENVQHFE	FTSYFSGFET	NRENPFYSDEE	180
KSTSIAYRLV	HDNLPIFIKN	IYIFEKLKEQ	FDAKTLSEIF	ENYKLYVAGS	SLDEVFSLEY	240
FNNLTLTQKI	DNYNNAVIGI	VKEDKQEIQG	LNEHINLYNQ	KHKDRRLPPF	ISLKKQILSD	300
REALSWLPDM	FKNDSEVIDA	LKGFYIEDGF	ENNVLTPLAT	LLSSLDKYNL	NGIFIRNNEA	360
LSSLSQNVYR	NFSIDEAIDA	QNAELQTFNN	YELIANALRA	KIKKETKQGR	KSFKEYEYEI	420
DKKVK AIDSL	SIQEINELVE	NYVSEFNSNS	GNMPRKVEDY	FSLMRKGDGF	SNDLIENIKT	480
KLAAAEKLLG	TKYQETAKDI	FKKDENSCLI	KELLDATKQF	QHFYKPLLGT	GEEADRDLVF	540
YGDFLPLYEK	FEELTLLYNK	VRNRLTQKPY	SKDKIRLCFN	KPKLMTGWVD	SKTEKSDNGT	600
QYGGYLFRRK	NEIGEYDYFL	GISSKAQLFR	KNEAVIGDYE	RLDYYQPKAN	TIYGSAYEGE	660
NSYKEDKKRL	NKVIIAYIEQ	IKQTNIKKSI	IESISKYPNI	SDDDKVTPSS	LLEKIKKSVI	720
DSYNGILSFK	SFQSVNKEVI	DNLLKTISPL	KNKAEFLDLI	NKDYQIFTEV	QAVIDEICKQ	780
KTFIYFPISN	VELEKEMGDK	DKPLCLFQIS	NKDLSFAKTF	SANLRKKRGA	ENLHTMLFKA	840
LMEGHQDNL	LGSGAIFYRA	KSLDGNKPTH	PANEAIKCRN	VANKDKVSLF	TYDIYKNRRY	900
MENKFLPHLS	IVQNYKAAND	SQALNSSATE	YIRKADDLHI	IGIDRGERNL	LYYSVIDMKG	960
NIVEQDSLNI	IRNNDLETDY	HDLLDKREKE	RKANRQNWEA	VEGIDKLKKG	YLSQAVHQIA	1020
QLMLKYNAL	ALEDLGGMFV	TRGQKIEKAV	YQQFEKSLVD	KLSYLVDDKR	PYNELGGILK	1080
AYQLASSITK	NNSDKQNGFL	FYVPAWNTSK	IDPVTGFTDL	LRPKAMTIKE	AQDFFGAFDN	1140
ISYNDKGYFE	FETNYDKFKI	RMKSAQTRWT	ICTFGNRIKR	KKDKNYWNYE	EVELTEEFKK	1200
LPKDSNIDYE	NCNLKEEIQN	KDNRKFFDDL	IKLLQLTLQM	RNSDDKGN DY	IISP VANAEG	1260
QFFDSRNGDK	KLPLDADANG	AYNIARKGLW	NIRQIKQTKN	KDDLNLNLISS	TEWLDFVREK	1320
PYLK						1324

SEQ ID NO: 129 moltype = AA length = 1484
 FEATURE Location/Qualifiers
 VARIANT 1073
 note = Xaa can be any naturally occurring amino acid
 source 1..1484
 mol_type = protein
 note = Peregrinibacteria bacterium
 organism = unidentified

SEQUENCE: 129

MSNFFKNFTN	LYELSKTLRF	ELKPVGDTLT	NMKDHLEYDE	KLQTF LKQDN	IDDAYQALKP	60
QFDEIHEEPI	TDSLESKKAK	EIDFSEYLDL	FOEKKELNDS	EKKLRNKIGE	TFNKAGEKWK	120
KEKYPQYEWK	KGSKIANGAD	ILSCQDMLQF	IKYKNPEDEK	IKNYIDDTLK	GFFTYFGGFN	180
QNRANYEYETK	KEASTAVATR	IVHENLPKFC	DNVIQFKHII	KRKKGDTVEK	TERKTEYLNA	240
YQYLKNNNKI	TQIKDAETEK	MIESTPIAEK	IPDVYFSSC	LSQKQIEEYN	RIIGHYNLLI	300
NLYNQAKRSE	GKHLSANEKK	YKDLPKFKTL	YKQIGCGKKK	DLFYTIKCDT	EEEAANKSRNE	360
GKESHSEVEI	LKQAQEAINK	YFKNNDNEN	INTVPDFINY	ILTKENYEGV	YWSKAAMNTI	420
SDKYFANYHD	IQDRLLKEAKV	FQKADKKSSED	DIKIPAEIEL	SGLFGVLDL	ADWQTTLFKS	480
SILSNEKLKI	ITDSQTPSEA	LLKMI FNDIE	KNMESFLKET	NDIITLKKYK	GNKEGTEKIK	540
QWFDYTLAIN	RMLKYFLVKE	NKIKGNSLDT	NISEALKTLI	YSDDAEWFKW	YDALRNYLTQ	600
KPQDEAKENK	LKLNDFNPSL	AGGWDVNKEC	SNFCVILKDK	NEKYLAMIK	KGENTLFOKE	660
WTEGRGKNLT	KKSNPLFEIN	NCEILSKMEY	DFWADVSKMI	PKCSTQLKAV	VNHFKQSDNE	720
FIFPIGYKVY	SGEKFRECEK	ISKQDFELNN	KVFNKNELSV	TAMRYDLSST	QEQYIKAFQ	780
KEYWELLFPK	EKRDTKL TNN	EIFNEWINFC	NKKYSELLSW	ERKYKDALTN	WINFCYFVLS	840
KYPKTTLFNY	SFKESENYNS	LDEFYRDVDI	CSYKLNINTT	INKSILDRLV	EEGKLYLFEI	900
KNQDSNDGKS	IGHKNNLHTI	YWN AIFENFD	NRPKLNGEAE	IFYRKAISKD	KLGI VKGKKT	960
KNGTWIKYNY	RFSKEKFILH	VPITLNFCSN	NEYVNDIVMT	KFYNFNSLHF	LGIDRGEKHL	1020
AYSLVNKNG	EIVDQGTLLN	PFTDKDGNQR	SIKKEKYFYN	KQEDKWEAKE	VDXWNYNDLL	1080
DAMASNRDMA	RKNWQRI GTI	KEAKNGYVSL	VIRKIADLAV	NNERPAPIVL	EDLNTGFKRS	1140

-continued

RQKIDKSVYQ	KFELALAKKL	NFLVDKNAKR	DEIGSPTKAL	QLTPPVNNYG	DIENKKQAGI	1200
MLYTRANYTS	QTDPATGWRK	TIYLKAGPEE	TTYKKDGKIK	NKSVKDQIIE	TFTDIGFDGK	1260
DYYFEYDKGE	FVDEKTGEIK	PKKWRLYSGE	NGKSLDRFRG	EREKDKYEWK	IDKIDIVKIL	1320
DDLFDVNFNFKN	ISLLKQLKEG	VELTRNNEHG	TGESLRFAIN	LIQQIRNTGN	NERDNDFILS	1380
PVRDENGKHF	DSREYWDKET	KGEKISMPSS	GDANGAFNIA	RKGIIMNAHI	LANSDSKDLS	1440
LFVSDEEWDL	HLNNKTEWKK	QLNIFSSRKA	MAKRKKKRP	ATKK		1484

SEQ ID NO: 130 moltype = AA length = 1245
 FEATURE Location/Qualifiers
 source 1..1245
 mol_type = protein
 organism = Porphyromonas macacae

SEQUENCE: 130

MKTQHFFEDF	TSLSLSKTI	RFELKPIGKT	LENIKKNGLI	RRDEQRLLDDY	EKLKKVIDEY	60
HEDFIANILS	SFSFSEELQ	SYIQNLSISE	ARAKIEKTMR	DTLAKAFSED	ERYKSIFKKE	120
LVKIDIPVWC	PAYKSLCKFF	DNFTTSLVPF	HENRKNLYTS	NEITASIPYR	IVHVNLPKFI	180
QNIERALCELQ	KKMADLYLE	MMENLRNVWP	SVFKTPDDLC	NLKTYNHLMV	QSSISEYNRF	240
VGGYSTEDGT	KHQGINWIN	IYRQRNKEMR	LPGLVFLHKQ	ILAKVDSSSF	ISDTLENDQ	300
VPCVLRQFRK	LFWNTVSSKE	DDAASLKDLF	CGLSGYDPEA	IYVSDAHLAT	ISKNIIDRW	360
YISDAIRRKT	EVLMPRKES	VERYAEKISK	QIKKRQSYSL	AELDDLLAHY	SEESLPAGFS	420
LLSYFTSLGG	QKYLVSDEGEV	ILYEEGSIW	DEVLIAFRDL	QVILDKDFTE	KKLGKDEEAV	480
SVIKKALDSA	LRLRKFFDLL	SGTGAEIRRD	SSFYALYTDR	MDKLKGLLKM	YDKVRNYLTK	540
KPYSIEKFKL	HFDNPSLLSG	WDKNKELNNL	SVIFRQNGYY	YLGIMTPKGG	NLFKTLPLKLG	600
AEEMFYEKME	YKQIAEPLML	LPKVFFPKKT	KPAPAPDQSV	VDIYNKTKFK	TGQKGFNKKD	660
LYRLIDFYKE	ALTVEHWKLF	NFSFSPTEQY	RNIGEFFDEV	REQAYKVMV	NVPASYIDEA	720
VENGKLYLFP	IYNKDFSPYS	KGIPNLHTLY	WKALFSEQNQ	SRVYKLCGGG	ELFYRKASLH	780
MQDTTVHPKG	ISIHKKNLNK	KGTSLFNYD	LVKDKRFTED	KFFPHVPIIS	NYKNKKITNV	840
NQMVRDYIAQ	NDDLQHGIDR	GERNLVYISR	IDTRGNLLEQ	FSLNVIESDK	GDLRTDYQKI	900
LGDRERQERL	RRQEWKSIES	IKDLKDGYS	QVVKICNMV	VEHKAIVVLE	NLNLSPMKGR	960
KKVEKSVYKE	FERMLVDKLN	YLVVDKKNLS	NEPGGLYAA	QLTNPLFSFE	ELHRYPQSGI	1020
LFVDPFWNTS	LTDPSTGFVN	LLGRINYTNV	GDARKEFFDR	NAIRYDGGKN	ILFDLDSRF	1080
DVRVETQRKL	WTLTTFGSRI	AKSKKSGKWM	VERIENLSLC	FLELFEQFNI	GYRVEKDLKK	1140
ATLSQDRKEF	YVRLIYLFNL	MMQIRNSDGE	EDYILSPALN	EKNLQFDSRL	IEAKDLPVDA	1200
DANGAYNVAR	KGLMVVQRIK	RGDHESIHR	GRAQWLRYVQ	EGIVE		1245

SEQ ID NO: 131 moltype = AA length = 1250
 FEATURE Location/Qualifiers
 source 1..1250
 mol_type = protein
 organism = Smithella sp.

SEQUENCE: 131

MQTLFENFTN	QYPVSKTLRF	ELIPQGGTKD	FIEQKGLLKK	DEDRAEKYKK	VKNIIDEYHK	60
DFIEKSLNGL	KLDGLEKYKT	LYLKQEKDDK	DKKAPDKEKE	NLRKQIANAF	RNNEKFKTLF	120
AKELIKNDLM	SFACEEDKKN	VKEFEAFTTY	FTGPHQNRAN	MYVADEKRTA	IASRLIHENL	180
PKFIDNIKIF	EKMKEAPEL	LSPFNQTLKD	MKDVIKGTTL	EEIFSLDYFN	KTLTQSGIDI	240
YNSVIGGRTP	BEGKTKIKGL	NEYINTDFNQ	KQTDKKKRQP	KFKQLYKQIL	SDRQSLSFIA	300
EAFKNDTEIL	EAIEKFYVNE	LLHFSNEGKS	TNVLDAIKNA	VSNLESFNLT	KMYFRSGASL	360
TDVSRKVFGE	WSIINRALDN	YYATTYPIKP	REKSEKYEER	KEKWLKQDFN	VSLIQTATIDE	420
YDNETVKGKN	SGKVIADYFA	KFCDDKETDL	IQKVNIEGYIA	VKDLLNTPCP	ENEKLGSKND	480
QVKQIKAFMD	SIMDIMHFVR	PLSLKDTDK	KDETFYSLFT	PLYDHLTQTI	ALYNKVRNYL	540
TQKPYSTEKI	KLNFNSTLL	GGWDLNKETD	NTAIIILKDN	LYYLIGIMDKR	HNRIFRNVPK	600
ADKDFCEYKE	MVYKLLPGAN	KMLPKVFFSQ	SRIQEFTPSA	KLLENYANET	HKKGDNFNLN	660
HCHKLIDFFK	DSINKHEDWK	NFDFRFSATS	TYADLSGFYH	EVEHQGYKIS	FQSVADSFID	720
DLVNEGKLYL	FQIYNKDFSP	FSKGKPNLHT	LYWKMLFDEN	NLKDVVYKLN	GEAEVFPYRK	780
SIAEKNTTIIH	KANESIINKN	PDNPKATSTF	NYDIVKDKRY	TIDKFQPHIP	ITMNFKAEGI	840
FMNQVRVNOF	LKANPDINII	GIDRGERHLL	YYALINQKGG	ILKQDTLNV	ANERQKVDYH	900
NLLDKKEGDR	ATARQEWGVI	ETIKELKEGY	LSQVIHKLTD	LMIENNAIIV	MEDLNFPGFKR	960
GRQKVEKQVY	QKFEKMLIDK	LNYLVDKNKK	ANELGGLLNA	FQLANKFESF	QKMGKQNGFI	1020
FYVPAWNTSK	TDPATGFIDF	LKPRYENLNQ	AKDPFEKFPDS	IRLNSKADYF	EFAFDPKNFT	1080
EKADGGRTKW	TVCTTNEEDRY	QWNRALNNNR	GSQEKYDITA	ELKSLFDGKV	DYKSGKDLKQ	1140
QIASQESADF	FKALMKNLIS	TLSLRHNGGE	KGDNEQDYIL	SPVADSKGRF	FDSRKADDDM	1200
PKNADANGAY	HIALKGLWCL	EQISKTDLLK	KVKLAISNKE	WLEFVQTLKG		1250

SEQ ID NO: 132 moltype = DNA length = 3987
 FEATURE Location/Qualifiers
 source 1..3987
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 132

atggccgggga	gcaagaagcg	cgggataaag	caggacacgc	agttcgaggg	cttcaccaac	60
ctgtaccaag	tctccaagac	gctccgggtc	gagcttatcc	cgcaaggga	gaccctgaaa	120
cacatccagg	aacaaggttt	catcgaggag	gacaaggccc	gcaacgacca	ctacacggag	180
ctcaagccca	taatcgatcg	gatctacaag	acgtacgccc	accagtgcct	ccaactgggtg	240
cagctcgact	gggagaacct	gagcgccgcc	attgacagct	accgcaagga	aaagacggag	300
gagacgcgca	acgcccctat	tgaggagcaa	gccacctacc	gcaacgccat	ccacgactac	360
ttcatcgggc	gcaccgacaa	cctgacggac	gcgatcaaca	agcgccacgc	ggaatctac	420

-continued

```

aagggccttt tcaaggccga gctcttcaac gggaagggtcc taaaacagct cgggactgtc 480
acgacaaccg agcatgagaa cgccctcctt cgcagcttcg acaagttcac cacatacttc 540
tcgggcttct accggaaccg caagaacgtt ttcagcgccg aggacatctc caccgccatc 600
ccgcacagga tcgtccagga caacttcccc aagttcaagg agaactgcca catcttcacg 660
cgccgtgatta cagccgtacc ttcaacttcgt gagcacttcg agaacgtcaa aaaggccatc 720
gggactcttcg tctccaagtc catcgaggag gtattctctt tcccgttcta taaccagctc 780
ctgacccaga cgcagatcga cctctacaac cagctactgg gcggcatcag ccgggaggcc 840
gggaccgaga aaataaaggg cctcaacgaa gttctcaacc tggccatcca gaagaacgac 900
gagaccgcgc atatcatcgc atccctgcgc catcgcttca ttcctttgtt caagcagata 960
ttgagcgacc ggaacacctt ctcggttcac ctgaagaat tcaagagcga cgaggagggtc 1020
attcagctctt tctgcaagta caagacgctc ctacggaatg agaattgtgt ggagaccgcg 1080
gaggcactct tcaatgagct gaactccatt gacctgacct acatcttcat tagccacaag 1140
aaactggaga cgtctctccg cgccctgtgc gacctctggg acactctccg caacgccttc 1200
tacgaacgcc ggatctccga acttacgcgc aagataacta agtcggctaa ggagaagggtg 1260
caacggagcc tcaagcagca ggacatcaac cttcaggaaa tcatctcagc cgcgggcaag 1320
gagctgagcg aggcgtttta gcagaaaaca tcggagatag tgagccacgc gcacgcggcc 1380
ctggatcaac cgtctgcgag gactctcaag aagcaagagg agaaggaaat ccttaagtcc 1440
cagctcgact cgtctgctcg cctctatcac ttgctcgact ggttcgcggt tgatgagtcc 1500
aacgaggtgg acccgaggtt ctccgcgcgc ctacgggta ttaagctgga gatggagcca 1560
agcttaagct tctacaacaa ggcccgcaac tacgcgacca aaaaaccgta ctcagtcgag 1620
aaatcaagc tgaatttcca gatgcctaca ttggcgaggg ggtgggagct gaaccgcgag 1680
aagaacaatg gagccatcct gttctgcaaa aatgggttgt actacctggg catcatgcc 1740
aagcagaagg gccgtttaca ggccctgtca ttcgagccta ccgagaagac ctcgaggggc 1800
ttcgacaaga tgtactacga ctatttcccg gacgcgcgca agatgatccc gaagtgtccc 1860
acgcagctca tagccgtcac ggccacttc cagacgcata ccacgccgat actctgagc 1920
aacaacttca ttgagcgct agagatcacg aaggagatat acgacctaaa caaccccgaa 1980
aaggagccca agaagttcca gacagcctac gctaagaaga caggtgatca gaagggat 2040
agggaggcac tctgcaagtg tatcgacttc acgcgcgact tctgtcgaa atatacaaa 2100
acgaccagca ttgacctaa gttctctccg ccatctctcc agtacaagga tctgggagag 2160
tattatcgcg agctgaaccc attgctgtac cacatcagct tccagaggat cgcgcgaga 2220
gagattatgg acgcggtgga gacggggaaa ctatacctgt tccaaatata taacaaggac 2280
ttcgtaaaag ggcaaccagg gaagcccaac ctgcacacac tctactggac gggcttgtt 2340
tcgcagaaaa atttggccaa gacttcgatc aagctcaacg gccaggcgga gttgttttac 2400
cgtcccaagt ctccgatgaa gcgcgatggc gatcgcttcg gagagaaaat gcttaacaag 2460
aagctcaagg atcagaagac gcccatacct gatacgttgt accaggaat gtacgactac 2520
gtgaaccacc gccctacgca cgacctctca gacgaggccc gcgcctcct cccaaacgtg 2580
attactaagg aggtttccca tgaataaatc aaggaccgac ggttaccag cgacaattt 2640
ttttccacg tgccctatcac gctcaattac caggcgccca actcccatc gaagttcaac 2700
cagcgctgta acgcctacct taaggagcac ccggagaccc caatcatcgg gatcgacct 2760
ggcgagcgga acctgatcta tattacggtg atcgatagca ccgggaagat cctggagcag 2820
cgctccctga acacaatcca gcagtttgac taccagaaga aactcgacaa ccgggagaa 2880
gagcgctcg cagcccgcca agcatggagt gtggtcgga ccataaagga cctgaaacag 2940
ggttacctaa gtcaagttat ccacgagatc gttgacctga tgatacacta tcaagccgta 3000
gtcgtgctgg agaacctcaa cttcggtttt aagttcaacg gcaccggcat cgcggagaag 3060
gcggtgtacc cgcagttcga gaagatgctg atcgacaagc tgaactgcct ggtgtcaca 3120
gactaccctg aggagaaggt cgccggggtc ttgaacccgt accagtaac cgaccagttc 3180
acgagcttcg ccaaaatggg cagcagttcc ggattctgtt tttatgtccc ggtccatat 3240
acaagtaaga tcgaccgctg gacagggttt gttgacctat cgtgtggaa gaccatcaag 3300
aaccacgaga gcaggaaaaca cttctagag ggcttcgact tctcgatta cgacgttaag 3360
acaggcgact tcactctgca cttcaagatg aaccgcaacc tgtcgttcca gaggggctcg 3420
cccgcttca tgcccgctg ggatatcgtc tttgagaaga atgagacgca gttcgacgcg 3480
aaagggagcg cgttcactgc tgaacccgag atcgtgcggg tcatcgagaa ccaccgctc 3540
acgggtcgct accgagatgt ataccccgcc aacgaactaa ttgcgctgct ggaggagaag 3600
gggatcgctg tccgagatgg cagcaacatt ctcccgaagc tgctggagaa cgacgactcg 3660
cacgctattg acacgatggg cgccctcata cggagcgtgc ttcagatgcg gaacagtaac 3720
gctgccacgg gcgaggacta cattaactcc ccggtccgag acctcaacgg ggtctgcttc 3780
gatagccgct tccagaaccc ggagtgccct atggatgcgg acgcgaacgg ggccctaccac 3840
atcgccctca agggccaact cctgtcaaac cacttgaagg aaagcaaaaga cctcaaatg 3900
cagaatggca tcagtaacca ggactggctc gcgtacatcc aggaactgag aaacgggttc 3960
aagaagcggc gtatcaagca agattga 3987

```

```

SEQ ID NO: 133      moltype = DNA length = 3987
FEATURE             Location/Qualifiers
source              1..3987
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 133
atggcgggaa gcaaaaaagcg ccggattaag caagacacgc agttcgaggg cttcacgaac 60
ctctaccaag tcagcaagac cctccgggtc gagctgatac cacagggaaa gacgtcaag 120
cacatccagg aacagggtct catcgaggag gacaaggcgc gcaacgacca ctacaaggag 180
ttgaaaccca tcatcgaccg catctacaag acgtacgcgc accagtgcct ccagctcgtg 240
cagctcgact gggagaacct ctccgcgcgc attgactcgt accggaagga gaagactgag 300
gagacccgca acgcccgtat cgaggagcaa gcaacctacc ggaacgccat ccacgactac 360
ttcatcgccc caccgcgcaa cctcaccgac gcgatcaaca agcggcacgc ggagatatac 420
aaagggctgt tcaaggcgga cgtgttcaac ggcaagggtg tcaagcagct agggacgggtg 480
accacgacgc agcacgagaa cgcgctctc cgcagcttcg acaagttcac cacctacttc 540

```

-continued

```

agcgggttct accggaaccc caagaatgtg ttcagcgcgg aggcacatcag caccggccatc 600
ccgcaccgca tcgtccagga caacttccc aagttcaagg agaactgcc catcttcacc 660
cgctcgataa ccgcgcgtcc ctccctgcgg gaggacttcg agaactgcaa aaaggcaatt 720
gggatcttcg tctcgaccag cattgaggag gtgttcagct tccccttcta caaccagctc 780
ctcaccagga cgcagatcga cctgtacaat cagttgctcg gcgggataag ccgcgaggcg 840
ggaaccgaaa aaatcaagggt gctgaacgaa gtgttgaacc tcgccatcca gaagaacgac 900
gagaccgcgc acatcatcgc ctccctgccc caccgggtca tcccgctgtt caagcagatc 960
ctctctgacc ggaacacccc gtccctcatt cttgaggagt tcaagtgcga cgaggaggctc 1020
atccagagct tctgcaagta caagacgctg ctacgggaac agaactgctt ggagacggcg 1080
gaggcactgt tcaacgagct aaacagcatc gacctcacgc acatcttcat cagtccaaag 1140
aaactggaga ccatctcttc cgcgctgtgc gacctctggg acacgctcag gaacgcgctc 1200
tacgagcgcc gaatcagtgat gctgacgggc aagatcacga agtcgcgaaa ggagaagggtg 1260
cagcgggtccc tcaagcacga ggacatcaac ctccaggaga tcatctcagc ggctgggaaa 1320
gagctgtccc aggcgttcaa gcagaaaacg agcgaatcc tgtcccacgc gcacgcggcc 1380
ctggatcagc ctctgcgcag gacctcaag aaacaagaag aaaggaaaat cctcaagtcg 1440
cagctcgact cgctgtgtgg cctgtacct ctctcgact ggttcggcgt ggacgagagc 1500
aacgaggtgg acccagagtt ctccgcgcgg cttacgggga tcaagctgga gatggagccc 1560
agcctgtcct tctacaacaa ggccgcgaac tacgccacca agaagcccta cagcgtggag 1620
aagttcaagc tcaacttcca gatgccact ctgcacgtg ggtgggacgt caaccgcgaa 1680
aaaaataatg gggcgatcct gtctgtcaag aacggcctgt actactggg catcatgccg 1740
aaacagaagg gccgtacaaa ggccctgagc ttgcaccca ccgagaaaac gagcgagggg 1800
ttcgacaaga tgtactacga ctactcccc gaacgcgcga agatgattcc aaagtgtccc 1860
acgcagctta aggcgctgac gggccacttc cagacgcaca cgaccctcat cctcctcagc 1920
aacaacttca tcgagccccc ggagatcacg aaggagatat acgacctgaa caacccgag 1980
aaggagccca agaaattcca gaccgcctac gccagaaga caggcgacca aaagggttac 2040
agggagggccc tctgcaagtg gatcgacttc actagggact tctgttccaa gtacaccaag 2100
actacctcta tcgacctgtc cagcctccgc ccgtcgtccc agtcaaaaag tttgggcgag 2160
tattacgcgg agctgaaccc actgctctac cacatcagct tccagcgcac cgcggagaag 2220
gagatcatgg acgcagtgga gacgggcaag ctatacctat ttcagatata caacaagac 2280
ttcgtaagg gacaccacgc caagcctaac ctgcacaccc tctactggac ggggctcttc 2340
agcccggaag accctgcgca gacctcgatc aagctcaacg gccaggccga gctgtttac 2400
cggcccaagt cccgcgtgaa gcggatggcc caccggctcg gggagaaaat gctcaacaag 2460
aaattgaagg accaaaaaac gccgatcccc gacaccctat accaggagct gtacgactat 2520
gtgaaccacc gcttgagcca gacctcagc gacgaggcgc gggccctcct gccgaacgtc 2580
atcacaaaag agtcagacca cgagatcatc aaggaccggc gcttcacctc cgacaagttt 2640
ttctttcacg tgcccatcac gctcaactac caggccgcca actcgccgtc caagttcaac 2700
cagcgcgtga acgctacact caaggagcac cccgagaccc cgatcatcgg gattgaccga 2760
ggggagcgga accctcatcta catcacgcgc atcgacagca ccgggaagat ccttgaacag 2820
cggctcgctc acaccatcca gcagttcgac taccagaaga aactcgacaa ccgggagaag 2880
gagagagtg cgcccccga ggttggttcc gtctcgggga cgattaagga cttgaaaaca 2940
ggttacctgt ccgaagtgtt ccacgagatc gttgacctga tgatccacta ccaagccgtc 3000
gtggtcctgg agaacctcaa cttcggttcc aagagcaaac gaaccggcat cgcggagaag 3060
gcggtgtacc ctaggttcga aaaaatgctg atcgacaagc tgaactgcct cgtgtcaag 3120
gactacccc ctagagaagg cgccgggggtg ctgaaccctg accagctcac tgaccagttc 3180
accagcttcg caaagatggg caccagctcc ggcttctcgt tctactgccc tgcgccatac 3240
acctcgaaga tcgaccgcgt caccgggttc tctgttcgga gacctcaag 3300
aaccacgaga gcccgcaagc cttcctggag ggcttcgact tcttccacta cgacgtcaag 3360
accggggact tcatctcgta cttcaagatg aaccgcaacc tcagtttcca gcgcggcctg 3420
ccgggggttca tgcccgcgtg ggatatagtc ttcgagaaga atgagacgca gttcgacg 3480
aagggcacc cgttcatcgc cgggaagcgc atcgtgcggg tcatcgagaa ccaccgggtc 3540
accgggcgtc accgcgacct ataccggcg aacgagttga tcgcccctcc ggaggagaag 3600
ggcatcgtgt tccgcgacgg ctccaacatc ctccgaagc tgcgcgaaaa cgacgactcc 3660
cacgccatcg acacgatggt cgcgctgac cggtcggtgc tccagatgcg gaactccaac 3720
gccgcgacgg gcgaggacta catcaacagt ccggtccgcg atctgaacgg cgtctgcttc 3780
gactcccggt tccagaaccc cagtgggcgg atggacggcg acgcgaacgg cgcataccac 3840
atcgccctaa aagggcaatt gctgtccaac caccctcaag aatccaaaga cctaaagctc 3900
cagaacggca tctccaacca ggactgggtg gcgtacatcc aggaactcgc gaacgggagc 3960
aaaaaacgtc ggatcaagca agattga 3987

```

```

SEQ ID NO: 134      moltype = DNA length = 3987
FEATURE             Location/Qualifiers
source               1..3987
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 134
atggcgggct ccaagaaacg ccggattaag caagataccc agttcgaggg gttcacgaac 60
ctctaccaag tgagcaagac cctccgattc gaactgattc ctcaggggaa gacctccaag 120
cacatccagg agcaagggtt catcgaggag gacaaggcgc ggaacgacca ctacaaggaa 180
ctcaaaccca tcatcgaccg catctacaag acctacgccg atcagtgctt ccagctcgtg 240
cagttggact gggagaacct cagcgcggcc attgactcct accggaagga gaaaacggag 300
gagacgcgca acgcgtctat cgaggaaacg gcaacctatc gcaacgccat ccacgactac 360
ttcatcggga ggactgacaa cctcactgac gcgattaaca agcgccacgc ggagatatac 420
aagggactct tcaagcgga gctgtttaac ggcaagggtt tcaagcaact cggcactgtg 480
accacgacct agcatgagaa gcgccgtgct cgctccttcg acaagttcac cactacttcc 540
tcggggttct accgcaacgg caagaatgtc ttacgcgcgg aggcacatcag caccggccatt 600
ccacatcgaa tcgtccaaga taacttccc aagttcaagg agaactgcca catcttcacc 660

```

-continued

```

cgactcatta ctgctgtacc gtcgttacgc gaacacttcg agaacgtcaa gaaggcaatt 720
ggaatctttcg tctctacgtc aatagaggag gtgttcagct tccctttcta caaccagctc 780
cttacgcaga cccagataga cctgtacaat cagctcctcg gtgggacag cggggaggcg 840
gggactgaga agattaaagg gctcaacgag gtcttgaacc tggccatcca aaaaaacgat 900
gagacggcgc acatcatcgc ctgctgccc caccggttca tcccgctgtt caagcagatc 960
ctcagtgaca ggaacacctt gagctttatc cttagaggag tcaagagcga cgaggagggtg 1020
atccagagct tctgcaagta caaaacctg ctgaggaacg agaacgtcct ggagacggcg 1080
gaggcgctgt tcaacgagct gaactctatc gacttaactc acatattcat ctgcacaaag 1140
aagctggaga ctattagctc tgcactctgc gaccactggg acacctcccg caacgcgctc 1200
tacgagcgcc gcctctcgga gctgaccggg aagatcacca aatccgcgaa ggaaaaggtc 1260
cagcgttccc tcaaacacga ggatattaac ttacaggaga ttatctcagc ggctgggaag 1320
gagttgtcag aggcgttcaa gcagaaaact tccgagatcc tgagccacgc gcacgcagcg 1380
ctcgaccacg ctctgcccac caccctcaaa aagcaggaag aaaaagagat cctcaagagc 1440
cagttggact cctctctggg gctctatcac cttctcgact gggtcgccgt cgatgagtcg 1500
aacgaggtgg accccgagtt ctccgcccgg ctgaccggca tcaagctaga gatggagcgg 1560
tccctcagct tctacaataa ggcccgcacac tacgcgacca aaaaacctta cagcgtggag 1620
aagttcaagc tgaacttcca gatgccgacc ttagcacgcg gttgggacgt aaacagggag 1680
aagaacaatg gaccatcct gtctgtcaag aacgggcttt actacctcgg gataatgcc 1740
aagcagaagg gccgctacaa ggccttttcc ttcgagccga cggagaaaaac ctccgagggg 1800
ttcgacaaga tgtactacga ctacttcccc gacgcgcgca agatgatccc gaagtgtctc 1860
acgcagctaa aagccgtgac cgcccacttc cagaccacca cgacgcgat cctgctgagc 1920
aacaacttca tcgagccctt tgagatcact aaggagatat acgacctgaa caaccccgag 1980
aaggagccca agaagtttca aaccgctcac gccaaaaaaa ctggcgacca aaagggttac 2040
agggaggcgc tgtgtaagtg gatcgacttc acacgcgact tcccttcgaa gtatacgaag 2100
acaacctcta ttgacctgag cagcctgcgt cctagctccc agtacaagaa tttgggcgag 2160
tactacgcgg agcttaatcc actactctac cacatctcat tccagcgcac cgctgagaag 2220
gaaatcatgg acgcggtgga gacaggcaaa ctgtacctct tccagatata caacaaagac 2280
ttcgctaagg ggcaaccagg gaagcccaac cttcatagc tctactggac ggcctattc 2340
agccccgaaa atctggccaa gacctccatc aagctgaacg gccaaagcga gctgttctac 2400
agacccaaga gccggatgaa gcggatggcc cacaggctcg gcgagaaat gcttaacaaa 2460
aagttgaagg accagaaaac ccttatcccc gacacctct accaggaact gtacgactac 2520
gtgaaccaca ggctctcgca cgacctttcc gacgaggccc gtgccctact cccgaacgtc 2580
attaccaaa aggttttcga cgagatcatc aaggaccggc gggtcacgag cgacaagttt 2640
ttctttcacg tccccatcac ccttaactac caggcgccca actccccatc caagttcaac 2700
cagcgtgtga atgcctacct caaggagcac ccagagaccc cgatcatagg gatcgaccgg 2760
ggcgagcggg acctgatcta catcacgcgt atcgactcga cgggcaagat tcttgagcag 2820
agatcgttga ataccataga gcagttcgac taccagaaga aactcgacaa ccgcgagaag 2880
gagcgcgtgg cggcccgcca ggcgtggtcc gtctgtggga cgattaagga cttgaaacaa 2940
ggttatctgt ccaagtcac ccacgagatc gttgatctga tgatccacta tcaggcagtg 3000
gtggtgctgg agaattctca cttcgcttc aagagtaagc ggacgggaat cgcgcgagaag 3060
gccgtgtacc agcagttcga gaagatgtg atcgacaagc tcaactgcct tgtgctgaaa 3120
gactaccggc ccgagaaggt cggcggcgtc ctcaaccgt accaacttac cgaccagttc 3180
acctccttcg ccaagatggg cactcagtc ggggttctgt tctacgtccc cgcaccttac 3240
acctctaaga tcgacctct gactggcttc gtagatccat tcgtgtggaa gaccattaa 3300
aaccacgaga gccgcaagca cttcctggag ggcttcgact tctcgacta cgacgtgaag 3360
accggggact tcatccttca cttcaagatg aaccggaacc tcagcttcca gcggggcctg 3420
ccgggggttca tgcccgctg ggacatcgtg ttcgagaaga acgagaccca gttcgacgcg 3480
aagggcacgc ccttcacgc cggaagcgt atcgtgccg tgatcgagaa ccatcgtttc 3540
acgggtcgct accgtgacct ctaccggcg aacgagctta tcgactcct ggaggagaag 3600
ggcatcgtct tccgggacgg ctccaacatc ctcccgaac tgctggaaaa cgacgactct 3660
cacgcatcgc acacgatggg ggcctcatc cgggtccgtg tccaaatgcg gaacagcaac 3720
gccgccaccg gtgaggacta catcaacagc ccgggtccgg atctgaacgg ggtgtgcttc 3780
gattcgcggt tccagaatcc tgagtggccg atggacgcgg atgcaaacgg ggcgtaccac 3840
atcgcgctca agggccagtt actctgaac caccttaagg agtctaaaga tttgaaactc 3900
cagaacggga tctcgaacca ggactggctg gcctacatcc aagagttgcg gaacggcagc 3960
aagaagcggc ggattaagca agattag 3987

```

```

SEQ ID NO: 135      moltype = AA length = 9
FEATURE            Location/Qualifiers
source              1..9
                    mol_type = protein
                    organism = synthetic construct

SEQUENCE: 135
LAGLIDADG

```

What is claimed is:

1. An isolated polynucleotide encoding an amino acid sequence having at least 95% sequence identity to the polypeptide sequence of SEQ ID NO:26, wherein the isolated polynucleotide is operably associated with a heterologous promoter.

2. A cell, plant part, or plant comprising the isolated polynucleotide of claim 1.

3. The cell, plant part, or plant of claim 2, further comprising a polynucleotide encoding gene editing machinery and/or a polynucleotide encoding a selectable marker.

4. An expression cassette and/or vector comprising the isolated polynucleotide of claim 1.

5. The expression cassette and/or vector of claim 4, further comprising a polynucleotide encoding gene editing machinery and/or a polynucleotide encoding a selectable marker.

* * * * *