



US 20250283128A1

(19) **United States**(12) **Patent Application Publication**
KIM et al.(10) **Pub. No.: US 2025/0283128 A1**(43) **Pub. Date: Sep. 11, 2025**(54) **A GENUS OF CORYNEBACTERIUM
MICROORGANISM FOR PRODUCING
L-ARGININE AND A METHOD FOR
PRODUCING L-ARGININE USING THE
SAME**(71) Applicant: **CJ CHEILJEDANG
CORPORATION**, Seoul (KR)(72) Inventors: **Hyo Kyung KIM**, Seoul (KR); **Jin
Sook CHANG**, Seoul (KR); **Sun
Hyoung CHOI**, Seoul (KR); **Haena
OH**, Seoul (KR); **Byoung Hoon
YOON**, Seoul (KR); **Hyung Joon
KIM**, Seoul (KR)(73) Assignee: **CJ CHEILJEDANG
CORPORATION**, Seoul (KR)(21) Appl. No.: **18/286,067**(22) PCT Filed: **Apr. 7, 2022**(86) PCT No.: **PCT/KR2022/005056**

§ 371 (c)(1),

(2) Date: **Dec. 20, 2023**(30) **Foreign Application Priority Data**

Apr. 7, 2021 (KR) 10-2021-0045262

Publication Classification(51) **Int. Cl.*****C12P 13/10*** (2006.01)***C12N 1/20*** (2006.01)***C12R 1/15*** (2006.01)(52) **U.S. Cl.**CPC ***C12P 13/10*** (2013.01); ***C12N 1/20***
(2013.01); ***C12R 2001/15*** (2021.05)

(57)

ABSTRACT

The present disclosure relates to an L-arginine-producing microorganism, in which a protein comprising an amino acid sequence of SEQ ID NO: 1 is weakened, and a method for producing L-arginine using the same.

Specification includes a Sequence Listing.

**A GENUS OF CORYNEBACTERIUM
MICROORGANISM FOR PRODUCING
L-ARGININE AND A METHOD FOR
PRODUCING L-ARGININE USING THE
SAME**

TECHNICAL FIELD

[0001] The present disclosure relates to an L-arginine-producing microorganism, in which a protein including an amino acid sequence of SEQ ID NO: 1 is weakened, and a method for producing L-arginine using the same.

BACKGROUND ART

[0002] L-Arginine has been widely used in medicines as a hepatic function-promoting agent, as a brain function-promoting agent, and as an ingredient of multiple amino acid supplements. Additionally, L-arginine has gained much interest in the food industry as a food additive for fish cakes and health beverages, and as a salt substitute for hypertension patients. Studies have been continuously conducted to use microorganisms for the production of industrially available high-concentration arginine, and examples thereof include a method of using a mutant strain induced from the microorganism belonging to the genus *Brevibacterium* or *Corynebacterium*, which is a glutamate-producing strain, or a method of using an amino acid-producing strain whose growth is improved through cell fusion. Meanwhile, *lysE* of the microorganism belonging to the genus *Corynebacterium* having an ability to export L-lysine has also been shown to export the same basic amino acid L-arginine (Bellmann A, et al., *Microbiology*, 147:1765-1774, 2001). Further, it has been disclosed a method for enhancing the production ability of L-arginine-producing strains through gene enhancement (U.S. Patent Application Publication No. US 2003-0113899 A1).

DISCLOSURE

Technical Problem

[0003] The present inventors have developed a microorganism of the genus *Corynebacterium* producing L-arginine and a method for producing L-arginine using the same, thereby completing the present disclosure.

Technical Solution

[0004] It is one object of the present disclosure to provide a recombinant microorganism of the genus *Corynebacterium* producing L-arginine, in which the activity of a protein including an amino acid sequence of SEQ ID NO: 1 is weakened.

[0005] It is another object of the present disclosure to provide a method for producing L-arginine, including: culturing a recombinant microorganism of the genus *Corynebacterium*, in which the activity of a protein including an amino acid sequence of SEQ ID NO: 1 is weakened, in a medium.

Advantageous Effects

[0006] The microorganism of the genus *Corynebacterium*, in which the activity of a protein including an amino acid sequence of SEQ ID NO: 1 is weakened, can produce

L-arginine in high yield, and thus can be effectively used for industrial production of L-arginine.

**DETAILED DESCRIPTION OF PREFERRED
EMBODIMENTS**

[0007] The present disclosure will be described in detail as follows. Meanwhile, each description and embodiment disclosed herein can be applied to other descriptions and embodiments, respectively. That is, all combinations of various elements disclosed herein fall within the scope of the present disclosure. Further, the scope of the present disclosure is not limited by the specific description described below. Additionally, a number of papers and patent documents have been cited throughout the present specification. The content of the cited papers and patent documents is incorporated herein by reference in their entirety and the level of technical field to which the present invention belongs and the contents of the present invention will be described more clearly.

[0008] One aspect of the present disclosure provides a recombinant microorganism of the genus *Corynebacterium*, in which the activity of a protein including an amino acid sequence of SEQ ID NO: 1 is weakened.

[0009] The protein of the present disclosure may have, include, consist or essentially consist of an amino acid represented by SEQ ID NO: 1.

[0010] In the present disclosure, the protein including the amino acid sequence of SEQ ID NO: 1 may be an endogenous protein of the microorganism of the present disclosure, but is not limited thereto.

[0011] The amino acid sequence of SEQ ID NO: 1 can be obtained from NIH GenBank, a known database. In the present disclosure, the amino acid sequence of SEQ ID NO: 1 may include an amino acid sequence having a homology or identity of at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, 99.5%, 99.7% or 99.9% with the amino acid sequence of SEQ ID NO: 1. Additionally, it is apparent that variants having an amino acid sequence, in which a part of the amino acid sequence is deleted, modified, substituted, conservatively substituted or added, may fall within the scope of the present disclosure, as long as the amino acid sequence has such homology or identity and shows an efficacy corresponding to that of the variant of the present disclosure.

[0012] For example, it may include sequence additions or deletions, naturally occurring mutations, silent mutations or conservative substitutions that do not alter the function of the variant of the present disclosure at the N-terminus, C-terminus and/or within the amino acid sequence.

[0013] As used herein, the term “conservative substitution” refers to substitution of an amino acid with another amino acid having similar structural and/or chemical properties. The variant may have, for example, one or more conservative substitutions while still retaining one or more biological activities. Such amino acid substitution may generally occur based on similarity of polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or amphipathic nature of a residue. For example, among the amino acids having an electrically charged side chain (electrically charged amino acid), positively charged (basic) amino acids include arginine, lysine, and histidine; negatively charged (acidic) amino acids include glutamic acid and aspartic acid. Further, among the amino acids having an uncharged side chain (uncharged amino acid), nonpolar amino acids include

glycine, alanine, valine, leucine, isoleucine, methionine, phenylalanine, tryptophan, and proline; polar or hydrophilic amino acids include serine, threonine, cysteine, tyrosine, asparagine, and glutamine; and among the nonpolar amino acids, aromatic amino acids include phenylalanine, tryptophan and tyrosine.

[0014] As used herein, the term “homology” or “identity” refers to a degree of relevance between two given amino acid sequences or nucleotide sequences, and may be expressed as a percentage. The terms homology and identity may often be used interchangeably with each other.

[0015] The sequence homology or identity of conserved polynucleotide or proteins may be determined by standard alignment algorithms and can be used with a default gap penalty established by the program being used. Substantially, homologous or identical sequences are generally expected to hybridize to all or part of the entire length of the sequences under moderate or high stringent conditions. It is apparent that hybridization with polynucleotides containing general codon or degenerate codons in hybridizing polynucleotides is also included.

[0016] Whether any two polynucleotide or protein sequences have a homology, similarity or identity may be, for example, determined by a known computer algorithm such as the “FASTA” program using default parameters (Pearson et al., (1988) *Proc. Natl. Acad. Sci. USA* 85:2444). Alternatively, it may be determined by the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, *J. Mol. Biol.* 48:443-453), which is performed using the Needleman program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice et al., 2000, *Trends Genet.* 16:276-277) (version 5.0.0 or later) (GCG program package (Devereux, J., et al., *Nucleic Acids Research* 12:387 (1984)), BLASTP, BLASTN, FASTA (Atschul, S. F., et al., *J MOLEC BIOL* 215:403 (1990); *Guide to Huge Computers*, Martin J. Bishop, ed., Academic Press, San Diego, 1994, and CARILLO et al. (1988) *SIAM J Applied Math* 48:1073). For example, the homology, similarity or identity may be determined using BLAST or ClustalW of the National Center for Biotechnology Information (NCBI).

[0017] The homology, similarity, or identity of polynucleotides or proteins may be determined by comparing sequence information using, for example, the GAP computer program, such as Needleman et al. (1970), *J Mol Biol.* 48:443 as disclosed in Smith and Waterman, *Adv. Appl. Math* (1981) 2:482. In summary, the GAP program defines the homology, similarity or identity as the value obtained by dividing the number of similarly aligned symbols (i.e., nucleotides or amino acids) by the total number of the symbols in the shorter of the two sequences. Default parameters for the GAP program may include (1) a unary comparison matrix (containing a value of 1 for identities and 0 for non-identities) and the weighted comparison matrix of Gribskov et al. (1986), *Nucl. Acids Res.* 14:6745, as disclosed in Schwartz and Dayhoff, eds., *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation, pp. 353-358 (1979) (or EDNAFULL substitution matrix (EMBOSS version of NCBI NUC4.4)); (2) a penalty of 3.0 for each gap and an additional 0.10 penalty for each symbol in each gap (or a gap opening penalty of 10 and a gap extension penalty of 0.5); and (3) no penalty for end gaps.

[0018] In the present disclosure, the polynucleotide encoding the protein including the amino acid sequence of SEQ ID NO: 1 may be named NCgl2608 gene.

[0019] As used herein, the term “polynucleotide”, which is a polymer of nucleotides composed of nucleotide monomers connected in a lengthy chain by a covalently bond, is a DNA or RNA strand having at least a certain length. More specifically, it may refer to a polynucleotide fragment encoding the variant.

[0020] The polynucleotide encoding the variant of the present disclosure may include a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 1. As an example of the present disclosure, the polynucleotide of the present disclosure may have or include a nucleotide sequence of SEQ ID NO: 2. In addition, the polynucleotide of the present disclosure may consist or consist essentially of a nucleotide sequence of SEQ ID NO: 2.

[0021] The polynucleotide of the present disclosure may undergo various modifications in the coding region within the scope that does not change the amino acid sequence of the variant, due to codon degeneracy or in consideration of the codons preferred in an organism in which the variant of the present disclosure is to be expressed. Specifically, the polynucleotide of the present disclosure may have or include a nucleotide sequence having a homology or identity of 70% or more, 75% or more, 80% or more, 85% or more, 90% or more, 95% or more, 96% or more, 97% or more, 98% or more and less than 100% with the sequence of SEQ ID NO: 2, or may consist or consist essentially of a nucleotide sequence having a homology or identity of 70% or more, 75% or more, 80% or more, 85% or more, 90% or more, 95% or more, 96% or more, 97% or more, 98% or more, and less than 100% with the sequence of SEQ ID NO: 2, but is not limited thereto.

[0022] Additionally, the polynucleotide of the present disclosure may include a probe that may be prepared from a known gene sequence, for example, any sequence which can hybridize with a sequence complementary to all or part of the polynucleotide sequence of the present disclosure under stringent without limitation. The “stringent conditions” refers to conditions under which specific hybridization between polynucleotides is allowed. Such conditions are specifically described in the literature (J. Sambrook et al., *Molecular Cloning, A Laboratory Manual*, 2nd Edition, Cold Spring Harbor Laboratory press, Cold Spring Harbor, New York, 1989; F. M. Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc., New York, 9.50-9.51, 11.7-11.8). For example, the stringent conditions may include conditions under which polynucleotides having a high homology or identity of 70% or more, 75% or more, 80% or more, 85% or more, 90% or more, 95% or more, 96% or more, 97% or more, 98% or more, or 99% or more are hybridized with each other and polynucleotides having a homology or identity lower than the above homologies or identities are not hybridized with each other, or washing conditions of Southern hybridization, that is, washing once, specifically, twice or three times at a salt concentration and a temperature corresponding to 60° C., 1×SSC, 0.1% SDS, specifically 60° C., 0.1×SSC, 0.1% SDS, and more specifically 68° C., 0.1×SSC, 0.1% SDS.

[0023] Hybridization requires that two nucleic acids contain complementary sequences, although mismatches between bases are possible depending on the stringency of the hybridization. The term “complementary” is used to

describe the relationship between nucleotide bases that can hybridize with each other. For example, with respect to DNA, adenine is complementary to thymine, and cytosine is complementary to guanine. Therefore, the polynucleotide of the present disclosure may include isolated nucleotide fragments complementary to the entire sequence as well as nucleic acid sequences substantially similar thereto.

[0024] Specifically, polynucleotides having a homology or identity with the polynucleotide of the present disclosure may be detected using the hybridization conditions including a hybridization step at a T_m value of 55° C. under the above-described conditions. Further, the T_m value may be 60° C., 63° C., or 65° C., but is not limited thereto, and may be appropriately adjusted by those skilled in the art depending on the purpose thereof.

[0025] The appropriate stringency for hybridizing the polynucleotides depends on the length of the polynucleotides and the degree of complementation, and these variables are well known in the art (e.g., Sambrook et al.).

[0026] As used herein, the term “microorganism (or strain)” includes all wild-type microorganisms, or naturally or artificially genetically modified microorganisms, and it may be a microorganism in which a particular mechanism is weakened or enhanced due to insertion of a foreign gene, or enhancement or inactivation of the activity of an endogenous gene, and may be a microorganism including genetic modification to produce a desired protein or product.

[0027] The microorganism of the present disclosure may be a microorganism in which the protein including the amino acid sequence of SEQ ID NO: 1 of the present disclosure or the polynucleotide encoding the protein is deleted; or a microorganism (i.e., a recombinant microorganism) which is genetically modified through a vector such that the protein including the amino acid sequence of SEQ ID NO: 1 of the present disclosure or the polynucleotide encoding the protein is deleted, but is not limited thereto.

[0028] The microorganism of the present disclosure may be a microorganism having the ability to produce L-arginine.

[0029] The microorganism of the present disclosure may be a microorganism having increased L-arginine producing ability compared to a parent strain.

[0030] The microorganism of the present disclosure may be a microorganism in which the ability to produce L-arginine has been imparted by weakening the protein including the amino acid sequence of SEQ ID NO: 1 of the present disclosure or deleting the polynucleotide encoding the protein into a parent strain (i.e., having the ability to naturally produce L-arginine or having no ability to produce L-arginine), but is not limited thereto.

[0031] In one example, the recombinant microorganism of the present disclosure is a microorganism which is transformed with a vector such that the protein including the amino acid sequence of SEQ ID NO: 1 of the present disclosure is weakened or the polynucleotide encoding the protein is deleted, and thus the protein including the amino acid sequence of SEQ ID NO: 1 of the present disclosure is weakened or the polynucleotide encoding the protein is deleted, and may be a microorganism having increased L-arginine producing ability as compared to a natural wild-type microorganism by weakening the protein including the amino acid sequence of SEQ ID NO: 1 of the present disclosure or deleting the polynucleotide encoding the protein in a natural wild-type microorganism or an L-arginine-producing microorganism, or a microorganism in which the

protein including the amino acid sequence of SEQ ID NO: 1 of the present disclosure or the polynucleotide encoding the protein is increased as compared to a non-modified microorganism, but is not limited thereto.

[0032] In one example, the non-modified microorganism, which is a target strain to be compared to determine the increase in the L-arginine producing ability, may be a *Corynebacterium glutamicum* KCCM10741P (van der Rest et al., *Appl Microbiol Biotechnol* 52:541-545, 1999), or a strain in which the argR gene is deleted in the wild-type *Corynebacterium glutamicum* ATCC 13869 (e.g., *Corynebacterium glutamicum* CJ1R, but is not limited thereto).

[0033] In one example, the recombinant strain having increased producing ability may have an increased L-arginine producing ability by about 1% or more, specifically about 1% or more, about 2.5% or more, about 5% or more, about 6% or more, about 7% or more, about 8% or more, about 9% or more, about 10% or more, about 10.5% or more, about 11% or more, about 11.5% or more, about 12% or more, about 12.5% or more, about 13% or more, about 13.5% or more, or about 14% or more (the upper limit is not particularly limited, for example, about 200% or less, about 150% or less, about 100% or less, about 50% or less, about 40% or less, about 30% or less, or about 25% or less), as compared to that of the parent strain before modification or a non-modified microorganism, but is not limited thereto, as long as it has an increased value compared to the production ability of the parent strain before modification or non-modified microorganism. In another example, the microorganism having increased production ability may have an increased L-arginine producing ability by about 1.1 times or more, about 1.12 times or more, about 1.13 times or more, or 1.14 times or more (the upper limit is not particularly limited, for example, about 10 times or less, about 5 times or less, about 3 times or less, or about 2 times or less), as compared to that of the parent strain before modification or non-modified microorganism, but is not limited thereto. As used herein, the term “about” refers to a range which includes all of ± 0.5 , ± 0.4 , ± 0.3 , ± 0.2 , ± 0.1 , etc. and includes all of the values that are equivalent or similar to those following the values, but the range is not limited thereto.

[0034] As used herein, the term “non-modified microorganism” does not exclude a strain containing a mutation that may occur naturally in a microorganism, and may refer to a wild-type strain or a natural-type strain itself, or a strain before the trait is altered due to genetic modification caused by natural or artificial factors. For example, the non-modified microorganism may refer to a microorganism before the protein including the amino acid sequence of SEQ ID NO: 1 of the present disclosure described in the present specification is weakened or the polynucleotide encoding the protein is deleted. The “non-modified microorganism” may be used interchangeably with “strain before modification”, “microorganism before modification”, “non-mutant strain”, “non-modified strain”, “non-mutant microorganism”, or “reference microorganism”.

[0035] In another embodiment, the microorganism of the genus *Corynebacterium* of the present disclosure may be *Corynebacterium glutamicum*, *Corynebacterium crudilactis*, *Corynebacterium deserti*, *Corynebacterium efficiens*, *Corynebacterium callunae*, *Corynebacterium stationis*, *Corynebacterium singulare*, *Corynebacterium halotolerans*, *Corynebacterium striatum*, *Corynebacterium ammoni-*

agenes, *Corynebacterium pollutisoli*, *Corynebacterium imitans*, *Corynebacterium testudinoris*, or *Corynebacterium flavesceus*.

[0036] Specifically, the recombinant microorganism of the present disclosure may be a microorganism in which all or part of the polynucleotide encoding the protein consisting of the amino acid sequence of SEQ ID NO: 1 is deleted.

[0037] Additionally, the microorganism of the present disclosure may be a microorganism in which the activity of arginine repressor (ArgR) is additionally weakened. Specifically, the microorganism of the present disclosure may be a microorganism in which all or part of the argR gene is additionally deleted.

[0038] The ArgR may include a polypeptide represented by the amino acid sequence of SEQ ID NO: 13. In addition, the argR gene may include a polynucleotide represented by the nucleotide sequence of SEQ ID NO: 14, but is not limited thereto.

[0039] In addition, the microorganism of the present disclosure may be a microorganism in which the activity of carbamoyl phosphate synthase small subunit (CarA) and/or carbamoyl phosphate synthase large subunit (CarB) are further enhanced, or which expresses normal ornithine carbamoyltransferase F (ArgF), or includes a combination thereof.

[0040] In one example, the microorganism of the present disclosure a *Corynebacterium glutamicum* microorganism in which carA and/or carB protein activity is further enhanced in NCgl2608 gene-deleted strain of the *Corynebacterium glutamicum* KCCM10741P (van der Rest et al., *Appl Microbiol Biotechnol* 52:541-545, 1999) or NCgl2608 gene-deleted strain of the *Corynebacterium glutamicum* CJIR.

[0041] In addition, the microorganism of the present disclosure may be a strain in which the NCgl2608 gene is deleted in *Corynebacterium glutamicum* KCCM10741P, or a microorganism that additionally expresses normal argF in *Corynebacterium glutamicum* CJIR, and the microorganism expressing the normal argF additionally be a *Corynebacterium glutamicum* microorganism with additionally enhanced carA and/or carB protein activity.

[0042] The argF protein may contain a polypeptide represented by the amino acid sequence of SEQ ID NO: 15, and the argF gene may contain a polynucleotide represented by the nucleotide sequence of SEQ ID NO: 16, but are not limited thereto. Further, the carA protein may contain a polypeptide represented by the amino acid sequence of SEQ ID NO: 17, the carB protein may contain a polypeptide represented by the amino acid sequence of SEQ ID NO: 19, the carA gene may contain a polynucleotide represented by the nucleotide sequence of SEQ ID NO: 18, and the carB gene may contain a polynucleotide represented by the nucleotide sequence of SEQ ID NO: 20.

[0043] As used herein, the term “weakening” of a protein is a comprehensive concept including both reduced or no activity compared to its endogenous activity. The weakening may be used interchangeably with terms such as inactivation, deficiency, down-regulation, decrease, reduce, attenuation, etc.

[0044] The weakening may also include a case where the protein activity itself is decreased or removed compared to the activity of the protein originally possessed by a microorganism due to a mutation of the polynucleotide encoding the protein; a case where the overall level of intracellular

protein activity and/or concentration (expression level) is decreased compared to a natural strain due to the inhibition of expression of the gene of the polynucleotide encoding the protein, or the inhibition of translation into the protein, etc.; a case where the polynucleotide is not expressed at all; and/or a case where no protein activity is observed even when the polynucleotide is expressed. As used herein, the term “endogenous activity” refers to the activity of a particular protein originally possessed by a parent strain before transformation, a wild-type or a non-modified microorganism, when a trait is altered through genetic modification caused by natural or artificial factors, and may be used interchangeably with “activity before modification”. The expression “the protein activity is ‘inactivated, deficient, decreased, down-regulated, reduced or attenuated’ compared to its endogenous activity” means that the protein activity is decreased compared to the activity of a particular protein originally possessed by a parent strain before transformation or a non-modified microorganism.

[0045] The weakening of the protein activity can be performed by any method known in the art, but the method is not limited thereto, and can be achieved by applying various methods well known in the art (e.g., Nakashima N et al., *Bacterial cellular engineering by genome editing and gene silencing*. *Int J Mol Sci*. 2014; 15(2):2773-2793; Sambrook et al. *Molecular Cloning* 2012; etc.).

[0046] Specifically, the weakening of the protein activity of the present disclosure may be achieved by:

[0047] 1) deleting a part or all of the gene encoding the protein;

[0048] 2) modifying the expression regulatory region (expression regulatory sequence) such that the expression of the gene encoding the protein is decreased;

[0049] 3) modifying the amino acid sequence constituting the protein such that the protein activity is removed or weakened (e.g., deletion/substitution/addition of one or more amino acids on the amino acid sequence);

[0050] 4) modifying the gene sequence encoding the protein such that the protein activity is removed or weakened (e.g., deletion/substitution/addition of one or more of nucleotides on the nucleotide sequence of the protein gene to encode a protein that has been modified to remove or weaken the activity of the protein);

[0051] 5) modifying the nucleotide sequence encoding the initiation codon or 5'-UTR of the gene transcript encoding the protein;

[0052] 6) introducing an antisense oligonucleotide (e.g., antisense RNA), which binds complementary to the gene transcript encoding the protein;

[0053] 7) adding a sequence complementary to the Shine-Dalgarno (SD) sequence on the front end of the SD sequence of the gene encoding the protein to form a secondary structure, thereby inhibiting the ribosomal attachment;

[0054] 8) a reverse transcription engineering (RTE), which adds a promoter, which is to be reversely transcribed, on the 3' terminus of the open reading frame (ORF) of the gene sequence encoding the protein; or

[0055] 9) a combination of two or more selected from the methods 1) to 8) above, but is not particularly limited thereto.

[0056] For example,

[0057] The 1) method of deleting a part or all of the gene encoding the protein may be achieved by deleting all of the

polynucleotide encoding the endogenous target protein within the chromosome, or by replacing the polynucleotide with a polynucleotide or a marker gene having partially deleted nucleotides.

[0058] The 2) method of modifying the expression regulatory region (expression regulatory sequence) may be achieved by inducing a modification on the expression regulatory region (expression regulatory sequence) through deletion, insertion, non-conservative substitution or conservative substitution, or a combination thereof; or by replacing the sequence with a sequence having a weaker activity. The expression regulatory region may include a promoter, an operator sequence, a sequence encoding a ribosome-binding site, and a sequence for regulating transcription and translation, but is not limited thereto.

[0059] The 3) method of modifying the nucleotide sequence encoding the initiation codon or 5'-UTR of the gene transcript encoding the protein may be achieved, for example, by substituting the nucleotide sequence with a nucleotide sequence encoding another initiation codon having a lower polypeptide expression rate than the endogenous initiation codon, but is not limited thereto.

[0060] The 4) and 5) methods of modifying the amino acid sequence or the polynucleotide sequence may be achieved by inducing a modification on the sequence through deletion, insertion, non-conservative or conservative substitution of the amino acid sequence of the protein or the polynucleotide sequence encoding the protein, or a combination thereof to weaken the activity of the protein, or by replacing the sequence with an amino acid sequence or a polynucleotide sequence modified to have a weaker activity, or an amino acid sequence or a polynucleotide sequence modified to have no activity, but are not limited thereto. For example, the expression of a gene may be inhibited or weakened by introducing a mutation into the polynucleotide sequence to form a stop codon, but is not limited thereto.

[0061] The 6) method of introducing an antisense oligonucleotide (e.g., antisense RNA), which binds complementary to the gene transcript encoding the protein can be found in the literature (Weintraub, H. et al., Antisense-RNA as a molecular tool for genetic analysis, *Reviews—Trends in Genetics*, Vol. 1(1) 1986).

[0062] The 7) method of adding a sequence complementary to the Shine-Dalgarno (SD) sequence on the front end of the SD sequence of the gene encoding the protein to form a secondary structure, thereby inhibiting the ribosome attachment may be achieved by inhibiting mRNA translation or reducing the speed thereof.

[0063] The 8) reverse transcription engineering (RTE), which adds a promoter, which is to be reversely transcribed, on the 3' terminus of the open reading frame (ORF) of the gene sequence encoding the protein may be achieved by forming an antisense nucleotide complementary to the gene transcript encoding the protein to weaken the activity.

[0064] As used herein, the term “enhancement” of a protein activity means that the activity of a protein is increased compared to its endogenous activity. The enhancement may be used interchangeably with terms such as activation, up-regulation, overexpression, increase, etc. In particular, the terms activation, enhancement, up-regulation, overexpression and increase may include both cases in which an activity not originally possessed is exhibited, or the activity is enhanced compared to the endogenous activity or the activity before modification. The “endogenous activity”

refers to the activity of a particular protein originally possessed by a parent strain before transformation or a non-modified microorganism, when a trait is altered through genetic modification caused by natural or artificial factors, and may be used interchangeably with “activity before modification”. The “enhancement”, “up-regulation”, “overexpression”, or “increase” in the activity of a protein compared to its endogenous activity means that the activity and/or concentration (expression level) of the protein is enhanced compared to that of a particular protein originally possessed by a parent strain before transformation or a non-modified microorganism.

[0065] The enhancement may be achieved by introducing a foreign protein, or by enhancing the activity and/or concentration (expression level) of the endogenous protein. The enhancement of the activity of the protein can be confirmed by the increase in the level of activity of the protein, expression level, or the amount of product excreted from the protein.

[0066] The enhancement of the activity of the protein can be applied by various methods well known in the art, and is not limited as long as it can enhance the activity of the target protein compared to that of the microorganism before modification. Specifically, genetic engineering and/or protein engineering well known to those skilled in the art, which is a common method of molecular biology, may be used, but the method is not limited thereto (e.g., Sitnicka et al. *Functional Analysis of Genes. Advances in Cell Biology*. 2010, Vol. 2. 1-16; Sambrook et al. *Molecular Cloning* 2012; etc.).

[0067] Specifically, the enhancement of the protein of the present disclosure may be achieved by:

[0068] 1) increasing the intracellular copy number of a polynucleotide encoding the protein;

[0069] 2) replacing the expression regulatory region of a gene encoding the protein on the chromosome with a sequence having a strong activity;

[0070] 3) modifying the nucleotide sequence encoding the initiation codon or 5'-UTR of the gene transcript encoding the protein;

[0071] 4) modifying the amino acid sequence of the protein such that the activity of the polypeptide is enhanced;

[0072] 5) modifying the polynucleotide sequence encoding the protein such that the activity of the protein is enhanced (e.g., modifying the polynucleotide sequence of the protein gene to encode a protein that has been modified to enhance the activity of the protein);

[0073] 6) introducing a foreign protein having the activity of the protein or a foreign polynucleotide encoding the same;

[0074] 7) codon-optimization of the polynucleotide encoding the protein;

[0075] 8) analyzing the tertiary structure of the protein and thereby selecting and modifying the exposed site, or chemically modifying the same; or

[0076] 9) a combination of two or more selected from above 1 to 8), but is not particularly limited thereto.

[0077] More specifically,

[0078] The 1) method of increasing the intracellular copy number of a polynucleotide encoding the protein may be achieved by introducing a vector, which is operably linked to the polynucleotide encoding the protein and is able to

replicate and function regardless of a host cell, into the host cell. Alternatively, the method may be achieved by introducing one copy or two copies of polynucleotides encoding the protein into the chromosome of a host cell. The introduction into the chromosome may be performed by introducing a vector, which is able to insert the polynucleotide into the chromosome of a host cell, into the host cell, but is not limited thereto. The vector is as described above.

[0079] The 2) method of replacing the expression regulatory region (or expression regulatory sequence) of a gene encoding the protein on the chromosome with a sequence having a strong activity may be achieved by inducing a modification on the sequence through deletion, insertion, non-conservative or conservative substitution, or a combination thereof to further enhance the activity of the expression regulatory region, or by replacing the sequence with a sequence having a stronger activity. The expression regulatory region may include, but is not particularly limited to, a promoter, an operator sequence, a sequence encoding a ribosome-binding site, and a sequence regulating the termination of transcription and translation. In one example, the method may include replacing the original promoter with a strong promoter, but is not limited thereto.

[0080] Examples of the strong promoter may include CJ1 to CJ7 promoters (U.S. Pat. No. 7,662,943 B2), lac promoter, trp promoter, trc promoter, tac promoter, lambda phage PR promoter, PL promoter, tet promoter, gapA promoter, SPL7 promoter, SPL13(sm3) promoter (U.S. Pat. No. 10,584,338 B2), O2 promoter (U.S. Pat. No. 10,273,491 B2), tkt promoter, yccA promoter, etc., but are not limited thereto.

[0081] The 3) method of modifying the nucleotide sequence encoding the initiation codon or 5'-UTR of the gene transcript encoding the protein may be achieved, for example, by substituting the nucleotide sequence with a nucleotide sequence encoding another initiation codon having a higher expression rate of the polypeptide compared to the endogenous initiation codon, but is not limited thereto.

[0082] The 4) and 5) methods of modifying the amino acid sequence or the polynucleotide sequence may be achieved by inducing a modification on the sequence through deletion, insertion, non-conservative or conservative substitution of the amino acid sequence of the protein or the polynucleotide sequence encoding the protein, or a combination thereof to enhance the activity of the protein, or by replacing the sequence with an amino acid sequence or polynucleotide sequence modified to have a stronger activity, or an amino acid sequence or polynucleotide sequence modified to enhance the activity, but are not limited thereto. The replacement may specifically be performed by inserting the polynucleotide into the chromosome by homologous recombination, but is not limited thereto. The vector used herein may further include a selection marker to confirm the insertion into the chromosome. The selection marker is the same as described above.

[0083] The 6) method of introducing a foreign polynucleotide having the activity of the protein may be achieved by introducing into a host cell a foreign polynucleotide encoding a protein that exhibits the same or similar activity to the protein. The foreign polynucleotide may be used without limitation regardless of its origin or sequence as long as it exhibits the same or similar activity to the protein. The introduction may be performed by those of ordinary skill in the art by appropriately selecting a transformation method

known in the art, and the expression of the introduced polynucleotide in the host cell enables to produce the protein, thereby increasing its activity.

[0084] The 7) method of codon-optimization of the polynucleotide encoding the protein may be achieved by codon-optimization of an endogenous polynucleotide to increase the transcription or translation within a host cell, or by optimizing the codons thereof such that the optimized transcription and translation of the foreign polynucleotide can be achieved within the host cell.

[0085] In addition, the 8) method of analyzing the tertiary structure of the protein and thereby selecting and modifying the exposed site, or chemically modifying the same may be achieved, for example, by comparing the sequence information of the protein to be analyzed with a database, in which the sequence information of known proteins is stored, to determine template protein candidates according to the degree of sequence similarity, and thus confirming the structure based on the information, thereby selecting and transforming or modifying the exposed site to be modified or chemically modified.

[0086] Such enhancement of the protein activity may mean that the activity or concentration (expression level) of the protein is increased relative to the activity or concentration of the protein expressed in a wild-type or a microorganism before modification, or that the amount of product produced from the protein is increased, but is not limited thereto.

[0087] As used herein, the term "vector" may include a DNA construct containing the nucleotide sequence of a polynucleotide encoding the target polypeptide operably linked to a suitable expression regulatory region (expression regulatory sequence) so as to be able to express the target polypeptide in a suitable host cell. The expression regulatory sequence may include a promoter capable of initiating transcription, any operator sequence for regulating the transcription, a sequence encoding a suitable mRNA ribosome-binding site, and a sequence for regulating termination of transcription and translation. Once transformed into a suitable host cell, the vector may replicate or function independently from the host genome, or may integrate into genome thereof.

[0088] The vector used in the present disclosure is not particularly limited, and any vector known in the art may be used. Examples of the vector typically used may include natural or recombinant plasmids, cosmids, viruses, and bacteriophages. For example, as a phage vector or cosmid vector, pWE15, M13, MBL3, MBL4, IXII, ASHII, APII, t10, t11, Charon4A, and Charon21A, etc. may be used; and as a plasmid vector, those based on pDC, pBR, pUC, pBluescriptII, pGEM, pTZ, pCL and pET, etc. may be used. Specifically, pDZ, pDC, pDCM2, pACYC177, pACYC184, pCL, pECCG117, pUC19, pBR322, pMW118, pCC1BAC vector, etc. may be used.

[0089] In one example, a polynucleotide encoding a target polypeptide may be inserted into the chromosome through a vector for intracellular chromosomal insertion. The insertion of the polynucleotide into the chromosome may be performed by any method known in the art, for example, by homologous recombination, but is not limited thereto. The vector may further include a selection marker to confirm the insertion into the chromosome. The selection marker is for selecting the cells transformed with the vector, that is, for confirming whether the target nucleic acid molecule has

been inserted, and markers that provide selectable phenotypes, such as drug resistance, auxotrophy, resistance to cell toxic agents, or expression of surface proteins, may be used. Only cells expressing the selection marker are able to survive or to show different phenotypes under the environment treated with the selective agent, and thus the transformed cells may be selected.

[0090] As used herein, the term “transformation” refers to the introduction of a vector containing a polynucleotide encoding a target polypeptide into a host cell or a microorganism so that the polypeptide encoded by the polynucleotide can be expressed in the host cell. As long as the transformed polynucleotide can be expressed in the host cell, it does not matter whether the transformed polynucleotide is integrated into the chromosome of the host cell and located therein or located extrachromosomally, and both cases can be included. Further, the polynucleotide may include DNA and/or RNA encoding the target polypeptide. The polynucleotide may be introduced in any form, as long as it can be introduced into the host cell and expressed therein. For example, the polynucleotide may be introduced into the host cell in the form of an expression cassette, which is a gene construct including all elements required for its autonomous expression. The expression cassette may commonly include a promoter operably linked to the polynucleotide, a transcription terminator, a ribosome-binding site, or a translation terminator. The expression cassette may be in the form of a self-replicable expression vector. Additionally, the polynucleotide may be introduced into the host cell as it is and operably linked to sequences required for expression in the host cell, but is not limited thereto.

[0091] Further, as used herein, the term “operably linked” means that the polynucleotide sequence is functionally linked to a promoter sequence that initiates and mediates transcription of the polynucleotide encoding the target variant of the present disclosure.

[0092] The modification of a part or all of the polynucleotide in the microorganism of the present disclosure may be achieved by (a) homologous recombination using a vector for chromosomal insertion in the microorganism or genome editing using engineered nuclease (e.g., CRISPR-Cas9) and/or (b) may be induced by light, such as ultraviolet rays and radiation, etc., and/or chemical treatments, but is not limited thereto. The method of modifying a part or all of the gene may include a method using DNA recombination technology. For example, a part or all of the gene may be deleted by injecting a nucleotide sequence or a vector containing a nucleotide sequence homologous to the target gene into a microorganism to induce homologous recombination. The injected nucleotide sequence or the vector may include a dominant selection marker, but is not limited thereto.

[0093] Another aspect of the present disclosure provides a method for producing L-arginine, including: culturing a recombinant microorganism of the genus *Corynebacterium*, in which the activity of a protein including an amino acid sequence of SEQ ID NO: 1 is weakened, in a medium.

[0094] The protein including the amino acid sequence of SEQ ID NO: 1, weakening, and microorganism, etc. are the same as described in the other aspects above.

[0095] The microorganism of the genus *Corynebacterium* may be *Corynebacterium glutamicum*, but is not limited thereto.

[0096] The microorganism may be a microorganism in which the activity of arginine repressor (ArgR) is further

weakened or the argR gene is further deleted, but is not limited thereto, and this is as described in the other aspects above.

[0097] As used herein, the term “cultivation” means that the microorganism of the genus *Corynebacterium* of the present disclosure is grown under appropriately controlled environmental conditions. The cultivation process of the present disclosure may be performed in a suitable culture medium and culture conditions known in the art. Such a cultivation process may be easily adjusted for use by those skilled in the art according to the strain to be selected. Specifically, the cultivation may be a batch culture, a continuous culture, and/or a fed-batch culture, but is not limited thereto.

[0098] As used herein, the term “medium” refers to a mixture of materials which contains nutrient materials required for the cultivation of the microorganism of the genus *Corynebacterium* of the present disclosure as a main ingredient, and it supplies nutrient materials and growth factors, along with water that is essential for survival and growth. Specifically, the medium and other culture conditions used for culturing the microorganism of the genus *Corynebacterium* of the present disclosure may be any medium used for conventional cultivation of microorganisms without any particular limitation. However, the microorganism of the genus *Corynebacterium* of the present disclosure may be cultured under aerobic conditions in a conventional medium containing an appropriate carbon source, nitrogen source, phosphorus source, inorganic compound, amino acid, and/or vitamin, while adjusting temperature, pH, etc.

[0099] Specifically, the culture medium for the microorganism of the genus *Corynebacterium* can be found in the literature (“Manual of Methods for General Bacteriology” by the American Society for Bacteriology (Washington D.C., USA, 1981)).

[0100] In the present disclosure, the carbon source may include carbohydrates, such as glucose, saccharose, lactose, fructose, sucrose, maltose, etc.; sugar alcohols, such as mannitol, sorbitol, etc.; organic acids, such as pyruvic acid, lactic acid, citric acid, etc.; amino acids, such as glutamic acid, methionine, lysine, etc. Additionally, the carbon source may include natural organic nutrients such as starch hydrolysate, molasses, blackstrap molasses, rice bran, cassava, sugar cane molasses, corn steep liquor, etc. Specifically, carbohydrates such as glucose and sterilized pretreated molasses (i.e., molasses converted to reducing sugar) may be used, and in addition, various other carbon sources in an appropriate amount may be used without limitation. These carbon sources may be used alone or in a combination of two or more kinds, but are not limited thereto.

[0101] The nitrogen source may include inorganic nitrogen sources, such as ammonia, ammonium sulfate, ammonium chloride, ammonium acetate, ammonium phosphate, ammonium carbonate, ammonium nitrate, etc.; amino acids, such as glutamic acid, methionine, glutamine, etc.; and organic nitrogen sources, such as peptone, NZ-amine, meat extract, yeast extract, malt extract, corn steep liquor, casein hydrolysate, fish or decomposition product thereof, defatted soybean cake or decomposition product thereof, etc. These nitrogen sources may be used alone or in a combination of two or more kinds, but are not limited thereto.

[0102] The phosphorus source may include monopotassium phosphate, dipotassium phosphate, or corresponding

sodium-containing salts, etc. Examples of the inorganic compound may include sodium chloride, calcium chloride, iron chloride, magnesium sulfate, iron sulfate, manganese sulfate, calcium carbonate, etc. Additionally, amino acids, vitamins, and/or appropriate precursors, etc. may be included. These constituting ingredients or precursors may be added to a medium in a batch or continuous manner, but these phosphorus sources are not limited thereto.

[0103] Further, the pH of a medium may be adjusted during the cultivation of the microorganism of the genus *Corynebacterium* of the present disclosure by adding a compound such as ammonium hydroxide, potassium hydroxide, ammonia, phosphoric acid, sulfuric acid, etc. to the medium in an appropriate manner. Additionally, during the cultivation, an antifoaming agent such as fatty acid polyglycol ester may be added to prevent foam generation. In addition, oxygen or oxygen-containing gas may be injected into the medium in order to maintain an aerobic state of the medium; or nitrogen, hydrogen, or carbon dioxide gas may be injected without the injection of gas in order to maintain an anaerobic or microaerobic state of the medium, but the gas is not limited thereto.

[0104] The medium temperature in the cultivation of the present disclosure may be in a range from 20° C. to 45° C., and specifically, from 25° C. to 40° C., and the cultivation may be carried out for about 10 to 160 hours, but is not limited thereto.

[0105] The L-arginine produced by the cultivation of the present disclosure may be released into the medium or remain in the cells.

[0106] The method for producing L-arginine of the present disclosure may further include a step of preparing the microorganism of the genus *Corynebacterium* of the present disclosure, a step of preparing a medium for culturing the microorganism, or a combination thereof (regardless of the order, in any order), for example, prior to the culturing step.

[0107] The method for producing L-arginine of the present disclosure may further include a step of recovering L-arginine from the culture medium (medium on which the culture was grown) or the cultured microorganism of the genus *Corynebacterium*. *The recovering step may be further included after the culturing step.*

[0108] In the recovering step, desired L-arginine may be collected using the method of culturing a microorganism of the present disclosure, for example, using a suitable method known in the art according to a batch culture, continuous culture, or fed-batch culture method. For example, methods such as centrifugation, filtration, treatment with a protein crystallizing precipitant (salting-out method), extraction, ultrasonic disruption, ultrafiltration, dialysis, various kinds of chromatography such as molecular sieve chromatography (gel filtration), adsorption chromatography, ion exchange chromatography, affinity chromatography, etc., HPLC, or a combination thereof may be used, and the desired L-arginine can be recovered from the medium or microorganisms using suitable methods known in the art.

[0109] Further, the method for producing L-arginine of the present disclosure may further include a purification process, which may be performed using an appropriate method known in the art. In one example, when the method for producing L-arginine of the present disclosure includes both a recovering step and a purification step, the recovering step and the purification step may be performed continuously or

intermittently regardless of the order, or simultaneously or may be integrated into one step, but the method is not limited thereto.

[0110] Still another aspect of the present disclosure provides a composition for producing L-arginine, including: a microorganism of the genus *Corynebacterium*, in which the activity of the protein including the amino acid sequence of SEQ ID NO: 1 of the present disclosure is weakened; a medium for culturing the same; or a combination thereof.

[0111] The composition of the present disclosure may further include any suitable excipient commonly used in the composition for producing amino acids, and such excipient may be, for example, a preservative, a wetting agent, a dispersing agent, a suspending agent, a buffering agent, a stabilizing agent, or an isotonic agent, etc., but is not limited thereto.

[0112] In the composition of the present disclosure, the protein including the amino acid sequence of SEQ ID NO: 1, weakening, microorganism, cultivation, and medium, etc. are the same as described in the other aspects above.

[0113] Yet another aspect of the present disclosure provides a method for producing a microorganism of the genus *Corynebacterium*, including weakening a protein including an amino acid sequence of SEQ ID NO: 1.

[0114] Even another aspect of the present disclosure provides the use of L-arginine production of a microorganism of the genus *Corynebacterium*, in which the activity of a protein including an amino acid sequence of SEQ ID NO: 1 is weakened.

[0115] In the method for producing the microorganism of the genus *Corynebacterium* and the use of L-arginine production of the microorganism of the genus *Corynebacterium*, the protein including the amino acid sequence of SEQ ID NO: 1, weakening, and microorganism, etc. are the same as described in the other aspects above.

MODE FOR CARRYING OUT THE INVENTION

[0116] Hereinafter, the present disclosure will be described in detail by way of Examples. However, these Examples are merely preferred Examples given for illustrative purposes, and thus, the scope of the present disclosure is not intended to be limited to or by these Examples. Meanwhile, technical features which are not described herein can be sufficiently understood and easily carried out by those skilled in the art in the technical field of the present disclosure or in a similar technical field.

Example 1: Construction of Random Mutation Library Using Transposon and Screening Thereof

[0117] In order to obtain a strain having increased L-arginine producing ability, the plasmids obtained using EZ-Tn5™<R6Kyor/KAN-2> Tnp Transposome™ Kit (Epicentre) were transformed based on the *Corynebacterium glutamicum* KCCM10741P (Korean Patent No. 0791659) as the parent strain by electroporation (*Appl. Microbiol. Biotechnol.* (1999) 52:541-545) and spread on a complex medium plate containing kanamycin (25 mg/L) to thereby obtain about 20,000 colonies.

<Complex Medium Plate (pH 7.0)>

[0118] Glucose 10 g, Peptone 10 g, Beef extract 5 g, Yeast extract 5 g, Brain Heart Infusion 18.5 g, NaCl 2.5 g, Urea 2 g, Sorbitol 91 g, Aga 20 g (per liter of distilled water)

[0119] The top 10 mutant strains with improved L-arginine producing ability compared to the *Corynebacterium glutamicum* KCCM10741P, the parent strain, were selected using the ninhydrin method (Moore, S., Stein, W. H., Photometric ninhydrin method for use in the chromatography of amino acids. *J. Biol. Chem.* 1948, 176, 367-388).

Example 2: Analysis of L-Arginine Producing Ability of Selected Random Mutant Strains

[0120] In order to finally select strains with reproducibly increased L-arginine producing ability among the 10 mutant strains selected in Example 1, flask culture was performed using the following medium. After completion of the culture, the L-arginine concentration in the culture medium was analyzed using HPLC, and the L-arginine production concentration of each mutant strain is shown in Table 1 below.

<Production Medium (pH 7.0)>

[0121] Glucose 6%, Ammonium sulfate 3%, Monopotassium phosphate 0.1%, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.2%, CSL (corn steep liquor) 1.5%, NaCl 1%, Yeast extract 0.5%, Biotin 100 mg/L, CaCO_3 3%, (per liter of distilled water).

TABLE 1

L-Arginine Production Concentration of 10 Selected Random Mutant Strains		L-Arginine (g/L)			
Strain		Batch 1	Batch 2	Batch 3	Average
Control	KCCM10741P	3.1	3.2	3	3.1
1	KCCM10741P/mt-1	2.9	3	2.8	2.9
2	KCCM10741P/mt-2	3	2.9	2.7	2.9
3	KCCM10741P/mt-3	3.1	3	3.1	3.1
4	KCCM10741P/mt-4	2.8	2.9	3.2	3.0
5	KCCM10741P/mt-5	3.5	3.5	3.4	3.5
6	KCCM10741P/mt-6	2.9	3.1	3.2	3.1
7	KCCM10741P/mt-7	3	3.2	2.8	3.0
8	KCCM10741P/mt-8	2.9	2.9	3	2.9
9	KCCM10741P/mt-9	3.1	3.1	2.8	3.0
10	KCCM10741P/mt-10	3	2.9	2.9	2.9

Among the 10 mutant strains selected, KCCM10741P/mt-5 was finally selected as a strain with significantly improved L-arginine producing ability.

Example 3: Identification of Causes of Increase in L-Arginine Producing Ability of Finally Selected Strains

[0122] The gene inactivated by random insertion of a transposon in the KCCM10741P/mt-5 of Example 2 was identified.

[0123] Specifically, the genomic DNA of KCCM10741P/mt-5 was extracted, cleaved and ligated to transform into *E. coli* DH5 α , and plated on LB solid medium containing kanamycin (25 mg/L). After selecting 20 types of transformed colonies, a plasmid containing part of an unknown gene was obtained. As a result of analyzing the nucleotide sequence using primer 1 (SEQ ID NO: 3) and primer 2 (SEQ ID NO: 4) of the EZ-Tn5™ <R6Kyor/KAN-2> Tnp Transposome™ Kit, it was found that the gene including the nucleotide sequence of SEQ ID NO: 2 was inactivated based on the nucleotide sequence registered in the NIH GenBank of the National Institutes of Health.

TABLE 2

Primers Used	Nucleotide Sequences
Primer 1 (SEQ ID NO: 3)	ACCTACAACAAAGCTCTCATCAACC
Primer 2 (SEQ ID NO: 4)	CTACCCTGTGGAACACCTACATCT

[0124] As a result of a BLAST search of SEQ ID NO: 2 of the wild-type *Corynebacterium glutamicum* ATCC 13869, it was confirmed that it was the NCgl2608 gene.

Example 4: Construction of Recombinant Vector for NCgl2608 Gene Deletion

[0125] In order to reconfirm the effect of inactivation of the gene including the nucleotide sequence of SEQ ID NO: 2 on L-arginine production, a recombinant vector was constructed for deletion of the NCgl2608 gene on the chromosome of a strain of the genus *Corynebacterium*.

[0126] First, primers 3 to 6 were synthesized to construct fragments for deletion of the above genes, and these are shown in Table 3.

TABLE 3

Primers 3 to 6 for Production of Fragments for Gene Deletion			
Primers Used	Nucleotide Sequences		
Primer 3 (SEQ ID NO: 5)	tccggtaccgcggggtaccTC GGTGGCGGTTTCGGC		
Primer 4 (SEQ ID NO: 6)	TTGCAGAAAGGGATACCAG GGCTAGCTCATAGAC		
Primer 5 (SEQ ID NO: 7)	GGTATCCCTTTCTGCAACG ATGTGCTGTGTGACG		
Primer 6 (SEQ ID NO: 8)	tgcaggtcgactctagaAG GACCTGGAAAAGGGC		

[0127] In order to delete the ORF region based on SEQ ID NO: 2, primer 3 (SEQ ID NO: 5), primer 4 (SEQ ID NO: 6), primer 5 (SEQ ID NO: No. 7), and primer 6 (SEQ ID NO: 8) (Table 2) were synthesized to have BamHI restriction enzyme site at the 5' end and XbaI restriction enzyme site at the 3' end, and PCR was performed based on the chromosomal DNA of the wild-type *Corynebacterium glutamicum* ATCC 13869 as a template (Sambrook et al., *Molecular Cloning, a Laboratory Manual* (1989), Cold Spring Harbor Laboratories). Based on this, a DNA fragment was obtained in which the top 600 bp and the bottom 600 bp of the portion encoding the protein encoded by the nucleotide sequence of NCgl2608 were connected. In particular, the PCR was performed under PCR conditions of 30 cycles of denaturation at 95° C. for 30 seconds; annealing at 50° C. for 30 seconds; and polymerization at 72° C. for 1 minutes, and then polymerization at 72° C. for 7 minutes. The pDZ vector (Korean Patent No. 10-0924065), which cannot replicate in *Corynebacterium glutamicum*, and the fragment amplified by PCR were treated with restriction enzymes BamHI and

XbaI and ligated using a DNA ligation enzyme, and transformed into *E. coli* DH5a and plated on LB solid medium containing kanamycin (25 mg/L).

[0128] After selecting colonies transformed with the plasmid into which the target gene was inserted through PCR, a plasmid was obtained using a plasmid extraction method, and this plasmid was named pDZ-ΔNCgl2608.

Example 5: Construction of Strain with Deleted NCgl2608 Gene Derived from *Corynebacterium glutamicum* KCCM10741P and Evaluation of L-Arginine Producing Ability

[0129] pDZ-ΔNCgl2608 was transformed into *Corynebacterium glutamicum* KCCM10741P, an L-arginine producing strain, by homologous recombination on the chromosome (van der Rest et al., *Appl Microbiol Biotechnol* 52:541-545, 1999).

[0130] Thereafter, secondary recombination was performed on a solid plate medium containing 4% sucrose. A strain with deleted gene of SEQ ID NO: 2 on the chromosome was identified through PCR using primers 3 and 6 based on the *Corynebacterium glutamicum* transformants for which secondary recombination was completed. The recombinant strain was named *Corynebacterium glutamicum* KCCM10741P-ΔNCgl2608.

[0131] In order to analyze the L-arginine producing ability of the *Corynebacterium glutamicum* KCCM10741P-ΔNCgl2608 strain constructed above, the strain was cultured with *Corynebacterium glutamicum* KCCM10741P, the parent strain, in the following manner.

[0132] The *Corynebacterium glutamicum* KCCM10741P, the parent strain, and the *Corynebacterium glutamicum* KCCM10741P-ΔNCgl2608 strain constructed in Example 6 were inoculated into a 250 mL corner-baffle flask containing 25 mL of the production medium below, and then cultured with shaking at 200 rpm at 30° C. for 20 hours. Then, 1 mL of seed culture was inoculated into a 250 mL corner-baffle flask containing 24 mL of production medium and cultured under shaking at 200 rpm at 30° C. for 72 hours. The compositions of the seed medium and production medium are as follows.

<Seed Medium (pH 7.0)>

[0133] Glucose 20 g, Pepetone 10 g, Yeast extract 5 g, Urea 1.5 g, KH₂PO₄ 4 g, K₂HPO₄ 8 g, MgSO₄·7H₂O 0.5 g, Biotine 100 μg, Thiamine-HCl 1000 μg, Calcium pantothenate 2000 μg, Nicotinamide 2000 μg (per liter of distilled water)

<Production Medium (pH 7.0)>

[0134] Glucose 6%, Ammonium sulfate 3%, Monopotassium phosphate 0.1%, MgSO₄·7H₂O 0.2%, CSL (corn steep liquor) 1.5%, NaCl 1%, Yeast extract 0.5%, Biotin 100 mg/L

[0135] After completion of the culture, the production amount of L-arginine was measured by HPLC (Waters 2478), and the concentration of L-arginine analyzed is shown in Table 4 below.

TABLE 4

Analysis of L-Arginine Producing Ability of KCCM10741P-derived KCCM10741P-ΔNCgl2608					
		L-Arginine (g/L)			
	Strains	Batch 1	Batch 2	Batch 3	Average
Control	KCCM10741P	3	2.9	3.1	3.0
Experimental group	KCCM10741P-ΔNCgl2608	3.5	3.7	3.6	3.6

[0136] As shown in the results above, when the gene including the nucleotide sequence of SEQ ID NO: 2 was deleted from the *Corynebacterium glutamicum* KCCM10741P, an L-arginine-producing strain, it was confirmed that the L-arginine producing ability was increased by an average of 20.0% compared to the parent strain.

[0137] Accordingly, it was confirmed that L-arginine producing ability can be improved by deleting the gene including the nucleotide sequence of SEQ ID NO: 2 in the microorganism of the genus *Corynebacterium*.

Example 6: Construction of Strain with Deleted NCgl2608 Gene Derived from *Corynebacterium glutamicum* CJ1R and Evaluation of L-Arginine Producing Ability

[0138] In order to confirm whether the same effect could be obtained in other strains of *Corynebacterium glutamicum* that produce L-arginine, an attempt was made to introduce one type of mutation (ΔargR) into the wild strain (ATCC 13869).

[0139] The recombinant vector for deletion of the argR gene was constructed in the same manner as Example 4, and the primers used to construct the vector are shown in Table 5

TABLE 5

Primers 7 to 10 for Production of Fragments for Deletion of argR gene	
Primers Used	Nucleotide Sequences
Primer 7 (SEQ ID NO: 9)	tccggtaccgccgggatccA GCAGGCCCTTAAGGGTAAG
Primer 8 (SEQ ID NO: 10)	AGGGGCGCTGTCTTACCT CGGCTGGTGGGCCAGC
Primer 9 (SEQ ID NO: 11)	GGTAAGACAGCGCCCTTA GTTCAGGCTTGTTAATC
Primer 10 (SEQ ID NO: 12)	tgcaggtcgactctagaA CCGTTGAACGTGTGCCAG

[0140] After selecting the plasmid into which the target gene was inserted through PCR, this plasmid was named pDZ-ΔargR. The wild-type *Corynebacterium glutamicum* strain was transformed using pDZ-ΔargR in the same manner as in Example 6, and the strain with argR deletion on the chromosome was identified through PCR using primers 7 and 10 based on the transformants. The recombinant strain was named CJ1R, and a strain with deleted NCgl2608 gene was constructed based on the *Corynebacterium glutamicum*

CJ1R, which had arginine producing ability, in the same manner as in Example 6, and then was named CJ1R-ΔNCgl2608.

[0141] The CJ1R-ΔNCgl2608 with deleted NCgl2608 gene was cultured in the same manner as in Example 5 above, and after completion of the culture, the production of L-arginine was measured by HPLC (Waters 2478), and the concentration of L-arginine analyzed is shown in Table 6 below.

TABLE 6

L-Arginine Producing Ability of CJ1R-Derived CJ1R-ΔNCgl2608					
Strains		L-Arginine (g/L)			
		Batch 1	Batch 2	Batch 3	Average
Control	CJ1R	1.3	1.3	1.4	1.33
Experimental group	CJ1R-ΔNCgl2608	1.6	1.5	1.6	1.56

[0142] As shown in the results above, it was confirmed that when the NCgl2608 gene was deleted from the *Coryne-*

bacterium glutamicum CJ1R, an L-arginine producing strain, the L-arginine producing ability was increased by an average of 17.3%.

[0143] Accordingly, as in the results of Example 5, it was confirmed that L-arginine producing ability can be improved by deleting the NCgl2608 gene in the microorganism of the genus *Corynebacterium*.

[0144] The CJ1R-ΔNCgl2608 strain was named CA06-7204 and deposited at the Korean Culture Center of Microorganisms (KCCM), a Depositary Authority under the Budapest Treaty, on Feb. 23, 2021, with Accession No. KCCM12961P.

[0145] From the foregoing, those of ordinary skill in the art to which the present disclosure belongs will recognize that the present disclosure may be embodied in other specific forms without departing from its spirit or essential characteristics. The described embodiments are to be considered in all respects only as illustrative and not restrictive. The scope of the present disclosure is therefore indicated by the appended claims rather than by the foregoing description. All changes which come within the meaning and range of equivalency of the claims are to be embraced within the scope of the present disclosure.

SEQUENCE LISTING

```

<160> NUMBER OF SEQ ID NOS: 20

<210> SEQ ID NO 1
<211> LENGTH: 513
<212> TYPE: PRT
<213> ORGANISM: Unknown
<220> FEATURE:
<223> OTHER INFORMATION: NCgl2608 amino acid

<400> SEQUENCE: 1

Met Ser Asp Leu Gly Pro Ile Trp Arg Trp Leu Leu Val Ser Val
1             5             10             15

Ser Ile Cys Ala Ala Ser Gly Leu Val Tyr Glu Leu Ala Leu Val Ser
20            25            30

Leu Ser Thr Ser Leu Asn Gly Gly Gly Ile Val Glu Thr Ser Leu Ile
35            40            45

Val Ala Gly Tyr Val Ala Ala Leu Gly Leu Gly Ala Leu Leu Val Lys
50            55            60

Pro Phe Leu Asn Trp Pro Ala Gln Thr Phe Leu Gly Val Glu Thr Leu
65            70            75            80

Leu Gly Leu Ile Gly Gly Cys Ser Ala Leu Val Leu Tyr Phe Thr Phe
85            90            95

Ala Thr Ile Gly Gln Ser Leu Trp Ile Leu Val Ile Ala Thr Ala Ala
100           105           110

Ile Gly Ile Leu Val Gly Ala Glu Leu Pro Leu Leu Met Thr Met Ile
115           120           125

Gln Gln Gly Arg Leu Ala Asp Ala Lys Thr Thr Gly Ser Leu Val Ala
130           135           140

Thr Leu Asn Ala Ala Asp Tyr Leu Gly Ala Leu Leu Gly Gly Leu Ala
145           150           155           160

Trp Pro Phe Val Leu Leu Pro Trp Leu Gly Met Met Arg Gly Ala Ala
165           170           175

Ala Ala Gly Met Ile Asn Leu Val Ala Ala Leu Phe Val Gly Cys Val

```

-continued

180				185				190							
Leu	Leu	Arg	His	Leu	Leu	Pro	Arg	Thr	His	Phe	Phe	Val	Ser	Val	Val
	195						200					205			
Ala	Leu	Leu	Leu	Ala	Ile	Ala	Ala	Leu	Ala	Thr	Val	Leu	Val	Lys	Ser
	210				215						220				
Asp	Gly	Ile	Val	Ala	Thr	Ala	Arg	Ala	Gln	Leu	Tyr	Arg	Asp	Pro	Val
	225				230					235					240
Ile	Tyr	Ser	His	Gln	Ser	Asp	Tyr	Gln	Asp	Ile	Val	Val	Thr	Glu	Arg
			245						250					255	
Gly	Lys	Asp	Arg	Arg	Leu	Tyr	Leu	Asn	Gly	Gly	Leu	Gln	Tyr	Ser	Thr
			260						265					270	
Arg	Asp	Gln	His	Arg	Tyr	Thr	Glu	Ser	Leu	Val	Tyr	Pro	Ser	Leu	Asn
			275				280							285	
Pro	Glu	Ala	Glu	Ser	Val	Leu	Ile	Ile	Gly	Gly	Gly	Asp	Gly	Leu	Ala
	290					295					300				
Ala	Arg	Glu	Leu	Leu	Arg	Phe	Pro	Ser	Met	Gln	Ile	Thr	Gln	Val	Glu
	305				310					315					320
Leu	Asp	Pro	Glu	Val	Ile	Glu	Val	Ala	Asn	Thr	Val	Leu	Arg	Ser	Asp
			325						330					335	
Asn	Gly	Gly	Ala	Met	Glu	Asp	Pro	Arg	Val	Ser	Ile	Ile	Val	Asp	Asp
			340						345					350	
Ala	Phe	Thr	Trp	Leu	Arg	Ser	Gly	Gly	Asn	Asn	Gly	Glu	Thr	Tyr	Asp
		355					360					365			
Ser	Ile	Ile	Ile	Asp	Leu	Pro	Asp	Pro	Asn	Asn	Asp	Thr	Met	Ala	Arg
	370				375						380				
Leu	Tyr	Ser	Glu	Glu	Phe	Tyr	Thr	Leu	Ala	Arg	Ala	Arg	Leu	Asn	Glu
	385				390					395					400
Gln	Gly	Arg	Met	Val	Val	Gln	Ser	Ser	Ser	Ala	Tyr	Thr	Thr	Pro	Asp
			405						410					415	
Val	Phe	Trp	Arg	Val	Gly	Ala	Thr	Leu	Lys	Ser	Ala	Gly	Cys	Glu	Gln
			420						425					430	
Val	Ile	Pro	Tyr	His	Val	His	Val	Pro	Thr	Phe	Gly	Asp	Trp	Gly	Phe
		435					440					445			
Gln	Leu	Cys	Gly	Pro	Ala	Asp	Met	Glu	Leu	Glu	Leu	Arg	Glu	Asp	Thr
	450					455					460				
Pro	Pro	Leu	Thr	Phe	Leu	Asn	Asp	Glu	Val	Leu	Val	Ala	Ala	Gly	Val
	465				470					475					480
Phe	Gly	Leu	Asp	Asn	Gln	Pro	Arg	Glu	Leu	Glu	Pro	Ser	Thr	Leu	Asp
			485						490					495	
His	Pro	Arg	Val	Val	Glu	Asp	Leu	Arg	Lys	Gly	Tyr	Arg	Glu	Ser	Gly
			500						505					510	

Asp

<210> SEQ ID NO 2

<211> LENGTH: 1542

<212> TYPE: DNA

<213> ORGANISM: Unknown

<220> FEATURE:

<223> OTHER INFORMATION: NCgl2608 nucleotide

<400> SEQUENCE: 2

atgtctgatt taggacccat ctggcgctgg ctgttattag tttcgtctc catttgtgcg

60

-continued

gcatcggggc tggctctatga gctagccctg gtatcccttt ccaccagctt gaacgggtggc	120
ggaattgtag aaacctccct catcgctgca ggttatgtag ctgcccttgg acttggtgca	180
ctgctgggtca agccgtttct caactggcct gcacaaacct tctcgggtgt ggaaaccttg	240
ctgggactca tcgggtggtg ctccgcctg gtgctatatt tcacctttgc gaccatcggc	300
cagtccttgt ggattctggt gattgccacc gctgcaattg gcactcctgg cggcgctgaa	360
cttcactgc tgatgacct gatccagcaa ggcgcctcg ccgacgcaa aactacagga	420
tctctggtg ccacctgaa tgctgctgat taccttggcg cacttttagg tggcctggcc	480
tggccttttg tgttgcctgc gtggcttggc atgatgcgcg gtgcgcgagc agccggaatg	540
atcaacctcg ttgcagcact atcgtgggc tgtgtgctgc tgcgacattt gcttcgcgc	600
acccacttct tcgtatccgt ggtggcgctt cttctcgca tcgcagcgtg tgccaccgtg	660
tgggtgaaat ccgacgggat cgttgccacc gccgcgcac agctctaccg cgaccccggtg	720
atctattcac accaatctga ctaccaagac atcgtagtga cagaacgagg caaagaccga	780
cgctctacc tcaatggcgg tttgcagtat tccactcgtg accagcatag atatacagaa	840
tcactggtgt atccaagcct taatccagag gcagaatcgg tgtaaatcat cggcggtggc	900
gatggcctcg cagcacggga actcctccga tccccatcaa tgcagatcac ccaagttgaa	960
ttagaccag aagtcacga agtagccaac actgtgctgc gttctgacaa tgggggagcg	1020
atggaagatc cccgcgtgct catcatcgct gacgacgcct tcacctggct tcgctccggc	1080
ggaaacaacg gcgaaactta cgattccatc atcatcgatc tccccgacc aaacaacgac	1140
accatggcca ggctgtattc agaagagttc tacaccttg cccgagcacg actgaatgaa	1200
caaggccgca tgggtggtca atcctccagc gcgtacacca cccagatgt gttctggcga	1260
gttgagacaa ccttgaaatc ggcgggctgt gaacaagtca tcccatatca cgtgcatgtt	1320
cccacatttg ggcactgggg cttccaactg tgtggcctg ccgacatgga attagagctt	1380
cgggaagaca ccccgccact gactttcctt aatgatgaag ttctggtggc tgctggggtg	1440
tttgggttgg ataatacagc tcgtgaattg gaaccttcca cgctggatca tccccgcgtg	1500
gtggaggatc tgcgcaagg atatcgcgaa tcaggcgact ag	1542

<210> SEQ ID NO 3
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer 1

<400> SEQUENCE: 3

acctacaaca aagctctcat caacc	25
-----------------------------	----

<210> SEQ ID NO 4
 <211> LENGTH: 24
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer 2

<400> SEQUENCE: 4

ctaccctgtg gaacacctac atct	24
----------------------------	----

<210> SEQ ID NO 5

-continued

<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer 3

<400> SEQUENCE: 5

tcggtaccgcg gggatccctcg gtggcggttt cggc 34

<210> SEQ ID NO 6
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer 4

<400> SEQUENCE: 6

ttgcagaaaag ggataccagg gctagctcat agac 34

<210> SEQ ID NO 7
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer 5

<400> SEQUENCE: 7

ggtatccctt tctgcaacga tgtgctgtgt gacg 34

<210> SEQ ID NO 8
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer 6

<400> SEQUENCE: 8

tgcaggtcga ctctagaagg acctggaaaa gggc 34

<210> SEQ ID NO 9
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer 7

<400> SEQUENCE: 9

tcggtaccgcg gggatccagc aggccttaag ggtaag 36

<210> SEQ ID NO 10
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer 8

<400> SEQUENCE: 10

aggggcgctg tcttacctcg gctggtagggc cagc 34

<210> SEQ ID NO 11
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: primer 9

<400> SEQUENCE: 11

ggtaagacag cgccctagtg tcaaggcttg ttaatc 36

<210> SEQ ID NO 12

<211> LENGTH: 37

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer 10

<400> SEQUENCE: 12

tgcaggtcga ctctagaacc gttgaactgc ttgccag 37

<210> SEQ ID NO 13

<211> LENGTH: 171

<212> TYPE: PRT

<213> ORGANISM: Unknown

<220> FEATURE:

<223> OTHER INFORMATION: argR amino acid

<400> SEQUENCE: 13

Met Ser Leu Gly Ser Thr Pro Ser Thr Pro Glu Asn Leu Asn Pro Val
1 5 10 15

Thr Arg Thr Ala Arg Gln Ala Leu Ile Leu Gln Ile Leu Asp Lys Gln
20 25 30

Lys Val Thr Ser Gln Val Gln Leu Ser Glu Leu Leu Leu Asp Glu Gly
35 40 45

Ile Asp Ile Thr Gln Ala Thr Leu Ser Arg Asp Leu Asp Glu Leu Gly
50 55 60

Ala Arg Lys Val Arg Pro Asp Gly Gly Arg Ala Tyr Tyr Ala Val Gly
65 70 75 80

Pro Val Asp Ser Ile Ala Arg Glu Asp Leu Arg Gly Pro Ser Glu Lys
85 90 95

Leu Arg Arg Met Leu Asp Glu Leu Leu Val Ser Thr Asp His Ser Gly
100 105 110

Asn Ile Ala Met Leu Arg Thr Pro Pro Gly Ala Ala Gln Tyr Leu Ala
115 120 125

Ser Phe Ile Asp Arg Val Gly Leu Lys Glu Val Val Gly Thr Ile Ala
130 135 140

Gly Asp Asp Thr Val Phe Val Leu Ala Arg Asp Pro Leu Thr Gly Lys
145 150 155 160

Glu Leu Gly Glu Leu Leu Ser Gly Arg Thr Thr
165 170

<210> SEQ ID NO 14

<211> LENGTH: 516

<212> TYPE: DNA

<213> ORGANISM: Unknown

<220> FEATURE:

<223> OTHER INFORMATION: argR nucleotide

<400> SEQUENCE: 14

atgtcccttg gctcaacccc gtcaacaccg gaaaaactaa atcccgtagc tcgcaactgca 60

cgccaagctc tcatttttgca gatttttgac aaacaaaaag tcaccagcca ggtacaactg 120

tctgaattgc tgctggatga aggcacgat atcaccacagg ccaccttgctc ccgagatctc 180

-continued

```

gatgaactcg gtgcacgcaa ggttcgcccc gatgggggac gcgcctacta cgcggtcggc 240
ccagtagata gcacgccccg cgaagatctc cgggggtccgt cggagaagct gcgccgcatg 300
cttgatgaac tgctgggttc tacagatcat tccggcaaca tcgcgatgct gcgcacccccg 360
ccgggagctg cccagtacct ggcaagtctc atcgataggg tggggctgaa agaagtcgtt 420
ggcaccatcg ctggcgatga caccgttttc gttctcgccc gtgatccgct cacaggtaaa 480
gaactagggtg aattactcag cgggcgcacc acttaa 516

```

```

<210> SEQ ID NO 15
<211> LENGTH: 319
<212> TYPE: PRT
<213> ORGANISM: Unknown
<220> FEATURE:
<223> OTHER INFORMATION: argF amino acid

```

```

<400> SEQUENCE: 15

```

```

Met Thr Ser Gln Pro Gln Val Arg His Phe Leu Ala Asp Asp Asp Leu
1          5          10          15

Thr Pro Ala Glu Gln Ala Glu Val Leu Thr Leu Ala Ala Lys Leu Lys
20          25          30

Ala Ala Pro Phe Ser Glu Arg Pro Leu Glu Gly Pro Lys Ser Val Ala
35          40          45

Val Leu Phe Asp Lys Thr Ser Thr Arg Thr Arg Phe Ser Phe Asp Ala
50          55          60

Gly Ile Ala His Leu Gly Gly His Ala Ile Val Val Asp Ser Gly Ser
65          70          75          80

Ser Gln Met Gly Lys Gly Glu Thr Leu Gln Asp Thr Ala Ala Val Leu
85          90          95

Ser Arg Tyr Val Glu Ala Ile Val Trp Arg Thr Tyr Ala His Ser Asn
100         105         110

Phe His Ala Met Ala Glu Thr Ser Thr Val Pro Leu Val Asn Ser Leu
115         120         125

Ser Asp Asp Leu His Pro Cys Gln Ile Leu Ala Asp Leu Gln Thr Ile
130         135         140

Val Glu Asn Leu Ser Pro Glu Glu Gly Pro Ala Gly Leu Lys Gly Lys
145         150         155         160

Lys Ala Val Tyr Leu Gly Asp Gly Asp Asn Asn Met Ala Asn Ser Tyr
165         170         175

Met Ile Gly Phe Ala Thr Ala Gly Met Asp Ile Ser Ile Ile Ala Pro
180         185         190

Glu Gly Phe Gln Pro Arg Ala Glu Phe Val Glu Arg Ala Glu Lys Arg
195         200         205

Gly Gln Glu Thr Gly Ala Lys Val Val Val Thr Asp Ser Leu Asp Glu
210         215         220

Val Ala Gly Ala Asp Val Val Ile Thr Asp Thr Trp Val Ser Met Gly
225         230         235         240

Met Glu Asn Asp Gly Ile Asp Arg Thr Thr Pro Phe Val Pro Tyr Gln
245         250         255

Val Asn Asp Glu Val Met Ala Lys Ala Asn Asp Gly Ala Ile Phe Leu
260         265         270

His Cys Leu Pro Ala Tyr Arg Gly Lys Glu Val Ala Ala Ser Val Ile
275         280         285

```

-continued

Asp Gly Pro Ala Ser Lys Val Phe Asp Glu Ala Glu Asn Arg Leu His
290 295 300

Ala Gln Lys Ala Leu Leu Val Trp Leu Leu Ala His Gln Pro Arg
305 310 315

<210> SEQ ID NO 16

<211> LENGTH: 960

<212> TYPE: DNA

<213> ORGANISM: Unknown

<220> FEATURE:

<223> OTHER INFORMATION: argF nucleotide

<400> SEQUENCE: 16

```
atgacttcac aaccacaggt tcgccatttc ctggctgatg atgatctcac ccctgcagag    60
caggcagagg ttttgacct agccgcaaag ctcaaggcag cgccgttttc ggagcgcca    120
ctcgagggac caaagtccgt tgcagttctt ttgataaga cttcaactcg tactcgcttc    180
tccttcgacg cgggcatcgc tcatttggtt ggacatgcc tctctgtgga ttccggcagc    240
tcacagatgg gtaagggcga gacctgcag gacaccgag ctgtattgtc ccgctacgtg    300
gaagcaattg tgtgggcgcac ctacgcacac agcaatttcc acgccatggc ggagacgtcc    360
actgtgccac tggtgaactc cttgtccgat gatctgcacc catgccagat tctggctgat    420
ctgcagacca tcgtggaaaa cctcagccct gaagaaggcc cagcaggcct taagggttaag    480
aaggctgtgt acctgggcga tggcgacaac aacatggcca actcctacat gattggcttt    540
gccaccgcgg gcattggatat ctccatcacc gctcctgaag gggtccagcc tcgtgcggaa    600
ttcgtggagc gcgcggaaaa gcgtggccag gaaaccggcg cgaaggttgt tgtcaccgac    660
agcctcgacg aggttgccgg cgccgatgtt gtcaccaccg atacctgggt atccatgggt    720
atggaaaacg acggcatcga tcgcaccaca ctttcgttc cctaccaggt caacgatgag    780
gtcatggcga aagctaacga cggcgccatc ttctcgact gccttcctgc ctaccgcggc    840
aaagaagtgg cagcctccgt gattgatgga ccagcgcca aagttttcga tgaagcagaa    900
aaccgcctcc acgctcagaa agcactgctg gtgtggctgc tggccacca gccgaggtaa    960
```

<210> SEQ ID NO 17

<211> LENGTH: 393

<212> TYPE: PRT

<213> ORGANISM: Unknown

<220> FEATURE:

<223> OTHER INFORMATION: carA amino acid

<400> SEQUENCE: 17

Met Ser Lys Asp Thr Thr Thr Tyr Gln Gly Val Thr Glu Ile Gly Ser
1 5 10 15

Val Pro Ala Tyr Leu Val Leu Ala Asp Gly Arg Thr Phe Thr Gly Phe
20 25 30

Gly Phe Gly Ala Ile Gly Thr Thr Leu Gly Glu Ala Val Phe Thr Thr
35 40 45

Ala Met Thr Gly Tyr Gln Glu Thr Met Thr Asp Pro Ser Tyr His Arg
50 55 60

Gln Ile Val Val Ala Thr Ala Pro Gln Ile Gly Asn Thr Gly Trp Asn
65 70 75 80

Asp Glu Asp Asn Glu Ser Arg Asp Gly Lys Ile Trp Val Ala Gly Leu
85 90 95

-continued

Val Ile Arg Asp Leu Ala Ala Arg Val Ser Asn Trp Arg Ala Thr Thr
 100 105 110
 Ser Leu Gln Gln Glu Met Ala Asp Gln Gly Ile Val Gly Ile Gly Gly
 115 120 125
 Ile Asp Thr Arg Ala Leu Val Arg His Leu Arg Asn Glu Gly Ser Ile
 130 135 140
 Ala Ala Gly Ile Phe Ser Gly Ala Asp Ala Gln Arg Pro Val Glu Glu
 145 150 155 160
 Leu Val Glu Ile Val Lys Asn Gln Pro Ala Met Thr Gly Ala Asn Leu
 165 170 175
 Ser Val Glu Val Ser Ala Asp Glu Thr Tyr Val Ile Glu Ala Glu Gly
 180 185 190
 Glu Glu Arg His Thr Val Val Ala Tyr Asp Leu Gly Ile Lys Gln Asn
 195 200 205
 Thr Pro Arg Arg Phe Ser Ala Arg Gly Val Arg Thr Val Ile Val Pro
 210 215 220
 Ala Glu Thr Pro Leu Glu Asp Ile Lys Gln Tyr Asn Pro Ser Gly Val
 225 230 235 240
 Phe Ile Ser Asn Gly Pro Gly Asp Pro Ala Ala Ala Asp Val Met Val
 245 250 255
 Asp Ile Val Arg Glu Val Leu Glu Ala Asp Ile Pro Phe Phe Gly Ile
 260 265 270
 Cys Phe Gly Asn Gln Ile Leu Gly Arg Ala Phe Gly Met Glu Thr Tyr
 275 280 285
 Lys Leu Lys Phe Gly His Arg Gly Ile Asn Val Pro Val Lys Asn His
 290 295 300
 Ile Thr Gly Lys Ile Asp Ile Thr Ala Gln Asn His Gly Phe Ala Leu
 305 310 315 320
 Lys Gly Glu Ala Gly Gln Glu Phe Glu Thr Asp Phe Gly Thr Ala Ile
 325 330 335
 Val Thr His Thr Cys Leu Asn Asp Gly Val Val Glu Gly Val Ala Leu
 340 345 350
 Lys Ser Gly Arg Ala Tyr Ser Val Gln Tyr His Pro Glu Ala Ala Ala
 355 360 365
 Gly Pro Asn Asp Ala Ser Pro Leu Phe Asp Gln Phe Val Glu Leu Met
 370 375 380
 Asp Ala Asp Ala Gln Lys Lys Gly Ala
 385 390

<210> SEQ ID NO 18

<211> LENGTH: 1182

<212> TYPE: DNA

<213> ORGANISM: Unknown

<220> FEATURE:

<223> OTHER INFORMATION: carA nucleotide

<400> SEQUENCE: 18

```

gtgagtaaag acaccaccac ctaccaggga gtcaccgaga tcggatccgt tccggcatac    60
ctggttcttg cagacggacg taccttcacc ggatttgget ttggagctat cggcaccacc    120
cttggtgagg cagtgttcac caccgcatg accggttacc aagaaaccat gaccgatcct    180
tcctatcacc gccagattgt tgtggctacc gcaccacaga tcggtaacac cggctggaac    240

```

-continued

gatgaggaca acgagtcccg cgacggcaag atttgggttg caggccttgt tatccgcgac	300
ctcgcagcac gtgtgtccaa ctggcgcgcc accacctcct tgcagcagga aatggcagac	360
caaggcatcg tcggcatcgg cggaatcgac acccgcgcac tggttcgcca cctgcgcaac	420
gaaggttcca tcgcagcggg catcttctcc ggcgctgacg cacagcgccc agtgaagaa	480
ctcgtagaga tcgtcaagaa tcagccagca atgaccggcg caaacctctc cgttgaggtc	540
tctgctgatg aaacctacgt catcgaagct gagggcgaag agcgccacac cgtcgtggcc	600
tacgacctgg gcattaagca aaacaccca cgtcgtttct ctgcacggcg tggtcgcacc	660
gtcctcgtgc ctgctgaaac ccatttgag gacatcaagc agtacaaccc atcaggcggtg	720
tttatctcca atggccctgg cgaccctgca gcagcagacg tcatggttga tatcgtccgc	780
gaagttctgg aagccgacat tccattcttt ggcactctgt tcggcaacca gatcctcggc	840
cgcgcattcg gcattggagac ctacaagctg aagttcggcc accgcggcat caacgttcca	900
gtgaagaacc acatcacggg caagatcgac atcacgccc agaaccacgg cttcgcactc	960
aaggggtgaag caggccagga attcgagaca gatttcggca ctgcgattgt caccacaccc	1020
tgccctaacg acggcgctgt tgaaggtgtt gcgctgaagt ccggacgcgc atactccgtt	1080
cagtaccacc cagaggcgcg tgccggccca aatgatgcaa gccccctgtt tgaccagttt	1140
gttgagctga tggatgcaga cgctcagaag aaaggcgcat aa	1182

<210> SEQ ID NO 19

<211> LENGTH: 1113

<212> TYPE: PRT

<213> ORGANISM: Unknown

<220> FEATURE:

<223> OTHER INFORMATION: carB amino acid

<400> SEQUENCE: 19

Met	Pro	Lys	Arg	Ser	Asp	Ile	Asn	His	Val	Leu	Val	Ile	Gly	Ser	Gly
1				5					10					15	

Pro	Ile	Val	Ile	Gly	Gln	Ala	Cys	Glu	Phe	Asp	Tyr	Ser	Gly	Thr	Gln
		20						25					30		

Ala	Cys	Arg	Val	Leu	Lys	Glu	Glu	Gly	Leu	Arg	Val	Thr	Leu	Ile	Asn
		35				40						45			

Ser	Asn	Pro	Ala	Thr	Ile	Met	Thr	Asp	Pro	Glu	Met	Ala	Asp	His	Thr
	50					55				60					

Tyr	Val	Glu	Pro	Ile	Glu	Pro	Glu	Tyr	Ile	Asp	Lys	Ile	Phe	Ala	Lys
65				70					75					80	

Glu	Ile	Glu	Gln	Gly	His	Pro	Ile	Asp	Ala	Val	Leu	Ala	Thr	Leu	Gly
			85					90						95	

Gly	Gln	Thr	Ala	Leu	Asn	Ala	Ala	Ile	Gln	Leu	Asp	Arg	Leu	Gly	Ile
			100					105					110		

Leu	Glu	Lys	Tyr	Gly	Val	Glu	Leu	Ile	Gly	Ala	Asp	Ile	Asp	Ala	Ile
		115				120					125				

Glu	Arg	Gly	Glu	Asp	Arg	Gln	Lys	Phe	Lys	Asp	Ile	Val	Thr	Thr	Ile
	130					135					140				

Gly	Gly	Glu	Ser	Ala	Arg	Ser	Arg	Val	Cys	His	Asn	Met	Asp	Glu	Val
145				150					155					160	

His	Glu	Thr	Val	Ala	Glu	Leu	Gly	Leu	Pro	Val	Val	Val	Arg	Pro	Ser
			165					170						175	

Phe	Thr	Met	Gly	Gly	Leu	Gly	Ser	Gly	Leu	Ala	Tyr	Asn	Thr	Glu	Asp
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

	180							185					190				
Leu	Glu	Arg	Ile	Ala	Gly	Gly	Gly	Leu	Ala	Ala	Ser	Pro	Glu	Ala	Asn		
		195					200					205					
Val	Leu	Ile	Glu	Glu	Ser	Ile	Leu	Gly	Trp	Lys	Glu	Phe	Glu	Leu	Glu		
	210					215					220						
Leu	Met	Arg	Asp	Thr	Ala	Asp	Asn	Val	Val	Val	Ile	Cys	Ser	Ile	Glu		
225					230					235					240		
Asn	Val	Asp	Ala	Leu	Gly	Val	His	Thr	Gly	Asp	Ser	Val	Thr	Val	Ala		
				245					250					255			
Pro	Ala	Leu	Thr	Leu	Thr	Asp	Arg	Glu	Phe	Gln	Lys	Met	Arg	Asp	Gln		
		260						265					270				
Gly	Ile	Ala	Ile	Ile	Arg	Glu	Val	Gly	Val	Asp	Thr	Gly	Gly	Cys	Asn		
	275						280					285					
Ile	Gln	Phe	Ala	Ile	Asn	Pro	Val	Asp	Gly	Arg	Ile	Ile	Thr	Ile	Glu		
	290					295					300						
Met	Asn	Pro	Arg	Val	Ser	Arg	Ser	Ser	Ala	Leu	Ala	Ser	Lys	Ala	Thr		
305					310					315					320		
Gly	Phe	Pro	Ile	Ala	Lys	Met	Ala	Ala	Lys	Leu	Ala	Ile	Gly	Tyr	Thr		
				325					330					335			
Leu	Asp	Glu	Ile	Thr	Asn	Asp	Ile	Thr	Gly	Glu	Thr	Pro	Ala	Ala	Phe		
		340						345					350				
Glu	Pro	Thr	Ile	Asp	Tyr	Val	Val	Val	Lys	Ala	Pro	Arg	Phe	Ala	Phe		
		355				360						365					
Glu	Lys	Phe	Val	Gly	Ala	Asp	Asp	Thr	Leu	Thr	Thr	Thr	Met	Lys	Ser		
	370					375						380					
Val	Gly	Glu	Val	Met	Ser	Leu	Gly	Arg	Asn	Tyr	Ile	Ala	Ala	Leu	Asn		
385					390				395						400		
Lys	Ala	Leu	Arg	Ser	Leu	Glu	Thr	Lys	Gln	Gln	Gly	Phe	Trp	Thr	Lys		
				405					410					415			
Pro	Asp	Glu	Phe	Phe	Ala	Gly	Glu	Arg	Ala	Thr	Asp	Lys	Ala	Ala	Val		
				420				425					430				
Leu	Glu	Asp	Leu	Lys	Arg	Pro	Thr	Glu	Gly	Arg	Leu	Tyr	Asp	Val	Glu		
		435					440					445					
Leu	Ala	Met	Arg	Leu	Gly	Ala	Ser	Val	Glu	Glu	Leu	Tyr	Glu	Ala	Ser		
		450				455					460						
Ser	Ile	Asp	Pro	Trp	Phe	Leu	Ala	Glu	Leu	Glu	Ala	Leu	Val	Gln	Phe		
465					470					475					480		
Arg	Gln	Lys	Leu	Val	Asp	Ala	Pro	Phe	Leu	Asn	Glu	Asp	Leu	Leu	Arg		
				485					490					495			
Glu	Ala	Lys	Phe	Met	Gly	Leu	Ser	Asp	Leu	Gln	Ile	Ala	Ala	Leu	Arg		
				500				505					510				
Pro	Glu	Phe	Ala	Gly	Glu	Asp	Gly	Val	Arg	Thr	Leu	Arg	Leu	Ser	Leu		
		515					520					525					
Gly	Ile	Arg	Pro	Val	Phe	Lys	Thr	Val	Asp	Thr	Cys	Ala	Ala	Glu	Phe		
	530					535					540						
Glu	Ala	Lys	Thr	Pro	Tyr	His	Tyr	Ser	Ala	Tyr	Glu	Leu	Asp	Pro	Ala		
545					550					555					560		
Ala	Glu	Ser	Glu	Val	Ala	Pro	Gln	Thr	Glu	Arg	Glu	Lys	Val	Leu	Ile		
				565					570					575			
Leu	Gly	Ser	Gly	Pro	Asn	Arg	Ile	Gly	Gln	Gly	Ile	Glu	Phe	Asp	Tyr		
			580					585					590				

-continued

Ser	Cys	Val	His	Ala	Ala	Leu	Glu	Leu	Ser	Arg	Val	Gly	Tyr	Glu	Thr	595	600	605	
Val	Met	Val	Asn	Cys	Asn	Pro	Glu	Thr	Val	Ser	Thr	Asp	Tyr	Asp	Thr	610	615	620	
Ala	Asp	Arg	Leu	Tyr	Phe	Glu	Pro	Leu	Thr	Phe	Glu	Asp	Val	Met	Glu	625	630	635	640
Val	Tyr	His	Ala	Glu	Ala	Gln	Ser	Gly	Thr	Val	Ala	Gly	Val	Ile	Val	645	650	655	
Gln	Leu	Gly	Gly	Gln	Thr	Pro	Leu	Gly	Leu	Ala	Asp	Arg	Leu	Lys	Lys	660	665	670	
Ala	Gly	Val	Pro	Val	Ile	Gly	Thr	Ser	Pro	Glu	Ala	Ile	Asp	Met	Ala	675	680	685	
Glu	Asp	Arg	Gly	Glu	Phe	Gly	Ala	Leu	Leu	Asn	Arg	Glu	Gln	Leu	Pro	690	695	700	
Ala	Pro	Ala	Phe	Gly	Thr	Ala	Thr	Ser	Phe	Glu	Glu	Ala	Arg	Thr	Val	705	710	715	720
Ala	Asp	Glu	Ile	Ser	Tyr	Pro	Val	Leu	Val	Arg	Pro	Ser	Tyr	Val	Leu	725	730	735	
Gly	Gly	Arg	Gly	Met	Glu	Ile	Val	Tyr	Asp	Glu	Ala	Ser	Leu	Glu	Asp	740	745	750	
Tyr	Ile	Asn	Arg	Ala	Thr	Glu	Leu	Ser	Ser	Asp	His	Pro	Val	Leu	Val	755	760	765	
Asp	Arg	Phe	Leu	Asp	Asn	Ala	Ile	Glu	Ile	Asp	Val	Asp	Ala	Leu	Cys	770	775	780	
Asp	Gly	Asp	Glu	Val	Tyr	Leu	Ala	Gly	Val	Met	Glu	His	Ile	Glu	Glu	785	790	795	800
Ala	Gly	Ile	His	Ser	Gly	Asp	Ser	Ala	Cys	Ala	Leu	Pro	Pro	Met	Thr	805	810	815	
Leu	Gly	Ala	Gln	Asp	Ile	Glu	Lys	Val	Arg	Glu	Ala	Thr	Lys	Lys	Leu	820	825	830	
Ala	Leu	Gly	Ile	Gly	Val	Gln	Gly	Leu	Met	Asn	Val	Gln	Tyr	Ala	Leu	835	840	845	
Lys	Asp	Asp	Ile	Leu	Tyr	Val	Ile	Glu	Ala	Asn	Pro	Arg	Ala	Ser	Arg	850	855	860	
Thr	Val	Pro	Phe	Val	Ser	Lys	Ala	Thr	Gly	Val	Asn	Leu	Ala	Lys	Ala	865	870	875	880
Ala	Ser	Arg	Ile	Ala	Val	Gly	Ala	Thr	Ile	Lys	Asp	Leu	Gln	Asp	Glu	885	890	895	
Gly	Met	Ile	Pro	Thr	Glu	Tyr	Asp	Gly	Gly	Ser	Leu	Pro	Leu	Asp	Ala	900	905	910	
Pro	Ile	Ala	Val	Lys	Glu	Ala	Val	Leu	Pro	Phe	Asn	Arg	Phe	Arg	Arg	915	920	925	
Pro	Asp	Gly	Lys	Thr	Leu	Asp	Thr	Leu	Leu	Ser	Pro	Glu	Met	Lys	Ser	930	935	940	
Thr	Gly	Glu	Val	Met	Gly	Leu	Ala	Asn	Asn	Phe	Gly	Ala	Ala	Tyr	Ala	945	950	955	960
Lys	Ala	Glu	Ala	Gly	Ala	Phe	Gly	Ala	Leu	Pro	Thr	Glu	Gly	Thr	Val	965	970	975	
Phe	Val	Thr	Val	Ala	Asn	Arg	Asp	Lys	Arg	Thr	Leu	Ile	Leu	Pro	Ile	980	985	990	

-continued

Gln	Arg	Leu	Ala	Ser	Met	Gly	Tyr	Lys	Ile	Leu	Ala	Thr	Glu	Gly	Thr
		995					1000					1005			
Ala	Gly	Met	Leu	Arg	Arg	Asn	Gly	Ile	Asp	Cys	Glu	Val	Val	Leu	
	1010					1015					1020				
Lys	Ala	Ser	Asp	Ile	Arg	Glu	Gly	Val	Glu	Gly	Lys	Ser	Ile	Val	
	1025					1030					1035				
Asp	Arg	Ile	Arg	Glu	Gly	Glu	Val	Asp	Leu	Ile	Leu	Asn	Thr	Pro	
	1040					1045					1050				
Ala	Gly	Ser	Ala	Gly	Ala	Arg	His	Asp	Gly	Tyr	Asp	Ile	Arg	Ala	
	1055					1060					1065				
Ala	Ala	Val	Thr	Val	Gly	Val	Pro	Leu	Ile	Thr	Thr	Val	Gln	Gly	
	1070					1075					1080				
Val	Thr	Ala	Ala	Val	Gln	Gly	Ile	Glu	Ala	Leu	Arg	Glu	Gly	Val	
	1085					1090					1095				
Val	Ser	Val	Arg	Ala	Leu	Gln	Glu	Leu	Asp	His	Ala	Val	Lys	Ala	
	1100					1105					1110				

<210> SEQ ID NO 20

<211> LENGTH: 3342

<212> TYPE: DNA

<213> ORGANISM: Unknown

<220> FEATURE:

<223> OTHER INFORMATION: carB nucleotide

<400> SEQUENCE: 20

atgccaaagc gttcagatat taaccacgtc ctcgtcatcg gttccggccc catcgtcatt	60
ggccaggcat gtgaattcga ctactccggc acccaggctt gccgcgtgct gaaggaagag	120
ggactgcgcg tcaccctcat caactccaac ccagcaacga tcatgaccga cccagaaatg	180
gctgaccaca cctacgtgga gccaatcgag ccggaatata tcgacaagat tttcgctaag	240
gagatcgagc agggccaccc aatcgacgcc gtcctggcaa cccttggtgg ccagactgca	300
cttaacgcag ctatccagct ggatcgccctc ggcatcctgg aaaagtacgg cgttgaactc	360
atcgggtgcag acatcgatgc cattgagcgc ggcgaagatc gccagaagtt caaggatatt	420
gtcaccacca tcggtggcga atccgcgcgt tcccgcgtct gccacaacat ggacgaagtc	480
catgagactg tcgcagaact tggccttcca gtagtcgtgc gtccatcctt cactatgggt	540
ggcctggggt ccggtcttgc atacaacacc gaagacctg agcgcacgc aggtggcgga	600
cttgctgcat ctctgaagc aaacgtcttg atcgaagaat ccacctctgg ttggaaggaa	660
ttcgagctcg agtcacgag cgataccgca gacaacgttg tggttatctg ctccattgaa	720
aacgtcgacg cactgggcgt gcacaccggc gactctgtca ccgtggcacc tgcctgacc	780
ctgactgacc gtgaattcca gaagatgcgc gatcagggtg tcgccatcat ccgcgaggtc	840
ggcgtggaca ccggtggatg taacatccag ttcgctatca acccagttga tggccgcatc	900
atcaccattg agatgaaccc acgtgtgtct cgttcctccg cgctggcatc caaggcaacg	960
ggcttcccaa ttgccaagat ggtgccaag ctggctatcg gatacacctt ggatgagatc	1020
accaacgaca tcaactgtga aacccagct gcgtttgagc ccaccatcga ctacgtcgtg	1080
gtcaaggccc cacgctttgc tttcgagaag tttgtcggcg ctgatgacac tttgaccacc	1140
accatgaagt ccgtcggtga ggatcatgtc ctgggccgta actacattgc agcactgaac	1200
aaggcactgc gttccctgga aaccaagcag cagggtttct ggaccaagcc tgatgagttc	1260

-continued

ttcgcagggg	agcgcgctac	cgataaggca	gctgtttctg	aagatctcaa	gcgccaacc	1320
gaaggccgcc	tctacgacgt	tgagctggca	atgcgccttg	gcgcaagcgt	ggaagaactc	1380
tacgaagcat	cttctattga	tccttggttc	ctcgcgcgagc	ttgaagctct	cgtgcagttc	1440
cgcacagaagc	tcgttgacgc	accattcctc	aacgaagatc	tcctgcgcga	agcaaagttc	1500
atgggtctgt	cgcacctgca	gatcgcagcc	cttcgcccag	agttcgctgg	cgaagacggc	1560
gtacgcacct	tgcgtctgtc	cctaggcctc	cgcgcagtat	tcaagactgt	ggatacctgt	1620
gcagcagagt	ttgaagctaa	gactccgtac	cactactccg	catacgagct	ggatccagca	1680
gctgagtctg	aggctgcacc	acagactgag	cgtgaaaagg	tcctgatctt	gggtccgggt	1740
ccaaaccgca	tcggccaggg	catcgagttc	gactattcct	gtgttcacgc	agctcttgag	1800
ctctcccgcg	tcggctacga	aactgtcatg	gtcaactgca	accagagac	cgtgtccacc	1860
gactacgaca	cgcgtgaccg	cctgtacttc	gagccactga	ccttcgaaga	cgtcatggag	1920
gtctaccacg	ctgaggcgca	gtccggcacc	gtcgcaggtg	ttatcgccca	gcttggtggc	1980
cagactcctc	tgggcttggc	agatcgtttg	aagaaggctg	gcgtccctgt	cattggtacc	2040
tccccagagg	caatcgacat	ggctgaggac	cgtggcagag	tcggtgcact	gctgaaccgc	2100
gagcagcttc	ctgtctcagc	attcggcacc	gcaacctctt	tcgaagaggc	tcgcacagta	2160
gccgatgaga	tcagctaccc	agtgtctggt	cgccttctct	acgtcttggg	tggccgtggc	2220
atggagattg	tctacgatga	ggcttccctc	gaggattaca	tcaaccgcgc	aactgagttg	2280
tcttctgacc	accagtgct	ggttgaccgc	ttcctggaca	acgtattga	gatcgacgtc	2340
gacgcactgt	gcgacggcga	cgaagtctac	ctggcgggcg	tcattggaaca	catcgaggaa	2400
gccggcatto	actcgggtga	ctcgcgatgt	gcacttcctc	caatgaactt	gggcgcacag	2460
gacatcgaga	aggctccgca	agcaaccaag	aagctggctc	tgggcacatcg	cgtacagggc	2520
ctgatgaacg	tccagtacgc	actcaaggac	gacatcctct	acgtcatcga	ggcaaaccga	2580
cgtgcatccc	gcaccgtgcc	gttcgtctcc	aaggcaacgg	gcgtcaacct	ggccaaggca	2640
gcatcccgtg	tcgcagtggg	cgcaccatc	aaggatctcc	aagatgaggg	catgattcct	2700
accgagtacg	acggcggtc	cttgccactg	gacgtctcaa	tcgctgtgaa	ggaagcagtg	2760
ttgccgttca	accgcttcgc	tcgcccagat	ggaaagaccc	tggacacct	gctttcccca	2820
gagatgaagt	ccactggcga	ggctcatggc	ttggccaaca	acttcggcgc	tgcatatgca	2880
aaggctgaag	ctggcgctgt	tggtgcattg	ccaaccgaag	gcacgctctt	cgtgacgtg	2940
gctaaccgcg	acaagcgcac	cctgatectg	ccaatccagc	gcctggcgct	gatgggtac	3000
aagatcctcg	ccaccgaagg	caccgcaggc	atgctgcgcc	gcaacggcat	tgattgtgaa	3060
gttgtgctca	aggcttccga	catccgcgaa	ggtgtagagg	gcaagtccat	cgtggatcgt	3120
atccgcgaag	gcgaagtga	cctcactcct	aacaccccag	ctggttctgc	tggcgctcgc	3180
cacgatggct	acgatatccg	cgcagcagca	gtgaccgtgg	gtgttccact	gatcaccact	3240
gtccaggggtg	tcaccgcagc	tgtccagggc	attgagggcc	tgcgtgaggg	cgttgtcagc	3300
gtccgcgcgc	tgcaggaact	cgaccacgca	gtcaaggctt	aa		3342

1. A recombinant microorganism of the genus *Corynebacterium* producing L-arginine, in which the activity of a protein consisting of an amino acid sequence of SEQ ID NO: 1 is weakened.

2. The microorganism of claim 1, wherein the microorganism has increased L-arginine producing ability compared to a parent strain.

3. The microorganism of claim 1, wherein the microorganism is *Corynebacterium glutamicum*.

4. The microorganism of claim 1, wherein the activity of an arginine repressor is further weakened.

5. The microorganism of claim 1, wherein a polynucleotide encoding the protein consisting of the amino acid sequence of SEQ ID NO: 1 is deleted.

6. A method for producing L-arginine, comprising: culturing a recombinant microorganism of the genus *Corynebacterium*, in which the activity of a protein comprising an amino acid sequence of SEQ ID NO: 1 is weakened, in a medium.

7. The method of claim 6, wherein the microorganism is *Corynebacterium glutamicum*.

8. The method of claim 6, wherein the activity of an arginine repressor is further weakened in the microorganism of the genus *Corynebacterium*.

9. The method of claim 6, wherein a nucleic acid sequence encoding the protein consisting of the amino acid sequence of SEQ ID NO: 1 is deleted in the microorganism.

* * * * *