



US 20250282762A1

(19) **United States**(12) **Patent Application Publication**
BONAZZI et al.(10) **Pub. No.: US 2025/0282762 A1**(43) **Pub. Date: Sep. 11, 2025**(54) **PYRAZOLOPYRIDINE DERIVATIVES AND
USES THEREOF**(71) Applicant: **Novartis AG**, Basel (CH)(72) Inventors: **Simone BONAZZI**, Cambridge, MA
(US); **Jennifer Stroka COBB**, Stow,
MA (US); **Natalie Alysia DALES**,
Arlington, MA (US); **Matthew James
HESSE**, Pittsburgh, PA (US); **Rama
JAIN**, Danville, CA (US); **John Ryan
KERRIGAN**, Wakefield, MA (US);
Hasnain Ahmed MALIK, Boston, MA
(US); **James R. MANNING**,
Emeryville, CA (US); **Pamela YF
TING**, Somerville, MA (US)division of application No. 17/693,759, filed on Mar.
14, 2022, now Pat. No. 11,787,785.(60) Provisional application No. 63/161,139, filed on Mar.
15, 2021, provisional application No. 63/164,130,
filed on Mar. 22, 2021.**Publication Classification**(51) **Int. Cl.****C07D 403/14** (2006.01)**A61P 7/00** (2006.01)**C07D 405/14** (2006.01)**C07D 417/14** (2006.01)(52) **U.S. Cl.**CPC **C07D 403/14** (2013.01); **A61P 7/00**
(2018.01); **C07D 405/14** (2013.01); **C07D**
417/14 (2013.01)(21) Appl. No.: **18/939,314**(22) Filed: **Nov. 6, 2024****Related U.S. Application Data**(60) Continuation of application No. 18/826,887, filed on
Sep. 6, 2024, now Pat. No. 12,252,482, which is a
continuation of application No. 18/460,428, filed on
Sep. 1, 2023, now Pat. No. 12,172,984, which is a

(57)

ABSTRACT

The present disclosure relates to compounds of formula (I) and pharmaceutical compositions and their use in reducing Widely Interspaced Zinc Finger Motifs (WIZ) expression levels, or inducing fetal hemoglobin (HbF) expression, and in the treatment of inherited blood disorders (e.g., hemoglobinopathies, e.g., beta-hemoglobinopathies), such as sickle cell disease and beta-thalassemia.



Specification includes a Sequence Listing.

Figure 1A




Figure 1B





Figure 1C



PYRAZOLOPYRIDINE DERIVATIVES AND USES THEREOF

CLAIM OF PRIORITY

[0001] This application is a continuation of U.S. patent application Ser. No. 18/826,887 filed Sep. 6, 2024, which is a continuation of U.S. patent application Ser. No. 18/460,428 filed Sep. 1, 2023, which is a divisional application of U.S. patent application Ser. No. 17/693,759 filed on Mar. 14, 2022, now U.S. Pat. No. 11,787,785; which claims the benefit of U.S. Provisional Application No. 63/161,139 filed on Mar. 15, 2021 and U.S. Provisional Application No. 63/164,130 filed on Mar. 22, 2021. All of these patent applications are incorporated by reference herein in their entirety.

SEQUENCE LISTING

[0002] The instant application contains a Sequence Listing which has been submitted electronically in XML format and is hereby incorporated by reference in its entirety. Said XML copy, created on Aug. 30, 2023 is named PAT059039-US-DIV_ST26 SQL.txt and is 20,480 bytes in size.

FIELD OF THE DISCLOSURE

[0003] The present disclosure relates to pyrazolopyridine derivatives compounds and pharmaceutical compositions and their use in reducing Widely Interspaced Zinc Finger Motifs (WIZ) protein expression levels and/or inducing fetal hemoglobin (HbF) protein expression levels, and in the treatment of inherited blood disorders (hemoglobinopathies, e.g., beta-hemoglobinopathies), such as sickle cell disease and beta-thalassemia.

BACKGROUND OF THE DISCLOSURE

[0004] Sickle cell disease (SCD) is a group of severe inherited blood disorders that cause red blood cells to contort into a sickle shape. These cells can cause blockages in blood flow, leading to intense pain, organ damage and premature death. Beta thalassemias are a group of inherited blood disorders that are caused by reduced or absent synthesis of beta globin, causing anemia.


[0005] Fetal hemoglobin (HbF) induction is known to ameliorate symptoms in SCD and beta-thalassemia patients, with both genetic (single nucleotide polymorphisms in the globin control locus & BCL11A) and pharmacologic (hydroxyurea) validation in the clinic (Vinjamur, D. S., et al. (2018), *The British Journal of Haematology*, 180(5), 630-643). Hydroxyurea is the current standard of care for SCD and is thought to provide benefit via induction of HbF, but is genotoxic, causes dose-limiting neutropenia and has a response rate of less than 40%. Other mechanisms being targeted clinically and preclinically include inhibition of HDAC1/2 (Shearstone et al., 2016, *PLoS One*, 11(4), e0153767), LSD1 (Rivers et al., 2018, *Experimental Hematology*, 67, 60-64), DNMT1, PDE9a (McArthur et al., 2019, *Haematologica*. doi:10.3324/haematol.2018.213462), HRI kinase (Grevet et al., 2018, *Science*, 361(6399), 285-290) and G9a/GLP (Krivega et al., 2015, *Blood*, 126(5), 665-672; Renneville et al., 2015, *Blood*, 126(16), 1930-1939). Additionally, the immunomodulators pomalidomide and lenalidomide induce HbF ex vivo in human primary erythroid cells (Moutouh-de Parseval, L. A. et al. (2008), *The Journal of Clinical Investigation*, 118(1), 248-258) and in

vivo (Meiler, S. E. et al. (2011), *Blood*, 118(4), 1109-1112). WIZ is ubiquitously expressed and plays a role in targeting the G9a/GLP histone methyltransferases to genomic loci to regulate chromatin structure and transcription (Bian, Chen, et al. (2015), *eLife* 2015; 4:e05606).

SUMMARY OF THE DISCLOSURE

[0006] The disclosure relates to a therapeutic agent, which is effective in reducing WIZ protein expression levels and/or inducing fetal hemoglobin (HbF) expression. The disclosure further relates to pyrazolopyridine compounds, which are effective in reducing WIZ protein expression levels and/or inducing fetal hemoglobin (HbF) expression, pharmaceutically acceptable salts thereof, compositions thereof, and their use in therapies for the conditions and purposes detailed above.

[0007] The disclosure provides, in a first aspect, a compound of formula (I') or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein:



[0008] == is a single bond or a double bond;

[0009] X is selected from CH, CF, and N;

[0010] R^x is selected from hydrogen, C₁-C₆alkyl, halo (e.g., F, Cl), C₁-C₆alkoxyl, and C₃-C₈cycloalkyl;

[0011] R¹ is selected from hydrogen and C₁-C₆alkyl;

[0012] R¹ is selected from hydrogen and C₁-C₆alkyl;

[0013] each R² is independently selected from C₁-C₆alkyl, C₁-C₆haloalkyl, halo, and oxo, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{2a}; or 2 R² on non-adjacent carbon atoms together with the non-adjacent carbon atoms to which they are attached form a bridging ring;

[0014] R^{2a} is selected from C₁-C₆alkoxyl and hydroxyl;

[0015] R³ is selected from hydrogen, C₁-C₈alkyl, CN₂-C₆alkenyl, —SO₂R⁴, C₁-C₆haloalkyl, —C(=O)—O—(R⁵), —C(=O)—(R⁶), C₃-C₁₀cycloalkyl, and a 4- to 10-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, wherein the C₁-C₈alkyl and C₁-C₆haloalkyl are each independently substituted with 0-3 occurrences of R^{3a}, and wherein the C₃-C₁₀cycloalkyl and 4- to 10-membered heterocyclyl are each independently substituted with 0-3 occurrences of R^{3b};

[0016] or

[0017] R³ together with the nitrogen atom to which it is attached and R² together with the carbon atom to which it is attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N,

O and S, which 5- or 6-membered heterocyclyl is substituted with 0-2 occurrences of an oxo group;

[0018] each R^{3a} is independently selected from C_3 - C_{10} cycloalkyl, a 4- to 10-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, a 5- to 10-membered heteroaryl comprising 1-4 heteroatoms independently selected from N, O, and S, C_6 - C_{10} aryl, C_1 - C_6 alkoxyl, hydroxyl, and $-C(=O)-NR^7R^8$, wherein the C_3 - C_{10} cycloalkyl, 4- to 6-membered heterocyclyl, 5- to 10-membered heteroaryl and C_6 - C_{10} aryl are substituted with 0-4 occurrences of R^{3b} ;

[0019] each R^{3b} is independently selected from C_1 - C_6 alkoxyl, halo, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkoxyl, C_1 - C_6 alkyl, $-CN$, $-SO_2NR^7R^8$, $-SO_2R^4$, and hydroxyl;

[0020] R^4 is selected from C_3 - C_8 cycloalkyl, C_1 - C_6 alkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, C_6 - C_{10} aryl, and $-NR^{4b}R^{4c}$, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{4a} ;

[0021] R^{4a} is selected from C_3 - C_8 cycloalkyl, C_6 - C_{10} aryl, and C_1 - C_6 alkoxyl;

[0022] R^{4b} is selected from hydrogen, and C_1 - C_6 alkyl;

[0023] R^{4c} is selected from hydrogen, C_1 - C_6 alkyl, and C_3 - C_8 cycloalkyl;

[0024] R^5 is selected from C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, and C_6 - C_{10} aryl;

[0025] R^6 is selected from C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, C_6 - C_{10} aryl, a 4- to 10-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, and $-NR^{4b}R^{4c}$, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{6a} , the C_3 - C_8 cycloalkyl is substituted with 0-1 occurrence of R^{6b} , and the 4- to 10-membered heterocyclyl is substituted with 0-1 occurrence of C_1 - C_6 alkyl;

[0026] R^{6a} is selected from C_6 - C_{10} aryl and C_3 - C_8 cycloalkyl;

[0027] R^{6b} is selected from halo, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkoxyl, and C_1 - C_6 alkyl;

[0028] R^7 is selected from hydrogen and C_1 - C_6 alkyl;

[0029] R^8 is selected from hydrogen and C_1 - C_6 alkyl;

[0030] or


[0031] R^7 and R^8 together with the nitrogen atom to which they are attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O, and S;

[0032] n is 0, 1, 2, 3 or 4;

[0033] m is 0, 1 or 2; and

[0034] p is 0 or 1.

[0035] The disclosure provides, in a further aspect, a compound of formula (I') or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein:



(I')

[0036] === is a single bond or a double bond;

[0037] X is selected from CH, CF, and N;

[0038] R^1 is selected from hydrogen and C_1 - C_6 alkyl;

[0039] R^1 is selected from hydrogen and C_1 - C_6 alkyl;

[0040] each R^2 is independently selected from C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, halo, and oxo, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{2a} ; or 2 R^2 on non-adjacent carbon atoms together with the non-adjacent carbon atoms to which they are attached form a bridging ring;

[0041] R^{2a} is selected from C_1 - C_6 alkoxyl and hydroxyl;

[0042] R^3 is selected from hydrogen, C_1 - C_8 alkyl, C_2 - C_6 alkenyl, $-SO_2R^4$, C_1 - C_6 haloalkyl, $-C(=O)-O-(R^5)$ and $-C(=O)-(R^6)$, wherein the C_1 - C_8 alkyl and C_1 - C_6 haloalkyl are independently substituted with 0-3 occurrences of R^{3a} ;

[0043] or

[0044] R^3 together with the nitrogen atom to which it is attached and R^2 together with the carbon atom to which it is attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O and S;

[0045] each R^{3a} is independently selected from C_3 - C_{10} cycloalkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, a 5- to 10-membered heteroaryl comprising 1-4 heteroatoms independently selected from N, O, and S, C_6 - C_{10} aryl, C_1 - C_6 alkoxyl, hydroxyl, and $-C(=O)-NR^7R^8$, wherein the C_3 - C_{10} cycloalkyl, 4- to 6-membered heterocyclyl, 5- to 10-membered heteroaryl and C_6 - C_{10} aryl are substituted with 0-4 occurrences of R^{3b} ;

[0046] each R^{3b} is independently selected from C_1 - C_6 alkoxyl, halo, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkoxyl, C_1 - C_6 alkyl, $-CN$, $-SO_2NR^7R^8$, $-SO_2R^4$, and hydroxyl;


[0047] R^4 is selected from C_3 - C_8 cycloalkyl, C_1 - C_6 alkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, and C_6 - C_{10} aryl, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{4a} ;

[0048] R^{4a} is selected from C_3 - C_8 cycloalkyl, C_6 - C_{10} aryl, and C_1 - C_6 alkoxyl;

[0049] R^5 is selected from C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, and C_6 - C_{10} aryl;

[0050] R^6 is selected from C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, and C_6 - C_{10} aryl, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{6a} and the C_3 - C_8 cycloalkyl is substituted with 0-1 occurrence of R^{6b} ;

- [0051] R^{6a} is selected from C_6 - C_{10} aryl and C_3 - C_8 cycloalkyl;
 [0052] R^{6b} is selected from halo, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkoxyl, and C_1 - C_6 alkyl;
 [0053] R^7 is selected from hydrogen and C_1 - C_6 alkyl;
 [0054] R^a is selected from hydrogen and C_1 - C_6 alkyl;
 [0055] or
 [0056] R^7 and R^8 together with the nitrogen atom to which they are attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O, and S;
 [0057] n is 0, 1, 2, 3 or 4;
 [0058] m is 0, 1 or 2; and
 [0059] p is 0 or 1.
 [0060] The disclosure provides, in a further aspect, a compound of formula (I) or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein:



- [0061] X is selected from CH, CF, and N;
 [0062] R' is selected from hydrogen and C_1 - C_6 alkyl;
 [0063] R^1 is selected from hydrogen and C_1 - C_6 alkyl;
 [0064] each R^2 is independently selected from C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, halo, and oxo, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{2a} ; or 2 R^2 on non-adjacent carbon atoms together with the non-adjacent carbon atoms to which they are attached form a bridging ring;
 [0065] R^{2a} is selected from C_1 - C_6 alkoxyl and hydroxyl;
 [0066] R^3 is selected from hydrogen, C_1 - C_8 alkyl, C_2 - C_6 alkenyl, $-\text{SO}_2\text{R}^4$, C_1 - C_6 haloalkyl, $-\text{C}(=\text{O})-\text{O}-(\text{R}^5)$ and $-\text{C}(=\text{O})-(\text{R}^6)$, wherein the C_1 - C_8 alkyl and C_1 - C_6 haloalkyl are independently substituted with 0-3 occurrences of R^{3a} ;
 [0067] or
 [0068] R^3 together with the nitrogen atom to which it is attached and R^2 together with the carbon atom to which it is attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O and S;
 [0069] each R^{3a} is independently selected from C_3 - C_{10} cycloalkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, a 5- to 10-membered heteroaryl comprising 1-4 heteroatoms independently selected from N, O, and S, C_6 - C_{10} aryl, C_1 - C_6 alkoxyl, hydroxyl, and $-\text{C}(=\text{O})-\text{NR}^7\text{R}^8$, wherein the C_3 - C_{10} cycloalkyl, 4- to 6-membered heterocyclyl, 5- to 10-membered heteroaryl and C_6 - C_{10} aryl are substituted with 0-4 occurrences of R^{3b} ;

- [0070] each R^{3b} is independently selected from C_1 - C_6 alkoxyl, halo, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkoxyl, C_1 - C_6 alkyl, $-\text{CN}$, $-\text{SO}_2\text{NR}^7\text{R}^8$, $-\text{SO}_2\text{R}^4$, and hydroxyl;
 [0071] R^4 is selected from C_3 - C_8 cycloalkyl, C_1 - C_6 alkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, and C_6 - C_{10} aryl, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{4a} ;
 [0072] R^{4a} is selected from C_3 - C_8 cycloalkyl, C_6 - C_{10} aryl, and C_1 - C_6 alkoxyl;
 [0073] R^5 is selected from C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, and C_6 - C_{10} aryl;
 [0074] R^6 is selected from C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, and C_6 - C_{10} aryl, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{6a} and the C_3 - C_8 cycloalkyl is substituted with 0-1 occurrence of R^{6b} ;
 [0075] R^{6a} is selected from C_6 - C_{10} aryl and C_3 - C_8 cycloalkyl;
 [0076] R^{6b} is selected from halo, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkoxyl, and C_1 - C_6 alkyl;
 [0077] R^7 is selected from hydrogen and C_1 - C_6 alkyl;
 [0078] R^8 is selected from hydrogen and C_1 - C_6 alkyl;
 [0079] or
 [0080] R^7 and R^8 together with the nitrogen atom to which they are attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O, and S;
 [0081] n is 0, 1, 2, 3 or 4;
 [0082] m is 0, 1 or 2; and
 [0083] p is 0 or 1.

[0084] In a further aspect, the disclosure provides a pharmaceutical composition comprising a therapeutically effective amount of a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, and a pharmaceutically acceptable carrier or excipient.

[0085] In a further aspect, the disclosure provides a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use as a medicament.

[0086] In a further aspect, the disclosure provides a method of treating or preventing a disease or disorder in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.

[0087] In a further aspect, the disclosure provides a method of treating or preventing a disorder that is affected by the reduction or modulation of WIZ protein levels, in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2),

[0089] In a further aspect, the disclosure provides a method of degrading WIZ protein in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of formula (I''), (I'), (I), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, pro-drug, stereoisomer, or tautomer thereof.

[0090] In an further aspect, the disclosure provides a method of inhibiting, reducing, or eliminating the activity of WIZ protein or WIZ protein expression, the method comprising administering to the subject a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, pro-drug, stereoisomer, or tautomer thereof.

[0091] In a further aspect, the disclosure provides a method of inducing or promoting fetal hemoglobin in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.

[0092] In a further aspect, the disclosure provides a method of reactivating fetal hemoglobin production or expression in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.

[0093] In an further aspect, the disclosure provides a method of increasing fetal hemoglobin expression in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.

[0094] In a further aspect, the disclosure provides a method of treating a hemoglobinopathy, e.g., a beta-hemoglobinopathy, in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.

[0095] In a further aspect, the disclosure provides a method of treating a sickle cell disease in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, pro-drug, stereoisomer, or tautomer thereof.

[0096] In a further aspect, the disclosure provides a method of treating beta-thalassemia in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, pro-drug, stereoisomer, or tautomer thereof.

[0097] In a further aspect, the disclosure provides a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in the treatment of a disease or disorder.

[0098] In a further aspect, the disclosure provides a compound of (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in the treatment of a disease or disorder selected from sickle cell disease and beta-thalassemia.

[0099] In a further aspect, the disclosure provides a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in the treatment or prevention of a disease or disorder that is affected by the reduction of WIZ protein levels.

[10100] In a further aspect, the disclosure provides a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in the treatment or prevention of a disease or disorder that is affected by the inhibition or reduction of WIZ protein expression.

[0101] In a further aspect, the disclosure provides a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in the treatment or prevention of a disease or disorder that is affected by the degradation of WIZ protein.

[10102] In a further aspect, the disclosure provides a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in inhibiting, reducing, or eliminating the activity of WIZ protein or WIZ protein expression.

[10103] In a further aspect, the disclosure provides a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in inducing or promoting fetal hemoglobin.

[0104] In a further aspect, the disclosure provides a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in reactivating fetal hemoglobin production or expression.

[0105] In a further aspect, the disclosure provides a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in increasing fetal hemoglobin expression.

[0106] In a further aspect, the disclosure provides a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in the treatment of a hemoglobinopathy.

[0107] In a further aspect, the disclosure provides a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in the treatment of a sickle cell disease.

[0108] In a further aspect, the disclosure provides a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in the treatment of beta-thalassemia.

[0109] In a further aspect, the disclosure provides a compound of (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in the treatment of a disease or disorder affected by an increase in fetal hemoglobin expression.

[0110] In a further aspect, the disclosure provides a compound of (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in the treatment of a disease or disorder affected by the inhibition, reduction, or elimination of the activity of WIZ protein or WIZ protein expression.

[0111] In a further aspect, the disclosure provides a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in the treatment of a disease or disorder affected by the induction or promotion of fetal hemoglobin.

[0112] In a further aspect, the disclosure provides a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in the treatment of a disease or disorder affected by the reactivation of fetal hemoglobin production or expression.

[0113] Various other aspects of the disclosure are described herein and in the claims.

[0114] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. In the specification and claims, the singular forms also include the plural unless the context clearly dictates otherwise. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the disclosure, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entireties for all purposes. The references cited herein are not admitted to be prior art to the claimed disclosure. In the case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and are not intended to be limiting.

[0115] Other features and advantages of compounds, compositions, and methods disclosed herein will be apparent from the following detailed description and claims.

BRIEF DESCRIPTION OF THE DRAWINGS

[0116] FIG. 1A depicts a volcano plot of differentially expressed genes from WIZ KO cells as compared to a scrambled gRNA control. Each dot represents a gene. HBG1/2 genes are differentially upregulated with WIZ_6 and WIZ_18 gRNA targeting WIZ KO.

[0117] FIG. 1B depicts a bar graph showing the frequency of HbF+ cells due to shRNA-mediated loss of WIZ in human mobilized peripheral blood CD34+ derived erythroid cells.

[0118] FIG. 1C depicts a bar graph showing the frequency of HbF+ cells due to CRISPR/Cas9-mediated loss of WIZ in human mobilized peripheral blood CD34+ derived erythroid cells.

DETAILED DESCRIPTION OF THE DISCLOSURE

[0119] The compounds disclosed herein are effective in reducing WIZ protein expression levels, or inducing fetal hemoglobin (HbF) expression. Without wishing to be bound by any theory, it is believed that the disclosed compounds may treat blood disorders, such as inherited blood disorders, e.g., sickle cell disease, and beta-thalassemia by inducing fetal hemoglobin HbF expression.

Definitions

[0120] Unless specified otherwise, the terms “compounds of the present disclosure,” “compounds of the disclosure,” or “compound of the disclosure” refer to compounds of formulae (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), and (Ie), exemplified compounds, salts thereof, particularly pharmaceutically acceptable salts thereof, hydrates, solvates, prodrugs, as well as all stereoisomers (including diastereoisomers and enantiomers), rotamers, tautomers, and isotopically labeled compounds (including deuterium substitutions), as well as inherently formed moieties.

[0121] In the groups, radicals, or moieties defined below, the number of carbon atoms is often specified preceding the group, for example, C₁-C₈alkyl means an alkyl group or radical having 1 to 8 carbon atoms. In general, for groups comprising two or more subgroups, the last named group is the radical attachment point, for example, “alkylaryl” means

a monovalent radical of the formula alkyl-aryl-, while “aryl-alkyl” means a monovalent radical of the formula aryl-alkyl-.

[0122] Furthermore, the use of a term designating a monovalent radical where a divalent radical is appropriate shall be construed to designate the respective divalent radical and vice versa. Unless otherwise specified, conventional definitions of terms control and conventional stable atom valences are presumed and achieved in all formulas and groups. The articles “a” and “an” refer to one or more than one (e.g., but at least one) of the grammatical object of the article. By way of example, “an element” means one element or more than one element.

[0123] The term “and/or” means either “and” or “or” unless indicated otherwise.

[0124] The term “substituted” means that the specified group or moiety bears one or more suitable substituents wherein the substituents may connect to the specified group or moiety at one or more positions. For example, an aryl substituted with a cycloalkyl may indicate that the cycloalkyl connects to one atom of the aryl with a bond or by fusing with the aryl and sharing two or more common atoms.

[0125] As used herein the term “C₁-C₈alkyl” refers to a straight or branched hydrocarbon chain radical consisting solely of carbon and hydrogen atoms, containing no unsaturation, having from one to eight carbon atoms, and which is attached to the rest of the molecule by a single bond. The terms “C₁-C₃alkyl”, “C₁-C₄alkyl”, “C₁-C₆alkyl”, are to be construed accordingly. Examples of C₁-C₈alkyl include, but are not limited to, methyl, ethyl, n-propyl, 1-methylethyl (iso-propyl), n-butyl, 1-methylpropyl (sec-butyl), 2-methylpropyl (iso-butyl or i-butyl), 1,1-dimethylethyl (t-butyl), n-pentyl, 3-pentyl, n-hexyl, n-heptyl, 4-heptyl, n-octyl, 2-isopropyl-3-methylbutyl.

[0126] As used herein, the term “C₁-C₆alkoxyl” refers to a radical of the formula —OR, where R_a is a C₁-C₆alkyl radical as generally defined above. Examples of C₁-C₆alkoxyl include, but are not limited to, methoxy, ethoxy, propoxy, iso-propoxy, butoxy, iso-butoxy, tert-butoxy, sec-butoxy, pentoxy, and hexoxy.

[0127] As used herein, the term “C₁-C₆haloalkyl” refers to C₁-C₆alkyl radical, as defined above, substituted by one or more halo radicals, as defined herein. Examples of C₁-C₆haloalkyl include, but are not limited to, trifluoromethyl, difluoromethyl, fluoromethyl, trichloromethyl, 1,1-difluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 2-fluoropropyl, 1,1,1-trifluoropropyl, 2,2-difluoropropyl, 3,3-difluoropropyl and 1-fluoromethyl-2-fluoroethyl, 1,3-dibromopropan-2-yl, 3-bromo-2-fluoropropyl, 1,1,2,2-tetrafluoropropyl, and 1,4,4-trifluorobutan-2-yl.

[0128] As used herein, the term “C₁-C₆haloalkoxyl” means a C₁-C₆alkoxyl group as defined herein substituted with one or more halo radicals. Examples of C₁-C₆haloalkoxyl groups include, but are not limited to, trifluoromethoxy, difluoromethoxy, fluoromethoxy, trichloromethoxy, 1,1-difluoroethoxy, 2,2-difluoroethoxy, 2,2,2-trifluoroethoxy, 1-fluoromethyl-2-fluoroethoxy, pentafluoroethoxy, 2-fluoropropoxy, 3,3-difluoropropoxy and 3-dibromopropoxy. Preferably, the one or more halo radicals of C₁-C₆haloalkoxyl is fluoro. Preferably, C₁-C₆haloalkoxyl is selected from trifluoromethoxy, difluoromethoxy, fluoromethoxy, 1,1-difluoroethoxy, 2,2-difluoroethoxy, 2,2,2-trifluoroethoxy, 1-fluoromethyl-2-fluoroethoxy, and pentafluoroethoxy.

[0129] The term “halogen” or “halo” means fluoro, chloro, bromo or iodo.

[0130] As used herein, the term “cycloalkyl” means a monocyclic or polycyclic saturated or partially unsaturated carbon ring containing 3-18 carbon atoms wherein there are no delocalized pi electrons (aromaticity) shared among the ring carbon. The terms “C₃-C₁₀cycloalkyl”, “C₃-C₈cycloalkyl”, “C₄-C₁₀cycloalkyl” and “C₄-C₇cycloalkyl” are to be construed accordingly. The term polycyclic encompasses bridged (e.g., norbornane), fused (e.g., decalin) and spirocyclic cycloalkyl. Preferably, cycloalkyl, e.g., C₃-C₁₀cycloalkyl, is a monocyclic, bridged or spirocyclic hydrocarbon group of 3 to 10 carbon atoms.

[0131] Examples of cycloalkyl groups include, without limitations, cyclopropenyl, cyclopropyl cyclobutyl, cyclobutenyl, cyclopentyl, cyclohexyl, cycloheptanyl, cyclooctanyl, norbornanyl, norbornenyl, spiro[3.3]heptanyl (e.g., spiro[3.3]heptan-6-yl), bicyclo[2.2.2]octanyl, bicyclo[2.2.2]octenyl, adamantyl and derivatives thereof. Preferably, the cycloalkyl group is saturated.

[0132] Preferred examples of C₃-C₁₀cycloalkyl include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, spiro[3.3]heptanyl (e.g., spiro[3.3]heptan-6-yl), bicyclo[1.1.1]pentyl, bicyclo[2.1.1]hexyl, bicyclo[2.1.1]heptyl, bicyclo[2.2.2]octyl and adamantyl.

[0133] “Heterocyclyl” means a saturated or partially saturated monocyclic or polycyclic ring containing carbon and at least one heteroatom selected from oxygen, nitrogen, and sulfur (O, N, and S) and wherein there are no delocalized pi electrons (aromaticity) shared among the ring carbon or heteroatoms. The terms “4- to 10-membered heterocyclyl”, “4- to 6-membered heterocyclyl” and “5- or 6-membered heterocyclyl” are to be construed accordingly. The heterocyclyl ring structure may be substituted by one or more substituents. The substituents can themselves be optionally substituted. The heterocyclyl may be bonded via a carbon atom or heteroatom. The term polycyclic encompasses bridged, fused and spirocyclic heterocyclyl.

[0134] Examples of heterocyclyl rings include, but are not limited to, oxetanyl, azetidiny, tetrahydrofuranyl, tetrahydropyranyl, pyrrolidiny, oxazoliny, isoxazoliny, oxazolidiny, thiazolidiny, pyranly, thiopyranly, tetrahydropyranyl, dioxaliny, piperidiny, morpholiny, thiomorpholiny, thiomorpholiny S-oxide, thiomorpholiny S-dioxide, piperazinyl, azepiny, oxepiny, diazepiny, tropanyl, oxazolidinonyl, 1,4-dioxanyl, dihydrofuranyl, 1,3-dioxolanyl, imidazolidiny, dihydroisoxazoliny, pyrroliny, pyrazoliny, oxazepiny, dithiolanyl, homotropanyl, dihydropyranyl (e.g., 3,6-dihydro-2H-pyranyl), oxaspiroheptanyl (e.g., 2-oxaspiro[3.3]heptan-6-yl), diazabicyclo[3.2.1]octan-3-yl), 2-azaspiro[3.3]heptanyl (e.g., 2-azaspiro[3.3]heptan-6-yl), and the like.

[0135] Preferred examples of heterocyclyl include, without limitations, oxetanyl, azetidiny, pyrrolidiny, tetrahydrofuranyl, tetrahydrothienyl, piperidiny, piperazinyl, dihydroisoxazoliny, tetrahydropyranyl, morpholiny, dihydropyranyl (e.g., 3,6-dihydro-2H-pyranyl) 2-azaspiro[3.3]heptanyl (e.g., 2-azaspiro[3.3]heptan-6-yl) and oxaspiroheptanyl (e.g., 2-oxaspiro[3.3]heptan-6-yl).

[0136] As used herein, the term “aryl” as used herein means monocyclic, bicyclic or polycyclic carbocyclic aromatic rings. Examples of aryl include, but are not limited to, phenyl, naphthyl (e.g., naphth-1-yl, naphth-2-yl), anthryl (e.g., anthr-1-yl, anthr-9-yl), phenanthryl (e.g., phenanthr-

1-yl, phenanthr-9-yl), and the like. Aryl is also intended to include monocyclic, bicyclic or polycyclic carbocyclic aromatic rings substituted with carbocyclic aromatic rings. Representative examples are biphenyl (e.g., biphenyl-2-yl, biphenyl-3-yl, biphenyl-4-yl), phenylnaphthyl (e.g., 1-phenylnaphth-2-yl, 2-phenylnaphth-1-yl), and the like. Aryl is also intended to include partially saturated bicyclic or polycyclic carbocyclic rings with at least one unsaturated moiety (e.g., a benzo moiety). Representative examples are, indanyl (e.g., indan-1-yl, indan-5-yl), indenyl (e.g., inden-1-yl, inden-5-yl), 1,2,3,4-tetrahydronaphthyl (e.g., 1,2,3,4-tetrahydronaphth-1-yl, 1,2,3,4-tetrahydronaphth-2-yl, 1,2,3,4-tetrahydronaphth-6-yl), 1,2-dihydronaphthyl (e.g., 1,2-dihydronaphth-1-yl, 1,2-dihydronaphth-4-yl, 1,2-dihydronaphth-6-yl), fluorenyl (e.g., fluoren-1-yl, fluoren-4-yl, fluoren-9-yl), and the like. Aryl is also intended to include partially saturated bicyclic or polycyclic carbocyclic aromatic rings containing one or two bridges. Representative examples are, benzonorbornyl (e.g., benzonorborn-3-yl, benzonorborn-6-yl), 1,4-ethano-1,2,3,4-tetrahydronaphthyl (e.g., 1,4-ethano-1,2,3,4-tetrahydronaphth-2-yl, 1,4-ethano-1,2,3,4-tetrahydronaphth-10-yl), and the like. The term “C₆-C₁₀aryl” is to be construed accordingly.

[0137] Preferred examples of aryl include, but are not limited to, indenyl, (e.g., inden-1-yl, inden-5-yl) phenyl (C₆H₅), naphthyl (C₁₀H₇) (e.g., naphth-1-yl, naphth-2-yl), indanyl (e.g., indan-1-yl, indan-5-yl), and tetrahydronaphthalenyl (e.g., 1,2,3,4-tetrahydronaphthalenyl).

[0138] Preferably, C₆-C₁₀aryl refers to a monocyclic or bicyclic carbocyclic aromatic ring.

[0139] Preferred examples of C₆-C₁₀aryl include, but are not limited to, phenyl and naphthyl. In an embodiment, C₆-C₁₀aryl is phenyl.

[0140] As used herein, the term “heteroaryl” as used herein is intended to include monocyclic heterocyclic aromatic rings containing one or more heteroatoms selected from oxygen, nitrogen, and sulfur (O, N, and S). Representative examples are pyrrolyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isothiazolyl, isooxazolyl, triazolyl, (e.g., 1,2,4-triazolyl), oxadiazolyl, (e.g., 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl), thiadiazolyl (e.g., 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl), tetrazolyl, pyranyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, 1,2,3-triazinyl, 1,2,4-triazinyl, 1,3,5-triazinyl, thiadiazinyl, azepinyl, azecinyl, and the like.

[0141] Heteroaryl is also intended to include bicyclic heterocyclic aromatic rings containing one or more heteroatoms selected from oxygen, nitrogen, and sulfur (O, N, and S). Representative examples are indolyl, isoindolyl, benzofuranyl, benzothiophenyl, indazolyl, benzopyranyl, benzimidazolyl, benzothiazolyl, benzisothiazolyl, benzoxazolyl, benzisoxazolyl, benzoxazinyl, benzotriazolyl, naphthyridinyl, phthalazinyl, pteridinyl, purinyl, quinazolinyl, cinnolinyl, quinolinyl, isoquinolinyl, quinoxalinyl, oxazolopyridinyl, isooxazolopyridinyl, pyrrolopyridinyl, furopyridinyl, thienopyridinyl, imidazopyridinyl, imidazopyrimidinyl, pyrazolopyrimidinyl, pyrazolopyrimidinyl, pyrazolotriazinyl, thiazolopyrimidinyl, thiazolopyrimidinyl, imidazothiazolyl, triazolopyrimidinyl, and the like.

[0142] Heteroaryl is also intended to include polycyclic heterocyclic aromatic rings containing one or more heteroatoms selected from oxygen, nitrogen, and sulfur (O, N, and

S). Representative examples are carbazolyl, phenoxazinyl, phenazinyl, acridinyl, phenothiazinyl, carbolinyl, phenanthrolinyl, and the like.

[0143] Heteroaryl is also intended to include partially saturated monocyclic, bicyclic or polycyclic heterocyclals containing one or more heteroatoms selected oxygen, nitrogen, and sulfur (O, N, and S). Representative examples are imidazolyl, indolyl, dihydrobenzofuranyl, dihydrobenzothienyl, dihydrobenzopyranyl, dihydropyridooxazinyl, dihydrobenzodioxinyl (e.g., 2,3-dihydrobenzo[b][1,4]dioxinyl), benzodioxolyl (e.g., benzo[d][1,3]dioxole), dihydrobenzooxazinyl (e.g., 3,4-dihydro-2H-benzo[b][1,4]oxazine), tetrahydroindazolyl, tetrahydrobenzimidazolyl, tetrahydroimidazo[4,5-c]pyridyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, tetrahydroquinoxalinyl, and the like.

[0144] The heteroaryl ring structure may be substituted by one or more substituents. The substituents can themselves be optionally substituted. The heteroaryl ring may be bonded via a carbon atom or heteroatom.

[0145] The term “5-10 membered heteroaryl” is to be construed accordingly.

[0146] Examples of 5-10 membered heteroaryl include, but are not limited to, indolyl, imidazopyridyl, isoquinolinyl, benzooxazolonyl, pyridinyl, pyrimidinyl, pyridinonyl, benzotriazolyl, pyridazinyl, pyrazolotriazinyl, indazolyl, benzimidazolyl, quinolinyl, triazolyl, (e.g., 1,2,4-triazolyl), pyrazolyl, thiazolyl, oxazolyl, isooxazolyl, pyrrolyl, oxadiazolyl, (e.g., 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl), imidazolyl, pyrrolopyridinyl, tetrahydroindazolyl, quinoxalinyl, thiadiazolyl (e.g., 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl), pyrazinyl, oxazolopyridinyl, pyrazolopyrimidinyl, benzoxazolyl, indolyl, isooxazolopyridinyl, dihydropyridooxazinyl, tetrazolyl, dihydrobenzodioxinyl (e.g., 2,3-dihydrobenzo[b][1,4]dioxinyl), benzodioxolyl (e.g., benzo[d][1,3]dioxole) and dihydrobenzooxazinyl (e.g., 3,4-dihydro-2H-benzo[b][1,4]oxazine).

[0147] As used herein, the term “oxo” refers to the radical =O.

[0148] “Cyano” or “—CN” means a substituent having a carbon atom joined to a nitrogen atom by a triple bond, e.g., C≡N.

[0149] The term “C₂-C₆alkenyl” as used herein represents a branched or straight hydrocarbon group having from 2 to 6 carbon atoms and at least one double bond. Representative examples are ethenyl (or vinyl), propenyl (e.g., prop-1-enyl, prop-2-enyl), 2-methylprop-1-enyl, 2-methylprop-2-enyl, 1,1-(dimethyl)prop-2-enyl, butadienyl (e.g., buta-1,3-dienyl), butenyl (e.g., but-1-en-1-yl, but-2-en-1-yl), 2-methylbut-1-enyl, pentenyl (e.g., pent-1-enyl, pent-2-enyl), hexenyl (e.g., hex-1-enyl, hex-2-enyl, hex-3-enyl), 2-methylpent-3-enyl, and the like.

[0150] As used herein, the term “bridging ring” refers to a ring formed at two non-adjacent carbon atoms of the heterocycloalkyl moiety of formula (I), linked to form a C₁-C₃ alkylene linker, wherein one of the carbon atoms of said linker is optionally replaced by a heteroatom selected from nitrogen, oxygen and sulfur. In a preferred embodiment, the alkylene linker comprises carbon atoms only.

[0151] As used herein, the term “C₁-C₃alkylene” refers to a straight hydrocarbon chain bivalent radical consisting solely of carbon and hydrogen atoms, containing no unsaturation, having from one to three carbon atoms.

[0152] As used herein, the term “optionally substituted” includes unsubstituted or substituted.

[0153] As used herein, “/” denotes the point of attachment to the other part of the molecule.

[0154] As used herein, the term nitrogen protecting group (PG) in a compound of Formula (X) or any intermediates in any of the general schemes 1 to 5 and subformulae thereof refers to a group that should protect the functional groups concerned against unwanted secondary reactions, such as acylations, etherifications, esterifications, oxidations, solvolysis and similar reactions. It may be removed under deprotection conditions. Depending on the protecting group employed, the skilled person would know how to remove the protecting group to obtain the free amine NH_2 group by reference to known procedures. These include reference to organic chemistry textbooks and literature procedures such as J. F. W. McOmie, “Protective Groups in Organic Chemistry”, Plenum Press, London and New York 1973; T. W. Greene and P. G. M. Wuts, “Greene’s Protective Groups in Organic Synthesis”, Fourth Edition, Wiley, New York 2007; in “The Peptides”; Volume 3 (editors: E. Gross and J. Meienhofer), Academic Press, London and New York 1981; P. J. Kocienski, “Protecting Groups”, Third Edition, Georg Thieme Verlag, Stuttgart and New York 2005; and in “Methoden der organischen Chemie” (Methods of Organic Chemistry), Houben Weyl, 4th edition, Volume 15/I, Georg Thieme Verlag, Stuttgart 1974.

[0155] Preferred nitrogen protecting groups generally comprise: $\text{C}_1\text{-C}_6$ alkyl (e.g., tert-butyl), preferably $\text{C}_1\text{-C}_4$ alkyl, more preferably $\text{C}_1\text{-C}_2$ alkyl, most preferably C_1 alkyl which is mono-, di- or tri-substituted by trialkylsilyl- $\text{C}_1\text{-C}_7$ alkoxy (e.g., trimethylsilyloxy), aryl, preferably phenyl, or a heterocyclic group (e.g., benzyl, cumyl, benzhydryl, pyrrolidinyl, trityl, pyrrolidinylmethyl, 1-methyl-1,1-dimethylbenzyl, (phenyl)methylbenzene) wherein the aryl ring or the heterocyclic group is unsubstituted or substituted by one or more, e.g., two or three, residues, e.g., selected from the group consisting of $\text{C}_1\text{-C}_7$ alkyl, hydroxy, $\text{C}_1\text{-C}_7$ alkoxy (e.g., para-methoxy benzyl (PMB)), $\text{C}_2\text{-C}_8$ -alkanoyl-oxy, halogen, nitro, cyano, and CF_3 , aryl- $\text{C}_1\text{-C}_2$ -alkoxycarbonyl (preferably phenyl- $\text{C}_1\text{-C}_2$ -alkoxycarbonyl (e.g., benzyloxycarbonyl (Cbz), benzyloxymethyl (BOM), pivaloyloxymethyl (POM)), $\text{C}_1\text{-C}_{10}$ -alkenyloxycarbonyl, $\text{C}_1\text{-C}_6$ alkylcarbonyl (e.g., acetyl or pivaloyl), $\text{C}_6\text{-C}_{10}$ -arylcarbonyl; $\text{C}_1\text{-C}_6$ -alkoxycarbonyl (e.g., tertbutoxycarbonyl (Boc), methylcarbonyl, trichloroethoxycarbonyl (Troc), pivaloyl (Piv), allyloxycarbonyl), $\text{C}_6\text{-C}_{10}$ -aryl- $\text{C}_1\text{-C}_6$ -alkoxycarbonyl (e.g., 9-fluorenylmethyloxycarbonyl (Fmoc)), allyl or cinnamyl, sulfonyl or sulfenyl, succinimidyl group, silyl groups (e.g., triarylsilyl, trialkylsilyl, triethylsilyl (TES), trimethylsilylethoxymethyl (SEM), trimethylsilyl (TMS), triisopropylsilyl or tertbutyldimethylsilyl).

[0156] According to the disclosure, the preferred nitrogen protecting group (PG) can be selected from the group comprising tert-butyloxycarbonyl (Boc), benzyloxycarbonyl (Cbz), para-methoxy benzyl (PMB), 2,4-dimethoxybenzyl (DMB), methyloxycarbonyl, trimethylsilylethoxymethyl (SEM) and benzyl. The nitrogen protecting group (PG) is preferably an acid labile protecting group, e.g., tert-butyloxycarbonyl (Boc), 2,4-dimethoxybenzyl (DMB).

[0157] In some embodiments, the compounds of the disclosure are selective over other proteins.

[0158] As used herein, the term “therapeutic agent” in connection with methods of reducing WIZ protein expres-

sion levels and/or inducing fetal hemoglobin (HbF) expression, refers to a substance that results in a detectably lower expression of WIZ gene or WIZ protein or lower activity level of WIZ proteins as compared to those levels without such substance.

[0159] As used herein “modulator” or “degrader”, means, for example, a compound of the disclosure, that effectively modulates, decreases, or reduces the levels of a specific protein (e.g., WIZ) or degrades a specific protein (e.g., WIZ). The amount of a specific protein (e.g., WIZ) degraded can be measured by comparing the amount of the specific protein (e.g., WIZ) remaining after treatment with a compound of the disclosure as compared to the initial amount or level of the specific protein (e.g., WIZ) present as measured prior to treatment with a compound of the disclosure.

[0160] As used herein “selective modulator”, “selective degrader”, or “selective compound” means, for example, a compound of the disclosure, that effectively modulates, decreases, or reduces the levels of a specific protein (e.g., WIZ) or degrades a specific protein (e.g., WIZ) to a greater extent than any other protein. A “selective modulator”, “selective degrader”, or “selective compound” can be identified, for example, by comparing the ability of a compound to modulate, decrease, or reduce the levels of or to degrade a specific protein (e.g., WIZ) to its ability to modulate, decrease, or reduce the levels of or to degrade other proteins. In some embodiments, the selectivity can be identified by measuring the EC_{50} or IC_{50} of the compounds. Degradation may be achieved through mediation of an E3 ligase, e.g., E3-ligase complexes comprising the protein Cereblon.

[0161] In one embodiment, the specific protein degraded is WIZ protein. In an embodiment, at least about 30% of WIZ is degraded compared to initial levels. In an embodiment, at least about 40% of WIZ is degraded compared to initial levels. In an embodiment, at least about 50% of WIZ is degraded compared to initial levels. In an embodiment, at least about 60% of WIZ is degraded compared to initial levels. In an embodiment, at least about 70% of WIZ is degraded compared to initial levels. In an embodiment, at least about 75% of WIZ is degraded compared to initial levels. In an embodiment, at least about 80% of WIZ is degraded compared to initial levels. In an embodiment, at least about 85% of WIZ is degraded compared to initial levels. In an embodiment, at least about 90% of WIZ is degraded compared to initial levels. In an embodiment, at least about 95% of WIZ is degraded compared to initial levels. In an embodiment, over 95% of WIZ is degraded compared to initial levels. In an embodiment, at least about 99% of WIZ is degraded compared to initial levels.

[0162] In an embodiment, the WIZ is degraded in an amount of from about 30% to about 99% compared to initial levels. In an embodiment, the WIZ is degraded in an amount of from about 40% to about 99% compared to initial levels. In an embodiment, the WIZ is degraded in an amount of from about 50% to about 99% compared to initial levels. In an embodiment, the WIZ is degraded in an amount of from about 60% to about 99% compared to initial levels. In an embodiment, the WIZ is degraded in an amount of from about 70% to about 99% compared to initial levels. In an embodiment, the WIZ is degraded in an amount of from about 80% to about 99% compared to initial levels. In an embodiment, the WIZ is degraded in an amount of from about 90% to about 99% compared to initial levels. In an embodiment, the WIZ is degraded in an amount of from

about 95% to about 99% compared to initial levels. In an embodiment, the WIZ is degraded in an amount of from about 90% to about 95% compared to initial levels.

[0163] As used herein, the terms “inducing fetal hemoglobin”, “fetal hemoglobin induction”, or “increasing fetal hemoglobin expression” refer to increasing the percentage of HbF in the blood of a subject. In an embodiment, the amount of total HbF in the blood of the subject increases. In an embodiment, the amount of total hemoglobin in the blood of the subject increases. In an embodiment, the amount of HbF is increased by at least about 10%, or at least about 20%, or at least about 30%, or at least about 40%, or at least about 50%, or at least about 60%, or at least about 70%, or at least about 80%, or at least about 90%, or at least about 100%, or more than 100%, for example, at least about 2-fold, or at least about 3-fold, or at least about 4-fold, or at least about 5-fold, or at least about 6-fold, or at least about 7-fold, or at least about 8-fold, or at least about 9-fold, or at least about 10-fold, or more than 10-fold as compared to either in the absence of a compound disclosed herein.

[0164] In an embodiment, the total hemoglobin in the blood, e.g., the blood in a subject, is increased by at least about 10%, or at least about 20%, or at least about 30%, or at least about 40%, or at least about 50%, or at least about 60%, or at least about 70%, or at least about 80%, or at least about 90%, or at least about 100%, or more than 100%, for example, at least about 2-fold, or at least about 3-fold, or at least about 4-fold, or at least about 5-fold, or at least about 6-fold, or at least about 7-fold, or at least about 8-fold, or at least about 9-fold, or at least about 10-fold, or more than 10-fold as compared to either in the absence of a compound disclosed herein.

[0165] The term “a therapeutically effective amount” of a compound of the disclosure refers to an amount of the compound of the disclosure that will elicit the biological or medical response of a subject, for example, reduction or inhibition of an enzyme or a protein activity, or ameliorate symptoms, alleviate conditions, slow or delay disease progression, or prevent a disease, etc. In one embodiment, the term “a therapeutically effective amount” refers to the amount of the compound of the disclosure that, when administered to a subject, is effective to (1) at least partially alleviate, prevent and/or ameliorate a condition, or a disorder or a disease (i) mediated by WIZ, or (ii) associated with WIZ activity, or (iii) characterized by activity (normal or abnormal) of WIZ; (2) reduce or inhibit the activity of WIZ; or (3) reduce or inhibit the expression of WIZ. In another embodiment, the term “a therapeutically effective amount” refers to the amount of the compound of the disclosure that, when administered to a cell, or a tissue, or a non-cellular biological material, or a medium, is effective to at least partially reducing or inhibiting the activity of WIZ; or at least partially reducing or inhibiting the expression of WIZ.

[0166] “HbF-dependent disease or disorder” means any disease or disorder which is directly or indirectly affected by the modulation of HbF protein levels.

[0167] As used herein, the term “subject” refers to primates (e.g., humans, male or female), dogs, rabbits, guinea pigs, pigs, rats and mice. In certain embodiments, the subject is a primate. In yet other embodiments, the subject is a human.

[0168] As used herein, the term “inhibit”, “inhibition” or “inhibiting” refers to the reduction or suppression of a given

condition, symptom, or disorder, or disease, or a significant decrease in the baseline activity of a biological activity or process.

[0169] As used herein, the term “treat”, “treating” or “treatment” of any disease or disorder refers to alleviating or ameliorating the disease or disorder (i.e., slowing or arresting the development of the disease or at least one of the clinical symptoms thereof); or alleviating or ameliorating at least one physical parameter or biomarker associated with the disease or disorder, including those which may not be discernible to the patient.

[0170] As used herein, the term “prevent”, “preventing” or “prevention” of any disease or disorder refers to the prophylactic treatment of the disease or disorder; or delaying the onset or progression of the disease or disorder


[0171] As used herein, a subject is “in need of” a treatment if such subject would benefit biologically, medically or in quality of life from such treatment.

[0172] As used herein, the term “a,” “an,” “the” and similar terms used in the context of the disclosure (especially in the context of the claims) are to be construed to cover both the singular and plural unless otherwise indicated herein or clearly contradicted by the context.

[0173] Various enumerated embodiments of the disclosure are described herein. It will be recognized that features specified in each embodiment may be combined with other specified features to provide further embodiments of the disclosure.

ENUMERATED EMBODIMENTS

[0174] Embodiment 1. A compound of Formula (I^a) or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein:



[0175] --- is a single bond or a double bond;

[0176] X is selected from CH, CF, and N;

[0177] R^x is selected from hydrogen, C₁-C₆alkyl, halo (e.g., F, Cl), C₁-C₆alkoxy, and C₃-C₈cycloalkyl;

[0178] R¹ is selected from hydrogen and C₁-C₆alkyl;

[0179] R¹ is selected from hydrogen and C₁-C₆alkyl;

[0180] each R² is independently selected from C₁-C₆alkyl, C₁-C₆haloalkyl, halo, and oxo, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{2a}; or 2 R² on non-adjacent carbon atoms together with the non-adjacent carbon atoms to which they are attached form a bridging ring;

[0181] R^{2a} is selected from C₁-C₆alkoxy and hydroxyl;

[0182] R³ is selected from hydrogen, C₁-C₈alkyl, C₂-C₆alkenyl, —SO₂R⁴, C₁-C₆haloalkyl, —C(=O)—O—(R⁵), —C(=O)—(R⁶), C₃-C₁₀cycloalkyl, and a 4- to 10-membered heterocyclyl comprising 1-2 heteroa-

toms independently selected from N, O and S, wherein the C₁-C₈alkyl and C₁-C₆haloalkyl are each independently substituted with 0-3 occurrences of R^{3a}, and wherein the C₃-C₁₀cycloalkyl and 4- to 10-membered heterocyclyl are each independently substituted with 0-3 occurrences of R^{3b};

[0183] or

[0184] R³ together with the nitrogen atom to which it is attached and R² together with the carbon atom to which it is attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O and S, which 5- or 6-membered heterocyclyl is substituted with 0-2 occurrences of an oxo group;

[0185] each R^{3a} is independently selected from C₃-C₁₀cycloalkyl, a 4- to 10-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, a 5- to 10-membered heteroaryl comprising 1-4 heteroatoms independently selected from N, O, and S, C₆-C₁₀aryl, C₁-C₆alkoxyl, hydroxyl, and —C(=O)—NR⁷R⁸, wherein the C₃-C₁₀cycloalkyl, 4- to 6-membered heterocyclyl, 5- to 10-membered heteroaryl and C₆-C₁₀aryl are substituted with 0-4 occurrences of R^{3b};

[0186] each R^{3b} is independently selected from C₁-C₆alkoxyl, halo, C₁-C₆haloalkyl, C₁-C₆haloalkoxyl, C₁-C₆alkyl, —CN, —SO₂NR⁷R⁸, —SO₂R⁴, and hydroxyl;

[0187] R⁴ is selected from C₃-C₈cycloalkyl, C₁-C₆alkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, C₆-C₁₀aryl, and —NR^{4b}R^{4c}, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{4a};

[0188] R^{4a} is selected from C₃-C₈cycloalkyl, C₆-C₁₀aryl, and C₁-C₆alkoxyl;

[0189] R^{4b} is selected from hydrogen, and C₁-C₆alkyl;

[0190] R^{4c} is selected from hydrogen, C₁-C₆alkyl, and C₃-C₈cycloalkyl;

[0191] R⁵ is selected from C₁-C₆alkyl, C₃-C₈cycloalkyl, and C₆-C₁₀aryl;

[0192] R⁶ is selected from C₁-C₆alkyl, C₃-C₈cycloalkyl, C₆-C₁₀aryl, a 4- to 10-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, and —NR^{4b}R^{4c}, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{6a}, the C₃-C₈cycloalkyl is substituted with 0-1 occurrence of R^{6b}, and the 4- to 10-membered heterocyclyl is substituted with 0-1 occurrence of C₁-C₆alkyl;

[0193] R^{6a} is selected from C₆-C₁₀aryl and C₃-C₈cycloalkyl;

[0194] R^{6b} is selected from halo, C₁-C₆haloalkyl, C₁-C₆haloalkoxyl, and C₁-C₆alkyl;

[0195] R⁷ is selected from hydrogen and C₁-C₆alkyl;

[0196] R⁸ is selected from hydrogen and C₁-C₆alkyl;

[0197] or


[0198] R⁷ and R⁸ together with the nitrogen atom to which they are attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O, and S;

[0199] n is 0, 1, 2, 3 or 4;

[0200] m is 0, 1 or 2; and

[0201] p is 0 or 1.

[0202] Embodiment 2. A compound of Formula (I') or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein:



[0203] — is a single bond or a double bond;

[0204] X is selected from CH, CF, and N;

[0205] R¹ is selected from hydrogen and C₁-C₆alkyl;

[0206] R¹ is selected from hydrogen and C₁-C₆alkyl;

[0207] each R² is independently selected from C₁-C₆alkyl, C₁-C₆haloalkyl, halo, and oxo, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{2a}; or 2 R² on non-adjacent carbon atoms together with the non-adjacent carbon atoms to which they are attached form a bridging ring;

[0208] R^{2a} is selected from C₁-C₆alkoxyl and hydroxyl;

[0209] R³ is selected from hydrogen, C₁-C₈alkyl, C₂-C₆alkenyl, —SO₂R⁴, C₁-C₆haloalkyl, —C(=O)—O—(R⁵) and —C(=O)—(R⁶), wherein the C₁-C₈alkyl and C₁-C₆haloalkyl are independently substituted with 0-3 occurrences of R^{3a};

[0210] or

[0211] R³ together with the nitrogen atom to which it is attached and R² together with the carbon atom to which it is attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O and S;

[0212] each R^{3a} is independently selected from C₃-C₁₀cycloalkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, a 5- to 10-membered heteroaryl comprising 1-4 heteroatoms independently selected from N, O, and S, C₆-C₁₀aryl, C₁-C₆alkoxyl, hydroxyl, and —C(=O)—NR⁷R⁸, wherein the C₃-C₁₀cycloalkyl, 4- to 6-membered heterocyclyl, 5- to 10-membered heteroaryl and C₆-C₁₀aryl are substituted with 0-4 occurrences of R^{3b};

[0213] each R^{3b} is independently selected from C₁-C₆alkoxyl, halo, C₁-C₆haloalkyl, C₁-C₆haloalkoxyl, C₁-C₆alkyl, —CN, —SO₂NR⁷R⁸, —SO₂R⁴, and hydroxyl;

[0214] R⁴ is selected from C₃-C₈cycloalkyl, C₁-C₆alkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, and C₆-C₁₀aryl, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{4a};

[0215] R^{4a} is selected from C₃-C₈cycloalkyl, C₆-C₁₀aryl, and C₁-C₆alkoxyl;

[0216] R⁵ is selected from C₁-C₆alkyl, C₃-C₈cycloalkyl, and C₆-C₁₀aryl;

[0217] R⁶ is selected from C₁-C₆alkyl, C₃-C₈cycloalkyl, and C₆-C₁₀aryl, wherein the

C₁-C₆alkyl is substituted with 0-1 occurrence of R^{6a} and the C₃-C₈cycloalkyl is substituted with 0-1 occurrence of R^{6b};

[0218] R^{6a} is selected from C₆-C₁₀aryl and C₃-C₈cycloalkyl;

[0219] R^{6b} is selected from halo, C₁-C₆haloalkyl, C₁-C₆haloalkoxyl, and C₁-C₆alkyl;

[0220] R⁷ is selected from hydrogen and C₁-C₆alkyl;

[0221] R^a is selected from hydrogen and C₁-C₆alkyl;

[0222] or


[0223] R⁷ and R⁸ together with the nitrogen atom to which they are attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O, and S;

[0224] n is 0, 1, 2, 3 or 4;

[0225] m is 0, 1 or 2; and

[0226] p is 0 or 1.

[0227] Embodiment 3. A compound of Formula (I) or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein:



[0228] X is selected from X is selected from CH, CF, and N;

[0229] R¹ is selected from hydrogen and C₁-C₆alkyl;

[0230] R¹ is selected from hydrogen and C₁-C₆alkyl;

[0231] each R² is independently selected from C₁-C₆alkyl, C₁-C₆haloalkyl, halo, and oxo, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{2a}; or 2 R² on non-adjacent carbon atoms together with the non-adjacent carbon atoms to which they are attached form a bridging ring;

[0232] R^{2a} is selected from C₁-C₆alkoxyl and hydroxyl;

[0233] R³ is selected from hydrogen, C₁-C₈alkyl, C₂-C₆alkenyl, —SO₂R⁴, C₁-C₆haloalkyl, —C(=O)—O—(R⁵) and —C(=O)—(R⁶), wherein the C₁-C₈alkyl and C₁-C₆haloalkyl are independently substituted with 0-3 occurrences of R^{3a};

[0234] or

[0235] R³ together with the nitrogen atom to which it is attached and R² together with the carbon atom to which it is attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O and S;

[0236] each R^{3a} is independently selected from C₃-C₁₀cycloalkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, a 5- to 10-membered heteroaryl comprising 1-4 heteroatoms independently selected from N, O, and S, C₆-C₁₀aryl, C₁-C₆alkoxyl, hydroxyl, and —C(=O)—NR⁷R⁸, wherein the C₃-C₁₀cycloalkyl, 4- to 6-membered heterocyclyl, 5- to

10-membered heteroaryl and C₆-C₁₀aryl are substituted with 0-4 occurrences of R^{3b};

[0237] each R^{3b} is independently selected from C₁-C₆alkoxyl, halo, C₁-C₆haloalkyl, C₁-C₆haloalkoxyl, C₁-C₆alkyl, —CN, —SO₂NR⁷R⁸, —SO₂R⁴, and hydroxyl;

[0238] R⁴ is selected from C₃-C₈cycloalkyl, C₁-C₆alkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, and C₆-C₁₀aryl, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{4a};

[0239] R^{4a} is selected from C₃-C₈cycloalkyl, C₆-C₁₀aryl, and C₁-C₆alkoxyl;

[0240] R⁵ is selected from C₁-C₆alkyl, C₃-C₈cycloalkyl, and C₆-C₁₀aryl;

[0241] R⁶ is selected from C₁-C₆alkyl, C₃-C₈cycloalkyl, and C₆-C₁₀aryl, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{6a} and the C₃-C₈cycloalkyl is substituted with 0-1 occurrence of R^{6b};

[0242] R^{6a} is selected from C₆-C₁₀aryl and C₃-C₈cycloalkyl;

[0243] R^{6b} is selected from halo, C₁-C₆haloalkyl, C₁-C₆haloalkoxyl, and C₁-C₆alkyl;

[0244] R⁷ is selected from hydrogen and C₁-C₆alkyl;

[0245] R⁸ is selected from hydrogen and C₁-C₆alkyl;

[0246] or

[0247] R⁷ and R⁸ together with the nitrogen atom to which they are attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O, and S;

[0248] n is 0, 1, 2, 3 or 4;

[0249] m is 0, 1 or 2; and

[0250] p is 0 or 1.

[0251] Embodiment 4. The compound of any one of Embodiments 1 to 3, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein X is selected from CH, CF, and N;

[0252] R¹ is selected from hydrogen and C₁-C₃alkyl;

[0253] R¹ is selected from hydrogen and C₁-C₃alkyl;

[0254] each R² is independently selected from unsubstituted C₁-C₆alkyl, C₁-C₆haloalkyl and halo; or 2 R² on non-adjacent carbon atoms together with the non-adjacent carbon atoms to which they are attached form a bridging ring;

[0255] R³ is selected from hydrogen, C₁-C₈alkyl, C₂-C₆alkenyl, —SO₂R⁴, C₁-C₆haloalkyl, —C(=O)—O—(R⁵) and —C(=O)—(R⁶), wherein the C₁-C₈alkyl and C₁-C₆haloalkyl are independently substituted with 0-3 occurrences of R^{3a};

[0256] or

[0257] R³ together with the nitrogen atom to which it is attached and R² together with the carbon atom to which it is attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N and O;

[0258] each R^{3a} is independently selected from C₃-C₁₀cycloalkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, a 5- to 10-membered heteroaryl comprising 1-4 heteroatoms independently selected from N, O, and S, C₆-C₁₀aryl, C₁-C₆alkoxyl, hydroxyl, and —C(=O)—NR⁷R⁸, wherein the C₃-C₁₀cycloalkyl, 4- to 6-membered heterocyclyl, 5- to

- 10-membered heteroaryl and C₆-C₁₀aryl are substituted with 0-4 occurrences of R^{3b};
- [0259] each R^{3b} is independently selected from C₁-C₆alkoxy, halo, C₁-C₆haloalkyl, C₁-C₆haloalkoxy, C₁-C₆alkyl, —CN, —SO₂NR⁷R⁸, —SO₂R⁴, and hydroxyl;
- [0260] R⁴ is selected from C₃-C₈cycloalkyl, C₁-C₆alkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, and C₆-C₁₀aryl, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{4a};
- [0261] R^{4a} is selected from C₃-C₈cycloalkyl, C₆-C₁₀aryl, and C₁-C₆alkoxy;
- [0262] R⁵ is selected from C₁-C₆alkyl, C₃-C₈cycloalkyl, and C₆-C₁₀aryl;
- [0263] R⁶ is selected from C₁-C₆alkyl, C₃-C₈cycloalkyl, and C₆-C₁₀aryl, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{6a} and the C₃-C₈cycloalkyl is substituted with 0-1 occurrence of R^{6b};
- [0264] R^{6a} is selected from C₆-C₁₀aryl and C₃-C₈cycloalkyl;
- [0265] R^{6b} is selected from chloro, fluoro, C₁-C₆haloalkyl, C₁-C₆haloalkoxy, and C₁-C₆alkyl;
- [0266] R⁷ is selected from hydrogen and C₁-C₆alkyl;
- [0267] R⁸ is selected from hydrogen and C₁-C₆alkyl;
- [0268] or
- [0269] R⁷ and R⁸ together with the nitrogen atom to which they are attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O, and S;
- [0270] n is 0, 1, 2 or 3;
- [0271] m is 0, 1 or 2; and
- [0272] p is 0 or 1.
- [0273] Embodiment 5. The compound of any one of the preceding Embodiments, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein
- [0274] X is selected from CH and N;
- [0275] R¹ is selected from hydrogen and methyl;
- [0276] R¹ is selected from hydrogen and methyl;
- [0277] each R² is independently selected from unsubstituted C₁-C₆alkyl and halo; or 2 R² on non-adjacent carbon atoms together with the non-adjacent carbon atoms to which they are attached form a C₁-C₃alkylene bridging ring;
- [0278] R³ is selected from hydrogen, C₁-C₈alkyl, C₂-C₆alkenyl, —SO₂R⁴, C₁-C₆haloalkyl, —C(=O)—O—(R⁵) and —C(=O)—(R⁶), wherein the C₁-C₈alkyl and C₁-C₆haloalkyl are independently substituted with 0-3 occurrences of R^{3a};
- [0279] or
- [0280] R³ together with the nitrogen atom to which it is attached and R² together with the carbon atom to which it is attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom;
- [0281] each R^{3a} is independently selected from C₃-C₁₀cycloalkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, a 5- to 10-membered heteroaryl comprising 1-4 heteroatoms independently selected from N, O, and S and phenyl, wherein the C₃-C₁₀cycloalkyl, 4- to 6-membered heterocyclyl, 5- to 10-membered heteroaryl and phenyl are substituted with 0-4 occurrences of R^{3b};
- [0282] each R^{3b} is independently selected from C₁-C₆alkoxy, halo, C₁-C₆haloalkyl, C₁-C₆haloalkoxy, C₁-C₆alkyl, —CN, —SO₂NR⁷R⁸, —SO₂R⁴, and hydroxyl;
- [0283] R⁴ is selected from C₃-C₈cycloalkyl, C₁-C₆alkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, and C₆-C₁₀aryl, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{4a};
- [0284] R^{4a} is selected from C₃-C₈cycloalkyl, C₆-C₁₀aryl, and C₁-C₆alkoxy;
- [0285] R⁵ is selected from C₁-C₆alkyl, C₃-C₈cycloalkyl, and C₆-C₁₀aryl;
- [0286] R⁶ is selected from C₁-C₆alkyl, C₃-C₈cycloalkyl, and C₆-C₁₀aryl, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{6a} and the C₃-C₈cycloalkyl is substituted with 0-1 occurrence of R^{6b};
- [0287] R^{6a} is selected from C₆-C₁₀aryl and C₃-C₈cycloalkyl;
- [0288] R^{6b} is selected from chloro, fluoro, C₁-C₆haloalkyl, C₁-C₆haloalkoxy, and C₁-C₆alkyl;
- [0289] R⁷ is selected from hydrogen and C₁-C₆alkyl;
- [0290] R⁸ is selected from hydrogen and C₁-C₆alkyl;
- [0291] or R⁷ and R⁸ together with the nitrogen atom to which they are attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O, and S;
- [0292] n is 0, 1, 2 or 3;
- [0293] m is 0, 1 or 2; and
- [0294] p is 0 or 1.
- [0295] Embodiment 6. The compound of any one of the preceding Embodiments, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein
- [0296] X is selected from CH and N;
- [0297] R¹ is hydrogen;
- [0298] R¹ is hydrogen;
- [0299] each R² is independently selected from unsubstituted C₁-C₆alkyl and fluoro; or 2 R² on non-adjacent carbon atoms together with the non-adjacent carbon atoms to which they are attached form a C₁-C₃alkylene bridging ring;
- [0300] R³ is selected from C₁-C₈alkyl, C₂-C₆alkenyl, —SO₂R⁴ and C₁-C₆haloalkyl, wherein the C₁-C₈alkyl is substituted with 0-2 occurrences of R^{3a} and the C₁-C₆haloalkyl is substituted with 0-1 occurrence of R^{3a};
- [0301] each R^{3a} is independently selected from C₃-C₁₀cycloalkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N and O, a 5- to 6-membered heteroaryl comprising 1-3 heteroatoms independently selected from N, O, and S and phenyl, wherein the C₃-C₁₀cycloalkyl, 4- to 6-membered heterocyclyl, 5- to 6-membered heteroaryl and phenyl are substituted with 0-4 occurrences of R^{3b};
- [0302] each R^{3b} is independently selected from halo, C₁-C₆haloalkyl, C₁-C₆haloalkoxy, C₁-C₆alkyl, and hydroxyl;
- [0303] R⁴ is selected from C₃-C₈cycloalkyl, C₁-C₆alkyl, a 4- to 6-membered heterocyclyl comprising

ing 1-2 heteroatoms independently selected from N, O and S, and C₆-C₁₀aryl, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{4a};

[0304] R^{4a} is selected from C₃-C₈cycloalkyl, C₆-C₁₀aryl, and C₁-C₆alkoxy;

[0305] n is 0, 1, 2 or 3;

[0306] m is 1 or 2; and

[0307] p is 0 or 1.

[0308] Embodiment 7. The compound of any one of the preceding Embodiments, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein

[0309] X is selected from CH and N;

[0310] R' is hydrogen;

[0311] R¹ is hydrogen;

[0312] each R² is independently selected from unsubstituted C₁-C₆alkyl; or 2 R² on non-adjacent carbon atoms together with the non-adjacent carbon atoms to which they are attached form a C₁-C₃alkylene bridging ring;

[0313] R³ is selected from C₁-C₈alkyl, C₂-C₆alkenyl, —SO₂R⁴ and unsubstituted C₁-C₆haloalkyl, wherein the C₁-C₈alkyl is substituted with 0-2 occurrences of R^{3a};

[0314] each R^{3a} is independently selected from C₃-C₁₀cycloalkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N and O, a 5- to 6-membered heteroaryl comprising 1-3 heteroatoms independently selected from N, O, and S and phenyl, wherein the C₃-C₁₀cycloalkyl, 4- to 6-membered heterocyclyl, 5- to 6-membered heteroaryl and phenyl are substituted with 0-3 occurrences of R^{3b};

[0315] each R^{3b} is independently selected from halo, C₁-C₆haloalkyl, C₁-C₆haloalkoxy, C₁-C₆alkyl, and hydroxyl;

[0316] R⁴ is selected from C₃-C₈cycloalkyl, C₁-C₆alkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, and C₆-C₁₀aryl, wherein the C₁-C₆alkyl is substituted with 1 occurrence of R^{4a};

[0317] R^{4a} is selected from C₃-C₈cycloalkyl, C₆-C₁₀aryl, and C₁-C₆alkoxy;

[0318] n is 0, 1 or 2;

[0319] m is 1 or 2; and

[0320] p is 1.

[0321] Embodiment 8. The compound of any one of the preceding Embodiments, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein

[0322] X is selected from CH and N;

[0323] R' is hydrogen;

[0324] R¹ is hydrogen;

[0325] each R² is independently selected from unsubstituted C₁-C₃alkyl;

[0326] R³ is selected from C₁-C₆alkyl, C₂-C₆alkenyl, —SO₂R⁴ and unsubstituted C₁-C₆haloalkyl, wherein the C₁-C₆alkyl is substituted with 0-2 occurrences of R^{3a};

[0327] each R^{3a} is independently selected from C₃-C₁₀cycloalkyl, a 4- to 6-membered heterocyclyl comprising 1 O heteroatom, a 6-membered heteroaryl comprising 1-2 N heteroatoms and phenyl, wherein the C₃-C₁₀cycloalkyl, 4- to 6-membered heterocyclyl, 6-membered heteroaryl and phenyl are substituted with 0-2 occurrences of R^{3b};

[0328] each R^{3b} is independently selected from chloro, fluoro, C₁-C₆haloalkyl, C₁-C₆haloalkoxy and C₁-C₆alkyl;

[0329] R⁴ is selected from C₃-C₈cycloalkyl, C₁-C₆alkyl, a 4- to 6-membered heterocyclyl comprising 1 O heteroatom and phenyl, wherein the C₁-C₆alkyl is substituted with 1 occurrence of R^{4a};


[0330] R^{4a} is selected from C₃-C₈cycloalkyl and phenyl;

[0331] n is 0, 1 or 2;


[0332] m is 1 or 2; and

[0333] p is 1.



[0334] Embodiment 9. The compound of Embodiment 1, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, of Formula (Ia''):



[0335] Embodiment 10. The compound of any one of Embodiments 1 and 2, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, of Formula (Ia'):




[0336] Embodiment 11. The compound of any one of the preceding Embodiments, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, of Formula (Ia):




[0337] Embodiment 12. The compound of any one of the preceding Embodiments, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein R^3 is selected from C_1 - C_6 alkyl and $-\text{CH}_2-\text{R}^{3a}$.

[0338] Embodiment 13. The compound of any one of Embodiments 1 and 9, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, of Formula (Ib''):




[0339] Embodiment 14. The compound of any one Embodiments 1, 2, 9 and 10, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, of Formula (Ib'):



[0340] Embodiment 15. The compound of any one of the preceding Embodiments, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, of Formula (Ib):

[0341] Embodiment 16. The compound of any one of Embodiments 1 and 9, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, of Formula (Ic'), wherein:



[0342] $\equiv$ is a single bond or a double bond;

[0343] X is selected from CH, CF and N;

[0344] R^x is selected from hydrogen, C_1 - C_6 alkyl, halo (e.g., F, Cl), C_1 - C_6 alkoxy, and C_3 - C_8 cycloalkyl;

[0345] R^{2b} is selected from hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, and halo, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{2a} ;

[0346] R^{2c} is selected from hydrogen and C_1 - C_6 alkyl, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{2a} ;

[0347] or R^{2b} and R^{2c} together with the carbon atoms to which they are attached form an oxo group;

[0348] each of R^{2d} and R^{2e} is independently selected from hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, halo, and oxo, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{2a} ;


[0349] R^{2f} is hydrogen;

[0350] or R^{2b} and R^{2e} or R^{2b} and R^{2f} together with the carbon atoms to which they are attached form a bridging ring;


[0351] R^{2a} is selected from C_1 - C_6 alkoxy and hydroxyl; and

[0352] R^3 is defined according to any one of the preceding Embodiments.

[0353] Embodiment 17. The compound of any one of Embodiments 1, 2, 9 and 10, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, of Formula (Ic'), wherein:



- [0354] $\equiv$ is a single bond or a double bond;
 [0355] X is selected from CH, CF and N;
 [0356] R^{2b} is selected from hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, and halo, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{2a} ;
 [0357] R^{2c} is selected from hydrogen and C_1 - C_6 alkyl, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{2a} ;
 [0358] or R^{2b} and R^{2c} together with the carbon atoms to which they are attached form an oxo group;
 [0359] each of R^{2d} and R^{2e} is independently selected from hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, halo, and oxo, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{2a} ; R^{2f} is hydrogen;
 [0360] or R^{2b} and R^{2e} or R^{2b} and R^{2f} together with the carbon atoms to which they are attached form a bridging ring;
 [0361] R^{2a} is selected from C_1 - C_6 alkoxy and hydroxyl; and
 [0362] R^3 is defined according to any one of the preceding Embodiments.
 [0363] Embodiment 18. The compound of any one of Embodiments 1 to 12, 16 and 17, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, of Formula (1c), wherein:



- [0364] X is selected from CH, CF and N;
 [0365] R^{2b} is selected from hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, and halo, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{2a} ;
 [0366] R^{2c} is selected from hydrogen and C_1 - C_6 alkyl, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{2a} ;
 [0367] or R^{2b} and R^{2c} together with the carbon atoms to which they are attached form an oxo group;

- [0368] each of R^{2d} and R^{2e} is independently selected from hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, halo, and oxo, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{2a} ;
 [0369] R^{2f} is hydrogen;
 [0370] or R^{2b} and R^{2e} or R^{2b} and R^{2f} together with the carbon atoms to which they are attached form a bridging ring;
 [0371] R^{2a} is selected from C_1 - C_6 alkoxy and hydroxyl; and
 [0372] R^3 is defined according to any one of the preceding Embodiments.
 [0373] Embodiment 19. The compound of any one of Embodiments 16 to 18, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein
 [0374] X is selected from CH and N;
 [0375] R^{2b} is selected from hydrogen, C_1 - C_3 alkyl, C_1 - C_3 haloalkyl, and halo, wherein the C_1 - C_3 alkyl is substituted with 0-1 occurrence of R^{2a} ;
 [0376] R^{2c} is selected from hydrogen and C_1 - C_3 alkyl, wherein the C_1 - C_3 alkyl is substituted with 0-1 occurrence of R^{2a} ;
 [0377] or R^{2b} and R^{2c} together with the carbon atoms to which they are attached form an oxo group;
 [0378] each of R^{2d} and R^{2e} is independently selected from hydrogen, C_1 - C_3 alkyl, C_1 - C_3 haloalkyl, halo, and oxo, wherein the C_1 - C_3 alkyl is substituted with 0-1 occurrence of R^{2a} ;
 [0379] R^{2f} is hydrogen;
 [0380] or R^{2b} and R^{2e} or R^{2b} and R^{2f} together with the carbon atoms to which they are attached form a bridging ring;
 [0381] R^{2a} is selected from C_1 - C_6 alkoxy and hydroxyl;
 [0382] R^3 is selected from C_1 - C_8 alkyl, C_2 - C_6 alkenyl, $-\text{SO}_2\text{R}^4$, C_1 - C_6 haloalkyl, $-\text{C}(=\text{O})-\text{O}-(\text{R}^5)$ and $-\text{C}(=\text{O})-(\text{R}^6)$, wherein the C_1 - C_8 alkyl and C_1 - C_6 haloalkyl are independently substituted with 0-3 occurrences of R^{3a} ;
 [0383] each R^{3a} is independently selected from C_3 - C_{10} cycloalkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, a 5- to 10-membered heteroaryl comprising 1-4 heteroatoms independently selected from N, O, and S, C_6 - C_{10} aryl, C_1 - C_6 alkoxy, hydroxyl, and $-\text{C}(=\text{O})-\text{NR}^7\text{R}^8$, wherein the C_3 - C_{10} cycloalkyl, 4- to 6-membered heterocyclyl, 5- to 10-membered heteroaryl and C_6 - C_{10} aryl are substituted with 0-4 occurrences of R^{3b} ;
 [0384] each R^{3b} is independently selected from C_1 - C_6 alkoxy, halo, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkoxy, C_1 - C_6 alkyl, $-\text{CN}$, $-\text{SO}_2\text{NR}^7\text{R}^8$, $-\text{SO}_2\text{R}^4$, and hydroxyl;
 [0385] R^4 is selected from C_3 - C_8 cycloalkyl, C_1 - C_6 alkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, and C_6 - C_{10} aryl, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{4a} ;
 [0386] R^{4a} is selected from C_3 - C_8 cycloalkyl, C_6 - C_{10} aryl, and C_1 - C_6 alkoxy;
 [0387] R^5 is selected from C_1 - C_6 alkyl and C_6 - C_{10} aryl;
 [0388] R^6 is selected from C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, and C_6 - C_{10} aryl, wherein the

C₁-C₆alkyl is substituted with 0-1 occurrence of R^{6a} and the C₃-C₈cycloalkyl is substituted with 0-1 occurrence of R^{6b};

[0389] R^{6a} is selected from C₆-C₁₀aryl and C₃-C₈cycloalkyl;

[0390] R^{6b} is selected from halo, C₁-C₆haloalkyl, C₁-C₆haloalkoxy, and C₁-C₆alkyl;

[0391] R⁷ is selected from hydrogen and C₁-C₆alkyl;

[0392] R⁸ is selected from hydrogen and C₁-C₆alkyl;

[0393] or

[0394] R⁷ and R⁸ together with the nitrogen atom to which they are attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O, and S; and

[0395] m is 1 or 2.

[0396] Embodiment 20. The compound of any one of Embodiments 16 to 19, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein

[0397] X is selected from CH and N;

[0398] each of R^{2b}, R^{2c}, R^{2d} and R^{2e} is independently selected from hydrogen and unsubstituted CO—C₃alkyl;

[0399] R^{2f} is hydrogen;

[0400] or R^{2b} and R^{2e} or R^{2b} and R^{2f} together with the carbon atoms to which they are attached form a C₁-C₃alkylene bridging ring;

[0401] R³ is selected from C₁-C₈alkyl, C₂-C₆alkenyl, —SO₂R⁴, and C₁-C₆haloalkyl, wherein the C₁-C₈alkyl and C₁-C₆haloalkyl are independently substituted with 0-3 occurrences of R^{3a};

[0402] each R^{3a} is independently selected from C₃-C₁₀cycloalkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, a 5- to 10-membered heteroaryl comprising 1-4 heteroatoms independently selected from N, O, and S, C₆-C₁₀aryl, C₁-C₆alkoxy, hydroxyl, and —C(=O)—NR⁷R⁸, wherein the C₃-C₁₀cycloalkyl, 4- to 6-membered heterocyclyl, 5- to 10-membered heteroaryl and C₆-C₁₀aryl are substituted with 0-4 occurrences of R^{3b};

[0403] each R^{3b} is independently selected from C₁-C₆alkoxy, halo, C₁-C₆haloalkyl, C₁-C₆haloalkoxy, C₁-C₆alkyl, —CN, —SO₂NR⁷R⁸, —SO₂R⁴, and hydroxyl;

[0404] R⁴ is selected from C₃-C₈cycloalkyl, C₁-C₆alkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, and C₆-C₁₀aryl, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{4a};

[0405] R^{4a} is selected from C₃-C₈cycloalkyl, C₆-C₁₀aryl, and C₁-C₆alkoxy;

[0406] R⁷ is selected from hydrogen and C₁-C₆alkyl;

[0407] R⁸ is selected from hydrogen and C₁-C₆alkyl;

[0408] or

[0409] R⁷ and R⁸ together with the nitrogen atom to which they are attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O, and S; and

[0410] m is 1 or 2.

[0411] Embodiment 21. The compound according to any of Embodiments 16 to 20, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein

[0412] X is selected from CH and N;

[0413] each of R^{2b}, R^{2c}, R^{2d} and R^{2e} is independently selected from hydrogen and unsubstituted CO—C₃alkyl;

[0414] R^{2f} is hydrogen;

[0415] R³ is selected from C₁-C₈alkyl, C₂-C₆alkenyl, —SO₂R⁴, and C₁-C₆haloalkyl, wherein the C₁-C₈alkyl and C₁-C₆haloalkyl are independently substituted with 0-3 occurrences of R^{3a};

[0416] each R^{3a} is independently selected from C₃-C₁₀cycloalkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, a 5- to 10-membered heteroaryl comprising 1-4 heteroatoms independently selected from N, O, and S and phenyl, wherein the C₃-C₁₀cycloalkyl, 4- to 6-membered heterocyclyl, 5- to 10-membered heteroaryl and phenyl are substituted with 0-4 occurrences of R^{3b};


[0417] each R^{3b} is independently selected from C₁-C₆alkoxy, halo, C₁-C₆haloalkyl, C₁-C₆haloalkoxy, C₁-C₆alkyl, and hydroxyl;

[0418] R⁴ is selected from C₃-C₈cycloalkyl, C₁-C₆alkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, and C₆-C₁₀aryl, wherein the C₁-C₆alkyl is substituted with 1 occurrence of R^{4a};

[0419] R^{4a} is selected from C₃-C₈cycloalkyl, C₆-C₁₀aryl, and C₁-C₆alkoxy; and


[0420] m is 1.

[0421] Embodiment 22. The compound of any one of Embodiments 1, 9 and 16, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, of Formula (Id''), wherein:




[0422] R^{2b}, R^{2c} and R^{2e} are defined according to any of Embodiments 16 to 21.

[0423] Embodiment 23. The compound of any one of Embodiments 1, 2, 9, 10, 16, 17 and 22, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, of Formula (Id'), wherein:




[0424] R^{2b} , R^{2c} and R^{2e} are defined according to any of Embodiments 16 to 21.

[0425] Embodiment 24. The compound of any one of Embodiments 1 to 12 and 16 to 23, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, of Formula (Id), wherein:



[0426] R^{2b} , R^{2c} and R^{2e} are defined according to any of Embodiments 16 to 21.

[0427] Embodiment 25. The compound of any one of Embodiments 1 to 12 and 16 to 24, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, of Formula (Id-1), wherein:



[0428] R^{2b} is selected from hydrogen and C_1 - C_4 alkyl; and

[0429] X and R^3 are defined according to any one of the preceding Embodiments.

[0430] Embodiment 26. The compound Embodiment 25, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein:

[0431] X is CH or N;



[0432] R^{2b} is selected from hydrogen and C_1 - C_4 alkyl;

[0433] R^3 is selected from C_1 - C_8 alkyl, $-\text{SO}_2\text{R}^4$, and $-\text{C}(=\text{O})-(\text{R}^6)$, wherein the C_1 - C_8 alkyl is independently substituted with 0-3 occurrences of R^{3a} ;


[0434] each R^{3a} is independently selected from C_3 - C_{10} cycloalkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, C_1 - C_6 alkoxy, and hydroxyl, wherein the C_3 - C_{10} cycloalkyl and 4- to 6-membered heterocyclyl are substituted with 0-2 occurrences of R^{3b} ;

[0435] each R^{3b} is independently selected from C_1 - C_6 alkoxy, halo, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkoxy, C_1 - C_6 alkyl, $-\text{CN}$, $-\text{SO}_2\text{R}^4$, and hydroxyl.

[0436] Embodiment 27. The compound of any one of Embodiments 25 and 26, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, of Formula (Id-2) or (Id-3):




[0437] Embodiment 28. The compound of any one of Embodiments 1, 9, 13, 16 and 22, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, of Formula (Ie''), wherein:




[0438] R^{2b} , R^{2c} and R^{2e} are defined according to any of Embodiments 16 to 21.

[0439] Embodiment 29. The compound of any one Embodiments 1, 2, 9, 10, 13, 14, 16, 17, 22, 23 and 28, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, of Formula (Ie'), wherein:



[0440] R^{2b} , R^{2c} and R^{2e} are defined according to any of Embodiments 16 to 21.

[0441] Embodiment 30. The compound of any one of the preceding Embodiments, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, of Formula (Ie), wherein:



[0442] R^{2b} , R^{2c} and R^{2e} are defined according to any of Embodiments 16 to 21.

[0443] Embodiment 31. The compound of any one of the preceding Embodiments, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein X is CH.

[0444] Embodiment 32. The compound of any one of the preceding Embodiments, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein X is N.

[0445] Embodiment 33. The compound of any one of Embodiments 1 to 15, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein n is selected from 0 and 1, and m is selected from 1 and 2.

[0446] Embodiment 34. The compound of any one of Embodiments 1 to 15, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein R^2 is unsubstituted C_1 - C_6 alkyl, e.g., methyl, and n is 1.


[0447] Embodiment 35. The compound of any one of Embodiments 1 to 20 and 33, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein m is 1.

[0448] Embodiment 36. The compound of any one of Embodiments 1 to 12, 16 to 27, and 31 to 35, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein R^3 is C_1 - C_6 alkyl, wherein the C_1 - C_6 alkyl is substituted with 1 occurrence of R^{3a} .

[0449] Embodiment 37. The compound of any one of Embodiments 1 to 12, 16 to 27, and 31 to 36, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein R^3 is selected from methyl, ethyl, n-propyl, i-propyl, 2-propanyl, butyl, i-butyl, 2-butanyl, 3-methyl-2-butanyl, i-pentyl, 3-pentanyl, neopentyl, 2,4-dimethylpentanyl, and $-\text{CH}_2-(\text{CH}_2)_{0-1}-R^{3a}$.

[0450] Embodiment 38. The compound of any one of the preceding Embodiments, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein R^{3a} is C_3 - C_{10} cycloalkyl, wherein the C_3 - C_{10} cycloalkyl is substituted with 0-4 occurrences of R^{3b} , wherein each R^{3b} is independently selected from C_1 - C_6 alkoxyl, chloro, fluoro, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkoxyl and C_1 - C_6 alkyl.

[0451] Embodiment 39. The compound of any one of the preceding Embodiments, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein R^{3a} is selected from cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, adamantyl,



[0452] Embodiment 40. The compound of any one of the preceding Embodiments, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein R^{3a} is C_3 - C_7 cycloalkyl, wherein the C_3 - C_7 cycloalkyl is substituted with 0-2 occurrences of fluoro.

[0453] Embodiment 41. The compound of any one of Embodiments 16 to 40, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein each of R^{2b} and R^{2e} is independently selected from hydrogen and unsubstituted C_1 - C_3 alkyl; and R^{2c} is hydrogen.

[0454] Embodiment 42. The compound of any one of Embodiments 16 to 41, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein each of R^{2b} and R^{2e} is independently selected from hydrogen and methyl; and R^{2c} is hydrogen.

[0455] Embodiment 43. The compound of any one of Embodiments 16 to 42, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein R^{2b} is unsubstituted C_1 - C_3 alkyl (e.g., methyl); R^{2c} is hydrogen; and R^{2e} is selected from hydrogen and unsubstituted C_1 - C_3 alkyl.


[0456] Embodiment 44. The compound of any one of Embodiments 16 to 43, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein R^{2b} is methyl and R^{2c} , R^{2d} , R^{2e} and R^{2f} are all hydrogen.

[0457] Embodiment 45. The compound of any one of Embodiments 1 to 15 and 31 to 44, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein R^2 is unsubstituted C_1 - C_3 alkyl, and n is 1.

[0458] Embodiment 46. The compound of any one of Embodiments 1, 2, 4 to 10, 12 to 14, 16, 17, 19 to 23, 28, 29 and 31 to 45, wherein $\text{---}\text{C}=\text{C}\text{---}$ is a double bond.


[0459] Embodiment 47. The compound of any one of the preceding Embodiments $\text{---}\text{C}\text{---}\text{C}\text{---}$ is a single bond.

[0460] Embodiment 48. The compound of Embodiment 1 or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, selected from:




1-(5-(((1s,4s)-4-methoxycyclohexyl)methyl)piperidin-4-yl)methylpyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued



1-(5-((1-(5-fluoro-2-methylbenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((1r,4r)-4-methoxycyclohexyl)methyl)piperidin-4-yl)methylpyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(R)-1-(5-((hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((1-(cyclohexylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



methyl 4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperidine-1-carboxylate




1-(5-((1-(cyclohexylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(acetyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-(((2S,4R)-1-((4,4-difluorocyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-(4-methoxybenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(cyclopentylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




phenyl 4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperidine-1-carboxylate


-continued




1-(5-((1-((tetrahydro-2H-pyran-4-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



1-(5-(((1R,5S)-8-isobutyl-azabicyclo[3.2.1]octan-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(cyclopentylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(thiazol-2-ylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione


-continued



1-(5-((1-((4,4-difluorocyclohexyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((3S,5S)-4-isobutyl-3,5-dimethylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-((tetrahydro-2H-pyran-4-yl)sulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione


-continued




(S)-1-(5-((3-methyl-4-(pyridin-3-ylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-(cyclopropylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((3-ethyl-4-((tetrahydro-2H-pyran-4-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-2-methyl-1-((tetrahydro-2H-pyran-4-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued



1-(5-((1-(2-oxo-2-(piperidin-1-yl)ethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(2-cyclohexylethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(isobutyryl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

-continued




1-(5-(((2S,4R)-1-(cyclohexylmethyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((1-(2-chlorobenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-fluoro-1-isobutylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-((cyclopropylmethyl)sulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-((cyclohexylmethyl)sulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-(2-methylbenzyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-(((2S,4R)-1-(cyclopentylmethyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-(((1R,5S)-8-(pyridin-3-ylmethyl)-8-azabicyclo[3.2.1]octan-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-((tetrahydro-2H-pyran-3-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-(4-fluorobenzyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued



1-(5-((4-(cyclohexylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(pyridin-2-ylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((4-((4,4-difluorocyclohexyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued



1-(5-((1-(2-methoxybenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(2-(tetrahydro-2H-pyran-4-yl)propan-2-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(1-cyclobutylmethyl)-4-fluoropiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued



1-(5-(((S)-4-(((1R,4S)-4-methoxycyclohexyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((3-ethyl-4-isobutylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((4-(cyclohexylmethyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued



1-(5-((4-(2-fluoro-2-methylpropyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((S)-4-((1S,4R)-4-methoxycyclohexyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-(3-methylbenzyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((1-(cycloheptylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-(thiazol-4-ylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4S)-1-(cyclohexylmethyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione








(S)-1-(5-((4-(cyclohexylmethyl)-2-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued



-continued




-continued




1-(5-((1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((1-(cyclohexanecarbonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(R)-1-(5-((4-isobutyl-3-(methoxymethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((2S,4R)-1-isobutyl-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((4-isobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((2-methyl-4-((tetrahydro-2H-pyran-4-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

-continued




(S)-1-((5-((4-(cyclopentylmethyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((1-(3-hydroxy-2-(hydroxymethyl)propyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-(pyridin-3-ylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-(2-cyclobutylethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(heptan-4-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-((1-methyl-1H-pyrazol-5-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

-continued




(S)-1-(5-((1-((tetrahydrofuran-3-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((4-isopropylsulfonyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((3,3-dimethyl-4-((tetrahydro-2H-pyran-4-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-(2-cyclopentylethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((4-isopentyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((4-((3,3-difluorocyclobutyl)methyl)-2-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued



(R)-1-(5-((1-((tetrahydrofuran-3-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((4-(cyclobutylmethyl)-2-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-((4,4-difluorocyclohexyl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((1-(isopropylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-(isopropylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione







1-(5-((4-(3,3,3-trifluoro-2,2-dimethylpropyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione






(S)-1-(5-((4-((3,3-difluorocyclobutyl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione


-continued




-continued




-continued




1-(5-((1-isobutyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((4-(2,2,3,3-tetrafluoropropyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-1-(cyclobutylmethyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(1-trifluoromethyl)cyclopropane-1-carbonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione


-continued



1-(5-((1-(3,3-difluorocyclobutyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




cyclohexyl 4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperidine-1-carboxylate




1-(5-((1-(oxetan-3-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((4-((1-methyl-1H-imidazol-4-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-((tetrahydrofuran-3-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-(isobutyl-3,3-dimethylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-((tetrahydro-2H-pyran-4-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((1-(3-phenylpropanoyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-(cyclobutylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-(thiazol-4-ylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(isopentyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione

-continued




1-(5-((4-(1-(pyridin-3-yl)ethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((4-(((1s,4s)-4-methoxycyclohexyl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((4-(cyclobutylmethyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((4-(pyridin-4-ylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-((1-methyl-1H-imidazol-4-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-((3,3-difluorocyclobutyl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione


-continued



1-(5-((1-(piperidin-4-ylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(((1R,4R)-4-ethoxycyclohexyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((1R,4R)-5-(pyridin-3-ylmethyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((4-(cyclopentylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-(4-methoxypyrimidin-2-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(sec-butylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

-continued




1-(5-((4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-(((2S)-2-methyl-4-((tetrahydrofuran-2-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-benzoylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((4-(2-cyclohexylethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-(1-(trifluoromethyl)cyclopropane-1-carbonyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued



1-(5-((4-((tetrahydro-2H-pyran-3-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(3-cyclohexylpropanoyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-isobutylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-(((1R,5S)-8-(pyridin-3-ylmethyl)-3-8-diazabicyclo[3.2.1]octan-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-(pentan-3-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(R)-1-(5-((3-methyl-4-(pyridin-3-ylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-neopentylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

-continued




1-(5-(((1S,4S)-5-(isopropylsulfonyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-(((3S,5R)-3,5-dimethyl-4-((tetrahydro-2H-pyran-4-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(isobutylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



(R)-1-(5-((3-(difluoromethyl)-4-isobutylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((1R,5S)-8-(3-fluorobenzyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((3-methyl-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione

-continued




1-(5-((4-(2-cyclohexylethyl)-3-oxopiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((1-(1-(pyridin-3-yl)ethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(1-(1-fluorocyclohexyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-(((1S,4S)-5-(pyridin-3-ylmethyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-(cyclohexylmethyl)-3-oxopiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(benzylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

-continued




1-(5-(((1S,4S)-5-(cyclohexylmethyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((1-(2-methyl-1-(tetrahydro-2H-pyran-4-yl)propan-2-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-(cyclobutylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



(R)-1-(5-((4-(cyclohexylmethyl)-3-(difluoromethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((4-isobutyl-3-isopropylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(3-fluorobenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

-continued




1-(5-(((1R,4R)-5-(isopropylsulfonyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((1-(cyclopropylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(4-fluorobenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



(S)-1-(5-((3-isopropyl-4-((tetrahydro-2H-pyran-4-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-isobutyl-2-oxopiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(methylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((4-(cyclohexylmethyl)-2-oxopiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-ethylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(R)-1-(5-((3-methyl-4-((tetrahydro-2H-pyran-4-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-((tetrahydrofuran-3-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-(((1-((3r,5r,7r)-adamantan-1-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-(3-(trifluoromethyl)benzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((4-(cycloheptylmethyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(1-phenylethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued



1-(5-((1-(2-cyclobutylpropan-2-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(3-(trifluoromethoxy)benzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((3S)-4-(((2R,6S)-2,6-dimethyltetrahydro-2H-pyran-4-yl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((4-propylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-(((2S,4R)-1-(cycloheptylmethyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-phenethylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(R)-1-(5-((4-isobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued



(R)-1-((5-((2-methyl-4-((tetrahydro-2H-pyran-4-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-((5-(((3S)-3-methyl-4-((tetrahydrofuran-2-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-((5-((1-(pyridin-4-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued



1-((5-(((4-(((2R,6S)-2,6-dimethyltetrahydro-2H-pyran-4-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-((5-(((3S,5R)-4-isobutyl-3,5-dimethylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-((5-((1-(1-isobutylpiperidin-4-yl)ethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued



1-(5-((1-benzylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-benzyl-4-fluoropiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(2,2,3,3-tetrafluoropropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-(((2S,4R)-1-((3,3-difluorocyclobutyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-(phenylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-(((3r,5r,7r)-adamantan-1-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-((5-fluoropyridin-3-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued



1-(5-(((2R,4R)-2-methyl-1-((tetrahydro-2H-pyran-4-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(2-methoxyethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(((2R,6S)-2,6-dimethyltetrahydro-2H-pyran-4-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((1-(3,3,3-trifluoro-2,2-dimethylpropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-(2,4-dimethylpentan-2-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(2-hydroxyethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2R,4S)-1-isobutyl-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((1-((5-methylpyridin-3-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-(((2S,4S)-1-isobutyl-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((1R,5S)-8-(cyclohexymethyl)-8-azabicyclo[3.2.1]octan-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((3S)-3-methyl-4-((tetrahydrofuran-2-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued



1-(5-((1-isopropylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-(cyclohexylmethyl)-3,3-dimethylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(2-fluorobenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-(((2S,4S)-2-methyl-1-((tetrahydro-2H-pyran-4-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-(2-fluoro-2-methylpropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2R,4S)-2-methyl-1-((tetrahydro-2H-pyran-4-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-((tetrahydrofuran-2-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-


-continued




(R)-1-(5-((hexahydropyrazino[2,1-c][1,4]oxazin-8(1H)-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-isopropylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-(cyclohexylmethyl)-3,3-dimethylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(2-fluorobenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-(((2S,4S)-2-methyl-1-((tetrahydro-2H-pyran-4-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-(2-fluoro-2-methylpropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2R,4S)-2-methyl-1-((tetrahydro-2H-pyran-4-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-((tetrahydrofuran-2-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione


-continued




(R)-1-(5-((hexahydropyrazino[2,1-c][1,4]oxazin-8(1H-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((4-(cyclopropylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2R,4R)-1-(cyclohexylmethyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




(R)-1-(5-((4-(cyclohexylmethyl)-2-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione


-continued




(S)-1-(5-((hexahydropyrazino[2,1-c][1,4]oxazin-8(1H)-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-(3-methoxybenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2R,4R)-1-isobutyl-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(2-methylbenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione


-continued



1-(5-(((S)-4-(((1R,4S)-4-hydroxycyclohexyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione







1-(5-((1-(3-fluoro-2-methylbenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione







1-(5-((1-(2-(4,4-difluorocyclohexyl)propan-2-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione


-continued



-continued




-continued




1-(5-((1-((6-methylpyridin-3-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-(((2S,4R)-2-methyl-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-(((4-((4,4-dimethylcyclohexyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



1-(5-(((1-(2-methoxyethyl)sulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(4-methylbenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2R,4R)-1-((4,4-difluorocyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-(((3S,5S)-4-(cyclohexylmethyl)-3,5-dimethylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((4-(2-(4,4-difluorocyclohexyl)propan-2-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(cyclohexylmethyl)piperidin-4-yl)methyl)-4-methylpyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-1-((4,4-dimethylcyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued



1-(5-((4-benzylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-2-methyl-1-(pyridin-3-ylmethyl)piperidin-4-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(cyclohexylmethyl)-4-fluoropiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-(((1-(1r,4r)-4-hydroxycyclohexyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



(S)-1-(5-((hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-1-(((1r,4S)-4-methoxycyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(2-methylallyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((1-(1,4-diazepan-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-(2-hydroxy-2-methylpropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-isobutyl-1,4-diazepan-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((octahydro-2H-pyrido[1,2-a]pyrazin-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((4-(cyclohexylmethyl)-1,4-diazepan-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-(3-fluorophenyl)ethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(azetidin-3-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-((1-methyl-1H-pyrazol-5-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((1-isobutylazetidin-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((4-(2-(tetrahydro-2H-pyran-4-yl)propan-2-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(cyclohexylmethyl)azetidin-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

-continued




1-(5-((1-(cyclohexylmethyl)pyrrolidin-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione


-continued




(R)-1-(5-((4-isobutyl-2-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(2,4-difluorobenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-(cyclobutylmethyl)pyrrolidin-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(pyridin-3-ylmethyl)pyrrolidin-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(3-fluoro-5-(trifluoromethyl)benzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione


-continued



1-(5-((1-(3-fluorobenzyl)pyrrolidin-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(2-(1H-imidazol-4-yl)ethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-(isobutyl)pyrrolidin-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((1-(2-cyclohexyl-2,2-difluoroethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



1-(5-(((2S,4R)-1-(((3R,4S)-4-difluorocyclopentyl)methyl)2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-isopropylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




(R)-1-(5-((octahydro-2H-pyrido[1,2-a]pyrazin-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((1-isobutyl-2,2-dimethylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-(((1r,4r)-4-hydroxycyclohexyl)methyl)piperazin-1-yl)methylpyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-diones




1-(5-((1-(2,3-difluorobenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-1-(isopropylsulfonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




(R)-1-(5-((4-(cyclohexylmethyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-(((2S,4R)-2-methyl-1-((tetrahydro-2H-pyran-4-yl)sulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((4-(2-hydroxy-2-methylpropyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-2-methyl-1-(2,2,3,3-tetrafluoropropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((1-((6-(trifluoromethyl)pyridin-3-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-(((2S,4R)-1-(((1S,3R)-3-methoxycyclobutyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4-(1H,3H)-dione




1-(5-((1-methyloxetan-3-yl)methyl)piperidin-4-yl)methylpyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((S)-4-(((1S,3R)-3-methoxycyclobutyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((1-((1-(trifluoromethyl)cyclopropyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-(((1S,3S)-3-methoxycyclobutyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione







1-(5-((1-(2-methoxy-2-methylpropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione







1-(5-(((2S,4R)-1-(((1R,3S)-3-methoxycyclobutyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




-continued




-continued




1-(5-(((2S,4R)-2-methyl-1-((tetrahydro-2H-pyran-4-yl)sulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-2-methyl-1-(2,2,3,3-tetrafluoropropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-1-(((1R,3S)-3-methoxycyclobutyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-1-(((1S,3R)-3-methoxycyclobutyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((S)-4-(((1R,3S)-3-methoxycyclobutyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4-(1H,3H)-dione



1-(5-(((S)-4-(((1R,3R)-3-methoxycyclobutyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4-(1H,3H)-dione




1-(5-(((2S,4R)-1-(((R)-3,3-difluorocyclopentyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4-(1H,3H)-dione




1-(5-(((2S,4R)-1-(((S)-3,3-difluorocyclopentyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4-(1H,3H)-dione


-continued




1-(5-(((S)-4-(((R)-3,3-difluorocyclopentyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4-(1H,3H)-dione



1-(5-(((S)-4-(((S)-3,3-difluorocyclopentyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4-(1H,3H)-dione




1-(5-(((2S,4R)-1-(((1R,3R,4S)-3,4-difluorocyclopentyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4-(1H,3H)-dione




1-(5-(((S)-4-(((1R,3R,4S)-3,4-difluorocyclopentyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4-(1H,3H)-dione


-continued




1-(5-(((2S,4R)-1-((4,4-difluorocyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4-(1H,3H)-dione



1-(5-(azepan-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4-(1H,3H)-dione




tert-butyl 4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)azepane-1-carboxylate




1-(5-((1-methylazepan-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4-(1H,3H)-dione

-continued




1-(5-((1-cyclohexylmethyl)azepan-4-yl)methyl)pyrazolo
[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4
(1H,3H)-dione


-continued




1-(5-(((2S,4R)-2-methyl-1-(3,3,3-trifluoropropyl)
piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)
dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-2-methyl-1-(2,2,3,3-tetrafluoropropyl)
piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)
dihydropyrimidine-2,4-(1H,3H)-dione




1-(5-(((2S,4R)-2-methyl-1-(oxetan-2-ylmethyl)piperidin-
4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-
2,4(1H,3H)-dione




1-(5-(((2S,4R)-1-(2,2-difluoroethyl)-2-methylpiperidin-4-yl)
methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-
2,4(1H,3H)-dione



1-(5-(((2S,4R)-1-(2,2-difluoro-3-methoxypropyl)-2-
methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-
yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-2-methyl-1-(2,2,2-trifluoroethyl)
piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-
yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-2-methyl-1-(oxetan-3-yl)piperidin-4-yl)
methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-
2,4(1H,3H)-dione


-continued




1-(5-(((2S,4R)-1-cyclobutyl-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-(oxetan-3-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-((1-cyclobutylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-1-(ethylsulfonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-(((2S,4R)-2-methyl-1-(methylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



1-(5-(((2S,4R)-1-(isopropylsulfonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-1-(cyclopropylsulfonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-1-(cyclopropanecarbonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione


-continued



1-(5-(((2S,4R)-1-isobutyryl-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-1-(cyclobutanecarbonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-2-methyl-1-(1-methylpiperidine-4-carbonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-(((2S,4R)-2-methyl-1-(pyrrolidin-1-ylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



(2S,4R)-4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-N,N,2-trimethylpiperidine-2-sulfonamide




(2S,4R)-N-cyclopentyl-4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperidine-1-sulfonamide




(2S,4R)-4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-N,N,2-trimethylpiperidine-1-carboxamide

-continued




1-(5-(((2S,4R)-2-methyl-1-(pyrrolidin-1-yl)carbamoyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione


-continued




(2S,4R)-4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-N-ethyl-N,2-dimethylpiperidine-1-carboxamide




(2S,4R)-N-cyclopentyl-4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperidine-1-carboxamide




1-(5-(((1s,3s)-3-methoxycyclobutyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((1-(pyrrolidin-1-yl)sulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



1-(5-(((1r,3R,4S)-3,4-difluorocyclopentyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




N-cyclopentyl-4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)piperidine-1-sulfonamide




1-(5-(((2S,4R)-1-(((1r,3S)-3-methoxycyclobutyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione

-continued




1-(5-(((2S,4R)-1-(((1S,3R)-3-methoxycyclobutyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-(((2S,4R)-1-(((S)-3,3-difluorocyclopentyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-1-(((1R,3R,4S)-3,4-difluorocyclopentyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4S)-1-(((4,4-difluorocyclohexyl)methyl)-4-fluoro-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-2-methyl-1-(((1R,4S)-4-(trifluoromethyl)cyclohexyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-(((2S,4R)-1-(((4,4-difluorocyclohexyl)methyl)-4-fluoro-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-1-(((R)-3,3-difluorocyclopentyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione










1-(5-(((2S,4S)-4-fluoro-2-methyl-1-(oxetan-3-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




-continued




-continued




1-(5-(((2S,4R)-4-fluoro-2-methyl-1-(oxetan-3-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-((1-isobutyl-2-(trifluoromethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(R)-1-(5-((octahydro-2H-pyrido[1,2-a]pyrazin-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(R)-1-(5-((4-oxohexahydropyrazino[2,1-c][1,4]oxazin-8(1H)-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




(S)-1-(5-(((1,1-dioxidohexahydro-5H-isothiazolo[2,3-a]pyrazin-5-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-(((S)-4-(((1R,3S)-3-methoxycyclobutyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((S)-4-(((1S,3R)-3-methoxycyclobutyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




1-(5-(((S)-4-(((1R,3R,4S)-3,4-difluorocyclopentyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-(((S)-4-(((R)-3,3-difluorocyclopentyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-(((S)-4-(((S)-3,3-difluorocyclopentyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-(((4-(cyclopropylmethyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-(((3-methyl-4-((1-methylcyclobutyl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-(((S)-3-methyl-4-(((1R,4S)-4-(trifluoromethyl)cyclohexyl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-(((3S)-3-methyl-4-(oxetan-2-ylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-(((4-((2-oxaspiro[3.3]heptan-6-yl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-(((4-((6,6-difluorospiro[3.3]heptan-2-yl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




(S)-1-(5-((4-(2,2-difluoroethyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



(S)-1-(5-((4-(2,2-difluoro-3-methoxypropyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((4-cyclobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((3-methyl-4-(tetrahydro-2H-pyran-4-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




(S)-1-(5-((3-methyl-4-(oxetan-3-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



(S)-1-(5-((3-methyl-4-(2-oxaspiro[3.3]heptan-6-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((4-cyclohexyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((4-(4,4-difluorocyclohexyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued



(S)-1-(5-((3-methyl-4-(spiro[3.3]heptan-2-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((3-methyl-4-(2-methyl-2-azaspiro[3.3]heptan-6-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((3-methyl-4-(2-(methylsulfonyl)-2-azaspiro[3.3]heptan-6-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-((4-cyclohexylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



(S)-1-(5-((4-(ethylsulfonyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((4-(cyclopropanecarbonyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((4-isobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued



(S)-1-(5-((4-(cyclohexanecarbonyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((4-(cyclopentanecarbonyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((4-(cyclobutanecarbonyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


-continued




1-(5-(1-(4-isobutylpiperazin-1-yl)ethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



1-(5-(((2S,4R)-1-isobutyl-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-1-isobutyl-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-5-methylpyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-1-(cyclopropylmethyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-5-methylpyrimidine-2,4(1H,3H)-dione


-continued




5-fluoro-1-(5-(((2S,4R)-1-isobutyl-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione



1-(5-(((2S,4R)-1-((4,4-difluorocyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-5-fluoropyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-1-((4,4-difluorocyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-5-methylpyrimidine-2,4(1H,3H)-dione




5-chloro-1-(5-(((2S,4R)-1-((4,4-difluorocyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione


-continued




1-(5-(((2S,4R)-1-((4,4-difluorocyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-5-methoxypyrimidine-2,4(1H,3H)-dione



5-cyclopropyl-1-(5-(((2S,4R)-1-isobutyl-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-1-(cyclopropanecarbonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione




1-(5-(((2S,4R)-1-(ethylsulfonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione


-continued




(S)-1-(5-((4-isobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione



(S)-5-fluoro-1-(5-((4-isobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((4-isobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-5-methylpyrimidine-2,4(1H,3H)-dione




(S)-1-(5-((4-isobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-5-methoxypyrimidine-2,4(1H,3H)-dione


-continued



(S)-5-cyclopropyl-1-(5-((4-isobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione



(S)-1-(5-((4-(cyclopropylmethyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-5-methylpyrimidine-2,4(1H,3H)-dione



(S)-1-(5-((4-(4,4-difluorocyclohexyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione

[0461] Embodiment 49. The compound of any one of the preceding Embodiments, or a pharmaceutically acceptable salt, thereof, wherein the pharmaceutically acceptable salt is an acid addition salt.

[0462] Embodiment 50. A pharmaceutical composition comprising a therapeutically effective amount of a compound of any of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, and a pharmaceutically acceptable carrier or excipient.

[0463] Embodiment 51. A compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use as a medicament.

[0464] Embodiment 52. A method of treating or preventing a disease or disorder in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, thereof.

[0465] Embodiment 53. A method of treating or preventing a disorder that is affected by the reduction of WIZ protein levels, in a subject in need thereof, the method comprising administering to the subject a therapeutically

effective amount of a compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.

[0466] Embodiment 54. A method of treating a disease or disorder that is affected by the modulation of WIZ protein levels comprising administering to the patient in need thereof a compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.

[0467] Embodiment 55. A method of inhibiting WIZ protein expression in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.

[0468] Embodiment 56. A method of degrading WIZ protein in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, thereof.

[0469] Embodiment 57. A method of inhibiting, reducing, or eliminating the activity of WIZ protein or WIZ protein expression, the method comprising administering to the subject a compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.

[0470] Embodiment 58. A method of inducing or promoting fetal hemoglobin in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.

[0471] Embodiment 59. A method of reactivating fetal hemoglobin production or expression in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.

[0472] Embodiment 60. A method of increasing fetal hemoglobin expression in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.

[0473] Embodiment 61. A method of treating a hemoglobinopathy, e.g., a beta-hemoglobinopathy, in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.

[0474] Embodiment 62. A method of treating a sickle cell disease in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.

[0475] Embodiment 63. A method of treating beta-thalassemia in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.

[0476] Embodiment 64. A method for reducing WIZ protein levels in a subject comprising the step of administering to a subject in need thereof a therapeutically effective amount of a compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.

[0477] Embodiment 65. A compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in treating or preventing a disease or disorder in a subject in need thereof.

[0478] Embodiment 66. A compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in the treatment of a disease or disorder selected from sickle cell disease and beta-thalassemia.

[0479] Embodiment 67. A compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in treating or preventing a disorder that is affected by the inhibition of WIZ protein levels, in a subject in need thereof.

[0480] Embodiment 68. A compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in treating or preventing a disorder that is affected by the reduction of WIZ protein levels, in a subject in need thereof.

[0481] Embodiment 69. A compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in the treatment or prevention of a disease or disorder that is affected by the degradation of WIZ protein.

[0482] Embodiment 70. A compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in inhibiting, reducing, or eliminating the activity of WIZ protein or WIZ protein expression in a subject in need thereof.

[0483] Embodiment 71. A compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in inducing or promoting fetal hemoglobin in a subject in need thereof.

[0484] Embodiment 72. A compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in reactivating fetal hemoglobin production or expression in a subject in need thereof.

[0485] Embodiment 73. A compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in increasing fetal hemoglobin expression in a subject in need thereof.

[0486] Embodiment 74. A compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in treating a hemoglobinopathy in a subject in need thereof.

[0487] Embodiment 75. A compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in treating a sickle cell disease in a subject in need thereof.

[0488] Embodiment 76. A compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in treating beta-thalassemia in a subject in need thereof.

[0489] Embodiment 77. A compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in the treatment of a disease or disorder affected by an increase in fetal hemoglobin expression.

[0490] Embodiment 78. A compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in the treatment of a disease or disorder affected by the inhibition, reduction, or elimination of the activity of WIZ protein or WIZ protein expression.

[0491] Embodiment 79. A compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in the treatment of a disease or disorder affected by the induction or promotion of fetal hemoglobin.

[0492] Embodiment 80. A compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in the treatment of a disease or disorder affected by the reactivation of fetal hemoglobin production or expression.

[0493] Embodiment 81. A compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in inhibiting WIZ protein expression in a subject in need thereof.

[0494] Embodiment 82. A compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in degrading WIZ protein in a subject in need thereof.

[0495] Embodiment 83. Use of a compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, in the manufacture of a medicament for treating a disease or disorder that is affected by the reduction of WIZ protein levels, inhibition of WIZ protein expression or degradation of WIZ protein.

[0496] Embodiment 84. Use of a compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, in the manufacture of a medicament for treating a disease or disorder that is affected by inducing or promoting fetal hemoglobin.

[0497] Embodiment 85. Use of a compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, in the manufacture of a medicament for treating a disease or disorder that is affected by reactivating fetal hemoglobin production or expression.

[0498] Embodiment 86. Use of a compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, in the manufacture of a medicament for treating a disease or disorder that is affected by increasing fetal hemoglobin expression.

[0499] Embodiment 87. The use of a compound of any one of Embodiments 83 to 86, wherein the disease or disorder is selected from sickle cell disease and beta-thalassemia.

[0500] Embodiment 88. Use of a compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, in the treatment of a disease or disorder that is affected by the reduction of WIZ protein levels, inhibition of WIZ protein expression or degradation of WIZ protein.

[0501] Embodiment 89. Use of a compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, in the treatment of a disease or disorder that is affected by inducing fetal hemoglobin, reactivating fetal hemoglobin production or expression, or increasing fetal hemoglobin expression.

[0502] Embodiment 90. The use of Embodiment 88 or 89, wherein the disease or disorder is selected from sickle cell disease and beta-thalassemia.

[0503] Embodiment 91. A pharmaceutical combination comprising a compound of any one of Embodiments 1 to 49, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, and one or more additional therapeutic agent(s).

[0504] Depending on the choice of the starting materials and procedures, the compounds can be present in the form of one of the possible isomers or as mixtures thereof, for example as pure optical isomers, or as isomer mixtures, such as racemates and diastereomeric mixtures, depending on the number of asymmetric centres. The disclosure is meant to include all such possible isomers, including racemic mixtures, enantiomerically enriched mixtures, diastereomeric mixtures and optically pure forms. Optically active (R)- and (S)-isomers may be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques. If the compound contains a disubstituted or trisubstituted cycloalkyl, the cycloalkyl substituent(s) may have a cis- or trans-configuration. The disclosure includes cis and trans configurations of substituted cycloalkyl groups as well as mixtures thereof. All tautomeric forms are also intended to be included. In particular, where a heteroaryl ring containing N as a ring atom is 2-pyridone, for example, tautomers where the carbonyl is depicted as a hydroxy (e.g., 2-hydroxypyridine) are included.

Pharmaceutically Acceptable Salts

[0505] As used herein, the terms “salt” or “salts” refers to an acid addition or base addition salt of a compound of the disclosure. “Salts” include in particular “pharmaceutically acceptable salts”. The term “pharmaceutically acceptable salts” refers to salts that retain the biological effectiveness and properties of the compounds of this disclosure and, which typically are not biologically or otherwise undesirable. The compounds of the disclosure may be capable of forming acid and/or base salts by virtue of the presence of amino and/or carboxyl groups or groups similar thereto.

[0506] Pharmaceutically acceptable acid addition salts can be formed with inorganic acids and organic acids. Inorganic acids from which salts can be derived include, for example, hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like. Organic acids from which salts can be derived include, for example, acetic acid, propionic acid, glycolic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, toluenesulfonic acid, sulfosalicylic acid, formic acid, trifluoroacetic acid, and the like.

[0507] Pharmaceutically acceptable base addition salts can be formed with inorganic and organic bases. Inorganic bases from which salts can be derived include, for example, ammonium salts and metals from columns I to XII of the periodic table. In certain embodiments, the salts are derived from sodium, potassium, ammonium, calcium, magnesium, iron, silver, zinc, and copper; particularly suitable salts include ammonium, potassium, sodium, calcium and magnesium salts.

[0508] Organic bases from which salts can be derived include, for example, primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines, basic ion exchange resins, and the like. Certain organic amines include isopropylamine, benzathine, choline, diethanolamine, diethylamine, lysine, meglumine, piperazine and tromethamine.

[0509] In another aspect, the disclosure provides compounds in acetate, ascorbate, adipate, aspartate, benzoate, besylate, bromide/hydrobromide, bicarbonate/carbonate, bisulfate/sulfate, camphorsulfonate, caprate, chloride/hydrochloride, chlorthalophyllonate, citrate, ethandisulfonate, formate, fumarate, gluceptate, gluconate, glucuronate, glutamate, glutarate, glycolate, hippurate, hydroiodide/iodide, isethionate, lactate, lactobionate, laurylsulfate, malate, maleate, malonate, mandelate, mesylate, methylsulphate, mucate, naphthoate, napsylate, nicotinate, nitrate, octadecanoate, oleate, oxalate, palmitate, pamoate, phosphate/hydrogen phosphate/dihydrogen phosphate, polygalacturonate, propionate, sebacate, stearate, succinate, sulfosalicylate, sulfate, tartrate, tosylate trifrenate, trifluoroacetate or xinafoate salt form.

[0510] In another aspect, the disclosure provides compounds in sodium, potassium, ammonium, calcium, magnesium, iron, silver, zinc, copper, isopropylamine, benzathine, choline, diethanolamine, diethylamine, lysine, meglumine, piperazine or tromethamine salt form.


[0511] Preferably, pharmaceutically acceptable salts of compounds of formulae (I), (Ia), (Ib), (Ic), (Id), and (Ie), are acid addition salts.

Isotopically Labeled Compounds

[0512] Any formula given herein is also intended to represent unlabeled forms as well as isotopically labeled forms of the compounds. Isotopically labeled compounds have structures depicted by the formulas given herein except that one or more atoms are replaced by an atom having a selected atomic mass or mass number. Examples of isotopes that can be incorporated into compounds of the disclosure include isotopes of hydrogen, carbon, nitrogen, oxygen, sulfur, fluorine, chlorine and iodine, such as ^2H , ^3H , ^{11}C , ^{13}C , ^{14}C , ^{18}O , ^{15}N , ^{18}F , ^{17}O , ^{18}O , ^{35}S , ^{36}Cl , ^{123}I , ^{124}I , ^{125}I respectively. The disclosure includes various isotopically labeled compounds as defined herein, for example those into which radioactive isotopes, such as ^3H and ^{14}C , or those into which non-radioactive isotopes, such as ^2H and ^{13}C are present. Such isotopically labeled compounds are useful in metabolic studies (with ^{14}C), reaction kinetic studies (with, for example ^2H or ^3H), detection or imaging techniques, such as positron emission tomography (PET) or single-photon emission computed tomography (SPECT) including drug or substrate tissue distribution assays, or in radioactive treatment of patients. In particular, an ^{18}F compound may be particularly desirable for PET or SPECT studies. Isotopically-labeled compounds of formulae (I''), (I'), (I), (Ia''),

(Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), and (Ie), can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described in the accompanying Examples and General Schemes (e.g., General Schemes 1 to 5) using an appropriate isotopically-labeled reagent in place of the non-labeled reagent previously employed.

[0513] In one embodiment of any aspect of the present disclosure, the hydrogens in the compound of Formula (I), Formula (I') or Formula (I'') (and subformulae thereof) are present in their normal isotopic abundances. In another embodiment, the hydrogens are isotopically enriched in deuterium (D), and in a particular embodiment of the disclosure the hydrogen(s) of the dihydrouracil (DHU) or the uracil portion in compounds of Formula (I) or Formula (I') are enriched in D, for example,



Deuterated dihydrouracil and uracil moieties can be prepared as described in Hill, R. K. et al., *Journal of Labelled Compounds and Radiopharmaceuticals*, Vol. XXII, No. 2, p. 143-148.

[0514] Further, substitution with heavier isotopes, particularly deuterium (i.e., ^2H or D) may afford certain therapeutic advantages resulting from greater metabolic stability, for example increased in vivo half-life or reduced dosage requirements or an improvement in therapeutic index. It is understood that deuterium in this context is regarded as a substituent of a compound of the formulae (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), and (Ie). The concentration of such a heavier isotope, specifically deuterium, may be defined by the isotopic enrichment factor. The term "isotopic enrichment factor" as used herein means the ratio between the isotopic abundance and the natural abundance of a specified isotope. If a substituent in a compound of this

disclosure is denoted deuterium, such compound has an isotopic enrichment factor for each designated deuterium atom of at least 3500 (52.5% deuterium incorporation at each designated deuterium atom), at least 4000 (60% deuterium incorporation), at least 4500 (67.5% deuterium incorporation), at least 5000 (75% deuterium incorporation), at least 5500 (82.5% deuterium incorporation), at least 6000 (90% deuterium incorporation), at least 6333.3 (95% deuterium incorporation), at least 6466.7 (97% deuterium incorporation), at least 6600 (99% deuterium incorporation), or at least 6633.3 (99.5% deuterium incorporation).

[0515] Pharmaceutically acceptable solvates in accordance with the disclosure include those wherein the solvent of crystallization may be isotopically substituted, e.g., D₂O, d₆-acetone, d₆-DMSO.

[0516] Compounds of the disclosure, i.e. compounds of formulae (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), (Ie), and (Ie), that contain groups capable of acting as donors and/or acceptors for hydrogen bonds may be capable of forming co-crystals with suitable co-crystal formers. These co-crystals may be prepared from compounds of (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), and (Ie) by known co-crystal forming procedures. Such procedures include grinding, heating, co-subliming, co-melting, or contacting in solution compounds of formulae (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), and (Ie) with the co-crystal former under crystallization conditions and isolating co-crystals thereby formed. Suitable co-crystal formers include those described in WO 2004/078163.

[0517] All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (e.g., "such as") provided herein is intended merely to better illuminate the disclosure and does not pose a limitation on the scope of the disclosure otherwise claimed.

[0518] Any asymmetric center (e.g., carbon or the like) of the compound(s) of the disclosure can be present in racemic or enantiomerically enriched, for example the (R)-, (S)- or (R,S)-configuration. In certain embodiments, for example, as a mixture of enantiomers, each asymmetric center is present in at least 10% enantiomeric excess, at least 20% enantiomeric excess, at least 30% enantiomeric excess, at least 40% enantiomeric excess, at least 50% enantiomeric excess, at least 60% enantiomeric excess, at least 70% enantiomeric excess, at least 80% enantiomeric excess, at least 90% enantiomeric excess, at least 95% enantiomeric excess, or at least 99% enantiomeric excess. In certain embodiments, for example, in enantiomerically enriched form, each asymmetric center is present in at least 50% enantiomeric excess, at least 60% enantiomeric excess, at least 70% enantiomeric excess, at least 80% enantiomeric excess, at least 90% enantiomeric excess, at least 95% enantiomeric excess, or at least 99% enantiomeric excess. Thus, compounds of the disclosure can be present in a racemic mixture or in enantiomerically enriched form or in an antipure form or as a mixture of diastereoisomers.

[0519] In the formulae of the present application the term " on a C-sp³ indicates the absolute stereochemistry, either (R) or (S). In the formulae of the present application the term " on a C-sp³ indicates the absolute stereochem-

istry, either (R) or (S). In the formulae of the present application the term " on a C-sp³ represents a covalent bond wherein the stereochemistry of the bond is not defined. This means that the term " on a C-sp³ comprises an (S) configuration or an (R) configuration of the respective chiral centre. Furthermore, mixtures may also be present. Therefore, mixtures of stereoisomers, e.g., mixtures of enantiomers, such as racemates, and/or mixtures of diastereoisomers are encompassed by the present disclosure.

[0520] For the avoidance of doubt, where compound structures are drawn with undefined stereochemistry with respect to any R group, for example, to R² in formula (I), as represented by a bond () , this means the asymmetric center has either a (R)- or (S)-configuration, or exists as a mixture thereof and stated as such.

[0521] Accordingly, as used herein a compound of the disclosure can be in the form of one of the possible stereoisomers, rotamers, atropisomers, tautomers or mixtures thereof, for example, as substantially pure geometric (cis or trans) stereoisomers, diastereomers, optical isomers, racemates or mixtures thereof.

[0522] Any resulting mixtures of stereoisomers can be separated on the basis of the physicochemical differences of the constituents, into the pure or substantially pure geometric or optical isomers, diastereomers, racemates, for example, by chromatography and/or fractional crystallization.

[0523] Any resulting racemates of compounds of the disclosure or of intermediates can be resolved into the optical isomers (enantiomers) by known methods, e.g., by separation of the diastereomeric salts thereof, obtained with an optically active acid or base, and liberating the optically active acidic or basic compound. In particular, a basic moiety may thus be employed to resolve the compounds of the disclosure into their optical antipodes, e.g., by fractional crystallization of a salt formed with an optically active acid, e.g., tartaric acid, dibenzoyl tartaric acid, diacetyl tartaric acid, di-O,O'-p-toluoyl tartaric acid, mandelic acid, malic acid or camphor-10-sulfonic acid. Racemic compounds of the disclosure or racemic intermediates can also be resolved by chiral chromatography, e.g., high pressure liquid chromatography (HPLC) using a chiral adsorbent.



[0524] Furthermore, the compounds of the disclosure, including their salts, can also be obtained in the form of their hydrates, or include other solvents used for their crystallization. The compounds of the disclosure may inherently or by design form solvates with pharmaceutically acceptable solvents (including water); therefore, it is intended that the disclosure embrace both solvated and unsolvated forms. The term "solvate" refers to a molecular complex of a compound of the disclosure (including pharmaceutically acceptable salts thereof) with one or more solvent molecules. Such solvent molecules are those commonly used in the pharmaceutical art, which are known to be innocuous to the recipient, e.g., water, ethanol, and the like. The term "hydrate" refers to the complex where the solvent molecule is water. The presence of solvates can be identified by a person of skill in the art with tools such as NMR.

[0525] The compounds of the disclosure, including salts, hydrates and solvates thereof, may inherently or by design form polymorphs.

Methods of Making

[0526] The compounds of the disclosure can be prepared in a number of ways well known to those skilled in the art of organic synthesis. By way of example, compounds of the present disclosure can be synthesized using the methods described below, together with synthetic methods known in the art of synthetic organic chemistry, or variations thereon as appreciated by those skilled in the art.

[0527] Generally, the compounds of formula (I''), formula (I) and formula (I') can be prepared according to the Schemes provided infra.





The starting materials for the above reaction scheme are commercially available or can be prepared according to methods known to one skilled in the art or by methods disclosed herein. In general, the compounds of the disclosure are prepared in the above reaction Scheme 1 as follows:

[0528] A cross-coupling reaction, such as a palladium (Pd)-catalysed coupling of I-1 with a boranyle coupling partner of formula I-2A (prepared by hydroboration of an appropriate alkene with 9-BBN, for example) in the presence of a polar solvent, such as N,N-dimethylformamide (DMF), a suitable ligand such as dppf, and a base such as potassium carbonate (K_2CO_3) can provide the cross-coupled product I-3 in Step 1, where X is CH. Removal of the protecting group (e.g., Boc) under acidic conditions at room temperature can provide the free amine I-4A, where Z=2, 4-dimethoxybenzyl (DMB). Alternatively, removal of the protecting groups under acidic conditions and heating can provide I-4B (Step 2). I-4A and I-4B can then be converted respectively to I-5A and I-5B via a reductive amination (Step 3-i) with an appropriate aldehyde in the presence of a borohydride reagent, such as sodium borohydride acetate. Alternatively, by an alkylation reaction (Step 3-ii) with an appropriate alkyl halide, mesylate, tosylate or triflate in the presence of an amine or carbonate base and polar solvent, such as diisopropylethylamine (DIPEA) or potassium carbonate (K_2CO_3) and dimethylformamide (DMF). Alternatively, by an amide coupling reaction (Step 3-iii) of the compound with an appropriate carboxylic acid, an activating agent, such as HATU, and a base such as DIPEA, when R^3 forms an amide with the nitrogen to which it is attached. Alternatively, by an acylation or sulfonylation reaction (Step 3-iv) with an appropriate acyl chloride or sulfonyl chloride and a base such as DIPEA or TEA, where R^3 forms an amide or sulfonamide with the nitrogen to which it is attached. Removal of the protecting group of I-5A under acidic conditions and heating can provide I-5B (Step 4).

-continued



General scheme 2



[0529] The starting materials for the above reaction scheme are commercially available or can be prepared according to methods known to one skilled in the art or by methods disclosed herein. In general, the compounds of the disclosure are prepared in the above reaction Scheme 2 as follows:

[0530] A cross-coupling reaction, such as a palladium (Pd)-catalysed coupling of I-1 with a trifluoroborate (potassium salt) coupling partner of formula II-2B in the presence of an organic solvent such as toluene, and water, a phosphine ligand such as RuPhos or Xphos, and a base such as cesium carbonate (Cs_2CO_3) can provide the cross-coupled product I-3 in Step 1, where X is N. Compound I-3 as prepared in this manner, can be converted to compounds of formula I-5B by the methods of General Scheme 1, steps 2-4.



General scheme 3




[0531] The starting materials for the above reaction scheme are commercially available or can be prepared according to methods known to one skilled in the art or by methods disclosed herein. In general, the compounds of the disclosure are prepared in the above reaction Scheme 3 as follows:

[0532] A cross-coupling reaction, such as a nickel (Ni)-catalysed coupling of I-1 with an alkyl bromide coupling partner of formula III-2C in the presence of a polar solvent such as DMA, a salt such as sodium iodide (NaI), zinc (Zn), and a ligand such as pyridine-2,6-bis(carboximidamide) dihydrochloride can provide the cross-coupled product I-3 in Step 1, where X is CH or CF. Compound I-3 as prepared in this manner, can be converted to compounds of formula I-5B by the methods of General Scheme 1, steps 2-4.

General Scheme 4




-continued



[0533] In General Scheme 4, a compound of formula I-5A or I-5B is subjected to oxidation conditions, e.g., MnO_2 , in a suitable solvent, such as toluene (e.g., at room temperature), or in the presence of N,O-bis(trimethylsilyl)trifluoroacetamide, to produce a compound of formula IV-5C (i.e., formula (I') when $\text{Z}=\text{H}$), followed by an optional deprotection step when the Z group represents a nitrogen protecting group, to give a compound of formula IV-5D (i.e., formula (I'')).



-continued



[0534] In General Scheme 5, a compound of formula I-5A undergoes a Claisen condensation followed by selenation/oxidation/elimination sequence to give a compound of formula V-5E. Compound formula V-5E undergoes hydrolysis followed by a copper-catalysed decarboxylation to give a compound of formula IV-5C. Subsequent deprotection, e.g., under acidic conditions and heating, provides a compound of formula IV-5D (i.e., formula (I') or (I'')).


[0535] For Schemes 1 to 5, X, R^2 , R^3 , n, m and p are as defined herein, in particular according to any one of Embodiments 1 to 49.

[0536] In a further embodiment, there is provided a compound of formula I-1, which is




[0537] 1-(5-bromopyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)di-hydropyrimidine-2,4(1H,3H)-dione.

[0538] In a further embodiment, there is provided a compound of formula (X-1) or a salt thereof,



- [0539] wherein:
[0540] === is a single bond or a double bond;
[0541] X is selected from CH, CF, and N;
[0542] Z is selected from hydrogen and 2,4-dimethoxybenzyl (DMB);
[0543] R^x is selected from hydrogen, C₁-C₆alkyl, halo (e.g., F, Cl), C₁-C₆alkoxyl, and C₃-C₈cycloalkyl;
[0544] R^N is selected from hydrogen and a nitrogen protecting group PG (e.g., tert-butyloxycarbonyl (Boc));
[0545] R' is selected from hydrogen and C₁-C₆alkyl;
[0546] R¹ is selected from hydrogen and C₁-C₆alkyl;
[0547] each R² is independently selected from C₁-C₆alkyl, C₁-C₆haloalkyl, halo, and oxo, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{2a}; or 2 R² on non-adjacent carbon atoms together with the non-adjacent carbon atoms to which they are attached form a bridging ring;
[0548] R^{2a} is selected from C₁-C₆alkoxyl and hydroxyl;
[0549] n is 0, 1, 2, 3 or 4;
[0550] m is 0, 1 or 2; and
[0551] p is 0 or 1.
- [0552] In a further embodiment, there is provided a compound of formula (X or a salt thereof,



- [0553]** wherein:
- [0554]** X is selected from CH, CF, and N;
- [0555]** Z is selected from hydrogen and 2,4-dimethoxybenzyl (DMB);
- [0556]** R^N is selected from hydrogen and a nitrogen protecting group PG (e.g., tert-butyloxycarbonyl (Boc))
- [0557]** each R² is independently selected from C₁-C₆alkyl, C₁-C₆haloalkyl, halo, and oxo, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{2a}; or 2 R² on non-adjacent carbon atoms together with the non-adjacent carbon atoms to which they are attached form a bridging ring;
- [0558]** R^{2a} is selected from C₁-C₆alkoxy and hydroxyl;
- [0559]** n is 0, 1, 2, 3 or 4;
- [0560]** m is 0, 1 or 2; and
- [0561]** p is 0 or 1.
- [0562]** In an embodiment of formula (X-1) or (X), PG is an acid labile protecting group.
- [0563]** In an embodiment of formula (X-1) or (X), PG is the Boc protecting group (tert-butyloxycarbonyl).
- [0564]** In a further embodiment of formula (X-1) or (X), there is provided a compound or a salt thereof, selected from:

- [0565] tert-butyl (2S,4R)-4-((3-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperidine-1-carboxylate;
- [0566] 1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione;
- [0567] (S)-5-methyl-1-(5-(((3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione);
- [0568] tert-butyl (S)-2-methyl-4-((3-(5-methyl-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperazine-1-carboxylate;
- [0569] tert-butyl (2S,4R)-2-methyl-4-((3-(5-methyl-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperidine-1-carboxylate;
- [0570] 5-methyl-1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione;
- [0571] 5-fluoro-1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione;
- [0572] tert-butyl (2S,4R)-4-((3-(5-fluoro-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperidine-1-carboxylate;
- [0573] (S)-5-fluoro-1-(5-(((3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione);
- [0574] tert-butyl (S)-4-((3-(5-fluoro-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperazine-1-carboxylate;
- [0575] 5-chloro-1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione;
- [0576] tert-butyl (2S,4R)-4-((3-(5-chloro-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperidine-1-carboxylate;
- [0577] 5-methoxy-1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione;
- [0578] tert-butyl (2S,4R)-4-((3-(5-methoxy-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperidine-1-carboxylate;
- [0579] (S)-5-methoxy-1-(5-(((3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione);
- [0580] tert-butyl (S)-4-((3-(5-methoxy-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperazine-1-carboxylate;
- [0581] (S)-5-cyclopropyl-1-(5-(((3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione);
- [0582] tert-butyl (S)-4-((3-(5-cyclopropyl-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperazine-1-carboxylate;
- [0583] 5-cyclopropyl-1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione;
- [0584] tert-butyl (2S,4R)-4-((3-(5-cyclopropyl-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperidine-1-carboxylate;
- [0585] (S)-1-(5-(((3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione);

- [0586] tert-butyl (S)-4-((3-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperazine-1-carboxylate;
- [0587] tert-butyl (2S,4R)-4-((3-(3-(3,4-dimethylbenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperidine-1-carboxylate;
- [0588] 3-(3,4-dimethylbenzyl)-1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione;
- [0589] 1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione;
- [0590] tert-butyl 4-((3-(3-(3,4-dimethylbenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperazine-1-carboxylate;
- [0591] 3-(3,4-dimethylbenzyl)-1-(5-(piperazin-1-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione;
- [0592] 1-(5-(piperazin-1-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione;
- [0593] tert-butyl 4-((3-(3-(3,4-dimethylbenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperidine-1-carboxylate;
- [0594] 3-(3,4-dimethylbenzyl)-1-(5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione;
- [0595] 1-(5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione;
- [0596] tert-butyl (S)-4-((3-(3-(3,4-dimethylbenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperazine-1-carboxylate;
- [0597] (S)-3-(3,4-dimethylbenzyl)-1-(5-((3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione; and
- [0598] (S)-1-(5-((3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione.
- [0599] In an embodiment of formula (X-1) or (X), the salt is selected from a HCl and TFA salt.
- [0600] In a further aspect, the disclosure provides to a process for the preparation of a compound of formula (I), (I'), (I'') or subformulae thereof, in free form or in pharmaceutically acceptable salt form, comprising the step of:
- [0601] 1) coupling an aryl bromide of formula (I-1) with a boranoyl coupling partner of formula I-2A or II-2B under cross coupling conditions, to give a compound of formula (I-3) as defined herein.
- [0602] The boranoyl coupling partner of step 1 may optionally be prepared by hydroboration of a precursor alkene, e.g., with 9-BBN.
- [0603] In a further aspect, the disclosure provides a process for the preparation of a compound of formula (I), (I'), (I'') or subformulae thereof, in free form or in pharmaceutically acceptable salt form, comprising the step of:
- [0604] 1) coupling an aryl bromide of formula (I-1) with an alkyl bromide of formula (III-2C) under cross coupling conditions, to give a compound of formula (I-3) as defined herein.
- [0605] Cross coupling reaction conditions for any of the aforementioned process steps or hereinafter involve the use of a Pd catalyst in the presence of a phosphine ligand, such as Pd(OAc)₂ and RuPhos or Xphos, and a base such as cesium carbonate (Cs₂CO₃), in the presence of a suitable solvent such as toluene, water, or a mixture thereof.

[0606] Cross coupling reaction conditions (e.g., in the case of an Sp²-Sp³ coupling) may alternatively involve the use of a Ni(II) complex, such as NiCl₂(DME), ligand, such as pyridine-2,6-bis(carboximidamide) dihydrochloride, additive such as NaI, a transmetalling agent such as Zn or Mn, a suitable solvent such as DMA, heating at a temperature of r.t. to 150° C., e.g., 70° C., for example, over a period of 12 hours.

[0607] In an embodiment of either process aspect described above, there is provided the further steps of:

[0608] 2) deprotecting a compound of formula (I)-3 to give a compound of formula (I)-4A or (I)-4B as defined herein;

[0609] 3-a) reacting a compound of formula (I)-4A or (I)-4B under reductive amination conditions to give a compound of formula (I)-5A or (I)-5B as defined herein; or

[0610] 3-b) reacting a compound of formula (I)-4A or (I)-4B under alkylation conditions to give a compound of formula (I)-5A or (I)-5B as defined herein; or

[0611] 3-c) reacting a compound of formula (I)-4A or (I)-4B under amide coupling conditions to give a compound of formula (I)-5A or (I)-5B as defined herein; or

[0612] 3-d) reacting a compound of formula (I)-4A or (I)-4B under acylation or sulfonylation conditions to give a compound of formula (I)-5A or (I)-5B as defined herein; and

[0613] 4) deprotecting the compound of formula (I)-5A to give a compound of formula (I) as described herein.

[0614] Reductive amination conditions for any of the aforementioned process steps or hereinafter involve the use of a corresponding aldehyde, a suitable hydride reagent, such as NaBH(OAc)₃, a suitable solvent, such as DMF, the reaction conducted at room temperature (r.t.).


[0615] Alkylation reaction conditions for any of the aforementioned process steps or hereinafter involve the use of a corresponding alkyl halide, mesylate, tosylate or triflate in the presence of a suitable base, such as DIPEA, or a carbonate base such as K₂CO₃, a polar solvent, such as DMF, the reaction conducted at a suitable temperature, such as r.t. to 100° C., e.g., 80° C., optionally, under microwave.

[0616] Amide coupling reaction conditions for any of the aforementioned process steps or hereinafter involve the use of a corresponding carboxylic acid, an activating agent, such as HATU, a suitable base, such as DIPEA or NMM, a suitable solvent, such as DMF, the reaction conducted at a suitable temperature, such as r.t., for a suitable amount of time, for example 12 hours.


[0617] Acylation or sulfonylation reaction conditions for any of the aforementioned process steps or hereinafter involve the use of a corresponding acyl chloride or sulfonyl chloride and a base such as DIPEA or TEA, in the presence of a suitable solvent such as DCM, the reaction conducted at a suitable temperature, such as r.t.

[0618] In a further embodiment there is provided a process for the preparation of a compound of formula (I') or (I''), comprising the step:


[0619] 1) coupling an aryl bromide of formula



wherein hal is halo, preferably I, with



under cross coupling reaction conditions, to give a compound of formula



as defined herein, wherein --- is a double bond or a single bond, R^1 , R' , X , R^2 , n , m , p , R^x are as defined herein, e.g., according to any one of enumerated Embodiments 1 to 49, PG is a nitrogen protecting group as defined herein, e.g., Boc, and R^N is selected from hydrogen and a nitrogen protecting group PG (e.g., tert-butyloxycarbonyl (Boc)). In an embodiment, R^1 and R' are both hydrogen.

[0620] Cross coupling conditions for the aforementioned process may involve the use of copper as a catalyst, e.g., Ullmann reaction conditions. For example, reaction conditions may employ copper(I) iodide as a catalyst, a ligand, such as N-(2-cyanophenyl)picolinamide, a base, such as K_3PO_4 , a suitable solvent, such as DMSO, heating at a temperature of r.t. to 130°C ., e.g., 70 to 120°C ., e.g., 110°C . The reaction may be heated over a period of 72 hours.

[0621] In a further embodiment there is provided a process for the preparation of a compound of formula (I), (I'), (I'') or subformulae thereof, in free form or in pharmaceutically acceptable salt form according to any of General Schemes 1 to 5.

[0622] Compounds of formulae (I)-1, (X-1) and (X) as defined herein are useful in the preparation of compounds of the disclosure, e.g., compounds of formulae (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), and (Ie). Thus, in an aspect, the disclosure relates to a compound of formula (I)-1, (X-1) or (X) or salts thereof. In another aspect, the disclosure relates to the use of a compound of formula (I)-1, (X-1) or (X) or salts thereof in the manufacture of a compound of

formulae (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), and (Ie). The disclosure further includes any variant of the present processes, in which an intermediate product obtainable at any stage thereof is used as starting material and the remaining steps are carried out, or in which the starting materials are formed in situ under the reaction conditions, or in which the reaction components are used in the form of their salts or optically pure material.

Pharmaceutical Compositions

[0623] In another aspect, the disclosure provides a pharmaceutical composition comprising one or more compounds of described herein or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, and one or more pharmaceutically acceptable carriers. As used herein, the term "pharmaceutical composition" refers to a compound of the disclosure, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, together with at least one pharmaceutically acceptable carrier, in a form suitable for oral or parenteral administration.

[0624] As used herein, the term "pharmaceutically acceptable carrier" refers to a substance useful in the preparation or use of a pharmaceutical composition and includes, for example, suitable diluents, solvents, dispersion media, surfactants, antioxidants, preservatives, isotonic agents, buffering agents, emulsifiers, absorption delaying agents, salts, drug stabilizers, binders, excipients, disintegration agents, lubricants, wetting agents, sweetening agents, flavoring agents, dyes, and combinations thereof, as would be known to those skilled in the art (see, for example, Remington The Science and Practice of Pharmacy, 22nd Ed. Pharmaceutical Press, 2013, pp. 1049-1070).

[0625] In another aspect, the disclosure provides a pharmaceutical composition comprising a compound of the disclosure, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, and a pharmaceutically acceptable carrier. In a further embodiment, the composition comprises at least two pharmaceutically acceptable carriers, such as those described herein. For purposes of the disclosure, unless designated otherwise, solvates and hydrates are generally considered compositions. Preferably, pharmaceutically acceptable carriers are sterile. The pharmaceutical composition can be formulated for particular routes of administration such as oral administration, parenteral administration, and rectal administration, etc. In addition, the pharmaceutical compositions of the disclosure can be made up in a solid form (including without limitation capsules, tablets, pills, granules, powders or suppositories), or in a liquid form (including without limitation solutions, suspensions or emulsions). The pharmaceutical compositions can be subjected to conventional pharmaceutical operations such as sterilization and/or can contain conventional inert diluents, lubricating agents, or buffering agents, as well as adjuvants, such as preservatives, stabilizers, wetting agents, emulsifiers and buffers, etc.

[0626] Typically, the pharmaceutical compositions are tablets or gelatin capsules comprising the active ingredient together with one or more of:

[0627] a) diluents, e.g., lactose, dextrose, sucrose, mannitol, sorbitol, cellulose and/or glycine;

[0628] b) lubricants, e.g., silica, talcum, stearic acid, its magnesium or calcium salt and/or polyethyleneglycol;

[0629] c) binders, e.g., magnesium aluminum silicate, starch paste, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose and/or polyvinylpyrrolidone;

[0630] d) disintegrants, e.g., starches, agar, alginic acid or its sodium salt, or effervescent mixtures; and

[0631] e) absorbents, colorants, flavors and sweeteners.

[0632] In an embodiment, the pharmaceutical compositions are capsules comprising the active ingredient only.

[0633] Tablets may be either film coated or enteric coated according to methods known in the art.

[0634] Suitable compositions for oral administration include an effective amount of a compound of the disclosure in the form of tablets, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsion, hard or soft capsules, or syrups or elixirs, solutions or solid dispersion. Compositions intended for oral use are prepared according to any method known in the art for the manufacture of pharmaceutical compositions and such compositions can contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets may contain the active ingredient in admixture with nontoxic pharmaceutically acceptable excipients, which are suitable for the manufacture of tablets. These excipients are, for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example, starch, gelatin or acacia; and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets are uncoated or coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate can be employed. Formulations for oral use can be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example, peanut oil, liquid paraffin or olive oil.

[0635] Certain injectable compositions are aqueous isotonic solutions or suspensions, and suppositories are advantageously prepared from fatty emulsions or suspensions. Said compositions may be sterilized and/or contain adjuvants, such as preserving, stabilizing, wetting or emulsifying agents, solution promoters, salts for regulating the osmotic pressure and/or buffers. In addition, they may also contain other therapeutically valuable substances. Said compositions are prepared according to conventional mixing, granulating or coating methods, respectively, and contain about 0.1-75%, or contain about 1-50%, of the active ingredient.

[0636] Suitable compositions for transdermal application include an effective amount of a compound of the disclosure with a suitable carrier. Carriers suitable for transdermal delivery include absorbable pharmacologically acceptable solvents to assist passage through the skin of the host. For example, transdermal devices are in the form of a bandage comprising a backing member, a reservoir containing the compound optionally with carriers, optionally a rate controlling barrier to deliver the compound of the skin of the

host at a controlled and predetermined rate over a prolonged period of time, and means to secure the device to the skin.

[0637] Suitable compositions for topical application, e.g., to the skin and eyes, include aqueous solutions, suspensions, ointments, creams, gels or sprayable formulations, e.g., for delivery by aerosol or the like. Such topical delivery systems will in particular be appropriate for dermal application, e.g., for the treatment of skin cancer, e.g., for prophylactic use in sun creams, lotions, sprays and the like. They are thus particularly suited for use in topical, including cosmetic, formulations well-known in the art. Such may contain solubilizers, stabilizers, tonicity enhancing agents, buffers and preservatives.

[0638] As used herein a topical application may also pertain to an inhalation or to an intranasal application. They may be conveniently delivered in the form of a dry powder (either alone, as a mixture, for example a dry blend with lactose, or a mixed component particle, for example with phospholipids) from a dry powder inhaler or an aerosol spray presentation from a pressurised container, pump, spray, atomizer or nebuliser, with or without the use of a suitable propellant.

[0639] The compounds of formulae (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), (Ie) in free form or in pharmaceutically acceptable salt form, exhibit valuable pharmacological properties, e.g., WIZ modulating properties or WIZ degrading properties or HbF inducing properties e.g., as indicated in the in vitro tests as provided in the examples, and are therefore indicated for therapy or for use as research chemicals, e.g., as tool compounds.

[0640] Additional properties of the disclosed compounds include having good potency in the biological assays described herein, favorable safety profile, and possess favorable pharmacokinetic properties.

Diseases and Disorders

[0641] In an embodiment of the present disclosure, there is provided a compound of the disclosure, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, which is effective in reducing WIZ protein expression levels and/or inducing fetal hemoglobin (HbF) expression.

[0642] The compounds of the disclosure can be used to treat one or more of the diseases or disorders described herein below. In one embodiment, the disease or disorder is affected by the reduction of WIZ protein expression levels and/or induction of fetal hemoglobin protein expression levels. In another embodiment, the disease or disorder is a hemoglobinopathy, e.g., beta hemoglobinopathy, including sickle cell disease (SCD) and beta-thalassemia.

Methods of Use

[0643] All the aforementioned embodiments and embodiments hereinafter relating to the methods of reducing WIZ protein expression levels and/or inducing fetal hemoglobin (HbF) expression are equally applicable to:

[0644] A compound of the disclosure, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in a method of reducing WIZ protein expression levels and/or inducing fetal hemoglobin (HbF) expression;

- [0645]** A compound of the disclosure, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in the treatment of the aforementioned diseases or disorders according to the present disclosure;
- [0646]** Use of a compound of the disclosure, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, in the treatment of the aforementioned diseases or disorders according to the present disclosure; and
- [0647]** A pharmaceutical composition comprising a compound of the disclosure, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, for use in the treatment of the aforementioned diseases or disorders according to the present disclosure.
- [0648]** Having regard to their activity as WIZ modulators or degraders, compounds of formulae (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), and (Ie) in free or pharmaceutically acceptable salt form, are useful in the treatment of conditions which may be treated by modulation of WIZ protein expression levels, reduction of WIZ protein expression levels, or induction of fetal hemoglobin (HbF), such as in a blood disorder, for example an inherited blood disorder, e.g., sickle cell disease, or beta-thalassemia. In one aspect, the disclosure provides a method of treating or preventing a disease or disorder in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.
- [0649]** In another aspect, the disclosure provides a method of treating or preventing a disorder that is affected by the reduction of WIZ protein levels, in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.
- [0650]** In another aspect, the disclosure provides a method of inhibiting WIZ protein expression in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.
- [0651]** In another aspect, the disclosure provides a method of degrading WIZ protein in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.
- [0652]** In another aspect, the disclosure provides a method of inhibiting, reducing, or eliminating the activity of WIZ protein or WIZ protein expression, the method comprising administering to the subject a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.
- (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.
- [0653]** In another aspect, the disclosure provides a method of inducing or promoting fetal hemoglobin in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.
- [0654]** In another aspect, the disclosure provides a method of reactivating fetal hemoglobin production or expression in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.
- [0655]** In another aspect, the disclosure provides a method of increasing fetal hemoglobin expression in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.
- [0656]** In another aspect, the disclosure provides a method of treating a hemoglobinopathy, e.g., a beta-hemoglobinopathy, in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.
- [0657]** In another aspect, the disclosure provides a method of treating a sickle cell disease in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.
- [0658]** In another aspect, the disclosure provides a method of treating beta-thalassemia in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.
- [0659]** In an embodiment, the beta-thalassemia major or intermedia is the result of homozygous null or compound heterozygous mutations resulting with beta-globin deficiency and the phenotypic complications of beta-thalassemia, whether transfusion-dependent or not.
- [0660]** In another aspect, the disclosure provides a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof for use in a method of treating or preventing a disease or

for use in a method of treating beta-thalassemia in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.

[0671] In an embodiment, the beta-thalassemia major or intermedia is the result of homozygous null or compound heterozygous mutations resulting with beta-globin deficiency and the phenotypic complications of beta-thalassemia, whether transfusion-dependent or not.

Dosage

[0672] The pharmaceutical composition or combination of the disclosure can be in unit dosage of about 1-1000 mg of active ingredient(s) for a subject of about 50-70 kg, or about 1-500 mg or about 1-250 mg or about 1-150 mg or about 0.5-100 mg, or about 1-50 mg of active ingredients. The therapeutically effective dosage of a compound, the pharmaceutical composition, or the combinations thereof, is dependent on the species of the subject, the body weight, age and individual condition, the disorder or disease or the severity thereof being treated.

[0673] The above-cited dosage properties are demonstrable in vitro and in vivo tests using advantageously mammals, e.g., mice, rats, dogs, monkeys or isolated organs, tissues and preparations thereof. The compounds of the disclosure can be applied in vitro in the form of solutions, e.g., aqueous solutions, and in vivo either enterally, parenterally, advantageously intravenously, e.g., as a suspension or in aqueous solution. The dosage in vitro may range between about 10⁻³ molar and 10⁻⁹ molar concentrations. A therapeutically effective amount in vivo may range depending on the route of administration, between about 0.1-500 mg/kg, or between about 1-100 mg/kg.

[0674] The activity of a compound according to the disclosure can be assessed by the in vitro methods described in the Examples.

Combination Therapy

[0675] In another aspect, the disclosure provides a pharmaceutical combination comprising a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, and one or more additional therapeutic agent(s) for simultaneous, separate or sequential use in therapy. In an embodiment, the additional therapeutic agent is a myelosuppressive agent, such as hydroxyurea.

[0676] Combination therapy includes the administration of the subject compounds in further combination with other biologically active ingredients (such as, but not limited to, a second and different antineoplastic agent or a therapeutic agent that targets HbF or another cancer target) and non-drug therapies (such as, but not limited to, surgery or radiation treatment). For instance, the compounds of the application can be used in combination with other pharmaceutically active compounds, preferably compounds that are able to enhance the effect of the compounds of the application.

[0677] The compound of the disclosure may be administered either simultaneously with, or before or after, one or more other therapeutic agents. The compound of the disclosure may be administered separately, by the same or different route of administration, or together in the same pharmaceutical composition as the other agents. A therapeutic agent is, for example, a chemical compound, peptide, antibody, antibody fragment or nucleic acid, which is therapeutically active or enhances the therapeutic activity when administered to a patient in combination with a compound of the disclosure. Thus, in one embodiment, the disclosure provides a combination comprising a therapeutically effective amount of a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof and one or more additional therapeutically active agents.

[0678] In one embodiment, the disclosure provides a product comprising a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), and at least one other therapeutic agent as a combined preparation for simultaneous, separate or sequential use in therapy. In one embodiment, the therapy is the treatment of a disease or condition modulated by WIZ. Products provided as a combined preparation include a composition comprising the compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), and the other therapeutic agent(s) together in the same pharmaceutical composition, or the compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), and the other therapeutic agent(s) in separate form, e.g., in the form of a kit.

[0679] In one embodiment, the disclosure provides a pharmaceutical composition comprising a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie), and another therapeutic agent(s). Optionally, the pharmaceutical composition may comprise a pharmaceutically acceptable carrier, as described above.

[0680] In one embodiment, the disclosure provides a kit comprising two or more separate pharmaceutical compositions, at least one of which contains a compound of formula (I''), (I'), (I), (Ia''), (Ia'), (Ia), (Ib''), (Ib'), (Ib), (Ic''), (Ic'), (Ic), (Id''), (Id'), (Id), (Id-1), (Id-2), (Id-3), (Ie''), (Ie'), or (Ie). In one embodiment, the kit comprises means for separately retaining said compositions, such as a container, divided bottle, or divided foil packet. An example of such a kit is a blister pack, as typically used for the packaging of tablets, capsules and the like.

[0681] The kit of the disclosure may be used for administering different dosage forms, for example, oral and parenteral, for administering the separate compositions at different dosage intervals, or for titrating the separate compositions against one another. To assist compliance, the kit of the disclosure typically comprises directions for administration.

[0682] In the combination therapies of the disclosure, the compound of the disclosure and the other therapeutic agent may be manufactured and/or formulated by the same or different manufacturers. Moreover, the compound of the disclosure and the other therapeutic may be brought together

into a combination therapy: (i) prior to release of the combination product to physicians (e.g., in the case of a kit comprising the compound of the disclosure and the other therapeutic agent); (ii) by the physician themselves (or under the guidance of the physician) shortly before administration; (iii) in the patient themselves, e.g., during sequential administration of the compound of the disclosure and the other therapeutic agent.

Preparation of Compounds

[0683] It is understood that in the following description, combinations of substituents and/or variables of the depicted formulae are permissible only if such combinations result in stable compounds.

[0684] It will also be appreciated by those skilled in the art that in the processes described below, the functional groups of intermediate compounds may need to be protected by suitable protecting groups. Such functional groups include hydroxy, phenol, amino and carboxylic acid. Suitable protecting groups for hydroxy or phenol include trialkylsilyl or diarylalkylsilyl (e.g., *t*-butyldimethylsilyl, *t*-butyldiphenylsilyl or trimethylsilyl), tetrahydropyranyl, benzyl, substituted benzyl, methyl, and the like. Suitable protecting groups for amino, amidino and guanidino include *t*-butoxycarbonyl, benzyloxycarbonyl, and the like. Suitable protecting groups for carboxylic acid include alkyl, aryl or arylalkyl esters.

[0685] Protecting groups may be added or removed in accordance with standard techniques, which are well-known to those skilled in the art and as described herein. The use of protecting groups is described in detail in J. F. W. McOmie, "Protective Groups in Organic Chemistry", Plenum Press, London and New York 1973; T. W. Greene and P. G. M. Wuts, "Greene's Protective Groups in Organic Synthesis", Fourth Edition, Wiley, New York 2007; P. J. Kocienski, "Protecting Groups", Third Edition, Georg Thieme Verlag, Stuttgart and New York 2005; and in "Methoden der organischen Chemie" (Methods of Organic Chemistry), Houben Weyl, 4th edition, Volume 15/I, Georg Thieme Verlag, Stuttgart 1974.

[0686] The protecting group may also be a polymer resin, such as a Wang resin or a 2-chlorotriptyl-chloride resin.

[0687] The following reaction schemes illustrate methods to make compounds of this disclosure. It is understood that one skilled in the art would be able to make these compounds by similar methods or by methods known to one skilled in the art. In general, starting components and reagents may be obtained from sources such as Sigma Aldrich, Lancaster Synthesis, Inc., Maybridge, Matrix Scientific, TCI, and Fluorochem USA, Strem, other commercial vendors, or synthesized according to sources known to those skilled in the art, or prepared as described in this disclosure.

Analytical Methods, Materials, and Instrumentation

[0688] Unless otherwise noted, reagents and solvents were used as received from commercial suppliers. Proton nuclear magnetic resonance (NMR) spectra were obtained on either Bruker Avance spectrometer or Varian Oxford 400 MHz spectrometer unless otherwise noted. Spectra are given in ppm (δ) and coupling constants, J, are reported in Hertz. Tetramethylsilane (TMS) was used as an internal standard. Chemical shifts are reported in ppm relative to dimethyl sulfoxide (δ 2.50), methanol (δ 3.31), chloroform (δ 7.26) or

other solvent as indicated in NMR spectral data. A small amount of the dry sample (2-5 mg) is dissolved in an appropriate deuterated solvent (1 mL). The chemical names were generated using ChemBioDraw Ultra v12 from CambridgeSoft.

[0689] Mass spectra (ESI-MS) were collected using a Waters System (Acquity UPLC and a Micromass ZQ mass spectrometer) or Agilent-1260 Infinity (6120 Quadrupole); all masses reported are the *m/z* of the protonated parent ions unless recorded otherwise. The sample was dissolved in a suitable solvent such as MeCN, DMSO, or MeOH and was injected directly into the column using an automated sample handler. The analysis is performed on Waters Acquity UPLC system (Column: Waters Acquity UPLC BEH C18 1.7 μ m, 2.1x30 mm; Flow rate: 1 mL/min; 55° C. (column temperature); Solvent A: 0.05% formic acid in water, Solvent B: 0.04% formic acid in MeOH; gradient 95% Solvent A from 0 to 0.10 min; 95% Solvent A to 20% Solvent A from 0.10 to 0.50 min; 20% Solvent A to 5% Solvent A from 0.50 to 0.60 min; hold at 5% Solvent A from 0.6 min to 0.8 min; 5% Solvent A to 95% Solvent A from 0.80 to 0.90 min; and hold 95% Solvent A from 0.90 to 1.15 min.

Abbreviations


[0690]	ACN acetonitrile
[0691]	AcOH acetic acid
[0692]	AIBN azobisisobutyronitrile
[0693]	aq. aqueous
[0694]	B ₂ pin ₂ bis(pinacolato)diboron
[0695]	9-BBN 9-borabicyclo[3.3.1]nonane
[0696]	Boc ₂ O di- <i>tert</i> -butyl dicarbonate
[0697]	Bn benzyl
[0698]	BnBr benzyl bromide
[0699]	br broad
[0700]	d doublet
[0701]	dd doublet of doublets
[0702]	ddd doublet of doublet of doublets
[0703]	ddq doublet of doublet of quartets
[0704]	ddt doublet of doublet of triplets
[0705]	dq doublet of quartets
[0706]	dt doublet of triplets
[0707]	dtbbpy 4,4'-di- <i>tert</i> -butyl-2,2'-dipyridyl
[0708]	dtd doublet of triplet of doublets
[0709]	Cs ₂ CO ₃ cesium carbonate
[0710]	DCE 1,2-dichloroethane
[0711]	DCM dichloromethane
[0712]	DHP dihydropyran
[0713]	DIBAL-H diisobutylaluminum hydride
[0714]	DIPEA (DIEA) diisopropylethylamine
[0715]	DIPEA N,N-diisopropylethylamine
[0716]	DMA N,N-dimethylacetamide
[0717]	DMAP 4-dimethylaminopyridine
[0718]	DMB 2,4-dimethoxybenzyl
[0719]	DME 1,2-dimethoxyethane
[0720]	DMF N,N-dimethylformamide
[0721]	DMP Dess-Martin periodinane or 1,1,1-Tris(acetyloxy)-1,1-dihydro-1,2-benziodoxol-3-(1H)-one
[0722]	DMSO dimethylsulfoxide
[0723]	EC ₅₀ half maximal effective concentration
[0724]	ELSD evaporative light scattering detector
[0725]	EtOH ethanol
[0726]	Et ₂ O diethyl ether
[0727]	Et ₃ N triethylamine
[0728]	EtOAc ethyl acetate

[0729] HATU 1-[bis(dimethylamino)methylene]-1H-1,2,3-triazolo[4,5-b]pyridinium 3-oxid hexafluorophosphate
 [0730] HCl hydrogen chloride
 [0731] hept heptet
 [0732] HPLC high performance liquid chromatography
 [0733] h or hr hour
 [0734] HRMS high resolution mass spectrometry
 [0735] g gram
 [0736] g/min gram per minute
 [0737] IC₅₀ half maximal inhibitory concentration
 [0738] IPA (iPrOH) isopropyl alcohol
 [0739] Ir[(dF(CF₃)ppy)₂dtbbpy]PF₆ [4,4'-Bis(1,1-dimethylethyl)-2,2'-bipyridine-N1,N1']bis[3,5-difluoro-2-[5-(trifluoromethyl)-2-pyridinyl-N]phenyl-C]Iridium (III) hexafluorophosphate
 [0740] K₂CO₃ potassium carbonate
 [0741] KI potassium iodide
 [0742] KOAc potassium Acetate
 [0743] K₃PO₄ tripotassium phosphate
 [0744] LCMS liquid chromatography mass spectrometry
 [0745] LDA lithium diisopropylamide
 [0746] m multiplet
 [0747] MeCN acetonitrile
 [0748] MeOH methanol
 [0749] mg milligram
 [0750] MHz megahertz
 [0751] min minutes
 [0752] mL milliliter
 [0753] mmol millimole
 [0754] M molar
 [0755] MS mass spectrometry
 [0756] NaH sodium hydride
 [0757] NaHCO₃ sodium bicarbonate
 [0758] NaBH(OAc)₃ sodium triacetoxyborohydride
 [0759] Na₂SO₄ sodium sulfate
 [0760] NBS N-bromosuccinimide
 [0761] NMM N-methylmorpholine
 [0762] NMP N-methyl-2-pyrrolidone
 [0763] NMR nuclear magnetic resonance
 [0764] on overnight
 [0765] Pd/C palladium on carbon
 [0766] PdCl₂(dppf)-DCM [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) complex with dichloromethane
 [0767] Pd(PPh₃)₄ tetrakis(triphenylphosphine)palladium(0)
 [0768] PMB para-methoxybenzyl
 [0769] q quartet
 [0770] qd quartet of doublets
 [0771] quint quintet
 [0772] quintd quintet of doublets
 [0773] rbf round bottom flask
 [0774] RockPhos G3 Pd [(2-di-tert-butylphosphino-3-methoxy-6-methyl-2',4',6'-triisopropyl-1,1'-biphenyl)-2-(2-aminobiphenyl)]palladium(II) methanesulfonate
 [0775] rt or r.t. room temperature
 [0776] Rt retention time
 [0777] RuPhos dicyclohexyl(2',6'-diisopropoxy-[1,1'-biphenyl]-2-yl)phosphane
 [0778] s singlet
 [0779] SEM 2-(trimethylsilyl)ethoxymethyl
 [0780] SnBu₃ tributyltin

[0781] t triplet
 [0782] td triplet of doublets
 [0783] tdd triplet of doublet of doublets
 [0784] TBAI tetrabutylammonium iodide
 [0785] TEA (NEt₃) triethylamine
 [0786] TFA trifluoroacetic acid
 [0787] TfOH triflic Acid
 [0788] THF tetrahydrofuran
 [0789] THP tetrahydropyran
 [0790] TMP 2,2,6,6-tetramethylpiperidine
 [0791] Ts tosyl
 [0792] tt triplet of triplets
 [0793] ttd triplet of triplet of doublets
 [0794] TLC thin-layer chromatography
 [0795] UPLC ultra-Performance liquid Chromatography
 [0796] XPhos Pd G2 chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II)
 [0797] μW or uW microwave

Preparation of Intermediates

Preparation of 3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione



Step 1. tert-butyl (3-((2,4-dimethoxybenzyl)amino)-3-oxopropyl)carbamate

[0798] To a solution of 3-((tert-butoxycarbonyl)amino)propanoic acid (200 g, 1.06 mol, 1.00 eq) in DCM (1200 mL) was added CDI (189 g, 1.16 mol, 1.10 eq). The reaction mixture was stirred at 20° C. for 2 h. The reaction mixture was slowly added to a solution of (2,4-dimethoxyphenyl)methanamine (212 g, 1.27 mol, 191 mL, 1.20 eq) and DMAP (12.9 g, 106 mmol, 0.10 eq) in DCM (1000 mL). The

solution was stirred at 20° C. for 12 h. The reaction mixture was slowly poured into water (2 L) and stirred at rt for 10 min. The organic phase was separated and the water phase was extracted with DCM (800 mL×2). The combined organic phase was washed with brine (1 L×2) and dried over Na₂SO₄, filtered and concentrated. The crude material was purified by silica gel chromatography (eluted with 50:1 to 2:1 petroleum ether:ethyl acetate) to give tert-butyl 3-((2,4-dimethoxybenzyl)amino)-3-oxopropyl)carbamate (210 g, 621 mmol, 59% yield) as a white solid. ¹H NMR (400 MHz, DMSO-d₆) δ 8.06 (t, J=5.6 Hz, 1H), 7.08-7.01 (m, 1H), 6.72 (br t, J=5.3 Hz, 1H), 6.55-6.51 (m, 1H), 6.46 (dd, J=2.4, 8.4 Hz, 1H), 4.13 (d, J=5.6 Hz, 2H), 3.77 (s, 3H), 3.73 (s, 3H), 3.13 (q, J=7.0 Hz, 2H), 2.28 (t, J=7.3 Hz, 2H), 1.37 (s, 9H).

Step 2.


3-amino-N-(2,4-dimethoxybenzyl)propanamide hydrochloride

[0799] To a solution of tert-butyl 3-((2,4-dimethoxybenzyl)amino)-3-oxopropyl)carbamate (210 g, 621 mmol, 1.00 eq) in DCM (1000 mL) was added HCl/dioxane (4 M, 1000 mL, 6.45 eq) and the solution was stirred at 20° C. for 5 h. The reaction mixture was concentrated to give 3-amino-N-(2,4-dimethoxybenzyl)propanamide hydrochloride (180 g, crude, HCl salt) as a white solid which was used in the next step without further purification. ¹H NMR (400 MHz, DMSO-d₆) δ 8.38 (br t, J=5.6 Hz, 1H), 8.07 (br s, 3H), 7.08 (d, J=8.3 Hz, 1H), 6.53 (d, J=2.4 Hz, 1H), 6.46 (dd, J=2.4, 8.3 Hz, 1H), 4.15 (d, J=5.6 Hz, 2H), 3.75 (d, J=15.6 Hz, 6H), 2.96 (sxt, J=6.3 Hz, 2H), 2.59-2.52 (m, 2H).


Step 3. 3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione

[0800] To a mixture of 3-amino-N-(2,4-dimethoxybenzyl)propanamide hydrochloride (180 g, 655 mmol, 1.00 eq) and DIPEA (212 g, 1.64 mol, 285 mL, 2.50 eq) in DCE (1200 mL) was added CDI (127 g, 786 mmol, 1.20 eq) at 0° C. The mixture was stirred at 0° C. for 0.5 h, then heated to 100° C. and stirred for 12 h. The reaction mixture was slowly poured into water (1000 mL) and stirred at 20° C. for 20 min. The organic phase was separated and the water phase was extracted with DCM (500 mL×2). The combined organic phases were washed with brine (500 mL×2), dried over Na₂SO₄, filtered and concentrated. The residue was purified by silica gel chromatography (45:1 to 0:1 petroleum ether:ethyl acetate) to give 3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione (120 g, 454 mmol, 69% yield) as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 6.99 (d, J=8.3 Hz, 1H), 6.48-6.37 (m, 2H), 5.79 (br s, 1H), 4.93 (s, 2H), 3.80 (d, J=15.3 Hz, 6H), 3.41 (dt, J=2.6, 6.8 Hz, 2H), 2.76 (t, J=6.8 Hz, 2H).

Preparation of potassium (R)-((4-(tert-butoxycarbonyl)-3-methylpiperazin-1-yl)methyl)trifluoroborate



-continued



Step 1. (((3R)-4-(tert-butoxycarbonyl)-3-methylpiperazin-1-ium-1-yl)methyl)trifluoroborate








[0801] To a solution of potassium (bromomethyl)trifluoroborate (2.00 g, 9.96 mmol) in THF (10 mL) was added tert-butyl (R)-2-methylpiperazine-1-carboxylate (2.09 g, 15.7 mmol). The reaction mixture was stirred at 80° C. for 16 h. The reaction mixture was filtered and the filter cake was washed with THF (2×10 mL), and the filter cake was collected and dried to give (((3R)-4-(tert-butoxycarbonyl)-3-methylpiperazin-1-ium-1-yl)methyl)trifluoroborate (4.3 g, crude) as a white solid. The crude was used in the next step without any other purification. ¹H NMR (400 MHz, DMSO-d₆) δ 8.45-8.44 (m, 1H), 4.31-2.92 (m, 1H), 3.87-3.82 (m, 1H), 3.67-3.54 (m, 1H), 3.27-3.04 (m, 2H), 2.99-2.77 (m, 2H), 1.99 (br s, 2H), 1.83-1.70 (m, 1H), 1.50-1.37 (m, 9H), 1.21 (br d, J=7.2 Hz, 3H).

Step 2. potassium (R)-((4-(tert-butoxycarbonyl)-3-methylpiperazin-1-yl)methyl)trifluoroborate






[0802] To a solution of (((3R)-4-(tert-butoxycarbonyl)-3-methylpiperazin-1-ium-1-yl)methyl)trifluoroborate (4.3 g, crude) in acetone (20 mL) was added K₂CO₃ (2.10 g, 15.2 mmol) and the reaction mixture was stirred at 25° C. for 16 h. The reaction mixture was filtered and the filter cake was washed with acetone (2×10 mL), and the filtrate was concentrated to give potassium (R)-((4-(tert-butoxycarbonyl)-3-methylpiperazin-1-yl)methyl)trifluoroborate (1.1 g, crude) as a white solid. The crude material was used in the next step without any other purification. ¹H NMR (400 MHz, DMSO-d₆) δ 4.09 (br s, 1H), 3.79-3.60 (m, 1H), 3.51-3.21 (m, 1H), 2.98 (br s, 3H), 1.71-1.46 (m, 2H), 1.39 (s, 9H), 1.15 (d, J=7.2 Hz, 3H).

Additional Borate Salts Prepared by the Method Above:







[0803] The borate salts in the following table were prepared by the method of potassium (R)-((4-(tert-butoxycarbonyl)-3-methylpiperazin-1-yl)methyl)trifluoroborate using the appropriate commercially available piperazine in step 1 except where noted.

Structure	Starting material
 potassium (S)-((4-(tert-butoxycarbonyl)-3-ethylpiperazin-1-yl)methyl)trifluoroborate	tert-butyl (S)-2-ethylpiperazine-1-carboxylate
 potassium (S)-((4-(tert-butoxycarbonyl)-3-isopropylpiperazin-1-yl)methyl)trifluoroborate	tert-butyl (S)-2-isopropylpiperazine-1-carboxylate
 potassium ((4-(tert-butoxycarbonyl)-3,3-dimethylpiperazin-1-yl)methyl)trifluoroborate	tert-butyl 2,2-dimethylpiperazine-1-carboxylate
 potassium (((1R,5S)-8-(tert-butoxycarbonyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)methyl)trifluoroborate	tert-butyl (1R,5S)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate
 potassium (((1R,4R)-5-(tert-butoxycarbonyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)trifluoroborate	tert-butyl (1R,4R)-2,5-diazabicyclo[2.2.1]heptane-2-carboxylate
 potassium (((1S,4S)-5-(tert-butoxycarbonyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)trifluoroborate	tert-butyl (1S,4S)-2,5-diazabicyclo[2.2.1]heptane-2-carboxylate
 potassium trifluoro((4-(3-methylbutan-2-yl)piperazin-1-yl)methyl)borate	1-(3-methylbutan-2-yl)piperazine



-continued

Structure	Starting material
	tert-butyl 1,4-diazepane-1-carboxylate
potassium ((4-(tert-butoxycarbonyl)-1,4-diazepan-1-yl)methyl)trifluoroborate	
	1-(cyclohexylmethyl)piperazin-2-one [see <i>RSC Advances</i> , 2018, 8, 11163-11176]
potassium ((4-(cyclohexylmethyl)-3-oxopiperazin-1-yl)methyl)trifluoroborate	
	1-(2-cyclohexylethyl)piperazin-2-one [see <i>J. Med. Chem.</i> 1990, 33, 2590-2595]
potassium ((4-(2-cyclohexylethyl)-3-oxopiperazin-1-yl)methyl)trifluoroborate	
	(R)-1-(cyclohexylmethyl)-2-(methoxymethyl)piperazine hydrochloride [vide infra]
potassium (R)-((4-(cyclohexylmethyl)-3-(methoxymethyl)piperazin-1-yl)methyl)trifluoroborate	
	(R)-1-isobutyl-2-(methoxymethyl)piperazine hydrochloride [vide infra]
potassium (R)-trifluoro((4-isobutyl-3-(methoxymethyl)piperazin-1-yl)methyl)borate	


-continued

Structure	Starting material
	(R)-1-(cyclohexylmethyl)-2-(difluoromethyl)piperazine hydrochloride [vide infra]
potassium (R)-((4-(cyclohexylmethyl)-3-(difluoromethyl)piperazin-1-yl)methyl)trifluoroborate	
	(R)-2-(difluoromethyl)-1-isobutylpiperazine hydrochloride [vide infra]
potassium (R)-((3-(difluoromethyl)-4-isobutylpiperazin-1-yl)methyl)trifluoroborate	
	1-(3,3,3-trifluoro-2,2-dimethylpropyl)piperazine hydrochloride [vide infra]
potassium trifluoro((4-(3,3,3-trifluoro-2,2-dimethylpropyl)piperazin-1-yl)methyl)borate	
	(S)-octahydropyrrolo[1,2-a]pyrazine
potassium (S)-trifluoro((hexahydro-pyrrolo[1,2-a]pyrazin-2(1H)-yl)methyl)borate	
	(R)-octahydropyrrolo[1,2-a]pyrazine
potassium (R)-trifluoro((hexahydro-pyrrolo[1,2-a]pyrazin-2(1H)-yl)methyl)borate	
	(S)-octahydro-2H-pyrido[1,2-a]pyrazine
potassium (S)-trifluoro((octahydro-2H-pyrido[1,2-a]pyrazin-2-yl)methyl)borate	

-continued

Structure	Starting material
 potassium (R)-trifluoro((hexahydropyrazino[2,1-c][1,4]oxazin-8(1H)-yl)methyl)borate	(R)-octahydropyrazino[2,1-c][1,4]oxazine
 potassium (S)-trifluoro((hexahydropyrazino[2,1-c][1,4]oxazin-8(1H)-yl)methyl)borate	(S)-octahydropyrazino[2,1-c][1,4]oxazine

Preparation of (R)-1-(cyclohexylmethyl)-2-(methoxymethyl)piperazine hydrochloride



Step 1. tert-butyl (R)-4-(cyclohexylmethyl)-3-(hydroxymethyl)piperazine-1-carboxylate

[0804] To a stirred solution of tert-butyl (R)-3-(hydroxymethyl)piperazine-1-carboxylate (6.0 g, 27.7 mmol) and cyclohexanecarbaldehyde (4.6 g, 41.6 mmol) dissolved in

DCM (70 mL) was added Et₃N (11.7 mL, 83.2 mmol). The reaction mixture was stirred for 30 min at rt and then sodium triacetoxy borohydride (11.7 g, 55.5 mmol) was added in portions at 0° C. The reaction mixture was allowed to stirred at rt for 16 h. The reaction was diluted with DCM and water and the organic layer was dried over Na₂SO₄, filtered and concentrated. The crude compound was purified by silica gel chromatography (eluting with 10-20% EtOAc in hexanes) to afford tert-butyl (R)-4-(cyclohexylmethyl)-3-(hydroxymethyl)piperazine-1-carboxylate (4.2 g, 13.4 mmol, 48% yield). LCMS [M+H-tBu]⁺: 257.2.


Step 2. tert-butyl (R)-4-(cyclohexylmethyl)-3-(methoxymethyl)piperazine-1-carboxylate

[0805] To a stirred solution of tert-butyl (R)-4-(cyclohexylmethyl)-3-(hydroxymethyl)piperazine-1-carboxylate (0.70 g, 2.2 mmol) in DMF (10 mL) cooled to 0° C. was added NaH (0.13 g, 3.36 mmol) under an inert atmosphere. The reaction mixture was stirred at 0° C. for 30 min and then MeI (0.47 g, 3.36 mmol) was added at 0° C. The reaction was diluted with EtOAc and water and the organic layer was dried over Na₂SO₄, filtered and concentrated. The crude compound was purified by silica gel chromatography (eluting with 10-20% EtOAc in hexanes) to afford tert-butyl (R)-4-(cyclohexylmethyl)-3-(methoxymethyl)piperazine-1-carboxylate (0.45 g, 1.37 mmol, 61%). LCMS [M+H]⁺: 327.1

Step 3. (R)-1-(cyclohexylmethyl)-2-(methoxymethyl)piperazine hydrochloride


[0806] To a stirred solution of tert-butyl (R)-4-(cyclohexylmethyl)-3-(methoxymethyl)piperazine-1-carboxylate (0.45 g, 1.37 mmol) in DCM (7.0 mL) cooled to 0° C. was added a solution of HCl (4.0 M in dioxane, 4.0 mL). The reaction mixture was stirred at rt for 3 h and then concentrated. The crude compound was washed with diethyl ether to afford (R)-1-(cyclohexylmethyl)-2-(methoxymethyl)piperazine hydrochloride (0.40 g, crude). LCMS [M+H]⁺: 227.1.

Preparation of
(R)-1-isobutyl-2-(methoxymethyl)piperazine
hydrochloride



[0807] Prepared by the method of (R)-1-(cyclohexylmethyl)-2-(methoxymethyl)piperazine hydrochloride, using isobutyraldehyde in step 1. LCMS $[M+H]^+$: 187.1.

Preparation of (R)-1-(cyclohexylmethyl)-2-(difluoromethyl)piperazine hydrochloride



Step 1. tert-butyl (R)-4-(cyclohexylmethyl)-3-(hydroxymethyl)piperazine-1-carboxylate

[0808] To a stirred solution of tert-butyl (R)-3-(hydroxymethyl)piperazine-1-carboxylate (6.0 g, 27.75 mmol) and cyclohexanecarbaldehyde (4.6 g, 41.62 mmol) in DCM (70 mL) was added Et_3N (11.69 mL, 83.25 mmol). The reaction mixture was stirred for 30 min at rt. Sodium triacetoxy borohydride (11.7 g, 55.50 mmol) was then added slowly at 0° C. The reaction mixture was stirred at rt for 16 h. After completion, the reaction was diluted with DCM and water and the organic layer was dried over Na_2SO_4 , filtered and concentrated. The crude compound was purified by silica gel chromatography (eluting with 10-20% EtOAc in hexanes) to afford tert-butyl (R)-4-(cyclohexylmethyl)-3-(hydroxymethyl)piperazine-1-carboxylate (4.2 g, 13.44 mmol, 48% yield). LCMS $[M+H-t\text{Bu}]^+$: 257.2.

Step 2. tert-butyl (R)-4-(cyclohexylmethyl)-3-formylpiperazine-1-carboxylate

[0809] To a stirred solution of oxalyl chloride (2.04 mL, 24.0 mmol) in DCM (25 mL) at -78° C. was added DMSO (3.41 mL, 48.0 mmol) dropwise under an inert atmosphere. The reaction mixture was stirred at -78° C. for 15 min and then a solution of tert-butyl (R)-4-(cyclohexylmethyl)-3-(hydroxymethyl)piperazine-1-carboxylate (2.5 g, 8.0 mmol) in DCM (5.0 mL) was added dropwise at -78° C. The reaction mixture was stirred at -78° C. for 1 h and Et_3N (11.24 mL, 80.01 mmol) was added slowly. The reaction mixture was stirred at -78° C. for 1 h and allowed to warm to rt. The reaction was diluted with DCM and water and the organic layer was dried over Na_2SO_4 , filtered and concentrated to afford crude tert-butyl (R)-4-(cyclohexylmethyl)-3-formylpiperazine-1-carboxylate (2.7 g, crude). LCMS $[M+H]^+$: 311.1.


Step 3. tert-butyl (R)-4-(cyclohexylmethyl)-3-(difluoromethyl)piperazine-1-carboxylate

[0810] To a stirred solution of tert-butyl (R)-4-(cyclohexylmethyl)-3-formylpiperazine-1-carboxylate (2.7 g, 8.69 mmol) in DCM (30 mL) at 0° C. was added DAST (2.29 mL, 17.4 mmol) under an inert atmosphere. The reaction mixture was stirred at 0° C. for 2 h. After completion, the reaction was quenched with saturated aqueous NaHCO_3 solution and diluted with DCM. The organic layer was dried over Na_2SO_4 , filtered and concentrated. The crude compound was purified by silica gel chromatography (eluting with 10-15% EtOAc in hexanes) to afford tert-butyl (R)-4-(cyclohexylmethyl)-3-(difluoromethyl)piperazine-1-carboxylate (0.41 g, 1.23 mmol, 14% yield). LCMS $[M+H]^+$: 333.5.

Step 4. (R)-1-(cyclohexylmethyl)-2-(difluoromethyl)piperazine hydrochloride

[0811] To a stirred solution of tert-butyl (R)-4-(cyclohexylmethyl)-3-(difluoromethyl)piperazine-1-carboxylate (0.41 g, 1.2 mmol) in DCM (7.0 mL) at 0° C. was added a solution of HCl in dioxane (4.0 M, 4.0 mL). The reaction mixture was stirred at rt for 3 h. After completion, the mixture was concentrated and the crude compound was washed with diethyl ether to afford (R)-1-(cyclohexylmethyl)-2-(difluoromethyl)piperazine hydrochloride (0.33 g, 1.2 mmol, 100% yield). LCMS $[M+H]^+$: 232.9.

Preparation of
1-(3,3,3-trifluoro-2,2-dimethylpropyl)piperazine
hydrochloride



Step 1. tert-butyl 4-(3,3,3-trifluoro-2,2-dimethylpropanoyl)piperazine-1-carboxylate

[0812] To a stirred solution of 3,3,3-trifluoro-2,2-dimethylpropanoic acid (1.0 g, 6.4 mmol) in DMF (15 mL) was added DIPEA (3.35 mL, 19.2 mmol) followed by HATU (3.65 g, 9.60 mmol) under inert atmosphere. The reaction mixture was stirred at rt for 15 min. After 15 min tert-butyl piperazine-1-carboxylate (1.43 g, 7.68 mmol) was added and the mixture was stirred at rt for 16 h. The reaction was poured into cold water and the precipitate was filtered and dried under vacuum to afford crude tert-butyl 4-(3,3,3-trifluoro-2,2-dimethylpropanoyl)piperazine-1-carboxylate (1.0 g, 3.08, 48% yield). LCMS $[M+H-tBu]^+$: 269.1.

Step 2. tert-butyl 4-(3,3,3-trifluoro-2,2-dimethylpropyl)piperazine-1-carboxylate

[0813] To a stirred solution of tert-butyl 4-(3,3,3-trifluoro-2,2-dimethylpropanoyl)piperazine-1-carboxylate (1.0 g, 3.08 mmol) in THF (15 mL) at 0° C. was added $BH_3 \cdot DMS$ (15.4 mL, 30.8 mmol, 1M in THF) under inert atmosphere. The reaction mixture was then stirred at 50° C. for 16 h. The reaction was quenched with MeOH and concentrate. The residue was dissolved in DCM and washed sequentially with a solution of aqueous 2M $NaHCO_3$ and brine. The organic layer was dried over Na_2SO_4 , filtered and concentrated. The crude compound was purified by silica gel chromatography (eluting with 10-12% EtOAc in hexanes) to afford tert-butyl 4-(3,3,3-trifluoro-2,2-dimethylpropyl)piperazine-1-carboxylate (0.5 g, 1.61 mmol, 52% yield). LCMS $[M+H-tBu]^+$: 254.9.


Step 3.

1-(3,3,3-trifluoro-2,2-dimethylpropyl)piperazine
hydrochloride

[0814] To stirred solution of tert-butyl 4-(3,3,3-trifluoro-2,2-dimethylpropyl)piperazine-1-carboxylate (500 mg, 1.61

mmol) in DCM (7 mL) was added 4M HCl in dioxane (3 mL) and the mixture was stirred at rt for 2 h. The mixture was concentrated and the residue was washed with diethyl ether to afford 1-(3,3,3-trifluoro-2,2-dimethylpropyl)piperazine hydrochloride (500 mg, crude). LCMS $[M+H]^+$: 211.2.

Preparation of (3-iodopyrazolo[1,5-a]pyridin-5-yl)
methyl 4-methylbenzenesulfonate



Step 1. 3-iodopyrazolo[1,5-a]pyridine-5-carbalde-
hyde

[0815] NIS (1.4 g, 5.92 mmol) was added to a solution of pyrazolo[1,5-a]pyridine-5-carbaldehyde (790 mg, 5.41 mmol) in DMF (10 mL) at 0° C. The mixture was then stirred at rt for 8 h. After completion, the reaction was quenched with water and the solid that precipitated was collected by filtration and dried under vacuum to afford 3-iodopyrazolo[1,5-a]pyridine-5-carbaldehyde (1.2 g, 4.4 mmol, 81% yield). LCMS $[M+H]^+$: 273.0.

Step 2. (3-iodopyrazolo[1,5-a]pyridin-5-yl)methanol


[0816] To a stirred solution of 3-iodopyrazolo[1,5-a]pyridine-5-carbaldehyde (1.2 g, 4.40 mmol) in MeOH:THF (2:1) (10 mL) was added $NaBH_4$ (250 mg, 6.60 mmol) at 0° C. The reaction mixture was stirred for 1 h and then concentrated. The residue was diluted with water and extracted with DCM. The organic layer was washed with brine, dried over Na_2SO_4 , filtered and concentrated to afford (3-iodopyrazolo[1,5-a]pyridin-5-yl)methanol (800 mg, crude). LCMS $[M+H]^+$: 274.7.

Step 3. (3-iodopyrazolo[1,5-a]pyridin-5-yl)methyl
4-methylbenzenesulfonate

[0817] To a stirred solution of (3-iodopyrazolo[1,5-a]pyridin-5-yl)methanol (700 mg, 2.55 mmol) in DCM (10 mL) at 0° C. was added TEA (0.8 mL, 3.66 mmol). The mixture was stirred for 10 min and then tosyl chloride (610 mg, 3.06 mmol) and DMAP (38 mg, 0.25 mmol) were added. The reaction was stirred at rt for 1 h. The mixture was then diluted with DCM and water and the organic layer was dried over Na_2SO_4 , filtered and concentrated to afford (3-iodopy-


razolo[1,5-a]pyridin-5-yl)methyl 4-methylbenzenesulfonate (1.0 g, crude). The crude material was used without further purification.

Preparation of
trans-3-methoxycyclobutane-1-carbaldehyde



[0818] To a solution of trans-3-methoxycyclobutylmethanol [see WO2021/124172, 2021, A1] (0.30 g, 2.6 mmol), 1.0 eq) in DCM (15 mL) at 0° C. was added DMP (1.2 g, 2.8 mmol, 1.1 eq). The reaction mixture was stirred at 0° C. for 2 h. The reaction mixture was then concentrated. The crude material was purified by neutral alumina chromatography (eluted with 15% EtOAc in hexane) to give trans-3-methoxycyclobutane-1-carbaldehyde (0.21 g, 1.8 mmol, 71% yield) as a colorless oil.

Preparation of
cis-3-methoxycyclobutane-1-carbaldehyde



[0819] To a solution of cis-3-methoxycyclobutylmethanol [see WO2021/124172, 2021, A1] (0.10 g, 0.86 mmol, 1.0 eq) in DCM (7 mL) at 0° C. was added DMP (0.40 g, 0.94 mmol, 1.1 eq). The reaction mixture was stirred at 0° C. for 2 h. The reaction mixture was then concentrated. The crude material was diluted with Et₂O (10 mL) and filtered through celite, washing with additional Et₂O. The filtrate was concentrated to give crude cis-3-methoxycyclobutane-1-carbaldehyde (0.12 g, 1.1 mmol). The crude material was used in the next step without any other purification.


Preparation of
(R)-3,3-difluorocyclopentane-1-carbaldehyde



[0820] To a solution of (R)-(3,3-difluorocyclopentyl)methanol (0.230 g, 1.69 mmol), 1.0 eq) in DCM (5 mL) at 0° C. was added DMP (0.788 g, 1.86 mmol, 1.1 eq). The reaction mixture was stirred at rt for 2 h. The reaction mixture was then concentrated. The crude material was diluted with EtOAc (10 mL) and filtered through celite. The filtrate was washed with water, dried over MgSO₄, filtered


and concentrated. The crude material was purified by neutral alumina chromatography (eluted with 20% EtOAc in hexane) to give (R)-3,3-difluorocyclopentane-1-carbaldehyde (0.13 g, 0.97 mmol, 57% yield) as a yellow oil.

Preparation of
(S)-3,3-difluorocyclopentane-1-carbaldehyde



[0821] To a solution of (S)-(3,3-difluorocyclopentyl)methanol (0.170 g, 1.24 mmol), 1.0 eq) in DCM (10 mL) at 0° C. was added DMP (0.582 g, 1.37 mmol, 1.1 eq). The reaction mixture was stirred at rt for 2 h. The reaction mixture was then concentrated. The crude material was purified by neutral alumina chromatography (eluted with 20% EtOAc in hexane) to give (S)-3,3-difluorocyclopentane-1-carbaldehyde (0.15 g) as a colorless oil.

Preparation of
(1r,3R,4S)-3,4-difluorocyclopentane-1-carbaldehyde





[0822] To a solution of ((1r,3R,4S)-3,4-difluorocyclopentyl)methanol [see EP2275414, 2011, A1] (0.170 g, 1.24 mmol), 1.0 eq) in DCM (10 mL) at 0° C. was added DMP (1.58 g, 3.74 mmol, 3.0 eq). The reaction mixture was stirred at rt for 2 h. The reaction mixture was then diluted with Et₂O and filtered through neutral alumina, washing with additional Et₂O. The filtrate containing crude (1r,3R,4S)-3,4-difluorocyclopentane-1-carbaldehyde was used in the next step without any other purification.

Preparation of
2-oxaspiro[3.3]heptane-6-carbaldehyde




[0823] To a solution of (2-oxaspiro[3.3]heptan-6-yl)methanol (0.25 g, 1.95 mmol), 1.0 eq) in DCM (10 mL) at 0° C. was added DMP (1.65 g, 3.90 mmol, 2 eq). The reaction mixture was stirred at rt for 1 h. The reaction mixture was then filtered through celite and the filtrate was diluted with NaHCO₃ solution. The mixture was extracted with DCM and the DCM layer was washed with water, brine, dried over Na₂SO₄ and concentrated to give crude 2-oxaspiro[3.3]heptane-6-carbaldehyde (0.1 g) as a colorless oil. The crude material was used without further purification.

[0824] The borate salts in the following table were prepared by the method of potassium (R)-((4-(tert-butoxycarbonyl)-3-methylpiperazin-1-yl)methyl)trifluoroborate using the appropriate commercially available piperazine in step 1.

Structure	Starting material
	(R)-octahydro-2H-pyrido[1,2-a]pyrazine
potassium (R)-trifluoro((octahydro-2H-pyrido[1,2-a]pyrazin-2-yl)methyl)borate	
	(R)-hexahydropyrazino[2,1-c][1,4]oxazin-4(3H)-one
potassium (R)-trifluoro((4-oxohexahydropyrazino[2,1-c][1,4]oxazin-8(1H)-yl)methyl)borate	

Preparation of potassium (S)-((1,1-dioxidohexahydro-5H-isothiazolo[2,3-a]pyrazin-5-yl)methyl)trifluoroborate



Step 1. tert-butyl (R)-4-(methanesulfonyl)-3-(((methanesulfonyl)oxy)methyl)piperazine-1-carboxylate

[0825] To a solution of tert-butyl (R)-3-(hydroxymethyl)piperazine-1-carboxylate (4.00 g, 18.5 mmol) and Et₃N (5.15 mL, 37.0 mmol) in DCM (10 mL) at -20° C. was added methanesulfonyl chloride (3.1 g, 27.7 mmol). The reaction mixture was stirred for 10 min and then diluted with a solution of saturated aqueous NaHCO₃. The mixture was extracted with DCM and the organic layer was dried over MgSO₄, filtered and concentrated. Silica gel column chromatography (eluting with 10% MeOH in DCM) provided tert-butyl (R)-4-(methanesulfonyl)-3-(((methanesulfonyl)oxy)methyl)piperazine-1-carboxylate (3.0 g, 8.1 mmol, 44% yield) as a gummy solid. The crude product was used in the next step without any other purification.

Step 2. tert-butyl (S)-hexahydro-5H-isothiazolo[2,3-a]pyrazine-5-carboxylate 1,1-dioxide

[0826] To a solution of tert-butyl (R)-4-(methanesulfonyl)-3-(((methanesulfonyl)oxy)methyl)piperazine-1-carboxylate (3.0 g, 8.1 mmol) in THF (10 mL) at -78° C. was added a solution of LHMDS (1.0 M in THF, 24.3 mL, 24.3 mmol). The reaction mixture was stirred at -78° C. for 3 h and then diluted with a solution of saturated aqueous NaHCO₃. The mixture was extracted with DCM and the organic layer was dried over MgSO₄, filtered and concentrated. Silica gel column chromatography (eluting with 50% EtOAc in hexane) provided tert-butyl (S)-hexahydro-5H-isothiazolo[2,3-a]pyrazine-5-carboxylate 1,1-dioxide (2.5 g) as a white solid. ¹H NMR (400 MHz, METHANOL-d₄) δ ppm 4.35-4.12 (m, 2H) 3.39-3.37 (m, 1H) 3.28-3.10 (m, 3H) 2.80-2.40 (m, 3H) 2.39-2.36 (m, 1H) 2.03-1.94 (m, 1H) 1.59 (s, 9H).

Step 3: (S)-hexahydro-2H-isothiazolo[2,3-a]pyrazine 1,1-dioxide hydrochloride


[0827] A solution of HCl (4.0 M in dioxane, 3 mL) was added to a solution of tert-butyl (S)-hexahydro-5H-isothiazolo[2,3-a]pyrazine-5-carboxylate 1,1-dioxide (2.5 g, 9.0

mmol) and the mixture was stirred for 2 h at rt. The reaction was then concentrated to give crude (S)-hexahydro-2H-isothiazolo[2,3-a]pyrazine 1,1-dioxide hydrochloride which was used without further purification.

Step 4. potassium (S)-((1,1-dioxidohexahydro-5H-isothiazolo[2,3-a]pyrazin-5-yl)methyl)trifluoroborate



[0828] To a solution of (S)-hexahydro-2H-isothiazolo[2,3-a]pyrazine 1,1-dioxide hydrochloride (1.7 g, 8.0 mmol) in THF (20 mL) was added NaH (0.461 g, 19.3 mmol), potassium (bromomethyl)trifluoroborate (1.9 g, 9.6 mmol) and tetrabutylammonium iodide (0.178 g, 0.482 mmol). The reaction mixture was stirred at rt for 16 h. The reaction mixture was then concentrated to give potassium (S)-((1,1-dioxidohexahydro-5H-isothiazolo[2,3-a]pyrazin-5-yl)methyl)trifluoroborate (3 g, crude) as a white solid.

Preparation of tert-butyl (2S)-4-(bromomethyl)-4-fluoro-2-methylpiperidine-1-carboxylate



[0829] To a solution of tert-butyl (S)-2-methyl-4-methylenepiperidine-1-carboxylate [see Example 71] (400 mg, 1.89 mmol) in DCM (10 mL) at 0° C. was added triethylamine trihydrofluoride (0.77 mL, 4.73 mmol). The reaction mixture was stirred at 0° C. for 30 min and then NBS (500 mg, 2.83 mmol) was added. The mixture was stirred at rt for 2 h. The reaction mixture was then basified with saturated aqueous NaHCO₃ solution and extracted with EtOAc. The organic layer was dried over sodium sulfate, filtered and concentrated. The crude product was purified by silica gel chromatography (eluted with 10% EtOAc/hexane) to give tert-butyl (2S)-4-(bromomethyl)-4-fluoro-2-methylpiperidine-1-carboxylate (0.35 g, 60% yield) as a mixture of diastereomers that was used without further purification.

Preparation of (cis)-4-(bromomethyl)-1-isobutyl-2-(trifluoromethyl)piperidine



Step 1. (cis)-methyl 1-isobutyl-2-(trifluoromethyl)piperidine-4-carboxylate

[0830] Isobutyraldehyde (0.767 g, 10.7 mmol) and triethylamine (3.07 mL, 21.3 mmol) were added to a solution of (cis)-methyl 2-(trifluoromethyl)piperidine-4-carboxylate [see WO2021/158948, 2021, A1] (1.5 g, 7.1 mmol) in DCM (15 mL). The reaction mixture was stirred at rt for 30 min and then sodium triacetoxyborohydride (4.51 g, 21.3 mmol) was added. The reaction mixture was stirred at rt for 4 h and then quenched with a solution of saturated aqueous NaHCO₃ and extracted three times with DCM. The combined organic extracts were washed with brine, dried over Na₂SO₄, filtered and concentrated to give crude (cis)-methyl 1-isobutyl-2-(trifluoromethyl)piperidine-4-carboxylate (1.0 g) which was used without further purification.

Step 2. (cis)-(1-isobutyl-2-(trifluoromethyl)piperidin-4-yl)methanol

[0831] A solution of LAH (2M in THF, 1.02 mL, 2.05 mmol) was added to a solution of (cis)-methyl 1-isobutyl-2-(trifluoromethyl)piperidine-4-carboxylate (0.5 g, 1.87 mmol) in THF (5 mL) at 0° C. The mixture was stirred at 0° C. for 2 h and then quenched with EtOAc. The mixture was washed with a solution of saturated aqueous ammonium chloride and the organic layer was dried over Na₂SO₄, filtered and concentrated to give crude (cis)-(1-isobutyl-2-(trifluoromethyl)piperidin-4-yl)methanol (150 mg) which was used without further purification.



Step 3. (cis)-4-(bromomethyl)-1-isobutyl-2-(trifluoromethyl)piperidine

[0832] Triphenylphosphine (427 mg, 1.62 mmol) and carbon tetrabromide (537 mg, 1.62 mmol) were added in portions to a solution of (cis)-(1-isobutyl-2-(trifluoromethyl)piperidin-4-yl)methanol (130 mg, 0.54 mmol) in DCM (4 mL) at 0° C. The reaction mixture was stirred at rt for 2 h. After completion of the reaction, the mixture was diluted with DCM and washed with water. The organic layer was

washed with brine, dried over Na_2SO_4 , filtered and concentrated. The crude material was purified by silica gel chromatography (eluted with 0-50% EtOAc in hexanes) to afford (cis)-4-(bromomethyl)-1-isobutyl-2-(trifluoromethyl)piperidine (60 mg, 0.20 mmol, 37% yield).

Preparation of Example Compounds

Example 1. Preparation of 1-(5-((1-(cyclohexylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 1)



Step 1: 1-(5-bromopyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione

[0833] A mixture of 3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione (400 mg, 1.51 mmol), 5-bromo-3-iodopyrazolo[1,5-a]pyridine, (499 mg, 1.54 mmol), cesium carbonate (986 mg, 3.03 mmol) and CuI (57.7 mg, 0.303 mmol) in 1,4-dioxane (12 mL) in a microwave vial was purged with nitrogen. ($\pm$)-trans-1,2-Diaminocyclohexane (0.036 mL, 0.30 mmol) was added and the mixture was purged again with nitrogen. The vial was capped and heated at 80° C. overnight in a heating block. The reaction mixture was diluted with ethyl acetate and washed with water and brine. The organic layer was dried over Na_2SO_4 , filtered and concentrated. Silica gel column chromatography (eluting with 0-100% EtOAc in heptane) provided 1-(5-bromopyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione as a light yellow, foamy solid. LCMS $[\text{M}+\text{H}]^+$: 459.2. ^1H NMR (500 MHz, CDCl_3) δ 8.23 (dd, $J=7.3, 0.8$ Hz, 1H), 7.90 (s, 1H), 7.49 (dd, $J=2.1, 0.8$ Hz, 1H), 7.14 (d, $J=8.9$ Hz, 1H), 6.86 (dd, $J=7.3, 2.1$ Hz, 1H), 6.49-6.39 (m, 2H), 5.03 (s, 2H), 3.83 (s, 3H), 3.80 (m, 5H), 2.96 (t, $J=6.6$ Hz, 2H).

Step 2: tert-butyl 4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperidine-1-carboxylate

[0834] a) A microwave vial containing tert-butyl 4-methylenepiperidine-1-carboxylate (500 mg, 2.53 mmol) was purged with nitrogen for 15 min and then a solution of 9-BBN (0.5M in THF, 5.07 mL, 2.53 mmol) was added. The vial was capped and the mixture was heated at 80° C. for 3.5 h and then cooled to rt.

[0835] b) The reaction mixture from part a) was added by syringe to a microwave vial containing a mixture of 1-(5-bromopyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione, (1048 mg, 2.281 mmol), K_2CO_3 (438 mg, 3.17 mmol) and $\text{PdCl}_2(\text{dppf}) \cdot \text{CH}_2\text{Cl}_2$ adduct (54 mg, 0.066 mmol) in DMF (14 mL) and water (1.4 mL). The vial was capped and the reaction

mixture was heated overnight at 60° C. The reaction mixture was then cooled to rt and diluted with ethyl acetate and washed sequentially with water and brine. The organic layer was dried over sodium sulfate, filtered and concentrated. Silica gel column chromatography (eluted with 0-100% EtOAc in heptane) provided tert-butyl 4-((3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperidine-1-carboxylate as a light yellow, foamy solid. LCMS [M+H]⁺: 578.4.

Step 3: 3-(2,4-dimethoxybenzyl)-1-(5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride

[0836] A solution of HCl (4.0 M in dioxane, 15 ml, 60 mmol) was added to tert-butyl 4-((3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperidine-1-carboxylate (1270 mg, 2.154 mmol) and the mixture was stirred for 2 h at rt. The reaction was then concentrated to give crude 3-(2,4-dimethoxybenzyl)-1-(5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride. LCMS [M+H]⁺: 478.4.

Step 4: 1-(5-((1-(cyclohexylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione

[0837] To a solution of 3-(2,4-dimethoxybenzyl)-1-(5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride (100 mg, 0.195 mmol) in DMF (3 mL) was added potassium carbonate (81 mg, 0.58 mmol) and (bromomethyl)cyclohexane (0.081 mL, 0.58 mmol). The mixture was heated at 80° C. for 4 h and then cooled to rt. The mixture was diluted with ethyl acetate and washed sequentially with water and brine. The organic


layer was dried over sodium sulfate, filtered and concentrated to give crude 1-(5-((1-(cyclohexylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione. LCMS [M+H]⁺: 574.4.

Step 5: 1-(5-((1-(cyclohexylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0838] TFA (2 mL, 26 mmol) was added to crude 1-(5-((1-(cyclohexylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione (30 mg, 0.0525 mmol) and the mixture was heated at 80° C. overnight. The mixture was then cooled to rt, concentrated and the residue was dissolved in toluene and concentrated again. The residue was dissolved in DMSO, filtered through a 1 micron filter and purified by reverse phase HPLC using ACN/Water/0.1% formic acid. The fractions containing the product were combined, frozen and lyophilized to afford a formate salt of 1-(5-((1-(cyclohexylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione.

LCMS [M+H]⁺: 424.3. ¹H NMR (500 MHz, DMSO-d₆) δ 10.44 (s, 1H), 8.58 (d, J=7.1 Hz, 1H), 8.16 (s, 1H), 8.00 (s, 1H), 7.37 (d, J=1.8 Hz, 1H), 6.79 (dd, J=7.2, 1.9 Hz, 1H), 3.77 (t, J=6.7 Hz, 2H), 3.598-3.26 (m, 2H), 3.19 (d, J=12.2 Hz, 3H), 2.79 (t, J=6.7 Hz, 2H), 2.60 (d, J=6.8 Hz, 2H), 2.56 (s, 1H), 1.82-1.58 (m, 9H), 1.41 (q, J=12.4 Hz, 2H), 1.30-1.09 (m, 3H), 0.96-0.82 (m, 2H).





[0839] The compounds in the following table were prepared by the method of Example 1, using the appropriate commercially available halide, mesylate, tosylate or triflate in step 4.

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
2	 1-(5-((1-(cyclopentylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	410.3	(500 MHz, DMSO-d ₆) δ 10.45 (d, J = 4.6 Hz, 1H), 8.60 (d, J = 7.1 Hz, 1H), 8.01 (s, 1H), 7.45-7.26 (m, 1H), 6.80 (dd, J = 7.1, 1.9 Hz, 1H), 3.78 (td, J = 6.7, 2.8 Hz, 2H), 3.57-3.39 (m, 2H), 3.30-3.12 (m, 1H), 3.02 (dd, J = 7.2, 5.4 Hz, 2H), 2.87 (dt, J = 14.1, 10.6 Hz, 2H), 2.79 (t, J = 6.7 Hz, 2H), 2.62 (d, J = 6.6 Hz, 2H), 2.20 (h, J = 7.4 Hz, 1H), 1.94-1.70 (m, 4H), 1.62 (tdq, J = 9.7, 6.8, 3.8, 3.1 Hz, 2H), 1.58-1.41 (m, 4H), 1.31-1.11 (m, 2H).




-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
3	 1-(5-((1-((tetrahydro-2H-pyran-4-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	426.3	(500 MHz, DMSO-d ₆) δ 10.45 (d, J = 5.4 Hz, 1H), 8.60 (dd, J = 7.2, 2.6 Hz, 1H), 8.02 (d, J = 2.3 Hz, 1H), 7.42-7.35 (m, 1H), 6.80 (dt, J = 7.2, 2.1 Hz, 1H), 3.85 (ddd, J = 11.6, 4.5, 1.9 Hz, 2H), 3.78 (td, J = 6.7, 3.5 Hz, 2H), 3.55-3.45 (m, 2H), 3.31 (td, J = 11.7, 2.1 Hz, 2H), 2.94 (t, J = 6.2 Hz, 2H), 2.87 (dt, J = 13.9, 10.9 Hz, 2H), 2.79 (t, J = 6.7 Hz, 2H), 2.62 (d, J = 6.6 Hz, 2H), 2.04 (ddt, J = 11.2, 7.3, 3.9 Hz, 1H), 1.92-1.71 (m, 3H), 1.69-1.58 (m, 2H), 1.56-1.42 (m, 2H), 1.22 (qd, J = 12.0, 4.5 Hz, 2H).
4	 1-(5-((1-((4,4-difluorocyclohexyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	460.4	(500 MHz, DMSO-d ₆) δ 10.36 (d, J = 5.1 Hz, 1H), 8.52 (d, J = 7.3 Hz, 1H), 7.93 (s, 1H), 7.29 (d, J = 15.4 Hz, 1H), 6.72 (d, J = 7.1 Hz, 1H), 3.69 (t, J = 6.8 Hz, 2H), 3.14 (s, 1H), 2.88 (t, J = 6.3 Hz, 2H), 2.79 (dd, J = 20.7, 8.9 Hz, 2H), 2.71 (t, J = 6.7 Hz, 2H), 2.54 (d, J = 6.5 Hz, 2H), 2.02-1.90 (m, 3H), 1.83-1.63 (m, 8H), 1.43 (q, J = 12.4, 11.6 Hz, 2H), 1.15 (q, J = 13.1 Hz, 2H).
5	 1-(5-((1-((cyclopropyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	382.2	(500 MHz, DMSO-d ₆) δ 10.45 (d, J = 5.4 Hz, 1H), 8.60 (dd, J = 7.1, 3.0 Hz, 1H), 8.02 (d, J = 2.3 Hz, 1H), 7.39 (d, J = 1.8 Hz, 1H), 6.80 (dd, J = 7.2, 1.9 Hz, 1H), 5.77 (ddt, J = 17.0, 10.3, 6.7 Hz, 1H), 5.31-5.08 (m, 2H), 3.78 (td, J = 6.7, 3.3 Hz, 2H), 3.58-3.38 (m, 2H), 3.25 (d, J = 6.3 Hz, 1H), 3.15-3.03 (m, 2H), 2.96-2.83 (m, 2H), 2.79 (t, J = 6.8 Hz, 2H), 2.61 (d, J = 6.5 Hz, 2H), 2.43 (dtd, J = 9.7, 6.4, 1.6 Hz, 2H), 1.83 (d, J = 13.8 Hz, 2H), 1.43 (q, J = 13.1 Hz, 2H).




-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
6	 1-(5-((1-(2-cyclohexylethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	438.4	(500 MHz, DMSO-d ₆) δ 10.45 (d, J = 5.6 Hz, 1H), 8.68-8.49 (m, 1H), 8.01 (d, J = 2.5 Hz, 1H), 7.46-7.24 (m, 1H), 6.80 (dd, J = 7.1, 1.9 Hz, 1H), 3.78 (t, J = 6.7 Hz, 2H), 3.53-3.41 (m, 2H), 3.28-3.10 (m, 1H), 3.03 (dt, J = 11.2, 5.3 Hz, 2H), 2.90-2.72 (m, 4H), 2.61 (d, J = 6.6 Hz, 2H), 1.83 (t, J = 15.3 Hz, 2H), 1.77-1.57 (m, 5H), 1.57-1.47 (m, 2H), 1.41 (q, J = 13.1 Hz, 2H), 1.18 (ddd, J = 26.1, 22.5, 12.0 Hz, 4H), 0.92 (q, J = 11.9 Hz, 2H).
7	 1-(5-((1-((tetrahydro-2H-pyran-3-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	426.3	¹ H NMR (500 MHz, DMSO-d ₆) δ 10.36 (d, J = 4.0 Hz, 1H), 8.52 (d, J = 7.2 Hz, 1H), 7.93 (d, J = 2.0 Hz, 1H), 7.30 (s, 1H), 6.71 (dd, J = 7.2, 1.9 Hz, 1H), 3.75-3.67 (m, 4H), 3.62 (dt, J = 11.5, 4.2 Hz, 2H), 3.44 (s, 2H), 3.19-3.02 (m, 1H), 2.92-2.74 (m, 3H), 2.71 (t, J = 6.7 Hz, 2H), 2.58-2.51 (m, 2H), 1.92 (s, 1H), 1.72 (d, J = 14.5 Hz, 3H), 1.59-1.47 (m, 1H), 1.41 (dt, J = 13.5, 9.7 Hz, 3H), 1.31-1.14 (m, 1H).
8	 1-(5-((1-(cycloheptylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	438.3	(500 MHz, DMSO-d ₆) δ 10.45 (d, J = 4.7 Hz, 1H), 8.60 (dt, J = 7.2, 1.3 Hz, 1H), 8.02 (d, J = 1.6 Hz, 1H), 7.46-7.28 (m, 1H), 6.80 (dd, J = 7.2, 1.8 Hz, 1H), 3.78 (td, J = 6.7, 2.8 Hz, 2H), 3.47 (d, J = 12.2 Hz, 2H), 3.21 (s, 1H), 2.92-2.83 (m, 3H), 2.79 (td, J = 6.7, 1.8 Hz, 2H), 2.62 (d, J = 6.6 Hz, 2H), 2.00-1.66 (m, 6H), 1.65-1.38 (m, 10H), 1.27-1.09 (m, 2H).
9	 1-(5-((1-(2-methylbutyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	398.3	(500 MHz, DMSO-d ₆) δ 10.44 (d, J = 4.6 Hz, 1H), 8.60 (s, 1H), 8.01 (s, 1H), 7.37 (d, J = 11.6 Hz, 1H), 6.79 (d, J = 7.2 Hz, 1H), 3.77 (t, J = 6.8 Hz, 2H), 3.46 (s, 2H), 3.21 (s, 1H), 3.00-2.81 (m, 3H), 2.79 (t, J = 6.6 Hz, 2H), 2.62 (d, J = 6.5 Hz, 2H), 1.92-1.71 (m, 4H), 1.65-1.33 (m, 3H), 1.16 (dq, J = 14.3, 7.2 Hz, 1H), 0.91 (dp, J = 23.0, 7.3, 6.9 Hz, 6H).





-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
10	 1-(5-((1-(2-cyclohexylpropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	452.3	(500 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 8.59 (d, J = 7.2 Hz, 1H), 8.01 (s, 1H), 7.36 (d, J = 11.5 Hz, 1H), 6.79 (d, J = 7.3 Hz, 1H), 3.77 (t, J = 6.6 Hz, 2H), 3.51 (d, J = 12.2 Hz, 4H), 3.21 (s, 1H), 3.01 (q, J = 6.3, 5.6 Hz, 1H), 2.97-2.70 (m, 4H), 2.62 (d, J = 6.5 Hz, 1H), 1.93-1.68 (m, 5H), 1.67-1.39 (m, 5H), 1.33-0.92 (m, 6H), 0.89 (d, J = 6.8 Hz, 3H).
11	 1-(5-((1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	440.2	(500 MHz, DMSO-d ₆) δ 10.44 (d, J = 5.6 Hz, 1H), 8.59 (d, J = 7.2 Hz, 1H), 8.00 (s, 1H), 7.37 (d, J = 13.5 Hz, 1H), 6.79 (d, J = 7.2 Hz, 1H), 3.88-3.79 (m, 2H), 3.77 (t, J = 6.7 Hz, 2H), 3.27 (q, J = 11.8 Hz, 4H), 3.04 (dt, J = 11.1, 5.4 Hz, 2H), 2.93-2.70 (m, 4H), 2.61 (d, J = 6.5 Hz, 2H), 1.83 (t, J = 14.6 Hz, 3H), 1.56 (d, J = 13.4 Hz, 5H), 1.40 (q, J = 13.0 Hz, 2H), 1.17 (qd, J = 12.2, 4.5 Hz, 2H).
12	 1-(5-((1-(heptan-4-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	426.3	(400 MHz, DMSO) δ 10.41 (s, 1H), 8.65 (s, 1H), 8.58 (d, J = 7.1 Hz, 1H), 7.99 (s, 1H), 7.36 (d, J = 1.8 Hz, 1H), 6.77 (dd, J = 7.2, 1.9 Hz, 1H), 3.76 (t, J = 6.7 Hz, 2H), 3.31 (d, J = 12.0 Hz, 2H), 3.05 (s, 1H), 3.02-2.92 (m, 2H), 2.78 (t, J = 6.7 Hz, 2H), 2.60 (d, J = 6.7 Hz, 2H), 1.92 (s, 1H), 1.84-1.61 (m, 4H), 1.56-1.39 (m, 4H), 1.34 (ddq, J = 13.6, 9.8, 7.0 Hz, 4H), 0.90 (t, J = 7.2 Hz, 6H).





-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
13	 1-(5-((1-(2-cyclobutylethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	410.3	(500 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 8.59 (d, J = 7.2 Hz, 1H), 8.00 (s, 1H), 7.37 (d, J = 11.6 Hz, 1H), 6.79 (d, J = 7.1 Hz, 1H), 3.77 (t, J = 6.7 Hz, 2H), 3.21 (s, 1H), 3.00 (d, J = 8.3 Hz, 1H), 2.94-2.71 (m, 6H), 2.60 (d, J = 6.5 Hz, 2H), 2.23 (dq, J = 15.5, 7.7 Hz, 1H), 2.13-1.97 (m, 2H), 1.83 (dd, J = 17.2, 10.7 Hz, 5H), 1.72 (dd, J = 10.5, 6.2 Hz, 2H), 1.63 (h, J = 9.4 Hz, 2H), 1.40 (q, J = 13.1, 12.7 Hz, 2H).
14	 1-(5-((1-((tetrahydrofuran-3-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	412.3	(500 MHz, Methanol-d ₄) δ 8.34 (dd, J = 7.2, 0.9 Hz, 1H), 7.91 (s, 1H), 7.27 (dd, J = 2.0, 1.0 Hz, 1H), 6.73 (dd, J = 7.2, 1.9 Hz, 1H), 3.84 (dd, J = 8.8, 7.1 Hz, 1H), 3.81-3.74 (m, 3H), 3.66 (dt, J = 8.5, 7.5 Hz, 1H), 3.51 (t, J = 15.4 Hz, 2H), 3.37 (dd, J = 8.8, 6.5 Hz, 1H), 3.11-3.03 (m, 2H), 2.85 (t, J = 12.8 Hz, 2H), 2.79 (t, J = 6.8 Hz, 2H), 2.60 (dd, J = 6.8, 3.7 Hz, 3H), 2.10 (dtd, J = 12.6, 7.8, 4.8 Hz, 1H), 1.89 (dd, J = 29.1, 13.1 Hz, 3H), 1.58 (dq, J = 12.5, 7.6 Hz, 1H), 1.46 (q, J = 13.4 Hz, 2H).
15	 1-(5-((1-((tetrahydrofuran-2-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	412.2	(400 MHz, Methanol-d ₄) δ 8.43 (d, J = 7.2 Hz, 1H), 8.35 (s, 1H), 8.00 (s, 1H), 7.36 (s, 1H), 6.82 (d, J = 7.1 Hz, 1H), 4.25 (ddt, J = 10.0, 7.1, 3.7 Hz, 1H), 3.96-3.84 (m, 3H), 3.81 (q, J = 7.4 Hz, 1H), 3.61 (d, J = 12.4 Hz, 2H), 3.37-3.14 (m, 2H), 3.08 (dd, J = 13.3, 10.4 Hz, 1H), 3.04-2.92 (m, 2H), 2.88 (t, J = 6.8 Hz, 2H), 2.70 (d, J = 7.0 Hz, 2H), 2.21-2.07 (m, 1H), 2.06-1.85 (m, 5H), 1.68-1.48 (m, 3H). NH protons were not observed due to solvent exchange





-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
16	 1-(5-((1-(2-cyclopentylethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	424.3	(500 MHz, DMSO-d ₆) δ 10.44 (d, J = 5.9 Hz, 1H), 8.59 (d, J = 7.2 Hz, 1H), 8.00 (s, 1H), J = 10.9 Hz, 1H), 6.79 (d, J = 7.2 Hz, 1H), 3.77 (t, J = 6.7 Hz, 2H), 3.22 (s, 1H), 2.99 (dt, J = 10.5, 5.0 Hz, 2H), 2.86 (t, J = 11.7 Hz, 2H), 2.82-2.72 (m, 3H), 2.60 (d, J = 6.6 Hz, 2H), 1.90-1.68 (m, 6H), 1.68-1.55 (m, 4H), 1.54-1.36 (m, 4H), 1.10 (d, J = 7.4 Hz, 2H).
17	 1-(5-((1-(pyridin-3-ylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	419.3	(500 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 8.69 (dd, J = 5.1, 1.7 Hz, 2H), 8.59 (d, J = 7.1 Hz, 1H), 8.01 (s, 1H), 7.96 (dt, J = 7.9, 2.0 Hz, 1H), 7.56 (dd, J = 7.9, 4.9 Hz, 1H), 7.37 (s, 1H), 6.79 (dd, J = 7.1, 1.9 Hz, 1H), 4.35 (d, J = 4.0 Hz, 2H), 3.77 (t, J = 6.7 Hz, 2H), 3.39 (d, J = 12.0 Hz, 2H), 2.94 (q, J = 11.5 Hz, 2H), 2.79 (t, J = 6.7 Hz, 2H), 2.60 (d, J = 6.5 Hz, 2H), 2.00-1.69 (m, 3H), 1.51-1.23 (m, 2H).
18	 1-(5-((1-(cyclobutylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	396.3	(500 MHz, DMSO-d ₆) δ 10.45 (d, J = 6.0 Hz, 1H), 8.63-8.55 (m, 1H), 8.01 (d, J = 2.6 Hz, 1H), 7.38 (d, J = 1.8 Hz, 1H), 6.79 (dd, J = 7.2, 1.9 Hz, 1H), 3.77 (t, J = 6.7 Hz, 2H), 3.35 (d, J = 12.0 Hz, 2H), 3.26-3.13 (m, 1H), 3.08 (dd, J = 7.1, 5.1 Hz, 2H), 2.86 (t, J = 11.8 Hz, 3H), 2.75-2.64 (m, 1H), 2.60 (d, J = 6.6 Hz, 2H), 2.07 (dtd, J = 10.6, 6.8, 3.7 Hz, 2H), 1.98-1.69 (m, 7H), 1.41 (q, J = 13.1 Hz, 2H).
19	 1-(5-((1-(isopentyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	398.4	(500 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 8.59 (d, J = 7.2 Hz, 1H), 8.00 (d, J = 2.8 Hz, 1H), 7.37 (d, J = 11.8 Hz, 1H), 6.79 (d, J = 7.1 Hz, 1H), 3.76 (dt, J = 7.0, 4.0 Hz, 2H), 3.27-3.10 (m, 2H), 3.01 (qd, J = 8.4, 5.1, 3.7 Hz, 2H), 2.93-2.67 (m, 4H), 2.68-2.55 (m, 2H), 1.80 (p, J = 20.8, 19.2 Hz, 3H), 1.59 (dt, J = 14.0, 7.9 Hz, 1H), 1.52 (t, J = 8.1 Hz, 2H), 1.40 (q, J = 13.5 Hz, 2H), 0.89 (d, J = 6.5 Hz, 6H).





-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
20	 1-(5-((1-(pentan-3-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	398.4	(400 MHz, DMSO-d ₆) δ 10.42 (s, 1H), 8.59 (d, J = 7.2 Hz, 1H), 8.00 (d, J = 1.7 Hz, 1H), 7.37 (s, 1H), 6.78 (dd, J = 7.1, 1.9 Hz, 1H), 3.77 (t, J = 6.8 Hz, 2H), 3.34 (d, J = 12.0 Hz, 2H), 2.98 (d, J = 11.5 Hz, 3H), 2.78 (td, J = 6.9, 1.9 Hz, 2H), 2.61 (d, J = 6.7 Hz, 2H), 2.00-1.70 (m, 5H), 1.66-1.33 (m, 4H), 0.95 (td, J = 7.4, 1.8 Hz, 6H).
21	 1-(5-((1-(1-(pyridin-3-yl)ethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	433.4	(500 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 8.78-8.65 (m, 2H), 8.58 (d, J = 7.1 Hz, 1H), 8.15-7.94 (m, 2H), 7.71-7.55 (m, 1H), 7.35 (d, J = 1.6 Hz, 1H), 6.76 (dd, J = 7.2, 1.8 Hz, 1H), 4.62 (d, J = 7.9 Hz, 2H), 3.76 (t, J = 6.7 Hz, 2H), 3.61 (d, J = 11.9 Hz, 1H), 3.30 (d, J = 12.1 Hz, 1H), 2.77 (q, J = 11.3, 9.0 Hz, 4H), 2.59 (d, J = 6.3 Hz, 1H), 1.81 (dd, J = 31.0, 11.2 Hz, 3H), 1.67 (d, J = 6.9 Hz, 3H), 1.45 (dt, J = 28.2, 13.6 Hz, 2H).
22	 1-(5-((1-(3-fluorobenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	436.3	(500 MHz, DMSO-d ₆) δ 10.45 (d, J = 14.1 Hz, 8.60 (dd, J = 10.0, 7.1 Hz, 1H), 8.01 (s, 1H), 7.54 (td, J = 8.0, 6.1 Hz, 1H), 7.49-7.26 (m, 4H), 6.79 (td, J = 7.2, 6.3, 1.9 Hz, 1H), 4.30 (d, J = 5.0 Hz, 2H), 3.77 (t, J = 6.7 Hz, 2H), 3.36 (d, J = 12.0 Hz, 2H), 2.91 (q, J = 11.6 Hz, 2H), 2.79 (t, J = 6.7 Hz, 2H), 2.60 (d, J = 6.5 Hz, 2H), 1.95-1.70 (m, 3H), 1.42 (q, J = 13.0 Hz, 2H).
23	 1-(5-((1-(4-fluorobenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	436.3	(500 MHz, DMSO-d ₆) δ 10.42 (s, 1H), 8.57 (d, J = 7.2 Hz, 1H), 7.99 (s, 1H), 7.52 (dd, J = 8.6, 5.4 Hz, 2H), 7.37-7.28 (m, 3H), 6.77 (dd, J = 7.2, 1.9 Hz, 1H), 4.25 (d, J = 5.0 Hz, 2H), 3.75 (q, J = 6.4 Hz, 2H), 3.33 (d, J = 12.1 Hz, 2H), 2.87 (d, J = 12.0 Hz, 2H), 2.77 (t, J = 6.7 Hz, 2H), 2.58 (m, 3H), 1.83-1.73 (m, 2H), 1.44-1.35 (m, 2H).





-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
24	 1-(5-((1-ethylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	356.3	(400 MHz, DMSO-d ₆) δ 10.42 (d, J = 4.5 Hz, 1H), 8.59 (dt, J = 7.1, 1.3 Hz, 1H), 8.00 (s, 1H), 7.45-7.31 (m, 1H), 6.79 (dd, J = 7.2, 1.9 Hz, 1H), 3.77 (t, J = 6.7 Hz, 2H), 3.45 (d, J = 12.2 Hz, 2H), 3.29-3.14 (m, 1H), 3.12-3.00 (m, 2H), 2.94-2.71 (m, 4H), 2.61 (d, J = 6.5 Hz, 2H), 1.82 (d, J = 13.5 Hz, 2H), 1.40 (q, J = 13.0 Hz, 2H), 1.19 (t, J = 7.3 Hz, 3H).
25	 1-(5-((1-(3-(trifluoromethyl)benzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	486.3	(500 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 8.59 (d, J = 7.1 Hz, 1H), 8.01 (s, 1H), 7.91 (s, 1H), 7.86 (d, J = 7.7 Hz, 1H), 7.79 (d, J = 7.7 Hz, 1H), 7.73 (q, J = 7.7, 7.0 Hz, 1H), 7.37 (d, J = 1.8 Hz, 1H), 6.84-6.75 (m, 1H), 4.39 (d, J = 4.8 Hz, 2H), 3.77 (t, J = 6.7 Hz, 2H), 3.37 (d, J = 12.0 Hz, 2H), 2.94 (q, J = 11.5 Hz, 2H), 2.78 (t, J = 6.7 Hz, 2H), 2.61 (d, J = 6.5 Hz, 2H), 2.02-1.66 (m, 3H), 1.42 (q, J = 13.1 Hz, 2H).
26	 1-(5-((1-(1-phenylethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	432.3	(500 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 8.57 (d, J = 7.1 Hz, 1H), 8.00 (s, 1H), 7.61-7.42 (m, 5H), 7.35 (s, 1H), 6.91-6.55 (m, 1H), 4.48 (dt, J = 11.9, 5.9 Hz, 1H), 3.76 (t, J = 6.7 Hz, 2H), 3.64 (d, J = 12.0 Hz, 1H), 3.29-3.10 (m, 1H), 2.87-2.63 (m, 4H), 2.59 (d, J = 6.3 Hz, 2H), 1.95-1.71 (m, 3H), 1.64 (d, J = 6.9 Hz, 3H), 1.44 (dq, J = 41.1, 12.9, 12.2 Hz, 2H).
27	 1-(5-((1-(3-(trifluoromethoxy)benzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	502.3	(500 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 8.59 (d, J = 7.1 Hz, 1H), 8.01 (s, 1H), 7.70-7.58 (m, 1H), 7.60-7.47 (m, 3H), 7.37 (s, 1H), 6.92-6.64 (m, 1H), 4.34 (d, J = 4.9 Hz, 2H), 3.77 (t, J = 6.7 Hz, 2H), 3.36 (d, J = 12.0 Hz, 2H), 2.92 (q, J = 11.6 Hz, 2H), 2.79 (t, J = 6.7 Hz, 2H), 2.60 (d, J = 6.5 Hz, 2H), 1.79 (dd, J = 34.7, 12.8 Hz, 3H), 1.42 (q, J = 13.1 Hz, 2H).





-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
28	 1-(5-((1-(pyridin-4-ylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	419.3	(500 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 8.71 (s, 2H), 8.58 (d, J = 7.2 Hz, 1H), 8.00 (s, 1H), 7.51 (d, J = 5.1 Hz, 2H), 7.36 (s, 1H), 6.78 (dd, J = 7.1, 1.9 Hz, 1H), 4.32 (d, J = 4.7 Hz, 2H), 3.76 (t, J = 6.7 Hz, 2H), 3.37 (d, J = 11.9 Hz, 2H), 2.94 (d, J = 12.0 Hz, 2H), 2.78 (t, J = 6.7 Hz, 2H), 2.60 (d, J = 6.5 Hz, 2H), 1.82 (d, J = 14.8 Hz, 3H), 1.41 (q, J = 13.1 Hz, 2H).
29	 1-(5-((1-benzylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	418.3	(400 MHz, DMSO-d ₆) δ 10.42 (d, J = 12.8 Hz, 1H), 8.63-8.52 (m, 1H), 8.00 (d, J = 6.5 Hz, 1H), 7.47 (s, 5H), 7.41-7.32 (m, 1H), 6.85-6.74 (m, 1H), 4.26 (d, J = 5.0 Hz, 2H), 3.76 (t, J = 6.7 Hz, 2H), 3.35 (d, J = 12.1 Hz, 2H), 2.90 (q, J = 11.7 Hz, 2H), 2.78 (t, J = 6.7 Hz, 2H), 2.59 (d, J = 6.4 Hz, 2H), 1.78 (dd, J = 25.2, 11.7 Hz, 3H), 1.41 (q, J = 13.1 Hz, 2H).
30	 1-(5-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	410.3	(500 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 8.57 (dd, J = 7.1, 0.9 Hz, 1H), 8.00 (s, 1H), 7.36 (dd, J = 1.9, 0.9 Hz, 1H), 6.79 (dd, J = 7.2, 1.9 Hz, 1H), 4.18 (d, 2H), 3.77 (t, J = 6.7 Hz, 2H), 3.11 (d, J = 62.3 Hz, 3H), 2.79 (t, J = 6.7 Hz, 2H), 2.59 (d, J = 6.7 Hz, 3H), 2.56 (s, 1H), 1.67 (d, J = 13.8 Hz, 3H), 1.36 (d, J = 13.1 Hz, 2H).
31	 1-(5-((1-(2,2,3,3-tetrafluoropropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	442.2	(500 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 8.56 (d, J = 7.4 Hz, 1H), 7.99 (s, 1H), 7.36 (s, 1H), 6.78 (dd, J = 7.1, 1.9 Hz, 1H), 6.50 (t, J = 52.4 Hz, 1H), 3.77 (t, J = 6.7 Hz, 2H), 3.59 (s, 4H), 2.93 (s, 2H), 2.79 (t, J = 6.7 Hz, 2H), 2.57 (d, J = 9.4 Hz, 2H), 2.45-2.05 (m, 1H), 1.61 (s, 3H), 1.37 (d, J = 56.0 Hz, 2H).




-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
32	 1-(5-((1-(3,3-difluorocyclobutyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	432.1	As formate salt-(400 MHz, CD ₃ OD) δ 8.44 (brs, 1H), 8.42 (d, J = 6.8 Hz, 1H), 7.99 (s, 1H), 7.36 (s, 1H), 6.82 (d, J = 7.2 Hz, 1H), 3.87 (t, J = 6.8 Hz, 2H), 3.45-3.42 (m, 2H), 3.20-3.19 (m, 2H), 2.89-2.40 (m, 1H), 1.92-1.88 (m, 3H), 1.55-1.53 (m, 2H) ppm. NH protons not observed due to solvent exchange
33	 1-(5-((1-(5-fluoropyridin-3-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	437.4	(500 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 8.71 (d, J = 2.8 Hz, 1H), 8.68-8.50 (m, 2H), 8.01 (s, 1H), 7.89 (ddd, J = 9.6, 2.8, 1.7 Hz, 1H), 7.37 (d, J = 1.8 Hz, 1H), 6.79 (dd, J = 7.1, 1.9 Hz, 1H), 4.38 (d, J = 3.3 Hz, 2H), 3.77 (t, J = 6.7 Hz, 2H), 3.41 (d, J = 12.0 Hz, 2H), 3.01-2.87 (m, 2H), 2.79 (t, J = 6.7 Hz, 2H), 2.61 (d, J = 6.5 Hz, 2H), 1.98-1.69 (m, 3H), 1.41 (q, J = 13.1 Hz, 2H).
34	 1-(5-((1-(2-methoxyethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	386.3	(500 MHz, DMSO-d ₆) δ 10.45 (d, J = 4.9 Hz, 1H), 8.60 (dd, J = 7.1, 3.2 Hz, 1H), 8.01 (d, J = 2.3 Hz, 1H), 7.42-7.32 (m, 1H), 6.80 (dd, J = 7.2, 1.9 Hz, 1H), 3.78 (td, J = 6.7, 2.8 Hz, 2H), 3.65 (dt, J = 10.0, 5.0 Hz, 2H), 3.47 (d, J = 12.0 Hz, 2H), 3.32 (d, J = 11.6 Hz, 3H), 3.24 (q, J = 5.1 Hz, 2H), 2.91 (q, J = 11.7 Hz, 2H), 2.79 (t, J = 6.7 Hz, 2H), 2.60 (d, J = 6.7 Hz, 2H), 1.81 (q, J = 12.7, 8.1 Hz, 3H), 1.48 (q, J = 12.9 Hz, 2H).
35	 1-(5-((1-(2-hydroxyethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	372.3	(500 MHz, DMSO-d ₆) δ 10.45 (d, J = 5.1 Hz, 1H), 9.10 (s, 1H), 8.60 (dd, J = 7.2, 2.8 Hz, 1H), 8.01 (d, J = 2.2 Hz, 1H), 7.38 (d, J = 12.6 Hz, 1H), 6.80 (dd, J = 7.1, 1.9 Hz, 1H), 3.78 (t, J = 6.7 Hz, 2H), 3.72 (t, J = 5.3 Hz, 2H), 3.49 (d, J = 12.1 Hz, 2H), 3.24 (dd, J = 13.4, 7.7 Hz, 1H), 3.11 (q, J = 5.3 Hz, 2H), 2.90 (t, J = 11.7 Hz, 1H), 2.79 (td, J = 6.8, 2.2 Hz, 2H), 2.60 (d, J = 6.7 Hz, 2H), 1.94-1.66 (m, 3H), 1.49 (q, J = 13.1, 12.4 Hz, 2H).





-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
36	 1-(5-((1-((5-methylpyridin-3-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	433.2	(500 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 8.59 (d, J = 7.1 Hz, 1H), 8.55 (d, J = 2.1 Hz, 1H), 8.50 (d, J = 2.0 Hz, 1H), 8.01 (s, 1H), 7.87-7.69 (m, 1H), 7.43-7.25 (m, 1H), 6.79 (dd, J = 7.2, 1.9 Hz, 1H), 4.30 (d, J = 3.6 Hz, 2H), 3.77 (t, J = 6.7 Hz, 2H), 3.38 (d, J = 12.0 Hz, 2H), 2.93 (d, J = 11.2 Hz, 2H), 2.79 (t, J = 6.7 Hz, 2H), 2.60 (d, J = 6.5 Hz, 2H), 2.36 (s, 3H), 1.82 (q, J = 18.9, 17.6 Hz, 3H), 1.41 (q, J = 13.1 Hz, 2H).
37	 1-(5-((1-isopropylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	370.3	(500 MHz, DMSO-d ₆) δ 10.45 (s, 1H), 8.60 (d, J = 7.0 Hz, 1H), 8.01 (s, 1H), 7.39 (d, J = 1.8 Hz, 1H), 6.79 (dd, J = 7.3, 1.9 Hz, 1H), 3.78 (t, J = 6.7 Hz, 2H), 3.34 (d, J = 12.4 Hz, 2H), 3.26-3.18 (m, 1H), 2.92 (dt, J = (t, J = 6.8 Hz, 2H), 2.62 (d, J = 6.7 Hz, 2H), 1.97-1.78 (m, 3H), 1.45 (qd, J = 13.6, 3.7 Hz, 2H), 1.22 (d, J = 6.6 Hz, 6H).
38	 1-(5-((1-(2-fluorobenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	436.3	(500 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 8.59 (d, J = 7.1 Hz, 1H), 8.01 (s, 1H), 7.57 (dtd, J = 13.7, 7.5, 1.8 Hz, 2H), 7.41-7.28 (m, 3H), 6.85-6.75 (m, 1H), 4.33 (d, J = 4.8 Hz, 2H), 3.77 (t, J = 6.7 Hz, 2H), 3.40 (d, J = 12.0 Hz, 2H), 2.99 (d, J = 11.8 Hz, 2H), 2.79 (q, J = 6.5 Hz, 2H), 2.60 (d, J = 6.6 Hz, 2H), 1.82 (q, J = 20.3, 18.4 Hz, 3H), 1.43 (d, J = 13.0 Hz, 2H).
39	 1-(5-((1-(2-fluoro-2-methylpropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	402.2	(400 MHz, DMSO-d ₆) δ 10.42 (s, 1H), 8.59 (d, J = 7.3 Hz, 1H), 8.00 (s, 1H), 7.36 (d, J = 8.6 Hz, 1H), 6.78 (d, J = 7.2 Hz, 1H), 3.77 (t, J = 6.7 Hz, 2H), 3.54 (s, 2H), 3.39 (d, J = 25.2 Hz, 2H), 3.01 (d, J = 11.9 Hz, 2H), 2.79 (t, J = 6.9 Hz, 2H), 2.65 (dd, J = 37.4, 7.6 Hz, 2H), 2.00-1.69 (m, 3H), 1.69-1.52 (m, 2H), 1.46 (dd, J = 21.5, 10.4 Hz, 6H).





-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
39a	 1-(5-((1-(2-methylallyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione Isolated during synthesis of Example 39	382.1	(500 MHz, DMSO-d ₆) δ 10.45 (d, J = 5.0 Hz, 1H), 8.60 (dt, J = 7.2, 0.9 Hz, 1H), 8.02 (d, J = 2.3 Hz, 1H), 7.43-7.27 (m, 1H), 6.80 (dt, J = 7.2, 1.8 Hz, 1H), 5.32-5.21 (m, 1H), 5.16 (d, J = 1.7 Hz, 1H), 3.83-3.74 (m, 2H), 3.66 (d, J = 5.6 Hz, 1H), 3.52 (d, J = 12.1 Hz, 1H), 3.46-3.33 (m, 2H), 3.21-2.96 (m, 1H), 2.87 (t, J = 11.9 Hz, 1H), 2.79 (t, J = 6.7 Hz, 2H), 2.67-2.59 (m, 2H), 1.91-1.72 (m, 4H), 1.60 (td, J = 27.4, 25.3, 11.8 Hz, 1H), 1.47 (dd, J = 21.5, 12.9 Hz, 3H).
40	 1-(5-((1-(3-methoxybenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	448.4	(500 MHz, DMSO-d ₆) δ 10.45 (d, J = 14.0 Hz, 1H), 8.60 (dd, J = 9.9, 7.1 Hz, 1H), 8.01 (d, J = 7.3 Hz, 1H), 7.56-7.30 (m, 2H), 7.19-6.96 (m, 3H), 6.80 (ddd, J = 12.4, 7.2, 1.9 Hz, 1H), 4.24 (d, J = 5.2 Hz, 2H), 3.78 (d, J = 12.4 Hz, 5H), 3.35 (d, J = 12.1 Hz, 2H), 2.91 (t, J = 11.8 Hz, 2H), 2.78 (t, J = 6.6 Hz, 2H), 2.60 (d, J = 6.5 Hz, 2H), 1.93-1.6 (m, 3H), 1.42 (q, J = 13.1 Hz, 2H).
41	 1-(5-((1-(2-methylbenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	432.3	(500 MHz, DMSO-d ₆) δ 10.45 (d, J = 16.5 Hz, 1H), 8.60 (dd, J = 11.1, 7.1 Hz, 1H), 8.02 (d, J = 8.6 Hz, 1H), 7.48 (dd, J = 15.1, 7.7 Hz, 1H), 7.43-7.21 (m, 4H), 6.87-6.73 (m, 1H), 4.28 (d, J = 5.3 Hz, 2H), 3.77 (t, J = 6.7 Hz, 2H), 3.37 (d, J = 12.0 Hz, 2H), 3.05 (q, J = 11.6 Hz, 2H), 2.79 (q, J = 6.8 Hz, 2H), 2.61 (d, J = 6.6 Hz, 2H), 2.39 (s, 3H), 1.96-1.67 (m, 3H), 1.45 (q, J = 12.9 Hz, 2H).




-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
42	 1-(5-((1-(3-fluoro-2-methylbenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	450.4	(500 MHz, DMSO-d ₆) δ 10.44 (d, J = 16.3 Hz, 1H), 8.59 (d, J = 7.6 Hz, 1H), 8.00 (d, J = 3.3 Hz, 1H), 7.54-7.21 (m, 4H), 6.78 (d, J = 7.6 Hz, 1H), 4.33 (t, J = 4.3 Hz, 2H), 3.77 (d, J = 8.0 Hz, 3H), 3.12-2.93 (m, 1H), 2.78 (d, J = 8.0 Hz, 2H), 2.61 (s, 2H), 2.35 (d, J = 17.2 Hz, 1H), 2.28 (s, 3H), 2.08 (d, J = 3.2 Hz, 1H), 1.96-1.67 (m, 3H), 1.42 (d, J = 13.4 Hz, 2H).
43	 1-(5-((1-(2,2-difluoroethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	392.3	(500 MHz, Methanol-d ₄) δ 8.34 (d, J = 7.2 Hz, 1H), 7.91 (s, 1H), 7.27 (t, J = 1.3 Hz, 1H), 6.73 (dd, J = 7.1, 1.9 Hz, 1H), 6.30 (tt, J = 53.4, 3.6 Hz, 1H), 3.79 (t, J = 6.8 Hz, 2H), 3.57 (q, J = 14.1, 13.3 Hz, 4H), 3.06 (s, 2H), 2.79 (t, J = 6.7 Hz, 2H), 2.61 (d, J = 6.9 Hz, 2H), 1.87 (q, J = 13.3, 8.7 Hz, 3H), 1.52 (d, J = 13.6 Hz, 2H).
44	 1-(5-((1-(3,5-difluorobenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	454.3	(500 MHz, DMSO-d ₆) δ 10.44 (d, J = 12.0 Hz, 1H), 8.59 (t, J = 7.9 Hz, 1H), 8.07-7.94 (m, 1H), 7.40 (t, J = 8.7 Hz, 1H), 7.36 (s, 1H), 7.28 (d, J = 7.0 Hz, 2H), 6.78 (d, J = 7.3 Hz, 1H), 4.29 (d, J = 5.0 Hz, 2H), 3.76 (t, J = 6.6 Hz, 2H), 3.18 (s, 1H), 2.89 (q, J = 11.4, 8.6 Hz, 2H), 2.78 (t, J = 6.7 Hz, 3H), 2.65-2.57 (m, 2H), 1.81 (q, J = 21.9, 19.6 Hz, 3H), 1.43 (q, J = 13.1 Hz, 2H).
45	 1-(5-((1-(3,4-difluorobenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	454.3	(500 MHz, DMSO-d ₆) δ 10.44 (d, J = 13.2 Hz, 1H), 8.58 (d, J = 7.3 Hz, 1H), 8.01 (d, J = 7.1 Hz, 1H), 7.58 (dt, J = 17.9, 8.9 Hz, 2H), 7.36 (s, 2H), 6.78 (d, J = 7.5 Hz, 1H), 4.27 (d, J = 5.0 Hz, 2H), 3.76 (t, J = 6.7 Hz, 2H), 3.34 (s, 2H), 2.89 (d, J = 11.7 Hz, 2H), 2.78 (t, J = 6.6 Hz, 2H), 2.66-2.57 (m, 2H), 1.98-1.69 (m, 3H), 1.50-1.29 (m, 2H).





-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
46	 1-(5-((1-((6-methylpyridin-3-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	433.2	(500 MHz, DMSO-d ₆) δ 10.42 (s, 1H), 8.66-8.46 (m, 2H), 8.00 (s, 1H), 7.82 (dd, J = 8.0, 2.3 Hz, 1H), 7.49-7.30 (m, 2H), 6.87-6.58 (m, 1H), 4.28 (d, J = 4.5 Hz, 2H), 3.76 (t, J = 6.7 Hz, 2H), 3.36 (d, J = 13.0 Hz, 3H), 3.18 (s, 2H), 2.89 (dd, J = 14.6, 8.6 Hz, 2H), 2.78 (t, J = 6.7 Hz, 2H), 2.59 (d, J = 6.5 Hz, 2H), 1.80 (dt, J = 32.7, 15.1 Hz, 3H), 1.38 (q, J = 12.2 Hz, 2H).
47	 1-(5-((1-(4-methylbenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	432.3	(500 MHz, DMSO-d ₆) δ 10.44 (d, J = 15.3 Hz, 1H), 9.18 (s, 1H), 8.59 (dd, J = 11.1, 7.2 Hz, 1H), 8.00 (d, J = 7.9 Hz, 1H), 7.41-7.35 (m, 2H), 7.28 (d, J = 7.8 Hz, 2H), 6.84-6.71 (m, 1H), 4.21 (d, J = 5.0 Hz, 2H), 3.76 (t, J = 6.7 Hz, 2H), 3.33 (d, J = 12.1 Hz, 2H), 2.87 (q, J = 11.7 Hz, 2H), 2.78 (q, J = 6.8 Hz, 2H), 2.59 (d, J = 6.5 Hz, 2H), 2.37-2.31 (m, 3H), 1.92-1.64 (m, 3H), 1.39 (q, J = 12.8 Hz, 2H).
48	 1-(5-((1-(1-(3-fluorophenyl)ethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	450.2	(500 MHz, DMSO-d ₆) δ 10.42 (s, 1H), 8.58 (dd, J = 11.5, 7.2 Hz, 1H), 8.00 (d, J = 7.0 Hz, 1H), 7.54 (dq, J = 14.8, 7.3 Hz, 1H), 7.46-7.24 (m, 4H), 6.85-6.60 (m, 1H), 4.56-4.46 (m, 1H), 3.76 (d, J = 6.7 Hz, 2H), 3.62 (d, J = 12.1 Hz, 1H), 3.22 (d, J = 13.1 Hz, 1H), 2.83-2.63 (m, 4H), 2.58 (d, J = 6.3 Hz, 2H), 1.92-1.70 (m, 3H), 1.62 (d, J = 6.9 Hz, 3H), 1.43 (dt, J = 40.7, 12.9 Hz, 2H).
49	 1-(5-((1-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	342.2	(500 MHz, DMSO-d ₆) δ 10.45 (d, J = 7.0 Hz, 1H), 8.60 (dt, J = 7.2, 2.1 Hz, 1H), 8.01 (d, J = 3.3 Hz, 1H), 7.55-7.08 (m, 1H), 6.80 (dd, J = 7.1, 1.9 Hz, 1H), 3.77 (t, J = 6.7 Hz, 2H), 3.40 (d, J = 12.1 Hz, 2H), 2.94-2.77 (m, 4H), 2.74 (d, J = 4.8 Hz, 3H), 2.60 (d, J = 6.5 Hz, 2H), 1.92-1.68 (m, 3H), 1.39 (q, J = 12.4, 11.7 Hz, 2H).





-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
50	 1-(5-((1-(2,4-difluorobenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	454.4	(500 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 8.59 (d, J = 7.1 Hz, 1H), 8.01 (s, 1H), 7.64 (td, J = 8.6, 6.5 Hz, 1H), 7.47-7.38 (m, 1H), 7.37 (s, 1H), 7.26 (td, J = 8.5, 2.7 Hz, 1H), 6.78 (dd, J = 7.1, 1.9 Hz, 1H), 4.31 (d, J = 4.6 Hz, 2H), 3.77 (t, J = 6.7 Hz, 2H), 3.39 (d, J = 11.8 Hz, 2H), 2.97 (q, J = 11.8 Hz, 2H), 2.79 (t, J = 6.7 Hz, 2H), 2.60 (d, J = 6.6 Hz, 2H), 1.82 (q, J = 18.9, 16.6 Hz, 3H), 1.57-1.31 (m, 2H)
51	 1-(5-((1-(3-fluoro-5-(trifluoromethyl)benzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	504.4	(500 MHz, DMSO-d ₆) δ 10.36 (d, J = 11.7 Hz, 1H), 8.60-8.41 (m, 1H), 7.92 (d, J = 2.9 Hz, 1H), 7.90-7.56 (m, 3H), 7.29 (s, 1H), 6.70 (d, J = 7.6 Hz, 1H), 4.31 (s, 2H), 3.69 (d, J = 7.8 Hz, 2H), 3.12 (d, J = 15.6 Hz, 1H), 2.84 (t, J = 11.8 Hz, 2H), 2.69 (d, J = 7.6 Hz, 2H), 2.59-2.48 (m, 2H), 1.86-1.56 (m, 4H), 1.33 (q, J = 13.4, 12.9 Hz, 2H).
52	 1-(5-((1-(2-oxo-2-(piperidin-1-ylethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	453.4	(400 MHz, CD ₃ OD) δ 8.43 (d, J = 7.2 Hz, 1H), 8.31 (s, 2H), 8.00 (s, 1H), 7.36 (s, 1H), 7.683 (d, J = 7.6 Hz, 1H), 4.10 (s, 2H), 3.88 (t, J = 6.4 Hz, 2H), 3.57-3.50 (m, 4H), 3.37-3.34 (m, 2H), 2.96-2.86 (m, 4H), 2.70 (d, J = 7.2 Hz, 2H), 1.93-1.89 (m, 3H), 1.69-1.56 (m, 7H).



-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
53	 1-(5-((1-(2,3-difluorobenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	454.4	(500 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 8.58 (d, J = 7.2 Hz, 1H), 8.00 (s, 1H), 7.68-7.55 (m, 1H), 7.37 (d, J = 16.6 Hz, 3H), 6.83-6.72 (m, 1H), 4.37 (s, 2H), 3.76 (t, J = 6.7 Hz, 2H), 3.42 (s, 3H), 3.00 (d, J = 11.6 Hz, 1H), 2.78 (t, J = 6.6 Hz, 2H), 2.59 (d, J = 6.5 Hz, 2H), 1.82 (d, J = 14.7 Hz, 3H), 1.41 (d, J = 13.0 Hz, 2H).
54	 1-(5-((1-((6-(trifluoromethyl)pyridin-3-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	487.1	(500 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 8.87 (d, J = 2.1 Hz, 1H), 8.59 (d, J = 7.1 Hz, 1H), 8.22 (dd, J = 8.1, 2.1 Hz, 1H), 8.06 (d, J = 8.1 Hz, 1H), 8.01 (s, 1H), 7.46-7.21 (m, 1H), 6.79 (dd, J = 7.2, 1.9 Hz, 1H), 4.45 (s, 2H), 3.77 (t, J = 6.7 Hz, 2H), 3.42 (d, J = 11.9 Hz, 2H), 2.96 (s, 2H), 2.79 (t, J = 6.7 Hz, 2H), 2.60 (d, J = 6.6 Hz, 2H), 1.98-1.73 (m, 3H), 1.53-1.21 (m, 2H).
55	 1-(5-((1-((1-(trifluoromethyl)cyclopropyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	450.1	(400 MHz, DMSO-d ₆) δ 10.42 (s, 1H), 8.59 (d, J = 7.1 Hz, 1H), 8.00 (d, J = 1.7 Hz, 1H), 7.38 (s, 1H), 6.87-6.47 (m, 1H), 3.77 (t, J = 6.7 Hz, 2H), 3.62-3.35 (m, 4H), 3.29 (s, 1H), 2.94 (d, J = 11.5 Hz, 1H), 2.78 (t, J = 6.7 Hz, 2H), 2.61 (d, J = 6.1 Hz, 2H), 1.92-1.71 (m, 3H), 1.52 (t, J = 13.3 Hz, 2H), 1.16 (d, J = 42.2 Hz, 4H).
56	 1-(5-((1-(4-(trifluoromethoxy)benzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	502.3	(500 MHz, DMSO-d ₆) δ 10.45 (d, J = 15.5 Hz, 1H), 8.59 (d, J = 7.1 Hz, 1H), 8.01 (s, 1H), 7.73-7.59 (m, 2H), 7.50 (d, J = 8.2 Hz, 2H), 7.37 (s, 1H), 6.87-6.71 (m, 1H), 4.32 (d, J = 5.0 Hz, 2H), 3.77 (t, J = 6.7 Hz, 2H), 3.36 (d, J = 12.0 Hz, 2H), 3.00-2.89 (m, 2H), 2.79 (t, J = 6.7 Hz, 2H), 2.60 (d, J = 6.5 Hz, 2H), 1.94-1.68 (m, 3H), 1.41 (q, J = 13.1 Hz, 2H).

-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
57	 1-(5-((1-(4-(trifluoromethyl)benzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	486.3	(500 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 8.59 (d, J = 7.1 Hz, 1H), 8.01 (s, 1H), 7.87 (d, J = 8.0 Hz, 2H), 7.81-7.69 (m, 2H), 7.37 (s, 1H), 6.89-6.68 (m, 1H), 4.39 (d, J = 4.8 Hz, 2H), 3.77 (t, J = 6.7 Hz, 2H), 3.37 (d, J = 12.0 Hz, 2H), 2.93 (q, J = 11.7 Hz, 2H), 2.78 (t, J = 6.7 Hz, 2H), 2.60 (d, J = 6.5 Hz, 2H), 1.96-1.30 (m, 3H), 1.52-1.30 (m, 2H).
58	 1-(5-((1-(5-fluoro-2-methylbenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	450.4	(500 MHz, DMSO-d ₆) δ 10.44 (d, J = 13.8 Hz, 1H), 8.59 (d, J = 7.7 Hz, 1H), 8.00 (d, J = 2.9 Hz, 1H), 7.35 (d, J = 10.7 Hz, 3H), 7.22 (d, J = 9.0 Hz, 1H), 6.78 (d, J = 7.2 Hz, 1H), 4.28 (s, 2H), 3.76 (t, J = 6.8 Hz, 2H), 3.05 (d, J = 12.0 Hz, 3H), 2.78 (d, J = 7.5 Hz, 2H), 2.61 (d, J = 6.5 Hz, 2H), 2.34 (s, 3H), 2.08 (d, J = 3.0 Hz, 1H), 1.81 (q, J = 18.9, 17.5 Hz, 3H), 1.56-1.27 (m, 2H).
59	 1-(5-((1-(2-chlorobenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	452.4	(500 MHz, DMSO-d ₆) δ 10.44 (d, J = 13.7 Hz, 1H), 8.58 (d, J = 7.8 Hz, 1H), 8.00 (d, J = 2.3 Hz, 1H), 7.64 (dd, J = 31.9, 7.8 Hz, 2H), 7.51 (q, J = 8.4 Hz, 2H), 7.35 (s, 1H), 6.88-6.61 (m, 1H), 4.40 (d, J = 4.9 Hz, 2H), 3.77 (d, J = 7.7 Hz, 3H), 3.08 (d, J = 11.7 Hz, 2H), 2.78 (d, J = 7.2 Hz, 2H), 2.69-2.55 (m, 2H), 2.08 (d, J = 2.7 Hz, 1H), 1.95-1.76 (m, 3H), 1.53-1.33 (m, 2H).
60	 1-(5-((1-(thiazol-2-ylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	425.4	(500 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 8.59 (d, J = 7.1 Hz, 1H), 8.00 (d, J = 4.0 Hz, 2H), 7.95 (d, J = 3.2 Hz, 1H), 7.38 (d, J = 1.4 Hz, 1H), 6.79 (dd, J = 7.2, 1.9 Hz, 1H), 4.72 (s, 2H), 3.77 (t, J = 6.7 Hz, 2H), 3.51 (d, J = 12.1 Hz, 2H), 3.02 (s, 2H), 2.79 (t, J = 6.7 Hz, 2H), 2.60 (d, J = 6.4 Hz, 2H), 1.83 (d, J = 14.2 Hz, 3H), 1.47 (d, J = 13.5 Hz, 2H).


-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
61	 1-(5-((1-(4-methoxybenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	448.2	(500 MHz, DMSO-d ₆) δ 10.45 (d, J = 16.0 Hz, 1H), 8.60 (dd, J = 11.3, 7.2 Hz, 1H), 8.01 (d, J = 8.0 Hz, 1H), 7.42-7.35 (m, 3H), 7.03 (dq, J = 9.5, 7.7, 7.1 Hz, 3H), 4.20 (d, J = 5.2 Hz, 2H), 3.82-3.74 (m, 4H), 3.34 (d, J = 12.0 Hz, 2H), 2.86 (q, J = 12.8, 12.0 Hz, 2H), 2.78 (t, J = 6.8 Hz, 2H), 2.60 (d, J = 6.4 Hz, 2H), 1.87-1.71 (m, 3H), 1.50-1.18 (m, 3H).
62	 1-(5-((1-(3,3,3-trifluoro-2,2-dimethylpropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	452.0	(400 MHz, CD ₃ OD) δ 8.44 (d, J = 7.2 Hz, 1H), 8.01 (s, 1H), 7.38 (s, 1H), 6.84 (d, J = 7.2 Hz, 1H), 3.89 (t, J = 6.8 Hz, 2H), 3.65 (s, 1H), 3.48 (s, 1H), 3.37-3.33 (m, 2H), 3.16-3.13 (m, 2H), 2.89 (t, J = 6.8 Hz, 2H), 2.77-2.69 (m, 2H), 2.01-1.90 (m, 3H), 1.70-1.67 (m, 2H), 1.36 (s, 6H) ppm. NH proton not observed due to solvent exchange.


Example 63. Preparation of 1-(5-((1-(piperidin-4-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 63)

2.94-2.82 (m, 4H), 2.79 (t, J=6.7 Hz, 2H), 2.62 (d, J=6.6 Hz, 2H), 2.09 (dt, J=7.7, 3.8 Hz, 1H), 1.84 (dq, J=29.1, 16.0, 14.5 Hz, 5H), 1.50 (q, J=13.1 Hz, 2H), 1.41-1.28 (in, 2H).

Examples 64 and 65. Preparation of 1-(5-((1-(((1*r*,4*r*)-4-methoxycyclohexyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 64) and 1-(5-((1-(((1*s*,4*s*)-4-methoxycyclohexyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 65)




[0840] Prepared by the method of Example 1 using tert-butyl 4-(bromomethyl)piperidine-1-carboxylate in step 4 in place of (bromomethyl)cyclohexane. LCMS [M+H]⁺: 425.2. ¹H NMR (500 MHz, DMSO-d₆) δ 10.45 (d, J=6.6 Hz, 1H), 8.61 (dd, J=7.2, 3.9 Hz, 1H), 8.40 (d, J=11.9 Hz, 1H), 8.02 (d, J=3.2 Hz, 1H), 7.47-7.35 (m, 1H), 6.80 (dd, J=7.2, 1.9 Hz, 1H), 3.78 (td, J=6.7, 4.0 Hz, 2H), 3.51 (d, J=11.8 Hz, 2H), 3.30 (d, J=12.8 Hz, 2H), 2.98 (t, J=6.1 Hz, 2H),



-continued

Example 65



[0841] Prepared by the method of Example 1 using a commercially available mixture of cis and trans 1-(bromomethyl)-4-methoxycyclohexane in step 4 in place of (bromomethyl)cyclohexane. The stereoisomers were purified after step 5 by reverse-phase HPLC (eluting with using ACN/Water/0.1% TFA).


Example 64. 1-(5-(((1r,4r)-4-methoxycyclohexyl)methyl)piperidin-4-yl)methylpyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

[0842] Eluted first, minor isomer. LCMS $[M+H]^+$: 454.3. 1H NMR (500 MHz, DMSO- d_6) δ 10.45 (d, $J=4.6$ Hz, 1H), 8.60 (d, $J=7.1$ Hz, 1H), 8.01 (s, 1H), 7.47-7.27 (m, 1H), 6.80 (dd, $J=7.1, 1.9$ Hz, 1H), 3.78 (td, $J=6.7, 3.0$ Hz, 2H), 3.47 (d, $J=12.1$ Hz, 2H), 3.24 (d, $J=5.2$ Hz, 4H), 3.05 (ddt, $J=16.9, 10.7, 5.3$ Hz, 1H), 2.93-2.83 (m, 3H), 2.79 (t, $J=6.7$ Hz, 2H), 2.62 (d, $J=6.6$ Hz, 2H), 2.00 (d, $J=12.3$ Hz, 2H), 1.92-1.63 (m, 6H), 1.49 (q, $J=13.1$ Hz, 2H), 1.19-1.07 (m, 2H), 0.98 (q, $J=12.9$ Hz, 2H).

Example 65. 1-(5-(((1s,4s)-4-methoxycyclohexyl)methyl)piperidin-4-yl)methylpyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

[0843] Eluted second, major isomer. LCMS $[M+H]^+$: 454.3. 1H NMR (500 MHz, Methanol- d_4) δ 8.45 (d, $J=7.1$ Hz, 1H), 8.02 (s, 1H), 7.49-7.25 (m, 1H), 6.85 (dd, $J=7.1, 1.8$ Hz, 1H), 3.90 (t, $J=6.8$ Hz, 2H), 3.61-3.44 (m, 3H), 3.35 (s, 3H), 2.92 (dt, $J=13.6, 6.9$ Hz, 6H), 2.71 (d, $J=7.1$ Hz, 2H), 2.09-1.82 (m, 6H), 1.70-1.46 (m, 6H), 1.44-1.31 (m, 2H).


Example 66. Preparation of 1-(5-(((1R,5S)-8-(cyclohexylmethyl)-8-azabicyclo[3.2.1]octan-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[0844] Prepared by the method of Example 1, steps 2-5 using tert-butyl (1R,5S)-3-methylene-8-azabicyclo[3.2.1]octane-8-carboxylate in step 2 in place of tert-butyl 4-meth-


ylenepiperidine-1-carboxylate. LCMS $[M+H]^+$: 450.4. 1H NMR (400 MHz, DMSO- d_6) δ 10.43 (s, 1H), 8.53-8.52 (m, 1H), 8.29-8.20 (m, 1H), 7.97 (d, $J=2.4$ Hz, 1H), 7.42-7.29 (m, 1H), 6.77 (d, $J=7.2$ Hz, 1H), 3.75-3.60 (m, 2H), 3.23 (br s, 2H), 2.84-2.70 (m, 3H), 2.24 (br s, 2H), 2.11-1.82 (m, 4H), 1.81-1.56 (m, 6H), 1.56-1.37 (m, 4H), 1.33-1.09 (m, 4H), 0.91-0.78 (m, 2H).

Example 67. Preparation of 1-(5-(((1R,5S)-8-isobutyl-8-azabicyclo[3.2.1]octan-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[0845] Prepared by the method of Example 1, steps 2-5 using tert-butyl (1R,5S)-3-methylene-8-azabicyclo[3.2.1]octane-8-carboxylate in step 2 in place of tert-butyl 4-methylenepiperidine-1-carboxylate, and 1-iodo-2-methylpropane in step 4 in place of (bromomethyl)cyclohexane. LCMS $[M+H]^+$: 410.3. 1H NMR (400 MHz, $CDCl_3$) ppm 11.33 (s, 1H), 8.40 (m, $J=6.8$ Hz, 1H), 7.94 (s, 1H), 7.75 (s, 1H), 6.67 (m, $J=7.3, 11.7$ Hz, 1H), 3.93-3.84 (m, 4H), 3.19 (s, 1H), 2.97-2.89 (m, 3H), 2.78-2.71 (m, 4H), 2.42-2.25 (m, 2H), 2.15 (s, 1H), 1.92 (m, $J=8.6$ Hz, 2H), 1.68 (s, 3H), 1.14 (m, $J=6.4$ Hz, 6H).


Example 68. Preparation of 1-(5-(((1R,5S)-8-(pyridin-3-ylmethyl)-8-azabicyclo[3.2.1]octan-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[0846] Prepared by the method of Example 1, steps 2-5 using tert-butyl (1R,5S)-3-methylene-8-azabicyclo[3.2.1]octane-8-carboxylate in step 2 in place of tert-butyl 4-methylenepiperidine-1-carboxylate, and 3-(bromomethyl)pyridine in step 4 in place of (bromomethyl)cyclohexane. LCMS $[M+H]^+$: 445.3. 1H NMR (400 MHz, METHANOL- d_4) ppm=8.54 (s, 1H), 8.46-8.34 (m, 2H), 8.00-7.96 (m, 1H), 7.89 (m, $J=7.8$ Hz, 1H), 7.44-7.30 (m, 2H), 6.81 (d, $J=2.1, 7.1$ Hz, 1H), 3.88 (dt, $J=4.6, 6.7$ Hz, 2H), 3.60 (d, $J=4.4$ Hz, 2H), 3.18 (s, 2H), 2.89 (m, 3H), 2.58 (d, $J=7.2$ Hz,

1H), 2.23-2.17 (m, 1H), 2.10-2.03 (m, 2H), 1.94-1.81 (m, 1H), 1.68-1.60 (m, 1H), 1.49 (d, J=2.7, 8.7 Hz, 2H), 1.41 (m, J=13.9 Hz, 1H).

Example 69. Preparation of 1-(5-((1-(4-methoxypyrimidin-2-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



Example 69

Step 1. 3-(2,4-dimethoxybenzyl)-1-(5-((1-(4-methoxypyrimidin-2-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[0847] To a stirred solution of 3-(2,4-dimethoxybenzyl)-1-(5-((piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride (100 mg, 0.209 mmol) and 2-chloro-4-methoxypyrimidine (30.2 mg,

0.209 mmol) in MeCN (1 mL) was added DIPEA (54 mL, 0.418 mmol). The mixture was stirred for 2 h at 120° C. After completion, the reaction was cooled to rt and diluted with EtOAc and water. The organic layer was dried over Na₂SO₄ and concentrate. The crude compound was purified by silica gel chromatography (eluting with 10% MeOH in DCM) to afford 3-(2,4-dimethoxybenzyl)-1-(5-((1-(4-methoxypyrimidin-2-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (90 mg, 0.153 mmol, 90% yield). LCMS [M+H]⁺: 586.3.

Step 2. 1-(5-((1-(4-methoxypyrimidin-2-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


[0848] To a stirred solution of 3-(2,4-dimethoxybenzyl)-1-(5-((1-(4-methoxypyrimidin-2-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (90 mg, 0.145 mmol) in TFA (1 mL) was added TfOH (0.1 mL) and the reaction mixture was stirred at 70° C. for 2 h. After completion, the reaction was concentrate. The crude compound was purified by PREP HPLC using: Mobile Phase: A=0.1% HCOOH in water, B=Acetonitrile, Column: JUPITER Phenomenex (250 mm×21.2 mm), 5.0 μm, Flow: 20 mL/min. the collected fraction were concentrated under reduced pressure to afford 1-(5-((1-(4-methoxypyrimidin-2-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (25 mg, 0.053 mmol, 27% yield) as an off-white solid. LCMS [M+H]⁺: 436.2; HPLC: Rt=4.794 min. ¹H NMR (400 MHz, Methanol-d₄) δ 8.43 (d, J=7.1 Hz, 1H), 8.00 (s, 1H), 7.96 (d, J=6.4 Hz, 1H), 7.37 (s, 1H), 6.85 (dd, J=7.1, 1.8 Hz, 1H), 6.21 (d, J=6.5 Hz, 1H), 4.56 (d, J=9.6 Hz, 2H), 3.89 (t, J=6.8 Hz, 2H), 3.05 (t, J=12.9 Hz, 2H), 2.89 (t, J=6.8 Hz, 2H), 2.68 (d, J=7.3 Hz, 2H), 2.10-1.97 (m, 1H), 1.83 (d, J=13.2 Hz, 2H), 1.33 (qd, J=12.3, 3.3 Hz, 3H), NH proton not observed due to solvent exchange.

Example 70. Preparation of 1-(5-((1-(3-methylbutan-2-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



See Example 1,
step 2 and 5
step 2

-continued



Example 70


Step 1.

1-(3-methylbutan-2-yl)-4-methylenepiperidine


[0849] To a solution of 4-methylenepiperidine hydrochloride (2.0 g, 15 mmol) in DCM (20 mL) was added TEA (6.25 mL 44.9 mmol), TiCl_4 (0.8 mL 7.48 mmol) and 3-methylbutan-2-one (1.4 g, 16.5 mmol). The mixture was stirred at rt for 12 h and then NaBH_3CN (2.8 g, 45 mmol) was added. The reaction was stirred for 4 h at rt. After completion, the mixture was diluted with DCM and washed with water. The organic layer was dried over Na_2SO_4 , filtered and concentrated to obtain 1-(3-methylbutan-2-yl)-4-methylenepiperidine (0.4 g, crude). ^1H NMR (300 MHz, CDCl_3) δ 4.60 (s, 2H), 2.61-2.53 (m, 2H), 2.36-2.31 (m, 2H), 2.25-2.09 (m, 4H), 1.65-1.59 (m, 1H), 1.11-1.06 (m, 3H), 0.96-0.92 (m, 3H), 0.89-0.85 (m, 3H).

[0850] Step 2. 1-(5-(((1-(3-methylbutan-2-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 70) was prepared from 1-(3-methylbutan-2-yl)-4-methylenepiperidine using the method of Example 1, steps 2 and 5, wherein 1-(3-methylbutan-2-yl)-4-methylenepiperidine was used in place of tert-butyl 4-methylenepiperidine-1-carboxylate. LCMS $[\text{M}+\text{H}]^+$: 398.3. ^1H NMR (300 MHz, CD_3OD) δ 8.42 (d, $J=7.2$ Hz, 1H), 8.00 (s, 1H), 7.35 (s, 1H), 6.82 (dd, $J=7.2$ Hz, 1.2 Hz, 1H), 3.87 (t, $J=6.4$ Hz, 2H), 3.58-3.47 (m, 2H), 3.13-2.96 (m, 3H), 2.87 (t, $J=6.8$ Hz, 2H), 2.69 (d, $J=6.4$ Hz, 2H), 2.24-2.20 (m, 1H), 1.98-1.94 (m, 3H), 1.59-1.55 (m, 2H), 1.27 (d, $J=6.8$ Hz, 3H), 1.04 (d, $J=6.8$ Hz, 3H), 0.98 (d, $J=6.8$ Hz, 3H), NH proton not observed due to solvent exchange.

Example 71. Preparation of 1-(5-(((2S,4S)-1-(cyclohexylmethyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



-continued



Example 71

Step 1. tert-butyl




(S)-2-methyl-4-methylenepiperidine-1-carboxylate

[0851] To dry t-BuOK (1.58 g, 14.1 mmol) in THF (20 mL) was added methyltriphenylphosphonium bromide (5.02 g, 14.07 mmol) at 0°C ., then the mixture was stirred at rt for 2 h. The mixture was cooled to 0°C . and a solution of tert-butyl (S)-2-methyl-4-oxopiperidine-1-carboxylate (2 g, 9.38 mmol) in THF (5 mL) was added slowly. The reaction mixture was stirred at rt for 14 h. The reaction mixture was quenched with a solution of saturated aqueous NH_4Cl (50 mL) and extracted with EtOAc (2 $\times$). The combined organic layers were concentrated to give the crude product. The crude product was purified by flash silica gel chromatography (eluted with 0-10% EtOAc/petroleum ether) to give tert-butyl (S)-2-methyl-4-methylenepiperidine-1-carboxylate (1.7 g, 8.1 mmol, 86% yield) as a yellow oil. ^1H NMR (400 MHz, CDCl_3) δ 4.85 (d, $J=1.6$ Hz, 1H), 4.74 (d, $J=1.6$ Hz, 1H), 4.51-4.48 (m, 1H), 4.04-4.01 (m, 1H), 2.89-2.82 (m, 1H), 2.42-2.37 (m, 1H), 2.17-2.13 (m, 2H), 2.03-2.00 (m, 1H), 1.47 (s, 9H), 1.07 (d, $J=6.8$ Hz, 3H).

Step 2. 1-(5-(((2S,4S)-1-(cyclohexylmethyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione





[0852] Example 71 was prepared from tert-butyl (S)-2-methyl-4-methylenepiperidine-1-carboxylate using the method of Example 1, steps 2 to 5, wherein tert-butyl (S)-2-methyl-4-methylenepiperidine-1-carboxylate was used in place of tert-butyl 4-methylenepiperidine-1-carboxylate. The final product contained a minor amount of the trans isomer which was removed by SFC purification: Column: Chiralpak IG-3 50 $\times$ 4.6 mm I.D., 3 μm Mobile phase: Phase A for CO_2 , and Phase B for IPA (0.05% DEA); Gradient elution: 40% IPA (0.05% DEA) in CO_2 Flow rate: 3 mL/min; Detector: PDA Column Temp: 35°C .; Back Pressure: 100 Bar. Product is peak 1, retention time 3.1 min. LCMS $[\text{M}+\text{H}]^+$: 438.3. ^1H NMR (400 MHz, $\text{METHANOL}-d_4$) δ 8.40 (d, $J=7.2$ Hz, 1H), 7.98 (s, 1H), 7.33 (s, 1H), 6.81-6.79 (m, 1H), 3.89-3.86 (m, 2H), 3.04 (br d, $J=12.0$ Hz, 1H), 2.89-2.87 (m, 2H), 2.66-2.56 (m, 3H), 2.20-1.84 (m, 4H), 1.75-1.44 (m, 8H), 1.37-1.11 (m, 5H), 1.10-1.03 (m, 3H), 0.99-0.81 (m, 2H).

[0853] The compounds in the following table were prepared by the method of Example 71, using the appropriate commercially available halide in the alkylation step.


Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
72	 1-(5-(((2S,4S)-1-isobutyl-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	398.3	(400 MHz, CD ₃ OD) δ 8.44 (d, J = 7.2 Hz, 1H), 8.37 (s, 1H), 8.01 (s, 1H), 7.37 (s, 1H), 6.84 (dd, J = 6.8 Hz, 1.2 Hz, 1H), 3.89 (t, J = 6.4 Hz, 2H), 3.62-3.60 (m, 1H), 3.20-3.14 (m, 3H), 2.92-2.87 (m, 3H), 2.84-2.79 (m, 1H), 2.70-2.66 (m, 3H), 2.10-2.05 (m, 2H), 1.93-1.90 (m, 2H), 1.55-1.45 (m, 2H), 1.37 (d, J = 6.4 Hz, 1H), 1.07-1.02 (m, 6H).
73	 1-(5-(((2S,4S)-2-methyl-1-((tetrahydro-2H-pyran-4-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	440.1	(400 MHz, CD ₃ OD) δ 8.44 (d, J = 7.2 Hz, 1H), 8.29 (s, 1H), 8.01 (s, 1H), 7.37 (s, 1H), 6.84 (dd, J = 7.2 Hz, 1.6 Hz, 1H), 3.98-3.94 (m, 2H), 3.89 (t, J = 6.4 Hz, 2H), 3.68-3.62 (m, 1H), 3.48-3.42 (m, 2H), 3.28-3.24 (m, 2H), 3.02-2.98 (m, 1H), 2.90-2.84 (m, 3H), 2.70-2.66 (m, 2H), 2.08-2.06 (m, 2H), 1.94-1.90 (m, 2H), 1.75-1.73 (m, 1H), 1.65-1.63 (m, 1H), 1.51-1.36 (m, 7H).
74	 1-(5-(((2S,4S)-1-((4,4-difluorocyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	474.2	(400 MHz, CD ₃ OD) δ 8.43 (d, J = 6.4 Hz, 2H), 7.99 (s, 1H), 7.35 (s, 1H), 6.82 (dd, J = 6.8 Hz, 1.6 Hz, 1H), 3.89 (t, J = 6.8 Hz, 2H), 3.62-3.58 (m, 1H), 3.24-3.20 (m, 2H), 2.98-2.85 (m, 4H), 2.68 (d, J = 6.8 Hz, 1H), 2.07-2.04 (m, 3H), 1.91-1.81 (m, 7H), 1.42-1.33 (m, 7H).

[0854] The compounds in the following table were prepared using the method of Example 71, wherein tert-butyl (R)-2-methyl-4-oxopiperidine-1-carboxylate was used in

place of tert-butyl (S)-2-methyl-4-oxopiperidine-1-carboxylate and, using the appropriate commercially available halide in the alkylation step.


Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
75	 1-(5-(((2R,4R)-2-methyl-1-((tetrahydro-2H-pyran-4-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	440.2	(400 MHz, Methanol-d ₄) δ 8.43 (d, J = 7.0 Hz, 1H), 8.35 (s, 1H), 8.00 (s, 1H), 7.36 (s, 1H), 6.82 (d, J = 7.1 Hz, 1H), 3.95 (dd, J = 11.8, 4.3 Hz, 2H), 3.88 (t, J = 6.7 Hz, 2H), 3.64 (s, 1H), 3.50-3.37 (m, 3H), 3.02 (d, J = 82.6 Hz, 5H), 2.88 (t, J = 6.8 Hz, 2H), 2.69 (d, J = 7.1 Hz, 2H), 1.99 (d, J = 53.6 Hz, 4H), 1.68 (dd, J = 37.8, 12.9 Hz, 2H), 1.56-1.31 (m, 8H). NH proton not observed due to solvent exchange
76	 1-(5-(((2R,4R)-1-(cyclohexylmethyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	438.2	(400 MHz, Methanol-d ₄) δ 8.43 (d, J = 7.2 Hz, 1H), 8.01 (s, 1H), 7.36 (d, J = 1.7 Hz, 1H), 6.83 (dd, J = 7.3, 1.9 Hz, 1H), 3.89 (t, J = 6.8 Hz, 2H), 3.59 (d, J = 12.6 Hz, 1H), 3.27-3.09 (m, 2H), 2.89 (t, J = 6.7 Hz, 3H), 2.77 (dd, J = 13.3, 5.2 Hz, 1H), 2.72-2.61 (m, 3H), 2.05 (ddq, J = 11.9, 7.6, 3.8 Hz, 1H), 1.98-1.65 (m, 7H), 1.64-1.16 (m, 8H), 1.05 (dd, J = 16.5, 7.2 Hz, 2H). NH proton not observed due to solvent exchange
77	 1-(5-(((2R,4R)-1-isobutyl-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	398.2	(400 MHz, CD ₃ OD) δ 8.50 (s, 1H), 8.43 (d, J = 7.2 Hz, 1H), 7.99 (s, 1H), 7.34 (s, 1H), 6.81 (dd, J = 7.2, 1.6 Hz, 1H), 3.87 (t, J = 7.2 Hz, 2H), 3.52-3.48 (m, 1H), 3.34 (m, 1H), 3.08-3.02 (m, 1H), 2.87 (t, J = 7.2 Hz, 2H), 2.80-2.74 (m, 1H), 2.67 (d, J = 6.8 Hz, 2H), 2.04-1.84 (m, 4H), 1.50-1.37 (m, 3H), 1.30 (d, J = 6.0 Hz, 3H), 1.01 (t, J = 6.4 Hz, 6H).
78	 1-(5-(((2R,4R)-1-((4,4-difluorocyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	474.4	(400 MHz, Methanol-d ₄) δ 8.40 (d, J = 7.1 Hz, 1H), 7.98 (s, 1H), 7.34 (s, 1H), 6.81 (dd, J = 7.3, 1.8 Hz, 1H), 3.88 (t, J = 6.7 Hz, 2H), 3.06 (d, J = 11.6 Hz, 1H), 2.89 (t, J = 6.8 Hz, 2H), 2.65 (d, J = 11.8 Hz, 1H), 2.60 (d, J = 7.1 Hz, 2H), 2.19 (t, J = 7.5 Hz, 1H), 2.12-1.90 (m, 5H), 1.84-1.54 (m, 6H), 1.40-1.11 (m, 5H), 1.08 (d, J = 6.3 Hz, 3H).

Example 79. Preparation of 1-(5-((1-isobutyl-2,2-dimethylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0855] Prepared using the method of Example 71 wherein tert-butyl 2,2-dimethyl-4-oxopiperidine-1-carboxylate was used in place of tert-butyl (S)-2-methyl-4-oxopiperidine-1-carboxylate. LCMS $[M+H]^+$: 412.6. 1H NMR (400 MHz, CD_3OD) δ 8.44 (d, $J=7.2$ Hz, 1H), 8.39 (s, 1H), 8.02 (s, 1H), 7.36 (s, 1H), 6.85-6.83 (m, 1H), 3.90 (t, $J=7.2$ Hz, 2H), 3.53-3.50 (m, 1H), 3.22-3.13 (m, 2H), 2.89 (t, $J=6.8$ Hz, 2H), 2.67-2.65 (m, 3H), 2.21 (brs, 1H), 2.03-1.82 (m, 3H), 1.65-1.52 (m, 2H), 1.43 (s, 3H), 1.36 (s, 3H), 1.10-1.06 (m, 6H).

Example 80. Preparation of 1-(5-((1-((cyclohexylmethyl)sulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



-continued






Example 80

Step 1: 1-(5-((1-((cyclohexylmethyl)sulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione




[0856] DIPEA (0.042 mL, 0.24 mmol) and cyclohexylmethanesulfonyl chloride (14 mg, 0.073 mmol) were added to a solution of 3-(2,4-dimethoxybenzyl)-1-(5-((1-((cyclohexylmethyl)sulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride (25 mg, 0.049 mmol) in DCM (1.5 mL) at rt. The mixture was stirred at rt for 1 h, then diluted with DCM and washed sequentially with water and brine. The organic layer was dried over sodium sulfate, filtered and concentrated to give crude 1-(5-((1-((cyclohexylmethyl)sulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione (31 mg, 0.049 mmol). LCMS $[M+H]^+$: 638.4.

[0857] Step 2: 1-(5-((1-((cyclohexylmethyl)sulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 80) was prepared from 1-(5-((1-((cyclohexylmethyl)sulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione by the method of Example 1, step 5, wherein 1-(5-((1-((cyclohexylmethyl)sulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione was used in place of 1-(5-((1-((cyclohexylmethyl)sulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione. LCMS $[M+H]^+$: 488.3. 1H NMR (500 MHz, $DMSO-d_6$) δ 10.44 (s, 1H), 8.57 (d, $J=7.1$ Hz, 1H), 8.00 (s, 1H), 7.38 (d, $J=1.7$ Hz, 1H), 6.80 (dd, $J=7.2, 1.9$ Hz, 1H), 3.77 (t, $J=6.7$ Hz, 2H), 3.56 (d, $J=12.2$ Hz, 2H), 2.85 (d, $J=6.2$ Hz, 2H), 2.79 (t, $J=6.7$ Hz, 2H), 2.70 (td, $J=12.1, 2.4$ Hz, 2H), 2.60 (d, $J=6.9$ Hz, 2H), 1.92-1.78 (m, 3H), 1.79-1.64 (m, 5H), 1.59 (dd, $J=10.3, 6.4$ Hz, 1H), 1.31-1.19 (m, 4H), 1.18-1.00 (m, 3H).




[0858] The compounds in the following table were prepared by the method of Example 80, using the appropriate commercially available sulfonyl chloride or chloroformate in step 1.

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
81	 1-(5-((1-(cyclohexylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	474.3	(500 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 8.56 (d, J = 7.0 Hz, 1H), 7.99 (s, 1H), 7.38 (t, J = 1.2 Hz, 1H), 6.79 (dd, J = 7.2, 1.9 Hz, 1H), 3.77 (t, J = 6.7 Hz, 2H), 3.70-3.49 (m, 2H), 3.06 (tt, J = 11.7, 3.4 Hz, 1H), 2.92-2.71 (m, 4H), 2.59 (d, J = 7.2 Hz, 2H), 2.09-1.92 (m, 2H), 1.88-1.72 (m, 3H), 1.72-1.55 (m, 3H), 1.43-0.96 (m, 7H).
82	 1-(5-((1-(cyclopentylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	460.4	(500 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 8.57 (d, J = 7.0 Hz, 1H), 7.99 (s, 1H), 7.37 (s, 1H), 6.79 (dd, J = 7.2, 1.9 Hz, 1H), 3.77 (t, J = 6.7 Hz, 2H), 3.62 (d, J = 13.3 Hz, 2H), 3.58 (d, J = 7.9 Hz, 1H), 2.82-2.74 (m, 4H), 2.60 (d, J = 7.1 Hz, 2H), 1.99-1.87 (m, 2H), 1.77 (dt, J = 14.1, 7.4 Hz, 3H), 1.67 (d, J = 13.0 Hz, 4H), 1.55 (t, J = 6.0 Hz, 2H), 1.21 (dt, J = 21.9, 10.9 Hz, 2H).
83	 1-(5-((1-((tetrahydro-2H-pyran-4-yl)sulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	476.3	(500 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 8.57 (d, J = 7.1 Hz, 1H), 7.99 (s, 1H), 7.43-7.30 (m, 1H), 6.79 (dd, J = 7.2, 1.9 Hz, 1H), 3.94-3.88 (m, 2H), 3.77 (t, J = 6.7 Hz, 2H), 3.63 (d, J = 12.5 Hz, 2H), 3.39 (tt, J = 12.0, 3.8 Hz, 1H), 3.33 (td, J = 11.8, 2.0 Hz, 2H), 2.92-2.83 (m, 2H), 2.79 (t, J = 6.7 Hz, 2H), 2.60 (d, J = 7.1 Hz, 2H), 1.92-1.71 (m, 3H), 1.62 (dtd, J = 29.5, 13.0, 12.4, 4.1 Hz, 4H), 1.32-1.08 (m, 2H).




-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
84	 1-(5-((1-(cyclopropylmethyl)sulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	446.3	(500 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 8.57 (d, J = 7.1 Hz, 1H), 7.99 (s, 1H), 7.38 (d, J = 1.8 Hz, 1H), 6.80 (dd, J = 7.1, 1.8 Hz, 1H), 3.77 (t, J = 6.7 Hz, 2H), 3.61 (d, J = 12.3 Hz, 2H), 2.97 (d, J = 7.1 Hz, 2H), 2.85-2.71 (m, 4H), 2.60 (d, J = 7.0 Hz, 2H), 1.90-1.59 (m, 3H), 1.22 (qd, J = 12.3, 4.2 Hz, 2H), 0.98 (dq, J = 15.1, 7.6, 4.8 Hz, 1H), 0.65-0.53 (m, 2H), 0.42-0.19 (m, 2H).
85	 1-(5-((1-(isopropylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	434.3	(500 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 8.56 (dd, J = 7.1, 0.9 Hz, 1H), 7.99 (s, 1H), 7.38 (d, J = 1.7 Hz, 1H), 6.79 (dd, J = 7.2, 1.9 Hz, 1H), 3.77 (t, J = 6.7 Hz, 2H), 3.63 (d, J = 12.5 Hz, 2H), 3.29 (p, J = 6.8 Hz, 1H), 2.87-2.76 (m, 4H), 2.60 (d, J = 7.2 Hz, 2H), 1.77 (t, J = 4.8 Hz, 1H), 1.66 (d, J = 13.1 Hz, 2H), 1.21 (d, J = 6.8 Hz, 8H).
86	 1-(5-((1-(sec-butylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	448.3	(500 MHz, Methanol-d ₄) δ 8.31 (dd, J = 7.2, 0.9 Hz, 1H), 7.88 (s, 1H), 7.26 (dd, J = 1.9, 0.9 Hz, 1H), 6.73 (dd, J = 7.2, 1.9 Hz, 1H), 3.78 (t, J = 6.8 Hz, 2H), 3.71-3.62 (m, 2H), 2.93 (ddd, J = 9.3, 6.8, 3.9 Hz, 1H), 2.84-2.73 (m, 4H), 2.55 (d, J = 7.3 Hz, 2H), 1.84 (dtd, J = 15.2, 7.6, 4.0 Hz, 1H), 1.73 (ddt, J = 11.4, 7.6, 3.8 Hz, 1H), 1.63 (d, J = 13.3 Hz, 2H), 1.40 (ddd, J = 13.7, 9.4, 7.4 Hz, 1H), 1.18 (d, J = 6.9 Hz, 5H), 0.91 (t, J = 7.5 Hz, 3H).




-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
87	 1-(5-((1-(isobutylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	448.3	(500 MHz, DMSO-d ₆) δ 10.42 (s, 1H), 8.56 (d, J = 7.1 Hz, 1H), 7.99 (d, J = 1.7 Hz, 1H), 7.37 (s, 1H), 6.79 (d, J = 7.2 Hz, 1H), 3.76 (t, J = 6.7 Hz, 2H), 3.65-3.57 (m, 2H), 2.85 (d, J = 6.6 Hz, 2H), 2.78 (t, J = 6.7 Hz, 2H), 2.70 (t, J = 11.9 Hz, 2H), 2.59 (d, J = 6.9 Hz, 2H), 2.09 (hept, J = 6.6 Hz, 1H), 1.80-1.62 (m, 3H), 1.30-1.16 (m, 2H), 1.02 (dd, J = 6.8, 1.7 Hz, 6H).
88	 1-(5-((1-(benzylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	482.3	(500 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 8.56 (d, J = 7.1 Hz, 1H), 8.00 (s, 1H), 7.50-7.26 (m, 6H), 6.78 (dd, J = 7.2, 1.9 Hz, 1H), 4.37 (s, 2H), 3.77 (t, J = 6.7 Hz, 2H), 3.54 (d, J = 12.4 Hz, 2H), 2.79 (t, J = 6.7 Hz, 2H), 2.67 (td, J = 12.4, 2.5 Hz, 2H), 2.57 (d, J = 7.1 Hz, 2H), 1.70 (tt, J = 7.4, 3.6 Hz, 1H), 1.63 (d, J = 13.2 Hz, 2H), 1.16 (qd, J = 12.2, 4.2 Hz, 2H).
89	 1-(5-((1-(cyclopropylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	432.3	(500 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 8.57 (d, J = 7.1 Hz, 1H), 8.00 (s, 1H), 7.46-7.27 (m, 1H), 6.80 (dd, J = 7.1, 1.9 Hz, 1H), 3.78 (t, J = 6.7 Hz, 2H), 3.59 (d, J = 12.2 Hz, 2H), 2.84-2.73 (m, 4H), 2.61 (d, J = 7.0 Hz, 2H), 2.59-2.54 (m, 1H), 1.85-1.59 (m, 3H), 1.25 (qd, J = 12.3, 4.0 Hz, 2H), 1.06-0.82 (m, 4H).


-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
90	 1-(5-((1-(methylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	406.3	(400 MHz, DMSO-d ₆) δ 10.41 (s, 1H), 8.56 (dd, J = 7.1, 0.9 Hz, 1H), 7.99 (s, 1H), 7.37 (dd, J = 1.9, 0.9 Hz, 1H), 6.79 (dd, J = 7.1, 1.9 Hz, 1H), 3.77 (t, J = 6.7 Hz, 2H), 3.54 (d, J = 11.6 Hz, 2H), 2.83 (s, 3H), 2.78 (t, J = 6.7 Hz, 2H), 2.71-2.57 (m, 4H), 1.77-1.64 (m, 3H), 1.27 (t, J = 11.8 Hz, 2H).
91	 1-(5-((1-(phenylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	468.3	(500 MHz, DMSO-d ₆) δ 10.42 (s, 1H), 8.52 (dd, J = 7.1, 0.9 Hz, 1H), 7.98 (s, 1H), 7.74-7.68 (m, 3H), 7.66-7.60 (m, 2H), 7.32 (t, J = 1.4 Hz, 1H), 6.72 (dd, J = 7.2, 1.9 Hz, 1H), 3.75 (t, J = 6.7 Hz, 2H), 3.65 (d, J = 12.0 Hz, 2H), 2.77 (t, J = 6.7 Hz, 2H), 2.56-2.53 (m, 2H), 2.19 (td, J = 12.0, 2.3 Hz, 2H), 1.77-1.42 (m, 3H), 1.24 (td, J = 12.3, 11.9, 4.0 Hz, 2H).
92	 1-(5-((1-((2-methoxyethyl)sulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	450.2	(500 MHz, Methanol-d ₄) δ 8.31 (dd, J = 7.2, 0.9 Hz, 1H), 7.88 (s, 1H), 7.26 (dd, J = 1.9, 0.9 Hz, 1H), 6.73 (dd, J = 7.2, 1.9 Hz, 1H), 3.78 (t, J = 6.8 Hz, 2H), 3.65-3.54 (m, 4H), 3.24 (s, 3H), 3.14 (t, J = 5.9 Hz, 2H), 2.79 (t, J = 6.8 Hz, 2H), 2.69 (td, J = 12.3, 2.4 Hz, 2H), 2.59-2.53 (m, 2H), 1.76-1.61 (m, 3H), 1.28-1.17 (m, 2H).


-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
93	 methyl 4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperidine-1-carboxylate	386.3	(400 MHz, DMSO-d ₆) δ 10.41 (s, 1H), 8.54 (dd, J = 7.1, 0.9 Hz, 1H), 7.98 (s, 1H), 7.35 (dd, J = 1.9, 1.0 Hz, 1H), 6.78 (dd, J = 7.1, 1.9 Hz, 1H), 3.94 (d, J = 13.0 Hz, 2H), 3.76 (t, J = 6.7 Hz, 2H), 3.57 (s, 3H), 2.78 (t, J = 6.7 Hz, 4H), 2.56 (d, J = 7.2 Hz, 2H), 1.85-1.76 (m, 1H), 1.59 (d, J = 13.0 Hz, 2H), 1.08 (qd, J = 12.3, 4.3 Hz, 2H).
94	 phenyl 4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperidine-1-carboxylate	448.3	(500 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 8.58 (dd, J = 7.1, 0.9 Hz, 1H), 8.00 (s, 1H), 7.46-7.35 (m, 3H), 7.27-7.19 (m, 1H), 7.15-7.06 (m, 2H), 6.82 (dd, J = 7.2, 1.9 Hz, 1H), 4.21-3.98 (m, 2H), 3.78 (t, J = 6.7 Hz, 2H), 2.99 (s, 1H), 2.80 (t, J = 6.7 Hz, 3H), 2.63 (d, J = 7.2 Hz, 2H), 1.88 (tt, J = 7.4, 3.7 Hz, 1H), 1.73-1.65 (m, 2H), 1.24 (s, 2H).
95	 isobutyl 4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperidine-1-carboxylate	428.3	(500 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 8.56 (d, J = 7.1 Hz, 1H), 7.99 (s, 1H), 7.36 (d, J = 1.8 Hz, 1H), 6.79 (dd, J = 7.1, 1.8 Hz, 1H), 3.97 (d, J = 13.2 Hz, 2H), 3.77 (dt, J = 6.7, 3.4 Hz, 5H), 2.79 (t, J = 6.7 Hz, 3H), 2.57 (d, J = 7.1 Hz, 2H), 1.96-1.73 (m, 2H), 1.66-1.54 (m, 2H), 1.10 (qd, J = 12.4, 4.3 Hz, 2H), 0.89 (d, J = 6.7 Hz, 6H).

-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
96	 cyclohexyl 4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperidine-1-carboxylate	454.3	(500 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 8.56 (d, J = 7.1 Hz, 1H), 7.99 (s, 1H), 7.36 (s, 1H), 6.79 (dd, J = 7.2, 1.9 Hz, 1H), 4.54 (dt, J = 8.6, 4.5 Hz, 1H), 3.97 (d, J = 13.1 Hz, 2H), 3.77 (t, J = 6.7 Hz, 2H), 2.79 (t, J = 6.7 Hz, 4H), 2.57 (d, J = 7.2 Hz, 2H), 1.76 (d, J = 7.9 Hz, 3H), 1.69-1.56 (m, 4H), 1.53-1.20 (m, 6H), 1.09 (qd, J = 12.3, 4.2 Hz, 2H).

Example 97. Preparation of tert-butyl 4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperidine-1-carboxylate (Example 97)




Step 1. 1-(5-bromopyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

[0859] TFA (1.5 mL, 19 mmol) was added to 1-(5-bromopyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl) dihydropyrimidine-2,4(1H,3H)-dione (95 mg, 0.21 mmol) and the mixture was heated at 80° C. overnight. The mixture was then cooled to rt, concentrated and the residue was dissolved in toluene and concentrated again to give crude 1-(5-bromopyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione as a TFA salt which was used without further purification. LCMS [M+H]⁺: 309.1.

[0860] Step 1. tert-butyl 4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperidine-1-carboxylate was prepared from 1-(5-bromopyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione using the method of Example 1, step 2, wherein 1-(5-bromopyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione was used in place of 1-(5-bromopyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione. Purified by reverse phase HPLC using ACN/Water/0.1% TFA. LCMS [M+H]⁺: 428.3. ¹H NMR (500 MHz, Methanol-d₄) δ 8.30 (d, J=7.1 Hz, 1H), 7.88 (s, 1H), 7.25 (s, 1H), 6.72 (d, J=7.2 Hz, 1H), 3.95 (d, J=13.3 Hz, 2H), 3.78 (t, J=6.7 Hz, 2H), 2.79 (t, J=6.8 Hz, 2H), 2.61 (d, J=15.8 Hz, 2H), 2.53 (d, J=7.2 Hz, 2H), 1.74 (s, 1H), 1.56 (d, J=13.3 Hz, 2H), 1.34 (s, 9H), 1.13-0.98 (m, 2H).

Example 98. Preparation of 1-(5-((1-isobutyrylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



Step 1: 3-(2,4-dimethoxybenzyl)-1-(5-((1-isobutyrylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0861] HATU (28 mg, 0.073 mmol) and isobutyric acid (6.2 μ l, 0.097 mmol) were added to a solution of 3-(2,4-dimethoxybenzyl)-1-(5-((1-isobutyrylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride (25 mg, 0.049 mmol) in DMF (1 mL) at rt. The mixture was stirred at rt for 5 min and then DIPEA (0.034 mL, 0.19 mmol) was added. The mixture was stirred at rt for 1 h and then diluted with ethyl acetate and washed sequentially with water and brine. The organic layer was dried over sodium sulfate, filtered and concentrated to give crude 3-(2,4-dimethoxybenzyl)-1-(5-((1-isobutyrylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (27 mg, 0.049 mmol). LCMS $[M+H]^+$: 548.3.

[0862] Step 2: 1-(5-((1-isobutyrylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 98) was prepared from 3-(2,4-dimethoxybenzyl)-1-(5-((1-isobutyrylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione using the method of Example 1, step 5, wherein 3-(2,4-dimethoxybenzyl)-1-(5-((1-isobutyrylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione was used in place of 1-(5-((1-(cyclohexylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione. LCMS $[M+H]^+$: 488.3. 1H NMR (500 MHz, DMSO- d_6) δ 10.43 (s, 1H), 8.56 (d, $J=7.1$ Hz, 1H), 7.99 (s, 1H), 7.37 (d, $J=1.7$ Hz, 1H), 6.80 (dd, $J=7.1, 1.9$ Hz, 1H), 4.39 (d, $J=13.1$ Hz, 1H), 3.93 (d, $J=13.6$ Hz, 1H), 3.77 (t, $J=6.7$ Hz, 2H), 2.97 (t, $J=12.8$ Hz, 1H), 2.86 (h, $J=6.7$ Hz, 1H), 2.79 (t, $J=6.7$ Hz, 2H), 2.58 (d, $J=7.2$ Hz, 2H), 2.47 (d, $J=12.9$ Hz, 1H), 1.86 (ddd, $J=11.2, 7.5, 3.8$ Hz, 1H), 1.73-1.56 (m, 2H), 1.22-1.01 (m, 2H), 0.99 (dd, $J=12.3, 6.7$ Hz, 6H).


[0863] The compounds in the following table were prepared by the method of Example 98, using the appropriate commercially available carboxylic acid in step 1.

Example No.	Structure	Mass $[M+H]^+$	1H NMR
99	1-(5-((1-(cyclohexanecarbonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	438.4	(500 MHz, DMSO- d_6) δ 10.43 (s, 1H), 8.56 (d, $J=7.1$ Hz, 1H), 7.99 (s, 1H), 7.37 (d, $J=1.8$ Hz, 1H), 6.80 (dd, $J=7.1, 1.9$ Hz, 1H), 4.38 (d, $J=13.2$ Hz, 2H), 3.98-3.87 (m, 2H), 3.77 (t, $J=6.7$ Hz, 2H), 2.95 (t, $J=12.8$ Hz, 1H), 2.79 (t, $J=6.7$ Hz, 2H), 2.57 (dd, $J=7.1, 4.1$ Hz, 2H), 1.85 (ddd, $J=11.1, 7.3, 3.7$ Hz, 1H), 1.76-1.52 (m, 7H), 1.30 (t, $J=10.4$ Hz, 4H), 1.22-0.93 (m, 3H).

-continued


Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
100	 1-(5-((1-(1-(trifluoromethyl)cyclopropane-1-carbonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	464.2	(400 MHz, DMSO-d ₆) δ 10.41 (s, 1H), 8.55 (dd, J = 7.5, 2.0 Hz, 1H), 7.98 (d, J = 2.0 Hz, 1H), 7.36 (s, 1H), 6.79 (dd, J = 7.2, 1.9 Hz, 1H), 4.23 (s, 2H), 3.76 (dd, J = 7.8, 5.8 Hz, 2H), 2.78 (dd, J = 7.7, 5.8 Hz, 2H), 2.58 (d, J = 6.9 Hz, 2H), 2.55 (d, J = 1.8 Hz, 2H), 1.87 (s, 1H), 1.67 (d, J = 13.1 Hz, 2H), 1.38-1.23 (m, 2H), 1.13 (d, J = 19.6 Hz, 4H).
101	 1-(5-((1-benzoylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	432.3	(500 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 8.62-8.39 (m, 1H), 7.99 (s, 1H), 7.48-7.42 (m, 3H), 7.37 (qt, J = 3.0, 1.7 Hz, 3H), 6.80 (dd, J = 7.2, 1.9 Hz, 1H), 4.47 (s, 2H), 3.77 (t, J = 6.7 Hz, 2H), 3.56 (s, 1H), 3.00 (s, 1H), 2.79 (t, J = 6.7 Hz, 3H), 2.60 (s, 2H), 2.03-1.46 (m, 2H), 1.19 (s, 2H).
102	 1-(5-((1-(3-phenylpropanoyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	460.3	(400 MHz, Methanol-d ₄) δ 8.40 (d, J = 7.1 Hz, 1H), 7.99 (s, 1H), 7.32 (s, 1H), 7.23 (ddd, J = 18.6, 13.0, 7.4 Hz, 5H), 6.80 (d, J = 7.2 Hz, 1H), 4.52 (d, J = 13.0 Hz, 1H), 3.87 (q, J = 9.5, 8.1 Hz, 3H), 2.97-2.85 (m, 5H), 2.79-2.45 (m, 5H), 1.86 (s, 1H), 1.69 (d, J = 13.4 Hz, 1H), 1.58 (d, J = 13.4 Hz, 1H), 1.30 (s, 1H), 1.17-1.00 (m, 1H), 0.93-0.76 (m, 1H). NH proton not observed due to solvent exchange.

-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
103	 1-(5-((1-(3-cyclohexylpropanoyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	466.2	(400 MHz, Methanol-d ₄) δ 8.40 (d, J = 7.0 Hz, 1H), 7.98 (s, 1H), 7.35 (s, 1H), 6.83 (d, J = 6.9 Hz, 1H), 4.50 (d, J = 13.2 Hz, 1H), 3.97-3.81 (m, 3H), 3.04 (t, J = 13.2 Hz, 1H), 2.88 (t, J = 6.6 Hz, 2H), 2.70-2.51 (m, 3H), 2.37 (dd, J = 9.5, 6.6 Hz, 2H), 1.93 (ddd, J = 13.5, 8.8, 5.1 Hz, 1H), 1.79-1.59 (m, 7H), 1.45 (q, J = 7.4 Hz, 2H), 1.33-1.06 (m, 6H), 0.91 (q, J = 13.7, 12.7 Hz, 2H). NH proton not observed due to solvent exchange.

Example 104. Preparation of 1-(5-((1-(3-cyclohexylpropanoyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

Step 1. 1-(5-((1-(3-cyclohexylpropanoyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl) dihydropyrimidine-2,4(1H,3H)-dione




Example 104

[0864] DIPEA (0.049 mL, 0.28 mmol) and acetic anhydride (0.016 mL, 0.17 mmol) were added to a solution of 3-(2,4-dimethoxybenzyl)-1-(5-((1-(3-cyclohexylpropanoyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione (27 mg, 0.057 mmol) in DCM (1.5 mL) at rt. The mixture was stirred at rt for 30 min and then partitioned between DCM and water. The organic layer separated, washed with brine, dried over sodium sulfate, filtered and concentrated to give crude 1-(5-((1-(3-cyclohexylpropanoyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl) dihydropyrimidine-2,4(1H,3H)-dione that was used without further purification. LCMS [M+H]: 520.4.

[0865] Step 2: 1-(5-((1-(3-cyclohexylpropanoyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione was prepared from 1-(5-((1-(3-cyclohexylpropanoyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl) dihydropyrimidine-2,4(1H,3H)-dione using the method of Example 1, step 5, wherein 1-(5-((1-(3-cyclohexylpropanoyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl) dihydropyrimidine-2,4(1H,3H)-dione was used in place of 1-(5-((1-(3-cyclohexylpropanoyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl) dihydropyrimidine-2,4(1H,3H)-dione. LCMS [M+H]: 370.3. ¹H NMR (400 MHz, DMSO-d₆) δ 10.41 (s, 1H), 8.55 (d, J=7.2 Hz, 1H), 7.98 (s, 1H), 7.35 (d, J=1.6 Hz, 1H), 6.78 (dd, J=7.1, 1.8 Hz, 1H), 4.34 (d, J=13.1 Hz, 1H), 3.77 (q, J=6.3 Hz, 3H), 3.03-2.90 (m, 1H), 2.78 (t, J=6.7 Hz, 2H), 2.57 (d, J=7.2 Hz, 2H), 2.49-2.41 (m, 1H), 1.97 (s, 3H), 1.83 (ddd, J=10.9, 7.4, 3.6 Hz, 1H), 1.61 (t, J=12.6 Hz, 2H), 1.08 (dq, J=47.6, 12.4, 4.2 Hz, 2H).

Example 105. Preparation of 1-(5-((1-(2-methoxy-2-methylpropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



Step 1. 3-(2,4-dimethoxybenzyl)-1-(5-((1-(2-hydroxy-2-methylpropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

[0866] 2,2-Dimethyloxirane (63 mg, 0.87 mmol) and DIPEA (0.15 mL, 0.87 mmol) were added to a solution of 3-(2,4-dimethoxybenzyl)-1-(5-((1-(2-hydroxy-2-methylpropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (150 mg, 0.29 mmol) in THF (2 mL) and MeOH (2 mL) at rt. The mixture was heated at 70° C. overnight, then cooled to rt and concentrated to give crude 3-(2,4-dime-


thoxybenzyl)-1-(5-((1-(2-hydroxy-2-methylpropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione which was used without further purification. LCMS $[M+H]^+$: 550.3.

Step 2. 3-(2,4-dimethoxybenzyl)-1-(5-((1-(2-methoxy-2-methylpropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

[0867] Trimethyloxonium tetrafluoroborate (40 mg, 0.27 mmol) and DIPEA (0.072 mL, 0.41 mmol) were added to a solution of 3-(2,4-dimethoxybenzyl)-1-(5-((1-(2-hydroxy-2-methylpropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (40 mg, 0.068 mmol) in DCM (2 mL) at rt. The mixture was stirred at rt overnight, diluted with DCM and then washed sequentially with saturated aqueous NaHCO_3 solution and brine. The organic layer was dried over Na_2SO_4 , filtered and concentrated to give crude 3-(2,4-dimethoxybenzyl)-1-(5-((1-(2-methoxy-2-methylpropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione which was used without further purification. LCMS $[M+H]^+$: 564.3.

[0868] Step 3: 1-(5-((1-(2-methoxy-2-methylpropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione was prepared from 3-(2,4-dimethoxybenzyl)-1-(5-((1-(2-methoxy-2-methylpropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione using the method of Example 1, step 5, wherein 3-(2,4-dimethoxybenzyl)-1-(5-((1-(2-methoxy-2-methylpropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione. LCMS $[M+H]^+$: 414.4. ^1H NMR (500 MHz, Methanol- d_4) δ 8.46 (d, $J=7.1$ Hz, 1H), 8.03 (s, 1H), 7.39 (s, 1H), 6.86 (d, $J=7.2$ Hz, 1H), 3.90 (q, $J=6.0$ Hz, 4H), 3.45 (s, 3H), 3.30-3.12 (m, 4H), 2.90 (t, $J=6.8$ Hz, 2H), 2.78 (d, $J=7.3$ Hz, 2H), 2.04 (d, $J=9.8$ Hz, 1H), 1.96-1.77 (m, 4H), 1.46 (s, 6H).


Example 106. Preparation of 1-(5-((1-(2-hydroxy-2-methylpropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[0869] Prepared from 3-(2,4-dimethoxybenzyl)-1-(5-((1-(2-hydroxy-2-methylpropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (from step 2 of Example 105) by the method of Example 1, step 5, wherein 3-(2,4-dimethoxybenzyl)-1-(5-((1-(2-hydroxy-2-methylpropyl)piperidin-4-yl)methyl)pyrazolo[1,5-

a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione was used in place of 1-(5-((1-(cyclohexylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione. LCMS $[M+H]^+$: 400.4. 1H NMR (400 MHz, DMSO- d_6) δ 10.42 (s, 1H), 8.58 (dd, $J=7.2, 2.8$ Hz, 1H), 8.00 (s, 1H), 7.36 (d, $J=3.2$ Hz, 1H), 6.78 (dd, $J=7.2, 1.8$ Hz, 1H), 4.00-3.62 (m, 2H), 3.57 (d, $J=12.8$ Hz, 2H), 3.30-3.10 (m, 3H), 3.07-2.89 (m, 3H), 2.79 (t, $J=6.6$ Hz, 2H), 2.63 (dd, $J=28.7, 7.0$ Hz, 1H), 1.91 (d, $J=53.7$ Hz, 1H), 1.68 (dt, $J=42.6, 14.6$ Hz, 4H), 1.23 (d, $J=5.3$ Hz, 6H).

Example 107. Preparation of 1-(5-((1-((1-fluorocyclohexyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



Step 1. 3-(2,4-dimethoxybenzyl)-1-(5-((1-((1-hydroxycyclohexyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


[0870] To a stirred solution of 3-(2,4-dimethoxybenzyl)-1-(5-((1-((1-hydroxycyclohexyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (250 mg, 0.523 mmol) in acetonitrile (5.0 mL) was added lithium perchlorate (110 mg, 1.046 mmol). The reaction was stirred for 10 min followed by the addition of 1-oxaspiro[2.5]octane (293 mg, 2.61 mmol). The mixture was stirred at 90°C for 4 h. After cooling to rt, the reaction was diluted with EtOAc and water. The organic layer was dried over Na_2SO_4 , filtered and concentrated to give crude 3-(2,4-dimethoxybenzyl)-1-(5-((1-((1-hydroxycyclohexyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (200 mg, crude). LCMS $[M+H]^+$: 590.3.

Step 2. 3-(2,4-dimethoxybenzyl)-1-(5-((1-((1-fluorocyclohexyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

[0871] To a stirred solution of 3-(2,4-dimethoxybenzyl)-1-(5-((1-((1-hydroxycyclohexyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (80 mg, 0.135 mmol) in DCM (10 mL) at 0°C was added DAST (43 mg, 0.271 mmol). The reaction was stirred at 0°C for 1 h. The reaction was then diluted with DCM and water. The organic layer was dried over Na_2SO_4 , filtered and concentrated to give crude 3-(2,4-dimethoxybenzyl)-1-(5-((1-((1-fluorocyclohexyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (50 mg, crude). LCMS $[M+H]^+$: 592.2.


[0872] Step 3: 1-(5-((1-((1-fluorocyclohexyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 107) was prepared from 3-(2,4-dimethoxybenzyl)-1-(5-((1-((1-fluorocyclohexyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (10 mg, 0.016 mmol) using the method of Example 1, step 5, wherein 3-(2,4-dimethoxybenzyl)-1-(5-((1-((1-fluorocyclohexyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione was used in place of 1-(5-((1-(cyclohexylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione. The crude compound was purified by chiral HPLC: COLUMN: CHIRALPAK IG, 250 mm $\times$ 20 mm $\times$ 5 μ m, MOBILE PHASE: HEXANE (A), 0.1% DEA in MeOH:EtOH, 1:1 (B), FLOW: 15 mL ISOCRATIC: 75(A):25(B). The collected fractions were concentrated to afford 1-(5-((1-((1-fluorocyclohexyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (3 mg, 0.006 mmol, 33% yield) as an off-white solid. LCMS $[M+H]^+$: 442.3. HPLC Rt=4.95 min. 1H NMR (300 MHz, Methanol- d_4) δ 8.40 (d, $J=7.1$ Hz, 1H), 7.98 (s, 1H), 7.34 (s, 1H), 6.81 (d, $J=7.2$ Hz, 1H), 3.88 (t, $J=6.6$ Hz, 2H), 2.98-2.84 (m, 4H), 2.61 (d, $J=6.5$ Hz, 2H), 2.42 (d, $J=23.5$ Hz, 2H), 2.07 (t, $J=11.2$ Hz, 2H), 1.82 (s, 2H), 1.69-1.47 (m, 10H), 1.43-1.24 (m, 3H). NH proton not observed due to solvent exchange.

Example 108. Preparation of 1-(5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione





[0873] TFA (25 mL, 5.28 mmol) was added to tert-butyl 4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperidine-1-carboxylate (3.05 g, 5.28 mmol). The mixture was heated overnight at in a sealed vial at 85° C. The mixture was then cooled to rt and, concentrated and azeotropically dried with toluene to provide crude product (2.33 g, 5.28 mmol). A portion (~20 mg) of the crude material was dissolved in DMSO, filtered through a 1 micron filter and purified by reverse phase HPLC using ACN/Water/0.1% TFA. The fractions containing the product were combined, frozen and lyophilized to afford a TFA salt of 1-(5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 108) (8 mg, 0.018 mmol). LCMS $[M+H]^+$: 328.3. 1H NMR (500 MHz, DMSO- d_6) δ 10.44 (s, 1H), 8.59 (d, $J=7.1$ Hz, 1H), 8.52 (d, $J=11.4$ Hz, 1H), 8.23 (d, $J=11.4$ Hz, 1H), 8.01 (s, 1H), 7.39 (d, $J=1.8$ Hz, 1H), 6.80 (dd, $J=7.2$, 1.9 Hz, 1H), 3.77 (t, $J=6.7$ Hz, 2H), 3.26 (d, $J=12.6$ Hz, 2H), 2.90-2.76 (m, 4H), 2.61 (d, $J=7.0$ Hz, 2H), 1.89 (ddh, $J=14.7$, 7.3, 3.6 Hz, 1H), 1.80-1.72 (m, 2H), 1.35 (tdd, $J=14.3$, 12.0, 4.0 Hz, 2H).

Example 109. Preparation of 1-(5-((1-isobutylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione





[0874] Isobutyraldehyde (1.90 g, 26.4 mmol) and triethylamine (1.10 mL, 7.92 mmol) were added to a solution of 1-(5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione trifluoroacetate (Example 108) (2.33 g, 5.28 mmol) in DCM (30 mL) and MeOH (2 mL). The reaction mixture was stirred at rt for 30 min and then sodium triacetoxyborohydride (5.59 g, 26.4 mmol) was added. The reaction mixture was stirred overnight at rt and then quenched with a solution of saturated aqueous $NaHCO_3$ and extracted three times with DCM. The combined organic extracts were washed with brine, dried over Na_2SO_4 , filtered and concentrated. Silica gel column chromatography (eluted with 0-100% EtOAc/EtOH (3:1), heptane and 0.1% TEA) provided a light brown solid which was triturated with diethyl ether to give 1-(5-((1-isobutylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione as an off white solid. (1055 mg, 2.738 mmol, 52% yield). LCMS $[M+H]^+$: 384.3. 1H NMR (500 MHz, DMSO- d_6) δ 10.42 (s, 1H), 8.53 (d, $J=7.1$ Hz, 1H), 7.97 (d, $J=1.4$ Hz, 1H), 7.34 (s, 1H), 6.83-6.63 (m, 1H), 3.76 (t, $J=6.6$ Hz, 2H), 2.78 (t, $J=6.8$ Hz, 4H), 2.55 (d, $J=6.5$ Hz, 2H), 1.97 (d, $J=7.4$ Hz, 2H), 1.75 (dt, $J=19.4$, 9.2 Hz, 3H), 1.56 (d, $J=11.5$ Hz, 3H), 1.31-1.09 (m, 2H), 0.83 (d, $J=6.5$ Hz, 6H).


[0875] The compounds in the following table were prepared by the method of Example 109, using the appropriate commercially available aldehyde.

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
110	 1-(5-((1-neopentyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione	398.4	(400 MHz, DMSO-d ₆) δ 10.42 (s, 1H), 8.69-8.52 (m, 1H), 8.00 (d, J = 1.6 Hz, 1H), 7.36 (s, 1H), 6.78 (dd, J = 7.2, 1.8 Hz, 1H), 3.77 (dd, J = 7.5, 6.1 Hz, 2H), 3.47 (d, J = 12.3 Hz, 1H), 3.17 (s, 1H), 3.10-2.99 (m, 2H), 2.94 (d, J = 4.0 Hz, 2H), 2.79 (dd, J = 7.5, 6.0 Hz, 2H), 2.67 (d, J = 7.1 Hz, 1H), 2.61 (d, J = 6.5 Hz, 2H), 1.92 (d, J = 55.2 Hz, 1H), 1.81-1.48 (m, 4H), 1.03 (d, J = 1.3 Hz, 9H).
111	 1-(5-((1-(pyridin-2-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione	419.4	(500 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 8.68 (dd, J = 4.9, 1.7 Hz, 1H), 8.58 (d, J = 7.1 Hz, 1H), 8.00 (s, 1H), 7.93 (td, J = 7.7, 1.8 Hz, 1H), 7.59-7.44 (m, 2H), 7.37 (s, 1H), 6.79 (dd, J = 7.2, 1.9 Hz, 1H), 4.45 (s, 2H), 3.76 (t, J = 6.7 Hz, 2H), 3.40 (d, J = 13.0 Hz, 2H), 3.03 (t, J = 12.5 Hz, 2H), 2.78 (t, J = 6.7 Hz, 2H), 2.62 (d, J = 6.9 Hz, 2H), 2.00-1.73 (m, 3H), 1.64-1.44 (m, 2H).

Example 112. Preparation of 1-(5-((1-(2-cyclohexyl-2,2-difluoroethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H, 3H)-dione



-continued



Example 112

Step 1. 1-(5-((1-(2-cyclohexyl-2,2-difluoroethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl) dihydropyrimidine-2,4(1H, 3H)-dione




[0876] To a solution of 3-(2,4-dimethoxybenzyl)-1-(5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione (200 mg, 0.419 mmol) and 2-cyclohexyl-2,2-difluoroacetaldehyde (1.36 g, 8.38 mmol) [see *Org. Lett.* 2009, 11, 943-946] in DOE (2 mL) was added NaBH(OAc)₃ (133 mg, 0.628 mmol) at rt, then the mixture was stirred for 16 h. The reaction mixture was diluted with water and then extracted with ethyl acetate. The organic layer was dried over Na₂SO₄, filtered and concentrated to give the crude product. The crude product was purified by

prep-HPLC (column: Waters Xbridge C18 150×25 mm×10 μm; mobile phase: [water (0.225% FA)-ACN]; B %: 24%-54%, 10 min), the eluent was concentrated to remove MeCN and lyophilized to give 1-(5-((1-(2-cyclohexyl-2,2-difluoroethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)di-hydropyrimidine-2,4(1H,3H)-dione (100 mg, 0.16 mmol, 38% yield) as a yellow solid. LCMS [M+H]⁺: 624.6.



[0877] Step 2: 1-(5-((1-(2-cyclohexyl-2,2-difluoroethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)di-hydropyrimidine-2,4(1H,3H)-dione (Example 1112) was prepared from 1-(5-((1-(2-cyclohexyl-2,2-difluoroethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)di-hydropyrimidine-2,4(1H,3H)-dione using the method of Example 1, step 5, wherein 1-(5-((1-(2-cyclo-

hexyl-2,2-difluoroethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)di-hydropyrimidine-2,4(1H,3H)-dione was used in place of 1-(5-((1-(cyclohexylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)di-hydropyrimidine-2,4(1H,3H)-dione. LCMS [M+H]⁺: 474.3. ¹H NMR (400 MHz, DMSO-d₆) δ 10.41 (s, 1H), 8.53 (d, J=6.8 Hz, 1H), 7.97 (=, 1H), 7.34 (m, 1H), 6.77-6.75 (m, 1H), 3.76-3.73 (m, 2H), 2.85-2.82 (m, 2H), 2.78-2.75 (m, 2H), 2.70-2.59 (m, 2H), 2.55-2.52 (m, 2H), 2.08-2.07 (m, 2H), 1.99-1.85 (3, 1H), 1.85-1.67 (m, 4H), 1.64-1.61 (m, 1H), 1.55-1.52 (m, 3H), 1.27-1.05 (in, 7H).


[0878] The compounds in the following table were prepared by the method of Example 112, using the appropriate commercially available aldehyde and TEA or DIPEA (2 equiv) in step 1.

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
113	 1-(5-((1-(((2R,6S)-2,6-dimethyltetrahydro-2H-pyran-4-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)di-hydropyrimidine-2,4(1H,3H)-dione	454.3	(300 MHz, Methanol-d ₄) δ 8.39 (d, J = 7.2 Hz, 1H), 8.31 (s, 1H), 7.96 (s, 1H), 7.34 (s, 1H), 6.79 (d, J = 7.2 Hz, 1H), 3.84 (t, J = 6.8 Hz, 2H), 3.68-3.38 (m, 4H), 3.33-3.19 (m, 1H), 2.98-2.79 (m, 5H), 2.65 (d, J = 6.7 Hz, 2H), 2.47-2.03 (m, 1H), 2.02-1.81 (m, 3H), 1.77-1.33 (m, 5H), 1.14 (d, J = 6.4 Hz, 3H), 1.09 (d, J = 6.0 Hz, 2H), 0.86 (q, J = 11.8 Hz, 1H). NH protons not observed due to solvent exchange
114	 1-(5-((1-((1-methyl-1H-pyrazol-5-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)di-hydropyrimidine-2,4(1H,3H)-dione	422.2	(400 MHz, CD ₃ OD): δ 8.44 (d, J = 7.2 Hz, 1H), 8.02 (s, 1H), 7.56 (s, 1H), 7.36 (s, 1H), 6.83 (d, J = 6.8 Hz, 1H), 6.56 (s, 1H), 4.47 (s, 2H), 3.95 (s, 3H), 3.89 (t, J = 6.8 Hz, 2H), 3.57-3.54 (m, 2H), 3.05 (m, 2H), 2.89 (t, J = 7.2 Hz, 2H), 2.71-2.69 (m, 2H), 2.01-1.97 (m, 3H), 1.53 (m, 2H) ppm. NH proton not observed due to solvent exchange
115	 1-(5-((1-(thiazol-4-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)di-hydropyrimidine-2,4(1H,3H)-dione	425.0	(400 MHz, Methanol-d ₄) δ 9.09 (d, J = 2.0 Hz, 1H), 8.42 (d, J = 7.1 Hz, 1H), 8.36 (s, 1H), 8.00 (s, 1H), 7.80 (d, J = 2.0 Hz, 1H), 7.36 (s, 1H), 6.82 (dd, J = 7.6, 1.6 Hz, 1H), 4.38 (s, 2H), 3.88 (t, J = 6.8 Hz, 2H), 3.45 (d, J = 12.5 Hz, 2H), 2.98-2.85 (m, 4H), 2.69 (d, J = 6.7 Hz, 2H), 1.91 (d, J = 15.0 Hz, 3H), 1.54 (q, J = 13.1 Hz, 2H). NH protons not observed due to solvent exchange


-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
116	 1-(5-(((1-methyl-1H-imidazol-4-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	422.2	(400 MHz, Methanol-d ₄) δ 9.04 (s, 1H), 8.43 (d, J = 6.9 Hz, 1H), 8.06-8.01 (m, 1H), 7.83 (s, 1H), 7.39 (d, J = 6.5 Hz, 1H), 6.88-6.80 (m, 1H), 4.47 (s, 2H), 3.97 (s, 3H), 3.88 (t, J = 6.8 Hz, 2H), 3.56 (d, J = 11.9 Hz, 2H), 3.06 (t, J = 12.8 Hz, 2H), 2.89 (t, J = 6.8 Hz, 3H), 2.70 (d, J = 6.7 Hz, 2H). NH proton not observed due to solvent exchange.
117	 1-(5-(((1-((3r,5r,7r)-adamantan-1-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	476.3	(400 MHz, CD ₃ OD) δ 8.42, 8.40 (m, 2H), 7.98 (s, 1H), 7.34 (s, 1H), 6.07 (d, J = 6.8 Hz, 1H), 3.86 (t, J = 7.2 Hz, 2H), 3.48, 3.38 (brs, 2H), 3.15 (s, 2H), 2.88-2.84 (m, 4H), 2.69 (d, J = 6.8 Hz, 2H), 2.01-1.65 (s, 17H). Two protons not integrated due to peak broadening. NH proton not observed due to solvent exchange.

Example 118. Preparation of 1-(5-(((1-((3-methyloxetan-3-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




-continued



[0879] Prepared from 1-(5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 108) using the method of Example 1, step 4, wherein 3-(bromomethyl)-3-methyloxetane was used in place of (bromomethyl)cyclohexane. LCMS [M+H]⁺: 412.3. ¹H NMR (400 MHz, DMSO-d₆) δ 10.42 (s, 1H), 8.70-8.39


(m, 1H), 8.00 (d, J=1.9 Hz, 1H), 7.37 (s, 1H), 6.78 (dd, J=7.2, 2.0 Hz, 1H), 4.45 (d, J=6.2 Hz, 2H), 4.24-4.20 (m, 2H), 3.82-3.67 (m, 3H), 3.60-3.31 (m, 3H), 3.22 (d, J=12.2 Hz, 2H), 3.15-2.88 (m, 2H), 2.78 (t, J=6.8 Hz, 2H), 2.60 (d, J=6.4 Hz, 1H), 1.78 (d, J=14.6 Hz, 3H), 1.70-1.38 (m, 4H).

Example 119. Preparation of 1-(5-((1-(oxetan-3-ylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0880] Prepared from 1-(5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 108) using the method of Example 1, step 4, wherein 3-(bromomethyl)oxetane was used in place of (bromomethyl)cyclohexane. LCMS $[M+H]^+$: 398.4. 1H NMR (500 MHz, Methanol- d_4) δ 8.45 (d, J=7.2 Hz, 1H), 8.03 (d, J=1.8 Hz, 1H), 7.38 (s, 1H), 6.84 (d, J=7.2 Hz, 1H), 4.84 (d, J=7.1 Hz, 2H), 4.50 (t, J=6.0 Hz, 1H), 3.90 (t, J=6.7 Hz, 2H), 3.77-3.67 (m, 1H), 3.67-3.42 (m, 4H), 3.29-3.23 (m, 1H), 3.04-2.88 (m, 4H), 2.77-2.65 (m, 2H), 2.12-1.92 (m, 3H), 1.53 (t, J=14.3 Hz, 2H).

Example 120. Preparation of 1-(5-((1-(2-(1H-imidazol-4-yl)ethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0881] Prepared from 1-(5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 108) using the method of Example 1, step 4, wherein 4-(2-chloroethyl)-1H-imidazole was used in place of (bromomethyl)cyclohexane and with the addition of KI (1.5 equiv). LCMS $[M+H]^+$: 422.4. 1H NMR (500 MHz, DMSO- d_6) δ 10.43 (s, 1H), 9.73-9.28 (m, 1H), 9.02 (s, 1H), 8.59 (d, J=7.1 Hz, 1H), 8.00 (s, 1H), 7.53 (s, 1H), 7.37 (s, 1H), 6.79 (dd, J=7.1, 1.9 Hz, 1H), 3.76 (t, J=6.7 Hz, 2H), 3.56-3.43 (m, 2H), 3.39-3.29 (m, 2H), 3.10 (t, J=7.9 Hz, 2H), 3.02-2.87 (m, 2H), 2.78 (t, J=6.7 Hz, 2H), 2.65-2.56 (m, 2H), 1.96-1.77 (m, 3H), 1.52-1.38 (m, 2H).

Example 121. Preparation of 1-(5-((1-(3-hydroxy-2-(hydroxymethyl)propyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[0882] Prepared from 1-(5-((1-(oxetan-3-ylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 119) using the method of Example 1, step 5, wherein 1-(5-((1-(oxetan-3-ylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione was used in place of 1-(5-((1-(cyclohexylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione. LCMS $[M+H]^+$: 416.3. 1H NMR (500 MHz, Methanol- d_4) δ 8.45 (d, J=7.2 Hz, 1H), 8.03 (d, J=1.6 Hz, 1H), 7.40 (d, J=11.5 Hz, 1H), 6.86 (dd, J=9.2, 7.1 Hz, 1H), 5.51 (d, J=1.5 Hz, 1H), 3.91 (t, J=6.6 Hz, 2H), 3.77-3.67 (m, 4H), 3.58 (dd, J=10.7, 7.6 Hz, 2H), 3.43-3.36 (m, 1H), 3.25 (d, J=6.8 Hz, 2H), 3.02-2.88 (m, 4H), 2.72 (d, J=6.8 Hz, 2H), 2.32 (s, 1H), 2.02 (t, J=18.4 Hz, 3H), 1.55 (q, J=13.2 Hz, 2H).

Example 122. Preparation of 1-(5-((1-(2-methoxybenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




Example 122


[0883] DIPEA (0.053 mL, 0.31 mmol) and 1-(chloromethyl)-2-methoxybenzene (11 mg, 0.073 mmol) were added to a solution of 1-(5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]

pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 108) (20 mg, 0.061 mmol) in DCM. The mixture was stirred at rt for 30 min. Additional 1-(chloromethyl)-2-methoxybenzene (11 mg, 0.073 mmol) and DIPEA (0.053 mL, 0.31 mmol) were added and the mixture was stirred for 2 h at rt. The reaction was then diluted with DCM and washed sequentially with water and brine. The organic layer was dried over sodium sulfate, filtered and concentrated. The residue was dissolved in DMSO, filtered through a 1 micron filter and purified by reverse phase HPLC using ACN/water/0.1% TFA. The fractions containing the product were combined, frozen and lyophilized to afford a TFA salt of 1-(5-((1-(2-methoxybenzyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (2.3 mg, 0.0039 mmol, 6% yield). LCMS $[M+H]^+$: 448.2. 1H NMR (500 MHz, DMSO- d_6) δ 10.45 (d, $J=14.6$ Hz, 1H), 8.60 (dd, $J=12.9, 7.2$ Hz, 1H), 8.01 (d, $J=8.5$ Hz, 1H), 7.54-7.42 (m, 2H), 7.42-7.33 (m, 1H), 7.14 (d, $J=8.3$ Hz, 1H), 7.05 (q, $J=7.8$ Hz, 1H), 6.78 (dd, $J=7.2, 1.9$ Hz, 1H), 4.22 (d, $J=4.9$ Hz, 2H), 3.84 (s, 3H), 3.77 (t, $J=6.7$ Hz, 2H), 3.36 (d, $J=12.1$ Hz, 2H), 2.95 (d, $J=11.6$ Hz, 2H), 2.78 (t, $J=6.8$ Hz, 2H), 2.59 (d, $J=6.7$ Hz, 2H), 1.95-1.70 (m, 3H), 1.44 (q, $J=12.2, 11.0$ Hz, 2H).

Example 123. Preparation of 1-(5-((1-(cyclohexylmethyl)-4-fluoropiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



-continued




Step 1. tert-butyl 4-((3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-4-fluoropiperidine-1-carboxylate

[0884] To an oven-dried vial was added tert-butyl 4-(bromomethyl)-4-fluoropiperidine-1-carboxylate (0.308 g, 1.04 mmol), 1-(5-bromopyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione (0.367 g, 0.8 mmol), $NiCl_2(DME)$ (8.8 mg, 0.040 mmol), pyridine-2,6-bis(carboximidamide) dihydrochloride (9.4 mg, 0.040 mmol), NaI (0.030 g, 0.20 mmol) and Zn (0.105 g, 1.60 mmol). The vial was sealed with a septum cap, evacuated and refilled with nitrogen 3 times. DMA (2.7 mL) and TFA (6 μ L, 0.08 mmol) were added and the reaction was stirred at rt for 2 min. The vial was carefully evacuated and refilled with nitrogen 3 times to remove any H_2 . The reaction was then heated overnight at 70°C, forming a brown reaction mixture. The reaction was cooled to rt, diluted with EtOAc and filtered through a plug of silica gel, eluting with EtOAc. The eluent was concentrated and the residue was purified by silica gel column chromatography (eluted with 0-100% EtOAc in heptane) to give tert-butyl 4-((3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-4-fluoropiperidine-1-carboxylate (0.47 g, 0.80 mmol, 98% yield, purity 70%). LCMS $[M+H]^+$: 596.4. The product was used without further purification.


[0885] Step 2: 1-(5-((1-(cyclohexylmethyl)-4-fluoropiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione was prepared from tert-butyl 4-((3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-4-fluoropiperidine-1-carboxylate using the method of Example 109, wherein cyclohexanecarbaldehyde was used in place of isobutyraldehyde. LCMS $[M+H]^+$: 442.2. 1H NMR (400 MHz, DMSO- d_6) δ 10.44 (s, 1H), 8.62 (dd, $J=7.1, 0.9$ Hz, 1H), 8.04 (d, $J=1.9$ Hz, 1H), 7.43 (d, $J=30.9$ Hz, 1H), 6.92-6.71 (m, 1H), 3.81-3.74 (m, 2H), 3.43 (d, $J=12.2$ Hz, 2H), 3.28-2.86 (m, 6H), 2.82-2.73 (m, 2H), 2.14-1.84 (m, 4H), 1.82-1.53 (m, 6H), 1.18 (dt, $J=30.0, 12.3$ Hz, 3H), 1.01-0.84 (m, 2H).

Example 124. Preparation of 1-(5-((4-fluoro-1-isobutylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0886] Prepared using the method of Example 123, wherein isobutyraldehyde was used in place of cyclohexanecarbaldehyde. LCMS $[M+H]^+$: 402.4. 1H NMR (400 MHz, DMSO- d_6) δ 10.44 (s, 1H), 8.62 (d, $J=7.0$ Hz, 1H), 8.04 (d, $J=1.9$ Hz, 1H), 7.47 (s, 1H), 6.82 (d, $J=7.3$ Hz, 1H), 3.78 (t, $J=6.8$ Hz, 2H), 3.44 (d, $J=12.5$ Hz, 2H), 3.15-2.89 (m, 6H), 2.83-2.73 (m, 2H), 2.17-1.87 (m, 5H), 0.93 (dd, $J=6.7, 2.0$ Hz, 6H).

Example 125. Preparation of 1-(5-((1-(cyclobutylmethyl)-4-fluoropiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0887] Prepared using the method of Example 123, wherein cyclobutanecarbaldehyde was used in place of cyclohexanecarbaldehyde. LCMS $[M+H]^+$: 402.4. 1H NMR (400 MHz, DMSO- d_6) δ 10.44 (s, 1H), 8.61 (dd, $J=7.2, 1.0$ Hz, 1H), 8.04 (s, 1H), 7.46 (d, $J=1.6$ Hz, 1H), 6.93-6.73 (m, 1H), 3.78 (t, $J=6.7$ Hz, 2H), 3.34 (d, $J=12.3$ Hz, 2H), 3.23-2.93 (m, 6H), 2.79 (t, $J=6.8$ Hz, 2H), 2.67 (p, $J=7.3, 6.9$ Hz, 1H), 2.15-1.92 (m, 5H), 1.91-1.71 (m, 5H).


Example 126. Preparation of 1-(5-((1-benzyl-4-fluoropiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[0888] Prepared from tert-butyl 4-((3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-4-fluoropiperidine-1-carboxylate using the method of Example 1, steps 3 to 5, wherein benzyl bromide was used in place of (bromomethyl)cyclohexane. LCMS $[M+H]^+$: 436.2. 1H NMR (500 MHz, DMSO- d_6) δ 10.44 (s, 1H), 8.61 (d, $J=7.1$ Hz, 1H), 8.03 (s, 1H), 7.46 (d, $J=16.6$ Hz, 6H), 6.80 (d, $J=7.3$ Hz, 1H), 4.34 (d, $J=5.1$ Hz, 2H), 3.77 (t, $J=6.6$ Hz, 2H), 3.21-3.01 (m, 6H), 2.77 (t, $J=6.7$ Hz, 2H), 2.64 (s, 1H), 2.07-1.80 (m, 3H).

Example 127. Preparation of 1-(5-((1-(cyclohexylmethyl)piperidin-4-yl)methyl)-4-methylpyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione





Step 1. 1-amino-4-((tert-butoxycarbonyl)amino)-3-methylpyridin-1-ium 2,4-dinitrophenolate

[0889] tert-Butyl (3-methylpyridin-4-yl)carbamate (2.447 g, 10.0 mmol) and O-(2,4-dinitrophenyl)hydroxylamine (2.19 g, 11.0 mmol) were added to a reaction flask. 2-MeTHF (20 mL) was added and the reaction was heated to at 40° C. for 1 h, then stirred at rt overnight. Additional O-(2,4-dinitrophenyl)hydroxylamine (600 mg, 3.00 mmol, 0.3 eq) was added and the reaction stirred for another 2 h at 40° C. The reaction was diluted with isopropanol and concentrated to give a yellow solid which was suspended in cold IPA, filtered and dried to give 1-amino-4-((tert-butoxycarbonyl)amino)-3-methylpyridin-1-ium 2,4-dinitrophenolate (5.5 g, crude) which was used without further purification. LCMS [M]⁺: 224.1.

Step 2. ethyl 5-((tert-butoxycarbonyl)amino)-4-methylpyrazolo[1,5-a]pyridine-3-carboxylate

[0890] Potassium carbonate (5.60 g, 40.5 mmol) was added to a mixture of 1-amino-4-((tert-butoxycarbonyl)amino)-3-methylpyridin-1-ium 2,4-dinitrophenolate (5.5 g, 13.5 mmol) in DMF (13.5 mL) at 0° C. After 5 min, ethyl

propionate (1.50 mL, 14.8 mmol) was added and the reaction was stirred at 0° C. and allowed to warm to rt overnight. Two regioisomers were present by LCMS. The reaction was concentrated and the residue was suspended in water, filtered and the solid was purified by silica gel chromatography to give ethyl 5-((tert-butoxycarbonyl)amino)-4-methylpyrazolo[1,5-a]pyridine-3-carboxylate (849 mg, 2.66 mmol, 20% yield) as a single regioisomer. LCMS [M+H]⁺: 320. ¹H NMR (500 MHz, CDCl₃) δ 8.37 (d, J=1.8 Hz, 1H), 8.34 (d, J=7.6 Hz, 1H), 7.80 (d, J=7.6 Hz, 1H), 6.58 (s, 1H), 4.37-4.27 (m, 2H), 2.77 (d, J=1.8 Hz, 3H), 1.55 (d, J=1.7 Hz, 9H), 1.39 (td, J=7.1, 1.8 Hz, 3H).

Step 3. ethyl 5-amino-4-methylpyrazolo[1,5-a]pyridine-3-carboxylate

[0891] TFA (3.3 mL) was added to a solution of ethyl 5-((tert-butoxycarbonyl)amino)-4-methylpyrazolo[1,5-a]pyridine-3-carboxylate (850 mg, 2.66 mmol) in DCM (10 mL) at rt. The reaction was stirred at rt for 1 h and then concentrated. The residue was azeotropically dried with toluene to give ethyl 5-amino-4-methylpyrazolo[1,5-a]pyridine-3-carboxylate (850 mg, 2.66 mmol) as a light yellow solid. LCMS [M+H]⁺: 220.

Step 4. ethyl 5-iodo-4-methylpyrazolo[1,5-a]pyridine-3-carboxylate

[0892] A solution of sodium nitrite (37.9 mg, 0.550 mmol) in water (0.50 mL) was added dropwise to a suspension of ethyl 5-amino-4-methylpyrazolo[1,5-a]pyridine-3-carboxylate (167 mg, 0.5 mmol) in MeCN (0.83 mL) and aqueous 6 M HCl (2.5 mL) at 0° C. The reaction turned bright yellow and was stirred at 0° C. for 1 h. A solution of potassium iodide (166 mg, 1.00 mmol) in water (0.50 mL) was added dropwise to the vigorously stirring reaction; the reaction turned dark brown and bubbled and a precipitate formed. After 15 min, the reaction was diluted with water, filtered and the solid was washed with water. The solid was then dissolved in EtOH/DCM and concentrated. The solid was suspended in cold methanol, filtered and dried to give ethyl 5-iodo-4-methylpyrazolo[1,5-a]pyridine-3-carboxylate (141 mg, 0.427 mmol, 85% yield) as a light yellow solid. LCMS [M+H]⁺: 331.

Step 5. ethyl 5-((1-(tert-butoxycarbonyl)piperidin-4-yl)methyl)-4-methylpyrazolo[1,5-a]pyridine-3-carboxylate

[0893] Prepared from ethyl 5-iodo-4-methylpyrazolo[1,5-a]pyridine-3-carboxylate by the method of Example 1, step 2, wherein ethyl 5-iodo-4-methylpyrazolo[1,5-a]pyridine-3-carboxylate was used in place of 1-(5-bromopyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione. LCMS [M+H-TBu]⁺: 346.1.

Step 6. 5-((1-(tert-butoxycarbonyl)piperidin-4-yl)methyl)-4-methylpyrazolo[1,5-a]pyridine-3-carboxylic acid

[0894] Sodium hydroxide (360 mg, 9.00 mmol) was added to a solution of ethyl 5-((1-(tert-butoxycarbonyl)piperidin-4-yl)methyl)-4-methylpyrazolo[1,5-a]pyridine-3-carboxylate (723 mg, 1.80 mmol) in EtOH (7.2 mL) and water (1.8 mL). The mixture was heated at 60° C. for 3 h, then cooled to rt and concentrated. The residue was dissolved in water, filtered and then aqueous 6 M HCl was added dropwise until

the product precipitated. The supernatant was decanted and the solid was washed with water, dried and purified by silica gel chromatography (eluted with 0-50% of (1% AcOH in 3:1 EtOAc/EtOH) in heptane) to give 5-((1-(tert-butoxycarbonyl)piperidin-4-yl)methyl)-4-methylpyrazolo[1,5-a]pyridine-3-carboxylic acid (636 mg, 1.703 mmol, 95% yield) as an off-white solid. LCMS [M+H-tBu]⁺: 318.

Step 7. tert-butyl 4-((3-((ethoxycarbonyl)amino)-4-methylpyrazolo[1,5-a]pyridin-5-yl)methyl)piperidine-1-carboxylate

[0895] DIPEA (675 μ l, 3.86 mmol) was added to a solution of 5-((1-(tert-butoxycarbonyl)piperidin-4-yl)methyl)-4-methylpyrazolo[1,5-a]pyridine-3-carboxylic acid (481 mg, 1.29 mmol) in dioxane (4.3 mL) at rt. The bright yellow mixture was stirred at rt for 3 h, then EtOH (1.5 mL, 25.8 mmol) was added and the reaction was heated at 100° C. for 15 min. The reaction was cooled to rt, diluted with water and brine and extracted with EtOAc. The organic layer was dried over Na₂SO₄, filtered and concentrated. The residue was purified by silica gel chromatography (eluted with 0-60% EtOAc/heptane) to give tert-butyl 4-((3-((ethoxycarbonyl)amino)-4-methylpyrazolo[1,5-a]pyridin-5-yl)methyl)piperidine-1-carboxylate (479 mg, 1.150 mmol, 89% yield) as a colorless oil. LCMS [M+H-tBu]⁺: 361.

Step 8. tert-butyl 4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-4-methylpyrazolo[1,5-a]pyridin-5-yl)methyl)piperidine-1-carboxylate

[0896] Acrylamide (14.2 mg, 0.200 mmol) and tert-butyl 4-((3-((ethoxycarbonyl)amino)-4-methylpyrazolo[1,5-a]pyridin-5-yl)methyl)piperidine-1-carboxylate (41.7 mg, 0.1 mmol) were added to a vial followed by tBuOH (0.5 mL) and potassium tert-butoxide (110 μ l, 0.110 mmol) (1.0 M in THF)—the reaction turned light yellow. The mixture was heated at 60° C. overnight. The reaction was quenched with

saturated aqueous NaHCO₃ and water. The mixture was extracted with EtOAc, dried over Na₂SO₄, filtered and concentrated to give crude tert-butyl 4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-4-methylpyrazolo[1,5-a]pyridin-5-yl)methyl)piperidine-1-carboxylate (39 mg, 0.088 mmol, 88% yield). LCMS [M+H-Boc]⁺: 342.


Step 9. 1-(4-methyl-5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

[0897] TFA (215 μ l) was added to a solution of tert-butyl 4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-4-methylpyrazolo[1,5-a]pyridin-5-yl)methyl)piperidine-1-carboxylate (38 mg, 0.086 mmol) in DCM (645 μ l) at rt. The reaction was stirred at rt for 30 min and was then concentrated to give crude 1-(4-methyl-5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione that was used without further purification. LCMS [M+H]⁺: 342.




Step 10. 1-(5-((1-(cyclohexylmethyl)piperidin-4-yl)methyl)-4-methylpyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

[0898] Prepared from 1-(4-methyl-5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione by the method of Example 109, wherein cyclohexanecarbaldehyde was used in place of isobutyraldehyde. LCMS [M+H]⁺: 438.5. ¹H NMR (500 MHz, DMSO-d₆) δ 10.45 (s, 1H), 8.60 (s, 1H), 8.44 (d, J=7.1 Hz, 1H), 7.98 (s, 1H), 6.73 (d, J=7.1 Hz, 1H), 3.79 (dt, J=13.3, 7.1 Hz, 1H), 3.65 (dt, J=12.6, 6.4 Hz, 2H), 2.88-2.80 (m, 3H), 2.76 (t, J=6.8 Hz, 2H), 2.68-2.59 (m, 2H), 2.35 (s, 3H), 1.86-1.58 (m, 10H), 1.53 (d, J=12.6 Hz, 2H), 1.33-1.07 (m, 4H), 0.92 (d, J=12.8 Hz, 2H).


[0899] The compounds in the following table were prepared by the method of Example 1, using tert-butyl 3-methylenepyrrolidine-1-carboxylate in step 2 and the appropriate commercially available halide in step 4.

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
128	 1-(5-((1-(cyclohexylmethyl)pyrrolidin-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	410.3	(500 MHz, Methanol-d ₄) δ 8.35 (d, J = 7.2 Hz, 1H), 7.93 (s, 1H), 7.30 (d, J = 6.9 Hz, 1H), 6.75 (d, J = 7.2 Hz, 1H), 3.80 (t, J = 6.5 Hz, 2H), 3.73-3.42 (m, 2H), 2.95 (t, J = 7.2 Hz, 2H), 2.87-2.71 (m, 5H), 2.70-2.58 (m, 1H), 2.21-2.02 (m, 1H), 1.84 (q, J = 10.8, 9.0 Hz, 1H), 1.65 (dd, J = 35.6, 13.5 Hz, 6H), 1.19 (ddd, J = 41.2, 25.5, 12.9 Hz, 4H), 0.93 (d, J = 12.3 Hz, 2H).


-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
129	 1-(5-((1-(pyridin-3-ylmethyl)pyrrolidin-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)methylpyrrolidine-2,4-dione	405.3	(500 MHz, DMSO-d ₆) δ 10.45 (d, J = 3.2 Hz, 1H), 8.73 (dd, J = 6.0, 2.1 Hz, 1H), 8.66 (d, J = 5.5 Hz, 1H), 8.60 (d, J = 7.2 Hz, 1H), 8.02 (d, J = 5.0 Hz, 1H), 7.96 (d, J = 7.9 Hz, 1H), 7.61-7.48 (m, 1H), 7.39 (d, J = 23.9 Hz, 1H), 6.81 (td, J = 5.1, 2.6 Hz, 1H), 4.45 (t, J = 5.1 Hz, 2H), 3.77 (q, J = 6.0 Hz, 2H), 3.58-3.25 (m, 3H), 3.16 (dd, J = 11.6, 6.1 Hz, 1H), 3.02-2.55 (m, 5H), 2.25-1.94 (m, 1H), 1.91-1.55 (m, 1H).
130	 1-(5-((1-(cyclobutylmethyl)pyrrolidin-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)methylpyrrolidine-2,4-dione	382.3	(500 MHz, DMSO-d ₆) δ 10.45 (s, 1H), 8.60 (d, J = 7.1 Hz, 1H), 8.02 (s, 1H), 7.39 (d, J = 16.8 Hz, 1H), 6.82 (ddd, J = 6.5, 4.2, 1.7 Hz, 1H), 3.83-3.72 (m, 2H), 3.50 (dq, J = 14.3, 6.4 Hz, 2H), 3.19 (q, J = 7.0 Hz, 3H), 3.15-2.95 (m, 1H), 2.85-2.69 (m, 5H), 2.60 (tt, J = 19.8, 9.2 Hz, 1H), 2.18-1.95 (m, 3H), 1.89 (ddt, J = 12.3, 8.4, 3.6 Hz, 1H), 1.83-1.71 (m, 4H).
131	 1-(5-((1-(3-fluorobenzyl)pyrrolidin-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)methylpyrrolidine-2,4-dione	422.3	(500 MHz, Methanol-d ₄) δ 8.34 (d, J = 7.1 Hz, 1H), 7.92 (s, 1H), 7.41 (q, J = 7.3 Hz, 1H), 7.30 (s, 1H), 7.21 (dd, J = 14.6, 8.6 Hz, 2H), 7.14 (t, J = 8.7 Hz, 1H), 6.74 (d, J = 7.2 Hz, 1H), 4.31 (s, 2H), 3.79 (t, J = 6.7 Hz, 2H), 3.45 (d, J = 26.2 Hz, 2H), 3.07 (s, 1H), 2.88 (t, J = 10.8 Hz, 1H), 2.79 (t, J = 6.7 Hz, 4H), 2.63 (s, 1H), 2.13 (d, J = 61.3 Hz, 1H), 1.93-1.56 (m, 1H).

-continued


Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
132	 1-(5-((1-isobutylpyrrolidin-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	370.3	(500 MHz, DMSO-d ₆) δ 10.45 (s, 1H), 8.61 (d, J = 7.2 Hz, 1H), 8.02 (s, 1H), 7.39 (d, J = 13.4 Hz, 1H), 6.82 (ddd, J = 7.2, 3.7, 1.9 Hz, 1H), 3.78 (t, J = 6.7 Hz, 2H), 3.50 (m, 1H), 3.30-3.13 (m, 2H), 3.00 (dt, J = 9.8, 6.6 Hz, 2H), 2.88-2.71 (m, 5H), 2.61 (dd, J = 18.5, 9.7 Hz, 1H), 2.19-1.88 (m, 2H), 1.86-1.55 (m, 1H), 0.95 (dd, J = 6.6, 4.1 Hz, 6H).

Example 133. Preparation of 1-(5-(azetidin-3-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[0900] Prepared using the method of Example 1, steps 2 and 5, wherein tert-butyl 3-methyleneazetidine-1-carboxylate was used in place of tert-butyl 4-methylenepiperidine-1-carboxylate. LCMS [M+H]⁺: 300.0. ¹H NMR (300 MHz, CD₃OD) δ 8.53 (s, 1H), 8.44 (d, J=7.2 Hz, 1H), 8.01 (s, 1H), 7.37 (s, 1H), 6.82-6.79 (m, 1H), 4.09 (t, J=8.7 Hz, 2H), 3.92-3.86 (m, 4H), 3.34 (m, 1H), 3.03 (d, J=8.1 Hz, 2H), 2.88 (t, J=7.2 Hz, 2H).


Example 134. Preparation of 1-(5-((1-isobutylazetidin-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[0901] Prepared from 1-(5-(azetidin-3-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


(Example 133) using the method of Example 109, wherein 1-(5-(azetidin-3-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione was used in place of 1-(5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione trifluoroacetate. LCMS [M+H]⁺: 356.3. ¹H NMR (300 MHz, CD₃OD) δ 8.55 (s, 1H), 8.43 (d, J=7.2 Hz, 1H), 8.01 (s, 1H), 7.37 (s, 1H), 6.81-6.79 (m, 1H), 3.96-3.87 (m, 4H), 3.63-3.61 (m, 2H), 3.13-3.09 (m, 1H), 3.00 (d, J=8.0 Hz, 2H), 2.89 (d, J=7.2 Hz, 2H), 2.79 (d, J=6.8 Hz, 2H), 1.84-1.81 (m, 1H), 0.96 (d, J=6.8 Hz, 6H), NH proton not observed due to solvent exchange.

Example 135. Preparation of 1-(5-((1-(cyclohexylmethyl)azetidin-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[0902] Prepared from 1-(5-(azetidin-3-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 133) using the method of Example 109, wherein 1-(5-(azetidin-3-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione was used in place of 1-(5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione trifluoroacetate and cyclohexanecarbaldehyde was used in place of isobutyraldehyde. LCMS [M+H]⁺: 396.1. ¹H NMR (300 MHz, CD₃OD) δ 8.54 (s, 1H), 8.42 (d, J=7.2 Hz, 1H), 8.00 (s, 1H), 7.35 (s, 1H), 6.80-6.77 (m, 1H), 3.95-3.86 (m, 4H), 3.62-3.56 (m, 2H), 3.09-3.07 (m, 1H), 2.99-2.97 (m, 2H), 2.88 (d, J=6.6 Hz, 2H), 2.80-2.78 (m, 2H), 1.73-1.70 (m, 5H), 1.51 (s, 1H), 1.30-0.95 (m, 5H).

Example 136. Preparation of 1-(5-((1-(2-(tetrahydro-2H-pyran-4-yl)propan-2-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



Step 1. (4-methylenepiperidin-1-yl)(tetrahydro-2H-pyran-4-yl)methanone


[0903] To a stirred solution of tetrahydro-2H-pyran-4-carboxylic acid (4.0 g, 27.7 mmol) in THF (80 mL) was added HATU (15.81 g, 41.60 mmol), DIPEA (14.2 mL, 83.2 mmol) at 0° C. The mixture was stirred for 10 min followed by the addition of a solution of 4-methylenepiperidine (4.41 g, 33.3 mmol) in THF (20 mL). The reaction mixture was then stirred at rt for 12 h. The reaction was then quenched with water and extracted with EtOAc. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated. The crude material was purified by silica gel chromatography (eluting with 50% EtOAc in hexanes) to afford (4-methylenepiperidin-1-yl)(tetrahydro-2H-pyran-4-yl)methanone (3.8 g, 18.1 mmol, 59% yield). LCMS [M+H]⁺: 210.0.

Step 2. 4-methylene-1-(2-(tetrahydro-2H-pyran-4-yl)propan-2-yl)piperidine

[0904] To a stirred solution of (4-methylenepiperidin-1-yl)(tetrahydro-2H-pyran-4-yl)methanone (1.0 g, 4.7 mmol) in THF (12 mL) was added ZrCl₄ (1.09 g, 4.7 mmol) at -20° C. and the mixture was stirred for 30 min. A solution of MeMgBr·Et₂O (9.4 mL, 28.2 mmol, 3.0 M) was added and the mixture was stirred for 10 min at -20° C. and then at rt for 2 h. After completion, the reaction was quenched with water (10 mL) and extracted with EtOAc. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated. The crude material was purified by silica gel chromatography (eluting with 50% EtOAc in hexanes) to afford 4-methylene-1-(2-(tetrahydro-2H-pyran-4-yl)propan-2-yl)piperidine (250 mg, 1.12 mmol, 24% yield). LCMS [M+H]⁺: 224.0.


[0905] Step 3: 1-(5-((1-(2-(tetrahydro-2H-pyran-4-yl)propan-2-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione was prepared from 4-methylene-1-(2-(tetrahydro-2H-pyran-4-yl)propan-2-yl)piperidine using the method of Example 1, steps 2 and 5, wherein was used in place of 4-methylene-1-(2-(tetrahydro-2H-pyran-4-yl)propan-2-yl)piperidine was used in place of tert-butyl 4-methylenepiperidine-1-carboxylate. LCMS [M+H]⁺: 454.2. ¹H NMR (400 MHz, DMSO-d₆): δ 10.44 (s, 1H), 8.59 (d, J=7.2 Hz, 1H), 8.00 (s, 1H), 7.83 (s, 1H), 7.35 (s, 1H), 6.78 (d, J=6.8 Hz, 1H), 3.91-3.88 (m, 2H), 3.74 (t, J=6.8 Hz, 2H), 3.28 (s, 2H), 2.96-2.93 (m, 2H), 2.78 (t, J=6.4 Hz, 2H), 2.61-2.59 (m, 2H), 2.42 (brs, 1H), 2.02-1.80 (m, 5H), 1.58-1.48 (m, 4H), 1.35-1.28 (m, 2H), 1.20 (s, 6H).

Example 137. Preparation of 1-(5-((1-(2-methyl-1-(tetrahydro-2H-pyran-4-yl)propan-2-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0906] Prepared using the method of Example 136, wherein 2-(tetrahydro-2H-pyran-4-yl)acetic acid was used in place of tetrahydro-2H-pyran-4-carboxylic acid. LCMS [M+H]⁺: 468.3. ¹H NMR (400 MHz, DMSO-d₆): δ 10.44 (s, 1H), 8.60 (d, J=7.1 Hz, 1H), 8.28 (t, J=8.3 Hz, 1H), 8.00 (s, 1H), 7.36 (s, 1H), 6.78 (d, J=7.1 Hz, 1H), 3.84-3.72 (m, 4H), 3.47 (s, 1H), 3.29 (t, J=11.4 Hz, 2H), 2.90 (q, J=11.7 Hz, 2H), 2.78 (t, J=6.7 Hz, 2H), 2.61 (d, J=6.4 Hz, 2H), 1.92 (s, 2H), 1.85 (d, J=14.6 Hz, 2H), 1.68-1.52 (m, 5H), 1.46 (t, J=12.9 Hz, 2H), 1.38-1.13 (m, 3H), 1.30 (s, 6H).

Example 138. Preparation of 1-(5-((1-(2-cyclobutylpropan-2-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




[0907] Prepared using the method of Example 136, wherein cyclobutanecarboxylic acids was used in place of tetrahydro-2H-pyran-4-carboxylic acid. LCMS $[M+H]^+$: 424.3. 1H NMR (400 MHz, Methanol- d_4) δ 8.43 (d, $J=7.2$ Hz, 1H), 8.01 (s, 1H), 7.36 (s, 1H), 7.08 (s, 1H), 6.83 (dd, $J=7.1, 1.9$ Hz, 1H), 3.89 (t, $J=6.8$ Hz, 2H), 3.58-3.48 (m, 2H), 2.97-2.86 (m, 3H), 2.82-2.74 (m, 1H), 2.68 (d, $J=6.7$ Hz, 1H), 2.64 (d, $J=5.9$ Hz, 2H), 2.09-1.90 (m, 6H), 1.50-1.26 (m, 10H).

Example 139. Preparation of 1-(5-((1-(2,4-dimethylpentan-2-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




[0908] Prepared using the method of Example 136, wherein 3-methylbutanoic acid was used in place of tetrahydro-2H-pyran-4-carboxylic acid. LCMS $[M+H]^+$: 426.3. 1H NMR (300 MHz, Methanol- d_4) δ 8.53 (s, 1H), 8.43 (d, $J=7.2$ Hz, 1H), 8.01 (s, 1H), 7.38 (s, 1H), 6.84 (d, $J=7.3$ Hz, 1H), 3.89 (t, $J=6.8$ Hz, 2H), 3.71-3.48 (m, 3H), 2.97 (t, $J=12.2$ Hz, 2H), 2.89 (t, $J=6.6$ Hz, 3H), 2.69 (d, $J=6.3$ Hz, 2H), 2.00 (d, $J=13.2$ Hz, 3H), 1.75 (hept, $J=6.3$ Hz, 1H), 1.63 (d, $J=5.4$ Hz, 2H), 1.61-1.49 (m, 3H), 1.41 (s, 6H), 1.03 (d, $J=6.6$ Hz, 6H). NH proton not observed due to solvent exchange.


Example 140. Preparation of 1-(5-((1-(2-(4,4-difluorocyclohexyl)propan-2-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



[0909] Prepared using the method of Example 136, wherein 4,4-difluorocyclohexane-1-carboxylic acid was used in place of tetrahydro-2H-pyran-4-carboxylic acid. LCMS $[M+H]^+$: 488.4. 1H NMR (400 MHz, CD_3OD): δ 8.43 (d, $J=7.2$ Hz, 1H), 8.00 (s, 1H), 7.35 (s, 1H), 6.82 (dd, $J=7.6$ Hz, 2.0 Hz, 1H), 3.87 (t, $J=6.8$ Hz, 2H), 3.59 (d, $J=12.4$ Hz, 2H), 3.06-2.98 (m, 2H), 2.87 (t, $J=6.8$ Hz, 2H), 2.70 (d, $J=6.8$ Hz, 2H), 2.08-2.10 (m, 2H), 2.01-1.75 (m, 8H), 1.64-1.46 (m, 4H), 1.33 (m, 6H). NH proton not observed due to solvent exchange.

Example 141. Preparation of 1-(5-(((2S,4R)-1-((4,4-difluorocyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione





Step 1. tert-butyl (S)-4-(methoxymethylene)-2-methylpiperidine-1-carboxylate

[0910] A solution of K_{OTf} (63.78 mL, 63.78 mmol, 1 M in THF) was added dropwise to a stirred suspension of (methoxymethyl)triphenylphosphonium chloride (21.86 g, 63.78 mmol) in THF (70 mL) at 0° C. The red solution was stirred for 30 min at rt and then cooled again to 0° C. A solution of tert-butyl (S)-2-methyl-4-oxopiperidine-1-carboxylate (8.0 g, 37.5 mmol) in THF (30 mL) was added and the reaction mixture was stirred at rt for 16 h. The reaction mixture was quenched with a solution of saturated aqueous NH₄Cl and extracted with EtOAc. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated. The crude material was purified by silica gel chromatography (eluting with 15-20% EtOAc in hexanes) to afford tert-butyl (S)-4-(methoxymethylene)-2-methylpiperidine-1-carboxylate (7.5 g, 31 mmol, 82% yield) as a mixture of E:Z isomers. LCMS [M+H-tBu]⁺: 186. ¹H NMR (300 MHz, CDCl₃) δ 5.96 (s, 1H), 5.78 (s, 1H), 4.46-4.39 (m, 2H), 4.00-3.89 (m, 2H), 3.56 (s, 3H), 3.53 (s, 3H), 2.86-2.74 (m, 2H), 2.62-2.47 (m, 2H), 2.31-2.24 (m, 1H), 2.02-1.94 (m, 4H), 1.90-1.81 (m, 3H), 1.46 (s, 18H), 1.04 (d, J=6.6 Hz, 3H).

Step 2. tert-butyl (2S)-4-formyl-2-methylpiperidine-1-carboxylate

[0911] A solution of HCl (2.0 M in water, 75 mL) was added to a solution of tert-butyl (S)-4-(methoxymethylene)-2-methylpiperidine-1-carboxylate (7.5 g, 31 mmol) in MeCN (220 mL) at rt. The reaction was heated to 40° C. and stirred for 40 min. The reaction was then cooled to rt and quenched by addition of solid NaHCO₃. Brine was added and the reaction was extracted with EtOAc 3 times. The combined organic layers were dried over Na₂SO₄, filtered and concentrated to give crude tert-butyl (2S)-4-formyl-2-methylpiperidine-1-carboxylate as a ~5:1 mixture of cis:trans isomers (7 g, 31 mmol). The crude material was used in the next reaction without further purification. LCMS [M+H-tBu]⁺: 172.

Step 3. tert-butyl (2S,4R)-4-(hydroxymethyl)-2-methylpiperidine-1-carboxylate

[0912] NaOMe (335 mg, 6.15 mmol) was added to a stirred solution of crude tert-butyl (2S)-4-formyl-2-methylpiperidine-1-carboxylate (7 g, 31 mmol) in MeOH (70

mL) at 0° C. The reaction mixture was kept at 4° C. for 24 h. After 24 h the reaction mixture was placed in an ice bath, NaBH₄ (4.65 g, 123 mmol) was added at 0° C. and the reaction was then stirred at rt for 10 min. The reaction mixture was quenched with water and extracted with EtOAc. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated. The crude material was purified by silica gel chromatography (eluted with 20% EtOAc in hexanes) to afford tert-butyl (2S,4R)-4-(hydroxymethyl)-2-methylpiperidine-1-carboxylate (5.4 g, 76% yield). LCMS [M+H-tBu]⁺: 174. ¹H NMR (400 MHz, CDCl₃) δ 4.45-4.41 (m, 1H), 3.99-3.94 (m, 1H), 3.38-3.35 (m, 2H), 2.90 (brs, 1H), 1.87-1.84 (m, 1H), 1.75-1.72 (m, 1H), 1.65-1.62 (m, 1H), 1.46 (s, 9H), 1.32-1.25 (m, 2H), 1.17 (d, J=6.4 Hz, 3H), 1.05-1.01 (s, 1H).

Step 4. tert-butyl (2S,4R)-4-(bromomethyl)-2-methylpiperidine-1-carboxylate

[0913] Triphenylphosphine dibromide (1.2 g, 28.5 mmol) was added to a solution of imidazole (2.10 g, 30.6 mmol) and tert-butyl (2S,4R)-4-(hydroxymethyl)-2-methylpiperidine-1-carboxylate (5.4 g, 23.5 mmol in DCM (50 mL) at 0° C. The reaction mixture was stirred at rt for 16 h. After completion of the reaction, the mixture was diluted with DCM and washed with water. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated. The crude material was purified by silica gel chromatography (eluted with 10% EtOAc in hexanes) to afford tert-butyl (2S,4R)-4-(bromomethyl)-2-methylpiperidine-1-carboxylate (3.5 g, 12.2 mmol, 50% yield). LCMS [M+H-tBu]⁺: 236. ¹H NMR (300 MHz, CDCl₃) δ 4.45 (brs, 1H), 4.01 (brs, 1H), 3.30-3.20 (m, 2H), 2.88-2.80 (m, 1H), 2.04-1.94 (m, 1H), 1.85-1.81 (m, 1H), 1.70-1.65 (m, 1H), 1.29-1.03 (m, 15H).

Step 5. tert-butyl (2S,4R)-4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperidine-1-carboxylate

[0914] Zn powder was activated by taking commercial material and stirring it vigorously in a solution of aqueous 1M HCl for 10 min. The material was then filtered and the large chunks were broken up with a spatula. The solids were washed with distilled water, followed by EtOH, followed by Et₂O. The solids were then heated at 50° C. overnight under vacuum.

[0915] To an oven-dried 2-necked flask was added tert-butyl (2S,4R)-4-(bromomethyl)-2-methylpiperidine-1-carboxylate (2.98 g, 10.2 mmol), 1-(5-bromopyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione (3.90 g, 8.5 mmol), NiCl₂(DME) (0.093 g, 0.425 mmol), pyridine-2,6-bis(carboximidamide) dihydrochloride (0.100 g, 0.425 mmol), Zn powder (1.11 g, 17.0 mmol) and sodium iodide (0.319 g, 2.125 mmol). The flask was sealed with a septum and evacuated and refilled with argon 3 times. DMA (34.0 mL, degassed by bubbling argon through for several min) was added and the reaction was stirred at 70° C. overnight. The reaction was cooled to rt and poured into water and a grey precipitate formed which was collected by filtration. The solid was diluted with EtOH (200 mL) and filtered through celite (washed with EtOH) to remove Zn solids. The filtrate was concentrated and the crude material was purified by silica gel chromatography

(eluted with 10-100% of (3:1 EtOAc/EtOH w/0.1% Et₃N)/heptane) to give tert-butyl (2S,4R)-4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperidine-1-carboxylate (4.30 g, 6.69 mmol, 79% yield) as an off-white solid. LCMS [M+H-Boc]⁺: 492.2.

Step 6. 3-(2,4-dimethoxybenzyl)-1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride

[0916] A solution of HCl (4.0 M in dioxane, 40 mL) was added to tert-butyl (2S,4R)-4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperidine-1-carboxylate (5.1 g, 8.61 mmol) and the mixture was stirred at rt for 3 h and then concentrated. The crude compound was triturated with diethyl ether to afford 3-(2,4-dimethoxybenzyl)-1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride (4.8 g, crude). LCMS [M+H]⁺: 492.3.

Step 7. 1-(5-(((2S,4R)-1-((4,4-difluorocyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione

[0917] TEA (3.3 mL, 23.7 mmol) and 4,4-difluorocyclohexane-1-carbaldehyde (1.4 g, 9.48 mmol) were added to a solution of 3-(2,4-dimethoxybenzyl)-1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride (2.5 g, 4.74 mmol) in DCM (25 mL) at rt and the mixture was stirred for 1.5 h. The reaction was then cooled to 0° C. and sodium triacetoxyborohydride (2.00 g, 9.48 mmol) was added. The reaction mixture was stirred for 16 h at rt. After completion, the reaction was diluted with DCM and water and the organic layer was dried over Na₂SO₄, filtered and concentrated to afford crude 1-(5-(((2S,4R)-1-((4,4-difluorocyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione (1.7 g, crude). LCMS [M+H]⁺: 624.5.




Step 8. 1-(5-(((2S,4R)-1-((4,4-difluorocyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

[0918] TFA (3 mL) was added to crude 1-(5-(((2S,4R)-1-((4,4-difluorocyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione. The reaction mixture was stirred for 16 h at 90° C. and then concentrated. The crude compound was purified by reverse-phase HPLC using: Mobile Phase: A=0.1% HCOOH in WATER, B=Acetonitrile, Column: X SELECT (250 mm×21.2 mm), 5.0 μm, Flow: 20 mL/min. The collected fractions were concentrated under reduced pressure to obtain the product as a formate salt. The product was dissolved in 20% MeOH in DCM, washed with a solution of saturated aqueous NaHCO₃ and concentrated to give 1-(5-(((2S,4R)-1-((4,4-difluorocyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (554 mg, 1.05 mmol, 48% yield) as an off-white solid.





HPLC: 98.40% [R_t=4.893 min]. LCMS [M+H]⁺: 474.5. ¹H NMR (500 MHz, DMSO-d₆) δ 10.42 (s, 1H), 8.54 (d, J=7.1 Hz, 1H), 7.98 (s, 1H), 7.34 (s, 1H), 6.78 (dd, J=7.3, 1.9 Hz, 1H), 3.76 (t, J=6.7 Hz, 2H), 2.91 (s, 1H), 2.79 (t, J=6.7 Hz, 2H), 2.53 (d, J=9.4 Hz, 2H), 2.45-2.34 (m, 2H), 2.27-2.12 (m, 2H), 2.04-1.92 (m, 2H), 1.92-1.85 (m, 1H), 1.85-1.67

(m, 4H), 1.60-1.47 (m, 2H), 1.47-1.35 (m, 2H), 1.24-1.14 (m, 1H), 1.13-1.00 (m, 2H), 0.88 (d, J=6.5 Hz, 3H).



[0919] The compounds in the following table were prepared by the method of Example 141, using the appropriate commercially available aldehyde in step 7.

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
142	 1-(5-(((2S,4R)-2-methyl-1-((tetrahydro-2H-pyran-4-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	440.5	(500 MHz, DMSO-d ₆) δ 10.42 (s, 1H), 8.54 (d, J = 7.1 Hz, 1H), 7.98 (s, 1H), 7.34 (s, 1H), 6.77 (dd, J = 7.2, 1.9 Hz, 1H), 3.82 (d, J = 11.3 Hz, 2H), 3.76 (t, J = 6.7 Hz, 2H), 3.26 (t, J = 11.5 Hz, 2H), 3.18 (d, J = 5.3 Hz, 1H), 2.91 (d, J = 6.2 Hz, 1H), 2.78 (t, J = 6.7 Hz, 2H), 2.54 (s, 1H), 2.46-2.33 (m, 2H), 2.19 (qd, J = 12.5, 6.9 Hz, 2H), 1.88 (s, 1H), 1.56 (dd, J = 44.9, 16.0 Hz, 4H), 1.47-1.35 (m, 2H), 1.25-1.14 (m, 1H), 1.14-1.00 (m, 2H), 0.88 (d, J = 6.5 Hz, 3H).
143	 1-(5-(((2S,4R)-1-(cyclohexylmethyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	438.3	(400 MHz, METHANOL-d ₄) δ = 8.40 (d, J = 7.2 Hz, 1H), 7.98 (s, 1H), 7.33 (s, 1H), 6.81-6.79 (m, 1H), 3.89-3.86 (m, 2H), 3.04 (br d, J = 12.0 Hz, 1H), 2.89-2.87 (m, 2H), 2.66-2.56 (m, 3H), 2.20-1.84 (m, 4H), 1.75-1.44 (m, 8H), 1.37-1.11 (m, 5H), 1.10-1.03 (m, 3H), 0.99-0.81 (m, 2H)
144	 1-(5-(((2S,4R)-1-(cyclopentylmethyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	424.5	(500 MHz, DMSO-d ₆) δ 10.41 (s, 1H), 8.52 (d, J = 7.1 Hz, 1H), 7.96 (s, 1H), 7.33 (s, 1H), 6.76 (d, J = 7.1 Hz, 1H), 3.75 (t, J = 6.7 Hz, 2H), 3.29 (s, 1H), 2.93 (s, 1H), 2.77 (t, J = 6.7 Hz, 2H), 2.50 (p, J = 1.8 Hz, 3H), 2.28-2.13 (m, 2H), 2.02-1.93 (m, 1H), 1.86 (s, 1H), 1.70-1.05 (m, 13H), 0.86 (d, J = 6.6 Hz, 3H).


-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
145	 1-(5-(((2S,4R)-1-isobutyl-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	398.4	(500 MHz, Methanol-d ₄) δ 8.41 (d, J = 7.2 Hz, 1H), 7.99 (s, 1H), 7.35 (d, J = 1.8 Hz, 1H), 6.82 (dd, J = 7.2, 1.8 Hz, 1H), 3.89 (t, J = 6.8 Hz, 2H), 3.09-2.98 (m, 1H), 2.90 (t, J = 6.8 Hz, 2H), 2.64-2.60 (m, 2H), 2.59-2.51 (m, 2H), 2.30-2.18 (m, 2H), 2.05-1.95 (m, 1H), 1.82-1.72 (m, 1H), 1.68-1.61 (m, 1H), 1.59-1.50 (m, 2H), 1.43-1.33 (m, 1H), 0.99 (d, J = 6.6 Hz, 3H), 0.93 (d, J = 4.0 Hz, 3H), 0.92 (d, J = 4.1 Hz, 3H).
146	 1-(5-(((2S,4R)-1-(cyclobutylmethyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	410.5	(400 MHz, DMSO-d ₆) δ 10.45 (d, J = 3.9 Hz, 1H), 8.58 (d, J = 7.1 Hz, 1H), 8.00 (d, J = 2.2 Hz, 1H), 7.35 (d, J = 5.1 Hz, 1H), 6.79 (d, J = 7.7 Hz, 1H), 3.76 (td, J = 6.8, 3.0 Hz, 2H), 3.54 (d, J = 47.4 Hz, 2H), 3.38-3.17 (m, 2H), 3.16-2.94 (m, 3H), 2.77 (t, J = 6.8 Hz, 2H), 2.73-2.59 (m, 2H), 2.18-1.98 (m, 3H), 1.95-1.53 (m, 7H), 1.25 (m, 3H).
147	 1-(5-(((2S,4R)-1-(cycloheptylmethyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	452.4	(400 MHz, CD ₃ OD) δ 8.43 (d, J = 6.8 Hz, 1H), 8.00 (s, 1H), 7.36 (s, 1H), 6.81 (dd, J = 7.2 Hz, 2.0 Hz, 1H), 3.89 (t, J = 6.4 Hz, 2H), 3.12 (bm, 2H), 2.90-2.68 (m, 5H), 2.20 (bm, 2H), 1.81-1.30 (m, 21H). NH proton not observed due to solvent exchange
148	 1-(5-(((2S,4R)-2-methyl-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	454.1	(300 MHz, CD ₃ OD) δ 8.49 (s, 1H), 8.43 (d, J = 7.2 Hz, 1H), 8.00 (s, 1H), 7.36 (s, 1H), 6.83 (d, J = 6.0 Hz, 1H), 3.93-3.86 (m, 4H), 3.60 (s, 1H), 3.46-3.36 (m, 3H), 3.23 (s, 2H), 3.08-3.04 (m, 3H), 2.88 (t, J = 6.6 Hz, 2H), 2.69-2.67 (m, 2H), 2.02-1.98 (s, 1H), 1.84-1.63 (m, 7H), 1.32-1.30 (m, 5H).

-continued


Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
149	 1-(5-(((2S,4R)-1-((4,4-dimethylcyclohexyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	466.2	(300 MHz, Methanol-d ₄) δ 8.43 (m, 2H), 8.01 (s, 1H), 7.36 (d, J = 1.8 Hz, 1H), 6.83 (dd, J = 7.2, 1.8 Hz, 1H), 3.88 (t, J = 6.8 Hz, 2H), 3.71 (s, 1H), 3.24 (m, 2H), 2.96 (s, 1H), 2.88 (t, J = 6.8 Hz, 2H), 2.69 (s, 2H), 2.22 (s, 1H), 1.73 (dd, J = 59.4, 16.0 Hz, 7H), 1.53-1.16 (m, 10H), 0.93 (d, J = 3.0 Hz, 6H).
150	 1-(5-(((2S,4R)-2-methyl-1-(pyridin-3-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	433.0	(300 MHz, CD ₃ OD) δ 8.60 (s, 1H), 8.53-8.51 (m, 1H), 8.42-8.39 (m, 2H), 7.98 (s, 1H), 7.96-7.94 (m, 1H), 7.48-7.46 (m, 1H), 7.34 (s, 1H), 6.84 (d, J = 7.5 Hz, 1.3 Hz, 1H), 4.04-4.02 (m, 2H), 3.87 (t, J = 6.9 Hz, 2H), 2.91-2.85 (m, 4H), 2.67-2.65 (m, 3H), 2.18-2.16 (s, 2H), 1.71-1.69 (m, 3H), 1.28 (s, 3H).

Example 151. Preparation of 1-(5-(((1-((1*r*,4*r*)-4-hydroxycyclohexyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione






[0920] Prepared using the method of Example 141, steps 7-8, wherein trans-4-(benzyloxy)cyclohexane-1-carbaldehyde [see WO2020/232470, 2020, A1] was used in place of 4,4-difluorocyclohexane-1-carbaldehyde. LCMS [M+H]⁺: 440.1. ¹H NMR (400 MHz, CD₃OD) δ 8.42 (s, 1H), 8.00 (s, 1H), 7.36 (s, 1H), 6.98 (d J=5.6 Hz, 1H), 3.87 (t, J=6.8 Hz, 2H), 3.56-3.48 (m, 2H), 2.93-2.85 (m, 6H), 2.68-2.65 (m, 2H), 1.97-1.80 (m, 7H), 1.58-1.55 (m, 2H), 1.29-1.27 (m, 4H), 1.11-1.08 (m, 2H), NH and OH protons not observed due to solvent exchange.

Example 152. Preparation of 1-(5-(((2S,4R)-1-((3,3-difluorocyclobutyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0921] Prepared from 3-(2,4-dimethoxybenzyl)-1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride (see Example 141, step 6) using the method of Example 1, steps 4 to 5, wherein 3-(bromomethyl)-1,1-difluorocyclobutane was used in place of (bromomethyl)cyclohexane. LCMS [M+H]⁺: 446.3. ¹H NMR (400 MHz, CD₃OD) δ 8.51 (s, 1H), 8.43 (d, J=7.1 Hz, 1H), 8.00 (s, 1H), 7.36 (s, 1H), 6.83 (dd, J=7.2, 1.9 Hz, 1H), 3.89 (t, J=6.8 Hz, 2H), 3.48 (s, 1H), 3.08 (d, J=41.8 Hz, 3H), 2.89 (t, J=6.8 Hz, 2H), 2.76 (d, J=7.6 Hz, 2H), 2.68 (d, J=7.2 Hz, 2H), 2.43 (d, J=29.2 Hz, 3H), 2.18 (d, J=10.5 Hz, 1H), 1.90-1.62 (m, 3H), 1.51 (s, 1H), 1.37-1.17 (m, 4H). NH proton not observed due to solvent exchange.

[0922] The compounds in the following table were prepared from tert-butyl (R)-2-methyl-4-oxopiperidine-1-carboxylate using the method of Example 141, wherein the appropriate commercially available aldehydes were used in step 7.

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
153	 1-(5-(((2R,4S)-1-isobutyl-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)methylpyrimidine-2,4-dione	398.2	(400 MHz, CD ₃ OD) δ 8.49 (s, 1H), 8.44 (d, J = 6.8 Hz, 1H), 7.98 (s, 1H), 7.35 (s, 1H), 6.81 (d, J = 6.0 Hz, 1H), 3.94-3.84 (m, 4H), 3.65 (s, 1H), 3.45-3.40 (m, 2H), 3.17 (s, 2H), 2.91-2.84 (m, 3H), 2.67-2.63 (m, 2H), 2.09 (s, 1H), 2.01 (s, 1H), 1.85-1.58 (m, 5H), 1.39-1.30 (m, 6H).
154	 1-(5-(((2R,4S)-2-methyl-1-((tetrahydro-2H-pyran-4-yl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)methylpyrimidine-2,4-dione	440.1	(400 MHz, CD ₃ OD) δ 8.51 (s, 1H), 8.41 (d, J = 7.2 Hz, 1H), 7.98 (s, 1H), 7.34 (s, 1H), 6.82-6.80 (m, 1H), 3.86 (t, J = 6.4 Hz, 2H), 3.68 (s, 1H), 3.19 (s, 2H), 2.88-2.84 (s, 4H), 2.68-2.66 (m, 2H), 2.19 (s, 1H), 2.08-2.02 (m, 1H), 1.86-1.72 (m, 4H), 1.64-1.60 (m, 2H), 1.33-1.27 (m, 3H), 1.04-1.00 (m, 6H).
155	 1-(5-(((2R,4S)-1-((4,4-difluorocyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)methylpyrimidine-2,4-dione	474.4	(400 MHz, CD ₃ OD) δ 8.52 (s, 1H), 8.41 (d, J = 6.8 Hz, 1H), 7.99 (s, 1H), 7.34 (s, 1H), 6.83-6.81 (m, 1H), 3.87 (t, J = 7.2 Hz, 2H), 2.89-2.86 (m, 3H), 2.66-2.65 (m, 2H), 2.12-2.04 (m, 4H), 1.88-1.70 (m, 7H), 1.50 (brs, 2H), 1.20 (s, 3H).

Example 156. Preparation of 1-(5-((4-(cyclohexylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



Example 156

Step 1: tert-butyl 4-((3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperazine-1-carboxylate

[0923] To a suspension of 1-(5-bromopyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione (45 mg, 0.098 mmol) in toluene (2 mL) and water (0.2 mL) at room temperature was added Cs₂CO₃ (128 mg, 0.392 mmol), potassium {[4-(tert-butoxycarbonyl)-1-piperazinyl]methyl}(trifluoro)borate (60.0 mg, 0.196 mmol) and RuPhos (9.14 mg, 0.020 mmol), followed by Pd(OAc)₂ (2.2 mg, 9.8 μmol). The mixture was stirred at 90° C. for 3 h, then cooled to rt and partitioned between EtOAc and water. The organic layer was separated, washed with brine, dried over sodium sulfate, filtered and concentrated to give crude tert-butyl 4-((3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperazine-1-carboxylate (56 mg, 0.098 mmol). LCMS [M+H]⁺: 579.4. The crude material was used without further purification.

Step 2: 3-(2,4-dimethoxybenzyl)-1-(5-(piperazin-1-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride

[0924] A solution of HCl (4.0 M in dioxane, 2 mL, 8 mmol) was added to tert-butyl 4-((3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperazine-1-carboxylate (55 mg, 0.095 mmol) and the mixture was stirred for 2 h at rt. The reaction was then concentrated to give crude 3-(2,4-dimethoxybenzyl)-1-(5-(piperazin-1-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride (46 mg, 0.095 mmol) which was used without further purification. LCMS [M+H]⁺: 479.4.

Step 3: 1-(5-((4-(cyclohexylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione

[0925] To a solution of 3-(2,4-dimethoxybenzyl)-1-(5-(piperazin-1-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride (22 mg, 0.043 mmol) in DMF (1 mL) was added potassium carbonate (30 mg, 0.21 mmol) and (bromomethyl)cyclohexane (0.012 mL, 0.085 mmol). The mixture was heated at 80° C. for 4 h and then cooled to rt. The mixture was diluted with ethyl acetate and washed sequentially with water and brine. The organic layer was dried over sodium sulfate, filtered and concentrated to give crude 1-(5-((4-(cyclohexylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione (25 mg, 0.043 mmol) which was used without further purification. LCMS [M+H]⁺: 575.4.





Step 4: 1-(5-((4-(cyclohexylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

[0926] TFA (1.5 mL, 19 mmol) was added to crude 1-(5-((4-(cyclohexylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione (24 mg, 0.042 mmol) and the mixture was heated at 80° C. overnight. The mixture was then cooled to rt, concentrated and the residue was dissolved in toluene and concentrated again. The residue was dis-





solved in DMSO, filtered through a 1 micron filter and purified by reverse phase HPLC using ACN/Water/0.1% TFA. The fractions containing the product were combined, frozen and lyophilized to afford a TFA salt of 1-(5-((4-(cyclohexylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione (5.5 mg, 10 μ mol, 24% yield). LCMS $[M+H]^+$: 425.3. 1H NMR (500 MHz, DMSO-d₆) δ 10.48 (s, 1H), 8.66 (d, J=7.2 Hz, 1H),

8.06 (s, 1H), 7.53 (s, 1H), 6.91 (d, J=6.9 Hz, 1H), 4.59 (s, 6H), 3.80 (t, J=6.7 Hz, 2H), 3.72 (s, 1H), 3.47 (s, 1H), 3.01 (s, 4H), 2.80 (t, J=6.7 Hz, 2H), 1.69 (td, J=29.6, 13.7 Hz, 6H), 1.21 (dq, J=36.0, 12.2 Hz, 3H), 0.95 (q, J=11.9 Hz, 2H).





[0927] The compounds in the following table were prepared by the method of Example 156, using the appropriate commercially available halide, mesylate or triflate in step 4.

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
157		461.4	(500 MHz, DMSO-d ₆) δ 10.48 (s, 1H), 8.67 (d, J = 7.2 Hz, 1H), 8.07 (s, 1H), 7.55 (s, 1H), 7.01-6.71 (m, 1H), 3.79 (d, J = 6.7 Hz, 4H), 3.49 (d, J = 29.6 Hz, 4H), 3.02 (s, 6H), 2.80 (t, J = 6.7 Hz, 2H), 2.04 (d, J = 9.1 Hz, 2H), 1.94-1.66 (m, 5H), 1.23 (d, J = 12.9 Hz, 2H).
158		439.3	(500 MHz, DMSO-d ₆) δ 10.48 (s, 1H), 8.66 (d, J = 7.2 Hz, 1H), 8.07 (s, 1H), 7.54 (s, 1H), 6.91 (dd, J = 7.1, 1.8 Hz, 1H), 4.27 (s, 4H), 3.80 (t, J = 6.7 Hz, 4H), 3.57-3.32 (m, 2H), 2.95 (d, J = 71.8 Hz, 5H), 2.80 (t, J = 6.7 Hz, 2H), 1.72 (ddt, J = 13.8, 6.8, 3.1 Hz, 2H), 1.67-1.51 (m, 4H), 1.52-1.38 (m, 4H), 1.27-1.13 (m, 2H).
159		411.2	(500 MHz, DMSO-d ₆) δ 10.48 (s, 1H), 8.67 (d, J = 7.1 Hz, 1H), 8.07 (s, 1H), 7.56 (s, 1H), 6.92 (dd, J = 7.1, 1.9 Hz, 1H), 4.85 (s, 4H), 3.80 (t, J = 6.7 Hz, 4H), 3.51 (s, 1H), 3.09 (s, 5H), 2.80 (t, J = 6.7 Hz, 2H), 2.20 (p, J = 7.7 Hz, 1H), 1.94-1.71 (m, 2H), 1.70-1.39 (m, 4H), 1.37-1.06 (m, 2H).
160		427.3	(500 MHz, DMSO-d ₆) δ 10.49 (s, 1H), 8.68 (dd, J = 7.1, 0.9 Hz, 1H), 8.08 (s, 1H), 7.58 (s, 1H), 6.93 (dd, J = 7.2, 1.9 Hz, 1H), 3.97-3.70 (m, 6H), 3.65-3.36 (m, 2H), 3.31 (td, J = 11.7, 2.0 Hz, 2H), 3.03 (d, J = 59.1 Hz, 6H), 2.80 (t, J = 6.7 Hz, 4H), 2.01 (s, 1H), 1.63 (ddd, J = 12.9, 4.1, 2.0 Hz, 2H), 1.34-1.07 (m, 2H).





-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
161	 1-(5-((4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	441.2	(500 MHz, DMSO-d ₆) δ 10.47 (s, 1H), 8.65 (dd, J = 7.2, 0.9 Hz, 1H), 8.06 (s, 1H), 7.52 (s, 1H), 6.90 (dd, J = 7.1, 1.8 Hz, 1H), 4.45 (t, J = 6.5 Hz, 1H), 3.86-3.78 (m, 4H), 3.45 (t, J = 6.6 Hz, 2H), 3.27 (tdd, J = 11.7, 4.2, 2.0 Hz, 4H), 3.08 (d, J = 47.4 Hz, 4H), 2.80 (t, J = 6.7 Hz, 2H), 1.62-1.51 (m, 5H), 1.36 (q, J = 6.7 Hz, 1H), 1.26-1.08 (m, 4H).
162	 1-(5-((4-((tetrahydro-2H-pyran-3-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	427.2	(500 MHz, DMSO-d ₆) δ 10.49 (s, 1H), 8.67 (d, J = 7.1 Hz, 1H), 8.08 (s, 1H), 7.57 (s, 1H), 6.92 (dd, J = 7.2, 1.8 Hz, 1H), 5.00 (s, 6H), 4.01-3.76 (m, 4H), 3.71 (dt, J = 11.3, 4.2 Hz, 1H), 3.36 (ddd, J = 11.2, 9.7, 3.0 Hz, 2H), 3.31-2.84 (m, 5H), 2.80 (t, J = 6.7 Hz, 2H), 1.98 (s, 1H), 1.87-1.76 (m, 1H), 1.66-1.38 (m, 2H), 1.38-1.17 (m, 1H).
163	 1-(5-((4-propyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	371.2	(400 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 8.63 (d, J = 7.1 Hz, 1H), 8.04 (d, J = 2.4 Hz, 1H), 7.51 (s, 1H), 6.89 (dd, J = 7.2, 1.8 Hz, 1H), 4.41 (s, 3H), 3.78 (dd, J = 7.8, 5.7 Hz, 2H), 3.69 (s, 2H), 3.45 (s, 2H), 3.02 (d, J = 8.5 Hz, 5H), 2.78 (td, J = 6.7, 2.2 Hz, 2H), 1.71-1.47 (m, 2H), 0.90 (t, J = 7.3 Hz, 3H).
164	 1-(5-((4-(pentan-3-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	399.3	(500 MHz, DMSO-d ₆) δ 10.48 (s, 1H), 8.65 (d, J = 6.7 Hz, 1H), 8.06 (d, J = 4.9 Hz, 1H), 7.57 (d, J = 37.0 Hz, 1H), 6.92 (d, J = 7.1 Hz, 1H), 4.56-3.26 (m, 8H), 3.05 (s, 5H), 2.80 (t, J = 6.7 Hz, 2H), 1.77 (s, 2H), 1.61 (s, 2H), 0.94 (t, J = 7.4 Hz, 6H).





-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
165	 1-(5-((4-benzylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4-dione	419.3	(500 MHz, DMSO-d ₆) δ 10.48 (s, 1H), 8.67 (d, J = 7.2 Hz, 1H), 8.08 (s, 1H), 7.56 (s, 1H), 7.48 (d, J = 4.8 Hz, 5H), 6.92 (dd, J = 7.2, 1.8 Hz, 1H), 4.24 (s, 2H), 3.79 (t, J = 6.7 Hz, 4H), 3.00 (d, J = 92.9 Hz, 5H), 2.79 (t, J = 6.7 Hz, 3H), 2.56 (s, 2H).
166	 1-(5-((4-isopropylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4-dione	371.2	(500 MHz, DMSO-d ₆) δ 10.47 (s, 1H), 8.65 (dd, J = 7.1, 0.9 Hz, 1H), 8.06 (s, 1H), 7.54 (s, 1H), 6.92 (dd, J = 7.2, 1.8 Hz, 1H), 4.56 (s, 2H), 3.88-3.65 (m, 4H), 3.60-3.33 (m, 3H), 3.07 (s, 4H), 2.80 (t, J = 6.7 Hz, 2H), 1.26 (d, J = 6.6 Hz, 6H).
167	 1-(5-((4-(2-methylbenzyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4-dione	433.4	(500 MHz, DMSO-d ₆) δ 10.49 (s, 1H), 8.69 (d, J = 7.1 Hz, 1H), 8.09 (s, 1H), 7.59 (s, 1H), 7.40 (s, 1H), 7.28 (dt, J = 14.0, 7.1 Hz, 3H), 7.01-6.83 (m, 1H), 4.27 (s, 4H), 3.80 (t, J = 6.7 Hz, 4H), 3.02 (d, J = 112.7 Hz, 6H), 2.79 (t, J = 6.7 Hz, 2H), 2.38 (s, 3H).
168	 1-(5-((4-(4-fluorobenzyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4-dione	437.2	(500 MHz, DMSO-d ₆) δ 10.48 (s, 1H), 8.67 (d, J = 7.2 Hz, 1H), 8.07 (s, 1H), 7.63-7.45 (m, 3H), 7.30 (t, J = 8.6 Hz, 2H), 6.92 (dd, J = 7.2, 1.8 Hz, 1H), 4.20 (s, 2H), 3.79 (t, J = 6.7 Hz, 6H), 3.08 (s, 6H), 2.79 (t, J = 6.7 Hz, 2H).



-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
169	 1-(5-((4-(2-fluoro-2-methylpropyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	403.2	(400 MHz, DMSO-d ₆) δ 10.50 (s, 1H), 8.73 (d, J = 7.1 Hz, 1H), 8.12 (s, 1H), 7.65 (d, J = 34.0 Hz, 1H), 6.98 (d, J = 7.2 Hz, 1H), 3.81 (q, J = 6.0, 5.3 Hz, 6H), 3.07 (s, 8H), 2.79 (dd, J = 7.4, 6.0 Hz, 2H), 1.34 (d, J = 21.4 Hz, 6H).
170	 1-(5-((4-(3-fluorobenzyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	437.3	(500 MHz, DMSO-d ₆) δ 10.49 (s, 1H), 8.69 (d, J = 7.1 Hz, 1H), 8.09 (s, 1H), 7.60 (s, 1H), 7.50 (td, J = 7.9, 6.1 Hz, 1H), 7.37-7.15 (m, 3H), 6.94 (dd, J = 7.3, 1.8 Hz, 1H), 4.67 (s, 2H), 4.05 (d, J = 80.5 Hz, 4H), 3.80 (t, J = 6.7 Hz, 2H), 2.98 (d, J = 72.5 Hz, 6H), 2.79 (t, J = 6.7 Hz, 2H).
171	 1-(5-((4-(2,2,2-trifluoroethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	411.3	(500 MHz, DMSO-d ₆) δ 10.54 (s, 1H), 8.77 (d, J = 7.3 Hz, 1H), 8.15 (s, 1H), 7.72 (s, 1H), 6.98 (d, J = 7.2 Hz, 1H), 4.39 (s, 2H), 3.82 (t, J = 6.7 Hz, 2H), 3.33 (s, 4H), 3.10 (dd, J = 47.4, 18.5 Hz, 4H), 2.79 (t, J = 6.7 Hz, 2H), 2.68 (d, J = 37.2 Hz, 2H).
172	 1-(5-((4-((1-(trifluoromethyl)cyclopropyl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	451.2	(500 MHz, DMSO-d ₆) δ 10.54 (s, 1H), 8.76 (d, J = 7.1 Hz, 1H), 8.15 (s, 1H), 7.73 (s, 1H), 7.00 (dd, J = 7.2, 1.8 Hz, 1H), 4.53 (d, J = 195.2 Hz, 8H), 3.83 (t, J = 6.7 Hz, 2H), 3.21 (d, J = 131.0 Hz, 4H), 2.80 (t, J = 6.7 Hz, 2H), 1.01 (s, 2H), 0.79 (s, 2H).

-continued


Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
173	 1-(5-((4-(2,2,3,3-tetrafluoropropyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	443.3	(500 MHz, DMSO-d ₆) δ 10.46 (s, 1H), 8.70 (d, J = 7.1 Hz, 1H), 8.07 (s, 1H), 7.66 (s, 1H), 6.91 (d, J = 7.2 Hz, 1H), 6.44 (tt, J = 52.3, 5.3 Hz, 1H), 4.31 (s, 2H), 3.75 (t, J = 6.7 Hz, 2H), 3.31 (s, 2H), 3.05 (t, J = 15.1 Hz, 4H), 2.93 (d, J = 12.8 Hz, 2H), 2.72 (t, J = 6.7 Hz, 2H), 2.67-2.53 (m, 2H).
174	 1-(5-((4-((3,3-difluorocyclobutyl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	433.4	(400 MHz, Methanol-d ₄) δ 8.48 (d, J = 7.2 Hz, 1H), 8.04 (s, 1H), 7.54 (s, 1H), 6.99 (dd, J = 7.2, 1.9 Hz, 1H), 4.88 (br. s, 2H), 3.90 (t, J = 6.8 Hz, 2H), 3.76 (s, 2H), 3.58-3.37 (m, 2H), 3.30-3.22 (m, 1H), 3.22-3.01 (m, 4H), 2.89 (t, J = 6.8 Hz, 2H), 2.87-2.74 (m, 2H), 2.67-2.52 (m, 2H), 2.52-2.36 (m, 2H). NH proton not observed due to solvent exchange
175	 1-(5-((4-(1-(pyridin-3-yl)ethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	434.3	(500 MHz, DMSO-d ₆) δ 10.51 (s, 1H), 8.79-8.60 (m, 3H), 8.14-7.98 (m, 2H), 7.76-7.58 (m, 2H), 6.95 (dd, J = 7.1, 1.9 Hz, 1H), 4.14 (s, 6H), 3.80 (t, J = 6.7 Hz, 2H), 3.52-2.84 (m, 5H), 2.79 (t, J = 6.7 Hz, 2H), 1.48 (s, 3H).
176	 1-(5-((4-(pyridin-4-ylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	420.3	(500 MHz, DMSO-d ₆) δ 10.53 (s, 1H), 8.80-8.61 (m, 3H), 8.14 (s, 1H), 7.70 (s, 3H), 6.99 (d, J = 7.0 Hz, 1H), 4.45 (s, 4H), 4.28 (s, 4H), 3.82 (t, J = 6.7 Hz, 2H), 3.00 (s, 4H), 2.80 (t, J = 6.7 Hz, 2H).

-continued


Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
177		413.4	(500 MHz, DMSO-d ₆) δ 10.48 (s, 1H), 8.67 (d, J = 7.1 Hz, 1H), 8.07 (s, 1H), 7.56 (s, 1H), 6.92 (dd, J = 7.2, 1.9 Hz, 1H), 4.61 (s, 4H), 3.92-3.71 (m, 6H), 3.64 (dt, J = 8.4, 7.4 Hz, 1H), 3.57-3.43 (m, 1H), 3.39 (dd, J = 8.6, 6.4 Hz, 1H), 3.08 (s, 5H), 2.80 (t, J = 6.7 Hz, 2H), 2.64-2.56 (m, 1H), 2.07 (td, J = 12.7, 7.7 Hz, 1H), 1.59 (dq, J = 12.3, 7.4 Hz, 1H).
178		433.3	(400 MHz, CDCl ₃) δ 8.42-8.29 (m, 1H), 7.93 (s, 1H), 7.66 (br s, 1H), 7.34-7.27 (m, 3H), 7.25-7.17 (m, 3H), 6.91-6.89 (m, 1H), 3.90 (t, J = 6.8 Hz, 2H), 3.56 (s, 2H), 2.97-2.86 (m, 4H), 2.86-2.51 (m, 10H).

Examples 179 and 180. Preparation of 1-(5-(((1*r*,4*r*)-4-methoxycyclohexyl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-*a*]pyridin-3-yl)dihydropyrimidine-2,4(1*H*,3*H*)-dione (Example 179) and 1-(5-(((1*s*,4*s*)-4-methoxycyclohexyl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-*a*]pyridin-3-yl)dihydropyrimidine-2,4(1*H*,3*H*)-dione (Example 180)

(Example 179)



(Example 180)



[0928] Prepared using the method of Example 156 using a commercially available mixture of *cis* and *trans* 1-(bromomethyl)-4-methoxycyclohexane in place of (bromomethyl)cyclohexane. The stereoisomers were purified after the final step by reverse-phase HPLC (eluting with using ACN/Water/0.1% TFA).


Example 179. 1-(5-(((1*r*,4*r*)-4-methoxycyclohexyl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-*a*]pyridin-3-yl)dihydropyrimidine-2,4(1*H*,3*H*)-dione

[0929] Eluted first, minor isomer. LCMS [M+H]⁺: 455.2. ¹H NMR (500 MHz, DMSO-d₆) δ 10.48 (s, 1H), 8.66 (d, J=7.2 Hz, 1H), 8.07 (s, 1H), 7.54 (s, 1H), 6.91 (dd, J=7.3, 1.8 Hz, 1H), 4.59 (s, 4H), 3.80 (t, J=6.7 Hz, 4H), 3.42 (d, J=45.9 Hz, 1H), 3.24 (s, 3H), 3.06 (ddd, J=14.6, 10.7, 4.2 Hz, 6H), 2.80 (t, J=6.7 Hz, 2H), 2.01 (d, J=12.1 Hz, 2H), 1.86-1.54 (m, 3H), 1.20-0.77 (m, 4H).

Example 180. 1-(5-(((1*s*,4*s*)-4-methoxycyclohexyl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-*a*]pyridin-3-yl)dihydropyrimidine-2,4(1*H*,3*H*)-dione

[0930] Eluted second, major isomer. LCMS [M+H]⁺: 455.2. ¹H NMR (500 MHz, DMSO-d₆) δ 10.48 (s, 1H), 8.66 (d, J=7.2 Hz, 1H), 8.07 (s, 1H), 7.54 (s, 1H), 6.91 (dd, J=7.1, 1.8 Hz, 1H), 4.60 (s, 4H), 3.80 (t, J=6.7 Hz, 4H), 3.47 (s, 1H), 3.38 (d, J=4.9 Hz, 1H), 3.21 (s, 3H), 3.02 (s, 5H), 2.80 (t, J=6.7 Hz, 2H), 1.91-1.62 (m, 3H), 1.56-1.36 (m, 4H), 1.32-1.18 (m, 2H).

Example 181. Preparation of 1-(5-((4-(2-(4,4-difluorocyclohexyl)propan-2-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



Example 181

Step 1. (4-benzylpiperazin-1-yl)(4,4-difluorocyclohexyl)methanone

[0931] HATU (8.26 g, 21.9 mmol) and DIPEA (9.53 mL, 54.75 mmol) were added to a solution of 4,4-difluorocyclohexanecarboxylic acid (3.0 g, 18.25 mmol) in THF (10 mL) at 0° C. The mixture was stirred for 10 min and then 1-benzylpiperazine (3.2 g, 18.3 mmol) was added and the reaction was stirred for 16 h at rt. The reaction mixture was quenched with water and extracted with EtOAc. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated. The crude material was purified by silica

gel chromatography (eluting with 50% EtOAc in hexane) to afford (4-benzylpiperazin-1-yl)(4,4-difluorocyclohexyl)methanone (1.5 g, 4.65 mmol, 25% yield). LCMS [M+H]⁺: 323.5.

Step 2. 1-benzyl-4-(2-(4,4-difluorocyclohexyl)propan-2-yl)piperazine

[0932] To a stirred solution of (4-benzylpiperazin-1-yl)(4,4-difluorocyclohexyl)methanone (1.5 g, 4.65 mmol) in THF (10 mL) was added ZrCl₄ (1.84 g, 4.65 mmol) at -20° C. and the mixture was stirred for 30 min. A solution of MeMgBr·Et₂O (9.4 mL, 28.2 mmol, 3.0 M) was added and the mixture was stirred for 10 min at -20° C. and then at rt for 16 h. After completion, the reaction was quenched with water (10 mL) and extracted with EtOAc. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated. The crude material was purified by silica gel chromatography (eluting with 10-15% EtOAc in hexanes) to afford 1-benzyl-4-(2-(4,4-difluorocyclohexyl)propan-2-yl)piperazine (500 mg, 1.49 mmol, 31% yield).

Step 3. 1-(2-(4,4-difluorocyclohexyl)propan-2-yl)piperazine


[0933] To a stirred solution of 1-benzyl-4-(2-(4,4-difluorocyclohexyl)propan-2-yl)piperazine (500 mg, 1.49 mmol) in EtOAc (15 mL) under an inert atmosphere was added Pd/C (100 mg) at rt. The flask was evacuated and refilled with hydrogen from a balloon and stirred at RT for 36 h. The reaction was then purged with argon and filtered through celite. The filtrate was concentrated to give crude 1-(2-(4,4-difluorocyclohexyl)propan-2-yl)piperazine (400 mg, crude). The material was used without further purification.

Step 4. potassium ((4-(2-(4,4-difluorocyclohexyl)propan-2-yl)piperazin-1-yl)methyl)trifluoroborate

[0934] To a stirred solution of 1-(2-(4,4-difluorocyclohexyl)propan-2-yl)piperazine (400 mg, 1.62 mmol) in THF (10 mL) at rt was added K₂CO₃ (448 mg, 3.25 mmol) and potassium (bromomethyl)trifluoroborate (326 mg, 1.62 mmol). The reaction was stirred for 12 h at 80° C. and then cooled to rt and concentrated to afford potassium ((4-(2-(4,4-difluorocyclohexyl)propan-2-yl)piperazin-1-yl)methyl)trifluoroborate (1.5 g, crude). The material was used without further purification.


[0935] Step 5: 1-(5-((4-(2-(4,4-difluorocyclohexyl)propan-2-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione was prepared from potassium ((4-(2-(4,4-difluorocyclohexyl)propan-2-yl)piperazin-1-yl)methyl)trifluoroborate by the method of Example 156, steps 1 and 4, wherein potassium ((4-(2-(4,4-difluorocyclohexyl)propan-2-yl)piperazin-1-yl)methyl)trifluoroborate was used in place of potassium {4-(tert-butoxycarbonyl)-1-piperazinyl)methyl}(trifluoro)borate. LCMS [M+H]⁺: 489.2. ¹H NMR (400 MHz, Methanol-d₄) δ 8.51 (d, J=6.9 Hz, 1H), 8.06 (s, 1H), 7.54 (d, J=14.9 Hz, 1H), 7.00 (dd, J=7.4, 1.8 Hz, 1H), 3.91 (t, J=6.7 Hz, 2H), 3.48 (s, 3H), 3.26-3.10 (m, 2H), 2.89 (t, J=6.7 Hz, 2H), 2.67 (s, 3H), 2.16 (d, J=23.5 Hz, 2H), 1.82 (d, J=14.0 Hz, 4H), 1.55-1.18 (m, 7H). 4 protons not integrated due to peak broadening. NH proton not observed due to solvent exchange.

Example 182. Preparation of 1-(5-((4-(2-(tetrahydro-2H-pyran-4-yl)propan-2-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0936] Prepared using the method of 181, wherein tetrahydro-2H-pyran-4-carboxylic acid was used in place of 4,4-difluorocyclohexane-1-carboxylic acid. LCMS [M+H]⁺: 474.8. ¹H NMR (400 MHz, DMSO-d₆) δ 10.46 (s, 1H), 8.74 (s, 1H), 8.62 (s, 1H), 8.13 (s, 1H), 8.03 (s, 1H), 7.48 (s, 1H), 6.90 (s, 1H), 3.89 (d, J=11.0 Hz, 2H), 3.78 (t, J=6.5 Hz, 2H), 3.63 (s, 1H), 3.32 (s, 9H), 2.78 (t, J=6.7 Hz, 2H), 1.85 (d, J=110.7 Hz, 2H), 1.52 (d, J=12.5 Hz, 2H), 1.43-1.07 (m, 6H), 0.86 (s, 3H).

Example 183. Preparation of 1-(5-((4-(2-hydroxy-2-methylpropyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0937] Prepared using the method of Example 106, wherein 3-(2,4-dimethoxybenzyl)-1-(5-(piperazin-1-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione was used in place of 3-(2,4-dimethoxybenzyl)-1-(5-((1-(2-hydroxy-2-methylpropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione. LCMS [M+H]⁺: 401.4. ¹H NMR (400 MHz, DMSO-d₆) δ 10.46 (s, 1H), 8.66 (d, J=7.1 Hz, 1H), 8.07 (d, J=1.7 Hz, 1H), 7.56 (s, 1H), 6.92 (dd, J=7.3, 1.9 Hz, 1H), 4.70 (s, 4H), 3.96-3.67 (m, 4H), 3.29 (s, 2H), 2.97 (d, J=36.7 Hz, 5H), 2.79 (t, J=6.8 Hz, 2H), 1.22 (d, J=1.8 Hz, 6H).

Example 184. Preparation of 1-(5-((4-(1-(trifluoromethyl)cyclopropane-1-carbonyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0938] Prepared using the method of Example 98, wherein 3-(2,4-dimethoxybenzyl)-1-(5-(piperazin-1-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride was used in place of 3-(2,4-dimethoxybenzyl)-1-(5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride and 1-(trifluoromethyl)cyclopropane-1-carboxylic acid was used in place of isobutyric acid. LCMS [M+H]⁺: 465.2. ¹H NMR (400 MHz, DMSO-d₆) δ 10.51 (d, J=2.2 Hz, 1H), 8.76 (d, J=7.1 Hz, 1H), 8.14 (d, J=1.9 Hz, 1H), 7.72 (s, 1H), 6.99 (dd, J=7.2, 2.0 Hz, 1H), 4.34 (s, 2H), 3.82 (td, J=6.4, 1.8 Hz, 2H), 3.64 (s, 4H), 3.19 (s, 4H), 2.79 (t, J=6.7 Hz, 2H), 1.47-0.84 (m, 4H).

Example 185. Preparation of 1-(5-((4-(isopropylsulfonyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[0939] Prepared using the method of Example 80, wherein 3-(2,4-dimethoxybenzyl)-1-(5-(piperazin-1-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride was used in place of 3-(2,4-dimethoxybenzyl)-1-(5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride and propane-2-sulfonyl chloride was used in place of cyclohexylmethanesulfonyl chloride. LCMS [M+H]⁺: 435.4. ¹H NMR (500 MHz, DMSO-d₆) δ 10.45 (s, 1H), 8.67 (d, J=6.7 Hz, 1H), 8.06 (s, 1H), 7.62 (s, 1H), 6.90 (d, J=7.4 Hz, 1H), 4.27 (bs, 4H), 3.74 (t, J=6.6 Hz, 2H), 3.23-2.83 (m, 4H), 2.72 (t, J=6.7 Hz, 2H), 1.16 (d, J=6.8 Hz, 6H) (missing proton obscured by water peak).

Example 186. Preparation of 1-(5-((4-isobutylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



Step 1. 1-(5-(piperazin-1-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

[0940] TFA (4 ml, 52 mmol) was added to tert-butyl 4-((3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-3-yl)methyl)piperazine-1-carboxylate (200 mg, 0.346 mmol). The mixture was heated overnight at in a sealed vial at 85° C. The mixture was then cooled to rt and, concentrated and azeotropically dried with toluene to provide crude 1-(5-(piperazin-1-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione trifluoroacetate which was used without further purification. LCMS [M+H]⁺: 329.2.





Step 2: 1-(5-((4-isobutylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

[0941] Isobutyraldehyde (10 mg, 0.14 mmol) and triethylamine (0.014 mL, 0.10 mmol) were added to a solution of 1-(5-(piperazin-1-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione trifluoroacetate (30 mg, 0.068 mmol) in DCM (2 mL) and MeOH (2 mL). The reaction mixture was stirred at rt for 10 min and then sodium triacetoxyborohydride (43 mg, 0.20 mmol) was added. The reaction mixture was stirred overnight at rt and then quenched with a solution of saturated aqueous NaHCO₃ and extracted three times with DCM. The combined organic extracts were washed with brine, dried over Na₂SO₄, filtered and concentrated. The residue was dissolved in DMSO, filtered through a 1 micron filter and purified by reverse phase HPLC using ACN/Water/0.1% TFA. The fractions containing the product were combined, frozen and lyophilized to afford a TFA salt of 1-(5-((4-isobutylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (12 mg, 0.023 mmol, 33% yield). LCMS [M+H]⁺: 385.3. ¹H NMR (500 MHz, DMSO-d₆) δ 10.46 (s, 1H), 8.56 (d, J=7.1 Hz, 1H), 8.05 (s, 1H), 7.50 (s, 1H), 6.89 (d, J=7.1 Hz, 1H), 3.79 (t, J=6.7 Hz, 2H), 3.72-3.55 (m, 4H), 3.41 (s, 4H), 3.00 (d, J=39.0 Hz, 4H), 2.79 (t, J=6.7 Hz, 2H), 2.11-1.98 (m, 1H), 0.94 (d, J=6.6 Hz, 6H).


[0942] The compounds in the following table were prepared by the method of Example 186, using the appropriate commercially available aldehyde in step 2.

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
187	1-(5-((4-(2-cyclohexylethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	439.3	(500 MHz, DMSO-d ₆) δ 10.38 (s, 1H), 8.56 (d, J = 7.1 Hz, 1H), 7.97 (s, 1H), 7.43 (s, 1H), 6.81 (dd, J = 7.2, 1.8 Hz, 1H), 3.71 (t, J = 6.7 Hz, 2H), 3.59 (s, 4H), 3.09-2.80 (m, 6H), 2.71 (t, J = 6.7 Hz, 2H), 2.37 (d, J = 31.0 Hz, 2H), 1.64-1.50 (m, 5H), 1.49-1.38 (m, 2H), 1.12 (tddd, J = 24.2, 21.4, 9.8, 6.5 Hz, 4H), 0.84 (q, J = 11.2 Hz, 2H).

-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
188	 1-(5-((4-(cyclobutylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	397.3	(500 MHz, DMSO-d ₆) δ 10.46 (s, 1H), 8.63 (d, J = 7.1 Hz, 1H), 8.04 (s, 1H), 7.50 (s, 1H), 6.88 (dd, J = 7.2, 1.8 Hz, 1H), 3.78 (t, J = 6.7 Hz, 4H), 3.35 (s, 2H), 3.14 (s, 2H), 2.99 (s, 4H), 2.79 (t, J = 6.7 Hz, 2H), 2.73-2.60 (m, 1H), 2.40 (d, J = 29.5 Hz, 2H), 2.13-2.01 (m, 2H), 1.95-1.84 (m, 1H), 1.84-1.69 (m, 3H).
189	 1-(5-((4-(cyclopropylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	383.4	(500 MHz, DMSO-d ₆) δ 10.48 (s, 1H), 8.66 (dd, J = 7.2, 0.9 Hz, 1H), 8.06 (s, 1H), 7.54 (s, 1H), 6.91 (dd, J = 7.2, 1.9 Hz, 1H), 3.84-3.72 (m, 5H), 3.55 (s, 3H), 3.25-2.94 (m, 6H), 2.80 (t, J = 6.7 Hz, 2H), 1.05 (dh, J = 10.5, 2.8 Hz, 1H), 0.74-0.54 (m, 2H), 0.44-0.21 (m, 2H).
190	 1-(5-((4-(3-methylbenzyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	433.2	(500 MHz, DMSO-d ₆) δ 10.39 (s, 1H), 8.57 (d, J = 7.2 Hz, 1H), 7.98 (s, 1H), 7.44 (s, 1H), 7.23 (d, J = 36.8 Hz, 4H), 6.82 (d, J = 7.3 Hz, 1H), 4.43-4.03 (m, 6H), 3.71 (t, J = 6.7 Hz, 5H), 3.19 (s, 1H), 2.95 (s, 2H), 2.71 (t, J = 6.7 Hz, 2H), 2.26 (s, 3H).
191	 1-(5-((4-(pyridin-3-ylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	420.3	(500 MHz, Methanol-d ₄) δ 8.66 (s, 1H), 8.61 (d, J = 5.4 Hz, 1H), 8.48 (d, J = 7.2 Hz, 1H), 8.23 (s, 1H), 8.01 (s, 1H), 7.74 (d, J = 6.8 Hz, 1H), 7.59 (s, 1H), 6.90 (dd, J = 7.3, 1.9 Hz, 1H), 4.14 (s, 2H), 3.88 (s, 2H), 3.82 (t, J = 6.7 Hz, 2H), 3.07 (s, 4H), 2.86 (s, 3H), 2.79 (t, J = 6.7 Hz, 3H).

Example 192. Preparation of (S)-1-(5-((4-(cyclohexylmethyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



Step 1. tert-butyl (S)-4-((3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperazine-1-carboxylate

[0943] To a suspension of 1-(5-bromopyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione (1.2 g, 2.61 mmol) in toluene (20 mL) and water (2 mL) at room temperature was added Cs_2CO_3 (2.55 g, 7.84 mmol), tert-butyl (S)-2-methyl-4-((trifluoro[4-borane]methyl)piperazine-1-carboxylate, potassium salt (3.35 g, 10.45 mmol) [see *ChemMedChem*, 2016, 11, 2640-2648] and RuPhos (242 mg, 0.52 mmol), followed by $\text{Pd}(\text{OAc})_2$ (59 mg, 0.26 mmol). The mixture was stirred at 100°C overnight, then cooled to rt and partitioned between EtOAc and water. The organic layer was separated, washed with brine, dried over sodium sulfate, filtered and concentrated. Silica gel column chromatography [eluted with 0-100% EtOAc/EtOH (3:1) in heptane] provided tert-butyl (S)-4-((3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperazine-1-carboxylate, (1.12 g, 2.61 mmol, 71% yield) as an off-white foamy solid. LCMS $[\text{M}+\text{H}]^+$: 593.4.

Step 2. (S)-3-(2,4-dimethoxybenzyl)-1-(5-((3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione trifluoroacetate

[0944] TFA (4 mL, 2 mmol) was added to a solution of tert-butyl (S)-4-((3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperazine-1-carboxylate (1.2 g, 2.0 mmol) in DCM (12 mL) and the mixture was stirred for 2 h at rt. The reaction was then concentrated and dried azeotropically with toluene to give crude (S)-3-(2,4-dimethoxybenzyl)-1-(5-((3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione trifluoroacetate (1.2 g, 1.4 mmol) which was used without further purification. LCMS $[\text{M}+\text{H}]^+$: 493.2.

Step 3. (S)-1-(5-((4-(cyclohexylmethyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione




[0945] To a solution of (S)-3-(2,4-dimethoxybenzyl)-1-(5-((3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (100 mg, 0.20 mmol) in DCE (2 mL) was added cyclohexanecarbaldehyde (21 mg, 0.20 mmol), $\text{NaBH}(\text{OAc})_3$ (120 mg, 0.60 mmol), 4 A MS (100 mg) and DIPEA (113 mg, 0.15 mL, 0.95 mmol). The reaction was stirred at rt for 2 h. The suspension was filtered through a pad of Celite and the filtrate was diluted with a solution of saturated aqueous NaHCO_3 and extracted with DCM. The organic layer was dried over Na_2SO_4 , filtered and concentrated to give (S)-1-(5-((4-(cyclohexylmethyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione (200 mg, 0.34 mmol, 72% purity). The crude product was used in the next step without any other purification. LCMS $[\text{M}+\text{H}]^+$: 589.2.

[0946] Step 4: (S)-1-(5-((4-(cyclohexylmethyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 192) was prepared using the method of Example 156, step 4, wherein



(S)-1-(5-((4-(cyclohexylmethyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione was used in place of 1-(5-((4-(cyclohexylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione. LCMS [M+H]⁺: 439.1. ¹H NMR (400 MHz, DMSO-d₆) δ 10.43 (s, 1H), 8.57-8.55 (m, 1H), 8.00 (s, 1H), 7.43 (s, 1H), 6.88-6.86 (m, 1H), 3.76 (s, 2H), 3.45-3.44 (m, 2H), 2.80-2.72 (m, 3H), 2.57 (br s, 1H),

2.47-2.40 (m, 1H), 2.35-2.26 (m, 1H), 2.23-2.08 (m, 2H), 2.03-1.77 (m, 3H), 1.63 (br s, 4H), 1.42-1.39 (m, 1H), 1.30-1.02 (m, 4H), 0.92-0.91 (m, 3H), 0.87-0.70 (s, 2H).


[0947] The compounds in the following table were prepared from 3-(2,4-dimethoxybenzyl)-1-(5-(piperazin-1-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride by the method of Example 192, steps 3-4 using the appropriate commercially available aldehyde in step 3.

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
193	 1-(5-((4-(((2R,6S)-2,6-dimethyltetrahydro-2H-pyran-4-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	455.2	(400 MHz, CD ₃ OD) δ 8.46 (d, J = 7.2 Hz, 1H), 8.44 (s, 1H), 8.02 (s, 1H), 7.51 (s, 1H), 6.99 (d, J = 4.4 Hz, 1H), 3.88 (t, J = 6.6 Hz, 2H), 3.69-3.48 (m, 4H), 3.08 (m, 4H), 2.90-2.77 (m, 7H), 2.31 (s, 1H), 2.19 (s, 1H), 1.74-1.42 (m, 3H), 1.18-1.11 (m, 6H), 0.88-0.84 (m, 1H).
194	 1-(5-((4-(((3r,5r,7r)-adamantan-1-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	477.2	(400 MHz, Methanol-d ₄) δ 8.49 (s, 1H), 8.46 (d, J = 7.2 Hz, 1H), 8.02 (s, 1H), 7.51 (s, 1H), 6.99 (dd, J = 7.4, 1.6 Hz, 1H), 3.89 (t, J = 6.7 Hz, 2H), 3.64 (s, 2H), 2.89 (t, J = 6.8 Hz, 2H), 2.78 (s, 4H), 2.63 (s, 4H), 2.23 (s, 2H), 1.96 (s, 3H), 1.76 (d, J = 12.4 Hz, 3H), 1.68 (d, J = 12.2 Hz, 3H), 1.57 (d, J = 3.0 Hz, 6H). NH proton not observed due to solvent exchange
195	 1-(5-((4-((1-methyl-1H-pyrazol-5-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	423.2	(400 MHz, CD ₃ OD) δ 8.58 (d, J = 7.6 Hz, 1H), 8.11 (s, 1H), 7.70 (s, 1H), 7.45 (d, J = 1.6 Hz, 1H), 7.02 (d, J = 7.6 Hz, 1H), 6.34 (d, J = 1.6 Hz, 1H), 4.24 (s, 2H), 3.94-3.91 (m, 7H), 3.23 (brs, 4H), 2.91-2.88 (m, 6H).



-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
196	 1-(5-((4-((1-methyl-1H-imidazol-4-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	422.8	(400 MHz, Methanol-d ₄) δ 8.86 (s, 1H), 8.61 (d, J = 7.2 Hz, 1H), 8.13 (s, 1H), 7.80 (d, J = 1.8 Hz, 1H), 7.51 (s, 1H), 7.01 (dd, J = 7.6, 1.9 Hz, 1H), 4.42 (s, 2H), 3.92 (m, 5H), 3.75 (s, 2H), 3.38 (bs, 3H), 2.90 (bm, 7H). NH proton not observed due to solvent exchange
197	 1-(5-((4-(thiazol-4-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	425.8	(400 MHz, Methanol-d ₄) δ 9.11 (d, J = 1.9 Hz, 1H), 8.50 (d, J = 7.1 Hz, 1H), 8.06 (s, 1H), 7.82 (d, J = 2.0 Hz, 1H), 7.58 (s, 1H), 6.99 (dd, J = 7.1, 1.9 Hz, 1H), 4.44 (s, 2H), 3.96-3.84 (m, 4H), 3.19-2.66 (m, 8H). 2 protons not integrated due to broadening. NH proton not observed due to solvent exchange.


[0948] The compounds in the following table were prepared by the method of Example 192, using the appropriate commercially available aldehyde in step 3.

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
198	 (S)-1-(5-((4-(cycloheptylmethyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	453.0	(400 MHz, CD ₃ OD): δ 8.45 (d, J = 6.8 Hz, 1H), 8.02 (s, 1H), 7.50 (s, 1H), 6.99 (d, J = 6.8 Hz, 1H), 3.89 (t, J = 7.2 Hz, 2H), 3.63-3.54 (s, 2H), 2.91-2.79 (m, 6H), 3.14-3.07 (m, 2H), 2.87 (t, J = 6.8 Hz, 2H), 2.48-2.16 (m, 1H), 1.86-1.47 (m, 12H), 1.21-1.14 (m, 5H), ppm. NH proton not observed due to solvent exchange.

-continued





Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
199	 1-(5-(((3S)-4-(((2R,6S)-2,6-dimethyltetrahydro-2H-pyran-4-yl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	469.3	(400 MHz, Methanol-d ₄) δ 8.56 (dd, J = 7.0, 2.7 Hz, 1H), 8.09 (d, J = 2.0 Hz, 1H), 7.67 (s, 1H), 7.00 (dd, J = 7.3, 1.8 Hz, 1H), 4.14 (s, 1H), 3.91 (t, J = 6.8 Hz, 2H), 3.99-3.37 (m, 8H), 3.12 (s, 3H), 2.89 (t, J = 6.8 Hz, 2H), 2.46-1.95 (m, 1H), 1.81-1.49 (m, 2H), 1.43 (d, J = 6.6 Hz, 3H), 1.29 (s, 1H), 1.22-1.10 (m, 6H), 1.04-0.84 (m, 2H). NH proton not observed due to solvent exchange.
200	 (S)-1-(5-(((4-(4,4-dimethylcyclohexyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	467.3	(400 MHz, CD ₃ OD) δ 8.46-8.43 (m, 2H), 8.02 (s, 1H), 7.48 (s, 1H), 6.99-6.97 (m, 1H), 3.88 (t, J = 7.2 Hz, 2H), 3.64-3.62 (s, 2H), 3.48-3.40 (m, 2H), 3.12-2.40 (m, 8H), 1.68-1.23 (m, 12H), 0.92 (d, J = 2.4 Hz, 6H). NH proton not observed due to solvent exchange.

Example 201. Preparation of 1-(5-(((S)-4-(((1R,4S)-4-hydroxycyclohexyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione







[0949] Prepared using the method of Example 192, steps 3-4, wherein trans-4-(benzyloxy)cyclohexane-1-carbaldehyde (see WO2020/232470, 2020, A1 which is incorporated herein by reference) was used in place of cyclohexanecarbaldehyde. LCMS [M+H]⁺: 455.2. ¹H NMR (400 MHz, Methanol-d₄) δ 8.44 (d, J=7.1 Hz, 1H), 8.00 (d, J=11.5 Hz, 1H), 7.38 (s, 1H), 6.83 (d, J=6.2 Hz, 1H), 3.88 (t, J=6.8 Hz, 2H), 3.63-3.44 (m, 3H), 2.96-2.85 (m, 6H), 2.73-2.64 (m, 2H), 2.08-1.72 (m, 7H), 1.58 (q, J=12.1, 11.1 Hz, 2H), 1.37-1.22 (m, 4H), 1.11 (q, J=12.5 Hz, 2H), NH and OH protons not observed due to solvent exchange.


[0950] The compounds in the following table were prepared from (S)-3-(2,4-dimethoxybenzyl)-1-(5-((3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione trifluoroacetate by the method of Example 156, steps 3-4 using the appropriate commercially available halide, mesylate or triflate in step 3.

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
202	 (S)-1-(5-((3-methyl-4-((tetrahydro-2H-pyran-4-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)methylpyrimidine-2,4(1H,3H)-dione	441.3	(500 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 8.57 (d, J = 7.1 Hz, 1H), 8.01 (s, 1H), 7.44 (s, 1H), 6.88 (dd, J = 7.2, 1.8 Hz, 1H), 3.81 (d, J = 11.6 Hz, 2H), 3.77 (t, J = 6.7 Hz, 2H), 3.49-3.38 (m, 2H), 3.30-3.19 (m, 2H), 2.78 (t, J = 6.7 Hz, 3H), 2.56 (d, J = 12.1 Hz, 2H), 2.46 (d, J = 12.2 Hz, 1H), 2.36 (d, J = 12.4 Hz, 1H), 2.18 (t, J = 10.5 Hz, 2H), 1.92 (s, 2H), 1.66 (d, J = 12.4 Hz, 2H), 1.51 (d, J = 13.0 Hz, 1H), 1.22-1.00 (m, 2H), 0.94 (d, J = 6.2 Hz, 3H).
203	 (S)-1-(5-((4-isobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)methylpyrimidine-2,4(1H,3H)-dione	399.4	(500 MHz, Methanol-d ₄) δ 8.46 (dd, J = 7.1, 0.9 Hz, 1H), 8.02 (s, 1H), 7.51 (dd, J = 1.9, 1.0 Hz, 1H), 7.01 (dd, J = 7.2, 1.8 Hz, 1H), 3.91 (t, J = 6.7 Hz, 2H), 3.63-3.49 (m, 2H), 2.91 (t, J = 6.8 Hz, 3H), 2.73 (dd, J = 28.7, 10.7 Hz, 2H), 2.39 (dd, J = 71.2, 21.3 Hz, 4H), 2.02 (d, J = 45.3 Hz, 2H), 1.81 (s, 1H), 1.04 (d, J = 6.2 Hz, 3H), 0.93 (dd, J = 7.7, 6.6 Hz, 6H).
204	 (S)-1-(5-((4-isopentyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)methylpyrimidine-2,4(1H,3H)-dione	413.4	(400 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 8.62 (d, J = 7.2 Hz, 1H), 8.04 (s, 1H), 7.49 (s, 1H), 6.89 (dd, J = 7.1, 1.8 Hz, 1H), 3.78 (t, J = 6.7 Hz, 2H), 3.62 (s, 3H), 3.29 (s, 2H), 3.00 (s, 4H), 2.79 (t, J = 6.7 Hz, 2H), 2.44-2.18 (m, 2H), 1.72-1.40 (m, 3H), 1.29 (dd, J = 29.4, 6.4 Hz, 3H), 0.92 (t, J = 6.1 Hz, 6H).
205	 (S)-1-(5-((4-((3,3-difluorocyclobutyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)methylpyrimidine-2,4(1H,3H)-dione	447.3	(500 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 8.57 (dd, J = 7.2, 0.9 Hz, 1H), 8.01 (s, 1H), 7.51-7.23 (m, 1H), 6.88 (dd, J = 7.2, 1.8 Hz, 1H), 3.77 (t, J = 6.7 Hz, 2H), 3.49-3.42 (m, 2H), 2.83-2.70 (m, 4H), 2.60 (q, J = 14.0, 11.8 Hz, 4H), 2.37 (s, 1H), 2.32-2.12 (m, 6H), 1.91 (s, 1H), 0.95 (d, J = 6.2 Hz, 3H).

-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
206	 (S)-1-(5-((3-methyl-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	455.4	(400 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 8.74-8.50 (m, 1H), 8.04 (s, 1H), 7.49 (s, 1H), 6.89(dd, J = 7.2, 1.8 Hz, 1H), 3.84 (dd, J = 11.0, 4.1 Hz, 2H), 3.78 (t, J = 6.7 Hz, 2H), 3.64 (s, 3H), 3.50 (d, J = 11.4 Hz, 1H), 3.41-3.19 (m, 4H), 3.00 (d, J = 14.8 Hz, 4H), 2.79 (t, J = 6.7 Hz, 2H), 2.26 (d, J = 11.8 Hz, 1H), 1.59 (d, J = 13.9 Hz, 5H), 1.22 (dd, J = 22.3, 9.4 Hz, 5H).
207	 (S)-1-(5-((4-((4,4-difluorocyclohexyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	475.2	(500 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 8.57 (d, J = 7.1 Hz, 1H), 8.01 (s, 1H), 7.44 (s, 1H), 6.88(dd, J = 7.2, 1.8 Hz, 1H), 3.77 (t, J = 6.7 Hz, 2H), 3.50-3.40 (m, 2H), 2.78 (t, J = 6.7 Hz, 2H), 2.68-2.53 (m, 2H), 2.50-2.28 (m, 4H), 2.17 (dt, J = 19.9, 10.0 Hz, 2H), 2.05-1.64 (m, 8H), 1.09 (dt, J = 45.2, 11.8 Hz, 2H), 0.94 (d, J = 6.1 Hz, 3H).
208	 (S)-1-(5-((3-methyl-4-(pyridin-3-ylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	434.4	(400 MHz, DMSO-d ₆) δ 10.41 (br s, 1H), 8.56 (d, J = 7.2 Hz, 1H), 8.48 (d, J = 1.6 Hz, 1H), 8.46-8.43 (m, 1H), 8.00 (s, 1H), 7.69 (d, J = 8.0 Hz, 1H), 7.43 (s, 1H), 7.37-7.31 (m, 1H), 6.90-6.84 (m, 1H), 3.95 (d, J = 14.0 Hz, 1H), 3.78-3.74 (m, 2H), 3.52-3.42 (m, 3H), 3.23 (d, J = 14.0 Hz, 1H), 2.79-2.75 (m, 2H), 2.64-2.55 (m, 3H), 2.17 (d, J = 8.0 Hz, 2H), 2.04-1.93 (m, 1H), 1.07 (d, J = 6.0 Hz, 3H).
209	 1-(5-(((3S)-3-methyl-4-((tetrahydrofuran-2-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	427.0	(400 MHz, Methanol-d ₄) δ 8.62 (d, J = 7.2 Hz, 1H), 8.15 (s, 1H), 7.89 (s, 1H), 7.05 (d, J = 7.3 Hz, 1H), 4.56 (s, 2H), 4.27 (q, J = 8.1 Hz, 1H), 4.14 (s, 1H), 4.05-3.74 (m, 7H), 3.74-3.50 (m, 3H), 3.50-3.20 (m, 3H), 2.91 (t, J = 6.8 Hz, 2H), 2.20 (dq, J = 12.8, 6.4 Hz, 1H), 2.09-1.87 (m, 2H), 1.77-1.57 (m, 1H), 1.48 (d, J = 6.6 Hz, 3H). NH proton not observed due to solvent exchange.

Example 203 (alternate synthesis). Preparation of (S)-1-(5-((4-isobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione (Example 203)




Step 1. (S)-3-(2,4-dimethoxybenzyl)-1-(5-((4-isobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione

[0951] To a solution of (S)-3-(2,4-dimethoxybenzyl)-1-(5-((3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione trifluoroacetate (2.2 g, 3.6 mmol) in DCM (20 mL) was added Et₃N (0.51 mL, 3.6 mmol) followed by isobutyraldehyde (1.31 g, 18.1 mmol). The mixture was stirred at rt for 30 min and then NaBH(OAc)₃ (3.84 g, 18.1 mmol) was added. The reaction was stirred overnight at rt. The reaction was quenched with a solution of saturated aqueous NaHCO₃ and extracted with DCM. The organic layer was dried over Na₂SO₄, filtered and concentrated. Silica gel column chromatography (eluting with 10-100% EtOAc (containing 25% EtOH) in hexane) provided (S)-3-(2,4-dimethoxybenzyl)-1-(5-((4-isobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione (1.25 g, 2.28 mmol, 63% yield) as a white solid. LCMS [M+H]⁺: 549.3.

Step 2. (S)-1-(5-((4-isobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione (Example 203)

[0952] TFA (10 mL) was added to (S)-3-(2,4-dimethoxybenzyl)-1-(5-((4-isobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione (1.25 g, 2.28 mmol). The reaction mixture was stirred for 16 h at 90° C. and then concentrated. The crude material was dissolved in DCM and washed with a solution of saturated aqueous NaHCO₃. The layers were separated and the aqueous layer was extracted with DCM. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered and concentrated. Silica gel column chromatography (eluting with 10-100% EtOAc (containing 25% EtOH) in hexane) provided (S)-1-(5-((4-isobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione (800 mg, 1.97 mmol, 86% yield). LCMS [M+H]⁺: 399.4. ¹H NMR (500 MHz, Methanol-d₄) δ 8.46 (dd, J=7.1, 0.9 Hz, 1H), 8.02 (s, 1H), 7.51 (dd, J=1.9, 1.0 Hz, 1H), 7.01 (dd, J=7.2, 1.8 Hz, 1H), 3.91 (t, J=6.7 Hz, 2H), 3.63-3.49 (m, 2H), 2.91 (t, J=6.8 Hz, 3H), 2.73 (dd, J=28.7, 10.7 Hz, 2H), 2.39 (dd, J=71.2, 21.3 Hz, 4H), 2.02 (d, J=45.3 Hz, 2H), 1.81 (s, 1H), 1.04 (d, J=6.2 Hz, 3H), 0.93 (dd, J=7.7, 6.6 Hz, 6H).

Examples 210 and 211. Preparation of 1-(5-(((S)-4-(((1*r*,4*S*)-4-methoxycyclohexyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione (Example 210) and 1-(5-(((S)-4-(((1*s*,4*R*)-4-methoxycyclohexyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione (Example 211)



[0953] Prepared using the method of Example 202, wherein a commercially available mixture of cis and trans 1-(bromomethyl)-4-methoxycyclohexane was used in place of trans-4-(benzyloxy)cyclohexane-1-carbaldehyde. The stereoisomers were purified after the final step by reverse-phase HPLC (eluting with using ACN/Water/0.1% TFA).


Example 210. 1-(5-(((S)-4-(((1R,4S)-4-methoxycyclohexyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

[0954] Eluted first, minor isomer. LCMS $[M+H]^+$: 469.3. 1H NMR (400 MHz, DMSO- d_6) δ 10.45 (s, 1H), 8.64 (d, $J=7.2$ Hz, 1H), 8.05 (s, 1H), 7.52 (s, 1H), 6.90 (dd, $J=7.2$, 1.8 Hz, 1H), 4.99 (s, 4H), 3.79 (t, $J=6.7$ Hz, 2H), 3.72 (s, 2H), 3.23 (s, 4H), 3.13-2.94 (m, 4H), 2.79 (t, $J=6.7$ Hz, 3H), 2.37 (d, $J=30.0$ Hz, 1H), 2.00 (s, 2H), 1.92-1.57 (m, 2H), 1.26 (s, 3H), 1.07 (dt, $J=37.0$, 13.5 Hz, 4H).

Example 211. 1-(5-(((S)-4-(((1S,4R)-4-methoxycyclohexyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

[0955] Eluted second, major isomer. LCMS $[M+H]^+$: 469.3. 1H NMR 400 MHz, DMSO- d_6) δ 10.44 (s, 1H), 8.63 (d, $J=7.2$ Hz, 1H), 8.04 (s, 1H), 7.50 (s, 1H), 6.90 (dd, $J=7.2$, 1.8 Hz, 1H), 3.78 (t, $J=6.7$ Hz, 2H), 3.67 (s, 5H), 3.23 (s, 3H), 3.20 (s, 3H), 3.17-2.84 (m, 4H), 2.79 (t, $J=6.7$ Hz, 2H), 2.46-2.21 (m, 1H), 1.92-1.69 (m, 3H), 1.55 (s, 1H), 1.51-1.29 (m, 4H), 1.25 (s, 3H).


Example 212. Preparation of (S)-1-(5-((4-(cyclopentylmethyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[0956] Prepared using the method of Example 186, wherein tert-butyl (S)-4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperazine-1-carboxylate was used in place of tert-butyl 4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperazine-1-carboxylate, and cyclopentanecarbaldehyde was used in place of isobutyraldehyde. LCMS

$[M+H]^+$: 425.3. 1H NMR (400 MHz, DMSO- d_6) δ 10.44 (s, 1H), 8.63 (d, $J=7.2$ Hz, 1H), 8.04 (s, 1H), 7.51 (s, 1H), 6.90 (dd, $J=7.2$, 1.9 Hz, 1H), 4.39 (s, 3H), 3.78 (t, $J=6.7$ Hz, 2H), 3.61 (d, $J=44.7$ Hz, 3H), 3.30 (d, $J=20.9$ Hz, 2H), 3.00 (d, $J=14.0$ Hz, 3H), 2.79 (t, $J=6.7$ Hz, 2H), 2.18 (s, 1H), 1.79 (d, $J=20.7$ Hz, 2H), 1.58 (ddd, $J=38.4$, 7.7, 4.0 Hz, 4H), 1.42-1.10 (m, 5H).

Example 213. Preparation of (S)-1-(5-((4-(cyclobutylmethyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione






[0957] Prepared using the method of Example 186, wherein tert-butyl (S)-4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperazine-1-carboxylate was used in place of tert-butyl 4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperazine-1-carboxylate, and cyclobutanecarbaldehyde was used in place of isobutyraldehyde. LCMS $[M+H]^+$: 411.5. 1H NMR (500 MHz, DMSO- d_6) δ 10.44 (s, 1H), 8.58 (d, $J=7.1$ Hz, 1H), 8.02 (s, 1H), 7.45 (s, 1H), 6.89 (dd, $J=7.1$, 1.8 Hz, 1H), 3.78 (t, $J=6.7$ Hz, 2H), 3.47 (s, 2H), 2.79 (t, $J=6.7$ Hz, 2H), 2.76-2.57 (m, 3H), 2.56 (s, 2H), 2.14 (d, $J=47.7$ Hz, 3H), 1.99 (s, 3H), 1.92-1.73 (m, 3H), 1.65 (s, 2H), 1.03 (d, $J=50.1$ Hz, 3H).

[0958] The compounds in the following table were prepared using the method of Example 192, wherein potassium (R)-((4-(tert-butoxycarbonyl)-3-methylpiperazin-1-yl)methyl)trifluoroborate was used in place of tert-butyl (S)-2-methyl-4-((trifluoro- I_4 -boraneyl)methyl)piperazine-1-carboxylate, potassium salt and the appropriate commercially available aldehyde was used in place of cyclohexanecarbaldehyde.




Example No.	Structure	Mass $[M+H]^+$	1H NMR
214	(R)-1-(5-((4-(cyclohexylmethyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	439.1	(400 MHz, DMSO- d_6) δ 10.43 (s, 1H), 8.60-8.54 (m, 1H), 8.00 (s, 1H), 7.44 (s, 1H), 6.89-6.86 (m, 1H), 3.79-3.75 (m, 2H), 3.51-3.39 (m, 2H), 2.81-2.71 (m, 3H), 2.58-2.52 (m, 2H), 2.45-2.39 (m, 1H), 2.36-2.26 (m, 1H), 2.24-2.06 (m, 2H), 2.00-1.75 (m, 3H), 1.63 (br s, 4H), 1.47-1.35 (m, 1H), 1.29-1.07 (m, 3H), 0.92 (d, $J=6.0$ Hz, 3H), 0.89-0.69 (m, 2H).

-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
215	 (R)-1-(5-((3-methyl-4-(pyridin-3-ylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	434.1	(400 MHz, CDCl ₃) δ 8.55 (d, J = 1.8 Hz, 1H), 8.49-8.47(m, 1H), 8.35 (d, J = 7.2 Hz, 1H), 8.09-7.88 (m, 2H), 7.67-7.64 (m, 1H), 7.29 (s, 1H), 7.27-7.25 (m, 1H), 6.89-6.87(m, 1H), 4.02-3.99 (m, 1H), 3.88 (t, J = 6.8 Hz, 2H), 3.48 (s, 2H), 3.24-3.21(m, 1H), 2.90 (t, J = 6.8 Hz, 2H), 2.71-2.50 (m, 4H), 2.25-2.23 (m, 2H), 2.07-2.06 (m, 1H), 1.14 (d, J = 6.4 Hz, 3H).
216	 (R)-1-(5-((4-isobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	399.0	(400 MHz, Methanol-d ₄) δ 8.53 (s, 1H), 8.45 (d, J = 7.3 Hz, 1H), 8.02 (s, 1H), 7.50 (s, 1H), 6.99 (dd, J = 7.2, 2.0 Hz, 1H), 4.73-4.50 (m, 1H), 3.89 (t, J = 6.8 Hz, 2H), 3.63 (d, J = 13.5 Hz, 1H), 3.58 (d, J = 13.5 Hz, 1H), 3.19 (d, J = 12.0 Hz, 1H), 2.89 (t, J = 6.8 Hz, 2H), 2.87-2.71 (m, 4H), 2.65 (s, 1H), 2.53-2.19 (m, 3H), 1.93 (hept, J = 6.1 Hz, 2H), 1.18 (d, J = 6.2 Hz, 3H), 0.98 (d, J = 6.0 Hz, 3H), 0.96 (s, 3H). NH proton not observed due to solvent exchange
217	 (R)-1-(5-((3-methyl-4-((tetrahydro-2H-pyran-4-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	441.3	(400 MHz, Methanol-d ₄) δ 8.53 (d, J = 7.2 Hz, 1H), 8.46 (d, J = 7.2 Hz, 1H), 8.03 (s, 1H), 7.51 (s, 1H), 7.00 (dd, J = 7.2, 1.8 Hz, 1H), 3.98-3.87 (m, 4H), 3.63 (d, J = 13.6 Hz, 1H), 3.58 (d, J = 13.6 Hz, 1H), 3.50-3.37 (m, 2H), 3.16 (dt, J = 12.2, 3.4 Hz, 1H), 2.90 (t, J = 6.8 Hz, 2H), 2.87-2.76 (m, 4H), 2.62 (t, J = 10.9 Hz, 1H), 2.45 (t, J = 10.7 Hz, 1H), 2.36 (dd, J = 11.2, 4.4 Hz, 1H), 2.29-2.18 (m, 1H), 1.97-1.84 (m, 1H), 1.78 (ddd, J = 13.3, 4.0, 2.1 Hz, 1H), 1.66-1.58 (m, 1H), 1.38-1.21 (m, 2H), 1.17 (d, J = 6.2 Hz, 3H). NH proton not observed due to solvent exchange





[0959] The compounds in the following table were prepared using the method of Example 192, wherein potassium (R)-((4-(tert-butoxycarbonyl)-2-methylpiperazin-1-yl)methyl)trifluoroborate [see *J. Med. Chem.* 2012, 55, 7796-

7816] was used in place of tert-butyl (S)-2-methyl-4-((trifluoro-14-boraneyl)methyl)piperazine-1-carboxylate, potassium salt, and the appropriate commercially available aldehyde was used in place of cyclohexanecarbaldehyde.


Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
218	 (R)-1-(5-((4-(cyclohexylmethyl)-2-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	439.1	(400 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 8.40 (d, J = 7.2 Hz, 1H), 7.99 (s, 1H), 7.43 (s, 1H), 6.88-6.84 (m, 1H), 3.96-3.91 (m, 1H), 3.76 (t, J = 6.8 Hz, 2H), 3.19-3.15 (m, 1H), 2.77 (t, J = 6.8 Hz, 2H), 2.62-2.51 (m, 3H), 2.44 (br s, 1H), 2.19-2.09 (m, 1H), 2.07-1.95 (m, 3H), 1.88-1.86 (m, 1H), 1.76-1.56 (m, 5H), 1.50-1.37 (m, 1H), 1.25-1.03 (m, 6H), 0.84-0.75 (m, 2H).
219	 (R)-1-(5-((2-methyl-4-((tetrahydro-2H-pyran-4-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	441.2	(400 MHz, Methanol-d ₄) δ 8.63 (d, J = 7.2 Hz, 1H), 8.14 (d, J = 2.2 Hz, 1H), 7.81 (d, J = 5.4 Hz, 1H), 7.05-6.94 (m, 1H), 4.89-4.88 (m, 2H), 4.29-4.16 (m, 1H), 4.00-3.85 (m, 5H), 3.84-3.66 (m, 2H), 3.66-3.55 (m, 1H), 3.50-3.41 (m, 2H), 3.26-3.05 (m, 4H), 2.90 (t, J = 6.8 Hz, 2H), 2.24-2.08 (m, 1H), 1.78-1.47 (m, 5H), 1.45-1.30 (m, 3H). NH protons not observed due to solvent exchange.
220	 (R)-1-(5-((4-isobutyl-2-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	399.3	(400 MHz, CD ₃ OD) δ 8.46 (s, 1H), 8.45 (d, J = 6.8 Hz, 1H), 8.01 (s, 1H), 7.50 (s, 1H), 6.99-6.96 (m, 1H), 4.18 (d, J = 14.0 Hz, 2H), 3.88 (d, J = 6.8 Hz, 2H), 3.12-3.08 (m, 2H), 2.88 (t, J = 6.4 Hz, 2H), 2.72-2.37 (m, 7H), 1.99-1.95 (m, 1H), 1.23 (d, J = 6.0 Hz, 3H), 0.97 (d, J = 6.4 Hz, 6H).

[0960] The compounds in the following table were prepared using the method of Example 192, wherein potassium (S)-((4-(tert-butoxycarbonyl)-2-methylpiperazin-1-yl)methyl)trifluoroborate [see *J. Med. Chem.* 2012, 55, 7796-


7816] was used in place of tert-butyl (S)-2-methyl-4-((trifluoro-14-boraneyl)methyl)piperazine-1-carboxylate, potassium salt and the appropriate commercially available aldehyde was used in place of cyclohexanecarbaldehyde.

Example No.	Structure	Mass [M + H] ⁺ ¹ H NMR
221	 (S)-1-(5-((4-(cyclohexylmethyl)-2-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	439.1 (400 MHz, DMSO-d ₆) δ = 10.43 (br s, 1H), 8.55 (d, J = 7.2 Hz, 1H), 7.99 (s, 1H), 7.43 (s, 1H), 6.93-6.81 (m, 1H), 3.94 (d, J = 13.8 Hz, 1H), 3.78-3.74 (m, 2H), 3.17 (d, J = 13.8 Hz, 1H), 2.78-2.75 (m, 2H), 2.63-2.52 (m, 3H), 2.47-2.41 (m, 1H), 2.19-1.96 (m, 4H), 1.87 (d, J = 8.6 Hz, 1H), 1.76-1.57 (m, 5H), 1.49-1.37 (m, 1H), 1.24-1.05 (m, 6H), 0.90-0.69 (m, 2H).
222	 (S)-1-(5-((4-isobutyl-2-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	399.0 (400 MHz, DMSO-D ₆) δ 10.44 (s, 1H), 8.60-8.57 (m, 1H), 8.31 (s, 1H), 8.13 (s, 1H), 8.01 (d, J = 2.4 Hz, 1H), 7.46 (s, 1H), 6.87-6.88 (m, 1H), 4.00 (m, 1H), 3.76 (t, J = 6.4 Hz, 2H), 3.55-3.50 (m, 1H), 2.92-2.08 (m, 6H), 2.32-2.10 (m, 3H), 1.85 (s, 1H), 1.13 (d, J = 6.0 Hz, 3H), 0.87 (d, J = 6.4 Hz, 6H).
223	 (S)-1-(5-((2-methyl-4-((tetrahydro-2H-pyran-4-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	440.8 (400 MHz, Methanol-d ₄) δ 8.48 (d, J = 7.4 Hz, 1H), 8.04 (s, 1H), 7.54 (s, 1H), 6.98 (d, J = 7.1 Hz, 1H), 4.30 (s, 1H), 4.00-3.85 (m, 4H), 3.45 (t, J = 12.0 Hz, 4H), 3.16-2.77 (m, 8H), 2.55 (s, 1H), 2.11 (s, 1H), 1.69 (d, J = 13.2 Hz, 3H), 1.34 (d, J = 17.8 Hz, 5H). NH protons not observed due to solvent exchange.
224	 (S)-1-(5-((4-(cyclobutylmethyl)-2-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	411.1 (400 MHz, Methanol-d ₄) δ 8.46 (d, J = 7.2 Hz, 1H), 8.37 (s, 1H), 8.02 (s, 1H), 7.50 (s, 1H), 6.97 (dd, J = 7.2, 1.9 Hz, 1H), 4.20 (d, J = 14.3 Hz, 1H), 3.89 (t, J = 6.8 Hz, 2H), 3.32-3.20 (m, 1H), 3.08 (s, 2H), 2.97-2.86 (m, 4H), 2.75 (d, J = 8.0 Hz, 3H), 2.41 (s, 1H), 2.18 (d, J = 7.8 Hz, 2H), 2.02 (d, J = 7.5 Hz, 1H), 1.87 (s, 3H), 1.35-1.28 (m, 2H), 1.26 (d, J = 4.9 Hz, 4H). NH protons not observed due to solvent exchange

-continued


Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
225	 1-(5-(((2S)-2-methyl-4-((tetrahydrofuran-2-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	427.1	(400 MHz, DMSO-d ₆) δ 10.44 (s, 1H), 8.58 (d, J = 7.3 Hz, 1H), 8.13 (s, 1H), 8.01 (s, 1H), 7.46 (s, 1H), 6.86 (d, J = 7.2 Hz, 1H), 4.06-3.94 (m, 2H), 3.82-3.71 (m, 3H), 3.63 (q, J = 8.6, 7.9 Hz, 1H), 3.10-2.85 (m, 3H), 2.77 (t, J = 6.7 Hz, 2H), 2.72-2.54 (m, 5H), 2.37-2.19 (m, 2H), 1.94 (qd, J = 12.4, 6.7 Hz, 1H), 1.86-1.72 (m, 2H), 1.46 (dq, J = 12.0, 7.8 Hz, 1H), 1.11 (d, J = 6.0 Hz, 3H). NH protons not observed due to solvent exchange.

Example 226. Preparation of (S)-1-(5-((4-((3,3-difluorocyclobutyl)methyl)-2-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0961] Prepared using the method of Example 156 wherein potassium (S)-((4-(tert-butoxycarbonyl)-2-methylpiperazin-1-yl)methyl)trifluoroborate [see J. Med. Chem. 2012, 55, 7796-7816] was used in place of potassium ((4-(tert-butoxycarbonyl)piperazin-1-yl)methyl)trifluoroborate and (3,3-difluorocyclobutyl)methyl 4-methylbenzenesulfonate was used in place of (bromomethyl)cyclohexane. LCMS [M+H]⁺: 447.0. ¹H NMR (400 MHz, Methanol-d₄) δ 8.45 (d, J=7.1 Hz, 1H), 8.37 (s, 1H), 8.01 (s, 1H), 7.51 (s, 1H), 6.98 (dd, J=7.1, 1.8 Hz, 1H), 4.19 (d, J=13.7 Hz, 1H), 3.89 (t, J=6.8 Hz, 2H), 3.34-3.27 (m, 1H), 3.07-2.94 (m, 2H), 2.89 (t, J=6.8 Hz, 2H), 2.86-2.63 (m, 6H), 2.53 (t, J=11.2 Hz, 1H), 2.49-2.24 (m, 5H), 1.23 (d, J=6.2 Hz, 3H), NH proton not observed due to solvent exchange.

Example 227. Preparation of (S)-1-(5-((3-ethyl-4-((tetrahydro-2H-pyran-4-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0962] Prepared using the method of Example 192, wherein potassium (S)-((4-(tert-butoxycarbonyl)-3-ethylpiperazin-1-yl)methyl)trifluoroborate was used in place of tert-butyl (S)-2-methyl-4-((trifluoro-1,4-boraneyl)methyl)piperazine-1-carboxylate, potassium salt and tetrahydro-2H-pyran-4-carbaldehyde was used in place of cyclohexanecarbaldehyde. LCMS [M+H]⁺: 455.5. ¹H NMR (400 MHz, Methanol-d₄) δ 8.48 (d, J=7.2 Hz, 1H), 8.23 (s, 1H), 8.04 (s, 1H), 7.52 (s, 1H), 7.00 (dd, J=7.3, 1.9 Hz, 1H), 3.96 (dd, J=11.5, 4.2 Hz, 2H), 3.90 (t, J=6.8 Hz, 2H), 3.86-3.50 (m, 4H), 3.50-3.39 (m, 2H), 3.28-2.51 (m, 9H), 1.96 (d, J=78.4 Hz, 3H), 1.75 (d, J=13.3 Hz, 2H), 1.66 (d, J=13.2 Hz, 1H), 1.49-1.31 (m, 2H), 0.96 (t, J=7.4 Hz, 3H), NH proton not observed due to solvent exchange.

Example 228. Preparation of (S)-1-(5-((3-ethyl-4-isobutylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0963] Prepared using the method of Example 192, wherein (S)-((4-(tert-butoxycarbonyl)-3-ethylpiperazin-1-yl)methyl)trifluoroborate was used in place of tert-butyl (S)-2-methyl-4-((trifluoro-1,4-boraneyl)methyl)piperazine-1-carboxylate, potassium salt and isobutyraldehyde was used in place of cyclohexanecarbaldehyde. LCMS [M+H]⁺: 413.3. ¹H NMR (400 MHz, Methanol-d₄) δ 8.48 (d, J=7.1 Hz, 1H), 8.37 (s, 1H), 8.04 (s, 1H), 7.51 (s, 1H), 7.00 (d, J=7.2 Hz, 1H), 3.90 (t, J=6.6 Hz, 2H), 3.74 (d, J=13.7 Hz, 1H), 3.63 (d, J=13.7 Hz, 1H), 3.48 (s, 1H), 3.23-3.01 (m, 3H), 2.89 (t, J=7.0 Hz, 5H), 2.75-2.50 (m, 2H), 2.12-1.99 (m, 1H), 1.95-1.69 (m, 2H), 1.05 (t, J=6.1 Hz, 6H), 0.96 (t, J=7.4 Hz, 3H), NH proton not observed due to solvent exchange.

Example 229. Preparation of (S)-1-(5-((4-isobutyl-3-isopropylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[0964] Prepared using the method of Example 192, wherein (S)-((4-(tert-butoxycarbonyl)-3-isopropylpiperazin-1-yl)methyl)trifluoroborate was used in place of tert-butyl (S)-2-methyl-4-((trifluoro-14-boranyl)methyl)piperazine-1-carboxylate, potassium salt and isobutyraldehyde was used in place of cyclohexanecarbaldehyde. LCMS $[M+H]^+$: 427.5. 1H NMR (400 MHz, Methanol- d_4) δ 8.48 (d, $J=7.4$ Hz, 1H), 8.07 (s, 1H), 8.04 (s, 1H), 7.53 (s, 1H), 7.00 (dd, $J=6.9, 1.8$ Hz, 1H), 3.90 (t, $J=6.8$ Hz, 2H), 3.86-3.53 (m, 3H), 3.41-2.95 (m, 4H), 2.89 (t, $J=6.8$ Hz, 2H), 2.55 (d, $J=44.4$ Hz, 4H), 2.12 (s, 1H), 1.11-1.01 (m, 9H), 1.00 (s, 3H), NH proton not observed due to solvent exchange.


Example 230. (Preparation of (S)-1-(5-((3-isopropyl-4-((tetrahydro-2H-pyran-4-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[0965] Prepared using the method of Example 192, wherein (S)-((4-(tert-butoxycarbonyl)-3-isopropylpiperazin-1-yl)methyl)trifluoroborate was used in place of tert-butyl (S)-2-methyl-4-((trifluoro-14-boranyl)methyl)piperazine-1-carboxylate, potassium salt and tetrahydro-2H-pyran-4-carbaldehyde was used in place of cyclohexanecarbaldehyde. LCMS $[M+H]^+$: 469.2. 1H NMR (400 MHz, Methanol- d_4) δ 8.49 (d, $J=7.2$ Hz, 1H), 8.30 (s, 1H), 8.04 (s, 1H), 7.55 (s, 1H), 7.00 (dd, $J=7.2, 1.8$ Hz, 1H), 3.95 (dd, $J=11.5, 4.2$ Hz, 2H), 3.90 (t, $J=6.8$ Hz, 2H), 3.85 (d, $J=13.7$ Hz, 1H), 3.72 (d, $J=13.6$ Hz, 1H), 3.53-3.38 (m, 3H), 3.18-2.92 (m, 5H), 2.89 (t, $J=6.8$ Hz, 2H), 2.59 (t,


$J=10.8$ Hz, 2H), 2.50 (dd, $J=12.6, 10.0$ Hz, 1H), 2.41-2.28 (m, 1H), 2.07-1.94 (m, 1H), 1.79 (d, $J=13.4$ Hz, 1H), 1.65 (d, $J=13.2$ Hz, 1H), 1.44-1.25 (m, 2H), 1.01 (d, $J=6.8$ Hz, 3H), 0.95 (d, $J=6.8$ Hz, 3H), NH proton not observed due to solvent exchange.

Example 231. Preparation of 1-(5-((4-(cyclohexylmethyl)-3,3-dimethylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0966] Prepared using the method of Example 192, wherein ((4-(tert-butoxycarbonyl)-3,3-dimethylpiperazin-1-yl)methyl)trifluoroborate was used in place of tert-butyl (S)-2-methyl-4-((trifluoro-14-boranyl)methyl)piperazine-1-carboxylate, potassium salt. LCMS $[M+H]^+$: 453.3. 1H NMR (400 MHz, Methanol- d_4) δ 8.52 (s, 1H), 8.46 (d, $J=7.2$ Hz, 1H), 8.02 (s, 1H), 7.49 (s, 1H), 7.00 (dd, $J=7.4, 1.7$ Hz, 1H), 3.89 (t, $J=6.8$ Hz, 2H), 3.61 (s, 2H), 2.89 (t, $J=6.8$ Hz, 2H), 2.99-2.32 (m, 5H), 1.92-1.59 (m, 6H), 1.42-1.15 (m, 10H), 1.04 (q, $J=11.4$ Hz, 2H), NH proton not observed due to solvent exchange.

Example 232. Preparation of 1-(5-((3,3-dimethyl-4-((tetrahydro-2H-pyran-4-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0967] Prepared using the method of Example 192, wherein ((4-(tert-butoxycarbonyl)-3,3-dimethylpiperazin-1-yl)methyl)trifluoroborate was used in place of tert-butyl (S)-2-methyl-4-((trifluoro-14-boranyl)methyl)piperazine-1-carboxylate, potassium salt and tetrahydro-2H-pyran-4-carbaldehyde was used in place of cyclohexanecarbaldehyde. LCMS $[M+H]^+$: 455.1. 1H NMR (300 MHz, Methanol- d_4) δ 8.45 (d, $J=7.2$ Hz, 1H), 8.27 (s, 1H), 8.01 (s, 1H), 7.47 (s, 1H), 6.98 (d, $J=7.3$ Hz, 1H), 3.94 (d, $J=12.8$ Hz, 1H), 3.88 (t, $J=6.8$ Hz, 2H), 3.71-3.31 (m, 9H), 3.05 (s, 5H), 2.87 (t, $J=6.7$ Hz, 2H), 2.08-1.58 (m, 3H), 1.32 (d, $J=24.2$ Hz, 7H). NH proton not observed due to solvent exchange.

Example 233. Preparation of 1-(5-((4-isobutyl-3,3-dimethylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0968] Prepared using the method of Example 192, wherein ((4-(tert-butoxycarbonyl)-3,3-dimethylpiperazin-1-yl)methyl)trifluoroborate was used in place of tert-butyl (S)-2-methyl-4-((trifluoro-14-boraneyl)methyl)piperazine-1-carboxylate, potassium salt and isobutyraldehyde was used in place of cyclohexanecarbaldehyde. LCMS [M+H]⁺: 413.3. ¹H NMR (300 MHz, Methanol-d₄) δ 8.47 (d, J=7.1 Hz, 1H), 8.03 (s, 1H), 7.50 (s, 1H), 7.00 (d, J=7.1 Hz, 1H), 3.90 (t, J=6.7 Hz, 2H), 3.81-3.36 (m, 2H), 3.11 (d, J=39.0 Hz, 2H), 2.89 (t, J=6.8 Hz, 2H), 2.83-2.42 (m, 3H), 2.01 (ddd, J=25.6, 12.6, 5.4 Hz, 2H), 1.59-1.20 (m, 8H), 1.08 (d, J=6.5 Hz, 6H).

Example 234. Preparation of 1-(5-(((1R,4R)-5-(cyclohexylmethyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0969] Prepared using the method of Example 192, wherein potassium (((1R,4R)-5-(tert-butoxycarbonyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)trifluoroborate was used in place of tert-butyl (S)-2-methyl-4-((trifluoro-14-boraneyl)methyl)piperazine-1-carboxylate. LCMS [M+H]⁺: 437.2.

Example 235. Preparation of 1-(5-(((1R,4R)-5-(pyridin-3-ylmethyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0970] Prepared using the method of Example 192, wherein potassium (((1R,4R)-5-(tert-butoxycarbonyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)trifluoroborate was used in place of tert-butyl (S)-2-methyl-4-((trifluoro-14-boraneyl)methyl)piperazine-1-carboxylate and using nicotinaldehyde was used in place of cyclohexanecarbaldehyde. LCMS [M+H]⁺: 432.2. ¹H NMR (400 MHz, CD₃OD) δ 8.80 (brs, 1H), 8.60 (d, J=7.2 Hz, 1H), 8.42 (m, 1H), 8.13 (s, 1H), 7.90 (brs, 1H), 7.77 (s, 1H), 7.07 (d, J=7.2 Hz, 1H), 4.49-4.44 (m, 1H), 4.32-4.20 (m, 3H), 4.08-3.87 (m, 5H), 3.62-3.61 (m, 3H), 3.04-3.01 (m, 1H), 2.90 (t, J=6.8 Hz, 2H), 2.31 (s, 2H), NH proton not observed due to solvent exchange.

Example 236. Preparation of 1-(5-(((1S,4S)-5-(cyclohexylmethyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[0971] Prepared using the method of Example 192, wherein potassium (((1S,4S)-5-(tert-butoxycarbonyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)trifluoroborate was used in place of tert-butyl (S)-2-methyl-4-((trifluoro-14-boraneyl)methyl)piperazine-1-carboxylate. LCMS [M+H]⁺: 437.2. ¹H NMR (400 MHz, Methanol-d₄) δ 8.46 (d, J=7.2 Hz, 1H), 8.03 (s, 1H), 7.52 (s, 1H), 7.02 (d, J=7.4 Hz, 1H), 4.23 (s, 1H), 3.97-3.86 (m, 2H), 3.83-3.54 (m, 2H), 3.28 (s, 1H), 3.19-2.95 (m, 3H), 2.95-2.84 (m, 2H), 2.84-2.61 (m, 1H), 2.23 (d, J=11.7 Hz, 1H), 2.13-2.04 (m, 1H), 1.91-1.69 (m, 6H), 1.59 (s, 1H), 1.42-1.20 (m, 4H), 1.15-0.95 (m, 2H), NH proton not observed due to solvent exchange.


Example 237. Preparation of 1-(5-(((1S,4S)-5-(pyridin-3-ylmethyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[0972] Prepared using the method of Example 192, wherein potassium (((1S,4S)-5-(tert-butoxycarbonyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)trifluoroborate was used in place of tert-butyl (S)-2-methyl-4-((trifluoro-14-boraneyl)methyl)piperazine-1-carboxylate and nicotinalde-

hyde was used in place of cyclohexanecarbaldehyde. LCMS $[M+H]^+$: 432.1. 1H NMR (400 MHz, Methanol- d_4) δ 8.83 (s, 1H), 8.75 (d, $J=5.6$ Hz, 1H), 8.59 (d, $J=7.3$ Hz, 1H), 8.52 (d, $J=8.1$ Hz, 1H), 8.12 (s, 1H), 7.96 (t, $J=6.9$ Hz, 1H), 7.78 (s, 1H), 7.07 (dd, $J=7.2$, 1.9 Hz, 1H), 4.49 (d, $J=13.3$ Hz, 1H), 4.33 (d, $J=13.2$ Hz, 1H), 4.28 (s, 1H), 4.20 (d, $J=14.7$ Hz, 1H), 4.05 (d, $J=14.7$ Hz, 1H), 3.93 (t, $J=6.8$ Hz, 2H), 3.84 (s, 1H), 3.62 (d, $J=11.8$ Hz, 1H), 3.38-3.19 (m, 2H), 3.01 (dd, $J=12.0$, 2.7 Hz, 1H), 2.89 (t, $J=6.8$ Hz, 2H), 2.31 (s, 2H), NH proton not observed due to solvent exchange.

Example 238. Preparation of 1-(5-(((1S,4S)-5-(isopropylsulfonyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



Example 238

Step 1. 1-(5-(((1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


[0973] To a stirred solution of 1-(5-(((1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione (175 mg, 0.32 mmol) [prepared in Example 236] in TFA (5 mL) was added TFOH (0.2 mL) and the reaction mixture was stirred for 2 h at 70° C. The reaction was

concentrated under to afford crude compound. The crude compound was dissolved in 10% MeOH in DCM and basified with Amberlyst-A21 (free base) resin and then filtered. The filtrate was concentrated to afford 1-(5-(((1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (150 mg, crude). LCMS $[M+H]^+$: 341.3.

Step 2. 1-(5-(((1S,4S)-5-(isopropylsulfonyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


[0974] To a stirred solution of 1-(5-(((1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (130 mg, 0.38 mmol) in DCM (5 mL) was added Et_3N (0.076 mL, 0.76 mmol) and propane-2-sulfonyl chloride (0.035 mL, 0.45 mmol) at -20° C. The reaction mixture was stirred for 2 h at 0° C. and then quenched with methanol and concentrated to afford the crude compound. The crude compound was purified by PREP HPLC using: Mobile Phase: A=0.1% TFA in water, B=Acetonitrile, Column: ATLANTIS (250 mm $\times$ 21.2 mm), 5.0 μ , Flow: 20 mL/min. The collected fraction were concentrated under reduced pressure to afford 1-(5-(((1S,4S)-5-(isopropylsulfonyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (12 mg, 0.022 mmol, 5.8% yield) as an off-white solid. LCMS $[M+H]^+$: 447.2. HPLC: R_t =5.406 min. 1H NMR (400 MHz, Methanol- d_4) δ 8.63 (dd, $J=7.2$, 3.1 Hz, 1H), 8.15 (d, $J=1.7$ Hz, 1H), 7.83 (d, $J=2.8$ Hz, 1H), 7.07 (dd, $J=7.4$, 2.1 Hz, 1H), 4.59 (dd, $J=13.5$, 7.5 Hz, 3H), 4.44 (d, $J=12.3$ Hz, 1H), 3.95 (t, $J=6.8$ Hz, 2H), 3.72 (s, 2H), 3.50 (d, $J=9.1$ Hz, 1H), 2.91 (t, $J=6.8$ Hz, 2H), 2.47 (d, $J=12.0$ Hz, 1H), 2.23 (d, $J=13.0$ Hz, 1H), 1.49 (d, $J=2.2$ Hz, 2H), 1.35 (d, $J=6.8$ Hz, 6H), NH proton not observed due to solvent exchange.

Example 239. Preparation of 1-(5-(((1R,4R)-5-(isopropylsulfonyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0975] Prepared from using the method of Example 238, wherein 1-(5-(((1R,4R)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione [prepared in Example 234] was used in place of 1-(5-(((1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione. LCMS $[M+H]^+$: 446.8.

Example 240. Preparation of 1-(5-((4-(3-methylbutan-2-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 240)




[0976] Prepared using the method of Example 156, steps 1 and 4, wherein potassium trifluoro((4-(3-methylbutan-2-yl)piperazin-1-yl)methyl)borate was used in place of potassium {[4-(tert-butoxycarbonyl)-1-piperazinyl]methyl}(trifluoro)borate. LCMS [M+H]⁺: 399.0. ¹H NMR (400 MHz, Methanol-d₄) δ 8.47 (d, J=7.1 Hz, 1H), 8.39 (s, 1H), 8.03 (s, 1H), 7.52 (s, 1H), 6.99 (d, J=7.2 Hz, 1H), 3.90 (t, J=6.7 Hz, 2H), 3.70 (s, 2H), 3.31 (m, 4H), 3.07 (d, J=6.5 Hz, 1H), 2.89 (t, J=6.8 Hz, 2H), 2.81 (bs, 4H), 2.30-2.16 (m, 1H), 1.25 (d, J=6.7 Hz, 3H), 1.04 (d, J=6.7 Hz, 3H), 0.97 (d, J=6.7 Hz, 3H).

Example 241. Preparation of 1-(5-((4-(3,3,3-trifluoro-2,2-dimethylpropyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0977] Prepared using the method of Example 156, steps 1 and 4, wherein potassium trifluoro((4-(3,3,3-trifluoro-2,2-dimethylpropyl)piperazin-1-yl)methyl)borate was used in place of potassium {[4-(tert-butoxycarbonyl)-1-piperazinyl]methyl}(trifluoro)borate. LCMS [M+H]⁺: 453.2. ¹H NMR (400 MHz, Methanol-d₄) δ 8.61 (d, J=7.0 Hz, 1H), 8.13 (s, 1H), 7.75 (s, 1H), 7.00 (dd, J=7.3, 1.9 Hz, 1H), 4.35 (s, 2H), 3.93 (t, J=6.8 Hz, 2H), 3.48 (d, J=12.2 Hz, 1H), 3.08 (t, J=11.8 Hz, 1H), 2.98-2.67 (m, 8H), 2.55 (d, J=10.8 Hz, 2H), 1.14 (d, J=2.1 Hz, 6H). NH proton not observed due to solvent exchange.

Example 242. Preparation of 1-(5-((4-(cyclohexylmethyl)-3-oxopiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0978] Prepared using the method of Example 156, steps 1 and 4, wherein potassium ((4-(cyclohexylmethyl)-3-oxopiperazin-1-yl)methyl)trifluoroborate was used in place of potassium {[4-(tert-butoxycarbonyl)-1-piperazinyl]methyl}(trifluoro)borate. LCMS [M+H]⁺: 439.2. ¹H NMR (400 MHz, Methanol-d₄) δ 8.50 (d, J=7.1 Hz, 1H), 8.04 (s, 1H), 7.57 (s, 1H), 7.00 (dd, J=7.1, 1.8 Hz, 1H), 3.90 (t, J=6.7 Hz, 2H), 3.81 (s, 2H), 3.43 (t, J=5.5 Hz, 2H), 3.38-3.31 (m, 2H), 3.25 (d, J=7.2 Hz, 2H), 2.89 (t, J=6.8 Hz, 4H), 1.80-1.59 (m, 6H), 1.24 (q, J=9.3, 6.0 Hz, 3H), 0.97 (q, J=11.9 Hz, 2H), NH proton not observed due to solvent exchange.

Example 243. Preparation of 1-(5-((4-(2-cyclohexylethyl)-3-oxopiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0979] Prepared using the method of Example 156, steps 1 and 4, wherein potassium ((4-(2-cyclohexylethyl)-3-oxopiperazin-1-yl)methyl)trifluoroborate was used in place of potassium {[4-(tert-butoxycarbonyl)-1-piperazinyl]methyl}(trifluoro)borate. LCMS [M+H]⁺: 452.9. ¹H NMR (400 MHz, Methanol-d₄) δ 8.52 (d, J=7.2 Hz, 1H), 8.06 (s, 1H), 7.62-7.58 (m, 1H), 7.00 (dd, J=7.1, 1.9 Hz, 1H), 4.01-3.78 (m, 3H), 3.51-3.38 (m, 5H), 3.12-2.96 (m, 2H), 2.89 (t, J=6.8 Hz, 2H), 1.83-1.60 (m, 5H), 1.46 (dt, J=9.7, 6.8 Hz, 2H), 1.37-1.15 (m, 5H), 0.96 (qd, J=14.1, 12.9, 4.4 Hz, 2H), NH proton not observed due to solvent exchange.

Example 244. Preparation of (R)-1-(5-((4-(cyclohexylmethyl)-3-(methoxymethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[0980] Prepared using the method of Example 156, steps 1 and 4, wherein potassium (R)-((4-(cyclohexylmethyl)-3-(methoxymethyl)piperazin-1-yl)methyl)trifluoroborate was used in place of potassium {[4-(tert-butoxycarbonyl)-1-piperazinyl]methyl}(trifluoro)borate. LCMS [M+H]⁺: 469. 4. ¹H NMR (400 MHz, Methanol-d₄) δ 8.46 (s, 1H), 8.45 (s, 1H), 8.02 (s, 1H), 7.49 (s, 1H), 3.89 (t, J=6.8 Hz, 2H), 3.70-3.53 (m, 4H), 3.36 (s, 4H), 3.19 (s, 1H), 3.03 (dd, J=12.0, 8.2 Hz, 1H), 2.97-2.82 (m, 5H), 2.67-2.47 (m, 3H), 1.90 (d, J=13.1 Hz, 1H), 1.82-1.63 (m, 5H), 1.41-1.18 (m, 4H), 1.09-0.92 (m, 2H), NH proton not observed due to solvent exchange.


Example 245. Preparation of (R)-1-(5-((4-isobutyl-3-(methoxymethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[0981] Prepared using the method of Example 156, steps 1 and 4, wherein potassium (R)-trifluoro((4-isobutyl-3-(methoxymethyl)piperazin-1-yl)methyl)borate was used in place of potassium {[4-(tert-butoxycarbonyl)-1-piperazinyl]methyl}(trifluoro)borate. LCMS [M+H]⁺: 429.3. ¹H NMR (400 MHz, Methanol-d₄) δ 8.46 (d, J=7.1 Hz, 1H), 8.42 (1H, s), 8.02 (s, 1H), 7.50 (s, 1H), 6.98 (d, J=7.1 Hz, 1H), 3.89 (t, J=6.7 Hz, 2H), 3.73-3.56 (m, 4H), 3.45 (d, J=13.1 Hz, 1H), 3.37 (s, 3H), 3.31 (s, 1H), 3.12-2.96 (m, 2H), 2.94-2.84 (m, 4H), 2.72 (dd, J=13.0, 5.5 Hz, 1H), 2.67-2.55 (m, 2H), 2.05


(h, J=6.8 Hz, 1H), 1.03 (d, J=6.8 Hz, 3H), 1.01 (d, J=6.8 Hz, 3H), NH proton not observed due to solvent exchange.

Example 246. Preparation of (R)-1-(5-((4-(cyclohexylmethyl)-3-(difluoromethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[0982] Prepared using the method of Example 156, steps 1 and 4, wherein potassium (R)-((4-(cyclohexylmethyl)-3-(difluoromethyl)piperazin-1-yl)methyl)trifluoroborate was used in place of potassium {[4-(tert-butoxycarbonyl)-1-piperazinyl]methyl}(trifluoro)borate. LCMS [M+H]⁺: 475. 4. ¹H NMR (400 MHz, Methanol-d₄) δ 8.56 (d, J=7.2 Hz, 1H), 8.09 (s, 1H), 7.66 (s, 1H), 7.01 (dd, J=7.4, 2.0 Hz, 1H), 6.35 (t, J=54.3 Hz, 1H), 4.12 (s, 1H), 3.92 (t, J=6.8 Hz, 2H), 3.53 (t, J=17.9 Hz, 1H), 3.10 (d, J=49.1 Hz, 4H), 2.89 (t, J=6.8 Hz, 3H), 2.59 (s, 1H), 2.12 (dd, J=35.5, 24.8 Hz, 1H), 1.98-1.45 (m, 7H), 1.26 (dq, J=21.1, 12.4, 11.8 Hz, 4H), 1.08-0.83 (m, 2H), NH proton not observed due to solvent exchange.


Example 247. Preparation of (R)-1-(5-((3-(difluoromethyl)-4-isobutylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[0983] Prepared using the method of Example 156, steps 1 and 4, wherein potassium (R)-((3-(difluoromethyl)-4-isobutylpiperazin-1-yl)methyl)trifluoroborate was used in place of potassium {[4-(tert-butoxycarbonyl)-1-piperazinyl]methyl}(trifluoro)borate. LCMS [M+H]⁺: 434.7. ¹H NMR (400 MHz, Methanol-d₄) δ 8.58 (d, J=7.2 Hz, 1H), 8.11 (s,


1H), 7.70 (s, 1H), 7.01 (dd, J=7.1, 1.9 Hz, 1H), 6.34 (t, J=54.1 Hz, 1H), 4.23 (d, J=24.8 Hz, 2H), 3.92 (t, J=6.7 Hz, 2H), 3.38 (s, 2H), 3.12 (m, 4H), 2.90 (t, J=6.7 Hz, 4H), 2.52 (s, 1H), 1.90 (s, 1H), 1.04-0.84 (m, 6H), NH proton not observed due to solvent exchange.

Example 248. Preparation of 1-(5-((1,4-diazepan-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0984] Prepared using the method of Example 192, steps 1 and 4, wherein potassium ((4-(tert-butoxycarbonyl)-1,4-diazepan-1-yl)methyl)trifluoroborate was used in place of tert-butyl (S)-2-methyl-4-((trifluoro-14-boraneyl)methyl)piperazine-1-carboxylate, potassium salt. LCMS [M+H]⁺: 343.1. ¹H NMR (400 MHz, Methanol-d₄) δ 8.56 (t, J=6.8 Hz, 1H), 8.10 (d, J=4.4 Hz, 1H), 7.71 (d, J=21.5 Hz, 1H), 7.14-6.99 (m, 1H), 4.27 (d, J=44.9 Hz, 2H), 3.92 (t, J=6.7 Hz, 2H), 3.68 (t, J=4.7 Hz, 2H), 3.60-3.50 (m, 3H), 3.45-3.37 (m, 2H), 3.26 (s, 1H), 2.89 (t, J=6.8 Hz, 2H), 2.29-2.16 (m, 2H), NH proton not observed due to solvent exchange.

Example 249. Preparation of 1-(5-((4-(cyclohexylmethyl)-1,4-diazepan-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0985] Prepared using the method of Example 192, wherein potassium ((4-(tert-butoxycarbonyl)-1,4-diazepan-1-yl)methyl)trifluoroborate was used in place of tert-butyl (S)-2-methyl-4-((trifluoro-14-boraneyl)methyl)piperazine-1-carboxylate, potassium salt. LCMS [M+H]⁺: 439.2. ¹H NMR (400 MHz, Methanol-d₄) δ 8.52 (d, J=7.2 Hz, 1H), 8.07 (s, 1H), 7.61 (s, 1H), 7.03 (dd, J=7.3, 2.0 Hz, 1H), 4.02 (s, 2H), 3.91 (t, J=6.8 Hz, 2H), 3.51 (s, 4H), 3.23 (s, 1H), 3.14-3.03 (m, 5H), 2.90 (t, J=6.8 Hz, 2H), 2.25-2.10 (m, 2H), 1.87-1.66 (m, 6H), 1.44-1.19 (m, 4H), 1.07 (t, J=11.7 Hz, 2H), NH proton not observed due to solvent exchange.

Example 250. Preparation of 1-(5-((4-isobutyl-1,4-diazepan-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0986] Prepared using the method of Example 192, wherein potassium ((4-(tert-butoxycarbonyl)-1,4-diazepan-1-yl)methyl)trifluoroborate was used in place of tert-butyl (S)-2-methyl-4-((trifluoro-14-boraneyl)methyl)piperazine-1-carboxylate, potassium salt and isobutyraldehyde was used in place of cyclohexanecarbaldehyde. LCMS [M+H]⁺: 399.3. ¹H NMR (400 MHz, CD₃OD) δ 8.44 (d, J=7.2 Hz, 1H), 8.01 (s, 1H), 7.49 (s, 1H), 7.02 (d, J=6.6 Hz, 1H), 3.88 (t, J=7.2 Hz, 2H), 3.73 (s, 2H), 3.14-3.07 (m, 4H), 2.90-2.69 (m, 8H), 1.97 (m, 3H), 0.97 (d, J=6.6 Hz, 6H), NH proton not observed due to solvent exchange.

Example 251. Preparation of 1-(5-(((1R,5S)-8-(3-fluorobenzyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



-continued




Example 251

Step 1. tert-butyl 1-((1R,5S)-3-((3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[0987] To a suspension of 1-(5-bromopyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)di-hydro-pyrimidine-2,4(1H,3H)-dione (600 mg, 1.30 mmol) in t-amyl-OH (6 mL) at rt was added Cs_2CO_3 (2.6 mL, 1.5 M aqueous solution), potassium (((1R,5S)-8-(tert-butoxycarbonyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)methyl)trifluoroborate (518 mg, 1.56 mmol) and $\text{Ad}_2\text{n-BuP-Pd-G}_3$ (44 mg, 0.06 mmol) in the glove-box. The reaction mixture was stirred at 90° C. for 16 h under inert atmosphere. The reaction mixture was then filtered and concentrated. The crude product was purified by silica gel chromatography (eluted with ethyl acetate in petroleum ether) to give tert-butyl 1-((1R,5S)-3-((3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a light yellow solid. LCMS $[\text{M}+\text{H}]^+$: 605.2.


[0988] Step 2: 1-(5-(((1R,5S)-8-(3-fluorobenzyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)di-hydro-pyrimidine-2,4(1H,3H)-dione was prepared using the method of Example 156, steps 2-4, wherein tert-butyl 1-((1R,5S)-3-((3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate was used in place of tert-butyl 4-((3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperazine-1-carboxylate and 1-(bromomethyl)-3-fluorobenzene was used in place of (bromomethyl)cyclohexane. LCMS $[\text{M}+\text{H}]^+$: 463.1. ^1H NMR (400 MHz, DMSO- d_6) δ =10.43 (br s, 1H), 8.57 (d, J =7.2 Hz, 1H), 7.99 (s, 1H), 7.41 (s, 1H), 7.38-7.28 (m, 1H), 7.20-7.17 (m, 2H), 7.07-6.98 (m, 1H), 6.91-6.88 (m, 1H), 3.76 (t, J =6.8 Hz, 2H), 3.49-3.47 (m, 4H), 3.04 (br s, 2H), 2.76 (t, J =6.8 Hz, 2H), 2.54-2.52 (m, 2H), 2.32-2.25 (m, 2H), 1.94-1.83 (m, 2H), 1.81-1.71 (m, 2H).

Example 252. Preparation of 1-(5-(((1R,5S)-8-(cyclohexylmethyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)di-hydro-pyrimidine-2,4(1H,3H)-dione




[0989] Prepared using the method of Example 156, steps 2-4, wherein tert-butyl 1-((1R,5S)-3-((3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate was used in place of tert-butyl 4-((3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperazine-1-carboxylate. LCMS $[\text{M}+\text{H}]^+$: 451.1. ^1H NMR (400 MHz, DMSO- d_6) δ 10.43 (br s, 1H), 8.56 (d, J =7.1 Hz, 1H), 7.99 (s, 1H), 7.40 (s, 1H), 6.89-6.87 (m, 1H), 3.75 (t, J =6.8 Hz, 2H), 3.45-3.42 (m, 2H), 3.01 (br s, 2H), 2.76 (t, J =6.8 Hz, 2H), 2.48 (br s, 2H), 2.24-2.21 (m, 2H), 2.06 (d, J =7.2 Hz, 2H), 1.87-1.53 (m, 9H), 1.09 (br s, 4H), 0.91-0.74 (m, 2H).


Example 253. Preparation of 1-(5-(((1R,5S)-8-(pyridin-3-ylmethyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)di-hydro-pyrimidine-2,4(1H,3H)-dione



[0990] Prepared using the method of Example 192, steps 2-4, wherein tert-butyl 1-((1R,5S)-3-((3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate was used in place of tert-butyl (S)-4-((3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperazine-1-carboxylate and nicotinaldehyde was used in place of cyclohexanecarbaldehyde. The reductive amination was carried out with NaBH_3CN , ZnCl_2 , and DIPEA in THF/EtOH. LCMS $[\text{M}+\text{H}]^+$: 446.1. ^1H NMR (400 MHz, CDCl_3) δ 8.58 (d, J =2.0 Hz, 1H), 8.50-8.48 (m, 1H), 8.35 (d, J =7.2 Hz, 1H), 7.92 (s, 1H), 7.79-7.67 (m, 2H), 7.27-7.24 (m, 2H), 6.91-6.90 (m, 1H), 3.88 (t, J =6.7 Hz, 2H), 3.51 (d, J =16.0 Hz, 4H), 3.10 (br s, 2H), 2.91 (t, J =6.8 Hz, 2H), 2.57-2.55 (m, 2H), 2.39-2.37 (m, 2H), 1.99-1.84 (m, 4H).

Example 254. Preparation of 1-(5-((4-isobutyl-2-oxopiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)di-hydro-pyrimidine-2,4(1H,3H)-dione





Step 1. 4-isobutylpiperazin-2-one

[0991] To a stirred solution of piperazin-2-one (500 mg, 4.99 mmol) in DCM (20 mL) was added TEA (2.0 mL 14.97 mmol) and isopropyl aldehyde (720 mg, 9.98 mmol) at rt. The mixture was stirred for 30 min and then $\text{NaBH}(\text{OAc})_3$ (2.1 g, 9.98 mmol) was added. The reaction was stirred at rt for 4 h and then diluted with DCM and water. The organic layer was dried over Na_2SO_4 , filtered and concentrated to afford 4-isobutylpiperazin-2-one (500 mg, crude). LCMS $[\text{M}+\text{H}]^+$: 157.0.


Step 2. 1-((3-iodopyrazolo[1,5-a]pyridin-5-yl)methyl)-4-isobutylpiperazin-2-one

[0992] To a stirred solution of 4-isobutylpiperazin-2-one (184 mg, 1.16 mmol) in THF (5 mL) at 0°C . was added NaH (88.0 mg, 2.23 mmol). The mixture was stirred for 30 min and allowed to warm to rt. A solution of (3-iodopyrazolo[1,5-a]pyridin-5-yl)methyl 4-methylbenzenesulfonate (500 mg, 1.16 mmol) in THF (5 mL) was added and the reaction was stirred for 1 h at rt. The reaction was diluted with EtOAc and water, the organic layer was dried over Na_2SO_4 , filtered and concentrated. The crude material was purified by silica gel chromatography (eluted with 60% EtOAc in hexanes) to afford 1-((3-iodopyrazolo[1,5-a]pyridin-5-yl)methyl)-4-isobutylpiperazin-2-one (130 mg). LCMS $[\text{M}+\text{H}]^+$: 413.0.

[0993] Step 3: 1-(5-((4-isobutyl-2-oxopiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione was prepared using the method of Example 1, steps 1 and 5, wherein 1-((3-iodopyrazolo[1,5-a]pyridin-5-yl)methyl)-4-isobutylpiperazin-2-one was used in place of 5-bromo-3-iodopyrazolo[1,5-a]pyridine. LCMS $[\text{M}+\text{H}]^+$: 399.1. ^1H NMR (400 MHz, DMSO- d_6): δ 10.45 (s, 1H), 8.60 (d, $J=7.6$ Hz, 1H), 8.16 (brs, 2H), 8.03 (s, 1H), 7.43 (s, 1H), 6.73 (d, $J=7.2$ Hz, 1H), 4.54 (s, 2H), 3.77 (t, $J=6.4$ Hz,


2H), 3.23-3.21 (m, 2H), 3.05 (s, 2H), 2.77 (t, $J=6.8$ Hz, 2H), 2.60-2.50 (m, 2H), 2.11-2.09 (m, 2H), 1.78-1.72 (m, 1H), 0.85 (d, $J=6.4$ Hz, 6H).


Example 255. Preparation of 1-(5-((4-(cyclohexylmethyl)-2-oxopiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[0994] Prepared using the method of Example 254, wherein cyclohexanecarbaldehyde was used in place of isopropyl aldehyde. LCMS $[\text{M}+\text{H}]^+$: 439.2. ^1H NMR (400 MHz, DMSO- d_6) δ 10.45 (s, 1H), 8.60 (d, $J=7.3$ Hz, 1H), 8.33 (s, 1H), 8.03 (s, 1H), 7.43 (s, 1H), 6.73 (dd, $J=7.3, 1.9$ Hz, 1H), 4.54 (s, 2H), 3.77 (t, $J=6.7$ Hz, 2H), 3.21 (t, $J=5.4$ Hz, 2H), 3.04 (s, 2H), 2.77 (t, $J=6.7$ Hz, 2H), 2.58 (t, $J=5.5$ Hz, 2H), 2.53-2.50 (m, 1H), 2.14 (d, $J=7.3$ Hz, 2H), 1.76-1.56 (m, 5H), 1.47 (ddt, $J=10.3, 6.2, 3.3$ Hz, 1H), 1.29-1.07 (m, 3H), 0.82 (q, $J=11.5$ Hz, 2H).

Example 256. Preparation of 1-(5-(((3S,5R)-3,5-dimethyl-4-((tetrahydro-2H-pyran-4-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione





Step 1. tert-butyl (3S,5R)-3,5-dimethyl-4-((tetrahydro-2H-pyran-4-yl)methyl)piperazine-1-carboxylate

[0995] To a stirred solution of tert-butyl (3S,5R)-3,5-dimethylpiperazine-1-carboxylate (200 mg, 0.933 mmol) in DCM (5 mL) 0° C. was added tetrahydro-2H-pyran-4-carbaldehyde (159 mg, 1.39 mmol) and TEA (0.39 mL, 2.79 mmol). The mixture was stirred for 30 min and then NaBH(OAc)₃ (395 mg, 1.86 mmol) was added and the mixture was stirred for 2 h at rt. The reaction was diluted with DCM and water the organic layer was dried over Na₂SO₄, filtered and concentrated. The crude compound was purified by silica gel chromatography (eluting with 10% MeOH in DCM) to obtain tert-butyl (3S,5R)-3,5-dimethyl-4-((tetrahydro-2H-pyran-4-yl)methyl)piperazine-1-carboxylate (200 mg, 0.64 mmol, 69% yield). LCMS [M+H]⁺: 313.2.

Step 2. (2S,6R)-2,6-dimethyl-1-((tetrahydro-2H-pyran-4-yl)methyl)piperazine hydrochloride

[0996] To a stirred solution of tert-butyl (3S,5R)-3,5-dimethyl-4-((tetrahydro-2H-pyran-4-yl)methyl)piperazine-1-carboxylate (200 mg, 0.64 mmol) in DCM (2 mL) was added 4M HCl in dioxane (2 mL) and the mixture was stirred for 2 h at rt. The reaction was then concentrated to afford crude (2S,6R)-2,6-dimethyl-1-((tetrahydro-2H-pyran-4-yl)methyl)piperazine hydrochloride (200 mg, crude) which was used without further purification. LCMS [M+H]⁺: 213.2.


Step 3. 5-(((3S,5R)-3,5-dimethyl-4-((tetrahydro-2H-pyran-4-yl)methyl)piperazin-1-yl)methyl)-3-iodopyrazolo[1,5-a]pyridine

[0997] To a stirred solution of (3-iodopyrazolo[1,5-a]pyridin-5-yl)methyl 4-methylbenzenesulfonate (200 mg, 0.467 mmol) in DMF (4.0 mL) was added Cs₂CO₃ (456 mg, 1.401 mmol). The mixture was stirred for 30 min at 0° C. and then a solution of (2S,6R)-2,6-dimethyl-1-((tetrahydro-2H-pyran-4-yl)methyl)piperazine hydrochloride (99.1 mg, 0.467 mmol) in DMF (1 mL) was added. The mixture was stirred for 1 h at rt. The reaction was diluted with EtOAc and water and the organic layer was dried over Na₂SO₄, filtered

and concentrated. The crude compound was purified by silica gel chromatography (eluting with 30% EtOAc in hexanes) to obtain 5-(((3S,5R)-3,5-dimethyl-4-((tetrahydro-2H-pyran-4-yl)methyl)piperazin-1-yl)methyl)-3-iodopyrazolo[1,5-a]pyridine (80 mg, 0.17 mmol, 40% yield). LCMS [M+H]⁺: 469.0.


[0998] Step 4: 1-(5-(((3S,5R)-3,5-dimethyl-4-((tetrahydro-2H-pyran-4-yl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione was prepared using the method of Example 1, steps 1 and 5, wherein 5-(((3S,5R)-3,5-dimethyl-4-((tetrahydro-2H-pyran-4-yl)methyl)piperazin-1-yl)methyl)-3-iodopyrazolo[1,5-a]pyridine was used in place of 5-bromo-3-iodopyrazolo[1,5-a]pyridine. LCMS [M+H]⁺: 455.4. ¹H NMR (400 MHz, CD₃OD): δ 8.46 (d, J=7.2 Hz, 1H), 8.02 (s, 1H), 7.49 (s, 1H), 6.99-6.97 (m, 1H), 3.97-3.87 (m, 4H), 3.63 (s, 2H), 3.42 (t, J=11.6 Hz, 4H), 3.05-2.98 (m, 4H), 2.88 (t, J=6.4 Hz, 2H), 2.30-2.25 (m, 2H), 1.94-1.93 (m, 2H), 1.79-1.75 (m, 2H), 1.42-1.28 (m, 7H).

Example 257. Preparation of 1-(5-(((3S,5R)-4-isobutyl-3,5-dimethylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[0999] Prepared using the method of Example 256, wherein isobutyraldehyde was used in place of tetrahydro-2H-pyran-4-carbaldehyde. LCMS [M+H]⁺: 413.2. ¹H NMR (400 MHz, Methanol-d₄) δ 8.46 (d, J=7.1 Hz, 1H), 8.03 (s, 1H), 7.50 (s, 1H), 6.99 (dd, J=7.1, 1.8 Hz, 1H), 3.90 (t, J=6.7 Hz, 2H), 3.60 (s, 2H), 3.30-3.13 (m, 2H), 3.00-2.77 (m, 6H), 2.21 (t, J=10.7 Hz, 2H), 2.00-1.85 (m, 1H), 1.26 (s, 2H), 1.25 (s, 2H), 1.05 (s, 2H), 1.04 (s, 2H). NH proton not observed due to solvent exchange.

Example 258. Preparation of 1-(5-(((3S,5S)-4-(cyclohexylmethyl)-3,5-dimethylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[1000] Prepared using the method of Example 256, wherein tert-butyl (3S,5S)-3,5-dimethylpiperazine-1-carboxylate was used in place of tert-butyl (3S,5R)-3,5-dimethylpiperazine-1-carboxylate and cyclohexanecarbaldehyde was used in place of tetrahydro-2H-pyran-4-carbaldehyde. LCMS $[M+H]^+$: 453.2. 1H NMR (400 MHz, Methanol- d_4) δ 8.51 (d, $J=7.3$ Hz, 1H), 8.06 (s, 1H), 7.56 (s, 1H), 7.01 (d, $J=7.7$ Hz, 1H), 3.95-3.64 (m, 6H), 3.20-2.49 (m, 8H), 1.91-1.65 (m, 6H), 1.59-1.17 (m, 9H), 1.08 (q, $J=12.0$ Hz, 2H). NH proton not observed due to solvent exchange.

Example 259. Preparation of 1-(5-(((3S,5S)-4-isobutyl-3,5-dimethylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 259)



[1001] Prepared using the method of Example 256, wherein tert-butyl (3S,5S)-3,5-dimethylpiperazine-1-carboxylate was used in place of tert-butyl (3S,5R)-3,5-dimethylpiperazine-1-carboxylate and isobutyraldehyde was used in place of tetrahydro-2H-pyran-4-carbaldehyde. LCMS $[M+H]^+$: 413.2. 1H NMR (400 MHz, Methanol- d_4) δ 8.47 (d, $J=7.3$ Hz, 1H), 8.07 (s, 1H), 8.03 (s, 1H), 7.49 (s, 1H), 7.00 (d, $J=7.1$ Hz, 1H), 3.90 (t, $J=6.7$ Hz, 2H), 3.82 (s, 1H), 3.76-3.51 (m, 3H), 3.09 (dd, $J=13.4, 9.7$ Hz, 1H), 3.04-2.93 (m, 2H), 2.93-2.75 (m, 3H), 2.62 (d, $J=12.5$ Hz, 1H), 2.46 (t, $J=11.2$ Hz, 1H), 2.16-2.01 (m, 1H), 1.50-1.33 (m, 7H), 1.06 (t, $J=6.9$ Hz, 7H). NH proton not observed due to solvent exchange.


Example 260. Preparation of (S)-1-(5-((octahydro-2H-pyrrolo[1,2-a]pyrazin-2(1H)-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 260)



[1002] Prepared using the method of Example 156, steps 1 and 4, wherein potassium (S)-trifluoro((octahydro-2H-pyrrolo[1,2-a]pyrazin-2(1H)-yl)methyl)borate was used in place of


potassium {[4-(tert-butoxycarbonyl)-1-piperazinyl]methyl} (trifluoro)borate. LCMS $[M+H]^+$: 369.3. 1H NMR (400 MHz, METHANOL- d_4) δ 8.46 (d, $J=7.2$ Hz, 1H), 8.03 (s, 1H), 7.52 (d, $J=0.4$ Hz, 1H), 7.03-7.01 (m, 1H), 3.93-3.89 (m, 2H), 3.74-3.57 (m, 2H), 3.12-2.99 (m, 3H), 2.95-2.82 (m, 3H), 2.42-2.29 (m, 2H), 2.25-2.19 (m, 2H), 2.05-1.95 (m, 1H), 1.91-1.78 (m, 3H), 1.50-1.36 (m, 1H).

Example 261. Preparation of (R)-1-(5-((hexahydro-pyrrolo[1,2-a]pyrazin-2(1H)-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[1003] Prepared using the method of Example 156, steps 1 and 4, wherein potassium (R)-trifluoro((hexahydro-pyrrolo[1,2-a]pyrazin-2(1H)-yl)methyl)borate was used in place of potassium {[4-(tert-butoxycarbonyl)-1-piperazinyl]methyl} (trifluoro)borate. LCMS $[M+H]^+$: 369.1. 1H NMR (400 MHz, METHANOL- d_4) δ 8.47-8.41 (m, 1H), 8.01 (s, 1H), 7.50 (d, $J=0.8$ Hz, 1H), 7.06-6.92 (m, 1H), 3.95-3.86 (m, 2H), 3.71-3.60 (m, 2H), 3.10-2.97 (m, 3H), 2.93-2.83 (m, 3H), 2.39-2.29 (m, 2H), 2.21 (d, $J=8.8$ Hz, 2H), 2.01-1.96 (m, 1H), 1.86-1.74 (m, 3H), 1.47-1.33 (m, 1H).

Example 262. Preparation of (S)-1-(5-((octahydro-2H-pyrrolo[1,2-a]pyrazin-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[1004] Prepared using the method of Example 156, steps 1 and 4, wherein potassium (S)-trifluoro((octahydro-2H-pyrrolo[1,2-a]pyrazin-2-yl)methyl)borate was used in place of potassium {[4-(tert-butoxycarbonyl)-1-piperazinyl]methyl} (trifluoro)borate. LCMS $[M+H]^+$: 383.1. 1H NMR (400 MHz, Methanol- d_4) δ 8.48 (d, $J=7.2$ Hz, 1H), 8.04 (s, 1H), 7.54 (s, 1H), 6.99 (dd, $J=7.2, 1.9$ Hz, 1H), 4.88 (br. s, 2H), 3.90 (t, $J=6.8$ Hz, 2H), 3.76 (s, 2H), 3.58-3.37 (m, 2H), 3.22-3.01 (m, 3H), 2.89 (t, $J=6.8$ Hz, 2H), 2.87-2.74 (m, 2H), 2.67-2.52 (m, 2H), 2.52-2.36 (m, 2H).

Example 263. Preparation of (R)-1-(5-((hexahydropyrazino[2,1-c][1,4]oxazin-8(1H)-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[1005] Prepared using the method of Example 156, steps 1 and 4, wherein potassium (R)-trifluoro((hexahydropyrazino[2,1-c][1,4]oxazin-8(1H)-yl)methyl)borate was used in place of potassium {[4-(tert-butoxycarbonyl)-1-piperazinyl]methyl}(trifluoro)borate. LCMS [M+H]⁺: 385.3. ¹H NMR (400 MHz, Methanol-d₄) δ 8.44 (d, J=7.1 Hz, 1H), 8.25 (s, 1H), 8.00 (s, 1H), 7.50 (s, 1H), 6.97 (dd, J=7.1, 1.8 Hz, 1H), 3.91-3.82 (m, 3H), 3.72-3.58 (m, 4H), 3.24 (d, J=11.1 Hz, 1H), 2.98-2.71 (m, 6H), 2.61-2.38 (m, 4H), 1.99 (t, J=10.9 Hz, 1H). NH proton not observed due to solvent exchange.

Example 264. Preparation of (S)-1-(5-((hexahydropyrazino[2,1-c][1,4]oxazin-8(1H)-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[1006] Prepared using the method of Example 156, steps 1 and 4, wherein potassium (S)-trifluoro((hexahydropyrazino[2,1-c][1,4]oxazin-8(1H)-yl)methyl)borate was used in place of potassium {[4-(tert-butoxycarbonyl)-1-piperazinyl]methyl}(trifluoro)borate. LCMS [M+H]⁺: 385.2. ¹H NMR (400 MHz, CD₃OD) δ 8.45 (d, J=7.2 Hz, 1H), 8.24 (brs, 1H), 8.01 (s, 1H), 7.50 (s, 1H), 6.98 (dd, J=7.2 Hz, 1.2 Hz, 1H), 3.89-3.84 (m, 3H), 3.72-3.61 (m, 4H), 3.27-3.24 (m, 1H), 2.97-2.78 (m, 6H), 2.62-2.44 (m, 4H), 2.03-1.98 (m, 1H).

Example 265. Preparation of 1-(5-(((2S,4R)-1-(((1R,4S)-4-methoxycyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[1007] Prepared using the method of Example 141, wherein trans-4-methoxycyclohexane-1-carbaldehyde was used in place of 4,4-difluorocyclohexane-1-carbaldehyde. LCMS [M+H]⁺: 468.1. ¹H NMR (400 MHz, CD₃OD) δ 8.55 (s, 1H), 8.43 (d, J=7.2 Hz, 1H), 8.01 (s, 1H), 7.36 (s, 1H), 6.83 (dd, J=7.1, 1.9 Hz, 1H), 3.89 (t, J=6.7 Hz, 2H), 3.62 (s, 1H), 3.35 (s, 3H), 3.16 (ddd, J=15.1, 7.5, 4.4 Hz, 3H), 2.89 (t, J=6.7 Hz, 4H), 2.68 (d, J=7.3 Hz, 1H), 2.25-2.07 (m, 3H), 1.96-1.56 (m, 7H), 1.38-1.04 (m, 8H).


Example 266. Preparation of 1-(5-(((1R,4R)-4-ethoxycyclohexyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[1008] Prepared using the method of Example 192, steps 3-4, wherein trans-4-ethoxycyclohexane-1-carbaldehyde [see US2016/122318, 2016, A1] was used in place of cyclohexanecarbaldehyde. LCMS [M+H]⁺: 468.2. ¹H NMR (500 MHz, DMSO-d₆) δ 10.45 (d, J=4.5 Hz, 1H), 8.60 (dt, J=7.1, 1.4 Hz, 1H), 8.02 (d, J=1.8 Hz, 1H), 7.46-7.28 (m, 1H), 6.80 (dd, J=7.2, 1.9 Hz, 1H), 3.78 (td, J=6.7, 2.8 Hz, 2H), 3.45 (p, J=6.6 Hz, 4H), 3.26-3.08 (m, 2H), 2.92-2.83 (m, 3H), 2.79 (td, J=6.7, 2.0 Hz, 2H), 2.61 (d, J=6.6 Hz, 2H), 1.98 (d, J=12.1 Hz, 2H), 1.92-1.65 (m, 6H), 1.49 (q, J=13.1 Hz, 2H), 1.11 (s, 2H), 1.09 (t, J=7.0 Hz, 3H), 0.99 (dd, J=14.4, 11.3 Hz, 2H).

Example 267. Preparation of 1-(5-(1-(1-isobutylpiperidin-4-yl)ethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione





Step 1. 5-(trimethylstannyl)pyrazolo[1,5-a]pyridine

[1009] To a stirred solution of 5-bromopyrazolo[1,5-a]pyridine (3 g, 15.2 mmol) in dioxane (40 mL) was added $\text{Pd}(\text{PPh}_3)_4$ (877 mg, 0.76 mmol), Hexamethylditin (4.97 g, 15.2 mmol) and the reaction was stirred for 4 h at 90° C. After completion, the reaction mixture was filtered through celite and concentrated to afford 5-(trimethylstannyl)pyrazolo[1,5-a]pyridine (3.2 g, crude). The material was used without further purification. LCMS $[\text{M}+\text{H}]^+$: 282.9.

Step 2. benzyl 4-(pyrazolo[1,5-a]pyridine-5-carboxylate)piperidine-1-carboxylate

[1010] To a stirred solution of 5-(trimethylstannyl)pyrazolo[1,5-a]pyridine (3.0 g, 10.7 mmol) in THF (30 mL) was added benzyl 4-(chlorocarbonyl)piperidine-1-carboxylate (3.0 g, 10.7 mmol) and the solution was de-gassed by

bubbling nitrogen through it for 10 min. Allylpalladium(II) chloride dimer (390 mg, 1.07 mmol) and molecular sieves (500 mg) were added and the reaction was stirred for 4 h at 60° C. After completion, the reaction mixture was concentrated. The crude compound was purified by silica gel chromatography (eluting with 50% EtOAc in hexanes) to afford benzyl 4-(pyrazolo[1,5-a]pyridine-5-carboxylate)piperidine-1-carboxylate (1.8 g, 4.95 mmol, 46% yield). LCMS $[\text{M}+\text{H}]^+$: 364.0.

Step 3. benzyl 4-(1-(pyrazolo[1,5-a]pyridin-5-yl)vinyl)piperidine-1-carboxylate

[1011] To a stirred solution of methyl triphenylphosphonium bromide (1.47 g, 4.12 mmol) in THF (10 mL) at 0° C., was slowly added 1M KOtBu (4.1 mL, 4.12 mmol) resulting in a yellow color. The reaction mixture was stirred at rt for 30 min and then a solution of benzyl 4-(pyrazolo[1,5-a]pyridine-5-carboxylate)piperidine-1-carboxylate (0.50 g, 1.37 mmol) in THF (2 mL) was added dropwise to at 0° C. The reaction mixture was allowed to stirred at rt for 3 h. After completion, the reaction was quenched with a solution of saturated aqueous NH_4Cl and the mixture was diluted with ethyl acetate and washed with water. The organic layer was dried over Na_2SO_4 , filtered and concentrated. The crude compound was purified by silica gel chromatography (eluting with 30% EtOAc in hexanes) to afford benzyl 4-(1-(pyrazolo[1,5-a]pyridin-5-yl)vinyl)piperidine-1-carboxylate (0.21 g, 0.58 mmol, 42% yield). ^1H NMR (400 MHz, CD_3OD) δ 8.41 (d, $J=7.2$ Hz, 1H), 7.93 (d, $J=2.1$ Hz, 1H), 7.45 (s, 1H), 7.36-7.30 (m, 4H), 6.76 (dd, $J=7.2$ Hz, 2.1 Hz, 1H), 6.49 (d, $J=2.1$ Hz, 1H), 5.34 (s, 1H), 5.13-5.11 (m, 3H), 4.28 (s, 2H), 2.89-2.82 (m, 2H), 2.63-2.56 (m, 2H), 1.85-1.81 (m, 2H), 1.43-1.40 (m, 2H).

Step 4. 5-(1-(piperidin-4-yl)ethyl)pyrazolo[1,5-a]pyridine

[1012] To a stirred solution of benzyl 4-(1-(pyrazolo[1,5-a]pyridin-5-yl)vinyl)piperidine-1-carboxylate (0.20 g, 0.55 mmol) in EtOH (10 mL) under a nitrogen atmosphere was added Pd/C (0.10 g). The flask was evacuated and refilled with hydrogen from a balloon and stirred for 16 h at rt. After completion, the reaction mixture was filtered through celite and washed through with EtOH. The filtrate was concentrated to afford 5-(1-(piperidin-4-yl)ethyl)pyrazolo[1,5-a]pyridine (0.18 g, crude). The compound was used in the next step without further purification.

Step 5. tert-butyl 4-(1-(pyrazolo[1,5-a]pyridin-5-yl)ethyl)piperidine-1-carboxylate

[1013] To a stirred solution of 5-(1-(piperidin-4-yl)ethyl)pyrazolo[1,5-a]pyridine (0.18 g, 0.78 mmol) in DCM (7 mL) at 0° C. was added Et_3N (0.33 mL, 2.4 mmol) followed by di-tert-butyl dicarbonate (0.26 g, 1.17 mmol) and DMAP (9.59 mg, 0.078 mmol). The reaction mixture was stirred at rt for 16 h. After completion, the reaction was diluted with water and extracted with EtOAc. The organic layer was washed with brine, dried over Na_2SO_4 , filtered and concentrated. The crude compound was purified by silica gel chromatography (eluting with 10-15% EtOAc in hexanes) to afford tert-butyl 4-(1-(pyrazolo[1,5-a]pyridin-5-yl)ethyl)piperidine-1-carboxylate (110 mg, 0.33 mmol, 42% yield). ^1H NMR (300 MHz, CD_3OD) δ 8.39 (d, $J=7.5$ Hz, 1H), 7.91 (s, 1H), 6.56 (dd, $J=7.2$ Hz, 2.1 Hz, 1H), 6.40 (d, $J=1.2$ Hz, 1H),


2.65-2.45 (m, 3H), 1.84-1.80 (m, 1H), 1.61-1.49 (m, 1H), 1.43-1.38 (m, 10H), 1.28-1.02 (m, 6H).

Step 6. tert-butyl 4-(1-(3-iodopyrazolo[1,5-a]pyridin-5-yl)ethyl)piperidine-1-carboxylate

[1014] To a stirred solution of tert-butyl 4-(1-(pyrazolo[1,5-a]pyridin-5-yl)ethyl)piperidine-1-carboxylate (0.10 g 0.30 mmol) in DMF (5 mL) at 0° C. was added NIS (68.2 mg, 0.30 mmol) under an inert atmosphere. The reaction mixture was stirred at rt for 3 h. After completion, the reaction was quenched with water and the yellow precipitate that formed was collected by filtration. The solid was washed with water and dried under vacuum to afford tert-butyl 4-(1-(3-iodopyrazolo[1,5-a]pyridin-5-yl)ethyl)piperidine-1-carboxylate (120 mg, 0.26 mmol, 86% yield). LCMS [M+H]⁺: 456.0.

[1015] Step 7: tert-butyl 4-(1-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)ethyl)piperidine-1-carboxylate was prepared using the method of Example 1, step 1, wherein tert-butyl 4-(1-(3-iodopyrazolo[1,5-a]pyridin-5-yl)ethyl)piperidine-1-carboxylate was used in place of 5-bromo-3-iodopyrazolo[1,5-a]pyridine. LCMS [M+H]⁺: 592.2. Step 8: 1-(5-(1-(1-isobutylpiperidin-4-yl)ethyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione was prepared using the method of Example 141, steps 6-8, wherein tert-butyl 4-(1-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)ethyl)piperidine-1-carboxylate was used in place of tert-butyl (2S,4R)-4-((3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperidine-1-carboxylate and isobutyraldehyde was used in place of 4,4-difluorocyclohexane-1-carbaldehyde. LCMS [M+H]⁺: 398.3. ¹H NMR (400 MHz, Methanol-d₄) δ 8.51 (s, 1H), 8.43 (d, J=7.2 Hz, 1H), 8.01 (s, 1H), 7.36 (s, 1H), 6.84 (dd, J=7.4, 1.9 Hz, 1H), 3.89 (t, J=6.8 Hz, 2H), 3.74-3.61 (m, 1H), 3.23-3.14 (m, 2H), 2.95 (s, 1H), 2.89 (t, J=6.7 Hz, 2H), 2.69 (d, J=7.2 Hz, 2H), 2.21 (s, 1H), 2.06 (d, J=19.5 Hz, 2H), 1.81 (dd, J=37.0, 15.2 Hz, 7H), 1.59 (s, 1H), 1.35 (t, J=11.2 Hz, 5H).


Example 268. Preparation of 1-(5-(((2S,4R)-1-((4,4-difluorocyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione



[1016] Methanesulfonic acid (2.0 mL, 31 mmol) was added to 1-(5-(((2S,4R)-1-((4,4-difluorocyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione (1.24 g, 1.99 mmol) in DCM (8 mL). The resulting pale red solution was stirred overnight at 40° C. The reaction mixture was quenched with 50% aqueous sodium bicarbonate solution and extracted with 4:1 dichlo-


romethane:trifluoroethanol three times. The combined organic phases were dried over Na₂SO₄, filtered and concentrated. The crude material was purified by silica gel chromatography (eluting with 15-100% 3:1 EtOAc:EtOH in heptane, 0.1% TEA as modifier) to afford impure product. The material was further purified by C18 reverse phase chromatography (eluting with 25-75% acetonitrile in water, 0.1% NH₄OH as modifier) and the product-containing fractions were assembled and poured into a pH 7 phosphate buffered solution. The aqueous phase was extracted with 4:1 dichloromethane:trifluoroethanol three times. The combined organic phases were concentrated in vacuo, diluted with 2:1 acetonitrile:water (3 mL) and lyophilized to afford 1-(5-(((2S,4R)-1-((4,4-difluorocyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione (3.5 mg, 0.0072 mmol, 0.36% yield) as a white solid. LCMS [M+H]⁺: 472.3. ¹H NMR (400 MHz, DMSO-d₆) δ 11.48 (s, 1H), 8.61 (d, J=7.1 Hz, 1H), 8.12 (s, 1H), 7.70 (d, J=7.8 Hz, 1H), 7.33 (s, 1H), 6.85 (dd, J=7.2, 1.8 Hz, 1H), 5.69 (d, J=7.8 Hz, 1H), 2.96-2.83 (m, 1H), 2.57-2.50 (m, 2H), 2.46-2.35 (m, 2H), 2.17 (qd, J=12.5, 7.2 Hz, 2H), 2.02-1.90 (m, 2H), 1.90-1.84 (m, 1H), 1.84-1.62 (m, 4H), 1.59-1.44 (m, 2H), 1.44-1.33 (m, 2H), 1.22-1.12 (m, 1H), 1.05 (q, J=13.7, 12.7 Hz, 2H), 0.86 (d, J=6.6 Hz, 3H).

Example 269. Preparation of 1-(5-(azepan-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[1017] Prepared using the method of Example 1, steps 2 and 5, wherein tert-butyl 4-methyleneazepane-1-carboxylate [see WO2021/158829, 2021, A1] was used in place of tert-butyl 4-methylenepiperidine-1-carboxylate in step 2. LCMS [M+H]⁺: 341.8. ¹H NMR (400 MHz, CD₃OD) δ 8.43 (s, 1H), 8.41 (s, 1H), 8.00 (d, J=2.2 Hz, 1H), 7.36 (s, 1H), 6.84 (dd, J=7.1, 2.0 Hz, 1H), 3.88 (td, J=6.8, 2.1 Hz, 2H), 3.28-3.21 (m, 2H), 3.20-3.04 (m, 2H), 2.89 (td, J=6.8, 1.8 Hz, 2H), 2.75-2.61 (m, 2H), 2.09-1.89 (m, 4H), 1.86-1.73 (m, 1H), 1.62 (ddt, J=18.5, 13.0, 6.5 Hz, 1H), 1.46-1.29 (m, 1H). Missing NH due to solvent exchange.

Example 270. Preparation of tert-butyl 4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)azepane-1-carboxylate




[1018] TEA (0.061 mL, 0.44 mmol) and di-tert-butyl dicarbonate (0.015 mL, 0.32 mmol) were added to a solution of 1-(5-(azepan-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione (Example 269) (100 mg, 0.21 mmol) in DCM (5 mL) at rt. The mixture was stirred at rt for 4 h and then partitioned between DCM and water. The organic layer was separated, washed with brine, dried over sodium sulfate, filtered and concentrated. The residue was dissolved in DMSO, filtered through a 1 micron filter and purified by reverse phase HPLC using ACN/Water/0.1% formic acid. The fractions containing the product were combined, frozen and lyophilized to afford tert-butyl 4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)azepane-1-carboxylate as an off-white solid. LCMS $[M+H]^+$: 385.9. 1H NMR (400 MHz, CD_3OD) δ 8.40 (d, $J=7.1$ Hz, 1H), 7.98 (d, $J=2.7$ Hz, 1H), 7.34 (s, 1H), 6.82 (dd, $J=7.1, 1.9$ Hz, 1H), 3.88 (t, $J=6.7$ Hz, 2H), 3.55 (dt, $J=16.6, 5.4$ Hz, 1H), 3.37 (dd, $J=15.6, 8.8$ Hz, 2H), 3.25-3.12 (m, 1H), 2.89 (t, $J=6.8$ Hz, 2H), 2.64 (d, $J=7.1$ Hz, 2H), 1.90-1.73 (m, 4H), 1.56 (s, 1H), 1.45 (d, $J=5.7$ Hz, 9H), 1.40-1.19 (m, 2H). Missing NH due to solvent exchange.

Example 271. Preparation of 1-(5-((1-methylazepan-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




[1019] Prepared from 1-(5-(azepan-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione (Example 269) using the method of Example 109, wherein paraformaldehyde was used in place of isobutyraldehyde. LCMS $[M+H]^+$: 355.9. 1H NMR (400 MHz, CD_3OD) δ 8.46-8.35 (m, 2H), 8.00 (s, 1H), 7.35 (s, 1H), 6.83 (dd, $J=7.0, 1.9$ Hz, 1H), 3.89 (t, $J=6.8$ Hz, 2H), 3.22 (d, $J=12.2$ Hz, 3H), 2.94-2.83 (m, 5H), 2.74-2.60 (m, 2H), 2.09 (ddt, $J=10.3, 7.2, 3.6$ Hz, 1H), 2.01-1.80 (m, 4H), 1.77-1.65 (m, 1H), 1.48-1.28 (m, 2H).

Example 272. Preparation of 1-(5-((1-(cyclohexylmethyl)azepan-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




[1020] Prepared from 1-(5-(azepan-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione (Example 269) using the method of Example 109, wherein cyclohexanecarbaldehyde was used in place of isobutyraldehyde. LCMS $[M+H]^+$: 438.2. 1H NMR (400 MHz, CD_3OD) δ 8.42 (d, $J=7.3$ Hz, 1H), 8.38 (s, 1H), 8.01 (s, 1H), 7.35 (s, 1H), 6.83 (dd, $J=7.1, 1.9$ Hz, 1H), 3.89 (t, $J=6.7$ Hz, 2H), 3.41 (s, 2H), 3.22 (s, 2H), 2.98 (d, $J=6.7$ Hz, 2H), 2.89 (t, $J=6.8$ Hz, 2H), 2.68 (qd, $J=13.5, 7.1$ Hz, 2H), 2.11-1.61 (m, 12H), 1.45-1.16 (m, 4H), 1.04 (q, $J=12.1$ Hz, 2H).

Example 273. Preparation of 1-(5-(((2S,4R)-2-methyl-1-(2,2,3,3-tetrafluoropropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



-continued




Step 1: 1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

[1021] TFA (2 mL) was added to 3-(2,4-dimethoxybenzyl)-1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (340 mg, 0.57 mmol). The reaction mixture was stirred for 16 h at 90° C. and then concentrated. The residue was taken up in DCM, stirred with basic resin, filtered and concentrated again. The crude material was triturated sequentially with pentane and diethyl ether to provide crude 1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (360 mg) as a yellow semi solid. LCMS $[M+H]^+$: 341.9.

Step 2: 1-(5-(((2S,4R)-2-methyl-1-(2,2,3,3-tetrafluoropropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione


[1022] To a solution of 1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (150 mg, 0.43 mmol) DCM (5 mL) was added triethylamine (131 mg, 1.29 mmol) and 2,2,3,3-tetrafluoropropyl trifluoromethanesulfonate (137 mg, 0.51 mmol). The reaction mixture was allowed to stir for 16 h at rt. The reaction mixture was diluted with water (20 mL) and extracted with DCM (2x30 mL). The organic layer was washed with brine solution (10 mL), dried over sodium sulfate and concentrated. The crude material was purified by reverse phase HPLC using ACN/Water/0.1% formic acid. The fractions containing the product were combined, frozen and lyophilized to afford 1-(5-(((2S,4R)-2-methyl-1-(2,2,3,3-tetrafluoropropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione as a white solid. LCMS $[M+H]^+$: 456.1. 1H NMR (300 MHz, CD_3OD) δ 8.40 (d, $J=7.1$ Hz, 1H), 8.23 (s, 1H), 7.98 (s, 1H), 7.33 (s, 1H), 6.81 (dd, $J=7.1$, 1.9 Hz, 1H), 6.39-5.96 (m, 1H), 3.88 (t, $J=6.8$ Hz, 2H), 3.04 (d, $J=14.0$ Hz, 2H), 2.89 (t, $J=6.8$ Hz, 3H), 2.81-2.69 (m, 1H), 2.62 (dd, $J=17.2$, 5.7 Hz, 3H), 2.00 (d, $J=23.6$ Hz, 1H), 1.65-1.47 (m, 3H), 1.35-1.27 (m, 1H), 1.02 (d, $J=6.7$ Hz, 3H).

Example 274. Preparation of 1-(5-(((2S,4R)-1-(2,2-difluoroethyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[1023] Prepared by the method of Example 273 wherein 2,2-difluoroethyl trifluoromethanesulfonate was used in place of 2,2,3,3-tetrafluoropropyl trifluoromethanesulfonate in step 2. LCMS $[M+H]^+$: 406.2. 1H NMR (400 MHz, MeOD) δ 8.41 (dd, $J=7.2$, 0.9 Hz, 1H), 8.17 (s, 1H), 7.99 (s, 1H), 7.34 (dd, $J=1.9$, 1.0 Hz, 1H), 6.82 (dd, $J=7.2$, 1.9 Hz, 1H), 5.98 (tt, $J=55.7$, 4.2 Hz, 1H), 3.88 (t, $J=6.8$ Hz, 2H), 3.21 (d, $J=6.5$ Hz, 1H), 3.05-2.92 (m, 2H), 2.89 (t, $J=6.8$ Hz, 2H), 2.81 (dd, $J=8.0$, 3.4 Hz, 2H), 2.64 (d, $J=7.3$ Hz, 2H), 2.10-1.99 (m, 1H), 1.69 (dd, $J=13.5$, 3.9 Hz, 1H), 1.60 (dd, $J=7.5$, 4.2 Hz, 2H), 1.47-1.36 (m, 1H), 1.10 (d, $J=6.7$ Hz, 3H). NH proton not observed due to solvent exchange.

Example 275. Preparation of 1-(5-(((2S,4R)-2-methyl-1-(2,2,2-trifluoroethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[1024] Prepared by the method of Example 273 wherein 2,2,2-trifluoroethyl trifluoromethanesulfonate was used in place of 2,2,3,3-tetrafluoropropyl trifluoromethanesulfonate in step 2. LCMS $[M+H]^+$: 424.0. 1H NMR (400 MHz, MeOD) δ 8.40 (dd, $J=7.2$, 0.9 Hz, 1H), 7.98 (s, 1H), 7.34 (dd, $J=1.9$, 0.9 Hz, 1H), 6.81 (dd, $J=7.2$, 1.8 Hz, 1H), 3.88 (t, $J=6.8$ Hz, 2H), 3.17-2.95 (m, 3H), 2.89 (t, $J=6.8$ Hz, 2H), 2.81-2.65 (m, 2H), 2.61 (d, $J=7.3$ Hz, 2H), 1.97 (h, $J=6.1$ Hz, 1H), 1.66-1.49 (m, 3H), 1.34 (dtd, $J=12.9$, 10.7, 4.4 Hz, 1H), 1.02 (d, $J=6.6$ Hz, 3H). NH proton not observed due to solvent exchange.

Example 276. Preparation of 1-(5-(((2S,4R)-2-methyl-1-(3,3,3-trifluoropropyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




[1025] Prepared by the method of Example 273 wherein 3,3,3-trifluoropropyl trifluoromethanesulfonate was used in place of 2,2,3,3-tetrafluoropropyl trifluoromethanesulfonate in step 2. LCMS $[M+H]^+$: 438.0. 1H NMR (400 MHz, MeOD) δ 8.42 (d, $J=7.2$ Hz, 1H), 7.99 (s, 1H), 7.35 (s, 1H), 6.83 (dd, $J=7.2, 1.9$ Hz, 1H), 3.88 (t, $J=6.8$ Hz, 2H), 2.98 (s, 2H), 2.89 (t, $J=6.8$ Hz, 2H), 2.66 (d, $J=7.3$ Hz, 2H), 2.53 (s, 2H), 2.15-2.00 (m, 2H), 1.84-1.56 (m, 4H), 1.51-1.39 (m, 2H), 1.16 (d, $J=6.7$ Hz, 3H). NH proton not observed due to solvent exchange.

Example 277. Preparation of 1-(5-(((2S,4R)-2-methyl-1-(oxetan-2-ylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



[1026] Prepared from 1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione by the method of Example 109 wherein oxetane-2-carbaldehyde was used in place of isobutyraldehyde. LCMS $[M+H]^+$: 412.0. 1H NMR (400 MHz, MeOD) δ 8.44 (s, 1H), 8.42 (s, 1H), 8.01 (s, 1H), 7.36 (s, 1H), 6.86-6.79 (m, 1H), 5.39-5.21 (m, 1H), 5.20-4.99 (m, 1H), 4.76-4.52 (m, 3H), 4.43-4.26 (m, 2H), 3.89 (t, $J=6.8$ Hz, 2H), 3.63 (s, 1H), 3.23-3.00 (m, 2H), 2.89 (t, $J=6.8$ Hz, 2H), 2.82-2.48 (m, 4H), 2.03 (d, $J=6.0$ Hz, 1H), 1.89-1.57 (m, 2H), 1.40-1.25 (m, 3H).

Example 278. Preparation of 1-(5-(((2S,4R)-1-(2,2-difluoro-3-methoxypropyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



Step 1: 1-(5-(((2S,4R)-1-(3-(((tert-butyl)diphenylsilyl)oxy)-2,2-difluoropropyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl) dihydropyrimidine-2,4(1H,3H)-dione

[1027] To a mixture of 3-(2,4-dimethoxybenzyl)-1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]

pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride (0.17 g, 0.32 mmol) and K_2CO_3 (0.133 g, 0.96 mmol) in dioxane (5 mL) was added 3-((tert-butyldiphenylsilyl)oxy)-2,2-difluoropropyl trifluoromethanesulfonate (0.156 g, 0.32 mmol). The reaction mixture was stirred at 90° C. for 16 h. The reaction mixture was cooled to rt and diluted with water and saturated aqueous $NaHCO_3$ solution. The mixture was extracted with DCM and the organic layer was dried over Na_2SO_4 , filtered, and concentrated to afford crude 1-(5-(((2S,4R)-1-(3-((tert-butyldiphenylsilyl)oxy)-2,2-difluoropropyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione (0.24 g) that was used without further purification. LCMS $[M+H]^+$: 824.6.

Step 2: 1-(5-(((2S,4R)-1-(2,2-difluoro-3-hydroxypropyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione

[1028] To a solution of 1-(5-(((2S,4R)-1-(3-((tert-butyldiphenylsilyl)oxy)-2,2-difluoropropyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione (0.24 g, 0.29 mmol) in THF (3 mL) was added TBAF (1.0 M in THF, 2 mL) and the reaction mixture was stirred at room temperature for 2 h. The reaction mixture was diluted with water and extracted with DCM. The organic layer was dried over Na_2SO_4 , filtered, and concentrated to afford crude 1-(5-(((2S,4R)-1-(2,2-difluoro-3-hydroxypropyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione (0.11 g, 0.15 mmol) as a brown solid. LCMS $[M+H]^+$: 586.1.


Step 3: 1-(5-(((2S,4R)-1-(2,2-difluoro-3-methoxypropyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione

[1029] To a stirred solution of 1-(5-(((2S,4R)-1-(2,2-difluoro-3-hydroxypropyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione (80 mg, 0.136 mmol) in DMF (2 mL) was added sodium hydride (10 mg, 0.41 mmol) at 0° C. After 5 min, methyl iodide (39 mg, 0.27 mmol) was added and the reaction mixture was stirred at room temperature for 10 min. The reaction mixture was diluted with water and extracted with EtOAc. The organic layer was washed with brine, dried over Na_2SO_4 , filtered, and concentrated. Silica gel column chromatography (eluting with 30% EtOAc in hexane) provided 1-(5-(((2S,4R)-1-(2,2-difluoro-3-methoxypropyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione (80 mg). LCMS $[M+H]^+$: 600.2.

[1030] Step 4: 1-(5-(((2S,4R)-1-(2,2-difluoro-3-methoxypropyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 278) was prepared from 1-(5-(((2S,4R)-1-(2,2-difluoro-3-methoxypropyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione by the method of Example 1, step 5, wherein 1-(5-(((2S,4R)-1-(2,2-difluoro-3-methoxypropyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione was used in place of 1-(5-(((2S,4R)-1-(2,2-difluoro-3-methoxypropyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione.

pyrimidine-2,4(1H,3H)-dione was used in place of 1-(5-(((2S,4R)-1-(2,2-difluoro-3-methoxypropyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione. LCMS $[M+H]^+$: 450.0. 1H NMR (400 MHz, MeOD) δ 8.42-8.36 (m, 1H), 8.24 (s, 1H), 7.98 (s, 1H), 7.33 (s, 1H), 6.81 (dd, $J=7.3, 1.9$ Hz, 1H), 3.88 (t, $J=6.8$ Hz, 2H), 3.65 (q, $J=12.9$ Hz, 2H), 3.41 (s, 3H), 2.95 (d, $J=15.0$ Hz, 1H), 2.89 (t, $J=6.8$ Hz, 2H), 2.85-2.68 (m, 3H), 2.60 (d, $J=7.2$ Hz, 2H), 2.02 (q, $J=7.7$ Hz, 2H), 1.61 (d, $J=12.3$ Hz, 2H), 1.54 (dd, $J=7.7, 4.0$ Hz, 2H), 1.04 (d, $J=6.7$ Hz, 3H).


Example 279. Preparation of 1-(5-(((2S,4R)-2-methyl-1-(oxetan-3-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[1031] To a solution of 1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride (180 mg, 0.41 mmol) in THF (10 mL) was added oxetan-3-one (85 mg, 1.2 mmol), dibutyltin dichloride (62 mg, 0.20 mmol), and triethylamine (0.2 mL, 1.2 mmol). The mixture was stirred at 80° C. for 1 h and then cooled to 0° C. and phenylsilane (45 mg, 0.41 mmol) was added. The reaction was stirred in a capped vial at 80° C. for 4 h. The reaction was cooled to rt, diluted with DCM and washed sequentially with water and brine. The organic layer was dried over Na_2SO_4 , filtered and concentrated. The crude material was purified by reverse phase HPLC using ACN/water/0.1% formic acid. The fractions containing the product were combined, frozen and lyophilized to afford 1-(5-(((2S,4R)-2-methyl-1-(oxetan-3-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione as white solid. LCMS $[M+H]^+$: 398.2. 1H NMR (400 MHz, MeOD) δ 8.44 (dd, $J=7.2, 0.9$ Hz, 1H), 8.01 (s, 1H), 7.36 (d, $J=1.8$ Hz, 1H), 6.83 (dd, $J=7.2, 1.9$ Hz, 1H), 4.84-4.74 (m, 3H), 4.52 (s, 1H), 3.89 (t, $J=6.8$ Hz, 2H), 3.81 (s, 1H), 3.24-3.02 (m, 2H), 2.89 (t,


J=6.8 Hz, 2H), 2.67 (d, J=9.1 Hz, 2H), 2.25 (d, J=19.4 Hz, 1H), 2.00-1.20 (m, 8H). NH proton not observed due to solvent exchange.

Example 280. Preparation of 1-(5-(((2S,4R)-1-cyclobutyl-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




[1032] Prepared using the method of Example 279 wherein cyclobutanone was used in place of oxetan-3-one. LCMS [M+H]⁺: 396.0. ¹H NMR (400 MHz, MeOD) δ 8.44 (s, 1H), 8.01 (s, 1H), 7.36 (s, 1H), 6.83 (dd, J=7.2, 1.9 Hz, 1H), 3.89 (t, J=6.8 Hz, 2H), 3.70 (s, 1H), 3.14 (s, 1H), 2.89 (t, J=6.8 Hz, 2H), 2.68 (d, J=7.0 Hz, 2H), 2.31 (s, 2H), 2.25-2.12 (m, 3H), 1.95-1.80 (m, 4H), 1.30 (s, 7H). NH proton not observed due to solvent exchange.

Example 281. Preparation of 1-(5-((1-(oxetan-3-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




[1033] Prepared using the method of Example 279 wherein 1-(5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione was used in place of 1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione. LCMS [M+H]⁺: 384.1. ¹H NMR (400 MHz, MeOD) δ 8.42 (dd, J=7.1, 0.9 Hz, 1H), 8.25 (s, 1H), 7.99 (s, 1H), 7.39-7.31 (m, 1H), 6.82 (dd, J=7.2, 1.9 Hz, 1H), 4.75 (t, J=7.1 Hz, 2H), 4.67 (t, J=6.6 Hz, 2H), 3.88 (t, J=6.8 Hz, 3H), 3.09 (d, J=11.7 Hz, 2H), 2.89 (t, J=6.8 Hz, 2H), 2.68 (d, J=6.8 Hz, 2H), 2.30 (t, J=12.1 Hz, 2H), 1.88-1.78 (m, 3H), 1.45 (td, J=14.3, 7.4 Hz, 2H).

Example 282. Preparation of 1-(5-((1-(cyclobutyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




[1034] Prepared using the method of Example 279 wherein 1-(5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione was used in place of 1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione and cyclobutanone was used in place of oxetan-3-one. LCMS [M+H]⁺: 382.0. ¹H NMR (400 MHz, MeOD) δ 8.43 (d, J=7.2 Hz, 1H), 8.01 (s, 1H), 7.37 (s, 1H), 6.83 (dd, J=7.2, 1.8 Hz, 1H), 3.88 (t, J=6.8 Hz, 2H), 3.59 (s, 1H), 3.41 (d, J=12.2 Hz, 2H), 3.35 (s, 2H), 2.89 (t, J=6.8 Hz, 2H), 2.78-2.65 (m, 3H), 2.36-2.26 (m, 2H), 2.17 (d, J=13.8 Hz, 2H), 2.04-1.79 (m, 4H), 1.49 (d, J=13.3 Hz, 2H). NH proton not observed due to solvent exchange.

Example 283. Preparation of 1-(5-(((2S,4R)-1-(ethylsulfonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



-continued



Step 1: 3-(2,4-dimethoxybenzyl)-1-(5-(((2S,4R)-1-(ethylsulfonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

[1035] Triethylamine (0.066 mL, 0.47 mmol) and ethylsulfonyl chloride (37 mg, 0.28 mmol) were added to a solution of 3-(2,4-dimethoxybenzyl)-1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione hydrochloride (50 mg, 0.095 mmol) in DCM (2 mL) at 0° C. The mixture was stirred at rt for 2 h, then diluted with DCM and washed with brine. The organic layer was dried over sodium sulfate,


filtered and concentrated to give crude 3-(2,4-dimethoxybenzyl)-1-(5-(((2S,4R)-1-(ethylsulfonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione, which was used without further purification. LCMS $[M+H]^+$: 584.4.

[1036] Step 2: 1-(5-(((2S,4R)-1-(ethylsulfonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione (Example 283) was prepared from 3-(2,4-dimethoxybenzyl)-1-(5-(((2S,4R)-1-(ethylsulfonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione by the method of Example 1, step 5, wherein 3-(2,4-dimethoxybenzyl)-1-(5-(((2S,4R)-1-(ethylsulfonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione was used in place of 1-(5-((1-(cyclohexylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl) dihydropyrimidine-2,4(1H,3H)-dione. LCMS $[M+H]^+$: 434.1. ^1H NMR (500 MHz, DMSO) δ 10.44 (s, 1H), 8.57 (dd, $J=7.1, 0.9$ Hz, 1H), 7.99 (s, 1H), 7.37 (dd, $J=1.9, 0.9$ Hz, 1H), 6.81 (dd, $J=7.2, 1.9$ Hz, 1H), 4.12-4.00 (m, 1H), 3.77 (t, $J=6.7$ Hz, 2H), 3.61-3.45 (m, 1H), 3.15-2.92 (m, 3H), 2.79 (t, $J=6.7$ Hz, 2H), 2.54 (d, $J=7.4$ Hz, 2H), 2.03 (dd, $J=7.6, 3.9$ Hz, 1H), 1.65-1.49 (m, 2H), 1.37 (td, $J=12.9, 5.4$ Hz, 1H), 1.25-1.15 (m, 6H), 1.11 (td, $J=12.6, 4.6$ Hz, 1H).


[1037] The compounds in the following table were prepared by the method of Example 283, using the appropriate commercially available sulfonyl chloride in step 1.

Example No	Structure	Mass $[M+H]^+$	^1H NMR
284	1-(5-(((2S,4R)-2-methyl-1-(methylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione	420.1	(500 MHz, DMSO- d_6) δ 10.44 (s, 1H), 8.57 (dd, $J = 7.1, 0.9$ Hz, 1H), 8.00 (s, 1H), 7.36 (dd, $J = 1.9, 0.9$ Hz, 1H), 6.81 (dd, $J = 7.1, 1.9$ Hz, 1H), 4.10 (t, $J = 6.6$ Hz, 1H), 3.77 (t, $J = 6.8$ Hz, 2H), 3.58-3.49 (m, 1H), 2.98 (td, $J = 13.2, 2.7$ Hz, 1H), 2.91 (s, 3H), 2.79 (t, $J = 6.7$ Hz, 2H), 2.57-2.53 (m, 2H), 2.10-1.94 (m, 1H), 1.60 (d, $J = 13.0$ Hz, 1H), 1.55-1.48 (m, 1H), 1.40 (td, $J = 12.9, 5.3$ Hz, 1H), 1.27-1.06 (m, 4H).
285	1-(5-(((2S,4R)-1-(isopropylsulfonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione	448.2	(500 MHz, DMSO- d_6) δ 10.44 (s, 1H), 8.57 (d, $J = 7.1$ Hz, 1H), 7.99 (s, 1H), 7.37 (s, 1H), 6.81 (dd, $J = 7.1, 1.8$ Hz, 1H), 4.12-3.99 (m, 1H), 3.77 (t, $J = 6.7$ Hz, 2H), 3.00 (ddp, $J = 21.1, 14.8, 6.7$ Hz, 2H), 2.79 (t, $J = 6.7$ Hz, 2H), 2.00 (d, $J = 22.1$ Hz, 2H), 1.71-1.47 (m, 4H), 1.37 (td, $J = 12.7, 5.3$ Hz, 2H), 1.26-1.07 (m, 5H), 0.98 (t, $J = 7.4$ Hz, 4H).

-continued

Example No	Structure	Mass [M + H] ⁺	¹ H NMR
286	 1-(5-(((2S,4R)-1-(cyclopropylsulfonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	446.1	(400 MHz, MeOD) δ 8.41 (dd, J = 7.2, 0.9 Hz, 1H), 7.99 (s, 1H), 7.35 (dd, J = 1.9, 0.9 Hz, 1H), 6.83 (dd, J = 7.2, 1.8 Hz, 1H), 4.20 (t, J = 7.0 Hz, 1H), 3.88 (t, J = 6.8 Hz, 2H), 3.70-3.60 (m, 1H), 3.13 (td, J = 13.3, 2.7 Hz, 1H), 2.89 (t, J = 6.8 Hz, 2H), 2.61 (d, J = 7.1 Hz, 2H), 2.53-2.37 (m, 1H), 2.21-2.05 (m, 1H), 1.75-1.59 (m, 2H), 1.51 (td, J = 12.8, 5.3 Hz, 1H), 1.33-1.23 (m, 4H), 1.07-0.96 (m, 4H). Missing NH due to solvent exchange.

Example 287. Preparation of 1-(5-(((2S,4R)-1-(cyclopropanecarbonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione






Step 1: 1-(5-(((2S,4R)-1-(cyclopropanecarbonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione


[1038] HATU (43 mg, 0.11 mmol) and cyclopropanecarboxylic acid (6.5 mg, 0.076 mmol) were added to a solution of 3-(2,4-dimethoxybenzyl)-1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride (40 mg, 0.076 mmol) in DMF (1 mL) at rt. The mixture was stirred at rt for 5 min and then DIPEA (0.040 mL, 0.23 mmol) was added. The mixture was stirred at rt for 12 h and then diluted with ethyl acetate and washed sequentially with brine. The organic layer was dried over sodium sulfate, filtered and concentrated to give crude 1-(5-(((2S,4R)-1-(cyclopropanecarbonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione, which was used without further purification. LCMS [M+H]⁺: 560.4.

[1039] Step 2: 1-(5-(((2S,4R)-1-(cyclopropanecarbonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 287) was prepared from 1-(5-(((2S,4R)-1-(cyclopropanecarbonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione using the method of Example 1, step 5, wherein 1-(5-(((2S,4R)-1-(cyclopropanecarbonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione was used in place of 1-(5-(((1-(cyclohexylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione. LCMS [M+H]⁺: 410.2. ¹H NMR (500 MHz, DMSO-d₆) δ 10.43 (s, 1H), 8.56 (d, J=7.1 Hz, 1H), 7.99 (s, 1H), 7.36 (s, 1H), 6.81 (dd, J=7.2, 1.8 Hz, 1H), 4.85-4.48 (m, 1H), 4.19 (dd, J=89.0, 13.8 Hz, 1H), 3.77 (t, J=6.7 Hz, 2H), 3.22-3.04 (m, 1H), 2.79 (t, J=6.7 Hz, 2H), 2.67-2.54 (m, 2H), 2.09 (d, J=11.9 Hz, 1H), 1.93 (ddt, J=16.7, 13.1, 5.8 Hz, 1H), 1.71-1.47 (m, 2H), 1.43-0.92 (m, 5H), 0.82-0.58 (m, 4H).

[1040] The compounds in the following table were prepared by the method of Example 287, using the appropriate commercially available carboxylic acid in step 1.

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
288	 1-(5-(((2S,4R)-1-isobutyryl-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	412.2	(500 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 8.55 (d, J = 7.1 Hz, 1H), 7.99 (s, 1H), 7.35 (s, 1H), 6.81 (dd, J = 7.1, 1.8 Hz, 1H), 4.79-4.73 (m, 1H), 4.37-4.30 (m, 1H), 4.28 (t, J = 6.4 Hz, 1H), 3.77 (d, J = 6.7 Hz, 2H), 3.06 (td, J = 13.5, 2.7 Hz, 1H), 2.80 (dt, J = 10.8, 6.7 Hz, 3H), 2.71-2.54 (m, 1H), 2.08 (d, J = 5.6 Hz, 1H), 1.72-1.47 (m, 2H), 1.34-1.18 (m, 1H), 1.17 (d, J = 6.8 Hz, 1H), 1.10-0.92 (m, 9H).
289	 1-(5-(((2S,4R)-1-(cyclobutanecarbonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	424.4	(500 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 8.55 (d, J = 7.1 Hz, 1H), 7.99 (s, 1H), 7.34 (s, 1H), 6.80 (dd, J = 7.1, 1.8 Hz, 1H), 4.77-4.64 (m, 1H), 4.28 (d, J = 13.6 Hz, 1H), 4.07-3.98 (m, 1H), 3.76 (t, J = 6.7 Hz, 2H), 3.28 (dt, J = 18.1, 8.7 Hz, 1H), 2.78 (t, J = 6.7 Hz, 2H), 2.69-2.57 (m, 1H), 2.24-1.96 (m, 5H), 1.88 (dq, J = 10.8, 8.8 Hz, 1H), 1.79-1.66 (m, 1H), 1.60 (d, J = 12.9 Hz, 1H), 1.51 (d, J = 12.2 Hz, 1H), 1.22 (dtd, J = 18.4, 12.8, 5.4 Hz, 1H), 1.12 (d, J = 6.8 Hz, 2H), 1.06-0.87 (m, 3H).
290	 1-(5-(((2S,4R)-2-methyl-1-(1-methylpiperidin-4-carbonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	467.1	(500 MHz, DMSO-d ₆) δ 10.43 (s, 1H), 8.56 (d, J = 7.1 Hz, 1H), 8.00 (s, 1H), 7.34 (s, 1H), 6.80 (dt, J = 7.3, 1.9 Hz, 1H), 4.92-4.55 (m, 1H), 4.31 (d, J = 13.2 Hz, 1H), 3.77 (t, J = 6.7 Hz, 2H), 3.18-3.04 (m, 1H), 3.05-2.90 (m, 2H), 2.84 (dt, J = 8.0, 3.9 Hz, 1H), 2.79 (t, J = 6.7 Hz, 2H), 2.75 (d, J = 4.5 Hz, 3H), 2.71-2.57 (m, 1H), 2.08 (s, 1H), 1.97-1.49 (m, 6H), 1.38-1.17 (m, 2H), 1.16-0.86 (m, 3H). 3 missing protons attributed to overlap with solvent peak.

Example 291. Preparation of 1-(5-(((2S,4R)-2-methyl-1-(pyrrolidin-1-ylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione






[1041] Triethylamine (89 mg, 0.87 mmol) and pyrrolidine-1-sulfonyl chloride (60 mg, 0.35 mmol) were added to a solution of 1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione (100 mg, 0.29 mmol) in DCM (4 mL) at rt. The mixture was stirred at rt for 16 h, then diluted with saturated aqueous NaHCO₃ solution. The mixture was extracted with DCM and the organic layer was washed with brine. The organic layer was dried over sodium sulfate, filtered and concentrated. The crude material was purified by reverse phase HPLC using ACN/water/0.1% formic acid. The fractions containing the product were combined, frozen and lyophilized to afford 1-(5-(((2S,4R)-2-methyl-1-(pyrrolidin-1-ylsulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione as white solid. LCMS [M+H]⁺: 475.2. ¹H NMR (400 MHz, MeOD) δ 8.41 (dd, J=7.2, 0.9 Hz, 1H), 7.98 (s, 1H), 7.37-7.32 (m, 1H), 6.83 (dd, J=7.2, 1.9 Hz, 1H), 4.10 (t, J=6.4 Hz, 1H), 3.88 (t, J=6.7 Hz, 2H), 3.52 (dt, J=13.2, 3.6 Hz, 1H), 3.24-3.18 (m, 4H), 3.05 (td, J=13.2, 2.7 Hz, 1H), 2.89 (t, J=6.8 Hz, 2H), 2.60 (d, J=7.1 Hz, 2H), 2.07 (ddd, J=11.9, 7.7, 4.0 Hz, 1H), 1.94-1.87 (m, 4H), 1.72-1.57 (m, 2H), 1.49 (td, J=12.7, 5.2 Hz, 1H), 1.23 (d, J=7.0 Hz, 4H). NH proton not observed due to solvent exchange.


[1042] The compounds in the following table were prepared by the method of Example 291, using the appropriate commercially available sulfamoyl chloride, carbamic chloride or isocyanate.

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
292		449.1	(400 MHz, MeOD) δ 8.41 (dd, J = 7.2, 0.9 Hz, 1H), 7.98 (s, 1H), 7.35 (dd, J = 1.9, 1.0 Hz, 1H), 6.83 (dd, J = 7.2, 1.9 Hz, 1H), 4.06 (t, J = 6.6 Hz, 1H), 3.88 (t, J = 6.8 Hz, 2H), 3.50 (d, J = 13.7 Hz, 1H), 3.06 (td, J = 13.2, 2.7 Hz, 1H), 2.89 (t, J = 6.8 Hz, 2H), 2.74 (s, 6H), 2.61 (d, J = 7.2 Hz, 2H), 2.13-2.00 (m, 1H), 1.72-1.45 (m, 3H), 1.27 (d, J = 4.6 Hz, 1H), 1.23 (d, J = 7.0 Hz, 3H). NH proton not observed due to solvent exchange.
	(2S,4R)-4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-N,N,2-trimethylpiperidine-1-sulfonamide		


-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
293	 (2S,4R)-N-cyclopentyl-4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperidine-1-sulfonamide	489.4	(400 MHz, MeOD) δ 8.41 (d, J = 7.2 Hz, 1H), 7.98 (s, 1H), 7.35 (s, 1H), 6.83 (dd, J = 7.2, 1.9 Hz, 1H), 4.10 (d, J = 6.4 Hz, 1H), 3.88 (t, J = 6.8 Hz, 2H), 3.55-3.46 (m, 2H), 3.05-2.95 (m, 1H), 2.89 (t, J = 6.8 Hz, 2H), 2.60 (d, J = 7.1 Hz, 2H), 2.06 (d, J = 17.9 Hz, 1H), 1.88 (t, J = 6.2 Hz, 2H), 1.74-1.47 (m, 9H), 1.27 (s, 1H), 1.23 (d, J = 6.9 Hz, 3H). Two NH protons not observed due to solvent exchange.
294	 (2S,4R)-4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-N,N,2-trimethylpiperidine-1-carboxamide	413.3	(400 MHz, MeOD) δ 8.41 (dd, J = 7.2, 0.9 Hz, 1H), 7.98 (s, 1H), 7.34 (dd, J = 1.9, 0.9 Hz, 1H), 6.83 (dd, J = 7.2, 1.9 Hz, 1H), 4.10-4.00 (m, 1H), 3.88 (t, J = 6.8 Hz, 2H), 3.44 (dt, J = 13.6, 3.7 Hz, 1H), 3.03 (td, J = 13.3, 2.7 Hz, 1H), 2.89 (t, J = 6.8 Hz, 2H), 2.80 (s, 6H), 2.59 (d, J = 7.1 Hz, 2H), 2.11 (dtd, J = 15.6, 8.1, 4.0 Hz, 1H), 1.66 (dt, J = 12.6, 3.1 Hz, 1H), 1.58 (ddt, J = 13.2, 4.0, 2.0 Hz, 1H), 1.43 (td, J = 12.8, 5.2 Hz, 1H), 1.28-1.20 (m, 1H), 1.18 (d, J = 7.0 Hz, 3H). NH proton not observed due to solvent exchange.
295	 1-(5-(((2S,4R)-2-methyl-1-(pyrrolidine-1-carbonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	439.3	(400 MHz, MeOD) δ 8.41 (dd, J = 7.2, 0.9 Hz, 1H), 7.98 (s, 1H), 7.34 (dd, J = 1.9, 1.0 Hz, 1H), 6.83 (dd, J = 7.2, 1.9 Hz, 1H), 4.17-4.09 (m, 1H), 3.88 (t, J = 6.8 Hz, 2H), 3.60-3.50 (m, 1H), 3.36-3.33 (m, 4H), 3.00 (td, J = 13.3, 2.7 Hz, 1H), 2.89 (t, J = 6.8 Hz, 2H), 2.59 (d, J = 7.2 Hz, 2H), 2.12 (ddd, J = 11.8, 7.9, 4.1 Hz, 1H), 1.84 (h, J = 3.4 Hz, 4H), 1.70-1.54 (m, 2H), 1.41 (td, J = 12.9, 5.3 Hz, 1H), 1.19 (d, J = 7.0 Hz, 4H). Missing NH due to solvent exchange.

-continued


Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
296	 (2S,4R)-N-cyclopentyl-4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperidine-1-carboxamide	453.3	(400 MHz, MeOD) δ 8.41 (dd, J = 7.2, 0.9 Hz, 1H), 7.98 (s, 1H), 7.40-7.25 (m, 1H), 6.83 (dd, J = 7.2, 1.9 Hz, 1H), 4.36 (t, J = 6.7 Hz, 1H), 4.01 (p, J = 7.2 Hz, 1H), 3.92-3.79 (m, 3H), 2.94-2.83 (m, 3H), 2.59 (d, J = 7.1 Hz, 2H), 2.15-2.03 (m, 1H), 1.90 (p, J = 6.3 Hz, 2H), 1.77-1.51 (m, 6H), 1.52-1.26 (m, 4H), 1.23-1.06 (m, 4H). Missing NH due to solvent exchange.

Example 297. Preparation of 1-(5-((1-(pyrrolidin-1-yl)sulfonyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




[1043] Prepared using the method of Example 291 wherein 1-(5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione was used in place of 1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione. LCMS [M+H]⁺: 461.3. ¹H NMR (400 MHz, MeOD) δ 8.40 (dd, J=7.2, 0.9 Hz, 1H), 7.98 (s, 1H), 7.36 (dd, J=1.9, 1.0 Hz, 1H), 6.82 (dd, J=7.2, 1.9 Hz, 1H), 3.88 (t, J=6.8 Hz, 2H), 3.66 (d, J=12.1 Hz, 2H), 3.29-3.24 (m, 3H), 2.89 (t, J=6.8 Hz, 2H), 2.77 (td, J=12.3, 2.4 Hz, 2H), 2.65 (d, J=7.0 Hz, 2H), 1.93-1.87 (m, 4H), 1.74 (d, J=13.9 Hz, 3H), 1.40-1.27 (m, 3H). NH proton not observed due to solvent exchange.

Example 298. Preparation of N-cyclopentyl-4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperidine-1-sulfonamide




[1044] Prepared using the method of Example 291 wherein cyclopentylsulfamoyl chloride was used in place of pyrrolidine-1-sulfonyl chloride and 1-(5-(piperidin-4-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione was used in place of 1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione. LCMS [M+H]⁺: 475.3. ¹H NMR (400 MHz, MeOD) δ 8.41 (dd, J=7.1, 0.9 Hz, 1H), 7.99 (s, 1H), 7.39-7.31 (m, 1H), 6.83 (dd, J=7.2, 1.9 Hz, 1H), 3.88 (t, J=6.7 Hz, 2H), 3.67-3.56 (m, 3H), 2.89 (t, J=6.7 Hz, 2H), 2.71-2.61 (m, 3H), 1.90 (q, J=5.5 Hz, 2H), 1.78-1.66 (m, 4H), 1.61-1.46 (m, 5H), 1.32 (dd, J=18.1, 6.0 Hz, 4H). NH proton not observed due to solvent exchange.

Example 299. Preparation of (2S,4R)-4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-N-ethyl-N,2-dimethylpiperidine-1-carboxamide



-continued




Step 1: (2S,4R)-4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-N-ethyl-N,2-dimethylpiperidine-1-carboxamide

[1045] Triethylamine (74 mg, 0.73 mmol) and triphosgene (213 mg, 0.73 mmol) were added to a solution of 3-(2,4-dimethoxybenzyl)-1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (120 mg, 0.24 mmol) in DCM (5 mL) at rt. The mixture was stirred at rt for 10 min, then N-methylethanamine (22 mg, 0.36 mmol) was added. The reaction was stirred at rt for 4 h then diluted with water and extracted with EtOAc. The organic layer was dried over sodium sulfate, filtered and concentrated. The crude product was purified by flash silica gel chromatography (eluted with 6.5% MeOH/DCM) to give (2S,4R)-4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-N-ethyl-N,2-dimethylpiperidine-1-carboxamide (0.10 g, 0.17 mmol, 71% yield) as an off-white solid. LCMS [M+H]⁺: 576.9.


[1046] Step 2: (2S,4R)-4-((3-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-N-ethyl-N,2-dimethylpiperidine-1-carboxamide (Example 299) was prepared from (2S,4R)-4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-N-ethyl-N,2-dimethylpiperidine-1-carboxamide by the method of Example 1, step 5, wherein (2S,4R)-4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-N-ethyl-N,2-dimethylpiperidine-1-carboxamide was used in place of 1-(5-((1-(cyclohexylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione. LCMS [M+H]⁺: 427.2. ¹H NMR (400 MHz, MeOD) δ 8.41 (dd, J=7.2, 0.9 Hz, 1H), 7.98 (s, 1H), 7.34 (d, J=1.6 Hz, 1H), 6.83 (dd, J=7.2, 1.9 Hz, 1H), 4.03 (t, J=6.5 Hz, 1H), 3.88 (t, J=6.8 Hz, 2H), 3.42 (d, J=13.8 Hz, 1H), 3.24-3.10 (m, 3H), 3.03 (td, J=13.2, 2.7 Hz, 1H), 2.89 (t, J=6.8 Hz, 2H), 2.79 (s, 3H), 2.59 (d, J=7.2 Hz, 2H), 2.11 (s, 1H), 1.62 (dd, J=29.5, 13.2 Hz, 2H), 1.49-1.39 (m, 1H), 1.17 (d, J=6.9 Hz, 3H), 1.13 (t, J=7.1 Hz, 3H). NH proton not observed due to solvent exchange.

Example 300. Preparation of 1-(5-(((1s,3s)-3-methoxycyclobutyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[1047] Prepared using the method of Example 112 wherein cis-3-methoxycyclobutane-1-carbaldehyde was used in place of 2-cyclohexyl-2,2-difluoroacetaldehyde in step 1. LCMS [M+H]⁺: 426.2. ¹H NMR (400 MHz, cd3od) δ 8.42 (d, J=7.2 Hz, 1H), 8.00 (s, 1H), 7.36 (s, 1H), 6.82 (dd, J=7.4, 1.8 Hz, 1H), 3.88 (t, J=6.8 Hz, 2H), 3.79 (p, J=7.3 Hz, 1H), 3.22 (d, J=4.2 Hz, 3H), 2.89 (t, J=6.8 Hz, 4H), 2.67 (d, J=6.7 Hz, 4H), 2.56-2.44 (m, 2H), 2.24-2.03 (m, 2H), 1.85 (d, J=14.0 Hz, 3H), 1.63 (dt, J=11.7, 9.2 Hz, 2H), 1.46 (d, J=13.0 Hz, 2H), 1.30 (s, 1H). NH proton not observed due to solvent exchange.

Example 301. Preparation of 1-(5-(((1r,3R,4S)-3,4-difluorocyclopentyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[1048] Prepared using the method of Example 112 wherein (1r,3R,4S)-3,4-difluorocyclopentane-1-carbaldehyde was used in place of 2-cyclohexyl-2,2-difluoroacetaldehyde in step 1. LCMS [M+H]⁺: 446.3. ¹H NMR (400 MHz, MeOD) δ 8.43 (dd, J=7.2, 0.9 Hz, 1H), 8.01 (s, 1H), 7.36 (dd, J=1.9, 0.9 Hz, 1H), 6.83 (dd, J=7.2, 1.9 Hz, 1H), 3.89 (t, J=6.8 Hz, 2H), 3.50-3.40 (m, 2H), 3.06 (d, J=7.2 Hz, 2H), 2.89 (t, J=6.8 Hz, 2H), 2.85-2.75 (m, 2H), 2.70 (d, J=6.8 Hz, 2H), 2.43 (dq, J=15.2, 7.9 Hz, 1H), 2.37-2.19 (m, 2H), 1.91 (d, J=14.4 Hz, 3H), 1.75 (ddq, J=26.1, 12.6, 6.0 Hz, 2H), 1.53 (q, J=13.1 Hz, 2H), 1.27 (d, J=24.2 Hz, 2H). NH proton not observed due to solvent exchange.

Example 302. Preparation of 1-(5-(((2S,4R)-1-(((1R,3S)-3-methoxycyclobutyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




[1049] Prepared using the method of Example 141 wherein trans-3-methoxycyclobutane-1-carbaldehyde was used in place of 4,4-difluorocyclohexane-1-carbaldehyde in step 7. LCMS [M+H]⁺: 440.3. ¹H NMR (400 MHz, cd3od) δ 8.43 (d, J=7.3 Hz, 1H), 8.01 (d, J=1.8 Hz, 1H), 7.37 (d, J=7.3 Hz, 1H), 6.87-6.78 (m, 1H), 4.05-3.65 (m, 4H), 3.63-3.36 (m, 1H), 3.26-3.07 (m, 6H), 2.89 (t, J=6.7 Hz, 2H), 2.80-2.47 (m, 4H), 2.19 (d, J=8.3 Hz, 3H), 1.95-1.65 (m, 4H), 1.38 (ddd, J=34.9, 6.8, 4.2 Hz, 4H). Missing NH due to solvent exchange.

Example 303. Preparation of 1-(5-(((2S,4R)-1-(((1S,3R)-3-methoxycyclobutyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




[1050] Prepared using the method of Example 141 wherein cis-3-methoxycyclobutane-1-carbaldehyde was used in place of 4,4-difluorocyclohexane-1-carbaldehyde in step 7. LCMS [M+H]⁺: 440.4. ¹H NMR (400 MHz, cd3od) δ 8.43 (dd, J=7.4, 1.8 Hz, 1H), 8.01 (d, J=1.6 Hz, 1H), 7.36 (s, 1H), 6.83 (dd, J=7.1, 1.9 Hz, 1H), 3.89 (td, J=6.8, 3.4 Hz, 2H), 3.75-3.67 (m, 1H), 3.26-3.21 (m, 4H), 3.19-3.06 (m, 3H), 2.89 (t, J=6.8 Hz, 2H), 2.66 (d, J=7.1 Hz, 2H), 2.60-2.47 (m, 2H), 2.30-2.12 (m, 3H), 1.87 (q, J=14.0 Hz, 2H), 1.70 (dddt, J=17.3, 12.0, 9.0, 3.6 Hz, 3H), 1.50 (dd, J=12.9, 5.0 Hz, 1H), 1.33 (dd, J=7.0, 3.5 Hz, 3H). NH proton not observed due to solvent exchange.

Example 304. Preparation of 1-(5-(((2S,4R)-1-(((1R,3R,4S)-3,4-difluorocyclopentyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




[1051] Prepared using the method of Example 141 wherein (1R,3R,4S)-3,4-difluorocyclopentane-1-carbaldehyde was used in place of 4,4-difluorocyclohexane-1-carbaldehyde in step 7. LCMS [M+H]⁺: 460.3. ¹H NMR (400 MHz, MeOD) δ 8.49 (s, 1H), 8.43 (dd, J=7.2, 0.9 Hz, 1H), 8.01 (s, 1H), 7.36 (dd, J=1.9, 0.9 Hz, 1H), 6.83 (dd, J=7.2, 1.9 Hz, 1H), 5.14-4.93 (m, 1H), 3.89 (t, J=6.8 Hz, 2H), 3.64 (s, 1H), 3.22-3.00 (m, 4H), 2.89 (t, J=6.8 Hz, 2H), 2.69 (d, J=7.2 Hz, 1H), 2.46-2.15 (m, 4H), 1.93-1.64 (m, 5H), 1.58 (s, 1H), 1.32 (q, J=6.3 Hz, 5H).

Example 305. Preparation of 1-(5-(((2S,4R)-2-methyl-1-(((1R,4S)-4-(trifluoromethyl)cyclohexyl)methyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




[1052] Prepared using the method of Example 141 wherein (1R,4R)-4-(trifluoromethyl)cyclohexane-1-carbaldehyde was used in place of 4,4-difluorocyclohexane-1-carbaldehyde in step 7. LCMS [M+H]⁺: 506.4. ¹H NMR (300 MHz, cd3od) δ 8.44 (d, J=7.1 Hz, 1H), 8.02 (s, 1H), 7.37 (s, 1H), 6.84 (d, J=6.9 Hz, 1H), 3.89 (t, J=6.7 Hz, 2H), 3.74 (s, 1H), 2.98 (s, 2H), 2.89 (t, J=6.7 Hz, 2H), 2.70 (d, J=7.1 Hz, 2H), 2.25-2.13 (m, 2H), 2.05-1.75 (m, 8H), 1.49-1.26 (m, 8H), 1.14 (d, J=12.7 Hz, 2H). NH proton not observed due to solvent exchange.

Example 306. Preparation of 1-(5-(((2S,4R)-1-(((R)-3,3-difluorocyclopentyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione





[1053] Prepared using the method of Example 141 wherein (R)-3,3-difluorocyclopentane-1-carbaldehyde was used in place of 4,4-difluorocyclohexane-1-carbaldehyde in step 7. LCMS $[M+H]^+$: 460.3. 1H NMR (400 MHz, MeOD) δ 8.44 (d, $J=7.1$ Hz, 1H), 8.01 (s, 1H), 7.36 (d, $J=2.3$ Hz, 1H), 6.84 (ddd, $J=7.1, 4.9, 1.9$ Hz, 1H), 3.89 (td, $J=6.8, 1.9$ Hz, 2H), 3.82 (s, 1H), 3.69-3.40 (m, 1H), 3.23-3.06 (m, 2H), 2.89 (t, $J=6.8$ Hz, 2H), 2.72 (dd, $J=42.6, 7.3$ Hz, 2H), 2.63-2.35 (m, 2H), 2.34-2.03 (m, 4H), 2.01-1.50 (m, 6H), 1.40 (dd, $J=41.6, 6.8$ Hz, 4H). NH proton not observed due to solvent exchange.

Example 307. Preparation of 1-(5-(((2S,4R)-1-(((S)-3,3-difluorocyclopentyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[1054] Prepared using the method of Example 141 wherein (S)-3,3-difluorocyclopentane-1-carbaldehyde was used in place of 4,4-difluorocyclohexane-1-carbaldehyde in step 7. LCMS $[M+H]^+$: 460.3. 1H NMR (400 MHz, MeOD) δ 8.43 (d, $J=7.2$ Hz, 1H), 8.01 (s, 1H), 7.36 (d, $J=2.1$ Hz, 1H), 6.88-6.80 (m, 1H), 3.89 (td, $J=6.8, 1.8$ Hz, 2H), 3.82 (s, 1H), 3.23-3.05 (m, 2H), 2.89 (t, $J=6.8$ Hz, 2H), 2.67 (d, $J=7.0$ Hz, 2H), 2.62-2.37 (m, 2H), 2.31-2.05 (m, 4H), 1.98-1.52 (m, 6H), 1.47-1.22 (m, 5H). NH proton not observed due to solvent exchange.

Examples 308 and 309. Preparation of 1-(5-(((2S)-1-((4,4-difluorocyclohexyl)methyl)-4-fluoro-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



Example 308



Example 309

Step 1. tert-butyl (2S)-4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-4-fluoro-2-methylpiperidine-1-carboxylate


[1055] To an oven-dried vial was added tert-butyl (2S)-4-(bromomethyl)-4-fluoro-2-methylpiperidine-1-carboxylate (0.270 g, 0.87 mmol), 1-(5-bromopyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione (0.250 g, 0.544 mmol), $\text{NiCl}_2(\text{DME})$ (5 mg, 0.027 mmol), pyridine-2,6-bis(carboximidamide) dihydrochloride (6 mg, 0.027 mmol), NaI (0.020 g, 0.14 mmol) and Zn (0.070 g, 1.1 mmol). The vial was sealed with a septum cap, evacuated and refilled with nitrogen 3 times. DMA (3 mL) was added and the reaction was stirred at 70° C. for 17 h. The reaction was cooled to rt, diluted with EtOAc and filtered through a plug of silica gel, eluting with EtOAc. The eluent was concentrated and the residue was purified by silica gel column chromatography (eluted with 50-70% EtOAc in heptane) to give tert-butyl (2S)-4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-4-fluoro-2-methylpiperidine-1-carboxylate as a mixture of diastereomers. The diastereomers were separated by chiral HPLC purification: Column: Lux Cellulose-4 250×21.2 mm I.D., 5 μm Mobile phase: Phase A for n-hexane, and Phase B for 1:1 EtOH:MeOH (0.1% HCOOH); Isocratic elution: 50% (A):50% (B) Flow rate: 15 mL/min; Peak 1 (145 mg), LCMS $[\text{M}+\text{H}-\text{Boc}]^+$: 510.3, HPLC rt =17.04 min; Peak 2 (245 mg), LCMS $[\text{M}+\text{H}-\text{Boc}]^+$: 510.3, HPLC rt =17.64 min.

[1056] Step 2: 1-(5-(((2S)-1-((4,4-difluorocyclohexyl)methyl)-4-fluoro-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 308) was prepared from tert-butyl (2S)-4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-4-fluoro-2-methylpiperidine-1-carboxylate (chiral peak 1) using the method of Example 141, steps 6-8 wherein tert-butyl (2S)-4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-4-fluoro-2-methylpiperidine-1-carboxylate (chiral peak 1) was used in place of tert-butyl (2S,4R)-4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperidine-1-carboxylate in step 6. LCMS $[\text{M}+\text{H}]^+$: 492.1. ^1H NMR (400 MHz, MeOD) δ 8.44 (dd, J =7.2, 0.9 Hz, 1H), 8.33 (s, 1H), 8.03 (s, 1H), 7.42 (s, 1H), 6.87 (d, J =7.2 Hz, 1H), 3.90 (t, J =6.8 Hz, 2H), 3.18-3.06 (m, 2H), 2.98 (s, 2H), 2.89 (t, J =6.8 Hz, 2H), 2.81 (s, 1H), 2.19-1.72 (m, 12H), 1.45-1.25 (m, 6H).

[1057] 1-(5-(((2S)-1-((4,4-difluorocyclohexyl)methyl)-4-fluoro-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 309) was prepared from tert-butyl (2S)-4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-4-fluoro-2-methylpiperidine-1-carboxylate (chiral peak 2) using the method of Example 141, steps 6-8 wherein tert-butyl (2S)-4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-4-fluoro-2-methylpiperidine-1-carboxylate (chiral peak 2) was used in place of tert-butyl (2S,4R)-4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperidine-1-carboxylate in step 6. LCMS $[\text{M}+\text{H}]^+$: 492.1. ^1H NMR (400 MHz, MeOD) δ 8.44 (d, J =7.2 Hz, 1H), 8.33 (s, 1H), 8.03 (s, 1H), 7.43 (s, 1H),

6.87 (d, J =7.2 Hz, 1H), 3.90 (t, J =6.7 Hz, 2H), 3.15-2.96 (m, 4H), 2.89 (t, J =6.8 Hz, 2H), 2.69 (s, 1H), 2.13-1.72 (m, 10H), 1.29 (t, J =5.0 Hz, 8H).

Example 310. Preparation of 1-(5-(((2S)-4-fluoro-2-methyl-1-(oxetan-3-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




Step 1. 1-(5-(((2S)-4-fluoro-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

[1058] TFA (2 mL) was added to tert-butyl (2S)-4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-4-fluoro-2-methylpiperidine-1-carboxylate (chiral peak 1 from Example 308, step 1) (350 mg, 0.574 mmol). The mixture was heated in a sealed vial for 24 h at 90° C. The mixture was then cooled to rt and, concentrated and azeotropically dried with toluene to provide crude 1-(5-(((2S)-4-fluoro-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione which was used without further purification. LCMS $[\text{M}+\text{H}]^+$: 360.0.

[1059] Step 2: 1-(5-(((2S)-4-fluoro-2-methyl-1-(oxetan-3-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 310) was prepared from 1-(5-(((2S)-4-fluoro-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione by the method of Example 279, wherein 1-(5-(((2S)-4-fluoro-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-di-




one was used in place of 1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride. LCMS $[M+H]^+$: 416.0. 1H NMR (400 MHz, MeOD) δ 8.41 (dd, $J=7.2, 0.9$ Hz, 1H), 8.00 (s, 1H), 7.40 (d, $J=1.7$ Hz, 1H), 6.90-6.79 (m, 1H), 4.71-4.60 (m, 4H), 3.96 (q, $J=7.0$ Hz, 1H), 3.89 (t, $J=6.8$ Hz, 2H), 3.06 (d, $J=24.6$ Hz, 2H), 2.89 (t, $J=6.8$ Hz, 2H), 2.79 (ddt, $J=17.7, 12.0, 5.9$ Hz, 2H), 2.37 (dt, $J=11.9, 5.7$ Hz, 1H), 1.95-1.79 (m, 3H), 1.72 (td, $J=13.1, 6.6$ Hz, 1H), 0.98 (dd, $J=6.7, 1.7$ Hz, 3H). NH proton not observed due to solvent exchange.

Example 311. Preparation of 1-(5-(((2S)-4-fluoro-2-methyl-1-(oxetan-3-yl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[1060] Prepared using the method of Example 310 wherein tert-butyl (2S)-4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-4-fluoro-2-methylpiperidine-1-carboxylate (chiral peak 2 from Example 308, step 1) was used in place of tert-butyl (2S)-4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-4-fluoro-2-methylpiperidine-1-carboxylate (chiral peak 1 from Example 308, step 1). LCMS $[M+H]^+$: 416.0. 1H NMR (400 MHz, MeOD) δ 8.41 (dd, $J=7.2, 0.9$ Hz, 1H), 8.00 (s, 1H), 7.42 (s, 1H), 6.87 (d, $J=7.2$ Hz, 1H), 4.71-4.54 (m, 4H), 3.89 (t, $J=6.8$ Hz, 2H), 2.99 (d, $J=22.4$ Hz, 2H), 2.89 (t, $J=6.8$ Hz, 2H), 2.58 (d, $J=11.9$ Hz, 1H), 2.40 (s, 1H), 2.20-2.10 (m, 1H), 1.85-1.65 (m, 3H), 1.64-1.37 (m, 2H), 0.87 (d, $J=6.4$ Hz, 3H). NH proton not observed due to solvent exchange.

Example 312. Preparation of (cis)-1-(5-((1-isobutyl-2-(trifluoromethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




Step 1. (cis)-3-(2,4-dimethoxybenzyl)-1-(5-((1-isobutyl-2-(trifluoromethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione

[1061] To an oven-dried vial was added (cis)-4-(bromomethyl)-1-isobutyl-2-(trifluoromethyl)piperidine (14 mg, 0.049 mmol), 1-(5-bromopyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione (17.5 mg, 0.038 mmol), $NiCl_2(DME)$ (1 mg, 0.001 mmol), pyridine-2,6-bis(carboximidamide) dihydrochloride (1 mg, 0.001 mmol), NaI (1 mg, 0.009 mmol) and Zn (4 mg, 0.076 mmol). The vial was sealed with a septum cap, evacuated and refilled with nitrogen 3 times. DMA (2 mL) was added and the reaction was stirred at 100°C for 17 h. The reaction was cooled to rt, diluted with EtOAc and filtered through a plug of silica gel, eluting with EtOAc. The eluent was concentrated to give crude (cis)-3-(2,4-dimethoxybenzyl)-1-(5-((1-isobutyl-2-(trifluoromethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione which was used without further purification. LCMS $[M+H]^+$: 602.2.


[1062] Step 2: (cis)-1-(5-((1-isobutyl-2-(trifluoromethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 312) was prepared from (cis)-3-(2,4-dimethoxybenzyl)-1-(5-((1-isobutyl-2-(trifluoromethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione using the method of Example 1, step 5 wherein (cis)-3-(2,4-dimethoxybenzyl)-1-(5-((1-isobutyl-2-(trifluoromethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione was used in place of 1-(5-((1-(cyclohexylmethyl)piperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione. LCMS $[M+H]^+$: 452.4. 1H NMR (400 MHz, MeOD) δ 8.40 (dd, $J=7.2, 0.9$ Hz, 1H), 7.98 (s, 1H), 7.37 (dd, $J=1.9, 1.0$ Hz, 1H), 6.84 (dd, $J=7.2, 1.9$ Hz, 1H), 3.87 (t, $J=6.8$ Hz, 2H), 3.27 (d, $J=8.1$ Hz, 1H), 3.24-3.12 (m, 1H), 2.89 (t, $J=6.8$ Hz, 2H), 2.72 (dt, $J=9.1, 6.4$ Hz, 2H), 2.46 (d, $J=7.4$ Hz, 2H), 2.25 (dq, $J=11.6, 5.6$ Hz, 1H), 2.10-2.00 (m, 2H), 1.73 (dddd, $J=13.3, 10.5, 7.9, 3.2$ Hz, 4H), 0.91 (d, $J=6.5$ Hz, 3H), 0.86 (d, $J=6.7$ Hz, 3H).

Example 313. Preparation of (R)-1-(5-((octahydro-2H-pyrido[1,2-a]pyrazin-2-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[1063] Prepared using the method of Example 156, steps 1 and 4, wherein potassium (R)-trifluoro((octahydro-2H-pyrido[1,2-a]pyrazin-2-yl)methyl)borate was used in place of potassium {[4-(tert-butoxycarbonyl)-1-piperazinyl]methyl}(trifluoro)borate. LCMS $[M+H]^+$: 383.0. 1H NMR (400 MHz, MeOD) δ 8.49 (d, $J=7.1$ Hz, 1H), 8.04 (s, 1H), 7.55 (s, 1H), 6.99 (dd, $J=7.2, 1.8$ Hz, 1H), 3.90 (t, $J=6.7$ Hz, 2H), 3.79 (s, 2H), 3.51-3.41 (m, 2H), 3.28-3.12 (m, 3H), 3.04 (td, $J=12.9, 3.2$ Hz, 1H), 2.89 (t, $J=6.8$ Hz, 2H), 2.63 (s, 1H), 2.37 (s, 1H), 1.94 (dd, $J=30.0, 13.6$ Hz, 3H), 1.77 (q, $J=13.4$ Hz, 1H), 1.69-1.43 (m, 2H), 1.37-1.27 (m, 1H). NH proton not observed due to solvent exchange.

Example 314. Preparation of (R)-1-(5-((4-oxohexahydropyrazino[2,1-c][1,4]oxazin-8(1H)-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[1064] Prepared using the method of Example 156, steps 1 and 4, wherein potassium (R)-trifluoro((4-oxohexahydropyrazino[2,1-c][1,4]oxazin-8(1H)-yl)methyl)borate was used in place of potassium {[4-(tert-butoxycarbonyl)-1-piperazinyl]methyl}(trifluoro)borate. LCMS $[M+H]^+$: 399.0. 1H NMR (400 MHz, MeOD) δ 8.45 (dd, $J=7.2, 0.9$ Hz, 1H), 8.01 (s, 1H), 7.51 (dd, $J=1.9, 1.0$ Hz, 1H), 7.01 (dd, $J=7.2, 1.8$ Hz, 1H), 4.49 (ddd, $J=13.3, 3.3, 1.8$ Hz, 1H), 4.10 (d, $J=1.7$ Hz, 2H), 3.98 (dd, $J=12.0, 4.6$ Hz, 1H), 3.90 (t, $J=6.8$ Hz, 2H), 3.74-3.51 (m, 4H), 2.98-2.82 (m, 5H), 2.14 (td, $J=11.6, 3.2$ Hz, 1H), 2.06-1.94 (m, 1H). NH proton not observed due to solvent exchange.

Example 315. Preparation of (S)-1-(5-((1,1-dioxido-hexahydro-5H-isothiazolo[2,3-a]pyrazin-5-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[1065] Prepared using the method of Example 156, steps 1 and 4, wherein potassium (S)-((1,1-dioxido-hexahydro-5H-isothiazolo[2,3-a]pyrazin-5-yl)methyl)trifluoroborate was used in place of potassium {[4-(tert-butoxycarbonyl)-1-piperazinyl]methyl}(trifluoro)borate. LCMS $[M+H]^+$: 419.0. 1H NMR (400 MHz, MeOD) δ 8.45 (d, $J=7.1$ Hz, 1H), 8.01 (s, 1H), 7.50 (s, 1H), 7.00 (dd, $J=7.2, 1.8$ Hz, 1H), 3.89 (t, $J=6.8$ Hz, 2H), 3.65 (q, $J=13.7$ Hz, 2H), 3.26-3.10 (m, 3H), 3.04 (d, $J=11.3$ Hz, 1H), 2.96 (d, $J=11.5$ Hz, 1H), 2.92-2.86 (m, 3H), 2.40-2.14 (m, 2H), 2.09-1.90 (m, 3H). NH proton not observed due to solvent exchange.


Example 316. Preparation of 1-(5-(((S)-4-(((1R,3S)-3-methoxycyclobutyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[1066] Prepared using the method of Example 192 wherein trans-3-methoxycyclobutane-1-carbaldehyde was used in place of cyclohexanecarbaldehyde in step 3. LCMS $[M+H]^+$: 441.1. 1H NMR (400 MHz, cd_3od) δ 8.45 (d, $J=7.2$ Hz, 1H), 8.01 (s, 1H), 7.48 (s, 1H), 6.97 (dd, $J=7.1, 1.9$ Hz, 1H), 4.03-3.94 (m, 1H), 3.88 (t, $J=6.8$ Hz, 2H), 3.69-3.56


(m, 2H), 3.40 (d, J=12.6 Hz, 3H), 3.22 (d, J=4.3 Hz, 3H), 3.10 (d, J=11.6 Hz, 2H), 3.02 (s, 1H), 2.87 (t, J=6.7 Hz, 2H), 2.72 (d, J=8.3 Hz, 2H), 2.52 (s, 2H), 2.16 (ddt, J=36.3, 12.7, 7.4 Hz, 4H), 1.34 (d, J=6.6 Hz, 3H). NH proton not observed due to solvent exchange.

Example 317. Preparation of 1-(5-(((S)-4-(((1s,3R)-3-methoxycyclobutyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




[1067] Prepared using the method of Example 192 wherein cis-3-methoxycyclobutane-1-carbaldehyde was used in place of cyclohexanecarbaldehyde in step 3. LCMS [M+H]⁺: 441.3. ¹H NMR (400 MHz, cd3od) δ 8.46 (d, J=7.2 Hz, 1H), 8.02 (s, 1H), 7.50 (s, 1H), 6.99 (dd, J=7.1, 1.9 Hz, 1H), 3.89 (t, J=6.8 Hz, 2H), 3.81 (p, J=7.2 Hz, 1H), 3.70-3.56 (m, 2H), 3.22 (s, 6H), 3.06-2.84 (m, 6H), 2.60-2.46 (m, 3H), 2.36 (s, 1H), 2.22 (qd, J=9.1, 6.6 Hz, 1H), 1.75-1.60 (m, 2H), 1.30 (d, J=6.3 Hz, 3H). NH proton not observed due to solvent exchange.

Example 318. Preparation of 1-(5-(((S)-4-(((1r,3R,4S)-3,4-difluorocyclopentyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




[1068] Prepared using the method of Example 192 wherein (1r,3R,4S)-3,4-difluorocyclopentane-1-carbaldehyde was used in place of cyclohexanecarbaldehyde in step 3. LCMS [M+H]⁺: 461.4. ¹H NMR (400 MHz, MeOD) δ 8.46 (dd, J=7.2, 0.9 Hz, 1H), 8.02 (s, 1H), 7.50 (d, J=1.8 Hz, 1H), 6.99 (dd, J=7.2, 1.8 Hz, 1H), 5.08-4.92 (m, 1H), 4.81 (s, 1H), 3.89 (t, J=6.8 Hz, 2H), 3.68-3.54 (m, 2H), 3.20-3.11 (m, 1H), 2.99 (t, J=11.1 Hz, 1H), 2.93-2.75 (m, 5H), 2.53 (d, J=75.2 Hz, 3H), 2.39-2.06 (m, 4H), 1.83-1.56 (m, 2H), 1.17 (d, J=6.1 Hz, 3H). NH proton not observed due to solvent exchange.

Example 319. Preparation of 1-(5-(((S)-4-(((R)-3,3-difluorocyclopentyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




[1069] Prepared using the method of Example 192 wherein (R)-3,3-difluorocyclopentane-1-carbaldehyde was used in place of cyclohexanecarbaldehyde in step 3. LCMS [M+H]⁺: 461.2. ¹H NMR (300 MHz, CD₃OD) δ 8.45 (d, J=7.2 Hz, 1H), 8.01 (s, 1H), 7.50 (s, 1H), 7.05-6.94 (m, 1H), 3.89 (t, J=6.7 Hz, 2H), 3.56 (d, J=3.6 Hz, 2H), 2.89 (t, J=6.7 Hz, 3H), 2.74 (t, J=12.3 Hz, 3H), 2.61-1.91 (m, 11H), 1.38-1.25 (m, 2H), 1.06 (d, J=6.1 Hz, 2H). NH proton not observed due to solvent exchange.

Example 320. Preparation of 1-(5-(((S)-4-(((S)-3,3-difluorocyclopentyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




[1070] Prepared using the method of Example 192 wherein (S)-3,3-difluorocyclopentane-1-carbaldehyde was used in place of cyclohexanecarbaldehyde in step 3. LCMS [M+H]⁺: 461.1. ¹H NMR (400 MHz, MeOD) δ 8.44 (d, J=7.1 Hz, 1H), 8.01 (s, 1H), 7.49 (s, 1H), 6.99 (dd, J=7.2, 1.8 Hz, 1H), 3.89 (t, J=6.8 Hz, 2H), 3.54 (d, J=6.3 Hz, 1H), 2.89 (t, J=6.8 Hz, 3H), 2.79-2.63 (m, 3H), 2.50-1.90 (m, 12H), 1.32 (d, J=16.3 Hz, 1H), 1.04 (dd, J=6.1, 4.1 Hz, 3H). NH proton not observed due to solvent exchange.

Example 321. Preparation of (S)-1-(5-((4-(cyclopropylmethyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione




[1071] Prepared using the method of Example 192 wherein cyclopropanecarbaldehyde was used in place of cyclohexanecarbaldehyde in step 3. LCMS $[M+H]^+$: 397.4. 1H NMR (500 MHz, DMSO) δ 10.47 (s, 1H), 8.64 (d, $J=7.2$ Hz, 1H), 8.05 (s, 1H), 7.52 (s, 1H), 6.91 (dd, $J=7.2, 1.8$ Hz, 1H), 3.79 (t, $J=6.7$ Hz, 2H), 3.68 (s, 4H), 3.36 (s, 1H), 3.25-3.12 (m, 2H), 3.04 (d, $J=14.6$ Hz, 3H), 2.80 (t, $J=6.7$ Hz, 2H), 2.31 (d, $J=11.9$ Hz, 1H), 1.25 (d, $J=6.4$ Hz, 3H), 1.05 (s, 1H), 0.65 (s, 2H), 0.39 (d, $J=29.5$ Hz, 2H).

Example 322. Preparation of (S)-1-(5-((3-methyl-4-((1-methylcyclobutyl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4 (1H,3H)-dione




[1072] Prepared using the method of Example 192 wherein 1-methylcyclobutane-1-carbaldehyde was used in place of cyclohexanecarbaldehyde in step 3. LCMS $[M+H]^+$: 425.2. 1H NMR (500 MHz, DMSO) δ 10.47 (s, 1H), 8.64 (d, $J=7.2$ Hz, 1H), 8.05 (s, 1H), 7.51 (s, 1H), 6.91 (d, $J=7.2$ Hz, 1H), 3.79 (t, $J=6.7$ Hz, 6H), 2.96 (s, 6H), 2.80 (t, $J=6.7$ Hz, 3H), 2.10 (d, $J=13.4$ Hz, 1H), 2.01 (d, $J=9.1$ Hz, 1H), 1.88 (s, 1H), 1.84-1.71 (m, 2H), 1.67 (s, 1H), 1.27 (s, 6H).

Example 323. Preparation of 1-(5-(((S)-3-methyl-4-(((1r,4S)-4-(trifluoromethyl)cyclohexyl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



-continued




Step 1. (S)-1-(5-((3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4 (1H,3H)-dione

[1073] TFA (5 mL) was added to tert-butyl (S)-4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperazine-1-carboxylate (0.24 g, 0.40 mmol). The mixture was heated in a sealed vial at 70° C. for 2 h. The mixture was then cooled to rt and, concentrated and azeotropically dried with toluene to provide crude product (0.3 g) which was used without further purification. LCMS $[M+H]^+$: 343.9.

Step 2. 1-(5-(((S)-3-methyl-4-(((1r,4S)-4-(trifluoromethyl)cyclohexyl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4 (1H,3H)-dione


[1074] (1r,4r)-4-(trifluoromethyl)cyclohexane-1-carbaldehyde (0.126 g, 0.70 mmol) and triethylamine (0.14 mL, 1.05 mmol) were added to a solution of (S)-1-(5-((3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione (0.12 g, 0.35 mmol) in DCM (4 mL). The reaction mixture was stirred at rt for 90 min and then sodium triacetoxyborohydride (0.148 g, 0.70 mmol) was added. The reaction mixture was stirred at rt for 3 h and then quenched with a solution of saturated aqueous $NaHCO_3$ and extracted three times with DCM. The combined organic extracts were washed with brine, dried over Na_2SO_4 , filtered and concentrated. The residue was dissolved in DMSO, filtered through a 1 micron filter and purified by reverse phase HPLC using ACN/Water/0.1% formic acid. The fractions containing the product were combined, frozen and lyophilized to afford a formate salt of 1-(5-(((S)-3-methyl-4-(((1r,4S)-4-(trifluoromethyl)cyclohexyl)methyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione. LCMS $[M+H]^+$: 507.3. 1H NMR (400 MHz, cd_3od) δ 8.46 (d, $J=7.3$ Hz, 1H), 8.03 (s, 1H), 7.51 (s, 1H), 6.99 (dd, $J=7.1, 1.9$ Hz, 1H), 3.90 (t, $J=6.7$ Hz, 2H), 3.70-3.59 (m, 2H), 3.38 (s, 1H), 3.21 (q, $J=7.4$ Hz, 2H), 3.04-2.95 (m, 1H), 2.89 (t, $J=6.7$ Hz, 3H), 2.60 (s, 2H), 2.37 (s, 1H), 2.15 (dd, $J=23.1, 8.1$ Hz, 1H), 1.98 (d, $J=12.3$ Hz, 2H), 1.86 (d, $J=13.1$ Hz, 1H), 1.72 (s, 1H), 1.43-1.23 (m, 7H), 1.10 (dt, $J=21.2, 11.5$ Hz, 2H). NH proton not observed due to solvent exchange.

Example 324. Preparation of 1-(5-(((3S)-3-methyl-4-(oxetan-2-ylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[1075] Prepared using the method of Example 323 wherein oxetane-2-carbaldehyde was used in place of (1*r*,4*r*)-4-(trifluoromethyl)cyclohexane-1-carbaldehyde in step 2. LCMS $[M+H]^+$: 413.0. 1H NMR (400 MHz, MeOD) δ 8.47 (dd, $J=7.2, 0.9$ Hz, 1H), 8.03 (s, 1H), 7.50 (s, 1H), 6.98 (dd, $J=7.2, 1.9$ Hz, 1H), 5.18 (q, $J=9.2$ Hz, 1H), 4.76-4.56 (m, 2H), 3.90 (t, $J=6.8$ Hz, 2H), 3.81-3.61 (m, 3H), 3.57-3.37 (m, 3H), 3.24-3.10 (m, 2H), 3.08-2.78 (m, 4H), 2.66-2.29 (m, 3H), 1.41-1.26 (m, 3H). NH proton not observed due to solvent exchange.

Example 325. Preparation of (S)-1-(5-(((4-((2-oxaspiro[3.3]heptan-6-yl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[1076] Prepared using the method of Example 323 wherein 2-oxaspiro[3.3]heptane-6-carbaldehyde was used in place of (1*r*,4*r*)-4-(trifluoromethyl)cyclohexane-1-carbaldehyde in step 2. LCMS $[M+H]^+$: 453.2. 1H NMR (300 MHz, cd_3od) δ 8.47 (s, 1H), 8.45 (s, 1H), 8.03 (s, 1H), 7.49 (s, 1H), 6.98 (dd, $J=7.2, 1.9$ Hz, 1H), 4.75 (s, 2H), 4.57 (s, 2H), 3.89 (t, $J=6.7$ Hz, 2H), 3.67-3.56 (m, 2H), 3.21 (d, $J=24.5$ Hz, 2H), 2.89 (t, $J=6.5$ Hz, 6H), 2.56-2.41 (m, 4H), 2.38-2.21 (m, 1H), 2.04 (d, $J=3.8$ Hz, 2H), 1.76 (td, $J=10.7, 5.4$ Hz, 1H), 1.27 (d, $J=6.5$ Hz, 3H).

Example 326. Preparation of (S)-1-(5-(((4-((6,6-difluorospiro[3.3]heptan-2-yl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[1077] Prepared using the method of Example 323 wherein 6,6-difluorospiro[3.3]heptane-2-carbaldehyde was used in place of (1*r*,4*r*)-4-(trifluoromethyl)cyclohexane-1-carbaldehyde in step 2. LCMS $[M+H]^+$: 487.3. 1H NMR (400 MHz, MeOD) δ 8.45 (dd, $J=7.2, 1.0$ Hz, 1H), 8.02 (d, $J=1.5$ Hz, 1H), 7.49 (d, $J=1.9$ Hz, 1H), 6.98 (dt, $J=7.3, 1.9$ Hz, 1H), 3.89 (t, $J=6.8$ Hz, 2H), 3.68 (dd, $J=5.5, 4.0$ Hz, 1H), 3.63-3.52 (m, 2H), 3.22-3.11 (m, 2H), 2.93-2.75 (m, 6H), 2.75-2.55 (m, 3H), 2.55-2.37 (m, 3H), 2.35-1.93 (m, 5H), 1.33-1.17 (m, 3H). NH proton not observed due to solvent exchange.

Example 327. Preparation of (S)-1-(5-(((4-(2,2-difluoroethyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[1078] Prepared from (S)-3-(2,4-dimethoxybenzyl)-1-(5-(((3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione trifluoroacetate by the method of Example 156, steps 3-4 using 2,2-difluoroethyl trifluoromethanesulfonate in step 3. LCMS $[M+H]^+$: 407.0. 1H NMR (400 MHz, MeOD) δ 8.47 (dd, $J=7.2, 0.9$ Hz, 1H), 8.02 (s, 1H), 7.53 (dd, $J=1.9, 1.0$ Hz, 1H), 7.00 (dd, $J=7.2, 1.8$ Hz, 1H), 5.93 (tt, $J=56.0, 4.2$ Hz, 1H), 3.90 (t, $J=6.8$ Hz, 2H), 3.72-3.62 (m, 2H), 3.15-2.95 (m, 2H), 2.89 (t, $J=6.7$ Hz, 2H), 2.82 (ddd, $J=9.4, 6.8, 3.7$ Hz, 2H), 2.76-2.58 (m, 3H), 2.44 (td, $J=10.9, 2.8$ Hz, 1H), 2.15 (dd, $J=12.0, 8.8$ Hz, 1H), 1.07 (d, $J=6.2$ Hz, 3H). NH proton not observed due to solvent exchange.

Example 328. Preparation of (S)-1-(5-(((4-(2,2-difluoro-3-methoxypropyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione





-continued




[1079] Prepared by the method of Example 278 wherein (S)-3-(2,4-dimethoxybenzyl)-1-(5-(((3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride was used in place of 3-(2,4-dimethoxybenzyl)-1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride in step 1. LCMS $[M+H]^+$: 451.1. 1H NMR (300 MHz, METHANOL- d_4) δ ppm 8.46 (d, $J=7.3$ Hz, 1H), 8.01 (s, 1H), 7.51 (s, 1H), 6.99 (br d, $J=6.9$ Hz, 1H), 3.89 (t, $J=6.8$ Hz, 2H), 3.73-3.53 (m, 3H), 3.39 (s, 3H), 3.18-2.99 (m, 2H), 2.88 (t, $J=6.76$ Hz, 2H), 2.38-2.80 (m, 7H), 2.09-2.24 (m, 1H), 1.05 (d, $J=6.3$ Hz, 3H). NH proton not observed due to solvent exchange.

Example 329. Preparation of (S)-1-(5-((4-cyclobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



-continued






Step 1. (S)-1-(5-((4-cyclobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione




[1080] To a solution of (S)-3-(2,4-dimethoxybenzyl)-1-(5-(((3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione hydrochloride (50 mg, 0.094 mmol) in THF (2 mL) was added cyclobutanone (20 mg, 0.28 mmol), dibutyltin dichloride (86 mg, 0.28 mmol), and triethylamine (48 mg, 0.47 mmol). The mixture was stirred at rt for 1 h and then phenylsilane (20 mg, 0.19 mmol) was added. The reaction was stirred in a capped vial at 80° C. for 12 h. The reaction was cooled to rt, diluted with DCM and washed sequentially with water and brine. The organic layer was dried over Na_2SO_4 , filtered and concentrated to give crude (S)-1-(5-((4-cyclobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione. The crude product was used in the next step without any other purification. LCMS $[M+H]^+$: 547.6.

[1081] Step 2: (S)-1-(5-((4-cyclobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione (Example 329) was prepared using the method of Example 156, step 4, wherein (S)-1-(5-((4-cyclobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione was used in place of 1-(5-((4-cyclohexylmethyl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione. LCMS $[M+H]^+$: 397.2. 1H NMR (500 MHz, DMSO) δ 10.47 (s, 1H), 8.64 (d, $J=7.1$ Hz, 1H), 8.05 (s, 1H), 7.50 (s, 1H), 6.90 (d, $J=7.0$ Hz, 1H), 3.79 (t, $J=6.7$ Hz, 6H), 3.12 (d, $J=33.9$ Hz, 4H), 3.03-2.85 (m, 2H), 2.79 (t, $J=6.7$ Hz, 2H), 2.17 (d, $J=8.0$ Hz, 4H), 1.83-1.63 (m, 2H), 1.26 (dd, $J=31.8, 6.6$ Hz, 3H).


[1082] The compounds in the following table were prepared by the method of Example 329, using the appropriate commercially available ketone in step 1.

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
330	 (S)-1-((3-methyl-4-((tetrahydro-2H-pyran-4-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	427.2	(400 MHz, MeOD) δ 8.46 (dd, J = 7.2, 0.9 Hz, 1H), 8.02 (s, 1H), 7.50 (dd, J = 1.9, 0.9 Hz, 1H), 6.99 (dd, J = 7.2, 1.8 Hz, 1H), 4.04 (d, J = 11.3 Hz, 2H), 3.90 (t, J = 6.8 Hz, 2H), 3.71-3.54 (m, 3H), 3.52-3.39 (m, 4H), 3.25-3.12 (m, 4H), 2.89 (t, J = 6.8 Hz, 2H), 2.81 (s, 1H), 2.03 (d, J = 6.1 Hz, 1H), 1.83 (d, J = 25.3 Hz, 2H), 1.71-1.58 (m, 1H), 1.36-1.27 (m, 3H). NH proton not observed due to solvent exchange.
331	 (S)-1-((3-methyl-4-((oxetan-3-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	399.0	(400 MHz, MeOD) δ 8.45 (dd, J = 7.2, 1.0 Hz, 1H), 8.25 (s, 1H), 8.01 (s, 1H), 7.51 (dd, J = 1.9, 0.9 Hz, 1H), 6.99 (dd, J = 7.2, 1.8 Hz, 1H), 4.69 (dd, J = 12.8, 6.7 Hz, 3H), 4.61 (dd, J = 7.4, 6.3 Hz, 1H), 3.86 (dt, J = 21.3, 6.9 Hz, 3H), 3.68-3.56 (m, 2H), 2.89 (t, J = 6.8 Hz, 2H), 2.86-2.68 (m, 3H), 2.45 (dd, J = 25.5, 14.8 Hz, 2H), 2.32-2.12 (m, 2H), 0.92 (d, J = 6.5 Hz, 3H).
332	 (S)-1-((3-methyl-4-((2-oxaspiro[3.3]heptan-6-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	439.2	(400 MHz, MeOD) δ 8.48 (dd, J = 7.2, 0.9 Hz, 1H), 8.03 (s, 1H), 7.52 (s, 1H), 7.01 (dd, J = 7.2, 1.9 Hz, 1H), 3.90 (t, J = 6.8 Hz, 3H), 3.79-3.51 (m, 9H), 3.16-3.07 (m, 1H), 2.99-2.84 (m, 3H), 2.72-2.57 (m, 2H), 2.17-2.00 (m, 4H), 1.57 (d, J = 6.7 Hz, 3H). NH proton not observed due to solvent exchange.


-continued

Example No.	Structure	Mass [M + H] ⁺	¹ H NMR
333	 (S)-1-(5-((4-cyclohexyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	425.2	(400 MHz, MeOD) δ 8.46 (d, J = 7.1 Hz, 1H), 8.02 (s, 1H), 7.50 (d, J = 1.7 Hz, 1H), 6.99 (dd, J = 7.1, 1.8 Hz, 1H), 3.89 (t, J = 6.7 Hz, 2H), 3.68-3.57 (m, 2H), 3.45 (d, J = 25.8 Hz, 1H), 3.26 (s, 2H), 2.89 (t, J = 6.8 Hz, 4H), 2.41 (d, J = 60.0 Hz, 2H), 1.97 (dd, J = 42.9, 11.3 Hz, 4H), 1.72 (d, J = 13.1 Hz, 1H), 1.57 (d, J = 12.0 Hz, 1H), 1.49-1.12 (m, 8H). NH proton not observed due to solvent exchange.
334	 (S)-1-(5-((4-(4,4-difluorocyclohexyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	461.2	(500 MHz, DMSO) δ 10.47 (s, 1H), 8.64 (d, J = 7.1 Hz, 1H), 8.05 (s, 1H), 7.50 (s, 1H), 6.91 (d, J = 7.3 Hz, 1H), 3.79 (t, J = 6.8 Hz, 3H), 3.62 (s, 4H), 3.03 (s, 4H), 2.80 (t, J = 6.7 Hz, 3H), 2.34 (s, 1H), 2.09 (t, J = 23.8 Hz, 5H), 1.77 (s, 1H), 1.63 (s, 1H), 1.38 (d, J = 6.6 Hz, 1H), 1.26 (s, 2H).
335	 (S)-1-(5-((3-methyl-4-(spiro[3.3]heptan-2-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione	437.3	(400 MHz, MeOD) δ 8.47-8.44 (m, 1H), 8.02 (s, 1H), 7.49 (s, 1H), 6.98 (dd, J = 7.2, 1.9 Hz, 1H), 3.89 (t, J = 6.8 Hz, 2H), 3.70-3.54 (m, 3H), 3.25 (d, J = 9.0 Hz, 1H), 2.89 (t, J = 6.8 Hz, 5H), 2.41 (ddt, J = 27.4, 12.4, 6.0 Hz, 3H), 2.23 (t, J = 10.1 Hz, 1H), 2.12 (dd, J = 8.8, 6.0 Hz, 3H), 2.07-1.96 (m, 3H), 1.96-1.84 (m, 2H), 1.74-1.55 (m, 1H), 1.32 (d, J = 6.6 Hz, 3H). NH proton not observed due to solvent exchange.

Example 336. Preparation of (S)-1-((3-methyl-4-(2-methyl-2-azaspiro[3.3]heptan-6-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione



-continued



[1083] Step 1. tert-butyl (S)-6-(4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperazin-1-yl)-2-azaspiro[3.3]heptane-2-carboxylate was prepared using the method of Example 329, step 1, wherein tert-butyl 6-oxo-2-azaspiro[3.3]heptane-2-carboxylate was used in place of cyclobutanone. LCMS $[M+H]^+$: 688.3.


Step 2. (S)-3-(2,4-dimethoxybenzyl)-1-(5-((3-methyl-4-(2-azaspiro[3.3]heptan-6-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione trifluoroacetate

[1084] TFA (4 mL) was added to a solution of tert-butyl (S)-6-(4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperazin-1-yl)-2-azaspiro[3.3]heptane-2-carboxylate (300 mg, 0.43 mmol) in DCM (2 mL) at rt. The reaction was stirred at rt for 2 h and then concentrated. The residue was azeotropically dried with toluene to give crude (S)-3-(2,4-dimethoxybenzyl)-1-(5-((3-methyl-4-(2-azaspiro[3.3]heptan-6-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione trifluoroacetate which was used without further purification.

[1085] Step 3. (S)-3-(2,4-dimethoxybenzyl)-1-(5-((3-methyl-4-(2-methyl-2-azaspiro[3.3]heptan-6-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione was prepared using the method of Example 329, step 1, wherein paraformaldehyde was used in place of cyclobutanone and triethylamine was omitted. LCMS $[M+H]^+$: 602.3.

[1086] Step 4. (S)-1-((3-methyl-4-(2-methyl-2-azaspiro[3.3]heptan-6-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione was prepared from (S)-3-(2,4-dimethoxybenzyl)-1-(5-((3-methyl-4-(2-methyl-2-azaspiro[3.3]heptan-6-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl) dihydropyrimidine-2,4(1H,3H)-dione by the method of Example 156, step 4. LCMS $[M+H]^+$: 452.2. 1H NMR (400 MHz, MeOD) δ 8.44 (d, $J=7.2$ Hz, 1H), 8.01 (s, 1H), 7.48 (s, 1H), 6.98 (dd, $J=7.1, 1.8$ Hz, 1H), 4.16 (s, 2H), 4.03 (s, 2H), 3.89 (t, $J=6.8$ Hz, 2H), 3.16-3.09 (m, 1H), 2.93-2.79 (m, 5H), 2.70 (s, 2H), 2.60-2.17 (m, 7H), 2.00 (s, 1H), 1.68 (s, 1H), 1.48-1.37 (m, 2H), 1.09 (d, $J=6.4$ Hz, 3H). NH proton not observed due to solvent exchange.

Example 337. Preparation of (S)-1-(5-((3-methyl-4-(2-(methylsulfonyl)-2-azaspiro[3.3]heptan-6-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




Step 1. (S)-1-(5-((3-methyl-4-(2-azaspiro[3.3]heptan-6-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione trifluoroacetate

[1087] TFA (5 mL) was added to tert-butyl (S)-6-(4-((3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperazin-1-yl)-2-azaspiro[3.3]heptane-2-carboxylate (152 mg, 0.22 mmol). The reaction was stirred at 90° C. for 16 h and then concentrated. The residue was azeotropically dried with toluene to give crude (S)-1-(5-((3-methyl-4-(2-azaspiro[3.3]heptan-6-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione trifluoroacetate which was used without further purification.

[1088] Step 2: (S)-1-(5-((3-methyl-4-(2-(methylsulfonyl)-2-azaspiro[3.3]heptan-6-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione was prepared from (S)-1-(5-((3-methyl-4-(2-azaspiro[3.3]heptan-6-yl)piperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione trifluoroacetate by the method of Example 283, step 1 wherein methane-


sulfonyl chloride was used in place of ethylsulfonyl chloride. LCMS [M+H]⁺: 516.3. ¹H NMR (400 MHz, DMSO) δ 10.42 (s, 1H), 8.60-8.49 (m, 1H), 8.13 (s, 1H), 8.00 (s, 1H), 7.43 (s, 1H), 6.86 (dd, J=7.2, 1.8 Hz, 1H), 3.88 (s, 2H), 3.80-3.71 (m, 4H), 3.45 (q, J=13.7 Hz, 1H), 2.93 (s, 4H), 2.77 (t, J=6.7 Hz, 2H), 2.60 (s, 1H), 2.41 (s, 1H), 2.37-2.28 (m, 3H), 2.28-1.92 (m, 6H), 0.94 (d, J=6.3 Hz, 3H).

Example 338. Preparation of 1-(5-((4-cyclohexylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[1089] Prepared using the method of Example 329 wherein 3-(2,4-dimethoxybenzyl)-1-(5-(piperazin-1-ylmethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione was used in place of (S)-3-(2,4-dimethoxybenzyl)-1-(5-((3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione and cyclohexanone was used in place of cyclobutanone in step 1. LCMS [M+H]⁺: 411.2. ¹H NMR (300 MHz, CD₃OD) δ 8.46 (d, J=7.1 Hz, 1H), 8.03 (s, 1H), 7.50 (s, 1H), 6.99 (dd, J=7.2, 1.8 Hz, 1H), 3.89 (t, J=6.8 Hz, 2H), 3.67 (s, 2H), 3.11 (s, 2H), 2.89 (t, J=6.8 Hz, 8H), 2.12 (d, J=9.2 Hz, 2H), 1.94 (d, J=10.7 Hz, 2H), 1.72 (d, J=12.7 Hz, 1H), 1.53-1.15 (m, 6H). NH proton not observed due to solvent exchange.


Example 339. Preparation of (S)-1-(5-((4-(ethylsulfonyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[1090] Prepared from (S)-3-(2,4-dimethoxybenzyl)-1-(5-((3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione by the method of Example 283. LCMS [M+H]⁺: 435.1. ¹H NMR (500 MHz, DMSO) δ 10.53 (s, 1H), 8.75 (d, J=7.2 Hz, 1H), 8.14 (s, 1H), 7.71 (s, 1H), 7.01 (d, J=7.2 Hz, 1H), 4.27 (d, J=76.7 Hz, 5H), 3.90-3.77 (m, 2H), 3.71 (s, 1H), 3.35 (s, 2H), 3.17 (tq,


J=14.3, 7.1 Hz, 3H), 2.80 (dd, J=7.4, 6.1 Hz, 2H), 1.33 (d, J=7.0 Hz, 3H), 1.21 (t, J=7.3 Hz, 3H).

Example 340. Preparation of (S)-1-(5-((4-(cyclopropanecarbonyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[1091] Prepared by the method of Example 287 wherein (S)-3-(2,4-dimethoxybenzyl)-1-(5-((3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione was used in place of 3-(2,4-dimethoxybenzyl)-1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione in step 1. LCMS [M+H]⁺: 411.1. ¹H NMR (500 MHz, DMSO) δ 10.54 (s, 1H), 8.77 (d, J=7.2 Hz, 1H), 8.15 (s, 1H), 7.73 (s, 1H), 7.10-6.93 (m, 1H), 4.80 (s, 1H), 4.37 (s, 2H), 3.84 (ddd, J=10.1, 8.3, 5.0 Hz, 4H), 3.37 (s, 2H), 3.01 (s, 1H), 2.80 (t, J=6.7 Hz, 2H), 1.97 (t, J=6.4 Hz, 1H), 1.35 (s, 1H), 1.19 (d, J=10.9 Hz, 2H), 0.75 (s, 5H).

Example 341. Preparation of (S)-1-(5-((4-isobutyryl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione




[1092] Prepared using the method of Example 340 wherein isobutyric acid was used in place of cyclopropanecarboxylic acid in step 1. LCMS [M+H]⁺: 413.2. ¹H NMR (500 MHz, DMSO) δ 10.54 (s, 1H), 8.77 (d, J=7.2 Hz, 1H), 8.15 (s, 1H), 7.73 (s, 1H), 7.02 (d, J=7.3 Hz, 1H), 4.82 (s, 2H), 4.06 (s, 2H), 3.92-3.73 (m, 3H), 3.39 (d, J=41.9 Hz, 2H), 2.83 (dt, J=28.3, 6.7 Hz, 3H), 2.56 (s, 1H), 1.34 (d, J=6.9 Hz, 1H), 1.18 (d, J=7.1 Hz, 2H), 0.99 (dd, J=24.2, 6.9 Hz, 7H).

Example 342. Preparation of (S)-1-(5-((4-(cyclohexanecarbonyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[1093] Prepared using the method of Example 340 wherein cyclohexanecarboxylic acid was used in place of cyclopropanecarboxylic acid in step 1. LCMS [M+H]⁺: 453.1. ¹H NMR (500 MHz, DMSO) δ 10.54 (s, 1H), 8.77 (d, J=7.2 Hz, 1H), 8.15 (s, 1H), 7.72 (s, 1H), 7.02 (d, J=7.2 Hz, 1H), 4.81 (s, 1H), 4.41 (d, J=69.3 Hz, 3H), 3.82 (dd, J=7.0, 5.0 Hz, 4H), 3.40 (dd, J=35.2, 21.7 Hz, 3H), 2.94 (d, J=30.7 Hz, 2H), 2.80 (dd, J=7.3, 6.1 Hz, 2H), 1.71 (d, J=13.2 Hz, 2H), 1.64 (d, J=14.5 Hz, 2H), 1.31 (td, J=24.7, 14.1 Hz, 5H), 1.17 (d, J=7.3 Hz, 3H).


Example 343. Preparation of (S)-1-(5-((4-(cyclopentanecarbonyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[1094] Prepared using the method of Example 340 wherein cyclopentanecarboxylic acid was used in place of cyclopropanecarboxylic acid in step 1. LCMS [M+H]⁺:



439.1. ^1H NMR (500 MHz, DMSO) δ 10.54 (s, 1H), 8.77 (s, 1H), 8.15 (s, 1H), 7.72 (s, 1H), 7.02 (s, 1H), 4.81 (s, 1H), 4.37 (s, 4H), 3.87-3.58 (m, 2H), 3.39 (d, $J=14.3$ Hz, 3H), 3.09-2.86 (m, 2H), 2.80 (t, $J=6.7$ Hz, 2H), 1.75 (d, $J=21.2$ Hz, 3H), 1.56 (d, $J=33.7$ Hz, 5H), 1.33 (d, $J=6.9$ Hz, 1H), 1.18 (d, $J=6.9$ Hz, 2H).

Example 344. Preparation of (S)-1-(5-((4-(cyclobutanecarbonyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



[1095] Prepared using the method of Example 340 wherein cyclobutanecarboxylic acid was used in place of cyclopropanecarboxylic acid in step 1. LCMS $[\text{M}+\text{H}]^+$: 425.1. ^1H NMR (500 MHz, DMSO) δ 10.53 (s, 1H), 8.76 (s, 1H), 8.14 (s, 1H), 7.71 (s, 1H), 7.00 (s, 1H), 4.58 (d, $J=188.4$ Hz, 5H), 3.90-3.72 (m, 3H), 3.50-3.17 (m, 3H), 2.97 (s, 1H), 2.80 (t, $J=6.7$ Hz, 2H), 2.27-2.01 (m, 4H), 1.91 (p, $J=8.7$ Hz, 1H), 1.76 (d, $J=9.3$ Hz, 1H), 1.29 (d, $J=6.9$ Hz, 1H), 1.18 (d, $J=7.1$ Hz, 2H).

Example 345. Preparation of 1-(5-(1-(4-isobutylpiperazin-1-yl)ethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione



Step 1. 1-(5-acetylpyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione

[1096] Tributyl(1-ethoxyvinyl)stannane (757 mg, 2.09 mmol) and $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ (122 mg, 0.174 mmol) were added to a solution of 1-(5-bromopyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione (800 mg, 1.74 mmol) in DMF (8 mL) at rt. The mixture was stirred at 90°C . for 6 h, then cooled to rt and acidified with aqueous 1 N HCl solution. The mixture was partitioned between EtOAc and water. The organic layer was separated, washed with brine, dried over sodium sulfate, filtered and concentrated. The crude product was purified by flash silica gel chromatography (eluted with 45% EtOAc/hexane) to give 1-(5-acetylpyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione as an orange solid. LCMS $[\text{M}+\text{H}]^+$: 423.2.



Step 2. tert-butyl 4-(1-(3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)ethyl)piperazine-1-carboxylate

[1097] To a solution of 1-(5-acetylpyrazolo[1,5-a]pyridin-3-yl)-3-(2,4-dimethoxybenzyl)dihydropyrimidine-2,4(1H,3H)-dione (170 mg, 0.402 mmol) and tert-butyl piperazine-1-carboxylate (89 mg, 0.48 mmol) in THF (5 mL) was added dibutyltin dichloride (244 mg, 0.804 mmol), and triethylamine (0.17 mL, 1.2 mmol). The mixture was stirred at 80°C . for 2 h and then phenylsilane (87 mg, 0.80 mmol) was added. The reaction was stirred in a capped vial at 80°C . for 12 h. The reaction was cooled to rt, diluted with EtOAc and washed with water. The organic layer was dried over Na_2SO_4 , filtered and concentrated to give crude tert-butyl 4-(1-(3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)ethyl)piperazine-1-carboxylate. The crude product was used in the next step without any other purification. LCMS $[\text{M}+\text{H}]^+$: 593.0.

[1098] Step 3. 1-(5-(1-(4-isobutylpiperazin-1-yl)ethyl)pyrazolo[1,5-a]pyridin-3-yl)dihydropyrimidine-2,4(1H,3H)-dione was prepared by the method of Example 192,

steps 2-4 wherein tert-butyl 4-(1-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperazine-1-carboxylate was used in place of tert-butyl 4-((3-(3-(2,4-dimethoxybenzyl)-2,4-dioxotetrahydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)piperazine-1-carboxylate in step 2 and isobutyraldehyde was used in place of cyclohexanecarbaldehyde in step 3. LCMS $[M+H]^+$: 399.2. 1H NMR (400 MHz, MeOD) δ 8.47 (d, $J=7.3$ Hz, 1H), 8.03 (s, 1H), 7.51-7.47 (m, 1H), 7.02 (dd, $J=7.3, 1.9$ Hz, 1H), 3.90 (t, $J=6.8$ Hz, 2H), 3.67 (q, $J=6.6$ Hz, 1H), 3.25-2.55 (m, 12H), 2.08 (dq, $J=15.4, 7.6$ Hz, 1H), 1.45 (d, $J=6.7$ Hz, 3H), 1.02 (d, $J=6.6$ Hz, 6H). NH proton not observed due to solvent exchange.

Example 346. Preparation of 1-(5-(((2S,4R)-1-isobutyl-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione



Step 1. tert-butyl (2S,4R)-2-methyl-4-(pyrazolo[1,5-a]pyridin-5-ylmethyl)piperidine-1-carboxylate

[1099] To an oven-dried vial was added 5-bromopyrazolo[1,5-a]pyridine (100 mg, 507 μ mol), nickel chloride, dime-thoxyethane adduct (5.6 mg, 25 μ mol), pyridine-2,6-bis(carboximidamide)-hydrochloride (6.0 mg, 25 μ mol), activated zinc (83 mg, 1.3 mmol), tert-butyl (2S,4R)-4-(bromomethyl)-2-methylpiperidine-1-carboxylate (178 mg, 609 μ mol) and sodium iodide (19 mg, 127 μ mol). The reaction was sealed with a septa-top cap and purged with N_2 via needle. DMA (2 mL) was added, and the reaction was heated overnight at 70° C. The reaction was cooled to rt, diluted with EtOAc and filtered through a plug of silica gel, eluting with EtOAc. The eluent was concentrated and the residue was purified by silica gel column chromatography (eluted with 0-100% EtOAc in heptane) to give tert-butyl (2S,4R)-2-methyl-4-(pyrazolo[1,5-a]pyridin-5-ylmethyl)piperidine-1-carboxylate as a sticky solid. LCMS $[M+H]^+$: 330.3.

Step 2: tert-butyl (2S,4R)-4-((3-iodopyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperidine-1-carboxylate

[1100] To a solution of tert-butyl (2S,4R)-2-methyl-4-(pyrazolo[1,5-a]pyridin-5-ylmethyl)piperidine-1-carboxylate (220 mg, 668 μ mol) in MeCN (5 mL) at 0° C. was added NIS (180 mg, 801 μ mol). The reaction was then stirred at rt for 1 h. The reaction was quenched by addition of aqueous $Na_2S_2O_3$ solution and extracted with EtOAc. The organic layer was washed sequentially with water and brine, dried over Na_2SO_4 , filtered and concentrated. Silica gel column chromatography (eluting with 0-100% EtOAc in heptane) provided tert-butyl (2S,4R)-4-((3-iodopyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperidine-1-carboxylate as a transparent sticky solid. LCMS $[M+H]^+$: 456.1.

Step 3: tert-butyl (2S,4R)-4-((3-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperidine-1-carboxylate

[1101] To an oven-dried vial was added a solution of tert-butyl (2S,4R)-4-((3-iodopyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperidine-1-carboxylate (80 mg, 176 μmol) in DMSO (0.5 mL), pyrimidine-2,4(1H,3H)-dione (uracil) (26 mg, 228 μmol), potassium phosphate (78 mg, 369 μmol), N-(2-cyanophenyl) picolinamide (16 mg, 70 μmol) and copper(I) iodide (6.7 mg, 35 μmol). The vial was sealed with a septa-top cap and purged with N_2 via needle. The reaction was heated at 110°C . for 72 h. The reaction was quenched with aqueous 1M KHSO_4 and extracted with EtOAc. The organic layer was dried over Na_2SO_4 , filtered and concentrated. Silica gel column chromatography (EtOAc/heptane) provided tert-butyl (2S,4R)-4-((3-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperidine-1-carboxylate. LCMS $[\text{M}+\text{H}]^+$: 440.2.


Step 4: 1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione trifluoroacetate

[1102] To a solution of tert-butyl (2S,4R)-4-((3-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)pyrazolo[1,5-a]pyridin-5-yl)methyl)-2-methylpiperidine-1-carboxylate (35 mg, 80 μmol) in DCM (2 mL) was added TFA (2 mL). The reaction was stirred at rt for 45 min. The reaction was concentrated and the crude material was azeotropically dried with toluene to give crude 1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione trifluoroacetate which used without further purification. LCMS $[\text{M}+\text{H}]^+$: 340.2.

Step 5: 1-(5-(((2S,4R)-1-isobutyl-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione (Example 346)


[1103] To a solution of 1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione trifluoroacetate (36 mg, 79 μmol) in DCM (2 mL) and MeOH (500 μL) was added isobutyraldehyde (14 μL , 159 μmol) and triethylamine (10 μL , 71 μmol). The reaction was stirred at rt for 10 min and then sodium triacetoxyborohydride (84 mg, 397 μmol) was added. The reaction was stirred at rt overnight. The reaction was quenched with one drop of TFA and then concentrated. The crude material was dissolved in DMSO, filtered through a 1 micron filter and purified by reverse phase HPLC using ACN/Water/0.1% TFA to afford 1-(5-(((2S,4R)-1-isobutyl-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione. LCMS $[\text{M}+\text{H}]^+$: 396.2. ^1H NMR (500 MHz, DMSO) δ 11.51 (s, 1H), 8.67 (t, $J=26.8$ Hz, 1H), 8.14 (s, 1H), 7.71 (d, $J=7.8$ Hz, 1H), 7.36 (d, $J=14.1$ Hz, 1H), 6.87 (s, 1H), 5.71 (s, 1H), 3.59 (d, $J=89.5$ Hz, 1H), 3.23-2.69 (m, 2H), 2.59 (d, $J=47.7$ Hz, 1H), 2.36 (s, 1H), 2.23-1.81 (m, 2H), 1.78-1.14 (m, 7H), 0.89 (d, $J=69.0$ Hz, 8H).

Example 347. Preparation of 1-(5-(((2S,4R)-1-isobutyl-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-5-methylpyrimidine-2,4(1H,3H)-dione




[1104] Prepared using the method of Example 346 wherein 5-methylpyrimidine-2,4(1H,3H)-dione was used in place of pyrimidine-2,4(1H,3H)-dione in step 3. LCMS $[\text{M}+\text{H}]^+$: 410.2. ^1H NMR (500 MHz, DMSO) δ 11.53 (d, $J=6.5$ Hz, 1H), 8.73-8.54 (m, 1H), 8.15 (d, $J=1.4$ Hz, 1H), 7.62 (dd, $J=8.8, 1.4$ Hz, 1H), 7.42-7.23 (m, 1H), 6.89 (dd, $J=7.2, 1.9$ Hz, 1H), 3.69 (s, 1H), 3.24-2.90 (m, 3H), 2.89-2.69 (m, 2H), 2.60 (h, $J=6.6$ Hz, 1H), 2.23-1.88 (m, 2H), 1.83 (d, $J=1.2$ Hz, 3H), 1.77-1.61 (m, 3H), 1.61-1.43 (m, 1H), 1.34 (d, $J=6.6$ Hz, 1H), 1.22 (d, $J=6.8$ Hz, 2H), 1.03-0.88 (m, 6H).

Example 348. Preparation of 1-(5-(((2S,4R)-1-(cyclopropylmethyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-5-methylpyrimidine-2,4(1H,3H)-dione




[1105] Prepared using the method of Example 346 wherein 5-methylpyrimidine-2,4(1H,3H)-dione was used in place of pyrimidine-2,4(1H,3H)-dione in step 3 and cyclopropanecarbaldehyde was used in place of isobutyraldehyde in step 5. LCMS $[\text{M}+\text{H}]^+$: 408.2. ^1H NMR (500 MHz, DMSO) δ 11.53 (d, $J=5.4$ Hz, 1H), 8.69 (dt, $J=7.2, 1.4$ Hz, 1H), 8.16 (d, $J=1.4$ Hz, 1H), 7.63 (dd, $J=9.4, 1.4$ Hz, 1H), 7.37 (dd, $J=14.6, 1.8$ Hz, 1H), 6.90 (ddd, $J=7.2, 3.6, 1.8$ Hz, 1H), 3.65 (s, 1H), 3.44 (s, 1H), 3.37-2.93 (m, 3H), 2.75 (dd, $J=14.9, 7.2$ Hz, 1H), 2.69-2.56 (m, 1H), 2.26-2.03 (m, 1H), 1.83 (d, $J=1.2$ Hz, 3H), 1.80-1.53 (m, 3H), 1.51-1.37 (m, 1H), 1.32 (d, $J=6.6$ Hz, 1H), 1.23 (d, $J=6.9$ Hz, 2H), 1.05 (dt, $J=27.2, 7.1$ Hz, 1H), 0.73-0.53 (m, 2H), 0.49-0.22 (m, 2H).

Example 349. Preparation of 5-fluoro-1-((5-(((2S, 4R)-1-isobutyl-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione




[1106] Prepared using the method of Example 346 wherein 5-fluoropyrimidine-2,4(1H,3H)-dione was used in place of pyrimidine-2,4(1H,3H)-dione in step 3. LCMS $[M+H]^+$: 414.4. 1H NMR (500 MHz, DMSO) δ 12.04 (dd, $J=8.4, 5.2$ Hz, 1H), 8.69 (dd, $J=7.3, 2.7$ Hz, 1H), 8.29-8.06 (m, 2H), 7.51-7.37 (m, 1H), 6.97-6.82 (m, 1H), 3.38 (d, $J=11.0$ Hz, 1H), 2.97 (dt, $J=12.7, 6.1$ Hz, 3H), 2.90-2.70 (m, 2H), 2.61 (tt, $J=13.0, 7.0$ Hz, 1H), 2.29-1.91 (m, 2H), 1.68 (d, $J=4.5$ Hz, 3H), 1.62-1.44 (m, 1H), 1.35 (d, $J=6.6$ Hz, 1H), 1.23 (d, $J=6.8$ Hz, 2H), 1.08-0.83 (m, 6H).

Example 350. Preparation of 1-((5-(((2S,4R)-1-((4,4-difluorocyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-5-fluoropyrimidine-2,4(1H,3H)-dione




[1107] Prepared using the method of Example 346 wherein 5-fluoropyrimidine-2,4(1H,3H)-dione was used in place of pyrimidine-2,4(1H,3H)-dione in step 3 and 4,4-difluorocyclohexane-1-carbaldehyde was used in place of isobutyraldehyde in step 5. LCMS $[M+H]^+$: 490.4. 1H NMR (500 MHz, DMSO) δ 8.92 (s, 1H), 8.68 (dd, $J=7.2, 3.2$ Hz, 1H), 8.22 (dd, $J=10.6, 6.5$ Hz, 1H), 8.14 (d, $J=2.0$ Hz, 1H), 7.43 (d, $J=17.7$ Hz, 1H), 6.89 (d, $J=7.2$ Hz, 1H), 3.40-2.98 (m, 4H), 2.90 (t, $J=6.2$ Hz, 1H), 2.76-2.53 (m, 2H), 2.23-1.98 (m, 3H), 1.96-1.42 (m, 9H), 1.28 (dd, $J=54.9, 6.8$ Hz, 5H).

Example 351. Preparation of 1-((5-(((2S,4R)-1-((4,4-difluorocyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-5-methylpyrimidine-2,4(1H,3H)-dione




[1108] Prepared using the method of Example 346 wherein 5-methylpyrimidine-2,4(1H,3H)-dione was used in place of pyrimidine-2,4(1H,3H)-dione in step 3 and 4,4-difluorocyclohexane-1-carbaldehyde was used in place of isobutyraldehyde in step 5. LCMS $[M+H]^+$: 486.4. 1H NMR (500 MHz, DMSO) δ 8.89 (s, 1H), 8.67 (dd, $J=7.2, 3.2$ Hz, 1H), 8.14 (d, $J=2.0$ Hz, 1H), 7.61 (dd, $J=9.8, 1.5$ Hz, 1H), 7.34 (d, $J=16.0$ Hz, 1H), 6.87 (dd, $J=7.2, 1.9$ Hz, 1H), 3.73 (s, 1H), 3.53 (s, 1H), 3.28-2.97 (m, 2H), 2.89 (t, $J=6.1$ Hz, 1H), 2.75-2.53 (m, 2H), 2.10 (d, $J=80.9$ Hz, 3H), 1.93-1.42 (m, 12H), 1.27 (dd, $J=53.7, 6.8$ Hz, 5H).

Example 352. Preparation of 5-chloro-1-((5-(((2S, 4R)-1-((4,4-difluorocyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione



[1109] Prepared using the method of Example 346 wherein 5-chloropyrimidine-2,4(1H,3H)-dione was used in place of pyrimidine-2,4(1H,3H)-dione in step 3 and 4,4-difluorocyclohexane-1-carbaldehyde was used in place of isobutyraldehyde in step 5. LCMS $[M+H]^+$: 506.4. 1H NMR (500 MHz, DMSO) δ 8.86 (s, 1H), 8.68 (dd, $J=7.2, 3.0$ Hz, 1H), 8.24 (d, $J=9.9$ Hz, 1H), 8.16 (d, $J=2.0$ Hz, 1H), 7.43 (d, $J=19.0$ Hz, 1H), 6.91-6.85 (m, 1H), 3.41-2.99 (m, 4H), 2.90 (t, $J=6.2$ Hz, 1H), 2.75-2.54 (m, 2H), 2.24-1.95 (m, 3H), 1.95-1.41 (m, 9H), 1.27 (dd, $J=54.4, 6.8$ Hz, 5H).


Example 353. Preparation of 1-((5-(((2S,4R)-1-((4,4-difluorocyclohexyl)methyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-5-methoxypyrimidine-2,4(1H,3H)-dione



[1110] Prepared using the method of Example 346 wherein 5-methoxypyrimidine-2,4(1H,3H)-dione was used in place of pyrimidine-2,4(1H,3H)-dione in step 3 and 4,4-difluorocyclohexane-1-carbaldehyde was used in place of isobutyraldehyde in step 5. LCMS $[M+H]^+$: 502.2. 1H NMR (500 MHz, DMSO) δ 11.67 (s, 1H), 8.63 (d, $J=7.1$ Hz, 1H), 8.15 (s, 1H), 7.36 (d, $J=4.2$ Hz, 2H), 6.97-6.76 (m, 1H), 3.63 (s, 3H), 2.91 (s, 1H), 2.44-2.32 (m, 3H), 2.19 (s, 2H), 2.05-1.68


(m, 8H), 1.47 (d, $J=52.8$ Hz, 4H), 1.22 (d, $J=25.8$ Hz, 1H), 1.08 (d, $J=12.7$ Hz, 2H), 0.88 (s, 3H).

Example 354. Preparation of 5-cyclopropyl-1-((2S,4R)-1-isobutyl-2-methylpiperidin-4-yl)methylpyrazolo[1,5-a]pyridin-3-ylpyrimidine-2,4(1H,3H)-dione



[1111] Prepared using the method of Example 346 wherein 5-cyclopropylpyrimidine-2,4(1H,3H)-dione was used in place of pyrimidine-2,4(1H,3H)-dione in step 3. LCMS $[M+H]^+$: 436.2. 1H NMR (500 MHz, DMSO) δ 11.52 (d, $J=7.0$ Hz, 1H), 8.68 (dd, $J=7.2, 2.7$ Hz, 1H), 8.15 (d, $J=1.5$ Hz, 1H), 7.44-7.24 (m, 2H), 6.88 (d, $J=7.1$ Hz, 1H), 3.70 (s, 2H), 3.24-2.90 (m, 2H), 2.90-2.70 (m, 2H), 2.69-2.56 (m, 3H), 2.30-1.92 (m, 1H), 1.85-1.42 (m, 4H), 1.34 (d, $J=6.5$ Hz, 1H), 1.22 (d, $J=6.8$ Hz, 2H), 1.09-0.87 (m, 6H), 0.71 (dt, $J=8.8, 2.9$ Hz, 2H), 0.68-0.57 (m, 2H).


Example 355. Preparation of 1-(5-(((2S,4R)-1-(cyclopropanecarbonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione



[1112] HATU (38 mg, 0.099 mmol) and cyclopropanecarboxylic acid (8.5 mg, 0.099 mmol) were added to a solution of 1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione trifluoro-


acetate (30 mg, 0.066 mmol) in DMF (1.5 mL) at rt. The mixture was stirred at rt for 5 min and then DIPEA (0.035 mL, 0.19 mmol) was added. The mixture was stirred at rt overnight and then filtered through a 1 micron filter and purified by reverse phase HPLC using ACN/Water/0.1% TFA to afford 1-(5-(((2S,4R)-1-(cyclopropanecarbonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione. LCMS $[M+H]^+$: 408.2. 1H NMR (500 MHz, DMSO) δ 11.52 (d, $J=2.3$ Hz, 1H), 8.66 (dd, $J=7.2, 0.9$ Hz, 1H), 8.15 (s, 1H), 7.73 (d, $J=7.8$ Hz, 1H), 7.37 (d, $J=1.6$ Hz, 1H), 6.90 (dd, $J=7.2, 1.9$ Hz, 1H), 5.72 (dd, $J=7.8, 2.3$ Hz, 1H), 4.68 (d, $J=66.7$ Hz, 1H), 4.19 (dd, $J=89.6, 13.7$ Hz, 1H), 3.12 (t, $J=13.1$ Hz, 1H), 2.86-2.56 (m, 2H), 2.19-1.82 (m, 2H), 1.74-1.45 (m, 2H), 1.43-1.22 (m, 1H), 1.21-0.90 (m, 4H), 0.80-0.46 (m, 4H).

Example 356. Preparation of 1-(5-(((2S,4R)-1-(ethylsulfonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione



[1113] Triethylamine (0.046 mL, 0.33 mmol) and ethylsulfonyl chloride (0.019 mL, 0.19 mmol) were added to a solution of 1-(5-(((2S,4R)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione trifluoroacetate (30 mg, 0.066 mmol) in DCM (2 mL) at 0° C. The mixture was stirred at rt overnight and then filtered through a 1 micron filter and purified by reverse phase HPLC using ACN/Water/0.1% TFA to afford 1-(5-(((2S,4R)-1-(ethylsulfonyl)-2-methylpiperidin-4-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione (8 mg, 14 μ mol, 21% yield). LCMS $[M+H]^+$: 432.5. 1H NMR (500 MHz, DMSO) δ 11.52 (d, $J=2.3$ Hz, 1H), 8.66 (d, $J=7.1$ Hz, 1H), 8.15 (s, 1H), 7.73 (dd, $J=7.8, 1.2$ Hz, 1H), 7.37 (s, 1H), 6.89 (dd, $J=7.2, 1.8$ Hz, 1H), 5.72 (dd, $J=7.8, 2.3$ Hz, 1H), 4.05 (t, $J=6.3$ Hz, 1H), 3.49 (s, 1H), 3.14-2.86 (m, 3H), 2.59-2.54 (m, 2H), 2.03 (s, 1H), 1.55 (dd, $J=32.4, 13.3$ Hz, 2H), 1.37 (td, $J=12.8, 5.3$ Hz, 1H), 1.18 (t, $J=7.3$ Hz, 6H), 1.14-1.07 (m, 1H).

Example 357. Preparation of (S)-1-(5-((4-isobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione




Step 1: tert-butyl (S)-2-methyl-4-(pyrazolo[1,5-a]pyridin-5-ylmethyl)piperazine-1-carboxylate

[1114] To a suspension of 5-bromopyrazolo[1,5-a]pyridine (500 mg, 2.54 mmol) in toluene (10 mL) and water (1 mL) at room temperature was added Cs_2CO_3 (2.48 g, 7.61 mmol), tert-butyl (S)-2-methyl-4-((trifluoromethyl)boran-1-yl)methyl)piperazine-1-carboxylate, potassium salt (2.44 g, 7.61 mmol) and RuPhos (237 mg, 0.508 mmol), followed by $\text{Pd}(\text{OAc})_2$ (57 mg, 0.25 μmol). The mixture was stirred at 100° C. overnight, then cooled to rt and partitioned between EtOAc and water. The organic layer was separated, washed with brine, dried over sodium sulfate, filtered and concentrated. Silica gel column chromatography (EtOAc/EtOH/heptane) provided tert-butyl (S)-2-methyl-4-(pyrazolo[1,5-a]pyridin-5-ylmethyl)piperazine-1-carboxylate (735 mg, 1.9 mmol, 75% yield). LCMS $[\text{M}+\text{H}]^+$: 331.4.


[1115] Step 2. (S)-1-(5-((4-isobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione was prepared by the method of Example 346, steps 2-5 wherein tert-butyl (S)-2-methyl-4-(pyrazolo[1,5-a]pyridin-5-ylmethyl)piperazine-1-carboxylate was used in place of tert-butyl (2S,4R)-2-methyl-4-(pyrazolo[1,5-a]pyridin-5-ylmethyl)piperidine-1-carboxylate in step 2. LCMS $[\text{M}+\text{H}]^+$: 397.2. ^1H NMR (500 MHz, DMSO) δ 11.54 (s, 1H), 8.80-8.57 (m, 1H), 8.19 (s, 1H), 7.74 (d, $J=7.8$ Hz, 1H), 7.48 (d, $J=22.7$ Hz, 1H), 6.98 (d, $J=7.1$ Hz, 1H), 5.73 (d, $J=8.2$ Hz, 1H), 3.64 (d, $J=14.1$ Hz, 1H), 3.52 (t, $J=15.9$ Hz, 1H), 3.21-2.99 (m, 1H), 2.92 (d, $J=13.0$ Hz, 2H), 2.80-2.64 (m, 1H), 2.49-2.28 (m, 2H), 2.29-1.58 (m, 2H), 1.29 (d, $J=6.4$ Hz, 2H), 0.98 (dd, $J=15.1$, 6.6 Hz, 5H), 0.85 (s, 2H). Two missing protons attributed to overlap with solvent.

Example 358. Preparation of (S)-5-fluoro-1-(5-((4-isobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione




[1116] Prepared using the method of Example 357 wherein 5-fluoropyrimidine-2,4(1H,3H)-dione was used in place of pyrimidine-2,4(1H,3H)-dione in step 2. LCMS $[\text{M}+\text{H}]^+$: 415.2. ^1H NMR (500 MHz, DMSO) δ 12.05 (d, $J=5.3$ Hz, 1H), 8.74 (d, $J=7.2$ Hz, 1H), 8.23 (d, $J=6.4$ Hz, 1H), 8.19 (s, 1H), 7.59 (s, 1H), 6.99 (dd, $J=7.1$, 1.8 Hz, 1H), 3.51 (s, 4H), 3.21-2.72 (m, 7H), 2.21-1.75 (m, 1H), 1.27 (s, 3H), 0.97 (t, $J=6.7$ Hz, 6H).

Example 359. Preparation of (S)-1-(5-((4-isobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-5-methylpyrimidine-2,4(1H,3H)-dione




[1117] Prepared using the method of Example 357 wherein 5-methylpyrimidine-2,4(1H,3H)-dione was used in place of pyrimidine-2,4(1H,3H)-dione in step 2. LCMS $[\text{M}+\text{H}]^+$: 411.2. ^1H NMR (500 MHz, DMSO) δ 11.52 (s, 1H), 8.71 (dd, $J=7.1$, 0.9 Hz, 1H), 8.17 (s, 1H), 7.61 (q, $J=1.2$ Hz, 1H), 7.50 (s, 1H), 6.97 (dd, $J=7.2$, 1.8 Hz, 1H), 3.72 (s, 5H), 3.05 (s, 4H), 2.75 (d, $J=59.3$ Hz, 2H), 2.01 (s, 1H), 1.81 (d, $J=1.3$ Hz, 3H), 1.26 (d, $J=6.4$ Hz, 3H), 0.94 (t, $J=6.7$ Hz, 6H).

Example 360. Preparation of (S)-1-(5-((4-isobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-5-methoxypyrimidine-2,4(1H,3H)-dione




[1118] Prepared using the method of Example 357 wherein 5-methoxypyrimidine-2,4(1H,3H)-dione was used in place of pyrimidine-2,4(1H,3H)-dione in step 2. LCMS $[M+H]^+$: 427.2. 1H NMR (500 MHz, DMSO) δ 11.71 (s, 1H), 8.73 (d, $J=7.2$ Hz, 1H), 8.21 (s, 1H), 7.53 (s, 1H), 7.37 (s, 1H), 6.98 (d, $J=7.2$ Hz, 1H), 3.64 (s, 8H), 2.93 (d, $J=47.3$ Hz, 5H), 2.38 (s, 1H), 2.03 (s, 1H), 1.49-1.14 (m, 3H), 0.96 (t, $J=6.3$ Hz, 6H).

Example 361. Preparation of (S)-5-cyclopropyl-1-((4-isobutyl-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-ylpyrimidine-2,4(1H,3H)-dione




[1119] Prepared using the method of Example 357 wherein 5-cyclopropylpyrimidine-2,4(1H,3H)-dione was used in place of pyrimidine-2,4(1H,3H)-dione in step 2. LCMS $[M+H]^+$: 437.2. 1H NMR (500 MHz, DMSO) δ 11.53 (s, 1H), 8.72 (d, $J=7.1$ Hz, 1H), 8.19 (s, 1H), 7.49 (d, $J=14.4$ Hz, 1H), 7.35 (d, $J=0.9$ Hz, 1H), 6.97 (dd, $J=7.2, 1.8$ Hz, 1H), 3.52 (s, 5H), 2.94 (d, $J=49.6$ Hz, 5H), 2.38 (s, 1H), 2.03 (s, 1H), 1.65 (ddd, $J=11.0, 8.6, 5.3$ Hz, 1H), 1.39-1.16 (m, 3H), 0.96 (t, $J=6.6$ Hz, 6H), 0.81-0.67 (m, 2H), 0.65-0.47 (m, 2H).

Example 362. Preparation of (S)-1-((4-(cyclopropylmethyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)-5-methylpyrimidine-2,4(1H,3H)-dione



[1120] Prepared using the method of Example 357 wherein 5-methylpyrimidine-2,4(1H,3H)-dione was used in place of pyrimidine-2,4(1H,3H)-dione and cyclopropanecarbaldehyde was used in place of isobutyraldehyde in step 2. LCMS $[M+H]^+$: 409.2. 1H NMR (500 MHz, DMSO) δ 11.54 (s, 1H), 8.73 (d, $J=7.3$ Hz, 1H), 8.19 (s, 1H), 7.64 (d, $J=1.4$ Hz, 1H), 7.50 (s, 1H), 6.98 (dd, $J=7.2, 1.8$ Hz, 1H), 3.67 (d, $J=13.9$ Hz, 2H), 3.35 (s, 2H), 3.17 (d, $J=11.8$ Hz, 2H), 3.01 (t, $J=13.6$ Hz, 3H), 2.25 (t, $J=11.8$ Hz, 1H), 1.83 (d, $J=1.3$ Hz, 3H), 1.32 (d, $J=6.6$ Hz, 1H), 1.24 (d, $J=6.4$ Hz, 3H), 1.05 (s, 1H), 0.65 (d, $J=13.3$ Hz, 2H), 0.50-0.15 (m, 2H).

Example 363. Preparation of (S)-1-((4-((4,4-difluorocyclohexyl)methyl)-3-methylpiperazin-1-yl)methyl)pyrazolo[1,5-a]pyridin-3-yl)pyrimidine-2,4(1H,3H)-dione



[1121] Prepared using the method of Example 357 wherein 4,4-difluorocyclohexane-1-carbaldehyde was used in place of isobutyraldehyde in step 2. LCMS $[M+H]^+$: 473.2. 1H NMR (500 MHz, DMSO) δ 11.56 (s, 1H), 8.74 (d, $J=7.2$ Hz, 1H), 8.21 (s, 1H), 7.74 (d, $J=7.7$ Hz, 1H), 7.52 (s, 1H), 7.14-6.62 (m, 1H), 5.82-5.52 (m, 1H), 4.19 (s, 2H), 3.68 (s, 3H), 3.29 (s, 2H), 2.99 (s, 3H), 2.36 (d, $J=19.6$ Hz, 1H), 2.04 (s, 2H), 1.96-1.59 (m, 5H), 1.26 (s, 5H).

Biological Data

Abbreviations

- [1122]** BSA bovine serum albumin
- [1123]** Cas9 CRISPR associated protein 9
- [1124]** CRISPR Clustered regularly interspaced short palindromic repeats
- [1125]** crRNA CRISPR RNA
- [1126]** DMEM Dulbecco's modified eagle media
- [1127]** DMSO Dimethyl sulfoxide
- [1128]** DTT Dithiothreitol
- [1129]** EDTA ethylenediaminetetraacetic acid
- [1130]** eGFP enhanced green fluorescent protein
- [1131]** FACS fluorescence-activated cell sorting
- [1132]** FBS fetal bovine serum
- [1133]** FITC fluorescein
- [1134]** Flt3L Fms-related tyrosine kinase 3 ligand, Flt3L
- [1135]** HbF Fetal hemoglobin
- [1136]** HEPES (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid)
- [1137]** IMDM Iscove's modified Dulbecco's medium
- [1138]** KCl potassium chloride
- [1139]** mPB mobilized peripheral blood
- [1140]** PBS phosphate buffered saline
- [1141]** rhEPO recombinant human erythropoietin
- [1142]** rhIL-3 recombinant human interleukin-3
- [1143]** rhIL-6 recombinant human interleukin-6
- [1144]** rhSCF recombinant human stem cell factor
- [1145]** rhTPO recombinant human thrombopoietin
- [1146]** RNP ribonucleoprotein
- [1147]** shRNA short hairpin RNA
- [1148]** tracrRNA trans-activating crRNA
- [1149]** WIZ Widely-Interspaced Zinc Finger Containing Protein

Materials and Methods

Example 364: Quantification of WIZ Protein Levels in HiBit Tag Fusion Protein Assay

[1150] The HiBit system from Promega was used to develop high-throughput and quantitative assays to measure changes in WIZ protein levels in response to compounds. The HiBit tag was derived from a split Nanoluciferase and has the following protein sequence: VSGWRLFKKIS (SEQ ID NO: 1). The complementary fragment of Nanoluciferase (known as LgBit, from Promega), was added to the HiBit tag to form an active Nanoluciferase enzyme whose activity can be precisely measured. In this way, the levels of a fusion protein with the HiBit tag can be quantified in cell lysates.

[1151] Lentiviral vectors, based on the Invitrogen™ pLenti6.2/V5 DEST backbone were constructed that places the HiBit tag upstream of WIZ and expressed the fusion protein from an HSVTK promoter.

[1152] To ensure moderate and consistent expression of the HiBit-WIZ fusion protein across all cells in the population, stable cell lines were constructed from cells harboring a single copy of the construct. Lentivirus packaged with the constructs were made using the ViraPower™ kit from Invitrogen™. 293T cells from ATCC (Catalog number: CRL-3216), were infected with the virus at low multiplicity of infection and selected by 5 µg/mL blasticidin in culture media for 2 weeks.

[1153] The levels of HiBit-WIZ tagged fusion proteins in compound-treated cell lines were measured as follows:

[1154] On day 1, cells were diluted to 1.0×10^6 cells/ml in normal growth medium. 20 µL of cell suspension were plated in each well of a solid white 384-well plate. Plates were incubated overnight in a 37° C. and 5% CO₂ humidified tissue culture incubator.

[1155] On day 2, serial dilutions of compounds were made in 384-well plates. Compound plates were set up with DMSO in columns 1, 2, 23, 24, and 10-point compound dilution series in column 3-12 and column 13-22. 10 mM stock solution of compound were placed into column 3 or 13 and a 1:5 serial dilution was carried out until there was a 10-point dilution series per compound. 50 nL of diluted compounds were transferred into the plated cells by Echo® (Labcyte) acoustic transfer.

[1156] The highest concentration of compound was 25 µM. Plates were incubated overnight (about 18 hours) in a 37° C. and 5% CO₂ humidified tissue culture incubator.

[1157] On day 3, plates were removed from the incubator and allowed to equilibrate at room temperature for 60 minutes. HiBit substrate (Nano-Glo® HiBit Lytic Detection System, Promega Catalogue number: N3050) was added as described by the manufacturers protocols. Plates were incubated at room temperature for 30 minutes and luminescence was read using an EnVision® reader (PerkinElmer®). Data was analyzed and visualized using the Spotfire® software package.

Wiz Degradation Activity of Compounds (Table 1)

[1158] Table 1 shows WIZ degradation activity of compounds of the disclosure in the WIZ HiBit assay in 293T cells. WIZ Amax reflects the DMSO-normalized, curve-fitted percentage of WIZ-HiBit remaining at 25 µM. It was calculated by normalizing DMSO controls to 100%, parametric curve fitting of the dose response data (10-point,

5-fold), followed by calculation of response at 25 µM using the fitted equation (nd=not determined).

TABLE 1

Cmpd No.	WIZ AC ₅₀ (µM)	WIZ Amax	% degradation of WIZ (100-Amax)
1	0.020	9	91
2	0.023	9	91
3	0.020	10	90
4	0.005	10	90
5	0.204	11	89
6	0.013	11	89
7	0.023	14	86
8	0.008	14	86
9	0.031	16	84
10	0.004	16	84
11	0.055	16	84
12	0.051	16	84
13	0.019	16	84
14	0.028	16	84
15	0.523	29	71
16	0.018	16	84
17	0.048	18	82
18	0.034	18	82
19	0.020	19	81
20	0.022	21	79
21	0.025	22	78
22	0.071	23	77
23	0.144	23	77
24	1.075	23	77
25	0.040	24	76
26	0.297	24	76
27	0.046	24	76
28	0.105	25	75
29	0.232	26	74
30	0.201	20	80
31	0.027	26	74
32	0.023	18	82
33	0.031	27	73
34	0.945	27	73
35	3.138	27	73
36	0.068	28	72
37	0.446	28	72
38	0.112	29	71
39	0.083	29	71
39a	0.087	32	68
40	0.052	30	70
41	0.027	30	70
42	0.026	30	70
43	0.034	30	70
44	0.043	30	70
45	0.023	30	70
46	0.058	31	69
47	0.157	31	69
48	0.061	32	68
49	2.787	34	66
50	0.113	34	66
51	0.039	34	66
52	0.987	41	59
53	0.037	35	65
54	0.044	36	64
55	0.103	36	64
56	0.204	37	63
57	0.107	37	63
58	0.033	37	63
59	0.098	42	58
60	0.034	39	61
61	1.739	38	62
62	0.049	27	73
63	>25	64	36
64	0.043	6	94
65	0.032	6	94
66	0.195	28	72
67	0.561	39	61

TABLE 1-continued

Cmpd No.	WIZ AC ₅₀ (μM)	WIZ Amax	% degradation of WIZ (100-Amax)
68	0.384	43	57
69	0.350	65	35
70	0.021	17	83
71	0.007	15	85
72	0.006	13	87
73	0.003	11	89
74	0.003	10	90
75	0.098	19	81
76	0.036	19	81
77	0.077	32	68
78	0.010	17	83
79	0.065	35	65
80	0.049	12	88
81	0.015	8	92
82	0.033	10	90
83	0.076	11	89
84	0.055	13	87
85	0.038	17	83
86	0.027	20	80
87	0.025	21	79
88	0.181	22	78
89	0.285	23	77
90	2.002	23	77
91	0.150	26	74
92	0.131	31	69
93	6.518	38	62
94	3.993	38	62
95	>25	57	43
96	>25	58	42
97	24.763	56	44
98	0.587	41	59
99	1.833	48	52
100	0.755	58	42
101	>25	67	33
102	0.165	60	40
103	0.082	72	28
104	7.365	38	62
105	2.156	36	64
106	0.283	32	68
107	0.031	22	78
108	1.341	66	34
109	0.039	18	82
110	0.044	21	79
111	0.304	44	56
112	0.016	34	66
113	0.004	12	88
114	0.090	32	68
115	0.267	46	54
116	1.700	61	39
117	0.001	10	90
118	0.166	36	64
119	0.109	18	82
120	1.434	34	66
121	>25	50	50
122	2.577	45	55
123	0.095	32	68
124	0.502	42	58
125	0.596	45	55
126	0.29	65	35
127	0.013	31	69
128	0.83	71	29
129	0.80	65	35
130	1.32	64	36
131	0.37	62	38
132	0.72	59	41
133	4.8	79	21
134	4.3	73	27
135	2.1	74	26
136	0.033	14	86
137	0.121	23	77
138	0.10	20	80
139	0.003	17	83

TABLE 1-continued

Cmpd No.	WIZ AC ₅₀ (μM)	WIZ Amax	% degradation of WIZ (100-Amax)
140	0.007	10	90
141	0.002	9	91
142	0.011	11	89
143	0.004	12	88
144	0.008	13	87
145	0.025	16	84
146	0.051	18	82
147	0.005	10	90
148	0.063	14	86
149	0.0004	10	90
150	0.0003	10	90
151	0.27	14	86
152	0.91	19	81
153	0.014	14	86
154	0.041	22	78
155	0.009	15	85
156	0.023	14	86
157	0.022	17	83
158	0.025	18	82
159	0.023	19	81
160	0.045	18	82
161	0.115	20	80
162	0.079	21	79
163	0.561	24	76
164	0.323	31	69
165	0.198	31	69
166	0.632	34	66
167	0.286	43	57
168	0.127	44	56
169	0.326	45	55
170	0.228	47	53
171	0.981	48	52
172	0.527	58	42
173	0.771	58	42
174	0.095	19	81
175	1.240	60	40
176	1.396	62	38
177	0.106	23	77
178	0.048	24	76
179	0.049	18	82
180	0.078	19	81
181	0.030	25	75
182	0.199	33	67
183	0.468	35	65
184	1.627	71	29
185	0.582	53	47
186	0.076	21	79
187	0.020	20	80
188	0.071	23	77
189	0.191	29	71
190	0.086	46	54
191	0.683	50	50
192	0.019	14	86
193	0.010	14	86
194	0.003	11	89
195	2.047	52	48
196	3.821	59	41
197	1.144	60	40
198	0.009	13	87
199	0.009	14	86
200	0.008	10	90
201	0.020	12	88
202	0.005	15	85
203	0.025	16	84
204	0.051	16	84
205	0.024	18	82
206	0.193	22	78
207	0.014	14	86
208	0.580	40	60
209	0.38	26	74
210	0.016	14	86
211	0.033	14	86

TABLE 1-continued

Cmpd No.	WIZ AC ₅₀ (μM)	WIZ Amax	% degradation of WIZ (100-Amax)
212	0.029	16	84
213	0.048	19	81
214	0.151	35	65
215	2.617	81	19
216	0.089	35	65
217	0.092	25	75
218	0.068	29	71
219	0.212	25	75
220	0.133	34	66
221	0.318	46	54
222	0.457	36	64
223	0.877	49	51
224	0.854	55	45
225	1.050	68	32
226	1.233	55	45
227	0.956	40	60
228	0.704	45	55
229	>25	100	0
230	>25	100	0
231	0.15	47	53
232	0.206	52	48
233	0.589	60	40
234	0.942	58	42
235	3.277	65	35
236	3.198	95	5
237	>25	94	6
238	0.754	81	19
239	>25	100	0
240	0.025	20	80
241	0.351	56	44
242	1.604	93	7
243	3.065	87	13
244	1.179	47	53
245	0.351	48	52
246	>25	100	0
247	1.371	85	15
248	>25	100	0
249	0.30	59	41
250	1.35	59	41
251	0.165	82	18
252	0.277	58	42
253	0.531	77	23
254	>25	100	0
255	>25	100	0
256	0.186	22	78
257	0.101	26	74
258	0.148	31	69
259	0.258	39	61
260	0.796	32	68
261	0.417	37	63
262	0.583	32	68
263	0.88	45	55
264	0.26	36	64
266	0.044	19	81
267	0.10	14	86
268	0.024	11	89
269	1.02	51	49
270	0.099	43	57
271	0.702	37	63
272	0.025	31	69
273	0.040	22	78
274	0.032	23	77
275	0.020	21	79
276	0.009	15	84
277	0.203	24	76
278	0.066	28	72
279	0.026	9	91
280	0.008	12	88
281	0.004	17	83
282	0.001	19	81
283	0.029	19	80
284	0.018	20	80

TABLE 1-continued

Cmpd No.	WIZ AC ₅₀ (μM)	WIZ Amax	% degradation of WIZ (100-Amax)
285	0.003	18	82
286	0.031	20	79
287	0.015	19	81
288	0.008	20	80
289	0.033	21	79
290	1.70	46	54
291	0.015	17	83
292	0.011	17	83
293	0.065	24	76
294	0.018	19	81
295	0.024	20	80
296	0.010	23	77
297	0.029	32	68
298	0.242	49	51
299	0.009	18	82
300	0.003	18	82
301	0.002	14	86
302	0.042	16	84
303	0.046	20	80
304	0.002	13	87
305	0.002	15	85
306	0.008	16	84
307	0.006	16	84
308	0.044	23	77
309	0.056	38	62
310	0.15	27	73
311	0.31	22	78
312	0.37	56	44
313	0.085	30	70
314	2.56	66	34
315	1.02	47	53
316	0.061	17	83
317	0.031	20	80
318	0.010	13	87
319	0.004	21	78
320	0.018	21	78
321	0.067	22	78
322	0.031	14	86
323	0.004	17	83
324	0.108	28	72
325	0.047	18	82
326	0.049	13	87
327	0.653	46	54
328	0.338	48	52
329	0.166	17	83
330	0.041	18	82
331	0.073	24	76
332	22	51	49
333	0.068	13	87
334	0.032	15	85
335	0.006	12	88
336	12	24	76
337	0.399	17	83
338	0.033	17	83
339	0.715	65	35
340	0.862	68	32
341	0.799	70	30
342	0.916	63	37
343	0.495	54	46
344	1.07	65	35
345	0.191	27	73
346	0.108	13	87
347	0.378	11	89
348	0.400	14	86
349	0.118	14	86
350	0.028	10	90
351	0.047	9	91
352	0.192	10	90
353	9.1	30	70
354	>25	100	0
355	0.183	18	82
356	0.162	17	83

TABLE 1-continued

Cmpd No.	WIZ AC ₅₀ (μM)	WIZ Amax	% degradation of WIZ (100-Amax)
357	0.124	15	85
358	0.128	14	86
359	0.365	11	89
360	>25	76	24
361	>25	100	0
362	0.765	16	84
363	0.109	15	85

Example 365: Small Molecule HbF Induction Assay

[1159] Cryopreserved primary human CD34⁺ hematopoietic stem and progenitor cells were obtained from AllCells, LLC. The CD34⁺ cells were isolated from the peripheral blood of healthy donors after mobilization by administration of granulocyte colony-stimulating factor. Cells were differentiated ex vivo toward the erythroid lineage using a 2-phase culture method. In the first phase, cells were cultured in StemSpan™ Serum-Free Expansion Media (SFEM) (STEMCELL Technologies Inc.) supplemented with rhSCF (50 ng/mL, Peprotechs, Inc.), rhIL-6 (50 ng/mL, Peprotech®, Inc.), rhIL-3 (50 ng/mL, Peprotech®, Inc.), and rhFlt3L (50 ng/mL, Peprotech®, Inc.), and 1× anti biotic-antimycotic (Life Technologies, Thermo Fisher Scientific) for 6 days at 37° C. with 5% 002. During the second phase, cells were cultured in erythroid differentiation media at 5,000 cells/mL in the presence of compound for 7 days at 37° C. with 5% 002. Erythroid Differentiation Media is comprised of IMDM (Life Technologies) supplemented with insulin (10 μg/mL, Sigma Aldrich), heparin (2 U/mL Sigma Aldrich), holo-transferrin (330 μg/mL, Sigma Aldrich), human serum AB (5%, Sigma Aldrich), hydrocortisone (1 μM, STEMCELL Technologies), rhSCF (100 ng/mL, Peprotech®, Inc.), rhIL-3 (5 ng/mL, Peprotech®, Inc.), rhEPO (3 U/mL, Peprotech®, Inc.), and 1× antibiotic-antimycotic. All compounds were dissolved and diluted into dimethylsulfoxide (DMSO) and were added to culture media for a final concentration of 0.3% DMSO for testing in a 7-point, 1:3 dilution series starting at 30 μM.

Staining and Flow Cytometry

[1160] For viability analysis, samples were washed and resuspended in phosphate-buffered saline (PBS) and stained with LIVE/DEAD™ Fixable Violet Dead Cell Stain Kit (Life Technologies, L34963) for 20 minutes. Cells were then washed again with PBS and resuspended in PBS supplemented with 2% fetal bovine serum (FBS), and 2 mM EDTA to prepare for cell surface marker analysis. Cells were labeled with allophycocyanin-conjugated CD235a (1:100, BD Biosciences, 551336) and Brilliant Violet-conjugated CD71 (1:100, BD Biosciences, 563767) antibodies for 20 minutes. For analysis of cytoplasmic Fetal Hemoglobin (HbF), cells were fixed and permeabilized using the Fixation (BioLegend®, 420801) and Permeabilization Wash (BioLegend®, 421002) Buffers according to the manufacturer's protocol. During the permeabilization step, cells were stained with phycoerythrin-conjugated or FITC-conjugated HbF-specific antibody (1:10-1:25, Invitrogen™, MHFH04-

4) for 30 minutes. Stained cells were washed with phosphate-buffered saline before analysis on the FACSCanto™ II flow cytometer or LSRFortessa™ (BD Biosciences). Data analysis was performed with FlowJo™ Software (BD Biosciences).

HbF Induction Activity of Compounds (Table 2)

[1161] mPB CD34⁺ cells were expanded for 6 days, then erythroid differentiated in the presence of compound for 7 days. Cells were fixed, stained and analyzed by flow cytometry. Table 2 shows HbF induction activity of the compounds. HbF Amax=the highest percentage of cells staining positive for HbF (% HbF+ cells) in the fitted dose-response curve. The baseline % HbF+ cells for DMSO-treated cells is approximately 30-40%.

TABLE 2

Cmpd no.	HbF EC ₅₀ (μM)	HbF Amax
17	0.266	80
109	0.129	77
141	0.006	90
142	0.038	89
145	0.030	76
156	0.012	81
160	0.999	84
179	0.074	79
203	0.045	82
207	0.089	85
210	0.064	86
283	0.127	61
287	0.060	63

Example 366: Cell Culture for shRNA and CRISPR Assays

[1162] HEK293T cells were maintained in DMEM high glucose complete media with sodium pyruvate, non-essential amino acids, 10% FBS, 2 mM L-glutamine, 100 U/mL pen/strep, 25 mM HEPES. Unless stated otherwise, all reagents for culturing HEK293T cells were obtained from Invitrogen™.

[1163] Mobilized peripheral blood (mPB) CD34⁺ cells (AllCells, LLC) were maintained in StemSpan™ serum-free expansion media (SFEM) (STEMCELL Technologies Inc.) supplemented with 50 ng/mL each of rhTPO, rhIL-6, rhFLT3L, rhSCF for 2-3 days prior to shRNA transduction or targeted ribonucleoprotein (RNP) electroporation targeting WIZ. All cytokines were obtained from Peprotech®, Inc. Cell cultures were maintained at 37° C. and 5% CO₂ in a humidified tissue culture incubator.

Generation of shRNA Lentiviral Clones Targeting WIZ

[1164] 5-phosphorylated sense and anti-sense complementary single-stranded DNA oligos of the respective shRNA against WIZ were synthesized by Integrated DNA Technologies, Inc. (IDT). Each DNA oligonucleotide was designed with PmeI/AscI restriction overhangs on 5'- and 3'-ends, respectively, for subsequent compatible ligation into the lentiviral vector backbone. Equimolar of each of the complementary oligonucleotides were annealed in NEB Buffer 2 (New England Biolabs® Inc.) by heating on a heating block at 98° C. for 5 minutes followed by cooling to room temperature on the bench top. Annealed double-stranded DNA oligonucleotides were ligated into pHAGE

lentiviral backbone digested with PmeI/AscI using T4 DNA ligase kit (New England Biolabs). Ligation reactions were transformed into chemically competent Stbl3 cells (Invitrogen™) according to the manufacturer’s protocol. Positive clones were verified using the sequencing primer (5'-ctacat-tttacatgatagg-3'; SEQ ID NO: 2) and plasmids were purified by Alta Biotech LLC.

[1165] Lentivirus particles for the respective shRNA constructs were generated by co-transfection of HEK293T cells with pCMV-dR8.91 and pCMV-VSV-G expressing envelope plasmid using Lipofectamine 3000 reagent in 150 mm tissue culture dish format as per manufacturer’s instructions (Invitrogen™). Lentivirus supernatant was harvested 48 hours after co-transfection, filtered through a 0.45 µm filter (Millipore) and concentrated using Amicon Ultra 15 with Ultracel-100 membrane (Millipore). Infectious units of each of the lentivirus particle was determined by flow cytometry using eGFP expression as marker of transduction after serial dilution and infection of HEK293T cells.

[1166] The shRNA sequences are as follows:

```
shWIZ_#1
                               (SEQ ID NO: 3)
5'-AGCCCAATGCCACGGAAT-3';

shWIZ_#2
                               (SEQ ID NO: 4)
5'-GCAACATCTACACCTCAAAT-3';

shWIZ_#4
                               (SEQ ID NO: 5)
5'-TGACCGAGTGGTACGTCAATG-3';

shWIZ_#5
                               (SEQ ID NO: 6)
5'-AGCGGCAGAACATCAACAAAT-3'.
```

Lentiviral shRNA Transduction and FACS of mPB CD34+ Cells

[1167] mPB CD34+ transduction was performed on retroNectin coated non-tissue culture treated 96 well-flat bottom plates (Corning, Inc.). Briefly, plates were coated with 100 µL of RetroNectin® (1 µg/mL) (TAKARABIO, Inc.), sealed and incubated at 4° C. overnight. RetroNectin® was then removed and plates were incubated with BSA (bovine

serum albumin) (1%) in PBS for 30 minutes at room temperature. Subsequently, BSA (bovine serum albumin) was aspirated and replaced with 100 µL of lentiviral concentrate and centrifuged at 2000×g for 2 hours at room temperature. Next, residual supernatant was gently pipetted out and ready for transductions of mPB CD34+ cells. Ten thousand cells were plated in 150 µL of StemSpan™ Serum-free Expansion Medium (SFEM) supplemented with 50 ng/mL each of rhTPO, rhIL-6, rhFLT3L, rhSCF to initiate transduction. Cells were cultured for 72 hours prior to assessing transduction efficiencies using eGFP expression as a marker.

[1168] eGFP-positive cells were sorted on an FACS Aria™ III (BD Biosciences). Briefly, the transduced mPB CD34+ cell population was washed and re-suspended with FACS buffer containing 1× Hank’s buffered saline solution, EDTA (1 mM) and FBS (2%). Sorted eGFP-positive cells were used for the erythroid differentiation assay.

Targeting CRISPR Knockout of WIZ

[1169] Alt-R CRISPR-Cas9 crRNA and tracrRNA (5'-AGCAUAGCAAGUAAAAUAAGGCUAGU-CCGUUAUCAACUUGAAAAAGUGGCACCGAGUCGGUGCUUU-3'; SEQ ID NO: 7) were purchased from Integrated DNA Technologies, Inc. Equimolar tracrRNA was annealed with WIZ targeting crRNA (Table 3) in Tris buffer (10 mM, pH 7.5) by heating at 95° C. for 5 minutes using a polymerase chain reaction (PCR) machine (Bio-Rad) followed by cooling to room temperature on the benchtop. Subsequently, a ribonucleoprotein (RNP) complex was generated by mixing annealed tracrRNA:crRNA with 6 µg of Cas9 at 37° C. for 5 minutes in 1× buffer containing HEPES (100 mM), KCl (50 mM), MgCl₂ (2.5 mM), glycerol (0.03%), DTT (1 mM) and Tris pH 7.5 (2 mM).

[1170] Electroporation of the RNP complex was performed on a 4D-Nucleofector™ (Lonza) as per manufacturer’s recommendation. Briefly, 50,000 mPB CD34+ cells resuspended in Primary Cell P3 Buffer with supplement (Lonza) were pre-mixed with 5 µL of RNP complex per well in nucleocuvettes and incubated for 5 minutes at room temperature. Subsequently, the mixture was electroporated using the CM-137 program. Cells were cultured for 72 hours post-RNP electroporation before initiating erythroid differentiation. The crRNA sequences are shown in Table 3 below.

TABLE 3

Name	Sequence (5' to 3')	Target genomic region	Strand	SEQ ID NO
rg_0111	ACGGAGGCTAAGCGTCGCAA	random guide, non-targeting		8
WIZ_6	AACATCTTTGGGCGTAGG	chr19:15427143-15427163	(+)	9
WIZ_9	GACATCCGCTGCGAGTTCTG	chr19:15427488-15427510	(-)	10
WIZ_12	TGCAGCGTCCCGGCAGAGC	chr19:15425751-15425773	(-)	11
WIZ_14	CAAGCCGTGCCTCATCAAGA	chr19:15425571-15425593	(-)	12
WIZ_15	CGGGCACACCTGCGGCAGTT	chr19:15424942-15424964	(-)	13
WIZ_18	AGTGGGTGCGGCACTTACAG	chr19:15423169-15423191	(-)	14

Erythroid Differentiation of shRNA Transduced or RNP Electroporated mPB CD34+ Cells

[1171] Erythroid differentiation was initiated by plating 8,000 RNP-electroporated or FACS sorted eGFP+ mPB CD34+ cells per well in 96-well tissue culture plate. Base differentiation media consists of IMDM (Iscove's Modified Dulbecco's Medium), human AB serum (5%), transferrin (330 µg/mL), Insulin (10 µg/mL) and Heparin (2 IU/mL). Differentiation media was supplemented with rhSCF (100 ng/mL), rhIL-3 (10 ng/mL), rhEPO (2.5 U/mL) and hydrocortisone (1 µM). After 4 days of differentiation, the cells were split (1:4) in fresh media to maintain optimal growth density. Cells were cultured for additional 3 days and utilized for assessment of fetal hemoglobin (HbF) expression.

Analysis of HbF Gene Expression by RNA-Seq

[1172] Two independent, targeted CRISPR/Cas9 knockout (KO) of WIZ was done using WIZ_6 and WIZ_18 gRNAs or a non-targeting scrambled gRNA negative control in mPB CD34+ HSCs. Cells from KO and negative control were then cultured for 7 days for erythroid differentiation and used for total RNA isolation (Zymo Research, catalogue #R1053). The quality of isolated RNA was determined before sequencing using Agilent RNA 6000 Pico Kit (Agilent, catalogue #5067-1513).

[1173] RNA sequencing libraries were prepared using the Illumina TruSeq Stranded mRNA Sample Prep protocol and sequenced using the Illumina NovaSeq6000 platform (Illumina). Samples were sequenced to a length of 2×76 base-pairs. For each sample, salmon version 0.8.2 (Patro et al. 2017; doi: 10.1038/nmeth.4197) was used to map sequenced fragments to annotated transcripts in the human reference genome hg38 provided by the ENSEMBL database. Per-gene expression levels were obtained by summing the counts of transcript-level counts using tximport (Soneson et al. 2015; doi: 10.12688/f1000research.7563.1). DESeq2 was used to normalize for library size and transcript length differences, and to test for differential expression between samples treated with the gRNAs targeting WIZ and the samples treated with the scrambled gRNA controls (Love et al. 2014; doi: 10.1186/s13059-014-0550-8). Data were visualized using ggplot2 (Wickham H (2016). ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York. ISBN 978-3-319-24277-4; <https://ggplot2.tidyverse.org>).

HbF Intracellular Staining

[1174] One hundred thousand cells were aliquoted into U-bottom 96-well plate and stained for 20 min in the dark with diluted LIVE/DEAD fixable violet viability dye as per manufacturer's recommendation (Invitrogen). Cells were washed with FACS staining buffer and subsequently stained with anti-CD71-BV711 (BD Biosciences) and anti-CD235a-APC (BD Biosciences) for 20 mins in the dark. After two rounds of washes with three volumes of 1×PBS, cells were

fixed and permeabilized with 1×BD Cytofix/Cytoperm (BD Biosciences) for 30 minutes at room temperature in the dark. Subsequently, cells were washed twice with three volumes of 1× Perm/wash buffer (BD Biosciences). Anti-HbF-FITC (ThermoScientific) was diluted (1:25) in 1× perm/wash buffer, added to permeabilized cells and incubated for 30 minutes at room temperature in the dark. Next, cells were washed twice with three volumes of 1× perm/wash buffer and analyzed by flow cytometry using LSR Fortessa (BD Biosciences). Data was analyzed with FlowJo software.

Results

Wiz KO Upregulates HBG1/2 Expression Upon Erythroid Differentiation

[1175] Targeted KO of WIZ using two independent gRNAs (WIZ_6 and WIZ_18) demonstrated upregulation of fetal hemoglobin genes (HBG1/2), as presented in FIG. 1A.

Loss of WIZ Induces Fetal Hemoglobin Expression in mPB CD34+ Derived Erythroid Cells

[1176] In order to validate whether WIZ is a negative regulator of HbF expression, shRNA and CRISPR-Cas9-mediated knockdown and knockout functional genetics approaches were employed. mPB CD34+ cells were treated with shRNA or CRISPR-Cas9 reagents and erythroid differentiated for 7 days prior to flow cytometry analysis. Targeted knockdown of WIZ transcript results in 78-91% HbF+ cells compared to 40% for the negative control scrambled shRNA. Error bars represent standard error of two biological replicates with three technical replicates each (FIG. 1B). CRISPR/Cas9-mediated targeted loss of WIZ results in 62-88% HbF+ cells compared to 39% for random guide crRNA. Error bars represent standard error of one biological sample with four technical replicates (FIG. 1C). To summarize, the results indicate that loss of WIZ induces HbF in human primary erythroid cells. As such, the zinc finger transcription factor Widely Interspaced Zinc Finger Motifs (WIZ) was identified as a novel target for HbF induction. These data provide genetic evidence that WIZ is a regulator of fetal hemoglobin expression and represents a novel target for the treatment of sickle cell disease and beta-thalassemia.

[1177] Having thus described several aspects of several embodiments, it is to be appreciated various alterations, modifications, and improvements will readily occur to those skilled in the art. Such alterations, modifications, and improvements are intended to be part of this disclosure, and are intended to be within the spirit and scope of the disclosure. Accordingly, the foregoing description and drawings are by way of example only.

[1178] Those skilled in the art will recognize, or be able to ascertain, using no more than routine experimentation, numerous equivalents to the specific embodiments described specifically herein. Such equivalents are intended to be encompassed in the scope of the following claims.

SEQUENCE LISTING

Sequence total quantity: 14
 SEQ ID NO: 1 moltype = AA length = 11
 FEATURE Location/Qualifiers
 REGION 1..11


-continued

	note = source = /note="Description of Artificial Sequence: Synthetic peptide"	
source	1..11 mol_type = protein organism = synthetic construct	
SEQUENCE: 1		
VSGWRLFKKI S		11
SEQ ID NO: 2	moltype = DNA length = 19 Location/Qualifiers	
FEATURE	1..19	
misc_feature	note = source = /note="Description of Artificial Sequence: Synthetic primer"	
source	1..19 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 2		
ctacatttta catgatagg		19
SEQ ID NO: 3	moltype = DNA length = 21 Location/Qualifiers	
FEATURE	1..21	
misc_feature	note = source = /note="Description of Artificial Sequence: Synthetic oligonucleotide"	
source	1..21 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 3		
agcccacaat gccacggaaa t		21
SEQ ID NO: 4	moltype = DNA length = 21 Location/Qualifiers	
FEATURE	1..21	
misc_feature	note = source = /note="Description of Artificial Sequence: Synthetic oligonucleotide"	
source	1..21 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 4		
gcaacatcta caccctcaaa t		21
SEQ ID NO: 5	moltype = DNA length = 21 Location/Qualifiers	
FEATURE	1..21	
misc_feature	note = source = /note="Description of Artificial Sequence: Synthetic oligonucleotide"	
source	1..21 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 5		
tgaccgagtg gtacgtcaat g		21
SEQ ID NO: 6	moltype = DNA length = 21 Location/Qualifiers	
FEATURE	1..21	
misc_feature	note = source = /note="Description of Artificial Sequence: Synthetic oligonucleotide"	
source	1..21 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 6		
agcggcagaa catcaacaaa t		21
SEQ ID NO: 7	moltype = RNA length = 67 Location/Qualifiers	
FEATURE	1..67	
misc_feature	note = source = /note="Description of Artificial Sequence: Synthetic oligonucleotide"	
source	1..67 mol_type = other RNA organism = synthetic construct	
SEQUENCE: 7		
agcatagcaa gttaaaataa ggctagtcg ttatcaactt gaaaaagtgg caccgagtcg		60
gtgcttt		67
SEQ ID NO: 8	moltype = DNA length = 20	

-continued

FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = source = /note="Description of Artificial Sequence: Synthetic oligonucleotide"	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 8		
acggaggcta agcgtcgcaa		20
SEQ ID NO: 9	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = source = /note="Description of Artificial Sequence: Synthetic oligonucleotide"	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 9		
aacatcttttc gggccgtagg		20
SEQ ID NO: 10	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = source = /note="Description of Artificial Sequence: Synthetic oligonucleotide"	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 10		
gacatccgct gcgagttctg		20
SEQ ID NO: 11	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = source = /note="Description of Artificial Sequence: Synthetic oligonucleotide"	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 11		
tgcagcgtcc cgggcagagc		20
SEQ ID NO: 12	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = source = /note="Description of Artificial Sequence: Synthetic oligonucleotide"	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 12		
caagccgtgc ctcacaaaga		20
SEQ ID NO: 13	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = source = /note="Description of Artificial Sequence: Synthetic oligonucleotide"	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 13		
cgggcacacc tgcggcagtt		20
SEQ ID NO: 14	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = source = /note="Description of Artificial Sequence: Synthetic oligonucleotide"	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 14		
agtgggtgcg gcacttacag		20

1. A compound of Formula (I'') or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein:



— is a single bond or a double bond;

X is selected from CH, CF, and N;

R^x is selected from hydrogen, C₁-C₆alkyl, halo (e.g., F, Cl), C₁-C₆alkoxy, and C₃-C₈cycloalkyl;

R' is selected from hydrogen and C₁-C₆alkyl;

R¹ is selected from hydrogen and C₁-C₆alkyl;

each R² is independently selected from C₁-C₆alkyl, C₁-C₆haloalkyl, halo, and oxo, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{2a}; or 2 R² on non-adjacent carbon atoms together with the non-adjacent carbon atoms to which they are attached form a bridging ring;

R^{2a} is selected from C₁-C₆alkoxy and hydroxyl;

R³ is selected from hydrogen, C₁-C₈alkyl, C₂-C₆alkenyl, —SO₂R⁴, C₁-C₆haloalkyl, —C(=O)—O—(R⁵), —C(=O)—(R⁶), C₃-C₁₀cycloalkyl, and a 4- to 10-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, wherein the C₁-C₈alkyl and C₁-C₆haloalkyl are each independently substituted with 0-3 occurrences of R^{3a}, and wherein the C₃-C₁₀cycloalkyl and 4- to 10-membered heterocyclyl are each independently substituted with 0-3 occurrences of R^{3b};

or

R³ together with the nitrogen atom to which it is attached and R² together with the carbon atom to which it is attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O and S, which 5- or 6-membered heterocyclyl is substituted with 0-2 occurrences of an oxo group;

each R^{3a} is independently selected from C₃-C₁₀cycloalkyl, a 4- to 10-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, a 5- to 10-membered heteroaryl comprising 1-4 heteroatoms independently selected from N, O, and S, C₆-C₁₀aryl, C₁-C₆alkoxy, hydroxyl, and —C(=O)—NR⁷R⁸, wherein the C₃-C₁₀cycloalkyl, 4- to 6-membered heterocyclyl, 5- to 10-membered heteroaryl and C₆-C₁₀aryl are substituted with 0-4 occurrences of R^{3b};

each R^{3b} is independently selected from C₁-C₆alkoxy, halo, C₁-C₆haloalkyl, C₁-C₆haloalkoxy, C₁-C₆alkyl, —CN, —SO₂NR⁷R⁸, —SO₂R⁴, and hydroxyl;

R⁴ is selected from C₃-C₈cycloalkyl, C₁-C₆alkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, C₆-C₁₀aryl,

and —NR^{4b}R^{4c}, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{4a};

R^{4a} is selected from C₃-C₈cycloalkyl, C₆-C₁₀aryl, and C₁-C₆alkoxy;

R^{4b} is selected from hydrogen, and C₁-C₆alkyl;

R^{4c} is selected from hydrogen, C₁-C₆alkyl, and C₃-C₈cycloalkyl;

R⁵ is selected from C₁-C₆alkyl, C₃-C₈cycloalkyl, and C₆-C₁₀aryl;

R⁶ is selected from C₁-C₆alkyl, C₃-C₈cycloalkyl, C₆-C₁₀aryl, a 4- to 10-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, and and —NR^{4b}R^{4c}, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{6a}, the C₃-C₈cycloalkyl is substituted with 0-1 occurrence of R^{6b}, and the 4- to 10-membered heterocyclyl is substituted with 0-1 occurrence of C₁-C₆alkyl;

R^{6a} is selected from C₆-C₁₀aryl and C₃-C₈cycloalkyl;

R^{6b} is selected from halo, C₁-C₆haloalkyl, C₁-C₆haloalkoxy, and C₁-C₆alkyl;

R⁷ is selected from hydrogen and C₁-C₆alkyl;

R⁸ is selected from hydrogen and C₁-C₆alkyl;

or

R⁷ and R⁸ together with the nitrogen atom to which they are attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O, and S;


n is 0, 1, 2, 3 or 4;

m is 0, 1 or 2; and

p is 0 or 1.

2. The compound according to claim 1 or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein R^x is selected from hydrogen, C₁-C₆alkyl, and halo.

3. The compound according to claim 1 having Formula (I') or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein:



— is a single bond or a double bond;

X is selected from CH, CF, and N;

R' is selected from hydrogen and C₁-C₆alkyl;

R¹ is selected from hydrogen and C₁-C₆alkyl;

each R² is independently selected from C₁-C₆alkyl, C₁-C₆haloalkyl, halo, and oxo, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{2a}; or 2 R² on non-adjacent carbon atoms together with the non-adjacent carbon atoms to which they are attached form a bridging ring;

R^{2a} is selected from C₁-C₆alkoxy and hydroxyl;

R³ is selected from hydrogen, C₁-C₈alkyl, C₂-C₆alkenyl, —SO₂R⁴, C₁-C₆haloalkyl, —C(=O)—O—(R⁵) and

—C(=O)—(R⁶), wherein the C₁-C₈alkyl and C₁-C₆haloalkyl are independently substituted with 0-3 occurrences of R^{3a};

or

R³ together with the nitrogen atom to which it is attached and R² together with the carbon atom to which it is attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O and S;

each R^{3a} is independently selected from C₃-C₁₀cycloalkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, a 5- to 10-membered heteroaryl comprising 1-4 heteroatoms independently selected from N, O, and S, C₆-C₁₀aryl, C₁-C₆alkoxyl, hydroxyl, and —C(=O)—NR⁷R⁸, wherein the C₃-C₁₀cycloalkyl, 4- to 6-membered heterocyclyl, 5- to 10-membered heteroaryl and C₆-C₁₀aryl are substituted with 0-4 occurrences of R^{3b};

each R^{3b} is independently selected from C₁-C₆alkoxyl, halo, C₁-C₆haloalkyl, C₁-C₆haloalkoxyl, C₁-C₆alkyl, —CN, —SO₂NR⁷R⁸, —SO₂R⁴, and hydroxyl;

R⁴ is selected from C₃-C₈cycloalkyl, C₁-C₆alkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, and C₆-C₁₀aryl, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{4a};

R^{4a} is selected from C₃-C₈cycloalkyl, C₆-C₁₀aryl, and C₁-C₆alkoxyl;

R⁵ is selected from C₁-C₆alkyl C₃-C₈cycloalkyl, and C₆-C₁₀aryl;

R⁶ is selected from C₁-C₆alkyl, C₃-C₈cycloalkyl, and C₆-C₁₀aryl, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{6a} and the C₃-C₈cycloalkyl is substituted with 0-1 occurrence of R^{6b};

R^{6a} is selected from C₆-C₁₀aryl and C₃-C₈cycloalkyl;

R^{6b} is selected from halo, C₁-C₆haloalkyl, C₁-C₆haloalkoxyl, and C₁-C₆alkyl;

R⁷ is selected from hydrogen and C₁-C₆alkyl;

R⁸ is selected from hydrogen and C₁-C₆alkyl;

or


R⁷ and R⁸ together with the nitrogen atom to which they are attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O, and S;

n is 0, 1, 2, 3 or 4;

m is 0, 1 or 2; and

p is 0 or 1.

4. The compound according to claim 3 having a Formula (I) or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein:



X is selected from CH, CF, and N;

R' is selected from hydrogen and C₁-C₆alkyl;

R¹ is selected from hydrogen and C₁-C₆alkyl;

each R² is independently selected from C₁-C₆alkyl, C₁-C₆haloalkyl, halo, and oxo, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{2a}; or 2 R² on non-adjacent carbon atoms together with the non-adjacent carbon atoms to which they are attached form a bridging ring;

R^{2a} is selected from C₁-C₆alkoxyl and hydroxyl;

R³ is selected from hydrogen, C₁-C₈alkyl, C₂-C₆alkenyl, —SO₂R⁴, C₁-C₆haloalkyl, —C(=O)—O—(R⁵) and —C(=O)—(R⁶), wherein the C₁-C₈alkyl and C₁-C₆haloalkyl are independently substituted with 0-3 occurrences of R^{3a};

or

R³ together with the nitrogen atom to which it is attached and R² together with the carbon atom to which it is attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O and S;

each R^{3a} is independently selected from C₃-C₁₀cycloalkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, a 5- to 10-membered heteroaryl comprising 1-4 heteroatoms independently selected from N, O, and S, C₆-C₁₀aryl, C₁-C₆alkoxyl, hydroxyl, and —C(=O)—NR⁷R⁸, wherein the C₃-C₁₀cycloalkyl, 4- to 6-membered heterocyclyl, 5- to 10-membered heteroaryl and C₆-C₁₀aryl are substituted with 0-4 occurrences of R^{3b};

each R^{3b} is independently selected from C₁-C₆alkoxyl, halo, C₁-C₆haloalkyl, C₁-C₆haloalkoxyl, C₁-C₆alkyl, —CN, —SO₂NR⁷R⁸, —SO₂R⁴, and hydroxyl;

R⁴ is selected from C₃-C₈cycloalkyl, C₁-C₆alkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, and C₆-C₁₀aryl, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{4a};

R^{4a} is selected from C₃-C₈cycloalkyl, C₆-C₁₀aryl, and C₁-C₆alkoxyl;

R⁵ is selected from C₁-C₆alkyl C₃-C₈cycloalkyl, and C₆-C₁₀aryl;

R⁶ is selected from C₁-C₆alkyl, C₃-C₈cycloalkyl, and C₆-C₁₀aryl, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{6a} and the C₃-C₈cycloalkyl is substituted with 0-1 occurrence of R^{6b};

R^{6a} is selected from C₆-C₁₀aryl and C₃-C₈cycloalkyl;

R^{6b} is selected from halo, C₁-C₆haloalkyl, C₁-C₆haloalkoxyl, and C₁-C₆alkyl;

R⁷ is selected from hydrogen and C₁-C₆alkyl;

R⁸ is selected from hydrogen and C₁-C₆alkyl;

or

R⁷ and R⁸ together with the nitrogen atom to which they are attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O, and S;

n is 0, 1, 2, 3 or 4;

m is 0, 1 or 2; and

p is 0 or 1.

5. The compound according to claim 1 or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein

X is selected from CH, CF, and N;

R' is selected from hydrogen and C₁-C₃alkyl;

R¹ is selected from hydrogen and C₁-C₃alkyl;

each R² is independently selected from unsubstituted C₁-C₆alkyl, C₁-C₆haloalkyl and halo; or 2 R² on non-adjacent carbon atoms together with the non-adjacent carbon atoms to which they are attached form a bridging ring;

R³ is selected from hydrogen, C₁-C₈alkyl, C₂-C₆alkenyl, —SO₂R⁴, C₁-C₆haloalkyl, —C(=O)—O—(R⁵) and —C(=O)—(R⁶), wherein the C₁-C₈alkyl and C₁-C₆haloalkyl are independently substituted with 0-3 occurrences of R^{3a};

or

R³ together with the nitrogen atom to which it is attached and R² together with the carbon atom to which it is attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N and O;

each R^{3a} is independently selected from C₃-C₁₀cycloalkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, a 5- to 10-membered heteroaryl comprising 1-4 heteroatoms independently selected from N, O, and S, C₆-C₁₀aryl, C₁-C₆alkoxy, hydroxyl, and —C(=O)—NR⁷R⁸, wherein the C₃-C₁₀cycloalkyl, 4- to 6-membered heterocyclyl, 5- to 10-membered heteroaryl and C₆-C₁₀aryl are substituted with 0-4 occurrences of R^{3b};

each R^{3b} is independently selected from C₁-C₆alkoxy, halo, C₁-C₆haloalkyl, C₁-C₆haloalkoxy, C₁-C₆alkyl, —CN, —SO₂NR⁷R⁸, —SO₂R⁴, and hydroxyl;

R⁴ is selected from C₃-C₈cycloalkyl, C₁-C₆alkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, and C₆-C₁₀aryl, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{4a};

R^{4a} is selected from C₃-C₈cycloalkyl, C₆-C₁₀aryl, and C₁-C₆alkoxy;

R⁵ is selected from C₁-C₆alkyl C₃-C₈cycloalkyl, and C₆-C₁₀aryl;

R⁶ is selected from C₁-C₆alkyl, C₃-C₈cycloalkyl, and C₆-C₁₀aryl, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{6a} and the C₃-C₈cycloalkyl is substituted with 0-1 occurrence of R^{6b};

R^{6a} is selected from C₆-C₁₀aryl and C₃-C₈cycloalkyl;

R^{6b} is selected from chloro, fluoro, C₁-C₆haloalkyl, C₁-C₆haloalkoxy, and C₁-C₆alkyl;

R⁷ is selected from hydrogen and C₁-C₆alkyl;

R⁸ is selected from hydrogen and C₁-C₆alkyl;

or

R⁷ and R⁸ together with the nitrogen atom to which they are attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O, and S;

n is 0, 1, 2 or 3;

m is 0, 1 or 2; and

p is 0 or 1.

6. The compound according to claim 1 or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein

X is selected from CH and N;

R' is selected from hydrogen and methyl;

R¹ is selected from hydrogen and methyl;

each R² is independently selected from unsubstituted C₁-C₆alkyl and halo; or 2 R² on non-adjacent carbon atoms together with the non-adjacent carbon atoms to which they are attached form a C₁-C₃alkylene bridging ring;

R³ is selected from hydrogen, C₁-C₈alkyl, C₂-C₆alkenyl, —SO₂R⁴, C₁-C₆haloalkyl, —C(=O)—O—(R⁵) and —C(=O)—(R⁶), wherein the C₁-C₈alkyl and C₁-C₆haloalkyl are independently substituted with 0-3 occurrences of R^{3a};

or

R³ together with the nitrogen atom to which it is attached and R² together with the carbon atom to which it is attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional O heteroatom;

each R^{3a} is independently selected from C₃-C₁₀cycloalkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, a 5- to 10-membered heteroaryl comprising 1-4 heteroatoms independently selected from N, O, and S and phenyl, wherein the C₃-C₁₀cycloalkyl, 4- to 6-membered heterocyclyl, 5- to 10-membered heteroaryl and phenyl are substituted with 0-4 occurrences of R^{3b};

each R^{3b} is independently selected from C₁-C₆alkoxy, halo, C₁-C₆haloalkyl, C₁-C₆haloalkoxy, C₁-C₆alkyl, —CN, —SO₂NR⁷R⁸, —SO₂R⁴, and hydroxyl;

R⁴ is selected from C₃-C₈cycloalkyl, C₁-C₆alkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, and C₆-C₁₀aryl, wherein the C₁-C₆alkyl is substituted with 1 occurrence of R^{4a};

R^{4a} is selected from C₃-C₈cycloalkyl, C₆-C₁₀aryl, and C₁-C₆alkoxy;

R⁵ is selected from C₁-C₆alkyl C₃-C₈cycloalkyl, and C₆-C₁₀aryl;

R⁶ is selected from C₁-C₆alkyl, C₃-C₈cycloalkyl, and C₆-C₁₀aryl, wherein the C₁-C₆alkyl is substituted with 0-1 occurrence of R^{6a} and the C₃-C₈cycloalkyl is substituted with 0-1 occurrence of R^{6b};

R^{6a} is selected from C₆-C₁₀aryl and C₃-C₈cycloalkyl;

R^{6b} is selected from chloro, fluoro, C₁-C₆haloalkyl, C₁-C₆haloalkoxy, and C₁-C₆alkyl;

R⁷ is selected from hydrogen and C₁-C₆alkyl;

R⁸ is selected from hydrogen and C₁-C₆alkyl;

or

R⁷ and R⁸ together with the nitrogen atom to which they are attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O, and S;

n is 0, 1, 2 or 3;

m is 0, 1 or 2; and

p is 0 or 1.

7. The compound according to claim 1 or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein

X is selected from CH and N;

R' is hydrogen;

R¹ is hydrogen;

each R² is independently selected from unsubstituted C₁-C₆alkyl and fluoro; or 2 R² on non-adjacent carbon

atoms together with the non-adjacent carbon atoms to which they are attached form a C₁-C₃alkylene bridging ring;

R³ is selected from C₁-C₈alkyl, C₂-C₆alkenyl, —SO₂R⁴ and C₁-C₆haloalkyl, wherein the C₁-C₈alkyl is substituted with 0-2 occurrences of R^{3a} and the C₁-C₆haloalkyl is substituted with 0-1 occurrence of R^{3a};

each R^{3a} is independently selected from C₃-C₁₀cycloalkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N and O, a 5- to 6-membered heteroaryl comprising 1-3 heteroatoms independently selected from N, O, and S and phenyl, wherein the C₃-C₁₀cycloalkyl, 4- to 6-membered heterocyclyl, 5- to 6-membered heteroaryl and phenyl are substituted with 0-4 occurrences of R^{3b};

each R^{3b} is independently selected from halo, C₁-C₆haloalkyl, C₁-C₆haloalkoxyl, C₁-C₆alkyl, and hydroxyl;

R⁴ is selected from C₃-C₈cycloalkyl, C₁-C₆alkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, and C₆-C₁₀aryl, wherein the C₁-C₆alkyl is substituted with 1 occurrence of R^{4a};

R^{4a} is selected from C₃-C₈cycloalkyl, C₆-C₁₀aryl, and C₁-C₆alkoxyl;

n is 0, 1, 2 or 3;

m is 1 or 2; and

p is 0 or 1.

8. The compound according to claim 1 or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein

X is selected from CH and N;

R' is hydrogen;

R¹ is hydrogen;

each R² is independently selected from unsubstituted C₁-C₆alkyl; or 2 R² on non-adjacent carbon atoms together with the non-adjacent carbon atoms to which they are attached form a C₁-C₃alkylene bridging ring;

R³ is selected from C₁-C₈alkyl, C₂-C₆alkenyl, —SO₂R⁴ and unsubstituted C₁-C₆haloalkyl, wherein the C₁-C₈alkyl is substituted with 0-2 occurrences of R^{3a};

each R^{3a} is independently selected from C₃-C₁₀cycloalkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N and O, a 5- to 6-membered heteroaryl comprising 1-3 heteroatoms independently selected from N, O, and S and phenyl, wherein the C₃-C₁₀cycloalkyl, 4- to 6-membered heterocyclyl, 5- to 6-membered heteroaryl and phenyl are substituted with 0-3 occurrences of R^{3b};

each R^{3b} is independently selected from halo, C₁-C₆haloalkyl, C₁-C₆haloalkoxyl, C₁-C₆alkyl, and hydroxyl;

R⁴ is selected from C₃-C₈cycloalkyl, C₁-C₆alkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, and C₆-C₁₀aryl, wherein the C₁-C₆alkyl is substituted with 1 occurrence of R^{4a};

R^{4a} is selected from C₃-C₈cycloalkyl, C₆-C₁₀aryl, and C₁-C₆alkoxyl;

n is 0, 1 or 2;

m is 1 or 2; and

p is 1.

9. The compound according to claim 1 or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein

X is selected from CH and N;

R' is hydrogen;

R¹ is hydrogen;

each R² is independently selected from unsubstituted C₁-C₆alkyl;

R³ is selected from C₁-C₆alkyl, C₂-C₆alkenyl, —SO₂R⁴ and unsubstituted C₁-C₆haloalkyl, wherein the C₁-C₆alkyl is substituted with 0-2 occurrences of R^{3a};

each R^{3a} is independently selected from C₃-C₁₀cycloalkyl, a 4- to 6-membered heterocyclyl comprising 1 O heteroatom, a 6-membered heteroaryl comprising 1-2 N heteroatoms and phenyl, wherein the C₃-C₁₀cycloalkyl, 4- to 6-membered heterocyclyl, 6-membered heteroaryl and phenyl are substituted with 0-2 occurrences of R^{3b};

each R^{3b} is independently selected from chloro, fluoro, C₁-C₆haloalkyl, C₁-C₆haloalkoxyl and C₁-C₆alkyl;

R⁴ is selected from C₃-C₈cycloalkyl, C₁-C₆alkyl, a 4- to 6-membered heterocyclyl comprising 1 O heteroatom and phenyl, wherein the C₁-C₆alkyl is substituted with 1 occurrence of R^{4a};

R^{4a} is selected from C₃-C₈cycloalkyl and phenyl;


n is 0, 1 or 2;

m is 1 or 2; and


p is 1.

10. The compound according to claim 1 or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein R³ is selected from C₁-C₆alkyl and —CH₂—R^{3a}.

11. The compound according to claim 1 or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, having a Formula (Ib):



12. The compound according to claim 1 or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, having a Formula (Ic), wherein:



X is selected from CH, CF and N;

R^{2b} is selected from hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, and halo, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{2a} ;

R^{2c} is selected from hydrogen and C_1 - C_6 alkyl, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{2a} ;

or R^{2b} and R^{2c} together with the carbon atoms to which they are attached form an oxo group;

each of R^{2d} and R^{2e} is independently selected from hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, halo, and oxo, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{2a} ;

R^{2f} is hydrogen;

or R^{2b} and R^{2e} or R^{2b} and R^{2f} together with the carbon atoms to which they are attached form a bridging ring;

R^{2a} is selected from C_1 - C_6 alkoxy and hydroxyl;

R^3 is defined according to any one of the preceding claims; and

m is 1 or 2.

13. The compound according to claim 12, or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein

X is selected from CH and N;

R^{2b} is selected from hydrogen, C_1 - C_3 alkyl, C_1 - C_3 haloalkyl, and halo, wherein the C_1 - C_3 alkyl is substituted with 0-1 occurrence of R^{2a} ;

R^{2c} is selected from hydrogen and C_1 - C_3 alkyl, wherein the C_1 - C_3 alkyl is substituted with 0-1 occurrence of R^{2a} ;

or R^{2b} and R^{2c} together with the carbon atoms to which they are attached form an oxo group;

each of R^{2d} and R^{2e} is independently selected from hydrogen, C_1 - C_3 alkyl, C_1 - C_3 haloalkyl, halo, and oxo, wherein the C_1 - C_3 alkyl is substituted with 0-1 occurrence of R^{2a} ;

R^{2f} is hydrogen;

or R^{2b} and R^{2e} or R^{2b} and R^{2f} together with the carbon atoms to which they are attached form a bridging ring;

R^{2a} is selected from C_1 - C_6 alkoxy and hydroxyl;

R^3 is selected from C_1 - C_8 alkyl, C_2 - C_6 alkenyl, $-\text{SO}_2\text{R}^4$, C_1 - C_6 haloalkyl, $-\text{C}(=\text{O})-\text{O}-(\text{R}^5)$ and $-\text{C}(=\text{O})-(\text{R}^6)$, wherein the C_1 - C_8 alkyl and C_1 - C_6 haloalkyl are independently substituted with 0-3 occurrences of R^{3a} ;

each R^{3a} is independently selected from C_3 - C_{10} cycloalkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, a 5- to 10-membered heteroaryl comprising 1-4 heteroatoms independently selected from N, O, and S, C_6 - C_{10} aryl, C_1 - C_6 alkoxy, hydroxyl, and $-\text{C}(=\text{O})-\text{NR}^7\text{R}^8$, wherein the C_3 - C_{10} cycloalkyl, 4- to 6-membered heterocyclyl, 5- to 10-membered heteroaryl and C_6 - C_{10} aryl are substituted with 0-4 occurrences of R^{3b} ;

each R^{3b} is independently selected from C_1 - C_6 alkoxy, halo, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkoxy, C_1 - C_6 alkyl, $-\text{CN}$, $-\text{SO}_2\text{NR}^7\text{R}^8$, $-\text{SO}_2\text{R}^4$, and hydroxyl;

R^4 is selected from C_3 - C_8 cycloalkyl, C_1 - C_6 alkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, and C_6 - C_{10} aryl, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{4a} ;

R^{4a} is selected from C_3 - C_8 cycloalkyl, C_6 - C_{10} aryl, and C_1 - C_6 alkoxy;

R^5 is selected from C_1 - C_6 alkyl and C_6 - C_{10} aryl;

R^6 is selected from C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, and C_6 - C_{10} aryl, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{6a} and the C_3 - C_8 cycloalkyl is substituted with 0-1 occurrence of R^{6b} ;

R^{6a} is selected from C_6 - C_{10} aryl and C_3 - C_8 cycloalkyl;

R^{6b} is selected from halo, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkoxy, and C_1 - C_6 alkyl;

R^7 is selected from hydrogen and C_1 - C_6 alkyl;

R^8 is selected from hydrogen and C_1 - C_6 alkyl;

or

R^7 and R^8 together with the nitrogen atom to which they are attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O, and S; and

m is 1 or 2.

14. The compound according to claim 12 or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein

X is selected from CH and N;

each of R^{2b} , R^{2c} , R^{2d} and R^{2e} is independently selected from hydrogen and unsubstituted C_1 - C_3 alkyl;

R^{2f} is hydrogen;

or R^{2b} and R^{2e} or R^{2b} and R^{2f} together with the carbon atoms to which they are attached form a C_1 - C_3 alkylene bridging ring;

R^3 is selected from C_1 - C_8 alkyl, C_2 - C_6 alkenyl, $-\text{SO}_2\text{R}^4$, C_1 - C_6 haloalkyl, wherein the C_1 - C_8 alkyl and C_1 - C_6 haloalkyl are independently substituted with 0-3 occurrences of R^{3a} ;

each R^{3a} is independently selected from C_3 - C_{10} cycloalkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, a 5- to 10-membered heteroaryl comprising 1-4 heteroatoms independently selected from N, O, and S, C_6 - C_{10} aryl, C_1 - C_6 alkoxy, hydroxyl, and $-\text{C}(=\text{O})-\text{NR}^7\text{R}^8$, wherein the C_3 - C_{10} cycloalkyl, 4- to 6-membered heterocyclyl, 5- to 10-membered heteroaryl and C_6 - C_{10} aryl are substituted with 0-4 occurrences of R^{3b} ;

each R^{3b} is independently selected from C_1 - C_6 alkoxy, halo, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkoxy, C_1 - C_6 alkyl, $-\text{CN}$, $-\text{SO}_2\text{NR}^7\text{R}^8$, $-\text{SO}_2\text{R}^4$, and hydroxyl;

R^4 is selected from C_3 - C_8 cycloalkyl, C_1 - C_6 alkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, and C_6 - C_{10} aryl, wherein the C_1 - C_6 alkyl is substituted with 0-1 occurrence of R^{4a} ;

R^{4a} is selected from C_3 - C_8 cycloalkyl, C_6 - C_{10} aryl, and C_1 - C_6 alkoxy;

R^7 is selected from hydrogen and C_1 - C_6 alkyl;

R^8 is selected from hydrogen and C_1 - C_6 alkyl;

or

R^7 and R^8 together with the nitrogen atom to which they are attached form a 5- or 6-membered heterocyclyl comprising 0-1 additional heteroatom selected from N, O, and S; and

m is 1 or 2.

15. The compound according to claim **12** or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein

X is selected from CH and N;

each of R^{2b} , R^{2c} , R^{2d} and R^{2e} is independently selected from hydrogen and unsubstituted C_1 - C_3 alkyl;

R^{2f} is hydrogen;

R^3 is selected from C_1 - C_8 alkyl, C_2 - C_6 alkenyl, $-\text{SO}_2R^4$, and C_1 - C_6 haloalkyl, wherein the C_1 - C_8 alkyl and C_1 - C_6 haloalkyl are independently substituted with 0-3 occurrences of R^{3a} ;

each R^{3a} is independently selected from C_3 - C_{10} cycloalkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, a 5- to 10-membered heteroaryl comprising 1-4 heteroatoms independently selected from N, O, and S and phenyl, wherein the C_3 - C_{10} cycloalkyl, 4- to 6-membered heterocyclyl, 5- to 10-membered heteroaryl and phenyl are substituted with 0-4 occurrences of R^{3b} ;


each R^{3b} is independently selected from C_1 - C_6 alkoxyl, halo, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkoxyl, C_1 - C_6 alkyl, and hydroxyl;

R^4 is selected from C_3 - C_8 cycloalkyl, C_1 - C_6 alkyl, a 4- to 6-membered heterocyclyl comprising 1-2 heteroatoms independently selected from N, O and S, and C_6 - C_{10} aryl, wherein the C_1 - C_6 alkyl is substituted with 1 occurrence of R^{4a} ;

R^{4a} is selected from C_3 - C_8 cycloalkyl, C_6 - C_{10} aryl, and C_1 - C_6 alkoxy; and

m is 1.

16. The compound according to claim **1** or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, having a Formula (Id),




wherein,

X is selected from N and CH;

R^{2b} , R^{2c} and R^{2e} are defined according to any one of claims **11** to **14**, e.g., R^{2b} is C_1 - C_3 alkyl and R^{2c} and R^{2e} are both hydrogen; and

R^3 is defined according to any one of the preceding claims.

17. The compound according to claim **1** or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, having a Formula (Ie),



wherein,

X is selected from N and CH;

R^{2b} , R^{2c} and R^{2e} are defined according to any one of claims **11** to **14**, e.g., R^{2b} is C_1 - C_3 alkyl and R^{2c} and R^{2e} are both hydrogen; and

R^3 is defined according to any one of the preceding claims.

18. The compound according to claim **1** or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein X is CH.

19. The compound according to claim **1** or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof, wherein R^2 is unsubstituted C_1 - C_6 alkyl and n is 1.

20. A method of treating a disease or disorder that is affected by the modulation of WIZ protein levels, comprising administering to a patient in need thereof a compound of claim **1** or a pharmaceutically acceptable salt, hydrate, solvate, prodrug, stereoisomer, or tautomer thereof.

* * * * *