

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent Application Publication

20250281448

Kind Code

A1

Publication Date

September 11, 2025

Inventor(s)

Druzgala; Pascal et al.

Cardioversion Methods Using Budiodarone

Abstract

Disclosed are cardioversion methods for patients with either paroxysmal or persistent AFib which methods use budiodarone or a pharmaceutical composition comprising budiodarone.

Inventors: Druzgala; Pascal (Santa Rosa, CA), Milner; Peter Gerard (Mountain View, CA)

Applicant: Xyra, LLC (Santa Rosa, CA)

Family ID: 92141868

Appl. No.: 19/056381

Filed: February 18, 2025

Related U.S. Application Data

parent US continuation 18351261 20230712 parent-grant-document US 12257228 child US 19056381

us-provisional-application US 63525014 20230705

Publication Classification

Int. Cl.: A61K31/343 (20060101); A61K31/4439 (20060101); A61K31/444 (20060101); A61K31/4545 (20060101); A61K31/5377 (20060101); A61P9/06 (20060101)

U.S. Cl.:

CPC A61K31/343 (20130101); A61K31/4439 (20130101); A61K31/444 (20130101); A61K31/4545 (20130101); A61K31/5377 (20130101); A61P9/06 (20180101);

Background/Summary

CROSS-REFERENCE TO RELATED APPLICATIONS [0001] This application is a continuation application of, and claims priority to, U.S. application Ser. No. 18,351,261, filed Jul. 12, 2023, now allowed, which is a non-provisional application of, and claims priority to U.S. Provisional Application 63/525,014 filed Jul. 5, 2023, all of which are incorporated herein by reference.

FIELD

[0002] This disclosure provides for methods to cardiovert patients with either paroxysmal or persistent atrial fibrillation (AFib) back to sinus rhythm which methods use a pharmaceutical composition comprising budiadarone.

BACKGROUND

[0003] Patients suffering from paroxysmal or persistent AFib experiencing a prolonged episode of AFib are at elevated risk of a stroke and/or congestive heart failure. These risks increase as the duration of the episode extends. Patients with symptomatic paroxysmal or persistent AFib are significantly more likely to recognize their condition whereas asymptomatic patients on a heart rhythm monitor that provides an alarm-based warning of prolonged AFib will become aware of their condition once the alarm has been activated. In either case, many such patients presenting themselves at a hospital emergency room or are in a critical care unit with such an extended episode of AFib are treated by having a cardioversion procedure typically by stopping the heart and restarting it in order to restore sinus rhythm. Currently, there are from about 70,000 to perhaps as high as 100,000 or more cardioversions done each year with the number increasing as the population ages. These numbers represent a fraction of patients who present themselves in emergency rooms with AFib. One reference suggested that about 25% of patients presenting at the emergency room with AFib were treated with cardioversion. Gulizia et al., *Europace*. 2019; 21 (2): 230-238. This suggests that some of the non-treated patients were contra-indicated for this procedure due to co-morbidities and some had their sinus rhythm restored without cardioversion.

[0004] The method of choice for cardioversion is electrical cardioversion, which relies upon stopping and restarting the heart to restore sinus rhythm. However, this procedure comes with several shortcomings. From the patient's perspective, there is a risk of dying during or within a few months after this procedure due to, for example, the release of blood clots from the heart during electrical cardioversion. From the clinician's perspective, electrical cardioversions require at least one or more nurses, an anesthesiologist, a cardiologist, the use of an anti-coagulant, and the like all at a significant cost to the health care system. From a cost perspective, cardioversion can cost thousands of dollars which, given the number of cardioversions conducted annually, imposes a significant cost burden on the healthcare system, particularly when the costs for treating patients who have a blood clot that releases during or shortly after the procedure or are included.

SUMMARY

[0005] Disclosed are methods to cardiovert a patient with an extended episode of paroxysmal or persistent AFib that utilizes a single high dose of budiadarone or a pharmaceutically acceptable salt thereof to effect cardioversion of the AFib sinus rhythm in said patient. The single high dose of budiadarone or a pharmaceutically acceptable salt thereof is preferably administered orally but other routes of administration are contemplated herein including intravenous, rectal, and pulmonary administration. The patient undergoing cardioversion is first deemed suitable for the procedure.

[0006] In one embodiment, there is provided a method for cardioverting a patient with paroxysmal or persistent AFib suffering from an extended episode of AFib, said method comprising: a) evaluating the patient to confirm that the patient has paroxysmal or persistent AFib and is suitable for cardioversion; b) administering budiadarone or a pharmaceutically acceptable salt thereof to said patient in an amount sufficient to effect cardioversion of said AFib back to sinus rhythm in said patient; and c) monitoring said patient subsequent to said administration to confirm that the patient is responsive to cardioversion.

[0007] In another embodiment, a method is provided for cardioverting a patient with paroxysmal or

persistent AFib suffering from an extended episode of AFib and deemed suitable for cardioversion, said method comprising: a) administering of budiiodarone or a pharmaceutically acceptable salt thereof to said patient in an amount sufficient to effect cardioversion of said AFib back to sinus rhythm, wherein the patient has been confirmed as having paroxysmal or persistent AFib as well as being deemed suitable for cardioversion; and b) monitoring said patient subsequent to said administration to confirm that the patient is responsive to cardioversion.

[0008] A further embodiment provides a method for cardioverting a patient with paroxysmal or persistent AFib suffering from an extended episode of AFib, said method comprising: a) administering a sufficient amount of budiiodarone or a pharmaceutically acceptable salt thereof to said patient to achieve a plasma concentration of at least about 13 ng/ml for a period of time to cardiovert said patient back to sinus rhythm, wherein the patient has been confirmed as having paroxysmal or persistent AFib and deemed suitable for cardioversion; and b) monitoring said patient subsequent to said administration to confirm that the patient is responsive to cardioversion

[0009] In another embodiment, a method is provided for cardioverting a patient with paroxysmal or persistent AFib suffering from an extended episode of AFib and deemed suitable for cardioversion, said method comprising: [0010] a) administering budiiodarone or a pharmaceutically acceptable salt thereof to said patient in an amount sufficient to effect cardioversion of said AFib back to sinus rhythm, wherein the patient has been confirmed as having paroxysmal or persistent AFib; and

[0011] b) monitoring said patient subsequent to said administration to confirm that the patient is responsive to cardioversion.

[0012] A further embodiment provides a method for cardioverting a patient with paroxysmal or persistent AFib suffering from an extended episode of AFib, said method comprising: [0013] a) administering a sufficient amount of budiiodarone or a pharmaceutically acceptable salt thereof to said patient to achieve a plasma concentration of at least about 13 ng/ml for a period of time to cardiovert said patient back to sinus rhythm, wherein the patient has been confirmed as having paroxysmal or persistent AFib and deemed suitable for cardioversion; and [0014] b) monitoring said patient subsequent to said administration to confirm that the patient is responsive to cardioversion.

[0015] In one embodiment, the amount of budiiodarone or a pharmaceutically acceptable salt thereof administered to said patient is sufficient to provide for sustained plasma concentration of at least about 13 ng/ml for a period of at least about 3 hours, at least about 4 hours, at least about 5 hours, or at least about 6 hours post-administration of budiiodarone or a pharmaceutically acceptable salt thereof. Monitoring of the patient's heart rhythm is preferably continued during the entire period when the plasma concentration of budiiodarone or a pharmaceutically acceptable salt thereof is maintained at about 13 ng/mL. In a preferred embodiment, the plasma concentration of budiiodarone or a pharmaceutically acceptable salt thereof ranges from about 13 ng/mL up to a C.sub.max of about 350 ng/mL. In another preferred embodiment, the plasma concentration of budiiodarone or a pharmaceutically acceptable salt thereof ranges from about 50 ng/mL to about 350 ng/ml and any subrange of about 10 ng within said range.

[0016] In one embodiment, budiiodarone or a pharmaceutically acceptable salt thereof is administered orally, sublingually, intravenously (i.v. or IV), rectally, as an aerosol (pulmonary), or as an intranasal spray, or a combination thereof. In one preferred embodiment, budiiodarone or a pharmaceutically acceptable salt thereof is administered orally using capsules, pills, or tablets. Given the efficiency of delivery, the amounts required to provide the plasma levels as presented above or herein, the dosing of budiiodarone or a pharmaceutically acceptable salt thereof when given intravenously, sublingually, or in a spray may be in a lower dosage than that required for oral dosing.

[0017] As per the above, the methods described herein include the administration of budiiodarone or a pharmaceutically acceptable salt thereof to the patient in an amount sufficient to cardiovert the AFib back to sinus rhythm when the plasma concentration of budiiodarone or a pharmaceutically

acceptable salt thereof is maintained at a level above as defined herein. In most cases, it is contemplated that cardioversion will be achieved for responsive patients within about 90 minutes, preferably within about 60 minutes, and more preferably within about 45, or within about 30 minutes after administration of budiiodarone or a pharmaceutically acceptable salt thereof.

[0018] In general, the amount of budiiodarone or a pharmaceutically acceptable salt thereof used to cardiovert the patient will be a single bolus oral dose of from about 400 mg to about 1,200 mg and preferably from about 800 mg to 1,200 mg. Preferably, each of these doses of budiiodarone or a pharmaceutically acceptable salt thereof is administered as a single pill or capsule. In one preferred embodiment, the budiiodarone or a pharmaceutically acceptable salt thereof dose to be used is selected by the clinician to achieve a C.sub.max of at least about 50 ng/ml or more which, based on its half-life data, is coupled with retaining a therapeutic plasma concentration of budiiodarone of at least about 13 ng/mL for about 6 hours in the patient. Preferred dosages of budiiodarone or a pharmaceutically acceptable salt thereof to achieve such prolonged retention of a therapeutic concentration include single doses of 800 mg, 1,000 mg, or 1,200 mg, or any dose between said 800 mg and 1,200 mg. Dosages of budiiodarone or a pharmaceutically acceptable salt thereof may also be about 50 mg or about 100 mg amount interval increases between 400 mg to 1,200 mg (e.g., 1,100 mg or 700 mg). Sublingual, intravenous, rectal, and spray dosage forms of budiiodarone would be less than 1,200 mg and are contemplated to range from about 10% to about 50% of the 1,200 mg dose used orally. Another aspect contemplates the administration of budiiodarone or a pharmaceutically acceptable salt thereof via one or more means to the patient, e.g., via IV and oral, via sublingual and oral, via sublingual and spray, via oral and spray, via oral and IV, via IV and spray, or via IV and sublingual. It is contemplated that intravenous administration may induce a more rapid conversion to a sinus rhythm in the patient in need thereof.

[0019] In one embodiment, the administration of budiiodarone or a pharmaceutically acceptable salt thereof is accompanied by the administration of a second drug such as a blood thinner (e.g., dabigatran, edoxaban, apixaban, or rivaroxaban—Xarelto®).

[0020] Also provided is a use of budiiodarone or a pharmaceutically acceptable salt thereof to cardiovert a patient with paroxysmal or persistent AFib suffering from an extended episode of AFib.

[0021] Another embodiment provides administering a dose of budiiodarone or a pharmaceutically acceptable salt thereof to a patient in need of cardioversion determined to have either paroxysmal or persistent AFib that is sufficient to produce an effective plasma level of >50 ng/ml in said patient within 60 min of administration, and maintaining a plasma level in said patient of >50 ng/ml for up to 6 hours thereby cardioverting a single sustained long episode of PAF, or a storm of long PAF episodes, or a persistent AFib episode back to normal sinus rhythm in said patient.

[0022] A further embodiment provides a use of budiiodarone or a pharmaceutically acceptable salt thereof to cardiovert a patient with paroxysmal or persistent AFib suffering from an extended episode of AFib.

[0023] Yet another embodiment provides an oral dose of budiiodarone or a pharmaceutically acceptable salt thereof for use in cardioverting a patient with paroxysmal or persistent AFib suffering from an extended AFib episode with an oral dose of budiiodarone or a pharmaceutically acceptable salt thereof comprising from about 1,000 mg to about 2,000 mg. The oral dose of budiiodarone or a pharmaceutically acceptable salt thereof can be delivered as a single bolus of about 1,000 mg, about 1,100 mg, about 1,200 mg, about 1,300 mg, about 1,400 mg, about 1,500 mg, about 1,600 mg, about 1,700 mg, about 1,800 mg, about 1,900 mg or about 2,000 and any 50 mg increment between 1,000 mg to about 2,000 mg. The dosage form of the administration of budiiodarone or a pharmaceutically acceptable salt thereof of 1,000 mg to about 2,000 mg as described in this paragraph can be as a pill, a tablet, or a capsule.

[0024] In another embodiment, the pharmaceutically acceptable salt of budiiodarone is budiiodarone tartrate.

Description

BRIEF DESCRIPTION OF DRAWINGS

[0025] The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawing(s) will be provided by the Office upon request and payment of the necessary fee as required under 37 C.F.R. § 1.84.

[0026] FIG. 1 illustrates the plasma concentration over time using different single-dose administrations of budiadarone. The data establishes that each single dose at or above 400 mg provided for a C.sub.max greater than 50 ng/mL and each maintained a concentration of greater than about 13 ng/ml for over 6 hours.

[0027] FIG. 2 illustrates the comparison of the plasma budiadarone concentrations between a fasting patient and a fed patient.

DETAILED DESCRIPTION

[0028] This disclosure is directed to methods for cardioverting AFib to sinus rhythm for patients with paroxysmal or persistent atrial fibrillation (AFib) and who are experiencing an extended period of AFib. The methods described herein employ budiadarone to cardiovert these patients. Terms that are not defined are given their definition in context or are given their medically acceptable definition.

[0029] The terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting. As used herein, the singular forms “a”, “an”, and “the” are intended to include the plural forms as well, unless the context clearly indicates otherwise.

Definitions

[0030] As used herein, the term “optional” or “optionally” means that the subsequently described event or circumstance may or may not occur and that the description includes instances where the event or circumstance occurs and instances where it does not.

[0031] As used herein, the term “about” when used before a numerical designation, e.g., temperature, time, amount, concentration, and such other, including a range, indicates approximations that may vary by (+) or (−) 15%, 10%, 5%, 1%, or any subrange and/or value therebetween. Preferably, the term “about” when used with regard to a dose amount means that the dose may vary by +/-10%.

[0032] As used herein, the term “comprising” or “comprises” is intended to mean that the compositions or methods include the recited elements, but do not exclude others.

[0033] As used herein, the term “consisting essentially of” when used to define compositions and methods, shall mean excluding other elements of any essential significance to the stated purpose of the combination. Thus, a composition consisting essentially of the elements as defined herein, or a method consisting essentially of the protocol defined herein, would not exclude other components that do not materially affect the basic and novel characteristic(s) of the claimed subject matter.

[0034] As used herein, the term “consisting of” shall mean excluding more than trace elements of other ingredients or substantial method steps. Embodiments defined by each of these transition terms are within the scope of this disclosure.

[0035] As used herein, the term “AFib” or “atrial fibrillation” refers to all variants of atrial fibrillation except permanent AFib. Such variants include but are not limited to paroxysmal AFib, persistent AFib, as well as paroxysmal AFib and persistent AFib with low CHA.sub.2DS.sub.2-VASc scores (2 or less) and high CHA.sub.2DS.sub.2-VASc scores (3 or higher).

CHA.sub.2DS.sub.2-VASc scores stand for congestive heart failure (C-1 pt), hypertension (H-1 pt), age ≥ 75 (A-2 pts), diabetes (D-1 pt), stroke (S-2 pt), vascular disease (V-1 pt), age 65 to 74 (A-1 pt), and sex category (Sc-1 pt for female) and are clinical prediction rules for estimating the risk of stroke in people with non-rheumatic atrial fibrillation (AF). A CHA.sub.2DS.sub.2-VASc score comprises 1 point or 2 points for each of the constituent risk factors, as indicated above

[0036] As used herein, the term “extended duration of AFib” relates to the length of time an episode of AFib lasts in a patient. This length relates to the risk of stroke and/or heart failure. As is apparent, the longer the length of time a patient is in AFib, the higher the risk becomes. So, in one embodiment, an extended duration of AFib is any AFib episode lasting over at least about 30 minutes or at least about 1 hour or at least 5 hours, provided that the patient is not diagnosed as having permanent AFib.

[0037] As used herein, the term “paroxysmal AFib” refers to sporadic and intermittent episodes of AFib that last no more than 7 days before resolving back into sinus rhythm. Typically, but not necessarily, patients with de minimis AFib are also characterized as having paroxysmal AFib.

[0038] As used herein, the term “persistent AFib” refers to sporadic and intermittent episodes of AFib that last more than 7 days and may not resolve back into sinus rhythm without medical intervention (e.g., cardioversion).

[0039] As used herein, the term “budiodarone” refers to (S)-sec-butyl2-(3-(4-(2-(diethyl amino)ethoxy)-3,5-diiodobenzoyl)benzofuran-2-yl) acetate as well as pharmaceutically acceptable salts thereof. In some cases, budiodarone tartrate salt is referred to as ATI-2042 or just “2042”.

Budiodarone, as the free base, is represented by the following formula:

##STR00001##

Pharmaceutically acceptable salts of the free base are included in the term “budiodarone”) and include all approved pharmaceutically acceptable salts. Inclusive of acceptable salts of budiodarone are budiodarone tartrate or budiodarone citrate. In an embodiment, the pharmaceutically acceptable salt may be a polycarboxylic salt. As is appreciated in the art, the salt disassociates from the free base in vivo. In addition, the major metabolite of budiodarone is the deacylated compound—the free acid. Accordingly, in calculating the serum plasma level of budiodarone, the molecular weight of the free acid and the free base is used to determine molarity and concentration. In addition, when a salt other than the tartrate salt is administered, the dose of budiodarone tartrate salt that is used herein will necessarily be changed to reflect the molecular weight change due to the different salt.

[0040] Budiodarone has a combined mode of action and has been shown to reduce episodes of long-duration AFib without a significant increase in the QT interval. Budiodarone has a pattern of cardiac ion channel effects that are like amiodarone by design with enhanced late Na^{sup.}+ channel blockade. Budiodarone's deliberately altered metabolism results in a much shorter half-life than amiodarone, however, enabling the avoidance of accumulation-related toxicities seen with amiodarone and complete inactivation and elimination from the body within hours to days of discontinuation.

[0041] As used herein, a “single high dose” refers to a dose of budiodarone that ranges from about 400 mg to about 1,200 mg which is administered in one or multiple dosing units or in a dosage form sufficient to achieve at least about 13 ng/ml budiodarone in a patient in need thereof. As noted above, a dosage form sufficient to achieve at least about 13 ng/ml budiodarone in a patient in need thereof could be less than about 1,200 mg when administered via inhalation, sublingually, or intravenously than when administered orally. A combination of these administration methods may be utilized and are contemplated when using the term “bolus”. When multiple dosing units are employed, they can be administered concurrently with the other units such that the entire dose is administered within about 30 minutes and preferably within about 10 minutes. It is further noted that the patient may be asked to eat food, possibly and preferably fatty food, at the time of ingesting the bolus of budiodarone to achieve a more rapid intake of the drug into the system.

[0042] As used herein, “C.sub.max” refers to the maximum plasma concentration of budiodarone after the administration of a single dose to a patient.

[0043] As used herein, a “suitable patient” or “patient” is a person who has been experiencing paroxysmal or persistent AFib for about 6 hours or less, or is a patient who has been experiencing paroxysmal or persistent AFib but has been determined to not have a blood clot through the use of a transesophageal echocardiogram or other means of confirming the absence of a blood clot in a

patient, or the patient is known to have persistent or paroxysmal AFib and is on a blood thinner.

Methodology

[0044] The methods recited herein utilize a single high dose or bolus of budiadarone to cardiovert the patient suffering from an extended duration of AFib back to sinus rhythm. These methods are utilized with patients diagnosed with paroxysmal or persistent AFib. Patients with permanent AFib are not responsive to budiadarone and should be excluded by the attending physician from treatment with this drug. For those patients diagnosed with paroxysmal or persistent AFib, having an extended duration of AFib, and who present themselves to the emergency room or other facility capable of properly treating the patient, the clinician or caregiver confirms that the heart rhythm irregularity is AFib and not another known irregular heart rhythm and also determines that the patient is suitable for cardioversion.

[0045] Once confirmed, the attending clinician administers a single bolus (high) dose of budiadarone to the patient. The dose administered is sufficient to provide for a plasma concentration of budiadarone of at least about 13 ng/mL is maintained for at least about 3 hours and preferably for at least about 4 hours, or at least about 5 hours, or at least about 6 hours. In one preferred embodiment, the amount of budiadarone administered to the patient ranges comprises a single bolus of budiadarone at about 800 mg, or at least about 1,000 mg, or at least about 1200 mg. In some embodiments, the use of a high single dose of budiadarone is employed to provide rapid uptake to a therapeutic concentration in vivo. When so used, the plasma concentration of such doses will reach a C.sub.max of about 350 ng/ml or less and will maintain at least about 40 ng/ml or about 50 ng/ml for about 5+ hours or, in some cases, at least about 6 hours (see, FIG. 1). Regardless, such high dosing levels ensure that the plasma concentration remains over about 13 ng/ml and, in fact, over about 40 ng/mL, for up to about 6 hours.

[0046] Once the bolus of budiadarone is administered, the patient remains under observation in the emergency room, hospital room, or another urgent care facility while monitoring heart rhythm for a sufficient period to determine if the patient is responsive to budiadarone. Once the heart is restored to sinus rhythm and the plasma concentration of budiadarone is reduced to a satisfactory level, in the opinion of the attending clinician, the patient can be discharged provided that there are no other co-morbidities requiring further observation of the patient. In one preferred embodiment, such a level is below about 20 ng/mL and preferably below about 15 ng/mL. In one preferred embodiment, the patient is required to remain under observation for a short period of time (e.g., about 30 minutes to about 6 hours) to ensure that sinus rhythm is retained.

[0047] For those patients who are responsive to said methods, the patient can be discharged with a wearable that records and optionally transmits the heart rhythm to ascertain the extent and duration of AFib. For those patients who were non-responsive to said methods, these patients can be treated with electrical cardioversion.

Pharmaceutical Compositions

[0048] The specific dosing of budiadarone is accomplished by any number of accepted modes of administration including oral, intravenous, rectal, and pulmonary delivery. Pharmaceutical compositions compatible with each such delivery are well known in the art as are the methods useful in such delivery. Preferably, oral delivery is preferred typically using tablets, pills, capsules, and the like. The particular form used for oral delivery is not critical.

[0049] Pharmaceutical dosage forms of budiadarone may be manufactured by any of the methods well-known in the art, such as conventional mixing, tableting, encapsulating, and the like. The compositions as disclosed herein can include one or more physiologically acceptable inactive ingredients that facilitate the processing of active molecules into preparations for pharmaceutical use.

[0050] The compositions can comprise the drug in combination with at least one pharmaceutically acceptable excipient. Acceptable excipients are non-toxic, aid administration, and do not adversely affect the therapeutic benefit of the claimed compounds. Such excipient may be any solid, liquid, or

semi-solid that is generally available to one of skill in the art.

[0051] Solid pharmaceutical excipients include starch, cellulose, talc, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, magnesium stearate, sodium stearate, glycerol monostearate, sodium chloride, dried skim milk and the like. Other suitable pharmaceutical excipients and their formulations are described in REMINGTON'S PHARMACEUTICAL SCIENCES, edited by E. W. Martin (Mack Publishing Company, 18th ed., 1990).

[0052] The compositions as disclosed herein may, if desired, be presented in a single-use package. Such a package may, for example, comprise metal or plastic foil, such as a blister pack, a vial, or any other type of containment. The pack or dispenser device may be accompanied by instructions for administration including, for example, instructions for use.

[0053] The amount of the drug in a formulation can vary depending on the number of subunits required for the daily or periodic dose of the drug. Typically, the formulation will contain, on a weight percent (wt. %) basis, from about 10 to 99 wt. % of the drug based on the total formulation, with the balance being one or more suitable pharmaceutical excipients. Preferably, the compound is present at a level of about 20 to 70 wt. %.

Examples

[0054] The present disclosure is further understood by reference to the following examples, which are intended to be purely exemplary of this invention. The present disclosure is not limited in scope by the exemplified embodiments, which are intended as illustrations of single aspects of this disclosure only. Any functionally equivalent methods are within the scope of this disclosure.

Various modifications of this invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description and accompanying figures. Such modifications fall within the scope of the appended claims. In these examples, the following terms are used herein and have the following meanings. If not defined, the abbreviation has its conventional medical meaning.

[0055] It is noted that budiodarone concentrations used in the examples below include both the free acid and the free base versions of budiodarone.

[0056] The following abbreviations used herein have the following meanings. If not defined, the abbreviation will have its accepted meaning:

TABLE-US-00001 AFib = Atrial Fibrillation b.i.d. or bid = twice a day Pk = pharmacokinetics Hrs = Hours mg = Milligrams mL = Milliliter ng = nanogram

Example 1—Determination of the Minimum Concentration of Budiodarone to Reduce the Duration of Episodes of AFib

[0057] In this example, patients were selected with long episodes of either paroxysmal or persistent AFib. Each of the patients was implanted with a pacemaker which monitored their heart rhythm. Patients were then evaluated for disease progression by monitoring their heart rhythm without budiodarone treatment in order to generate a baseline level of the duration of episodes of AFib. The baseline data was collected over two weeks. Patients were then given incremental doses of budiodarone at 200 mg b.i.d.; 400 mg b.i.d.; 600 mg b.i.d.; and 800 mg b.i.d. where each dose was maintained for 2 weeks for a total of 4 treatment periods. A two-week washout period was employed after the completion of the 800 mg b.i.d. treatment period, wherein a washout period is the time after the last treatment to avoid misinterpreting observations about study-related treatments that may have been due to prior therapies. Pacemaker data were downloaded on days 8 and 14 of each two-week cycle and averages for each test period are shown. Blood was collected during each cycle and tested for the concentration of budiodarone. The results of this test are as follows:

TABLE-US-00002 Dose Baseline 200 mg bid 400 mg bid 600 mg bid 800 mg bid washout Average
4.8 hrs 1.7 hrs 0.6 hrs 0.1 hrs 0.5 hrs 2.4 Episode duration Trough — 2.4 +/- 0.9 5.2 +/- 1.7 13.1
+/- 5.6 19.8 +/- 17.9 0.3 +/- 0.4 Pk levels ng/mL in plasma

Note that the duration of AFib in all dosing levels from 400 mg b.i.d. to 800 mg b.i.d. provided for

maximum duration AFib levels of less than 90 minutes and, in fact, less than 60 and less than 45 minutes. In the case of 600 mg provided b.i.d., the longest average duration of AFib was 6 minutes (less than 30 minutes). It is also noted that the trough Pk levels of budiodarone are predicated on dosing approximately 12 hours apart and are not relevant to the trough levels at 6 hours post-administration.

[0058] This data supports the underlying basis that the administration of a bolus of budiodarone would rapidly reach C.sub.max plasma concentrations of at least 50 ng/ml and maintain as much for a sufficient period of time to effect cardioversion of a patient's AFib back to sinus rhythm. Note also the rapid removal of budiodarone from the blood such that the trough concentration in the plasma at 2 weeks post 800 mg budiodarone administered b.i.d. was 0.3 ng/ml in plasma or about 1.5% of the trough concentration of 19.8 ng/ml of budiodarone at the 800 mg dose b.i.d. showing rapid washout of the drug once stopped.

Example 2—Determination of Budiodarone Concentrations in Plasma Over Time Using Different Dosing Amounts of the Drug

[0059] In this example, plasma concentrations of budiodarone were evaluated using different single-dosing amounts of the drug. Specifically, as illustrated in FIG. 1, budiodarone was administered to patients as a single dose at 50 mg, 100 mg, 200 mg, 400 mg, 600 mg, 800 mg, and 1000 mg. Each of the patients had blood draws taken and the concentration of budiodarone or its deacylated metabolite which is inactive. The resulting data was placed on a graph and the data for each concentration provided for both a C.sub.max for the dose employed as well as a trough value at 6 hours.

[0060] As to specifics, FIG. 1 establishes that the plasma concentration of a single dose of budiodarone at or above 400 mg provides for a C.sub.max greater than 50 ng/ml within 60 min or giving a single oral dose, and each maintained a plasma concentration of greater than about 13 ng/ml for over 6 hours. Still, further, a single dose of budiodarone at 800 mg or 1,000 mg will provide a plasma concentration of about 50 ng/ml at 6 hours and at least about 15 ng/ml at 8 hours.

[0061] The data also suggests that the use of a bolus of budiodarone such as 1,000 mg will provide for rapid uptake to a C.sub.max concentration of over 300 ng/ml in plasma which rapidly is eliminated to less than about 20 ng/ml in about 8 hours. In combination, the rapid uptake coupled with elimination in a short period of time renders such a bolus suitable for cardioversion as the high dosing of budiodarone will rapidly reach efficacious levels within 60 min of administration in vivo which is maintained at a plasma level above about 13 ng/ml for at least 6 hours followed by clearance from the body shortly thereafter. In combination, this data suggests that budiodarone will rapidly transition a long-duration episode of AFib to a significantly shorter episode or eliminates AFib, thereby cardioverting the patient into sinus rhythm within 90 minutes or less from the time the drug was administered.

[0062] As budiodarone has been established in clinical studies as being safe for in vivo administration, the use of budiodarone to cardiovert a patient is an alternative to electrical cardioversion and the corresponding issues associated therewith.

Example 3—The Impact of Food on the Concentration of Budiodarone

[0063] This example evaluates the impact on plasma budiodarone concentrations based on a fasting or fed patient. In this example, patients were either fasting or fed before the administration of budiodarone. After budiodarone administration, blood was collected from the patients, and the plasma concentration of budiodarone was determined. FIG. 2 illustrates that in the same dosing of budiodarone, the uptake of budiodarone increased as evidenced by the C.sub.max values. Note, however, that C.sub.max in the fed group is shifted to about an hour later in time than the C.sub.max in fasted patients. This suggests that fasting patients who are to undergo budiodarone cardioversion are preferably fed within about one hour (about 60 minutes) before the start of said cardioversion. This data shows that whether or not the drug is taken with food a single oral dose of 400-1,000 mg will achieve an effective plasma level of >50 ng/ml within 60 min and maintain that,

and higher levels, for up to 6 hours thereby cardioverting back to normal sinus rhythm a single of a sustained long episode of PAF (paroxysmal atrial fibrillation) or persistent, or a storm of long AFib episodes that caused symptoms that brought the patient to the ER.

Embodiments

[0064] The following embodiments are contemplated.

[0065] [1] A method for cardioverting a patient with paroxysmal or persistent AFib suffering from an extended episode of AFib and deemed suitable for cardioversion, said method comprising: a) administering budiolarone or a pharmaceutically acceptable salt thereof to said patient in an amount sufficient to effect cardioversion of said AFib back to sinus rhythm, wherein the patient has been confirmed as having paroxysmal or persistent AFib; and b) monitoring said patient subsequent to said administration to confirm that the patient is responsive to cardioversion.

[0066] [2] A method for cardioverting a patient with paroxysmal or persistent AFib suffering from an extended episode of AFib, said method comprising: a) administering a sufficient amount of budiolarone or a pharmaceutically acceptable salt thereof to said patient to achieve a plasma concentration of said budiolarone or the salt thereof at least about 13 ng/ml for a period of time to cardiovert said patient back to sinus rhythm, wherein the patient has been confirmed as having paroxysmal or persistent AFib and deemed suitable for cardioversion; and b) monitoring said patient subsequent to said administration to confirm that the patient is responsive to cardioversion.

[0067] [3] The method of any one of Embodiments [1] or [2], wherein the amount of budiolarone or a salt thereof administered to said patient is sufficient to provide for a plasma concentration of from about 13 ng/mL to about 350 ng/mL in said patient, provided that the plasma concentration of budiolarone remains greater than at least about 13 ng/ml for at least 6 hours post-administration.

[0068] [4] The method of any one of Embodiments [1] or [3], wherein the amount of budiolarone or a salt thereof administered to said patient is sufficient to provide a plasma concentration of budiolarone or a salt thereof in said patient to about 50 ng/ml to about 350 ng/ml for at least about 5 hours post-administration.

[0069] [5] The method of Embodiment [4], wherein the amount of budiolarone or a salt thereof administered to said patient is sufficient to provide a plasma concentration of budiolarone or a salt thereof in said patient of about 50 ng/mL to about 350 ng/ml for at least about 6 hours.

[0070] [6] The method of any one of Embodiments [1]-[5], wherein said administration of the budiolarone or a salt thereof is oral, sublingual, intravenous, rectal, or an aerosol.

[0071] [7] The method of any one of Embodiments [1]-[6], wherein budiolarone or a salt thereof is administered orally using capsules, pills, or tablets.

[0072] [8] The method of any one of Embodiments [1]-[7], wherein said dose of budiolarone or a salt thereof is administered as a single dose of about 400 mg to about 2,000 mg.

[0073] [9] The method of Embodiment [8], wherein said dose of budiolarone or a salt thereof is administered as a single dose of about 800 mg to about 1,200 mg.

[0074] [10] The method of Embodiment [8], wherein said dose of budiolarone or a salt thereof is administered as a single dose of about 1,000 mg.

[0075] [11] The method of Embodiment [8], wherein said dose of budiolarone or a salt thereof is administered as a single dose of about 1,200 mg.

[0076] [12] The method of any one of Embodiments [1]-[11], which further comprises administering a blood thinner with the budiolarone or salt thereof to said patient, wherein the blood thinner is dabigatran, edoxaban, apixaban, or rivaroxaban.

[0077] [13] The method of any one of Embodiments [1]-[12], wherein cardioversion of said patient is accomplished in about 90 minutes or less from administering budiolarone or a salt thereof.

[0078] [14] The method of any one of Embodiments [1]-[12], wherein cardioversion is accomplished in about 60 minutes or less from administering budiolarone or a salt thereof.

[0079] [15] The method of any one of Embodiments [1]-[12], wherein cardioversion is accomplished in about 45 minutes or less from administering budiolarone or a salt thereof.

[0080] [16] The method of any one of Embodiments [1]-[12], wherein cardioversion is accomplished in about 30 minutes or less from administering budiadarone or a salt thereof.

[0081] [17] The method of any one of Embodiments [1]-[12], wherein said patient is fed about 1 hour or less prior to administering budiadarone or a salt thereof.

[0082] [18] A kit comprising a single dose of budiadarone or a pharmaceutically effective salt thereof that is effective in cardioverting a patient and instructions for post-administration monitoring of the patient.

[0083] [19] Administering a dose of budiadarone or a pharmaceutically acceptable salt thereof to cardiovert a patient determined to have either paroxysmal or persistent Afib, wherein said administered dose produces an effective plasma level of >50 ng/ml in said patient within about 60 min of administration, and maintaining a plasma level in said patient of >50 ng/ml for up to about 6 hours thereby cardioverting a single sustained long episode of PAF, or a storm of long PAF episodes, or a persistent AFib episode that caused symptoms back to normal sinus rhythm in said patient.

[0084] [20] A use of budiadarone or a pharmaceutically acceptable salt thereof to cardiovert a patient with paroxysmal or persistent AFib suffering from an extended episode of AFib by administering a dosage of budiadarone according to any one of the methods of Embodiments [1]-[19].

[0085] [21] An oral dose of budiadarone or a pharmaceutically acceptable salt thereof of Embodiment [20] comprising from about 1,000 mg to about 2,000 mg in a single dosage unit.

[0086] [22] An oral dose of budiadarone or a pharmaceutically acceptable salt thereof of Embodiment [20] comprising about 900 mg, about 1,000 mg, about 1,100 mg, or about 1,200 mg in a single dosing unit.

[0087] [23] The oral dose of budiadarone of any of Embodiments [21]-[22], wherein said dosing unit is a pill, a capsule, or a tablet.

[0088] [24] The oral dose of budiadarone or a pharmaceutically acceptable salt of any of Embodiments [20] to [23], wherein the budiadarone salt is budiadarone tartrate.

Claims

1. A method comprising cardioverting a patient to normal sinus rhythm, the method comprising: administering, to the patient based on a determination that the patient is in need of cardioversion, a single oral dose comprising at least about 400 mg of budiadarone or a pharmaceutically acceptable salt thereof, wherein the patient is fed about 1 hour or less before the administering of the budiadarone or the pharmaceutically acceptable salt thereof and wherein the patient has either paroxysmal or persistent atrial fibrillation (AFib).
2. The method of claim 1, wherein the single oral dose comprises between 400 mg to 1000 mg budiadarone or pharmaceutically acceptable salt thereof.
3. The method of claim 1, wherein the determination that the patient is suitable for cardioversion comprises determining one or more of: that the patient is experiencing an AFib episode deemed suitable for cardioversion; that data confirms an absence of blood clots in the patient; or that the patient is on a blood thinner.
4. The method of claim 1, further comprising: monitoring a heart rhythm of the patient, subsequent to the administering, until a budiadarone blood plasma concentration in the patient is measured to be below about 20 ng/mL.
5. The method of claim 1, further comprising: monitoring a heart rhythm of the patient, subsequent to the administering, to confirm that the patient converts to and maintains normal sinus rhythm.
6. The method of claim 1, wherein the single oral dose comprises the budiadarone or the pharmaceutically acceptable salt thereof in an amount that is sufficient to: provide for a peak budiadarone blood plasma concentration in the patient of at least about 50 ng/mL; maintain a

- budiodarone blood plasma concentration in the patient of at least about 13 ng/ml for at least 6 hours; and convert a heart rhythm of the patient from AFib to sinus rhythm.
- 7.** The method of claim 1, wherein the single oral dose comprises one or more unit doses and the administering of the single oral dose is performed within a 30 minute time period.
- 8.** The method of claim 1, wherein an amount of the budiodarone or the pharmaceutically acceptable salt thereof in the single oral dose is sufficient to provide for a peak budiodarone blood plasma concentration of at least about 300 ng/ml in the patient.
- 9.** The method of claim 1, wherein an amount of the budiodarone or the pharmaceutically acceptable salt thereof in the single oral dose is sufficient to provide a budiodarone blood plasma concentration in the patient of at least about 50 ng/ml over at least about 5 hours.
- 10.** The method of claim 1, wherein an amount of the budiodarone or the pharmaceutically acceptable salt thereof in the single oral dose is sufficient to provide a budiodarone blood plasma concentration in the patient of at least about 50 ng/mL over at least about 6 hours.
- 11.** The method of claim 1, wherein the budiodarone or the pharmaceutically acceptable salt thereof is administered orally using one or more of: a capsule, a pill, or a tablet.
- 12.** The method of claim 3, which further comprises administering a blood thinner with the budiodarone or the pharmaceutically acceptable salt thereof, wherein the blood thinner comprises one or more of dabigatran, edoxaban, apixaban, or rivaroxaban.
- 13.** The method of claim 1, wherein cardioversion is achieved in about 90 minutes or less from the administering of the budiodarone or the pharmaceutically acceptable salt thereof.
- 14.** The method of claim 1, wherein the budiodarone or the pharmaceutically acceptable salt thereof comprises budiodarone tartrate.
-