



US 20250282757A1


(19) **United States**(12) **Patent Application Publication**
ZHANG et al.(10) **Pub. No.: US 2025/0282757 A1**(43) **Pub. Date: Sep. 11, 2025**(54) **ARYL ETHER-SUBSTITUTED
HETEROCYCLIC COMPOUNDS AS GLP1R
AGONISTS**(71) Applicant: **MINDRANK AI LTD.**, Hangzhou
(CN)(72) Inventors: **Long ZHANG**, Hangzhou (CN);
Zhangming NIU, Hangzhou (CN);
Bowen TANG, Hangzhou (CN)(21) Appl. No.: **19/218,087**(22) Filed: **May 23, 2025****Publication Classification**

- (51) **Int. Cl.**
C07D 401/14 (2006.01)
C07D 405/14 (2006.01)
C07D 471/04 (2006.01)
- (52) **U.S. Cl.**
CPC **C07D 401/14** (2013.01); **C07D 405/14**
(2013.01); **C07D 471/04** (2013.01)


(57) **ABSTRACT**

The present invention relates to a novel aryl ether substituted heterocyclic compound having GLP1R agonist activity. Specifically, disclosed are a compound as a GLP1R agonist represented by formula (I), or a pharmaceutically acceptable salt, solvate, hydrate, isotope substituent or isomer thereof.

Formula (I)

**Related U.S. Application Data**(63) Continuation of application No. 18/255,247, filed on
May 31, 2023, filed as application No. PCT/CN2022/
075295 on Jan. 31, 2022.**Foreign Application Priority Data**

Aug. 30, 2021 (CN) 202111017657.5
Sep. 29, 2021 (CN) 202111168512.5

IPGTT




FIG. 1




FIG. 2




FIG. 3

ARYL ETHER-SUBSTITUTED HETEROCYCLIC COMPOUNDS AS GLP1R AGONISTS

[0001] This application claims the priority of the previous applications submitted to the State Intellectual Property Office of China on Aug. 30, 2021, with the patent application number of 202111017657.5 and the name of “novel aromatic ether substituted heterocyclic compound as GLP1R agonist” and submitted to the State Intellectual Property Office of China on Sep. 29, 2021 with the patent application number of 202111168512.5 and the name of “novel aromatic ether substituted heterocyclic compound as GLP1R agonist”. The full text of the earlier applications are incorporated herein by reference.

TECHNICAL FIELD

[0002] The invention belongs to the field of medicinal chemistry, specifically comprising novel aryl ether substituted heterocyclic compounds having GLP1R agonistic activity, compositions comprising the class of compounds and methods for applying the class of compounds to the preparation of drugs for the treatment or prevention of diseases associated with GLP1/GLP1R.

BACKGROUND OF THE INVENTION

[0003] With the improvement of living standards, overweight or obesity is becoming more and more common in modern society. The number of patients with obesity related complications such as diabetes and fatty liver is increasing. Reports from the World Health Organization (WHO) and ZhiYan advisory predict that the number of obese patients in the world will reach 3.26 billion by 2030; By 2029, the number of global patients with diabetes will exceed 500 million; The number of patients with nonalcoholic fatty liver disease in the world will exceed 1.5 billion. At present, there is no effective drug for fatty liver; There are only 6 drugs approved by FDA for the treatment of obesity, and most of them are controlled drugs with weak efficacy and strong side effects; Although many drugs for the treatment of type 2 diabetes have been approved for marketing, achieved optimal diabetes control-rate (HbA1c<7%), and the achieved optimal diabetes control-rate of even the most active combination drug is only about 45%. Therefore, for fatty liver, obesity or diabetes, new drugs need to be developed to meet the unmet needs of more patients.

[0004] Glucagon-like peptide-1 (GLP-1) is a long peptide hormone containing 30 or 31 amino acids. It is produced and secreted by enteroendocrine L cells and certain neurons in the nucleus tractus solitarius of the brainstem during feeding. GLP-1 stimulates insulin secretion, reduces glucagon secretion, inhibits gastric emptying, reduces appetite, and stimulates beta cell proliferation in a physiological and glucose-dependent manner. In non-clinical experiments, GLP-1 promotes β -cell persistence by stimulating transcription of genes important for glucose-dependent insulin secretion and promoting β -cell regeneration (Meier, et al. *Biodrugs*. 2003; 17(2): 93-102). The GLP1 receptor is a proven ideal target for the treatment of metabolic diseases such as obesity, diabetes, fatty liver, etc., and several GLP1R agonist peptide drugs such as dulaglutide, somalutamide have been approved abroad for the treatment of diabetes and weight loss.

[0005] However, these peptides need to be injected with poor compliance, high cost, poor accessibility, and heavy social medical burden. These peptides need to be refrigerated, which is inconvenient to carry and store. In addition, these peptides are difficult for combination use with existing oral small molecule drugs for diseases having complex causes and requiring treatment of multiple drug combinations, such as nonalcoholic fatty liver disease. Therefore, there is an urgent need to develop small-molecule oral GLP1R agonists.

[0006] Oral small molecule GLP1 agonists have been reported in a few literatures or patents: for example, Pfizer's PF-06882961 can achieve similar or better efficacy as GLP1 polypeptides (<https://doi.org/10.1101/2020.09.29.319483>). Although the efficacy and safety have been preliminarily verified, PF-06882961 has some druggability deficiencies, such as poor oral absorption, extremely low bioavailability, high clinical dosage, and heavy burden on gastrointestinal tract of patients (GI toxicity), which cannot achieve better glucose- and weight-lowering effect by further increasing the drug amount, and so on. Therefore, it is necessary to develop new small-molecule GLP1 agonists with better druggabilities to meet the needs of more patients.


Technical Effects

[0007] The inventors unexpectedly found that some of the novel aryl ether-substituted heterocyclic compounds of formula (I) of the present invention not only have significant GLP1R agonistic activity, but also have better pharmacokinetics parameters (including longer T1/2, higher exposure) and bioavailability than the reference compound PF-06882961 with a known structure. These compounds are expected to have better human PK, and are more suitable as drug candidates for preventing or treating diseases related to GLP1/GLP1R target or signaling pathway.

SUMMARY OF INVENTION

[0008] The object of the present invention is to provide the compound represented by Formula (I) or its pharmaceutically acceptable salt, solvate, enantiomer and isotopic variations thereof.


Formula (I)



[0009] Wherein,

[0010] Ring A and ring B are optionally and independently selected from monocyclic or polycyclic structures of 3 to 18 carbons atoms, and the monocyclic or polycyclic structures can be optionally selected from aromatic rings, heteroaromatic rings, aliphatic rings, heterocyclic rings, fused, spirocyclic or bridged ring structures;

- [0011] X and X' are each independently selected from $-\text{C}(\text{R}_{d1})(\text{R}_{d2})-$, $-\text{C}(=\text{O})\text{N}(\text{R}_{d3})-$, $-\text{N}(\text{R}_{d4})-$, $-\text{C}(=\text{NR}_{d5})-$, $-\text{S}(=\text{O})_2\text{N}(\text{R}_{d6})-$, $-\text{N}(\text{R}_{d7})-$, $-\text{O}-$, $-\text{S}-$, $-\text{C}(=\text{O})\text{O}-$, $-\text{OC}(=\text{O})-$, $-\text{C}(=\text{O})-$, $-\text{C}(=\text{S})-$, $-\text{S}(=\text{O})-$, or $-\text{S}(=\text{O})_2-$;
- [0012] L is independently selected from $-\text{C}(\text{R}_{d1})(\text{R}_{d2})-$, $-\text{OC}(\text{R}_{d1})(\text{R}_{d2})-$, $-\text{C}(\text{R}_{d1})(\text{R}_{d2})\text{O}-$, $-\text{C}(=\text{O})\text{N}(\text{R}_{d3})-$, $-\text{N}(\text{R}_{d4})-$, $-\text{C}(=\text{NR}_{d5})-$, $-\text{S}(=\text{O})_2\text{N}(\text{R}_{d6})-$, $-\text{N}(\text{R}_{d7})-$, $-\text{O}-$, $-\text{S}-$, $-\text{C}(=\text{O})\text{O}-$, $-\text{OC}(=\text{O})-$, $-\text{C}(=\text{O})-$, $-\text{C}(=\text{S})-$, $-\text{S}(=\text{O})-$, or $-\text{S}(=\text{O})_2-$;
- [0013] X_1 , X_2 , X_3 , X_4 , X_5 , X_8 and X_9 are each independently selected from $-\text{CR}_5-$ or $-\text{N}-$;
- [0014] R_0 is independently selected from hydrogen, deuterium, halogen, $-\text{CN}$, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} alkoxy, $-\text{NR}_{d8}\text{R}_{d9}$, 6- to 10-membered aryl, 5- to 8-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl and 3- to 8-membered saturated or partially saturated heterocyclyl, wherein R_0 representing C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} alkoxy, 6- to 10-membered aryl, 5- to 8-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl and 3- to 8-membered saturated or partially saturated heterocyclyl are optionally optimally substituted with one to multiple substituent groups, the substituent groups are optionally selected from hydrogen, deuterium, halogen, alkyl, haloalkyl, alkoxy, alkylamino, $\text{O}=\text{CN}$, OH , $-\text{NR}_{d8}\text{R}_{d9}$, C_{3-10} saturated or partially saturated cycloalkyl, C_{3-10} saturated or partially saturated heterocyclic group, 6- to 10-aryl and 5- to 8-membered heteroaryl; Wherein the aryl, heteroaryl, saturated or partially saturated cycloalkyl, saturated or partially saturated heterocyclyl are further optionally substituted with one to multiple substituents, and the substituents are optionally selected from hydrogen, deuterium, Halogen, alkyl, haloalkyl, cyano, cyanoethyl, $\text{O}=\text{CN}$, OH , C_{1-3} alkyl, C_{1-3} alkoxy, saturated or partially saturated cycloalkyl or saturated or partially saturated heterocyclyl, wherein the C_{1-3} alkyl group, C_{1-3} alkoxy group, saturated or partially saturated cycloalkyl group or saturated or partially saturated heterocyclic group are optionally substituted with 1 to 3 substituent groups, the substituent groups are optionally selected from H, deuterium, halogen, haloalkyl, cyano, OCH_3 and OH .
- [0015] Each R_1 is the same or different and is independently selected from the group consisting of hydrogen, deuterium, halogen, $-\text{CN}$, $-\text{OH}$, $-\text{SH}$ and $-\text{NH}_2$, $-\text{COOH}$ or from C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy, C_{2-10} heteroalkyl, C_{3-10} saturated or partially saturated cycloalkyl, C_{3-10} saturated or partially saturated heterocycloalkyl, C_{1-10} alkyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl, C_{2-10} heteroalkyl substituted by C_{3-10} cycloalkyl, C_{3-10} heterocyclyl, C_{1-10} alkyl substituted with carboxyl or carboxyl surrogate, preferably, the carboxyl surrogate is



Wherein C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy, C_{2-10} heteroalkyl, C_{3-10} saturated or partially saturated cycloalkyl, C_{3-10} saturated or partially saturated heterocycloalkyl, C_{1-10} alkyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl, C_{2-10} heteroalkyl substituted by C_{3-10} cycloalkyl, C_{3-10} heterocyclyl, C_{1-10} alkylcarboxyl or carboxyl surrogates are optionally optimally substituted with one to more substituents selected from H, deuterium, halogen, OCH_3 , Carboxyl, OH , CN and $\text{NR}_{d8}\text{R}_{d9}$; Or any two adjacent R_1 together with the carbon to which they are attached form a 5- to 6-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl, 3- to 8-membered saturated or partially saturated heterocyclyl, the hydrogen on the aryl, saturated or the partially saturated cycloalkyl, heterocycloalkyl is optionally substituted with hydrogen, deuterium, halogen, $-\text{CN}$, $-\text{OH}$, CF_3 , C_{1-6} alkyl, C_{1-6} alkoxy, $-\text{NH}_2$, $-\text{NHC}_{1-6}$ alkyl, $-\text{N}(\text{C}_{1-6}$ alkyl) $_2$, $=\text{O}$, and saturated or partially saturated C_{3-6} cycloalkyl, and the C_{1-6} alkyl and C_{1-6} alkoxy are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN , CF_3 , OH , OCH_3 , OCH_2CH_3 , saturated or partially saturated C_{3-6} cycloalkyl.

[0016] Each R_2 , R_2' and R_{d1} , R_{d2} is the same or different and is independently selected from hydrogen, deuterium, halogen, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} alkylamino, N , N -di(C_{1-10} alkyl)amino, C_{1-10} alkoxy, C_{1-10} alkylacyl, C_{1-10} alkoxy, C_{1-10} alkylsulfonyl, C_{1-10} alkylsulfinyl, C_{3-10} cycloalkylamino, C_{3-10} heterocycloalkylamino, C_{3-10} cycloalkoxy, C_{3-10} cycloalkylacyl, C_{3-10} cycloalkoxyacetyl, C_{3-10} cycloalkylsulfonyl and C_{3-10} cycloalkylsulfinyl; and the alkyl, alkenyl, alkynyl, aryl, saturated or partially saturated cycloalkyl, heterocycloalkyl are optionally substituted with one to more selected from hydrogen, deuterium, halogen, $-\text{CN}$, $-\text{OH}$, CF_3 , C_{1-6} alkyl, C_{1-6} alkoxy, $-\text{NH}_2$, $-\text{NHC}_{1-6}$ alkyl, $-\text{N}(\text{C}_{1-6}$ alkyl) $_2$, oxy, and saturated or partially saturated C_{3-6} cycloalkyl, and the C_{1-6} alkyl and C_{1-6} alkoxy are optionally further substituted with one to more selected from hydrogen, deuterium, halogen, oxo, CN , CF_3 , OH , OCH_3 , OCH_2CH_3 and saturated or partially saturated C_{3-6} cycloalkyl; Or optionally R_2 and R_2' or R_{d1} and R_{d2} may together with the carbon to which they are attached form 5- to 6-membered aryl or heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl, 3- to 8-membered saturated or partially saturated heterocyclic group, wherein the cycloalkyl and heterocycloalkyl are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN , CF_3 , OH , OCH_3 , OCH_2CH_3 ;

[0017] Each R_4 is the same or different and is optionally and independently selected from hydrogen, deuterium, halogen, CN , OH , SH and NH_2 , $-\text{COOH}$ or C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy, C_{2-10} heteroalkyl, C_{3-10} cycloalkyl, C_{3-10} heterocycloalkyl, C_{1-10} alkyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl, and C_{1-10} heteroalkyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl; wherein, the C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN , OH , and C_{3-10} saturated or partially saturated cycloalkyl or heterocyclyl;

[0018] Each R_5 and R_6 are the same or different and is independently selected from hydrogen, deuterium, halogen, CN , OH , SH , $\text{NR}_{d8}\text{R}_{d9}$, NH_2 , $-\text{COOH}$ or

C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl, or C₃₋₁₀ heterocyclyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl; wherein, the C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl, or C₃₋₁₀ heterocycloalkyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, OH and C₃₋₁₀ saturated or partially substituted saturated cycloalkyl or heterocyclyl; Or any adjacent two R₅ or R₆ together with the carbon to which they are attached form 5- to 6-heteroaryl, 3- to 8-saturated or partially saturated cycloalkyl, 3- to 8-saturated or partially saturated heterocyclyl, and wherein the aryl, saturated or partially saturated cycloalkyl, heterocycloalkyl are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, —CN, —OH, CF₃, C₁₋₆ alkyl, C₁₋₆ alkoxy, —NH₂, —NHC₁₋₆ alkyl, —N(C₁₋₆ alkyl)₂, oxy, and saturated or partially saturated C₃₋₆ cycloalkyl, and the C₁₋₆ alkyl and C₁₋₆ alkoxy are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, CF₃, OH, OCH₃, OCH₂CH₃, saturated or partially saturated C₃₋₆ cycloalkyl;

[0019] R_{d3}, R_{d4}, R_{d5}, R_{d6}, R_{d7}, R_{d8}, R_{d9} and R_{d10} are the same or different, and are optionally and independently selected from hydrogen, deuterium, NH₂, C₁₋₁₀ alkyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₁₋₁₀ alkylacyl, C₁₋₁₀ alkylsulfonyl, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl, or C₃₋₁₀ heterocyclyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl; wherein, the C₁₋₁₀ alkyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₁₋₁₀ alkylacyl, C₁₋₁₀ alkylsulfonyl, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl, or C₃₋₁₀ heterocyclyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl are optionally substituted with one to multiple groups selected from hydrogen, deuterium, halogen, oxo, CN, OH and C₃₋₁₀ saturated or partially substituted saturated cycloalkyl or heterocyclyl;

[0020] the hetero represents any heteroatom optionally and independently selected from O, N, S, P and isotopes thereof;

[0021] the halogen is optionally and independently selected from F, Cl, Br, I and isotopes thereof;


[0022] m is an integer optionally selected from 1, 2, 3 and 4;

[0023] n is an integer optionally selected from 0, 1, 2, 3, 4 and 5;

[0024] q is an integer optionally selected from 0, 1, 2, 3, 4 and 5.

[0025] In one embodiment of the present invention, the compound or a pharmaceutically acceptable salt, isotopic variations or isomer thereof has the structure of Formula (IA),

Formula (IA)



[0026] Wherein,

[0027] Ring A and ring B are optionally and independently selected from monocyclic or polycyclic structures of 3 to 18 carbons atoms, and the monocyclic or polycyclic structures can be optionally selected from aromatic rings, heteroaromatic rings, aliphatic rings, heterocyclic rings, fused, spirocyclic or bridged ring structures;

[0028] X and X' are each independently selected from —C(R_{d1})(R_{d2})—, —C(=O)N(R_{d3})—, —N(R_{d4})—, —C(=NR_{d5})—, —S(=O)₂N(R_{d6})—, —N(R_{d7})—, —O—, —S—, —C(=O)O—, —OC(=O)—, —C(=O)—, —C(=S)—, —S(=O)—, or —S(=O)₂—;


[0029] L is independently selected from —C(R_{d1})(R_{d2})—, —OC(R_{d1})(R_{d2})—, —C(R_{d1})(R_{d2})O—, —C(=O)N(R_{d3})—, —N(R_{d4})—, —C(=NR_{d5})—, —S(=O)₂N(R_{d6})—, —N(R_{d7})—, —O—, —S—, —C(=O)O—, —OC(=O)—, —C(=O)—, —C(=S)—, —S(=O)—, or —S(=O)₂—;

[0030] X₁, X₃, X₈ and X₉ are each independently selected from —CR₅— or —N—;

[0031] R₀ is independently selected from hydrogen, deuterium, halogen, —CN, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ alkoxy, —NR_{d8}R_{d9}, 6- to 10-membered aryl, 5- to 8-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl and 3- to 8-membered saturated or partially saturated heterocyclyl, wherein R₀ representing C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ alkoxy, 6- to 10-membered aryl, 5- to 8-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl and 3- to 8-membered saturated or partially saturated heterocyclyl are optionally optimally substituted with one to multiple substituent groups, the substituent groups are optionally selected from hydrogen, deuterium, halogen, alkyl, haloalkyl, alkoxy, alkylamino, O=, CN, OH, —NR_{d8}R_{d9}, C₃₋₁₀ saturated or partially saturated cycloalkyl, C₃₋₁₀ saturated or partially saturated heterocyclic group, 6- to 10-aryl and 5- to 8-membered heteroaryl; Wherein the aryl, heteroaryl, saturated or partially saturated cycloalkyl, saturated or partially saturated heterocyclyl are further optionally substituted with one to multiple substituents, and the substituents are optionally selected from hydrogen, deuterium, halogen, alkyl, haloalkyl, cyano, cyanoethyl, O=, OH, C₁₋₃ alkyl, C₁₋₃ alkoxy, saturated or partially saturated cycloalkyl or saturated or partially saturated heterocyclyl, wherein the C₁₋₃ alkyl group, C₁₋₃ alkoxy group,

saturated or partially saturated cycloalkyl group or saturated or partially saturated heterocyclic group are optionally optimally substituted with 1 to 3 substituent groups, the substituent groups are optionally selected from H, deuterium, halogen, haloalkyl, cyano, OCH_3 and OH.

[0032] Each R_1 is the same or different and is independently selected from the group consisting of hydrogen, deuterium, halogen, $-\text{CN}$, $-\text{OH}$, $-\text{SH}$ and $-\text{NH}_2$, $-\text{COOH}$ or from C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy, C_{2-10} heteroalkyl, C_{3-10} saturated or partially saturated cycloalkyl, C_{3-10} saturated or partially saturated heterocycloalkyl, C_{1-10} alkyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl, C_{2-10} heteroalkyl substituted by C_{3-10} cycloalkyl, C_{3-10} heterocyclyl, C_{1-10} alkyl substituted with carboxyl or carboxyl surrogate, preferably, the carboxyl surrogate is



Wherein C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy, C_{2-10} heteroalkyl, C_{3-10} saturated or partially saturated cycloalkyl, C_{3-10} saturated or partially saturated heterocycloalkyl, C_{1-10} alkyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl, C_{2-10} heteroalkyl substituted by C_{3-10} cycloalkyl, C_{3-10} heterocyclyl, C_{1-10} alkylcarboxyl or carboxyl surrogates are optionally optimally substituted with one to more substituents selected from H, Deuterium, Halogen, OCH_3 , Carboxyl, OH, CN and $\text{NR}_{d8}\text{R}_{d9}$; Or any two adjacent R_1 together with the carbon to which they are attached form a 5- to 6-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl, 3- to 8-membered saturated or partially saturated heterocyclyl, hydrogen on the aryl, saturated or the partially saturated cycloalkyl, heterocycloalkyl is optionally substituted with hydrogen, deuterium, halogen, $-\text{CN}$, $-\text{OH}$, CF_3 , C_{1-6} alkyl, C_{1-6} alkoxy, $-\text{NH}_2$, $-\text{NHC}_{1-6}$ alkyl, $-\text{N}(\text{C}_{1-6}$ alkyl) $_2$, $=\text{O}$, and saturated or partially saturated C_{3-6} cycloalkyl, and the C_{1-6} alkyl and C_{1-6} alkoxy are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, CF_3 , OH, OCH_3 , OCH_2CH_3 , saturated or partially saturated C_{3-6} cycloalkyl.

[0033] Each R_2 , R_2 and R_{d1} , R_{d2} is the same or different and is independently selected from hydrogen, deuterium, halogen, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} alkylamino, N, N-di(C_{1-10} alkyl)amino, C_{1-10} alkoxy, C_{1-10} alkylacyl, C_{1-10} alkoxy, C_{1-10} alkylsulfonyl, C_{1-10} alkylsulfinyl, C_{3-10} cycloalkylamino, C_{3-10} heterocycloalkylamino, C_{3-10} cycloalkoxy, C_{3-10} cycloalkylacyl, C_{3-10} cycloalkoxyacetyl, C_{3-10} cycloalkylsulfonyl and C_{3-10} cycloalkylsulfinyl; and the alkyl, alkenyl, alkynyl, aryl, saturated or partially saturated cycloalkyl, heterocycloalkyl are optionally substituted with one to more selected from hydrogen, deuterium, halogen, $-\text{CN}$, $-\text{OH}$, CF_3 , C_{1-6} alkyl, C_{1-6} alkoxy, $-\text{NH}_2$, $-\text{NHC}_{1-6}$ alkyl, $-\text{N}(\text{C}_{1-6}$ alkyl) $_2$, oxy, and saturated or partially saturated C_{3-6} cycloalkyl, and the C_{1-6} alkyl and C_{1-6} alkoxy are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, CF_3 , OH, OCH_3 , OCH_2CH_3 and saturated or partially

saturated C_{3-6} cycloalkyl; Or optionally R_2 and R_2 , or R_{d1} and R_{d2} may together with the carbon to which they are attached form 5- to 6-membered aryl or heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl, 3- to 8-membered saturated or partially saturated heterocyclic group, wherein the cycloalkyl and heterocycloalkyl are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, CF_3 , OH, OCH_3 , OCH_2CH_3 ;

[0034] Each R_4 is the same or different and is optionally and independently selected from hydrogen, deuterium, halogen, CN, OH, SH and NH_2 , $-\text{COOH}$ or C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy, C_{2-10} heteroalkyl, C_{3-10} cycloalkyl, C_{3-10} heterocycloalkyl, C_{1-10} alkyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl, and C_{1-10} heteroalkyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl; wherein, the C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, OH, and C_{3-10} saturated or partially saturated cycloalkyl or heterocyclyl;

[0035] Each R_5 and R_6 is the same or different and independently selected from the group consisting of hydrogen, deuterium, halogen, CN, OH, SH, $\text{NR}_{d8}\text{R}_{d9}$, NH_2 , $-\text{COOH}$ or C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy, C_{2-10} heteroalkyl, C_{3-10} cycloalkyl, C_{3-10} heterocycloalkyl, C_{1-10} alkyl substituted by C_{3-10} cycloalkyl, or C_{3-10} heterocyclyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl, wherein, the C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy, C_{2-10} heteroalkyl, C_{3-10} cycloalkyl, C_{3-10} heterocycloalkyl, C_{1-10} alkyl substituted by C_{3-10} cycloalkyl, or C_{3-10} heterocycloalkyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, OH and C_{3-10} saturated or partially substituted saturated cycloalkyl or Heterocyclyl; Or any adjacent two R_5 or R_6 together with the carbon to which they are attached form 5- to 6-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl, 3- to 8-membered saturated or partially saturated heterocyclyl, and the aryl, saturated or partially saturated cycloalkyl, heterocycloalkyl are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, $-\text{CN}$, $-\text{OH}$, CF_3 , C_{1-6} alkyl, C_{1-6} alkoxy, $-\text{NH}_2$, $-\text{NHC}_{1-6}$ alkyl, $-\text{N}(\text{C}_{1-6}$ alkyl) $_2$, oxy, and saturated or partially saturated C_{3-6} cycloalkyl, and the C_{1-6} alkyl and C_{1-6} alkoxy is optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, CF_3 , OH, OCH_3 , OCH_2CH_3 , saturated or partially saturated C_{3-6} cycloalkyl;

[0036] Each R_{d3} , R_{d4} , R_{d4} , R_{d6} , R_{d7} , R_{d8} , R_{d9} and R_{d10} is the same or different, and is optionally and independently selected from hydrogen, deuterium, NH_2 , C_{1-10} alkyl, C_{2-10} alkynyl or C_{1-10} alkoxy, C_{1-10} alkylacyl, C_{1-10} alkylsulfonyl, C_{2-10} heteroalkyl, C_{3-10} cycloalkyl, C_{3-10} heterocycloalkyl, C_{1-10} alkyl substituted by C_{3-10} cycloalkyl, or C_{3-10} heterocyclyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl; wherein, the C_{1-10} alkyl, C_{2-10} alkynyl or C_{1-10} alkoxy, C_{1-10} alkylacyl, C_{1-10} alkylsulfonyl, C_{2-10} heteroalkyl, C_{3-10} cycloalkyl, C_{3-10} heterocycloalkyl, C_{1-10} alkyl substituted by C_{3-10}

cycloalkyl, or C₃₋₁₀ heterocyclcyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl are optionally substituted with one to multiple groups selected from hydrogen, deuterium, halogen, oxo, CN, OH and C₃₋₁₀ saturated or partially substituted saturated cycloalkyl or heterocyclcyl;

[0037] The hetero represents any heteroatom independently selected from O, N, S, P and isotopes thereof;

[0038] The halogen is optionally and independently selected from F, Cl, Br, I and isotopes thereof;


[0039] m is an integer optionally selected from 1, 2, 3 and 4;

[0040] n is an integer optionally selected from 0, 1, 2, 3, 4 and 5;

[0041] q is an integer optionally selected from 0, 1, 2, 3, 4 and 5;

[0042] t is an integer optionally selected from 0, 1, 2, 3, and 4.

[0043] In one embodiment of the present invention, the compound or a pharmaceutically acceptable salt, isotopic variations or isomer thereof has the structure of Formula (IB),



[0044] Wherein,

[0045] Ring B is optionally and independently selected from monocyclic or polycyclic structures of 3 to 18 carbons atoms, and the monocyclic or polycyclic structures can be optionally selected from aromatic rings, heteroaromatic rings, aliphatic rings, heterocyclic rings, fused, spirocyclic or bridged ring structures;

[0046] X and X' are independently selected from
 $-\text{C}(\text{R}_{d1})(\text{R}_{d2})-$, $-\text{C}(=\text{O})\text{N}(\text{R}_{d3})-$, $-\text{N}(\text{R}_{d4})-$,
 $-\text{C}(=\text{NR}_{d3})-$, $-\text{S}(=\text{O})_2\text{N}(\text{R}_{d6})-$, $-\text{N}(\text{R}_{d7})-$,
 $-\text{O}-$, $-\text{S}-$, $-\text{C}(=\text{O})\text{O}-$, $-\text{OC}(=\text{O})-$,
 $-\text{C}(=\text{O})-$, $-\text{C}(=\text{S})-$, $-\text{S}(=\text{O})-$, or $-\text{S}(=\text{O})$
 $^-$;


[0047] L is independently selected from $-C(R_{d1})$, $(R_{d2})-$, $-OC(R_{d1})(R_{d2})-$, $-C(R_{d1})(R_{d2})O-$, $-C(=O)N(R_{d3})-$, $-N(R_{d4})-$, $-C(=NR_{d5})-$, $-S(=O)_2N(R_{d6})-$, $-N(R_{d7})-$, $-O-$, $-S-$, $-C(=O)O-$, $-OC(=O)-$, $-C(=O)-$, $-C(S)-$, $-S(=O)-$, or $-S(=O)_2-$;

[0048] $X_1, X_3, X_8, X_9, X_{10}, X_{11}$ and X_{12} are independently selected from $-\text{CR}_5-$ or $-\text{N}-$;

[0049] R₀ is independently selected from hydrogen, deuterium, halogen, —CN, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ alkoxy, —NR₄₈R₄₉, 6- to 10-membered aryl, 5- to 8-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl and 3- to 8-membered saturated or partially saturated heterocyclyl, wherein R₀ representing C₁₋₁₀ alkyl, C₂₋₁₀ al-

enyl, C₂₋₁₀ alkynyl, C₁₋₁₀ alkoxy, 6- to 10-membered aryl, 5- to 8-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl and 3- to 8-membered saturated or partially saturated heterocyclyl are optionally substituted with one to multiple substituent groups, the substituent groups are optionally selected from hydrogen, deuterium, halogen, alkyl, Haloalkyl, carboxyl, alkoxy, alkylamino, O=, CN, OH, —NR₁₈R₁₉, C₃₋₁₀ saturated or partially saturated cycloalkyl, C₃₋₁₀ saturated or partially saturated heterocyclic group, 6- to 10-aryl and 5- to 8-membered heteroaryl; Wherein the aryl, heteroaryl, saturated or partially saturated cycloalkyl, saturated or partially saturated heterocyclyl are optionally substituted with one to multiple substituents, and the substituents are optionally selected from hydrogen, deuterium, Halogen, alkyl, haloalkyl, cyano, cyanoethyl, O=, OH, C₁₋₃ alkyl, C₁₋₃ alkoxy, saturated or partially saturated cycloalkyl or saturated or partially saturated heterocyclyl, wherein the C₁₋₃ alkyl group, C₁₋₃ alkoxy group, saturated or partially saturated cycloalkyl group or saturated or partially saturated heterocyclic group are optionally substituted with one to three substituent groups, the substituent groups are optionally selected from H, deuterium, halogen, haloalkyl, cyano, OCH₃ and OH.

[0050] Each R₁ is the same or different and is independently selected from the group consisting of hydrogen, deuterium, halogen, —CN, —OH, —SH and —NH₂, —COOH or from C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ Alkynyl or C₁₋₁₀ alkoxy, C₂₋₁₀ heteroalkyl, C₃₋₁₀ saturated or partially saturated cycloalkyl, C₃₋₁₀ saturated or partially saturated heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl, C₂₋₁₀ heteroalkyl substituted by C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocyclyl, C₁₋₁₀ alkyl substituted with carboxyl or carboxyl surrogate, preferably, the carboxyl surrogate is



Wherein C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₂₋₁₀ heteroalkyl, C₃₋₁₀ saturated or partially saturated cycloalkyl, C₃₋₁₀ saturated or partially saturated heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl, C₂₋₁₀ heteroalkyl substituted by C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocyclyl, C₁₋₁₀ alkylcarboxyl or carboxyl surrogates are optionally substituted with one to more groups selected from H, deuterium, Halogen, OCH₃, Carboxyl, OH, CN and NR_{ab}R_{ab}; Or any two adjacent R₁ together with the carbon to which they are attached form a 5- to 6-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl, 3- to 8-membered saturated or partially saturated heterocyclyl, and the aryl, saturated or the partially saturated cycloalkyl, heterocycloalkyl is optionally substituted with hydrogen, deuterium, halogen, —CN, —OH, CF₃, C₁₋₆ alkyl, C₁₋₆ alkoxy, —NH₂, —NHC₁₋₆ alkyl, —N(C₁₋₆ alkyl)₂, =O, and saturated or partially saturated C₃₋₆ cycloalkyl, and the C₁₋₆ alkyl and C₁₋₆ alkoxy are optionally substituted with one to more groups selected

from hydrogen, deuterium, halogen, oxo, CN, CF₃, OH, OCH₃, OCH₂CH₃, saturated or partially saturated C₃₋₆ cycloalkyl.

[0051] Each R₂, R₂, and R_{d1}, R_{d2} is the same or different and is independently selected from hydrogen, deuterium, halogen, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ alkylamino, N, N-di(C₁₋₁₀ alkyl)amino, C₁₋₁₀ alkoxy, C₁₋₁₀ alkylacyl, C₁₋₁₀ alkoxy, C₁₋₁₀ alkylsulfonyl, C₁₋₁₀ alkylsulfinyl, C₃₋₁₀ cycloalkylamino, C₃₋₁₀ heterocycloalkylamino, C₃₋₁₀ cycloalkoxy, C₃₋₁₀ cycloalkylacyl, C₃₋₁₀ cycloalkoxyacetyl, C₃₋₁₀ cycloalkylsulfonyl and C₃₋₁₀ cycloalkylsulfinyl; and the alkyl, alkenyl, alkynyl, aryl, saturated or partially saturated cycloalkyl, heterocycloalkyl are optionally substituted with one to more selected from hydrogen, deuterium, halogen, —CN, —OH, CF₃, C₁₋₆ alkyl, C₁₋₆ alkoxy, —NH₂, —NHC₁₋₆ alkyl, —N(C₁₋₆ alkyl)₂, oxy, and saturated or partially saturated C₃₋₆ cycloalkyl, and the C₁₋₆ alkyl and C₁₋₆ alkoxy are optionally substituted with one to more selected from hydrogen, deuterium, halogen, oxo, CN, CF₃, OH, OCH₃, OCH₂CH₃ and saturated or partially saturated C₃₋₆ cycloalkyl; Or optionally R₂ and R₂, or R_{d1} and R_{d2} may together with the carbon to which they are attached form 5- to 6-membered aryl or heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl, 3- to 8-membered saturated or partially saturated heterocyclic group, wherein the cycloalkyl and heterocycloalkyl are optionally substituted with one to more selected from hydrogen, deuterium, halogen, oxo, CN, CF₃, OH, OCH₃, OCH₂CH₃;

[0052] Each R₄ is the same or different and is optionally and independently selected from hydrogen, deuterium, halogen, CN, OH, SH and NH₂, —COOH or C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl, and C₁₋₁₀ heteroalkyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl; wherein, the C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, OH, and C₃₋₁₀ saturated or partially saturated cycloalkyl or heterocyclyl;

[0053] Each R₅ and R₆ is the same or different and is independently hydrogen, deuterium, halogen, CN, OH, SH, NR_{d8}R_{d9}, NH₂, —COOH or C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl, or C₃₋₁₀ heterocyclyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl; wherein, the C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl, or C₃₋₁₀ heterocycloalkyl are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, OH and C₃₋₁₀ saturated or partially substituted saturated cycloalkyl or heterocyclyl; Or any adjacent two R₅ or R₆ together with the carbon to which they are attached form 5- to 6-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl, 3- to 8-membered saturated or partially saturated heterocyclyl, and the aryl, saturated or partially saturated cycloalkyl, heterocycloalkyl are optionally substituted with one to more groups selected from hydrogen,

deuterium, halogen, —CN, —OH, CF₃, C₁₋₆ alkyl, C₁₋₆ alkoxy, —NH₂, —NHC₁₋₆ alkyl, —N(C₁₋₆ alkyl)₂, oxy, and saturated or partially saturated C₃₋₆ cycloalkyl, and the C₁₋₆ alkyl and C₁₋₆ alkoxy are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, CF₃, OH, OCH₃, OCH₂CH₃, saturated or partially saturated C₃₋₆ cycloalkyl;

[0054] Each R_{d3}, R_{d4}, R_{d5}, R_{d6}, R_{d7}, R_{d8}, R_{d9} and R_{d10} is the same or different, and is optionally and independently selected from hydrogen, deuterium, NH₂, C₁₋₁₀ alkyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₁₋₁₀ alkylacyl, C₁₋₁₀ alkylsulfonyl, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl, or C₃₋₁₀ heterocyclyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl; wherein, the C₁₋₁₀ alkyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₁₋₁₀ alkylacyl, C₁₋₁₀ alkylsulfonyl, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl, or C₃₋₁₀ heterocyclyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl are optionally substituted with one to multiple groups selected from hydrogen, deuterium, halogen, oxo, CN, OH and C₃₋₁₀ saturated or partially substituted saturated cycloalkyl or heterocyclyl;

[0055] The hetero represents any heteroatom independently selected from O, N, S, P and isotopes thereof;

[0056] The halogen is optionally and independently selected from F, Cl, Br, I and isotopes thereof;

[0057] m is an integer optionally selected from 1, 2, 3 and 4;


[0058] n is an integer optionally selected from 0, 1, 2, 3, 4 and 5;

[0059] q is an integer optionally selected from 0, 1, 2, 3, 4 and 5;

[0060] t is an integer optionally selected from 0, 1, 2, 3, and 4.

[0061] In one embodiment of the present invention, the compound or a pharmaceutically acceptable salt, isotopic variations or isomer thereof has the structure of Formula (IC),

Formula(IC)



[0062] Wherein,

[0063] ——— optionally represents a single bond or a double bond;


[0064] X and X' are independently selected from —C(R_{d1})(R_{d2})—, —C(=O)N(R_{d3})—, —N(R_{d4})—, —C(=NR_{d5})—, —S(=O)₂N(R_{d6})—, —N(R_{d7})—, —O—, —S—, —C(=O)O—, —OC(=O)—, —C(=O)—, —C(=S)—, —S(=O)—, or —S(=O)₂—;

[0065] L is independently selected from $-\text{C}(\text{R}_{d1})(\text{R}_{d2})-$, $-\text{OC}(\text{R}_{d1})(\text{R}_{d2})-$, $-\text{C}(\text{R}_{d1})(\text{R}_{d2})\text{O}-$, $-\text{C}(=\text{O})\text{N}(\text{R}_{d3})-$, $-\text{N}(\text{R}_{d4})-$, $-\text{C}(=\text{NR}_{d5})-$, $-\text{S}(=\text{O})_2\text{N}(\text{R}_{d6})-$, $-\text{N}(\text{R}_{d7})-$, $-\text{O}-$, $-\text{S}-$, $-\text{C}(=\text{O})\text{O}-$, $-\text{OC}(=\text{O})-$, $-\text{C}(=\text{O})-$, $-\text{C}(=\text{S})-$, $-\text{S}(=\text{O})-$, or $-\text{S}(=\text{O})_2-$;

[0066] X_1 , X_3 , X_8 , X_9 , X_{10} , X_{11} , X_{12} , X_{13} or X_{14} are independently selected from $-\text{CR}_5-$ or $-\text{N}-$;

[0067] R_0 is independently selected from hydrogen, deuterium, halogen, $-\text{CN}$, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} alkoxy, $-\text{NR}_{d8}\text{R}_{d9}$, 6- to 10-membered aryl, 5- to 8-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl and 3- to 8-membered saturated or partially saturated heterocyclyl, wherein R_0 representing C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} alkoxy, 6- to 10-membered aryl, 5- to 8-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl and 3- to 8-membered saturated or partially saturated heterocyclyl are optionally substituted with one to multiple substituent groups, the substituent groups are optionally selected from hydrogen, deuterium, halogen, alkyl, haloalkyl, alkoxy, alkylamino, $\text{O}=-$, CN , OH , $-\text{NR}_{d8}\text{R}_{d9}$, C_{3-10} saturated or partially saturated cycloalkyl, C_{3-10} saturated or partially saturated heterocyclic group, 6- to 10-aryl and 5- to 8-membered heteroaryl; Wherein the aryl, heteroaryl, saturated or partially saturated cycloalkyl, saturated or partially saturated heterocyclyl are further optionally substituted with one to multiple substituents, and the substituents are optionally selected from hydrogen, deuterium, halogen, alkyl, haloalkyl, cyano, cyanoethyl, $\text{O}=-$, OH , C_{1-3} alkyl, C_{1-3} alkoxy, saturated or partially saturated cycloalkyl or saturated or partially saturated heterocyclyl, wherein the C_{1-3} alkyl group, C_{1-3} alkoxy group, saturated or partially saturated cycloalkyl group or saturated or partially saturated heterocyclic group are optionally substituted with one to three substituent groups, the substituent groups are optionally selected from H, deuterium, halogen, haloalkyl, cyano, OCH_3 and OH .

[0068] Each R_1 is the same or different and is independently selected from the group consisting of hydrogen, deuterium, halogen, $-\text{CN}$, $-\text{OH}$, $-\text{SH}$ and $-\text{NH}_2$, $-\text{COOH}$ or from C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} Alkynyl or C_{1-10} alkoxy, C_{2-10} heteroalkyl, C_{3-10} saturated or partially saturated cycloalkyl, C_{3-10} saturated or partially saturated heterocycloalkyl, C_{1-10} alkyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl, C_{2-10} heteroalkyl substituted by C_{3-10} cycloalkyl, C_{3-10} heterocyclyl, C_{1-10} alkyl substituted with carboxyl or carboxyl surrogate, preferably, the carboxyl surrogate is



Wherein C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy, C_{2-10} heteroalkyl, C_{3-10} saturated or partially saturated cycloalkyl, C_{3-10} saturated or partially saturated heterocycloalkyl, C_{1-10} alkyl substituted by C_{3-10} cycloalkyl or

C_{3-10} heterocycloalkyl, C_{2-10} heteroalkyl substituted by C_{3-10} cycloalkyl, C_{3-10} heterocyclyl, C_{1-10} alkylcarboxyl or carboxyl surrogates are optionally substituted with one to more substituents selected from H, Deuterium, Halogen, OCH_3 , Carboxyl, OH , CN and $\text{NR}_{d8}\text{R}_{d9}$; Or any two adjacent R_1 together with the carbon to which they are attached form a 5- to 6-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl, 3- to 8-membered saturated or partially saturated heterocyclyl, and the hydrogen on the aryl, saturated or the partially saturated cycloalkyl, heterocycloalkyl is optionally substituted with hydrogen, deuterium, halogen, $-\text{CN}$, $-\text{OH}$, CF_3 , C_{1-6} alkyl, C_{1-6} alkoxy, $-\text{NH}_2$, $-\text{NHC}_{1-6}$ alkyl, $-\text{N}(\text{C}_{1-6}\text{ alkyl})_2$, $=\text{O}$, and saturated or partially saturated C_{3-6} cycloalkyl, and the C_{1-6} alkyl and C_{1-6} alkoxy are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN , CF_3 , OH , OCH_3 , OCH_2CH_3 , saturated or partially saturated C_{3-6} cycloalkyl.

[0069] Each R_2 , R_2' and R_{d1} , R_{d2} is the same or different and is independently selected from hydrogen, deuterium, halogen, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} alkylamino, N, N-di(C_{1-10} alkyl)amino, C_{1-10} alkoxy, C_{1-10} alkylacyl, C_{1-10} alkoxy, C_{1-10} alkylsulfonfyl, C_{1-10} alkylsulfiny, C_{3-10} cycloalkylamino, C_{3-10} heterocycloalkylamino, C_{3-10} cycloalkoxy, C_{3-10} cycloalkylacyl, C_{3-10} cycloalkoxyacetyl, C_{3-10} cycloalkylsulfonyl and C_{3-10} cycloalkylsulfiny; and the alkyl, alkenyl, alkynyl, aryl, saturated or partially saturated cycloalkyl, heterocycloalkyl are optionally substituted with one to more selected from hydrogen, deuterium, halogen, $-\text{CN}$, $-\text{OH}$, CF_3 , C_{1-6} alkyl, C_{1-6} alkoxy, $-\text{NH}_2$, $-\text{NHC}_{1-6}$ alkyl, $-\text{N}(\text{C}_{1-6}\text{ alkyl})_2$, oxy, and saturated or partially saturated C_{3-6} cycloalkyl, and the C_{1-6} alkyl and C_{1-6} alkoxy are optionally substituted with one to more selected from hydrogen, deuterium, halogen, oxo, CN , CF_3 , OH , OCH_3 , OCH_2CH_3 and saturated or partially saturated C_{3-6} cycloalkyl; Or optionally R_2 and R_2' or R_{d1} and R_{d2} may together with the carbon to which they are attached form 5- to 6-membered aryl or heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl, 3- to 8-membered saturated or partially saturated heterocyclic group, wherein the cycloalkyl and heterocycloalkyl are optionally substituted with one to more selected from hydrogen, deuterium, halogen, oxo, CN , CF_3 , OH , OCH_3 , OCH_2CH_3 ;

[0070] Each R_4 is the same or different and is optionally and independently selected from hydrogen, deuterium, halogen, CN , OH , SH and NH_2 , $-\text{COOH}$ or C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy, C_{2-10} heteroalkyl, C_{3-10} cycloalkyl, C_{3-10} heterocycloalkyl, C_{1-10} alkyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl, and C_{1-10} heteroalkyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl; wherein, the C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN , OH , and C_{3-10} saturated or partially saturated cycloalkyl or heterocyclyl;

[0071] Each R_5 and R_6 is the same or different and is independently selected from hydrogen, deuterium, halogen, CN , OH , SH , $\text{NR}_{d8}\text{R}_{d9}$, NH_2 , $-\text{COOH}$ or C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy, C_{2-10} heteroalkyl, C_{3-10} cycloalkyl, C_{3-10} heterocycloalkyl, C_{1-10} alkyl substituted by C_{3-10} cycloalkyl, or C_{3-10} heterocyclyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl; wherein, the C_{1-10} alkyl, C_{2-10}

alkenyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl, or C₃₋₁₀ heterocycloalkyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, OH and C₃₋₁₀ saturated or partially substituted saturated cycloalkyl or heterocyclyl; Or any adjacent two R₅ or R₆ together with the carbon to which they are attached form 5- to 6-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl, 3- to 8-membered saturated or partially saturated heterocyclyl, and the aryl, saturated or partially saturated cycloalkyl, heterocycloalkyl are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, —CN, —OH, CF₃, C₁₋₆ alkyl, C₁₋₆ alkoxy, —NH₂, —NHC₁₋₆ alkyl, —N(C₁₋₆ alkyl)₂, oxy, and saturated or partially saturated C₃₋₆ cycloalkyl, and the C₁₋₆ alkyl and C₁₋₆ alkoxy is optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, CF₃, OH, OCH₃, OCH₂CH₃, saturated or partially saturated C₃₋₆ cycloalkyl;

[0072] Each R_{d3}, R_{d4}, R_{d5}, R_{d6}, R_{d7}, R_{d8}, R_{d9} and R_{d10} is the same or different, and is optionally and independently selected from hydrogen, deuterium, NH₂, C₁₋₁₀ alkyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₁₋₁₀ alkylacyl, C₁₋₁₀ alkylsulfonyl, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl, or C₃₋₁₀ heterocyclyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl; wherein, the C₁₋₁₀ alkyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₁₋₁₀ alkylacyl, C₁₋₁₀ alkylsulfonyl, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl, or C₃₋₁₀ heterocyclyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl are optionally substituted with one to multiple groups selected from hydrogen, deuterium, halogen, oxo, CN, OH and C₃₋₁₀ saturated or partially substituted saturated cycloalkyl or heterocyclyl;

[0073] The hetero represents any heteroatom independently selected from O, N, S, P and isotopes thereof;

[0074] The halogen is optionally and independently selected from F, Cl, Br, I and isotopes thereof;

[0075] m is an integer optionally selected from 1, 2, 3 and 4;


[0076] n is an integer optionally selected from 0, 1, 2, 3, 4 and 5;

[0077] q is an integer optionally selected from 0, 1, 2, 3, 4 and 5;

[0078] t is an integer optionally selected from 0, 1, 2, 3, and 4.

[0079] In one embodiment of the present invention, the compound or a pharmaceutically acceptable salt, isotopic variations or isomer thereof has the structure of Formula (ID),

Formula(ID)



[0080] Wherein,

[0081] ——— optionally represents a single bond or a double bond;

[0082] X and X' are independently selected from —C(R_{d1})(R_{d2})—, —C(=O)N(R_{d3})—, —N(R_{d4})—, —C(=NR_{d5})—, —S(=O)₂N(R_{d6})—, —N(R_{d7})—, —O—, —S—, —C(=O)O—, —OC(=O)—, —C(=O)—, —C(S)—, —S(=O)—, or —S(=O)₂—; preferred —C(R_{d1})(R_{d2})—, —N(R_{d4})—, —O—, —S—, —S(=O)— or —S(=O)₂—;

[0083] L is independently selected from —C(R_{d1})(R_{d2})—, —OC(R_{d1})(R_{d2})—, —C(R_{d1})(R_{d2})O—, —C(=O)N(R_{d3})—, —N(R_{d4})—, —C(=NR_{d5})—, —S(=O)₂N(R_{d6})—, —N(R_{d7})—, —O—, —S—, —C(=O)O—, —OC(=O)—, —C(=O)—, —C(S)—, —S(=O)—, or —S(=O)₂—; preferred —OC(R_{d1})(R_{d2})—, —C(R_{d1})(R_{d2})O—, —O—, —S— or —N(R_{d4})—;

[0084] X₁, X₃, X₈ and X₁₃ are independently selected from —CR₅— or —N—;

[0085] R₀ is independently selected from hydrogen, deuterium, halogen, —CN, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ alkoxy, —NR_{d8}R_{d9}, 6- to 10-membered aryl, 5- to 8-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl and 3- to 8-membered saturated or partially saturated heterocyclyl, wherein R₀ representing C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ alkoxy, 6- to 10-membered aryl, 5- to 8-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl and 3- to 8-membered saturated or partially saturated heterocyclyl are optionally substituted with one to multiple substituent groups, the substituent groups are optionally selected from hydrogen, deuterium, halogen, alkyl, haloalkyl, alkoxy, alkylamino, O=, CN, OH, —NR_{d8}R_{d9}, C₃₋₁₀ saturated or partially saturated cycloalkyl, C₃₋₁₀ saturated or partially saturated heterocyclic group, 6- to 10-aryl and 5- to 8-membered heteroaryl; wherein the aryl, heteroaryl, saturated or partially saturated cycloalkyl, saturated or partially saturated heterocyclyl are further optionally substituted with one to multiple substituents, and the substituents are optionally selected from hydrogen, deuterium, Halogen, alkyl, haloalkyl, cyano, cyanoethyl, O=, OH, C₁₋₃ alkyl, C₁₋₃ alkoxy, saturated or partially saturated cycloalkyl or saturated or partially saturated heterocyclyl, wherein the C₁₋₃ alkyl group, C₁₋₃ alkoxy group, saturated or partially saturated cycloalkyl group or saturated or partially saturated heterocyclic group are

optionally substituted with one to three substituent groups, the substituent groups are optionally selected from H, deuterium, halogen, haloalkyl, cyano, OCH₃ and OH.

[0086] Each R₁ is the same or different and is independently selected from the group consisting of hydrogen, deuterium, halogen, —CN, —OH, —SH and —NH₂, —COOH or from C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₂₋₁₀ heteroalkyl, C₃₋₁₀ saturated or partially saturated cycloalkyl, C₃₋₁₀ saturated or partially saturated heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl, C₂₋₁₀ heteroalkyl substituted by C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocyclyl, C₁₋₁₀ alkyl substituted with carboxyl or carboxyl surrogate; wherein C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₂₋₁₀ heteroalkyl, C₃₋₁₀ saturated or partially saturated cycloalkyl, C₃₋₁₀ to saturated or partially saturated heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl, C₂₋₁₀ heteroalkyl substituted by C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocyclyl, C₁₋₁₀ alkylcarboxyl or carboxyl surrogates are optionally substituted with one to more substituents selected from H, deuterium, halogen, OCH₃, carboxyl, OH, CN and NR_{d8}R_{d9}; Or any two adjacent R₁ together with the carbon to which they are attached form a 5- to 6-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl, 3- to 8-membered saturated or partially saturated heterocyclyl, a hydrogen on the aryl, saturated or the partially saturated cycloalkyl, or heterocycloalkyl is optionally substituted by hydrogen, deuterium, halogen, —CN, —OH, CF₃, C₁₋₆ alkyl, C₁₋₆ alkoxy, —NH₂, —NHC₁₋₆ alkyl, —N(C₁₋₆ alkyl)₂, =O, and saturated or partially saturated C₃₋₆ cycloalkyl, and the C₁₋₆ alkyl and C₁₋₆ alkoxy are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, CF₃, OH, OCH₃, OCH₂CH₃, saturated or partially saturated C₃₋₆ cycloalkyl.

[0087] Each R₂, R₂, and R_{d1}, R_{d2} is the same or different and is independently selected from hydrogen, deuterium, halogen, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ alkylamino, N, N-di(C₁₋₁₀ alkyl)amino, C₁₋₁₀ alkoxy, C₁₋₁₀ alkylacyl, C₁₋₁₀ alkoxy, C₁₋₁₀ alkylsulfonyl, C₁₋₁₀ alkylsulfinyl, C₃₋₁₀ cycloalkylamino, C₃₋₁₀ heterocycloalkylamino, C₃₋₁₀ cycloalkoxy, C₃₋₁₀ cycloalkylacyl, C₃₋₁₀ cycloalkoxyacetyl, C₃₋₁₀ cycloalkylsulfonyl and C₃₋₁₀ cycloalkylsulfinyl; and the alkyl, alkenyl, alkynyl, aryl, saturated or partially saturated cycloalkyl, heterocycloalkyl are optionally substituted with one to more selected from hydrogen, deuterium, halogen, —CN, —OH, CF₃, C₁₋₆ alkyl, C₁₋₆ alkoxy, —NH₂, —NHC₁₋₆ alkyl, —N(C₁₋₆ alkyl)₂, oxy, and saturated or partially saturated C₃₋₆ cycloalkyl, and the C₁₋₆ alkyl and C₁₋₆ alkoxy are optionally further substituted with one to more selected from hydrogen, deuterium, halogen, oxo, CN, CF₃, OH, OCH₃, OCH₂CH₃ and saturated or partially saturated C₃₋₆ cycloalkyl; Or optionally R₂ and R₂, or R_{d1} and R_{d2} may together with the carbon to which they are attached form 5- to 6-membered aryl or heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl, 3- to 8-membered saturated or partially saturated heterocyclic group, wherein the cycloalkyl and heterocycloalkyl are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, CF₃, OH, OCH₃, OCH₂CH₃;

[0088] Each R₄ is the same or different and is optionally and independently selected from hydrogen, deuterium, halogen, CN, OH, SH and NH₂, —COOH or C₁₋₁₀

alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl, and C₁₋₁₀ heteroalkyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl; wherein, the C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, OH, and C₃₋₁₀ saturated or partially saturated cycloalkyl or heterocyclyl;

[0089] Each R₅ and R₆ is the same or different and is independently hydrogen, deuterium, halogen, CN, OH, SH, NR_{d8}R_{d9}, NH₂, —COOH or C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl, or C₃₋₁₀ heterocyclyl substituted by C₃₋₁₀ Cycloalkyl or C₃₋₁₀ heterocycloalkyl; wherein, the C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl, or C₃₋₁₀ heterocycloalkyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, OH and C₃₋₁₀ saturated or partially substituted saturated cycloalkyl or Heterocyclyl; Or any adjacent two R₅ or R₆ together with the carbon to which they are attached form 5- to 6-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl, 3- to 8-membered saturated or partially saturated heterocyclyl, and the aryl, saturated or partially saturated cycloalkyl, heterocycloalkyl are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, —CN, —OH, CF₃, C₁₋₆ alkyl, C₁₋₆ alkoxy, —NH₂, —NHC₁₋₆ alkyl, —N(C₁₋₆ alkyl)₂, oxy, and saturated or partially saturated C₃₋₆ cycloalkyl, and the C₁₋₆ alkyl and C₁₋₆ alkoxy is optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, CF₃, OH, OCH₃, OCH₂CH₃, saturated or partially saturated C₃₋₆ cycloalkyl;


[0090] Each R_{d4}, R_{d8}, and R_{d9} is the same or different, and is optionally and independently selected from hydrogen, deuterium, NH₂, C₁₋₁₀ alkyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₁₋₁₀ alkylacyl, C₁₋₁₀ alkylsulfonyl, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl, or C₃₋₁₀ heterocyclyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl; wherein, the C₁₋₁₀ alkyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₁₋₁₀ alkylacyl, C₁₋₁₀ alkylsulfonyl, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl, or C₃₋₁₀ heterocyclyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl are optionally substituted with one to multiple groups selected from hydrogen, deuterium, halogen, oxo, CN, OH and C₃₋₁₀ saturated or partially substituted saturated cycloalkyl or heterocyclyl;

[0091] The hetero represents any heteroatom independently selected from O, N, S, P and isotopes thereof;

[0092] The halogen is optionally and independently selected from F, Cl, Br, I and isotopes thereof;

[0093] m is an integer optionally selected from 1, 2, 3 and 4;

- [0094] n is an integer optionally selected from 0, 1, 2, 3, 4 and 5;
- [0095] q is an integer optionally selected from 0, 1, 2, 3, 4 and 5;
- [0096] t is an integer optionally selected from 0, 1, 2, 3, and 4.
- [0097] In one embodiment of the present invention, the compound or a pharmaceutically acceptable salt, isotopic variations or isomer thereof has the structure of Formula (IE),



[0098] Wherein,

[0099] ----- optionally represents a single bond or a double bond;

[0100] X and X' are independently selected from $-\text{C}(\text{R}_{d1})(\text{R}_{d2})-$, $-\text{N}(\text{R}_{d4})-$, $-\text{N}(\text{R}_{d7})-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(=\text{O})-$ or $-\text{S}(=\text{O})_2-$; preferred $-\text{C}(\text{R}_{d1})(\text{R}_{d2})-$, $-\text{N}(\text{R}_{d4})-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(=\text{O})-$ or $-\text{S}(=\text{O})_2-$;

[0101] L is independently selected from $-\text{C}(\text{R}_{d1})(\text{R}_{d2})-$, $-\text{OC}(\text{R}_{d1})(\text{R}_{d2})-$, $-\text{C}(\text{R}_{d1})(\text{R}_{d2})\text{O}-$, $-\text{C}(=\text{O})\text{N}(\text{R}_{d3})-$, $-\text{N}(\text{R}_{d4})-$, $-\text{C}(=\text{NR}_{d5})-$, $-\text{S}(=\text{O})_2\text{N}(\text{R}_{d6})-$, $-\text{N}(\text{R}_{d7})-$, $-\text{O}-$, $-\text{S}-$, $-\text{C}(=\text{O})\text{O}-$, $-\text{OC}(=\text{O})-$, $-\text{C}(=\text{O})-$, $-\text{C}(=\text{S})-$, $-\text{S}(=\text{O})-$ or $-\text{S}(=\text{O})_2-$; preferred $-\text{OC}(\text{R}_{d1})(\text{R}_{d2})-$, $-\text{C}(\text{R}_{d1})(\text{R}_{d2})\text{O}-$, $-\text{O}-$, $-\text{S}-$ or $-\text{N}(\text{R}_{d4})-$;

[0102] X_1 , X_3 , X_8 and X_{13} are independently selected from $-\text{CR}_5-$ or $-\text{N}-$;

[0103] R is independently selected from hydrogen, deuterium, halogen, $-\text{CN}$, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} alkoxy, $-\text{NR}_{d8}\text{R}_{d9}$, 6- to 10-membered aryl, 5- to 8-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl and 3- to 8-membered saturated or partially saturated heterocyclyl, wherein R representing C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} alkoxy, 6- to 10-membered aryl, 5- to 8-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl and 3- to 8-membered saturated or partially saturated heterocyclyl are optionally substituted with one to multiple substituent groups, the substituent groups are optionally selected from hydrogen, deuterium, halogen, alkyl, haloalkyl, alkoxy, alkylamino, $\text{O}=\text{}$, CN , OH , $-\text{NR}_{d8}\text{R}_{d9}$, C_{3-10} saturated or partially saturated cycloalkyl, C_{3-10} saturated or partially saturated heterocyclic group, 6- to 10-aryl and 5- to 8-membered heteroaryl; wherein the aryl, heteroaryl, saturated or partially saturated cycloalkyl, saturated or partially saturated heterocyclyl are optionally substituted with one to multiple substituents, and the substituents are optionally selected from hydrogen,

deuterium, halogen, alkyl, haloalkyl, cyano, cyanoethyl, $\text{O}=\text{}$, OH , C_{1-3} alkyl, C_{1-3} alkoxy, saturated or partially saturated cycloalkyl or saturated or partially saturated heterocyclyl, wherein the C_{1-3} alkyl group, C_{1-3} alkoxy group, saturated or partially saturated cycloalkyl group or saturated or partially saturated heterocyclic group are optionally substituted with one to three substituent groups, the substituent groups are optionally selected from H , deuterium, halogen, haloalkyl, cyano, OCH_3 and OH .

[0104] Each R_1 is the same or different and is independently selected from the group consisting of hydrogen, deuterium, halogen, $-\text{CN}$, $-\text{OH}$, $-\text{SH}$ and $-\text{NH}_2$, $-\text{COOH}$ or from C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy, C_{2-10} heteroalkyl, C_{3-10} saturated or partially saturated cycloalkyl, C_{3-10} saturated or partially saturated heterocycloalkyl, C_{1-10} alkyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl, C_{2-10} heteroalkyl substituted by C_{3-10} cycloalkyl, C_{3-10} heterocyclyl, C_{1-10} alkyl substituted with carboxyl or carboxyl surrogate; or any two adjacent R_1 together with the carbon to which they are attached form a 5- to 6-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl, 3- to 8-membered saturated or partially saturated heterocyclyl, and the hydrogen on the aryl, saturated or the partially saturated cycloalkyl, heterocycloalkyl is optionally substituted with hydrogen, deuterium, halogen, $-\text{CN}$, $-\text{OH}$, CF_3 , C_{1-6} alkyl, C_{1-6} alkoxy, $-\text{NH}_2$, $-\text{NHC}_{1-6}$ alkyl, $-\text{N}(\text{C}_{1-6}\text{ alkyl})_2$, $=\text{O}$, and saturated or partially saturated C_{3-6} cycloalkyl, and the C_{1-6} alkyl and C_{1-6} alkoxy are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN , CF_3 , OH , OCH_3 , OCH_2CH_3 , saturated or partially saturated C_{3-6} cycloalkyl.

[0105] Each R_4 is the same or different and is optionally and independently selected from hydrogen, deuterium, halogen, CN , OH , SH and NH_2 , $-\text{COOH}$ or C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy, C_{2-10} heteroalkyl, C_{3-10} cycloalkyl, C_{3-10} heterocycloalkyl, C_{1-10} alkyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl, and C_{1-10} heteroalkyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl; wherein, the C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN , OH , and C_{3-10} saturated or partially saturated cycloalkyl or heterocyclyl;

[0106] Each R_5 and R_6 is the same or different and is independently selected from hydrogen, deuterium, halogen, CN , OH , SH , $\text{NR}_{d8}\text{R}_{d9}$, NH_2 , $-\text{COOH}$ or C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy, C_{2-10} heteroalkyl, C_{3-10} cycloalkyl, C_{3-10} heterocycloalkyl, C_{1-10} alkyl substituted by C_{3-10} cycloalkyl, or C_{3-10} heterocyclyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl; wherein, the C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy, C_{2-10} heteroalkyl, C_{3-10} cycloalkyl, C_{3-10} heterocycloalkyl, C_{1-10} alkyl substituted by C_{3-10} cycloalkyl, or C_{3-10} heterocycloalkyl are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN , OH and C_{3-10} saturated or partially substituted saturated cycloalkyl or heterocyclyl; or any adjacent two R_5 or R_6 together with the carbon to which they are attached form 5- to 6-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl

kyl, 3- to 8-membered saturated or partially saturated heterocyclyl, and the aryl, saturated or partially saturated cycloalkyl, heterocycloalkyl are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, —CN, —OH, CF₃, C₁₋₆ alkyl, C₁₋₆ alkoxy, —NH₂, —NHC₁₋₆ alkyl, —N(C₁₋₆ alkyl)₂, oxy, and saturated or partially saturated C₃₋₆ cycloalkyl, and the C₁₋₆ alkyl and C₁₋₆ alkoxy are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, CF₃, OH, OCH₃, OCH₂CH₃, saturated or partially saturated C₃₋₆ cycloalkyl;

[0107] Each R_{d1}, R_{d2}, R_{d4}, R_{d8} and R_d is the same or different, and is optionally and independently selected from hydrogen, deuterium, NH₂, C₁₋₁₀ alkyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₁₋₁₀ alkylacyl, C₁₋₁₀ alkylsulfonyl, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl, or C₃₋₁₀ heterocyclyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl; wherein, the C₁₋₁₀ alkyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₁₋₁₀ alkylacyl, C₁₋₁₀ alkylsulfonyl, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl, or C₃₋₁₀ heterocyclyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl are optionally substituted with one to multiple groups selected from hydrogen, deuterium, halogen, oxo, CN, OH and C₃₋₁₀ saturated or partially substituted saturated cycloalkyl or heterocyclyl;

[0108] The hetero represents any heteroatom independently selected from O, N, S, P and isotopes thereof;

[0109] The halogen is optionally and independently selected from F, Cl, Br, I and isotopes thereof;


[0110] m is an integer optionally selected from 1, 2, 3 and 4;

[0111] n is an integer optionally selected from 0, 1, 2, 3, 4 and 5;

[0112] q is an integer optionally selected from 0, 1, 2, 3, 4 and 5;


[0113] t is an integer optionally selected from 0, 1, 2, 3, and 4.

[0114] In one embodiment of the present invention, the compound or a pharmaceutically acceptable salt, isotopic variations or isomer thereof has the structure of Formula (IF-1) or (IF-2),



-continued

Formula (IF-2)



[0115] Wherein,

[0116] ——— optionally represents a single bond or a double bond;

[0117] X and X' are independently selected from —C(R_{d1})(R_{d2})—, —N(R_{d4})—, —O—, —S—, —S(=O)— or —S(=O)₂—;

[0118] L is independently selected from —OC(R_{d1})(R_{d2})—, —C(R_{d1})(R_{d2})O—, —O—, —S— or —N(R_{d4})—;

[0119] X₁ and X₈ are independently selected from —CR₅— or —N—;

[0120] R is independently selected from hydrogen, deuterium, halogen, —CN, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ alkoxy, —NR_{d8}R_{d9}, 6- to 10-membered aryl, 5- to 8-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl and 3- to 8-membered saturated or partially saturated heterocyclyl, wherein R representing C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ alkoxy, 6- to 10-membered aryl, 5- to 8-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl and 3- to 8-membered saturated or partially saturated heterocyclyl are optionally substituted with one to multiple substituent groups, the substituent groups are optionally selected from hydrogen, deuterium, halogen, alkyl, haloalkyl, alkoxy, alkylamino, O=, CN, OH, —NR_{d8}R_{d9}, C₃₋₁₀ saturated or partially saturated cycloalkyl, C₃₋₁₀ saturated or partially saturated heterocyclic group, 6- to 10-aryl and 5- to 8-membered heteroaryl; Wherein the aryl, heteroaryl, saturated or partially saturated cycloalkyl, saturated or partially saturated heterocyclyl are further optionally substituted with one to multiple substituents, and the substituents are optionally selected from hydrogen, deuterium, halogen, alkyl, haloalkyl, cyano, cyanoethyl, O=, OH, C₁₋₃ alkyl, C₁₋₃ alkoxy, saturated or partially saturated cycloalkyl or saturated or partially saturated heterocyclyl, wherein the C₁₋₃ alkyl group, C₁₋₃ alkoxy group, saturated or partially saturated cycloalkyl group or saturated or partially saturated heterocyclic group are optionally substituted with one to three substituent groups, the substituent groups are optionally selected from H, deuterium, halogen, haloalkyl, cyano, OCH₃ and OH.

[0121] Each R₁ is the same or different and is independently selected from the group consisting of hydrogen, deuterium, halogen, —CN, —OH, —SH and —NH₂, —COOH or from C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ Alkynyl or C₁₋₁₀ alkoxy, C₂₋₁₀ heteroalkyl, C₃₋₁₀ saturated or partially saturated cycloalkyl, C₃₋₁₀ saturated or partially saturated heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl, C₂₋₁₀ heteroalkyl substituted by

C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocyclyl, C₁₋₁₀ alkyl substituted with carboxyl or carboxyl surrogate;

[0122] Each R₄ is the same or different and is optionally and independently selected from hydrogen, deuterium, halogen, CN, OH, SH and NH₂, —COOH or C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl, and C₁₋₁₀ heteroalkyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl; wherein, the C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, OH, and C₃₋₁₀ saturated or partially saturated cycloalkyl or heterocyclyl;

[0123] Each R₅ and R₆ is the same or different and is independently hydrogen, deuterium, halogen, CN, OH, SH, NR_{d8}R_{d9}, NH₂, —COOH or C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl, or C₃₋₁₀ heterocycloalkyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl; wherein, the C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl, or C₃₋₁₀ heterocycloalkyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, OH and C₃₋₁₀ saturated or partially substituted saturated cycloalkyl or heterocyclyl; or any adjacent two R₅ or R₆ together with the carbon to which they are attached form 5- to 6-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl, 3- to 8-membered saturated or partially saturated heterocyclyl, and where in the aryl, saturated or partially saturated cycloalkyl, heterocycloalkyl are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, —CN, —OH, CF₃, C₁₋₆ alkyl, C₁₋₆ alkoxy, —NH₂, —NHC₁₋₆ alkyl, —N(C₁₋₆ alkyl)₂, oxy, and saturated or partially saturated C₃₋₆ cycloalkyl, and the C₁₋₆ alkyl and C₁₋₆ alkoxy is optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, CF₃, OH, OCH₃, OCH₂CH₃, saturated or partially saturated C₃₋₆ cycloalkyl;

[0124] Each R_{d1}, R_{d2}, R_{d4}, R_{d8} and R_d is the same or different, and is optionally and independently selected from hydrogen, deuterium, NH₂, C₁₋₁₀ alkyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₁₋₁₀ alkylacyl, C₁₋₁₀ alkylsulfonyl, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl, or C₃₋₁₀ heterocyclyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl; wherein, the C₁₋₁₀ alkyl, C₂₋₁₀ alkynyl or C₁₋₁₀ alkoxy, C₁₋₁₀ alkylacyl, C₁₋₁₀ alkylsulfonyl, C₂₋₁₀ heteroalkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, C₁₋₁₀ alkyl substituted by C₃₋₁₀ cycloalkyl, or C₃₋₁₀ heterocyclyl substituted by C₃₋₁₀ cycloalkyl or C₃₋₁₀ heterocycloalkyl are optionally substituted with one to multiple groups selected from hydrogen, deuterium, halogen, oxo, CN, OH and C₃₋₁₀ saturated or partially substituted saturated cycloalkyl or heterocyclyl;

[0125] The hetero represents any heteroatom independently selected from O, N, S, P and isotopes thereof;

[0126] The halogen is optionally and independently selected from F, Cl, Br, I and isotopes thereof;


[0127] m is an integer optionally selected from 1, 2, 3 and 4;

[0128] n is an integer optionally selected from 0, 1, 2, 3, 4 and 5;

[0129] q is an integer optionally selected from 0, 1, 2, 3, 4 and 5;

[0130] t is an integer optionally selected from 0, 1, 2, 3, and 4.

[0131] In one embodiment of the present invention, the compound or a pharmaceutically acceptable salt, isotopic variations or isomer thereof has the structure of Formula (IG),



[0132] Wherein,

[0133] X and X' are independently selected from —C(R_{d1})(R_{d2})—, —N(R_{d4})—, —O—, —S—, —S(=O)— or —S(=O)₂—;

[0134] L is independently selected from —OC(R_{d1})(R_{d2})—, —C(R_{d1})(R_{d2})O—, —O—, —S— or —N(R_{d4})—;

[0135] X₁ and X₈ are each independently selected from —CR₅— or —N—;

[0136] R is independently hydrogen, deuterium, halogen, —CN, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ alkoxy, —NR_{d8}R_{d9}, 6- to 10-membered aryl, 5- to 8-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl and 3- to 8-membered saturated or partially saturated heterocyclyl, wherein R representing C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ alkoxy, 6- to 10-membered aryl, 5- to 8-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl and 3- to 8-membered saturated or partially saturated heterocyclyl are optionally substituted with one to multiple substituent groups, the substituent groups are optionally selected from hydrogen, deuterium, halogen, alkyl, haloalkyl, alkoxy, alkylamino, O=, CN, OH, —NR_{d8}R_{d9}, C₃₋₁₀ saturated or partially saturated cycloalkyl, C₃₋₁₀ saturated or partially saturated heterocyclic group, 6- to 10-aryl and 5- to 8-membered heteroaryl; wherein the aryl, heteroaryl, saturated or partially saturated cycloalkyl, saturated or partially saturated heterocyclyl are optionally substituted with one to multiple substituents, and the substituents are optionally selected from hydrogen, deuterium, halogen, alkyl, haloalkyl, cyano, cyanoethyl, O=, OH, C₁₋₃ alkyl, C₁₋₃ alkoxy, saturated or partially

saturated cycloalkyl or saturated or partially saturated heterocyclyl, wherein the C_{1-3} alkyl group, C_{1-3} alkoxy group, saturated or partially saturated cycloalkyl group or saturated or partially saturated heterocyclic group are optionally substituted with one to three substituent groups, the substituent groups are optionally selected from H, deuterium, halogen, haloalkyl, cyano, OCH_3 and OH.

[0137] Each R_1 is the same or different and is independently selected from the group consisting of hydrogen, deuterium, halogen, $-CN$, $-OH$, $-SH$ and $-NH_2$, $-COOH$ or from C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy, C_{2-10} heteroalkyl, C_{3-10} saturated or partially saturated cycloalkyl, C_{3-10} saturated or partially saturated heterocycloalkyl, C_{1-10} alkyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl, C_{2-10} heteroalkyl substituted by C_{3-10} cycloalkyl, C_{3-10} heterocyclyl, C_{1-10} alkyl substituted with carboxyl or carboxyl surrogate;

[0138] Each R_4 is the same or different and is optionally and independently selected from hydrogen, deuterium, halogen, CN, OH, SH and NH_2 , $-COOH$ or C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy, C_{2-10} heteroalkyl, C_{3-10} cycloalkyl, C_{3-10} heterocycloalkyl, C_{1-10} alkyl substituted C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl, and C_{1-10} heteroalkyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl; wherein, the C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, OH, and C_{3-10} saturated or partially saturated cycloalkyl or heterocyclyl;

[0139] Each R_5 and R_6 is the same or different and is independently hydrogen, deuterium, halogen, CN, OH, SH, $NR_{d8}R_{d9}$, NH_2 , $-COOH$ or C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy, C_{2-10} heteroalkyl, C_{3-10} cycloalkyl, C_{3-10} heterocycloalkyl, C_{1-10} alkyl substituted by C_{3-10} cycloalkyl, or C_{3-10} heterocyclyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl; wherein, the C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl or C_{1-10} alkoxy, C_{2-10} heteroalkyl, C_{3-10} cycloalkyl, C_{3-10} heterocycloalkyl, C_{1-10} alkyl substituted by C_{3-10} cycloalkyl, or C_{3-10} heterocycloalkyl are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, OH and C_{3-10} saturated or partially substituted saturated cycloalkyl or Heterocyclyl; Or any adjacent two R_5 or R_6 together with the carbon to which they are attached form 5- to 6-membered heteroaryl, 3- to 8-membered saturated or partially saturated cycloalkyl, 3- to 8-membered saturated or partially saturated heterocyclyl, and the aryl, saturated or partially saturated cycloalkyl, heterocycloalkyl are optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, $-CN$, $-OH$, CF_3 , C_{1-6} alkyl, C_{1-6} alkoxy, $-NH_2$, $-NHC_{1-6}$ alkyl, $-N(C_{1-6} \text{ alkyl})_2$, oxy, and saturated or partially saturated C_{3-6} cycloalkyl, and the C_{1-6} alkyl and C_{1-6} alkoxy is optionally substituted with one to more groups selected from hydrogen, deuterium, halogen, oxo, CN, CF_3 , OH, OCH_3 , OCH_2CH_3 , saturated or partially saturated C_{3-6} cycloalkyl;

[0140] Each R_{d1} , R_{d2} , R_{d4} , R_{d8} and R_d is the same or different, and is optionally and independently selected

from hydrogen, deuterium, NH_2 , C_{1-10} alkyl, C_{2-10} alkynyl or C_{1-10} alkoxy, C_{1-10} alkylacyl, C_{1-10} alkylsulfonyl, C_{2-10} heteroalkyl, C_{3-10} cycloalkyl, C_{3-10} heterocycloalkyl, C_{1-10} alkyl substituted by C_{3-10} cycloalkyl, or C_{3-10} heterocyclyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl; wherein, the C_{1-10} alkyl, C_{2-10} alkynyl or C_{1-10} alkoxy, C_{1-10} alkylacyl, C_{1-10} alkylsulfonyl, C_{2-10} heteroalkyl, C_{3-10} cycloalkyl, C_{3-10} heterocycloalkyl, C_{1-10} alkyl substituted by C_{3-10} cycloalkyl, or C_{3-10} heterocyclyl substituted by C_{3-10} cycloalkyl or C_{3-10} heterocycloalkyl are optionally substituted with one to multiple groups selected from hydrogen, deuterium, halogen, oxo, CN, OH and C_{3-10} saturated or partially substituted saturated cycloalkyl or heterocyclyl;

[0141] The hetero represents any heteroatom independently selected from O, N, S, P and isotopes thereof;

[0142] The halogen is optionally and independently selected from F, Cl, Br, I and isotopes thereof;

[0143] m is an integer optionally selected from 1, 2, 3 and 4;

[0144] n is an integer optionally selected from 0, 1, 2, 3, 4 and 5;

[0145] q is an integer optionally selected from 0, 1, 2, 3, 4 and 5;

[0146] t is an integer optionally selected from 0, 1, 2, 3, and 4.

[0147] According to an embodiment of the present invention,

[0148] A is selected from unsubstituted or substituted C_{6-20} aryl, 5- to 20-membered heteroaryl;

[0149] B is selected from unsubstituted or substituted 3- to 20-membered heterocyclyl, 6- to 18-membered spirocyclyl or bridged cyclyl, C_{6-20} aryl;

[0150] X and X' are the same or different and are independently selected from $-C(R_{d1})(R_{d2})-$, $-O-$, $-S-$, $-C(=O)O-$, $-OC(=O)-$, $-C(=O)-$, $-S(=O)-$ or $-S(=O)_2-$;


[0151] L is selected from $-C(R_{d1})(R_{d2})-$, $OC(R_{d1})(R_{d2})-$, $-C(R_{d1})(R_{d2})O-$, $-O-$, $-S-$, $-NH-$, $-C(=O)O-$, $-OC(=O)-$, $-C(=O)-$, $-S(=O)-$ or $-S(=O)_2-$; R_{d1} and R_{d2} are the same or different and optionally independently selected from hydrogen, deuterium, halogen, C_{1-10} alkyl, C_{1-10} alkoxy;

[0152] X_1 , X_2 , X_3 , X_4 , X_5 , X_8 and X_9 are the same or different and are independently selected from $-CR_5-$ or $-N-$; each R_5 is the same or different and is independently selected from hydrogen, deuterium, halogen, CN, OH, SH, NH_2 , $-COOH$, C_{1-10} alkyl, C_{1-10} alkoxy;


[0153] R_0 is selected from C_{1-10} alkyl unsubstituted or optionally substituted with one, two or more R_{oi} ; each R_{oi} is the same or different and is independently selected from C_{3-20} cycloalkyl, 3- to 20-membered heterocyclyl and 5- to 20-membered heteroaryl unsubstituted or optionally substituted with one, two or more R_{o2} ; each R_{o2} is the same or different, and is independently selected from halogen, deuterium, CN, oxo ($=O$), C_{1-10} alkyl, halogenated C_{1-10} alkyl, $CN-C_{1-10}$ alkyl, C_{3-6} cycloalkyl- C_{1-10} alkyl;

[0154] Each R_1 is the same or different and is independently selected from hydrogen, deuterium, halogen, $-CN$, $-OH$, $-SH$, $-NH_2$, $COOH$, C_{1-10} alkyl-


- COOH or $-C_{2-10}$ alkenyl-COOH unsubstituted or optionally substituted with one, two or more Ru; each Ru is the same or different and is independently selected from H, deuterium, halogen, C_{1-10} alkyl, C_{1-10} alkoxy;
- [0155] Each R_2 , R_2 , is the same or different, and is independently selected from hydrogen, deuterium, halogen, oxo ($=O$), C_{1-10} alkyl, C_{1-10} alkoxy;
- [0156] Each R_4 is the same or different and is optionally and independently selected from hydrogen, deuterium, halogen, oxo ($=O$), CN, OH, SH and NH_2 , $-COOH$, C_{1-10} alkyl, C_{1-10} alkoxy;
- [0157] Each R_6 is the same or different and is independently selected from the group consisting of hydrogen, deuterium, halogen, CN, OH, SH, NH_2 , $-COOH$, C_{1-10} alkyl, C_{1-10} alkoxy, C_{2-10} alkynyl, 5- to 14-membered heteroaryl;
- [0158] R_{d3} , R_{d4} , R_{d5} , R_{d6} , R_{d7} , R_{d8} , R_{d9} and R_{d10} are the same or different and are optionally and independently selected from hydrogen, deuterium, CN, OH, SH and NH_2 , $-COOH$, C_{1-10} alkyl, C_{1-10} alkoxy;
- [0159] m is an integer optionally selected from 1, 2, 3 and 4;
- [0160] n is an integer optionally selected from 0, 1, 2, 3, 4 and 5;
- [0161] q is an integer optionally selected from 0, 1, 2, 3, 4 and 5.
- [0162] According to an embodiment of the present invention, A is selected from unsubstituted or substituted C_{6-14} aryl, 5- to 14-membered heteroaryl; B is selected from unsubstituted or substituted 3- to 14-membered heterocycl, 6- to 18-membered spirocycl or bridged cycl, C_{6-14} aryl;
- [0163] X and X' are the same or different and are independently selected from $-C(R_{d1})(R_{d2})-$, $-O-$, $-S-$;
- [0164] L is selected from $-C(R_{d1})(R_{d2})-$, $-OC(R_{d1})(R_{d2})-$, $-C(R_{d1})(R_{d2})O-$, $-O-$, $-S-$, $-NH-$; R_{d1} and R_{d2} are the same or different, and are optionally and independently selected from hydrogen, deuterium, C_{1-10} alkyl;
- [0165] X_1 , X_2 , X_3 , X_4 , X_5 , X_8 and X_9 are the same or different, independently selected from $-CR_5-$ or $-N-$; each R_5 is the same or different, independently selected from hydrogen, deuterium, halogen, C_{1-10} alkyl, C_{1-10} alkoxy;
- [0166] R_0 is selected from C_{1-10} alkyl unsubstituted or optionally substituted with one, two or more Roi; each Roi is the same or different and is independently selected from C_{3-20} cycloalkyl, 3- to 20-membered heterocycl and 5- to 20-membered heteroaryl unsubstituted or optionally substituted with one, two or more R_{02} ; each R_{02} is the same or different, independently selected from CN, deuterium, oxo ($=O$), C_{1-10} alkyl, CN- C_{1-10} alkyl, C_{3-6} cycloalkyl- C_{1-10} alkyl;
- [0167] Each R_1 is the same or different and independently selected from, halogen, COOH, C_{1-10} alkyl-COOH, $-C_{2-10}$ alkenyl-COOH unsubstituted or optionally substituted with one, two or more Ru; each Ru is the same or different, independently selected from H, deuterium, halogen, C_{1-10} alkyl, C_{1-10} alkoxy;
- [0168] R_2 , R_2 , are the same or different, and are independently selected from hydrogen, deuterium, halogen, oxo ($=O$), C_{1-10} alkyl, C_{1-10} alkoxy;
- [0169] Each R_4 is the same or different and is optionally and independently selected from hydrogen, deuterium, halogen, oxo ($=O$), CN, OH, SH and NH_2 , $-COOH$, C_{1-10} alkyl, C_{1-10} alkoxy;
- [0170] Each R_6 is the same or different and is independently selected from the group consisting of hydrogen, deuterium, halogen, CN, C_{1-10} alkyl, C_{1-10} alkoxy, C_{2-10} alkynyl, 5- to 14-membered heteroaryl;
- [0171] m is an integer optionally selected from an integer of 1, 2 and 3;
- [0172] n is an integer optionally selected from 0, 1, 2, 3 and 4;
- [0173] q is an integer optionally selected from 0, 1, 2, 3 and 4.
- [0174] According to an embodiment of the present invention, A is selected from phenyl or pyridyl; B is piperidinyl, azetidiny, phenyl, (1R,5S)-3-azabicyclo[3.2.1]octyl, spirocyclic group or bridged cyclic group; the spirocyclic or the bridged ring contains one or more heteroatoms optionally and independently selected from N, O or S;
- [0175] X and X' are the same or different and are independently selected from CH_2O or S;
- [0176] L can be selected from O, S, NH, CH_2 , OCH_2 , CH_2O ;
- [0177] X_1 , X_2 , X_3 , X_4 , X_5 , X_8 and X_9 are the same or different and are independently selected from CH, C-F or N
- [0178] R_0 is selected from C_{1-3} alkyl unsubstituted or substituted with imidazolyl, pyrazolyl, pyrrolyl, azetidiny, oxetanyl, pyrrolidinyl or cyclopropyl; the imidazolyl, pyrazolyl, pyrrolyl, azetidiny, oxetanyl, pyrrolidinyl or cyclopropyl may be unsubstituted or substituted with oxo ($=O$) or C_{1-3} alkyl, CN- C_{1-3} alkyl, C_{3-6} cycloalkyl- C_{1-3} alkyl;
- [0179] Each R_1 is the same or different and is independently selected from COOH, F, $-C_{1-3}$ alkyl-COOH, $-C_{2-3}$ alkenyl-COOH;
- [0180] R_2 , R_2 , are the same or different, and are independently selected from hydrogen, deuterium, halogen, oxo ($=O$), C_{1-3} alkyl;
- [0181] Each R_4 is the same or different and is optionally and independently selected from hydrogen, deuterium, halogen, oxo ($=O$), C_{1-3} alkyl;
- [0182] Each R_6 is the same or different and is independently selected from hydrogen, deuterium, F, Cl, Br, CN, ethynyl, imidazolyl;
- [0183] m is optionally selected from 1 or 2;
- [0184] n is optionally selected from 0, 1 or 2;
- [0185] q is optionally selected from 0, 1 or 2.
- [0186] According to an embodiment of the present invention,




can be selected from the structures shown below:




-continued




[0187] B can have the following structure:



[0188] L can be selected from O, S, NH, CH₂, OCH₂, CH₂O;








can be selected from the structures shown below:





[0189] R_0 can be selected from the structures shown below:


[0190] Methyl,



-continued



[0191] R_1 can be selected from COOH, F,




[0192] Both R_2 and R_2' are H;




[0193] R_4 is H, F, methyl; X_8 is selected from CH or N; X_9 is selected from CH or N;

[0194] m is selected from 1 or 2; n is selected from 1 or 2.

[0195] In one embodiment of the present invention, the above-mentioned compound or a pharmaceutically acceptable salt thereof, or an enantiomer or isotopic variations thereof, is a compound selected from the following structures:



-continued


Number	Structure
Compound 2	
Compound 3	
Compound 4	

-continued


Number

Structure


Compound 5



Compound 6



Compound 7




-continued


Number

Structure


Compound 8



Compound 9



Compound 10




-continued


Number

Structure


Compound 11






Compound 12



Compound 13



-continued


Number	Structure
Compound 14	
Compound 15	
Compound 16	

-continued


Number

Structure


Compound 17



Compound 18



Compound 19




-continued


Number

Structure


Compound 20



Compound 21



Compound 22




-continued


Number

Structure


Compound 23



Compound 24



Compound 25




-continued


Number

Structure


Compound 26



Compound 27



Compound 28



-continued


Number	Structure
Compound 29	
Compound 30	
Compound 31	

-continued


Number

Structure


Compound 32



Compound 33



Compound 34




-continued


Number

Structure


Compound 35



Compound 36



Compound 37




-continued


Number

Structure


Compound 38



Compound 39



Compound 40




-continued


Number

Structure


Compound 41



Compound 42



Compound 43




-continued


Number

Structure


Compound 44



Compound 45



Compound 46




-continued


Number

Structure


Compound 47



Compound 48



Compound 49




-continued


Number

Structure


Compound 50



Compound 51



Compound 52




-continued


Number

Structure


Compound 53



Compound 54



Compound 55




-continued


Number

Structure


Compound 56



Compound 57



Compound 58




-continued


Number

Structure


Compound 59



Compound 60



Compound 61




-continued


Number

Structure


Compound 62



Compound 63



Compound 64




-continued


Number

Structure


Compound 65



Compound 66



Compound 67




-continued


Number

Structure


Compound 68



Compound 69



Compound 70




-continued


Number

Structure


Compound 71



Compound 72



Compound 73




-continued


Number

Structure


Compound 74



Compound 75



Compound 76




-continued


Number

Structure


Compound 77



Compound 78



Compound 79




-continued


Number

Structure


Compound 80



Compound 81



Compound 82




-continued


Number

Structure


Compound 83



Compound 84



Compound 85




-continued


Number

Structure


Compound 86



Compound 87



Compound 88




-continued


Number

Structure


Compound 89



Compound 90



Compound 91




-continued


Number

Structure


Compound 92



Compound 93



Compound 94




-continued


Number

Structure


Compound 95



Compound 96



Compound 97




-continued


Number

Structure


Compound 98



Compound 99



Compound 100




-continued


Number

Structure


Compound 101



Compound 102



Compound 103




-continued


Number

Structure


Compound 104



Compound 105



Compound 106




-continued


Number

Structure


Compound 107



Compound 108



Compound 109




-continued


Number

Structure


Compound 110



Compound 111



Compound 112




-continued


Number

Structure


Compound 113



Compound 114



Compound 115




-continued


Number

Structure


Compound 116



Compound 117



Compound 118




-continued


Number

Structure


Compound 119



Compound 120



Compound 121




-continued


Number

Structure


Compound 122



Compound 123



Compound 124




-continued


Number

Structure


Compound 125



Compound 126



Compound 127




-continued


Number

Structure


Compound 128



Compound 129



Compound 130




-continued


Number

Structure


Compound 131



Compound 132



Compound 133




-continued


Number

Structure


Compound 134



Compound 135



Compound 136




-continued


Number

Structure


Compound 137



Compound 138



Compound 139




-continued


Number

Structure


Compound 140



Compound 141



Compound 142




-continued


Number

Structure


Compound 143



Compound 144



Compound 145




-continued


Number

Structure


Compound 146



Compound 147



Compound 148




-continued


Number

Structure


Compound 149



Compound 150



Compound 151




-continued


Number

Structure


Compound 152



Compound 153



Compound 154




-continued


Number

Structure


Compound 155



Compound 156



Compound 157




-continued


Number

Structure


Compound 158



Compound 159



Compound 160




-continued


Number

Structure


Compound 161



Compound 162



Compound 163




-continued


Number

Structure


Compound 164



Compound 165



Compound 166




-continued


Number

Structure


Compound 167



Compound 168



Compound 169




-continued


Number

Structure


Compound 170



Compound 171



Compound 172




-continued


Number

Structure


Compound 173



Compound 174



Compound 175




-continued


Number

Structure


Compound 176



Compound 177



Compound 178




-continued


Number

Structure


Compound 179



Compound 180



Compound 181




-continued


Number

Structure


Compound 182



Compound 183



Compound 184




-continued


Number

Structure


Compound 185



Compound 186



Compound 187




-continued


Number

Structure


Compound 188



Compound 189



Compound 190




-continued


Number

Structure


Compound 191



Compound 192



Compound 193




-continued


Number

Structure


Compound 194



Compound 195



Compound 196




-continued


Number

Structure


Compound 197



Compound 198



Compound 199




-continued


Number

Structure


Compound 200



Compound 201



Compound 202




-continued


Number

Structure


Compound 203



Compound 204



Compound 205




-continued


Number

Structure


Compound 206



Compound 207



Compound 208




-continued


Number

Structure


Compound 209



Compound 210



Compound 211




-continued


Number

Structure


Compound 212



Compound 213



Compound 214




-continued


Number

Structure


Compound 215



Compound 216



Compound S1




-continued


Number

Structure


Compound S2



Compound S3



Compound S4




-continued


Number

Structure


Compound S5



Compound S6



Compound S7




-continued


Number

Structure


Compound S8



Compound S9



Compound S10




-continued


Number

Structure


Compound S11



Compound S14



Compound S15



-continued


Structure

-continued


Number

Structure


Compound S21



Compound S22



Compound S23




-continued


Number

Structure


Compound S24



Compound S25



Compound S26




-continued


Number

Structure


Compound S27



Compound S28



Compound S29




-continued


Number

Structure


Compound S32



Compound S33



Compound S34




-continued


Number

Structure


Compound S35



Compound S36



Compound S37




-continued


Number

Structure


Compound S40



Compound S41



Compound S42




-continued


Number

Structure


Compound S43



Compound S44



Compound S45




-continued


Number

Structure


Compound S48



Compound S49



Compound S50




-continued


Number

Structure


Compound S51



Compound S52



Compound S53




-continued


Number

Structure


Compound S56



Compound S57



Compound S58




-continued


Number

Structure


Compound S59



Compound S60



Compound S61




-continued


Number

Structure


Compound S62



Compound S63



Compound S64




-continued


Number

Structure


Compound S65



Compound S66



Compound C1




-continued


Number

Structure


Compound C2



Compound C3



Compound C4




-continued


Number

Structure


Compound C5



Compound C6



Compound C7




-continued


Number

Structure


Compound C8



Compound C9



Compound C10




-continued


Number

Structure


Compound C11



Compound C12



Compound C13




-continued


Number

Structure


Compound C14



Compound C15



Compound C16




-continued


Number

Structure


Compound C17



Compound C18



Compound C19



-continued


Number	Structure
Compound C20	
Compound C21	
Compound C22	

-continued


Number

Structure


Compound C23



Compound C24



Compound C25




-continued


Number

Structure


Compound C26



Compound C27



Compound C28




-continued


Number

Structure


Compound C29



Compound C30



Compound C31




-continued


Number

Structure


Compound C32



Compound C33



Compound C34




-continued


Number

Structure


Compound C35



Compound C36



Compound C37




-continued


Number

Structure


Compound C38



Compound C39



Compound C40




-continued


Number

Structure


Compound C41



Compound C42



Compound C43




-continued


Number

Structure


Compound C44



Compound C45



Compound C46




-continued


Number

Structure


Compound C47



Compound C48



Compound C49




-continued


Number

Structure


Compound C50



Compound C51



Compound C52




-continued


Number

Structure


Compound C53



Compound C54



Compound C55




-continued


Number

Structure


Compound C56



Compound C57



Compound C58




-continued


Number

Structure


Compound C59



Compound C60



Compound C61




-continued


Number

Structure


Compound C62



Compound C63



Compound C64




-continued


Number

Structure


Compound C65



Compound C66



Compound C67




-continued


Number

Structure


Compound C68



Compound C69



Compound C70




-continued


Number

Structure


Compound C71



Compound C72



Compound C73




-continued


Number

Structure


Compound C74



Compound C75



Compound C76




-continued


Number

Structure


Compound C77



Compound C78



Compound C79




-continued


Number

Structure


Compound C80





Compound C81



Compound C82



-continued

Number	Structure
Compound C83	
Compound C84	

[0196] The present invention also provides a pharmaceutical composition comprising a therapeutically effective amount of at least one of the compounds represented by Formula (I), its pharmaceutically acceptable salts, solvates, enantiomers, and isotopic variations.

[0197] According to an embodiment of the invention, the pharmaceutical composition is formulated for administration by a route selected from the group consisting of oral, injection, rectal, nasal, pulmonary, topical, buccal and sublingual, vaginal, parenteral, subcutaneous, intramuscular, intravenous, intradermal, intrathecal, and epidural.

[0198] According to an embodiment of the present invention, the pharmaceutical composition is preferably administered orally.

[0199] The oral dosage form is not particularly limited, and any oral dosage form well known in the art can be used, preferably including tablets, capsules, suspensions or oral solutions and other oral dosage forms known in the art. As an oral dosage form, the dosage standard used is, for example, 500-1500 mg/day, preferably 700-1200 mg/day, preferably 800-1000 mg/day, most preferably 1000 mg/day.

[0200] The administration time of the pharmaceutical composition according to the present invention may depend on the degree of the disease, preferably at least 1 month, for

example, 1, 2, 3, 4, 5, or 6 months, and the longest may be lifelong medication due to the needs of the disease.

[0201] According to an embodiment of the present invention, the pharmaceutical composition may further comprise pharmaceutically acceptable excipients selected from at least one including but not limited to the following excipients: filler, disintegrant, binder, lubricant, surfactants, flavoring agents, wetting agents, pH regulators, solubilizers or cosolvents, osmotic pressure regulators. Those skilled in the art can easily determine how to select the corresponding excipients and their corresponding amounts according to the needs of specific dosage forms.

[0202] According to embodiments of the present invention, the pharmaceutical composition may further contain one or more additional therapeutic agents.

[0203] Another object of the present invention is to provide use of the above-mentioned compounds for manufacturing a medicament for preventing or treating GLP1/GLP1R signaling pathway related diseases. The GLP1/GLP1R signaling pathway related diseases include but are not limited to overweight, obesity, diabetes (T1 D and/or T2DM, including prediabetes), idiopathic T1 D (type 1B), latent autoimmune diabetes in adults (LADA), early-onset T2DM (EOD), youthful atypical diabetes mellitus (YOAD), maturity-onset diabetes mellitus (MODY), malnutrition-re-

lated diabetes mellitus, gestational diabetes mellitus, hyperglycemia, insulin resistance, hepatic insulin resistance, glucose tolerance Impairment, diabetic neuropathy, diabetic nephropathy, renal disease (e.g., acute kidney disease, renal tubular dysfunction, proinflammatory changes in proximal tubules), diabetic retinopathy, adipocyte dysfunction, visceral fat deposition, sleep apnea suspension, obesity (including hypothalamic and monogenic obesity) and associated comorbidities (e.g., osteoarthritis and urinary incontinence), eating disorders (including binge drinking syndrome, bulimia nervosa, and syndromic obesity such as Prader-Willi and Bardet-Biedl syndrome), weight gain due to use of other drugs (e.g., steroid and antipsychotic use), hyperglycemia, dyslipidemia (including hyperlipidemia, hypertriglyceridemia, total Increased cholesterol, high-density lipoprotein cholesterol, low-density lipoprotein cholesterol, hyperinsulinemia, NAFLD (including steatosis, NASH, fibrosis, liver cirrhosis, hepatocellular carcinoma and other related diseases), cardiovascular disease, atherosclerosis (including coronary artery disease), peripheral vascular disease, hypertension, endothelial dysfunction, impaired vascular compliance, congestive heart failure, myocardial infarction (e.g., necrosis and apoptosis), stroke, hemorrhagic stroke, ischemic stroke, Traumatic brain injury, pulmonary hypertension, restenosis after angioplasty, intermittent claudication, postprandial lipids, metabolic acidosis, ketosis, arthritis, osteoporosis, Parkinson's disease, left ventricular hypertrophy, peripheral artery disease, macular degeneration, cataract, glomerulosclerosis, chronic renal failure, metabolic syndrome, syndrome X, premenstrual syndrome, angina pectoris, thrombosis, atherosclerosis, transient ischemic attack, restenosis, sugar Impaired metabolism, impaired fasting glucose, hyperuricemia, gout, erectile dysfunction, skin and connective tissue disorders, psoriasis, foot ulcers, ulcerative colitis, hyperapolipoprotein B lipoproteinemia, Alzheimer's disease Prophylaxis or treatment of silent disease, schizophrenia, cognitive impairment, inflammatory bowel disease, short bowel syndrome, Crohn's disease, colitis, irritable bowel syndrome, polycystic ovary syndrome, and addiction treatment (e.g., alcohol and/or drug abuse) and other conditions.

[0204] The present invention also provides the use of the compound represented by the Formula (I), its pharmaceutically acceptable salts, solvates, enantiomers and isotopic variations, and the pharmaceutical compositions in the prevention and/or treatment of diseases related to GLP1/GLP1R signaling pathway. The GLP1/GLP1R signaling pathway-related diseases have the above-mentioned definitions.

[0205] The present invention also provides a method for preventing and/or treating diseases related to the GLP1/GLP1R signaling pathway, comprising administering to a patient a preventive or therapeutically effective amount of the compound represented by Formula (I), a pharmaceutically acceptable salt, a solvate, enantiomers and isotopic variations, thereof at least one of the compounds, or administering to the patient a prophylactically or therapeutically effective amount of the above-mentioned pharmaceutical composition. The GLP1/GLP1R signaling pathway-related diseases have the above-mentioned definitions.

[0206] In some embodiments, the patient is a mammal, preferably a human.

BRIEF DESCRIPTION OF THE DRAWINGS

[0207] FIG. 1 shows _comparison curve of the effects of the compounds of the present invention and known drugs on the IPGTT blood sugar of hGLP1R mice.

[0208] FIG. 2 shows comparison of the effects of the compounds of the present invention and known drugs on the insulin-stimulating release of IPGTT in hGLP1R mice.

[0209] FIG. 3 shows comparison curve of the effect of the compounds of the present invention and known drugs on blood glucose of cynomolgus monkey IVGTT.

DETAILED DESCRIPTION OF THE INVENTION

Definition and Description

[0210] C_{1-10} is selected from $C_1, C_2, C_3, C_4, C_5, C_6, C_7, C_8, C_9$ and C_{10} ; C_{2-10} is selected from $C_2, C_3, C_4, C_5, C_6, C_7, C_8, C_9$ and C_{10} ; C_{3-10} is selected from $C_3, C_4, C_5, C_6, C_7, C_8, C_9$ and C_{10} ;

[0211] The term 'alkyl' as used herein refers to a linear or branched chain monovalent hydrocarbon group. Examples include, but are not limited to, methyl, ethyl, propyl, butyl, 2-methyl-propyl, 1,1-dimethylethyl, pentyl, and hexyl.

[0212] The term 'alkylene' as used herein refers to a linear or branched chain divalent hydrocarbon group of formula $-(CH_2)_n-$. Examples include, but are not limited to, ethylene and propylene.

[0213] The term 'one to more' or 'one to multiple' as used herein refers to more than 1, such as 1, 2, 3, 4, 5 or more.

[0214] The term 'carbocycle (radical)' or "cycloalkyl" refers to a saturated or partially unsaturated monocyclic or polycyclic hydrocarbon group, the carbocycle may contain 3 to 20 carbon atoms, preferably 3 to 12 (e.g. 3, 4, 5, 6, 7, 8, 9, 10, 11, 12) carbon atoms, more preferably 3 to 6 carbon atoms. The carbocycle may be monocyclic or polycyclic, it may be a saturated cycloalkyl or may optionally contain one, two or more double and/or triple bonds in its ring, thereby forming so-called cycloalkenyl or cycloalkynyl. Where carbocycles have multiple rings, the rings can form spiro, fused and bridged ring structures. For example, non-limiting examples of monocyclic carbocycles include cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, cyclohexadienyl, cycloheptyl, cycloheptyl alkenyl, cyclooctyl, cyclooctatetraenyl, etc.; non-limiting examples of polycyclic carbocycles include decalin or isobornyl.

[0215] The term 'heterocycle (radical)' refers to a saturated or partially unsaturated monocyclic or polycyclic cyclic hydrocarbon substituent containing 3 to 20 ring atoms, wherein one or more of the ring atoms is a heteroatom or group of atoms selected from N, O, NH, S, S(O) or S(O)₂, but excluding ring parts of $-O-O-$, $-O-S-$ or $-S-S-$, the remaining ring atoms are carbon. Preferably 3 to 12 ring atoms, of which 1 to 4 are heteroatoms, are comprised (e.g. 1, 2, 3 and 4); more preferably 3 to 6 ring atoms (e.g. 3, 4, 5, 6). A heterocyclyl group can be attached to the remainder of the molecule through any of the carbon atoms or a nitrogen atom (if present) or an oxygen or sulfur atom (especially in the case of an onium salt formation). The heterocyclyl group may include fused or bridged rings and/or spiro rings. Non-limiting examples of monocyclic heterocyclyl groups include azetidiny, oxetanyl, pyrrolidinyl, imidazolidinyl, tetrahydrofuranly, tetrahydrothienyl,

dihydroimidazolyl, dihydrofuranyl, dihydropyrazolyl, dihydropyrrolyl, dioxolyl, tetrahydropyranyl, pyrroline, piperidinyl, piperazinyl, morpholinyl, thiomorpholinyl, dithianyl trithianyl, homopiperazinyl, diazepanyl, etc., preferably piperidinyl and pyrrolidinyl.

[0216] Polycyclic heterocyclyls include spiro, fused and bridged heterocyclyls, and may also be benzo-fused heterocyclyls such as dihydroisoquinolinyl. The heterocyclyl may be bicyclic, non-limiting examples of which include hexahydrocyclopento[c]pyrrol-2(1H)-yl, hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl. Heterocyclyl can also be partially unsaturated, i.e. it can contain one or more double bonds, non-limiting examples of which include dihydrofuranyl, dihydropyranyl, 2,5-dihydro-1H-pyrrolyl, 4H-[1,3,4]thiadiazinyl, 4,5-dihydrooxazolyl or 4H-[1,4]thiazinyl.

[0217] Heterocyclyl may be optionally substituted or unsubstituted, and when substituted, the substituents are preferably one or more of the following groups independently selected from alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, halogen, mercapto, hydroxyl, nitro, cyano, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, cycloalkoxy, heterocycloalkoxy, cycloalkaneithio, heterocycloalkylthio, oxo, carboxyl or carboxylate.

[0218] As used herein, the term 'heteroaryl/heteroaromatic ring' refers to a heteroaromatic system comprising 1 to 4 heteroatoms, 5 to 20 ring atoms, wherein the heteroatoms are selected from oxygen, sulfur, nitrogen and phosphorus. Heteroaryl is preferably 5- to 10-membered (e.g. 5-, 6-, 7-, 8-, 9- or 10-membered), more preferably 5- or 6-membered. Non-limiting examples of heteroaryl groups include, but are not limited to, thienyl, furyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl oxazolyl, thiadiazolyl, thi-4H-pyrazolyl, etc. and their benzo derivatives such as benzofuranyl, benzothienyl, benzoxazolyl, benzisoxazolyl, benzoyl imidazolyl, benzotriazolyl, indazolyl, indolyl, isindolyl, etc.; or pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, etc., and their benzo derivatives, such as quinolinyl, quinazolinyl, isoquinolinyl, etc; or azinyl, indozinyl, purinyl, etc., as well as their benzo derivatives; or cinolinyl, phthalazinyl, quinazolinyl, quinoxalinyl, naphthyridinyl, pteridinyl, carbazolyl, acridinyl, phenazinyl, phenothiazinyl, and/or phenoxazinyl, etc.

[0219] Heteroaryl/heteroaryl rings may be optionally substituted or unsubstituted, when substituted, the substituents are preferably one, two or more groups independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, halogen, mercapto, hydroxyl, nitro, cyano, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, cycloalkoxy, heterocycloalkoxy, cycloalkylthio, heterocycloalkylthio, carboxyl or carboxylate.

[0220] Unless otherwise stated, a heterocyclyl, heteroaryl or heteroaryl ring includes all possible isomeric forms thereof, e.g. positional isomers thereof. Thus, for some illustrative non-limiting examples, can be included in its 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8-, 9-, 10-, 11-, 12- and etc. (if present) substituted at one, two or more positions, or bonded to other groups, including pyridin-2-yl, pyridin-2-yl, pyridin-3-yl, Pyridin-3-yl, pyridin-4-yl and pyridin-4-yl; thienyl or thienylene includes thien-2-yl, thien-2-yl, thien-3-yl and thien-3-base; pyrazol-1-yl, pyrazol-3-yl, pyrazol-4-yl, pyrazol-5-yl.

[0221] As used herein, the term 'pharmaceutically acceptable' refers to those compounds, materials, compositions

and/or dosage forms that, within the scope of sound medical judgment, are suitable for use in contact with human and animal tissue, without excessive toxicity, irritation, allergic reactions or other problems or complications, commensurate with a reasonable benefit/risk ratio.

[0222] The term 'pharmaceutically acceptable salts' refers to salts of the compounds of the present invention, prepared from compounds with specific substituents discovered by the present invention and relatively non-toxic acids or bases. When the compounds of the present invention contain relatively acidic functional groups, base addition salts can be obtained by contacting the neutral forms of such compounds with a sufficient amount of base in neat solution or in a suitable inert solvent. Pharmaceutically acceptable base addition salts include sodium, potassium, calcium, ammonium, organic ammonia or magnesium salts or similar salts. When the compounds of the present invention contain relatively basic functional groups, acid addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of acid in neat solution or in a suitable inert solvent. Examples of pharmaceutically acceptable acid addition salts include inorganic acid salts including, for example, hydrochloric acid, hydrobromic acid, nitric acid, carbonic acid, bicarbonate, phosphoric acid, monohydrogen phosphate, dihydrogen phosphate, sulfuric acid, hydrogen sulfate, hydroiodic acid, phosphorous acid, etc.; and organic acid salts including, for example, acetic acid, propionic acid, isobutyric acid, maleic acid, malonic acid, benzoic acid, succinic acid, suberic acid, fumaric acid, lactic acid, mandelic acid, ortho Phthalic acid, benzenesulfonic acid, p-toluenesulfonic acid, citric acid, tartaric acid, methanesulfonic acid and similar acids; also includes salts of amino acids such as arginine, etc., and salts of organic acids such as glucuronic acid (See Berge et al., "Pharmaceutical Salts", Journal of Pharmaceutical Science 66: 1-19 (1977)). Certain specific compounds of the present invention contain both basic and acidic functional groups and thus can be converted into either base or acid addition salts.

[0223] The neutral forms of the compounds are preferably regenerated by contacting the salt with a base or acid and isolating the parent compounds in the conventional manner. The parent form of the compound differs from the various salt forms in certain physical properties, such as solubility in polar solvents.

[0224] As used herein, "pharmaceutically acceptable salts" pertain to derivatives of compounds of the present invention wherein the parent compound is modified by salt formation with an acid or salt formation with a base. Examples of pharmaceutically acceptable salts include, but are not limited to, inorganic or organic acid salts of bases such as amines, alkali metal or organic salts of acid groups such as carboxylic acids, and the like. Pharmaceutically acceptable salts include conventional non-toxic salts such as Na salts, potassium salts, amine salts, quaternary ammonium salts of the parent compound, and the like. Conventional non-toxic salts include, but are not limited to, those derived from inorganic and organic acids, inorganic and organic bases selected from the group consisting of 2-acetoxybenzoic acid, 2-hydroxyethanesulfonic acid, acetic acid, ascorbic acid, benzenesulfonic acid, benzoic acid, bicarbonate, carbonic acid, citric acid, edetic acid, ethanedisulfonic acid, ethanesulfonic acid, fumaric acid, glucoheptose, gluconic acid, glutaric acid Amino acid, glycolic acid, hydrobromic acid, hydrochloric acid, hydroiodide, hydroxy,

hydroxynaphthalene, isethionic acid, lactic acid, lactose, dodecylsulfonic acid, maleic acid, malic acid, mandelic acid, methanesulfonic acid, nitric acid, oxalic acid, pamoic acid, pantothenic acid, phenylacetic acid, phosphoric acid, polygalacturonic acid, propionic acid, salicylic acid, stearic acid, acetic acid, succinic acid, sulfamic acid, p-aminobenzenesulfonic acid, sulfuric acid, tannin, tartaric acid and p-toluenesulfonic acid etc. described inorganic base and organic base are selected from Na, potassium, magnesium, calcium etc. or amine, diethylamine, triethylamine, ethanolamine etc.

[0225] The pharmaceutically acceptable salts of the present invention can be synthesized from the acid or base containing parent compound by conventional chemical methods. Generally, such salts are prepared by reacting the free acid or base form of these compounds with a stoichiometric amount of the appropriate base or acid in water or an organic solvent or a mixture of the two. Generally, non-aqueous media such as ether, ethyl acetate, ethanol, isopropanol or acetonitrile are preferred.

[0226] In addition to salt forms, the compounds provided herein also exist in prodrug forms. Prodrugs of the compounds described herein are readily chemically altered under physiological conditions to convert to the compounds of the present invention. In addition, prodrugs can be converted to the compounds of the present invention by chemical or biochemical methods in an in vivo environment.

[0227] Certain compounds of the present invention may exist in unsolvated as well as solvated forms, including hydrated forms. In general, solvated and unsolvated forms are equivalent and are intended to be included within the scope of the present invention. Certain compounds of the present invention may exist in multiple crystalline or amorphous forms.

[0228] As used herein, the term 'solvate' refers to an association of one or more solvent molecules with a compound of the present invention. Solvate-forming solvents include, but are not limited to, water, isopropanol, ethanol, methanol, dimethyl sulfoxide, ethyl acetate, acetic acid, and aminoethanol. Thus, the term 'hydrate' refers to an association in which the solvent molecule is water.

[0229] Certain compounds of the present invention may have asymmetric carbon atoms (optical centers) or double bonds. Racemates, diastereomers, geometric isomers and individual isomers are included within the scope of the present invention.

[0230] Schematic representations of racemates, ambiscalemic and scalemic or enantiomerically pure compounds herein are from Maehr, J. Chem. Ed. 1985, 62: 114-120. 1985, 62: 114-120. Unless otherwise stated, wedge and dashed bonds are used to indicate the absolute configuration of a stereocenter. When the compounds described herein contain olefinic double bonds or other centers of geometric asymmetry, they include E, Z geometric isomers unless otherwise specified. Likewise, all tautomeric forms are included within the scope of the present invention.

[0231] The compounds of the present invention may exist in specific geometric or stereoisomeric forms. The present invention contemplates all such compounds, including cis and trans isomers, (-)- and (+)-enantiomers, (R)- and (S)-enantiomers, diastereomers isomers, (D)-isomers, (L)-isomers, and racemic and other mixtures thereof, such as enantiomerically or diastereomerically enriched mixtures, all of which belong to within the scope of the present invention. Additional asymmetric carbon atoms may be

present in substituents such as alkyl. All such isomers, as well as mixtures thereof, are included within the scope of the present invention.

[0232] Optically active (R)- and (S)-isomers as well as D and L isomers can be prepared by chiral synthesis or chiral reagents or other conventional techniques. If one enantiomer of a compound of the present invention is desired, it can be prepared by asymmetric synthesis or derivatization with a chiral auxiliary, wherein the resulting mixture of diastereomers is separated and the auxiliary group is cleaved to provide pure desired enantiomer. Alternatively, when the molecule contains basic functional groups (such as amino groups) or acidic functional groups (such as carboxyl groups), diastereomeric salts are formed with appropriate optically active acids or bases, followed by stepwise steps well known in the art. The diastereoisomers are resolved by crystallization or chromatography, followed by recovery of the pure enantiomers. In addition, separation of enantiomers and diastereomers is usually accomplished by the use of chromatography using a chiral stationary phase, optionally in combination with chemical derivatization (e.g., carbamate formation from amines).

[0233] The compounds of the present invention may contain unnatural proportions of atomic isotopes at one or more of the atoms that constitute the compound. For example, compounds may be labeled with radioactive isotopes, such as tritium (^3H), iodine-125 (^{125}I) or C-14 (^{14}C). All isotopic variations of the compounds of the invention, whether radioactive or not, are intended to be encompassed within the scope of the invention.

[0234] The term "pharmaceutically acceptable carrier" refers to any formulation or carrier medium that provides the appropriate delivery of an effective amount of an active agent as defined herein, does not negatively interfere with the effectiveness of the biological activity of the active agent, and that is sufficiently non-toxic to the host. Representative carriers include water, oils, both vegetable and mineral, cream bases, lotion bases, ointment bases and the like. These bases include suspending agents, thickeners, penetration enhancers, and the like. Their formulation is well known to those in the art of topical pharmaceuticals. Additional information concerning carriers can be found in Remington: The Science and Practice of Pharmacy. 21st Ed., Lippincott, Williams & Wilkins (2005) which is incorporated herein by reference.

[0235] When any variable (e.g., R) occurs more than once in the composition or structure of the compound, its definition in each case is independent. Thus, for example, if a group is substituted with 0-2 R, the group may optionally be substituted with up to two R, and in each case the R has an independent option. In addition, the combination of substituents and/or their variants is allowed only if such a combination will produce a stable compound.

[0236] When a bond of a substituent could be cross-linked to two atoms on a ring, such substituent may be bonded to any atoms in the ring. Where the listed substituent does not specify through which atom it is connected to the general structure formula including the compound that is not specifically mentioned, the substituent may be bonded through any one of its atoms. The combination of substituents and/or variant thereof is allowed only if such a combination results in a stable compound.

[0237] The term "halo" or 'halogen' refers to fluorine, chlorine, bromine and iodine.

[0238] In order to illustrate the disclosure in more detail, the following examples are given, but the scope of the disclosure is not limited thereto. The compounds of the disclosure can be prepared by various synthetic methods well-known to those skilled in the art, including the embodiments described below, the embodiments combining the embodiments described below with other synthetic methods, and equivalent alternatives known to those skilled in the art. Preferred embodiments include but are not limited to those embodiments of the present invention.

[0239] Unless otherwise specified, all solvents used in the present invention are commercially available, and no further purification is required for use. The reaction is usually carried out using an anhydrous solvent under an inert atmosphere of nitrogen. NMR spectra were measured on a Bruker-Avance-400 (400 MHz) spectrometer and chemical shifts are reported in δ (ppm). Mass spectrometry was performed with an Agilent 1200 series (plus 6110/and 1956A) LC/MS or a Shimadzu MS (DAD: SPD-M20A (LC)) and a Shimadzu Micromass 2020 detector. The mass spectrometer is equipped with an electrospray ionization (ESI) source operating in positive and negative modes.

[0240] Abbreviations used are as follows: aq is aqueous; TLC is thin layer chromatography; RT is room temperature; MeOH is methanol; EtOH is ethanol; EtOAc is ethyl acetate; THF is tetrahydrofuran; eq for equivalent or equivalents; CDI is carbonyldiimidazole; DCM is Dichloromethane; PE is petroleum ether; DIAD is diisopropyl azodicarboxylate; DMF is N,N-dimethylformamide; DMSO is dimethyl sulfoxide; CBz is benzyloxycarbonyl; Boc is tert-butyl; HOAc is acetic acid; Ms is methylsulfonyl; NMP is N-methylpyrrolidone; DMAP is 4-(dimethylamino)pyridine; Boc₂O is di-tert-butyl dicarbonate; TFA is trifluoroacetic acid; DIPEA is diisopropylethylamine; SOCl₂ is thionyl chloride; CS₂ is carbon disulfide; TsOH is 4-toluenesulfonic acid; MTBE is tert-butyl methyl ether; FA is formic acid; ACN is acetonitrile; i-PrOH is 2-propanol.


[0241] Compounds can be named manually or by ChemDraw® software, or if purchased commercially the supplier's catalog name is used. Usually, TLC or LC-MS is used to determine whether the reaction is completed.

EXAMPLES

[0242] In order to illustrate the present invention in more detail, the following examples are given, but the scope of the present invention is not limited thereto.

Example 1-1. Synthesis of methyl (S,E)-3-(2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazol-6-yl)acrylate (intermediate A-1)


1). (E) Preparation of methyl (E)-3-(3-fluoro-4-nitrophenyl)acrylate



[0243] To a solution of methyl diethylphosphonoacetate (1 g, 4.76 mmol) in dry THF (10 mL) at 0° C. sodium hydride (60% w/w dispersed in mineral oil, 210 mg, 5.24 mmol) was added. The mixture was stirred at 0° C. for 30 mins, then 3-fluoro-4-nitrobenzaldehyde (885 mg, 5.24 mmol) was added slowly to above mixture, maintaining the reaction temperature around 0° C. After the addition was completed, the mixture was naturally warmed to room temperature and stirred for 16 hours. Cool to 0° C., the mixture was quenched with water (50 mL) and extracted with ethyl acetate (2×30 mL). The combined organic layer was concentrated in vacuo to give a residue. The residue was subjected to silica gel column chromatography (PE/EA=3/1) to obtain methyl (E)-3-(3-fluoro-4-nitrophenyl)acrylate (800 mg, 74.8% yield).


[0244] ¹H NMR (400 MHz, DMSO-d₆): δ 8.18 (t, J=8.2 Hz, 1H), 8.05 (d, J=12.8 Hz, 1H), 7.80 (d, J=8.6 Hz, 1H), 7.73 (d, J=16.0 Hz, 1H), 6.93 (d, J=16.0 Hz, 1H), 3.76 (s, 3H).

2). Synthesis of methyl (S,E)-3-(4-nitro-3-((oxetan-2-ylmethyl)amino)phenyl)acrylate




[0245] The mixture of methyl (E)-3-(3-fluoro-4-nitrophenyl)acrylate (300 mg, 1.33 mmol), (S)-oxetan-2-ylmethanamine (116 mg, 1.33 mmol) and K₂CO₃ (368 mg, 2.67 mmol) in N,N-dimethylformamide (5 mL) was stirred at room temperature for 16 hours. The mixture was diluted with water (50 mL) and extracted with EA (3×10 mL). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered, and concentrated in vacuo to give a residue. The residue was purified by pre-TLC (PE/EA=1/1) to give methyl (S,E)-3-(4-nitro-3-((oxetan-2-ylmethyl)amino)phenyl)acrylate (320 mg, 82.4% yield), LC-MS m/z: 293 [M+H]⁺.

3). Synthesis of methyl (S,E)-3-(4-amino-3-((oxetan-2-ylmethyl)amino)phenyl)acrylate




-continued



[0246] The mixture of methyl (S,E) (4-nitro-3-((oxetan-2-ylmethyl)amino)phenyl)acrylate (Intermediate 2) (320 mg, 1.1 mmol), Fe (173 mg) and NH_4Cl (164 mg) in ethanol/water (v:v=10/1, 5 mL) was heated to 80° C. and stirred for 16 hours. The reaction mixture was cooled to room temperature, poured into saturated NaHCO_3 (30 mL), and then extracted with EA (3×10 mL). The combined organic layer was concentrated in vacuo to give a residue. The residue was purified by silica gel column chromatography (DCM/MeOH=20/1) to obtain methyl (S, E)-3-(4-amino-3-((oxetan-2-ylmethyl)amino)phenyl)acrylate (270 mg, 93.7% yield). LC-MS m/z: 263 $[\text{M}+\text{H}]^+$.


4). Synthesis of methyl (S,E)-3-(2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazol-6-yl)acrylate (intermediate A-1)



[0247] To a solution of methyl (S, E)-3-(4-amino-3-((oxetan-2-ylmethyl)amino)phenyl)acrylate (intermediate 3) in dry THF (5 mL) chloroacetic anhydride (173 mg, 1.01 mmol) was slowly added at 0° C. The reaction mixture was stirred at 0° C. for 30 mins, after which it was heated to 60° C. and stirred for 3 hours. Cooled to room temperature, quenched with water (20 ml) and extracted with EA (2×10 ml). The combined organic layer was directly concentrated in vacuo to give a residue. The residue was purified by silica gel column chromatography (DCM/MeOH=50/1) to obtain methyl (S, E)-3-(2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazol-6-yl)acrylate (intermediate A-1) (240 mg, 81.9% yield). LC-MS m/z: 321 $[\text{M}+\text{H}]^+$.


Example 1-2. Synthesis of methyl 4-(2-chloroacetamide)-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)-benzoate (Intermediate A-2)

1). Synthesis of methyl 3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)-4-nitrobenzoate




[0248] To a mixture of (1-ethyl-1H-imidazol-5-yl)methanamine (240 mg, 1.21 mmol) and methyl 3-fluoro-4-nitrobenzoate (241.2 mg, 1.21 mmol) in THF (6 mL) and MeOH (4 mL) TEA (1.2 g, 12.1 mmol) was added. The reaction mixture was warmed to 60° C. and stirred over weekend. The reaction mixture was poured into saturated brine (100 mL) and extracted with EA (2×80 mL). The organic layer was concentrated in vacuo to give a residue. The residue was purified by flash column chromatography on silica gel (eluted with EA/PE=3/1) to obtain methyl 3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)-4-nitrobenzoate (320 mg, 87.0% yield). LC-MS m/z: 305 $[\text{M}+\text{H}]^+$.

2). Synthesis of methyl 4-amino-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate




-continued



[0249] To a solution of methyl 3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)-4-nitrobenzoate (320 mg, 1.05 mmol) in MeOH (10 mL) wet Pd/C (50 mg) was added under N₂ atmosphere. The reaction solution was degassed with H₂ three times, and then raised to 45° C. for 3 hours under H₂ atmosphere. The mixture was filtered, and the filtrate was concentrated in vacuo to give a residue. The residue was purified by flash column chromatography on silica gel (eluted with DCM/MeOH=10/1) to obtain methyl 4-amino-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)methyl)benzoate (270 mg, 93.8% yield). LC-MS m/z: 275 [M+H]⁺.


3). Synthesis of methyl 4-(2-chloroacetamide)-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (Intermediate A-2)



[0250] The solution of methyl 4-amino-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (230.0 mg, 0.84 mmol) and 2-chloroacetic anhydride (285.4 mg, 1.68 mmol) in THF (10 mL) was stirred at room temperature for 16 hours. The reaction solution was poured into saturated brine (100 mL) and extracted with DCM (2×80 mL). The combined organic layers were concentrated in vacuo to obtain methyl 4-(2-chloroacetamide)-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (230.0 mg, 78.2% yield), LC-MS m/z: 351 [M+H]⁺.


Example 1-3. Synthesis of methyl (S)-2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (Intermediate A-3)

1). Synthesis of methyl (S)-4-nitro-3-((oxetan-2-ylmethyl)amino)benzoate



[0251] A solution of methyl 3-fluoro-4-nitrobenzoate (4.0 g, 20 mmol) and (S)-oxetan-2-ylmethanamine (1.7 g, 20 mmol) in DMF (40 mL) was added K₂CO₃ (5.5 g, 40 mmol). The reaction was stirred at room temperature for 12 hours. The reaction solution was poured into saturated brine (200 mL) and extracted with ethyl acetate (2×80 mL). The combined organic layers were washed with saturated brine (2×200 mL), dried over anhydrous sodium sulfate, and concentrated in vacuo to obtain crude product methyl (S)-4-nitro-3-((oxetan-2-ylmethyl)amino)benzoate (5.2 g). LC-MS m/z: 267 [M+H]⁺.


2). Synthesis of methyl (S)-4-amino-3-((oxetan-2-ylmethyl)amino)benzoate



[0252] To a solution of methyl (S)-4-nitro-3-((oxetan-2-ylmethyl)amino)benzoate (5.2 g, 20 mmol) in methanol (50 mL) wet Pd/C (520 mg) was added under N₂ atmosphere. The mixture was degassed with H₂ three times, the reaction mixture was stirred at room temperature for 12 hours under H₂ atmosphere. The catalyst was removed by filtration, and the filtrate was concentrated in vacuo to obtain methyl

(S)-4-amino-3-((oxetan-2-ylmethyl)amino)benzoate (3.9 g, 82.3% yield), LC-MS m/z : 237 $[M+H]^+$.


3). Synthesis of methyl (S)-2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (Intermediate A-3)



[0253] To a solution of methyl (S)-4-amino-3-((oxetan-2-ylmethyl)amino)benzoate (1.5 g, 6 mmol) in THF (12 mL) at 0° C. 2-chloroacetic anhydride (1.1 g, 6.6 mmol) was added. The reaction mixture was stirred at room temperature for 12 hours. The mixture was concentrated in vacuo and purified by silica gel column chromatography (EA/PE=1/1) to obtain methyl (S)-2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (Intermediate A-3) (1.6 g, 90% yield). LC-MS m/z : 295 $[M+H]^+$.

Example 1-4. Synthesis of methyl(E)-3-(4-(2-chloroacetamide)-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)-phenyl)acrylate (Intermediate A-4)


1). Synthesis of methyl (E)-3-(3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)-4-nitrophenyl)acrylate



[0254] TEA (1.80 g, 17.80 mmol, 10.0 eq) was added to a mixture of methyl (E)-3-(3-fluoro-4-nitrophenyl)acrylate (Intermediate 1) (400 mg, 1.78 mmol) and (1-Ethyl-1H-imidazol-5-yl)methanamine (354 mg, 1.78 mmol) in THF (6


mL) and MeOH (4 mL). After stirring uniformly, the reaction mixture was heated to 60° C. and stirred for 48 hours. The resulting mixture was poured into brine (50 mL) and extracted with EA (2×20 mL). The combined organic layer was concentrated in vacuo and then purified by silica gel column chromatography (DCM/MeOH=10/1) to obtain methyl (E)-3-(3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)-4-nitrophenyl)acrylate (400 mg, 68.1% yield). LC-MS m/z : 331 $[M+H]^+$.


2) Synthesis of methyl (E)-3-(3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)-4-aminophenyl)acrylate



[0255] The mixture of methyl (E)-3-(3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)-4-nitrophenyl)acrylate (400 mg, 1.21 mmol), iron powder (203 mg, 3.63 mmol) and NH_4Cl (192 mg, 3.62 mmol) in EtOH/ H_2O (v/v=10/1, 5 mL) was raised to 80° C. and stirred for 5 hours. The reaction was cooled to room temperature, poured into saturated NaHCO_3 solution (30 mL) and extracted with EA (3×10 mL). The combined organic layers were washed with brine (3×10 mL), dried over Na_2SO_4 , filtered, and concentrated in vacuo to give a residue. The residue was purified by silica gel column chromatography (DCM/MeOH=20/1) to obtain methyl (E)-3-(3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)-4-aminophenyl)acrylate (60 mg, 16.5% yield). LC-MS m/z : 301 $[M+H]^+$.

3). Synthesis of methyl (E)-3-(4-(2-chloroacetamide)-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)phenyl)-acrylate (Intermediate A-4)






[0256] The mixture of methyl (E)-3-(3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)-4-aminophenyl)acrylate (50 mg, 0.17 mmol) and 2-chloroacetic anhydride (58 mg, 0.34 mmol) in THF (2 mL) was stirred at room temperature for 16 hours. The resulting mixture was poured into brine (10 mL) and extracted with DCM (2×5 mL). The combined organic layer was directly concentrated to give methyl (E)-3-(4-(2-chloroacetamide)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)phenyl)acrylate (Intermediate A-4) (50 mg, 79.8% yield). LC-MS m/z : 377 $[M+H]^+$.

Example 1-5. Synthesis of methyl (S)-2-(2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazol-6-yl)acetate (intermediate A-5)


1). Synthesis of methyl 2-(3-fluoro-4-nitrophenyl)acetate




[0257] To a solution of 2-(3-fluoro-4-nitrophenyl)acetic acid (800 mg, 4.02 mmol) in methanol (10 mL) at 0° C. thionyl chloride (957 mg, 8.04 mmol) was slowly added. The mixture was stirred at room temperature for 3 hours, then the reaction mixture was quenched with water (50 mL) at 0° C. and then extracted with ethyl acetate (2×20 mL). The combined organic layers were dried over anhydrous sodium sulfate and filtered to give a filtrate. The filtrate was concentrated in vacuo to obtain methyl 2-(3-fluoro-4-nitrophenyl)acetate (840 mg, 98.1% yield).

[0258] ^1H NMR (400 MHz, DMSO- d_6): δ 8.13 (t, $J=8.2$ Hz, 1H), 7.55 (dd, $J=12.4$, 1.6 Hz, 1H), 7.37 (dd, $J=8.4$, 1.0 Hz, 1H), 3.90 (s, 2H), 3.65 (s, 3H).

2). Synthesis of methyl (S)-2-(4-nitro-3-((oxetan-2-ylmethyl)amino)phenyl)acetate




-continued




[0259] The mixture of methyl 2-(3-fluoro-4-nitrophenyl)acetate (1.05 g, 4.93 mmol), (S)-oxetan-2-ylmethanamine (468 mg, 5.38 mmol) and potassium carbonate (1.35 g, 9.78 mmol) in N, N-dimethylformamide (10 mL) was stirred at room temperature for 3 hours. The resulting mixture was diluted with water (100 mL) and extracted with dichloromethane (3×10 mL). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo to give a residue. The residue was purified by silica gel column chromatography (PE/EA=2/1) to obtain methyl (S)-2-(4-nitro-3-((oxetan-2-ylmethyl)amino)phenyl)acetate (300 mg, 21.9% yield). LC-MS m/z : 280 $[M+H]^+$.

3). Synthesis of methyl (S)-2-(4-amino-3-((oxetan-2-ylmethyl)amino)phenyl)acetate



[0260] To a solution of methyl (S)-2-(4-nitro-3-((oxetan-2-ylmethyl)amino)phenyl)acetate (300 mg, 1.07 mmol) in methanol (10 mL) wet Pd/C (116 mg) was added under N_2 atmosphere solution. The mixture was degassed with H_2 three times and stirred at room temperature under H_2 atmosphere (1 atm) for 16 hours. The reaction solution was filtered, and the filter cake was washed with methanol (50 mL). The filtrate was concentrated in vacuo and then purified by silica gel column chromatography (PE/EA=1/1) to obtain methyl (S)-2-(4-amino-3-((oxetan-2-ylmethyl)amino)phenyl)acetate (120 mg, 45.0% yield). LC-MS m/z : 251 $[M+H]^+$.


4). Synthesis of Methyl (S)-2-(2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazol-6-yl)acetate (Intermediate A-5)



[0261] To a solution of methyl (S)-2-(4-amino-3-((oxetan-2-ylmethyl)amino)phenyl)acetate (120 mg, 0.48 mmol) in anhydrous THF (5 mL) 2-chloroacetic anhydride (91 mg, 0.53 mmol) was slowly added at 0° C. The mixture was stirred at 0° C. for 30 min, then heated to 60° C. for 3 h. Cooled to room temperature and quenched with water (20 mL), then extracted with ethyl acetate (2×10 mL). The combined organic layers were dried over Na₂SO₄, filtered, and concentrated in vacuo to give a residue. The residue was purified by silica gel column chromatography (DCM/MeOH=50/1) to obtain methyl (S)-2-(2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazol-6-yl)acetate (Intermediate A-5) (100 mg, 67.6% yield). LC-MS m/z: 309 [M+H]⁺.

Example 1-6. Synthesis of methyl 2-(chloromethyl)-1-(1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]-imidazole-6-carboxylate (Intermediate A-6)


1). Synthesis of (1-(cyanomethyl)cyclopropyl)methanesulfonate



[0262] To a mixture of 2-(1-(hydroxymethyl)cyclopropyl)acetonitrile (2.00 g, 18.00 mmol) and triethylamine (4.00 g, 39.25 mmol) in DCM (20 mL) MsCl (3.12 g, 27.23 mmol) was added dropwise at 0° C. Then the mixture was stirred at 0° C. for 1 hour and then stirred at room temperature for 2 hours. The mixture was diluted with DCM (50 mL) and washed with brine (25 mL). The organic layer were dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo to give (1-(cyanomethyl)cyclopropyl)methanesulfonate (3.26 g, 17.23 mmol).

[0263] ¹H NMR (400 MHz, CDCl₃) δ 4.15 (s, 2H), 3.08 (s, 3H), 2.59 (s, 2H), 0.82 (d, J=4.0 Hz, 4H).


2). Synthesis of 2-(1-(azidomethyl)cyclopropyl)acetonitrile



[0264] A solution of (1-(cyanomethyl)cyclopropyl)methanesulfonate (3.32 g, 17.54 mmol) and sodium azide (4.81 g, 73.99 mmol) in N, N-dimethylformamide (20 mL) was heated to 120° C. for 16 hours. The mixture was cooled to room temperature and quenched by the addition of water, extracted with dichloromethane (30 mL×3). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo to obtain 2-(1-(azidomethyl)cyclopropyl)acetonitrile (1.60 g, 11.75 mmol, 66.9% yield).

[0265] ¹H NMR (400 MHz, CDCl₃) δ 3.31 (s, 2H), 2.53 (s, 2H), 0.70 (d, J=1.2 Hz, 4H).


3). Synthesis of 2-(1-(aminomethyl)cyclopropyl)acetonitrile



[0266] To a mixture of 2-(1-(azidomethyl)cyclopropyl)acetonitrile (1.60 g, 11.75 mmol) in methanol (15 mL)/water (9 mL) tributyl phosphine (7.13 g, 35.24 mmol) was added dropwise at room temperature. The resulting mixture was heated to 65° C. and stirred for 3 hours. The mixture was filtered, and the filter cake was washed with methanol (10 mL). The filtrate was concentrated in vacuo to obtain 2-(1-(aminomethyl)cyclopropyl)acetonitrile (821 mg, 7.45 mmol, 63.5% yield).


[0267] ¹H NMR (400 MHz, DMSO-*d*₆) δ 2.39 (s, 2H), 2.25 (s, 2H), 0.22 (t, J=5.2 Hz, 2H), 0.17 (t, J=5.2 Hz, 2H).

4). Synthesis of methyl 3-((1-(1-(cyanomethyl)cyclopropyl)methyl)amino)-4-nitrobenzoate




[0268] To a mixture of 3-fluoro-4-nitrobenzoate (300 mg, 1.51 mmol) and 2-(1-(aminomethyl)cyclopropyl)acetonitrile (166 mg, 1.51 mmol) in N,N-dimethylformamide (5 mL) potassium carbonate (625 mg, 4.53 mmol) was added. The mixture was stirred at room temperature for 16 hours. The reaction mixture was diluted with water (15 mL) and then extracted with ethyl acetate (3×30 mL). The combined organic layers were washed with brine (2×30 mL), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo to give a residue. The residue was purified by silica gel column chromatography (PE/EA=5/1) to give methyl 3-((1-(cyanomethyl)cyclopropyl)methyl)amino)-4-nitrobenzoate (250 mg, 0.86 mmol, 57.6% yield). LC-MS *m/z*: 290 [M+H]⁺.

5). Synthesis of methyl 4-amino-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)benzoate



[0269] To a mixture of methyl 3-((1-(cyanomethyl)cyclopropyl)methyl)amino)-4-nitrobenzoate (200 mg, 0.69 mmol) dissolved in methanol (8 mL) zinc powder (449 mg, 6.87 mmol) and acetic acid (415 mg, 6.91 mmol) were added at room temperature. The mixture was stirred at room temperature for 1 hour. The mixture was diluted water (15 mL) and extracted with ethyl acetate (3×30 mL). The combined organic layers were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated to give a residue in vacuo. The residue was purified by reverse phase column chromatography with the following conditions: Column: Spherical C18, 20-40 μm, 120 g; Mobile Phase A: water (10 mM FA); Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 40% B-60% B within 20 minutes; detector: 254 nm. The fractions containing desired product were collected at 52% B and concentrated in vacuo to afford compound methyl 4-amino-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)benzoate (141 mg, 0.54 mmol, 78.6% yield). LC-MS *m/z*: 260 [M+H]⁺.


6). Synthesis of methyl 2-(chloromethyl)-1-(1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylate (Intermediate A-6)



[0270] To a solution of methyl 4-amino-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)benzoate (140 mg, 0.54 mmol) in dry THF (5 mL) 2-chloroacetic anhydride (92 mg, 0.54 mmol) was added batchwise at room temperature. The mixture was stirred at 60° C. for 16 hours. The reaction mixture was quenched by water (5 mL) and extracted with ethyl acetate (3×20 mL). The combined organic layers were combined and washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated to give a residue in vacuo. The residue was purified by reverse phase flash chromatography with the following conditions: column: Spherical C18, 20-40 μm, 120 g; mobile phase A: water (10 mM FA); mobile phase B: acetonitrile; flow rate: 80 mL/min; gradient: 20 minutes 40%-60% B; detector: 254 nm). The mobile phase containing the desired product was collected under 50% B mobile phase and then concentrated in vacuo to obtain methyl 2-(chloromethyl)-1-(1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylate (Intermediate A-6) (90 mg, 0.54 mmol, 52.5% yield). LC-MS *m/z*: 318 [M+H]⁺.


Example 1-7. Synthesis of methyl 4-(2-chloroacetamide)-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)-5-fluorobenzoate (Intermediate A-7)

1). Synthesis of 2-(1-((5-Bromo-3-fluoro-2-nitrophenyl)amino)methyl)cyclopropyl)acetonitrile





[0271] To a mixture of 5-bromo-1,3-difluoro-2-nitrobenzene (1.00 g, 4.20 mmol) and 2-(1-(aminomethyl)-cyclopropyl)acetonitrile (464.2 mg, 4.21 mmol) in dimethyl sulfoxide (20 mL) *N,N*-diisopropylethylamine (1.63 g, 12.61 mmol) was added. The mixture was stirred at 70° C. for 4 hours. The reaction mixture was diluted with water (15 mL) and extracted with ethyl acetate (3×30 mL). The combined organic layers were washed with brine (2×30 mL), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (PE/EA=5/1) to obtain 2-(1-((5-bromo-3-fluoro-2-nitrophenyl)amino)methyl)-cyclopropyl)acetonitrile (542.0 mg, 1.65 mmol, 39.3% yield). LC-MS *m/z*: 328, 330 [M+H]⁺.

2). Synthesis of 2-(1-((2-amino-5-bromo-3-fluorophenyl)amino)methyl)cyclopropyl)acetonitrile




[0272] To a solution of 2-(1-((5-bromo-3-fluoro-2-nitrophenyl)amino)methyl)cyclopropyl)acetonitrile (542.0 mg, 1.65 mmol) in methanol (10 mL) was added zinc powder (1.08 g, 16.52 mmol) and acetic acid (996.0 mg, 16.59 mmol). The mixture was stirred at room temperature for 30 minutes. The reaction mixture was diluted with water (10 mL) and extracted with ethyl acetate (3×20 mL). The combined organic layers were washed with brine (2×20 mL), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (PE/EA=2/1) to obtain 2-(1-((2-amino-5-bromo-3-fluorophenyl)amino)methyl)-cyclopropyl)acetonitrile (358.0 mg, 1.21 mmol, 72.9% yield). LC-MS *m/z*: 298 [M+H]⁺.

3). Synthesis of methyl 4-amino-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)-5-fluorobenzoate



[0273] To a mixture of 2-(1-((2-amino-5-bromo-3-fluorophenyl)amino)methyl)cyclopropyl)acetonitrile (358.0 mg, 1.21 mmol) in *N,N*-dimethylformamide (3 mL)/methanol (10 mL) triethylamine (366.6 mg, 3.62 mmol) and 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (88.5 mg, 0.12 mmol) were added at room temperature. The mixture was warmed to 90° C. for 16 hours under carbon monoxide atmosphere. Cooled to room temperature, the reaction mixture was diluted with water (15 mL) and extracted with ethyl acetate (30 mL×3). The combined organic layers were washed with brine (50 mL), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (PE/EA=1/1) to obtain methyl 4-amino-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)-5-fluorobenzoate (100.0 mg, 0.36 mmol, 29.8% yield), LC-MS *m/z*: 278 [M+H]⁺.

4). Synthesis of methyl 4-(2-chloroacetamide)-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)-5-fluorobenzoate (Intermediate A-7)




A solution of methyl 4-amino-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)-5-fluorobenzoate (85.0 mg, 0.31 mmol) in THF (4 mL) 2-chloroacetic anhydride (52.5 mg, 0.31 mmol) was added dropwise at room temperature. The mixture was stirred at 60° C. for 16 hours. The mixture was quenched by adding water (5 mL) and extracted with ethyl acetate (3×20 mL). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by reverse phase flash chromatography under the following conditions (column: Spherical C18, 20-40 μm, 120 g; mobile phase A: water (10 mM FA); mobile phase B: acetonitrile; flow rate: 80 mL/min; Gradient: 40% B-60% B in 20 minutes; detector: 254 nm). The mobile phase containing the desired product was collected under 52% B mobile phase and concentrated in vacuo to

obtain methyl 4-(2-chloroacetamide)-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)-5-fluorobenzoate (Intermediate A-7) (141 mg, 0.54 mmol, 78.6% yield). LC-MS m/z : 354 $[M+H]^+$.


Example 1-8. Synthesis of methyl 2-(chloromethyl)-1-(oxetan-3-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (Intermediate A-8)

1). Synthesis of methyl 4-nitro-3-((oxetan-3-ylmethyl)amino)benzoate




[0274] The mixture of methyl 3-fluoro-4-nitrobenzoate (1.60 g, 8.05 mmol), oxetan-3-ylmethanamine (700 mg, 8.05 mmol) and potassium carbonate (2.22 g, 16.10 mmol) in N, N-dimethylformamide (20 mL) was stirred at room temperature for 16 hours. The resulting mixture was diluted with water (200 mL) and extracted with ethyl acetate (50 mL $\times$ 3). The combined organic layers were washed with brine (50 mL $\times$ 3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=1/2) to obtain methyl 4-nitro-3-((oxetan-3-ylmethyl)amino)benzoate (1.50 g, 70.1% yield). LC-MS m/z : 267 $[M+H]^+$.

2). Synthesis of methyl 4-amino-3-((oxetan-3-ylmethyl)amino)benzoate



[0275] To a solution of methyl 4-nitro-3-((oxetan-3-ylmethyl)amino)benzoate (1.50 g, 5.64 mmol) in methanol (30 mL) palladium on carbon (200 mg) was added under N_2 atmosphere. The resulting mixture was degassed H_2 three times and stirred at room temperature for 4 hours under H_2 atmosphere. The reaction was filtered, and the filter cake was washed with methanol (20 mL). The filtrate was concentrated to obtain methyl 4-amino-3-((oxetan-3-ylmethyl)amino)benzoate (1.20 g, 90.2% yield), which was used in the next reaction without purification. LC-MS m/z : 237 $[M+H]^+$.


3). Synthesis of methyl 2-(chloromethyl)-1-(oxetan-3-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate




[0276] To a solution of methyl 4-amino-3-((oxetan-3-ylmethyl)amino)benzoate (1.20 g, 5.08 mmol) in dry THF (20 mL) 2-chloroacetic anhydride (869 mg, 5.08 mmol) was added slowly at 0° C. After stirred at 0° C. for 30 minutes, the mixture was heated to 70° C. and stirred for 3 hours. The mixture was quenched with water (100 mL) at room temperature and extracted with ethyl acetate (2 $\times$ 0 mL). The organic layer was concentrated in vacuo and purified by silica gel column chromatography (petroleum ether/ethyl acetate=1/1) to obtain methyl 2-(chloromethyl)-1-(oxetan-3-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (1.00 g, 67.0% yield), LC-MS m/z : 295 $[M+H]^+$.

Example 1-9. Synthesis of methyl 2-(chloromethyl)-1-((3-methyloxetan-3-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (Intermediate A-9)

1). Synthesis of methyl 3-((3-methoxy-3-yl)methyl)amino)-4-nitrobenzoate





-continued





[0277] The mixture of methyl 3-fluoro-4-nitrobenzoate (2.00 g, 10.00 mmol), (3-methyloxan-3-yl)carboxamide (1.00 g, 10.00 mmol) and potassium carbonate (4.14 g, 30.00 mmol) in N,N-dimethylformamide (20 mL) was stirred at room temperature for 16 hours. The resulting mixture was diluted with water (200 mL) and extracted with ethyl acetate (50 mL $\times$ 3). The combined organic layers were washed with brine (50 mL $\times$ 3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=2/1) to obtain methyl 3-((3-methoxy-3-yl)methyl)amino-4-nitrobenzoate (1.96 g, 70.4% yield). LC-MS m/z : 281 [M+H]⁺.

2). Synthesis of methyl 4-amino-3-((3-methyloxetan-3-yl)methyl)amino)benzoate



[0278] A solution of methyl 3-((3-methoxy-3-yl)methyl)amino-4-nitrobenzoate (1.96 g, 7.04 mmol) in methanol (20 mL) was added palladium on carbon (400 mg). The resulting mixture was degassed and flushed with hydrogen three times, and then stirred at room temperature for 4 hours. The reaction was filtered, and the filter cake was washed with methanol (20 mL). The filtrate was concentrated to give methyl 4-amino-3-((3-methyloxetan-3-yl)methyl)amino)benzoate (1.55 g, 85.2% yield), which was directly used for the next step without further purification, LC-MS m/z : 251 [M+H]⁺.


3). Synthesis of methyl 2-(chloromethyl)-1-((3-methyloxetan-3-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (Intermediate A-10)



[0279] To a solution of methyl 4-amino-3-((3-methyloxetan-3-yl)methyl)amino)benzoate (200 mg, 0.80 mmol) in dry THF (5 mL) 2-chloroacetic anhydride (137 mg, 0.80 mmol) was added dropwise at 0° C. The mixture was stirred at room temperature for 1 hour. The solvent was removed by concentration in vacuo to give a residue. The residue was added dioxane (5 mL) then heated to 100° C. for 3 hours. The mixture was quenched with water (20 mL) at room temperature and extracted with ethyl acetate (2 $\times$ 10 mL). The combined organic layer was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=1/1) to obtain methyl 2-(chloromethyl)-1-((3-methyloxetan-3-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (Intermediate A-10) (200 mg, 81.2% yield). LC-MS m/z : 309 [M+H]⁺.

Example 1-10. Synthesis of methyl 4-(2-chloroacetamide)-3-((1,2-dimethyl-1H-imidazol-5-yl)methyl)amino)-benzoate (Intermediate A-10)


1). Synthesis of 1,2-dimethyl-1H-imidazole-5-carbaldehyde



[0280] To a solution of 5-bromo-1,2-dimethyl-1H-imidazole (500 mg, 2.86 mmol) in dry tetrahydrofuran (10 mL) *n*-butyllithium (1.26 mL, 3.15 mmol, 2.5M in hexane) was added at -78° C. The mixture was stirred at -78° C. for 30 min, then N, N-dimethylformamide (626 mg, 8.58 mmol)


was slowly added to the mixture at 0° C. The mixture was stirred at room temperature for 2 hours. When completion, the reaction was quenched with water (50 mL) at 0° C. and extracted with ethyl acetate (2×30 mL). The combined organic layer was concentrated in vacuo and purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain 1,2-dimethyl-1H-imidazole-5-carbaldehyde (300 mg, 84.6 yield %), LC-MS *m/z*: 125 [M+H]⁺.

2). Synthesis of (E)-N-((1,2-dimethyl-1H-imidazol-5-yl)methylene)-2-methylpropane-2-sulfinamide




[0281] The mixture of 1,2-dimethyl-1H-imidazole-5-carbaldehyde (300 mg, 2.42 mmol), 2-methylpropane-2-sulfoxide (439 mg, 3.63 mmol) and tetraisopropyl titanate (2.06 g, 7.26 mmol) in THF (5 mL) was stirred at room temperature for 16 hours. The resulting mixture was diluted with water (2 mL) and filtered. The filtrate was concentrated in vacuo and purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain (E)-N-((1,2-dimethyl-1H-imidazol-5-yl)methylene)-2-methylpropane-2-sulfoxide amide (500 mg, 91.0% yield). LC-MS *m/z*: 228 [M+H]⁺.

3). Synthesis of N-((1,2-Dimethyl-1H-imidazol-5-yl)methyl)-2-methylpropane-2-sulfoxide amide



[0282] The mixture of (E)-N-((1,2-dimethyl-1H-imidazol-5-yl)methylene)-2-methylpropane-2-sulfoxide amide (500 mg, 2.20 mmol) and boron sodium hydride (167 mg, 4.40 mmol) in methanol (10 mL) was stirred at room temperature for 2 hours. When completion, the reaction was quenched with water (50 mL) at room temperature and extracted with dichloromethane (2×20 mL). The combined organic layer was concentrated in vacuo to obtain N-((1,2-dimethyl-1H-imidazol-5-yl)methyl)-2-methylpropane-2-sulfoxide amide (400 mg; 55.3% yield), which was directly used in the next step without further purification. LC-MS *m/z*: 230 [M+H]⁺.


4). Synthesis of 1,2-Dimethyl-1H-imidazol-5-yl)methanamine hydrochloride



[0283] The mixture of N-((1,2-Dimethyl-1H-imidazol-5-yl)methyl)-2-methylpropane-2-sulfoxide amide (400 mg, 1.75 mmol) in HCl (5 mL, 3M in MeOH) was stirred at room temperature for 1 hour. The reaction was filtered to give (1,2-dimethyl-1H-imidazol-5-yl)methanamine hydrochloride (200 mg, 57.8% yield).


[0284] ¹H NMR (400 MHz, DMSO-d₆): δ 8.86 (s, 2H), 7.63 (s, 1H), 4.19 (s, 2H), 3.77 (s, 3H), 2.63 (s, 3H).

5). Synthesis of methyl 3-((1,2-dimethyl-1H-imidazol-5-yl)methyl)amino)-4-nitrobenzoate




[0285] To a mixture of (1,2-dimethyl-1H-imidazol-5-yl)methanamine hydrochloride (200 mg, 1.01 mmol) and methyl 3-fluoro-4-nitrobenzoate (201 mg, 1.01 mmol) in THF (3 mL) and methanol (2 mL) triethylamine (510 mg, 5.05 mmol) was added. The mixture was stirred at 60° C. for 3 h. The resulting mixture was poured into brine (50 mL) and extracted with ethyl acetate (2×20 mL). The combined organic layer was concentrated in vacuo and purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain 3-((1,2-dimethyl-1H-imidazol-5-yl)methyl)amino)-4-Nitrobenzoic acid methyl ester (200 mg, 65.1% yield). LC-MS *m/z*: 305 [M+H]⁺.

6). Synthesis of methyl 4-amino-3-((1,2-dimethyl-1H-imidazol-5-yl)methyl)amino)benzoate



[0286] To a solution of methyl 3-((1,2-dimethyl-1H-imidazol-5-yl)methyl)amino)-4-nitrobenzoate (200 mg, 0.66 mmol) in methanol (10 mL) palladium on carbon (20 mg) was added. The resulting mixture was degassed and flushed with hydrogen three times and stirred at room temperature for 3 hours. The reaction mixture was filtered, and washed with methanol (50 mL). The filtrate was concentrated in vacuo to obtain methyl 4-amino-3-((1,2-dimethyl-1H-imidazol-5-yl)methyl)amino)benzoate (150 mg, 82.9% yield) that was directly used in the next step without purification. LC-MS *m/z*: 275 [M+H]⁺.

7). Synthesis of methyl 4-(2-chloroacetamide)-3-((1,2-dimethyl-1H-imidazol-5-yl)methyl)amino)benzoate




[0287] A solution of methyl 4-amino-3-((1,2-dimethyl-1H-imidazol-5-yl)methyl)amino)benzoate (150 mg, 0.55

mmol) and 2-chloroacetic anhydride (142 mg, 0.83 mmol) in THF (3 mL) was stirred at room temperature for 2 hours. The resulting mixture was concentrated in vacuo to give a residue. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 4-(2-chloroacetamide)-3-((1,2-dimethyl-1H-imidazol-5-yl)methyl)amino)benzoate (150 mg, 77.9% yield), LC-MS *m/z*: 351 [M+H]⁺.


Example 1-11. Synthesis of methyl 5-(2-chloroacetamide)-6-(1-(cyanomethyl)cyclopropyl)methyl)amino-picolinate

1). Synthesis of methyl 6-((1-(cyanomethyl)cyclopropyl)methyl)amino)-5-nitropicolinate



[0288] To a solution of methyl 6-chloro-5-nitropicolinate (850 mg, 3.94 mmol) and 2-(1-(aminomethyl) cyclopropyl) acetonitrile (433 mg, 3.94 mmol) in DMSO (10 mL) *N,N*-diisopropylethylamine (1.52 g, 11.81 mmol) was added. The reaction was stirred at 60° C. for 16 hours. The resulting mixture was poured into brine (20 mL) and extracted with ethyl acetate (20 mL×2). The combined organic layer was concentrated and purified by silica gel column chromatography (petroleum ether/ethyl acetate=1/1) to obtain methyl 6-((1-(cyanomethyl)cyclopropyl)methyl)amino)-5-nitropicolinate (190 mg, 16.7% yield). LC-MS *m/z*: 291 [M+H]⁺.


2). Synthesis of methyl 5-amino-6-((1-(cyanomethyl)cyclopropyl)methyl)amino)picolinate



[0289] To a solution of methyl 6-((1-(cyanomethyl)cyclopropyl)methyl)amino)-5-nitropicolinate (190 mg, 0.65 mmol) in methanol (5 mL) zinc powder (428 mg, 6.55 mmol) and acetic acid (393 mg, 6.55 mmol) were added at room temperature and the resulting solution was stirred at


room temperature for 2 hours. After completion, the resulting mixture was filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=2/1) to obtain methyl 5-amino-6-((1-(cyanomethyl)cyclopropyl)methyl)amino)picolinate (130 mg, 53.7% yield). LC-MS m/z : 261 $[M+H]^+$.

3). Synthesis of methyl 5-(2-chloroacetamide)-6-(1-(cyanomethyl)cyclopropyl)methyl)aminopicolinate (Intermediate A-11)



[0290] To a solution of methyl 5-amino-6-((1-(cyanomethyl)cyclopropyl)methyl)amino)picolinate (80 mg, 0.31 mmol) in tetrahydrofuran (5 mL) 2-chloroacetic anhydride (79 mg, 0.46 mmol) was added. The resulting mixture was stirred at room temperature for 2 hours. After completion, the mixture was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=1/1) to obtain methyl 5-(2-chloroacetamido)-6-((1-(cyanomethyl)cyclopropyl)methyl)amino)picolinate (Intermediate A-11) (150 mg, 88% yield). LC-MS m/z : 337 $[M+H]^+$.


Example 1-12. Synthesis of methyl (S)-2-(chloromethyl)-3-(oxetan-2-ylmethyl)-3H-imidazo[4,5-b]pyridine-5-carboxylate (Intermediate A-12)



[0291] To a solution of methyl (S)-5-amino-6-((oxetan-2-ylmethyl)amino)picolinate (100 mg, 0.42 mmol) in tetrahydrofuran (10 mL) 2-chloroacetic anhydride (101 mg, 0.63


mmol) was added. The reaction mixture was stirred at 60° C. for 3 hours. The resulting mixture was directly concentrated in vacuo to give methyl (S)-2-(chloromethyl)-3-(oxetan-2-ylmethyl)-3H-imidazo[4,5-b]pyridine-5-carboxylate (Intermediate A-12) (140 mg).

Example 1-13. Synthesis of methyl 5-(2-chloroacetamide)-6-(1-ethyl-1H-imidazol-5-yl)methyl)aminopicolinate (Intermediate A-13)



[0292] To a solution of methyl 5-amino-6-(1-ethyl-1H-imidazol-5-yl)methyl)aminopicolinate (150 mg, 0.54 mmol) in THF (5 mL) 2-chloroacetic anhydride (138 mg, 0.82 mmol) was added dropwise. The mixture was stirred at 25° C. for 16 hours. Then the mixture was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 5-(2-chloroacetamide)-6-(1-ethyl-1H-imidazol-5-yl)methyl)aminopicolinate (180 mg, 94.7% yield). LC-MS m/z : 352 $[M+H]^+$.


Example 1-14. Synthesis of methyl 4-(2-chloroacetamide)-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)-benzoate (Intermediate A-14)



[0293] To a solution of methyl 4-amino-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)benzoate (500 mg, 1.93 mmol) in THF (5 mL) 2-chloroacetic anhydride (494 mg,


2.89 mmol) was added. The mixture was stirred at room temperature for 2 hours. After completion, the mixture was concentrated and purified by silica gel column chromatography (petroleum ether/ethyl acetate=2/1) to obtain methyl 4-(2-chloroacetamide)-3-((1-(cyanomethyl)cyclopropyl)methylamino)-benzoate (Intermediate A-14) (400 mg, 61.8% yield). LC-MS m/z : 336 $[M+H]^+$.

Example 1-15. Synthesis of (S)-N-(4-Bromo-2-fluoro-6-((oxetan-2-ylmethyl)amino)phenyl)-2-chloroacetamide (Intermediate A-15)




[0294] To a solution of (S)-5-bromo-3-fluoro-N¹-(oxetan-2-ylmethyl)benzene-1,2-diamine (300 mg, 1.09 mmol) in dry THF (5 mL) 2-chloroacetic anhydride (241 mg, 1.42 mmol) was added. The mixture was stirred at room temperature for 1 hour. The mixture was diluted with water (10 mL) then extracted with ethyl acetate (10 mL×3). The combined organic layer was concentrated in vacuo and purified by silica gel column chromatography (petroleum ether/ethyl acetate=1/1) to obtain (S)-N-(4-bromo-2-fluoro-6-((oxetan-2-ylmethyl)amino)phenyl)-2-chloroacetamide (230 mg, 59.9% yield). LC-MS m/z : 353 $[M+H]^+$.

Example 1-16. Synthesis of N-(4-bromo-2-((1-(cyanomethyl)cyclopropyl)methylamino)-6-fluorophenyl)-2-chloroacetamide (Intermediate A-16)




-continued



[0295] To a solution of 2-(1-((2-amino-5-bromo-3-fluorophenyl)amino)methyl)cyclopropyl)acetonitrile (150 mg, 0.50 mmol) in dry THF (5 mL) 2-chloroacetic anhydride (129 mg, 0.76 mmol) was slowly added. The mixture was stirred at room temperature for 1 hour. The mixture was diluted with water (10 mL) then extracted with ethyl acetate (3×10 mL). The combined organic layer was concentrated in vacuo and purified by silica gel column chromatography (petroleum ether/ethyl acetate=3/1) to give N-(4-bromo-2-((1-(cyanomethyl)cyclopropyl)methylamino)-6-fluorophenyl)-2-chloroacetamide (218 mg, crude). LC-MS m/z : 376 $[M+H]^+$.

Example 1-17. Synthesis of methyl 4-(2-chloroacetamide)-3-((1-ethyl-1H-imidazol-5-yl)methylamino)-5-fluorobenzoate (Intermediate A-17)


1). Synthesis of methyl 4-amino-3-((1-ethyl-1H-imidazol-5-yl)methylamino)-5-fluorobenzoate



[0296] Potassium acetate (470 mg, 4.80 mmol) and 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (117 mg, 0.16 mmol) were added to a mixture of 5-bromo-N¹-(1-ethyl-1H-imidazol-5-yl)methyl)-3-fluorobenzene-1,2-diamine (500 mg, 1.60 mmol) in methanol (5 mL) and N,N-dimethylformamide (5 mL). The mixture was degassed with carbon monoxide three times and stirred at 90°C for 16 hours under carbon monoxide atmosphere. After completion, the mixture was diluted with water (30 mL) and


extracted with ethyl acetate (30 mL×2). The combined organic layers were washed with brine (20 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to give methyl 4-amino-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)-5-fluorobenzoate (150 mg, 32.1% yield). LC-MS m/z : 293 $[M+H]^+$.

2). Synthesis of methyl 4-(2-chloroacetamide)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)-5-fluorobenzoate




[0297] To a solution of methyl 4-amino-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)-5-fluorobenzoate (150 mg, 0.51 mmol) in THF (5 mL) 2-chloroacetic anhydride (176 mg, 1.02 mmol) was slowly added at room temperature. The reaction was stirred at room temperature for 3 hours. After completion, the reaction was quenched with water (30 mL) and extracted with ethyl acetate (2×20 mL). The organic phase was concentrated in vacuo and then purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 4-(2-chloroacetamide)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)-5-fluorobenzoate (130 mg, 69.3% yield). LC-MS m/z : 369 $[M+H]^+$.

Example 1-18. Synthesis of methyl 4-(2-chloroacetamido)-3-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)aminobenzoate (Intermediate A-18)




-continued



[0298] To a solution of methyl 4-amino-3-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)aminobenzoate (500 mg, 1.67 mmol) in dry THF (10 mL) chloroacetic anhydride (715 mg, 4.18 mmol) was slowly added at room temperature. The resulting mixture was stirred at room temperature for 2 hours. After completion, the solvent was removed by concentration in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 4-(2-chloroacetamide)-3-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)aminobenzoate (Intermediate A-18) (400 mg, 63.5% yield). LC-MS m/z : 377 $[M+H]^+$.

Example 1-19. Synthesis of methyl 2-(chloromethyl)-1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (Intermediate A-19)




[0299] To a solution of methyl 4-amino-3-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)aminobenzoate (500 mg, 1.67 mmol) in dry THF (10 mL) 2-chloroacetic anhydride (715 mg, 4.18 mmol) was added at room temperature solution. The resulting mixture was stirred at room temperature for 2 hours then raised to 60° C. stirred overnight. After completion, the solvent was removed by concentration in vacuo and the residue was purified by silica gel column

chromatography (dichloromethane/methanol=20/1) to obtain methyl 2-(chloromethyl)-1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 16.7% yield). LC-MS m/z : 359 $[M+H]^+$.


Example 1-20. Synthesis of methyl 2-(chloromethyl)-1-methyl-1H-benzo[d]imidazole-6-carboxylate (Intermediate A-20)

1). Synthesis of methyl 3-(methylamino)-4-nitrobenzoate




[0300] The solution of methyl 3-fluoro-4-nitrobenzoate (2.00 g, 10.00 mmol) and methylamine (10 mL, 2 M in tetrahydrofuran) in THF (30 mL) was stirred at room temperature for 2 hours. After completion, it was concentrated in vacuo and was purified by silica gel column chromatography (petroleum ether/ethyl acetate=1/1) to obtain methyl 3-(methylamino)-4-nitrobenzoate (1.80 g, 85.0% yield). LC-MS m/z : 211 $[M+H]^+$.

2). Synthesis of methyl 4-amino-3-(methylamino)benzoate



[0301] To a solution of methyl 3-(methylamino)-4-nitrobenzoate (1.80 g, 8.57 mmol) in methanol (20 mL) was added palladium on carbon (900 mg) at room temperature under N_2 atmosphere. The resulting mixture was degassed and flushed with hydrogen three times and stirred at room temperature for 3 hours. After completion, the reaction solution was filtered, and the filter cake was washed with methanol (20 mL). The filtrate was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=16/9) to obtain methyl 4-amino-3-(methylamino)benzoate (1.2 g, 77.8% yield). LC-MS m/z : 181 $[M+H]^+$.


3). Synthesis of methyl 2-(chloromethyl)-1-methyl-1H-benzo[d]imidazole-6-carboxylate



[0302] To a solution of methyl 4-amino-3-(methylamino)benzoate (1.2 g, 2.83 mmol) in dry THF (20 mL) was slowly added chloroacetic anhydride (2.27 g, 5.66 mmol) at room temperature. The reaction was stirred at room temperature for 2 hours, then the reaction mixture was heated to 60° C. for 16 hours. After completion, it was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=3/1) to obtain methyl 2-(chloromethyl)-1-methyl-1H-benzo[d]imidazole-6-carboxylate (1.4 g, 88.2% yield). LC-MS m/z : 239 $[M+H]^+$.

Example 1-21. Synthesis of methyl 4-(2-chloroacetamide)-3-((1-isopropyl-1H-imidazol-5-yl)methyl)amino)-benzoate (Intermediate A-21)


1). Synthesis of (E)-N-((1-isopropyl-1H-imidazol-5-yl)methylene)-2-methylpropane-2-sulfoxide amide



[0303] The mixture of (E)-N-((1H-imidazol-5-yl)methylene)-2-methylpropane-2-sulfoxide amide (5.00 g, 25.12 mmol), potassium carbonate (10.40 g, 75.36 mmol) and 2-iodopropane (8.54 g, 50.25 mmol) in N,N-dimethylacetamide (100 mL) was stirred at room temperature for 16 hours. After completion, it was diluted with water (200 mL)


and extracted with ethyl acetate (50 mL×3). The combined organic layers were washed with brine (50 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=12/1) to give (E)-N-((1-isopropyl-1H-imidazol-5-yl)methylene)-2-methylpropane-2-sulfoxide amide (4.8 g, 73.4% yield). LC-MS *m/z*: 242 [M+H]⁺.

2). Synthesis of N-((1-isopropyl-1H-imidazol-5-yl)methyl)-2-methylpropane-2-sulfoxide amide




[0304] To a solution of (E)-N-((1-isopropyl-1H-imidazol-5-yl)methylene)-2-methylpropane-2-sulfoxide amide (4.80 g, 19.92 mmol) in methanol (30 mL) sodium borohydride (2.28 g, 59.75 mmol) was slowly added at room temperature. The resulting mixture was stirred at room temperature for 2 hours. After completion, water (200 mL×2) was added to quench the reaction, and extracted with ethyl acetate (200 mL). The combined organic layers were washed with brine (200 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo to obtain N-((1-isopropyl-1H-imidazol-5-yl)methyl)-2-methylpropane-2-sulfoxide amide (4.5 g, 87.5% yield), LC-MS *m/z*: 244 [M+H]⁺.

3). Synthesis of (1-isopropyl-1H-imidazol-5-yl)methanamine hydrochloride




[0305] The mixture of N-((1-isopropyl-1H-imidazol-5-yl)methyl)-2-methylpropane-2-sulfoxide amide (4.50 g, 18.44 mmol) and HCl (50 mL, 3 M in methanol) was stirred for 16 hours at room temperature. After completion, the reaction mixture was concentrated in vacuo to obtain (1-isopropyl-1H-imidazol-5-yl)methanamine hydrochloride (2.4 g, 93.6% yield). LC-MS *m/z*: 140 [M+H]⁺.

4). Synthesis of methyl 3-((1-isopropyl-1H-imidazol-5-yl)methylamino)-4-nitrobenzoate




[0306] The mixture of (1-isopropyl-1H-imidazol-5-yl)methanamine hydrochloride (3.0 g, 17.27 mmol), methyl 3-fluoro-4-nitrobenzoate (4.14 g, 20.72 mmol) and potassium carbonate (4.79 g, 34.53 mmol) in N,N-dimethylformamide (50 mL) was stirred at room temperature for 16 hours. After completion, it was diluted with water (200 mL) and extracted with ethyl acetate (50 mL×3). The combined organic layers were washed with brine (50 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=12/1) to obtain methyl 3-((1-isopropyl-1H-imidazol-5-yl)methylamino)-4-nitrobenzoate (1.2 g, 21.9% yield). LC-MS *m/z*: 319 [M+H]⁺.

5). Synthesis of methyl 4-amino-3-((1-isopropyl-1H-imidazol-5-yl)methylamino)benzoate




-continued



[0307] To a solution of methyl 3-((1-isopropyl-1H-imidazol-5-yl)methyl)amino-4-nitrobenzoate (1.2 g, 3.78 mmol) in methanol (30 mL) palladium on carbon (600 mg, 5.6 mmol) was added at room temperature. The resulting mixture was degassed and flushed with hydrogen three times, and the reaction was stirred at room temperature for 3 hours under H₂ atmosphere. After completion, filter and wash the filter cake with methanol (50 mL). After the filtrate was concentrated to give a residue in vacuo, the residue was purified by silica gel column chromatography (dichloromethane/methanol=13/1) to obtain 4-amino-3-((1-isopropyl-1H-imidazol-5-yl)methyl)amino) methyl benzoate (750 mg, 69.0% yield). LC-MS m/z: 289 [M+H]⁺.


6). Synthesis of methyl 4-(2-chloroacetamido)-3-((1-isopropyl-1H-imidazol-5-yl)methyl)amino)benzoate



[0308] To a solution of methyl 4-amino-3-((1-isopropyl-1H-imidazol-5-yl)methyl)amino)benzoate (400 mg, 1.39 mmol) in dry THF (8 mL) added chloroacetic anhydride (595 mg, 3.48 mmol) was slowly at room temperature. The reaction was stirred at for 3 hours. After completion, the reaction solution was concentrated in vacuo, The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain 4-(2-chloroacetamido)-3-((1-isopropyl-1H-imidazol-5-yl)methyl)amino) methyl benzoate (450 mg, 88.8% yield). LC-MS m/z: 365 [M+H]⁺.


Example 1-22. Synthesis of methyl 4-(2-chloroacetamido)-3-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)amino)benzoate (Intermediate A-22)

1). Synthesis of (E)-N-((1H-imidazol-5-yl)methylene)-2-methylpropane-2-sulfoxide amide




[0309] To a mixture of 1H-imidazole-5-carbaldehyde (5.00 g, 52.08 mmol) and 2-methylpropane-2-sulfoxide amide (9.39 g, 78.12 mmol) in THF (100 mL) titanium tetraisopropoxide (44.40 g, 156.25 mmol) was added at room temperature. The resulting mixture was stirred at room temperature for 16 hours. After completion, it was diluted with water (100 mL), filtered, and the filter cake was washed with ethyl acetate (150 mL). And the aqueous layer was extracted with ethyl acetate (100 mL×2). The combined organic layers were washed with brine (250 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=12/1) to obtain (E)-N-((1H-imidazol-5-yl)methylene)-2-methylpropane-2-sulfoxide amide (9.00 g, 86.4% yield). LC-MS m/z: 200 [M+H]⁺.

2). Synthesis of (E)-N-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methylene)-2-methylpropane-2-sulfoxide amide




[0310] To a mixture of (E)-N-((1H-imidazol-5-yl)methylene)-2-methylpropane-2-sulfoxide amide (3.00 g, 15.10 mmol) and potassium carbonate (6.25 g, 45.30 mmol) in N,N-dimethylformamide (50 mL) (bromomethyl)cyclopropane (4.08 g, 30.20 mmol) was added. The resulting mixture was stirred at room temperature for 16 hours. After completion, the resulting mixture was diluted with water (200 mL) and extracted with ethyl acetate (150 mL×3). The combined organic layers were washed with brine (150 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=13/1) to give (E)-N-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methylene)-2-methylpropane-2-sulfoxide amide (2.8 g, yield 73.4%). LC-MS m/z: 254 [M+H]⁺.

3). Synthesis of N-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-2-methylpropane-2-sulfoxide amide




[0311] To a solution of (E)-N-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methylene)-2-methylpropane-2-sulfoxide amide (2.80 g, 11.60 mmol) in methanol (20 mL) sodium borohydride (1.26 g, 33.2 mmol) was added slowly. The reaction was stirred at room temperature for 2 hours. After completion, water (200 mL) was added to quench the reaction, and extracted with ethyl acetate (60 mL×3). The combined organic layers were washed with brine (200 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo to obtain N-((1-(Cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-2-methylpropane-2-sulfoxide amide (2.55 g, 90.9% yield). LC-MS m/z: 256 [M+H]⁺.

4). Synthesis of (1-(cyclopropylmethyl)-1H-imidazol-5-yl)methylamine hydrochloride




[0312] The mixture of N-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-2-methylpropane-2-sulfoxide amide (2.55 g, 10.00 mmol) and HCl (30 mL, 3 M in methanol) was stirred for 16 hours at room temperature. After completion, the mixture was concentrated to obtain (1-(cyclopropylmethyl)-1H-imidazol-5-yl)methanamine hydrochloride (1.87 g, 99.3% yield). LC-MS m/z: 152 [M+H]⁺.

5). Synthesis of methyl 3-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)amino)-4-nitrobenzoate




[0313] The mixture of (1-(cyclopropylmethyl)-1H-imidazol-5-yl)methanamine hydrochloride (1.87 g, 9.93 mmol), methyl 3-fluoro-4-nitrobenzoate (2.98 g, 14.90 mmol) and potassium carbonate (2.76 g, 19.87 mmol) in N,N-dimethylformamide (30 mL) was stirred at room temperature for 16 hours. After completion, the resulting mixture was diluted with water (200 mL) and extracted with ethyl acetate (50 mL×3). The combined organic layers were washed with brine (50 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=13/1) to obtain methyl 3-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)amino)-4-nitrobenzoate (1.5 g, 45.7% yield). LC-MS m/z: 331 [M+H]⁺.

6). Synthesis of methyl 4-amino-3-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)aminobenzoate



[0314] To a solution of methyl 3-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)amino)-4-nitrobenzoate (1.50 g, 4.55 mmol) in methanol (40 mL) palladium on carbon (750 mg) was added at room temperature. The resulting mixture was degassed and flushed with hydrogen three times, and the reaction was stirred at room temperature for 4 hours. After completion, the reaction was filtered, and the filter cake was washed with methanol (50 mL). The filtrate was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=13/1) to obtain 4-amino-3-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)aminobenzoate (1.1 g, 80.7% yield). LC-MS m/z: 301 [M+H]⁺.


7). Synthesis of methyl 4-(2-chloroacetamide)-3-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)aminobenzoate



[0315] To a solution of methyl 4-amino-3-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)aminobenzoate (150 mg, 0.50 mmol) in dry THF (5 mL) chloroacetic anhydride (257 mg, 1.50 mmol) was added at room temperature. The reaction was stirred at room temperature for 16 hours. After completion, the reaction solution was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=15/1) to obtain methyl 4-(2-chloroacetamide)-3-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)aminobenzoate (180 mg, 95.5% yield). LC-MS m/z: 377 [M+H]⁺.

Example 1-23. Synthesis of methyl 4-(2-chloroacetamide)-3-((1-ethyl-5-oxopyrrolidin-2-yl)methyl)aminobenzoate (Intermediate A-23)

1). Synthesis of dimethyl ethyl-L-glutamate




[0316] The mixture of dimethyl L-glutamate hydrochloride (5.00 g, 23.70 mmol) and potassium hydroxide (770 mg, 26.07 mmol) in methanol (100 mL) was stirred for 15

minutes at room temperature. Then acetaldehyde (1.56 g, 35.55 mmol) was added to the above mixture. The resulting mixture was stirred at room temperature for 2 hours. After completion, the reaction was quenched with water (200 mL) at room temperature, and extracted with ethyl acetate (100 mL×2). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo to give dimethyl N-ethyl-L-glutamate (3.50 g, 72.7% yield).


[0317] ^1H NMR (400 MHz, CDCl_3): δ 3.73 (s, 3H), 3.67 (s, 3H), 2.65-12.59 (m, 1H), 2.50-2.42 (m, 3H), 2.01-1.86 (m, 2H), 1.07 (t, $J=7.2$ Hz, 3H).

2). Synthesis of methyl 1-ethyl-5-oxopyrrolidine-2-carboxylate




[0318] The solution of dimethyl ethyl-L-glutamate (3.50 g, 17.24 mmol) in toluene (50 mL) was stirred at 110° C. for 16 hours at room temperature. After completion, the solvent was removed in vacuo to obtain methyl 1-ethyl-5-oxopyrrolidine-2-carboxylate (2.00 g, 67.8% yield). LC-MS m/z : 172 $[\text{M}+\text{H}]^+$.

3). Synthesis of 1-ethyl-5-(hydroxymethyl)pyrrolidin-2-one




[0319] To a solution of methyl 1-ethyl-5-oxopyrrolidine-2-carboxylate (1.00 g, 5.85 mmol) in tetrahydrofuran (10 mL) at 0° C. lithium aluminum hydride (222 mg, 5.85 mmol) was added. The resulting mixture was stirred at room temperature for 1 hour. Upon completion of the reaction, the reaction mixture was quenched by adding water (2 mL) and aqueous sodium hydroxide (2 mL, 15%). The reaction was filtered and concentrated under reduced pressure to give 1-ethyl-5-(hydroxymethyl)pyrrolidin-2-one (600 mg, 71.7% yield). LC-MS m/z : 144 $[\text{M}+\text{H}]^+$.

4). Synthesis of methyl (1-ethyl-5-oxopyrrolidin-2-yl)sulfonate




[0320] To a solution of 1-ethyl-5-(hydroxymethyl)pyrrolidin-2-one (600 mg, 4.20 mmol) in dichloromethane (10 mL) triethylamine (848 mg, 8.40 mmol) and methanesulfonyl chloride (580 mg, 5.04 mmol) was added at 0° C. The resulting mixture was stirred at room temperature for 1 hour. After completion, the reaction mixture was quenched with water (20 mL) at room temperature, and then extracted with dichloromethane (10 mL×2). The combined organic layers were dried over with anhydrous sodium sulfate, filtered, and concentrated in vacuo to give methyl (1-ethyl-5-oxopyrrolidin-2-yl)sulfonate (500 mg, 53.9% yield). LC-MS m/z : 222 $[\text{M}+\text{H}]^+$.

5). Synthesis of 5-(azidomethyl)-1-ethylpyrrolidin-2-one



[0321] The mixture of methyl (1-ethyl-5-oxopyrrolidin-2-yl)sulfonate (500 mg, 2.26 mmol) and sodium azide (220 mg, 3.39 mmol) in N,N -dimethylformamide (10 mL) was stirred at 120° C. for 16 hours. After completion, it was diluted with saturated sodium bicarbonate (50 mL) solution at room temperature and extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (20 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo to obtain 5-(azidomethyl)-1-ethylpyrrolidin-2-one (500 mg, crude). LC-MS m/z : 169 $[\text{M}+\text{H}]^+$.


6). Synthesis of 5-(aminomethyl)-1-ethylpyrrolidin-2-one



[0322] To a mixture of 5-(azidomethyl)-1-ethylpyrrolidin-2-one (500 mg, 2.98 mmol) in THF (10 mL) and water (2 mL) triphenylphosphine (1.56 g, 5.96 mmol) was added at


room temperature. The resulting mixture was stirred at room temperature for 2 hours. After completion, the solvent was removed by concentration. The residue was purified by silica gel column chromatography to obtain 5-(aminomethyl)-1-ethylpyrrolidin-2-one (265 mg, 99.9% yield). LC-MS m/z : 143 $[M+H]^+$.

7). Synthesis of methyl 3-((1-ethyl-5-oxypyrrolidin-2-yl)methyl)amino)-4-nitrobenzoate



[0323] The mixture of 5-(aminomethyl)-1-ethylpyrrolidin-2-one (500 mg, 3.52 mmol), methyl 3-fluoro-4-nitrobenzoate (700 mg, 3.52 mmol) and potassium carbonate (972 mg, 7.04 mmol) in N, N-dimethylformamide (10 mL) was stirred at room temperature for 16 hours. After completion, the resulting mixture was diluted with water (100 mL) and extracted with ethyl acetate (20 mL $\times$ 2). The combined organic layers were washed with brine (50 mL $\times$ 3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 3-((1-ethyl-5-oxypyrrolidin-2-yl)methyl)amino)-4-nitrobenzoate (120 mg, 10.6% yield).


8). Synthesis of methyl 4-amino-3-((1-ethyl-5-oxypyrrolidin-2-yl)methyl)amino)benzoate



[0324] To a solution of methyl 3-((1-ethyl-5-oxypyrrolidin-2-yl)methyl)amino)-4-nitrobenzoate (120

mg, 0.37 mmol) in methanol (10 mL) Pd/C (20 mg) was added at room temperature. The resulting mixture was degassed and flushed with hydrogen three times, and the reaction was stirred at room temperature for 16 hours. The reaction solution was filtered by celite with washed with MeOH (30 mL) and concentrated in vacuo to obtain methyl 4-amino-3-((1-ethyl-5-oxypyrrolidin-2-yl)methyl)amino)benzoate (100 mg, 92.9% yield). LC-MS m/z : 292 $[M+H]^+$.


9). Synthesis of methyl methyl 4-(2-chloroacetamide)-3-((1-ethyl-5-oxypyrrolidin-2-yl)methyl)aminobenzoate




[0325] To a solution of methyl 4-amino-3-((1-ethyl-5-oxypyrrolidin-2-yl)methyl)amino)benzoate (100 mg, 0.31 mmol) in dry THF (3 mL) chloroacetic anhydride (106 mg, 0.62 mmol) was slowly added. The reaction was stirred at room temperature for 2 hours. After completion, the reaction mixture was directly concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 4-(2-chloroacetamide)-3-((1-ethyl-5-oxypyrrolidin-2-yl)methyl)aminobenzoate (100 mg, 87.9% yield). LC-MS m/z : 368 $[M+H]^+$.

Example 1-24. Synthesis of methyl (S)-4-(2-chloroacetamide)-3-((1-ethylpyrrolidin-2-yl)methyl)aminobenzoate (Intermediate A-24)

1). Synthesis of methyl (S)-3-((1-ethylpyrrolidin-2-yl)methyl)amino)-4-nitrobenzoate




-continued



[0326] The mixture of methyl 3-fluoro-4-nitrobenzoate (7.77 g, 39.04 mmol), (S)-(1-ethylpyrrolidin-2-yl)carboxamide (5.00 g, 39.04 mmol) and potassium carbonate (16.16 g, 117.12 mmol) in N,N-dimethylformamide (70 mL) was stirred at room temperature for 16 hours. After completion, it was diluted with water (200 mL) and extracted with ethyl acetate (50 mL×3). The combined organic layers were washed with brine (50 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane:methanol=10:1) to obtain methyl (S)-3-((1-ethylpyrrolidin-2-yl)methyl)amino-4-nitrobenzoate (9.30 g, 77.7% yield). LC-MS m/z: 308 [M+H]⁺.


2). Synthesis of methyl (S)-4-amino-3-((1-ethylpyrrolidin-2-yl)methyl)aminobenzoate



[0327] To a solution of methyl (S)-3-((1-ethylpyrrolidin-2-yl)methyl)amino-4-nitrobenzoate (3.07 g, 10.00 mmol) in methanol (50 mL) palladium on carbon (1.50 g, 14.00 mmol) was added at room temperature. The resulting mixture was degassed and flushed with hydrogen three times, and the reaction was stirred at room temperature for 4 hours. After completion, filtered, and washed the filter cake with methanol (50 mL). The filtrate was concentrated in vacuo to give a residue. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to

obtain methyl (S)-4-amino-3-((1-ethylpyrrolidin-2-yl)methyl)aminobenzoate (2.3 g, 80.0% yield). LC-MS m/z: 278 [M+H]⁺.


3). Synthesis of methyl (S)-4-(2-chloroacetamide)-3-((1-ethylpyrrolidin-2-yl)methyl)aminobenzoate




[0328] To a solution of methyl (S)-4-amino-3-((1-ethylpyrrolidin-2-yl)methyl)aminobenzoate (120 mg, 0.43 mmol) in dry THF (20 mL) chloroacetic anhydride (111 mg, 0.65 mmol) was slowly added at room temperature. The mixture was stirred at room temperature for 2 hours. After completion, the mixture was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain (S)-4-(2-chloroacetamide)-3-((1-ethylpyrrolidin-2-yl)methyl)aminobenzoate (140 mg, 91.5% yield). LC-MS m/z: 354 [M+H]⁺.

Example 1-25. Synthesis of (S)-N-(4-Bromo-2-methyl-6-((oxetan-2-ylmethyl)amino)phenyl)-2-chloroacetamide (Intermediate A-25)

1). Synthesis of (S)-5-bromo-3-methyl-2-nitro-N-(oxetan-2-ylmethyl) aniline




-continued




[0329] The mixture of 5-bromo-1-fluoro-3-methyl-2-nitrobenzene (500 mg, 2.14 mmol), (S)-oxetan-2-ylmethanamine (186 mg, 2.14 mmol) and potassium carbonate (590 mg, 4.28 mmol) in N, N-dimethylformamide (5 mL) was stirred at room temperature for 16 hours. After completion, water (5 mL) was added to quench the reaction, and it was extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (20 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=4/1) to obtain (S)-5-bromo-3-methyl-2-nitro-N-((oxetan-2-ylmethyl)amino)aniline (490 mg, 6.2% yield). LC-MS m/z : 301 $[M+H]^+$.

2). Synthesis of (S)-5-bromo-3-methyl-N-((oxetan-2-ylmethyl)amino)benzene-1,2-diamine



[0330] To a solution of (S)-5-bromo-3-methyl-2-nitro-N-((oxetan-2-ylmethyl)amino)aniline (490 mg, 1.63 mmol) in methanol (5 mL) was added zinc dust (1.06 g, 16.30 mmol). The resulting mixture was stirred at room temperature for 2 hours. After completion, the reaction solution was filtered, and then the filtrate was adjusted to pH=7-8 with aqueous ammonia. After this, the mixture was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain (S)-5-bromo-3-methyl-N-((oxetan-2-ylmethane)yl)benzene-1,2-diamine (269 mg, 60.7% yield), LC-MS m/z : 271 $[M+H]^+$.


3). Synthesis of (S)-N-(4-bromo-2-methyl-6-((oxetan-2-ylmethyl)amino)phenyl)-2-chloroacetamide



[0331] To a solution of (S)-5-bromo-3-methyl-N-((oxetan-2-ylmethyl)amino)benzene-1,2-diamine (109 mg, 0.40 mmol) in THF (3 mL) chloroacetic anhydride (75 mg, 0.44 mmol) was added dropwise at room temperature. The resulting mixed solution was stirred and reacted at room temperature for 1 hour. After completion, the reaction solution was concentrated, and then was purified by silica gel column chromatography (petroleum ether/ethyl acetate=1/1) to obtain (S)-N-(4-bromo-2-methyl-6-((oxetan-2-ylmethyl)amino)phenyl)-2-chloroacetamide (98 mg, 70.4% yield), LC-MS m/z : 349 $[M+H]^+$.


Example 1-26. Synthesis of methyl (S)-2-(chloromethyl)-4-methyl-1-((oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (intermediate A-26)

1). Synthesis of methyl (S)-4-amino-3-methyl-5-((oxetan-2-ylmethyl)amino)benzoate



[0332] The solution of (S)-5-bromo-3-methyl-N-(oxetan-2-ylmethyl)benzene-1,2-diamine (130 mg, 0.48 mmol), 1, 1'-bis(diphenylphosphino)ferrocene palladium dichloride (37 mg, 0.05 mmol) and potassium acetate (141 mg, 1.44 mmol) in N,N-dimethylformamide (2 mL) and methanol (2 mL) was stirred at 90° C. for 16 hours under CO atmosphere (58.76 psi). After completion, the reaction solution was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl (S)-4-amino-3-methyl-5-((oxetan-2-ylmethyl)amino)benzoate (50 mg, 41.5% yield). LC-MS m/z: 251 [M+H]⁺.


2). Synthesis of methyl (S)-2-(chloromethyl)-4-methyl-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate



[0333] To a solution of methyl (S)-4-amino-3-methyl-5-((oxetan-2-ylmethyl)amino)benzoate (50 mg, 0.20 mmol) in THF (2 mL) chloroacetic anhydride (51 mg, 0.30 mmol) was added dropwise. The resulting mixture solution was stirred at room temperature for 1 hour. After completion, the reaction solution was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=1/1) to obtain (S)-2-(chloromethyl)-4-methyl-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (20 mg, 32.6% yield). LC-MS m/z: 309 [M+H]⁺.

Example 1-27. Synthesis of methyl 4-(2-chloroacetamide)-3-(1-ethyl-1H-pyrazol-5-yl)methyl)amino-benzoate (Intermediate A-27)


1). Synthesis of methyl 1-ethyl-1H-pyrazole-5-carboxylate



[0334] To a solution of 1-Ethyl-1H-pyrazole-5-carboxylic acid (770 mg, 5.50 mmol) in methanol (8 mL) (diazomethyl)


trimethylsilane (27.5 mL, 55.00 mmol, 2.0 M in tetrahydrofuran) was dropwise added at 0° C. The resulting mixture was stirred at room temperature for 1 hour, then quenched with water (100 mL) and extracted with ethyl acetate (50 mL×2). The combined organic layers were washed with brine (50 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo to obtain methyl 1-ethyl-1H-pyrazole-5-carboxylate (600 mg, 70.8% yield), LC-MS m/z: 155 [M+H]⁺.

2). Synthesis of (1-ethyl-1H-pyrazol-5-yl)methanol




[0335] To a solution methyl 1-ethyl-1H-pyrazole-5-carboxylate (600 mg, 3.90 mmol) in dry THF (8 mL) lithium aluminum hydride powder (296 mg, 7.80 mmol) was added at 0° C. The reaction was stirred at room temperature for 1 hour. After completion, the reaction was quenched with water (2 mL) and aqueous sodium hydroxide (2 mL, 15%). The mixture was filtered, and concentrated in vacuo to give (1-ethyl-1H-pyrazol-5-yl)methanol (400 mg, 81.4% yield). LC-MS m/z: 127 [M+H]⁺.

3). Synthesis of 5-(chloromethyl)-1-ethyl-1H-pyrazole




[0336] To a solution of (1-Ethyl-1H-pyrazol-5-yl)methanol (400 mg, 3.17 mmol) in dichloromethane (20 mL) triethylamine (640 mg, 6.34 mmol) and methanesulfonyl chloride (729 mg, 6.34 mmol) were added at 0° C. The reaction was stirred at room temperature for 2 hours. After completion, the reaction was quenched with water (100 mL) and extracted with dichloromethane (50 mL×2). The combined organic layer was washed with brine (50 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo to obtain 5-(chloromethyl)-1-ethyl-1H-pyrazole (400 mg, 87.0% yield), LC-MS m/z: 145 [M+H]⁺.

4). Synthesis of 5-(azidomethyl)-1-ethyl-1H-pyrazole




[0337] The mixture of 5-(chloromethyl)-1-ethyl-1H-pyrazole (400 mg, 2.76 mmol) and sodium azide (359 mg, 5.52 mmol) in N, N-dimethyl formamide was stirred at 120° C. for 16 hours. After completion, it was diluted with saturated sodium bicarbonate solution (10 mL) and extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (10 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo to obtain 5-(azidomethyl)-1-ethyl-1H-pyrazole (370 mg, 88.8% yield). LC-MS m/z: 152 [M+H]⁺.

5). Synthesis of
(1-ethyl-1H-pyrazol-5-yl)methanamine




[0338] To a mixture of 5-(Azidomethyl)-1-ethyl-1H-pyrazole (320 mg, 2.12 mmol) in THF (4 mL) and water (0.4 mL) triphenylphosphine (1.1 g, 4.24 mmol) was added at room temperature. The reaction was stirred at room temperature for 5 hours. After completion, the solvent was removed by concentration in vacuo to obtain (1-ethyl-1H-pyrazol-5-yl)methanamine (265 mg, 99.9% yield).

6). Synthesis of methyl 3-((1-ethyl-1H-pyrazol-5-yl)methyl)amino)-4-nitrobenzoate




[0339] The mixture of (1-ethyl-1H-pyrazol-5-yl)methanamine (265 mg, 2.12 mmol), methyl 3-fluoro-4-nitrobenzoate (422 mg, 2.12 mmol) and potassium carbonate (585 mg, 4.24 mmol) in N,N-dimethylformamide (7 mL) was stirred at room temperature for 16 hours. After completion, it was diluted with water (60 mL) and extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (20 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to give methyl 3-((1-ethyl-1H-pyrazol-5-yl)methyl)amino)-4-nitrobenzoate (100 mg, 15.5% yield). LC-MS m/z: 346 [M+H]⁺.

7). Synthesis of methyl 4-amino-3-((1-ethyl-1H-pyrazol-5-yl)methyl)amino)benzoate



[0340] To solution of methyl 3-((1-ethyl-1H-pyrazol-5-yl)methyl)amino)-4-nitrobenzoate (130 mg, 0.43 mmol) in methanol (12 mL) palladium on carbon (50 mg, 30.42 mmol) was added at room temperature. The resulting mixture was degassed and flushed with hydrogen three times, and the reaction was stirred at room temperature for 16 hours. After completion, the reaction solution was filtered, the filter cake was rinsed with methanol. The filtrate was concentrated in vacuo to obtain methyl 4-amino-3-((1-ethyl-1H-pyrazol-5-yl)methyl)amino)benzoate (70 mg, 59.4% yield), LC-MS m/z: 275 [M+H]⁺.


8). Synthesis of methyl 4-(2-chloroacetamide)-3-(1-ethyl-1H-pyrazol-5-yl)methyl)aminobenzoate



[0341] To a solution of methyl 4-amino-3-((1-ethyl-1H-pyrazol-5-yl)methyl)amino)benzoate (70 mg, 0.26 mmol) in tetrahydrofuran (3 mL) chloroacetic anhydride (111 mg, 0.65 mmol) was slowly added. The reaction was stirred at room temperature for 2 hours. After completion, the solvent was removed by concentration in vacuo to give a residue. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 4-(2-chloroacetamido)-3-(1-ethyl-1H-pyrazol-5-yl)methyl)aminobenzoate (60 mg, 65.7% yield). LC-MS m/z: 351 [M+H]⁺.


Example 1-28. Synthesis of methyl 4-(2-chloroacetamide)-3-((1-ethyl-3-methyl-1H-pyrazol-5-yl)methyl)-amino)benzoate (Intermediate A-28)

1). Synthesis of methyl 1-ethyl-3-methyl-1H-pyrazole-5-carboxylate




[0342] To a solution of 1-Ethyl-3-methyl-1H-pyrazole-5-carboxylic acid (1.00 g, 6.49 mmol) in methanol (10 mL) (diazomethyl)trimethylsilane (32.45 mL, 64.90 mmol, 2.0 M in THF) was added dropwise at 0° C. The resulting mixture was stirred at room temperature for 1 hour. After completion, the mixture was quenched with water (300 mL) and extracted with ethyl acetate (100 mL×3). The combined organic layers were washed with brine (100 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo to give methyl 1-ethyl-3-methyl-1H-pyrazole-5-carboxylate (800 mg, 73.4% yield). LC-MS m/z: 169 [M+H]⁺.

2). Synthesis of (1-ethyl-3-methyl-1H-pyrazol-5-yl)methanol




[0343] To solution of methyl 1-ethyl-3-methyl-1H-pyrazole-5-carboxylate (800 mg, 4.76 mmol) in dry THF (10 mL) Aluminum lithium hydride (543 mg, 14.28 mmol) was added at 0° C. The reaction was stirred at room temperature for 1 hour. After completion, the mixture was quenched with water (2 mL) and aqueous sodium hydroxide solution (2 mL, 15% w/w). The mixture was filtered, and the filtrate was concentration in vacuo to give (1-ethyl-3-methyl-1H-pyrazol-5-yl)methanol (550 mg, 82.5% yield). LC-MS m/z: 141. [M+H]⁺.

3). Synthesis of 5-(chloromethyl)-1-ethyl-3-methyl-1H-pyrazole




[0344] To a mixture of (1-ethyl-3-methyl-1H-pyrazol-5-yl)methanol (550 mg, 3.93 mmol) in dichloromethane (20 mL) triethylamine (794 mg, 7.86 mmol) and methanesulfonyl chloride (679 mg, 5.90 mmol) was added at 0° C. The reaction was stirred at room temperature for 2 hours. After completion, the mixture was quenched with water (100 mL) and extracted with dichloromethane (50 mL×2). The combined organic layer was washed with brine (50 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo to obtain 5-(chloromethyl)-1-ethyl-3-methyl-1H-pyrazole (500 mg, 80.0% yield). LC-MS m/z: 159 [M+H]⁺.

4). Synthesis of 5-(azidomethyl)-1-ethyl-3-methyl-1H-pyrazole




[0345] The mixture of 5-(chloromethyl)-1-ethyl-3-methyl-1H-pyrazole (450 mg, 2.83 mmol) and sodium azide (368 mg, 5.66 mmol) in N, N-dimethylformamide (4 mL) was stirred at 120° C. for 16 hours. After completion, it was diluted with saturated sodium bicarbonate solution (10 mL) and extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (10 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo to obtain 5-(azidomethyl)-1-ethyl-3-methyl-1H-pyrazole (350 mg, 75.0% yield). LC-MS m/z: 166 [M+H]⁺.


5). Synthesis of (1-ethyl-3-methyl-1H-pyrazol-5-yl)methanamine



[0346] To a mixture of 5-(azidomethyl)-1-ethyl-3-methyl-1H-pyrazole (350 mg, 2.12 mmol) in THF (7 mL) and water (0.7 mL) triphenylphosphine (1.1 g, 4.24 mmol) was added, then stirred at room temperature for 5 hours. After completion, the solvent was removed in vacuo to obtain a colorless oil compound (1-ethyl-3-methyl-1H-pyrazol-5-yl)methanamine (286 mg, 97.1% yield). LC-MS m/z: 140 [M+H]⁺.


6) Synthesis of methyl 3-((1-ethyl-3-methyl-1H-pyrazol-5-yl)methyl)amino)-4-nitrobenzoate





[0347] The mixture of (1-ethyl-3-methyl-1H-pyrazol-5-yl)methanamine (286 mg, 2.06 mmol), methyl 3-fluoro-4-nitrobenzoate (410 mg, 2.06 mmol) and potassium carbonate (569 mg, 4.12 mmol) in N,N-dimethylformamide (8 mL) was stirred at room temperature for 16 hours. After completion, it was diluted with water (50 mL) and extracted with ethyl acetate (25 mL×2). The combined organic layers were washed with brine (25 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to give methyl 3-((1-ethyl-3-methyl-1H-pyrazol-5-yl)methyl)amino)-4-nitrobenzoate (300 mg, 45.8% yield). LC-MS *m/z*: 319 [M+H]⁺.


7). Synthesis of methyl 4-amino-3-((1-ethyl-3-methyl-1H-pyrazol-5-yl)methyl)amino)benzoate



[0348] To a solution of methyl 3-((1-ethyl-3-methyl-1H-pyrazol-5-yl)methyl)amino)-4-nitrobenzoate (330 mg, 1.04 mmol) dissolved in methanol (24 mL) palladium on carbon (50 mg, 30.42 mmol) was added at room temperature. The resulting mixture was degassed and flushed with hydrogen

three times and stirred at room temperature for 16 hours. After completion, the reaction solution was filtered, the filter cake was rinsed with methanol. The filtrate was concentrated in vacuo to obtain methyl 4-amino-3-((1-ethyl-3-methyl-1H-pyrazol-5-yl)methyl)amino)benzoate (200 mg, 67.1% yield), LC-MS *m/z*: 289 [M+H]⁺.

8). Synthesis of methyl 4-(2-chloroacetamide)-3-((1-ethyl-3-methyl-1H-pyrazol-5-yl)methyl)amino)benzoate








[0349] To a solution of methyl 4-amino-3-((1-ethyl-3-methyl-1H-pyrazol-5-yl)methyl)amino)benzoate (200 mg, 0.69 mmol) in THF (4 mL) chloroacetic anhydride (296 mg, 1.73 mmol) was slowly added at room temperature. The reaction was stirred at room temperature for 2 hours. After completion, the solvent was removed in vacuo to give a residue. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 4-(2-chloroacetamide)-3-((1-ethyl-3-methyl-1H-pyrazol-5-yl)methyl)amino)benzoate (200 mg, 79.4% yield). LC-MS *m/z*: 365 [M+H]⁺.

Example 1-29. Synthesis of Other Part a Intermediates






[0350] The following intermediates can be obtained with similar to the synthetic steps of each intermediate from Example 1-1 to Example 1-28

Number	Structure	Name	MS <i>m/z</i> : [M + H] ⁺
Intermediate A-29		Methyl (E)-3-(4-(2-chloroacetamido)-3-((1-cyanomethyl)cyclopropyl)methyl)amino)phenyl)acrylate	362






-continued

Number	Structure	Name	MS m/z: [M + H] ⁺
IntermediateA-30		Methyl (E)-3-(5-(2-chloroacetamido)-6-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)pyridin-2-yl)acrylate	378
IntermediateA-31		Methyl 2-(chloromethyl)-1-((1-ethyl-1H-imidazol-4-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate	333
IntermediateA-32		Methyl (E)-3-(4-(2-chloroacetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)-5-fluorophenyl)acrylate	395
IntermediateA-33		Methyl (E)-3-(5-(2-chloroacetamido)-6-(((1-(cyanomethyl)cyclopropyl)methyl)amino)pyridin-2-yl)acrylate	363
IntermediateA-34		Methyl (E)-3-(4-(2-chloroacetamido)-3-(((1-(cyanomethyl)cyclopropyl)methyl)amino)-5-fluorophenyl)acrylate	380

-continued

Number	Structure	Name	MS m/z: [M + H] ⁺
IntermediateA-35		Methyl (S,E)-3-(5-(2-chloroacetamido)-6-((oxetan-2-ylmethyl)amino)pyridin-2-yl)acrylate	340
IntermediateA-36		Methyl (S,E)-3-(4-(2-chloroacetamido)-3-fluoro-5-((oxetan-2-ylmethyl)amino)phenyl)acrylate	357
IntermediateA-37		Methyl (E)-3-(5-(2-chloroacetamido)-6-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)pyridin-2-yl)acrylate	378
IntermediateA-38		Methyl 2-(5-(2-chloroacetamido)-6-(((1-ethyl-1H-imidazol-5-yl)methyl)aminopyridin-2-yl)acetate	366
IntermediateA-39		Methyl 2-(4-(2-chloroacetamido)-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)phenyl)acetate	365

-continued

Number	Structure	Name	MS m/z: [M + H] ⁺
IntermediateA- 40		Methyl 2-(4-(2-chloroacetamido)-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)-5-fluorophenyl)acetate	383
IntermediateA- 41		(S)-Methyl 5-(2-chloroacetamido)-6-((oxetan-2-ylmethyl)amino)-picolinate	314
IntermediateA- 42		Methyl (S)-4-(2-chloroacetamido)-3-fluoro-5-((oxetan-2-ylmethyl)amino)-benzoate	331
IntermediateA- 43		Methyl (S)-2-(4-(2-chloroacetamido)-3-fluoro-5-((oxetan-2-ylmethyl)amino)phenyl)-acetate	345
IntermediateA- 44		Methyl (S)-2-(5-(2-chloroacetamido)-6-((oxetan-2-ylmethyl)amino)pyridin-2-yl)acetate	328

-continued

Number	Structure	Name	MS m/z: [M + H] ⁺
IntermediateA-45		Methyl 2-(4-(2-chloroacetamido)-3-(((1-(cyanomethyl)cyclopropyl)methyl)amino)phenyl)acetate	350
IntermediateA-46		Methyl 2-(4-(2-chloroacetamido)-3-(((1-(cyanomethyl)cyclopropyl)methyl)amino)-5-fluorophenyl)acetate	368
IntermediateA-47		Methyl 2-(5-(2-chloroacetamido)-6-(((1-(cyanomethyl)cyclopropyl)methyl)amino)pyridin-2-yl)acetate	351
IntermediateA-48		Methyl (R)-2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate	295
IntermediateA-49		2-(chloromethyl)-4-fluoro-1-((1-isopropyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid	351


-continued

Number	Structure	Name	MS m/z: [M + H] ⁺
Intermediate A-50		2-(chloromethyl)-1-(1-isopropyl-1H-imidazol-5-yl)methyl-1H-benzo[d]imidazole-6-carboxylic acid	333

② indicates text missing or illegible when filed


Example 2-1. Synthesis of 3-fluoro-4-((6-(piperidin-4-oxy)pyridin-2-yl)methoxy)benzonitrile (Intermediate B-1)

1). Synthesis of tert-butyl 4-(6-bromopyridin-2-yl)oxy)piperidine-1-carboxylate




[0351] To a solution of compound tert-butyl 4-hydroxypiperidine-1-carboxylate (2.00 g, 9.95 mmol) in tetrahydrofuran (30 mL) sodium hydride (6000 w/w dispersed in mineral oil, 597 mg, 14.93 mmol) was added at 0° C. The mixture was stirred at 0° C. for 30 min, then 2-bromo-6-fluoropyridine (2.09 g, 11.94 mmol) was added slowly to the mixture. The reaction mixture was stirred at 70° C. for 4 hours. After completion, the reaction was quenched with water (30 mL) at room temperature and extracted with ethyl acetate (50 mL×3). The combined organic layers were washed with brine (30 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=10/1) to obtain tert-butyl 4-(6-bromopyridin-2-yl)oxy)piperidine-1-carboxylate (1.80 g, 5.06 mmol, 50.9% yield). LC-MS m/z: 356.9, 358.9 [M+H]⁺.

2). Synthesis of methyl 6-((1-(tert-butoxycarbonyl)piperidin-4-yl)oxy)picolinate




-continued



[0352] To a solution of 4-(6-bromopyridin-2-yl)oxy)piperidine-1-carboxylic acid tert-butyl ester (1.80 g, 5.06 mmol) in methanol (5 mL) potassium acetate (1.49 g, 15.18 mmol) and 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (373 mg, 0.51 mmol) were added. The mixture was stirred at 90° C. for 16 h under CO (58.76 psi) atmosphere. The reaction was quenched by water (15 mL) and then extracted ethyl acetate (30 mL×3). The combined organic layers were washed with brine (30 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. The residue was purified by reverse-phase flash chromatography under the following conditions: (Column: spherical C18, 20-40 μm, 120 g; mobile phase A: 10 mM NH₄OH in water; Mobile Phase B: Acetonitrile; Flow rate: 80 mL/min; Gradient: 60% B-80% B in 20 minutes; detector: 254 nm.) The mobile phase containing the desired product was collected at 70% B and then concentrated in vacuo to give methyl 6-((1-(tert-butoxycarbonyl)piperidin-4-yl)oxy)picolinate (1.50 g, 4.46 mmol, 88.1% yield). LC-MS m/z: 337 [M+H]⁺.


3). Synthesis of tert-butyl 4-(6-(hydroxymethyl)pyridin-2-yl)oxy)piperidine-1-carboxylate



[0353] To a solution of methyl 6-((1-(tert-butoxycarbonyl)piperidin-4-yl)oxy)picolinate (500 mg, 1.49 mmol) in


methanol (5 mL) sodium borohydride (170 mg, 4.47 mmol) and lithium chloride (6 mg, 0.15 mmol) were added at 0° C. The mixture was stirred at 50° C. for 16 hours. The reaction was quenched by water (5 mL) and then extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (10 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by reverse-phase flash chromatography under the following conditions: (Column: spherical C18, 20-40 μm, 120 g; Mobile Phase A: 10 mM NH₄OH in water; Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 60% B-80% B in 20 minutes; detector: 254 nm) The mobile phase containing the desired product was collected and concentrated in vacuo to give tert-butyl 4-(6-(hydroxymethyl)pyridin-2-yl)oxy)piperidine-1-carboxylate (200 mg, 0.65 mmol, 43.6% yield). LC-MS m/z: 309 [M+H]⁺.

4). Synthesis of tert-butyl 4-(6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidine-1-carboxylate




[0354] To a solution of tert-butyl 4-(6-(hydroxymethyl)pyridin-2-yl)oxy)piperidine-1-carboxylate (200 mg, 0.65 mmol) and 3-fluoro-4-hydroxybenzonitrile (89 mg, 0.65 mmol) in tetrahydrofuran (10 mL) triphenylphosphine (257 mg, 0.98 mmol) was added at 0° C. Diisopropyl azodicarboxylate (198 mg, 0.98 mmol) was added dropwise at 0° C. The mixture was then stirred at room temperature for 16 hours. The resulting mixture was concentrated and purified by silica gel column chromatography (ethyl acetate/petroleum ether=1/1) to obtain tert-butyl 4-(6-((4-cyano-2-fluorophenoxy)methyl)pyridine-2-yl)oxy)piperidine-1-carboxylate (250 mg, 0.59 mmol, 90.8% yield). LC-MS m/z: 428 [M+H]⁺.

5). Synthesis of 3-fluoro-4-((6-(piperidin-4-oxymethyl)pyridin-2-yl)methoxy)benzonitrile (Intermediate B-1)




-continued



[0355] To a solution of tert-butyl 4-(6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidine-1-carboxylate (250 mg, 0.59 mmol) in dichloromethane (15 mL) dropwise trifluoroacetic acid (3 mL) was added at room temperature. The mixture stirred at room temperature for 2 hours. After completion, the mixture was concentrated in vacuo to give the crude product, which was further purified by reverse-phase flash chromatography under the following conditions: (Column: spherical C18, 20-40 μm, 120 g; Mobile Phase A: 10 mM NH₄OH in water; Mobile phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 40% B-60% B in 20 minutes; Detector: 254 nm.) The mobile phase containing the desired product at 52% B was collected and concentrated in vacuo to obtain 3-fluoro-4-((6-(piperidin-4-oxymethyl)pyridin-2-yl)methoxy)benzonitrile (180 mg, 0.55 mmol, 93.2% yield), LC-MS m/z: 328 [M+H]⁺.


Example 2-2. Synthesis of 3-fluoro-4-((6-((piperidin-4-yloxy)methyl)pyridin-2-yloxy)methyl)benzonitrile (Intermediate B-2)

1). Synthesis of tert-butyl 4-(6-(fluoropyridin-2-yl)methoxy)piperidine-1-carboxylate




[0356] To a solution of tert-butyl 4-hydroxypiperidine-1-carboxylate (1.06 g, 5.26 mmol) in tetrahydrofuran (15 mL) sodium hydride (252 mg, 6.31 mmol, 60% w/w dispersed in mineral oil) was added at 0° C. Then the mixture was stirred for 30 min. After this, the 2-(bromomethyl)-6-fluoropyridine (500 mg, 2.63 mmol) was added slowly to above mixture. The mixture was stirred at room temperature for 16 hours, quenched with methanol (10 mL) at room temperature and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to give tert-butyl 4-(6-(fluoropyridin-2-yl)methoxy)piperidine-1-carboxylate (447 mg, 54.8% yield), LC-MS m/z: 311 [M+H]⁺.

2). Synthesis of tert-butyl 4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)methoxy)piperidine-1-carboxylate



[0357] To a solution of 3-fluoro-4-(hydroxymethyl)benzonitrile (581 mg, 3.85 mmol) in tetrahydrofuran (10 mL) sodium hydride (185 mg, 4.62 mmol, 60% w/w dispersed in mineral oil) was added at 0° C., the mixture was stirred for 30 min, then the tert-butyl 4-((6-fluoropyridin-2-yl)methoxy)piperidine-1-carboxylate (240 mg, 0.77 mmol) was added slowly at 0° C. The mixture was stirred at 70° C. for 16 hours. After completion, the mixture was quenched with water (20 mL) at room temperature and extracted with ethyl acetate (20 mL×2). The combined organic layers were washed with brine (10 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to give tert-butyl 4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)methoxy)piperidine-1-carboxylate (281 mg, 82.4% yield). LC-MS m/z : 442 $[M+H]^+$

3). Synthesis of 3-fluoro-4-((6-((piperidin-4-yloxy)methyl)pyridin-2-yloxy)methyl)benzonitrile




[0358] To a solution of tert-butyl 4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)methoxy)piperidine-1-carboxylate (400 mg, 0.91 mmol) in dichloromethane (2 mL) trifluoroacetic acid (1 mL) was added. The mixture was stirred at room temperature for 16 hours. After complete, the mixture was concentrated to remove the dichloromethane and then adjusted pH to 8-9 with $NH_3/MeOH$ (7 M). After this, the mixture was directly concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to give

3-fluoro-4-((6-((piperidin-4-yloxy)methyl)pyridin-2-yl oxy)methyl)benzonitrile (301 mg, 97.4% yield). LC-MS m/z : 342 $[M+H]^+$.


Example 2-3. Synthesis of 3-fluoro-4-((6-((piperidin-4-ylmethoxy)pyridin-2-yl)oxy)methyl)benzonitrile (Intermediate B-3)

1). Synthesis of tert-butyl 4-((6-fluoropyridin-2-yl)oxy)methyl)piperidine-1-carboxylate



[0359] To a solution of tert-butyl 4-(hydroxymethyl)piperidine-1-carboxylate (3.0 g, 13.93 mmol) in dry THF (50 mL) sodium hydride (837 mg, 20.93 mmol, 60% w/w dispersion in mineral oil) was added at 0° C. The mixture was stirred at 0° C. for 30 minutes, then 2,6-difluoropyridine (2.41 g, 20.94 mmol) was slowly added at 0° C. The mixture was stirred at 70° C. for 5 hours. After completion, the mixture was quenched with water (200 mL) at 0° C. and extracted with dichloromethane (50 mL×3). The combined organic layer was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (PE/EA=4/1) to give tert-butyl 4-((6-fluoropyridin-2-yl)oxy)methyl)piperidine-1-carboxylate (2.0 g, 6.44 mmol, 46.2% yield). LC-MS m/z : 311 $[M+H]^+$.


2). Synthesis of tert-butyl 4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)methyl)piperidine-1-carboxylate



[0360] To a solution of 3-fluoro-4-(hydroxymethyl)benzonitrile (365 mg, 2.41 mmol) in dry THF (50 mL) sodium hydride (97 mg, 2.42 mmol, 60% w/w dispersion in mineral oil) was added at 0° C. The mixture was stirred at 0° C. for 30 min, then the tert-butyl 4-((6-fluoropyridin-2-yl)oxy)methyl)piperidine-1-carboxylate (500 mg, 1.61 mmol) was

slowly added to above mixture at 0° C. The mixture was stirred at 70° C. for 5 hours. After this, the mixture was quenched with water (30 mL) at 0° C. and extracted with dichloromethane (10 mL×3). The combined organic layer was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (PE/EA=5/1) to give tert-butyl 4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)methyl)piperidine-1-carboxylate (400 mg, 0.91 mmol, 56.3% yield). LC-MS m/z: 442 [M+H]⁺.



3). Synthesis of 3-fluoro-4-((6-(piperidin-4-ylmethoxy)pyridin-2-yl)oxy)methyl)benzonitrile



[0361] To a solution of tert-butyl 4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)methyl)piperidine-1-carboxylate (400 mg, 0.91 mmol) in dichloromethane (5 mL) trifluoroacetic acid (1 mL) was added and stirred at room temperature for 16 hours. When completed, The reaction was quenched with saturated sodium bicarbonate (30 mL) solution and then extracted with dichloromethane (10 mL×2). The combined organic layer was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (DCM/MeOH=10/1) to give 3-fluoro-4-((6-(piperidin-4-ylmethoxy)pyridin-2-yl)oxy)methyl)benzonitrile (250 mg, 0.73 mmol, 80.6% yield). LC-MS m/z: 342 [M+H]⁺.

Example 2-4. Synthesis of 3-fluoro-4-((6-(piperidin-4-ylmethoxy)pyridin-2-yl)oxy)methyl)benzonitrile (Intermediate B-4)


1). Synthesis of tert-butyl 4-((6-fluoropyridin-2-yl)oxy)piperidine-1-carboxylate



[0362] To a solution of tert-butyl 4-hydroxypiperidine-1-carboxylate (3.50 g, 17.39 mmol) in dry tetrahydrofuran (20 mL) sodium hydride (870 mg, 21.74 mmol, 60% w/w dispersion in mineral oil) was added at 0° C. The mixture was stirred at 0° C. for 30 min, then 2,6-difluoropyridine (2.00 g, 17.38 mmol) was slowly added to above mixture at 0° C. The mixture was warmed to 80° C. and stirred for 4 hours. The mixture was quenched with water (20 mL) at room temperature and then extracted with ethyl acetate (20 mL×2). The combined organic layers were washed with brine (10 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (PE/EA=10/1) to obtain tert-butyl 4-((6-fluoropyridin-2-yl)oxy)piperidine-1-carboxylate (5.10 g, 17.21 mmol, 99.2% yield).


[0363] ¹H NMR (400 MHz, DMSO-d₆) δ 7.86 (dd, J=16.8, 8.0 Hz, 1H), 6.73 (dd, J=8.0, 1.6 Hz, 1H), 6.68 (dd, J=7.6, 2.4 Hz, 1H), 5.06-5.02 (m, 1H), 3.70-3.64 (m, 2H), 3.20-3.15 (m, 2H), 1.96-1.90 (m, 2H), 1.58-1.49 (m, 2H), 1.40 (s, 9H).

2). Synthesis of tert-butyl 4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidine-1-carboxylate



[0364] Sodium hydride (541 mg, 13.52 mmol, 60% w/w dispersion in mineral oil) was added to a solution of 3-fluoro-4-((hydroxymethyl)benzonitrile (1.22 g, 8.07 mmol) in dry THF (20 mL). The mixture was stirred at 0° C. for 30 minutes, then the tert-butyl 4-((6-fluoropyridin-2-yl)oxy)piperidine-1-carboxylate (2.00 g, 6.75 mmol) was slowly added to the above mixture at 0° C. The reaction mixture was warmed to 70° C. and stirred for 3 hours. After completion, the reaction solution was cooled to room temperature, quenched with water (20 mL), and extracted with EA (20 mL×2). The combined organic layers were washed with brine (10 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (PE/EA=10/1) to give tert-butyl 4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidine-1-carboxylate (1.52 g, 3.56 mmol, 55.6% yield). LC-MS m/z: 428 [M+H]⁺.


3). Synthesis of 3-fluoro-4-((6-(piperidin-4-oxy)pyridin-2-yl)oxy)methyl)benzonitrile



[0365] To a solution of tert-butyl 4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidine-1-carboxylate (1.52 g, 3.56 mmol) in dichloromethane (15 mL) trifluoroacetic acid (3 mL) was added. The reaction solution was stirred at room temperature for 16 hours. After completion, the reaction mixture was adjusted pH to 8-9 with NH_3/MeOH (7M) and then concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (DCM/MeOH=10/1) to give 3-fluoro-4-((6-(piperidin-4-oxy)pyridin-2-yl)oxy)methyl)benzonitrile (1.00 g, 3.05 mmol, 86.2% yield). LC-MS m/z : 328 $[\text{M}+\text{H}]^+$


Example 2-5. Synthesis of 3-fluoro-4-((6-(piperidin-4-oxy)methyl)pyridin-2-yl)methoxy)benzonitrile (Intermediate B-5)

1). Synthesis of methyl 6-(chloromethyl)picolinate




[0366] To a solution of methyl 6-(hydroxymethyl)picolinate (2.00 g, 11.98 mmol) in dichloromethane (20 mL) triethylamine (2.42 g, 23.96 mmol) and methanesulfonyl chloride (1.65 g, 14.38 mmol) was sequentially added at 0°C. slowly. The mixture was stirred at room temperature for 3 hours. The mixture was quenched with water (10 mL) and then extracted with dichloromethane (20 mL $\times$ 2). The combined organic layers were washed with brine (10 mL $\times$ 2), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=4/1) to obtain methyl 6-(chloromethyl)picolinate (1.2 g, 54.2% yield). LC-MS m/z : 188 $[\text{M}+\text{H}]^+$.

2). Synthesis of 6-((1-(tert-butoxycarbonyl)piperidin-4-yl)oxy)methyl)picolinic acid



[0367] To a solution of tert-butyl 4-hydroxypiperidine-1-carboxylate (4.92 g, 24.50 mmol) in tetrahydrofuran (50 mL) sodium hydride (1.18 g, 29.40 mmol, 60% w/w dispersed in mineral oil) was added at 0°C. The mixture was stirred for 30 min, then the methyl 6-(chloromethyl)picolinate (454 mg, 2.45 mmol) was slowly added above mixture at 0°C. The mixture was stirred at room temperature for 16 hours. The reaction was then quenched with water (10 mL) at room temperature and extracted with ethyl acetate (20 mL $\times$ 2). The combined organic layers were washed with brine (10 mL $\times$ 2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to give 6-((1-(tert-butoxycarbonyl)piperidin-4-yl)oxy)methyl)picolinic acid (356 mg, 43.2% yield). LC-MS m/z : 337 $[\text{M}+\text{H}]^+$.


3). Synthesis of methyl 6-((piperidine-4-oxy)methyl)picolinate



[0368] To a solution of 6-((1-(tert-butoxycarbonyl)piperidin-4-yl)oxy)methyl)picolinic acid (356 mg, 1.06 mmol) in methanol (5 mL) thionyl chloride (189 mg, 1.59 mmol) was added dropwise at 0°C. The mixture was stirred at 60°C. for 2 hours. The mixture was quenched with water (10 mL) at room temperature and extracted with ethyl acetate (10 mL $\times$ 2). The combined organic layers were washed with saturated sodium bicarbonate (10 mL), brine (10 mL), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to give methyl 6-((piperidine-4-oxy)methyl)picolinate (1.2 g, 54.2% yield). LC-MS m/z : 188 $[\text{M}+\text{H}]^+$.


nol=10/1) to obtain methyl 6-((piperidin-4-oxy)methyl)picolinate (260 mg, 98.1% yield). LC-MS m/z : 251 $[M+H]^+$.

4). Synthesis of methyl 6-((1-(tert-butoxycarbonyl)piperidin-4-yl)oxy)methyl)picolinate



[0369] To a mixture of methyl 6-((piperidin-4-oxy)methyl)picolinate (260 mg, 1.04 mmol) and triethylamine (315 mg, 3.12 mmol) in THF (3 mL) di-tert-butyl dicarbonate (453 mg, 2.08 mmol) was added at room temperature. The mixture was stirred at room temperature for 16 hours. The mixture was quenched with water (10 mL) at room temperature and then extracted with ethyl acetate (10 mL $\times$ 2). The combined organic layers were washed with brine (10 mL $\times$ 2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=1/1) to obtain methyl 6-((1-(tert-butoxycarbonyl)piperidin-4-yl)oxy)methyl)picolinate ester (255 mg, 70.1% yield). LC-MS m/z : 351 $[M+H]^+$.


5). Synthesis of tert-butyl 4-(6-(hydroxymethyl)pyridin-2-yl)methoxy)piperidine-1-carboxylate



[0370] To a solution of methyl 6-((1-(tert-butoxycarbonyl)piperidin-4-yl)oxy)methyl)picolinate (244 mg, 0.70 mmol) in tetrahydrofuran (3 mL) Aluminum lithium hydride (27 mg, 0.70 mmol) was added at 0° C. The mixture was stirred at room temperature for 16 hours. The mixture was quenched with water (10 mL) at room temperature and extracted with ethyl acetate (10 mL $\times$ 2). The combined organic layers were washed with brine (10 mL $\times$ 2), dried over anhydrous sodium sulfate, filtered, and concentrated to give residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=1/1)


to obtain tert-butyl 4-(6-(hydroxymethyl)pyridin-2-yl)methoxy)piperidine-1-carboxylate (107 mg, 39.6% yield). LC-MS m/z : 323 $[M+H]^+$.

6). Synthesis of tert-butyl 4-(6-(chloromethyl)pyridin-2-yl)methoxy)piperidine-1-carboxylate



[0371] To a solution of tert-butyl 4-(6-(hydroxymethyl)pyridin-2-yl)methoxy)piperidine-1-carboxylate (47 mg, 0.15 mmol) in dichloromethane (1 mL) triethylamine (30 mg, 0.30 mmol) and methanesulfonyl chloride (21 mg, 0.18 mmol) was added sequentially at 0° C. The mixture was stirred at room temperature for 16 hours. After completion, water (10 mL) was added to mixture and then extracted with DCM (20 mL $\times$ 2). The combined organic layers were washed with brine (10 mL $\times$ 2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue under reduced pressure. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=1/1) to obtain tert-butyl 4-(6-(chloromethyl)pyridin-2-yl)methoxy)piperidine-1-carboxylate (40 mg, 80.6% yield). LC-MS m/z : 341 $[M+H]^+$.


7). Synthesis of tert-butyl 4-(6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)methoxy)piperidine-1-carboxylate



[0372] To a mixture of tert-butyl 4-(6-(chloromethyl)pyridin-2-yl)methoxy)piperidine-1-carboxylate (40 mg, 0.12 mmol) and 3-fluoro-4-hydroxybenzonitrile (33 mg, 0.24 mmol) in N,N-dimethylformamide (1 mL) potassium carbonate (541 mg, 13.52 mmol) was added at room temperature. The mixture was stirred at 60° C. for 16 hours. The mixture was quenched with water (10 mL) at room tem-

perature and extracted with ethyl acetate (10 mL×2). The combined organic layers were washed with saturated ammonium chloride (20 mL×3), brine (10 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=4/1) to obtain tert-butyl 4-(6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)methoxy)piperidine-1-carboxylate (35 mg, 67.4% yield). LC-MS m/z: 442 [M+H]⁺.


8). Synthesis of 3-fluoro-4-((6-((piperidin-4-oxy)methyl)pyridin-2-yl)methoxy)benzonitrile



[0373] To a solution of tert-butyl 4-(6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)methoxy)piperidine-1-carboxylate (35 mg, 0.08 mmol) in dichloromethane (2 mL) trifluoroacetic acid (1 mL) was added. The mixture was stirred at room temperature for 16 hours. The reaction mixture was adjusted to pH=8-9 with NH₃/MeOH (7 M) and then concentrated to give a residue under reduced pressure. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to give 3-fluoro-4-((6-((piperidin-4-oxy)methyl)pyridin-2-yl)methoxy)benzonitrile (27 mg, 99% yield). LC-MS m/z: 342 [M+H]⁺.

Example 2-6. Synthesis of 4-((6-(azetidin-3-yloxy)pyridin-2-yloxy)methyl)-3-fluorobenzonitrile (Intermediate B-6)


1). Synthesis of tert-butyl 3-(6-fluoropyridin-2-yl)oxy)azetidine-1-carboxylate



[0374] To a solution of tert-butyl 3-hydroxyazetidine-1-carboxylate (2.00 g, 11.55 mmol) in dry THF (30 mL) sodium hydride (694 mg, 17.34 mmol, 60% w/w dispersion in mineral oil) was added at 0° C. The mixture was stirred at 0° C. for 30 min, then the 2,6-difluoropyridine (1.60 g, 13.90 mmol) was added slowly to the mixture at 0° C. The


mixture was warmed to 70° C. and stirred for 4 hours. After completion, it was quenched with water (20 mL) at room temperature and extracted with ethyl acetate (2×20 mL). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=10/1) to obtain tert-butyl 3-(6-fluoropyridin-2-yl)oxy)azetidine-1-carboxylate (3.00 g, 11.18 g mmol, 96.7% yield). LC-MS m/z: 269 [M+H]⁺.

2). Synthesis of tert-butyl 3-(6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)azetidine-1-carboxylate




[0375] To a solution of 3-fluoro-4-(hydroxymethyl)benzonitrile (1.00 g, 6.62 mmol) in dry THF (20 mL) sodium hydride (397 mg, 9.93 mmol, 60% w/w dispersion in mineral oil) was added at 0° C. The mixture was stirred at 0° C. for 30 minutes, then the tert-butyl 3-(6-fluoropyridin-2-yl)oxy)azetidine-1-carboxylate (1.33 g, 4.96 mmol) was added to above mixture. The mixture was stirred at 70° C. for 4 hours. The reaction mixture was quenched with water (20 mL) at room temperature and extracted with ethyl acetate (2×20 mL). The combined organic layers were washed with brine (2×20 mL), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=5/1) to give 3-(6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)azetidine-1-carboxylate tert-butyl ester (1.52 g, 3.81 mmol, 57.4% yield), LC-MS m/z: 400 [M+H]⁺.

3). Synthesis of 4-((6-(azetidin-3-yloxy)pyridin-2-yloxy)methyl)-3-fluorobenzonitrile




-continued



[0376] To a solution of tert-butyl 3-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)azetidine-1-carboxylate (900 mg, 2.25 mmol) in dichloromethane (10 mL) trifluoroacetic acid (2 mL) was added. The reaction mixture was stirred at room temperature for 16 hours. The reaction mixture was adjusted to pH=8-9 with NH_3/MeOH (7M) and then concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (DCM/MeOH=10/1) to obtain 4-((6-(azetidin-3-yloxy)pyridin-2-yloxy)methyl)-3-fluorobenzonitrile (650 mg, 2.17 mmol, 96.0% yield). LC-MS m/z : 300 $[\text{M}+\text{H}]^+$.


Example 2-7. Synthesis of 3-fluoro-4-((3-(piperidin-4-oxy)phenoxy)methyl)benzonitrile (Intermediate B-7)

1). Synthesis of 3-fluoro-4-((3-(hydroxyphenoxy)methyl)benzonitrile




[0377] The mixture of resorcinol (1.04 g, 9.52 mmol), potassium carbonate (1.31 g, 9.52 mmol) and 4-bromoethyl-3-fluorobenzonitrile (1.00 g, 4.76 mmol) in acetonitrile (10 mL) was stirred at 100° C. for 16 hours and cooled to room temperature. The mixture was diluted with water (20 mL) and then extracted with ethyl acetate (30 mL $\times$ 3). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum/ethyl acetate=4/1) to obtain 3-fluoro-4-((3-(hydroxyphenoxy)methyl)benzonitrile (514 mg, 44.0% yield). LC-MS m/z : 244 $[\text{M}+\text{H}]^+$.

2). Synthesis of tert-butyl 4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)piperidine-1-carboxylate




-continued



[0378] The mixture of 3-fluoro-4-((3-(hydroxyphenoxy)methyl)benzonitrile (200 mg, 0.82 mmol), tert-butyl 4-((methylsulfonyl)oxy)piperidine-1-carboxylate (919 mg, 3.29 mmol) and Cs_2CO_3 (805 mg, 2.47 mmol) in N,N-dimethylformamide (5 mL) was stirred at 100° C. for 16 hours. The mixture was concentrated in vacuo and purified by reverse-phase flash chromatography (Column: Spherical C18, 20-40 μm , 120 g; Mobile Phase A: water 10 mM NH_4OH in water, Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 40% B-80% B in 20 minutes; Detector: 254 nm). The fractions containing the desired product were collected at 65% B and concentrated in vacuo to give tert-butyl 4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)piperidine-1-carboxylate (180.0 mg, 79.8% yield). LC-MS m/z : 371 $[\text{M}+\text{H}]^+$.


3). Synthesis of 3-fluoro-4-((3-(piperidin-4-oxy)phenoxy)methyl)benzonitrile



[0379] To a solution of tert-butyl 4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)piperidine-1-carboxylate (120.0 mg, 0.282 mmol) in DCM (3 mL) trifluoroacetic acid (1 mL) was added dropwise. The reaction mixture was stirred at room temperature for 2 hours. The mixture was quenched with saturated sodium bicarbonate (30 mL), then extracted with DCM (2 $\times$ 10 mL). The combined organic layers were concentrated in vacuo and purified by silica gel column chromatography (DCM/MeOH=20/1) to obtain 3-fluoro-4-((3-(piperidin-4-oxy)phenoxy)methyl)benzonitrile (89.0 mg, 96.8% yield). LC-MS m/z : 327 $[\text{M}+\text{H}]^+$.


Example 2-8. Synthesis of 2-((4-Chloro-2-fluorobenzyl)oxy)-3-fluoro-6-(piperidin-4-oxy)pyridine (Intermediate B-8)

1). Synthesis of tert-butyl 4-(6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxypiperidine-1-carboxylate



[0380] The sodium hydride (264 mg, 0.56 mmol) was added to solution of tert-butyl 4-hydroxypiperidine-1-carboxylate (1.10 g, 5.50 mmol) in tetrahydrofuran (15 mL) at 0° C. The mixture was stirred at 0° C. for 30 minutes. Then 2-((4-chloro-2-fluorobenzyl)oxy)-3,6-difluoropyridine (100 mg, 0.28 mmol) was added to above mixture at 0° C. The mixture was warmed to 70° C. and stirred for 16 hours. The desired product was detected by LC-MS, then the reaction was quenched with methanol (10 mL) and extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (20 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue under reduced pressure. The residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate=5:1) to obtain tert-butyl 4-(6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridine-2-yl)oxypiperidine-1-carboxylate (447 mg, 53.7% yield). LC-MS m/z: 399 [M+H]⁺.

2). Synthesis of 2-((4-Chloro-2-fluorobenzyl)oxy)-3-fluoro-6-(piperidin-4-oxy)pyridine




[0381] To a solution of tert-butyl 4-(6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxypiperidine-1-carboxylate (447 mg, 0.98 mmol) in dichloromethane (10 mL) trifluoroacetic acid (3 mL) was added. The mixture was

stirred at room temperature for 1 hour. The desired product was detected by LC-MS, then the mixture was quenched with water (5 mL) and extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (20 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane:methanol=10:1) to give 2-((4-chloro-2-fluorobenzyl)oxy)-3-fluoro-6-(piperidin-4-oxy)pyridine (199 mg, 57.2% yield). LC-MS m/z: 355 [M+H]⁺.


Example 2-9. Synthesis of 3-fluoro-4-((3-fluoro-6-(piperidin-4-oxy)pyridin-2-yl)methoxy)benzonitrile (Intermediate B-9)

1). Synthesis of methyl 3,6-difluoropicolinate




[0382] To a solution of 3,6-difluoropicolinic acid (3.6 g, 22.64 mmol) in methanol (80 mL) thionyl chloride (4.0 g, 33.96 mmol) was added. The mixture was stirred at 60° C. for 2 hours. The reaction mixture was directly concentrated in vacuo and then purification by silica gel column chromatography (petroleum ether/ethyl acetate=4/1) to obtain methyl 3,6-difluoropicolinate (3.15 g, 80.4% yield). LC-MS m/z: 174 [M+H]⁺.

2). Synthesis of (3,6-difluoropyridin-2-yl)methanol




[0383] To a solution of methyl 3,6-difluoropicolinate (3.15 g, 18.21 mmol) in methanol (55 mL) sodium borohydride (2.07 g, 54.63 mmol) and lithium chloride (153 mg, 3.64 mmol) was added. The solution was stirred at 50° C. for 4 hours. The reaction mixture was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=2:1) to obtain (3,6-difluoropyridin-2-yl)methanol (2.0 g, 91.1% yield). LC-MS m/z: 146 [M+H]⁺.

3). Synthesis of methyl (3,6-difluoropyridin-2-yl) methanesulfonate




[0384] To a solution of (3,6-difluoropyridin-2-yl)methanol (300 mg, 2.07 mmol) in dry dichloromethane (20 mL) triethylamine (460 mg, 4.55 mmol) and methanesulfonyl chloride (358 mg, 3.11 mmol) was added slowly. The mixture was stirred at 0° C. for 1 hour and then stirred at room temperature for another 2 hours. The mixture was diluted with dichloromethane (50 mL) and washed with brine (25 mL). The organic layer was dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=2/1) to obtain methyl (3,6-difluoropyridin-2-yl) methanesulfonate (272 mg, 58.9% yield). LC-MS m/z : 224 $[M+H]^+$.

4). Synthesis of 4-((3,6-difluoropyridin-2-yl) methoxy)-3-fluorobenzonitrile




[0385] To a mixture of methyl (3,6-difluoropyridin-2-yl) methanesulfonate (272 mg, 1.22 mmol) and 3-fluoro-4-hydroxybenzonitrile (167 mg, 1.22 mmol) in *N,N*-dimethylformamide (5 mL) potassium carbonate (505 mg, 3.66 mmol) was added. The reaction mixture was stirred at 60° C. for 2 hours. The reaction mixture was quenched by adding water (15 mL), then was extracted with ethyl acetate (30 mL×3). The combined organic layers were washed with brine (30 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=2/1) to obtain 4-((3,6-difluoropyridin-2-yl)methoxy)-3-fluorobenzonitrile (270 mg, 83.6% yield). LC-MS m/z : 265 $[M+H]^+$.

5). Synthesis of tert-butyl 4-(6-((4-cyano-2-fluorophenoxy)methyl)-5-fluoropyridin-2-yl)oxypiperidine-1-carboxylate



[0386] To a solution of tert-butyl 4-hydroxypiperidine-1-carboxylate (246 mg, 1.22 mmol) in THF (30 mL) sodium hydride (61 mg, 1.53 mmol, 60% w/w dispersed in mineral oil) was added at 0° C. The mixture was stirred at 0° C. for 30 minutes, then 4-((3,6-difluoropyridin-2-yl)methoxy)-3-fluorobenzonitrile (270 mg, 1.02 mmol) was added to above mixture. The mixture was stirred at 70° C. for 2 hours. After this, the mixture was quenched with water (20 mL) and extracted with ethyl acetate (20 mL×2). The combined organic layers were washed with brine (10 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=2/1) to obtain tert-butyl 4-(6-((4-cyano-2-fluorophenoxy)methyl)-5-fluoropyridin-2-yl)oxypiperidine-1-carboxylate (200 mg, 36.9% yield). LC-MS m/z : 390 $[M+H]^+$.

6). Synthesis of 3-fluoro-4-((3-fluoro-6-(piperidin-4-oxy)pyridin-2-yl)methoxy)benzonitrile




[0387] To a solution of tert-butyl 4-(6-((4-cyano-2-fluorophenoxy)methyl)-5-fluoropyridin-2-yl)oxypiperidine-1-carboxylate (200 mg, 0.45 mmol) in dichloromethane (10 mL) trifluoroacetic acid (2 mL) was added. The mixture was stirred at room temperature for 16 hours. The reaction mixture was adjusted to pH=8-9 with NH_3/MeOH (7M) and then concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain 3-fluoro-4-((3-fluoro-6-(piperidin-4-oxy)pyridin-2-yl)methoxy)benzonitrile.

6-(piperidin-4-oxy)pyridin-2-yl)methoxy)benzonitrile (140 mg, 91.1% yield). LC-MS m/z : 346 $[M+H]^+$.


Example 2-10. Synthesis of 2-((4-chloro-2-fluorophenylthio)methyl)-6-(piperidin-4-oxy)pyridine (Intermediate B-10)

1) Synthesis of
1,2-bis(4-chloro-2-fluorophenyl)disulfane




[0388] The mixture of 4-chloro-1,2-difluorobenzene (1.00 g, 6.76 mmol) and sodium sulfide (1.05 g, 13.52 mmol) in dimethylsulfoxide (10 mL) was stirred at 80° C. for 2 hours. The mixture was quenched with water (100 mL) and then extracted with dichloromethane (50 mL×2). The aqueous phase was acidified with HCl to pH=2, and extracted with ethyl acetate (50 mL×3). The combined organic layers were washed with brine (100 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo to give 1,2-bis(4-chloro-2-fluorophenyl)disulfane (700 mg, 32.1% yield).

2). Synthesis of 4-chloro-2-fluorobenzenethiol




[0389] The mixture of 1,2-bis(4-chloro-2-fluorophenyl)disulfane (700 mg, 2.17 mmol) and zinc powder (423 mg, 6.51 mmol) in methanol (10 mL) and HCl (10 mL, 10% in H₂O) was as stirred at room temperature for 2 hours. The mixture was diluted with water (50 mL) and then extracted with dichloromethane (20 mL×3). The combined organic layer was concentrated in vacuo to obtain 4-chloro-2-fluorobenzenethiol (400 mg, 56.9% yield), LC-MS m/z : 161 $[M+H]^+$.

3). Synthesis of tert-butyl 4-(6-((4-chloro-2-fluorophenylthio)methyl)pyridin-2-yl)oxypiperidine-1-carboxylate



[0390] The mixture of 4-chloro-2-fluorobenzenethiol (400 mg, 2.47 mmol), potassium carbonate (682 mg, 4.94 mmol) and tert-butyl 4-((6-((methylsulfonyl)oxy)methyl)pyridine-2-yl)oxypiperidine-1-carboxylate (956 mg, 2.47 mmol) in N,N-dimethylformamide (20 mL) was stirred at room temperature for 16 hours. The resulting mixture was diluted with water (200 mL) and then extracted with ethyl acetate (80 mL×3). The combined organic layers were washed with saturated brine (100 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (ethyl acetate/petroleum ether=1/4) to obtain 4-(6-((4-chloro-2-fluorophenylthio)methyl)pyridin-2-yl) Oxypiperidine-1-carboxylate tert-butyl ester (1.00 g, 89.6% yield), LC-MS m/z : 453 $[M+H]^+$.


4). Synthesis of 2-((4-chloro-2-fluorophenylthio)methyl)-6-(piperidin-4-oxy)pyridine



[0391] To a solution of tert-butyl 4-(6-((4-chloro-2-fluorophenylthio)methyl)pyridin-2-yl)oxypiperidine-1-carboxylate (1.00 g, 2.21 mmol) in dichloromethane (10 mL) trifluoroacetic acid (3 mL) was added. The solution was stirred at room temperature for 2 hours. The mixture was quenched with saturated sodium bicarbonate (50 mL), then extracted with dichloromethane (20 mL×3). The combined organic layer was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain 2-((4-


chloro-2-fluorophenylthio)methyl)-6-(piperidin-4-oxy)pyridine (700 mg, 90.0% yield). LC-MS m/z : 353 $[M+H]^+$.

Example 2-11. Synthesis of 3-fluoro-4-((6-(piperidin-4-oxy)pyridin-2-yl)methylthio)benzonitrile (Intermediate B-11)




[0392] To a solution of 3,4-difluorobenzonitrile (10.00 g, 71.89 mmol) in dimethylsulfoxide (100 mL) was added sodium sulfide (11.22 g, 143.88 mmol). The resulting mixture was stirred at 80°C for 2 hours. After completion, the reaction was quenched with water (500 mL) and extracted with dichloromethane (200 mL $\times$ 2). The aqueous phase was acidified to pH=2 with aqueous hydrochloric acid, and extracted with ethyl acetate (200 mL $\times$ 3). The combined organic layers were washed with brine (400 mL $\times$ 3), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo to give 4,4'-dithiobis(3-fluorobenzonitrile) (5.70 g, 26.1% yield).

2). Synthesis of 3-fluoro-4-mercaptobenzonitrile




[0393] To a mixture of 4,4'-dithionylbis(3-fluorobenzonitrile) (2.00 g, 6.57 mmol) in methanol (10 mL) and hydrochloric acid (10% in water, 10 mL) zinc powder (1.28 g, 19.58 mmol) was added and the mixture was stirred at room temperature for 2 hours. After this, the mixture was diluted with water (200 mL) and then extracted with dichloromethane (80 mL $\times$ 3). The organic layer was concentrated to give 3-fluoro-4-mercaptobenzonitrile (1.20 g, 59.6% yield).

3). Synthesis of tert-butyl 4-((6-((4-cyano-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidine-1-carboxylate



[0394] To a mixture of 3-fluoro-4-mercaptobenzonitrile (474 mg, 3.10 mmol) and tert-butyl 4-((6-((methylsulfonyl)methyl)pyridin-2-yl)oxy)piperidine-1-carboxylate in N,N-dimethylformamide (20 mL) potassium carbonate (856 mg, 6.20 mmol) was added. The mixture was stirred at room temperature for 16 hours. The mixture was diluted with water (200 mL) and then extracted with ethyl acetate (80 mL $\times$ 3). The combined organic layers were washed with brine (100 mL $\times$ 3), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. The residue was purified by silica gel column chromatography (ethyl acetate/petroleum ether=1/4) to obtain 4-((6-((4-cyano-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidine-1-carboxylate tert-butyl ester (600 mg, 43.7% yield). LC-MS m/z : 444 $[M+H]^+$.

4). Synthesis of 3-fluoro-4-((6-(piperidin-4-oxy)pyridin-2-yl)methylthio)benzonitrile




[0395] To a solution of tert-butyl 4-((6-((4-cyano-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidine-1-carboxylate (500 mg, 1.13 mmol) in dichloromethane (5 mL) 2,2,2-trifluoroacetic acid (1 mL) was added. The mixture was stirred at room temperature for 2 hours. After completion, the reaction mixture was quenched with saturated sodium bicarbonate (10 mL) and extracted with dichloromethane (5 mL $\times$ 3). The combined layer was concentrated and then purified by silica gel column chromatography (dichloromethane/methanol=20/1) to give 3-fluoro-4-((6-

(piperidin-4-oxy)pyridin-2-yl)methylthio)benzonitrile (334 mg, 86.4% yield). LC-MS m/z : 344 $[M+H]^+$.


Example 2-12. Synthesis of 3-fluoro-4-((3-(piperidin-4-oxy)benzyl)oxy)benzonitrile (Intermediate B-12)

1). Synthesis of tert-butyl 4-(3-(methoxycarbonyl)phenoxy)piperidine-1-carboxylate



[0396] To a mixture of methyl 3-hydroxybenzoate (3.00 g, 19.73 mmol), tert-butyl 4-hydroxypiperidine-1-carboxylate (3.97 g, 19.73 mmol) and triphenylphosphine (7.76 g, 29.60 mmol) in tetrahydrofuran (30 mL) diisopropyl azodicarboxylate (5.98 g, 29.60 mmol) was added at 0° C. The mixture was stirred at room temperature for 3 hours. After completion, water (50 mL) was added to quench the reaction, and extracted with ethyl acetate (50 mL $\times$ 2). The combined organic layers were washed with brine (20 mL $\times$ 2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=5/1) to give tert-butyl 4-(3-(methoxycarbonyl)phenoxy)piperidine-1-carboxylate (2.44 g, 36.9% yield). LC-MS m/z : 358 $[M+H]^+$.


2). Synthesis of tert-butyl 4-(3-(hydroxymethyl)phenoxy)piperidine-1-carboxylate



[0397] To a solution of tert-butyl 4-(3-(methoxycarbonyl)phenoxy)piperidine-1-carboxylate (1.40 g, 4.18 mmol) in tetrahydrofuran (15 mL) lithium aluminum hydride (238 mg, 6.27 mmol) was slowly added at 0° C. The resulting mixture was stirred at room temperature for 3 hours. After completion, water (5 mL) was added to quench the reaction, then extracted with ethyl acetate (20 mL $\times$ 2). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The


residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=5/1) to obtain tert-butyl 4-(3-(hydroxymethyl)phenoxy)piperidine-1-carboxylate (400 mg, 31.2% yield).

3). Synthesis of tert-butyl 4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidine-1-carboxylate



[0398] To a mixture of 3-fluoro-4-hydroxybenzonitrile (178 mg, 1.30 mmol), triphenylphosphine (409 mg, 1.56 mmol) and tert-butyl 4-(3-(hydroxymethyl)phenoxy)piperidine-1-carboxylate (400 mg, 1.30 mmol) in tetrahydrofuran (15 mL) azodicarboxylate (315 mg, 1.56 mmol) was added diisopropyl at 0° C. The mixture was stirred at room temperature for 3 hours. After completion, the mixture was diluted with water (20 mL) and then extracted with ethyl acetate (30 mL $\times$ 2). The combined organic layers were washed with brine (20 mL $\times$ 2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain tert-butyl 4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidine-1-carboxylate (312 mg, 56.3% yield). LC-MS m/z : 371 $[M+H]^+$.

4). Synthesis of 3-fluoro-4-((3-(piperidin-4-oxy)benzyl)oxy)benzonitrile




[0399] To a solution of tert-butyl 4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidine-1-carboxylate (312 mg, 0.73 mmol) in dichloromethane (5 mL) trifluoroacetic acid (1 mL) was added at room temperature. The mixture was stirred at room temperature for 3 hours. After completion, the mixture was diluted with water (20 mL), and then extracted with ethyl acetate (20 mL $\times$ 2). The combined organic layers were washed with brine (20 mL $\times$ 2), dried

over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to give 3-fluoro-4-((3-(piperidin-4-oxy)benzyl)oxy)benzonitrile (230 mg, 96.6% yield). LC-MS m/z : 327 $[M+H]^+$.


Example 2-13. Synthesis of 4-(3-((4-chloro-2-fluorophenoxy)methyl)phenoxy)piperidine (Intermediate B-13)

1). Synthesis of tert-butyl 4-(3-((4-chloro-2-fluorophenoxy)methyl)phenoxy)piperidine-1-carboxylate



[0400] To a mixture of 4-chloro-2-fluorophenol (144 mg, 0.98 mmol), triphenylphosphine (308 mg, 1.18 mmol) and tert-butyl 4-(3-(hydroxymethyl)phenoxy)piperidine-1-formate (300 mg, 0.98 mmol) in tetrahydrofuran (8 mL) diisopropyl azodicarboxylate (238 mg, 1.18 mmol) was added slowly at 0° C. The mixture was stirred at room temperature for 3 hours. After completion, water (20 mL) was added to quench the reaction. The mixture was extracted with ethyl acetate (20 mL×2). The combined organic layers were washed with brine (20 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=5/1) to obtain tert-butyl 4-(3-((4-chloro-2-fluorophenoxy)methyl)phenoxy)piperidine-1-carboxylate (381 mg, 89.2% yield).

2). Synthesis of 4-(3-((4-chloro-2-fluorophenoxy)methyl)phenoxy)piperidine




[0401] To a solution of tert-butyl 4-(3-((4-chloro-2-fluorophenoxy)methyl)phenoxy)piperidine-1-carboxylate (381

mg, 0.88 mmol) in dichloromethane (5 mL) trifluoroacetic acid (1 mL) was added at room temperature. The resulting mixture was stirred at room temperature for 1 hour. After completion, the mixture was adjusted to pH=7-8 with $NH_3/MeOH$ (7 M). The reaction solution was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain 4-(3-((4-chloro-2-fluorophenoxy)methyl)phenoxy)piperidine (271 mg, 91.6% yield). LC-MS m/z : 336 $[M+H]^+$.


Example 2-14. Synthesis of methyl 2-(4-((6-((methylsulfonyl)oxy)methyl)pyridin-2-yl)oxy)phenyl)acetate (Intermediate B-14)

1). Synthesis of methyl 2-(4-((6-bromopyridin-2-yl)oxy)phenyl)acetate



[0402] To a solution of methyl 2-(4-hydroxyphenyl)acetate (2.00 g, 12.05 mmol) in acetonitrile (20 mL) added 2-bromo-6-fluoropyridine (2.53 g, 14.46 mmol) and cesium carbonate (5.89 g, 18.07 mmol) was at room temperature. The mixture was stirred at 90° C. for 16 hours. After completion, it was diluted by water (5 mL) and then extracted with ethyl acetate (40 mL×3). The combined organic layers were washed with brine (30 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=10/1) to give methyl 2-(4-((6-bromopyridin-2-yl)oxy)phenyl)acetate (3.17 g: 81.7% yield). LC-MS m/z : 322, 324 $[M+H]^+$.


2). Synthesis of 2-(4-((6-bromopyridin-2-yl)oxy)phenyl)acetic acid



[0403] To a mixture of methyl 2-(4-((6-bromopyridin-2-yl)oxy)phenyl)acetate (3.17 g, 9.84 mmol) in tetrahydrofuran (20 mL) and water (15 mL) lithium hydroxide monohydrate (4.13 g, 98.40 mmol) was added at room temperature. The mixture was stirred at room temperature


for 5 hours. After completion, the mixture was diluted with water (5 mL), and then extracted with ethyl acetate (40 mL $\times$ 3). The combined organic layers were washed with brine (30 mL $\times$ 2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to give 2-(4-((6-bromopyridin-2-yl)oxy)phenyl)acetic acid (1.8 g, 59.4% yield). LC-MS m/z : 308, 310 $[M+H]^+$.

3). Synthesis of 2-(4-((6-(methylcarboxylate)pyridin-2-yl)oxy)phenyl)acetic acid



[0404] The mixture of 2-(4-((6-bromopyridin-2-yl)oxy)phenyl)acetic acid (900 mg, 2.92 mmol), 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (213 mg, 0.29 mmol) and potassium acetate (859 mg, 8.76 mmol) in methanol (10 mL) was degassed with CO three times at room temperature. The mixture was warmed up to 90° C. and stirred for 16 hours under CO (56 Psi) atmosphere. After completion, water (5 mL) was added to quench the reaction. The mixture was extracted with ethyl acetate (40 mL $\times$ 3). The combined organic layers were washed with brine (30 mL $\times$ 2), dried over anhydrous sodium sulfate, filtered, and concentrated to give residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to give 2-(4-((6-(methylcarboxylate)pyridin-2-yl)oxy)phenyl)acetic acid (385 mg, 45.9% yield). LC-MS m/z : 288 $[M+H]^+$.


4). Synthesis of 2-(4-((6-(hydroxymethyl)pyridin-2-yl)oxy)phenyl)acetic acid



[0405] To a solution of 2-(4-((6-(methylcarboxylate)pyridin-2-yl)oxy)phenyl)acetic acid (385 mg, 1.34 mmol) in tetrahydrofuran (5 mL) lithium aluminum hydride (76 mg, 2.01 mmol) was added at 0° C. The mixture was stirred at room temperature for an additional 2 hours. After completion, the mixture was adjusted to pH=5-6 with hydrochloric acid (1M), and then extracted with ethyl acetate (20 mL $\times$ 3). The combined organic layer were washed with brine (20 mL), dried over anhydrous sodium sulfate, filtered, and


concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain 2-(4-((6-(hydroxymethyl)pyridin-2-yl)oxy)phenyl)acetic acid (118 mg, 33.9% yield). LC-MS m/z : 260 $[M+H]^+$.

5). Synthesis of methyl 2-(4-((6-(hydroxymethyl)pyridin-2-yl)oxy)phenyl)acetate



[0406] To a solution of 2-(4-((6-(Hydroxymethyl)pyridin-2-yl)oxy)phenyl)acetic acid (118 mg, 0.45 mmol) in methanol (3 mL) p-toluenesulfonic acid (12 mg, 0.32 mmol) was added at room temperature. The resulting mixture was stirred at 60° C. for 1 hour. When completion, the reaction mixture was quenched by adding water (5 mL) and then was extracted with ethyl acetate (20 mL $\times$ 3). The combined organic layers were washed with brine (20 mL $\times$ 2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=15/1) to obtain methyl 2-(4-((6-(hydroxymethyl)pyridin-2-yl)oxy)phenyl)acetate (89 mg, 72.4% yield). LC-MS m/z : 274 $[M+H]^+$.






6). Synthesis of methyl 2-(4-((6-((methylsulfonyl)oxy)methyl)pyridin-2-yl)oxy)phenyl)acetate








[0407] To a mixture of 2-(4-((6-(hydroxymethyl)pyridin-2-yl)oxy)phenyl)acetate (89 mg, 0.33 mmol) and triethylamine (74 mg, 0.73 mmol) in dichloromethane (3 mL), methanesulfonyl chloride was added slowly at 0° C. The reaction mixture was stirred at room temperature for 3 hours. After completion, the reaction was quenched by adding water (5 mL) and extracted with ethyl acetate (20 mL $\times$ 3). The combined organic layers were washed with brine (20 mL $\times$ 2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=4/1) to obtain 2-(4-((6-((methylsulfonyl)oxy)methyl)pyridin-2-yl)oxy)phenyl)acetic acid (56 mg, 48.3% yield). LC-MS m/z : 352 $[M+H]^+$.

Example 2-15. Synthesis of Other B Intermediates

[0408] The following intermediates can be obtained with similar to the synthetic steps of each intermediate from example 2-1 to Example 2-14

Number	Structure	Name	MS m/z: [M + H] ⁺
IntermediateB-15		3-Fluoro-4-(((3-fluoro-6-(piperidin-4-oxy)pyridin-2-yl)oxy)methyl)benzonitrile	346
IntermediateB-16		3-Fluoro-4-((2-fluoro-5-(piperidin-4-oxy)phenoxy)methyl)benzonitrile	345
IntermediateB-17		2-((4-Chloro-2-fluorobenzyl)oxy)-6-((piperidin-4-oxy)methyl)pyridine	351
IntermediateB-18		4-(3-((4-Chloro-2-fluorobenzyl)oxy)-4-fluorophenoxy)piperidine	354
IntermediateB-19		2-((4-Chloro-2-fluorophenoxy)methyl)-3-fluoro-6-(piperidin-4-oxy)pyridine	355






-continued

Number	Structure	Name	MS
			m/z: [M + H] ⁺
IntermediateB-20		2-((4-Chloro-2-fluorophenoxy)methyl)-6-(piperidin-4-oxy)pyridine	337
IntermediateB-21		4-(3-((4-Chloro-2-fluorophenoxy)methyl)-4-fluorophenoxy)piperidine	354
IntermediateB-22		3-Fluoro-4-((2-fluoro-5-(piperidin-4-oxy)benzyl)oxy)-benzonitrile	345
IntermediateB-23		4-(((3-azabicyclo[3.2.1]octan-8-yl)oxy)pyridin-2-yl)methoxy)-3-fluorobenzonitrile	354
IntermediateB-24		8-(((4-Chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)-3-azabicyclo[3.2.1]octane	363





-continued

Number	Structure	Name	MS m/z: [M + H] ⁺
IntermediateB-25		3-Fluoro-4-((6-((3-methylpiperidin-4-yl)oxy)pyridin-2-yl)methoxy)benzonitrile	342
IntermediateB-26		2-((4-Chloro-2-fluorophenoxy)methyl)-6-((3-methylpiperidin-4-yl)oxy)pyridine	351
IntermediateB-27		3-Fluoro-4-((6-((3-fluoropiperidin-4-yl)oxy)pyridin-2-yl)methoxy)benzonitrile	346
IntermediateB-28		2-((4-Chloro-2-fluorophenoxy)methyl)-6-((3-fluoropiperidin-4-yl)oxy)pyridine	355
IntermediateB-29		3-Fluoro-4-((3-(piperidin-4-oxy)pyridin-2-yl)methoxy)benzonitrile	328
IntermediateB-30		2-((4-Chloro-2-fluorophenoxy)methyl)-3-(piperidin-4-oxy)pyridine	337

-continued


Number	Structure	Name	MS
			m/z: [M + H] ⁺
IntermediateB-31		3-Fluoro-4-((2-(piperidin-4-oxy)pyridin-3-yl)methoxy)benzonitrile	328
IntermediateB-32		3-((4-Chloro-2-fluorophenoxy)methyl)-2-(piperidin-4-oxy)pyridine	337
IntermediateB-33		3-Fluoro-4-((5-(piperidin-4-oxy)pyridin-3-yl)methoxy)benzonitrile	328
IntermediateB-34		3-((4-Chloro-2-fluorophenoxy)methyl)-5-(piperidin-4-oxy)pyridine	337
IntermediateB-35		4-(3-((2-Fluoro-4-(1H-imidazol-1-yl)phenoxy)methyl)phenoxy)piperidine	368

-continued

Number	Structure	Name	MS
			m/z: [M + H] ⁺
IntermediateB-36		2-((2-Fluoro-4-(1H-imidazol-1-yl)phenoxy)methyl)-6-(piperidin-4-oxy)pyridine	369
IntermediateB-37		2-((2,4-Difluoro-phenoxy)methyl)-6-(piperidin-4-oxy)pyridine	321
IntermediateB-38		6-((6-(Piperidin-4-oxy)pyridin-2-yl)methoxy)-nicotinonitrile	311
IntermediateB-39		5-Chloro-2-((6-(piperidin-4-oxy)-pyridin-2-yl)methoxy)-pyridine	320

Example 3. Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-LH-benzo[d]imidazole-6-carboxylic acid (Compound 1)


1). Synthesis of methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamide)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate




[0409] To a mixture of 3-fluoro-4-((6-((piperidin-4-oxy)pyridin-2-yl)methoxy)benzonitrile (100 mg, 0.31 mmol) and methyl 4-(2-chloroacetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (109 mg, 0.31 mmol) in N,N-dimethylformamide (2 mL) potassium carbonate (86 mg, 0.62 mmol) was added. The mixture was stirred at 60° C. for 3 hours. The resulting mixture was poured into brine (20 mL) and extracted with dichloromethane (5 mL×2). The combined organic layer was concentrated in vacuo and purified by reverse-phase flash chromatography under the following conditions (Column: spherical C18, 20-40 m, 120 g; Mobile Phase A: 0.1% NH₄OH in Water; Mobile Phase B:

Acetonitrile; Flow rate: 80 mL/min; Gradient: 0% B-95% B over 30 minutes; Detector: 254 nm). The fractions containing desired product were collected at 72% B and concentrated in vacuo to afford methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamide)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (40 mg, 20.1% yield). LC-MS m/z: 642.5 [M+H]⁺.

2). Synthesis of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate




-continued




[0410] To a solution of methyl 4-(2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (40 mg, 0.06 mmol) in toluene (1 mL) acetic acid (0.1 mL) was added dropwise. The reaction mixture was stirred at 110° C. for 3 hours. After completion, the mixture was poured into brine (10 mL) and extracted with dichloromethane (5 mL). The combined organic layer was concentrated in vacuo to give methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (30 mg, 80.3% yield). LC-MS m/z : 624 $[M+H]^+$.

noxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (30 mg, 80.3% yield). LC-MS m/z : 624 $[M+H]^+$.

3). Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid






[0411] To a mixture of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)methoxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (30 mg, 0.05 mmol) in THF (1 mL) and water (1 mL) lithium hydroxide (2.4 mg, 0.10 mmol) was added. The mixture was stirred at room temperature for 16 hours. The reaction mixture was directly concentrated in vacuo to remove the solvent, and then purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19x 250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)methoxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (7.67 mg, 25.2% yield). LC-MS *m/z*: 610 [M+H]⁺.

[0412] ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.04 (s, 1H), 7.89-7.80 (m, 2H), 7.74-7.60 (m, 4H), 7.43 (m, 1H), 7.04 (d, *J*=7.2 Hz, 1H), 6.72 (d, *J*=8.0 Hz, 1H), 6.39 (s, 1H), 5.69 (s, 2H), 5.30 (s, 2H), 4.88-4.83 (m, 1H), 4.01-4.00 (m, 2H), 3.79 (s, 2H), 2.68-2.65 (m, 2H), 2.25-2.20 (m, 2H), 1.81-1.76 (m, 2H), 1.47-1.44 (m, 2H), 1.16 (t, *J*=7.2 Hz, 3H).

Example 4. Synthesis of (S)-2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)methoxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound S22)

1). Synthesis of methyl (S)-2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)methoxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate




[0413] The mixture of methyl (S)-2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (150 mg, 0.51 mmol), 3-fluoro-4-((6-((piperidin-4-yloxy)methyl)pyridin-2-yloxy)methyl)benzonitrile (173 mg, 0.51 mmol) and potassium carbonate (140 mg, 1.02 mmol) in N,N-dimethylformamide (2 mL) was stirred at 60° C. for 3 hours. The mixture was diluted with water (10 mL) and extracted with ethyl acetate (20 mL×2). The combined organic layers were washed successively with saturated ammonium chloride (20 mL×3) and brine (10 mL×2) in turn, dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to give methyl (S)-2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)methoxy)-piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 32.9% yield). LC-MS m/z: 600 [M+H]⁺.

2). Synthesis of (S)-2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)methoxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid


[0414] To a mixture of methyl (S)-2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)methoxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 0.17 mmol) in water (4 mL) and THF (4 mL) lithium hydroxide monohydrate (12 mg, 0.51 mmol) was added. The reaction mixture was stirred at 40° C. for 16 hours. The reaction mixture was adjusted to pH=5-6 with formic acid. The solvent was removed in vacuo and the residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μm 19×250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain (S)-2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridine-2-yl)methoxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound S22) (19.26 mg, 19.7% yield). LC-MS m/z: 586 [M+H]⁺.

[0415] ¹H NMR (400 MHz, DMSO-d₆): 8.08-8.06 (m, 1H), 7.90 (d, J=10.0 Hz, 1H), 7.77-7.70 (m, 4H), 7.46-7.44 (m, 1H), 7.07 (d, J=7.2 Hz, 1H), 6.80 (d, J=8.4 Hz, 1H), 5.44 (s, 2H), 5.10-5.05 (m, 1H), 4.73-4.68 (m, 1H), 4.59-4.51 (m, 1H), 4.49-4.46 (m, 3H), 4.39-4.34 (m, 1H), 3.86 (d, J=13.2 Hz, 1H), 3.73-3.69 (m, 1H), 3.47-3.41 (m, 1H), 2.76-2.64 (m, 3H), 2.49-2.41 (m, 1H), 2.23-2.17 (m, 2H), 1.89-1.85 (m, 2H), 1.56-1.49 (m, 2H).



Example 5. Synthesis of (S)-2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)methyl)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound S1)


1). Synthesis of methyl (S)-2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)methyl)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate




[0416] The mixture of 3-fluoro-4-((6-(piperidin-4-ylmethoxy)pyridin-2-yl)oxy)methylbenzonitrile (100 mg, 0.29 mmol), methyl (S)-2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (85 mg, 0.29 mmol) and potassium carbonate (80 mg, 0.58 mmol) in N,N-dimethylformamide (2 mL) was heated to $50^\circ C$ and stirred for 3 hours. After completion, the resulting mixture was diluted with water (20 mL) and extracted with ethyl acetate (10 mL $\times$ 2). The combined organic layers were washed with brine (30 mL), dried over sodium sulfate, filtered, and concentrated to give a residue in vacuo. The

residue was purified by silica gel column chromatography (DCM/MeOH=20/1) to give methyl (S)-2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridine-2-yl)oxy)methyl)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 0.17 mmol, 57.6% yield). LC-MS m/z : 600 $[M+H]^+$.

2). Synthesis of (S)-2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)methyl)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid






[0417] To a mixture of methyl (S)-2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)methyl)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 0.17 mmol) in water (1.0 mL) and THF (1.0 mL) lithium hydroxide (8 mg, 0.33 mmol) was added. The reaction mixture was stirred at room temperature for 16 hours. The solvent was removed in vacuo and the residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelengths: 254 nm/214 nm; Flow rate: mL/min) to give (S)-2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)yl)methyl)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (15.48 mg, 15.6% yield). LC-MS m/z: 586 [M+H]⁺.

[0418] ¹H NMR (400 MHz, DMSO-d₆): δ 8.28 (s, 1H), 8.24 (s, 1H), 7.92-7.87 (m, 1H), 7.80 (d, J=8.4 Hz, 1H), 7.71

(dd, J=7.6, 1.2 Hz, 1H), 7.66 (d, J=6.4 Hz, 1H), 7.64-7.60 (m, 2H), 6.46 (d, J=8.0 Hz, 1H), 6.37 (d, J=8.0 Hz, 1H), 5.45 (s, 2H), 5.11-5.03 (m, 1H), 4.80-4.72 (m, 1H), 4.66-4.58 (m, 1H), 4.52-4.45 (m, 1H), 4.40-4.33 (m, 1H), 4.00 (d, J=6.0 Hz, 2H), 3.89 (d, J=13.6 Hz, 1H), 3.73 (d, J=13.6 Hz, 1H), 2.90 (d, J=10.4 Hz, 1H), 2.80-2.70 (m, 1H), 2.72-2.64 (m, 1H), 2.45-2.37 (m, 1H), 2.13-1.95 (m, 2H), 1.70-1.60 (m, 3H), 1.27-1.13 (m, 2H).


Example 6. Synthesis of (S)-2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound S2)

1). Synthesis of methyl (S)-2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate




[0419] The mixture of methyl (S)-2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (200 mg, 0.68 mmol), 3-fluoro-4-((6-(piperidin-4-oxy)pyridin-2-yl)oxy)methyl)benzonitrile (245 mg, 0.75 mmol) and potassium carbonate (188 mg, 1.36 mmol) in N,N-dimethylformamide (2 mL) was stirred at 60° C. for 3 hours. The resulting mixture was diluted with water (10 mL) and extracted with ethyl acetate (20 mL×2). The combined organic layers were washed successively with saturated ammonium chloride (20 mL×3) and brine (10 mL×2), then dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (DCM/MeOH=20/1) to give methyl (S)-2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (205 mg, 0.35 mmol, 51.5% yield). LC-MS m/z: 586 [M+H]⁺.

2). Synthesis of (S)-2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid



-continued





[0420] To a mixture of methyl (S)-2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (150 mg, 0.26 mmol) in water (1 mL) and THF (1 mL) lithium hydroxide (62 mg, 2.59 mmol) was added. The reaction mixture was stirred at room temperature for 16 hours. The reaction mixture was adjusted to pH=5-6 with formic acid. Then the solvent was removed by concentration in vacuo to give a residue. The residue was purified prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μm 19×250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength 254 nm/214 nm; Flow rate: 20 mL/min) to give (S)-2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (38.01 mg, 0.07 mmol, 26.0% yield). LC-MS m/z: 572 [M+H]⁺.

[0421] ¹H NMR (400 MHz, DMSO-d₆): δ 8.26 (s, 1H), 7.93-7.90 (m, 1H), 7.80 (d, J=8.0 Hz, 1H), 7.72-7.69 (m, 1H), 7.65-7.61 (m, 3H), 6.46 (d, J=7.6 Hz, 1H), 6.36 (d, J=8.0 Hz, 1H), 5.45 (s, 2H), 5.09-5.07 (m, 1H), 4.81-4.75 (m, 2H), 4.66-4.61 (m, 1H), 4.52-4.47 (m, 1H), 4.40-4.36 (m, 1H), 3.94 (d, J=13.6 Hz, 1H), 3.78 (d, J=13.6 Hz, 1H), 2.78-2.67 (m, 3H), 2.45-2.41 (m, 1H), 2.31-2.27 (m, 2H), 1.86-1.83 (m, 2H), 1.58-1.53 (m, 2H).

Example 7. Synthesis of (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)methoxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C1)

1): Synthesis of methyl (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)methoxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate






[0422] The mixture of 3-fluoro-4-((6-((piperidin-4-oxy)methyl)pyridin-2-yl)methoxy)benzonitrile (27 mg, 0.08 mmol), methyl (S)-2-(chloromethyl)-1-(oxetane-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (24 mg, 0.08 mmol) and potassium carbonate (22 mg, 0.16 mmol) in N,N-dimethylformamide (1 mL) was stirred at 60° C. for 3 hours. The resulting mixture was diluted with water (10 mL) and extracted with ethyl acetate (20 mL×2). The combined organic layers were combined, washed with saturated ammonium chloride (20 mL×3) and brine (10 mL×2) in turn, dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by

silica gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)methoxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (23 mg, 47.0% yield). LC-MS m/z : 600 $[M+H]^+$.

2). Synthesis of (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)methoxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid




[0423] To a mixture of methyl (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)methoxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (23 mg, 0.04 mmol) in water (1 mL) and THF (1 mL) lithium hydroxide (10 mg, 0.40 mmol) was added. The reaction mixture was stirred at room temperature for 16 hours. The reaction mixture was adjusted to pH=5-6 with formic acid. The solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to give (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)methoxy)piperidine-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (14.73 mg, 65.5% yield). LC-MS m/z: 586 [M+H]⁺.

[0424] ¹H NMR (400 MHz, DMSO-d₆) δ 8.26 (s, 1H), 8.22 (s, 1H), 7.91-7.86 (m, 2H), 7.82-7.79 (m, 1H), 7.69 (d, J=8.8 Hz, 1H), 7.63 (d, J=8.4 Hz, 1H), 7.46-7.42 (m, 3H), 5.35 (s, 2H), 5.10-5.08 (m, 1H), 4.81-4.76 (m, 1H), 4.66-4.59 (m, 3H), 4.52-4.47 (m, 1H), 4.40-4.35 (m, 1H), 3.92 (d, J=13.6 Hz, 1H), 3.75 (d, J=13.6 Hz, 1H), 3.51-3.46 (m, 1H), 2.79-2.67 (m, 3H), 2.46-2.41 (m, 1H), 2.27-2.24 (m, 2H), 1.91-1.89 (m, 2H), 1.56-1.51 (m, 2H).

Example 8. Synthesis of (S)-2-((3-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)azetidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound S3)


1). Synthesis of methyl (S)-2-((3-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)azetidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate



[0425] The mixture of methyl (S)-2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (150 mg, 0.51 mmol), 4-((6-(azetidin-3-yloxy)pyridin-2-yloxy)methyl)-3-fluorobenzonitrile (153 mg, 0.51 mmol) and potassium carbonate (211 mg, 1.530 mmol) in N,N-dimethylformamide (2 mL) was warmed to 60° C. for 4

hours. The resulting mixture was diluted with water (10 mL) and extracted with ethyl acetate (2 $\times$ 20 mL). The combined organic layers was washed sequentially with saturated ammonium chloride (30 mL) and brine (2 $\times$ 20 mL), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (DCM/MeOH=20/1) to give methyl (S)-2-((3-((6-((4-cyano-2-fluorobenzyl)oxy)pyridine-2-yl)oxy)azetidine-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (180.0 mg, 0.32 mmol, 63.3% yield). LC-MS m/z: 558 [M+H]⁺.

2). Synthesis of (S)-2-((3-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)azetidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid




[0426] To a mixture of methyl (S)-2-((3-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)azetidine-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (90 mg, 0.16 mmol) in water (4 mL) and THF (4 mL) lithium hydroxide monohydrate (68 mg, 1.63 mmol) was added. The reaction mixture was stirred at room temperature for 16 hours. The reaction mixture was adjusted to pH=5-6 with formic acid. The solvent was removed to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% NH₄OH in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain (S)-2-((3-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)azetidine-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (compound S3) (10.23 mg, 11.7% yield). LC-MS m/z: 544 [M+H]⁺.

[0427] ¹H NMR (400 MHz, DMSO-d₆) δ 12.56 (brs, 1H), 8.24 (d, J=0.8 Hz, 1H), 7.86-7.78 (m, 2H), 7.71-7.63 (m, 4H), 6.49 (d, J=7.6 Hz, 1H), 6.42 (d, J=7.6 Hz, 1H), 5.41 (s, 2H), 5.08-5.02 (m, 2H), 4.72 (dd, J=15.4, 7.2 Hz, 1H), 4.59 (dd, J=15.4, 2.8 Hz, 1H), 4.47-4.43 (m, 1H), 4.33-4.28 (m, 1H), 4.05 (d, J=13.6 Hz, 1H), 3.95 (d, J=13.6 Hz, 1H), 3.77-3.69 (m, 2H), 3.20-3.16 (m, 2H), 2.70-2.66 (m, 1H), 2.40-2.35 (m, 1H).


Example 9. Synthesis of (S)-2-((4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (compound S5)


1). Synthesis of methyl (S)-2-((4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate



[0428] To a mixture of 3-fluoro-4-((6-(piperidin-4-oxymethyl)benzonitrile (89.0 mg, 0.273 mmol) and methyl (S)-2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (80.3 mg, 0.273 mmol) in N,N-dimethylformamide (4 mL) potassium carbonate (113.0 mg, 0.819 mmol) was added. The mixture was heated to 60° C. for 3 hours. The mixture was diluted with water (5 mL) and extracted with EA (3×20 mL). The combined organic layers were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated to give a

residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to give methyl (S)-2-((4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy))piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (94 mg, 59.0% yield). LC-MS m/z: 585 [M+H]⁺. 2). (S)-2-((4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid






[0429] To a mixture of methyl (S)-2-((4-(3-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)methyl)piperidin-1-yl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (94 mg, 0.16 mmol) in THE (4 mL) and H₂O (4 mL) LiOH·H₂O (68 mg, 1.60 mmol) was added. The mixture was warmed to 40° C. and stirred for 16 hours. The mixture was adjusted to pH=5-6 with hydrochloric acid (1 M) and extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (10 mL), dried over anhydrous Na₂SO₄, filtered, and concentrated to give a residue in vacuo. The residue was purified by reverse phase column chromatography (Spherical C18 column, 20-40 μm, 40 g; Mobile Phase A: 10 mM in NH₃H₂O water; Mobile

3.90 (d, J=13.6 Hz, 1H), 3.76 (d, J=13.5 Hz, 1H), 2.78-2.68 (m, 3H), 2.50-2.33 (m, 3H), 1.93-1.91 (m, 2H), 1.62-1.58 (m, 2H).

Example 10. Synthesis of 2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)methyl)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound S4)

1). Synthesis of methyl 4-(2-(4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)methyl)piperidin-1-yl)acetamido)-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate




Phase B: acetonitrile; Flow rate: 40 mL/min; Gradient: 20% B-50% B in 20 minutes; Detector: 254 nm).

[0430] The mobile phase containing the desired product was collected under 32% B and then concentrated in vacuo to give (S)-2-((4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)yl)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (26.27 mg, 28.6% yield). LC-MS m/z: 571 [M+H]⁺

[0431] ¹H NMR (400 MHz, DMSO-d₆) δ 8.16 (s, 1H), 7.92 (d, J=8.0 Hz, 1H), 7.80-7.77 (m, 1H), 7.76-7.75 (m, 2H), 7.53-7.51 (m, 1H), 7.17 (t, J=8.0 Hz, 1H), 6.62-6.57 (m, 3H), 5.21 (s, 2H), 5.11-5.06 (m, 1H), 4.77-4.71 (m, 1H), 4.63-4.59 (m, 1H), 4.52-4.47 (m, 1H), 4.40-4.36 (m, 2H),

[0432] To a mixture of methyl 4-(2-chloroacetamido)-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (140 mg, 0.40 mmol) and 3-fluoro-4-(((6-(piperidin-4-yloxy)pyridin-2-yl)methyl)methyl)benzonitrile (131 mg, 0.40 mmol) in N,N-dimethylformamide (3 mL) potassium carbonate (110 mg, 0.80 mmol) was added. The mixture was stirred at 60° C. for 3 hours. The resulting mixture was poured into brine (50 mL) and extracted with DCM (10 mL×2). After the combined organic phases were concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (DCM/MeOH=10/1) to obtain methyl 4-(2-(4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)methyl)piperidin-1-yl)acetamido)-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (140 mg, 54.6% yield). LC-MS m/z: 642 [M+H]⁺.


2). Synthesis of 4-(2-(4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoic acid



[0433] To a mixture of methyl 4-(2-(4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (140 mg, 0.22 mmol) in THF (1 mL) and H₂O (1 mL) lithium hydroxide (11 mg, 0.46 mmol) was added. The mixture was stirred at room temperature for 5 hours. The resulting mixture was poured into brine (10 mL) and extracted with DCM (5 mL×2). The combined organic layer was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (DCM/

MeOH=10/1) to obtain 4-(2-(4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoic acid (80 mg, 58.0% yield). LC-MS m/z: 628 [M+H]⁺.

3). Synthesis of 2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound 4)




[0434] The mixture of 4-(2-(4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoic acid (80 mg, 0.13 mmol) in toluene (1 mL) and AcOH (0.2 mL), stirred at 110° C. for 3 hours. The solvent was removed in vacuo to give a residue. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient eluted ACN/0.1% NH₄OH in H₂O solvent system; Detection Wavelength 254 nm/214 nm; Flow rate 20 mL/min) to obtain 2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (compound S4) (20.72 mg, 26.2% yield). LC-MS m/z: 610 [M+H]⁺.

[0435] ¹H NMR (400 MHz, DMSO-d₆): δ 8.07 (s, 1H), 7.92-7.89 (m, 1H), 7.83-7.80 (m, 1H), 7.71-7.67 (m, 3H), 7.64-7.59 (m, 2H), 6.45 (d, J=8.0 Hz, 1H), 6.41 (s, 1H), 6.34 (d, J=8.0 Hz, 1H), 5.71 (s, 2H), 5.43 (s, 2H), 4.73-4.71 (m, 1H), 4.03-3.97 (m, 2H), 3.81 (s, 2H), 2.68-2.65 (m, 2H), 2.25-2.20 (m, 2H), 1.75-1.71 (m, 2H), 1.43-1.39 (m, 2H), 1.16 (t, J=7.2 Hz, 3H).

Example 11. Synthesis of 2-((4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound S23)


1). Synthesis of methyl 4-(2-(4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate



[0436] To a mixture of methyl 4-(2-chloroacetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (100 mg, 0.29 mmol) and 3-fluoro-4-((3-(piperidin-4-oxy)phenoxy)methyl)benzonitrile (95 mg, 0.29 mmol) in N,N-dimethylformamide (5 mL) potassium carbonate (120 mg, 0.87 mmol) was added at room temperature. The mixture was stirred at 60° C. for 3 hours. The reaction was diluted with water (5 mL) and extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (20 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=15/1) to obtain methyl 4-(2-(4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (45 mg, 24.1% yield). LC-MS m/z: 641 [M+H]⁺.


2). Synthesis of methyl 2-((4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate

[0437] To a solution of methyl 4-(2-(4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (45 mg, 0.07 mmol) in toluene (10 mL) acetic acid (2 mL) was added. The mixture was stirred at 110° C. for 16 hours. The mixture was concentrated to give the crude product, which was further purified by reverse-phase flash chromatography under the following conditions: (Column: spherical C18, 20-40, 120 g; Mobile Phase A: 10 mM NH₃r H₂O in water; Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 60% B-80% B in 20 minutes; Monitor: 254 nm.). The mobile phase containing desired product was collected at 72% B and concentrated in vacuo to obtain methyl 2-((4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazole-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (22 mg, 57.1% yield). LC-MS m/z: 623 [M+H]⁺.



3). Synthesis of 2-((4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid

C18, 20-40 μm , 40 g; Mobile Phase A: 10 mM $\text{NH}_3 \cdot \text{H}_2\text{O}$ in water; Mobile Phase B: acetonitrile; Flow rate: 40 mL/min; Gradient: 20% B-50% B in 20 minutes; Detector: 254 nm.). The mobile phase containing the desired product was col-




[0438] To a mixture of 2-((4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (22 mg, 0.04 mmol) in THF (4 mL) and water (4 mL) lithium hydroxide monohydrate (17 mg, 0.40 mmol) was added. The mixture was stirred at room temperature for 3 hours. The resulting mixture was adjusted to pH=5-6 with hydrochloric acid solution (1 N) and extracted with ethyl acetate (10 mL $\times$ 3). The combined organic layers were washed with brine (10 mL), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by reverse-phase flash chromatography under the following conditions: (Column: spherical

lected at 29% B and concentrated in vacuo to obtain 2-((4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)yl)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (11.42 mg, 0.019 mmol, 47.5% yield). LC-MS m/z : 609 $[\text{M}+\text{H}]^+$.

[0439] ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.00 (s, 1H), 7.91 (d, $J=9.6$ Hz, 1H), 7.80-7.73 (m, 3H), 7.66 (s, 1H), 7.52 (d, $J=8.0$ Hz, 1H), 7.16 (t, $J=8.0$ Hz, 1H), 6.60-6.54 (m, 3H), 6.37 (s, 1H), 5.66 (s, 2H), 5.20 (s, 2H), 4.36-4.32 (m, 1H), 4.00 (q, $J=7.2$ Hz, 2H), 3.76 (s, 2H), 2.67-2.65 (m, 2H), 2.31-2.26 (m, 2H), 1.82-1.79 (m, 2H), 1.49-1.37 (m, 2H), 1.17 (t, $J=7.2$ Hz, 3H).

Example 12. Synthesis of (S,E)-3-(2-((4-((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazol-6-yl)acrylic acid formate


1). Synthesis of methyl (S,E)-3-(2-((4-((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazol-6-yl)acrylate




[0440] To a mixture of methyl (methyl (S, E)-3-(2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazol-6-yl)acrylate (90 mg, 0.28 mmol) and 2-((4-chloro-2-fluorobenzyl)oxy)-3-fluoro-6-(piperidin-4-yl)pyridine (100 mg, 0.28 mmol) in N,N-dimethylformamide (3 mL) potassium carbonate (78 mg, 0.56 mmol) was added. The mixture was stirred at 60° C. for 3 hours. The desired product was detected by LC-MS. The mixture was quenched by adding water (5 mL) and extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (20 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was

purified by silica gel column chromatography (dichloromethane:methanol=10:1) to obtain methyl (S, E)-3-(2-((4-((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazol-6-yl)acrylate (60 mg, 33.5% yield). LC-MS m/z: 639 [M+H]⁺.

2). Synthesis of (S,E)-3-(2-((4-((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)-methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazol-6-yl)acrylic acid formate salt






[0441] To a mixture of methyl (S,E)-3-(2-((4-((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazol-6-yl)acrylate (60 mg, 0.094 mmol) in THF (2 mL) and water (2 mL) lithium hydroxide monohydrate (6 mg, 0.24 mmol) was added. The mixture was stirred at room temperature for 5 hours. The resulting mixture was adjusted to pH=5-6 with formic acid. The mixture was concentrated to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: Sun-Fire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with

Example 13. Synthesis of (E)-3-(2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazol-6-yl)acrylic acid (Compound C2)

1). Synthesis of methyl (E)-3-(4-(2-(4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)phenyl)acrylate




ACN/0.1% FA in H₂O Solvent system; Detection Wavelength 254 nm/214 nm; Flow rate: 20 mL/min) to obtain (S,E)-3-(2-((4-((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazol-6-yl)acrylic acid formate salt (13.10 mg, 27% yield). LC-MS m/z: 625 [M+H]⁺.

[0442] ¹H NMR (400 MHz, DMSO-d₆): δ 8.27 (s, 1H), 8.00 (s, 1H), 7.66-7.47 (m, 6H), 7.32 (dd, J=10.0 Hz, 2.0 Hz, 1H), 6.54 (d, J=15.6 Hz, 1H), 6.34 (dd, J=8.8 Hz, 2.0 Hz, 1H), 5.44 (s, 2H), 5.11-5.09 (m, 1H), 4.79-4.75 (m, 2H), 4.62-4.58 (m, 1H), 4.51-4.40 (m, 2H), 3.95-3.92 (m, 1H), 3.80-3.76 (m, 1H), 2.80-2.68 (m, 3H), 2.50-2.41 (m, 1H), 2.36-2.29 (m, 2H), 1.91-1.89 (m, 2H), 1.62-1.57 (m, 2H).

[0443] To a mixture of methyl (E)-3-(4-(2-chloroacetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)phenyl)acrylate (100 mg, 0.27 mmol) and 3-fluoro-4-((6-((piperidin-4-yloxy)pyridin-2-yl)oxy)methyl)benzonitrile (88 mg, 0.27 mmol) in N,N-dimethylformamide (5 mL) potassium carbonate (75 mg, 0.54 mmol) was added. The mixture was stirred at room temperature for 16 hours. The resulting mixture was poured into brine (50 mL) and extracted with ethyl acetate (2 $\times$ 15 mL). The combined organic was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl (E)-3-(4-(2-(4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)

acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)phenyl)acrylate (100 mg, 55.5% yield). LC-MS m/z : 668 $[M+H]^+$.


2). Synthesis of methyl (E)-3-(2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazol-6-yl)acrylate




[0444] The mixture of methyl (E)-3-(4-(2-(4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidine-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)phenyl)acrylate (100 mg, 0.15 mmol) in toluene (2 mL) and acetic acid (0.2 mL) was stirred at 110° C. for 3 hours. The resulting mixture was poured into brine (20 mL) and extracted with dichloromethane (3×5 mL). The combined organic layer was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to give methyl (E)-

3-(2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazol-6-yl) acrylate (70 mg, 71.9% yield). LC-MS m/z : 650 $[M+H]^+$.

3). Synthesis of (E)-3-(2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazol-6-yl)acrylic acid




-continued



[0445] To A mixture of methyl (E)-3-(2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazol-6-yl)acrylate (70 mg, 0.11 mmol) in THF (2 mL) and water (2 mL) lithium hydroxide (5 mg, 0.21 mmol) was added. The mixture was stirred at room temperature for 16 hours. The mixture was concentrated to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10

Example 14. Synthesis of 2-((4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C3)

1). Synthesis of methyl 4-(2-(4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate




μm 19 $\times$ 250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to give (E)-3-(2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazol-6-yl)acrylic acid (13.86 mg, 19.8% yield). LC-MS m/z : 636 [M+H]⁺.

[0446] ¹H NMR (400 MHz, DMSO-*d*₆): δ 7.91 (d, *J*=9.6 Hz, 1H), 7.81 (s, 1H), 7.71-7.69 (m, 2H), 7.64-7.59 (m, 4H), 7.52-7.50 (m, 1H), 6.50 (d, *J*=8.4 Hz, 1H), 6.45 (d, *J*=7.6 Hz, 1H), 6.34 (d, *J*=8.0 Hz, 1H), 5.66 (s, 2H), 5.43 (s, 2H), 4.74-4.70 (m, 1H), 4.00-3.95 (m, 2H), 3.76 (s, 2H), 2.67-2.64 (m, 2H), 2.23-2.19 (m, 2H), 1.76-1.73 (m, 2H), 1.44-1.38 (m, 2H), 1.14-1.10 (m, 3H).

[0447] The mixture of methyl 4-(2-chloroacetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (540 mg, 1.54 mmol), 2-((4-bromo-2-fluorophenoxy)methyl)-6-(piperidin-4-yl)pyridine (586 mg, 1.54 mmol) and potassium carbonate (637 mg, 4.62 mmol) in dry N,N-dimethylformamide (15 mL) was stirred at room temperature for 16 hours. After completion, the reaction was quenched with water (10 mL) and extracted with ethyl acetate (3 $\times$ 10 mL). The combined organic layer was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 4-(2-(4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (500 mg, 46.8% yield). LC-MS m/z : 695 [M+H]⁺.


2). Synthesis of methyl 2-((4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate




[0448] The mixture of methyl 4-(2-((4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (500 mg, 0.72 mmol) in toluene (6 mL) and acetic acid (1 mL) was stirred at 110° C. for 3 hours. The mixture was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 2-((4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperi-

din-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (470 mg, 96.5% yield). LC-MS m/z: 679 [M+H]⁺.

3). Synthesis of 2-((4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid






[0449] To a mixture of methyl 2-((4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (80 mg, 0.12 mmol) in THF (1 mL) and water (1 mL) lithium hydroxide (9 mg, 0.36 mmol) was added. The reaction mixture was stirred at room temperature for 16 hours. The reaction mixture was adjusted to pH=5-6 with formic acid. The mixture was concentrated to give a residue in vacuo. The residue was eluted by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1%

2.33-2.23 (m, 2H), 1.81-1.78 (m, 2H), 1.47-1.44 (m, 2H), 1.16 (t, J=7.2 Hz, 3H).

Example 15. Synthesis of (S)-2-((4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C4)

1). Synthesis of methyl (S)-2-((4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate




FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to give 2-((4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (30.96 mg, 39% yield). LC-MS m/z: 665 [M+H]⁺.

[0450] ¹H NMR (400 MHz, DMSO-d₆): δ 8.19 (s, 1H), 8.07 (s, 1H), 7.81 (dd, J=8.8 Hz, 1.6 Hz, 1H), 7.72-7.67 (m, 3H), 7.54 (dd, J=6.8 Hz, 2.4 Hz, 1H), 7.30 (d, J=8.8 Hz, 1H), 7.21 (t, J=8.8 Hz, 1H), 7.03 (d, J=7.2 Hz, 1H), 6.70 (d, J=8.4 Hz, 1H), 6.41 (s, 1H), 5.72 (s, 2H), 5.17 (s, 2H), 4.88-4.86 (m, 1H), 4.03-3.98 (m, 2H), 3.81 (s, 2H), 2.69-2.67 (m, 2H),

[0451] The mixture of methyl (S)-2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 0.34 mmol), 2-((4-bromo-2-fluorophenoxy)methyl)-6-(piperidin-4-yl)pyridine (130 mg, 0.34 mmol) and potassium carbonate (140 mg, 1.02 mmol) in N,N-dimethylformamide (4 mL) was stirred at 60° C. for 3 hours. The mixture was quenched with water (10 mL) and extracted with ethyl acetate (3 $\times$ 10 mL). The combined organic layer was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl (S)-2-((4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 46.1% yield). LC-MS m/z: 639 [M+H]⁺.

2). Synthesis of (S)-2-((4-(((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate




[0452] To a mixture of methyl (S)-2-((4-(((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 0.16 mmol) in THF (2 mL) and water (2 mL) lithium hydroxide (12 mg, 0.48 mmol) was added. The reaction mixture was stirred at room temperature for 16 hours. The reaction mixture was adjusted to pH=5-6 with formic acid. The solvent was removed by concentration to give residue in vacuo. The residue purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 1 Oum 19×250 mm; Gradient elution with ACN/0.10% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain (S)-2-((4-(((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (20.08 mg, 20.5% yield). LC-MS m/z: 627 [M+H]⁺.


[0453] ¹H NMR (400 MHz, DMSO-d₆): δ 8.28-8.23 (m, 1H), 7.80 (d, J=8.8 Hz, 1H), 7.71 (t, J=7.6 Hz, 1H), 7.63 (d, J=8.4 Hz, 1H), 7.54 (dd, J=10.4 Hz, 2.0 Hz, 1H), 7.31 (d, J=9.2 Hz, 1H), 7.22 (t, J=8.8 Hz, 1H), 7.04 (d, J=7.2 Hz, 1H), 6.72 (d, J=8.4 Hz, 1H), 5.18 (s, 2H), 5.13-5.06 (m, 1H), 4.97-4.89 (m, 1H), 4.83-4.73 (m, 1H), 4.68-4.60 (m, 1H), 4.54-4.46 (m, 1H), 4.41-4.33 (m, 1H), 3.98-3.90 (m, 1H), 3.82-3.75 (m, 1H), 2.82-2.66 (m, 4H), 2.46-2.36 (m, 1H), 2.33-2.26 (m, 2H), 1.97-1.88 (m, 2H), 1.66-1.56 (m, 2H).

Example 16. Synthesis of 2-((4-(((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-3-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C82)

1). Synthesis of methyl 2-((4-(((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-3-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate



-continued




[0454] The mixture of methyl 2-(chloromethyl)-1-(oxetan-3-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 0.34 mmol), 3-fluoro-4-((6-(piperidin-4-oxy)pyridin-2-yl)methoxy)benzonitrile (111 mg, 0.34 mmol) and potassium carbonate (94 mg, 0.68 mmol) in N,N-dimethylformamide (2 mL) was stirred at 60° C. for 3 hours. The resulting mixture was diluted with water (30 mL) and extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (30 mL), concentrated under to give a residue reduced pressure. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-3-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 50.3% yield). LC-MS m/z: 586 [M+H]⁺.

2). Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-3-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid


[0455] To a mixture of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-3-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 0.17 mmol) in THF (1.0 mL) and water (1.0 mL) lithium hydroxide (8 mg, 0.34 mmol) was added. The reaction mixture was stirred at room temperature for 16 hours. The solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μm 19×250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-3-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (25.81 mg, 26.6% yield). LC-MS m/z: 572 [M+H]⁺.

[0456] ¹H NMR (400 MHz, DMSO-d₆): δ 8.22 (s, 1H), 7.88 (dd, J=11.2, 2.0 Hz, 1H), 7.81 (dd, J=8.4, 1.2 Hz, 1H), 7.73 (t, J=8.0 Hz, 1H), 7.67-7.61 (m, 2H), 7.45 (t, J=8.4 Hz, 1H), 7.06 (d, 0.1=7.2 Hz, 1H), 6.74 (d, J=8.4 Hz, 1H), 5.31 (s, 2H), 4.94-4.91 (m, 1H), 4.72 (d, 0.1=7.6 Hz, 2H), 4.64-4.61 (m, 2H), 4.56-4.53 (m, 2H), 3.81 (s, 2H), 3.67-3.61 (m, 1H), 2.80-2.77 (m, 2H), 2.33-2.27 (m, 2H), 1.93-1.90 (m, 2H), 1.64-1.59 (m, 2H).





Example 17. Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((3-methyloxetan-3-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C5)

1). Synthesis of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((3-methyloxetan-3-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate



[0457] The mixture of methyl 2-(chloromethyl)-1-((3-methyloxetan-3-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 0.32 mmol), 3-fluoro-4-((6-(piperidin-4-oxymethyl)pyridin-2-yl)methoxy)benzonitrile (105 mg, 0.32 mmol) and potassium carbonate (88 mg, 0.64 mmol) in N,N-dimethylformamide (2 mL) was stirred at 60° C. for 3 hours. The resulting mixture was diluted with water (30 mL) and extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with saturated brine (30 mL) and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)-pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((3-methyloxetan-3-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (50 mg, 26.0% yield). LC-MS m/z: 600 [M+H]⁺.

2). Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((3-methyloxetan-3-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid




[0458] To a mixture of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)-pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((3-methyloxetan-3-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (50 mg, 0.08 mmol) in THF (5 mL) and water (5 mL) lithium hydroxide (10 mg, 0.42 mmol) was added. The mixture was stirred at room temperature for 16 hours. The solvent was removed to give a residue by concentration in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μm 19×250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((3-methyloxetan-3-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (9.50 mg, 20.3% yield). LC-MS m/z: 586 [M+H]⁺.

[0459] ¹H NMR (400 MHz, DMSO-d₆): δ 8.21 (s, 1H), 7.88 (dd, J=11.2, 1.6 Hz, 1H), 7.81 (d, J=9.2 Hz, 1H), 7.73 (t, J=8.0 Hz, 1H), 7.66 (d, J=8.4 Hz, 2H), 7.44 (t, J=8.8 Hz, 1H), 7.05 (d, J=7.2 Hz, 1H), 6.73 (d, J=8.4 Hz, 1H), 5.30 (s, 2H), 4.92-4.90 (m, 1H), 4.65-4.61 (m, 4H), 4.22 (d, J=6.0 Hz, 1H), 3.78 (s, 2H), 2.76-2.74 (m, 2H), 2.31-2.26 (m, 2H), 1.92-1.87 (m, 2H), 1.62-1.58 (m, 2H), 1.25 (s, 3H).



Example 18. Synthesis of (S)-2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound 67)

1). Synthesis of methyl (S)-2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate



[0460] To A mixture of methyl (S)-2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 0.34 mmol) and 3-fluoro-4-((3-(piperidin-4-oxy)benzyl)oxy)benzonitrile (122 mg, 0.37 mmol) in N,N-dimethylformamide (5 mL) potassium carbonate (94 mg, 0.68 mmol) was added. The mixture was stirred at room temperature for 16 hours. The mixture was quenched by adding water (5 mL) and extracted with ethyl acetate (40 mL $\times$ 3). The combined organic layers were washed with brine (30 mL $\times$ 2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl (S)-2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (71 mg, 35.8% yield). LC-MS m/z : 585 $[M+H]^+$.

2). Synthesis of (S)-2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid




[0461] To a mixture of methyl (S)-2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (71 mg, 0.12 mmol) in THF (2 mL) and water (2 mL) lithium hydroxide monohydrate (50 mg, 1.20 mmol) was added at room temperature. The reaction was stirred at room temperature for 5 hours. The resulting mixture was adjusted to pH=5-6 with hydrochloric acid (1M), and then extracted with ethyl acetate (10 mL $\times$ 3). The combined organic layers were washed with brine (10 mL), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by reverse-phase flash chromatography under the following conditions (Column: Spherical C18, 20-40 μm , 40 g; Mobile Phase A: 10 mM NH_3H_2O in water; Mobile Phase B: acetonitrile; Flow rate: 40 mL/min; Gradient: 20% B-50% B in 20 minutes; Detector: 254 nm) to obtain (S)-2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (4.69 mg, 7.3% yield). LC-MS m/z : 571 $[M+H]^+$.

[0462] ^1H NMR (400 MHz, $\text{DMSO}-d_6$) 8.25 (s, 1H), 7.88-7.80 (m, 2H), 7.68 (d, $J=8.4$ Hz, 1H), 7.61 (d, $J=8.0$ Hz, 1H), 7.42 (t, $J=8.4$ Hz, 1H), 7.30 (t, $J=7.6$ Hz, 1H), 7.05-6.93 (m, 3H), 5.25 (s, 2H), 5.10-5.07 (m, 1H), 4.80-4.75 (m, 1H), 4.63 (d, $J=13.2$ Hz, 1H), 4.52-4.47 (m, 1H), 4.42-4.35 (m, 2H), 3.94 (d, $J=13.2$ Hz, 1H), 3.78 (d, $J=13.2$ Hz, 1H), 2.79-2.67 (m, 3H), 2.46-2.35 (m, 3H), 1.95-1.93 (m, 2H), 1.64-1.60 (m, 2H).

Example 19. Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1,2-dimethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C6)


1). Synthesis of methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1,2-dimethyl-1H-imidazol-5-yl)methyl)amino)benzoate




[0463] To a mixture of methyl 4-(2-chloroacetamido)-3-((1,2-dimethyl-1H-imidazol-5-yl)methyl)amino)benzoate (150 mg, 0.43 mmol) and 3-fluoro-4-((6-(piperidin-4-oxy)pyridin-2-yl)methoxy)benzonitrile (141 mg, 0.43 mmol) in N,N -dimethylformamide (5 mL) potassium carbonate (119 mg, 0.86 mmol) was added. The mixture was stirred at room temperature for 16 hours. After completion, the mixture was poured into brine (50 mL) and extracted with ethyl acetate (2x20 mL). The combined organic layer was concentrated to give a residue in vacuo. The residue was purified by silica

gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1,2-dimethyl-1H-imidazol-5-yl)methyl)amino)benzoate (150 mg, 54.3% yield). LC-MS m/z : 642 $[\text{M}+\text{H}]^+$.

2). Synthesis of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1,2-dimethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate




-continued




[0464] The mixture of methyl 4-(2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1,2-dimethyl-1H-imidazol-5-yl)methyl)amino)benzoate (150 mg, 0.08 mmol) in toluene (5 mL) and acetic acid (1 mL) was stirred at 110° C. for 3 hours. The mixture was poured into brine (20 mL) and extracted with dichloromethane (2×10 mL). The combined organic layer was concentrated in vacuo to give methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1,2-dimethyl-1H-imidazol-5-yl)methyl)-1H-benzo-[d]imidazole-6-carboxylate (100 mg, 69.8% yield). LC-MS *m/z*: 624 [M+H]⁺.

-continued



3). Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1,2-dimethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid formate salt




[0465] To a mixture of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1,2-dimethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 0.17 mmol) in THF (5 mL) and water (5 mL) lithium hydroxide (8 mg, 0.34 mmol) was added. The reaction mixture was stirred at room temperature for 16 hours. The solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19×250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtained 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidine-1-yl)methyl)-1-((1,2-dimethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid formate salt (71.07 mg, 68.6% yield). LC-MS *m/z*: 610 [M+H]⁺.

[0466] ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.16 (s, 2H), 8.09 (s, 1H), 7.88 (dd, *J*=11.2, 1.6 Hz, 1H), 7.81 (dd, *J*=8.4, 1.2 Hz, 1H), 7.74-7.65 (m, 3H), 7.44 (t, *J*=8.4 Hz, 1H), 7.05 (d, *J*=7.2 Hz, 1H), 6.72 (d, *J*=8.4 Hz, 1H), 6.41 (s, 1H), 5.67 (s, 2H), 5.30 (s, 2H), 4.90-4.86 (m, 1H), 3.82 (s, 2H), 3.44 (s, 3H), 2.72-2.69 (m, 2H), 2.33-2.24 (m, 5H), 1.84-1.82 (m, 2H), 1.54-1.47 (m, 2H).

Example 20. Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound 135)

1). Synthesis of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylate

benzonitrile (173 mg, 0.50 mmol) in N,N-dimethylformamide (4 mL) potassium carbonate (207 mg, 1.50 mmol) was added at room temperature. The mixture was stirred at 60° C. for 2 hours. The mixture was quenched with water (5 mL) and extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (20 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 2-((4-




[0467] To a mixture of methyl 2-(chloromethyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylate (160 mg, 0.50 mmol) and 3-fluoro-4-((3-fluoro-6-(piperidin-4-oxypyridin-2-yl)methoxy)

((6-((4-cyano-2-fluorophenoxy)methyl)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylate (17 mg, 6.0% yield). LC-MS m/z: 627 [M+H]⁺.

2). Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid

trated to give a residue in vacuo. The residue was purified by reverse-phase flash chromatography under the following conditions (Column: Spherical C18, 20-40 μm , 120 g; Mobile Phase A: 10 mM $\text{NH}_3\text{H}_2\text{O}$ in water); Mobile Phase



[0468] To a mixture of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylate (17 mg, 0.03 mmol) in THF (3 mL) and water (10 mL) lithium hydroxide monohydrate (12 mg, 0.30 mmol) was added. The mixture was stirred at room temperature for 3 hours. The resulting mixture was adjusted to pH=5-6 with hydrochloric acid (1 M), and then extracted with ethyl acetate (10 mL $\times$ 3). The combined organic layers were washed with brine (10 mL), dried over anhydrous sodium sulfate, filtered, and concen-


B: acetonitrile; Flow rate: 80 mL/min; Gradient: 30% B-50% B in 20 minutes; Detector: 254 nm) to obtain 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (7.02 mg, 33.3% yield). LC-MS m/z : 613 $[\text{M}+\text{H}]^+$.

[0469] ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.23 (s, 1H), 7.87 (dd, $J=11.2, 1.6$ Hz, H), 7.82-7.80 (m, 1H), 7.74-7.63 (m, 3H), 7.51 (t, $J=8.4$ Hz, 1H), 6.84 (dd, $J=8.8, 2.8$ Hz, 1H), 5.40 (s, 2H), 4.74-4.72 (m, 1H), 4.59 (s, 2H), 3.83 (s, 2H), 2.78-2.75 (m, 2H), 2.68 (s, 2H), 2.24 (t, $J=9.2$ Hz, 2H), 1.84 (d, $J=10.0$ Hz, 2H), 1.60-1.55 (m, 2H), 0.74-0.67 (m, 4H).

Example 21. Synthesis of (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C83)

1). Synthesis of methyl (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate

ture. The mixture was stirred at room temperature for 16 hours. The reaction was quenched with water (5 mL) and extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (20 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by reverse-phase flash chromatography under the following conditions (Column: Spherical C18, 20-40 μ m, 120 g; Mobile Phase A: 10 mM hexachlorocyclohexane in water; Mobile Phase B:




[0470] To a mixture of 3-fluoro-4-((3-fluoro-6-(piperidin-4-oxy)pyridin-2-yl)methoxy)benzonitrile (90 mg, 0.26 mmol) and methyl (S)-2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (81 mg, 0.26 mmol) in N,N-dimethylformamide (4 mL) potassium carbonate (108 mg, 0.78 mmol) was added at room tempera-

ture. The mixture was stirred at room temperature for 16 hours. The reaction was quenched with water (5 mL) and extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (20 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by reverse-phase flash chromatography under the following conditions (Column: Spherical C18, 20-40 μ m, 120 g; Mobile Phase A: 10 mM hexachlorocyclohexane in water; Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 30% B-50% B in 20 minutes; Detector: 254 nm) to obtain methyl (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)-5-fluoropyridin-2-yl)oxy)piperidine-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (85 mg, 53.8% yield). LC-MS m/z: 604 [M+H]⁺.

2). (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid

Spherical C18, 20-40 μm , 120 g; Mobile Phase A: water 10 mM $\text{NH}_3 \cdot \text{H}_2\text{O}$ in water; Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 30% B-50% B in 20 minutes; Detector: 254 nm) to obtain (S)-2-((4-((6-((4-cyano-2-fluo-




[0471] To a mixture of methyl (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)-5-fluoropyridin-2-yl)oxy)piperidine-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (85 mg, 0.14 mmol) in THF (3 mL) and water (3 mL) lithium hydroxide monohydrate (56 mg, 1.4 mmol) was added. The reaction was stirred at room temperature for 3 hours. The resulting mixture was adjusted to pH=5-6 with hydrochloric acid (1M), and then extracted with ethyl acetate (10 mL $\times$ 3). The combined organic layers were washed with brine (10 mL), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by reverse-phase flash chromatography under the following conditions (Column:

rophenoxy)methyl)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (50.26 mg, 64.3% yield). LC-MS m/z : 590 $[\text{M}+\text{H}]^+$.

[0472] ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.25 (s, 1H), 7.88 (dd, $J=11.2, 1.6$ Hz, 1H), 7.79 (dd, $J=8.4, 1.6$ Hz, 1H), 7.74-7.67 (m, 2H), 7.63 (d, $J=8.4$ Hz, 1H), 7.50 (t, $J=8.8$ Hz, 1H), 6.84 (dd, $J=8.8, 3.2$ Hz, 1H), 5.41 (s, 2H), 5.09-5.07 (m, 1H), 4.80-4.70 (m, 2H), 4.62 (dd, $J=16.0, 2.8$ Hz, 1H), 4.49 (q, $J=7.6$, 1H), 4.39-4.34 (m, 1H), 3.92 (d, $J=13.6$ Hz, 1H), 3.76 (d, $J=13.6$ Hz, 1H), 2.76-2.66 (m, 3H), 2.45-2.40 (m, 1H), 2.25-2.20 (m, 1H), 1.84-1.81 (m, 2H), 1.57-1.53 (m, 2H).

Example 22. Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H benzo[d]imidazole-6-carboxylic acid (Compound 3)


1). methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate




[0473] To a mixture of methyl 4-(2-chloroacetamido)-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (130 mg, 0.37 mmol) and 3-fluoro-4-((3-fluoro-6-(piperidin-4-oxypyridin-2-yl)methoxy)benzonitrile) (128 mg, 0.37 mmol) in N,N-dimethylformamide (4 mL) potassium carbonate (153 mg, 1.11 mmol) was added at room temperature. The mixture was stirred at room temperature for 16 hours. The reaction was quenched with water (5 mL) and extracted with ethyl acetate (20 mL $\times$ 3). The combined organic layers were washed with brine (20 mL $\times$ 2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in

vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to give methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (116 mg, 48.6% yield). LC-MS m/z : 660 $[\text{M}+\text{H}]^+$.

2). methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate




-continued




[0474] To a solution of methyl 4-(2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (116 mg, 0.18 mmol) in dioxane (6 mL) acetic acid (2 mL) was added at room temperature. The mixture was stirred at 100° C. for 3 hours. The reaction was quenched with water (5 mL) and extracted with ethyl acetate (40 mL×3). The combined organic layers were washed with brine (40 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography

(dichloromethane/methanol=10/1) to obtain methyl 2-((6-((4((4-cyano-2-fluorophenoxy)methyl)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (18 mg, 15.6% yield). LC-MS m/z : 642 $[M+H]^+$.

3). Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid






[0475] To a mixture of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (18 mg, 0.028 mmol) in THF (3 mL) and water (3 mL) lithium hydroxide monohydrate (11 mg, 0.28 mmol) was added. The mixture was stirred at room temperature for 3 hours. The resulting mixture was adjusted to pH=5-6 with hydrochloric acid 1 (1 M) and then extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (10 mL), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by reversed-phase flash chromatography under the following conditions (Column: Spherical C18, 20-40 μ m, 120 g; Mobile Phase A: 10 mM $\text{NH}_3\text{H}_2\text{O}$ in water; Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 30% B-50% B in 20 min; Detector: 254 nm) to obtain 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (5.37 mg, 32% yield). LC-MS m/z : 628 $[\text{M}+\text{H}]^+$.

[0476] ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.06 (s, 1H), 7.87 (dd, $J=11.6$, 2.0 Hz, 1H), 7.81 (d, $J=8.4$ Hz, 1H), 7.73-7.66 (m, 4H), 7.49 (t, $J=8.4$ Hz, 1H), 6.83 (dd, $J=9.2$, 2.8 Hz, 1H), 6.40 (s, 1H), 5.70 (s, 2H), 5.40 (s, 2H), 4.67-4.65 (m, 1H), 4.02-3.97 (m, 2H), 3.80 (s, 2H), 2.67-2.63 (m, 2H), 2.19-2.15 (m, 2H), 1.72-1.70 (m, 2H), 1.43-1.34 (m, 2H), 1.16 (t, $J=7.2$ Hz, 3H).

Example 23. Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-3-((1-(cyanomethyl)cyclopropyl)methyl)-3H-imidazo[4,5-b]pyridine-5-carboxylic acid (Compound 138)


1). Synthesis of methyl 5-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-6-((1-(cyanomethyl)cyclopropyl)methyl)amino)picolinate



[0477] To a mixture of methyl 5-(2-chloroacetamido)-6-(1-(cyanomethyl)cyclopropyl)methylamino)picolinate (150 mg, 0.45 mmol) and 3-fluoro-4-((6-(piperidin-4-oxy)pyridin-2-yl)methoxy)benzonitrile (175 mg, 0.54 mmol) in N,N-dimethylformamide (5 mL) potassium carbonate (185 mg, 1.34 mmol) was added. The mixture was stirred at room temperature for 16 hours. The resulting mixture was poured into brine (20 mL) and extracted with ethyl acetate (20 mL $\times$ 2). The combined organic layers were concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/metha-


nol=20/1) to obtain methyl 5-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-6-((1-(cyanomethyl)cyclopropyl)methylamino)picolinate: (220 mg, 78.6% yield). LC-MS m/z : 628 $[M+H]^+$.

2). Synthesis of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-3-((1-(cyanomethyl)cyclopropyl)methyl)-3H-imidazo[4,5-b]pyridine-5-carboxylate




[0478] The mixture of methyl 5-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-6-((1-(cyanomethyl)cyclopropyl)methylamino)picolinate (220 mg, 0.35 mmol) in toluene (10 mL) and acetic acid (10 mL) was stirred at 110° C. for 16 h under nitrogen atmosphere. After completion, the resulting mixture was filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=40/1) to obtain methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)-methyl)pyridin-2-yl)oxy)-piperidin-1-yl)methyl)-3-((1-(cyanomethyl)cyclopropyl)methyl)-3H-imidazo[4,5-b]pyridine-5-carboxylate (200 mg, 93.5% yield). LC-MS m/z : 610 $[M+H]^+$.

3). Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-3-((1-(cyanomethyl)cyclopropyl)methyl)-3H-imidazo[4,5-b]pyridine-5-carboxylic acid



-continued




[0479] To a mixture of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-3-((1-(cyanomethyl)cyclopropyl)methyl)-3H-imidazo[4,5-b]pyridine-5-carboxylate (100 mg, 0.16 mmol) in THF (5 mL) and water (5 mL) lithium hydroxide monohydrate (69 mg, 1.60 mmol) was added. The mixture was stirred at 40° C. for 3 hours. The solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 um 19x250 mm; Gradient elution with ACN/0.05% NH₄OH in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidine-1-yl)methyl)-3-((1-(cyanomethyl)cyclopropyl)methyl)-3H-imidazo[4,5-b]pyridine-5-carboxylic acid (48.10 mg, 49.2% yield). LC-MS m/z: 596 [M+H]⁺.

[0480] ¹H NMR (400 MHz, DMSO-d₆) δ 8.03 (d, J=8.4 Hz, 1H), 7.94-7.86 (m, 2H), 7.73 (t, J=8.0 Hz, 1H), 7.66 (d, J=8.8 Hz, 1H), 7.45 (t, J=8.8 Hz, 1H), 7.05 (d, J=7.2 Hz, 1H), 6.73 (d, J=8.4 Hz, 1H), 5.31 (s, 2H), 4.94-4.90 (m, 1H), 4.52 (s, 2H), 3.86 (s, 2H), 2.83-2.76 (m, 4H), 2.33-2.29 (m, 2H), 1.93-1.91 (m, 2H), 1.66-1.59 (m, 2H), 1.05 (s, 2H), 0.62-0.61 (m, 2H).

Example 24. Synthesis of (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-3-(oxetan-2-ylmethyl)-3H-imidazo[4,5-b]pyridine-5-carboxylic acid (Compound 60)

1). Synthesis of methyl (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-3-(oxetan-2-ylmethyl)-3H-imidazo[4,5-b]pyridine-5-carboxylate




[0481] To a mixture of methyl (S)-2-(chloromethyl)-3-(oxetan-2-ylmethyl)-3H-imidazo[4,5-b]pyridine-5-carboxylate (180 mg, 0.61 mmol) and potassium carbonate (248 mg, 1.83 mmol) in N,N-dimethylformamide (5 mL) 3-fluoro-4-((6-(piperidine-4-oxy)pyridin-2-yl)methoxy)benzonitrile (216 mg, 0.67 mmol) was added. The mixture was stirred at 60° C. for 3 hours under nitrogen atmosphere. After completion, the mixture was poured into water (10 mL) and extracted with ethyl acetate (10 mL×2). The combined organic layers were washed with saturated brine (10 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)-methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-3-(oxetan-2-ylmethyl)-3H-imidazo[4,5-b]pyridine-5-carboxylate (190 mg, 48.7% yield). LCMS *m/z*: 587 [M+H]⁺.

2). Synthesis of (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)-methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-3-(oxetan-2-ylmethyl)-3H-imidazo[4,5-b]pyridine-5-carboxylic acid

(3 mL) with lithium hydroxide (36 mg, 1.50 mmol) was added. The reaction was stirred at 25° C. for 16 hours. The resulting mixture was adjusted to pH=5-6 with formic acid, and then extracted with ethyl acetate (10 mL×2). The combined organic layers were washed with saturated brine (5 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by reverse-phase flash chromatography under the following conditions (Column: Spherical C18, 20-m, 120 g; Mobile Phase A: Water 10 mM formic acid in Water); Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 30% B-50% B in 16 min; Detector: 254 nm) to obtain (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)-methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-3-(oxetan-2-ylmethyl)-3H-imidazo[4,5-b]pyridine-5-carboxylic acid (10.82 mg, 13.8% yield). LC-MS *m/z*: 573 [M+H]⁺.

[0483] ¹H NMR (400 MHz, DMSO-d₆) δ 8.12 (d, J=8.4 Hz, 1H), 7.98 (d, J=8.4 Hz, 1H), 7.88 (dd, J=11.6 Hz, 2.0 Hz, 1H), 7.73 (t, J=8.0 Hz, 1H), 7.66 (d, J=8.8 Hz, 1H), 7.45 (t, J=8.4 Hz, 1H), 7.05 (d, J=7.6 Hz, 1H), 6.74 (d, J=8.4 Hz,




[0482] To a solution of methyl (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)-methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-3-(oxetan-2-ylmethyl)-3H-imidazo[4,5-b]pyridine-5-carboxylate (90 mg, 0.15 mmol) in THF (3 mL) and water

1H), 5.31 (s, 2H), 5.18-5.15 (m, 1H), 4.94-4.81 (m, 2H), 4.74-4.69 (m, 1H), 4.52-4.47 (m, 1H), 4.38-4.33 (m, 1H), 4.00-3.90 (m, 2H), 2.76-2.67 (m, 4H), 2.35-2.31 (m, 2H), 1.93-1.91 (m, 2H), 1.66-1.62 (m, 2H).

Example 25. Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-3-((1-ethyl-1H-imidazol-5-yl)methyl)-3H-imidazo[4,5-b]pyridine-5-carboxylic acid (Compound 6)


1). Synthesis of methyl 5-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-6-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)picolinate




[0484] To a mixture of methyl 5-(2-chloroacetamido)-6-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)picolinate (180 mg, 0.51 mmol) and 3-fluoro-4-((6-(piperidin-4-oxy)pyridin-2-yl)methoxy)benzonitrile (167 mg, 0.51 mmol) in N,N-dimethylformamide (10 mL) potassium carbonate (211 mg, 1.53 mmol) was added at room temperature. The mixture was stirred at 25° C. for 16 hours. The reaction mixture was quenched with H₂O (5 mL) and extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with saturated brine (20 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give residue in vacuo.

The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 5-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-6-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)picolinate (230 mg, 71.6% yield). LC-MS m/z: 643 [M+H]⁺.

2). methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-3-((1-ethyl-1H-imidazol-5-yl)methyl)-3H-imidazo[4,5-b]pyridine-5-carboxylate




-continued



[0485] The mixture of methyl 5-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-6-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)picolinate (230 mg, 0.36 mmol) in toluene (3 mL) and acetic acid (0.5 mL) was stirred at 100° C. for 16 hours. The resulting mixture was poured into saturated brine (10 mL) and extracted with dichloromethane (5 mL×2). The combined organic layer was concentrated to give methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-3-((1-ethyl-1H-imidazol-5-yl)methyl)-3H-imidazo[4,5-b]pyridine-5-carboxylate (180 mg, 81% yield). LC-MS *m/z*: 625 [M+H]⁺.

3). Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-3-((1-ethyl-1H-imidazol-5-yl)methyl)-3H-imidazo[4,5-b]pyridine-5-carboxylic acid


[0486] To a mixture of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-3-((1-ethyl-1H-imidazol-5-yl)methyl)-3H-imidazo[4,5-b]pyridine-5-carboxylate (130 mg, 0.20 mmol) in THF (3 mL) and water (3 mL) lithium hydroxide (32 mg, 2.0 mmol) was added. The mixture was stirred at room temperature for 16 hours. The resulting mixture was adjusted to pH 5-6 with hydrochloric acid (1 M) and extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with saturated brine (10 mL), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by reversed-phase flash chromatography (Column: sphere C18, 20-40 μm, 120 g; Mobile phase A: 10 mM NH₃·H₂O in water; Mobile phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 30% B-50% B in 20 min; Detector: 254 nm) to obtain 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-3-((1-ethyl-1H-imidazol-5-yl)methyl)-3H-imidazo[4,5-b]pyridine-5-carboxylic acid (30.28 mg, 24.5% yield). LC-MS *m/z*: 611 [M+H]⁺.



[0487] ^1H NMR (400 MHz, DMSO-d_6) δ 8.16-8.13 (m, 1H), 8.02-7.99 (m, 1H), 7.88 (d, $J=11.6$ Hz, 1H), 7.74-7.70 (m, 1H), 7.67-7.65 (m, 2H), 7.46-7.42 (m, 1H), 7.05 (d, $J=7.6$ Hz, 1H), 6.74-6.71 (m, 2H), 5.72 (s, 2H), 5.30 (s, 2H), 4.90-4.87 (m, 1H), 4.17-4.12 (m, 2H), 3.80 (s, 2H), 2.73-2.67 (m, 2H), 2.28-2.24 (m, 2H), 1.87-1.84 (m, 2H), 1.57-1.55 (m, 2H), 1.12-1.08 (m, 3H).

Example 26. Synthesis of 2-((4-((6-((4-chloro-2-fluorophenylthio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C7)


1). Synthesis of methyl 4-(2-(4-((6-((4-chloro-2-fluorophenylthio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)benzoate




[0488] To a mixture of methyl 4-(2-chloroacetamido)-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)benzoate (200 mg, 0.60 mmol) and 2-((4-chloro-2-fluorophenylthio)methyl)pyridine (210 mg, 0.60 mmol) in N,N -dimethylformamide (4 mL) potassium carbonate (247 mg, 1.80 mmol) was added. The reaction was stirred at room temperature for 16 hours. After completion, the resulting mixture was poured into brine (10 mL) and extracted with ethyl acetate (2x5 mL). The combined organic layer was concentrated to give residue in vacuo. The residue was purified by silica gel column chromatography (dichlo-

romethane/methanol=20/1) to obtain methyl 4-(2-(4-((6-((4-chloro-2-fluorophenylthio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)benzoate (300 mg, 77.3% yield). LC-MS m/z : 652 $[\text{M}+\text{H}]^+$.

2). Synthesis of methyl 2-((4-((6-((4-chloro-2-fluorophenylthio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylate



-continued




[0489] The mixture of methyl 4-(2-((4-((6-((4-chloro-2-fluorophenylthio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)benzoate (300 mg, 0.46 mmol) in toluene (10 mL) and acetic acid (4 mL) was stirred at 110° C. for 16 hours under nitrogen atmosphere. After completion, the mixture was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 2-((4-((6-((4-chloro-2-fluorophenylthio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylate (210 mg, 80.0% yield). LC-MS m/z: 634 [M+H]⁺.

3). Synthesis of 2-((4-((6-((4-Chloro-2-fluorophenylthio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid

methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylate (210 mg, 0.33 mmol) in THF (1 mL) and water (1 mL) lithium hydroxide (79 mg, 3.30 mmol) was added. The mixture was stirred at 40° C. for 16 hours. The solvent was removed by concentration in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.05% NH₄OH in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain 2-((4-((6-((4-chloro-2-fluorophenylthio)methyl)pyridin-2-yl)oxy)piperidine-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (19.36 mg, 9.4% yield). LC-MS m/z: 620 [M+H]⁺.

[0491] ¹H NMR (400 MHz, DMSO-d₆) δ 8.23 (s, 1H), 7.81 (d, J=8.8 Hz, 1H), 7.65-7.61 (m, 3H), 7.29-7.21 (m, 2H), 6.98 (d, J=7.2 Hz 1H), 6.63 (d, J=8.4 Hz, 1H), 4.87-4.85 (m, 1H), 4.59 (s, 2H), 4.30 (s, 2H), 3.83 (s, 2H), 2.78-2.75 (m, 2H), 2.68 (d, J=6.8 Hz, 2H), 2.32-2.28 (m,




[0490] To a mixture of methyl 2-((4-((6-((4-chloro-2-fluorophenylthio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)

2H), 1.90-1.88 (m, 2H), 1.60-1.57 (m, 2H), 0.74-0.67 (m, 4H).

Example 27. Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound 133)


1). Synthesis of methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)benzoate




[0492] To a mixture of methyl 4-(2-chloroacetamido)-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)benzoate (160 mg, 0.48 mmol) and 3-fluoro-4-((6-(piperidin-4-yl)pyridin-2-yl)methoxy)benzonitrile (187 mg, 0.57 mmol) in N,N-dimethylformamide (10 mL) potassium carbonate (197 mg, 1.43 mmol) was added. The reaction mixture was stirred at room temperature for 16 hours. The mixture was poured into brine (20 mL) and extracted with ethyl acetate (2x20 mL). The combined organic layer was concentrated to give a residue in vacuo. The residue was purified by silica gel

column chromatography (dichloromethane/methanol=20/1) to give methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)benzoate (290 mg, 94.2% yield). LC-MS m/z : 627 [M+H]⁺.

2). Synthesis of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylate






[0493] The mixture of methyl 4-(2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)benzoate (290 mg, 0.46 mmol) in toluene (10 mL) and acetic acid (10 mL) was stirred at 110° C. for 16 hours under nitrogen atmosphere. The mixture was concentrated and purified by silica gel column chromatography (dichloromethane/methanol=40/1) to obtain methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)-methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylate (250 mg, 88.9% yield). LC-MS *m/z*: 609 [M+H]⁺.

3). Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid

THF (5 mL) and water (5 mL) lithium hydroxide monohydrate (86 mg, 2.10 mmol) was added. The mixture was stirred at 40° C. for 3 hours. The solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.05% NH₄OH in 1H₂O solvent system; Detection Wavelength 254 nm/214 nm; Flow rate 20 mL/min) to obtain 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidine-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (40.09 mg, 32.8% yield). LC-MS *m/z*: 595 [M+H]⁺.

[0495] ¹H NMR (400 MHz, DMSO-d₆) 8.22 (s, 1H), 7.89-7.86 (m, 1H), 7.81 (d, J=8.4 Hz, 1H), 7.73 (t, J=7.6 Hz, 1H), 7.67-7.61 (m, 2H), 7.45 (t, J=8.8 Hz, 1H), 7.05 (d,




[0494] To a mixture of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)-methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylate (125 mg, 0.21 mmol) in

J=7.6 Hz, 1H), 6.73 (d, J=8.0 Hz, 1H), 5.31 (s, 2), 4.94-4.90 (m, 1H), 4.60 (s, 2H), 3.84 (s, 2H), 2.80-2.77 (m, 2H), 2.70 (s, 2H), 2.33-2.28 (m, 2H), 1.92-1.90 (m, 2H), 1.65-1.58 (m, 2H), 0.74-0.69 (m, 4H).

Example 28. Synthesis of (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-4-fluoro-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound 56)


1). Synthesis of (S)-N-(4-bromo-2-fluoro-6-((oxetan-2-ylmethyl)amino)phenyl)-2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido




[0496] The mixture of (S)-N-(4-bromo-2-fluoro-6-((oxetan-2-ylmethyl)amino)phenyl)-2-chloroacetamido (230 mg, 0.65 mmol), 3-fluoro-4-((6-(piperidin-4-oxy)pyridin-2-yl)methoxy)benzonitrile (214 mg, 0.65 mmol) and potassium carbonate (271 mg, 1.95 mmol) in dry N,N-dimethylformamide (7 mL) was stirred at room temperature for 16 hours. After completion, the reaction mixture was quenched with water (10 mL) and extracted with ethyl acetate (3×10 mL). The combined organic layer was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/

methanol=10/1) to give (S)-N-(4-bromo-2-fluoro-6-((oxetan-2-ylmethyl)amino)phenyl)-2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido (300 mg, 71.9% yield). LC-MS m/z : 644 $[M+H]^+$.

2). Synthesis of (S)-4-((6-((1-((6-bromo-4-fluoro-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-2-yl)methyl)piperidin-4-yl)oxy)pyridin-2-yl)methoxy)-3-fluorobenzonitrile






[0497] The mixture of (S)—N-(4-bromo-2-fluoro-6-((oxoalk-2-ylmethyl)amino)phenyl)-2-(4-(((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido (230 mg, 0.36 mmol) in dioxane (5 mL) and acetic acid (0.5 mL) was stirred at 110° C. for 16 hours. The mixture was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain (S)-4-(((1-((6-bromo-4-fluoro-1)-(oxetan-2-ylmethyl)-1H-benzo[d]

imidazol-2-yl)methyl)piperidin-4-yl)oxy)pyridin-2-yl)methoxy)-3-fluorobenzonitrile (80 mg, 35.7% yield). LC-MS m/z : 626 $[M+H]^+$.

3). Synthesis of methyl (S)-2-(((4-(((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-4-fluoro-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate




[0498] To a mixture of (S)-4-((6-((1-((6-bromo-4-fluoro-1-(oxetan-2-ylmethyl))-1H-benzo[d]imidazol-2-yl)methyl)piperidin-4-yl)oxy)pyridin-2-yl)methoxy)-3-fluorobenzonitrile (61 mg, 0.10 mmol) in N,N-dimethylformamide (2 mL) and methanol (2 mL) triethylamine (30 mg, 0.30 mmol) and 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (7 mg, 0.01 mmol) were added. The mixture was stirred at 80° C. for 16 hours under carbon monoxide atmosphere. The mixture was diluted with water (5 mL) and extracted with ethyl acetate (10 mL×2). The combined organic layers were washed with saturated brine (10 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=1/1) to obtain methyl (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-4-fluoro-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (30 mg, 50.8% yield). LC-MS m/z: 604 [M+H]⁺.

4). Synthesis of (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-4-fluoro-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid

zole-6-carboxylate (37 mg, 0.06 mmol) in water (1 mL) and THF (1 mL) lithium hydroxide (14 mg, 0.60 mmol) was added. The reaction mixture was stirred at room temperature for 16 hours. The reaction solution was adjusted to pH=5-6 with formic acid and extracted with ethyl acetate (10 mL×2). The combined organic layers were washed saturated brine (10 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by reverse-phase flash chromatography under the following conditions (Column: Spherical C18, 20-40 m, 120 g; Mobile Phase A: 10 mmol FA in water; Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 30% B-50% B in 20 min; Detector: 254 nm) to obtain (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-4-fluoro-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (6.63 mg, 18.3% yield). LC-MS m/z: 590 [M+H]⁺.

[0500] ¹H NMR (400 MHz, DMSO-d₆): δ 8.09 (s, 1H), 7.89 (dd, J=11.2 Hz, 1.6 Hz 1H), 7.75-7.66 (m, 2H), 7.52-7.43 (m, 2H), 7.06 (d, J=7.6 Hz, 1H), 6.74 (d, J=8.4 Hz, 1H), 5.31 (s, 2H), 5.11-5.06 (m, 1H), 4.95-4.89 (m, 1H), 4.81-4.76 (m, 1H), 4.66-4.63 (m, 1H), 4.52-4.47 (m, 1H),




[0499] To a mixture of methyl (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-4-fluoro-1-(oxetan-2-ylmethyl)-1H-benzo[d]imida-

4.40-4.34 (m, 1H), 3.96-3.92 (m, 1H), 3.81-3.78 (m, 1H), 2.79-2.67 (m, 3H), 2.46-2.41 (m, 1H), 2.33-2.27 (m, 2H), 1.93-1.89 (m, 2H), 1.64-1.57 (m, 2H).

Example 29. Synthesis of (S)-2-((4-((6-((4-chloro-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C8)


1). Synthesis of methyl (S)-2-((4-((6-((4-chloro-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate




[0501] The mixture of 2-((4-((6-((4-chloro-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (144 mg, 0.41 mmol), methyl (S)-2-((4-((6-((4-chloro-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (120 mg, 0.41 mmol) and potassium carbonate (168 mg, 1.23 mmol) in N,N-dimethylformamide (4 mL) was stirred at 60° C. for 3 hours. The mixture was diluted with water (30 mL) and extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (10 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated to give residue in vacuo. The residue was purified by silica gel column chromatog-

raphy (dichloromethane/methanol=20/1) to give methyl (S)-2-((4-((6-((4-chloro-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (80 mg, 32.0% yield). LC-MS m/z: 611 [M+H]⁺.

2). Synthesis of (S)-2-((4-((6-((4-chloro-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid



-continued




[0502] To a mixture of methyl (S)-2-((4-(((6-((4-chloro-2-fluorophenylthio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (80 mg, 0.13 mmol) in water (1.0 mL) and THF (1.0 mL) lithium hydroxide (9 mg, 0.39 mmol) was added. The reaction mixture was stirred at room temperature for 16 hours. The solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain (S)-2-((4-(((6-((4-chloro-2-fluorophenylthio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (20.29 mg, 26.2% yield). LC-MS *m/z*: 597 [M+H]⁺.


[0503] ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.26 (s, 1H), 7.81-7.79 (m, 1H), 7.65-7.61 (m, 3H), 7.28-7.22 (m, 2H), 6.98 (d, *J*=7.2 Hz, 1H), 6.64 (d, *J*=8.0 Hz, 1H), 5.10-5.07 (m, 1H), 4.86 (s, 1H), 4.81-4.75 (m, 1H), 4.65-4.61 (m, 1H), 4.51-4.49 (m, 1H), 4.38-4.34 (m, 1H), 4.30 (s, 2H), 3.93 (d, *J*=13.6 Hz, 1H), 3.77 (d, *J*=13.6 Hz, 1H), 2.75-2.67 (m, 3H), 2.46-2.39 (m, 1H), 2.31-2.26 (m, 2H), 1.88-1.87 (m, 2H), 1.60-1.56 (m, 2H).

Example 30. Synthesis of (S)-2-((4-(((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound 55)

1). Synthesis of methyl (S)-2-((4-(((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate




-continued




[0504] The mixture methyl (S)-2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (150 mg, 0.51 mmol), 3-fluoro-4-((6-(piperidin-4-oxy)pyridin-2-yl)methoxy)benzonitrile (184 mg, 0.56 mmol) and potassium carbonate (176 mg, 1.27 mmol) in N,N-dimethylformamide (8 mL) was stirred at 60° C. for 3 hours. The mixture was quenched with water (10 mL) and extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (10 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chro-

matography (dichloromethane/methanol=20/1) to obtain methyl (S)-2-((4-(((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 33.5% yield). LCMS m/z: 586 [M+H]⁺.

2). Synthesis of (S)-2-((4-(((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid






[0505] To a mixture of methyl (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 0.17 mmol) in THF (2 mL) and water (2 mL) lithium hydroxide (13 mg, 0.51 mmol) was added. The reaction was stirred at room temperature for 16 hours. The resulting mixture was adjusted to pH=5-6 with formic acid. The solvent was removed by concentration to give a residue in vacuo. The residue was eluted by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength 254 nm/214 nm; Flow rate: 20 mL/min) to obtain (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (56.80 mg, 58.6% yield). LC-MS m/z: 572 [M+H]⁺.

[0506] ¹H NMR (400 MHz, DMSO-d₆): δ 8.23 (s, 1H), 7.88 (dd, J=2.0 Hz, 11.6 Hz, 1H), 7.80 (dd, J=1.6, 8.8 Hz,

1H), 7.73 (t, J=8.0 Hz, 1H), 7.67 (d, J=8.4 Hz, 1H), 7.60 (d, J=8.8 Hz, 1H), 7.45 (t, J=8.4 Hz, 1H), 7.06 (d, J=7.6 Hz, 1H), 6.74 (d, J=8.4 Hz, 1H), 5.31 (s, 2H), 5.10-5.08 (m, 1H), 4.92-4.90 (m, 1H), 4.80-4.74 (m, 1H), 4.65-4.61 (m, 1H), 4.50-4.47 (m, 1H), 4.40-4.35 (m, 1H), 3.93 (d, J=13.6 Hz, 1H), 3.78 (d, J=13.6 Hz, 1H), 2.79-2.67 (m, 3H), 2.51-2.41 (m, 1H), 2.32-2.27 (m, 2H), 1.92-1.89 (m, 2H) 1.63-1.58 (m, 2H).


Example 31. Synthesis of 2-((4-((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylic acid (Compound S41)

1). Synthesis of N-(4-bromo-2-((1-(cyanomethyl)cyclopropyl)methyl)amino)-6-fluorophenyl)-2-(4-((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)acetamide




[0507] The mixture of N-(4-bromo-2-((1-(cyanomethyl)cyclopropyl)methyl)amino)-6-fluorophenyl)-2-chloroacetamide (189 mg, 0.50 mmol), 2-((4-chloro-2-fluorobenzyl)oxy)-3-fluoro-6-(piperidin-4-oxypyridine (90 mg, 0.50 mmol) and potassium carbonate (140 mg, 1.00 mmol) in dry N,N-dimethylformamide (4 mL) was stirred at room temperature for 16 hours. When completion, it was quenched with water (10 mL) and extracted with dichloromethane (3×10 mL). The combined organic layer was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to give N-(4-bromo-2-((1-(cyanomethyl)cyclopropyl)methyl)amino)-6-fluorophenyl)-2-((4-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidine-1-yl)methyl)-4-fluoro-1H-benzo[d]imidazol-1-yl)methyl)cyclopropyl)acetamide (218 mg, 62.6% yield). LC-MS m/z : 694 $[M+H]^+$.

-continued




2). Synthesis of 2-(1-((6-bromo-2-((4-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidine-1-yl)methyl)-4-fluoro-1H-benzo[d]imidazol-1-yl)methyl)cyclopropyl)acetoneitrile



[0509] To a mixture of 2-(1-((6-bromo-2-((4-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidine-1-yl)methyl)-4-fluoro-1H-benzo[d]imidazol-1-yl)methyl)cyclopropyl)acetoneitrile (220 mg, 0.33 mmol) in N,N-dimethylformamide (10 mL) and methanol (10 mL) triethylamine (60 mg, 0.6 mmol) and 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (20 mg, 0.03 mmol) were added under carbon monoxide atmosphere. The mixture was stirred at 80° C. for 16 hours, diluted with water (50 mL) and extracted with ethyl acetate (100 mL×2). The combined organic layers were washed with brine (50 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=2/1) to obtain methyl 2-((4-((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidine-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylate (93.7 mg, 43.5% yield). LC-MS m/z : 654 $[M+H]^+$.


[0508] The mixture of N-(4-bromo-2-((1-(cyanomethyl)cyclopropyl)methyl)amino)-6-fluorophenyl)-2-((4-((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidine-1-yl)acetamide (218 mg, 3.17 mmol) in toluene (4 mL) and acetic acid (1 mL) was stirred at 110° C. for 16 hours. The solvent was removed by concentration to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain 2-(1-((6-bromo-2-((4-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidine-1-yl)methyl)-4-fluoro-1H-benzo[d]imidazol-1-yl)methyl)cyclopropyl)acetoneitrile (45 mg, 21.2% yield). LC-MS m/z : 676 $[M+H]^+$.

3). Synthesis of methyl 2-((4-((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidine-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylate



methyl)cyclopropyl)acetoneitrile (220 mg, 0.33 mmol) in N,N-dimethylformamide (10 mL) and methanol (10 mL) triethylamine (60 mg, 0.6 mmol) and 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (20 mg, 0.03 mmol) were added under carbon monoxide atmosphere. The mixture was stirred at 80° C. for 16 hours, diluted with water (50 mL) and extracted with ethyl acetate (100 mL×2). The combined organic layers were washed with brine (50 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=2/1) to obtain methyl 2-((4-((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidine-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylate (93.7 mg, 43.5% yield). LC-MS m/z : 654 $[M+H]^+$.

4). Synthesis of 2-((4-(((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylic acid





[0510] To a mixture of methyl 2-((4-(((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylate (18 mg, 0.03 mmol) in THF (1 mL) and water (1 mL) lithium hydroxide (7 mg, 0.30 mmol) was added. The reaction mixture was stirred at room temperature for 16 hours. The reaction mixture was adjusted to pH=5-6 with formic acid. The solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient eluted ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: mL/min) to obtain 2-((4-(((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidine-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylic acid (4.84 mg, 27.5% yield). LC-MS m/z : 640 $[M+H]^+$.

[0511] ¹H NMR (400 MHz, DMSO- d_6): δ 8.07 (s, 1H), 7.66-7.61 (m, 1H), 7.54-7.47 (m, 3H), 7.32 (dd, J =8.4 Hz, 2.0 Hz 1H), 6.34 (dd, J =8.8 Hz, 2.0 Hz 1H), 5.44 (s, 2H), 4.81-4.76 (m, 1H), 4.60 (s, 2H), 3.86 (s, 2H), 2.81-2.78 (m, 2H), 2.70 (s, 2H), 2.36-2.32 (m, 2H), 1.92-1.90 (m, 2H), 1.64-1.56 (m, 2H), 0.76-0.69 (m, 4H).

Example 32. Synthesis of 2-((4-(((6-((4-chloro-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C9)

1). Synthesis of 4-(2-(4-(((6-((4-chloro-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate






[0512] To a mixture of methyl 4-(2-chloroacetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (190 mg, 0.54 mmol) and 2-((4-chloro-2-fluorophenylthio)methyl)-6-(piperidin-4-oxy)pyridine (190 mg, 0.54 mmol) in N,N-dimethylformamide (5 mL) potassium carbonate (224 mg, 1.62 mmol) was added. The reaction was stirred at room temperature for 16 hours. After completion, the mixture was quenched with water (5 mL) and extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (20 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 4-(2-(4-((6-(((4-chloro-2-fluorophenyl)thio)methyl)pyridine-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (149 mg, 41.5% yield). LC-MS *m/z*: 667 [M+H]⁺.


2). Synthesis of methyl 2-((4-((6-(((4-chloro-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzod[*d*]imidazole-6-carboxylate

[0513] The mixture of methyl 4-(2-(4-((6-(((4-chloro-2-fluorophenyl)thio)methyl)pyridine-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (149 mg, 0.22 mmol) in toluene (6 mL) and acetic acid (1 mL) was stirred at 110° C. for 2 hours. The resulting mixture was poured into water (10 mL) and extracted with ethyl acetate (10 mL×2). The combined organic layers were washed successively with saturated sodium bicarbonate (10 mL×2), brine (10 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=15/1) to obtain methyl 2-((4-((6-(((4-chloro-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzod[*d*]imidazole-6-carboxylate (129 mg, 90.5% yield). LC-MS *m/z*: 649 [M+H]⁺.



3). Synthesis of 2-((4-((6-((4-chloro-2-fluorophenylthio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid

purified by reverse-phase flash chromatography under the following conditions (Column: spherical C18, 20-40 μm , 120 g; Mobile Phase A: 10 mM NH_4OH in water; Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 30%




[0514] To a mixture of methyl 2-((4-((4-chloro-2-fluorophenylthio)methyl)pyridin-2-yl)oxypiperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (129 mg, 0.20 mmol) in THF (4 mL) and water (4 mL) lithium hydroxide monohydrate (47 mg, 1.99 mmol) was added. The reaction was stirred at room temperature for 16 hours. The resulting mixture was adjusted to pH=5-6 with hydrochloric acid (1M) and extracted with ethyl acetate (10 mL $\times$ 3). The combined organic layers were washed with brine (10 mL), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. The residue was

B-50% B in 20 min; Detector: 254 nm) to obtain 2-((4-((6-((4-chloro-2-fluorophenylthio)methyl)pyridine-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (58.36 mg, 0.09 mmol, 46% yield). LC-MS m/z : 635 $[\text{M}+\text{H}]^+$.

[0515] ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.08 (s, 1H), 7.82 (dd, $J=8.4$ Hz, 1.2 Hz, 1H), 7.69-7.67 (m, 2H), 7.64-7.60 (m, 2H), 7.30-7.22 (m, 2H), 6.98 (d, $J=7.2$ Hz, 1H), 6.62 (d, $J=8.4$ Hz, 1H), 6.41 (s, 1H), 5.72 (s, 2H), 4.83-4.78 (m, 1H), 4.29 (s, 2H), 4.03-3.98 (m, 2H), 3.80 (s, 2H), 2.66-2.63 (m, 2H), 2.25-2.21 (m, 2H), 1.74 (s, 2H), 1.45-1.42 (m, 2H), 1.16 (t, $J=7.2$ Hz, 3H).

Example 33. Synthesis of 2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound 43)


1) Synthesis of methyl 4-(2-(4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate




[0516] To a mixture of methyl 4-(2-chloroacetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (190 mg, 0.54 mmol) and 2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridine-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (181 mg, 0.54 mmol) in N,N-dimethylformamide (4 mL) potassium carbonate (224 mg, 1.62 mmol) was added at room temperature. The mixture was stirred for 16 hours at room temperature. After completion, the water (5 mL) was added to quench the reaction mixture, and extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (20 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. The residue was purified

by silica gel column chromatography (dichloromethane/methanol=15/1) to obtain methyl 4-(2-(4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridine-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (241 mg, 0.37 mmol, 68.5% yield). LC-MS m/z : 651 $[\text{M}+\text{H}]^+$.

2). Synthesis of methyl 2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate




-continued




[0517] To a mixture of methyl 4-(2-(4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (241 mg, 0.37 mmol) in toluene (10 mL) acetic acid (1 mL) was added, and the reaction mixture was stirred at 110° C. for 16 hours. After completion, the obtained mixture was poured into water (10 mL) and extracted with ethyl acetate (10 mL×2). The organic layer were washed successively with saturated sodium bicarbonate (10 mL×2) and brine (10 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was

purified by silica gel column chromatography (dichloromethane/methanol=15/1) to obtain methyl 2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (87 mg, 37.8% yield). LC-MS m/z : 633 $[M+H]^+$.

3). Synthesis of 2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid



-continued





[0518] To a mixture of methyl 2-((4-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (87 mg, 0.14 mmol) in THF (3 mL) and water (3 mL) lithium hydroxide monohydrate (56 mg, 1.40 mmol) was added and the reaction was stirred at room temperature for 16 hours. The resulting mixture was adjusted to pH=5-6 with hydrochloric acid (1 N) and extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (10 mL), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. The residue was purified by reverse-phase flash chromatography under the following conditions (Column: Spherical C18, 20-40 m, 120 g; Mobile Phase A: 10 mM NH₄OH in water; Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 30% B-50% B in 20 min; Detector: 254 nm) to obtain 2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridine-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (46.87 mg, 55.4% yield). LC-MS m/z: 619 [M+H]⁺.

[0519] ¹H NMR (400 MHz, DMSO-d₆) δ 8.02 (s, 1H), 7.80 (d, J=8.4 Hz, 1H), 7.72-7.67 (m, 2H), 7.56 (d, J=8.4 Hz, 1H), 7.44 (dd, J=10.8, 2.0 Hz, 1H), 7.27 (t, J=9.2 Hz, 1H), 7.18 (d, J=8.8 Hz, 1H), 7.03 (d, J=7.2 Hz, 1H), 6.70 (d, J=8.4 Hz, 1H), 6.38 (s, 1H), 5.67 (s, 2H), 5.17 (s, 2H), 4.89-4.86 (m, 1H), 4.00 (q, J=7.2 Hz, 2H), 3.78 (s, 2H), 2.67 (s, 2H), 2.24 (t, J=8.0 Hz, 2H), 1.79 (d, J=9.6 Hz, 2H), 1.46 (m, 2H), 1.17 (t, J=7.2 Hz, 3H).

Example 34. Synthesis of (S)-2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound 49)

1). Synthesis of methyl (S)-2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate






[0520] The mixture of 2-((4-chloro-2-fluorophenoxy)methyl)-6-(piperidin-4-oxy)pyridine (115 mg, 0.34 mmol), (S)-2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 0.34 mmol) and potassium carbonate (140 mg, 1.02 mmol) in N,N-dimethylformamide (4 mL) was stirred at 60° C. for 16 hours. After completion, the resulting mixture was diluted with water (40 mL) and extracted with ethyl acetate (20 mL×2). The organic layers were washed with brine (20 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. The residue was purified by silica gel column chro-

matography (dichloromethane/methanol=20/1) to obtain methyl (S)-2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (110 mg, 54.5% yield). LC-MS m/z: 595 [M+H]⁺.


2). Synthesis of (S)-2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate



[0521] To a mixture of methyl (S)-2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (110 mg, 0.19 mmol) in THF (2 mL) and water (2 mL) lithium hydroxide (9 mg, 0.38 mmol) was added, and the reaction mixture was stirred at room temperature for 16 hours. After completion, the solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20

Example 35. Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylic acid (Compound 2)

1). Synthesis of methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)-5-fluorobenzoate




mL/min) to obtain (S)-2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (52.13 mg, 47.2% yield). LC-MS m/z: 581 [M+H]⁺.

[0522] ¹H NMR (400 MHz, DMSO-d₆): δ 8.27 (s, 1H), 7.80 (dd, J=8.4 Hz, 1.2 Hz, 1H), 7.72 (t, J=7.6 Hz, 1H), 7.64 (d, J=8.4 Hz, 1H), 7.44 (dd, J=11.2 Hz, 2.0 Hz, 1H), 7.28 (t, J=8.8 Hz, 1H), 7.18 (d, J=8.4 Hz, 1H), 7.04 (d, J=7.2 Hz, 1H), 6.72 (d, J=8.0 Hz, 1H), 5.18 (s, 2H), 5.12-5.06 (m, 1H), 4.95-4.93 (m, 1H), 4.81-4.76 (m, 1H), 4.66-4.62 (m, 1H), 4.51-4.49 (m, 1H), 4.38-4.36 (m, 1H), 3.94 (d, J=13.6 Hz, 1H), 3.78 (d, J=13.6 Hz, 1H), 2.79-2.67 (m, 2H), 2.46-2.41 (m, 1H), 2.32 (s, 2H), 1.92-1.91 (m, 2H), 1.63-1.59 (m, 2H).

[0523] The mixture of methyl 4-(2-chloroacetamido)-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)-5-fluorobenzoate (130 mg, 0.35 mmol), 3-fluoro-4-((6-(piperidin-4-oxypyridin-2-yl)methoxy)benzonitrile (115 mg, 0.35 mmol) and potassium carbonate (146 mg, 1.05 mmol) in N,N-dimethylformamide (5 mL) was stirred at room temperature for 16 hours. The resulting mixture was diluted with water (50 mL) and extracted with ethyl acetate (10 mL $\times$ 3). The organic layer were washed with brine (10 mL $\times$ 3), and concentrated to give a residue in vacuo. The residue was purified by column chromatography (dichloromethane/methanol=20/1) to obtain methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-(((1-ethyl-1H-imidazol-5-yl)methyl)amino)-5-fluorobenzoate (70 mg, 30.3% yield). LC-MS m/z: 660 [M+H]⁺.


2). Synthesis of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylate




[0524] The mixture of methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)-5-fluorobenzoate (70 mg, 0.11 mmol) in toluene (2 mL) and acetic acid (0.5 mL) was stirred at 110° C. for 3 hours. After completion, The mixture was diluted with water (20 mL) and extracted with dichloromethane (2×25 mL). The organic phase was concentrated to give methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)

methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylate (40 mg, 56.7% yield). LC-MS m/z: 642 [M+H]⁺.

3). Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylic acid






[0525] To a mixture of methyl 2-((4-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-4-fluoro-H-benzo[d]imidazole-6-carboxylate (40 mg, 0.062 mmol) in THF (4 mL) and water (4 mL) lithium hydroxide (4 mg) was added, and the reaction was stirred at room temperature for 16 hours. The solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidine-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylic acid (8.49 mg, 22.6% yield). LC-MS *m/z*: 628 [M+H]⁺.

[0526] ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.20 (s, 1H), 7.93-7.86 (m, 2H), 7.74-7.65 (m, 3H), 7.52 (d, *J*=11.6 Hz

1H), 7.43 (t, *J*=8.4 Hz 1H), 7.04 (d, *J*=7.2 Hz 1H), 6.72 (d, *J*=8.0 Hz 1H), 6.42 (s, 1H), 5.74 (s, 2H), 5.30 (s, 2H), 4.86-4.84 (m, 1H), 4.03-3.98 (m, 2H), 3.83 (s, 2H), 2.67 (s, 2H), 2.27-2.22 (m, 2H), 1.79-1.76 (m, 2H), 1.46-1.44 (m, 2H), 1.17 (t, *J*=7.2 Hz, 3H).

Example 36. Synthesis of (S)-2-((4-((6-((4-cyano-2-fluorophenylthio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C10)

1). Synthesis of methyl (S)-2-((4-((6-((4-cyano-2-fluorophenylthio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate




[0527] The mixture of 3-fluoro-4-((6-(piperidin-4-oxy)pyridin-2-yl)methylthio)benzonitrile (152 mg, 0.44 mmol), methyl (S)-2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 0.34 mmol) and potassium carbonate (140 mg, 1.02 mmol) in N,N-dimethylformamide (4 mL) was stirred at 60° C. for 3 hours. The resulting mixture was diluted with water (30 mL) and extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (10 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to give methyl (S)-2-((4-((6-((4-cyano-2-fluorophenylthio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (150 mg, 56.6% yield). LC-MS *m/z*: 602 [M+H]⁺.

2). Synthesis of (S)-2-((4-((6-((4-cyano-2-fluorophenylthio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid


[0528] To a mixture of methyl (S)-2-((4-((6-((4-cyano-2-fluorophenylthio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (150 mg, 0.25 mmol) in THF (2 mL) and water (2 mL) lithium hydroxide (10 mg, 0.75 mmol) was added. The mixture was stirred at room temperature for 16 hours. The solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μm 19×250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: mL/min) to obtain (S)-2-((4-((6-((4-cyano-2-fluorophenylthio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (40.33 mg, 27.4% yield). LC-MS *m/z*: 588 [M+H]⁺.

[0529] ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.26 (s, 1H), 7.84-7.75 (m, 3H), 7.68-7.62 (m, 3H), 7.02 (d, *J*=7.2 Hz, 1H), 6.65 (d, *J*=8.4 Hz, 1H), 5.10-5.08 (m, 1H), 4.81-4.75 (m, 2H), 4.66-4.61 (m, 1H), 4.53-4.47 (m, 1H), 4.41 (s, 2H), 4.39-4.36 (m, 1H), 3.93 (d, *J*=13.6, 1H), 3.77 (d, *J*=13.6 Hz, 1H), 2.77-2.67 (m, 4H), 2.46-2.41 (m, 1H), 2.28-2.23 (m, 2H), 1.91-1.86 (m, 2H), 1.59-1.55 (m, 2H).



Example 37. Synthesis of 2-((4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)yl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylic acid (Compound S6)


1). Synthesis of N-(4-bromo-2-(((1-(cyanomethyl)cyclopropyl)methyl)amino)-6-fluorophenyl)-2-(4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)piperidin-1-yl)acetamido




[0530] To a mixture of 3-fluoro-4-((3-(piperidin-4-oxymethyl)benzonitrile (130 mg, 0.40 mmol) and N-(4-bromo-2-(((1-(cyanomethyl)cyclopropyl)methyl)amino)-6-fluorophenyl)-2-chloroacetamido (149 mg, 0.40 mmol) in N,N-dimethylformamide (4 mL) K_2CO_3 (165 mg, 1.20 mmol) was added. The mixture was stirred at 60° C. for 2 hours. The mixture was diluted with water (5 mL) and extracted with EA (2×30 mL). The combined organic phases were washed with brine (20 mL), dried over anhydrous Na_2SO_4 , filtered, and concentrated in vacuo to give N-(4-

bromo-2-(((1-(cyanomethyl)cyclopropyl)methyl)amino)-6-fluorophenyl)-2-(4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)piperidin-1-yl)acetamido (180 mg, 67.9% yield). LC-MS m/z: 666, 664 $[M+H]^+$.

2). Synthesis of 4-((3-((1-((6-bromo-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-2-yl)methyl)piperidin-4-yl)oxy)phenoxy)methyl)-3-fluorobenzonitrile




-continued



[0531] To a mixture of N-(4-bromo-2-((1-(cyanomethyl)cyclopropyl)methyl)amino)-6-fluorophenyl)-2-(4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)piperidin-1-yl)acetamido (160 mg, 0.24 mmol) in toluene (6 mL) AcOH (1 mL) was added, and the mixture was stirred at 110° C. for 32 hours. The resulting mixture was poured into H₂O (10 mL) and extracted with EA (2×10 mL). The combined organic layers were sequentially washed with saturated NaHCO₃ (2×10 mL), brine (10 mL×2), dried over anhydrous Na₂SO₄, filtered, and concentrated to give residue in vacuo. The residue was purified by reverse-phase column chromatography (Column: Spherical C18 column, 20-40 μm, 120 g; Mobile Phase A: 10 mM NH₄OH in water; Mobile Phase B:

Acetonitrile; Flow Rate: 80 mL/min; Gradient: 60% B-80% B in 20 min; Detector: 254 nm). The fractions containing the desired product were collected at 69% B and concentrated in vacuo to obtain 4-((3-((1-((6-bromo-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazol-2-yl)methyl)piperidin-4-yl)oxy)phenoxy)methyl)-3-fluorobenzonitrile (142 mg, 91.3% yield). LC-MS m/z: 646, 648 [M+H]⁺.

3). Synthesis of methyl 2-((4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)-cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylate




[0532] To a mixture of 4-((3-((1-((6-bromo-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazol-2-yl)methyl)piperidin-4-yl)oxy)phenoxy)methyl)-3-fluorobenzonitrile (130 mg, 0.20 mmol) in methanol (5 mL)) KOAc (59 mg, 0.60 mmol) and Pd(dppf)Cl₂ (15 mg, 0.02 mmol) were added at room temperature under carbon monoxide atmosphere. The mixture was stirred at 90° C. for 16 hours. The mixture was diluted with water (15 mL) and extract with ethyl acetate (3×30 mL). The combined organic layers were washed with brine (30 mL), dried over anhydrous Na₂SO₄, filtered, and concentrated to give a residue in vacuo. The residue was purified by reverse phase column chromatography (Column: Spherical C18 column, 20-40 um, 120 g; Mobile Phase A: 10 mM NH₄OH in water; Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 60% B-80% B in 20 min; Detector: 254 nm) to obtain methyl 2-((4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylate (60 mg, 47.5% yield). LC-MS m/z: 626 [M+H]⁺.

4). Synthesis of 2-((4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylic acid


[0533] To a mixture of methyl 2-((4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylate (60 mg, 0.10 mmol) in THF (2 mL) and H₂O (2 mL) lithium hydroxide monohydrate (40 mg, 1.00 mmol) was added and stirred at 40° C. for 3 hours. The resulting mixture was adjusted to pH=5-6 with HCl (1 M) and extracted with EA (3×10 mL). The combined organic layers were washed with brine (10 mL), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by reverse-phase flash chromatography under the following conditions (Column: Spherical C18, 20-40 m, 40 g; Mobile Phase A: 10 mM NH₄OH in water; Mobile Phase B: acetonitrile; Flow rate: 40 ml/min; Gradient: 20% B-50% B in 20 min; Detection Wavelength: 254 nm) to obtain 2-((4-(3-((4-cyano-2-fluorobenzyl)oxy)phenoxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylic acid (26.23 mg, 43.5% yield). LC-MS m/z: 612 [M+H]⁺.

[0534] ¹H NMR (400 MHz, DMSO-d₆) δ 8.04 (s, 1H), 7.91 (d, J=10.4 Hz, 1H), 7.75 (d, J=3.2 Hz, 2H), 7.49 (d, J=11.2 Hz, 1H), 7.17 (t, J=8.0 Hz, 1H), 6.62-6.57 (m, 3H), 5.21 (s, 2H), 4.60 (s, 2H), 4.42-4.40 (m, 1H), 3.85 (s, 2H), 2.79-2.77 (m, 2H), 2.70 (s, 2H), 2.38 (t, J=10.0 Hz, 2H), 1.93 (d, J=10.8 Hz, 2H), 1.64-1.60 (m, 2H), 0.75-0.67 (m, 4H).



Example 38. Synthesis of 2-((4-((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C11)


1). Synthesis of methyl 4-(2-(4-((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate




[0535] The mixture of 2-((4-((6-((4-chloro-2-fluorobenzyl)oxy)-3-fluoro-6-(piperidin-4-oxy)pyridine (175 mg, 0.49 mmol), methyl 4-(2-chloroacetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (184 mg, 0.49 mmol) and potassium carbonate (144 mg, 1.04 mmol) in N,N-dimethylformamide (5 mL) was stirred at room temperature for 16 hours. The resulting mixture was diluted with water (50 mL) and extracted with ethyl acetate (10 mL $\times$ 3). The combined organic layers were washed with brine (10 mL $\times$ 3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica

gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 4-(2-(4-((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (180 mg, 51.82% yield). LC-MS m/z : 669 $[M+H]^+$.

2). Synthesis of methyl 2-((4-((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate




-continued



[0536] The mixture of methyl 4-(2-((4-((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (180 mg, 0.27 mmol) in toluene (3 mL) and acetic acid (1 mL) was stirred at 110° C. for 2 hours. The resulting mixture was poured into brine (20 mL) and extracted with dichloromethane (2×10 mL). The combined organic layer was concentrated to give a residue in vacuo to obtain methyl 2-((4-((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-

yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (140 mg, 77.9% yield). LC-MS m/z : 651 $[M+H]^+$.

3). Synthesis of 2-((4-((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid




[0537] To a mixture of methyl 2-((4-((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (70 mg, 0.11 mmol) in THF (4 mL) and water (4 mL) lithium hydroxide (8 mg, 0.33 mmol) was added. The mixture was stirred at room temperature for 16 hours. The solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength 254 nm/214 nm; Flow rate 20 mL/min) to give 2-((4-((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidine-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (37.86 mg, 54.1% yield). LC-MS m/z: 637 [M+H]⁺.

[0538] ¹H NMR (400 MHz, DMSO-d₆): δ 8.08 (s, 1H), 7.82 (d, J=8.4 Hz, 1H), 7.69-7.60 (m, 3H), 7.53-7.46 (m, 2H), 7.31 (dd, J=8.4 Hz, 2.0 Hz, 1H), 6.42 (s, 1H), 6.33 (dd, J=8.8 Hz, 2.0 Hz, 1H), 5.72 (s, 2H), 5.43 (s, 2H), 4.74-4.71 (m, 1H), 4.03-3.98 (m, 2H), 3.82 (s, 2H), 2.69-2.67 (m, 2H), 2.29-2.24 (m, 2H), 1.79-1.77 (m, 2H), 1.46-1.42 (m, 2H), 1.16 (t, J=7.2 Hz, 3H).


Example 39. Synthesis of 2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound S66)

1). Synthesis of 2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate

[0539] To a mixture of methyl 2-((4-((6-((4-chloro-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (132 mg, 0.20 mmol) and zinc cyanide (234 mg, 2.00 mmol) in N-methylpyrrolidone (2 mL) methanesulfonato (2-dicyclohexylphosphino-2',6'-di-1-propoxy-1,1'-biphenyl)(2'-amino-1,1'-biphenyl-2-yl)palladium(II) (17 mg, 0.02 mmol) and 2-dicyclohexylphosphine-2',4',6'-triisopropylbiphenyl (10 mg, 0.02 mmol) were added. The reaction mixture was stirred at 130° C. for 3 hours. After completion, the mixture was diluted with water (10 mL) and extracted with ethyl acetate (10 mL $\times$ 2). The combined organic layers were washed with brine (10 mL $\times$ 2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by pre-TLC chromatography (dichloromethane/methanol=20/1) to give methyl 2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)-5-fluoropyridine-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (40 mg, 31.2% yield). LC-MS m/z: 642 [M+H]⁺.



2). Synthesis of 2-((4-(((6-((4-cyano-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid





[0540] To a mixture of methyl 2-((4-(((6-((4-cyano-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (40 mg, 0.06 mmol) in THF (4 mL) and water (4 mL) lithium hydroxide (4 mg, 0.17 mmol) was added. The mixture was stirred at room temperature for 16 hours. The solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength 254 nm/214 nm; Flow rate: 20 mL/min) to give 2-((4-(((6-((4-cyano-2-fluorobenzyl)oxy)-5-fluoropyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (6.35 mg, 16.9% yield). LC-MS *m/z*: 628 [M+H]⁺.

[0541] ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.07 (s, 1H), 7.93 (dd, *J*=9.6 Hz, 0.8 Hz, 1H), 7.81 (dd, *J*=8.4 Hz, 1.2 Hz, 1H), 7.73-7.62 (m, 5H), 6.40 (s, 1H), 6.34 (dd, *J*=8.4 Hz, 2.0 Hz, 1H), 5.71 (s, 2H), 5.52 (s, 2H), 4.65-4.63 (m, 1H), 4.03-3.97 (m, 2H), 3.81 (s, 2H), 2.67-2.65 (m, 2H), 2.25-2.20 (m, 2H), 1.74-1.71 (m, 2H), 1.43-1.36 (m, 2H), 1.16 (t, *J*=7.2 Hz, 3H).

Example 40. Synthesis of 2-((4-(((6-((4-cyano-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C12)

1). Synthesis of methyl 4-(2-(4-(((6-((4-cyano-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate






[0542] To a solution of methyl 4-(2-chloroacetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (176 mg, 0.50 mmol) in N,N-dimethylformamide (5 mL) potassium carbonate (138 mg, 1.00 mmol) and 3-fluoro-4-((6-(piperidin-4-oxy)pyridin-2-yl)methyl)sulfanyl)benzonitrile (223 mg, 0.65 mmol). The mixture was stirred at room temperature for 3 hours. After completion, water (10 mL) was added to quench the reaction, and then it was extracted with ethyl acetate (15 mL×2). The combined organic layers were washed with brine (15 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 4-(2-((4-((6-((4-cyano-2-fluorophenylthio)-methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (141 mg, 42.9% yield). LC-MS m/z : 658 $[M+H]^+$.


2). Synthesis of methyl 2-((4-((6-((4-cyano-2-fluorophenylthio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate

[0543] To a mixture of 4-(2-((4-((6-((4-cyano-2-fluorophenylthio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (141 mg, 0.21 mmol) in dioxane (5 mL) acetic acid (0.5 mL) was added at room temperature. The mixture was stirred at 100° C. for 2 hours. After completion, the mixture was diluted with water (15 mL) and extracted with ethyl acetate (20 mL×2). The combined organic layers were washed with brine (20 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 2-((4-((6-((4-cyano-2-fluorophenylthio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (130 mg, 96.7% yield). LC-MS m/z : 640 $[M+H]^+$.



3). Synthesis of 2-((4-((6-(((4-cyano-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid

reverse-phase flash chromatography under the following conditions (Column: Spherical C18, 20-40 μm , 40 g; Mobile Phase A: 10 mM NH_4OH in water; Mobile Phase B: acetonitrile; Flow rate: 40 mL/min; Gradient: 20% B-50% B




[0544] To a mixture of methyl 2-((4-((6-(((4-cyano-2-fluorophenyl)thio)methyl)pyridine-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (130 mg, 0.20 mmol) in THF (2 mL) and water (2 mL) lithium hydroxide monohydrate (85 mg, 2.00 mmol) was added at room temperature. The mixture was stirred at room temperature for 5 hours. After completion, the mixture was adjusted to pH=5-6 with formic acid, and extracted with ethyl acetate (10 mL $\times$ 3). The combined organic layers were washed with brine (10 mL), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by

in 20 min; Detector: 254 nm) to obtain 2-((4-((6-(((4-cyano-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (29.92 mg, 23.9% yield). LC-MS m/z : 626 $[\text{M}+\text{H}]^+$.

[0545] ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.07 (s, 1H), 7.83 (d, $J=8.8$ Hz, 2H), 7.76 (t, $J=8.0$ Hz, 1H), 7.69-7.61 (m, 4H), 7.02 (d, $J=7.2$ Hz, 1H), 6.64 (d, $J=8.4$ Hz, 1H), 6.41 (s, 1H), 5.71 (s, 2H), 4.75-4.74 (m, 1H), 4.40 (s, 2H), 4.04-3.98 (m, 2H), 3.80 (s, 2H), 2.67-2.64 (m, 2H), 2.21-2.17 (m, 2H), 1.76-1.74 (m, 2H), 1.44-1.41 (m, 2H), 1.16 (t, $J=7.2$ Hz, 3H).

Example 41. Synthesis of 2-((4-((6-(((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylic acid (Compound S20)


1). Synthesis of N-(4-bromo-2-(((1-(cyanomethyl)cyclopropyl)methyl)amino)-6-fluorophenyl)-2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)acetamide




[0546] To a solution of N-(4-bromo-2-(((1-(cyanomethyl)cyclopropyl)methyl)amino)-6-fluorophenyl)-2-chloroacetamide (100 mg, 0.27 mmol) in N,N-dimethylformamide (5 mL) 3-fluoro-4-((6-((piperidin-4-yl)oxy)pyridin-2-yl)oxy)methyl)benzonitrile (88 mg, 0.27 mmol) and potassium carbonate (112 mg, 0.81 mmol) was added at room temperature. The mixture was stirred at room temperature for 16 hours. LCMS indicated the mixture was completed. The reaction was quenched by adding water (5 mL) and extracted with ethyl acetate (20 mL $\times$ 3). The combined organic layers were washed with brine (20 mL $\times$ 2), dried

over anhydrous sodium sulfate, filtered, and concentrated to give N-(4-bromo-2-(((1-(cyano-methyl)cyclopropyl)methyl)amino)-6-fluorophenyl)-2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)-piperidine-1-yl)acetamide (120 mg, 66.7% yield). LC-MS m/z : 665, 667 $[M+H]^+$.

2). Synthesis of 4-((6-((1-((6-bromo-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-2-yl)methyl)piperidin-4-yl)oxy)pyridin-2-yl)oxy)methyl)-3-fluorobenzonitrile







[0547] To a mixture of N-(4-bromo-2-((1-(cyanomethyl)cyclopropyl)methyl)amino)-6-fluorophenyl)-2-(4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)acetamide (120 mg, 0.18 mmol) in toluene (6 mL) acetic acid (1 mL) was added. The mixture was stirred at 120° C. for overnight. After completion, the mixture was poured into water (10 mL) and extracted with ethyl acetate (10 mL×2). The combined organic layer was washed with saturated sodium bicarbonate (10 mL×2) and brine (10 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by reverse-phase flash chromatography under the following conditions (Column: Spherical C18, 20-40 m, 120 g; Mobile Phase A: 10 mM NH₄OH in water; Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 60% B-80% B in 20 min; Detector: 254 nm) to obtain 4-((6-((1-(6-bromo-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazol-2-yl)methyl)piperidin-4-yl)oxy)pyridin-2-yl)oxy)methyl)-3-fluorobenzonitrile (100 mg, 83.3% yield). LC-MS m/z: 647, 649 [M+H]⁺.

3). Synthesis of methyl 2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylate

[0548] To a solution of 4-((6-((1-(6-bromo-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazol-2-yl)methyl)piperidin-4-yl)oxy)pyridin-2-yl)oxy)methyl)-3-fluorobenzonitrile (100 mg, 0.15 mmol) in methanol/N,N-dimethylformamide (5 mL/5 mL) triethylamine (45 mg, 0.45 mmol) and 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (15 mg, 0.02 mmol) were added at room temperature. The mixture was stirred at 90° C. for 16 hours under carbon monoxide atmosphere. After completion, water (15 mL) was added to quench the reaction. The mixture was extracted with ethyl acetate (30 mL×3). The combined organic layers were washed with brine (30 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated under reduced pressure. The residue was purified by reverse-phase flash chromatography under the following conditions (Column: Spherical C18, 20-40 m, 120 g; Mobile Phase A: water 10 mM NH₄OH in water; Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 60% B-80% B in 20 min; Detector: 254 nm) to obtain methyl 2-((4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylate (60 mg, 66.7% yield). LC-MS m/z: 627 [M+H]⁺.



4). Synthesis of 2-((4-(((6-((4-cyano-2-fluorobenzyl)oxy)pyridine-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyano-methyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylic acid




[0549] To a mixture of methyl 2-((4-(((6-((4-cyano-2-fluorobenzyl)oxy)pyridine-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylate (60 mg, 0.10 mmol) in THF (3 mL) and water (3 mL) lithium hydroxide monohydrate (32 mg, 1.0 mmol) was added. The mixture was stirred at room temperature for 5 hours. After completion, the resulting mixture was adjusted to pH=5-6 with hydrochloric acid (1 M) and extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (10 mL), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by reverse-phase flash chromatography under the following conditions (Column: Spherical C18, 20-40 μ m, 120 g; Mobile Phase A: 10 mM NH_4OH in water; Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 30% B-50% B in 20 min; Detector: 254 nm) to obtain 2-((4-(((6-((4-cyano-2-fluorobenzyl)oxy)pyridine-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyano-methyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylic acid (19.01 mg, 30% yield). LC-MS m/z : 613 $[\text{M}+\text{H}]^+$.


[0550] ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 8.10 (s, 1H), 7.91 (dd, $J=10.4, 1.6$ Hz, 1H), 7.70 (dd, $J=8.0, 1.2$ Hz, 1H), 7.63 (t, $J=8.0$ Hz, 2H), 7.53 (d, $J=11.2$ Hz, 1H), 6.45 (d, $J=8.0$ Hz, 1H), 6.35 (d, $J=8.0$ Hz, 1H), 5.45 (s, 2H), 4.81-4.77 (m, 1H), 4.61 (s, 2H), 3.87 (s, 2H), 2.80-2.77 (m, 2H), 2.70 (s, 2H), 2.30 (t, $J=9.6$ Hz, 2H), 1.87 (d, $J=9.6$ Hz, 2H), 1.60-1.54 (m, 2H), 0.77-0.68 (m, 4H).

Example 42. Synthesis of 2-((4-(((6-((4-cyano-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C13)

1). Synthesis of methyl 4-(2-(4-(((6-((4-cyano-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)benzoate




-continued




[0551] The mixture of methyl 4-(2-chloroacetamido)-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)benzoate (210 mg, 0.63 mmol), 3-fluoro-4-((6-(piperidin-4-oxy)pyridin-2-yl)methyl)thio)benzonitrile (214 mg, 0.63 mmol) and potassium carbonate (259 mg, 1.89 mmol) in N,N-dimethylformamide (5 mL) was stirred at room temperature for 16 hours. After completion, the resulting mixture was diluted with water (40 mL) and extracted with ethyl acetate (20 mL×2). The combined organic layers were washed brine (20 mL×2) and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 4-(2-(4-(((6-(((4-cyano-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)amino)-benzoate (300 mg, 28.6% yield). LC-MS m/z : 643[M+H]⁺.

2). Synthesis of methyl 2-((4-(((6-(((4-cyano-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylate

[0552] The mixture of methyl 4-(2-(4-(((6-(((4-cyano-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)amino)-benzoate (300 mg, 0.47 mmol) in toluene (5 mL) and acetic acid (0.5 mL) was stirred at 110° C. for 3 hours. After completion, the resulting mixture was diluted with water (60 mL) and extracted with ethyl acetate (30 mL×2). The combined organic layers were washed with brine (30 mL×2), and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 2-((4-(((6-(((4-cyano-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylate (200 mg, 68.1% yield). LC-MS m/z : 625 [M+H]⁺.



3). Synthesis of 2-((4-((6-(((4-cyano-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid




[0553] To a mixture of methyl 2-((4-((6-(((4-cyano-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylate (200 mg, 0.33 mmol) in THF (2 mL) and water (2 mL) lithium hydroxide (24 mg, 0.99 mmol) was added at room temperature. The mixture was stirred at room temperature for 16 hours. After completion, the solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19x250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) purification to obtain 2-((4-((6-(((4-cyano-2-fluorophenyl)thio)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid formic salt (51.79 mg, 25.7% yield). LC-MS m/z: 611[M+H]⁺.


[0554] ¹H NMR (400 MHz, DMSO-d₆): δ 8.25 (s, 1H), 8.19 (s, 1H), 7.84-7.75 (m, 3H), 7.68-7.62 (m, 3H), 7.02 (d, J=6.8 Hz, 1H), 6.65 (d, J=8.0 Hz, 1H), 4.83-4.79 (m, 1H), 4.60 (s, 2H), 4.41 (s, 2H), 3.85 (s, 2H), 2.79-2.76 (m, 2H), 2.68 (s, 2H), 2.29-2.24 (m, 2H), 1.89-1.87 (m, 2H), 1.61-1.55 (m, 2H), 0.75-0.68 (m, 4H).

Example 43. Synthesis of 1-((1-ethyl-1H-imidazol-5-yl)methyl)-2-((4-((4-ethynyl-2-fluorophenoxy)methyl)pyridin-2-yl)oxypiperidin-1-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C14)

1). Synthesis of 1-((1-ethyl-1H-imidazol-5-yl)methyl)-2-((4-((6-((2-fluoro-4-((trimethylsilyl)ethynyl)benzene)methyl)oxy)methyl)pyridin-2-yl)oxypiperidin-1-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate




-continued



[0555] The mixture of methyl 2-((4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (240 mg, 0.35 mmol), ethynyltrimethylsilane (343 mg, 3.50 mmol), 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (26 mg, 0.035 mmol) and triethylamine (354 mg, 3.50 mmol) was stirred at 90° C. for 2 hours. After the reaction mixture was completed, it was diluted with water (10 mL) and extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (20 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo.

The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 1-((1-ethyl-1H-imidazol-5-yl)methyl)-2-((4-((6-((2-fluoro-4-((trimethylsilyl)ethynyl)phenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (190 mg, 77.2% yield). LC-MS m/z : 695[M+H]⁺.


2). Synthesis of 1-((1-ethyl-1H-imidazol-5-yl)methyl)-2-((4-((4-ethynyl-2-fluorophenoxy)methyl)pyridin-2-yl)oxypiperidin-1-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid



[0556] To a mixture of methyl 1-((1-ethyl-1H-imidazol-5-yl)methyl)-2-((4-((6-((2-fluoro-4-((trimethylsilyl) ethynyl)phenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (160 mg, 0.23 mmol) in methanol (1.0 mL), water (1.0 mL) and THF (1.0 mL) lithium hydroxide (44 mg, 1.84 mmol) was added. The reaction mixture was stirred at room temperature for 5 hours. After the completion of the reaction, the reaction mixture was adjusted to pH=5-6 by with formic acid. The

Example 44. Synthesis of (S)-2-((4-((6-((2-fluoro-4-((trimethylsilyl)ethynyl)phenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C15)

1). Synthesis of methyl (S)-2-((4-((6-((2-fluoro-4-((trimethylsilyl)ethynyl)phenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate




solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 um 19x250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to give 1-((1-ethyl-1H-imidazol-5-yl)methyl)-2-((4-((4-ethynyl-2-fluorophenoxy)methyl)pyridin-2-yl)oxypiperidin-1-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (14.25 mg, 10.2% yield). LC-MS m/z: 609 [M+H]⁺.

[0557] ¹H NMR (400 MHz, DMSO-d₆): δ 8.06 (s, 1H), 7.81 (dd, J=8.4 Hz, J=1.2 Hz, 1H), 7.73-7.65 (m, 3H), 7.39-7.36 (m, 1H), 7.24-7.23 (m, 2H), 7.03 (d, J=7.2 Hz, 1H), 6.71 (d, J=8.0 Hz, 1H), 6.41 (s, 1H), 5.71 (s, 2H), 5.20 (s, 2H), 4.92-4.84 (m, 1H), 4.11 (s, 1H), 4.01-3.99 (m, 2H), 3.81 (s, 2H), 2.68-2.66 (m, 2H), 2.26-2.20 (m, 2H), 1.83-1.75 (m, 2H), 1.52-1.42 (m, 2H), 1.16 (t, J=7.2 Hz, 3H).

[0558] The mixture of ethynyltrimethylsilane (304 mg, 3.10 mmol), triethylamine (313 mg, 3.10 mmol) and 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (22 mg, 0.03 mmol) and methyl (S)-2-((4-((6-((2-fluoro-4-((trimethylsilyl)ethynyl)phenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (200 mg, 0.31 mmol) in DMSO (6 mL) was stirred at 90° C. for 2 hours. After completion, the reaction was quenched with water (10 mL) at room temperature, and extracted with ethyl acetate (10 mLx3). The combined organic layers were washed with brine (10 mLx2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl (S)-2-((4-((6-((2-fluoro-4-((trimethylsilyl)ethynyl)phenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (260 mg, crude). LC-MS m/z: 657 [M+H]⁺.

2). Synthesis of (S)-2-((4-((6-((4-ethynyl-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid





[0559] To a mixture of methyl (S)-2-((4-((6-((2-fluoro-4-((trimethylsilyl)ethynyl)phenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (210 mg, 0.32 mmol) in water (1 mL), THF (1 mL) and methanol (1 mL) lithium hydroxide (61 mg, 2.56 mmol) was added. The reaction mixture was stirred at room temperature for 5 hours. After the completion of the reaction, the reaction mixture was adjusted to pH=5-6 with formic acid. The solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: mL/min) to obtain (S)-2-((4-((6-((4-ethynyl-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (10.76 mg, 5.9% yield). LC-MS m/z: 571 [M+H]⁺.

[0560] ¹H NMR (400 MHz, DMSO-d₆): δ 8.27 (s, 1H), 7.80 (d, J=8.0 Hz, 1H), 7.74-7.70 (m, 1H), 7.64 (d, J=8.8 Hz, 1H), 7.38 (d, J=11.2 Hz, 1H), 7.25 (s, 2H), 7.05 (d, J=6.8 Hz, 1H), 6.73 (d, J=8.0 Hz, 1H), 5.21 (s, 2H), 5.10-5.08 (m, 1H), 4.96-4.94 (m, 1H), 4.79-4.75 (m, 1H), 4.64 (d, J=15.2 Hz, 1H), 4.52-4.49 (m, 1H), 4.38-4.36 (m, 1H), 4.13 (d, J=2.8 Hz, 1H), 3.94 (d, J=14.0 Hz, 1H), 3.78 (d, J=13.2 Hz, 1H), 2.79-2.69 (m, 3H), 2.44-2.32 (m, 3H), 1.93-1.87 (m, 2H), 1.63-1.61 (m, 2H).

Example 45. Synthesis of 2-((4-((3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound 13)

1). Synthesis of methyl 4-(2-(4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)acetamido)-3-((1-ethyl)-1H-imidazol-5-yl)methyl)amino)benzoate






[0561] To a solution of methyl 4-(2-chloroacetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (190 mg, 0.54 mmol) in N,N-dimethylformamide (5 mL) 3-fluoro-4-((3-(piperidin-4-oxybenzyl)oxy)benzonitrile (196 mg, 0.60 mmol) and potassium carbonate (149 mg, 1.08 mmol) were added at room temperature. The mixture was stirred at room temperature for 16 hours. After completion, the mixture was diluted with water (20 mL) and extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (20 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 4-(2-(4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (60 mg, 17.4% yield). LC-MS m/z: 641[M+H]⁺.


2). Synthesis of methyl 2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate

[0562] To a mixture of methyl 4-(2-(4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (60 mg, 0.09 mmol,) in dioxane (5 mL) acetic acid (0.5 mL) was added. The resulting mixture was stirred and reacted at 100° C. for 3 hours. After completion, the mixture was diluted with water (20 mL) and extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (20 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (50 mg, 89.2% yield). LC-MS m/z: 623[M+H]⁺.



3). Synthesis of 2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid

phase flash chromatography under the following condition (Column: Spherical C18, 20-40 μ m, 40 g; Mobile Phase A: 10 mM NH_4OH in water); Mobile Phase B: acetonitrile; Flow rate: 40 mL/min; Gradient: 20% B-50% B in 20 min;




[0563] To a mixture of methyl 2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (50 mg, 0.08 mmol) in THF (5 mL) and water (4 mL) lithium hydroxide monohydrate (34 mg, 0.80 mmol) was added. The resulting mixture was stirred at room temperature for 5 hours. After completion, the resulting mixture was adjusted to pH=5-6 with hydrochloric acid (1 N) and extracted with ethyl acetate (10 mL $\times$ 3). The combined organic layers were washed with brine (10 mL), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by reverse-

Detector: 254 nm) to obtain 2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (10.33 mg, 21.2% yield). LC-MS m/z : 609 $[\text{M}+\text{H}]^+$.

[0564] ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.06 (s, 1H), 7.88-7.80 (m, 2H), 7.69-7.64 (m, 3H), 7.42 (t, $J=8.8$ Hz, 1H), 7.29 (t, $J=8.0$ Hz, 1H), 7.02-6.98 (m, 2H), 6.92 (dd, $J=8.0$ Hz, 2.0 Hz, 1H), 6.41 (s, 1H), 5.71 (s, 2H), 5.24 (s, 2H), 4.36-4.35 (m, 1H), 4.03-3.97 (m, 2H), 3.81 (s, 2H), 2.69-2.67 (m, 2H), 2.33-2.28 (m, 2H), 1.83-1.81 (m, 2H), 1.49-1.47 (m, 2H), 1.15 (t, $J=7.2$ Hz, 3H).

Example 46. Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzimidazole-6-carboxylic acid (Compound 134)


1). Synthesis of N-(4-bromo-2-(((1-(cyanomethyl)cyclopropyl)methyl)amino)-6-fluorophenyl)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamide




[0565] The mixture of N-(4-bromo-2-(((1-(cyanomethyl)cyclopropyl)methyl)amino)-6-fluorophenyl)-2-chloroacetamide (350 mg, 1.04 mmol), 3-fluoro-4-((6-(piperidin-4-oxy)pyridin-2-yl)methoxy)benzonitrile (389 mg, 1.04 mmol) and potassium carbonate (287 mg, 2.08 mmol) in N,N-dimethylformamide (8 mL) was stirred at room temperature for 16 hours. Upon completion of the reaction, the resulting mixture was poured into brine (20 mL) and extracted with ethyl acetate (2×20 mL). The combined organic layers were concentrated to give a residue in vacuo.

The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain N-(4-bromo-2-(((1-(cyanomethyl)cyclopropyl)methyl)amino)-6-fluorophenyl)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidine-1-yl)acetamide (693 mg, 99.0% yield). LC-MS m/z : 667 $[M+H]^+$.

2). Synthesis of 4-((6-((1-((6-bromo-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-2-yl)methyl)piperidin-4-yl)oxy)pyridin-2-yl)methoxy)-3-fluorobenzonitrile




-continued



[0566] The mixture of N-(4-bromo-2-(((1-(cyanomethyl)cyclopropyl)methyl)amino)-6-fluorophenyl)-2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamide (690 mg, 1.04 mmol) in toluene (12 mL) and acetic acid (2 mL) was warmed to 110° C. and stirred for 3 hours under nitrogen atmosphere. After completion, the resulting mixture was concentrated to give a residue. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=1/1) to obtain 4-((6-((1-((6-

bromo-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazol-2-yl)methyl)piperidin-4-yl)oxy)pyridin-2-yl)methoxy)-3-fluorobenzonitrile (330 mg, 49.1% yield). LC-MS m/z: 649 [M+H]⁺.

3). Synthesis of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylate




[0567] To a solution of 4-((6-((1-((6-bromo-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-2-yl)methyl)-piperidin-4-yl)oxy)pyridin-2-yl)methoxy)-3-fluorobenzonitrile (200 mg, 0.31 mmol) in N,N-dimethylformamide (10 mL) and methanol (10 mL) triethylamine (94 mg, 0.93 mmol) and 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (25 mg, 0.03 mmol) were added at room temperature under carbon monoxide atmosphere. The mixture was stirred at 80° C. for 16 hours. After completion, the mixture was diluted with water (5 mL) and extracted with ethyl acetate (10 mL×2). The combined organic layers were washed with brine (10 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=1/1) to obtain methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylate (150 mg, 77.5% yield). LC-MS m/z: 627 [M+H]⁺.

4). Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzimidazole-6-carboxylic acid


[0568] To a mixture of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzo[d]imidazole-6-carboxylate (150 mg, 0.24 mmol) in THF (3 mL) water (3 mL) lithium hydroxide (57 mg, 2.40 mmol) was added at room temperature. The resulting mixture was stirred and reacted at 40° C. for 3 hours. After completion, the solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μm 19×250 mm; Gradient elution with ACN/0.1% NH₄OH in H₂O solvent system; Detection Wavelength 254 nm/214 nm; Flow rate: 20 mL/min) purification to give 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-4-fluoro-1H-benzimidazole-6-carboxylic acid (16.90 mg, 11.5% yield). LC-MS m/z: 613 [M+H]⁺.

[0569] ¹H NMR (400 MHz, DMSO-d₆) δ 8.07 (s, 1H), 7.88 (dd, J=11.2 Hz, 1.6 Hz, 1H), 7.73 (d, J=8.0 Hz, 1H), 7.66 (d, J=8.8 Hz, 1H), 7.52-7.43 (m, 2H), 7.05 (d, J=7.2 Hz, 1H), 6.73 (d, J=8.4 Hz, 1H), 5.31 (s, 2H), 4.95-4.90 (m, 1H), 4.61 (s, 2H), 3.86 (s, 2H), 2.81-2.78 (m, 2H), 2.70 (s, 2H), 2.34-2.29 (m, 2H), 1.94-1.91 (m, 2H), 1.66-1.61 (m, 2H), 0.76-0.67 (m, 4H).



Example 47. Synthesis of 2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C84)


1). Synthesis of methyl 2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylate




[0570] The mixture of methyl 2-(chloromethyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylate (57 mg, 0.18 mmol), 2-((4-chloro-2-fluorophenoxy)methyl)-6-(piperidin-4-acyloxy)pyridine (61 mg, 0.18 mmol), and potassium carbonate (50 mg, 0.36 mmol) in N, N-dimethylformamide (5 mL) was stirred at to 60° C. for 2 hours. After completion, the resulting mixture was poured into brine (20 mL) and extracted with ethyl acetate (2×20 mL). The combined organic layer was concentrated to give a residue in vacuo. The residue was purified by silica gel

column chromatography (dichloromethane/methanol=20/1) to methyl 2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridine-2-yl)oxypiperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylate (20 mg, 18.0% yield). LC-MS m/z: 618 [M+H]⁺.

2). Synthesis of 2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid






[0571] The mixture of methyl 2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylate (20 mg, 0.03 mmol) and lithium hydroxide (7 mg, 0.30 mmol) in THE (2 mL) and (2 mL) was stirred at 40° C. and stirred for 3 hours. After completion, the solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% NH_4OH in H_2O solvent system; Detection Wavelength 254 nm/214 nm; Flow rate: 20 ml/min) to obtain 2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (16.58 mg, 84.8% yield). LC-MS m/z : 604 $[\text{M}+\text{H}]^+$.

[0572] ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.26 (s, 1H), 7.82 (d, $J=8.4$ Hz, 1H), 7.74-7.66 (m, 2H), 7.44 (d, $J=11.2$ Hz, 1H), 7.28 (t, $J=8.4$ Hz, 1H), 7.19 (d, $J=8.8$ Hz, 1H), 7.05 (d, $J=6.8$ Hz, 1H), 6.72 (d, $J=8.4$ Hz, 1H), 5.19 (s, 2H), 4.98-4.92 (m, 1H), 4.61 (s, 2H), 3.86 (s, 2H), 2.81-2.79 (m, 2H), 2.69 (s, 2H), 2.36-2.31 (m, 2H), 1.94-1.92 (m, 2H), 1.66-1.59 (m, 2H), 0.76-0.70 (m, 4H).

Example 48. Synthesis of 2-((4-((3-((4-chloro-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound 7)


1). Synthesis of methyl 4-(2-(4-(3-((4-chloro-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino methyl benzoate



[0573] The mixture of 4-(3-((4-chloro-2-fluorophenoxy)methyl)phenoxy)piperidine (198 mg, 0.59 mmol), methyl 4-(2-chloroacetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (190 mg, 0.54 mmol) and potassium carbonate (149 mg, 1.08 mmol) in N,N-dimethylformamide (7 mL) was stirred at room temperature for 16 hours. After completion, water (20 mL) was added to quench the reaction. The mixture was extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (20 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The

residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 4-(2-(4-(3-((4-chloro-2-fluorophenoxy)methyl)phenoxy))piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (210 mg, 59.8% yield). LC-MS m/z: 650 [M+H]⁺.

2). Synthesis of methyl 2-((4-(3-((4-chloro-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate




[0574] The mixture of methyl 4-(2-(4-(3-((4-chloro-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (210 mg, 0.32 mmol) was dioxane (10 mL) and acetic acid (1 mL) was stirred at 100° C. for 2 hours. After completion, the reaction solution was concentrated to give a residue. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 2-((4-(3-((4-chloro-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (158 mg, 78.1% yield). LC-MS m/z: 632 [M+H]⁺.

3). Synthesis of 2-((4-(3-((4-chloro-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid

After completion, the resulting mixture was adjusted to pH=5-6 with hydrochloric acid (1M) and extracted with ethyl acetate (10 mL×3). The combined organic phases were washed with brine (10 mL), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by reverse-phase flash chromatography under the following conditions (Column: Spherical C18, 20-40 μm, 40 g; Mobile Phase A: 10 mM NH₄OH in water; Mobile Phase B: Acetonitrile; Flow rate: 40 mL/min; Gradient: 20% B-50% B in 20 min; Detector: 254 nm) to obtain 2-((4-(3-((4-chloro-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid formic acid (10.32 mg, 6.7% yield). LC-MS m/z: 618 [M+H]⁺.

[0576] ¹H NMR (400 MHz, DMSO-d₆) δ 8.17 (s, 1H), 8.07 (s, 1H), 7.81 (dd, J=8.4, 1.2 Hz, 1H), 7.68 (d, J=8.4 Hz,




[0575] The mixture of lithium hydroxide monohydrate (105 mg, 2.50 mmol) and methyl 2-((4-(3-((4-chloro-2-fluorophenoxy)methyl)phenoxy)piperidine-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (158 mg, 0.25 mmol) in THF (5 mL) and water (4 mL) was stirred at room temperature for 16 hours.

2H), 7.43 (dd, J=11.6, 2.8 Hz, 1H), 7.30-7.18 (m, 3H), 6.99-6.96 (m, 2H), 6.90 (dd, J=8.0, 1.6 Hz, 1H), 6.42 (s, 1H), 5.72 (s, 2H), 5.14 (s, 2H), 4.37-4.34 (m, 1H), 4.03-3.97 (m, 2H), 3.81 (s, 2H), 2.70-2.67 (m, 2H), 2.33-2.29 (m, 2H), 1.83-1.80 (m, 2H), 1.49-1.46 (m, 2H), 1.16 (t, J=7.2 Hz, 3H).



Example 49. Synthesis of (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)amino)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C16)

1). Synthesis of methyl (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)amino)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate



[0577] The mixture of methyl (S)-2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 0.34 mmol), 3-fluoro-4-((6-(piperidin-4-ylamino)pyridin-2-yl)methoxy)benzonitrile (110 mg, 0.34 mmol) and potassium carbonate (94 mg, 0.68 mmol) in N,N-dimethylformamide (4 mL) was stirred and reacted at 60° C. for 3 hours. After completion, the mixture was diluted with water (10 mL) and extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (10 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)amino)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (120 mg, 63.2% yield). LC-MS m/z : 585 $[M+H]^+$.

2). Synthesis of (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)amino)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid




[0578] To a mixture of methyl (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)amino)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (120 mg, 0.21 mmol) in THF (3 mL) and water (3 mL) lithium hydroxide (25 mg, 1.05 mmol) was added. The reaction was stirred at room temperature for 16 hours. After completion, the resulting mixture was adjusted to pH=5-6 with formic acid. The solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19×250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: mL/min) to give (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)amino)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (16.75 mg, 14.1% yield). LC-MS m/z : 571 $[M+H]^+$.

[0579] ¹H NMR (400 MHz, DMSO-d₆): δ 8.27 (s, 1H), 7.88-7.79 (m, 2H), 7.66-7.64 (m, 2H), 7.45-7.34 (m, 2H),

6.55-6.51 (m, 1H), 6.39 (d, J=8.4 Hz, 1H), 5.15 (s, 2H), 5.10-5.09 (m, 1H), 4.81-4.76 (m, 1H), 4.64 (d, J=15.6 Hz, 1H), 4.51-4.49 (m, 1H), 4.40-4.38 (m, 1H), 3.93 (d, J=13.2 Hz, 1H), 3.75 (d, J=12.8 Hz, 1H), 3.63-3.62 (m, 1H), 2.87-2.84 (m, 1H), 2.74-2.67 (m, 2H), 2.46-2.41 (m, 1H), 2.21-2.11 (m, 2H), 1.85-1.82 (m, 2H), 1.40-1.34 (m, 2H).

Example 50. Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)amino)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C17)


1). Synthesis of methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)amino)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate




[0580] To a mixture of methyl 4-(2-chloroacetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (170 mg, 0.48 mmol) and 3-fluoro-4-((6-(piperidin-4-ylamino)pyridin-2-yl)methoxy)benzonitrile (158 mg, 0.48 mmol) in N,N-dimethylformamide (5 mL) potassium carbonate (132 mg, 0.96 mmol) was added at room temperature. The mixture was stirred at room temperature for 16 hours. After completion, the mixture was extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (20 mL), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was

purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)amino)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (65 mg, 41.8% yield). LC-MS m/z: 641 [M+H]⁺.

2). Synthesis of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)amino)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate




-continued



[0581] The mixture of methyl 4-(2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)amino)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (65 mg, 0.10 mmol) in toluene (3 mL) and acetic acid (0.5 mL) was stirred at 110° C. for 3 hours. After the completion of the reaction, the reaction solution was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 2-((4-((6-((4-cyano-2-

fluorophenoxy)methyl)pyridin-2-yl)amino)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazol-6-carboxylate (45 mg, 71.4% yield). LC-MS m/z : 623 $[M+H]^+$.

3) Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)amino)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid




[0582] To a mixture of methyl 2-((4-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)aminopiperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (45 mg, 0.07 mmol) in water (3.0 mL) and THF (3.0 mL) lithium hydroxide (7 mg, 0.28 mmol) was added. The reaction mixture was stirred at room temperature for 5 hours. After the completion of the reaction, the reaction mixture was adjusted to pH=5-6 with formic acid.

[0583] The solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)amino)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (14.70 mg, 33.5% yield). LC-MS m/z: 609 [M+H]⁺.

[0584] ¹H NMR (400 MHz, DMSO-d₆): δ 8.05 (s, 1H), 7.87-7.79 (m, 2H), 7.68-7.63 (m, 3H), 7.44-7.34 (m, 2H), 6.53 (d, J=7.2 Hz, 1H), 6.47-6.44 (m, 2H), 6.40-6.38 (m, 1H), 5.72-5.69 (m, 2H), 5.14 (s, 2H), 4.02-3.97 (m, 2H), 3.78-3.76 (m, 2H), 3.59-3.54 (m, 1H), 2.76-2.67 (m, 2H), 2.14-2.07 (m, 2H), 1.77-1.72 (m, 2H), 1.29-1.20 (m, 2H), 1.14 (t, J=7.2 Hz, 3H).


Example 51. Synthesis of 1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-2-(((4-((6-((4-ethyl-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C18)

1). Synthesis of methyl 4-(2-(4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-(((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)amino)benzoate



[0585] The mixture of methyl 4-(2-chloroacetamido)-3-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)amino-benzoate (300 mg, 0.80 mmol), 2-((4-bromo-2-fluorophenoxy)methyl)-6-(piperidin-4-oxy)pyridine (305 mg, 0.80 mmol) and potassium carbonate (221 mg, 1.60 mmol) in N,N-dimethylformamide (8 mL) was stirred at room temperature for 16 hours. After completion, the resulting mixture was diluted with water (30 mL) and extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (10 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 4-(2-(4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)amino)benzoate (350 mg, 60.6% yield). LC-MS m/z : 723 $[M+H]^+$.


2). Synthesis of methyl 2-((4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate



[0586] The mixture of methyl 4-(2-(4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)amino)benzoate (350 mg, 0.48 mmol) in toluene (4 mL) and acetic acid (0.4 mL) was stirred at 110° for 3 hours. After completion, the solvent was removed by concentration

to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 2-((4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (300 mg, 89.6% yield). LC-MS m/z : 705 $[M+H]^+$.


3). Synthesis of methyl 1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-2-(((4-((6-(2-fluoro-4-((trimethylsilyl)ethynyl)phenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate



[0587] The mixture of methyl 2-(((4-((6-(2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (150 mg, 0.21 mmol), ethynyltrimethylsilane (206 mg, 2.10 mmol), triethylamine (212 mg, 2.10 mmol) and 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (15 mg, 0.02 mmol) in dry DMSO (5 mL) was purged with nitrogen at room temperature. The resulting mixture was warmed to 90° C. and stirred at 90° C. for 2 hours. After completion, the reaction solution was cooled to room temperature and

quenched with water (60 mL), and then extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (20 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated to give residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=15/1) to give methyl 1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-2-(((4-((6-(2-fluoro-4-((trimethylsilyl)ethynyl)phenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (80 mg, 52.8% yield). LC-MS *m/z*: 721 [M+H]⁺.

4). Synthesis of 1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-2-((4-((6-((4-ethynyl-2-fluorophenoxy)-methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid




[0588] To a mixture of methyl 1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-2-((4-((6-((2-fluoro-4-((trimethylsilyl)ethynyl)phenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (80 mg, 0.11 mmol) in THF (2 mL), water (2 mL) and MeOH (2 mL) lithium hydroxide (8 mg, 0.33 mmol) was added at room temperature. The resulting mixture was stirred at room temperature for 16 hours. After completion, the solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214

nm; Flow rate: mL/min) to obtain 1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-2-((4-((6-((4-ethynyl-2-fluorophenoxy)-methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (3.59 mg, 5.1% yield). LC-MS m/z : 635 [M+H]⁺.

[0589] ¹H NMR (400 MHz, DMSO-d₆): δ 8.05 (s, 1H), 7.82 (dd, J=8.4 Hz, 1.2 Hz, 1H), 7.73-7.65 (m, 3H), 7.39-7.36 (m, 1H), 7.24-7.23 (m, 2H), 7.03 (d, J=7.2 Hz, 1H), 6.71 (d, J=8.0 Hz, 1H), 6.36 (s, 1H), 5.74 (s, 2H), 5.20 (s, 2H), 4.88 (s, 1H), 4.11 (s, 1H), 3.89-3.82 (m, 4H), 2.70-2.67 (m, 2H), 2.28-2.24 (m, 2H), 1.81-1.79 (m, 2H), 1.48-1.46 (m, 2H), 1.06-0.97 (m, 1H), 0.49-0.46 (m, 2H), 0.34-0.31 (m, 2H).

Example 52. Synthesis of 2-((4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C19)




[0590] To a mixture of methyl 2-((4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (60 mg, 0.09 mmol) in THF (4 mL) and water (4 mL) lithium hydroxide (6 mg, 0.27 mmol) was added at room temperature. The mixture was stirred at room temperature for 16 hours. After completion, the solvent was removed by concentration to give a residue in vacuo. The residue was purified prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214

nm; flow rate 20 ml/min) to obtain 2-((4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (19.27 mg, 31.0% yield). LC-MS *m/z*: 691 [M+H]⁺.

[0591] ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.17 (s, 1H), 8.06 (s, 1H), 7.81 (dd, *J*=8.4 Hz, 1.2 Hz, 1H), 7.72-7.67 (m, 3H), 7.54 (dd, *J*=10.8 Hz, 2.0 Hz, 1H), 7.30 (d, *J*=9.2 Hz, 1H), 7.21 (t, *J*=9.2 Hz, 1H), 7.03 (d, *J*=7.2 Hz, 1H), 6.70 (d, *J*=8.4 Hz, 1H), 6.36 (s, 1H), 5.75 (s, 2H), 5.17 (s, 2H), 4.88-4.87 (m, 1H), 3.87-3.83 (m, 4H), 2.70-2.67 (m, 2H), 2.28-2.23 (m, 2H), 1.80-1.78 (m, 2H), 1.47-1.45 (m, 2H), 1.06-0.97 (m, 1H), 0.51-0.46 (m, 2H), 0.34-0.32 (m, 2H).

Example 53. Synthesis of 2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C20)


1). Synthesis of methyl 2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate




[0592] The mixture of methyl 2-(chloromethyl)-1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 0.28 mmol), 3-fluoro-4-((3-(piperidin-4-oxy)benzyl)oxy)benzonitrile (92 mg, 0.28 mmol) and potassium carbonate (77 mg, 0.56 mmol) in N, N-dimethylformamide (3 mL) was warmed to 60° C. and stirred at 60° C. for 3 hours. After completion, the mixture was diluted with water (20 mL) and extracted with ethyl acetate (10 mL×2). The combined organic layers were washed with brine (10 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo.

The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (30 mg, 16.5% yield). LC-MS m/z : 649 $[M+H]^+$.

2). Synthesis of 2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid



-continued




[0593] To a mixture of methyl 2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (30 mg, 0.046 mmol) in THF (2 mL) and water (2 mL) lithium hydroxide (4 mg, 0.15 mmol) was added. The reaction was stirred at room temperature for 16 hours. After completion, the solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain 2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (5.16 mg, 16.3% yield). LC-MS *m/z*: 635 [M+H]⁺.

[0594] ¹H NMR (400 MHz, DMSO-d₆): δ 8.19 (s, 1H), 8.05 (s, 1H), 7.88-7.80 (m, 2H), 7.72-7.66 (m, 3H), 7.42 (t, J=8.8 Hz, 1H), 7.29 (t, J=8.4 Hz, 1H), 7.02-6.98 (m, 2H), 6.93-6.91 (m, 1H), 6.36 (s, 1H), 5.75 (s, 2H), 5.24 (s, 2H), 4.36 (s, 1H), 3.86-3.82 (m, 4H), 2.70-2.67 (m, 2H), 2.33-2.29 (m, 2H), 1.83-1.80 (m, 2H), 1.48-1.46 (m, 2H), 1.06-0.99 (m, 1H), 0.49-0.46 (m, 2H), 0.34-0.31 (m, 2H).

Example 54. Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-methyl-1H-benzo[d]imidazole-6-carboxylic acid (compound

[0595] C21)

1). Synthesis of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-methyl-1H-benzo[d]imidazole-6-carboxylate




[0596] The mixture of methyl 2-(chloromethyl)-1-methyl-1H-benzo[d]imidazole-6-carboxylate (120 mg, 0.50 mmol), 3-fluoro-4-((6-(piperidin-4-oxy)pyridin-2-yl)methoxy)benzonitrile (163 mg, 0.50 mmol) and potassium carbonate (138 mg, 1.0 mmol) in N,N-dimethylformamide (5 mL) was stirred at room temperature for 16 hours. After completion, the mixture was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-methyl-1H-benzo[d]imidazole-6-carboxylate (135 mg, 50.7% yield). LC-MS m/z : 530 $[M+H]^+$.

2). Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-methyl-1H-benzo[d]imidazole-6-carboxylic acid

(135 mg, 0.25 mmol) in water (5 mL) and THF (5 mL) hydroxide lithium (60 mg, 2.50 mmol) was added at room temperature. The resulting mixture was stirred at room temperature for 16 hours. After completion, the solvent was removed by concentration to give a residue in vacuo. The residue was purified by Prep HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) purification to give 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-methyl-1H-benzo[d]imidazole-6-carboxylic acid (72.01 mg; 55.9% yield). LC-MS m/z : 516 $[M+H]^+$.

[0598] ¹H NMR (400 MHz, DMSO- d_6): δ 12.75 (s, 1H), 8.16-8.15 (m, 1H), 7.90-7.87 (m, 1H), 7.81 (dd, J =8.4, 1.6 Hz, 1H), 7.73 (t, J =8.0 Hz, 1H), 7.68-7.64 (m, 2H), 7.45 (t, J =8.4 Hz, 1H), 7.06 (d, J =7.2 Hz, 1H), 6.74 (d, J =8.4 Hz,




[0597] To a mixture of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-methyl-1H-benzo[d]imidazole-6-carboxylate

1H), 5.31 (s, 2H), 4.94-4.89 (m, 1H), 3.92 (s, 3H), 3.84 (s, 2H), 2.77-2.74 (m, 2H), 2.33-2.29 (m, 2H), 1.92-1.90 (m, 2H), 1.65-1.59 (m, 2H).


Example 55: Synthesis of 2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-methyl-1H-benzo[d]imidazole-6-carboxylic acid (compound C22)

1). Synthesis of methyl 2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-methyl-1H-benzo[d]imidazole-6-carboxylate



[0599] The mixture of methyl 2-(chloromethyl)-1-methyl-1H-benzo[d]imidazole-6-carboxylate (120 mg, 0.50 mmol), 2-((4-chloro-2-fluorophenoxy)methyl)-6-(piperidin-4-oxy)pyridine (170 mg, 0.50 mmol) and potassium carbonate (210 mg, 1.05 mmol) in N,N-dimethylformamide (10 mL) was stirred at room temperature for 16 hours. After completion, the mixture was concentrated to give a residue in vacuo. The residue was separated and purified by silica gel column (dichloromethane/methanol=19/1) to obtain methyl 2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-methyl-1H-benzo[d]imidazole-6-carboxylate (120 mg, 44.3% yield). LC-MS m/z: 539 [M+H]⁺.

2). Synthesis of 2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-methyl-1H-benzo[d]imidazole-6-carboxylic acid




[0600] To a mixture of methyl 2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-methyl-1H-benzo[d]imidazole-6-carboxylate (120 mg, 0.22 mmol) in water (4 mL) and THF (4 mL) lithium hydroxide (53 mg, 2.20 mmol) was added at room temperature. The reaction mixture was stirred at room temperature for 16 hours. After completion, the solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain 2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidine pyridin-1-yl)methyl)-1-methyl-1H-benzo[d]imidazole-6-carboxylic acid (89.02 mg, 77.2% yield). LC-MS *m/z*: 525 [M+H]⁺.

[0601] ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.15 (s, 1H), 7.81 (dd, *J*=8.4 Hz, 1.6 Hz, 1H), 7.71 (t, *J*=8.0 Hz, 1H), 7.64 (d, *J*=8.4 Hz, 1H), 7.44 (dd, *J*=11.2 Hz, 2.4 Hz, 1H), 7.28 (t, *J*=9.2 Hz, 2H), 7.18 (d, *J*=8.8 Hz, 1H), 7.04 (d, *J*=7.2 Hz, 1H), 6.71 (d, *J*=8.4 Hz, 1H), 5.18 (s, 2H), 4.94-4.92 (m, 1H), 3.92 (s, 3H), 3.83 (s, 2H), 2.77-2.74 (m, 2H), 2.34-2.29 (m, 2H), 1.93-1.91 (m, 2H), 1.65-1.61 (m, 2H).

Example 56. Synthesis of (S)-2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-4-fluoro-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C23)


1). Synthesis of (S)-N-(4-bromo-2-fluoro-6-((oxetan-2-ylmethyl)amino)phenyl)-2-(4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamide



[0602] The mixture of 2-((4-chloro-2-fluorophenoxy)methyl)-6-(piperidin-4-oxy)pyridine (1.63 g, 4.85 mmol), (S)-N-(4-bromo-2-fluoro-6-((oxetan-2-yl-methyl)amino)phenyl)-2-chloroacetamide (1.70 g, 4.85 mmol) and potassium carbonate (1.34 g, 9.70 mmol) in N,N-dimethylformamide (20 mL) was stirred at room temperature for 16 hours. After completion, the mixture was diluted with ethyl acetate (100 mL) and washed with saturated ammonium chloride

(20 mL $\times$ 3) and brine (20 mL $\times$ 3), dried over anhydrous sodium sulfate, filtered, and reduced in pressure concentrate. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=1/1) to obtain (S)-N-(4-bromo-2-fluoro-6-((oxetan-2-yl-methyl)amino)phenyl)-2-(4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamide (2.10 g, 66.6% yield). LC-MS *m/z*: 653 [M+H]⁺.


2). Synthesis of methyl (S)-6-Bromo-2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl))-4-fluoro-1-(oxetan-2-yl-methyl)-1H-benzo[d]imidazole




[0603] To a solution of (S)—N-(4-bromo-2-fluoro-6-((oxetan-2-yl-methyl)amino)phenyl)-2-(4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamide (500 mg, 0.77 mmol) in dioxane (10 mL) acetic acid (1 mL) was added at room temperature. The resulting mixture was stirred for at 100° C. 16 hours. After completion, the reaction solution was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=1/1) to

obtain (S)-6-bromo-2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-4-fluoro-1-(oxetan-2-yl-methyl)-1H-benzo[d]imidazole (220 mg, 45.2% yield). LC-MS m/z : 635 $[M+H]^+$.

3). Synthesis of methyl (S)-2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-4-fluoro-1-(oxetan-2-yl-methyl)-1H-benzo[d]imidazole-6-carboxylate






[0604] To a mixture of (S)-6-bromo-2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-4-fluoro-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole (100 mg, 0.16 mmol) in N,N-dimethylformamide (5 mL) and methanol (5 mL) triethylamine (48 mg, 0.48 mmol) and 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (15 mg, 0.02 mmol) were added at room temperature under carbon monoxide atmosphere. The mixture was stirred at 80° C. for 16 hours. After completion, the mixture was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum

ether/ethyl acetate=1/1) to obtain methyl (S)-2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-4-fluoro-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (48 mg, 50.0% yield). LC-MS m/z: 613 [M+H]⁺.

4). Synthesis of (S)-2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-4-fluoro-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid




[0605] To a mixture of methyl (S)-2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-4-fluoro-1-(oxetan-2-yl-methyl)-1H-benzo[d]imidazole-6-carboxylate (48 mg, 0.08 mmol) in THF (2 mL) and water (2 mL) lithium hydroxide monohydrate (34 mg, 0.80 mmol) was added at room temperature. The resulting mixture was stirred at room temperature for 5 hours. After completion, the resulting mixture was adjusted to pH=5-6 with hydrochloric acid (1 M) and extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (10 mL), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by reverse-phase flash chromatography under the following conditions (Column: Spherical C18, 20-40 m, 40 g; Mobile Phase A: 10 mM NH₄OH in water; Mobile Phase B: acetonitrile; Flow rate: 40 mL/min; Gradient: 20% B-50% B in 20 min; Detector: 254 nm) to give (S)-2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-4-fluoro-1-(oxetan-2-

ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (20.87 mg, 43.6% yield). LC-MS m/z: 599 [M+H]⁺.

[0606] ¹H NMR (400 MHz, DMSO-d₆) δ 8.11 (s, 1H), 7.72 (t, J=7.6 Hz, 1H), 7.54 (d, J=11.6 Hz, 1H), 7.45 (dd, J=11.2 Hz, 2.8 Hz, 1H), 7.28 (t, J=9.2 Hz, 1H), 7.20-7.17 (m, 1H), 7.04 (d, J=7.2 Hz, 1H), 6.72 (d, J=8.4 Hz, 1H), 5.19 (s, 2H), 5.12-5.07 (m, 1H), 4.97-4.92 (m, 1H), 4.82-4.76 (m, 1H), 4.65 (d, J=12.8 Hz, 1H), 4.52-4.47 (m, 1H), 4.39-4.34 (m, 1H), 3.94 (d, J=13.6 Hz, 1H), 3.79 (d, J=13.6 Hz, 1H), 2.80-2.67 (m, 3H), 2.46-2.41 (m, 1H), 2.35-2.32 (m, 2H), 1.93-1.91 (m, 2H), 1.63-1.59 (m, 2H).


Example 57. Synthesis of 2-(4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)benzyl)-1-((1-ethyl-1H-imidazole-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C24)

1). Synthesis of methyl 2-(4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)phenyl)acetate




[0607] The mixture of methyl 2-(4-((6-((methylsulfonyl)oxy)methyl)pyridin-2-yl)oxy)phenyl)acetate (410 mg, 1.17 mmol), 3-fluoro-4-hydroxybenzonitrile (171 mg, 1.17 mmol) and potassium carbonate (323 mg, 2.34 mmol) in N,N-dimethylformamide (6 mL) was stirred at room temperature for 16 hours. After completion, water (5 mL) was added to dilute the mixture, and extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (15 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 2-(4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)phenyl)acetate (406 mg, 86.4% yield). LC-MS m/z: 402 [M+H]⁺.

2). Synthesis of 2-(4-((6-((4-Chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)phenyl)acetic acid



[0608] To a mixture of methyl 2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)phenyl)acetate (406 mg, 1.01 mmol) in THF (5 mL) and water (5 mL) lithium hydroxide monohydrate (424 mg, 10.10 mmol) was added at room temperature. The resulting mixture was stirred at room temperature for 5 hours. After completion, the resulting mixture was adjusted to pH=5-6 with hydrochloric acid (1 M). The mixture was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain 2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)phenyl)acetic acid (248 mg, 63.3% yield). LC-MS m/z : 388 $[M+H]^+$.


3. Synthesis of methyl 4-(2-(4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)phenyl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate




[0609] To a mixture of 2-((4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)phenyl)acetic acid (248 mg, 0.64 mmol), 2-(3H-[1,2,3]triazolo[4,5-b]pyridin-3-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate(V) (365 mg, 0.96 mmol) and N,N-diisopropylethylamine (330 mg, 2.56 mmol) in N,N-dimethylformamide (3 mL) methyl 4-amino-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (175 mg, 0.64 mmol) was added at room temperature. The mixture was stirred at room temperature for 16 hours. After completion, the reaction was quenched by adding water (5 mL) and extracted with ethyl acetate (20 mL $\times$ 2). The combined organic layers washed with brine (20 mL $\times$ 3),

dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 4-(2-(4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)phenyl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (232 mg, 56.3% yield). LC-MS m/z : 644 $[M+H]^+$.

4. Synthesis of methyl 2-(4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)benzyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate




-continued



[0610] To a solution of methyl 4-(2-(4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)phenyl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (232 mg, 0.36 mmol) in dioxane (5 mL) acetic acid (0.5 mL) was added at room temperature. The resulting mixture was stirred at 100° C. for 2 hours. After completion, the reaction solution was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain

methyl 2-(4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)benzyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (190 mg, 84.3% yield). LC-MS m/z : 626 $[M+H]^+$.

5). Synthesis of -2-(4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)benzyl)-1-((1-ethyl-1H-imidazole-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid




[0611] To a solution of methyl 2-(4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)benzyl)-1-((1-ethyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (190 mg, 0.30 mmol) dissolved in THF (2 mL) and water (2 mL) lithium hydroxide monohydrate (128 mg, 3.00 mmol) was added at room temperature. The resulting mixture was stirred at room temperature for 5 hours. After completion, the resulting mixture was adjusted to pH=5-6 with hydrochloric acid (1 M) and extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (10 mL), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by prep-HPLC (Gilson CA 258 system; Column: Agilent Prep C18 OBD 10 um 21.2×250 mm; Gradient elution with ACN/0.1% NH₄OH solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain 2-(4-((6-((4-chloro-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)benzyl)-1-((1-ethyl)-1H-imidazol-

5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (20.10 mg, 10.9% yield). LC-MS m/z: 612 [M+H]⁺.

[0612] ¹H NMR (400 MHz, DMSO-d₆) δ 8.03 (s, 1H), 7.88 (t, J=8.0 Hz, 1H), 7.79 (dd, J=8.4 Hz, 1.6 Hz, 1H), 7.63-7.58 (m, 2H), 7.43 (dd, J=11.2 Hz, 2.4 Hz, 1H), 7.30 (d, J=8.4 Hz, 2H), 7.24-7.14 (m, 3H), 7.04 (d, J=8.8 Hz, 2H), 6.91 (d, J=8.4 Hz, 1H), 6.35 (s, 1H), 5.62 (s, 2H), 5.10 (s, 2H), 4.35 (s, 2H), 3.93-3.87 (m, 2H), 1.11 (t, J=7.6 Hz, 3H).


Example 58. Synthesis of 1-((1-(cyanomethyl)cyclopropyl)methyl)-2-((4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridine-2-yl)oxy)piperidin-1-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C25)

1). Synthesis of methyl 4-(2-(4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)benzoate




[0613] The mixture of methyl 4-(2-chloroacetamido)-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)benzoate (513 mg, 2.50 mmol), 2-((4-bromo-2-fluorophenoxy)methyl)-6-(piperidin-4-oxo)pyridine (950 mg, 2.50 mmol) and potassium carbonate (1.00 g, 7.50 mmol) in N,N-dimethylformamide (10 mL) was stirred at room temperature for 16 hours. Upon completion of the reaction, the resulting mixture was poured into brine (20 mL) and extracted with ethyl acetate (2×20 mL). The combined organic layer was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=1/1) to obtain methyl 4-(2-(4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)benzoate (347 mg, 28.0% yield). LC-MS m/z: 680 [M+H]⁺.

2). Synthesis of methyl 2-((4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylate




-continued



[0614] To a solution of methyl 4-(2-(4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-(cyanomethyl)cyclopropyl)methyl)amino)benzoate (347 mg, 0.51 mmol) in toluene (6 mL) acetic acid (1 mL) was added at room temperature. The resulting mixture was stirred for 2 hours at 110° C. under N₂ atmosphere. After completion, the reaction solution was concentrated to give a residue. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 2-((4-((6-((4-bromo-2-fluorophenoxy))

methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]-imidazole-6-carboxylate (336 mg, 99.7% yield). LC-MS m/z: 664 [M+H]⁺.

3). Synthesis of 1-((1-(cyanomethyl)cyclopropyl)methyl)-2-((4-((6-((2-fluoro-4-((trimethylsilyl)ethynyl)phenoxy)methyl)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate




[0615] The mixture of methyl 2-((4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylate (336 mg, 0.51 mmol), ethynyltrimethylsilane (500 mg, 5.10 mmol), triethylamine (515 mg, 5.10 mmol) and 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (37 mg, 0.05 mmol) in DMSO (5 mL) was stirred for 2 hours at 90° C. under nitrogen protection. After completion, the resulting mixture was diluted with water (10 mL) at room temperature, and extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (10 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 1-((1-(cyanomethyl)cyclopropyl)methyl)-2-((4-((6-((2-fluoro-4-((trimethylsilyl)ethynyl)phenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (258 mg, 76.1% yield). LC-MS *m/z*: 680 [M+H]⁺.


4). Synthesis of 1-((1-(cyanomethyl)cyclopropyl)methyl)-2-((4-((6-((4-ethynyl-2-fluorophenoxy)methyl)pyridine-2-yl)oxy)piperidin-1-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid

[0616] To a mixture of methyl 1-((1-(cyanomethyl)cyclopropyl)methyl)-2-((4-((6-((2-fluoro-4-((trimethylsilyl)acetylene)yl)phenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (50 mg, 0.08 mmol) in THF (1 mL), water (1 mL) and methanol (1 mL) lithium hydroxide (19 mg, 0.80 mmol) was added at room temperature. The resulting mixture was stirred at room temperature for 16 hours. After completion, the solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 um 19×250 mm Column; Gradient elution with ACN/0.05% NH₄OH in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain 1-((1-(cyanomethyl)cyclopropyl)methyl)-2-((4-((6-((4-ethynyl)-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (3.20 mg, 6.7% yield). LC-MS *m/z*: 594 [M+H]⁺.

[0617] ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.24 (s, 1H), 7.82 (d, *J*=8.0 Hz, 1H), 7.72 (t, *J*=8.4 Hz, 1H), 7.64 (d, *J*=8.4 Hz, 1H), 7.40-7.37 (m, 1H), 7.26-7.25 (m, 2H), 7.05 (d, *J*=7.6 Hz, 1H), 6.73 (d, *J*=8.0 Hz, 1H), 5.22 (s, 2H), 4.98-4.94 (m, 1H), 4.61 (s, 2H), 4.12 (s, 1H), 3.86 (s, 2H), 2.82-2.79 (m, 2H), 2.69 (s, 2H), 2.37-2.32 (m, 2H), 1.95-1.91 (m, 2H), 1.66-1.59 (m, 2H), 0.76-0.68 (m, 4H).



Example 59. Synthesis of 2-((4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C26)



[0618] To a mixture of 2-((4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylate (50 mg, 0.08 mmol) in THF (3 mL) and water (3 mL) lithium hydroxide (19 mg, 0.80 mmol) was added at room temperature. The resulting mixture was stirred at room temperature for 16 hours. After completion, the residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19x250 mm; Gradient elution with ACN/0.05% NH_4OH in H_2O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain 2-((4-((6-((4-bromo-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyanomethyl)cyclopropyl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (16.81 mg, 32.5% yield). LC-MS m/z : 650 $[\text{M}+\text{H}]^+$.


[0619] ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.22 (s, 1H), 7.81 (d, $J=8.4$ Hz, 1H), 7.71 (t, $J=8.0$ Hz, 1H), 7.61 (d, $J=8.4$


Hz, 1H), 7.54 (dd, $J=11.2$ Hz, 2.0 Hz, 1H), 7.30 (d, $J=8.4$ Hz, 1H), 7.22 (t, $J=8.8$ Hz, 1H), 7.03 (d, $J=7.2$ Hz, 1H), 6.72 (d, $J=8.0$ Hz, 1H), 5.18 (s, 2H), 4.96-4.92 (m, 1H), 4.60 (s, 2H), 3.85 (s, 2H), 2.81-2.78 (m, 2H), 2.69 (s, 2H), 2.35-2.30 (m, 2H), 1.94-1.91 (m, 2H), 1.66-1.61 (m, 2H), 0.72-0.67 (m, 4H).

Example 60. Synthesis of (S)-2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)benzyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound

[0620] C27)

1). Synthesis of methyl (S)-4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)phenyl)acetamido)-3-((oxetan-2-ylmethyl)amino)benzoate







[0621] To a mixture of 2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)phenyl)acetic acid (288 mg, 0.76 mmol) 2-(3H-[1,2,3]triazolo[4,5-b]pyridin-3-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate(V) (433 mg, 1.14 mmol) and N,N-diisopropylethylamine (392 mg, 3.04 mmol) in N,N-dimethylformamide (3 mL) methyl (S)-4-amino-3-((oxetan-2-ylmethyl)amino)benzoate (179 mg, 0.76 mmol) was added at room temperature. The resulting mixture was stirred at room temperature for 16 hours. After completion, the mixture was quenched by adding water (5 mL) and extracted with ethyl acetate (20 mL $\times$ 2). The combined organic layers were washed with brine (20 mL $\times$ 3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl (S)-2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)phenyl)acetamido-3-((oxetan-2-ylmethyl)amino)benzoate (301 mg, 66.6% yield). LC-MS m/z : 597 $[M+H]^+$.


2). Synthesis of methyl (S)-2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)benzyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate

[0622] To a mixture of (S)-4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)phenyl)acetamido)-3-((oxetan-2-ylmethyl)amino)benzoate (301 mg, 0.51 mmol) in dioxane (10 mL) acetic acid (1 mL) was added at room temperature. The resulting mixture was stirred and reacted at 100° C. for 2 hours. After completion, the reaction solution was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl (S)-2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)benzyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (223 mg, 75.6% yield). LC-MS m/z : 579 $[M+H]^+$.

3). Synthesis of (S)-2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)benzyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid



-continued




[0623] To a mixture of methyl (S)-2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridine-2-yl)oxy)benzyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (223 mg, 0.39 mmol) in THF (2 mL) and water (2 mL) lithium hydroxide (93 mg, 3.90 mmol) was added. The resulting mixture was stirred at room temperature for 5 hours. After completion, the obtained mixture was adjusted to pH=5-6 with formic acid, and extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (10 mL), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. The residue was purified by prep-HPLC (Gilson CA 258 system; Column: Agilent Prep C18 OBD 10 μ m 21.2×250 mm; Gradient elution with ACN/0.1% $\text{NH}_3\cdot\text{H}_2\text{O}$ solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain (S)-2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridine-2-yl)oxy)benzyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (75.28 mg, 34.2% yield). LC-MS m/z : 565 $[\text{M}+\text{H}]^+$.


[0624] ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.14 (s, 1H), 7.90 (t, $J=7.6$ Hz, 1H), 7.84 (dd, $J=11.2$ Hz, 2.0 Hz, 1H), 7.79 (dd, $J=8.4$ Hz, 1.2 Hz, 1H), 7.62 (d, $J=8.8$ Hz, 1H), 7.53 (d, $J=8.4$ Hz, 1H), 7.38-7.33 (m, 3H), 7.25 (d, $J=7.2$ Hz, 1H), 7.07 (d, $J=8.8$ Hz, 2H), 6.97 (d, $J=8.0$ Hz, 1H), 5.21 (s, 2H), 4.97-4.92 (m, 1H), 4.65-4.59 (m, 1H), 4.53-4.43 (m, 2H), 4.39-4.32 (m, 3H), 2.68-2.58 (m, 1H), 2.38-2.30 (m, 1H).

Example 61. Synthesis of 2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridine-2-yl)oxy)benzyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C28)

1). Synthesis of methyl 2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridine-2-yl)oxy)phenyl)acetate





-continued




[0625] The mixture of 3-fluoro-4-hydroxybenzonitrile (22 mg, 0.16 mmol), potassium carbonate (44 mg, 0.32 mmol) and methyl 2-(4-((6-((methylsulfonyl)oxy)methyl)pyridine-2-yl)oxy)phenyl)acetate in N,N -dimethylformamide (2 mL) was stirred and reacted at 60°C . for 3 hours. After completion, the mixture was diluted with water (5 mL) and extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (15 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=7/3) to obtain 2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridine-2-yl)oxy)phenyl)acetate (31 mg, 49.4% yield). LC-MS m/z : 393 $[\text{M}+\text{H}]^+$.

2). Synthesis of 2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridine-2-yl)oxy)phenyl)acetic acid



[0626] To a solution of methyl 2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridine-2-yl)oxy)phenyl)acetate (31 mg, 0.08 mmol) in THF (1.5 mL) and water (1.5 mL) lithium hydroxide monohydrate (33 mg, 0.80 mmol) was added at room temperature. The resulting mixture was stirred at room temperature for 5 hours. After completion, the resulting mixture was adjusted to pH=5-6 with hydrochloric acid (1 M), then concentrated to give a residue. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain 2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridine-2-yl)oxy)phenyl)acetic acid (30 mg, 99.2% yield). LC-MS m/z : 379 $[\text{M}+\text{H}]^+$.


3). Synthesis of methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)phenyl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate




[0627] To a mixture of 2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)phenyl)acetic acid (30 mg, 0.08 mmol), 2-(3H-[1,2,3]triazolo[4,5-b]pyridin-3-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate(V) (65 mg, 0.12 mmol) and N,N-diisopropylethylamine (57 mg, 0.32 mmol) in N,N-dimethylformamide (2 mL) methyl 4-amino-3-((1-ethyl-1H-imidazol-5-yl)methyl)aminobenzoate (22 mg, 0.08 mmol) was added at room temperature. The resulting mixture was stirred at room temperature for 16 hours. When completion, the mixture was quenched by adding water (5 mL) and extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (15 mL×2), dried over anhydrous sodium sulfate,

filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether: ethyl acetate=1/1) to obtain methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)phenyl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (27 mg, 54.7% yield). LC-MS m/z : 635 $[M+H]^+$.

4). Synthesis of methyl 2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)benzyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate




-continued



[0628] To a solution of methyl 4-(2-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)phenyl)acetamido-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (27 mg, 0.04 mmol) in dioxane (3 mL) acetic acid (0.3 mL) was added at room temperature. The resulting mixture solution was stirred at 100° C. for 3 hours. After completion, the reaction solution was concentrated to give a residue under reduced pressure. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)benzyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (20 mg, 76.3% yield). LC-MS *m/z*: 617 [M+H]⁺.

5). Synthesis of 2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)benzyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid


[0629] To a mixture of (4 mg, 0.09 mmol) methyl 2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridine-2-yl)oxy)benzyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (20 mg, 0.03 mmol) in THF (2 mL) and water (2 mL) lithium hydroxide monohydrate was added at room temperature. The resulting mixture solution was stirred at room temperature for 5 hours. After completion, the resulting mixture was adjusted to pH=5-6 with hydrochloric acid (1 M) and extracted with ethyl acetate (10 mL×3). The combined organic phases were washed with brine (10 mL), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by reverse-phase flash chromatography under the following conditions (Column: Spherical C18, 20-40 μm, 40 g; mobile phase A: 10 mM NH₄OH in water; Mobile Phase B: acetonitrile; Flow rate: 40 mL/min; Gradient: 20% B-50% B in 20 min; Detector: 254 nm to obtain 2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridine-2-yl)oxy)benzyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (2.77 mg, 15.1% yield). LC-MS *m/z*: 603 [M+H]⁺.



[0630] ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.93 (s, 1H), 7.89 (t, $J=8.0$ Hz, 1H), 7.84 (dd, $J=11.2$ Hz, 2.0 Hz, 1H), 7.78 (dd, $J=8.0$ Hz, 0.8 Hz, 1H), 7.64-7.61 (m, 2H), 7.43 (d, $J=8.0$ Hz, 1H), 7.37 (t, $J=8.4$ Hz, 1H), 7.29 (d, $J=8.4$ Hz, 2H), 7.24 (d, $J=7.6$ Hz, 1H), 7.02 (d, $J=8.8$ Hz, 2H), 6.92 (d, $J=8.4$ Hz, 1H), 6.29 (s, 1H), 5.53 (s, 2H), 5.22 (s, 2H), 4.30 (s, 2H), 3.95-3.90 (m, 2H), 1.14 (t, $J=7.2$ Hz, 3H).

Example 62. Synthesis of 2-((4-(3-((4-ethynyl-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-isopropyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C29)


1). Synthesis of methyl 4-(2-(4-(3-((4-bromo-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)acetamido)-3-((1-isopropyl-1H-imidazol-5-yl)methyl)amino)benzoate




[0631] The mixture of methyl 4-(2-chloroacetamido)-3-((1-isopropyl-1H-imidazol-5-yl)methyl)amino)benzoate (450 mg, 1.23 mmol), 4-(3-((4-bromo-2-fluorophenoxy)methyl)phenoxy)piperidine (467 mg, 1.23 mmol) and potassium carbonate (339 mg, 2.46 mmol) in dry N,N-dimethylformamide (10 mL) was stirred at room temperature for 16 hours. After completion, the reaction was quenched with water (10 mL) and extracted with ethyl acetate (3×10 mL). The combined organic layers were washed with brine (10 mL $\times 2$), dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. The residue was purified by silica gel

column chromatography (dichloromethane/methanol=10/1) to obtain methyl 4-(2-(4-(3-((4-bromo-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)acetamido)-3-((1-isopropyl-1H-imidazol-5-yl)methyl)amino)benzoate (400 mg, 51.4% yield). LC-MS m/z : 710 $[\text{M}+\text{H}]^+$.

2). Synthesis of methyl 2-((4-(3-((4-bromo-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)-1-((1-isopropyl-1H-imidazole-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate




-continued



[0632] To a solution of methyl 4-(2-(4-(3-((4-bromo-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)acetamido)-3-((1-isopropyl-1H-imidazol-5-yl)methyl)amino)benzoate (400 mg, 0.56 mmol) in toluene (5 mL) acetic acid (0.5 mL) was added at room temperature. The resulting mixed solution was stirred and reacted at 110° C. for 2 hours. After completion, the reaction solution was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1)

to obtain methyl 2-((4-(3-((4-bromo-2-fluorophenoxy)yl)methyl)phenoxy)piperidin-1-yl)-1-((1-isopropyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazol-6-carboxylate (380 mg, 97.4% yield). LC-MS m/z : 692 $[M+H]^+$.

3). Synthesis of methyl 2-((4-(3-((2-fluoro-4-((trimethylsilyl)ethynyl)phenoxy)phenoxy)piperidin-1-yl)methyl)-1-((1-isopropyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate




[0633] The mixture of methyl 2-((4-(3-((4-bromo-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)-1-(((1-isopropyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (280 mg, 0.41 mmol), ethynyltrimethylsilane (402 mg, 4.10 mmol), triethylamine (414 mg, 4.10 mmol) and 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (29 mg, 0.04 mmol) in DMSO (6 mL) was stirred at 90° C. for 2 hours under N₂ atmosphere. After completion, the reaction was quenched with water (10 mL) and extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (10 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 2-((4-(3-((2-fluoro-4-((trimethylsilyl)ethynyl)phenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-isopropyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (420 mg, crude). LC-MS m/z: 708 [M+H]⁺.


4). Synthesis of 2-((4-(3-((4-ethynyl-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-isopropyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid

[0634] To a mixture of methyl 2-((4-(3-((2-fluoro-4-((trimethylsilyl)ethynyl)phenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-isopropyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (420 mg, 0.59 mmol) in methanol (1 mL), water (1 mL) and THF (1 mL) lithium hydroxide (113 mg, 4.72 mmol) was added at room temperature. The reaction mixture was stirred at room temperature for 5 hours. After completion, the reaction mixture was adjusted to pH with formic acid=5-6. The mixture was concentrated to give a residue in vacuo. The residue was purified by reverse-phase flash chromatography under the following conditions (Column: Spherical C18, 20-40 μm, 40 g; Mobile Phase A: 10 mM NH₄OH in water; Phase B: acetonitrile; Flow rate: 40 mL/min; gradient: 20% B-50% B in 20 min; Detector: 254 nm) to obtain 2-((4-(3-((4-ethynyl-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-isopropyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (23.6 mg, 6.4% yield). LC-MS m/z: 622 [M+H]⁺.

[0635] ¹H NMR (400 MHz, DMSO-d₆): δ 8.02 (s, 1H), 7.81-7.78 (m, 2H), 7.65 (d, J=7.6 Hz, 1H), 7.38-7.35 (m, 1H), 7.30-7.20 (m, 3H), 7.01-6.97 (m, 2H), 6.92-6.89 (m, 1H), 6.37 (s, 1H), 5.72 (s, 2H), 5.16 (s, 2H), 4.45-4.35 (m, 2H), 4.12 (s, 1H), 3.80 (s, 2H), 2.70-2.65 (m, 2H), 2.35-2.27 (m, 2H), 1.86-1.79 (m, 2H), 1.51-1.44 (m, 2H), 1.33-1.27 (m, 6H).



Example 63. Synthesis of 2-((4-(3-((4-bromo-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-isopropyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C30)




[0636] To a solution of 2-((4-(3-((4-bromo-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)-1-((1-isopropyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 0.16 mmol) in water (1 mL) and THF (1 mL), lithium hydroxide (10 mg, 0.42 mmol) was added at room temperature. The reaction mixture was stirred at room temperature for 16 hours. After completion, the reaction mixture was adjusted to pH=5-6 with formic acid. The mixture was concentrated to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: mL/min) to obtain 2-((4-(3-((4-bromo-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-isopropyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (13.21 mg, 12.2% yield). LC-MS m/z : 678 [M+H]⁺.

[0637] ¹H NMR (400 MHz, DMSO- d_6): δ 8.02 (s, 1H), 7.81-7.79 (m, 2H), 7.65 (d, J=8.4 Hz, 1H), 7.54 (dd, J=10.8 Hz, 2.4 Hz, 1H), 7.33-7.26 (m, 2H), 7.22-7.18 (m, 1H), 6.99-6.96 (m, 2H), 6.91-6.89 (m, 1H), 6.37 (s, 1H), 5.72 (s, 2H), 5.14 (s, 2H), 4.46-4.36 (m, 2H), 3.80 (s, 2H), 2.70-2.65 (m, 2H), 2.33-2.29 (m, 2H), 1.82-1.80 (m, 2H), 1.49-1.43 (m, 2H), 1.36-1.30 (m, 6H).


Example 64. Synthesis of 2-((4-(6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-isopropyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C31)

1). Synthesis of methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-isopropyl-1H-imidazol-5-yl)methyl)amino)benzoate



[0638] The mixture of 3-fluoro-4-((6-(piperidin-4-oxy)pyridin-2-yl)methoxy)benzonitrile (160 mg, 0.49 mmol), methyl 4-(2-(chloroacetamido))-3-((1-isopropyl-1H-imidazol-5-yl)methyl)amino)benzoate (178 mg, 0.49 mmol) and potassium carbonate (135 mg, 0.98 mmol) in N,N-dimethylformamide (5 mL) was stirred at room temperature for 16 hours. After completion, water (5 mL) was added to quench the reaction, and extracted with ethyl acetate (40 mL×3). The combined organic layers were washed with brine (30 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)-pyridine)-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-isopropyl-1H-imidazol-5-yl)methyl)amino)benzoate (155 mg, 48.3% yield). LC-MS m/z : 656 $[M+H]^+$.


2). Synthesis of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-isopropyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate



[0639] To a solution of methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-isopropyl-1H-imidazol-5-yl)methyl)amino)benzoate (155 mg, 0.27 mmol) in toluene (3 mL) acetic acid (0.5 mL) was added at room temperature. The reaction was stirred at 110°C. for 2 hours. After completion, water (5 mL) was added to quench the reaction, and extracted with ethyl acetate (40 mL×3). The combined organic layers were

washed with brine (40 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-isopropyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (131 mg, yield: 87.3%). LC-MS m/z : 638 $[M+H]^+$.

3). Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-isopropyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid




[0640] To a mixture of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-isopropyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (131 mg, 0.20 mmol) in THF (2 mL) and water (2 mL) lithium hydroxide (15 mg, 0.60 mmol) was added at room temperature. The mixture was stirred at room temperature for 16 hours. After completion, the resulting mixture was adjusted to pH=5-6 with acetic acid (1M), and then extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (10 mL), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by reverse-phase flash chromatography under the following conditions (Column: Spherical C18, 20-40 μm, 40

g; Mobile Phase A: 10 mM NH₄OH in water; Mobile phase B: acetonitrile; Flow rate: 40 mL/min; Gradient: 20% B-50% B in 20 min; Detector: 254 nm) to give 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-isopropyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (35.02 mg, 28.0% yield). LC-MS m/z: 624 [M+H]⁺.

[0641] ¹H NMR (400 MHz, DMSO-d₆) δ 8.03 (s, 1H), 7.89-7.78 (m, 3H), 7.74-7.64 (m, 3H), 7.44 (t, J=8.4 Hz, 1H), 7.05 (d, J=7.2 Hz, 1H), 6.72 (d, J=8.4 Hz, 1H), 6.37 (s, 1H), 5.73 (s, 2H), 5.30 (s, 2H), 4.88-4.82 (m, 1H), 4.46-4.40 (m, 1H), 3.81 (s, 2H), 2.74-2.57 (m, 2H), 2.27-2.22 (m, 2H), 1.86-1.79 (m, 2H), 1.48-1.46 (m, 2H), 1.34-1.29 (m, 6H).

Example 65. Synthesis of 2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C32)


1). Synthesis of methyl 4-(2-(4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)acetamido-3-((1-isopropyl-1H-imidazol-5-yl)methyl)amino)benzoate




[0642] The mixture of methyl 4-(2-chloroacetamido)-3-((1-isopropyl-1H-imidazol-5-yl)methyl)amino)benzoate (120 mg, 0.33 mmol), 3-fluoro-4-(3-(piperidin-4-oxymethyl)phenoxy)benzonitrile (108 mg, 0.33 mmol) and potassium carbonate (91 mg, 0.66 mmol) in dry N,N-dimethylformamide (5 mL) was stirred at room temperature for 16 hours. After completion, the reaction was quenched with water (10 mL) and extracted with ethyl acetate (3×10 mL). The combined organic layers were washed with brine (10 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was

purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 4-(2-(4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)acetamido-3-((1-isopropyl-1H-imidazol-5-yl)methyl)amino)benzoate (130 mg, 60.5% yield). LC-MS m/z : 655 $[M+H]^+$.

2). Synthesis of methyl 2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)-1-((1-isopropyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate




-continued



[0643] To a solution of methyl 4-(2-(4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)acetamido-3-((1-isopropyl-1H-imidazol-5-yl)methyl)amino)benzoate (130 mg, 0.20 mmol) in toluene (3 mL) acetic acid (0.5 mL) was added at room temperature. The mixture was stirred at 110° C. for 2 hours. After completion, the reaction solution was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 2-((4-(3-((4-

cyano-2-fluorobenzene oxy)methyl)phenoxy)piperidin-1-yl)-1-((1-isopropyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazol-6-carboxylate (100 mg, 79.4% yield). LC-MS *m/z*: 637 [M+H]⁺.

3). Synthesis of 2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid




[0644] To a mixture of methyl 2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)-1-((1-isopropyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 0.16 mmol) in water (2 mL) and THF (2 mL) lithium hydroxide (12 mg, 0.48 mmol) was added at room temperature. The reaction mixture was stirred at room temperature for 16 hours. After completion, the reaction mixture was adjusted pH to 5-6 with formic acid. The mixture was concentrated to give a residue in vacuo. The residue was purified by prep-PLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: mL/min) to obtain 2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-1-((1-ethyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (17.60 mg, 18.0% yield). LC-MS m/z: 623 [M+H]⁺.

[0645] ¹H NMR (400 MHz, DMSO-d₆): δ 8.03 (s, 1H), 7.88-7.80 (m, 3H), 7.67 (t, J=8.8 Hz, 2H), 7.42 (t, J=8.4 Hz, 1H), 7.29 (t, J=7.6 Hz, 1H), 7.03-6.98 (m, 2H), 6.93 (d, J=8.4 Hz, 1H), 6.38 (s, 1H), 5.73 (s, 2H), 5.24 (s, 2H), 4.45-4.37 (m, 2H), 3.81 (s, 2H), 2.69-2.68 (m, 2H), 2.34-2.29 (m, 2H), 1.82 (s, 2H), 1.49-1.47 (m, 2H), 1.34-1.32 (m, 6H).


Example 66. Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C33)

1). Synthesis of methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)amino)benzoate



[0646] The mixture of methyl 4-(2-chloroacetamido)-3-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)amino-benzoate (180 mg, 0.48 mmol), 3-fluoro-4-((6-(piperidin-4-oxy)pyridin-2-yl)methoxy)benzonitrile (156 mg, 0.48 mmol) and potassium carbonate (133 mg, 0.96 mmol) in N,N-dimethylformamide (5 mL) was stirred at room temperature for 16 hours. After completion, it was diluted with water (30 mL) and extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (30 mL) and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=15/1) to obtain methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)amino) benzoate (200 mg, 62.8% yield). LC-MS m/z : 668 $[M+H]^+$.


2). Synthesis of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate



[0647] To a solution of methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)amino)benzoate (200 mg, 0.30 mmol) in dioxane (10 mL) acetic acid (1 mL) was added at room temperature. The resulting mixture was stirred and reacted at 100° C. for 3 hours. After completion, the reaction solution was concen-

trated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=15/1) to obtain methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (150 mg, 77.1% yield). LC-MS m/z : 650 $[M+H]^+$.

3). Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid




[0648] To a mixture of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (150 mg, 0.23 mmol) in water (5 mL) and THF (5 mL) lithium hydroxide (56 mg, 2.30 mmol) was added at room temperature. The reaction mixture was stirred at room temperature for 16 hours. After completion, the reaction was adjusted to pH=5-6 with formic acid. The mixture was concentrated to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% FA in

H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-(cyclopropylmethyl)-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (77.69 mg, 53.1% yield). LC-MS *m/z*: 636 [M+H]⁺.

[0649] ¹H NMR (400 MHz, DMSO-*d*₆): δ 12.76 (br, 1H), 8.06 (s, 1H), 7.90-7.81 (m, 2H), 7.74-7.65 (m, 4H), 7.44 (t, *J*=8.4 Hz, 1H), 7.05 (d, *J*=7.2 Hz, 1H), 6.72 (d, *J*=8.0 Hz, 1H), 6.37 (s, 1H), 5.75 (s, 2H), 5.30 (s, 2H), 4.85-4.84 (m, 1H), 3.87-3.83 (m, 4H), 2.69-2.67 (m, 2H), 2.27-2.22 (m, 2H), 1.79-1.77 (m, 2H), 1.47-1.41 (m, 2H), 1.04-1.00 (m, 1H), 0.51-0.46 (m, 2H), 0.35-0.31 (m, 2H).

Example 67. Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-5-oxopyrrolidin-2-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C34)


1). Synthesis of methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-(((1-ethyl)-5-oxopyrrolidin-2-yl)methyl)amino)benzoate




[0650] The mixture of methyl 4-(2-chloroacetamido)-3-(((1-ethyl-5-oxopyrrolidin-2-yl)methyl)amino)benzoate (100 mg, 0.27 mmol), 3-fluoro-4-((6-(piperidin-4-oxo)pyridin-2-yl)methoxy)benzonitrile (88 mg, 0.27 mmol) and potassium carbonate (75 mg, 0.54 mmol) in N,N-dimethylformamide (2 mL) was stirred at room temperature for 16 hours. After completion, the mixture was diluted with water (30 mL) and extracted with ethyl acetate (10 mL×2). The combined organic layers were washed with brine (20 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica

gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-(((1-ethyl)-5-oxopyrrolidin-2-yl)methyl)amino)benzoate (90 mg, 50.7% yield). LC-MS m/z : 659 $[M+H]^+$.

2). Synthesis of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-5-oxopyrrolidin-2-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate




-continued



[0651] The mixture of 4-(2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-(((1-ethyl-5-oxopyrrolidin-2-yl)methyl)amino)benzoate (90 mg, 0.14 mmol) in toluene (3 mL) and acetic acid (0.5 mL) was stirred at 110° C. for 3 hours. After completion, the solvent was removed by concentration to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyri-

din-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-5-oxopyrrolidin-2-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (70 mg, 78.1% yield). LC-MS m/z : 641 $[M+H]^+$.

3). Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-5-oxopyrrolidin-2-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid




[0652] To a mixture of methyl 2-((4-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxypiperidin-1-yl)methyl)-1-(((1-ethyl-5-oxopyrrolidin-2-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (70 mg, 0.11 mmol) in THF (5 mL) and water (5 mL) lithium hydroxide (8 mg, 0.33 mmol) was added at room temperature. The reaction was stirred at room temperature for 16 hours. After completion and concentrated to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: Sun-Fire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-5-oxopyrrolidin-2-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (32.33 mg, 46.9% yield). LC-MS *m/z*: 627 [M+H]⁺.

[0653] ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.27 (s, 1H), 7.90-7.82 (m, 2H), 7.75-7.66 (m, 3H), 7.45 (t, *J*=8.8 Hz, 1H), 7.06 (d, *J*=7.2 Hz, 1H), 6.74 (d, *J*=8.4 Hz, 1H), 5.31 (s, 2H), 4.93-4.91 (m, 1H), 4.77-4.72 (m, 1H), 4.51-4.46 (m, 1H), 4.33-4.31 (m, 1H), 3.91-3.81 (m, 2H), 3.51-3.46 (m, 1H), 2.76-2.67 (m, 3H), 2.47-2.39 (m, 1H), 2.34-2.30 (m, 2H), 2.16-2.14 (m, 1H), 1.94-1.79 (m, 4H), 1.62-1.55 (m, 2H), 1.00 (t, *J*=7.2 Hz, 3H).

Example 68. Synthesis of (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethylpyrrolidin-2-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C35)


1). Synthesis of methyl (S)-4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-ethylpyrrolidin-2-yl)methyl)amino)benzoate



[0654] The mixture of methyl (S)-4-(2-chloroacetamido)-3-((1-ethylpyrrolidin-2-yl)methyl)aminobenzoate (140 mg, 0.40 mmol), 3-fluoro-4-((6-((piperidin-4-oxy)pyridin-2-yl)methoxy)benzonitrile (142 mg, 0.44 mmol) and potassium carbonate (165 mg, 1.20 mmol) in N,N-dimethylformamide (5 mL) was stirred at room temperature for 16 hours. After completion, it was diluted with water (50 mL) and extracted with ethyl acetate (50 mL×3). The combined organic layers washed with brine (120 mL×2). The organic layer was concentrated in vacuo and purified by silica gel column


chromatography (dichloromethane/methanol=13/1) to give methyl (S)-4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-ethylpyrrolidin-2-yl)methyl)amino)benzoate (180 mg, 70.4% yield). LC-MS m/z: 645 [M+H]⁺.

2). Synthesis of methyl (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethylpyrrolidin-2-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate




[0655] To a mixture of methyl (S)-4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-ethylpyrrolidin-2-yl)methyl)amino)benzoate (180 mg, 0.28 mmol) in dioxane (10 mL) acetic acid (2 mL) was added. The resulting mixture was stirred and reacted at 100° C. for 3 hours. After completion, the reaction mixture was concentrated to give a residue in vacuo. The residue was purified by silica gel (dichloromethane/methanol=10/1) to obtain methyl (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethylpyrrolidin-2-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (80 mg, 45.7% yield). LC-MS m/z: 627 [M+H]⁺.

3). Synthesis of (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethylpyrrolidin-2-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid



-continued




[0656] To a mixture of methyl (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxypiperidin-1-yl)methyl)-1-((1-ethylpyrrolidin-2-yl)methyl)-1H-benzod[1,2-b]imidazole-6-carboxylate (80 mg, 0.13 mmol) was dissolved in H₂O (4 mL) and THF (4 mL) lithium hydroxide (32 mg, 1.30 mmol) was added at room temperature. The reaction mixture was stirred at room temperature for 16 hours. After completion, the reaction solution was adjusted to pH=5-6 with formic acid. The mixture was concentrated to give a residue in vacuo to remove the solvent. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethylpyrrolidin-2-yl)methyl)-1H-benzod[1,2-b]imidazole-6-carboxylic acid (19.20 mg, 24.6% yield). LC-MS m/z: 613 [M+H]⁺.

[0657] ¹H NMR (400 MHz, DMSO-d₆): δ 8.21 (s, 1H), 7.88 (dd, J=11.2 Hz, 2.0 Hz, 1H), 7.79 (dd, J=8.8 Hz, 1.2 Hz, 1H), 7.73 (t, J=8.0 Hz, 1H), 7.68-7.62 (m, 2H), 7.45 (t, J=8.8 Hz, 1H), 7.06 (d, J=7.6 Hz, 1H), 6.74 (d, J=8.4 Hz, 1H), 5.31 (s, 2H), 4.93-4.89 (m, 1H), 4.44-4.39 (m, 1H), 4.33-4.28 (m, 1H), 4.04 (d, J=13.2 Hz, 1H), 3.76 (d, J=13.2 Hz, 1H), 3.11-3.07 (m, 2H), 2.77-2.70 (m, 2H), 2.34-2.26 (m, 2H), 2.22-2.14 (m, 2H), 1.91-1.89 (m, 2H), 1.80-1.62 (m, 3H), 1.61-1.55 (m, 3H), 0.80 (t, J=7.2 Hz, 3H).

Example 69. Synthesis of (S)-2-((4-((3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-4-methyl-1-((oxetan-2-ylmethyl)amino)phenyl)-1H-benzod[1,2-b]imidazole-6-carboxylic acid (Compound C36)


1). Synthesis of (S)-N-(4-bromo-2-methyl-6-((oxetan-2-ylmethyl)amino)phenyl)-2-((4-((3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)acetamide



[0658] The mixture of (S)—N-(4-bromo-2-methyl-6-((oxetan-2-ylmethyl)amino)phenyl)-2-chloroacetamide (98 mg, 0.28 mmol), 3-fluoro-4-((3-(piperidin-4-oxy)benzyl)oxy)benzonitrile (119 mg, 0.37 mmol) and potassium carbonate (77 mg, 0.56 mmol) in N,N-dimethylformamide (3 mL) was stirred at room temperature for 16 hours. After completion, the reaction was quenched by adding water (5 mL) and extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (20 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was


purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain (S)—N-(4-bromo-2-methyl-6-((oxetan-2-ylmethyl)amino)phenyl)-2-(4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)acetamide (65 mg, 35.7% yield) as yellow solid. LC-MS m/z : 639 $[M+H]^+$.

2). Synthesis of (S)-4-((3-((1-((6-bromo-4-methyl-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-2-yl)methyl)-piperidin-4-yl)oxy)benzyl)oxy)-3-fluorobenzonitrile




[0659] To a solution of (S)—N-(4-bromo-2-methyl-6-((oxetan-2-ylmethyl)amino)phenyl)-2-(4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)acetamide (65 mg, 0.10 mmol) in dioxane (5 mL) acetic acid (0.5 mL) was added at room temperature. The resulting mixed solution was stirred at 100° C. for 2 hours. After completion, the reaction solution was concentrated to give a residue under reduced pressure. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain compound (S)-4-((3-((1-((6-bromo-4-methyl-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazol-2-yl)methyl)piperidin-4-yl)oxy)benzyl)oxy)-3-fluorobenzonitrile (60 mg; 96.9% yield) as yellow solid. LC-MS m/z : 621 $[M+H]^+$.

3). Synthesis of methyl (S)-2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-4-methyl-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate





-continued



[0660] The mixture of (S)-4-((3-((1-((6-bromo-4-methyl-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazol-2-yl)methyl)piperidin-4-yl)oxy)benzyl)oxy)-3-fluorobenzonitrile (60 mg, 0.10 mmol), 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (7 mg, 0.01 mmol) and potassium acetate (29 mg, 0.30 mmol) in N,N-dimethylformamide (1 mL) and methanol (1 mL) was stirred at 90° C. for 16 hours under carbon monoxide (58.76 psi) atmosphere. After completion, the reaction mixture was concentrated to give a residue. The residue was purified by silica gel column chromatography


(dichloromethane/methanol=20/1) to obtain methyl (S)-2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-4-methyl-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (11 mg, 17.8% yield). LC-MS m/z: 599 [M+H]⁺.

4. Synthesis of (S)-2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidine-1-yl)methyl)-4-methyl-1-(oxocyclobutan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid methyl ester



[0661] The mixture of (S)-2-(chloromethyl)-4-methyl-1-(oxocyclobutan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid methyl ester (20 mg, 0.07 mmol), 3-fluoro-4-(3-(piperidin-4-oxy benzyl oxy) benzonitrile (30 mg, 0.09 mmol) and potassium carbonate (29 mg, 0.21 mmol) in N, N-dimethylformamide (1 mL) was stirred at 60° C. for 2 hours. After completion, the mixture was diluted with water (3 mL), and extract with ethyl acetate (15 mL×3). The combined organic layer was washed with brine (10 mL×3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain (S)-2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidine-1-yl)methyl)-4-methyl-1-(oxocyclobutan-2-yl methyl)-1H-benzo[d]imidazole-6-carboxylic acid methyl ester (11 mg, yield: 28.6%). LC-MS m/z: 599 [M+H]⁺.

5). Synthesis of (S)-2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-4-methyl-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid





[0662] To a mixture of methyl (S)-2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-4-methyl-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (22 mg, 0.04 mmol) in THF (4 mL) and water (4 mL) lithium hydroxide (16 mg, 0.40 mmol) was added at room temperature. The resulting mixture solution was stirred and reacted at room temperature for 5 hours. After completion, the resulting mixture was adjusted to pH=5-6 with acetic acid (1 M) and extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (10 mL), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by reverse-phase flash chromatography under the following conditions (Column: Spherical C18, 20-40 μm, 40 g; Mobile Phase A: 10 mM NH₄OH in water; Mobile Phase B: acetonitrile; Flow rate: 40 mL/min; Gradient: 20% B-50% B in 20 min; Detector: 254 nm) to obtain (S)-2-((4-(3-((4-cyano-2-fluorophenoxy)methyl)phenoxy)piperidin-1-yl)methyl)-4-methyl-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (9.33 mg, 39.9% yield). LC-MS m/z: 585 [M+H]⁺.

[0663] ¹H NMR (400 MHz, DMSO-d₆) δ 7.90-7.85 (m, 2H), 7.69-7.63 (m, 2H), 7.42 (t, J=8.4 Hz, 1H), 7.30 (t, J=8.0 Hz, 1H), 7.05 (m, 1H), 7.00 (d, J=7.2 Hz, 1H), 6.95 (dd, J=8.0 Hz, 1.6 Hz, 1H), 5.25 (s, 2H), 5.11-5.05 (m, 1H), 4.70-4.65 (m, 1H), 4.58-4.54 (m, 1H), 4.51-4.45 (m, 1H), 4.41-4.33 (m, 2H), 3.89 (d, J=13.6 Hz, 1H), 3.77 (d, J=13.6 Hz, 1H), 2.74-2.63 (m, 3H), 2.49 (s, 3H), 2.45-2.35 (m, 3H), 1.93 (s, 2H), 1.64-1.60 (m, 2H).

Example 70. Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)thio)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C37)

1). Synthesis of methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)mercapto)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate







[0664] The mixture of methyl 4-(2-chloroacetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (100 mg, 0.29 mmol), 3-fluoro-4-((6-(piperidin-4-ylthio)pyridin-2-yl)methoxy)benzonitrile (99 mg, 0.29 mmol) and potassium carbonate (80 mg, 0.58 mmol) in N,N-dimethylformamide (5 mL) was stirred at room temperature for 16 hours. After completion, the resulting mixture was poured into saturated ammonium chloride solution (20 mL) and extracted with ethyl acetate (10 mL $\times$ 2). The combined organic layers were washed with brine (20 mL $\times$ 3) and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 4-(2-(4-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)mercapto)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (70 mg, 38.1% yield). LC-MS m/z : 658 $[M+H]^+$.

2). Synthesis of methyl 12-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)mercapto)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate

[0665] To a mixture of methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)mercapto)piperidin-1-yl)acetamido)-methyl 3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (70 mg, 0.11 mmol) in toluene (5 mL) acetic acid (1 mL) was added at room temperature. The resulting mixture was stirred at 110° C. for 3 hours. After completion, the solvent was removed by concentration to give a residue in vacuo. The residue was purified by preparative thin layer chromatography (dichloromethane/methanol=30/1) to give methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)mercapto)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (60 mg, 85.4% yield). LC-MS m/z : 640 $[M+H]^+$.



3). Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)mercapto)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid




[0666] To a solution of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)mercapto)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (60 mg, 0.09 mmol) in H₂O (5 mL) and THF (5 mL) lithium hydroxide (7 mg, 0.27 mmol) was added at room temperature. The reaction mixture was stirred at room temperature for 16 hours; after completion concentrated to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: Sun-Fire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% NH₄OH in H₂O solvent system; Detection

Wavelength 254 nm/214 nm; Flow rate 20 mL/min to obtain 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)mercapto)piperidin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (28.45 mg, 50.6% yield). LC-MS m/z: 626 [M+H]⁺.

[0667] ¹H NMR (400 MHz, DMSO-d₆): δ 8.06 (s, 1H), 7.89 (dd, J=11.2 Hz, 2.0 Hz, 1H), 7.81 (dd, J=8.8 Hz, 1.6 Hz, 1H), 7.69-7.65 (m, 4H), 7.41 (t, J=8.8 Hz, 1H), 7.22-7.19 (m, 2H), 6.38 (s, 1H), 5.69 (s, 1H), 5.36 (s, 2H), 4.02-3.96 (m, 2H), 3.79 (s, 2H), 3.68-3.60 (m, 1H), 2.71-2.68 (m, 2H), 2.19-2.14 (m, 2H), 1.86-1.84 (m, 2H), 1.41-1.32 (m, 2H), 1.16 (t, J=7.2 Hz, 3H).

Example 71. Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)methyl)piperazin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C38)


1). Synthesis of methyl 4-(2-(4-((6-((4-Cyano-2-fluorophenoxy)methyl)pyridin-2-yl)methyl)piperazin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate




[0668] The mixture of potassium carbonate (135 mg, 0.98 mmol), methyl 4-(2-chloroacetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (172 mg, 0.49 mmol) and 3-fluoro-4-((6-(piperazin-1-yl)methyl)pyridin-2-yl)methoxy)benzonitrile (176 mg, 0.54 mmol) in *N,N*-dimethylformamide (5 mL) was stirred and reacted at room temperature for 3 hours. After completion, the mixture was diluted with water (5 mL) and extracted with ethyl acetate (40 mL $\times$ 3). The combined organic layers were washed with brine (30 mL $\times$ 2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The

residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)methyl)piperazin-1-yl)acetamido)-3-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (120 mg, 38.3% yield). LC-MS *m/z*: 641 [M+H]⁺.

2). Synthesis of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)methyl)piperazin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate




-continued



[0669] To a solution of methyl 4-(2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)methyl)piperazin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)amino)benzoate (120 mg, 0.19 mmol) in toluene (6 mL) acetic acid (1 mL) was added at room temperature. The resulting mixture solution was stirred and reacted at 110° C. for 2 hours. After completion, the reaction solution was concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain methyl 2-((4-((6-((4-cyano-2-fluorophe-

noxy)methyl)pyridin-2-yl)methyl)piperazin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 84.5% yield). LC-MS m/z : 623 $[M+H]^+$.

3). Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)methyl)piperazin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid




[0670] To a mixture of methyl 2-((4-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)methyl)piperazin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 0.16 mmol) in water (3.0 mL) and THF (3.0 mL) lithium hydroxide monohydrate (67 mg, 1.60 mmol) was added at room temperature. The resulting mixture solution was stirred and reacted at room temperature for 5 hours. After completion, the resulting mixture was adjusted to pH=5-6 using hydrochloric acid (1M) and concentrated to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm; Gradient elution with ACN/0.1% NH₄OH in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)methyl)piperazin-1-yl)methyl)-1-((1-ethyl-1H-imidazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (18.66 mg, 19.2% yield). LC-MS m/z: 609 [M+H]⁺.

[0671] ¹H NMR (400 MHz, DMSO-d₆) δ 8.05 (s, 1H), 7.90-7.79 (m, 3H), 7.68-7.62 (m, 3H), 7.46-7.38 (m, 3H), 6.38 (s, 1H), 5.68 (s, 2H), 5.33 (s, 2H), 4.00-3.95 (m, 2H), 3.79 (s, 2H), 3.53 (s, 2H), 2.50-2.42 (m, 4H), 2.33-2.28 (m, 4H), 1.15 (t, J=7.2 Hz, 3H).

Example 72. Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-pyrazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C39)


1). Synthesis of 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-pyrazol-5-yl)methyl)amino) methyl benzoate




[0672] The mixture of methyl 4-(2-chloroacetamide)-3-(1-ethyl-1H-pyrazol-5-yl)methylaminobenzoate (50 mg, 0.14 mmol), 3-fluoro-4-((6-((piperidin-4-oxo)pyridin-2-yl)methoxy)benzonitrile (46 mg, 0.14 mmol) and potassium carbonate (39 mg, 0.28 mmol) in N,N-dimethylformamide (2 mL) was stirred at room temperature for 16 hours. After completion, it was diluted with water (10 mL) and extracted with ethyl acetate (5 mL $\times$ 2). The combined organic layers

were washed with brine (5 mL $\times$ 2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamide)-3-((1-ethyl-1H-pyrazol-5-yl)methyl)amino)benzoate (60 mg, 66.8% yield). LC-MS m/z: 642 [M+H]⁺.


2). Synthesis of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-pyrazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate



[0673] The mixture of methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-ethyl-1H-pyrazol-5-yl)methyl)amino)benzoate (60 mg, 0.09 mmol) in toluene (2 mL) and acetic acid (0.5 mL) was stirred at 110° C. for 3 hours. After completion, the solvent was removed to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-pyrazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (50 mg, 89.0% yield). LC-MS m/z: 624 [M+H]⁺.



3). Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-pyrazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid




[0674] To a mixture of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-pyrazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (50 mg, 0.08 mmol) in THF (5 mL) and water (5 mL) lithium hydroxide (6 mg, 0.24 mmol) was added at room temperature. The reaction was stirred at room temperature for 16 hours. After completion, the resulting mixture was adjusted to pH=5-6 with formic acid. The mixture was concentrated to give a residue. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μm 19×250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-1H-pyrazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (14.38 mg, 29.5% yield). LC-MS m/z: 610 [M+H]⁺.

[0675] ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 8.06 (s, 1H), 7.89-7.82 (m, 2H), 7.73-7.64 (m, 3H), 7.43 (t, $J=8.4$ Hz, 1H), 7.29 (s, 1H), 7.04 (d, $J=7.2$ Hz, 1H), 6.71 (d, $J=8.4$ Hz, 1H), 5.80 (s, 2H), 5.51 (s, 1H), 5.29 (s, 2H), 4.81 (s, 1H), 4.25-4.20 (m, 2H), 3.82 (s, 2H), 2.65-2.62 (m, 2H), 2.22-2.17 (m, 2H), 1.73-1.71 (m, 2H), 1.39-1.37 (m, 2H), 1.30 (t, $J=7.2$ Hz, 3H).

Example 73. Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-3-methyl-1H-pyrazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C40)


1). Synthesis of methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-ethyl-3-methyl-1H-pyrazol-5-yl)methyl)amino)benzoate




[0676] The mixture of methyl 4-(2-chloroacetamide)-3-((1-ethyl-3-methyl-1H-pyrazol-5-yl)methyl)amino)benzoate (200 mg, 0.55 mmol), 3-fluoro-4-((6-(piperidin-4-oxy)pyridin-2-yl)methoxy)benzonitrile (180 mg, 0.55 mmol) and potassium carbonate (152 mg, 1.10 mmol) in N,N -dimethylformamide (5 mL) was stirred at room temperature for 16 hours. After completion, it was diluted with water (30 mL) and extracted with ethyl acetate (10 mL $\times$ 3). The combined organic layers were washed with brine (10 mL $\times$ 3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica

gel column chromatography (dichloromethane/methanol=20/1) to obtain methyl 4-(2-(4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-ethyl-3-methyl-1H-pyrazol-5-yl)methyl)amino)benzoate (200 mg, 55.4% yield). LC-MS m/z : 656 $[\text{M}+\text{H}]^+$.

2). Synthesis of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-3-methyl-1H-pyrazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate





-continued



[0677] The mixture of methyl 4-(2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)acetamido)-3-((1-ethyl-3-methyl-1H-pyrazol-5-yl)methyl)amino)benzoate (200 mg, 0.30 mmol) in toluene (2 mL) and acetic acid (0.5 mL) was stirred at 110° C. for 3 hours. After completion, the solvent was removed by concentration to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=20/1) to obtain 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-3-methyl-1H-pyrazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (150 mg, 78.4% yield). LC-MS m/z: 638 [M+H]⁺.

3). Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-3-methyl-1H-pyrazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid




[0678] To a mixture of methyl 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxypiperidin-1-yl)methyl)-1-((1-ethyl-3-methyl-1H-pyrazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylate (150 mg, 0.24 mmol) in THF (5 mL) and water (5 mL) lithium hydroxide (6 mg, 0.24 mmol) was added at room temperature and the reaction was stirred at room temperature for 16 hours. After completion, the resulting mixture was adjusted to pH=5-6 with formic acid. The solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 um 19x250 mm; Gradient elution with ACN/0.1% FA in H₂O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-((1-ethyl-3-methyl-1H-pyrazol-5-yl)methyl)-1H-benzo[d]imidazole-6-carboxylic acid (59.70 mg, 39.9% yield). LC-MS m/z: 624 [M+H]⁺.

[0679] ¹H NMR (400 MHz, DMSO-d₆): δ 8.09 (s, 1H), 7.89-7.82 (m, 2H), 7.74-7.64 (m, 3H), 7.43 (t, J=8.8 Hz, 1H), 7.05 (d, J=7.2 Hz, 1H), 6.72 (d, J=8.0 Hz, 1H), 5.75 (s, 2H), 5.30 (s, 2H), 5.26 (s, 1H), 4.83 (s, 1H), 4.16-4.11 (m, 2H), 3.81 (s, 2H), 2.66-2.63 (m, 2H), 2.23-2.19 (m, 2H), 2.00 (s, 3H), 1.75-1.73 (m, 2H), 1.43-1.40 (m, 2H), 1.29 (t, J=7.2 Hz, 3H).



Example 74. Synthesis of (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)thio)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C41)

1). Synthesis of methyl (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)thio)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate



[0680] A mixture of 3-fluoro-4-((6-(piperidin-4-ylmercapto)pyridin-2-yl)methoxy)benzonitrile (86 mg, 0.29 mmol) and methyl (S)-2-(chloromethyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 0.29 mmol) in N,N-dimethylformamide (5 mL) potassium carbonate (80 mg, 0.58 mmol) was added at room temperature. The reaction was stirred at room temperature for 16 hours. Upon completion of the reaction, the resulting mixture was poured into brine (20 mL) and extracted with ethyl acetate (20 mL×2). The combined organic layers were concentrated and then purified by silica gel column chromatography (dichloromethane/methanol=10/1) to give methyl (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)thio)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 57.5% yield). LC-MS m/z : 602 $[M+H]^+$.

2). Synthesis of (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)thio)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid




[0681] To a mixture of methyl (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)mercapto)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (100 mg, 0.17 mmol) in THF (5 mL) and water (5 mL) lithium hydroxide (40 mg, 1.70 mmol) was added at room temperature. The reaction was stirred at room temperature for 16 hours. After completion, the resulting mixture was adjusted to pH=5-6 using hydrochloric acid (1M). The solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19×250 mm; Gradient elution with ACN/0.1% NH_4OH in H_2O solvent system; Detection Wavelength: 254 nm/214 nm; Flow rate: 20 mL/min) to obtain (S)-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)thio)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (37.32 mg, 37.4% yield). LC-MS m/z : 588 $[M+H]^+$.

[0682] ^1H NMR (400 MHz, DMSO- d_6) δ 8.26 (s, 1H), 7.89 (dd, $J=11.2$ Hz, 1.6 Hz, 1H), 7.81 (d, $J=8.4$ Hz, 1H), 7.70-7.63 (m, 3H), 7.43 (t, $J=8.4$ Hz, 1H), 7.22 (t, $J=8.4$ Hz, 2H), 5.37 (s, 2H), 5.08-5.06 (m, 1H), 4.80-4.75 (m, 1H), 4.65-4.61 (m, 1H), 4.50-4.48 (m, 1H), 4.38-4.35 (m, 1H), 3.92 (d, $J=13.6$ Hz, 1H), 3.78-3.70 (m, 2H), 2.82-2.68 (m, 3H), 2.42-2.38 (m, 1H), 2.30-2.23 (m, 2H), 1.94 (s, 2H), 1.58-1.53 (m, 2H).

Example 75. Synthesis of 2-(((4-((6-(2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-indole-6-carboxylic acid (Compound C42)


1). Synthesis of oxetan-2-ylmethanesulfonate



[0683] To a mixture of oxetan-2-ylmethanol (250 mg, 2.84 mmol) and triethylamine (574 mg, 5.68 mmol) in anhydrous dichloromethane (15 mL) methanesulfonyl chloride (557 mg, 5.68 mmol) was added dropwise at 0°C under nitrogen atmosphere. The mixture was stirred at for 3 hours. After completion, water (10 mL) was added to quench the reaction, and it was extracted with ethyl acetate (20 mL $\times$ 3). The combined organic layers were washed with brine (20 mL $\times$ 3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo to obtain oxetan-2-ylmethanesulfonate (300 mg, yield: 63.7%).

[0684] ^1H -NMR (400 MHz, CDCl_3) δ 5.05-5.01 (m, 1H), 4.70-4.66 (m, 1H), 4.61-4.56 (m, 1H), 4.36 (d, $J=4.0$ Hz, 2H), 3.11 (s, 3H), 2.81-2.74 (m, 1H), 2.66-2.61 (m, 1H).


2). Synthesis of 1-(tert-butyl) 2-methyl 6-bromo-1H-indole-1,2-dicarboxylate



[0685] To a solution of methyl 6-bromo-1H-indole-2-carboxylate (2.00 g, 7.91 mmol) in dichloromethane (30 mL) 4-dimethylaminopyridine (193 mg, 1.58 mmol) and di-tert-butyl dicarbonate (2.41 g, 11.07 mmol) were added at room temperature. The resulting mixture was stirred at room temperature for 16 hours. After completion, the mixture was concentrated in vacuo to obtain a crude product. The crude product was further purified by silica gel column chromatography (petroleum ether/ethyl acetate=5/1) to obtain 1-(tert-butyl) 2-methyl 6-bromo-1H-indole-1,2-dicarboxylate (2.2 g, 78.8% yield).

[0686] ^1H NMR (400 MHz, CDCl_3) δ 8.31 (s, 1H), 7.46 (d, $J=8.4$ Hz, 1H), 7.38 (dd, $J=8.4$, 2.0 Hz, 1H), 7.05 (d, $J=0.8$ Hz, 1H), 3.92 (s, 3H), 1.62 (s, 9H).


3). Synthesis of tert-butyl 6-bromo-2-(hydroxymethyl)-1H-indole-1-carboxylate



[0687] To a solution of 1-(tert-butyl) 2-methyl 6-bromo-1H-indole-1,2-dicarboxylate (1.00 g, 2.83 mmol) in dichloromethane (15 mL) diisobutylaluminum hydride (7 mL, 7.08 mol) was slowly added at -78°C under nitrogen atmosphere. The resulting mixture was stirred at -40°C for 2 hours. After completion, the reaction was quenched by adding methanol (15 mL) and water (9 mL) and then extracted with ethyl acetate (30 mL $\times$ 3). The combined organic layers were washed with brine (50 mL $\times$ 2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=1/1) to obtain tert-butyl 6-bromo-2-(hydroxymethyl)-1H-indole-1-carboxylate (650 mg, 70.7% yield).

[0688] ^1H NMR (400 MHz, CDCl_3) δ 8.18 (d, $J=0.8$ Hz, 1H), 7.37-7.32 (m, 2H), 6.54 (s, 1H), 4.79 (d, $J=5.6$ Hz, 2H), 3.62 (s, 1H), 1.73 (s, 9H).


4). Synthesis of tert-butyl ester 6-bromo-2-(chloromethyl)-1H-indole-1-carboxylate



[0689] To a mixture of tert-butyl 6-bromo-2-(hydroxymethyl)-1H-indole-1-carboxylate (650 mg, 2.00 mmol), triethylamine (343 mg, 3.40 mmol), and lithium chloride (840 mg, 20.00 mmol) in dichloromethane (20 mL) methanesulfonyl chloride (333 mg, 3.40 mmol) was slowly added at room temperature. The resulting mixture was stirred at room temperature for 16 hours. After completion, water (10 mL) was added to quench the reaction, and the mixture was extracted with ethyl acetate (30 mL $\times$ 3). The combined organic layers were washed with brine (50 mL $\times$ 2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=15/1) to obtain tert-butyl 6-bromo-2-(chloromethyl)-1H-indole-1-carboxylate (500 mg, 73.0% yield).

[0690] ^1H NMR (400 MHz, CDCl_3) δ 8.39 (s, 1H), 7.36-7.35 (m, 2H), 6.66 (d, $J=0.8$ Hz, 1H), 4.95 (dd, $J=0.4$ Hz, 2H), 1.73 (s, 9H).


5). Synthesis of tert-butyl 6-bromo-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1H-indole-1-carboxylate




[0691] The mixture of tert-butyl 6-bromo-2-(chloromethyl)-1H-indole-1-carboxylate (300 mg, 0.87 mmol), 3-fluoro-4-((6-(piperidine-4-oxy)pyridin-2-yl)methoxy) benzonitrile (284 mg, 0.87 mmol), and potassium carbonate (360 mg, 2.61 mmol) in N,N-dimethylformamide (5 mL) was stirred and reacted at 60° C. for 3 hours. After completion, the resulting mixture was cooled to room temperature, then poured into brine (5 mL) and extracted with ethyl acetate (2×20 mL). The combined organic layers were washed with brine (2×20 mL) and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=1/1) to obtain tert-butyl 6-bromo-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridine-2-yl)oxy)piperidin-1-yl)methyl)-1H-indole-1-carboxylate (400 mg, 63.1% yield). LC-MS m/z : 635 $[M+H]^+$.

[0692] 1H NMR (400 MHz, $CDCl_3$) δ 8.26 (s, 1H), 7.58 (t, $J=8.0$ Hz, 1H), 7.40-7.30 (m, 4H), 7.10 (t, $J=8.0$ Hz, 1H), 7.00 (d, $1=7.2$ Hz, 1H), 6.66 (d, $1=8.0$ Hz, 1H), 6.55 (s, 1H), 5.18 (s, 2H), 5.05-5.03 (m, 1H), 3.86 (s, 2H), 2.85-2.82 (m, 2H), 2.40-2.36 (m, 2H), 2.05-2.00 (m, 2H), 1.82-1.78 (m, 2H), 1.70 (s, 9H).

6). Synthesis of 4-((6-((1-((6-bromo-1H-indol-2-yl)methyl)piperidin-4-yl)oxy)pyridin-2-yl)methoxy)-3-fluorobenzonitrile




-continued



[0693] To a solution of tert-butyl 6-bromo-2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1H-indole-1-carboxylate (400 mg, 0.63 mmol) in dichloromethane (10 mL) trifluoroacetic acid (4 mL) was added at room temperature. And the mixture was stirred at room temperature for 3 hours. After completion, the reaction mixture was concentrated in vacuo to obtain a crude product. The crude product was further purified by silica gel column chromatography (dichloromethane/methanol=15/1) to obtain 4-((6-((1-((6-bromo-1H-indol-2-yl)methyl)piperidin-4-yl)oxy)pyridin-2-yl)methoxy)-3-fluorobenzonitrile (280 mg, 82.5% yield).

[0694] 1H NMR (400 MHz, $CDCl_3$) δ 11.09 (s, 1H), 7.67-7.62 (m, 2H), 7.42-7.37 (m, 3H), 7.20 (dd, $J=8.4, 1.6$ Hz, 1H), 7.10-7.03 (m, 2H), 6.70 (d, $J=8.4$ Hz, 1H), 6.48 (s, 1H), 5.34 (s, 1H), 5.15 (s, 2H), 4.18 (s, 2H), 3.08 (s, 2H), 2.96 (s, 1H), 2.88 (s, 1H), 2.31-2.22 (m, 2H), 1.31-1.24 (m, 2H).



7). Synthesis of 4-((6-((1-((6-bromo-1-(oxetan-2-ylmethyl)-1H-indol-2-yl)methyl)piperidin-4-yl)oxy)pyridin-2-yl)methoxy)-3-fluorobenzonitrile



[0695] The mixture of 4-((6-((1-((6-bromo-1H-indol-2-yl)methyl)piperidin-4-yl)oxy)pyridin-2-yl)methoxy)-3-fluorobenzonitrile (280 mg, 0.52 mmol), oxetan-2-ylmethanesulfonate (259 mg, 1.56 mmol) and cesium carbonate (509 mg, 1.56 mmol) in N,N-dimethylformamide (5 mL) was stirred at 110° C. for 16 hours. Upon completion of the reaction, the resulting mixture was cooled to room temperature, poured into brine (5 mL) and then extracted with ethyl acetate (2×20 mL). The combined organic layers were washed with brine (2×20 mL) and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (dichloromethane/methanol=10/1) to obtain 4-((6-((1-((6-bromo-1-(oxetan-2-ylmethyl)-1H-indol-2-yl)methyl)piperidin-4-yl)oxy)pyridin-2-yl)methoxy)-3-fluorobenzonitrile (150 mg, 48.0% yield). LC-MS *m/z*: 607 [M+H]⁺.

[0696] ¹H NMR (400 MHz, CDCl₃) δ 7.60-7.55 (m, 2H), 7.41-7.36 (m, 3H), 7.17 (dd, J=8.4, 1.6 Hz, 1H), 7.09 (t, J=8.0 Hz, 1H), 7.00 (d, J=7.2 Hz, 1H), 6.65 (d, J=8.0 Hz, 1H), 6.36 (s, 1H), 5.18-5.14 (m, 3H), 5.02 (s, 1H), 4.64-4.48 (m, 3H), 4.43-4.38 (m, 1H), 3.75-3.67 (m, 2H), 2.75-2.64 (m, 3H), 2.47-2.42 (m, 1H), 2.32-2.24 (m, 2H), 1.95 (s, 2H), 1.74 (d, J=8.8 Hz, 2H).

8). Synthesis of 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-indole-6-carboxylic acid




[0697] To a mixture of 4-((6-((1-((6-bromo-1-(oxetan-2-ylmethyl)-1H-indole-2-yl)methyl)piperidin-4-yl)oxy)pyridin-2-yl)methoxy)-3-fluorobenzonitrile (150 mg, 0.25 mmol) in N,N-dimethylformamide (5 mL) and water (5 mL) triethylamine (76 mg, 0.75 mmol) and 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (18 mg, 0.025 mmol) were added at room temperature. The resulting mixture was heated to 90° C. under carbon monoxide gas atmosphere. The reaction was stirred for 16 hours. After completion, water (15 mL) was added to quench the reaction, and extracted with ethyl acetate (30 mL×3). The combined organic layers were washed with brine (30 mL×2), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by reverse-phase flash chromatography under the following conditions (Column: Spherical C18, 20-40 m, 40 g; Mobile Phase A: 10 mM NH₄OH in water; Mobile Phase B: acetonitrile; Flow rate: 40 mL/min; Gradient: 20% B-50% B in 20 min; Detector: 254 nm) to obtain 2-((4-((6-((4-cyano-2-fluorophenoxy)methyl)pyridin-2-yl)oxy)piperidin-1-yl)methyl)-1-(oxetan-2-ylmethyl)-1H-indole-6-carboxylic acid (9.07 mg, 6.4% yield). LC-MS *m/z*: 571 [M+H]⁺.

[0698] ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.13 (s, 1H), 7.89 (dd, $J=11.2, 2.0$ Hz, 1H), 7.73 (t, $J=8.0$ Hz, 1H), 7.67 (d, $J=8.4$ Hz, 1H), 7.61 (d, $J=8.4$ Hz, 1H), 7.52-7.43 (m, 2H), 7.05 (d, $J=7.2$ Hz, 1H), 6.73 (d, $J=8.4$ Hz, 1H), 6.45 (s, 1H), 5.32 (s, 2H), 5.05-5.02 (m, 1H), 4.91-4.87 (m, 1H), 4.71-4.66 (m, 1H), 4.56-4.45 (m, 2H), 4.37-4.32 (m, 1H), 3.80 (d, $J=13.6$ Hz, 1H), 3.63 (d, $J=13.6$ Hz, 1H), 2.73-2.65 (m, 3H), 2.43-2.38 (m, 1H), 2.23-2.18 (m, 2H), 1.89-1.88 (m, 2H), 1.62-1.57 (m, 2H).


Example 76. Synthesis of (S)-2-(4-(((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)-2,5-difluorobenzyl)-4-fluoro-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid (Compound C43)

1). Synthesis of
2-(4-Bromo-2,5-difluorophenoxy)-6-fluoropyridine




[0699] 4-Bromo-2,5-difluorophenol (2.00 g, 9.57 mmol), 2,6-difluoropyridine (2.20 g, 19.14 mmol) and cesium carbonate (6.24 g, 19.14 mmol) were dissolved in acetonitrile (50 mL) at room temperature. The resulting mixture was warmed to 80° C. and stirred for 16 hours. After completion, the solvent was removed by concentration to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=4/1) to obtain 2-(4-bromo-2,5-difluorophenoxy)-6-fluoropyridine (2.00 g, 68.7% yield). LC-MS m/z : 304 $[\text{M}+\text{H}]^+$.

2). Synthesis of 4-((6-((4-bromo-2,5-difluorophenoxy)pyridin-2-yl)oxy)methyl)-3-fluorobenzonitrile





[0700] To a solution of 3-fluoro-4-(hydroxymethyl)benzonitrile (1.49 g, 9.87 mmol) in dry tetrahydrofuran (50 mL) sodium hydride (395 mg, 9.87 mmol 60% w/w in mineral oil) was slowly added at 0° C. The resulting mixture was stirred at 0° C. for 30 min, then 2-(4-bromo-2,5-difluorophenoxy)-6-fluoropyridine (2.00 g, 6.58 mmol) was slowly added to above mixture at 0° C. The resulting mixture was stirred for 3 hours at 70° C. After completion, the reaction was quenched with water (100 mL) at 0° C. and then extracted with dichloromethane (3x50 mL). The combined organic layers were washed with brine (20 mLx3), dried over anhydrous sodium sulfate, filtered, and concentrated to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=4/1) to obtain 4-((6-((4-bromo-2,5-difluorophenoxy)pyridin-2-yl)oxy)methyl)-3-fluorobenzonitrile (2.00 g, 69.9% yield). LC-MS m/z : 437 $[\text{M}+\text{H}]^+$.

3). Synthesis of tert-butyl 2-(4-(((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)-2,5-difluorophenyl)acetate




[0701] To a mixture of (2-(tert-butoxy)-2-oxoethyl)zinc bromide (1.20 g, 4.60 mmol), 1,2,3,4,5-pentylphenyl-1'-(di-tert-butylphosphonium)ferrocene (164 mg, 0.23 mmol) and tris(dibenzylideneacetone)dipalladium (211 mg, 0.23 mmol) in tetrahydrofuran (20 mL) 4-(((6-(4-bromo-2,5-difluorophenoxy)pyridin-2-yl)oxy)methyl)-3-fluorobenzonitrile (1.00 g, 2.30 mmol) was added at room temperature. The resulting mixture was stirred at 70° C. for 16 hours. After completion, the solvent was removed by concentration to give a residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=5/1) to obtain tert-butyl 2-(4-(((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)-2,5-difluorophenyl)acetate (1.00 g, 92.5% yield). LC-MS *m/z*: 471 [M+H]⁺.

4). Synthesis of 2-(4-(((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)-2,5-difluorophenyl)acetic acid



[0702] To a solution of tert-butyl 2-(4-(((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)-2,5-difluorophenyl)acetate (1.00 g, 2.13 mmol) in dichloromethane (10 mL) trifluoroacetic acid (5 mL) was added at room temperature. The resulting mixture was stirred at room temperature for 2 hours. After completion, the solvent was removed by concentration to give a residue in vacuo. The residue was purified by reverse-phase flash chromatography under the following conditions (Column: Spherical C18, 20-40 μ m, 120 g; Mobile Phase A: 10 mM hexachlorocyclohexane in water; Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 5% B-95% B in 30 minutes; Detector: 254 nm). The mobile phase containing the desired product was collected at 82% B and then concentrated in vacuo to give 2-(4-(((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)-2,5-difluorophenyl)acetic acid (800 mg, 90.7% yield). LC-MS *m/z*: 415 [M+H]⁺.


5). Synthesis of methyl (S)-4-(2-(4-(((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)-2,5-difluorophenyl)-acetamido)-3-fluoro-5-((oxetan-2-yl)methyl)amino) benzoate



[0703] The mixture of 2-(4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)-2,5-difluorophenyl)acetic acid (400 mg, 0.97 mmol), methyl (S)-4-amino-3-fluoro-5-((oxetan-2-ylmethyl)amino)benzoate (246 mg, 0.97 mmol), N,N-diisopropylethylamine (250 mg, 1.94 mmol) and 2,4,6-tripropyl-1,3,5,2,4,6-trioxatriphosphinane 2,4,6-trioxide (954 mg, 1.50 mmol, 50% in ethyl acetate) in THF (10 mL) was stirred at room temperature for 16 hours. After completion, the solvent was removed by concentration to give a residue in vacuo. The residue was purified by silica gel


column chromatography (petroleum ether/ethyl acetate=1/1) to obtain methyl (S)-4-(2-(4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)-2,5-difluorophenyl)acetamido)-3-fluoro-5-((oxetan-2-ylmethyl)amino)benzoate (400 mg, 63.4% yield). LC-MS m/z : 651 $[M+H]^+$.

6). Synthesis of methyl (S)-2-(4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)-2,5-difluorobenzyl)-4-fluoro-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate




[0704] The mixture of methyl (S)-4-(2-(4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)-2,5-difluorophenyl)acetamido)-3-fluoro-5-((oxetan-2-ylmethyl)amino)benzoate (200 mg, 0.31 mmol) in toluene (5 mL) and acetic acid (0.5 mL) was stirred at 110°C. for 3 hours. After completion, the solvent was removed by concentration to give residue in vacuo. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=1/1) to obtain methyl (S)-2-(4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)-2,5-difluorobenzyl)-4-fluoro-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (180 mg, 91.9% yield). LC-MS m/z : 633 $[M+H]^+$.

7). Synthesis of (S)-2-(4-((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)-2,5-difluorobenzyl)-4-fluoro-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylic acid



-continued



[0705] To a mixture of methyl (S)-2-(4-(((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)-2,5-difluorobenzyl)-4-fluoro-1-(oxetan-2-ylmethyl)-1H-benzo[d]imidazole-6-carboxylate (180 mg, 0.28 mmol) in water (5 mL) and THF (5 mL), lithium hydroxide (20 mg, 0.84 mmol) was added at room temperature. The resulting mixture was stirred at room temperature for 16 hours. After completion, the reaction mixture was adjusted to pH=5-6 with formic acid. The solvent was removed by concentration to give a residue in vacuo. The residue was purified by prep-HPLC (Waters 2767/2545/2489 system; Column: SunFire Prep C8 OBD 10 μ m 19 $\times$ 250 mm Column; Gradient elution with ACN/0.1% FA in H₂O solvent system); Detection Wavelength 254 nm/214 nm; Flow rate: 20 mL/min) to obtain (S)-2-(4-(((6-((4-cyano-2-fluorobenzyl)oxy)pyridin-2-yl)oxy)-2,5-difluo-

robenzyl)-4-fluoro-1-(oxetan-2-ylmethyl)-1H-benzo-[d]imidazole-6-carboxylic acid (148.49 mg, 85.8% yield). LC-MS m/z: 619 [M+H]⁺.




[0706] ¹H NMR (400 MHz, DMSO-d₆): δ 8.13 (d, J=1.2 Hz, 1H), 7.83 (t, J=8.0 Hz, 1H), 7.76 (dd, J=10.0 Hz, 1.2 Hz, TH), 7.54 (dd, J=8.0 Hz, 1.2 Hz, 1H), 7.47 (dd, J=11.2 Hz, 1.2 Hz, 1H), 7.42-7.38 (m, 1H), 7.26 (t, J=8.0 Hz, 2H), 6.72-6.67 (m, 2H), 5.19 (s, 2H), 5.09-5.04 (m, 1H), 4.81-4.76 (m, 1H), 4.67-4.63 (m, 1H), 4.54-4.42 (m, 3H), 4.38-4.33 (m, 1H), 2.74-2.67 (m, 1H), 2.45-2.33 (in, 1H).

Example 77. Synthesis of Other Compounds




[0707] Similar to the synthesis steps in Example 3-74, the following compounds can be obtained from the A-type intermediates and the B-type intermediates

Number	Structure	MS m/z: [M + H] ⁺
Compound 4		646




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 5		629
Compound 8		636
Compound 9		636




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 10		654
Compound 11		637
Compound 12		619




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 14		627
Compound 15		627
Compound 16		645




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 17		628
Compound 18		610
Compound 19		645




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 20		663
Compound 21		663
Compound 22		681




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 23		664
Compound 24		646
Compound 25		636




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 26		654
Compound 27		654
Compound 28		672




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 29		655
Compound 30		637
Compound 31		644




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 32		662
Compound 33		662
Compound 34		680




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 35		663
Compound 36		645
Compound 37		635




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 38		653
Compound 39		653
Compound 40		671




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 41		654
Compound 42		636
Compound 44		637




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 45		637
Compound 46		655
Compound 47		638




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 48		620
Compound 51		599
Compound 52		617




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 53		600
Compound 54		582
Compound 58		608




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 59		591
Compound 61		580
Compound 62		598




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 63		598
Compound 64		616
Compound 65		599




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 66		581
Compound 68		589
Compound 69		589




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 70		607
Compound 71		590
Compound 72		572




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 73		607
Compound 74		625
Compound 75		625




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 76		643
Compound 77		626
Compound 78		608




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 79		598
Compound 80		616
Compound 81		616




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 82		634
Compound 83		617
Compound 84		599

-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 85		606
Compound 86		624
Compound 87		624




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 88		642
Compound 89		625
Compound 90		607




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 91	 <chem>O=C(O)/C=C/c1ccc2nc3c(c1)nc(CN4CCCCC4Oc5ccc(cc5)COc6ccc(cc6)C#N)cc23[C@H]4OCCO4</chem>	597
Compound 92	 <chem>O=C(O)/C=C/c1cc(F)c2nc3c(c1)nc(CN4CCCCC4Oc5ccc(cc5)COc6ccc(cc6)C#N)cc23[C@H]4OCCO4</chem>	615
Compound 93	 <chem>O=C(O)/C=C/c1cc(F)c2nc3c(c1)nc(CN4CCCCC4Oc5ccc(F)cc5COc6ccc(cc6)C#N)cc23[C@H]4OCCO4</chem>	615




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 94		633
Compound 95		616
Compound 96		598





-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 97		594
Compound 98		612
Compound 99		612





-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 100		630
Compound 101		613
Compound 102		595





-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 103		630
Compound 104		648
Compound 105		648
Compound 106		666

-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 107		649
Compound 108		631
Compound 109		621
Compound 110		639





-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 111		639
Compound 112		657
Compound 113		640
Compound 114		622

-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 115	 <chem>O=C(O)/C=C/c1ccc2nc(CN3CCCCC3Oc4ccc(COc5cc(F)cc(Cl)c5)cc4)nc2c1F</chem>	629
Compound 116	 <chem>O=C(O)/C=C/c1cc(F)cc2nc(CN3CCCCC3Oc4ccc(COc5cc(F)cc(Cl)c5)cc4)nc2c1F</chem>	647
Compound 117	 <chem>O=C(O)/C=C/c1ccc2nc(CN3CCCCC3Oc4ccc(COc5cc(F)cc(Cl)c5)cc4)nc2c1F</chem>	647
Compound 118	 <chem>O=C(O)/C=C/c1cc(F)cc2nc(CN3CCCCC3Oc4ccc(COc5cc(F)cc(Cl)c5)cc4)nc2c1F</chem>	665




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 119		648
Compound 120		630
Compound 121		620
Compound 122		638




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 123	 <chem>O=C(O)/C=C/c1ccc2nc3c(ncn3C#CC4CC4)c5cc(F)ccc5N2</chem>	638
Compound 124	 <chem>O=C(O)/C=C/c1ccc2nc3c(ncn3C#CC4CC4)c5cc(F)ccc5N2</chem>	656
Compound 125	 <chem>O=C(O)/C=C/c1ccc2nc3c(ncn3C#CC4CC4)c5cc(F)ccc5N2</chem>	639
Compound 126	 <chem>O=C(O)/C=C/c1ccc2nc3c(ncn3C#CC4CC4)c5cc(F)ccc5N2</chem>	621




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 128		622
Compound 129		622
Compound 130		640




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 131		623
Compound 132		605
Compound 136		631




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 137		614
Compound 139		603
Compound 140		621




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 141		621
Compound 142		639
Compound 143		622




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 144		604
Compound 145		608
Compound 146		626




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 147		626
Compound 148		644
Compound 149		627





-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 150		609
Compound 151		618
Compound 152		636




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 153		636
Compound 154		654
Compound 155		637




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 156		619
Compound 157		609
Compound 158		627
Compound 159		627




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 160		645
Compound 161		628
Compound 162		610




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 163		617
Compound 164		635
Compound 165		635




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 166		653
Compound 167		636
Compound 168		618




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 169		585
Compound 170		603
Compound 171		603




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 172		621
Compound 173		604
Compound 174		586




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 175		595
Compound 176		613
Compound 177		613




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 179		614
Compound 180		596
Compound 181		586




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 182		604
Compound 183		604
Compound 184		622




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 185		605
Compound 186		587
Compound 187		594




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 188		612
Compound 189		612
Compound 190		630




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 191		613
Compound 192		595
Compound 193		633




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 194		651
Compound 195		651
Compound 196		669




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 197		652
Compound 198		634
Compound 199		624




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 200		642
Compound 201		642
Compound 202		660




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 203		643
Compound 204		625
Compound 205		632




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 206		650
Compound 207		650
Compound 208		668




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 209		651
Compound 210		633
Compound 211		623




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 212		641
Compound 213		641
Compound 214		659




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound 215		642
Compound 216		624
Compound S7		589




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound S8		607
Compound S9		590
Compound S10		616




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound S11		597
Compound S14		608
Compound S15		631




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound S16		630
Compound S17		639
Compound S21		620




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound S24		627
Compound S25		646
Compound S26		645

-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound S27		628
Compound S28		654
Compound S29		635




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound S32		598
Compound S33		617
Compound S34		616




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound S35		599
Compound S37		606
Compound S40		621




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound S42		639
Compound S43		622
Compound S44		648




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound S45		629
Compound S48		636
Compound S49		655




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound S50		654
Compound S51		637
Compound S52		663




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound S53		644
Compound S56		595
Compound S57		611




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound S58		614
Compound S59		573
Compound S60		573




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound S61		609
Compound S62		627
Compound S63		624




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound S64		642
Compound C44		641
Compound C45		624




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound C46		640
Compound C47		642
Compound C48		581




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound C49		572
Compound C50		598
Compound C51		607




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound C52		586
Compound C53		595
Compound C54		590




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound C55		599
Compound C56		624
Compound C57		633




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound C58		628
Compound C59		637
Compound C60		645




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound C61		636
Compound C62		572
Compound C63		581




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound C64		581
Compound C65		572
Compound C66		581




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound C67		572
Compound C68		619
Compound C69		610




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound C70		619
Compound C71		610
Compound C72		621




-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound C73		630
Compound C74		595
Compound C75		604

-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound C76		604
Compound C77		595
Compound C78		555

-continued

Number	Structure	MS m/z: [M + H] ⁺
Compound C79		564
Compound C80		613
Compound C81		565

Test Example 1. Detection of Agonist Activity of
the Test Compounds on GLP-1R

1). Reagents

Name	Manufacturer
cAMP Detection Kit	Cisbio
Bovine Serum Albumin(BSA)	Sigma

-continued

Name	Manufacturer
Hanks' Balanced Salt Solution 1X(HBSS)	Gibco
1M HEPES	Invitrogen
3-Isobutyl-1-methylxanthine(IBMx)	Invitrogen
PP-384 well plate	Greiner

-continued

Name	Manufacturer
OptiPlate-384 GLP-1(7-37)	PerkinElmer Hao Yuan

2). Equipment

Instrument	Model	Factory
En Vision Vi-cell counter	Envision2014 Vi-CELL™ XR Cell Viability Analyzer	PerkinElmer Beckman
Bravo Centrifuge	Bravo V11 Allegro™ 25R Centrifuge	Agilent Beckman
ECHO	ECHO 555	Labcyte

3). Cell Line

Target	Host cell	Source
GLP-1R	HEK293	WuXi

4). Methods and Procedures

[0708] 1) The test compound was dissolved in DMSO and diluted 4-fold with Bravo for a total of 10 concentration gradients (the starting concentration was 1 mM). The reference compound GLP-1(7-37) was dissolved in DMSO and diluted 4-fold with Bravo for a total of 10 concentration gradients (starting at 500 nM).

[0709] 2) The stable transfection GLP-1R/HEK293 cells were resuscitated in a sterile water bath at 37° C. and shaken gently until ice cubes were completely melted.

[0710] 3) The recovered cells were transferred to a 15 mL sterile centrifuge tube to remove DMSO. HBSS (10 mL) pre-warmed at 37° C. was added, mixed gently, and centrifuged at 1000 rpm for 5 minutes. The supernatant was discarded and another 10 mL of HBSS was added and mixed gently. The cells were counted to measure viability with Vi-cell counter.

[0711] 4) The cells were re-suspended with assay buffer, counted with a Vi-cell counter, and diluted to 1.0×10^5 /mL.

[0712] 5) 10 μ L of assay buffer containing cells was added to a 384-well plate with electronic multichannel pipette and the plate was shaken at 1000 rpm for 5 seconds.

[0713] 6) The diluted test compound solution was transfer to the 384 cell plate with ECHO by 100 nL of transfer volume. The plate was shaken to mix at 2000 rpm for 60 seconds.

[0714] 7) The cells in the 384-well plate were incubated in CO₂ cell incubator at 37° C. for 30 min.

[0715] 8) 10 μ L of HTRF cAMP detection solution was added to the 384-well plate, stick with sealing film, and incubated at room temperature for 1 hour.

[0716] 9) After 1 hour of incubation, the sealing film was removed. The plate was put into the EnVision microplate reader to read the values of OD665 nm and OD615 nm.

4). Data Analysis

[0717] a) Calculation formula:

$$\% \text{ Activity} = (\text{cAMP level of testing sample} - \text{average cAMP level of LC}) / (\text{average cAMP level of HC} - \text{average cAMP level of LC}) \times 100\%$$

[0718] Z factor: $1 - 3 \times (\text{STD_cAMP level of HC} + \text{STD_cAMP level of LC}) / (\text{average cAMP level of HC} - \text{average cAMP level of LC})$

[0719] Assay Window: HC/LC

[0720] HC: wells with adding the top concentration of reference compound

[0721] LC: wells with adding assay buffer

[0722] b) Using “log(agonist) vs. response—Variable slope” mode of GraphPad Prism 5.0 to fit the dose response curve of each sample and calculate the EC50 value.

TABLE 1

Data And Analysis of Some Representative Compounds		
Numer	EC ₅₀ (nM)	Max activity(%)
GLP-1(7-37)	0.02	98.1
PF-06882961	0.229	99.7
Compound 1	0.088	116.8
Compound 2	0.0529	98.6
Compound 3	0.317	128.3
Compound 6	0.133	114.8
Compound 7	0.145	91.3
Compound 13	0.114	99.4
Compound 43	0.110	109.7
Compound 49	0.193	129.3
Compound 55	0.149	111.5
Compound 56	0.210	124.9
Compound 60	0.455	112.0
Compound 67	1.086	106.4
Compound 133	0.264	126.6
Compound 134	0.133	117.3
Compound 135	0.244	109.6
Compound 138	0.669	125.0
Compound S1	1401	93.7
Compound S2	5.578	139.9
Compound S3	2170	70.1
Compound S4	0.29	106.9
Compound S5	15.44	125.2
Compound S6	2.6	114.9
Compound S20	0.962	129.1
Compound S22	730.8	92.7
Compound S23	1.472	98.0
Compound S36	150.2	114.2
Compound S41	21.68	94.8
Compound S66	1.253	106.7
Compound C1	1194	76.6
Compound C2	0.614	123.1
Compound C3	0.278	122.3
Compound C4	0.560	147.9
Compound C5	2.409	112.6
Compound C6	13.77	99.8

TABLE 1-continued

Data And Analysis of Some Representative Compounds		
Numer	EC ₅₀ (nM)	Max activity(%)
Compound C7	23.48	90.1
Compound C8	25.58	117.4
Compound C9	3.743	97.2
Compound C10	4.157	110.2
Compound C11	1.484	114.8
Compound C12	0.233	125.5
Compound C13	1.819	141.7
Compound C14	0.315	101.9
Compound C15	0.807	109.1
Compound C16	2.544	99.2
Compound C17	0.668	127.7
Compound C18	0.472	103.8
Compound C19	1.067	134.3
Compound C20	0.425	137.0
Compound C21	46.24	N/A
Compound C22	30.37	N/A
Compound C23	0.249	109.1
Compound C24	1.67	76.88
Compound C25	0.512	133.4
Compound C26	0.215	117.3
Compound C27	8.03	N/A
Compound C28	5.051	N/A
Compound C29	6.831	121.6
Compound C30	5.851	121.2
Compound C31	0.833	106.6
Compound C32	0.768	114.3
Compound C33	0.068	99.1
Compound C34	10.98	113.6
Compound C35	>100	N/A
Compound C36	3.919	104.0
Compound C37	0.223	121.0
Compound C38	>100	N/A
Compound C39	7.93	142.8
Compound C40	6.50	90.7
Compound C41	4.164	122.3
Compound C42	>100	N/A
Compound C43	5.306	88.8
Compound C82	7.483	90.7

TABLE 1-continued

Data And Analysis of Some Representative Compounds		
Numer	EC ₅₀ (nM)	Max activity(%)
Compound C83	0.705	111.1
Compound C84	0.246	136.4

[0723] Data in Table 1 indicated that most of the compounds of the present invention had potent agonistic effects on hGLP1R receptors, and were excellent hGLP1R agonists. Its maximal agonistic effect could reach or even exceed that of GLP1 polypeptide or the control molecule PF-06882961 currently under clinical research.

Test Example 2. Pharmacokinetic Study of the Test Compounds in Mice

[0724] The pharmacokinetic properties of some compounds of the present invention were evaluated in mice pharmacokinetic experiments. SPF male CD-1 mice from Beijing Vital River Laboratory Animal Technology Co., Ltd were used with 6 mice in each group. Compounds were dissolved in 10% solutol/90% saline. A single intravenous administration was given at a dose of 1 mg/kg in a volume of 1 mL/kg. A single intragastric administration was given at a dose of 5 mg/kg in a volume of 1 mL/kg. Animals were fasted overnight prior to the administration and resumed feeding 4 hours after administration, ad libitum to water. Blood was collected at 0.0830, 0.25, 0.5, 1, 2, 4, 6, 8 and 24 hours after dosing. About 0.4 mL of whole blood was collected from the saphenous vein of animals and placed in an EDTA-K2 anticoagulation tube. The samples were centrifuged at 4° C. and 4200 rpm for 5 min, and the plasma was transferred into a centrifuge tube and stored at -80° C. until analysis. For plasma sample analysis, the test compounds and internal standards (Labetalol & tolbutamide & Verapamil & dexamethasone & glyburide & Celecoxib) were extracted from mice plasma by protein precipitation method with acetonitrile, and the extracts were analyzed by LC/MS/MS. The measured plasma concentration-time data of individual animals were analyzed with the non-compartmental model of Phoenix WinNonlin 7.0 (Pharsight, USA) software, and the pharmacokinetic parameters of mice were obtained as follows: maximum (peak) plasma drug concentration C_{max}; Peak time T_{max}; half-life T_{1/2} and area under the plasma concentration-time curve extrapolated to infinity AUC_{0-inf}.

TABLE 2

Pharmacokinetic Parameters of Representative Compounds in Mice							
Compound	Dose(mg/kg)	C _{max} (ng/mL)	AUC _{0-inf} (ng · h/mL)	Cl(mL/min/kg)	V _{dss} (L/kg)	T _{1/2} (h)	F (%)
PF-06882961	1(iv)	—	370	45.1	1	0.3	—
Compound 55	—	—	637	26.2	1.3	1.5	—
Compound 49	—	—	1350	12	1	1	—
Compound 56	—	—	771	21.7	1.2	2.3	—
Compound C23	—	—	1141	15	1	1	—
PF-06882961	5(po)	299	366	—	—	2.0	19.8
Compound 55	—	483	1042	—	—	1.7	32.7
Compound 49	—	515	1760	—	—	3.2	20.7
Compound 56	—	1640	2097	—	—	1.0	54.4
Compound C23	—	2193	2575	—	—	1.4	45.1

[0725] The results in Table 2 indicated that compared with the reference molecule PF-06882961, some representative compounds of the present invention such as compounds 56, 49, 55 and C23 had lower clearance rates higher exposure and larger oral bioavailability, and therefore more suitable as candidates for drug development.

Test Example 3. Pharmacokinetic Study of the Test Compounds in Monkeys

[0726] The pharmacokinetic properties of compounds of the present invention were evaluated in monkey pharmacokinetic experiments with 3 male Non-naive cynomolgus monkeys per group. Compounds were dissolved in 10% solutol/90% saline. For single intravenous administration, the dose was 1 mg/kg, and the administration volume was 1 mL/kg. For single intragastric administration, the dose was 5 mg/kg, and the administration volume was 1 mL/kg. Animals were fasted overnight with free water before the experiment and resumed feeding 4 hours after administration. Blood was collected at 0.0830, 0.25, 0.5, 1, 2, 4, 6, 8 and 24 hours after dosing. About 0.4 mL of whole blood was collected from the animal's forearm vein and placed in an EDTA-K2 anticoagulant tube. The samples were centrifuged at 4° C. and 4200 rpm for 5 min, and the plasma was transferred to a centrifuge tube and stored at -80° C. until analysis. For plasma sample analysis, the test compounds and internal standards (Labetalol & tolbutamide & Verapamil & dexamethasone & glyburide & Celecoxib) were extracted from monkey plasma by protein precipitation method with acetonitrile, and the extracts were analyzed by LC/MS/MS. Measured plasma concentration-time data in individual animals were analyzed using the non-compartmental model of Phoenix WinNonlin 7.0 (Pharsight, USA) software to obtain the following pharmacokinetic parameters: maximum (peak) plasma drug concentration C_{max}; time to peak T_{max}; Half-life T_{1/2} and area under the plasma concentration-time curve extrapolated to infinity AUC_{0-inf}.

TABLE 3

Pharmacokinetic Parameters of Some Representative Compounds in Monkeys						
Compound	Dose(mg/kg)	C _{max} (ng/mL)	AUC _{0-inf} (ng · h/mL)	Cl(mL/min/kg)	V _{dss} (L/kg)	T _{1/2} (h) F (%)
PF-06882961	1(iv)	—	1167	14.4	0.3	1.1 —
Compound 55	—	—	1425	12.0	0.4	2.2 —
Compound 49	—	—	2381	7.26	0.29	1.34 —
PF-06882961	5(po)	58.8	344	—	—	2.8 5.9
Compound 55	—	214	614	—	—	5.4 8.6
Compound 49	—	235	1244	—	—	4.01 10.4

[0727] The results in Table 3 showed that, compared with the reference molecule PF-06882961, some representative compounds of the present invention, such as compound 49 and compound 55, had lower clearance rates, higher exposure, and greater oral bioavailability and half-life, thus more suitable as candidates for drug development.

Test Example 4. Intraperitoneal Glucose Tolerance (IPGTT) Test in h-GLP1-R C57BL6J Mice

1) Formulation

[0728] The compounds to be tested were dissolved in 10% Solutol HS15+90% Saline, and vortexed by ultrasonic to dissolve.

2) Animals

[0729] Male SPF h-GLP1-R C57BL6J mice, 6-9 weeks old, weighing 20 g-25 g.

3) Animal Acclimation and Grouping

[0730] Upon arrival, animals were acclimated for 7 days before experiments. The body weight and random blood sugar of the animals were measured one day before administration. Animals were divided into 4 groups based on body weight and random blood glucose.

[0731] The animal grouping information was as follows:

TABLE 4

Animal Grouping Information				
Group	Dose level (mg/kg)	Dose (ml/kg)	Route of administration	Frequency
Vehicle	—	10	PO	QD
PF-06882961	0.3	10	PO	QD
Compound 55	0.3	10	PO	QD

4) IPGTT Test

[0732] Mice were overnight fasted before IPGTT. Baseline fasting blood glucose and insulin level were measured before vehicle or compound dosing. 30 min post of compound dosing, glucose were injected intraperitoneally (2 g/kg, 10 ml/kg). Blood glucose levels were measured at -30, 0, 15, 30, 60, 120 min. Plasma 15 ul were collected at 0 and 15 min for insulin analysis by Rat/Mouse Insulin Elisa Kit.

5) Data Analysis

[0733] AUC_{0-120min} was the area under the blood glucose-time curve. Insulin change was the difference between 15

min and 0 min plasma insulin concentration. Statistical analysis of data was performed with Graphpad Prism 6, and the statistical method was one-way ANOVA Dunnett test. Compared with the vehicle group, p<0.05 was considered a significant difference. * means p<0.05, ** means p<0.01, *** means p<0.001.

6) Experimental Results

[0734] As shown in FIGS. 1 and 2, a single oral dose of compound 55 significantly reduced the blood glucose enduring IPGTT. AUC_{0-120min} results showed that 0.3 mpk had a significant difference compared with the vehicle group. Insulin change results showed that 0.3 mpk compound 55 could significantly increase insulin secretion. In conclusion,

compound 55 showed similar or slightly better hypoglycemic and insulin-releasing effects compared with the control molecule PF-06882961.

Test Example 5. Intravenous Glucose Tolerance (IVGTT) Experiment in Cynomolgus Monkey

1) Formulation

[0735] The compounds were dissolved in 10% Solutol HS15+90% Saline by vortex ultrasound. Vehicle was 10% Solutol HS15+90% Saline.

2) Animals

[0736] Male cynomolgus monkeys, 10-15 years old, weighing 6-13 kg. Healthy animals were selected for efficacy studies according to alanine aminotransferase (ALT), aspartate aminotransferase (AST), triglyceride (TG), total cholesterol (TC), high density lipoprotein (HDL), low density lipoprotein (LDL), glycosylated hemoglobin (HbA1c), body weight (BW).

3) Animal Acclimation and Grouping

[0737] After 1 week of acclimation, the IVGTT baseline before administration was measured, and the animals were grouped according to the insulin and blood glucose results of the IVGTT. The detailed information of treatment was as follows:

TABLE 5

Animal Grouping Information				
Group	Dose level (mg/kg)	Dose (ml/kg)	Route of administration	Frequency
Vehicle	—	5	PO	QD
PF-06882961	100	5	PO	QD
Compound 49	60	5	PO	QD
Compound 49	100	5	PO	QD
Compound 55	100	5	PO	QD

4) IVGTT Test

[0738] IVGTT was started after 2 hours post of compound dosing. Blood were collected at -15 min, 0 min (before sugar administration), and 2, 5, 7, 10, 20, and 30 min after sugar administration. And the blood glucose level were measured.

5) Data Analysis

[0739] Data was analyzed by Graphpad Prism 6 and the statistical method was one-way ANOVA Dunnett's test. Compared with the vehicle group, $p < 0.05$ was considered significant difference. * means $p < 0.05$, ** means $p < 0.01$, *** means $p < 0.001$.


6) Experimental Results

[0740] As shown in FIG. 3, in the IVGTT experiment, a single oral dose of 100 mg/kg compound 49, 55 or PF-06882961 could significantly lower the blood glucose level. Compared with PF-06882961, compound 49 was slightly more effective, and the effect of compound 55 was equivalent. Further more, compound 49 still had a significant hypoglycemic effect in the 60 mg/kg dose group.

[0741] The embodiments of the present invention have been described above. However, the present invention is not limited to the above-described embodiments. Any modification, equivalent replacement, improvement, etc. made within the spirit and principle of the present invention shall be included within the protection scope of the present invention.


1-12. (canceled)

13. A compound, which is:




or a pharmaceutically acceptable salt, solvate, enantiomer, or isotopic variation thereof.


14. The compound of claim 13, which is:



15. The compound of claim 13, which is a pharmaceutical acceptable salt of:




16. A compound, which is:




or a pharmaceutically acceptable salt, solvate, enantiomer, or isotopic variation thereof.

17. The compound of claim 16, which is:



18. The compound of claim 16, which is a pharmaceutical acceptable salt of:



19. A pharmaceutical composition comprising a therapeutically effective amount of the compound of claim 13.

20. The pharmaceutical composition of claim 19, further comprising one or more pharmaceutically acceptable excipients.

21. The pharmaceutical composition of claim 19, further comprising one or more additional therapeutic agents.

22. A pharmaceutical composition comprising a therapeutically effective amount of the compound of claim 16.

23. The pharmaceutical composition of claim 22, further comprising one or more pharmaceutically acceptable excipients.

24. The pharmaceutical composition of claim 22, further comprising one or more additional therapeutic agents.

25. A method of inducing insulin secretion in a subject in need thereof, comprising administering a therapeutic effective amount of the compound of claim 13 to the subject.

26. A method of reducing blood sugar level in a subject in need thereof, comprising administering a therapeutic effective amount of the compound of claim 13 to the subject.

27. A method of treating or preventing obesity in a subject in need thereof, comprising administering a preventive or therapeutically effective amount of the compound of claim 13 to the subject.

28. A method of treating or preventing diabetes in a subject in need thereof, comprising administering a preventive or therapeutically effective amount of the compound of claim 13 to the subject.

29. The method of claim 28, wherein the diabetes is Type 2 diabetes.

30. A method of inducing insulin secretion in a subject in need thereof, comprising administering a therapeutic effective amount of the compound of claim 16 to the subject.

31. A method of reducing blood sugar level in a subject in need thereof, comprising administering a therapeutic effective amount of the compound of claim 16 to the subject.

32. A method of treating or preventing obesity in a subject in need thereof, comprising administering a preventive or therapeutically effective amount of the compound of claim 16 to the subject.

33. A method of treating or preventing diabetes in a subject in need thereof, comprising administering a preventive or therapeutically effective amount of the compound of claim 16 to the subject.

34. The method of claim 33, wherein the diabetes is Type 2 diabetes.

* * * * *