



US 20250283106A1

(19) **United States**

(12) **Patent Application Publication**
WANG

(10) **Pub. No.: US 2025/0283106 A1**

(43) **Pub. Date: Sep. 11, 2025**

(54) **METHODS AND COMPOSITIONS FOR
PLANT PATHOGEN RESISTANCE IN
PLANTS**

(71) Applicant: **UNIVERSITY OF FLORIDA
RESEARCH FOUNDATION, INC.**,
Gainesville, FL (US)

(72) Inventor: **Nian WANG**, Auburndale, FL (US)

(21) Appl. No.: **19/220,933**

(22) Filed: **May 28, 2025**

Related U.S. Application Data

(62) Division of application No. 18/089,985, filed on Dec. 28, 2022, now Pat. No. 12,344,854, which is a division of application No. 16/761,409, filed on May 4, 2020, filed as application No. PCT/US2018/059269 on Nov. 5, 2018, now Pat. No. 11,634,725.

(60) Provisional application No. 62/581,491, filed on Nov. 3, 2017, provisional application No. 62/627,496, filed on Feb. 7, 2018.

Publication Classification

(51) **Int. Cl.**
C12N 15/82 (2006.01)
(52) **U.S. Cl.**
CPC **C12N 15/8281** (2013.01)

(57) **ABSTRACT**

The disclosure relates to a plant that is tolerant or resistant to species of *Ca. Liberibacter*. Specifically exemplified are *Citrus* and solanaceous plants. Provided by the disclosure is a modified *Citrus* or solanaceous plant that is resistant or tolerant to Sec-dependent effectors secreted by bacteria. Also provided by the disclosure are methods of modifying a plant genome plant to provide tolerance or resistance to species of *Ca. Liberibacter*. Still further provided by the disclosure are methods conferring a population of plants with tolerance or resistance to species of *Ca. Liberibacter* and screening that population for the plants that are tolerant or resistant to species of *Ca. Liberibacter*.

Specification includes a Sequence Listing.

FIG. 1A




FIG. 1C




FIG. 1B

BD	AD	Mating control (DDO)	Selection (DDO/X)	Selection (DDO/X/A)	Selection (QDO)	Selection (QDO/X)	Selection (QDO/X/A)
4025	RLK2						
EV	RLK2						
4025	PR10						
EV	PR10						
4025	PP2B2						
EV	PP2B2						
Positive							
Negative							

FIGURE 2




FIG. 3A




FIG. 3B


**FIGURE 4**

FIG. 5A

BD	AD	Mating control (DDO)	Selection (QDO/X/A)
SDE15	CtACD2		
SDE15	EV		
EV	CtACD2		

FIG 5B





FIG. 5C




FIG. 5D




FIG. 5E

Figure 6A

MTISKNQAILFFITGMILSSCGDTLSDSKQHKNKINNTKNHLDLLFPIDDSHNQKPTEKKPN

TSSIKIKNNIIEPQPGPSRWEGGWNGERYVREWER




Figure 6B




Figure 6E




Figure 6C

Figure 6D




Figure 6F




Figure 7A




Figure 7B




Figure 8

Las Ct Value (per 100ng DNA)					
	0 mpi	1 mpi	2 mpi	3 mpi	4 mpi
SDE15	37.17±0.46	37.43±0.53	35.38±2.09	32.54±1.38**	32.11±0.68**
EV	36.51±0.50	37.06±0.73	36.60±0.66	34.88±1.05	34.17±0.90




Figure 9A




Figure 9B

Figure 10A

BD	AD	Mating control {BDO}	Selection {BDO/X/A}
SDE15	CsACD2		
SDE15	EV		
EV	CsACD2		

Figure 10B





Figure 10C




Figure 10D




Figure 10E




Figure 11

Figure 12A





Figure 12B




Figure 12C

METHODS AND COMPOSITIONS FOR PLANT PATHOGEN RESISTANCE IN PLANTS

REFERENCE TO ELECTRONIC SEQUENCE LISTING

[0001] The application contains a Sequence Listing which has been submitted electronically in .XML format and is hereby incorporated by reference in its entirety. Said .XML copy, created on May 2, 2023, is named “10457373US4_SequenceListing.xml” and is 134,000 bytes in size. The sequence listing contained in this .XML file is part of the specification and is hereby incorporated by reference herein in its entirety.

FIELD OF THE INVENTION

[0002] The present disclosure relates to the field of biotechnology. More specifically, the disclosure relates to compositions and methods for producing plants that are resistant to *Ca. Liberibacter* infection in plants, such as Huanglongbing (HLB), also known as *Citrus* greening disease.

BACKGROUND

[0003] Currently available commercial *Citrus* plants lack tolerance or resistance to Huanglongbing (HLB), also known as *Citrus* greening disease. HLB is caused by species of the phloem-limited, gram-negative bacteria of genus *Ca. Liberibacter*. In the U.S., the predominant pathogenic species is *Ca. Liberibacter asiaticus* (Las); whereas, *Ca. Liberibacter africanus* (Laf) and *Ca. Liberibacter americanus* (Lam) are the predominant pathogenic species in South Africa and Brazil, respectively. *Ca. Liberibacter* is a vector-transmitted pathogen. The vector organisms are the Asian *Citrus* psyllid (ACP), *Diaphorina citri*, and African *Citrus* psyllid, *Trioza erythrae*. HLB was first detected in the United States in August 2005 and has rapidly moved into several *Citrus* producing areas. All commercial *Citrus* plants are susceptible to HLB, and infected *Citrus* plants will irrevocably decline. Thus, HLB has resulted in a severe decline in fruit production in Florida, where HLB has become endemic. Currently, HLB management consists of preventing trees from becoming infected, which includes protecting young flush from the *Ca. Liberibacter* vector organisms and destroying infected plant material. However, due to the lack of rapid curative methods that control HLB, new methods to prevent infection are required to stop the spread of infection and further decline of the U.S. *Citrus* industry.

SUMMARY

[0004] Certain embodiments of the disclosure relate to increasing plant resistance to infection by a bacterial species from the genus *Ca. Liberibacter*. One aspect of the present disclosure relates to modified *Citrus* plants comprising genomes in which endogenous genes or regulatory elements thereof may be modified, wherein the modification confers resistance to HLB to the modified *Citrus* plant relative to a plant of the same variety lacking the modification. The *Citrus* plant in certain embodiments may be a grapefruit tree, orange tree, sweet orange tree, or mandarin tree. Further provided are plant parts and seeds of the modified *Citrus* plant. Another aspect of the disclosure is a method of producing a commodity plant product, from the modified

Citrus plant. In certain embodiments this method comprises collecting the commodity plant product from the modified *Citrus* plant. Further provided are commodity plant products produced by this method. In addition to modified *Citrus* plants, other plants known to be infected by *Ca. Liberibacter* such as solanaceous crops may be genomically modified to disrupted to confer resistance to such infection.

[0005] In certain embodiments, modified endogenous genes may encode any polypeptide that interacts with any Sec-dependent effector (SDE) secreted by a bacterial species from the genus *Ca. Liberibacter*. In certain embodiments, modified regulatory elements regulate any endogenous gene that may encode any polypeptide that interacts with any SDE secreted by a bacterial species from the genus *Ca. Liberibacter*. An endogenous gene in particular embodiments may encode PP2-B2/12, Lectin, Cysteine protease, Cysteine protease 15A-like, Papain-like cysteine proteases, Myb family transcription factor, YLS9-like, Cell death suppressor protein Lls1, Acd1-Like, Acd1, accelerated cell death 2 (ACD2) protein, red chlorophyll catabolite reductase-like, NDR1/HIN1-like protein 13 (Cs8g01640), PHL5-like (Cs7g01290), and PHL5 (orange 1.1t02259). An SDE in particular embodiments may be Las4025, Las470, Las4065, Las5150, or Las4250.

[0006] Still a further aspect of the disclosure is a method of generating a modified plant comprising resistance to *Ca. Liberibacter* infection. In one embodiment, the method comprises the following steps: (a) introducing a genetic modification into the genome of a plant cell, wherein the modification is to an endogenous gene or regulatory element thereof, wherein a polypeptide encoded by the endogenous gene interacts with an SDE secreted by a bacterial species from the genus *Ca. Liberibacter*; (b) regenerating the modified plant from the plant cell or a progenitor cell thereof, wherein the plant comprises the modification (i.e. comprises cells that possess the modification); and (c) identifying a plant comprising the modification and the resistance to *Ca. Liberibacter* infection. In a specific example, the plant is a *Citrus* plant or a solanaceous crop.

[0007] In certain embodiments, modified endogenous genes may encode any polypeptide that interacts with any SDE secreted by a bacterial species from the genus *Ca. Liberibacter*. In certain embodiments, modified regulatory elements regulate any endogenous gene that may encode any polypeptide that interacts with any SDE secreted by a bacterial species from the genus *Ca. Liberibacter*. An endogenous gene in particular embodiments may encode PP2-B2/12, Lectin, Cysteine protease, Cysteine protease 15A-like, Papain-like cysteine proteases, Myb family transcription factor, YLS9-like, Cell death suppressor protein Lls1, Acd1-Like, Acd1, accelerated cell death 2 (ACD2) protein, red chlorophyll catabolite reductase-like, NDR1/HIN1-like protein 13 (Cs8g01640), PHL5-like (Cs7g01290), and PHL5 (orange 1.1102259). An SDE in particular embodiments may be Las4025, Las470, Las4065, Las5150, or Las4250. In certain embodiments, step (a) comprises a genome-editing technique. In certain embodiments, the genome-editing technique comprises a nuclease, wherein the nuclease introduces a single-strand DNA break or a double-strand DNA break. In certain embodiments, the genome-editing technique comprises a TALEN, a ZFN, meganuclease, or a CRISPR/Cas system. The disclosure still further provides a *Citrus* plant produced by this and the foregoing methods.

[0008] Still yet another aspect of the disclosure is a method for conferring a plurality of plants with a resistance to *Ca. Liberibacter* infection. In one embodiment, the method comprises the following steps: (a) introducing a genetic modification into a plurality of plants, wherein the modification is to an endogenous gene or regulatory element thereof, wherein a polypeptide encoded by the endogenous gene interacts with an SDE secreted by a bacteria species from the genus *Ca. Liberibacter*; and (b) screening the plurality of plants for the modification and a resistance to *Ca. Liberibacter* infection. The plurality of plants may include *Citrus* plants or solanaceous plants. In certain embodiments, modified endogenous genes may encode any polypeptide that interacts with any SDE secreted by a bacterial species from the genus *Ca. Liberibacter*. In certain embodiments, modified regulatory elements regulate any endogenous gene that may encode any polypeptide that interacts with any SDE secreted by a bacterial species from the genus *Ca. Liberibacter*. An endogenous gene in particular embodiments may encode PP2-B2/12, Lectin, Cysteine protease, Cysteine protease 15A-like, Papain-like cysteine proteases, Myb family transcription factor, YLS9-like, Cell death suppressor protein Lls1, Acd1-Like, Acd1, accelerated cell death 2 (ACD2) protein, red chlorophyll catabolite reductase-like, NDR1/HIN1-like protein 13 (Cs8g01640), PHL5-like (Cs7g01290), and PHL5 (orange1.1102259). An SDE in particular embodiments may be Las4025, Las470, Las4065, Las5150, or Las4250. In certain embodiments, step (a) comprises a genome-editing technique. In certain embodiments, the genome-editing technique comprises a nuclease, wherein the nuclease introduces a single-strand DNA break or a double-strand DNA break. In certain embodiments, the genome-editing technique comprises a TALEN, a ZFN, meganuclease, or a CRISPR/Cas system. The disclosure still further provides a *Citrus* plant produced by this and the foregoing methods.

BRIEF DESCRIPTION OF THE DRAWINGS

[0009] FIGS. 1A-C show transgenic expression of CLIBASIA_04025 (Las4025) in grapefruit (*Citrus paradisi*) caused HLB-like symptoms. FIG. 1A shows binary plasmid used for transgenic expression of CLIBASIA_04025 (Las4025) in *Citrus*. CsVMV: Cassava vein mosaic virus promoter; CaMV 35S: cauliflower mosaic virus 35S promoter; 35T: cauliflower mosaic virus terminator; NosT: nopaline synthase gene terminator; LB: left border of the T-DNA region; RB: right border of the T-DNA region; GFP: green fluorescent protein; NptII: coding sequence of neomycin phosphotransferase II; and Mature protein: nucleotide sequence encoding the mature CLIBASIA_04025 (Las4025) protein without its signal peptide. FIG. 1B shows transgenic grapefruit plants transformed with empty vector (control) or the vector that expressed CLIBASIA_04025 (Las4025) (T1, T2, and T3). FIG. 1C shows Western blots that confirmed transgenic grapefruit lines T1, T2, and T3 expressed CLIBASIA_04025. For these Western blots, a CLIBASIA_04025 (Las4025)-specific antibody was used; the polypeptide with SEQ ID NO:1 was the antigen used to produce this antibody.

[0010] FIG. 2 shows interactions between CLIBASIA_04025 (Las4025) and targets proteins demonstrated by yeast two-hybrid assay. Yeast cells were co-transformed with GAL4 DNA-binding domain (BD) expression vector pGBKT7 and GAL4 activation domain (AD) expression

vector pGADT7. The pGBKT7 vector was either empty (EV) or contained CLIBASIA_04025 (Las4025) without its signal peptide cloned in-frame with the BD; and the pGADT7 vector contained either RLK2, Pathogenesis-related protein 10 (Cs9g03630) (PR10), or PP2-B2 (orange1.1t04174) (PP2B2) cloned in-frame with the AD. The negative control was performed with empty-AD and -BD vectors, and the positive control was performed with p53-BD and T-antigen-AD vectors. Cultures of the co-transformed yeast clones were adjusted to an OD₆₀₀ of 0.2 and then spotted at three dilutions ($1/10$, $1/100$, and $1/1000$) on DDO, DDO/X, DDO/X/A, QDO, QDO/X, and QDO/X/A media. The images were captured after 4 days of growth. SD: synthetic defined yeast minimal medium; DDO: double dropout media, SD-Leu/-Trp; QDO: quadruple dropout media, SD/-Ade/-His/-Leu/-Trp; X: X- α -Gal; and A: Aureobasidin A.

[0011] FIGS. 3A-B show CLIBASIA_04025 interacted with cysteine protease (Cs4g07410). FIG. 3A shows the interaction between CLIBASIA_04025 (4025) and cysteine protease (CP, Cs4g07410) detected with a Glutathione-S-transferase (GST) pull-down assay. GST/CP was expressed in *E. coli*, immobilized on glutathione sepharose beads, and incubated with *E. coli* lysates containing Maltose-Binding Protein (MBP)-4025. The bound proteins were immunoblotted using anti-GST and anti-MBP antibodies. FIG. 3B shows the interaction between 4025 and CP detected using Bimolecular Fluorescence Complementation (BiFC), in which EYFP fluorescence represents a protein-protein interaction. Expression vectors with 4025 without its signal peptide or CP cloned in-frame with either the N- or C-terminal fragments of EYFP (EYFPN and EYFPC) were synthesized. Citrus protoplasts were co-transformed with combinations of EYFPC or EYFPN with 4025 or CP expression vectors. The co-expression of 4025-EYFPN and EYFPC/CP and the co-expression of EYFPC-4025 and CP-EYFPN produced strong signal in the *Citrus* protoplasts; whereas, no detectable fluorescence signal was produced by the negative control combinations.

[0012] FIG. 4 shows overexpression of Myb family transcription factor (MYTL, XP_015380888.1) in Duncan grapefruit induced HLB-like symptoms. FIG. 4A shows expression of MYTL in grapefruit driven by the 35S promoter reduced plant growth. FIG. 4B shows the normal amount of starch particles, red stain, that accumulated in wildtype grapefruit. FIG. 4C shows overexpression of MYTL in grapefruit induced greater starch accumulation than displayed by wildtype grapefruits.

[0013] FIGS. 5A-E show SDE15 (identified as CLIBASIA_04025) interacts with CtACD2 protein, which negatively regulate hypersensitive reaction. FIG. 5A. Yeast-two hybrid (Y2H) assay. Full-length SDE15 fused to the GAL4 DNA binding domain (BD) was expressed in combination with full-length CtACD2 fused to the GAL4 activation domain (AD) in the yeast strain Y2HGold. Strains were grown on double dropout medium (DDO) with -Trp and -Leu and screened on quadruple dropout medium (QDO) with -Trp, -Leu, -Ade and -His supplemented with X- α -Gal and Aureobasidin A (QDO/X/A). The empty BD and AD vectors were used as the negative controls. FIG. 5B. Bimolecular fluorescence complementation (BiFC) assay. Coding sequence of SDE15 without signal peptide fused to N-terminal or C-terminal fragments of EYFP vectors pSAT6-nEYFP-C1 and pSAT6-cEYFP-C1-B were co-transformed

with full-length CtCtACD2 and CtCtACD1 protein which fused to C-terminal and N-terminal fragments of EYFP vectors pSAT6-nEYFP-C1 and pSAT6-cEYFP-C1-B into *Citrus* protoplasts. The EYFP fluorescence of protoplasts were imaged 1 day after incubation using a Leica fluorescence microscope. For genuine interaction between SDE15 and CtACD2, co-expression of SDE15-EYFP^N and EYFP^C-CtACD2 or EYFP^C-SDE15 and CtACD2-EYFP^N should give strong fluorescent signals in protoplast. Co-transformations of SDE15-EYFP^N and EYFP^C, EYFP^N and EYFP^C-SDE15, SDE15-EYFP^N and EYFP^C-SDE15, CtACD2-EYFP^N and EYFP^C, EYFP^N and EYFP^C-CtACD2, CtACD2-EYFP^N and EYFP^C-CtACD2, EYFP^N+EYFP^C were used as negative controls which didn't produce any detectable fluorescence signal. FIG. 5C. Glutathione-S-transferase (GST) pull-down assay. GST-SDE15 and GST empty vectors were expressed in *E. coli*, immobilized on glutathione sepharose beads, and incubated with *E. coli* lysate containing MBP-CtACD2. Total cell extract (Input) and eluted protein (Elute) were immunoblotted using the anti-MBP and anti-GST antibody. FIG. 5D. Hypersensitive response (HR) assay. *Agrobacterium tumefaciens* strain GV2260 harboring binary vectors containing SDE15 and CtACD2 were infiltrated into leaves of *N. benthamiana* at the concentration of 10⁸ CFU ml⁻¹. Two days later, another *Agrobacterium tumefaciens* strain GV2260 harboring the binary vector containing AvrBsT protein that can trigger HR was infiltrated on the same area of the leaves treated before. HR induction was observed and photographed 2-3 days post-inoculation. All experiments were repeated three times with the similar results, and only one leaf was presented. FIG. 5E. Electrolyte leakage associated with HR induced by AvrBsT 2 days post infiltration. Leaf discs of

[0014] AvrBsT infiltrated plants were floated on deionized water with shaking. The conductivity of the solution was measured after 4 h shaking. Error bars indicate standard error of mean (n=3). Alphabets represent significant differences in different types of samples.

[0015] FIGS. 6A-F Characterization of SDE15. FIG. 6A. Sequence analysis of SDE15. Amino acid sequence of SDE15 (96 aa) with N-terminal signal peptide (highlighted in yellow) predicted using SignalP V4.1. The cleavage site localizes between the 22nd and 23rd aa (SCG-DT). FIG. 6B. Yellowing and mottling of the leaf were observed in transgenic *Citrus* cultivar 'Duncan' plants constitutively expressing SDE15 compared with the leaf of empty-vector (EV) transgenic *Citrus*. FIG. 6C. SDE15 detection in phloem sap. Phloem sap was isolated from the bark of both healthy and HLB infected *Citrus*. T1: total bark proteins; T2: total bark proteins after phloem sap isolation; P: phloem sap. FIG. 6D. Subcellular localization of SDE15. SDE15-EYFP was co-expressed with the plasma membrane localization-marker PM-CFP or the nucleus localization-marker CFP-nucleus in leaves of *N. benthamiana*. *Agrobacterium* strains carrying the corresponding expression plasmids were infiltrated at the optical density (OD₆₀₀) of 0.2. Subcellular localization of SDE15-EYFP was inspected and photographed 1 day post infiltration. Scale bars: 10 μm. FIG. 6E, FIG. 6F. qRT-PCR analysis of SDE15 expression in different Las hosts (FIG. 6E) and in different stages of Las infection (FIG. 6F). Relative transcript abundances were determined using gyrase subunit A of Las (CLIBASIA_00325) and *Citrus* house keeping gene encoding glyceraldehyde-3-phosphate dehydrogenase-C (GAPDH-C) as endogenous controls.

Bars represent the mean of eight replicates. Asterisks represent significant differences in the transcript abundance between *Citrus* and psyllids (**p-Value<0.01). Alphabets represent significant differences in samples of different Las infection stages. Error bars indicate standard error of mean (n=6). All experiments were repeated three times with the similar results.

[0016] FIGS. 7A-B. Hypersensitive reaction (HR) was repressed in SDE15-transgenic *Citrus*. FIG. 7A A strong HR, a form of programmed cell death (PCD), was observed in wild type Duncan grapefruit at 3 days after inoculation with *Xanthomonas citri* subsp. *citri* strain A^W (XccA^W). Only slight cell death was observed on the XccA^W-infiltrated leaves of SDE15-transgenic *Citrus* at 5 days post inoculation. XccA^W cells were infiltrated into *Citrus* leaves at a concentration of 10⁸ CFU/ml. FIG. 7B. qRT-PCR analysis of PR genes. Expression of PR1, PR3 and PR5 was repressed in SDE15-transgenic *Citrus* compared to that in wild type Duncan after HR induction by XccA^W. The house keeping gene encoding glyceraldehyde-3-phosphate dehydrogenase-C (GAPDH-C) was used as an endogenous control. Bars represent the mean of four replicates. Error bars indicate standard error of mean. All experiments were repeated three times with the similar results.

[0017] FIG. 8. Transgenic SDE15 *Citrus* plants are more susceptible to HLB. Leaf images taken 3 months post HLB infection via budding grafting. The Las titer in SDE15-transgenic *Citrus* and EV-transgenic control *Citrus* were determined by TaqMan qPCR 0, 1, 2, and 3 months post HLB infection. Each Ct value was represented by Means±standard error (n_{SDE15}=7, n_{EV}=5). Asterisks represent significant differences in the Las titer between SDE15-transgenic *Citrus* and non-transgenic control (**p-Value<0.01).

[0018] FIGS. 9A-B SDE15-transgenic *Citrus* became more susceptible to *Citrus* canker caused by a virulent strain (XacA 306) of *Xanthomonas citri* pv. *citri*. FIG. 9A. Water-soak symptom (grey color) was observed on SDE15-transgenic *Citrus* at 5 days post XacA 306 inoculation. FIG. 9B. Bacterial population increase of XacA 306 in SDE15-transgenic *Citrus* was faster than in non-transgenic Duncan grapefruit. Bacterial cells were infiltrated into *Citrus* leaves at a concentration of 10⁶ CFU/ml. Error bars indicate standard error of mean (n=4). Asterisks represent significant differences in the bacteria population between SDE15-transgenic *Citrus* and non-transgenic control (**p-Value<0.01, ***p-Value<0.001).

[0019] FIGS. 10A-E. SDE15 interacts with CsACD2 protein. FIG. 10A. Yeast-two hybrid (Y2H) assay using SDE15 as the bait and full-length CsACD2 protein as prey. Full-length SDE15 fused to the GAL4 DNA binding domain (BD) was expressed in combination with full-length CsACD2 fused to the GAL4 activation domain (AD) in the yeast strain Y2HGold. Strains were grown on double dropout medium (DDO) with -Trp and -Leu and screened on quadruple dropout medium (QDO) with -Trp, -Leu, -Ade and -His supplemented with X-α-Gal and Aurcobasidin A (QDO/X/A). The empty BD and AD vectors were used as the negative controls. FIG. 10B Bimolecular fluorescence complementation (BiFC) assay. The coding sequence of SDE15 (without its signal peptide) fused to that of the N-terminal or C-terminal fragment of EYFP in vectors pSAT6-nEYFP-C1 and pSAT6-cEYFP-C1-B, respectively, was co-transformed into *Citrus* leaf protoplasts with full-

length CsACD2 or CsACD2 protein which fused to the C-terminal or N-terminal fragment of EYFP in vectors pSAT6-nEYFP-C1 and pSAT6-cEYFP-C1-B. The EYFP fluorescence of protoplasts were imaged, 1 day after incubation, using a Leica fluorescence microscope. Co-transformations of SDE15-EYFP^N and EYFP^C, EYFP^N and EYFP^C-SDE15, SDE15-EYFP^N and EYFP^C-SDE15, CSACD2-EYFP^N and EYFP^C, EYFP^N and EYFP^C-CsACD2, CSACD2-EYFP^N and EYFP^C-CsACD2, EYFP^N+EYFP^C were used as negative controls, which did not produce any detectable fluorescence signal. FIG. 10C, FIG. 10D, FIG. 10E. Glutathione-S-transferase (GST) pull-down assay. GST-SDE15 and GST empty vectors were expressed in *E. coli*, immobilized on glutathione sepharose beads, and incubated with *E. coli* lysate containing MBP-CsACD2. Total cell extract (Input) and eluted protein (Elute) were immunoblotted using the anti-MBP and anti-GST antibody.

[0020] FIG. 11. Subcellular localization of CsACD2 and co-localization of SDE15 and CsACD2. CsACD2-EYFP was co-expressed with the plasma membrane localization-marker PM-CFP, the nucleus-marker CFP-nucleus or SDE15-CFP in leaves of *N. benthamiana*. *A. tumefaciens* strain GV2260 harboring the corresponding plasmids were infiltrated into leaves at OD₆₀₀ of 0.2. Subcellular localization was inspected and photographed 1 day post infiltration. Scale bars: 10 μ m.

[0021] FIGS. 12A-C. SDE15 represses the hypersensitive response in tobacco and promotes the RCCR activity of CsACD2 in vitro. FIG. 12A. Hypersensitive response (HR) assay. *A. tumefaciens* strain GV2260 harboring binary vectors that are designed to express SDE15, CsACD2 (Left) or truncated SDE15 (Right) were co-infiltrated into leaves of *N. benthamiana* at the concentration of 10⁸ CFU ml⁻¹. Two days later, another *A. tumefaciens* strain GV2260 harboring the binary vector that is designed to express AvrBsT protein, which can trigger an HR was infiltrated on the same area of the leaves. HR induction was observed and photographed 2-3 days post-inoculation. All experiments were repeated three times with the similar results. FIG. 12B. Electrolyte leakage associated with the HR induced by AvrBsT 2 days post infiltration. Leaf discs were floated on deionized water with shaking. The conductivity of the solution was measured after 4 h shaking. Error bars indicate standard error of mean (n=3). Alphabets represent significant differences in different types of samples. FIG. 12C. Coupled PAO/RCCR assay to measure CsACD2 activity. Activity of purified recombinant CsACD2 was assessed in a coupled assay using purified PAO and co-factors. pFCC as the product was measured by HPLC. Purified GST-SDE15, SDE15^{AN} or SDE15^{AC} were added to the reaction mixture to examine whether full-length SDE15 and truncated SDE15 proteins affect the activity of CsACD2. As negative controls, purified GST protein or mock purification of the vector alone without CsACD2 was added to the reaction system. Error bars represent SD (n=3). This experiment was done twice with similar results.

BRIEF DESCRIPTION OF THE SEQUENCES

[0022] SEQ ID NO:1 Antigen sequence used to produce CLIBASIA_04025 (Las4025)-specific antibody from *Ca. Liberibacter asiaticus*.

[0023] SEQ ID NO:2 CLIBASIA_04025 cDNA sequence from *Ca. Liberibacter asiaticus*.

[0024] SEQ ID NO:3 CLIBASIA_00470 cDNA sequence from *Ca. Liberibacter asiaticus*.

[0025] SEQ ID NO:4 CLIBASIA_04065 cDNA sequence from *Ca. Liberibacter asiaticus*.

[0026] SEQ ID NO:5 CLIBASIA_05150 cDNA sequence from *Ca. Liberibacter asiaticus*.

[0027] SEQ ID NO:6 CLIBASIA_04250 cDNA sequence from *Ca. Liberibacter asiaticus*.

[0028] SEQ ID NO:7 ACD2 Cs1g22670 gene sequence from *Citrus sinensis* cultivar Valencia.

[0029] SEQ ID NO:8 ACD2 Cs1g22670.1 cDNA sequence from *Citrus sinensis* cultivar Valencia.

[0030] SEQ ID NO:9 AT4G37000 DNA sequence from *Arabidopsis thaliana*.

[0031] SEQ ID NO:10 AT4G37000 protein sequence from *Arabidopsis thaliana*.

[0032] SEQ ID NO:11 Cysteine Protease Cs4g07410 gene sequence from *Citrus sinensis* cultivar Valencia.

[0033] SEQ ID NO:12 Cysteine Protease Cs4g07410.1 cDNA variant sequence from *Citrus sinensis* cultivar Valencia.

[0034] SEQ ID NO:13 Cysteine Protease CS4g07410.2 cDNA variant sequence from *Citrus sinensis* cultivar Valencia.

[0035] SEQ ID NO:14 RCCR-like Cs1g22680 gene sequence from *Citrus sinensis* cultivar Valencia.

[0036] SEQ ID NO:15 RCCR-like Cs1g22680.1 cDNA sequence from *Citrus sinensis* cultivar Valencia.

[0037] SEQ ID NO:16 Lls1 Cs9g02990 gene sequence from *Citrus sinensis* cultivar Valencia.

[0038] SEQ ID NO:17 Lls1 Cs9g02990.1 cDNA sequence from *Citrus sinensis* cultivar Valencia.

[0039] SEQ ID NO:18 ACD1-like Cs9g03000 gene sequence from *Citrus sinensis* cultivar Valencia.

[0040] SEQ ID NO:19 ACD1-like Cs9g03000.1 cDNA sequence from *Citrus sinensis* cultivar Valencia.

[0041] SEQ ID NO:20 ACD1 Cs8g15480 gene sequence from *Citrus sinensis* cultivar Valencia.

[0042] SEQ ID NO:21 ACD1 Cs8g15480.1 cDNA sequence from *Citrus sinensis* cultivar Valencia.

[0043] SEQ ID NO:22 Cysteine Proteinase 15A-like Cs3g25530 gene sequence from *Citrus sinensis* cultivar Valencia.

[0044] SEQ ID NO:23 Cysteine Proteinase 15A-like Cs3g25530.1 cDNA sequence from *Citrus sinensis* cultivar Valencia.

[0045] SEQ ID NO:24 Myb orange1.1t02260 gene sequence from *Citrus sinensis* cultivar Valencia.

[0046] SEQ ID NO:25 Myb orange1.1t02260.1 cDNA sequence from *Citrus sinensis* cultivar Valencia.

[0047] SEQ ID NO:26 YLS9-like Cs2g29120 gene sequence from *Citrus sinensis* cultivar Valencia.

[0048] SEQ ID NO:27 YLS9-like Cs2g29120.1 cDNA sequence from *Citrus sinensis* cultivar Valencia.

[0049] SEQ ID NO:28 Lectin orange1.1t05126 gene sequence from *Citrus sinensis* cultivar Valencia.

[0050] SEQ ID NO:29 Lectin orange1.1t05126.1 cDNA sequence from *Citrus sinensis* cultivar Valencia.

[0051] SEQ ID NO:30 PP2-B2/12 orange1.1t04174 gene sequence from *Citrus sinensis* cultivar Valencia.

[0052] SEQ ID NO:31 PP2-B2/12 orange1.1t04174.1 cDNA sequence from *Citrus sinensis* cultivar Valencia.

[0053] SEQ ID NO:32 NDR1/HIN1-like protein 13 Cs8g01640 gene sequence from *Citrus sinensis* cultivar Valencia.

[0054] SEQ ID NO:33 NDR1/HIN1-like protein 13 Cs8g01640.1 cDNA sequence from *Citrus sinensis* cultivar Valencia.

[0055] SEQ ID NO:34 PHL5 orange1.1t02259 gene sequence from *Citrus sinensis* cultivar Valencia.

[0056] SEQ ID NO:35 PHL5 orange1.1t02259.1 cDNA variant sequences from *Citrus sinensis* cultivar Valencia.

[0057] SEQ ID NO:36 PHL5 orange1.1t02259.2 cDNA variant sequences from *Citrus sinensis* cultivar Valencia.

[0058] SEQ ID NO:37 PHL5-like Cs7g01290 gene sequence from *Citrus sinensis* cultivar Valencia.

[0059] SEQ ID NO:38 PHL5-like Cs7g01290.1 cDNA sequence from *Citrus sinensis* cultivar Valencia.

[0060] SEQ ID NO:39 ACD2 LOC102591737 gene sequence from *Solanum tuberosum*.

[0061] SEQ ID NO:40 ACD2 NM_001318612.1 CDS sequence from *Solanum tuberosum*.

[0062] SEQ ID NO:41 Cysteine Proteinase 3 LOC102578939 gene sequence from *Solanum tuberosum*.

[0063] SEQ ID NO:42 Cysteine Proteinase 3 XM_006342320.2 CDS sequence from *Solanum tuberosum*.

[0064] SEQ ID NO:43 Lls1 LOC102597185 gene sequence from *Solanum tuberosum* (bp 734-7789).

[0065] SEQ ID NO:44 Lls1 XM_006364374.2 CDS sequence from *Solanum tuberosum*.

[0066] SEQ ID NO:45 Lls1/ACD1-like LOC102604461 gene sequence from *Solanum tuberosum*.

[0067] SEQ ID NO:46 Lls1/ACD1-like XM_006340026.2 CDS sequence from *Solanum tuberosum*.

[0068] SEQ ID NO:47 YLS9-like LOC102602250 gene sequence from *Solanum tuberosum*.

[0069] SEQ ID NO:48 YLS9-like NM_001289011.1 CDS sequence from *Solanum tuberosum*.

[0070] SEQ ID NO:49 Myb family transcription factor LOC102578723 gene sequence from *Solanum tuberosum*.

[0071] SEQ ID NO:50 Myb family transcription factor XM_006362170.2 CDS sequence from *Solanum tuberosum*.

[0072] SEQ ID NO:51 Lls1 LOC102597185 gene sequence from *Solanum tuberosum* (bp 1-144).

DETAILED DESCRIPTION

Introduction

[0073] The disclosure provides a modified plant comprising a genetic modification to an endogenous gene or regulatory element thereof, wherein it is believed that the polypeptide encoded by said endogenous gene interacts with Sec-dependent pathway effector polypeptides secreted by pathogenic species of *Ca. Liberibacter*. The cysteine protease gene may be modified such that expression of the endogenous gene is knocked-down or reduced, or otherwise modified such that interaction with SDE is reduced. In specific examples, the endogenous gene is cysteine protease and the modified plant is *Citrus*, wherein the *Citrus* plant exhibits increased resistance to HLB as a result of the modification. Also provided are seeds, fruit, and plant parts of such plants. In another embodiment, methods are provided for generating a modified plant that is tolerant to *Ca. Liberibacter* infection, such as *Citrus* plant that is tolerant to HLB. Methods are also provided for conferring plants with resistance to *Ca. Liberibacter* infection, such as conferring *Citrus* plants with a resistance to HLB, and screening that plurality of plants for said resistance. In specific examples this is accomplished using nucleic acid modification tech-

niques, genome recombination techniques, genome editing techniques, or a combination thereof.

Definitions

[0074] Expression: The term “expression” as used herein refers to the transcription of a particular nucleic acid sequence to produce sense or antisense RNA or mRNA, and/or the translation of an mRNA molecule to produce a polypeptide, with or without subsequent post-translational events. Expression also encompasses production of a functional nucleic acid (e.g., an RNAi, antisense molecule, ribozyme, aptamer, etc.).

[0075] Genome editing: Modifying a genome with techniques that employ targeted mutagenesis to activate DNA repair pathways. These techniques include, but are not limited to, those that utilize endonucleases to generate single-strand and double-strand DNA breaks that activate DNA repair pathways. Genome editing techniques may also comprise systems that enable targeted editing at any genomic locus. These targeting systems include, but are not limited to, polypeptides, such as, Transcription Activator-Like Effectors (TALEs) and zinc fingers (ZFs), or nucleic acids, such as, Clustered Regularly Interspaced Short Palindromic Repeats/Cas (CRISPR/CAS) single guide RNAs or NgAgo (Argonaute) single strand DNAs. As used herein, “genome editing” and “genome-engineering” are interchangeable.

[0076] Genetic modification: A DNA sequence difference, epigenetic difference, or combination thereof between two genomes of the same species in which one genome is identified as the modified genome and the other is identified as the unmodified genome and the DNA sequence or epigenetic difference is the result of applying genome modifying techniques to the unmodified genome to yield the modified genome. A genetic modification, as used herein, encompasses any insertion, deletion, or substitution of a nucleotide sequence of any size and nucleotide content, any epigenetic modification to any number of nucleotides, or a combination thereof. A genetic modification, as used herein, may also encompass introduction of one or more exogenous coding nucleic acids that do not integrate into the unmodified genome, yet are capable of autonomous replication. In certain embodiments, a modification to an endogenous gene or regulatory element thereof may be a deletion, a substitution, or an insertion that reduces expression of the endogenous gene or the polypeptide for which it encodes. In specific embodiments, the modification may be an indel, wherein the indel may cause a frameshift mutation, a missense mutation, a nonsense mutation, a neutral mutation, or a silent mutation. In specific embodiments, a modification to a regulatory element of an endogenous gene may alter or eliminate a function of the regulatory element. In further contemplated embodiments, the modification may comprise a nucleic acid sequence that provides exogenous control of endogenous gene, mRNA, or polypeptide expression levels. In specific embodiments, the modification may also disrupt a post-translational process of a polypeptide encoded by an endogenous gene. Post-translational processes in certain embodiments may be post-translational modification, protein sorting, or proteasomal degradation.

[0077] Genetically modified cell: A cell in which the endogenous genome has been genetically modified; a cell in which one or more exogenous, coding nucleic acids have

been introduced that do not integrate into the genome, yet are capable of autonomous replication; or a combination thereof.

[0078] Genetically modified plant: A plant comprising at least one genetically modified cell. A genetically modified plant may be regenerated from a genetically modified cell or plant part comprising genetically modified cells, and thus the genetic modification may be heritable and inherited by progeny thereof. The progeny thereof that inherit the genetic modification are also considered genetically modified plants. A genetically modified plant, as used herein, also refers to a plant in which at least one genetically modified cell is introduced to a plant or arises as a result of genetic modification techniques directly applied to the plant.

[0079] Genetic modification techniques: Any technique known to those in the art that can modify the genome of a cell including, but not limited to, genome editing, site-specific genetic recombination, epigenetic modifications, and genetic transformation.

[0080] Genetic Transformation: A process of introducing a DNA sequence or construct (e.g., a vector or expression cassette) into a cell or protoplast in which that exogenous DNA is incorporated into a chromosome or is capable of autonomous replication.

[0081] Heterologous: A sequence which is not normally present in a given host genome in the genetic context in which the sequence is currently found. In this respect, the sequence may be from another species, organism, plant, tree, or variety, or may be native to the host genome, but be rearranged with respect to other genetic sequences within the host sequence. For example, a regulatory sequence may be heterologous in that it is linked to a different coding sequence relative to the native regulatory sequence. In addition, a particular recombinant DNA molecule may be heterologous with respect to a cell or organism into which it is inserted when it would not naturally occur in that particular cell or organism.

[0082] Overexpress: As used herein, “overexpress” refers to increased expression of a gene or coding sequence over that found in nature or a control plant or tissue. In some embodiments, “overexpress” may refer to greater expression of a gene or coding sequence in a genetically modified plant, when compared to a plant lacking the genetic modification.

[0083] Plant: As used herein, the term “plant” refers to *Citrus* or solanaceous plant, or any other plant that can be infected by a *Ca Liberibacter* species.

[0084] Plant part: The term “plant part” refer to cells, tissues, organs, seeds, and severed parts (e.g., roots, leaves, and flowers) that retain the distinguishing characteristics of the parent plant. “Seed” refers to any plant structure that is formed by continued differentiation of the ovule of the plant, following its normal maturation point at flower opening, irrespective of whether it is formed in the presence or absence of fertilization and irrespective of whether or not the seed structure is fertile or infertile. A plant part may be any part of the plant from which another plant may arise.

[0085] Promoter: A recognition site on a DNA sequence or group of DNA sequences that provides an expression control element for a structural gene and to which RNA polymerase specifically binds and initiates RNA synthesis (transcription) of that gene.

[0086] R_0 genetically modified plant: A plant that has been genetically modified or has been regenerated from a plant cell or cells that have been genetically modified.

[0087] Reduction of Expression: The term “Reduc(e), (es) or (ing) the expression” of a gene or polypeptide in a plant or a plant cell includes inhibiting, interrupting, knocking-out, or knocking-down the gene or polypeptide, such that transcription of the gene and/or translation of the encoded polypeptide is reduced as compared to a corresponding control plant, plant cell, or population of plants or plant cells in which expression of the gene or polypeptide is not inhibited, interrupted, knocked-out, or knocked-down. “Reduced expression” encompasses any decrease in expression level (e.g., a decrease of 10% or more, 20% or more, 30% or more, 40% or more, 50% or more, 60% or more, 70% or more, 80% or more, 90% or more, or even 100%) as compared to the corresponding control plant, plant cell, or population of plants or plant cells. In some embodiments, reducing expression by 50% or more may be particularly useful. Expression levels can be measured using methods such as, for example, reverse transcription-polymerase chain reaction (RT-PCR), Northern blotting, dot-blot hybridization, in situ hybridization, nuclear run-on and/or nuclear run-off, RNase protection, or immunological and enzymatic methods such as ELISA, radioimmunoassay, and western blotting.

[0088] Regeneration: The process of growing a plant from a plant cell (e.g., plant protoplast, callus, or explant).

[0089] Rootstock: As used herein, a “rootstock” refers to underground plant parts such as roots, from which new above-ground growth of a plant or tree can be produced. In accordance with the disclosure, a rootstock may be used to grow a different variety through asexual propagation or reproduction such as grafting. As used herein, a “scion” refers to a plant part that is grafted onto a rootstock variety. A scion may be from the same or a different plant type or variety.

[0090] Site-specific genome modification: Any genome modification technique that employs an enzyme that can modify a nucleotide sequence in a sequence-specific manner. Site-specific genome modification enzymes include, but are not limited to, nucleases, endonucleases, recombinases, invertases, transposases, methyltransferases, demethylases, aminases, deaminases, helicases, and any combination thereof.

[0091] Transformation construct: A chimeric DNA molecule which is designed for introduction into a host cell by genetic transformation. Preferred transformation constructs will comprise all of the genetic elements necessary to direct the expression of one or more exogenous nucleic acid sequences. In particular embodiments of the instant disclosure, it may be desirable to introduce a transformation construct into a host cell in the form of an expression cassette.

[0092] Transgene: A segment of DNA which has been incorporated into a host genome or is capable of autonomous replication in a host cell and is capable of causing the expression of one or more nucleic acid sequences. Exemplary transgenes will provide the host cell, or plants regenerated therefrom, with a novel phenotype relative to the corresponding non-transformed cell or plant. Transgenes may be directly introduced into a plant by genetic transformation, or may be inherited from a plant of any previous generation which was modified with the DNA segment.

[0093] Vector: A DNA molecule designed for transformation into a host cell. Some vectors may be capable of

replication in a host cell. A plasmid is an exemplary vector, as are expression cassettes isolated therefrom.

[0094] Tolerance or resistance: Tolerance encompasses any relief from, reduced presentation of, improvement of, or any combination thereof of any symptom of an infection by a *Ca. Liberibacter* species. Resistance encompasses tolerance as well as a reduction of bacteria upon infection or reduction of ability to infect by a *Ca. Liberibacter* species. In specific embodiments of the disclosure, *Citrus* plant may be provided that are defined as comprising a complete or less than complete resistance or tolerance to HLB. This may be assessed, for example, relative to a *Citrus* plant not comprising a genetic modification according to the disclosure.

[0095] Hypersensitive Response (or Reaction): The hypersensitive response (or sometimes referred to a hypersensitive reaction) (HR) is plant defense mechanism that protects a plant against infection by a plant pathogen. HR is a form of cell death often associated with plant resistance to pathogen infection to prevent the spread of the potential pathogen from infected to uninfected tissues. Cell death is activated by recognition of pathogen-derived molecules by the resistance (R) gene products, and is associated with the massive accumulation of reactive oxygen species (ROS), salicylic acid (SA), and other pro-death signals such as nitric oxide (NO). *Ca. Liberibacter* species inhibit hypersensitive response, which inhibits the plant from defending itself against the *Ca. Liberibacter*, *Xanthomonas* species, and other pathogens. It is shown herein that secretion of SDEs by a bacterial species inhibit HR. The genomic modifications described herein prevent or minimize inhibition of HR by SDES.

Detailed Description of Embodiments

[0096] The disclosure provides a significant improvement over the art due to the lack of agronomically acceptable *Citrus* plants with tolerance or resistance to HLB. HLB is a disease caused by species of the phloem-limited, gram-negative bacteria of genus *Ca. Liberibacter*. In the U.S., the predominant pathogenic species is *Ca. Liberibacter asiaticus* (Las); whereas *Ca. Liberibacter africanus* (Laf) and *Ca. Liberibacter americanus* (Lam) are the predominant pathogenic species in South Africa and Brazil, respectively. *Ca. Liberibacter* is a vector-transmitted pathogen. The vector organisms are the Asian *Citrus* psyllid, *Diaphorina citri*, and African *Citrus* psyllid, *Trioza erythrae*. HLB was first detected in the United States in August 2005 and has rapidly moved into several *Citrus* producing areas. All commercial *Citrus* plants are susceptible to HLB, and infected *Citrus* plants will irrevocably decline. Plant decline is usually preceded by a decline in the quality of the fruit and fruit drop. Fruit from infected plants are smaller, yield less juice, and have higher acidity, lower sugar and greener peel color than those from uninfected plants.

[0097] HLB has resulted in a severe decline in fruit production in Florida, where it has become endemic. However, due to the lack of rapid curative methods that control HLB, prevention of new infections is essential in HLB management. Currently, HLB management consists of preventing trees from becoming infected, which includes protecting young flush from HLB vector organisms and destroying infected plant material.

[0098] New infections could be prevented, and the disease could be managed, by planting trees that are tolerant or resistant to the disease. However, utilization of resistant

germplasm to slow the spread of HLB is difficult due to the lack of commercially available resistant rootstock/scion combinations. Identification and incorporation of resistance traits from tolerant *Citrus* species and relatives is also a potential disease management strategy, but applying conventional plant breeding methods to *Citrus* plants is difficult and time consuming due to their level of nucellar embryony and long juvenile phases.

[0099] Genetically modifying *Citrus* plants is a viable alternative to conventional plant breeding. It is a relatively rapid process and some techniques allow for targeted modification of genetic locus without significant off-target effects. In such cases, genetic modification of existing cultivars has been a key component to combat HLB. In some embodiments, the disclosure employs genetic modification to render the modified *Citrus* plant tolerant to pathogenic *Ca. Liberibacter* species. In specific embodiments, the disclosure provides a *Citrus* plant that is tolerant to *Ca. Liberibacter* effector proteins. As will be understood to those of skill in the art, once a genetic modification conferring resistance to HLB is generated this could readily be introduced into any other cultivar by crossing.

[0100] Zebra Chip (ZC) is an economically important disease that occurs in commercial potato fields in the United States, Mexico, Central America, and New Zealand (Munyanca, J. E. Am. J. Pot Res (2012) 89: 329). ZC was first found in Mexico in 1994 then spread to the United States in 2000 (Rondon, S., Schreiber, A., Hamm, P., Olsen, N., Wenninger, E., Wohleb, C., Waters, T., Cooper, R., Walenta, D., and Reitz, S. 2017. Potato Psyllid Vector of Zebra Chip Disease in the Pacific Northwest. A Pacific Northwest Extension Publication. pp. 1-8.). Similar to Huanglongbing (HLB), ZC and diseases of other solanaceous crops are associated with a fastidious alpha-proteobacterium belonging to the ‘*Candidatus*’ genus *Liberibacter*, ‘*Candidatus Liberibacter solanacearum*’ (CLso), that is transmitted by a phloem-feeding psyllid vector, *Bactericera cockerelli* (Jagoueix, S., et al. 1994, Int. J. Syst. Bacteriol. 44:379-386.; Bové, J. M. 2006, J. Plant Pathol. 88:7-37; Pelz-Stelinski et al. 2010, J. Econ. Entomol. 103:1531-1541). CLso vectored by *B. cockerelli* results in a severe decline of potato, tomato, and pepper production. Current management of CLso consists of chemical controls using insecticides (Rondon et al. 2017). Due to the rapid spread of CLso, new methods to prevent infections are required.

[0101] Exemplary *Ca. Liberibacter* effector proteins contemplated by this disclosure are those secreted via the Sec-dependent pathway. Sec-dependent effector (SDE), as used herein, refers to any bacterial effector protein secreted from a bacterium via the Sec-dependent pathway. Pathogenic *Ca. Liberibacter* species secrete SDEs into the phloem of host plants, such as *Citrus* and solanaceous crops. As used herein, the terms “solanaceous crop” or “solanaceous plant” are used interchangeably and are directed plants of the Solanacea family including tomato (*Solanum lycopersicum* and *Solanum pennellii*); potato (*Solanum tuberosum*); eggplant (*Solanum melongena*), bell/chili peppers (*Capsicum annuum*, *Capsicum baccatum*, and *Capsicum chinense*). These SDEs interact with endogenous proteins and nucleic acids in the phloem and companion cells, disrupting normal physiology and inducing the symptoms of HLB in *Citrus*. Moreover, this same interaction can occur with SDEs related to *Ca. Liberibacter* species (e.g. *Ca. Liberibacter solan-*

acearum, 'CLso') infection in solanaceous crops such as tomato, pepper eggplant, tamarillo and potato (e.g. zebra chips).

[0102] Here we show that one of these secreted proteins, SDE15 also known as Las4025, targets a well-known negative regulator of plant programmed cell death (PCD) to promote infection. Las4025 could be detected in the phloem sap of Las-infected plants. Transgenic expression of Las4025 in *Citrus* promotes Las multiplication and HLB symptom development. SDE15 suppresses not only PCD induced by *Xanthomonas citri* subsp. *citri* (Xcc) in *Citrus*, but also PCD induced by AvrBsT (a PCD-eliciting *Xanthomonas* effector) in tobacco, suggesting that Las4025 is a broad-spectrum bacterial suppressor of plant PCD. Yeast two-hybrid, in vitro protein pull-down and in vivo bimolecular fluorescence complementation assays showed that SDE15 interacts with ACD2 (ACCELERATED CELL DEATH 2), a repressor of plant PCD and that it enhances the red chlorophyll catabolite reductase (RCCR) activity of ACD2 to remove porphyrin-related molecules, accumulation of which causes PCD. Las4025 promotes the chlorophyll break-down in planta and contributes to the development of yellowing symptom associated with HLB. Characterization of Las4025 unravels an elusive aspect of the mechanism of a major plant disease.

[0103] In some embodiments, a modified plant no longer expresses endogenous molecules, for example, polypeptides and nucleic acids, capable of interacting with *Ca. Liberibacter* SDEs. In specific embodiments, the SDEs are those secreted by Las. In more specific embodiments, an Las SDE is selected from CLIBASIA_04025 (Las4025), CLIBASIA_00470 (Las470), CLIBASIA_04065 (Las4065), CLIBASIA_05150 (Las5150), and CLIBASIA_04250 (Las4250), which are encoded by the cDNA sequences corresponding to SEQ ID NOs: 2-6.

[0104] In other embodiments, a modified plant no longer expresses an endogenous molecule, for example, a polypeptide or nucleic acid, and which may be capable of interacting with a *Ca. Liberibacter* Sec-dependent effector (SDE), that is modified. A susceptibility protein or S-protein, as used

herein, refers to an endogenous host polypeptide targeted by an SDE. A susceptibility gene or S-gene, as used herein, refers to an endogenous host gene encoding an S-protein. An S-protein-SDE complex, as used herein refers to an S-protein interacting with an SDE. An S-protein-SDE interaction, as used herein refers to a protein-protein interaction between an S-protein and an SDE. In some embodiments, an S-gene is modified such that the encoded S-protein is no longer capable of interacting with an SDE. In other embodiments, an S-gene is modified such that the encoded S-protein may interact with an SDE, but not disrupt normal physiology to an extent that a deleterious mechanism of action is triggered, for a non-limiting example, a modified S-gene that promotes proteasomal degradation of an SDE-S-protein complex before the complex activates a deleterious mechanism of action. In specific embodiments, an S-protein is selected from the group consisting of [accession numbers for *Citrus* provided in parentheses for each S-protein group] PP2-B2/12 (orange1.1t04174), Lectin (orange1.1t05126), Cysteine protease (Cs4g07410), Cysteine protease 15A-like (Cs3g25530), Papain-like cysteine proteases, Myb family transcription factor (orange1.1t02260), YLS9-like (Cs2g29120), Cell death suppressor protein Lls1 (Cs9g02990.1), Acd1-Like Cs9g03000, Acd1 Cs8g15480, accelerated cell death 2 (ACD2) protein (AT4G37000.1, Cs1g22670), red chlorophyll catabolite reductase-like (Cs1g22680), NDR1/HIN1-like protein 13 (Cs8g01640), PHL5-like (Cs7g01290), and PHL5 (orange1.1t02259), for which cDNA examples of the *Citrus* versions are encoded by the cDNA sequences corresponding to SEQ ID NOs: 7-29 and SEQ ID NO:31. Potato orthologs are encoded by SEQ ID Nos 39-51. Provided below in Table 1 are accession numbers for select *Citrus* S-protein sequences and S-genes encoding such S-proteins, as well as orthologs in solanaceous plants, that may be modified as taught herein (Cs or orange1= gene id (*Citrus* genome database citrusgenomedb.org); NC or NW= genome sequence (NCBI database); XM= cDNA accession no. (NCBI database); LOC= gene accession no (NCBI database); and XP, NP, PHT or PHU= polypeptide accession no. (NCBI database)):

TABLE 1

Species	Myb family transcription	Cell death suppressor protein Lls1	ACD2	Lectin
<i>Citrus sinensis</i>	1. orange1.1t02260	1. Cs9g02990.1	1. Cs1g22680	1. orange1.1t05126
	LOC102621262	LOC102615553	XM_006466545	LOC102630138
	XM_025093288	XM_006488849	LOC102623285	XM_006495169
	XP_024949056	XP_006488912	XP_006466608	XP_006495232
	NW_006257094.1	NC_023054.1	NC_023046.1	NW_006257465.1
	(602674 ... 604650)	(1382910 ... 1386383)	(25356426 ... 25358668)	(8465 ... 9844)
	2. XP_024949055	2. Cs9g03000	2. Cs1g22670	2. LOC107177625
	3. orange1.1t02259	XM_006488848	NC_023046.1	XM_015531689
	LOC102608693	LOC102615272	XM_006466544	XP_015387175
	XM_015525401	XP_006488911	(LOC102622999)	3. XP_015387176
	XP_015380887	NC_023054.1	XP_006466607	LOC107177626
	NW_006257094.1	(1390097 ... 1395026)	NC_023046.1	4. XP_015387172
	(597073 ... 599506)	3. Cs8g15480,	(25352463 ... 25354069)	LOC107177622
	4. LOC102608059	XM_006487933		6. XP_006475932
	XM_015531655	XP_006487996		LOC102628131
<i>Capsicum annuum</i>	XP_015387141	NC_023053.1		7. XP_015387174,
	5. XP_015380888.1	(18675482 ... 18679902)		LOC107177624
	LOC107843940,	PHT80565,	LOC107868112,	PHT71355,
	XP_016543872,	XP_016571811,	PHT83236,	PHT71353
	XP_016543873,	XP_016571812,	XP_16570190,	
	XP_016576061,	XP_016571813	NP_001311893,	
	PHT77744,		XP_016557361	
	XP_016576060,			

TABLE 1-continued

<i>Capsicum baccatum</i>	XP_016565589, PHT82561, XP_016565591, XP_016573871, XP_016544890, PHT71362 PHT37283, PHT44431, PHT52438, PHT48923, PHT33349, PHT60401, PHT39991	PHT54548, PHT54549, PHT33802, PHT33064, PHT45532	PHT50387, PHT58680	PHT37058
<i>Capsicum chinense</i>	PHU06047, PHU13451, PHU22243, PHU18686, PHU05798	PHU16680, PHU16678, PHU16679, PHU03790, PHU02689, PHU01515	PHU19541	PHT99312, PHU_05788
<i>Solanum lycopersicum</i>	1. LOC101251632 NC_015447.3 (60577349 . . . 60580659 complement)	1. LOC101255583, NC_015441.3 (11974361 . . . 11979604) XP_004237332, AAL32300, NP_001234535	1. LOC778267 NC_015440.3 (9353792 . . . 9357403)	N/a
<i>Solanum pennellii</i>	LOC107032497(XP_015089588), XP_015088029, XP_015088022, XP_015078237, XP_015078236, XP_015076100, XP_015072629, XP_015055275	XP_015073606, XP_015058211, XP_015072446	LOC107014711, XP_015070234	N/a
<i>Solanum tuberosum</i>	1. LOC102578723 NW_006239309.1 (222724 . . . 226103)	1. LOC102597185 NW_006239415.1 (360276 . . . 368064) 2. LOC102604461 NW_006238942.1 (19429 . . . 23712)	1. LOC102591737, NW_006239292.1 (123266 . . . 127905 complement) NP_001305541	N/a
Species	Cysteine protease	PP2-B12	YLS9-like	
<i>Citrus sinensis</i>	1. Cs4g07410 LOC102578016 XM_006474664 NM_001288897 NP_001275826 NC_023049.1 (4697175 . . . 4700328) 2. Cs3g25530 XM_006473521 XP_006473584 LOC102608509 NC_023048.1 (27116634 . . . 27118954)	1. orange1.1t04174 LOC102626181 XM_025094054 XP_024949822 NW_006257165.1 (68010 . . . 79424)	1. Cs2g29120 LOC102624273 XM_006470378 XP_006470441 NC_023047.1 (28676278 . . . 28677278) 2. Cs2g29120 LOC107174220 NC_023053.1 (385414 . . . 385806 complement)	
<i>Capsicum annuum</i>	XP_016580127, XP_016557040, PHT84613, XP_016539529, PHT90352, XP_016561024, PHT70914	PHT66823, XP_016552209, XP_016552365, PHT66822, XP_016573353, XP_016552297, XP_016573817	XP_016552935, XP_016561811, XP_016560224, XP_016563876, PHT87859, XP_016562568, PHT62176, PHT71162, XP_016562568	
<i>Capsicum baccatum</i>	PHT42963, PHT30454, PHT41774, PHT_57030, PHT_36437, PHT30404, PHT30386, PHT59073	PHT32781, PHT32780, PHT32035, PHT39101, PHT32776, PHT32782, PHT48962	PHT59305, PHT54760, PHT27977, PHT53910, PHT29394, PHT43389, PHT36895	

TABLE 1-continued

<i>Capsicum chinense</i>	PHU11729, PHU20763, PHU10453, PHU27226, PHU05443	PHU01437, PHU01440, PHU00690, PHU21394, PHU01435, PHU18677, PHT98988, PHU10308	PHU29409, PHU24981, PHT98447, PHU24513, PHU12324, BAD11071
<i>Solanum lycopersicum</i>	1. LOC101252505 NC_015444.3 (54626241 . . . 54628525 complement)	XP_004239687, XP_004253004, XP_004237494, XP_004237583, XP_010314855, XP_004252380	1. LOC101250915 NC_015438.3 (3106210 . . . 3109693 complement)
<i>Solanum pennellii</i>	XP_015082349, XP_015081027, XP_015074247, XP_015068628, XP_015061093, XP_015058018, XP_015063485, XP_015069437	XP_015074926, XP_015071726, XP_015059542, XP_015073190, XP_01505954, XP_015084179, XP_015060715	XP_015084054, XP_015086729, XP_015065001, XP_015067199, XP_015070124, XP_015081836
<i>Solanum tuberosum</i>	1. LOC102578939 NW_006238961.1 (2218619 . . . 2220963 complement)	XP_006349935, XP_006345814, XP_006340500, XP_006366161, XP_015165222, XP_006344703, XP_006361502, XP_006351708	1. LOC102602250 NW_006238997.1 (685210 . . . 686001 complement)

[0105] This disclosure also contemplates embodiments in which a genetic modification anywhere in the genome disrupts expression, and in turn, may disrupt an S-protein-SDE interaction from activating a deleterious mechanism of action. In some embodiments, an S-gene or a regulatory element thereof is modified. In some embodiments, a modification may be made elsewhere in the genome, but ultimately disrupts an S-protein-SDE interaction, for a non-limiting example, inserting genetic material encoding a polypeptide capable of disrupting an S-protein-SDE interaction anywhere in the genome. In specific embodiments, an SDE is secreted the *Ca. Liberibacter* species selected from the group consisting of Las, Laf, and Lam. In some specific embodiments, the SDEs are CLIBASIA_0402 (Las4025), CLIBASIA_00470 (Las470), CLIBASIA_04065 (Las4065), CLIBASIA_05150 (Las5150), and CLIBASIA_04250 (Las4250). In specific embodiments, an S-protein is selected from the group consisting of PP2-B2/12 (orange 1.1t04174), Lectin (orange1.1t05126), Cysteine protease (Cs4g07410), Cysteine protease 15A-like (Cs3g25530), Papain-like cysteine proteases, Myb family transcription factor (orange 1.1t02260), YLS9-like (Cs2g29120), Cell death suppressor protein Lls 1 (Cs9g02990.1), Acd1-Like Cs9g03000, Acd1 Cs8g15480, accelerated cell death 2 (ACD2) protein (AT4G37000.1, Cs1g22670), red chlorophyll catabolite reductase-like (Cs1g22680), NDR1/HIN1-like protein 13 (Cs8g01640), PHL5-like (Cs7g01290), and PHL5 (orange1.1t02259).

[0106] In some cases, a modification is conducted at a target sequence as set forth in Table 1, or at a target sequence that is at least 95 percent (e.g., at least 96 percent, at least 97 percent, at least 98 percent, or at least 99 percent) identical to the sequence set forth in Table 1. In a more specific example, a modification is conducted at a target sequence set forth in SEQ ID Nos 7-38 or 39-51, or at a target sequence that is at least 95 percent (e.g., at least 96 percent, at least 97

percent, at least 98 percent, or at least 99 percent) identical to a sequence set forth in SEQ ID Nos 7-38 or 39-51.

[0107] The percent sequence identity between a particular nucleic acid or amino acid sequence and a sequence referenced by a particular sequence identification number may be determined by techniques known in the art. In one example, sequence identity is determined as follows. First, a nucleic acid or amino acid sequence is compared to the sequence set forth in a particular sequence identification number using the BLAST 2 Sequences (Bl2seq) program from the stand-alone version of BLASTZ containing BLASTN version 2.0.14 and BLASTP version 2.0.14. This stand-alone version of BLASTZ can be obtained online at fr.com/blast or at ncbi.nlm.nih.gov. Instructions explaining how to use the Bl2 seq program can be found in the readme file accompanying BLASTZ. Bl2seq performs a comparison between two sequences using either the BLASTN or BLASTP algorithm. BLASTN is used to compare nucleic acid sequences, while BLASTP is used to compare amino acid sequences. To compare two nucleic acid sequences, the options are set as follows: -i is set to a file containing the first nucleic acid sequence to be compared (e.g., C:\seq1.txt); -j is set to a file containing the second nucleic acid sequence to be compared (e.g., C:\seq2.txt); -p is set to blastn; -o is set to any desired file name (e.g., C:\output.txt); -q is set to -1; -r is set to 2; and all other options are left at their default setting. For example, the following command can be used to generate an output file containing a comparison between two sequences: C:\Bl2seq -i c:\seq1.txt -j c:\seq2.txt -p blastn -o c:\output.txt -q -1 -r 2. To compare two amino acid sequences, the options of Bl2seq are set as follows: -i is set to a file containing the first amino acid sequence to be compared (e.g., C:\seq1.txt); -j is set to a file containing the second amino acid sequence to be compared (e.g., C:\seq2.txt); -p is set to blastp; -o is set to any desired file name (e.g., C:\output.txt); and all other options are left at their default

setting. For example, the following command can be used to generate an output file containing a comparison between two amino acid sequences: C:\BI2seq -i c:\seq2.txt -j c:\seq2.txt -p blastp -o c:\output.txt. If the two compared sequences share homology, then the designated output file will present those regions of homology as aligned sequences. If the two compared sequences do not share homology, then the designated output file will not present aligned sequences.

[0108] Once aligned, the number of matches is determined by counting the number of positions where an identical nucleotide or amino acid residue is presented in both sequences. The percent sequence identity is determined by dividing the number of matches either by the length of the sequence set forth in the identified sequence (e.g., SEQ ID NO:8), or by an articulated length (e.g., 100 consecutive nucleotides or amino acid residues from a sequence set forth in an identified sequence), followed by multiplying the resulting value by 100. For example, a nucleic acid sequence that has 1200 matches when aligned with the sequence set forth in SEQ ID NO:8 is 83.7 percent identical to the sequence set forth in SEQ ID NO:1 (i.e., $1200 \div 1434 \times 100 = 83.7$). It is noted that the percent sequence identity value is rounded to the nearest tenth. For example, 75.11, 75.12, 75.13, and 75.14 are rounded down to 75.1, while 75.15, 75.16, 75.17, 75.18, and 75.19 are rounded up to 75.2. It also is noted that the length value will always be an integer.

[0109] The embodiments described herein are not limited to a particular *Citrus* or solanaceous plant or variety but rather encompass any *Citrus* or solanaceous plant or hybrid thereof that may be useful in accordance with the disclosure. *Citrus* varieties contemplated by this disclosure include, but are not limited to, cultivated *Citrus* types such as sweet orange, bitter orange, blood orange, grapefruit, pomelo, citron, clementine, naval orange, lemon, lime, mandarin, tangerine, tangelo, or the like.

I. Genome Editing

[0110] Certain aspects of the present disclosure relate to methods of modifying the genome of a *Citrus* or solanaceous plant using genome editing techniques. As used herein, “genome editing” and “genome-engineering” are terms used interchangeably and refer to the modification of a genome through mutagenesis. For example, in plant genome engineering, endonucleases may be used to generate double-strand DNA breaks (DSBs) and activate genome repair pathways. These DSB repair pathways may repair the break cleanly, i.e., without altering the starting sequence, or, alternatively, induce a mutation through an error in repair. In some embodiments, genome editing is used to insert, delete, or substitute one or more base pairs at one or any combination of genetic loci. In some embodiments, a genome editing technique is used to create a mutation, for example, a point mutation or single nucleotide polymorphism.

[0111] In some embodiments the DSB repair pathway is non-homologous end-joining (NHEJ) or microhomology mediated end joining (MMEJ). During NHEJ, any nucleotide overhangs on the break ends are either resected or filled to form blunt ends that are ligated. During MMEJ, the break ends are processed to reveal overhangs comprising microhomology sequences that are then ligated together. The insertions or deletions resulting from the terminal end processing in both the NHEJ and MMEJ pathways can be referred to as indels. In some embodiments, the NHEJ or

MHEJ that occurs can be relied upon to introduce a genome modification including, but not limited to, a silent mutation, a neutral mutation, a missense mutation, a nonsense mutation, or a frameshift mutation.

[0112] In other embodiments, the DSB repair pathway is homologous recombination (HR). During HR, a DSB is repaired using a template with sequences with homology to the DNA flanking the break, i.e., a homologous chromosome. In plant genome editing, a linear DNA polynucleotide flanked by sequences (e.g., of 50 base pairs or more) homologous to those flanking a targeted genomic locus, may be introduced into the genome when a DSB is repaired by HR. In some embodiments, this approach is used to introduce, substitute, or delete a DNA sequence at a genomic locus. Any DNA sequence of interest may be introduced, deleted, or substituted. An introduced or substituted DNA sequence may encode an RNA molecule with a specific activity or function, a DNA molecule with a specific activity or function (e.g., encoding a polypeptide, representing a detectable marker, etc.), a DNA molecule comprising cis-regulatory elements, or a DNA molecule encoding a polypeptide, a motif thereof, or domain thereof. In some embodiments, the nucleic acid encoding the linear DNA sequence that will act as the HR template is encoded by an expression vector. In some embodiments, the nucleic acid encoding the linear DNA sequence of interest is encoded by a DNA sequence separate from the expression vector. For example, and without limitation, the nucleic acid encoding a DNA sequence of interest may be a linear DNA polynucleotide that is co-transformed with an expression vector.

[0113] In some embodiments, single-strand breaks or “nicks” are introduced into the target DNA sequence. As used herein, the term “single-strand break inducing agent” or “nickase” refers to any agent that can induce a single-strand break (SSB) in a DNA molecule. In some embodiments two SSBs are introduced into the target DNA to generate a DSB. These breaks may also be repaired by HR, NHEJ, or MMEJ. In some embodiments, sequence modifications occur at or near the SSB sites, which can include deletions or insertions that result in modification of the nucleic acid sequence, or integration of exogenous nucleic acids by HR or NHEJ.

[0114] In one aspect, a “modification” comprises the insertion of at least 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 15, at least 25, at least 50, at least 100, at least 200, at least 300, at least 400, at least 500, at least 750, at least 1000, at least 1500, at least 2000, at least 3000, at least 4000, at least 5000, or at least 10,000 nucleotides. In another aspect, a “modification” comprises the deletion of at least 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 15, at least 25, at least 50, at least 100, at least 200, at least 300, at least 400, at least 500, at least 750, at least 1000, at least 1500, at least 2000, at least 3000, at least 4000, at least 5000, or at least 10,000 nucleotides. In a further aspect, a “modification” comprises the inversion of at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 15, at least 25, at least 50, at least 100, at least 200, at least 300, at least 400, at least 500, at least 750, at least 1000, at least 1500, at least 2000, at least 3000, at least 4000, at least 5000, or at least 10,000 nucleotides. In still another aspect, a “modification” comprises the substitution of at least 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 15, at least 25, at least

50, at least 100, at least 200, at least 300, at least 400, at least 500, at least 750, at least 1000, at least 1500, at least 2000, at least 3000, at least 4000, at least 5000, or at least 10,000 nucleotides. In some embodiments, a “modification” comprises the substitution of an “A” for a “C,” “G” or “T” in a nucleic acid sequence. In some embodiments, a “modification” comprises the substitution of a “C” for an “A,” “G” or “T” in a nucleic acid sequence. In some embodiments, a “modification” comprises the substitution of a “G” for an “A,” “C” or “T” in a nucleic acid sequence. In some embodiments, a “modification” comprises the substitution of a “T” for an “A,” “C” or “G” in a nucleic acid sequence. In some embodiments, a “modification” comprises the substitution of a “C” for an “U” in a nucleic acid sequence. In some embodiments, a “modification” comprises the substitution of a “G” for an “A” in a nucleic acid sequence. In some embodiments, a “modification” comprises the substitution of an “A” for a “G” in a nucleic acid sequence. In some embodiments, a “modification” comprises the substitution of a “T” for a “C” in a nucleic acid sequence.

[0115] In some embodiments, genome editing of a *Citrus* plant as described herein may encompass techniques that employ methods of targeting endonucleases to one or more genetic loci. In some embodiments, synthetic polypeptides, for example, Transcription Activator-Like Effectors (TALEs) and zinc fingers (ZFs), or nucleic acids, for example, Clustered Regularly Interspaced Short Palindromic Repeats/Cas (CRISPR/CAS) single guide RNAs or NgAgo (Argonaute) single strand DNAs, are used to target endonucleases to any genomic locus. The targeted endonucleases may catalyze a DSB at a target locus. Upon detecting these breaks, a cell may initiate any DSB repair pathway. In some embodiments, genome editing is carried out at more than one genomic locus simultaneously (i.e., multiplex genome engineering). In some embodiments, multiplex genome engineering may be used to remove a sequence of any size from the genome. In some embodiments, any combination and number of endonuclease targeting techniques may be used to target one or more genetic loci.

A. RNA- and DNA-Guided Genome Editing Systems

[0116] In some embodiments, genome engineering of a *Citrus* plant as described herein may employ RNA-guided endonucleases including, but not limited to CRISPR/Cas systems. CRISPR/Cas systems have been described in U.S. Patent Application Publication Nos. 2017/0191082 and 2017/0106025, each of which are incorporated herein by reference in their entirety. In some embodiments, a targeted genome modification as described herein comprises the use of at least one, at least two, at least three, at least four, at least five, at least six, at least seven, at least eight, at least nine, or at least ten RNA-guided nucleases. In some embodiments, a CRISPR/Cas9 system, a CRISPR/Cpf1 system, a CRISPR/CasX system, or a CRISPR/CasY system are alternatives that may be used to generate modifications to target sequences as described herein.

[0117] The CRISPR systems are based on RNA-guided endonucleases that use complementary base pairing to recognize DNA sequences at target sites. CRISPR/Cas systems are part of the adaptive immune system of bacteria and archaea, protecting them against invading DNA, such as viral DNA, by cleaving the foreign DNA in a sequence-dependent manner. The immunity is acquired by the inte-

gration of short fragments of the invading DNA known as spacers between two adjacent repeats at the proximal end of a CRISPR locus. The CRISPR arrays, including the spacers, are transcribed during subsequent encounters with invasive DNA and are processed into small interfering CRISPR RNAs (crRNAs) approximately 40 nt in length, which combine with the trans-activating CRISPR RNA (tracrRNA) to activate and guide the Cas9 nuclease. This cleaves homologous double-stranded DNA sequences known as protospacers in the invading DNA.

[0118] A prerequisite for cleavage is the presence of a conserved protospacer-adjacent motif (PAM) downstream of the target DNA, which usually has the sequence 5'-NGG-3' but less frequently NAG. Specificity is provided by the so-called “seed sequence” approximately 12 bases upstream of the PAM, which must match between the RNA and target DNA. Cpf1 acts in a similar manner to Cas9, but Cpf1 does not require a tracrRNA. Specificity of the CRISPR/Cas system is based on an RNA-guide that use complementary base pairing to recognize target DNA sequences. In some embodiments, the site-specific genome modification enzyme is a CRISPR/Cas system. In an aspect, a site-specific genome modification enzyme provided herein can comprise any RNA-guided Cas endonuclease (non-limiting examples of RNA-guided nucleases include Cas1, Cas1B, Cas2, Cas3, Cas4, Cas5, Cas6, Cas7, Cas8, Cas9 (also known as Csn1 and Csx12), Cas10, Csy1, Csy2, Csy3, Cse1, Cse2, Cse1, Csc2, Csa5, Csn2, Csm2, Csm3, Csm4, Csm5, Csm6, Cmr1, Cmr3, Cmr4, Cmr5, Cmr6, Csb1, Csb2, Csb3, Csx17, Csx14, Csx10, Csx16, CsaX, Csx3, Csx1, Csx15, Csf1, Csf2, Csf3, Csf4, Cpf1, homologs thereof, or modified versions thereof); and, optionally, the guide RNA necessary for targeting the respective nucleases.

[0119] In some embodiments, an RNA-guided endonuclease is the DNA cleavage domain of a restriction enzyme fused to a deactivated Cas9 (dCas9), for example dCas9-FokI. As used herein, a “dCas9” refers to an endonuclease protein with one or more amino acid mutations that result in a Cas9 protein without endonuclease activity, but retaining RNA-guided site-specific DNA binding. As used herein, a “dCas9-restriction enzyme fusion protein” is a dCas9 with a protein fused to the dCas9 in such a manner that the restriction enzyme is catalytically active on the DNA.

[0120] In some embodiments, genome editing of a *Citrus* or solanaceous plant as described herein may employ DNA-guided endonucleases including, but not limited to, NgAgo systems.

[0121] In one aspect, a method and/or composition provided herein comprises one or more, two or more, three or more, four or more, or five or more guide RNAs or DNAs. In another aspect, a CRISPR/CAS system, dCas9-restriction enzyme fusion protein, NgAgo system provided herein is capable of generating a targeted DSB in a target sequence as described herein. In one aspect, vectors comprising nucleic acids encoding one or more, two or more, three or more, four or more, or five or more guide RNAs or DNAs and the corresponding CRISPR/CAS system, dCas9-restriction enzyme fusion protein, NgAgo system are provided to a cell by transformation methods known in the art (e.g., without being limiting, viral transfection, particle bombardment, PEG-mediated protoplast transfection or *Agrobacterium*-mediated transformation).

B. Transcription Activator-Like Effector Nucleases

[0122] In some embodiments, genome editing of a *Citrus* plant as described herein may employ Transcription Activator-Like Effector Nucleases (TALENs). TALENs have been described in U.S. Patent Application Publication Nos. 2016/0369301 and 2015/0203871 (both of which are incorporated herein by reference in their entirety) and are well known in the art. TALENs are artificial restriction enzymes generated by fusing the transcription activator-like effector (TALE) DNA binding domain to an endonuclease domain. In one aspect, the nuclease is selected from a group consisting of PvuII, MthI, TefI and FokI, AlwI, MlyI, SbfI, SdaI, SttI, CleDORF, Clo51, Pcp071. The term TALEN, as used herein, is broad and includes a monomeric TALEN that can cleave double stranded DNA without assistance from another TALEN. The term TALEN is also used to refer to one or both members of a pair of TALENs that work together to cleave DNA at the same site.

[0123] TALEs can be engineered to bind practically any DNA sequence, such as a target sequence as described herein. TALE proteins are DNA-binding domains derived from various plant bacterial pathogens of the genus *Xanthomonas*. The X pathogens secrete TALEs into the host plant cell during infection. The TALE moves to the nucleus, where it recognizes and binds to a specific DNA sequence in the promoter region of a specific DNA sequence in the promoter region of a specific gene in the host genome. TALE has a central DNA-binding domain composed of 13-28 repeat monomers of 33-34 amino acids. The amino acids of each monomer are highly conserved, except for hypervariable amino acid residues at positions 12 and 13. The two variable amino acids are called repeat-variable diresidues (RVDs). The amino acid pairs NI, NG, HD, and NN of RVDs preferentially recognize adenine, thymine, cytosine, and guanine/adenine, respectively, and modulation of RVDs can recognize consecutive DNA bases. This simple relationship between amino acid sequence and DNA recognition has allowed for the engineering of specific DNA binding domains by selecting a combination of repeat segments containing the appropriate RVDs.

[0124] In one aspect, a method and/or composition provided herein comprises one or more, two or more, three or more, four or more, or five or more TALENs. In another aspect, a TALEN provided herein is capable of generating a targeted DSB in a target sequence as described herein. In one aspect, vectors comprising polynucleotides encoding one or more, two or more, three or more, four or more, or five or more TALENs are provided to a cell by transformation methods known in the art (e.g., without being limiting, viral transfection, particle bombardment, PEG-mediated protoplast transfection or *Agrobacterium*-mediated transformation).

C. Zinc Finger Nucleases

[0125] In some embodiments, genome engineering of a *Citrus* or solanaceous plant as described herein may employ Zinc Finger Nucleases (ZFNs). ZFNs have been described in U.S. Pat. No. 9,322,006 (incorporated herein by reference in its entirety) and are well known in the art. ZFNs are synthetic proteins consisting of an engineered zinc finger DNA-binding domain fused to the cleavage domain of an endonuclease, for example, FokI. ZFNs can be designed to cleave almost any long stretch of double-stranded DNA by

the modification of the zinc finger DNA-binding domain. ZFNs form dimers from monomers composed of a non-specific DNA cleavage domain of FokI nuclease fused to a zinc finger array engineered to bind a target DNA sequence. The DNA-binding domain of a ZFN is typically composed of 3-4 zinc-finger arrays. The amino acids at positions -1, +2, +3, and +6 relative to the start of the zinc finger α -helix, which contribute to site-specific binding to the target DNA, can be changed and customized to fit specific target sequences. The other amino acids form the consensus backbone to generate ZFNs with different sequence specificities. Rules for selecting target sequences for ZFNs are known in the art. The FokI nuclease domain requires dimerization to cleave DNA and therefore two ZFNs with their C-terminal regions are needed to bind opposite DNA strands of the cleavage site (separated by 5-7 nt). The ZFN monomer can cut the target site if the two-ZF-binding sites are palindromic. The term ZFN, as used herein, is broad and includes a monomeric ZFN that can cleave double stranded DNA without assistance from another ZFN. The term ZFN is also used to refer to one or both members of a pair of ZFNs that are engineered to work together to cleave DNA at the same site.

[0126] Without being limited by any scientific theory, because the DNA-binding specificities of zinc finger domains can in principle be re-engineered using one of various methods, customized ZFNs can theoretically be constructed to target nearly any gene sequence. Publicly available methods for engineering zinc finger domains include Context-dependent Assembly (CoDA), Oligomerized Pool Engineering (OPEN), and Modular Assembly.

[0127] Several embodiments relate to a method and/or composition provided herein comprising one or more, two or more, three or more, four or more, or five or more ZFNs directed to a target sequence as described herein. In another aspect, a ZFN provided herein is capable of generating a targeted DSB. In one aspect, vectors comprising polynucleotides encoding one or more, two or more, three or more, four or more, or five or more ZFNs are provided to a cell by transformation methods known in the art (e.g., without being limiting, viral transfection, particle bombardment, PEG-mediated protoplast transfection or *Agrobacterium*-mediated transformation).

D. Meganucleases

[0128] In some embodiments, genome engineering of a *Citrus* or solanaceous plant as described herein may employ a meganuclease. Meganucleases, which are commonly identified in microbes, are unique enzymes with high activity and long recognition sequences (>14 nt) resulting in site-specific digestion of target DNA. Engineered versions of naturally occurring meganucleases typically have extended DNA recognition sequences (for example, 14 to 40 nt). The engineering of meganucleases can be more challenging than that of ZFNs and TALENs because the DNA recognition and cleavage functions of meganucleases are intertwined in a single domain. Specialized methods of mutagenesis and high-throughput screening have been used to create novel meganuclease variants that recognize unique sequences and possess improved nuclease activity.

[0129] In one aspect, a method and/or composition provided herein comprises one or more, two or more, three or more, four or more, or five or more meganucleases directed to a target sequence as described herein. In some embodi-

ments, a meganuclease provided herein is capable of generating a targeted DSB. In some embodiments, vectors comprising polynucleotides encoding one or more, two or more, three or more, four or more, or five or more meganucleases are provided to a cell by transformation methods known in the art (e.g., without being limiting, viral transfection, particle bombardment, PEG-mediated protoplast transfection or *Agrobacterium*-mediated transformation).

II. Site-Specific Genome Modification

[0130] Certain aspects of the present disclosure relate to methods of modifying the genome of a *Citrus* plant using site-specific genome modification techniques. In some embodiments, site-specific genome modification of a *Citrus* plant as described herein may employ any site-specific genome modification enzyme. As used herein, the term “site-specific genome modification enzyme” refers to any enzyme that can modify a nucleotide sequence in a sequence-specific manner. In some embodiments, a site-specific genome modification enzyme modifies the genome by inducing a single-strand break. In some embodiments, a site-specific genome modification enzyme modifies the genome by inducing a double-strand break. In some embodiments, a site-specific genome modification enzyme is a recombinase. In some embodiments, a site-specific genome modification enzyme is a transposase. In the present disclosure, site-specific genome modification enzymes include, but are not limited to, nucleases, endonucleases, recombinases, invertases, transposases, methyltransferase, demethylases, aminases, deaminases, helicases, and any combination thereof.

[0131] In some embodiments, the site-specific genome modification enzyme is a recombinase. Non-limiting examples of recombinases include a tyrosine and serine recombinases and coupled with a DNA recognition motifs, for example, a Cre recombinase, a Gin recombinase, a Flp recombinase, and a Tnp1 recombinase. In another aspect, a serine recombinase coupled with a DNA recognition motif, for example, a PhiC31 integrase, an R4 integrase, and a TP-901 integrase. In an aspect, a recombinase is tethered to a zinc-finger DNA-binding domain, or a TALE DNA-binding domain, or a Cas9 nuclease.

[0132] The FLP-FRT site-directed recombination system comes from the 2 μ plasmid from the baker's yeast *Saccharomyces cerevisiae*. In this system, FLP recombinase (flippase) recombines sequences between flippase recognition target (FRT) sites. FRT sites comprise 34 nucleotides. FLP binds to the “arms” of the FRT sites (one arm is in reverse orientation) and cleaves the FRT site at either end of an intervening nucleic acid sequence. After cleavage, FLP recombines nucleic acid sequences between two FRT sites.

[0133] Cre-lox is a site-directed recombination system derived from the bacteriophage P1 that is similar to the FLP-FRT recombination system. Cre-lox can be used to invert a nucleic acid sequence, delete a nucleic acid sequence, or translocate a nucleic acid sequence. In this system, Cre recombinase recombines a pair of lox nucleic acid sequences. Lox sites comprise 34 nucleotides, with the first and last 13 nucleotides (arms) being palindromic. During recombination, Cre recombinase protein binds to two lox sites on different nucleic acids and cleaves at the lox sites. The cleaved nucleic acids are spliced together (reciprocally translocated) and recombination is complete. In

another aspect, a lox site provided herein is a loxP, lox 2272, loxN, lox 511, lox 5171, lox71, lox66, M2, M3, M7, or M11 site.

[0134] In another aspect, the site-specific genome modification enzyme is a dCas9-recombinase fusion protein. As used herein, a “dCas9-recombinase fusion protein” is a dCas9 with a protein fused to the dCas9 in such a manner that the recombinase is catalytically active on the DNA. In some embodiments, dCas9 may be fused with the catalytic domain of any enzyme such that the catalytic domain is catalytically active on DNA. In another aspect, a DNA transposase is attached to a DNA binding domain for example, a TALE-piggyBac and TALE-Mutator.

[0135] Several embodiments relate to promoting DNA recombination by providing a site-specific genome modification enzyme to a plant cell. In some embodiments, recombination is promoted by providing a strand separation inducing reagent. In one aspect, the site-specific genome modification enzyme is selected from an endonuclease, a recombinase, an invertase, a transposase, a helicase or any combination thereof. In some embodiments, recombination occurs between B chromosomes. In some embodiments, recombination occurs between a B chromosome and an A chromosome.

[0136] Several embodiments relate to promoting integration of one or more DNAs of interest by providing a site-specific genome modification enzyme. In some embodiments, integration of one or more DNAs of interest is promoted by providing a strand separation inducing reagent. In one aspect, the site-specific genome modification enzyme is selected from an endonuclease, a recombinase, a transposase, a helicase or any combination thereof. Any DNA sequence can be integrated into a target site of a chromosome sequence by introducing the DNA sequence and the provided site-specific genome modification enzymes. Any method provided herein can utilize any site-specific genome modification enzyme provided herein.

[0137] Several embodiments relate to a method and/or a composition provided herein comprising at least one, at least two, at least three, at least four, at least five, at least six, at least seven, at least eight, at least nine, or at least ten site-specific genome modification enzymes. In yet another aspect, a method and/or a composition provided herein comprises at least one, at least two, at least three, at least four, at least five, at least six, at least seven, at least eight, at least nine, or at least ten polynucleotides encoding at least one, at least two, at least three, at least four, at least five, at least six, at least seven, at least eight, at least nine, or at least ten site-specific genome modification enzymes.

III. Plant Transformation Constructs

[0138] Vectors used for plant transformation may include, for example, plasmids, cosmids, YACs (yeast artificial chromosomes), BACs (bacterial artificial chromosomes) or any other suitable cloning system, as well as fragments of DNA therefrom. Thus when the term “vector” or “expression vector” is used, all of the foregoing types of vectors, as well as nucleic acid sequences isolated therefrom, are included. In some embodiments, a viral vector based on a plant virus such as a *Citrus tristeza* Virus may be used in accordance with the disclosure. It is contemplated that utilization of cloning systems with large insert capacities will allow introduction of large genetic sequences comprising more than one selected gene. In accordance with the disclosure,

this could be used to introduce genetic material corresponding to an entire biosynthetic pathway into a plant. Introduction of such sequences may be facilitated by use of bacterial or yeast artificial chromosomes (BACs or YACs, respectively), or even plant artificial chromosomes. For example, the use of BACs for *Agrobacterium*-mediated transformation was disclosed by Hamilton et al. (1996).

[0139] Particularly useful for transformation are expression cassettes that have been isolated from such vectors. DNA segments used for transforming plant cells will, of course, generally comprise the cDNA, gene or genes which one desires to introduce into and have expressed in the host cells. These DNA segments can further include structures such as promoters, enhancers, polylinkers, or even regulatory genes as desired. The DNA segment or gene chosen for cellular introduction will often encode a protein which will be expressed in the resultant genetically modified cells resulting in a screenable or selectable trait and/or will impart an improved phenotype to the resulting genetically modified plant. However, this may not always be the case, and the present disclosure also encompasses genetically modified plants incorporating non-expressed transgenes.

[0140] In accordance with the disclosure, a nucleic acid vector comprising a coding sequence may be introduced into a plant such as a *Citrus* tree or variety, such that, when the vector is transformed into a *Citrus* variety or plant as described herein, the coding sequence is expressed in the plant. In some embodiments the coding sequence may be expressed in, for example, the phloem or roots of the plant, or any other part of the plant. Expression of the coding sequence in the resulting genetically modified *Citrus* tree or variety results in the tree exhibiting increased tolerance or resistance to HLB when compared to a tree lacking expression of the coding sequence.

A. Proteins and Recombinant DNA Molecules

[0141] As used herein, a “protein/Coding DNA molecule” or “polypeptide/Coding DNA molecule” refers to a DNA molecule comprising a nucleotide sequence that encodes a protein or polypeptide. A “coding sequence” or “protein/Coding sequence” or “polypeptide/Coding sequence” means a DNA sequence that encodes a protein or polypeptide. A “sequence” means a sequential arrangement of nucleotides or amino acids. The boundaries of a protein/Coding sequence or polypeptide/Coding sequence are usually determined by a translation start codon at the 5'-terminus and a translation stop codon at the 3'-terminus. A protein/Coding molecule or polypeptide/Coding molecule may comprise a DNA sequence encoding a protein or polypeptide sequence. As used herein, “transgene expression,” “expressing a transgene,” “protein expression,” “polypeptide expression,” “expressing a protein,” and “expressing a polypeptide” mean the production of a protein or polypeptide through the process of transcribing a DNA molecule into messenger RNA (mRNA) and translating the mRNA into polypeptide chains, which may be ultimately folded into proteins. A protein/Coding DNA molecule or polypeptide/Coding DNA molecule may be operably linked to a heterologous promoter in a DNA construct for use in expressing the protein or polypeptide in a cell transformed with the recombinant DNA molecule. As used herein, “operably linked” means two DNA molecules linked in manner so that one may affect the function of the other. Operably-linked DNA molecules may be part of a single contiguous molecule and may or may not

be adjacent. For example, a promoter is operably linked with a protein/Coding DNA molecule or polypeptide/Coding DNA molecule in a DNA construct where the two DNA molecules are so arranged that the promoter may affect the expression of the transgene.

[0142] As used herein, a “DNA construct” is a recombinant DNA molecule comprising two or more heterologous DNA sequences. DNA constructs are useful for transgene expression and may be comprised in vectors and plasmids. DNA constructs may be used in vectors for the purpose of genome modification, that is the introduction of heterologous DNA into a host cell, in order to produce genetically modified plants and cells, and as such may also be contained in the plastid DNA or genomic DNA of a genetically modified plant, seed, cell, or plant part. As used herein, a “vector” means any recombinant DNA molecule that may be used for the purpose of genetically modifying a plant or plant cell. Recombinant DNA molecules as set forth in the sequence listing, can, for example, be inserted into a vector as part of a construct having the recombinant DNA molecule operably linked to a promoter that functions in a plant to drive expression of the protein encoded by the recombinant DNA molecule. Methods for constructing DNA constructs and vectors are well known in the art. The components for a DNA construct, or a vector comprising a DNA construct, generally include, but are not limited to, one or more of the following: a suitable promoter for the expression of an operably linked DNA, an operably linked protein/Coding DNA molecule, and a 3' untranslated region (3'-UTR). Promoters useful in practicing the present disclosure include those that function in a plant for expression of an operably linked polynucleotide. Such promoters are varied and well known in the art and include those that are inducible, viral, synthetic, constitutive, temporally regulated, spatially regulated, and/or spatio-temporally regulated. Additional optional components include, but are not limited to, one or more of the following elements: 5'-UTR, enhancer, leader, cis-acting element, intron, chloroplast transit peptides (CTP), and one or more selectable marker transgenes.

[0143] Recombinant DNA molecules of the present disclosure may be synthesized and modified by methods known in the art, either completely or in part, especially where it is desirable to provide sequences useful for DNA manipulation (such as restriction enzyme recognition sites or recombination-based cloning sites), plant-preferred sequences (such as plant/Codon usage or Kozak consensus sequences), or sequences useful for DNA construct design (such as spacer or linker sequences). The present disclosure includes recombinant DNA molecules and proteins having at least about 80% (percent) sequence identity, about 81% sequence identity, about 82% sequence identity, about 83% sequence identity, about 84% sequence identity, about 85% sequence identity, about 86% sequence identity, about 87% sequence identity, about 88% sequence identity, about 89% sequence identity, about 90% sequence identity, about 91% sequence identity, about 92% sequence identity, about 93% sequence identity, about 94% sequence identity, about 95% sequence identity, about 96% sequence identity, about 97% sequence identity, about 98% sequence identity, about 99% sequence identity, or about 100% sequence identity to a coding sequence provided herein, for instance the sequences set forth as SEQ ID NOs: 2-9-11-51. As used herein, the term “percent sequence identity” or “% sequence identity” refers to the percentage of identical nucleotides or amino acids in

a linear polynucleotide or polypeptide sequence of a reference (“query”) sequence (or its complementary strand) as compared to a test (“subject”) sequence (or its complementary strand) when the two sequences are optimally aligned (with appropriate nucleotide or amino acid insertions, deletions, or gaps totaling less than 20 percent of the reference sequence over the window of comparison). Optimal alignment of sequences for aligning a comparison window are well known to those skilled in the art and may be conducted by tools such as the local homology algorithm of Smith and Waterman, the homology alignment algorithm of Needleman and Wunsch, the search for similarity method of Pearson and Lipman, and by computerized implementations of these algorithms such as GAP, BESTFIT, FASTA, and TFASTA available as part of the Sequence Analysis software package of the GCG® Wisconsin Package® (Accelrys Inc., San Diego, CA), MEGAlign (DNASar, Inc., Madison, WI), and MUSCLE (version 3.6) (Edgar, *Nucl. Acids Res.* 32:1792-1797, 2004) with default parameters. An “identity fraction” for aligned segments of a test sequence and a reference sequence is the number of identical components which are shared by the two aligned sequences divided by the total number of components in the reference sequence segment, that is, the entire reference sequence or a smaller defined part of the reference sequence. Percent sequence identity is represented as the identity fraction multiplied by 100. The comparison of one or more sequences may be to a full-length sequence or a portion thereof, or to a longer sequence.

[0144] Proteins in accordance with the disclosure may be produced by changing (that is, modifying) a wild-type protein to produce a new protein with a novel combination of useful protein characteristics, such as altered Vmax, Km, substrate specificity, substrate selectivity, and protein stability. Modifications may be made at specific amino acid positions in a protein and may be a substitution of the amino acid found at that position in nature (that is, in the wild-type protein) with a different amino acid. Proteins provided by the disclosure thus provide a new protein with one or more altered protein characteristics relative to the wild-type protein found in nature. In one embodiment of the disclosure, a protein may have altered protein characteristics such as improved or decreased activity against one or more herbicides or improved protein stability as compared to a similar wild-type protein, or any combination of such characteristics. In one embodiment, the disclosure provides a protein, and the DNA molecule or coding sequence encoding it, having at least about 80% sequence identity, about 81% sequence identity, about 82% sequence identity, about 83% sequence identity, about 84% sequence identity, about 85% sequence identity, about 86% sequence identity, about 87% sequence identity, about 88% sequence identity, about 89% sequence identity, about 90% sequence identity, about 91% sequence identity, about 92% sequence identity, about 93% sequence identity, about 94% sequence identity, about 95% sequence identity, about 96% sequence identity, about 97% sequence identity, about 98% sequence identity, about 99% sequence identity, or about 100% sequence identity to a protein sequence such as set forth as SEQ ID NOs: 2-9 and SEQ ID NO:11-51. Amino acid mutations may be made as a single amino acid substitution in the protein or in combination with one or more other mutation(s), such as one or more other amino acid substitution(s), deletions, or addi-

tions. Mutations may be made as described herein or by any other method known to those of skill in the art

B. Regulatory Elements

[0145] The plants and methods of the present disclosure can utilize a vector comprising a coding sequence that, when the vector is transfected into a plant, the coding sequence is expressed in the plant. The site and conditions under which the first selected DNA is expressed can be controlled to a great extent by selecting a promoter element in the vector that causes expression under the desired conditions.

[0146] In some embodiments, the coding sequence is expressed primarily in the roots of the plant, or in the phloem tissue of the plant. In this case, the coding sequence may be expressed in a greater quantity in roots or phloem than in other tissues of the plant. In some embodiments, more than one copy of an coding sequence may be expressed in a plant such that expression in the roots or phloem may be at least twice as much as in any other individual plant tissue (e.g., leaves, flowers, etc).

[0147] Limiting expression of the coding sequence primarily to the roots or phloem of a plant may be accomplished by operably linking the coding sequence to a heterologous promoter active in plant tissues, such as a root-specific or phloem-specific promoter. In other embodiments, a constitutive promoter may be preferred such that the coding sequence is expressed in all tissues of the plant. In some embodiments, a phloem-specific promoter in accordance with the disclosure may comprise an *Arabidopsis* sucrose-proton symporter 2 (AtSUC2) promoter, or a constitutive promoter may comprise a CaMV 35S promoter. Any root-specific or phloem-specific promoter known in the art may potentially be utilized to direct expression of the coding sequence to the roots or the phloem tissue. Examples of these may include, but are not limited to, an RB7, RPE15, RPE14, RPE19, RPE29, RPE60, RPE2, RPE39, RPE61, SHR, ELG3, EXP7, EXP18 or Atlg73160 promoter (Vijaybhaskar et al., 2008; Kurata et al., 2005; PCT Publication WO 01/53502; U.S. Pat. No. 5,459,252; Cho and Cosgrove, 2002).

[0148] In some embodiments, a coding sequence as described herein may be expressed at any level in the plant such that it may be detected in the plant using techniques known in the art. A coding sequence may be expressed in a greater quantity in a genetically modified *Citrus* plant or variety than in a plant not expressing the coding sequence as described herein. In some embodiments, the coding sequence is expressed at least twice as much as in a plant not expressing a coding sequence. In further embodiments, the coding sequence is expressed at least three, or four, or five times, or more, as much as in a plant not expressing a coding sequence. In yet another embodiment, there is no detectable expression of the coding sequence in a plant not expressing a coding sequence.

[0149] The DNA sequence between the transcription initiation site and the start of the coding sequence, i.e., the untranslated leader sequence, can also influence gene expression. One may thus wish to employ a particular leader sequence with a transformation construct of the disclosure. Useful leader sequences are contemplated to include those which comprise sequences predicted to direct optimum expression of the attached gene, i.e., to include a consensus leader sequence which may increase or maintain mRNA stability and prevent inappropriate initiation of translation.

The choice of such sequences will be known to those of skill in the art in light of the present disclosure.

[0150] It is contemplated that vectors for use in accordance with the present disclosure may be constructed to include an ocs enhancer element. This element was first identified as a 16-bp palindromic enhancer from the octopine synthase (ocs) gene of *Agrobacterium* (Ellis et al., 1987), and is present in at least 10 other promoters (Bouchez et al., 1989). The use of an enhancer element, such as the ocs element and particularly multiple copies of the element, may act to increase the level of transcription from adjacent promoters when applied in the context of plant transformation.

C. Terminators

[0151] Transformation constructs prepared in accordance with the disclosure will typically include a 3' end DNA sequence that acts as a signal to terminate transcription and allow for the poly-adenylation of the mRNA produced by coding sequences operably linked to a promoter. In one embodiment of the disclosure, the native terminator of a coding sequence coding sequence may be used. Alternatively, a heterologous 3' end may enhance the expression of coding sequences. Examples of terminators that are deemed to be useful in this context include those from the nopaline synthase gene of *Agrobacterium tumefaciens* (nos 3' end) (Bevan et al., 1983), the terminator for the T7 transcript from the octopine synthase gene of *Agrobacterium tumefaciens*, and the 3' end of the protease inhibitor I or II genes from potato or tomato. Regulatory elements such as an Adh intron (Callis et al., 1987), sucrose synthase intron (Vasil et al., 1989) or TMV omega element (Gallie et al., 1989), may further be included where desired.

D. Transit or Signal Peptides

[0152] Sequences that are joined to the coding sequence of an expressed gene, which are removed post-translationally from the initial translation product and which facilitate the transport of the protein into or through intracellular or extracellular membranes, are termed transit (usually into vacuoles, vesicles, plastids and other intracellular organelles) and signal sequences (usually to the endoplasmic reticulum, Golgi apparatus, and outside of the cellular membrane). By facilitating the transport of the protein into compartments inside and outside the cell, these sequences may increase the accumulation of gene product protecting them from proteolytic degradation. These sequences also allow for additional mRNA sequences from highly expressed genes to be attached to the coding sequence of the genes. Since mRNA being translated by ribosomes is more stable than naked mRNA, the presence of translatable mRNA in front of the gene may increase the overall stability of the mRNA transcript from the gene and thereby increase synthesis of the gene product. Since transit and signal sequences are usually post-translationally removed from the initial translation product, the use of these sequences allows for the addition of extra translated sequences that may not appear on the final polypeptide. It further is contemplated that targeting of certain proteins may be desirable in order to enhance the stability of the protein (U.S. Pat. No. 5,545,818, incorporated herein by reference in its entirety).

[0153] Additionally, vectors may be constructed and employed in the intracellular targeting of a specific gene

product within the cells of a genetically modified plant or in directing a protein to the extracellular environment. This generally will be achieved by joining a DNA sequence encoding a transit or signal peptide sequence to the coding sequence of a particular gene. The resultant transit, or signal, peptide will transport the protein to a particular intracellular, or extracellular destination, respectively, and will then be post-translationally removed.

E. Marker Genes

[0154] By employing a selectable or screenable marker protein, one can provide or enhance the ability to identify transformants. "Marker genes" are genes that impart a distinct phenotype to cells expressing the marker protein and thus allow such transformed cells to be distinguished from cells that do not have the marker. Such genes may encode either a selectable or screenable marker, depending on whether the marker confers a trait which one can "select" for by chemical means, i.e., through the use of a selective agent (e.g., a herbicide, antibiotic, or the like), or whether it is simply a trait that one can identify through observation or testing, i.e., by "screening" (e.g., the green fluorescent protein). Many examples of suitable marker proteins are known to the art and can be employed in the practice of the disclosure. Examples include, but not limited to, neo (Potrykus et al., 1985), bar (Hinchey et al., 1988), bxn (Stalker et al., 1988); a mutant acetolactate synthase (ALS) (European Patent Application 154, 204, 1985) a methotrexate resistant DHFR (Thillet et al., 1988), β -glucuronidase (GUS); R-locus (Dellaporta et al., 1988), β -lactamase (Sutcliffe, 1978), xy/E (Zukowsky et al., 1983), α -amylase (Ikuta et al., 1990), tyrosinase (Katz et al., 1983), β -galactosidase, luciferase (lux) (Ow et al., 1986), acquirin (Prasher et al., 1985), and green fluorescent protein (Sheen et al., 1995; Haseloff et al., 1997; Reichel et al., 1996; Tian et al., 1997; WO 97/41228).

[0155] Included within the terms "selectable" or "screenable" markers also are genes which encode a "secretable marker" whose secretion can be detected as a means of identifying or selecting for genetically modified cells. Examples include markers which are secretable antigens that can be identified by antibody interaction, or even secretable enzymes which can be detected by their catalytic activity. Secretable proteins fall into a number of classes, including small, diffusible proteins detectable, e.g., by ELISA; small active enzymes detectable in extracellular solution (e.g., α -amylase, β -lactamase, phosphinothricin acetyltransferase); and proteins that are inserted or trapped in the cell wall (e.g., proteins that include a leader sequence such as that found in the expression unit of extensin or tobacco PR-S).

IV. Antisense and RNAi Constructs

[0156] In the methods and compositions of the present disclosure, endogenous gene activity can be down-regulated by any means known in the art, including through the use of ribozymes or aptamers. Endogenous gene activity can also be down-regulated with an antisense or RNAi molecule.

[0157] In particular, constructs comprising a coding sequence, including fragments thereof, in antisense orientation, or combinations of sense and antisense orientation, may be used to decrease or effectively eliminate the expression of the gene in a plant such as a *Citrus* tree or variety.

Accordingly, this may be used to “knock-out” the function of the coding sequence or homologous sequences thereof.

[0158] Techniques for RNAi are well known in the art and are described in, for example, Lehner et al., (2004) and Downward (2004). The technique is based on the ability of double stranded RNA to direct the degradation of messenger RNA with sequence complementary to one or the other strand (Fire et al., 1998). Therefore, by expression of a particular coding sequence in sense and antisense orientation, either as a fragment or longer portion of the corresponding coding sequence, the expression of that coding sequence can be down-regulated.

[0159] Antisense, and in some aspects RNAi, methodology takes advantage of the fact that nucleic acids tend to pair with “complementary” sequences. By complementary, it is meant that polynucleotides are those which are capable of base-pairing according to the standard Watson/Crick complementarity rules. That is, the larger purines will base pair with the smaller pyrimidines to form combinations of guanine paired with cytosine (G:C) and adenine paired with either thymine (A:T) in the case of DNA, or adenine paired with uracil (A:U) in the case of RNA. Inclusion of less common bases such as inosine, 5-methylcytosine, 6-methyladenine, hypoxanthine and others in hybridizing sequences does not interfere with pairing.

[0160] Targeting double-stranded (ds) DNA with polynucleotides leads to triple-helix formation; targeting RNA will lead to double-helix formation. Antisense oligonucleotides, when introduced into a target cell, specifically bind to their target polynucleotide and interfere with transcription, RNA processing, transport, translation and/or stability. Antisense and RNAi constructs, or DNA encoding such RNA's, may be employed to inhibit gene transcription or translation or both within a host cell, either in vitro or in vivo, such as within a host plant cell. In certain embodiments of the disclosure, such an oligonucleotide may comprise any unique portion of a nucleic acid sequence provided herein. In certain embodiments of the disclosure, such a sequence comprises at least 18, 30, 50, 75, or 100 or more contiguous nucleic acids of the nucleic acid sequence of a gene, and/or complements thereof, which may be in sense and/or antisense orientation. By including sequences in both sense and antisense orientation, increased suppression of the corresponding coding sequence may be achieved.

[0161] Constructs may be designed that are complementary to all or part of the promoter and other control regions, exons, introns or even exon-intron boundaries of a gene. It is contemplated that the most effective constructs may include regions complementary to intron/exon splice junctions. Thus, it is proposed that an embodiment includes a construct with complementarity to regions within 50-200 bases of an intron-exon splice junction. It has been observed that some exon sequences can be included in the construct without seriously affecting the target selectivity thereof. The amount of exonic material included will vary depending on the particular exon and intron sequences used. One can readily test whether too much exon DNA is included simply by testing the constructs in vitro to determine whether normal cellular function is affected or whether the expression of related genes having complementary sequences is affected.

[0162] As stated above, “complementary” or “antisense” means polynucleotide sequences that are substantially complementary over their entire length and have very few

base mismatches. For example, sequences of fifteen bases in length may be termed complementary when they have complementary nucleotides at thirteen or fourteen positions. Naturally, sequences which are completely complementary will be sequences which are entirely complementary throughout their entire length and have no base mismatches. Other sequences with lower degrees of homology also are contemplated. For example, an RNAi or antisense construct which has limited regions of high homology, but also contains a non-homologous region (e.g., as in a ribozyme) could be designed. Methods for selection and design of sequences that generate RNAi are well known in the art (e.g. Reynolds, 2004). These molecules, though having less than 50% homology, would bind to target sequences under appropriate conditions.

[0163] It may be advantageous to combine portions of genomic DNA with cDNA or synthetic sequences to generate specific constructs. For example, where an intron is desired in the ultimate construct, a genomic clone will need to be used. The cDNA or a synthesized polynucleotide may provide more convenient restriction sites for the remaining portion of the construct and, therefore, would be used for the rest of the sequence. Constructs useful for generating RNAi may also comprise concatemers of sub-sequences that display gene regulating activity.

V. Methods for Genetic Transformation

[0164] Suitable methods for transformation of plant or other cells for use with the current disclosure are believed to include virtually any method by which DNA can be introduced into a cell, such as by direct delivery of DNA such as by PEG-mediated transformation of protoplasts (Omirulch et al., 1993), by desiccation/inhibition-mediated DNA uptake (Potrykus et al., 1985), by electroporation (U.S. Pat. No. 5,384,253, specifically incorporated herein by reference in its entirety), by *Agrobacterium*-mediated transformation (U.S. Pat. Nos. 5,591,616 and 5,563,055; both specifically incorporated herein by reference) and by acceleration of DNA coated particles (U.S. Pat. Nos. 5,550,318; 5,538,877; and 5,538,880; each specifically incorporated herein by reference in its entirety), etc. Through the application of techniques such as these, the cells of virtually any plant species may be stably transformed, and these cells developed into genetically modified plants.

[0165] *Agrobacterium*-mediated transfer is a widely applicable system for introducing genes into plant cells because the DNA can be introduced into whole plant tissues, thereby bypassing the need for regeneration of an intact plant from a protoplast. The use of *Agrobacterium*-mediated plant integrating vectors to introduce DNA into plant cells is well known in the art. See, for example, the methods described by Fraley et al., (1985), Rogers et al., (1987) and U.S. Pat. No. 5,563,055, specifically incorporated herein by reference in its entirety.

[0166] Another method for delivering transforming DNA segments to plant cells in accordance with the disclosure is microprojectile bombardment (U.S. Pat. Nos. 5,550,318; 5,538,880; 5,610,042; and PCT Application WO 94/09699; each of which is specifically incorporated herein by reference in its entirety). In this method, particles may be coated with nucleic acids and delivered into cells by a propelling force.

VI. Production and Characterization of Genetically Modified Plants

[0167] After effecting delivery of exogenous DNA to recipient cells, the next steps generally concern first identifying and selecting the transformed cells and from those cells identifying the selecting the genetically modified cells for further culturing and plant regeneration. In order to improve the ability to identify transformed and genetically modified cells, one may desire to employ one or more selectable or screenable marker genes with a transformation vector prepared in accordance with the disclosure. In this case, one would then generally assay the potentially transformed and modified cell population by exposing the cells to a selective agent or agents, or one would screen the cells for the desired marker gene trait.

[0168] It is believed that DNA is introduced into only a small percentage of target cells in any one study. In order to provide an efficient system for identification of those cells that are transformed and predisposed to genetic modification one may employ a means for selecting those cells that are stably transformed. One exemplary embodiment of such a method is to introduce, into the host cell, a marker gene which confers resistance to some normally inhibitory agent, such as an antibiotic or herbicide. Potentially transformed cells then are exposed to the selective agent. In the population of surviving cells will be those cells where, generally, the resistance/Conferring gene has been integrated and expressed at sufficient levels to permit cell survival. Cells may be tested further to confirm stable integration of the exogenous DNA.

[0169] Cells that survive the exposure to the selective agent, or cells that have been scored positive in a screening assay, may then be selected again using a second, distinct selection paradigm that detects those cells that contain the genetic modification. Cells that survive the exposure to the second selective agent, or cells that have been scored positive in the second screening assay, may be cultured in media that supports regeneration of plants. The genetically modified cells, identified by selection or screening and cultured in an appropriate medium that supports regeneration, will then be allowed to mature into plants. Developing plantlets are transferred to soilless plant growth mix, and hardened, e.g., in an environmentally controlled chamber, for example, at about 85% relative humidity, 600 ppm CO₂, and 25-250 microeinsteins m⁻²s⁻¹ of light. Plants may be matured in a growth chamber or greenhouse. Plants can be regenerated from about 6 wk to 10 months after a genetically modified cell is identified, depending on the initial tissue.

[0170] To confirm the presence of the genetic modification in the regenerating plants, a variety of assays may be performed. Such assays include, for example, “molecular biological” assays, such as Southern and northern blotting and polymerase chain reaction (PCR); “biochemical” assays, such as detecting the absence or presence of a protein product, e.g., by immunological means (ELISAs and western blots) or by enzymatic function; plant part assays, such as leaf or root assays; and also, by analyzing the phenotype of the whole regenerated plant. Modification of the host genome and the independent identities of genetically modified plants may be determined using, e.g., Southern hybridization or PCR. Genetic modifications that affect, for example, protein or gene expression may then be evaluated by specifically measuring the expression of those affected

molecules or evaluating the phenotypic changes brought about by their expression change.

VII. Breeding Plants of the Disclosure

[0171] In addition to direct transformation of a particular plant genotype with a construct prepared according to the current disclosure, genetically modified plants may be made by crossing a plant having a selected genetic modification of the disclosure to a second plant lacking the construct. For example, a selected lignin biosynthesis coding sequence can be introduced into a particular plant variety by crossing, without the need for ever directly transforming a plant of that given variety. Therefore, the current disclosure not only encompasses a plant directly modified or regenerated from cells which have been modified in accordance with the current disclosure, but also the progeny of such plants.

[0172] As used herein the term “progeny” denotes the offspring of any generation of a parent plant prepared in accordance with the instant disclosure, wherein the progeny comprises a selected DNA construct. “Crossing” a plant to provide a plant line having one or more added transgenes relative to a starting plant line, as disclosed herein, is defined as the techniques that result in a coding sequence of the disclosure being introduced into a plant line by crossing a starting line with a donor plant line that comprises a first selected DNA of the disclosure. To achieve this in a plant such as a *Citrus* tree one could, for example, perform the following steps:

[0173] (a) plant seeds of the first (starting line) and second (donor plant line that comprises a first selected DNA of the disclosure) parent plants;

[0174] (b) grow the seeds of the first and second parent plants into plants that bear flowers;

[0175] (c) pollinate a flower from the first parent plant with pollen from the second parent plant; and

[0176] (d) harvest seeds produced on the parent plant bearing the fertilized flower.

[0177] Backcrossing is herein defined as the process including the steps of:

[0178] (a) crossing a plant of a first genotype containing a desired gene, DNA sequence or element to a plant of a second genotype lacking the desired gene, DNA sequence or element;

[0179] (b) selecting one or more progeny plant containing the desired gene, DNA sequence or element;

[0180] (c) crossing the progeny plant to a plant of the second genotype; and

[0181] (d) repeating steps (b) and (c) for the purpose of transferring a desired DNA sequence from a plant of a first genotype to a plant of a second genotype.

[0182] Introgression of a DNA element into a plant genotype is defined as the result of the process of backcross conversion. A plant genotype into which a DNA sequence has been introgressed may be referred to as a backcross converted genotype, line, inbred, or hybrid. Similarly a plant genotype lacking the desired DNA sequence may be referred to as an unconverted genotype, line, inbred, or hybrid.

[0183] In some embodiments, asexual reproduction or propagation may be used to obtain a progeny plant in accordance with the disclosure. Techniques to achieve asexual propagation or reproduction in *Citrus* trees or varieties may include, for example, grafting, budding, top-working, layering, runner division, cuttings, rooting, T-budding, and the like. In some embodiments, one *Citrus* variety

into which a coding sequence has been introduced may be grafted onto the rootstock of another variety. In other embodiments, a coding sequence may be introduced into the rootstock. In either of these situations, one or both of the plant varieties may exhibit increased tolerance or resistance to HLB.

EXAMPLES

Example 1

Identification of Las SDEs

[0184] Las has been predicted to produce at least 166 SDEs, 36 of which have been shown to be highly expressed in plants. These 36 Las SDEs were stably or transiently expressed in *Citrus* plants and *Nicotiana benthamiana* to determine if they are HLB effectors. Transgenic expression of CLIBASIA_04025 (Las4025), CLIBASIA_00470 (Las470), CLIBASIA_04065 (Las4065), CLIBASIA_05150 (Las5150), and CLIBASIA_04250 (Las4250) induced symptoms consistent with HLB. For example, transgenic expression of CLIBASIA_04025 (Las4025) without its signal peptide stunted leaf growth, delayed plant growth, and induced leaf yellowing (FIG. 1A and FIG. 1B). The CLIBASIA_04025 expression in transgenic lines was

confirmed using an antibody against CLIBASIA_04025 (FIG. 1C). Transgenic expression of CLIBASIA_00470 also delayed plant growth.

Example 2

Las SDE and Host Target Proteins Interaction Assays

[0185] Yeast two-hybrid (Y2H) screening was performed to identify putative target proteins of the SDEs identified above. A cDNA library was generated from mRNA isolated from ‘Valencia’ sweet orange plants infected with Las. The mRNA was obtained from these plants during the early stage of infection in which the Las Ct value was between 28-30. The cDNA library was constructed using the Make Your Own Mate & Plate™ Library System (Clontech) following the manufacturer’s instructions and had a titer greater than 3×10^8 cfu.

[0186] The coding sequences of CLIBASIA_04025 (Las4025), CLIBASIA_00470 (Las470), CLIBASIA_04065 (Las4065), CLIBASIA_05150 (Las5150), and CLIBASIA_04250 (Las4250), without their signal peptides, were cloned in-frame with the GAL4 DNA-binding domain (BD) of the bait vector pGBKT7. The Y2H screen was performed using the Matchmaker® Gold Yeast Two-Hybrid System (Clontech) following the manufacturer’s instructions. The SDE target proteins identified in the screen are summarized in Table 1B.

TABLE 1B

Y2H-confirmed Las SDE target proteins.	
SDEs	Target proteins
CLIBASIA_04025 (Las4025)	PP2-B2/12 (orange1.1t04174) Lectin (orange1.1t05126) Cysteine protease (Cs4g07410) Cysteine protease 15A-like (Cs3g25530) Myb family transcription factor (orange1.1t02260) YLS9-like (Cs2g29120) Cell death suppressor protein Lls1 (Cs9g02990) Red chlorophyll catabolite reductase; Accelerated cell death 2 (Cs1g22670) Homolog to Acd2, red chlorophyll catabolite reductase-like (Cs1g22680), which is 64% similar Homolog to Lls1 - Acd1-Like Cs9g03000, which is 93.4% similar; Acd1 Cs8g15480, which is 51% similar, but is also work in same chlorophyll catabolism pathway to Acd2. Homologs to Cysteine protease (Cs3g25530) and Cysteine Protease (Cs4g07410), all Papain-like cysteine proteases Homolog YLS9-like, NDR1/HIN1-like protein 13, Cs8g01640, which is 75% similar Homologs to Myb family transcription factor, PHL5-like, Cs7g01290, 56% similar; PHL5, orange1.1t02259, which is 76% similar Galactinol-sucrose galactosyltransferase 2 (Cs9g12460) Vacuolar protein sorting-associated protein 36 (Cs7g24050) DnaJ protein homolog (Cs7g23510) Plastid-specific ribosomal protein 4 (Cs6g08000) Pathogenesis-related protein 10 (Cs9g03630) Glucan endo-1,3-beta-D-glucosidase-like protein (orange1.1t00643) Core-2/I-branching beta-1,6-N-acetylglucosaminyltransferase family protein (Cs7g07430) Leucyl-tRNA synthetase bacterial/mitochondrial, class Ia (Cs2g02720) Annexin D1 (Cs3g18360) Pentatricopeptide repeat-containing protein (Cs5g26120) Probable fructose-bisphosphate aldolase 2, chloroplastic (Cs8g08710) Arginine/serine-rich splicing factor, putative, expressed (Cs3g18350) SVP1-like protein 2 (Cs5g32770) Alanine aminotransferase 2 (Cs7g09270) BEL1-like homeodomain protein 1 (Cs6g13660) AT-rich interactive domain-containing protein 4 (Cs4g06750)

TABLE 1B-continued

Y2H-confirmed Las SDE target proteins.	
SDEs	Target proteins
CLIBASIA_00470 (Las470)	Heat shock factor protein HSF8 (Cs7g24140)
	Plasma membrane ATPase 1 (Cs6g03480)
	Gag-pol polyprotein (Cs7g14770)
	Phospholipid: diacylglycerol acyltransferase (Cs1g17750)
	Aconitate hydratase, cytoplasmic (Cs2g21430)
	DNA-directed RNA polymerase subunit alpha (orange1.1t03665)
	Polyubiquitin 10 (Cs4g11190)
	Diacylglycerol kinase theta (Cs4g02800)
	Chloroplast methionine sulfoxide reductase B2 (Cs9g05400)
	Leucine-rich repeat receptor protein kinase EXS (Cs7g18050)
	Lateral organ boundaries-domain 29 (orange1.1t00246.1)
	Formamidase (Cs1g21820)
	Signal peptidase complex subunit 3B (Cs3g13460)
	Probable plastid-lipid-associated protein 8 (Cs7g07440)
	Stress responsive gene 6 protein (orange1.1t01091)
	UBX domain-containing protein (Cs5g01690)
	Protein SRG1 (Cs5g13180)
	Thioredoxin H-type (Cs1g24740)
	Glycoside hydrolase (Cs8g12020)
	Thioredoxin F2 (Cs6g02830)
	RNA pseudourine synthase 7 (orange1.1t02625)
	Kunitz-type protease inhibitor KPI-D2.2 (Cs5g16850)
	Aspartate aminotransferase (Cs4g19830)
	UDP-glucuronate decarboxylase 4 (Cs6g05450)
	Nitrate transporter 1.5 (orange1.1t00223)
	FKBP-like peptidyl-prolyl cis-trans isomerase family protein (orange1.1t00062)
	Zinc-binding alcohol dehydrogenase domain-containing protein 2 (Cs8g05790)
	Mannitol dehydrogenase (Cs1g20600)
	RNA recognition motif family protein (Cs2g07940)
	alpha/beta-Hydrolases superfamily protein (Cs2g21120)
	Protein argonaute 1 (Cs5g16710)
	Progesterone 5-beta-reductase (Cs3g11840)
	NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 6 (Cs1g16240)
	Tetratricopeptide repeat (TPR)-like superfamily protein (Cs6g03690)
	Mitochondrial carrier domain-containing protein (Cs6g03800)
	Cell division control protein 48 homolog C (Cs3g01650)
	Subtilisin-like protease (Cs8g02780)
	U6 snRNA-associated Sm-like protein LSm6 (orange1.1t02120)
	Subtilisin-like protease (Cs8g06090)
	Histone H4 (Cs8g18120)
	Chromatin-associated protein Dek (Cs4g08790)
	Serine carboxypeptidase-like 49 (Cs7g24460)
	RING/U-box superfamily protein (Cs8g16720)
	Linoleate 13S-lipoxygenase 2-1 (orange1.1t04376)
	Isoflavone reductase-like (Cs2g16260)
	alpha-like protein (Cs6g16290.1)
	THO complex subunit 3 (Cs7g18110)
	Uncharacterized protein Sb07g024435 (Cs7g26460)
	Ubiquitin-activating enzyme E1 2 (Cs8g20660)
	Structure-specific endonuclease subunit SLX1 (Cs2g09350)
	Putative uncharacterized protein P0458H05.117 (Cs4g05300)
	Aconitate hydratase 1 (Cs1g26040)
	DnaJ homolog subfamily B member 13 (Cs3g04780)
	Plastocyanin (Cs3g26730)
	Putative uncharacterized protein Sb03g000880 (Cs5g31880)
	50S ribosomal protein L14 (orange1.1t04817)
	Oligopeptidase A (Cs1g20720)
	Maturase K (Cs2g09070)
	Protein FRA10AC1 (Cs7g01730)
	Putative Uncharacterized protein AINc14C124G6763 (Cs8g09030)
	Putative Uncharacterized protein Sb10g020525 (orange1.1t00482)
	Putative Uncharacterized protein OSJNBb0021O11.27 (Cs4g16820)
	Lectin (orange1.1t05126)
	Galactinol-sucrose galactosyltransferase 2-like (Cs9g12460)
	DnaJ homolog 1 like (Cs1g19720)
	YLS9-like (Cs2g29120)
	8-hydroxygeraniol dehydrogenase (XM_006466284.2)

TABLE 1B-continued

Y2H-confirmed Las SDE target proteins.	
SDEs	Target proteins
CLIBASIA_04065 (Las4065)	Hypothetical protein (orange1.1t00563)
CLIBASIA_05150 (Las5150)	Cysteine protease (Cs4g07410)
CLIBASIA_04250 (Las4250)	Translation initiation factor IF-3 (orange1.1g044576m)

[0187] Multiple proteins interacted with CLIBASIA_04025 (Las4025) and CLIBASIA_00470 (Las470); whereas, CLIBASIA_04065 (Las4065), CLIBASIA_05150 (Las5150), and CLIBASIA_04250 (Las4250) each interacted with a single protein. CLIBASIA_04025 (Las4025) interacted with PP2-B2 (orange1.1t04174), a phloem protein. Phloem protein encoding genes are known to be involved in phloem blockage and are suggested to contribute to HLB symptom onset. In addition, SDE15 was shown to interact with the CtACD2 protein by the Y2H assay (FIG. 5A).

[0188] To confirm the initial Y2H screen results, full-length sequences of the SDE target proteins were cloned in-frame with the GALA activation domain (AD) of the prey vector pGADT7. Following the manufacturer's instructions, the Y2HGOLD yeast strain was co-transformed with relevant bait and prey vector pairs. For negative controls, a prey vector co-transformed with an empty bait vector was used. Exemplary results that were achieved in the Y2H assay using the CLIBASIA_04025 (Las4025) bait vector are shown in FIG. 2. CLIBASIA_04025 (Las4025) interacted with PP2-B2 (orange1.1t04174), and Pathogenesis-related protein 10 (Cs9g03630).

[0189] Glutathione S-transferase (GST) pull-down and Bimolecular fluorescence complementation (BiFC) assays were performed to further confirm Y2H results. Las genomic DNA was extracted from infected *Citrus* leaves and the coding sequences of CLIBASIA_04025 (Las4025), CLIBASIA_00470 (Las470), CLIBASIA_04065 (Las4065), CLIBASIA_05150 (Las5150), and CLIBASIA_04250 (Las4250) were PCR-amplified for use in both assays. For the GST pull-down assay, the respective fragments were cloned in-frame with Maltose-binding protein (MBP) in the pMAL™/C5X vector (NEB, USA) to generate MBP-SDE fusion proteins. The coding sequences of the SDE target proteins were PCR-amplified using *Citrus* leaf cDNA as a template. The respective fragments were cloned in-frame with GST in the pGEX-4T-1 vector (GE Healthcare, USA) to generate GST-target fusion proteins. For the BiFC assays, the coding sequences of the SDE target proteins were PCR-amplified using *Citrus* leaf cDNA as a template. The respective SDEs and SDE target proteins were cloned in-frame with either N-terminal or C-terminal fragments of EYFP using pSAT6-nEYFP/C1 and pSAT6/CEYFP/C1-B vectors, respectively. This was done with an In-Fusion cloning kit (Clontech, USA) and produced SDE-EYFPN, EYFPC-SDE, SDE target-EYFPN, and EYFPC-SDE target fusion proteins. *Citrus* protoplasts isolated from grapefruit epicotyl segments were co-transformed with pairs of EYFPC and EYFPN vectors. Additionally, SDE15 was shown to interact with the CtACD2 protein by the GST pull-down assay (FIG. 5C) and the BiFC assay (FIG. 5B).

[0190] Exemplary results obtained in the GST pull-down and BiFC assays are provided in FIG. 3. In both assays, the CLIBASIA_04025 (Las4025) directly interacted with cysteine protease (Cs4g07410), therefore confirming the interaction observed between the two proteins in the Y2H assays. No interactions were detected in any of the negative controls (FIG. 3A and FIG. 3B).

[0191] SDE15 was also shown to interact with the CtACD2 protein that negatively regulates the hypersensitive reaction by the hypersensitive response (HR) assay (FIG. 5D). The electrolyte leakage associated with HR induced by AvrBsT protein is shown in FIG. 5E.

Example 3

Overexpression of SDE Target Proteins in Citrus Plants

[0192] Overexpression of Myb family transcription factor (orange1.1t02260), a CLIBASIA_04025 (Las4025) target protein, in Duncan grapefruits induced symptoms similar to those observed in *Citrus* plants infected with *Candidatus Liberibacter asiaticus*. Plants that overexpressed MYTL displayed stunted plant growth and greater starch accumulation when compared to wild type controls (FIG. 4A, FIG. 4B, and FIG. 4C).

Example 4

Modification of Susceptibility Genes in Citrus Plants

[0193] Based on the identification of susceptibility genes that encode proteins that are targets of SDEs secreted by Las, *Citrus* plants and regulatory elements thereof may be modified to exhibit resistance to infection by Las. In particular, plants may be modified to exhibit reduced expression of one or more of the susceptibility genes identified herein. These susceptibility genes and regulatory elements thereof may themselves be modified using targeted genome editing (e.g., single-stranded oligonucleotide-mediated genome editing, meganucleases, zinc finger nucleases, TALENs, or RNA-guided nuclease (e.g., CRISPR technology). Modified *Citrus* plants may be produced using a plant transformation vector comprising an expression cassette comprising a promoter and 3' UTR operably linked to the coding sequence for a modified target protein. The plant transformation vector may be used with *Agrobacterium tumefaciens* and standard methods for plant transformation known in the art.

[0194] Alternatively, plants comprising modified endogenous genes or regulatory elements thereof may be produced by inserting DNA directly into the plant genome at a specified targeted location. Any site or locus within the plant

genome may potentially be chosen for site-directed integration of a transgene or construct of the present disclosure. For site-directed integration, a double-strand break (DSB) or nick may first be made at a selected genomic locus with a site-specific nuclease, such as, for example, a zinc-finger nuclease, an engineered or native meganuclease, a TALE-endonuclease, or an RNA-guided endonuclease (e.g., Cas9, Cpf1, CasX or CasY). Any method known in the art for site-directed integration may be used. In the presence of a donor template molecule, the DSB or nick may then be repaired by homologous recombination between the homology arm(s) of the donor template and the plant genome, or by non-homologous end joining (NHEJ), resulting in site-directed integration of the insertion sequence into the plant genome to create the targeted insertion event at the site of the DSB or nick.

[0195] Plants having one or more mutations in the genomic susceptibility genes or regulatory elements thereof may be produced using a double-strand break (DSB) or nick made at the genomic locus with a site-specific nuclease, such as, a zinc-finger nuclease, an engineered or native meganuclease, a TALE-endonuclease, or an RNA-guided endonuclease (e.g., Cas9, Cpf1, CasX or CasY). Any method known in the art for genome editing or non-templated editing may be used. Delivery methods for the nuclease and gRNA include, but are not limited to, delivering by *Agrobacterium*-mediated methods, delivering as a protein or RNA using transfection or biolistics, and delivery by expression from a virus. One or more nucleases or gRNA may be used. Donor molecules to deliver the desired changes may include, but are not limited to, double-stranded DNA, single-stranded DNA oligonucleotides, RNA or viral DNA. Donor molecules may be delivered by *Agrobacterium*, virus, biolistic delivery, or transfection. In the presence of a donor template molecule, the one or more DSBs or nicks may then be repaired by homologous recombination between the homology arm(s) of the donor template and the plant genome, by non-homologous end joining (NHEJ), by single-strand annealing pathway or other DNA repair mechanisms resulting in modification of the native sequence in the plant genome to that contained by the donor to create the desired mutation.

[0196] Modified *Citrus* plants may be tested for resistance to HLB. *Citrus* plants that are modified to exhibit reduced expression of one or more of the susceptibility genes identified herein may be inoculated with a strain from the genus *Ca. Liberibacter* and evaluated for symptoms of HLB infection compared to *Citrus* plants that do not comprise modified susceptibility genes or regulatory elements thereof. Modified plants that test negative for HLB may be used for further greenhouse and field testing and breeding of HLB resistant *Citrus* lines.

Example 5

Overexpression of SDE15 in *Nicotiana tabacum* and *Citrus*

[0197] SDE15 (CLIBASIA_04025, 11.17 kDa) is a hypothetical protein. Sequence analysis of SDE15 did not reveal any insights into its function except the presence of a typical signal peptide (amino acids 1-22) predicted by SignalP v4.1 (FIG. 6A). Interestingly, transgenic expression of SDE15 without the signal peptide (8.68 kDa) in *N. tabacum* caused markedly reduced plant growth and a slightly yellow leaf

phenotype, suggestive of biological effects of SDE15 on host biology. These interesting phenotypes, observed in a heterologous plant (*N. tabacum*), prompted us to overexpress SDE15 in the native host plant *Citrus* (cv. Duncan grapefruit) via stable *Agrobacterium* mediated transformation to determine whether SDE15 is involved in inducing HLB-like symptoms, suppressing plant defenses and/or promoting Las growth. Indeed, SDE15-transgenic Duncan grapefruit plants showed some HLB-like symptoms including slight yellowing, smaller leaves, and reduced growth at the early stage of development (FIG. 6B). Interestingly, these HLB-like symptoms disappeared approximately one year after transgenic plants were grown under greenhouse conditions.

[0198] Las-infected plants show characteristic gene expression patterns (Albrecht, U., and Bowman, K. (2008). *PLANT SCIENCE* 175, 291-306; Kim, J. S., Sagaram, U. S., Burns, J. K., Li, J. L., and Wang, N. (2009). *Phytopathology* 99, 50-57; Albrecht, U., and Bowman, K. D. (2012). *Plant Sci* 185-186, 118-130; Rawat, N., Kiran, S. P., Du, D., Gmitter, F. G., and Deng, Z. (2015). *BMC Plant Biol* 15, 184). To determine whether SDE15 also modulates host gene expression, we compared the expression patterns of HLB-associated marker genes in SDE15-transgenic and EV (empty vector)-transgenic *Citrus* plants. We found that expression of the genes encoding phloem protein 2 (PP2)-B15, WRKY40, and Zn transporter 5 (ZIP5) was highly induced in SDE15-transgenic *Citrus*, whereas expression of the genes encoding sugar transporter protein Sweet7 and Sweet15 and chalcone synthase were reduced compared to control, resembling those in HLB plants for these marker genes (Albrecht and Bowman, 2008; Kim et al., 2009; Albrecht and Bowman, 2012; Rawat et al., 2015). Taken together, both the partial HLB-like symptom, albeit transient, and the expression pattern of HLB marker genes in the SDE15-transgenic *Citrus* indicate that SDE15 is involved in the outcome of the Las-*Citrus* interaction.

Example 6

Detection of SDE15 Inside Phloem Sap and Subcellular Localization of SDE15

[0199] To function as an effector, we expect that SDE15 would be secreted into the cytosol of phloem cells and may be detected in the phloem sap of Las-infected plants. We therefore developed a SDE15-specific antibody and examined whether SDE15 can be detected in the phloem sap by western blot. We could clearly detect SDE15 in the phloem sap extract of Las-infected barks but not in healthy *Citrus* barks by western blot (FIG. 6C). To determine the subcellular locations of SDE15 in the plant cell, we performed subcellular localization of SDE15 by transient expression of SDE15-EYFP fusion protein in *N. benthamiana*. We found that SDE15 is localized to the cytoplasm and the nucleus (FIG. 6D).

[0200] We next analyzed the expression pattern of the SDE15 transcript in Las-infected *Citrus* and psyllids using quantitative reverse transcription-PCR (qRT-PCR). SDE15 showed a higher expression level in *Citrus* than in psyllids (FIG. 6E). Furthermore, SDE15 expression level was higher in the early stage (asymptomatic) than the later stage (symptomatic) of Las infection (FIG. 6F), indicating that SDE15 likely functions at a relatively early stage of Las infection before the appearance of visible HLB symptoms.

Example 7

SDE15 Suppresses PCD and Promotes Las Growth in Planta

[0201] Many effectors of extracellular pathogens have been reported to suppress plant defenses (Jones, J. D., and Dangl, J. L. (2006). *Nature* 444, 323-329.; Toruño, T. Y., Stergiopoulos, I., and Coaker, G. (2016). *Annu Rev Phytopathol* 54, 419-441). To investigate whether SDE15 (from a cytosolic bacterial pathogen) is involved in suppressing plant defenses, we inoculated SDE15-transgenic and non-transgenic Duncan grapefruit with *Xanthomonas citri* subsp. *citri* strain A^w (XccA^w), which triggers the immunity-associated hypersensitive reaction (HR, a form of PCD) in Duncan grapefruit. Intriguingly, we found XccA^w did not induce an HR in the leaves of SDE15-transgenic Duncan grapefruit, whereas an HR was observed in the nontransgenic Duncan grapefruit (FIG. 7A). We next examined immunity-associated gene expression in SDE15 transgenic and nontransgenic plants after inoculation with XccA^w. The expression of PR1, PR3, and PR5 genes was markedly reduced in the SDE15-transgenic than nontransgenic Duncan grapefruit (FIG. 7B). Taken together, these results suggest that SDE15 is a potent suppressor of PCD in *Citrus*.

[0202] The ability of SDE15 to suppress PCD and defense gene expression prompted us to examine whether the SDE15 transgenic plants could promote Las growth. For this purpose, we graft-inoculated SDE15-transgenic and EV-transgenic Duncan grapefruit plants with Las. Significant increase of Las titers was observed over a four-month period in the SDE15-transgenic *Citrus* compared to the EV-transgenic Duncan grapefruit plants, as estimated by qPCR assays (FIG. 8). In accordance, visible HLB symptoms were observed earlier on leaves of SDE15-transgenic *Citrus* than control plants after Las graft-inoculation (FIG. 8).

[0203] Because SDE15 suppressed PCD triggered by XccA^w, we also tested whether SDE15 transgenic plants become more susceptible to virulent Xcc strain XacA 306. Indeed, the canker symptom caused by XacA 306 appeared faster in SDE15 transgenic *Citrus* compared to EV-transgenic plants. In addition, higher XacA 306 populations were detected in SDE15 transgenic grapefruit than that in the EV-transgenic Duncan grapefruit (FIG. 9). Thus, transgenic expression of SDE15 causes broad susceptibility to both HLB and *Citrus* canker, two of the most important diseases in *Citrus*.

Example 8

SDE15 Interacts With the PCD Repressing Protein CsACD2

[0204] Because SDE15 does not contain obvious sequence domains that suggest enzymatic activities, we hypothesized that SDE15 might exert its suppressive effect on PCD and defense gene expression via protein-protein interactions. To identify the targets of SDE15, we performed yeast two-hybrid (Y2H) screening with SDE15 without its signal peptide using cDNA libraries generated with mRNAs isolated from the Valencia sweet orange at different Las infection stages (healthy (H), asymptomatic (AS) and symptomatic(S)). We identified 20 SDE15-interacting proteins in the H library, 60 in the AS library and 6 in the S library (Table 1B).

[0205] Among the SDE15-interacting proteins identified by Y2H (Table 1B), CsACD2 (Accession No. Cs1g22670) (FIG. 10) was of special interest. It shares 55% similarity with the *Arabidopsis* ACD2 protein (Greenberg et al., 1994; Wüthrich et al., 2000). In *Arabidopsis*, ACD2 protects cells from PCD caused by endogenous porphyrin-related molecules including red chlorophyll catabolite (RCC) or exogenous protoporphyrin IX (Mach, J. M., Castillo, A. R., Hoogstraten, R., and Greenberg, J. T. (2001). *T Proc Natl Acad Sci U S A* 98, 771-776; Yao, N., Eisfelder, B. J., Marvin, J., and Greenberg, J. T. (2004). *Plant J* 40, 596-610). We hypothesized that SDE15 may target CsACD2 to suppress PCD as a mechanism to promote Las growth in the phloem. We confirmed the SDE15-CsACD2 interaction via pair-wise Y2H assay (FIG. 10A). To further confirm the interaction between SDE15 and CsACD2, we conducted bimolecular fluorescence complementation (BiFC) assay in vivo and glutathione-S-transferase (GST) pull-down assay in vitro. Co-transformation of SDE15-EYFPN and EYFPC-CsACD2 or EYFPC-SDE15 and CsACD2-EYFP^N into *Citrus* leaf protoplasts displayed strong signals in the cytoplasm, but not in negative controls (FIG. 10B). GST pull-down assay confirmed direct interaction between GST-SDE15 and MBP-CsACD2 fusion proteins (FIG. 10C). These results demonstrate a genuine interaction between SDE15 and CsACD2.

Example 9

C-Terminal is Essential for SDE15 to Interact With RCCR Domain of CsACD2

[0206] To further define the regions of SDE15 and CsACD2 involved in interaction, truncated proteins GST-SDE15^N (SDE15 protein without the signal peptide and additional 25 aa at the N-terminus), GST-SDE15^C (SDE15 protein without the signal peptide and additional 25 aa at the C-terminus) and MBP-RCCR (the RCCR domain of CsACD2 with deletion of 71 aa at the N-terminus) were expressed in *E. coli*, purified and used in GST pull-down assays (FIG. 10D, E). Results showed that the RCCR domain of CsACD2 was sufficient to interact with GST-SDE15 (FIG. 10D). In addition, deletion of the C-terminal, but not the N-terminal of SDE15 abolished its interaction with CsACD2 (FIG. 10E).

[0207] Next, we examined the subcellular localization of CsACD2 in *Nicotiana benthamiana* leaf cells. CsACD2 was mostly found in the chloroplast and partially in the cytosol and the nucleus (FIG. 11). Co-localization of SDE15-CFP and CsACD2-EYFP could be detected in the cytoplasm and the nucleus (FIG. 11).

Example 10

SDE15 Promotes the RCCR Activity of CsACD2

[0208] It has been shown that ACD2 in *Arabidopsis* represses PCD by functioning as a red chlorophyll catabolite reductase (RCCR) to catalyzing porphyrin-related molecules such as RCC (Pruzinska, A., Tanner, G., Aubry, S., Anders, I., Moser, S., Müller, T., Ongania, K. H., Kräutler, B., Youn, J. Y., Liljegren, S. J., and Hörtensteiner, S. (2005). *Plant Physiol* 139, 52-63). To determine whether CsACD2 is involved in SDE15-mediated suppression of PCD, we infiltrated the leaves of *N. benthamiana* with *Agrobacterium* harboring the binary vector that expresses CsACD2-YFP

fusion protein and/or SDE15, followed by infiltration with *Agrobacterium* containing a binary vector that expresses AvrBsT 2 days after CsACD2 infiltration (FIG. 12A, B). As expected, AvrBsT triggered an HR in the absence of SDE15 but that HR was repressed in the presence of SDE15. On the other hand, we found neither SDE15^{ΔN} nor SDE151^{ΔC} could repress HR induced by AvrBsT. Intriguingly, however, we found that transient overexpression of CsACD2-YFP protein alone was also sufficient to suppress AvrBsT-elicited HR in *N. benthamiana* (FIG. 12A, B), suggesting that (i) SDE15 could interact with the endogenous homolog of ACD2 in *N. benthamiana* (NbACD2 hereinafter) and (ii) it enhances the stability or activity of CsACD2. To test the first possibility, we performed GST pull-down assays. Indeed, SDE15 interacted strongly with NbACD2. As in the case of CsACD2, the NbACD2-YFP fusion protein alone, when transiently expressed, was sufficient to suppress HR induction by AvrBsT. To examine the second hypothesis, we directly performed RCCR activity assay. SDE15 significantly increased the RCCR activity of CsACD2 to catabolize RCC to primary fluorescent catabolite (pFCC), while SDE15 1.N and SDE151C lost the ability to promote the RCCR activity of CsACD2 (FIG. 12C).

Example 11

SDE15 Promotes the Chlorophyll Break-Down in *Planta*

[0209] HLB is associated with tissue chlorosis and vein mottling, both are indicative of chlorophyll break down and previous studies have shown that the levels of chlorophyll a and b were significantly reduced in HLB-diseased *Citrus* trees (Killiny, N., and Nehela, Y. (2017). Mol Plant Microbe Interact 30, 543-556). As ACD2 is critical for alleviating the accumulation of toxic intermediates in the chlorophyll breakdown pathway, we surmise that SDE15 promotes the RCCR activity during Las infection by lowering toxic intermediates that induce PCD. We quantified the concentration of pheophorbide a, which is upstream of RCC in the chlorophyll break-down pathway. The concentration of pheophorbide a was significantly lower in the leaves of SDE15 transgenic *Citrus* than that of the EV transgenic *Citrus* under HLB-free conditions. In addition, the levels of chlorophyll a and chlorophyll b were also lower in SDE15 transgenic *Citrus* than those in the EV transgenic *Citrus*. Furthermore, the concentrations of the three compounds were lower in the leaves of Las-infected *Citrus* (both SDE15 and EV transgenic plants) than those in the healthy EV transgenic plants. Our results suggest that, by promoting the RCCR activity of CsACD2, SDE15 likely prevents accumulation of PCD-eliciting intermediates during the breakdown of chlorophylls. This activity also likely contributes to the development of yellowing symptom associated with HLB and explained the yellowing symptoms observed in the SDE15 transgenic *Citrus* (FIG. 6B).

Example 12

Methods For Examples 5-11

Vectors Construction

[0210] To generate the construct for plant transformation, Las genomic DNA was isolated from HLB diseased *Citrus* leaf by CTAB method. The coding sequence of SDE15 (222

bp) without signal peptide was PCR-amplified using gene-specific primers (Table S2). A BamHI recognition sequence and a KpnI recognition site with two protecting nucleotides were added to the 5' end of primers. The PCR product was purified and cloned into pGEM-T Easy vector (Promega, Madison, WI, USA) and then cloned into the binary vector erGFP-1380N at the BamHI and KpnI sites to generate SDE15-overexpression vector. The resulting binary vector was transferred into *Agrobacterium tumefaciens* strain EHA105 and LBA4404 for *Citrus* and tobacco transformation. Empty Vector (EV) without SDE15 fragment was used for *Citrus* and tobacco transformation as negative controls.

[0211] For Y2H, the coding region of SDE15 (minus the putative signal peptide) was amplified and cloned in-frame with the GAL4 DNA-binding domain (BD) of the bait vector pGBKT7 to generate BD-SDE15 for Y2H library screening and co-transformation in yeast. The 996 bp coding sequence of CsACD2 was PCR-amplified from *Citrus* leaf cDNA and cloned in-frame with the GAL4 DNA-activating domain (AD) of the prey vector pGADT7 to generate AD-CsACD2 for co-transformation with bait vector in yeast to confirm the interaction. BD and AD vectors were constructed by using the In-Fusion cloning kit (Clontech, Mountain View, CA, USA).

[0212] Transient expression in *Citrus* protoplast were used for subcellular localization and BiFC assays. For the subcellular localization assay, the coding region of SDE15 without signal peptide was inserted into EcoRI-digested C-terminal EYFP containing vector pSAT6-EYFP-NI by using In-Fusion cloning kit to generate SDE15-EYFP fusion proteins. For the BiFC assay, SDE15 and CsACD2 were inserted into SalI-digested BiFC vectors pSAT6-nEYFP-CI and pSAT6-cEYFP-C1-B by using In-Fusion cloning kit to produce SDE15-EYFPN, EYFPC-SDE15, CSACD2-EYFPN and EYFPC-CsACD2 fusion proteins. All the vectors were subsequently used for *Citrus* protoplast transformation.

[0213] To generate recombinant protein constructs for GST pull-down assay and red chlorophyll catabolite reductase (RCCR) assay, the coding region of SDE15 (without signal peptide) was inserted between EcoRI and XhoI sites of pGEX-4T-1 vector (GE Healthcare, Chicago, IL, USA) to generate GST-SDE15 fusion protein vector as bait. The coding sequence of CsACD2 was inserted between BamHI and EcoRI sites of pMALTM-c5X vector (NEB, Ipswich, MA, USA) to generate MBP-CsACD2 fusion protein vector as prey and source of RCCR for enzyme assay. The truncated sequences of SDE15 fragments (SDE15^{ΔN} and SDE15^{ΔC}) were inserted into EcoRI-digested pGEX-4T-1 vector by using In-Fusion cloning kit to generate GST-SDE15^{ΔN} and GST-SDE15^{ΔC} fusion protein vectors as bait. The coding sequence of RCCR domain of CsACD2 was amplified and inserted between BamHI and EcoRI sites of pMALTM-c5X vector by using In-Fusion cloning kit to generate MBP-RCCR fusion protein vector as prey.

[0214] To generate the constructs for agro-infiltration assay in *N. benthamiana*, modified pCambia1380 vectors were constructed by inserting cauliflower mosaic virus promoter (CaMV 35S) and EYFP/CFP coding sequence to create the transient expression vectors with C-terminal EYFP reporter protein (pCambia1380-35S-EYFP) or C-terminal CFP reporter protein (pCambia1380-35S-CFP). The coding sequence of SDE15 without signal peptide was PCR-amplified and inserted between BamHI and KpnI sites

of pCambia1380-35S-EYFP and pCambia1380-35S-CFP to generate 35S-SDE15-EYFP and 35S-SDE15-CFP individually. The truncated sequences of SDE15 fragments (SDE15^{AN} and SDE15^{AC}) were inserted between BamH I and Kpn I sites of pCambia1380-35S-EYFP by using In-Fusion cloning kit to generate 35S-SDE15^{AN}-EYFP and 35S-SDE15^{AC}-EYFP. The coding sequence of CsACD2 was inserted between BamH I and Kpn I sites of pCambia1380-35S-EYFP to generate 35S-CsACD2-EYFP vector. The coding sequence of NbACD2 was inserted between BamH I and Kpn I sites of pCambia1380-35S-EYFP by using In-Fusion cloning kit to generate 35S-NbACD2-EYFP vector. All the vectors were then transferred into *Agrobacterium tumefaciens* strain GV2260 for agro-infiltration assay.

[0215] All the primers used for vector construction were listed in Table S2.

Lee, L. Y., Vyas, S., Glick, E., Chen, M. H., Vainstein, A., Gafni, Y., Gelvin, S. B., and Tzfira, T. (2006). J Mol Biol 362, 1120-1131; Lee, L. Y., Fang, M. J., Kuang, L. Y., and Gelvin, S. B. (2008). Plant Methods 4, 24). For the BiFC assay, the coding sequence of SDE15 without signal peptide and full-length CsACD2 were cloned into either N-terminal or C-terminal fragments of EYFP vectors pSAT6-nEYFP-C1 and pSAT6-cEYFP-C1-B (Citovsky et al., 2006). The combinations of SDE15-EYFP^N/EYFP^C-CsACD2 and CsACD2-EYFP^N/EYFP^C-SDE15 were transiently co-transformed into protoplasts. Other combinations, such as SDE15-EYFP^N/EYFP^C, EYFP^N/EYFP^C-SDE15, SDE15-EYFP^N/EYFP^C-SDE15, CsACD2-EYFP^N/EYFP^C, EYFP^N/EYFP^C-CsACD2, CsACD2-EYFP^N/EYFP^C-CsACD2, EYFP^N/EYFP^C were also transformed into *Citrus* protoplasts as controls. After incubation in dark overnight,

TABLE S2

primers used for vector construction		
	Forward (5'-3')	Reverse (5'-3')
SDE15-OE	GGGGATCCATGGATACTCTCTGACTC	GGGGTACCTCTTTCCCATCTCTAAC
BD-SDE15	CATGGAGGCCGAATTCATGGATACTCTCTGACTC	GGATCCCCGGGAATTCCTTTCCCATCTCTAAC
AD-CsACD2	GGAGGCCAGTGAATTCATGGCTGTGAACCACTTATG	CACCCGGGTGGAATTCGGCAGTAAAAACCTTCTGTA
SDE15-BiFC	GAATTCTGCAGTCGACATGGATACTCTCTGACTC	CCGCGGTACCGTCGACTCTTTCCCATCTCTAAC
CsACD2-BiFC	GAATTCTGCAGTCGACATGGCTGTGAACCACTTATG	CCGCGGTACCGTCGACGGCAGTAAAAACCTTCTGTA
GST-SDE15	GGGAATTCATGGATACTCTCTGACTC	GGCTCGAGTCTTTCCCATCTCTAAC
GST-SDE15 ^{AN}	TGGATCCCCGGAATTCATGGACGACTCCCATATCAA	GTCGACCCGGGAATTCCTTTCCCATCTCTAAC
GST-SDE15 ^{AC}	TGGATCCCCGGAATTCATGGATACTCTCTGACTC	GTCGACCCGGGAATTCATATTGTTCTTTATCTTTAT
MBP-CsACD2	TATCGTCGACGGATCCATGGCTGTGAACCACTTATG	TACCTGCAGGGAATTCGGCAGTAAAAACCTTCTGTA
MBP-RCCR	TATCGTCGACGGATCCATGCCTGTTAGGCAGCTGAT	TACCTGCAGGGAATTCGGCAGTAAAAACCTTCTGTA
MBP-NbACD2	TATCGTCGACGGATCCATGGCTATTTCAATATCCT	TACCTGCAGGGAATTCAGCATTGTAGATTCTCC
SDE15-YFP	TTGGATCCATGGATACTCTCTGACTC	GGGGTACCTCTTTCCCATCTCTAAC
SDE15-CFP	TTGGATCCATGGATACTCTCTGACTC	GGGGTACCTCTTTCCCATCTCTAAC
SDE15 ^{AN} -EYFP	GGACTCTAGAGGATCCATGGACGACTCCCATATCAA	CCCTTGCTCACCATTGGTACCTCTTTCCCATCTCTAAC
SDE15 ^{AC} -EYFP	GGACTCTAGAGGATCCATGGATACTCTCTGACTC	CCCTTGCTCACCATTGGTACCTATATTGTTCTTTATCTTTAT
CsACD2-YFP	TTGGATCCATGGCTGTGAACCACTTATG	GGGGTACCGGCAGTAAAAACCTTCTGTA
NbACD2-YFP	GGACTCTAGAGGATCCATGGCTATTTCAATATCCT	CCCTTGCTCACCATTGGTACCGCATTGTAGATTCTCC

*Nucleotides underline is the restriction enzyme cutting site

Transient Gene Expression in Citrus Protoplasts

[0216] Protoplasts were isolated from etiolated Duncan grapefruit epicotyl segments by following the protocol of transient gene expression in *Arabidopsis* mesophyll protoplasts with modifications (Yoo, S. D., Cho, Y. H., and Sheen, J. (2007). Nat Protoc 2, 1565-1572). Briefly, epicotyl segments of Duncan grapefruit cultured in dark were cut to small pieces and digested in Cellulose "Onozuka" R-10 and MACEROZYME R-10 (Yakult Pharmaceutical, Japan) enzyme solution overnight. Protoplasts were harvested and used for plasmid transformation. Plasmids were transformed into *Citrus* protoplasts by the polyethylene glycol 4000 (PEG4000)-mediated transformation method (Citovsky, V.,

the EYFP signals were examined and photographed under a fluorescence microscope for BiFC assay with excitation wavelength 514 nm (Olympus, Tokyo, Japan).

Plant Transformation and Pathogen Inoculation

[0217] *Agrobacterium* mediated transformation of etiolated epicotyl segments of Duncan grapefruit were carried out as described previously (Orbović, V., and Grosser, J. W. (2015). Methods Mol Biol 1224, 245-257). *Agrobacterium tumefaciens* EHA105 harboring the recombinant plasmid was used for *Citrus* transformation. Transgenic lines showing kanamycin-resistance and erGFP-specific fluorescence were selected and then micro-grafted in vitro onto 1-month

old Carrizo citrange nucellar rootstock seedlings. After a month of growth in vitro, the grafted shoots were potted into a peat based commercial potting medium and acclimated under greenhouse conditions.

[0218] *N. tabacum* cv. Petite Havana SR1 seeds were sown on MS medium (Sigma-Aldrich, St. Louis, MO, USA) containing 3% sucrose and 0.8% agar and allowed to germinate at $22 \pm 1^\circ$ C. (16h light and 8 h darkness). Subsequently, plants were grown and maintained in MS medium. Fresh tobacco leaf discs were infected with *A. tumefaciens* strain LBA4404 harboring the recombinant plasmid. The regenerated shoots were maintained on MS medium supplemented with 0.2 mg L^{-1} NAA and 3 mg L^{-1} 6-BA along with 100 mg L^{-1} kanamycin and 500 mg L^{-1} cephotaxime. Kanamycin-resistant, eGFP and PCR positive shoots of T0 transgenic plants were selected, transferred to the greenhouse and maintained up to T2 generations, which were used for phenotype inspection and further analysis.

[0219] For the HLB pathogenicity assay, the SDE15-transgenic and EV-transgenic trees was inoculated with Las via grafting as previously reported (Li, J., Pang, Z., Trivedi, P., Zhou, X., Ying, X., Jia, H., and Wang, N. (2017). *Mol Plant Microbe Interact* 30, 620-630). Midrib DNA was isolated from the grafted trees monthly after grafting up to 4-month post grafting and used to quantify Las by Taqman qPCR with Primer/probe combination (CQUA04F-CQUA04R) as described previously (Wang, Z., Yin, Y., Hu, H., Yuan, Q., Peng, G., and Xia, Y. (2006). *Plant Pathology* 55, 630-638). The Ct value of each amplicon represents the Las genomic copy numbers in 100 ng *Citrus* midrib DNA. The test was repeated three times.

[0220] For the *Xanthomonas citri* subsp. *citri* (Xcc) pathogenicity and hypersensitive reaction (HR) assays in *Citrus*, SDE15-transgenic and non-transgenic Duncan grapefruit plants were used for inoculation in a quarantine greenhouse. The wild-type strain Xac306 causes disease on grapefruit whereas the Xcc A^w strain triggers hypersensitive reaction in grapefruit leaves⁷. Xcc strains were grown with shaking overnight at 28° C. in NB, centrifuged down, and suspended in sterile tap water, and the concentrations were adjusted to 10^6 CFU/ml (for Xac306) and 10^8 CFU/ml (for Xcc A^w) individually. Bacterial solution was infiltrated into fully expanded, immature leaves with needleless syringes (Yan, Q., and Wang, N. (2012). *Mol Plant Microbe Interact* 25, 69-84.). The tests were repeated three times with similar results. Disease symptoms and HR phenotype were photographed at 3, 5, 7, 9 and 11 days post inoculation. Growth curve assay of Xac 306 was conducted at 0, 1, 3, 5, 7, 9 and 11 days post inoculation.

Agro-Infiltration Assay in *N. benthamiana*

[0221] *A. tumefaciens* strain GV2260 cells containing binary vectors were cultured overnight in LB medium with $50 \mu\text{g ml}^{-1}$ of rifampicin and $50 \mu\text{g ml}^{-1}$ kanamycin and re-suspended in induction medium (10 mM MgCl₂, 10 mM MES pH 5.6, 200 uM acetosyringone), and incubated at 25° C. with shaking for 4 h. The cultures were diluted to OD600 of 0.1 or 0.2. For each vector, three leaves of young *N. benthamiana* plants were infiltrated with diluted *A. tumefaciens* suspension as triplicates.

[0222] For the HR assay, young leaves of *N. benthamiana* were first infiltrated with *A. tumefaciens* cells containing binary vectors for SDE15-EYFP and/or CsACD2-EYFP by using a needleless syringe, kept in a greenhouse for 2 days

and then infiltrated with another *A. tumefaciens* strain harboring the binary vector carrying the AvrBsT protein which can trigger HR as reported previously Kim, N. H., Choi, H. W., and Hwang, B. K. (2010). *Mol Plant Microbe Interact* 23, 1069-1082). Agro-infiltrated plants were kept in a greenhouse and HR were examined and photographed at 3 days post AvrBsT inoculation. For the electrolyte leakage assay, leaf discs of AvrBsT infiltrated plants at 2 days post infiltration were floated on deionized water with shaking. The conductivity of the solution was measured 4 h later using an Oakton™ Conductivity Benchtop Meters (Thermo Fisher, Waltham, MA, USA). The *A. tumefaciens* transformant cells harboring an empty vector were infiltrated into the leaves of *N. benthamiana* as controls.

[0223] For the localization and co-localization assay, CsACD2-EYFP was co-expressed with the plasma membrane localization-marker PM-CFP, the nucleus-marker CFP-nucleus or SDE15-CFP in leaves of *N. benthamiana*. *A. tumefaciens* strain GV2260 harboring the corresponding plasmids were infiltrated into leaves at OD600 of 0.2. Subcellular localization was inspected and photographed 1 day post infiltration.

Extraction of Phloem Sap Proteins

[0224] An optimized method of protein extraction from phloem sap was performed by combining two methods reported before (Hijaz, F., and Killiny, N. (2014). Collection and chemical composition of phloem sap from *Citrus sinensis* L. Osbeck (sweet orange). *PLOS One* 9, e101830; O'Leary, B. M., Rico, A., McCraw, S., Fones, H. N., and Preston, G. M. (2014). *J Vis Exp.*). Briefly, 10-20 cm (0.5 cm diameter) stems from Las infected and uninfected trees were collected. The bark area was stripped into two pieces and was manually removed from the twig. The inner part of the bark was rinsed with deionized water and dried with Kim wipes. Then the bark strips were cut into about 1-cm pieces using a sterile razor blade and placed in a 60-mL syringe filled with distilled water. Vacuum was applied for 5-15 seconds repeatedly to let water penetrated barks. Then the barks were dried with Kim wipes and placed in a 20-mL syringe, centrifuged in 50 mL falcon tube for 10 min at 4,000 g, at 4° C. The collected phloem sap was centrifuged at 15,000 g for 5 min. The supernatant was heating for 5 min at 95° C. in SDS gel-loading buffer for SDE15 detection with specific antibody.

RNA Isolation and Expression Analysis of HLB Associated Genes

[0225] qRT-PCR was performed to detect the expression of SDE15 in SDE15-transgenic plants (both in *Citrus* and tobacco) and in non-transgenic *Citrus* plants and psyllids. We also examined the expression of HLB marker genes in the SDE15-transgenic *Citrus* and PR genes in SDE15-transgenic plants after HR induction. Total RNA of transgenic *Citrus*, transgenic tobacco and psyllids were extracted by Trizol reagent (Thermo Fisher) and digested with DNase I (Promega) followed by the manufacturers' instructions. First-strand cDNA was synthesized from purified RNA with ImProm-II™ Reverse Transcription System (Promega) and diluted 10 times for RT-qPCR to detect related genes with specific primers (Table S3). 20 μL of qPCR reaction consisted of 10 μL of 2xKiCqStart® SYBR® Green qPCR ReadyMix™ (Sigma-Aldrich), 1 μL of each primer (5 μM),

2 µl of diluted cDNA template, and 6 µl of DNase/RNase free water. The PCR cycling consisted an initial activation step at 95° C. for 3 min, followed by 40 cycles of 95° C. for 15 s and 60° C. for 40 s. All cDNA samples were run in triplicates. Citrus GAPDH gene, tobacco Actin gene and Las gryA gene were used as endogenous controls wherever appropriate.

[0226] The qPCR primer sequences of specific genes and endogenous control genes are listed in Table S3.

(Clontech). Standard positive controls (pGBKT7-53 and pGADT7-T; Clontech) and standard negative control (pGBKT7-Lam and pGADT7-T) were included. After mating between the Gold strain transformed with BD: SDE15 and the Y187 libraries, diploid yeasts were plated on synthetic dropout (SD)/-Leu/-Trp (DDO), SD/-Leu/-Trp/-Ade/-His (QDO) and SD/-Leu/-Trp/-Ade/-His plus X-α-gal and Aurcobasidin A (AbA) (QDO/A/X) agar plates to detect the activation of reporter genes HIS3, ADE2, MEL1 (for α-ga-

TABLE S3

primers used for qRT-PCR and Tagman probe PCR analysis		
	Forward (5'-3')	Reverse (5'-3')
qSDE15	ACTCCCATATCAAAAGCCTACG	CGTATCTTTCACCATCCATCCTC
qCsACD2	GGCTAAATCAGTGTGCTGTG	ATCAACCCATCCCTCTTTTCC
PR1	AAATGTGGGTGAATGAGAAAGC	ATTATTGTTGCACGTCACCTTG
PR2	TTCCACTGCCATCGAAACTG	GTAATCTTGTTTAAATGAGCCTCTTG
PR3	GGCTCAAACCTTCACATGAAACTAC	GTTGACAATAATCTCCAGGGTTTC
PR5	CACCATTGCCAATAACCCCTAATG	GGGACAGTTACCGTTAAGATCAG
PP2-B15	TCGTTGCCATCAGAAGTATCAC	CCAACGCAAATAAACTGTCCC
WRKY40	CTCCTGTTCCAAATGCCAAG	CCGAGGTGAGGGATTATCTTTAG
ZIP5	TGAATATGCTGGTGAATCGGAG	GCTGCAACCAAAGGCTTAATAG
Sweet7	GCTAACCCCTACTTCACCTCCAC	GGCATATACTCCACGCTCTTG
Sweet15	GTGTTGCCGTTTCTGTTAGTG	GCGAACCACATAATTGCACTC
Chalcone synthase	GCGTTCTAGTCGTATGCTCTG	GCCAATGATTAAAGCTGCGG
CsGAPDH	GGAAGGTCAAGATCGGAATCAA	GGAAGGTCAAGATCGGAATCAA
NbActin	CCTGAGGTCCTTTTCCAACCA	GGATTCCGCGAGCTTCCATT
gryA	CAATGTGCTGGTCAATGGTG	AATCTCCATCAAGGCATCCAG
CQULA04 (β operon primer)	TGGAGGTGTAAAAGTTGCCAAA	CCAACGAAAAGATCAGATATTCTCTA
CQULAP10 (β operon probe, 5'-3')	FAM-ATCGTCTCGTCAAGATTGCTATCCGTGATACTAG-TAMRA	

Yeast Two-Hybrid Library Screening and Interaction Analysis

[0227] Total RNA was extracted from leaves of Valencia sweet orange (HLB symptomatic (S, Las Ct value 25~26 per 100 ng DNA), HLB asymptomatic (AS, Las Ct value 28~30 per 100 ng DNA) and healthy (H, Las free)) by Trizol reagent (Thermo Fisher), digested by DNase I (Promega). mRNA samples were purified using the NucleoSpin® RNA kit (Clontech). ALL three types of mRNA samples were used to construct yeast two-hybrid libraries in the pGADT7-Rec vector using the Make Your Own “mate and plate” library system (Clontech) following the manufacturer’s instructions and transformed in the yeast strain Y187 by using Yeast-maker™ Yeast Transformation System 2 (Clontech). The titer of each constructed library is more than 3×10⁸ which represents the good transformation efficiency. BD:SDE15 construct was transformed into Y2HGOLD yeast strain (Clontech) Library screening was performed according to the Matchmaker Gold yeast two-hybrid system protocol

(lactosidase activity) and AbA^r (for Aurcobasidin A resistance).The fragments of positive diploid yeast were amplified by colony PCR with Matchmaker® Insert Check PCR Mix 2 (Clontech) and analyzed by electrophoresis on a 0.8% TAE Agarose/EtBr gel. The PCR products with single band were purified and sent for sequencing. The PCR products with multiple bands indicate the presence of more than one prey plasmid in a heterozygote cell. For this situation, plasmids were isolated from the heterozygote cells with multiple plasmids with Easy Yeast Plasmid Isolation Kit (Clontech) and transferred into *E. coli* for sequencing. BLAST was used to compare the inserts nucleotide sequences to the genome of sweet orange to identify corresponding proteins which interact with SDE15.

Recombinant Proteins Expression and GST Pull-Down Assay

[0228] *E. coli* cells expressing GST or GST fusion proteins were washed in PBS buffer and suspended with Cel-

Lytic B Cell Lysis Reagent (Sigma-Aldrich) to generate the cell lysates. After centrifugation, the cell lysates were incubated with glutathione agarose beads in accordance with the GST Protein Interaction Pull-Down Kit instructions (Thermo Scientific). The beads were washed to remove the unbound proteins and incubated with *E. coli* cell lysates expressing either MBP or MBP fusion protein for 1 to 2 h at 4° C. After washing four times, the beads were eluted with 10 mM glutathione, and the eluates were collected and immunoblotted using anti-MBP (NEB) and anti-GST (Abcam, Cambridge, UK) antibodies.

Enzyme Assays

[0229] Coupled Pheophorbide a oxygenase (PaO)/RCCR assay was performed to test CsACD2 activity according to published procedures (Hortensteiner, S., Vicentini, F., and Matile, P. (1995). Chlorophyll breakdown in senescent cotyledons of rape, *Brassica napus* L.: Enzymatic cleavage of phaeophorbide a in vitro. *New Phytol.* 129, 237-246; Wüthrich, K. L., Bovet, L., Hunziker, P. E., Donnison, I. S., and Hortensteiner, S. (2000). *Plant J* 21, 189-198; Pruzinska et al., 2005). Thylakoids containing PaO were isolated and solubilized from senescent *Citrus* leaves as described previously (Hortensteiner et al., 1995). PaO was partially purified from solubilized membranes and used for enzyme assay (Rodoni, S., Vicentini, F., Schellenberg, M., Matile, P., and Hortensteiner, S. (1997). *Plant Physiol* 115, 677-682.). MBP-CsACD2 fused protein was expressed and purified with the pMAL protein fusion & purification system (NEB) as the source of RCCR. Briefly, assays (total volume of 50 µl) contained different combinations of PaO (equivalent to 0.5 g of tissue), *E. coli* (50 µg) protein extracts as a source of RCC-forming factor (RFF), and purified MBP-CsACD2

(1.5 µg) as the source of RCCR. The assays were supplemented with 0.5 mM pheide a, 10 µg ferredoxin (Fd), and a Fd-reducing system consisting of 2 mM Glc-6-P, 1 mM NADPH, 50 milliunits of Glc-6-P dehydrogenase, and 5 milliunits of Fd-NADP⁺ oxidoreductase. After 1 hour incubation at 25° C., reactions were terminated by the addition of 80 mL methanol. Formation of primary fluorescent chlorophyll catabolite (pFCC) was followed by reversed-phase HPLC with 36% (v/v) 50 mM potassium phosphate buffer, pH 7.0, in methanol as solvent. Activities are determined as integrated fluorescence units (320/450 nm) of pFCCs.

Quantification of Compounds Participating Chlorophyll Break-Down Pathway

[0230] Three compounds (chlorophylls a, b and pheophorbide a) in Chlorophyll break-down pathway were extracted and quantified as previously described (Garrido, J. L., Rodríguez, F., Campaña, E., and Zapata, M. (2003). *J Chromatogr A* 994, 85-92.), with modification. Briefly, leaf samples of SDE15 transgenic *Citrus*, EV transgenic *Citrus*, SDE15 transgenic *Citrus* infected with HLB and EV transgenic *Citrus* infected with HLB were collected and homogenized with 8 ml of 90% acetone and left for 16 hours at -10° C. All extracts were filtered through 25 mm, 0.2 µm GHP Acrodisc filters (Sigma-Aldrich) prior to injection. All sample preparations were done under subdued light. The standards of chlorophylls a, b and pheophorbide a were obtained from Sigma-Aldrich. All the standards and samples were followed by reversed-phase HPLC. Mobile phase consisted of (A) methanol, (B) 0.025M ammonium acetate and (C) acetone. A linear gradient from A-B (80:20, v/v) to A-C (80:20, v/v) was pumped during 15 min, followed by an isocratic hold at A-C (80:20, v/v) during a further 5 min. The flow-rate was 1 ml/min.

SEQUENCE LISTING

```
Sequence total quantity: 51
SEQ ID NO: 1          moltype = AA  length = 14
FEATURE              Location/Qualifiers
source                1..14
                     mol_type = protein
                     organism = Liberobacter asiaticum sp.

SEQUENCE: 1
DDSHNQKPTK KKPNN                                         14

SEQ ID NO: 2          moltype = DNA  length = 291
FEATURE              Location/Qualifiers
source                1..291
                     mol_type = genomic DNA
                     organism = Liberobacter asiaticum sp.

SEQUENCE: 2
atgacaatat caaaaaaatca agccattctt ttctttatta caggcatgat actttcttct 60
tgtggtgata ctctctctga ctctaagcaa cataataaaa tcaacaatac aaaaaatcat 120
cttgatcttc ttttcccaat agacgactcc cataatcaaa agcctacgga aaaaaaacca 180
aatatcatcat ccataaagat aaagaacaat ataatagaac cacaaccggy ccctagtcgc 240
tggaagaggag gatggaatgg tgaaagatac gttagagaat gggaaagata a          291

SEQ ID NO: 3          moltype = DNA  length = 156
FEATURE              Location/Qualifiers
source                1..156
                     mol_type = genomic DNA
                     organism = Liberobacter asiaticum sp.

SEQUENCE: 3
atgtatacca aaagtttgtt aatggtagct tatttgttat ctctcggttc aatatcggtt 60
gggctatgct ttaaccgccc aaaaggcccg tctaaagaag aacaagcacg aatagaacaa 120
atcagagcag aagcaagaga aaggcgctac caataa          156

SEQ ID NO: 4          moltype = DNA  length = 1227
```

-continued

FEATURE
source

Location/Qualifiers
1..1227
mol_type = genomic DNA
organism = *Liberobacter asiaticum* sp.

SEQUENCE: 4

atgtttcaaa	ataagaattt	tttattaggc	gttttgcgct	taaaaaagtg	cacgcgtggt	60
gtttttttag	ttattacagc	tattttattg	tcttcttttg	tcgccatagt	agatgttggtg	120
gtagatcagg	tgacgggtgat	gcagaaaaa	gcttggtctc	aagaggtcct	ggatcatggt	180
atttatcgca	cttctccaaa	aaatttatat	gatttaaggg	aagcgggtag	agataatttt	240
ataagacatc	aaattgaaaa	ggctttaaat	acttataatt	ccagagatct	ttcaaatatt	300
gggaggtatag	aatcaatcgt	taaagatgct	gttattttta	cgaaaaatgt	taattctttg	360
cctttacaat	ttactgtcga	catcgccctt	tcgacaacgg	ttcaattgcg	aggatctctt	420
ttgcagatgt	tttcgcaatc	gaagggtaaa	gttgatattt	ctcgtcgaaa	aaaagttatg	480
tataaaca	aatattggtt	aatgattatg	ccttttgcat	gggatggcta	ttggctagct	540
tcaagaggaa	agggttgaga	ctcaaaagta	catcctccaa	agtatttgga	gtattctcat	600
tattatcagc	agtatcttaa	tagaaatcgc	cttgtgaaaa	atttctctc	acaaattcct	660
tataaaaa	tctgtatggc	tccttatcat	tattctttct	ttctttattg	ggcagtcgga	720
accttgacgt	atagcgtgga	taacaagacg	actacgcgcg	aatattataa	ggatccttac	780
tatgctacat	gggactattt	tccttattca	tttatcaaga	acgtatttga	catgacttcc	840
aatcaatttg	gagatgggca	ggtattgact	aataccaatc	attgctttcc	tcattggagca	900
agtcagaata	aataatgtgt	gatgcttgcg	ataggaaatc	agctatctcg	aagttcagta	960
gagaaagaaa	aaatagaaaa	ggttctgcaa	gactgccatt	acatgcataa	gcgtcataga	1020
actggaagag	atgcaattac	catattttct	gtaggatttt	ctccggatca	agatacgcga	1080
tatactttta	gacaatgtgc	aagtgatcca	tctaataatt	atgaaattaa	ttctgacgag	1140
aatgtcatgc	ctatagccaa	aagtccttga	cgtaattgta	taactaattg	gtttagttag	1200
tttacaatta	ctgtgggtga	tagctag				1227

SEQ ID NO: 5

FEATURE
source

Location/Qualifiers
1..678
mol_type = genomic DNA
organism = *Liberobacter asiaticum* sp.

SEQUENCE: 5

atgagagata	taagaaaaat	tagaaattat	tttaggaata	ctgctaaaa	tatattgagt	60
gggttatctc	taggggtttt	ttcttctgct	gcaatggcag	actatgggta	ttctccccag	120
tttcagccga	ctataatggt	gtccaatttt	gcaaaattta	aagggttata	tgttgctgct	180
gattttttca	aaatagatca	tcagtcgcct	gttcgtttgc	aaaatctttc	tttaaatggg	240
gtgtccattg	gtcttgatgg	tcaagatgga	acccttggtt	atggtgcttc	tttgggtgtc	300
gagggatttc	atcttgaaac	acgaggggga	attgatgggg	ataaggtagc	gggaacactc	360
ttgtttcgta	ccggtttttac	gtttgataat	aataattcct	ctattctcca	aaatactcct	420
atttatgggt	ttggtggagc	tctgtataga	aatattatgt	ctgttgaatc	tgctgacaca	480
gcaaaatcca	caatacgaaa	cattgtagca	aacggttttt	tagataaagt	tattggtgtg	540
gggattgaaa	agaaacttgc	tagcatgctc	tcgattcgtg	gtgagtatcg	ttatgtcgct	600
tgttatgacc	agccttgggg	tgtcagcaag	tggagagaaa	aaggtgactt	cacagctggt	660
gtgggtttac	gctttttaa					678

SEQ ID NO: 6

FEATURE
source

Location/Qualifiers
1..150
mol_type = genomic DNA
organism = *Liberobacter asiaticum* sp.

SEQUENCE: 6

atgaatacaa	gaataatagg	aaccgtatta	atgcttgcta	ttagtcctct	tttattttagc	60
tgttcttcta	aaaaaggagg	tgaaaaaaaa	ggtgaggaga	aaaaaacttc	tgctccgttg	120
aaaaatggta	aaaatcaatc	cagaagatag				150

SEQ ID NO: 7

FEATURE
source

Location/Qualifiers
1..1607
mol_type = genomic DNA
organism = *Citrus sinensis*

SEQUENCE: 7

ataagcaagt	ggccaagtag	tgatttgacg	gagtgattaa	gaacgaatgg	ctgtgaacca	60
cttatgccag	tggcagttat	tacgcttcca	gctctctcat	ccatcggtc	cggcttgacg	120
atatttatct	ccttcogagac	caaaagtcct	aacgtcgta	accgccaag	tcaattgttc	180
tgccgcacca	tcgtcgtctc	cgatggactc	gcacaacgaa	ggccgtaaga	agttcatgga	240
attcccttac	gcttcaggcc	ctgttaggca	gctgatgggt	gatctcgat	caacggtgga	300
gaataccctc	gattcgcagc	tactcccttg	cactctgcca	ccagatgtac	agtattacga	360
gaaccaa	ggcactgctc	aagcttctct	tcaaatcaga	tccgggctca	agtcctcact	420
ggtagtctct	tcttttacat	aattggcggt	tgtttaattg	cgagtgtcaa	cttgagttaa	480
taagtctcat	aacttgattt	aactgaaac	taggcgcctc	tcttaaacat	tggaaactgaa	540
cgcatttatg	tctgaaatag	attttctttt	cttagccctag	gattgcacta	ctctgcttag	600
ttgcctaagc	ataagaaaca	aatgattcaa	aatttactag	gctctagtgt	gacattcaag	660
ttcgaggcgt	acatggtctt	caataatttg	atgataatgg	atgcgtgtca	gattgatttc	720
atactgggaa	gttgggtaca	cagtgagcta	ccaacaggag	cagcattgaa	cataacaagc	780
ctttcagcat	atctaaactc	ttccactgat	gcaccaaac	tgctaattga	gctcatccag	840

-continued

```

agtagcccta cttctctagt cctcatcctt gacttgccct ctcgaaagga tcttgctctc 900
catcccgact atcttcacac tttctatgaa agcacacggg tggatgaata taggcaaatg 960
cttgagaaag tacctgaagt tagaccctac ttctcttctt ccctttactt aagatgtgtc 1020
gtctctcctt cagcaattat ggtccgtgta gatactgaaa ctgaaactgg ggcaggtgaa 1080
tcaacacggt tggactatat tataacaaat catgtgcatc ctgttgctaa gcaagttatt 1140
ggaatctggc taaatcagtg tgccttggtg gggagacatg taggggagtc agacaaggct 1200
tatctggaag agagggatgg gttgattaag aacaaaacta ttgagattga tctcggtctc 1260
agctttccga gattgtttgg accgcaggta gcaagccggg tattaggcga gatacagaag 1320
gtttttactg cctgaggttg gtatttgaat ttgaggttgg gaatgtacaa agaattggag 1380
ttgattgacc ttaattttag tgtgtgtatg aacatcattg tccccctttt tatgcacaag 1440
ttctttgatt tcttctgtga attgatattg cacttaaat actgttgctt tctaatttta 1500
ttcgaaattg ttaaaatttc gtgcattgat ggtgtttcta atcttgtaaa gcatgaagaa 1560
aggaacttac aacaatctca attaggaatg gaattgaatc ccttgaa 1607

```

```

SEQ ID NO: 8      moltype = DNA length = 1434
FEATURE          Location/Qualifiers
source           1..1434
                 mol_type = genomic DNA
                 organism = Citrus sinensis

```

```

SEQUENCE: 8
atttgacgga gtgattaaga acgaatggct gtgaaccact tatgccagtg gcagtattta 60
cgctccagc tctctcatcc atcggctccg gcttgccagat atttatctcc ttcgagacca 120
aagtcctcaa cgtcgtcaac cgccaaagtc aattgttctg ccgcaccatc gtcgtctccg 180
atggactcgc acaacgaagg ccgtaagaag ttcattggaat tccccatcgc ttcaggccct 240
gttaggcagc tgatggttga tctcgtatca acggtggaga ataccctcga ttcgcagcta 300
ctcccttgca cctcgccacc agatgtacag tattacgaga accaaaatgg cactgctcaa 360
gcttctcttc aaatcagatc cgggctcaag tctcactga ttgatttcat actgggaagt 420
tgggtacaca gtgagctagc aacaggagca gcattgaaca taacaagcct ttcagcatat 480
ctaaactctt ccactgatgc accaaacttg ctaattgagc tcatccagag tagccctact 540
tctctagctc tcatccttga cttgcctcct cgaaaggatc ttgtctccca tcccactat 600
cttcacactt tcatgaaaag cacacgggtg gatgaatata ggcaaatgct tgagaaagta 660
cctgaagtta gacctactt ctcttcttcc ctttacttaa gatgtgtcgt ctctccttca 720
gcaattatgg tccgtgtaga tactgaaact gaaactgggg caggtgaatc aacacgtttg 780
gactatatta taacaaatca tctgcatcct gttgctaagc aagttatttg aatctggcta 840
aatcagtggt cttgtggagg gagacatgta ggggagtcag acaaggctta tctggaaaag 900
agggatgggt tgattaagaa caaaactatt gagattgatc tcggctctag ctttccgaga 960
ttgtttggac cgcaggtagc aagccgggta ttaggcgaga tacagaaggt tttactgcc 1020
tgaggttggt attggaattt gaggttggga atgtacaaag aattggagtt gattgacctt 1080
aattttagtg tgtgtatgaa catcattgtc ccccttttta tgcacaagtt ctttgatttc 1140
tctctgtaat tgatattgga cttaaattac tgttgcttcc taattctatt cgaattgggt 1200
aaaaattcgt gcattgtatg tgtttctaatt cttgtaaaag atgaagaaag gaacttacaa 1260
caatctcaat taggaatgga attgaatccc ttgaacatct tgttgaactg ttggtagatt 1320
aattaaactg ctaagaatgg tgtgattgat gcactgaaga atatgaaatt tatttctgca 1380
tgattctact ctcactctat cataatgggt tgatcactct gctgagcctt aaag 1434

```

```

SEQ ID NO: 9      moltype = DNA length = 960
FEATURE          Location/Qualifiers
source           1..960
                 mol_type = genomic DNA
                 organism = Arabidopsis thaliana

```

```

SEQUENCE: 9
atggcgatga tattttgcaa cactctctac tcttcttctt ctccatcata tctctcgccg 60
ttaacttcaa aacgctcgcg attctcaaag aatctcagac ctcgagctca attccagtc 120
atggaagacc acgacgatca cctccgccga aaatttatgg agttcccgta tgtgtcacc 180
acgcggaagc agctcattgt tgatctcatg tcgacgggtg agaatcgctt ccaatcaca 240
ctcttccctt gtaacctccc tccagatgta cgaaacttca ataaccctaa cggttccgcc 300
gaagcatctc ttcatatcag atccggcgac aaatcttctc cgattgattt tgttatagga 360
agttgggatac attgcaagat cccaacagga gtatctttga atatacaag catctctgga 420
ttcttaaaat catcaacaaa agctccaaac tttgtggtcg aactaataca gacagttcc 480
aagtgcgttg tgctaactct tgacctccca catcgtaaa atcttgttct taaccggat 540
tatctcaagg agtattacca agacactgct cttgattctc atcgacaatc tctccttaag 600
ctacctgaag ttaaccctta tgtgtctcct tctctctttg tccgttctgc tttctctcct 660
actgcttcga tgcttaagat tgatgcggag gaagaggata agttggagga gatattgaga 720
gatcatgtta gtcagctgac taaggagggt ctcgaggttt ggttggagcg gtgtgtgaag 780
gaagaagaag agaagattgt ggttgggaaa gaagagagaa tggagttgga gagaagagat 840
aaaagcttta gaaggagag catagaggac gatttggatt tgcagtttcc gagaatgttt 900
ggtgaagaag tttctcccg tgtgttacac gctattaaag aagctttcgg tgttctctag 960

```

```

SEQ ID NO: 10     moltype = AA length = 318
FEATURE          Location/Qualifiers
source           1..318
                 mol_type = protein
                 organism = Arabidopsis thaliana

```

```

SEQUENCE: 10
MAMIFCNTLY SSSPSYLS LTKPSRFSK NLRPRAQFS MEDHDDLRR KFMEFPYVSP 60
TRKQLMVDLM STVENRLQSQ LLPCNLPPDV RNFNNPNSA EASLHIRSGD KSSPIDFVIG 120

```

-continued

SWIHKIPTG	VSLNITSISG	FLNSSTKAPN	FVVELIQSSS	KSLVLILDLP	HRKDLVLNPD	180
YLKEYYQDTA	LDSHRQSLK	LPEVNPYVSP	SLFVRSAPSP	TASMLKIDAE	EEDKLEEILR	240
DHVSPAAKEV	LEVWLERCVK	EEEEKIVVGE	EERMELERRD	KSPFRKSIED	DLDLQFPRMF	300
GEEVSSRVVH	AIKEAFGV					318

SEQ ID NO: 11 moltype = DNA length = 3154
 FEATURE Location/Qualifiers
 source 1..3154
 mol_type = genomic DNA
 organism = Citrus sinensis

SEQUENCE: 11

cagtggtgctc	tctccaaaca	aagcagacac	aatcattagg	catggctcgc	cccgtagagt	60
tggtctcgctc	cgctcatcttg	ttgctttgct	gcgctgccgc	agcatcagca	tcagcatcaa	120
gcttcgacga	ctccaatccg	atcagattgg	tatcatcgga	cggtctccgt	gacttcgaga	180
ctccgctcct	ccaggtagtc	ggccaagccc	gccatgctct	ctcctttgcc	cgttttgctc	240
gcaggtatgg	gaagatttac	gagtcctgtg	aggagatgaa	gctccggttc	gcgactttct	300
ccaagaacct	ggattttaac	agatctacca	attgtaaagg	cctatcttac	aggctcgggt	360
tgaacagtaa	gtttttcatt	ttgaatattg	gtctgtagct	caggaggcca	gtcaagccat	420
ctgattgcct	agattcctga	ttggtctgta	gttcacgtta	gtcacgtgct	ttctataaac	480
aatctacgca	ctagaaatga	atttttacga	ttttattatt	attactgtta	ttgttatgcc	540
catattttta	ccaaaactca	acttcgatga	cataaaatga	aagtgtgagg	gcccatttta	600
attaggataa	aacaatacaa	actctaactc	tcatcaaat	cagtattacg	cgataagaga	660
taaaaatttt	ctaattaatt	atactgttta	cttcaattat	atatattttt	ttttcatat	720
acgcgtgttc	ttagttattc	tatttgattt	gtatagggat	ttgtacaaa	tatatatttt	780
ctcatcatag	tactagtagt	ttgtgatct	tcatttcttg	ggaaccaatc	atggattgat	840
gatgcttag	cttatgtagt	gattctgtag	ttccctcagt	agattaatca	cggatgggta	900
taggttataa	ttggataatc	aagcttcaaa	cttatatttt	tgcattcatt	tgtgtatctc	960
gtgcaatcta	gtagtaataa	tttatagtga	aaatgaaaga	ggtattaaaa	ctgtcttatt	1020
tctactattg	ctagtgtaca	aataacgaga	atttgtataa	aaaatttaag	atcgtattta	1080
ctaaagttac	actagtatta	tgtattgcta	attaactcta	tacaattggt	gttgatggcg	1140
tttaaatgta	attaaactaa	cacgcagagt	ttgctgactg	gagctgggaa	gagttccaaa	1200
ggcacaggtt	gggagctcgc	caaaaactgct	ctgccactac	aaaaggaaat	cacaagctta	1260
ctgctgatgt	gcttcacaga	acggtaattc	tacgaaatc	tattctcgat	tgaacaacca	1320
agatgacct	cagtttctat	aagcttgat	tttgtattac	atagagagga	aaagactgat	1380
ttttctgctg	ctatatgtgt	agaaaactcg	gagggaaatc	ggcatagtaa	gcccagttaa	1440
agaccaaggt	cactgtggat	ctgctgggac	tttcaggcca	gcttgatttg	gaatgaaatc	1500
agaatttcta	aactgagttt	tcaattttag	tgctaaataa	ttaccttttg	cagcacgact	1560
ggatctcttg	aggtgcttta	ccaccaggcc	tttggaagg	gtatctcttt	gtcagaacag	1620
cagcttgtag	actgtgcccc	agctttcaac	aaccagggat	gcaatggtag	gttgccatcc	1680
caagcctttg	ataacatcaa	gtacaatggt	ggccttgata	ctgaggaagc	ataccctac	1740
accggaaaag	atggtgtctg	caaatctcca	tctgaaaatg	ttggcgtcca	agtcctggac	1800
tcagtcacaa	ttaccttggt	gagttttatg	ctgaattttc	attttaaatg	agagttagagt	1860
ctgcagtagt	ggactcacta	catgccaaac	aaatgaaaat	caaccataaa	taaatcaatc	1920
ataatctaga	aactctggat	atgtatatgt	ttcaagtttc	atgacttgta	agacaatgca	1980
gggtgctgaa	gatgaattac	agcatgcagt	tggtcttggt	cgacctgtga	gtgtggcatt	2040
tgaagttagta	gatggattcc	gattttacaa	gagtggaagt	tacagcagca	ctaaatgtgg	2100
aaatactccc	atggtgagtc	ttattgactt	ggaaaaacgat	acattatttt	gataggcatt	2160
gaatgcgaat	ctgttaaaaa	atgttttgat	ttatgttcaa	aatcaaaatt	taacagggca	2220
ttcaagtttt	gaagtttttg	aaaaaaaata	aatgcctaac	attatgggtc	aaactctcca	2280
aacaacattt	ctcattttct	tttctgaact	tctcagttgg	tcaacatggt	actcacgcaa	2340
ggaatgctta	atgtagacgt	tgtgtaaatg	atccttcact	tggaattatc	acccaagcat	2400
taaccactca	tccgatgcgt	ttattgcatt	tctaactttc	gggattttat	ggcaggatgt	2460
gaaccatgcc	gtcgttgctg	ttgggtacgg	agttgaagat	gggtgtccat	attggctgat	2520
taagaactct	tggggagaga	actggggcga	tcatgggtac	tttaagatgg	agatggggaa	2580
gaacatgtgt	ggtaagttac	tgttacatct	agattgtcag	taccagatca	ttgctcacia	2640
cttaaaattat	cagtcacgtg	caattttcgt	catcatcaat	tttcagtatg	acctaagata	2700
tgatgtcaat	aaaagaacaa	gaataatgta	gtatacatcc	gataagctct	gataagttaa	2760
cagagtagta	acagatctac	gcatgaaatt	tgaaacattt	aaataggtga	tatatatttg	2820
tgtttgctga	ttttactgat	ttgattgtag	gtattgcaac	ttgtgcatca	taccagttg	2880
tggcttagtc	tgctcctgaa	gaaatagttg	gatctggcta	tcagcaagtc	atttgctcat	2940
aaaaactata	tattttcact	caagaatgat	agcagaatgg	ttggctttat	gtacgaaata	3000
aattcggaga	ttaatgtcca	tataatctac	aatagcaatg	catggctgct	tgatgttcaa	3060
aataaattct	gagatctatg	tccatgtaaa	cagtcattgt	gactaggaca	ccaacgatgt	3120
tatatatatt	ttgtcaatgc	aaggtagttg	ttat			3154

SEQ ID NO: 12 moltype = DNA length = 2059
 FEATURE Location/Qualifiers
 source 1..2059
 mol_type = genomic DNA
 organism = Citrus sinensis

SEQUENCE: 12

cctcttttgt	gataaaaaatg	taagtaaaat	attttgtacg	attacaaatc	attcctatat	60
taacagaaat	gaacgctata	acggcagaga	aaaaaacctg	tacgacgacg	tgaaggaaat	120
tcacagaaac	gtcatcgctc	gcgtcagagt	tagttccctat	cgctacgat	acgacaaagc	180
cagcttagac	gcccggtcta	atctacaaat	ttccaatcca	tccctcagac	tgaaaatgat	240
acggaatctt	gagtcctgac	ttatcttgca	cccgacgagg	catctcataa	taataataat	300

-continued

```

aataataata ataataataa taataataca caactcattt caaccacata aattatgaat 360
ctcataaattt tataataattg agatgaatta tgtaagttac attgaagata acttagcact 420
tgccgaataa taatagtggt catagactgc attcatttaa tccgaagccc ttgagtgag 480
tgtgtctctc ccaaacaaag cagacacaat cattagggcat ggctcgcccc gtacagttgg 540
tctcgtccgt catcttggtg ctttgctgcg ctgcgcgagc atcagcatca gcatcaagct 600
tcgacgactc caatccgac agattgggtat catcggaagg tctccgtgac ttcgagacct 660
ccgtcctcca ggtgatcggc caagcccgcc atgctctctc ctttgcccggt ttgtctcgca 720
ggatggggaa gatttacgag tccgttgagg agatgaagct ccggttcgcg actttctcca 780
agaacttgga tttaatcaga tctaccaatt gtaaaggcct atcttacagg ctcggttgga 840
acaagtttgc tgactggagc tgggaagagt tccaaaggca caggttgagg gctgcgcaaa 900
actgctctgc cactacaaaa ggaaatcaca agcttactgc tgatgtgctt ccagaaacga 960
aagactggag ggaatctggc atagtgaagg caggtaaaga ccaaggtcac tgtggatctt 1020
gctggacttt cagcagcact ggaatctctg aggtgcttta ccaccaggcc ttggaaagg 1080
gtatctcttt gtcagaacag cagcttgtag actgtgcccc agctttcaac aaccagggat 1140
gcaatgggtg gttgccatcc caagcctttg aatacatcaa gtacaatggt ggcttgata 1200
ctgaggaagc atatccctac accggaagg atggtgtctg caaattctca tctgaaaatg 1260
ttggcgctca agtccctggac tcagtcaaca ttaccttgca tgcagttggt cttgtctgac 1320
ctgtgagtggt ggcatttgaa gtatgagatg gattccgatt ttacaagagt ggagtttaca 1380
gcagcactaa atgtggaagt actcccatgg atgtgaacca tgcctgctgt gctgttggtg 1440
acggagttga agatgggtgt ccatattggc tgattaagaa ctcttgaggga gagaactggg 1500
gcatcatggc ctactttaag atggagatgg ggaagaacat gtgtggtatt gcaactgtg 1560
catcatacc agtgtgtggt tagtctgctc ctgaagaaat agttgagatc ggctatcagc 1620
aagtcatttg ctcataaaac ttatattatt tcaactcaaga atgatagcag aatggttggc 1680
tttatgtacg aaataaattt ggagattaat gtccatataa tctacaatag caatgcatgg 1740
ctgcttgatg ttcaaaataa attctgagat ctatgtccat gtaaacagtc attgtgacta 1800
ggacaccaac gatgttatat atattttgtc aatgcaaggt agttgttata tggaaagctt 1860
aggcaaatat caatgcattg cttaaaaaat ttggttgtct tctgccctaa aaaggaactg 1920
agaaacttgc gtgagaatga tctgtgtgtc atgttgacat ctgcctacta gatgcattc 1980
aattcatgct cttcacaggc cttttcatta tcataatttg ttgctgaaaa ataaggggac 2040
ttagcagctt aacctctta 2059

```

```

SEQ ID NO: 13      moltype = DNA length = 2080
FEATURE            Location/Qualifiers
source              1..2080
                    mol_type = genomic DNA
                    organism = Citrus sinensis

```

```

SEQUENCE: 13
cctcttttgt gataaaaatg taagtaaaat atttgtgacg attacaaatc attcctatat 60
taacagaaat gaacgctata acggcagaga aaaaaaccgt tacgacgacg tgaaggaatt 120
tcacgagaac gtcacgcgcg gcgtcagagt tagttcctat cgtctacgat acgacaaaagc 180
cagcttagac gcccgcttct atctacaaat ttccaatcca tccctcagac tgaatatgat 240
acggaatctt gagtcttgac ttatcttgca cccgacgagg catctcataa taataataat 300
aataataata ataataataa taataataca caactcattt caaccacata aattatgaat 360
ctcataaatt tataataattg agatgaatta tgtaagttac attgaagata acttagcact 420
tgccgaataa taatagtggt catagactgc attcatttaa tccgaagccc ttgagtgag 480
tgtgtctctc ccaaacaaag cagacacaat cattagggcat ggctcgcccc gtacagttgg 540
tctcgtccgt catcttggtg ctttgctgcg ctgcgcgagc atcagcatca gcatcaagct 600
tcgacgactc caatccgac agattgggtat catcggaagg tctccgtgac ttcgagacct 660
ccgtcctcca ggtgatcggc caagcccgcc atgctctctc ctttgcccggt ttgtctcgca 720
ggatggggaa gatttacgag tccgttgagg agatgaagct ccggttcgcg actttctcca 780
agaacttgga tttaatcaga tctaccaatt gtaaaggcct atcttacagg ctcggttgga 840
acaagtttgc tgactggagc tgggaagagt tccaaaggca caggttgagg gctgcgcaaa 900
actgctctgc cactacaaaa ggaaatcaca agcttactgc tgatgtgctt ccagaaacga 960
aagactggag ggaatctggc atagtgaagg caggtaaaga ccaaggtcac tgtggatctt 1020
gctggacttt cagcagcact ggaatctctg aggtgcttta ccaccaggcc ttggaaagg 1080
gtatctcttt gtcagaacag cagcttgtag actgtgcccc agctttcaac aaccagggat 1140
gcaatgggtg gttgccatcc caagcctttg aatacatcaa gtacaatggt ggcttgata 1200
ctgaggaagc atatccctac accggaagg atggtgtctg caaattctca tctgaaaatg 1260
ttggcgctca agtccctggac tcagtcaaca ttaccttgga tgcagagtat gaattacagc 1320
atgcagttgg tctgttgcca cctgtgagtg tggcatttga agtagtagat ggattccgat 1380
tttacaagag ttgagtttac agcagcacta aatgtggaat tactccatg gatgtgaacc 1440
atgccgtcgt tgctgttggtg tacggagttg aagatggtgt tccatattgg ctgattaaga 1500
actcttgagg agagaactgg gccgacatg gctactttaa gatggagatg ggaagaacaa 1560
tgtgtggtat tggcaactgt gcatcatacc cagttgtggc ttagtctgct cctgaagaaa 1620
tagttggtat tggctatcag caagtcatct gctcataaaa cttatattat ttcaactcaag 1680
aatgatagca gaatggttgg ctttatgtac gaaataaatt cggagattaa tgtccatata 1740
atctacaata gcaatgcatt gctgcttgat gttcaaaata aattctgaga tctatgtcca 1800
tgtaaacagt cattgtgact aggacacca cgaatgttata tatattttgt caatgcaagg 1860
tagttgttat atggaagctt taggcaaaata tcaatgcatt gcttaaaaaa ttggttgtc 1920
tctgccctaa aaaaggaact gagaacttgc tgtgagaatg atcgtgtgtt cattgtgaca 1980
tctgcctact agatgcatt caattcatgc ccttttcatt atcataattt 2040
gttctgaaa aataagggca cttagcagct taacctctta 2080

```

```

SEQ ID NO: 14      moltype = DNA length = 2243
FEATURE            Location/Qualifiers
source              1..2243

```

-continued

```

mol_type = genomic DNA
organism = Citrus sinensis

SEQUENCE: 14
caaaactcacc gaaaattaat taaatctaata ctataataca aatacaaaaa cattaactac 60
cacgattgta taatgatatt tttgaatatt cttccact gaaagctctg tgcgaaaggg 120
ggaagtcaaa agtggttattc gttgtgtgtct atggctatga gttcgccact ttgggtagcc 180
ccaacttttc tttctcgctc atcgtcaaat atatcgagat caattactta caaacgttgc 240
aggctctcgg caagtaactt tttctcacaa tcagacatgg agcagccgct gatgaagggg 300
cagaagctga tggaaattccc tcacctgacg gcggcgacaca aagatctgat ggttagctta 360
atctcagcct tggagactag gcttgactct catcttcttg cttcttcggt agtcctcct 420
gatgttgagt tttatcagaa tgaccaaggc acttctcagg gttctcttca tatcagacgt 480
ggccttcctt cctctcatgt actcgcttcc tctccctctc gtgtccagat tcaaattcac 540
tggtgcttat tcatatttat attattgggc attttaatct cggtaaatct ggatatatat 600
atataatata atataatata agggcactgt tccaatccgg attggaacag cctccgggaa 660
tggtttgggg atcattatata atataatata atataatata ttttaagtgt ggacgctcgt 720
tttttttggg tttttatata gacctacggg tacagcagtt tttatacaga cactcttaaa 780
tcgtgtggtt taaaaaatta gtttttaaat aaactatata tatatatata tatattttat 840
tgcgaccgtc cttatttttt ttattcatgc tgatatactt taaaattaca gggtttaaga 900
gagttagttt ttaacacttt taaactgcac ggttttataa ttcattcaat atataaaaa 960
ttatttttag gtgcggacat ccgcattttt ttaactcatga acacactttt aaaccgaaaa 1020
gattagttat tgaatgaact ataaaatcat gtaattttta agagtgtccg cgtaaaaaaa 1080
aaaaataaagg catcctctag agaagacta tatatatata taatggtttt tggaaattga 1140
atgattaata accaccatgg agtcaattat caactctaat gtttaattggg atactgctga 1200
tcgcttttct cagcatgctt atagtagcta gctacatctt tccaaaactt atgtaaacat 1260
cctcctcttt tatgaacaaat atactaggaa gtggtactag tgtactactc taaataattg 1320
actacctata tttttttcca aaaaatctat caaaatctct cttcttcatg aacaatttaa 1380
atgggttaat attacgagag cccacttggt ttcaagtgac gcttgctgga tattacactt 1440
tgtataataa atgcttgtgg gaatgggaat cttttttctt ttttttttta acttttgcct 1500
tggcatggtt aacagattga ttctgtatta gcaagttggc ttcacttgaa ggtaccaacg 1560
ggaagtgcc a tgaacataac caactctcaa gcttacctaa aatcatcaac cgatgtacca 1620
cattttcaat tcgagcttgt ccaatgcagc cccacatatt tcattctctt cctagatata 1680
actcctagaa aagaccttgt tctataccca aattacctca aaacatttta cgaagaagct 1740
cagcttgaaa cattgagaca gagacttgag caagtcaccg aaaccaaacc ctacttgagt 1800
tctctctctt actttcgtgg tgtggtctcc ccaactggga ttttggctag cataaaatgt 1860
gaggaagtgt gtggaacgga tcgctgtgaa gagattatac gtgaacatgt gagccctata 1920
gctcatgatg tgatgggatg ttggttgagg aagatttttt ctggagcaac tgttggggta 1980
actgagagag ctgaattgga gaagagagat cttttgggta agactagagc catagagatg 2040
gacctgagtt ccagctctgc cttgcagttt ggacaagaag tggcaaatcg agttttgagt 2100
gttattaaag gtgttttggg tgtgtagggt agaaaagtgg ggagctccag attgcaagga 2160
attatagaat atattaaggt ggtgtttggg ttgggaagaga gactagagga aatgaaagta 2220
atttaaatc tcgttttttt tta 2243

```

```

SEQ ID NO: 15      moltype = DNA length = 1085
FEATURE            Location/Qualifiers
source              1..1085
                    mol_type = genomic DNA
                    organism = Citrus sinensis

```

```

SEQUENCE: 15
tgcgaaaggg ggaagtcaaa agtggttattc gttgtgtgtct atggctatga gttcgccact 60
ttgggtagcc ccaacttttc tttctcgctc atcgtcaaat atatcgagat caattactta 120
caaacgttgc aggtcttcgg caagtaactt tttctcacaa tcagacatgg agcagccgct 180
gatgaagggg cagaagctga tggaaattccc tcacctgacg gcggcgacaca aagatctgat 240
ggttagctta atctcagcct tggagactag gcttgactct catcttcttg cttcttcggt 300
agtcctcct gatgttgagt tttatcagaa tgaccaaggc acttctcagg gttctcttca 360
tatcagacgt ggccttccct cctctcatat tgatttcgta ttgcaaggt ggcttcaact 420
gaaggtacca acgggaagtg ccatgaacat aaccaatctt caagcttacc taaaatcatc 480
aacgatgta ccacattttc aattcgagct tgtccaatgc agccccacat atttcattct 540
cttctcatag ataactccta gaaaagacct tgttctatac ccaaatacc tcaaaacatt 600
ttacgaagaa gctcagcttg aaacattgag acagagactt gagcaagtcc cagaaaccaa 660
accctacttg agttcctctc tttactttcg tgggtgtggt tccccaaact ggattttggt 720
cagcataaaa tgtgaggaag ttggtggaac ggtatcgctg gaagagatta tacgtgaaca 780
tgtgagccct atagctcatg atgtgatggg gatttggttg gagaagtatt tttctggagc 840
aactgttggg gtaactgaga gagctgaatt ggagaagaga gatcttttgg ttaagactag 900
agccatagag atggacctga gttccagctc gcccttcgag tttggacaag aagtggcaaa 960
tcgagttttg agtgttatta aagggttttt ggggtgtgag ggtagaaaaa tggggagctc 1020
cagattgcaa ggaattatag aatatattaa ggtggtgttt ggttgggaag agagactaga 1080
ggaaa 1085

```

```

SEQ ID NO: 16      moltype = DNA length = 3474
FEATURE            Location/Qualifiers
source              1..3474
                    mol_type = genomic DNA
                    organism = Citrus sinensis

```

```

SEQUENCE: 16
agaaaacaac aaataaaaat tgcgtcattc cttcatgttg ggcaatttgc agcctcaagc 60
tcacggttcc ctgatcatga ctcatggcat gactcgtgtt cgtttcacaa ctatcaataa 120

```

-continued

```

ttgectcaag ttttaattctt gcttgatctc tagtttgtgt tgagaacggt gctgcatgca 180
gcoatggaag ctcttagact ctctcgggtt tcaccttttt gtaacgctac attcaaaactt 240
gagtagacaata agaccactt cactaagccc aagttcttaa gctttcagtt cagttccctg 300
tccactttat catctttctc atcaaaacca taaaaatct tcaccacctt atccaccatcg 360
tcacaagttt caactgaagc cacagaccca ccagaaagag agcttgaaac taactcacia 420
gaggagaaat ttgattgggt ctacacagtg tatccattga tgccggtgtg tgatttggac 480
aagagagtcct ctcatgcaaa gaaagtgttg gggcttgatg tcgtggtttg gtgggacagg 540
aatgagaatg aatggagagt ctttgctgat gcttgctctc acagattggc tcctttgtca 600
gaaggagaagaa ttgatcaatg ggggaaggctt cagtgtgtgt atcatggctg gtgcttcagt 660
ggctcgggtg actgtaaatat tatccctcag gctccccag acggccccc ggtaacatct 720
attgcttcac ttttgcatca catcatatg ttttattatg ttttggtttg ctatagatgc 780
ttttcggtt gtctcgctta tgtgtagagt gagatgctat agttttgtcc aattagaatc 840
tcttctaggc atagagtttg tttgtttata attgttgctg ttcagcgctt tttttgccc 900
aaagctatcc cggagagact tacagactgg aagtaataaa atagggtcaag ggtctaaattc 960
caaggggata cggttaattc tgaagagaca gtatgtaatt aaatggttgg taaagccgcc 1020
acacctctca tcttcagttt aattgtgaag gaacaatgaa agataagggg aatacatgaa 1080
ttatgattcc attgcttgct tgataaaatt ttgttttgat tttctgcac tctagttttg 1140
tttttttgtt cttgcatgca gttgtgtgta agatttctac tttgctgggt ggctgattgc 1200
tttaattttt accaacttat ttgtcaaaat atgtaggctc acacattcga gaaagcatgt 1260
gcagcagttt atccaagtac cgtccagcat gacatcgat gggtttggcc aaatattgct 1320
ctcgggtaca aggatattat caagactaag aaacctctc acatcccgga actagatgac 1380
ccgtcattca caaaattggt tgggaacaga gatatacctt atgggtatgc aatgatccag 1440
tcttccattt taatttgagc gagttagcac acgaaattct agattgaatt tgtactatac 1500
tgatatttgg caagagtcac taagaaaact agttgaactt aagtgtaatg acaacctggt 1560
caatgcaagt taacaacttt ttctaattgc aagtaagtgc cattttcagt tatgaggtct 1620
tattggaaaaa tcttatggac cctgctcatg ttccatagc acattacgga ttgatgcgta 1680
caaggaaacc caaaggttga taaattccta gataaacctt gttctacggt taagtcat 1740
atattctctt cgtgaataaa acctattagt taataactaaa tttctttac agtgaagctc 1800
gatagagaag ggggaagacc agttgaaatg agtgtaata aaatagacat aaatggtttc 1860
attggaagac aggaagggtt aagtagcaaa tttttggcac cttgtatctt tttgtcttat 1920
actgatctta tgaaggatca agaaaatgga tctgcatcat cagcaggagc cgaagggtta 1980
aaggtaataa tgtttcatta tgcacagctg acagtagaat tttgcactct tttcagttct 2040
tattctatgc tgtttgggtt ttcttttatc catatacatt tcggttggtg cagaagctgg 2100
agcaacaaag agcagctctg atttttattt gtgttccagt tagtccctgt cagcagat 2160
taatatgggc gttcccaaga aactttcaaa cttggataga caaagttggt ccgcggtgga 2220
tatttcatat tggacagata ctaattcttg attcagattt atacctgct cactgtgagg 2280
tgattctctg ctctgtatgc taaaataaatt ttaacctaaa gttttttgaa aatgctgagt 2340
ggattggcgt tcaagctctac tgcagatcta ggccctcaaac tcagatttga gactcgtcga 2400
ccatttcatt ttactagatt ggttgtgaca ccatccaaaa agcatatttt tgtttttctt 2460
ctcaatagaa ggtattcact taaatactac tctctacaaa tagaagaatt taatagattc 2520
tttcttggtt ctggctgtga attacctgac ttctgtttct ctctgcagga gcgaaagata 2580
atggatgttg gccctgctaa ttggcagaaa gctgtttttg tgccaacaaa agccgatgcc 2640
ctggtagttg gtttcaggag gtggttaaaa aaatatgccg gtggccaatt caattgggga 2700
gggaaattca atgcagctct tccaccaaca ccgcccaggg aacagctcat ggacagggtat 2760
ctgggcttca cttttataaa ctggaacccc ttaacacat gtcatgggat ttacagtag 2820
ttccagttt tgaactctat ataaagagca gaggtttatg gatattttaa agaagcatta 2880
tttgtttctt caagcttgag aaacattttc tttgtgttt tgccaaattt cttatgcaat 2940
gcttgaaaaa cggaacactc gtgaagtttg ttgtgacccc attaaaacat aagatttcgt 3000
acttaattaa aggaatttta ttcaattttg caggtaactg tctcatgttg tgaattgcaa 3060
aagttgcaat gcgacacaca agagtctcag cgcacttgag gtcacgtctc aagtcgtctc 3120
cattgcttca attgggattg ttgctacaac caagcagaat gccatgtcaa tggctacaag 3180
aaactacaatc atctcatttg cagtaattctg ctttgcgctc tcaaaatggt tgtctcactt 3240
catctacaaa acctttcatt atcatgacta caatcatgcc cttcgctgag cttagcattt 3300
aacgtcgaaa attagaatat gtaaatacaa cttatttttc tgcactgaaa tactggaatg 3360
tagcttgat gcaaacattt tgatcaagtg aaattagaaa gtgcagttgt aatagaaaac 3420
atataattat catcaccatt agcagttgta attgtaagta cattatcatc acca 3474

```

```

SEQ ID NO: 17      moltype = DNA length = 2093
FEATURE            Location/Qualifiers
source              1..2093
                    mol_type = genomic DNA
                    organism = Citrus sinensis

```

```

SEQUENCE: 17
atgcttcaga aaacaacaaa taaaaattgc gtcattcctt catgttgggc aatttgcagc 60
tcaagctca cgggtccctg atcatgactc atggcatgac tcgtgttcgt ttcacaacta 120
tcaataattg cctcaagttt taatcttgct tgatctctag tttgtgttga gaacggtgct 180
gcatgcagcc atggaagctc tttagactctc ctcggtttca ccttttttga acgctacatt 240
caaaacttgag tacaataaga cccacttcac taagcccaag ttcttaagct ttcagttcag 300
ttccctgtcc actttatcat ctttctcacc aaaaccatac aaaatcttca ccaccttatc 360
accatcgta caagtttcaa ctgaagccac agaccacca gaaagagagc ttgaaactaa 420
ctcaacagag gagaattttg attggttctc acagtgggat ccattgatgc cgggtgttga 480
tttgacaag agagtccttc atgcaaaagaa agtgttgggg cttgatgtcg tggtttggg 540
ggacaggaat gagaatgaat ggagagctct tgcgtatgct tgcctcaca gattggtctc 600
tttgtcagaa ggaagaattg atcaatgggg aaggcttcag tgtgtgtatc atggctgggtg 660
ttcagtggc tcgggtgact gtaaaattat cctcaggct cccccagacg gcccccgggt 720
ccacacattc gagaagcat gtgcagcagt ttatccaagt accgtccagc atgacatcgt 780

```

-continued

atgggttttg	ccaaatattg	ctcctcggtg	caaggatatt	atcaagacta	agaacctcc	840
tcacatccc	gaactagatg	accgtcatt	cacaaaattg	tttgggaaca	gagatatacc	900
ttatggttat	gaggtcttat	tggaaaatct	tatggaccct	gctcatgttc	catatgcaca	960
ttacggattg	atgcgtacaa	ggaaccccaa	agtgaagctc	gatagagaag	ggggaagacc	1020
agttgaaatg	agtggtcaata	aaatagacat	aaatggtttc	attggaaagc	aggagtgggg	1080
aagtagcaaa	tttttggcac	cttgatcttt	ttttgcttat	actgatctta	tgaagatca	1140
agaaaatgga	tctgcatcat	cagcaggagc	cgaaaagaag	ctggagcaac	aaagagcagc	1200
tctgattttt	atttgtgttc	cagttagtc	tggtcacagc	agattaatat	gggcgttccc	1260
aagaaacttt	caaacttgga	tagacaaagt	tggtccgcgg	tggatatctc	atattggaca	1320
gaatctaatt	cttgattcag	atttatacct	gcttcacgtt	gaggagcgaa	agataatgga	1380
tgttggccct	gctaattggc	agaaaagctg	ttttgtgcca	acaaaagccg	atgccctggg	1440
agttgggttc	aggaggtggg	taaaaaata	tgcgggtggc	caattcaatt	ggggagggaa	1500
attcaatgcg	actcttcac	caacaccgcc	caggaacacg	ctcatggaca	ggtactgggc	1560
tcatgtggg	aattgcacaa	gttgcaatgc	ggcacacaag	agtctcagcg	cacttgaggt	1620
cacgtacaaa	gtcgtctcca	ttgcttcaat	tgggattggt	gctacaccca	agcagaatgc	1680
catgtcaatg	gtcacaagaa	ctacaatcat	ctcatttgca	gtaatctgct	ttcgggcttc	1740
aaaaatgggtg	tctcacttca	tctacaaaac	ctttcattat	catgactaca	atcatgccct	1800
tcgctgagct	tagcatttaa	cgtcgaaaat	tagaataatg	aaatacaact	tatttttctg	1860
tacgtaaata	ctggaatgta	gcttgtatgc	aaacattttg	atcaagtga	attagaaagt	1920
gcagttgtaa	tagaaaacat	ataattatca	tcaccattag	cagttgtaat	tgtaagtaca	1980
ttatcatcac	cattaacaat	tgcagagaaa	aacgtataca	caactgaacc	acggcttcag	2040
aatgaacaca	agaagaacat	ttttactagt	tttcacaatg	gtatctagtg	atg	2093

SEQ ID NO: 18 moltype = DNA length = 4930
 FEATURE Location/Qualifiers
 source 1..4930
 mol_type = genomic DNA
 organism = Citrus sinensis

SEQUENCE: 18

ccttgcttga	tctcgatttt	tgtgcagagt	aaggtgttgc	atgcagccat	ggaagctctc	60
ttactctcct	cagttttcac	attttataac	actccattaa	aacttaagta	caacagaacc	120
cacttccactg	ctaagcccaa	gctcttaagc	ttccacttca	gtccactatc	cactttatca	180
tctttctcat	caaaaccatc	caaaactctc	accaccttat	caccatcatc	tcaagtttca	240
actgaagcca	cagaccacac	agagacagag	cctgaaacta	actcacaaaga	ggaaaaattt	300
gattgggtct	cacagtggtg	tcatttgatg	ccggtgtgtg	atttggaaca	gagagtcccc	360
catgcaaaaga	aagtgttggg	gcttgatgtc	gtggtttggg	gggacaggaa	tgagaatgaa	420
tgagagtgct	ttgcccagtc	ttgtcctcac	agattggctc	ctttgtcaga	aggaaagaatt	480
gatcaatggg	gacgggttca	gtgtccgtat	catggctggg	gcttcagtgg	ctcgggtgac	540
tgtaaattta	tcctccaggc	tcctccagac	ggccccccgg	taacatcgat	tgcttcactt	600
ttgcatcaca	ttcatatggt	ttattatttt	ttatgctgct	atgtatgctt	ttcgggtttt	660
cttgattacg	tctagagtga	gatgctactt	ccttgctata	ttttggtcta	attagattcc	720
ctgaaggact	aaaagactgg	aagtaataaa	ataggtcaag	ggcctaattc	caatgggata	780
cgtttaattc	tgaagaactc	gtagctaatt	taatgtttga	taaagccacc	acacctctcg	840
tcttcagttt	gattgtgagg	aaacaacgaa	agataagcag	aatacatgag	ttatgattcc	900
atttcctgct	tgataaaatt	ttgtagtgat	ttttctgcaa	tctagtattg	attttcggtc	960
ttgcatgcag	aagttgttaa	gaattctact	ttgctgggtg	gctgattgct	ttaattttta	1020
ctgacttatt	tatcaaaata	cgcagggtca	cacatccaa	aaagcatgtg	cagctgttta	1080
tccaagtgc	gtacagaagt	gcactctatg	gttttggcca	gatattgtct	ctcagtgcaa	1140
ggatattatc	aagactaaga	aacctcctca	catcccggaa	ctcgtatgac	cgctatttac	1200
aaaaatgttt	ggaagcaggg	atgttcctta	tgggtatgca	gtgatccggt	tttccatttt	1260
taatttgaaa	ttccagtttg	aatttgtacc	atatgtatat	ttggcaacaa	tcattaagaa	1320
aaactagtga	actcaagtgt	aatgacagcc	gggtaaaatg	aagttaacag	ctttttctaa	1380
gtgtaagtta	attgcatttt	cagatatgag	gtcctaattg	aaaaatctat	ggatcctgct	1440
catcttacat	atgcacatta	cggaaatgat	cgtacgagga	aacccaaagg	ttgataaatt	1500
cctggataaa	ccttgtttta	cgtttacgtc	aactatatta	ataaaaccca	ttagctaattg	1560
ctaaattttc	tttacagtga	tgctcgatag	agaaggggga	agaccaatca	aaataagttt	1620
cgagaaaata	gacataaagt	gtttcattgc	aaagcaggat	tcggaaaagt	ccaaattttt	1680
ggcaccttgt	gtctttgttg	tttattttga	tcttctggag	aatcaggaaa	atggatctgc	1740
atcatcggga	ggagccgaag	aggtaagagt	caatatgttt	cgtaatgtct	agctgacgga	1800
tagaattttg	cactcttttc	acccccctat	ctatgttgct	cccgcatgtt	ttgtccatat	1860
aaatttcgat	cgggtgcagaa	gctgaagcaa	cgaagagtag	ctatgatttt	tatttgtgct	1920
ccagtaagtc	cgggtaacag	cagagtaatc	tgggccttcc	caagaaactt	ccagatttgg	1980
atagacaaa	ttgttcgcgg	gtggatattt	catattggac	agaacctaat	tctggattca	2040
gatttatgcc	tgtctcacgt	tgaggtgatt	cctatctctg	tatgctaaaa	taattttaat	2100
ctaaagtgtt	tggaaactgc	cgggtggatt	ggcggttaaa	tggagtgtca	ggtagcattt	2160
attttatttg	tctgaaattt	caatgagtga	ataaatctct	ctctctctct	ctctctctct	2220
ctctctatat	atatatatat	atnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	2280
atatatgaat	aacatgttta	tttttcttat	tttgatatat	gaataaagt	gttattaaaa	2340
aaaaaaatga	catcttctat	atctatagag	aagccaattt	cgttgtagat	tatatggcga	2400
acctcacttt	ttctctctct	tttggccttc	ttgtgtacac	tgccccacct	ctagggtgta	2460
ggctctcttt	gcttcaatac	ttttacgggg	tttctcacc	cgttctgtt	cttctttagc	2520
tctttttaga	gccccagttc	aataaaataa	taaaataata	aaatatgata	ttttgaaata	2580
tataacatca	taaatattaa	caatacttat	attttttttt	attaaacaca	tgacaagttg	2640
taataaaatt	atttgattag	taaaaaaaat	ttaataatgt	tttccttaac	agagcacgaa	2700
aaataaaaata	tgtatctaac	acacaatata	tgatacttat	gttggttatt	ttagtataat	2760
atattgattt	ttttaattac	atatgtcaat	taaagattgt	ttaatcattt	gaattaaatt	2820

-continued

taatgtatta	aataaaatc	aaaatctatt	ataaaaatgg	caaatataaa	taccaagaga	2880
taggtccagt	ggcggattta	aaaaataat	ttaccggggg	ctaaattaga	taataataaa	2940
atatgaaaaa	ttaaaatfff	aaatatataa	aatatttgct	agtacattta	taattattct	3000
ctgcgcattt	ttatatfita	aaaatgttat	acaattgact	cattatcaat	attattaaat	3060
ttattgttcc	gtatataaac	aatcaagttc	aaatttaaat	tttatggaga	tttataattt	3120
atagtgattt	ttttaataaa	ttttttatgg	gggttaatta	aaaaactaaa	atatttttct	3180
tttttaattt	taatttttta	ttttggcagt	gggggctcaa	gccccacta	gtcacatgct	3240
agatccgccc	ctggataggt	ctattataaa	acacactcat	ctaaagtctg	attatgttca	3300
aaattctaata	tcacaatata	tatatatata	tagttgaata	catgtatata	accaaaaaaa	3360
ggtaaaaaaa	aaataataaa	attatttata	tgatagggaa	taaatfittt	taatatgaat	3420
attaacgcaga	gagaatatta	aaattacatt	taaatatttt	acaattttaga	taaaagttyt	3480
agtttttaac	tttttgattt	tagatctaaa	catcttataa	ttagatataa	gtaataaaaa	3540
aaagacagta	aatatcataa	aacaattaag	attcaaaactg	agaaaaaaa	aatcaaaaaa	3600
cccttgctcc	tccagtctag	gcctcaaaact	cagatattag	actcgtcaac	catttcattt	3660
tactagattg	attgtgacac	catccgaaaa	gcataatttt	gtttttcttc	tcaatagaaa	3720
gtgttgatta	aaataacact	ttgttgattt	aaataactact	ttcctacgat	aaaagaaatg	3780
aatagctttc	ccgttgtttc	tagccgcgaa	taaaattctc	tgcaacatat	tgttacctga	3840
ctttggtttc	tccctgcagg	agcgaagat	aatggctggt	ggccctgcta	attggcagaa	3900
agcttgtttt	gtgccaacaa	aatccgataa	cctggtagtt	ggtttcagga	tgtggttaaa	3960
aaaaatattc	ggtggccaat	tcaattgggg	aggaaaaatc	gatgcaactc	ttccaccaac	4020
actgccaaga	gaacagctca	tggaacagta	tttgggcttc	acttcaataa	acttgaatcc	4080
cttaacgcgc	tgtagtctct	gggtttgaag	aaagagcagg	gatttatgga	tttttaaaag	4140
aaagcattatt	tatttcttca	ggcttgggaa	acattttctc	ttgtgttttt	gccaaaaatc	4200
ttatgcaatg	ccttgaatac	gggaacaata	taaagtttgc	tgtgacccca	ttaaaacgta	4260
tgatcttggt	cttaattaaa	ggaactttat	tcaaattttc	aggtaactgt	ctcatgtggt	4320
gaactgcaaaa	attgtgcaatg	ctgcacacaa	gagttctcaat	gcacttgagg	tcatactgca	4380
agtcgtctct	gttggttccag	ttgggattgt	tgtgcacacc	aagcagaacg	ccatgtcaat	4440
ggctacaaga	gctacgatct	tgtcatttgc	agtaactctgc	tttgcagctt	caaaatggtt	4500
gtctcacttt	gtctacaaaa	cttttcatta	tcatgactac	aatcatgctc	ttcgtcaagt	4560
tagactctgg	taatactgta	acttttaaaa	taattgctat	tacttatagc	gttgaaataa	4620
tctgccgtga	gcaaaaatcaa	tttaaaaact	gataaaatftt	attttaaaaa	atattgtgtc	4680
ttaaaaaata	gtcgtatata	agtttacatt	ggtcataatg	ttgaataaaa	tttaggaaaa	4740
ttatcattcg	tgtaccctaa	agatgcactt	ttatcaaaaa	tattacaaca	ctttcaaggt	4800
gtatcactca	tccaccaaaa	aataccaaaa	tatatctacc	caccactatt	ccgttagcca	4860
ccgtttgcaa	actaacagaa	ttgttgcgaa	atgacaaaaa	tgccctttaa	actaaaaaaa	4920
aacgcaaaaa						4930

SEQ ID NO: 19 moltype = DNA length = 2237
 FEATURE Location/Qualifiers
 source 1..2237
 mol_type = genomic DNA
 organism = Citrus sinensis

SEQUENCE: 19

cacaaacata	agagacgcat	taattttatgt	tggacttttc	agcgtcacgg	tcagagtcca	60
ctgatctgat	ctcatgtggc	gcacgttcat	tcgcaattag	ctataattat	ctcgagttct	120
aaaccttgct	gatctcgatt	tttctgcaga	gtaagggtgt	gcatgcagcc	atggaagctc	180
tcttactctc	ctcagtttca	ccattttata	acactccatt	aaaacttaag	tacaacagaa	240
cccacttcac	tgttaagccc	aagctcttaa	gcttccactt	cagtccacta	tccactttat	300
catctttctc	atcaaaaacca	tccaaactct	tcaccacctt	atcaccatca	tctcaagttt	360
caactgaagc	cacagaccca	ccagagacag	agcctgaaac	taactcacia	gaggaaaaat	420
ttgattgggt	ctcacagtgg	tatccattga	tgcgggtgtg	tgatttggac	aagagagtcc	480
cccatgcaaa	gaaagtgtgt	gggcttgatg	tcgtgggttg	gtgggacagg	aatgagaatg	540
aatggagagt	ctttgcgat	gcttgtcttc	acagattggc	tcctttgtca	gaaggagaaa	600
ttgatcaatg	gggacggcct	cagtgctccg	atcatggctg	gtgcttcagt	ggctcgggtg	660
actgtaaatt	tatccctcag	gctcccccag	acggccccc	gggtccacaca	tccaagaaag	720
catgtgcagc	tgtttatcca	agtgccgtac	agaatggcat	cctatggttt	tggccagata	780
ttgtcctcca	gtgcaaggat	attatcaaga	ctaagaaacc	tcctcacatc	ccggaactcg	840
atgacccgtc	atttacaata	atgtttggaa	gcagggatgt	tccttatgga	tatgagggtc	900
taatggaaaa	tcttatggat	cctgctcctc	ttacatatgc	acattacgga	atgatgcgta	960
cgaggaaacc	caaagtgatg	ctcगतagag	aagggggaag	accaatcaaa	ataagtttcg	1020
agaaaaatga	cataaatggt	ttcattgcaa	agcaggattc	ggaaagtgcc	aaattttttg	1080
caccttgtgt	ctttgtgtgt	tattttgatc	ttctggagaa	tcaggaaaaa	ggatctgcat	1140
catcgggagg	agccgaagag	aagctgaagc	aacgaagagt	agctatgatt	tttattttgt	1200
ctccagtaag	tccgggtaac	agcagagtaa	tctgggcctt	cccaagaaac	ttccagattt	1260
ggatagacaa	agttgttccg	cgggtgatat	ttcatattgg	acagaaccta	attctggatt	1320
cagatttatg	cctgcttcac	gttgaggagc	gaaagataat	ggctgttggt	cctgctaatt	1380
ggcagaaaag	tgtttttgtg	ccaacaaaaa	ccgataacct	ggtagttggt	ttcaggatgt	1440
ggttaaaaaa	atatcccggt	ggccaattca	attggggagg	aaaattcgat	gcaactcttc	1500
caccaaacact	gccaagagaa	cagctcatgg	acaggtactg	gtctcatgtg	gtgaactgca	1560
aaagttgcaa	tgtcgcacac	aagagtctca	atgcacttga	ggctacactg	caagtctgct	1620
ctgtgttttc	agttgggatt	gttctgtcaa	ccaagcagaa	cgccatgtca	atgggtacaa	1680
gagctacgat	cgtgtcattt	gcagtaaatc	gctttgcagc	ttcaaaatgg	ttgtctcact	1740
ttgtctacaa	aacctttcat	tatcatgact	acaatcatgc	tcttcgctaa	gttttagcatt	1800
ggtaataactg	taacttttaa	aataattgct	attacttata	gcgttgaat	aatctgccgt	1860
gagcaaaatc	aatttaaaaa	ctgataaaat	ttatttttaa	aaataattgt	tcttaaaaaa	1920
tagtctgata	taagtttaca	ttggtcataa	tgttgaataa	aatttaggaa	aattatcatt	1980

-continued

cgtgtaccct	aaagatgcac	ttttatcaaa	catattacaa	cactttcaag	gtgtatcact	2040
catccaccca	aaaataccaa	aatatatcta	cccaccacta	tcccgtagc	caccgtttgc	2100
aaactaacag	aattgttgcg	aaatgacaaa	tatgccctta	aaactaaaaa	aaaacgcaaa	2160
aagacataaa	tacccacaaa	accaagcaaa	gaaaaataga	tccgttagag	acaataacag	2220
ttgtatTTTT	tggtcat					2237

SEQ ID NO: 20 moltype = DNA length = 4421
 FEATURE Location/Qualifiers
 source 1..4421
 mol_type = genomic DNA
 organism = Citrus sinensis

SEQUENCE: 20

ccggtttttc	ttcttttttc	cgtttttctga	attctggaag	cgtgaatata	gacagacagc	60
tgggagaatt	ggaggggcaa	attgcaaaact	tgcacgcaaa	gtcctcttct	ttttctgtct	120
ctctctcgaa	agctcacttt	atttttcgctt	cactagaatt	ggaaaaatca	tacaaaaatt	180
ttcattaaac	tcaaaagcaa	aggacatggc	gctacttctt	tctactactg	ctaatacac	240
cacatcacca	agaaaaaccc	ttccattttt	ggccacagga	accccgaaag	gacaaatcac	300
ggtaaaaagc	ttgcaaaaga	gaagcaagaa	tttgtctcca	ctacgagtgg	cagctcctcc	360
ttcagacctc	gcagcatcag	atgaagaaac	gatgagaaaa	gatgagaaag	aagattatgg	420
atcattggtc	gatgatgagt	atggtaaaaga	gagttcggat	tctaagtttt	cttggaggga	480
tcattgggtac	ccagtttctt	tagttgaaga	tttggacccg	aacttgcccta	caccgtttca	540
gcttcttggg	agagacttag	ttctttgggt	tgataacaat	tctaataaat	gggttgcat	600
tgatgataaa	tgccctcata	gactcgcctc	attatcggtg	atctattcta	ttgttctctc	660
tgatctctgt	aattttgtga	tcgaggtgaa	acatagaaga	taaaattgta	atttgctcat	720
tttttatTTA	attttcgata	ttatagtggt	aatggaagaa	gtaatatctg	ttgttgtatc	780
ttattttctt	tttaagaaat	gaaatttgaa	tggagggttt	tctaggatct	agagtgaagg	840
atatgaaatc	taattttcaa	attcatatct	tgttctgttc	ttgatcagtt	tatccaagga	900
aagctcacca	tatagtccat	tgaatgacca	attttttgtc	gttttcgata	tagaatccga	960
aattcagaat	tttgggagct	gtttgtttga	ccattgatgt	ttaactctgg	tagctaaagt	1020
gggaattatg	tttcttagat	catgtgattg	atgggttccct	taaaatttta	taggaagggc	1080
ggatcgatga	aaatgggcat	ttgcagtggt	cataccatgg	atggtctttt	gacgggtgtg	1140
gatcttgca	tcgcattccc	caagcagcat	ctgaagggtcc	tgaagctcgt	gcaattcagt	1200
ctcccagagc	gtgtgctacg	aggtttcccta	caatgggtgc	tcaaggtctg	ctattcggtt	1260
ggccagatga	gaatgggtcag	gaaagagcca	atgccaccaa	gccaccaatg	taattgactc	1320
tattctcttc	ttctgttttt	agattatgca	ttgactgaat	taccgacttt	gccgggttcat	1380
taaacctgtg	tggtaatggt	taggttgccct	gatgactttg	acaaacctga	gttctcatcg	1440
gtcacaaatc	agcgggcatc	attttatggc	tatgacactc	tcatggaaaa	tgtctcagat	1500
cttcccccac	ttgatTTTgc	acatcacaa	gtacatactg	aatttcaatg	gtagtgtcgc	1560
tgggtgcaag	atggaaatgt	catttgaaca	atagctggtt	gggttaagat	aattttgata	1620
tgaattaaat	atggtatgca	gctcttggtc	gatatgtctc	attaactaaa	ctagaaacac	1680
agttatcgtc	tcttgtaag	ttctttgggt	ttcagtagtg	tcagggaagt	cctgcaggta	1740
ttgttttctt	agtggactga	ttgaacacat	agtggcgca	ctttttgttt	acttggccat	1800
gttggtgctt	ctctctcaat	cacttggaa	ctcgtaacat	cctagtactc	atatacaggt	1860
catgttggtc	tcattcatgg	atatgtcctt	gaacaaggag	taaagtagtg	aattctatag	1920
ccctatctcc	tcatttagat	gctataggaa	tgtgacacta	attgctctct	ctgtggaac	1980
aaatctccat	gtactctatt	gagagacatc	gtttgcaggt	taccggcaga	agggacaggg	2040
ccaagccttt	accgtttaag	ttggagtcct	gtggacattg	gggatttgct	ggagccaatg	2100
atggaaaccc	aaggataagt	gctaaatttg	ttgtcctctg	ttattacatg	aacaagtaag	2160
ttctccactg	cttggacagc	caaaattaat	acgtctagat	agactatctc	cttctcgtct	2220
tatgctctct	cagtgtactc	taaaattggt	gattttttca	agaactggag	atttgaataa	2280
tttcccaaac	ttacttcaaa	agcgtttagt	attcctgatg	aaaacattgt	gtgataaagt	2340
gtgaacctta	accatccaat	atttctcctc	accaagactc	aagttcattt	tgtaagcca	2400
atcaatggag	gcttgtatTT	caaagttaga	ggttgacttt	tggttaactg	gaggatttat	2460
gcacaaacgt	agaactgagt	tgTTTTgaca	tcagctgtag	ttcaaggaa	ccgatgaaac	2520
atcatttcat	gtcttatttg	tgTTTTaaat	tgaatacaac	tgtagacgaa	cttctctctg	2580
agttagggtg	tagtttttagt	tgatttagta	ctatatTTgt	ttatcagttc	attctgcttc	2640
cgaaacttaa	ctatccgtgt	attgcttttt	ttccaggata	gagatagata	caaaacttcc	2700
tgtagttggt	gataaaaaat	ggataatatg	gatttgttcc	ttcaacgtac	caatggcacc	2760
agggaaaaac	cgctcaattg	tttgtagtgc	acgaaaactc	ttccagttca	ctatgccagg	2820
acctgcctgg	tggcaggtaa	gaacatctca	gttgtgttcc	tgatgaaatt	tgtataatgt	2880
gacacttgca	tggaaacttt	agctcaaaaga	gaacactcaa	atgctgttga	catgttcaag	2940
catacggaaa	gtttatcttg	ttaacacgct	tgtgctagtg	tgacacacac	tggttttagc	3000
ttccaaattc	ggtagttgat	tgttcttatc	ttggatacca	tactgtccca	tttgggaagg	3060
ggaagtTTtg	gatgtctggc	tgaatgtgg	agagaaaact	ccttgatttg	gagaagaaaa	3120
cttagttgag	atgtatgctt	aaaagttttc	tccctgtttg	agggcctttt	cttagatatt	3180
ggcatccgtg	tcttgtaaac	tttctaata	atattattca	tatttttcat	gttgatgcca	3240
aagttccaat	ctaaggggtt	tgtgtgcaga	tttgataaga	ctttgttgag	ctgtattggt	3300
gatgattttc	ctccacatga	aatttagttt	ttacggaggt	tatgtcctat	ctccacctta	3360
gaaaactaga	aatttaatat	ttttacttcg	taaattgggt	aattatctac	aaaatggctt	3420
cgagttttac	caggttggtt	ctagatggca	cgagcattgg	acttcaaaata	aggtctatga	3480
tggagacatg	atgtctctcc	aaggtcaaga	gaagatcttt	ctttcaaaat	tgatggaagg	3540
ttctgaagat	gttaacaagg	agtaacacaa	aattacgttt	acacccacac	aggcagatcg	3600
gcttggtgtg	gcatttagaa	attggctgag	gcgacacggc	aacagccaac	ctgaatgggt	3660
cggcttcagc	agccaacaac	cttccctctc	aacagctctg	tcgaaatgtc	aggtacgaaa	3720
cccttctccc	attatttcaa	gaaataaaat	atgaatatgt	ggattgaatt	tagattttta	3780
aagtagatgg	attcataatt	tgtgttaagt	gggacaatgt	cacatttata	ataagctcga	3840

-continued

```

ttaatatgtt attacacctt ctgggtttga tttcttcctt acgttttagt tttctgttg 3900
ttgggttttg tttcagatgc tcgatagatt tgagcagcac accctcaagt gttcatcatg 3960
tagagaagct tattcagcat tccagacggg ccaaaagtgt ctcattggcg cgaccgttgc 4020
attctgcgca acagctggga ttccttcaga ttgcaatca cggattgttt tggetgggct 4080
tgcactagtg agcgtgcct tggcttatgc ttgcatgaa ctacaaaaga attttgtgtt 4140
tgttgattat gtgcatgctg aaatcgatta gagaggagtg agatatgctg ctgaaagaat 4200
caatgtgtcc agaggtataa gaaagatatg gtgataaatc ttgtcaaaat ttgcgagttt 4260
gtatatatct attagataga aatcagtggt atagctaagc ttagagtggt tctcagtat 4320
tccctttgtt tttgtgcca atgaatgtca tcagataaat atgtgcagac atgcacccaa 4380
attcatgagt aaatggatta aacaatatat actttgtcaa a 4421

```

```

SEQ ID NO: 21      moltype = DNA length = 2334
FEATURE            Location/Qualifiers
source              1..2334
                    mol_type = genomic DNA
                    organism = Citrus sinensis

```

```

SEQUENCE: 21
cacgaaacac gccattgcca aagggcatta cgggaatttt gtgccagaaa atgtgacagt 60
aaatgaccag cagccccttg actttctatt ctgttagaac ctttaattgg tgagaccaat 120
catacaaaact aaaccacaaa gtacttgatc gcaaatcgga caatcagaaa aatcgagcta 180
acgggttttt cttctttttt cggttttctg aattctggaa gcgtgaatac agacagacag 240
ctggggagaat tggaggggga aattgcaaac ttgcaacgaa agtcctcttc tttttctgtc 300
tctctctcga aagctcactt tattttcgct tcactagaat tggaaaaatc atacaaaaat 360
tttcattaaa ctcaaaagcaa aaggacatgg cgctacttct ttctactact gctaatatca 420
ccacatcacc aagaaaaacc ctccatttt tggccacagg aaccccgagg cgacaaatca 480
cggtaaaaaa cttgcaaaag agaagcaaga atttgtctcc actacgagtg gcagctcctc 540
cttcagaccc tgcagcatca gatgaagaaa cgatgagaaa agatgagaaa gaagattatg 600
gatcattggt cgatgatgag tatggttaaag agagtccgga ttctaagtgt tcttgagggg 660
atcatttggt atcttggtt ttagttgaag atttggacc gaacttgctt acaccgtttc 720
agcttcttgg gagagactta gttctttggt ttgataacaa ttctaataaa tgggttgcat 780
ttgatgataa atgccctcat agactcgccc cattatcgga agggcggatc gatgaaaatg 840
ggcattttga gtgttcatac catgggatgt cttttgacgg gtgtggatct tgcaactgca 900
ttccccaagc agcatctgaa ggtcctgaag ctctgccaat tcagtctccc agagcgtgtg 960
ctacgaggtt tcttacaagt gtgtctcaag gtctgtctatt cgtttggcca gatgagaatg 1020
gtcaggaaaag agccaatgcc accaagccac caatgttgcc tgatgacttt gacaaaacctg 1080
agttctcatc ggtcacaaat cagcgggatc tattttatgg ctatgacact ctcatggaaa 1140
atgtctcaga tctctccac attgattttg cacatcaciaa ggttaccggc agaagggaca 1200
gggccaagcc tttaccgttt aagttggagt ctagtggaac ttggggattt gctggagcca 1260
atgatggaaa cccaaggata agtctgaaat ttgttgctcc ttgttattac atgaacaaga 1320
tagagataga tacaaaaact ctgtagtgtg gtgataaaaa atggataata tggatttgtt 1380
cctcaacgtt accaatggca ccagggaataa cccgctcaat tgttttagt gcacgaaact 1440
tcttcaggtt cactatgcca ggaacctgct ggtggcagg gtgtcctaga tggcacgagc 1500
attggacttc aataaaggtc tatgatggag acatgattgt cctccaaggt caagagaaga 1560
tctttcttcc aaaattgatg gaaggttctg aagatgttaa caaggagtac acaaaaatta 1620
cgtttacacc cacacaggca gatcggttg ttgtggcatt tagaaattgg ctgaggcgac 1680
acggcaacag ccaacctgaa tggttcggct tcagcagcca acaaccttcc ccttcaacag 1740
tcttctcga atgtcagatg ctcgatagat ttgagcagca caccctcaag tgttcatcat 1800
gtagagaagc ttattcagca ttccagacgg gccaaaagt ttctattggc gcgaccgttg 1860
cattctgcgc aacagctggg attccttcag atttgcaatc acggattgtt ttggctgggc 1920
ttgcactagt gagcgctgcc ttggcttatg ctttgcata actacaaaag aattttgtgt 1980
ttgttgatta tgtgcactgt gaaatcgatt agagaggag tagatatgct gctgaaagaa 2040
tcaatgtgtc cagaggtata agaaagatat ggtgataaat cttgtcaaaa ttgcgagtt 2100
tgtatatatc tattagatag aaatcagtg gatagctaa cttagagtgt ttctcagta 2160
ttccctttgt tttgtgcca aatgaatgtc atcagataaa tatgtgcaga catgcattca 2220
aattcatgag taaatggatt aaacaatata cactttgtca aatggaaaag agcgcatttc 2280
ctttaggaac aagcaaccac aaagttagt catgaatcag tttgaaaaaa caat 2334

```

```

SEQ ID NO: 22      moltype = DNA length = 2321
FEATURE            Location/Qualifiers
source              1..2321
                    mol_type = genomic DNA
                    organism = Citrus sinensis

```

```

SEQUENCE: 22
cgtcatgcaa tgaggagaaa ttaggtgtcg aaatctcata cgcggaccaca accacaagt 60
tcttctgtct ttttggaaac atttattatc tgcttcaccc tgctttgcag tccactgaaa 120
caaatcgaaa atggagcgcc ttattctctc ctctctctcc ctctctctcc tatcttctgt 180
gcttgcgctc gctgtagctg tcaacgacga cgatgctatg atcagacagg tctgtccgctc 240
agacggcgaa caatccgaag atcatctcc gaacgcggag caccacttct cctcttccaa 300
gtccaaatcc tccaagactt acgccacca ggaggagcac gattaccgat tccgctgtgt 360
caaggctaata ctgcgcgag caaagcgagc ccagctctctg gacccactct cgtgccacgg 420
tgtaccaagc ttctccgag taacgccctc tgaattccgc cgtcagttcc ttggcttgaa 480
tagggcggtt cggctgcag ctgacgctca aaaggctcct attctcccca ccaacgatct 540
tctactgac tttgactggc gtgatcacgg cgccgttact ggctgcaaa accaggtacg 600
ttagagaaag tgtctttttt ttttttcaaa aataaatttg aaaaaataa ttgatttttg 660
ttgatatttt tttattaatg tatgtagggc cgtgtgggtc gtttagtgca 720
accggcgctt tggaggggag gcactcttta tcgacgggag agcttctcag cctcagtgag 780

```

-continued

```

caacagcttg tggactgtga tcaacaggtt tgttcaattg tttgttatat ttttaacgtt 840
agttaaatga tgaaaattac agattttgtg tgattttgtt tgatagctag gatgagttgt 900
ataattttcta atcccaaaaca gttagttatg acggtaaaatc tcaaagttag attgggttct 960
cactctctga atttatgata aagaatttat atgtcagttt ccttttttct ttttaattct 1020
aatgttggtt cttcattatg ttaactctct tttttgttg gtttgatatt gttggatgcc 1080
ataatttgaa gttttgtggt gatataata atcatataac tatccacaat atgttgggct 1140
tgtgcttttt gtattctcaa tattgactgt ttgagttggt tggacgtttc cctctcaagt 1200
ctgaactcac atctcacatg atcaaaagtgt atgatttgac aaataaatt atgtttatta 1260
atgcaagttt tgtttcattt ttcatgtgta tccagaggaa tctggttcat gtgactctgg 1320
gtgcaatggt gggctaataa actctgcctt tgagtacata ctcaaggctg gtggggttga 1380
gcgagagaag gactaccctt acactgggac cgacgggtgt tctcgcaaat ttgacaaaag 1440
caaaattgct gcagctgtat ctaatttcag tgttatttcc tctgatgaag atcaaatggc 1500
tgcaaatgtg gtgaaacatg gccctctggc aggtaagtga gcttcgatac aattacctca 1560
tatttcgttt tccgtttctt ggcttttctc ttcactgtga gctctccaaa ataactttg 1620
gaaaagttag ttaatttaatt aatttctttt gagatgttgg taattttttt attaaacgga 1680
atggatagaa tgatgacaga atttgtgctg atcttgcctg ttgcttttgc agtgggtatc 1740
aatgccgttt ggatgcaaac atataattgga ggagtttcat gccatacat ttgcgggaag 1800
tatttggatc atggagtgct tatcgtgggc tatggatctt caggtttcgc cccgatccgt 1860
ttcaaggaga agccttactg gatcataaag aactcctggg gagagaactg gggagagaat 1920
ggatattata agcttgcgat gggtcgcaat gtctgtgggg tgcactccat ggtctcatct 1980
gtagctgctg tccatacaac ctcaagctag acattatgga ggttgtgcta ggcaagtgga 2040
gcttatatac gaagatatta taggatatcc ttttaaatag cgtctgcaa ttataaggat 2100
gcctacatgc gtgggctgag gcatgaactt tatatgctct tgtaatttt aagcatatgt 2160
catgtcagaa tgtaatatat atccatttta tagttaacca tgctacagaa ttgttattga 2220
agatggtatt aatattttct ttttatattg ggcaggcttg tagaaatatt ataattgtat 2280
attttctttt tatgtaactc aaaatagtag aacttcacgt a 2321

```

```

SEQ ID NO: 23      moltype = DNA length = 1890
FEATURE           Location/Qualifiers
source            1..1890
                  mol_type = genomic DNA
                  organism = Citrus sinensis

```

```

SEQUENCE: 23
atttattata ataaaaaatt gttaccaata aattaataac ctaaaaaaaa aaattccaat 60
ggatgtatgg taaagggtgag tacgatcgat gcagtcaccc gtgacgcggg tttcttttgt 120
tttatagttc caatttgcag acttatatga ctaatcaaat ccgtcatgca atgaggagaa 180
attaggtgtc gaaatctcat acgcggaccc aaccacaagt ttcttctgtc tttttggaaa 240
catttattat ctgcttcacg ctgctttgca gtccactgaa acaaatcgaa aatggagcgc 300
cttattctct cctctctcct cctcctcctc ctatcttctg tgettgcgtc cgctgtagct 360
gtcaacgcagc acgatgtcat gatcagacag gtctgtccgt cagacggcga acaatcgcaa 420
gatcatctcc tgaacgcgga gcaccacttc tccctcttca agtccaaatt ctccaagact 480
tacgccaccc agggaggagca cgattaccga ttccgcgtgt tcaaggctaa tctgcgcgca 540
gcaaaagcagc gccagctcct ggaccccaact gctgtccacg gtgtccacaa gttctccgac 600
ttaacgcctg ctgaattccg ccgtcagttc cttggcttga ataggcggtt tccggtgccca 660
gctgacgctc aaaaggctcc tatttctccc accaacgacg ttctactgca ctttgactgg 720
cgtgatcacg gcgcctgtac tggcgtcaaa gaccaggcgc catgtggatc gtgctggctg 780
tttagtgcaa cggcgctttt ggaggggagcg cacttcttat cgacgggcga gcttgtcagc 840
ctcagtgagc aacagcttgt ggaactgtgat cacgagtggt atccagagga atctggttca 900
tgtgactctg ggtgcaatgg ttgggctaag aactctgcct ttgagtacat actcaaggct 960
ggtgggggtg agcagagagaa ggaactaccct tacactggga ccgacgggtg ttctgcgaaa 1020
tttgacaaaa gcaaaattgc tgcagctgta tctaatttca gtgttatctc ctctgatgaa 1080
gatcaaatgg ctgcaaatat ggtgaaacat ggccctctg cagtgggtat caatgcogtt 1140
tggatgcaaa catatattgg aggagtttca tgcccataca tttgcgggaa gtatttggat 1200
catggagtgct ttatcgtggg ctatggatct tcaggtttgc ccccgatccg tttcaaggag 1260
aagccttact ggaatcataaa gaactcctgg ggagagaact ggggagagaa tggatattat 1320
aagatctgca tgggtcgcaa tgtctgtggg gtgcactcca tggctctcat tgtagctgct 1380
gtccatacaa cctcaagcta gacattatgg aggttgtgct agggcaagtg agcttatata 1440
cgaaagatatt ataggatatt cttttaaata gccgtctgca attataagga tgccatcatg 1500
cgtgggtgta ggcatgaact ttatatgctc ttgtaatat taagcatatg tcatgtcaga 1560
atgtaatat ttatccatttt atagttaacc atgctacaga attgttattg aagatgggtat 1620
taatattttc tttttatat ttggcaggct gttagaaatat tataatgtta tattttcttt 1680
ttatgtaact caaaaatagta gaacttcacg tatgcttaga cgcttgtatt ttcttattat 1740
atttgaagtt agctctatct agctctgttc atattaagaa atttcacaca actgccaaagt 1800
ttgtgactgc ctgaccagtg tcaactggtt aatagtctta tatgcttcta agtagcctaa 1860
gattacttag actctctttt ataaactggt 1890

```

```

SEQ ID NO: 24      moltype = DNA length = 1977
FEATURE           Location/Qualifiers
source            1..1977
                  mol_type = genomic DNA
                  organism = Citrus sinensis

```

```

SEQUENCE: 24
gaaattgatt tcagaaccag taaccaatat tcttcccaaa tatatcagtg ttaagtta 60
aacacttctt gaaggatggg catttgcact ccgacaataa atactatcca tgggaagcaca 120
atcacaagaa aagcaaaacc ttggagcctg taaaatgcca tcaagcatca tttctagtta 180
tcatttcaaa gaactggcct ctgcttttta tctaacagat caatgtttct ttggatttca 240

```

-continued

tcaaaatcaa	cacaaaaata	atgaagttat	agaagcttca	ttaccaccat	caaatcaatt	300
ttccggggat	aattttttcca	agaagttgtc	tgagctagac	actttggaat	cattggtatt	360
atcgagcaac	cgcaacagta	aattttcccag	aaaaaatagc	agctttccca	ctccttctga	420
gagcagccaa	aataactaaag	taagtttctt	gtgttacctt	acattatata	atatatgtaa	480
ttattacatc	aatcgattat	tttcttttct	gaattttgac	ctgtgtatat	aacctttttt	540
ttttttttgc	atcactatag	aatatgtctg	aattttgacc	tgtgtatata	accttctttt	600
ctgaattttg	acctgtgtat	ataacctttt	ttttttttt	gcacactat	agaatatgag	660
cattttttct	tcagaagaaa	agcattcttg	tgggttgatt	tctgattctt	atcgacacat	720
tttgcgaat	aaaaaaagaa	ttacgtggac	taaggatctg	catgaacatt	ttgtcagtg	780
tgtaaatcgc	cttggagggt	ctgagagtga	gtaaattgat	atgatcaata	atttatatag	840
gatctagcta	atgattttat	gtacttactt	attttaattt	ttgatgatca	ttgtttgtga	900
attgattaat	cctagaggca	acaccaagg	cgatactgaa	actgatgaaa	tcgaaagaat	960
tgagtatcct	acaagtaaaa	agtcatttgc	aggtttctat	taattaatta	aacaaactgct	1020
attgctgttt	ttttttttct	ttttacttca	tgaaatttta	attgtgattt	atctcaacat	1080
gagatctgat	attctctgtt	ttattctctt	cttgcgtgct	gtcagaaata	tcgatccgag	1140
aagctcatat	cagaccagtc	ttacaagggt	aattaatcaa	ttctgtttaa	aaaaatata	1200
ttctatttaa	tttgaatc	tgtaatttaa	aaaaaaaaaa	ttgcaggatt	tcccagagaaa	1260
acagtttgta	tcaatgat	atcctcagct	tacatgaaaa	tgtacgcaaa	ctcttcttga	1320
ttaatttctc	tctacattct	tgatattgtt	cataacaatt	ccagaaatta	tgtgcttaaa	1380
tttaattttat	gcaggggcat	gcaaataaga	gaggcacttc	aattgcagct	agaactcgag	1440
aaagcatctc	atgatcaatt	agaggcatgt	atttctatag	aactttaatt	tctattataa	1500
acttttaact	tttgggtacg	atattttttt	tcaaaaaaga	aaaaagagtt	aaaatcgtct	1560
tactacttta	aaaattgtaa	aataatagtg	tccaatttat	tattatattt	tacaaaaata	1620
tttactactt	ttctgttaat	gccataaatt	gacctgtaaa	atactaatac	taaaaaaaat	1680
tgggttattt	tgtggaacaa	ttaacaatga	ttatcatctt	catgaaaatg	attaatttga	1740
ttcacctcac	gtttaattta	ttactttctt	gctacagatg	caaatgaatt	tacaaaagct	1800
gattgaggat	caagggaagc	agggtgaagat	gatgttagag	aagcaattaa	aatcaaacca	1860
gaaataaatt	gagcttttacg	attataatta	tgtcgacaga	gatcatgtta	gaaaaggatt	1920
aattgtagtt	tattgacaac	ataatcacaa	gaaaaacaaa	aatgattgta	gtaataa	1977

SEQ ID NO: 25 moltype = DNA length = 693
 FEATURE Location/Qualifiers
 source 1..693
 mol_type = genomic DNA
 organism = Citrus sinensis

SEQUENCE: 25

atggaagcac	aatcacaga	aaagcaaac	ctggagcctg	gtaaaatgcc	atcaagcatc	60
atttctagtt	atcatttaca	agaactggct	tctgcttttt	atctaacaga	tcaatgtttc	120
tttgatttc	atcaaaatca	acacaaaaat	aatgaagtta	tagaagcttc	attaccacca	180
tcaaatcaat	tttccgggga	taatttttcc	aagaagtgtg	ctgagctaga	cactttggaa	240
tcattggtat	tatcgagcaa	ccgcaacagt	aaatttccca	gaaaaaatag	cagctttccc	300
actccttctg	agagcagcca	aaatactaaa	aaatatcgat	ccgagaagct	catatcagac	360
cagtctttac	agggatttcc	cgagaaaaaca	gtttgtatca	atgatatacc	tcagctttac	420
atgaaaaatg	gcattgcaat	aagagaggca	cttcaattgc	agctagaact	cgagaagcat	480
cttcatgatc	aattagagat	gcaaatgaat	ttacaaaagc	tgattgagga	tcaagggaag	540
agggtgaaga	tgatgttaga	gaagcaatta	aaatcaaac	agaaataatt	tgagctttac	600
gattataatt	atgtcgacag	agatcatgtt	agaaaaggat	taattgtagt	ttattgacaa	660
cataatcaca	agaaaaacaa	aatgattgt	agt			693

SEQ ID NO: 26 moltype = DNA length = 1001
 FEATURE Location/Qualifiers
 source 1..1001
 mol_type = genomic DNA
 organism = Citrus sinensis

SEQUENCE: 26

aacataaacg	aatttactct	ctgataaac	tttttaatat	atattttcac	ttacctgcag	60
ttgtagcgct	atctgctgac	tgctgctgct	gctgctactc	aaaaatggctg	aaaggatcca	120
ccccgaaacg	acaccggcca	acgaacaaga	gccctctcat	ccgcccggcg	ccgcccggcg	180
aggaacctac	gtcatccaaa	tcccgaagga	tcaaatctac	cgagttccgc	cccccgagaa	240
cgccgaccgc	atcaagggcc	tctcccgcgc	ccgcaagtcc	cgagcacta	cctgctgctg	300
cttcggttct	tgtctgtgct	cgctgcttct	cctcgctcct	ctcttgccca	tcgcccgcgc	360
cgtcttctac	ctcgtcttcc	gtcccgaatc	ccccaaactac	tccgtcgacg	gcgtctccat	420
cgccggcctc	aacctcactc	cgccgtcctc	cgctgtctct	ccccgggtcg	acgtctccgt	480
caccgcccgc	aatccgaacg	acaagatcgg	aatctactac	gagagaggca	gctcgggtgga	540
ggctctctac	aaggacgtcg	ccttatgcga	cgccgaatgg	cctcagtttt	accagccgag	600
caacaatgtc	acggttttca	agacctcgct	gaaaggatcg	tccatcgagt	tgaccagcgc	660
tatgcgcaaa	gacctggttg	ctgctcagac	gagtggcga	acggtgcctg	ttaagggtgaa	720
cttaagagtg	ccggttaaaa	taaaagtggg	gtcggttaag	acgtggacga	ttaagggtaaa	780
agtgagatgt	gatctgacgg	tggataagct	gacgtctcag	tccaagatcg	tatctaagga	840
ctgtgattac	tctgtcaaac	tttggttaaaa	aaagttaaaa	aaatttcaaa	tcaaaaggat	900
tcatgttaat	tgtaggatta	gattatacaa	ttaattatta	taatttggg	tgtatttgtt	960
acaaatacac	actattattt	atacttggtt	ttagtctgtt	t		1001

SEQ ID NO: 27 moltype = DNA length = 1141
 FEATURE Location/Qualifiers
 source 1..1141

-continued

```

mol_type = genomic DNA
organism = Citrus sinensis

SEQUENCE: 27
gtcagacaca caccggccaaa gtttaggcat acgcgttagc gcgtacgcgg tttctttatc 60
aaaaatatag ctatgctgca tgcataatgat ataataacat aaacgaatct actctctgat 120
aacacttttt aatatatatt ttcaacttacc tgcagttgta gcgctatctg ctgactgctg 180
ctgctgctgc tactcaaaat ggctgaaagg atccaccccg aaacgacacc gcgcaacgaa 240
caagagccct ctcatccgcc ggccgccgcg gccgcaggaa cctacgcat ccaaatcccg 300
aaggatcaaa tctaccgagt tccgcccccc gagaacgcgg accgcatcaa gggcctctcc 360
cgccgccgca agtcccgag cactacctgc tgcctgctcc gttctgctg ctgctcgctg 420
cttctctctg tctctctctt ggccatcgcc gccggcgctc tctacctgt cttccgtccc 480
gaatccccc actactccgt cgacggcgctc tccatcgccg gcctcaacct cactcgccg 540
tctctcgctg tctctcccc ggctcgactc tccgtcaccg ccgacaatcc gaacgacaag 600
atcggaatc actacgagag aggcagctcg gtggaggctc cctacaagga cgtcgcccta 660
tgcgacggcg aatggcctca gttttaccag ccgagcaaca atgtcacggt ttcaagacc 720
tcgctgaaag gatcgctcat cgagttgacc agcgctatgc gcaaagacct ggttgctgct 780
cagacgagtg gcaagacggt gccgtttaag gtgaacttaa gactgcccgt taaaaataaa 840
gtggggctcg ttaagacgtg gacgattaag gtaaaagtga gatgtgatct gacggtggtg 900
aagctgacgt ctacgtcgaa gatcgatct aaggactgtg attactctgt caaaccttgg 960
taaaaaaagt taaaaaaatt tcaaatcaaa aggattcatt gtaattgtag gattagatta 1020
tacaattaat tattataatt tgtggtgat ttgtacaaa tacacactta ttattatact 1080
tgttattagt ctgttttgta aaattcttgt gtggaacaaa gatatacaat taattaatta 1140
a

```

```

SEQ ID NO: 28      moltype = DNA length = 1380
FEATURE           Location/Qualifiers
source            1..1380
                  mol_type = genomic DNA
                  organism = Citrus sinensis

SEQUENCE: 28
caagggtttc aaaggagagt gacctttaaa aaaaaatgtc accgacacac aagaaaaaaa 60
taattaaaaa aaataaataa aaaaatgtct aagtccccta tataataagt gcatgtgaag 120
ctgagcgatg ccattcaata agtagccctc gcgaacaaa aaatgatttt cctactactc 180
ctctcagttt ttcttcgagg agttcttcca tcaatccttt ccgaagacac acctattagc 240
ttctcatttc cctcatctgc caaagacagt tgtgacaata agaccctcat ttgctacgga 300
gcgattgaaa gttccggcgc cttaagcatc acaccaggcc ctccaccaa cctgccgatc 360
agaaagggtg gacgggtttt atacggcaag cctctgagtt tacagcgatc ttttattgat 420
accaccatca ccattaagat ctcacgccat cagaattaca ctgacgtgac cggagatggc 480
atgacgttca tttttgcaag cgataaaaac ggtccatcag caaagggcgt cggcgaaat 540
cttggaactg agtcttcacc aggtatgatt atcaatgtga agaattaaga atttatgcag 600
ataccattga aagtactgac aaatgcattg atggggcgat tttatttata atggcagtca 660
aatgataaac aatatgatga tatatcatat gttctatata taacacttat acatatatat 720
tcatgatcat gcatgtaaat tatgcaggcg atgcagggcg tccatttagc gtggagctgg 780
acacatgcct gaacaagaac ctgaatgac cagatgataa ccatattggc atcgacataa 840
acggaatcga atcaaatcca gttaaatgtc tgcctgacgt tgatctcaaa agtggacgag 900
caatccagggt tcgaatttat tacaatccag actttggaca actctctatt tatgcggcat 960
attcggggga aacacttgtg aagggtgattg aaaaacccat taacctgtca gatataattc 1020
caacgcccggt ctatgttgga ttacacagcag ctacggggga ctttttagaa agccatgagg 1080
ttataaattg gacttccag tcgttcccag tgccgccttc tctcaaggag aaaaacctgg 1140
tgatgccaat aataattctaa accgtccttg aaaaaccatg tcacataaat tataatcaat 1200
aataattaat atcaccaata aagtgcacat gctcttgcat taaaattatt aataaaatga 1260
tagcagcgac taggattaat actgttttgt tgcctgtctga gatgtactgt atgtgcttat 1320
gtagaatgac tcatattcag cttctattga ttacgttgca gctttgagct tctgatgttt 1380

```

```

SEQ ID NO: 29      moltype = DNA length = 1512
FEATURE           Location/Qualifiers
source            1..1512
                  mol_type = genomic DNA
                  organism = Citrus sinensis

SEQUENCE: 29
acacaaggtt ttcaaaggag agtgaccttt aaaaaaaaat gtcaccgaca cacaagaaaa 60
aaataaataa aataaataaa taaaaaaatg tctaagtccc ctatataata agtgcattgt 120
aagctgagcg atgccattca ataagtagcc ctccgagaac aaaaatgat tttcctacta 180
ctcctctcag tttttcttcg cggagcttct tcatcaatcc ttccgaaga cacacctatt 240
agcttctcat ttccctcatt cgccaaagac agttgtgaca ataagacctt catttgcac 300
ggagcgattg aaagtcccg cgcttaagc atcacaccag gccctccacc aaacctgccg 360
atcagaaaag ttggacgggt tttatacggc aagcctctga gtttacagcg atcttttatt 420
gataccacca tcaccattaa gatctcacgc catcagaatt acactgatcg tgcggagat 480
ggcatgacgt tcatttttgc aagcgataaa aacggtccat cagcaaaggg cgtcggcgaa 540
tatcttgga tgcagcttc accagcgcat aaattctctc cattagccgt ggagctggac 600
acatgcctga acaagaacct gaatgatcca gatgataacc atattggcat cgacataaac 660
ggaatcgaat caaatccagt taatagtctg cttgacgttg atctcaaaag tggacgagca 720
atccagggtc gaatttatta caatccagac ttggacaac tctctattta tgcggcatat 780
tcgggggaaa cacttgtgaa ggtgattgaa aaacccatta acctgtcaga tataattcca 840
acgcccgctc atgttggtg acagcagct acgggggact ttttagaaag ccatgagggt 900
ataaattgga ccttcaactc gttcccgatg ccgccttctc tcaaggagaa aaacctgggt 960

```

-continued

atgccaatat	aattctaacc	cgtccttgaa	aaaccatgtc	acataaatta	taatcaataa	1020
taattaatat	caccaataaa	gtgacatggc	tcttgcatga	aaatattgaa	taaaatgata	1080
gcagcgacta	ggattaatac	tgtttgcttg	ctgtctgaga	tgtactgtat	gtgcttatgt	1140
agaaatgctc	atattcagct	tctattgatt	acggtgcagc	tttgagcttc	tgatgttttc	1200
ttaccctttt	cctcggtctc	ttctttaagt	ttatcggaac	aaaattctct	tctcacgggtg	1260
gatatcatag	cataaaattt	aaccatagga	aacaatgcat	gcacgctat	agatattaat	1320
tacatgtgga	gtcctttttc	ctgcggaacg	tctgtacgaa	ttttgtgctg	atggaactca	1380
gattttttaga	acgtcgtaga	tttgccctca	gatgtatatg	acaataataa	ctctattaga	1440
aatctgaatt	catacattta	ttgcgtagtc	catagtgtcta	tcgataataa	atatgaaata	1500
tgtgaaaaga	at					1512

SEQ ID NO: 30 moltype = DNA length = 7810
 FEATURE Location/Qualifiers
 source 1..7810
 mol_type = genomic DNA
 organism = Citrus sinensis

SEQUENCE: 30

atggagaaat	taattaagca	aatcatgaat	caaatacgag	aagaatcgaa	agaaattcaa	60
gaaccgtttc	aacaatcacc	tccactggta	agtgattcta	tttttccact	tttactccct	120
ccttcctttt	acctcatata	atttcaggta	aacgctaatt	tggtccatat	aatttaagag	180
aattattccc	tataaatttt	gtcttattag	ataaactctt	acaatcagaa	cactaggcca	240
agaagacaaa	ggcttcagcg	aaatggacaa	atatggccag	gtcaattctt	ttttgttttt	300
tttttggtt	cttttttcac	aattaaatcg	gtacagcagt	cagagtcagt	tgatgagatc	360
ttgcatttta	catcttttgc	tttaagcaca	tggaaatgaa	ttgttggtat	atgaatttta	420
gaattttctg	ctgtaccgta	catccgctt	ggccgtacag	cagaagtgta	gtacaaaatt	480
ccaatccatt	gtgattatat	atactataat	ttatggatgc	catttaatta	attcacgtag	540
gaaacagaa	atttgaggct	gaagattgat	gaattggcag	aggaagtaa	gaatggagaa	600
atataatag	caaatcatga	atcaaatcag	agaagaatcg	aaagaaattc	aagaaccgtt	660
ccaacaatca	cccactctgg	taagtgatgc	tattttttca	cttttacttc	ctcttccctt	720
ttacctcata	taatttcagg	taaacgctaa	tttggtccat	ataatttaag	agaattattc	780
cctataaatt	ttgtcttatt	agataaaact	ttacaatcag	aacactaggc	caagaagaca	840
aaggcttcag	cgaatgggac	aaatatggcc	aggccaattc	ttttttgttt	tttttttggt	900
ttcttttttc	acaattaatg	ctttcttttt	ctcgttggtt	tagctgtttt	agttgatata	960
gttgattcca	ttatttattt	tatagttttt	cttttccctt	agtgagtcgg	ccgacagctt	1020
atttactcgg	ttgagctttt	aataaataaa	ttttatttct	ttttgttaaa	aaaaattgcc	1080
caacccccca	aggggttcac	caagtagttt	ttttaattga	tcaccagaac	ttaaattttg	1140
agcaggttgg	agggctaaaa	aaaaaaatta	attttaaact	ttaaaattaa	tttatttgta	1200
aattaatttg	tgagggtgtc	tcagaaatca	gggttttcag	aaaattttta	gggggtgccag	1260
tagcaccat	cttaattttt	agagtttttc	tctgacgaag	atatcgaaag	agtttataag	1320
cagactaaaa	tatatctctg	atttagcttg	aaaactttga	attaaactca	ctagcgagaa	1380
aattctaaat	taactattta	agatccgata	cttatgaaga	cagtaattaa	ccaaacatgt	1440
cttggtcttt	ctattttttt	tatccttttt	gtttaatcag	tttttctcac	tctcctttct	1500
ggtttatagg	ttatgaggtg	atcggtgagg	taaggaaaca	atgttgactg	aaaagtctat	1560
caaaattttg	tttctctctc	tctctctctc	tattttcttc	ttctaagaca	ctgtctttata	1620
accagaaaa	taactcagga	cgacaaaggc	ttcaacgaaa	ggaaaaaatt	tggccagggtc	1680
aagatacata	tttgtgtctc	tttgttttcc	cgttaatcac	aatattttta	ttctaatttc	1740
gaattttgta	aaatctggat	gttcaaacct	gacaaacaata	tatatatata	tattattttat	1800
ttctttttat	taaaataaaa	atactcttat	aaatagaaaa	agaactccaa	tgtattgaaa	1860
attttatttt	tttatatttc	agtaacattg	agtttcgcat	tttttttaaa	tttcaaatgt	1920
gagactaagg	tggcgtttgt	tttttaactt	aataacttaa	agtgacttaa	cttaattaat	1980
taagttaatt	agagggtgtt	gtttttataa	cttaatgaga	cttttttagat	aactttgact	2040
taataaaaaa	agtttaattt	tttgactttt	tacttaattg	agaaatctga	attaagtcaa	2100
ttacttgcta	atgtcaaaaa	tatccctatt	ttataattta	tttttcatgt	atatcccaaa	2160
actcaccat	agatattttc	cccacataaa	aatatccctg	ttttttcttt	cttatattat	2220
atcagataaa	atttgaaatt	tatggtattt	tatatattaa	attctaaaaa	aattatgtat	2280
tcacttgaat	atagttttac	ttataaatat	taaaatattg	tggattttta	tattattttt	2340
atttgacatt	caaatttaaa	aaggatataa	atgtaaaagt	acataatttc	aactttttta	2400
gttgaaaaaa	ataaacaaat	taatacttat	tttccgagat	tcagacgaaa	aaataaatat	2460
attattttaga	ttcagacatt	cagacctatt	cagaatttag	acctattcag	gtttattcag	2520
atttcagaca	aaaaaacaaa	cgtcacctaa	gtctggagat	agtccttttt	attttcgcat	2580
tttttttata	ataaaaaaaa	ttttattttt	tgttcatctt	ttttcacatt	ttttttataa	2640
acaaatattt	accttttttc	atttgtcttc	ctatattttc	attcttttat	gtttaattta	2700
tccttttttc	tccttttttg	ttctagggat	ctgaagaggt	aattgggtgac	ggttgctcat	2760
tttcgattcg	tacggaaact	aggaaaaaac	ttaaagatgt	agggatttgt	gctcttcata	2820
attcaaaaata	taattttctc	taaaaaaaga	aaaaattttc	ctttataatt	cataataata	2880
atttattttt	tctaataata	taccagattc	gcggaaccaa	tccataccca	ggatccatgg	2940
tcaaggataa	agcaaaaaga	attgttgatg	acaataaggt	cagtacaaaa	ttatttgcaa	3000
acatcaatcc	aaaacaaaat	tcattactgt	cttatatgtg	ttattttatt	attataggaa	3060
acagttgcta	attcaattag	tgaaggcttt	gaagtgattg	gcgacgaggt	agatccagga	3120
ttaaaattat	ttttatttta	aatttattag	taggaaaaag	aacacttaca	tccttaaggt	3180
cggagaaaata	attaaataga	cccaaaattt	tctaattcca	agatattttt	tttatatcat	3240
aatatctttt	tatttttgtt	aactcaaaaa	atattgacaa	aataaaaaata	taatttttaa	3300
ataaatgtaa	taataaatat	attttaaaat	aactttaata	ttaaaaaata	ttttaatact	3360
taacaatcta	ttagacattt	tttattatat	caaaaataaa	ttttatacta	aatagaacaa	3420
taatgtttta	acaaatatat	taaaatctaa	attattttta	atattatgta	aaataataat	3480
tgggatattt	taattttttc	actagaagtt	ttttaatata	tcacgttatt	atcaagattt	3540

-continued

cttgaaggtt	gcatttttact	acctaattgtt	taggctatta	tcccttaacc	acccaaatgt	3600
taatatagtt	aaaaatttac	tgatgtcata	agggcttttaa	agaaattttc	attaaaacca	3660
atatgcacgt	ttctcaatta	tgaaataaac	tgctaaataa	taaaaaaatt	aaaaaattcc	3720
taaccggaaa	tgaaaattaa	atggaaattt	gaagaaattg	agagtgaaga	gaggaagaaa	3780
ggcaaatgct	cttgagaatt	tgagaagggt	agttcttagt	tgagaatctg	aagatgaagg	3840
agaaatggaa	aagcagaaga	aacgaagaag	aagacaaaaa	gggggaaaaa	agttaaggaa	3900
aaaaaaaaa	agaataaacg	aagaggggaaa	gaaaaagaaa	aggaaaaatg	agagacattt	3960
gaagatgaag	aagaaacaga	gaagccggag	aaatgaagaa	gaagagaaaa	aagggggaaa	4020
aaaaaaatca	gaaacacatt	aaagccctac	ccgcctcagc	tgccgcgaac	atccatcttc	4080
ttttcctcga	tcatcaccac	tagaaaacac	tggccatcat	aaccagcaaa	tcaacaccat	4140
gaaccgcgcg	cttaagtacc	ataaatacca	tcaaacagct	acgcagccaa	acaacacac	4200
tcgggtttgt	tatttctttt	ttttttcaaa	ggacaagatt	gtcttttgta	ctctagggac	4260
aaaattgaat	tttaacttat	attctgttaa	atatttactt	attttttaat	ggaatagtgg	4320
tacagtaata	accgcataaa	tattaaagtg	taaaatgcag	ttttcacgaa	tccttaattg	4380
taacttttgg	tagaaaaatg	ataatgtccc	taataattta	agcaccatta	gtaagtaaga	4440
tattttacta	attgtaaaat	atttttaatc	taaagattta	tataaaatgt	ctgtaataaa	4500
tacaaaata	tttttcaata	aatataatac	ttttatatat	tcaaaattat	tttagatact	4560
acaaaatttt	taattattatt	cgtcaactat	attcacaagag	tattatggta	aaaaaatatc	4620
ttagtggtct	tttctaaaaa	cgtgagcatt	tacatgtcta	tttccaaaaa	cttgatataa	4680
actatccatt	tctcctatta	ggctcctaaa	aatataattt	tgatagttaa	atatgctagg	4740
ttacattgta	atatacctta	catcaattgc	aaaaggatct	gattagctcc	aagctcatgt	4800
catttttttt	attcttttaa	actcttttgt	tcttagacaa	gctctatcaa	ccaagaaatt	4860
aggccacgac	gacaaaggct	tcagcgaaac	ggaaacattt	ggccagggtc	attcgttttt	4920
tttccctttt	attattttct	ttcttttttg	ttttcccttt	tctaattaat	catatctttt	4980
ccatttctta	gtttcatttt	attacgggtc	aacatttgaa	tttatttttt	gtgtgtgtgt	5040
atgtatgtgt	gtgtatatag	agttctgcca	tcatttttact	ttaatgcaga	cttaattaaag	5100
tagtaactaa	gtgtttttta	atagtataaa	aaaaatcaca	aactctcatg	tattatttat	5160
taatggatta	tgatgacaaa	tgctactttg	agtgaaaaaa	agtgactgag	aatttttagtg	5220
agtttagaaa	gagaccaaa	tacctttact	gattcttttg	ttattattat	tattttttaa	5280
aggtaagggt	ggataaagat	aaaattttat	tttaattctat	actcttttta	aaattttaa	5340
tcattgtatc	agttaaagaa	tatttaattaa	aaaccattta	aaaaaatacc	attaatgatt	5400
tcaactttta	attaaagttaa	tttaataatta	aatctctgag	tagctgttta	tgaatggaga	5460
cgtgcatgaa	ggctgtaact	aaacaaaatt	tgtcttggtc	ctctctgttt	tactcatgca	5520
gtctcctttt	cggtttctag	gctctgaggt	gattgttgac	gatgttcaag	attcgggtcc	5580
tgtggattct	gggaccaaca	atgtagggtc	gtgacctctt	ttccatacga	cattatatgc	5640
atacgtgtca	tttttcaatt	tttcggcact	gggattttat	taattttttc	gattatttta	5700
tttctgatta	gacaccgaaa	aaaaaaaata	caatcatttt	gccagttaaa	taaaattgga	5760
aaagaaacaa	atattaaact	caaaatctgt	cttttttttt	tttctttttg	tcttttttgt	5820
ttaattcttt	tttttttttt	tctattttct	tcttcttaga	tgagctctta	cgcccgagaa	5880
actaaggcag	gacggcaaa	gcttaagcga	aaagaataca	tttggccagg	tcaatatata	5940
tattttttct	tataaactaa	tctcatgttt	tctcagatg	caggtttatt	ttttaattta	6000
ttttaatctt	tagttaaaca	taattgtcag	acctcacta	gaaaataata	taaaaaatat	6060
catatatgta	agaatgaaaa	attaaatcac	gaactaatat	gtatgctgat	ttaattcaga	6120
gtgatgtcac	ttgacggaac	aaaaagtagg	aaaaaaatct	aacttcgatt	ttgaaagagt	6180
ttgtataata	ccgtgcttaa	tttgggtaac	tttttttttt	ttttttggg	tgaggatcac	6240
tgattttcac	ctcattttcc	ccactaaggc	tcgaacttag	tacttgccc	taagaagaca	6300
atgctcttac	tatatgaact	aagctctcgc	atccacttaa	tttggatact	taggtaggat	6360
atctactgat	gtgtcaccac	ttatataata	ttatgtatat	ggatataatt	aattttatta	6420
tcaatttata	acctactact	tttaaaattt	ctcaataaat	gatataaag	atgctatatc	6480
agtaagtatc	tcaattgagt	atccaaagt	aataagcaca	atattactgc	tttaatttat	6540
atatcttgat	ctttaatttt	ttctttcttt	tctgtttaac	ccatttttct	tattctcctt	6600
tttggtttct	aggttctgag	gtgattggtg	agagtgggtc	gatttttact	tatcaaaaaga	6660
aaaaactcga	agatgtacga	atttgagctc	tttctattat	tgatattaga	tacatgtcat	6720
ttacaaaatt	ttaccactat	gattaaatct	atttttgatt	ttcagattgg	ttagtatgaa	6780
taaatatccc	aaaattttat	ccaaaaataa	ttgggcattg	gatgtatgat	tcaatcaaat	6840
agttgataaa	tcttataggt	cccaagtaaa	ttagttataa	tatttcacta	tccaaacaaa	6900
tactacatca	tttaggatca	aagtttggtg	taggaaaaaa	aaatattttg	cttgaaaaat	6960
attttttgatt	ttgtccagtc	gggtataaatt	ttcggacact	gttgtagtgc	tcttttgtgat	7020
gtattttgag	ataaattcag	tacattgtct	tacatttgtg	cagctacctg	caaagtgcac	7080
ttcgcgcac	atttctctta	cgacacctcg	cgtatgcaag	aggttggtac	tggtatgtcc	7140
cgcttcaga	tcggtcgcg	attcagattc	cgtatgggag	aagtttttgc	cgtccgatta	7200
cgaagggttc	atttcgaact	catctttgat	cgacaggaag	aagaaggatc	tttattttca	7260
tctatgtcgc	aaccccatcc	tcttcgacaa	taataccgcg	agctttgggc	tagagcaaga	7320
gagtggtaaa	aaatgttaca	tggttggtgc	aaaatggatt	tatgaaaatt	cgggaatttc	7380
acaccgagat	tgcgaaatga	ttccttcac	agctggatct	aggtttctctg	aagtgtatga	7440
acttaagctt	atgtcagatt	tagaaatcga	agcaagattt	ggtacacaaa	ttttttcacc	7500
caaaaccaat	tatgcagctt	actttgtgtt	caagtttgcg	gaattcagag	aagggcctga	7560
aaactagctc	atagattttg	aagttctatt	tgagggaagc	cataatggca	aaaagcgtag	7620
agagtttctt	gacctcaac	tatctcaaga	ccgagggaat	aggtggatag	agattaagat	7680
gggtgagttc	tctattgaaa	atggagatga	aggaacagta	gtttgtaggc	tgtccgaacc	7740
agaaacctta	tctaagcgtg	gcactattat	tttccaaggt	attgaggtta	ggcctgaata	7800
tggcaggttaa						7810

SEQ ID NO: 31

moltype = DNA length = 1434

FEATURE

Location/Qualifiers

source

1..1434

-continued

```

mol_type = genomic DNA
organism = Citrus sinensis

SEQUENCE: 31
atggagaaa taattaagca aatcatgaat caaatacgag aagaatcgaa agaaattcaa 60
gaaccgttcc aacaatcacc tccactgaac actaggccaa gaagacaaag gcttcagcga 120
aatggacaaa tatggccagg ttatgaggtg atcgggtgaga cactgtctta taaccagaaa 180
actaatctag gacgacaaag gcttcaacga aaggaaaaaa tttggccagg atctgaagag 240
gtaattgggtg acggtttgtca tgtttcgatt cgtacggaaa ctaggaaaaa acttaaagat 300
attcgcggaa acaatccata ccaggatcc atggtcaagg ataaagcaaa agaaattggt 360
gatgacaata aggaacacgt tgctaattca attagtgaag gctttgaagt gattggcgac 420
gagacaagct ctatcaacca agaaattagg ccacgacgac aaaggcttca gcgaaacgga 480
aacatttggc caggctctga ggtgattgtt gacgatgttc aagattcggg tctgtggat 540
tctgggacca acaatatgag ctcttacggc cagaaaaact agccaggacg gcaaaggctt 600
aagcgaaaaa aatacatttg gccaggttct gaggtgattg gtgagagtgg ttcgatttct 660
acttatcaaa agaaaaaact cgaagatcta cctgcaaatg gcatttcggc catcatttct 720
cttacgcac ctcgcgatgc aagtaggttg gcactggtat gtcccgcctt cagatcggt 780
gcggttcag attcogtat ggagaagttt ttgccgtccg attacgaagg gttcatttct 840
aactcatctt tgatcgacag gaagaagaag gatctttatt ttcactatg tgcgaacccc 900
atcctcttcg acaataatc cgcgagcttt gggctagagc aagagagtgg taaaaaatgt 960
tacatggctg gtgcaaaatg gatttatgaa aattcgggaa ttccacaccg agattgcgaa 1020
atgattcctt catcagctgg atctaggttt cctgaagtga ttgaacttaa gcttatgtcg 1080
agttagaaaa tcgaagcaag atttggtaga acaatttttt cacccaaaac caattatgca 1140
gcttactttg tgttcaagtt tgcggaattc agagaagggc ctgaaactag tctatagat 1200
tttgaagctt attttgaggg aagccataat ggcaaaaagc gtagagagt tcttgatcct 1260
caactatctc aagaccgagg aaataggttg atagagatta agatgggtga gttctctatt 1320
gaaaatggag atgaaggaac agtagtttgt aggctgtccg aaccagaacc cttatctaag 1380
cgtggcacta ttattttcca aggtatttag gttaggcctg aatatggcag gtaa 1434

```

```

SEQ ID NO: 32      moltype = DNA length = 393
FEATURE            Location/Qualifiers
source              1..393
                    mol_type = genomic DNA
                    organism = Citrus sinensis

```

```

SEQUENCE: 32
atggctgaaa ggatccaccc cgaacgaca cagcgcaacg aacaagagcc ctacatccg 60
ccggcgcccc cgcccgcaag aaactacgtc atccaaatcc cgaaggatca aatctaccga 120
gttatgcccc cgatgaacgc gcaccgcac aagggcctct ccgcgcgcg caaatcccgt 180
agcactacct gctgtctgct ctgtttctgc tgcgtctcgc tgccttctct cgtcctcctc 240
ttggccatcg ccgcgcagct cttctacctc gtcttcgcgc ccgaagcccc caactactcc 300
gtcgacgaca agatcgccat ctactacgag agaggcagct cgatggaggt ctactacaag 360
gacgtcggct tatcgacagg cgtctggcct tag 393

```

```

SEQ ID NO: 33      moltype = DNA length = 393
FEATURE            Location/Qualifiers
source              1..393
                    mol_type = genomic DNA
                    organism = Citrus sinensis

```

```

SEQUENCE: 33
atggctgaaa ggatccaccc cgaacgaca cagcgcaacg aacaagagcc ctacatccg 60
ccggcgcccc cgcccgcaag aaactacgtc atccaaatcc cgaaggatca aatctaccga 120
gttatgcccc cgatgaacgc gcaccgcac aagggcctct ccgcgcgcg caaatcccgt 180
agcactacct gctgtctgct ctgtttctgc tgcgtctcgc tgccttctct cgtcctcctc 240
ttggccatcg ccgcgcagct cttctacctc gtcttcgcgc ccgaagcccc caactactcc 300
gtcgacgaca agatcgccat ctactacgag agaggcagct cgatggaggt ctactacaag 360
gacgtcggct tatcgacagg cgtctggcct tag 393

```

```

SEQ ID NO: 34      moltype = DNA length = 2434
FEATURE            Location/Qualifiers
source              1..2434
                    mol_type = genomic DNA
                    organism = Citrus sinensis

```

```

SEQUENCE: 34
tggctcttgg ttttgtcaat tagatcaatg ttcatttttc caatatatat atatacatc 60
acacacacat gaagtcttac cccgaagat ttatggaata taagagagaa ttttaacca 120
taagtttgaa ttcaacgaca tgcctgttca ttatttatcc attcaatctg catcttctc 180
ttttcagatt ttccgtgttt gttgcgcggt tccttctctt ttaacaaatt agggctcgaa 240
taattttgat tctaccattt aaaaactgaa tgcatagcag gctagcaaga agttcttgtt 300
cataaatttt tgcagtacct tccggaaatg aatactcaaa agcttaattt tcaggaaaca 360
tttcaaaaaa agcatttggg tttcggggca ccatcccaat atttcggcga tatccaccac 420
cagcagccct ggatgatgag aacaaccacc cagcagcacc aaaaatctga tcatgtctga 480
tctccgagca caatcttgag ccggtttgag tctccagctt cagcttttta cgcaactgaa 540
agatacatgg ggttttgtca gtagtattcc caagctgctg gtaacaactg ctcaaatatt 600
tccaggactt gtgattcttc acaacagttt catttgtatc agtcccctgg agaaaatttt 660
tctgttttat cagctgaaca agctgtccct ctagaaatcc cctgggaactt ttacaaatcc 720
cctgaagctt cgtgtatcaa tccccttggg aaacaatatt caggtccatt tgatgaacat 780
caagatcata gagtaagtag tttccctccc tggagacttt tcattcattt tttttcttt 840

```

-continued

```

ttttcattca gaatttggtt ttgattgtgg gtagtttttg tttttttttt ctaccttcag 900
gtctctaagt atgggttatgg attaacttca ctttcacaaac agggctacgc ttcacatcaa 960
gagaagcaat ctccaagatt ttcttctagt agttcttttt caactggacc tgtgatcacg 1020
aacaaaaact gaattagatg gactcaggat cttcatgaaa aatttgtcga atgtgtgaat 1080
cgactagggg gagcagacag taagtgttga tatttattga attttgtcag ttgaaggagc 1140
aaaaatTTTT aattatttgt ttttcacttt tttttttttt ttttgaattg gttgattaat 1200
tgcaaggagg acgcccgaagg caatttgaa gctgatggat tctgaaggat tgacaatttt 1260
tcatgtgaaa agtcatttgc agaaatatcg aatggccaag tacgtcccag aatttcctga 1320
aggatatatta aatctgcact ggttttggtt aaattgattt ttttttcttt ttaggggtcaa 1380
aattctgata ggaactctgt ttttttttgc aggaaaatta gagaaaagaa gtagcttgaa 1440
tgatttgctc caaatcgatg tcaaaagcgt aattcattat attttggaat tttacagaaa 1500
taattgttta gaagtctctc taacaactcg taacttccgt ctatggcagc actctgcaaa 1560
tcaaaaggagc attacaactt caattagatg tccaaaggcg actgcatgaa caactagagg 1620
tacattccag taatttttgt taatataatt tctatgaaaa tctcaatgc caaacattcc 1680
cttaatagcc atggtccaaa aatgtaattc ttgttctctc catgcaattc aatgttattc 1740
ttgaatggca gtagtaattt gattgatatt ctttttgggg tcagattcag agaaaattac 1800
agttgagaat tgaagagcaa ggggaagcat tcaagatggt gtttgatcaa caacaaaaag 1860
caagtaaggga tcaactcgaa cctcaaaatt tggaaaaagt accagaagt gaccccccat 1920
ttaattttga agggatcgaa ttttcaactt cagagaattc gggaaactcc catttcacgt 1980
aaaagataag ttagtttcat ttaactgaag ctgaaatcgt ttgaaaattt tatacgagaa 2040
gacttggggt tgaagcgaag attattacag ttcgtgcca tgaatcaaaa atagctgctt 2100
actgttacag agtgaagtat ttacattatg attctacaca cagaagaagt gattacaaa 2160
aagaagtaaaa taattacaaa gaagaagtaa atatatattt tacttgtaa taaatcatc 2220
aatgggttgt gtataaaatt tagatctaca ttattgaatc tagtacggat taattcaagc 2280
tccatcatct tgaacagat acagtgcgac agttttgatt tttgctgctt ggtctgtgta 2340
aagtaggttt caaattttga tttcatggtt tcatcagacg atgagtaggt ggagaaacag 2400
agttgaatac catgatcatt gtatctcttc ttaa 2434

```

```

SEQ ID NO: 35      moltype = DNA length = 1445
FEATURE            Location/Qualifiers
source              1..1445
                    mol_type = genomic DNA
                    organism = Citrus sinensis

```

```

SEQUENCE: 35
atgaatactc aaaagcttaa ttttcaggaa acatttcaaa aaaagcattt ggatttcggg 60
ccaccatccc aatatttcgg cgatatccac caccagcagc cctggatgat gagaacaacc 120
acccagcagc atcaaaatct tgatcatgct cgatctccga gcacaatctt gagccgggtt 180
gagtcctcag cttcagcttt ttacgcaact gaaagataca tgggggtttg tcagtatgat 240
tcccagctg ctggttaacaa ctgctcacaa ttttccagga cttgtgattc ttcacaacag 300
tttcatttgt atcagctccc tggagaaaaa ttttctgttt tatcagctga acaagctgtc 360
cctctagaaa ttcctggaa cttttacaaa tcccctgaag cttcgtgtat caatccccct 420
ggaaaacaat attcaggtcc atttgatgaa catcaagatc atagagtctc taatgatggt 480
tatggattaa cttcactttc acaacagggc tacgcttcac atcaagagaa gcaatctcca 540
agattttctt ctagtagttc tttttcaact ggacctgtga tcacgaacaa aactcgaatt 600
agatggactc aggtatcttc tgaaaaattt gtcgaatgtg tgaatcgact agggggagca 660
gacaaggcga cgccgaagtc aatattgaag ctgatggatt ctgaaggatt gacaattttt 720
catgtgaaaa gtcatttgca gaaatatcga atggccaagt acgtcccaga atttcttgaa 780
ggaaaattag agaaaagaag tagcttgaat gatttgcctc aaatcgatca ctcgtcaaat 840
caaaagagca ttacaacttc aattagatgt ccaaaggcga ctgcatgaac aactagagat 900
tcagagaaaa ttacagttga gaattgaaga gcaagggaag catctcaaga tgttgtttga 960
tcaacaacaa aaagcaagta aggtacactc gaagcctcaa aatttggaag aagtaccaga 1020
agatgacccc ccatttaatt ttgaagggtt cgaattttca acttcagaga attcgggaaa 1080
ctcccatttc acgtaaaaga tgaagttagt tcaatttaact gaagctgaaa tctgttgaaa 1140
attttatacg agaagacttg gggttgaagc aaagattatt acagttcgtg ccaatgaatc 1200
aaaaatagct gcttactgtt acagagtga gttattacat tatgattcta cacacagaag 1260
aagtgattac aaagaagaag taaataatta caaagaagaa gtaaatatat attttacttg 1320
ttaataaatc atacaatggt ttgtgtataa aatttagatc tacattattg aatctagtac 1380
ggattaattc aagctccatc atcttgtaac agatacagtg cgacagtttt gatttttgct 1440
gcttg 1445

```

```

SEQ ID NO: 36      moltype = DNA length = 1309
FEATURE            Location/Qualifiers
source              1..1309
                    mol_type = genomic DNA
                    organism = Citrus sinensis

```

```

SEQUENCE: 36
atgaatactc aaaagcttaa ttttcaggaa acatttcaaa aaaagcattt ggatttcggg 60
ccaccatccc aatatttcgg cgatatccac caccagcagc cctggatgat gagaacaacc 120
acccagcagc atcaaaatct tgatcatgct cgatctccga gcacaatctt gagccgggtt 180
gagtcctcag cttcagcttt ttacgcaact gaaagataca tgggggtttg tcagtatgat 240
tcccagctg ctggttaacaa ctgctcacaa ttttccagga cttgtgattc ttcacaacag 300
tttcatttgt atcagctccc tggagaaaaa ttttctgttt tatcagctga acaagctgtc 360
cctctagaaa ttcctggaa cttttacaaa tcccctgaag cttcgtgtat caatccccct 420
ggaaaacaat attcaggtcc atttgatgaa catcaagatc atagagtctc taatgatggt 480
tatggattaa cttcactttc acaacagggc tacgcttcac atcaagagaa gcaatctcca 540
agattttctt ctagtagttc tttttcaact ggacctgtga tcacgaacaa aactcgaatt 600

```

-continued

agatggactc	aggatcttca	tgaataattt	gtcgaatgtg	tgaatcgact	agggggagca	660
gacaaggcga	cgccgaaggc	aattattgaag	ctgatggatt	ctgaaggatt	gacaattttt	720
catgtgaaaa	gtcatttgca	gaaatatcga	atggccaagt	acgtcccaga	atttcctgaa	780
ggaaaattag	agaaaagaag	taccttgaat	gatttgcctc	aaatcgatgt	caaagccact	840
ctgcaaatca	aagaggcatt	acaacttcaa	ttagatgtcc	aaaggcgact	gcatgaacaa	900
ctagagattc	agagaaaaatt	acagttgaga	attgaagagc	aagggaagca	tctcaagatg	960
ttgtttgatc	aacaacaaaa	agcaagtaag	gatcactcga	agcctcaaaa	tttggaaaaa	1020
gtaccagaag	atgaccccc	atttaatttt	gaagggatcg	aattttcaac	ttcagagaat	1080
tcgggaaact	cccatttcac	gtaaaagata	agttagtctc	atttaactga	agctgaaatc	1140
gtttgaaat	tttatcacag	aagacttggg	gttgaagcaa	agattattac	agttcgtgcc	1200
aatgaatcaa	aaatagctgc	ttactgttac	agagtgaagt	atttacatta	tgattctaca	1260
cacagaagaa	gtgattacaa	agaagaagta	aataattaca	aagaagaag		1309

SEQ ID NO: 37 moltype = DNA length = 4122
 FEATURE Location/Qualifiers
 source 1..4122
 mol_type = genomic DNA
 organism = Citrus sinensis

SEQUENCE: 37

tgaagtgtca	atacttatat	agttcccttt	tacaggaaaa	agaaatgaac	atataagtaa	60
cgctagataa	taaaggggaa	gtgggaagag	ggaatgggga	actggttcag	gaaaagagag	120
gattctgttc	cttcaattaa	atagaataag	gagtccttta	agcttgcaaa	ttggccacca	180
aaaacccctg	aaatgcaccc	cccgtaata	ttgaattaaa	cgaaacatgc	atgcttcgat	240
gagtcgatct	caaaactcaat	gggtcccccga	cattcatact	ttaagctggg	aaagagcatc	300
aaaatttcgg	gaaaatgatg	ctcattctag	ctagttgctc	atgcgcgtgg	cgtaattaag	360
ccagaccaat	tttgtagatc	tgctattgag	tgctagctga	tgatggcaga	acttattccc	420
atataaataa	ccaccaatgg	caataattta	tacatacagg	aggtttacaa	tttctggaag	480
tgataccaag	cttttgattt	aagcatgaaa	gcgataccta	acggttaaaa		540
tttttgacaa	agaaataaac	ccatgcataa	taagaaaata	ggtttaattc	gtgcaactaa	600
tatacatcac	agttatcatc	caacaataag	actagtattt	ttagctgctt	aatatgcaaa	660
aattattgac	taataatagt	aagtaattgt	tttgacctgc	taacgtagga	atttgatcgg	720
tcagcgggaa	agttatcaat	taatcgcttg	aggccaaatt	accctttcga	tacatatagc	780
ataggtgaag	gcattatgta	tatgctgtcc	cttcaaaact	ctaagggtca	tttttagata	840
gtgtccacct	aaatggaaac	aaaagaaaaa	cctttgttta	ttagtggatc	tcagtcttac	900
cctgatgcaa	agagtcgtct	tcatcatcac	gcaagaattg	tttgtttatt	atcggtgtcc	960
ctcacaaatt	ataaagttat	taacaatttt	gcttgagcca	ttttggaagt	atcgagtact	1020
ccctgtttta	attagatcat	gttcaatttg	ccgctgacag	aatgatgatc	gtaaaatata	1080
atgcttaacc	tatgcattct	gattaattta	taattgggatt	cacaatgtat	gcgtaacacc	1140
gttataacta	agcaaacctt	cccagatact	actgataatg	acaagaactt	tcctaaaatg	1200
taaaacaacc	atgaagcgtc	taggatagaa	aatgacttgt	aacctaccaa	gtgtttaaga	1260
tcagtgatga	tttgatgagc	aacaaagtca	agatttatct	tagtaaatgg	ctcgttgtaa	1320
tgtatgatta	tttttggaag	tgtatgaatg	tcagttaatt	atatgctgta	agtgactaag	1380
taagtttgga	actcgggtgc	tatcaaacgc	aaacaagatt	aatgttacct	ttctctctga	1440
cttggaactga	ttaaataaga	tcaagcattg	atgtcgacac	cttatgctga	agctttacac	1500
atcagatata	cggaagaaga	gtctcctaatt	ttcttttata	agtaaacgaa	attcataact	1560
tactacattt	tgttgctcct	cctgatcaga	tcattgtctc	gacatcttct	atgcaataaa	1620
aataaagaaa	gaaagaaaga	aagaaaagatt	aaagttagcg	ccattctgag	cattcatcaa	1680
catataaatt	gaaagctagc	tatacaaaatt	tatttttggt	tgcatcattt	taaaatgaaa	1740
actgagaaga	gaaaggtgct	agaacagaga	tacagttata	gatcttcaca	ggaacagctg	1800
ccatgcaagg	tggaactcag	cttccattct	caattaccac	acgtagaggg	agaaggtctc	1860
ttcaagctaa	atccagggag	ttgtaatgca	tctggattca	tttgttctag	caatttcgca	1920
atgccaattt	cagtattcta	tgcaactgag	aattgcattg	acttttcaca	agatttggac	1980
aattttgatc	ttcaatcatc	cgttaaagtt	cacctgcaat	acaatcagaa	ttccagttta	2040
cctaaaaagc	agccgcatca	agatgcttat	cgaatttcac	cagcaagtgt	ttctcatttt	2100
atgcaggacc	cggcagaaga	agaagccctc	ctgaacgaaa	gacaaaaatg	tgttagtttc	2160
agtgaatata	agaagcatca	agtttagtaca	taaatatgtg	cccttggttg	tatatatata	2220
tgagctgca	gggtgaaggt	ggtttaactc	cttgcccttg	aactgcccga	ttctgaaacc	2280
aagctcatat	catgttcaga	ctcaccatga	gaagcagact	ccaataacta	tgactagtca	2340
tagcaaaaacc	agaataagat	ggactcagca	tcttcacaac	cgatttggtg	aatgcgtaga	2400
gtttcttggt	gggtgctgaa	gtactgcaaaa	accctttttt	ttttttcccc	agttactggc	2460
tttattttatt	gtgttactac	tcatcaaat	ttgggtattt	gtttgtagag	gctactccta	2520
agggaaatcct	gaaactgatg	gacatcgatg	gattgaccat	ctttcacggt	aaaagtcact	2580
tgacgggttg	atttgtgtgt	taaatatttt	atggctgtca	aatataatca	taactcaaag	2640
cccttagttt	ttttcttttc	ttctctctcc	ttttattttt	tgtatttcca	gaaatatcga	2700
acggcaaggc	acatttcaga	aggttatatta	ttttcaagtc	cctgataatt	gatattggtc	2760
ttaaactata	aatttacctg	tttgcatctg	atttcatcct	tgtccatgcc	aaatgccagt	2820
gocataaact	catgaatttt	gacgcttttg	ttgttttgca	tcaaatatca	acattgattg	2880
gtgtcacaga	caagaacaat	taacctgaca	tcaaacgaaa	aatatatatt	tccaaaaaat	2940
aaaaataaaa	aaggccatta	atatgccttt	ttgaagagta	ctttgtgtat	acttggaccc	3000
agcatataga	ttatttgatt	cttatacttt	atattcggaa	ttcgcatagt	taattcagct	3060
atcgtacgat	taatgaaaca	tgtctttgtc	ccatgacatt	ttgttggtcca	aatgctatat	3120
atataatgag	ctgatgtaat	tgtcgatctt	ccctttaatc	tgaatttaatt	agacctcaaa	3180
agaacaggtg	ctcgccaggt	aacgaaacta	atatatcatt	tctattttga	gatgaccatc	3240
atgggaatag	gtttcagcac	aaaaggggca	ctagattcta	ttatgtgaag	taagacatcc	3300
ataaaaaatca	tccatttaatt	acccttaggc	tatcaacctg	ggattttcag	acacaacaat	3360
cgtcttcata	ttagacccta	atgtaatact	aatagaatta	gccggggccg	cttagaggct	3420

-continued

```

cagttagaaa tgggggggaaa aagttacgtc tcagtaaatt tgctgtgcat tttatgcttt 3480
aattatataat gcgtgtgagt ttctgtatgca ggaaaatcaa agcgtgagag gacaaccgac 3540
ctgaatgcaa tagtaaggct cgactcagaa tcgtgaagctt ccggtgctga tcatacatat 3600
attctccctt taaatatctc aatccttgca ttgacctcaa gtcatttcc tgttttcaac 3660
aaagagagct ataggggaagt gatattctaca tgtatgcaat gcaaccatct aaaaatttcta 3720
aatatgcata cttgatctca gcttcattaa cagaggcatg cagcttgttg aaacattgaa 3780
attgcagcta gatgtccaga agcgttaca cgaccaactg gaggtaccct tttatcccta 3840
ttgtgattaa ggaattgata ctgatatatg ttcaccacat tataatccct taacagtttc 3900
gtttcacttg tctgggttag gtccaaagaa atctacagtt gcagattgaa gaacaaggga 3960
agcagcttac acagatgcta gaccagcaac taaagccaaa caaatctctc gttgattcca 4020
acaacgtgga tatcgagttg caagataacc aaccaaatga tctcaaagac acgcgtccct 4080
tcaacatata aggtttcaag gatgccttt atccttgcaa tg 4122

```

```

SEQ ID NO: 38      moltype = DNA length = 1008
FEATURE           Location/Qualifiers
source            1..1008
                  mol_type = genomic DNA
                  organism = Citrus sinensis

```

```

SEQUENCE: 38
atgaaaaactg agaagagaaa ggtgctagaa cagagatata gttatagatc ttcacaggaa 60
cagctgcccac gcaagggtggg actcagcttc cattctcaat taccacaact agaggagaaa 120
ggctcattca agctaaatcc agggagttgt aatgcatctg gattcatttg ttctagcaat 180
ttcgcaatgc caaattcagt attctatgca gctgagaatt gcatggactt ttcacaagat 240
ttggacaatt ttgatcttca atcatccgtt aaagtccacc tgcaatacaa tcagaatccc 300
agtttaccta aaaagcagcc gcatcaagat gcttatcgaa attcaccagc aagtgttttc 360
tcatttatgc aggaccggcg agaagaagaa gccctccctga acgaaagaca aaaatcggtt 420
agtttcagtg aatatcagaa gcatcaaat ctgaaaccaa gctcatatca tgttcagact 480
caccatgaga agcagactcc caatactatg actagtata gcaaaaccag aataagatgg 540
actcagcatc ttcacaaccg atttgttgaa tgcgtagagt ttcttggttg tgcgtgaaa 600
gctactccta agggaaatcc gaaactgatg gacatcgatg gattgaccat ctttcacgtt 660
aaaagtccat tgcagaaaata tcgaacggca aggcacattc cagaaggaaa atcaaaagcgt 720
gagaggacaa ccgacctgaa gccaatagta aggcctcgact cagaatcagg catgcagctt 780
gtggaaacat tgaattgca gctagatgtc cagaagcgtt tacacgacca actggagggtc 840
caaaagaaatc tacagtgtga gattgaagaa caagggaagc agcttacaca gatgttagac 900
cagcaactaa agccaaacaa atctctcggt gattccaaca acgtggatat cgagttgcaa 960
gataaccaac caaatgatct caaagacacg cgtcccttca acatataa 1008

```

```

SEQ ID NO: 39      moltype = DNA length = 4640
FEATURE           Location/Qualifiers
source            1..4640
                  mol_type = genomic DNA
                  organism = Solanum tuberosum

```

```

SEQUENCE: 39
ccaaaaagca gcgagatagg agtggacctg gcagtcctcc tgctatggcc gttccactct 60
tctctgcccac attcacaact tcaattctca cacaaccact tccctctctc cgctcttttt 120
ccgtcggaata actcttgctc tgctcgctcg cgtcaacttc catggaacat catgaatcga 180
aattcaagga atttccctat gcatctgttc cgcatagaga gttaatgggt gaacttgat 240
cgactgtgga gaatcgctct ggagaatctc tacttccctg tactctgctt tctcatgtgc 300
agtattttga gaattgaatc gctactgctc atgtctctct ctatgtcaga tctggaaatt 360
cctcttctca ggtactctcc gactgattct actacttatt acctattcat tcatctctgc 420
acttgtgcca attatatgta ttgagggtat atataagaa tacttctctg tggatgtttt 480
gttgaaatag tatataggaa ttgggttact ctatgttata aaagcttcta tggaaactca 540
agcggataat ccaattcagg caactatatt tagcaagact gagattcgag tgaataact 600
ccttctcatc ccattttatc ttaattttgt gttgttgcat cacaatgtta tgaaaaacga 660
aaagcgcaaaa aaaactttaa agtaggcggg gactttaagc ctgaaatata aataaagcgt 720
gagctttaat gaaaaaaaaga cacaataggg aaaaaaatc aaatatatta tatgtatcca 780
agactaataa atataagcat gaataacaaa tctatggaca aagaaattga tctttttata 840
ataaagtgtg ataccaatag cttagaaccc ttattcgcaa ggaaaagatt gtcttaggga 900
ctgaatgacg acaatgaagt gcacattaag cgaggcaag tgctcaacat attttgagcc 960
ttgcttctgg gattgagcat tcttaaagca cgcctttgac aacattgttg catcactcct 1020
acgtgagacc aaaaaaatgc gccagaat ttctcctaact accctccaat atgaatgatg 1080
tactgtctat cttgtacatg gtgaagataa agaaaaaag aaattattat gtactataga 1140
ataatgattg agaagctag atcttgcct ctaattttag tggatttgcc ttatacagaa 1200
gagttaatcc attacgaatt gtgcttttgt tgattggcct ttttactatg tatgctaaat 1260
atatcagctc ttttaataata gtaacatgaa ttgaaatgcc aaatacttaa ttgtataatg 1320
tgctgatggg agttaagaag atcagcagta cttgtattac tctgtgcatc tgtgggtttc 1380
tccatccaat gaagtccagc aagatcttta tcaaatctgt ttttttttag ctaatttcaa 1440
atttagtatt tcttggttcaa gccatatatc ttgccccttc tagtaataaa cagtcttata 1500
attcctttta attggacatg gctctaagaa aggtgttttg attatagctt attttaagta 1560
gcttttggtc ttttaagtcc tttttatttt tgagggtgtt gggaaagaca aaaagtgcct 1620
ttaagcactt atttttaggc gaaaatagta caaaaataag ttaaaaacca aaaaatgggc 1680
atgaccaact tatgactttt agcttttagc ttataagcta cttaaaaaaa gtcaatccaa 1740
tcaacccctc tgtagatgaa aaaaagcatc ctaattctat aataggttat tatatgggtt 1800
gttagtatcc aaaaatttaa ctttctcaaa aaaaaaatga tataagccaa atggcgaatg 1860
taaaaatagg cccctaaagc aatgtgtgtg ctcccttttg gttttaagtc tatcaatgga 1920
ttctggtgtc tagagtctaa agggcatgtt ttgaaagaca ataaacaaaa ggggtgtgca 1980

```

-continued

```

taaaaatcac atacccaaag gggttggtt tgaattgtac tgcaaaactgc catatttttc 2040
acggacgacc cctaactgcc atttggtaca tttaaaaaat ataaaaacaaa cgcaattcgt 2100
gaaatgtggt gccaattccc actaattttt taatagtgtg tagcatgttc tgtgaactgc 2160
atctcaacta gttttgctgt gatcttgta aagtagttg atctctgctt gttgcccctc 2220
ccctagggtg aggagctgta aggggaaaaat tcttatgac tgcaatctca ccttctcaca 2280
taagctttcc caacttcaat atctggtggc actgagttcc catctgaata ggtgccgagg 2340
tagtttgta actagatct aaactggaac aatagaagtt tcttgctctg gtttaaaatt 2400
cagtgcgcat ctccaggaatt tcatagtgta tgacccaatg tgaatggatg acttgaaaat 2460
atgtgattga agtctgtttt gttacagaaa atcaatgtaa acagtgtctg agttgtagta 2520
tgcaatcttg actcttcctt tcaattgaaa ttgactcaca aaaattttcc ttagtttttt 2580
catagtaggc tcccttgatg aatttaccga caatagtcac ttgcaattat tggccgaatt 2640
cacgaaatag tcatcagcag agaaaaatat ggttggctta ccgagatgag tctcatgcct 2700
gatttcaata ggagcttcgg cggttgaggg aatcgccaaa ctgactgaa cctttctgcg 2760
aagggttggt tcgatggagc aacaccctta cctttccctg ctaaatggg attttctgt 2820
tgaggattga tatcggtgt tctgatctgg gcttgaaaga ttggttatag caggattttt 2880
atcagtgga caggggttat gatcagtttc attgatgttg gtcgaggcaa ccgatttcaa 2940
ttttgaagag ggtgtgtcga aatctgagga caattttggc tgagggggtg tatgattttg 3000
gtgggtggat ttcaaatggg cagtgttttc ctgttgagggt gttgcatggt gatcgaaaaa 3060
agagtaggct gagtgttaatt ggcattctgg aggttctgga gaattgttg gggcaagctt 3120
taggagctta ttactgtcca tgtgtgttc ctctgaagga ttggaagata taggggatga 3180
tgttttgttg ttgttggtg tgttggtgct agaaggagtt gattttggtg agtttgggat 3240
ttctcgggtg gatgtgttca aactagcaat tgggggtgtg aggagagcag gtgtgttttg 3300
ctgagttgtc ggtgtaggag aggtgcagc tgggtgagat gagtatggat tgtggatgta 3360
taactagtgt tttagggtgg ttccgatgca ccatttgggt gtttgcgtga aagatcaata 3420
gatcttcata gagagaagc agagagaata tgcatccaaa attgaccaac caatctaaca 3480
gctcgattca agaattatcc ttgtgcataa tttgtagttt actgcatttc tgctagtgtc 3540
gttccaccga tatcttcttg gggaaagttat acataaggac tgcaaaagca tcatggagat 3600
agcaagtagg aagcttaatt agttcatcaa tctaactctc ctatgcttgt tcattcttga 3660
atttggatct ggtctctgtt cttttattct tgtccagttc catctctctg cttatagtta 3720
ctagtctcac tgagagttct ttgacaagtc ttgtttctct ttgagattta aagttctttt 3780
tctaaccgct ataataaaag aaaaactggg aacacacagt taacgatttc aactttctca 3840
tcaggagtat atgcaagggt aatatttttg ccttttctaa attgtgacag gttgatttca 3900
tacttggtag ttgggttcac tgcaacctac ccacaggcgg agcgttgaat attacaagcc 3960
tttcagcata tttagagcct tcaactgatg caccaaaact cttaatgaa gttatccgca 4020
gcagtcacac aactctcatc cttattcttg atctacctcc gcgaaaggac cttgtccaac 4080
atcctgatta cctcaagacc ttttatgagg aaacacaatt agacaagcag agacaacttc 4140
tcgagaaatt acctgagcta aagccttact tctcttcgtc tctatatatt cgatccctag 4200
tctctccatt ggtctacttg gtttctatag aaaccgaacc ttcccaggcc attcgcattg 4260
atgagattat tcaggatcac ataagtcctg ttgctaagggt aatgctggat acatggttgg 4320
atctgtgtgc ttgtactgag agaagattga cagatgatga aagtgcagat ctggctaaga 4380
gggatcgaaat aattaagaat aagactatcg agatagatct tgaatcaagc ttccctaggc 4440
gtttcgggca agaagtagcg aaccagggtt taggagtagt aagggaaatc tacaacagtt 4500
gaatttcttg cttctgctgc tgttttattg tgtgattatt gtatgtaatc tttataattc 4560
ttcaacatat aatacattta aaaagatgta aattgagagt aactataaaa gttgcattct 4620
tctattttaga gttcttgtca 4640

```

```

SEQ ID NO: 40      moltype = DNA length = 939
FEATURE            Location/Qualifiers
source              1..939
                    mol_type = genomic DNA
                    organism = Solanum tuberosum

```

```

SEQUENCE: 40
atggccggttc cactcttctc tgcccaattt cacacttcaa ttctcacaca accacttccc 60
tcttctcgct ctttttcctg cggaaaaactc ttgtcttgct cgtcgtcgtc aacttccatg 120
gaacatcatg aatcgaaaatt caaggaaattt ccttatgcat ctgttccgca tagagagtta 180
atgggtggaac ttgtatcgac tgtggagaat cgtcttgag aatctctact tcttctact 240
ctgccttctc atgtgcagta ttttgagaat gaactctgta ctgctcatgc ttctctctat 300
gtcagatctg gaaatttctc ttctcaggtt gatttcatac ttggtagtgt gtttcaactgc 360
aacctaccca caggcggagc gttgaatatt acaagccttt cagcatattt gagaccttca 420
actgatgcac caaactctct aattgaagtt atccgcagca gtccaacaac tctcatcctt 480
attcttgatc tacctccgag aaaggacctt gtccaacatc ctgattacct caagaccttt 540
tatgaggaaa cgcaattaga caagcagaga caacttctcg agaaattacc tgaggctctg 600
tcttatctct cttcgctctc atatatcca ccccaacca ctccatgggt tatattgaat 660
tctatacaaa ccgaaccttc ccaggccatt cgcattgatg atattattca ggatcacata 720
agtcctgttg ctaaggtaac gctggataca tgggtggatc tgtgtgcttg tactgagaga 780
agattgacag atgatgaaag tgcatgctg gctaagaggg atcgaaat taagaataag 840
actatcgaga tagatcttga atcaagcttc cctaggcttt tcgggcaaga agtagcgaa 900
caggttttag gagtactaag ggaaatctac aacagttga 939

```

```

SEQ ID NO: 41      moltype = DNA length = 2345
FEATURE            Location/Qualifiers
source              1..2345
                    mol_type = genomic DNA
                    organism = Solanum tuberosum

```

```

SEQUENCE: 41
ctattcttac aatcttccc acattacaat aattacaatg tcaagtctct cactctatt 60

```

-continued

```

ggtttctgtgc gccggccttt tcgctgtgtgc acttgccgga ccggcgacct tcgccgatga 120
gaatccgatc aggcgaagtag tagtttccga agagctggag aacggaattc ttcaagtcgt 180
cgcccgagct cgcaatgctc tctccttcgc tcgctttgct atcagggtac taaattcgaa 240
aatgagatcg atttctctat cgttttgctt ttataattaa actaatattg ttctattttt 300
gaatgtttat aaaaggcatc ggaaaaggta cgagtcggtt gaggagatca agcaaagggt 360
cgagatattt ttggacaatc tgaagatgat ccgatcgcat aacagcaaa gactatcata 420
caaaactcgt gtcaatggta tactttctta tcttcaacta ggaaaacaag ttcacactta 480
ctaattagtt atttagatta taaaaactca gtatgaacta cttattcatg atgtataaaa 540
aaaacacttt gtagaagtta agttctttct ttattttact tagtaactaa tcaagctaaa 600
tgaacaattc ttaattgaat tcttatattc atcatatatg cttggataag tgtttgcaac 660
ttatgatcaa tctaattggt tatcgattac ttttatgttt catggcagag ttaccgacc 720
taacatggga tgagttccgt agacacaagt tgggggcac tcaaaactgt tctgccacta 780
caaggggcaa tctcaagcta actaacgtcg ttctgccaga gacggtatat ccaatctgaa 840
tgaactccga tcttttatgg ttatatattt ctggaggtac tcattagagt taattaaact 900
agtttgatc taatgcttta ttatttccaa gatggtagag tgcactgagt tgaattttgc 960
tataataaag atagaaacac taacacatca tccaccgtgt gctgcgttaa ttagtgtgtt 1020
caattgggtg cagaaagatt atttttttga tcttaacgaa caccaaaatt ccaattttgt 1080
gattatagaa ggactggagg gaagtcggtt ttgttagccc agtgaaggca cagggcaagt 1140
gcggatcttg ctggacattc aggtgagaat tagttagaat catgttggac tctaaaatt 1200
gaaatctaat ggagcaggca tatatatgtg ggggttttgc agcactactg gtgcactaga 1260
ggcagcatat gcccaagcat ttgggaaggg aatctctctg tcagagcagc agcttgttga 1320
ctgtgctgga gcttttaata actttggctg caatgggggg ttgccatcac aagccttga 1380
gtacattaaa tctaatggtg gtcttgacac tgaagaagca tatccataca ccggcaagaa 1440
tggcatatgt aaattctcac aagcaaatat tgggtgtcaa gtcacatggt ctgtcaatat 1500
taacctggta attaagatct ctttagtttc cttgggatgg aaccaacttt ttgccagtgt 1560
tattcagccc atttggttaa cttattgagc tgctgctttt accaattaca catatggact 1620
cctgattaac atgtgttatt acagggtgct gaagatgaac tgaataacgc agttgcattg 1680
gttaggcctg ttagtggtgc ttttgagggt gtaaaagggt tcaaacagta taagagcgga 1740
gtttacacca gcaactgaat tggcgacact cccatggtaa gtcactctg ccgagtaacc 1800
tgagaagatg caattatcta ttatcaccta aataggccta tatggacaat attacaaaca 1860
ctgactgttt catgggcagg acgttaacca tgctgttctt gctgtgggtt acggtgttga 1920
aaatggcgtt cctcactggc tcataaagaa ctcatgggga gcagattggg gtgaggatgg 1980
atacttcaaa ttggagatgg gaaagaacat gtgtgggtgt gcgacttgcg catcctaccc 2040
aatcgttgcc taagctttgg agttttgtga aaaaattatg cataaatccg tgttgtccca 2100
gttaaatgatg cagcagcagc attcagggtc cattctcaga tttatattca gaacatgtat 2160
ggatcggtat acatacaaaa atgggttagg ctacttatat gaaagaaaca ataagatcaa 2220
aatatttagt tcacagagat tattatgcag gaaaagtccc catgtaattt atacattata 2280
agtaatgaaa gggaggaaga aattcttatt gtaagcatta ttaatccact gttgtcctta 2340
gttta 2345

```

```

SEQ ID NO: 42      moltype = DNA length = 1071
FEATURE           Location/Qualifiers
source            1..1071
                  mol_type = genomic DNA
                  organism = Solanum tuberosum

```

```

SEQUENCE: 42
atgtcaagtc tctcactcct attggttctt gtcgccggcc ttttctgtgc tgcacttgcc 60
ggaccggcga ccttcgccga tgagaatccg atcaggcgaag tagtagtttc cgaagagctg 120
gagaacggaa ttcttcaagt cgtcgccag actcgcaatg ctctctcctt cgctcgcttt 180
gctatcaggc atcggaaaag gtacgagtc cgttgaggaga tcaagcaaa gttcgagata 240
tttttgaca atctgaagat gatccgatcg cataacagca aaggactatc atacaaactc 300
ggtgtcaatg agtttaccca cctaacatgg gatgagttcc gtagacacaa gttgggggca 360
tctcaaaact gttctgccac tacaaaaggc aatctcaagc taactaacgt cgttctgcca 420
gagacgaagg actggaggga agtcggtatt gttagcccag tgaaggcaca gggcaagtgc 480
ggatcttgct ggacattcag cactactggt gcactagagg cagcatatgc ccaagcattt 540
gggaagggaa tctctctgtc agagcagcag cttgtggact gtgctggagc ttttaataac 600
tttgctgca atgggggggt gccatcacaa gcctttgagt acattaaatt caatggtggt 660
cttgacactg aagaagcata tccatacacc ggcaagaatg gcatatgtaa attctcacaa 720
gcaaatattg gtgtcaaaag catcagttct gtcaatatta ccctgggtgc tgaagatgaa 780
ctgaaatacg cagttgcatt ggttaggcct gtttaggttg cttttgaggt ggtaaaagg 840
ttcaaacagt ataagagcgg agtttacacc agcactgaat gtggcgacac tcccattggc 900
gtaaaccatg ctgttcttgc tgtgggttac ggtgttgaaa atggcgcttc ctactggctc 960
ataaagaact catggggagc agattggggt gaggatggat acttcaaaat ggagatggga 1020
aagaacatgt gtggtgttgc gacttgcgca tcctacccaa tcgttgcccta a 1071

```

```

SEQ ID NO: 43      moltype = length =
SEQUENCE: 43
000

```

```

SEQ ID NO: 44      moltype = DNA length = 1632
FEATURE           Location/Qualifiers
source            1..1632
                  mol_type = genomic DNA
                  organism = Solanum tuberosum

```

```

SEQUENCE: 44
atgcaaaacta tcaaagcttc ttcttttca ccatttcacc tcaacttgaa ctcaactagt 60

```

-continued

```

tcatttccca aaattacca cttgtacatt caacaaaatt atgaaaaccc cattttctgt 120
tttccctcaa ttcagagcca aaatgcaaaa ttcaagggtt ttactgctat ttccccaagt 180
gttttcaactg aatcagaaac cccattttgat gaaaggactg aaaaatgaaa tcaagaagag 240
aaatttgagt ggtatgctga gtggtacca ataatgcca tttgtgatct tgataagagg 300
aggccacatg ggaagaaagt gatgggtatt gatgtggttg tgtgggtgga taagaatgag 360
aaagaatgga aagtaatgga tgattcttgt cctcatagat atgctccact ttctgaagga 420
agaattgatc aatggggaag attgcagtgt gtgtatcatg gttgggtgctt taatggagct 480
ggtgattgta agtttatccc tcaagctcct agggatgggc ctccggttca tacgtccaaa 540
agagcttggt caactgttta tccaagttgt gtgcaaaatg acattctcttg gttttggcca 600
aactctgata ctctatacaa ggacatatat ttgacgaaaa ggctccctta tatacctgaa 660
cttgatgaca gttcgttttc gaaaaacctt atagtcagag atatatcata tgggtatgag 720
cttctgattg aaaaacctat ggaccagct catgtccaat attcacacta tggcattatg 780
aatgttccag tagccccc aaagtgtgaaa gctgatagag aagggggaag accacttgac 840
ataactgtca cgaagtggga tgtaaatgtc attactgcaa accagggacc tggacggaa 900
acatttgttc cgccttggtg gtattatagt tattttgctt tcggaggacc tcagggaaaa 960
acatctgctg tatcatctgg aactgtacag gaaaaacctt cagctgagaa gcagaaaaaa 1020
gcacttctag tttcatctg tattccggtt agtccaggtc atagcagaat tatatttgca 1080
tctccaagaa actttgccac ttgggcagat cgaataatct cagcttggtt atttcacctg 1140
ggacaaaatc taattctgga ttctgatttg tatcttcttc atgtggagga gcgcaagcta 1200
aaggaaaatt gctcttataa ttggcataaa gcttgctatg tgccaacaaa ggcagatgcc 1260
attgttgttg cttttagaag gtggctaaac aaatatgcag gtggtcaagt tgattggcgt 1320
ggaaagtaca atggggacct ccgcgcaact cctccaaggg agcagctgct ggacaggtat 1380
tggactcata cagtgaaatt cacaagttgc aatcttgcat ataaagttct caatgctctt 1440
gaagtgttac tgcagatcgc ctccattggt gtgctcggaa ttgttgctgc tgc aaagcag 1500
ggcacattgt cagtggtggc taggtattct ttggtcacca ttgcattact atgcttctg 1560
gcctcgagat ggttatctca ttttatatac aaaaatttcc atttccacga ttatgatcac 1620
gccttctgct ga                                     1632

```

```

SEQ ID NO: 45      moltype = DNA length = 4284
FEATURE           Location/Qualifiers
source            1..4284
                  mol_type = genomic DNA
                  organism = Solanum tuberosum

```

```

SEQUENCE: 45
gattgacaca agaaaaagaa gcatacactt gaagactagc tagctatatg gaagctctta 60
aaatttctac ttgttttcca ccatctctct tcaacttgaa aacacctaga ttttcaagaa 120
ttatatgtga gaagaaacga aacttttctt ttctctaaaa tcatcaacaa ccccaaga 180
caagattcaa tcttttctac accaatatct ttaattcaac taatgaacca caacaactac 240
tccaaaatga tgaacaagaa attagtacta aaaaatgaaa aaatcaagaa aaagagaaat 300
ttgattggta tgcacaatgg tatccaataa tgccactaag tgaacttgat aagagaaagg 360
cacatgggaa aaaagtgtag ggaattgatt tagtagtggt gtgggataaa aatttggagg 420
aatggagagt gatggatgat gcttgttctc atagattggc tccactttct caagggagaa 480
ttgatcaatg gggaaagatt cagtggtgtg atcatggttg gtgttttagt ggttctggtg 540
attgcaagtt catctctcag gctcctagag acaagcttca tgtaattact actccttttt 600
tatctgtcgt ttaacatatt gacacatcta ttaagaaaaa catttgataa tatatgtaac 660
attttttttt ttctgtttta cgttatttag ttcatttcag ctcaaggctc aaacacgaaa 720
tttctgatta ggggtgaaag tattttgtcc atcctatcat aatccttgga ttttctttt 780
agatgttttt caaagtgaat cacactttaa acttttttta acttctcatt ttacatttag 840
gttaatgacg tacattattt tccattgaca ttataaataa acatgtcttt taacttggct 900
tcagctagac acacacatcc catgtgacgt cctacatgat atttcacaac ctatgtaagt 960
tccacttgtt attatgccaa gtaggacatg tgtgtctatg tgttctctt tatgcaagtt 1020
taagtatcta cttgtgcaca ctcaaattag ttgaagagca tagatgacaa ctgaggccaa 1080
taaaagactc atttatgcat ttacgctcag atccggaatt ccaactgtcc gtacggagtg 1140
tgcataataa ttgacctggg gtatttccac ccccttcaa tcagtcccg ggagtttcca 1200
cacacacata tccgggagcc cgcattgaat cggatagac tctactatca tatcaaat 1260
agatttactc ctaactcatt attgcatata ttcaaggcca atgagttttc caaataagg 1320
tgtgtgtaaa tgatcattaa ttaaatgttt cgacttttca cataattttg tacagttttt 1380
tttatttact tatgacaagg agtgactata tattaatttt ttgtaggttc acacatccaa 1440
aagagcatgt gtgacagttg atccaagttt tgtgcaaaat gacattcttt ggttttggcc 1500
caacactgat cctttatata aggacataca cttgagcaaa accccacctt atattccacc 1560
tttagatgat ttaacttcat atgcaaaaac aacacttggt agagacatcc catatgggt 1620
agtacccttg taaatatatt tgggtgttta ttttggcatg acaaaaaata attttggaa 1680
aaatattttt aagaaaataa gtcatttttt tgaaaaaaga aaaagttaat aagtcatttt 1740
ctgaagtttg gttaaaatta tcaataagta ctttcaagga aaaacatttt aaatgactt 1800
ctctcacttt aggcataaagt cattttccca cacaattatt tcaactctag aaaagtttgc 1860
attagcctat ttattattac ttggtagaat taatatttaa tcagagtttg aaaaatttaa 1920
ttgagaaata attatttttt caggtagtag tttttgattg aaaacctcat ggacctatct 1980
catgtcaatt atgcacatca tggcataaag aaaattggga aaatagaagt tccaaacaga 2040
taagaaaatt aaaccttaac atatatatac atacttgatt tgaattttta ttgtatttgg 2100
atatttttac caaactaata atataataat aatagtgcga ctgtgaaggg tgatagagaa 2160
ggaggaaaac cacttgatata aagccttgaa aaattagaca taaatggatt tattgcaag 2220
caaggacatg atgaacacaa atttattgcc ccttgtgtgt actatgggtc atttgggtgt 2280
caaaactatt tggataatta tgaatcaaag gtaattatct ttttctgaat tttctctata 2340
atatcatcgt ttaacaaaat accttttgat tgttataaca tgctattttg ttatagagaa 2400
catataatat aacataaaaa ttgatctaaa taaattggct gctattttta atgtcctacc 2460
taccatgt attcgaggga gaggttagaa tcgaaaaaga tcaggttcca cacatataag 2520

```

-continued

```

ataggcagaa ggtatgggac gagtagtatg atgtacggag aaccaagacc agtttaatat 2580
taattaattg ctggatcaaa acaaaaatta acccccccgc gcccggaag ttaaaactgca 2640
ttgattatat gaagggataa gacactagta ccccatgtga caccttttgg ttacgtggc 2700
acactctgtg actccacgtg gttgaggcgc gtaggatatg tttggatgcc acgtaagcca 2760
aaaagatgta caaaattaca aataaataaa taaatttaga ataatagaac cttagtttaa 2820
ttaagggtgtg cctctagatt ttgatcatga tctagaggga tacctgtgct ttacgtgtaa 2880
tataaaagat ctttactatg aaagtgtact tctaagattt attttatttg tgatgtgatg 2940
caggaagaat catcatcaaa tgatacaaat agaataattc tagtatttat atgtgttcca 3000
gtaagtccag gtaattgcag attgatgatg acatccttca gaaactttgc tggttgggag 3060
tataaactat ttccaccatg gaaatttcac cttgacaata acctaatacat tgattctgat 3120
ttataattac ttcatcttca ggtaattaat tatcctctta caattttttg attttactag 3180
tagtcgtaac attaatcttg ttctttgatt ttattataaa ataaaatatt attttttcaa 3240
aatgatttat catgctatct ttagtatttc gtctgtgact ttctactttt gttatgttag 3300
tttgatattt taattattaat aaaaatgtat catatgtcac gttatgtatc atcacaaact 3360
aactgattgt ttattttatt ttgtcaattt gttttcaact gtaatctatt ataatttatt 3420
gtatttatgg tatgttacac caacacttta ttatgttttt cgttgagtgg agggcctagt 3480
gggaacaatc ttcttacctt caaggtaggg gtaagatctg tgataacact acctcctca 3540
atcttcactt gtggattata ctggatatata tatgttgtat ttcttaaaaa agttgattag 3600
tagaaattaa ttatgttggt catcagactc tccaaaaata ttatcgctac aatgtgtaat 3660
atttagttcc ttttgaagga gcacaagcta agggaaaaag gtccacagaa ttggcaaaaa 3720
atthtattat taccacaataa ggcatatgca cttgtgggtg gttttagaag atgggtgacc 3780
aaatatggag gtgcccagtgt gatttggggc acaaaattta ctggtgactt gcaaccaact 3840
cctgctaggg aacaactttt ggacagggtg gtatatagct cgtactttaa ttgtgtgaat 3900
cgtcttatct aaattattata cttgttatat gaatgcatag gtactggaca catacaataa 3960
attgtagcag ttgtagcaga gcatataaaa gtctaaatgt ccttgaatc attatgcaaa 4020
ttatctctgt tgcctcaatt ggaattgctg ctgcagcaaa ggagatgtgc atgtcaattg 4080
ctgcaagata ttcatgtgtc ttcttggcat tactatgctt catggcttcc agatggttat 4140
ccaaatttat atacaaaagt ttcattttcc atgattatga tcatgccttt tgttaaatgg 4200
tgtactatgt aatagatttg gagatcctaa acattatgta ttattgagga tattgtgtta 4260
tgaataagat ttctccgtca gtaa 4284

```

```

SEQ ID NO: 46      moltype = DNA length = 1632
FEATURE           Location/Qualifiers
source            1..1632
                  mol_type = genomic DNA
                  organism = Solanum tuberosum

```

```

SEQUENCE: 46
atggaagctc ttaaaatttc tacttgtttt ccaccatttc tcttcaactt gaaaacacct 60
agattttcaa gaatttatat tgagaagaaa cgaacttttt ctttttctct aaatcatcaa 120
caaccocaca agtcaagatt caactctttc actaccaata ttttaatttc aactaatgaa 180
ccacaacaac tacttccaaa tgatgaacaa gaaattagta ctaaaaatga aaaaaatcaa 240
gaaaagagaa aatttgattg gtatgcacaa tggatcccaa taatgccact aagtgaactt 300
gataagagaa ggcacacatg gaaaaaagtg atgggaattg atttagagt gtggtgggag 360
aaaaatttgg aggaattggag agtgatggat gatgcttgtt ctcatagatt ggctccactt 420
tctcaaggga gaattgatca atggggaaga ttgcagtgtg tgatcatagg ttggtgtttt 480
agtggttctg gtgattgcaa gttcattcct caggctccta gagacaagct tcatgttcac 540
acatccaaaa gagcatgtgt agcagtgtat ccaagttttg tgcaaaatga cattcttttg 600
ttttggccca acactgatct ttatatacag gacatacact tgagcaaaac cccaccttat 660
attccacctt tagatgattt aacttcatat gcaaaaacaa cacttgtag agacatccca 720
tatgggtatg agtttttgat tgaaaacctc atggacccat ctcatgtcaa ttatgcacat 780
catggcataa tgaaaatttg gaaaatagaa gttccaaaca gtgtgaaggg tgatagagaa 840
ggaggaaaac cacttgatat aagccttgaa aaattagaca taaatggatt tattgcaaa 900
caaggacatg atgaacacaa atttattgcc ccttgtgtgt actatggtcc atttgggtgt 960
caaagctatt tggataatta tgaatcaaag gaagaatcat catcaaatga tacaataga 1020
atatttctag tttttatg tggtccagta agtccaggtt attgcagatt gatgatgaca 1080
tccttcagaa actttgctgg ttgggagtat aaactatttc caccatggaa atttcacctt 1140
gacaataacc taatcattga ttctgattha tattacttcc atcttcagga gcacaagcta 1200
agggaaaaag gtccacagaa ttggcaaaaa atttgttatg taccaacaaa ggcagatgca 1260
cttggtggtg gttttagaag atgggtgacc aaatatggag gtgcccaggt tgattggggc 1320
acaaaattta ctggtgactt gcaaccaact cctgctaggg aacaactttt ggacagggtac 1380
tggaacacata caataaattg tagcagttgt agcagagcat ataaaagtct aaatgtcctt 1440
gaaatcatta tgcaaaattat ctctgtgtgt tcaattggaa ttgctgctgc agcaaggag 1500
agtgtcatgt caattgtgtc aagatattca ttggtcttct tggcattact atgcttcatg 1560
gcttccagat gggtatccaa atttatatac aaaagtttcc atttccatga ttatgatcat 1620
gcctttgttt aa 1632

```

```

SEQ ID NO: 47      moltype = DNA length = 792
FEATURE           Location/Qualifiers
source            1..792
                  mol_type = genomic DNA
                  organism = Solanum tuberosum

```

```

SEQUENCE: 47
atggctgato gcgtacatcc tcgagattca tcgccggcga gttcaccgcc atcgtaaat 60
aactccggcg aagtagccgc gggaaactaa acaaagcatg tgcggctgcg ggaacggtat 120
gtgtttcaag taccaaaaga tcaaatctac cggatccctc caccggggaa ttctgcctg 180
tacgaagctt tgagaaaaac aaagcctcgt cggagcttct gttgccgggt cgtttgctac 240

```


-continued

```

actttctctc tcttctaat tctcattatc gcacttggaa tcaactgtgc cgttctctac 300
ctcgtcttcc gtcctgaagc tccaaaatac actatatcca acgtcgcgat taagaatttc 360
aacttaactt cgtcgtctcc agtatcgccg gaattcgacg ttactgtccg agctgaaaat 420
cctaacaata agatcggaat ttactaccgg aaaggtagct ccgtcaccgt attctactcc 480
gatgtccgcc tctctaacgg cgaattgccg gcgttctatc agccaacgaa taacgtaacg 540
gtttttcaga cgccgttaaa aggatcaaac gtcttgcttg gtaacgccgt taagacggcg 600
ttaaggaatg aacagttgaa agggaaagt ccggttaagg ttaacatcaa agcgcccggt 660
aaagttaaaag ttggcgccgt taagatgtgg gaaatcacgg ttaagggtta gtgtgacata 720
acgggtgaata cattaacggc caaatcaaaa ataatttctg aagattgtaa ataatgtgtt 780
aggctttggt ag 792

```

```

SEQ ID NO: 48      moltype = DNA length = 792
FEATURE
source            Location/Qualifiers
                  1..792
                  mol_type = genomic DNA
                  organism = Solanum tuberosum

```

```

SEQUENCE: 48
atggctgatac ggcgtacatcc tgcgagattca tgcgcggcga gttcaccgcc atcgtaaaat 60
aaactccggcg aagtagccgc gggaaactaat acaaagcatg tgcggtcgcc gggaaactat 120
gttgttcaag taccaaaaga tcaaatctac cggatccctc caccggggaa ttctcgccgt 180
tacgaagctt tgagaaaacg aaagcctcgt cggagcttct gttgcgggtg cgtttgctac 240
actttctctc tcttctaat tctcattatc gcacttggaa tcaactgtgc cgttctctac 300
ctcgtcttcc gtcctgaagc tccaaaatac actatatcca acgtcgcgat taagaatttc 360
aacttaactt cgtcgtctcc agtatcgccg gaattcgacg ttactgtccg agctgaaaat 420
cctaacaata agatcggaat ttactaccgg aaaggtagct ccgtcaccgt attctactcc 480
gatgtccgcc tctctaacgg cgaattgccg gcgttctatc agccaacgaa taacgtaacg 540
gtttttcaga cgccgttaaa aggatcaaac gtcttgcttg gtaacgccgt taagacggcg 600
ttaaggaatg aacagttgaa agggaaagt ccggttaagg ttaacatcaa agcgcccggt 660
aaagttaaaag ttggcgccgt taagatgtgg gaaatcacgg ttaagggtta gtgtgacata 720
acgggtgaata cattaacggc caaatcaaaa ataatttctg aagattgtaa ataatgtgtt 780
aggctttggt ag 792

```

```

SEQ ID NO: 49      moltype = DNA length = 3380
FEATURE
source            Location/Qualifiers
                  1..3380
                  mol_type = genomic DNA
                  organism = Solanum tuberosum

```

```

SEQUENCE: 49
gaggagggtt gactgtattt ctttgttga aggggcaaaa tccgtgtata tatagtagta 60
gtaaattgtt caccttgtta aatgttatgg tcttatcatt tgttcattcc ttatctacgt 120
tacagaatat ttaaaatata gtccgcatat ggcacttaca tgttgacata ttcatattct 180
ttatattatg tgtatggagt catttagcat ataacaaaaa gaagaagaag aaattaggag 240
caaaagattca atttttattt ttcaagaatc ttcacocctt tagtttcttg atttttcg 300
ctttttaatt aagacattca ttcaagaaat gcagtttgtt tgttgaaaaa attaaaaacg 360
aatcacccct gtttgtgtat ttgatagaa gttgtattt tcttgatctt gatcttgact 420
agtcgttttg gcgaaaaaat gagtattcag aactatgaat tagcaagtga ttgcaactta 480
gagttcccac agatgggatt ttgtttccag cctgaaaaat ctgcagaaaa tggttgtcaa 540
cagcagcagc aacagaattt ttggcctagt actgattcat catcatcgag aacgattata 600
agtcgaatag gatcatcacc ttctgctttt ttgtctacag agaggtaact tggattaaca 660
caatatgaaa accaagacaa caataatagt tgttctcaac tatccaagaa tcttggctct 720
caaaactacg cgtttactca gcaatgttga aatggattct tggcagattc atcagcacga 780
gttgacacgg attttcctaa gatttcaatg ccatcattca tcagatcaca gttttcaagt 840
agtcaaccat ttggtcttga aggactctat ggaaatccct ttagtaatct atcagagaaa 900
gagaggattt tgcttcttaa gagcaagtgt tttagagaaa ttgactcttc aaataggcag 960
cctgcttcaa tcccttttca aggaaatcaa gactatggtg taagtacttg attcttgtt 1020
acataataca gtcaaatctc tctgtaacga ttttgtttct cctgatatgt ttgggttgc 1080
atagctagat gttgttatcg agaacatcta atataacgta atatgaaagt ttgttccaaa 1140
gaaaacttgg ctggtataga gaggtctgac tgaagtatag tttttcaagt atttgaattt 1200
aagtaacttc aattgcttgt tgtctacata atacagtcaa acctctctat aacagtgttg 1260
tttgctcctga tatgttttag ttgctatagc gagatgttat tgaaaacat ctaataattc 1320
gtaatatgaa agttcttctc aaagaaaact tggccgttat agagaggcct gactgtagta 1380
ttgtttttca agtttttatt ctgactcgag ttatgttcca acagggtctc aataatacat 1440
gtggttttaa cttggtagat ataaggcaac aatctggaag tcaatcagca aatagtttta 1500
acaactctgg atgttcttga ggaatcttat cgagtaaggc acgaatcagg tggactcagg 1560
atcttcatga tcaattgttt gagtgtgtaa atcgtcttgg aggagctgac agtaagttaa 1620
tttacacatt tttttagctt ttgtttcttt ggaggtgata attttggcta tggattaatt 1680
gtttcttact tcaattgcca gaggcgacgc caaaggcaat actaaagctg atggattcag 1740
aaggattaac aatttttcat gtaaaaagtc atttacaggt acttggtta atgaaagaaa 1800
tactttcttg gaacactttt tgtatttaga caaaattctg aatcaaatgt tttttccttc 1860
ttgttttgac tagaaatcat gaaatgcaaa gttcatccct gaatcgacag aaggtagta 1920
ttatccgaaa taagcttctc gtttattatg taagagatat tcatccaca gcttgaaccc 1980
gtgacttgta gttcatcag agacaatttt atcgtttctc caaggctctc ttcattactc 2040
aagcatcagt acagatgtat gctgcttgtt ttatgctttt ttgtgtta atttctgagca 2100
aaaaaaatca aatatgtttc tccaaatata agatccttaa ccacataata gtatcctcgt 2160
cttgataaac agttaataca ccttatttcc aaacaagttg gactccgtta tatgaatcct 2220
cacgaacat gttctaaacc ttcttcgaaa aatcttctct gctttcattt ttatcattt 2280

```

-continued

```

acgcaacagg tagatccaac atgatgaaaa acacaagaag agactatttc cttaaacaaa 2340
tatgctatta gtaaatctgt tgaagtcttt gaaatctcag aagattgaaa gaacattttc 2400
atcttacttc ttatacaaat cactttaatt ttatttcgta tgatgatgat atgaaatgag 2460
ctttaaatga aggagtgggg gattcgata gcggaaccca acttgttga gactgaggcg 2520
tagttgtgtg ttaaatttga tatttacagc aaactaaaaa cttaatttta actctgcttg 2580
tggaaaaatg ctctctgaga tatactcaaa tctgtcatta catgctaagt acttaatctt 2640
aaagtttcca tctctttctg gaaatgagag ttccctcctt cctatgatgc agggagatct 2700
ggaaaaacag acagcccga taatgtgtca cagatcgaca gcaaaacgta tgttctttgc 2760
aactatcttg acaaattttc gtgactattt tagctggaaa gttaccttaa tacagatttc 2820
ttttcagtgg aatgcaaatc aaagaagcat tgcatacgca gctagaagtc cagaggcgctc 2880
ttcacgagca actagaggta catttagtag atgaaaagat ataaatttaa acacttgtag 2940
gcatgcttag agctgaaaca gtatcaaggc aatctttcta aatcttattg tctttctgtt 3000
gttagattca gcggaagtta caattgagga tcgaagaaca aggggagcag ttgaagaaga 3060
tatttgaaaca acaacaacaa acaactagga gtctcttgga gacacgaaat tcaagcattt 3120
cgtctcctgc tgatcagttc accccgcacg aagatgaagt ttttgctgca gaaagcttca 3180
ataatactca ttccaatct aatataagtt acaatgacat gtaaacacaa ttagttttac 3240
attttttcag ctagtttttg aaagagagtc gacgttagca tttctgtaaa gataattttg 3300
cctcccaagc aaactacaca aaaaaaaat gtatattaca aagtgaagac ataaatatca 3360
tgcaaaaact taaagtactt                                     3380

```

```

SEQ ID NO: 50      moltype = DNA length = 1209
FEATURE           Location/Qualifiers
source            1..1209
                  mol_type = genomic DNA
                  organism = Solanum tuberosum

```

```

SEQUENCE: 50
atgagtattc agaactatga attagcaagt gattgcaact tagagttccc acagatggga 60
ttttgtttcc agcctgaaaa ctctgcagaa aatggttgtc aacagcagca gcaacagaat 120
ttttggccta gactgatctc atcatcatcg agaacgatta taagtgcgaat aggatcatca 180
ccttctgctt tttttgctac agagaggtag ctgtgattaa cacaatatga aaaccaagac 240
aacaataata gttgtttcca actatccaag aatcttggtc ctcaaaactac gtcgtttact 300
cagcaatgtg gaaatggatt cttggcagat tcatcagcac gagttgacac cgatttttct 360
aagattttca tgccatcatt catcagatca cagttttcaa gtagtcaacc atttggtcct 420
gaaggactct atggaatccc ctttagtaat ctatcagaga aagagaggat ttgcttctt 480
aagagcaagt tgtttagaga aattgactct tcaaataggc agcctgcttc aatccctttt 540
caaggaaatc aagactatgg tgtctcaaat aatacatgtg gtttttaact ggtacatata 600
aggcaacaat ctggaagtca atcagcaaat agttttaaca actctggatg ttctggagga 660
tctttatcga gtaaggcacg aatcaggtgg actcaggatc ttcagatcg atttggttag 720
tgtgtaaatc gtcttgagg agctgacaag gcgacgcca aggcaactact aaagctgatg 780
gattcagaag gattaacaat ttttcagtga aaaagtcatt tacagaaata tcgaaatgca 840
aagttcatcc ctgaatcgac agaagggaga tctggaaaaa cagacagccc gaataatgtg 900
tcacagatcg acagcaaaac tgggaatgcaa atcaaaagag cattgcatat gcagctagaa 960
gtccagagggc gtccttcacga gcaactagag attcagcgga agttacaatt gaggatcgaa 1020
gaacaagggg agcagttgaa gaagatattt gaacaacaac aacaacaac taggagtctc 1080
ttggagacac gaaattcaag catctcgtct cctgctgac agttcacccc gcacgaagat 1140
gaagtttttg ctgcagaag cttcaataat actcatttcc aatctaatat aagttacaat 1200
gacatgtaa                                     1209

```

```

SEQ ID NO: 51      moltype = DNA length = 144
FEATURE           Location/Qualifiers
source            1..144
                  mol_type = genomic DNA
                  organism = Solanum tuberosum

```

```

SEQUENCE: 51
aagattttta gtatttgatt tatgaatgaa tttttttttt tgagaaatat tttttatttt 60
tactagagta gaaaataatt ttgaaattg aaaatagttt ttaaaaaaca acttaaatTT 120
ttttattttt ttttgggggt gggc                                     144

```

What is claimed is:

1. A plant comprising plant cells comprising a modification to an endogenous gene, wherein the polypeptide encoded by the endogenous gene interacts with a Sec-dependent effector (SDE) secreted by a bacterial species from the genus *Ca. Liberibacter*, wherein the modification knocks-down or reduces expression of the endogenous gene and/or interrupts interaction of the polypeptide with an SDE and confers resistance or tolerance to *Ca Liberibacter* infection in the plant relative to a plant of the same variety lacking the modification, and wherein the endogenous gene encodes PP2-B2/12 and the modification comprises a modification to the endogenous gene comprising;

insertion of at least 1 nucleotide,
a deletion of at least 1 nucleotide, and/or
a substitution of at least 1 nucleotide.

2. The plant of claim 1, wherein the plant is *Citrus*.

3. The plant of claim 1, wherein the modification is made to at least one of SEQ ID NO: 30, 31, or a sequence comprising at least 95% identity therewith.

4. The plant of claim 1, wherein the plant is a solanaceous plant.

5. The plant of claim 4, wherein the plant is *Solanum tuberosum*.

6. The plant of claim 1, wherein the plant is a grapefruit tree, an orange tree, a sweet orange tree, a lime tree,

citrumelo tree, trifoliolate tree, reticulata tree, aurantium tree, lemon tree, a papeda tree, a pummelo tree or a mandarin orange tree.

7. The plant of claim 1, wherein the SDE is Las4025.

8. A seed that produces the plant of claim 1.

9. A plant part of the plant of claim 1, wherein the plant part comprises the modification.

10. A method of generating a modified *Citrus* plant comprising resistance or tolerance to infection by a bacterial species from the genus *Ca. Liberibacter*; the method comprising the steps of:

(a) modifying a PP2-B2/12 gene of a *Citrus* plant cell such that expression of the PP2-B2/12 gene is knocked-down or reduced and/or interaction of the polypeptide encoded by said PP2-B2/12 gene with a Sec-dependent effector (SDE) secreted by a bacteria species from the genus *Ca. Liberibacter* is reduced; and

(b) regenerating the modified plant from said plant cell or a progenitor cell thereof, wherein said the plant comprises said modification.

11. The method of claim 10, wherein the plant is *Citrus*.

12. The method of claim 11, wherein the plant is a solanaceous plant.

13. A plant comprising resistance to *Ca. Liberibacter* infection produced by the method of claim 10.

14. The method of claim 10, effects of a Sec-dependent effector (SDE) secreted by a bacteria species from the genus *Ca. Liberibacter* are reduced.

15. The method of claim 14, wherein the SDE comprises Las4025.

16. The method of claim 10, wherein step (a) comprises a genome-editing technique.

17. The method of claim 16, wherein the genome-editing technique comprises a use of a nuclease, wherein the nuclease introduces a single-strand DNA break or a double-strand DNA break.

18. The method of claim 16, wherein the genome-editing technique comprises use of a Talen, a ZFN, meganuclease, or a CRISP/Cas system.

* * * * *