



US 20250276002A1

(19) **United States**

(12) **Patent Application Publication**
Salaita et al.

(10) **Pub. No.: US 2025/0276002 A1**

(43) **Pub. Date: Sep. 4, 2025**

(54) **PARTICLES WITH RNA CLEAVING
NUCLEOBASE POLYMERS AND USES FOR
MANAGING INFLAMMATORY DISORDERS**

(71) Applicants: **Emory University**, Atlanta, GA (US);
**The United States Government as
represented by the Department of
Veterans Affairs**, Washington, DC (US)

(72) Inventors: **Khalid Salaita**, Decatur, GA (US);
Cherry Wongtrakool, Atlanta, GA
(US); **Kornelia Galior**, Atlanta, GA
(US)

(21) Appl. No.: **19/211,934**

(22) Filed: **May 19, 2025**

Related U.S. Application Data

(60) Division of application No. 17/122,946, filed on Dec. 15, 2020, which is a continuation of application No. 16/304,549, filed on Nov. 26, 2018, filed as application No. PCT/US2017/034268 on May 24, 2017, now Pat. No. 10,905,710.

(60) Provisional application No. 62/340,586, filed on May 24, 2016.

Publication Classification

(51) **Int. Cl.**

A61K 31/713 (2006.01)

A61K 9/00 (2006.01)

A61K 9/14 (2006.01)

A61K 9/51 (2006.01)

A61K 31/712 (2006.01)

A61K 45/06 (2006.01)

A61P 11/06 (2006.01)

B82Y 5/00 (2011.01)

C07H 21/04 (2006.01)

(52) **U.S. Cl.**


CPC *A61K 31/713* (2013.01); *A61K 9/007*
(2013.01); *A61K 9/0073* (2013.01); *A61K*
9/0075 (2013.01); *A61K 9/0078* (2013.01);
A61K 9/008 (2013.01); *A61K 9/141* (2013.01);
A61K 9/51 (2013.01); *A61K 31/712* (2013.01);
A61K 45/06 (2013.01); *A61P 11/06* (2018.01);
B82Y 5/00 (2013.01); *C07H 21/04* (2013.01)

(57)

ABSTRACT

This disclosure relates to nucleobase polymers useful for degrading GATA-3 mRNA. In certain embodiments, this disclosure relates to nucleobase polymers and nanoparticles conjugated to nucleobase polymers disclosed herein. In certain embodiments, the nucleobase polymers or nanoparticles can be used in methods of managing disorders associated with excessive GATA-3 expression in inflammatory disorders and respiratory disorders such as asthma.

Specification includes a Sequence Listing.






FIG. 1




FIG. 2




FIG. 3A




FIG. 3B




FIG. 4A




FIG. 4B


Average $2^{-\Delta\Delta C_t}$ for Dz-Xs on AuNP compared to Hgd40-DzNP

FIG. 5




FIG. 6A




FIG. 6B




FIG. 6C




FIG. 7

PARTICLES WITH RNA CLEAVING NUCLEOBASE POLYMERS AND USES FOR MANAGING INFLAMMATORY DISORDERS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application is a division of U.S. application Ser. No. 17/122,946 filed Dec. 15, 2020, which is a continuation of U.S. application Ser. No. 16/304,549 filed Nov. 26, 2018 that granted as U.S. Pat. No. 10,905,710 on Feb. 2, 2021, which is the National Stage of International Application No. PCT/US2017/034268 filed May 24, 2017, which claims the benefit of U.S. Provisional Application No. 62/340,586 filed May 24, 2016. The entirety of each of these applications is hereby incorporated by reference for all purposes.

INCORPORATION-BY-REFERENCE OF MATERIAL SUBMITTED AS AN XML FILE VIA THE OFFICE ELECTRONIC FILING SYSTEM

[0002] The Sequence Listing associated with this application is provided in XML format and is hereby incorporated by reference into the specification. The name of the XML file containing the Sequence Listing is 16116USDIV.xml. The XML file is 53,036 bytes, was created on May 15, 2025, and is being submitted electronically via the USPTO patent center.

BACKGROUND

[0003] In asthma patients, airway obstruction is due to mucus secretion and airway inflammation. A number of medications can be used to treat asthma attacks, but managing persistent severe allergic asthma is problematic. Omalizumab is a recombinant humanized monoclonal antibody that specifically binds to human immunoglobulin E (IgE) and used in patients with severe persistent allergic asthma. However, systemic omalizumab administration may result in anaphylaxis or the generation of inactivating antibodies. Thus, there is a need to identify improved therapeutic methods to control persistent allergic asthma.

[0004] Allergic asthma results in T-helper (TH)-2 driven responses. Activation of TH2 cells produces cytokines such as IL-4, IL-5, and IL-13. Expression of these cytokines are dependent on the zinc finger transcription factor GATA-3. GATA-3 overexpression is observed in patients with severe asthma. Molecules that inactivate GATA-3 have been reported. See Sel et al. Effective prevention and therapy of experimental allergic asthma using a GATA-3-specific DNzyme, *J Allergy Clin Immunol.* 2008, 121 (4): 910-916.

[0005] Krug et al. report allergen-induced asthmatic responses modified by a GATA-3-specific DNzyme. *The New England Journal of Medicine*, 2015, 372 (21): 1987-1995. See also WO/2016/184556, WO/2014/040891, and WO/2005/033314.

[0006] Somasuntharam et al. report the knockdown of TNF-alpha by DNzyme gold nanoparticles as an anti-inflammatory therapy for myocardial infarction. *Biomaterials*, 2016, 83:12-22. Yehl et al. report catalytic deoxyribozyme-modified nanoparticles for RNAi-independent gene regulation. *ACS Nano*, 2012, 6 (10): 9150-7.

[0007] References cited herein are not an admission of prior art.

SUMMARY

[0008] This disclosure relates to nucleobase polymers useful for degrading GATA-3 mRNA. In certain embodiments, this disclosure relates to nucleobase polymers and nanoparticles conjugated to nucleobase polymers disclosed herein. In certain embodiments, the nucleobase polymers or nanoparticles can be used in methods of managing disorders associated with excessive GATA-3 expression in inflammatory disorders and respiratory disorders such as asthma.

[0009] In certain embodiments, the disclosure relates to nucleobase polymers comprising or consisting of SEQ ID NO: 1-49, 53-55 or variants thereof.

[0010] In certain embodiments, the nucleobase polymer comprises

[0011] SEQ ID NO: 10 (GGCTTATTCAGGCTAGCTACAACGAAGATGGGG), SEQ ID NO: 20 (ATTCTTAAAGGCTAGCTACAACGATTCTTGCC), or SEQ ID NO: 30 (TCTTTCTTAGGCTAGCTACAACGATTGCTGC) or variants thereof.

[0012] In certain embodiments, the disclosure contemplates a particle coated with or conjugated to a nucleobase polymer disclosed herein, e.g., SEQ ID NO: 1-51, 53-55 or variants thereof. In certain embodiments, the nucleobase polymers of disclose herein have an RNA cleaving sequence such as the 10-23 DNzyme with SEQ ID NO: 51 (GGCTAGCTACAACGA), e.g., contained within SEQ ID NO: 1-50, 53-55. In certain embodiments, the cleaving sequence has monomers of 2-deoxyribose.

[0013] In certain embodiments, the variant is a nucleobase polymer comprising one nucleobase substitution, insertion, or deletion. In certain embodiments, the variant is a nucleobase polymer comprising two nucleobase substitutions, insertions, or deletions. In certain embodiments, the variant is a nucleobase polymer comprising three or nucleobase substitutions, insertions, or deletions.

[0014] In certain embodiments, the variant is a nucleobase polymer comprising one 5' end nucleobase substitution or deletion. In certain embodiments, the variant is a nucleobase polymer comprising two 5' end nucleobase substitutions or deletions. In certain embodiments, the variant is a nucleobase polymer comprising three 5' end nucleobase substitutions or deletion.

[0015] In certain embodiments, the variant is a nucleobase polymer comprising one 3' end nucleobase substitutions or deletions. In certain embodiments, the variant is a nucleobase polymer comprising two 3' end nucleobase substitutions or deletions. In certain embodiments, the variant is a nucleobase polymer comprising three 3' end nucleobase substitutions or deletions.

[0016] In certain embodiments, the disclosure relates to particles coated with a nucleobase polymer comprising an RNA cleaving sequence such as a nucleobase polymer comprising SEQ ID NO: 51 (GGCTAGCTACAACGA) and linked to flanking 5' and 3' nucleobases that hybridize with SEQ ID NO: 52.

[0017] In certain embodiments, the particle or nanoparticle has a core or hydrodynamic diameter between 5 nm and 500 nm or 5 nm and 200 nm. In certain embodiments, the particle may contain a metal or inorganic or polymer core. In certain embodiments, conjugation to a particle can be accomplished by using linking groups with ligands, e.g., mono or polydentate ligands, capable of binding the metal in the core or by direct conjugation through covalent bonds of a polymer surrounding and encapsulating the core. In certain

embodiments, the nucleobase polymer may further contain a targeting nucleobase polymer sequence or targeting nucleic acid sequence that can hybridize with a target sequence coated on the exterior of the particle.

[0018] In certain embodiments, the disclosure relates to aerosols, liquid particles, mixtures or gas and liquid particles, comprising a nucleobase polymer or a particle disclosed herein. In certain embodiments, the liquid particle has a diameter of between 1 micron and 5 microns, 0.5 microns and 10 microns, 0.1 microns and 50 microns, or 0.5 microns and 100 microns.

[0019] In certain embodiments, this disclosure relates to a nucleobase polymer or a particle disclosed herein in the form of a micronized powder, e.g., solid particles of a diameter between 1 microns and 5 microns, 0.5 microns and 10 microns, 0.1 microns and 50 microns, or 0.5 microns and 100 microns.

[0020] In certain embodiments, this disclosure relates to pharmaceutical composition comprising a nucleobase polymer disclosed herein or a particle disclosed herein and a pharmaceutically acceptable excipient. In certain embodiments, the pharmaceutical composition comprises a sterilized pH buffered aqueous salt solution or an isotonic aqueous buffer solution.

[0021] In certain embodiments, the disclosure relates to a container, optionally sealed gas tight, comprising a nucleobase polymer or particle disclosed herein. In certain embodiments, the container further comprises a propellant. In certain embodiments, the container is configured with a spraying or misting apparatus, mouthpiece, or facemask such as a nebulizer or inhaler. In certain embodiments, the inhaler comprises a liquefied gas propellant. In certain embodiments, the nucleobase polymer or particle is dissolved or suspended in the propellant. In certain embodiments, the inhaler is configured with an actuator or mouthpiece, which allows the patient to operate the device, and directs the aerosol, a chamber, and metering valve that controls a metered quantity of a formulation to be dispensed with an actuation.

[0022] In certain embodiments, the disclosure relates to methods of treating an inflammatory disease comprising administering an effective amount of a nucleobase polymer or particle disclosed herein to a subject in need thereof. In certain embodiments, the inflammatory disease is a respiratory disorder, asthma, COPD, atopic dermatitis, psoriasis, or ulcerative colitis. In certain embodiments, the subject is diagnosed with, exhibiting symptoms of or at risk of asthma, COPD, bronchitis, emphysema, chronic obstructive pulmonary disease, laryngitis, or cystic fibrosis.

[0023] In certain embodiments, administration is in combination with a second respiratory agent. In certain embodiments, the second respiratory agent is a corticosteroid, bronchodilator, albuterol, ipratropium, or combinations thereof.

BRIEF DESCRIPTION OF THE DRAWINGS

[0024] FIG. 1 shows a plot quantifying inhaled delivery of GATA-3 DzNPs using fluorescence imaging. GATA-3 DzNPs were nebulized with even and distal distribution in the lung. Vehicle (phosphate buffered saline, 50 μ l of) and 50 μ l of fluorescently tagged DzNPs (100 nM, two lungs on right) were administered to mice by nebulization into 4-6 um

sized aerosol droplets using a bias flow of 2 liters/min of room air. Imaging was performed 24 hours following delivery.

[0025] FIG. 2 shows a plot quantifying the uptake of DzNPs in lung resident cell lines. The data was collected using flow cytometry following incubation with 10 nM concentrations of Cy5-labeled DzNPs in epithelial, macrophage, and smooth muscle cells. Cells were incubated in standard serum-containing media for 24 hours without the use of any transfection agent. Flow cytometry confirms resident lung cell uptake of DzNPs.

[0026] FIG. 3A shows the results of performing a pulmonary function test on four groups of animals. Two groups received daily doses of the house dust mite (HDM) allergen for two weeks, while two groups were control animals and did not receive the HDM. This allergen induced airway resistance in mice and recapitulates the pathophysiology of asthma. Animals received either daily doses of the active DzNPs or an inactive DzNP. The plot shows the raw data measuring the airway resistance as a function of a methacholine challenge. There were three animals in each group.

[0027] FIG. 3B shows data summarizing the efficacy of active GATA-3 Dz compared to non-specific (NS) DzNPs when challenged with 100 mg/ml methacholine. House dust mite (HDM) extract (50 ug) was inhaled via nebulization daily for 5 days. AuNP (20 μ l) administered intranasally (100 nM of DzNPs).

[0028] FIG. 4A shows RT-PCR quantification of GATA-3 expression levels in cell lines following DNase treatment. The screen shows DNases that are more active than HGD40. The dotted line indicates the GATA-3 levels in HGD40 sequence treated cells. Arrows indicate the DNase sequences that showed lower levels of GATA-3 compared to the soluble HGD40 sequence is SEQ ID NO: 50. All of the DNases have a 3'-end T10 modification.

[0029] FIG. 4B shows RT-PCR quantification of GATA-3 expression levels in cells treated with with HGD40 sequence with certain modifications. HGD40 9 methyl refers to SEQ ID NO: 53, (G*-T*-G*-G*-dA-dT-dG-dG-dA-dG-dG-dC-dT-dA-dG-dC-dT-dA-dC-dA-dA-dC-dG-dA-dG-dT-dC-dT-dT-G*-G*-A*-G*-T10-3'-3'T) wherein d is deoxy and * is 2'-O-methyl ribose. HGD40 8 methyl refers to SEQ ID NO: 54, (T*-G*-G*-A*-dT-dG-dG-dA-dG-dG-dC-dT-dA-dG-dC-dT-dA-dC-dA-dA-dC-dG-dA-dG-dT-dC-dT-T*-G*-G*-A*-T10-3'-3'T) wherein d is deoxy and * is 2'-O-methyl ribose. HGD40 7 methyl refers to SEQ ID NO: 55, (G*-G*-A*-T*-dG-dG-dA-dG-dG-dC-dT-dA-dG-dC-dT-dA-dC-dA-dA-dC-dG-dA-dG-dT-dC-T*-T*-G*-G*-T10-3'-3'T) wherein d is deoxy and * is 2'-O-methyl ribose.

[0030] FIG. 5 shows RT-PCR data on GATA-3 expression levels following DNase-AuNP treatment (10 nM for 24 hrs). Several DzNPs were shown to be more active than HGD40-NPs.

[0031] FIG. 6A shows the results of performing a pulmonary function test on the HDM model of mouse asthma. There were four groups of animals with three animals within each group. One group did not receive the HDM allergen and was considered as a control. The second group received the HDM allergen for 3 weeks and was not rescued using the DzNP drug treatment. The third group received the HGD40-NP along with the HDM allergen. The final group received the DzNP with Dz2251 along with the HDM allergen. The animals were challenged using 25 mg/mL methacholine.

Airway resistance was measured in mouse model of asthma. The DzNP were delivered intranasally for 2 weeks (100 nM, 20 uL daily).

[0032] FIG. 6B shows data using 50 mg/mL methacholine.

[0033] FIG. 6C shows data using 100 mg/mL methacholine.

[0034] FIG. 7 plots RT-PCR levels of GATA-3 mRNA in the lungs of the animals (from FIG. 6).

DETAILED DISCUSSION

[0035] Before the present disclosure is described in greater detail, it is to be understood that this disclosure is not limited to particular embodiments described, and as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present disclosure will be limited only by the appended claims.

[0036] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. Although any methods and materials similar or equivalent to those described herein can also be used in the practice or testing of the present disclosure, the preferred methods and materials are now described.

[0037] All publications and patents cited in this specification are herein incorporated by reference as if each individual publication or patent were specifically and individually indicated to be incorporated by reference and are incorporated herein by reference to disclose and describe the methods and/or materials in connection with which the publications are cited. The citation of any publication is for its disclosure prior to the filing date and should not be construed as an admission that the present disclosure is not entitled to antedate such publication by virtue of prior disclosure. Further, the dates of publication provided could be different from the actual publication dates that may need to be independently confirmed.

[0038] As will be apparent to those of skill in the art upon reading this disclosure, each of the individual embodiments described and illustrated herein has discrete components and features which may be readily separated from or combined with the features of any of the other several embodiments without departing from the scope or spirit of the present disclosure. Any recited method can be carried out in the order of events recited or in any other order that is logically possible.

[0039] Embodiments of the present disclosure will employ, unless otherwise indicated, techniques of medicine, organic chemistry, biochemistry, molecular biology, pharmacology, and the like, which are within the skill of the art. Such techniques are explained fully in the literature.

[0040] It must be noted that, as used in the specification and the appended claims, the singular forms “a,” “an,” and “the” include plural referents unless the context clearly dictates otherwise.

[0041] The transitional term “comprising,” which is synonymous with “including,” “containing,” or “characterized by,” is inclusive or open-ended and does not exclude additional, unrecited elements or steps, e.g., does not exclude the presence of terminal nucleotides. The transitional phrase “consisting of” excludes any additional nucleotides, elements, steps, or ingredients not specified in the claim.

[0042] As used herein, the term “combination with” when used to describe administration with an additional treatment means that the agent may be administered prior to, together with, or after the additional treatment, or a combination thereof.

[0043] As used herein, the terms “prevent” and “preventing” include the prevention of the recurrence, spread or onset. It is not intended that the present disclosure be limited to complete prevention. In some embodiments, the onset is delayed, or the severity is reduced.

[0044] As used herein, the terms “treat” and “treating” are not limited to the case where the subject (e.g., patient) is cured and the condition or disease is eradicated. Rather, embodiments, of the present disclosure also contemplate treatment that merely reduces symptoms, and/or delays conditions or disease progression.

[0045] As used herein, the term “nucleic acid” is intended to mean a ribonucleic or deoxyribonucleic acid or analog thereof, including a nucleic acid analyte presented in any context; for example, a probe, target or primer. A nucleic acid can include native or non-native bases. In this regard, a native deoxyribonucleic acid can have one or more bases selected from the group consisting of adenine, thymine, cytosine or guanine and a ribonucleic acid can have one or more bases selected from the group consisting of uracil, adenine, cytosine or guanine. It will be understood that a deoxyribonucleic acid used in the methods or compositions set forth herein can include uracil bases and a ribonucleic acid can include a thymine base. Exemplary non-native bases that can be included in a nucleic acid, whether having a native backbone or analog structure, include, without limitation, inosine, xanthine, hypoxanthine, isocytosine, isoguanine, 2-aminopurine, 5-methylcytosine, 5-hydroxymethyl cytosine, 2-aminoadenine, 6-methyl adenine, 6-methyl guanine, 2-propyl guanine, 2-propyl adenine, 2-thioLiracil, 2-thiothymine, 2-thiocytosine, 15-halouracil, 15-halocytosine, 5-propynyl uracil, 5-propynyl cytosine, 6-azo uracil, 6-azo cytosine, 6-azo thymine, 5-uracil, 4-thiouracil, 8-halo adenine or guanine, 8-amino adenine or guanine, 8-thiol adenine or guanine, 8-thioalkyl adenine or guanine, 8-hydroxyl adenine or guanine, 5-halo substituted uracil or cytosine, 7-methylguanine, 7-methyladenine, 8-azaguanine, 8-azaadenine, 7-deazaguanine, 7-deazaadenine, 3-deazaguanine, 3-deazaadenine or the like. A particular embodiment can utilize isocytosine and isoguanine in a nucleic acid in order to reduce non-specific hybridization, as generally described in U.S. Pat. No. 5,681,702.

[0046] A non-native base used in a nucleic acid can have universal base pairing activity, wherein it is capable of base pairing with any other naturally occurring base. Exemplary bases having universal base pairing activity include 3-nitropyrrole and 5-nitroindole. Other bases that can be used include those that have base pairing activity with a subset of the naturally occurring bases such as inosine, which base-pairs with cytosine, adenine or uracil. Alternatively, or additionally, oligonucleotides, nucleotides or nucleosides including the above-described non-native bases can further include reversible blocking groups on the 2', 3' or 4' hydroxyl of the sugar moiety.

[0047] The terms “binding,” “binds,” “recognition,” or “recognize” as used herein are meant to include interactions between molecules that may be detected using, for example, a hybridization assay. When hybridization occurs in an antiparallel configuration between two single-stranded poly-

nucleotides, the reaction is called “annealing” and those polynucleotides are described as “complementary”. A double-stranded polynucleotide can be complementary or homologous to another polynucleotide, if hybridization can occur between one of the strands of the first polynucleotide and the second. Complementarity or homology (the degree that one polynucleotide is complementary with another) is quantifiable in terms of the proportion of bases in opposing strands that are expected to form hydrogen bonding with each other, according to generally accepted base-pairing rules.

[0048] A “linking group” refers to any variety of molecular arrangements that can be used to bridge to molecular moieties together. An example formula may be —Rm— wherein R is selected individually and independently at each occurrence as: —CRmRm—, —CHmRm—, —CH—, —C—, —CH₂—, —C(OH)Rm—, —C(OH)(OH)—, —C(OH)H—, —C(Hal)Rm—, —C(Hal)(Hal)—, —C(Hal)H—, —C(N₃)Rm—, —C(CN)Rm—, —C(CN)(CN)—, —C(CN)H—, —C(N₃)(N₃)—, —C(N₃)H—, —O—, —S—, —N—, —NH—, —NRm—,

[0049] —(C=O)—, —(C=NH)—, —(C=S)—, —(C=CH₂)—, which may contain single, double, or triple bonds individually and independently between the R groups. If an R is branched with an Rm it may be terminated with a group such as —CH₃, —H, —CH=CH₂, —CCH, —OH, —SH, —NH₂, —N₃, —CN, or —Hal, or two branched Rs may form a cyclic structure. It is contemplated that in certain instances, the total Rs or “m” may be less than 100 or 50 or 25 or 10. Examples of linking groups include bridging amide, alkyl, and alkoxyalkyl groups.

Targeting the Transcription Factor GATA-3 to Modulate the Inflammatory Response

[0050] The most prevalent type of asthma (50% of patients) displays a type 2 helper T cell (Th2) endotype, which is characterized by an overabundance of Th2 cells that express inflammatory cytokines such as Il-4, Il-5, and Il-13. GATA-3 is the major transcription factor involved in driving differentiation of helper T cells toward the Th2 fate. Moreover, GATA-3 has been found to be upregulated in biopsies and serum from patients with severe asthma, even while on oral corticosteroids. GATA-3 is expressed in resident lung cell types, such as mast cells, eosinophils, macrophages, fibroblasts, and airway epithelial cells.

[0051] Inhibiting GATA-3 production by gene knockdown has been shown to suppress upregulation of cytokines and prevent the induction of airway hyper-responsiveness amongst other asthma symptoms. A recent phase II clinical trial demonstrated that administering GATA-3 DNazymes (Dzs) by inhalation can reduce the impact of an allergen challenge on FEV1 (forced expiratory volume) in mild asthmatics. After treatment with Dz targeting GATA-3, both the early and late asthmatic responses are attenuated, improving overall airway function. This emphasized the importance of GATA-3 in regulating the asthmatic response. However, allowing an oligonucleotide to pass across the plasma membrane is challenging. Accordingly, clinical trials require 2 mg doses of oligonucleotides per patient per day.

[0052] Dzs are synthetic, catalytically active DNA anti-sense molecules that bind to and cleave specific mRNA. They possess a central catalytic domain that typically ranges from 8-15 nucleotides. Flanking this site are two variable domains, typically 8-12 nucleotides each that can be

designed to hybridize to the mRNA molecule of interest. By screening sites in the GATA-3 mRNA, a highly active Dz was discovered that target and cleave the GATA-3 gene. Cytoplasmic RNAses further degrade the cleavage products. **[0053]** To overcome the challenges of delivering Dzs as gene regulation agents, Dz nanoparticle conjugates were developed for gene regulation. Copies of Dz molecules immobilized onto a 14 nm gold nanoparticle (DzNP) through the 3' terminus are highly active for mRNA knock down. These particles can rapidly enter cells, protect the DNazyme against nucleases, and regulate gene expression in vitro. DzNPs show minimal off-target effects and did not require the use of any transfection agents. Therefore, DzNPs offer an attractive approach for delivering catalytic nucleic acids to treat disease.

Gata-3 mRNA and Nucleobase Polymers Containing DNazyme Sequences

[0054] *Homo sapiens* GATA binding protein 3 (GATA-3), transcript variant 1, mRNA has the NCBI Reference Sequence NM_001002295.1 (SEQ ID NO: 52):

```
GGCGCCGTCCTTGATACTTTCAGAAAGAAATGCATTCCCTGTAAAAA
AAAAAAATACTGAGAGAGGGAGAGAGAGAGAAGAAGAGAGAGAGA
CGGAGGGAGAGCGAGACAGAGCGAGCAACGCAATCTGACCGAGCAGGT
CGTACGCCGCCGCCCTCCTCCTCCTCTCTGCTCTTCGCTACCCAGGTGA
CCCAGGAGGGGACTCCGCTCCGAGCGGCTGAGGACCCCGGTGCAGAG
GAGCCTGGCTCGCAGAATTGCAGAGTCGTCGCCCTTTTACAACCTG
GTCCCGTTTATTCTGCCGTACCCAGTTTTTGGATTTTTGTCTTCCCC
TTCTTCTCTTTGCTAAACGACCCCTCCAAGATAATTTTAAAAAACCT
TCTCCTTTGCTCACCTTTGCTTCCCAGCCTTCCCATCCCCCACCAGAA
AGCAATATTCATCAACGACCCCGACCCCTCCGACGGCAGGAGCCCCCG
ACCTCCAGGCGGACCGCCCTCCCTCCCGCGCGCGGTTCCGGGCC
GGCGAGAGGGCGCGAGCACAGCCGAGGCCATGGAGGTGACGGCGGACC
AGCCGCGCTGGGTGAGCCACCACCCCGCCGCTGCTCAACGGGCGAGC
ACCCGGACACGACCAACCCGGGCTCAGCCACTCTACATGGACGCGG
CGCAGTACCCGCTGCCGAGGAGGTGGATGTGCTTTTAAACATCGACG
GTCAAGGCAACCAGCTCCCGCCTACTACGGAACTCGGTACGGGCCA
CGGTGACAGAGGTACCCTCCGACCCACACGGGAGCAGGTGTGCCGCC
CGCCTCTGCTTCATGGATCCCTACCCCTGGCTGGACGGCGGCAAAGCCC
TGGGCGACCCACACCCGCTCCCGCTGGAATCTCAGCCCTTCTCCA
AGACGTCATCCACACGCTCCCGGGGGCCCTCTCCGTCTACCCCC
CGGCCTCGTCTCTCTTGTGCGGGGGCCACGCCAGCCCGCACCTCT
TCACCTTCCCGCCACCCCGCGAAGGAGCTCTCCCGGACCCATCGC
TGTCACCCCGAGGCTCGGCCGCTCGGCCCGGACGAGAGAAAGAGT
GCCTCAAGTACCAGGTGCCCTGCCGACAGCATGAAGCTGGAGTCGT
CCCCTCCGCTGGCAGCATGACCGCCCTGGGTGGAGCTCTCTGTCGA
CCCACACCCCATCACCACTACCCGCCCTACGTGCCCGAGTACAGCT
```

-continued

CCGGACTCTTCCCCCAGCAGCCTGCTGGGCGGCTCCCCACCGGCT
 TCGGATGCAAGTCCAGGCCAAGGCCGGTCCAGCACAGAAGGCAGGG
 AGTGTGTGAACGTGGGGCAACCTCGACCCACTGTGGCGGCGAGATG
 GCACGGGACACTACCTGTGCAACGCCTGCGGGCTCTATCACAAAATGA
 ACGGACAGAACCGGCCCTCATTAAAGCCCAAGCGAAGGCTGTCTGCAG
 CCAGGAGAGCAGGGACGTCCTGTGCGAACTGTCAGACCAACACAACCA
 CACTCTGGAGGAGGAATGCCAATGGGGACCTGTCTGCAATGCCTGTG
 GGCTCTACTACAAGCTTCACATATTAACAGACCCCTGACTATGAAGA
 AGGAAGGCATCCAGACCAGAAACCGAAAAATGTCTAGCAATCCAAAA
 AGTGCAAAAAAGTGCATGACTCACTGGAGGACTTCCCAAGAACAGCT
 CGTTTAACCCGGCCGCTCTCCAGACACATGTCTCCCTGAGCCACA
 TCTCGCCCTTCAGCCACTCCAGCCACATGCTGACCACGCCACGCCGA
 TGCACCCGCCATCCAGCCTGTCTTTGGACCACACCACCCCTCCAGCA
 TGGTCAACCCATGGGTAGAGCCCTGCTCGATGCTCAGAGGCCCCC
 AGCGAGAGTCCCTGCAGTCCCTTCGACTTGCAATTTTGCAGGAGCAG
 TATCATGAAGCCTAAACGCGATGGATATATGTTTTGAAGGCAGAAAG
 CAAAATTATGTTTGCCACTTTGCAAGGAGCTCACTGTGGTGTCTGTG
 TTCCAACCACTGAATCTGGACCCCTCTGTGAATAAGCCATTCTGACT
 CATATCCCTATTTAACAGGGTCTCTAGTGTGTGAAAAAAAATGC
 TGAACATTGCATATAACTTATATTGTAAGAAATACTGTACAATGACTT
 TATTGCATCTGGTAGCTGTAAGGCATGAAGGATGCCAAGAGTTTAA
 GGAATATGGGAGAAATAGTGTGGAATTAAGAAGAACTAGTCTGAT
 ATTCAAATGGACAACTGCCAGTTTTGTTTCTTTCACTGCCCACAGT
 TGTTTGATGCATTAAAGAAAAATAAAAAAGAAAAAGAGAAAAAGAA
 AAAAAAGAAAAAGTTGTAGGCGAATCATTTGTTCAAAGCTGTTGGC
 CTCTGCAAAGGAAATACCACTTCTGGGCAATCAGTGTACCGTTCCACC
 AGTTGCCGTTGAGGGTTTCAGAGAGCCTTTTCTAGGCCTACATGCTT
 TGTGAACAAGTCCCTGTAATTGTTGTTGTATGTATAATTCAAAGCAC
 CAAAATAAGAAAGATGTAGATTTATTTTCATCATATTATACAGACCGA
 ACTGTTGTATAATTTATTACTGCTAGTCTTAAGAACTGCTTTCTTTC
 GTTTGTTGTTTCAATATTTTCTTCTCTCAATTTTGGTTGAATA
 AACTAGATTACATTCACTTGGCCTAAGGTGGTGTGCTCGGAGGGTTT
 CTGTTTCTTTTCCATTTTGTGTTTGGATGATATTTATTAATAGCTT
 CTAAGAGTCCGGCGCATCTGTCTTGTCCCTATTCTGCAGCCTGTGC
 TGAGGGTAGCAGTGTATGAGCTACCAGCGTGCATGTACGCGACCCCTGG
 CCCGACAGGCCACGTCTGCAATCGGCCCGGCTGCCTCTTCGCCCTGT
 CGTGTCTGTGTTAGTGATCACTGCCTTTAATACAGTCTGTTGGAATA
 ATATTATAAGCATAATAATAAGTGAAATATTTTAAACTACAA.
 The GATA-3 protein is encoded by nucleotide
 558-1892.

[0055] In certain embodiments, methods and compositions disclosed herein may be implemented with nucleobase polymers comprising sequences that hybridize or are the reverse complement to GATA-3 mRNA and contain an RNA cleaving nucleobase sequence such as a cleaving DNAzyme sequence. DNAzymes are catalytically active DNA molecules. DNAzyme 10-23 is DNA with two binding domains flanking a central catalytic domain. After binding of a DNAzyme to the corresponding sequence in the target mRNA via the binding domains, the catalytic domain cleaves the target mRNA molecule. The binding domains and catalytic domain may contain modifications provided they are capable of specifically binding the corresponding target mRNA of GATA-3. Schubert et al. report DNAzymes against the same target site that are stabilized by the use of a 3'-3'-inverted thymidine, phosphorothioate linkages, 2'-O-methyl RNA and locked nucleic acids. *Nucleic Acids Res.* 2003, 31 (20): 5982-5992. In certain embodiments, methods and compositions disclosed herein may be implemented with a nucleobase polymers comprising units of a ribose, 2'-deoxyribose, locked nucleic acids (1-(hydroxymethyl)-2, 5-dioxabicyclo[2.2.1]heptan-7-ol), 2'-O-methyl groups, a 3'-3'-inverted thymidine, phosphorothioate linkages, or combinations thereof.

[0056] The term "nucleobase polymer" refers to a molecule having nucleobase monomers capable of hybridizing to a single-stranded nucleic acid target. The nucleobase polymers that target GATA-3 mRNA typically comprise a sequence that is the reverse complement of, more than 7, 8, 9, 10, 11, 12, 13, 14, 15, 20, or more nucleotides or nucleobases or continuous nucleotide nucleobases of SEQ ID NO: 52. The targeting sequence of nucleobases is typically separated by an RNA cleaving sequence such as a DNAzyme sequence capable of forming a loop-like structure. The nucleobase polymer may be a single stranded nucleic acid or analog containing a sufficiently small number of target mismatches, additions, or deletions as long as the targeting sequences retain the ability to bind to the target RNA.

[0057] In certain embodiments, the nucleobase polymer including the targeting sequence and the RNA cleaving DNAzyme may be less than 500, 200, 100, 50, or 35 nucleotides or nucleobases. In certain embodiments, the disclosure contemplates the use of nucleobase polymers disclosed herein for disruption of GATA-3 expression.

[0058] Nucleobase monomers are typically nitrogen containing aromatic or heterocyclic bases that bind to naturally occurring nucleic acids through hydrogen bonding otherwise known as base pairing. A typical nucleobase polymer is a nucleic acid, RNA, DNA, or chemically modified form thereof. A nucleobase polymer may be single or double stranded or both, e.g., they may contain overhangs. Nucleobase polymers may contain naturally occurring or synthetically modified bases and backbones. In certain embodiments, a nucleobase polymer need not be entirely complementary, e.g., may contain one or more insertions, deletions, or be in a hairpin structure provided that there is sufficient selective binding.

[0059] With regard to the nucleobases, it is contemplated that the term encompasses isobases, otherwise known as modified bases, e.g., are isoelectronic or have other substitutes configured to mimic naturally occurring hydrogen bonding base-pairs, e.g., within any of the sequences herein U may be substituted for T, or T may be substituted for U.

Examples of nucleotides with modified adenosine or guanosine include, but are not limited to, hypoxanthine, xanthine, 7-methylguanine.

[0060] Examples of nucleotides with modified cytidine, thymidine, or uridine include 5,6-dihydrouracil, 5-methylcytosine, 5-hydroxymethylcytosine. Contemplated isobases include 2'-deoxy-5-methylisocytidine (iC) and 2'-deoxy-isoguanosine (iG) (see U.S. Pat. Nos. 6,001,983; 6,037,120; 6,617,106; 6,977,161). In another embodiment, a removable base (such as uracil or 8-oxoguanine) is contemplated so that treatment by uracil-DNA glycosylase (UDG) or formamidopyrimidine-DNA glycosylase (FPG), can lead to cleavage and degradation of unwanted sequences.

[0061] In order to prevent in vivo breakdown nucleic acids may be chemically modified, e.g., within the sugar backbone or on the 5' or 3' ends. As such, in certain embodiments, nucleobase polymers disclosed herein may contain monomers of phosphodiester, phosphorothioate, methylphosphonate, phosphorodiamidate, piperazine phosphorodiamidate, ribose, 2'-O-methyl ribose, 2'-O-methoxyethyl ribose, 2'-fluororibose, deoxyribose, 1-(hydroxymethyl)-2,5-dioxabicyclo[2.2.1]heptan-7-ol, P-(2-(hydroxymethyl) morpholino)-N,N-dimethylphosphonamide, morpholin-2-ylmethanol, (2-(hydroxymethyl) morpholino)(piperazin-1-yl) phosphinate, or peptide nucleic acids or combinations thereof.

[0062] Within any of the sequences disclosed herein, U may be T or T may be U.

[0063] In certain embodiments, the nucleotide base polymer is single or double stranded DNA that is 3' end capped with one, two, or more thymidine nucleotides and/or a 5' end polyphosphorylated, e.g., di-phosphate, tri-phosphate.

[0064] In certain embodiments, the nucleobase polymer can be modified to contain a 3' end thiol group for direct absorption on gold or silver surfaces and particles. In certain embodiments, the nucleobase polymer is conjugated to a poly T sequence. In certain embodiments, the poly T sequence is on the 3' end of the nucleobase polymer. In certain embodiments, the poly T sequence has four or more repeating thymine (T) bases followed by a 3' end thiol group. In certain embodiments, the nucleobase polymer can be modified to contain a phosphodiester bond. Replacing one of the non-bridging oxygen by sulfur imparts resistance to nuclease degradation.

[0065] In certain embodiments, the nucleobase polymer can be modified to contain a C-5 propyne substitution of dC and/or dT in the target sequences. In certain embodiments, the nucleobase polymer can be modified to contain a 2'-5' linkages and 2'-5' linked ends synthesized using 3'-deoxy-2'-phosphoramidites and 2'-deoxy-3'-phosphoramidites. In certain embodiments, the nucleobase polymer can be modified to contain a terminal 5'-5' or 3'-3' linkage. For the terminal 5'-5' linkage, the appropriate 5'-phosphoramidite is incorporated at the 5'-end in a synthesis cycle. For the terminal 3'-3' linkage, the appropriate deoxynucleoside-5'-CPG is used as the solid support for the 3'-end, followed by synthesis of the oligo in the standard 3'-5' direction to make the terminal 3'-3' linkage.

[0066] In certain embodiments, the disclosure relates to compounds, compositions, and methods disclosed herein using nucleobase polymers. In certain embodiment, nucleobase polymers may contain any of the sequences provided in the table 1 of the experimental section (SEQ ID NO: 1-49)

[0067] In certain embodiments, the disclosure contemplates that DNAzyme 10-23, SEQ ID NO: 51 (GGCTAGC-TACAACGA) can be substituted with other DNAzymes sequences. The DNAzyme 10-23 is comprised of a sequence of DNA that will cleave mRNA strands that contain an unpaired purine-pyrimidine pair. The DNAzyme 10-23 is flanked by recognition sequences that hybridized to the regions of the target mRNA sequences adjacent to the unpaired purine-pyrimidine pair. Therefore, the DNAzyme hybridizes and cleaves at the purine-pyrimidine site.

[0068] In certain embodiments, this disclosure contemplates that the cleaving nucleic acids comprise sequences of DNAzymes 8-17 and 10-23. Santoro & Joyce disclosed a general purpose RNA-cleaving DNAzymes 8-17 and 10-23. See PNSA, 1997, 94 (9), 4262-4266.

[0069] In certain embodiments, this disclosure contemplates that the cleaving nucleic acids comprise amine, guanidine, imidazole modifications such as 8-imidazolyl modified deoxy adenosines. Perrin et al., disclose modified DNAzymes 20-49 containing amine, guanidine, and imidazole-modified dNTPs. Org Biomol Chem 2011, 9 (7), 2266-2273.

Synthesis of Nucleobases Polymers

[0070] Small nucleobase polymers and nucleic acid motifs ("small" refers to nucleic acid motifs no more than 100 nucleotides in length, preferably no more than 80 nucleotides in length, and most preferably no more than 50 nucleotides in length; e.g., individual oligonucleotide sequences or sequences synthesized in tandem) are preferably used for exogenous delivery. Exemplary molecules of the instant disclosure may be chemically synthesized or by the use of cellular or non-cellular expression systems. Nucleic acids made by cellular or non-cellular expression systems can be further modified.

[0071] One synthesizes oligonucleotides (e.g., certain modified oligonucleotides or portions of oligonucleotides) using protocols known in the art as, for example, described in U.S. Pat. No. 6,001,311. The synthesis of oligonucleotides makes use of common nucleic acid protecting and coupling groups, such as dimethoxytrityl at the 5'-end and phosphoramidites at the 3'-end.

[0072] In a non-limiting example, small scale syntheses are conducted on a 394 Applied Biosystems, Inc. synthesizer using a 0.2 micro mol scale protocol with a 2.5 min coupling step for 2'-O-methylated nucleotides and a 45 second coupling step for 2'-deoxy nucleotides or 2'-deoxy-2'-fluoro nucleotides. Alternatively, syntheses at the 0.2 micro mol scale can be performed on a 96-well plate synthesizer, such as the instrument produced by Protogene (Palo Alto, Calif.) with minimal modification to the cycle. A 33-fold excess of 2'-O-methyl phosphoramidite and a 105-fold excess of S-ethyl tetrazole can be used in each coupling cycle of 2'-O-methyl residues relative to polymer-bound 5'-hydroxyl. A 22-fold excess of deoxy phosphoramidite and a 70-fold excess of S-ethyl tetrazole mop can be used in each coupling cycle of deoxy residues relative to polymer-bound 5'-hydroxyl. Other oligonucleotide synthesis reagents for the 394 Applied Biosystems, Inc. synthesizer include the following: detritylation solution is 3% TCA in methylene chloride (ABI); capping is performed with 16% N-methyl imidazole in THF (ABI) and 10% acetic anhydride/10% 2,6-lutidine in THF (ABI); and oxidation solution is 16.9 mM 12, 49 mM pyridine, 9% water in THF (PerSeptive Biosystems, Inc.).

S-Ethyltetrazole solution (0.25 M in acetonitrile) is made up from the solid obtained from American International Chemical, Inc. Alternately, for the introduction of phosphorothioate linkages, Beaucage reagent (3H-1,2-benzodithiol-3-one 1,1-dioxide, 0.05 M in acetonitrile) is used.

[0073] Deprotection of the DNA-based oligonucleotides is performed as follows: the polymer-bound trityl-on oligonucleotide is transferred to a 4 mL glass screw top vial and suspended in a solution of 40% aqueous methylamine (1 mL) at 65 degrees for 10 minutes. After cooling to -20 degrees, the supernatant is removed from the polymer support. The support is washed three times with 1.0 mL of EtOH: MeCN: H₂O/3:1:1, vortexed and the supernatant is then added to the first supernatant. The combined supernatants, containing the oligonucleotide, are dried.

[0074] Alternately, the nucleic acid molecules can be synthesized separately and joined together post-synthetically, for example, by ligation or by hybridization following synthesis and/or deprotection.

[0075] Nucleic acids can also be assembled from two distinct nucleic acid strands or fragments wherein one fragment includes the sense region and the second fragment includes the antisense region of the RNA molecule.

[0076] The nucleic acid molecules can be modified extensively to enhance stability by modification with nuclease resistant groups, for example, 2'-amino, 2'-C-allyl, 2'-fluoro, 2'-O-methyl, 2'-H). Constructs can be purified by gel electrophoresis using general methods or can be purified by high pressure liquid chromatography and re-suspended in water.

[0077] Chemically synthesizing nucleic acid molecules with modifications (base, sugar and/or phosphate) can prevent their degradation by serum ribonucleases, which can increase their potency. See e.g., U.S. Pat. Nos. 5,652,094, 5,334,711, and 6,300,074. All of the above references describe various chemical modifications that can be made to the base, phosphate and/or sugar moieties of the nucleic acid molecules described herein. Modifications that enhance their efficacy in cells, and removal of bases from nucleic acid molecules to shorten oligonucleotide synthesis times and reduce chemical requirements are desired.

[0078] In one embodiment, nucleic acid molecules include one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) G-clamp nucleotides. A G-clamp is a tricyclic aminoethylphenoxazine 2'-deoxycytidine or analogue. See Lin & Matteucci, J Am Chem Soc, 1998, 120, 8531-8532; Flanagan, et al., Proc Nat Acad Sci USA, 1999, 96, 3513-3518; and Maier, et al., Biochemistry, 2002, 41, 1323-1327. A single G-clamp analog substitution within an oligonucleotide can result in substantially enhanced helical thermal stability and mismatch discrimination when hybridized to complementary oligonucleotides. The inclusion of such nucleotides in nucleic acid molecules results in both enhanced affinity and specificity to nucleic acid targets, complementary sequences, or template strands.

[0079] In another embodiment, nucleic acid molecules include one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) LNA "locked nucleic acid" nucleotides such as a 2',4'-C methylene bicyclo nucleotide (see for example U.S. Pat. Nos. 6,639,059, 6,670,461, 7,053,207).

[0080] In another embodiment, the disclosure features conjugates and/or complexes of nucleobase polymers. Such conjugates and/or complexes can be used to facilitate delivery of polymers into a biological system, such as a cell. Contemplated conjugates include those with cell penetrating

peptide. The conjugates and complexes provided may impart therapeutic activity by transferring therapeutic compounds across cellular membranes, altering the pharmacokinetics, and/or modulating the localization of nucleic acid molecules. In general, the transporters described are designed to be used either individually or as part of a multi-component system, with or without degradable linkers. These compounds improve delivery and/or localization of nucleic acid molecules into a number of cell types originating from different tissues, in the presence or absence of serum (see U.S. Pat. No. 5,854,038). Conjugates of the molecules described herein can be attached to biologically active molecules via linkers that are biodegradable, such as biodegradable nucleic acid linker molecules.

[0081] In another aspect, a nucleobase polymer comprises one or more 5' and/or 3'-cap structures. A "cap structure" refers to chemical modifications, which have been incorporated at either terminus of the oligonucleotide. See, for example, Adamic et al., U.S. Pat. No. 5,998,203. These terminal modifications protect the nucleic acid molecule from exonuclease degradation and may help in delivery and/or localization within a cell. The cap may be present at the 5'-terminus (5'-cap) or at the 3'-terminal (3'-cap) or may be present on both termini. In non-limiting examples, the 5'-cap includes, but is not limited to, glyceryl, inverted deoxy abasic residue (moiety); 4',5'-methylene nucleotide; 1-(beta-D-erythrofuransyl) nucleotide, 4'-thio nucleotide; carbocyclic nucleotide; 1,5-anhydrohexitol nucleotide; L-nucleotides; alpha-nucleotides; modified base nucleotide; phosphorodithioate linkage; threo-pentofuransyl nucleotide; acyclic 3',4'-seco nucleotide; acyclic 3,4-dihydroxybutyl nucleotide; acyclic 3,5-dihydroxypentyl nucleotide, 3'-3'-inverted nucleotide moiety; 3'-3'-inverted abasic moiety; 3'-2'-inverted nucleotide moiety; 3'-2'-inverted abasic moiety; 1,4-butanediol phosphate; 3'-phosphoramidate; hexylphosphate; aminohexyl phosphate; 3'-phosphate; 3'-phosphorothioate; phosphorodithioate; or bridging or non-bridging methylphosphonate moiety.

[0082] Non-limiting examples of the 3'-cap include, but are not limited to, glyceryl, polyethylene glycol, alkyl, inverted deoxy abasic residue (moiety), 4',5'-methylene nucleotide; 1-(beta-D-erythrofuransyl) nucleotide; 4'-thio nucleotide, carbocyclic nucleotide; 5'-amino-alkyl phosphate; 1,3-diamino-2-propyl phosphate; 3-aminopropyl phosphate; 6-aminohexyl phosphate; 1,2-aminododecyl phosphate; hydroxypropyl phosphate; 1,5-anhydrohexitol nucleotide; L-nucleotide; alpha-nucleotide; modified base nucleotide; phosphorodithioate; threo-pentofuransyl nucleotide; acyclic 3',4'-seco nucleotide; 3,4-dihydroxybutyl nucleotide; 3,5-dihydroxypentyl nucleotide, 5'-5'-inverted nucleotide moiety; 5'-5'-inverted abasic moiety; 5'-phosphoramidate; 5'-phosphorothioate; 1,4-butanediol phosphate; 5'-amino; bridging and/or non-bridging 5'-phosphoramidate, phosphorothioate and/or phosphorodithioate, bridging or non-bridging methylphosphonate and 5'-mercapto moieties (for more details see Beaucage and Iyer, 1993, Tetrahedron 49, 1925).

[0083] In one embodiment, the disclosure features modified nucleobase polymer, with phosphate backbone modifications comprising one or more phosphorothioate, phosphorodithioate, methylphosphonate, phosphotriester, morpholino, amidate carbamate, carboxymethyl, acetamidate, polyamide, sulfonate, sulfonamide, sulfamate, formacetal, thioformacetal, and/or alkylsilyl, substitutions.

Pharmaceutical Compositions

[0084] In certain embodiments, the disclosure contemplates pharmaceutical composition comprising a nucleobase polymer disclosed herein and a pharmaceutically acceptable excipient.

[0085] In certain embodiments, the pharmaceutical composition is in the form of a sterilized pH buffered aqueous salt solution or an isotonic aqueous buffer solution.

[0086] In certain embodiments, the pharmaceutically acceptable excipient is a propellant or aerosolizing agent. In certain embodiments, the propellant or aerosolizing agent is a hydrofluoroalkane, 1,1,1,2-tetrafluoroethane, 1,1,1,2,3,3,3-heptafluoropropane, propane, n-butane, isobutene, carbon dioxide, compressed air, nitrogen, nitrous oxide, dimethyl ether, trans-1,3,3,3-tetrafluoroprop-1-ene, or combinations thereof.

[0087] In certain embodiments, the pharmaceutically acceptable excipient is a lipid, fatty acid, phospholipid, diacyl phospholipid, phosphatidyl choline, sorbitan monopalmitate, polyethoxylated sorbitan monopalmitate, sucrose stearate (mono- and di-ester), alginate, copolymer of poly(lactide-co-glycolide)(PLGA), poly(vinyl alcohol), or poly(L-lysine)(PLL).

[0088] In certain embodiments, the pharmaceutical compositions may be stored in a nebulizer, inhaler, or other container optionally sealed or under a pressure for propelling the pharmaceutical agent(s). The container may contain a spraying apparatus that is manually-actuated or pressurized. Metered dose inhalers (MDIs) typically have a hand-held aerosol canister that, upon being pushed, releases an amount of medicine to inhale. Dry powder inhalers (DPIs) do not use a propellant to release the medicine. Instead, a dry powder form of the peptide or agent is drawn into your lungs after a breath. In certain configurations, a container comprising the peptide or agent is inserted a device. Pressing a button or section on the device pierces the container. One can breathe in the powder contained in the container through a mouthpiece on the device.

[0089] In certain embodiments, the pharmaceutical compositions disclosed herein further comprise a respiratory agent selected from a glucocorticoid receptor agonist (steroidal and non-steroidal) such as triamcinolone, triamcinolone acetonide, prednisone, mometasone furoate, loteprednol etabonate, fluticasone propionate, fluticasone furoate, flucinolone acetonide, dexamethasone cipeclate, desisobutyl ciclesonide, clobetasol propionate, ciclesonide, butixocort propionate, budesonide, beclomethasone dipropionate, alclometasone dipropionate; a p38 antagonist such as losmapimod; a phosphodiesterase (PDE) inhibitor such as a methylxanthanine, theophylline, and aminophylline; a selective PDE isoenzyme inhibitor, a PDE4 inhibitor and the isoform PDE4D, such as tetomilast, roflumilast, oglemilast, ibudilast, ronmilast; a modulator of chemokine receptor function such as vicriviroc, maraviroc, cenicriviroc, navarixin; a leukotriene biosynthesis inhibitor, 5-lipoxygenase (5-LO) inhibitor, and 5-lipoxygenase activating protein (FLAP) antagonist such as TA270 (4-hydroxy-1-methyl-3-octyloxy-7-sinapinoylamino-2 (1H)-quinolinone) such as setileuton, licofelone, quiflapon, zileuton, zafirlukast, or montelukast; and a myeloperoxidase antagonist such as resveratrol and piceatannol.

[0090] Effective doses may be extrapolated from dose-response curves derived from in vitro or animal model test systems. For nucleobase polymers or particles disclosed

herein, the dosage administered to a patient is typically 0.0001 mg/kg to 100 mg/kg of the patient's body weight. Preferably, the dosage administered to a patient is between 0.0001 mg/kg and 20 mg/kg, 0.0001 mg/kg and 10 mg/kg, 0.0001 mg/kg and 5 mg/kg, 0.0001 and 2 mg/kg, 0.0001 and 1 mg/kg, 0.0001 mg/kg and 0.75 mg/kg, 0.0001 mg/kg and 0.5 mg/kg, 0.0001 mg/kg to 0.25 mg/kg, 0.0001 to 0.15 mg/kg, 0.0001 to 0.10 mg/kg, 0.001 to 0.5 mg/kg, 0.01 to 0.25 mg/kg or 0.01 to 0.10 mg/kg of the patient's body weight. Further, the dosage and frequency of administration of nucleobase polymers or particles disclosed herein may be reduced by enhancing uptake and tissue penetration of the nucleobase polymers or particles disclosed herein by modifications such as, for example, lipidation.

[0091] In a specific embodiment, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term "carrier" refers to a diluent, adjuvant (e.g., Freund's adjuvant (complete and incomplete), excipient, or vehicle with which the therapeutic is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like.

[0092] Generally, the ingredients of compositions are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

[0093] The compositions can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include, but are not limited to, those formed with anions such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

[0094] One embodiment provides a pharmaceutical pack or kit comprising one or more containers filled with nucleobase polymers or particles disclosed herein. Additionally, one or more other prophylactic or therapeutic agents useful for the treatment of a disease can also be included in the pharmaceutical pack or kit. One embodiment provides a

pharmaceutical pack or kit including one or more containers filled with one or more of the ingredients of the pharmaceutical compositions.

[0095] Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

[0096] In certain embodiment, this disclosure contemplates pharmaceutical compositions comprising nucleobase polymers or particles disclosed herein and pharmaceutically acceptable excipient. In certain embodiments, this disclosure contemplates the production of a medicament comprising nucleobase polymers or particles disclosed herein and uses for methods disclosed herein.

[0097] In certain embodiments, the disclosure relates to pharmaceutical compositions comprising nucleobase polymers or particles disclosed herein and a pharmaceutically acceptable excipient. In certain embodiments, the composition is a pill or in a capsule or the composition is an aqueous buffer, e.g., a pH between 6 and 8. In certain embodiments, the pharmaceutically acceptable excipient is selected from a filler, glidant, binder, disintegrant, lubricant, and saccharide.

[0098] Compositions suitable for parenteral injection may comprise physiologically acceptable sterile aqueous or non-aqueous solutions, dispersions, suspensions or emulsions, and sterile powders for reconstitution into sterile injectable solutions or dispersions. Examples of suitable aqueous and nonaqueous carriers, diluents solvents or vehicles include water, ethanol, polyols (propylene glycol, polyethylene glycol, glycerol, and the like), suitable mixtures thereof, vegetable (such as olive oil, sesame oil) and injectable organic esters such as ethyl oleate.

[0099] Prevention of the action of microorganisms may be controlled by addition of any of various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, and the like. It may also be desirable to include isotonic agents, for example sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form can be brought about by the use of agents delaying absorption, for example, aluminum monostearate and gelatin.

[0100] Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs. In addition to the nucleobase polymers or particles disclosed herein, the liquid dosage forms may contain inert diluents commonly used in the art, such as water or other solvents, solubilizing agents and emulsifiers, for example, ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils, in particular, cottonseed oil, groundnut oil, corn germ oil, olive oil, castor oil and sesame oil, glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan or mixtures of these substances, and the like.

[0101] In certain embodiments, production processes are contemplated which two components, nucleobase polymers or particles disclosed herein and a pharmaceutical carrier, are provided already in a combined dry form ready to be reconstituted together. In other embodiments, it is contemplated

that nucleobase polymers or particles disclosed herein and a pharmaceutical carrier are admixed to provide a pharmaceutical composition.

[0102] Providing a pharmaceutical composition is possible in a one-step process, simply by adding a suitable pharmaceutically acceptable diluent to the composition in a container. In certain embodiments, the container is preferably a syringe for administering the reconstituted pharmaceutical composition after contact with the diluent. In certain embodiments, the nucleobase polymers or particles disclosed herein can be filled into a syringe, and the syringe can then be closed with the stopper. A diluent is used in an amount to achieve the desired end-concentration. The pharmaceutical composition may contain other useful component, such as ions, buffers, excipients, stabilizers, etc.

[0103] A “dry” pharmaceutical composition typically has only a residual content of moisture, which may approximately correspond to the moisture content of comparable commercial products, for example, has about 12% moisture as a dry product. Usually, the dry pharmaceutical composition according to the present invention has a residual moisture content preferably below 10% moisture, more preferred below 5% moisture, especially below 1% moisture. The pharmaceutical composition can also have lower moisture content, e.g. 0.1% or even below. In certain embodiments, the pharmaceutical composition is provided in dry in order to prevent degradation and enable storage stability.

[0104] A container can be any container suitable for housing (and storing) pharmaceutically compositions such as inhalers, syringes, vials, tubes, etc. The pharmaceutical composition may then be applied via actuation or specific needles of the syringe or via suitable catheters. A typical diluent comprises water for injection, and NaCl (preferably 50 to 150 mM, especially 110 mM), CaCl_2 (preferably 10 to 80 mM, especially 40 mM), sodium acetate (preferably 0 to 50 mM, especially 20 mM) and mannitol (preferably up to 10% w/w, especially 2% w/w). Preferably, the diluent can also include a buffer or buffer system so as to buffer the pH of the reconstituted dry composition, preferably at a pH of 6.2 to 7.5, especially at pH of 6.9 to 7.1.

[0105] In certain embodiments, this disclosure contemplates a kit comprising a pharmaceutical composition disclosed herein such as a peptide or agent and a container optionally with a suitable diluent. Further components of the kit may be instructions for use, administration means, such as inhalers, syringes, catheters, brushes, etc. (if the compositions are not already provided in the administration means) or other components necessary for use in medical (surgical) practice, such as substitute needles or catheters, extra vials or further wound cover means. In certain embodiments, the kit comprises a syringe housing the dry and stable hemostatic composition and a syringe containing the diluent (or provided to take up the diluent from another diluent container).

[0106] In certain embodiments, the diluent is provided in a separate container. This can preferably be a syringe. The diluent in the syringe can then easily be applied to the container for reconstitution of the dry compositions. If the container is also a syringe, both syringes can be finished together in a pack. It is therefore preferred to provide the dry compositions in a syringe, which is finished with a diluent syringe with a pharmaceutically acceptable diluent for reconstituting, said dry and stable composition.

Methods of Use

[0107] Chronic inflammations constitute an increasing medical problem area of high socioeconomic significance. In certain embodiments, the disclosure relates to methods of treating an inflammatory disease or chronic inflammation comprising administering an effective amount of a nucleobase polymer disclosed herein to a subject in need thereof. In certain embodiments, the inflammatory disease is a respiratory disorder. In certain embodiments, the subject is diagnosed with, exhibiting symptoms of or at risk of asthma, bronchitis, emphysema, chronic obstructive pulmonary disease, laryngitis, or cystic fibrosis.

[0108] In certain embodiments, the inflammatory disease is ulcerative colitis.

[0109] In certain embodiments, the chronic inflammation is due to autoimmune diseases and diseases from the area of rheumatic diseases (manifestations among others on the skin, lungs, kidneys, vascular system, nervous system, connective tissue, locomotor system, endocrine system), immediate-type allergic reactions and asthma, chronic obstructive lung diseases (COPD), arteriosclerosis, psoriasis and contact eczema and chronic rejection reactions after organ and bone marrow transplants.

Examples

[0110] Dz-conjugated to nanoparticles readily enter cells in vitro and in vivo To test the activity of GATA-3 DzNPs, 3'thiol modified GATA-3 Dzs were synthesized. Approximately 100 copies of the Dzs were functionalized onto a gold nanoparticle. PAGE demonstrated that these DzNPs were highly active. To test whether DzNPs could be delivered to the lung, the Dzs were tagged with Cy5 dye to generate fluorescent DzNPs. These particles were administered by nebulization into 4-6 micrometer sized aerosol droplets using a bias flow commercial nebulization system. Fluorescence imaging of the mouse lung showed even and distal distribution 24 hrs after nebulization (FIG. 1).

[0111] To confirm resident lung cells can internalize DzNP conjugates, uptake in cultured lung cell lines was tested. The lung cell lines tested were macrophages, smooth muscle cells, as well as epithelial cells. DzNP particles were prepared that were modified with the Cy5 dye, and these particles were incubated at 10 nM concentration for 36 hrs. The cells were cultured in standard serum-containing media. Cells were washed and analyzed using flow cytometry (FIG. 2). The results clearly show a significant level of cellular uptake, with epithelial cells showing the highest level of uptake. Fluorescence microscopy confirmed that the particles were internalized.

Dznps for GATA-3 Gene Regulation are Active In Vivo

[0112] A model of mouse asthma was used to test the efficacy of the DzNPs and showed a significant improve-

ment in airway function for the treated animals as well as reduced cytokine levels in the blood plasma (FIGS. 3A and B). In this model, animals are treated with house dust mite ion a daily basis. The house dust mite antigen exposure model mimics the Th2 inflammation and airway hyper-responsiveness seen in asthma. Der pl house dust mite extract (50 ug) was administered for 2 weeks. DzNP treatment (100 nM, 20 uL) was delivered intranasally for 5 days (the last week of the experiment). The results of the animal experiments demonstrate that the control group did not show any negative effects due to the nanoparticle treatment. The group receiving the active DzNP showed airway resistance that is similar to that of the untreated animal group. Importantly, the oligonucleotide dose with the DzNP construct was one order of magnitude smaller than that used for soluble Dz experiments. These results are promising and suggest that DzNP based targeting of GATA-3 in the lung offers potential for treating asthma. This experiment was conducted using the HGD40 sequence (site 917 on the human mRNA transcript for GATA-3, SEQ ID NO: 50, (TGGATG-GAGGCTAGCTACAACGAGTCTTGGAG) that was previously identified and also tested in phase II human trials. DNzyme sequences were screened to identify ones that could be more active than the HGD40 sequence. In vivo efficacy of DzNP were tested for improvements.

In Silico Screen to Discover Active DNzyme

[0113] A computational algorithm was created to predict the most efficient DNzyme sequences to destroy any given mRNA sequence. The algorithm first scans the entire gene transcript identifying regions of minimal secondary structure by leveraging the mFold software package. NCBI Reference Sequence: NM_001002295.1 for the human mRNA and NCBI Reference Sequence: NM_008091.3 was used for the mouse gene. Next, purine-pyrimidine junctions were identified where the 10-23 DNzyme is most active. Subsequently, the left and right arms of the DNzyme are tuned such that the delta G of hybridization is between -8 and -10 kcal/mol. For example, G-C rich regions (~5-7 mer) will tend to have shorter arms, while A-T rich regions require longer arms (~10-12 mer). This screen generates a list of potential DNzyme targets. The list can then be further filtered based on comparing the human and murine genomes and also various splice variants of a given transcript.

[0114] A screen was performed on the potential DNzyme sites. The top ranked sequences were determined using the algorithm, and these were custom synthesized (see table below). The first column is the location of purine-uracil site that will be cleaved in the human transcript. All of these sequences are shared between mouse and human (albeit at different locations). The sequences in the far right are the DNzyme sequences used to generate the DzNP conjugates

TABLE 1

DNase hits that target GATA-3 mRNA				
Human GATA-3 cleavage site (b)	3'b	5'b	Sequence	
1461	7	9	5-TGGGCTTAAGGCTAGCTACAACGAGAGGGGC-3	
SEQ ID NO: 1				

TABLE 1-continued

DNAzyme hits that target GATA--3 mRNA			
Human GATA-3 cleavage site (b)	3'b	5'b	Sequence
1519	8	8	5-TGGTCTGAGGCTAGCTACAACGAAGTTCGCA-3 SEQ ID NO: 2
1606	8	11	5-GTCTGTTAATAGGCTAGCTACAACGATGTGAAGC-3 SEQ ID NO: 3
1662	8	9	5-TGCTAGACAGGCTAGCTACAACGATTTTCGGT-3 SEQ ID NO: 4
1664	9	9	5-TTTGCTAGAGGCTAGCTACAACGAATTTTCGG-3 SEQ ID NO: 5
1673	9	9	5-CTTTTGGAGGCTAGCTACAACGATTGCTAGAC-3 SEQ ID NO: 6
1682	10	9	5-TTTTTGCAGGCTAGCTACAACGATTTTGGATT-3 SEQ ID NO: 7
1970	8	9	5-GCTTCATGAGGCTAGCTACAACGAAGTCTCC-3 SEQ ID NO: 8
1973	9	8	5-AGGCTTCAGGCTAGCTACAACGAGATACTGCT-3 SEQ ID NO: 9
2093	8	10	5-GGCTTATTCAGGCTAGCTACAACGAAGATGGGG-3 SEQ ID NO: 10
2097	9	9	5-AATGGCTTAGGCTAGCTACAACGATCACAGATG-3 SEQ ID NO: 11
2114	9	9	5-TAGGGGATAGGCTAGCTACAACGAGAGTCAGAA-3 SEQ ID NO: 12
2116	9	9	5-AATAGGGGAGGCTAGCTACAACGAATGAGTCAG-3 SEQ ID NO: 13
2133	10	9	5-CACTAGAGAGGCTAGCTACAACGACCTGTTAAAT-3 SEQ ID NO: 14
2140	9	8	5-TCACAGCAGGCTAGCTACAACGATAGAGACCC-3 SEQ ID NO: 15
2145	9	10	5-TTTTTTTTCAGGCTAGCTACAACGAAGCACTAGA-3 SEQ ID NO: 16
2157	11	8	5-GTTCAGCAGGCTAGCTACAACGATTTTTTTTCA-3 SEQ ID NO: 17
2171	9	12	5-ATATAAGTTATAGGCTAGCTACAACGAGCAATGTTC-3 SEQ ID NO: 18
2173	9	12	5-CAATATAAGTTAGGCTAGCTACAACGAATGCAATGT-3 SEQ ID NO: 19
2251	8	10	5-ATTCCTTAAAGGCTAGCTACAACGATTCTTGGC-3 SEQ ID NO: 20
2326	8	10	5-GGAAACAAAAGGCTAGCTACAACGATGGCAGTT-3 SEQ ID NO: 21
2331	9	9	5-GAAAGGAAAAGGCTAGCTACAACGAAAACTGGC-3 SEQ ID NO: 22
2363	9	11	5-TTTCCTTTAAGGCTAGCTACAACGAGCATCAAAC-3 SEQ ID NO: 23
2560	8	10	5-ACAACAATTAGGCTAGCTACAACGAAGGGACTT-3 SEQ ID NO: 24

TABLE 1-continued

DNAzyme hits that target GATA--3 mRNA			
Human GATA-3 cleavage site (b)	3'b	5'b	Sequence
2563	9	10	5-CAAACAACAAGGCTAGCTACAACGATACAGGGAC-3 SEQ ID NO: 25
2566	10	10	5-ATACAACAAGGCTAGCTACAACGAAATTACAGGG-3 SEQ ID NO: 26
2569	10	10	5-TACATACAAGGCTAGCTACAACGAAACAATTACA-3 SEQ ID NO: 27
2575	9	11	5-TGAATTATACAGGCTAGCTACAACGAACAACAAC-3 SEQ ID NO: 28
2577	10	11	5-TTTGAATTATAGGCTAGCTACAACGAATACAACAA-3 SEQ ID NO: 29
2597	8	10	5-TCTTTTCTTAGGCTAGCTACAACGATTGGTGC-3 SEQ ID NO: 30
2607	11	11	5-ATAAATCTACAGGCTAGCTACAACGACTTTTCTTATT-3 SEQ ID NO: 31
2609	10	11	5-AAATAAATCTAGGCTAGCTACAACGAATCTTTTCTT-3 SEQ ID NO: 32
2622	11	12	5-TGTATAATATGAGGCTAGCTACAACGAGAAATAAATCT-3 SEQ ID NO: 33
2625	11	11	5-TCTGTATAATAGGCTAGCTACAACGAGATGAAATAAA-3 SEQ ID NO: 34
2627	10	10	5-GTCTGTATAAGGCTAGCTACAACGAATGATGAAAT-3 SEQ ID NO: 35
2644	8	11	5-AATTATACAAGGCTAGCTACAACGAAGTTCGGT-3 SEQ ID NO: 36
2647	9	13	5-AAATAAATTTATAGGCTAGCTACAACGAAACAGTTCG-3 SEQ ID NO: 37
2649	9	12	5-TAAATAAATTTAGGCTAGCTACAACGAACAACAGTT-3 SEQ ID NO: 38
2653	10	11	5-CAGTAAATAAGGCTAGCTACAACGATTATACAACA-3 SEQ ID NO: 39
2657	11	9	5-AGCAGTAAAGGCTAGCTACAACGAAATTTATACA-3 SEQ ID NO: 40
2668	9	10	5-GTTCTTAAGAGGCTAGCTACAACGATAGCAGTAA-3 SEQ ID NO: 41
2723	9	10	5-CAACCAAAAAGGCTAGCTACAACGATGAGAGAGA-3 SEQ ID NO: 42
2827	12	10	5-TTAGAAGCTAGGCTAGCTACAACGATTAATAAATATC-3 SEQ ID NO: 43
2910	8	9	5-TGACATGCAGGCTAGCTACAACGAGCTGGTAG-3 SEQ ID NO: 44
2986	8	9	5-TCACTAACAGGCTAGCTACAACGAAGAACACG-3 SEQ ID NO: 45
2988	9	10	5-TGATCACTAAGGCTAGCTACAACGAACAGAACAC-3 SEQ ID NO: 46
3017	10	10	5-ATTATTCCAAGGCTAGCTACAACGAAGACTGTATT-3 SEQ ID NO: 47

TABLE 1-continued

DNAzyme hits that target GATA-3 mRNA			
Human GATA-3 cleavage site (b)	3'b	5'b	Sequence
3048	12	11	5-AAATATTTTCAGGCTAGCTACAACGATTATTATTATG-3 SEQ ID NO: 48
3056	10	11	5-TAGTTTAAAAAGGCTAGCTACAACGAATTTTCACTT-3 SEQ ID NO: 49

Experimental Screening of DNAzymes.

[0115] DNAzyme sequences were screened for activity using RT-PCR in T47D breast cancer cell lines that are known to express high levels of the GATA-3 mRNA (FIG. 4). Each oligonucleotide was incubated with the cells for 24 hours prior to collect mRNA, reverse transcription and then running RT-PCR. Lipofectamine delivery was performed using of the soluble oligonucleotides. The expression levels were normalized and the data analyzed using standard methods. Each experiment was run in triplicate, and each well was analyzed several times to obtain the mean GATA-3 expression level in each sample. Several of the tested sequences showed activity that was greater than HGD-40.

Screening of Gold Nanoparticle DNAzymes Conjugates

[0116] Activity of the top DzNP conjugates were tested and compared against a HGD-40 DzNP. FIG. 5 show the results of the screen and confirm that several of the hits were indeed more potent than HGD40. All the DzNP samples

were prepared at 10 nM concentration and incubated in standard serum-containing cell media for 24 hrs prior to measuring mRNA levels.

[0117] Dz2251-NP conjugates is active in vivo Animal studies were performed to compare the activity of HGD40 DzNPs against Dz2251 DzNPs (FIG. 6A-C and FIG. 7). The house dust mite antigen model of mouse asthma was used. Four groups of animals were tested. The first group were untreated with antigen and show baseline airway resistance when the animals were challenged with methacholine. The second group received the house dust mite antigen but did not receive any DNAzyme treatments. The third and fourth groups were treated with both the allergen as well as the HGD40-DzNP and the Dz2251 DzNP, respectively. The results are consistent with findings that indicate that HGD40 DzNP is active in improvement airway function. Importantly, Dz2251 showed a higher level of efficacy at all levels of methacholine challenge. Dz2251 also showed greater GATA-3 mRNA knockdown in vivo. These results indicate that Dz2251-DzNPs are more active in treating the Th2 endotype of asthma as well as other inflammatory diseases such as ulcerative colitis.

SEQUENCE LISTING

```

Sequence total quantity: 55
SEQ ID NO: 1          moltype = DNA length = 31
FEATURE              Location/Qualifiers
source                1..31
                     mol_type = other DNA
                     organism = synthetic construct

SEQUENCE: 1
tgggcttaag gctagctaca acgagagggg c                               31

SEQ ID NO: 2          moltype = DNA length = 31
FEATURE              Location/Qualifiers
source                1..31
                     mol_type = other DNA
                     organism = synthetic construct

SEQUENCE: 2
tggctctgagg ctactgtacaa cgaagttcgc a                               31

SEQ ID NO: 3          moltype = DNA length = 34
FEATURE              Location/Qualifiers
source                1..34
                     mol_type = other DNA
                     organism = synthetic construct

SEQUENCE: 3
gtctgttaat aggctagcta caacgatgtg aagc                               34

SEQ ID NO: 4          moltype = DNA length = 32
FEATURE              Location/Qualifiers
source                1..32
                     mol_type = other DNA
                     organism = synthetic construct

SEQUENCE: 4

```

-continued

tgctagacag gctagctaca acgattttcg gt	32
SEQ ID NO: 5	moltype = DNA length = 33
FEATURE	Location/Qualifiers
source	1..33
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 5	
tttgctagag gctagctaca acgaattttt cgg	33
SEQ ID NO: 6	moltype = DNA length = 33
FEATURE	Location/Qualifiers
source	1..33
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 6	
ctttttggag gctagctaca acgattgcta gac	33
SEQ ID NO: 7	moltype = DNA length = 34
FEATURE	Location/Qualifiers
source	1..34
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 7	
ttttttgcag gctagctaca acgatttttg gatt	34
SEQ ID NO: 8	moltype = DNA length = 32
FEATURE	Location/Qualifiers
source	1..32
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 8	
gcttcagtag gctagctaca acgaactgct cc	32
SEQ ID NO: 9	moltype = DNA length = 32
FEATURE	Location/Qualifiers
source	1..32
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 9	
aggcttcagg ctagctacaa cgagatactg ct	32
SEQ ID NO: 10	moltype = DNA length = 33
FEATURE	Location/Qualifiers
source	1..33
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 10	
ggcttattca ggctagctac aacgaagatg ggg	33
SEQ ID NO: 11	moltype = DNA length = 33
FEATURE	Location/Qualifiers
source	1..33
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 11	
aatggcttag gctagctaca acgatcacag atg	33
SEQ ID NO: 12	moltype = DNA length = 33
FEATURE	Location/Qualifiers
source	1..33
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 12	
taggggtag gctagctaca acgagagtca gaa	33
SEQ ID NO: 13	moltype = DNA length = 33
FEATURE	Location/Qualifiers
source	1..33
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 13	
aataggggag gctagctaca acgaatgagt cag	33
SEQ ID NO: 14	moltype = DNA length = 34
FEATURE	Location/Qualifiers

-continued

source 1..34
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 14
 cactagagag gctagctaca acgacctgtt aaat 34

SEQ ID NO: 15 moltype = DNA length = 32
 FEATURE Location/Qualifiers
 source 1..32
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 15
 tcacagcagg ctgctacaa cgatagagac cc 32

SEQ ID NO: 16 moltype = DNA length = 34
 FEATURE Location/Qualifiers
 source 1..34
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 16
 ttttttttca ggctagctac aacgaagcac taga 34

SEQ ID NO: 17 moltype = DNA length = 34
 FEATURE Location/Qualifiers
 source 1..34
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 17
 gttcagcagg ctgctacaa cgattttttt ttca 34

SEQ ID NO: 18 moltype = DNA length = 36
 FEATURE Location/Qualifiers
 source 1..36
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 18
 atataagtta taggctagct acaacgagca atgttc 36

SEQ ID NO: 19 moltype = DNA length = 36
 FEATURE Location/Qualifiers
 source 1..36
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 19
 caatataagt taggctagct acaacgaatg caatgt 36

SEQ ID NO: 20 moltype = DNA length = 33
 FEATURE Location/Qualifiers
 source 1..33
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 20
 attccttaaa ggctagctac aacgattctt ggc 33

SEQ ID NO: 21 moltype = DNA length = 33
 FEATURE Location/Qualifiers
 source 1..33
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 21
 ggaaacaaaa ggctagctac aacgatggca gtt 33

SEQ ID NO: 22 moltype = DNA length = 33
 FEATURE Location/Qualifiers
 source 1..33
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 22
 gaaaggaaag gctagctaca acgaaaaact ggc 33

SEQ ID NO: 23 moltype = DNA length = 35
 FEATURE Location/Qualifiers
 source 1..35
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 23

-continued

ttttctttta aggctagcta caacgagcat caaac	35
SEQ ID NO: 24	moltype = DNA length = 33
FEATURE	Location/Qualifiers
source	1..33
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 24	
acaacaatta ggctagctac aacgaaggga ctt	33
SEQ ID NO: 25	moltype = DNA length = 34
FEATURE	Location/Qualifiers
source	1..34
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 25	
caaacacaag ggctagctac aacgatacag ggac	34
SEQ ID NO: 26	moltype = DNA length = 35
FEATURE	Location/Qualifiers
source	1..35
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 26	
atacaaacaa ggctagctac aacgaaatta caggg	35
SEQ ID NO: 27	moltype = DNA length = 35
FEATURE	Location/Qualifiers
source	1..35
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 27	
tacatacaaa ggctagctac aacgaacaa ttaca	35
SEQ ID NO: 28	moltype = DNA length = 35
FEATURE	Location/Qualifiers
source	1..35
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 28	
tgaattatac aggctagcta caacgaacaa acaac	35
SEQ ID NO: 29	moltype = DNA length = 36
FEATURE	Location/Qualifiers
source	1..36
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 29	
tttgaattat aggctagcta caacgaatac aaacaa	36
SEQ ID NO: 30	moltype = DNA length = 33
FEATURE	Location/Qualifiers
source	1..33
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 30	
tcctttctta ggctagctac aacgatttgg tgc	33
SEQ ID NO: 31	moltype = DNA length = 37
FEATURE	Location/Qualifiers
source	1..37
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 31	
ataaatctac aggctagcta caacgacttt tcttatt	37
SEQ ID NO: 32	moltype = DNA length = 36
FEATURE	Location/Qualifiers
source	1..36
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 32	
aaataaatct aggctagcta caacgaatct tttctt	36
SEQ ID NO: 33	moltype = DNA length = 38
FEATURE	Location/Qualifiers

-continued

```

source                1..38
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 33
tgtataatat gaggctagct acaacgagaa ataaatct                38

SEQ ID NO: 34          moltype = DNA length = 37
FEATURE               Location/Qualifiers
source                1..37
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 34
tctgtataat aggctagcta caacgagatg aaataaa                37

SEQ ID NO: 35          moltype = DNA length = 35
FEATURE               Location/Qualifiers
source                1..35
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 35
gtctgtataa ggctagctac aacgaatgat gaaat                35

SEQ ID NO: 36          moltype = DNA length = 34
FEATURE               Location/Qualifiers
source                1..34
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 36
aatttataca aggctagcta caacgaagtt cggt                34

SEQ ID NO: 37          moltype = DNA length = 37
FEATURE               Location/Qualifiers
source                1..37
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 37
aaataaattt ataggctagc tacaacgaaa cagttcg                37

SEQ ID NO: 38          moltype = DNA length = 36
FEATURE               Location/Qualifiers
source                1..36
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 38
taaataaatt taggctagct acaacgaaca acagtt                36

SEQ ID NO: 39          moltype = DNA length = 36
FEATURE               Location/Qualifiers
source                1..36
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 39
cagtaaataa aggctagcta caacgattat acaaca                36

SEQ ID NO: 40          moltype = DNA length = 35
FEATURE               Location/Qualifiers
source                1..35
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 40
agcagtaaag gctagctaca acgaaaattt ataca                35

SEQ ID NO: 41          moltype = DNA length = 34
FEATURE               Location/Qualifiers
source                1..34
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 41
gttcttaaga ggctagctac aacgatagca gtaa                34

SEQ ID NO: 42          moltype = DNA length = 34
FEATURE               Location/Qualifiers
source                1..34
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 42

```

-continued

caacccaaaaa ggctagctac aacgatgaga gaga	34
SEQ ID NO: 43	moltype = DNA length = 37
FEATURE	Location/Qualifiers
source	1..37
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 43	
ttagaagcta ggctagctac aacgattaat aaatatac	37
SEQ ID NO: 44	moltype = DNA length = 32
FEATURE	Location/Qualifiers
source	1..32
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 44	
tgacatgcag gctagctaca acgagctggt ag	32
SEQ ID NO: 45	moltype = DNA length = 32
FEATURE	Location/Qualifiers
source	1..32
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 45	
tcactaacag gctagctaca acgaagaaca cg	32
SEQ ID NO: 46	moltype = DNA length = 34
FEATURE	Location/Qualifiers
source	1..34
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 46	
tgatcactaa ggctagctac aacgaacaga acac	34
SEQ ID NO: 47	moltype = DNA length = 35
FEATURE	Location/Qualifiers
source	1..35
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 47	
attattccaa ggctagctac aacgaagact gtatt	35
SEQ ID NO: 48	moltype = DNA length = 38
FEATURE	Location/Qualifiers
source	1..38
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 48	
aaatatatttc aggctagcta caacgattta ttattatg	38
SEQ ID NO: 49	moltype = DNA length = 36
FEATURE	Location/Qualifiers
source	1..36
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 49	
tagttttaaa aggctagcta caacgaattt tcactt	36
SEQ ID NO: 50	moltype = DNA length = 32
FEATURE	Location/Qualifiers
source	1..32
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 50	
tggatggagg ctagctacaa cgagtcttgg ag	32
SEQ ID NO: 51	moltype = DNA length = 15
FEATURE	Location/Qualifiers
source	1..15
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 51	
ggctagctac aacga	15
SEQ ID NO: 52	moltype = RNA length = 3069
FEATURE	Location/Qualifiers

-continued

```

source          1..3069
                 mol_type = genomic RNA
                 organism = Homo sapiens

SEQUENCE: 52
ggcgccgtct  tgatactttc  agaagaatg  cattccctgt  aaaaaaaaaa  aaaaaaact  60
gagagagggga  gagagagaga  gaagaagaga  gagagacgga  gggagagcga  gacagagcga  120
gcaacgcaat  ctgaccgagc  aggtcgtacg  ccgcccctc  ctccctctct  ctgctcttcg  180
ctaccaggt  gaccogagga  gggactccgc  ctccgagcgg  ctgaggaccc  cggtgcagag  240
gagcctggct  cgcagaattg  cagagtcgtc  gcccttttt  acaacctgtt  cccgttttat  300
tctgccgtac  ccagtttttg  gatttttgtc  ttcccttct  tctctttgt  aaacgacccc  360
tccaagataa  tttttaaaaa  accttctcct  ttgtccacct  ttgtttccca  gccttcccat  420
ccccccaccg  aaagcaaatc  attcaacgac  ccccgaccct  ccgacggcag  gagccccccg  480
acctcccagg  cggaccgccc  tccctccccg  cgcgcgggtt  ccgggcccgg  cgagagggcg  540
cgagcacagc  cgaggccatg  gaggtgacgg  cggaccagcc  gcgctgggtg  agccaccacc  600
accccgccgt  gctcaacggg  cagcaccccg  acacgcacca  cccgggcctc  agccactcct  660
acatggacgc  ggcgcagtc  ccctgccgg  aggaggtgga  tgtgttttt  aacatcgacg  720
gtcaaggcaa  ccacgtcccc  cctactacg  gaaactcgg  cagggccacg  gtgcagaggt  780
acctctcgac  ccaccacggg  agccaggtgt  gcccccgc  tctgttctat  ggatccctac  840
cctggctgga  cggcgggcaa  gccctgggca  gccaccacac  cgctccccc  tggaatctca  900
gccccttctc  caagacgttc  atccaccacg  gctccccggg  gccctctcc  gtctaccccc  960
gggctctctc  ctctctcttg  tcgggggggc  acgcagcccc  gcactcttc  acctccccgc  1020
ccaccgcc  gaaggacgtc  tccccggacc  catcgtgtc  caccocaggc  tcggccggct  1080
cggcccgga  ggacgagaaa  gagtgcctca  agtaccaggt  gccctgccc  gacagcatga  1140
agctggagtc  gtccactccc  cgtggcagca  tgaccgccct  gggtaggacc  tctctgtcga  1200
ccaccaccc  catccacacc  tacccgccct  acgtgccga  gtacagctcc  ggactcttcc  1260
ccccagcag  cctgctgggc  ggtcccccca  ccggttcgg  atgcaagtc  aggcccaagg  1320
cccggtccag  cacagaaggc  agggagtggt  tgaactgtgg  ggcaacctcg  accccactgt  1380
ggcggcgaga  tggcacggga  cactacgtgt  gcaacgcctg  cgggctctat  cacaaaatga  1440
acggacagaa  ccggcccctc  attaagccca  agcgaaggct  gtctgcagcc  aggagagcag  1500
ggacgtcctg  tgcgaaactgt  cagaccacca  caaccacact  ctggaggagg  aatgccaatg  1560
gggaccctgt  ctgcaatgcc  ctggggtct  actacaagct  tcacaatatt  aacagacccc  1620
tgactatgaa  gaagggaagg  atccagacca  gaaaccgaaa  aatgtctagc  aaatccaaaa  1680
agtgcataaa  agtgcatgac  tcaactggagg  acttcccaaa  gaacagctcg  tttaccccg  1740
ccgcctctc  cagacacatg  tctccctga  gccacatctc  gcccttcagc  cactccagcc  1800
acatgctgac  cagccccacg  ccgatgcacc  cggcatccag  cctgtcttt  ggaccacacc  1860
acccctccag  catggtcacc  gccatgggtt  agagccctgc  tcgatgtcga  caggggcccc  1920
agcgagagtc  cctgcagtc  ctctcgactt  gcatttttgc  aggagcagta  tcatgaagcc  1980
taaacgcgat  ggatatatgt  ttttgaaggc  agaaagcaaa  attatgtttg  ccactttgca  2040
aaggagctca  ctgtggtgtc  tgtgttccaa  ccaactgaatc  tggaccccat  ctgtgaataa  2100
gcattctga  ctcatatccc  ctatttaaca  ggggtctctag  tgcgttgaaa  aaaaaaatgc  2160
tgaacattgc  atataactta  tattgtaaga  aatactgtac  aatgacttta  ttgcactctg  2220
gtagctgtaa  ggcataagg  atgccaagaa  gtttaaggaa  tatgggagaa  atagtgtgga  2280
aattaagaag  aaactagatt  tgatattcaa  atggacaaac  tgcagtttt  gtttcctttc  2340
actggccaca  gttgtttgat  gcattaaaag  aaaaataaaa  aaagaaaaaa  gagaaaaaaa  2400
aaaaaaagaa  aaaagttgta  ggcgaatcat  ttgtccaaag  ctgttgccct  ctgcaaaagg  2460
aataccagtt  ctgggcaatc  agtggtaccg  ttcaccagtt  gccgttgagg  gtttcagaga  2520
gcctttttct  aggcctacat  gctttgtgaa  caagtcctgt  taattgttgt  ttgtatgtat  2580
aattcaaacg  accaaaataa  gaaaagatgt  agatttattt  catcatatta  tacagaccga  2640
actgttgat  aatttattta  ctgtagctct  taagaactgc  tttctttcgt  ttgtttgttt  2700
caatattttc  cttctctctc  aattttttgt  tgaataaaact  agattacatt  cagttggcct  2760
aaggtggttg  tgctcggagg  gtttcttgtt  tcttttccat  tttgtttttg  gatgatattt  2820
attaaatagc  ttctaagagt  ccggcggcat  ctgtcttgtc  cctattcctg  cagcctgtgc  2880
tgagggtagc  agtgtatgag  ctaccagcgt  gcatgtcagc  gacctgggcc  cgacaggcca  2940
cgtcctgcaa  tcggcccggc  tgctctctcg  ccctgtctgt  ttctgtgtta  gtgatcactg  3000
cctttaatac  agtctgttgg  aataatatta  taagcataat  aataaagtga  aaatatttta  3060
aaactacaa  3069

SEQ ID NO: 53      moltype = DNA  length = 43
FEATURE
source            Location/Qualifiers
                 1..43
                 mol_type = other DNA
                 organism = synthetic construct

SEQUENCE: 53
gtggatggag  gctagctaca  acgagtcttg  gagttttttt  ttt  43

SEQ ID NO: 54      moltype = DNA  length = 41
FEATURE
source            Location/Qualifiers
                 1..41
                 mol_type = other DNA
                 organism = synthetic construct

SEQUENCE: 54
tgatggagg  ctactacaa  cgagtcttg  attttttttt  t  41

SEQ ID NO: 55      moltype = DNA  length = 39
FEATURE
source            Location/Qualifiers
                 1..39

```

-continued

	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 55		
ggatggaggc tagctacaac gagtcttggt ttttttttt		39

1. A method of treating asthma comprising administering an effective amount of a nanoparticle coated with a nucleobase polymer comprising ATTCCTTAAAGGCTAGCTACAACGATTCT TGGC-3 (SEQ ID NO: 20) to a subject in need thereof.

2. The method of claim 1, wherein administration of the nanoparticle is in combination with a respiratory agent.
3. The method of claim 2, wherein the respiratory agent is a corticosteroid, bronchodilator, albuterol, ipratropium, or combinations thereof.

4. The method of claim 1, wherein the subject is a human subject.

* * * * *