



US 20250276997A1

(19) **United States**(12) **Patent Application Publication**
ANSARI et al.(10) **Pub. No.: US 2025/0276997 A1**(43) **Pub. Date: Sep. 4, 2025**(54) **METHODS AND COMPOUNDS FOR THE
TREATMENT OF GENETIC DISEASE***C07D 401/14* (2006.01)*C07D 403/14* (2006.01)*C07D 487/04* (2006.01)(71) Applicant: **Design Therapeutics, Inc.**, Carlsbad,
CA (US)(52) **U.S. Cl.**CPC *C07K 7/02* (2013.01); *A61K 47/545*(2017.08); *A61K 47/60* (2017.08); *A61P 21/00*(2018.01); *C07D 401/14* (2013.01); *C07D**403/14* (2013.01); *C07D 487/04* (2013.01)(72) Inventors: **Aseem ANSARI**, Carlsbad, CA (US);
Pratik SHAH, Carlsbad, CA (US)(21) Appl. No.: **18/600,510**

(57)

ABSTRACT(22) Filed: **Mar. 8, 2024****Related U.S. Application Data**(63) Continuation of application No. 17/056,566, filed on
Nov. 18, 2020, filed as application No. PCT/US2019/
033624 on May 22, 2019, now abandoned.(60) Provisional application No. 62/674,940, filed on May
22, 2018.**Publication Classification**(51) **Int. Cl.***C07K 7/02* (2006.01)*A61K 47/54* (2017.01)*A61K 47/60* (2017.01)*A61P 21/00* (2006.01)

The present disclosure relates to compounds and methods for modulating the expression of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1 and treating diseases and conditions in which dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1 plays an active role. The compound can be a transcription modulator molecule having a first terminus, a second terminus, and oligomeric backbone, wherein: a) the first terminus comprises a DNA-binding moiety capable of non-covalently binding to a nucleotide repeat sequence CAG or CTG; b) the second terminus comprises a protein-binding moiety binding to a regulatory molecule that modulates an expression of a gene comprising the nucleotide repeat sequence CAG or CTG; and c) the oligomeric backbone comprising a linker between the first terminus and the second terminus.

Specification includes a Sequence Listing.

METHODS AND COMPOUNDS FOR THE TREATMENT OF GENETIC DISEASE

CROSS REFERENCE

[0001] This application is a continuation of U.S. application Ser. No. 17/056,566, filed Nov. 18, 2020, which is the National Phase application of PCT/US2019/033624, filed May 22, 2019, which claims the benefit of U.S. Application No. 62/674,940 filed May 22, 2018, which are hereby incorporated by reference in their entirety.

SEQUENCE LISTING

[0002] The instant application contains a Sequence Listing which has been submitted electronically in XML format and is hereby incorporated by reference in its entirety. Said XML copy, created on May 19, 2025, is named 56009-706_301_SL.xml and is 93,184 bytes in size.

FIELD OF INVENTION

[0003] Disclosed herein are new chimeric heterocyclic polyamide compounds and compositions and their application as pharmaceuticals for the treatment of disease. Methods to modulate the expression of a target gene comprising the CAG trinucleotide repeat sequence in a human or animal subject are also provided for the treatment diseases such as myotonic dystrophy type 1, spinocerebellar ataxia, Huntington's disease, Huntington's disease-like syndrome, spinobulbar muscular atrophy, and dentatorubral-pallidoluysian atrophy.

BACKGROUND

[0004] The disclosure relates to the treatment of inherited genetic diseases characterized by overproduction of mRNA.

[0005] Myotonic dystrophy ("DM"), a member of the class of ailments known as muscular dystrophy, affects approximately 1 in 8000 people. DM is the most common form of muscular dystrophy among adult-onset patients, with most DM cases being diagnosed after age 20. DM is characterized by the persistence of muscular contraction, and is associated with several symptoms, including muscular disorders and cataracts, and cardiac and respiratory disorders, both of which typically are seen later in the progression of the disease. Although treatment is available for the amelioration of associated symptoms, no cure is currently employed that can stop or reverse the progression of DM. Respiratory failure and cardiac dysrhythmia account for the most common causes of death amongst DM patients.

[0006] The most severe form of DM is DM1, or myotonic dystrophy type 1. DM1 is an autosomal dominant genetic disease, caused by a mutation of the dmpk gene. This gene codes for the myotonic dystrophy protein kinase ("MDPK") protein, also known as myotonin-protein kinase. The MDPK protein can be found in muscular, cardiac, and neural tissue.

[0007] DM1 is induced by transcription of the defective dmpk gene in DM1 subjects. Normally, this gene contains a 3' untranslated region with a count of 5-37 CTG trinucleotide repeats. In the DM1 genotype, this trinucleotide is expanded to a count of 50 to over 3,000 repeats, with most having over 1,000 repeats of the CTG sequence. The count tends to increase in descendants, resulting in an earlier age of onset for later generations. Furthermore, the count has been observed to increase in a subject's lifetime, due possibly to aberrant DNA repair.

[0008] The progression of DM1 is attributed to "RNA toxicity" from dmpk mRNA having the expanded CTG region. This mRNA forms aggregates with certain proteins, and these aggregates interfere with the normal cellular function. Binding of defective mRNA to muscle blind proteins is perhaps a mechanism leading to the symptoms of DM1, particularly since muscle blind protein activity is required for proper muscle development in flies.

[0009] Spinocerebellar ataxia refers to a family of genetic diseases that is characterized by neuronal degeneration, particularly in the cerebellum. Symptoms are generally related to loss of motor function, and include incoordination of gait, poor coordination of manual and eye movements, dysarthria (unclear speech) and related complications such as poor nutrition due to dysphagia.

[0010] Several subclasses of spinocerebellar ataxia have been identified, of which a number are linked with the presence of oligonucleotide repeat sequences. Of these, several are associated with multiple copies of the CAG trinucleotide repeat sequence. In six of the subclasses: SCA1, SCA2, SCA3, SCA6, SCA7, and SCA17, the CAG trinucleotide repeat sequences give rise to polyglutamine ("poly-Q") repeat sequences in the coded proteins. These subclasses of SCA, along with Huntington's disease, dentatorubral-pallidoluysian atrophy, and spinobulbar muscular atrophy, have been collectively termed "polyglutamine expansion disorders." The exact mechanism that links the defective poly-Q proteins to the observed pathology is not always clear; aggregation of the protein, as well as formation of ubiquitinated inclusion bodies, has been proposed.

[0011] In contrast to the above group of diseases, the CAG trinucleotide repeat sequence of SCA12 is located outside of the coding region of the gene. Thus, although the mRNA contains the CAG trinucleotide repeat sequence, translation of the mRNA does not produce a poly-Q tract. The pathology associated with this defect may be due to the failure of normal cellular mechanisms to break down the abnormal mRNA, perhaps due to the presence of stable hairpin structures, leading to accumulation of the mRNA in the cell.

[0012] Spinocerebellar ataxia type 1 ("SCA1") is associated with the presence of the CAG trinucleotide repeat sequence in the atxn1 gene. Afflicted individuals have 39 or more of the trinucleotide repeat sequence; age of onset of symptoms is inversely correlated with a higher count of trinucleotide repeat sequences. The condition is generally fatal within 10-30 years; no curative treatment is currently available. The CAG trinucleotide repeat sequence is observed in mRNA as well as in genomic DNA. The gene codes for a protein termed ATXN1 which contains a poly-Q tract from the CAG trinucleotide repeat sequences. Animal studies indicate that protein toxicity, and not loss of function, is the primary mechanism responsible for the pathology of defective ATXN1. Degradation of defective ATXN1 by the proteasome is impaired, leading to accumulation of the protein.

[0013] Spinocerebellar ataxia type 2 ("SCA2") is associated with the presence of the CAG trinucleotide repeat sequence in the atxn2 gene. Afflicted individuals have 32 or more of the trinucleotide repeat sequences; age of onset of symptoms is inversely correlated with a higher count of trinucleotide repeat sequences. The gene codes for a protein termed ATXN2 which contains a poly-Q tract from the CAG trinucleotide repeat sequences. The function of the ATXN2 protein is not well understood: it is cytoplasmic and asso-

ciated with Golgi bodies and the endoplasmic reticulum. Regulation of mRNA translation is suggested by the RNA binding property of ATXN2.

[0014] Spinocerebellar ataxia type 3 (“SCA3”) is associated with the presence of the CAG trinucleotide repeat sequence in the atxn3 gene. Afflicted individuals have 50 or more copies of the trinucleotide repeat sequence; age of onset of symptoms is inversely correlated with a higher count of trinucleotide repeat sequences. The gene codes for a protein termed ATXN3 which contains a poly-Q tract from the CAG trinucleotide repeat sequences. The ATXN3 protein plays a role in the ubiquitin/proteasome mechanism for the metabolism of proteins: after a protein is marked for metabolism by ubiquitination, and before degradation of the protein by the proteasome, ATXN3 removes the ubiquitin for recycling. Defective ATXN3 containing a poly-Q tract loses this catalytic property, thus leading to a build-up of unwanted proteins.

[0015] Spinocerebellar ataxia type 6 (“SCA6”) is associated with the presence of the CAG trinucleotide repeat sequence in the cacna1a gene. Afflicted individuals have 20 or more of the trinucleotide repeat sequences. Average onset of symptoms is 45 years; the disease progresses slowly, and the duration of the disease can span over 25 years. Treatment for the disease is supportive, with acetazolamide providing relief from ataxia. The gene codes for the alpha-1 subunit of the CaV2.1 calcium channel, which is essential for proper neuronal function. The alpha-1 subunit produced by the defective cacna1a gene in afflicted individuals migrates to the cytoplasm as well as the cell membrane, where it forms aggregates. The mechanism that leads to the observed symptoms is unclear, although malfunction of the calcium channel is suspected, as well as the formation of a toxic C-terminal segment from posttranslational cleavage of the expanded protein.

[0016] Spinocerebellar ataxia type 7 (“SCA7”) is associated with the presence of the CAG trinucleotide repeat sequence in the atxn7 gene. Afflicted individuals have from 36 to over 300 of the trinucleotide repeat sequences. Onset of symptoms is typically observed in the second through fourth decade, with earlier onset correlating with more severe symptoms. In addition to the symptoms observed for the SCA class of diseases, subjects with SCA7, particularly subjects with earlier onset, can experience degradation of vision and blindness. Treatment for the disease is supportive only. The gene codes for the ataxin-7 protein, a nuclear protein that plays a role in transcription. In addition, the defective gene product interferes with cone-rod homeobox protein, providing an explanation for the retinopathy observed for this syndrome. Proteolytic cleavage of mutant ataxin-7 and transneuronal responses may be responsible for the pathogenesis of SCA7.

[0017] Spinocerebellar ataxia type 17 (“SCA17”) is associated with the presence of the CAG trinucleotide repeat sequence, with CAA interruptions, in the TATA box-binding protein (“TBP”) gene. The TBP gene product plays a role in the initiation of transcription. Afflicted individuals typically have 47 or more of the trinucleotide repeat sequences. Onset of symptoms is typically observed by age 50, with dysphagia often leading to aspiration and death. The link between the expanded CAG sequence and the observed pathology is unclear, with both gain-of-function and loss-of-function being suggested at different repeat lengths.

[0018] Huntington’s disease (“HD”) was first identified in the late 19th century as an autosomal dominant, neurodegenerative disorder. The symptoms of HD, which include a range of movement, cognitive and psychiatric disorders, generally appear in adulthood. HD is associated with the presence of the CAG trinucleotide repeat sequence in the htt gene, which codes for a protein termed huntingtin. Subjects with more than about 36 trinucleotide repeat sequences generally present with symptoms of HD, with a larger number of trinucleotide repeat sequences associated with an earlier onset of symptoms. Pathology stems from a cascade of steps: production of poly-Q huntingtin, followed by fragmentation of the elongated huntingtin into smaller peptides, which bind together and accumulate in neurons. The effects of this cascade are pronounced in the basal ganglia and cortex of the brain.

[0019] Huntington’s disease-like syndrome refers to a group of ailments whose symptoms are similar to those of Huntington’s disease, but which lack the characteristic mutation in the htt gene. Huntington’s disease-like 2 syndrome (“HDL2”) is associated with count of about 40 or more CAG trinucleotide repeat sequences in the juncophilin 3 (“jph3”) gene. HDL2 is a genetic disorder that has been seen in subjects with African lineage. Age of onset is inversely correlated with the number of trinucleotide repeat sequences. Symptoms of this syndrome include dystonia and chorea (uncontrolled movements), emotional disruptions, dysarthria, bradykinesia, inability to incorporate new learning, and difficulty in making decisions. Life expectancy can range from a few years post diagnosis to over a decade. The current theory holds that a poly-Q protein that is coded by the jph3 gene forms aggregates in neuronal cells that is responsible for the pathology of the disease. However, evidence suggesting toxic gain-of-function of mRNA has also been uncovered, indicating a possible dual pathway for pathology.

[0020] Spinobulbar muscular atrophy, also known as Kennedy disease, is an X-linked genetic disease observed in males whose symptoms include muscle atrophy, dysarthria and dysphagia due to bulbar muscles in the face and throat, fasciculations (involuntary twitches), and infertility. This disease is linked to the presence of the CAG trinucleotide repeat sequences in the androgen receptor (“ar”) gene. Pathology is thought to be due to the accumulation of fragments of the androgen receptor protein in nerve cells of the brain and spinal cord. Treatment is limited to management of symptoms; neither anti-androgen drugs nor testosterone or analogues display efficacy. Recent studies suggest that pathology of the poly-Q androgen receptor is due to inhibition of the ubiquitin ligase anaphase-promoting complex/cyclosome (APC/C), followed by disruptions in neurite formation and in the cell cycle.

[0021] Dentatorubral-pallidolysian atrophy (“DRPLA”) is an autosomal dominant genetic disorder observed mostly in subjects of Japanese descent whose symptoms, which typically appear in adulthood, include seizures, ataxia, and myoclonus (involuntary spasmodic muscular contractions). Also observed in adult subjects are dementia and psychiatric disorders. DRPLA is linked with a CAG trinucleotide repeat sequence in the atrophin-1 (“atn1”) gene. Healthy individuals have fewer than about 34 trinucleotide repeat sequences; afflicted individuals generally have more than about 50 trinucleotide repeat sequences. The atrophin-1 protein (“ATN1”) is expressed in all tissue, but is proteolytically

cleaved in neurons, suggesting a role in neural activity. Toxicity is thought to be due to accumulation of the ATN1 protein.

[0022] In certain embodiments, the mechanism set forth above will provide an effective treatment for a disease or disorder which is characterized by the presence of an excessive count of CAG or CTG trinucleotide repeat sequences in a target gene. In some embodiments, the pathology of the disease or disorder is due to the presence of mRNA containing an excessive count of CAG or GTG trinucleotide repeat sequences. In some embodiments, the pathology of the disease or disorder is due to the presence of a translation product containing an excessive count of glutamine amino acid residues. In some embodiments, the pathology of the disease or disorder is due to a loss of function in the translation product. In some embodiments, the pathology of the disease or disorder is due to a gain of function in the translation product. In some embodiments, the pathology of the disease or disorder can be alleviated by increasing the rate of transcription of the defective gene. In some embodiments, the pathology of the disease or disorder can be alleviated by decreasing the rate of transcription of the defective gene.

[0023] In certain embodiments, the mechanism set forth above will provide an effective treatment for DM1, which is caused by the overexpression of dmpk. Correction of the overexpression of the defective dmpk gene thus represents a promising method for the treatment of DM1.

[0024] In certain embodiments, the mechanism set forth above will provide an effective treatment for spinocerebellar ataxia type 1, which is caused by expression of a defective atxn1 gene. In certain embodiments, the mechanism set forth above will provide an effective treatment for spinocerebellar ataxia type 2, which is caused by expression of a defective atxn2 gene. In certain embodiments, the mechanism set forth above will provide an effective treatment for spinocerebellar ataxia type 3, which is caused by expression of a defective atxn3 gene. In certain embodiments, the mechanism set forth above will provide an effective treatment for spinocerebellar ataxia type 6, which is caused by expression of a defective cacna1a gene. In certain embodiments, the mechanism set forth above will provide an effective treatment for spinocerebellar ataxia type 7, which is caused by expression of a defective atxn7 gene. In certain embodiments, the mechanism set forth above will provide an effective treatment for spinocerebellar ataxia type 12, which is caused by expression of a defective ppp2r2b gene. In certain embodiments, the mechanism set forth above will provide an effective treatment for spinocerebellar ataxia type 17, which is caused by expression of a defective tbp gene. In certain embodiments, the mechanism set forth above will provide an effective treatment for Huntington's disease, which is caused by expression of a defective htt gene. In certain embodiments, the mechanism set forth above will provide an effective treatment for Huntington's disease-like 2 syndrome, which is caused by expression of a defective jph3 gene. In certain embodiments, the mechanism set forth above will provide an effective treatment for spinobulbar muscular atrophy, which is caused by expression of a defective ar gene. In certain embodiments, the mechanism set forth above will provide an effective treatment for dentatorubral-pallidoluysian atrophy, which is caused by expression of a defective atn1 gene.

SUMMARY

[0025] This disclosure utilizes regulatory molecules present in cell nuclei that control gene expression. Eukaryotic cells provide several mechanisms for controlling gene replication, transcription, and/or translation. Regulatory molecules that are produced by various biochemical mechanisms within the cell can modulate the various processes involved in the conversion of genetic information to cellular components. Several regulatory molecules are known to modulate the production of mRNA and, if directed to the target gene (such as, dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1), would modulate the production of the target gene mRNA that causes diseases such as, for example, spinocerebellar ataxia, Huntington's disease, Huntington's disease-like syndrome, spinobulbar muscular atrophy, and dentatorubral-pallidoluysian atrophy, Fuchs Endothelial Corneal Dystrophy, and thus reverse the progress of these diseases.

[0026] The disclosure provides compounds and methods for recruiting a regulatory molecule into close proximity to the target gene comprising a CAG or CTG trinucleotide repeat sequence. The compounds disclosed herein contain: (a) a recruiting moiety that will bind to a regulatory molecule, linked to (b) a DNA binding moiety that will selectively bind to the target gene. The compounds will counteract the expression of defective target gene in the following manner:

[0027] (1) The DNA binding moiety will bind selectively the characteristic CAG trinucleotide repeat sequence of the target gene;

[0028] (2) The recruiting moiety, linked to the DNA binding moiety, will thus be held in proximity to the target gene;

[0029] (3) The recruiting moiety, now in proximity to the target gene, will recruit the regulatory molecule into proximity with the gene; and

[0030] (4) The regulatory molecule will modulate expression of the target gene and therefore counteract the expression of defective mRNA, by direct interaction with the gene.

[0031] The DNA binding moiety will bind selectively the characteristic CAG trinucleotide repeat sequence of atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1 or the DNA binding moiety will bind selectively the characteristic CTG trinucleotide repeat sequence of dmpk. The recruiting moiety, linked to the DNA binding moiety, will thus be held in proximity to the target gene; will recruit the regulatory molecule into proximity with the gene; and the regulatory molecule will modulate expression, and therefore counteract the production of defective target gene by direct interaction with the target gene.

[0032] The mechanism set forth above will provide an effective treatment for DM1, which is caused by the expression of defective dmpk. Additionally, the mechanism set forth above will provide an effective treatment for spinocerebellar ataxia, Huntington's disease, Huntington's disease-like syndrome, spinobulbar muscular atrophy, and dentatorubral-pallidoluysian atrophy, which is caused by the expression of defective atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1. Correction of the expression of the defective target gene thus represents an effective method for the treatment for these diseases.

[0033] The disclosure provides recruiting moieties that will bind to regulatory molecules. Small molecule inhibitors

of regulatory molecules serve as templates for the design of recruiting moieties, since these inhibitors generally act via noncovalent binding to the regulatory molecules.

[0034] The disclosure further provides for DNA binding moieties that will selectively bind to one or more copies of the CAG or CTG trinucleotide repeat that are characteristic of the defective target gene. Selective binding of the DNA binding moiety to the target gene, made possible due to the high CAG or CTG count associated with the defective target gene, will direct the recruiting moiety into proximity of the gene, and recruit the regulatory molecule into position to modulate gene transcription.

[0035] The DNA binding moiety will comprise a polyamide segment that will bind selectively to the target CAG or CTG sequence. Polyamides have been designed by Derivan and others that can selectively bind to selected DNA sequences. These polyamides sit in the minor groove of double helical DNA and form hydrogen bonding interactions with the Watson-Crick base pairs. Polyamides that selectively bind to particular DNA sequences can be designed by linking monoamide building blocks according to established chemical rules. One building block is provided for each DNA base pair, with each building block binding noncovalently and selectively to one of the DNA base pairs: A/T, T/A, G/C, and C/G. Following this guideline, trinucleotides will bind to molecules with three amide units, i.e. triamides. In general, these polyamides will orient in either direction of a DNA sequence.

[0036] In principle, longer DNA sequences can be targeted with higher specificity and/or higher affinity by combining a larger number of monoamide building blocks into longer polyamide chains. Ideally, the binding affinity for a polyamide would simply be equal to the sum of each individual monoamide/DNA base pair interaction. In practice, however, due to the geometric mismatch between the fairly rigid polyamide and DNA structures, longer polyamide sequences do not bind to longer DNA sequences as tightly as would be expected from a simple additive contribution. The geometric mismatch between longer polyamide sequences and longer DNA sequences induces an unfavorable geometric strain that subtracts from the binding affinity that would be otherwise expected.

[0037] The disclosure therefore provides DNA moieties that comprise triamine subunits that are connected by flexible spacers. The disclosure therefore provides DNA moieties that comprise multi-amide subunits that are connected by flexible spacers. The spacers alleviate the geometric strain that would otherwise decrease binding affinity of a larger polyamide sequence.

[0038] Disclosed herein are polyamide compounds that can bind to one or more copies of the CAG or CTG trinucleotide repeat sequence, and can modulate the expression of a target gene comprising a CAG or CTG trinucleotide repeat sequence. Treatment of a subject with these compounds will modulate expression of the defective target gene, and this can reduce the occurrence, severity, or frequency of symptoms associated with disease. Certain compounds disclosed herein will provide higher binding affinity and selectivity than has been observed previously for this class of compounds.

INCORPORATION BY REFERENCE

[0039] All publications, patents, and patent applications mentioned in this specification are herein incorporated by

reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference.

DETAILED DESCRIPTION

[0040] The transcription modulator molecule described herein represents an interface of chemistry, biology and precision medicine in that the molecule can be programmed to regulate the expression of a target gene containing nucleotide repeat CAG or CTG. As described herein, “CAG or CTG” as used herein refers to the nucleotide CAG and its complementary sequence CTG. A person skilled in the art would understand that a sequence containing CAG trinucleotide (5'-3' direction) also has CTG trinucleotide on its complementary strand; and a sequence having multiple repeats of CAG in one strand also has multiple repeats of CTG on the complementary strand. Therefore, a polyamide binding to “CAG or CTG” repeat can mean a polyamide binding to CAG and/or its complementary sequence CTG.

[0041] The transcription modulator molecule contains DNA binding moieties that will selectively bind to one or more copies of the CAG or CTG trinucleotide repeat that is characteristic of the defective target gene (e.g., *dmpk*, *atxn1*, *atxn2*, *atxn3*, *cacna1a*, *atxn7*, *ppp2r2b*, *tbp*, *htt*, *jph3*, *ar*, or *atn1*). The transcription modulator molecule also contains moieties that bind to regulatory proteins. The selective binding of the target gene will bring the regulatory protein into proximity to the target gene and thus downregulates transcription of the target gene. The molecules and compounds disclosed herein provide higher binding affinity and selectivity than has been observed previously for this class of compounds and can be more effective in treating diseases associated with the defective target gene.

[0042] Treatment of a subject with these compounds will modulate the expression of the defective target gene, and this can reduce the occurrence, severity, or frequency of symptoms associated with genetic disease (such as DM1). The transcription modulator molecules described herein recruits the regulatory molecule to modulate the expression of the defective target gene and effectively treats and alleviates the symptoms associated with diseases.

Transcription Modulator Molecule

[0043] The transcription modulator molecules disclosed herein possess useful activity for modulating the transcription of a target gene having one or more CAG or CTG repeats (e.g., *dmpk* or *atxn1*), and may be used in the treatment or prophylaxis of a disease or condition in which the target gene (e.g., *dmpk* or *atxn1*) plays an active role. Thus, in broad aspect, certain embodiments also provide pharmaceutical compositions comprising one or more compounds disclosed herein together with a pharmaceutically acceptable carrier, as well as methods of making and using the compounds and compositions. Certain embodiments provide methods for modulating the expression of *dmpk*, *atxn1*, *atxn2*, *atxn3*, *cacna1a*, *atxn7*, *ppp2r2b*, *tbp*, *htt*, *jph3*, *ar*, or *atn1*. Other embodiments provide methods for treating a *dmpk*, *atxn1*, *atxn2*, *atxn3*, *cacna1a*, *atxn7*, *ppp2r2b*, *tbp*, *htt*, *jph3*, *ar*, or *atn1*-mediated disorder in a patient in need of such treatment, comprising administering to said patient a therapeutically effective amount of a compound or composition according to the present disclosure. Also provided is the use of certain compounds disclosed herein for use in

the manufacture of a medicament for the treatment of a disease or condition ameliorated by the modulation of the expression of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1.

[0044] Some embodiments relate to a transcription modulator molecule or compound having a first terminus, a second terminus, and oligomeric backbone, wherein: a) the first terminus comprises a DNA-binding moiety capable of noncovalently binding to a nucleotide repeat sequence CAG or CTG; b) the second terminus comprises a protein-binding moiety binding to a regulatory molecule that modulates an expression of a gene comprising the nucleotide repeat sequence CAG or CTG; and c) the oligomeric backbone comprising a linker between the first terminus and the second terminus. In some embodiments, the second terminus is not a Brd4 binding moiety. In some embodiments, the nucleotide is CAG. In some embodiments, the nucleotide is CTG.

[0045] In certain embodiments, the compounds have structural Formula I:



or a salt thereof, wherein:

[0046] X comprises a recruiting moiety that is capable of noncovalent binding to a regulatory moiety within the nucleus;

[0047] Y comprises a DNA recognition moiety that is capable of noncovalent binding to one or more copies of the trinucleotide repeat sequence CAG or CTG; and

[0048] L is a linker.

[0049] Certain compounds disclosed herein may possess useful activity for modulating the transcription of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1, and may be used in the treatment and/or prophylaxis of a disease or condition in which dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1 plays an active role. Thus, in broad aspect, certain embodiments also provide pharmaceutical compositions comprising one or more compounds disclosed herein together with a pharmaceutically acceptable carrier, as well as methods of making and using the compounds and compositions. Certain embodiments provide methods for modulating the expression of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1. Other embodiments provide methods for treating a dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1-mediated disorder in a patient in need of such treatment, comprising administering to said patient a therapeutically effective amount of a compound or composition according to the present disclosure. Also provided is the use of certain compounds disclosed herein for use in the manufacture of a medicament for the treatment of a disease or condition ameliorated by the modulation of the expression of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1.

[0050] In certain embodiments, the regulatory molecule is chosen from a bromodomain-containing protein, a nucleosome remodeling factor ("NURF"), a bromodomain PHD finger transcription factor ("BPTF"), a ten-eleven translocation enzyme ("TET"), methylcytosine dioxygenase ("TET1"), a DNA demethylase, a helicase, an acetyltransferase, and a histone deacetylase ("HDAC").

[0051] In some embodiments, the first terminus is Y, and the second terminus is X, and the oligomeric backbone is L.

[0052] In certain embodiments, the compounds have structural Formula II:



[0053] or a salt thereof, wherein:

[0054] X comprises a recruiting moiety that is capable of noncovalent binding to a regulatory molecule within the nucleus;

[0055] L is a linker;

[0056] Y_1 , Y_2 , and Y_3 are internal subunits, each of which comprises a moiety chosen from a heterocyclic ring or a C_{1-6} straight chain aliphatic segment, and each of which is chemically linked to its two neighbors;

[0057] Y_0 is an end subunit which comprises a moiety chosen from a heterocyclic ring or a straight chain aliphatic segment, which is chemically linked to its single neighbor; each subunit can noncovalently bind to an individual nucleotide in the CAG or CTG repeat sequences;

[0058] n is an integer between 1 and 15, inclusive; and

[0059] $(Y_1-Y_2-Y_3)_n-Y_0$ combine to form a DNA recognition moiety that is capable of noncovalent binding to one or more copies of the trinucleotide sequences CAG or CTG.

[0060] In certain embodiments, the compounds of structural Formula II comprise a subunit for each individual nucleotide in the CAG or CTG repeat sequence. In certain embodiments, the compounds of structural Formula II comprise a subunit for each individual nucleotide in the CAG sequence. In certain embodiments, the compounds of structural Formula II comprise a subunit for each individual nucleotide in the CTG repeat sequence.

[0061] In certain embodiment, each internal subunit has an amino ($-NH-$) group and a carboxy ($-CO-$) group.

[0062] In certain embodiments, the compounds of structural Formula II comprise amide ($-NHCO-$) bonds between each pair of internal subunits.

[0063] In certain embodiments, the compounds of structural Formula II comprise an amide ($-NHCO-$) bond between L and the leftmost internal subunit.



[0064] In certain embodiments, the compounds of structural Formula II comprise an amide bond between the rightmost internal subunit and the end subunit.




[0065] In certain embodiments, each subunit comprises a moiety that is independently chosen from a heterocycle and an aliphatic chain.

[0066] In certain embodiments, the heterocycle is a monocyclic heterocycle. In certain embodiments, the heterocycle is a monocyclic 5-membered heterocycle. In certain embodiments, each heterocycle contains a heteroatom independently chosen from N, O, or S. In certain embodiments, each heterocycle is independently chosen from pyrrole, imidazole, thiazole, oxazole, thiophene, and furan.


[0067] In certain embodiments, the aliphatic chain is a C_{1-6} straight chain aliphatic chain. In certain embodiments, the aliphatic chain has structural formula $-(CH_2)_m-$, for m chosen from 1, 2, 3, 4, and 5. In certain embodiments, the aliphatic chain is $-CH_2CH_2-$.

[0068] In certain embodiments, each subunit comprises a moiety independently chosen from


















—NH-benzopyrazinylene-CO—, —NH-phenylene-CO—,
 —NH-pyridinylene-CO—, —NH-piperidinylene-CO—,
 —NH-pyrimidinylene-CO—, —NH-anthracenylene-CO—,
 —NH-quinolinylene-CO—, and




wherein Z is H, NH₂, C₁₋₆ alkyl, C₁₋₆ haloalkyl or C₁₋₆ alkyl-NH₂.

[0069] In some embodiments,







[0070] In certain embodiments, n is an integer between 1 and 5, inclusive.

[0071] In certain embodiments, n is an integer between 1 and 3, inclusive.


[0072] In certain embodiments, n is an integer between 1 and 2, inclusive.

[0073] In certain embodiments, n is 1.

[0074] In certain embodiments, L comprises a C₁₋₆ straight chain aliphatic segment.

[0075] In certain embodiments, L comprises (CH₂OCH₂)_m; and m is an integer between 1 to 20, inclusive. In certain further embodiments, m is an integer between 1 to 10, inclusive. In certain further embodiments, m is an integer between 1 to 5, inclusive.

[0076] In certain embodiments, the compounds have structural Formula III:



[0077] or a salt thereof, wherein:

[0078] X comprises a recruiting moiety that is capable of noncovalent binding to a regulatory molecule within the nucleus;

[0079] L is a linker;

[0080] Y₁, Y₂, and Y₃ are internal subunits, each of which comprises a moiety chosen from a heterocyclic ring or a C₁₋₆ straight chain aliphatic segment, and each of which is chemically linked to its two neighbors;

[0081] Y₀ is an end subunit which comprises a moiety chosen from a heterocyclic ring or a straight chain aliphatic segment, which is chemically linked to its single neighbor;


[0082] each subunit can noncovalently bind to an individual nucleotide in the CAG or CTG repeat sequence;

[0083] W is a spacer;

[0084] n is an integer between 1 and 10, inclusive; and


[0085] (Y₁-Y₂-Y₃)-(W-Y₁-Y₂-Y₃)_n-Y₀ combine to form a DNA recognition moiety that is capable of noncovalent binding to one or more copies of the trinucleotide repeat sequence CAG or CTG.

[0086] In certain embodiments, Y₁-Y₂-Y₃ is:



[0087] In certain embodiments, Y₁-Y₂-Y₃ is Im-β-Py.

[0088] In certain embodiments, the compounds have structural Formula IV:



[0089] or a salt thereof, wherein:

[0090] X comprises a recruiting moiety that is capable of noncovalent binding to a regulatory molecule within the nucleus;

[0091] Y₁, Y₂, Y₃, Y₄, Y₅, and Y₆ are internal subunits, each of which comprises a moiety chosen from a heterocyclic ring or a C₁₋₆ straight chain aliphatic segment, and each of which is chemically linked to its two neighbors;

[0092] Y₀ is an end subunit which comprises a moiety chosen from a heterocyclic ring or a straight chain aliphatic segment, which is chemically linked to its single neighbor;

[0093] each subunit can noncovalently bind to an individual nucleotide in the CAG or CTG repeat sequence;


[0094] L is a linker;

[0095] V is a turn component for forming a hairpin turn; and

[0096] (Y₁-Y₂-Y₃)-V-(Y₄-Y₅-Y₆)-Y₀ combine to form a DNA recognition moiety that is capable of noncovalent binding to one or more copies of the trinucleotide repeat sequence CAG or CTG.

[0097] In certain embodiments, V is —HN—CH₂CH₂CH₂—CO—.

[0098] In certain embodiments, the compounds have structural Formula Va:




[0099] or a salt thereof, wherein:

[0100] X comprises a recruiting moiety that is capable of noncovalent binding to a regulatory molecule within the nucleus;

[0101] Y₀ is an end subunit which comprises a moiety chosen from a heterocyclic ring or a straight chain aliphatic segment, which is chemically linked to its single neighbor; and

[0102] n is an integer between 1 and 5, inclusive.

[0103] In certain embodiments, the compounds have structural Formula VI:




[0104] or a salt thereof, wherein:

[0105] X comprises a recruiting moiety that is capable of noncovalent binding to a regulatory molecule within the nucleus;

[0106] Y₀ is an end subunit which comprises a moiety chosen from a heterocyclic ring or a straight chain aliphatic segment, which is chemically linked to its single neighbor; and

[0107] n is an integer between 1 and 5, inclusive.

[0108] In certain embodiments, the compounds have structural Formula VII:



[0109] or a salt thereof, wherein:

[0110] X comprises a recruiting moiety that is capable of noncovalent binding to a regulatory molecule within the nucleus; and

[0111] W is a spacer;

[0112] Y₀ is an end subunit which comprises a moiety chosen from a heterocyclic ring or a straight chain aliphatic segment, which is chemically linked to its single neighbor; and

[0113] n is an integer between 1 and 200, inclusive.

[0114] In certain embodiments of the compounds of structural Formula VII,


[0115] W is $\text{—NHCH}_2\text{—(CH}_2\text{OCH}_2\text{)}_p\text{—CH}_2\text{CO—}$; and

[0116] p is an integer between 1 and 4, inclusive.

[0117] In certain embodiments, the compounds have structural Formula VIII:

[0123] In certain embodiments of the compounds of structural Formula VIII, V is $\text{—(CH}_2\text{)}_q\text{—NH—(CH}_2\text{)}_q\text{—}$; and q is an integer between 2 and 4, inclusive.

[0124] In some embodiments, V is $\text{—(CH}_2\text{)}_a\text{—NR}^1\text{—(CH}_2\text{)}_b\text{—}$, $\text{—(CH}_2\text{)}_a\text{—}$, $\text{—(CH}_2\text{)}_a\text{—O—(CH}_2\text{)}_b\text{—}$, $\text{—(CH}_2\text{)}_a\text{—CH(NHR}^1\text{)—}$, $\text{—(CH}_2\text{)}_a\text{—CH(NHR}^1\text{)—}$, $\text{—(CR}^2\text{R}^3\text{)}_a\text{—}$, or $\text{—(CH}_2\text{)}_a\text{—CH(NR}^1_3\text{)}^+\text{—(CH}_2\text{)}_b\text{—}$, wherein each a is independently an integer between 2 and 4; R¹ is H, an optionally substituted C₁₋₆ alkyl, an optionally substituted C₃₋₁₀ cycloalkyl, an optionally substituted C₆₋₁₀ aryl, an optionally substituted 4-10 membered heterocyclyl, or an optionally substituted 5-10 membered heteroaryl; each R² and R³ are independently H, halogen, OH, NHAc, or C₁₋₄ alkyl. In some embodiments, R¹ is H. In some embodiments, R¹ is C₁₋₆ alkyl optionally substituted by 1-3 substituents selected from —C(O)—phenyl . In some embodiments, V is



[0118] or a salt thereof, wherein:

[0119] X comprises a recruiting moiety that is capable of noncovalent binding to a regulatory molecule within the nucleus;

[0120] V is a turn component for forming a hairpin turn;

[0121] Y₀ is an end subunit which comprises a moiety chosen from a heterocyclic ring or a straight chain aliphatic segment, which is chemically linked to its single neighbor; and

[0122] n is an integer between 1 and 200, inclusive.

$\text{—(CR}^2\text{R}^3\text{)—(CH}_2\text{)}_a\text{—}$ or $\text{—(CH}_2\text{)}_a\text{—(CR}^2\text{R}^3\text{)—(CH}_2\text{)}_b\text{—}$, wherein each a is independently 1-3, b is 0-3, and each R² and R³ are independently H, halogen, OH, NHAc, or C₁₋₄ alkyl. In some embodiments, V is $\text{—(CH}_2\text{)—CH(NH}_3\text{)}^+\text{—(CH}_2\text{)—}$ or $\text{—(CH}_2\text{)—CH}_2\text{CH(NH}_3\text{)—}$.

[0125] In one aspect, the compounds of the present disclosure bind to the CAG or CTG of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, ttp, htt, jph3, ar, or atn1 and recruit a regulatory moiety to the vicinity of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, ttp, htt, jph3, ar, or atn1. The regulatory moiety, due to its proximity to the gene,

will be more likely to modulate the expression of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1.

[0126] Also provided are embodiments wherein any compound disclosed above, including compounds of Formulas I-VIII, are singly, partially, or fully deuterated. Methods for accomplishing deuterium exchange for hydrogen are known in the art.

[0127] Also provided are embodiments wherein any embodiment above may be combined with any one or more of these embodiments, provided the combination is not mutually exclusive.

[0128] As used herein, two embodiments are “mutually exclusive” when one is defined to be something which is different than the other. For example, an embodiment wherein two groups combine to form a cycloalkyl is mutually exclusive with an embodiment in which one group is ethyl the other group is hydrogen. Similarly, an embodiment wherein one group is CH₂ is mutually exclusive with an embodiment wherein the same group is NH.

[0129] In one aspect, the compounds of the present disclosure bind to the CAG or CTG of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1 and recruit a regulatory moiety to the vicinity of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1. The regulatory moiety, due to its proximity to the gene, will be more likely to modulate the expression of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1.

[0130] In some embodiments, the molecules described herein bind to the CAG of atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, and atn1. In some embodiments, the molecules described herein bind to the CAG of the gene encoding TCF4. In some embodiments, the molecules of the present disclosure bind to the CTG of dmpk. In some embodiments, the molecules of the present disclosure bind to the CAG of TCF4 gene.

[0131] In one aspect, the molecules of the present disclosure provide a polyamide sequence for interaction of a single polyamide subunit to each base pair in the CAG or CTG repeat sequence. In one aspect, the molecules of the present disclosure provide a turn component V, in order to enable hairpin binding of the molecule to the CAG or CTG, in which each nucleotide pair interacts with two subunits of the polyamide.

[0132] In one aspect, the molecules of the present disclosure provide more than one copy of the polyamide sequence for noncovalent binding to the CAG or CTG. In one aspect, the molecules of the present disclosure bind to dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1 with an affinity that is greater than a corresponding molecule that contains a single polyamide sequence.

[0133] In one aspect, the molecules of the present disclosure provide more than one copy of the polyamide sequence for noncovalent binding to the CAT or CTG, and the individual polyamide sequences in this molecule are linked by a spacer W, as defined above. The spacer W allows this molecule to adjust its geometry as needed to alleviate the geometric strain that otherwise affects the noncovalent binding of longer polyamide sequences.

[0134] In certain embodiments, the molecules comprise a cell-penetrating ligand moiety.

[0135] In certain embodiments, the cell-penetrating ligand moiety is a polypeptide.

[0136] In certain embodiments, the cell-penetrating ligand moiety is a polypeptide containing fewer than 30 amino acid residues.

[0137] In certain embodiments, the polypeptide is chosen from any one of SEQ ID NO. 1 to SEQ ID NO. 37, inclusive.

First Terminus—DNA Binding Moiety

[0138] The first terminus interacts and binds with the gene, particularly with the minor grooves of the CAG or CTG sequence. In one aspect, the molecules of the present disclosure provide a polyamide sequence for interaction of a single polyamide subunit to each base pair in the CAG or CTG repeat sequence. In one aspect, the molecules of the present disclosure provide a turn component (e.g., aliphatic amino acid moiety), in order to enable hairpin binding of the molecule to the CAG or CTG, in which each nucleotide pair interacts with two subunits of the polyamide.

[0139] In one aspect, the molecules of the present disclosure provide more than one copy of the polyamide sequence for noncovalent binding to CAG or CTG. In one aspect, the molecules of the present disclosure bind to dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1 with an affinity that is greater than a corresponding molecule that contains a single polyamide sequence. In one aspect, the molecules of the present disclosure bind to dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1 with an affinity that is greater than a corresponding molecule that contains a single polyamide sequence.

[0140] In one aspect, the molecules of the present disclosure provide more than one copy of the polyamide sequence for noncovalent binding to the CAG or CTG, and the individual polyamide sequences in this molecule are linked by a spacer W, as defined above. The spacer W allows this molecule to adjust its geometry as needed to alleviate the geometric strain that otherwise affects the noncovalent binding of longer polyamide sequences.

[0141] In certain embodiments, the DNA recognition or binding moiety binds in the minor groove of DNA.

[0142] In certain embodiments, the DNA recognition or binding moiety comprises a polymeric sequence of monomers, wherein each monomer in the polymer selectively binds to a certain DNA base pair.

[0143] In certain embodiments, the DNA recognition or binding moiety comprises a polyamide moiety.

[0144] In certain embodiments, the DNA recognition or binding moiety comprises a polyamide moiety comprising heteroaromatic monomers, wherein each heteroaromatic monomer binds noncovalently to a specific nucleotide, and each heteroaromatic monomer is attached to its neighbor or neighbors via amide bonds.

[0145] In certain embodiments, the DNA recognition moiety binds to a sequence comprising at least 1000 nucleotide repeats. In certain embodiments, the DNA recognition moiety binds to a sequence comprising at least 500 nucleotide repeats. In certain embodiments, the DNA recognition moiety binds to a sequence comprising at least 200 nucleotide repeats. In certain embodiments, the DNA recognition moiety binds to a sequence comprising at least 100 nucleotide repeats. In certain embodiments, the DNA recognition moiety binds to a sequence comprising at least 50 nucleotide repeats. In certain embodiments, the DNA recognition moiety binds to a sequence comprising at least 20 nucleotide repeats.

[0146] In some embodiments, the nucleotide is CAG. In some embodiments, the nucleotide is CTG.

[0147] In certain embodiments, the molecules have structural Formula II:



[0148] or a salt thereof, wherein:

[0149] X comprises a recruiting moiety that is capable of noncovalent binding to a regulatory molecule within the nucleus;

[0150] L is a linker;

[0151] Y_1 , Y_2 , and Y_3 are internal subunits, each of which comprises a moiety chosen from a heterocyclic ring or a C_{1-6} straight chain aliphatic segment, and each of which is chemically linked to its two neighbors;

[0152] Y_0 is an end subunit which comprises a moiety chosen from a heterocyclic ring or a straight chain aliphatic segment, which is chemically linked to its single neighbor;

[0153] each subunit can noncovalently bind to an individual nucleotide in the CAG or CTG repeat sequence;

[0154] n is an integer between 1 and 15, inclusive; and

[0155] $(Y_1-Y_2-Y_3)_n-Y_0$ combine to form a DNA recognition moiety that is capable of noncovalent binding to one or more copies of the trinucleotide repeat sequence CAG or CTG.

[0156] In certain embodiments, the molecules of structural Formula II comprise a subunit for each individual nucleotide in the CAG or CTG repeat sequence.

[0157] In certain embodiment, each internal subunit has an amino ($-\text{NH}-$) group and a carboxy ($-\text{CO}-$) group.

[0158] In certain embodiments, the molecules of structural Formula II comprise amide ($-\text{NHCO}-$) bonds between each pair of internal subunits.

[0159] In certain embodiments, the molecules of structural Formula II comprise an amide ($-\text{NHCO}-$) bond between L and the leftmost internal subunit.

[0160] In certain embodiments, the molecules of structural Formula II comprise an amide bond between the rightmost internal subunit and the end subunit.

[0161] In certain embodiments, each subunit comprises a moiety that is independently chosen from a heterocycle and an aliphatic chain.

[0162] In certain embodiments, the heterocycle is a monocyclic heterocycle. In certain embodiments, the heterocycle is a monocyclic 5-membered heterocycle. In certain embodiments, each heterocycle contains a heteroatom independently chosen from N, O, or S. In certain embodiments, each heterocycle is independently chosen from pyrrole, imidazole, thiazole, oxazole, thiophene, and furan.

[0163] In certain embodiments, the aliphatic chain is a C_{1-6} straight chain aliphatic chain. In certain embodiments, the aliphatic chain has structural formula $-(\text{CH}_2)_m-$, for m chosen from 1, 2, 3, 4, and 5. In certain embodiments, the aliphatic chain is $-\text{CH}_2\text{CH}_2-$.

[0164] The form of the polyamide selected can vary based on the target gene. The first terminus can include a polyamide selected from the group consisting of a linear polyamide, a hairpin polyamide, a H-pin polyamide, an overlapped polyamide, a slipped polyamide, a cyclic polyamide, a tandem polyamide, and an extended polyamide. In some embodiments, the first terminus comprises a linear polyamide. In some embodiments, the first terminus comprises a hairpin polyamide.

[0165] The binding affinity between the polyamide and the target gene can be adjusted based on the composition of the polyamide. In some embodiments, the polyamide is capable of binding the DNA with an affinity of less than about 600 nM, about 500 nM, about 400 nM, about 300 nM, about 250 nM, about 200 nM, about 150 nM, about 100 nM, or about 50 nM. In some embodiments, the polyamide is capable of binding the DNA with an affinity of less than about 300 nM. In some embodiments, the polyamide is capable of binding the DNA with an affinity of less than about 200 nM. In some embodiments, the polyamide is capable of binding the DNA with an affinity of greater than about 200 nM, about 150 nM, about 100 nM, about 50 nM, about 10 nM, or about 1 nM. In some embodiments, the polyamide is capable of binding the DNA with an affinity in the range of about 1-600 nM, 10-500 nM, 20-500 nM, 50-400 nM, or 100-300 nM.

[0166] The binding affinity between the polyamide and the target DNA can be determined using a quantitative footprint titration experiment. The experiment involve measuring the dissociation constant K_d of the polyamide for target sequence at either 24° C. or 37° C., and using either standard polyamide assay solution conditions or approximate intracellular solution conditions.

[0167] The binding affinity between the regulatory protein and the ligand on the second terminus can be determined using an assay suitable for the specific protein. The experiment involve measuring the dissociation constant K_d of the ligand for protein and using either standard protein assay solution conditions or approximate intracellular solution conditions.

[0168] In some embodiments, the first terminus comprises $-\text{NH-Q-C(O)}-$, wherein Q is an optionally substituted C_{6-10} arylene group, optionally substituted 4-10 membered heterocyclene, optionally substituted 5-10 membered heteroarylene group, or an optionally substituted alkylene group. In some embodiments, Q is an optionally substituted C_{6-10} arylene group or optionally substituted 5-10 membered heteroarylene group. In some embodiments, Q is an optionally substituted 5-10 membered heteroarylene group. In some embodiments, the 5-10 membered heteroarylene group is optionally substituted with 1-4 substituents selected from H, OH, halogen, C_{1-10} alkyl, NO_2 , CN, NR'' , C_{1-6} haloalkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, $(C_{1-6}$ alkoxy) C_{1-6} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{3-7} carbocyclyl, 4-10 membered heterocyclyl, C_{6-10} aryl, 5-10 membered heteroaryl, $(C_{3-7}$ carbocyclyl) C_{1-6} alkyl, (4-10 membered heterocyclyl) C_{1-6} alkyl, $(C_{6-10}$ aryl) C_{1-6} alkyl, $(C_{6-10}$ aryl) C_{1-6} alkoxy, (5-10 membered heteroaryl) C_{1-6} alkyl, $(C_{3-7}$ carbocyclyl)amine, (4-10 membered heterocyclyl)amine, $(C_{6-10}$ aryl)amine, (5-10 membered heteroaryl)amine, acyl, C-carboxy, O-carboxy, C-amido, N-amido, S-sulfonamido, N-sulfonamido, $-\text{SR}'$, COOH, or CONR'' ; wherein each R' and R'' are independently H, C_{1-10} alkyl, C_{1-10} haloalkyl, C_{1-10} alkoxy.

[0169] In some embodiments, the first terminus comprises at least three aromatic carboxamide moieties selected to correspond to the nucleotide repeat sequence CAG or CTG and at least one aliphatic amino acid residue chosen from the group consisting of glycine, β -alanine, γ -aminobutyric acid, 2,4-diaminobutyric acid, and 5-aminovaleic acid. In some embodiments, the first terminus comprises at least one β -alanine subunit.

[0170] In some embodiments, the monomer element is independently selected from the group consisting of option-

ally substituted pyrrole carboxamide monomer, optionally substituted imidazole carboxamide monomer, optionally substituted C—C linked heteromonocyclic/heterobicyclic moiety, and β -alanine.

[0171] In some embodiments, the first terminus comprises a structure of Formula (A-1):




[0172] wherein:

[0173] each $[A-M]$ appears p times and p is an integer in the range of 1 to 10,

[0174] L_{1a} is a bond, a C_{1-6} alkylene, $-NR^a-C_{1-6}$ alkylene- $C(O)-$, $-NR^aC(O)-$, $-NR^a-C_{1-6}$ alkylene, $-O-$, or $-O-C_{1-6}$ alkylene;

[0175] each A is selected from the group consisting of a bond, C_{1-10} alkylene, optionally substituted C_{6-10} arylene group, optionally substituted 4-10 membered heterocyclene, optionally substituted 5-10 membered heteroarylene group, $-C_{1-10}$ alkylene- $C(O)-$, $-C_{1-10}$ alkylene- NR^a- , $-CO-$, $-NR^a-$, $CONR^a-$, $-CONR^aC_{1-4}$ alkylene-, $-NR^aCO-C_{1-4}$ alkylene-, $-C(O)O-$, $-O-$, $-S-$, $-S(O)-$, $-S(O)_2-$, $-C(=S)-NH-$, $-C(O)-NH-NH-$, $-C(O)-N=N-$, $-C(O)-CH=CH-$, $(CH_2)_{0-4}-CH=CH-(CH_2)_{0-4}$, $-N(CH_3)-C_{1-6}$ alkylene, and



$-NH-C_{1-6}$ alkylene- $NH-$, $-O-C_{1-6}$ alkylene- $O-$, $-NH-N=N-$, $-NH-C(O)-NH-$, and any combinations thereof, and at least one A is $-CONH-$;

[0176] each M is an optionally substituted C_{6-10} arylene group, optionally substituted 4-10 membered heterocyclene, optionally substituted 5-10 membered heteroarylene group, or an optionally substituted alkylene;


[0177] E_1 is H or $-A^E-G$;

[0178] A^E is absent or $-NHCO-$;

[0179] G is selected from the group consisting of optionally substituted C_{6-10} aryl, optionally substituted 4-10 membered heterocyclyl, optionally substituted 5-10 membered heteroaryl, an optionally substituted C_{1-6} alkyl, C_{0-4} alkylene-NHC(=NH)NH, $-CN$, $-C_{0-4}$ alkylene- $C(=NH)(NR^aR^b)$, $-C_{0-4}$ alkylene- $C(=N^+H_2)(NR^aR^b)C_{1-5}$ alkylene- NR^aR^b , C_{0-4} alkylene-NHC(=NH) R^a , and optionally substituted amine; and


[0180] each R^a and R^b are independently selected from the group consisting of H, an optionally substituted C_{1-6} alkyl, an optionally substituted C_{3-10} cycloalkyl, optionally substituted C_{6-10} aryl, optionally substituted 4-10 membered heterocyclyl, and optionally substituted 5-10 membered heteroaryl.

[0181] In some embodiments, the first terminus can comprise a structure of Formula (A-2):



[0182] wherein:

[0183] L_{2a} is a linker selected from $-C_{1-12}$ alkylene- CR^a , $-CH$, N , $-C_{1-6}$ alkylene- N , $-C(O)N$, $-NR^a-C_{1-6}$ alkylene- CH , $-O-C_{0-6}$ alkylene- CH ,



[0184] each p and q are independently an integer in the range of 1 to 10;

[0185] each m and n are independently an integer in the range of 0 to 10;

[0186] each A is independently selected from a bond, C_{1-10} alkylene, $-C_{1-10}$ alkylene- $C(O)-$, $-C_{1-10}$ alkylene- NR^a- , $-CO-$, $-NR^a-$, $-CONR^a-$, $-CONR^aC_{1-4}$ alkylene-, $-NR^aCO-C_{1-4}$ alkylene-, $-C(O)O-$, $-O-$, $-S-$, $-S(O)-$, $-S(O)_2-$, $-C(=S)-NH-$, $-C(O)-NH-NH-$, $-C(O)-N=N-$, or $-C(O)-CH=CH-$, and at least one A is $CONH-$;

[0187] each M is independently an optionally substituted C_{6-10} arylene group, optionally substituted 4-10 membered heterocyclene, optionally substituted 5-10 membered heteroarylene group, or an optionally substituted alkylene;

[0188] each E_1 and E_2 are independently H or $-A^E-G$;

[0189] each A^E is independently absent or $NHCO$;


[0190] each G is independently selected from the group consisting of C_{6-10} aryl, optionally substituted 4-10 membered heterocyclyl, optionally substituted 5-10 membered heteroaryl, an optionally substituted C_{1-6} alkyl, C_{0-4} alkylene-NHC(=NH)NH, $-CN$, $-C_{0-4}$ alkylene- $C(=NH)(NR^aR^b)$, $-C_{0-4}$ alkylene- $C(=N^+H_2)(NR^aR^b)C_{1-5}$ alkylene- NR^aR^b , C_{0-4} alkylene-NHC(=NH) R^a , and optionally substituted amine; and

[0191] each R^a and R^b are independently selected from the group consisting of H, an optionally substituted C_{1-6} alkyl, an optionally substituted C_{3-10} cycloalkyl, optionally substituted C_{6-10} aryl, optionally substituted 4-10 membered heterocyclyl, and an optionally substituted 5-10 membered heteroaryl; and


[0192] each R^{1a} and R^{1b} is independently H, or C_{1-6} alkyl.

[0193] In certain embodiments, the integers p and q are $2 \leq p+q \leq 20$. In some embodiments, p is in the range of about 2 to 10. In some embodiments, p is in the range of about 4 to 8. In some embodiments, q is in the range of about 2 to 10. In some embodiments, q is in the range of about 4 to 8.


[0194] In certain embodiments, L^{2a} is $-C_{2-8}$ alkylene-CH,




and wherein each m and n is independently an integer in the range of 0 to 10. In certain embodiments, L^{2a} is




In some embodiments, L^{2a} is $-C_{2-8}$ alkylene-CH. In some embodiments, L^{2a} is



wherein (m+n) is in the range of about 1 to 4. In some embodiments, L^{2a} is



and (m+n) is in the range of about 2 to 5. In some embodiments, L^{2a} is



wherein (m+n) is in the range of about 1 to 6.

[0195] The transcription modulator molecule of claim 1, wherein the first terminus comprises a structure of Formula (A-3):



[0196] wherein:

[0197] L^{1a} is a bond, a C_{1-6} alkylene, $-\text{NH}-C_{0-6}$ alkylene- $\text{C}(\text{O})-$, $-\text{N}(\text{CH}_3)-C_{0-6}$ alkylene, or $-\text{O}-C_{0-6}$ alkylene;

[0198] L^{3a} is a bond, C_{1-6} alkylene, $-\text{NH}-C_{0-6}$ alkylene- $\text{C}(\text{O})-$, $-\text{N}(\text{CH}_3)-C_{0-6}$ alkylene, $-\text{O}-C_{0-6}$ alkylene, $-(\text{CH}_2)_a-\text{NR}^a-(\text{CH}_2)_b-$, $-(\text{CH}_2)_a-$, $-(\text{CH}_2)_a-\text{O}-(\text{CH}_2)_b-$, $-(\text{CH}_2)_a-\text{CH}(\text{NHR}^a)-$, $-(\text{CH}_2)_a-\text{CH}(\text{NHR}^a)-$, $-(\text{CR}^{1a}\text{R}^{1b})_a-$, or $-(\text{CH}_2)_a-\text{CH}(\text{NR}^a\text{R}^b)-(\text{CH}_2)_b-$;

[0199] each a and b are independently an integer between 2 and 4;


[0200] each R^a and R^b are independently selected from H, an optionally substituted C_{1-6} alkyl, an optionally substituted C_{3-10} cycloalkyl, optionally substituted C_{6-10} aryl, optionally substituted 4-10 membered heterocyclyl, and an optionally substituted 5-10 membered heteroaryl;

[0201] each R^{1a} and R^{1b} is independently H, halogen, OH, NHAc, or C_{1-4} alkyl;

[0202] each [A-M] appears p^1 times and p^1 is an integer in the range of 1 to 10;

[0203] each [M-A] appears q^1 times and q^1 is an integer in the range of 1 to 10;

[0204] each A is selected from a bond, C_{1-10} alkylene, optionally substituted C_{6-10} arylene group, optionally substituted 4-10 membered heterocyclene, optionally substituted 5-10 membered heteroarylene group, $-C_{1-10}$ alkylene- $\text{C}(\text{O})-$, $-C_{1-10}$ alkylene- NR^a- , $-\text{CO}-$, $-\text{NR}^a-$, $-\text{CONR}^a-$, $-\text{CONR}^a\text{C}_{1-4}$ alkylene-, $-\text{NR}^a\text{CO}-C_{1-4}$ alkylene-, $-\text{C}(\text{O})\text{O}-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O})-$, $-\text{S}(\text{O})_2-$, $-\text{C}(=\text{S})-\text{NH}-$, $-\text{C}(\text{O})-\text{NH}-\text{NH}-$, $-\text{C}(\text{O})-\text{N}=\text{N}-$, $-\text{C}(\text{O})-\text{CH}=\text{CH}-$, $(\text{CH}_2)_{0-4}-\text{CH}=\text{CH}-(\text{CH}_2)_{0-4}$, $-\text{N}(\text{CH}_3)-C_{1-6}$ alkylene, and



$-\text{NH}-C_{1-6}$ alkylene- $\text{NH}-$, $-\text{O}-C_{1-6}$ alkylene- $\text{O}-$, $-\text{NH}-\text{N}=\text{N}-$, $-\text{NH}-\text{C}(\text{O})-\text{NH}-$, and any combination thereof, and at least one A is NHCO;


[0205] each M in each [A-M] and [M-A] unit is independently an optionally substituted C_{6-10} arylene group, optionally substituted 4-10 membered heterocyclene, optionally substituted 5-10 membered heteroarylene group, or an optionally substituted alkylene; and

[0206] E_1 is selected from the group consisting of optionally substituted C_{6-10} aryl, optionally substituted 4-10 membered heterocyclyl, optionally substituted 5-10 membered heteroaryl, an optionally substituted C_{1-6} alkyl, C_{0-4} alkylene- $\text{NHC}(=\text{NH})\text{NH}$, $-\text{CN}$, $-C_{0-4}$ alkylene- $\text{C}(=\text{NH})(\text{NR}^a\text{R}_2)$, $-C_{0-4}$ alkylene- $\text{C}(=\text{N}^+\text{H}_2)(\text{NR}^a\text{R}^b)$, C_{1-5} alkylene- NR^aR^b , and C_{0-4} alkylene- $\text{NHC}(=\text{NH})\text{R}$.

[0207] In certain embodiments, the integers p^1 and q^1 are $2 \leq p^1+q^1 \leq 20$.

[0208] In some embodiments, for Formula (A-1) to (A-4), each A is independently a bond, C_{1-6} alkylene, optionally substituted phenylene, optionally substituted thiophenylene,


optionally substituted furanylene, $-\text{C}_{1-10}\text{ alkylene-C(O)-}$, $-\text{C}_{1-10}\text{ alkylene-NH-}$, $-\text{CO-}$, $-\text{NR}^a-$, $-\text{CONR}^a-$, $-\text{CONR}^a\text{C}_{1-4}\text{ alkylene-}$, $-\text{NR}^a\text{CO-C}_{1-4}\text{ alkylene-}$, $-\text{C(O)O-}$, $-\text{O-}$, $-\text{S-}$, $-\text{S(O)-}$, $-\text{S(O)}_2-$, $-\text{C(=S)-NH-}$, $-\text{C(O)-NH-NH-}$, $-\text{C(O)-N=N-}$, $-\text{C(O)-CH=CH-}$, $-\text{CH=CH-}$, $-\text{NH-N=N-}$, $-\text{NH-C(O)-NH-}$, $-\text{N(CH}_3\text{)-C}_{1-6}\text{ alkylene}$, and



$-\text{NH-C}_{1-6}\text{ alkylene-NH-}$, $-\text{O-C}_{1-6}\text{ alkylene-O-}$, and any combinations optionally substituted 5-10 membered heteroarylene group. In some embodiments, in Formula (A-1) and (A-3), L_{1a} is a bond. In some embodiments, in Formula (A-1) and (A-3), L_{1a} is a $\text{C}_{1-6}\text{ alkylene}$. In some embodiments, in Formula (A-1) and (A-3), L_{1a} is $-\text{NH-C}_{1-6}\text{ alkylene-C(O)-}$. In some embodiments, in Formula (A-1) and (A-3), L_{1a} is $-\text{N(CH}_3\text{)-C}_{1-6}\text{ alkylene-}$. In some embodiments, in Formula (A-1) and (A-3), L_{1a} is $-\text{O-C}_{0-6}\text{ alkylene-}$.

[0209] In some embodiments, L_{3a} is a bond. In some embodiments, L_{3a} is $\text{C}_{1-6}\text{ alkylene}$. In some embodiments, L_{3a} is $-\text{NH-C}_{1-6}\text{ alkylene-C(O)-}$. In some embodiments, L_{3a} is $-\text{N(CH}_3\text{)-C}_{1-6}\text{ alkylene C(O)-}$. In some embodiments, L_{3a} is $-\text{O-C}_{0-6}\text{ alkylene}$. In some embodiments, L_{3a} is $-(\text{CH}_2)_a-\text{NR}^a-(\text{CH}_2)_b-$. In some embodiments, L_{3a} is $-(\text{CH}_2)_a-\text{O}-(\text{CH}_2)_b-$. In some embodiments, L_{3a} is $-(\text{CH}_2)_a-\text{CH(NHR}^a\text{)-}$. In some embodiments, L_{3a} is $-(\text{CH}_2)_a-\text{CH(NHR}^a\text{)-}$. In some embodiments, L_{3a} is $-(\text{CR}^{1a}\text{R}^{1b})_a-$. In some embodiments, L_{3a} is $-(\text{CH}_2)_a-\text{CH(NR}^a\text{R}^b\text{)-}-(\text{CH}_2)_b-$.

[0210] In some embodiments, for Formula (A-1) to (A-4), at least one A is NH and at least one A is C(O). In some embodiments, for Formula (A-1) to (A-4), at least two A is NH and at least two A is C(O). In some embodiments, when M is a bicyclic ring, A is a bond. In some embodiments, at least one A is a phenylene optionally substituted with one or more alkyl. In some embodiments, at least one A is thiophenylene optionally substituted with one or more alkyl. In some embodiments, at least one A is a furanylene optionally substituted with one or more alkyl. In some embodiments, at least one A is $(\text{CH}_2)_{0-4}-\text{CH=CH-}(\text{CH}_2)_{0-4}$, preferably $-\text{CH=CH-}$. In some embodiments, at least one A is $-\text{NH-N=N-}$. In some embodiments, at least one A is $-\text{NH-C(O)-NH-}$. In some embodiments, at least one A is $-\text{N(CH}_3\text{)-C}_{1-6}\text{ alkylene}$. In some embodiments, at least one A is



In some embodiments, at least one A is $-\text{NH-C}_1\text{ alkylene-NH-}$. In some embodiments, at least one A is $-\text{O-C}_{1-6}\text{ alkylene-O-}$.


[0211] In some embodiments, each M in [A-M] of Formula (A-1) to (A-4) is $\text{C}_{6-10}\text{ arylene}$ group, 4-10 membered

heterocyclene, optionally substituted 5-10 membered heteroarylene group, or $\text{C}_{1-6}\text{ alkylene}$; each optionally substituted by 1-3 substituents selected from H, OH, halogen, $\text{C}_{1-10}\text{ alkyl}$, NO_2 , CN, NR^aR^b , $\text{C}_{1-6}\text{ haloalkyl}$, $-\text{C}_{1-6}\text{ alkoxy}$, $\text{C}_{1-6}\text{ haloalkoxy}$, $(\text{C}_{1-6}\text{ alkoxy})\text{C}_{1-6}\text{ alkyl}$, $\text{C}_{2-10}\text{ alkenyl}$, $\text{C}_{2-10}\text{ alkynyl}$, $\text{C}_{3-7}\text{ carbocyclyl}$, 4-10 membered heterocyclyl, $\text{C}_{6-10}\text{ aryl}$, 5-10 membered heteroaryl, $-(\text{C}_{3-7}\text{ carbocyclyl})\text{C}_{1-6}\text{ alkyl}$, (4-10 membered heterocyclyl) $\text{C}_{1-6}\text{ alkyl}$, $(\text{C}_{6-10}\text{ aryl})\text{C}_{1-6}\text{ alkyl}$, $(\text{C}_{6-10}\text{ aryl})\text{C}_{1-6}\text{ alkoxy}$, (5-10 membered heteroaryl) $\text{C}_{1-6}\text{ alkyl}$, $-(\text{C}_{3-7}\text{ carbocyclyl})\text{-amine}$, (4-10 membered heterocyclyl)amine, $(\text{C}_{6-10}\text{ aryl})\text{amine}$, (5-10 membered heteroaryl)amine, acyl, C-carboxy, O-carboxy, C-amido, N-amido, S-sulfonamido, N-sulfonamido, $-\text{SR}$, COOH, or CONR^aR^b ; wherein each R^a and R^b are independently H, $\text{C}_{1-10}\text{ alkyl}$, $\text{C}_{1-10}\text{ haloalkyl}$, $-\text{C}_{1-10}\text{ alkoxy}$. In some embodiments, each M in [A-M] of Formula (A-1) to (A-3) is a 5-10 membered heteroarylene containing at least one heteroatoms selected from O, S, and N or a $\text{C}_{1-6}\text{ alkylene}$, and the heteroarylene or the a $\text{C}_{1-6}\text{ alkylene}$ is optionally substituted with 1-3 substituents selected from OH, halogen, $\text{C}_{1-10}\text{ alkyl}$, NO_2 , CN, NR^aR^b , $\text{C}_{1-6}\text{ haloalkyl}$, $-\text{C}_{1-6}\text{ alkoxy}$, $\text{C}_{1-6}\text{ haloalkoxy}$, $\text{C}_{3-7}\text{ carbocyclyl}$, 4-10 membered heterocyclyl, $\text{C}_{6-10}\text{ aryl}$, 5-10 membered heteroaryl, $-\text{SR}$, COOH, or CONR^aR^b ; wherein each R^a and R^b are independently H, $\text{C}_{1-10}\text{ alkyl}$, $\text{C}_{1-10}\text{ haloalkyl}$, $-\text{C}_{1-10}\text{ alkoxy}$. In some embodiments, each R in [A-R] of Formula (A-1) to (A-3) is a 5-10 membered heteroarylene containing at least one heteroatoms selected from O, S, and N, and the heteroarylene is optionally substituted with 1-3 substituents selected from OH, $\text{C}_{1-6}\text{ alkyl}$, halogen, and $\text{C}_{1-6}\text{ alkoxy}$.

[0212] In some embodiments, for Formula (A-1) to (A-4), at least one M is a 5 membered heteroarylene having at least one heteroatom selected from O, N, S and optionally substituted with one or more $\text{C}_{1-10}\text{ alkyl}$. In some embodiments, at least one M is a pyrrole optionally substituted with one or more $\text{C}_{1-10}\text{ alkyl}$. In some embodiments, at least one M is a imidazole optionally substituted with one or more $\text{C}_{1-10}\text{ alkyl}$. In some embodiments, for Formula (A-1) to (A-4), at least one M is a $\text{C}_{2-6}\text{ alkylene}$ optionally substituted with one or more $\text{C}_{1-10}\text{ alkyl}$. In some embodiments, at least one M is a pyrrole optionally substituted with one or more $\text{C}_{1-10}\text{ alkyl}$. In some embodiments, for Formula (A-1) to (A-4), at least one M is a bicyclic heteroarylene or arylene. In some embodiments, at least one M is a phenylene optionally substituted with one or more $\text{C}_{1-10}\text{ alkyl}$. In some embodiments, at least one M is a benzimidazole optionally substituted with one or more $\text{C}_{1-10}\text{ alkyl}$.


[0213] In some embodiments, the first terminus comprises a structure of Formula (A-4):

(A-4)




[0214] wherein:

[0215] L_{1c} is a bivalent or trivalent group selected from



a C_{1-10} alkylene, $-\text{NH}-C_{0-6}$ alkylene- $\text{C}(\text{O})-$, $-\text{N}(\text{CH}_3)-C_{0-6}$ alkylene, and




[0216] p is an integer in the range of 3 to 10;

[0217] $2 \leq q \leq (p-1)$;

[0218] $2 \leq r \leq (p-1)$;

[0219] m and n are each independently an integer in the range of 0 to 10;

[0220] each A^2 through A^p is independently selected from the group consisting of a bond, C_{1-10} alkylene, optionally substituted C_{6-10} arylene group, optionally substituted 4-10 membered heterocyclene, optionally substituted 5-10 membered heteroarylene group, $-\text{C}_{1-10}$ alkylene- $\text{C}(\text{O})-$, $-\text{C}_{1-10}$ alkylene- NR^a- , $-\text{CO}-$, $-\text{NR}^a-$, $-\text{CONR}^a-$, $-\text{CONR}^a\text{C}_{1-4}$ alkylene-, $-\text{NR}^a\text{CO}-\text{C}_{1-4}$ alkylene-, $-\text{C}(\text{O})\text{O}-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O})-$, $-\text{S}(\text{O})_2-$, $-\text{C}(=\text{S})-\text{NH}-$, $-\text{C}(\text{O})-\text{NH}-\text{NH}-$, $-\text{C}(\text{O})-\text{N}=\text{N}-$, $-\text{C}(\text{O})-\text{CH}=\text{CH}-$, $(\text{CH}_2)_{0-4}-\text{CH}=\text{CH}-(\text{CH}_2)_{0-4}$, $-\text{N}(\text{CH}_3)-C_{1-6}$ alkylene,




$-\text{NH}-C_{1-6}$ alkylene- $\text{NH}-$, $-\text{O}-C_{1-6}$ alkylene- $\text{O}-$, $-\text{NH}-\text{N}=\text{N}-$, $-\text{NH}-\text{C}(\text{O})-\text{NH}-$, and any combinations thereof, and at least one A^2 through A^p is NHCO ;

[0221] each M^1 through M^p is an optionally substituted C_{6-10} arylene group, optionally substituted 4-10 membered heterocyclene, optionally substituted 5-10 membered heteroarylene group, or an optionally substituted alkylene;

[0222] each T^2 through T^p is independently selected from the group consisting of a bond, C_{1-10} alkylene, optionally substituted C_{6-10} arylene group, optionally

substituted 4-10 membered heterocyclene, optionally substituted 5-10 membered heteroarylene group, $-\text{C}_{1-10}$ alkylene- $\text{C}(\text{O})-$, $-\text{C}_{1-10}$ alkylene- NR^a- , $-\text{CO}-$, $-\text{NR}^a-$, $-\text{CONR}^a-$, $-\text{CONR}^a\text{C}_{1-4}$ alkylene-, $-\text{NR}^a\text{CO}-\text{C}_{1-4}$ alkylene-, $-\text{C}(\text{O})\text{O}-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O})-$, $-\text{S}(\text{O})_2-$, $-\text{C}(=\text{S})-\text{NH}-$, $-\text{C}(\text{O})-\text{NH}-\text{NH}-$, $-\text{C}(\text{O})-\text{N}=\text{N}-$, $-\text{C}(\text{O})-\text{CH}=\text{CH}-$, $(\text{CH}_2)_{0-4}-\text{CH}=\text{CH}-(\text{CH}_2)_{0-4}$, $-\text{N}(\text{CH}_3)-C_{1-6}$ alkylene,



$-\text{NH}-C_{1-6}$ alkylene- $\text{NH}-$, $-\text{O}-C_{1-6}$ alkylene- $\text{O}-$, $-\text{NH}-\text{N}=\text{N}-$, and $-\text{NH}-\text{C}(\text{O})-\text{NH}-$, and any combinations thereof;

[0223] each Q^1 to Q^p is an optionally substituted C_{6-10} arylene group, optionally substituted 4-10 membered heterocyclene, optionally substituted 5-10 membered heteroarylene group, or an optionally substituted alkylene;

[0224] each A^1 , A^2 , E_1 , and E_2 are independently H or $-\text{A}^E-\text{G}$;

[0225] each A^E is independently absent or NHCO ;

[0226] each G is independently selected from the group consisting of optionally substituted H, C_{6-10} aryl, optionally substituted 4-10 membered heterocyclyl, optionally substituted 5-10 membered heteroaryl, an optionally substituted C_{1-6} alkyl, C_{0-4} alkylene- $\text{NHC}(=\text{NH})\text{NH}$, $-\text{CN}$, $-\text{C}_{0-4}$ alkylene- $\text{C}(=\text{NH})(\text{NR}^a\text{R}^b)$, $-\text{C}_{0-4}$ alkylene- $\text{C}(=\text{N}^+\text{H}_2)(\text{NR}^a\text{R}^b)\text{C}_{1-5}$ alkylene- NR^aR^b , C_{0-4} alkylene- $\text{NHC}(=\text{NH})\text{R}^a$, and optionally substituted amine;

[0227] when L_{1c} is a trivalent group, the oligomeric backbone is attached to the first terminus through L_{1c} , and each G is an end group independently selected from the group consisting of optionally substituted C_{6-10} aryl, optionally substituted 4-10 membered heterocyclyl, optionally substituted 5-10 membered heteroaryl, an optionally substituted C_{1-6} alkyl, C_{0-4} alkylene- $\text{NHC}(=\text{NH})\text{NH}$, $-\text{CN}$, $-\text{C}_{0-4}$ alkylene- $\text{C}(=\text{NH})(\text{NR}^a\text{R}^b)$, $-\text{C}_{0-4}$ alkylene- $\text{C}(=\text{N}^+\text{H}_2)(\text{NR}^a\text{R}^b)\text{C}_{1-5}$ alkylene- NR^aR^b , C_{0-4} alkylene- $\text{NHC}(=\text{NH})\text{R}^a$, $-\text{CO}$ -halogen, and optionally substituted amine;

[0228] when L_{1c} is a divalent group, the oligomeric backbone is attached to the first terminus through one of A^1 , T^1 , E_1 , and E_2 , and each G is independently selected from the group consisting of a bond, a $-\text{C}_{1-6}$ alkylene-, $-\text{NH}-C_{0-6}$ alkylene- $\text{C}(\text{O})-$, $-\text{N}(\text{CH}_3)-C_{0-6}$ alkylene-, $-\text{C}(\text{O})-$, $-\text{C}(\text{O})-C_{1-6}$ alkylene, and $-\text{O}-C_{0-6}$ alkylene, optionally substituted C_{6-10} aryl, optionally substituted 4-10 membered heterocyclyl, optionally substituted 5-10 membered heteroaryl, an optionally substituted C_{1-6} alkyl, C_{0-4} alkylene- $\text{NHC}(=\text{NH})\text{NH}$, $-\text{CN}$, $-\text{C}_{0-4}$ alkylene- $\text{C}(=\text{NH})(\text{NR}^a\text{R}^b)$, $-\text{C}_{0-4}$ alkylene- $\text{C}(=\text{N}^+\text{H}_2)(\text{NR}^a\text{R}^b)\text{C}_{1-5}$ alkylene- NR^aR^b , C_{0-4} alkylene- $\text{NHC}(=\text{NH})\text{R}^a$, and optionally substituted amine; or





[0229] when L_{1c} is a bivalent group, the oligomeric backbone is attached to the first terminus through a nitrogen or carbon atom on one of M^1 , M^2 , $\dots$ M^{p-1} , M^p , T^1 , T^2 , $\dots$ T^{p-1} , and T^p , and each G is an end group independently selected from the group consisting of optionally substituted C_{6-10} aryl, optionally substituted

4-10 membered heterocyclyl, optionally substituted 5-10 membered heteroaryl, an optionally substituted C_{1-6} alkyl, C_{0-4} alkylene-NHC(=NH)NH, —CN, — C_{0-4} alkylene-C(=NH)(NR^aR^b), — C_{0-4} alkylene-C(=N⁺H₂)(NR^aR^b) C_{1-5} alkylene-NR^aR^b, C_{0-4} alkylene-NHC(=NH)R^a, and optionally substituted, and

[0230] each R^a and R^b are independently H, an optionally substituted C_{1-6} alkyl, an optionally substituted C_{3-10} cycloalkyl, optionally substituted C_{6-10} aryl, optionally substituted 4-10 membered heterocyclyl, or an optionally substituted 5-10 membered heteroaryl;

[0231] each R^{1a} and R^{1b} are independently H or an optionally substituted C_{1-6} alkyl.

[0232] In some embodiments, the first terminus comprises a structure of Formula (A-4a) or (A-4b):




Formula (A-4a)


Formula (A-4b)

[0233] wherein:

[0234] L_{1c} is a bivalent or trivalent group selected from



a C_{1-10} alkylene, —NH— C_{0-6} alkylene-C(O)—, —N(CH₃)— C_{0-6} alkylene, and



[0235] p is an integer in the range of 2 to 10;


[0236] p' is an integer in the range of 2 to 10;

[0237] $2 \leq q \leq (p-1)$;

[0238] $2 \leq r \leq (p-1)$;

[0239] m and n are each independently an integer in the range of 0 to 10;


[0240] each A² through A^p is independently selected from the group consisting of a bond, C_{1-10} alkylene, optionally substituted C_{6-10} arylene group, optionally substituted 4-10 membered heterocyclene, optionally substituted 5-10 membered heteroarylene group, — C_{1-10} alkylene-C(O)—, — C_{1-10} alkylene-NR^a—, —CO—, —NR^a—, —CONR^a—, —CONR^a C_{1-4} alkylene—, —NR^aCO— C_{1-4} alkylene—, —C(O)O—, —O—, —S—, —S(O)—, —S(O)₂—, —C(=S)—NH—, —C(O)—NH—NH—, —C(O)—N=N—, —C(O)—CH=CH—, (CH₂)₀₋₄—CH=CH—(CH₂)₀₋₄—, —N(CH₃)— C_{1-6} alkylene, and



—NH— C_{1-6} alkylene-NH—, —O— C_{1-6} alkylene-O—, —NH—N=N—, —NH—C(O)—NH—, and any combinations thereof, and at least one of A² through A^p is —CONH—;

[0241] each M¹ through M^p is an optionally substituted C_{6-10} arylene group, optionally substituted 4-10 membered heterocyclene, optionally substituted 5-10 membered heteroarylene group, or an optionally substituted alkylene;

[0242] each T² through T^{p'} in formula (A-4a) is independently selected from the group consisting of a bond, C_{1-10} alkylene, optionally substituted C_{6-10} arylene group, optionally substituted 4-10 membered heterocyclene, optionally substituted 5-10 membered heteroarylene group, — C_{1-10} alkylene-C(O)—, — C_{1-10} alkylene-NR^a—, —CO—, —NR^a—, —CONR^a—, —CONR^a C_{1-4} alkylene—, —NR^aCO— C_{1-4} alkylene—, —C(O)O—, —O—, —S—, —, —S(O)—, —S(O)₂—, —C(=S)—NH—, —C(O)—NH—NH—, —C(O)—N=N—, —C(O)—CH=CH—, (CH₂)₀₋₄—CH=CH—(CH₂)₀₋₄—, —N(CH₃)— C_{1-6} alkylene, and



—NH—C₁₋₆ alkylene—NH—, —O—C₁₋₆ alkylene—O—, —NH—N=N—, —NH—C(O)—NH—, and any combinations thereof, and at least one of T² through T^p is —CONH—;

[0243] each Q¹ to Q^{p'} is an optionally substituted C₆₋₁₀ arylene group, optionally substituted 4-10 membered heterocyclene, optionally substituted 5-10 membered heteroarylene group, or an optionally substituted alkylene;

[0244] each A¹, T¹, E₁, and E₂ are independently H or —A^E—G,

[0245] each A^E is independently absent or NHCO,


[0246] each G is independently selected from the group consisting of optionally substituted H, C₆₋₁₀ aryl, optionally substituted 4-10 membered heterocyclyl, optionally substituted 5-10 membered heteroaryl, an optionally substituted C₁₋₆ alkyl, C₀₋₄ alkylene-NHC(=NH)NH, —CN, —C₀₋₄alkylene-C(=NH)(NR^aR^b), —C₀₋₄alkylene-C(=N⁺H₂)(NR^aR^b)C₁₋₅alkylene-NR^aR^b, C₀₋₄ alkylene-NHC(=NH) R^a, and optionally substituted amine;

[0247] when L_{1c} is a trivalent group, the oligomeric backbone is attached to the first terminus through L_{1c}, when L_{1c} is a bivalent group, the oligomeric backbone is attached to the first terminus through one of A¹, T¹, E₁, and E₂, or the oligomeric backbone is attached to the first terminus through a nitrogen or carbon atom on one of M¹, M², . . . M^{p-1}, M^p, T¹, T², . . . T^{p-1}, and T^p and


[0248] each R^a and R^b are independently H, an optionally substituted C₁₋₆ alkyl, an optionally substituted C₃₋₁₀ cycloalkyl, optionally substituted C₆₋₁₀ aryl, optionally substituted 4-10 membered heterocyclyl, or an optionally substituted 5-10 membered heteroaryl; and

[0249] each R^{1a} and R^{1b} are independently H or an optionally substituted C₁₋₆ alkyl.


[0250] In certain embodiments, L_{1c} is



C₁₋₁₀ alkylene, or



In certain embodiments, L_{1c} is C₃₋₈ alkylene. In certain embodiments, L_{1c} is



and wherein 2 ≤ m + n ≤ 10. In some embodiments, L_{1c} is C₂₋₈ alkylene. In some embodiments, L_{1c} is C₃₋₈ alkylene. In some embodiments, L_{1c} is C₄₋₈ alkylene. In some embodiments, L_{1c} is C₃ alkylene, C₄ alkylene, C₅ alkylene, C₆ alkylene, C₇ alkylene, C₈ alkylene, or C₉ alkylene.

[0251] In certain embodiments, 3 ≤ m + n ≤ 7. In certain embodiments, (m + n) is 3, 4, 5, 6, 7, 8, or 9. In certain embodiments, m is in the range of 3 to 8. In certain embodiments, m is 3, 4, 5, 6, 7, 8, or 9.

[0252] In certain embodiments, M^q is a five to 10 membered heteroaryl ring comprising at least one nitrogen; Q^q is a five to 10 membered heteroaryl ring comprising at least one nitrogen; and M^q is linked to Q^q through L_{1c}. In certain embodiments, M^q is a five membered heteroaryl ring comprising at least one nitrogen; Q^q is a five membered heteroaryl ring comprising at least one nitrogen; M^q is linked to Q^q through L_{1c}, and L_{1c} is attached to the nitrogen atom on M^q and L_{1c} is attached to the nitrogen atom on Q^q.

[0253] In certain embodiments, each M¹ through M^p is independently selected from an optionally substituted pyrrolylene, an optionally substituted imidazolylene, an optionally substituted pyrazolylene, an optionally substituted thiazolylene, an optionally substituted diazolylene, an optionally substituted benzopyridazinylene, an optionally substituted benzopyrazinylene, an optionally substituted phenylene, an optionally substituted pyridinylene, an optionally substituted thiophenylene, an optionally substituted furanylene, an optionally substituted piperidinylene, an optionally substituted pyrimidinylene, an optionally substituted anthracenylenylene, an optionally substituted quinolinylene, and an optionally substituted C₁₋₆ alkylene.


[0254] In certain embodiments, at least one M of M¹ through M^p is a 5 membered heteroarylene having at least one heteroatom selected from O, N, S and optionally substituted with one or more C₁₋₁₀ alkyl. In certain embodiments, at least two M of M¹ through M^p is a 5 membered heteroarylene having at least one heteroatom selected from O, N, S and optionally substituted with one or more C₁₋₁₀ alkyl. In some embodiments, at least three, four, five, or six M of M¹ through M^p is a 5 membered heteroarylene having at least one heteroatom selected from O, N, S and optionally substituted with one or more C₁₋₁₀ alkyl. In some embodiments, at least one of M¹ through M^p is a pyrrole optionally substituted with one or more C₁₋₁₀ alkyl. In some embodiments, at least one of M¹ through M^p is an imidazole optionally substituted with one or more C₁₋₁₀ alkyl. In some embodiments, at least one of M¹ through M^p is a C₂₋₆ alkylene optionally substituted with one or more C₁₋₁₀ alkyl. In some embodiments, at least one of M¹ through M^p is a phenyl optionally substituted with one or more C₁₋₁₀ alkyl. In some embodiments, at least one of M¹ through M^p is a bicyclic heteroarylene or arylene. In some embodiments, at least one of M¹ through M^p is a phenylene optionally substituted with one or more C₁₋₁₀ alkyl. In some embodiments, at least one of M¹ through M^p is a benzimidazole optionally substituted with one or more C₁₋₁₀ alkyl.

[0255] In certain embodiments, each Q¹ to Q^p is independently selected from an optionally substituted pyrrolylene,

an optionally substituted imidazolylen, an optionally substituted pyrazolylen, an optionally substituted thiazolylen, an optionally substituted diazolylen, an optionally substituted benzopyridazinylen, an optionally substituted benzopyrazinylen, an optionally substituted phenylene, an optionally substituted pyridinylen, an optionally substituted thiophenylen, an optionally substituted furanylen, an optionally substituted piperidinylen, an optionally substituted pyrimidinylen, an optionally substituted anthracenylen, an optionally substituted quinolinylene, and an optionally substituted C_{1-6} alkylene.


[0256] In certain embodiments, at least one Q of Q^1 through Q^p is a 5 membered heteroarylene having at least one heteroatom selected from O, N, S and optionally substituted with one or more C_{1-10} alkyl. In certain embodiments, at least two Q of Q^1 through Q^p is a 5 membered heteroarylene having at least one heteroatom selected from O, N, S and optionally substituted with one or more C_{1-10} alkyl. In certain embodiments, at least three, four, five, or six Q of Q^1 through Q^p is a 5 membered heteroarylene having at least one heteroatom selected from O, N, S and optionally substituted with one or more C_{1-10} alkyl. In some embodiments, at least one of Q^1 through Q^p is a pyrrole optionally substituted with one or more C_{1-10} alkyl. In some embodiments, at least one of Q^1 through Q^p is a imidazole optionally substituted with one or more C_{1-10} alkyl. In some embodiments, at least one of Q^1 through Q^p is a C_{2-6} alkylene optionally substituted with one or more C_{1-10} alkyl. In some embodiments, at least one of Q^1 through Q^p is a phenyl optionally substituted with one or more C_{1-10} alkyl. In some embodiments, at least one of Q^1 through Q^p is a bicyclic heteroarylene or arylen. In some embodiments, at least one of Q^1 through Q^p is a phenylene optionally substituted with one or more C_{1-10} alkyl. In some embodiments, at least one of Q^1 through Q^p is a benzimidazole optionally substituted with one or more C_{1-10} alkyl.

[0257] In some embodiments, at least one of A^2 through A^p is NH and at least one of A^2 through A^p is C(O). In some embodiments, at least two of A^2 through A^p is NH and at least two of A^2 through A^p is C(O). In some embodiments, when one of M^2 through M^p is a bicyclic ring, the adjacent A is a bond. In some embodiments, one of A^2 through A^p is a phenylene optionally substituted with one or more alkyl. In some embodiments, one of A^2 through A^p is thiophenylene optionally substituted with one or more alkyl. In some embodiments, one of A^2 through A^p is a furanylene optionally substituted with one or more alkyl. In some embodiments, one of A^2 through A^p is $(CH_2)_{0-4}-CH=CH-(CH_2)_{0-4}$, preferably $-CH=CH-$. In some embodiments, one of A^2 through A^p is $-NH-N=N-$. In some embodiments, one of A^2 through A^p is $-NH-C(O)-NH-$. In some embodiments, one of A^2 through A^p is $-N(CH_3)-C_{1-6}$ alkylene. In some embodiments, one of A^2 through A^p is




In some embodiments, one of A^2 through A^p is $-NH-C_{1-6}$ alkylene-NH—. In some embodiments, one of A^2 through A^p is $-O-C_{1-6}$ alkylene-O—.

[0258] In certain embodiments, each A^2 through A^p is independently selected from a bond, C_{1-10} alkylene, optionally substituted phenylene, optionally substituted thiophenylene, optionally substituted furanylene, $-C_{1-10}$ alkylene-C(O)—, $-C_{1-10}$ alkylene-NH—, $-CO-$, $-NR^a-$, $-CONR^a-$, $-CON^aC_{1-4}$ alkylene-, $-NR^aCO-C_{1-4}$ alkylene-, $-C(O)O-$, $-O-$, $-S-$, $-S(O)-$, $-S(O)_2-$, $-C(=S)-NH-$, $-C(O)-NH-NH-$, $-C(O)-N=N-$, $-C(O)-CH=CH-$, $-CH=CH-$, $-NH-N=N-$, $-NH-C(O)-NH-$, $-N(CH_3)-C_{1-6}$ alkylene,




$-NH-C_{1-6}$ alkylene-NH—, and $-O-C_{1-6}$ alkylene-O—, and any combinations thereof.

[0259] In some embodiments, at least one T of T^2 through P is NH and at least one of T of T^2 through P is C(O). In some embodiments, at least two T of T^2 through P is NH and at least two T of T^2 through P is C(O). In some embodiments, when one Q of Q^2 through Q^p is a bicyclic ring, the adjacent T is a bond. In some embodiments, one T of T^2 through T^p is a phenylene optionally substituted with one or more alkyl. In some embodiments, one T of T^2 through T^p is thiophenylene optionally substituted with one or more alkyl. In some embodiments, one T of T^2 through T^p is a furanylene optionally substituted with one or more alkyl. In some embodiments, one T of T^2 through T^p is $(CH_2)_{0-4}-CH=CH-(CH_2)_{0-4}$, preferably $-CH=CH-$. In some embodiments, one T of T^2 through T^p is $-NH-N=N-$. In some embodiments, one T of T^2 through T^p is $-NH-C(O)-NH-$. In some embodiments, one T of T^2 through T^p is $-N(CH_3)-C_{1-6}$ alkylene. In some embodiments, one T of T^2 through T^p is



In some embodiments, one T of T^2 through T^p is $-NH-C_{1-6}$ alkylene-NH—. In some embodiments, one T of T^2 through T^p is $-O-C_{1-6}$ alkylene-O—.

[0260] In certain embodiments, each T^2 through T^p is independently selected from a bond, C_{1-10} alkylene, optionally substituted phenylene, optionally substituted thiophenylene, optionally substituted furanylene, $-C_{1-10}$ alkylene-C(O)—, $-C_{1-10}$ alkylene-NH—, $-CO-$, $-NR^a-$, $-CONR^a-$, $-CON^aC_{1-4}$ alkylene-, $-NR^aCO-C_{1-4}$ alkylene-, $-C(O)O-$, $-O-$, $-S-$, $-C(=S)-NH-$, $-C(O)-NH-NH-$, $-C(O)-N=N-$, $-C(O)-CH=CH-$, $-CH=CH-$, $-NH-N=N-$, $-NH-C(O)-NH-$, $-N(CH_3)-C_{1-6}$ alkylene, and




—NH—C₁₋₆ alkylene-NH—, —O—C₁₋₆ alkylene-O—, and any combinations thereof.


[0261] In certain embodiments, each A^1 , T^1 , E_1 , and E_2 are independently $-A^E-G$, and each A^E is independently absent or NHCO. In certain embodiments, each A^1 , T^1 , E_1 , and E_2 are independently $-A^E-G$ and each A^E is independently NHCO.

[0262] In certain embodiments, for Formula (A-1) to (A-4), each end group G independently comprises a moiety selected from the group consisting of optionally substituted C₆₋₁₀ aryl, optionally substituted 4-10 membered heterocyclyl, a 5-10 membered heteroaryl optionally substituted with 1-3 substituents selected from C₁₋₆ alkyl, —NHCOH, halogen, —NR^aR^b, an optionally substituted C₁₋₆ alkyl, C₀₋₄ alkylene-NHC(=NH)NH, C₀₋₄ alkylene-NHC(=NH)—R_E, —C₁₋₄ alkylene-R_E, —CN, —C₀₋₄alkylene-C(=NH)(NR^aR^b), —C₀₋₄alkylene-C(=N⁺H₂)(NR^aR^b)C₁₋₅ alkylene-NR^aR^b, C₀₋₄ alkylene-NHC(=NH) R^a, and optionally substituted amine, wherein each R^a and R^b are independently H, an optionally substituted C₁₋₆ alkyl, an optionally substituted C₃₋₁₀ cycloalkyl, optionally substituted C₆₋₁₀ aryl, optionally substituted 4-10 membered heterocyclyl, or an optionally substituted 5-10 membered heteroaryl. In certain embodiments, for Formula (A-1) to (A-4), each end group G independently comprises a NH or CO group. In certain embodiments, each R^a and R^b are independently H or C₁₋₆ alkyl. In certain embodiments, for formula (A-1) to (A-4), at least one of the end groups is H. In certain embodiments, for Formula (A-1) to (A-4), at least two of the end groups are H. In certain embodiments, for formula (A-1) to (A-4), at least one of the end groups is H. In certain embodiments, for Formula (A-1) to (A-4), at least one of the end groups is —NH—5-10 membered heteroaryl ring optionally substituted with one or more alkyl or —CO—5-10 membered heteroaryl ring optionally substituted with one or more alkyl.

[0263] In certain embodiments, for Formula (A-1) to (A-4), each end group G is independently selected from



-continued



[0264] In certain embodiments, for Formula (A-1) to (A-4), each E_1 independently comprises an optionally substituted thiophene-containing moiety, optionally substituted pyrrole containing moiety, optionally substituted imidazole containing moiety, or optionally substituted amine.

[0265] In certain embodiments, for Formula (A-1) to (A-4), each E_2 independently comprises an optionally substituted thiophene-containing moiety, optionally substituted pyrrole containing moiety, optionally substituted imidazole containing moiety, or optionally substituted amine.

[0266] In certain embodiments, for Formula (A-1) to (A-4), each E_1 and E_2 independently comprises a moiety selected from the group consisting of optionally substituted N-methylpyrrole, optionally substituted N-methylimidazole, optionally substituted benzimidazole moiety, and optionally substituted 3-(dimethylamino)propanamidy. In certain embodiments, each E_1 and E_2 independently comprises thiophene, benzothiophene, C—C linked benzimidazole/thiophene-containing moiety, or C—C linked hydroxybenzimidazole/thiophene-containing moiety. In certain


embodiments, for Formula (A-1) to (A-4), each E_1 and E_2 independently also comprises NH or CO group.

[0267] In certain embodiments, for Formula (A-1) to (A-4), each E_1 or E_2 independently comprises a moiety selected from the group consisting of isophthalic acid; phthalic acid; terephthalic acid; morpholine; N,N-dimethylbenzamide; N,N-bis(trifluoromethyl)benzamide; fluorobenzene; (trifluoromethyl)benzene; nitrobenzene; phenyl acetate; phenyl 2,2,2-trifluoroacetate; phenyl dihydrogen phosphate; 2H-pyran; 2H-thiopyran; benzoic acid; isonicotinic acid; and nicotinic acid; wherein one, two, or three ring members in any of the end-group candidates can be independently substituted with C, N, S or O; and where any one, two, three, four or five of the hydrogens bound to the ring can be substituted with R^{3a} , wherein R_5 may be independently selected from H, OH, halogen, C_{1-10} alkyl, NO_2 , NH_2 , C_{1-10} haloalkyl, $-OC_{1-10}$ haloalkyl, $COOH$, and $CONR^{1c}R^{1d}$, wherein each R^{1c} and R^{1d} are independently H, C_{1-10} alkyl, C_{1-10} haloalkyl, or $-C_{1-10}$ alkoxy.

[0268] In some embodiments, the first terminus comprises the structure of Formula (A-5a) or Formula (A-5b):



or



[0269] wherein:

[0270] each Q^1 , Q^2 , $Q^3 \dots$ through Q^p are independently an optionally substituted C_{6-10} arylene group, optionally substituted 4-10 membered heterocyclene, optionally substituted 5-10 membered heteroarylene group, or an optionally substituted alkylene;

[0271] each A^{1a} and T^{1a} are independently a bond, H, a $-C_{1-6}$ alkylene-, $-NH-C_{0-6}$ alkylene- $C(O)-$, $-N(CH_3)-C_{0-6}$ alkylene-, $-C(O)-$, $-C(O)-C_{1-10}$ alkylene, and $-O-C_{0-6}$ alkylene, optionally substituted C_{6-10} aryl, optionally substituted 4-10 membered heterocyclyl, optionally substituted 5-10 membered heteroaryl, an optionally substituted C_{1-6} alkyl, C_{0-4} alkylene-NHC(=NH)NH, $-CN$, $-C_{0-4}$ alkylene- $C(=NH)(NR^aR^b)$, $-C_{0-4}$ alkylene- $C(=N^+H_2)(NR^aR^b)$, C_{1-5} alkylene- NR^aR^b , C_{0-4} alkylene-NHC(=NH) R^a , and optionally substituted amine;


[0272] p is an integer between 2 and 10; and

[0273] G is selected from the group consisting of optionally substituted C_{6-10} aryl, optionally substituted 4-10 membered heterocyclyl, optionally substituted 5-10 membered heteroaryl, or an optionally substituted alkyl, C_{0-4} alkylene-NHC(=NH)NH, $-CN$, $-C_{0-4}$ alkylene- $C(=NH)(NR^aR^b)$, $-C_{0-4}$ alkylene- $C(=N^+H_2)(NR^aR^b)$, C_{1-5} alkylene- NR^aR^b , C_{0-4} alkylene-NHC(=NH) R^a , and optionally substituted amine;

[0274] each R^a and R^b are independently H, an optionally substituted C_{1-6} alkyl, an optionally substituted C_{3-10} cycloalkyl, optionally substituted C_{6-10} aryl, optionally substituted 4-10 membered heterocyclyl, or an optionally substituted 5-10 membered heteroaryl; and

[0275] wherein the first terminus is connected to the oligomeric backbone through either A¹ or T¹, or a nitrogen or carbon atom on one of Q¹ through Q^p.

[0276] In certain embodiments, the first terminus comprises the structure of Formula (A-5c):



[0277] wherein:

[0278] each Q_a^1 , Q_a^2 . . . Q_a^q . . . through Q_a^p are independently an optionally substituted C₆₋₁₀ arylene group, optionally substituted 4-10 membered heterocycle, optionally substituted 5-10 membered heteroarylene group, or an optionally substituted alkylene;




[0279] each Q_b^1 , Q_b^2 . . . Q_b^r . . . through Q_b^p are independently an optionally substituted C₆₋₁₀ arylene group, optionally substituted 4-10 membered heterocyclene, optionally substituted 5-10 membered heteroarylene group, or an optionally substituted alkylene;

[0280] p is an integer between 3 and 10;


$$2 \leq q \leq (p-1);$$

$$2 \leq r \leq (p-1);$$

[0281] L_a is selected from a divalent or trivalent group
selected from the group consisting of



[0282] a C₁₋₁₀ alkylene, —NH—C₀₋₆ alkylene-C(O)—, —N(CH₃)—C₀₋₆ alkylene, and



[0283] each m and n are independently an integer in the range of 1 to 10;

[0284] n is an integer in the range of 1 to 10;

[0285] each R^{1a} and R^{1b} are independently H, or C_{1-6} alkyl;


[0286] when L_a is a trivalent group, the oligomeric backbone is attached to the first terminus through L_a , and each W_a^1 , G_a , G_b , and W_b^1 are end groups independently selected from the group consisting of optionally substituted C_{6-10} aryl, optionally substituted 4-10 membered heterocyclyl, optionally substituted 5-10 membered heteroaryl, an optionally substituted C_{1-6} alkyl, C_{0-4} alkylene-NHC(=NH)NH, —CN, — C_{0-4} alkylene-C(=NH)(NR^aR^b), — C_{0-4} alkylene-C(=N⁺H₂)(NR^aR^b)C₁₋₅alkylene-NR^aR^b, C_{0-4} alkylene-NHC(=NH) R^a, and optionally substituted amine;

[0287] when L_a is a divalent group, the oligomeric backbone is attached to the first terminus through one of W_a^1 , G_a , G_b , and W_b^1 , and each W_a^1 , G_a , G_b , and W_b^1 are independently selected from the group consisting of a bond, a $-C_{1-6}$ alkylene-, $-NH-C_{0-6}$ alkylene-C(O)-, $-N(CH_3)-C_{0-6}$ alkylene-, $-C(O)-$, $-C(O)-C_{1-10}$ alkylene, and $-O-C_{0-6}$ alkylene, optionally substituted C_{6-10} aryl, optionally substituted 4-10 membered heterocycl, optionally substituted 5-10 membered heteroaryl, an optionally substituted C_{1-6} alkyl, C_{0-4} alkylene-NHC(=NH)NH, $-CN$, $-C_{0-4}$ alkylene-C(=NH)(NR^aR^b), $-C_{0-4}$ alkylene-C(=N⁺H₂)(NR^aR^b) $C_{1-5alkylene}-NR^aR^b$, C_{0-4} alkylene-NHC(=NH) R^a, and optionally substituted amine; or

[0288] when L_a is a bivalent group, the oligomeric backbone is attached to the first terminus through a nitrogen or carbon atom on one of $Q_a^1, Q_{a2}, \dots, Q_{ap}^1, Q_{ap}, Q_b^1, Q_{a2}, \dots, Q_{bp-1},$ and Q_b^p , and each $W_a^1, G_a, G_b,$ and W_b^1 are end groups independently selected from the group consisting of optionally substituted C_{6-10} aryl, optionally substituted 4-10 membered heterocyclyl, optionally substituted 5-10 membered heteroaryl, an optionally substituted C_{1-6} alkyl, C_{0-4} alkylene-NHC(=NH)NH, —CN, — C_{0-4} alkylene-C(=NH)(NR^aR^b), — C_{0-4} alkylene-C(=N⁺H₂)(NR^aR^b), C_{1-5} alkylene-NR^aR^b, C_{0-4} alkylene-NHC(=NH) R^a, and optionally substituted amine, and

[0289] each R^a and R^b are independently H, an optionally substituted C_{1-6} alkyl, an optionally substituted C_{3-10} cycloalkyl, optionally substituted C_{6-10} aryl, optionally substituted 4-10 membered heterocyclyl, or an optionally substituted 5-10 membered heteroaryl.

[0290] In some embodiments, the first terminus comprises the structure of Formula (A-5c) or (A-5d):



[0304] In certain embodiments, Q_a^q is a five to 10 membered heteroaryl ring comprising at least one nitrogen; $Q_b^{q'}$ is a five to 10 membered heteroaryl ring comprising at least one nitrogen; and Q_a^q is linked to $Q_b^{q'}$ through L_a . In certain embodiments, Q_a^q is a five membered heteroaryl ring comprising at least one nitrogen; $Q_b^{q'}$ is a five membered heteroaryl ring comprising at least one nitrogen; Q_a^q is linked to $Q_b^{q'}$ through L_a , and L_a is attached to the nitrogen atom on Q_a^q and L_{1c} is attached to the nitrogen atom on $Q_b^{q'}$.

[0305] In certain embodiments, each Q_a^1 through Q_a^p is independently selected from an optionally substituted pyrrolylene, an optionally substituted imidazolylene, an optionally substituted pyrazolylene, an optionally substituted thiazolylene, an optionally substituted diazolylene, an optionally substituted benzopyridazinylene, an optionally substituted benzopyrazinylene, an optionally substituted phenylene, an optionally substituted pyridinylene, an optionally substituted thiophenylene, an optionally substituted furanylene, an optionally substituted piperidinylene, an optionally substituted pyrimidinylene, an optionally substituted anthracenylenylene, an optionally substituted quinolinylene, and an optionally substituted C_{1-6} alkylene.

[0306] In certain embodiments, at least one Q of Q_a^1 through Q_a^p is a 5 membered heteroarylene having at least one heteroatom selected from O, N, S and optionally substituted with one or more C_{1-10} alkyl. In certain embodiments, at least two Q of Q_a^1 through Q_a^p is a 5 membered heteroarylene having at least one heteroatom selected from O, N, S and optionally substituted with one or more C_{1-10} alkyl. In certain embodiments, at least three, four, five, or six Q of Q_a^1 through Q_a^p is a 5 membered heteroarylene having at least one heteroatom selected from O, N, S and optionally substituted with one or more C_{1-10} alkyl. In some embodiments, at least one Q of Q_a^1 through Q_a^p is a pyrrole optionally substituted with one or more C_{1-10} alkyl. In some embodiments, at least one of Q of Q_a^1 through Q_a^p is a imidazole optionally substituted with one or more C_{1-10} alkyl. In some embodiments, at least one Q of Q_a^1 through Q_a^p is a C_{2-6} alkylene optionally substituted with one or more C_{1-10} alkyl. In some embodiments, at least one Q of Q_a^1 through Q_a^p is a phenyl optionally substituted with one or more C_{1-10} alkyl. In some embodiments, at least one Q of Q_a^1 through Q_a^p is a bicyclic heteroarylene or arylenylene. In some embodiments, at least one Q of Q_a^1 through Q_a^p is a phenylene optionally substituted with one or more C_{1-10} alkyl. In some embodiments, at least one Q of Q_a^1 through Q_a^p is a benzimidazole optionally substituted with one or more C_{1-10} alkyl.

[0307] In certain embodiments, each Q_b^1 through Q_b^p is independently selected from an optionally substituted pyrrolylene, an optionally substituted imidazolylene, an optionally substituted pyrazolylene, an optionally substituted thiazolylene, an optionally substituted diazolylene, an optionally substituted benzopyridazinylene, an optionally substituted benzopyrazinylene, an optionally substituted phenylene, an optionally substituted pyridinylene, an optionally substituted thiophenylene, an optionally substituted furanylene, an optionally substituted piperidinylene, an optionally substituted pyrimidinylene, an optionally substituted anthracenylenylene, an optionally substituted quinolinylene, and an optionally substituted C_{1-6} alkylene.

[0308] In certain embodiments, at least one Q of Q_b^1 through Q_b^p is a 5 membered heteroarylene having at least one heteroatom selected from O, N, S and optionally sub-

stituted with one or more C_{1-10} alkyl. In certain embodiments, at least two Q of Q_b^1 through Q_b^p is a 5 membered heteroarylene having at least one heteroatom selected from O, N, S and optionally substituted with one or more C_{1-10} alkyl. In certain embodiments, at least three, four, five, or six Q of Q_b^1 through Q_b^p is a 5 membered heteroarylene having at least one heteroatom selected from O, N, S and optionally substituted with one or more C_{1-10} alkyl. In some embodiments, at least one of Q_b^1 through Q_b^p is a pyrrole optionally substituted with one or more C_{1-10} alkyl. In some embodiments, at least one of Q_b^1 through Q_b^p is a imidazole optionally substituted with one or more C_{1-10} alkyl. In some embodiments, at least one of Q_b^1 through Q_b^p is a C_{2-6} alkylene optionally substituted with one or more C_{1-10} alkyl. In some embodiments, at least one of Q_b^1 through Q_b^p is a phenyl optionally substituted with one or more C_{1-10} alkyl. In some embodiments, at least one of Q_b^1 through Q_b^p is a bicyclic heteroarylene or arylenylene. In some embodiments, at least one of Q_b^1 through Q_b^p is a phenylene optionally substituted with one or more C_{1-10} alkyl. In some embodiments, at least one of Q_b^1 through Q_b^p is a benzimidazole optionally substituted with one or more C_{1-10} alkyl.


[0309] In certain embodiments, for Formula (A-5c), each end group G_a , G_b , W_a^1 , and W_b^1 is independently selected from the group consisting of optionally substituted C_{6-10} aryl, optionally substituted 4-10 membered heterocyclyl, a 5-10 membered heteroaryl optionally substituted with 1-3 substituents selected from C_{1-6} alkyl, $-\text{NHCOR}^b$, halogen, $-\text{NR}^a\text{R}^b$, an optionally substituted C_{1-6} alkyl, C_{0-4} alkylene-NHC(=NH)NH, C_{0-4} alkylene-NHC(=NH)— R^a , $-\text{C}_{1-4}$ alkylene- R^a , $-\text{CN}$, $-\text{C}_{0-4}$ alkylene-C(=NH)(NR^aR^b), $-\text{C}_{0-4}$ alkylene-C(=N $^+$ H $_2$)(NR^aR^b) C_{1-5} alkylene- NR^aR^b , C_{0-4} alkylene-NHC(=NH) R^a , and optionally substituted amine, wherein each R^a and R^b are independently H, an optionally substituted C_{1-6} alkyl, an optionally substituted C_{3-10} cycloalkyl, optionally substituted C_{6-10} aryl, optionally substituted 4-10 membered heterocyclyl, or an optionally substituted 5-10 membered heteroaryl. In certain embodiments, each R^a and R^b are independently H or C_{1-6} alkyl. In certain embodiments, at least one of the end groups is 5-10 membered heteroaryl optionally substituted with C_{1-6} alkyl, COOH, or OH. In certain embodiments, at least two of the end groups are 5-10 membered heteroaryl optionally substituted with C_{1-6} alkyl, COOH, or OH. In certain embodiments, for Formula (A-1) to (A-5d), at least one of the end groups is 5-10 membered heteroaryl optionally substituted with C_{1-6} alkyl, COOH, or OH. In certain embodiments, at least one of the end groups is 5-10 membered heteroaryl ring optionally substituted with one or more alkyl.

[0310] In some embodiments, A^E is absent. In some embodiments, A^E is $-\text{NHCO}-$.

[0311] In some embodiments, the first terminus comprises at least one C_{3-5} achiral aliphatic or heteroaliphatic amino acid.

[0312] In some embodiments, the first terminus comprises one or more subunits selected from the group consisting of optionally substituted pyrrole, optionally substituted imidazole, optionally substituted thiophene, optionally substituted furan, optionally substituted beta-alanine, γ -aminobutyric acid, (2-aminoethoxy)-propanoic acid, 3((2-aminoethyl)(2-oxo-2-phenyl-1 λ^2 -ethyl)amino)-propanoic acid, or dimethylaminopropylamide monomer.

[0313] In some embodiments, the first terminus comprises a polyamide having the structure of Formula (A-6):



[0314] wherein:

[0315] each A¹ is —NH— or —NH—(CH₂)_m—CH₂—C(O)—NH—;

[0316] each M is an optionally substituted C₆₋₁₀ arylene group, optionally substituted 4-10 membered heterocyclene, optionally substituted 5-10 membered heteroarylene group, or optionally substituted alkylene;

[0317] m is an integer between 1 to 10; and

[0318] n is an integer between 1 and 6.

[0319] In some embodiments, each M¹ in [A¹-M¹] of Formula (A-6) is a C₆₋₁₀ arylene group, 4-10 membered heterocyclene, optionally substituted 5-10 membered heteroarylene group, or C₁₋₆ alkylene; each optionally substituted by 1-3 substituents selected from H, OH, halogen, C₁₋₁₀ alkyl, NO₂, CN, NR'R'', C₁₋₆ haloalkyl, —C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, (C₁₋₆ alkoxy)C₁₋₆ alkyl, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl, C₃₋₇ carbocyclyl, 4-10 membered heterocyclyl, 4-10 membered heterocyclyl, C₆₋₁₀aryl, 5-10 membered heteroaryl, —(C₃₋₇ carbocyclyl)C₁₋₆alkyl, (4-10 membered heterocyclyl)C₁₋₆alkyl, (C₆₋₁₀aryl)C₁₋₆alkyl, (C₆₋₁₀aryl)C₁₋₆alkoxy, (5-10 membered heteroaryl)C₁₋₆alkyl, —(C₃₋₇carbocyclyl)-amine, (4-10 membered heterocyclyl)amine, (C₆₋₁₀aryl)amine, (5-10 membered heteroaryl)amine, acyl, C-carboxy, O-carboxy, C-amido, N-amido, S-sulfonamido, N-sulfonamido, —SR', COOH, or CONR'R'', wherein each R' and R'' are independently H, C₁₋₁₀ alkyl, C₁₋₁₀ haloalkyl, —C₁₋₁₀ alkoxy. In some embodiments, each R¹ in [A¹-R¹] of Formula (A-6) is a 5-10 membered heteroarylene containing at least one heteroatoms selected from O, S, and N or a C₁₋₆ alkylene, and the heteroarylene or the a C₁₋₆ alkylene is optionally substituted with 1-3 substituents selected from OH, halogen, C₁₋₁₀ alkyl, NO₂, CN, NR'R'', C₁₋₆ haloalkyl, —C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, C₃₋₇ carbocyclyl, 4-10 membered heterocyclyl, C₆₋₁₀aryl, 5-10 membered heteroaryl, —SR', COOH, or CONR'R'', wherein each R' and R'' are independently H, C₁₋₁₀ alkyl, C₁₋₁₀ haloalkyl, —C₁₋₁₀ alkoxy. In some embodiments, each R¹ in [A¹-R¹] of Formula (A-6) is a 5-10 membered heteroarylene containing at least one heteroatoms selected from O, S, and N, and the heteroarylene is optionally substituted with 1-3 substituents selected from OH, C₁₋₆ alkyl, halogen, and C₁₋₆ alkoxy.

[0320] In some embodiments, the first terminus has a structure of Formula (A-7):

or a salt thereof, wherein:

[0321] E is an end subunit which comprises a moiety chosen from a heterocyclic group or a straight chain aliphatic group, which is chemically linked to its single neighbor;

[0322] X¹, Y¹, and Z¹ in each m¹ unit are independently selected from CR⁴, N, NR⁵, O, or S;

[0323] X², Y², and Z² in each m³ unit are independently selected from CR⁴, N, NR⁵, O, or S;

[0324] X³, Y³, and Z³ in each m⁵ unit are independently selected from CR⁴, N, NR⁵, O, or S;

[0325] X⁴, Y⁴, and Z⁴ in each m¹ unit are independently selected from CR⁴, N, NR⁵, O, or S;

[0326] each R⁴ is independently H, —OH, halogen, C₁₋₆ alkyl, C₁₋₆ alkoxy;

[0327] each R⁵ is independently H, C₁₋₆ alkyl or C₁₋₆alkylamine;

[0328] each m¹, m³, m⁵ and m⁷ are independently an integer between 0 and 5;


[0329] each m², m⁴ and m⁶ are independently an integer between 0 and 3; and

[0330] m¹+m²+m³+m⁴+m⁵+m⁶+m⁷ is between 3 and 15.

[0331] In some embodiments, m¹ is 3, and X¹, Y¹, and Z¹ in the first unit is respectively CH, N(CH₃), and CH; X¹, Y¹, and Z¹ in the second unit is respectively CH, N(CH₃), and N; and X¹, Y¹, and Z¹ in the third unit is respectively CH, N(CH₃), and N. In some embodiments, m³ is 1, and X², Y², and Z² in the first unit is respectively CH, N(CH₃), and CH. In some embodiments, m⁵ is 2, and X³, Y³, and Z³ in the first unit is respectively CH, N(CH₃), and N; X³, Y³, and Z³ in the second unit is respectively CH, N(CH₃), and N. In some embodiments, m⁷ is 2, and X⁴, Y⁴, and Z⁴ in the first unit is respectively CH, N(CH₃), and CH; X⁴, Y⁴, and Z⁴ in the second unit is respectively CH, N(CH₃), and CH. In some embodiments, each m², m⁴ and m⁶ are independently 0 or 1. In some embodiments, each of the X¹, Y¹, and Z¹ in each m¹ unit are independently selected from CH, N, or N(CH₃). In some embodiments, each of the X², Y², and Z² in each m³ unit are independently selected from CH, N, or N(CH₃). In some embodiments, each of the X³, Y³, and Z³ in each m⁵ unit are independently selected from CH, N, or N(CH₃). In some embodiments, each of the X⁴, Y⁴, and Z⁴ in each m¹ unit are independently selected from CH, N, or N(CH₃). In some embodiments, each Z¹ in each m¹ unit is independently selected from CR⁴ or NRS. In some embodiments, each Z² in each m³ unit is independently selected from CR⁴ or NRS. In some embodiments, each Z³ in each m⁵ unit is independently selected from CR⁴ or NRS. In some embodiments, each Z⁴ in each m¹ unit is independently selected from CR⁴ or NRS. In some embodiments, R⁴ is H, CH₃, or OH. In some embodiments, R⁵ is H or CH₃.

[0332] In some embodiments, for Formula (A-7), the sum of m², m⁴ and m⁶ is between 1 and 6. In some embodiments, for formula (A-7), the sum of m², m⁴ and m⁶ is between 2 and 6. In some embodiments, for Formula (A-7), the sum of


(A-7)



m^1 , m^3 , m^5 and m^1 is between 2 and 10. In some embodiments, the sum of m^1 , m^3 , m^5 and m^1 is between 3 and 8. In some embodiments, for Formula (A-7), $(m^1+m^2+m^3+m^4+m^5+m^6+m^7)$ is between 3 and 12. In some embodiments, $(m^1+m^2+m^3+m^4+m^5+m^6+m^7)$ is between 4 and 10.

[0333] In some embodiments, for Formula (A-1) to (A-7), the first terminus comprises at least one beta-alanine moiety. In some embodiments, for Formula (A-1) to (A-7), the first terminus comprises at least two beta-alanine moieties. In some embodiments, for Formula (A-1) to (A-7), the first terminus comprises at least three or four beta-alanine moieties.


[0334] In some embodiments, the first terminus has the structure of Formula (A-8):



or a salt thereof, wherein:

[0335] E is an end subunit which comprises a moiety chosen from a heterocyclic group or a straight chain aliphatic group, which is chemically linked to its single neighbor;

[0336] W is C_{1-6} alkylene,



[0337] $X^{1'}$, $Y^{1'}$, and $Z^{1'}$ in each n^1 unit are independently selected from CR^4 , N, NR^5 , O, or S;

[0338] $X^{2'}$, $Y^{2'}$, and $Z^{2'}$ in each n^3 unit are independently selected from CR^4 , N, NR^5 , O, or S;

[0339] $X^{3'}$, $Y^{3'}$, and $Z^{3'}$ in each n^5 unit are independently selected from CR^4 , N, NR^5 , O, or S;

[0340] $X^{4'}$, $Y^{4'}$, and $Z^{4'}$ in each n^6 unit are independently selected from CR^4 , N, NR^5 , O, or S;

[0341] $X^{5'}$, $Y^{5'}$, and $Z^{5'}$ in each n^8 unit are independently selected from CR^4 , N, NR^5 , O, or S;

[0342] $X^{6'}$, $Y^{6'}$, and $Z^{6'}$ in each n^{10} unit are independently selected from CR^4 , N, NR^5 , O, or S;

[0343] each R^4 is independently H, —OH, halogen, C_{1-6} alkyl, C_{1-6} alkoxy;

[0344] each R^5 is independently H, C_{1-6} alkyl or C_{1-6} alkylaminine is an integer between 1 and 5;

[0345] each n^1 , n^3 , n^5 , n^6 , n^8 and n^{10} are independently an integer between 0 and 5;

[0346] each n^2 , n^4 , n^7 and n^9 are independently an integer between 0 and 3, and



[0347] $n^1+n^2+n^3+n^4+n^5+n^6+n^7+n^8+n^9+n^{10}$ is between 3 and 15.

[0348] In some embodiments, for Formula (A-8), the sum of n^2 , n^4 , n^7 and n^9 is between 1 and 6. In some embodiments, for Formula (A-8), the sum of n^2 , n^4 , n^1 and n^9 is between 2 and 6. In some embodiments, for Formula (A-8), the sum of n^1 , n^3 , n^5 , n^6 , n^8 and n^{10} is between 3 and 13. In some embodiments, the sum of n^1 , n^3 , n^5 , n^6 , n^8 and n^{10} is between 4 and 10. In some embodiments, for Formula (A-8), $(n^1+n^2+n^3+n^4+n^5+n^6+n^7+n^8+n^9+n^{10})$ is between 3 and 12. In some embodiments, $(n^1+n^2+n^3+n^4+n^5+n^6+n^7+n^8+n^9+n^{10})$ is between 4 and 10.

[0349] In some embodiments, n^1 is 3, and $X^{1'}$, $Y^{1'}$, and $Z^{1'}$ in the first unit is respectively CH, N(CH_3), and CH; $X^{1'}$, $Y^{1'}$,

and $Z^{1'}$ in the second unit is respectively CH, N(CH_3), and N; and $X^{1'}$, $Y^{1'}$, and $Z^{1'}$ in the third unit is respectively CH, N(CH_3), and N. In some embodiments, n^3 is 1, and $X^{2'}$, $Y^{2'}$, and $Z^{2'}$ in the first unit is respectively CH, N(CH_3), and CH. In some embodiments, n^5 is 2, and $X^{3'}$, $Y^{3'}$, and $Z^{3'}$ in the first unit is respectively CH, N(CH_3), and N; $X^{3'}$, $Y^{3'}$, and $Z^{3'}$ in the second unit is respectively CH, N(CH_3), and N. In some embodiments, n^6 is 2, and $X^{4'}$, $Y^{4'}$, and $Z^{4'}$ in the first unit is respectively CH, N(CH_3), and N; $X^{4'}$, $Y^{4'}$, and $Z^{4'}$ in the second unit is respectively CH, N(CH_3), and N. In some embodiments, the $X^{1'}$, $Y^{1'}$, and $Z^{1'}$ in each n^1 unit are independently selected from CH, N, or N(CH_3). In some embodiments, the $X^{2'}$, $Y^{2'}$, and $Z^{2'}$ in each n^3 unit are independently selected from CH, N, or N(CH_3). In some embodiments, the $X^{3'}$, $Y^{3'}$, and $Z^{3'}$ in each n^5 unit are independently selected from CH, N, or N(CH_3). In some embodiments, the $X^{4'}$, $Y^{4'}$, and $Z^{4'}$ in each n^6 unit are independently selected from CH, N, or N(CH_3). In some embodiments, the $X^{5'}$, $Y^{5'}$, and $Z^{5'}$ in each n^8 unit are independently selected from CH, N, or N(CH_3). In some embodiments, the $X^{6'}$, $Y^{6'}$, and $Z^{6'}$ in each n^{10} unit are independently selected from CH, N, or N(CH_3). In some embodiments, each $Z^{1'}$ in each n^1 unit is independently selected from CR^4 or NR^5 . In some embodiments, each $Z^{2'}$ in each n^3 unit is independently selected from CR^4 or NR^5 . In some embodiments, each $Z^{3'}$ in each n^5 unit is independently selected from CR^4 or NR^5 . In some embodiments, each $Z^{4'}$ in each n^6 unit is independently selected from CR^4 or NR^5 . In some embodiments, each $Z^{5'}$ in each n^8 unit is independently selected from CR^4 or NR^5 . In some embodiments, each $Z^{6'}$ in each n^{10} unit is independently selected from CR^4 or NR^6 . In some embodiments, R^4 is H, CH_3 , or OH. In some embodiments, R^5 is H or CH_3 .

[0350] In some embodiments, the first terminus has the structure of Formula (A-9):



[0351] or a salt thereof, wherein:

[0352] X^1 , Y^1 , and Z^1 in each n^1 unit are independently selected from CR^4 , N, NR^5 , O, or S;

[0353] X^2 , Y^2 , and Z^2 in each n^3 unit are independently selected from CR^4 , N, NR^5 , O, or S;

[0354] X^3 , Y^3 , and Z^3 are independently selected from CR^4 , N, NR^5 , O, or S;

[0355] X^4 , Y^4 , and Z^4 in each n^6 unit are independently selected from CR^4 , N, NR^5 , O, or S;

[0356] X^5 , Y^5 , and Z^5 in each n^1 unit are independently selected from CR^4 , N, NR^5 , O, or S;

[0357] X^6 , Y^6 , and Z^6 in each n^9 unit are independently selected from CR^4 , N, NR^5 , O, or S;

[0358] X^7 , Y^7 , and Z^7 in each n^{11} unit are independently selected from CR^4 , N, NR^5 , O, or S;

[0359] X^8 , Y^8 , and Z^8 are independently selected from CR^4 , N, NR^5 , O, or S;

[0360] X^9 , Y^9 , and Z^9 in each n^{14} unit are independently selected from CR^4 , N, NR^5 , O, or S;

[0361] X^{10} , Y^{10} , and Z^{10} in each n^{16} unit are independently selected from CR^4 , N, NR^5 , O, or S;

[0362] each R^4 is independently H, —OH, halogen, C_{1-6} alkyl, C_{1-6} alkoxy;

[0363] each R^5 is independently H, C_{1-6} alkyl or C_{1-6} alkylamine;


[0364] each n^1 , n^3 , n^6 , n^8 , n^9 , n^{11} , n^{14} , and n^{16} are independently an integer between 0 and 5;



[0365] each n^2 , n^4 , n^5 , n^7 , n^{10} , n^{13} , and n^{15} are independently an integer between 0 and 3,

[0366] $n^1+n^2+n^3+n^4+n^5+n^6+n^7+n^8+n^9+n^{10}+n^{11}+n^{12}+n^{13}+n^{14}+n^{15}+n^{16}$ is between 3 and 18


[0367] or a salt thereof, wherein:

[0368] L_a is selected from a divalent or trivalent group selected from the group consisting of





- [0369] each R^{1a} and R^{1b} are independently H, or an C₁₋₆ alkyl;
- [0370] each m and n are independently an integer between 1 and 10;
- [0371] when L_a is a trivalent group, the oligomeric backbone is attached to the first terminus through L_a, and each E_{1a}, E_{2a}, E_{1b}, and E_{2b} are end groups independently selected from the group consisting of optionally substituted C₆₋₁₀ aryl, optionally substituted 4-10 membered heterocyclyl, optionally substituted 5-10 membered heteroaryl, an optionally substituted C₁₋₆ alkyl, and optionally substituted amine;
- [0372] when L_a is a divalent group, the oligomeric backbone is attached to the first terminus through one of E_{1a}, E_{2a}, E_{1b}, and E_{2b}, and each E_{1a}, E_{2a}, E_{1b}, and E_{2b} are independently selected from the group consisting of a bond, a —C₁₋₆ alkylene-, —NH—C₀₋₆ alkylene-C(O)—, —N(CH₃)—C₀₋₆ alkylene, —C(O)—, —C(O)—C₁₋₁₀ alkylene, and —O—C₀₋₆ alkylene, optionally substituted C₆₋₁₀ aryl, optionally substituted 4-10 membered heterocyclyl, optionally substituted 5-10 membered heteroaryl, an optionally substituted C₁₋₆ alkyl, and optionally substituted amine; or
- [0373] when L_a is a bivalent group, the oligomeric backbone is attached to the first terminus through a nitrogen or carbon atom on one of five-membered heteroaryl rings, and each E_{1a}, E_{2a}, E_{1b}, and E_{2b} are end groups independently selected from the group consisting of optionally substituted C₆₋₁₀ aryl, optionally substituted 4-10 membered heterocyclyl, optionally substituted 5-10 membered heteroaryl, an optionally substituted C₁₋₆ alkyl, and optionally substituted amine.
- [0374] In some embodiments, the first terminus comprises a polyamide having the structure of Formula (A-10):



- [0375] wherein:
- [0376] each Y¹, Y², Z¹, and Z² are independently CR⁴, N, NR, O, or S;
- [0377] each R⁴ is independently H, —OH, halogen, C₁₋₆ alkyl, or C₁₋₆ alkoxy;

[0378] each R⁵ is independently H, C₁₋₆ alkyl, or C₁₋₆ alkylamine;

[0379] each W¹ and W² are independently a bond, NH, a C₁₋₆ alkylene, —NH—C₁₋₆ alkylene, —NH—5-10

[0380] membered heteroaryl, —NH—5-10 membered heterocyclene, —N(CH₃)—C₀₋₆ alkylene, —C(O)—, —C(O)—C₁₋₁₀ alkylene, or —O—C₀₋₆ alkylene; and

[0381] n is an integer between 2 and 11.

[0382] In some embodiments, each R⁴ is independently H, —OH, halogen, C₁₋₆ alkyl, C₁₋₆ alkoxy; and each R² is independently H, C₁₋₆ alkyl or C₁₋₆ alkylamine. In some embodiments, each R⁴ is selected from the group consisting of H, COH, Cl, NO, N-acetyl, benzyl, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkenyl, C₁₋₆ alkynyl, C₁₋₆ alkylamine, —C(O)NH—(CH₂)₁₋₄—C(O)NH—(CH₂)₁₋₄—NR^aR^b; and each R^a and R^b are independently hydrogen or C₁₋₆ alkyl.

[0383] In some embodiments, R⁵ is independently selected from the group consisting of H, C₁₋₆ alkyl, and C₁₋₆ alkylNH₂, preferably H, methyl, or isopropyl.


[0384] In some embodiments, R⁴ in Formula (A-7) to (A-8) is independently selected from H, OH, C₁₋₆ alkyl, halogen, and C₁₋₆ alkoxy. In some embodiments, R⁴ in Formula (A-7) to (A-8) is selected from H, OH, halogen, C₁₋₁₀ alkyl, NO₂, CN, NR^aR^b, C₁₋₆ haloalkyl, —C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, (C₁₋₆ alkoxy)C₁₋₆ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₃₋₇ carbocyclyl, 4-10 membered heterocyclyl, C₆₋₁₀ aryl, 5-10 membered heteroaryl, —(C₃₋₇ carbocyclyl)C₁₋₆ alkyl, (4-10 membered heterocyclyl)C₁₋₆ alkyl, (C₆₋₁₀ aryl)C₁₋₆ alkyl, (C₆₋₁₀ aryl)C₁₋₆ alkoxy, (5-10 membered heteroaryl)C₁₋₆ alkyl, —(C₃₋₇ carbocyclyl)-amine, (4-10 membered heterocyclyl)amine, (C₆₋₁₀ aryl)amine, (5-10 membered heteroaryl)amine, acyl, C-carboxy, O-carboxy, C-amido, N-amido, S-sulfonamido, N-sulfonamido, —SR', COOH, or CONR'R"; wherein each R' and R" are independently H, C₁₋₁₀ alkyl, C₁₋₁₀ haloalkyl, —C₁₋₁₀ alkoxy. In some embodiments, R⁴ in Formula (A-7) to (A-8) is selected from O, S, and N or a C₁₋₆ alkylene, and the heteroaryl or the a C₁₋₆ alkylene is optionally substituted with 1-3 substituents selected from OH, halogen, C₁₋₁₀ alkyl, NO₂, CN, NR^aR^b, C₁₋₆ haloalkyl, —C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, C₃₋₇ carbocyclyl, 4-10 membered heterocyclyl, C₆₋₁₀ aryl, 5-10 membered heteroaryl, —SR, COOH, or CONR'R"; wherein each R' and R" are independently H, C₁₋₁₀ alkyl, C₁₋₁₀ haloalkyl, —C₁₋₁₀ alkoxy.

[0385] For the chemical Formula (A-1) to (A-9), each E₁ and E₂ independently are optionally substituted thiophene-containing moiety, optionally substituted pyrrole containing moiety, optionally substituted imidazole containing moiety, and optionally substituted amine. In some embodiments, each E₁ and E₂ are independently selected from the group consisting of N-methylpyrrole, N-methylimidazole, benzimidazole moiety, and 3-(dimethylamino)propanimidyl, each group optionally substituted by 1-3 substituents selected from the group consisting of H, OH, halogen, C₁₋₁₀ alkyl, NO₂, CN, NR^aR^b, C₁₋₆ haloalkyl, —C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, (C₁₋₆ alkoxy)C₁₋₆ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₃₋₇ carbocyclyl, 4-10 membered heterocyclyl, C₆₋₁₀ aryl, 5-10 membered heteroaryl, amine, acyl, C-carboxy, O-carboxy, C-amido, N-amido, S-sulfonamido, N-sulfonamido, —SR, COOH, or CONR'R"; wherein each R' and R" are independently H, C₁₋₁₀ alkyl, C₁₋₁₀ haloalkyl, —C₁₋₁₀ alkoxy. In some embodiments, each E₁ and E₂


independently comprises thiophene, benzothiophene, C—C linked benzimidazole/thiophene-containing moiety, or C—C linked hydroxybenzimidazole/thiophene-containing moiety, wherein each R' and R'' are independently H, C₁₋₁₀ alkyl, C₁₋₁₀ haloalkyl, —C₁₋₁₀alkoxyl.

[0386] In some embodiments, each E, E₁ or E₂ are independently selected from the group consisting of isophthalic acid; phthalic acid; terephthalic acid; morpholine; N,N-dimethylbenzamide; N,N-bis(trifluoromethyl)benzamide; fluorobenzene; (trifluoromethyl)benzene; nitrobenzene; phenyl acetate; phenyl 2,2,2-trifluoroacetate; phenyl dihydrogen phosphate; 2H-pyran; 2H-thiopyran; benzoic acid; isonicotinic acid; and nicotinic acid; wherein one, two or three ring members in any of these end-group candidates can be independently substituted with C, N, S or O; and where any one, two, three, four or five of the hydrogens bound to the ring can be substituted with R₅, wherein R₅ may be independently selected for any substitution from H, OH, halogen, C₁₋₁₀ alkyl, NO₂, NH₂, C₁₋₁₀ haloalkyl, —OC₁₋₁₀ haloalkyl, COOH, CONR'R'', wherein each R' and R'' are independently H, C₁₋₁₀ alkyl, C₁₋₁₀ haloalkyl, —C₁₋₁₀alkoxyl.


[0387] The DNA recognition or binding moiety can include one or more subunits selected from the group consisting of:




(Py)




(Im)




(Hp)




(Th)




(Pz)




(Nt)




(Ht)




(Tn)




(Nh)




(Fr)




(Tp)




(iNt)




(B)




(gAB)



(PyT)




(ImT)




(CTh)

-continued




(iIm)


-continued




(HpBi)






(ImBi)




(PyBi)



("Dp")









—NH-benzopyrazinylene-CO—, —NH-phenylene-CO—,
 —NH-pyridinylene-CO—, —NH-piperidinylene-CO—,
 —NH-pyrimidinylene-CO—, —NH-anthracenylene-CO—,
 —NH-quinolinyne-CO—, and











wherein Z is H, NH₂, C₁₋₆ alkyl, or C₁₋₆ alkylNH₂.

[0388] In some embodiments,




-continued




-continued


—NH-benzopyrazinylene-CO— is




—NH-phenylene-CO— is




—NH-pyridinylene-CO— is




—NH-piperidinylene-CO— is




—NH-pyrazinylene-CO— is



—NH-anthracenylene-CO— is




—NH-quinolinyne-CO— is



[0389] In some embodiments, the first terminus comprises one or more subunits selected from the group consisting of optionally substituted N-methylpyrrole, optionally substituted N-methylimidazole, and β -alanine (β).

[0390] In some embodiments, the first terminus does not have a structure of



[0393] In one aspect, the molecules of the present disclosure preferentially bind to the repeated CAG or CTG of

[0391] The first terminus in the molecules described herein has a high binding affinity to a sequence having multiple repeats of CAG or CTG and binds to the target nucleotide repeats preferentially over other nucleotide repeats or other nucleotide sequences. In some embodiments, the first terminus has a higher binding affinity to a sequence having multiple repeats of CAG or CTG than to a sequence having repeats of CGG. In some embodiments, the first terminus has a higher binding affinity to a sequence having multiple repeats of CAG or CTG than to a sequence having repeats of CCG. In some embodiments, the first terminus has a higher binding affinity to a sequence having multiple repeats of CAG or CTG than to a sequence having repeats of CCTG. In some embodiments, the first terminus has a higher binding affinity to a sequence having multiple repeats of CAG or CTG than to a sequence having repeats of TGGAA. In some embodiments, the first terminus has a higher binding affinity to a sequence having multiple repeats of CAG or CTG than to a sequence having repeats of GGGGCC. In some embodiments, the first terminus has a higher binding affinity to a sequence having multiple repeats of CAG or CTG than to a sequence having repeats of GAA.

[0392] Due to the preferential binding between the first terminus and the target nucleotide repeat, the transcription modulation molecules described herein become localized around regions having multiple repeats of CAG or CTG. In some embodiments, the local concentration of the first terminus or the molecules described herein is higher near a sequence having multiple repeats of CAG or CTG than near a sequence having repeats of CGG. In some embodiments, the local concentration of the first terminus or the molecules described herein is higher near a sequence having multiple repeats of CAG or CTG than near a sequence having repeats of CCTG. In some embodiments, the local concentration of the first terminus or the molecules described herein is higher near a sequence having multiple repeats of CAG or CTG than near a sequence having repeats of TGGAA. In some embodiments, the local concentration of the first terminus or the molecules described herein is higher near a sequence having multiple repeats of CAG or CTG than near a sequence having repeats of GGGGCC. In some embodiments, the local concentration of the first terminus or the molecules described herein is higher near a sequence having multiple repeats of CAG or CTG than near a sequence having repeats of GAA.

dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1 than to CAG or CTG elsewhere in the subject's DNA, due to the high number of CAG or CTG repeats associated with dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1. In one aspect, the molecules of the present disclosure are more likely to bind to the repeated CTG of dmpk than to CTG elsewhere in the subject's DNA due to the high number of CTG repeats associated with dmpk. In one aspect, the molecules of the present disclosure are more likely to bind to the repeated CAG of atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1 than to CAG elsewhere in the subject's DNA, due to the high number of CAG repeats associated with atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1. In one aspect, the molecules of the present disclosure are more likely to bind to the repeated CAG of TCF4 gene than to CAG elsewhere in the subject's DNA, due to the high number of CAG repeats associated with TCF4.

[0394] The first terminus is localized to a sequence having multiple repeats of CAG or CTG and binds to the target nucleotide repeats preferentially over other nucleotide repeats. In some embodiments, the sequence has at least 2, 3, 4, 5, 8, 10, 12, 15, 20, 25, 30, 40, 50, 100, 200, 300, 400, or 500 repeats of CAG or CTG. In certain embodiments, the sequence comprises at least 1000 nucleotide repeats of CAG or CTG. In certain embodiments, the sequence comprises at least 500 nucleotide repeats of CAG or CTG. In certain embodiments, the sequence comprises at least 200 nucleotide repeats of CAG or CTG. In certain embodiments, the sequence comprises at least 100 nucleotide repeats of CAG or CTG. In certain embodiments, the sequence comprises at least 50 nucleotide repeats of CAG or CTG. In certain embodiments, the sequence comprises at least 20 nucleotide repeats of CAG or CTG.

[0395] The polyamide composed of a pre-selected combination of subunits can selectively bind to the DNA in the minor groove. In their hairpin structure, antiparallel side-by-side pairings of two aromatic amino acids bind to DNA sequences, with a polyamide ring packed specifically against each DNA base. N-Methylpyrrole (Py) favors T, A, and C bases, excluding G; N-methylimidazole (Im) is a G-reader; and 3-hydroxyl-N-methylpyrrol (Hp) is specific for thymine base. The nucleotide base pairs can be recognized using different pairings of the amino acid subunits using the pairing principle shown in Table 1A and 1B below. For example, an Im/Py pairing reads G•C by symmetry, a Py/Im pairing reads C•G, an Hp/Py pairing can distinguish T•A from A•T, G•C,

and C•G, and a Py/Py pairing nonspecifically discriminates both A•T and T•A from G•C and C•G.

[0396] In some embodiments, the first terminus comprises Im corresponding to the nucleotide G; Im or Nt corresponding to the nucleotide pair G; Py corresponding to the nucleotide C, wherein Im is N-alkyl imidazole, Py is N-alkyl

pyrrole, Hp is 3-hydroxy N-methyl pyrrole, and β-alanine. In some embodiments, the first terminus comprises Im/Py to correspond to the nucleotide pair G/C, Py/Im to correspond to the nucleotide pair C/G, and wherein Im is N-alkyl imidazole (e.g., N-methyl imidazole), Py is N-alkyl pyrrole (e.g., N-methyl pyrrole), and Hp is 3-hydroxy N-methyl pyrrole.

TABLE 1A







Base pairing for single amino acid subunit (Favored (+), disfavored (-))				
Subunit	G	C	A	T
Py	-	+	+	+
Im	+	-	-	
	-	-	-	+
(Hp)				
	-	-	+	+
(Th)				
	-	-	+	+
(Pz)				
	-	-	+	+
(Tp)				
	+	-	-	-
(Nt)				
	-	-	-	+
(Ht)				

TABLE 1A-continued







Base paring for single amino acid subunit (Favored (+), disfavored (-))				
Subunit	G	C	A	T
 (iPTA)	+	-	-	-
 ("CTh")	-	-	-	+
 PEG	-	+	+	+
 iIm	+	-	-	-
 Ip	+	-	-	-
 Hz	-	-	-	+

TABLE 1A-continued









Base pairing for single amino acid subunit (Favored (+), disfavored (-))				
Subunit	G	C	A	T
 Bi	-	-	-	+
 (gly)	-	-	-	-
 (β)	-	-	+	+
 (gAB)	-	-	+ (as a part of the turn)	+ (as a part of the turn)
 (Alx)	-	+	-	-
 (Da)	-	-	+	+
 (Dp)	-	-	+	+
 (iPP)	-	-	+	+

TABLE 1A-continued







Base pairing for single amino acid subunit (Favored (+), disfavored (-))				
Subunit	G	C	A	T
 (CTh)	+	+	-	-
 (Dab)	-	-	+	+
 (gAH)	-	-	+	+
 HpBi	WW* (bind to two nucleotides with same selectivity as Hp-Py)			
 PyBi	WW* (bind to two nucleotides with same selectivity as Py-Py)			

TABLE 1A-continued

Base paring for single amino acid subunit (Favored (+), disfavored (-))				
Subunit	G	C	A	T
GW* (bind to two nucleotides with same selectivity as Im-Py)				
				
ImBi				

*The subunit HpBi, ImBi, and PyBi function as a conjugate of two monomer subunits and bind to two nucleotides. The binding property of HpBi, ImBi, and PyBi corresponds to Hp-Py, Im-Py, and Py-Py respectively.

TABLE 1B

Base paring for hairpin polyamide				
	G · C	C · G	T · A	A · T
Im/β	+	-	-	-
β/Im	-	+	-	-
Py/β	-	-	+	+
β/Py	-	-	+	+
β/β	-	-	+	+
Py/Py	-	-	+	+
Im/Im	-	-	-	-
Im/Py	+	-	-	-
Py/Im	-	+	-	-
Th/Py	-	-	+	-
Py/Th	-	-	-	+
Th/Im	+	-	-	-
Im/Th	-	+	-	-
β/Th	-	-	+	-
Th/β	-	-	-	+
Hp/Py,	-	-	+	-
Py/Hp,	-	-	-	+
Hp/Im	+	-	-	-
Im/Hp	-	+	-	-
Tn/Py	-	-	+	+
Py/Tn,	-	-	+	+
Ht/Py,	-	-	+	+
Py/Ht,	-	-	+	+
Bi/Py,	-	-	+	+
Py/Bi,	-	-	+	+
β/Bi	-	-	+	+
Bi/β	-	-	+	+
Bi/Im,	-	+	-	-
Im/Bi,	+	-	-	-
Tp/Py,	-	-	+	+
Py/Tp,	-	-	+	+
β/Tp	-	-	+	+
Tp/β	-	-	+	+
Tp/Im,	-	+	-	-
Im/Tp	+	-	-	-
Tp/Tp	-	-	+	+
Tp/Tn	-	-	+	+
Tn/Tp	-	-	+	+
Ht/Py,	-	-	+	-
Py/Ht,	-	-	-	+
Ip/Py	+	-	-	-
Py/Ip,	-	+	-	-

TABLE 1B-continued

Base paring for hairpin polyamide				
	G · C	C · G	T · A	A · T
Bi/Ht,	-	-	+	+
Ht/Bi,	-	-	+	+
Bi/Bi	-	+	+	+
Th/Py,	-	-	+	+
Py/Th	-	-	+	+
Im/gAB	+	-	-	-
gAB/Im	-	+	-	-
Py/gAB	+	-	-	-
gAB/Py	-	+	-	-
gAB/β	-	-	+	+
β/gAB	-	-	+	+
Im/Dp	+	-	-	-
Dp/Im	-	+	-	-
Py/Dp	-	-	+	+
Dp/Py	-	-	+	+
Dp/β	-	-	+	+

Each of HpBi, ImBi, and PyBi can bind to two nucleotides and have binding properties corresponding to Hp-Py, Im-Py, and Py-Py respectively. HpBi, ImBi, and PyBi can be paired with two monomer subunits or with themselves in a hairpin structure to bind to two nucleotide pairs.

[0397] The monomer subunits of the polyamide can be strung together based on the paring principles shown in Table 1A and Table 1B. The monomer subunits of the polyamide can be strung together based on the paring principles shown in Table 1C and Table 1D.


[0398] Table 1C shows an example of the monomer subunits that can bind to the specific nucleotide. The first terminus can include a polyamide described having several monomer subunits stung together, with a monomer subunit selected from each row. For example, the polyamide can include Py-Py-Im that binds to CAG, with Py is selected from the C column, Py is selected from the A column, and Im selected from the first G column. The polyamide can be any combinations of the subunits of CAGCAG, with a subunit selected from each column in Table 1C, wherein the subunits are strung together following the CAG binding order. In another example, the polyamide can include Py-β-

Im that binds to CTG, with Py selected from the C column, β from the T column, and Im from the G column.


[0399] In addition, the polyamide can also include a partial or multiple sets of the five subunits, such as 1.5, 2, 2.5, 3, 3.5, or 4 sets of the three subunits. The polyamide can include 3, 4, 5, 6, 7, 8, 9, 10, 12, 14, and 16 monomer subunits. The multiple sets can be joined together by W. In addition to the five subunits or ten subunits, the polyamide can also include 1-4 additional subunits that can link multiple sets of the five subunits.

[0400] The polyamide can include monomer subunits that bind to 2, 3, 4, or 5 nucleotides of CAG or CTG. For example, the polyamide can bind to CA, CAG, AGC, CAGC, CAGCA, CAGCAG. The polyamide can include monomer subunits that bind to 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides of CAG repeat. For example, the polyamide can bind to CT, CTG, TGC, CTGC, CTGCT, CTGCTG, CTGCTGC. The polyamide can include monomer subunits that bind to 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides of CTG repeat. The nucleotides can be joined by W.



[0401] The monomer subunit, when positioned as a terminal unit, does not have an amine, carbonyl, or a carboxylic acid group at the terminal. The amine or carboxylic acid group in the terminal is replaced by a hydrogen. For example, Py, when used as a terminal unit, is understood to have the structure of



and Im, when positioned as a terminal unit, is understood to have the structure of



In addition, when Py or Im is used as a terminal unit, Py and Im can be respectively replaced by



[0402] The linear polyamide can have nonlimiting examples including but not limited to Py-Py-Im-Py-Py-Im-Py-Py-Im, β -Im-Py- β -Im-Py- β -Im, Im-Py-Py-Im-Py-Py-Im, Im-Py-Py-Im-Py- β , β -Im-Py-Py-Im-Py- β , Py-Py-Im- β - β -Im-Py-Py-Im, and any combinations thereof.

TABLE 1C

Examples of monomer subunits in a linear polyamide that binds to CAG or CTG.						
Nucleotide	C	A	G	or C	T	G
Subunit that selectively binds to nucleotide	Py or PyT ilm or ilmT PEG CTh Alx	Py Th Pz Tp PEG β iPP Da Dab gAH	Im or ImT ilm or ilmT iImT Pz CTh Nt iPTA Ip CTh PEG Hz Bi Da Dp iPP Dab gAH	Py or PyT ilm or ilmT PEG CTh Alx	β Py Hp Th Pz Tp Ht CTh PEG Hz Bi Da Dp iPP Dab gAH	Im or ImT ilm or ilmT iImT PEG CTh Nt iPTA Ip CTh PEG Hz Bi Da Dp iPP Dab gAH

[0403] The DNA-binding moiety can also include a hairpin polyamide having subunits that are strung together based on the pairing principle shown in Table 1B. Table 1D shows some examples of the monomer subunit pairs that selectively bind to the nucleotide pair. The hairpin polyamide can include 2n monomer subunits (n is an integer in the range of 2-8), and the polyamide also includes a W in the center of the 2n monomer subunits. W can be $-(CH_2)_a-NR^1-(CH_2)_b-$, $-(CH_2)_a-$, $-(CH_2)_a-O(CH_2)_b-$, $-(CH_2)_a-CH(NHR^1)-$, $-(CH_2)_a-CH(NHR^1)-$, $-(CR^2R^3)_a-$ or $-(CH_2)_a-CH(NR^1_3)^+- (CH_2)_b-$, wherein each a is independently an integer between 2 and 4; R^1 is H, an optionally substituted C_{1-6} alkyl, an optionally substituted C_{3-10} cycloalkyl, an optionally substituted C_{6-10} aryl, an optionally substituted 4-10 membered heterocyclyl, or an optionally substituted 5-10 membered heteroaryl; each R^2

and R^3 are independently H, halogen, OH, NHAc, or C_{1-4} alkyl. In some embodiments, W is $-(CH_2)-CH(NH_3)^+-$ $(CH_2)-$ or $-(CH_2)-CH_2CH(NH_3)^+-$. In some embodiments, R^1 is H. In some embodiments, R^1 is C_{1-6} alkyl optionally substituted by 1-3 substituents selected from $-C(O)-$ phenyl. In some embodiments, W is $-(CR^2R^3)-(CH_2)_a-$ or $-(CH_2)_a-(CR^2W)-(CH_2)_b-$, wherein each a is independently 1-3, b is 0-3, and each R^2 and R^3 are independently H, halogen, OH, NHAc, or C_{1-4} alkyl. W can be an aliphatic amino acid residue shown in Table 4 such as gAB.

[0404] When n is 2, the polyamide includes 4 monomer subunits, and the polyamide also includes a W joining the first set of two subunits with the second set of two subunits, Q1-Q2-W-Q3-Q4, and Q1/Q4 correspond to a first nucleotide pair on the DNA double strand, Q2/Q3 correspond to a second nucleotide pair, and the first and the second nucleotide pair is a part of the GGGGCC repeat. When n is 3, the polyamide includes 6 monomer subunits, and the polyamide also includes a W joining the first set of three subunits with the second set of three subunits, Q1-Q2-Q3-W-Q4-Q5-Q6, and Q1/Q6 correspond to a first nucleotide pair on the DNA double strand, Q2/Q5 correspond to a second nucleotide pair, Q3/Q4 correspond to a third nucleotide pair, and the first and the second nucleotide pair is a part of the A repeat. When n is 4, the polyamide includes 8 monomer subunits, and the polyamide also includes a W joining the first set of four subunits with the second set of four subunits, Q1-Q2-Q3-Q4-W-Q5-Q6-Q7-Q8, and Q1/Q8 correspond to a first nucleotide pair on the DNA double strand, Q2/Q7 correspond to a second nucleotide pair, Q3/Q6 correspond to a third nucleotide pair, and Q4/Q5 correspond to a fourth nucleotide pair on the DNA double strand. When n is 5, the polyamide includes 10 monomer subunits, and the polyamide also includes a W joining a first set of five subunits with a second set of five subunits, Q1-Q2-Q3-Q4-Q5-W-Q6-Q7-Q8-Q9-Q10, and Q1/Q10, Q2/Q9, Q3/Q8, Q4/Q7, Q5/Q6 respectively correspond to the first to the fifth nucleotide pair on the DNA double strand. When n is 6, the polyamide includes 12 monomer subunits, and the polyamide also includes a W joining a first set of six subunits with a second set of six subunits, Q1-Q2-Q3-Q4-Q5-Q6-W-Q7-Q8-Q9-Q10-Q11-Q12, and Q1/Q12, Q2/Q11, Q3/Q10, Q4/Q9, Q5/Q8, Q6/Q7 respectively correspond to the first to the six nucleotide pair on the DNA double strand. When n is 8, the polyamide includes 16 monomer subunits, and the polyamide also includes a W joining a first set of eight subunits with a second set of eight subunits, Q1-Q2-Q3-Q4-Q5-Q6-Q7-Q8-W-Q9-Q10-Q11-Q12-Q13-Q14-Q15-Q16, and Q1/Q16, Q2/Q15, Q3/Q14, Q4/Q13, Q5/Q12, Q6/Q11, Q7/Q10, and Q8/Q9 respectively correspond to the first to the eight nucleotide pair on the DNA double strand. When n is 9, the polyamide includes 18 monomer subunits, and the polyamide also includes a W joining a first set of eight subunits with a second set of eight subunits, Q1-Q2-Q3-Q4-Q5-Q6-Q7-Q8-Q9-W-Q10-Q11-Q12-Q13-Q14-Q15-Q16-Q17-Q18, and Q1/Q18, Q2/Q17, Q3/Q16, Q4/Q15, Q5/Q14, Q6/Q13, Q7/Q12, Q8/Q11, and Q9/Q10 respectively correspond to the first to the eight nucleotide pair on the DNA double strand. In some hairpin polyamide structures, the number of monomer subunits on each side of W can be different, and one side of the hairpin can partial pair with the other side of the hairpin to bind the nucleotide pairs on a double strand DNA based on the binding principle in

Table 1B and 1D, while the rest of the unpaired monomer subunit(s) can bind to the nucleotide based on the binding principle in Table 1A and 1C but does not pair with the monomer subunit on the other side. The hairpin polyamide can have one or more overhanging monomer subunit that binds to the nucleotide but does not pair with the monomer subunit on the antiparallel strand. For example, the hairpin structure can include 5 monomer subunits on one side of W and 4 monomer subunits on the other side of W, Q1-Q2-Q3-Q4-Q5-W-Q6-Q7-Q8-Q9, and Q2/Q9, Q3/Q8, Q4/Q7, Q5/Q6 respectively correspond to the first to the fourth nucleotide pair on the DNA double strand, and Q1 binds to a single nucleotide but does not pair with a monomer subunit on the other strand to bind with a nucleotide pair. W can be an aliphatic amino acid residue such as gAB or other appropriate spacers as shown in Table 4. When W is gAB, it favors binding to T.

[0405] Because the target gene can include multiple repeats of CAG or CTG, the subunits can be strung together to bind at least two, three, four, five, six, seven, eight, nine, or ten nucleotides in one or more CAG or CTG repeat (e.g., CAGCAG, or CTGCTGCTG). For example, the polyamide can bind to the CAG or CTG repeat by binding to a partial copy, a full copy, or a multiple repeats of CAG or CTG such as CA, CAG, AGC, CAGC, CAGCA, CAGCAG, CT, CTG, TGC, CTGC, CTGCT, CTGCTG. For example, the polyamide can include Im-Im-Im-Im- β - β -W-Im-Im- β -Py- β -Py that binds to GGGGCC and its complementary nucleotides on a double strand DNA, in which the Im/Py pair binds to the G•C, the Im/P pair binds to G C, the Im/Py pair binds to G•C, the Im/ β binds to G•C, and β /Im binds to C•G; and β /Im binds to C•G. In one example Py- β -Im- β -W-Im-Py-Py-Im that binds to CTGC and its complementary nucleotides on a double strand DNA, in which Py/Im pair binds to C•G, β /Py pair binds to T•A, Im/Im pair binds to C•G and β /Py pair binds to C•G. W can be an aliphatic amino acid residue such as gAB or other appropriate spacers as shown in Table 4. In another example, the polyamide can include Im-Py-Py-Im-Py-gAB-Im-Py-Py-Im- β that binds to GCTGC and its complementary nucleotides on a double strand DNA, in which the Im/ β pair binds to G•C, the Py/Im pair binds to C•G the Py/Py binds to T•A, Im/Py pair binds to the G•C, and Py/Im binds to G. In another example, Im-Py-Py-Im-Py-gAB-Im-Py-Py binds to GCTGC with a part of the complementary nucleotides (ACG) on the double strand DNA, in which Im binds to G, Py binds to C, Py/Py binds to TA, Im/Py binds to the G•C, and Py/Im binds to the BG.

[0406] Some additional examples of the polyamide include but are not limited to Py- β -Im- β -gAB-Im-Py-Py-Im, Im-Py-Py-Im-Py-gAB-Im-Py-Py-Im- β , Py-Py-Im-Py-gAB-Im-Py-Py-Im- β , Py-Py-Im-Py-gAB-Im-Py-Py-Im, Py-Py-Im-Py-gAB-Im-Py-Py, Im-Py-Py-Im-Py-gAB-Im-Py-Py-Im, Im-Py-Py-Im-Py-gAB-Im-Py-Py, and any combinations thereof.

TABLE 1D

Examples of monomer pairs in a hairpin or H-pin polyamide that binds to CAG or CTG.							
Nucleotide	C•G	A•T	G•C	or	C•G	T•A	G•C
Subunit pairs that	β /Im Py/Im	Py/ β β /Py	Im/ β Im/Py		β /Im Py/Im	Py/ β β /Py	Im/ β Im/Py

TABLE 1D-continued

Examples of monomer pairs in a hairpin or H-pin polyamide that binds to CAG or CTG.						
Nucleotide	C•G	A•T	G•C	or	C•G	T•A
selectively binds to nucleotide	Im/Th	β/β	Th/Im		Im/Th	β/β
	Im/Hp	Py/Py	Hp/Im		Im/Hp	Py/Py
	Bi/Im	Py/Th	Im/Bi		Bi/Im	Th/Py
	Tp/Im	Th/β	Im/Tp		Tp/Im	β/Th
	Py/Ip	Py/Hp,	Ip/Py		Py/Ip	Hp/Py
	Bi/Bi	Tn/Py	Im/gAB		Bi/Bi	Bi/Py
	gAB/Im	Py/Tn,	Py/gAB		gAB/Im	Py/Bi
	gAB/Py	Ht/Py,	Im/Dp		gAB/Py	β/Bi
	Dp/Im	Py/Ht,			Dp/Im	Bi/β
		Bi/Py,				Tp/Py
		Py/Bi,				Py/Tp
		β/Bi				β/Tp
		Bi/β				Tp/β
		Tp/Py,				Tp/Tp
		Py/Tp,				Tp/Tn
		β/Tp				Tn/Tp
		Tp/β				Ht/Py
		Tp/Tp				Bi/Ht,
		Tp/Tn				Ht/Bi,
		Tn/Tp				Bi/Bi
		Py/Ht,				Th/Py,
		Bi/Ht,				Py/Th
		Ht/Bi,				gAB/β
		Bi/Bi				β/gAB
		Th/Py,				Py/Dp
		Py/Th				Dp/Py
		gAB/β				Dp/β
		β/gAB				β/Dp
		Py/Dp				
		Dp/Py				
		Dp/β				


[0407] Recognition of a nucleotide repeat or DNA sequence by two antiparallel polyamide strands depends on a code of side-by-side aromatic amino acid pairs in the minor groove, usually oriented N to C with respect to the 5' to 3' direction of the DNA helix. Enhanced affinity and specificity of polyamide nucleotide binding is accomplished by covalently linking the antiparallel strands. The “hairpin motif” connects the N and C termini of the two strands with a W (e.g., gamma-aminobutyric acid unit (gamma-turn)) to form a folded linear chain. The “H-pin motif” connects the antiparallel strands across a central or near central ring/ring pairs by a short, flexible bridge.

[0408] The DNA-binding moiety can also include a H-pin polyamide having subunits that are strung together based on the pairing principles shown in Table 1A and/or Table 1B. Table 1C shows some examples of the monomer subunit that selectively binds to the nucleotide, and Table 1D shows some examples of the monomer subunit pairs that selectively bind to the nucleotide pair. The h-pin polyamide can include 2 strands and each strand can have a number of monomer subunits (each strand can include 2-8 monomer subunits), and the polyamide also includes a bridge L₁ to connect the two strands in the center or near the center of each strand. At least one or two of the monomer subunits on each strand are paired with the corresponding monomer subunits on the other stand following the pairing principle in Table 1D to favor binding of either G•C or C•G pair, and these monomer subunit pairs are often positioned in the center, close to center region, at or close to the bridge that connects the two strands. In some instances, the H-pin polyamide can have all of the monomer subunits be paired with the corresponding monomer subunits on the antiparallel


strand based on the pairing principle in Table 1B and 1D to bind to the nucleotide pairs on the double strand DNA. In some instances, the H-pin polyamide can have a part of the monomer subunits (2, 3, 4, 5, or 6) be paired with the corresponding monomer subunits on the antiparallel strand based on the binding principle in Table 1B and 1D to bind to the nucleotide pairs on the double strand DNA, while the rest of the monomer subunit binds to the nucleotide based on the binding principle in Table 1A and 1C but does not pair with the monomer subunit on the antiparallel strand. The h-pin polyamide can have one or more overhanging monomer subunit that binds to the nucleotide but does not pair with the monomer subunit on the antiparallel strand.

[0409] Another polyamide structure that derives from the h-pin structure is to connect the two antiparallel strands at the end through a bridge, while only the two monomer subunits that are connected by the bridge form a pair that bind to the nucleotide pair G•C or C•G based on the binding principle in Table 1B/1D, but the rest of the monomer subunits on the strand form an overhang, bind to the nucleotide based on the binding principle in Table 1A and/or 1C and do not pair with the monomer subunit on the other strand.

[0410] The bridge can be is a bivalent or trivalent group selected from



a C₁₋₁₀ alkylene, —NH—C₀₋₆ alkylene—C(O)—, —N(CH₃)—C₀₋₆ alkylene, and



—(CH₂N^R—(CH₂)_b—, —(CH₂)_a—, —(CH₂)_a—O—(CH₂)_b—, —(CH₂)_a—CH(NHR¹)—, —(CH₂)_a—CH(NHR¹)—, —(CR²R³)_a— or —(CH₂)_a—CH(NR¹₃)—(CH₂)_b—, wherein m is an integer in the range of 0 to 10; n is an integer in the range of 0 to 10; each a is independently an integer between 2 and 4; R¹ is H, an optionally substituted C₁₋₆ alkyl, an optionally substituted C₃₋₁₀ cycloalkyl, an optionally substituted C₆₋₁₀ aryl, an optionally substituted 4-10 membered heterocyclyl, or an optionally substituted 5-10 membered heteroaryl; each R² and R³ are independently H,

halogen, OH, NHAc, or C_{1-4} alkyl. In some embodiments, W is $-(CH_2)-CH(NH_3^+)-(CH_2)-$ or $-(CH_2)-CH_2CH(NH_3^+)-$. In some embodiments, R^1 is H. In some embodiments, R^1 is C_{1-6} alkyl optionally substituted by 1-3 substituents selected from $-C(O)-phenyl$. In some embodiments, L_1 is $-(CR^2R^3)-(CH_2)_a-$ or $-(CH_2)_a-(CR^2R^3)-(CH_2)_b-$, wherein each a is independently 1-3, b is 0-3, and each R^2 and R^3 are independently H, halogen, OH, NHAc, or C_{1-4} alkyl. L_1 can be a C_{2-9} alkylene or $(PEG)_{2-8}$.

[0411] When n is 3, the polyamide includes 6 monomer subunits, and the polyamide also includes a bridge L_1 joining the first set of three subunits with the second set of three subunits, and Q1-Q2-Q3 can be joined to Q4-Q5-Q6 through L_1 at the center Q2 and Q5, and Q1/Q4 correspond to a first nucleotide pair on the DNA double strand, Q2/Q5 correspond to a second nucleotide pair, Q3/Q6 correspond to a third nucleotide pair. When n is 4, the polyamide includes 8 monomer subunits, and the polyamide also includes a bridge L_1 joining the first set of four subunits with the second set of four subunits, Q1-Q2-Q3-Q4 can be joined to Q5-Q6-Q7-Q8 through L_1 at Q2 and Q6 Q2 and Q7, Q3 and Q6, or Q3 and Q7 positions; Q1/Q5 may correspond to a nucleotide pair on the DNA double strand, and Q3/Q8 may correspond to another nucleotide pair; or Q1 and Q8 form overhangs on each strand, or Q and Q5 form overhangs on each strand. When n is 5, the polyamide includes 10 monomer subunits, and the polyamide also includes a bridge L_1 joining a first set of five subunits with a second set of five subunits, and Q1-Q2-Q3-Q4-Q5 can be joined to Q6-Q7-Q8-Q9-Q10 through a bridge L_1 at non-terminal positions (any position except for Q1, Q5, Q6 and Q10); if the two strands are linked at Q3 and Q8 by the bridge, Q1/Q6, Q2/Q7, Q3/Q8, Q4/Q9, and Q5/Q10 can be paired to bind to the nucleotide pairs; if the two strands are linked at Q2 and Q9 by the bridge, then Q1/Q8, Q3/Q10 can be paired to bind to the nucleotide pairs, Q4 and Q5 form an overhang on one strand and Q6 and Q7 form an overhang on the other strand.

[0412] In some embodiments, the monomer subunit at the central or near the central ($n/2$, ($n\pm 1$)/2) on one strand is paired with the corresponding one on the other strand to bind to the nucleotide pairs on the double stranded DNA. In some embodiments, the monomer subunit at the central or near the central ($n/2$, ($n\pm 1$)/2) on one strand is connected with the corresponding one on the other strand through a bridge L_1 .

[0413] When n is 4, the polyamide includes 8 monomer subunits, and the polyamide also includes a bridge L_1 joining the first set of four subunits with the second set of four subunits, Q1-Q2-Q3-Q4 can be joined to Q5-Q6-Q7-Q8 at the end Q4 and Q5 through L_1 , while Q4/Q5 can be paired to bind to the nucleotide pairs, Q1-Q2-Q3 form an overhang on one strand and Q6-Q7-Q8 form an overhang on the other strand.

[0414] Some additional examples of the polyamide include but are not limited to Py-Py-Im-Py (linked in the middle—either position 2 or 3) to Im-Py-Py-Im, Py- β -Im- β (linked in the middle—either position 2 or 3) Im-Py-Py-Im, Im-Py-Py-Im-Py (linked in the middle—either position 2, 3, or 4) Im-Py-Py-Im- β , Py-Py-Im-Py (middle position 2 or 3 of Py-Py-Im-Py linked with position 2, 3, or 4 of Im-Py-Py-Im- β) Im-Py-Py-Im- β , Py-Py-Im-Py (linked in the middle—either position 2 or 3) Im-Py-Py-Im, Py-Py-Im-Py (middle position 2 or 3 of Py-Py-Im-Py linked with position 2 of Im-Py-Py) Im-Py-Py, Im-Py-Py-Im-Py (either middle

position 2, 3, or 4 of Im-Py-Py-Im-Py linked with middle position 2 or 3 of Im-Py-Py-Im) Im-Py-Py-Im, Im-Py-Py-Im-Py (middle position 2, 3, or 4 linked with middle position of Im-Py-Py) Im-Py-Py.

Second Terminus—Regulatory Protein Binding Moiety

[0415] In certain embodiments, the regulatory molecule is chosen from a nucleosome remodeling factor (“NURF”), a bromodomain PHD finger transcription factor (“BPTF”), a ten-eleven translocation enzyme (“TET”), methylcytosine dioxygenase (“TET1”), a DNA demethylase, a helicase, an acetyltransferase, and a histone deacetylase (“HDAC”).

[0416] The binding affinity between the regulatory protein and the second terminus can be adjusted based on the composition of the molecule or type of protein. In some embodiments, the second terminus binds the regulatory molecule with an affinity of less than about 600 nM, about 500 nM, about 400 nM, about 300 nM, about 250 nM, about 200 nM, about 150 nM, about 100 nM, or about 50 nM. In some embodiments, the second terminus binds the regulatory molecule with an affinity of less than about 300 nM. In some embodiments, the second terminus binds the regulatory molecule with an affinity of less than about 200 nM. In some embodiments, the polyamide is capable of binding the DNA with an affinity of greater than about 200 nM, about 150 nM, about 100 nM, about 50 nM, about 10 nM, or about 1 nM. In some embodiments, the polyamide is capable of binding the DNA with an affinity in the range of about 1-600 nM, 10-500 nM, 20-500 nM, 50-400 nM, 100-300 nM, or 50-200 nM.

[0417] In some embodiments, the second terminus comprises one or more optionally substituted C_{6-10} aryl, optionally substituted C_{4-10} carbocyclic, optionally substituted 4 to 10 membered heterocyclic, or optionally substituted 5 to 10 membered heteroaryl.

[0418] In some embodiments, the protein-binding moiety binds to the regulatory molecule that is selected from the group consisting of a CREB binding protein (CBP), a P300, an O-linked β -N-acetylglucosamine-transferase-(OGT-), a P300-CBP-associated-factor- (PCAF-), histone methyltransferase, histone demethylase, chromodomain, a cyclin-dependent-kinase-9- (CDK9-), a nucleosome-remodeling-factor-(NURF-), a bromodomain-PHD-finger-transcription-factor-(BPTF-), a ten-eleven-translocation-enzyme-(TET-), a methylcytosine-dioxygenase-(TET1-), histone acetyltransferase (HAT), a histone deacetylase (HDAC), a host-cell-factor-1(HCF1-), an octamer-binding-transcription-factor-(OCT1-), a P-TEFb-, a cyclin-T1-, a PRC2-, a DNA-demethylase, a helicase, an acetyltransferase, a histone-deacetylase, methylated histone lysine protein.

[0419] In some embodiments, the second terminus comprises a moiety that binds to an O-linked β -N-acetylglucosamine-transferase (OGT), or CREB binding protein (CBP). In some embodiments, the protein binding moiety is a residue of a molecule that binds to an O-linked β -N-acetylglucosamine-transferase (OGT), or CREB binding protein (CBP).

[0420] In some embodiments, the second terminus comprises a moiety that binds to p300/CBP HAT (histone acetyltransferase).

[0421] In some embodiments, the second terminus does not comprises JQ1, iBET762, OTX015, RVX208, or AU1. In some embodiments, the second terminus does not com-

prises JQ1. In some embodiments, the second terminus does not comprise a moiety that binds to a bromodomain protein.

[0422] In some embodiments, the second terminus comprises a diazine or diazepine ring, wherein the diazine or diazepine ring is fused with a C₆₋₁₀ aryl or a 5-10 membered heteroaryl ring comprising one or more heteroatom selected from S, N and O.


[0423] In some embodiments, the second terminus comprises an optionally substituted bicyclic or tricyclic structure. In some embodiments, the optionally substituted bicyclic or tricyclic structure comprises a diazepine ring fused with a thiophene ring.

[0424] In some embodiments, the second terminus does not comprise an optionally substituted bicyclic structure, wherein the bicyclic structure comprises a diazepine ring fused with a thiophene ring.

[0425] In some embodiments, the second terminus does not comprise an optionally substituted tricyclic structure, wherein the tricyclic structure is a diazepine ring that is fused with a thiophene and a triazole.

[0426] In some embodiments, the second terminus does not comprise an optionally substituted diazine ring.

[0427] In some embodiments, the second terminus does not comprise a structure of Formula (C-11):



(C-11)

[0428] wherein:

[0429] each of A^{1p} and B^{1p} is independently an optionally substituted aryl or heteroaryl ring;

[0430] X^{1p} is CH or N;





[0431] R^{1p} is hydrogen, halogen, or an optionally substituted C₁₋₆ alkyl group; and

[0432] R^{2p} is an optionally substituted C₁₋₆ alkyl, cycloalkyl, C₆₋₁₀ aryl, or heteroaryl.


[0433] In some embodiments, X^{1p} is N. In some embodiments, A^{1p} is an aryl or heteroaryl substituted with one or more substituents. In some embodiments, A^{1p} is an aryl or heteroaryl substituted with one or more substituents selected from halogen, C₁₋₆alkyl, hydroxyl, C₁₋₆alkoxy, and C₁₋₆haloalkyl. In some embodiments, B^{1p} is an optionally substituted aryl or heteroaryl substituted with one or more substituents selected from halogen, C₁₋₆alkyl, hydroxyl, C₁₋₆alkoxy, and C₁₋₆haloalkyl.

[0434] In some embodiments, A^{1p} is an optionally substituted thiophene or phenyl. In some embodiments, A^{1p} is a thiophene or phenyl, each substituted with one or more substituents selected from halogen, C₁₋₆alkyl, hydroxyl, C₁₋₆alkoxy, and C₁₋₆haloalkyl. In some embodiments, B^{1p} is an optionally substituted triazole. In some embodiments, B^{1p} is a triazole substituted with one or more substituents selected from halogen, C₁₋₆alkyl, hydroxyl, C₁₋₆alkoxy, and C₁₋₆haloalkyl.


[0435] In some embodiments, the protein binding moiety is not



[0436] In some embodiments, the protein binding moiety is not



[0437] In some embodiments, the protein binding moiety does not have the structure of Formula (C-12):



(C-12)

[0438] wherein:

[0439] R_{1q} is a hydrogen or an optionally substituted alkyl, hydroxyalkyl, aminoalkyl, alkoxyalkyl, halogenated alkyl, hydroxyl, alkoxy, or $-\text{COOR}_{4q}$;

[0440] R_{4q} is hydrogen, or an optionally substituted aryl, aralkyl, cycloalkyl, heteroaryl, heteroaralkyl, heterocycloalkyl, alkyl, alkenyl, alkynyl, or cycloalkylalkyl group, optionally containing one or more heteroatoms;

[0441] R_{2q} is an optionally substituted aryl, alkyl, cycloalkyl, or aralkyl group;

[0442] R_{3q} is hydrogen, halogen, or an optionally substituted alkyl group, preferably $(\text{CH}_2)_x-\text{C}(\text{O})(\text{R}_{20})(\text{R}_{21})$, or $(\text{CH}_2)_x-\text{N}(\text{R}_{20})-\text{C}(\text{O})\text{R}_{21}$; or halogenated alkyl group;

[0443] wherein x is an integer from 1 to 10; and R_{20} and R_{21} are each independently hydrogen or C_1 - C_6 alkyl group, preferably R_{20} is hydrogen and R_{21} is methyl; and

[0444] Ring E is an optionally substituted aryl or heteroaryl group.

[0445] The protein binding moiety can include a residue of a compound that binds to a regulatory protein. In some embodiments, the protein binding moiety can be a residue of a compound shown in Table 2. Exemplary residues include, but are not limited to, amides, carboxylic acid esters, thioesters, primary amines, and secondary amines of any of the compounds shown in Table 2.

TABLE 2





A list of compounds that bind to regulator proteins.	
Target protein	Compound
p300/CBP HAT (histone acetyltransferase)	 Lys-CoA
p300/CBP HAT (histone acetyltransferase)	 H3-CoA-20
p300/CBP HAT (histone acetyltransferase)	 anacardic acid (AA)
p300/CBP HAT (histone acetyltransferase)	 curcumin
p300/CBP HAT (histone acetyltransferase)	 MB-3

TABLE 2-continued




A list of compounds that bind to regulator proteins.	
Target protein	Compound
p300/CBP HAT (histone acetyltransferase)	 isothiazolones X = H, Cl R = NO ₂ , Cl, CF ₃ , OCH ₃ , COOC ₂ H ₅
p300/CBP HAT (histone acetyltransferase)	 garcinol
p300/CBP HAT (histone acetyltransferase)	 MC1823 (4)

TABLE 2-continued




A list of compounds that bind to regulator proteins.	
Target protein	Compound
p300/CBP HAT (histone acetyltransferase)	 MC1626 (R = CH ₃) MC1752 (R = H)
	
p300/CBP HAT (histone acetyltransferase)	1 (R = OC ₂ H ₅ ; R' = CH ₃) 2 (R = OH; R' = CH ₃) 3 (R = OC ₂ H ₅ ; R' = C ₃ H ₁₁) 5 (R = OC ₂ H ₅ ; R' = C ₁₀ H ₂₁) 6 (R = OH; R' = C ₁₀ H ₂₁) 7 (R = OC ₂ H ₅ ; R' = C ₁₆ H ₃₁) 8 (R = OH; R' = C ₁₆ H ₃₁)
	
	CBP30 Time (min)

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	
p300/CBP HAT (histone acetyltransferase)	
p300/CBP HAT (histone acetyltransferase)	

TABLE 2-continued





Target protein	Compound
p300/CBP HAT (histone acetyltransferase)	
p300/CBP HAT (histone acetyltransferase)	
p300/CBP HAT (histone acetyltransferase)	
p300/CBP HAT (histone acetyltransferase)	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
p300/CBP HAT (histone acetyltransferase)	
p300/CBP HAT (histone acetyltransferase)	
p300/CBP HAT (histone acetyltransferase)	

TABLE 2-continued

A list of compounds that bind to regulator proteins.







Target protein	Compound
p300/CBP HAT (histone acetyltransferase)	      Ph Me i-Pr

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
p300/CBP HAT (histone acetyltransferase)	
p300/CBP HAT (histone acetyltransferase)	
p300/CBP HAT (histone acetyltransferase)	   R H 3-Me 2-CH ₂ NH ₂ R
p300/CBP HAT (histone acetyltransferase)	 Ph i-Pr i-Pr

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
p300/CBP HAT (histone acetyltransferase)	
p300/CBP HAT (histone acetyltransferase)	 X = Cl, (R,R)-31 X = Br, (R,R)-32
p300/CBP HAT (histone acetyltransferase)	 X = Cl, (S,S)-31 X = Br, (S,S)-32
p300/CBP HAT (histone acetyltransferase)	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	
	
p300/CBP HAT (histone acetyltransferase)	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	
	
p300/CBP HAT (histone acetyltransferase)	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
p300/CBP HAT (histone acetyltransferase)	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.

Target protein	Compound
p300/CBP HAT (histone acetyltransferase)	 Garcinol
p300/CBP HAT (histone acetyltransferase)	 C646

② indicates text missing or illegible when filed

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	
	
	

TABLE 2-continued



A list of compounds that bind to regulator proteins.	
Target protein	Compound
p300/CBP HAT (histone acetyltransferase)	
	
	3a R = H 3b R = Me

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound										
											
											
p300/CBP HAT (histone acetyltransferase)	 * stereochemistry <table><tr><td>R1</td><td>R2</td></tr><tr><td>H</td><td>H</td></tr><tr><td>CN</td><td>H</td></tr><tr><td>H</td><td>H</td></tr><tr><td>CONH2</td><td>H</td></tr></table>	R1	R2	H	H	CN	H	H	H	CONH2	H
R1	R2										
H	H										
CN	H										
H	H										
CONH2	H										

TABLE 2-continued

A list of compounds that bind to regulator proteins.

Target protein	Compound
	R,S
	H
	R,S
	H
	OMe
	CH ₂ OH
	R,S
	H
	R,S
	cyclopropyl
	H
	H
	S
	H
	S
	H
	S
	H

p300/CBP HAT
(histone
acetyltransferase)





TABLE 2-continued

A list of compounds that bind to regulator proteins.

Target protein	Compound
p300/CBP HAT (histone acetyltransferase)	② indicates text missing or illegible when filed
	② indicates text missing or illegible when filed
	② indicates text missing or illegible when filed
	② indicates text missing or illegible when filed
p300/CBP HAT	② indicates text missing or illegible when filed
	② indicates text missing or illegible when filed
	② indicates text missing or illegible when filed
	② indicates text missing or illegible when filed

TABLE 2-continued





A list of compounds that bind to regulator proteins.	
Target protein	Compound
	<chem>CN1CCCCC1</chem> indicates text missing or illegible when filed
	<chem>CN1CCCCC1C2=CC=CC=C2</chem> indicates text missing or illegible when filed
	<chem>CN1CCCCC1[C@H](C2=CC=CC=C2)C3=CC=CC=C3</chem> indicates text missing or illegible when filed
	<chem>CN1CCCCC1C2=CC=CC=C2C3=CC=CC=C3</chem> indicates text missing or illegible when filed

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
Br	 ② indicates text missing or illegible when filed
Br	 ② indicates text missing or illegible when filed
OGT	
OGT	

TABLE 2-continued

A list of compounds that bind to regulator proteins.








Target protein	Compound
OGT	 <chem>O=C1OC(=O)N1C2=CC=C(Cl)C=C2OC(=O)c3ccccc3</chem>
	 <chem>N#CCCC1=CC=C(C=C1)SC2=NC(=C3C(=C2)C(=CC4CCCCC4)N=C3C5=CC=CC=C5S5)C(=O)O</chem>
	 <chem>CC1=CC=C2C(=C1)C(=CC(=C2)S(=O)(=O)N3C(C)C(=CC=C3)C4=CC=CC=C4OCC)C(=O)N5C=CC=CC=C5</chem>
	 <chem>O=C1C(=CC2=CC=CC=C2N1C(=O)C3=CC=CC=C3)S(=O)(=O)NC(C(=O)Nc4ccccc4)C5=CC=CC=C5</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.



OGT

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
OGT	  R_1 is H or C₁₋₆ alkyl; R_2 is H or C₁₋₆ alkyl; R_3 is H or C₁₋₆ alkyl

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
OGT	  

TABLE 2-continued




A list of compounds that bind to regulator proteins.	
Target protein	Compound
LFA-1/ICAM-1	
	
LFA-1/ICAM-1	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	 <chem>Cc1ccc2c(c1)c(c[nH]2)C(=O)NCc3ccccc3</chem>
LFA-1/ICAM-1	 <chem>CCN(CC)CCc1ccc(NC2=NC(C3=CC=C(C=C3)OCC)C=C2C4=CC=CC=C4Cl)cc1Cl</chem>
	 <chem>Oc1ncnc(N1C2=CC=C(C=C2)SC3=CC=C(C=C3)[N+](=O)[O-])c1</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
LFA-1/ICAM-1	   

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
LFA-1/ICAM-1	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	
Methyllysine binding/ L3MBTL1	 

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	
	
	
	UNC280
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	
	UNC669
	
	UNC926
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	
	
	
	
	8: R = H 9: R = Me

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
Methyllysine binding/ L3MBTL3	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
Methyllysine binding/ L3MBTL3	 UNC1215
	 UNC1679
	 UNC1679
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
	
	
	
	
Methyllysine binding/ L3MBTL3	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
Methyllysine binding/ L3MTL3	 <chem>CN(C)CCCCNC(=O)c1ccc(Br)cc1</chem>
	 <chem>CN(C)CCCCNC(=O)c1ccc(R)cc1</chem>
	 <chem>CN1CCCC1COc2cc3c(nc4c3ccccc45C6CC7C(C6)CC7)nn5CO</chem>
	 <chem>CN(C)CCCCOc1cc2c(nc3c1ccccc34)nn4C5C(CN(C)C)CC5</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
Chromodomain	
	

TABLE 2-continued



A list of compounds that bind to regulator proteins.	
Target protein	Compound
Chromodomain	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
	 <chem>COc1ccc(cc1)C(=O)N[C@@H](Cc2ccccc2)C(=O)N[C@@H](C3CCCC3)C(=O)N[C@@H](CCO)C(=O)NCCCC[N+](C)(C)C</chem>
	 <chem>CO[C@H](O)C(=O)N[C@@H](C(C)C)C(=O)N[C@@H](Cc1ccccc1)C(=O)N[C@@H](C(C)C)C(=O)N[C@@H](Cc2ccccc2)C(=O)NCCCCN(CC)CC</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	
	
Chromodomain	

TABLE 2-continued

A list of compounds that bind to regulator proteins.


Target protein	Compound
Chromodomain	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	
Chromodomain	 

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	
	
	

TABLE 2-continued







A list of compounds that bind to regulator proteins.	
Target protein	Compound
Chromodomain	
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	 <chem>N#Cc1ccccc1C(=O)Nc2ccccc2</chem>
	 <chem>COc1ccc2c(c1)CC[C@H]3[C@@H](C[C@H]4[C@@H](C[C@@H](C4)CN(C)C[C@H]5C[C@@H](C[C@H](C5)C[C@@H](C[C@H]6C[C@@H](C[C@H](C6)COC)CC7=CC=CC=C7C)C)C)C2</chem>
	 <chem>COC1=CC=C(C=C1)CC2CCNCC2</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
Chromodomain/ CBX7	
Chromodomain	
Chromodomain	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
Chromodomain	 <chem>CC1(CCCC1)C(=O)N[C@@H](CO)C(=O)N[C@@H](CCCC[N+](C)(C)C)C(=O)N[C@@H](Cc2ccccc2)C(=O)N</chem>
	 <chem>CC1(CCCC1[NH])C(=O)N[C@@H](CO)C(=O)N[C@@H](CCCC[N+](C)(C)C)C(=O)N[C@@H](Cc2ccccc2)C(=O)N[C@@H](C(C)C)C(=O)N</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
Chromodomain	
	

TABLE 2-continued



A list of compounds that bind to regulator proteins.	
Target protein	Compound
Methyl transferase	
	EPZ004777 (ref. 21), EPZ-5676 (ref. 24), SGC0946 (ref. 86)
	EZH2 GSK126 (ref. 37), GSK343 (refs 87, 88), EPZ005687 (ref. 38), EPZ-6438 (ref. 44), EII (ref. 39), UNC1999 (ref. 89)
	G9A BIX01294 (ref. 90), UNC0321 (ref. 91), UNC0638 (ref. 92), NC0642 (ref. 88), BRD4770 (ref. 93)
	PRMT3 14a (ref. 94)
	PRMT4 (CARM1) 17b (Bristol-Myers Squibb) (rets 95, 96), Methylgene (ref. 97)
Methyl transferase	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	 <chem>Nc1nc2c(ncn2C1SC[C@H](N)C(=O)O)c3cc(Br)cn3</chem>
Methyl transferase	 <chem>Nc1nc2nc3c(ncn2c1O[C@H]3C[C@@H](NCCNCCc4ccc(Cl)cc4)CC[C@H](N)C(=O)O)O</chem>
	 <chem>Nc1nc2c(ncn2[C@H]1C[C@H](O)C[C@@H](O)CO</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
Methyl transferase	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
	
Methyl transferase	

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
Chromodomain	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.

Target protein	Chromodomain	Compound

TABLE 2-continued

A list of compounds that bind to regulator proteins.







Target protein	Compound
	 <chem>COc1ccc(cc1)C(=O)O</chem>
	 <chem>CC(C)CC(C(=O)O)N</chem>
	 <chem>CC1(CCN1)C(=O)O</chem>
	 <chem>OC[C@H](C(=O)O)N</chem>
	 <chem>C1CCNCC1C(=O)O</chem>
	 <chem>BrC1=CC=C(C=C1)C(=O)O</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	 <chem>N[C@@H](Cc1ccccc1)C(=O)O</chem>
	 <chem>N[C@@H](Cc1cc(F)c(F)c(F)c1)C(=O)O</chem>
	 <chem>CC(N)C(=O)O</chem>
	 <chem>CCN(CC)CCCC(N)C(=O)O</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
	 <chem>CCN(CC)CCCCC(N)C(=O)O</chem>
	 <chem>CC(C(C(N)=O)O)C(O)C(N)=O</chem>
	 <chem>CC1(CCN1C(=O)O)C(O)C</chem>
	 <chem>NC1(CCN1C(=O)O)CCCCN</chem>
	 <chem>c1ccoc1CC(=O)O</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	 <chem>CCN(CC)CCCC(N)C(=O)O</chem>
	 <chem>CN(C)CCCC(N)C(=O)O</chem>
	 <chem>CC(C)C(N)C(=O)O</chem>
	 <chem>CC(C)C(N)C(=O)O</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
Chromodomain	 The chemical structure shows a diethylamino group (N(CH2CH3)2) connected via a four-carbon chain to a cyclohexyl ring. This cyclohexyl ring is part of a larger molecule that includes a benzyl group and a linker connected to a bead (represented by a shaded sphere). The molecule is a complex amide derivative.
	 This is a second chemical structure, very similar to the first one, also featuring a diethylamino group, a cyclohexyl ring, a benzyl group, and a linker to a bead. The stereochemistry and connectivity of the amide bonds differ slightly from the first structure.

TABLE 2-continued

A list of compounds that bind to regulator proteins.

Target protein	Compound
	
	

4 redundant hits

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
	 4 redundant hits
	 4 redundant hits

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	 <chem>CC(C)C(=O)NC[C@H](c1cc(F)c(F)c(F)c1)C(=O)N[C@@H](CO)C(=O)N[C@@H](CCCCN(C)C)C(=O)N[C@@H](CO)C(=O)NLinker</chem>
	4 redundant hits
	 <chem>CC(C)[C@H](C(=O)N[C@@H](Cc1ccccc1)C(=O)N[C@@H](Cc2ccncc2)C(=O)N)C(=O)N[C@@H](CCCCN(CC)CC)C(=O)N[C@@H](CO)C(=O)NLinker</chem>
	4 redundant hits

TABLE 2-continued

A list of compounds that bind to regulator proteins.


Target protein	Compound
Chromodomain	a)



UNC4979

TABLE 2-continued

A list of compounds that bind to regulator proteins.

Target protein	Compound
	 The chemical structure of UNC4980 is a complex molecule. It features a central chain of amide bonds. On the left, there is a benzamide group. Moving right, there is a chiral center with a phenyl group and a methyl group. This is followed by another chiral center with a hydroxyl group and a methyl group. Further right is a chiral center with a phenyl group and a methyl group. The chain continues with a chiral center having a hydroxyl group and a methyl group, and finally a chiral center with a phenyl group and a methyl group. The rightmost part of the molecule includes a triazole ring system connected to a chain ending in an amide group with an NH₂ group. A circled question mark (?) is located near the end of the chain.

UNC4980

TABLE 2-continued

A list of compounds that bind to regulator proteins.


Target protein	Compound
	 The chemical structure of UNC4981 is a complex molecule. It features a central core with several side chains. On the left, there is a side chain with a phenyl ring and a methyl group. In the middle, there is a side chain with a cyclohexyl ring and a hydroxyl group. On the right, there is a side chain with a pyrazole ring and a terminal amine group. The molecule is labeled UNC4981.

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
	
Chromodomain	 UNC4982

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
	 <chem>COc1ccc(cc1)C(=O)O</chem>
	 <chem>COc1ccc(cc1)C(=O)O</chem>
	 <chem>FC(F)(F)c1ccc(cc1)C(=O)O</chem>
	 <chem>CC(C)(C)c1ccc(cc1)C(=O)O</chem>
	 <chem>Brc1ccncc1C(=O)O</chem> Targeting extended β -groove
	CpG -1

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
Kme3 S	
	
	
	
Chromodomain	 Targeting Aromatic cage
	
	R =

TABLE 2-continued

A list of compounds that bind to regulator proteins.







Target protein	Compound
	 <chem>O=C(c1ccc(Br)cc1)C#N</chem>
	 <chem>O=C(c1cccc(Br)c1)C#N</chem>
	 <chem>O=C(c1ccccc1Br)C#N</chem>
	 <chem>O=C(c1ccc(OC)cc1)C#N</chem>
	 <chem>O=C(c1cccc(OC)c1)C#N</chem>
	 <chem>O=C(c1ccccc1OC)C#N</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	 <chem>O=C(Cl)c1ccc(I)cc1</chem>
	 <chem>O=C(Cl)c1ccc(C(F)(F)F)cc1</chem>
	 <chem>O=C(Cl)c1ccc(C(C)(C)C)cc1</chem>
	 <chem>O=C(Cc1ccncc1Br)</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
Chormodonain	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
Chormodonain	a)  Chemical structure of UNC3866, a complex molecule featuring a central amide backbone with various side chains including a benzyl group, a methyl group, a hydroxymethyl group, and a diethylamino group. b  Chemical structure of UNC3567 (1), a complex molecule featuring a central amide backbone with various side chains including a benzyl group, a methyl group, a hydroxymethyl group, and a diethylamino group. UNC3866 UNC3567 (1)

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
Chormodomain	 UNC4219 (3)
Chormodomain	 UNC4195 (4)

TABLE 2-continued






A list of compounds that bind to regulator proteins.	
Target protein	Compound
Methyl lysine binding domain	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	 <chem>OC(=O)c1ccc(cc1)C(=O)N2CCCC2</chem>
	 <chem>CC(=O)c1ccc(cc1)C(=O)N2CCCC2</chem>
	 <chem>Brc1ccc(cc1)C(=O)N2CCCC2</chem>
	 <chem>Brc1ccc(cc1)C(=O)N2CCCCC2</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	 <chem>BrC1=CC=C(C=C1)C(=O)NCCN2CCCC2</chem>
	 <chem>BrC1=CC=C(C=C1)C(=O)N2CCCCCN2CN3CCCC3</chem>
	 <chem>BrC1=CC=C(C=C1)C(=O)N2C3CC4C(C2)C1CC4CN3C5CCCC5</chem>
	 <chem>BrC1=CC=C(C=C1)C(=O)N2CCCCCN2CN3CCCC3</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
Methyl lysine binding domain	 UNC1215
	 UNC2533 (1)
	 UNC669
	 UNC1079

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	
	UNC1215
Methyl lysine binding domain	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.







Target protein	Compound
	
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.








Target protein	Compound
	
	
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.








Target protein	Compound
	
	
	
Methyl lysine binding domain	   

TABLE 2-continued

A list of compounds that bind to regulator proteins.








Target protein	Compound
Methyl lysine binding domain	
	
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	
	14
	
Methyl lysine binding domain	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
Methyl lysine binding domain	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
Methyl lysine binding domain	
	Ar
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	 <chem>CC(=O)c1ccc2ccccc2c1C(=O)C</chem>
	 <chem>CC(=O)c1cc(s1)C(=O)C</chem>
	 <chem>CC(=O)c1cc(s1)C(=O)C</chem>
	 <chem>CC(=O)c1ccc(Nc2ccccc2)cc1C(=O)C</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
Methyl lysine binding domain	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
Methyl lysine binding domain	 <chem>BrC1=CC=C(C=C1)C(=O)N2CCCC2C3CCCC3</chem>
	 <chem>CCN1CCCC1C(=O)Nc2ccc(cc2)C(=O)N3CCCC3</chem>
	 <chem>CCN1CCCC1C(=O)Nc2ccc(cc2)C(=O)N3C4C5C6C7C8C9C10C11C12C13C14C15C16C17C18C19C20C21C22C23C24C25C26C27C28C29C30C31C32C33C34C35C36C37C38C39C40C41C42C43C44C45C46C47C48C49C50C51C52C53C54C55C56C57C58C59C60C61C62C63C64C65C66C67C68C69C70C71C72C73C74C75C76C77C78C79C80C81C82C83C84C85C86C87C88C89C90C91C92C93C94C95C96C97C98C99C100C101C102C103C104C105C106C107C108C109C110C111C112C113C114C115C116C117C118C119C120C121C122C123C124C125C126C127C128C129C130C131C132C133C134C135C136C137C138C139C140C141C142C143C144C145C146C147C148C149C150C151C152C153C154C155C156C157C158C159C160C161C162C163C164C165C166C167C168C169C170C171C172C173C174C175C176C177C178C179C180C181C182C183C184C185C186C187C188C189C190C191C192C193C194C195C196C197C198C199C200C201C202C203C204C205C206C207C208C209C210C211C212C213C214C215C216C217C218C219C220C221C222C223C224C225C226C227C228C229C230C231C232C233C234C235C236C237C238C239C240C241C242C243C244C245C246C247C248C249C250C251C252C253C254C255C256C257C258C259C260C261C262C263C264C265C266C267C268C269C270C271C272C273C274C275C276C277C278C279C280C281C282C283C284C285C286C287C288C289C290C291C292C293C294C295C296C297C298C299C300C301C302C303C304C305C306C307C308C309C310C311C312C313C314C315C316C317C318C319C320C321C322C323C324C325C326C327C328C329C330C331C332C333C334C335C336C337C338C339C340C341C342C343C344C345C346C347C348C349C350C351C352C353C354C355C356C357C358C359C360C361C362C363C364C365C366C367C368C369C370C371C372C373C374C375C376C377C378C379C380C381C382C383C384C385C386C387C388C389C390C391C392C393C394C395C396C397C398C399C400C401C402C403C404C405C406C407C408C409C410C411C412C413C414C415C416C417C418C419C420C421C422C423C424C425C426C427C428C429C430C431C432C433C434C435C436C437C438C439C440C441C442C443C444C445C446C447C448C449C450C451C452C453C454C455C456C457C458C459C460C461C462C463C464C465C466C467C468C469C470C471C472C473C474C475C476C477C478C479C480C481C482C483C484C485C486C487C488C489C490C491C492C493C494C495C496C497C498C499C500C501C502C503C504C505C506C507C508C509C510C511C512C513C514C515C516C517C518C519C520C521C522C523C524C525C526C527C528C529C530C531C532C533C534C535C536C537C538C539C540C541C542C543C544C545C546C547C548C549C550C551C552C553C554C555C556C557C558C559C560C561C562C563C564C565C566C567C568C569C570C571C572C573C574C575C576C577C578C579C580C581C582C583C584C585C586C587C588C589C590C591C592C593C594C595C596C597C598C599C600C601C602C603C604C605C606C607C608C609C610C611C612C613C614C615C616C617C618C619C620C621C622C623C624C625C626C627C628C629C630C631C632C633C634C635C636C637C638C639C640C641C642C643C644C645C646C647C648C649C650C651C652C653C654C655C656C657C658C659C660C661C662C663C664C665C666C667C668C669C670C671C672C673C674C675C676C677C678C679C680C681C682C683C684C685C686C687C688C689C690C691C692C693C694C695C696C697C698C699C700C701C702C703C704C705C706C707C708C709C710C711C712C713C714C715C716C717C718C719C720C721C722C723C724C725C726C727C728C729C730C731C732C733C734C735C736C737C738C739C740C741C742C743C744C745C746C747C748C749C750C751C752C753C754C755C756C757C758C759C760C761C762C763C764C765C766C767C768C769C770C771C772C773C774C775C776C777C778C779C780C781C782C783C784C785C786C787C788C789C790C791C792C793C794C795C796C797C798C799C800C801C802C803C804C805C806C807C808C809C810C811C812C813C814C815C816C817C818C819C820C821C822C823C824C825C826C827C828C829C830C831C832C833C834C835C836C837C838C839C840C841C842C843C844C845C846C847C848C849C850C851C852C853C854C855C856C857C858C859C860C861C862C863C864C865C866C867C868C869C870C871C872C873C874C875C876C877C878C879C880C881C882C883C884C885C886C887C888C889C890C891C892C893C894C895C896C897C898C899C900C901C902C903C904C905C906C907C908C909C910C911C912C913C914C915C916C917C918C919C920C921C922C923C924C925C926C927C928C929C930C931C932C933C934C935C936C937C938C939C940C941C942C943C944C945C946C947C948C949C950C951C952C953C954C955C956C957C958C959C960C961C962C963C964C965C966C967C968C969C970C971C972C973C974C975C976C977C978C979C980C981C982C983C984C985C986C987C988C989C990C991C992C993C994C995C996C997C998C999C1000C1001C1002C1003C1004C1005C1006C1007C1008C1009C1010C1011C1012C1013C1014C1015C1016C1017C1018C1019C1020C1021C1022C1023C1024C1025C1026C1027C1028C1029C1030C1031C1032C1033C1034C1035C1036C1037C1038C1039C1040C1041C1042C1043C1044C1045C1046C1047C1048C1049C1050C1051C1052C1053C1054C1055C1056C1057C1058C1059C1060C1061C1062C1063C1064C1065C1066C1067C1068C1069C1070C1071C1072C1073C1074C1075C1076C1077C1078C1079C1080C1081C1082C1083C1084C1085C1086C1087C1088C1089C1090C1091C1092C1093C1094C1095C1096C1097C1098C1099C1100C1101C1102C1103C1104C1105C1106C1107C1108C1109C1110C1111C1112C1113C1114C1115C1116C1117C1118C1119C1120C1121C1122C1123C1124C1125C1126C1127C1128C1129C1130C1131C1132C1133C1134C1135C1136C1137C1138C1139C1140C1141C1142C1143C1144C1145C1146C1147C1148C1149C1150C1151C1152C1153C1154C1155C1156C1157C1158C1159C1160C1161C1162C1163C1164C1165C1166C1167C1168C1169C1170C1171C1172C1173C1174C1175C1176C1177C1178C1179C1180C1181C1182C1183C1184C1185C1186C1187C1188C1189C1190C1191C1192C1193C1194C1195C1196C1197C1198C1199C1200C1201C1202C1203C1204C1205C1206C1207C1208C1209C1210C1211C1212C1213C1214C1215C1216C1217C1218C1219C1220C1221C1222C1223C1224C1225C1226C1227C1228C1229C1230C1231C1232C1233C1234C1235C1236C1237C1238C1239C1240C1241C1242C1243C1244C1245C1246C1247C1248C1249C1250C1251C1252C1253C1254C1255C1256C1257C1258C1259C1260C1261C1262C1263C1264C1265C1266C1267C1268C1269C1270C1271C1272C1273C1274C1275C1276C1277C1278C1279C1280C1281C1282C1283C1284C1285C1286C1287C1288C1289C1290C1291C1292C1293C1294C1295C1296C1297C1298C1299C1300C1301C1302C1303C1304C1305C1306C1307C1308C1309C1310C1311C1312C1313C1314C1315C1316C1317C1318C1319C1320C1321C1322C1323C1324C1325C1326C1327C1328C1329C1330C1331C1332C1333C1334C1335C1336C1337C1338C1339C1340C1341C1342C1343C1344C1345C1346C1347C1348C1349C1350C1351C1352C1353C1354C1355C1356C1357C1358C1359C1360C1361C1362C1363C1364C1365C1366C1367C1368C1369C1370C1371C1372C1373C1374C1375C1376C1377C1378C1379C1380C1381C1382C1383C1384C1385C1386C1387C1388C1389C1390C1391C1392C1393C1394C1395C1396C1397C1398C1399C1400C1401C1402C1403C1404C1405C1406C1407C1408C1409C1410C1411C1412C1413C1414C1415C1416C1417C1418C1419C1420C1421C1422C1423C1424C1425C1426C1427C1428C1429C1430C1431C1432C1433C1434C1435C1436C1437C1438C1439C1440C1441C1442C1443C1444C1445C1446C1447C1448C1449C1450C1451C1452C1453C1454C1455C1456C1457C1458C1459C1460C1461C1462C1463C1464C1465C1466C1467C1468C1469C1470C1471C1472C1473C1474C1475C1476C1477C1478C1479C1480C1481C1482C1483C1484C1485C1486C1487C1488C1489C1490C1491C1492C1493C1494C1495C1496C1497C1498C1499C1500C1501C1502C1503C1504C1505C1506C1507C1508C1509C1510C1511C1512C1513C1514C1515C1516C1517C1518C1519C1520C1521C1522C1523C1524C1525C1526C1527C1528C1529C1530C1531C1532C1533C1534C1535C1536C1537C1538C1539C1540C1541C1542C1543C1544C1545C1546C1547C1548C1549C1550C1551C1552C1553C1554C1555C1556C1557C1558C1559C1560C1561C1562C1563C1564C1565C1566C1567C1568C1569C1570C1571C1572C1573C1574C1575C1576C1577C1578C1579C1580C1581C1582C1583C1584C1585C1586C1587C1588C1589C1590C1591C1592C1593C1594C1595C1596C1597C1598C1599C1600C1601C1602C1603C1604C1605C1606C1607C1608C1609C1610C1611C1612C1613C1614C1615C1616C1617C1618C1619C1620C1621C1622C1623C1624C1625C1626C1627C1628C1629C1630C1631C1632C1633C1634C1635C1636C1637C1638C1639C1640C1641C1642C1643C1644C1645C1646C1647C1648C1649C1650C1651C1652C1653C1654C1655C1656C1657C1658C1659C1660C1661C1662C1663C1664C1665C1666C1667C1668C1669C1670C1671C1672C1673C1674C1675C1676C1677C1678C1679C1680C1681C1682C1683C1684C1685C1686C1687C1688C1689C1690C1691C1692C1693C1694C1695C1696C1697C1698C1699C1700C1701C1702C1703C1704C1705C1706C1707C1708C1709C1710C1711C1712C1713C1714C1715C1716C1717C1718C1719C1720C1721C1722C1723C1724C1725C1726C1727C1728C1729C1730C1731C1732C1733C1734C1735C1736C1737C1738C1739C1740C1741C1742C1743C1744C1745C1746C1747C1748C1749C1750C1751C1752C1753C1754C1755C1756C1757C1758C1759C1760C1761C1762C1763C1764C1765C1766C1767C1768C1769C1770C1771C1772C1773C1774C1775C1776C1777C1778C1779C1780C1781C1782C1783C1784C1785C1786C1787C1788C1789C1790C1791C1792C1793C1794C1795C1796C1797C1798C1799C1800C1801C1802C1803C1804C1805C1806C1807C1808C1809C1810C1811C1812C1813C1814C1815C1816C1817C1818C1819C1820C1821C1822C1823C1824C1825C1826C1827C1828C1829C1830C1831C1832C1833C1834C1835C1836C1837C1838C1839C1840C1841C1842C1843C1844C1845C1846C1847C1848C1849C1850C1851C1852C1853C1854C1855C1856C1857C1858C1859C1860C1861C1862C1863C1864C1865C1866C1867C1868C1869C1870C1871C1872C1873C1874C1875C1876C1877C1878C1879C1880C1881C1882C1883C1884C1885C1886C1887C1888C1889C1890C1891C1892C1893C1894C1895C1896C1897C1898C1899C1900C1901C1902C1903C1904C1905C1906C1907C1908C1909C1910C1911C1912C1913C1914C1915C1916C1917C1918C1919C1920C1921C1922C1923C1924C1925C1926C1927C1928C1929C1930C1931C1932C1933C1934C1935C1936C1937C1938C1939C1940C1941C1942C1943C1944C1945C1946C1947C1948C1949C1950C1951C1952C1953C1954C1955C1956C1957C1958C1959C1960C1961C1962C1963C1964C1965C1966C1967C1968C1969C1970C1971C1972C1973C1974C1975C1976C1977C1978C1979C1980C1981C1982C1983C1984C1985C1986C1987C1988C1989C1990C1991C1992C1993C1994C1995C1996C1997C1998C1999C2000C2001C2002C2003C2004C2005C2006C2007C2008C2009C2010C2011C2012C2013C2014C2015C2016C2017C2018C2019C2020C2021C2022C2023C2024C2025C2026C2027C2028C2029C2030C2031C2032C2033C2034C2035C2036C2037C2038C2039C2040C2041C2042C2043C2044C2045C2046C2047C2048C2049C2050C2051C2052C2053C2054C2055C2056C2057C2058C2059C2060C2061C2062C2063C2064C2065C2066C2067C2068C2069C2070C2071C2072C2073C2074C2075C2076C2077C2078C2079C2080C2081C2082C2083C2084C2085C2086C2087C2088C2089C2090C2091C2092C2093C2094C2095C2096C2097C2098C2099C2100C2101C2102C2103C2104C2105C2106C2107C2108C2109C2110C2111C2112C2113C2114C2115C2116C2117C2118C2119C2120C2121C2122C2123C2124C2125C2126C2127C2128C2129C2130C2131C2132C2133C2134C2135C2136C2137C2138C2139C2140C2141C2142C2143C2144C2145C2146C2147C2148C2149C2150C2151C2152C2153C2154C2155C2156C2157C2158C2159C2160C2161C2162C2163C2164C2165C2166C2167C2168C2169C2170C2171C2172C2173C2174C2175C2176C2177C2178C2179C2180C2181C2182C2183C2184C2185C2186C2187C2188C2189C2190C2191C2192C2193C2194C2195C2196C2197C2198C2199C2200C2201C2202C2203C2204C2205C2206C2207C2208C2209C2210C2211C2212C2213C2214C2215C2216C2217C2218C2219C2220C2221C2222C2223C2224C2225C2226C2227C2228C2229C2230C2231C2232C2233C2234C2235C2236C2237C2238C2239C2240C2241C2242C2243C2244C2245C2246C2247C2248C2249C2250C2251C2252C2253C2254C2255C2256C2257C2258C2259C2260C2261C2262C2263C2264C2265C2266C2267C2268C2269C2270C2271C2272C2273C2274C2275C2276C2277C2278C2279C2280C2281C2282C2283C2284C2285C2286C2287C2288C2289C2290C2291C2292C2293C2294C2295C2296C2297C2298C2299C2300C2301C2302C2303C2304C2305C2306C2307C2308C2309C2310C2311C2312C2313C2314C2315C2316C2317C2318C2319C2320C2321C2322C2323C2324C2325C2326C2327C2328C2329C2330C2331C2332C2333C2334C2335C2336C2337C2338C2339C2340C2341C2342C2343C2344C2345C2346C2347C2348C2349C2350C2351C2352C2353C2354C2355C2356C2357C2358C2359C2360C2361C2362C2363C2364C2365C2366C2367C2368C2369C2370C2371C2372C2373C2374C2375C2376C2377C2378C2379C2380C2381C2382C2383C2384C2385C2386C2387C2388C2389C2390C2391C2392C2393C2394C2395C2396C2397C2398C2399C2400C2401C2402C2403C2404C2405C2406C2407C2408C2409C2410C2411C2412C2413C2414C2415C2416C2417C2418C2419C2420C2421C2422C2423C2424C2425C2426C2427C2428C2429C2430C2431C2432C2433C2434C2435C2436C2437C2438C2439C2440C2441C2442C2443C2444C2445C2446C2447C2448C2449C2450C2451C2452C2453C2454C2455C2456C2457C2458C2459C2460C2461C2462C2463C2464C2465C2466C2467C2468C2469C2470C2471C2472C2473C2474C2475C2476C2477C2478C2479C2480C2481C2482C2483C2484C2485C2486C2487C2488C2489C2490C2491C2492C2493C2494C2495C2496C2497C2498C2499C2500C2501C2502C2503C2504C2505C2506C2507C2508C2509C2510C2511C2512C2513C2514C2515C2516C2517C2518C2519C2520C2521C2522C2523C2524C2525C2526C2527C2528C2529C2530C2531C2532C2533C2534C2535C2536C2537C2538C2539C2540C2541C2542C2543C2544C2545C2546C2547C2548C2549C2550C2551C2552C2553C2554C2555C2556C2557C2558C2559C2560C2561C2562C2563C2564C2565C2566C2567C2568C2569C2570C2571C2572C2573C2574C2575C2576C2577C2578C2579C2580C2581C2582C2583C2584C2585</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.







Target protein	Compound
	
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	
	
	R'' H
	
	H H H
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	
Methyl lysine binding domain	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
	 Amiodarone HCl
	 Tegaserod maleate

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
Methyl lysine binding domain	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
	
	

TABLE 2-continued



A list of compounds that bind to regulator proteins.	
Target protein	Compound
Methyl lysine binding domain	 WAG-003 (n = 2, trimethyl)
	 WAG-004 (n = 2, dimethyl)

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
	 WAG-005 (n = 3, trimethyl)
	 WAG-006 (n = 3, dimethyl)

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
Methyl lysine binding domain	 IS1
	 IS2

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
	 <chem>CC1=CC=C(C=C1)CCNC(=O)CCC2=NC(=O)N(CCCO)C3=CC=CC=C23</chem> IS3
	 <chem>CC1=CC=C(C=C1)S(=O)(=O)N2CCN(CC2)C3CCCC3C4=CC=CC=C4NC(=O)CC5=CC=CC=C5S4</chem> IS5

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
	 IS12
	 IS15

TABLE 2-continued

A list of compounds that bind to regulator proteins.









Target protein	Compound
	
	
Methyl lysine binding domain	
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
	
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
Methyl lysine binding domain	
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.











Target protein	Compound
	 <chem>Fc1ccccc1C(F)(F)F</chem>
	 <chem>Brc1ccccc1</chem>
	 <chem>Fc1ccc(cc1)C(F)(F)F</chem>
	 <chem>Clc1ccccc1</chem>
	 <chem>[O-][N+](=O)c1ccccc1</chem>
	 <chem>Fc1ccccc1</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.

Target protein	Compound
	
	
	
Methyl lysine binding domain	

group 1-3

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
	(b)
	
group 4	
Methyl lysine binding domain	

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
	 MM-401
	 OICR-9429

TABLE 2-continued



A list of compounds that bind to regulator proteins.	
Target protein	Compound
	
Methyl lysine binding domain	 WDR5-47 18 R₁ = 3-COOH—Ph 19 R₁ = 4-COOH—Ph 20 R₁ = 4-CN—Ph 21 R₁ = —Ph 22 R₁ = 4-F—Ph 23 R₁ = 4-Pyridinyl 24 R₁ = 5-Pyridinyl 25 R₁ = 4-NO₂—Ph 26 R₁ = 4-NH₂—Ph h




TABLE 2-continued

A list of compounds that bind to regulator proteins.

Target protein	Compound
	$R_2 = \text{---H}$ $29 R_2 = 4\text{-COOH---Ph}$ $R_1 = \text{---H}$ $R_2 = \text{---H}$ $30 R_2 = 4\text{-Pyridinyl}$ $R_1 = \text{---H}$ $R_2 = \text{---H}$ $31 R_2 = 4\text{-NO}_2\text{---Ph}$ $R_1 = \text{---H}$ $R_2 = \text{---H}$ $32 R_2 = 4\text{-NH}_2\text{---Ph}$ $R_1 = \text{---H}$ h $R_2 = \text{---H}$ $R_2 = \text{---H}$ $R_2 = \text{---H}$ $R_2 = \text{---H}$ $R_2 = \text{---H}$ $R_2 = \text{---H}$
Methyl lysine binding domain	

TABLE 2-continued

A list of compounds that bind to regulator proteins.

Target protein	Compound
	
	33 R = 4-fluoro 34 R = 4-methoxyl 35 R = 3,4-dimethoxyl 36 R = 2,4,6-trimethyl
Methyl lysine binding domain	 

A list of compounds that bind to regulator proteins.



Target protein	Compound
Methyl lysine binding domain	 37a R = 4-fluoro-2-chloro-3-methyl 38a R = 3-methoxy 39a R = 2,4-difluoro 40a R = 2-chloro 37 R = 4-fluoro-2-chloro-3-methyl 38 R = 3-methoxy 39 R = 2,4-difluoro 40 R = 2-chloro
Methyl lysine binding domain	 X $\begin{matrix} \text{R}_4 \\ \text{—NHSO}_2\text{—} \\ \text{—NHSO}_2\text{—} \\ \text{—NHSO}_2\text{—} \end{matrix}$ $\begin{matrix} \text{R}_4 \\ \text{—4-fluoro} \\ \text{—4-methoxy} \\ \text{—3,4-dimethoxy} \end{matrix}$

TABLE 2-continued

A list of compounds that bind to regulator proteins.	
Target protein	Compound

- NHSO₂— 2,4,6-trimethyl
- CONH— 4-fluoro-2-chloro-3-methyl
- CONH— 3-methoxyl
- CONH— 2,4-difluoro
- CONH— 2-chloro
- NHSO₂— 4-fluoro-2-chloro-3-methyl

Methyl lysine
binding domain




TABLE 2-continued

A list of compounds that bind to regulator proteins.

Target protein	Compound
	$R = -CH_3$
	$R = -CH_3$ $R = -CH(CH_3)_2$ $R = -CH_2CH_2CH_3$ $R = -CH_2NH-Boc$ $R = -CH(C_3)NH-Boc$ $R = -CH_2CH_2NH-Boc$ $R = -C(CH_3)_2NH-Boc$
	$R = -(CH_2)_3NH-Boc$ $R = -CH_2CH(CH_3)_2$

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
Methyl lysine binding domain	 R =
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound																														
																															
																															
																															
Methyl lysine binding domain																															
	<table><tr><td>R₁</td><td>R₂</td></tr><tr><td>—Ph</td><td>—H</td></tr><tr><td>4-Pyridyl</td><td>—H</td></tr><tr><td>4-NH₂—Ph</td><td>—H</td></tr><tr><td>—Ph</td><td>—NO₂</td></tr><tr><td>4-NO₂—Ph</td><td>—NHCOCH₃</td></tr><tr><td>4-Pyridyl</td><td>—NO₂</td></tr><tr><td>4-COOCH₃—Ph</td><td>—NO₂</td></tr><tr><td>—Ph</td><td>—NH₂</td></tr><tr><td>4-Pyridyl</td><td>—NH₂</td></tr><tr><td>4-COOCH₃—Ph</td><td>—NH₂</td></tr><tr><td>4-NH₂—Ph</td><td>—NHCOCH₃</td></tr><tr><td>4-Pyridyl</td><td>—NHCOCH₃</td></tr><tr><td>4-NO₂—Ph</td><td>—NO₂</td></tr><tr><td>4-COOCH₃—Ph</td><td>—NH₂</td></tr></table>	R ₁	R ₂	—Ph	—H	4-Pyridyl	—H	4-NH ₂ —Ph	—H	—Ph	—NO ₂	4-NO ₂ —Ph	—NHCOCH ₃	4-Pyridyl	—NO ₂	4-COOCH ₃ —Ph	—NO ₂	—Ph	—NH ₂	4-Pyridyl	—NH ₂	4-COOCH ₃ —Ph	—NH ₂	4-NH ₂ —Ph	—NHCOCH ₃	4-Pyridyl	—NHCOCH ₃	4-NO ₂ —Ph	—NO ₂	4-COOCH ₃ —Ph	—NH ₂
R ₁	R ₂																														
—Ph	—H																														
4-Pyridyl	—H																														
4-NH ₂ —Ph	—H																														
—Ph	—NO ₂																														
4-NO ₂ —Ph	—NHCOCH ₃																														
4-Pyridyl	—NO ₂																														
4-COOCH ₃ —Ph	—NO ₂																														
—Ph	—NH ₂																														
4-Pyridyl	—NH ₂																														
4-COOCH ₃ —Ph	—NH ₂																														
4-NH ₂ —Ph	—NHCOCH ₃																														
4-Pyridyl	—NHCOCH ₃																														
4-NO ₂ —Ph	—NO ₂																														
4-COOCH ₃ —Ph	—NH ₂																														

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
Methyl lysine binding domain	
Methyl lysine binding domain	 

TABLE 2-continued

A list of compounds that bind to regulator proteins.

Target protein	Compound
	$\text{—NHCO(CH}_2\text{)}_3\text{NH}_2$ $\text{—NHCO(CH}_2\text{)}_3\text{NHBoc}$ $\text{—NHCOCH}_2\text{CH(CH}_3\text{)}_2$
	—NHCOCH_3
	—NHCOPh $\text{—NHCOCH}_2\text{CH}_3$ $\text{—NHCOCH(CH}_3\text{)}_2$ $\text{—NHCOCH}_2\text{CH}_2\text{CH}_3$ $\text{—NHCOCH}_2\text{NH}_2$ $\text{—NHCOCH}_2\text{NHBoc}$ $\text{—NHCOCH(CH}_3\text{)NH}_2$ $\text{—NHCOCH(CH}_3\text{)NHBoc}$
Methyl lysine binding domain	

WDR5-0101

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
	 WDR5-0102
	 WDR5-0103

TABLE 2-continued

A list of compounds that bind to regulator proteins.

Target protein	Compound
Methyl lysine binding domain	 <chem>CN1CCN(CC1)C(=O)c2ccc(cc2)[N+](=O)[O-]</chem> <chem>CN1CCN(CC1)C(=O)c2ccc(cc2)Cc3ccc(cc3)C(=O)Nc4cc(C(F)(F)F)cc(=O)n4</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
Methyl lysine binding domain	 <chem>O=C(O)c1ccc2ccccc2c1[C@H]3C(=O)Oc4ccccc43</chem>
	 <chem>O=C(O)c1ccc2ccccc2c1[C@@H]3C(=O)Oc4ccccc43</chem>
	 <chem>CC(C)C(=O)N[C@@H](C)C(=O)N[C@H](C)C(=O)N[C@@H](C)C(=O)N[C@@H](Cc1ccccc1)C(=O)N2CCCC2</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
Methyl lysine binding domain	 <chem>CC(C)C(=O)N[C@@H](C[C@H](N)C(=O)N[C@@H](CCCCNC(=O)C1=CC=CC=C1)C(=O)N[C@@H](C)C(=O)N</chem>
	 <chem>R</chem>
	 <chem>Br</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	
	
Methyl lysine binding domain	
	R 2-CF₃, 5-F 2-CF₃, 4-OH 2-Cl, 4-CF₃ 2-Cl, 5-CF₃ 2-Cl, 5-Me 2-Cl, 6-F 3-CF₃, 4-OMe 3-Me, 5-Me

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
Methyl lysine binding domain	3-Me, 5-CF ₃
	3-Cl, 5-Cl
	3-OH, 5-CF ₃
	2-F, 5-SO ₂ NH ₂
	2-F, 3-F, 5-OH
	2-F, 3-Cl, 5-CF ₃
	2-Cl, 3-Me, 6-F
	2-F, 3-Me, 4-F
	2-Me, 3-F, 5-F
	3-Me, 4-F, 5-Me
	2-F, 3-Me, 5-Me, 6-F
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
	 <chem>Oc1ccc2ccccc2n1</chem>
	 <chem>Oc1ccc2cc(C)ccn2c1</chem>
	 <chem>c1ccc2nc3ccccc3n2c1</chem>
	 <chem>Fc1ccc2nc3ccccc3n2c1N</chem>
	 <chem>c1ccc2nc3ccccc3n2c1</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	
	
	
Methyl lysine binding domain	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	 <chem>CC(=O)Nc1ccc(cc1N)~</chem>
	 <chem>Nc1ccc(cc1N)~</chem>
	 <chem>Nc1ccc(cc1)~</chem>
	 <chem>C1CCN(C1)CCc2cc(ccn2)~</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.	
Target protein	Compound
Methyl lysine binding domain	
Methyl lysine binding domain	
	② X = N, R¹ = Me, R² = H, n = 1 X = N, R¹ = Me, R² = Me, n = 1 X = N, R¹ = Me, R² = H, n = 2 X = O, R² = H, n = 1

TABLE 2-continued


Target protein	Compound
Methyl lysine binding domain	 NO₂ R¹ 2-Cl-phenyl Me 3-Cl-phenyl 3-Me-phenyl 2-Cl-3-Me-phenyl 3-OH-phenyl 3-OMe-phenyl 4-F-phenyl 2-Cl-4-F-phenyl 3-Me-4-F-phenyl 3-OMe-4-phenyl 2-Cl-3-Me-4-F-phenyl phenyl cyclohexyl 1-naphthyl 5-quinolyl benzyl 3-pyridyl 2-furanyl R¹ = Me R¹ = 3-Cl-Ph R¹ = 3-Me-Ph R¹ = 2-Cl,3-Me-Ph R¹ = 3-OH-Ph R¹ = 3-OMe-Ph R¹ = 4-F-Ph R¹ = 2-Cl, 4-F-Ph R¹ = 3-Me,4-F-Ph 46 R¹ = 3-OMe, 4-F-Ph 47 R¹ = 2-Cl, 3-Me, 4-F-Ph 48 R¹ = phenyl 49 R¹ = cyclohexyl

TABLE 2-continued


A list of compounds that bind to regulator proteins.	
Target protein	Compound
Methyl lysine binding domain	50 R ¹ = 1-naphthyl
	51 R ¹ = 5-quinolyl
	52 R ¹ = benzyl
	53 R ¹ = 3-pyridyl
	54 R ¹ = 2-furanyl
	
	② indicates text missing or illegible when filed
	R ¹ = NO ₂
	R ¹ = NH ₂
	R ¹ = CO ₂ Me
	R ¹ = CO ₂ H
	R ¹ = CF ₃
	R ¹ = Br
	R ¹ = cyclopropyl
	R ¹ = 2-furanyl
	R ¹ = 4-pyridyl
	R ¹
	NO ₂
	CO ₂ Me
	CF ₃
	Br
	NH ₂
	CO ₂ H
	cyclopropyl
	2-furanyl
	4-pyridyl

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
Methyl lysine binding domain	 <chem>CN1CCN(C)CC1c2ccc(cc2)Nc3ccc(cc3)C(=O)NCC(=O)c4cc(C(F)(F)F)c(=O)[nH]4</chem>
CDK2	 <chem>Clc1ccc(cc1)Oc2cc(=O)oc3c(O)c(O)c(C4CCN(C)CC4)c32</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
CDK2	
CDK2	
CDK2	

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
CDK2	
CDK2	

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
CDK1, 2, or 4	
CDK2, CDK1, or CDK5	

TABLE 2-continued




A list of compounds that bind to regulator proteins.	
Target protein	Compound
CDK2, CDK4, CDK5, CDK1, CDK7	
CDK2, CDK1, CDK4	
CDK2, CDK4, CDK5, or CDK1	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
CDK2, CDK5, or CDK7	 <chem>CC(C)CC(=O)Nc1nc2c(c1)C(C)(C)N(C2)C(=O)C3CCN(C)CC3</chem>
CDK2 or CDK4	 <chem>COCCNS(=O)(=O)c1ccc(Nc2nc3c(nc2)C(C)N(C)C3)cc1</chem>
CDK2	 <chem>CC(C)c1nc2c(nc1)C(C)N(C)C2c3ccnc3Nc4ccc(cc4)N5CCN(CC5)C(=O)CO</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
CDK2 or CDK1	
CDK1, CDK2, CDK4 or CDK9	
CDK2	

TABLE 2-continued




A list of compounds that bind to regulator proteins.	
Target protein	Compound
CDK1 or CDK2	
CDK1 or CDK2	
CDK5 or GSK3beta	

TABLE 2-continued




A list of compounds that bind to regulator proteins.	
Target protein	Compound
CDK1, CDK5, or GSK3 alpha/beta	 <chem>CCCC1=Cc2ncnc2C1c3ccc(O)cc3</chem>
CDK4 or FLT3	 <chem>O=C1C(=O)c2cc(O)c(O)c2Oc3cc(O)c(O)cc3O1</chem>
CDK8	 <chem>CN1C=NC2=C(N1)N=CN=C2C3CCNC3C(=O)Nc4ccc(Cc5ccccc5N6CCOCC6)c(F)(F)F4</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
CDK8	
CDK8 or CDK19	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
CDK8	
CDK8	
CDK8	

TABLE 2-continued





A list of compounds that bind to regulator proteins.	
Target protein	Compound
CDK8 or CDK19	
CDK9	  

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
CDK7/9	
CDK9	
CDK12/13	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
CDK12	
CDK12/2	
CDK1/2/5/9 (Dinacelilb)	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
CDK9/4/1/2/6 (P276-00)	 <chem>CC1(C)CCN1[C@H](C2=CC(=C(C=C2)C3=C(C(=O)O)C(=C(C=C3)O)OC4=CC(=C(C=C4)C5=CC=CC=C5Cl)O)O</chem> HCl
CDK9 (voniclib)	 <chem>CC1(C)CCN1[C@H](C2=CC(=C(C=C2)C3=C(C(=O)O)C(=C(C=C3)O)OC4=CC(=C(C=C4)C5=CC(=CC=C5)C(=C(C=C5)C)C(F)(F)F)Cl)O)O</chem>
CDK1/2/4/5/9 (AT7519M)	 <chem>Clc1cc(Cl)ccc1C(=O)Nc2cc(C(=O)Nc3ccnnc3)cc2</chem>

TABLE 2-continued




A list of compounds that bind to regulator proteins.	
Target protein	Compound
CDK9/2/7/GSK3 alpha (SNS-032)	
CDK2	 SCH 727965
CDK1/2/4	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





TABLE 2-continued



A list of compounds that bind to regulator proteins.	
Target protein	Compound
CDK12/13 (THZ531)	
CDK9/2/7/GSK3 alpha	

TABLE 2-continued




Target protein	A list of compounds that bind to regulator proteins.
Compound	
CDK2 (roscovitine)	
CDK2 (NU2058)	
CDK2 (R457)	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
CDK2 (Flavopiridol)	 Flavopiridol
CDK1/2/4/5/7/9 (R547)	
H3K4 lysine methyltransferase KMT7 (PF1-2)	

TABLE 2-continued








A list of compounds that bind to regulator proteins.	
Target protein	Compound
H3K4 lysine methyltransferase KMT7 (cypiohepatadien)	
KDM1A/B (RN1)	
KDM1A (GSK2879552)	
KDM5 (CPI-455)	

TABLE 2-continued

A list of compounds that bind to regulator proteins.	
Target protein	Compound
KDM5 (KDM-C49)	
KDM5 (aminodarone)	
KDM5 (Disulfuram)	

Ⓢ indicates text missing or illegible when filed

TABLE 2-continued




A list of compounds that bind to regulator proteins.	
Target protein	Compound
EHMT2 aka G9a	 1 BIX-01294
	 2 UNC0638
	 3

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	
	4
	
	5
	
	6
	
	7

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
	
	
EHMT2 aka G9a	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.








Target protein	Compound
	
	
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	
	
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	
	
	
	
	

TABLE 2-continued




A list of compounds that bind to regulator proteins.	
Target protein	Compound
EHMT2 or GLP methyltransferase	
	(UNC0638)
G9a or HDAC	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.








Target protein	Compound
	
	
	Me Me
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.








Target protein	Compound
	
	
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.

Target protein	Compound
SMYD2	
DOT1L	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
DOT1L	 EPZ015666 (GSK3235025) PRMT5
PRMT5	 Methylstat pan-jmjC
Pan-jmjC	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
JMJD3/UTX/JARID	 GSK-J1 JMJD3/ UTX/ JARID
JARID	 KDM5-C49 JARID
LSD1	 ORY-1101 LSD1

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
LSD1	
OGT	
OGT	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
OGT	 

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
OGT	
TET1, TET2	

TABLE 2-continued





A list of compounds that bind to regulator proteins.	
Target protein	Compound
TET1	
	
TET1	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
CBP BRD	 <chem>CCOC(=O)c1cc(Cl)c2c(c1)nc(=O)n2C1=CC=C(C=C1)N1=NC(=C(C=C1)NS(=O)(=O)O)NCC(=O)O</chem>
	 <chem>CC1=CC=C(C=C1)C(=O)O1=CC=C(C=C1)NC(=O)c2cc3c(cc2)nc(=O)[n+]3C</chem>
	 <chem>CC(=O)Nc1c[nH]c2cc(Br)ccc12</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	 <chem>COC(=O)c1ccc2c(c1)oc(=O)[nH]2</chem>
	 <chem>CC(=O)c1c[nH]c2ccccc12</chem>
	 <chem>CC(=O)c1c[nH]c2ccccc12C3=CN=CN=C3C4=CC=CC=C4</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	 <chem>CCOC1=CC=C(C=C1C(=O)NCC2=CC=CC=C2O2)C(=O)NCC3=CC=CC=C3C(=O)NCC4=CC=C(C=C4)C(=O)NCC5=CC=CC=C5O5</chem>
	 <chem>CCOC1=CC=C(C=C1C(=O)NCC2=CC=CC=C2O2)C(=O)NCC3=CC=CC=C3C(=O)NCC4=CC=C(C=C4)C(=O)NCC5=CC=CC=C5O5</chem>
	 <chem>CCOC1=CC=C(C=C1C(=O)NCC2=CC=CC=C2O2)C(=O)NCC3=CC=CC=C3C(=O)NCC4=CC=C(C=C4)C(=O)NCC5=CC=CC=C5O5</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	 <chem>CC(=O)c1ccc(NC(=O)c2ccccc2C(=O)O)cc1</chem>
	 <chem>CC(=O)c1ccc(NC(=O)c2cc(OC)c3ccccc3OCCN4CCNCC4)cc1</chem>
	 <chem>CC(=O)c1ccc(NC(=O)c2cc(ccc2c3nn[nH]3)c4ccsc4)cc1</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	 <chem>CC(=O)Nc1ccc(NC(=O)c2ccoc2)cc1NC(=O)c3ccc(OC)cc3C(=O)C</chem>
	 <chem>CC(=O)C1=CC=C(C=C1C2=CC=CC=C2N3CCF(F)CC3)C=C1C=C(C=C1)OCC</chem>
	 <chem>CC(=O)Nc1ccc(NC(=O)c2ccoc2)cc1C2=C(C(=O)N2C)O</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.


















































































































Target protein	Compound
CBP BRD	                                                                                                                

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
CBP BRD	 <chem>CC1=CN=C(C)O1c2ccc(cc2-c3cc4c(c3)nn4CC(C)(C)C)N(C)CC</chem>
	 <chem>CC1=CN=C(C)O1c2ccc(cc2-c3cc4c(c3)nn4CCc5ccccc5)N(C)CC</chem>
CBP BRD	 <chem>CC1=CN=C(C)O1c2ccc(cc2-c3cc4c(c3)nn4CCc5ccccc5)N(R)CC</chem>  

TABLE 2-continued

A list of compounds that bind to regulator proteins.








Target protein	Compound
	
	
	
	
	
	
CBP BRD	

TABLE 2-continued

A list of compounds that bind to regulator proteins.









Target protein	Compound
	R 
	
	
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.







Target protein	Compound
	
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.







Target protein	Compound
CBP BRD	     

TABLE 2-continued

A list of compounds that bind to regulator proteins.







Target protein	Compound
	
	
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
CBP BRD	
	
CBP BRD	

TABLE 2-continued






Target protein	Compound																																					
R ¹		A																																				
		B																																				
		C																																				
	<table><tr><th colspan="3">R</th></tr><tr><td>1</td><td>2</td><td>3</td></tr><tr><td>A</td><td>CH₃</td><td>H</td></tr><tr><td>A</td><td>H</td><td>CH₃</td></tr><tr><td>B</td><td>CH₃</td><td>H</td></tr><tr><td>B</td><td>H</td><td>CH₃</td></tr><tr><td>A</td><td>H</td><td>(R)-CH₃</td></tr><tr><td>A</td><td>H</td><td>(S)-CH₃</td></tr><tr><td>B</td><td>H</td><td>(R)-CH₃</td></tr><tr><td>B</td><td>H</td><td>(S)-CH₃</td></tr><tr><td>C</td><td>H</td><td>(R)-CH₃</td></tr><tr><td>C</td><td>H</td><td>(S)-CH₃</td></tr></table>	R			1	2	3	A	CH ₃	H	A	H	CH ₃	B	CH ₃	H	B	H	CH ₃	A	H	(R)-CH ₃	A	H	(S)-CH ₃	B	H	(R)-CH ₃	B	H	(S)-CH ₃	C	H	(R)-CH ₃	C	H	(S)-CH ₃	
	R																																					
	1	2	3																																			
	A	CH ₃	H																																			
	A	H	CH ₃																																			
	B	CH ₃	H																																			
	B	H	CH ₃																																			
	A	H	(R)-CH ₃																																			
	A	H	(S)-CH ₃																																			
B	H	(R)-CH ₃																																				
B	H	(S)-CH ₃																																				
C	H	(R)-CH ₃																																				
C	H	(S)-CH ₃																																				
HDAC		R																																				
																																						

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
	
	
	
	

TABLE 2-continued






A list of compounds that bind to regulator proteins.	
Target protein	Compound
HDAC	
HDAC	
HDAC1, HDAC2, HDAC3	
HDAC2, HDAC3	
HDAC1, HDAC3	

TABLE 2-continued




A list of compounds that bind to regulator proteins.	
Target protein	Compound
HDAC	 <chem>CN(C)c1ccc(cc1)/N=N/c2ccc(cc2)C(=O)Nc3cc(N)ccc3c4ccsc4</chem>
HDAC1, HDAC2, HDAC3	 <chem>C1CCNCC1c2ccc3cc4nc5cc(NC(=O)c6cc(N)ccc6c7ccsc7)ccc5n4</chem>
HDAC1, HDAC2, HDAC3	 <chem>CCCC(O)C(=O)Nc1ccc2cc3cc(NC(=O)c4cc(N)ccc4c5ccsc5)ccc3n2</chem>

TABLE 2-continued






A list of compounds that bind to regulator proteins.	
Target protein	Compound
HDAC6, HDAC8	
HDAC6	
HDAC6	
	
	

TABLE 2-continued





A list of compounds that bind to regulator proteins.	
Target protein	Compound
HDAC	
HDAC6	
HDAC1, HDAC2, HDAC3, HDAC6	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
HDAC1, HDAC2, HDAC3, HDAC6	
HDAC4	
HDAC6, HDAC8	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
HDAC6	
HDAC6	
HDAC	
HDAC6	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
HDAC6	
HDAC6	
HDAC1, HDAC6	
HDAC1, HDAC6	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
HDAC6, HDAC8	
HDAC1, HDAC6	
HDAC5, HDAC5, HDAC6, HDAC8	
HDAC6	

TABLE 2-continued




A list of compounds that bind to regulator proteins.	
Target protein	Compound
HDAC1, HDAC6	
HDAC1, HDAC6	
HDAC	

TABLE 2-continued






A list of compounds that bind to regulator proteins.	
Target protein	Compound
HDAC1, HDAC2, HDAC3, HDAC5, HDAC 6	
	
HDAC1, HDAC6	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
HDAC8, HDAC11	
	
HDAC8	
HDAC1, HDAC6	

TABLE 2-continued





A list of compounds that bind to regulator proteins.	
Target protein	Compound
HDAC1, HDAC6	
	
	
HDAC	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
HDAC1	
HDAC1, HDAC2, HDAC3, HDAC6, HDAC8, HDAC10, HDAC11	
HDAC1, HDAC2, HDAC3, HDAC6, HDAC8, HDAC10, HDAC11	

TABLE 2-continued




A list of compounds that bind to regulator proteins.	
Target protein	Compound
HDAC4, HDAC5, HDAC7, HDAC9	
	
HDAC4	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
HDAC4	 <chem>Fc1ccncc1[C@H]2C[C@@H]2C(=O)Nc3ccccc3</chem>
	 <chem>Cc1ccncc1[C@H]2C[C@@H]2C(=O)Nc3ccccc3</chem>
	 <chem>COc1ccncc1[C@H]2C[C@@H]2C(=O)Nc3ccccc3C(F)F</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	 <chem>FC1=CC=C(C=C1C2=CC=CC=C2C(=O)N(O)C(F)C3=CC=CC=C3)C4=CN=CC=C4C(F)(F)F</chem>
HDAC4	 <chem>FC1=CC=C(C=C1C2=CC=CC=C2C(=O)N(O)C(F)C3=CC=CC=C3)C4=CC=C(C=C4)C5=C(C=C(C=C5)OC6=CC=CC=C6C7CC7)O</chem>
HDAC4	 <chem>FC1=CC=C(C=C1C2=CC=CC=C2C(=O)N(O)C(F)C3=CC=CC=C3)C4=CC=C(C=C4)SC5=CC=NC=C5C6=CC=CC=C6F</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
HDAC4	
HDAC4	
HDAC5, HDAC8	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
HDAC4, HDAC8	
HDAC	
HDAC4	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
HDAC1, HDAC6, HDAC9	
HDAC2, HDAC6	
P300/CBP	

TABLE 2-continued

A list of compounds that bind to regulator proteins.	
Target protein	Compound
p300, PCAF	
p300, PCAF	
p300	
HAT	
Tip60	

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
p300/CBP, PCAF, Tip60	
p300 activator	
PCAF	
Tip60	
PCAF	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
p300	
p300, PCAF	
p300	

TABLE 2-continued





A list of compounds that bind to regulator proteins.	
Target protein	Compound
p300	 <chem>Oc1ccc(Oc2ccc(O)c(O)c2)cc1Oc3ccc(O)c(O)c3</chem>
p300/CBP	 <chem>COc1ccc(O)cc1/C=C/C(=O)C(=O)/C=C/c2ccc(O)c(OC)c2</chem>
p300	 <chem>CC[C@H](C)C1=CC=CC=C1</chem>
p300	 <chem>CC(=O)C(O)C(=O)N[C@@H](C(=O)O)CCOc1ccc(O)cc1C(=O)N[C@@H](C(=O)O)CC(=O)N</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
p300	
p300/CBP	
PCAF	
GCN5	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
p300	 <chem>Cc1cc(C)cc([N+](=O)[O-])cc1Oc1ccoc1C2=CN(C(=O)c3ccc(C(=O)O)cc3)N2</chem>
Tip60	 <chem>C1=CN=C(S2=NC=CC2)S3=NC=CC3</chem>
Tip60	 <chem>COc1ccc(OC)cc1C2=NN(C2C3=CC=CC=C3C(=O)N4C(=O)c5ccccc54)C6=CC=CC=C6S7=CC=CC=C7</chem>
p300	 <chem>COc1ccc(cc1)S(=O)(=O)OC(=O)N2C=CC(=O)CC2</chem>

TABLE 2-continued





A list of compounds that bind to regulator proteins.	
Target protein	Compound
Tip60	
HDAC1, HDAC2, HDAC3, HDAC8	
HDAC1, HDAC2, HDAC3, HDAC8	
HDAC1, HDAC2, HDAC3, HDAC8	

TABLE 2-continued





A list of compounds that bind to regulator proteins.	
Target protein	Compound
HDAC1, HDAC2, HDAC3	 <chem>CCN(CC)Cc1ccc2cc3c(cc12)C(=O)OCC(=O)Nc4ccc(NC(=O)O)cc4</chem>
HDAC1, HDAC2, HDAC3, HDAC8	 <chem>O=C(O)/C=C/c1ccc(NC(=O)S(=O)(=O)c2ccccc2)cc1</chem>
HDAC1, HDAC2, HDAC3, HDAC8	 <chem>O=C(O)/C=C/c1ccc(CN(CC1)CCc2c[nH]c3ccccc23)cc1</chem>
HDAC1, HDAC2, HDAC3, HDAC8	 <chem>O=C(O)CCc1c[nH]c2ccccc12</chem>

TABLE 2-continued





A list of compounds that bind to regulator proteins.	
Target protein	Compound
HDAC1, HDAC2, HDAC3	
HDAC1, HDAC2, HDAC3	
HDAC2, HDAC3	
CDK2	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
CDK2	 <chem>Nc1nc2c(ncn2COC3CCCCC3)c1</chem>
CDK2	 <chem>Nc1nc2c(ncn2COC3CCCCC3)c(R)c1</chem> 2; R = H 3; R = SO ₂ NH ₂
CDK2	 <chem>CC(C)OC1=NC(=C(N)C=C1C2=CC=C(S(=O)(=O)N)C2)C=O</chem>

TABLE 2-continued




A list of compounds that bind to regulator proteins.	
Target protein	Compound
CDK2, CDK, CDK9	
CDK2, CDK7, CDK9	
CDK2, CDK7, CDK9	

TABLE 2-continued

A list of compounds that bind to regulator proteins.







Target protein	Compound
CDK2	
	
	
	
	
	
	
	
	
	
	
	
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
	
	$C\equiv CSi(i-Pr)_3$
	$HC\equiv C$
	R^2
	H
	SO_2NH_2
	H
	SO_2NH_2
	SO_2NH_2
	SO_2NH_2
	SO_2NH_2
	SO_2NH_2
	SO_2NH_2
	SO_2NH_2
	SO_2NH_2
	SO_2NH_2
	SO_2NH_2
	H
CDK2	
	R^1
	$C\equiv CH$
	$C\equiv CMe$
	Et
	
	Ph
	

TABLE 2-continued

A list of compounds that bind to regulator proteins.






Target protein	Compound
	
	
	
	
	
	R^2 SO_2NH_2 H H SO_2NH_2 SO_2NH_2 SO_2NH_2 SO_2NH_2

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
CDK2	SO_2NH_2
	SO_2NH_2
	SO_2NH_2
	SO_2NH_2
	SO_2NH_2
CDK2	
	
	H
	
	
CDK2	Ph
	
	Structure

TABLE 2-continued

A list of compounds that bind to regulator proteins.











Target protein	Compound	H
		
		
		
		
		

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	 H SO ₂ NH ₂
	
CDK	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
	
	
	

TABLE 2-continued




A list of compounds that bind to regulator proteins.	
Target protein	Compound
PCAF BRD, L3MBTL3	
PCAF BRD, L3MBTL3	
CBP/p300	

TABLE 2-continued





A list of compounds that bind to regulator proteins.	
Target protein	Compound
PRMT5	
HDAC	
2-oxoglutarate dependent KDM5 demethylases	
CDK4, CDK6	

TABLE 2-continued





A list of compounds that bind to regulator proteins.	
Target protein	Compound
CDK4, CDK6	
CDK4, CDK6	
HDAC	
HDAC	

TABLE 2-continued






A list of compounds that bind to regulator proteins.	
Target protein	Compound
HDAC	
Pan-HDAC	
HDAC	
HDAC1, HDAC3	
HDAC	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
Pan-HDAC	
HDAC6	
Class I HDAC	

TABLE 2-continued

A list of compounds that bind to regulator proteins.	
Target protein	Compound
Class I HDAC	
Class I HDAC	
Class IIa HDAC	

TABLE 2-continued





A list of compounds that bind to regulator proteins.	
Target protein	Compound
HDAC3	
HDAC3	
HDAC6	
HDAC6	

TABLE 2-continued





A list of compounds that bind to regulator proteins.	
Target protein	Compound
HDAC6	
HDAC8	
HDAC8	
HDAC1, HDAC2	

TABLE 2-continued




A list of compounds that bind to regulator proteins.	
Target protein	Compound
HDAC1, HDAC2	
HDAC1	
HDAC	

TABLE 2-continued




A list of compounds that bind to regulator proteins.	
Target protein	Compound
HDAC, PI3K	
HDAC, EGFR, HER2	
HDAC	

TABLE 2-continued




A list of compounds that bind to regulator proteins.	
Target protein	Compound
HDAC1, HDAC6, ER	
Class I HDACs, ZEB1	
HDAC, Akt	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
HDAC	
HDAC	
HDAC1	
Class I HDACs	

TABLE 2-continued





A list of compounds that bind to regulator proteins.	
Target protein	Compound
HDAC6	
HDAC6	
HDAC3, HDAC6, HDAC8	
HDAC6	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
HDAC2	
HDAC2	
HDAC4	
HDAC1, HDAC2	

TABLE 2-continued



A list of compounds that bind to regulator proteins.	
Target protein	Compound
Pan-HDAC	
HDAC4	

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
HDAC6	
G9a, GLP	

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
SMYD2	
EZH2	

TABLE 2-continued




A list of compounds that bind to regulator proteins.	
Target protein	Compound
DOT1L	
PRMT5	
Pan-jmjC	

TABLE 2-continued





A list of compounds that bind to regulator proteins.	
Target protein	Compound
JARID	
JMJD3, UTX, JARID	
LSD1	
L3MBTL1-MBT	

TABLE 2-continued

A list of compounds that bind to regulator proteins.

Target protein	Compound
L3MBTL1-MBT	

TABLE 2-continued




A list of compounds that bind to regulator proteins.	
Target protein	Compound
L3MBTL3-MBT	
	
	

TABLE 2-continued




A list of compounds that bind to regulator proteins.	
Target protein	Compound
CBX7	
53BP1	 

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
JARID1A-PHD3	
	
	
Pygo-PHD	

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
WDR5-MML	 <chem>CN1CCN(CC1)c2ccc(cc2)N(=O)c3ccc(cc3)NC(=O)c4ccccc4</chem>
	 <chem>CN1CCN(CC1)c2ccc(cc2)N(=O)c3ccc(cc3)NC(=O)c4ccc(cc4)Cl</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.

Target protein	Compound
	<chem>COC(=O)c1ccc(NC(=O)c2ccc(OC)cc2)cc1N1CCN(C)CC1</chem>
	<chem>CCCCNC(=O)[C@H](Cc1ccc(F)cc1)NC(=O)N[C@@H]2CCC(CC2)(C)C(=O)C(C)(C)C3CCCC3C3=CC=CC=C3F</chem>

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
CDK1, CDK2, CDK4, CDK5, CDK6, CDK7, CDK9	 

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
CDK1, CDK2, CDK4, CDK6, CDK9	
CDK1, CDK2, CDK5, CDK7	

TABLE 2-continued




A list of compounds that bind to regulator proteins.	
Target protein	Compound
CDK1, CDK2, CDK5, CDK9	 <chem>CC1=CC=C(C(=C1)N+)CCNc2nc3c(ncn3C4CCCCC4N2)C(O)CC</chem>
CDK1, CDK2, CDK4, CDK5, CDK6, CDK7	 <chem>CC1(C)S(=O)(=O)N1CCN(C1)Nc2nc(N)c(C(=O)c3cc(F)c(F)cc3O)c2</chem>
CDK1, CDK2, CDK4, CDK5, CDK7, CDK9	 <chem>CC1(C)S(=O)(=O)N1c2ccc(Nc3nc(O[C@@H](C)C)nc3C(F)(F)F)cc2</chem>

TABLE 2-continued



A list of compounds that bind to regulator proteins.	
Target protein	Compound
CDK1, CDK2, CDK5, CDK7, CDK9	
CDK4, CDK6	

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
CDK1, CDK2, CDK4, CDK5	
CDK4, CDK6	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
CKD1, CDK2, CDK5, CDK6, CDK7, CDK9	
CDK2, CDK4, CDK5, CDK6, CDK9	
CDK1, CDK2, CDK4, CDK7, CDK9	

TABLE 2-continued




A list of compounds that bind to regulator proteins.	
Target protein	Compound
CDK1, CDK2, CDK4, CDK5, CDK6, CDK9	
CDK4	
CDK1, CDK4	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
CDK4, CDK6	
CDK4	
CDK2, CDK9	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
CDK5	
CDK8	
CDK1, CDK2, CDK5, CDK7, CDK9	

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
CDK	
CDK	

TABLE 2-continued



A list of compounds that bind to regulator proteins.	
Target protein	Compound
CDK	
CDK1, CDK2, CDK5, CDK9	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
CDK7	
CDK7	
CDK2	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
CDK2, HDAC	
CDK3	
CDK5	

TABLE 2-continued

A list of compounds that bind to regulator proteins.





Target protein	Compound
CDK4	
CDK4	
CDK8	
CDK4	

TABLE 2-continued






A list of compounds that bind to regulator proteins.	
Target protein	Compound
CDK2, CDK9	
CDK2, CDK9	
CDK2, CDK9	
CDK2, CDK9	
CDK2, CDK9	

TABLE 2-continued





A list of compounds that bind to regulator proteins.	
Target protein	Compound
CDK2, CDK9	
CDK2, CDK9	
CDK2, CDK9	
CDK2, CDK9	

TABLE 2-continued




A list of compounds that bind to regulator proteins.	
Target protein	Compound
CDK9	
CDK2, HDAC	
CDK7	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
CDK2, CDK9	
CDK1, CDK2, CDK5, CDK9	
CDK2, HDAC1	

TABLE 2-continued

A list of compounds that bind to regulator proteins.



Target protein	Compound
CDK9	
CDK9	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
CDK9	
CDK9	
CDK9	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
CDK9	
CDK, CDC7	
CDK8, CDK19	

TABLE 2-continued

A list of compounds that bind to regulator proteins.




Target protein	Compound
CDK8, CDK19	 <chem>N#Cc1ccc2nc3c(ncn3CNCc4ccc(Cl)cc4)c2</chem>
CDK8, CDK19, MAP4K2, YSK4	 <chem>CN1CCN(C)CC1C(=O)c2ccc3cc(NC4=NC5=C(N=CN=C5C#N)C=C4)ccc3</chem>
CDK8, CDK19	 <chem>N#Cc1ccc2nc3c(ncn3NCNCc4ccccc4)c2.[Cl-]</chem>

TABLE 2-continued



A list of compounds that bind to regulator proteins.	
Target protein	Compound
CDK4, CDK6	
CDI9, CK2, PIM1	

TABLE 2-continued






A list of compounds that bind to regulator proteins.	
Target protein	Compound
CDK1, CDK2, CDK5	 <chem>Fc1ccc(cc1Nc2ccc3ncnc3c2)Oc4ccccc4N5CCNCC5</chem>
CDK1, CDK2, CDK3, CDK4, CDK6, CDK7, CDK9, HDAC	 <chem>CC(C)(C)c1cc(oc1)CSCc2ncnc2NC(=O)Cc3ccc(OCCOC3=CC=CC=C3)cc3</chem>
CDK2	 <chem>Nc1ccc(cc1)NC(=O)c2ccsc2C(=O)Nc3ccc(N4CCNCC4)cc3</chem>

TABLE 2-continued


A list of compounds that bind to regulator proteins.

Target protein	Compound
CDK2	 <chem>CN1CCN(CC1)C(=O)c2sc(N)n2c3cc(Cl)cc(Cl)cc3</chem>
	 <chem>ClCCN(CC1=CC=C(C=C1)CC(=O)Nc2ncsc2SCc3nc(C(C)(C)C)co3)CC1</chem>

② indicates text missing or illegible when filed

[0446] In some embodiments, the second terminus does not comprises JQ1, JQ-1, OTX015, RVX208 acid, or RVX208 hydroxyl.

[0447] In certain embodiments, the protein binding moiety is a residue of a compound having a structure of Formula (C-1):



[0448] wherein:

[0449] X^a is $-\text{NHC(O)}-$, $-\text{C(O)}-\text{NH}-$, $-\text{NHSO}_2-$, or $-\text{SO}_2\text{NH}-$;

[0450] A^a is selected from an optionally substituted $-\text{C}_{1-12}$ alkyl, optionally substituted $-\text{C}_{2-10}$ alkenyl, optionally substituted $-\text{C}_{2-10}$ alkynyl, optionally substituted $-\text{C}_{1-12}$ alkoxy, optionally substituted $-\text{C}_{1-12}$ haloalkyl, optionally substituted C_{6-10} aryl, optionally substituted C_{3-7} cycloalkyl, optionally substituted 5- to 10 membered heteroaryl, and optionally substituted 5- to 10-membered heterocycloalkyl;


[0451] X^b is a bond, NH , $\text{NH}-\text{C}_{1-10}$ alkylene, $-\text{C}_{1-12}$ alkyl, $-\text{NHC(O)}-$, or $-\text{C(O)}-\text{NH}-$;

[0452] A^b is selected from an optionally substituted $-\text{C}_{1-12}$ alkyl, optionally substituted $-\text{C}_{2-10}$ alkenyl, optionally substituted $-\text{C}_{2-10}$ alkynyl, optionally substituted $-\text{C}_{1-12}$ alkoxy, optionally substituted $-\text{C}_{1-12}$ haloalkyl, optionally substituted C_{6-10} aryl, optionally substituted C_{3-7} cycloalkyl, optionally substituted 5- to 10 membered heteroaryl, and optionally substituted 4- to 10-membered heterocycloalkyl; and

[0453] each R^{1a} , R^{2e} , R^{3e} , R^{4e} are independently selected from the group consisting of H, OH, $-\text{NO}_2$, halogen, amine, COOH , COOC_{1-10} alkyl, $-\text{NHC(O)}$ -optionally substituted $-\text{C}_{1-12}$ alkyl, $-\text{NHC(O)}(\text{CH}_2)_{1-4}\text{NR}'\text{R}^g$, $-\text{NHC(O)}(\text{CH}_2)_{0-4}\text{CHR}'(\text{NR}'\text{R}^g)$, $-\text{NHC(O)}(\text{CH}_2)_{0-4}\text{CHR}^9$, $-\text{NHC(O)}(\text{CH}_2)_{0-4}-\text{C}_{3-7}$ cycloalkyl, $-\text{NHC(O)}(\text{CH}_2)_{0-4}$ -5- to 10-membered heterocycloalkyl, $\text{NHC(O)}(\text{CH}_2)_{0-4}\text{C}_{6-10}$ aryl, $-\text{NHC(O)}(\text{CH}_2)_{0-4}$ -5- to 10-membered heteroaryl, $-(\text{CH}_2)_{1-4}-\text{C}_{3-7}$ cycloalkyl, $-(\text{CH}_2)_{1-4}$ -5- to 10-membered heterocycloalkyl, $-(\text{CH}_2)_{1-4}\text{C}_{6-10}$ aryl, $-(\text{CH}_2)_{1-4}$ -5- to 10-membered heteroaryl, optionally substituted $-\text{C}_{2-10}$ alkenyl, optionally substituted $-\text{C}_{2-10}$ alkynyl, optionally substituted $-\text{C}_{1-12}$ alkoxy, optionally substituted $-\text{C}_{1-12}$ haloalkyl, optionally substituted C_{6-10} aryl, optionally substituted C_{3-7} cycloalkyl, optionally substituted 5- to 10-membered heteroaryl, and optionally substituted 4- to 10-membered heterocycloalkyl, and

[0454] wherein each R and R^g are independently H or C_{1-6} alkyl.


[0455] In certain embodiments, the protein binding moiety is a residue of a compound having a structure of Formula (C-2):



wherein R^{5c} is independently selected from the group consisting of H, COOC_{1-10} alkyl, $-\text{NHC(O)}$ -optionally substituted $-\text{C}_{1-12}$ alkyl, optionally substituted $-\text{C}_{2-10}$ alkenyl, optionally substituted $-\text{C}_{2-10}$ alkynyl, optionally substituted $-\text{C}_{1-12}$ alkoxy, optionally substituted $-\text{C}_{1-12}$ haloalkyl, optionally substituted C_{6-10} aryl, optionally substituted C_{3-7} cycloalkyl, optionally substituted 5- to 10-membered heteroaryl, and optionally substituted 5- to 10-membered heterocycloalkyl substituted $-\text{C}_{2-10}$ alkenyl, optionally substituted $-\text{C}_{2-10}$ alkynyl, optionally substituted $-\text{C}_{1-12}$ alkoxy, optionally substituted $-\text{C}_{1-12}$ haloalkyl, optionally substituted C_{6-10} aryl, optionally substituted C_{3-7} cycloalkyl, optionally substituted 5- to 10-membered heteroaryl, and optionally substituted 5- to 10-membered heterocycloalkyl.

[0456] In certain embodiments, A^a is selected from an optionally substituted C_{6-10} aryl, optionally substituted C_{3-7} cycloalkyl, optionally substituted 5- to 10 membered heteroaryl, and optionally substituted 5- to 10-membered heterocycloalkyl. In certain embodiments, A^a is an optionally substituted C_{6-10} aryl.

[0457] In certain embodiments, the protein binding moiety is a residue of a compound having a structure of Formula (C-3):



[0458] wherein:

[0459] M^{1c} is CR^{2h} or N; and

[0460] each R^{1h} , R^{2h} , R^{3h} , R^{4h} , and R^{5h} are independently selected from the group consisting of H, OH, $-\text{NO}_2$, halogen, amine, COOH , COOC_{1-10} alkyl, $-\text{NHC(O)}$ -optionally substituted $-\text{C}_{1-12}$ alkyl, $-\text{NHC(O)}(\text{CH}_2)_{1-4}\text{NR}'\text{R}^g$, $-\text{NHC(O)}(\text{CH}_2)_{0-4}\text{CHR}'(\text{NR}'\text{R}^g)$, $-\text{NHC(O)}(\text{CH}_2)_{0-4}\text{CHR}^g$, $-\text{NHC(O)}(\text{CH}_2)_{0-4}-\text{C}_{3-7}$ cycloalkyl, $-\text{NHC(O)}(\text{CH}_2)_{0-4}$ -5- to 10-membered heterocycloalkyl, $\text{NHC(O)}(\text{CH}_2)_{0-4}\text{C}_{6-10}$ aryl, $-\text{NHC(O)}(\text{CH}_2)_{0-4}$ -5- to 10-membered heteroaryl, and optionally substituted 4- to 10-membered heterocycloalkyl, and

10-membered heterocycloalkyl, $\text{NHC(O)(CH}_2\text{)}_{0-4}\text{C}_{6-10}$ aryl, $\text{—NHC(O)(CH}_2\text{)}_{0-4}$ -5- to 10-membered heteroaryl, $\text{—(CH}_2\text{)}_{1-4}\text{—C}_{3-7}$ cycloalkyl, $\text{—(CH}_2\text{)}_{1-4}$ -5- to 10-membered heterocycloalkyl, $\text{—(CH}_2\text{)}_{1-4}\text{C}_{6-10}$ aryl, $\text{—(CH}_2\text{)}_{1-4}$ -5- to 10-membered heteroaryl, optionally substituted —C_{2-10} alkenyl, optionally substituted —C_{2-10} alkynyl, optionally substituted —C_{1-12} alkoxy, optionally substituted —C_{1-12} haloalkyl, optionally substituted C_{6-10} aryl, optionally substituted C_{3-7} cycloalkyl, optionally substituted 5- to 10-membered heteroaryl, and optionally substituted 5- to 10-membered heterocycloalkyl, wherein each R and R^g are independently H or C_{1-6} alkyl.

[0461] In certain embodiments, each R^{1h} and R^{5h} are independently hydrogen, halogen, or C_{1-6} alkyl. In certain embodiments, each R^{2h} and R^{3h} are independently H, OH, —NO_2 , halogen, C_{1-4} haloalkyl, amine, COOH, COOC_{1-10} alkyl, —NHC(O)— optionally substituted —C_{1-12} alkyl, $\text{—NHC(O)(CH}_2\text{)}_{1-4}\text{NR}^g$, $\text{—NHC(O)(CH}_2\text{)}_{0-4}$ CHR' (NR^g), $\text{—NHC(O)(CH}_2\text{)}_{0-4}$ CHR'/ R^g , $\text{—NHC(O)(CH}_2\text{)}_{0-4}\text{—C}_{3-7}$ cycloalkyl, $\text{—NHC(O)(CH}_2\text{)}_{0-4}$ -5- to 10-membered heterocycloalkyl, $\text{NHC(O)(CH}_2\text{)}_{0-4}\text{C}_{6-10}$ aryl, $\text{—NHC(O)(CH}_2\text{)}_{0-4}$ -5- to 10-membered heteroaryl, $\text{—(CH}_2\text{)}_{1-4}\text{—C}_{3-7}$ cycloalkyl, $\text{—(CH}_2\text{)}_{1-4}$ -5- to 10-membered heterocycloalkyl, $\text{—(CH}_2\text{)}_{1-4}\text{C}_{6-10}$ aryl, $\text{—(CH}_2\text{)}_{1-4}$ -5- to 10-membered heteroaryl, optionally substituted —C_{2-10} alkenyl, optionally substituted —C_{2-10} alkynyl, optionally substituted —C_{1-12} alkoxy, optionally substituted C_{6-10} aryl, optionally substituted C_{3-7} cycloalkyl, optionally substituted 5- to 10-membered heteroaryl, and optionally substituted 5- to 10-membered heterocycloalkyl. In certain embodiments, R^{1e} , R^{3e} , and R^{4e} are hydrogen.


[0462] In certain embodiments, R^{2e} is selected from the group consisting of H, OH, —NO_2 , halogen, amine, COOH, COOC_{1-10} alkyl, —NHC(O)— optionally substituted —C_{1-12} alkyl, $\text{—NHC(O)(CH}_2\text{)}_{1-4}\text{NR}^g$, $\text{—NHC(O)(CH}_2\text{)}_{0-4}$ CHR (NR^g), $\text{—NHC(O)(CH}_2\text{)}_{0-4}$ CHR'/ R^g , $\text{—NHC(O)(CH}_2\text{)}_{0-4}\text{—C}_{3-7}$ cycloalkyl, $\text{—NHC(O)(CH}_2\text{)}_{0-4}$ -5- to 10-membered heterocycloalkyl, $\text{NHC(O)(CH}_2\text{)}_{0-4}\text{C}_{6-10}$ aryl, $\text{—NHC(O)(CH}_2\text{)}_{0-4}$ -5- to 10-membered heteroaryl, $\text{—(CH}_2\text{)}_{1-4}\text{—C}_{3-7}$ cycloalkyl, $\text{—(CH}_2\text{)}_{1-4}$ -5- to 10-membered heterocycloalkyl, $\text{—(CH}_2\text{)}_{1-4}\text{C}_{6-10}$ aryl, $\text{—(CH}_2\text{)}_{1-4}$ -5- to 10-membered heteroaryl, optionally substituted —C_{1-12} alkyl, optionally substituted —C_{2-10} alkenyl, optionally substituted —C_{2-10} alkynyl, optionally substituted —C_{1-12} alkoxy, optionally substituted —C_{1-12} haloalkyl, optionally substituted C_{6-10} aryl, optionally substituted C_{3-7} cycloalkyl, optionally substituted 5- to 10-membered heteroaryl, and optionally substituted 5- to 10-membered heterocycloalkyl, wherein each R and R^g are independently H or C_{1-6} alkyl.

[0463] In certain embodiments, R^{2e} is an phenyl or pyridinyl optionally substituted with 1-3 substituents, wherein the substituent is independently selected from the group consisting of OH, —NO_2 , halogen, amine, COOH, COOC_{1-10} alkyl, —NHC(O)— C_{1-12} alkyl, $\text{—NHC(O)(CH}_2\text{)}_{1-4}\text{NR}^g$, $\text{—NHC(O)(CH}_2\text{)}_{0-4}$ CHR (NR^g), $\text{—NHC(O)(CH}_2\text{)}_{0-4}$ CHR'/ R^g , $\text{—NHC(O)(CH}_2\text{)}_{0-4}\text{—C}_{3-7}$ cycloalkyl, $\text{—NHC(O)(CH}_2\text{)}_{0-4}$ -5- to 10-membered heterocycloalkyl, $\text{NHC(O)(CH}_2\text{)}_{0-4}\text{C}_{6-10}$ aryl, $\text{—NHC(O)(CH}_2\text{)}_{0-4}$ -5- to 10-membered heteroaryl, $\text{—(CH}_2\text{)}_{1-4}\text{—C}_{3-7}$ cycloalkyl, $\text{—(CH}_2\text{)}_{1-4}$ -5- to 10-membered heterocycloalkyl, $\text{—(CH}_2\text{)}_{1-4}\text{C}_{6-10}$ aryl, $\text{—(CH}_2\text{)}_{1-4}$ -5- to 10-membered heteroaryl, —C_{1-12} alkoxy, C_{1-12} haloalkyl, C_{6-10} aryl, C_{3-7} cycloalkyl, 5- to 10-membered heteroaryl, and 5- to 10-membered heterocycloalkyl,

wherein each R and R^g are independently H or C_{1-6} alkyl

[0464] In certain embodiments, A^a is a C_{6-10} aryl substituted with 1-4 substituents, and each substituent is independently selected from halogen, OH, NO_2 , an optionally substituted —C_{1-12} alkyl, optionally substituted —C_{2-10} alkenyl, optionally substituted —C_{2-10} alkynyl, optionally substituted —C_{1-12} alkoxy, optionally substituted —C_{1-12} haloalkyl, optionally substituted C_{6-10} aryl, optionally substituted C_{3-7} cycloalkyl, optionally substituted 5- to 10-membered heteroaryl, and optionally substituted 5- to 10-membered heterocycloalkyl.

[0465] In certain embodiments, the protein binding moiety is a residue of a compound having the structure of Formula (C-4):



wherein:

[0467] R^{1e} is an optionally substituted C_{6-10} aryl or an optionally substituted 5- to 10-membered heteroaryl,

[0468] X^c is —C(O)NH— , —C(O)— , $\text{—S(O)}_2\text{—}$, —NH— , or —C_{1-4} alkyl-NH,


[0469] n is 0-10,

[0470] R^{2f} is $\text{—NR}^3\text{R}^{4f}$, optionally substituted C_{6-10} aryl, optionally substituted C_{3-7} cycloalkyl, optionally substituted 5- to 10-membered heteroaryl, or optionally substituted 4- to 10-membered heterocycloalkyl; and

[0471] each R^{3f} and R^{4f} are independently H or optionally substituted —C_{1-12} alkyl.

[0472] In some embodiments, R^{2f} is $\text{—NHC(CH}_3\text{)}_3$, or a 4- to 10-membered heterocycloalkyl substituted with C_{1-12} alkyl.

[0473] In certain embodiments, the protein binding moiety is a residue of a compound having the structure of Formula (C-5):



wherein:

[0475] X^{2c} is a bond, C(O), SO_2 , or CHR^{3c} ;

[0476] M^{2c} is CH or N;

[0477] n is 0-10,

[0478] R^{2f} is $\text{—NR}^3\text{R}^{4f}$, optionally substituted C_{6-10} aryl, optionally substituted C_{3-7} cycloalkyl, optionally substituted 5- to 10-membered heteroaryl, or optionally substituted 4- to 10-membered heterocycloalkyl;


[0479] each R^{5f} is independently $\text{—NR}^3\text{R}^{4f}$, —C(O)R^{3f} , —COOH , —C(O)NHC_{1-6} alkyl, an optionally substituted C_{6-10} aryl, or an optionally substituted 5- to 10-membered heteroaryl;

[0480] R^{6f} is $\text{—NR}^3\text{R}^{4f}$, —C(O)R^{3f} , an optionally substituted C_{6-10} aryl, or an optionally substituted 5- to 10-membered heteroaryl; and

[0481] each R^{3j} and R^{4j} are independently H, an optionally substituted C_{6-10} aryl, optionally substituted 4- to 10-membered heterocycloalkyl, or optionally substituted $-C_{1-12}$ alkyl.

[0482] In certain embodiments, R^{2j} is a 4- to 10-membered heterocycloalkyl substituted by a 4- to 10-membered heterocycloalkyl. In certain embodiments, R^{6j} is $-C(O)R^{3j}$, and R^{3j} is a 4- to 10-membered heterocycloalkyl substituted by a 4- to 10-membered heterocycloalkyl. In certain embodiments, each R^{3j} is independently H, $-C(O)R^{3j}$, $-COOH$, $-C(O)NHC_{1-6}$ alkyl, $-NH-C_{6-10}$ aryl, or optionally substituted C_{6-10} aryl.

[0483] In certain embodiments, the protein binding moiety is a residue of a compound having the structure of Formula (C-6):



(C-6)

[0484] wherein:

[0485] X^{3c} is a bond, NH, C_{1-4} alkylene, or NC_{1-4} alkyl;


[0486] R^{7j} is an optionally substituted C_{1-6} alkyl, an optionally substituted cyclic amine, an optionally substituted aryl, an optionally substituted 5- to 10-membered heteroaryl, or optionally substituted 4- to 10-membered heterocycloalkyl,

[0487] R^{8j} is H, halogen, or C_{1-6} alkyl; and

[0488] R^{9j} is H, or C_{1-6} alkyl.

[0489] In certain embodiments, R^{7j} is an optionally substituted cyclic secondary or tertiary amine. In certain embodiments, R^{73} is a tetrahydroisoquinoline optionally substituted with C_{1-4} alkyl.

[0490] In certain embodiments, the protein binding moiety is a residue of a compound having the structure of Formula (C-7):



(C-7)

[0491] wherein:


[0492] A^{1a} is an optionally substituted aryl or heteroaryl;

[0493] X^2 is a bond, $(CH_2)_{1-4}$, or NH; and

[0494] A^{2a} is an optionally substituted aryl, heterocyclic, or heteroaryl, linked to an amide group.

[0495] In certain embodiments, A^{1a} is an aryl substituted with one or more halogen, C_{1-6} alkyl, hydroxyl, C_{1-6} alkoxy, or C_{1-6} haloalkyl. In certain embodiments, X^2 is NH. In certain embodiments, A^{2a} is a heterocyclic group. In certain embodiments, A^{2a} is a pyrrolidine. In certain embodiments, A^{2a} is an optionally substituted phenyl. In certain embodiments, A^{2a} is a phenyl optionally substituted with one or more halogen, C_{1-6} alkyl, hydroxyl, C_{1-6} alkoxy, or C_{1-6} haloalkyl.


[0496] In certain embodiments, the protein binding moiety is a residue of a compound having the structure of Formula (C-8):



(C-8)

[0497] wherein R^{1k} is H or C_{1-25} alkyl and R^{2k} is OH or $-OC_{1-12}$ alkyl.

[0498] In certain embodiments, the protein binding moiety is a residue of a compound having the structure of Formula (C-9):




(C-9)

[0499] wherein R_{1m} is H, OH, $-CONH_2$, $-COOH$, $-NHC(O)-C_{1-6}$ alkyl, $-NHC(O)O-C_{1-6}$ alkyl, $-NHS(O)_2-C_{1-6}$ alkyl, $-C_{1-6}$ alkyl, $-C_{1-6}$ alkoxy, or $-NHC(O)NH-C_{1-6}$ alkyl;

[0500] R_{2m} is H, CN, or $CONH_2$; and

[0501] R_{3m} is an optionally substituted C_{6-10} aryl.

[0502] In certain embodiments, the protein binding moiety is a residue of a compound having the structure of Formula (C-10):



(C-10)




[0503] wherein R_{1n} is an optionally substituted C_{6-10} aryl or optionally substituted 5- to 10-membered heteroaryl, and

[0504] each R_{2n} and R_{3n} are independently H, $-C_{1-4}$ alkyl- C_{6-10} aryl, $-C_{1-4}$ alkyl-5- to 10-membered heteroaryl, C_{6-10} aryl, or -5- to 10-membered heteroaryl, or


[0505] R_{2n} and R_{3n} together with N form an optionally substituted 4-10 membered heterocyclic or heteroaryl group.

[0506] In certain embodiments, the regulatory molecule is not a bromodomain-containing protein chosen from BRD2, BRD3, BRD4, and BRDT. In certain embodiments, the regulatory molecule is not BRD2, BRD3, BRD4, or BRDT.


[0507] In certain embodiments, the regulatory molecule is not BRD4. In certain embodiments, the recruiting moiety is not a BRD4 activator. In certain embodiments, the BRD4 activator is not chosen from JQ-1, OTX015, RVX208 acid, and RVX208 hydroxyl.



[0508] In certain embodiments, the regulatory molecule is BPTF. In certain embodiments, the recruiting moiety is a BPTF activator. In certain embodiments, the BPTF activator is AU1.




[0509] In certain embodiments, the regulatory molecule is histone acetyltransferase ("HAT"). In certain embodiments, the recruiting moiety is a HAT activator. In certain embodiments, the HAT activator is a oxopiperazine helix mimetic OHM. In certain embodiments, the HAT activator is selected from OHM1, OHM2, OHM3, and OHM4 (BB Lao et al., PNAS USA 2014, 111(21), 7531-7536). In certain embodiments, the HAT activator is OHM4.




[0510] In certain embodiments, the regulatory molecule is histone deacetylase ("HDAC"). In certain embodiments, the recruiting moiety is an HDAC activator. In certain embodiments, the HDAC activator is chosen from SAHA and 109 (Soragni E Front. Neurol. 2015, 6, 44, and references therein).

SAHA




109




[0511] In certain embodiments, the regulatory molecule is histone deacetylase (“HDAC”). In certain embodiments, the recruiting moiety is an HDAC inhibitor. In certain embodiments, the HDAC inhibitor is an inositol phosphate.

[0512] In certain embodiments, the regulatory molecules is O-linked β -N-acetylglucosamine transferase (“OGT”). In certain embodiments, the recruiting moiety is an OGT activator. In certain embodiments, the OGT activator is chosen from ST045849, ST078925, and ST060266 (Itkonen HM, “Inhibition of O-GlcNAc transferase activity reprograms prostate cancer cell metabolism”, *Oncotarget* 2016, 7(11), 12464-12476).

ST045849




ST078925




-continued

ST060266




[0513] In certain embodiments, the regulatory molecule is chosen from host cell factor 1 (“HCF1”) and octamer binding transcription factor (“OCT1”). In certain embodiments, the recruiting moiety is chosen from an HCF1 activator and an OCT1 activator. In certain embodiments, the recruiting moiety is chosen from VP16 and VP64.

[0514] In certain embodiments, the regulatory molecule is chosen from CBP and P300. In certain embodiments, the recruiting moiety is chosen from a CBP activator and a P300 activator. In certain embodiments, the recruiting moiety is CTPB.



[0515] In certain embodiments, the regulatory molecule is P300/CBP-associated factor (“PCAF”). In certain embodiments, the recruiting moiety is a PCAF activator. In certain embodiments, the PCAF activator is embelin.



[0516] In certain embodiments, the regulatory molecule modulates the rearrangement of histones.

[0517] In certain embodiments, the regulatory molecule modulates the glycosylation, phosphorylation, alkylation, or acylation of histones.

[0518] In certain embodiments, the regulatory molecule is a transcription factor.

[0519] In certain embodiments, the regulatory molecule is an RNA polymerase.

[0520] In certain embodiments, the regulatory molecule is a moiety that regulates the activity of RNA polymerase.

[0521] In certain embodiments, the regulatory molecule interacts with TATA binding protein.

[0522] In certain embodiments, the regulatory molecule interacts with transcription factor II D.

[0523] In certain embodiments, the regulatory molecule comprises a CDK9 subunit.

[0524] In certain embodiments, the regulatory molecule is P-TEFb.


[0525] In certain embodiments, X binds to the regulatory molecule but does not inhibit the activity of the regulatory molecule. In certain embodiments, X binds to the regulatory molecule and inhibits the activity of the regulatory molecule. In certain embodiments, X binds to the regulatory molecule and increases the activity of the regulatory molecule.

[0526] In certain embodiments, X binds to the active site of the regulatory molecule. In certain embodiments, X binds to a regulatory site of the regulatory molecule.


[0527] In certain embodiments, the recruiting moiety is chosen from a CDK-9 inhibitor, a cyclin T1 inhibitor, and a PRC2 inhibitor.

[0528] In certain embodiments, the recruiting moiety is a CDK-9 inhibitor. In certain embodiments, the CDK-9 inhibitor is chosen from flavopiridol, CR8, indirubin-3'-monoxime, a 5-fluoro-N2,N4-diphenylpyrimidine-2,4-diamine, a 4-(thiazol-5-yl)-2-(phenylamino)pyrimidine, TG02, CDKI-73, a 2,4,5-trisubstituted pyrimidine derivatives, LCD000067, Wogonin, BAY-1000394 (Roniciclib), AZD5438, and DRB (F Morales et al. "Overview of CDK9 as a target in cancer research", Cell Cycle 2016, 15(4), 519-527, and references therein).


-continued




CDKI-73




LCD000067




Wogonin




BAY 1000394




AZD5438



DRB




CR8




TG02

[0529] In certain embodiments, the regulatory molecule is a histone demethylase. In certain embodiments, the histone demethylase is a lysine demethylase. In certain embodiments, the lysine demethylase is KDM5B. In certain embodiments, the recruiting moiety is a KDM5B inhibitor. In certain embodiments, the KDM5B inhibitor is AS-8351 (N. Cao, Y. Huang, J. Zheng, et al., “Conversion of human fibroblasts into functional cardiomyocytes by small molecules”, *Science* 2016, 352(6290), 1216-1220, and references therein.)




RG108




AS-8351

[0530] In certain embodiments, the regulatory molecule is the complex between the histone lysine methyltransferases (“HKMT”) GLP and G9A (“GLP/G9A”). In certain embodiments, the recruiting moiety is a GLP/G9A inhibitor. In certain embodiments, the GLP/G9A inhibitor is BIX-01294 (Chang Y, “Structural basis for G9a-like protein lysine methyltransferase inhibition by BIX-01294”, *Nature Struct. Mol. Biol.* 2009, 16, 312-317, and references therein).




1149




BIX-01294

[0531] In certain embodiments, the regulatory molecule is a DNA methyltransferase (“DNMT”). In certain embodiments, the regulatory moiety is DNMT1. In certain embodiments, the recruiting moiety is a DNMT1 inhibitor. In certain embodiments, the DNMT1 inhibitor is chosen from RG108 and the RG108 analogues 1149, T1, and G6. (B Zhu et al. *BioorgMed Chem* 2015, 23(12), 2917-2927 and references therein).




T1




G6

[0532] In certain embodiments, the recruiting moiety is a PRC1 inhibitor. In certain embodiments, the PRC1 inhibitor is chosen from UNC4991, UNC3866, and UNC3567 (J I Stuckey et al. *Nature Chem Biol* 2016, 12(3), 180-187 and references therein; K D Barnash et al. *ACS Chem. Biol.* 2016, 11(9), 2475-2483, and references therein).


UNC4991




UNC3866




UNC3567




[0533] In certain embodiments, the recruiting moiety is a PRC2 inhibitor. In certain embodiments, the PRC2 inhibitor is chosen from A-395, MS37452, MAK683, DZNep, EPZ005687, Eli, GSK126, and UNC1999 (Konze K D ACS Chem Biol 2013, 8(6), 1324-1334, and references therein).




A-395



MS37452




MAK683




DZNep


-continued




EPZ005687



Eli-1




GSK126



UNC1999

[0534] In certain embodiments, the recruiting moiety is rohitukine or a derivative of rohitukine.

[0535] In certain embodiments, the recruiting moiety is DB08045 or a derivative of DB08045.




DB08045

[0536] In certain embodiments, the recruiting moiety is A-395 or a derivative of A-395.

[0537] In certain embodiments, the regulatory molecule is chosen from a bromodomain-containing protein, a nucleosome remodeling factor (NURF), a bromodomain PHD finger transcription factor (BPTF), a ten-eleven translocation enzyme (TET), methylcytosine dioxygenase (TET1), a DNA demethylase, a helicase, an acetyltransferase, and a histone deacetylase ("HDAC").

[0538] In certain embodiments, the regulatory molecule is a bromodomain-containing protein chosen from BRD2, BRD3, BRD4, and BRDT.



[0539] In certain embodiments, the regulatory molecule is BRD4. In certain embodiments, the recruiting moiety is a BRD4 activator. In certain embodiments, the BRD4 activator is chosen from JQ-1, OTX015, RVX208 acid, and RVX208 hydroxyl.



JQ1

-continued


OTX015




RVX208

[0540] In certain embodiments, the regulatory molecule is BPTF. In certain embodiments, the recruiting moiety is a BPTF activator. In certain embodiments, the BPTF activator is AU1.

AU1




[0541] In certain embodiments, the regulatory molecule is histone acetyltransferase ("HAT"). In certain embodiments, the recruiting moiety is a HAT activator. In certain embodiments, the HAT activator is a oxopiperazine helix mimetic OHM. In certain embodiments, the HAT activator is selected from OHM1, OHM2, OHM3, and OHM4 (B B Lao et al., PNAS USA 2014, 111(21), 7531-7536). In certain embodiments, the HAT activator is OHM4.




OHM1: R¹ = *i*-Bu; R² = *i*-Bu; R³ = Me; R⁴ = CH₂CH₂CONH₂.
 OHM2: R¹ = *i*-Bu; R² = *i*-Bu; R³ = Me; R⁴ = Me.
 OHM3: R¹ = *i*-Bu; R² = Me; R³ = Me; R⁴ = CH₂CH₂CONH₂.
 OHM4: R¹ = Me; R² = Me; R³ = Me; R⁴ = Me.

[0542] In certain embodiments, the regulatory molecule is histone deacetylase (“HDAC”). In certain embodiments, the recruiting moiety is an HDAC activator. In certain embodiments, the HDAC activator is chosen from SAHA and 109 (Soragni E Front. Neurol. 2015, 6, 44, and references therein).




SAHA

109



[0543] In certain embodiments, the regulatory molecule is histone deacetylase (“HDAC”). In certain embodiments, the recruiting moiety is an HDAC inhibitor. In certain embodiments, the HDAC inhibitor is an inositol phosphate.


[0544] In certain embodiments, the regulatory molecules is O-linked β-N-acetylglucosamine transferase (“OGT”). In certain embodiments, the recruiting moiety is an OGT activator. In certain embodiments, the OGT activator is chosen from ST045849, ST078925, and ST060266 (Itkonen HM, “Inhibition of O-GlcNAc transferase activity reprograms prostate cancer cell metabolism”, Oncotarget 2016, 7(11), 12464-12476).




ST045849

-continued

ST078925




ST060266




[0545] In certain embodiments, the regulatory molecule is chosen from host cell factor 1 (“HCF1”) and octamer binding transcription factor (“OCT1”). In certain embodiments, the recruiting moiety is chosen from an HCF1 activator and an OCT1 activator. In certain embodiments, the recruiting moiety is chosen from VP16 and VP64.

[0546] In certain embodiments, the regulatory molecule is chosen from CBP and P300. In certain embodiments, the recruiting moiety is chosen from a CBP activator and a P300 activator. In certain embodiments, the recruiting moiety is CTPB.



CTPB

[0547] In certain embodiments, the regulatory molecule is P300/CBP-associated factor (“PCAF”). In certain embodiments, the recruiting moiety is a PCAF activator. In certain embodiments, the PCAF activator is embelin.



[0548] In certain embodiments, the regulatory molecule modulates the rearrangement of histones.

[0549] In certain embodiments, the regulatory molecule modulates the glycosylation, phosphorylation, alkylation, or acylation of histones.

[0550] In certain embodiments, the regulatory molecule is a transcription factor.

[0551] In certain embodiments, the regulatory molecule is an RNA polymerase.

[0552] In certain embodiments, the regulatory molecule is a moiety that regulates the activity of RNA polymerase.

[0553] In certain embodiments, the regulatory molecule interacts with TATA binding protein.

[0554] In certain embodiments, the regulatory molecule interacts with transcription factor II D.

[0555] In certain embodiments, the regulatory molecule comprises a CDK9 subunit.


[0556] In certain embodiments, the regulatory molecule is P-TEFb.

[0557] In certain embodiments, the recruiting moiety binds to the regulatory molecule but does not inhibit the activity of the regulatory molecule. In certain embodiments, the recruiting moiety binds to the regulatory molecule and inhibits the activity of the regulatory molecule. In certain embodiments, the recruiting moiety binds to the regulatory molecule and increases the activity of the regulatory molecule.







[0558] In certain embodiments, the recruiting moiety binds to the active site of the regulatory molecule. In certain embodiments, the recruiting moiety binds to a regulatory site of the regulatory molecule.

[0559] In certain embodiments, the recruiting moiety is chosen from a CDK-9 inhibitor, a cyclin T1 inhibitor, and a PRC2 inhibitor.


[0560] In certain embodiments, the recruiting moiety is a CDK-9 inhibitor. In certain embodiments, the CDK-9 inhibitor is chosen from flavopiridol, CR8, indirubin-3'-monoxime, a 5-fluoro-N2,N4-diphenylpyrimidine-2,4-diamine, a 4-(thiazol-5-yl)-2-(phenylamino)pyrimidine, TG02, CDKI-73, a 2,4,5-trisubstituted pyrimidine derivatives, LCD000067, Wogonin, BAY-1000394 (Roniciclib), AZD5438, and DRB (F Morales et al. "Overview of CDK9 as a target in cancer research", *Cell Cycle* 2016, 15(4), 519-527, and references therein).



-continued




-continued



DRB


ments, the recruiting moiety is a DNMT1 inhibitor. In certain embodiments, the DNMT1 inhibitor is chosen from RG108 and the RG108 analogues 1149, T1, and G6. (B Zhu et al. *Bioorg Med Chem* 2015, 23(12), 2917-2927 and references therein).

RG108




[0561] In certain embodiments, the regulatory molecule is a histone demethylase. In certain embodiments, the histone demethylase is a lysine demethylase. In certain embodiments, the lysine demethylase is KDM5B. In certain embodiments, the recruiting moiety is a KDM5B inhibitor. In certain embodiments, the KDM5B inhibitor is AS-8351 (N. Cao, Y. Huang, J. Zheng, et al., "Conversion of human fibroblasts into functional cardiomyocytes by small molecules", *Science* 2016, 352(6290), 1216-1220, and references therein.)

AS-8351





1149




T1

[0562] In certain embodiments, the regulatory molecule is the complex between the histone lysine methyltransferases ("HKMT") GLP and G9A ("GLP/G9A"). In certain embodiments, the recruiting moiety is a GLP/G9A inhibitor. In certain embodiments, the GLP/G9A inhibitor is BIX-01294 (Chang Y, "Structural basis for G9a-like protein lysine methyltransferase inhibition by BIX-01294", *Nature Struct. Mol. Biol.* 2009, 16, 312-317, and references therein).

BIX-01294



G6




[0563] In certain embodiments, the regulatory molecule is a DNA methyltransferase ("DNMT"). In certain embodiments, the regulatory moiety is DNMT1. In certain embodi-


[0564] In certain embodiments, the recruiting moiety is a PRC1 inhibitor. In certain embodiments, the PRC1 inhibitor is chosen from UNC4991, UNC3866, and UNC3567 (J I

Stuckey et al. *Nature Chem Biol* 2016, 12(3), 180-187 and references therein; K D Barnash et al. *ACS Chem. Biol.* 2016, 11(9), 2475-2483, and references therein).


UNC4991




UNC3866




UNC3567




[0565] In certain embodiments, the recruiting moiety is a PRC2 inhibitor. In certain embodiments, the PRC2 inhibitor is chosen from A-395, MS37452, MAK683, DZNep, EPZ005687, Eli, GSK126, and UNC1999 (Konze K D ACS Chem Biol 2013, 8(6), 1324-1334, and references therein).




A-395



MS37452




MAK683




DZNep


-continued




EPZ005687



Eli-1




GSK126



UNC1999

[0566] In certain embodiments, the recruiting moiety is rohitukine or a derivative of rohitukine.

[0567] In certain embodiments, the recruiting moiety is DB08045 or a derivative of DB08045.



[0568] In certain embodiments, the recruiting moiety is A-395 or a derivative of A-395.

Oligomeric Backbone and Linker

[0569] The Oligomeric backbone contains a linker that connects the first terminus and the second terminus and brings the regulatory molecule in proximity to the target gene to modulate gene expression.

[0570] The length of the linker depends on the type of regulatory protein and also the target gene. In some embodiments, the linker has a length of less than about 50 Angstroms. In some embodiments, the linker has a length of about 20 to 30 Angstroms.

[0571] In some embodiments, the linker comprises between 5 and 50 chain atoms.

[0572] In some embodiments, the linker comprises a multimer having 2 to 50 spacing moieties, wherein the spacing moiety is independently selected from the group consisting of $-(\text{CR}^{3a}\text{R}^{3b})_x\text{O}-$, $-(\text{CR}^{3a}\text{R}^{3b})_x\text{NR}^{4a}-$, $-(\text{CR}^{3a}\text{R}^{3b})_x\text{CH}=\text{CH}-(\text{CR}^{3a}\text{R}^{3b})_x\text{O}-$, optionally substituted C_{1-12} alkyl, optionally substituted C_{2-10} alkenyl, optionally substituted C_{2-10} alkynyl, optionally substituted C_{6-10} arylene, optionally substituted C_{3-7} cycloalkylene, optionally substituted 5- to 10-membered heteroarylene, optionally substituted 4- to 10-membered heterocycloalkylene, amino acid residue, $-\text{O}-$, $-\text{C}(\text{O})\text{NR}^{4a}-$, $\text{NR}^{4a}\text{C}(\text{O})-$, $-\text{C}(\text{O})-$, $-\text{NR}^1-$, $-\text{C}(\text{O})\text{O}-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O})-$, $-\text{SO}_2-$, $-\text{SO}_2\text{NR}^{4a}-$, $-\text{NR}^{4a}\text{SO}_2-$, and $-\text{P}(\text{O})\text{OH}-$, and any combinations thereof; wherein

[0573] each x is independently 2-4;

[0574] each y is independently 1-10;

[0575] each R^{3a} and R^{3b} are independently selected from hydrogen, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, optionally substituted alkylamide, sulfonyl, optionally substituted thioalkoxy, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocyclyl; and

[0576] each R^{4a} is independently a hydrogen or an optionally substituted C_{1-6} alkyl.

[0577] In some embodiments, the oligomeric backbone comprises $-(\text{T}^1\text{-V}^1)_a-(\text{T}^2\text{V}^2)_b-(\text{T}^3\text{V}^3)_c-(\text{T}^4\text{V}^4)_d-(\text{T}^5\text{V}^5)_e-$,

[0578] wherein a , b , c , d and e are each independently 0 or 1, and where the sum of a , b , c , d and e is 1 to 5;

[0579] T^1 , T^2 , T^3 , T^4 and T^5 are each independently selected from an optionally substituted ($\text{C}_1\text{-C}_{12}$)alkylene, optionally substituted alkenylene, optionally substituted alkynylene, $(\text{EA})_w$, $(\text{EDA})_m$, $(\text{PEG})_n$, (modified $\text{PEG})_n$, $(\text{AA})_p$, $-(\text{CR}^{2a}\text{OH})_n-$, optionally substituted ($\text{C}_6\text{-C}_{10}$) arylene, optionally substituted C_{3-7} cycloalkylene, optionally substituted 5- to 10 membered heteroarylene, optionally substituted 4- to 10-membered heterocycloalkylene, an acetal group, a disulfide, a hydrazine, a carbohydrate, a beta-lactam, and an ester,

[0580] (a) w is an integer from 1 to 20;


[0581] (b) m is an integer from 1 to 20;

[0582] (c) n is an integer from 1 to 30;


[0583] (d) p is an integer from 1 to 20;

[0584] (e) h is an integer from 1 to 12;

[0585] (f) EA has the following structure



[0586] (g) EDA has the following structure:



[0587] wherein each q is independently an integer from 1 to 6, each x is independently an integer from 1 to 4, and each r is independently 0 or 1;

[0588] (h) (PEG), has the structure of $-(\text{CR}^{2a}\text{R}^{2b}-\text{CR}^{2a}\text{R}^{2b}-\text{O})_n-\text{CR}^{2a}\text{R}^{2b}-$;

[0589] (i) (modified PEG), has the structure of replacing at least one $-(\text{CR}^{2a}\text{R}^{2b}-\text{CR}^{2a}\text{R}^{2b}-\text{O})-$ in (PEG), with $-(\text{CH}_2-\text{CR}^{2a}=\text{CR}^a-\text{CH}_2-\text{O})-$ or $-(\text{CR}^{2a}\text{R}^{2b}-\text{CR}^{2a}\text{R}^{2b}-\text{S})-$;

[0590] (j) AA is an amino acid residue;

[0591] (k) V^1 , V^2 , V^3 , V^4 and V^5 are each independently selected from the group consisting of a bond, $\text{CO}-$, $-\text{NR}^{1a}-$, $-\text{CONR}^{1a}-$, $-\text{NR}^{1a}\text{CO}-$, $-\text{CONR}^{1a}\text{C}_{1-4}\text{alkyl}-$, $-\text{NR}^{1a}\text{CO}-\text{C}_{1-4}\text{alkyl}-$, $-\text{C}(\text{O})\text{O}-$, $-\text{OC}(\text{O})-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O})-$, $-\text{SO}_2-$, $-\text{SO}_2\text{NR}^{1a}-$, $-\text{NR}^{1a}\text{SO}_2-$ and $-\text{P}(\text{O})\text{OH}-$;

[0592] (l) each R^{1a} is independently hydrogen or an optionally substituted C_{1-6} alkyl; and


[0593] (m) each R^{2a} and R^{2b} are independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, halogen, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substi-

tuted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl.

[0594] In some embodiments, the a, b, c, d and e are each independently 0 or 1, where the sum of a, b, c, d and e is 1. In some embodiments, the a, b, c, d and e are each independently 0 or 1, where the sum of a, b, c, d and e is 2. In some embodiments, the a, b, c, d and e are each independently 0 or 1, where the sum of a, b, c, d and e is 3. In some embodiments, the a, b, c, d and e are each independently 0 or 1, where the sum of a, b, c, d and e is 4. In some embodiments, the a, b, c, d and e are each independently 0 or 1, where the sum of a, b, c, d and e is 5.

[0595] In some embodiments, n is 3-9. In some embodiments, n is 4-8. In some embodiments, n is 5 or 6.


[0596] In some embodiments, T¹, T², T³, and T⁴, and T⁵ are each independently selected from (C₁-C₁₂)alkyl, substituted (C₁-C₁₂)alkyl, (EA)_w, (EDA)_m, (PEG)_n, (modified PEG)_n, (AA)_p, —(CR^{2a}OH)_n—, phenyl, substituted phenyl, piperidin-4-amino (P4A), para-amino-benzoyloxycarbonyl (PABC), meta-amino-benzoyloxycarbonyl (MABC), para-amino-benzoyloxy (PABO), meta-amino-benzoyloxy (MABO), para-aminobenzyl, an acetal group, a disulfide, a hydrazine, a carbohydrate, a beta-lactam, an ester, (AA)_p-MABC-(AA)_p, (AA)_p-MABO-(AA)_p, (AA)_p-PABO-(AA)_p and (AA)_p-PABC-(AA)_p. In some embodiments, piperidin-4-amino (P4A) is



wherein R^{1a} is H or C₁₋₆ alkyl.


[0597] In some embodiments, T¹, T², T³, T⁴ and T⁵ are each independently selected from (C₁-C₁₂)alkyl, substituted (C₁-C₁₂)alkyl, (EA)_w, (EDA)_m, (PEG)_n, (modified PEG)_n,

(AA)_p, —(CR^{2a}OH)_n—, optionally substituted (C₆-C₁₀) arylene, 4-10 membered heterocycloalkene, optionally substituted 5-10 membered heteroarylene. In some embodiments, EA has the following structure:



and

[0598] EDA has the following structure:



[0599] In some embodiments, x is 2-3 and q is 1-3 for EA and EDA. In some embodiments, R^{1a} is H or C₁₋₆ alkyl.

[0600] In some embodiments, T⁴ or T⁵ is an optionally substituted (C₆-C₁₀) arylene.


[0601] In some embodiments, T⁴ or T⁵ is phenylene or substituted phenylene. In some embodiments, T⁴ or T⁵ is phenylene or phenylene substituted with 1-3 substituents selected from —C₁₋₆ alkyl, halogen, OH or amine. In some embodiments, T⁴ or T⁵ is 5-10 membered heteroarylene or substituted heteroarylene. In some embodiments, T⁴ or T⁵ is 4-10 membered heterocyclylene or substituted heterocyclylene. In some embodiments, T⁴ or T⁵ is heteroarylene or heterocyclylene optionally substituted with 1-3 substituents selected from —C₁₋₆ alkyl, halogen, OH or amine.

[0602] In some embodiments, T¹, T², T³, T⁴ and T⁵ and V¹, V², V³, V⁴ and V⁵ are selected from the following

TABLE 6


T ¹	V ¹	T ²	V ²	T ³	V ³	T ⁴	V ⁴	T ⁵	V ⁵
(C ₁ -C ₁₂) alkylene	CONR ^{1a}	(EA) _w	CO	(PEG) _n	NR ¹¹ CO	—	—	—	—
(C ₁ -C ₁₂) alkylene	CONR ^{1a}	(EA) _w	CO	(PEG) _n	O	arylene	NR ¹¹ CO	—	—
(C ₁ -C ₁₂) alkylene	CONR ^{1a}	(EA) _w	CO	(PEG) _n	O	Subst. arylene	NR ¹¹ CO	—	—
(C ₁ -C ₁₂) alkylene	CONR ^{1a}	(EA) _w	CO	(PEG) _n	O	NR ¹¹ CO	(C ₁ -C ₁₂) alkyl	Subst. arylene	NR ¹¹ CO
(C ₁ -C ₁₂) alkylene	CONR ^{1a}	(EA) _w	CO	(C ₁ -C ₁₂) alkyl	NR ¹¹ CO- C ₁₋₄ alkyl	Subst. arylene	NR ¹¹	—	—
(C ₁ -C ₁₂) alkylene	CONR ^{1a}	(EA) _w	CO	(PEG) _n	O	Subst. arylene	—	—	—
(PEG) _n	CONR ^{1a} - C ₁₋₄ alkyl	—	—	—	—	—	—	—	—
(EA) _w	CO	(C ₁ -C ₁₂) alkyl	CONR ¹¹ - C ₁₋₄ alkyl	—	—	—	—	—	—
(C ₁ -C ₁₂) alkylene	CONR ^{1a}	(EA) _w	CO	(PEG) _n	NR ¹¹ CO- C ₁₋₄ alkyl	—	—	—	—
(EA) _w	CO	(PEG) _n	O	phenyl	NR ¹¹ CO- C ₁₋₄ alkyl	—	—	—	—
(C ₁ -C ₁₂) alkylene	CONR ^{1a}	(PEG) _n	CO	—	—	—	—	—	—
(C ₁ -C ₁₂) alkylene	CONR ^{1a}	(EA) _w	CO	modifd. (PEG) _n	O	arylene	NR ¹¹ CO	—	—

[0603] In some embodiments, the linker comprises



or any combinations thereof, wherein r is an integer between 1 and 10, preferably between 3 and 7; and X is O, S, or NR^{1a} . In some embodiments, X is O or NR^{1a} . In some embodiments, X is O.


[0604] In some embodiments, the linker comprise a



or any combinations thereof; wherein at least one $-(\text{CH}_2-\text{CH}_2-\text{O})-$ is replaced with $-((\text{CR}^{1a}\text{R}^{1b})_x-\text{CH}=\text{CH}-\text{CH}_2-\text{O})-$, or any combinations thereof; W' is absent, $(\text{CH}_2)_{1-5}$, $-(\text{CH}_2)_{1-5}\text{O}$, $(\text{CH}_2)_{1-5}\text{C}(\text{O})\text{NH}-$, $(\text{CH}_2)_{1-5}\text{O}$, $(\text{CH}_2)_{1-5}\text{C}(\text{O})\text{NH}-(\text{CH}_2)_{1-5}$, $-(\text{CH}_2)_{1-5}\text{NHC}(\text{O})-(\text{CH}_2)_{1-5}\text{O}$, or $-(\text{CH}_2)_{1-5}\text{NHC}(\text{O})-(\text{CH}_2)_{1-5}$; E^3 is an optionally substituted C_{6-10} arylene group, optionally substituted 4-10 membered heterocycloalkylene, or optionally substituted 5-10 membered heteroarylene; X is O, S, or NH; each R^{1a} and R^{1b} are independently H or C_{1-6} alkyl; r is an integer between 1 and 10; and x is an integer between 1 and 15. In some embodiments, X is O. In some embodiments, X is NH. In some embodiments, E^3 is a C_{6-10} arylene group optionally substituted with 1-3 substituents selected from $-\text{C}_{1-6}$ alkyl, halogen, OH or amine.


[0605] In some embodiments, E^3 is a phenylene or substituted phenylene.

[0606] In some embodiments, the linker comprise a



[0607] In some embodiments, the linker comprises $-\text{X}(\text{CH}_2)_m(\text{CH}_2\text{CH}_2\text{O})_n-$, wherein X is $-\text{O}-$, $-\text{NH}-$, or $-\text{S}-$, wherein m is 0 or greater and n is at least 1.

[0608] In some embodiments, the linker comprises




following the second terminus, wherein R_c is selected from a bond, $-\text{N}(\text{R}^{1a})-$, $-\text{O}-$, and $-\text{S}-$; R_d is selected from $-\text{N}(\text{R}^{1a})-$, $-\text{O}-$, and $-\text{S}-$; and R_e is independently selected from hydrogen and optionally substituted C_{1-6} alkyl

[0609] In some embodiments, the linker comprises one or more structures selected from



$-\text{C}_{1-12}$ alkyl, arylene, cycloalkylene, heteroarylene, heterocycloalkylene, $-\text{O}-$, $-\text{C}(\text{O})\text{NR}^{1a}-$, $-\text{C}(\text{O})-$, $-\text{NR}^{1a}-$, $-(\text{CH}_2\text{CH}_2\text{CH}_2\text{O})_y-$, and $-(\text{CH}_2\text{CH}_2\text{CH}_2\text{NR}^{1a})_y-$, wherein each d and y are independently 1-10, and each R^{1a} is independently hydrogen or C_{1-6} alkyl. In some embodiments, d is 4-8.

[0610] In some embodiments, the linker comprises



and each d is independently 3-7. In some embodiments, d is 4-6.

[0611] In some embodiments, the linker comprises $\text{N}(\text{R}^{1a})(\text{CH}_2)_x\text{N}(\text{R}^{1b})(\text{CH}_2)_y\text{N}-$, wherein R^{1a} and R^{1b} are each independently selected from hydrogen or optionally substituted C_{1-6} alkyl; and each x is independently an integer in the range of 1-6.

[0612] In some embodiments, the linker comprises the linker comprises $-(\text{CH}_2-\text{C}(\text{O})\text{N}(\text{R}'')-(\text{CH}_2)_q-\text{N}(\text{R}')-(\text{CH}_2)_q-\text{N}(\text{R}'')\text{C}(\text{O})-(\text{CH}_2)_x-\text{C}(\text{O})\text{N}(\text{R}'')-\text{A}-, -(\text{CH}_2)_x-\text{C}(\text{O})\text{N}(\text{R}'')-(\text{CH}_2\text{CH}_2\text{O})_y-(\text{CH}_2)_x-\text{C}(\text{O})\text{N}(\text{R}'')-\text{A}-, -\text{C}(\text{O})\text{N}(\text{R}'')-(\text{CH}_2)_q-\text{N}(\text{R}')-(\text{CH}_2)_q-\text{N}(\text{R}'')\text{C}(\text{O})-(\text{CH}_2)_x-\text{A}-, -(\text{CH}_2)_x-\text{O}-(\text{CH}_2\text{CH}_2\text{O})_y-(\text{CH}_2)_x-\text{N}(\text{R}'')\text{C}(\text{O})-(\text{CH}_2)_x-\text{A}-, or $-\text{N}(\text{R}'')\text{C}(\text{O})-(\text{CH}_2)-\text{C}(\text{O})\text{N}(\text{R}'')-(\text{CH}_2)_x-\text{O}-(\text{CH}_2\text{CH}_2\text{O})_y-(\text{CH}_2)_x-\text{A}-$; wherein R' is methyl; R'' is hydrogen; each x and y are independently an integer from 1 to 10; each q is independently an integer from 2 to 10; and each A is independently selected from a bond, an optionally substituted C_{1-12} alkyl, an optionally substituted C_{6-10} arylene, optionally substituted C_{3-7} cycloalkylene, optionally substituted 5- to 10-membered heteroarylene, and optionally substituted 4- to 10-membered heterocycloalkylene.$

[0613] In some embodiments, the linker is joined with the first terminus with a group selected from $-\text{CO}-$, $-\text{NR}^{1a}-$, $-\text{CONR}^{1a}-$, $-\text{NR}^{1a}\text{CO}-$, $-\text{CONR}^{1a}\text{C}_{1-4}\text{al}-$

kyl-, $-\text{NR}^{1a}\text{CO}-\text{C}_{1-4}\text{alkyl}-$, $-\text{C}(\text{O})\text{O}-$, $-\text{OC}(\text{O})-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O})-$, $-\text{SO}_2-$, $-\text{SO}_2\text{NR}^{1a}-$, $-\text{NR}^1\text{SO}_2-$, $-\text{P}(\text{O})\text{OH}-$, $-\text{((CH}_2\text{)}_x\text{O)}-$, $-\text{((CH}_2\text{)}_y\text{NR}^{1a})-$, optionally substituted $-\text{C}_{1-12}$ alkylene, optionally substituted C_{2-10} alkenylene, optionally substituted C_{2-10} alkynylene, optionally substituted C_{6-10} arylene, optionally substituted C_{3-7} cycloalkylene, optionally substituted 5- to 10-membered heteroarylene, and optionally substituted 4- to 10-membered heterocycloalkylene, wherein each x is independently 1-4, each y is independently 1-4, and each R^{1a} is independently a hydrogen or optionally substituted C_{1-6} alkyl.

[0614] In some embodiments, the linker is joined with the first terminus with a group selected from $-\text{CO}-$, $-\text{NR}^{1a}-$, C_{1-12} alkyl, $-\text{CONR}^{1a}-$, and $-\text{NR}^{1a}\text{CO}-$.

[0615] In some embodiments, the linker is joined with second terminus with a group selected from $-\text{CO}-$, $-\text{NR}^{1a}-$, $-\text{CONR}^{1a}-$, $-\text{NR}^{1a}\text{CO}-$, $-\text{CONR}^{1a}\text{C}_{1-4}\text{alkyl}-$, $-\text{NR}^{1a}\text{CO}-\text{C}_{1-4}\text{alkyl}-$, $-\text{C}(\text{O})\text{O}-$, $-\text{OC}(\text{O})-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O})-$, $-\text{SO}_2-$, $-\text{SO}_2\text{NR}^{1a}-$, $-\text{NR}^1\text{SO}_2-$, $-\text{P}(\text{O})\text{OH}-$, $-\text{((CH}_2\text{)}_x\text{O)}-$, $-\text{((CH}_2\text{)}_y\text{NR}^{1a})-$, optionally substituted $-\text{C}_{1-12}$ alkylene, optionally substituted C_{2-10} alkenylene, optionally substituted C_{2-10} alkynylene, optionally substituted C_{6-10} arylene, optionally substituted C_{3-7} cycloalkylene, optionally substituted 5- to 10-membered heteroarylene, and optionally substituted 4- to 10-membered heterocycloalkylene, wherein each x is independently 1-4, each y is independently 1-4, and each R^{1a} is independently a hydrogen or optionally substituted C_{1-6} alkyl.

[0616] In some embodiments, the linker is joined with second terminus with a group selected from $-\text{CO}-$, $-\text{NR}^{1a}-$, $-\text{CONR}^{1a}-$, $-\text{NR}^{1a}\text{CO}-$, $-\text{((CH}_2\text{)}_x\text{O)}-$, $-\text{((CH}_2\text{)}_y\text{NR}^{1a})-$, $-\text{O}-$, optionally substituted $-\text{C}_{1-12}$ alkyl, optionally substituted C_{6-10} arylene, optionally substituted C_{3-7} cycloalkylene, optionally substituted 5- to 10-membered heteroarylene, and optionally substituted 4- to 10-membered heterocycloalkylene, wherein each x is independently 1-4, each y is independently 1-4, and each R^1 is independently a hydrogen or optionally substituted C_{1-6} alkyl.

Cell-Penetrating Ligand


[0617] In certain embodiments, the compounds comprise a cell-penetrating ligand moiety.

[0618] In certain embodiments, the cell-penetrating ligand moiety is a polypeptide.

[0619] In certain embodiments, the cell-penetrating ligand moiety is a polypeptide containing fewer than 30 amino acid residues.

[0620] In certain embodiments, the polypeptide is chosen from any one of SEQ ID NO. 1 to SEQ ID NO. 37, inclusive.

[0621] In some embodiments, the second terminus does not comprise a structure of Formula (C-11):



(C-11)

[0622] wherein:


[0623] each of A^{1p} and B^{1p} is independently an optionally substituted aryl or heteroaryl ring;

[0624] X^{1p} is CH or N;

[0625] R^{1p} is hydrogen, halogen, or an optionally substituted C_{1-6} alkyl group; and

[0626] R^{2p} is an optionally substituted C_{1-6} alkyl, cycloalkyl, C_{6-10} aryl, or heteroaryl.

[0627] In some embodiments, the protein binding moiety does not have the structure of Formula (C-12):



(C-12)

[0628] wherein:

[0629] R_{1q} is a hydrogen or an optionally substituted alkyl, hydroxyalkyl, aminoalkyl, alkoxyalkyl, halogenated alkyl, hydroxyl, alkoxy, or $-\text{COOR}_{4q}$;

[0630] R_{4q} is hydrogen, or an optionally substituted aryl, aralkyl, cycloalkyl, heteroaryl, heteroaralkyl, heterocycloalkyl, alkyl, alkenyl, alkynyl, or cycloalkylalkyl group, optionally containing one or more heteroatoms;

[0631] R_{2q} is an optionally substituted aryl, alkyl, cycloalkyl, or aralkyl group;

[0632] R_{3q} is hydrogen, halogen, or an optionally substituted alkyl group, preferably $(\text{CH}_2)_x-\text{C}(\text{O})\text{N}(\text{R}_{20})(\text{R}_{21})$, or $(\text{CH}_2)_x-\text{N}(\text{R}_{20})-\text{C}(\text{O})\text{R}_{21}$; or halogenated alkyl group;

[0633] wherein x is an integer from 1 to 10; and R_{20} and R_{21} are each independently hydrogen or C_{1-6} alkyl group, preferably R_{20} is hydrogen and R_{21} is methyl; and

[0634] Ring E is an optionally substituted aryl or heteroaryl group.

[0635] Also provided are embodiments wherein any molecule disclosed above, including molecules of Formulas I-VIII, are singly, partially, or fully deuterated. Methods for accomplishing deuterium exchange for hydrogen are known in the art.

[0636] Also provided are embodiments wherein any embodiment above may be combined with any one or more of these embodiments, provided the combination is not mutually exclusive.

[0637] As used herein, two embodiments are “mutually exclusive” when one is defined to be something which is different than the other. For example, an embodiment wherein two groups combine to form a cycloalkyl is mutually exclusive with an embodiment in which one group is ethyl the other group is hydrogen. Similarly, an embodiment wherein one group is CH_2 is mutually exclusive with an embodiment wherein the same group is NH.

Method of Treatment

[0638] The present disclosure also relates to a method of modulating the transcription of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1, the method comprising the step of contacting dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1 with a molecule as described herein.

[0639] The cell phenotype, cell proliferation, transcription of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1, production of mRNA from transcription of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1, translation of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1, change in biochemical output produced by the protein coded by dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1, or noncovalent binding of the protein coded by dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1 with a natural binding partner may be monitored. Such methods may be modes of treatment of disease, biological assays, cellular assays, biochemical assays, or the like. In some embodiments, the gene is dmpk. In some embodiments, the gene is atxn1.

[0640] In some embodiments, the gene is atxn2. In some embodiments, the gene is atxn3. In some embodiments, the gene is cacna1a. In some embodiments, the gene is atxn7. In some embodiments, the gene is ppp2r2. In some embodiments, the gene is tbp. In some embodiments, the gene is htt. In some embodiments, the gene is jph3. In some embodiments, the gene is ar. In some embodiments, the gene is atn1.

[0641] Also provided herein is a method of treatment of a disease mediated by transcription of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1 comprising the administration of a therapeutically effective amount of a molecule as disclosed herein, or a salt thereof, to a patient in need thereof.

[0642] In certain embodiments, the disease is chosen from DM1, spinocerebellar ataxia, Huntington's disease, a Huntington's disease-like syndrome, spinobulbar muscular atrophy, and dentatorubral-pallidoluysian atrophy.

[0643] In certain embodiments, the disease is chosen from DM1.

[0644] In certain embodiments, the disease is spinocerebellar ataxia. In certain embodiments, the spinocerebellar ataxia is chosen from SCA1, SCA2, SCA3, SCA6, SCA7, SCA12, and SCA17. In certain embodiments, the spinocerebellar ataxia is chosen from SCA1, SCA2, SCA3, SCA6, SCA7, and SCA17.

[0645] In certain embodiments, the disease is chosen from Huntington's disease and a Huntington's disease-like syndrome. In certain embodiments, the disease is chosen from Huntington's disease and Huntington's disease-like 2 syndrome.

[0646] In certain embodiments, the disease is spinobulbar muscular atrophy.

[0647] In certain embodiments, the disease is dentatorubral-pallidoluysian atrophy.

[0648] Also provided herein is a molecule as disclosed herein for use as a medicament.

[0649] Also provided herein is a molecule as disclosed herein for use as a medicament for the treatment of a disease mediated by transcription of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1.

[0650] Also provided is the use of a molecule as disclosed herein as a medicament.

[0651] Also provided is the use of a molecule as disclosed herein as a medicament for the treatment of a disease mediated by transcription of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1.

[0652] Also provided is a molecule as disclosed herein for use in the manufacture of a medicament for the treatment of a disease mediated by transcription of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1.

[0653] Also provided is the use of a molecule as disclosed herein for the treatment of a disease mediated by transcription of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1.

[0654] Also provided herein is a method of modulation of transcription of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1 comprising contacting dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1 with a molecule as disclosed herein, or a salt thereof.

[0655] Also provided herein is a method for achieving an effect in a patient comprising the administration of a therapeutically effective amount of a molecule as disclosed herein, or a salt thereof, to a patient, wherein the effect is chosen from muscular atrophy, ataxia, fasciculations, dementia, dysarthria, and dysphagia.

[0656] Also provided herein is a method for the prevention or treatment of a disease or condition mediated by or associated with the transcription of TCF4.

[0657] A method of modulation of the expression of the TCF4 comprising contacting TCF4 with a molecule described herein.

[0658] A method of treatment of a disease caused by transcription of TCF4 comprising the administration of a therapeutically effective amount of a molecule described herein to a patient in need thereof.

[0659] In some embodiments, the disease is Fuchs Endothelial Corneal Dystrophy.

[0660] Some embodiments relate to a method of treatment of a disease caused by transcription of TCF4 comprising the administration of: a therapeutically effective amount of a molecule described herein; and another therapeutic agent.

[0661] Provided herein is a method for achieving an effect in a patient comprising the administration of a therapeutically effective amount of a molecule as disclosed herein, or a salt thereof, to a patient, wherein the effect is chosen from glare, blurred vision, pain or grittiness on cornea.

[0662] Certain molecules of the present disclosure may be effective for treatment of subjects whose genotype has 3 or more repeats of CAG or CTG. Certain molecules of the present disclosure may be effective for treatment of subjects whose genotype has 5 or more repeats of CAG or CTG. Certain molecules of the present disclosure may be effective for treatment of subjects whose genotype has 10 or more repeats of CAG or CTG. Certain molecules of the present disclosure may be effective for treatment of subjects whose genotype has 20 or more repeats of CAG or CTG. Certain molecules of the present disclosure may be effective for treatment of subjects whose genotype has 50 or more repeats of CAG or CTG. Certain molecules of the present disclosure may be effective for treatment of subjects whose genotype has 100 or more repeats of CAG or CTG. Certain molecules of the present disclosure may be effective for treatment of subjects whose genotype has 200 or more repeats of CAG or CTG. Certain compounds or molecules of the present dis-

closure may be effective for treatment of subjects whose genotype has 500 or more repeats of CAG or CTG.

[0663] Also provided is a method of modulation of a dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1-mediated function in a subject comprising the administration of a therapeutically effective amount of a compound as disclosed herein.

[0664] Also provided is a pharmaceutical composition comprising a compound as disclosed herein, together with a pharmaceutically acceptable carrier.

[0665] In certain embodiments, the pharmaceutical composition is formulated for oral administration.

[0666] In certain embodiments, the pharmaceutical composition is formulated for intravenous injection and/or infusion.

[0667] In certain embodiments, the oral pharmaceutical composition is chosen from a tablet and a capsule.

[0668] In certain embodiments, ex vivo methods of treatment are provided. Ex vivo methods typically include cells, organs, and/or tissues removed from the subject. The cells, organs and/or tissues can, for example, be incubated with the agent under appropriate conditions. The contacted cells, organs, and/or tissues are typically returned to the donor, placed in a recipient, or stored for future use. Thus, the compound is generally in a pharmaceutically acceptable carrier.

[0669] In certain embodiments, administration of the pharmaceutical composition modulates expression of the target gene within 6 hours of treatment. In certain embodiments, administration of the pharmaceutical composition modulates expression of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1 within 24 hours of treatment. In certain embodiments, administration of the pharmaceutical composition modulates expression of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1 within 72 hours of treatment.

[0670] In certain embodiments, administration of the pharmaceutical composition causes a 2-fold increase in expression of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1. In certain embodiments, administration of the pharmaceutical composition causes a 5-fold increase in expression of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1. In certain embodiments, administration of the pharmaceutical composition causes a 10-fold increase in expression of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1. In certain embodiments, administration of the pharmaceutical composition causes a 20-fold increase in expression of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1.

[0671] In certain embodiments, administration of the pharmaceutical composition causes a 20% decrease in expression of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1. In certain embodiments, administration of the pharmaceutical composition causes a 50% decrease in expression of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1. In certain embodiments, administration of the pharmaceutical composition causes a 80% decrease in expression of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1. In certain embodiments, administration of the pharmaceutical composition causes a 90% decrease in expression of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1. In certain embodiments, admin-

istration of the pharmaceutical composition causes a 95% decrease in expression of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1. In certain embodiments, administration of the pharmaceutical composition causes a 99% decrease in expression of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1.

[0672] In certain embodiments, administration of the pharmaceutical composition causes expression of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1 to fall within 25% of the level of expression observed for healthy individuals. In certain embodiments, administration of the pharmaceutical composition causes expression of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1 to fall within 50% of the level of expression observed for healthy individuals. In certain embodiments, administration of the pharmaceutical composition causes expression of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1 to fall within 75% of the level of expression observed for healthy individuals. In certain embodiments, administration of the pharmaceutical composition causes expression of dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1 to fall within 90% of the level of expression observed for healthy individuals.

Pharmaceutical Composition and Administration

[0673] Also provided is a method of modulation of a dmpk, atxn1, atxn2, atxn3, cacna1a, atxn7, ppp2r2b, tbp, htt, jph3, ar, or atn1-mediated function in a subject comprising the administration of a therapeutically effective amount of a compound or molecule as disclosed herein.

[0674] Also provided is a pharmaceutical composition comprising a compound as disclosed herein, together with a pharmaceutically acceptable carrier.

[0675] In certain embodiments, the pharmaceutical composition is formulated for oral administration.

[0676] In certain embodiments, the pharmaceutical composition is formulated for intravenous injection or infusion.

[0677] In certain embodiments, the oral pharmaceutical composition is chosen from a tablet and a capsule.

[0678] In certain embodiments, ex vivo methods of treatment are provided. Ex vivo methods typically include cells, organs, or tissues removed from the subject. The cells, organs or tissues can, for example, be incubated with the agent under appropriate conditions. The contacted cells, organs, or tissues are typically returned to the donor, placed in a recipient, or stored for future use. Thus, the compound is generally in a pharmaceutically acceptable carrier.

[0679] In certain embodiments, the compound or molecule is effective at a concentration less than about 5 μ M. In certain embodiments, the compound or molecule is effective at a concentration less than about 1 μ M. In certain embodiments, the compound or molecule is effective at a concentration less than about 400 nM. In certain embodiments, the compound or molecule is effective at a concentration less than about 200 nM. In certain embodiments, the compound or molecule is effective at a concentration less than about 100 nM. In certain embodiments, the compound or molecule is effective at a concentration less than about 50 nM. In certain embodiments, the compound or molecule is effective at a concentration less than about 20 nM. In certain embodiments, the compound or molecule is effective at a concentration less than about 10 nM.

Abbreviations and Definitions


[0680] As used herein, the terms below have the meanings indicated.

[0681] It is to be understood that certain radical naming conventions can include either a mono-radical or a di-radical, depending on the context. For example, where a substituent requires two points of attachment to the rest of the molecule, it is understood that the substituent is a di-radical. For example, a substituent identified as alkyl that requires two points of attachment includes di-radicals such as $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2-$, and the like. Other radical naming conventions clearly indicate that the radical is a di-radical such as “alkylene,” “alkenylene,” “arylene,” “heteroarylene.”

[0682] When two R groups are said to form a ring (e.g., a carbocyclyl, heterocyclyl, aryl, or heteroaryl ring) “together with the atom to which they are attached,” it is meant that the collective unit of the atom and the two R groups are the recited ring. The ring is not otherwise limited by the definition of each R group when taken individually. For example, when the following substructure is present:




[0683] and R^1 and R^2 are defined as selected from the group consisting of hydrogen and alkyl, or R^1 and R^2 together with the nitrogen to which they are attached form a heterocyclyl, it is meant that R^1 and R^2 can be selected from hydrogen or alkyl, or alternatively, the substructure has structure:




[0684] where ring A is a heteroaryl ring containing the depicted nitrogen.

[0685] Similarly, when two “adjacent” R groups are said to form a ring “together with the atom to which they are attached,” it is meant that the collective unit of the atoms, intervening bonds, and the two R groups are the recited ring. For example, when the following substructure is present:




[0686] and R^1 and R^2 are defined as selected from the group consisting of hydrogen and alkyl, or R^1 and R^2 together with the atoms to which they are attached form an aryl or carbocyclyl, it is meant that R^1 and R^2 can be selected from hydrogen or alkyl, or alternatively, the substructure has structure:



[0687] where A is an aryl ring or a carbocyclyl containing the depicted double bond.

[0688] Wherever a substituent is depicted as a di-radical (i.e., has two points of attachment to the rest of the molecule), it is to be understood that the substituent can be attached in any directional configuration unless otherwise indicated. Thus, for example, a substituent depicted as -AE- or



includes the substituent being oriented such that the A is attached at the leftmost attachment point of the molecule as well as the case in which A is attached at the rightmost attachment point of the molecule.

[0689] When ranges of values are disclosed, and the notation “from n_1 . . . to n_2 ” or “between n_1 . . . and n_2 ” is used, where n_1 and n_2 are the numbers, then unless otherwise specified, this notation is intended to include the numbers themselves and the range between them. This range may be integral or continuous between and including the end values. By way of example, the range “from 2 to 6 carbons” is intended to include two, three, four, five, and six carbons, since carbons come in integer units. Compare, by way of example, the range “from 1 to 3 μM (micromolar),” which is intended to include 1 μM , 3 μM , and everything in between to any number of significant figures (e.g., 1.255 μM , 2.1 μM , 2.9999 μM , etc.).

[0690] The term “about,” as used herein, is intended to qualify the numerical values which it modifies, denoting such a value as variable within a margin of error. When no particular margin of error, such as a standard deviation to a mean value given in a chart or table of data, is recited, the term “about” should be understood to mean that range which would encompass the recited value and the range which would be included by rounding up or down to that figure as well, taking into account significant figures.

[0691] The term “polyamide” refers to polymers of linkable units chemically bound by amide (i.e., CONH) linkages; optionally, polyamides include chemical probes conjugated therewith. Polyamides may be synthesized by stepwise condensation of carboxylic acids (COOH) with amines ($\text{RR}'\text{NH}$) using methods known in the art. Alternatively, polyamides may be formed using enzymatic reactions in vitro, or by employing fermentation with microorganisms.

[0692] The term “linkable unit” refers to methylimidazoles, methylpyrroles, and straight and branched chain aliphatic functionalities (e.g., methylene, ethylene, propylene, butylene, and the like) which optionally contain nitrogen Substituents, and chemical derivatives thereof. The aliphatic functionalities of linkable units can be provided, for

example, by condensation of B-alanine or dimethylamino-propyl amine during synthesis of the polyamide by methods well known in the art.

[0693] The term “linker” refers to a chain of at least 10 contiguous atoms. In certain embodiments, the linker contains no more than 20 non-hydrogen atoms. In certain embodiments, the linker contains no more than 40 non-hydrogen atoms. In certain embodiments, the linker contains no more than 60 non-hydrogen atoms. In certain embodiments, the linker contains atoms chosen from C, H, N, O, and S. In certain embodiments, every non-hydrogen atom is chemically bonded either to 2 neighboring atoms in the linker, or one neighboring atom in the linker and a terminus of the linker. In certain embodiments, the linker forms an amide bond with at least one of the two other groups to which it is attached. In certain embodiments, the linker forms an ester or ether bond with at least one of the two other groups to which it is attached. In certain embodiments, the linker forms a thioester or thioether bond with at least one of the two other groups to which it is attached. In certain embodiments, the linker forms a direct carbon-carbon bond with at least one of the two other groups to which it is attached. In certain embodiments, the linker forms an amine or amide bond with at least one of the two other groups to which it is attached. In certain embodiments, the linker comprises $-(CH_2OCH_2)-$ units. In certain embodiments, the linker comprises $-(CH(CH_3)OCH_2)-$ units. In certain embodiments, the linker comprises $-(CH_2NR_NCH_2)-$ units, for $R_N=C_{1-4}$ alkyl. In certain embodiments, the linker comprises an arylene, cycloalkylene, or heterocycloalkylene moiety.

[0694] The term “spacer” refers to a chain of at least 5 contiguous atoms. In certain embodiments, the spacer contains no more than 10 non-hydrogen atoms. In certain embodiments, the spacer contains atoms chosen from C, H, N, O, and S. In certain embodiments, the spacer forms amide bonds with the two other groups to which it is attached. In certain embodiments, the spacer comprises $-(CH_2OCH_2)-$ units. In certain embodiments, the spacer comprises $-(CH_2NR_NCH_2)-$ units, for $R_N=C_{1-4}$ alkyl. In certain embodiments, the spacer contains at least one positive charge at physiological pH.

[0695] The term “turn component” refers to a chain of about 4 to 10 contiguous atoms. In certain embodiments, the turn component contains atoms chosen from C, H, N, O, and S. In certain embodiments, the turn component forms amide bonds with the two other groups to which it is attached. In certain embodiments, the turn component contains at least one positive charge at physiological pH.

[0696] The terms “nucleic acid” and “nucleotide” refer to ribonucleotide and deoxyribonucleotide, and analogs thereof, well known in the art.

[0697] The term “oligonucleotide sequence” refers to a plurality of nucleic acids having a defined sequence and length (e.g., 2, 3, 4, 5, 6, or even more nucleotides). The term “oligonucleotide repeat sequence” refers to a contiguous expansion of oligonucleotide sequences.

[0698] The term “transcription,” well known in the art, refers to the synthesis of RNA (i.e., ribonucleic acid) by DNA-directed RNA polymerase. The term “modulate transcription” refers to a change in transcriptional level which can be measured by methods well known in the art, for example, assay of mRNA, the product of transcription. In

certain embodiments, modulation is an increase in transcription. In other embodiments, modulation is a decrease in transcription.

[0699] The term “acyl,” as used herein, alone or in combination, refers to a carbonyl attached to an alkenyl, alkyl, aryl, cycloalkyl, heteroaryl, heterocycle, or any other moiety where the atom attached to the carbonyl is carbon. An “acetyl” group refers to a $-C(O)CH_3$ group. An “alkylcarbonyl” or “alkanoyl” group refers to an alkyl group attached to the parent molecular moiety through a carbonyl group. Examples of such groups include methylcarbonyl and ethylcarbonyl. Examples of acyl groups include formyl, alkanoyl and aroyl.

[0700] The term “alkenyl,” as used herein, alone or in combination, refers to a straight-chain or branched-chain hydrocarbon radical having one or more double bonds and containing from 2 to 20 carbon atoms. In certain embodiments, said alkenyl will comprise from 2 to 6 carbon atoms. The term “alkenylene” refers to a carbon-carbon double bond system attached at two or more positions such as ethenylene $[(-CH=CH-), (-C::C-)]$. Examples of suitable alkenyl radicals include ethenyl, propenyl, 2-methylpropenyl, 1,4-butadienyl and the like. Unless otherwise specified, the term “alkenyl” may include “alkenylene” groups.

[0701] The term “alkoxy,” as used herein, alone or in combination, refers to an alkyl ether radical, wherein the term alkyl is as defined below. Examples of suitable alkyl ether radicals include methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, iso-butoxy, sec-butoxy, tert-butoxy, and the like.

[0702] The term “alkyl,” as used herein, alone or in combination, refers to a straight-chain or branched-chain alkyl radical containing from 1 to 20 carbon atoms. In certain embodiments, said alkyl will comprise from 1 to 10 carbon atoms. In further embodiments, said alkyl will comprise from 1 to 8 carbon atoms. Alkyl groups may be optionally substituted as defined herein. Examples of alkyl radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, iso-amyl, hexyl, octyl, nonyl and the like. The term “alkylene,” as used herein, alone or in combination, refers to a saturated aliphatic group derived from a straight or branched chain saturated hydrocarbon attached at two or more positions, such as methylene $(-CH_2-)$. Unless otherwise specified, the term “alkyl” may include “alkylene” groups.

[0703] The term “alkylamino,” as used herein, alone or in combination, refers to an alkyl group attached to the parent molecular moiety through an amino group. Suitable alkylamino groups may be mono- or dialkylated, forming groups such as, for example, N-methylamino, N-ethylamino, N,N-dimethylamino, N,N-ethylmethylamino and the like.

[0704] The term “alkylidene,” as used herein, alone or in combination, refers to an alkenyl group in which one carbon atom of the carbon-carbon double bond belongs to the moiety to which the alkenyl group is attached.

[0705] The term “alkylthio,” as used herein, alone or in combination, refers to an alkyl thioether (R-S-) radical wherein the term alkyl is as defined above and wherein the sulfur may be singly or doubly oxidized. Examples of suitable alkyl thioether radicals include methylthio, ethylthio, n-propylthio, isopropylthio, n-butylthio, iso-butylthio, sec-butylthio, tert-butylthio, methanesulfonyl, ethanesulfonyl, and the like.

[0706] The term “alkynyl,” as used herein, alone or in combination, refers to a straight-chain or branched chain hydrocarbon radical having one or more triple bonds and containing from 2 to 20 carbon atoms. In certain embodiments, said alkynyl comprises from 2 to 6 carbon atoms. In further embodiments, said alkynyl comprises from 2 to 4 carbon atoms. The term “alkynylene” refers to a carbon-carbon triple bond attached at two positions such as ethynylene ($-\text{C}::\text{C}-$, $-\text{C}\equiv\text{C}-$). Examples of alkynyl radicals include ethynyl, propynyl, hydroxypropynyl, butyn-1-yl, butyn-2-yl, pentyn-1-yl, 3-methylbutyn-1-yl, hexyn-2-yl, and the like. Unless otherwise specified, the term “alkynyl” may include “alkynylene” groups.

[0707] The terms “amido” and “carbamoyl,” as used herein, alone or in combination, refer to an amino group as described below attached to the parent molecular moiety through a carbonyl group, or vice versa. The term “C-amido” as used herein, alone or in combination, refers to a $-\text{C}(\text{O})\text{N}(\text{RR}')$ group with R and R' as defined herein or as defined by the specifically enumerated “R” groups designated. The term “N-amido” as used herein, alone or in combination, refers to a $\text{RC}(\text{O})\text{N}(\text{R})-$ group, with R and R' as defined herein or as defined by the specifically enumerated “R” groups designated. The term “acylamino” as used herein, alone or in combination, embraces an acyl group attached to the parent moiety through an amino group. An example of an “acylamino” group is acetylamino ($\text{CH}_3\text{C}(\text{O})\text{NH}-$).

[0708] The term “amide,” as used herein, alone in combination, refers to $-\text{C}(\text{O})\text{NRR}'$, wherein R and R' are independently chosen from hydrogen, alkyl, acyl, heteroalkyl, aryl, cycloalkyl, heteroaryl, and heterocycloalkyl, any of which may themselves be optionally substituted. Additionally, R and R' may combine to form heterocycloalkyl, either of which may be optionally substituted. Amides may be formed by direct condensation of carboxylic acids with amines, or by using acid chlorides. In addition, coupling reagents are known in the art, including carbodiimide-based compounds such as DCC and EDCI.

[0709] The term “amino,” as used herein, alone or in combination, refers to $-\text{NRR}'$, wherein R and R' are independently chosen from hydrogen, alkyl, acyl, heteroalkyl, aryl, cycloalkyl, heteroaryl, and heterocycloalkyl, any of which may themselves be optionally substituted. Additionally, R and R' may combine to form heterocycloalkyl, either of which may be optionally substituted.

[0710] The term “aryl,” as used herein, alone or in combination, means a carbocyclic aromatic system containing one, two or three rings wherein such polycyclic ring systems are fused together. The term “aryl” embraces aromatic groups such as phenyl, naphthyl, anthracenyl, and phenanthryl. The term “arylene” embraces aromatic groups such as phenylene, naphthylene, anthracenylene, and phenanthrylene.

[0711] The term “arylkalkenyl” or “aralkenyl,” as used herein, alone or in combination, refers to an aryl group attached to the parent molecular moiety through an alkenyl group.

[0712] The term “arylkalkoxy” or “aralkoxy,” as used herein, alone or in combination, refers to an aryl group attached to the parent molecular moiety through an alkoxy group.

[0713] The term “arylkalkyl” or “aralkyl,” as used herein, alone or in combination, refers to an aryl group attached to the parent molecular moiety through an alkyl group.

[0714] The term “arylkalkynyl” or “aralkynyl,” as used herein, alone or in combination, refers to an aryl group attached to the parent molecular moiety through an alkynyl group.

[0715] The term “arylkalkanoyl” or “aralkanoyl” or “aroyl,” as used herein, alone or in combination, refers to an acyl radical derived from an aryl-substituted alkane carboxylic acid such as benzoyl, naphthoyl, phenylacetyl, 3-phenylpropionyl (hydrocinnamoyl), 4-phenylbutyryl, (2-naphthyl) acetyl, 4-chlorohydrocinnamoyl, and the like.

[0716] The term aryloxy as used herein, alone or in combination, refers to an aryl group attached to the parent molecular moiety through an oxy.

[0717] The terms “benzo” and “benz,” as used herein, alone or in combination, refer to the divalent radical $\text{C}_6\text{H}_4=$ derived from benzene. Examples include benzothioephene and benzimidazole.

[0718] The term “carbamate,” as used herein, alone or in combination, refers to an ester of carbamic acid ($-\text{NHCOO}-$) which may be attached to the parent molecular moiety from either the nitrogen or acid end, and which may be optionally substituted as defined herein.

[0719] The term “O-carbamyl” as used herein, alone or in combination, refers to a $-\text{OC}(\text{O})\text{NRR}'$ group with R and R' as defined herein.

[0720] The term “N-carbamyl” as used herein, alone or in combination, refers to a $\text{ROC}(\text{O})\text{NR}'-$ group, with R and R' as defined herein.

[0721] The term “carbonyl,” as used herein, when alone includes formyl [$-\text{C}(\text{O})\text{H}$] and in combination is a $-\text{C}(\text{O})-$ group.

[0722] The term “carboxyl” or “carboxy,” as used herein, refers to $-\text{C}(\text{O})\text{OH}$ or the corresponding “carboxylate” anion, such as is in a carboxylic acid salt. An “O-carboxy” group refers to a $\text{RC}(\text{O})\text{O}-$ group, where R is as defined herein. A “C-carboxy” group refers to a $-\text{C}(\text{O})\text{OR}$ groups where R is as defined herein.

[0723] The term “cyano,” as used herein, alone or in combination, refers to $-\text{CN}$.

[0724] The term “cycloalkyl,” or, alternatively, “carbocycle,” as used herein, alone or in combination, refers to a saturated or partially saturated monocyclic, bicyclic or tricyclic alkyl group wherein each cyclic moiety contains from 3 to 12 carbon atom ring members and which may optionally be a benzo fused ring system which is optionally substituted as defined herein. In certain embodiments, said cycloalkyl will comprise from 5 to 7 carbon atoms. Examples of such cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, tetrahydronaphthyl, indanyl, octahydronaphthyl, 2,3-dihydro-1H-indenyl, adamantyl and the like. “Bicyclic” and “tricyclic” as used herein are intended to include both fused ring systems, such as decahydronaphthalene, octahydronaphthalene as well as the multicyclic (multicentered) saturated or partially unsaturated type. The latter type of isomer is exemplified in general by, bicyclo[1,1,1]pentane, camphor, adamantane, and bicyclo[3,2,1]octane.

[0725] The term “ester,” as used herein, alone or in combination, refers to a carboxy group bridging two moieties linked at carbon atoms.

[0726] The term “ether,” as used herein, alone or in combination, refers to an oxy group bridging two moieties linked at carbon atoms.

[0727] The term “halo,” or “halogen,” as used herein, alone or in combination, refers to fluorine, chlorine, bromine, or iodine.

[0728] The term “haloalkoxy,” as used herein, alone or in combination, refers to a haloalkyl group attached to the parent molecular moiety through an oxygen atom.

[0729] The term “haloalkyl,” as used herein, alone or in combination, refers to an alkyl radical having the meaning as defined above wherein one or more hydrogens are replaced with a halogen. Specifically embraced are mono-haloalkyl, dihaloalkyl and polyhaloalkyl radicals. A mono-haloalkyl radical, for one example, may have an iodo, bromo, chloro or fluoro atom within the radical. Dihalo and polyhaloalkyl radicals may have two or more of the same halo atoms or a combination of different halo radicals. Examples of haloalkyl radicals include fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl and dichloropropyl. “haloalkylene” refers to a haloalkyl group attached at two or more positions. Examples include fluoromethylene ($-\text{CFH}-$), difluoromethylene ($-\text{CF}_2-$), chloromethylene ($-\text{CHCl}-$) and the like.

[0730] The term “heteroalkyl,” as used herein, alone or in combination, refers to a stable straight or branched chain, or combinations thereof, fully saturated or containing from 1 to 3 degrees of unsaturation, consisting of the stated number of carbon atoms and from one to three heteroatoms chosen from N, O, and S, and wherein the N and S atoms may optionally be oxidized and the N heteroatom may optionally be quaternized. The heteroatom(s) may be placed at any interior position of the heteroalkyl group. Up to two heteroatoms may be consecutive, such as, for example, $-\text{CH}_2-\text{NH}-\text{OCH}_3$.

[0731] The term “heteroaryl,” as used herein, alone or in combination, refers to a 3 to 15 membered unsaturated heteromonocyclic ring, or a fused monocyclic, bicyclic, or tricyclic ring system in which at least one of the fused rings is aromatic, which contains at least one atom chosen from N, O, and S. In certain embodiments, said heteroaryl will comprise from 1 to 4 heteroatoms as ring members. In further embodiments, said heteroaryl will comprise from 1 to 2 heteroatoms as ring members. In certain embodiments, said heteroaryl will comprise from 5 to 7 atoms. The term also embraces fused polycyclic groups wherein heterocyclic rings are fused with aryl rings, wherein heteroaryl rings are fused with other heteroaryl rings, wherein heteroaryl rings are fused with heterocycloalkyl rings, or wherein heteroaryl rings are fused with cycloalkyl rings. Examples of heteroaryl groups include pyrrolyl, pyrrolinyl, imidazolyl, pyrazolyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazolyl, pyranyl, furyl, thienyl, oxazolyl, isoxazolyl, oxadiazolyl, thiazolyl, thiadiazolyl, isothiazolyl, indolyl, isoindolyl, indoliziny, benzimidazolyl, quinolyl, isoquinolyl, quinoxaliny, quinazolinyl, indazolyl, benzotriazolyl, benzodioxolyl, benzopyranyl, benzoxazolyl, benzoxadiazolyl, benzothiazolyl, benzothiadiazolyl, benzofuryl, benzothienyl, chromonyl, coumarinyl, benzopyranyl, tetrahydroquinoliny, tetrazolopyridazinyl, tetrahydroisoquinoliny, thienopyridinyl, furopyridinyl, pyrrolopyridinyl and the

like. Exemplary tricyclic heterocyclic groups include carbazolyl, benzidolyl, phenanthrolinyl, dibenzofuranyl, acridinyl, phenanthridinyl, xanthenyl and the like.

[0732] The terms “heterocycloalkyl” and, interchangeably, “heterocycle,” as used herein, alone or in combination, each refer to a saturated, partially unsaturated, or fully unsaturated (but nonaromatic) monocyclic, bicyclic, or tricyclic heterocyclic group containing at least one heteroatom as a ring member, wherein each said heteroatom may be independently chosen from nitrogen, oxygen, and sulfur. In certain embodiments, said heterocycloalkyl will comprise from 1 to 4 heteroatoms as ring members. In further embodiments, said heterocycloalkyl will comprise from 1 to 2 heteroatoms as ring members. In certain embodiments, said heterocycloalkyl will comprise from 3 to 8 ring members in each ring. In further embodiments, said heterocycloalkyl will comprise from 3 to 7 ring members in each ring. In yet further embodiments, said heterocycloalkyl will comprise from 5 to 6 ring members in each ring. “Heterocycloalkyl” and “heterocycle” are intended to include sulfones, sulfoxides, N-oxides of tertiary nitrogen ring members, and carbocyclic fused and benzo fused ring systems; additionally, both terms also include systems where a heterocycle ring is fused to an aryl group, as defined herein, or an additional heterocycle group. Examples of heterocycle groups include tetrahydroisoquinoline, aziridinyl, azetidiny, 1,3-benzodioxolyl, dihydroisoindolyl, dihydroisoquinoliny, dihydrocinolinyl, dihydrobenzodioxinyl, dihydro[1,3]oxazolo[4,5-b]pyridinyl, benzothiazolyl, dihydroindolyl, dihydropyridinyl, 1,3-dioxanyl, 1,4-dioxanyl, 1,3-dioxolanyl, isoindoliny, morpholinyl, piperazinyl, pyrrolidinyl, tetrahydropyridinyl, piperidinyl, thiomorpholinyl, and the like. The heterocycle groups may be optionally substituted unless specifically prohibited.

[0733] The term “hydrazinyl” as used herein, alone or in combination, refers to two amino groups joined by a single bond, i.e., $-\text{N}-\text{N}-$.

[0734] The term “hydroxy,” as used herein, alone or in combination, refers to $-\text{OH}$.

[0735] The term “hydroxyalkyl,” as used herein, alone or in combination, refers to a hydroxy group attached to the parent molecular moiety through an alkyl group.

[0736] The term “imino,” as used herein, alone or in combination, refers to $=\text{N}-$.

[0737] The term “iminohydroxy,” as used herein, alone or in combination, refers to $=\text{N}(\text{OH})$ and $=\text{N}-\text{O}-$.

[0738] The phrase “in the main chain” refers to the longest contiguous or adjacent chain of carbon atoms starting at the point of attachment of a group to the compounds or molecules of any one of the formulas disclosed herein.

[0739] The term “isocyanato” refers to a $-\text{NCO}$ group.

[0740] The term “isothiocyanato” refers to a $-\text{NCS}$ group.

[0741] The phrase “linear chain of atoms” refers to the longest straight chain of atoms independently selected from carbon, nitrogen, oxygen and sulfur.

[0742] The term “lower,” as used herein, alone or in a combination, where not otherwise specifically defined, means containing from 1 to and including 6 carbon atoms (i.e., C_1-C_6 alkyl).

[0743] The term “lower aryl,” as used herein, alone or in combination, means phenyl or naphthyl, either of which may be optionally substituted as provided.

[0744] The term “lower heteroaryl,” as used herein, alone or in combination, means either 1) monocyclic heteroaryl comprising five or six ring members, of which between one and four said members may be heteroatoms chosen from N, O, and S, or 2) bicyclic heteroaryl, wherein each of the fused rings comprises five or six ring members, comprising between them one to four heteroatoms chosen from N, O, and S.

[0745] The term “lower cycloalkyl,” as used herein, alone or in combination, means a monocyclic cycloalkyl having between three and six ring members (i.e., C₃-C₆ cycloalkyl). Lower cycloalkyls may be unsaturated. Examples of lower cycloalkyl include cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl.

[0746] The term “lower heterocycloalkyl,” as used herein, alone or in combination, means a monocyclic heterocycloalkyl having between three and six ring members, of which between one and four may be heteroatoms chosen from N, O, and S (i.e., C₃-C₆ heterocycloalkyl). Examples of lower heterocycloalkyls include pyrrolidinyl, imidazolidinyl, pyrazolidinyl, piperidinyl, piperazinyl, and morpholinyl. Lower heterocycloalkyls may be unsaturated.

[0747] The term “lower amino,” as used herein, alone or in combination, refers to —NRR, wherein R and R' are independently chosen from hydrogen and lower alkyl, either of which may be optionally substituted.

[0748] The term “mercaptyl” as used herein, alone or in combination, refers to an RS-group, where R is as defined herein.

[0749] The term “nitro,” as used herein, alone or in combination, refers to —NO₂.

[0750] The terms “oxy” or “oxa,” as used herein, alone or in combination, refer to —O—.

[0751] The term “oxo,” as used herein, alone or in combination, refers to =O.

[0752] The term “perhaloalkoxy” refers to an alkoxy group where all of the hydrogen atoms are replaced by halogen atoms.

[0753] The term “perhaloalkyl” as used herein, alone or in combination, refers to an alkyl group where all of the hydrogen atoms are replaced by halogen atoms.

[0754] The terms “sulfonate,” “sulfonic acid,” and “sulfonic,” as used herein, alone or in combination, refer the —SO₃H group and its anion as the sulfonic acid is used in salt formation.

[0755] The term “sulfanyl,” as used herein, alone or in combination, refers to —S—.

[0756] The term “sulfinyl,” as used herein, alone or in combination, refers to —S(O)—.

[0757] The term “sulfonyl,” as used herein, alone or in combination, refers to —S(O)₂—.

[0758] The term “N-sulfonamido” refers to a RS(=O)₂NR'— group with R and R' as defined herein.

[0759] The term “S-sulfonamido” refers to a —S(=O)₂NRR', group, with R and R' as defined herein.

[0760] The terms “thia” and “thio,” as used herein, alone or in combination, refer to a —S— group or an ether wherein the oxygen is replaced with sulfur. The oxidized derivatives of the thio group, namely sulfinyl and sulfonyl, are included in the definition of thia and thio.

[0761] The term “thiol,” as used herein, alone or in combination, refers to an —SH group.

[0762] The term “thiocarbonyl,” as used herein, when alone includes thioformyl —C(S)H and in combination is a —C(S)— group.

[0763] The term “N-thiocarbamyl” refers to an ROC(S)NR'— group, with R and R' as defined herein.

[0764] The term “O-thiocarbamyl” refers to a —OC(S)NRR', group with R and R' as defined herein.

[0765] The term “thiocyanato” refers to a —CNS group.

[0766] The term “trihalomethanesulfonamido” refers to a X₃CS(O)₂NR— group with X is a halogen and R as defined herein.

[0767] The term “trihalomethanesulfonyl” refers to a X₃CS(O)₂— group where X is a halogen.

[0768] The term “trihalomethoxy” refers to a X₃CO— group where X is a halogen.

[0769] The term “trisubstituted silyl,” as used herein, alone or in combination, refers to a silicone group substituted at its three free valences with groups as listed herein under the definition of substituted amino. Examples include trimethylsilyl, tert-butyldimethylsilyl, triphenylsilyl and the like.

[0770] Any definition herein may be used in combination with any other definition to describe a composite structural group. By convention, the trailing element of any such definition is that which attaches to the parent moiety. For example, the composite group alkylamido would represent an alkyl group attached to the parent molecule through an amido group, and the term alkoxyalkyl would represent an alkoxy group attached to the parent molecule through an alkyl group.

[0771] When a group is defined to be “null,” what is meant is that said group is absent.

[0772] The term “optionally substituted” means the antecedent group may be substituted or unsubstituted. When substituted, the substituents of an “optionally substituted” group may include, without limitation, one or more substituents independently selected from the following groups or a particular designated set of groups, alone or in combination: lower alkyl, lower alkenyl, lower alkynyl, lower alkanoyl, lower heteroalkyl, lower heterocycloalkyl, lower haloalkyl, lower haloalkenyl, lower haloalkynyl, lower perhaloalkyl, lower perhaloalkoxy, lower cycloalkyl, phenyl, aryl, aryloxy, lower alkoxy, lower haloalkoxy, oxo, lower acyloxy, carbonyl, carboxyl, lower alkylcarbonyl, lower carboxyester, lower carboxamido, cyano, hydrogen, halogen, hydroxy, amino, lower alkylamino, arylamino, amido, nitro, thiol, lower alkylthio, lower haloalkylthio, lower perhaloalkylthio, arylthio, sulfonate, sulfonic acid, trisubstituted silyl, N₃, SH, SCH₃, C(O)CH₃, CO₂CH₃, CO₂H, pyridinyl, thiophene, furanyl, lower carbamate, and lower urea. Where structurally feasible, two substituents may be joined together to form a fused five-, six-, or seven-membered carbocyclic or heterocyclic ring consisting of zero to three heteroatoms, for example forming methylenedioxy or ethylenedioxy. An optionally substituted group may be unsubstituted (e.g., —CH₂CH₃), fully substituted (e.g., —CF₂CF₃), monosubstituted (e.g., —CH₂CH₂F) or substituted at a level anywhere in-between fully substituted and monosubstituted (e.g., —CH₂CF₃). Where substituents are recited without qualification as to substitution, both substituted and unsubstituted forms are encompassed. Where a substituent is qualified as “substituted,” the substituted form is specifically intended. Additionally, different sets of optional substituents to a particular moiety may be defined

as needed; in these cases, the optional substitution will be as defined, often immediately following the phrase, “optionally substituted with.”

[0773] As used herein, a substituted group is derived from the unsubstituted parent group in which there has been an exchange of one or more hydrogen atoms for another atom or group. Unless otherwise indicated, when a group is deemed to be “substituted,” it is meant that the group is substituted with one or more substituents independently selected from C₁-C₆ alkyl, C₁-C₆ alkenyl, C₁-C₆ alkynyl, C₁-C₆ heteroalkyl, C₃-C₇ carbocyclyl (optionally substituted with halo, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkyl, and C₁-C₆ haloalkoxy), C₃-C₇-carbocyclyl-C₁-C₆-alkyl (optionally substituted with halo, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkyl, and C₁-C₆ haloalkoxy), 3-10 membered heterocyclyl (optionally substituted with halo, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkyl, and C₁-C₆ haloalkoxy), and C₁-C₆ haloalkoxy), 3-10 membered heterocyclyl-C₁-C₆-alkyl (optionally substituted with halo, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkyl, and C₁-C₆ haloalkoxy), aryl (optionally substituted with halo, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkyl, and C₁-C₆ haloalkoxy), aryl(C₁-C₆)alkyl (optionally substituted with halo, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkyl, and C₁-C₆ haloalkoxy), 5-10 membered heteroaryl (optionally substituted with halo, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkyl, and C₁-C₆ haloalkoxy), 5-10 membered heteroaryl(C₁-C₆)alkyl (optionally substituted with halo, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkyl, and C₁-C₆ haloalkoxy), halo, cyano, hydroxy, C₁-C₆ alkoxy, C₁-C₆ alkoxy(C₁-C₆)alkyl (i.e., ether), aryloxy, sulfhydryl (mercapto), halo(C₁-C₆)alkyl (e.g., —CF₃), halo(C₁-C₆)alkoxy (e.g., —OCF₃), C₁-C₆ alkylthio, arylthio, amino, amino(C₁-C₆)alkyl, nitro, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, S-sulfonamido, N-sulfonamido, C-carboxy, O-carboxy, acyl, cyanato, isocyanato, thiocyanato, isothiocyanato, sulfinyl, sulfonyl, and oxo (=O). Wherever a group is described as “optionally substituted” that group can be substituted with the above substituents.

[0774] The term R or the term R', appearing by itself and without a number designation, unless otherwise defined, refers to a moiety chosen from hydrogen, alkyl, cycloalkyl, heteroalkyl, aryl, heteroaryl and heterocycloalkyl, any of which may be optionally substituted. Such R and R' groups should be understood to be optionally substituted as defined herein. Whether an R group has a number designation or not, every R group, including R, R' and R" where n=(1, 2, 3, . . . n), every substituent, and every term should be understood to be independent of every other in terms of selection from a group. Should any variable, substituent, or term (e.g. aryl, heterocycle, R, etc.) occur more than one time in a formula or generic structure, its definition at each occurrence is independent of the definition at every other occurrence. Those of skill in the art will further recognize that certain groups may be attached to a parent molecule or may occupy a position in a chain of elements from either end as written. For example, an unsymmetrical group such as —C(O)N(R)— may be attached to the parent moiety at either the carbon or the nitrogen.

[0775] Asymmetric centers exist in the compounds or molecules disclosed herein. These centers are designated by the symbols “R” or “S,” depending on the configuration of substituents around the chiral carbon atom. It should be understood that the disclosure encompasses all stereochemical isomeric forms, including diastereomeric, enantiomeric,

and epimeric forms, as well as d-isomers and l-isomers, and mixtures thereof. Individual stereoisomers of compounds or molecules can be prepared synthetically from commercially available starting materials which contain chiral centers or by preparation of mixtures of enantiomeric products followed by separation such as conversion to a mixture of diastereomers followed by separation or recrystallization, chromatographic techniques, direct separation of enantiomers on chiral chromatographic columns, or any other appropriate method known in the art. Starting compounds or molecules of particular stereochemistry are either commercially available or can be made and resolved by techniques known in the art. Additionally, the compounds or molecules disclosed herein may exist as geometric isomers. The present disclosure includes all cis, trans, syn, anti, entgegen (E), and zusammen (Z) isomers as well as the appropriate mixtures thereof. Additionally, compounds or molecules may exist as tautomers; all tautomeric isomers are provided by this disclosure. Additionally, the compounds or molecules disclosed herein can exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like. In general, the solvated forms are considered equivalent to the unsolvated forms.

[0776] The term “bond” refers to a covalent linkage between two atoms, or two moieties when the atoms joined by the bond are considered to be part of larger substructure. A bond may be single, double, or triple unless otherwise specified. A dashed line between two atoms in a drawing of a molecule indicates that an additional bond may be present or absent at that position.

[0777] The term “disease” as used herein is intended to be generally synonymous, and is used interchangeably with, the terms “disorder,” “syndrome,” and “condition” (as in medical condition), in that all reflect an abnormal condition of the human or animal body or of one of its parts that impairs normal functioning, is typically manifested by distinguishing signs and symptoms, and causes the human or animal to have a reduced duration or quality of life.

[0778] The term “combination therapy” means the administration of two or more therapeutic agents to treat a therapeutic condition or disorder described in the present disclosure. Such administration encompasses co-administration of these therapeutic agents in a substantially simultaneous manner, such as in a single capsule having a fixed ratio of active ingredients or in multiple, separate capsules for each active ingredient. In addition, such administration also encompasses use of each type of therapeutic agent in a sequential manner. In either case, the treatment regimen will provide beneficial effects of the drug combination in treating the conditions or disorders described herein.

[0779] The phrase “therapeutically effective” is intended to qualify the amount of active ingredients used in the treatment of a disease or disorder or on the effecting of a clinical endpoint.

[0780] The term “therapeutically acceptable” refers to those compounds or molecules (or salts, prodrugs, tautomers, zwitterionic forms, etc.) which are suitable for use in contact with the tissues of patients without undue toxicity, irritation, and allergic response, are commensurate with a reasonable benefit/risk ratio, and are effective for their intended use.

[0781] As used herein, reference to “treatment” of a patient is intended to include prophylaxis. Treatment may also be preemptive in nature, i.e., it may include prevention

of disease. Prevention of a disease may involve complete protection from disease, for example as in the case of prevention of infection with a pathogen, or may involve prevention of disease progression. For example, prevention of a disease may not mean complete foreclosure of any effect related to the diseases at any level, but instead may mean prevention of the symptoms of a disease to a clinically significant or detectable level. Prevention of diseases may also mean prevention of progression of a disease to a later stage of the disease.

[0782] The term “patient” is generally synonymous with the term “subject” and includes all mammals including humans. Examples of patients include humans, livestock such as cows, goats, sheep, pigs, and rabbits, and companion animals such as dogs, cats, rabbits, and horses. Preferably, the patient is a human.

[0783] The term “prodrug” refers to a compound or molecule that is made more active in vivo. Certain compounds or molecules disclosed herein may also exist as prodrugs, as described in *Hydrolysis in Drug and Prodrug Metabolism: Chemistry, Biochemistry, and Enzymology* (Testa, Bernard and Mayer, Joachim M. Wiley-VHCA, Zurich, Switzerland 2003). Prodrugs of the compounds described herein are structurally modified forms of the compound that readily undergo chemical changes under physiological conditions to provide the compound. Additionally, prodrugs can be converted to the compound by chemical or biochemical methods in an ex vivo environment. For example, prodrugs can be slowly converted to a compound when placed in a transdermal patch reservoir with a suitable enzyme or chemical reagent. Prodrugs are often useful because, in some situations, they may be easier to administer than the compound, or parent drug. They may, for instance, be bioavailable by oral administration whereas the parent drug is not. The prodrug may also have improved solubility in pharmaceutical compositions over the parent drug. A wide variety of prodrug derivatives are known in the art, such as those that rely on hydrolytic cleavage or oxidative activation of the prodrug. An example, without limitation, of a prodrug would be a compound which is administered as an ester (the “prodrug”), but then is metabolically hydrolyzed to the carboxylic acid, the active entity. Additional examples include peptidyl derivatives of a compound.

[0784] The compounds or molecules disclosed herein can exist as therapeutically acceptable salts. The present disclosure includes compounds or molecules listed above in the form of salts, including acid addition salts. Suitable salts include those formed with both organic and inorganic acids. Such acid addition salts will normally be pharmaceutically acceptable. However, salts of non-pharmaceutically acceptable salts may be of utility in the preparation and purification of the compound or molecule in question. Basic addition salts may also be formed and be pharmaceutically acceptable. For a more complete discussion of the preparation and selection of salts, refer to *Pharmaceutical Salts: Properties, Selection, and Use* (Stahl, P. Heinrich. Wiley-VCHA, Zurich, Switzerland, 2002).

[0785] Basic addition salts can be prepared during the final isolation and purification of the compounds or molecules by reacting a carboxy group with a suitable base such as the hydroxide, carbonate, or bicarbonate of a metal cation or with ammonia or an organic primary, secondary, or tertiary amine. The cations of therapeutically acceptable salts include lithium, sodium, potassium, calcium, magnesium,

and aluminum, as well as nontoxic quaternary amine cations such as ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, diethylamine, ethylamine, tributylamine, pyridine, N,N-dimethylaniline, N-methylpiperidine, N-methylmorpholine, dicyclohexylamine, procaine, dibenzylamine, N,N-dibenzylphenethylamine, 1-phenamine, and N,N-dibenzylethylenediamine. Other representative organic amines useful for the formation of base addition salts include ethylenediamine, ethanolamine, diethanolamine, piperidine, and piperazine.

[0786] Other carrier materials and modes of administration known in the pharmaceutical art may also be used. Pharmaceutical compositions of the disclosure may be prepared by any of the well-known techniques of pharmacy, such as effective formulation and administration procedures. Preferred unit dosage formulations are those containing an effective dose, as herein below recited, or an appropriate fraction thereof, of the active ingredient.

[0787] It should be understood that in addition to the ingredients particularly mentioned above, the formulations described above may include other agents conventional in the art having regard to the type of formulation in question, for example those suitable for oral administration may include flavoring agents.

[0788] The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration.

[0789] The compounds or molecules can be administered in various modes, e.g. orally, topically, or by injection. The precise amount of compound administered to a patient will be the responsibility of the attendant physician. The specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diets, time of administration, route of administration, rate of excretion, drug combination, the precise disorder being treated, and the severity of the indication or condition being treated. In addition, the route of administration may vary depending on the condition and its severity. The above considerations concerning effective formulations and administration procedures are well known in the art and are described in standard textbooks.

Combinations and Combination Therapy

[0790] In certain instances, it may be appropriate to administer at least one of the compounds or molecules described herein (or a pharmaceutically acceptable salt thereof) in combination with another therapeutic agent. By way of example only, if one of the side effects experienced by a patient upon receiving one of the compounds herein is hypertension, then it may be appropriate to administer an anti-hypertensive agent in combination with the initial therapeutic agent. Or, by way of example only, the therapeutic effectiveness of one of the compounds described herein may be enhanced by administration of an adjuvant (i.e., by itself the adjuvant may only have minimal therapeutic benefit, but in combination with another therapeutic agent, the overall therapeutic benefit to the patient is enhanced). Or, by way of example only, the benefit of experienced by a patient may be increased by administering one of the compounds described herein with another therapeutic agent (which also includes a therapeutic regimen) that also has therapeutic benefit. By

way of example only, in a treatment for diabetes involving administration of one of the compounds described herein, increased therapeutic benefit may result by also providing the patient with another therapeutic agent for diabetes. In any case, regardless of the disease, disorder or condition being treated, the overall benefit experienced by the patient may simply be additive of the two therapeutic agents or the patient may experience a synergistic benefit.

[0791] In any case, the multiple therapeutic agents (at least one of which is a compound disclosed herein) may be administered in any order or even simultaneously. If simultaneously, the multiple therapeutic agents may be provided in a single, unified form, or in multiple forms (by way of example only, either as a single pill or as two separate pills). One of the therapeutic agents may be given in multiple doses, or both may be given as multiple doses. If not simultaneous, the timing between the multiple doses may be any duration of time ranging from a few minutes to four weeks.

[0792] Thus, in another aspect, certain embodiments provide methods for treating disorders described herein in a human or animal subject in need of such treatment comprising administering to said subject an amount of a compound disclosed herein effective to reduce or prevent said disorder in the subject, in combination with at least one additional agent for the treatment of said disorder that is known in the art. In a related aspect, certain embodiments provide therapeutic compositions comprising at least one compound disclosed herein in combination with one or more additional agents for the treatment of disorders. The disorders can be associated with the expression of defective genes such as *dmpk*, *atxn1*, *atxn2*, *atxn3*, *cacna1a*, *atxn7*, *ppp2r2b*, *tbp*, *htt*, *jph3*, *ar*, *atn*, and gene encoding TCF4.

[0793] Besides being useful for human treatment, certain compounds and formulations disclosed herein may also be useful for veterinary treatment of companion animals, exotic animals and farm animals, including mammals, rodents, and the like. More preferred animals include horses, dogs, and cats.

Compound Synthesis

[0794] Compounds of the present disclosure can be prepared using methods illustrated in general synthetic schemes and experimental procedures detailed below. General synthetic schemes and experimental procedures are presented for purposes of illustration and are not intended to be limiting. Starting materials used to prepare compounds of the present disclosure are commercially available or can be prepared using routine methods known in the art.

List of Abbreviations


[0795] Ac_2O =acetic anhydride; AcCl =acetyl chloride; AcOH =acetic acid; AIBN =azobisisobutyronitrile; aq. =aqueous; Bu_3SnH =tributyltin hydride; CD_3OD =deuterated methanol; CDCl_3 =deuterated chloroform; CDI =1,1'-Carbonyldiimidazole; DBU =1,8-diazabicyclo[5.4.0]undec-7-ene; DCM =dichloromethane; DEAD =diethyl azodicarboxylate; DIBAL-H =di-iso-butyl aluminium hydride; DIEA =DIPEA= N,N -diisopropylethylamine; DMAP =4-dimethylaminopyridine; DMF = N,N -dimethylformamide; DMSO-d_6 =deuterated dimethyl sulfoxide; DMSO =dimethyl sulfoxide; DPPA =diphenylphosphoryl azide; $\text{EDC}\cdot\text{HCl}$ = $\text{EDCI}\cdot\text{HCl}$ =1-ethyl-3-(3-dimethylaminopropyl)


carbodiimide hydrochloride; Et_2O =diethyl ether; EtOAc =ethyl acetate; EtOH =ethanol; h =hour; HATU =2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium; HMDS =hexamethyldisilazane; HOBT =1-hydroxybenzotriazole; i-PrOH =isopropanol; LAH =lithium aluminium hydride; LiHMDS =Lithium bis(trimethylsilyl)amide; MeCN =acetonitrile; MeOH =methanol; MP =carbonate resin=macroporous triethylammonium methylpoly styrene carbonate resin; MsCl =mesyl chloride; MTBE =methyl tertiary butyl ether; MW =microwave irradiation; n-BuLi =n-butyllithium; NaHMDS =Sodium bis(trimethylsilyl)amide; NaOMe =sodium methoxide; NaOtBu =sodium t-butoxide; NBS =N-bromosuccinimide; NCS =N-chlorosuccinimide; NMP =N-Methyl-2-pyrrolidone; $\text{Pd}(\text{Ph}_3)_4$ =tetrakis(triphenylphosphine)palladium(O); $\text{Pd}_2(\text{dba})_3$ =tris(dibenzylidenecyclopentadienyl)dipalladium(O); $\text{PdCl}_2(\text{PPh}_3)_2$ =bis(triphenylphosphine)palladium(II) dichloride; PG =protecting group; prep-HPLC =preparative high-performance liquid chromatography; PyBop =(benzotriazol-1-yloxy)-tripyrrolidinophosphonium hexafluorophosphate; Pyr =pyridine; RT =room temperature; RuPhos =2-dicyclohexylphosphino-2',6'-diisopropoxybiphenyl; sat. =saturated; ss =saturated solution; t-BuOH =tert-butanol; T3P =Propylphosphonic Anhydride; TBS = TBDMS =tert-butyldimethylsilyl; TBSCl = TBDMSCl =tert-butyldimethylchlorosilane; TEA = Et_3N =triethylamine; TFA =trifluoroacetic acid; TFAA =trifluoroacetic anhydride; THF =tetrahydrofuran; Tol =toluene; TsCl =tosyl chloride; XPhos =2-dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl.

General Synthetic Methods for Preparing Compounds

[0796] In general, polyamides of the present disclosure may be synthesized by solid supported synthetic methods, using compounds such as Boc-protected straight chain aliphatic and heteroaromatic amino acids, and alkylated derivatives thereof, which are cleaved from the support by aminolysis, deprotected (e.g., with sodium thiophenoxide), and purified by reverse-phase HPLC, as well known in the art. The identity and purity of the polyamides may be verified using any of a variety of analytical techniques available to one skilled in the art such as $^1\text{H-NMR}$, analytical HPLC, or mass spectrometry.

[0797] The following scheme can be used to practice the present disclosure.





[0798] The compounds disclosed herein can be synthesized using Scheme I. For clarity and compactness, the scheme depicts the synthesis of a diamide comprising subunits “C” and “D”, both of which are represented as unspecified five-membered rings having amino and carboxy moieties. The amino group of subunit “D” is protected with a protecting group “PG” such as a Boc or CBz carbamate to give 101. The free carboxylic acid is then reacted with a solid support, using a coupling reagent such as EDC, to give the supported compound 103. Removal of PG under acidic conditions gives the free amine 104, which is coupled with the nitrogen-protected carboxylic acid 105 to give amide 106. Removal of PG under acidic conditions gives the free amine 107. In this example, the free amine is reacted with acetic anhydride to form an acetamide (not shown). The molecule is then cleaved from the solid support under basic conditions to give carboxylic acid 108. Methods for attachment of the linker L and recruiting moiety X are disclosed below.

[0799] The person of skill will appreciate that many variations of the above scheme are available to provide a wide range of compounds:

[0800] 1) The sequence 104-106-107 can be repeated as often as desired, in order to form longer polyamine sequences.

[0801] 2) A variety of amino heterocycle carboxylic acids can be used, to form different subunits. Table 3, while not intended to be limiting, provides several heterocycle amino acids that are contemplated for the synthesis of the compounds in this disclosure. Carbamate protecting groups PG can be incorporated using techniques that are well established in the art.

TABLE 3

Heterocyclic amino acids.
Structure




TABLE 3-continued

Heterocyclic amino acids.
Structure











TABLE 3-continued
















Heterocyclic amino acids.
Structure(Z is H, C₁₋₆ alkyl, amine, or halogen)(Z is H, C₁₋₆ alkyl, amine, or halogen)

TABLE 3-continued

Heterocyclic amino acids.
Structure


[0802] 3) Hydroxy-containing heterocyclic amino acids can be incorporated into Scheme I as their TBS ethers. While not intended to be limiting, Scheme II provides the synthesis of TBS-protected heterocyclic amino acids contemplated for the synthesis of the molecules in this disclosure.

Scheme II: Synthesis of TBS-protected heterocyclic amino acids



X = N(CH₃); S

-continued





[0803] 4) Aliphatic amino acids can be used in the above synthesis for the formation of spacer units “W” and subunits for recognition of DNA nucleotides. Table 4, while not intended to be limiting, provides several aliphatic amino acids contemplated for the synthesis of the or molecules in this disclosure.

TABLE 4


Aliphatic amino acids.	
Structure	
	beta-alanine (B)
	gamma-aminobutyric acid (“gAB” or γ)
	3-(2-aminoethoxy)propanoic acid
	3-((2-aminoethyl)(2-oxo-2-phenyl-1λ ² -ethyl)amino)propanoic acid
	Dp
	(R is H, C ₁₋₆ alkyl)
	(R is H, C ₁₋₆ alkyl, aryl, or heteroaryl)

TABLE 4-continued


Aliphatic amino acids.	
Structure	
	X is F or OH



[0804] Attachment of the linker L and recruiting moiety X can be accomplished with the methods disclosed in Scheme III, which uses a triethylene glycol moiety for the linker L. The mono-TBS ether of triethylene glycol 301 is converted to the bromo compound 302 under Mitsunobu conditions. The recruiting moiety X is attached by displacement of the bromine with a hydroxyl moiety, affording ether 303. The TBS group is then removed by treatment with fluoride, to provide alcohol 304, which will be suitable for coupling with the polyamide moiety. Other methods will be apparent to the person of skill in the art for inclusion of alternate linkers L, including but not limited to propylene glycol or polyamine linkers, or alternate points of attachment of the recruiting moiety X, including but not limited to the use of amines and thiols.




378




[0805] Synthesis of the X-L-Y molecule can be completed with the methods set forth in Scheme IV. Carboxylic acid 108 is converted to the acid chloride 401. Reaction with the alcohol functionality of 301 under basic conditions provides the coupled product 402. Other methods will be apparent to the person of skill in the art for performing the coupling procedure, including but not limited to the use of carbodiimide reagents. For instance, the amide coupling reagents can be used, but not limited to, are carbodiimides such as dicyclohexylcarbodiimide (DCC), diisopropylcarbodiimide (DIC), ethyl-(N',N'-dimethylamino)propylcarbodiimide hydrochloride (EDC), in combination with reagents such as 1-hydroxybenzotriazole (HOBt), 4-(N,N-dimethylamino)pyridine (DMAP) and diisopropylethylamine (DIEA). Other reagents are also often used depending the actual coupling reactions are (Benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (BOP), (Benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyBOP), (7-Azabenzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyAOP), Bromotripyrrolidinophosphonium hexafluorophosphate (PyBrOP), Bis(2-oxo-3-oxazolidinyl)phosphinic chloride (BOP-Cl), O-(Benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU), O-(Benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU), O-(7-Azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU), O-(7-Azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TATU), O-(6-Chlorobenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HCTU), Carbonyldiimidazole (CDI), and N,N,N',N'-Tetramethylchloroformamidinium Hexafluorophosphate (TCFH).

-continued




Scheme V: Proposed synthesis of rohitukine-based CDK9 inhibitor



[0806] A proposed synthesis of a rohitukine-based CDK9 inhibitor is set forth in Scheme V. Synthesis begins with the natural product rohitukine, which is a naturally available compound that has been used as a precursor for CDK9-active drugs such as Alvocidib. The existing hydroxy groups are protected as TBS ethers, the methyl group is brominated, and the bromo compound is coupled with a suitably functionalized linker reagent such as 501 to afford the linked compound 502. Variants of this procedure will be apparent to the person of skill.


Scheme VI: Proposed synthesis of DB08045-based cyclin T1 inhibitor




[0807] Proposed syntheses of DB08045-based cyclin T1 inhibitors are set forth in Scheme VI. Synthesis begins with DB08045, which contains a primary amino group that is available for functionalization. Coupling of the amino group

with a carboxylic acid under conventional conditions gives amide 601. Alternatively, reductive amination with a carboxaldehyde gives amine 602. Variants of this procedure will be apparent to the person of skill.

Scheme VII: Proposed synthesis of A-395 based PRC2 inhibitor



-continued




[0808] A proposed synthesis of an A-395 based PRC2 inhibitor is set forth in Scheme VII. The piperidine compound 701, a precursor to A-395, can be reacted with methanesulfonyl chloride 702 to give A-395. In a variation of this synthesis, 701 is reacted with linked sulfonyl chloride 703, to provide linked A-395 inhibitor 704


Attaching Protein Binding Molecules to Oligomeric Backbone

[0809] Generally the oligomeric backbone is functionalized to adapt to the type of chemical reactions can be performed to link the oligomers to the attaching position in protein binding moieties. The type reactions are suitable but not limited to, are amide coupling reactions, ether formation reactions (O-alkylation reactions), amine formation reactions (N-alkylation reactions), and sometimes carbon-carbon coupling reactions. The general reactions used to link oligomers and protein binders are shown in below schemes (VIII through X). The compounds and structures shown in Table 2 can be attached to the oligomeric backbone described herein at any position that is chemically feasible while not interfering with the hydrogen bond between the compound and the regulatory protein.


Scheme VIII. Amide Couplings



or





-continued



[0810] Either the oligomer or the protein binder can be functionalized to have a carboxylic acid and the other coupling counterpart being functionalized with an amino group so the moieties can be conjugated together mediated by amide coupling reagents. The amide coupling reagents can be used, but not limited to, are carbodiimides such as dicyclohexylcarbodiimide (DCC), diisopropylcarbodiimide (DIC), ethyl-(N',N'-dimethylamino)propylcarbodiimide hydrochloride (EDC), in combination with reagents such as 1-hydroxybenzotriazole (HOBt), 4-(N,N-dimethylamino) pyridine (DMAP) and diisopropylethylamine (DIEA). Other reagents are also often used depending the actual coupling reactions are (Benzotriazol-1-yloxy)tris(dimethylamino) phosphonium hexafluorophosphate (BOP), (Benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (Py-BOP), (7-Azabenzotriazol-1-yloxy) tripyrrolidinophosphonium hexafluorophosphate (PyAOP), Bromotripyrrolidinophosphonium hexafluorophosphate (PyBrOP), Bis(2-oxo-3-oxazolidinyl)phosphinic chloride (BOP-Cl), O-(Benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU), O-(Benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU), O-(7-Azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU), O-(7-Azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TATU), O-(6-Chlorobenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HCTU), Carbonyldiimidazole (CDI), and N,N,N',N'-Tetramethylchloroformamidinium Hexafluorophosphate (TCFH).


Scheme IX. Ether Formation Reactions (O-alkylation reactions)





[0811] In an ether formation reaction, either the oligomer or the protein binder can be functionalized to have an hydroxyl group (phenol or alcohol) and the other coupling counterpart being functionalized with a leaving group such as halide, tosylate and mesylate so the moieties can be conjugated together mediated by a base or catalyst. The bases can be selected from, but not limited to, sodium hydride, potassium hydride, sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate. The catalyst can be selected from silver oxide, phase transfer reagents, iodide salts, and crown ethers.

Scheme X. Amine Formation Reactions (N-alkylation reactions)



[0812] In an N-alkylation reaction, either the oligomer or the protein binder can be functionalized to have an amino group (arylamine or alkylamine) and the other coupling counterpart being functionalized with a leaving group such as halide, tosylate and mesylate so the moieties can be conjugated together directly or with a base or catalyst. The bases can be selected from, but not limited to, sodium hydride, potassium hydride, sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate. The catalyst can be selected from silver oxide, phase transfer reagents, iodide salts, and crown ethers. The alkylation of amines can also be achieved through reductive amination

reactions, where in either the oligomer or the protein binder can be functionalized to have an amino group (arylamine or alkylamine) and the other coupling counterpart being functionalized with an aldehyde or ketone group so the moieties can be conjugated together with the treatment of a reducing reagent (hydride source) directly or in combination with a dehydration agent. The reducing reagents can be selected from, but not limited to, NaBH_4 , $\text{NaHB}(\text{OAc})_3$, NaBH_3CN , and dehydration agents are normally $\text{Ti}(\text{iPrO})_4$, $\text{Ti}(\text{OEt})_4$, $\text{Al}(\text{iPrO})_3$, orthoformates and activated molecular sieves.

Cell-Penetrating Ligand

[0813] In one aspect, the molecules of the present disclosure comprises a cell-penetrating ligand moiety. The cell-penetrating ligand moiety serves to facilitate transport of the compound across cell membranes. In certain embodiments, the cell-penetrating ligand moiety is a polypeptide. Several peptide sequences can facilitate passage into the cell, including polycationic sequences such as poly-R; arginine-rich sequences interspersed with spacers such as $(\text{RXR})_n$ (X =6-aminohexanoic acid) and $(\text{RXRRBR})_n$ (B =beta-alanine); sequences derived from the Penetratin peptide; and sequences derived from the PNA/PMO internalisation peptide (Pip). The Pip5 series is characterized by the sequence ILFQY.

[0814] In certain embodiments, the cell-penetrating polypeptide comprises an N-terminal cationic sequence $\text{H}_2\text{N}-(\text{R})_n-\text{COOH}$, with $n=5-10$, inclusive. In certain embodiments, the N-terminal cationic sequence contains 1, 2, or 3 substitutions of R for amino acid residues independently chosen from beta-alanine and 6-aminohexanoic acid.

[0815] In certain embodiments, the cell-penetrating polypeptide comprises the ILFQY sequence. In certain embodiments, the cell-penetrating polypeptide comprises the QFLY sequence. In certain embodiments, the cell-penetrating polypeptide comprises the QFL sequence.

[0816] In certain embodiments, the cell-penetrating polypeptide comprises a C-terminal cationic sequence $-\text{HN}-(\text{R})_n-\text{COOH}$, with $n=5-10$, inclusive. In certain embodiments, the C-terminal cationic sequence contains 1, 2, or 3 substitutions of R for amino acid residues independently chosen from beta-alanine and 6-aminohexanoic acid. In certain embodiments, the C-terminal cationic sequence is substituted at every other position with an amino acid residue independently chosen from beta-alanine and 6-aminohexanoic acid. In certain embodiments, the C-terminal cationic sequence is $-\text{HN}-\text{RXRBRXRB}-\text{COOH}$.


TABLE 5

Cell-penetrating peptides	
SEQ ID NO.	Sequence
SEQ ID NO. 1	GRKKRRQRRRPPQ
SEQ ID NO. 2	RQKIWFQNRMRKWK
SEQ ID NO. 3	KLALKLALKALKALKA
SEQ ID NO. 4	GWTLNS/AGYLLGKINLKALAALAKKIL
SEQ ID NO. 5	NAKTRRHERRRKLAIER
SEQ ID NO. 6	RRRRRRRR
SEQ ID NO. 7	RRRRRRRR
SEQ ID NO. 8	GALFLGFLGAAGSTMGA
SEQ ID NO. 9	KETWWETWWTEWSQPKKKRKV
SEQ ID NO. 10	LLILRRRIKQAHHSK
SEQ ID NO. 11	YTAIAWVKAFIRKLK
SEQ ID NO. 12	IAWVKAFIRKLKGPLG

TABLE 5-continued

Cell-penetrating peptides	
SEQ ID NO.	Sequence
SEQ ID NO. 13	MVTVLFRRIRRRACGPPRVRV
SEQ ID NO. 14	GLWRALWRLRLSLWRLWRA
SEQ ID NO. 15	RRRRRRR QIKIWFQNNRMKWKKGG
SEQ ID NO. 16	RXRRXRRXRIKILFQNNRMKWKK
SEQ ID NO. 17	RXRRXRRXRIdKILFQNdRRMKWHKB
SEQ ID NO. 18	RXRRXRRXRIHILFQNdRRMKWHKB
SEQ ID NO. 19	RXRRBRXRILFQYRXRBXRBB
SEQ ID NO. 20	RXRRBRXRILFQYRXRXRXRB
SEQ ID NO. 21	RXRRXRILFQYRXRXRXR
SEQ ID NO. 22	RBRRXRBRILFQYRBXRBRB
SEQ ID NO. 23	RBRRXRBRILFQYRXRBXRBB
SEQ ID NO. 24	RBRRXRBRILFQYRXRXRXRB
SEQ ID NO. 25	RBRRXRBRILFQYRXRBXRBB
SEQ ID NO. 26	RXRRBRXRILFQYRXRXRXRB
SEQ ID NO. 27	RXRRBRXRILFQYRXRBXRBB
SEQ ID NO. 28	RXRRBRXRQFLIRXRBRXRBB
SEQ ID NO. 29	RXRRBRXRQFLIRXRBRXRBB
SEQ ID NO. 30	RXRRBRXRQFLIRXRBRXRBB
SEQ ID NO. 31	RXRRBRXRQFLIRXRBRXRBB
SEQ ID NO. 32	RXRRBRXRQFLIRXRBRXRBB
SEQ ID NO. 33	RXRRBRXRQFLIRXRBRXRBB
SEQ ID NO. 34	RXRRBRXRQFLIRXRBRXRBB
SEQ ID NO. 35	RXRRBRXRILFRYRXRBXRBB
SEQ ID NO. 36	Ac-RRLSYRRRFXBpgG
SEQ ID NO. 37	Ac-RRLSYRRRFPFVYLIXBpgG

Ac = acetyl;
Bpg = L-bis-homopropargylglycine =



B = beta-alanine;
X = 6-aminoheptanoic acid;
dK/dR = corresponding D-amino acid.

EXAMPLES

[0817] The following examples are given for the purpose of illustrating various embodiments of the invention and are not meant to limit the present invention in any fashion. The present examples, along with the methods described herein are presently representative of preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention. Changes therein and other uses which are encompassed within the spirit of the invention as defined by the scope of the claims will occur to those skilled in the art.

[0818] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments described herein may be employed. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

Example 1

[0819] Scheme A describes the steps involved for preparing the polyamide, attaching the polyamide to the oligomeric backbone, and then attaching the ligand to the other end of the oligomeric backbone. The second terminus can include any structure in Table 2. The oligomeric backbone can be selected from the various combinations of linkers shown in Table 6. The transcription modulator molecule such as those listed in Table 7 below can be prepared using the synthesis scheme shown below.

TABLE 6

Examples of oligomeric backbone as represented by $-(T^1-V^1)_a-(T^2-V^2)_b-(T^3-V^3)_c-(T^4-V^4)_d-(T^5-V^5)_e-$									
T ¹	V ¹	T ²	V ²	T ³	V ³	T ⁴	V ⁴	T ⁵	V ⁵
(C ₁ -C ₁₂) alkylene	CONR ^{1a}	(EA) _w	CO	(PEG) _n	NR ¹¹ CO	—	—	—	—
(C ₁ -C ₁₂) alkylene	CONR ^{1a}	(EA) _w	CO	(PEG) _n	O	arylene	NR ¹¹ CO	—	—
(C ₁ -C ₁₂) alkylene	CONR ^{1a}	(EA) _w	CO	(PEG) _n	O	Subst. arylene	NR ¹¹ CO	—	—
(C ₁ -C ₁₂) alkylene	CONR ^{1a}	(EA) _w	CO	(PEG) _n	O	NR ¹¹ CO	(C ₁ -C ₁₂) alkyl	Subst. arylene	NR ¹¹ CO
(C ₁ -C ₁₂) alkylene	CONR ^{1a}	(EA) _w	CO	(C ₁ -C ₁₂) alkyl	NR ¹¹ CO- C ₁₋₄ alkyl	Subst. arylene	NR ¹¹	—	—
(C ₁ -C ₁₂) alkylene	CONR ^{1a}	(EA) _w	CO	(PEG) _n	O	Subst. arylene	—	—	—
(PEG) _n	CONR ^{1a} - C ₁₋₄ alkyl	—	—	—	—	—	—	—	—
(EA) _w	CO	(C ₁ -C ₁₂) alkyl	CONR ¹¹ - C ₁₋₄ alkyl	—	—	—	—	—	—
(C ₁ -C ₁₂) alkylene	CONR ^{1a}	(EA) _w	CO	(PEG) _n	NR ¹¹ CO- C ₁₋₄ alkyl	—	—	—	—
(EA) _w	CO	(PEG) _n	O	phenyl	NR ¹¹ CO- C ₁₋₄ alkyl	—	—	—	—
(C ₁ -C ₁₂) alkylene	CONR ^{1a}	(PEG) _n	CO	—	—	—	—	—	—
(C ₁ -C ₁₂) alkylene	CONR ^{1a}	(EA) _w	CO	modifd. (PEG) _n	O	arylene	NR ¹¹ CO	—	—

Examples of transcription modulator molecules

First terminus	Oligomeric backbone	Second terminus
Py- β -Im-Py		
Py- β -Im-Py		
Py- β -Im-Py		
Py- β -Im-Py		
Py- β -Im-Py		

TABLE 7-continued











Examples of transcription modulator molecules		
First terminus	Oligomeric backbone	Second terminus
Py- β -Im-Py		 (OGT binder)
Py- β -Im-Py		 (methyl lysine binder)
Py- β -Im-Py		 (methyl lysine binder)
Py- β -Im-Py		 (methyl lysine binder)
Py- β -Im- β -W-Im-Py-Im		 (CDK 1, 2, 5)

TABLE 7-continued

Examples of transcription modulator molecules		
First terminus	Oligomeric backbone	Second terminus
Py- β - Im- β - W-Im- Py- Py-Im		 (CDK 1, 2, 5)
Py- β - Im- β - W-Im- Py- Py-Im		 (CDK 1, 2, 5)
Py- β - Im- β - W-Im- Py- Py-Im		 (methyl transferase)
Py- β - Im- β - W-Im- Py- Py-Im		 (methyl transferase)

TABLE 7-continued











Examples of transcription modulator molecules		
First terminus	Oligomeric backbone	Second terminus
Py- β - Im- β - W-Im- Py- Py-Im		 (methyl transferase)
Py- β - Im- β - W-Im- Py- Py-Im		 (CBP binder)
Py- β - Im- β - W-Im- Py- Py-Im		 (CBP binder)
Py- β - Im- β - W-Im- Py- Py-Im		 (CBP binder)
Py- β - Im- β - W-Im- Py- Py-Im		 (OGT binder)

TABLE 7-continued











Examples of transcription modulator molecules		
First terminus	Oligomeric backbone	Second terminus
Py- β - Im- β - W-Im- Py- Py-Im		 (OGT binder)
Py- β - Im- β - W-Im- Py- Py-Im		 (OGT binder)
Py- β - Im- β - W-Im- Py- Py-Im		 (methyl lysine binder)
Py- β - Im- β - W-Im- Py- Py-Im		 (methyl lysine binder)
Py- β - Im- β - W-Im- Py- Py-Im		 (methyl lysine binder)

TABLE 7-continued










Examples of transcription modulator molecules		
First terminus	Oligomeric backbone	Second terminus
Py-β- Im-β- W-Im- Py- Py-Im		 (CDK 1, 2, 5)
Py-β- Im-β- W-Im- Py- Py-Im		 (CDK 1, 2, 5)
Py-β- Im-β- W-Im- Py- Py-Im		 (CDK 1, 2, 5)
Py-Py- Im-Py (Linked in the middle - either position 2 or 3) to Im-Py- Py-Im		

TABLE 7-continued

Examples of transcription modulator molecules		
First terminus	Oligomeric backbone	Second terminus
Py-Py-Im-Py (Linked in the middle - either position 2 or 3) to Im-Py-Py-Im		
Py-Py-Im-Py (Linked in the middle - either position 2 or 3) to Im-Py-Py-Im		

Scheme A: Synthesis of first terminus/second terminus/linker conjugate.




② indicates text missing or illegible when filed


[0820] The ligand or protein binder can be attached to the oligomeric backbone using the schemes described below. The oligomeric backbone can be linked to the protein binder at any position on the protein binder that is chemically feasible while not interfering with the binding between the protein binder and the regulatory protein. The protein binder binds to the regulatory protein often through hydrogen bonds, and linking the oligomeric backbone and the regulatory protein should not interfere the hydrogen bond formation. The protein binder is attached to the oligomeric backbone through an amide or ether bond.

[0821] Scheme B through Scheme D demonstrate several examples of linking the oligomeric backbone and protein binder.

Scheme B. Example for Amide Coupling


Amide coupling Reaction:






Scheme C. Example for Ether Formation Reaction (O-alkylation reaction)

Ether Formation Reaction (O-alkylation):



Scheme D. Example for Amine Formation Reaction (N-alkylation reaction)

Amine Formation Reaction (N-alkylation):



Example 2. Biological Activity Assays

[0822] The methods as set forth below will be used to demonstrate the binding of the disclosed molecules and the efficacy in treatment. In general, the assays are directed at evaluating the effect of the disclosed molecules on the level of expression of the target gene containing CAG or CTG repeats (e.g., *dmpk*, *atxn1*, *atxn2*, *atxn3*, *cacna1a*, *atxn7*, *ppp2r2b*, *tbp*, *htt*, *jph3*, *ar*, *atn*, and gene encoding TCF4).

Gene Expression

[0823] Expression of a target gene containing CAG or CTG repeats will be assayed by techniques known in the field. These assays include, but are not limited to quantitative reverse transcription polymerase chain reaction (RT-PCR), microarray, or multiplexed RNA sequencing (RNA-seq), with the chosen assay measuring either total expression, or the allele specific expression of the target gene. Exemplary assays are found at: Freeman W M et al., "Quantitative RT-PCR: pitfalls and potential", *BioTechniques* 1999, 26, 112-125; Dudley A M et al., "Measuring absolute expression with microarrays with a calibrated reference sample and an extended signal intensity range", *PNAS USA* 2002, 99(11), 7554-7559; Wang Z et al., "RNA-Seq: a revolutionary tool for transcriptomics" *Nature Rev. Genetics* 2009, 10, 57-63.

[0824] Production of the translation product of the target gene will be assayed by techniques known in the field. These assays include, but are not limited to Western blot assay, with the chosen assay measuring either total protein expression, or allele specific expression of the target gene.

[0825] For use in assay, two tissue models and two animal models are contemplated.

Disease Model I: Human Cell Culture

[0826] This model will constitute patient-derived cells, including fibroblasts, induced pluripotent stem cells and cells differentiated from stem cells. Attention will be made in particular to cell types that show impacts of the disease, e.g., neuronal cell types.

Disease Model I: Murine Cell Culture

[0827] This model will constitute cell cultures from mice from tissues that are particularly responsible for disease symptoms, which will include fibroblasts, induced pluripotent stem cells and cells differentiated from stem cells and primary cells that show impacts of the disease, e.g., neuronal cell types.

Disease Model III: Murine

[0828] This model will constitute mice whose genotypes contain the relevant number of repeats for the disease phenotype—these models should show the expected altered gene expression (e.g., modulation of expression of the target gene).

Disease Model IV: Murine

[0829] This model will constitute mice whose genotypes contain a knock in of the human genetic locus from a diseased patient—these models should show the expected altered gene expression (e.g., modulation of expression of the target gene).

[0830] All references, patents or applications, U.S. or foreign, cited in the application are hereby incorporated by reference as if written herein in their entireties. Where any inconsistencies arise, material literally disclosed herein controls.

[0831] From the foregoing description, one skilled in the art can easily ascertain the essential characteristics of this disclosure, and without departing from the spirit and scope thereof, can make various changes and modifications of the disclosure to adapt it to various usages and conditions.

SEQUENCE LISTING

Sequence total quantity: 48

SEQ ID NO: 1	moltype = AA length = 13	
FEATURE	Location/Qualifiers	
source	1..13	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 1		
GRKKRRQRRR PPQ		13
SEQ ID NO: 2	moltype = AA length = 16	
FEATURE	Location/Qualifiers	
source	1..16	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 2		
RQIKIWFQNR RMKWKK		16
SEQ ID NO: 3	moltype = AA length = 18	
FEATURE	Location/Qualifiers	
source	1..18	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 3		
KLALKLALKL LKAALKLA		18
SEQ ID NO: 4	moltype = AA length = 26	
FEATURE	Location/Qualifiers	
VARIANT	6	
	note = S or A	
source	1..26	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 4		
GWTLNXYGLL GKINLKALAA LAKKIL		26
SEQ ID NO: 5	moltype = AA length = 17	
FEATURE	Location/Qualifiers	
source	1..17	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 5		
NAKTRRHERR RKLAIER		17
SEQ ID NO: 6	moltype = AA length = 8	
FEATURE	Location/Qualifiers	
source	1..8	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 6		
RRRRRRRR		8
SEQ ID NO: 7	moltype = AA length = 9	
FEATURE	Location/Qualifiers	
source	1..9	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 7		
RRRRRRRRR		9
SEQ ID NO: 8	moltype = AA length = 17	
FEATURE	Location/Qualifiers	
source	1..17	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 8		
GALFLGFLGA AGSTMGA		17
SEQ ID NO: 9	moltype = AA length = 21	

-continued

FEATURE	Location/Qualifiers	
source	1..21	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 9		
KETWETWWT EWSQPKKKRK V		21
SEQ ID NO: 10	moltype = AA length = 18	
FEATURE	Location/Qualifiers	
source	1..18	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 10		
LLIILRRRIR KQAHASK		18
SEQ ID NO: 11	moltype = AA length = 16	
FEATURE	Location/Qualifiers	
source	1..16	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 11		
YTAIAWKAF IRKLK		16
SEQ ID NO: 12	moltype = AA length = 17	
FEATURE	Location/Qualifiers	
source	1..17	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 12		
IAWKAFIRK LRKGPLG		17
SEQ ID NO: 13	moltype = AA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 13		
MVTVLFRRLR IRRACGPPRV RV		22
SEQ ID NO: 14	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 14		
GLWRALWRL RSLWRLWRA		20
SEQ ID NO: 15	moltype = AA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 15		
RRRRRRQIK IWFQNRMMK KKG		24
SEQ ID NO: 16	moltype = AA length = 23	
FEATURE	Location/Qualifiers	
SITE	2	
	note = Acp	
SITE	5	
	note = Acp	
SITE	8	
	note = Acp	
source	1..23	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 16		
RXRRXRXRI KILFQNRMMK WKK		23
SEQ ID NO: 17	moltype = AA length = 24	
FEATURE	Location/Qualifiers	
SITE	2	
	note = Acp	
SITE	5	
	note = Acp	
SITE	8	

-continued

SITE	note = Acp 11	
	note = D-Lys	
SITE	17	
	note = D-Arg	
SITE	24	
	note = bAla	
source	1..24	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 17		
RXRRXRRXRI KILFQNRMMK WHKA		24
SEQ ID NO: 18	moltype = AA length = 24	
FEATURE	Location/Qualifiers	
SITE	2	
	note = Acp	
SITE	5	
	note = Acp	
SITE	8	
	note = Acp	
SITE	17	
	note = D-Arg	
SITE	24	
	note = bAla	
source	1..24	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 18		
RXRRXRRXRI HILFQNRMMK WHKA		24
SEQ ID NO: 19	moltype = AA length = 22	
FEATURE	Location/Qualifiers	
SITE	2	
	note = Acp	
SITE	5	
	note = bAla	
SITE	8	
	note = Acp	
SITE	16	
	note = Acp	
SITE	18	
	note = bAla	
SITE	20	
	note = Acp	
SITE	22	
	note = bAla	
source	1..22	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 19		
RXRRARRXRI LFQYRXRAX RA		22
SEQ ID NO: 20	moltype = AA length = 22	
FEATURE	Location/Qualifiers	
SITE	2	
	note = Acp	
SITE	5	
	note = bAla	
SITE	8	
	note = Acp	
SITE	16	
	note = Acp	
SITE	18	
	note = Acp	
SITE	20	
	note = Acp	
SITE	22	
	note = bAla	
source	1..22	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 20		
RXRRARRXRI LFQYRXRXRX RA		22
SEQ ID NO: 21	moltype = AA length = 17	

-continued

FEATURE	Location/Qualifiers	
SITE	2	
	note = Acp	
SITE	5	
	note = Acp	
SITE	13	
	note = Acp	
SITE	16	
	note = Acp	
source	1..17	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 21		
RXRRXRILFQ YRXXRXR		17
SEQ ID NO: 22	moltype = AA length = 22	
FEATURE	Location/Qualifiers	
SITE	2	
	note = bAla	
SITE	5	
	note = Acp	
SITE	8	
	note = bAla	
SITE	16	
	note = bAla	
SITE	18	
	note = Acp	
SITE	20	
	note = bAla	
SITE	22	
	note = bAla	
source	1..22	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 22		
RARRXRRARI LFQYRAXRA RA		22
SEQ ID NO: 23	moltype = AA length = 22	
FEATURE	Location/Qualifiers	
SITE	2	
	note = bAla	
SITE	5	
	note = Acp	
SITE	8	
	note = bAla	
SITE	16	
	note = Acp	
SITE	18	
	note = bAla	
SITE	20	
	note = Acp	
SITE	22	
	note = bAla	
source	1..22	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 23		
RARRXRRARI LFQYRXRAX RA		22
SEQ ID NO: 24	moltype = AA length = 21	
FEATURE	Location/Qualifiers	
SITE	2	
	note = bAla	
SITE	5	
	note = Acp	
SITE	8	
	note = bAla	
SITE	16	
	note = Acp	
SITE	19	
	note = Acp	
SITE	21	
	note = bAla	
source	1..21	
	mol_type = protein	
	organism = synthetic construct	

-continued

SEQUENCE: 24
RARRXRRARI LFQYRXXRXR A 21

SEQ ID NO: 25 moltype = AA length = 21
FEATURE Location/Qualifiers
SITE 2
 note = bAla
SITE 5
 note = Acp
SITE 8
 note = bAla
SITE 16
 note = Acp
SITE 18
 note = bAla
SITE 20
 note = Acp
SITE 21
 note = bAla
source 1..21
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 25
RARRXRRARI LFQYRXRARX A 21

SEQ ID NO: 26 moltype = AA length = 21
FEATURE Location/Qualifiers
SITE 2
 note = Acp
SITE 5
 note = bAla
SITE 8
 note = Acp
SITE 16
 note = Acp
SITE 19
 note = Acp
SITE 21
 note = bAla
source 1..21
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 26
RXRRARRXRI LFQYRXXRXR A 21

SEQ ID NO: 27 moltype = AA length = 21
FEATURE Location/Qualifiers
SITE 2
 note = Acp
SITE 5
 note = bAla
SITE 8
 note = Acp
SITE 16
 note = Acp
SITE 18
 note = bAla
SITE 20
 note = Acp
SITE 21
 note = bAla
source 1..21
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 27
RXRRARRXRI LFQYRXRARX A 21

SEQ ID NO: 28 moltype = AA length = 22
FEATURE Location/Qualifiers
SITE 2
 note = Acp
SITE 5
 note = bAla
SITE 8
 note = Acp
SITE 16

-continued

SITE	note = Acp 18	
	note = bAla	
SITE	20	
	note = Acp	
SITE	22	
	note = bAla	
source	1..22	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 28		
RXRRARRXY QFLIRXRAX		22
SEQ ID NO: 29	moltype = AA length = 22	
FEATURE	Location/Qualifiers	
SITE	2	
	note = Acp	
SITE	5	
	note = bAla	
SITE	8	
	note = Acp	
SITE	16	
	note = Acp	
SITE	18	
	note = bAla	
SITE	20	
	note = Acp	
SITE	22	
	note = bAla	
source	1..22	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 29		
RXRRARRXI QFLIRXRAX		22
SEQ ID NO: 30	moltype = AA length = 21	
FEATURE	Location/Qualifiers	
SITE	2	
	note = Acp	
SITE	5	
	note = bAla	
SITE	8	
	note = Acp	
SITE	15	
	note = Acp	
SITE	17	
	note = bAla	
SITE	19	
	note = Acp	
SITE	21	
	note = bAla	
source	1..21	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 30		
RXRRARRXQ FLIRXRAXR		21
SEQ ID NO: 31	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
SITE	2	
	note = Acp	
SITE	5	
	note = bAla	
SITE	8	
	note = Acp	
SITE	14	
	note = Acp	
SITE	16	
	note = bAla	
SITE	18	
	note = Acp	
SITE	20	
	note = bAla	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	

-continued

SEQUENCE: 31
RXRRARRXQ FLRXRARXRA 20

SEQ ID NO: 32 moltype = AA length = 21
FEATURE Location/Qualifiers
SITE 2
 note = Acp
SITE 5
 note = bAla
SITE 8
 note = Acp
SITE 15
 note = Acp
SITE 17
 note = bAla
SITE 19
 note = Acp
SITE 21
 note = bAla
source 1..21
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 32
RXRRARRXYR FLIRXRARXR A 21

SEQ ID NO: 33 moltype = AA length = 22
FEATURE Location/Qualifiers
SITE 2
 note = Acp
SITE 5
 note = bAla
SITE 8
 note = Acp
SITE 16
 note = Acp
SITE 18
 note = bAla
SITE 20
 note = Acp
SITE 22
 note = bAla
source 1..22
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 33
RXRRARRXRF QILYXRARX RA 22

SEQ ID NO: 34 moltype = AA length = 21
FEATURE Location/Qualifiers
SITE 2
 note = Acp
SITE 5
 note = bAla
SITE 8
 note = Acp
SITE 15
 note = Acp
SITE 17
 note = bAla
SITE 19
 note = Acp
SITE 21
 note = bAla
source 1..21
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 34
RXRRARRXYR FRLIXRARXR A 21

SEQ ID NO: 35 moltype = AA length = 21
FEATURE Location/Qualifiers
SITE 2
 note = Acp
SITE 5
 note = bAla
SITE 8

-continued

SITE	note = Acp 15	
SITE	note = Acp 17	
SITE	note = bAla 19	
SITE	note = Acp 21	
SITE	note = bAla 21	
source	1..21 mol_type = protein organism = synthetic construct	
SEQUENCE: 35		
RXRRARRXIL FRYRXRARXR A		21
SEQ ID NO: 36	moltype = AA length = 13	
FEATURE	Location/Qualifiers	
SITE	11	
SITE	note = Acp 12	
SITE	note = L-bis-homopropargylglycine 1..13	
source	mol_type = protein organism = synthetic construct	
SEQUENCE: 36		
RRLSYSRRRF XXG		13
SEQ ID NO: 37	moltype = AA length = 19	
FEATURE	Location/Qualifiers	
SITE	17	
SITE	note = Acp 18	
SITE	note = L-bis-homopropargylglycine 1..19	
source	mol_type = protein organism = synthetic construct	
SEQUENCE: 37		
RRLSYSRRRF PFVYLIXXG		19
SEQ ID NO: 38	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
source	1..20 mol_type = protein organism = synthetic construct	
SEQUENCE: 38		
ARTKQTARKS TGGKAPRKQL		20
SEQ ID NO: 39	moltype = AA length = 8	
FEATURE	Location/Qualifiers	
source	1..8 mol_type = protein organism = synthetic construct	
SEQUENCE: 39		
CWKKLCFK		8
SEQ ID NO: 40	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
source	1..20 mol_type = protein organism = synthetic construct	
SEQUENCE: 40		
ARTKQTARKS TGGKAPRKQL		20
SEQ ID NO: 41	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
source	1..20 mol_type = protein organism = synthetic construct	
SEQUENCE: 41		
ARTKQTARKS TGGKAPRKQL		20
SEQ ID NO: 42	moltype = AA length = 6	
FEATURE	Location/Qualifiers	
SITE	2	
SITE	note = Acp 5	

-continued

source	note = bAla 1..6 mol_type = protein organism = synthetic construct	
SEQUENCE: 42 RXRRAR		6
SEQ ID NO: 43 FEATURE source	moltype = AA length = 5 Location/Qualifiers 1..5 mol_type = protein organism = synthetic construct	
SEQUENCE: 43 ILFQY		5
SEQ ID NO: 44 FEATURE REGION source	moltype = AA length = 10 Location/Qualifiers 1..10 note = This sequence may encompass 5-10 residues 1..10 mol_type = protein organism = synthetic construct	
SEQUENCE: 44 RRRRRRRRRR		10
SEQ ID NO: 45 FEATURE source	moltype = AA length = 4 Location/Qualifiers 1..4 mol_type = protein organism = synthetic construct	
SEQUENCE: 45 QFLY		4
SEQ ID NO: 46 FEATURE SITE SITE SITE SITE source	moltype = AA length = 8 Location/Qualifiers 2 note = Acp 4 note = bAla 6 note = Acp 8 note = bAla 1..8 mol_type = protein organism = synthetic construct	
SEQUENCE: 46 RXRARXRA		8
SEQ ID NO: 47 FEATURE source	moltype = AA length = 4 Location/Qualifiers 1..4 mol_type = protein organism = synthetic construct	
SEQUENCE: 47 PWWP		4
SEQ ID NO: 48 FEATURE source	moltype = AA length = 5 Location/Qualifiers 1..5 mol_type = protein organism = synthetic construct	
SEQUENCE: 48 FALKS		5

1.-177. (canceled)

178. A method for modulating transcription of a gene comprising a trinucleotide repeat sequence CTG the method comprising contacting a cell comprising the gene with an agent having a first terminus, a second terminus, and an oligomeric backbone, wherein:

(a) the first terminus comprises a DNA-binding moiety capable of noncovalently binding to the trinucleotide repeat sequence CTG;

(b) the second terminus comprises a protein-binding moiety capable of binding to a regulatory molecule that modulates an expression of the gene comprising the trinucleotide repeat sequence CTG; and

(c) the oligomeric backbone comprises a linker between the first terminus and the second terminus.

179. The method of claim **178**, wherein the DNA-binding moiety is a polyamide selected from of a linear polyamide, a hairpin polyamide, a H-pin polyamide, an overlapped

polyamide, a slipped polyamide, a cyclic polyamide, a tandem polyamide, and an extended polyamide.

180. The method of claim **178**, wherein the trinucleotide repeat comprises at least 20 repeats, at least 50 repeats, at least 100 repeats, at least 200 repeats, at least 500 repeats, or at least 1000 repeats.

181. The method of claim **178**, wherein the protein-binding moiety is capable of binding to a regulatory molecule that is selected from the group consisting of a CREB binding protein (CBP), a P300, an O-linked β -N-acetylglucosamine-transferase (OGT), a P300-CBP-associated-factor (PCAF), a histone methyltransferase, a histone demethylase, a chromodomain, a cyclin-dependent-kinase-9 (CDK9), a nucleosome-remodeling-factor (NURF), a bromodomain-PHD-finger-transcription-factor (BPTF), a ten-eleven-translocation-enzyme (TET), a methylcytosine-dioxygenase (TET1), a histone acetyltransferase (HAT), a histone deacetylase (HDAC), a host-cell-factor-1 (HCF1), an octamer-binding-transcription-factor (OCT1), a P-TEFb, a cyclin-T1, a PRC2, a DNA-demethylase, a helicase, an acetyltransferase, a histone-deacetylase, a bromodomain-containing protein and a methylated histone lysine protein.

182. The method of claim **178**, wherein the protein-binding moiety is selected from the group consisting of a bromodomain inhibitor, a BPTF inhibitor, a methylcytosine dioxygenase inhibitor, a DNA demethylase inhibitor, a helicase inhibitor, an acetyltransferase inhibitor, a histone deacetylase inhibitor, a CDK-9 inhibitor, a positive transcription elongation factor inhibitor, and a polycomb repressive complex inhibitor.

183. The method of claim **178**, wherein the second terminus does not comprise a moiety that binds to a bromodomain protein.

184. The method of claim **178**, wherein the protein-binding moiety does not comprise JQ1, iBET762, OTX015, RVX208, or AU1.

185. The method of claim **178**, wherein the protein-binding moiety binds the regulatory molecule with an affinity of less than 200 nM.

186. The method of claim **178**, wherein the linker has a length of less than about 50 Angstroms.

187. The method of claim **178**, wherein the gene is chosen from DMPK or TCF4.

188. The method of claim **187**, wherein the DNA-binding moiety is capable of selectively binding to a CTG trinucleotide repeat sequence of DMPK or TCF4.

189. The method of claim **188**, wherein the method comprises decreasing DMPK or TCF4 expression.

190. The method of claim **188**, wherein the method comprises a 20%, 50%, 80%, 90%, 95%, or 99% decrease in expression of DMPK or TCF4.

191. The method of claim **178**, wherein the method further comprises treating a disease mediated by transcription of an allele of DMPK or TCF4 comprising the CTG trinucleotide repeat sequence in a patient in need thereof.

192. The method of claim **191**, wherein the disease is myotonic dystrophy type 1 (DM1).

193. The method of claim **191**, wherein the disease is Fuchs Endothelial Corneal Dystrophy

* * * * *