



US 20250277269A1

(19) **United States**

(12) **Patent Application Publication**
HATCHWELL et al.

(10) **Pub. No.: US 2025/0277269 A1**

(43) **Pub. Date: Sep. 4, 2025**

(54) **METHODS FOR ASSESSING RISK OF
DEVELOPING A VIRAL DISEASE USING A
GENETIC TEST**

(60) Provisional application No. 62/524,324, filed on Jun.
23, 2017, provisional application No. 62/454,676,
filed on Feb. 3, 2017.

(71) Applicants: **PML Screening, LLC**, Newport Beach,
CA (US); **Université Paris-Saclay**, Gif
Sur Yvette (FR); **The Assistance
Publique – Hôpitaux de Paris
(APHP)**, Paris (FR); **The Institut
National de la Santé et de la
Recherche Médicale (INSERM)**, Paris
Cedex 13 (FR)

(72) Inventors: **Eli HATCHWELL**, Winchester (GB);
Peggy S. EIS, Fitchburg, WI (US);
Edward B. SMITH, III, New York,
NY (US); **Yassine TAOUFIK**, Paris
(FR)

(21) Appl. No.: **19/199,930**

(22) Filed: **May 6, 2025**

Related U.S. Application Data

(60) Division of application No. 17/141,885, filed on Jan.
5, 2021, now Pat. No. 12,331,357, which is a division
of application No. 16/245,849, filed on Jan. 11, 2019,
now Pat. No. 10,941,448, which is a continuation of
application No. 15/639,591, filed on Jun. 30, 2017,
now Pat. No. 10,240,205.

Publication Classification

(51) **Int. Cl.**

C12Q 1/6883 (2018.01)

A61K 39/00 (2006.01)

C07K 16/28 (2006.01)

G01N 33/569 (2006.01)

(52) **U.S. Cl.**

CPC **C12Q 1/6883** (2013.01); **C07K 16/2839**

(2013.01); **G01N 33/56983** (2013.01); **A61K**

2039/505 (2013.01); **A61K 2039/54** (2013.01);

A61K 2039/545 (2013.01); **C12Q 2600/106**

(2013.01); **C12Q 2600/118** (2013.01); **C12Q**

2600/156 (2013.01); **G01N 2333/025**

(2013.01); **G01N 2800/065** (2013.01); **G01N**

2800/285 (2013.01); **G01N 2800/52** (2013.01)

(57)

ABSTRACT

This document provides methods and materials related to
treating a disease. For example, this document provides
methods for treating a subject's disease based on identifying
the risk of progressive multifocal leukoencephalopathy
PML using a genetic test.

Specification includes a Sequence Listing.




FIG. 1




FIG. 1 (Cont.)




FIG. 2 (Cont. 1)




FIG. 2 (Cont. 2)







FIG. 2 (Cont. 3)










FIG. 6




FIG. 7





FIG. 8








FIG. 10








FIG. 13

METHODS FOR ASSESSING RISK OF DEVELOPING A VIRAL DISEASE USING A GENETIC TEST

CROSS-REFERENCE

[0001] This application is a divisional of U.S. application Ser. No. 17/141,885, filed Jan. 5, 2021, which is a divisional of U.S. application Ser. No. 16/245,849, filed Jan. 11, 2019, now U.S. Pat. No. 10,941,448, issued Mar. 9, 2021, which is a continuation of U.S. application Ser. No. 15/639,591 filed Jun. 30, 2017, now U.S. Pat. No. 10,240,205, issued Mar. 26, 2019, which claims the benefit of U.S. Provisional Application No. 62/524,324, filed Jun. 23, 2017, and U.S. Provisional Application No. 62/454,676, filed Feb. 3, 2017, both of which are incorporated herein by reference in their entireties.

REFERENCE TO A SEQUENCE LISTING

[0002] The instant application contains a Sequence Listing which has been submitted electronically in XML format and is hereby incorporated by reference in its entirety. Said XML copy, created on Apr. 22, 2025, is named 56969-702_404_SL.xml and is 80,955,165 bytes in size.

BACKGROUND OF THE DISCLOSURE

[0003] Progressive multifocal leukoencephalopathy (PML) is a rare and potentially fatal opportunistic infection of the central nervous system that is caused by a ubiquitous polyomavirus, the JC virus (JCV). While JCV is present at very high rates in the general population, PML remains a rare disorder, albeit an important one because of the poor survival and the severe neurological sequelae, and the recently demonstrated association with a variety of useful therapies, for example, natalizumab in multiple sclerosis (MS). A number of risk factors for PML have been described but these are better viewed as necessary but not sufficient. While these risk factors are highly relevant, they do not, on their own, predict who will develop PML, since the vast majority of individuals with these risk factors will not develop the disorder. Other factors need to be considered and there is growing evidence for the role of host genetic factors in susceptibility to PML.

[0004] The ability to more accurately predict who is at risk of developing PML will be of enormous benefit in the context of drug treatment with compounds that are highly effective in their disease context (natalizumab in MS, for example) but carry a small risk of a devastating disorder. There is a need to develop a companion diagnostic testing, in order to effectively exclude those that were at risk of PML, in the process reassuring those with negative tests about their dramatically reduced risk of developing PML.

INCORPORATION BY REFERENCE

[0005] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference. In the event of a conflict between a term herein and a term incorporated by reference, the term herein controls.

BRIEF DESCRIPTION OF THE DRAWINGS

[0006] The novel features of the disclosure are set forth with particularity in the appended claims. A better understanding of the features and advantages of the present disclosure will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the disclosure are utilized, and the accompanying drawings.

[0007] FIG. 1 represents an example of a gene (PRKCB) impacted by germline and acquired CNVs.

[0008] FIG. 2 represents an example of genes (TNFRSF13C and CENPM) impacted by acquired CNVs.

[0009] FIG. 3 represents an example of a gene (PKHD1) impacted by germline and acquired CNVs.

[0010] FIG. 4 represents an example of a gene (BMPR2) impacted by a recurrent CNV (homozygous and heterozygous losses).

[0011] FIG. 5 represents an example of a gene (COMMD6) impacted by a recurrent CNV (e.g., homozygous duplication).

[0012] FIG. 6 represents an example of genes (KCTD7, RABGEF1) directly and potentially impacted by a recurrent CNV (e.g., homozygous duplication).

[0013] FIG. 7 represents an example of a gene (FPR2) impacted by a recurrent CNV (e.g., homozygous duplication).

[0014] FIG. 8 represents an example of a gene (PIK3CD) impacted by a CNV (e.g., homozygous loss).

[0015] FIG. 9 represents an example of a gene (CD180) potentially impacted by an intergenic CNV gain (e.g., homozygous duplication).

[0016] FIG. 10 represents an example of a gene (VDAC1) potentially impacted by an intergenic CNV (homozygous loss).

[0017] FIG. 11 represents an example of genes (EGR1 and ETF1) potentially impacted by an intergenic CNV (homozygous loss).

[0018] FIG. 12 represents an example of a gene (ITSN2) potentially impacted by an intergenic CNV (homozygous loss).

[0019] FIG. 13 represents an example of known and/or predicted protein interactions using the String database for 21 of 43 genes (non-redundant list) reported in Table 7. The number of PML cases found to harbor variants impacting a given gene is indicated next to each gene.

SUMMARY OF THE INVENTION

[0020] Provided herein is a method of treating a condition in a subject in need thereof, comprising: administering a therapeutically effective amount of one or more immunosuppressive medications to the subject, wherein the subject is identified as not having a risk of developing progressive multifocal leukoencephalopathy (PML) by a genetic test. In some embodiments, the subject is identified as not having a high risk of developing PML by a genetic test.

[0021] In some embodiments, the condition is a cancer, an organ transplant, or an autoimmune disease.

[0022] In some embodiments, the condition is an autoimmune disease.

[0023] In some embodiments, the autoimmune disease is selected from the group consisting of Addison disease, Anti-NMDA receptor encephalitis, antisynthetase syndrome, Aplastic anemia, autoimmune anemias, Autoimmune

hemolytic anemia, Autoimmune pancreatitis, Behcet's Disease, bullous skin disorders, Celiac disease—sprue (gluten-sensitive enteropathy), chronic fatigue syndrome, Chronic inflammatory demyelinating polyneuropathy, chronic lymphocytic leukemia, Crohn's disease, Dermatomyositis, Devic's disease, Erythroblastopenia, Evans syndrome, Focal segmental glomerulosclerosis, Granulomatosis with polyangiitis, Graves disease, Graves' ophthalmopathy, Guillain-Barre syndrome, Hashimoto thyroiditis, idiopathic thrombocytopenic purpura (ITP), IgA nephropathy, IgA-mediated autoimmune diseases, IgG4-related disease, Inflammatory bowel disease, Juvenile idiopathic arthritis, Multiple sclerosis, Myasthenia gravis, myeloma, non-Hodgkin's lymphoma, Opsoclonus myoclonus syndrome (OMS), Pemphigoid, Pemphigus, pemphigus vulgaris, Pernicious anemia, polymyositis, Psoriasis, pure red cell aplasia, Reactive arthritis, Rheumatoid arthritis, Sarcoidosis, scleroderma, Sjögren syndrome, Systemic lupus erythematosus, Thrombocytopenic purpura, Thrombotic thrombocytopenic purpura, Type I diabetes, Ulcerative colitis, Vasculitis (e.g., vasculitis associated with anti-neutrophil cytoplasmic antibody), Vitiligo, and combinations thereof.

[0024] In some embodiments, the autoimmune disease is multiple sclerosis or Crohn's disease. In some embodiments, the autoimmune disease is multiple sclerosis. In some embodiments, the multiple sclerosis is a relapsing form of multiple sclerosis. In some embodiments, the multiple sclerosis is relapsing-remitting multiple sclerosis (RRMS). In some embodiments, the multiple sclerosis is primary progressive multiple sclerosis (PPMS). In some embodiments, the multiple sclerosis is secondary progressive multiple sclerosis (SPMS).

[0025] In some embodiments, the one or more immunosuppressive medications comprise a glucocorticoid, cytostatic, antibody, drug acting on immunophilins, interferon, opioid, TNF binding protein, mycophenolate, small biological agent, small molecule, organic compound, or any combination thereof.

[0026] In some embodiments, the one or more immunosuppressive medications comprise abatacept, adalimumab, alefacept, alemtuzumab, anakinra, azathioprine, belimumab, bendamustine, bevacizumab, bortezomib (e.g., Velcade), eculizumab (e.g., Soliris), leflunomide, brentuximab vedotin, capecitabine, carboplatin, cetuximab, chlorambucil, cladribine, cyclophosphamide, cyclosporine, daclizumab, doxorubicin, efalizumab, etanercept, etoposide, fludarabine, gemcitabine, ibritumomab tiuxetan, imatinib, infliximab, lenalidomide, methotrexate, mycophenolate mofetil, natalizumab, oxaliplatin, rituximab, tocilizumab, tofacitinib, ustekinumab, vedolizumab, vincristine, belatacept, cytotoxic chemotherapy, corticosteroids, antithymocyte Ig, basiliximab, muromonab-CD3, mycophenolic acid, prednisone/prednisolone, sirolimus (rapamycin), tacrolimus, dimethyl fumarate, fingolimod, ruxolitinib, interferon beta-1a, interferon beta-1b, glatiramer acetate, peginterferon beta-1a, teriflunomide, mitoxantrone, ocrelizumab, asparaginase, bleomycin, busulfan, carmustine, certolizumab, ibrutinib, idarubicin, idelalisib, hydrocortisone, ifosfamide, levamisole, mercaptopurine, mizoribine, obinutuzumab, ofatumumab, tegafur/gimeracil/oteracil, thiopeta, vinblastine, or any combination thereof.

[0027] In some embodiments, the one or more immunosuppressive medications comprise interferon beta-1a, interferon beta-1b, glatiramer acetate, peginterferon beta-1a,

teriflunomide, fingolimod, dimethyl fumarate, alemtuzumab, mitoxantrone, natalizumab, daclizumab, ocrelizumab, or any combination thereof.

[0028] In some embodiments, the subject has not taken the one or more immunosuppressive medications. In some embodiments, the subject has taken the one or more immunosuppressive medications. In some embodiments, the subject is taking the one or more immunosuppressive medications.

[0029] In some embodiments, the one or more immunosuppressive medications comprise natalizumab (Tysabri). In some embodiments, at least about 10 mg of the natalizumab is administered, for example, at least about 10 mg, at least about 15 mg, at least about 20 mg, at least about 30 mg, at least about 40 mg, at least about 50 mg, at least about 60 mg, at least about 70 mg, at least about 80 mg, at least about 90 mg, at least about 100 mg, at least about 150 mg, at least about 200 mg, at least about 250 mg, or at least about 300 mg of the natalizumab is administered. In some embodiments, at least about 10 mg of the natalizumab is administered via intravenous infusion. In some embodiments, at least about 10 mg of the natalizumab is administered via intravenous infusion in four weeks.

[0030] In some embodiments, about 100 mg to about 500 mg of the natalizumab is administered, for example, about 100 mg to about 200 mg, about 100 mg to about 300 mg, about 100 mg to about 400 mg, about 100 mg to about 500 mg, about 200 mg to about 300 mg, about 200 mg to about 400 mg, about 200 mg to about 500 mg, about 300 mg to about 400 mg, about 300 mg to about 500 mg, or about 400 mg to about 500 mg of the natalizumab is administered. In some embodiments, about 100 mg to about 500 mg of the natalizumab is administered via intravenous infusion. In some embodiments, about 100 mg to about 500 mg of the natalizumab is administered via intravenous infusion in four weeks. In some embodiments, about 300 mg of the natalizumab is administered. In some embodiments, about 300 mg of the natalizumab is administered via intravenous infusion. In some embodiments, about 300 mg of the natalizumab is administered via intravenous infusion in four weeks.

[0031] In some embodiments, the subject does not have one or more genetic variations associated with a risk of developing PML. In some embodiments, the subject does not have one or more genetic variations associated with a high risk of developing PML.

[0032] In some embodiments, the genetic test comprises detecting one or more genetic variations associated with a risk of developing PML in a polynucleic acid sample from the subject. In some embodiments, the genetic test comprises detecting one or more genetic variations associated with a high risk of developing PML in a polynucleic acid sample from the subject.

[0033] In some embodiments, the one or more genetic variations comprise a point mutation, polymorphism, single nucleotide polymorphism (SNP), single nucleotide variation (SNV), translocation, insertion, deletion, amplification, inversion, interstitial deletion, copy number variation (CNV), loss of heterozygosity, or any combination thereof.

[0034] In some embodiments, the one or more genetic variations disrupt or modulate a corresponding gene according to Tables 3 and 6.

[0035] Provided herein is a method of treating a condition in a subject in need of natalizumab therapy, comprising: administering a therapeutically effective amount of natali-

zumab to the subject, wherein the subject is identified as not having one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6.

[0036] Provided herein is a method of reducing a risk of a subject developing progressive multifocal leukoencephalopathy (PML) comprising administering a therapeutically effective amount of natalizumab to the subject, wherein the subject is identified as not having one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6.

[0037] In some embodiments, the condition is multiple sclerosis.

[0038] In some embodiments, the condition is Crohn's disease.

[0039] Provided herein is a method of treating multiple sclerosis comprising administering natalizumab to a subject with multiple sclerosis, wherein the subject is identified as not having one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6.

[0040] Provided herein is a method of treating Crohn's disease comprising administering natalizumab to a subject with Crohn's disease, wherein the subject is identified as not having one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6.

[0041] Provided herein is a method of treating multiple sclerosis comprising testing a subject with multiple sclerosis for the presence of one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6, determining that the subject does not have the one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6, and

[0042] administering natalizumab to the subject that was determined not to have the one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6.

[0043] Provided herein is a method of treating Crohn's disease comprising testing a subject with Crohn's disease for the presence of one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6, determining that the subject does not have the one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6, and administering natalizumab to the subject that was determined not to have the one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6.

[0044] Provided herein is a method of reducing a risk of a subject developing progressive multifocal leukoencephalopathy (PML) comprising testing a subject for the presence of one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6, determining that the subject has at least one of the one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6, and advising against administering natalizumab to the subject that was determined to have at least one of the one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6.

[0045] In some embodiments, the subject has multiple sclerosis.

[0046] In some embodiments, the subject has Crohn's disease.

[0047] Provided herein is a method of treating multiple sclerosis comprising testing a subject with multiple sclerosis for the presence of one or more genetic variations that

disrupt or modulate a corresponding gene according to Tables 3 and 6, determining that the subject has at least one of the one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6, and advising against administering natalizumab to the subject that was determined to have at least one of the one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6.

[0048] Provided herein is a method of treating Crohn's disease comprising testing a subject with Crohn's disease for the presence of one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6, determining that the subject has at least one of the one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6, and advising against administering natalizumab to the subject that was determined to have at least one of the one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6.

[0049] In some embodiments, the advising comprises advising that administering natalizumab is contraindicated.

[0050] In some embodiments, the advising comprises advising that administering natalizumab increases the risk of the subject developing progressive multifocal leukoencephalopathy (PML).

[0051] In some embodiments, the advising comprises advising that administering natalizumab is a factor that increases the risk of the subject developing progressive multifocal leukoencephalopathy (PML).

[0052] In some embodiments, the testing comprises testing the subject for the presence of one or more genetic variations that disrupt or modulate a corresponding gene according to Table 13.

[0053] In some embodiments, the testing comprises testing the subject for the presence of one or more genetic variations that disrupt or modulate a corresponding gene according to Table 14.

[0054] In some embodiments, the testing comprises testing the subject for the presence of one or more genetic variations that disrupt or modulate a corresponding gene according to Table 15.

[0055] In some embodiments, the testing comprises testing the subject for the presence of one or more genetic variations that disrupt or modulate a corresponding gene according to Table 16.

[0056] In some embodiments, the testing comprises testing the subject for the presence of one or more genetic variations that disrupt or modulate a corresponding gene according to Table 17.

[0057] In some embodiments, the testing comprises testing the subject for the presence of one or more genetic variations that disrupt or modulate a corresponding gene according to Table 18.

[0058] In some embodiments, the testing comprises testing the subject for the presence of one or more genetic variations that disrupt or modulate a corresponding gene selected from the group consisting of ALG12, AP3B1, ASH1L, ATL2, ATM, ATR, BACH1, BLM, CHD7, CLCN7, CR2, CX3CR1, DOCK2, DOCK8, EHF, EPG5, FAS, FUK, GF11, GOLGB1, GTPBP4, HIVEP1, HIVEP2, HIVEP3, IFIH1, IGLL1, IL10, IL12B, IL17F, ITK, ITSN2, JAGN1, KITLG, LRBA, LYST, MALT1, MAVS, MCEE, NHEJ1, NOD2, NRIP1, ORAI1, PGM3, PIK3CD, PLCG2, PNP, POLE, PRF1, RBCK1, RBFOX1, RNASEL, RTE1L,

SALL2, SHARPIN, SNAP29, STIM2, STXBP2, TAP1, TBC1D16, TCIRG1, TICAM1, TMEM173, TNFRSF10A, TTC7A, VPS13B, and combinations thereof.

[0059] In some embodiments, the testing comprises testing the subject for the presence of one or more genetic variations that disrupt or modulate a corresponding gene selected from the group consisting of PLCG2, RBCK1, EPG5, IL17F, SHARPIN, PRF1, JAGN1, TAP1, POLE, LRBA, EHF, IL12B, ATL2, NHEJ1, LYST, HIVEP1, AP3B1, TNFRSF10A, PIK3CD, PNP, MCEE, DOCK2, ALG12, and combinations thereof.

[0060] In some embodiments, the testing comprises testing the subject for the presence of one or more genetic variations that disrupt or modulate a corresponding gene selected from the group consisting of PLCG2, IFIH1, TCIRG1, IGLL1, MAVS, SHARPIN, CHD7, CX3CR1, LRBA, HIVEP3, RNASEL, and combinations thereof.

[0061] In some embodiments, the testing comprises testing the subject for the presence of one or more genetic variations that disrupt or modulate a corresponding gene selected from the group consisting of SHARPIN, RTE11, PGM3, TMEM173, CLCN7, MAVS, ORAI1, RBFOX1, MALT1, GFII1, DOCK2, ATM, SNAP29, TICAM1, GTPBP4, BACH1, STXBP2, FAS, GOLGB1, FUK, IL10, ITK, STIM2, ASH1L, TBC1D16, LYST, SALL2, CHD7, BLM, NOD2, IGLL1, TTC7A, KITLG, ATR, ATM, CR2, HIVEP2, ITSN2, DOCK8, VPS13B, NRIP1, and combinations thereof.

[0062] In some embodiments, the testing comprises testing the subject for the presence of one or more genetic variations that disrupt or modulate a corresponding gene selected from the group consisting of SHARPIN, IFIH1, PLCG2, CHD7, and combinations thereof.

[0063] In some embodiments, the testing comprises testing the subject for the presence of one or more genetic variations that disrupt or modulate a corresponding gene selected from the group consisting of PLCG2, POLE, LRBA, EPG5, SHARPIN, and combinations thereof.

[0064] In some embodiments, the testing comprises testing the subject for the presence of one or more genetic variations that disrupt or modulate a corresponding gene selected from the group consisting of PLCG2, CHD7, IFIH1, AP3B1, EPG5, PIK3CD, LRBA, SHARPIN, and combinations thereof.

[0065] In some embodiments, the subject is identified as not having a risk of developing progressive multifocal leukoencephalopathy (PML) by a genetic test. In some embodiments, the subject is identified as not having a high risk of developing progressive multifocal leukoencephalopathy (PML) by a genetic test.

[0066] In some embodiments, the testing comprises assaying a polynucleic acid sample from the subject for the one or more genetic variations.

[0067] In some embodiments, the one or more genetic variations result in a loss of function of the corresponding gene.

[0068] In some embodiments, the corresponding gene comprises a gene selected from the group consisting of gene numbers (GNs) GN1-GN490.

[0069] In some embodiments, the corresponding gene comprises a gene selected from the group consisting of gene numbers (GNs) 1-156 (in Table 3).

[0070] In some embodiments, the corresponding gene comprises a gene selected from the group consisting of gene numbers (GNs) in Table 6.

[0071] In some embodiments, the corresponding gene comprises a gene selected from the group consisting of PLCG2, RBCK1, EPG5, IL17F, SHARPIN, PRF1, JAGN1, TAP1, POLE, LRBA, EHF, IL12B, ATL2, NHEJ1, LYST, HIVEP1, AP3B1, TNFRSF10A, PIK3CD, PNP, MCEE, DOCK2 and ALG12 (see Table 13).

[0072] In some embodiments, the one or more genetic variations are encoded by a sequence with at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to SEQ ID NOs 1-172 or SRN1-SRN363, with 100% sequence identity to SEQ ID NOs 1000-1329, or with at least 80% and less than 100% sequence identity to GN1-GN490, or complements thereof.

[0073] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a CNV with at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to SEQ ID NOs 1-172, or complements thereof.

[0074] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a CNV sub-region (SRN) with at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to SRN1-SRN363, or complements thereof.

[0075] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a single nucleotide variation (SNV) with a sequence of any one of SEQ ID NOs: 1000-1329, or complements thereof.

[0076] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a sequence with at least 80% and less than 100% sequence identity to GN1-GN490, or complements thereof.

[0077] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a single nucleotide variation (SNV) with a sequence of any one of SEQ ID NO: 1000, 1001, 1002, 1009, 1010, 1011, 1012, 1014, 1016, 1017, 1019, 1020, 1028, 1032, 1033, 1034, 1035, 1036, 1037, 1040, 1041, 1043, 1051, 1054, 1056, 1057, 1058, 1059, 1061, 1062, 1063, 1066, 1068, 1069, 1070, 1071, 1073, 1074, 1075, 1076, 1077, 1078, 1080, 1082, 1084, 1090, 1092, 1098, 1099, 1100, 1101, 1104, 1107, 1114, 1116, 1118, 1121, 1122, 1123, 1125, 1126, 1127, 1128, 1129, 1130, 1131, 1133, 1135, 1136, 1137, 1138, 1142, 1146, 1147, 1148, 1150, 1152, 1154, 1157, 1160, 1161, 1165, 1166, 1167, 1168, 1169, 1171, 1174, 1175, 1176, 1177, 1178, 1179, 1180, 1181, 1182, 1183, 1184, 1193, 1194, 1200, 1201, 1202, 1203, 1204, 1208, 1219, 1220, 1221, 1222, 1226, 1227, 1228, 1229, 1230, 1231, 1232, 1235, 1239, 1247, 1248, 1249, 1250, 1251, 1252, 1254, 1255, 1256, 1259, 1260, 1261, 1263, 1264, 1266, 1267, 1273, 1278, 1279, 1283, 1284, 1286, 1287, 1289, 1290, 1291, 1299, 1300, 1301, 1304, 1311, 1327 or 1328 (see Tables 7 and 8), or complements thereof.

[0078] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a single nucleotide variation (SNV) with a sequence of any one of SEQ ID NO: 1011, 1020, 1028, 1032, 1034, 1035, 1036, 1040, 1056, 1069, 1073, 1077, 1101, 1114, 1123, 1125, 1126, 1127, 1135, 1142, 1146, 1147, 1148, 1152, 1154, 1157, 1167, 1174, 1184, 1193, 1194, 1203, 1208, 1221, 1222, 1229, 1235, 1252, 1255, 1256, 1259, 1260, 1261, 1263,

1273, 1278, 1279, 1284, 1287, 1289, 1299 or 1311 (see Table 7), or complements thereof.

[0079] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a single nucleotide variation (SNV) with a sequence of any one of SEQ ID NO: 1000, 1001, 1002, 1009, 1010, 1012, 1014, 1016, 1017, 1019, 1033, 1037, 1041, 1043, 1051, 1054, 1057, 1058, 1059, 1061, 1062, 1063, 1066, 1068, 1070, 1071, 1074, 1075, 1076, 1078, 1080, 1082, 1084, 1090, 1092, 1098, 1099, 1100, 1104, 1107, 1116, 1118, 1121, 1122, 1128, 1129, 1130, 1131, 1133, 1136, 1137, 1138, 1146, 1147, 1150, 1152, 1160, 1161, 1165, 1166, 1168, 1169, 1171, 1175, 1176, 1177, 1178, 1179, 1180, 1181, 1182, 1183, 1200, 1201, 1202, 1204, 1219, 1220, 1226, 1227, 1228, 1230, 1231, 1232, 1239, 1247, 1248, 1249, 1250, 1251, 1252, 1254, 1264, 1266, 1267, 1278, 1279, 1283, 1286, 1290, 1291, 1300, 1301, 1304, 1327 or 1328 (see Table 8), or complements thereof.

[0080] In some embodiments, the one or more genetic variations comprise a genetic variation selected from the group consisting of chr16:81942175 A>G, chr2:163136505 C>G, chr11:67818269 G>A, chr22:23917192 G>T, chr20:3846397 C>T, chr8:145154222, G>A chr8:61654298 T>A, chr3:39323163 A>C, chr4:151199080 G>A, chr1:42047208 C>G, chr2:163124051 C>T, chr1:182554557 C>T, chr8:145154824 A>C, chr20:62305450 C>T, chr22:23915745 G>A, chr6:83884161 C>G, chr11:108202772 G>T, chr5:138856923 C>T, chr16:1510535 C>T, chr20:3843027 C>A, chr12:122064788 G>GT, chr16:7714909 C>T, chr18:56401523 C>T, chr1:92946625 G>C, chr5:169081453 G>C, chr11:108117787 C>T, chr22:21235389 A>G, chr19:4817657 C>T, chr10:1060218 G>A, chr21:30698953 T>G, chr9:304628 G>A, chr19:7712287 G>C, chr10:90771767 G>A, chr3:121415370 T>C, chr16:70503095 A>G, chr1:206945738 C>T, chr5:156593120 C>T, chr4:27019452 C>T, chr1:155317682 C>T, chr17:77926526 C>T, chr1:235840495 G>T, chr14:21993359 G>A, chr8:61757805 C>T, chr15:91306241 G>A, chr16:50741791 C>T, chr22:23915583 T>C, chr2:47205921 C>T, chr12:88900891 C>A, chr3:142281353 C>G, chr11:108123551 C>T, chr1:207641950 C>T, chr6:143092151 T>C, chr2:24431184 C>T, chr2:24432937 C>T, chr9:312134 G>A, chr8:100205255 G>A, chr21:16339852 T>C, and any combination thereof (see Tables 14 and 15).

[0081] In some embodiments, the one or more genetic variations comprise a genetic variation selected from the group consisting of chr16:81942175 A>G, chr2:163136505 C>G, chr11:67818269 G>A, chr22:23917192 G>T, chr20:3846397 C>T, chr8:145154222, G>A chr8:61654298 T>A, chr3:39323163 A>C, chr4:151199080 G>A, chr1:42047208 C>G, chr2:163124051 C>T, chr1:182554557 C>T, and any combination thereof (see Table 14).

[0082] In some embodiments, the one or more genetic variations comprise a genetic variation selected from the group consisting of chr8:145154824 A>C, chr20:62305450 C>T, chr22:23915745 G>A, chr6:83884161 C>G, chr11:108202772 G>T, chr5:138856923 C>T, chr16:1510535 C>T, chr20:3843027 C>A, chr12:122064788 G>GT, chr16:7714909 C>T, chr18:56401523 C>T, chr1:92946625 G>C, chr5:169081453 G>C, chr11:108117787 C>T, chr22:21235389 A>G, chr19:4817657 C>T, chr10:1060218 G>A, chr21:30698953 T>G, chr9:304628 G>A, chr19:7712287 G>C, chr10:90771767 G>A, chr3:121415370 T>C, chr16:70503095 A>G, chr1:206945738 C>T, chr5:156593120

C>T, chr4:27019452 C>T, chr1:155317682 C>T, chr17:77926526 C>T, chr1:235840495 G>T, chr14:21993359 G>A, chr8:61757805 C>T, chr15:91306241 G>A, chr16:50741791 C>T, chr22:23915583 T>C, chr2:47205921 C>T, chr12:88900891 C>A, chr3:142281353 C>G, chr11:108123551 C>T, chr1:207641950 C>T, chr6:143092151 T>C, chr2:24431184 C>T, chr2:24432937 C>T, chr9:312134 G>A, chr8:100205255 G>A, chr21:16339852 T>C, and any combination thereof (see Table 15).

[0083] In some embodiments, the SNV is a heterozygous SNV.

[0084] In some embodiments, the SNV is a homozygous SNV.

[0085] In some embodiments, the one or more genetic variations comprise a pair of single nucleotide variations (SNVs), wherein the pair of SNVs are encoded by any one of SEQ ID NO pairs: 1003 and 1004, 1003 and 1005, 1006 and 1007, 1024 and 1025, 1030 and 1031, 1047 and 1048, 1049 and 1050, 1063 and 1064, 1063 and 1065, 1063 and 1066, 1075 and 1076, 1091 and 1093, 1091 and 1096, 1093 and 1095, 1094 and 1097, 1098 and 1099, 1098 and 1100, 1099 and 1100, 1102 and 1103, 1104 and 1106, 1104 and 1107, 1104 and 1108, 1104 and 1109, 1104 and 1110, 1104 and 1111, 1104 and 1112, 1110 and 1111, 1112 and 1113, 1119 and 1120, 1124 and 1125, 1124 and 1126, 1125 and 1126, 1140 and 1141, 1142 and 1144, 1146 and 1151, 1147 and 1148, 1147 and 1149, 1153 and 1146, 1153 and 1147, 1155 and 1156, 1160 and 1161, 1165 and 1166, 1186 and 1187, 1188 and 1193, 1189 and 1193, 1191 and 1192, 1191 and 1193, 1191 and 1195, 1192 and 1193, 1192 and 1195, 1196 and 1197, 1206 and 1207, 1210 and 1218, 1211 and 1213, 1212 and 1213, 1213 and 1215, 1213 and 1216, 1213 and 1217, 1233 and 1238, 1242 and 1243, 1245 and 1246, 1263 and 1260, 1269 and 1279, 1270 and 1279, 1270 and 1282, 1271 and 1279, 1274 and 1279, 1278 and 1279, 1278 and 1281, 1279 and 1280, 1279 and 1281, 1279 and 1282, 1292 and 1293, 1296 and 1297, 1305 and 1314, 1306 and 1310, 1313 and 1321 or 1315 and 1322 (see Table 9 or Tables 9 and 7 for a subset), or complements thereof.

[0086] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a CNV with at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to any one of SEQ ID NOs 157, 2, 140, 65, 26, 14 or 45 (see Tables 7 and 8), or complements thereof.

[0087] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a CNV with at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to any one of SEQ ID NOs 2, 140, 65, 26, 14 or 45 (see Table 7), or complements thereof.

[0088] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a CNV with at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to SEQ ID NO 157 (see Table 8), or a complement thereof.

[0089] In some embodiments, the one or more genetic variations comprise a CNV-SNV pair comprising a CNV and a single nucleotide variation (SNV), wherein the SNV of the CNV-SNV pair is encoded by any one of SEQ ID NO pairs: 146 and 1301, 85 and 1173, 58 and 1107, 58 and 1104, 91 and 1199, 103 and 1225, 103 and 1086 or 41 and 1223 (see Tables 1 and 10), or complements thereof.

[0090] In some embodiments, the one or more genetic variations comprise a genetic variation selected from the group consisting of: chr8:145154222 G>A, chr2:163136505 C>G, chr16:81942175 A>G, chr8:61654298 T>A, and combinations thereof (see Tables 14 and 16).

[0091] In some embodiments, the one or more genetic variations disrupt or modulate one or more of the following genes: PLCG2, POLE, LRBA, EPG5 and SHARPIN (see Table 17).

[0092] In some embodiments, the one or more genetic variations disrupt or modulate one or more of the following genes: PLCG2, CHD7, IFIH1, AP3B1, EPG5, PIK3CD, LRBA and SHARPIN (see Table 18).

[0093] In some embodiments, the corresponding gene encodes a transcript with a sequence that has at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to any one of SEQ ID NOS 173-455 or 1500-2177 (see Tables 4 and 12), or complements thereof.

[0094] In some embodiments, the corresponding gene encodes a transcript with a sequence that has at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to any one of SEQ ID NOS 173-455 (see Table 4), or complements thereof.

[0095] In some embodiments, the corresponding gene encodes a transcript with a sequence that has at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to any one of SEQ ID NOS 1500-2177 (see Table 12), or complements thereof.

[0096] In some embodiments, the one or more genetic variations comprise 2 or 3 or 4 or 5 or more genetic variations.

[0097] In some embodiments, the one or more genetic variations comprise 10 or more genetic variations.

[0098] In some embodiments, the one or more genetic variations comprise 20 or more genetic variations.

[0099] In some embodiments, the one or more genetic variations comprise 50 or more genetic variations.

[0100] In some embodiments, the genetic test or the testing comprises microarray analysis, PCR, sequencing, nucleic acid hybridization, or any combination thereof.

[0101] In some embodiments, the genetic test or the testing comprises microarray analysis selected from the group consisting of a Comparative Genomic Hybridization (CGH) array analysis and an SNP array analysis.

[0102] In some embodiments, the genetic test or the testing comprises sequencing, wherein the sequencing is selected from the group consisting of Massively Parallel Signature Sequencing (MPSS), polony sequencing, 454 pyrosequencing, Illumina sequencing, Illumina (Solexa) sequencing using 10x Genomics library preparation, SOLiD sequencing, ion semiconductor sequencing, DNA nanoball sequencing, heliscope single molecule sequencing, single molecule real time (SMRT) sequencing, RNAP sequencing, Nanopore DNA sequencing, sequencing by hybridization, and microfluidic Sanger sequencing.

[0103] In some embodiments, the genetic test or the testing comprises analyzing a whole genome of the subject.

[0104] In some embodiments, the genetic test or the testing comprises analyzing a whole exome of the subject.

[0105] In some embodiments, the genetic test or the testing comprises analyzing nucleic acid information that has already been obtained for a whole genome or a whole exome of the subject.

[0106] In some embodiments, the nucleic acid information is obtained from an in silico analysis.

[0107] In some embodiments, the subject is a human subject.

[0108] In some embodiments, the polynucleic acid sample comprises a polynucleic acid from blood, saliva, urine, serum, tears, skin, tissue, or hair of the subject.

[0109] In some embodiments, the method further comprises treating the subject with an agent that reduces a viral load in the subject.

[0110] In some embodiments, the immunosuppressive agent is administered after the viral load is reduced.

[0111] In some embodiments, the viral load is a JCV viral load.

[0112] In some embodiments, the agent that reduces the viral load is an agent that targets JCV.

[0113] In some embodiments, the method further comprises analyzing for a presence of JCV in a biological sample from the subject. In some embodiments, the method comprises a JCV-antibody test. In some embodiments, the JCV-antibody test has a negative result. In some embodiments, the JCV-antibody test does not detect a presence of JCV in the biological sample from the subject. In some embodiments, the JCV-antibody test detects a presence of JCV in the biological sample from the subject.

[0114] In some embodiments, the analyzing for a presence of JCV comprises contacting a JCV detection reagent to the biological sample.

[0115] In some embodiments, the JCV detection reagent is selected from the group consisting of an anti-JCV antibody, a JCV specific primer, and combinations thereof.

[0116] Provided herein is a method of treating a condition in a subject in need thereof, comprising: administering a therapeutically effective amount of one or more immunosuppressive medications to the subject, and one or more agents that reduce a viral load in the subject, wherein the subject is identified as not having a risk of developing progressive multifocal leukoencephalopathy (PML) by a genetic test. In some embodiments, the subject is identified as not having a high risk of developing progressive multifocal leukoencephalopathy (PML) by a genetic test.

[0117] Provided herein is a method of treating a condition in a subject in need thereof, comprising: analyzing a polynucleic acid sample from the subject for one or more genetic variations that disrupt or modulate a gene of GN1-GN490, wherein a genetic variation of the one or more genetic variations that disrupt or modulate a gene of GN1-GN490 is not present in the polynucleic acid sample; identifying the subject as not having a risk of developing PML; administering a therapeutically effective amount of one or more immunosuppressive medications to the subject. In some embodiments, the method comprises identifying the subject as not having a high risk of developing PML.

[0118] Provided herein is a method of identifying a subject as having a risk of developing PML, comprising: analyzing a polynucleic acid sample from the subject for one or more genetic variations that disrupt or modulate a gene of GN1-GN490, wherein a genetic variation of the one or more genetic variations that disrupt or modulate a gene of GN1-GN490 is not present in the polynucleic acid sample; identifying the subject as not having a risk of developing PML. In some embodiments, the method comprises identifying the subject as not having a high risk of developing PML.

[0119] Provided herein is a method of identifying a subject as having a risk of developing progressive multifocal leukoencephalopathy (PML) comprising obtaining a genetic test result from a polynucleic acid sample from a subject, and identifying the subject as having a risk of developing PML based on the genetic test result; wherein the subject is immunosuppressed.

[0120] Provided herein is a method of monitoring a subject as having a risk of developing progressive multifocal leukoencephalopathy (PML) comprising obtaining a genetic test result from a polynucleic acid sample from a subject, and identifying the subject as having an increased risk of developing PML based on the genetic test result; wherein the subject is immunosuppressed.

[0121] In some embodiments, the subject is on an immunosuppressive therapy.

[0122] Provided herein is a method of identifying a subject as having a risk of developing progressive multifocal leukoencephalopathy (PML) comprising detecting one or more genetic variations that disrupt or modulate a gene of GN1-GN490 in a polynucleic acid sample from a subject, and identifying the subject as having a risk of developing PML; wherein the subject is immunosuppressed.

[0123] Provided herein is a method of identifying a subject as having a risk of developing progressive multifocal leukoencephalopathy (PML) comprising: analyzing a polynucleic acid sample from the subject for one or more genetic variations that disrupt or modulate a gene of GN1-GN490, wherein a genetic variation of the one or more genetic variations that disrupt or modulate a gene of GN1-GN490 is present in the polynucleic acid sample; identifying the subject as having a risk of developing PML; wherein the subject is immunosuppressed. In some embodiments, the method comprises identifying the subject as having a high risk of developing PML.

[0124] In some embodiments, the subject has HIV. In some embodiments, the subject has HIV infection. In some embodiments, the subject is at risk of HIV infection.

[0125] In some embodiments, the condition is a cancer, a hematologic malignancy, an organ transplant, or an autoimmune disease. In some embodiments, the condition is idiopathic CD4⁺ lymphocytopenia (ICL).

[0126] In some embodiments, the condition is an autoimmune disease.

[0127] In some embodiments, the autoimmune disease is selected from the group consisting of Addison disease, Behcet's Disease, Inflammatory bowel disease, Celiac disease—sprue (gluten-sensitive enteropathy), Crohn's disease, Dermatomyositis, Focal segmental glomerulosclerosis, Graves disease, Hashimoto thyroiditis, Multiple sclerosis, Myasthenia gravis, Pemphigus, Pemphigoid, Aplastic anemia, Pernicious anemia, Autoimmune hemolytic anemia, Erythroblastopenia, Thrombocytopenic purpura, Evans syndrome, Vasculitis, Granulomatosis with polyangiitis, Chronic inflammatory demyelinating polyneuropathy, Guillain-Barre syndrome, Anti-NMDA receptor encephalitis, Devic's disease, Autoimmune pancreatitis, Opsoclonus myoclonus syndrome, IgG4-related disease, Psoriasis, Reactive arthritis, Rheumatoid arthritis, Juvenile idiopathic arthritis, Sarcoidosis, Sjögren syndrome, Systemic lupus erythematosus, Type I diabetes, Vitiligo, or Ulcerative colitis.

[0128] In some embodiments, the autoimmune disease is multiple sclerosis or Crohn's disease.

[0129] In some embodiments, the one or more immunosuppressive medications comprise a glucocorticoid, cytostatic, antibody, drug acting on immunophilins, interferon, opioid, TNF binding protein, mycophenolate, small biological agent, small molecule, organic compound, or any combination thereof.

[0130] In some embodiments, the one or more immunosuppressive medications comprise a interferon beta-1a, interferon beta-1b, glatiramer acetate, peginterferon beta-1a, teriflunomide, fingolimod, dimethyl fumarate, alemtuzumab, mitoxantrone, natalizumab, daclizumab, ocrelizumab, or any combination thereof.

[0131] In some embodiments, the one or more immunosuppressive medications comprise natalizumab (Tysabri).

[0132] In some embodiments, the one or more genetic variations comprise a point mutation, polymorphism, single nucleotide polymorphisms (SNP), single nucleotide variation (SNV), translocation, insertion, deletion, amplification, inversion, interstitial deletion, copy number variation (CNV), loss of heterozygosity, or any combination thereof.

[0133] In some embodiments, the one or more genetic variations result in a loss of function of the corresponding gene.

[0134] In some embodiments, the corresponding gene comprises a gene selected from the group consisting of gene numbers (GNs) GN1-GN490.

[0135] In some embodiments, the gene comprises a gene selected from the group consisting of gene numbers (GNs) 1-156 (in Table 3).

[0136] In some embodiments, the gene comprises a gene selected from the group consisting of gene numbers (GNs) in Table 6.

[0137] In some embodiments, the gene comprises a gene selected from the group consisting of PLCG2, RBCK1, EPG5, IL17F, SHARPIN, PRF1, JAGN1, TAP1, POLE, LRBA, EHF, IL12B, ATL2, NHEJ1, LYST, HIVEP1, AP3B1, TNFRSF10A, PIK3CD, PNP, MCEE, DOCK2 and ALG12 (see Table 13).

[0138] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a sequence with at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to SEQ ID NOs 1-172 or SRN1-SRN363, with 100% sequence identity to SEQ ID NOs 1000-1329, or with at least 80% and less than 100% sequence identity to GN1-GN490, or complements thereof.

[0139] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a CNV with at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to SEQ ID NOs 1-172, or complements thereof.

[0140] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a CNV sub-region (SRN) with at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to SRN1-SRN363, or complements thereof.

[0141] In some embodiments, the one or more genetic variations are encoded by a single nucleotide variation (SNV) with a sequence of any one of SEQ ID NOs: 1000-1329, or complements thereof.

[0142] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a sequence with at least 80% and less than 100% sequence identity to GN1-GN490, or complements thereof.

[0143] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a single nucleotide variation (SNV) with a sequence of any one of SEQ ID NO: 1000, 1001, 1002, 1009, 1010, 1011, 1012, 1014, 1016, 1017, 1019, 1020, 1028, 1032, 1033, 1034, 1035, 1036, 1037, 1040, 1041, 1043, 1051, 1054, 1056, 1057, 1058, 1059, 1061, 1062, 1063, 1066, 1068, 1069, 1070, 1071, 1073, 1074, 1075, 1076, 1077, 1078, 1080, 1082, 1084, 1090, 1092, 1098, 1099, 1100, 1101, 1104, 1107, 1114, 1116, 1118, 1121, 1122, 1123, 1125, 1126, 1127, 1128, 1129, 1130, 1131, 1133, 1135, 1136, 1137, 1138, 1142, 1146, 1147, 1148, 1150, 1152, 1154, 1157, 1160, 1161, 1165, 1166, 1167, 1168, 1169, 1171, 1174, 1175, 1176, 1177, 1178, 1179, 1180, 1181, 1182, 1183, 1184, 1193, 1194, 1200, 1201, 1202, 1203, 1204, 1208, 1219, 1220, 1221, 1222, 1226, 1227, 1228, 1229, 1230, 1231, 1232, 1235, 1239, 1247, 1248, 1249, 1250, 1251, 1252, 1254, 1255, 1256, 1259, 1260, 1261, 1263, 1264, 1266, 1267, 1273, 1278, 1279, 1283, 1284, 1286, 1287, 1289, 1290, 1291, 1299, 1300, 1301, 1304, 1311, 1327 or 1328 (see Tables 7 and 8), or complements thereof.

[0144] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a single nucleotide variation (SNV) with a sequence of any one of SEQ ID NO: 1011, 1020, 1028, 1032, 1034, 1035, 1036, 1040, 1056, 1069, 1073, 1077, 1101, 1114, 1123, 1125, 1126, 1127, 1135, 1142, 1146, 1147, 1148, 1152, 1154, 1157, 1167, 1174, 1184, 1193, 1194, 1203, 1208, 1221, 1222, 1229, 1235, 1252, 1255, 1256, 1259, 1260, 1261, 1263, 1273, 1278, 1279, 1284, 1287, 1289, 1299 or 1311 (see Table 7), or complements thereof.

[0145] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a single nucleotide variation (SNV) with a sequence of any one of SEQ ID NO: 1000, 1001, 1002, 1009, 1010, 1012, 1014, 1016, 1017, 1019, 1033, 1037, 1041, 1043, 1051, 1054, 1057, 1058, 1059, 1061, 1062, 1063, 1066, 1068, 1070, 1071, 1074, 1075, 1076, 1078, 1080, 1082, 1084, 1090, 1092, 1098, 1099, 1100, 1104, 1107, 1116, 1118, 1121, 1122, 1128, 1129, 1130, 1131, 1133, 1136, 1137, 1138, 1146, 1147, 1150, 1152, 1160, 1161, 1165, 1166, 1168, 1169, 1171, 1175, 1176, 1177, 1178, 1179, 1180, 1181, 1182, 1183, 1200, 1201, 1202, 1204, 1219, 1220, 1226, 1227, 1228, 1230, 1231, 1232, 1239, 1247, 1248, 1249, 1250, 1251, 1252, 1254, 1264, 1266, 1267, 1278, 1279, 1283, 1286, 1290, 1291, 1300, 1301, 1304, 1327 or 1328 (see Table 8), or complements thereof.

[0146] In some embodiments, the one or more genetic variations comprise a genetic variation selected from the group consisting of chr16:81942175 A>G, chr2:163136505 C>G, chr11:67818269 G>A, chr22:23917192 G>T, chr20:3846397 C>T, chr8:145154222, G>A chr8:61654298 T>A, chr3:39323163 A>C, chr4:151199080 G>A, chr1:42047208 C>G, chr2:163124051 C>T, chr1:182554557 C>T, chr8:145154824 A>C, chr20:62305450 C>T, chr22:23915745 G>A, chr6:83884161 C>G, chr11:108202772 G>T, chr5:138856923 C>T, chr16:1510535 C>T, chr20:3843027 C>A, chr12:122064788 G>GT, chr16:7714909 C>T, chr18:56401523 C>T, chr1:92946625 G>C, chr5:169081453 G>C, chr11:108117787 C>T, chr22:21235389 A>G, chr19:4817657 C>T, chr10:1060218 G>A, chr21:30698953 T>G, chr9:304628 G>A, chr19:7712287 G>C, chr10:90771767 G>A, chr3:121415370 T>C, chr16:70503095 A>G, chr1:206945738 C>T, chr5:156593120 C>T, chr4:27019452

C>T, chr1:155317682 C>T, chr17:77926526 C>T, chr1:235840495 G>T, chr14:21993359 G>A, chr8:61757805 C>T, chr15:91306241 G>A, chr16:50741791 C>T, chr22:23915583 T>C, chr2:47205921 C>T, chr12:88900891 C>A, chr3:142281353 C>G, chr11:108123551 C>T, chr1:207641950 C>T, chr6:143092151 T>C, chr2:24431184 C>T, chr2:24432937 C>T, chr9:312134 G>A, chr8:100205255 G>A, chr21:16339852 T>C, and any combination thereof (see Tables 14 and 15).

[0147] In some embodiments, the one or more genetic variations comprise a genetic variation selected from the group consisting of chr16:81942175 A>G, chr2:163136505 C>G, chr11:67818269 G>A, chr22:23917192 G>T, chr20:3846397 C>T, chr8:145154222, G>A chr8:61654298 T>A, chr3:39323163 A>C, chr4:151199080 G>A, chr1:42047208 C>G, chr2:163124051 C>T, chr1:182554557 C>T, and any combination thereof (see Table 14).

[0148] In some embodiments, the one or more genetic variations comprise a genetic variation selected from the group consisting of chr8:145154824 A>C, chr20:62305450 C>T, chr22:23915745 G>A, chr6:83884161 C>G, chr11:108202772 G>T, chr5:138856923 C>T, chr16:1510535 C>T, chr20:3843027 C>A, chr12:122064788 G>GT, chr16:7714909 C>T, chr18:56401523 C>T, chr1:92946625 G>C, chr5:169081453 G>C, chr11:108117787 C>T, chr22:21235389 A>G, chr19:4817657 C>T, chr10:1060218 G>A, chr21:30698953 T>G, chr9:304628 G>A, chr19:7712287 G>C, chr10:90771767 G>A, chr3:121415370 T>C, chr16:70503095 A>G, chr1:206945738 C>T, chr5:156593120 C>T, chr4:27019452 C>T, chr1:155317682 C>T, chr17:77926526 C>T, chr1:235840495 G>T, chr14:21993359 G>A, chr8:61757805 C>T, chr15:91306241 G>A, chr16:50741791 C>T, chr22:23915583 T>C, chr2:47205921 C>T, chr12:88900891 C>A, chr3:142281353 C>G, chr11:108123551 C>T, chr1:207641950 C>T, chr6:143092151 T>C, chr2:24431184 C>T, chr2:24432937 C>T, chr9:312134 G>A, chr8:100205255 G>A, chr21:16339852 T>C, and any combination thereof (see Table 15).

[0149] In some embodiments, the SNV is a heterozygous SNV.

[0150] In some embodiments, the SNV is a homozygous SNV.

[0151] In some embodiments, the one or more genetic variations comprise a pair of single nucleotide variations (SNVs), wherein the pair of SNVs are encoded by any one of SEQ ID NO pairs: 1003 and 1004, 1003 and 1005, 1006 and 1007, 1024 and 1025, 1030 and 1031, 1047 and 1048, 1049 and 1050, 1063 and 1064, 1063 and 1065, 1063 and 1066, 1075 and 1076, 1091 and 1093, 1091 and 1096, 1093 and 1095, 1094 and 1097, 1098 and 1099, 1098 and 1100, 1099 and 1100, 1102 and 1103, 1104 and 1106, 1104 and 1107, 1104 and 1108, 1104 and 1109, 1104 and 1110, 1104 and 1111, 1104 and 1112, 1110 and 1111, 1112 and 1113, 1119 and 1120, 1124 and 1125, 1124 and 1126, 1125 and 1126, 1140 and 1141, 1142 and 1144, 1146 and 1151, 1147 and 1148, 1147 and 1149, 1153 and 1146, 1153 and 1147, 1155 and 1156, 1160 and 1161, 1165 and 1166, 1186 and 1187, 1188 and 1193, 1189 and 1193, 1191 and 1192, 1191 and 1193, 1191 and 1195, 1192 and 1193, 1192 and 1195, 1196 and 1197, 1206 and 1207, 1210 and 1218, 1211 and 1213, 1212 and 1213, 1213 and 1215, 1213 and 1216, 1213 and 1217, 1233 and 1238, 1242 and 1243, 1245 and 1246, 1263 and 1260, 1269 and 1279, 1270 and 1279, 1270 and 1282, 1271 and 1279, 1274 and 1279, 1278 and 1279, 1278

and 1281, 1279 and 1280, 1279 and 1281, 1279 and 1282, 1292 and 1293, 1296 and 1297, 1305 and 1314, 1306 and 1310, 1313 and 1321 or 1315 and 1322 (see Table 9 or Tables 9 and 7 for a subset), or complements thereof.

[0152] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a CNV with at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to any one of SEQ ID NOs 157, 2, 140, 65, 26, 14 or 45 (see Tables 7 and 8), or complements thereof.

[0153] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a CNV with at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to any one of SEQ ID NOs 2, 140, 65, 26, 14 or 45 (see Table 7), or complements thereof.

[0154] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a CNV with at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to SEQ ID NO 157 (see Table 8), or a complement thereof.

[0155] In some embodiments, the one or more genetic variations comprise a CNV-SNV pair comprising a CNV and a single nucleotide variation (SNV), wherein the SNV of the CNV-SNV pair is encoded by any one of SEQ ID NOs 1301, 1173, 1107, 1104, 1199, 1225, 1086 or 1223 (see Table 10), or complements thereof.

[0156] In some embodiments, the one or more genetic variations comprise a genetic variation selected from the group consisting of one or more of the following: chr8:145154222 G>A, chr2:163136505 C>G, chr16:81942175 A>G, and chr8:61654298 T>A (see Tables 14 and 16).

[0157] In some embodiments, the one or more genetic variations disrupt or modulate one or more of the following genes: PLCG2, POLE, LRBA, EPG5 and SHARPIN (see Table 17).

[0158] In some embodiments, the one or more genetic variations disrupt or modulate one or more of the following genes: PLCG2, CHD7, IFIH1, AP3B1, EPG5, PIK3CD, LRBA and SHARPIN (see Table 18).

[0159] In some embodiments, the gene encodes a transcript with a sequence that has at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to any one of SEQ ID NOs 173-455 or 1500-2177 (see Tables 4 and 12), or complements thereof.

[0160] In some embodiments, the gene encodes a transcript with a sequence that has at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to any one of SEQ ID NOs 173-455 (see Table 4), or complements thereof.

[0161] In some embodiments, the gene encodes a transcript with a sequence that has at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to any one of SEQ ID NOs 1500-2177 (see Table 12), or complements thereof.

[0162] In some embodiments, the one or more genetic variations comprise 2 or 3 or 4 or 5 or more genetic variations.

[0163] In some embodiments, the one or more genetic variations comprise 10 or more genetic variations.

[0164] In some embodiments, the one or more genetic variations comprise 20 or more genetic variations.

[0165] In some embodiments, the one or more genetic variations comprise 50 or more genetic variations.

[0166] In some embodiments, the analyzing comprises microarray analysis, PCR, sequencing, nucleic acid hybridization, or any combination thereof.

[0167] In some embodiments, the genetic test result comprises a genetic test result from a microarray analysis, PCR, sequencing, nucleic acid hybridization, or any combination thereof.

[0168] In some embodiments, the detecting comprises a microarray analysis, PCR, sequencing, nucleic acid hybridization, or any combination thereof.

[0169] In some embodiments, the microarray analysis selected from the group consisting of a Comparative Genomic Hybridization (CGH) array analysis and an SNP array analysis.

[0170] In some embodiments, the sequencing is selected from the group consisting of Massively Parallel Signature Sequencing (MPSS), polony sequencing, 454 pyrosequencing, Illumina sequencing, Illumina (Solexa) sequencing using 10x Genomics library preparation, SOLiD sequencing, ion semiconductor sequencing, DNA nanoball sequencing, heliscope single molecule sequencing, single molecule real time (SMRT) sequencing, RNAP sequencing, Nanopore DNA sequencing, sequencing by hybridization, and microfluidic Sanger sequencing.

[0171] In some embodiments, the analyzing comprises analyzing a whole genome or a whole exome of the subject.

[0172] In some embodiments, the analyzing comprises analyzing nucleic acid information that has already been obtained for a whole genome or a whole exome of the subject.

[0173] In some embodiments, the nucleic acid information is obtained from an in silico analysis.

[0174] In some embodiments, the analyzing comprises analyzing a whole genome or a whole exome of the subject.

[0175] In some embodiments, the analyzing comprises analyzing nucleic acid information that has already been obtained for a whole genome or a whole exome of the subject.

[0176] In some embodiments, the nucleic acid information is obtained from an in silico analysis.

[0177] In some embodiments, the detecting comprises analyzing a whole genome or a whole exome of the subject.

[0178] In some embodiments, the detecting comprises analyzing nucleic acid information that has already been obtained for a whole genome or a whole exome of the subject.

[0179] In some embodiments, the nucleic acid information is obtained from an in silico analysis.

[0180] In some embodiments, the subject is a human subject.

[0181] In some embodiments, the polynucleic acid sample comprises a polynucleic acid from blood, saliva, urine, serum, tears, skin, tissue, or hair of the subject.

[0182] In some embodiments, the method further comprises analyzing for a presence of JCV in a biological sample from the subject.

[0183] In some embodiments, the analyzing for a presence of JCV comprises contacting a JCV detection reagent to the biological sample.

[0184] In some embodiments, the JCV detection reagent is selected from the group consisting of an anti-JCV antibody, a JCV specific primer, and combinations thereof.

[0185] Provided herein is a kit, comprising reagents for assaying a polynucleic acid sample from a subject in need

thereof for the presence of one or more genetic variations that disrupt or modulate a gene of GN1-GN490.

[0186] In some embodiments, the reagents comprise at least one contiguous oligonucleotide that hybridizes to a fragment of the polynucleic acid sample.

[0187] In some embodiments, the reagents comprise at least one pair of oligonucleotides that hybridize to opposite strands of a fragment of the polynucleic acid sample.

[0188] In some embodiments, the kit further comprises one or more immunosuppressive medications.

[0189] In some embodiments, the one or more immunosuppressive medications comprise a glucocorticoid, cytostatic, antibody, drug acting on immunophilins, interferon, opioid, TNF binding protein, mycophenolate, small biological agent, or any combination thereof.

[0190] In some embodiments, the one or more immunosuppressive medications comprise a interferon beta-1a, interferon beta-1b, glatiramer acetate, peginterferon beta-1a, teriflunomide, fingolimod, dimethyl fumarate, alemtuzumab, mitoxantrone, natalizumab, daclizumab, ocrelizumab, or any combination thereof.

[0191] In some embodiments, the one or more immunosuppressive medications comprise natalizumab (Tysabri).

[0192] In some embodiments, the kit further comprises a JCV detection reagent.

[0193] In some embodiments, the JCV detection reagent is selected from the group consisting of an anti-JCV antibody, a JCV specific primer, and combinations thereof.

[0194] In some embodiments, the kit further comprises a set of instructions for administration of the one or more immunosuppressive medications.

[0195] In some embodiments, the one or more genetic variations comprise a point mutation, polymorphism, single nucleotide polymorphisms (SNP), single nucleotide variation (SNV), translocation, insertion, deletion, amplification, inversion, interstitial deletion, copy number variation (CNV), loss of heterozygosity, or any combination thereof.

[0196] In some embodiments, the one or more genetic variations result in a loss of function of the corresponding gene.

[0197] In some embodiments, the one or more genetic variations comprise 5 or more genetic variations.

[0198] In some embodiments, the one or more genetic variations comprise 10 or more genetic variations.

[0199] In some embodiments, the one or more genetic variations comprise 20 or more genetic variations.

[0200] In some embodiments, the one or more genetic variations comprise 50 or more genetic variations.

[0201] In some embodiments, the subject is a human subject.

[0202] In some embodiments, the polynucleic acid sample comprises a polynucleic acid from blood, saliva, urine, serum, tears, skin, tissue, or hair of the subject.

[0203] Provided herein is a panel of polynucleic acids for detecting one or more genetic variations that disrupt or modulate a gene of GN1-GN490, wherein each polynucleic acid of the panel comprises a sequence complementary to a sequence of one or more genetic variation or complements thereof that disrupts or modulates a gene selected from the group consisting of GN1-GN490.

[0204] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a sequence with at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to

SEQ ID NOs 1-172 or SRN1-SRN363, with 100% sequence identity to SEQ ID NOs 1000-1329, or with at least 80% and less than 100% sequence identity to GN1-GN490, or complements thereof.

[0205] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a CNV with at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to SEQ ID NOs 1-172, or complements thereof.

[0206] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a CNV sub-region (SRN) with at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to SRN1-SRN363, or complements thereof.

[0207] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a single nucleotide variation (SNV) with a sequence of any one of SEQ ID NOs: 1000-1329, or complements thereof.

[0208] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a sequence with at least 80% and less than 100% sequence identity to GN1-GN490, or complements thereof.

[0209] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a single nucleotide variation (SNV) with a sequence of any one of SEQ ID NO: 1000, 1001, 1002, 1009, 1010, 1011, 1012, 1014, 1016, 1017, 1019, 1020, 1028, 1032, 1033, 1034, 1035, 1036, 1037, 1040, 1041, 1043, 1051, 1054, 1056, 1057, 1058, 1059, 1061, 1062, 1063, 1066, 1068, 1069, 1070, 1071, 1073, 1074, 1075, 1076, 1077, 1078, 1080, 1082, 1084, 1090, 1092, 1098, 1099, 1100, 1101, 1104, 1107, 1114, 1116, 1118, 1121, 1122, 1123, 1125, 1126, 1127, 1128, 1129, 1130, 1131, 1133, 1135, 1136, 1137, 1138, 1142, 1146, 1147, 1148, 1150, 1152, 1154, 1157, 1160, 1161, 1165, 1166, 1167, 1168, 1169, 1171, 1174, 1175, 1176, 1177, 1178, 1179, 1180, 1181, 1182, 1183, 1184, 1193, 1194, 1200, 1201, 1202, 1203, 1204, 1208, 1219, 1220, 1221, 1222, 1226, 1227, 1228, 1229, 1230, 1231, 1232, 1235, 1239, 1247, 1248, 1249, 1250, 1251, 1252, 1254, 1255, 1256, 1259, 1260, 1261, 1263, 1264, 1266, 1267, 1273, 1278, 1279, 1283, 1284, 1286, 1287, 1289, 1290, 1291, 1299, 1300, 1301, 1304, 1311, 1327 or 1328 (see Tables 7 and 8), or complements thereof.

[0210] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a single nucleotide variation (SNV) with a sequence of any one of SEQ ID NO: 1011, 1020, 1028, 1032, 1034, 1035, 1036, 1040, 1056, 1069, 1073, 1077, 1101, 1114, 1123, 1125, 1126, 1127, 1135, 1142, 1146, 1147, 1148, 1152, 1154, 1157, 1167, 1174, 1184, 1193, 1194, 1203, 1208, 1221, 1222, 1229, 1235, 1252, 1255, 1256, 1259, 1260, 1261, 1263, 1273, 1278, 1279, 1284, 1287, 1289, 1299 or 1311 (see Table 7), or complements thereof.

[0211] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a single nucleotide variation (SNV) with a sequence of any one of SEQ ID NO: 1000, 1001, 1002, 1009, 1010, 1012, 1014, 1016, 1017, 1019, 1033, 1037, 1041, 1043, 1051, 1054, 1057, 1058, 1059, 1061, 1062, 1063, 1066, 1068, 1070, 1071, 1074, 1075, 1076, 1078, 1080, 1082, 1084, 1090, 1092, 1098, 1099, 1100, 1104, 1107, 1116, 1118, 1121, 1122, 1128, 1129, 1130, 1131, 1133, 1136, 1137, 1138, 1146, 1147, 1150, 1152, 1160, 1161, 1165, 1166, 1168, 1169, 1171, 1175, 1176, 1177, 1178, 1179, 1180, 1181, 1182, 1183, 1200, 1201,

1202, 1204, 1219, 1220, 1226, 1227, 1228, 1230, 1231, 1232, 1239, 1247, 1248, 1249, 1250, 1251, 1252, 1254, 1264, 1266, 1267, 1278, 1279, 1283, 1286, 1290, 1291, 1300, 1301, 1304, 1327 or 1328 (see Table 8), or complements thereof.

[0212] In some embodiments, the one or more genetic variations comprise a genetic variation selected from the group consisting of chr16:81942175 A>G, chr2:163136505 C>G, chr11:67818269 G>A, chr22:23917192 G>T, chr20:3846397 C>T, chr8:145154222, G>A chr8:61654298 T>A, chr3:39323163 A>C, chr4:151199080 G>A, chr1:42047208 C>G, chr2:163124051 C>T, chr1:182554557 C>T, chr8:145154824 A>C, chr20:62305450 C>T, chr22:23915745 G>A, chr6:83884161 C>G, chr11:108202772 G>T, chr5:138856923 C>T, chr16:1510535 C>T, chr20:3843027 C>A, chr12:122064788 G>GT, chr16:7714909 C>T, chr18:56401523 C>T, chr1:92946625 G>C, chr5:169081453 G>C, chr11:108117787 C>T, chr22:21235389 A>G, chr19:4817657 C>T, chr10:1060218 G>A, chr21:30698953 T>G, chr9:304628 G>A, chr19:7712287 G>C, chr10:90771767 G>A, chr3:121415370 T>C, chr16:70503095 A>G, chr1:206945738 C>T, chr5:156593120 C>T, chr4:27019452 C>T, chr1:155317682 C>T, chr17:77926526 C>T, chr1:235840495 G>T, chr14:21993359 G>A, chr8:61757805 C>T, chr15:91306241 G>A, chr16:50741791 C>T, chr22:23915583 T>C, chr2:47205921 C>T, chr12:88900891 C>A, chr3:142281353 C>G, chr11:108123551 C>T, chr1:207641950 C>T, chr6:143092151 T>C, chr2:24431184 C>T, chr2:24432937 C>T, chr9:312134 G>A, chr8:100205255 G>A, chr21:16339852 T>C, and any combination thereof (see Tables 14 and 15).

[0213] In some embodiments, the one or more genetic variations comprise a genetic variation selected from the group consisting of chr16:81942175 A>G, chr2:163136505 C>G, chr11:67818269 G>A, chr22:23917192 G>T, chr20:3846397 C>T, chr8:145154222, G>A chr8:61654298 T>A, chr3:39323163 A>C, chr4:151199080 G>A, chr1:42047208 C>G, chr2:163124051 C>T, chr1:182554557 C>T, and any combination thereof (see Table 14).

[0214] In some embodiments, the one or more genetic variations comprise a genetic variation selected from the group consisting of chr8:145154824 A>C, chr20:62305450 C>T, chr22:23915745 G>A, chr6:83884161 C>G, chr11:108202772 G>T, chr5:138856923 C>T, chr16:1510535 C>T, chr20:3843027 C>A, chr12:122064788 G>GT, chr16:7714909 C>T, chr18:56401523 C>T, chr1:92946625 G>C, chr5:169081453 G>C, chr11:108117787 C>T, chr22:21235389 A>G, chr19:4817657 C>T, chr10:1060218 G>A, chr21:30698953 T>G, chr9:304628 G>A, chr19:7712287 G>C, chr10:90771767 G>A, chr3:121415370 T>C, chr16:70503095 A>G, chr1:206945738 C>T, chr5:156593120 C>T, chr4:27019452 C>T, chr1:155317682 C>T, chr17:77926526 C>T, chr1:235840495 G>T, chr14:21993359 G>A, chr8:61757805 C>T, chr15:91306241 G>A, chr16:50741791 C>T, chr22:23915583 T>C, chr2:47205921 C>T, chr12:88900891 C>A, chr3:142281353 C>G, chr11:108123551 C>T, chr1:207641950 C>T, chr6:143092151 T>C, chr2:24431184 C>T, chr2:24432937 C>T, chr9:312134 G>A, chr8:100205255 G>A, chr21:16339852 T>C, and any combination thereof (see Table 15).

[0215] In some embodiments, the SNV is a heterozygous SNV.

[0216] In some embodiments, the SNV is a homozygous SNV.

[0217] In some embodiments, the one or more genetic variations comprise a pair of single nucleotide variations (SNVs), wherein the pair of SNVs are encoded by any one of SEQ ID NO pairs: 1003 and 1004, 1003 and 1005, 1006 and 1007, 1024 and 1025, 1030 and 1031, 1047 and 1048, 1049 and 1050, 1063 and 1064, 1063 and 1065, 1063 and 1066, 1075 and 1076, 1091 and 1093, 1091 and 1096, 1093 and 1095, 1094 and 1097, 1098 and 1099, 1098 and 1100, 1099 and 1100, 1102 and 1103, 1104 and 1106, 1104 and 1107, 1104 and 1108, 1104 and 1109, 1104 and 1110, 1104 and 1111, 1104 and 1112, 1110 and 1111, 1112 and 1113, 1119 and 1120, 1124 and 1125, 1124 and 1126, 1125 and 1126, 1140 and 1141, 1142 and 1144, 1146 and 1151, 1147 and 1148, 1147 and 1149, 1153 and 1146, 1153 and 1147, 1155 and 1156, 1160 and 1161, 1165 and 1166, 1186 and 1187, 1188 and 1193, 1189 and 1193, 1191 and 1192, 1191 and 1193, 1191 and 1195, 1192 and 1193, 1192 and 1195, 1196 and 1197, 1206 and 1207, 1210 and 1218, 1211 and 1213, 1212 and 1213, 1213 and 1215, 1213 and 1216, 1213 and 1217, 1233 and 1238, 1242 and 1243, 1245 and 1246, 1263 and 1260, 1269 and 1279, 1270 and 1279, 1270 and 1282, 1271 and 1279, 1274 and 1279, 1278 and 1279, 1278 and 1281, 1279 and 1280, 1279 and 1281, 1279 and 1282, 1292 and 1293, 1296 and 1297, 1305 and 1314, 1306 and 1310, 1313 and 1321 or 1315 and 1322 (see Table 9 or Tables 9 and 7 for a subset), or complements thereof.

[0218] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a CNV with at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to any one of SEQ ID NOs 157, 2, 140, 65, 26, 14 or 45 (see Tables 7 and 8), or complements thereof.

[0219] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a CNV with at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to any one of SEQ ID NOs 2, 140, 65, 26, 14 or 45 (see Table 7), or complements thereof.

[0220] In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a CNV with at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to SEQ ID NO 157 (see Table 8), or a complement thereof.

[0221] In some embodiments, the one or more genetic variations comprise a CNV and a single nucleotide variations (SNV), wherein SNVs is encoded by any one of SEQ ID NOs 1301, 1173, 1107, 1104, 1199, 1225, 1086 or 1223 (see Table 10), or complements thereof.

[0222] In some embodiments, the one or more genetic variations comprise a genetic variation selected from the group consisting of one or more of the following: chr8:145154222 G>A, chr2:163136505 C>G, chr16:81942175 A>G, and chr8:61654298 T>A (see Tables 14 and 16).

[0223] In some embodiments, the one or more genetic variations disrupt or modulate one or more of the following genes: PLCG2, POLE, LRBA, EPG5 and SHARPIN (see Table 17).

[0224] In some embodiments, the one or more genetic variations disrupt or modulate one or more of the following genes: PLCG2, CHD7, IFIH1, AP3B1, EPG5, PIK3CD, LRBA and SHARPIN (see Table 18).

[0225] In some embodiments, the gene encodes a transcript with a sequence that has at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99%

sequence identity to any one of SEQ ID NOs 173-455 or 1500-2177 (see Tables 4 and 12), or complements thereof.

[0226] In some embodiments, the gene encodes a transcript with a sequence that has at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to any one of SEQ ID NOs 173-455 (see Table 4), or complements thereof.

[0227] In some embodiments, the gene encodes a transcript with a sequence that has at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence identity to any one of SEQ ID NOs 1500-2177 (see Table 12), or complements thereof.

[0228] In some embodiments, the one or more genetic variations comprise at least 5, at least 10, at least 20, or at least 50 genetic variations.

[0229] In some embodiments, panel of polynucleic acids comprises at least 5, at least 10, at least 20, or at least 50 polynucleic acids.

[0230] In some embodiments, the gene comprises a gene selected from the group consisting of gene numbers (GNs) 1-156 (in Table 3).

[0231] In some embodiments, the gene comprises a gene selected from the group consisting of gene numbers (GNs) in Table 6.

[0232] In some embodiments, the gene comprises a gene selected from the group consisting of PLCG2, RBCK1, EPG5, IL17F, SHARPIN, PRF1, JAGN1, TAP1, POLE, LRBA, EHF, IL12B, ATL2, NHEJ1, LYST, HIVEP1, AP3B1, TNFRSF10A, PIK3CD, PNP, MCEE, DOCK2 and ALG12 (see Table 13).

[0233] Provided herein is a method to predict an adverse responsiveness of a subject to a therapy, the method comprising detecting one or more genetic variations that disrupt or modulate a gene of GN1-GN490 in a polynucleic acid sample from the subject; and using that detection as a biomarker for predicting a response of the subject to the therapy to be adverse, wherein the therapy is an immunosuppressive therapy.

[0234] Provided herein is a method of screening for a PML biomarker comprising obtaining biological samples from subjects with PML; screening the biological samples to obtain nucleic acid information; detecting one or more genetic variations that disrupt or modulate a gene of GN1-GN490 in a polynucleic acid sample from a subject suspected of having PML; and using that detection as a biomarker for predicting a response of the subject to the therapy to be adverse, wherein the therapy is an immunosuppressive therapy.

[0235] Provided herein is a method of screening for a PML biomarker comprising obtaining biological samples from subjects with PML; screening the biological samples to obtain nucleic acid information; confirming each biological sample is not a duplicate of any other biological sample based on the nucleic acid information; detecting one or more genetic variations that disrupt or modulate a gene of GN1-GN490 in a polynucleic acid sample from a subject suspected of having PML; and using that detection as a biomarker for predicting a response of the subject to the therapy to be adverse, wherein the therapy is an immunosuppressive therapy.

[0236] Provided herein is a method of screening for a PML biomarker comprising obtaining biological samples from subjects with PML; screening the biological samples to obtain nucleic acid information; determining a sex genotype

for each biological sample based on the nucleic acid information; confirming the sex genotype of each sample is the same as a sex phenotype of the subject from the subjects with PML; detecting one or more genetic variations that disrupt or modulate a gene of GN1-GN490 in a polynucleic acid sample from a subject suspected of having PML; and using that detection as a biomarker for predicting a response of the subject to the therapy to be adverse, wherein the therapy is an immunosuppressive therapy.

[0237] Provided herein is a method of treating a condition in a subject in need of natalizumab therapy, comprising: administering a therapeutically effective amount of natalizumab to the subject, wherein the subject has a decreased risk of progressive multifocal leukoencephalopathy (PML) due to an infection of the brain by John Cunningham virus (JCV), wherein the subject's decreased risk is due to the absence of one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6.

[0238] In some embodiments, the subject is identified as not having one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6. In some embodiments, the subject is known as not having one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6. In some embodiments, the subject is identified in a report (e.g., health report) as not having one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6.

[0239] In some embodiments, the condition is multiple sclerosis or Crohn's disease. In some embodiments, the condition is a relapsing form of multiple sclerosis. In some embodiments, the natalizumab is administered via intravenous infusion.

[0240] In some embodiments, about 100 mg to about 500 mg of the natalizumab is administered. In some embodiments, about 100 mg to about 500 mg of the natalizumab is administered, for example, about 100 mg to about 200 mg, about 100 mg to about 300 mg, about 100 mg to about 400 mg, about 100 mg to about 500 mg, about 200 mg to about 300 mg, about 200 mg to about 400 mg, about 200 mg to about 500 mg, about 300 mg to about 400 mg, about 300 mg to about 500 mg, or about 400 mg to about 500 mg of the natalizumab is administered. In some embodiments, about 100 mg to about 500 mg of the natalizumab is administered via intravenous infusion. In some embodiments, about 100 mg to about 500 mg of the natalizumab is administered via intravenous infusion in four weeks. In some embodiments, about 300 mg of the natalizumab is administered. In some embodiments, about 300 mg of the natalizumab is administered via intravenous infusion. In some embodiments, about 300 mg of the natalizumab is administered via intravenous infusion in four weeks.

[0241] In some embodiments, the one or more genetic variations are associated with a risk of developing PML in a polynucleic acid sample from the subject. In some embodiments, the one or more genetic variations comprises a first genetic variation and a second genetic variation, wherein the first genetic variation disrupts or modulates a corresponding gene according to Tables 3 and 6, and wherein the second genetic variation disrupts or modulates a corresponding gene according to Tables 25A, 25B, and 26.

[0242] In some embodiments, the method comprises testing the subject for a genetic predisposition for PML with a genetic assay. In some embodiments, the genetic assay has

a diagnostic yield of at least 5%. In some cases, the genetic assay has a diagnostic yield of at least about 5%, 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99%. In some cases, the genetic assay has a diagnostic yield of about 1%-5%, 1%-10%, 1%-20%, 5%-10%, 5%-20%, 10%-20%, 10%-30%, 20%-30%, 20%-40%, 30%-40%, 30%-50%, 40%-50%, 40%-60%, 50%-60%, 50%-70%, 60%-70%, 60%-80%, 70%-80%, 70%-90%, 80%-90%, 80%-95%, 90%-95%, 90%-99%, 90%-100%, 95%-99%, or 99%-100%. In some embodiments, the genetic assay has a diagnostic yield of at least 20%.

[0243] In some embodiments, the one or more genetic variations disrupt or modulate a corresponding gene according to Tables 13-18. In some embodiments, the one or more genetic variations disrupt or modulate a corresponding gene according to Tables 19-24.

[0244] In some embodiments, the subject's decreased risk is further due to the absence of one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 25A, 25B, and 26.

[0245] In some embodiments, the one or more genetic variations disrupt or modulate a corresponding gene selected from the group consisting of *Homo sapiens* chromodomain helicase DNA binding protein 7 (CHD7), *Homo sapiens* interferon induced with helicase C domain 1 (IFIH1), *Homo sapiens* immunoglobulin lambda like polypeptide 1 (IGLL1), *Homo sapiens* mitochondrial antiviral signaling protein (MAVS), *Homo sapiens* phospholipase C gamma 2 (PLCG2), *Homo sapiens* SHANK-associated RH domain interactor (SHARPIN), *Homo sapiens* T-cell immune regulator 1, ATPase H⁺ transporting V0 subunit a3 (TCIRG1), and any combination thereof. In some embodiments, the one or more genetic variations comprise chr8:61654298 T>A, chr2:163136505 C>G, chr22:23917192 G>T, chr20:3846397 C>T, chr16:81942175 A>G, chr8:145154222 G>A, chr11:67818269 G>A, chr8:145154824 A>C, chr22:23915745 G>A, chr20:3843027 C>A, or any combination thereof.

[0246] In some embodiments, the corresponding gene comprises a gene selected from the group consisting of gene numbers (GNs) GN1-GN490. In some embodiments, the corresponding gene comprises a gene selected from the group consisting of gene numbers (GNs) GN1-GN241, GN243-GN369, and GN371-GN490.

[0247] In some embodiments, the one or more genetic variations are encoded by a sequence with at least 60% sequence identity to SEQ ID NOs 1-172 or SRN1-SRN363, with 100% sequence identity to SEQ ID NOs 1000-1329, or with at least 80% and less than 100% sequence identity to GN1-GN490, or complements thereof. In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a CNV with at least 60% sequence identity to SEQ ID NOs 1-172, or complements thereof. In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a CNV sub-region (SRN) with at least 60% sequence identity to SRN1-SRN363, or complements thereof. In some embodiments, the one or more genetic variations comprise a genetic variation encoded by a single nucleotide variation (SNV) with a sequence of any one of SEQ ID NOs: 1000-1329, or complements thereof. In some embodiments, the one or more genetic variations are encoded by a sequence with at least 40% sequence identity to SEQ ID NOs 1-172 or

SRN1-SRN363, for example, at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to SEQ ID NOs 1-172 or SRN1-SRN363, or complements thereof. In some embodiments, the one or more genetic variations are encoded by a sequence with at least 40% sequence identity to SEQ ID NOs 1000-1329, for example, at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to SEQ ID NOs 1000-1329, or complements thereof. In some embodiments, the one or more genetic variations are encoded by a sequence with at least 40% and less than 100% sequence identity to GN1-GN490, for example, at least 40% and less than 50%, at least 50% and less than 60%, at least 60% and less than 70%, at least 70% and less than 80%, at least 80% and less than 90%, or at least 90% and less than 100% sequence identity to GN1-GN490, or complements thereof.

[0248] In some embodiments, the genetic assay comprises microarray analysis, PCR, sequencing, nucleic acid hybridization, or any combination thereof.

[0249] In some embodiments, the method comprises testing the subject with a JCV-antibody test, a CD62L test, or a CSF IgM oligoclonal bands test. In some embodiments, the method comprises testing the subject with the JCV-antibody test, wherein the JCV-antibody test does not detect a presence of JCV. In some embodiments, the method comprises testing the subject with the JCV-antibody test, wherein the JCV-antibody test detects a presence of JCV. In some embodiments, the JCV-antibody test comprises contacting a JCV detection reagent to a biological sample from the subject. In some embodiments, the JCV detection reagent is selected from the group consisting of an anti-JCV antibody, a JCV specific primer, and combinations thereof.

[0250] In some embodiments, the subject is identified as not having one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6.

[0251] Provided herein is a kit, comprising reagents for assaying a polynucleic acid sample from a subject in need thereof for the presence of one or more genetic variations that disrupt or modulate a gene of GN1-GN490. In some embodiments, the one or more genetic variations that disrupt or modulate a gene of GN1-GN241, GN243-GN369, and GN371-GN490.

[0252] Provided herein is a method of treating multiple sclerosis or Crohn's disease comprising: (a) testing a subject with multiple sclerosis or Crohn's disease for a genetic predisposition for PML with a genetic assay, wherein the genetic assay has a diagnostic yield of at least 20%, and (b) administering a therapeutically effective amount of natalizumab to the subject, wherein the testing does not identify the subject as having the genetic predisposition for PML.

[0253] In some embodiments, the method further comprises testing the subject with a JCV-antibody test. In some embodiments, the JCV-antibody test does not detect a presence of JCV. In some embodiments, the JCV-antibody test detects a presence of JCV. In some embodiments, the genetic assay tests the subject for the presence of one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6.

[0254] Provided herein is a method of identifying a subject as not having a risk of developing PML, comprising: (a) analyzing a polynucleic acid sample from the subject for one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6, wherein a

genetic variation of the one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6 is not present in the polynucleic acid sample; and (b) identifying the subject as not having a risk of developing PML.

DETAILED DESCRIPTION OF THE DISCLOSURE

[0255] The details of one or more inventive embodiments are set forth in the accompanying drawings, the claims, and in the description herein. Other features, objects, and advantages of inventive embodiments disclosed and contemplated herein will be apparent from the description and drawings, and from the claims. As used herein, unless otherwise indicated, the article “a” means one or more unless explicitly otherwise provided for. As used herein, unless otherwise indicated, terms such as “contain,” “containing,” “include,” “including,” and the like mean “comprising.” As used herein, unless otherwise indicated, the term “or” can be conjunctive or disjunctive. As used herein, unless otherwise indicated, any embodiment can be combined with any other embodiment. As used herein, unless otherwise indicated, some inventive embodiments herein contemplate numerical ranges. When ranges are present, the ranges include the range endpoints. Additionally, every subrange and value within the range is present as if explicitly written out. The term “about” and its grammatical equivalents in relation to a reference numerical value and its grammatical equivalents as used herein can include a range of values plus or minus 10% from that value, such as a range of values plus or minus 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, or 1% from that value. For example, the amount “about 10” includes amounts from 9 to 11. Progressive Multifocal Leukoencephalopathy (PML)

[0256] Progressive multifocal leukoencephalopathy (PML) is a rare and usually fatal viral disease characterized by progressive damage or inflammation of the white matter of the brain at multiple locations. The cause of PML can be a type of polyomavirus called the John Cunningham (JC) virus (or JCV), which can be harmless except in cases of weakened immune systems. While JCV is present at very high rates in the general population, PML remains a rare disorder, albeit an important one because of the clinical sequelae.

[0257] PML can occur in patients with severe immune deficiency, which allows reactivation of the JC virus, such as: 1) most commonly among patients with acquired immune deficiency syndrome (AIDS) that results from infection with human immunodeficiency virus (HIV), 2) patients on immunosuppressive medications like corticosteroids for organ transplant (e.g., renal, liver, lung, and heart) and in people with cancer (e.g., Hodgkin’s disease, leukemia, or lymphoma, and myeloproliferative neoplasms such as myelofibrosis), and 3) individuals with autoimmune diseases (e.g., multiple sclerosis, rheumatoid arthritis, psoriasis, and systemic lupus erythematosus) with therapies that depress the immune response. Several immunosuppressive drugs have been reported in the context of drug-induced PML or drug-associated PML. For example, see: Melis et al. *CNS Drugs*. 2015; 29(10):879-91; Maas et al. *J Neurol*. 2016 October; 263(10):2004-21; Colin et al. *Fundam Clin Pharmacol*. 2016 Oct. 13. Immunosuppressive medications can include, but are not limited to, interferon beta-1a, interferon beta-1b, glatiramer acetate, peginterferon beta-1a,

teriflunomide, mitoxantrone, ocrelizumab, abatacept, adalimumab, alefacept, alemtuzumab, anakinra, bortezomib (e.g., Velcade), ecuzumab (e.g., Soliris), leflunomide, and various other transplant drugs such as antithymocyte Ig, asparaginase, azathioprine, basiliximab, belatacept, belimumab, bendamustine, bevacizumab, bleomycin, brentuximab vedotin, busulfan, capecitabine, carboplatin, carmustine, certolizumab, cetuximab, chlorambucil, cladribine, corticosteroids, cyclophosphamide, cyclosporine, cytotoxic chemotherapy, daclizumab, dimethyl fumarate, doxorubicin, efalizumab, etanercept, etoposide, fingolimod, fludarabine, gemcitabine, hydrocortisone, ibritumomab tiuxetan, ibrutinib, idarubicin, idelalisib, ifosfamide, imatinib, infliximab, lenalidomide, levamisole, mercaptopurine, methotrexate, mizoribine, muromonab-CD3, mycophenolate mofetil, mycophenolic acid, natalizumab, obinutuzumab, ofatumumab, oxaliplatin, prednisone/prednisolone, rituximab, ruxolitinib, sirolimus (also known as rapamycin), tacrolimus, tegafur/gimeracil/oteracil, thiotepa, tocilizumab, tofacitinib, ustekinumab, vedolizumab, vinblastine and vincristine. Exemplary small molecule immunosuppressive medications include dimethyl fumarate, fingolimod, and ruxolitinib. In some embodiments, an immunosuppressive therapy is classified as a Class 1 (high risk) therapeutic agent, such as efalizumab and natalizumab as reported in Calabrese L. H. et al., *Nat Rev Rheumatol*. (2015).

[0258] PML can be diagnosed in a patient with a progressive course of the disease, finding JC virus DNA in spinal fluid together with consistent white matter lesions on brain magnetic resonance imaging (MRI); alternatively, a brain biopsy can be diagnostic when the typical histopathology of demyelination, bizarre astrocytes, and enlarged oligodendroglial nuclei are present, coupled with techniques showing the presence of JC virus. Characteristic evidence of PML on brain CT scan images can be multifocal, non-contrast enhancing hypodense lesions without mass effect, but MRI can be more sensitive than CT. The most common area of involvement can be the cortical white matter of frontal and parieto-occipital lobes, but lesions may occur anywhere in the brain, like the basal ganglia, external capsule, and posterior cranial fossa structures like the brainstem and cerebellum.

[0259] In general, treatment of PML aims at reversing the immune deficiency to slow or stop the disease progress. Patients on an immunosuppression regime can stop taking the immunosuppressive medication or plasma exchange (PLEX) can be used to accelerate the removal of the immunosuppressive medication that put the person at risk for PML. HIV-infected patients can start highly active antiretroviral therapy (HAART). Occurrence of PML can also occur in the context of immune reconstitution inflammatory syndrome (IRIS), wherein onset of PML can occur or PML symptoms may get worse after cessation of immunosuppression (e.g., as reviewed by Pavlovic et al. *Ther Adv Neurol Disord*. 2015 November; 8(6):255-73 and Bowen et al. *Nat Rev Neurol*. 2016 Oct. 27; 12(11):662-674). For example, in MS patients that develop PML during treatment with natalizumab, IRIS often results when treatment is stopped and PLEX is used to remove natalizumab from the patient’s circulation. Treatment of IRIS in PML patients can include administration of corticosteroids. Other potential treatments of PML can include cidofovir, cytarabine, anti-malaria drug mefloquine, interleukin-2, and 1-O-hexadecyloxypropyl-cidofovir (CMX001, aka brincidofovir). As

reviewed by Pavlovic (Ther Adv Neurol Disord. 2015 November; 8(6):255-73), potential treatments for PML include antiviral agents (e.g., chlorpromazine, citalopram, mirtazapine, risperidone, ziprasidone, retro-2cycl, brefeldin A, cidofovir, brincidofovir, cytarabine, ganciclovir, leflunomide, topotecan, mefloquine, 3-aminobenzamide, imatinib, and Ag122), immune response modulators (e.g., IFN-alpha, IL-2, IL-7, maraviroc, and glucocorticoids), and immunization (e.g., recombinant human anti-JCV VP-1 monoclonal antibodies, JCV-specific cytotoxic T lymphocyte therapy, IL-7 plus JCV VP1 vaccine, and JCV oral vaccine).

[0260] The term “diagnostic yield” as used herein refers to the percentage of cases that would identify the presence of one or more genetic variations (e.g., CNV, SNV) in a PML cohort using an assay. For example, if 40 cases would identify the presence of one or more genetic variations (e.g., CNV, SNV) in a cohort of 100 PML patients, the diagnostic yield of the assay is 40%. In some cases, the patients in the PML cohort are clinically diagnosed with PML. In some cases, a patient is clinically diagnosed with PML when JC virus DNA is present in spinal fluid and consistent white matter lesions is present on brain magnetic resonance imaging (MRI). In some cases, a patient is clinically diagnosed with PML when typical histopathology of demyelination, bizarre astrocytes, and enlarged oligodendroglial nuclei are present in a brain biopsy, coupled with the presence of JC virus. In some cases, the PML cohort has at least 5 PML cases, for example, at least 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, or 1000 PML cases. In some cases, the PML cohort is a cohort listed herein. For example, the PML cohort is the PML patient cohort listed in Table 7. In some cases, the assay is JCV-antibody assay. In some cases, the assay is not JCV-antibody assay. In some cases, the assay is a genetic assay. In some cases, the genetic assay tests the genetic predisposition for PML.

[0261] The genetic assay can comprise any method disclosed herein. In some cases, the genetic assay has a diagnostic yield of at least about 5%, 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99%. In some cases, the genetic assay has a diagnostic yield of about 1%-5%, 1%-10%, 1%-20%, 5%-10%, 5%-20%, 10%-20%, 10%-30%, 20%-30%, 20%-40%, 30%-40%, 30%-50%, 40%-50%, 40%-60%, 50%-60%, 50%-70%, 60%-70%, 60%-80%, 70%-80%, 70%-90%, 80%-90%, 80%-95%, 90%-95%, 90%-99%, 90%-100%, 95%-99%, or 99%-100%.

Genetic Variations Associated with PML

[0262] Described herein, are methods that can be used to detect genetic variations. Detecting specific genetic variations, for example polymorphic markers and/or haplotypes, copy number, absence or presence of an allele, or genotype associated with a condition (e.g., disease or disorder) as described herein, can be accomplished by methods known in the art for analyzing nucleic acids and/or detecting sequences at polymorphic or genetically variable sites, for example, amplification techniques, hybridization techniques, sequencing, microarrays/arrays, or any combination thereof. Thus, by use of these methods disclosed herein or other methods available to the person skilled in the art, one or more alleles at polymorphic markers, including microsatellites, single nucleotide polymorphisms (SNPs), single nucleotide variations (SNVs), insertions/deletions (indels),

copy number variations (CNVs), or other types of genetic variations, can be identified in a sample obtained from a subject.

[0263] Genomic sequences within populations exhibit variability between individuals at many locations in the genome. For example, the human genome exhibits sequence variations that occur on average every 500 base pairs. Such genetic variations in polynucleic acid sequences are commonly referred to as polymorphisms or polymorphic sites. As used herein, a polymorphism, e.g., genetic variation, includes a variation in the sequence of the genome amongst a population, such as allelic variations and other variations that arise or are observed. Thus, a polymorphism refers to the occurrence of two or more genetically determined alternative sequences or alleles in a population. These differences can occur in coding (e.g., exonic) and non-coding (e.g., intronic or intergenic) portions of the genome, and can be manifested or detected as differences in polynucleic acid sequences, gene expression, including, for example transcription, processing, translation, transport, protein processing, trafficking, DNA synthesis; expressed proteins, other gene products or products of biochemical pathways or in post-translational modifications and any other differences manifested amongst members of a population. Polymorphisms that arise as the result of a single base change, such as single nucleotide polymorphisms (SNPs) or single nucleotide variations (SNVs), can include an insertion, deletion or change in one nucleotide. A polymorphic marker or site is the locus at which divergence occurs. Such sites can be as small as one base pair (an SNP or SNV). Polymorphic markers include, but are not limited to, restriction fragment length polymorphisms (RFLPs), variable number of tandem repeats (VNTRs), hypervariable regions, minisatellites, dinucleotide repeats, trinucleotide repeats, tetranucleotide repeats and other repeating patterns, simple sequence repeats and insertional elements, such as Alu. Polymorphic forms also are manifested as different mendelian alleles for a gene. Polymorphisms can be observed by differences in proteins, protein modifications, RNA expression modification, DNA and RNA methylation, regulatory factors that alter gene expression and DNA replication, and any other manifestation of alterations in genomic polynucleic acid or organelle polynucleic acids. Those skilled in the art can appreciate that polymorphisms are sometimes considered to be a subclass of variations, defined on the basis of a particular frequency cutoff in a population. For example, in some embodiments, polymorphisms are considered to genetic variants/variations that occur at >1%, or >5%, frequency in the population.

[0264] In some embodiments, these genetic variations can be found to be associated with one or more disorders and/or diseases using the methods disclosed herein. In some embodiments, these genetic variations can be found to be associated with absence of one or more disorders and/or diseases (i.e. the one or more variants are protective against development of the disorder and/or diseases) using the methods disclosed herein.

[0265] In some embodiments, these genetic variations comprise point mutations, polymorphisms, single nucleotide polymorphisms (SNPs), single nucleotide variations (SNVs), translocations, insertions, deletions, amplifications, inversions, interstitial deletions, copy number variations (CNVs), loss of heterozygosity, or any combination thereof. As genetic variation includes any deletion, insertion or base

substitution of the genomic DNA of one or more individuals in a first portion of a total population which thereby results in a difference at the site of the deletion, insertion or base substitution relative to one or more individuals in a second portion of the total population. Thus, the term “genetic variation” encompasses “wild type” or the most frequently occurring variation, and also includes “mutant,” or the less frequently occurring variation. In some embodiments, a wild type allele may be referred to as an ancestral allele.

[0266] As used herein, a target molecule that is “associated with” or “correlates with” a particular genetic variation is a molecule that can be functionally distinguished in its structure, activity, concentration, compartmentalization, degradation, secretion, and the like, as a result of such genetic variation. In some embodiments polymorphisms (e.g., polymorphic markers, genetic variations, or genetic variants) can comprise any nucleotide position at which two or more sequences are possible in a subject population. In some embodiments, each version of a nucleotide sequence, with respect to the polymorphism/variation, can represent a specific allele of the polymorphism/variation. In some embodiments, genomic DNA from a subject can contain two alleles for any given polymorphic marker, representative of each copy of the marker on each chromosome. In some embodiments, an allele can be a nucleotide sequence of a given location on a chromosome. Polymorphisms/variations can comprise any number of specific alleles. In some embodiments of the disclosure, a polymorphism/variation can be characterized by the presence of two or more alleles in a population. In some embodiments, the polymorphism/variation can be characterized by the presence of three or more alleles. In some embodiments, the polymorphism/variation can be characterized by four or more alleles, five or more alleles, six or more alleles, seven or more alleles, nine or more alleles, or ten or more alleles. In some embodiments an allele can be associated with one or more diseases or disorders, for example, a PML risk allele can be an allele that is associated with increased or decreased risk of developing PML. In some embodiments, genetic variations and alleles can be used to associate an inherited phenotype with a responsible genotype. In some embodiments, a PML risk allele can be a variant allele that is statistically associated with a screening of PML. In some embodiments, genetic variations can be of any measurable frequency in the population, for example, a frequency higher than 10%, a frequency from 5-10%, a frequency from 1-5%, a frequency from 0.1-1%, or a frequency below 0.1%. As used herein, variant alleles can be alleles that differ from a reference allele. As used herein, a variant can be a segment of DNA that differs from the reference DNA, such as a genetic variation. In some embodiments, genetic variations can be used to track the inheritance of a gene that has not yet been identified, but whose approximate location is known.

[0267] As used herein, a “haplotype” can be information regarding the presence or absence of one or more genetic markers in a given chromosomal region in a subject. In some embodiments, a haplotype can be a segment of DNA characterized by one or more alleles arranged along the segment, for example, a haplotype can comprise one member of the pair of alleles for each genetic variation or locus. In some embodiments, the haplotype can comprise two or more alleles, three or more alleles, four or more alleles, five or

more alleles, or any combination thereof, wherein, each allele can comprise one or more genetic variations along the segment.

[0268] In some embodiments, a genetic variation can be a functional aberration that can alter gene function, gene expression, polypeptide expression, polypeptide function, or any combination thereof. In some embodiments, a genetic variation can be a loss-of-function mutation, gain-of-function mutation, dominant negative mutation, or reversion. In some embodiments, a genetic variation can be part of a gene’s coding region or regulatory region. Regulatory regions can control gene expression and thus polypeptide expression. In some embodiments, a regulatory region can be a segment of DNA wherein regulatory polypeptides, for example, transcription or splicing factors, can bind. In some embodiments a regulatory region can be positioned near the gene being regulated, for example, positions upstream or downstream of the gene being regulated. In some embodiments, a regulatory region (e.g., enhancer element) can be several thousands of base pairs upstream or downstream of a gene.

[0269] In some embodiments, variants can include changes that affect a polypeptide, such as a change in expression level, sequence, function, localization, binding partners, or any combination thereof. In some embodiments, a genetic variation can be a frameshift mutation, nonsense mutation, missense mutation, neutral mutation, or silent mutation. For example, sequence differences, when compared to a reference nucleotide sequence, can include the insertion or deletion of a single nucleotide, or of more than one nucleotide, resulting in a frame shift; the change of at least one nucleotide, resulting in a change in the encoded amino acid; the change of at least one nucleotide, resulting in the generation of a premature stop codon; the deletion of several nucleotides, resulting in a deletion of one or more amino acids encoded by the nucleotides; the insertion of one or several nucleotides, such as by unequal recombination or gene conversion, resulting in an interruption of the coding sequence of a reading frame; duplication of all or a part of a sequence; transposition; or a rearrangement of a nucleotide sequence. Such sequence changes can alter the polypeptide encoded by the nucleic acid, for example, if the change in the nucleic acid sequence causes a frame shift, the frame shift can result in a change in the encoded amino acids, and/or can result in the generation of a premature stop codon, causing generation of a truncated polypeptide. In some embodiments, a genetic variation associated with PML can be a synonymous change in one or more nucleotides, for example, a change that does not result in a change in the amino acid sequence. Such a polymorphism can, for example, alter splice sites, affect the stability or transport of mRNA, or otherwise affect the transcription or translation of an encoded polypeptide. In some embodiments, a synonymous mutation can result in the polypeptide product having an altered structure due to rare codon usage that impacts polypeptide folding during translation, which in some cases may alter its function and/or drug binding properties if it is a drug target. In some embodiments, the changes that can alter DNA increase the possibility that structural changes, such as amplifications or deletions, occur at the somatic level. A polypeptide encoded by the reference nucleotide sequence can be a reference polypeptide with a particular reference amino acid sequence, and polypeptides encoded

by variant nucleotide sequences can be variant polypeptides with variant amino acid sequences.

[0270] The most common sequence variants comprise base variations at a single base position in the genome, and such sequence variants, or polymorphisms, are commonly called single nucleotide polymorphisms (SNPs) or single nucleotide variants (SNVs). In some embodiments, a SNP represents a genetic variant present at greater than or equal to 1% occurrence in a population and in some embodiments a SNP or an SNV can represent a genetic variant present at any frequency level in a population. A SNP can be a nucleotide sequence variation occurring when a single nucleotide at a location in the genome differs between members of a species or between paired chromosomes in a subject. SNPs can include variants of a single nucleotide, for example, at a given nucleotide position, some subjects can have a 'G', while others can have a 'C'. SNPs can occur in a single mutational event, and therefore there can be two possible alleles possible at each SNP site; the original allele and the mutated allele. SNPs that are found to have two different bases in a single nucleotide position are referred to as biallelic SNPs, those with three are referred to as triallelic, and those with all four bases represented in the population are quadallelic. In some embodiments, SNPs can be considered neutral. In some embodiments SNPs can affect susceptibility to a condition (e.g., PML). SNP polymorphisms can have two alleles, for example, a subject can be homozygous for one allele of the polymorphism wherein both chromosomal copies of the individual have the same nucleotide at the SNP location, or a subject can be heterozygous wherein the two sister chromosomes of the subject contain different nucleotides. The SNP nomenclature as reported herein is the official Reference SNP (rs) ID identification tag as assigned to each unique SNP by the National Center for Biotechnological Information (NCBI).

[0271] Another genetic variation of the disclosure can be copy number variations (CNVs). As used herein, "CNVs" include alterations of the DNA of a genome that results in an abnormal number of copies of one or more sections of DNA. In some embodiments, a CNV comprises a CNV-subregion. As used herein, a "CNV-subregion" includes a continuous nucleotide sequence within a CNV. In some embodiments, the nucleotide sequence of a CNV-subregion can be shorter than the nucleotide sequence of the CNV, and in another embodiment the CNV-subregion can be equivalent to the CNV (e.g., such as for some recurrent CNVs). CNVs can be inherited or caused by de novo mutation and can be responsible for a substantial amount of human phenotypic variability, behavioral traits, and disease susceptibility. In some embodiments, CNVs of the current disclosure can be associated with susceptibility to one or more conditions, for example, PML. In some embodiments, CNVs can include a single gene or include a contiguous set of genes. In some embodiments, CNVs can be caused by structural rearrangements of the genome, for example, unbalanced translocations or inversions, insertions, deletions, amplifications, and interstitial deletions. In some embodiments, these structural rearrangements occur on one or more chromosomes. Low copy repeats (LCRs), which are region-specific repeat sequences (also known as segmental duplications), can be susceptible to these structural rearrangements, resulting in CNVs. Factors such as size, orientation, percentage similarity and the distance between the copies can influence the susceptibility of LCRs to genomic rearrangement. In addition,

rearrangements may be mediated by the presence of high copy number repeats, such as long interspersed elements (LINEs) and short interspersed elements (SINEs), often via non-homologous recombination. For example, chromosomal rearrangements can arise from non-allelic homologous recombination during meiosis or via a replication-based mechanism such as fork stalling and template switching (FoSTeS) (Zhang F. et al., *Nat. Genet.* (2009)) or microhomology-mediated break-induced repair (MMBIR) (Hastings P. J. et al., *PLoS Genetics* (2009)). In some embodiments, CNVs are referred to as structural variants, which are a broader class of variant that also includes copy number neutral alterations such as balanced inversions and balanced translocations.

[0272] CNVs can account for genetic variation affecting a substantial proportion of the human genome, for example, known CNVs can cover over 15% of the human genome sequence (Estivill and Armengol, *PLoS Genetics* (2007)). CNVs can affect gene expression, phenotypic variation and adaptation by disrupting or impairing gene dosage, and can cause disease, for example, microdeletion and microduplication disorders, and can confer susceptibility to diseases and disorders. Updated information about the location, type, and size of known CNVs can be found in one or more databases, for example, the Database of Genomic Variants (See, MacDonald JR et al., *Nucleic Acids Res.*, 42, D986-92 (2014), which currently contains data for over 500,000 CNVs (as of May, 2016).

[0273] Other types of sequence variants can be found in the human genome and can be associated with a disease or disorder, including but not limited to, microsatellites. Microsatellite markers are stable, polymorphic, easily analyzed, and can occur regularly throughout the genome, making them especially suitable for genetic analysis. A polymorphic microsatellite can comprise multiple small repeats of bases, for example, CA repeats, at a particular site wherein the number of repeat lengths varies in a population. In some embodiments, microsatellites, for example, variable number of tandem repeats (VNTRs), can be short segments of DNA that have one or more repeated sequences, for example, about 2 to 5 nucleotides long, that can occur in non-coding DNA. In some embodiments, changes in microsatellites can occur during genetic recombination of sexual reproduction, increasing or decreasing the number of repeats found at an allele, or changing allele length.

[0274] The genetic variations disclosed herein can be associated with a risk of developing PML in a subject. In some cases, the subject can have a decreased risk due to the absence of one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 1 to 26. For example, the subject can have a decreased risk due to the absence of one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6. In some cases, the subject can have an increased risk due to the presence of one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 1 to 26. For example, the subject can have an increased risk due to the presence of one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6. In some cases, one or more genes listed in Tables 25A, 25B, and 26 can be removed from any one of the Tables 1-24. In some cases, one or more genes listed in Tables 25A, 25B, and 26 can be added to any one of the Tables 1-24.

TABLE 25A

exemplary 8-gene panel				
RefSeq Gene Symbol	Disease Model	Gene Source	Source Annotation	Gene Number (GN)
BAG3	AR	Public_db	PMID: 19229298, 19282432, 22984599, 27042682	175
BTK	XLR	Public_db	PMID: 18281276, 23765059, 25930993, 26029204	180
CD40LG	XLR	Public_db	PMID: 17360404, 21455173, 23765059, 26008899, 26029204	206
DOCK8	AR	Public_db	PMID: 23765059, 23887241, 26029204, 26454313	242
MAGT1	XLR	Public_db	PMID: 23887241, 25504528, 27873163	326
RAG1	AD_AR	Public_db	PMID: 23122631, 23765059, 23887241, 25976673, 26029204, 26454313, 27484032, 27808398	370
STAT1	AD_AR	Public_db	PMID: 23887241, 25645939, 26029204, 26513235, 26743090, 27821552, 27873163	436
WAS	XLR	Both	PMID: 12874226, 14647476, 19782549, 20008220, 24753205, 26029204, 26371186	483

TABLE 25B

exemplary 16-gene panel				
RefSeq Gene Symbol	Disease Model	Gene Source	Source Annotation	Gene Number (GN)
ADA	AR	Both	PMID: 23765059, 24135998, 25930993, 26029204, 26454313	1
BAG3	AR	Public_db	PMID: 19229298, 19282432, 22984599, 27042682	175
BTK	XLR	Public_db	PMID: 18281276, 23765059, 25930993, 26029204	180
CD40LG	XLR	Public_db	PMID: 14647476, 17360404, 21455173, 23765059, 26008899, 26029204	206
DNMT3B	AR	Public_db	PMID: 23486536, 23765059, 26029204, 26851945	240
DOCK8	AR	Public_db	PMID: 23765059, 23887241, 26029204, 26454313	242
ITK	AR	Public_db	PMID: 14647476, 23765059, 26029204, 26454313	308
LCK	AR	Public_db	PMID: 14647476, 23765059, 26029204, 26454313	316
PNP	AR	Both	PMID: 26029204, 26454313	354
RAG1	AD_AR	Public_db	PMID: 23122631, 23765059, 23887241, 25976673, 26029204, 26454313, 27484032, 27808398	370
STAT1	AD_AR	Public_db	PMID: 23887241, 25645939, 26029204, 26513235, 26743090, 27821552, 27873163	436
STAT3	AD	Public_db	PMID: 23765059, 23887241, 25645939, 25930993, 26029204, 27658964, 27873163	438
STK3	unknown	Both	PMID: 26029204	135
TYK2	AR	Public_db	PMID: 26029204, 26513235, 27821552	144
WAS	XLR	Both	PMID: 12874226, 19782549, 20008220, 24753205, 26029204, 26371186	483
WIPF1	AR	Public_db	PMID: 23765059, 26029204, 26453379	485

TABLE 26

exemplary 2-gene panel				
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary
ADA	intronic	100	adenosine deaminase	This gene encodes an enzyme that catalyzes the hydrolysis of adenosine to inosine. Various mutations have been described for this gene and have been linked to human diseases. Deficiency in this enzyme causes a form of severe combined immunodeficiency disease (SCID), in which there is dysfunction of both B and T lymphocytes with impaired cellular immunity and decreased production of immunoglobulins, whereas elevated levels of this enzyme have been associated with congenital hemolytic anemia. [provided by RefSeq, July 2008]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript

TABLE 26-continued

exemplary 2-gene panel					
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary	Gene # (GN)
STK3	intronic	6788	serine/ threonine- protein kinase 3 isoform 1	exon combination :: BC040226.1, X02994.1 [ECO:0000332] RNAseq introns :: mixed/partial sample support ERS025081, ERS025082 [ECO:0000350] ##Evidence-Data-END## This gene encodes a serine/threonine protein kinase activated by proapoptotic molecules indicating the encoded protein functions as a growth suppressor. Cleavage of the protein product by caspase removes the inhibitory C-terminal portion. The N-terminal portion is transported to the nucleus where it homodimerizes to form the active kinase which promotes the condensation of chromatin during apoptosis. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, January 2012]. Transcript Variant: This variant (1) encodes isoform 1. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: U26424.1, BC010640.2 [ECO:0000332] RNAseq introns :: single sample supports all introns ERS025084, ERS025088 [ECO:0000348] ##Evidence-Data-END##	135

[0275] A “subject”, as used herein, can be an individual of any age or sex from whom a sample containing polynucleotides is obtained for analysis by one or more methods described herein so as to obtain polynucleic acid information; for example, a male or female adult, child, newborn, or fetus. In some embodiments, a subject can be any target of therapeutic administration. In some embodiments, a subject can be a test subject or a reference subject.

[0276] As used herein, a “cohort” can represent an ethnic group, a patient group, a particular age group, a group not associated with a particular condition (e.g., disease or disorder), a group associated with a particular condition (e.g., disease or disorder), a group of asymptomatic subjects, a group of symptomatic subjects, or a group or subgroup of subjects associated with a particular response to a treatment regimen or enrolled in a clinical trial. In some embodiments, a patient can be a subject afflicted with a condition (e.g., disease or disorder). In some embodiments, a patient can be a subject not afflicted with a condition (e.g., disease or disorder) and is considered apparently healthy, or a normal or control subject. In some embodiments, a subject can be a test subject, a patient or a candidate for a therapeutic, wherein genomic DNA from the subject, patient, or candidate is obtained for analysis by one or more methods of the present disclosure herein, so as to obtain genetic variation information of the subject, patient or candidate.

[0277] In some embodiments, the polynucleic acid sample can be obtained prenatally from a fetus or embryo or from the mother, for example, from fetal or embryonic cells in the maternal circulation. In some embodiments, the polynucleic acid sample can be obtained with the assistance of a health care provider, for example, to draw blood. In some embodiments, the polynucleic acid sample can be obtained without the assistance of a health care provider, for example, where the polynucleic acid sample is obtained non-invasively, such

as a saliva sample, or a sample comprising buccal cells that is obtained using a buccal swab or brush, or a mouthwash sample.

[0278] The present disclosure also provides methods for assessing genetic variations in subjects who are members of a target population. Such a target population is in some embodiments a population or group of subjects at risk of developing the condition (e.g., disease or disorder), based on, for example, other genetic factors, biomarkers, biophysical parameters, diagnostic testing such as magnetic resonance imaging (MRI), family history of the condition, previous screening or medical history, or any combination thereof.

[0279] The genetic variations of the present disclosure found to be associated with a condition (e.g., disease or disorder) can show similar association in other human populations. Particular embodiments comprising subject human populations are thus also contemplated and within the scope of the disclosure. Such embodiments relate to human subjects that are from one or more human populations including, but not limited to, Caucasian, Ashkenazi Jewish, Sephardi Jewish, European, American, Eurasian, Asian, Central/South Asian, East Asian, Middle Eastern, African, Hispanic, Caribbean, and Oceanic populations. European populations include, but are not limited to, Swedish, Norwegian, Finnish, Russian, Danish, Icelandic, Irish, Kelt, English, Scottish, Dutch, Belgian, French, German, Spanish, Portuguese, Italian, Polish, Bulgarian, Slavic, Serbian, Bosnian, Czech, Greek and Turkish populations. The ethnic contribution in subjects can also be determined by genetic analysis, for example, genetic analysis of ancestry can be carried out using unlinked microsatellite markers or single nucleotide polymorphisms (SNPs) such as those set out in Smith et al., (Smith M. W. et al., Am. J. Hum. Genet., 74:1001 (2004)).

[0280] Certain genetic variations can have different population frequencies in different populations, or are polymor-

phic in one population but not in another. The methods available and as thought herein can be applied to practice the present disclosure in any given human population. This can include assessment of genetic variations of the present disclosure, so as to identify those markers that give strongest association within the specific population. Thus, the at-risk variants of the present disclosure can reside on different haplotype background and in different frequencies in various human populations.

Conditions and Immunosuppressive Medications

[0281] In some embodiments, a subject can be diagnosed or undiagnosed with a condition (e.g., disease or disorder), can be asymptomatic or symptomatic, can have increased or decreased susceptibility to a condition (e.g., disease or disorder), can be currently under or previously under or not under a treatment for a condition (e.g., disease or disorder), or any combination thereof. In some embodiments, the condition can be AIDS, cancer, organ transplant, or an autoimmune disease. In some embodiments, the condition is PML.

[0282] In some embodiments, a subject can be diagnosed or undiagnosed with PML, can be asymptomatic or symptomatic, can have increased or decreased susceptibility to PML, can be currently under or previously under or not under a treatment for PML, or any combination thereof. In some embodiments, a subject can be diagnosed or undiagnosed with AIDS (e.g., individuals infected with HIV), can be asymptomatic or symptomatic, can have increased or decreased susceptibility to AIDS, can be currently under or previously under or not under a treatment for AIDS, or any combination thereof. In some embodiments, a subject can be diagnosed or undiagnosed with cancer (e.g., Hodgkin's disease, leukemia, lymphoma, or myelofibrosis), can be asymptomatic or symptomatic, can have increased or decreased susceptibility to cancer, can be currently under or previously under or not under a treatment for cancer, or any combination thereof. In some embodiments, a subject can be currently diagnosed or previously diagnosed or undiagnosed with an autoimmune disease (e.g., multiple sclerosis, rheumatoid arthritis, psoriasis, systemic lupus erythematosus), can be asymptomatic or symptomatic, can have increased or decreased susceptibility to an autoimmune disease, can be currently under or previously under or not under a treatment for an autoimmune disease, or any combination thereof.

[0283] The term "cancer" is meant to include all types of cancerous growths or oncogenic processes, metastatic tissues or malignantly transformed cells, tissues, or organs, irrespective of histopathologic type or stage of invasiveness. A metastatic tumor can arise from a multitude of primary tumor types, including but not limited to those of breast, lung, liver, colon and ovarian origin. Examples of cancers include, but are not limited to, a fibrosarcoma, myosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, gastric cancer, esophageal cancer, rectal cancer, pancreatic cancer, ovarian cancer, prostate cancer, uterine cancer, cancer of the head and neck, skin cancer, brain cancer, squamous cell carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinoma, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carci-

noma, choriocarcinoma, seminoma, embryonal carcinoma, Wilm's tumor, cervical cancer, testicular cancer, small cell lung carcinoma, non-small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma, retinoblastoma, leukemia, lymphoma, myelofibrosis, or Kaposi sarcoma.

[0284] The term "autoimmune disease" is meant to include all types of pathological states arising from abnormal immune responses of the body to substances and tissues that are normally present in the body. Examples of autoimmune diseases include, but are not limited to, Addison disease, Anti-NMDA receptor encephalitis, antisyntetase syndrome, Aplastic anemia, autoimmune anemias, Autoimmune hemolytic anemia, Autoimmune pancreatitis, Behcet's Disease, bullous skin disorders, Celiac disease—sprue (gluten-sensitive enteropathy), chronic fatigue syndrome, Chronic inflammatory demyelinating polyneuropathy, chronic lymphocytic leukemia, Crohn's disease, Dermatomyositis, Devic's disease, Erythroblastopenia, Evans syndrome, Focal segmental glomerulosclerosis, Granulomatosis with polyangiitis, Graves disease, Graves' ophthalmopathy, Guillain-Barre syndrome, Hashimoto thyroiditis, idiopathic thrombocytopenic purpura (ITP), IgA nephropathy, IgA-mediated autoimmune diseases, IgG4-related disease, Inflammatory bowel disease, Juvenile idiopathic arthritis, Multiple sclerosis, Myasthenia gravis, myeloma, non-Hodgkin's lymphoma, Opsoclonus myoclonus syndrome (OMS), Pemphigoid, Pemphigus, pemphigus vulgaris, Pernicious anemia, polymyositis, Psoriasis, pure red cell aplasia, Reactive arthritis, Rheumatoid arthritis, Sarcoidosis, scleroderma, Sjögren syndrome, Systemic lupus erythematosus, Thrombocytopenic purpura, Thrombotic thrombocytopenic purpura, Type I diabetes, Ulcerative colitis, Vasculitis (e.g., vasculitis associated with anti-neutrophil cytoplasmic antibody) and Vitiligo.

[0285] In some embodiments, a subject can be currently treated with an immunosuppressive medication. In some embodiments, a subject can be previously treated with an immunosuppressive medication. In some embodiments, a subject can be not yet treated with an immunosuppressive medication. The immunosuppressive medication can include but not limited to glucocorticoids, cytostatics, antibodies, drugs acting on immunophilins, interferons, opioids, TNF binding proteins, mycophenolate, or other small biological agents. For example, glucocorticoids can include but not limited to cortisol (hydrocortisone), cortisone, prednisone, prednisolone, methylprednisolone, dexamethasone, betamethasone, triamcinolone, beclometasone, fludrocortisone acetate, deoxycorticosterone acetate (DOCA), or aldosterone. Cytostatics can include but not limited to nitrogen mustards (cyclophosphamide), nitrosoureas, platinum compounds, folic acid analogues such as methotrexate, purine analogues such as azathioprine and mercaptopurine, pyrimidine analogues such as fluorouracil, protein synthesis inhibitors, cytotoxic antibiotics such as dactinomycin, anthracyclines, mitomycin C, bleomycin, or mithramycin. Antibodies can include but not limited to polyclonal antibodies such as atgam and thymoglobuline, monoclonal antibodies such as CD25- and CD3-directed antibodies, muromonab-CD3, basiliximab (Simulect), and daclizumab (Zenapax). Drugs acting on immunophilins can include but

not limited to ciclosporin, tacrolimus, sirolimus, or everolimus. TNF binding proteins can include but not limited to infliximab (Remicade), etanercept (Enbrel), or adalimumab (Humira). Other small biological agents can include but not limited to fingolimod and myriocin.

[0286] In some embodiments, the immunosuppressive medication can be drugs for treating multiple sclerosis include but not limited to interferon beta-1a (e.g., Avonex, Rebif), interferon beta-1b (e.g., Betaseron, Extavia), glatiramer acetate (Copaxone, Glatopa), peginterferon beta-1a (e.g., Plegridy), teriflunomide (Aubagio), fingolimod (Gilenya), dimethyl fumarate (Tecfidera), alemtuzumab (Lemtrada), mitoxantrone (e.g., Novantrone), natalizumab (e.g., Tysabri), daclizumab (e.g., Zinbryta), or ocrelizumab (e.g., Ocrevus).

[0287] In some embodiments, the immunosuppressive medication can be adalimumab (e.g., Humira), alemtuzumab (e.g., Lemtrada), alemtuzumab (e.g., Campath), azathioprine (e.g., Imuran), belimumab (e.g., Benlysta), bevacizumab (e.g., Avastatin), bortezomib (e.g., Velcade), ecilizumab (e.g., Soliris), leflunomide, brentuximab vedotin (e.g., Adcetris), cetuximab (e.g., Erbitux), cyclophosphamid, cimetidyl fumarate (e.g., Tecfidera), efalizumab (e.g., Raptiva), fingolimod (e.g., Gilenya), fludarabine (e.g., Fludara), fumaric acid, imatinib (e.g., Gleevec, Glivec), infliximab (e.g., Remicade), methotrexate (e.g., Trexall, Rheumatrex), mycophenolate mofetil (e.g., Cellcept), natalizumab (e.g., Tysabri), daclizumab (e.g., Zinbryta), rituximab (e.g., Rituxin), vedolizumab (Entyvio), ruxolitinib (e.g., Jakafi, Jakavi), or ocrelizumab (e.g., Ocrevus).

[0288] In some embodiments, a method of treating a condition in a subject in need of natalizumab therapy, comprises administering a therapeutically effective amount of natalizumab to the subject, wherein the subject is identified as not having one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6. In some embodiments, a method of reducing a risk of a subject developing PML comprises administering a therapeutically effective amount of natalizumab to the subject, wherein the subject is identified as not having one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6. In some embodiments, the condition is multiple sclerosis. In some embodiments, the condition is Crohn's disease. In some embodiments, a method of treating multiple sclerosis comprises administering natalizumab to a subject with multiple sclerosis, wherein the subject is identified as not having one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6. In some embodiments, a method of treating Crohn's disease comprises administering natalizumab to a subject with Crohn's disease, wherein the subject is identified as not having one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6. In some embodiments, a method of treating multiple sclerosis comprises testing a subject with multiple sclerosis for the presence of one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6, determining that the subject does not have the one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6, and administering natalizumab to the subject that was determined not to have the one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6. In some embodiments, a

method of treating Crohn's disease comprises testing a subject with Crohn's disease for the presence of one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6, determining that the subject does not have the one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6, and administering natalizumab to the subject that was determined not to have the one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6. In some embodiments, a method of reducing a risk of a subject developing PML comprises testing a subject for the presence of one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6, determining that the subject has at least one of the one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6, and advising against administering natalizumab to the subject that was determined to have at least one of the one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6. In some embodiments, the subject has multiple sclerosis. In some embodiments, the subject has Crohn's disease. In some embodiments, a method of treating multiple sclerosis comprises testing a subject with multiple sclerosis for the presence of one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6, determining that the subject has at least one of the one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6, and advising against administering natalizumab to the subject that was determined to have at least one of the one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6. In some embodiments, a method of treating Crohn's disease comprises testing a subject with Crohn's disease for the presence of one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6, determining that the subject has at least one of the one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6, and advising against administering natalizumab to the subject that was determined to have at least one of the one or more genetic variations that disrupt or modulate a corresponding gene according to Tables 3 and 6. In some embodiments, the advising comprises advising that administering natalizumab is contraindicated. In some embodiments, the advising comprises advising that administering natalizumab increases the risk of the subject developing PML. In some embodiments, the advising comprises advising that administering natalizumab is a factor that increases the risk of the subject developing PML.

Samples

[0289] Samples that are suitable for use in the methods described herein can be polynucleic acid samples from a subject. A "polynucleic acid sample" as used herein can include RNA or DNA, or a combination thereof. In another embodiment, a "polypeptide sample" (e.g., peptides or proteins, or fragments therefrom) can be used to ascertain information that an amino acid change has occurred, which is the result of a genetic variant. Polynucleic acids and polypeptides can be extracted from one or more samples including but not limited to, blood, saliva, urine, mucosal scrapings of the lining of the mouth, expectorant, serum, tears, skin, tissue, or hair. A polynucleic acid sample can be

assayed for polynucleic acid information. “Polynucleic acid information,” as used herein, includes a polynucleic acid sequence itself, the presence/absence of genetic variation in the polynucleic acid sequence, a physical property which varies depending on the polynucleic acid sequence (e.g., T_m), and the amount of the polynucleic acid (e.g., number of mRNA copies). A “polynucleic acid” means any one of DNA, RNA, DNA including artificial nucleotides, or RNA including artificial nucleotides. As used herein, a “purified polynucleic acid” includes cDNAs, fragments of genomic polynucleic acids, polynucleic acids produced using the polymerase chain reaction (PCR), polynucleic acids formed by restriction enzyme treatment of genomic polynucleic acids, recombinant polynucleic acids, and chemically synthesized polynucleic acid molecules. A “recombinant” polynucleic acid molecule includes a polynucleic acid molecule made by an artificial combination of two otherwise separated segments of sequence, e.g., by chemical synthesis or by the manipulation of isolated segments of polynucleic acids by genetic engineering techniques. As used herein, a “polypeptide” includes proteins, fragments of proteins, and peptides, whether isolated from natural sources, produced by recombinant techniques, or chemically synthesized. A polypeptide may have one or more modifications, such as a post-translational modification (e.g., glycosylation, phosphorylation, etc.) or any other modification (e.g., pegylation, etc.). The polypeptide may contain one or more non-naturally-occurring amino acids (e.g., such as an amino acid with a side chain modification).

[0290] In some embodiments, the polynucleic acid sample can comprise cells or tissue, for example, cell lines. Exemplary cell types from which nucleic acids can be obtained using the methods described herein include, but are not limited to, the following: a blood cell such as a B lymphocyte, T lymphocyte, leukocyte, erythrocyte, macrophage, or neutrophil; a muscle cell such as a skeletal cell, smooth muscle cell or cardiac muscle cell; a germ cell, such as a sperm or egg; an epithelial cell; a connective tissue cell, such as an adipocyte, chondrocyte; fibroblast or osteoblast; a neuron; an astrocyte; a stromal cell; an organ specific cell, such as a kidney cell, pancreatic cell, liver cell, or a keratinocyte; a stem cell; or any cell that develops therefrom. A cell from which nucleic acids can be obtained can be a blood cell or a particular type of blood cell including, for example, a hematopoietic stem cell or a cell that arises from a hematopoietic stem cell such as a red blood cell, B lymphocyte, T lymphocyte, natural killer cell, neutrophil, basophil, eosinophil, monocyte, macrophage, or platelet. Generally, any type of stem cell can be used including, without limitation, an embryonic stem cell, adult stem cell, or pluripotent stem cell.

[0291] In some embodiments, a polynucleic acid sample can be processed for RNA or DNA isolation, for example, RNA or DNA in a cell or tissue sample can be separated from other components of the polynucleic acid sample. Cells can be harvested from a polynucleic acid sample using standard techniques, for example, by centrifuging a cell sample and resuspending the pelleted cells, for example, in a buffered solution, for example, phosphate-buffered saline (PBS). In some embodiments, after centrifuging the cell suspension to obtain a cell pellet, the cells can be lysed to extract DNA. In some embodiments, the nucleic acid sample can be concentrated and/or purified to isolate DNA. All nucleic acid samples obtained from a subject, including

those subjected to any sort of further processing, are considered to be obtained from the subject. In some embodiments, standard techniques and kits known in the art can be used to extract RNA or DNA from a nucleic acid sample, including, for example, phenol extraction, a QIAamp® Tissue Kit (Qiagen, Chatsworth, Calif.), a Wizard® Genomic DNA purification kit (Promega), or a Qiagen Autopure method using Puregene chemistry, which can enable purification of highly stable DNA well-suited for archiving.

[0292] In some embodiments, determining the identity of an allele or determining copy number can, but need not, include obtaining a polynucleic acid sample comprising RNA and/or DNA from a subject, and/or assessing the identity, copy number, presence or absence of one or more genetic variations and their chromosomal locations within the genomic DNA (i.e. subject’s genome) derived from the polynucleic acid sample.

[0293] The individual or organization that performs the determination need not actually carry out the physical analysis of a nucleic acid sample from a subject. In some embodiments, the methods can include using information obtained by analysis of the polynucleic acid sample by a third party. In some embodiments, the methods can include steps that occur at more than one site. For example, a polynucleic acid sample can be obtained from a subject at a first site, such as at a health care provider or at the subject’s home in the case of a self-testing kit. The polynucleic acid sample can be analyzed at the same or a second site, for example, at a laboratory or other testing facility.

Nucleic Acids

[0294] The nucleic acids and polypeptides described herein can be used in methods and kits of the present disclosure. In some embodiments, aptamers that specifically bind the nucleic acids and polypeptides described herein can be used in methods and kits of the present disclosure. As used herein, a nucleic acid can comprise a deoxyribonucleotide (DNA) or ribonucleotide (RNA), whether singular or in polymers, naturally occurring or non-naturally occurring, double-stranded or single-stranded, coding, for example a translated gene, or non-coding, for example a regulatory region, or any fragments, derivatives, mimetics or complements thereof. In some embodiments, nucleic acids can comprise oligonucleotides, nucleotides, polynucleotides, nucleic acid sequences, genomic sequences, complementary DNA (cDNA), antisense nucleic acids, DNA regions, probes, primers, genes, regulatory regions, introns, exons, open-reading frames, binding sites, target nucleic acids and allele-specific nucleic acids.

[0295] A “probe,” as used herein, includes a nucleic acid fragment for examining a nucleic acid in a specimen using the hybridization reaction based on the complementarity of nucleic acid.

[0296] A “hybrid” as used herein, includes a double strand formed between any one of the abovementioned nucleic acid, within the same type, or across different types, including DNA-DNA, DNA-RNA, RNA-RNA or the like.

[0297] “Isolated” nucleic acids, as used herein, are separated from nucleic acids that normally flank the gene or nucleotide sequence (as in genomic sequences) and/or has been completely or partially purified from other transcribed sequences (e.g., as in an RNA library). For example, isolated nucleic acids of the disclosure can be substantially isolated with respect to the complex cellular milieu in which it

naturally occurs, or culture medium when produced by recombinant techniques, or chemical precursors or other chemicals when chemically synthesized. In some instances, the isolated material can form part of a composition, for example, a crude extract containing other substances, buffer system or reagent mix. In some embodiments, the material can be purified to essential homogeneity using methods known in the art, for example, by polyacrylamide gel electrophoresis (PAGE) or column chromatography (e.g., HPLC). With regard to genomic DNA (gDNA), the term “isolated” also can refer to nucleic acids that are separated from the chromosome with which the genomic DNA is naturally associated. For example, the isolated nucleic acid molecule can contain less than about 250 kb, 200 kb, 150 kb, 100 kb, 75 kb, 50 kb, 25 kb, 10 kb, 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb or 0.1 kb of the nucleotides that flank the nucleic acid molecule in the gDNA of the cell from which the nucleic acid molecule is derived.

[0298] Nucleic acids can be fused to other coding or regulatory sequences can be considered isolated. For example, recombinant DNA contained in a vector is included in the definition of “isolated” as used herein. In some embodiments, isolated nucleic acids can include recombinant DNA molecules in heterologous host cells or heterologous organisms, as well as partially or substantially purified DNA molecules in solution. Isolated nucleic acids also encompass in vivo and in vitro RNA transcripts of the DNA molecules of the present disclosure. An isolated nucleic acid molecule or nucleotide sequence can be synthesized chemically or by recombinant means. Such isolated nucleotide sequences can be useful, for example, in the manufacture of the encoded polypeptide, as probes for isolating homologous sequences (e.g., from other mammalian species), for gene mapping (e.g., by in situ hybridization with chromosomes), or for detecting expression of the gene, in tissue (e.g., human tissue), such as by Northern blot analysis or other hybridization techniques disclosed herein. The disclosure also pertains to nucleic acid sequences that hybridize under high stringency hybridization conditions, such as for selective hybridization, to a nucleotide sequence described herein. Such nucleic acid sequences can be detected and/or isolated by allele- or sequence-specific hybridization (e.g., under high stringency conditions). Stringency conditions and methods for nucleic acid hybridizations are well known to the skilled person (see, e.g., *Current Protocols in Molecular Biology*, Ausubel, F. et al., John Wiley & Sons, (1998), and Kraus, M. and Aaronson, S., *Methods Enzymol.*, 200:546-556 (1991), the entire teachings of which are incorporated by reference herein.

[0299] Calculations of “identity” or “percent identity” between two or more nucleotide or amino acid sequences can be determined by aligning the sequences for optimal comparison purposes (e.g., gaps can be introduced in the sequence of a first sequence). The nucleotides at corresponding positions are then compared, and the percent identity between the two sequences is a function of the number of identical positions shared by the sequences (i.e. $\% \text{ identity} = \frac{\# \text{ of identical positions}}{\text{total } \# \text{ of positions}} \times 100$). For example, a position in the first sequence is occupied by the same nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the

length of each gap, which need to be introduced for optimal alignment of the two sequences.

[0300] In some embodiments, the length of a sequence aligned for comparison purposes is at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, or at least 95%, of the length of the reference sequence. The actual comparison of the two sequences can be accomplished by well-known methods, for example, using a mathematical algorithm. A non-limiting example of such a mathematical algorithm is described in Karlin, S. and Altschul, S., *Proc. Natl. Acad. Sci. USA*, 90:5873-5877 (1993). Such an algorithm is incorporated into the NBLAST and XBLAST programs (version 2.0), as described in Altschul, S. et al., *Nucleic Acids Res.*, 25:3389-3402 (1997). When utilizing BLAST and Gapped BLAST programs, any relevant parameters of the respective programs (e.g., NBLAST) can be used. For example, parameters for sequence comparison can be set at score=100, word length=12, or can be varied (e.g., W=5 or W=20). Other examples include the algorithm of Myers and Miller, CABIOS (1989), ADVANCE, ADAM, BLAT, and FASTA. In some embodiments, the percent identity between two amino acid sequences can be accomplished using, for example, the GAP program in the GCG software package (Accelrys, Cambridge, UK).

[0301] “Probes” or “primers” can be oligonucleotides that hybridize in a base-specific manner to a complementary strand of a nucleic acid molecule. Probes can include primers, which can be a single-stranded oligonucleotide probe that can act as a point of initiation of template-directed DNA synthesis using methods including but not limited to, polymerase chain reaction (PCR) and ligase chain reaction (LCR) for amplification of a target sequence. Oligonucleotides, as described herein, can include segments or fragments of nucleic acid sequences, or their complements. In some embodiments, DNA segments can be between 5 and 10,000 contiguous bases, and can range from 5, 10, 12, 15, 20, or 25 nucleotides to 10, 15, 20, 25, 30, 40, 50, 100, 200, 500, 1000 or 10,000 nucleotides. In addition to DNA and RNA, probes and primers can include polypeptide nucleic acids (PNA), as described in Nielsen, P. et al., *Science* 254: 1497-1500 (1991). A probe or primer can comprise a region of nucleotide sequence that hybridizes to at least about 15, typically about 20-25, and in certain embodiments about 40, 50, 60 or 75, consecutive nucleotides of a nucleic acid molecule.

[0302] The present disclosure also provides isolated nucleic acids, for example, probes or primers, that contain a fragment or portion that can selectively hybridize to a nucleic acid that comprises, or consists of, a nucleotide sequence, wherein the nucleotide sequence can comprise at least one polymorphism or polymorphic allele contained in the genetic variations described herein or the wild-type nucleotide that is located at the same position, or the complements thereof. In some embodiments, the probe or primer can be at least 70% identical, at least 80% identical, at least 85% identical, at least 90% identical, or at least 95% identical, to the contiguous nucleotide sequence or to the complement of the contiguous nucleotide sequence.

[0303] In some embodiments, a nucleic acid probe can be an oligonucleotide capable of hybridizing with a complementary region of a gene associated with a condition (e.g., PML) containing a genetic variation described herein. The

nucleic acid fragments of the disclosure can be used as probes or primers in assays such as those described herein.

[0304] The nucleic acids of the disclosure, such as those described above, can be identified and isolated using standard molecular biology techniques well known to the skilled person. In some embodiments, DNA can be amplified and/or can be labeled (e.g., radiolabeled, fluorescently labeled) and used as a probe for screening, for example, a cDNA library derived from an organism. cDNA can be derived from mRNA and can be contained in a suitable vector. For example, corresponding clones can be isolated, DNA obtained following *in vivo* excision, and the cloned insert can be sequenced in either or both orientations by art-recognized methods to identify the correct reading frame encoding a polypeptide of the appropriate molecular weight. Using these or similar methods, the polypeptide and the DNA encoding the polypeptide can be isolated, sequenced and further characterized.

[0305] In some embodiments, nucleic acid can comprise one or more polymorphisms, variations, or mutations, for example, single nucleotide polymorphisms (SNPs), single nucleotide variations (SNVs), copy number variations (CNVs), for example, insertions, deletions, inversions, and translocations. In some embodiments, nucleic acids can comprise analogs, for example, phosphorothioates, phosphoramidates, methyl phosphonate, chiral methyl phosphonates, 2-O-methyl ribonucleotides, or modified nucleic acids, for example, modified backbone residues or linkages, or nucleic acids combined with carbohydrates, lipids, polypeptide or other materials, or peptide nucleic acids (PNAs), for example, chromatin, ribosomes, and transcriptosomes. In some embodiments nucleic acids can comprise nucleic acids in various structures, for example, A DNA, B DNA, Z-form DNA, siRNA, tRNA, and ribozymes. In some embodiments, the nucleic acid may be naturally or non-naturally polymorphic, for example, having one or more sequence differences, for example, additions, deletions and/or substitutions, as compared to a reference sequence. In some embodiments, a reference sequence can be based on publicly available information, for example, the U.C. Santa Cruz Human Genome Browser Gateway (genome.ucsc.edu/cgi-bin/hgGateway) or the NCBI website (www.ncbi.nlm.nih.gov). In some embodiments, a reference sequence can be determined by a practitioner of the present disclosure using methods well known in the art, for example, by sequencing a reference nucleic acid.

[0306] In some embodiment a probe can hybridize to an allele, SNP, SNV, or CNV as described herein. In some embodiments, the probe can bind to another marker sequence associated with PML as described herein.

[0307] One of skill in the art would know how to design a probe so that sequence specific hybridization can occur only if a particular allele is present in a genomic sequence from a test nucleic acid sample. The disclosure can also be reduced to practice using any convenient genotyping method, including commercially available technologies and methods for genotyping particular genetic variations

[0308] Control probes can also be used, for example, a probe that binds a less variable sequence, for example, a repetitive DNA associated with a centromere of a chromosome, can be used as a control. In some embodiments, probes can be obtained from commercial sources. In some embodiments, probes can be synthesized, for example, chemically or *in vitro*, or made from chromosomal or

genomic DNA through standard techniques. In some embodiments sources of DNA that can be used include genomic DNA, cloned DNA sequences, somatic cell hybrids that contain one, or a part of one, human chromosome along with the normal chromosome complement of the host, and chromosomes purified by flow cytometry or microdissection. The region of interest can be isolated through cloning, or by site-specific amplification using PCR.

[0309] One or more nucleic acids for example, a probe or primer, can also be labeled, for example, by direct labeling, to comprise a detectable label. A detectable label can comprise any label capable of detection by a physical, chemical, or a biological process for example, a radioactive label, such as ^{32}P or ^3H , a fluorescent label, such as FITC, a chromophore label, an affinity-ligand label, an enzyme label, such as alkaline phosphatase, horseradish peroxidase, or β -galactosidase, an enzyme cofactor label, a hapten conjugate label, such as digoxigenin or dinitrophenyl, a Raman signal generating label, a magnetic label, a spin label, an epitope label, such as the FLAG or HA epitope, a luminescent label, a heavy atom label, a nanoparticle label, an electrochemical label, a light scattering label, a spherical shell label, semiconductor nanocrystal label, such as quantum dots (described in U.S. Pat. No. 6,207,392), and probes labeled with any other signal generating label known to those of skill in the art, wherein a label can allow the probe to be visualized with or without a secondary detection molecule. A nucleotide can be directly incorporated into a probe with standard techniques, for example, nick translation, random priming, and PCR labeling. A "signal," as used herein, include a signal suitably detectable and measurable by appropriate means, including fluorescence, radioactivity, chemiluminescence, and the like.

[0310] Non-limiting examples of label moieties useful for detection include, without limitation, suitable enzymes such as horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; members of a binding pair that are capable of forming complexes such as streptavidin/biotin, avidin/biotin or an antigen/antibody complex including, for example, rabbit IgG and anti-rabbit IgG; fluorophores such as umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, tetramethyl rhodamine, eosin, green fluorescent protein, erythrosin, coumarin, methyl coumarin, pyrene, malachite green, stilbene, lucifer yellow, Cascade Blue, Texas Red, dichlorotriazinylamine fluorescein, dansyl chloride, phycoerythrin, fluorescent lanthanide complexes such as those including Europium and Terbium, cyanine dye family members, such as Cy3 and Cy5, molecular beacons and fluorescent derivatives thereof, as well as others known in the art as described, for example, in *Principles of Fluorescence Spectroscopy*, Joseph R. Lakowicz (Editor), Plenum Pub Corp, 2nd edition (July 1999) and the 6th Edition of the *Molecular Probes Handbook* by Richard P. Hoagland; a luminescent material such as luminol; light scattering or plasmon resonant materials such as gold or silver particles or quantum dots; or radioactive material include ^{14}C , ^{123}I , ^{124}I , ^{121}I , $^{99\text{m}}\text{Tc}$, ^{32}P , ^{33}P , ^{31}S or ^3H .

[0311] Other labels can also be used in the methods of the present disclosure, for example, backbone labels. Backbone labels comprise nucleic acid stains that bind nucleic acids in a sequence independent manner. Non-limiting examples include intercalating dyes such as phenanthridines and acridines (e.g., ethidium bromide, propidium iodide, hexidium

iodide, dihydroethidium, ethidium homodimer-1 and -2, ethidium monoazide, and ACMA); some minor groove binders such as indoles and imidazoles (e.g., Hoechst 33258, Hoechst 33342, Hoechst 34580 and DAPI); and miscellaneous nucleic acid stains such as acridine orange (also capable of intercalating), 7-AAD, actinomycin D, LDS751, and hydroxystilbamidine. All of the aforementioned nucleic acid stains are commercially available from suppliers such as Molecular Probes, Inc. Still other examples of nucleic acid stains include the following dyes from Molecular Probes: cyanine dyes such as SYTOX Blue, SYTOX Green, SYTOX Orange, POPO-1, POPO-3, YOYO-1, YOYO-3, TOTO-1, TOTO-3, JOJO-1, LOLO-1, BOBO-1, BOBO-3, PO-PRO-1, PO-PRO-3, BO-PRO-1, BO-PRO-3, TO-PRO-1, TO-PRO-3, TO-PRO-S, JO-PRO-1, LO-PRO-1, YO-PRO-1, YO-PRO-3, PicoGreen, OliGreen, RiboGreen, SYBR Gold, SYBR Green I, SYBR Green II, SYBR DX, SYTO-40, -41, -42, -43, -44, -45 (blue), SYTO-13, -16, -24, -21, -23, -12, -11, -20, -22, -15, -14, -25 (green), SYTO-81, -80, -82, -83, -84, -85 (orange), SYTO-64, -17, -59, -61, -62, -60, -63 (red).

[0312] In some embodiments, fluorophores of different colors can be chosen, for example, 7-amino-4-methylcoumarin-3-acetic acid (AMCA), 5-(and -6)-carboxy-X-rhodamine, lissamine rhodamine B, 5-(and -6)-carboxyfluorescein, fluorescein-5-isothiocyanate (FITC), 7-diethylaminocoumarin-3-carboxylic acid, tetramethylrhodamine-5-(and -6)-isothiocyanate, 5-(and -6)-carboxy-tetramethylrhodamine, 7-hydroxycoumarin-3-carboxylic acid, 6-[fluorescein 5-(and -6)-carboxamido]hexanoic acid, N-(4,4-difluoro-5,7-dimethyl-4-bora-3a,4a-diaza-3-indacenepropanoic acid, eosin-5-isothiocyanate, erythrosin-5-isothiocyanate, TRITC, rhodamine, tetramethylrhodamine, R-phycoerythrin, Cy-3, Cy-5, Cy-7, Texas Red, Phar-Red, allophycocyanin (APC), and CASCADE™ blue acetylazide, such that each probe in or not in a set can be distinctly visualized. In some embodiments, fluorescently labeled probes can be viewed with a fluorescence microscope and an appropriate filter for each fluorophore, or by using dual or triple band-pass filter sets to observe multiple fluorophores.

In some embodiments, techniques such as flow cytometry can be used to examine the hybridization pattern of the probes.

[0313] In other embodiments, the probes can be indirectly labeled, for example, with biotin or digoxigenin, or labeled with radioactive isotopes such as ^{32}P and/or ^3H . As a non-limiting example, a probe indirectly labeled with biotin can be detected by avidin conjugated to a detectable marker. For example, avidin can be conjugated to an enzymatic marker such as alkaline phosphatase or horseradish peroxidase. In some embodiments, enzymatic markers can be detected using colorimetric reactions using a substrate and/or a catalyst for the enzyme. In some embodiments, catalysts for alkaline phosphatase can be used, for example, 5-bromo-4-chloro-3-indolylphosphate and nitro blue tetrazolium. In some embodiments, a catalyst can be used for horseradish peroxidase, for example, diaminobenzoate.

[0314] One or more genes disclosed herein can be in conditions or molecular pathways related to various aspects of immune function including, but not limited to, Type I interferon response (e.g., PMID 26052098), B cell receptor pathway (e.g., Wikipathways WP23; PMID 22566564), RANKL/RANK signaling pathway (e.g., Wikipathways WP2 b18), TCR signaling pathway (e.g., Wikipathways WP69), NF- κ B signaling (e.g., PMID 28362430), JAK-STAT pathway (e.g., PMID 28255960), post-translational modification biology such as ubiquitination via LUBAC (e.g., PMID 23104095, 24958845, 25086647, 26085218, 26111062, 26525107, 26848516, 26877205, 27178468, 27786304, 27892465), Aicardi-Goutieres syndrome (e.g., PMID 26052098), eosinophilia (e.g., PMID 27222657), congenital neutropenia (e.g., PMID 24753205), T cell receptor defects (e.g., PMID 25452106, 25636200, 26246585, 26379669, 26453379, 28400082), and autophagy defects (e.g., 19229298, 22984599, 23222957, 26917586, 26953272, 27588602). In some embodiments, one or more genes disclosed herein can be related to JC virus biology (e.g., PMID 15327898, 19282432, 19903823, 22984599, 25910481). In some embodiments, one or more genes disclosed herein can be antibiral immune response genes.

TABLE 27

Exemplary pathways and biology for PML risk genes (96-gene panel)*							
Genes	Autoinflammatory disease	Autophagy defects	B cell defects	B cell receptor pathway	Deubiquitinase pathway	Eosinophilia-associated immuno deficiency	JC virus biology
AP3B1							
APOL1							27042682
ASH1L							
ATM			23765059			27222657	19903823, 25910481, 27042682
ATR							19903823, 25910481
BLM							
CARD11			23765059	23765059, WP23		27222657	
CDKN1B							
CHD7						27222657	
CLCN7							
DCLRE1C							
DDX58							
DOCK8			23765059			27222657	
EGR1							

TABLE 27-continued

Exemplary pathways and biology for PML risk genes (96-gene panel)*					
EPG5	23222957, 26917586				
ETF1					
FPR2					
GATA2		23765059			
GFI1					
HIVEP1					
HIVEP2					
HTR2A					27042682
IDO2					
IFIH1					
IFNGR2					
IFNLR1					
IGLL1		23765059	23765059		
IKBKB	28469620	23765059	23765059, WP23	26877205, 28362430	
IL17F					
IL1B				27892465	15327898
IL21R		23765059			
IRAK4		23765059	23765059	28362430	
ITSN2					
JUN			WP23		27042682
KAT6B					
KCTD7					
LIG4					27222657
LRBA	26707784	23765059			
MALL					
MAPK3					27042682
MAVS					
MCEE					
MKL1					
MYD88		23765059		28362430	
NBN					
NFKB1	28469620	23765059	WP23	26877205, 28362430	
NOD2	28421071	26953272		28362430	27222657
NRIP1					
PIAS1					
PIAS2					
PIK3CD					
PIK3CD-AS1					
PK3R1		23765059	WP23		
PKHD1					
PLCG2		23765059	WP23		
PNPT1					
POLA1					
POLE		23765059			
PRF1					
PRKCB			WP23		
PRKCD			WP23		
PRKCH					
PRKDC					
PSTPIP1	28421071				
PTEN					
PTPRC			WP23		
RABGEF1					
RAD51					27042682
RAG1		23765059			27222657
RAG2		23765059			27222657
RIPK1	28469620			26877205, 27892465, 28362430	
RIPK3				26877205, 27892465, 28362430	
RNF168					
RTEL1					
SHARPIN	28469620	23765059		26877205, 27892465, 28362430	
SKIV2L					
SMAD4					27042682
STIM1		23765059			
STIM2					
STXBP2					

TABLE 27-continued

Exemplary pathways and biology for PML risk genes (96-gene panel)*							
TAP2							
TBK1							
TCIRG1							
TICAM1							28362430
TLR3	28469620						28362430
TLR4	28469620						28362430
TNFRSF11A	28421071						28362430
TNFRSF13B			23765059				
TNFRSF8							
TP53							
TRAF3							
TRAFIG1							
TRPM2							
VPS45							
WEE1							27042682
ZAP70			23765059			27222657	
Genes	Osteopetrosis	PI3K signaling	RANKL/ RANK pathway	T cell defects	T cell receptor pathway	TLR signaling	Type I interferon pathway
AP3B1	19782549, 24753205						
APOL1							
ASH1L							
ATM							25692705
ATR							24799566
BLM							
CARD11					WP69	25930993	
CDKN1B							
CHD7							
CLCN7	23877423, 24753205						
DCLRE1C							
DDX58							25145756, 26052098, 26763980, 27821552
DOCK8				26379669			
EGR1							
EPG5							
ETF1							
FPR2							
GATA2							
GF11	19782549, 24753205						
HIVEP1							
HIVEP2							
HTR2A							
IDO2							
IFIH1							26052098, 27821552
IFNGR2							
IFNLR1							
IGLL1							
IKBKB	21079651		WP2018		WP69	20404851, 25930993	17047224, 25145756
IL17F							
IL1B					WP69	25930993	25145756, 26763980
IL21R							
IRAK4						20404851, 25930993	25737587
ITSN2							
JUN							25888367
KAT6B							
KCTD7							
LIG4							
LRBA				26707784			
MALL							

TABLE 27-continued

Exemplary pathways and biology for PML risk genes (96-gene panel)*									
MAPK3			WP2018		WP69	25930993			
MAVS								19120474, 22626058, 22901541, 25145756, 26763980	
MCEE									
MKL1								26098208, 26098211	
MYD88						20404851, 25930993		16474425, 18573338, 25145756	
NBN									
NFKB1	21079651	27616589	WP2018		WP69	20404851, 25930993		17047224, 25145756, 26763980 26763980	
NOD2									
NRIP1									
PIAS1								24036127	
PIAS2								21156324, 24036127	
PIK3CD		27616589						20231019	
PIK3CD-AS1		27616589							
PK3R1		27616589	WP2018		WP69			26196376	
PKHD1									
PLCG2		27616589	WP2018	26379669	WP69	25930993			
PNPT1									
POLA1								27019227, 27821552	
POLE									
PRF1									
PRKCB									
PRKCD					WP69			11839738	
PRKCH									
PRKDC								23251783	
PSTPIP1									
PTEN		27616589							
PTPRC		27616589						19673688, 25869642	
RABGEF1									
RAD51									
RAG1		27616589							
RAG2		27616589							
RIPK1						20404851		25145756	
RIPK3								25145756	
RNF168									
RTEL1									
SHARPIN			25930993			20404851		22901541, 25145756	
SKIV2L								25064072, 27821552	
SMAD4									
STIM1								22144678	
STIM2									
STXBP2									
TAP2									
TBK1						25930993		18573338, 22626058, 25145756, 26763980, 28049150	
TCIRG1	23877423, 24753205								
TICAM1						20404851, 25930993		19120474, 25145756, 28049150	
TLR3						20404851, 25930993		19120474, 25145756, 28049150	
TLR4						20404851, 25930993		25145756	

TABLE 27-continued

Exemplary pathways and biology for PML risk genes (96-gene panel)*				
TNFRSF11A	21079651, 23877423	WP2018	25930993	21527253, 25407789
TNFRSF13B			25930993	
TNFRSF8				
TP53				
TRAF3	21079651	WP2018	25930993	22901541, 25723057, 26763980
TRAFD1	25992615		16221674	18849341
TRPM2				
VPS45				
WEE1				
ZAP70	27616589	WP69		

*PMID numbers are listed for curated PubMed references or Wikipathway ID number

[0315] Table 27 contains an exemplary pathways and biology for PML risk genes based on the 96-gene panel listed in Table 19. The genes disclosed herein, such as the genes in the 96-gene panel, can be grouped based on the pathway or biological processes they are involved in.

Methods of Screening

[0316] As used herein, screening a subject comprises diagnosing or determining, theranosing, or determining the susceptibility to developing (prognosing) a condition, for example, PML. In particular embodiments, the disclosure is a method of determining a presence of, or a susceptibility to, PML, by detecting at least one genetic variation in a sample from a subject as described herein. In some embodiments, detection of particular alleles, markers, variations, or haplotypes is indicative of a presence or susceptibility to a condition (e.g., PML).

[0317] While means for screening PML using a JCV antibody test exist, PML risk is not adequately assessed by the JCV antibody test alone. Thus there exists a need for an improved screening test for assessing the risk of developing PML. Described herein are methods of screening an individual for a risk of developing PML, including but not limited to, determining the identity and location of genetic variations, such as variations in nucleotide sequence and copy number, and the presence or absence of alleles or genotypes in one or more samples from one or more subjects using any of the methods described herein. In some embodiments, determining an association to having or developing PML can be performed by detecting particular variations that appear more frequently in test subjects compared to reference subjects and analyzing the molecular and physiological pathways these variations can affect.

[0318] Within any given population, there can be an absolute susceptibility of developing a disease or trait, defined as the chance of a person developing the specific disease or trait over a specified time-period. Susceptibility (e.g., being at-risk) is typically measured by looking at very large numbers of people, rather than at a particular individual. As described herein, certain copy number variations (genetic variations) and/or single nucleotide variations are found to be useful for susceptibility assessment of PML. Susceptibility assessment can involve detecting particular genetic variations in the genome of individuals undergoing assessment. Particular genetic variations are found more frequently in individuals with PML, than in individuals without PML. Therefore, these genetic variations have pre-

dictive value for detecting PML, or a susceptibility to PML, in an individual. Without intending to be limited by theory, it is believed that the genetic variations described herein to be associated with susceptibility of PML represent functional variants predisposing to the disease. In some embodiments, a genetic variation can confer a susceptibility of the condition, for example carriers of the genetic variation are at a different risk of the condition than non-carriers. In some embodiments, the presence of a genetic variation is indicative of increased susceptibility to PML.

[0319] In some embodiments, screening can be performed using any of the methods disclosed, alone or in combination. In some embodiments, screening can be performed using Polymerase Chain Reaction (PCR). In some embodiments screening can be performed using Array Comparative Genomic Hybridization (aCGH) to detect CNVs. In another preferred embodiment screening can be performed using exome sequencing to detect SNVs, indels, and in some cases CNVs using appropriate analysis algorithms. In another preferred embodiment screening is performed using high-throughput (also known as next generation) whole genome sequencing methods and appropriate algorithms to detect all or nearly all genetic variations present in a genomic DNA sample. In some embodiments, the genetic variation information as it relates to the current disclosure can be used in conjunction with any of the above mentioned symptomatic screening tests to screen a subject for PML, for example, using a combination of aCGH and/or sequencing with a JCV screening test, such as the JCV antibody test, CD62L test, or CSF IgM oligoclonal band test. In some embodiments, the L-selectin (CD62L) expressed by CD3+CD4+T cells in, for example, cryopreserved peripheral blood mononuclear cells (PBMCs), can be a biomarker for JCV screening. A CD62L expression can be correlated with the risk of PML.

[0320] In some embodiments, information from any of the above screening methods (e.g., specific symptoms, scoring matrix, or genetic variation data) can be used to define a subject as a test subject or reference subject. In some embodiments, information from any of the above screening methods can be used to associate a subject with a test or reference population, for example, a subject in a population.

[0321] In one embodiment, an association with PML can be determined by the statistical likelihood of the presence of a genetic variation in a subject with PML, for example, an unrelated individual or a first or second-degree relation of the subject. In some embodiments, an association with PML can be decided by determining the statistical likelihood of

the absence of a genetic variation in an unaffected reference subject, for example, an unrelated individual or a first or second-degree relation of the subject. The methods described herein can include obtaining and analyzing a nucleic acid sample from one or more suitable reference subjects.

[0322] In the present context, the term screening comprises diagnosis, prognosis, and theragnosis. Screening can refer to any available screening method, including those mentioned herein. As used herein, susceptibility can be proneness of a subject towards the development of PML, or towards being less able to resist PML than one or more control subjects. In some embodiments, susceptibility can encompass increased susceptibility. For example, particular nucleic acid variations of the disclosure as described herein can be characteristic of increased susceptibility to PML. In some embodiments, particular nucleic acid variations can confer decreased susceptibility, for example particular nucleic variations of the disclosure as described herein can be characteristic of decreased susceptibility to development of PML.

[0323] As described herein, a genetic variation predictive of susceptibility to or presence of PML can be one where the particular genetic variation is more frequently present in a group of subjects with the condition (affected), compared to the frequency of its presence in a reference group (control), such that the presence of the genetic variation is indicative of susceptibility to or presence of PML. In some embodiments, the reference group can be a population nucleic acid sample, for example, a random nucleic acid sample from the general population or a mixture of two or more nucleic acid samples from a population. In some embodiments, disease-free controls can be characterized by the absence of one or more specific disease-associated symptoms, for example, individuals who have not experienced symptoms associated with PML. In some embodiments, the disease-free control group is characterized by the absence of one or more disease-specific risk factors, for example, at least one genetic and/or environmental risk factor. In some embodiments, a reference sequence can be referred to for a particular site of genetic variation. In some embodiments, a reference allele can be a wild-type allele and can be chosen as either the first sequenced allele or as the allele from a control individual. In some embodiments, one or more reference subjects can be characteristically matched with one or more affected subjects, for example, with matched aged, gender or ethnicity.

[0324] A person skilled in the art can appreciate that for genetic variations with two or more alleles present in the population being studied, and wherein one allele can be found in increased frequency in a group of individuals with PML in the population, compared with controls, the other allele of the marker can be found in decreased frequency in the group of individuals with the trait or disease, compared with controls. In such a case, one allele of the marker, for example, the allele found in increased frequency in individuals with PML, can be the at-risk allele, while the other allele(s) can be a neutral or protective allele.

[0325] A genetic variant associated with PML can be used to predict the susceptibility of the disease for a given genotype. For any genetic variation, there can be one or more possible genotypes, for example, homozygote for the at-risk variant (e.g., in autosomal recessive disorders), heterozygote, and non-carrier of the at-risk variant. Autosomal

recessive disorders can also result from two distinct genetic variants impacting the same gene such that the individual is a compound heterozygote (e.g., the maternal allele contains a different mutation than the paternal allele). Compound heterozygosity may result from two different SNVs, two different CNVs, an SNV and a CNV, or any combination of two different genetic variants but each present on a different allele for the gene. For X-linked genes, males who possess one copy of a variant-containing gene may be affected, while carrier females, who also possess a wild-type gene, may remain unaffected. In some embodiments, susceptibility associated with variants at multiple loci can be used to estimate overall susceptibility. For multiple genetic variants, there can be k ($k=3^n \cdot 2^p$) possible genotypes; wherein n can be the number of autosomal loci and p can be the number of gonosomal (sex chromosomal) loci. Overall susceptibility assessment calculations can assume that the relative susceptibilities of different genetic variants multiply, for example, the overall susceptibility associated with a particular genotype combination can be the product of the susceptibility values for the genotype at each locus. If the susceptibility presented is the relative susceptibility for a person, or a specific genotype for a person, compared to a reference population, then the combined susceptibility can be the product of the locus specific susceptibility values and can correspond to an overall susceptibility estimate compared with a population. If the susceptibility for a person is based on a comparison to non-carriers of the at-risk allele, then the combined susceptibility can correspond to an estimate that compares the person with a given combination of genotypes at all loci to a group of individuals who do not carry at-risk variants at any of those loci. The group of non-carriers of any at-risk variant can have the lowest estimated susceptibility and can have a combined susceptibility, compared with itself, for example, non-carriers, of 1.0, but can have an overall susceptibility, compared with the population, of less than 1.0.

[0326] Overall risk for multiple risk variants can be performed using standard methodology. Genetic variations described herein can form the basis of risk analysis that combines other genetic variations known to increase risk of PML, or other genetic risk variants for PML. In certain embodiments of the disclosure, a plurality of variants (genetic variations, variant alleles, and/or haplotypes) can be used for overall risk assessment. These variants are in some embodiments selected from the genetic variations as disclosed herein. Other embodiments include the use of the variants of the present disclosure in combination with other variants known to be useful for screening a susceptibility to PML. In such embodiments, the genotype status of a plurality of genetic variations, markers and/or haplotypes is determined in an individual, and the status of the individual compared with the population frequency of the associated variants, or the frequency of the variants in clinically healthy subjects, such as age-matched and sex-matched subjects.

[0327] Methods such as the use of available algorithms and software can be used to identify, or call, significant genetic variations, including but not limited to, algorithms of DNA Analytics or DNACopy, iPattern and/or QuantiSNP. In some embodiments, a threshold log ratio value can be used to determine losses and gains. For example, using DNA Analytics, a $\log_2$ ratio cutoff of >0.5 and <0.5 to classify CNV gains and losses respectively can be used. For example, using DNA Analytics, a $\log_2$ ratio cutoff of >0.25

and <0.25 to classify CNV gains and losses respectively can be used. As a further example, using DNACopy, a $\log_2$ ratio cutoff of >0.35 and <0.35 to classify CNV gains and losses respectively can be used. For example, an Aberration Detection Module 2 (ADM2) algorithm, such as that of DNA Analytics 4.0.85 can be used to identify, or call, significant genetic variations. In some embodiments, two or more algorithms can be used to identify, or call, significant genetic variations. For example, 2, 3, 4, 5, 6, 7, 8, 9, or 10 or more algorithms can be used to identify, or call, significant genetic variations. In another embodiment, the $\log_2$ ratio of one or more individual probes on a microarray can be used to identify significant genetic variations, such as the presence of homozygously deleted regions in a subject's genome. In some embodiments, significant genetic variations can be CNVs.

[0328] CNVs detected by two or more algorithms can be defined as stringent and can be utilized for further analyses. In some embodiments, the information and calls from two or more of the methods described herein can be compared to each other to identify significant genetic variations more or less stringently. For example, CNV calls generated by two or more of DNA Analytics, Aberration Detection Module 2 (ADM2) algorithms, and DNACopy algorithms can be defined as stringent CNVs. In some embodiments significant or stringent genetic variations can be tagged as identified or called if it can be found to have a minimal reciprocal overlap to a genetic variation detected by one or more platforms and/or methods described herein. For example, a minimum of 50% reciprocal overlap can be used to tag the CNVs as identified or called. For example, significant or stringent genetic variations can be tagged as identified or called if it can be found to have a reciprocal overlap of more than about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95%, 99%, or equal to 100%, to a genetic variation detected by one or more platforms and/or methods described herein. For example, significant or stringent genetic variations can be tagged as identified or called if it can be found to have a reciprocal overlap of more than about 50% reciprocal overlap to a genetic variation detected by one or more platforms and/or methods described herein. In another embodiment, genetic variations can be detected from the $\log_2$ ratio values calculated for individual probes present on an aCGH microarray via a statistical comparison of the probe's $\log_2$ ratio value in a cohort of subjects with PML to the probe's $\log_2$ ratio value in a cohort of subjects without PML.

[0329] In some embodiments, a threshold log ratio value can be used to determine losses and gains. A log ratio value can be any log ratio value; for example, a log ratio value can be a $\log_2$ ratio or a $\log_{10}$ ratio. In some embodiments, a CNV segment whose median $\log_2$ ratio is less than or equal to a $\log_2$ ratio threshold value can be classified as a loss. For example, any segment whose median $\log_2$ ratio is less than or equal to $-0.1, -0.11, -0.12, -0.13, -0.14, -0.15, -0.16, -0.17, -0.18, -0.19, -0.2, -0.21, -0.22, -0.23, -0.24, -0.25, -0.26, -0.27, -0.28, -0.29, -0.3, -0.31, -0.32, -0.33, -0.34, -0.35, -0.36, -0.37, -0.38, -0.39, -0.4, -0.41, -0.42, -0.43, -0.44, -0.45, -0.46, -0.47, -0.48, -0.49, -0.5, -0.55, -0.6, -0.65, -0.7, -0.75, -0.8, -0.85, -0.9, -0.95, -1, -1.1, -1.2, -1.3, -1.4, -1.5, -1.6, -1.7, -1.8, -1.9, -2, -2.1, -2.2, -2.3, -2.4, -2.5, -2.6, -2.7, -2.8, -2.9, -3, -3.1, -3.2, -3.3, -3.4, -3.5, -3.6, -3.7, -3.8, -3.9, -4, -4.1, -4.2, -4.3, -4.4, -4.5, -4.6, -4.7, -4.8, -4.9, -5,$

$-5.5, -6, -6.5, -7, -7.5, -8, -8.5, -9, -9.5, -10, -11, -12, -13, -14, -15, -16, -17, -18, -19, -20$ or less, can be classified as a loss.

[0330] In some embodiments, one algorithm can be used to call or identify significant genetic variations, wherein any segment whose median $\log_2$ ratio was less than or equal to $-0.1, -0.11, -0.12, -0.13, -0.14, -0.15, -0.16, -0.17, -0.18, -0.19, -0.2, -0.21, -0.22, -0.23, -0.24, -0.25, -0.26, -0.27, -0.28, -0.29, -0.3, -0.31, -0.32, -0.33, -0.34, -0.35, -0.36, -0.37, -0.38, -0.39, -0.4, -0.41, -0.42, -0.43, -0.44, -0.45, -0.46, -0.47, -0.48, -0.49, -0.5, -0.55, -0.6, -0.65, -0.7, -0.75, -0.8, -0.85, -0.9, -0.95, -1, -1.1, -1.2, -1.3, -1.4, -1.5, -1.6, -1.7, -1.8, -1.9, -2, -2.1, -2.2, -2.3, -2.4, -2.5, -2.6, -2.7, -2.8, -2.9, -3, -3.1, -3.2, -3.3, -3.4, -3.5, -3.6, -3.7, -3.8, -3.9, -4, -4.1, -4.2, -4.3, -4.4, -4.5, -4.6, -4.7, -4.8, -4.9, -5, -5.5, -6, -6.5, -7, -7.5, -8, -8.5, -9, -9.5, -10, -11, -12, -13, -14, -15, -16, -17, -18, -19, -20$ or less, can be classified as a loss. For example, any CNV segment whose median $\log_2$ ratio is less than -0.35 as determined by DNACopy can be classified as a loss. For example, losses can be determined according to a threshold $\log_2$ ratio, which can be set at -0.35 . In another embodiment, losses can be determined according to a threshold $\log_2$ ratio, which can be set at -0.5 .

[0331] In some embodiments, two algorithms can be used to call or identify significant genetic variations, wherein any segment whose median $\log_2$ ratio is less than or equal to $-0.1, -0.11, -0.12, -0.13, -0.14, -0.15, -0.16, -0.17, -0.18, -0.19, -0.2, -0.21, -0.22, -0.23, -0.24, -0.25, -0.26, -0.27, -0.28, -0.29, -0.3, -0.31, -0.32, -0.33, -0.34, -0.35, -0.36, -0.37, -0.38, -0.39, -0.4, -0.41, -0.42, -0.43, -0.44, -0.45, -0.46, -0.47, -0.48, -0.49, -0.5, -0.55, -0.6, -0.65, -0.7, -0.75, -0.8, -0.85, -0.9, -0.95, -1, -1.1, -1.2, -1.3, -1.4, -1.5, -1.6, -1.7, -1.8, -1.9, -2, -2.1, -2.2, -2.3, -2.4, -2.5, -2.6, -2.7, -2.8, -2.9, -3, -3.1, -3.2, -3.3, -3.4, -3.5, -3.6, -3.7, -3.8, -3.9, -4, -4.1, -4.2, -4.3, -4.4, -4.5, -4.6, -4.7, -4.8, -4.9, -5, -5.5, -6, -6.5, -7, -7.5, -8, -8.5, -9, -9.5, -10, -11, -12, -13, -14, -15, -16, -17, -18, -19, -20$ or less, as determined by one algorithm, and wherein any segment whose median $\log_2$ ratio is less than or equal to $-0.1, -0.11, -0.12, -0.13, -0.14, -0.15, -0.16, -0.17, -0.18, -0.19, -0.2, -0.21, -0.22, -0.23, -0.24, -0.25, -0.26, -0.27, -0.28, -0.29, -0.3, -0.31, -0.32, -0.33, -0.34, -0.35, -0.36, -0.37, -0.38, -0.39, -0.4, -0.41, -0.42, -0.43, -0.44, -0.45, -0.46, -0.47, -0.48, -0.49, -0.5, -0.55, -0.6, -0.65, -0.7, -0.75, -0.8, -0.85, -0.9, -0.95, -1, -1.1, -1.2, -1.3, -1.4, -1.5, -1.6, -1.7, -1.8, -1.9, -2, -2.1, -2.2, -2.3, -2.4, -2.5, -2.6, -2.7, -2.8, -2.9, -3, -3.1, -3.2, -3.3, -3.4, -3.5, -3.6, -3.7, -3.8, -3.9, -4, -4.1, -4.2, -4.3, -4.4, -4.5, -4.6, -4.7, -4.8, -4.9, -5, -5.5, -6, -6.5, -7, -7.5, -8, -8.5, -9, -9.5, -10, -11, -12, -13, -14, -15, -16, -17, -18, -19, -20$ or less, as determined by the other algorithm can be classified as a loss. For example, CNV calling can comprise using the Aberration Detection Module 2 (ADM2) algorithm and the DNACopy algorithm, wherein losses can be determined according to a two threshold $\log_2$ ratios, wherein the Aberration Detection Module 2 (ADM2) algorithm $\log_2$ ratio can be -0.25 and the DNACopy algorithm $\log_2$ ratio can be -0.41 .

[0332] In some embodiments, the use of two algorithms to call or identify significant genetic variations can be a stringent method. In some embodiments, the use of two algorithms to call or identify significant genetic variations can be

a more stringent method compared to the use of one algorithm to call or identify significant genetic variations.

[0333] In some embodiments, any CNV segment whose median log 2 ratio is greater than a log 2 ratio threshold value can be classified as a gain. For example, any segment whose median log 2 ratio is greater than 0.1, 0.11, 0.12, 0.13, 0.14, 0.15, 0.16, 0.17, 0.18, 0.19, 0.2, 0.21, 0.22, 0.23, 0.24, 0.25, 0.26, 0.27, 0.28, 0.29, 0.3, 0.31, 0.32, 0.33, 0.34, 0.35, 0.36, 0.37, 0.38, 0.39, 0.4, 0.41, 0.42, 0.43, 0.44, 0.45, 0.46, 0.47, 0.48, 0.49, 0.5, 0.55, 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, 0.95, 1, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3, or more can be classified as a gain.

[0334] In some embodiments, one algorithm can be used to call or identify significant genetic variations, wherein any segment whose median log 2 ratio is greater than or equal to 0.1, 0.11, 0.12, 0.13, 0.14, 0.15, 0.16, 0.17, 0.18, 0.19, 0.2, 0.21, 0.22, 0.23, 0.24, 0.25, 0.26, 0.27, 0.28, 0.29, 0.3, 0.31, 0.32, 0.33, 0.34, 0.35, 0.36, 0.37, 0.38, 0.39, 0.4, 0.41, 0.42, 0.43, 0.44, 0.45, 0.46, 0.47, 0.48, 0.49, 0.5, 0.55, 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, 0.95, 1, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3, or more can be classified as a gain. For example, any CNV segment whose median log 2 ratio is greater than 0.35 as determined by DNACopy can be classified as a gain. For example, gains can be determined according to a threshold log 2 ratio, which can be set at 0.35. In another embodiment, gains can be determined according to a threshold log 2 ratio, which can be set at 0.5.

[0335] In some embodiments, two algorithms can be used to call or identify significant genetic variations, wherein any segment whose median log 2 ratio is greater than or equal to 0.1, 0.11, 0.12, 0.13, 0.14, 0.15, 0.16, 0.17, 0.18, 0.19, 0.2, 0.21, 0.22, 0.23, 0.24, 0.25, 0.26, 0.27, 0.28, 0.29, 0.3, 0.31, 0.32, 0.33, 0.34, 0.35, 0.36, 0.37, 0.38, 0.39, 0.4, 0.41, 0.42, 0.43, 0.44, 0.45, 0.46, 0.47, 0.48, 0.49, 0.5, 0.55, 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, 0.95, 1, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3 or more, as determined by one algorithm, and wherein any segment whose median log 2 ratio is greater than or equal to 0.1, 0.11, 0.12, 0.13, 0.14, 0.15, 0.16, 0.17, 0.18, 0.19, 0.2, 0.21, 0.22, 0.23, 0.24, 0.25, 0.26, 0.27, 0.28, 0.29, 0.3, 0.31, 0.32, 0.33, 0.34, 0.35, 0.36, 0.37, 0.38, 0.39, 0.4, 0.41, 0.42, 0.43, 0.44, 0.45, 0.46, 0.47, 0.48, 0.49, or 0.5, 0.55, 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, 0.95, 1, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3, or more, as determined by the other algorithm the can be classified as a gain. For example, CNV calling can comprise using the Aberration Detection Module 2 (ADM2) algorithm and the DNACopy algorithm, wherein gains can be determined according to a two threshold log 2 ratios, wherein the Aberration Detection Module 2 (ADM2) algorithm log 2 ratio can be 0.25 and the DNACopy algorithm log 2 ratio can be 0.32.

[0336] Any CNV segment whose absolute (median log-ratio/mad) value is less than 2 can be excluded (not identified as a significant genetic variation). For example, any CNV segment whose absolute (median log-ratio/mad) value is less than 2, 1.9, 1.8, 1.7, 1.6, 1.5, 1.4, 1.3, 1.2, 1.1, 1, 0.9, 0.8, 0.7, 0.6, or 0.5 or less can be excluded.

[0337] In some embodiments, multivariate analyses or joint risk analyses, including the use of multiplicative model for overall risk assessment, can subsequently be used to determine the overall risk conferred based on the genotype

status at the multiple loci. Use of a multiplicative model, for example, assuming that the risk of individual risk variants multiply to establish the overall effect, allows for a straightforward calculation of the overall risk for multiple markers. The multiplicative model is a parsimonious model that usually fits the data of complex traits reasonably well. Deviations from multiplicity have been rarely described in the context of common variants for common diseases, and if reported are usually only suggestive since very large sample sizes can be required to be able to demonstrate statistical interactions between loci. Assessment of risk based on such analysis can subsequently be used in the methods, uses and kits of the disclosure, as described herein.

[0338] In some embodiments, the significance of increased or decreased susceptibility can be measured by a percentage. In some embodiments, a significant increased susceptibility can be measured as a relative susceptibility of at least 1.2, including but not limited to: at least 1.3, at least 1.4, at least 1.5, at least 1.6, at least 1.7, at least 1.8, at least 1.9, at least 2.0, at least 2.5, at least 3.0, at least 4.0, at least 5.0, at least 6.0, at least 7.0, at least 8.0, at least 9.0, at least 10.0, and at least 15.0. In some embodiments, a relative susceptibility of at least 2.0, at least 3.0, at least 4.0, at least 5.0, at least 6.0, or at least 10.0 is significant. Other values for significant susceptibility are also contemplated, for example, at least 2.5, 3.5, 4.5, 5.5, or any suitable other numerical values, wherein the values are also within scope of the present disclosure. In some embodiments, a significant increase in susceptibility is at least about 20%, including but not limited to about 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 100%, 150%, 200%, 300%, 400%, 500%, 600%, 700%, 800%, 900%, 1000%, and 1500%. In one particular embodiment, a significant increase in susceptibility is at least 100%. In other embodiments, a significant increase in susceptibility is at least 200%, at least 300%, at least 400%, at least 500%, at least 700%, at least 800%, at least 900% and at least 1000%. Other cutoffs or ranges as deemed suitable by the person skilled in the art to characterize the disclosure are also contemplated, and those are also within scope of the present disclosure. In certain embodiments, a significant increase in susceptibility is characterized by a p-value, such as a p-value of less than 0.5, less than 0.4, less than 0.3, less than 0.2, less than 0.1, less than 0.05, less than 0.01, less than 0.001, less than 0.0001, less than 0.00001, less than 0.000001, less than 0.0000001, or less than 0.000000001.

[0339] In some embodiments, an individual who is at a decreased susceptibility for or the lack of presence of a condition (e.g., PML) can be an individual in whom at least one genetic variation, conferring decreased susceptibility for or the lack of presence of the condition is identified. In some embodiments, the genetic variations conferring decreased susceptibility are also protective. In one aspect, the genetic variations can confer a significant decreased susceptibility of or lack of presence of PML.

[0340] In some embodiments, significant decreased susceptibility can be measured as a relative susceptibility of less than 0.9, including but not limited to less than 0.9, less than 0.8, less than 0.7, less than 0.6, less than 0.5, less than 0.4, less than 0.3, less than 0.2 and less than 0.1. In some embodiments, the decrease in susceptibility is at least 20%, including but not limited to at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least

55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95% and at least 98%. Other cutoffs or ranges as deemed suitable by the person, skilled in the art to characterize the disclosure are however also contemplated, and those are also within scope of the present disclosure. In certain embodiments, a significant decrease in susceptibility is characterized by a p-value, such as a p-value of less than 0.05, less than 0.01, less than 0.001, less than 0.0001, less than 0.00001, less than 0.000001, less than 0.0000001, or less than 0.000000001. Other tests for significance can be used, for example, a Fisher-exact test. Other statistical tests of significance known to the skilled person are also contemplated and are also within scope of the disclosure.

[0341] In some preferred embodiments, the significance of increased or decreased susceptibility can be determined according to the ratio of measurements from a test subject to a reference subject. In some embodiments, losses or gains of one or more CNVs can be determined according to a threshold $\log_2$ ratio determined by these measurements. In some embodiments, a $\log_2$ ratio value greater than 0.35, or 0.5, is indicative of a gain of one or more CNVs. In some embodiments, a $\log_2$ ratio value less than -0.35, or -0.5, is indicative of a loss of one or more CNVs. In some embodiments, the ratio of measurements from a test subject to a reference subject may be inverted such that the $\log_2$ ratios of copy number gains are negative and the $\log_2$ ratios of copy number losses are positive.

[0342] In some embodiments, the combined or overall susceptibility associated with a plurality of variants associated with PML can also be assessed; for example, the genetic variations described herein to be associated with susceptibility to PML can be combined with other common genetic risk factors. Combined risk for such genetic variants can be estimated in an analogous fashion to the methods described herein.

[0343] Calculating risk conferred by a particular genotype for the individual can be based on comparing the genotype of the individual to previously determined risk expressed, for example, as a relative risk (RR) or an odds ratio (OR), for the genotype, for example, for a heterozygous carrier of an at-risk variant for PML. An odds ratio can be a statistical measure used as a metric of causality. For example, in genetic disease research it can be used to convey the significance of a variant in a disease cohort relative to an unaffected/normal cohort. The calculated risk for the individual can be the relative risk for a subject, or for a specific genotype of a subject, compared to the average population. The average population risk can be expressed as a weighted average of the risks of different genotypes, using results from a reference population, and the appropriate calculations to calculate the risk of a genotype group relative to the population can then be performed. Alternatively, the risk for an individual can be based on a comparison of particular genotypes, for example, heterozygous and/or homozygous carriers of an at-risk allele of a marker compared with non-carriers of the at-risk allele (or pair of alleles in the instance of compound heterozygous variants, wherein one variant impacts the maternally inherited allele and the other impacts the paternally inherited allele). Using the population average can, in certain embodiments, be more convenient, since it provides a measure that can be easy to interpret for

the user, for example, a measure that gives the risk for the individual, based on his/her genotype, compared with the average in the population.

[0344] In some embodiments, the OR value can be calculated as follows: $OR = (A/(N1-A))/(U/(N2-U))$, where A=number of affected cases with variant, N1=total number of affected cases, U=number of unaffected cases with variant and N2=total number of unaffected cases. In circumstances where U=0, it is conventional to set U=1, so as to avoid infinities. In some preferred embodiments, the OR can be calculated essentially as above, except that where U or A=0, 0.5 is added to all of A, N1, U, N2. In another embodiment, a Fisher's Exact Test (FET) can be calculated using standard methods. In another embodiment, the p-values can be corrected for false discovery rate (FDR) using the Benjamini-Hochberg method (Benjamini Y. and Hochberg Y., *J. Royal Statistical Society* 57:289 (1995); Osborne J. A. and Barker C. A. (2007)).

[0345] In certain embodiments of the disclosure, a genetic variation is correlated to PML by referencing genetic variation data to a look-up table that comprises correlations between the genetic variation and PML. The genetic variation in certain embodiments comprises at least one indication of the genetic variation. In some embodiments, the table comprises a correlation for one genetic variation. In other embodiments, the table comprises a correlation for a plurality of genetic variations in both scenarios, by referencing to a look-up table that gives an indication of a correlation between a genetic variation and PML, a risk for PML, or a susceptibility to PML, can be identified in the individual from whom the nucleic acid sample is derived.

[0346] The present disclosure also pertains to methods of clinical screening, for example, diagnosis, prognosis, or theragnosis of a subject performed by a medical professional using the methods disclosed herein. In other embodiments, the disclosure pertains to methods of screening performed by a layman. The layman can be a customer of a genotyping, microarray, exome sequencing, or whole genome sequencing service provider. The layman can also be a genotype, microarray, exome sequencing, or whole genome sequencing service provider, who performs genetic analysis on a DNA sample from an individual, in order to provide service related to genetic risk factors for particular traits or diseases, based on the genotype status of the subject obtained from use of the methods described herein. The resulting genotype or genetic information can be made available to the individual and can be compared to information about PML or risk of developing PML associated with one or various genetic variations, including but not limited to, information from public or private genetic variation databases or literature and scientific publications. The screening applications of PML-associated genetic variations, as described herein, can, for example, be performed by an individual, a health professional, or a third party, for example a service provider who interprets genotype information from the subject. In some embodiments the genetic analysis is performed in a CLIA-certified laboratory (i.e. the federal regulatory standards the U.S. that are specified in the Clinical Laboratory Improvement Amendments, administered by the Centers for Medicare and Medicaid Services) or equivalent laboratories in Europe and elsewhere in the world.

[0347] The information derived from analyzing sequence data can be communicated to any particular body, including the individual from which the nucleic acid sample or

sequence data is derived, a guardian or representative of the individual, clinician, research professional, medical professional, service provider, and medical insurer or insurance company. Medical professionals can be, for example, doctors, nurses, medical laboratory technologists, and pharmacists. Research professionals can be, for example, principle investigators, research technicians, postdoctoral trainees, and graduate students.

[0348] In some embodiments, a professional can be assisted by determining whether specific genetic variants are present in a nucleic acid sample from a subject, and communicating information about genetic variants to a professional. After information about specific genetic variants is reported, a medical professional can take one or more actions that can affect subject care. For example, a medical professional can record information in the subject's medical record (e.g., electronic health record or electronic medical record, including, but not limited to, country-scale health services such as the National Health Service in the United Kingdom) regarding the subject's risk of developing PML. In some embodiments, a medical professional can record information regarding risk assessment, or otherwise transform the subject's medical record, to reflect the subject's current medical condition. In some embodiments, a medical professional can review and evaluate a subject's entire medical record and assess multiple treatment strategies for clinical intervention of a subject's condition. In another embodiment, information can be recorded in the context of the system developed by the World Health Organization (WHO), the International Statistical Classification of Diseases and Related Health Problems (ICD), which is currently using the 10th revision (ICD-10 codes). For example, the ICD-10 code for PML is A81.2, whereas the ICD-10 code for multiple sclerosis is G35.

[0349] A medical professional can initiate or modify treatment after receiving information regarding a subject's screening for PML, for example. In some embodiments, a medical professional can recommend a change in therapy or exclude a therapy. In some embodiments, a medical professional can enroll a subject in a clinical trial for, by way of example, detecting correlations between a haplotype as described herein and any measurable or quantifiable parameter relating to the outcome of the treatment as described above.

[0350] In some embodiments, a medical professional can communicate information regarding a subject's screening of developing PML to a subject or a subject's family. In some embodiments, a medical professional can provide a subject and/or a subject's family with information regarding PML and risk assessment information, including treatment options, and referrals to specialists. In some embodiments, a medical professional can provide a copy of a subject's medical records to a specialist. In some embodiments, a research professional can apply information regarding a subject's risk of developing PML to advance scientific research. In some embodiments, a research professional can obtain a subject's haplotype as described herein to evaluate a subject's enrollment, or continued participation, in a research study or clinical trial. In some embodiments, a research professional can communicate information regarding a subject's screening of PML to a medical professional. In some embodiments, a research professional can refer a subject to a medical professional.

[0351] Any appropriate method can be used to communicate information to another person. For example, information can be given directly or indirectly to a professional and a laboratory technician can input a subject's genetic variation as described herein into a computer-based record. In some embodiments, information is communicated by making a physical alteration to medical or research records. For example, a medical professional can make a permanent notation or flag a medical record for communicating the risk assessment to other medical professionals reviewing the record. In addition, any type of communication can be used to communicate the risk assessment information. For example, mail, e-mail, telephone, and face-to-face interactions can be used. The information also can be communicated to a professional by making that information electronically available to the professional. For example, the information can be communicated to a professional by placing the information on a computer database such that the professional can access the information. In addition, the information can be communicated to a hospital, clinic, or research facility serving as an agent for the professional.

[0352] Results of these tests, and optionally interpretive information, can be returned to the subject, the health care provider or to a third party. The results can be communicated to the tested subject, for example, with a prognosis and optionally interpretive materials that can help the subject understand the test results and prognosis; used by a health care provider, for example, to determine whether to administer a specific drug, or whether a subject should be assigned to a specific category, for example, a category associated with a specific disease endophenotype, or with drug response or non-response; used by a third party such as a healthcare payer, for example, an insurance company or HMO, or other agency, to determine whether or not to reimburse a health care provider for services to the subject, or whether to approve the provision of services to the subject. For example, the healthcare payer can decide to reimburse a health care provider for treatments for PML if the subject has PML or has an increased risk of developing PML.

[0353] Also provided herein are databases that include a list of genetic variations as described herein, and wherein the list can be largely or entirely limited to genetic variations identified as useful for screening PML as described herein. The list can be stored, for example, on a flat file or computer-readable medium. The databases can further include information regarding one or more subjects, for example, whether a subject is affected or unaffected, clinical information such as endophenotype, age of onset of symptoms, any treatments administered and outcomes, for example, data relevant to pharmacogenomics, diagnostics, prognostics or theranostics, and other details, for example, data about the disorder in the subject, or environmental (e.g., including, but not limited to, infection or a history of infection with HIV or JCV) or other genetic factors. The databases can be used to detect correlations between a particular haplotype and the information regarding the subject.

[0354] The methods described herein can also include the generation of reports for use, for example, by a subject, care giver, or researcher, that include information regarding a subject's genetic variations, and optionally further information such as treatments administered, treatment history, medical history, predicted response, and actual response.

The reports can be recorded in a tangible medium, e.g., a computer-readable disk, a solid state memory device, or an optical storage device.

Methods of Screening Using Variations in RNA and/or Polypeptides

[0355] In some embodiments of the disclosure, screening of PML can be made by examining or comparing changes in expression, localization, binding partners, and composition of a polypeptide encoded by a nucleic acid variant associated with PML, for example, in those instances where the genetic variations of the present disclosure results in a change in the composition or expression of the polypeptide and/or RNA, for example, mRNAs, microRNAs (miRNAs), and other noncoding RNAs (ncRNAs). Thus, screening of PML can be made by examining expression and/or composition of one of these polypeptides and/or RNA, or another polypeptide and/or RNA encoded by a nucleic acid associated with PML, in those instances where the genetic variation of the present disclosure results in a change in the expression, localization, binding partners, and/or composition of the polypeptide and/or RNA. In some embodiments, screening can comprise diagnosing a subject. In some embodiments, screening can comprise determining a prognosis of a subject, for example determining the susceptibility of developing PML. In some embodiments, screening can comprise theranosing a subject.

[0356] The genetic variations described herein that show association to PML can play a role through their effect on one or more of these genes, either by directly impacting one or more genes or influencing the expression of one or more nearby genes. For example, while not intending to be limited by theory, it is generally expected that a deletion of a chromosomal segment comprising a particular gene, or a fragment of a gene, can either result in an altered composition or expression, or both, of the encoded polypeptide and/or mRNA. Likewise, duplications, or high number copy number variations, are in general expected to result in increased expression of encoded polypeptide and/or RNA if the gene they are expressed from is fully encompassed within the duplicated (or triplicated, or even higher copy number gains) genomic segment, or conversely can result in decreased expression or a disrupted RNA or polypeptide if one or both breakpoints of the copy number gain disrupt a given gene. Other possible mechanisms affecting genes within a genetic variation region include, for example, effects on transcription, effects on RNA splicing, alterations in relative amounts of alternative splice forms of mRNA, effects on RNA stability, effects on transport from the nucleus to cytoplasm, and effects on the efficiency and accuracy of translation. Thus, DNA variations can be detected directly, using the subjects unamplified or amplified genomic DNA, or indirectly, using RNA or DNA obtained from the subject's tissue(s) that are present in an aberrant form or expression level as a result of the genetic variations of the disclosure showing association to PML. In another embodiment, DNA variations can be detected indirectly using a polypeptide or protein obtained from the subject's tissue(s) that is present in an aberrant form or expression level as a result of genetic variations of the disclosure showing association to the PML. In another embodiment, an aberrant form or expression level of a polypeptide or protein that results from one or more genetic variations of the disclosure showing association to PML can be detected indirectly via another polypeptide or protein present in the

same biological/cellular pathway that is modulated or interacts with said polypeptide or protein that results from one or more genetic variations of the disclosure. In some embodiments, the genetic variations of the disclosure showing association to PML can affect the expression of a gene within the genetic variation region. In some embodiments, a genetic variation affecting an exonic region of a gene can affect, disrupt, or modulate the expression of the gene. In some embodiments, a genetic variation affecting an intronic or intergenic region of a gene can affect, disrupt, or modulate the expression of the gene.

[0357] Certain genetic variation regions can have flanking duplicated segments, and genes within such segments can have altered expression and/or composition as a result of such genomic alterations. Regulatory elements affecting gene expression can be located far away, even as far as tens or hundreds of kilobases away, from the gene that is regulated by said regulatory elements. Thus, in some embodiments, regulatory elements for genes that are located outside the gene (e.g., upstream or downstream of the gene) can be located within the genetic variation, and thus be affected by the genetic variation. It is thus contemplated that the detection of the genetic variations described herein, can be used for assessing expression for one or more of associated genes not directly impacted by the genetic variations. In some embodiments, a genetic variation affecting an intergenic region of a gene can affect, disrupt, or modulate the expression of a gene located elsewhere in the genome, such as described above. For example, a genetic variation affecting an intergenic region of a gene can affect, disrupt, or modulate the expression of a transcription factor, located elsewhere in the genome, which regulates the gene. Regulatory elements can also be located within a gene, such as within intronic regions, and similarly impact the expression level of the gene and ultimately the protein expression level without changing the structure of the protein. The effects of genetic variants on regulatory elements can manifest in a tissue-specific manner; for example, one or more transcription factors that bind to the regulatory element that is impacted by one or more genetic variations may be expressed at higher concentration in neurons as compared to skin cells (i.e., the impact of the one or more genetic variations may be primarily evident in neuronal cells).

[0358] In some embodiments, genetic variations of the disclosure showing association to PML can affect protein expression at the translational level. It can be appreciated by those skilled in the art that this can occur by increased or decreased expression of one or more microRNAs (miRNAs) that regulates expression of a protein known to be important, or implicated, in the cause, onset, or progression of PML. Increased or decreased expression of the one or more miRNAs can result from gain or loss of the whole miRNA gene, disruption or impairment of a portion of the gene (e.g., by an indel or CNV), or even a single base change (SNP or SNV) that produces an altered, non-functional or aberrant functioning miRNA sequence. It can also be appreciated by those skilled in the art that the expression of protein, for example, one known to cause PML by increased or decreased expression, can result due to a genetic variation that results in alteration of an existing miRNA binding site within the polypeptide's mRNA transcript, or even creates a new miRNA binding site that leads to aberrant polypeptide expression.

[0359] A variety of methods can be used for detecting polypeptide composition and/or expression levels, including but not limited to enzyme linked immunosorbent assays (ELISA), Western blots, spectroscopy, mass spectrometry, peptide arrays, colorimetry, electrophoresis, isoelectric focusing, immunoprecipitations, immunoassays, and immunofluorescence and other methods well-known in the art. A test nucleic acid sample from a subject can be assessed for the presence of an alteration in the expression and/or an alteration in composition of the polypeptide encoded by a nucleic acid associated with PML. An “alteration” in the polypeptide expression or composition, as used herein, refers to an alteration in expression or composition in a test nucleic acid sample, as compared to the expression or composition of the polypeptide in a control nucleic acid sample. Such alteration can, for example, be an alteration in the quantitative polypeptide expression or can be an alteration in the qualitative polypeptide expression, for example, expression of a mutant polypeptide or of a different splicing variant, or a combination thereof. In some embodiments, screening of PML can be made by detecting a particular splicing variant encoded by a nucleic acid associated with PML, or a particular pattern of splicing variants.

[0360] Antibodies can be polyclonal or monoclonal and can be labeled or unlabeled. An intact antibody or a fragment thereof can be used. The term “labeled”, with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly labeled as previously described herein. Other non-limiting examples of indirect labeling include detection of a primary antibody using a labeled secondary antibody, for example, a fluorescently-labeled secondary antibody and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently-labeled streptavidin.

Methods of Detecting Genetic Variations

[0361] In some embodiments, standard techniques for genotyping for the presence genetic variations, for example, amplification, can be used. Amplification of nucleic acids can be accomplished using methods known in the art. Generally, sequence information from the region of interest can be used to design oligonucleotide primers that can be identical or similar in sequence to opposite strands of a template to be amplified. In some embodiments, amplification methods can include but are not limited to, fluorescence-based techniques utilizing PCR, for example, ligase chain reaction (LCR), Nested PCR, transcription amplification, self-sustained sequence replication, nucleic acid based sequence amplification (NASBA), and multiplex ligation-dependent probe amplification (MLPA). Guidelines for selecting primers for PCR amplification are well known in the art. In some embodiments, a computer program can be used to design primers, for example, Oligo (National Biosciences, Inc, Plymouth Minn), MacVector (Kodak/IBI), and GCG suite of sequence analysis programs.

[0362] In some embodiments, commercial methodologies available for genotyping, for example, SNP genotyping, can be used, but are not limited to, TaqMan genotyping assays (Applied Biosystems), SNplex platforms (Applied Biosystems), gel electrophoresis, capillary electrophoresis, size exclusion chromatography, mass spectrometry, for example, MassARRAY system (Sequenom), minisequencing meth-

ods, real-time Polymerase Chain Reaction (PCR), Bio-Plex system (BioRad), CEQ and SNPstream systems (Beckman), array hybridization technology, for example, Affymetrix GeneChip (Perlegen), BeadArray Technologies, for example, Illumina GoldenGate and Infinium assays, array tag technology, Multiplex Ligation-dependent Probe Amplification (MLPA), and endonuclease-based fluorescence hybridization technology (Invader assay, either using unamplified or amplified genomic DNA, or unamplified total RNA, or unamplified or amplified cDNA; Third Wave/Hologic). PCR can be a procedure in which target nucleic acid is amplified in a manner similar to that described in U.S. Pat. No. 4,683,195 and subsequent modifications of the procedure described therein. PCR can include a three phase temperature cycle of denaturation of DNA into single strands, annealing of primers to the denatured strands, and extension of the primers by a thermostable DNA polymerase enzyme. This cycle can be repeated so that there are enough copies to be detected and analyzed. In some embodiments, real-time quantitative PCR can be used to determine genetic variations, wherein quantitative PCR can permit both detection and quantification of a DNA sequence in a nucleic acid sample, for example, as an absolute number of copies or as a relative amount when normalized to DNA input or other normalizing genes. In some embodiments, methods of quantification can include the use of fluorescent dyes that can intercalate with double-stranded DNA, and modified DNA oligonucleotide probes that can fluoresce when hybridized with a complementary DNA.

[0363] In some embodiments of the disclosure, a nucleic acid sample obtained from the subject can be collected and PCR can be used to amplify a fragment of nucleic acid that comprises one or more genetic variations that can be indicative of a susceptibility to PML. In some embodiments, detection of genetic variations can be accomplished by expression analysis, for example, by using quantitative PCR. In some embodiments, this technique can assess the presence or absence of a genetic alteration in the expression or composition of one or more polypeptides or splicing variants encoded by a nucleic acid associated with PML.

[0364] In some embodiments, the nucleic acid sample from a subject containing a SNP can be amplified by PCR prior to detection with a probe. In such an embodiment, the amplified DNA serves as the template for a detection probe and, in some embodiments, an enhancer probe. Certain embodiments of the detection probe, the enhancer probe, and/or the primers used for amplification of the template by PCR can comprise the use of modified bases, for example, modified A, T, C, G, and U, wherein the use of modified bases can be useful for adjusting the melting temperature of the nucleotide probe and/or primer to the template DNA. In some embodiments, modified bases are used in the design of the detection nucleotide probe. Any modified base known to the skilled person can be selected in these methods, and the selection of suitable bases is well within the scope of the skilled person based on the teachings herein and known bases available from commercial sources as known to the skilled person.

[0365] In some embodiments, identification of genetic variations can be accomplished using hybridization methods. The presence of a specific marker allele or a particular genomic segment comprising a genetic variation, or representative of a genetic variation, can be indicated by sequence-specific hybridization of a nucleic acid probe

specific for the particular allele or the genetic variation in a nucleic acid sample that has or has not been amplified but methods described herein. The presence of more than one specific marker allele or several genetic variations can be indicated by using two or more sequence-specific nucleic acid probes, wherein each is specific for a particular allele and/or genetic variation.

[0366] Hybridization can be performed by methods well known to the person skilled in the art, for example, hybridization techniques such as fluorescent in situ hybridization (FISH), Southern analysis, Northern analysis, or in situ hybridization. In some embodiments, hybridization refers to specific hybridization, wherein hybridization can be performed with no mismatches. Specific hybridization, if present, can be using standard methods. In some embodiments, if specific hybridization occurs between a nucleic acid probe and the nucleic acid in the nucleic acid sample, the nucleic acid sample can contain a sequence that can be complementary to a nucleotide present in the nucleic acid probe. In some embodiments, if a nucleic acid probe can contain a particular allele of a polymorphic marker, or particular alleles for a plurality of markers, specific hybridization is indicative of the nucleic acid being completely complementary to the nucleic acid probe, including the particular alleles at polymorphic markers within the probe. In some embodiments a probe can contain more than one marker alleles of a particular haplotype, for example, a probe can contain alleles complementary to 2, 3, 4, 5 or all of the markers that make up a particular haplotype. In some embodiments detection of one or more particular markers of the haplotype in the nucleic acid sample is indicative that the source of the nucleic acid sample has the particular haplotype.

[0367] In some embodiments, PCR conditions and primers can be developed that amplify a product only when the variant allele is present or only when the wild type allele is present, for example, allele-specific PCR. In some embodiments of allele-specific PCR, a method utilizing a detection oligonucleotide probe comprising a fluorescent moiety or group at its 3' terminus and a quencher at its 5' terminus, and an enhancer oligonucleotide, can be employed (see e.g., Kutuyavin et al., *Nucleic Acid Res.* 34:e128 (2006)).

[0368] An allele-specific primer/probe can be an oligonucleotide that is specific for particular a polymorphism can be prepared using standard methods. In some embodiments, allele-specific oligonucleotide probes can specifically hybridize to a nucleic acid region that contains a genetic variation. In some embodiments, hybridization conditions can be selected such that a nucleic acid probe can specifically bind to the sequence of interest, for example, the variant nucleic acid sequence.

[0369] In some embodiments, allele-specific restriction digest analysis can be used to detect the existence of a polymorphic variant of a polymorphism, if alternate polymorphic variants of the polymorphism can result in the creation or elimination of a restriction site. Allele-specific restriction digests can be performed, for example, with the particular restriction enzyme that can differentiate the alleles. In some embodiments, PCR can be used to amplify a region comprising the polymorphic site, and restriction fragment length polymorphism analysis can be conducted. In some embodiments, for sequence variants that do not alter a common restriction site, mutagenic primers can be

designed that can introduce one or more restriction sites when the variant allele is present or when the wild type allele is present.

[0370] In some embodiments, fluorescence polarization template-directed dye-terminator incorporation (FP-TDI) can be used to determine which of multiple polymorphic variants of a polymorphism can be present in a subject. Unlike the use of allele-specific probes or primers, this method can employ primers that can terminate adjacent to a polymorphic site, so that extension of the primer by a single nucleotide can result in incorporation of a nucleotide complementary to the polymorphic variant at the polymorphic site.

[0371] In some embodiments, DNA containing an amplified portion can be dot-blotted, using standard methods and the blot contacted with the oligonucleotide probe. The presence of specific hybridization of the probe to the DNA can then be detected. The methods can include determining the genotype of a subject with respect to both copies of the polymorphic site present in the genome, wherein if multiple polymorphic variants exist at a site, this can be appropriately indicated by specifying which variants are present in a subject. Any of the detection means described herein can be used to determine the genotype of a subject with respect to one or both copies of the polymorphism present in the subject's genome.

[0372] In some embodiments, a peptide nucleic acid (PNA) probe can be used in addition to, or instead of, a nucleic acid probe in the methods described herein. A PNA can be a DNA mimic having a peptide-like, inorganic backbone, for example, N-(2-aminoethyl) glycine units with an organic base (A, G, C, T or U) attached to the glycine nitrogen via a methylene carbonyl linker.

[0373] Nucleic acid sequence analysis can also be used to detect genetic variations, for example, genetic variations can be detected by sequencing exons, introns, 5' untranslated sequences, or 3' untranslated sequences. One or more methods of nucleic acid analysis that are available to those skilled in the art can be used to detect genetic variations, including but not limited to, direct manual sequencing, automated fluorescent sequencing, single-stranded conformation polymorphism assays (SSCP); clamped denaturing gel electrophoresis (CDGE); denaturing gradient gel electrophoresis (DGGE), two-dimensional gel electrophoresis (2DGE or TDGE); conformational sensitive gel electrophoresis (CSGE); denaturing high performance liquid chromatography (DHPLC), infrared matrix-assisted laser desorption/ionization (IR-MALDI) mass spectrometry, mobility shift analysis, quantitative real-time PCR, restriction enzyme analysis, heteroduplex analysis; chemical mismatch cleavage (CMC), RNase protection assays, use of polypeptides that recognize nucleotide mismatches, allele-specific PCR, real-time pyrophosphate DNA sequencing, PCR amplification in combination with denaturing high performance liquid chromatography (dHPLC), and combinations of such methods.

[0374] Sequencing can be accomplished through classic Sanger sequencing methods, which are known in the art. In some embodiments sequencing can be performed using high-throughput sequencing methods some of which allow detection of a sequenced nucleotide immediately after or upon its incorporation into a growing strand, for example, detection of sequence in substantially real time or real time. In some cases, high throughput sequencing generates at least

1,000, at least 5,000, at least 10,000, at least 20,000, at least 30,000, at least 40,000, at least 50,000, at least 100,000 or at least 500,000 sequence reads per hour; with each read being at least 50, at least 60, at least 70, at least 80, at least 90, at least 100, at least 120 or at least 150 bases per read (or 500-1,000 bases per read for 454).

[0375] High-throughput sequencing methods can include but are not limited to, Massively Parallel Signature Sequencing (MPSS, Lynx Therapeutics), Polony sequencing, 454 pyrosequencing, Illumina (Solexa) sequencing, Illumina (Solexa) sequencing using 10× Genomics library preparation, SOLiD sequencing, on semiconductor sequencing, DNA nanoball sequencing, Helioscope™ single molecule sequencing, Single Molecule SMRT™ sequencing, Single Molecule real time (RNAP) sequencing, Nanopore DNA sequencing, and/or sequencing by hybridization, for example, a non-enzymatic method that uses a DNA microarray, or microfluidic Sanger sequencing.

[0376] In some embodiments, high-throughput sequencing can involve the use of technology available by Helicos BioSciences Corporation (Cambridge, Mass.) such as the Single Molecule Sequencing by Synthesis (SMSS) method. SMSS is unique because it allows for sequencing the entire human genome in up to 24 hours. This fast sequencing method also allows for detection of a SNP/nucleotide in a sequence in substantially real time or real time. Finally, SMSS is powerful because, like the MIP technology, it does not use a pre-amplification step prior to hybridization. SMSS does not use any amplification. SMSS is described in US Publication Application Nos. 20060024711; 20060024678; 20060012793; 20060012784; and 20050100932. In some embodiments, high-throughput sequencing involves the use of technology available by 454 Life Sciences, Inc. (a Roche company, Branford, Conn.) such as the PicoTiterPlate device which includes a fiber optic plate that transmits chemiluminescent signal generated by the sequencing reaction to be recorded by a CCD camera in the instrument. This use of fiber optics allows for the detection of a minimum of 20 million base pairs in 4.5 hours.

[0377] In some embodiments, PCR-amplified single-strand nucleic acid can be hybridized to a primer and incubated with a polymerase, ATP sulfurylase, luciferase, apyrase, and the substrates luciferin and adenosine 5' phosphosulfate. Next, deoxynucleotide triphosphates corresponding to the bases A, C, G, and T (U) can be added sequentially. A base incorporation can be accompanied by release of pyrophosphate, which can be converted to ATP by sulfurylase, which can drive synthesis of oxyluciferin and the release of visible light. Since pyrophosphate release can be equimolar with the number of incorporated bases, the light given off can be proportional to the number of nucleotides adding in any one step. The process can repeat until the entire sequence can be determined. In some embodiments, pyrosequencing can be utilized to analyze amplicons to determine whether breakpoints are present. In some embodiments, pyrosequencing can map surrounding sequences as an internal quality control.

[0378] Pyrosequencing analysis methods are known in the art. Sequence analysis can include a four-color sequencing by ligation scheme (degenerate ligation), which involves hybridizing an anchor primer to one of four positions. Then an enzymatic ligation reaction of the anchor primer to a population of degenerate nonamers that are labeled with fluorescent dyes can be performed. At any given cycle, the

population of nonamers that is used can be structured such that the identity of one of its positions can be correlated with the identity of the fluorophore attached to that nonamer. To the extent that the ligase discriminates for complementarity at that queried position, the fluorescent signal can allow the inference of the identity of the base. After performing the ligation and four-color imaging, the anchor primer: nonamer complexes can be stripped and a new cycle begins. Methods to image sequence information after performing ligation are known in the art.

[0379] In some embodiments, analysis by restriction enzyme digestion can be used to detect a particular genetic variation if the genetic variation results in creation or elimination of one or more restriction sites relative to a reference sequence. In some embodiments, restriction fragment length polymorphism (RFLP) analysis can be conducted, wherein the digestion pattern of the relevant DNA fragment indicates the presence or absence of the particular genetic variation in the nucleic acid sample.

[0380] In some embodiments, arrays of oligonucleotide probes that can be complementary to target nucleic acid sequence segments from a subject can be used to identify genetic variations. In some embodiments, an array of oligonucleotide probes comprises an oligonucleotide array, for example, a microarray. In some embodiments, the present disclosure features arrays that include a substrate having a plurality of addressable areas, and methods of using them. At least one area of the plurality includes a nucleic acid probe that binds specifically to a sequence comprising a genetic variation, and can be used to detect the absence or presence of the genetic variation, for example, one or more SNPs, microsatellites, or CNVs, as described herein, to determine or identify an allele or genotype. For example, the array can include one or more nucleic acid probes that can be used to detect a genetic variation associated with a gene and/or gene product. In some embodiments, the array can further comprise at least one area that includes a nucleic acid probe that can be used to specifically detect another marker associated with PML as described herein.

[0381] Microarray hybridization can be performed by hybridizing a nucleic acid of interest, for example, a nucleic acid encompassing a genetic variation, with the array and detecting hybridization using nucleic acid probes. In some embodiments, the nucleic acid of interest is amplified prior to hybridization. Hybridization and detecting can be carried out according to standard methods described in Published PCT Applications: WO 92/10092 and WO 95/11995, and U.S. Pat. No. 5,424,186. For example, an array can be scanned to determine the position on the array to which the nucleic acid hybridizes. The hybridization data obtained from the scan can be, for example, in the form of fluorescence intensities as a function of location on the array.

[0382] Arrays can be formed on substrates fabricated with materials such as paper; glass; plastic, for example, polypropylene, nylon, or polystyrene; polyacrylamide; nitrocellulose; silicon; optical fiber; or any other suitable solid or semisolid support; and can be configured in a planar, for example, glass plates or silicon chips); or three dimensional, for example, pins, fibers, beads, particles, microtiter wells, and capillaries, configuration.

[0383] Methods for generating arrays are known in the art and can include for example; photolithographic methods (U.S. Pat. Nos. 5,143,854, 5,510,270 and 5,527,681); mechanical methods, for example, directed-flow methods

(U.S. Pat. No. 5,384,261); pin-based methods (U.S. Pat. No. 5,288,514); bead-based techniques (PCT US/93/04145); solid phase oligonucleotide synthesis methods; or by other methods known to a person skilled in the art (see, e.g., Bier, F. F., et al., *Adv Biochem Eng Biotechnol* 109:433-53 (2008); Hoheisel, J. D., *Nat Rev Genet* 7: 200-10 (2006); Fan, J. B., et al., *Methods Enzymol* 410:57-73 (2006); Raquoussis, J. & Elvidge, G., *Expert Rev Mol Design* 6: 145-52 (2006); Mockler, T. C., et al., *Genomics* 85: 1-15 (2005), and references cited therein, the entire teachings of each of which are incorporated by reference herein). Many additional descriptions of the preparation and use of oligonucleotide arrays for detection of polymorphisms can be found, for example, in U.S. Pat. Nos. 6,858,394, 6,429,027, 5,445,934, 5,700,637, 5,744,305, 5,945,334, 6,054,270, 6,300,063, 6,733,977, 7,364,858, EP 619 321, and EP 373 203, the entire teachings of which are incorporated by reference herein. Methods for array production, hybridization, and analysis are also described in Snijders et al., *Nat. Genetics* 29:263-264 (2001); Klein et al., *Proc. Natl. Acad. Sci. USA* 96:4494-4499 (1999); Albertson et al., *Breast Cancer Research and Treatment* 78:289-298 (2003); and Snijders et al., "BAC microarray based comparative genomic hybridization," in: Zhao et al., (eds), *Bacterial Artificial Chromosomes: Methods and Protocols*, Methods in Molecular Biology, Humana Press (2002).

[0384] In some embodiments, oligonucleotide probes forming an array can be attached to a substrate by any number of techniques, including, but not limited to, in situ synthesis, for example, high-density oligonucleotide arrays, using photolithographic techniques; spotting/printing a medium to low density on glass, nylon, or nitrocellulose; by masking; and by dot-blotting on a nylon or nitrocellulose hybridization membrane. In some embodiments, oligonucleotides can be immobilized via a linker, including but not limited to, by covalent, ionic, or physical linkage. Linkers for immobilizing nucleic acids and polypeptides, including reversible or cleavable linkers, are known in the art (U.S. Pat. No. 5,451,683 and WO98/20019). In some embodiments, oligonucleotides can be non-covalently immobilized on a substrate by hybridization to anchors, by means of magnetic beads, or in a fluid phase, for example, in wells or capillaries.

[0385] An array can comprise oligonucleotide hybridization probes capable of specifically hybridizing to different genetic variations. In some embodiments, oligonucleotide arrays can comprise a plurality of different oligonucleotide probes coupled to a surface of a substrate in different known locations. In some embodiments, oligonucleotide probes can exhibit differential or selective binding to polymorphic sites, and can be readily designed by one of ordinary skill in the art, for example, an oligonucleotide that is perfectly complementary to a sequence that encompasses a polymorphic site, for example, a sequence that includes the polymorphic site, within it, or at one end, can hybridize preferentially to a nucleic acid comprising that sequence, as opposed to a nucleic acid comprising an alternate polymorphic variant.

[0386] In some embodiments, arrays can include multiple detection blocks, for example, multiple groups of probes designed for detection of particular polymorphisms. In some embodiments, these arrays can be used to analyze multiple different polymorphisms. In some embodiments, detection blocks can be grouped within a single array or in multiple, separate arrays, wherein varying conditions, for example,

conditions optimized for particular polymorphisms, can be used during hybridization. General descriptions of using oligonucleotide arrays for detection of polymorphisms can be found, for example, in U.S. Pat. Nos. 5,858,659 and 5,837,832. In addition to oligonucleotide arrays, cDNA arrays can be used similarly in certain embodiments.

[0387] The methods described herein can include but are not limited to providing an array as described herein; contacting the array with a nucleic acid sample, and detecting binding of a nucleic acid from the nucleic acid sample to the array. In some embodiments, the method can comprise amplifying nucleic acid from the nucleic acid sample, for example, a region associated with PML or a region that includes another region associated with PML. In some embodiments, the methods described herein can include using an array that can identify differential expression patterns or copy numbers of one or more genes in nucleic acid samples from control and affected individuals. For example, arrays of probes to a marker described herein can be used to identify genetic variations between DNA from an affected subject, and control DNA obtained from an individual that does not have PML. Since the nucleotides on the array can contain sequence tags, their positions on the array can be accurately known relative to the genomic sequence.

[0388] In some embodiments, it can be desirable to employ methods that can detect the presence of multiple genetic variations, for example, polymorphic variants at a plurality of polymorphic sites, in parallel or substantially simultaneously. In some embodiments, these methods can comprise oligonucleotide arrays and other methods, including methods in which reactions, for example, amplification and hybridization, can be performed in individual vessels, for example, within individual wells of a multi-well plate or other vessel.

[0389] Determining the identity of a genetic variation can also include or consist of reviewing a subject's medical history, where the medical history includes information regarding the identity, copy number, presence or absence of one or more alleles or SNPs in the subject, e.g., results of a genetic test.

[0390] In some embodiments extended runs of homozygosity (ROH) may be useful to map recessive disease genes in outbred populations. Furthermore, even in complex disorders, a high number of affected individuals may have the same haplotype in the region surrounding a disease mutation. Therefore, a rare pathogenic variant and surrounding haplotype can be enriched in frequency in a group of affected individuals compared with the haplotype frequency in a cohort of unaffected controls. Homozygous haplotypes (HH) that are shared by multiple affected individuals can be important for the discovery of recessive disease genes in a condition such as PML. In some embodiments, the traditional homozygosity mapping method can be extended by analyzing the haplotype within shared ROH regions to identify homozygous segments of identical haplotype that are present uniquely or at a higher frequency in PML probands compared to parental controls. Such regions are termed risk homozygous haplotypes (rHH), which may contain low-frequency recessive variants that contribute to PML risk in a subset of PML patients.

[0391] Genetic variations can also be identified using any of a number of methods well known in the art. For example, genetic variations available in public databases, which can be searched using methods and custom algorithms or algo-

gorithms known in the art, can be used. In some embodiments, a reference sequence can be from, for example, the human draft genome sequence, publicly available in various databases, or a sequence deposited in a database such as GenBank.

[0392] A comparison of one or more genomes relative to one or more other genomes with array CGH, or a variety of other genetic variation detection methods, can reveal the set of genetic variations between two genomes, between one genome in comparison to multiple genomes, or between one set of genomes in comparison to another set of genomes. In some embodiments, an array CGH experiment can be performed by hybridizing a single test genome against a pooled nucleic acid sample of two or more genomes, which can result in minimizing the detection of higher frequency variants in the experiment. In some embodiments, a test genome can be hybridized alone (i.e., one-color detection) to a microarray, for example, using array CGH or SNP genotyping methods, and the comparison step to one or more reference genomes can be performed in silico to reveal the set of genetic variations in the test genome relative to the one or more reference genomes. In one embodiment, a single test genome is compared to a single reference genome in a 2-color experiment wherein both genomes are cohybridized to the microarray. In some embodiments, the whole genome or whole exome from one or more subjects is analyzed. In some embodiments, nucleic acid information has already been obtained for the whole genome or whole exome from one or more individuals and the nucleic acid information is obtained from in silico analysis.

[0393] Any of the polynucleotides described, including polynucleotides comprising a genetic variation, can be made synthetically using methods known in the art.

Methods of Detecting CNVs

[0394] Detection of genetic variations, specifically CNVs, can be accomplished by one or more suitable techniques described herein. Generally, techniques that can selectively determine whether a particular chromosomal segment is present or absent in an individual can be used for genotyping CNVs. Identification of novel copy number variations can be done by methods for assessing genomic copy number changes.

[0395] In some embodiments, methods include but are not limited to, methods that can quantitatively estimate the number of copies of a particular genomic segment, but can also include methods that indicate whether a particular segment is present in a nucleic acid sample or not. In some embodiments, the technique to be used can quantify the amount of segment present, for example, determining whether a DNA segment is deleted, duplicated, or triplicated in subject, for example, Fluorescent In Situ Hybridization (FISH) techniques, and other methods described herein. In some embodiments, methods include detection of copy number variation from array intensity and sequencing read depth using a stepwise Bayesian model (Zhang, et al., *BMC Bioinformatics*, 11:539 (2010)). In some embodiments, methods include detecting copy number variations using shotgun sequencing, CNV-seq (Xie C., et al., *BMC Bioinformatics*, 10:80 (2009)). In some embodiments, methods include analyzing next-generation sequencing (NGS) data for CNV detection using any one of several algorithms developed for each of the four broad methods for CNV detection using NGS, namely the depth of coverage (DOC),

read-pair (RP), split-read (SR) and assembly-based (AS) methods. (Teo et al., *Bioinformatics* (2012)). In some embodiments, methods include combining coverage with map information for the identification of deletions and duplications in targeted sequence data (Nord et al., *BMC Genomics*, 12:184 (2011)).

[0396] In some embodiments, other genotyping technologies can be used for detection of CNVs, including but not limited to, karyotype analysis, Molecular Inversion Probe array technology, for example, Affymetrix SNP Array 6.0, and BeadArray Technologies, for example, Illumina GoldenGate and Infinium assays, as can other platforms such as NimbleGen HD2.1 or HD4.2, High-Definition Comparative Genomic Hybridization (CGH) arrays (Agilent Technologies), tiling array technology (Affymetrix), multiplex ligation-dependent probe amplification (MLPA), Invader assay, fluorescence in situ hybridization, and, in one embodiment, Array Comparative Genomic Hybridization (aCGH) methods. As described herein, karyotype analysis can be a method to determine the content and structure of chromosomes in a nucleic acid sample. In some embodiments, karyotyping can be used, in lieu of aCGH, to detect translocations or inversions, which can be copy number neutral, and, therefore, not detectable by aCGH. Information about amplitude of particular probes, which can be representative of particular alleles, can provide quantitative dosage information for the particular allele, and by consequence, dosage information about the CNV in question, since the marker can be selected as a marker representative of the CNV and can be located within the CNV. In some embodiments, if the CNV is a deletion, the absence of particular marker allele is representative of the deletion. In some embodiments, if the CNV is a duplication or a higher order copy number variation, the signal intensity representative of the allele correlating with the CNV can represent the copy number. A summary of methodologies commonly used is provided in Perkel (Perkel J. *Nature Methods* 5:447-453 (2008)).

[0397] PCR assays can be utilized to detect CNVs and can provide an alternative to array analysis. In particular, PCR assays can enable detection of precise boundaries of gene/chromosome variants, at the molecular level, and which boundaries are identical in different individuals. PCR assays can be based on the amplification of a junction fragment present only in individuals that carry a deletion. This assay can convert the detection of a loss by array CGH to one of a gain by PCR.

[0398] Examples of PCR techniques that can be used in the present disclosure include, but are not limited to quantitative PCR, real-time quantitative PCR (qPCR), quantitative fluorescent PCR (QF-PCR), multiplex fluorescent PCR (MF-PCR), real time PCR (RT-PCR), single cell PCR, PCR-RFLP/RT-PCR-RFLP, hot start PCR and Nested PCR. Other suitable amplification methods include the ligase chain reaction (LCR), ligation mediated PCR (LM-PCR), degenerate oligonucleotide probe PCR (DOP-PCR), transcription amplification, self-sustained sequence replication, selective amplification of target polynucleotide sequences, consensus sequence primed polymerase chain reaction (CP-PCR), arbitrarily primed polymerase chain reaction (AP-PCR) and nucleic acid sequence based amplification (NASBA).

[0399] Alternative methods for the simultaneous interrogation of multiple regions include quantitative multiplex PCR of short fluorescent fragments (QMPSF), multiplex

amplifiable probe hybridization (MAPH) and multiplex ligation-dependent probe amplification (MLPA), in which copy-number differences for up to 40 regions can be scored in one experiment. Another approach can be to specifically target regions that harbor known segmental duplications, which are often sites of copy-number variation. By targeting the variable nucleotides between two copies of a segmental duplication (called paralogous sequence variants) using a SNP-genotyping method that provides independent fluorescence intensities for the two alleles, it is possible to detect an increase in intensity of one allele compared with the other.

[0400] In some embodiments, the amplified piece of DNA can be bound to beads using the sequencing element of the nucleic acid tag under conditions that favor a single amplified piece of DNA molecule to bind a different bead and amplification occurs on each bead. In some embodiments, such amplification can occur by PCR. Each bead can be placed in a separate well, which can be a picoliter-sized well. In some embodiments, each bead is captured within a droplet of a PCR-reaction-mixture-in-oil-emulsion and PCR amplification occurs within each droplet. The amplification on the bead results in each bead carrying at least one million, at least 5 million, or at least 10 million copies of the single amplified piece of DNA molecule.

[0401] In embodiments where PCR occurs in oil-emulsion mixtures, the emulsion droplets are broken, the DNA is denatured and the beads carrying single-stranded nucleic acids clones are deposited into a well, such as a picoliter-sized well, for further analysis according to the methods described herein. These amplification methods allow for the analysis of genomic DNA regions. Methods for using bead amplification followed by fiber optics detection are described in Margulies et al., *Nature*, 15; 437(7057):376-80 (2005), and as well as in US Publication Application Nos. 20020012930; 20030068629; 20030100102; 20030148344; 20040248161; 20050079510, 20050124022; and 20060078909.

[0402] Another variation on the array-based approach can be to use the hybridization signal intensities that are obtained from the oligonucleotides employed on Affymetrix SNP arrays or in Illumina Bead Arrays. Here hybridization intensities are compared with average values that are derived from controls, such that deviations from these averages indicate a change in copy number. As well as providing information about copy number, SNP arrays have the added advantage of providing genotype information. For example, they can reveal loss of heterozygosity, which could provide supporting evidence for the presence of a deletion, or might indicate segmental uniparental disomy (which can recapitulate the effects of structural variation in some genomic regions—Prader-Willi and Angelman syndromes, for example).

[0403] Many of the basic procedures followed in microarray-based genome profiling are similar, if not identical, to those followed in expression profiling and SNP analysis, including the use of specialized microarray equipment and data-analysis tools. Since microarray-based expression profiling has been well established in the last decade, much can be learned from the technical advances made in this area. Examples of the use of microarrays in nucleic acid analysis that can be used are described in U.S. Pat. Nos. 6,300,063, 5,837,832, 6,969,589, 6,040,138, 6,858,412, U.S. application Ser. No. 08/529,115, U.S. application Ser. No. 10/272,

384, U.S. application Ser. No. 10/045,575, U.S. application Ser. No. 10/264,571 and U.S. application Ser. No. 10/264,574. It should be noted that there are also distinct differences such as target and probe complexity, stability of DNA over RNA, the presence of repetitive DNA and the need to identify single copy number alterations in genome profiling.

[0404] In some embodiments, the genetic variations detected comprise CNVs and can be detected using array CGH. In some embodiments, array CGH can be implemented using a wide variety of techniques. The initial approaches used arrays produced from large-insert genomic clones such as bacterial artificial chromosomes (BACs). Producing sufficient BAC DNA of adequate purity to make arrays is arduous, so several techniques to amplify small amounts of starting material have been employed. These techniques include ligation-mediated PCR (Snijders et al., *Nat. Genet.* 29:263-64), degenerate primer PCR using one or several sets of primers, and rolling circle amplification. BAC arrays that provide complete genome tiling paths are also available. Arrays made from less complex nucleic acids such as cDNAs, selected PCR products, and oligonucleotides can also be used. Although most CGH procedures employ hybridization with total genomic DNA, it is possible to use reduced complexity representations of the genome produced by PCR techniques. Computational analysis of the genome sequence can be used to design array elements complementary to the sequences contained in the representation. Various SNP genotyping platforms, some of which use reduced complexity genomic representations, can be useful for their ability to determine both DNA copy number and allelic content across the genome. In some embodiments, small amounts of genomic DNA can be amplified with a variety of whole genome or whole exome amplification methods prior to CGH analysis of the nucleic acid sample. A “whole exome,” as used herein, includes exons throughout the whole genome that are expressed in genes. Since exon selection has tissue and cell type specificity, these positions may be different in the various cell types resulting from a splice variant or alternative splicing. A “whole genome,” as used herein, includes the entire genetic code of a genome.

[0405] The different basic approaches to array CGH provide different levels of performance, so some are more suitable for particular applications than others. The factors that determine performance include the magnitudes of the copy number changes, their genomic extents, the state and composition of the specimen, how much material is available for analysis, and how the results of the analysis can be used. Many applications use reliable detection of copy number changes of much less than 50%, a more stringent requirement than for other microarray technologies. Note that technical details are extremely important and different implementations of methods using the same array CGH approach can yield different levels of performance. Various CGH methods are known in the art and are equally applicable to one or more methods of the present disclosure. For example, CGH methods are disclosed in U.S. Pat. Nos. 7,030,231; 7,011,949; 7,014,997; 6,977,148; 6,951,761; and 6,916,621, the disclosure from each of which is incorporated by reference herein in its entirety.

[0406] The data provided by array CGH are quantitative measures of DNA sequence dosage. Array CGH provides high-resolution estimates of copy number aberrations, and can be performed efficiently on many nucleic acid samples. The advent of array CGH technology makes it possible to

monitor DNA copy number changes on a genomic scale and many projects have been launched for studying the genome in specific diseases.

[0407] In some embodiments, whole genome array-based comparative genome hybridization (array CGH) analysis, or array CGH on a subset of genomic regions, can be used to efficiently interrogate human genomes for genomic imbalances at multiple loci within a single assay. The development of comparative genomic hybridization (CGH) (Kallioniemi et al., *Science* 258: 818-21 (1992)) provided the first efficient approach to scanning entire genomes for variations in DNA copy number. The importance of normal copy number variation involving large segments of DNA has been unappreciated. Array CGH is a breakthrough technique in human genetics, which is attracting interest from clinicians working in fields as diverse as cancer and IVF (In Vitro Fertilization). The use of CGH microarrays in the clinic holds great promise for identifying regions of genomic imbalance associated with disease. Advances from identifying chromosomal critical regions associated with specific phenotypes to identifying the specific dosage sensitive genes can lead to therapeutic opportunities of benefit to patients. Array CGH is a specific, sensitive and rapid technique that can enable the screening of the whole genome in a single test. It can facilitate and accelerate the screening process in human genetics and is expected to have a profound impact on the screening and counseling of patients with genetic disorders. It is now possible to identify the exact location on the chromosome where an aberration has occurred and it is possible to map these changes directly onto the genomic sequence.

[0408] An array CGH approach provides a robust method for carrying out a genome-wide scan to find novel copy number variants (CNVs). The array CGH methods can use labeled fragments from a genome of interest, which can be competitively hybridized with a second differentially labeled genome to arrays that are spotted with cloned DNA fragments, revealing copy-number differences between the two genomes. Genomic clones (for example, BACs), cDNAs, PCR products and oligonucleotides, can all be used as array targets. The use of array CGH with BACs was one of the earliest employed methods and is popular, owing to the extensive coverage of the genome it provides, the availability of reliable mapping data and ready access to clones. The last of these factors is important both for the array experiments themselves, and for confirmatory FISH experiments.

[0409] In a typical CGH measurement, total genomic DNA is isolated from control and reference subjects, differentially labeled, and hybridized to a representation of the genome that allows the binding of sequences at different genomic locations to be distinguished. More than two genomes can be compared simultaneously with suitable labels. Hybridization of highly repetitive sequences is typically suppressed by the inclusion of unlabeled Cot-1 DNA in the reaction. In some embodiments of array CGH, it is beneficial to mechanically shear the genomic DNA in a nucleic acid sample, for example, with sonication, prior to its labeling and hybridization step. In another embodiment, array CGH may be performed without use of Cot-1 DNA or a sonication step in the preparation of the genomic DNA in a nucleic acid sample. The relative hybridization intensity of the test and reference signals at a given location can be proportional to the relative copy number of those sequences

in the test and reference genomes. If the reference genome is normal then increases and decreases in signal intensity ratios directly indicate DNA copy number variation within the genome of the test cells. Data are typically normalized so that the modal ratio for the genome is set to some standard value, typically 1.0 on a linear scale or 0.0 on a logarithmic scale. Additional measurements such as FISH or flow cytometry can be used to determine the actual copy number associated with a ratio level.

[0410] In some embodiments, an array CGH procedure can include the following steps. First, large-insert clones, for example, BACs can be obtained from a supplier of clone libraries. Then, small amounts of clone DNA can be amplified, for example, by degenerate oligonucleotide-primed (DOP) PCR or ligation-mediated PCR in order to obtain sufficient quantities needed for spotting. Next, PCR products can be spotted onto glass slides using, for example, microarray robots equipped with high-precision printing pins. Depending on the number of clones to be spotted and the space available on the microarray slide, clones can either be spotted once per array or in replicate. Repeated spotting of the same clone on an array can increase precision of the measurements if the spot intensities are averaged, and allows for a detailed statistical analysis of the quality of the experiments. Subject and control DNAs can be labeled, for example, with either Cy3 or Cy5-dUTP using random priming and can be subsequently hybridized onto the microarray in a solution containing an excess of Cot1-DNA to block repetitive sequences. Hybridizations can either be performed manually under a coverslip, in a gasket with gentle rocking or, automatically using commercially available hybridization stations. These automated hybridization stations can allow for an active hybridization process, thereby improving the reproducibility as well as reducing the actual hybridization time, which increases throughput. The hybridized DNAs can be detected through the two different fluorochromes using standard microarray scanning equipment with either a scanning confocal laser or a charge coupled device (CCD) camera-based reader, followed by spot identification using commercially or freely available software packages.

[0411] The use of CGH with arrays that comprise long oligonucleotides (60-100 bp) can improve the detection resolution (in some embodiments, as small as ~3-5 kb sized CNVs on arrays designed for interrogation of human whole genomes) over that achieved using BACs (limited to 50-100 kb or larger sized CNVs due to the large size of BAC clones). In some embodiments, the resolution of oligonucleotide CGH arrays is achieved via in situ synthesis of 1-2 million unique features/probes per microarray, which can include microarrays available from Roche NimbleGen and Agilent Technologies. In addition to array CGH methods for copy number detection, other embodiments for partial or whole genome analysis of CNVs within a genome include, but are not limited to, use of SNP genotyping microarrays and sequencing methods.

[0412] Another method for copy number detection that uses oligonucleotides can be representational oligonucleotide microarray analysis (ROMA). It is similar to that applied in the use of BAC and CGH arrays, but to increase the signal-to-noise ratio, the 'complexity' of the input DNA is reduced by a method called representation or whole-genome sampling. Here the DNA that is to be hybridized to the array can be treated by restriction digestion and then ligated to adapters, which results in the PCR-based ampli-

fication of fragments in a specific size-range. As a result, the amplified DNA can make up a fraction of the entire genomic sequence—that is, it is a representation of the input DNA that has significantly reduced complexity, which can lead to a reduction in background noise. Other suitable methods available to the skilled person can also be used, and are within scope of the present disclosure.

[0413] A comparison of one or more genomes relative to one or more other genomes with array CGH, or a variety of other CNV detection methods, can reveal the set of CNVs between two genomes, between one genome in comparison to multiple genomes, or between one set of genomes in comparison to another set of genomes. In some embodiments, an array CGH experiment can be performed by hybridizing a single test genome against a pooled nucleic acid sample of two or more genomes, which can result in minimizing the detection of higher frequency variants in the experiment. In some embodiments, a test genome can be hybridized alone (i.e. one-color detection) to a microarray, for example, using array CGH or SNP genotyping methods, and the comparison step to one or more reference genomes can be performed in silico to reveal the set of CNVs in the test genome relative to the one or more reference genomes. In one preferred embodiment, a single test genome is compared to a single reference genome in a 2-color experiment wherein both genomes are cohybridized to the microarray.

[0414] Array CGH can be used to identify genes that are causative or associated with a particular phenotype, condition, or disease by comparing the set of CNVs found in the affected cohort to the set of CNVs found in an unaffected cohort. An unaffected cohort may consist of any individual unaffected by the phenotype, condition, or disease of interest, but in one preferred embodiment is comprised of individuals or subjects that are apparently healthy (normal). Methods employed for such analyses are described in U.S. Pat. Nos. 7,702,468 and 7,957,913. In some embodiments of CNV comparison methods, candidate genes that are causative or associated (i.e. potentially serving as a biomarker) with a phenotype, condition, or disease will be identified by CNVs that occur in the affected cohort but not in the unaffected cohort. In some embodiments of CNV comparison methods, candidate genes that are causative or associated (i.e. potentially serving as a biomarker) with a phenotype, condition, or disease will be identified by CNVs that occur at a statistically significant higher frequency in the affected cohort as compared their frequency in the unaffected cohort. Thus, CNVs preferentially detected in the affected cohort as compared to the unaffected cohort can serve as beacons of genes that are causative or associated with a particular phenotype, condition, or disease. Methods employed for such analyses are described in U.S. Pat. No. 8,862,410. In some embodiments, CNV detection and comparison methods can result in direct identification of the gene that is causative or associated with phenotype, condition, or disease if the CNVs are found to overlap with or encompass the gene(s). In some embodiments, CNV detection and comparison methods can result in identification of regulatory regions of the genome (e.g., promoters, enhancers, transcription factor binding sites) that regulate the expression of one or more genes that are causative or associated with the phenotype, condition, or disease of interest. In some embodiments, CNV detection and comparison methods can result in identification of a region in the

genome in linkage disequilibrium with a genetic variant that is causative or associated with the phenotype, condition, or disease of interest. In another embodiment, CNV detection and comparison methods can result in identification of a region in the genome in linkage disequilibrium with a genetic variant that is protective against the condition or disease of interest.

[0415] Due to the large amount of genetic variation between any two genomes, or two sets (cohorts) of genomes, being compared, one preferred embodiment is to reduce the genetic variation search space by interrogating only CNVs, as opposed to the full set of genetic variants that can be identified in an individual's genome or exome. The set of CNVs that occur only, or at a statistically higher frequency, in the affected cohort as compared to the unaffected cohort can then be further investigated in targeted sequencing experiments to reveal the full set of genetic variants (of any size or type) that are causative or associated (i.e. potentially serving as a biomarker) with a phenotype, condition, or disease. It can be appreciated to those skilled in the art that the targeted sequencing experiments are performed in both the affected and unaffected cohorts in order to identify the genetic variants (e.g., SNVs and indels) that occur only, or at a statistically significant higher frequency, in the affected individual or cohort as compared to the unaffected cohort. Methods employed for such analyses are described in U.S. Pat. No. 8,862,410.

[0416] A method of screening a subject for a disease or disorder can comprise assaying a nucleic acid sample from the subject to detect sequence information for more than one genetic locus and comparing the sequence information to a panel of nucleic acid biomarkers and screening the subject for the presence or absence of the disease or disorder if one or more of low frequency biomarkers in the panel are present in the sequence information.

[0417] The panel can comprise at least one nucleic acid biomarker for each of the more than one genetic loci. For example, the panel can comprise 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 75, 100, 150, 200 or more nucleic acid biomarkers for each of the more than one genetic locus. In some embodiments, the panel can comprise from about 2-1000 nucleic acid biomarkers. For example, the panel can comprise from about 2-900, 2-800, 2-700, 2-600, 2-500, 2-400, 2-300, 2-200, 2-100, 25-900, 25-800, 25-700, 25-600, 25-500, 25-400, 25-300, 25-200, 25-100, 100-1000, 100-900, 100-800, 100-700, 100-600, 100-500, 100-400, 100-300, 100-200, 200-1000, 200-900, 200-800, 200-700, 200-600, 200-500, 200-400, 200-300, 300-1000, 300-900, 300-800, 300-700, 300-600, 300-500, 300-400, 400-1000, 400-900, 400-800, 400-700, 400-600, 400-500, 500-1000, 500-900, 500-800, 500-700, 500-600, 600-1000, 600-900, 600-800, 600-700, 700-1000, 700-900, 700-800, 800-1000, 800-900, or 900-1000 nucleic acid biomarkers.

[0418] In some embodiments, a biomarker can occur at a frequency of 1% or more in a population of subjects without a diagnosis of the disease or disorder. For example, a biomarker can occur at a frequency of 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, or more in a population of subjects without a diagnosis of the disease or disorder. In some embodiments, a biomarker can occur at a frequency from about 1%-20% in a population of subjects without a diagnosis of the disease or disorder. For example, a bio-

marker can occur at a frequency of from about 1%-5% or 1%-10%, in a population of subjects without a diagnosis of the disease or disorder.

[0419] The panel can comprise at least 2 low frequency biomarkers. For example, the panel can comprise at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 125, 150, 175, 200, 250, 500, or 1000 or more low frequency biomarkers. In some embodiments, the panel can comprise from about 2-1000 low frequency biomarkers. For example, the panel can comprise from about 2-900, 2-800, 2-700, 2-600, 2-500, 2-400, 2-300, 2-200, 2-100, 25-900, 25-800, 25-700, 25-600, 25-500, 25-400, 25-300, 25-200, 25-100, 100-1000, 100-900, 100-800, 100-700, 100-600, 100-500, 100-400, 100-300, 100-200, 200-1000, 200-900, 200-800, 200-700, 200-600, 200-500, 200-400, 200-300, 300-1000, 300-900, 300-800, 300-700, 300-600, 300-500, 300-400, 400-1000, 400-900, 400-800, 400-700, 400-600, 400-500, 500-1000, 500-900, 500-800, 500-700, 500-600, 600-1000, 600-900, 600-800, 600-700, 700-1000, 700-900, 700-800, 800-1000, 800-900, or 900-1000 low frequency biomarkers.

[0420] In some embodiments, a low frequency biomarker can occur at a frequency of 1% or less in a population of subjects without a diagnosis of the disease or disorder. For example, a low frequency biomarker can occur at a frequency of 0.5%, 0.1%, 0.05%, 0.01%, 0.005%, 0.001%, 0.0005%, or 0.0001% or less in a population of subjects without a diagnosis of the disease or disorder. In some embodiments, a low frequency biomarker can occur at a frequency from about 0.00010%-0.1% in a population of subjects without a diagnosis of the disease or disorder. For example, a low frequency biomarker can occur at a frequency of from about 0.0001%-0.0005%, 0.0001%-0.001%, 0.0001%-0.005%, 0.0001%-0.01%, 0.0001%-0.05%, 0.0001%-0.1%, 0.0001%-0.5%, 0.0005%-0.001%, 0.0005%-0.005%, 0.0005%-0.01%, 0.0005%-0.05%, 0.0005%-0.1%, 0.0005%-0.5%, 0.0005%-1%, 0.001%-0.005%, 0.001%-0.01%, 0.001%-0.05%, 0.001%-0.1%, 0.001%-0.5%, 0.001%-1%, 0.005%-0.01%, 0.005%-0.05%, 0.005%-0.1%, 0.005%-0.5%, 0.005%-1%, 0.01%-0.05%, 0.01%-0.1%, 0.01%-0.5%, 0.01%-1%, 0.05%-0.1%, 0.05%-0.5%, 0.05%-1%, 0.1%-0.5%, 0.1%-1%, or 0.5%-1% in a population of subjects without a diagnosis of the disease or disorder. In another embodiment, genetic biomarker frequencies can range higher (e.g., 0.5% to 5%) and have utility for diagnostic testing or drug development targeting the genes that harbor such variants. Genetic variants of appreciable frequency and phenotypic effect in the general population are sometimes described as goldilocks variants (e.g., see Cohen J Clin Lipidol. 2013 May-June; 7(3 Suppl):S1-5 and Price et al. Am J Hum Genet. 2010 Jun. 11; 86(6):832-8).

[0421] In some embodiments, the presence or absence of the disease or disorder in the subject can be determined with at least 50% confidence. For example, the presence or absence of the disease or disorder in the subject can be determined with at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99%, or 100% confidence. In some embodiments, the presence or absence of the disease or disorder in the subject can be determined with a 50%-100% confidence. For example, the presence or absence of the disease or disorder in the subject can be determined with a 60%-100%, 70%-100%, 80%-100%, 90%-100%, 50%-90%, 50%-80%, 50%-70%, 50%-60%, 60%-90%, 60%-

80%, 60%-70%, 70%-90%, 70%-80%, or 80%-90%. In one embodiment, PML candidate CNVs and genes or regulatory loci associated with these CNVs can be determined or identified by comparing genetic data from a cohort of normal individuals to that of an individual or a cohort of individuals known to have, or be susceptible to PML.

[0422] In one embodiment, PML candidate CNV-subregions and genes associated with these regions can be determined or identified by comparing genetic data from a cohort of normal individuals, such as a pre-existing database of CNVs found in normal individuals termed the Normal Variation Engine (NVE), to that of a cohort of individual known to have, or be susceptible to PML.

[0423] In some embodiments, a nucleic acid sample from one individual or nucleic acid samples from a pool of 2 or more individuals without PML can serve as the reference nucleic acid sample(s) and the nucleic acid sample from an individual known to have PML or being tested to determine if they have PML can serve as the test nucleic acid sample. In one preferred embodiment, the reference and test nucleic acid samples are sex-matched and co-hybridized on the CGH array. For example, reference nucleic acid samples can be labeled with a fluorophore such as Cy5, using methods described herein, and test subject nucleic acid samples can be labeled with a different fluorophore, such as Cy3. After labeling, nucleic acid samples can be combined and can be co-hybridized to a microarray and analyzed using any of the methods described herein, such as aCGH. Arrays can then be scanned and the data can be analyzed with software. Genetic alterations, such as CNVs, can be called using any of the methods described herein. A list of the genetic alterations, such as CNVs, can be generated for one or more test subjects and/or for one or more reference subjects. Such lists of CNVs can be used to generate a master list of non-redundant CNVs and/or CNV-subregions for each type of cohort. In one embodiment, a cohort of test nucleic acid samples, such as individuals known to have or suspected to have PML, can be cohybridized with an identical sex-matched reference individual or sex-matched pool of reference individuals to generate a list of redundant or non-redundant CNVs. Such lists can be based on the presence or absence of one or more CNVs and/or CNV subregions present in individuals within the cohort. In this manner, a master list can contain a number of distinct CNVs and/or CNV-subregions, some of which are uniquely present in a single individual and some of which are present in multiple individuals.

[0424] In some embodiments, CNVs and/or CNV-subregions of interest can be obtained by annotation of each CNV and/or CNV-subregion with relevant information, such as overlap with known genes and/or exons or intergenic regulatory regions such as transcription factor binding sites. In some embodiments, CNVs and/or CNV-subregions of interest can be obtained by calculating the OR for a CNV and/or CNV-subregion according to the following formula: $OR = (PML / ((\# \text{individuals in PML cohort}) - PML)) / (NVE / ((\# \text{individuals in NVE cohort}) - NVE))$, where: PML=number of PML individuals with a CNV-subregion of interest and NVE=number of NVE subjects with the CNV-subregion of interest. If NVE=0, it can be set to 1 to avoid dealing with infinities in cases where no CNVs are seen in the NVE. In some embodiments, a set of publicly available CNVs (e.g., the Database of Genomic Variants) can be used as the Normal cohort for comparison to the affected cohort CNVs. In another embodiment, the set of Normal cohort CNVs may

comprise a private database generated by the same CNV detection method, such as array CGH, or by a plurality of CNV detection methods that include, but are not limited to, array CGH, SNP genotyping arrays, custom CGH arrays, custom genotyping arrays, exome sequencing, whole genome sequencing, targeted sequencing, FISH, q-PCR, or MLPA.

[0425] The number of individuals in any given cohort can be at least about 10, 50, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2500, 5000, 7500, 10,000, 100,000, or more. In some embodiments, the number of individuals in any given cohort can be from 25-900, 25-800, 25-700, 25-600, 25-500, 25-400, 25-300, 25-200, 25-100, 100-1000, 100-900, 100-800, 100-700, 100-600, 100-500, 100-400, 100-300, 100-200, 200-1000, 200-900, 200-800, 200-700, 200-600, 200-500, 200-400, 200-300, 300-1000, 300-900, 300-800, 300-700, 300-600, 300-500, 300-400, 400-1000, 400-900, 400-800, 400-700, 400-600, 400-500, 500-1000, 500-900, 500-800, 500-700, 500-600, 600-1000, 600-900, 600-800, 600-700, 700-1000, 700-900, 700-800, 800-1000, 800-900, or 900-1000.

[0426] In some embodiments, a method of determining relevance or statistical significance of a genetic variant in a human subject to a disease or a condition associated with a genotype comprising screening a genome of a human subject with the disease or condition, such as by array Comparative Genomic Hybridization, sequencing, or SNP genotyping, to provide information on one or more genetic variants, such as those in Tables 1 and 2. The method can further comprise comparing, such as via a computer, information of said one or more genetic variants from the genome of said subject to a compilation of data comprising frequencies of genetic variants in at least 100 normal human subjects, such as those without the disease or condition. The method can further comprise determining a statistical significance or relevance of said one or more genetic variants from said comparison to the condition or disease or determining whether a genetic variant is present in said human subject but not present in said compilation of data from said comparison, or an algorithm can be used to call or identify significant genetic variations, such as a genetic variation whose median log 2 ratio is above or below a computed value. A computer can comprise computer executable logic that provides instructions for executing said comparison.

[0427] Different categories for CNVs of interest can be defined. In some embodiments, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions occur within intergenic regions and are associated with an OR of at least 0.7. For example, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions occur within intergenic regions and are associated with an OR of at least 0.7, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 175, or more. In some embodiments, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions occur within intergenic regions and are associated with an OR from about 0.7-200, 0.7-200, 0.7-90, 0.7-80, 0.7-70, 0.7-60, 0.7-50, 0.7-40, 0.7-30, 0.7-20, 0.7-10, 0.7-5, 10-200, 10-180, 10-160, 10-140, 10-120, 10-100, 10-80, 10-60, 10-40, 10-20, 20-200, 20-180, 20-160, 20-140, 20-120, 20-100, 20-80, 20-60, 20-40, 30-200, 30-180, 30-160, 30-140, 30-120, 30-100, 30-80, 30-60, 30-40, 40-200, 40-180, 40-160, 40-140, 40-120, 40-100, 40-90, 40-80, 40-70, 40-60, 40-50, 50-200, 50-180, 50-160,

50-140, 50-120, 50-100, 50-90, 50-80, 50-70, 50-60, 60-200, 60-180, 60-160, 60-140, 60-120, 60-100, 60-90, 60-80, 60-70, 70-200, 70-180, 70-160, 70-140, 70-120, 70-100, 70-90, 70-80, 80-200, 80-180, 80-160, 80-140, 80-120, 80-100, 80-90, 90-200, 90-180, 90-160, 90-140, 90-120, or 90-100.

[0428] In some embodiments, CNVs/CNV-subregions can be of interest if the CNV/CNV-subregion overlaps a known gene, and is associated with an OR of at least 1.8. For example, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions occur within intergenic regions and are associated with an OR of at least 1.8, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 175, or more. In some embodiments, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions occur within exonic regions and are associated with an OR from about 1.8-200, 1.8-200, 1.8-90, 1.8-80, 1.8-70, 1.8-60, 1.8-50, 1.8-40, 1.8-30, 1.8-20, 1.8-10, 1.8-5, 10-200, 10-180, 10-160, 10-140, 10-120, 10-100, 10-80, 10-60, 10-40, 10-20, 20-200, 20-180, 20-160, 20-140, 20-120, 20-100, 20-80, 20-60, 20-40, 30-200, 30-180, 30-160, 30-140, 30-120, 30-100, 30-80, 30-60, 30-40, 40-200, 40-180, 40-160, 40-140, 40-120, 40-100, 40-90, 40-80, 40-70, 40-60, 40-50, 50-200, 50-180, 50-160, 50-140, 50-120, 50-100, 50-90, 50-80, 50-70, 50-60, 60-200, 60-180, 60-160, 60-140, 60-120, 60-100, 60-90, 60-80, 60-70, 70-200, 70-180, 70-160, 70-140, 70-120, 70-100, 70-90, 70-80, 80-200, 80-180, 80-160, 80-140, 80-120, 80-100, 80-90, 90-200, 90-180, 90-160, 90-140, 90-120, or 90-100.

[0429] In some embodiments, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions are overlapping and/or non-overlapping, impact an exon, and they affect 1 or more PML cases but only 0 Normal subjects. In some embodiments, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions are overlapping and/or non-overlapping, impact an exon, and they affect 2 or more PML cases but only 0 or 1 Normal subjects. In some embodiments, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions are overlapping and/or non-overlapping, impact an exon, and they affect 1-5 PML cases but only 0 or 1 Normal subjects. For example, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions are overlapping and/or non-overlapping, impact an exon, and they affect 1 PML case but only 0 or 1 Normal subjects. This can enable identification of rarer CNVs in cases with PML. In some embodiments, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions are overlapping and/or non-overlapping, impact an exon, and they affect 1 PML case but only 0 or 1 Normal subjects, and are associated with an OR greater than 0.7, such as 1.8. In some embodiments, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions are overlapping and/or non-overlapping, impact an exon, and they affect 2 PML cases but only 0 or 1 Normal subjects. In some embodiments, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions are overlapping and/or non-overlapping, impact an exon, and they affect 3 PML cases but only 0 or 1 Normal subjects. In some embodiments, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions are overlapping and/or non-overlapping, impact an exon, and they affect 4 PML cases but only 0 or 1 Normal subjects.

[0430] In some embodiments, CNVs/CNV-subregions can be of interest if the OR associated with the sum of PML

cases and the sum of NVE subjects affecting the same gene (including distinct CNVs/CNV-subregions) is at least 0.67. For example, a CNV/CNV-subregion can be of interest if the OR associated with the sum of PML cases and the sum of NVE subjects affecting the same gene (including distinct CNVs/CNV-subregions) is at least 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 175, or more. In some embodiments, a CNVs/CNV-subregions can be of interest if the OR associated with the sum of PML cases and the sum of NVE subjects affecting the same gene (including distinct CNVs/CNV-subregions) is from about 0.7-200, 0.7-200, 0.7-90, 0.7-80, 0.7-70, 0.7-60, 0.7-50, 0.7-40, 0.7-30, 0.7-20, 0.7-10, 0.7-5, 10-200, 10-180, 10-160, 10-140, 10-120, 10-100, 10-80, 10-60, 10-40, 10-20, 20-200, 20-180, 20-160, 20-140, 20-120, 20-100, 20-80, 20-60, 20-40, 30-200, 30-180, 30-160, 30-140, 30-120, 30-100, 30-80, 30-60, 30-40, 40-200, 40-180, 40-160, 40-140, 40-120, 40-100, 40-90, 40-80, 40-70, 40-60, 40-50, 50-200, 50-180, 50-160, 50-140, 50-120, 50-100, 50-90, 50-80, 50-70, 50-60, 60-200, 60-180, 60-160, 60-140, 60-120, 60-100, 60-90, 60-80, 60-70, 70-200, 70-180, 70-160, 70-140, 70-120, 70-100, 70-90, 70-80, 80-200, 80-180, 80-160, 80-140, 80-120, 80-100, 80-90, 90-200, 90-180, 90-160, 90-140, 90-120, or 90-100.

[0431] In some embodiments, CNVs/CNV-subregions can be of interest if the OR associated with the sum of PML cases and the sum of NVE subjects affecting the same gene (including distinct CNVs/CNV-subregions) is at least 1.8. For example, a CNV/CNV-subregion can be of interest if the OR associated with the sum of PML cases and the sum of NVE subjects affecting the same gene (including distinct CNVs/CNV-subregions) is at least 1.8, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 175, or more. In some embodiments, a CNVs/CNV-subregions can be of interest if the OR associated with the sum of PML cases and the sum of NVE subjects affecting the same gene (including distinct CNVs/CNV-subregions) is from about 1.8-200, 1.8-200, 1.8-90, 1.8-80, 1.8-70, 1.8-60, 1.8-50, 1.8-40, 1.8-30, 1.8-20, 1.8-10, 1.8-5, 10-200, 10-180, 10-160, 10-140, 10-120, 10-100, 10-80, 10-60, 10-40, 10-20, 20-200, 20-180, 20-160, 20-140, 20-120, 20-100, 20-80, 20-60, 20-40, 30-200, 30-180, 30-160, 30-140, 30-120, 30-100, 30-80, 30-60, 30-40, 40-200, 40-180, 40-160, 40-140, 40-120, 40-100, 40-90, 40-80, 40-70, 40-60, 40-50, 50-200, 50-180, 50-160, 50-140, 50-120, 50-100, 50-90, 50-80, 50-70, 50-60, 60-200, 60-180, 60-160, 60-140, 60-120, 60-100, 60-90, 60-80, 60-70, 70-200, 70-180, 70-160, 70-140, 70-120, 70-100, 70-90, 70-80, 80-200, 80-180, 80-160, 80-140, 80-120, 80-100, 80-90, 90-200, 90-180, 90-160, 90-140, 90-120, or 90-100.

[0432] In some embodiments, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions do not overlap (distinct CNV/CNV-subregion), but impact the same gene (or regulatory locus) and are associated with an OR of at least 6 (Genic (distinct CNV-subregions); OR>6). For example, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions do not overlap, but impact the same gene (or regulatory locus), and are associated with an OR of at 7, 8, 9, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 45, 50, or more. In some embodiments, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions do not overlap, but impact the same gene (or regulatory locus), and are associ-

ated with an OR from about 6-100, 6-50, 6-40, 6-30, 6-20, 6-10, 6-9, 6-8, 6-7, 8-100, 8-50, 8-40, 8-30, 8-20, 8-10, 10-100, 10-50, 10-40, 10-30, 10-20, 20-100, 20-50, 20-40, 20-30, 30-100, 30-50, 30-40, 40-100, 40-50, 50-100, or 5-7. The CNV-subregion/gene can be an exonic or intronic part of the gene, or both.

[0433] In some embodiments, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions do not overlap a known gene (e.g., are non-genic or intergenic) and they are associated with an OR of at least 7 (Exon+ve, PML>4, NVE<2). For example, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregion does not overlap a known gene (e.g., is non-genic or intergenic) and/or non-overlapping, impact an exon, affect 2 or more PML cases but only 0 or 1 Normal subjects and are associated with an OR of at least 8, 9, 10, 11, 12, 14, 16, 18, 20, 25, 30, 35, 40, 45, 50, or more. In some embodiments, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions are overlapping and/or non-overlapping, impact an exon, affect 2 or more PML cases but only 0 or 1 Normal subjects and are associated with an OR from about 7-100, 7-50, 7-40, 7-30, 7-20, 20-100, 20-50, 20-40, 20-30, 30-100, 30-50, 30-40, 40-100, 40-50, 50-100, or 7-11.

[0434] In some embodiments, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions are overlapping and/or non-overlapping, impact an exon, and they affect 1-5 PML cases but only 0 or 1 Normal subjects. This can enable identification of rarer CNVs in cases with PML. In some embodiments, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions are overlapping and/or non-overlapping, impact an exon, and they affect 1 PML case but only 0 or 1 Normal subjects, and are associated with an OR greater than 1, such as 1.47, or from 1-2.5. In some embodiments, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions are overlapping and/or non-overlapping, impact an exon, and they affect 2 PML cases but only 0 or 1 Normal subjects and are associated with an OR greater than 2.5, such as 2.95, or from 2.5-4. In some embodiments, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions are overlapping and/or non-overlapping, impact an exon, and they affect 3 PML cases but only 0 or 1 Normal subjects and are associated with an OR greater than 4, such as 4.44, or from 4-5.5. In some embodiments, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions are overlapping and/or non-overlapping, impact an exon, and they affect 4 PML cases but only 0 or 1 Normal subjects and are associated with an OR greater than 5.5, such as 5.92, or from 5.5-6.8.

[0435] In some embodiments, CNVs/CNV-subregions can be of interest if the OR associated with the sum of PML cases and the sum of NVE subjects affecting the same gene (including distinct CNVs/CNV-subregions) is at least 6. For example, a CNV/CNV-subregion can be of interest if the OR associated with the sum of PML cases and the sum of NVE subjects affecting the same gene (including distinct CNVs/CNV-subregions) is at least 7, 8, 9, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 45, 50, or more. In some embodiments, a CNVs/CNV-subregions can be of interest if the OR associated with the sum of PML cases and the sum of NVE subjects affecting the same gene (including distinct CNVs/CNV-subregions) is from about 6-100, 6-50, 6-40, 6-30, 6-20, 6-10, 6-9, 6-8, 6-7, 8-100, 8-50, 8-40, 8-30, 8-20, 8-10, 10-100, 10-50, 10-40, 10-30, 10-20, 20-100, 20-50, 20-40, 20-30, 30-100, 30-50, 30-40, 40-100, 40-50, 50-100, or 5-7.

[0436] In some embodiments, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions impact an intron and they affect 5 or more PML cases but only 0 or 1 Normal subjects and they are associated with an OR of at least 7 (Intron+ve, PML>4, Normals<2). For example, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions impact an intron and they affect 5 or more PML cases but only 0 or 1 Normal subjects and they are associated with an OR of at least 8, 9, 10, 11, 12, 14, 16, 18, 20, 25, 30, 35, 40, 45, 50, or more. In some embodiments, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions impact an intron and they affect 5 or more PML cases but only 0 or 1 Normal subjects and they are associated with an OR from about 7-100, 7-50, 7-40, 7-30, 7-20, 20-100, 20-50, 20-40, 20-30, 30-100, 30-50, 30-40, 40-100, 40-50, 50-100, or 7-11. CNVs/CNV-subregions impacting introns can be pathogenic (e.g., such variants can result in alternatively spliced mRNAs or loss of a microRNA binding site, which may deleteriously impact the resulting protein's structure or expression level).

[0437] In some embodiments, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions occur within intergenic regions and are associated with an OR of greater than 30 (High OR intergenic (OR>30)). For example, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions occur within intergenic regions and are associated with an OR of greater than 31, 32, 33, 34, 35, 40, 45, 50, 66, 60, 65, 70, 75, 80, 85, 90, 95, 100 or more. In some embodiments, CNVs/CNV-subregions can be of interest if the CNVs/CNV-subregions impact occur within intergenic regions and are associated with an OR from about 30-100, 30-90, 30-80, 30-70, 30-60, 30-50, 30-40, 40-100, 40-90, 40-80, 40-70, 40-60, 40-50, 50-100, 50-90, 50-80, 50-70, 50-60, 60-100, 60-90, 60-80, 60-70, 70-100, 70-90, 70-80, 80-100, 80-90, or 90-100.

[0438] In some embodiments, a CNV/CNV-subregion can be of interest if the CNV/CNV-subregion overlaps a known gene, and is associated with an OR of at least 10. In some embodiments, a CNV/CNV-subregion can be of interest if the CNV/CNV-subregion overlaps a known gene, is associated with an OR of at least 6, and if the OR associated with the sum of PML cases and the sum of NVE subjects affecting the same gene (including distinct CNV-subregions) is at least 6.

Methods of Treatment

[0439] One embodiment of the present disclosure provides methods, pharmaceutical compositions, and kits for the treatment of a condition in animal subjects. The condition can be HIV/AIDS, cancer, or an autoimmune disease. In some embodiments, the condition can be PML. For example, the condition can be multiple sclerosis. In some embodiments, the methods comprise administering one or more immunosuppressive medications. In some embodiments, the pharmaceutical compositions and kits comprise one or more immunosuppressive medications. The one or more immunosuppressive medications can be adalimumab (e.g., Humira), alemtuzumab (e.g., Lemtrada), alemtuzumab (e.g., Campath), azathioprine (e.g., Imuran), belimumab (e.g., Benlysta), bevacizumab (e.g., Avastatin), bortezomib (e.g., Velcade), ecilizumab (e.g., Soliris), leflunomide, brentuximab vedotin (e.g., Adcetris), cetuximab (e.g., Erbitux), cyclophosphamid, dimethyl fumarate (e.g., Tecfidera), efalizumab (e.g., Raptiva), fingolimod (e.g., Gilenya), fludara-

bine (e.g., Fludara), fumaric acid, imatinib (e.g., Gleevec, Glivec), infliximab (e.g., Remicade), methotrexate (e.g., Trexall, Rheumatrex), mycophenolate mofetil (e.g., Cellcept), natalizumab (e.g., Tysabri), rituximab (e.g., Rituxin), daclizumab (e.g., Zinbryta), vedolizumab (Entyvio), ruxolitinib (e.g., Jakafi, Jakavi), ocrelizumab (e.g., Ocrevus), or any combinations thereof. The term "animal subject" as used herein includes humans as well as other mammals. The term "treating" as used herein includes achieving a therapeutic benefit and/or a prophylactic benefit. By therapeutic benefit is meant eradication or amelioration of the underlying viral infection (e.g., HIV), cancer, or autoimmune disease.

[0440] In some embodiments, a subject can be currently treated with an antiretroviral medication. In some embodiments, a subject can be previously treated with an antiretroviral medication. In some embodiments, a subject can be not yet treated with an antiretroviral medication. The antiretroviral medication can include but not limited to Nucleoside Reverse Transcriptase Inhibitors (NRTIs), Non-Nucleoside Reverse Transcriptase Inhibitors (NNRTIs), Protease Inhibitors (PIs), Fusion Inhibitors, Entry Inhibitors, Integrase Inhibitors, Pharmacokinetic Enhancers, and Combination HIV Medicines. In some cases, the Nucleoside Reverse Transcriptase Inhibitors can include but not limited to abacavir, didanosine, emtricitabine, lamivudine, stavudine, tenofovir disoproxil fumarate, and zidovudine. In some cases, the Non-Nucleoside Reverse Transcriptase Inhibitors can include but not limited to efavirenz, etravirine, nevirapine, and rilpivirine. In some cases, the Protease Inhibitors can include but not limited to atazanavir, darunavir, fosamprenavir, indinavir, nelfinavir, ritonavir, saquinavir, and tipranavir. In some cases, the Fusion Inhibitors can include but not limited to enfuvirtide. In some cases, the Entry Inhibitors can include but not limited to maraviroc. In some cases, the Integrase Inhibitors can include but not limited to dolutegravir, elvitegravir, and raltegravir. In some cases, the Pharmacokinetic Enhancers can include but not limited to cobicistat. In some cases, the Combination HIV Medicines can include but not limited to abacavir and lamivudine, abacavir, dolutegravir, and lamivudine, abacavir, lamivudine, and zidovudine, atazanavir and cobicistat, darunavir and cobicistat, efavirenz, emtricitabine, and tenofovir disoproxil fumarate, elvitegravir, cobicistat, emtricitabine, and tenofovir disoproxil fumarate, emtricitabine, rilpivirine, and tenofovir alafenamide, emtricitabine, rilpivirine, and tenofovir disoproxil fumarate, emtricitabine and tenofovir alafenamide, emtricitabine and tenofovir disoproxil fumarate, lamivudine and zidovudine, lopinavir and ritonavir, and any combination of antiretroviral medications listed above.

[0441] In some embodiments, such as when a subject is identified as having at least one of the genetic variants described herein, an agent targeting the JC Virus can be administered to the subject. In some embodiments, a medication can be administered to a subject that prevents PML from developing, or it can reduce, lessen, shorten and/or otherwise ameliorate the progression of PML, or symptoms that develop. The pharmaceutical composition can modulate or target JC Virus. In some embodiments, a subject identified as having PML can be administered an agent that reduces a viral load in the subject. In some embodiments, an immunosuppressive agent can be administered prior to, or in conjunction with, an agent that reduces a viral load in the

subject. In some embodiments, a subject identified as having a risk of developing PML can be administered an agent that prevents an increase in a viral load in the subject. In some embodiments, a subject identified as having a high risk of developing PML can be administered an agent that prevents an increase in a viral load in the subject. In some embodiments, an immunosuppressive agent can be administered prior to, or in conjunction with, an agent that prevents an increase in a viral load in the subject. The agent that reduces a viral load in the subject or that prevents an increase in a viral load in the subject can be, for example, an agent that targets JC Virus. Exemplary agents include antibodies, such as broadly neutralizing JCV antibodies. For example, an agent can be a broadly neutralizing human monoclonal JC polyomavirus VP-1 specific antibody (See, e.g., Jelcic et al., *Science Translational Medicine*, Vol. 7, Issue 306, pp. 306ra150 (2015) and Ray et al., *Science Translational Medicine*, Vol. 7, Issue 306, pp. 306ra151 (2015)). Additional exemplary agents include antiretroviral agents, cidofovir, hexadecyloxypropyl-cidofovir (a lipid-ester derivative), cytarabine (cytosine arabinoside), agents that block the 5HT_{2a} receptor (e.g., olanzapine, ziprasidone, mirtazapine, cyproheptadine, and risperidone), topoisomerase inhibitors (e.g., topotecan), and mefloquine.

[0442] In some embodiments, a pharmaceutical composition of the disclosure can be administered to a subject at risk of developing PML, or to a subject reporting one or more of the physiological symptoms of PML, even though a screening of the condition cannot have been made. In some embodiments, a pharmaceutical composition of the disclosure can be administered to a subject not identified as having a risk of developing PML, or to a subject not identified as having one or more of the physiological symptoms of PML, even though a screening of the condition cannot have been made.

[0443] The present disclosure also includes kits that can be used to treat a condition in animal subjects. These kits comprise one or more immunosuppressive medications and in some embodiments instructions teaching the use of the kit according to the various methods and approaches described herein. Such kits can also include information, such as scientific literature references, package insert materials, clinical trial results, and/or summaries of these and the like, which indicate or establish the activities and/or advantages (or risks and/or disadvantages) of the agent. Such information can be based on the results of various studies, for example, studies using experimental animals involving in vivo models and studies based on human clinical trials. Kits described herein can be provided, marketed and/or promoted to health providers, including physicians, nurses, pharmacists, formulary officials, and the like.

[0444] In some aspects a host cell can be used for testing or administering therapeutics. In some embodiments, a host cell can comprise a nucleic acid comprising expression control sequences operably-linked to a coding region. The host cell can be natural or non-natural. The non-natural host used in aspects of the method can be any cell capable of expressing a nucleic acid of the disclosure including, bacterial cells, fungal cells, insect cells, mammalian cells and plant cells. In some aspects the natural host is a mammalian tissue cell and the non-natural host is a different mammalian tissue cell. Other aspects of the method include a natural host that is a first cell normally residing in a first mammalian species and the non-natural host is a second cell normally

residing in a second mammalian species. In another alternative aspect, the method uses a first cell and the second cell that are from the same tissue type. In those aspects of the method where the coding region encodes a mammalian polypeptide, the mammalian polypeptide may be a hormone. In other aspects the coding region may encode a neuropeptide, an antibody, an antimetabolite, or a polypeptide or nucleotide therapeutic.

[0445] Expression control sequences can be those nucleotide sequences, both 5' and 3' to a coding region, that are required for the transcription and translation of the coding region in a host organism. Regulatory sequences include a promoter, ribosome binding site, optional inducible elements and sequence elements required for efficient 3' processing, including polyadenylation. When the structural gene has been isolated from genomic DNA, the regulatory sequences also include those intronic sequences required for splicing of the introns as part of mRNA formation in the target host.

Formulations, Routes of Administration, and Effective Doses

[0446] Yet another aspect of the present disclosure relates to formulations, routes of administration and effective doses for pharmaceutical compositions comprising an agent or combination of agents of the instant disclosure. Such pharmaceutical compositions can be used to treat a condition (e.g., multiple sclerosis) as described above.

[0447] Compounds of the disclosure can be administered as pharmaceutical formulations including those suitable for oral (including buccal and sub-lingual), rectal, nasal, topical, transdermal patch, pulmonary, vaginal, suppository, or parenteral (including intramuscular, intraarterial, intrathecal, intradermal, intraperitoneal, subcutaneous and intravenous) administration or in a form suitable for administration by aerosolization, inhalation or insufflation. General information on drug delivery systems can be found in Ansel et al., *Pharmaceutical Dosage Forms and Drug Delivery Systems* (Lippencott Williams & Wilkins, Baltimore Md. (1999)).

[0448] In various embodiments, the pharmaceutical composition includes carriers and excipients (including but not limited to buffers, carbohydrates, mannitol, polypeptides, amino acids, antioxidants, bacteriostats, chelating agents, suspending agents, thickening agents and/or preservatives), water, oils including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like, saline solutions, aqueous dextrose and glycerol solutions, flavoring agents, coloring agents, detackifiers and other acceptable additives, adjuvants, or binders, other pharmaceutically acceptable auxiliary substances to approximate physiological conditions, such as pH buffering agents, tonicity adjusting agents, emulsifying agents, wetting agents and the like. Examples of excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. In some embodiments, the pharmaceutical preparation is substantially free of preservatives. In other embodiments, the pharmaceutical preparation can contain at least one preservative. General methodology on pharmaceutical dosage forms is found in Ansel et al., *Pharmaceutical Dosage Forms and Drug Delivery Systems* (Lippencott, Williams, & Wilkins, Baltimore Md. (1999)). It can be recognized that, while any

suitable carrier known to those of ordinary skill in the art can be employed to administer the compositions of this disclosure, the type of carrier can vary depending on the mode of administration.

[0449] Compounds can also be encapsulated within liposomes using well-known technology. Biodegradable microspheres can also be employed as carriers for the pharmaceutical compositions of this disclosure. Suitable biodegradable microspheres are disclosed, for example, in U.S. Pat. Nos. 4,897,268, 5,075,109, 5,928,647, 5,811,128, 5,820,883, 5,853,763, 5,814,344 and 5,942,252.

[0450] The compound can be administered in liposomes or microspheres (or microparticles). Methods for preparing liposomes and microspheres for administration to a subject are well known to those of skill in the art. U.S. Pat. No. 4,789,734, the contents of which are hereby incorporated by reference, describes methods for encapsulating biological materials in liposomes. Essentially, the material is dissolved in an aqueous solution, the appropriate phospholipids and lipids added, and along with surfactants if required, and the material dialyzed or sonicated, as necessary. A review of known methods is provided by G. Gregoriadis, Chapter 14, "Liposomes," *Drug Carriers in Biology and Medicine*, pp. 2.sup.87-341 (Academic Press, 1979).

[0451] Microspheres formed of polymers or polypeptides are well known to those skilled in the art, and can be tailored for passage through the gastrointestinal tract directly into the blood stream. Alternatively, the compound can be incorporated and the microspheres, or composite of microspheres, implanted for slow release over a period of time ranging from days to months. See, for example, U.S. Pat. Nos. 4,906,474, 4,925,673 and 3,625,214, and Jain, *TIPS* 19:155-157 (1998), the contents of which are hereby incorporated by reference.

[0452] The concentration of drug can be adjusted, the pH of the solution buffered and the isotonicity adjusted to be compatible with intravenous injection, as is well known in the art.

[0453] The compounds of the disclosure can be formulated as a sterile solution or suspension, in suitable vehicles, well known in the art. The pharmaceutical compositions can be sterilized by conventional, well-known sterilization techniques, or can be sterile filtered. The resulting aqueous solutions can be packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile solution prior to administration. Suitable formulations and additional carriers are described in Remington "The Science and Practice of Pharmacy" (20th Ed., Lippincott Williams & Wilkins, Baltimore MD), the teachings of which are incorporated by reference in their entirety herein.

[0454] The agents or their pharmaceutically acceptable salts can be provided alone or in combination with one or more other agents or with one or more other forms. For example, a formulation can comprise one or more agents in particular proportions, depending on the relative potencies of each agent and the intended indication. For example, in compositions for targeting two different host targets, and where potencies are similar, about a 1:1 ratio of agents can be used. The two forms can be formulated together, in the same dosage unit e.g., in one cream, suppository, tablet, capsule, aerosol spray, or packet of powder to be dissolved in a beverage; or each form can be formulated in a separate unit, e.g., two creams, two suppositories, two tablets, two

capsules, a tablet and a liquid for dissolving the tablet, two aerosol sprays, or a packet of powder and a liquid for dissolving the powder, etc.

[0455] The term "pharmaceutically acceptable salt" means those salts which retain the biological effectiveness and properties of the agents used in the present disclosure, and which are not biologically or otherwise undesirable.

[0456] Typical salts are those of the inorganic ions, such as, for example, sodium, potassium, calcium, magnesium ions, and the like. Such salts include salts with inorganic or organic acids, such as hydrochloric acid, hydrobromic acid, phosphoric acid, nitric acid, sulfuric acid, methanesulfonic acid, p toluenesulfonic acid, acetic acid, fumaric acid, succinic acid, lactic acid, mandelic acid, malic acid, citric acid, tartaric acid or maleic acid. In addition, if the agent(s) contain a carboxy group or other acidic group, it can be converted into a pharmaceutically acceptable addition salt with inorganic or organic bases. Examples of suitable bases include sodium hydroxide, potassium hydroxide, ammonia, cyclohexylamine, dicyclohexylamine, ethanolamine, diethanolamine, triethanolamine, and the like.

[0457] A pharmaceutically acceptable ester or amide refers to those which retain biological effectiveness and properties of the agents used in the present disclosure, and which are not biologically or otherwise undesirable. Typical esters include ethyl, methyl, isobutyl, ethylene glycol, and the like. Typical amides include unsubstituted amides, alkyl amides, dialkyl amides, and the like.

[0458] In some embodiments, an agent can be administered in combination with one or more other compounds, forms, and/or agents, e.g., as described above. Pharmaceutical compositions with one or more other active agents can be formulated to comprise certain molar ratios. For example, molar ratios of about 99:1 to about 1:99 of a first active agent to the other active agent can be used. In some subset of the embodiments, the range of molar ratios of a first active agent: other active agents are selected from about 80:20 to about 20:80; about 75:25 to about 25:75, about 70:30 to about 30:70, about 66:33 to about 33:66, about 60:40 to about 40:60; about 50:50; and about 90:10 to about 10:90. The molar ratio of a first active: other active agents can be about 1:9, and in some embodiments can be about 1:1. The two agents, forms and/or compounds can be formulated together, in the same dosage unit e.g., in one cream, suppository, tablet, capsule, or packet of powder to be dissolved in a beverage; or each agent, form, and/or compound can be formulated in separate units, e.g., two creams, suppositories, tablets, two capsules, a tablet and a liquid for dissolving the tablet, an aerosol spray a packet of powder and a liquid for dissolving the powder, etc.

[0459] If necessary or desirable, the agents and/or combinations of agents can be administered with still other agents. The choice of agents that can be co-administered with the agents and/or combinations of agents of the instant disclosure can depend, at least in part, on the condition being treated. Agents of particular use in the formulations of the present disclosure include, for example, any agent having a therapeutic effect for a viral infection, including, e.g., drugs used to treat inflammatory conditions. For example, in treatments for influenza, in some embodiments formulations of the instant disclosure can additionally contain one or more conventional anti-inflammatory drugs, such as an NSAID, e.g., ibuprofen, naproxen, acetaminophen, ketoprofen, or aspirin. In some alternative embodiments for the

treatment of influenza formulations of the instant disclosure can additionally contain one or more conventional influenza antiviral agents, such as amantadine, rimantadine, zanamivir, and oseltamivir. In treatments for retroviral infections, such as HIV, formulations of the instant disclosure can additionally contain one or more conventional antiviral drug, such as protease inhibitors (lopinavir/ritonavir {Kaletra}, indinavir {Crixivan}, ritonavir {Norvir}, nelfinavir {Viracept}, saquinavir hard gel capsules {Invirase}, atazanavir {Reyataz}, amprenavir {Agenerase}, fosamprenavir {Telzir}, tipranavir {Aptivus}), reverse transcriptase inhibitors, including non-Nucleoside and Nucleoside/nucleotide inhibitors (AZT {zidovudine, Retrovir}, ddI {didanosine, Videx}, 3TC {lamivudine, Epivir}, d4T {stavudine, Zerit}, abacavir {Ziagen}, FTC {emtricitabine, Emtriva}, tenofovir {Viread}, efavirenz {Sustiva} and nevirapine {Viramune}), fusion inhibitors T20 {enfuvirtide, Fuzeon}, integrase inhibitors (MK-0518 and GS-9137), and maturation inhibitors (PA-457 {Bevirimat}). As another example, formulations can additionally contain one or more supplements, such as vitamin C, E or other anti-oxidants.

[0460] The agent(s) (or pharmaceutically acceptable salts, esters or amides thereof) can be administered per se or in the form of a pharmaceutical composition wherein the active agent(s) is in an admixture or mixture with one or more pharmaceutically acceptable carriers. A pharmaceutical composition, as used herein, can be any composition prepared for administration to a subject. Pharmaceutical compositions for use in accordance with the present disclosure can be formulated in conventional manner using one or more physiologically acceptable carriers, comprising excipients, diluents, and/or auxiliaries, e.g., which facilitate processing of the active agents into preparations that can be administered. Proper formulation can depend at least in part upon the route of administration chosen. The agent(s) useful in the present disclosure, or pharmaceutically acceptable salts, esters, or amides thereof, can be delivered to a subject using a number of routes or modes of administration, including oral, buccal, topical, rectal, transdermal, transmucosal, subcutaneous, intravenous, and intramuscular applications, as well as by inhalation.

[0461] For oral administration, the agents can be formulated readily by combining the active agent(s) with pharmaceutically acceptable carriers well known in the art. Such carriers enable the agents of the disclosure to be formulated as tablets, including chewable tablets, pills, dragees, capsules, lozenges, hard candy, liquids, gels, syrups, slurries, powders, suspensions, elixirs, wafers, and the like, for oral ingestion by a subject to be treated. Such formulations can comprise pharmaceutically acceptable carriers including solid diluents or fillers, sterile aqueous media and various non-toxic organic solvents. A solid carrier can be one or more substances which can also act as diluents, flavoring agents, solubilizers, lubricants, suspending agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material. In powders, the carrier generally is a finely divided solid which is a mixture with the finely divided active component. In tablets, the active component generally is mixed with the carrier having the necessary binding capacity in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain from about one (1) to about seventy (70) percent of the active compound. Suitable carriers include but are not limited to magnesium carbonate, magnesium stearate, talc,

sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. Generally, the agents of the disclosure can be included at concentration levels ranging from about 0.5%, about 5%, about 10%, about 20%, or about 30% to about 50%, about 60%, about 70%, about 80% or about 90% by weight of the total composition of oral dosage forms, in an amount sufficient to provide a desired unit of dosage.

[0462] Aqueous suspensions for oral use can contain agent (s) of this disclosure with pharmaceutically acceptable excipients, such as a suspending agent (e.g., methyl cellulose), a wetting agent (e.g., lecithin, lysolecithin and/or a long-chain fatty alcohol), as well as coloring agents, preservatives, flavoring agents, and the like.

[0463] In some embodiments, oils or non-aqueous solvents can be used to bring the agents into solution, due to, for example, the presence of large lipophilic moieties. Alternatively, emulsions, suspensions, or other preparations, for example, liposomal preparations, can be used. With respect to liposomal preparations, any known methods for preparing liposomes for treatment of a condition can be used. See, for example, Bangham et al., *J. Mol. Biol.* 23: 238-252 (1965) and Szoka et al., *Proc. Natl Acad. Sci. USA* 75: 4194-4198 (1978), incorporated herein by reference. Ligands can also be attached to the liposomes to direct these compositions to particular sites of action. Agents of this disclosure can also be integrated into foodstuffs, e.g., cream cheese, butter, salad dressing, or ice cream to facilitate solubilization, administration, and/or compliance in certain subject populations.

[0464] Pharmaceutical preparations for oral use can be obtained as a solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; flavoring elements, cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl cellulose, sodium carboxymethylcellulose, and/or polyvinyl pyrrolidone (PVP). If desired, disintegrating agents can be added, such as the cross linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate. The agents can also be formulated as a sustained release preparation.

[0465] Dragee cores can be provided with suitable coatings. For this purpose, concentrated sugar solutions can be used, which can optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments can be added to the tablets or dragee coatings for identification or to characterize different combinations of active agents.

[0466] Pharmaceutical preparations that can be used orally include push fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active agents can be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers can be added.

All formulations for oral administration should be in dosages suitable for administration.

[0467] Other forms suitable for oral administration include liquid form preparations including emulsions, syrups, elixirs, aqueous solutions, aqueous suspensions, or solid form preparations which are intended to be converted shortly before use to liquid form preparations. Emulsions can be prepared in solutions, for example, in aqueous propylene glycol solutions or can contain emulsifying agents, for example, such as lecithin, sorbitan monooleate, or acacia. Aqueous solutions can be prepared by dissolving the active component in water and adding suitable colorants, flavors, stabilizers, and thickening agents. Aqueous suspensions can be prepared by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well known suspending agents. Suitable fillers or carriers with which the compositions can be administered include agar, alcohol, fats, lactose, starch, cellulose derivatives, polysaccharides, polyvinylpyrrolidone, silica, sterile saline and the like, or mixtures thereof used in suitable amounts. Solid form preparations include solutions, suspensions, and emulsions, and can contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

[0468] A syrup or suspension can be made by adding the active compound to a concentrated, aqueous solution of a sugar, e.g., sucrose, to which can also be added any accessory ingredients. Such accessory ingredients can include flavoring, an agent to retard crystallization of the sugar or an agent to increase the solubility of any other ingredient, e.g., as a polyhydric alcohol, for example, glycerol or sorbitol.

[0469] When formulating compounds of the disclosure for oral administration, it can be desirable to utilize gastroretentive formulations to enhance absorption from the gastrointestinal (GI) tract. A formulation which is retained in the stomach for several hours can release compounds of the disclosure slowly and provide a sustained release that can be preferred in some embodiments of the disclosure. Disclosure of such gastro-retentive formulations are found in Klausner E. A., et al., *Pharm. Res.* 20, 1466-73 (2003); Hoffman, A. et al., *Int. J. Pharm.* 11, 141-53 (2004), Streubel, A., et al. *Expert Opin. Drug Deliver.* 3, 217-3, and Chavanpatil, M. D. et al., *Int. J. Pharm.* (2006). Expandable, floating and bioadhesive techniques can be utilized to maximize absorption of the compounds of the disclosure.

[0470] The compounds of the disclosure can be formulated for parenteral administration (e.g., by injection, for example, bolus injection or continuous infusion) and can be presented in unit dose form in ampoules, pre-filled syringes, small volume infusion or in multi-dose containers with an added preservative. The compositions can take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, for example, solutions in aqueous polyethylene glycol.

[0471] For injectable formulations, the vehicle can be chosen from those known in art to be suitable, including aqueous solutions or cotton suspensions, or emulsions, with sesame oil, corn oil, cottonseed oil, or peanut oil, as well as elixirs, mannitol, dextrose, or a sterile aqueous solution, and similar pharmaceutical vehicles. The formulation can also comprise polymer compositions which are biocompatible, biodegradable, such as poly(lactic-co-glycolic)acid. These

materials can be made into micro or nanospheres, loaded with drug and further coated or derivatized to provide superior sustained release performance. Vehicles suitable for periocular or intraocular injection include, for example, suspensions of therapeutic agent in injection grade water, liposomes and vehicles suitable for lipophilic substances. Other vehicles for periocular or intraocular injection are well known in the art.

[0472] In some embodiments, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition can also include a solubilizing agent and a local anesthetic such as lidocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients can be mixed prior to administration.

[0473] When administration is by injection, the active compound can be formulated in aqueous solutions, specifically in physiologically compatible buffers such as Hanks solution, Ringer's solution, or physiological saline buffer. The solution can contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active compound can be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use. In some embodiments, the pharmaceutical composition does not comprise an adjuvant or any other substance added to enhance the immune response stimulated by the peptide. In some embodiments, the pharmaceutical composition comprises a substance that inhibits an immune response to the peptide. Methods of formulation are known in the art, for example, as disclosed in Remington's *Pharmaceutical Sciences*, latest edition, Mack Publishing Co., Easton P.

[0474] In addition to the formulations described previously, the agents can also be formulated as a depot preparation. Such long acting formulations can be administered by implantation or transcutaneous delivery (for example, subcutaneously or intramuscularly), intramuscular injection or use of a transdermal patch. Thus, for example, the agents can be formulated with suitable polymeric or hydrophobic materials (for example, as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

[0475] In some embodiments, pharmaceutical compositions comprising one or more agents of the present disclosure exert local and regional effects when administered topically or injected at or near particular sites of infection. Direct topical application, e.g., of a viscous liquid, solution, suspension, dimethylsulfoxide (DMSO)-based solutions, liposomal formulations, gel, jelly, cream, lotion, ointment, suppository, foam, or aerosol spray, can be used for local administration, to produce for example, local and/or regional effects. Pharmaceutically appropriate vehicles for such formulation include, for example, lower aliphatic alcohols,

polyglycols (e.g., glycerol or polyethylene glycol), esters of fatty acids, oils, fats, silicones, and the like. Such preparations can also include preservatives (e.g., p-hydroxybenzoic acid esters) and/or antioxidants (e.g., ascorbic acid and tocopherol). See also *Dermatological Formulations: Percutaneous absorption*, Barry (Ed.), Marcel Dekker Incl, 1983.

[0476] Pharmaceutical compositions of the present disclosure can contain a cosmetically or dermatologically acceptable carrier. Such carriers are compatible with skin, nails, mucous membranes, tissues and/or hair, and can include any conventionally used cosmetic or dermatological carrier meeting these requirements. Such carriers can be readily selected by one of ordinary skill in the art. In formulating skin ointments, an agent or combination of agents of the instant disclosure can be formulated in an oleaginous hydrocarbon base, an anhydrous absorption base, a water-in-oil absorption base, an oil-in-water water-removable base and/or a water-soluble base. Examples of such carriers and excipients include, but are not limited to, humectants (e.g., urea), glycols (e.g., propylene glycol), alcohols (e.g., ethanol), fatty acids (e.g., oleic acid), surfactants (e.g., isopropyl myristate and sodium lauryl sulfate), pyrrolidones, glycerol monolaurate, sulfonates, terpenes (e.g., menthol), amines, amides, alkanes, alkanols, water, calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols.

[0477] Ointments and creams can, for example, be formulated with an aqueous or oily base with the addition of suitable thickening and/or gelling agents. Lotions can be formulated with an aqueous or oily base and can in general also contain one or more emulsifying agents, stabilizing agents, dispersing agents, suspending agents, thickening agents, or coloring agents. The construction and use of transdermal patches for the delivery of pharmaceutical agents is well known in the art. See, e.g., U.S. Pat. Nos. 5,023,252, 4,992,445 and 5,001,139. Such patches can be constructed for continuous, pulsatile, or on demand delivery of pharmaceutical agents.

[0478] Lubricants which can be used to form pharmaceutical compositions and dosage forms of the disclosure include, but are not limited to, calcium stearate, magnesium stearate, mineral oil, light mineral oil, glycerin, sorbitol, mannitol, polyethylene glycol, other glycols, stearic acid, sodium lauryl sulfate, talc, hydrogenated vegetable oil (e.g., peanut oil, cottonseed oil, sunflower oil, sesame oil, olive oil, corn oil, and soybean oil), zinc stearate, ethyl oleate, ethyl laureate, agar, or mixtures thereof. Additional lubricants include, for example, a syloid silica gel, a coagulated aerosol of synthetic silica, or mixtures thereof. A lubricant can optionally be added, in an amount of less than about 1 weight percent of the pharmaceutical composition.

[0479] The compositions according to the present disclosure can be in any form suitable for topical application, including aqueous, aqueous-alcoholic or oily solutions, lotion or serum dispersions, aqueous, anhydrous or oily gels, emulsions obtained by dispersion of a fatty phase in an aqueous phase (O/W or oil in water) or, conversely, (W/O or water in oil), microemulsions or alternatively microcapsules, microparticles or lipid vesicle dispersions of ionic and/or nonionic type. These compositions can be prepared according to conventional methods. Other than the agents of the disclosure, the amounts of the various constituents of the compositions according to the disclosure are those conventionally used in the art. These compositions in particular

constitute protection, treatment or care creams, milks, lotions, gels or foams for the face, for the hands, for the body and/or for the mucous membranes, or for cleansing the skin. The compositions can also consist of solid preparations constituting soaps or cleansing bars.

[0480] Compositions of the present disclosure can also contain adjuvants common to the cosmetic and dermatological fields, such as hydrophilic or lipophilic gelling agents, hydrophilic or lipophilic active agents, preserving agents, antioxidants, solvents, fragrances, fillers, sunscreens, odor-absorbers and dyestuffs. The amounts of these various adjuvants are those conventionally used in the fields considered and, for example, are from about 0.010% to about 20% of the total weight of the composition. Depending on their nature, these adjuvants can be introduced into the fatty phase, into the aqueous phase and/or into the lipid vesicles.

[0481] In some embodiments, ocular viral infections can be effectively treated with ophthalmic solutions, suspensions, ointments or inserts comprising an agent or combination of agents of the present disclosure. Eye drops can be prepared by dissolving the active ingredient in a sterile aqueous solution such as physiological saline, buffering solution, etc., or by combining powder compositions to be dissolved before use. Other vehicles can be chosen, as is known in the art, including but not limited to: balance salt solution, saline solution, water soluble polyethers such as polyethylene glycol, polyvinyls, such as polyvinyl alcohol and povidone, cellulose derivatives such as methylcellulose and hydroxypropyl methylcellulose, petroleum derivatives such as mineral oil and white petrolatum, animal fats such as lanolin, polymers of acrylic acid such as carboxypolyethylene gel, vegetable fats such as peanut oil and polysaccharides such as dextrans, and glycosaminoglycans such as sodium hyaluronate. If desired, additives ordinarily used in the eye drops can be added. Such additives include isotonicizing agents (e.g., sodium chloride, etc.), buffer agent (e.g., boric acid, sodium monohydrogen phosphate, sodium dihydrogen phosphate, etc.), preservatives (e.g., benzalkonium chloride, benzethonium chloride, chlorobutanol, etc.), thickeners (e.g., saccharide such as lactose, mannitol, maltose, etc.; e.g., hyaluronic acid or its salt such as sodium hyaluronate, potassium hyaluronate, etc.; e.g., mucopolysaccharide such as chondroitin sulfate, etc.; e.g., sodium polyacrylate, carboxyvinyl polymer, crosslinked polyacrylate, polyvinyl alcohol, polyvinyl pyrrolidone, methyl cellulose, hydroxypropyl methylcellulose, hydroxyethyl cellulose, carboxymethyl cellulose, hydroxypropyl cellulose or other agents known to those skilled in the art).

[0482] The solubility of the components of the present compositions can be enhanced by a surfactant or other appropriate co-solvent in the composition. Such cosolvents include polysorbate 20, 60, and 80, Pluronic F68, F-84 and P-103, cyclodextrin, or other agents known to those skilled in the art. Such cosolvents can be employed at a level of from about 0.010% to 2% by weight.

[0483] The compositions of the disclosure can be packaged in multidose form. Preservatives can be preferred to prevent microbial contamination during use. Suitable preservatives include: benzalkonium chloride, thimerosal, chlorobutanol, methyl paraben, propyl paraben, phenylethyl alcohol, edetate disodium, sorbic acid, Onamer M, or other agents known to those skilled in the art. In the prior art ophthalmic products, such preservatives can be employed at a level of from 0.004% to 0.02%. In the compositions of the

present application the preservative, preferably benzalkonium chloride, can be employed at a level of from 0.001% to less than 0.01%, e.g., from 0.001% to 0.008%, preferably about 0.005% by weight. It has been found that a concentration of benzalkonium chloride of 0.005% can be sufficient to preserve the compositions of the present disclosure from microbial attack.

[0484] In some embodiments, the agents of the present disclosure are delivered in soluble rather than suspension form, which allows for more rapid and quantitative absorption to the sites of action. In general, formulations such as jellies, creams, lotions, suppositories and ointments can provide an area with more extended exposure to the agents of the present disclosure, while formulations in solution, e.g., sprays, provide more immediate, short-term exposure.

[0485] In some embodiments relating to topical/local application, the pharmaceutical compositions can include one or more penetration enhancers. For example, the formulations can comprise suitable solid or gel phase carriers or excipients that increase penetration or help delivery of agents or combinations of agents of the disclosure across a permeability barrier, e.g., the skin. Many of these penetration-enhancing compounds are known in the art of topical formulation, and include, e.g., water, alcohols (e.g., terpenes like methanol, ethanol, 2-propanol), sulfoxides (e.g., dimethyl sulfoxide, decylmethyl sulfoxide, tetradecylmethyl sulfoxide), pyrrolidones (e.g., 2-pyrrolidone, N-methyl-2-pyrrolidone, N-(2-hydroxyethyl)pyrrolidone), laurocapram, acetone, dimethylacetamide, dimethylformamide, tetrahydrofurfuryl alcohol, L- α -amino acids, anionic, cationic, amphoteric or nonionic surfactants (e.g., isopropyl myristate and sodium lauryl sulfate), fatty acids, fatty alcohols (e.g., oleic acid), amines, amides, clofibric acid amides, hexamethylene lauramide, proteolytic enzymes, α -bisabolol, d-limonene, urea and N,N-diethyl-m-toluamide, and the like. Additional examples include humectants (e.g., urea), glycols (e.g., propylene glycol and polyethylene glycol), glycerol monolaurate, alkanes, alkanols, ORGELASE, calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and/or other polymers. In some embodiments, the pharmaceutical compositions can include one or more such penetration enhancers.

[0486] In some embodiments, the pharmaceutical compositions for local/topical application can include one or more antimicrobial preservatives such as quaternary ammonium compounds, organic mercurials, p-hydroxy benzoates, aromatic alcohols, chlorobutanol, and the like.

[0487] In some embodiments, the pharmaceutical compositions can be orally- or rectally delivered solutions, suspensions, ointments, enemas and/or suppositories comprising an agent or combination of agents of the present disclosure.

[0488] In some embodiments, the pharmaceutical compositions can be aerosol solutions, suspensions or dry powders comprising an agent or combination of agents of the present disclosure. The aerosol can be administered through the respiratory system or nasal passages. For example, one skilled in the art can recognize that a composition of the present disclosure can be suspended or dissolved in an appropriate carrier, e.g., a pharmaceutically acceptable propellant, and administered directly into the lungs using a nasal spray or inhalant. For example, an aerosol formulation comprising an agent can be dissolved, suspended or emulsified in a propellant or a mixture of solvent and propellant,

e.g., for administration as a nasal spray or inhalant. Aerosol formulations can contain any acceptable propellant under pressure, such as a cosmetically or dermatologically or pharmaceutically acceptable propellant, as conventionally used in the art.

[0489] An aerosol formulation for nasal administration is generally an aqueous solution designed to be administered to the nasal passages in drops or sprays. Nasal solutions can be similar to nasal secretions in that they are generally isotonic and slightly buffered to maintain a pH of about 5.5 to about 6.5, although pH values outside of this range can additionally be used. Antimicrobial agents or preservatives can also be included in the formulation.

[0490] An aerosol formulation for inhalations and inhalants can be designed so that the agent or combination of agents of the present disclosure is carried into the respiratory tree of the subject when administered by the nasal or oral respiratory route. Inhalation solutions can be administered, for example, by a nebulizer. Inhalations or insufflations, comprising finely powdered or liquid drugs, can be delivered to the respiratory system as a pharmaceutical aerosol of a solution or suspension of the agent or combination of agents in a propellant, e.g., to aid in disbursement. Propellants can be liquefied gases, including halocarbons, for example, fluorocarbons such as fluorinated chlorinated hydrocarbons, hydrochlorofluorocarbons, and hydrochlorocarbons, as well as hydrocarbons and hydrocarbon ethers.

[0491] Halocarbon propellants useful in the present disclosure include fluorocarbon propellants in which all hydrogens are replaced with fluorine, chlorofluorocarbon propellants in which all hydrogens are replaced with chlorine and at least one fluorine, hydrogen-containing fluorocarbon propellants, and hydrogen-containing chlorofluorocarbon propellants. Halocarbon propellants are described in Johnson, U.S. Pat. No. 5,376,359; Byron et al., U.S. Pat. No. 5,190,029; and Purewal et al., U.S. Pat. No. 5,776,434. Hydrocarbon propellants useful in the disclosure include, for example, propane, isobutane, n-butane, pentane, isopentane and neopentane. A blend of hydrocarbons can also be used as a propellant. Ether propellants include, for example, dimethyl ether as well as the ethers. An aerosol formulation of the disclosure can also comprise more than one propellant. For example, the aerosol formulation can comprise more than one propellant from the same class, such as two or more fluorocarbons; or more than one, more than two, more than three propellants from different classes, such as a fluorohydrocarbon and a hydrocarbon. Pharmaceutical compositions of the present disclosure can also be dispensed with a compressed gas, e.g., an inert gas such as carbon dioxide, nitrous oxide or nitrogen.

[0492] Aerosol formulations can also include other components, for example, ethanol, isopropanol, propylene glycol, as well as surfactants or other components such as oils and detergents. These components can serve to stabilize the formulation and/or lubricate valve components.

[0493] The aerosol formulation can be packaged under pressure and can be formulated as an aerosol using solutions, suspensions, emulsions, powders and semisolid preparations. For example, a solution aerosol formulation can comprise a solution of an agent of the disclosure in (substantially) pure propellant or as a mixture of propellant and solvent. The solvent can be used to dissolve the agent and/or retard the evaporation of the propellant. Solvents useful in the disclosure include, for example, water, ethanol and

glycols. Any combination of suitable solvents can be used, optionally combined with preservatives, antioxidants, and/or other aerosol components.

[0494] An aerosol formulation can also be a dispersion or suspension. A suspension aerosol formulation can comprise a suspension of an agent or combination of agents of the instant disclosure. Dispersing agents useful in the disclosure include, for example, sorbitan trioleate, oleyl alcohol, oleic acid, lecithin and corn oil. A suspension aerosol formulation can also include lubricants, preservatives, antioxidant, and/or other aerosol components.

[0495] An aerosol formulation can similarly be formulated as an emulsion. An emulsion aerosol formulation can include, for example, an alcohol such as ethanol, a surfactant, water and a propellant, as well as an agent or combination of agents of the disclosure. The surfactant used can be nonionic, anionic or cationic. One example of an emulsion aerosol formulation comprises, for example, ethanol, surfactant, water and propellant. Another example of an emulsion aerosol formulation comprises, for example, vegetable oil, glyceryl monostearate and propane.

[0496] The compounds of the disclosure can be formulated for administration as suppositories. A low melting wax, such as a mixture of triglycerides, fatty acid glycerides, Witapol S55 (trademark of Dynamite Nobel Chemical, Germany), or cocoa butter is first melted and the active component is dispersed homogeneously, for example, by stirring. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool, and to solidify.

[0497] The compounds of the disclosure can be formulated for vaginal administration. Pessaries, tampons, creams, gels, pastes, foams or sprays containing in addition to the active ingredient such carriers as are known in the art to be appropriate.

[0498] It is envisioned additionally, that the compounds of the disclosure can be attached releasably to biocompatible polymers for use in sustained release formulations on, in or attached to inserts for topical, intraocular, periocular, or systemic administration. The controlled release from a biocompatible polymer can be utilized with a water soluble polymer to form an instillable formulation, as well. The controlled release from a biocompatible polymer, such as for example, PLGA microspheres or nanospheres, can be utilized in a formulation suitable for intra ocular implantation or injection for sustained release administration, as well as any suitable biodegradable and biocompatible polymer can be used.

[0499] In one aspect of the disclosure, the subject's carrier status of any of the genetic variation risk variants described herein, or genetic variants identified via other analysis methods within the genes or regulatory loci that are identified by the CNVs or SNVs described herein, can be used to help determine whether a particular treatment modality, such as any one of the above, or a combination thereof, should be administered. Whether a treatment option such as any of the above mentioned treatment options is administered can be determined based on the presence or absence of a particular genetic variation risk variant in the individual, or by monitoring expression of genes that are associated with the variants of the present disclosure. Expression levels and/or mRNA levels can thus be determined before and during treatment to monitor its effectiveness. Alternatively, or concomitantly, the status with respect to a genetic variation, and/or genotype and/or haplotype status of at least one risk

variant for PML presented herein can be determined before and during treatment to monitor its effectiveness. It can also be appreciated by those skilled in the art that aberrant expression levels of a gene impacted by a CNV or other mutations found as a consequence of targeted sequencing of the CNV-identified gene can be assayed or diagnostically tested for by measuring the polypeptide expression level of said aberrantly expressed gene. In another embodiment, aberrant expression levels of a gene may result from a CNV impacting a DNA sequence (e.g., transcription factor binding site) that regulates a gene whose aberrant expression level is involved in or causes PML, or other mutations found as a consequence of targeted sequencing of the CNV-identified gene regulatory sequence, can be assayed or diagnostically tested for by measuring the polypeptide expression level of the gene involved in or causative of PML. In some embodiments, a specific CNV mutation within a gene, or other specific mutations found upon targeted sequencing of a CNV-identified gene found to be involved in or causative of PML, may cause an aberrant structural change in the expressed polypeptide that results from said gene mutations and the altered polypeptide structure(s) can be assayed via various methods known to those skilled in the art.

[0500] Alternatively, biological networks or metabolic pathways related to the genes within, or associated with, the genetic variations described herein can be monitored by determining mRNA and/or polypeptide levels. This can be done for example, by monitoring expression levels of polypeptides for several genes belonging to the network and/or pathway in nucleic acid samples taken before and during treatment. Alternatively, metabolites belonging to the biological network or metabolic pathway can be determined before and during treatment. Effectiveness of the treatment is determined by comparing observed changes in expression levels/metabolite levels during treatment to corresponding data from healthy subjects.

[0501] In some embodiments, the genetic variations described herein and/or those subsequently found (e.g., via other genetic analysis methods such as sequencing) via targeted analysis of those genes initially identified by the genetic variations described herein, can be used to prevent adverse effects associated with a therapeutic agent, such as during clinical trials. For example, individuals who are carriers of at least one at-risk genetic variation can be more likely to respond negatively to a therapeutic agent, such as an immunosuppressive agent. For example, carriers of certain genetic variants may be more likely to show an adverse response to the therapeutic agent. In some embodiments, one or more of the genetic variations employed during clinical trials for a given therapeutic agent can be used in a companion diagnostic test that is administered to the patient prior to administration of the therapeutic agent to determine if the patient is likely to have a favorable or an adverse response to the therapeutic agent.

[0502] The genetic variations described herein can be used for determining whether a subject is administered a pharmaceutical agent, such as an immunosuppressive drug. Certain combinations of variants, including those described herein, but also combinations with other risk variants for PML, can be suitable for one selection of treatment options, while other variant combinations can be suitable for selection of other treatment options. Such combinations of variants can include one variant, two variants, three variants, or

four or more variants, as needed to determine with clinically reliable accuracy the selection of treatment module. In another embodiment, information from testing for the genetic variations described herein, or other rare genetic variations in or near the genes described herein, may be combined with information from other types of testing (e.g., a JCV antibody test, CD62L test, or CSF IgM oligoclonal bands test) for selection of treatment options.

Kits

[0503] Kits useful in the methods of the disclosure comprise components useful in any of the methods described herein, including for example, primers for nucleic acid amplification, hybridization probes for detecting genetic variation, or other marker detection, restriction enzymes, nucleic acid probes, optionally labeled with suitable labels, allele-specific oligonucleotides, antibodies that bind to an altered polypeptide encoded by a nucleic acid of the disclosure as described herein or to a wild type polypeptide encoded by a nucleic acid of the disclosure as described herein, means for amplification of genetic variations or fragments thereof, means for analyzing the nucleic acid sequence of nucleic acids comprising genetic variations as described herein, means for analyzing the amino acid sequence of a polypeptide encoded by a genetic variation, or a nucleic acid associated with a genetic variation, etc. The kits can for example, include necessary buffers, nucleic acid primers for amplifying nucleic acids, and reagents for allele-specific detection of the fragments amplified using such primers and necessary enzymes (e.g., DNA polymerase). Additionally, kits can provide reagents for assays to be used in combination with the methods of the present disclosure, for example, reagents for use with other screening assays for PML.

[0504] In some embodiments, the disclosure pertains to a kit for assaying a nucleic acid sample from a subject to detect the presence of a genetic variation, wherein the kit comprises reagents necessary for selectively detecting at least one particular genetic variation in the genome of the individual. In some embodiments, the disclosure pertains to a kit for assaying a nucleic acid sample from a subject to detect the presence of at least one particular allele of at least one polymorphism associated with a genetic variation in the genome of the subject. In some embodiments, the reagents comprise at least one contiguous oligonucleotide that hybridizes to a fragment of the genome of the individual comprising at least genetic variation. In some embodiments, the reagents comprise at least one pair of oligonucleotides that hybridize to opposite strands of a genomic segment obtained from a subject, wherein each oligonucleotide primer pair is designed to selectively amplify a fragment of the genome of the individual that includes at least one genetic variation, or a fragment of a genetic variation. Such oligonucleotides or nucleic acids can be designed using the methods described herein. In some embodiments, the kit comprises one or more labeled nucleic acids capable of allele-specific detection of one or more specific polymorphic markers or haplotypes with a genetic variation, and reagents for detection of the label. In some embodiments, a kit for detecting SNP markers can comprise a detection oligonucleotide probe, that hybridizes to a segment of template DNA containing a SNP polymorphism to be detected, an enhancer oligonucleotide probe, detection probe, primer and/or an endonuclease, for example, as described by Kutuyavin et al.,

(Nucleic Acid Res. 34:e128 (2006)). In other embodiments, the kit can contain reagents for detecting SNVs and/or CNVs.

[0505] In some embodiments, the DNA template is amplified by any means of the present disclosure, prior to assessment for the presence of specific genetic variations as described herein. Standard methods well known to the skilled person for performing these methods can be utilized, and are within scope of the disclosure. In one such embodiment, reagents for performing these methods can be included in the reagent kit.

[0506] In a further aspect of the present disclosure, a pharmaceutical pack (kit) is provided, the pack comprising a therapeutic agent and a set of instructions for administration of the therapeutic agent to humans screened for one or more variants of the present disclosure, as disclosed herein. The therapeutic agent can be a small molecule drug, an antibody, a peptide, an antisense or RNAi molecule, or other therapeutic molecules as described herein. In some embodiments, an individual identified as a non-carrier of at least one variant of the present disclosure is instructed to take the therapeutic agent. In one such embodiment, an individual identified as a non-carrier of at least one variant of the present disclosure is instructed to take a prescribed dose of the therapeutic agent. In some embodiments, an individual identified as a carrier of at least one variant of the present disclosure is instructed not to take the therapeutic agent. In some embodiments, an individual identified as a carrier of at least one variant of the present disclosure is instructed not to take a prescribed dose of the therapeutic agent. In some embodiments, an individual identified as a carrier of at least one variant of the present disclosure is instructed to take an agent that targets the JC Virus. For example, an individual identified as a carrier of at least one variant of the present disclosure can be instructed to take an agent that targets the JC Virus prior to or in conjunction with, taking an immunosuppressive agent.

[0507] Also provided herein are articles of manufacture, comprising a probe that hybridizes with a region of human chromosome as described herein and can be used to detect a polymorphism described herein. For example, any of the probes for detecting polymorphisms or genetic variations described herein can be combined with packaging material to generate articles of manufacture or kits. The kit can include one or more other elements including: instructions for use; and other reagents such as a label or an agent useful for attaching a label to the probe. Instructions for use can include instructions for screening applications of the probe for making a diagnosis, prognosis, or theranosis to PML in a method described herein. Other instructions can include instructions for attaching a label to the probe, instructions for performing in situ analysis with the probe, and/or instructions for obtaining a nucleic acid sample to be analyzed from a subject. In some cases, the kit can include a labeled probe that hybridizes to a region of human chromosome as described herein.

[0508] The kit can also include one or more additional reference or control probes that hybridize to the same chromosome or another chromosome or portion thereof that can have an abnormality associated with a particular endophenotype. A kit that includes additional probes can further include labels, e.g., one or more of the same or different labels for the probes. In other embodiments, the additional probe or probes provided with the kit can be a labeled probe

or probes. When the kit further includes one or more additional probe or probes, the kit can further provide instructions for the use of the additional probe or probes. Kits for use in self-testing can also be provided. Such test kits can include devices and instructions that a subject can use to obtain a nucleic acid sample (e.g., buccal cells, blood) without the aid of a health care provider. For example, buccal cells can be obtained using a buccal swab or brush, or using mouthwash.

[0509] Kits as provided herein can also include a mailer (e.g., a postage paid envelope or mailing pack) that can be used to return the nucleic acid sample for analysis, e.g., to a laboratory. The kit can include one or more containers for the nucleic acid sample, or the nucleic acid sample can be in a standard blood collection vial. The kit can also include one or more of an informed consent form, a test requisition form, and instructions on how to use the kit in a method described herein. Methods for using such kits are also included herein. One or more of the forms (e.g., the test requisition form) and the container holding the nucleic acid sample can be coded, for example, with a bar code for identifying the subject who provided the nucleic acid sample.

[0510] In some embodiments, an in vitro screening test can comprise one or more devices, tools, and equipment configured to collect a nucleic acid sample from an individual. In some embodiments of an in vitro screening test, tools to collect a nucleic acid sample can include one or more of a swab, a scalpel, a syringe, a scraper, a container, and other devices and reagents designed to facilitate the collection, storage, and transport of a nucleic acid sample. In some embodiments, an in vitro screening test can include reagents or solutions for collecting, stabilizing, storing, and processing a nucleic acid sample.

[0511] Such reagents and solutions for nucleotide collecting, stabilizing, storing, and processing are well known by those of skill in the art and can be indicated by specific methods used by an in vitro screening test as described herein. In some embodiments, an in vitro screening test as disclosed herein, can comprise a microarray apparatus and reagents, a flow cell apparatus and reagents, a multiplex nucleotide sequencer and reagents, and additional hardware and software necessary to assay a nucleic acid sample for certain genetic markers and to detect and visualize certain genetic markers.

[0512] The present disclosure further relates to kits for using antibodies in the methods described herein. This includes, but is not limited to, kits for detecting the presence of a variant polypeptide in a test nucleic acid sample. One preferred embodiment comprises antibodies such as a labeled or labelable antibody and a compound or agent for detecting variant polypeptides in a nucleic acid sample, means for determining the amount or the presence and/or absence of variant polypeptide in the nucleic acid sample, and means for comparing the amount of variant polypeptide in the nucleic acid sample with a standard, as well as instructions for use of the kit. In certain embodiments, the kit further comprises a set of instructions for using the reagents comprising the kit.

Computer-Implemented Aspects

[0513] As understood by those of ordinary skill in the art, the methods and information described herein (genetic variation association with PML) can be implemented, in all or in part, as computer executable instructions on known

computer readable media. For example, the methods described herein can be implemented in hardware. Alternatively, the method can be implemented in software stored in, for example, one or more memories or other computer readable medium and implemented on one or more processors. As is known, the processors can be associated with one or more controllers, calculation units and/or other units of a computer system, or implanted in firmware as desired. If implemented in software, the routines can be stored in any computer readable memory such as in RAM, ROM, flash memory, a magnetic disk, a laser disk, or other storage medium, as is also known. Likewise, this software can be delivered to a computing device via any known delivery method including, for example, over a communication channel such as a telephone line, the Internet, a wireless connection, etc., or via a transportable medium, such as a computer readable disk, flash drive, etc.

[0514] More generally, and as understood by those of ordinary skill in the art, the various steps described above can be implemented as various blocks, operations, tools, modules and techniques which, in turn, can be implemented in hardware, firmware, software, or any combination of hardware, firmware, and/or software. When implemented in hardware, some or all of the blocks, operations, techniques, etc. can be implemented in, for example, a custom integrated circuit (IC), an application specific integrated circuit (ASIC), a field programmable logic array (FPGA), a programmable logic array (PLA), etc.

[0515] Results from such genotyping can be stored in a data storage unit, such as a data carrier, including computer databases, data storage disks, or by other convenient data storage means. In certain embodiments, the computer database is an object database, a relational database or a post-relational database. Data can be retrieved from the data storage unit using any convenient data query method.

[0516] When implemented in software, the software can be stored in any known computer readable medium such as on a magnetic disk, an optical disk, or other storage medium, in a RAM or ROM or flash memory of a computer, processor, hard disk drive, optical disk drive, tape drive, etc. Likewise, the software can be delivered to a user or a computing system via any known delivery method including, for example, on a computer readable disk or other transportable computer storage mechanism.

[0517] The steps of the claimed methods can be operational with numerous other general purpose or special purpose computing system environments or configurations. Examples of well known computing systems, environments, and/or configurations that can be suitable for use with the methods or system of the claims include, but are not limited to, personal computers, server computers, hand-held or laptop devices, multiprocessor systems, microprocessor-based systems, set top boxes, programmable consumer electronics, network PCs, minicomputers, mainframe computers, distributed computing environments that include any of the above systems or devices, and the like.

[0518] The steps of the claimed method and system can be described in the general context of computer-executable instructions, such as program modules, being executed by a computer. Generally, program modules include routines, programs, objects, components, and/or data structures that perform particular tasks or implement particular abstract data types. The methods and apparatus can also be practiced in distributed computing environments where tasks are

performed by remote processing devices that are linked through a communications network. In both integrated and distributed computing environments, program modules can be located in both local and remote computer storage media including memory storage devices. Numerous alternative embodiments could be implemented, using either current technology or technology developed after the filing date of this application, which would still fall within the scope of the claims defining the disclosure.

[0519] While the risk evaluation system and method, and other elements, have been described as preferably being implemented in software, they can be implemented in hardware, firmware, etc., and can be implemented by any other processor. Thus, the elements described herein can be implemented in a standard multi-purpose CPU or on specifically designed hardware or firmware such as an application-specific integrated circuit (ASIC) or other hard-wired device as desired. When implemented in software, the software routine can be stored in any computer readable memory such as on a magnetic disk, a laser disk, or other storage medium, in a RAM or ROM of a computer or processor, in any database, etc. Likewise, this software can be delivered to a user or a screening system via any known or desired delivery method including, for example, on a computer readable disk or other transportable computer storage mechanism or over a communication channel, for example, a telephone line, the internet, or wireless communication. Modifications and variations can be made in the techniques and structures described and illustrated herein without departing from the spirit and scope of the present disclosure.

[0520] Unless otherwise explained, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. The following references contain embodiments of the methods and compositions that can be used herein: The Merck Manual of Diagnosis and Therapy, 18th Edition, published by Merck Research Laboratories, 2006 (ISBN 0-911910-18-2); Benjamin Lewin, *Genes IX*, published by Jones & Bartlett Publishing, 2007 (ISBN-13: 9780763740634); Kendrew et al., (eds.), *The Encyclopedia of Molecular Biology*, published by Blackwell Science Ltd., 1994 (ISBN 0-632-02182-9); and Robert A. Meyers (ed.), *Molecular Biology and Biotechnology: a Comprehensive Desk Reference*, published by VCH Publishers, Inc., 1995 (ISBN 1-56081-569-8).

[0521] Standard procedures of the present disclosure are described, e.g., in Maniatis et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., USA (1982); Sambrook et al., *Molecular Cloning: A Laboratory Manual* (2 ed.), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., USA (1989); Davis et al., *Basic Methods in Molecular Biology*, Elsevier Science Publishing, Inc., New York, USA (1986); or *Methods in Enzymology: Guide to Molecular Cloning Techniques Vol. 152*, S. L. Berger and A. R. Kimmmerl (eds.), Academic Press Inc., San Diego, USA (1987). *Current Protocols in Molecular Biology (CPMB)* (Fred M. Ausubel, et al., ed., John Wiley and Sons, Inc.), *Current Protocols in Protein Science (CPPS)* (John E. Coligan, et al., ed., John Wiley and Sons, Inc.), *Current Protocols in Immunology (CPI)* (John E. Coligan, et al., ed. John Wiley and Sons, Inc.), *Current Protocols in Cell Biology (CPCB)* (Juan S. Bonifacino et al., ed., John Wiley and Sons, Inc.), *Culture of Animal Cells: A Manual of Basic Technique*

by R. Ian Freshney, Publisher: Wiley-Liss; 5th edition (2005), and *Animal Cell Culture Methods (Methods in Cell Biology, Vol. 57, Jennie P. Mather and David Barnes editors, Academic Press, 1st edition, 1998)*, which are all incorporated by reference herein in their entireties.

[0522] It should be understood that the following examples should not be construed as being limiting to the particular methodology, protocols, and compositions, etc., described herein and, as such, can vary. The following terms used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the embodiments disclosed herein.

[0523] Disclosed herein are molecules, materials, compositions, and components that can be used for, can be used in conjunction with, can be used in preparation for, or are products of methods and compositions disclosed herein. It is understood that when combinations, subsets, interactions, groups, etc. of these materials are disclosed and while specific reference of each various individual and collective combinations and permutation of these molecules and compounds cannot be explicitly disclosed, each is specifically contemplated and described herein. For example, if a nucleotide or nucleic acid is disclosed and discussed and a number of modifications that can be made to a number of molecules including the nucleotide or nucleic acid are discussed, each and every combination and permutation of nucleotide or nucleic acid and the modifications that are possible are specifically contemplated unless specifically indicated to the contrary. This concept applies to all aspects of this application including, but not limited to, steps in methods of making and using the disclosed molecules and compositions. Thus, if there are a variety of additional steps that can be performed it is understood that each of these additional steps can be performed with any specific embodiment or combination of embodiments of the disclosed methods, and that each such combination is specifically contemplated and should be considered disclosed.

[0524] Those skilled in the art can recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the method and compositions described herein. Such equivalents are intended to be encompassed by the following claims.

[0525] It is understood that the disclosed methods and compositions are not limited to the particular methodology, protocols, and reagents described as these can vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present disclosure which can be limited only by the appended claims.

[0526] Unless defined otherwise, all technical and scientific terms used herein have the meanings that would be commonly understood by one of skill in the art in the context of the present specification.

[0527] It should be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, reference to "a nucleotide" includes a plurality of such nucleotides; reference to "the nucleotide" is a reference to one or more nucleotides and equivalents thereof known to those skilled in the art, and so forth.

[0528] The term "and/or" shall in the present context be understood to indicate that either or both of the items connected by it are involved. While preferred embodiments

of the present disclosure have been shown and described herein, it can be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions can now occur to those skilled in the art without departing from the disclosure. It should be understood that various alternatives to the embodiments of the disclosure described herein can be employed in practicing the disclosure. It is intended that the following claims define the scope of the disclosure and that methods and structures within the scope of these claims and their equivalents be covered thereby.

EXAMPLES

Example 1—Experimental Approach

[0529] In the present study, a set of genes were identified, deleterious variants within which increase susceptibility to PML. The relevant genes were discovered on the basis of a combined CNV plus sequence analysis approach. Two sets of genes were compiled (see Table 6 and corresponding description):

[0530] A. A set based on a detailed literature review of genes involved in the immune system and JC virus biology, along with genes described in the context of PML via case reports.

[0531] B. A set based on the observation of rare CNVs within the PML cohort.

[0532] A non-redundant list of 419 genes was generated (see Table 6), which contains 245 curated from immune deficiency (immunodeficiency) reviews (Table 6, ‘Public_db’), 169 identified via rare CNVs using the methods described herein (Table 6, ‘PBio’), and 6 genes that were found using both methods (Table 6, ‘Both’). See Table 6 and description below for further information).

[0533] Using this set of 419 genes, it was determined whether:

[0534] Rare CNVs were present that might explain the susceptibility to PML;

[0535] Rare sequence variants (determined via whole exome sequencing analysis—WES) were present that might explain the susceptibility to PML;

[0536] Combinations of CNVs, SNVs and/or CNVs and SNVs might explain the susceptibility;

[0537] Individual variants might be present at higher frequency in the PML cohort (variant burden analysis—Tables 14, 15);

[0538] Total numbers of heterozygous, damaging variants were high for any specific genes (gene burden analysis—Table 13).

[0539] In all cases, due consideration was given to:

[0540] Pathogenic/deleterious nature of the variants observed (e.g., whether gene function was highly likely to be affected);

[0541] Rarity of the variants or variant combinations (e.g., those that would be expected to be present in 1% or less of the normal population were considered);

[0542] Ethnicity of the PML cases to account for potential frequency differences in one population subgroup vs. another. Ethnicities (e.g., ancestry) for the PML patients are reported in Table 7. For Sample_ID identifiers beginning with ‘MVGS’, ethnicities were not reported but all patients were from the USA and their ethnicities were assumed to be of European (EUR) ancestry. However, PML case MVGS811-13a is poten-

tially of African (AFR) ancestry on the basis of common SNVs that are also found in PML cases known to be of AFR ancestry. In one embodiment, ethnic-specific frequency data from the ExAC database was used to assess relative frequencies of variants found in PML patients vs. an unselected population (ExAC subjects). ExAC ethnicities were designated as follows: African/African American (AFR), Latino (LAT, also known as AMR), East Asian (EAS), Finnish (FIN), Non-Finnish European (EUR, also known as NFE), South Asian (SAS), and Other (OTH). For some PML cases reported in Table 7, the ethnicities were alternately reported as Subsaharan, North African (MGB), Caribbean (CAR), or Hispanic (HISP). For interpretation of variants found in these patients, the assignments of ancestry using ExAC db designations were as follows: AFR=MGB or Subsaharan; LAT=CAR or HISP. Ancestry was unknown for two PML cases (PML02 and PML28) and, for frequency interpretation purposes (using ExAC db), they were assumed to be of European (EUR) ancestry.

[0543] While the primary genetic mechanism that was considered was autosomal recessive (AR) inheritance, a number of solutions were based on autosomal dominant (AD) inheritance but only in cases for which prior evidence was found that heterozygous variants in the relevant gene had previously been associated with an immune deficiency syndrome. It can be appreciated by those skilled in the art that some genes may contain both AR and AD model pathogenic variants (e.g., see Table 6 entries marked as ‘AD_AR’ in the ‘Disease_Model’ column).

[0544] For AR inheritance (~40% of genes in Table 6 fall into this category, AR or AD_AR), the following were considered:

[0545] Homozygous or compound heterozygous gene-disruptive CNVs;

[0546] Homozygous or compound heterozygous sequence variants; i.e. single nucleotide variants (SNVs). Compound heterozygosity was only inferred when either phasing was available or one of the pairs of SNVs was itself homozygous;

[0547] Compound heterozygosity for a CNV and SNV. Such calls were only possible in cases for which the SNV was in trans to a deletion (e.g., DUSP16 SNV in Table 10 and the CNV in Table 1).

Example 2—Copy Number Variant (CNV) Analysis

[0548] The data presented herein was generated on the basis of a comparison of copy number variants (CNVs) identified in 2 cohorts:

[0549] 1) 1,005 Normal individuals (Normal Variation Engine—NVE);

[0550] 2) 71 Progressive Multifocal Leukoencephalopathy (PML) cases along with 6 Human Immunodeficiency Virus (HIV) cases without a diagnosis of PML (in order to aid in distinguishing germline variants vs. acquired variants that result from HIV infection). Total cohort size=77.

Genomic DNA Sample Hybridization—NVE and PML, HIV Cohorts

[0551] Genomic DNA samples from individuals within the Normal cohort (NVE ‘test’ subjects, also referred to as

'NVE cases' in some tables herein) and from the PML, HIV cohort (PML, HIV 'test' subjects) were hybridized against a single, sex-matched reference individual. Reference DNA samples were labeled with Cy5 and test subject DNA samples were labeled with Cy3. After labeling, samples were combined and co-hybridized to Agilent 1M feature oligonucleotide microarrays, design ID 021529 (Agilent Product Number G4447A) using standard conditions (array Comparative Genomic Hybridization—aCGH). Post-hybridization, arrays were scanned at 2 μ m resolution, using Agilent's DNA microarray scanner, generating tiff images for later analysis.

[0552] All tiff images were analyzed using Agilent Feature Extraction (FE) software, with the following settings:

Human Genome Freeze:	hg18:NCBI36:Mar2006
FE version:	10.7.3.1
Grid/design file:	021529_D_F_20091001
Protocol:	CGH_107_Sep09

[0553] This procedure generates a variety of output files, one of which is a text-tab delimited file, containing ~1,000,000 rows of data, each corresponding to a specific feature on the array. This *.txt file was used to perform CNV calling using DNACopy, an open source software package implemented in R via BioConductor (<http://www.bioconductor.org/packages/release/bioc/html/DNACopy.html>). Heterozygous losses (het_loss), homozygous losses (hom_loss) or gains were determined according to a threshold log 2ratio, which was set at:

```
hom_loss min = -1000;
hom_loss max = -2;
het_loss min = -2;
het_loss max = -0.5;
gain min = 0.5;
gain max = 1000;
```

[0554] With very few exceptions, all CNVs with a log 2ratio value between -0.5 and +0.5 were not considered. All log 2ratio values were determined according to Cy3/Cy5 (Test/Reference). A minimum probe threshold for CNV-calling was set at 2 (2 consecutive probes were sufficient to call a CNV). A CNV list was generated for each individual in the 3 cohorts (NVE, PML, and HIV).

[0555] Using custom scripts, CNVs identified in the NVE and PML cohorts (many of which appeared in multiple individuals) were (separately) 'merged' into master lists of non-redundant CNV-subregions, according to the presence or absence of the CNV-subregion in individuals within the cohort. Using this approach, the NVE-master lists have:

[0556] 7778 het_loss
 [0557] 653 hom_loss
 [0558] 4862 gain

distinct CNV-subregions, respectively. The PML+HIV cohort of 77 individuals master lists contained:

[0559] 2523 het_loss
 [0560] 314 hom_loss
 [0561] 1639 gain

distinct CNV-subregions, respectively.

[0562] Those skilled in the art can appreciate that CNVs can be acquired in an individual's genome that are not inherited. Such 'acquired CNVs' often occur in a tissue specific manner, such as in solid tumors compared to a patient's normal tissue. In blood-derived genomic DNA samples, which are what was used for both the NVE and PML subjects in the studies described herein, acquired CNVs can be the result of blood cancers such as leukemia and lymphoma, but also due to HIV infection. Many of the PML cases in this study had HIV as their primary disease (see Table 7). In order to aid in the interpretation of acquired vs. germline CNVs, an HIV sub-cohort of 6 cases was included in the primary, genome-wide CNV comparison but rare CNVs in the 6 HIV (non-PML) cases were not considered as relevant to PML susceptibility. The purpose of generating data on the 6 HIV cases was to determine whether some changes seen in PML patients who developed the disorder on a background of HIV (PML/HIV) were likely related to the underlying HIV and not the PML susceptibility itself. In other words, the HIV cases served as a general control for the large number of PML/HIV cases.

[0563] For example, consider 3 individuals within the NVE cohort with the following hypothetical CNVs:

[0564] Chr1:1-100,000; Chr1:10,001-100,000; and Chr1:1-89,999. In the master list, these would be merged into 3 distinct CNV subregions, as follows:

CNV-subregion 1	Chr1:1-10,000	Subjects A, C
CNV-subregion 2	Chr1: 10,001-89,999	Subjects A, B, C
CNV-subregion 3	Chr90,000: 1-100,000	Subjects A, B

[0565] Comparison of the corresponding NVE and PML master lists of CNV-subregions was performed (het_loss versus het_loss, hom_loss versus hom_loss and gain versus gain), resulting in a combined file with totals for NVE and PML for each distinct CNV-subregion in the study.

[0566] The data are subsequently curated as follows (The example calculation below was based on an original PML cohort of 80 cases, of which 6 are non-PML HIV controls and 3 PML cases that were duplicate samples. In some instances, the OR and FET values reported in Table 2 were used as 'relative' guidelines when considering the relevance of a CNV. In nearly all instances, a CNV was considered as a potential cause or contributing factor to PML if it was absent from the NVE database of CNVs).

[0567] Annotation using custom designed scripts in order to attach relevant information to each CNV region regarding overlap with known genes and exons, overlap with genes involved in the immune system and overlap with regulatory regions, including transcription factor binding sites.

[0568] A calculation of the odds ratio (OR) and Fishers Exact test (FET) for each CNV-subregion, according to the following formula:

$$OR = (PML / (77 - PML)) / (NVE / (1005 - NVE))$$

where:

PML = number of PML individuals with CNV-subregion of interest

NVE = number of NVE individuals with CNV-subregion of interest

[0569] As an illustrative example, consider the CNV sub-region gain involving chr2:55764753-55771586, which is found in 3 individuals in the PML cohort and 1 individual in the NVE cohort (see Table 2). The OR is: $(3/74)/(1/1004) = 40.7$

[0570] Note that, by one convention, if either of NVE or PML=0, a value of 0.5 is added to all 4 entries in the main formula above, in order to avoid dealing with infinities (see Deeks and Higgins, Statistical algorithms in Review Manager 5, Statistical Methods Group of The Cochrane Collaboration, (2010)). This has the effect of artificially lowering OR values in cases where no individuals within the NVE have the CNV. This method is applicable to all the calculations in Table 2. This method is also used when calculating the Fisher's 2-tailed Exact Test (FET) in the event that any one of the variables is zero. For convenience in analysis, the sub-cohort of 6 HIV (non-PML) cases were retained in Table 2. Therefore, the OR values reported in Table 2 are slightly different from the OR calculations for the actual number of PML cases (n=71). Using the example above for a CNV-subregion gain involving chr2:55764753-55771586, the actual OR using 71 PML cases vs. 1005 NVE subjects was: $(3/68)/(1/(1004))=44.29$. In some instances, a non-PML HIV control (see Table 11, identified as 3280, 3281, 3283, 3284, 3285, and 3286) is found to have a CNV of potential relevance in PML subjects. This can also impact the OR calculation. For example, for CNV-subregion loss chr19:55247874-55250186 the OR in Table 2 is listed as 17.38 but one case is a non-PML HIV control (Table 11, PML70_control=3280). For this example, the actual OR using 71 PML cases vs. 1005 NVE subjects, and excluding the non-PML HIV case, was: $(4/67)/(4/(1001))=14.94$.

[0571] The CNV-subregions/genes that are listed herein (e.g., in one or more of Tables 1-4), fulfill one of the following criteria:

[0572] Strong biology linking the gene that a CNV-subregion impacts or is near, with known immune deficiency pathways/mechanisms or biology in PML (e.g., JC virus related biology). That is, in some cases, statistical evidence is lacking but does not exclude the CNV-subregion as a candidate;

[0573] Statistical analysis combined with medium to strong biology (e.g., links in the peer-reviewed literature to PML, JC virus, host defense, immune deficiency, or neuropathology) without obvious biological connection (best FET in this category was $3.25E-10$);

[0574] It can be appreciated by those skilled in the art that the number of PML candidate CNV-subregions, irrespective of category, may increase or decrease as additional PML cohorts are analyzed.

Example 3—Whole Exome Sequencing (WES) and Case Level Analysis

[0575] WES data was obtained on a total of 70 PML cases (non-PML HIV cases were not sequenced—they were used simply to help in the interpretation of complex CNVs observed in PML patients who also had HIV).

[0576] Variant annotation reports were further interrogated against the full set of genes detailed above. Synonymous variants and variants predicted to be modifiers (outside coding regions) were not considered. For all other variants, further filtering was performed so that only those predicted by at least one in silico prediction algorithm (e.g., Polyphen2, SIFT, MutationTaster) to be pathogenic were considered for further evaluation. Finally, only variants or variant combinations that would be expected to be present in 1% or less of the normal population were evaluated for case level analysis (Tables 7-10). Data from the Exome Aggregation Consortium (ExAC) was used to obtain ethnic-specific frequency data for variants under consideration (see, Lek et al., Nature, 17; 536(7616):285-91) (2016)).

Example 4—Description of Sequence Data

[0577] The sequence file 33655-710.101_ST25.txt contains genomic sequence information for (in the following order):

[0578] A. All distinct CNVs listed in Table 1;

[0579] B. The full genomic extent of the transcripts listed in Table 4;

[0580] C. Sequence variants detailed in Table 5.

[0581] D. The full genomic extent of the transcripts listed in Table 12

[0582] Note that:

[0583] 1. SEQ_ID 1-172 are the CNV sequences from Table 1;

[0584] 2. SEQ_ID 173-455 are the transcript sequences from Table 4;

[0585] 3. SEQ_ID 1000-1329 are the sequence variants from Table 5;

[0586] 4. SEQ_ID 1500-2177 are the transcript sequences from Table 12.

Examples of Sequences Submitted

Sequence entry starts:

TABLE 1

```
SEQ_ID 1 = 49,653bp CNV (het_loss) at chr1: 1086119-1135772
involving genes MIR200A, MIR200B, MIR429, TNFRSF18, TTL10:
<210> 1

<211> 49654

<212> DNA

<213> Homo sapiens
```

TABLE 1-continued

```

<400> 1
cttctggggt ctaaggccag aagtgacctt tcttctcag gaggcacccc cacatcacag    60
gccccaaagt cccaccagga gtccccaggc agcagggttt ccaccacagc cgggaagagc   120
cccgcttca ccaccacca ccagccaatc ccgagaccac cgaagccccc agaccgggccc   180
.....(sequence truncated for brevity)
gattcccgca cggccgggga cggccccagg gccttgggag cgtctgtgga cacctgtggt 49560
gtgggcccag gagctgggag ctcatctgaa cagccagca ctccgcacac cacgtgtgtg 49620
gcggatgcct gggtttctcc actgtggggc cacg                                49654

```

Sequence entry ends.

Sequence entry starts:

TABLE 4

```

SEQ_ID 173 = MIR200B, transcript NR_029639, which is 95bp in length:
<210> 173

```

<211> 95

<212> DNA

<213> *Homo sapiens*

<400> 173

```

ccagctcggg cagccgtggc catcttactg ggcagcattg gatggagtca ggtctctaat    60
actgcctggt aatgatgacg geggagccct gcacg                                95

```

Sequence entry ends.

Sequence entry starts:

TABLE 5

```

SEQ ID 1148 = chr9:304628 reference allele = G;
alternate allele = A

```

<210> 1148

<211> 40

<212> DNA

<213> *Homo sapiens*

TABLE 5-continued

<220>

<221> variant

<222> (20) . . . (20)

<223> G->A

<400> 1148

```

tttaaaaaga ctggatctog aaaagatttt cacaagacgc    40

```

Sequence entry ends.

Sequence entry starts:

TABLE 12

```

SEQ_ID 1500 = ACADM, transcript NM_000016, which is 39,
313 bp in length:
<210> 1500

```

<211> 39313

<212> DNA

<213> *Homo sapiens*

<400> 1500

```

cgcaagtccc cccaccgttc agcgcaaccg ggccctccca gccccgccgc cgtccccctc    60

```

TABLE 12-continued

```

ccccgccctg getctctttc cgcgctgcgg tcagcctcgg cgtcccacag agagggccag 120
.....(sequence truncated for brevity)
gtaatagtgt atatttcttg tatttactat gatgaaaaaa ggctggttta attttgaatt 39240
gaataaagtt acctgttcat tttttattag atattttaaa gacttcagaa aatataaata 39300
tgaaataatt taa 39313

```

Sequence entry ends.

Example 5

[0587] Those skilled in the art can appreciate that genes can be impacted by acquired or germline genetic variants (e.g., CNVs), wherein each gene has the potential to contain genetic variants that are acquired (e.g., via a disease process such as HIV infection, or cancers such as leukemia and lymphoma) or present in the germ line (e.g., inherited from a parent or are de novo, i.e. not inherited from a parent). In FIG. 1, the PRKCB gene was impacted by germ line variants in 2 PML cases and acquired variants in 6 PML cases. The invention described herein is focused on detection of germline variants that are present in PML patient genomes. Therefore, no solutions/explanations for a given patient's PML was based on an acquired CNV, although another PML patient could potentially be 'solved' by one or two germline rare variants impacting the gene.

[0588] For this PRKCB example, no CNV-based solutions were found (an AR model was assumed), but 1 SNV solution is reported in Table 8 (het SNV, an AD model is assumed for this PML case). Further supporting evidence was assessed for the PRKCB gene by performing String analysis (high confidence=0.7, 1st shell=up to 10 interactors; string-db.org; see Szklarczyk et al., (2015), and references therein). String analysis showed that PRKCB interacts with PML-419 genes CARD11, IKBKB, and RBCK1 (see Table 6).

[0589] In FIG. 2, both TNFRSF13C and CENPM are disrupted and/or gained by a set of acquired CNV gains. Acquired CNVs can be very complex, such as the high copy number gains often identified in tumor-derived DNA samples (as compared to the patient's normal genome). In the PML gene discovery described herein, blood-derived genomic DNA obtained from several PML-diagnosed HIV patients, or PML cases with a primary disease of leukemia and lymphoma (reported as 'Other' in Table 7), showed complex genomic changes (e.g., gains exhibiting a dup-trip-dup pattern). In some PML cases, the acquired gains passed the log 2 ratio cutoff (>0.5) that was selected for this study, but in other PML cases the log 2 ratios for the gains were <0.5 and this data was filtered out from the main analyses that were performed to ascertain rare germline CNVs.

[0590] In one embodiment of the invention, a set of 6 non-PML HIV cases (3 African ancestry, 3 European ancestry) were used to aid in the interpretation of whether a CNV was an acquired or germline event. The non-PML 'PML cases' are labeled with '_control' in Table 11 and correspond to 'PML_Case_ID' numbers 3280, 3281, 3283, 3284, 3285, and 3286. While some CNVs are reported in Tables 1 and 2 for this set of non-PML control HIV subjects, none of these genetic findings were used to nominate a gene discovered on the basis of rare CNVs (as compared to the NVE db) as a

potential PML gene (PBio genes reported in Table 6). In other words, these rare CNVs were only used to aid in determining if a particular genomic region containing multiple overlapping CNVs was potentially due to an acquired genetic event. Those skilled in the art can appreciate that the set of experiments described herein do not necessarily fully rule in or out that a given genomic region contained only acquired CNVs vs. only germline CNVs (i.e. it's possible that the same region can contain an acquired CNV in one individual and a germline CNV in another).

[0591] For the CNV data shown in FIG. 2, both the TNFRSF13C and CENPM genes were included in PML-419 gene list (Table 6) on the basis of their immune or neurological related biology reported in the literature. No CNV or SNV PML solutions were found for these two genes, but String analysis (high confidence=0.7, 1st shell=up to 10 interactors) shows that TNFRSF13C interacts with PML-419 genes TRAF3 (Table 7 solution) and TNFRSF13B (Table 8 solution), as well as BTK (a known PML gene, see Table 6).

[0592] FIG. 3 shows another example of a gene that is impacted by both germline and acquired CNVs. While no PML cases were solved on the basis of the acquired or germline CNVs shown to impact the PKHD1 gene, nomination of this gene to Table 6 on the basis of its biology resulted in finding 3 potential alternate solutions (AR model) for 3 other PML cases (see Table 8). However, String analysis (high confidence=0.7, 1st shell=up to 10 interactors) did not reveal any PML-419 gene interactions with PKHD1.

Example 6

[0593] Those skilled in the art can appreciate that an AR disease model would involve ascertaining whether both alleles (for a gene or genetic locus) are impacted by a genetic variant in individuals affected by the disorder. The types of genetic variants can be SNVs, CNVs, indels, etc. In the study describe herein, if an AR disease model was invoked for a gene (see Table 6), we assessed the PML patient's CGH data for CNVs (heterozygous or homozygous) and their exome data for SNVs (heterozygous or homozygous). Thus, each patient may be solved for one of the PML-419 genes (Table 6) with one of the following scenarios: homozygous deletion, homozygous duplication (log 2 ratio will appear comparable to that typically found for triplications), homozygous SNV, compound heterozygous SNVs, compound heterozygous CNVs, or compound heterozygous SNV and CNV. Those skilled in the art know that, for an AR disease mechanism, a pathogenic SNV or CNV may have appreciable frequency in the general population (e.g., up to 1% frequency) with little to no impact on the individual's

health, but when present with a second pathogenic variant on the other allele, can cause disease.

[0594] FIG. 4 shows an example of a recurrent intronic loss impacting the BMPR2 gene. Patient PML29 was found to have a homozygous deletion, whereas as patients PML58 and MVGS811-13a have a heterozygous deletion. Assuming an AR disease model, no SNV solutions were found for this gene; however, PML29 is potentially solved due to the homozygous deletion that was detected. While immune-related biology is reported for studies on BMPR2 (see Table 6), String analysis (high confidence=0.7, 1st shell=up to 10 interactors) did not reveal any PML-419 gene interactions with BMPR2.

[0595] FIG. 5 shows an example of a recurrent exonic gain that disrupts the COMMD6 gene. Two PML patients were found to have homozygous duplications of this CNV. Interestingly, while String analysis (high confidence=0.7, 1st shell=up to 10 interactors) did not reveal any PML-419 gene interactions with COMMD6, recent studies (see Table 6, PMIDs 25355947 and 27441653) show a potential link between COMMD6 and known PML gene WAS via the WASH gene.

[0596] FIG. 6 shows an example of a recurrent exonic gain that disrupts the KCTD7 gene and its right breakpoint is upstream of RABGEF1 (i.e. one or both genes may be causing/contributing to PML). A recently annotated non-coding RNA (see hg19 assembly, LOC100996437) may also be impacted by this CNV. Both genes have immune and neurological links (see Table 6) and since patient PML29 has a homozygous duplication, it was added as a PML solution in Table 7. String analysis (high confidence=0.7, 1st shell=up to 10 interactors) did not reveal any PML-419 gene interactions for either gene, but they are linked together in a joint String analysis.

[0597] FIG. 7 shows an example of a recurrent gain that disrupts FPR2 (left breakpoint) and ZNF616 (right breakpoint, gene not labeled), and other genes are fully encompassed by this CNV. There is strong supporting biology for FPR2 (see Table 6) and it is listed as a PML solution in Table 7. String analysis (high confidence=0.7, 1st shell=up to 10 interactors) did not reveal any PML-419 gene interactions for FPR2, but a joint analysis of Table 7 genes did reveal an interaction (see FIG. 13).

[0598] FIG. 8 shows an example of an exonic loss impacting the PIK3CD and PIK3CD-AS1 genes. Patient MVGS811-13a has a homozygous deletion and is reported as a solution in Table 7 based on the strong immune-related biology for PIK3CD (see Table 6). String analysis (high confidence=0.7, 1st shell=up to 10 interactors) reveals PML-419 gene interactions for PTEN and PIK3R1.

Example 7

[0599] A subset of the rare CNVs found in our PML study were located in intergenic regions. While those skilled in the art can appreciate that intergenic variants (CNVs, SNVs, etc.) can have long range effects on the expression of genes (e.g., gene regulatory elements can be located several kilobases away from the genes under their influence), in our study we assumed that intergenic CNVs were potentially impacting one or both adjacent genes if they were located <~100 Kb away, either upstream or downstream. The ENCODE project has revealed a wealth of information, such as transcription factor binding sites, and rare CNVs that were identified in the study herein were checked for their

potential impact on these sites (hg19 assembly ENCODE annotation was checked) and were often found to impact transcription factor binding sites and/or were located in conserved DNA regions.

[0600] FIG. 9 shows an intergenic, recurrent gain that is upstream of CD180. Patient MVGS995-4a has a homozygous duplication and, while not considered as a PML solution in Table 7, is potentially an alternate solution that may be causing or contributing to the patient's PML based on altered expression of CD180. The gene has immune-related biology (see Table 6) and String analysis (high confidence=0.7, 1st shell=up to 10 interactors) reveals a PML-419 gene interaction with PLCG2 (see Table 7, 2 PML cases have a solution for this gene).

[0601] FIG. 10 shows an intergenic, recurrent loss that is upstream of VDAC1. Patient PML30 has a homozygous deletion and, while not considered as a PML solution in Table 7, is potentially an alternate solution that may be causing or contributing to the patient's PML based on altered expression of VDAC1. String analysis (high confidence=0.7, 1st shell=up to 10 interactors) did not reveal any PML-419 gene interactions for VDAC1.

[0602] FIG. 11 shows an intergenic, recurrent loss that is downstream of EGR1 and ETF1. Patient PML69 has a homozygous deletion and, based on links for EGR1 to PML-419 genes (Table 6) and its proximity to EGR1 (~4 Kb away), it was added as a potential PML solution in Table 7. String analysis (high confidence=0.7, 1st shell=up to 10 interactors) reveals PML-419 gene interactions with JUN, PTEN, and TP53, but nothing of note was found for String analysis of ETF1.

[0603] FIG. 12 shows an intergenic, recurrent loss that is upstream of ITSN2. Patient PML65 has a homozygous deletion and, based on links for ITSN2 to a known PML gene (WAS) in the PML-419 gene list (Table 6), it was added as a potential PML solution in Table 7. Interestingly, another PML case was found to have a rare homozygous SNV in ITSN2, so this gene has 2 PML solutions reported in Table 7. String analysis (high confidence=0.7, 1st shell=up to 10 interactors) did not reveal any PML-419 gene interactions.

Example 8

[0604] Pathway analyses, such as protein-protein interactions, are providing valuable insights into the underlying biology for complex diseases. While PML is a very rare disease that requires several concurrent factors (e.g., infection by the JC virus), multiple genes may be independently causing or increasing the risk of developing this neurodegenerative disorder based on the presence of a genetic variant in a given gene (e.g., a heterozygous variant wherein one deleterious variant is present on the maternally or paternally inherited allele, a homozygous variant wherein the same deleterious variant is present on both alleles, or compound heterozygous variants wherein a pair of deleterious variants are present but one is found on the maternally inherited allele and the other is found on the paternally inherited allele). As hypothesized, presence of an immune deficiency genetic disorder was another prerequisite. Indeed, in the PML study described herein, 43 genes were proposed as solutions for 61 of 71 PML cases (see Table 7) that were assessed using array CGH and whole exome sequencing. Numerous algorithms and associated databases have been developed to investigate molecular pathways, such as String (see, Szklarczyk et al., (2015), and references therein).

[0605] FIG. 13 shows an example of String analysis performed on the 43 genes considered as PML solutions on the basis of an AD or AR disease model. A series of interactions were found for 21 of 43 genes, and in several instances this included interactions for genes implicated in 2 or more PML cases that are reported in Table 7 (9 cases for TNFRSF11A, 4 cases for PLCG2, 3 cases for ZAP70 and NOD2, and 2 PML cases for TICAM1).

Example 9

[0606] To determine the likelihood that a randomly selected individual would harbor one of the variants described herein, the following analysis was performed: For each variant or combination of variants, the ethnic-specific frequency quoted in Table 7 was used to determine the probability that a randomly selected individual of the same ethnicity would be expected not to harbor the variant or combination of variants. The product of all such probabilities was calculated (e.g., the probability that a randomly selected individual would not harbor any of the variants) and subtracted from 1, yielding the probability that a random individual would harbor at least one of the variants. It was found that, for HIV cases, the probability of a random individual harboring at least one of the variants was—which

is consistent with the pre-HAART risk of PML in the context of HIV. For non-HIV cases (mostly MS/NTZ), the risk was ~1%, which, again, is consistent with the risk of PML in MS/NTZ, especially after long-term therapy.

[0607] These analyses support the notion that the frequencies of the variants identified as relevant to PML risk are consistent with the actual observed risks for unselected individuals. The analyses are predicated on the reasonable assumption that there is no PML-relevant connection with the risk of developing HIV (an acquired infection) and/or MS (e.g., this implies that treatment of healthy individuals with Natalizumab, for example, would result in similar risks of PML). Any deviations (e.g., variants found in a slightly higher number of normal individuals than expected according to the numbers actually observed to be affected by PML) may be due to: penetrance (e.g., not everyone with the variants will be at maximal risk of PML); the assumption that individuals with MS, HIV and other underlying conditions represented a normal (e.g., with respect to PML risk) cross-section of the general population, prior to developing the underlying disorders HIV, MS etc; and under ascertainment of PML, even in patients with HIV, MS/NTZ.

Example 10—Tables Referenced in this Study

TABLE 1

CNVs of interest in this study							
Chr	Original CNV Start	Original CNV Stop	Original CNV Size	CNV Type	PML Case ID	RefSeq Gene Symbol	SEQ ID
1	1086119	1135772	49653	het_loss	3009	MIR200A, MIR200B, MIR429, TNFRSF18, TTL10	1
1	9634094	9635206	1112	hom_loss	3009	PIK3CD	2
1	12018512	12032581	14069	gain	3205		3
1	19593401	19602807	9406	het_loss	3203	CAPZB	4
1	21695957	21700243	4286	het_loss	3161		5
1	24364786	24391166	26380	gain	3199	IFNLR1	6
1	28666669	28737671	71002	gain	3161	PHACTR4, RCC1, SNHG3	7
1	49372054	49380088	8034	het_loss	3145	AGBL4	8
1	153816159	153827698	11539	het_loss	3168		9
1	205607255	205610341	3086	gain	3007		10
1	215760485	215762451	1966	het_loss	3117	GPATCH2	11
1	215866737	215869900	3163	het_loss	3151	GPATCH2	12
2	10352668	10356083	3415	het_loss	3007		13
2	24457024	24462631	5607	hom_loss	3204		14
2	38468717	38471950	3233	het_loss	3175		15
2	38516138	38524237	8099	het_loss	3151		16
2	38726517	38731845	5328	het_loss	3159		17
2	40620890	40624089	3199	het_loss	3202		18
2	46631006	46643501	12495	gain	3145	RHOQ	19
2	55764753	55790559	25806	gain	3143	PNPT1	20
2	55764753	55790559	25806	gain	3193	PNPT1	20
2	55764753	55790559	25806	gain	3282	PNPT1	20
2	55764753	55790559	25806	gain	3143	PNPT1	20
2	55764753	55790559	25806	gain	3193	PNPT1	20
2	55764753	55790559	25806	gain	3282	PNPT1	20
2	55764753	55790559	25806	gain	3143	PNPT1	20
2	55764753	55790559	25806	gain	3193	PNPT1	20
2	55764753	55790559	25806	gain	3282	PNPT1	20
2	71190677	71200120	9443	het_loss	3175	MCEE	21
2	71190677	71200120	9443	het_loss	3175	MCEE	21
2	71191311	71200120	8809	het_loss	3204	MCEE	22
2	71198108	71200120	2012	het_loss	3143	MCEE	23
2	71190677	71200120	9443	het_loss	3175	MCEE	21
2	71198108	71200120	2012	het_loss	3193	MCEE	23
2	71198108	71200120	2012	het_loss	3200	MCEE	23
2	71191311	71200120	8809	het_loss	3204	MCEE	22
2	74773432	74913493	140061	gain	3118	HK2	24
2	105418748	105435274	16526	het_loss	3193	FHL2	25
2	110182348	110210249	27901	gain	3174	MALL, MIR4267, MIR4436B1, MIR 4436B2	26

TABLE 1-continued

CNVs of interest in this study							
Chr	Original CNV Start	Original CNV Stop	Original CNV Size	CNV Type	PML Case ID	RefSeq Gene Symbol	SEQ ID
2	127823042	127828410	5368	het_loss	3273		27
2	134911636	134914254	2618	het_loss	3273	MGAT5	28
2	203005216	203019933	14717	het_loss	3009	BMPR2	29
2	203005216	203019933	14717	het_loss	3192	BMPR2	29
2	203005216	203019933	14717	hom_loss	3152	BMPR2	29
2	230212897	230216339	3442	het_loss	3154	DNER	30
3	122979920	122994402	14482	gain	3202	IQCB1	31
4	26565071	26566345	1274	het_loss	3010	STIM2	32
4	26565071	26566345	1274	het_loss	3125	STIM2	32
4	26565071	26566345	1274	het_loss	3168	STIM2	32
4	26565071	26566345	1274	het_loss	3282	STIM2	32
4	26565071	26566345	1274	het_loss	3284	STIM2	32
4	26565071	26566345	1274	hom_loss	3273	STIM2	32
4	54838623	54873909	35286	gain	3153	PDGFRA	33
4	90791460	90843887	52427	gain	3168		34
4	90800863	90808258	7395	het_loss	3009		35
4	90800863	90808258	7395	het_loss	3284		35
5	45331278	46150784	819506	gain	3157	HCN1	36
5	49771219	49774457	3238	gain	3273	EMB	37
5	66619415	66636116	16701	gain	3010		38
5	78480194	78497296	17102	gain	3205		39
5	78497296	78531091	33795	gain	3132		40
5	78497296	78521408	24112	gain	3185		41
5	78497296	78531091	33795	gain	3132		40
5	78497296	78521408	24112	gain	3185		41
5	78500552	78526637	26085	gain	3205		42
5	78497296	78531091	33795	gain	3132		40
5	78500552	78526637	26085	gain	3205		42
5	78497296	78531091	33795	gain	3132		40
5	83490494	83495169	4675	het_loss	3204	EDIL3	43
5	133372071	133379727	7656	hom_loss	3153		44
5	137836466	137843309	6843	hom_loss	3279		45
5	150159466	150202601	43135	het_loss	3117		46
5	150159466	150204134	44668	het_loss	3180		47
5	150159466	150202601	43135	het_loss	3199		46
5	150159466	150204134	44668	het_loss	3278		47
5	150159466	150202601	43135	het_loss	3117		46
5	150159466	150204134	44668	het_loss	3180		47
5	150159466	150202601	43135	het_loss	3199		46
5	150159466	150204134	44668	het_loss	3278		47
5	150159466	150202601	43135	het_loss	3117		46
5	150159466	150204134	44668	het_loss	3180		47
5	150159466	150202601	43135	het_loss	3199		46
5	150159466	150204134	44668	het_loss	3278		47
5	150185190	150201145	15955	hom_loss	3009		48
5	150185190	150201145	15955	hom_loss	3143		48
5	150185190	150202601	17411	hom_loss	3152		49
5	150185190	150202601	17411	hom_loss	3154		49
5	150185190	150202601	17411	hom_loss	3193		49
5	150159466	150201145	41679	hom_loss	3196		50
5	150185190	150201145	15955	hom_loss	3281		48
5	150185190	150201145	15955	hom_loss	3009		48
5	150185190	150201145	15955	hom_loss	3143		48
5	150185190	150202601	17411	hom_loss	3152		49
5	150185190	150202601	17411	hom_loss	3154		49
5	150185190	150202601	17411	hom_loss	3193		49
5	150159466	150201145	41679	hom_loss	3196		50
5	150185190	150201145	15955	hom_loss	3281		48
5	150185190	150202601	17411	hom_loss	3152		49
5	150185190	150202601	17411	hom_loss	3154		49
5	150185190	150202601	17411	hom_loss	3193		49
5	150185190	150204134	18944	het_loss	3132		51
5	150159466	150204134	44668	het_loss	3180		47
5	150202601	150204134	1533	het_loss	3196		52
5	150191322	150204134	12812	het_loss	3273		53
5	150185190	150204134	18944	het_loss	3277		51
5	150159466	150204134	44668	het_loss	3278		47
5	150185190	150204134	18944	het_loss	3280		51
5	150185190	150204134	18944	het_loss	3282		51
5	179590681	179626660	35979	het_loss	3172	MAPK9	54
6	2882577	2947403	64826	het_loss	3196	DKFZP686I15217, NQO2, SERPINB6	55

TABLE 1-continued

CNVs of interest in this study							
Chr	Original CNV Start	Original CNV Stop	Original CNV Size	CNV Type	PML Case ID	RefSeq Gene Symbol	SEQ ID
6	2964646	2966011	1365	het_loss	3193	HTATSF1P2, NQO2	56
6	51766024	51773250	7226	het_loss	3167	PKHD1	57
6	51952217	51969378	17161	gain	3127	PKHD1	58
6	51952217	51969378	17161	gain	3127	PKHD1	58
6	51953476	51965723	12247	gain	3205	PKHD1	59
6	51952217	51969378	17161	gain	3127	PKHD1	58
6	74396294	74404837	8543	het_loss	3009	SLC17A5	60
6	74396294	74398409	2115	het_loss	3160	SLC17A5	61
6	74396294	74404837	8543	het_loss	3009	SLC17A5	60
6	86416979	86431527	14548	het_loss	3197		62
6	91131823	91135670	3847	het_loss	3171		63
6	107882367	107890605	8238	het_loss	3201	PDSS2	64
6	166418511	166422386	3875	het_loss	3125		65
6	166418511	166422386	3875	het_loss	3163		65
6	166418511	166422386	3875	het_loss	3192		65
6	166418511	166422386	3875	het_loss	3193		65
6	166418511	166422386	3875	het_loss	3194		65
6	166418511	166422386	3875	het_loss	3200		65
6	166418511	166422386	3875	het_loss	3205		65
6	166418511	166422386	3875	het_loss	3280		65
6	166418511	166422386	3875	het_loss	3281		65
6	166418511	166422386	3875	het_loss	3284		65
6	166418511	166422386	3875	hom_loss	3009		65
6	166418511	166422386	3875	hom_loss	3152		65
6	166418511	166422386	3875	hom_loss	3175		65
7	65741238	65768682	27444	gain	3152	KCTD7	66
7	65741238	65768682	27444	gain	3202	KCTD7	66
7	157174966	157177843	2877	het_loss	3009	PTPRN2	67
7	157425841	157496238	70397	gain	3189	PTPRN2	68
7	158000082	158024569	24487	het_loss	3279	PTPRN2	69
7	158000082	158024569	24487	het_loss	3279	PTPRN2	69
7	158000082	158024569	24487	het_loss	3279	MIR595, PTPRN2	69
8	23103186	23125443	22257	het_loss	3140	TNFRSF10A	70
8	39914488	39919594	5106	het_loss	3126	IDO2	71
8	79905654	79910286	4632	het_loss	3159		72
8	99790200	99799839	9639	het_loss	3006	STK3	73
8	102049360	102064431	15071	het_loss	3173		74
8	102049360	102064431	15071	het_loss	3175		74
8	102049360	102064431	15071	het_loss	3282		74
9	571398	584647	13249	het_loss	3006	KANK1	75
9	571398	584647	13249	het_loss	3006	KANK1	75
9	580722	598488	17766	het_loss	3200	KANK1	76
9	580722	598488	17766	het_loss	3282	KANK1	76
9	580722	598488	17766	het_loss	3200	KANK1	76
9	580722	598488	17766	het_loss	3282	KANK1	76
9	634039	637589	3550	het_loss	3273	KANK1	77
9	634039	637589	3550	het_loss	3282	KANK1	77
9	74050088	74059447	9359	het_loss	3165	GDA	78
9	93140394	93447826	307432	gain	3198	AUH, MIR3163, MIR3910-1, MIR3910-2, NFIL3	79
9	118564159	118575633	11474	gain	3193	ASTN2	80
9	118612694	118664593	51899	het_loss	3144	ASTN2	81
9	119220847	119233078	12231	gain	3005		82
10	899657	1071401	171744	gain	3161	GTPBP4, IDI2, IDI2-AS1, LARP4B	83
10	76217585	76411591	194006	gain	3179	KAT6B	84
10	116000069	116004388	4319	gain	3010	VWA2	85
11	14677012	14689025	12013	het_loss	3199	PDE3B	86
11	34608313	34615878	7565	het_loss	3117	EHF	87
11	62382087	62398462	16375	het_loss	3205	SLC3A2	88
11	76631014	76643625	12611	het_loss	3193	GDPD4	89
12	11616557	12422129	805572	het_loss	3126	ETV6	90
12	12435301	12778142	342841	het_loss	3126	APOLD1, CDKN1B, CREBL2, DUSP16, GPR19, LOH12CR1	91
12	12968705	12971310	2605	gain	3127		92
12	91786998	94313682	2526684	het_loss	3126	EEA1, LOC643339	93
12	91786998	94313682	2526684	het_loss	3126	LOC643339, MRPL42, NUDT4, NUDT4P1, SOCS2, SOCS2-AS1, UBE2N	93
12	91786998	94313682	2526684	het_loss	3126	CCDC41, CRADD, PLXNC1	93

TABLE 1-continued

CNVs of interest in this study							
Chr	Original CNV Start	Original CNV Stop	Original CNV Size	CNV Type	PML Case ID	RefSeq Gene Symbol	SEQ ID
12	111061085	111064486	3401	het_loss	3004	TRAFD1	94
13	40939924	41026908	86984	gain	3140	RGCC	95
13	75006025	75016304	10279	gain	3009	COMMD6	96
13	75006025	75016304	10279	gain	3152	COMMD6	96
13	91811087	91814369	3282	het_loss	3143	GPC5	97
13	91811087	91811118	31	hom_loss	3173	GPC5	98
13	110754499	110778301	23802	gain	3006	ARHGEF7, TEX29	99
14	20021118	20055469	34351	gain	3205	RNASE10	100
14	20426824	20481852	55028	hom_loss	3200	ECRP,RNASE3	101
14	20430810	20490129	59319	het_loss	3192	ECRP	102
14	20430810	20490129	59319	het_loss	3192		102
14	21096689	21105611	8922	het_loss	3125		103
14	21096689	21105611	8922	het_loss	3175		103
14	21096689	21105611	8922	het_loss	3194		103
14	21096689	21105611	8922	het_loss	3204		103
14	21096689	21105611	8922	het_loss	3273		103
14	21120750	21125513	4763	gain	3143		104
14	21120750	21125513	4763	gain	3173		104
14	60901636	60909492	7856	het_loss	3193	PRKCH	105
14	60912874	60921269	8395	het_loss	3174	PRKCH	106
14	63937192	63944459	7267	gain	3205	MTHFD1	107
14	95754535	95759056	4521	het_loss	3009	BDKRB2	108
14	95754535	95759056	4521	het_loss	3173	BDKRB2	108
14	95754535	95759056	4521	het_loss	3202	BDKRB2	108
15	66065925	66082418	16493	het_loss	3010		109
15	70432627	70443017	10390	gain	3169	HEXA	110
15	75096101	75128723	32622	gain	3200	PSTPIP1	111
15	75101524	75115806	14282	gain	3132	PSTPIP1	112
15	75096101	75128723	32622	gain	3200	PSTPIP1	111
15	75105789	75115806	10017	gain	3127	PSTPIP1	113
15	75101524	75115806	14282	gain	3132	PSTPIP1	112
15	75105789	75115806	10017	gain	3199	PSTPIP1	113
15	75096101	75128723	32622	gain	3200	PSTPIP1	111
15	75105789	75115806	10017	gain	3279	PSTPIP1	113
15	75105789	75115806	10017	gain	3127	PSTPIP1	113
15	75101524	75115806	14282	gain	3132	PSTPIP1	112
15	75105789	75115806	10017	gain	3199	PSTPIP1	113
15	75096101	75128723	32622	gain	3200	PSTPIP1	111
15	75105789	75115806	10017	gain	3279	PSTPIP1	113
15	75096101	75128723	32622	gain	3200	PSTPIP1	111
15	88999998	89016848	16850	het_loss	3172		114
16	6823677	6932753	109076	het_loss	3126	RBFOX1	115
16	6823677	6932753	09076	het_loss	3126	RBFOX1	115
16	6942078	6945539	3461	gain	3173	RBFOX1	116
16	6942078	6945539	3461	gain	3175	RBFOX1	116
16	6942078	6945539	3461	gain	3282	RBFOX1	116
16	23842653	23848772	6119	het_loss	3198	PRKCB	117
16	23892842	23903495	10653	gain	3199	PRKCB	118
16	23892842	23903495	10653	gain	3199	PRKCB	118
16	23893969	23908248	14279	gain	3205	PRKCB	119
16	23893969	23908248	14279	gain	3205	PRKCB	119
16	69044235	69050151	5916	gain	3174	FUK	120
16	69044235	69050151	5916	gain	3185	FUK	120
16	69052450	69081640	29190	het_loss	3197	COG4, FUK	121
16	70653499	70665447	11948	gain	3143	HPR	122
16	70653499	70665447	11948	gain	3152	HPR	122
16	70653499	70665447	11948	gain	3192	HPR	122
16	70653499	70665447	11948	gain	3200	HPR	122
16	70653499	70665447	11948	gain	3282	HPR	122
16	70653499	70665447	11948	gain	3284	HPR	122
17	69341925	70202523	860598	gain	3183	BTBD17, C17orf77, CD300A, CD300C, CD300E, CD300LB, CD300LD, CD300LF, DNAI2, GPR142,GPRC5C,KIF19, MGC16275, RAB37, RPL38, TTYH2	123
17	75608151	75615433	7282	het_loss	3144	TBC1D16	124
17	75608151	75615433	7282	het_loss	3152	TBC1D16	124
17	75608151	75615433	7282	het_loss	3163	TBC1D16	124
17	75608151	75611602	3451	het_loss	3192	TBC1D16	125
17	75608151	75615433	7282	het_loss	3200	TBC1D16	124

TABLE 1-continued

CNVs of interest in this study							
Chr	Original CNV Start	Original CNV Stop	Original CNV Size	CNV Type	PML Case ID	RefSeq Gene Symbol	SEQ ID
17	75608151	75611602	3451	het_loss	3204	TBC1D16	125
17	75608151	75611602	3451	het_loss	3284	TBC1D16	125
17	75608151	75611602	3451	hom_loss	3009	TBC1D16	125
17	75611602	75615433	3831	hom_loss	3175	TBC1D16	126
17	75608151	75615433	7282	het_loss	3144	TBC1D16	124
17	75608151	75615433	7282	het_loss	3152	TBC1D16	124
17	75608151	75615433	7282	het_loss	3163	TBC1D16	124
17	75608151	75615433	7282	het_loss	3200	TBC1D16	124
17	76241510	76267844	26334	gain	3205	RPTOR	127
17	76247305	76265683	18378	gain	3127	RPTOR	128
17	76241510	76267844	26334	gain	3205	RPTOR	127
17	76241510	76267844	26334	gain	3205	RPTOR	127
18	9985530	10125331	139801	gain	3175		129
18	12764095	12781985	17890	gain	3191	PTPN2	130
18	27026203	27029351	3148	het_loss	3125		131
18	27026203	27029351	3148	het_loss	3143		131
18	27026203	27029351	3148	het_loss	3175		131
18	42537949	42663605	125656	gain	3125	PIAS2, ST8SIA5	132
18	46917195	46945018	27823	het_loss	3161		133
18	59457622	59465699	8077	het_loss	3145	SERPINB4	134
19	3270755	3291144	20389	gain	3205		135
19	46386511	46388364	1853	hom_loss	3175		136
19	52496536	52501292	4756	gain	3124		137
19	55247874	55252420	4546	het_loss	3163	FLJ26850	138
19	55247874	55252420	4546	het_loss	3173	FLJ26850	138
19	55247874	55252420	4546	het_loss	3192	FLJ26850	138
19	55247874	55252420	4546	het_loss	3200	FLJ26850	138
19	55247874	55252420	4546	het_loss	3280	FLJ26850	138
19	55247874	55252420	4546	het_loss	3163	FLJ26850	138
19	55247874	55252420	4546	het_loss	3173	FLJ26850	138
19	55247874	55252420	4546	het_loss	3192	FLJ26850	138
19	55250187	55252420	2233	het_loss	3194	FLJ26850	139
19	55247874	55252420	4546	het_loss	3200	FLJ26850	138
19	55247874	55252420	4546	het_loss	3280	FLJ26850	138
19	55250187	55252420	2233	hom_loss	3175	FLJ26850	139
19	55250187	55252420	2233	hom_loss	3202	FLJ26850	139
19	56964168	57308449	344281	gain	3155	FPR2, FPR3, ZNF350, ZNF432, ZNF577, ZNF613, ZNF614, ZNF615, ZNF649, ZNF841	140
19	56964168	57308449	344281	gain	3157	FPR2, FPR3, ZNF350, ZNF432, ZNF577, ZNF613, ZNF614, ZNF615, ZNF649, ZNF841	140
19	59013780	59023850	10070	het_loss	3117	NLRP12	141
19	59249279	59251831	2552	hom_loss	3160	VSTM1	142
19	59249279	59251831	2552	hom_loss	3164	VSTM1	142
19	59250742	59251831	1089	hom_loss	3117	VSTM1	143
19	59249279	59251831	2552	hom_loss	3160	VSTM1	142
19	59249279	59251831	2552	hom_loss	3164	VSTM1	142
19	59250742	59251831	1089	hom_loss	3277	VSTM1	143
20	17844577	17954650	110073	gain	3166	MGME1, OVOL2, SNORD17, SNX5	144
20	42706680	42711434	4754	het_loss	3125	ADA	145
21	15234620	15312960	78340	gain	3009	NRIP1	146
21	29643302	29647950	4648	het_loss	3202	BACH1	147
21	44634707	44666832	32125	gain	3200	TRPM2	148
21	44634707	44641658	6951	gain	3205	TRPM2	149
21	44634707	44671482	36775	gain	3279	TRPM2	150
21	44637544	44669596	32052	gain	3127	TRPM2	151
21	44637544	44657372	19828	gain	3185	TRPM2	152
21	44634707	44666832	32125	gain	3200	TRPM2	148
21	44634707	44641658	6951	gain	3205	TRPM2	149
21	44634707	44671482	36775	gain	3279	TRPM2	150
21	44637544	44669596	32052	gain	3127	TRPM2	151
21	44637544	44657372	19828	gain	3185	TRPM2	152
21	44634707	44666832	32125	gain	3200	TRPM2	148
21	44634707	44671482	36775	gain	3279	TRPM2	150
21	44643974	44657372	13398	het_loss	3161	TRPM2	153
21	44637544	44669596	32052	gain	3127	TRPM2	151
21	44637544	44657372	19828	gain	3185	TRPM2	152
21	44634707	44666832	32125	gain	3200	TRPM2	148
21	44643974	44657372	13398	gain	3205	TRPM2	153

TABLE 1-continued

CNVs of interest in this study							
Chr	Original CNV Start	Original CNV Stop	Original CNV Size	CNV Type	PML Case ID	RefSeq Gene Symbol	SEQ ID
21	44634707	44671482	36775	gain	3279	TRPM2	150
21	44637544	44669596	32052	gain	3127	TRPM2	151
21	44634707	44666832	32125	gain	3200	TRPM2	148
21	44634707	44671482	36775	gain	3279	TRPM2	150
21	44637544	44669596	32052	gain	3127	TRPM2	151
21	44634707	44666832	32125	gain	3200	TRPM2	148
21	44660199	44681194	20995	gain	3205	TRPM2	154
21	44634707	44671482	36775	gain	3279	TRPM2	150
21	44637544	44669596	32052	gain	3127	TRPM2	151
21	44660199	44681194	20995	gain	3205	TRPM2	154
21	44634707	44671482	36775	gain	3279	TRPM2	150
21	44637544	44669596	32052	gain	3127	TRPM2	151
21	44660199	44681194	20995	gain	3205	TRPM2	154
21	44634707	44671482	36775	gain	3279	TRPM2	150
21	44660199	44681194	20995	gain	3205	TRPM2	154
21	45348895	45354820	5925	het_loss	3179	ADARB1	155
22	37689058	37715385	26327	gain	3169	APOBEC3A, APOBEC3A_B, APOBEC3B	156
22	39257585	39261621	4036	het_loss	3005	MKL1	157
22	40642402	40655210	12808	gain	3205	TNFRSF13C	158
22	40655820	40673250	17430	gain	3185		159
22	40655820	40675788	19968	gain	3205		160
22	40659633	40671866	12233	gain	3127		161
22	40655820	40673250	17430	gain	3185		159
22	40655820	40675788	19968	gain	3205		160
22	40659633	40671866	12233	gain	3127	CENPM	161
22	40655820	40673250	17430	gain	3185	CENPM	159
22	40663050	40668079	5029	gain	3190	CENPM	162
22	40663050	40668079	5029	gain	3202	CENPM	162
22	40655820	40675788	19968	gain	3205	CENPM	160
22	40659633	40671866	12233	gain	3127	CENPM	161
22	40655820	40673250	17430	gain	3185	CENPM	159
22	40655820	40675788	19968	gain	3205	CENPM	160
22	40655820	40673250	17430	gain	3185	CENPM	159
22	40655820	40675788	19968	gain	3205	CENPM	160
22	40655820	40675788	19968	gain	3205	CENPM	160
23	232907	244684	11777	het_loss	3007	PPP2R3B	163
23	7585301	7830994	245693	gain	3172		164
23	7585301	7830994	245693	gain	3172	VCX	164
23	7769323	7779354	10031	het_loss	3132		165
23	6465033	8093113	1628080	het_loss	3171		166
23	7769323	7779354	10031	het_loss	3204		165
23	7585301	7830994	245693	gain	3172		164
23	7585301	7830994	245693	gain	3172		164
23	6465033	8093113	1628080	het_loss	3171	MIR651, PNPLA4	166
23	7585301	7830994	245693	gain	3172	PNPLA4	164
23	48358646	48408854	50208	het_loss	3009		167
23	64710574	64725828	15254	gain	3125		168
23	73083877	73086192	2315	hom_loss	3193	JPX	169
23	73083877	73086192	2315	hom_loss	3200	JPX	169
23	122337025	122340879	3854	hom_loss	3125	GRIA3	170
23	148452844	148461889	9045	het_loss	3163		171
23	148452844	148461889	9045	het_loss	3205		171
23	148452844	148461889	9045	hom_loss	3144		171
23	148452844	148461889	9045	hom_loss	3193		171
23	149901706	149904265	2559	gain	3117	HMGB3	172
23	149901706	149904265	2559	gain	3118	HMGB3	172

[0608] Table 1 lists all CNVs of interest, obtained as described in the text, with the exception that, for each entry, the original CNV start and stop positions are noted, along with original CNV size, type (heterozygous loss, homozygous loss or gain), Case_ID and gene annotation (for the CNV-subregion NOT original CNV). The final column

contains SEQ_ID numbers. Standard chromosomal numbering used by those skilled in the art is used in Table 1 for the autosomal chromosomes (1-22) but, for convenience with analysis methods, chromosome X is designated as chromosome 23 herein. All coordinates are based on hg 18.

TABLE 2

CNV -subregions of interest in this study												
Chr	CNV Subregion Start	CNV Subregion Stop	CNV Sub-region Size	CNV Type	PML Case ID	RefSeq Gene Symbol	Exon overlap	NVE cases	PML cases	FET	OR	CNV Sub-region No (SRN)
1	1086119	1135772	49653	het_loss	3009	MIR200A, MIR200B, MIR429, TNFRSF18, TLL10	Y	0	1	0.005115965	39.43	1
1	9634094	9635206	1112	hom_loss	3009	PIK3CD	Y	0	1	0.005115965	39.43	2
1	12018512	12032581	14069	gain	3205		N	0	1	0.005115965	39.43	3
1	19593401	19602807	9406	het_loss	3203	CAPZB	N	0	1	0.005115965	39.43	4
1	21698753	21700243	1490	het_loss	3161		N	0	1	0.005115965	39.43	5
1	24364786	24391166	26380	gain	3199	IFNLR1	Y	0	1	0.005115965	39.43	6
1	28666669	28737671	71002	gain	3161	PHACTR4, RCC1, SNHG3	Y	0	1	0.005115965	39.43	7
1	49372054	49380088	8034	het_loss	3145	AGBL4	N	0	1	0.005115965	39.43	8
1	153816159	153827698	11539	het_loss	3168		N	0	1	0.005115965	39.43	9
1	205607255	205610341	3086	gain	3007		N	0	1	0.005115965	39.43	10
1	215760485	215762451	1966	het_loss	3117	GPATCH2	N	0	1	0.005115965	39.43	11
1	215866737	215869900	3163	het_loss	3151	GPATCH2	N	0	1	0.005115965	39.43	12
2	10352668	10356083	3415	het_loss	3007		N	0	1	0.005115965	39.43	13
2	24457024	24462631	5607	hom_loss	3204		N	0	1	0.005115965	39.43	14
2	38468717	38471950	3233	het_loss	3175		N	0	1	0.005115965	39.43	15
2	38516138	38524237	8099	het_loss	3151		N	0	1	0.005115965	39.43	16
2	38726517	38731845	5328	het_loss	3159		N	0	1	0.005115965	39.43	17
2	40620890	40624089	3199	het_loss	3202		N	0	1	0.005115965	39.43	18
2	46631006	46643501	12495	gain	3145	RHOQ	N	0	1	0.005115965	39.43	19
2	55764753	55771586	6833	gain	3143	PNPT1	Y	1	3	0.001318303	40.7	20
2	55764753	55771586	6833	gain	3193	PNPT1	Y	1	3	0.001318303	40.7	21
2	55764753	55771586	6833	gain	3282	PNPT1	Y	1	3	0.001318303	40.7	22
2	55771587	55772965	1378	gain	3143	PNPT1	N	2	3	0.003126725	20.33	23
2	55771587	55772965	1378	gain	3193	PNPT1	N	2	3	0.003126725	20.33	24
2	55771587	55772965	1378	gain	3282	PNPT1	N	2	3	0.003126725	20.33	25
2	55772966	55790559	17593	gain	3143	PNPT1	Y	1	3	0.001318303	40.7	26
2	55772966	55790559	17593	gain	3193	PNPT1	Y	1	3	0.001318303	40.7	27
2	55772966	55790559	17593	gain	3282	PNPT1	Y	1	3	0.001318303	40.7	28
2	71190677	71191310	633	het_loss	3175	MCEE	Y	0	1	0.005115965	39.43	29
2	71191311	71198107	6796	het_loss	3175	MCEE	N	1	2	0.014314826	26.77	30
2	71191311	71198107	6796	het_loss	3204	MCEE	N	1	2	0.014314826	26.77	31
2	71198108	71200120	2012	het_loss	3143	MCEE	N	2	5	3.02E-05	34.83	32
2	71198108	71200120	2012	het_loss	3175	MCEE	N	2	5	3.02E-05	34.83	33
2	71198108	71200120	2012	het_loss	3193	MCEE	N	2	5	3.02E-05	34.83	34
2	71198108	71200120	2012	het_loss	3200	MCEE	N	2	5	3.02E-05	34.83	35
2	71198108	71200120	2012	het_loss	3204	MCEE	N	2	5	3.02E-05	34.83	36
2	74827730	74913493	85763	gain	3118	HK2	Y	0	1	0.005115965	39.43	37
2	105418748	105435274	16526	het_loss	3193	FHL2	Y	0	1	0.005115965	39.43	38
2	110182348	110210249	27901	gain	3174	MALL, MIR4267, MIR4436B1, MIR4436B2	Y	2	1	0.198831257	6.6	39
2	127823042	127828410	5368	het_loss	3273		N	0	1	0.005115965	39.43	40
2	134911636	134914254	2618	het_loss	3273	MGAT5	N	0	1	0.005115965	39.43	41
2	203005216	203019933	14717	het_loss	3009	BMPT2	N	2	2	0.02731135	13.37	42
2	203005216	203019933	14717	het_loss	3192	BMPT2	N	2	2	0.02731135	13.37	43
2	203005216	203019933	14717	hom_loss	3152	BMPT2	N	0	1	0.005115965	39.43	44
2	230212897	230216339	3442	het_loss	3154	DNER	N	0	1	0.005115965	39.43	45
3	122979920	122994402	14482	gain	3202	IQCB1	Y	0	1	0.005115965	39.43	46
4	26565071	26566345	1274	het_loss	3010	STIM2	N	85	5	0.671895631	0.75	47
4	26565071	26566345	1274	het_loss	3125	STIM2	N	85	5	0.671895631	0.75	48
4	26565071	26566345	1274	het_loss	3168	STIM2	N	85	5	0.671895631	0.75	49
4	26565071	26566345	1274	het_loss	3282	STIM2	N	85	5	0.671895631	0.75	50
4	26565071	26566345	1274	het_loss	3284	STIM2	N	85	5	0.671895631	0.75	51
4	26565071	26566345	1274	hom_loss	3273	STIM2	N	1	1	0.13732578	13.21	52

TABLE 2-continued

CNV -subregions of interest in this study												
Chr	CNV Subregion Start	CNV Subregion Stop	CNV Sub-region Size	CNV Type	PML Case ID	RefSeq Gene Symbol	Exon overlap	NVE cases	PML cases	FET	OR	CNV Sub-region No (SRN)
4	54838623	54873909	35286	gain	3153	PDGFRA	Y	0	1	0.005115965	39.43	53
4	90791460	90843887	52427	gain	3168		N	0	1	0.005115965	39.43	54
4	90800863	90808258	7395	het_loss	3009		N	0	2	0.005115965	66.59	55
1	90800863	90808258	7395	het_loss	3284		N	0	2	0.005115965	66.59	56
5	45331278	45785151	453873	gain	3157	HCN1	Y	0	1	0.005115965	39.43	57
5	49771219	49774457	3238	gain	3273	EMB	Y	0	1	0.005115965	39.43	58
5	66619415	66636116	16701	gain	3010		N	0	1	0.005115965	39.43	59
5	78480194	78497296	17102	gain	3205		N	0	1	0.005115965	39.43	60
5	78497296	78500551	3255	gain	3132		N	0	2	0.005115965	66.59	61
5	78497296	78500551	3255	gain	3185		N	0	2	0.005115965	66.59	62
5	78500552	78521408	20856	gain	3132		N	0	3	2.49E-05	94.48	63
5	78500552	78521408	20856	gain	3185		N	0	3	2.49E-05	94.48	64
5	78500552	78521408	20856	gain	3205		N	0	3	2.49E-05	94.48	65
5	78521409	78526637	5228	gain	3132		N	0	2	0.005115965	66.59	66
5	78521409	78526637	5228	gain	3205		N	0	2	0.005115965	66.59	67
5	78526638	78531091	4453	gain	3132		N	0	1	0.005115965	39.43	68
5	83490494	83495169	4675	het_loss	3204	EDIL3	N	0	1	0.005115965	39.43	69
5	133372071	133379727	7656	hom_loss	3153		N	0	1	0.005115965	39.43	70
5	137836466	137843309	6843	hom_loss	3279		N	1	1	0.13732578	13.21	71
5	150159466	150161037	1571	het_loss	3117		N	15	4	0.040487703	3.62	72
5	150159466	150161037	1571	het_loss	3180		N	15	4	0.040487703	3.62	73
5	150159466	150161037	1571	het_loss	3199		N	15	4	0.040487703	3.62	74
5	150159466	150161037	1571	het_loss	3278		N	15	4	0.040487703	3.62	75
5	150161038	150181399	20361	het_loss	3117		N	14	4	0.033744017	3.88	76
5	150161038	150181399	20361	het_loss	3180		N	14	4	0.033744017	3.88	77
5	150161038	150181399	20361	het_loss	3199		N	14	4	0.033744017	3.88	78
5	150161038	150181399	20361	het_loss	3278		N	14	4	0.033744017	3.88	79
5	150181400	150185189	3789	het_loss	3117		N	13	4	0.027710312	4.18	80
5	150181400	150185189	3789	het_loss	3180		N	13	4	0.027710312	4.18	81
5	150181400	150185189	3789	het_loss	3199		N	13	4	0.027710312	4.18	82
5	150181400	150185189	3789	het_loss	3278		N	13	4	0.027710312	4.18	83
5	150185190	150191626	6436	hom_loss	3009		N	6	7	8.59E-06	16.65	84
5	150185190	150191626	6436	hom_loss	3143		N	6	7	8.59E-06	16.65	85
5	150185190	150191626	6436	hom_loss	3152		N	6	7	8.59E-06	16.65	86
5	150185190	150191626	6436	hom_loss	3154		N	6	7	8.59E-06	16.65	87
5	150185190	150191626	6436	hom_loss	3193		N	6	7	8.59E-06	16.65	88
5	150185190	150191626	6436	hom_loss	3196		N	6	7	8.59E-06	16.65	89
5	150185190	150191626	6436	hom_loss	3281		N	6	7	8.59E-06	16.65	90
5	150191627	150201145	9518	hom_loss	3009		N	6	7	8.59E-06	16.65	91
5	150191627	150201145	9518	hom_loss	3143		N	6	7	8.59E-06	16.65	92
5	150191627	150201145	9518	hom_loss	3152		N	6	7	8.59E-06	16.65	93
5	150191627	150201145	9518	hom_loss	3154		N	6	7	8.59E-06	16.65	94
5	150191627	150201145	9518	hom_loss	3193		N	6	7	8.59E-06	16.65	95
5	150191627	150201145	9518	hom_loss	3196		N	6	7	8.59E-06	16.65	96
5	150191627	150201145	9518	hom_loss	3281		N	6	7	8.59E-06	16.65	97
5	150201146	150202601	1455	hom_loss	3152		N	1	3	0.001318303	40.7	98
5	150201146	150202601	1455	hom_loss	3154		N	1	3	0.001318303	40.7	99
5	150201146	150202601	1455	hom_loss	3193		N	1	3	0.001318303	40.7	100
5	150202602	150204134	1532	het_loss	3132		N	51	8	0.062987683	2.17	101
5	150202602	150204134	1532	het_loss	3180		N	51	8	0.062987683	2.17	102
5	150202602	150204134	1532	het_loss	3196		N	51	8	0.062987683	2.17	103
5	150202602	150204134	1532	het_loss	3273		N	51	8	0.062987683	2.17	104
5	150202602	150204134	1532	het_loss	3277		N	51	8	0.062987683	2.17	105
5	150202602	150204134	1532	het_loss	3278		N	51	8	0.062987683	2.17	106
5	150202602	150204134	1532	het_loss	3280		N	51	8	0.062987683	2.17	107
5	150202602	150204134	1532	het_loss	3282		N	51	8	0.062987683	2.17	108
5	179590681	179626660	35979	het_loss	3172	MAPK9	Y	0	1	0.005115965	39.43	109
6	2882577	2947403	64826	het_loss	3196	DKFZP686I15217, NQO2, SERPINB6	Y	0	1	0.005115965	39.43	110
6	2964646	2966011	1365	het_loss	3193	HTATSF1P2, NQO2	Y	0	1	0.005115965	39.43	111
6	51766024	51773250	7226	het_loss	3167	PKHD1	N	0	1	0.005115965	39.43	112
6	51952217	51953475	1258	gain	3127	PKHD1	N	0	1	0.005115965	39.43	113
6	51953476	51965723	12247	gain	3127	PKHD1	N	0	2	0.005115965	66.59	114
6	51953476	51965723	12247	gain	3205	PKHD1	N	0	2	0.005115965	66.59	115
6	51965724	51969378	3654	gain	3127	PKHD1	N	0	1	0.005115965	39.43	116
6	74396294	74398409	2115	het_loss	3009	SLC17A5	N	0	2	0.005115965	66.59	117
6	74396294	74398409	2115	het_loss	3160	SLC17A5	N	0	2	0.005115965	66.59	118

TABLE 2-continued

CNV -subregions of interest in this study												
Chr	CNV Subregion Start	CNV Subregion Stop	CNV Sub-region Size	CNV Type	PML Case ID	RefSeq Gene Symbol	Exon overlap	NVE cases	PML cases	FET	OR	CNV Sub-region No (SRN)
6	74398410	74404837	6427	het_loss	3009	SLC17A5	Y	0	1	0.005115965	39.43	119
6	86416979	86431527	14548	het_loss	3197		N	0	1	0.005115965	39.43	120
6	91131823	91135670	3847	het_loss	3171		N	0	1	0.005115965	39.43	121
6	107882367	107890605	8238	het_loss	3201	PDSS2	Y	0	1	0.005115965	39.43	122
6	166418511	166422386	3875	het_loss	3125		N	11	10	3.49E-07	13.49	123
6	166418511	166422386	3875	het_loss	3163		N	11	10	3.49E-07	13.49	124
6	166418511	166422386	3875	het_loss	3192		N	11	10	3.49E-07	13.49	125
6	166418511	166422386	3875	het_loss	3193		N	11	10	3.49E-07	13.49	126
6	166418511	166422386	3875	het_loss	3194		N	11	10	3.49E-07	13.49	127
6	166418511	166422386	3875	het_loss	3200		N	11	10	3.49E-07	13.49	128
6	166418511	166422386	3875	het_loss	3205		N	11	10	3.49E-07	13.49	129
6	166418511	166422386	3875	het_loss	3280		N	11	10	3.49E-07	13.49	130
6	166418511	166422386	3875	het_loss	3281		N	11	10	3.49E-07	13.49	131
6	166418511	166422386	3875	het_loss	3284		N	11	10	3.49E-07	13.49	132
6	166418511	166422386	3875	hom_loss	3009		N	0	3	2.49E-05	94.48	133
6	166418511	166422386	3875	hom_loss	3152		N	0	3	2.49E-05	94.48	134
6	166418511	166422386	3875	hom_loss	3175		N	0	3	2.49E-05	94.48	135
7	65741238	65768682	27444	gain	3152	KCTD7	Y	0	2	0.005115965	66.59	136
7	65741238	65768682	27444	gain	3202	KCTD7	Y	0	2	0.005115965	66.59	137
7	157174966	157177843	2877	het_loss	3009	PTPRN2	N	0	1	0.005115965	39.43	138
7	157425841	157496238	70397	gain	3189	PTPRN2	N	1	1	0.13732578	13.21	139
7	158000082	158007892	7810	het_loss	3279	PTPRN2	N	1	1	0.13732578	13.21	140
7	158007893	158010672	2779	het_loss	3279	PTPRN2	N	5	1	0.358539546	2.63	141
7	158010673	158024569	13896	het_loss	3279	MIR595, PTPRN2	Y	1	1	0.13732578	13.21	142
8	23103186	23125443	22257	het_loss	3140	TNFRSF10A	Y	0	1	0.005115965	39.43	143
8	39914488	39919594	5106	het_loss	3126	IDO2	N	0	1	0.005115965	39.43	144
8	79905654	79910286	4632	het_loss	3159		N	0	1	0.005115965	39.43	145
8	99790200	99799839	9639	het_loss	3006	STK3	N	0	1	0.005115965	39.43	146
8	102049360	102064431	15071	het_loss	3173		N	0	3	2.49E-05	94.48	147
8	102049360	102064431	15071	het_loss	3175		N	0	3	2.49E-05	94.48	148
8	102049360	102064431	15071	het_loss	3282		N	0	3	2.49E-05	94.48	149
9	571398	580721	9323	het_loss	3006	KANK1	N	2	1	0.198831257	6.6	150
9	580722	584647	3925	het_loss	3006	KANK1	N	3	3	0.005933668	13.54	151
9	580722	584647	3925	het_loss	3200	KANK1	N	3	3	0.005933668	13.54	152
9	580722	584647	3925	het_loss	3282	KANK1	N	3	3	0.005933668	13.54	153
9	584648	598488	13840	het_loss	3200	KANK1	N	2	2	0.02731135	13.37	154
9	584648	598488	13840	het_loss	3282	KANK1	N	2	2	0.02731135	13.37	155
9	634039	637589	3550	het_loss	3273	KANK1	N	0	2	0.005115965	66.59	156
9	634039	637589	3550	het_loss	3282	KANK1	N	0	2	0.005115965	66.59	157
9	74050088	74059447	9359	het_loss	3165	GDA	Y	0	1	0.005115965	39.43	158
9	93140394	93447826	307432	gain	3198	AUH, MIR3163, MIR3910-1, MIR3910-2, NFIL3	Y	0	1	0.005115965	39.43	159
9	118564159	118575633	11474	gain	3193	ASTN2	N	0	1	0.005115965	39.43	160
9	118657526	118664593	7067	het_loss	3144	ASTN2	N	0	1	0.005115965	39.43	161
9	119220847	119233078	12231	gain	3005		N	0	1	0.005115965	39.43	162
10	899657	1071401	171744	gain	3161	GTPBP4, IDI2, IDI2-AS1, LARP4B	Y	0	1	0.005115965	39.43	163
10	76217585	76411591	194006	gain	3179	KAT6B	Y	0	1	0.005115965	39.43	164
10	116000069	116004388	4319	gain	3010	VWA2	Y	0	1	0.005115965	39.43	165
11	14677012	14689025	12013	het_loss	3199	PDE3B	N	0	1	0.005115965	39.43	166
11	34608313	34615878	7565	het_loss	3117	EHF	Y	0	1	0.005115965	39.43	167
11	62382087	62398462	16375	het_loss	3205	SLC3A2	Y	0	1	0.005115965	39.43	168
11	76631014	76643625	12611	het_loss	3193	GDPD4	Y	0	1	0.005115965	39.43	169
12	11616557	12114030	497473	het_loss	3126	ETV6	Y	0	1	0.005115965	39.43	170
12	12438904	12778142	339238	het_loss	3126	APOLD1, CDKN1B, CREBL2, DUSP16, GPR19, LOH12CR1	Y	0	1	0.005115965	39.43	171
12	12968705	12971310	2605	gain	3127		N	0	1	0.005115965	39.43	172
12	91845527	92201342	355815	het_loss	3126	EEA1, LOC643339	Y	0	1	0.005115965	39.43	173
12	92215898	92567120	351222	het_loss	3126	LOC643339, MRPL42, NUDT4, NUDT4P1,	Y	0	1	0.005115965	39.43	174

TABLE 2-continued

CNV -subregions of interest in this study												
Chr	CNV Subregion Start	CNV Subregion Stop	CNV Sub-region Size	CNV Type	PML Case ID	RefSeq Gene Symbol	Exon overlap	NVE cases	PML cases	FET	OR	CNV Sub-region No (SRN)
12	92568362	93307172	738810	het_loss	3126	SOCS2, SOCS2-AS1, UBE2N						
						CCDC41, CRADD, PLXNC1	Y	0	1	0.005115965	39.43	175
12	111061085	111064486	3401	het_loss	3004	TRAFD1	Y	0	1	0.005115965	39.43	176
13	40939924	41026908	86984	gain	3140	RGCC	Y	0	1	0.005115965	39.43	177
13	75006025	75016304	10279	gain	3009	COMMD6	Y	0	2	0.005115965	66.59	178
13	75006025	75016304	10279	gain	3152	COMMD6	Y	0	2	0.005115965	66.59	179
13	91811087	91814369	3282	het_loss	3143	GPC5	N	1	1	0.13732578	13.21	180
13	91811087	91811118	31	hom_loss	3173	GPC5	N	0	1	0.005115965	39.43	181
13	110754499	110778301	23802	gain	3006	ARHGEF7, TEX29	Y	0	1	0.005115965	39.43	182
14	20021118	20055469	34351	gain	3205	RNASE10	Y	0	1	0.005115965	39.43	183
14	20426824	20481852	55028	hom_loss	3200	ECRP, RNASE3	Y	0	1	0.005115965	39.43	184
14	20430810	20458350	27540	het_loss	3192	ECRP	Y	3	1	0.256004559	4.39	185
14	20458351	20481852	23501	het_loss	3192		N	4	1	0.309147091	3.29	186
14	20481853	20490129	8276	het_loss	3192		N	1	1	0.13732578	13.21	187
14	21096689	21105611	8922	het_loss	3125		N	0	5	1.16E-07	152.56	188
14	21096689	21105611	8922	het_loss	3175		N	0	5	1.16E-07	152.56	189
14	21096689	21105611	8922	het_loss	3194		N	0	5	1.16E-07	152.56	190
14	21096689	21105611	8922	het_loss	3204		N	0	5	1.16E-07	152.56	191
14	21096689	21105611	8922	het_loss	3273		N	0	5	1.16E-07	152.56	192
14	21120750	21125513	4763	gain	3143		N	1	2	0.014314826	26.77	193
14	21120750	21125513	4763	gain	3173		N	1	2	0.014314826	26.77	194
14	60901636	60909492	7856	het_loss	3193	PRKCH	N	0	1	0.005115965	39.43	195
14	60912874	60921269	8395	het_loss	3174	PRKCH	N	0	1	0.005115965	39.43	196
14	63937192	63944459	7267	gain	3205	MTHFD1	Y	0	1	0.005115965	39.43	197
14	95754535	95759056	4521	het_loss	3009	BDKRB2	N	0	3	2.49E-05	94.48	198
14	95754535	95759056	4521	het_loss	3173	BDKRB2	N	0	3	2.49E-05	94.48	199
14	95754535	95759056	4521	het_loss	3202	BDKRB2	N	0	3	2.49E-05	94.48	200
15	66065925	66082418	16493	het_loss	3010		N	0	1	0.005115965	39.43	201
15	70432627	70443017	10390	gain	3169	HEXA	Y	0	1	0.005115965	39.43	202
15	75096101	75101523	5422	gain	3200	PSTPIP1	Y	0	1	0.005115965	39.43	203
15	75101524	75105788	4264	gain	3132	PSTPIP1	Y	0	2	0.005115965	66.59	204
15	75101524	75105788	4264	gain	3200	PSTPIP1	Y	0	2	0.005115965	66.59	205
15	75105789	75109086	3297	gain	3127	PSTPIP1	Y	0	5	1.16E-07	152.56	206
15	75105789	75109086	3297	gain	3132	PSTPIP1	Y	0	5	1.16E-07	152.56	207
15	75105789	75109086	3297	gain	3199	PSTPIP1	Y	0	5	1.16E-07	152.56	208
15	75105789	75109086	3297	gain	3200	PSTPIP1	Y	0	5	1.16E-07	152.56	209
15	75105789	75109086	3297	gain	3279	PSTPIP1	Y	0	5	1.16E-07	152.56	210
15	75109087	75115806	6719	gain	3127	PSTPIP1	Y	1	5	9.14E-06	69.72	211
15	75109087	75115806	6719	gain	3132	PSTPIP1	Y	1	5	9.14E-06	69.72	212
15	75109087	75115806	6719	gain	3199	PSTPIP1	Y	1	5	9.14E-06	69.72	213
15	75109087	75115806	6719	gain	3200	PSTPIP1	Y	1	5	9.14E-06	69.72	214
15	75109087	75115806	6719	gain	3279	PSTPIP1	Y	1	5	9.14E-06	69.72	215
15	75115807	75117798	1991	gain	3200	PSTPIP1	Y	1	1	0.13732578	13.21	216
15	88999998	89016848	16850	het_loss	3172		N	0	1	0.005115965	39.43	217
16	6823677	6884976	61299	het_loss	3126	RBFOX1	N	0	1	0.005115965	39.43	218
16	6886815	6896330	9515	het_loss	3126	RBFOX1	N	0	1	0.005115965	39.43	219
16	6942078	6945539	3461	gain	3173	RBFOX1	N	1	3	0.001318303	40.7	220
16	6942078	6945539	3461	gain	3175	RBFOX1	N	1	3	0.001318303	40.7	221
16	6942078	6945539	3461	gain	3282	RBFOX1	N	1	3	0.001318303	40.7	222
16	23844022	23848772	4750	het_loss	3198	PRKCB	N	7	1	0.447101793	1.88	223
16	23892842	23893968	1126	gain	3199	PRKCB	N	0	1	0.005115965	39.43	224
16	23893969	23903495	9526	gain	3199	PRKCB	N	0	2	0.005115965	66.59	225
16	23893969	23903495	9526	gain	3205	PRKCB	N	0	2	0.005115965	66.59	226
16	23903496	23908248	4752	gain	3205	PRKCB	Y	0	1	0.005115965	39.43	227
16	69047888	69050151	2263	gain	3174	FUK	N	0	2	0.005115965	66.59	228
16	69047888	69050151	2263	gain	3185	FUK	N	0	2	0.005115965	66.59	229
16	69052450	69081640	29190	het_loss	3197	COG4, FUK	Y	0	1	0.005115965	39.43	230
16	70653499	70665447	11948	gain	3143	HPR	Y	0	6	1.16E-07	182.82	231
16	70653499	70665447	11948	gain	3152	HPR	Y	0	6	1.16E-07	182.82	232
16	70653499	70665447	11948	gain	3192	HPR	Y	0	6	1.16E-07	182.82	233
16	70653499	70665447	11948	gain	3200	HPR	Y	0	6	1.16E-07	182.82	234
16	70653499	70665447	11948	gain	3282	HPR	Y	0	6	1.16E-07	182.82	235
16	70653499	70665447	11948	gain	3284	HPR	Y	0	6	1.16E-07	182.82	236
17	69341925	70202523	860598	gain	3183	BTBD17,	Y	1	1	0.13732578	13.21	237

TABLE 2-continued

CNV -subregions of interest in this study												
Chr	CNV Subregion Start	CNV Subregion Stop	CNV Sub-region Size	CNV Type	PML Case ID	RefSeq Gene Symbol	Exon overlap	NVE cases	PML cases	FET	OR	CNV Sub-region No (SRN)
						C17orf77, CD300A, CD300C, CD300E, CD300LB, CD300LD, CD300LF, DNAI2, GPR142, GPRC5C, KIF19, MGC16275, RAB37, RPL38, TTYH2						
17	75608151	75611602	3451	het_loss	3144	TBC1D16	N	1	7	5.37E-08	100.4	238
17	75608151	75611602	3451	het_loss	3152	TBC1D16	N	1	7	5.37E-08	100.4	239
17	75608151	75611602	3451	het_loss	3163	TBC1D16	N	1	7	5.37E-08	100.4	240
17	75608151	75611602	3451	het_loss	3192	TBC1D16	N	1	7	5.37E-08	100.4	241
17	75608151	75611602	3451	het_loss	3200	TBC1D16	N	1	7	5.37E-08	100.4	242
17	75608151	75611602	3451	het_loss	3204	TBC1D16	N	1	7	5.37E-08	100.4	243
17	75608151	75611602	3451	het_loss	3284	TBC1D16	N	1	7	5.37E-08	100.4	244
17	75608151	75611602	3451	hom_loss	3009	TBC1D16	N	0	1	0.005115965	39.43	245
17	75611602	75615433	3831	hom_loss	3175	TBC1D16	N	0	1	0.005115965	39.43	246
17	75611603	75615433	3830	het_loss	3144	TBC1D16	N	1	4	0.000112689	55.01	247
17	75611603	75615433	3830	het_loss	3152	TBC1D16	N	1	4	0.000112689	55.01	248
17	75611603	75615433	3830	het_loss	3163	TBC1D16	N	1	4	0.000112689	55.01	249
17	75611603	75615433	3830	het_loss	3200	TBC1D16	N	1	4	0.000112689	55.01	250
17	76241510	76247304	5794	gain	3205	RPTOR	N	10	1	0.005115965	39.43	251
17	76247305	76265683	18378	gain	3127	RPTOR	N	0	2	0.005115965	66.59	252
17	76247305	76265683	18378	gain	3205	RPTOR	N	0	2	0.005115965	66.59	253
17	76265684	76267844	2160	gain	3205	RPTOR	N	0	1	0.005115965	39.43	254
18	9985530	10125331	139801	gain	3175		N	0	1	0.005115965	39.43	255
18	12764095	12781985	17890	gain	3191	PTPN2	Y	0	1	0.005115965	39.43	256
18	27026203	27029351	3148	het_loss	3125		N	0	3	2.49E-05	94.48	257
18	27026203	27029351	3148	het_loss	3143		N	0	3	2.49E-05	94.48	258
18	27026203	27029351	3148	het_loss	3175		N	0	3	2.49E-05	94.48	259
18	42537949	42663605	125656	gain	3125	PIAS2, ST8SIA5	Y	0	1	0.005115965	39.43	260
18	46917195	46945018	27823	het_loss	3161		N	0	1	0.005115965	39.43	261
18	59457622	59465699	8077	het_loss	3145	SERPINB4	Y	0	1	0.005115965	39.43	262
19	3270755	3291144	20389	gain	3205		N	0	1	0.005115965	39.43	263
19	46386511	46388364	1853	hom_loss	3175		N	0	1	0.005115965	39.43	264
19	52496536	52501292	4756	gain	3124		N	0	1	0.005115965	39.43	265
19	55247874	55250186	2312	het_loss	3163	FLJ26850	N	4	5	0.000161709	17.38	266
19	55247874	55250186	2312	het_loss	3173	FLJ26850	N	4	5	0.000161709	17.38	267
19	55247874	55250186	2312	het_loss	3192	FLJ26850	N	4	5	0.000161709	17.38	268
19	55247874	55250186	2312	het_loss	3200	FLJ26850	N	4	5	0.000161709	17.38	269
19	55247874	55250186	2312	het_loss	3280	FLJ26850	N	4	5	0.000161709	17.38	270
19	55250187	55252420	2233	het_loss	3163	FLJ26850	N	4	6	1.80E-05	21.15	271
19	55250187	55252420	2233	het_loss	3173	FLJ26850	N	4	6	1.80E-05	21.15	272
19	55250187	55252420	2233	het_loss	3192	FLJ26850	N	4	6	1.80E-05	21.15	273
19	55250187	55252420	2233	het_loss	3194	FLJ26850	N	4	6	1.80E-05	21.15	274
19	55250187	55252420	2233	het_loss	3200	FLJ26850	N	4	6	1.80E-05	21.15	275
19	55250187	55252420	2233	het_loss	3280	FLJ26850	N	4	6	1.80E-05	21.15	276
19	55250187	55252420	2233	hom_loss	3175	FLJ26850	N	0	2	0.005115965	66.59	277
19	55250187	55252420	2233	hom_loss	3202	FLJ26850	N	0	2	0.005115965	66.59	278
19	56964168	57308449	344281	gain	3155	FPR2, FPR3, ZNF350, ZNF432, ZNF577, ZNF613, ZNF614, ZNF615, ZNF649, ZNF841	Y	3	2	0.043434433	8.91	279
19	56964168	57308449	344281	gain	3157	FPR2, FPR3, ZNF350, ZNF432, ZNF577, ZNF613, ZNF614, ZNF615, ZNF649, ZNF841	Y	3	2	0.043434433	8.91	280
19	59016855	59023850	6995	het_loss	3117	NLRP12	Y	0	1	0.005115965	39.43	281
19	59249279	59250741	1462	hom_loss	3160	VSTM1	N	37	2	1	0.7	282
19	59249279	59250741	1462	hom_loss	3164	VSTM1	N	37	2	1	0.7	283

TABLE 2-continued

CNV -subregions of interest in this study												
Chr	CNV Subregion Start	CNV Subregion Stop	CNV Sub-region Size	CNV Type	PML Case ID	RefSeq Gene Symbol	Exon overlap	NVE cases	PML cases	FET	OR	CNV Sub-region No (SRN)
19	59250742	59251831	1089	hom_loss	3117	VSTM1	N	38	4	0.533838399	1.39	284
19	59250742	59251831	1089	hom_loss	3160	VSTM1	N	38	1	0.533838399	1.39	285
19	59250742	59251831	1089	hom_loss	3164	VSTM1	N	38	4	0.533838399	1.39	286
19	59250742	59251831	1089	hom_loss	3277	VSTM1	N	38	4	0.533838399	1.39	287
20	17844577	17954650	110073	gain	3166	MGME1, OVOL2, SNORD17, SNX5	Y	0	1	0.005115965	39.43	288
20	42706680	42711434	4754	het_loss	3125	ADA	N	0	1	0.005115965	39.43	289
21	15237071	15312960	75889	gain	3009	NRIP1	Y	0	1	0.005115965	39.43	290
21	29643302	29647950	4648	het_loss	3202	BACH1	Y	0	1	0.005115965	39.43	291
21	44634707	44637543	2836	gain	3200	TRPM2	Y	1	3	0.001318303	40.7	292
21	44634707	44637543	2836	gain	3205	TRPM2	Y	1	3	0.001318303	40.7	293
21	44634707	44637543	2836	gain	3279	TRPM2	Y	1	3	0.001318303	40.7	294
21	44637544	44641658	4114	gain	3127	TRPM2	Y	1	5	9.14E-06	69.72	295
21	44637544	44641658	4114	gain	3185	TRPM2	Y	1	5	9.14E-06	69.72	296
21	44637544	44641658	4114	gain	3200	TRPM2	Y	1	5	9.14E-06	69.72	297
21	44637544	44641658	4114	gain	3205	TRPM2	Y	1	5	9.14E-06	69.72	298
21	44637544	44641658	4114	gain	3279	TRPM2	Y	1	5	9.14E-06	69.72	299
21	44641659	44643973	2314	gain	3127	TRPM2	Y	1	1	0.000112689	55.01	300
21	44641659	44643973	2314	gain	3185	TRPM2	Y	1	4	0.000112689	55.01	301
21	44641659	44643973	2314	gain	3200	TRPM2	Y	1	4	0.000112689	55.01	302
21	44641659	44643973	2314	gain	3279	TRPM2	Y	1	4	0.000112689	55.01	303
21	44643974	44657372	13398	het_loss	3161	TRPM2	Y	1	1	0.13732578	13.21	304
21	44643975	44657372	13397	gain	3127	TRPM2	Y	0	5	1.16E-07	152.56	305
21	44643975	44657372	13397	gain	3185	TRPM2	Y	0	5	1.16E-07	152.56	306
21	44643975	44657372	13397	gain	3200	TRPM2	Y	0	5	1.16E-07	152.56	307
21	44643975	44657372	13397	gain	3205	TRPM2	Y	0	5	1.16E-07	152.56	308
21	44643975	44657372	13397	gain	3279	TRPM2	Y	0	5	1.16E-07	152.56	309
21	44657373	44660198	2825	gain	3127	TRPM2	Y	0	3	2.49E-05	94.48	310
21	44657373	44660198	2825	gain	3200	TRPM2	Y	0	3	2.49E-05	94.48	311
21	44657373	44660198	2825	gain	3279	TRPM2	Y	0	3	2.49E-05	94.48	312
21	44660199	44666832	6633	gain	3127	TRPM2	Y	0	4	2.49E-05	123.12	313
21	44660199	44666832	6633	gain	3200	TRPM2	Y	0	4	2.49E-05	123.12	314
21	44660199	44666832	6633	gain	3205	TRPM2	Y	0	4	2.49E-05	123.12	315
21	44660199	44666832	6633	gain	3279	TRPM2	Y	0	4	2.49E-05	123.12	316
21	44666833	44669596	2763	gain	3127	TRPM2	Y	0	3	2.49E-05	94.48	317
21	44666833	44669596	2763	gain	3205	TRPM2	Y	0	3	2.49E-05	94.48	318
21	44666833	44669596	2763	gain	3279	TRPM2	Y	0	3	2.49E-05	94.48	319
21	44669597	44671482	1885	gain	3205	TRPM2	Y	0	2	0.005115965	66.59	320
21	44669597	44671482	1885	gain	3279	TRPM2	Y	0	2	0.005115965	66.59	321
21	44671483	44681194	9711	gain	3205	TRPM2	Y	0	1	0.005115965	39.43	322
21	45348895	45354820	5925	het_loss	3179	ADARB1	N	0	1	0.005115965	39.43	323
22	37689058	37715385	26327	gain	3169	APOBEC3A, APOBEC3A_B, APOBEC3B	Y	0	1	0.005115965	39.43	324
22	39257585	39261621	4036	het_loss	3005	MKL1	N	0	1	0.005115965	39.43	325
22	40642402	40655210	12808	gain	3205	TNFRSF13C	Y	0	1	0.005115965	39.43	326
22	40655820	40659632	3812	gain	3185		N	0	2	0.005115965	66.59	327
22	40655820	40659632	3812	gain	3205		N	0	2	0.005115965	66.59	328
22	40659633	40663049	3416	gain	3127		N	0	3	2.49E-05	94.48	329
22	40659633	40663049	3416	gain	3185		N	0	3	2.49E-05	94.48	330
22	40659633	40663049	3416	gain	3205		N	0	3	2.49E-05	94.48	331
22	40663050	40668079	5029	gain	3127	CENPM	Y	0	5	1.16E-07	152.56	332
22	40663050	40668079	5029	gain	3185	CENPM	Y	0	5	1.16E-07	152.56	333
22	40663050	40668079	5029	gain	3190	CENPM	Y	0	5	1.16E-07	152.56	334
22	40663050	40668079	5029	gain	3202	CENPM	Y	0	5	1.16E-07	152.56	335
22	40663050	40668079	5029	gain	3205	CENPM	Y	0	5	1.16E-07	152.56	336
22	40668080	40671866	3786	gain	3127	CENPM	Y	0	3	2.49E-05	94.48	337
22	40668080	40671866	3786	gain	3185	CENPM	Y	0	3	2.49E-05	94.48	338
22	40668080	40671866	3786	gain	3205	CENPM	Y	0	3	2.49E-05	94.48	339
22	40671867	40673250	1383	gain	3185	CENPM	Y	0	2	0.005115965	66.59	340
22	40671867	40673250	1383	gain	3205	CENPM	Y	0	2	0.005115965	66.59	341
22	40673251	40675788	2537	gain	3205		N	0	1	0.005115965	39.43	342
23	2329207	234429	1522	het_loss	3007	PPP2R3B	N	0	1	0.005115965	39.43	343
23	7585301	7769322	184021	gain	3172		N	5	1	0.358539546	2.63	344
23	7769323	7773949	4626	gain	3172	VCX	Y	7	1	0.447101793	1.88	345
23	7773982	7779354	5372	het_loss	3132		N	0	3	2.49E-05	94.48	346
23	7773982	7779354	5372	het_loss	3171		N	0	3	2.49E-05	94.48	347
23	7773982	7779354	5372	het_loss	3204		N	0	3	2.49E-05	94.48	348
23	7773982	7779353	5371	gain	3172		N	5	1	0.358539546	2.63	349

TABLE 2-continued

CNV -subregions of interest in this study												
Chr	CNV Subregion Start	CNV Subregion Stop	CNV Sub-region Size	CNV Type	PML Case ID	RefSeq Gene Symbol	Exon overlap	NVE cases	PML cases	FET	OR	CNV Sub-region No (SRN)
23	7779354	7815400	36046	gain	3172	MIR651, PNPLA4	N	6	1	0.404443314	2.19	350
23	7779355	8093113	313758	het_loss	3171		Y	0	1	0.005115965	39.43	351
23	7815401	7830994	15593	gain	3172		Y	7	1	0.447101793	1.88	352
23	48358646	48408854	50208	het_loss	3009		N	0	1	0.005115965	39.43	353
23	64710574	64725828	15254	gain	3125		N	0	1	0.005115965	39.43	354
23	73083877	73086192	2315	hom_loss	3193	JPX	N	1	2	0.014314826	26.77	355
23	73083877	73086192	2315	hom_loss	3200	JPX	N	1	2	0.014314826	26.77	356
23	122337025	122340879	3854	hom_loss	3125	GRIA3	N	0	1	0.005115965	39.43	357
23	148452844	148461889	9045	het_loss	3163		N	7	2	0.129983268	3.8	358
23	148452844	148461889	9045	het_loss	3205		N	7	2	0.129983268	3.8	359
23	148459108	148461889	2781	hom_loss	3144		N	0	2	0.005115965	66.59	360
23	148459108	148461889	2781	hom_loss	3193		N	0	2	0.005115965	66.59	361
23	149901706	149902701	995	gain	3117	HMGB3	Y	0	2	0.005115965	66.59	362
23	149901706	149902701	995	gain	3118	HMGB3	Y	0	2	0.005115965	66.59	363

[0609] Table 2 is identical to Table 1, with a number of exceptions. Firstly, the CNV coordinates listed refer to the actual CNV-subregions found to be unique or significantly different between the disease and normal cohorts, as opposed to Table 1, which lists the original CNVs. Secondly, an extra column details whether genic CNV-subregions of interest overlap an exon or not. Third and fourth, 2 extra columns detail the number of normal cases and the number

of disease cases that harbor the relevant CNV-subregion. Finally, 2 columns report Fisher's 2-tailed Exact Test (FET) and the odds ratio (OR). Standard chromosomal numbering used by those skilled in the art is used in Table 2 for the autosomal chromosomes (1-22) but, for convenience with analysis methods, chromosome X is designated as chromosome 23 herein. All coordinates are in hg18.

TABLE 3
A non-redundant list of genes listed in Table 2

RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary	Gene # (GN)
ADARB1	intronic	104	double-stranded RNA-specific editase 1 isoform 1	This gene encodes the enzyme responsible for pre-mRNA editing of the glutamate receptor subunit B by site-specific deamination of adenosines. Studies in rat found that this enzyme acted on its own pre-mRNA molecules to convert an AA dinucleotide to an AI dinucleotide which resulted in a new splice site. Alternative splicing of this gene results in several transcript variants, some of which have been characterized by the presence or absence of an ALU cassette insert and a short or long C-terminal region. [provided by RefSeq, July 2008]. Transcript Variant: This variant (1), also known as ADAR2a-L1 or DRADA2a, lacks the ALU cassette insert and contains the long C-terminal region, as compared to variant 2. The resulting isoform (1), also known as hRED1-Short, lacks an internal segment, compared to isoform 2. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##RefSeq-Attributes-START## undergoes RNA editing :: PMID: 11717408, 12045112 ##RefSeq-Attributes-END## ##Evidence-Data-START## Transcript exon combination :: AB194370.1, U76420.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025084, ERS025085 [ECO: 0000348] ##Evidence-Data-END## N/A	2
AGBL4	intronic	84871	cytosolic carboxypeptidase 6	This gene is a member of the cytidine deaminase gene family. It is one of seven related genes or pseudogenes found in a cluster, thought to result from gene duplication, on chromosome 22. Members of the cluster encode proteins that are structurally and functionally related to the C to U RNA-editing cytidine deaminase APOBEC1. The protein encoded by this gene lacks the zinc binding activity of other family members. The protein plays a role in immunity, by restricting transmission of foreign DNA such as viruses. One mechanism of foreign DNA restriction is deamination of foreign double-stranded DNA cytidines to uridines, which leads to DNA degradation. However, other mechanisms are also thought to be involved, as anti-viral effect is not dependent on deaminase activity. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, July 2012]. Transcript Variant: This variant (1) represents the longer transcript and encodes the longer isoform (a). Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: U03891.2, BC126416.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025081, ERS025084 [ECO: 0000348] ##Evidence-Data-END##	3
APOBEC3A	exonic	200315	DNA dC->dU-editing enzyme APOBEC-3A isoform a		4
APOBEC3A_B	intronic	100913187	probable DNA dC->dU-editing enzyme APOBEC-3A		5

TABLE 3-continued

A non-redundant list of genes listed in Table 2				
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary
APOBEC3B	exonic	9582	DNA dC->dU- editing enzyme APOBEC-3B isoform a	cytidines to uridines, which leads to DNA degradation. However, other mechanisms are also thought to be involved, as anti-viral effect is not dependent on deaminase activity. The protein encoded by this gene is the same as that encoded by APOBEC3A; however, this gene is a hybrid gene that results from the deletion of approximately 29.5 kb of sequence between the APOBEC3A gene and the adjacent gene APOBEC3B. The breakpoints of the deletion are within the two genes, so the deletion hybrid is predicted to have the promoter and coding region of APOBEC3A, but the 3' UTR of APOBEC3B. [provided by RefSeq, July 2012]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## RNAseq introns :: single sample supports all introns ERS025081, ERS025084 [ECO: 0000348] ##Evidence-Data-END## This gene is a member of the cytidine deaminase gene family. It is one of seven related genes or pseudogenes found in a cluster, thought to result from gene duplication, on chromosome 22. Members of the cluster encode proteins that are structurally and functionally related to the C to U RNA-editing cytidine deaminase APOBEC1. It is thought that the proteins may be RNA editing enzymes and have roles in growth or cell cycle control. A hybrid gene results from the deletion of approximately 29.5 kb of sequence between this gene, APOBEC3B, and the adjacent gene APOBEC3A. The breakpoints of the deletion are within the two genes, so the deletion allele is predicted to have the promoter and coding region of APOBEC3A, but the 3' UTR of APOBEC3B. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, July 2012]. Transcript Variant: This variant (1) represents the longer transcript and encodes the longer isoform (a). Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: AY743217.1 [ECO: 0000332] RNAseq introns :: mixed/partial sample support ERS025081, ERS025082 [ECO: 0000350] ##Evidence-Data-END## APOLD1 is an endothelial cell early response protein that may play a role in regulation of endothelial cell signaling and vascular function (Regard et al. 2004 [PubMed 15102925]). [supplied by OMIM, December 2008]. Transcript Variant: This variant (1) represents the longer transcript and encodes the longer isoform (1). Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The extent of this transcript is supported by transcript alignments. ##Evidence-Data-START## Transcript exon combination :: BC042478.1, DR000985.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025086 [ECO: 0000348] ##Evidence-Data-END## Rho GTPases play a fundamental role in numerous cellular processes triggered by extracellular stimuli that work through G protein coupled receptors. The encoded protein belongs to a family of cytoplasmic proteins that activate the Ras-like family of Rho proteins by exchanging bound GDP for GTP. It forms a complex with the small GTP binding protein Rac1 and recruits Rac1 to membrane ruffles and to focal adhesions. This protein can induce membrane ruffling. Multiple alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, July 2008].
APOLD1	exonic	81575	apolipoprotein L domain-containing protein 1 isoform 1	
ARHGEF7	exonic	8874	rho guanine nucleotide exchange factor 7 isoform a	

TABLE 3-continued

A non-redundant list of genes listed in Table 2				
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary
ASTN2	intronic	23245	astrotactin-2 isoform a precursor	Transcript Variant: This variant (1) differs in the 5' UTR, 3' UTR, coding region, and uses a downstream start codon, compared to variant 3. Both variants 1 and 5 encode isoform a, which has a shorter N-terminus and a longer and distinct C-terminus, compared to isoform c. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: D63476.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025081, ERS025082 [ECO: 0000348] ##Evidence-Data-END##
AUH	exonic	549	methylglutaconyl-CoA hydratase, mitochondrial precursor	This gene encodes a protein that is expressed in the brain and may function in neuronal migration, based on functional studies of the related astrotactin 1 gene in human and mouse. A deletion at this locus has been associated with schizophrenia. Multiple transcript variants encoding different proteins have been found for this locus. [provided by RefSeq, May 2010]. Transcript Variant: This variant (1) represents the longest transcript and encodes the longest isoform (a). ##Evidence-Data-START## Transcript exon combination :: BC146756.1, AB014534.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025082 [ECO: 0000348] ##Evidence-Data-END## The methylglutaconyl-CoA hydratase, mitochondrial protein binds to the AU-rich element (ARE), a common element found in the 3' UTR of rapidly decaying mRNA such as c-fos, c-myc, and granulocyte/macrophage colony stimulating factor. ARE elements are involved in directing RNA to rapid degradation and deadenylation. AUH is also homologous to enol-CoA hydratase, an enzyme involved in fatty acid degradation, and has been shown to have intrinsic hydratase enzymatic activity. AUH is thus a bifunctional chimera between RNA binding and metabolic enzyme activity. A possible subcellular localization in the mitochondria has been demonstrated for the mouse homolog of this protein which shares 92% identity with the human protein. It has been suggested that AUH may have a novel role as a mitochondrial located AU-binding protein. Human AUH is expressed as a single mRNA species of 1.8 kb, and translated as a 40-kDa precursor protein which is subsequently processed to a 32-kDa mature form. [provided by RefSeq, May 2010]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##RefSeq-Attributes-START## gene product(s) localized to mito. :: reported by MitoCarta ##RefSeq-Attributes-END## ##Evidence-Data-START## Transcript exon combination :: X79888.1, AL533438.3 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025084, ERS025088 [ECO: 0000348] ##Evidence-Data-END##
BACH1	exonic	571	BTB Domain And CNC Homolog 1	This gene encodes a transcription factor that belongs to the cap'n collar type of basic region leucine zipper factor family (CNC-bZip). The encoded protein contains broad complex, tramtrack, birc-a-brac/poxvirus and zinc finger (BTB/POZ) domains, which is atypical of CNC-bZip family members. These BTB/POZ domains facilitate protein-protein interactions and formation of homo- and/or hetero-oligomers. When this encoded protein forms a heterodimer with MafK, it functions as a repressor of Maf recognition element (MARE) and transcription is repressed. Multiple alternatively spliced transcript variants have been identified for this gene. [provided by RefSeq, May 2009]. Transcript Variant: This variant (3), also named BACHlt, differs in the 5'

TABLE 3-continued

A non-redundant list of genes listed in Table 2					
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary	Gene # (GN)
BDKRB2	intronic	624	B2 bradykinin receptor	UTR, 3' coding region and 3' UTR (compared to variant 1). This variant is represented as non-coding because the use of the 5'-most supported translational start codon, as used in variant 1, renders the transcript a candidate for nonsense-mediated mRNA decay (NMD). This transcript represents the splice variant reported by Kanezaki et al. (PMID: 11069897). Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## RNAseq introns :: mixed/partial sample support ERS025084, ERS025088 [ECO: 0000350] ##Evidence-Data-END## This gene encodes a receptor for bradykinin. The 9 aa bradykinin peptide elicits many responses including vasodilation, edema, smooth muscle spasm and pain fiber stimulation. This receptor associates with G proteins that stimulate a phosphatidylinositol-calcium second messenger system. Alternate start codons result in two isoforms of the protein. [provided by RefSeq, July 2008]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: DC369062.1, DC417219.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025090 [ECO: 0000348] ##Evidence-Data-END##	12
BMPR2	intronic	659	bone morphogenetic protein receptor type-2 precursor	This gene encodes a member of the bone morphogenetic protein (BMP) receptor family of transmembrane serine/threonine kinases. The ligands of this receptor are BMPs, which are members of the TGF-beta superfamily. BMPs are involved in endochondral bone formation and embryogenesis. These proteins transduce their signals through the formation of heteromeric complexes of two different types of serine (threonine) kinase receptors: type I receptors of about 50-55 kD and type II receptors of about 70-80 kD. Type II receptors bind ligands in the absence of type I receptors, but they require their respective type I receptors for signaling, whereas type I receptors require their respective type II receptors for ligand binding. Mutations in this gene have been associated with primary pulmonary hypertension, both familial and fenfluramine-associated, and with pulmonary venoocclusive disease. [provided by RefSeq, July 2008]. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The extent of this transcript is supported by transcript alignments. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: BC052985.2, AK292430.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025081, ERS025082 [ECO: 0000348] ##Evidence-Data-END##	13
BTBD17	exonic	388419	BTB/POZ domain-containing protein 17 precursor	N/A	14
C17orf77	exonic	146723	protein C17orf77 precursor	N/A	15

TABLE 3-continued

A non-redundant list of genes listed in Table 2					
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary	Gene # (GN)
CAPZB	intronic	832	F-actin-capping protein subunit beta isoform 1	This gene encodes the beta subunit of the barbed-end actin binding protein, which belongs to the F-actin capping protein family. The capping protein is a heterodimeric actin capping protein that blocks actin filament assembly and disassembly at the fast growing (barbed) filament ends and functions in regulating actin filament dynamics as well as in stabilizing actin filament lengths in muscle and nonmuscle cells. A pseudogene of this gene is located on the long arm of chromosome 2. Multiple alternatively spliced transcript variants encoding different isoforms have been found. [provided by RefSeq, August 2013]. Transcript Variant: This variant (1) encodes isoform 1. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: BC107752.1, BM451686.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025088 [ECO: 0000348] ##Evidence-Data-END## N/A	16
CCDC41	exonic	51134	centrosomal protein of 83 kDa	This gene encodes a member of the CD300 glycoprotein family of cell surface proteins found on leukocytes involved in immune response signaling pathways. This gene is located on chromosome 17 in a cluster with all but one of the other family members. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, February 2012]. Transcript Variant: This variant (1) represents the longer transcript and encodes the longer protein (isoform 1), also referred to as IRC1a. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: BC02352.1, AL531420.3 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025081, ERS025083 [ECO: 0000348] ##Evidence-Data-END##	17
CD300A	exonic	11314	CMRF35-like molecule 8 isoform 1 precursor		18
CD300C	exonic	10871	CMRF35-like molecule 6 precursor	The CMRF35 antigen, which was identified by reactivity with a monoclonal antibody, is present on monocytes, neutrophils, and some T and B lymphocytes (Jackson et al., 1992 [PubMed 1349552]). [supplied by OMIM, March 2008]. ##Evidence-Data-START## Transcript exon combination :: BC022279.1, BM922826.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025084, ERS025087 [ECO: 0000348] ##Evidence-Data-END## This gene encodes a member of the CD300 glycoprotein family of cell surface proteins expressed on myeloid cells. The protein interacts with the TYRO protein tyrosine kinase-binding protein and is thought to act as an activating receptor. [provided by RefSeq, November 2012]. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments. An in-frame AUG is located 41 codons upstream of the annotated translation start site but is not being annotated as a start site since it is not conserved and is in a weak Kozak sequence context. ##RefSeq-Attributes-START## CDS uses downstream in-frame AUG :: downstream AUG is associated with N-terminal localization signal ##RefSeq-Attributes-END## ##Evidence-Data-START## Transcript exon combination :: AK303545.1, BX648376.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025084, ERS025088 [ECO: 0000348] ##Evidence-Data-END##	19
CD300E	exonic	342510	CMRF35-like molecule 2 precursor		20

TABLE 3-continued

A non-redundant list of genes listed in Table 2					
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary	Gene # (GN)
CD300LB	exonic	124599	CMRF35-like molecule 7 precursor	CD300LB is a nonclassical activating receptor of the immunoglobulin (Ig) superfamily expressed on myeloid cells (Martinez-Barrioacanal and Sayos, 2006 [PubMed 16920917]). [supplied by OMIM, March 2008]. CCDS Note: The coding region has been updated to shorten the N-terminus to one that is more supported by available conservation data and paralogous family members. The update has a predicted N-terminal signal peptide, which is consistent with functional support for the protein (e.g., PMIDs 16920917, 19359216). ##Evidence-Data-START## Transcript exon combination :: BC028091.1, AY359025.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025084, ERS025088 [ECO: 0000348] ##Evidence-Data-END## ##RefSeq-Attributes-START## CDS uses downstream in-frame AUG :: downstream AUG is associated with N-terminal localization signal ##RefSeq-Attributes-END## N/A	21
CD300LD	exonic	100131439	CMRF35-like molecule 4 precursor	CD300LF is an inhibitory receptor of the Ig superfamily expressed on myeloid cells. It mediates negative regulatory signals by recruiting SHP1 (PTPN6; MIM 176883) or SHP (INPP5D; MIM 601582) (Sui et al., 2004 [PubMed 15184070]; Alvarez-Errico et al., 2004 [PubMed 15549731]). [supplied by OMIM, March 2008]. Sequence Note: The RefSeq transcript and protein were derived from genomic sequence to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on alignments. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: AF251706.1, AY358545.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025084 [ECO: 0000348] ##Evidence-Data-END##	22
CD300LF	exonic	146722	CMRF35-like molecule 1 precursor	This gene encodes a cyclin-dependent kinase inhibitor, which shares a limited similarity with CDK inhibitor CDKN1A/p21. The encoded protein binds to and prevents the activation of cyclin E-CDK2 or cyclin D-CDK4 complexes, and thus controls the cell cycle progression at G1. The degradation of this protein, which is triggered by its CDK dependent phosphorylation and subsequent ubiquitination by SCF complexes, is required for the cellular transition from quiescence to the proliferative state. [provided by RefSeq, July 2008]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: BC001971.1, AY004255.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025081, ERS025082 [ECO: 0000348] ##Evidence-Data-END##	23
CDKN1B	exonic	1027	cyclin-dependent kinase inhibitor 1B	The centromere is a specialized chromatin domain, present throughout the cell cycle, that acts as a platform on which the transient assembly of the kinetochore occurs during mitosis. All active centromeres are characterized by the presence of long arrays of nucleosomes in which CENPA (MIM 117139) replaces histone H3 (see MIM 601128). CENPM is an additional factor required for centromere assembly (Foltz et al., 2006 [PubMed 16622419]). [supplied by OMIM, March 2008]. Transcript Variant: This variant	24
CENPM	exonic	79019	centromere protein M isoform a		25

TABLE 3-continued

A non-redundant list of genes listed in Table 2				
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary
COG4	exonic	25839	conserved oligomeric Golgi complex subunit 4 isoform 1	(1) represents the longer transcript and encodes the longer isoform (a). Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: BC000705.2, BC007495.2 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025085, ERS025088 [ECO: 0000348] ##Evidence-Data-END## The protein encoded by this gene is a component of an oligomeric protein complex involved in the structure and function of the Golgi apparatus. Defects in this gene may be a cause of congenital disorder of glycosylation type IIj. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, August 2010]. Transcript Variant: This variant (1) represents the longer transcript and encodes the longer isoform (1). ##Evidence-Data-START## Transcript exon combination :: BC072438.1, AK022874.1 [ECO: 0000332] RNAseq introns :: mixed/partial sample support ERS025081, ERS025082 [ECO: 0000350] ##Evidence-Data-END## COMMD6 belongs to a family of NF-kappa-B (see RELA; MIM 164014)-inhibiting proteins characterized by the presence of a COMM domain (see COMMD1; MIM 607238) (de Bie et al., 2006 [PubMed 16573520]). [supplied by OMIM, March 2009]. ##Evidence-Data-START## Transcript exon combination :: HY028175.1, DW440523.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025088 [ECO: 0000348] ##Evidence-Data-END##
COMMD6	exonic	170622	COMM domain-containing protein 6 isoform a	The protein encoded by this gene is a death domain (CARD/DD)-containing protein and has been shown to induce cell apoptosis. Through its CARD domain, this protein interacts with, and thus recruits, caspase 2/ICH1 to the cell death signal transduction complex that includes tumor necrosis factor receptor 1 (TNFR1A), RIPK1/RIP kinase, and numbers of other CARD domain-containing proteins. [provided by RefSeq, July 2008]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: BX480215.1, BC017042.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025081, ERS025083 [ECO: 0000348] ##Evidence-Data-END## cAMP response element (CRE)-binding protein-like-2 (CREBL2) was identified in a search to find genes in a commonly deleted region on chromosome 12p13 flanked by ETV6 and CDKN1B genes, frequently associated with hematopoietic malignancies, as well as breast, non-small-cell lung and ovarian cancers. CREBL2 shares a 41% identity with CRE-binding protein (CREB) over a 48-base long region which encodes the bZip domain of CREB. The bZip domain consists of about 30 amino acids rich in basic residues involved in DNA binding, followed by a leucine zipper motif involved in protein dimerization. This suggests that CREBL2 encodes a protein with DNA binding capabilities. The occurrence of CREBL2 deletion in malignancy suggests that CREBL2 may act as a tumor suppressor gene. [provided by RefSeq, July 2008]. ##Evidence-Data-START## Transcript exon combination :: BC106021.1, AF039081.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025081, ERS025082 [ECO: 0000348] ##Evidence-Data-END##
CRADD	exonic	8738	death domain-containing protein CRADD	
CREBL2	exonic	1389	cAMP-responsive element-binding protein-like 2	

Gene # (GN)

TABLE 3-continued

A non-redundant list of genes listed in Table 2					
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary	Gene # (GN)
DNAI2	exonic	64446	dynein intermediate chain 2, axonemal isoform 1	The protein encoded by this gene belongs to the dynein intermediate chain family, and is part of the dynein complex of respiratory cilia and sperm flagella. Mutations in this gene are associated with primary ciliary dyskinesia type 9. Alternatively spliced transcript variants encoding different isoforms have been noted for this gene. [provided by RefSeq, March 2010]. Transcript Variant: This variant (1) encodes the longer isoform (1). ##Evidence-Data-START## Transcript exon combination :: AF250288.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025085 [ECO: 0000348] ##Evidence-Data-END## ##RefSeq-Attributes-START## NMD candidate :: translation inferred from conservation ##RefSeq-Attributes-END## N/A	30
DNER	intronic	92737	delta and Notch-like epidermal growth factor-related receptor precursor		31
DUSP16	exonic	80824	dual specificity protein phosphatase 16	This gene encodes a mitogen-activated protein kinase phosphatase that is a member of the dual specificity protein phosphatase subfamily. These phosphatases inactivate their target kinases by dephosphorylating both the phosphoserine/threonine and phosphotyrosine residues. The encoded protein specifically regulates the c-Jun amino-terminal kinase (JNK) and extracellular signal-regulated kinase (ERK) pathways.[provided by RefSeq, May 2010]. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: AF506796.1, AB052156.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025084, ERS025088 [ECO: 0000348] ##Evidence-Data-END## N/A	32
ECRP	exonic	643332	N/A		33
EDIL3	intronic	10085	EGF-like repeat and discoidin I-like domain-containing protein 3 isoform 1 precursor	The protein encoded by this gene is an integrin ligand. It plays an important role in mediating angiogenesis and may be important in vessel wall remodeling and development. It also influences endothelial cell behavior. [provided by RefSeq, July 2008]. Transcript Variant: This variant (1) encodes the longer isoform (1). Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: BC030828.1, U70312.1 [ECO: 0000332] RNAseq introns :: mixed/partial sample support ERS025081, ERS025082 [ECO: 0000350] ##Evidence-Data-END## N/A	34
EEA1	exonic	8411	early endosome antigen 1		35

TABLE 3-continued

A non-redundant list of genes listed in Table 2					
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary	Gene # (GN)
EHF	both	26298	ETS homologous factor isoform 1 precursor	This gene encodes a protein that belongs to an ETS transcription factor subfamily characterized by epithelial-specific expression (ESEs). The encoded protein acts as a transcriptional repressor and may be involved in epithelial differentiation and carcinogenesis. Three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, June 2011]. Transcript Variant: This variant (1) encodes the longest isoform (1). Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: AK310867.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025084, ERS025086 [ECO: 0000348] ##Evidence-Data-END##	36
EMB	exonic	133418	embigin precursor	This gene encodes a transmembrane glycoprotein that is a member of the immunoglobulin superfamily. The encoded protein may be involved in cell growth and development by mediating interactions between the cell and extracellular matrix. A pseudogene of this gene is found on chromosome 1. [provided by RefSeq, January 2009]. ##Evidence-Data-START## Transcript exon combination :: BC059398.1, AK300860.1 [ECO: 0000332] RNAseq introns :: mixed/partial sample support ERS025081, ERS025082 [ECO: 0000350] ##Evidence-Data-END##	37
ETV6	exonic	2120	transcription factor ETV6	This gene encodes an ETS family transcription factor. The product of this gene contains two functional domains: a N-terminal pointed (PNT) domain that is involved in protein-protein interactions with itself and other proteins, and a C-terminal DNA-binding domain. Gene knockout studies in mice suggest that it is required for hematopoiesis and maintenance of the developing vascular network. This gene is known to be involved in a large number of chromosomal rearrangements associated with leukemia and congenital fibrosarcoma. [provided by RefSeq, September 2008]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: BC043399.1, U11732.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025081, ERS025082 [ECO: 0000348] ##Evidence-Data-END##	38
FHL2	exonic	2274	four and a half LIM domains protein 2	This gene encodes a member of the four-and-a-half-LIM-only protein family. Family members contain two highly conserved, tandemly arranged, zinc finger domains with four highly conserved cysteines binding a zinc atom in each zinc finger. This protein is thought to have a role in the assembly of extracellular membranes. Also, this gene is down-regulated during transformation of normal myoblasts to rhabdomyosarcoma cells and the encoded protein may function as a link between presenilin-2 and an intracellular signaling pathway. Multiple alternatively spliced variants, encoding the same protein, have been identified. [provided by RefSeq, August 2011]. Transcript Variant: This variant (1) differs in the 5' UTR compared to variant 2. Variants 1, 2, 4 and 5 encode the same isoform. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##RefSeq-Attributes-START## CDS uses	39

TABLE 3-continued

A non-redundant list of genes listed in Table 2					
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary	Gene # (GN)
FLJ26850 FPR2 FPR3 FUK	intronic exonic	400710 2358	N/A N-formyl peptide receptor 2	downstream in-frame AUG :: lack of evidence for use of upstream AUG ##RefSeq-Attributes-END## ##Evidence-Data-START## Transcript exon combination :: BC093049.1, AL523628.3 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025084, ERS025088 [ECO: 0000348] ##Evidence-Data-END##	40 41
	exonic	2359	N-formyl peptide receptor 3	N/A	42
	both	197258	L-fucose kinase	The protein encoded by this gene belongs to the GHMP (galacto-, homoserine, mevalonate and phosphomevalonate) kinase family and catalyzes the phosphorylation of L-fucose to form beta-L-fucose 1-phosphate. This enzyme catalyzes the first step in the utilization of free L-fucose in glycoprotein and glycolipid synthesis. L-fucose may be important in mediating a number of cell- cell interactions such as blood group antigen recognition, inflammation, and metastasis. While several transcript variants may exist for this gene, the full- length nature of only one has been described to date. [provided by RefSeq, July 2008]. ##Evidence-Data-START## Transcript exon combination :: AJ441184.1, BC032542.1 [ECO: 0000332] RNAseq introns :: mixed/partial sample support ERS025081, ERS025082 [ECO: 0000350] ##Evidence-Data- END##	43
	GDA	exonic	9615	guanine deaminase isoform a	This gene encodes an enzyme responsible for the hydrolytic deamination of guanine. Studies in rat ortholog suggest this gene plays a role in microtubule assembly. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, November 2011]. Transcript Variant: This variant (1) encodes the longest isoform (a). Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments. ##Evidence-Data- START## RNAseq introns :: mixed/partial sample support ERS025082, ERS025083 [ECO: 0000350] ##Evidence-Data-END##
GDPD4	exonic	220032	glycerophosphodiester phosphodiesterase domain-containing protein 4	N/A	45
GPATCH2	intronic	55105	G patch domain- containing protein 2	N/A	46
GPC5	intronic	2262	glypican-5 precursor	Cell surface heparan sulfate proteoglycans are composed of a membrane- associated protein core substituted with a variable number of heparan sulfate chains. Members of the glypican-related integral membrane proteoglycan family (GRIPs) contain a core protein anchored to the cytoplasmic membrane via a glycosyl phosphatidylinositol linkage. These proteins may play a role in the control of cell division and growth regulation. [provided by RefSeq, July 2008]. Publication Note: This RefSeq record includes a subset of the	47

TABLE 3-continued

A non-redundant list of genes listed in Table 2				
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary
				publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: BC030584.1, BC039730.1 [ECO: 0000332] RNAseq introns :: mixed/partial sample support ERS025082, ERS025083 [ECO: 0000350] ##Evidence-Data-END##
GPR19	exonic	2842	probable G-protein coupled receptor 19	N/A
GPR142	exonic	350383	probable G-protein coupled receptor 142	48
GPRC5C	exonic	55890	G-protein coupled receptor family C group 5 member C isoform a	49
				50
GRIA3	intronic	2892	glutamate receptor 3 isoform 1 precursor	51
				52
GTPBP4	exonic	23560	nucleolar GTP-binding protein 1	

TABLE 3-continued

A non-redundant list of genes listed in Table 2				
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary
HCN1	exonic	348980	potassium/sodium hyperpolarization-activated cyclic nucleotide-gated channel 1	occurs; the active protein has an opportunity to diffuse away from the receptor and deliver its message for a prolonged period to its downstream target. [provided by RefSeq, July 2008]. ##Evidence-Data-START## Transcript exon combination :: AK001552.1, AK222861.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025081, ERS025082 [ECO: 0000348] ##Evidence-Data-END## The membrane protein encoded by this gene is a hyperpolarization-activated cation channel that contributes to the native pacemaker currents in heart and neurons. The encoded protein can homodimerize or heterodimerize with other pore-forming subunits to form a potassium channel. This channel may act as a receptor for sour tastes. [provided by RefSeq, October 2011]. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: AF488549.1, AF064876.1 [ECO: 0000332] RNAseq introns :: mixed/partial sample support ERS025081, ERS025082 [ECO: 0000350] ##Evidence-Data-END##
HEXA	exonic	3073	beta-hexosaminidase subunit alpha preproprotein	This gene encodes the alpha subunit of the lysosomal enzyme beta-hexosaminidase that, together with the cofactor GM2 activator protein, catalyzes the degradation of the ganglioside GM2, and other molecules containing terminal N-acetyl hexosamines. Beta-hexosaminidase is composed of two subunits, alpha and beta, which are encoded by separate genes. Both beta-hexosaminidase alpha and beta subunits are members of family 20 of glycosyl hydrolases. Mutations in the alpha or beta subunit genes lead to an accumulation of GM2 ganglioside in neurons and neurodegenerative disorders termed the GM2 gangliosidosis. Alpha subunit gene mutations lead to Tay-Sachs disease (GM2-gangliosidosis type I). [provided by RefSeq, July 2009]. Sequence Note: This RefSeq record was created from transcript and genomic sequence data because no single transcript was available for the full length of the gene. The extent of this transcript is supported by transcript alignments. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: M13520.1, CR627386.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025084, ERS025088 [ECO: 0000348] ##Evidence-Data-END##
HK2	exonic	3099	hexokinase-2	Hexokinases phosphorylate glucose to produce glucose-6-phosphate, the first step in most glucose metabolism pathways. This gene encodes hexokinase 2, the predominant form found in skeletal muscle. It localizes to the outer membrane of mitochondria. Expression of this gene is insulin-responsive, and studies in rat suggest that it is involved in the increased rate of glycolysis seen in rapidly growing cancer cells. [provided by RefSeq, April 2009]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: BC064369.1, AF148513.1 [ECO: 0000332] RNAseq introns :: single sample

TABLE 3-continued

A non-redundant list of genes listed in Table 2				
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary
HMGB3	exonic	3149	high mobility group protein B3	supports all introns ERS025083, ERS025084 [ECO: 0000348] ##Evidence-Data-END## HMGB3 belongs to the high mobility group (HMG) protein superfamily. Like HMGI (MIM 163905) and HMGI2 (MIM 163906), HMGB3 contains DNA-binding HMG box domains and is classified into the HMG box subfamily. Members of the HMG box subfamily are thought to play a fundamental role in DNA replication, nucleosome assembly and transcription (Wilke et al., 1997 [PubMed 9370291]; Nemeth et al., 2006 [PubMed 16945912]). [supplied by OMIM, March 2008]. ##Evidence-Data-START## Transcript exon combination :: Y10043.1, BG176733.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025081, ERS025082 [ECO: 0000348] ##Evidence-Data-END##
56				
HPR	exonic	3250	haptoglobin-related protein precursor	This gene encodes a haptoglobin-related protein that binds hemoglobin as efficiently as haptoglobin. Unlike haptoglobin, plasma concentration of this protein is unaffected in patients with sickle cell anemia and extensive intravascular hemolysis, suggesting a difference in binding between haptoglobin-hemoglobin and haptoglobin-related protein-hemoglobin complexes to CD163, the hemoglobin scavenger receptor. This protein may also be a clinically important predictor of recurrence of breast cancer. [provided by RefSeq, October 2011]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: CB147217.1, CB122261.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025084, ERS025088 [ECO: 0000348] ##Evidence-Data-END##
57				
HTATSF1P2 IDI2	exonic exonic	401233 91734	N/A isopentenyl-diphosphate Delta-isomerase 2	N/A N/A
58 59				
IDI2-AS1 IDO2	exonic intronic	55853 169355	N/A indoleamine 2,3-dioxygenase 2	N/A Along with the enzymes encoded by the INDO (MIM 147435) and TDO2 (MIM 191070) genes, the enzyme encoded by the INDOL1 gene metabolizes tryptophan in the kynurenine pathway (Ball et al., 2007 [PubMed 17499941]). [supplied by OMIM, February 2011]. Sequence Note: The RefSeq transcript 3' UTR was derived from genomic sequence to make the sequence consistent with the reference genome assembly. The genomic coordinates used were based on transcript alignments.
60 61				
IFNLR1	exonic	163702	interferon lambda receptor 1 isoform 1 precursor	The protein complex has been shown to interact with three beta (IL10RB). The receptor complex belongs to the class II cytokine receptor family. This protein forms a receptor complex with interleukin 10 receptor, closely related cytokines, including interleukin 28A (IL28A), interleukin 28B (IL28B), and interleukin 29 (IL29). The expression of all three cytokines can be induced by viral infection. The cells overexpressing this protein have been found to have enhanced responses to IL28A and IL29, but decreased response to IL28B. Three alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, July 2008]. Transcript Variant: This variant (1) represents the longest transcript and it encodes the longest protein (isoform 1). Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene
62				

TABLE 3-continued

A non-redundant list of genes listed in Table 2				
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary
IQCB1	exonic	9657	IQ calmodulin-binding motif-containing protein 1 isoform a	record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: AF439325.1, AK160364.1 [ECO: 0000332] RNASeq introns :: single sample supports all introns ERS025084 [ECO: 0000348] ##Evidence-Data-END## This gene encodes a nephrocystin protein that interacts with calmodulin and the retinitis pigmentosa GTPase regulator protein. The encoded protein has a central coiled-coil region and two calmodulin-binding IQ domains. It is localized to the primary cilia of renal epithelial cells and connecting cilia of photoreceptor cells. The protein is thought to play a role in ciliary function. Defects in this gene result in Senior-Løken syndrome type 5. Alternative splicing results in multiple transcript variants. [provided by RefSeq, November 2009]. Transcript Variant: This variant (1) encodes the longer isoform (a). Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: D25278.1, AY714228.1 [ECO: 0000332] RNASeq introns :: mixed/partial sample support ERS025081, ERS025082 [ECO: 0000350] ##Evidence-Data-END##
JPX	intronic	554203		JPX is a nonprotein-coding RNA transcribed from a gene within the X-inactivation center (XIC; MIM 314670) that appears to participate in X-chromosome inactivation (Tian et al., 2010 [PubMed 21029862]). [supplied by OMIM, February 2011]. ##Evidence-Data-START## Transcript exon combination :: BC017761.1 [ECO: 0000332] RNASeq introns :: mixed/partial sample support ERS025081, ERS025082 [ECO: 0000350] ##Evidence-Data-END##
KANK1	intronic	23189	KN motif and ankyrin repeat domain-containing protein 1 isoform a	The protein encoded by this gene belongs to the Kank family of proteins, which contain multiple ankyrin repeat domains. This family member functions in cytoskeleton formation by regulating actin polymerization. This gene is a candidate tumor suppressor for renal cell carcinoma. Mutations in this gene cause cerebral palsy spastic quadriplegic type 2, a central nervous system development disorder. A t(5; 9) translocation results in fusion of the platelet-derived growth factor receptor beta gene (PDGFRB) on chromosome 5 with this gene in a myeloproliferative neoplasm featuring severe thrombocytopenia. Alternative splicing of this gene results in multiple transcript variants. A related pseudogene has been identified on chromosome 20. [provided by RefSeq, March 2012]. Transcript Variant: This variant (1) represents the shortest transcript but encodes the longer isoform (a, also known as Kank-L). Variants 1, 3 and 4 all encode isoform a. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: AL833161.1, AK292989.1 [ECO: 0000332] RNASeq introns :: single sample supports all introns ERS025084, ERS025085 [ECO: 0000348] ##Evidence-Data-END##
KAT6B	exonic	23522	histone acetyltransferase KAT6B isoform 1	The protein encoded by this gene is a histone acetyltransferase and component of the MOZ/MORF protein complex. In addition to its acetyltransferase activity, the encoded protein has transcriptional activation activity in its N-terminal end and transcriptional repression activity in its C-terminal end. This

TABLE 3-continued

A non-redundant list of genes listed in Table 2				
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary
KCTD7	exonic	154881	BTB/POZ domain-containing protein KCTD7 isoform 1	protein is necessary for RUNX2-dependent transcriptional activation and could be involved in brain development. Mutations have been found in patients with genitopatellar syndrome. A translocation of this gene and the CREBBP gene results in acute myeloid leukemias. Three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, March 2012]. Transcript Variant: This variant (1) represents the longest transcript and encodes the longest isoform (1). Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: AF217500.1, BC150618.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025083, ERS025084 [TECO: 00003481] ##Evidence-Data-END##
				This gene encodes a member of the potassium channel tetramerization domain-containing protein family. Family members are identified on a structural basis and contain an amino-terminal domain similar to the T1 domain present in the voltage-gated potassium channel. Mutations in this gene have been associated with progressive myoclonic epilepsy-3. Alternative splicing results in multiple transcript variants. [provided by RefSeq, January 2011]. Transcript Variant: This variant (1) represents the longer transcript and encodes the longer isoform (1). Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments. ##Evidence-Data-START## Transcript exon combination :: AK056631.1, BU902852.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025081, ERS025083 [ECO: 0000348] ##Evidence-Data-END##
KIF19	exonic	124602	kinesin-like protein KIF19	
LARP4B	exonic	23185	la-related protein 4B	
				This gene encodes a member of an evolutionarily conserved protein family implicated in RNA metabolism and translation. Members of this family are characterized by the presence of an La motif, which is often located adjacent to one or more RNA recognition motifs (RRM). Together, the two motifs constitute the functional region of the protein and enable its interaction with the RNA substrate. This protein family is divided into five sub-families: the genuine La proteins and four La-related protein (LARP) sub-families. The protein encoded by this gene belongs to LARP sub-family 4. It is a cytoplasmic protein that may play a stimulatory role in translation. [provided by RefSeq, October 2012]. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments. ##Evidence-Data-START## CDS exon combination :: BC152443.1, D86971.2 [ECO: 0000331] RNAseq introns :: mixed/partial sample support ERS025088 [ECO: 0000350] ##Evidence-Data-END##
LOC643339	exonic	643339	N/A	
LOH12CR1	exonic	118426	loss of heterozygosity 12 chromosomal region 1 protein	

67

68

69

70

71

TABLE 3-continued

A non-redundant list of genes listed in Table 2				
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary
MALL	exonic	7851	MAL-like protein	This gene encodes an element of the machinery for raft-mediated trafficking in endothelial cells. The encoded protein, a member of the MAL protolipid family, predominantly localizes in glycolipid- and cholesterol-enriched membrane (GEM) rafts. It interacts with caveolin-1. [provided by RefSeq, July 2008]. ##Evidence-Data-START## Transcript exon combination :: AK125647.1, AK056616.1 [ECO: 0000332] RNaseq introns :: single sample supports all introns ERS025084, ERS025088 [ECO: 0000348] ##Evidence-Data-END##
MAPK9	exonic	5601	mitogen-activated protein kinase 9 isoform alpha1	The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase targets specific transcription factors, and thus mediates immediate-early gene expression in response to various cell stimuli. It is most closely related to MAPK8, both of which are involved in UV radiation induced apoptosis, thought to be related to the cytochrome c-mediated cell death pathway. This gene and MAPK8 are also known as c-Jun N-terminal kinases. This kinase blocks the ubiquitination of tumor suppressor p53, and thus it increases the stability of p53 in nonstressed cells. Studies of this gene's mouse counterpart suggest a key role in T-cell differentiation. Several alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, September 2008]. Transcript Variant: This variant (JNK2-a1) uses a different acceptor splice site in the last coding exon compared to transcript variant JNK2-a2, resulting in a frameshift and a shorter isoform (JNK2 alpha1) with a different C-terminus, compared to isoform JNK2 alpha2. The JNK2-a1 variant differs from the JNK2-b1 variant in the use of an alternate internal coding exon of the same length. Thus, JNK2 alpha1 isoform is the same length as JNK2 beta1 isoform, with a few aa differences in an internal protein segment. Sequence Note: This RefSeq record was created from transcript and genomic sequence data because no single transcript was available for the full length of the gene. The extent of this transcript is supported by transcript alignments. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## CDS exon combination :: U34821.1 [ECO: 0000331] RNaseq introns :: mixed/partial sample support ERS025081, ERS025082 [ECO: 0000350] ##Evidence-Data-END##
MCEE	both	84693	methylmalonyl-CoA epimerase, mitochondrial precursor	The product of this gene catalyzes the interconversion of D- and L-methylmalonyl-CoA during the degradation of branched chain amino acids, odd chain-length fatty acids, and other metabolites. Mutations in this gene result in methylmalonyl-CoA epimerase deficiency, which is presented as mild to moderate methylmalonic aciduria. [provided by RefSeq, July 2008]. ##Evidence-Data-START## Transcript exon combination :: BC020825.1, BG567074.1 [ECO: 0000332] RNaseq introns :: single sample supports all introns ERS025081, ERS025082 [ECO: 0000348] ##Evidence-Data-END##
MGAT5	intronic	4249	alpha-1,6-mannosylglycoprotein 6-beta-N-acetylglucosaminyltransferase A	The protein encoded by this gene belongs to the glycosyltransferase family. It catalyzes the addition of beta-1,6-N-acetylglucosamine to the alpha-linked mannose of biantennary N-linked oligosaccharides present on the newly synthesized glycoproteins. It is one of the most important enzymes involved in the regulation of the biosynthesis of glycoprotein oligosaccharides. Alterations
				72
				73
				74
				75

TABLE 3-continued

A non-redundant list of genes listed in Table 2				
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary
MGC16275 MGME1	exonic exonic	85001 92667	N/A mitochondrial genome maintenance exonuclease 1	of the oligosaccharides on cell surface glycoproteins cause significant changes in the adhesive or migratory behavior of a cell. Increase in the activity of this enzyme has been correlated with the progression of invasive malignancies. [provided by RefSeq, October 2011]. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: D17116.1, AF113921.1 [ECO: 0000332] RNaseq introns :: single sample supports all introns ERS025081, ERS025082 [ECO: 0000348] ##Evidence-Data-END## N/A N/A
MIR200A	exonic	406983		microRNAs (miRNAs) are short (20-24 nt) non-coding RNAs that are involved in post-transcriptional regulation of gene expression in multicellular organisms by affecting both the stability and translation of mRNAs. miRNAs are transcribed by RNA polymerase II as part of capped and polyadenylated primary transcripts (pri-miRNAs) that can be either protein-coding or non-coding. The primary transcript is cleaved by the Drosha ribonuclease III enzyme to produce an approximately 70-nt stem-loop precursor miRNA (pre-miRNA), which is further cleaved by the cytoplasmic Dicer ribonuclease to generate the mature miRNA and antisense miRNA star (miRNA*) products. The mature miRNA is incorporated into a RNA-induced silencing complex (RISC), which recognizes target mRNAs through imperfect base pairing with the miRNA and most commonly results in translational inhibition or destabilization of the target mRNA. The RefSeq represents the predicted microRNA stem-loop. [provided by RefSeq, September 2009]. Sequence Note: This record represents a predicted microRNA stem-loop as defined by miRBase. Some sequence at the 5' and 3' ends may not be included in the intermediate precursor miRNA produced by Drosha cleavage. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications.
MIR200B	exonic	406984		microRNAs (miRNAs) are short (20-24 nt) non-coding RNAs that are involved in post-transcriptional regulation of gene expression in multicellular organisms by affecting both the stability and translation of mRNAs. miRNAs are transcribed by RNA polymerase II as part of capped and polyadenylated primary transcripts (pri-miRNAs) that can be either protein-coding or non-coding. The primary transcript is cleaved by the Drosha ribonuclease III enzyme to produce an approximately 70-nt stem-loop precursor miRNA (pre-miRNA), which is further cleaved by the cytoplasmic Dicer ribonuclease to generate the mature miRNA and antisense miRNA star (miRNA*) products. The mature miRNA is incorporated into a RNA-induced silencing complex (RISC), which recognizes target mRNAs through imperfect base pairing with the miRNA and most commonly results in translational inhibition or destabilization of the target mRNA. The RefSeq represents the predicted

76
77

78

79

TABLE 3-continued

A non-redundant list of genes listed in Table 2				
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary
MIR429	exonic	554210		microRNA stem-loop. [provided by RefSeq, September 2009]. Sequence Note: This record represents a predicted microRNA stem-loop as defined by miRBase. Some sequence at the 5' and 3' ends may not be included in the intermediate precursor miRNA produced by Drosha cleavage. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. microRNAs (miRNAs) are short (20-24 nt) non-coding RNAs that are involved in post-transcriptional regulation of gene expression in multicellular organisms by affecting both the stability and translation of mRNAs. miRNAs are transcribed by RNA polymerase II as part of capped and polyadenylated primary transcripts (pri-miRNAs) that can be either protein-coding or non-coding. The primary transcript is cleaved by the Drosha ribonuclease III enzyme to produce an approximately 70-nt stem-loop precursor miRNA (pre-miRNA), which is further cleaved by the cytoplasmic Dicer ribonuclease to generate the mature miRNA and antisense miRNA star (miRNA*) products. The mature miRNA is incorporated into a RNA-induced silencing complex (RISC), which recognizes target mRNAs through imperfect base pairing with the miRNA and most commonly results in translational inhibition or destabilization of the target mRNA. The RefSeq represents the predicted microRNA stem-loop. [provided by RefSeq, September 2009]. Sequence Note: This record represents a predicted microRNA stem-loop as defined by miRBase. Some sequence at the 5' and 3' ends may not be included in the intermediate precursor miRNA produced by Drosha cleavage. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. microRNAs (miRNAs) are short (20-24 nt) non-coding RNAs that are involved in post-transcriptional regulation of gene expression in multicellular organisms by affecting both the stability and translation of mRNAs. miRNAs are transcribed by RNA polymerase II as part of capped and polyadenylated primary transcripts (pri-miRNAs) that can be either protein-coding or non-coding. The primary transcript is cleaved by the Drosha ribonuclease III enzyme to produce an approximately 70-nt stem-loop precursor miRNA (pre-miRNA), which is further cleaved by the cytoplasmic Dicer ribonuclease to generate the mature miRNA and antisense miRNA star (miRNA*) products. The mature miRNA is incorporated into a RNA-induced silencing complex (RISC), which recognizes target mRNAs through imperfect base pairing with the miRNA and most commonly results in translational inhibition or destabilization of the target mRNA. The RefSeq represents the predicted microRNA stem-loop. [provided by RefSeq, September 2009]. Sequence Note: This record represents a predicted microRNA stem-loop as defined by miRBase. Some sequence at the 5' and 3' ends may not be included in the intermediate precursor miRNA produced by Drosha cleavage.
MIR595	exonic	693180		microRNAs (miRNAs) are short (20-24 nt) non-coding RNAs that are involved in post-transcriptional regulation of gene expression in multicellular organisms by affecting both the stability and translation of mRNAs. miRNAs are transcribed by RNA polymerase II as part of capped and polyadenylated primary transcripts (pri-miRNAs) that can be either protein-coding or non-coding. The primary transcript is cleaved by the Drosha ribonuclease III enzyme to produce an approximately 70-nt stem-loop precursor miRNA (pre-miRNA), which is further cleaved by the cytoplasmic Dicer ribonuclease to generate the mature miRNA and antisense miRNA star (miRNA*) products. The mature miRNA is incorporated into a RNA-induced silencing complex (RISC), which recognizes target mRNAs through imperfect base pairing with the miRNA and most commonly results in translational inhibition or destabilization of the target mRNA. The RefSeq represents the predicted microRNA stem-loop. [provided by RefSeq, September 2009]. Sequence Note: This record represents a predicted microRNA stem-loop as defined by miRBase. Some sequence at the 5' and 3' ends may not be included in the intermediate precursor miRNA produced by Drosha cleavage.
MIR651	exonic	723779		microRNAs (miRNAs) are short (20-24 nt) non-coding RNAs that are involved in post-transcriptional regulation of gene expression in multicellular organisms by affecting both the stability and translation of mRNAs. miRNAs are transcribed by RNA polymerase II as part of capped and polyadenylated primary transcripts (pri-miRNAs) that can be either protein-coding or non-coding. The primary transcript is cleaved by the Drosha ribonuclease III enzyme to produce an approximately 70-nt stem-loop precursor miRNA (pre-

TABLE 3-continued

A non-redundant list of genes listed in Table 2				
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary
MIR3163	exonic	100423029		miRNA), which is further cleaved by the cytoplasmic Dicer ribonuclease to generate the mature miRNA and antisense miRNA star (miRNA*) products. The mature miRNA is incorporated into a RNA-induced silencing complex (RISC), which recognizes target mRNAs through imperfect base pairing with the miRNA and most commonly results in translational inhibition or destabilization of the target mRNA. The RefSeq represents the predicted microRNA stem-loop. [provided by RefSeq, September 2009]. Sequence Note: This record represents a predicted microRNA stem-loop as defined by miRBase. Some sequence at the 5' and 3' ends may not be included in the intermediate precursor miRNA produced by Drosha cleavage. microRNAs (miRNAs) are short (20-24 nt) non-coding RNAs that are involved in post-transcriptional regulation of gene expression in multicellular organisms by affecting both the stability and translation of mRNAs. miRNAs are transcribed by RNA polymerase II as part of capped and polyadenylated primary transcripts (pri-miRNAs) that can be either protein-coding or non-coding. The primary transcript is cleaved by the Drosha ribonuclease III enzyme to produce an approximately 70-nt stem-loop precursor miRNA (pre-miRNA), which is further cleaved by the cytoplasmic Dicer ribonuclease to generate the mature miRNA and antisense miRNA star (miRNA*) products. The mature miRNA is incorporated into a RNA-induced silencing complex (RISC), which recognizes target mRNAs through imperfect base pairing with the miRNA and most commonly results in translational inhibition or destabilization of the target mRNA. The RefSeq represents the predicted microRNA stem-loop. [provided by RefSeq, September 2009]. Sequence Note: This record represents a predicted microRNA stem-loop as defined by miRBase. Some sequence at the 5' and 3' ends may not be included in the intermediate precursor miRNA produced by Drosha cleavage. microRNAs (miRNAs) are short (20-24 nt) non-coding RNAs that are involved in post-transcriptional regulation of gene expression in multicellular organisms by affecting both the stability and translation of mRNAs. miRNAs are transcribed by RNA polymerase II as part of capped and polyadenylated primary transcripts (pri-miRNAs) that can be either protein-coding or non-coding. The primary transcript is cleaved by the Drosha ribonuclease III enzyme to produce an approximately 70-nt stem-loop precursor miRNA (pre-miRNA), which is further cleaved by the cytoplasmic Dicer ribonuclease to generate the mature miRNA and antisense miRNA star (miRNA*) products. The mature miRNA is incorporated into a RNA-induced silencing complex (RISC), which recognizes target mRNAs through imperfect base pairing with the miRNA and most commonly results in translational inhibition or destabilization of the target mRNA. The RefSeq represents the predicted microRNA stem-loop. [provided by RefSeq, September 2009]. Sequence Note: This record represents a predicted microRNA stem-loop as defined by miRBase. Some sequence at the 5' and 3' ends may not be included in the intermediate precursor miRNA produced by Drosha cleavage.
MIR3910-1	exonic	100500821		microRNAs (miRNAs) are short (20-24 nt) non-coding RNAs that are involved in post-transcriptional regulation of gene expression in multicellular organisms by affecting both the stability and translation of mRNAs. miRNAs are transcribed by RNA polymerase II as part of capped and polyadenylated primary transcripts (pri-miRNAs) that can be either protein-coding or non-coding. The primary transcript is cleaved by the Drosha ribonuclease III enzyme to produce an approximately 70-nt stem-loop precursor miRNA (pre-miRNA), which is further cleaved by the cytoplasmic Dicer ribonuclease to generate the mature miRNA and antisense miRNA star (miRNA*) products. The mature miRNA is incorporated into a RNA-induced silencing complex (RISC), which recognizes target mRNAs through imperfect base pairing with the miRNA and most commonly results in translational inhibition or destabilization of the target mRNA. The RefSeq represents the predicted microRNA stem-loop. [provided by RefSeq, September 2009]. Sequence Note: This record represents a predicted microRNA stem-loop as defined by miRBase. Some sequence at the 5' and 3' ends may not be included in the intermediate precursor miRNA produced by Drosha cleavage. microRNAs (miRNAs) are short (20-24 nt) non-coding RNAs that are involved in post-transcriptional regulation of gene expression in multicellular organisms by affecting both the stability and translation of mRNAs. miRNAs are transcribed by RNA polymerase II as part of capped and polyadenylated primary transcripts (pri-miRNAs) that can be either protein-coding or non-
MIR3910-2	exonic	100500902		microRNAs (miRNAs) are short (20-24 nt) non-coding RNAs that are involved in post-transcriptional regulation of gene expression in multicellular organisms by affecting both the stability and translation of mRNAs. miRNAs are transcribed by RNA polymerase II as part of capped and polyadenylated primary transcripts (pri-miRNAs) that can be either protein-coding or non-

TABLE 3-continued

A non-redundant list of genes listed in Table 2

RefSeq	Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary	Gene # (GN)
MIR4267		exonic	100422994		coding. The primary transcript is cleaved by the Drosha ribonuclease III enzyme to produce an approximately 70-nt stem-loop precursor miRNA (pre-miRNA), which is further cleaved by the cytoplasmic Dicer ribonuclease to generate the mature miRNA and antisense miRNA star (miRNA*) products. The mature miRNA is incorporated into a RNA-induced silencing complex (RISC), which recognizes target mRNAs through imperfect base pairing with the miRNA and most commonly results in translational inhibition or destabilization of the target mRNA. The RefSeq represents the predicted microRNA stem-loop. [provided by RefSeq, September 2009]. Sequence Note: This record represents a predicted microRNA stem-loop as defined by miRBase. Some sequence at the 5' and 3' ends may not be included in the intermediate precursor miRNA produced by Drosha cleavage.	86
MIR4436B1		exonic	100616123		microRNAs (miRNAs) are short (20-24 nt) non-coding RNAs that are involved in post-transcriptional regulation of gene expression in multicellular organisms by affecting both the stability and translation of mRNAs. miRNAs are transcribed by RNA polymerase II as part of capped and polyadenylated primary transcripts (pri-miRNAs) that can be either protein-coding or non-coding. The primary transcript is cleaved by the Drosha ribonuclease III enzyme to produce an approximately 70-nt stem-loop precursor miRNA (pre-miRNA), which is further cleaved by the cytoplasmic Dicer ribonuclease to generate the mature miRNA and antisense miRNA star (miRNA*) products. The mature miRNA is incorporated into a RNA-induced silencing complex (RISC), which recognizes target mRNAs through imperfect base pairing with the miRNA and most commonly results in translational inhibition or destabilization of the target mRNA. The RefSeq represents the predicted microRNA stem-loop. [provided by RefSeq, September 2009]. Sequence Note: This record represents a predicted microRNA stem-loop as defined by miRBase. Some sequence at the 5' and 3' ends may not be included in the intermediate precursor miRNA produced by Drosha cleavage.	87
MIR4436B2		exonic	100847033		microRNAs (miRNAs) are short (20-24 nt) non-coding RNAs that are involved in post-transcriptional regulation of gene expression in multicellular organisms by affecting both the stability and translation of mRNAs. miRNAs are transcribed by RNA polymerase II as part of capped and polyadenylated primary transcripts (pri-miRNAs) that can be either protein-coding or non-coding. The primary transcript is cleaved by the Drosha ribonuclease III enzyme to produce an approximately 70-nt stem-loop precursor miRNA (pre-miRNA), which is further cleaved by the cytoplasmic Dicer ribonuclease to generate the mature miRNA and antisense miRNA star (miRNA*) products. The mature miRNA is incorporated into a RNA-induced silencing complex (RISC), which recognizes target mRNAs through imperfect base pairing with the miRNA and most commonly results in translational inhibition or destabilization of the target mRNA. The RefSeq represents the predicted microRNA stem-loop. [provided by RefSeq, September 2009]. Sequence Note: This record represents a predicted microRNA stem-loop as defined by miRBase. Some sequence at the 5' and 3' ends may not be included in the intermediate precursor miRNA produced by Drosha cleavage.	88

A non-redundant list of genes listed in Table 2

RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary	Gene # (GN)
MRPL42	exonic	28977	39S ribosomal protein L42, mitochondrial precursor	transcribed by RNA polymerase II as part of capped and polyadenylated primary transcripts (pri-miRNAs) that can be either protein-coding or non-coding. The primary transcript is cleaved by the Drosha ribonuclease III enzyme to produce an approximately 70-nt stem-loop precursor miRNA (pre-miRNA), which is further cleaved by the cytoplasmic Dicer ribonuclease to generate the mature miRNA and antisense miRNA star (miRNA*) products. The mature miRNA is incorporated into a RNA-induced silencing complex (RISC), which recognizes target mRNAs through imperfect base pairing with the miRNA and most commonly results in translational inhibition or destabilization of the target mRNA. The RefSeq represents the predicted microRNA stem-loop. [provided by RefSeq, September 2009] Sequence Note: This record represents a predicted microRNA stem-loop as defined by miRBase. Some sequence at the 5' and 3' ends may not be included in the intermediate precursor miRNA produced by Drosha cleavage.	89
MRPL42	exonic	28977	39S ribosomal protein L42, mitochondrial precursor	The protein encoded by this gene interacts with the transcription factor myocardin, a key regulator of smooth muscle cell differentiation. The encoded protein is predominantly nuclear and may help transduce signals from the cytoskeleton to the nucleus. This gene is involved in a specific translocation event that creates a fusion of this gene and the RNA-binding motif protein-15 gene. This translocation has been associated with acute megakaryocytic leukemia. [provided by RefSeq, July 2008] Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: AB037859.2, AJ297258.1 [ECO: 0000332] RNAseq introns :: mixed/partial sample support ERS025081, ERS025082 [ECO: 0000350] ##Evidence-Data-END## Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a protein identified as belonging to both the 28S and the 39S subunits. Alternative splicing results in multiple transcript variants. Pseudogenes corresponding to this gene are found on chromosomes 4q, 6p, 6q, 7p, and 15q. [provided by RefSeq, May 2011]. Transcript Variant: This variant (1) encodes the supported protein. Both variants 1 and 2 encode the same protein. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments. ##RefSeq-Attributes-START## gene product(s) localized to mito. :: reported by MitoCarta ##RefSeq-Attributes-END## ##Evidence-Data-START## Transcript exon combination :: AK000285.1, AF151038.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025081, ERS025082 [ECO: 0000348] ##Evidence-Data-END##	90

TABLE 3-continued

A non-redundant list of genes listed in Table 2					
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary	Gene # (GN)
MTHFD1	exonic	4522	C-1-tetrahydrofolate synthase, cytoplasmic	This gene encodes a protein that possesses three distinct enzymatic activities, 5,10-methylenetetrahydrofolate dehydrogenase, 5,10-methylenetetrahydrofolate cyclohydrolase and 10-formyltetrahydrofolate synthetase. Each of these activities catalyzes one of three sequential reactions in the interconversion of 1-carbon derivatives of tetrahydrofolate, which are substrates for methionine, thymidylate, and de novo purine syntheses. The trifunctional enzymatic activities are conferred by two major domains, an aminoterminal portion containing the dehydrogenase and cyclohydrolase activities and a larger synthetase domain. [provided by RefSeq, July 2008]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##RefSeq-Attributes-START## CDS uses downstream in-frame AUG :: experimental evidence (PMID:3053686) ##RefSeq-Attributes-END## ##Evidence-Data-START## Transcript exon combination :: BC050420.1, J04031.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025081, ERS025082 [ECO: 0000348] ##Evidence-Data-END## Expression of interleukin-3 (IL3; MIM 147740) is restricted to activated T cells, natural killer (NK) cells, and mast cell lines. Transcription initiation depends on the activating capacity of specific protein factors, such as NFIL3, that bind to regulatory regions of the gene, usually upstream of the transcription start site (Zhang et al., 1995 [PubMed 7565758]). [supplied by OMIM, February 2009]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: S79880.1, U26173.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025081, ERS025082 [ECO: 0000348] ##Evidence-Data-END##	91
NFIL3	exonic	4783	nuclear factor interleukin-3-regulated protein	This gene encodes a member of the CATERPILLER family of cytoplasmic proteins. The encoded protein, which contains an N-terminal pyrin domain, a NACHT domain, a NACHT-associated domain, and a C-terminus leucine-rich repeat region, functions as an attenuating factor of inflammation by suppressing inflammatory responses in activated monocytes. Mutations in this gene cause familial cold autoinflammatory syndrome type 2. Alternative splicing results in multiple transcript variants. [provided by RefSeq, March 2013]. Transcript Variant: This variant (2) uses an alternate splice site in the central coding region, compared to variant 3, resulting in an isoform (2) that is 1 aa shorter than isoform 3. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: AY095146.1, BC028069.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025089 [ECO: 0000348] ##Evidence-Data-END##	92
NLRP12	exonic	91662	NACHT, LRR and PYD domains-containing protein 12 isoform 2	This gene encodes a member of the CATERPILLER family of cytoplasmic proteins. The encoded protein, which contains an N-terminal pyrin domain, a NACHT domain, a NACHT-associated domain, and a C-terminus leucine-rich repeat region, functions as an attenuating factor of inflammation by suppressing inflammatory responses in activated monocytes. Mutations in this gene cause familial cold autoinflammatory syndrome type 2. Alternative splicing results in multiple transcript variants. [provided by RefSeq, March 2013]. Transcript Variant: This variant (2) uses an alternate splice site in the central coding region, compared to variant 3, resulting in an isoform (2) that is 1 aa shorter than isoform 3. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: AY095146.1, BC028069.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025089 [ECO: 0000348] ##Evidence-Data-END##	93
NQO2	exonic	4835	ribosyl-dihydroquinone dehydrogenase [quinone]	NQO2 (EC 1.10.99.2) is a flavoprotein that catalyzes the 2-electron reduction of various quinones, redox dyes, and the vitamin K menadione. NQO2 predominantly uses dihydronicotinamide riboside (NRH) as the electron donor (summary by Wu et al., 1997 [PubMed 9367528]). [supplied by OMIM, July 2010]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon	94

TABLE 3-continued

A non-redundant list of genes listed in Table 2				
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary
NRIP1	exonic	8204	nuclear receptor-interacting protein 1	combination :: J02888.1, AK311746.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025081, ERS025082 [ECO: 0000348] ##Evidence-Data-END## Nuclear receptor interacting protein 1 (NRIP1) is a nuclear protein that specifically interacts with the hormone-dependent activation domain AF2 of nuclear receptors. Also known as RIP140, this protein modulates transcriptional activity of the estrogen receptor. [provided by RefSeq, July 2008]. Sequence Note: The RefSeq transcript and protein were derived from transcript and genomic sequence to make the sequence consistent with the reference genome assembly. The extent of this RefSeq transcript is supported by transcript alignments. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: AK289786.1, DA230125.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025098 [ECO: 0000348] ##Evidence-Data-END## The protein encoded by this gene regulates the turnover of dihydrophosphoinositol polyphosphates. The turnover of these high-energy dihydrophosphoinositol polyphosphates represents a molecular switching activity with important regulatory consequences. Molecular switching by dihydrophosphoinositol polyphosphates may contribute to regulating intracellular trafficking. Several alternatively spliced transcript variants have been described, but the full-length nature of some variants has not been determined. Isoforms DIPP2alpha and DIPP2beta are distinguishable from each other solely by DIPP2beta possessing one additional amino acid due to intron boundary skidding in alternate splicing. [provided by RefSeq, July 2008]. Transcript Variant: This variant (1) encodes the predominant isoform (alpha). ##Evidence-Data-START## Transcript exon combination :: AF191651.1, AF191650.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025081, ERS025082 [ECO: 0000348] ##Evidence-Data-END##
NUDT4	exonic	11163	diphosphoinositol polyphosphate phosphohydrolase 2 isoform alpha	
NUDT4P1	exonic	440672	N/A	
OVOL2	exonic	58495	transcription factor Ovo-like 2	
PDE3B	intronic	5140	cGMP-inhibited 3',5'-cyclic phosphodiesterase B	
PDGFRA	exonic	5156	platelet-derived growth factor receptor alpha precursor	This gene encodes a cell surface tyrosine kinase receptor for members of the platelet-derived growth factor family. These growth factors are mitogens for cells of mesenchymal origin. The identity of the growth factor bound to a receptor monomer determines whether the functional receptor is a homodimer or a heterodimer, composed of both platelet-derived growth factor receptor alpha and beta polypeptides. Studies suggest that this gene plays a role in organ development, wound healing, and tumor progression. Mutations in this gene have been associated with idiopathic hyperostosis syndrome, somatic and familial gastrointestinal stromal tumors, and a variety of other cancers. [provided by RefSeq, March 2012]. Sequence Note: This RefSeq record was created from transcript and genomic sequence data because no single transcript was available for the full length of the gene. The extent of this transcript is supported by transcript alignments and orthologous data. Publication Note:
		97		
		98		
		99		
		100		

TABLE 3-continued

A non-redundant list of genes listed in Table 2				
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary
PDSS2	exonic	57107	decaprenyl-diphosphate synthase subunit 2	This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: M21574.1, M22734.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025084 [ECO: 0000348] ##Evidence-Data-END## The protein encoded by this gene is an enzyme that synthesizes the prenyl side-chain of coenzyme Q, or ubiquinone, one of the key elements in the respiratory chain. The gene product catalyzes the formation of all trans-polyisoprenyl pyrophosphates from isopentenyl diphosphate in the assembly of polyisoprenoid side chains, the first step in coenzyme Q biosynthesis. Defects in this gene are a cause of coenzyme Q10 deficiency. [provided by RefSeq, October 2009]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: BC039906.1, AF254956.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025084, ERS025088 [ECO: 0000348] ##Evidence-Data-END##
PHACTR4	exonic	65979	phosphatase and actin regulator 4 isoform 1	This gene encodes a member of the phosphatase and actin regulator (PHACTR) family. Other PHACTR family members have been shown to inhibit protein phosphatase 1 (PP1) activity, and the homolog of this gene in the mouse has been shown to interact with actin and PP1. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, July 2008]. Transcript Variant: This variant (1) represents the longer transcript but encodes the shorter isoform (1). Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments. ##Evidence-Data-START## Transcript exon combination :: CR749449.1, BC029266.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025084, ERS025087 [ECO: 0000348] ##Evidence-Data-END##
PIAS2	exonic	9063	E3 SUMO-protein ligase PIAS2 isoform alpha	This gene encodes a member of the protein inhibitor of activated STAT (PIAS) family. PIAS proteins function as SUMO E3 ligases and play important roles in many cellular processes by mediating the sumoylation of target proteins. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. Isoforms of the encoded protein enhance the sumoylation of specific target proteins including the p53 tumor suppressor protein, c-Jun, and the androgen receptor. A pseudogene of this gene is located on the short arm of chromosome 4. The symbol MIZ1 has also been associated with ZBTB17 which is a different gene located on chromosome 1. [provided by RefSeq, August 2011]. Transcript Variant: This variant (alpha) utilizes an alternate 3' coding exon, compared to variant beta, resulting in a shorter isoform (alpha) that has a unique C-terminus compared to isoform beta. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: BC015190.1 [ECO: 0000332] RNAseq introns ::

TABLE 3-continued

A non-redundant list of genes listed in Table 2				
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary
PIK3CD	exonic	5293	phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit delta isoform	single sample supports all introns ERS025088 [ECO: 0000348] ##Evidence-Data-END## Phosphoinositide 3-kinases (PI3Ks) phosphorylate inositol lipids and are involved in the immune response. The protein encoded by this gene is a class I PI3K found primarily in leukocytes. Like other class I PI3Ks (p110-alpha, p110-beta, and p110-gamma), the encoded protein binds p85 adapter proteins and GTP-bound RAS. However, unlike the other class I PI3Ks, this protein phosphorylates itself, not p85 protein. [provided by RefSeq, July 2010]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: U86453.1, Y10055.2 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025089 [ECO: 0000348] ##Evidence-Data-END## The protein encoded by this gene is predicted to have a single transmembrane (TM)-spanning domain and multiple copies of an immunoglobulin-like plexin-transcription-factor domain. Alternative splicing results in two transcript variants encoding different isoforms. Other alternatively spliced transcripts have been described, but the full length sequences have not been determined. Several of these transcripts are predicted to encode truncated products which lack the TM and may be secreted. Mutations in this gene cause autosomal recessive polycystic kidney disease, also known as polycystic kidney and hepatic disease-1. [provided by RefSeq, July 2008]. Transcript Variant: This variant (1) encodes the longer isoform of this protein. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: AY074797.1, AF480664.1 [ECO: 0000332] RNAseq introns :: mixed/partial sample support ERS025084, ERS025085 [ECO: 0000350] ##Evidence-Data-END## This gene encodes a member of the plexin family. Plexins are transmembrane receptors for semaphorins, a large family of proteins that regulate axon guidance, cell motility and migration, and the immune response. The encoded protein and its ligand regulate melanocyte adhesion, and viral semaphorins may modulate the immune response by binding to this receptor. The encoded protein may be a tumor suppressor protein for melanoma. Alternatively spliced transcript variants have been observed for this gene. [provided by RefSeq, January 2011]. Transcript Variant: This variant (2) lacks multiple 5' exons but contains an alternate 5' exon, compared to variant 1. This variant is represented as non-coding due to the presence of an upstream ORF that is predicted to interfere with translation of the longest in-frame ORF. Translation of the upstream ORF renders the transcript a candidate for nonsense-mediated mRNA decay (NMD). Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## RNAseq introns :: single sample supports all introns ERS025084, ERS025088 [ECO: 0000348] ##Evidence-Data-END##
PKHD1	intronic	5314	fibrocystin isoform 1 precursor	
PLXNC1	exonic	10154	Plexin C1	

TABLE 3-continued

A non-redundant list of genes listed in Table 2				
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary
PNPLA4	exonic	8228	patatin-like phospholipase domain-containing protein 4 isoform 1 precursor	This gene encodes a member of the patatin-like family of phospholipases. The encoded enzyme has both triacylglycerol lipase and transacylase activities and may be involved in adipocyte triglyceride homeostasis. Alternate splicing results in multiple transcript variants. A pseudogene of this gene is found on chromosome Y. [provided by RefSeq, February 2010]. Transcript Variant: This variant (1) represents the longest transcript and encodes the longer isoform (1). Variants 1 and 2 encode the same isoform (1). Sequence Note: The RefSeq transcript and protein were derived from transcript and genomic sequence to make the sequence consistent with the reference genome assembly. The extent of this transcript is supported by transcript alignments. ##Evidence-Data-START## Transcript exon combination :: U03886.1, AK289888.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns [ECO: 0000332] RNAseq introns [ECO: 0000348] ##Evidence-Data-END##
PNPT1	both	87178	polynucleotide phosphatase nucleotidyltransferase 1, mitochondrial precursor	The protein encoded by this gene belongs to the evolutionary conserved polynucleotide phosphatase family comprised of phosphate dependent 3'-to-5' exonucleases implicated in RNA processing and degradation. This enzyme is predominantly localized in the mitochondrial intermembrane space and is involved in import of RNA to mitochondria. Mutations in this gene have been associated with combined oxidative phosphorylation deficiency-13 and autosomal recessive nonsyndromic deafness-70. Related pseudogenes are found on chromosomes 3 and 7. [provided by RefSeq, December 2012]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: BC053660.1, AJ458465.1 [ECO: 0000332] RNAseq introns :: mixed/partial sample support ERS025081, ERS025082 [ECO: 0000350] ##Evidence-Data-END## ##RefSeq-Attributes-START## gene product(s) localized to mito. :: PMID: 12798676; reported by MitoCarta ##RefSeq-Attributes-END## Protein phosphatase 2 (formerly named type 2A) is one of the four major Ser/Thr phosphatases and is implicated in the negative control of cell growth and division. Protein phosphatase 2 holoenzymes are heterotrimeric proteins composed of a structural subunit A, a catalytic subunit C, and a regulatory subunit B. The regulatory subunit is encoded by a diverse set of genes that have been grouped into the B/PR55, B/PR61, and B/PR72 families. These different regulatory subunits confer distinct enzymatic specificities and intracellular localizations to the holoenzyme. The product of this gene belongs to the B" family. The B" family has been further divided into subfamilies. The product of this gene belongs to the beta subfamily of regulatory subunit B". [provided by RefSeq, April 2010]. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments. ##Evidence-Data-START## Transcript exon combination :: BK000521.1, BC063429.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns [ECO: 0000332] RNAseq introns [ECO: 0000348] ##Evidence-Data-END##
PPP2R3B	intronic	28227	serine/threonine-protein phosphatase 2A regulatory subunit B" subunit beta	Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family
PRKCB	both	5579	protein kinase C beta type isoform 1	

A non-redundant list of genes listed in Table 2

RefSeq Gene Symbol Exon overlap NCBI Gene ID Gene Description RefSeq_Summary Gene # (GN)

PNPLA4 8228 patatin-like phospholipase domain-containing protein 4 isoform 1 precursor This gene encodes a member of the patatin-like family of phospholipases. The encoded enzyme has both triacylglycerol lipase and transacylase activities and may be involved in adipocyte triglyceride homeostasis. Alternate splicing results in multiple transcript variants. A pseudogene of this gene is found on chromosome Y. [provided by RefSeq, February 2010]. Transcript Variant: This variant (1) represents the longest transcript and encodes the longer isoform (1). Variants 1 and 2 encode the same isoform (1). Sequence Note: The RefSeq transcript and protein were derived from transcript and genomic sequence to make the sequence consistent with the reference genome assembly. The extent of this transcript is supported by transcript alignments. ##Evidence-Data-START## Transcript exon combination :: U03886.1, AK289888.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns [ECO: 0000332] RNAseq introns [ECO: 0000348] ##Evidence-Data-END## 107

PNPT1 87178 polynucleotide phosphatase nucleotidyltransferase 1, mitochondrial precursor The protein encoded by this gene belongs to the evolutionary conserved polynucleotide phosphatase family comprised of phosphate dependent 3'-to-5' exonucleases implicated in RNA processing and degradation. This enzyme is predominantly localized in the mitochondrial intermembrane space and is involved in import of RNA to mitochondria. Mutations in this gene have been associated with combined oxidative phosphorylation deficiency-13 and autosomal recessive nonsyndromic deafness-70. Related pseudogenes are found on chromosomes 3 and 7. [provided by RefSeq, December 2012]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: BC053660.1, AJ458465.1 [ECO: 0000332] RNAseq introns :: mixed/partial sample support ERS025081, ERS025082 [ECO: 0000350] ##Evidence-Data-END## ##RefSeq-Attributes-START## gene product(s) localized to mito. :: PMID: 12798676; reported by MitoCarta ##RefSeq-Attributes-END## 108

PPP2R3B 28227 serine/threonine-protein phosphatase 2A regulatory subunit B" subunit beta Protein phosphatase 2 (formerly named type 2A) is one of the four major Ser/Thr phosphatases and is implicated in the negative control of cell growth and division. Protein phosphatase 2 holoenzymes are heterotrimeric proteins composed of a structural subunit A, a catalytic subunit C, and a regulatory subunit B. The regulatory subunit is encoded by a diverse set of genes that have been grouped into the B/PR55, B/PR61, and B/PR72 families. These different regulatory subunits confer distinct enzymatic specificities and intracellular localizations to the holoenzyme. The product of this gene belongs to the B" family. The B" family has been further divided into subfamilies. The product of this gene belongs to the beta subfamily of regulatory subunit B". [provided by RefSeq, April 2010]. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments. ##Evidence-Data-START## Transcript exon combination :: BK000521.1, BC063429.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns [ECO: 0000332] RNAseq introns [ECO: 0000348] ##Evidence-Data-END## 109

PRKCB 5579 protein kinase C beta type isoform 1 Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family 110

TABLE 3-continued

A non-redundant list of genes listed in Table 2				
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary
				members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role in cells. The protein encoded by this gene is one of the PKC family members. This protein kinase has been reported to be involved in many different cellular functions, such as B cell activation, apoptosis induction, endothelial cell proliferation, and intestinal sugar absorption. Studies in mice also suggest that this kinase may also regulate neuronal functions and correlate fear-induced conflict behavior after stress. Alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, July 2008]. Transcript Variant: This variant (1) uses an alternate splice junction at the 5' end of the last exon compared to variant 2. The resulting isoform (1) has a distinct and shorter C-terminus compared to isoform 2. Sequence Note: This RefSeq record was created from transcript and genomic sequence data because no single transcript was available for the full length of the gene. The extent of this transcript is supported by transcript alignments. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: X06318.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025082, ERS025084 [ECO: 0000348] ##Evidence-Data-END##
PRKCH	intronic	5583	protein kinase C eta type	Protein Kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role in cells. The protein encoded by this gene is one of the PKC family members. It is a calcium-independent and phospholipids-dependent protein kinase. It is predominantly expressed in epithelial tissues and has been shown to reside specifically in the cell nucleus. This protein kinase can regulate keratinocyte differentiation by activating the MAP kinase MAPK13 (p38delta)-activated protein kinase cascade that targets CCAAT/enhancer-binding protein alpha (CEBPA). It is also found to mediate the transcription activation of the transglutaminase 1 (TGM1) gene. [provided by RefSeq, July 2008]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: BC037268.1, AK290183.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025081, ERS025083 [ECO: 0000348] ##Evidence-Data-END##
PSTPIP1	exonic	9051	proline-serine-threonine phosphatase-interacting protein 1	The protein encoded by this gene binds to the cytoplasmic tail of CD2, an effector of T cell activation and adhesion, negatively affecting CD2-triggered T cell activation. The encoded protein appears to be a scaffold protein and a regulator of the actin cytoskeleton. It has also been shown to bind ABL1, PTPN18, WAS, CD2AP, and PTPN12. Mutations in this gene are a cause of PAPA syndrome. [provided by RefSeq, July 2008]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-
				111
				112

TABLE 3-continued

A non-redundant list of genes listed in Table 2				
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary
PTPN2	exonic	5771	tyrosine-protein phosphatase non-receptor type 2 isoform 1	Data-START## Transcript exon combination :: BC008602.1, U94778.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025084, ERS025088 [ECO: 0000348] ##Evidence-Data-END## The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. Members of the PTP family share a highly conserved catalytic motif, which is essential for the catalytic activity. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. Epidermal growth factor receptor and the adaptor protein Shc were reported to be substrates of this PTP, which suggested the roles in growth factor mediated cell signaling. Multiple alternatively spliced transcript variants encoding different isoforms have been found. Two highly related but distinctly processed pseudogenes that localize to chromosomes 1 and 13, respectively, have been reported. [provided by RefSeq, May 2011]. Transcript Variant: This variant (1) encodes the longest isoform (1). Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: M25393.1, AK292570.1 [ECO: 0000332] RNAseq introns :: mixed/partial sample support ERS025081, ERS025082 [ECO: 0000350] ##Evidence-Data-END## The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP possesses an extracellular region, a single transmembrane region, and a single intracellular catalytic domain, and thus represents a receptor-type PTP. The catalytic domain of this PTP is most closely related to PTPRN/IA-2beta. This PTP and PTPRN are both found to be major autoantigens associated with insulin-dependent diabetes mellitus. Three alternatively spliced transcript variants of this gene, which encode distinct proteins, have been reported. [provided by RefSeq, July 2008]. Transcript Variant: This variant (1) encodes the longest isoform (1). Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: U66702.1, AF007555.1 [ECO: 0000332] RNAseq introns :: mixed/partial sample support ERS025081, ERS025082 [ECO: 0000350] ##Evidence-Data-END## Rab proteins are low molecular mass GTPases that are critical regulators of vesicle trafficking. For additional background information on Rab proteins, see MIM 179508. [supplied by OMIM, April 2006]. Transcript Variant: This variant (2) represents use of an alternate promoter, 5' UTR, and alternate start codon, and includes an alternate coding exon, compared to variant 3. The resulting isoform (2) has a distinct and longer N-terminus, compared to isoform 3. ##Evidence-Data-START## Transcript exon combination :: AK098068.1, BX332255.2 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025084, ERS025088 [ECO: 0000348] ##Evidence-Data-END##
PTPRN2	intronic	5799	receptor-type tyrosine-protein phosphatase N2 isoform 1 precursor	113
RAB37	exonic	326624	ras-related protein Rab-37 isoform 2	114
				115

TABLE 3-continued

A non-redundant list of genes listed in Table 2					
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary	Gene # (GN)
RBFOX1	intronic	54715	RNA binding protein fox-1 homolog 1 isoform 1	The Fox-1 family of RNA-binding proteins is evolutionarily conserved, and regulates tissue-specific alternative splicing in metazoa. Fox-1 recognizes a (U)GCAUG stretch in regulated exons or in flanking introns. The protein binds to the C-terminus of ataxin-2 and may contribute to the restricted pathology of spinocerebellar ataxia type 2 (SCA2). Ataxin-2 is the product of the SCA2 gene which causes familial neurodegenerative diseases. Fox-1 and ataxin-2 are both localized in the trans-Golgi network. Several alternatively spliced transcript variants encoding different isoforms have been found for this gene, [provided by RefSeq, November 2011]. Transcript Variant: This variant (1), also known as gamma, encodes the longest isoform (1). Sequence Note: This RefSeq record was created from transcript and genomic sequence data because no single transcript was available for the full length of the gene. The extent of this transcript is supported by transcript alignments. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: AF229057.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025083, ERS025099 [ECO: 0000348] ##Evidence-Data-END##	116
RCCL RGCC	exonic exonic	1104 28984	N/A regulator of cell cycle RGCC	This gene is thought to regulate cell cycle progression. It is induced by p53 in response to DNA damage, or by sublytic levels of complement system proteins that result in activation of the cell cycle. The encoded protein localizes to the cytoplasm during interphase and to centrosomes during mitosis. The protein forms a complex with polo-like kinase 1. The protein also translocates to the nucleus in response to treatment with complement system proteins, and can associate with and increase the kinase activity of cell division cycle 2 protein. In different assays and cell types, overexpression of this protein has been shown to activate or suppress cell cycle progression. [provided by RefSeq, July 2008]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: BC066334.1, BG037019.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025081, ERS025092 [ECO: 0000348] ##Evidence-Data-END##	117 118
RHOQ	intronic	23433	rho-related GTP-binding protein RhoQ precursor	This gene encodes a member of the Rho family of small GTPases, which cycle between inactive GDP-bound and active GTP-bound states and function as molecular switches in signal transduction cascades. Rho proteins promote reorganization of the actin cytoskeleton and regulate cell shape, attachment, and motility. The encoded protein is an important signalling protein for sarcomere assembly and has been shown to play a significant role in the exocytosis of the solute carrier family 2, facilitated glucose transporter member 4 and other proteins, possibly acting as the signal that turns on the membrane fusion machinery. Three related pseudogene have been identified on chromosomes 2 and 14. [provided by RefSeq, August 2011]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: BX428852.2, BC013151.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025081, ERS025082 [ECO: 0000348] ##Evidence-Data-END##	119

TABLE 3-continued

A non-redundant list of genes listed in Table 2					
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary	Gene # (GN)
RNASE3	exonic	6037	eosinophil cationic protein precursor	N/A	120
RNASE10	exonic	338879	inactive ribonuclease-like protein 10 precursor	N/A	121
RPL38	exonic	6169	60S ribosomal protein L38	Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L38E family of ribosomal proteins. It is located in the cytoplasm. Alternative splice variants have been identified, both encoding the same protein. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome, including one located in the promoter region of the type 1 angiotensin II receptor gene. [provided by RefSeq, July 2008]. Transcript Variant: This variant (1) is the longer and predominant transcript. Variants 1 and 2 encode the same protein. ##Evidence-Data-START## Transcript exon combination :: BQ276548.1, BU569438.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025081, ERS025082 [ECO: 0000348] ##Evidence-Data-END##	122
RPTOR	intronic	57521	regulatory-associated protein of mTOR isoform 1	This gene encodes a component of a signaling pathway that regulates cell growth in response to nutrient and insulin levels. The encoded protein forms a stoichiometric complex with the mTOR kinase, and also associates with eukaryotic initiation factor 4E-binding protein-1 and ribosomal protein S6 kinase. The protein positively regulates the downstream effector ribosomal protein S6 kinase, and negatively regulates the mTOR kinase. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, September 2009]. Transcript Variant: This variant (1) represents the longer transcript and encodes the longer isoform (1). Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: AY090663.1, BC136652.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025083, ERS025085 [ECO: 0000348] ##Evidence-Data-END##	123
SERPINB4	exonic	6318	serpin B4	N/A	124
SERPINB6	exonic	5269	serpin B6 isoform a	The protein encoded by this gene is a member of the serpin (serine proteinase inhibitor) superfamily, and ovalbumin(ov)-serpin subfamily. It was originally discovered as a placental thrombin inhibitor. The mouse homolog was found to be expressed in the hair cells of the inner ear. Mutations in this gene are associated with nonsyndromic progressive hearing loss, suggesting that this serpin plays an important role in the inner ear in the protection against leakage of lysosomal content during stress, and that loss of this protection results in cell death and sensorineural hearing loss. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, September 2010]. Transcript Variant: This variant (1) represents the predominant transcript. Variants 1, 5 and 6 encode the same isoform (a). Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START##	125

TABLE 3-continued

A non-redundant list of genes listed in Table 2				
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary
SLC3A2	both	6520	4F2 cell-surface antigen heavy chain isoform b	Transcript exon combination :: AK314578.1, BC098564.1 [ECO: 0000332] RNAseq introns :: mixed/partial sample support ERS025081, ERS025082 [ECO: 0000350] ##Evidence-Data-END## This gene is a member of the solute carrier family and encodes a cell surface, transmembrane protein. The protein exists as the heavy chain of a heterodimer, covalently bound through di-sulfide bonds to one of several possible light chains. The encoded transporter plays a role in regulation of intracellular calcium levels and transports L-type amino acids. Alternatively spliced transcript variants, encoding different isoforms, have been characterized. [provided by RefSeq, November 2010]. Transcript Variant: This variant (2) represents the longest transcript and encodes the longest isoform (b). Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: AK025584.1 [ECO: 0000332] RNAseq introns :: mixed/partial sample support ERS025082, ERS025084 [ECO: 0000350] ##Evidence-Data-END## This gene encodes a membrane transporter that exports free sialic acids that have been cleaved off of cell surface lipids and proteins from lysosomes. Mutations in this gene cause sialic acid storage diseases, including infantile sialic acid storage disorder and Salla disease, an adult form. [provided by RefSeq, July 2008]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: BC020961.2, AJ387747.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025081, ERS025082 [ECO: 0000348] ##Evidence-Data-END##
SLC17A5	both	26503	sialin	
SNHG3	exonic	8420	N/A	N/A
SNORD17	exonic	692086	N/A	N/A
SNX5	exonic	27131	sorting nexin-5 isoform a	This gene encodes a member of the sorting nexin family. Members of this family contain a phox (PX) domain, which is a phosphoinositide binding domain, and are involved in intracellular trafficking. This protein functions in endosomal sorting, the phosphoinositide-signaling pathway, and macropinocytosis. This gene may play a role in the tumorigenesis of papillary thyroid carcinoma. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, September 2013]. Transcript Variant: This variant (1) differs in the 5' UTR, compared to variant 2. Variants 1 and 2 encode the same protein (isoform a). ##Evidence-Data-START## Transcript exon combination :: BC000100.3, AF121855.1 [ECO: 0000332] RNAseq introns :: mixed/partial sample support ERS025081, ERS025082 [ECO: 0000350] ##Evidence-Data-END##
SOCS2	exonic	8835	suppressor of cytokine signaling 2	This gene encodes a member of the suppressor of cytokine signaling (SOCS) family. SOCS family members are cytokine-inducible negative regulators of cytokine receptor signaling via the Janus kinase/signal transducer and activation of transcription pathway (the JAK/STAT pathway). SOCS family proteins interact with major molecules of signaling complexes to block further signal transduction, in part, by proteasomal depletion of receptors or signal-transducing proteins via ubiquitination. The expression of this gene can be induced by a subset of cytokines, including erythropoietin, GM-CSF, IL10, interferon (IFN)-gamma and by cytokine receptors such as growth hormone
				126
				127
				128
				129
				130
				131

TABLE 3-continued

A non-redundant list of genes listed in Table 2					
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary	Gene # (GN)
SOCS2-AS1 ST8SIA5	exonic	144481	N/A	receptor. The protein encoded by this gene interacts with the cytoplasmic domain of insulin-like growth factor-1 receptor (IGF1R) and is thought to be involved in the regulation of IGF1R mediated cell signaling. This gene has pseudogenes on chromosomes 20 and 22. Alternative splicing results in multiple transcript variants. [provided by RefSeq, July 2012]. Transcript Variant: This variant (1) differs in the 5' UTR, compared to variant 5. Variants 1-6 encode the same protein. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: AK313165.1, AL522912.3 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025081, ERS025082 [ECO: 0000348] ##Evidence-Data-END## N/A	132
	exonic	29906	alpha-2,8-sialyltransferase 8E		133
STIM2	intronic	57620	stromal interaction molecule 2 isoform 1 precursor	The protein encoded by this gene is a type II membrane protein that may be present in the Golgi apparatus. The encoded protein, which is a member of glycosyltransferase family 29, may be involved in the synthesis of gangliosides GD1c, GT1a, GQ1b, and GT3 from GD1a, GT1b, GM1b, and GD3, respectively. [provided by RefSeq, July 2008]. ##Evidence-Data-START## Transcript exon combination :: AK056270.1, BC108910.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025082, ERS025084 [ECO: 0000348] ##Evidence-Data-END## This gene is a member of the stromal interaction molecule (STIM) family and likely arose, along with related family member STIM1, from a common ancestral gene. The encoded protein functions to regulate calcium concentrations in the cytosol and endoplasmic reticulum, and is involved in the activation of plasma membrane Orai Ca(2+) entry channels. This gene initiates translation from a non-AUG (UUG) start site. A signal peptide is cleaved from the resulting protein. Multiple transcript variants result from alternative splicing. [provided by RefSeq, December 2009]. Transcript Variant: This variant (1) encodes the longest isoform (1). Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: BC136449.1, AK096846.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025081, ERS025084 [ECO: 0000348] ##Evidence-Data-END## ##RefSeq-Attributes-START## CDS uses downstream in-frame AUG :: experimental evidence (PMID:11463338) non-AUG initiation codon :: PMID: 11463338 ##RefSeq-Attributes-END## N/A	134
TBC1D16	intronic	125058	TBC1 domain family member 16 isoform a	N/A	136
TEX29	exonic	121793	testis-expressed sequence 29 protein		137

TABLE 3-continued

A non-redundant list of genes listed in Table 2					
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary	Gene # (GN)
TNFRSF10A	exonic	8797	tumor necrosis factor receptor superfamily member 10A	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10/TRAIL), and thus transduces cell death signal and induces cell apoptosis. Studies with FADD-deficient mice suggested that FADD, a death domain containing adaptor protein, is required for the apoptosis mediated by this protein. [provided by RefSeq, July 2008]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: BC012866.1, AK291299.1 [ECO: 0000332] RNaseq introns :: single sample supports all introns ERS025081, ERS025084 [ECO: 0000348] ##Evidence-Data-END##	138
TNFRSF13C	exonic	115650	tumor necrosis factor receptor superfamily member 13C	B cell-activating factor (BAFF) enhances B-cell survival in vitro and is a regulator of the peripheral B-cell population. Overexpression of BAFF in mice results in mature B-cell hyperplasia and symptoms of systemic lupus erythematosus (SLE). Also, some SLE patients have increased levels of BAFF in serum. Therefore, it has been proposed that abnormally high levels of BAFF may contribute to the pathogenesis of autoimmune diseases by enhancing the survival of autoreactive B cells. The protein encoded by this gene is a receptor for BAFF and is a type III transmembrane protein containing a single extracellular cysteine-rich domain. It is thought that this receptor is the principal receptor required for BAFF-mediated mature B-cell survival. [provided by RefSeq, July 2008]. Sequence Note: The RefSeq transcript and protein were derived from genomic sequence to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on alignments. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: AF373846.1, BC112030.1 [ECO: 0000332] RNaseq introns :: single sample supports all introns ERS025089 [ECO: 0000348] ##Evidence-Data-END##	139
TNFRSF18	exonic	8784	tumor necrosis factor receptor superfamily member 18 isoform 1 precursor	This gene encodes a member of the TNF-receptor superfamily. The encoded receptor has been shown to have increased expression upon T-cell activation, and it is thought to play a key role in dominant immunological self-tolerance maintained by CD25(+)CD4(+) regulatory T cells. Knockout studies in mice also suggest the role of this receptor is in the regulation of CD3-driven T-cell activation and programmed cell death. Three alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq, February 2011]. Transcript Variant: This variant (1) represents the longest transcript. It contains an extra coding segment, which leads to a frame shift, compared to variant 2. The resulting protein (isoform 1) contains a distinct and shorter C-terminus, as compared to isoform 2. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: AY358877.1, AF125304.1 [ECO: 0000332] RNaseq introns ::	140

TABLE 3-continued

A non-redundant list of genes listed in Table 2				
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary
TRAFD1	exonic	10906	TRAF-type zinc finger domain-containing protein 1	single sample supports all introns ERS025089, ERS025093 [ECO: 0000348] ##Evidence-Data-END## The innate immune system confers host defense against viral and microbial infection, and TRAFD1 is a negative feedback regulator that controls excessive immune responses (Sanada et al., 2008 [PubMed 18849341]). [supplied by OMIM, December 2009]. Transcript Variant: This variant (1) represents the longer transcript. Variants 1 and 2 both encode the same protein. ##Evidence-Data-START## Transcript exon combination :: AK122620.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025083, ERS025084 [ECO: 0000348] ##Evidence-Data-END## The protein encoded by this gene is a calcium-permeable cation channel that is regulated by free intracellular ADP-ribose. The encoded protein is activated by oxidative stress and confers susceptibility to cell death. Several alternatively spliced transcript variants of this gene have been described, but their full-length nature is not known. [provided by RefSeq, July 2008]. Transcript Variant: This variant (2) uses an alternate in-frame splice junction at the 5' end of an exon compared to variant 1. This results in the introduction of a premature stop codon and renders the transcript a nonsense-mediated mRNA decay (NMD) candidate. Therefore, this transcript is not thought to be protein-coding. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## RNAseq introns :: mixed/partial sample support ERS025081, ERS025082 [ECO: 0000350] ##Evidence-Data-END## N/A
TRPM2	exonic	7226	Transient Receptor Potential Cation Channel Subfamily M Member 2	This gene encodes a member of the twenty family of proteins. Members of this family function as chloride anion channels. The encoded protein functions as a calcium(2+)-activated large conductance chloride(-) channel, and may play a role in kidney tumorigenesis. Two transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq, July 2008]. Transcript Variant: This variant (1) represents the longer transcript, and encodes the longer isoform (1). ##Evidence-Data-START## Transcript exon combination :: AF319952.1, BC107492.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025082, ERS025083 [ECO: 0000348] ##Evidence-Data-END## The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, or E1s, ubiquitin-conjugating enzymes, or E2s, and ubiquitin-protein ligases, or E3s. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. Studies in mouse suggest that this protein plays a role in DNA postreplication repair. [provided by RefSeq, July 2008]. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Gene record to access additional publications. ##Evidence-Data-START## Transcript exon combination :: BC000396.2, D83004.1 [ECO: 0000332] RNAseq introns :: single sample supports all introns ERS025084 [ECO: 0000348] ##Evidence-Data-END##
TTLL10	exonic	254173	inactive polyglycylase TTLL10 isoform 1	
TTYH2	exonic	94015	protein twenty homolog 2 isoform 1	
UBE2N	exonic	7334	ubiquitin-conjugating enzyme E2 N	

Gene # (GN)

141

142

143

144

145

TABLE 3-continued

A non-redundant list of genes listed in Table 2					
RefSeq Gene Symbol	Exon overlap	NCBI Gene ID	Gene Description	RefSeq_Summary	Gene # (GN)
VCX	exonic	26609	variable charge X-linked protein 1	This gene belongs to the VCXY gene family, which has multiple members on both X and Y chromosomes, and all are expressed exclusively in male germ cells. The X-linked members are clustered on chromosome Xp22 and Y-linked members are two identical copies of the gene within a palindromic region on Yq11. The family members share a high degree of sequence identity, with the exception that a 30-bp unit is tandemly repeated in X-linked members but occurs only once in Y-linked members. The VCX gene cluster is polymorphic in terms of copy number; different individuals may have a different number of VCX genes. VCXY genes encode small and highly charged proteins of unknown function. The presence of a putative bipartite nuclear localization signal suggests that VCXY members are nuclear proteins. This gene contains 10 repeats of the 30-bp unit. [provided by RefSeq, July 2008]. ##Evidence-Data-START## Transcript exon combination :: AF167081.2 [ECO: 0000332] ##Evidence-Data-END##	146
VSTM1	intronic	284415	V-set and transmembrane domain-containing protein 1 precursor	N/A	147
VWA2	exonic	340706	von Willebrand factor A domain-containing protein 2 precursor	This gene encodes a member of the von Willebrand factor A-like domain protein superfamily. The encoded protein is localized to the extracellular matrix and may serve as a structural component in basement membranes or in anchoring structures on scaffolds of collagen VII or fibrillin. This gene has been linked to type 1A diabetes and is a candidate serological marker for colon cancer. [provided by RefSeq, January 2013]. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments. CCDS Note: The coding region has been updated to represent an alternative 3' splicing pattern that is more supported by the available transcript and protein data. ##Evidence-Data-START## Transcript exon combination :: AY572972.1, AJ536328.2 [ECO: 0000332] RNAseq introns :: mixed/partial sample support ERS025081, ERS025084 [ECO: 0000350] ##Evidence-Data-END##	148
ZNF350	exonic	59348	zinc finger protein 350	N/A	149
ZNF432	exonic	9668	zinc finger protein 432	N/A	150
ZNF577	exonic	84765	N/A	N/A	151
ZNF613	exonic	79898	zinc finger protein 613 isoform 1	N/A	152
ZNF614	exonic	80110	zinc finger protein 614	N/A	153
ZNF615	exonic	284370	zinc finger protein 615 isoform 1	N/A	154
ZNF649	exonic	65251	zinc finger protein 649	N/A	155
ZNF841	exonic	284371	zinc finger protein 841	N/A	156

[0610] For all genes listed in Table 2 (namely, those relevant to CNV-subregions of interest), Table 3 represents a non-redundant list.

TABLE 4

A non-redundant list of transcript variants that correspond to the genes in Table 3				
RefSeq Gene Symbol	Exon overlap	RefSeq Accession Number	mRNA_Description	SEQ ID
MIR200B	exonic	NR_029639	<i>Homo sapiens</i> microRNA 200b (MIR200B), microRNA.	173
MIR200A	exonic	NR_029834	<i>Homo sapiens</i> microRNA 200a (MIR200A), microRNA.	174
MIR429	exonic	NR_029957	<i>Homo sapiens</i> microRNA 429 (MIR429), microRNA.	175
TTLL10	exonic	NM_001130045	<i>Homo sapiens</i> tubulin tyrosine ligase-like family, member 10 (TTLL10), transcript variant 1, mRNA.	176
TTLL10	exonic	NM_153254	<i>Homo sapiens</i> tubulin tyrosine ligase-like family, member 10 (TTLL10), transcript variant 2, mRNA.	177
TNFRSF18	exonic	NM_004195	<i>Homo sapiens</i> tumor necrosis factor receptor superfamily, member 18 (TNFRSF18), transcript variant 1, mRNA.	178
TNFRSF18	exonic	NM_148901	<i>Homo sapiens</i> tumor necrosis factor receptor superfamily, member 18 (TNFRSF18), transcript variant 2, mRNA.	179
TNFRSF18	exonic	NM_148902	<i>Homo sapiens</i> tumor necrosis factor receptor superfamily, member 18 (TNFRSF18), transcript variant 3, mRNA.	180
PIK3CD	exonic	NM_005026	<i>Homo sapiens</i> phosphatidylinositol-4,5-bisphosphate 3-kinase, catalytic subunit delta (PIK3CD), mRNA.	181
CAPZB	intronic	NR_038125	<i>Homo sapiens</i> capping protein (actin filament) muscle Z-line, beta (CAPZB), transcript variant 4, non-coding RNA.	182
CAPZB	intronic	NM_001206540	<i>Homo sapiens</i> capping protein (actin filament) muscle Z-line, beta (CAPZB), transcript variant 2, mRNA.	183
CAPZB	intronic	NM_004930	<i>Homo sapiens</i> capping protein (actin filament) muscle Z-line, beta (CAPZB), transcript variant 1, mRNA.	184
IFNLR1	exonic	NM_170743	<i>Homo sapiens</i> interferon, lambda receptor 1 (IFNLR1), transcript variant 1, mRNA.	185
IFNLR1	exonic	NM_173064	<i>Homo sapiens</i> interferon, lambda receptor 1 (IFNLR1), transcript variant 2, mRNA.	186
IFNLR1	exonic	NM_173065	<i>Homo sapiens</i> interferon, lambda receptor 1 (IFNLR1), transcript variant 3, mRNA.	187
PHACTR4	exonic	NM_001048183	<i>Homo sapiens</i> phosphatase and actin regulator 4 (PHACTR4), transcript variant 1, mRNA.	188
PHACTR4	exonic	NM_023923	<i>Homo sapiens</i> phosphatase and actin regulator 4 (PHACTR4), transcript variant 2, mRNA.	189
SNHG3	exonic	NR_002909	<i>Homo sapiens</i> small nucleolar RNA host gene 3 (non-protein coding) (SNHG3), transcript variant 2, non-coding RNA.	190
SNHG3	exonic	NR_036473	<i>Homo sapiens</i> small nucleolar RNA host gene 3 (non-protein coding) (SNHG3), transcript variant 1, non-coding RNA.	191
RCC1	exonic	NM_001048199	<i>Homo sapiens</i> regulator of chromosome condensation 1 (RCC1), transcript variant 4, mRNA.	192
RCC1	exonic	NR_030725	<i>Homo sapiens</i> regulator of chromosome condensation 1 (RCC1), transcript variant 5, non-coding RNA.	193
RCC1	exonic	NR_030726	<i>Homo sapiens</i> regulator of chromosome condensation 1 (RCC1), transcript variant 6, non-coding RNA.	194
RCC1	exonic	NM_001048194	<i>Homo sapiens</i> regulator of chromosome condensation 1 (RCC1), transcript variant 1, mRNA.	195
RCC1	exonic	NM_001048195	<i>Homo sapiens</i> regulator of chromosome condensation 1 (RCC1), transcript variant 2, mRNA.	196
RCC1	exonic	NM_001269	<i>Homo sapiens</i> regulator of chromosome condensation 1 (RCC1), transcript variant 3, mRNA.	197
AGBL4	intronic	NM_032785	<i>Homo sapiens</i> ATP/GTP binding protein-like 4 (AGBL4), mRNA.	198
GPATCH2	intronic	NM_018040	<i>Homo sapiens</i> G patch domain containing 2 (GPATCH2), mRNA.	199
RHOQ	intronic	NM_012249	<i>Homo sapiens</i> ras homolog family member Q (RHOQ), mRNA.	200
PNPT1	both	NM_033109	<i>Homo sapiens</i> polyribonucleotide nucleotidyltransferase 1 (PNPT1), mRNA.	201
MCEE	both	NM_032601	<i>Homo sapiens</i> methylmalonyl CoA epimerase (MCEE), mRNA.	202
HK2	exonic	NM_000189	<i>Homo sapiens</i> hexokinase 2 (HK2), mRNA.	203
FHL2	exonic	NM_201557	<i>Homo sapiens</i> four and a half LIM domains 2 (FHL2), transcript variant 4, mRNA.	204
FHL2	intronic	NM_001039492	<i>Homo sapiens</i> four and a half LIM domains 2 (FHL2), transcript variant 5, mRNA.	205
FHL2	intronic	NM_001450	<i>Homo sapiens</i> four and a half LIM domains 2 (FHL2), transcript variant 1, mRNA.	206
FHL2	intronic	NM_201555	<i>Homo sapiens</i> four and a half LIM domains 2 (FHL2), transcript variant 2, mRNA.	207
MIR 4267	exonic	NR_036225	<i>Homo sapiens</i> microRNA 4267 (MIR4267), microRNA.	208
MALL	exonic	NM_005434	<i>Homo sapiens</i> mal, T-cell differentiation protein-like (MALL), mRNA.	209
MIR 4436B1	exonic	NR_039941	<i>Homo sapiens</i> microRNA 4436b-1 (MIR4436B1), microRNA.	210
MIR4436B2	exonic	NR_049830	<i>Homo sapiens</i> microRNA 4436b-2 (MIR4436B2), microRNA.	211
MGAT5	intronic	NM_002410	<i>Homo sapiens</i> mannosyl (alpha-1,6-)-glycoprotein beta-1,6-N-acetyl-glucosaminyltransferase (MGAT5), mRNA.	212
BMPR2	intronic	NM_001204	<i>Homo sapiens</i> bone morphogenetic protein receptor, type II (serine/threonine kinase) (BMPR2), mRNA.	213
DNER	intronic	NM_139072	<i>Homo sapiens</i> delta/notch-like EGF repeat containing (DNER), mRNA.	214
IQCB1	exonic	NM_001023570	<i>Homo sapiens</i> IQ motif containing B1 (IQCB1), transcript variant 1, mRNA.	215
IQCB1	exonic	NM_001023571	<i>Homo sapiens</i> IQ motif containing B1 (IQCB1), transcript variant 3, mRNA.	216
STIM2	intronic	NM_001169117	<i>Homo sapiens</i> stromal interaction molecule 2 (STIM2), transcript variant 3, mRNA.	217
STIM2	intronic	NM_001169118	<i>Homo sapiens</i> stromal interaction molecule 2 (STIM2), transcript variant 1, mRNA.	218
STIM2	intronic	NM_020860	<i>Homo sapiens</i> stromal interaction molecule 2 (STIM2), transcript variant 2, mRNA.	219
PDGFRA	exonic	NM_006206	<i>Homo sapiens</i> platelet-derived growth factor receptor, alpha polypeptide (PDGFRA), mRNA.	220
HCN1	exonic	NM_021072	<i>Homo sapiens</i> hyperpolarization activated cyclic nucleotide-gated potassium channel 1 (HCN1), mRNA.	221
EMB	exonic	NM_198449	<i>Homo sapiens</i> embigin (EMB), mRNA.	222

TABLE 4-continued

A non-redundant list of transcript variants that correspond to the genes in Table 3				
RefSeq Gene Symbol	Exon overlap	RefSeq Accession Number	mRNA_Description	SEQ ID
EDIL3	intronic	NM_005711	<i>Homo sapiens</i> EGF-like repeats and discoidin I-like domains 3 (EDIL3), transcript variant 1, mRNA.	223
MAPK9	exonic	NM_002752	<i>Homo sapiens</i> mitogen-activated protein kinase 9 (MAPK9), transcript variant JNK2-a2, mRNA.	224
MAPK9	exonic	NM_139068	<i>Homo sapiens</i> mitogen-activated protein kinase 9 (MAPK9), transcript variant JNK2-a1, mRNA.	225
MAPK9	exonic	NM_139069	<i>Homo sapiens</i> mitogen-activated protein kinase 9 (MAPK9), transcript variant JNK2-b1, mRNA.	226
MAPK9	exonic	NM_139070	<i>Homo sapiens</i> mitogen-activated protein kinase 9 (MAPK9), transcript variant JNK2-b2, mRNA.	227
MAPK9	exonic	NM_001135044	<i>Homo sapiens</i> mitogen-activated protein kinase 9 (MAPK9), transcript variant JNK2-g, mRNA.	228
SERPINB6	exonic	NM_001271825	<i>Homo sapiens</i> serpin peptidase inhibitor, clade B (ovalbumin), member 6 (SERPINB6), transcript variant 6, mRNA.	229
SERPINB6	exonic	NM_001271823	<i>Homo sapiens</i> serpin peptidase inhibitor, clade B (ovalbumin), member 6 (SERPINB6), transcript variant 4, mRNA.	230
SERPINB6	exonic	NM_001271822	<i>Homo sapiens</i> serpin peptidase inhibitor, clade B (ovalbumin), member 6 (SERPINB6), transcript variant 3, mRNA.	231
SERPINB6	exonic	NM_001195291	<i>Homo sapiens</i> serpin peptidase inhibitor, clade B (ovalbumin), member 6 (SERPINB6), transcript variant 2, mRNA.	232
SERPINB6	exonic	NM_001271824	<i>Homo sapiens</i> serpin peptidase inhibitor, clade B (ovalbumin), member 6 (SERPINB6), transcript variant 5, mRNA.	233
SERPINB6	exonic	NM_004568	<i>Homo sapiens</i> serpin peptidase inhibitor, clade B (ovalbumin), member 6 (SERPINB6), transcript variant 1, mRNA.	234
DKFZP686I15217	exonic	NR_026855	<i>Homo sapiens</i> long intergenic non-protein coding RNA 1011 (LINC01011), transcript variant 1, non-coding RNA.	235
DKFZP686I15217	exonic	NR_026856	<i>Homo sapiens</i> long intergenic non-protein coding RNA 1011 (LINC01011), transcript variant 2, non-coding RNA.	236
NQO2	exonic	NM_000904	<i>Homo sapiens</i> NAD(P)H dehydrogenase, quinone 2 (NQO2), mRNA.	237
HTATSF1P2	exonic	NR_033884	<i>Homo sapiens</i> HIV-1 Tat specific factor 1 pseudogene 2 (HTATSF1P2), non-coding RNA.	238
PKHD1	intronic	NM_138694	<i>Homo sapiens</i> polycystic kidney and hepatic disease 1 (autosomal recessive) (PKHD1), transcript variant 1, mRNA.	239
PKHD1	intronic	NM_170724	<i>Homo sapiens</i> polycystic kidney and hepatic disease 1 (autosomal recessive) (PKHD1), transcript variant 2, mRNA.	240
SLC17A5	both	NM_012434	<i>Homo sapiens</i> solute carrier family 17 (acidic sugar transporter), member 5 (SLC17A5), mRNA.	241
PDSS2	exonic	NM_020381	<i>Homo sapiens</i> prenyl (decaprenyl) diphosphate synthase, subunit 2 (PDSS2), mRNA.	242
KCTD7	exonic	NM_001167961	<i>Homo sapiens</i> potassium channel tetramerization domain containing 7 (KCTD7), transcript variant 2, mRNA.	243
KCTD7	exonic	NM_153033	<i>Homo sapiens</i> potassium channel tetramerization domain containing 7 (KCTD7), transcript variant 1, mRNA.	244
PTPRN2	intronic	NM_002847	<i>Homo sapiens</i> protein tyrosine phosphatase, receptor type, N polypeptide 2 (PTPRN2), transcript variant 1, mRNA.	245
PTPRN2	intronic	NM_130842	<i>Homo sapiens</i> protein tyrosine phosphatase, receptor type, N polypeptide 2 (PTPRN2), transcript variant 2, mRNA.	246
PTPRN2	intronic	NM_130843	<i>Homo sapiens</i> protein tyrosine phosphatase, receptor type, N polypeptide 2 (PTPRN2), transcript variant 3, mRNA.	247
MIR595	exonic	NR_030325	<i>Homo sapiens</i> microRNA 595 (MIR595), microRNA.	248
TNFRSF10A	exonic	NM_003844	<i>Homo sapiens</i> tumor necrosis factor receptor superfamily, member 10a (TNFRSF10A), mRNA.	249
IDO2	intronic	NM_194294	<i>Homo sapiens</i> indoleamine 2,3-dioxygenase 2 (IDO2), mRNA.	250
STK3	intronic	NM_001256313	<i>Homo sapiens</i> serine/threonine kinase 3 (STK3), transcript variant 3, mRNA.	251
STK3	intronic	NM_006281	<i>Homo sapiens</i> serine/threonine kinase 3 (STK3), transcript variant 1, mRNA.	252
STK3	intronic	NM_001256312	<i>Homo sapiens</i> serine/threonine kinase 3 (STK3), transcript variant 2, mRNA.	253
KANK1	intronic	NM_001256876	<i>Homo sapiens</i> KN motif and ankyrin repeat domains 1 (KANK1), transcript variant 3, mRNA.	254
KANK1	intronic	NM_001256877	<i>Homo sapiens</i> KN motif and ankyrin repeat domains 1 (KANK1), transcript variant 4, mRNA.	255
KANK1	intronic	NM_015158	<i>Homo sapiens</i> KN motif and ankyrin repeat domains 1 (KANK1), transcript variant 1, mRNA.	256
KANK1	intronic	NM_153186	<i>Homo sapiens</i> KN motif and ankyrin repeat domains 1 (KANK1), transcript variant 2, mRNA.	257
GDA	exonic	NM_001242507	<i>Homo sapiens</i> guanine deaminase (GDA), transcript variant 4, mRNA.	258
GDA	exonic	NM_001242505	<i>Homo sapiens</i> guanine deaminase (GDA), transcript variant 1, mRNA.	259
GDA	exonic	NM_001242506	<i>Homo sapiens</i> guanine deaminase (GDA), transcript variant 3, mRNA.	260
GDA	exonic	NM_004293	<i>Homo sapiens</i> guanine deaminase (GDA), transcript variant 2, mRNA.	261
AUH	exonic	NM_001698	<i>Homo sapiens</i> AU RNA binding protein/enoyl-CoA hydratase (AUH), mRNA.	262
MIR3163	exonic	NR_036121	<i>Homo sapiens</i> microRNA 3163 (MIR3163), microRNA.	263
NFIL3	exonic	NM_005384	<i>Homo sapiens</i> nuclear factor, interleukin 3 regulated (NFIL3), mRNA.	264
MIR3910-1	exonic	NR_037472	<i>Homo sapiens</i> microRNA 3910-1 (MIR3910-1), microRNA.	265
MIR3910-2	exonic	NR_037489	<i>Homo sapiens</i> microRNA 3910-2 (MIR3910-2), microRNA.	266

TABLE 4-continued

A non-redundant list of transcript variants that correspond to the genes in Table 3				
RefSeq Gene Symbol	Exon overlap	RefSeq Accession Number	mRNA_Description	SEQ ID
ASTN2	intronic	NM_014010	<i>Homo sapiens</i> astrotactin 2 (ASTN2), transcript variant 1, mRNA.	267
ASTN2	intronic	NM_198186	<i>Homo sapiens</i> astrotactin 2 (ASTN2), transcript variant 2, mRNA.	268
ASTN2	intronic	NM_001184734	<i>Homo sapiens</i> astrotactin 2 (ASTN2), transcript variant 5, mRNA.	269
ASTN2	intronic	NM_198187	<i>Homo sapiens</i> astrotactin 2 (ASTN2), transcript variant 3, mRNA.	270
ASTN2	intronic	NM_198188	<i>Homo sapiens</i> astrotactin 2 (ASTN2), transcript variant 4, mRNA.	271
ASTN2	intronic	NM_001184735	<i>Homo sapiens</i> astrotactin 2 (ASTN2), transcript variant 6, mRNA.	272
LARP4B	exonic	NM_015155	<i>Homo sapiens</i> La ribonucleoprotein domain family, member 4B (LARP4B), mRNA.	273
GTPBP4	exonic	NM_012341	<i>Homo sapiens</i> GTP binding protein 4 (GTPBP4), mRNA.	274
IDI2	exonic	NM_033261	<i>Homo sapiens</i> isopentenyl-diphosphate delta isomerase 2 (IDI2), mRNA.	275
IDI2-AS1	exonic	NR_024628	<i>Homo sapiens</i> IDI2 antisense RNA 1 (IDI2-AS1), transcript variant 1, non-coding RNA.	276
IDI2-AS1	exonic	NR_024629	<i>Homo sapiens</i> IDI2 antisense RNA 1 (IDI2-AS1), transcript variant 2, non-coding RNA.	277
IDI2-AS1	exonic	NR_027708	<i>Homo sapiens</i> IDI2 antisense RNA 1 (IDI2-AS1), transcript variant 3, non-coding RNA.	278
IDI2-AS1	exonic	NR_027709	<i>Homo sapiens</i> IDI2 antisense RNA 1 (IDI2-AS1), transcript variant 4, non-coding RNA.	279
KAT6B	exonic	NM_001256468	<i>Homo sapiens</i> K(lysine) acetyltransferase 6B (KAT6B), transcript variant 2, mRNA.	280
KAT6B	exonic	NM_001256469	<i>Homo sapiens</i> K(lysine) acetyltransferase 6B (KAT6B), transcript variant 3, mRNA.	281
KAT6B	exonic	NM_012330	<i>Homo sapiens</i> K(lysine) acetyltransferase 6B (KAT6B), transcript variant 1, mRNA.	282
VWA2	exonic	NM_001272046	<i>Homo sapiens</i> von Willebrand factor A domain containing 2 (VWA2), mRNA.	283
PDE3B	intronic	NM_000922	<i>Homo sapiens</i> phosphodiesterase 3B, cGMP-inhibited (PDE3B), mRNA.	284
EHF	intronic	NM_001206615	<i>Homo sapiens</i> ets homologous factor (EHF), transcript variant 3, mRNA.	285
EHF	intronic	NM_012153	<i>Homo sapiens</i> ets homologous factor (EHF), transcript variant 2, mRNA.	286
EHF	exonic	NM_001206616	<i>Homo sapiens</i> ets homologous factor (EHF), transcript variant 1, mRNA.	287
SLC3A2	exonic	NM_001012662	<i>Homo sapiens</i> solute carrier family 3 (amino acid transporter heavy chain), member 2 (SLC3A2), transcript variant 2, mRNA.	288
SLC3A2	intronic	NM_001012664	<i>Homo sapiens</i> solute carrier family 3 (amino acid transporter heavy chain), member 2 (SLC3A2), transcript variant 5, mRNA.	289
SLC3A2	exonic	NM_002394	<i>Homo sapiens</i> solute carrier family 3 (amino acid transporter heavy chain), member 2 (SLC3A2), transcript variant 3, mRNA.	290
SLC3A2	intronic	NM_001013251	<i>Homo sapiens</i> solute carrier family 3 (amino acid transporter heavy chain), member 2 (SLC3A2), transcript variant 6, mRNA.	291
SLC3A2	intronic	NR_037193	<i>Homo sapiens</i> solute carrier family 3 (amino acid transporter heavy chain), member 2 (SLC3A2), transcript variant 7, non-coding RNA.	292
GDPD4	exonic	NM_182833	<i>Homo sapiens</i> glycerophosphodiester phosphodiesterase domain containing 4 (GDPD4), mRNA.	293
ETV6	exonic	NM_001987	<i>Homo sapiens</i> ets variant 6 (ETV6), mRNA.	294
LOH12CR1	exonic	NM_058169	<i>Homo sapiens</i> loss of heterozygosity, 12, chromosomal region 1 (LOH12CR1), mRNA.	295
DUSP16	exonic	NM_030640	<i>Homo sapiens</i> dual specificity phosphatase 16 (DUSP16), mRNA.	296
CREBL2	exonic	NM_001310	<i>Homo sapiens</i> cAMP responsive element binding protein-like 2 (CREBL2), mRNA.	297
GPR19	exonic	NM_006143	<i>Homo sapiens</i> G protein-coupled receptor 19 (GPR19), mRNA.	298
CDKN1B	exonic	NM_004064	<i>Homo sapiens</i> cyclin-dependent kinase inhibitor 1B (p27, Kip1) (CDKN1B), mRNA.	299
APOLD1	exonic	NM_001130415	<i>Homo sapiens</i> apolipoprotein L domain containing 1 (APOLD1), transcript variant 1, mRNA.	300
APOLD1	intronic	NM_030817	<i>Homo sapiens</i> apolipoprotein L domain containing 1 (APOLD1), transcript variant 2, mRNA.	301
EEA1	exonic	NM_003566	<i>Homo sapiens</i> early endosome antigen 1 (EEA1), mRNA.	302
LOC643339	exonic	NR_040096	<i>Homo sapiens</i> uncharacterized LOC643339 (LOC643339), non-coding RNA.	303
NUDT4	exonic	NM_019094	<i>Homo sapiens</i> nudix (nucleoside diphosphate linked moiety X)-type motif 4 (NUDT4), transcript variant 1, mRNA.	304
NUDT4	exonic	NM_199040	<i>Homo sapiens</i> nudix (nucleoside diphosphate linked moiety X)-type motif 4 (NUDT4), transcript variant 2, mRNA.	305
NUDT4P1	exonic	NR_002212	<i>Homo sapiens</i> nudix (nucleoside diphosphate linked moiety X)-type motif 4 pseudogene 1 (NUDT4P1), non-coding RNA.	306
UBE2N	exonic	NM_003348	<i>Homo sapiens</i> ubiquitin-conjugating enzyme E2N (UBE2N), mRNA.	307
MRPL42	exonic	NM_014050	<i>Homo sapiens</i> mitochondrial ribosomal protein L42 (MRPL42), transcript variant 1, mRNA.	308
MRPL42	exonic	NM_172177	<i>Homo sapiens</i> mitochondrial ribosomal protein L42 (MRPL42), transcript variant 2, mRNA.	309
MRPL42	exonic	NR_038159	<i>Homo sapiens</i> mitochondrial ribosomal protein L42 (MRPL42), transcript variant 3, non-coding RNA.	310
MRPL42	exonic	NR_038160	<i>Homo sapiens</i> mitochondrial ribosomal protein L42 (MRPL42), transcript variant 4, non-coding RNA.	311
MRPL42	exonic	NR_038161	<i>Homo sapiens</i> mitochondrial ribosomal protein L42 (MRPL42), transcript variant 5, non-coding RNA.	312
SOCS2-AS1	exonic	NR_038263	<i>Homo sapiens</i> SOCS2 antisense RNA 1 (SOCS2-AS1), non-coding RNA.	313
SOCS2	exonic	NM_003877	<i>Homo sapiens</i> suppressor of cytokine signaling 2 (SOCS2), transcript variant 1, mRNA.	314
SOCS2	exonic	NM_001270467	<i>Homo sapiens</i> suppressor of cytokine signaling 2 (SOCS2), transcript variant 2, mRNA.	315
SOCS2	exonic	NM_001270468	<i>Homo sapiens</i> suppressor of cytokine signaling 2 (SOCS2), transcript variant 3, mRNA.	316
SOCS2	exonic	NM_001270469	<i>Homo sapiens</i> suppressor of cytokine signaling 2 (SOCS2), transcript variant 4, mRNA.	317
SOCS2	exonic	NM_001270470	<i>Homo sapiens</i> suppressor of cytokine signaling 2 (SOCS2), transcript variant 5, mRNA.	318
SOCS2	exonic	NM_001270471	<i>Homo sapiens</i> suppressor of cytokine signaling 2 (SOCS2), transcript variant 6, mRNA.	319
CRADD	exonic	NM_003805	<i>Homo sapiens</i> CASP2 and RIPK1 domain containing adaptor with death domain (CRADD), mRNA.	320
PLXNC1	exonic	NM_005761	<i>Homo sapiens</i> plexin C1 (PLXNC1), transcript variant 1, mRNA.	321
PLXNC1	exonic	NR_037687	<i>Homo sapiens</i> plexin C1 (PLXNC1), transcript variant 2, non-coding RNA.	322
CCDC41	exonic	NM_001042399	<i>Homo sapiens</i> coiled-coil domain containing 41 (CCDC41), transcript variant 2, mRNA.	323
CCDC41	exonic	NM_016122	<i>Homo sapiens</i> coiled-coil domain containing 41 (CCDC41), transcript variant 1, mRNA.	324

TABLE 4-continued

A non-redundant list of transcript variants that correspond to the genes in Table 3				
RefSeq Gene Symbol	Exon overlap	RefSeq Accession Number	mRNA_Description	SEQ ID
TRAFD1	exonic	NM_001143906	<i>Homo sapiens</i> TRAF-type zinc finger domain containing 1 (TRAFD1), transcript variant 1, mRNA.	325
TRAFD1	exonic	NM_006700	<i>Homo sapiens</i> TRAF-type zinc finger domain containing 1 (TRAFD1), transcript variant 2, mRNA.	326
RGCC	exonic	NM_014059	<i>Homo sapiens</i> regulator of cell cycle (RGCC), mRNA.	327
COMMD6	exonic	NM_203495	<i>Homo sapiens</i> COMM domain containing 6 (COMMD6), transcript variant 2, mRNA.	328
COMMD6	exonic	NM_203497	<i>Homo sapiens</i> COMM domain containing 6 (COMMD6), transcript variant 1, mRNA.	329
GPC5	intronic	NM_004466	<i>Homo sapiens</i> glypican 5 (GPC5), mRNA.	330
ARHGEF7	exonic	NM_003899	<i>Homo sapiens</i> Rho guanine nucleotide exchange factor (GEF) 7 (ARHGEF7), transcript variant 1, mRNA.	331
ARHGEF7	exonic	NM_001113513	<i>Homo sapiens</i> Rho guanine nucleotide exchange factor (GEF) 7 (ARHGEF7), transcript variant 5, mRNA.	332
TEX29	exonic	NM_152324	<i>Homo sapiens</i> testis expressed 29 (TEX29), mRNA.	333
ARHGEF7	intronic	NM_001113511	<i>Homo sapiens</i> Rho guanine nucleotide exchange factor (GEF) 7 (ARHGEF7), transcript variant 3, mRNA.	334
ARHGEF7	intronic	NM_001113512	<i>Homo sapiens</i> Rho guanine nucleotide exchange factor (GEF) 7 (ARHGEF7), transcript variant 4, mRNA.	335
ARHGEF7	intronic	NM_145735	<i>Homo sapiens</i> Rho guanine nucleotide exchange factor (GEF) 7 (ARHGEF7), transcript variant 2, mRNA.	336
RNASE10	exonic	NM_001012975	<i>Homo sapiens</i> ribonuclease, RNase A family, 10 (non-active) (RNASE10), mRNA.	337
RNASE3	exonic	NM_002935	<i>Homo sapiens</i> ribonuclease, RNase A family, 3 (RNASE3), mRNA.	338
ECRP	exonic	NR_033909	<i>Homo sapiens</i> ribonuclease, RNase A family, 2 (liver, eosinophil-derived neurotoxin) pseudogene (ECRP), non-coding RNA.	339
PRKCH	intronic	NM_006255	<i>Homo sapiens</i> protein kinase C, eta (PRKCH), mRNA.	340
MTHFD1	exonic	NM_005956	<i>Homo sapiens</i> methyltetrahydrofolate dehydrogenase (NADP+ dependent) 1, methenyltetrahydrofolate cyclohydrolase, formyltetrahydrofolate synthetase (MTHFD1), mRNA.	341
BDKRB2	intronic	NM_000623	<i>Homo sapiens</i> bradykinin receptor B2 (BDKRB2), mRNA.	342
HEXA	exonic	NM_000520	<i>Homo sapiens</i> hexosaminidase A (alpha polypeptide) (HEXA), mRNA.	343
PSTPIP1	exonic	NM_003978	<i>Homo sapiens</i> proline-serine-threonine phosphatase interacting protein 1 (PSTPIP1), mRNA.	344
RBFOX1	intronic	NM_001142333	<i>Homo sapiens</i> RNA binding protein, fox-1 homolog (<i>C. elegans</i>) 1 (RBFOX1), transcript variant 5, mRNA.	345
RBFOX1	intronic	NM_018723	<i>Homo sapiens</i> RNA binding protein, fox-1 homolog (<i>C. elegans</i>) 1 (RBFOX1), transcript variant 4, mRNA.	346
RBFOX1	intronic	NM_001142334	<i>Homo sapiens</i> RNA binding protein, fox-1 homolog (<i>C. elegans</i>) 1 (RBFOX1), transcript variant 6, mRNA.	347
RBFOX1	intronic	NM_145891	<i>Homo sapiens</i> RNA binding protein, fox-1 homolog (<i>C. elegans</i>) 1 (RBFOX1), transcript variant 1, mRNA.	348
RBFOX1	intronic	NM_145892	<i>Homo sapiens</i> RNA binding protein, fox-1 homolog (<i>C. elegans</i>) 1 (RBFOX1), transcript variant 2, mRNA.	349
RBFOX1	intronic	NM_145893	<i>Homo sapiens</i> RNA binding protein, fox-1 homolog (<i>C. elegans</i>) 1 (RBFOX1), transcript variant 3, mRNA.	350
PRKCB	both	NM_002738	<i>Homo sapiens</i> protein kinase C, beta (PRKCB), transcript variant 2, mRNA.	351
PRKCB	both	NM_212535	<i>Homo sapiens</i> protein kinase C, beta (PRKCB), transcript variant 1, mRNA.	352
FUK	both	NM_145059	<i>Homo sapiens</i> fucokinase (FUK), mRNA.	353
COG4	exonic	NM_001195139	<i>Homo sapiens</i> component of oligomeric golgi complex 4 (COG4), transcript variant 2, mRNA.	354
COG4	exonic	NM_015386	<i>Homo sapiens</i> component of oligomeric golgi complex 4 (COG4), transcript variant 1, mRNA.	355
HPR	exonic	NM_020995	<i>Homo sapiens</i> haptoglobin-related protein (HPR), mRNA.	356
RPL38	exonic	NM_000999	<i>Homo sapiens</i> ribosomal protein L38 (RPL38), transcript variant 1, mRNA.	357
RPL38	exonic	NM_001035258	<i>Homo sapiens</i> ribosomal protein L38 (RPL38), transcript variant 2, mRNA.	358
MGC16275	exonic	NR_026914	<i>Homo sapiens</i> uncharacterized protein MGC16275 (MGC16275), non-coding RNA.	359
TTYH2	exonic	NM_032646	<i>Homo sapiens</i> tweety family member 2 (TTYH2), transcript variant 1, mRNA.	360
TTYH2	exonic	NM_052869	<i>Homo sapiens</i> tweety family member 2 (TTYH2), transcript variant 2, mRNA.	361
DNAI2	exonic	NM_001172810	<i>Homo sapiens</i> dynein, axonemal, intermediate chain 2 (DNAI2), transcript variant 2, mRNA.	362
DNAI2	exonic	NM_023036	<i>Homo sapiens</i> dynein, axonemal, intermediate chain 2 (DNAI2), transcript variant 1, mRNA.	363
KIF19	exonic	NM_153209	<i>Homo sapiens</i> kinesin family member 19 (KIF19), mRNA.	364
BTBD17	exonic	NM_001080466	<i>Homo sapiens</i> BTB (POZ) domain containing 17 (BTBD17), mRNA.	365
GPR142	exonic	NM_181790	<i>Homo sapiens</i> G protein-coupled receptor 142 (GPR142), mRNA.	366
GPRC5C	exonic	NM_022036	<i>Homo sapiens</i> G protein-coupled receptor, family C, group 5, member C (GPRC5C), transcript variant 1, mRNA.	367
GPRC5C	exonic	NM_018653	<i>Homo sapiens</i> G protein-coupled receptor, family C, group 5, member C (GPRC5C), transcript variant 2, mRNA.	368
CD300A	exonic	NM_001256841	<i>Homo sapiens</i> CD300a molecule (CD300A), transcript variant 2, mRNA.	369
CD300A	exonic	NM_007261	<i>Homo sapiens</i> CD300a molecule (CD300A), transcript variant 1, mRNA.	370
CD300LB	exonic	NM_174892	<i>Homo sapiens</i> CD300 molecule-like family member b (CD300LB), mRNA.	371
CD300C	exonic	NM_006678	<i>Homo sapiens</i> CD300c molecule (CD300C), mRNA.	372
CD300LD	exonic	NM_001115152	<i>Homo sapiens</i> CD300 molecule-like family member d (CD300LD), mRNA.	373
C17orf77	exonic	NM_152460	<i>Homo sapiens</i> chromosome 17 open reading frame 77 (C17orf77), mRNA.	374
CD300E	exonic	NM_181449	<i>Homo sapiens</i> CD300e molecule (CD300E), mRNA.	375

TABLE 4-continued

A non-redundant list of transcript variants that correspond to the genes in Table 3				
RefSeq Gene Symbol	Exon overlap	RefSeq Accession Number	mRNA_Description	SEQ ID
RAB37	exonic	NM_175738	<i>Homo sapiens</i> RAB37, member RAS oncogene family (RAB37), transcript variant 3, mRNA.	376
CD300LF	exonic	NM_139018	<i>Homo sapiens</i> CD300 molecule-like family member f (CD300LF), mRNA.	377
RAB37	intronic	NM_001163989	<i>Homo sapiens</i> RAB37, member RAS oncogene family (RAB37), transcript variant 4, mRNA.	378
RAB37	intronic	NM_001006638	<i>Homo sapiens</i> RAB37, member RAS oncogene family (RAB37), transcript variant 2, mRNA.	379
RAB37	intronic	NM_001163990	<i>Homo sapiens</i> RAB37, member RAS oncogene family (RAB37), transcript variant 5, mRNA.	380
TBC1D16	intronic	NM_019020	<i>Homo sapiens</i> TBC1 domain family, member 16 (TBC1D16), transcript variant 1, mRNA.	381
TBC1D16	intronic	NM_001271844	<i>Homo sapiens</i> TBC1 domain family, member 16 (TBC1D16), transcript variant 2, mRNA.	382
TBC1D16	intronic	NM_001271845	<i>Homo sapiens</i> TBC1 domain family, member 16 (TBC1D16), transcript variant 3, mRNA.	383
TBC1D16	intronic	NM_001271846	<i>Homo sapiens</i> TBC1 domain family, member 16 (TBC1D16), transcript variant 4, mRNA.	384
RPTOR	intronic	NM_001163034	<i>Homo sapiens</i> regulatory associated protein of MTOR, complex 1 (RPTOR), transcript variant 2, mRNA.	385
RPTOR	intronic	NM_020761	<i>Homo sapiens</i> regulatory associated protein of MTOR, complex 1 (RPTOR), transcript variant 1, mRNA.	386
PTPN2	exonic	NM_001207013	<i>Homo sapiens</i> protein tyrosine phosphatase, non-receptor type 2 (PTPN2), transcript variant 4, mRNA.	387
PTPN2	exonic	NM_080422	<i>Homo sapiens</i> protein tyrosine phosphatase, non-receptor type 2 (PTPN2), transcript variant 2, mRNA.	388
PTPN2	exonic	NM_080423	<i>Homo sapiens</i> protein tyrosine phosphatase, non-receptor type 2 (PTPN2), transcript variant 3, mRNA.	389
PTPN2	intronic	NM_002828	<i>Homo sapiens</i> protein tyrosine phosphatase, non-receptor type 2 (PTPN2), transcript variant 1, mRNA.	390
ST8SIA5	exonic	NM_013305	<i>Homo sapiens</i> ST8 alpha-N-acetyl-neuraminide alpha-2,8-sialyltransferase 5 (ST8SIA5), mRNA.	391
PIAS2	exonic	NM_004671	<i>Homo sapiens</i> protein inhibitor of activated STAT, 2 (PIAS2), transcript variant beta, mRNA.	392
PIAS2	exonic	NM_173206	<i>Homo sapiens</i> protein inhibitor of activated STAT, 2 (PIAS2), transcript variant alpha, mRNA.	393
SERPINB4	exonic	NM_002974	<i>Homo sapiens</i> serpin peptidase inhibitor, clade B (ovalbumin), member 4 (SERPINB4), mRNA.	394
FLJ26850	intronic	NR_027257	<i>Homo sapiens</i> FLJ26850 protein (FLJ26850), non-coding RNA.	395
FPR2	exonic	NM_001005738	<i>Homo sapiens</i> formyl peptide receptor 2 (FPR2), transcript variant 2, mRNA.	396
FPR2	exonic	NM_001462	<i>Homo sapiens</i> formyl peptide receptor 2 (FPR2), transcript variant 1, mRNA.	397
FPR3	exonic	NM_002030	<i>Homo sapiens</i> formyl peptide receptor 3 (FPR3), mRNA.	398
ZNF577	exonic	NR_024181	<i>Homo sapiens</i> zinc finger protein 577 (ZNF577), transcript variant 3, non-coding RNA.	399
ZNF577	exonic	NM_001135590	<i>Homo sapiens</i> zinc finger protein 577 (ZNF577), transcript variant 2, mRNA.	400
ZNF577	exonic	NM_032679	<i>Homo sapiens</i> zinc finger protein 577 (ZNF577), transcript variant 1, mRNA.	401
ZNF649	exonic	NM_023074	<i>Homo sapiens</i> zinc finger protein 649 (ZNF649), mRNA.	402
ZNF613	exonic	NM_001031721	<i>Homo sapiens</i> zinc finger protein 613 (ZNF613), transcript variant 1, mRNA.	403
ZNF613	exonic	NM_024840	<i>Homo sapiens</i> zinc finger protein 613 (ZNF613), transcript variant 2, mRNA.	404
ZNF350	exonic	NM_021632	<i>Homo sapiens</i> zinc finger protein 350 (ZNF350), mRNA.	405
ZNF615	exonic	NM_001199324	<i>Homo sapiens</i> zinc finger protein 615 (ZNF615), transcript variant 1, mRNA.	406
ZNF615	exonic	NM_198480	<i>Homo sapiens</i> zinc finger protein 615 (ZNF615), transcript variant 2, mRNA.	407
ZNF614	exonic	NM_025040	<i>Homo sapiens</i> zinc finger protein 614 (ZNF614), mRNA.	408
ZNF432	exonic	NM_014650	<i>Homo sapiens</i> zinc finger protein 432 (ZNF432), mRNA.	409
ZNF841	exonic	NM_001136499	<i>Homo sapiens</i> zinc finger protein 841 (ZNF841), mRNA.	410
NLRP12	exonic	NM_001277126	<i>Homo sapiens</i> NLR family, pyrin domain containing 12 (NLRP12), transcript variant 3, mRNA.	411
NLRP12	exonic	NM_001277129	<i>Homo sapiens</i> NLR family, pyrin domain containing 12 (NLRP12), transcript variant 4, mRNA.	412
NLRP12	exonic	NM_144687	<i>Homo sapiens</i> NLR family, pyrin domain containing 12 (NLRP12), transcript variant 2, mRNA.	413
VSTM1	intronic	NM_198481	<i>Homo sapiens</i> V-set and transmembrane domain containing 1 (VSTM1), mRNA.	414
SNX5	exonic	NM_014426	<i>Homo sapiens</i> sorting nexin 5 (SNX5), transcript variant 2, mRNA.	415
SNX5	exonic	NM_152227	<i>Homo sapiens</i> sorting nexin 5 (SNX5), transcript variant 1, mRNA.	416
SNORD17	exonic	NR_003045	<i>Homo sapiens</i> small nucleolar RNA, C/D box 17 (SNORD17), small nucleolar RNA.	417
MGME1	exonic	NM_052865	<i>Homo sapiens</i> mitochondrial genome maintenance exonuclease 1 (MGME1), mRNA.	418
OVOL2	exonic	NM_021220	<i>Homo sapiens</i> ovo-like 2 (<i>Drosophila</i>) (OVOL2), mRNA.	419
ADA	intronic	NM_000022	<i>Homo sapiens</i> adenosine deaminase (ADA), mRNA.	420
NRIP1	exonic	NM_003489	<i>Homo sapiens</i> nuclear receptor interacting protein 1 (NRIP1), mRNA.	421
BACH1	exonic	NR_027655	<i>Homo sapiens</i> BTB and CNC homology 1, basic leucine zipper transcription factor 1 (BACH1), transcript variant 3, non-coding RNA.	422
BACH1	intronic	NM_001186	<i>Homo sapiens</i> BTB and CNC homology 1, basic leucine zipper transcription factor 1 (BACH1), transcript variant 2, mRNA.	423
BACH1	intronic	NM_206866	<i>Homo sapiens</i> BTB and CNC homology 1, basic leucine zipper transcription factor 1 (BACH1), transcript variant 1, mRNA.	424
TRPM2	exonic	NM_003307	<i>Homo sapiens</i> transient receptor potential cation channel, subfamily M, member 2 (TRPM2), transcript variant 1, mRNA.	425
TRPM2	exonic	NR_038257	<i>Homo sapiens</i> transient receptor potential cation channel, subfamily M, member 2 (TRPM2), transcript variant 2, non-coding RNA.	426
ADARB1	intronic	NM_001112	<i>Homo sapiens</i> adenosine deaminase, RNA-specific, B1 (ADARB1), transcript variant 1, mRNA.	427

TABLE 4-continued

A non-redundant list of transcript variants that correspond to the genes in Table 3				
RefSeq Gene Symbol	Exon overlap	RefSeq Accession Number	mRNA_Description	SEQ ID
ADARB1	intronic	NM_001160230	<i>Homo sapiens</i> adenosine deaminase, RNA-specific, B1 (ADARB1), transcript variant 7, mRNA.	428
ADARB1	intronic	NM_015833	<i>Homo sapiens</i> adenosine deaminase, RNA-specific, B1 (ADARB1), transcript variant 2, mRNA.	429
ADARB1	intronic	NM_015834	<i>Homo sapiens</i> adenosine deaminase, RNA-specific, B1 (ADARB1), transcript variant 3, mRNA.	430
ADARB1	intronic	NR_027672	<i>Homo sapiens</i> adenosine deaminase, RNA-specific, B1 (ADARB1), transcript variant 5, non-coding RNA.	431
ADARB1	intronic	NR_027673	<i>Homo sapiens</i> adenosine deaminase, RNA-specific, B1 (ADARB1), transcript variant 4, non-coding RNA.	432
ADARB1	intronic	NR_027674	<i>Homo sapiens</i> adenosine deaminase, RNA-specific, B1 (ADARB1), transcript variant 6, non-coding RNA.	433
ADARB1	intronic	NR_073200	<i>Homo sapiens</i> adenosine deaminase, RNA-specific, B1 (ADARB1), transcript variant 8, non-coding RNA.	434
APOBEC3A	exonic	NM_001270406	<i>Homo sapiens</i> apolipoprotein B mRNA editing enzyme, catalytic polypeptide-like 3A (APOBEC3A), transcript variant 3, mRNA.	435
APOBEC3A	exonic	NM_145699	<i>Homo sapiens</i> apolipoprotein B mRNA editing enzyme, catalytic polypeptide-like 3A (APOBEC3A), transcript variant 1, mRNA.	436
APOBEC3A_B	intronic	NM_001193289	<i>Homo sapiens</i> APOBEC3A and APOBEC3B deletion hybrid (APOBEC3A_B), mRNA.	437
APOBEC3B	exonic	NM_001270411	<i>Homo sapiens</i> apolipoprotein B mRNA editing enzyme, catalytic polypeptide-like 3B (APOBEC3B), transcript variant 2, mRNA.	438
APOBEC3B	exonic	NM_004900	<i>Homo sapiens</i> apolipoprotein B mRNA editing enzyme, catalytic polypeptide-like 3B (APOBEC3B), transcript variant 1, mRNA.	439
MKL1	intronic	NM_020831	<i>Homo sapiens</i> megakaryoblastic leukemia (translocation) 1 (MKL1), mRNA.	440
TNFRSF13C	lexonic	NM_052945	<i>Homo sapiens</i> tumor necrosis factor receptor superfamily, member 13C (TNFRSF13C), mRNA.	441
CENPM	exonic	NM_001110215	<i>Homo sapiens</i> centromere protein M (CENPM), transcript variant 3, mRNA.	442
CENPM	exonic	NM_001002876	<i>Homo sapiens</i> centromere protein M (CENPM), transcript variant 2, mRNA.	443
CENPM	exonic	NM_024053	<i>Homo sapiens</i> centromere protein M (CENPM), transcript variant 1, mRNA.	444
PPP2R3B	intronic	NM_013239	<i>Homo sapiens</i> protein phosphatase 2, regulatory subunit B", beta (PPP2R3B), mRNA.	445
VCX	exonic	NM_013452	<i>Homo sapiens</i> variable charge, X-linked (VCX), mRNA.	446
PNPLA4	exonic	NM_004650	<i>Homo sapiens</i> patatin-like phospholipase domain containing 4 (PNPLA4), transcript variant 1, mRNA.	447
PNPLA4	exonic	NM_001142389	<i>Homo sapiens</i> patatin-like phospholipase domain containing 4 (PNPLA4), transcript variant 2, mRNA.	448
PNPLA4	exonic	NM_001172672	<i>Homo sapiens</i> patatin-like phospholipase domain containing 4 (PNPLA4), transcript variant 3, mRNA.	449
MIR651	exonic	NR_030380	<i>Homo sapiens</i> microRNA 651 (MIR651), microRNA.	450
JPX	intronic	NR_024582	<i>Homo sapiens</i> JPX transcript, XIST activator (non-protein coding) (JPX), non-coding RNA.	451
GRIA3	intronic	NM_000828	<i>Homo sapiens</i> glutamate receptor, ionotropic, AMPA 3 (GRIA3), transcript variant 2, mRNA.	452
GRIA3	intronic	NM_007325	<i>Homo sapiens</i> glutamate receptor, ionotropic, AMPA 3 (GRIA3), transcript variant 1, mRNA.	453
GRIA3	intronic	NM_001256743	<i>Homo sapiens</i> glutamate receptor, ionotropic, AMPA 3 (GRIA3), transcript variant 3, mRNA.	454
HMGB3	exonic	NM_005342	<i>Homo sapiens</i> high mobility group box 3 (HMGB3), mRNA.	455

[0611] For all genes listed in Table 2 (namely, those relevant to CNV-subregions of interest), Table 4 represents a non-redundant list.

TABLE 5

The set of SNVs reported in Tables 7-10, 14, or 15 that were found in the 70 PML cases in this study for which WES data were generated.				
Chromosome	Position_hg19	REF	ALT	SEQ_ID
1	9777599	C	G	1000
1	12172008	T	C	1001
1	24486004	G	T	1002
1	33476435	C	A	1003
1	33478900	T	A	1004
1	33487007	C	T	1005
1	36932047	C	T	1006
1	36933715	A	G	1007
1	42047208	C	G	1008

TABLE 5-continued

The set of SNVs reported in Tables 7-10, 14, or 15 that were found in the 70 PML cases in this study for which WES data were generated.					
Chromosome	Position_hg19	REF	ALT	SEQ_ID	
1	59248085	G	C	1009	
1	59248339	T	C	1010	
1	92941660	C	T	1011	
1	92946625	G	C	1012	
1	92946625	G	C	1013	
1	150053494	C	T	1014	
1	155317682	C	T	1015	
1	155449630	T	G	1016	
1	155450331	C	T	1017	
1	182554557	C	T	1018	
1	198717250	T	G	1019	
1	198717272	A	T	1020	
1	206945738	C	T	1021	

TABLE 5-continued

The set of SNVs reported in Tables 7-10, 14, or 15 that were found in the 70 PML cases in this study for which WES data were generated.				
Chromosome	Position_hg19	REF	ALT	SEQ_ID
1	207641950	C	T	1022
1	235840495	G	T	1023
1	235897907	C	T	1024
1	235909815	A	T	1025
2	24431184	C	T	1026
2	24432937	C	T	1027
2	24435599	G	A	1028
2	47205921	C	T	1029
2	47273468	A	G	1030
2	47277182	T	C	1031
2	55910961	T	C	1032
2	71337203	C	T	1033
2	98351032	C	G	1034
2	98351066	C	T	1035
2	98351081	C	T	1036
2	113589000	C	T	1037
2	163124051	C	T	1038
2	163133194	T	C	1039
2	163134203	G	T	1040
2	163136505	C	G	1041
2	163139025	C	T	1042
2	163139085	A	T	1043
2	163144899	G	A	1044
2	163174327	C	A	1045
2	163174328	T	G	1046
2	219942026	T	A	1047
2	220023045	C	T	1048
2	230231632	C	T	1049
2	230450646	T	A	1050
3	38181899	G	T	1051
3	39323163	A	C	1052
3	53213691	G	C	1053
3	53221390	T	C	1054
3	121415370	T	C	1055
3	128204761	C	T	1056
3	128205808	C	T	1057
3	142272098	A	G	1058
3	142274880	G	C	1059
3	142281353	C	G	1060
3	142286928	C	T	1061
3	196199032	A	T	1062
3	196199204	G	T	1063
3	196210704	G	A	1064
3	196210764	T	C	1065
3	196214320	C	T	1066
4	27019452	C	T	1067
4	27024170	A	G	1068
4	103522068	A	G	1069
4	103522150	G	A	1070
4	103528328	C	T	1071
4	151199080	G	A	1072
4	151520216	G	A	1073
4	187003729	C	G	1074
4	187004074	C	T	1075
4	187005854	A	C	1076
5	67591018	A	C	1077
5	77311370	C	T	1078
5	77311370	C	T	1079
5	77334964	T	C	1080
5	77334964	T	C	1081
5	77335015	G	T	1082
5	77335015	G	T	1083
5	77437092	G	C	1084
5	77437092	G	C	1085
5	78596018	G	C	1086
5	138856923	C	T	1087
5	156593120	C	T	1088
5	169081453	G	C	1089
6	3077139	T	C	1090
6	12121113	C	T	1091
6	12122102	T	G	1092

TABLE 5-continued

The set of SNVs reported in Tables 7-10, 14, or 15 that were found in the 70 PML cases in this study for which WES data were generated.				
Chromosome	Position_hg19	REF	ALT	SEQ_ID
6	12123538	G	T	1093
6	12124215	C	T	1094
6	12125232	C	T	1095
6	12162068	C	T	1096
6	12163657	C	T	1097
6	31928306	A	G	1098
6	31935750	G	A	1099
6	31936679	C	T	1100
6	32797809	C	T	1101
6	32810794	T	A	1102
6	32811752	C	T	1103
6	51483961	T	C	1104
6	51484077	G	C	1105
6	51491885	G	A	1106
6	51497503	C	A	1107
6	51524339	C	G	1108
6	51524409	G	T	1109
6	51612746	G	A	1110
6	51712759	T	C	1111
6	51747943	T	A	1112
6	51798908	C	T	1113
6	52101833	C	T	1114
6	83884161	C	G	1115
6	143081232	T	C	1116
6	143092151	T	C	1117
6	143092673	G	A	1118
6	144508353	G	A	1119
6	144508563	G	A	1120
7	2959240	C	T	1121
7	2962933	C	T	1122
7	2983958	T	C	1123
8	39840234	A	G	1124
8	39862881	C	T	1125
8	39862893	T	A	1126
8	42176189	G	A	1127
8	48690299	A	G	1128
8	48773526	T	C	1129
8	48798507	T	C	1130
8	48826575	C	G	1131
8	61654298	T	A	1132
8	61732632	A	G	1133
8	61757805	C	T	1134
8	61769428	A	G	1135
8	61777914	C	G	1136
8	61777922	C	G	1137
8	90990521	T	C	1138
8	100205255	G	A	1139
8	100791158	G	A	1140
8	100865941	G	A	1141
8	145154222	G	A	1142
8	145154222	G	A	1143
8	145154257	C	G	1144
8	145154824	A	C	1145
9	286491	G	A	1146
9	286593	C	A	1147
9	304628	G	A	1148
9	312134	G	A	1149
9	328047	T	A	1150
9	334277	G	A	1151
9	368128	C	T	1152
9	399233	A	G	1153
9	446401	A	G	1154
9	711359	C	T	1155
9	713132	G	T	1156
9	32526077	C	T	1157
9	32526077	C	T	1158
9	120466814	A	G	1159
9	120475302	A	G	1160
9	120475602	C	T	1161
9	120476568	A	G	1162
9	120476816	C	T	1163

TABLE 5-continued

The set of SNVs reported in Tables 7-10, 14, or 15 that were found in the 70 PML cases in this study for which WES data were generated.				
Chromosome	Position_hg19	REF	ALT	SEQ_ID
10	1060218	G	A	1164
10	14974905	T	C	1165
10	14976727	G	C	1166
10	14977469	C	A, T	1167
10	72358167	G	A	1168
10	76602923	G	T	1169
10	76748831	C	T	1170
10	89720659	G	T	1171
10	90771767	G	A	1172
10	116045796	G	A	1173
11	4104626	C	A	1174
11	4112582	C	T	1175
11	9598696	G	A	1176
11	9608330	G	A	1177
11	36595321	C	T	1178
11	36596528	G	C	1179
11	36596863	C	T	1180
11	36597513	G	A	1181
11	36614561	G	T	1182
11	36615033	C	T	1183
11	67814983	G	A	1184
11	67818269	G	A	1185
11	76954833	G	A	1186
11	76979511	A	G	1187
11	108117787	C	T	1188
11	108119823	T	C	1189
11	108123551	C	T	1190
11	108138003	T	C	1191
11	108143456	C	G	1192
11	108175462	G	A	1193
11	108181014	A	G	1194
11	108186610	G	A	1195
11	108186631	A	G	1196
11	108198384	C	G	1197
11	108202772	G	T	1198
12	12673965	G	A	1199
12	12870798	G	A	1200
12	44166753	A	G	1201
12	44167821	A	T	1202
12	64878241	G	A	1203
12	64879775	C	T	1204
12	88900891	C	A	1205
12	93196332	C	T	1206
12	93205148	T	G	1207
12	112583447	A	C	1208
12	122064788	G	GT	1209
12	133201381	T	A	1210
12	133202816	C	T	1211
12	133209020	G	C	1212
12	133220526	T	C	1213
12	133220544	C	T	1214
12	133237658	T	G	1215
12	133245026	G	A	1216
12	133252406	C	A	1217
12	133253971	C	T	1218
12	133253995	G	A	1219
13	47466549	T	C	1220
13	108861092	G	T	1221
13	108863591	G	A	1222
14	21992397	T	C	1223
14	21993359	G	A	1224
14	22004996	G	T	1225
14	24805463	G	T	1226
14	24806303	G	A	1227
14	61924007	C	G	1228
14	103369593	G	A	1229
15	41011016	G	A	1230
15	68378781	A	C	1231
15	77329479	C	T	1232
15	91306241	G	A	1233
15	91310209	A	G	1234

TABLE 5-continued

The set of SNVs reported in Tables 7-10, 14, or 15 that were found in the 70 PML cases in this study for which WES data were generated.				
Chromosome	Position_hg19	REF	ALT	SEQ_ID
15	91326099	C	T	1235
15	91328219	G	T	1236
15	91328310	A	G	1237
15	91341543	A	C	1238
16	1498408	G	A	1239
16	1510535	C	T	1240
16	1524855	C	G	1241
16	7568296	C	T	1242
16	7703891	A	G	1243
16	7714909	C	T	1244
16	7759119	G	A	1245
16	7759496	C	T	1246
16	24124365	A	G	1247
16	27460020	G	A	1248
16	30133233	T	C	1249
16	30134529	A	C	1250
16	50733536	T	C	1251
16	50741791	C	T	1252
16	50741791	C	T	1253
16	50744688	A	G	1254
16	50745021	C	T	1255
16	50753867	G	T	1256
16	70503095	A	G	1257
16	81819605	C	T	1258
16	81902826	C	T	1259
16	81904539	C	T	1260
16	81939089	T	C	1261
16	81942028	C	G	1262
16	81942175	A	G	1263
16	81946278	A	G	1264
16	81960772	C	A	1265
17	7577069	C	T	1266
17	16852187	A	G	1267
17	77926526	C	T	1268
18	43445580	C	T	1269
18	43445601	T	G	1270
18	43456296	C	T	1271
18	43458306	G	A	1272
18	43460105	C	A	1273
18	43464763	C	T	1274
18	43479473	T	C	1275
18	43488030	T	C	1276
18	43496370	G	A	1277
18	43496539	G	A	1278
18	43497710	A	G	1279
18	43523240	C	T	1280
18	43529551	C	T	1281
18	43531186	C	T	1282
18	44392443	T	C	1283
18	48584504	C	T	1284
18	56401523	C	T	1285
18	60036429	G	A	1286
18	60052034	A	C	1287
19	4817657	C	T	1288
19	4817852	G	A	1289
19	7705818	C	T	1290
19	7712287	G	C	1291
19	48631258	G	A	1292
19	48639022	T	C	1293
20	3843027	C	A	1294
20	3846397	C	T	1295
20	31383307	G	A	1296
20	31384614	G	T	1297
20	62305450	C	T	1298
20	62309621	T	C	1299
20	62326964	C	G	1300
21	16338814	T	C	1301
21	16339852	T	C	1302
21	30698953	T	G	1303
21	34809232	C	T	1304
21	45786650	C	T	1305

TABLE 5-continued

The set of SNVs reported in Tables 7-10, 14, or 15 that were found in the 70 PML cases in this study for which WES data were generated.				
Chromosome	Position_hg19	REF	ALT	SEQ_ID
21	45795833	G	T	1306
21	45795877	G	T	1307
21	45811411	G	T	1308
21	45811438	C	T	1309
21	45815307	T	C	1310
21	45815331	G	A	1311
21	45815343	A	G	1312
21	45815425	C	G	1313
21	45820196	C	T	1314
21	45826486	G	A	1315
21	45826616	C	T	1316
21	45838333	C	T	1317
21	45844780	C	T	1318
21	45845528	G	A	1319
21	45845661	A	G	1320
21	45845699	G	A	1321

TABLE 5-continued

The set of SNVs reported in Tables 7-10, 14, or 15 that were found in the 70 PML cases in this study for which WES data were generated.				
Chromosome	Position_hg19	REF	ALT	SEQ_ID
21	45855099	C	T	1322
22	21235389	A	G	1323
22	23915583	T	C	1324
22	23915745	G	A	1325
22	23917192	G	T	1326
22	36661354	C	T	1327
X	24759574	G	T	1328
X	24759574	G	T	1329

[0612] Table 5 lists, in order of genomic coordinates, all single nucleotide variants (SNVs) that are relevant to the present study, whether as case-level solutions (Tables 7, 8) or potential solutions (Tables 9, 10), or at the level of variant burden analysis (Tables 14, 15). All genome coordinates are based on hg 19.

TABLE 6

Non-redundant list of 419 genes involved in the immune system and/or linked to PML via a CNV				
RefSeq Gene Symbol	Disease Model	Gene Source	Source Annotation	Gene Number (GN)
ACADM	AR	Public_db	MySQL	157
ACKR1	AD	Public_db	MySQL	158
ACP5	AR	Public_db	PMID: 26052098, 27260006, 27821552	159
ADAR	AD_AR	Public_db	PMID: 26052098, 27260006, 27821552	160
ADARB1	unknown	PBio	PMID: 16227093, 17376196, 19482597, 20220309, 21682836, 21809195, 22001568, 22085847, 22113393, 24586166, 24725957, 24760760, 25826567	2
ADK	AR	PBio	PMID: 17205396, 23592612, 25654762, 25720338, 25979489, 26341819, 26642971	161
AGBL4	unknown	PBio	PMID: 17244818, 21074048, 23085998, 25416787, 25332286, 26502776	3
AICDA	AD_AR	Public_db	MySQL; PMID: 23765059	162
AK2	AR	Public_db	PMID: 19043417, 19782549, 20008220, 23765059, 24135998, 24753205, 26454313	163
ALG12	AR	Public_db	MySQL	164
ALPL	AD_AR	PBio	PMID: 18821074, 20049532, 20977932, 21191615, 21289095, 23091474, 23454488, 23860646, 26219705, 26219711, 26219717	165
AP3B1	AR	Public_db	MySQL; PMID: 11590544, 19782549, 24302998, 24753205, 24916509, 25980904, 27889060	166
AP3B2	AR	Public_db	PMID: 26377319, 27889060	167
AP3D1	AR	Public_db	PMID: 26744459, 27889060	168
APOBEC3A	unknown	PBio	PMID: 16720547, 17303427, 20062055, 20615867, 22896697, 23344558, 23640892, 25262471, 25576866, 26416327, 26489798, 26678087	4
APOBEC3B	unknown	PBio	PMID: 16720547, 17303427, 20062055, 20615867, 22896697, 23344558, 23640892, 25262471, 25576866, 26416327, 26489798, 26678087	6
APOL1	association	Public_db	PMID: 27042682	169
ARHGEF7	unknown	PBio	PMID: 11160719, 16983070, 18378701, 19861492, 25284783, 25500533	8
ASH1L	unknown	PBio	PMID: 17923682, 17981149, 22541069, 24012418, 24244179, 25866973, 26002201, 27154821, 27229316, 27434206	170
ASTN2	unknown	PBio	PMID: 2223091, 8602532, 20573900, 24357807, 24381304, 25146927, 25410587, 26514622, 26586575	9
ATL2	unknown	PBio	PMID: 18270207, 19665976, 25773277	171
ATM	AR	Public_db	MySQL; PMID: 19903823, 20301790, 23765059, 24799566, 25692705, 27042682, 27484032, 27884168, 27895165	172
ATR	AD_AR	Public_db	PMID: 17564965, 17151099, 19903823, 20506465, 21615334, 24799566, 25910481	173
AUH	AR	PBio	PMID: 12434311, 12655555, 17130438, 20855850, 25280001, 25597510	10
BACH1	unknown	PBio	PMID: 15068237, 18555605, 22024395, 22791292, 23456643, 23562577, 24752012, 25344725, 25391381, 24752012, 26045540, 26894991	11
BACH2	unknown	PBio	PMID: 17262715, 17991429, 18769450, 22791292, 23728300, 24367030, 24608439, 24681888, 24694524, 25123280, 25344725, 25665584, 25686607, 26444573, 26620562, 26731475, 26894991, 26981933	174
BCL10	AR	Public_db	MySQL	176
BDKRB2	unknown	PBio	PMID: 7787759, 18930543, 22047990, 22095814, 24925394	12
BLM	AR	PBio	PMID: 15137905, 15493327, 17210642, 17321898, 19109166, 19709744, 2032252, 23572515, 24606147	177
BLNK	AR	Public_db	PMID: 23765059	178
BLOC1S6	AR	Public_db	MySQL	179
BMPR2	AD	PBio	PMID: 15877825, 19191909, 23733693, 24334027	13

TABLE 6-continued

Non-redundant list of 419 genes involved in the immune system and/or linked to PML via a CNV				
RefSeq Gene Symbol	Disease Model	Gene Source	Source Annotation	Gene Number (GN)
C11orf65	unknown	Public_db	MySql	181
C1QA	AR	Public_db	PMID: 27821552	182
C1QB	AR	Public_db	PMID: 27821552	183
C1QC	AR	Public_db	PMID: 27821552	184
C5AR1	unknown	PBio	PMID: 1847994, 22964232, 25041699, 25174320, 25455139, 25539817, 25769922, 26059553, 26283482, 26537334	185
CAPZB	unknown	PBio	PMID: 99354614, 19806181, 22493691, 22706086, 22710966, 22918941, 23178720, 26758871	186
CARD11	AD_AR	Public_db	MySql; PMID: 23765059, 25645939, 26525107	187
CARD9	AR	Public_db	PMID: 27222657	188
CASP8	AR	Public_db	PMID: 22365665, 26454313, 27873163, 27999438	189
CCL11	AD	Public_db	MySql	190
CCL2	association	Public_db	MySql	191
CCL5	association	Public_db	MySql	192
CCR2	association	Public_db	MySql	193
CCR5	association	Public_db	MySql	194
CD180	unknown	PBio	PMID: 9763566, 10880523, 21918197, 21959264, 22484241, 23103284, 23483427, 24019553, 25749095, 26371254, 26384474, 26482097, 26555723, 26371254	195
CD19	AR	Public_db	MySql; PMID: 23765059, 26453379	196
CD209	association	Public_db	MySql	197
CD247	AR	Public_db	PMID: 26454313	198
CD27	AR	Public_db	MySql; PMID: 23765059	199
CD27-AS1	unknown	Public_db	MySql	200
CD300LF	unknown	PBio	PMID: 15184070, 15549731, 17202342, 18688020, 19592130, 22288587, 23072861, 23293083, 24035150	23
CD34	unknown	Public_db	PMID: 27042682	201
CD3D	AR	Public_db	PMID: 23765059, 26454313	202
CD3E	AR	Public_db	PMID: 23765059, 26454313	203
CD3G	AR	Public_db	PMID: 23765059, 26454313	204
CD40	AR	Public_db	MySql; PMID: 23765059, 26453379	205
CD55	unknown	PBio	PMID: 12417446, 1385527, 16406700, 16503113, 17678954, 18424707, 19660813, 21143144, 22795896, 24588829, 24639397, 25156074, 25954012, 26423932	207
CD59	AR	Public_db	MySql	208
CD79A	AR	Public_db	PMID: 23765059	209
CD79B	AR	Public_db	PMID: 23765059	210
CD81	AR	Public_db	MySql	211
CD8A	AR	Public_db	PMID: 26454313	212
CDCA7	AR	Public_db	PMID: 26216346	213
CDKN1B	AD	PBio	PMID: 10799578, 10825149, 10916090, 11123298, 11123306, 15557280, 16410832, 17273559, 20854895, 21078910, 22454463, 24317118, 25213837	24
CEBPB	unknown	Public_db	PMID: 27042682	214
CENPM	unknown	PBio	PMID: 15183305, 16391015, 19711193, 25006165	25
CHD7	AD	Public_db	PMID: 18505430, 18976358, 26454313, 27484032	215
CHEK1	unknown	Public_db	PMID: 19903823, 27042682	216
CIITA	AR	Public_db	PMID: 23765059, 26454313, 27484032	217
CLCN7	AD	Public_db	PMID: 21107136, 25992615	218
COG4	AR	PBio	PMID: 18086915, 18256213, 20065092, 20143049, 21421995, 23462996, 23865579, 24784932, 26125015	26
COG6	AR	Public_db	MySql	219
COMMMD6	unknown	PBio	PMID: 14685242, 15799966, 16573520, 20126548, 25355947, 27441653	27
CORO1A	AR	Public_db	PMID: 23887241, 26454313	220
CR2	AR	Public_db	MySql	221
CRADD	AR	PBio	PMID: 11573962, 21242994, 22323537, 24958727, 26190521, 27135977	28
CRTC3	unknown	PBio	PMID: 15466468, 2032252, 21536665, 23033494, 23241891, 25114223, 25316186, 25351958, 26937622	222
CSF3R	AR	Public_db	PMID: 24753537, 26324699, 27789332	223
CTLA4	AD	Public_db	PMID: 18219311, 25213377, 25329329	224
CTPS1	AR	Public_db	MySql	225
CTSC	AR	Public_db	PMID: 27222657	226
CX3CR1	association	Public_db	MySql	227
CXCL12	association	Public_db	MySql	228
CXCL9	unknown	Public_db	PMID: 27042682	229
CXCR1	association	Public_db	MySql	230
CXCR4	AD	Public_db	PMID: 19782549, 19950235, 23765059, 24753205, 25645939, 28009639	231
CXorf40A	unknown	PBio	PMID: 15541360, 24916366, 26881174	232
CYBB	XLR	Public_db	PMID: 27222657	233
CYP2S1	unknown	PBio	PMID: 15681441, 23933117	234
DCI1RE1C	AR	Public_db	PMID: 26454313, 26476407, 27484032	235
DDX1	unknown	Public_db	PMID: 27042682	236
DDX58	AD	Public_db	PMID: 23592984, 25692705, 25794939, 26052098, 26748340, 26848516, 26987611, 27260006, 27821552	237

TABLE 6-continued

Non-redundant list of 419 genes involved in the immune system and/or linked to PML via a CNV				
RefSeq Gene Symbol	Disease Model	Gene Source	Source Annotation	Gene Number (GN)
DHX58	association	Public_db	PMID: 25794939, 26748340, 26848516	238
DKC1	XLR	Public_db	MySQL; PMID: 23765059	239
DNER	unknown	PBio	PMID: 15965470, 16298139, 16997755, 17765022, 18474614, 20058045, 20367751, 22447725, 23041955, 23328254, 24248099, 24935874, 26869529	31
DOCK2	AR	Public_db	MySQL	241
DSC1	unknown	PBio	PMID: 16048752, 20222919, 22692770, 24460202, 24680560, 25078507, 25244249, 26043694, 26758100	243
DUSP16	unknown	PBio	PMID: 15284860, 21613215, 24311790, 25716993, 26381291, 27162525	32
ECRP	unknown	PBio	PMID: 9826755, 12855582, 25271100, 26184157	33
EDIL3	unknown	PBio	PMID: 22601780, 23518061, 24060278, 24504014, 25385367, 26038125	34
EEA1	unknown	PBio	PMID: 16670179, 22591512, 24491918, 24561067, 26478006, 26909655, 27077111	35
EGR1	unknown	PBio	PMID: 15308739, 19050264, 19812322, 20414733, 21368226, 21622185, 22554935, 25613134, 26052046, 26980486, 11910893, 14647476, 18203138, 24627779, 25368162, 27192563	244
EHF	unknown	PBio	PMID: 16380452, 17027647, 19801549, 20879862, 21617703, 24219556, 25217163	36
ELANE	AD	Public_db	PMID: 20008220, 24145314, 27222657	245
EMB	unknown	PBio	PMID: 8432389, 15917240, 18209069, 19164284, 25773908	37
EPG5	AR	Public_db	MySQL; PMID: 21965116, 23222957, 23838600, 26917586, 26927810, 27588602	246
ETF1	unknown	PBio	PMID: 20418372, 25606970, 26384426, 26833392, 27387891	247
ETV6	AD	PBio	PMID: 19264918, 20350664, 21714648, 22438058, 25581430, 25807284, 26718572, 27365488	38
F9	XLR	Public_db	MySQL	248
FAS	AD	Public_db	PMID: 27222657	249
FASLG	AD	Public_db	PMID: 27222657	250
FCGR2A	AD_AR	Public_db	MySQL	251
FCGR3A	AR	Public_db	MySQL	252
FCN3	AR	Public_db	MySQL	253
FEZ1	unknown	Public_db	PMID: 27042682	254
FHL2	unknown	PBio	PMID: 16389449, 20592280, 22417706, 22633286, 23212909	39
FOS	unknown	Public_db	PMID: 27042682	255
FOXH1	unknown	Public_db	PMID: 27042682	256
FOXP1	AR	Public_db	MySQL	257
FOXP3	XLR	Public_db	PMID: 26454313	258
FPR1	unknown	PBio	PMID: 8994115, 10229829, 10611407, 17084101, 22934745, 23230437, 25605714, 25826286, 26101324, 26701131, 27034344, 27100350, 27131862, 27154726	259
FPR2	unknown	PBio	PMID: 8994115, 10229829, 10611407, 17084101, 22934745, 23230437, 25605714, 25826286, 26101324, 26701131, 27034344, 27100350, 27131862, 27154726	41
FPR3	unknown	PBio	PMID: 8994115, 10229829, 10611407, 17084101, 22934745, 23230437, 25605714, 25826286, 26101324, 26701131, 27034344, 27100350, 27131862, 27154726	42
FUK	unknown	PBio	PMID: 11753075, 12651883, 15774760, 19394435, 19647987, 20363321, 22134107, 22203233, 22276660, 22461019, 24239607	43
G6PC3	AR	Public_db	PMID: 20008220, 24145314, 25879134, 26479985	260
GATA2	AD	Public_db	PMID: 23765059, 23887241	261
GDA	unknown	PBio	PMID: 10595517, 18600524, 20826431, 23838888, 24834013	44
GDPD4	unknown	PBio	PMID: 24373430, 24977479, 24977485, 24977489, 25528375, 25596343	45
GFI1	AD	Public_db	PMID: 20008220, 24145314	262
GOLGB1	unknown	PBio	PMID: 17475246, 21217069, 22034594, 23555793, 24046448	263
GPATCH2	unknown	PBio	PMID: 19432882, 25353171, 25376275	46
GPC5	unknown	PBio	PMID: 24130709, 24943672, 25354479, 26224662, 26349477	47
GPRC5A	unknown	PBio	PMID: 19593893, 20959490, 22239913, 25621293, 25714996, 26165721	264
GRAP2	unknown	Public_db	PMID: 25452106, 25636200, 26246585	265
GRIA3	XLR	PBio	PMID: 10441169, 12682273, 17202328, 18590483, 25904555, 26648591	51
GTPBP4	unknown	PBio	PMID: 17785438, 26015807	52
HAX1	AR	Public_db	PMID: 20008220, 24145314	266
HCN1	AD	PBio	PMID: 9405696, 9630217, 9634236, 9921901, 11133998, 23042740, 23077068, 23319474, 24403084, 24747641, 24756635, 25580535, 26578877	53
HELLS	AR	Public_db	PMID: 26216346	267
HEXA	AR	PBio	PMID: 20301397, 21997228, 23727835, 24445368	54
HIVEP1	association	Public_db	MySQL; PMID 20226436, 26117544	268
HIVEP2	AD	Public_db	MySQL; PMID: 21475200, 21936769, 23389689, 24366360, 26153216, 26483320, 27003583	269
HIVEP3	unknown	Public_db	MySQL	270
HK2	unknown	PBio	PMID: 2749583, 4265132, 19519254, 2496891, 23874603, 25525876, 25602755	55
HMGB3	unknown	PBio	PMID: 12714519, 15259015, 15358624, 16945912, 22014684, 23994280, 26553261	56
HNRNPLL	unknown	PBio	PMID: 18669861, 18719244, 19100700, 20505149, 22073166, 23934048, 24476532, 25825742	271
HP	unknown	PBio	PMID: 16046400, 19380867, 19795414, 26445729	272
HPCAL1	unknown	PBio	PMID: 12445467, 24699524, 25519916, 26659654, 26729710	273
HPR	unknown	PBio	PMID: 16046400, 19380867, 19795414, 26445729	57
HTR2A	association	Public_db	PMID: 19204164, 24089568, 25078361, 26056932, 27003757, 27042682	274
ICOS	AR	Public_db	MySQL; PMID: 19380800, 23765059, 27250108	275

TABLE 6-continued

Non-redundant list of 419 genes involved in the immune system and/or linked to PML via a CNV				
RefSeq Gene Symbol	Disease Model	Gene Source	Source Annotation	Gene Number (GN)
IDI1	unknown	PBio	PMID: 14629038, 17202134, 19454010, 20955688, 22579571, 23585482, 25950736	276
IDI2	unknown	PBio	PMID: 14629038, 17202134, 19454010, 20955688, 22579571, 23585482, 25950736	59
IDI2-AS1	unknown	PBio	See IDI2	60
IDO2	unknown	PBio	PMID: 18219311, 18364004, 19487973, 19799997, 20197554, 20484729, 20693847, 21084489, 21406395, 21990421, 22754762, 24391212, 24402311, 24844751, 25477879, 25478733, 25541686, 25949913, 26308414, 27183624	61
IFIH1	AD	Public_db	PMID: 21156324, 24686847, 24995871, 25794939, 26052098, 26748340, 27260006, 27821552	277
IFNAR1	association	Public_db	PMID: 27821552	278
IFNAR2	AR	Public_db	PMID: 26424569, 27821552	279
IFNG	association	Public_db	MySql	280
IFNGR1	AD_AR	Public_db	MySql	281
IFNGR2	AD_AR	Public_db	MySql; PMID: 15356149, 23161749	282
IFNLR1	unknown	PBio	PMID: 12469119, 12483210, 15166220, 22386267, 22891284, 25634147, 25904743, 25941255	62
IGLL1	AR	Public_db	PMID: 25502423	283
IKBKB	AD_AR	Public_db	MySql; PMID: 17047224, 17072332, 25764117, 25930993, 26117626, 26525107	284
IKBKG	XLD_XLR	Public_db	MySql; PMID: 17047224, 21455173, 21455181, 23765059, 25764117, 25886387, 25930993, 26117626, 26525107	285
IKZF1	AD	Public_db	PMID: 26454313, 26981933	286
IL10	AR	Public_db	PMID: 23887241	287
IL10RA	AR	Public_db	PMID: 23887241	288
IL10RB	AR	Public_db	PMID: 23887241	289
IL12B	AR	Public_db	MySql	290
IL12RB1	AR	Public_db	MySql	291
IL17F	AD	Public_db	PMID: 22284928, 23887241, 24240291, 24690400, 25890879, 27144517	292
IL17RA	AR	Public_db	PMID: 23887241	293
IL1B	AD	Public_db	PMID: 15327898, 20543597, 24248593, 26525107, 27730320, 27873163, 27999438	294
IL21	AR	Public_db	MySql	295
IL21R	AD_AR	Public_db	PMID: 23765059, 23887241	296
IL2RA	AR	Public_db	MySql	297
IL2RG	XLR	Public_db	PMID: 23765059, 26454313, 27484032	298
IL4R	association	Public_db	MySql	299
IL7	unknown	PBio	PMID: 21508983, 22288682, 24507157, 24979548, 25130296, 25214510, 25411246, 25734144, 26537673, 26675348, 26908786	300
IL7R	AR	Public_db	PMID: 26454313, 27484032	301
IRAK4	AD_AR	Public_db	PMID: 23766853, 25232776, 25344726, 25764117, 25886387, 25930993, 26785681, 27845762	302
IRF3	AD	Public_db	PMID: 23388631, 26513235, 26748340	303
IRF7	AR	Public_db	MySql; PMID: 26748340	304
IRF8	AD_AR	Public_db	PMID: 23887241	305
IRGM	association	PBio	PMID: 14707092, 17911638, 22174682, 22722598, 23084913, 23335927	306
ISG15	AR	Public_db	MySql; PMID: 26052098, 27260006, 27821552	307
ITSN2	unknown	PBio	PMID: 11748279, 15020715, 17696400, 17696407, 22558309, 22975684, 23986746, 24097067, 24284073, 25797047, 26479042	309
JAGN1	AR	Public_db	PMID: 25129144	310
JAK3	AR	Public_db	PMID: 23765059, 26454313	311
JMY	unknown	PBio	PMID: 19287377, 20573979, 20574148, 20888769, 21965285, 23291261, 25015719, 26223951, 26305109	312
JPX	association	PBio	PMID: 21029862, 23791181, 23943155	64
JUN	unknown	Public_db	PMID: 16928756, 27042682	313
KANK1	association	PBio	PMID: 18458160, 20164854, 21685469, 24399197, 25961457, 26656975	65
KAT6B	AD	PBio	PMID: 17460191, 17694082, 22715153, 25920810	66
KCTD7	AR	PBio	PMID: 17455289, 20301601, 21710140, 22606975, 22638565, 22748208, 25060828, 27629772, 27742667	67
KITLG	AD	Public_db	PMID: 27042682	314
LAMTOR2	AR	Public_db	MySql; PMID 19782549, 20008220, 24753205	315
LARP4B	unknown	PBio	PMID: 20573744, 23815932, 25534202, 26001795, 26501340, 26644407	69
LCP2	unknown	Public_db	PMID: 12874226, 18219311, 19056831, 23494777, 26246585	317
LIG1	AR	Public_db	MySql	318
LIG4	AR	Public_db	MySql	319
LOC102724297	unknown	Public_db	MySql	320
LOC400710	unknown	PBio	ncRNA, limited gene information; see SNAR gene family (adjacent locus)	321
LRBA	AR	Public_db	MySql; PMID: 23765059, 27873163, 27192563	322
LYST	AR	Public_db	PMID: 19302049, 19782549, 20008220, 24753205, 24916509, 26454313, 27881733	323
IMAGEA9	unknown	PBio	PMID: 15222021, 15900605, 21093980, 21791470, 25315972, 25445503, 25755744	324
MAGEA9B	unknown	PBio	PMID: 15222021, 15900605, 21093980, 21791470, 25315972, 25445503, 25755744	325
MAGT1	XLR	Public_db	PMID: 23887241, 27873163, 25504528	326
MALL	unknown	PBio	PMID: 11294831, 19064697, 24101378, 24746959, 26109641, 26622604, 26641089, 26772392, 27583248, 27846891	72
MALT1	AR	Public_db	MySql; PMID: 26525107	327

TABLE 6-continued

Non-redundant list of 419 genes involved in the immune system and/or linked to PML via a CNV				
RefSeq Gene Symbol	Disease Model	Gene Source	Source Annotation	Gene Number (GN)
MAP3K2	unknown	PBio	PMID: 11032806, 11278622, 12138187, 14734742, 16430878, 21333552, 2437584, 24847879, 25012295, 26056008	328
MAPK1	unknown	Public_db	PMID: 14671106, 27042682	329
MAPK3	unknown	Public_db	PMID: 14671106, 27042682	330
MAPK9	unknown	PBio	PMID: 15023353, 23685277, 24673683, 25762148, 26141991	73
MAVS	association	Public_db	PMID: 23582325, 26513235, 26987611	331
MCEE	AR	PBio	PMID: 17846917, 20301409, 21365456, 23726524, 24532006, 25763508, 26725562	74
MECP2	XLD_XLR	Public_db	PMID: 27042682	332
MEX3C	unknown	PBio	PMID: 18779327, 22357625, 22658931, 22863774, 22927639, 23140835, 23446422, 23999169, 24706898, 24741071	333
MGAT5	unknown	PBio	PMID: 12417426, 15585841, 18292539, 20089585, 20117844, 25768892, 26972830	75
MKL1	AR	PBio	PMID: 12944485, 22626970, 26098208, 26098211, 26221020, 26241940, 26405212, 26224645, 26554816	89
MRE11A	AR	Public_db	PMID: 23388631, 23765059	334
MS4A1	AR	Public_db	MySql; PMID: 23765059	335
MSN	unknown	PBio	PMID: 9070665, 10444190, 11777944, 12445265, 14758359, 16368573, 17110458, 18025306, 18725395, 21486194, 23526587, 23613524, 24250818, 24358210, 24760896, 25746045	336
MTHFD1	AR	Both	PMID: 26454313	337
MYD88	AD_AR	Public_db	PMID: 23766853, 25344726, 25764117, 25886387, 25930993, 26371186, 27435819	338
NBN	AD_AR	Public_db	MySql; PMID: 23765059	339
INFC	unknown	PBio	PMID: 11559801, 15327898, 16928756, 18474555, 19058033, 22205750	340
NFIL3	unknown	PBio	PMID: 20080759, 20697558, 22075207, 23453631, 24070385, 24277151, 24280221, 24442434, 24909887, 25092873, 25113970, 25310240, 25611557, 25614966, 25801035, 25249449, 25620184, 26379372, 26806130, 26880402	92
NFKB1	AD	Public_db	PMID: 22081022, 26279205	341
NFKB2	AD	Public_db	MySql; PMID: 25764117	342
NFKBIA	AD	Public_db	MySql; PMID: 23765059, 25645939, 25764117	343
NHEJ1	AR	Public_db	MySql; PMID: 23765059	344
NLRP12	AD	PBio	PMID: 17947705, 18230725, 20861596, 21978668, 23318142, 23970817, 24282415, 24347638, 25249449, 25620184, 25902475, 26083549, 26343520, 26386126, 26521018	93
NLRP3	AD	Public_db	PMID: 16724804, 19302049, 23592984, 26848516, 27999438	345
NOD2	AD	Public_db	PMID: 16724804, 19302049, 23584365, 26509073, 26848516, 26953272	346
NQO2	unknown	PBio	PMID: 16253210, 16905546, 17720881, 18552348, 26046590	94
NRIP1	unknown	PBio	PMID: 18267075, 23241901, 24969109, 25066731, 25697398, 25879677, 26937622	95
ORAI1	AD_AR	Public_db	PMID: 19075015, 20004786, 21790973, 22144678, 23765059, 26454313, 26469693	347
OSTM1	AR	Public_db	PMID: 16813530, 19507210, 21107136, 23685543	348
OVOL2	AD	PBio	PMID: 16423343, 25267199, 26619963, 26749309	98
PDE3B	unknown	PBio	PMID: 17220874, 23276671, 25816736, 26203135, 26297880, 26374610	99
PDGFRA	association	PBio	PMID: 12660384, 18634583, 18701889, 19246520, 19839938, 20032375, 20569695, 21123584, 21975205, 22449623, 22523564, 23771592, 25319708, 25940087	100
PDSS2	AR	PBio	PMID: 17186472, 18437205, 18784258, 21567994, 21871565, 21983691, 23150520	101
PGM3	AR	Public_db	MySql; PMID: 25502423	349
PHACTR4	unknown	PBio	PMID: 15107502, 17609112, 22215804, 22215812, 22766235, 23076051, 23203801, 23319639, 24748504, 26850007	102
PIAS1	unknown	PBio	PMID: 10805787, 10858346, 14644436, 15297606, 15311277, 17065208, 17540171, 18056374, 19857525, 20966256, 22969086, 22982248, 23299081, 24036127	103
PIAS2	unknown	PBio	PMID: 9724754, 11117529, 12077349, 12764129, 14514699, 15582666, 16460827, 19549844, 21156324, 21779164, 22210188, 22982248, 24344134, 25484205, 25434787, 26223632	350
PIK3CD	AD	Both	MySql; PMID: 24165795, 25133419, 25645939, 26437962, 26453379, 27379089, 27426521, 27873163, 14647476, 27192563	104
PIK3R1	AD_AR	Public_db	PMID: 23765059, 23887241, 25645939, 26246585, 26453379, 27076228, 14647476, 27192563	351
PKHD1	AR	PBio	PMID: 8178487, 15052665, 17450421, 23423256, 24964219, 24984783, 25186187, 26502924	105
PLCG2	AD	Public_db	PMID: 19056831, 23000145, 23765059, 23887241, 25452106, 25636200, 25645939, 26246585, 27192563	352
PMS2	AR	Public_db	MySql; PMID: 23765059	353
PNPLA4	unknown	PBio	PMID: 22289388, 23741432, 26017929, 26164793, 26713677, 26741492, 26968210	107
PNPT1	AR	PBio	PMID: 14563561, 15492272, 16410805, 16687933, 17983748, 19580345, 23084291, 23221631, 24143183, 24729470, 25457163	108
POLA1	XLR	Public_db	PMID: 27019227, 27821552	355
POLE	AR	Public_db	MySql; PMID: 23230001, 23765059, 25948378	356
PPP2R3B	unknown	PBio	PMID: 9847399, 10629059, 11593413, 18353419, 20485545, 26683421	109
PRF1	AD_AR	Public_db	PMID: 17311987, 19302049, 21881043, 24916509, 25776844, 26454313, 26864340, 27391055	357
PRKCB	unknown	PBio	PMID: 10872892, 15488737, 16935002, 17060474, 17395590, 19907441, 21997316, 22994860, 23959874, 24550541, 25548371, 24550541, 25808972, 26509731, 26510741	110

TABLE 6-continued

Non-redundant list of 419 genes involved in the immune system and/or linked to PML via a CNV				
RefSeq Gene Symbol	Disease Model	Gene Source	Source Annotation	Gene Number (GN)
PRKCD	AR	Public_db	MySQL; PMID: 23319571, 27250108, 27873163	358
PRKCH	unknown	PBio	PMID: 15327898, 16571806, 18353419, 22114277, 22155788, 22892130, 23868949, 24705298, 25617472, 25889880	111
PRKDC	AD_AR	Public_db	PMID: 12847277, 23722905, 26454313, 26838362, 27980111	359
PROC	AD_AR	PBio	PMID: 2437584, 18751723, 21114396, 22447930, 24162617	360
PSMB8	AR	Public_db	PMID: 26052098, 27260006, 27821552	361
PSTPIP1	AD	PBio	PMID: 9488710, 11313252, 12530983, 14707117, 16724804, 19290936, 19302049, 24421327, 25040622, 25645939, 25814341, 26386126, 26919742	112
PTEN	AD	Public_db	PMID: 26246517, 27426521	362
PTPN2	unknown	PBio	PMID: 11909529, 12359225, 12847239, 19290937, 19825843, 19930043, 20473312, 20564182, 20848498, 21220691, 22080861, 22080863, 22671594, 24442435, 24445916, 24608439, 24849651, 24997008, 25548153, 25581833	113
PTPRC	AR	Public_db	PMID: 26454313	363
PTPRN2	unknown	PBio	PMID: 9714834, 10426369, 11086001, 11086294, 11793386, 15114673, 19361477, 23595248, 24988487, 26141787, 26609326	114
PURA	AD	Public_db	PMID: 27042682	364
RAB27A	AR	Public_db	PMID: 19302049, 20008220, 21881043, 23810987	365
RAB37	unknown	PBio	PMID: 21805469, 22899725, 26931073, 27798165	115
RAB7A	AD	PBio	PMID: 25992615, 27588602	366
RABGEF1	unknown	PBio	PMID: 12505986, 15143060, 15235600, 16499958, 16533754, 16605131, 17341663, 20829437, 22846990, 23552075, 24569883, 24957337, 25427001, 26567216, 26588713, 27791468	367
RAC2	AD	Public_db	MySQL	368
RAD51	AD	Public_db	PMID: 25310191, 27042682	369
RAG2	AR	Public_db	PMID: 23765059, 23887241, 26454313, 27808398	371
RBCK1	AR	Public_db	MySQL; PMID: 21455173, 21455181, 23765059, 23969028, 24958845, 25764117, 25930993, 26008899, 26525107, 27810922	372
RBFOX1	unknown	PBio	PMID: 23350840, 24039908, 25043849, 26500751, 26687839	116
RCC1	unknown	PBio	PMID: 1961752, 18442486, 19060893, 20347844, 23536659, 25452301, 26864624	117
RFX5	AR	Public_db	PMID: 23765059, 26454313	373
RFXANK	AR	Public_db	PMID: 23765059, 26454313	374
RFXAP	AR	Public_db	PMID: 23765059, 26454313	375
RGCC	unknown	PBio	PMID: 19158077, 19652095, 23000427, 24973210, 25770350, 26134570	118
RHOQ	unknown	PBio	PMID: 10490598, 12456725, 14734537, 16246732, 17016434, 19258391, 22916134, 24223996, 24297911, 24663214, 24667291	119
RIPK1	association	Public_db	PMID: 21455173, 27999438	376
RIPK3	association	Public_db	PMID: 22365665, 27999438	377
RMRP	AR	Public_db	MySQL; PMID: 19782549, 20008220, 24753205	378
RNASE3	unknown	PBio	PMID: 19515815, 26184157	120
RNASEH2A	AR	Public_db	PMID: 26052098, 27260006, 27821552	379
RNASEH2B	AR	Public_db	PMID: 26052098, 27260006, 27821552	380
RNASEH2C	AR	Public_db	PMID: 26052098, 27260006, 27821552	381
RNASEL	association	Public_db	PMID: 24995003, 27525044	382
RNF168	AR	Public_db	MySQL; PMID: 23765059	383
RNF31	AR	Public_db	PMID: 21455173, 21455181, 23969028, 24958845, 26008899, 26525107, 27810922	384
RNU4ATAC	AR	Public_db	PMID: 27222657	385
RPTOR	unknown	PBio	PMID: 16959881, 22810227, 23349361, 23812589, 24287405, 24303063, 24671993, 24948799, 26678875	123
RTKL1	AR	Public_db	MySQL; PMID: 23329068, 23765059, 24009516, 25607374, 26810774	386
RTKL1-TNFRSF6B	unknown	Public_db	MySQL; PMID: 23329068, 23765059, 25607374	387
SALL2	AR	PBio	PMID: 11734654, 15082782, 18818376, 19076363, 19131967, 21362508, 21689070, 21791360, 22074632, 22978642, 23029531, 24040083, 24412933, 24903482, 25360671, 25580951, 25608837, 26181197	388
SAMHD1	AR	Public_db	PMID: 26052098, 27260006, 27821552	389
SBDS	AR	Public_db	PMID: 20008220, 21062271, 27418648, 27658964	390
SERPINB4	unknown	PBio	PMID: 15203215, 19070595, 21857942, 22451727, 22808225, 24560885, 24635038, 25111616, 25133778, 25213322	124
SERPINB6	AR	PBio	PMID: 14670919, 20451170, 24172014, 24359430	125
SH2D1A	XLR	Public_db	MySQL; PMID: 19302049, 23765059, 25744037	391
SHARPIN	unknown	Public_db	PMID: 21455181, 22901541, 23969028, 24958845, 26525107, 26848516, 27810922, 27892465	392
SKIV2L	AR	Public_db	PMID: 27260006, 27821552	393
SLC17A5	AR	PBio	PMID: 14742248, 15006695, 15172005, 16575519, 18399798, 20007460, 20951965, 21628664, 22778404, 23760462, 23889254, 25494612, 25855729, 25879139	127
SLC37A4	AR	Public_db	PMID: 20008220, 20301489	394
SLC3A2	unknown	PBio	PMID: 22588539, 22624878, 23297381, 24491544, 25002078, 26172215, 26439699, 26444422	126
SLC46A1	AR	Public_db	PMID: 26454313	395
SLC8A1	unknown	PBio	PMID: 23224883, 23224887, 23224890, 23224891, 26045217, 26418956, 26775040, 26859825, 26924806	396

TABLE 6-continued

Non-redundant list of 419 genes involved in the immune system and/or linked to PML via a CNV				
RefSeq Gene Symbol	Disease Model	Gene Source	Source Annotation	Gene Number (GN)
SMAD2	unknown	Public_db	PMID: 27042682	397
SMAD3	AD	Public_db	PMID: 27042682	398
SMAD4	AD	Both	PMID: 12202226, 14987161, 16800882, 19420158, 25637015, 25705527, 26454313, 27042682	399
SNAP29	AR	Public_db	PMID: 15968592, 21073448, 27588602	400
SNAR-A1	unknown	PBio	PMID: 25327818, 25447144	401
SNAR-A10	unknown	PBio	PMID: 25327818, 25447144	402
SNAR-A11	unknown	PBio	PMID: 25327818, 25447144	403
SNAR-A12	unknown	PBio	PMID: 25327818, 25447144	404
SNAR-A13	unknown	PBio	PMID: 25327818, 25447144	405
SNAR-A14	unknown	PBio	PMID: 25327818, 25447144	406
SNAR-A2	unknown	PBio	PMID: 25327818, 25447144	407
SNAR-A3	unknown	PBio	PMID: 25327818, 25447144	408
SNAR-A4	unknown	PBio	PMID: 25327818, 25447144	409
SNAR-A5	unknown	PBio	PMID: 25327818, 25447144	410
SNAR-A6	unknown	PBio	PMID: 25327818, 25447144	411
SNAR-A7	unknown	PBio	PMID: 25327818, 25447144	412
SNAR-A8	unknown	PBio	PMID: 25327818, 25447144	413
SNAR-A9	unknown	PBio	PMID: 25327818, 25447144	414
SNAR-B1	unknown	PBio	PMID: 25327818, 25447144	415
SNAR-B2	unknown	PBio	PMID: 25327818, 25447144	416
SNAR-C1	unknown	PBio	PMID: 25327818, 25447144	417
SNAR-C2	unknown	PBio	PMID: 25327818, 25447144	418
SNAR-C3	unknown	PBio	PMID: 25327818, 25447144	419
SNAR-C4	unknown	PBio	PMID: 25327818, 25447144	420
SNAR-C5	unknown	PBio	PMID: 25327818, 25447144	421
SNAR-D	unknown	PBio	PMID: 25327818, 25447144	422
SNAR-E	unknown	PBio	PMID: 25327818, 25447144	423
SNAR-F	unknown	PBio	PMID: 25327818, 25447144	424
SNAR-G1	unknown	PBio	PMID: 25327818, 25447144	425
SNAR-G2	unknown	PBio	PMID: 25327818, 25447144	426
SNAR-H	unknown	PBio	PMID: 25327818, 25447144	427
SNAR-I	unknown	PBio	PMID: 25327818, 25447144	428
SNCA	AD	PBio	PMID: 12406186, 14648159, 16953112, 19115126, 19432400, 19652146, 22209147, 23378275, 23771222, 24586351, 24593806, 25092570, 25450953, 25522431, 25635231, 25866630, 26087293, 26272943, 26342897, 26646749	429
SNHG3	unknown	PBio	PMID: 22308462, 22843687, 26373735	128
SNX10	AR	Public_db	PMID: 22499339, 23123320	430
SNX5	unknown	PBio	PMID: 10600472, 11128621, 14499622, 15133132, 15561769, 16857196, 18596235, 21725319, 21903422, 21943487, 23213485, 24820351, 26220253	130
SOCS2	unknown	PBio	PMID: 19279332, 21403007, 22693634, 22795647, 23455506, 24400794, 26216515, 26709655, 26765997, 27071013, 27158906, 27330188, 27338192	131
SP110	AR	Public_db	MySQL	431
SP140	unknown	Public_db	MySQL	432
SPINK5	AR	Public_db	PMID: 19683336, 26865388, 27222657, 27905021	433
SQSTM1	AD_AR	Public_db	PMID: 19229298, 27715390	434
SRSF1	unknown	Public_db	PMID: 27042682	435
ST8SIA5	unknown	PBio	PMID: 11089916, 15829700	133
STAT2	AR	Public_db	PMID: 23391734, 26122121, 27821552	437
STAT5B	AR	Public_db	MySQL	439
STIM1	AD_AR	Public_db	PMID: 20004786, 21790973, 23765059, 26454313, 26469693	440
STIM2	unknown	PBio	PMID: 20004786, 21790973, 21880262, 22129055, 22477146, 22914293, 25157823, 26109647, 26469693	134
STK4	AR	Public_db	PMID: 19782549, 23765059, 23887241, 24753205, 26029204	441
STX11	AR	Public_db	PMID: 19302049, 21881043, 24916509, 26454313	442
STXBP2	AD_AR	Public_db	PMID: 21881043, 24916509, 25564401, 26454313	443
SYNCRIP	unknown	PBio	PMID: 10734137, 18045242, 19331829, 19232660, 22493061, 22935615, 23679954, 23700384, 24844655, 25100733, 26641092	444
T	AD_AR	PBio	PMID: 11897834, 17438107, 23064415, 23662285, 24253444, 24556085, 25186612, 26210634, 26919728	445
TAP1	AR	Public_db	PMID: 26454313	446
TAP2	AR	Public_db	PMID: 26454313	447
TAPBP	unknown	Public_db	PMID: 26454313	448
TAZ	XLR	Public_db	PMID: 20008220	449
TBC1D16	unknown	PBio	PMID: 16923123, 19077034, 21250943, 23019362, 23485563, 23812537, 24513270, 26030178	136
TBK1	AD	Public_db	PMID: 23887241, 25930993, 26513235, 28049150	450
TBX1	AD	Public_db	PMID: 26454313	451
TCIRG1	AD_AR	Public_db	MySQL; PMID 19507210, 19782549, 24753205, 27233968	452
TICAM1	AD_AR	Public_db	PMID: 22105173, 23887241, 25764117, 25930993, 26513235, 28049150	453
TLR3	AD	Public_db	PMID: 23592984, 23887241, 25930993, 26513235, 27810922, 27873163, 27881733	454

TABLE 6-continued

Non-redundant list of 419 genes involved in the immune system and/or linked to PML via a CNV				
RefSeq Gene Symbol	Disease Model	Gene Source	Source Annotation	Gene Number (GN)
TLR4	association	Both	PMID: 12124407, 17893200, 18946062, 19843948, 20521908, 21677132, 22474023, 22962435, 23055527, 23890253, 25365308, 25454804, 25930993, 26189680, 26453379, 27881733	455
TMEM173	AD	Public_db	PMID: 23388631, 25645939, 25692705, 26052098, 27260006, 27801882, 27821552	456
TNF	association	Public_db	MySQL; PMID: 27042682	457
TNFAIP3	AD	Public_db	PMID: 23969028, 26642243, 27845235	458
TNFRSF10A	Unknown	PBio	PMID: 10889508, 11602752, 11704827, 11777550, 11844843, 12390973, 12694389, 14975593, 15007095, 16394652, 16554480, 17671142, 19690337, 20921531	138
TNFRSF11A	AD_AR	Public_db	PMID: 17088646, 17360404, 18281276, 18606301, 19380800, 19507210, 25102334, 25393853, 27003757, 27016605	459
TNFRSF11B	AR	Public_db	PMID: 19507210, 25102334, 25393853, 27003757	460
TNFRSF13B	AD_AR	Public_db	MySQL; PMID: 17467261, 17492055, 18978466, 18981294, 19629655, 20889194, 21458042, 22697072, 23765059, 25454804, 25930993, 26727773, 27123465	461
TNFRSF13C	AR	Both	MySQL; PMID: 16769579, 17785824, 18784835, 18813230, 19136305, 19406831, 20547827, 20547828, 20817206, 21897850, 22028296, 22030463, 23684423, 24101550, 24953530, 25454804, 25637018, 25724205, 26419927, 26453379, 26600308, 26888554	139
TNFRSF18	unknown	PBio	PMID: 16439533, 19162554, 19363449, 22017440, 23432692, 24484736, 25738498	140
TNFRSF4	AR	Public_db	MySQL	462
TNFRSF8	unknown	PBio	PMID: 10921351, 15990453, 16472805, 18852356, 20141444, 20378007, 21933041, 23115213, 23307550, 23654079, 24809535, 25999451	463
TNFSF11	AR	Public_db	PMID: 17088646, 17360404, 18281276, 18606301, 19507210, 25992615, 27003757	464
TNFSF12	association	Public_db	PMID: 23765059	465
TP53	AD_AR	Public_db	MySQL; PMID: 11048806, 11079782, 12009037, 19282432, 26870672	466
TRAF3	AD	Public_db	PMID: 20832341, 23887241, 25764117, 25930993, 28049150	467
TRAF6	unknown	Public_d b	PMID: 10215628, 10421844, 25200954, 27808398, 27999438	468
TRAFD1	unknown	PBio	PMID: 16221674, 18849341, 23913580, 25909814, 25992615, 26283173	141
TREX1	AD_AR	Public_db	PMID: 26052098, 27260006, 27821552	469
TRNT1	AR	Public_db	MySQL; PMID: 25193871	470
TRPM2	unknown	PBio	PMID: 9806837, 16585058, 18569867, 19411837, 20107186, 25012489, 25049394, 25088676, 26300888, 26558786, 26679996, 26942016, 26969190, 27405665, 27872485	142
TTC7A	AR	Public_db	MySQL; PMID: 27873163	471
UBE2N	unknown	PBio	PMID: 21512573, 23159053, 24906799, 25343992, 25503582, 25548215, 26085214, 26150489, 26212332, 26518362	145
UNC119	AD	Public_db	MySQL	472
UNC13D	AR	Public_db	PMID: 19302049, 21881043, 24916509, 25564401, 25980904, 26454313	473
UNC93B1	association	Public_db	PMID: 23810987, 23887241, 25930993, 27873163	474
UNG	AR	Public_db	MySQL; PMID: 23765059	475
USP18	AR	Public_db	PMID: 27016605, 27325888, 27801882, 27821552	476
USP20	unknown	Public_db	PMID: 27801882	477
VAPA	unknown	PBio	PMID: 9657962, 10523508, 10655491, 11511104, 12931207, 18713837, 23536298, 24076421, 24569996, 25015719	478
VCP	AD	Public_db	PMID: 24248593, 27730320	479
VDAC1	unknown	PBio	PMID: 10620603, 25874870, 26322231, 26542804, 26616244, 26758954, 26878172	480
VPS13B	AR	Public_db	PMID: 20008220, 20301655	481
VPS45	AR	Public_db	MySQL; PMID: 23738510, 24145314, 24164830, 26358756	482
VSTM1	unknown	PBio	PMID: 22960280, 23436183, 24205237, 25351446, 25887911, 26760041	147
VWA2	unknown	PBio	PMID: 14506275, 18434322, 21385852, 23443151, 23960233, 26121272	148
WEE1	unknown	Public_db	PMID: 19903823, 25088202, 26598692, 26881506, 27042682	484
WIPF1	AR	Public_db	PMID: 23765059, 26029204, 26453379	485
XIAP	XLD_XLR	Public_db	MySQL; PMID: 22365665, 25744037, 26953272	486
YBX1	unknown	Public_db	PMID: 27042682	487
YWHAZ	unknown	Both	PMID: 25894827, 27042682	488
ZAP70	AD_AR	Public_db	PMID: 18219311, 23494777, 23765059, 24164480, 26454313	489
ZBTB24	AR	Public_db	MySQL; PMID: 23486536, 23765059, 26851945, 27098601	490

[0613] Table 6 is a comprehensive list of 419 exemplary genes (referred to herein as ‘PML-419 genes’ or ‘PML-419 gene list’) interrogated in the present study, along with information related to the inheritance pattern assumed for analysis and the reason for inclusion of the gene. Gene sources for Table 6 (column heading ‘Gene_Source’): 1) nominated on the basis of being linked to immune deficiency, as curated from public databases (indicated by ‘Public_db’) such as PubMed and ClinVar, 2) PBio CNV-identified genes (‘PBio’, see Table 6 column heading ‘Gene_

Source’) from a genome-wide array CGH gene discovery study of 71 PML cases, or 3) curated from public databases and identified in PBio’s PML gene discovery study (indicated by ‘Both’). A genetic predisposition to PML on the basis of the host’s genome was proposed; that is, germline genetic variant(s) in the PML patient’s genome, rather than genetic variants that are present in the JC virus, are the cause of the patient’s PML (Hatchwell, Front Immunol., 6:216 (2015). Details on the source of the genes in the PML-419 gene list can be found in the following immunodeficiency

and immune-related gene sources: Durandy et al., *Nat Rev Immunol.*, 13(7):519-33 (2013); Milner et al., *Nat Rev Immunol.*, 13(9):635-48 (2013); Paciolla et al., *Genes Immun.*, 16(4):239-46 (2015); Hatchwell, *Front Immunol.*, 6:216 (2015); Thijssen et al., *Nat Commun.*, 6:7870 (2015); Chinn et al., *Immunol Allergy Clin North Am.*, 35(4):671-94 (2015); Zhou et al., *Nat Genet.*, 48(1):67-73 (2015); Navabi et al., *Allergy Asthma Clin Immunol.*, 12:27 (2016); and Tsujita et al., *J Allergy Clin Immunol.* (2016). MySQL genes are derived from the ClinVar database. ClinVar was searched using the terms “immune deficiency” and “immunodeficiency.” Entries that described large genomic rearrangements, containing multiple genes, were excluded. A non-redundant list of 125 genes was compiled by combining the output of the two searches and deposited into a MySQL database. NOTE: A subset of these genes are not flagged as ‘MySQL’ if they appeared in one or more of the immune gene review papers noted above. van der Kolk et al., *Ann Clin Transl Neurol.*; 3(3):226-32 (2016) was the source of known BAG3 PML gene (see below) and 28 candidate PML genes on the basis of connection to JCV. Van der Kolk et al., cite

a method as follows: “the latter was performed by searching for JCV in NCBI, and selecting for genes in humans.” This yielded 30 human genes, 5 of which overlapped with the PML gene list and 2 genes (HLA-DQB1, HLA-DRB1) were excluded because HLA loci are difficult to interpret. The genes ADA, BAG3, BTK, CD40LG, DOCK8, STAT1, WAS, and WIPF1 were derived from Hatchwell, *Front Immunol.*, 6:216 (2015) (see Table 1 for primary references); van der Kolk et al., *Ann Clin Transl Neurol.*, 3(3):226-32 (2016); and Zerbe et al., *Clin Infect Dis.*, 62(8):986-94 (2016). PBio genes are based on CNV studies and a subset overlap the immune review gene lists (annotated as ‘Both’ in column heading ‘Gene_Source’). Tier 1 genes were used as potential solutions for PML cases. Determination of Autosomal Dominant (AD), Autosomal Recessive (AR), X-linked dominant (XLD), or X-linked (XLR) disease model for each gene was derived from the immunodeficiency review papers and/or OMIM annotations. Entries marked ‘association’ denotes variants were found to be associated with an immune-related condition; ‘unknown’ denotes no evidence reported in the literature for an AD or AR model.

TABLE 7

Potential cause of PML in each patient in the study								
Sample_ID	Ethnicity	Gender	Primary Disease	RefSeq Gene Symbol	Variant_Type	Variant Frequency Details (Ethnic specific)	Frequency (Reciprocal) (Ethnic specific)	SEQ_ID
MVGS1116-8a	EUR	F	MS (NZ Rx)	DOCK8	SNV hom:SNV het	0.499:0.00447	1 in 1,792	1147:1148
MVGS1359	EUR	F	MS (NZ Rx)	IL17F	SNV het	0.00024	1 in 4,170	1114
MVGS1368	EUR	F	MS (NZ Rx)	IDO2	SNV hom:SNV het	0.508:0.065	1 in 121	1125:1126
MVGS540-374b	EUR	M	MS (NZ Rx)	SHARPIN	SNV hom	0.00217	1 in 461	1142
MVGS540-393b	EUR	F	MS (NZ Rx)	DOCK8	SNV hom:SNV het:SNV het	0.499:0.00153:0.194	1 in 5,246	1147:1154:1152
MVGS694-6a	EUR	F	Other	CHD7	SNV het	0.00028	1 in 3,528	1135
MVGS811-13a	EUR	M	HIV	PIK3CD_ PIK3CD-AS1	CNV hom	novel	0	2
MVGS995-4a	EUR	M	MS (NZ Rx)	EPG5	SNV hom:SNV het	0.495:0.000251	1 in 32,224	1279:1273
PML01	EUR	F	HIV	ITSN2	SNV hom	0.00183	1 in 547	1028
PML02	EUR	M	Other	IKBKKB	SNV het	novel	0	1127
PML03	EUR	F	MS (NZ Rx)	FPR2	CNV hom	2.23E-06	1 in 448,833	140
PML04	EUR	M	HIV	unsolved	n/a	n/a	n/a	n/a
PML05	LAT	M	HIV	TBK1	SNV het	novel	0	1203
PML06	AFR	M	HIV	TICAM1	SNV het	0.000777	1 in 1,287	1289
PML09	EUR	M	HIV	LIG4	SNV:SNV comp het	0.00399:0.287	1 in 3497	1221:1222
PML10	EUR	F	HIV	TNFRSF11A	SNV het	novel	0	1287
PML12	LAT	F	HIV	BLM	SNV hom	0.000874	1 in 1,144	1235
PML13	AFR	M	HIV	PLCG2	SNV:SNV comp het	0.00167:0.0187	1 in 128,105	1261:1263
PML14	EUR	M	HIV	PLCG2	SNV:SNV comp het	0.00998:0.0159	1 in 25,259	1261:1263
PML15	LAT	M	HIV	NOD2	SNV het	novel	0	1255
PML16	AFR	F	HIV	TNFRSF11A	SNV het	novel	0	1287
PML17	EUR	M	HIV	ZAP70	SNV het	0.00009	1 in 11,110	1035
PML18	EUR	M	HIV	unsolved	n/a	n/a	n/a	n/a
PML19	AFR	M	HIV	ATM	SNV:SNV comp het	0.0479:novel	0	1193:1194
PML20	AFR	M	HIV	NFKB1	SNV het	0.00173	1 in 577	1069
PML21	EUR	M	HIV	ZAP70	SNV het	0.0000602	1 in 16,623	1034
PML22	EUR	M	HIV	unsolved	n/a	n/a	n/a	n/a
PML23	EUR	F	HIV	DCLRE1C	SNV hom	novel	0	1167
PML25	EUR	F	HIV	PLCG2	SNV het	0.000150	1 in 6,672	1259
PML26	EUR	M	HIV	TRAFD1	SNV hom	0.000689	1 in 1,451	1208
PML27	EUR	M	HIV	TAP2	SNV hom	0.00837	1 in 120	1101
PML28	EUR	F	MS (NZ Rx)	TRPM2	SNV hom	novel	0	1311
PML29	AFR	M	HIV	KCTD7_ RABGEF1	CNV hom	0.000387	1 in 2,584	65
PML30	EUR	M	HIV	TNFRSF11A	SNV het	novel	0	1287
PML31	AFR	F	HIV	DDX58	SNV het	0.000779	1 in 1,283	1157
PML32	EUR	M	HIV	unsolved	n/a	n/a	n/a	n/a
PML33	EUR	M	HIV	TNFRSF11A	SNV het	novel	0	1287
PML35	EUR	F	HIV	TNFRSF11A	SNV het	novel	0	1287
PML36	AFR	F	HIV	TCIRG1	SNV het	0.002134	1 in 469	1184

TABLE 7-continued

Potential cause of PML in each patient in the study								
Sample_ID	Ethnicity	Gender	Primary Disease	RefSeq Gene Symbol	Variant_Type	Variant Frequency Details (Ethnic specific)	Frequency (Reciprocal) (Ethnic specific)	SEQ_ID
PML37	AFR	M	HIV	GATA2	SNV het	novel	0	1056
PML38	EUR	M	HIV	MALL	CNV hom	3.95E-06	1 in 253,036	26
PML39	AFR	M	HIV	unsolved	n/a	n/a	n/a	n/a
PML40	LAT	F	HIV	PNPT1	SNV hom	novel	0	1032
PML41	AFR	M	HIV	ZAP70	SNV het	novel	0	1036
PML43	EUR	M	HIV	PTPRC	SNV hom	novel	0	1020
PML44	EUR	M	HIV	TNFRSF11A	SNV het	novel	0	1287
PML45	EUR	F	Other	CARD11	SNV het	0.0024	1 in 417	1123
PML46	LAT	M	HIV	EPG5	SNV:SNV comp het	0.0123:0.436	1 in 745	1278:1279
PML48	EUR	M	HIV	SMAD4	SNV het	0.000901	1 in 11,100	1284
PML49	EUR	M	HIV	STIM1	SNV het	novel	0	1174
PML50	AFR	M	HIV	NOD2	SNV het	novel	0	1256
PML51	EUR	M	HIV	TICAM1	SNV het	0.00265	1 in 377	1289
PML52	EUR	F	Other	unsolved	n/a	n/a	n/a	n/a
PML53	EUR	M	Other	GFI1	SNV het	0.00003	1 in 32,635	1011
PML54	EUR	F	HIV	TNFRSF11A	SNV het	novel	0	1287
PML55	EUR	F	HIV	RTEL1	SNV het	0.00326	1 in 307	1299
PML56	EUR	M	HIV	TNFRSF11A	SNV het	novel	0	1287
PML57	EUR	F	Other	TRAF3	SNV het	0.00093	1 in 1,075	1229
PML58	AFR	M	HIV	DOCK8	SNV:SNV comp het	0.0575:0.478	1 in 146	1146:1147
PML59	AFR	M	HIV	IFIH1	SNV het	0.00281	1 in 356	1040
PML60	EUR	M	HIV	unsolved	n/a	n/a	n/a	n/a
PML61	AFR	F	HIV	TNFRSF11A	SNV het	novel	0	1287
PML62	AFR	F	HIV	unsolved	n/a	n/a	n/a	n/a
PML63	AFR	M	HIV	PLCG2	SNV het	0.00195	1 in 514	1260
PML64	AFR	M	HIV	PIK3R1	SNV het	novel	0	1077
PML65	AFR	M	HIV	ITSN2	CNV hom	0.00313	1 in 319	14
PML66	AFR	M	HIV	unsolved	n/a	n/a	n/a	n/a
PML67	EUR	F	MS (NZ Rx)	unsolved (CNV data, no WES data)	n/a	n/a	n/a	n/a
PML68	EUR	F	MS (NZ Rx)	LRBA	SNV hom	0.00162	1 in 618	1073
PML69	EUR	M	Other	EGR1_ETF1	CNV hom	0.001	1 in 1,005	45
PML72	AFR	F	HIV	NOD2	SNV het	0.004036	1 in 248	1252

[0614] Table 7 contains a single genetic solution/explanation that is the potential cause of PML in each patient in the study (71 cases were assessed with genome-wide array CGH and 71 were also assessed by whole exome sequencing), with the exception of 19 ‘unsolved’ cases. Solutions are based on a combination of CNV and SNV variants, connected by SEQ IDs to tables 1, 4 and 5. For homozygous or compound heterozygous variant solutions, expected population frequencies were calculated as follows:

[0615] Expected population frequency for variant a (freq p) and variant b (freq q)=pq/4.

[0616] For example, PML09 has 2 variants, SEQID 1221 and 1222, with individual frequencies in the normal population of 0.00399, 0.287. The expected frequency in an ethnically-matched normal population for this combination is $(0.00399 \times 0.287) \times 0.25 = 0.000286283 = 1/3,497$.

[0617] The Primary_Disease identifiers in Table 7 are: HIV, infection with human immunodeficiency virus; MS (NZ Rx), multiple sclerosis treated with natalizumab; Other, which includes a variety of disorders/conditions (MVGS694-6a had aplastic anemia, PML02 and PML52 had lymphoma, PML45 and PML 57 had chronic lymphocytic leukemia, PML53 had sarcoidosis, and PML69 is a kidney transplant patient who was on belatacept).

[0618] Solutions were considered on the basis of presence of rare variants (CNVs and/or SNVs) in or near genes that are listed in Table 6. Both autosomal recessive (AR) and autosomal dominant (AD) disease models comprise this set of solutions, based on finding homozygous SNVs, homozy-

gous CNVs, compound heterozygous SNVs, or heterozygous SNVs. Nine PML cases in Table 7 were considered ‘unsolved’ on the basis of analyzing both CNV and SNV data, and one case (PML67) was assessed for CNVs only since WES data were unavailable. In some instances, a case was considered unsolved for a best solution (Table 7) but alternate solutions were reported in Table 8 (see below).

[0619] For PML cases that had more than one potential solution. In these instances, the ‘best’ solution (Table 7) was determined on the basis of rarity of the genetic variant(s) and the relative strength of the biology for the PML-419 genes (Table 6). Alternate solutions are reported in Table 8. For example, for PML case MVGS1116-8a, three solutions were found, which impacted genes DOCK8, HIVEP2, and RNF168. In this example, DOCK8 compound heterozygous SNVs (Table 7, SNV hom:SNV het) were selected as the best solution because DOCK8 is a known PML gene. In another example, PML case MVGS1359 has IL17F (het SNV) listed as the best solution in Table 7 because it is rarer than alternate solutions for the ATR and STXBP2 genes.

[0620] While some PML patients may have multiple genes/variants causing and/or contributing to their PML, in many PML patients only a single gene will be the primary cause analogous to patients diagnosed with primary immunodeficiency disorders. In addition to the alternate solutions reported in Table 8, which are based on SNV genetic findings only, additional alternate solutions based on CNV genetic findings are reported in Table 1.

TABLE 8

Alternate genetic solutions/explanations as the potential cause of PML in the study					
Sample ID	RefSeq Gene Symbol	Variant_Type	Variant Frequency Details (Ethnic specific)	Frequency (Reciprocal) (Ethnic specific)	SEQ ID
MVGS1116-8a	HIVEP2	SNV het	novel	0	1118
MVGS1116-8a	RNF168	SNV hom:SNV het	0.469:0.00818	1 in 1,041	1063:1066
MVGS1359	ATR	SNV het	0.00393	1 in 254	1058
MVGS1359	STXBP2	SNV het	0.00501	1 in 199	1291
MVGS540-374b	MKL1	CNV hom	3.99E-08	1 in 25,081,515	157
MVGS540-393b	PRKDC	SNV het	0.00097	1 in 1,031	1130
MVGS811-13a	CLCN7	SNV het	0.00028	1 in 3,571	1239
MVGS995-4a	KAT6B	SNV het	0.00003	1 in 33,357	1169
MVGS995-4a	PRF1	SNV het	0.00243	1 in 412	1168
PML03	CDKN1B	SNV het	0.00003	1 in 32,209	1200
PML05	ATR	SNV het	novel	0	1061
PML05	NFKB1	SNV het	0.00501	1 in 200	1070
PML06	CHD7	SNV het	0.00797	1 in 125	1136
PML06	DOCK8	SNV hom:SNV het	0.478:0.0313	1 in 267	1147:1152
PML09	RIPK3	SNV het	0.00398	1 in 251	1227
PML10	JUN	SNV het	0.00103	1 in 968	1009
PML10	RAG1	SNV het	0.00039	1 in 2,566	1179
PML12	CARD11	SNV het	novel	0	1122
PML12	PRKDC	SNV het	novel	0	1128
PML13	DOCK8	SNV hom:SNV het	0.478:0.0313	1 in 267	1147:1152
PML13	IRAK4	SNV het	novel	0	1202
PML13	PIK3CD	SNV het	0.00679	1 in 147	1000
PML14	NBN	SNV het	0.0039	1 in 256	1138
PML14	NFKB1	SNV het	novel	0	1071
PML15	ASH1L	SNV:SNV comp het	novel:0.0019	0	1016:1017
PML15	CHD7	SNV het	0.00176	1 in 568	1133
PML15	HIVEP2	SNV het	novel	0	1116
PML15	STIM1	SNV het	0.00587	1 in 170	1175
PML16	TBK1	SNV het	novel	0	1204
PML16	TLR3	SNV het	0.00136	1 in 738	1076
PML17	APOL1	SNV het	0.0021	1 in 475	1327
PML18	PKHD1	SNV hom:SNV het	0.498:0.0471	1 in 171	1104:1107
PML19	DOCK8	SNV:SNV comp het	0.0575:0.478	1 in 146	1146:1147
PML19	IFIH1	SNV het	0.00444	1 in 225	1041
PML20	JUN	SNV het	0.00535	1 in 187	1010
PML21	PRKCH	SNV het	novel	0	1228
PML21	PSTPIP1	SNV het	0.00093	1 in 1,074	1232
PML21	RAG2	SNV het	novel	0	1182
PML22	RIPK3	SNV hom	0.00309	1 in 324	1226
PML22	VPS45	SNV het	0.00114	1 in 878	1014
PML23	NOD2	SNV het	novel	0	1251
PML23	RAG1	SNV het	0.00003	1 in 33,317	1180
PML28	PKHD1	SNV hom:SNV het	0.498:0.0471	1 in 171	1104:1107
PML28	TNFRSF13B	SNV het	0.00929	1 in 108	1267
PML30	RTKL1	SNV het	0.000124	1 in 8,068	1300
PML31	AP3B1	SNV het	novel	0	1084
PML31	PRKDC	SNV het	novel	0	1129
PML33	STIM2	SNV het	0.00003	1 in 32,688	1068
PML33	TLR3	SNV hom:SNV het	0.413:0.00435	1 in 2,227	1075:1074
PML33	TLR4	SNV hom:SNV hom	0.00283:0.00285	1 in 354	1161:1160
PML35	PRKCB	SNV het	0.00276	1 in 362	1247
PML36	NOD2	SNV het	0.00871	1 in 115	1254
PML36	PIK3CD	SNV het	0.00679	1 in 147	1000
PML37	AP3B1	SNV het	novel	0	1080
PML37	ATR	SNV het	0.00038	1 in 2,601	1059
PML37	WEE1	SNV het	0.00825	1 in 121	1177
PML38	MYD88	SNV het	novel	0	1051
PML40	MCEE	SNV hom	0.01	1 in 100	1033
PML41	AP3B1	SNV het	0.00173	1 in 577	1082
PML41	CHD7	SNV het	novel	0	1137
PML41	DOCK8	SNV:SNV comp het	0.0575:0.478	1 in 146	1146:1147
PML41	POLE	SNV hom	0.00019	1 in 5,203	1219
PML41	RNF168	SNV:SNV comp het	0.412:novel	0	1063:1062
PML43	DOCK8	SNV hom:SNV het	0.499:novel	0	1147:1150
PML44	DCLRE1C	SNV hom:SNV hom	0.0287:0.00575	1 in 174	1166:1165
PML44	GFI1	SNV het	0.00708	1 in 141	1012
PML45	POLA1	SNV het	novel	0	1328
PML46	AP3B1	SNV het	0.00587	1 in 170	1082
PML46	IL21R	SNV het	0.00573	1 in 175	1248
PML46	PRKDC	SNV het	0.00017	1 in 5,781	1131
PML48	TNFRSF11A	SNV het	0.00233	1 in 429	1286

TABLE 8-continued

Alternate genetic solutions/explanations as the potential cause of PML in the study					
Sample ID	RefSeq Gene Symbol	Variant_Type	Variant Frequency Details (Ethnic specific)	Frequency (Reciprocal) (Ethnic specific)	SEQ ID
PML49	DCLRE1C	SNV hom:SNV hom	0.00575:0.0287	1 in 174	1166:1165
PML49	PTEN	SNV het	novel	0	1171
PML49	RIPK1	SNV het	0.00090	1 in 1,112	1090
PML50	AP3B1	SNV het	0.00387	1 in 259	1078
PML50	PIAS2	SNV het	0.00357	1 in 280	1283
PML50	STXBP2	SNV het	0.00038	1 in 2,598	1290
PML52	GFI1	SNV het	0.00708	1 in 141	1012
PML53	IL1B	SNV het	novel	0	1037
PML53	STXBP2	SNV het	0.00501	1 in 199	1291
PML54	EPG5	SNV:SNV comp het	0.0638:0.495	1 in 127	1278:1279
PML54	IFNGR2	SNV het	0.00009	1 in 11,096	1304
PML54	RAG1	SNV het	0.00003	1 in 33,352	1178
PML54	RAG2	SNV het	novel	0	1183
PML57	PIAS1	SNV het	novel	0	1231
PML57	PKHD1	SNV hom:SNV het	0.498:0.0471	1 in 171	1104:1107
PML57	SKIV2L	SNV hom:SNV hom:SNV het	0.157:0.214:0.0471	1 in 538	1098:1100:1099
PML58	GFI1	SNV het	0.00144	1 in 693	1012
PML59	IFNLR1	SNV het	novel	0	1002
PML59	NOD2	SNV het	0.00404	1 in 248	1252
PML59	NRIP1	SNV hom	0.00711	1 in 141	1301
PML59	RAD51	SNV het	0.00865	1 in 116	1230
PML60	MAPK3	SNV het	novel	0	1250
PML60	TP53	SNV het	0.00048	1 in 2,085	1266
PML61	GATA2	SNV het	0.00024	1 in 4,139	1057
PML61	PTPRC	SNV hom	novel	0	1019
PML61	TNFRSF8	SNV het	novel	0	1001
PML62	PRKCD	SNV het	novel	0	1054
PML63	HTR2A	SNV hom	0.00519	1 in 193	1220
PML63	MAPK3	SNV het	0.00193	1 in 518	1249
PML64	PLCG2	SNV het	0.00044	1 in 2,276	1264
PML64	WEE1	SNV het	novel	0	1176
PML65	IRAK4	SNV het	0.00118	1 in 850	1201
PML66	PIK3CD	SNV het	0.00679	1 in 147	1000
PML68	RAG1	SNV het	0.00586	1 in 171	1181
PML72	CARD11	SNV het	0.00242	1 in 413	1121
PML72	HIVEP1	SNV hom	0.00164	1 in 610	1092
PML72	IFIH1	SNV het	0.00843	1 in 119	1043

[0621] Table 8 contains analogous information to Table 7, with the exception that Ethnicity, Gender and Primary Disease are not repeated. Table 8 contains alternate genetic solutions/explanations as the potential cause of PML for the patients in the study (71 cases were assessed with genome-wide array CGH and 70 were also assessed by whole exome sequencing). Solutions in Table 8 are also case-level and represent secondary, alternative solutions for the cases listed (using the same criteria used to identify potential solutions reported in Table 7). In other words, for some individuals,

more than one reasonable solution was identified and, while those in Table 7 are considered the most likely, those in Table 8 are also potential solutions. It can be appreciated by those skilled in the art that further data on new PML cases, patients with genetic-based immunodeficiency disorders, or functional studies on a given gene may support selection of a Table 8 solution as the ‘best’ single solution (i.e., a current Table 7 solution could be considered instead as a Table 8 solution, and vice versa).

TABLE 9

Pairs of SNVs impacting the same gene								
Sample ID	RefSeq Gene Symbol	Variant Type	Chromosome	Position	Ref Allele	Alt Allele	Amino Acid Change	Variant Frequency Details (Ethnic specific) SEQ ID NO
MVGS1359	TTC7A	SNV het	2	47273468	A	G	K252R	0.00684 1030
MVGS1359	TTC7A	SNV het	2	47277182	T	C	S318P	0.00683 1031
MVGS1368	RNF168	SNV het	3	196199204	G	T	P401Q	0.46947 1063
MVGS1368	RNF168	SNV het	3	196210764	T	C	n/a	0.00003 1065
MVGS1368	TLR4	SNV het	9	120475302	A	G	D259G	0.10251 1160
MVGS1368	TLR4	SNV het	9	120475602	C	T	T359I	0.10560 1161
MVGS811-13a	HIVEP1	SNV het	6	12121113	C	T	P362L	0.00024 1091
MVGS811-13a	HIVEP1	SNV het	6	12123538	G	T	K1170N	0.08730 1093

TABLE 9-continued

Pairs of SNVs impacting the same gene									
Sample ID	RefSeq Gene Symbol	Variant Type	Chromosome	Position	Ref Allele	Alt Allele	Amino Acid Change	Variant Frequency Details (Ethnic specific)	SEQ ID NO
MVGS995-4a	EEA1	SNV het	12	93196332	C	T	E840K	0.01949	1206
MVGS995-4a	EEA1	SNV het	12	93205148	T	G	E702D	0.00003	1207
PML02	RBFOX1	SNV het	16	7759119	G	A	G326S	0.00504	1245
PML02	RBFOX1	SNV het	16	7759496	C	T	P401S	novel	1246
PML04	POLE	SNV het	12	133220526	T	C	N1369S	0.22363	1213
PML04	POLE	SNV het	12	133237658	T	G	Q766P	novel	1215
PML05	TLR4	SNV het	9	120475302	A	G	D259G	0.04628	1160
PML05	TLR4	SNV het	9	120475602	C	T	T359I	0.04180	1161
PML05	POLE	SNV het	12	133220526	T	C	N1369S	0.12669	1213
PML05	POLE	SNV het	12	133252406	C	A	A121S	novel	1217
PML10	TLR4	SNV het	9	120475302	A	G	D259G	0.10251	1160
PML10	TLR4	SNV het	9	120475602	C	T	T359I	0.10560	1161
PML12	IDO2	SNV het	8	39840234	A	G	I127V	0.38971	1124
PML12	IDO2	SNV het	8	39862881	C	T	R235W	0.50282	1125
PML12	IDO2	SNV het	8	39862893	T	A	S239T	0.02384	1126
PML13	STX11	SNV het	6	144508353	G	A	V197M	novel	1119
PML13	STX11	SNV het	6	144508563	G	A	V267M	0.00202	1120
PML13	DCLRE1C	SNV het	10	14974905	T	C	H123R	0.16298	1165
PML13	DCLRE1C	SNV het	10	14976727	G	C	P171R	0.22295	1166
PML13	EPG5	SNV het	18	43497710	A	G	V1058A	0.42740	1279
PML13	EPG5	SNV het	18	43531186	C	T	S424N	0.00600	1282
PML14	ATM	SNV het	11	108117787	C	T	S333F	0.00280	1188
PML14	ATM	SNV het	11	108175462	G	A	D1853N	0.24654	1193
PML14	TRPM2	SNV het	21	45815425	C	G	I621M	novel	1313
PML14	TRPM2	SNV het	21	45845699	G	A	V1242M	0.00537	1321
PML16	TLR3	SNV het	4	187004074	C	T	L135F	0.12378	1075
PML16	TLR3	SNV het	4	187005854	A	C	I571L	0.00136	1076
PML16	HIVEP1	SNV het	6	12121113	C	T	P362L	0.07856	1091
PML16	HIVEP1	SNV het	6	12162068	C	T	S160F	0.01979	1096
PML16	PKHD1	SNV het	6	51483961	T	C	Q4048R	0.50029	1104
PML16	PKHD1	SNV het	6	51747943	T	A	D2433V	0.07153	1112
PML16	POLE	SNV het	12	133209020	G	C	Q2044E	novel	1212
PML16	POLE	SNV het	12	133220526	T	C	N1369S	0.24889	1213
PML17	RNF168	SNV het	3	196199204	G	T	P401Q	0.46947	1063
PML17	RNF168	SNV het	3	196210704	G	A	P206L	0.00003	1064
PML17	HIVEP1	SNV het	6	12123538	G	T	K1170N	0.08730	1093
PML17	HIVEP1	SNV het	6	12125232	C	T	S1735F	0.00027	1095
PML17	PKHD1	SNV het	6	51483961	T	C	Q4048R	0.49837	1104
PML17	PKHD1	SNV het	6	51497503	C	A	R3842L	0.04707	1107
PML17	DCLRE1C	SNV het	10	14974905	T	C	H123R	0.27332	1165
PML17	DCLRE1C	SNV het	10	14976727	G	C	P171R	0.13896	1166
PML17	ATM	SNV het	11	108119823	T	C	V410A	0.00643	1189
PML17	ATM	SNV het	11	108175462	G	A	D1853N	0.24654	1193
PML17	EPG5	SNV het	18	43464763	C	T	G1708D	0.00013	1274
PML17	EPG5	SNV het	18	43497710	A	G	V1058A	0.49513	1279
PML18	TLR4	SNV het	9	120475302	A	G	D259G	0.10251	1160
PML18	TLR4	SNV het	9	120475602	C	T	T359I	0.10560	1161
PML20	AK2	SNV het	1	33476435	C	A	n/a	novel	1003
PML20	AK2	SNV het	1	33478900	T	A	Y159F	0.04954	1004
PML20	HIVEP1	SNV het	6	12124215	C	T	P1396L	0.06774	1094
PML20	HIVEP1	SNV het	6	12163657	C	T	P2374S	0.06733	1097
PML20	KANK1	SNV het	9	711359	C	T	S198F	0.11985	1155
PML20	KANK1	SNV het	9	713132	G	T	G631V	0.00136	1156
PML21	DOCK8	SNV het	9	286593	C	A	P29T	0.49889	1147
PML21	DOCK8	SNV het	9	286593	C	A	P29T	0.49889	1147
PML21	DOCK8	SNV het	9	312134	G	A	E169K	0.06358	1149
PML21	DOCK8	SNV het	9	312134	G	A	E169K	0.06358	1149
PML21	TLR4	SNV het	9	120475302	A	G	D259G	0.10251	1160
PML21	TLR4	SNV het	9	120475302	A	G	D259G	0.10251	1160
PML21	TLR4	SNV het	9	120475602	C	T	T359I	0.10560	1161
PML21	TLR4	SNV het	9	120475602	C	T	T359I	0.10560	1161
PML21	ATM	SNV het	11	108138003	T	C	F858L	0.02864	1191
PML21	ATM	SNV het	11	108138003	T	C	F858L	0.02864	1191
PML21	ATM	SNV het	11	108143456	C	G	P1054R	0.05069	1192
PML21	ATM	SNV het	11	108143456	C	G	P1054R	0.05069	1192
PML21	TRPM2	SNV het	21	45786650	C	T	S146F	0.00072	1305
PML21	TRPM2	SNV het	21	45786650	C	T	S146F	0.00072	1305
PML21	TRPM2	SNV het	21	45820196	C	T	R735C	0.10374	1314
PML21	TRPM2	SNV het	21	45820196	C	T	R735C	0.10374	1314

TABLE 9-continued

Pairs of SNVs impacting the same gene									
Sample ID	RefSeq Gene Symbol	Variant Type	Chromosome	Position	Ref Allele	Alt Allele	Amino Acid Change	Variant Frequency Details (Ethnic specific)	SEQ ID NO
PML22	SKIV2L	SNV het	6	31928306	A	G	Q151R	0.15759	1098
PML22	SKIV2L	SNV het	6	31935750	G	A	V724M	0.04718	1099
PML22	SKIV2L	SNV het	6	31936679	C	T	A1071V	0.21419	1100
PML22	DOCK8	SNV het	9	286593	C	A	P29T	0.49889	1147
PML22	DOCK8	SNV het	9	304628	G	A	R151Q	0.00447	1148
PML22	GDPD4	SNV het	11	76954833	G	A	H383Y	0.44867	1186
PML22	GDPD4	SNV het	11	76979511	A	G	I233T	0.00504	1187
PML22	ATM	SNV het	11	108117787	C	T	S333F	0.00280	1188
PML22	ATM	SNV het	11	108175462	G	A	D1853N	0.24654	1193
PML22	BLM	SNV het	15	91306241	G	A	R643H	0.00799	1233
PML22	BLM	SNV het	15	91341543	A	C	N1112H	novel	1238
PML23	PKHD1	SNV het	6	51483961	T	C	Q4048R	0.49837	1104
PML23	PKHD1	SNV het	6	51497503	C	A	R3842L	0.04707	1107
PML23	SHARPIN	SNV het	8	145154222	G	A	P294S	0.08789	1142
PML23	SHARPIN	SNV het	8	145154257	C	G	S282T	0.14880	1144
PML23	DOCK8	SNV het	9	286491	G	A	D63N	0.27362	1146
PML23	DOCK8	SNV het	9	334277	G	A	R325H	0.00015	1151
PML25	SKIV2L	SNV het	6	31928306	A	G	Q151R	0.15759	1098
PML25	SKIV2L	SNV het	6	31935750	G	A	V724M	0.04718	1099
PML25	SKIV2L	SNV het	6	31936679	C	T	A1071V	0.21419	1100
PML25	PKHD1	SNV het	6	51483961	T	C	Q4048R	0.49837	1104
PML25	PKHD1	SNV het	6	51524409	G	T	S3505R	0.02049	1109
PML25	EPG5	SNV het	18	43445601	T	G	I174L	novel	1270
PML25	EPG5	SNV het	18	43531186	C	T	S424N	0.02391	1282
PML27	LYST	SNV het	1	235897907	C	T	G2804D	0.00114	1024
PML27	LYST	SNV het	1	235909815	A	T	F165Y	0.00102	1025
PML27	EPG5	SNV het	18	43445601	T	G	I174L	novel	1270
PML27	EPG5	SNV het	18	43497710	A	G	V1058A	0.49513	1279
PML29	LIG1	SNV het	19	48631258	G	A	T546I	0.07515	1292
PML29	LIG1	SNV het	19	48639022	T	C	M412V	0.05385	1293
PML30	DCLRE1C	SNV het	10	14974905	T	C	H123R	0.27332	1165
PML30	DCLRE1C	SNV het	10	14976727	G	C	P171R	0.13896	1166
PML30	ATM	SNV het	11	108138003	T	C	F858L	0.02864	1191
PML30	ATM	SNV het	11	108143456	C	G	P1054R	0.05069	1192
PML30	ATM	SNV het	11	108186610	G	A	G2023R	0.00465	1195
PML31	LYST	SNV het	1	235897907	C	T	G2804D	0.23000	1024
PML31	LYST	SNV het	1	235909815	A	T	F165Y	0.15155	1025
PML31	PKHD1	SNV het	6	51483961	T	C	Q4048R	0.50029	1104
PML31	PKHD1	SNV het	6	51524339	C	G	E3529Q	0.07244	1108
PML31	PKHD1	SNV het	6	51747943	T	A	D2433V	0.07153	1112
PML31	PKHD1	SNV het	6	51798908	C	T	G2041S	0.00173	1113
PML32	PKHD1	SNV het	6	51483961	T	C	Q4048R	0.49837	1104
PML32	PKHD1	SNV het	6	51491885	G	A	Q3899*	novel	1106
PML32	EPG5	SNV het	18	43496539	G	A	S1083L	0.06375	1278
PML32	EPG5	SNV het	18	43497710	A	G	V1058A	0.49513	1279
PML32	EPG5	SNV het	18	43529551	C	T	V466M	0.00006	1281
PML33	AK2	SNV het	1	33476435	C	A	n/a	novel	1003
PML33	AK2	SNV het	1	33487007	C	T	S129N	0.01100	1005
PML33	EPG5	SNV het	18	43497710	A	G	V1058A	0.49513	1279
PML33	EPG5	SNV het	18	43523240	C	T	M610I	0.00066	1280
PML35	RNF168	SNV het	3	196199204	G	T	P401Q	0.46947	1063
PML35	RNF168	SNV het	3	196214320	C	T	E170K	0.00818	1066
PML36	IDO2	SNV het	8	39862881	C	T	R235W	0.46108	1125
PML36	IDO2	SNV het	8	39862893	T	A	S239T	0.01135	1126
PML39	LYST	SNV het	1	235897907	C	T	G2804D	0.23000	1024
PML39	LYST	SNV het	1	235909815	A	T	F165Y	0.15155	1025
PML39	NHEJ1	SNV het	2	219942026	T	A	Q181L	0.06324	1047
PML39	NHEJ1	SNV het	2	220023045	C	T	A14T	0.23543	1048
PML40	ATM	SNV het	11	108186631	A	G	I2030V	0.00173	1196
PML40	ATM	SNV het	11	108186631	A	G	I2030V	0.03446	1196
PML40	ATM	SNV het	11	108198384	C	G	L2330V	0.00035	1197
PML40	ATM	SNV het	11	108198384	C	G	L2330V	0.00491	1197
PML41	PKHD1	SNV het	6	51483961	T	C	Q4048R	0.50029	1104
PML41	PKHD1	SNV het	6	51497503	C	A	R3842L	0.00654	1107
PML41	IDO2	SNV het	8	39840234	A	G	I127V	0.06350	1124
PML41	IDO2	SNV het	8	39862881	C	T	R235W	0.46108	1125
PML45	VPS13B	SNV het	8	100791158	G	A	E2560K	0.00964	1140
PML45	VPS13B	SNV het	8	100865941	G	A	A3442T	novel	1141
PML48	EPG5	SNV het	18	43497710	A	G	V1058A	0.49513	1279

TABLE 9-continued

Pairs of SNVs impacting the same gene									
Sample ID	RefSeq Gene Symbol	Variant Type	Chromosome	Position	Ref Allele	Alt Allele	Amino Acid Change	Variant Frequency Details (Ethnic specific)	SEQ ID NO
PML48	EPG5	SNV het	18	43531186	C	T	S424N	0.02391	1282
PML51	TRPM2	SNV het	21	45826486	G	A	V914I	novel	1315
PML51	TRPM2	SNV het	21	45855099	C	T	R1300W	0.00021	1322
PML53	EPG5	SNV het	18	43445580	C	T	D181N	novel	1269
PML53	EPG5	SNV het	18	43497710	A	G	V1058A	0.49513	1279
PML56	TLR4	SNV het	9	120475302	A	G	D259G	0.10251	1160
PML56	TLR4	SNV het	9	120475602	C	T	T359I	0.10560	1161
PML56	DCLRE1C	SNV het	10	14974905	T	C	H123R	0.27332	1165
PML56	DCLRE1C	SNV het	10	14976727	G	C	P171R	0.13896	1166
PML57	CSF3R	SNV het	1	36932047	C	T	E359K	0.01706	1006
PML57	CSF3R	SNV het	1	36933715	A	G	Y113H	0.00087	1007
PML57	TLR4	SNV het	9	120475302	A	G	D259G	0.10251	1160
PML57	TLR4	SNV het	9	120475602	C	T	T359I	0.10560	1161
PML57	ATM	SNV het	11	108138003	T	C	F858L	0.02864	1191
PML57	ATM	SNV het	11	108143456	C	G	P1054R	0.05069	1192
PML57	ATM	SNV het	11	108175462	G	A	D1853N	0.24654	1193
PML58	DOCK8	SNV het	9	3992233	A	G	N1002D	0.19737	1153
PML58	DCLRE1C	SNV het	10	14974905	T	C	H123R	0.16298	1165
PML58	DCLRE1C	SNV het	10	14976727	G	C	P171R	0.22295	1166
PML58	DNMT3B	SNV het	20	31383307	G	A	G311S	0.00192	1296
PML58	DNMT3B	SNV het	20	31384614	G	T	G343V	novel	1297
PML59	LYST	SNV het	1	235897907	C	T	G2804D	0.23000	1024
PML59	LYST	SNV het	1	235897907	C	T	G2804D	0.23000	1024
PML59	LYST	SNV het	1	235909815	A	T	F165Y	0.15155	1025
PML59	LYST	SNV het	1	235909815	A	T	F165Y	0.15155	1025
PML59	LIG1	SNV het	19	48631258	G	A	T546I	0.07515	1292
PML59	LIG1	SNV het	19	48631258	G	A	T546I	0.07515	1292
PML59	LIG1	SNV het	19	48639022	T	C	M412V	0.05385	1293
PML59	LIG1	SNV het	19	48639022	T	C	M412V	0.05385	1293
PML60	DCLRE1C	SNV het	10	14974905	T	C	H123R	0.27332	1165
PML60	DCLRE1C	SNV het	10	14976727	G	C	P171R	0.13896	1166
PML60	POLE	SNV het	12	133202816	C	T	E2113K	0.04686	1211
PML60	POLE	SNV het	12	133220526	T	C	N1369S	0.22363	1213
PML62	TLR4	SNV het	9	120475302	A	G	D259G	0.13066	1160
PML62	TLR4	SNV het	9	120475602	C	T	T359I	0.02672	1161
PML63	HIVEP1	SNV het	6	12124215	C	T	P1396L	0.06774	1094
PML63	HIVEP1	SNV het	6	12163657	C	T	P2374S	0.06733	1097
PML63	PLCG2	SNV het	16	81942175	A	G	N571S	0.01870	1263
PML63	TRPM2	SNV het	21	45795833	G	T	V297L	0.00097	1306
PML63	TRPM2	SNV het	21	45815307	T	C	V582A	0.00724	1310
PML64	DNER	SNV het	2	230231632	C	T	D687N	0.00058	1049
PML64	DNER	SNV het	2	230450646	T	A	T259S	0.00692	1050
PML64	IDO2	SNV het	8	39862881	C	T	R235W	0.46108	1125
PML64	IDO2	SNV het	8	39862893	T	A	S239T	0.01135	1126
PML65	POLE	SNV het	12	133201381	T	A	I2228F	0.00232	1210
PML65	POLE	SNV het	12	133253971	C	T	R233Q	0.02037	1218
PML66	PKHD1	SNV het	6	51483961	T	C	Q4048R	0.50029	1104
PML66	PKHD1	SNV het	6	51612746	G	A	S3223L	0.00000	1110
PML66	PKHD1	SNV het	6	51712759	T	C	T2641A	0.04812	1111
PML66	EPG5	SNV het	18	43456296	C	T	R1985Q	0.07733	1271
PML66	EPG5	SNV het	18	43497710	A	G	V1058A	0.42740	1279
PML68	DCLRE1C	SNV het	10	14974905	T	C	H123R	0.27332	1165
PML68	DCLRE1C	SNV het	10	14976727	G	C	P171R	0.13896	1166
PML72	PSMB8	SNV het	6	32810794	T	A	T70S	0.04224	1102
PML72	PSMB8	SNV het	6	32811752	C	T	G8R	0.04845	1103
PML72	POLE	SNV het	12	133220526	T	C	N1369S	0.24889	1213
PML72	POLE	SNV het	12	133245026	G	A	P477S	0.02332	1216
PML72	RBFOX1	SNV het	16	7568296	C	T	P102S	0.00692	1242
PML72	RBFOX1	SNV het	16	7703891	A	G	T235A	novel	1243

[0622] Table 9 lists, for each case (in multiple rows), variants for which it was not possible, using the whole exome sequencing (WES) data available, to determine phase (i.e., whether two variants are in cis—on the same chromosome—or trans—on opposite chromosomes). Determining phase is an important consideration when dealing with disorders that are being evaluated on an autosomal recessive

(AR) basis. If two variants are known to be present but it is impossible to determine whether they are in cis or trans, then it is impossible to conclude that both gene copies are affected, as opposed to only one (albeit with 2 variants). This problem does not arise in the case of homozygous variants, for which it is obvious that the variants must be in trans (i.e.,

it is only an issue for non-identical variants). All genome coordinates are based on hg19 build.

[0623] In summary, Table 9 lists all unphased case-level compound heterozygous SNV solutions, which might represent further case-level solutions, were phasing to have been possible. Furthermore, it can be appreciated by those skilled in the art that unphased solutions reported in Table 9 (2 het SNVs per gene) or Table 10 (see below, which reports het SNVs in patients that also have a CNV reported in Table 1) can potentially cause or contribute to the patient's PML if follow up genetic analysis reveals the pair of variants are on different alleles (i.e., each gene copy impacted by a variant). Variants reported in Tables 1, 9, or 10 may also be found to be significantly deleterious on their own (e.g., in functional studies on patient-derived cells, animal models, etc.) and thus constitute an AD model solution (i.e., genes presently listed as 'AR' model in Table 6) may be causal or contributing to disease via an AD or AR model, like several genes already known to be AD or AR (Table 6, 'AD_AR' disease model).

TABLE 10

SNVs found in genes suspected of being impacted by acquired CNVs									
Sample_ID	RefSeq Gene Symbol	Variant Type	Chr	Position	Ref Allele	Alt Allele	Amino Acid Change	Variant Frequency Details (Ethnic specific)	SEQ ID NO
MVGS811-13a	NRIP1	SNV het	21	16338814	T	C	N567S	0.00060	1301
MVGS995-4a	VWA2	SNV het	10	116045796	G	A	V366M	0.02392	1173
PML01	PKHD1	SNV het	6	51497503	C	A	R3842L	0.04707	1107
PML01	PKHD1	SNV het	6	51483961	T	C	Q4048R	0.49837	1104
PML02	DUSP16	SNV het	12	12673965	G	A	T23M	0.00015	1199
PML39	SALL2	SNV het	14	22004996	G	T	S13R	0.00231	1225
PML51	JMY	SNV het	5	78596018	G	C	D524H	novel	1086
PML65	SALL2	SNV het	14	21992397	T	C	S347G	0.07709	1223

NOTE:

These are het SNVs that are potentially compound heterozygotes with a CNV on the allele. See text for description.

The DUSP16 SNV (chr12: 12673965) was in trans with a chr12 deletion of DUSP16 in this patient (PML02), whose primary diagnosis was lymphoma.

[0624] Table 10 is a list of all heterozygous SNVs that are potentially compound heterozygotes with a CNV on the allele. See text for a fuller explanation. All genome coordinates are based on hg19 build.

TABLE 11

Key that maps Sample_ID for the PML cases to the PML_Case ID numbers	
Sample_ID	PML_Case_ID
MVGS1116-8a	3006
MVGS1359	3117
MVGS1368	3118
MVGS540-374b	3005
MVGS540-393b	3004
MVGS694-6a	3007
MVGS811-13a	3009
MVGS995-4a	3010
PML01	3127
PML02	3126
PML03	3155
PML04	3156
PML05	3125
PML06	3124
PML09	3132
PML10	3157
PML12	3159

TABLE 11-continued

Key that maps Sample_ID for the PML cases to the PML_Case ID numbers	
Sample_ID	PML_Case_ID
PML13	3160
PML14	3161
PML15	3194
PML16	3163
PML17	3140
PML18	3141
PML19	3164
PML20	3143
PML21	3144
PML22	3145
PML23	3165
PML25	3166
PML26	3167
PML27	3168
PML28	3151
PML29	3152
PML30	3153
PML31	3154
PML32	3169
PML33	3170
PML35	3171
PML36	3172
PML37	3173

TABLE 11-continued

Key that maps Sample_ID for the PML cases to the PML_Case ID numbers	
Sample_ID	PML_Case_ID
PML38	3174
PML39	3175
PML40	3273
PML41	3177
PML43	3178
PML44	3179
PML45	3180
PML46	3196
PML48	3197
PML49	3183
PML50	3198
PML51	3185
PML52	3186
PML53	3187
PML54	3188
PML55	3189
PML56	3190
PML57	3191
PML58	3192
PML59	3193
PML60	3199

TABLE 11-continued

Key that maps Sample_ID for the PML cases to the PML_Case ID numbers	
Sample_ID	PML_Case_ID
PML61	3200
PML62	3201
PML63	3202
PML64	3203
PML65	3204
PML66	3205
PML67	3277
PML68	3278
PML69	3279
PML72	3282
PML70_control	3280
PML71_control	3281
PML73_control	3283
PML74_control	3284
PML75_control	3285
PML76_control	3286

[0625] Table 11 provides the Sample_ID and PML_Case_ID (experimental ID for CGH data) for 77 ‘PML cases’ (includes 6 non-PML HIV cases listed as controls).

TABLE 12

Non-redundant list of transcript variants that correspond to the set of genes that no CNV ‘solutions’ have been reported in the 71 PML cases			
RefSeq_Gene_Symbol	RefSeq_Accession_Number	mRNA_Description	SEQ ID
ACADM	NM_000016	<i>Homo sapiens</i> acyl-CoA dehydrogenase, C-4 to C-12 straight chain (ACADM), transcript variant 1, mRNA.	1500
ACADM	NM_001127328	<i>Homo sapiens</i> atypical chemokine receptor 1 (Duffy blood group) (ACKR1), transcript variant 1, mRNA.	1501
ACKR1	NM_002036	<i>Homo sapiens</i> atypical chemokine receptor 1 (Duffy blood group) (ACKR1), transcript variant 2, mRNA.	1502
ACKR1	NM_001122951	<i>Homo sapiens</i> atypical chemokine receptor 1 isoform a	1503
ACP5	NM_001611	<i>Homo sapiens</i> acid phosphatase 5, tartrate resistant (ACP5), transcript variant 4, mRNA.	1504
ACP5	NM_001111034	<i>Homo sapiens</i> acid phosphatase 5, tartrate resistant (ACP5), transcript variant 2, mRNA.	1505
ACP5	NM_001111035	<i>Homo sapiens</i> acid phosphatase 5, tartrate resistant (ACP5), transcript variant 1, mRNA.	1506
ACP5	NM_001111036	<i>Homo sapiens</i> acid phosphatase 5, tartrate resistant (ACP5), transcript variant 3, mRNA.	1507
ADAR	NM_001111	<i>Homo sapiens</i> adenosine deaminase, RNA-specific (ADAR), transcript variant 1, mRNA.	1508
ADAR	NM_015840	<i>Homo sapiens</i> adenosine deaminase, RNA-specific (ADAR), transcript variant 2, mRNA.	1509
ADAR	NM_015841	<i>Homo sapiens</i> adenosine deaminase, RNA-specific (ADAR), transcript variant 3, mRNA.	1510
ADAR	NM_001025107	<i>Homo sapiens</i> adenosine deaminase, RNA-specific (ADAR), transcript variant 4, mRNA.	1511
ADAR	NM_001193495	<i>Homo sapiens</i> adenosine deaminase, RNA-specific (ADAR), transcript variant 5, mRNA.	1512
ADK	NM_001202450	<i>Homo sapiens</i> adenosine kinase (ADK), transcript variant 4, mRNA.	1513
ADK	NM_006721	<i>Homo sapiens</i> adenosine kinase (ADK), transcript variant 2, mRNA.	1514
ADK	NM_001123	<i>Homo sapiens</i> adenosine kinase (ADK), transcript variant 1, mRNA.	1515
ADK	NM_001202449	<i>Homo sapiens</i> adenosine kinase (ADK), transcript variant 3, mRNA.	1516
AICDA	NM_020661	<i>Homo sapiens</i> activation-induced cytidine deaminase (AICDA), mRNA.	1517
AK2	NM_001199199	<i>Homo sapiens</i> adenylate kinase 2 (AK2), transcript variant 3, mRNA.	1518
AK2	NM_013411	<i>Homo sapiens</i> adenylate kinase 2 (AK2), transcript variant 2, mRNA.	1519
AK2	NM_001625	<i>Homo sapiens</i> adenylate kinase 2 (AK2), transcript variant 1, mRNA.	1520
ALG12	NM_024105	<i>Homo sapiens</i> ALG12, alpha-1,6-mannosyltransferase (ALG12), mRNA.	1521
ALPL	NM_000478	<i>Homo sapiens</i> alkaline phosphatase, liver/bone/kidney (ALPL), transcript variant 1, mRNA.	1522
ALPL	NM_001127501	<i>Homo sapiens</i> alkaline phosphatase, liver/bone/kidney (ALPL), transcript variant 2, mRNA.	1523
ALPL	NM_001177520	<i>Homo sapiens</i> alkaline phosphatase, liver/bone/kidney (ALPL), transcript variant 3, mRNA.	1524
AP3B1	NM_001271769	<i>Homo sapiens</i> adaptor related protein complex 3 beta 1 subunit (AP3B1), transcript variant 2, mRNA.	1525
AP3B1	NM_003664	<i>Homo sapiens</i> adaptor related protein complex 3 beta 1 subunit (AP3B1), transcript variant 1, mRNA.	1526
AP3B2	NM_004644	<i>Homo sapiens</i> adaptor-related protein complex 3, beta 2 subunit (AP3B2), transcript variant 2, mRNA.	1527
AP3D1	NM_003938	<i>Homo sapiens</i> adaptor-related protein complex 3, delta 1 subunit (AP3D1), transcript variant 2, mRNA.	1528
AP3D1	NM_001261826	<i>Homo sapiens</i> adaptor-related protein complex 3, delta 1 subunit (AP3D1), transcript variant 3, mRNA.	1529

TABLE 12-continued

Non-redundant list of transcript variants that correspond to the set of genes that no CNV 'solutions' have been reported in the 71 PML cases			
RefSeq_Gene_ Symbol	RefSeq_Accession_ Number	mRNA_Description	SEQ ID
APOL1	NM_001136540	<i>Homo sapiens</i> apolipoprotein L1 (APOL1), transcript variant 3, mRNA.	1530
APOL1	NM_001136541	<i>Homo sapiens</i> apolipoprotein L1 (APOL1), transcript variant 4, mRNA.	1531
APOL1	NM_003661	<i>Homo sapiens</i> apolipoprotein L1 (APOL1), transcript variant 1, mRNA.	1532
APOL1	NM_145343	<i>Homo sapiens</i> apolipoprotein L1 (APOL1), transcript variant 2, mRNA.	1533
ASH1L	NM_018489	<i>Homo sapiens</i> ASH1 like histone lysine methyltransferase (ASH1L), mRNA.	1534
ATL2	NM_001135673	<i>Homo sapiens</i> atlastin GTPase 2 (ATL2), transcript variant 2, mRNA.	1535
ATL2	NM_022374	<i>Homo sapiens</i> atlastin GTPase 2 (ATL2), transcript variant 1, mRNA.	1536
ATL2	NR_024191	<i>Homo sapiens</i> atlastin GTPase 2 (ATL2), transcript variant 3, non-coding RNA.	1537
ATM	NM_000051	<i>Homo sapiens</i> ATM serine/threonine kinase (ATM), mRNA.	1538
ATR	NM_001184	<i>Homo sapiens</i> ATR serine/threonine kinase (ATR), mRNA.	1539
BACH2	NM_001170794	<i>Homo sapiens</i> BTB domain and CNC homolog 2 (BACH2), transcript variant 2, mRNA.	1540
BACH2	NM_021813	<i>Homo sapiens</i> BTB domain and CNC homolog 2 (BACH2), transcript variant 1, mRNA.	1541
BAG3	NM_004281	<i>Homo sapiens</i> BCL2 associated athanogene 3 (BAG3), mRNA.	1542
BCL10	NM_003921	<i>Homo sapiens</i> B-cell CLL/lymphoma 10 (BCL10), transcript variant 1, mRNA.	1543
BLM	NM_000057	<i>Homo sapiens</i> Bloom syndrome RecQ like helicase (BLM), transcript variant 1, mRNA.	1544
BLNK	NM_001114094	<i>Homo sapiens</i> B-cell linker (BLNK), transcript variant 2, mRNA.	1545
BLNK	NM_001258440	<i>Homo sapiens</i> B-cell linker (BLNK), transcript variant 3, mRNA.	1546
BLNK	NM_001258441	<i>Homo sapiens</i> B-cell linker (BLNK), transcript variant 4, mRNA.	1547
BLNK	NM_001258442	<i>Homo sapiens</i> B-cell linker (BLNK), transcript variant 5, mRNA.	1548
BLNK	NM_013314	<i>Homo sapiens</i> B-cell linker (BLNK), transcript variant 1, mRNA.	1549
BLNK	NR_047680	<i>Homo sapiens</i> B-cell linker (BLNK), transcript variant 6, non-coding RNA.	1550
BLNK	NR_047681	<i>Homo sapiens</i> B-cell linker (BLNK), transcript variant 7, non-coding RNA.	1551
BLNK	NR_047682	<i>Homo sapiens</i> B-cell linker (BLNK), transcript variant 8, non-coding RNA.	1552
BLNK	NR_047683	<i>Homo sapiens</i> B-cell linker (BLNK), transcript variant 9, non-coding RNA.	1553
BLOC1S6	NM_012388	<i>Homo sapiens</i> biogenesis of lysosomal organelles complex 1 subunit 6 (BLOC1S6), transcript variant 2, mRNA.	1554
BTK	NM_000061	<i>Homo sapiens</i> Bruton tyrosine kinase (BTK), transcript variant 1, mRNA.	1555
C11orf65	NM_152587	<i>Homo sapiens</i> chromosome 11 open reading frame 65 (C11orf65), mRNA.	1556
C1QA	NM_015991	<i>Homo sapiens</i> complement component 1, q subcomponent, A chain (C1QA), mRNA.	1557
C1QB	NM_000491	<i>Homo sapiens</i> complement component 1, q subcomponent, B chain (C1QB), mRNA.	1558
C1QC	NM_001114101	<i>Homo sapiens</i> complement component 1, q subcomponent, C chain (C1QC), transcript variant 1, mRNA.	1559
C1QC	NM_172369	<i>Homo sapiens</i> complement component 1, q subcomponent, C chain (C1QC), transcript variant 2, mRNA.	1560
C5AR1	NM_001736	<i>Homo sapiens</i> complement component 5a receptor 1 (C5AR1), mRNA.	1561
CARD11	NM_032415	<i>Homo sapiens</i> caspase recruitment domain family member 11 (CARD11), transcript variant 2, mRNA.	1562
CARD9	NM_052813	<i>Homo sapiens</i> caspase recruitment domain family, member 9 (CARD9), transcript variant 1, mRNA.	1563
CARD9	NM_052814	<i>Homo sapiens</i> caspase recruitment domain family, member 9 (CARD9), transcript variant 2, mRNA.	1564
CASP8	NM_001080124	<i>Homo sapiens</i> caspase 8 (CASP8), transcript variant F, mRNA.	1565
CASP8	NM_001228	<i>Homo sapiens</i> caspase 8 (CASP8), transcript variant A, mRNA.	1566
CASP8	NM_033355	<i>Homo sapiens</i> caspase 8 (CASP8), transcript variant B, mRNA.	1567
CASP8	NM_033358	<i>Homo sapiens</i> caspase 8 (CASP8), transcript variant E, mRNA.	1568
CASP8	NM_001080125	<i>Homo sapiens</i> caspase 8 (CASP8), transcript variant G, mRNA.	1569
CASP8	NM_033356	<i>Homo sapiens</i> caspase 8 (CASP8), transcript variant C, mRNA.	1570
CCL11	NM_002986	<i>Homo sapiens</i> C-C motif chemokine ligand 11 (CCL11), mRNA.	1571
CCL2	NM_002982	<i>Homo sapiens</i> C-C motif chemokine ligand 2 (CCL2), mRNA.	1572
CCL5	NM_002985	<i>Homo sapiens</i> C-C motif chemokine ligand 5 (CCL5), transcript variant 1, mRNA.	1573
CCR2	NM_001123041	<i>Homo sapiens</i> C-C motif chemokine receptor 2 (CCR2), transcript variant A, mRNA.	1574
CCR2	NM_001123396	<i>Homo sapiens</i> C-C motif chemokine receptor 2 (CCR2), transcript variant B, mRNA.	1575
CCR5	NM_000579	<i>Homo sapiens</i> C-C motif chemokine receptor 5 (gene/pseudogene) (CCR5), transcript variant A, mRNA.	1576
CCR5	NM_001100168	<i>Homo sapiens</i> C-C motif chemokine receptor 5 (gene/pseudogene) (CCR5), transcript variant B, mRNA.	1577
CD180	NM_005582	<i>Homo sapiens</i> CD180 molecule (CD180), mRNA.	1578
CD19	NM_001178098	<i>Homo sapiens</i> CD19 molecule (CD19), transcript variant 1, mRNA.	1579
CD19	NM_0011770	<i>Homo sapiens</i> CD19 molecule (CD19), transcript variant 2, mRNA.	1580
CD209	NM_001144893	<i>Homo sapiens</i> CD209 molecule (CD209), transcript variant 5, mRNA.	1581
CD209	NM_001144894	<i>Homo sapiens</i> CD209 molecule (CD209), transcript variant 6, mRNA.	1582
CD209	NM_001144895	<i>Homo sapiens</i> CD209 molecule (CD209), transcript variant 7, mRNA.	1583
CD209	NM_001144896	<i>Homo sapiens</i> CD209 molecule (CD209), transcript variant 3, mRNA.	1584
CD209	NM_001144897	<i>Homo sapiens</i> CD209 molecule (CD209), transcript variant 4, mRNA.	1585
CD209	NM_001144899	<i>Homo sapiens</i> CD209 molecule (CD209), transcript variant 8, mRNA.	1586
CD209	NM_021155	<i>Homo sapiens</i> CD209 molecule (CD209), transcript variant 1, mRNA.	1587
CD209	NR_026692	<i>Homo sapiens</i> CD209 molecule (CD209), transcript variant 2, non-coding RNA.	1588
CD247	NM_000734	<i>Homo sapiens</i> CD247 molecule (CD247), transcript variant 2, mRNA.	1589
CD247	NM_198053	<i>Homo sapiens</i> CD247 molecule (CD247), transcript variant 1, mRNA.	1590
CD27	NM_001242	<i>Homo sapiens</i> CD27 molecule (CD27), mRNA.	1591
CD27-AS1	NR_015382	<i>Homo sapiens</i> CD27 antisense RNA 1 (CD27-AS1), long non-coding RNA.	1592
CD34	NM_001025109	<i>Homo sapiens</i> CD34 molecule (CD34), transcript variant 1, mRNA.	1593

TABLE 12-continued

Non-redundant list of transcript variants that correspond to the set of genes that no CNV 'solutions' have been reported in the 71 PML cases			
RefSeq_Gene_ Symbol	RefSeq_Accession_ Number	mRNA_Description	SEQ ID
CD34	NM_001773	<i>Homo sapiens</i> CD34 molecule (CD34), transcript variant 2, mRNA.	1594
CD3D	NM_000732	<i>Homo sapiens</i> CD3d molecule (CD3D), transcript variant 1, mRNA.	1595
CD3D	NM_001040651	<i>Homo sapiens</i> CD3d molecule (CD3D), transcript variant 2, mRNA.	1596
CD3E	NM_000733	<i>Homo sapiens</i> CD3e molecule (CD3E), mRNA.	1597
CD3G	NM_000073	<i>Homo sapiens</i> CD3g molecule (CD3G), mRNA.	1598
CD40	NM_001250	<i>Homo sapiens</i> CD40 molecule (CD40), transcript variant 1, mRNA.	1599
CD40	NM_152854	<i>Homo sapiens</i> CD40 molecule (CD40), transcript variant 2, mRNA.	1600
CD40LG	NM_000074	<i>Homo sapiens</i> CD40 ligand (CD40LG), mRNA.	1601
CD55	NM_000574	<i>Homo sapiens</i> CD55 molecule (Cromer blood group) (CD55), transcript variant 1, mRNA.	1602
CD55	NM_001114752	<i>Homo sapiens</i> CD55 molecule (Cromer blood group) (CD55), transcript variant 2, mRNA.	1603
CD59	NM_000611	<i>Homo sapiens</i> CD59 molecule (CD59), transcript variant 2, mRNA.	1604
CD59	NM_001127223	<i>Homo sapiens</i> CD59 molecule (CD59), transcript variant 5, mRNA.	1605
CD59	NM_001127225	<i>Homo sapiens</i> CD59 molecule (CD59), transcript variant 6, mRNA.	1606
CD59	NM_001127226	<i>Homo sapiens</i> CD59 molecule (CD59), transcript variant 7, mRNA.	1607
CD59	NM_001127227	<i>Homo sapiens</i> CD59 molecule (CD59), transcript variant 8, mRNA.	1608
CD59	NM_203329	<i>Homo sapiens</i> CD59 molecule (CD59), transcript variant 3, mRNA.	1609
CD59	NM_203330	<i>Homo sapiens</i> CD59 molecule (CD59), transcript variant 1, mRNA.	1610
CD59	NM_203331	<i>Homo sapiens</i> CD59 molecule (CD59), transcript variant 4, mRNA.	1611
CD79A	NM_001783	<i>Homo sapiens</i> CD79a molecule (CD79A), transcript variant 1, mRNA.	1612
CD79A	NM_021601	<i>Homo sapiens</i> CD79a molecule (CD79A), transcript variant 2, mRNA.	1613
CD79B	NM_000626	<i>Homo sapiens</i> CD79b molecule (CD79B), transcript variant 1, mRNA.	1614
CD79B	NM_001039933	<i>Homo sapiens</i> CD79b molecule (CD79B), transcript variant 3, mRNA.	1615
CD79B	NM_021602	<i>Homo sapiens</i> CD79b molecule (CD79B), transcript variant 2, mRNA.	1616
CD81	NM_004356	<i>Homo sapiens</i> CD81 molecule (CD81), transcript variant 1, mRNA.	1617
CD8A	NM_001145873	<i>Homo sapiens</i> CD8a molecule (CD8A), transcript variant 3, mRNA.	1618
CD8A	NM_001768	<i>Homo sapiens</i> CD8a molecule (CD8A), transcript variant 1, mRNA.	1619
CD8A	NM_171827	<i>Homo sapiens</i> CD8a molecule (CD8A), transcript variant 2, mRNA.	1620
CD8A	NR_027353	<i>Homo sapiens</i> CD8a molecule (CD8A), transcript variant 4, non-coding RNA.	1621
CDCA7	NM_031942	<i>Homo sapiens</i> cell division cycle associated 7 (CDCA7), transcript variant 1, mRNA.	1622
CDCA7	NM_145810	<i>Homo sapiens</i> cell division cycle associated 7 (CDCA7), transcript variant 2, mRNA.	1623
CEBPB	NM_005194	<i>Homo sapiens</i> CCAAT/enhancer binding protein beta (CEBPB), transcript variant 1, mRNA.	1624
CHD7	NM_017780	<i>Homo sapiens</i> chromodomain helicase DNA binding protein 7 (CHD7), transcript variant 1, mRNA.	1625
CHEK1	NM_001114121	<i>Homo sapiens</i> checkpoint kinase 1 (CHEK1), transcript variant 2, mRNA.	1626
CHEK1	NM_001114122	<i>Homo sapiens</i> checkpoint kinase 1 (CHEK1), transcript variant 1, mRNA.	1627
CHEK1	NM_001244846	<i>Homo sapiens</i> checkpoint kinase 1 (CHEK1), transcript variant 4, mRNA.	1628
CHEK1	NR_045204	<i>Homo sapiens</i> checkpoint kinase 1 (CHEK1), transcript variant 5, non-coding RNA.	1629
CHEK1	NR_045205	<i>Homo sapiens</i> checkpoint kinase 1 (CHEK1), transcript variant 6, non-coding RNA.	1630
CHEK1	NM_001274	<i>Homo sapiens</i> checkpoint kinase 1 (CHEK1), transcript variant 3, mRNA.	1631
CIITA	NM_000246	<i>Homo sapiens</i> class II major histocompatibility complex transactivator (CIITA), transcript variant 2, mRNA.	1632
CLCN7	NM_001114331	<i>Homo sapiens</i> chloride channel, voltage-sensitive 7 (CLCN7), transcript variant 2, mRNA.	1633
CLCN7	NM_001287	<i>Homo sapiens</i> chloride channel, voltage-sensitive 7 (CLCN7), transcript variant 1, mRNA.	1634
COG6	NM_001145079	<i>Homo sapiens</i> component of oligomeric golgi complex 6 (COG6), transcript variant 2, mRNA.	1635
COG6	NM_020751	<i>Homo sapiens</i> component of oligomeric golgi complex 6 (COG6), transcript variant 1, mRNA.	1636
COG6	NR_026745	<i>Homo sapiens</i> component of oligomeric golgi complex 6 (COG6), transcript variant 3, non-coding RNA.	1637
CORO1A	NM_001193333	<i>Homo sapiens</i> coronin 1A (CORO1A), transcript variant 1, mRNA.	1638
CORO1A	NM_007074	<i>Homo sapiens</i> coronin 1A (CORO1A), transcript variant 2, mRNA.	1639
CR2	NM_001006658	<i>Homo sapiens</i> complement component 3d receptor 2 (CR2), transcript variant 1, mRNA.	1640
CR2	NM_001877	<i>Homo sapiens</i> complement component 3d receptor 2 (CR2), transcript variant 2, mRNA.	1641
CRTC3	NM_001042574	<i>Homo sapiens</i> CREB regulated transcription coactivator 3 (CRTC3), transcript variant 2, mRNA.	1642
CRTC3	NM_022769	<i>Homo sapiens</i> CREB regulated transcription coactivator 3 (CRTC3), transcript variant 1, mRNA.	1643
CSF3R	NM_000760	<i>Homo sapiens</i> colony stimulating factor 3 receptor (granulocyte) (CSF3R), transcript variant 1, mRNA.	1644
CSF3R	NM_156039	<i>Homo sapiens</i> colony stimulating factor 3 receptor (granulocyte) (CSF3R), transcript variant 3, mRNA.	1645
CSF3R	NM_172313	<i>Homo sapiens</i> colony stimulating factor 3 receptor (granulocyte) (CSF3R), transcript variant 4, mRNA.	1646
CTLA4	NM_005214	<i>Homo sapiens</i> cytotoxic T-lymphocyte-associated protein 4 (CTLA4), transcript variant 1, mRNA.	1647
CTLA4	NM_001037631	<i>Homo sapiens</i> cytotoxic T-lymphocyte-associated protein 4 (CTLA4), transcript variant 2, mRNA.	1648
CTPS1	NM_001905	<i>Homo sapiens</i> CTP synthase 1 (CTPS1), transcript variant 1, mRNA.	1649
CTSC	NM_148170	<i>Homo sapiens</i> cathepsin C (CTSC), transcript variant 2, mRNA.	1650
CTSC	NM_001114173	<i>Homo sapiens</i> cathepsin C (CTSC), transcript variant 3, mRNA.	1651
CTSC	NM_001814	<i>Homo sapiens</i> cathepsin C (CTSC), transcript variant 1, mRNA.	1652

TABLE 12-continued

Non-redundant list of transcript variants that correspond to the set of genes that no CNV 'solutions' have been reported in the 71 PML cases			
RefSeq_Gene_ Symbol	RefSeq_Accession_ Number	mRNA_Description	SEQ ID
CX3CR1	NM_001171171	<i>Homo sapiens</i> C-X3-C motif chemokine receptor 1 (CX3CR1), transcript variant 2, mRNA.	1653
CX3CR1	NM_001171172	<i>Homo sapiens</i> C-X3-C motif chemokine receptor 1 (CX3CR1), transcript variant 3, mRNA.	1654
CX3CR1	NM_001171174	<i>Homo sapiens</i> C-X3-C motif chemokine receptor 1 (CX3CR1), transcript variant 1, mRNA.	1655
CX3CR1	NM_001337	<i>Homo sapiens</i> C-X3-C motif chemokine receptor 1 (CX3CR1), transcript variant 4, mRNA.	1656
CXCL12	NM_000609	<i>Homo sapiens</i> C-X-C motif chemokine ligand 12 (CXCL12), transcript variant 2, mRNA.	1657
CXCL12	NM_001033886	<i>Homo sapiens</i> C-X-C motif chemokine ligand 12 (CXCL12), transcript variant 3, mRNA.	1658
CXCL12	NM_001178134	<i>Homo sapiens</i> C-X-C motif chemokine ligand 12 (CXCL12), transcript variant 4, mRNA.	1659
CXCL12	NM_199168	<i>Homo sapiens</i> C-X-C motif chemokine ligand 12 (CXCL12), transcript variant 1, mRNA.	1660
CXCL9	NM_002416	<i>Homo sapiens</i> C-X-C motif chemokine ligand 9 (CXCL9), mRNA.	1661
CXCR1	NM_000634	<i>Homo sapiens</i> C-X-C motif chemokine receptor 1 (CXCR1), mRNA.	1662
CXCR4	NM_001008540	<i>Homo sapiens</i> C-X-C motif chemokine receptor 4 (CXCR4), transcript variant 1, mRNA.	1663
CXCR4	NM_003467	<i>Homo sapiens</i> C-X-C motif chemokine receptor 4 (CXCR4), transcript variant 2, mRNA.	1664
CXorf40A	NM_001171907	<i>Homo sapiens</i> chromosome X open reading frame 40A (CXorf40A), transcript variant 2, mRNA.	1665
CXorf40A	NM_001171908	<i>Homo sapiens</i> chromosome X open reading frame 40A (CXorf40A), transcript variant 3, mRNA.	1666
CXorf40A	NM_178124	<i>Homo sapiens</i> chromosome X open reading frame 40A (CXorf40A), transcript variant 1, mRNA.	1667
CXorf40A	NM_001171909	<i>Homo sapiens</i> chromosome X open reading frame 40A (CXorf40A), transcript variant 4, mRNA.	1668
CYBB	NM_000397	<i>Homo sapiens</i> cytochrome b-245, beta polypeptide (CYBB), mRNA.	1669
CYP2S1	NM_030622	<i>Homo sapiens</i> cytochrome P450 family 2 subfamily S member 1 (CYP2S1), mRNA.	1670
DCLRE1C	NM_001033855	<i>Homo sapiens</i> DNA cross-link repair 1C (DCLRE1C), transcript variant a, mRNA.	1671
DCLRE1C	NM_001033857	<i>Homo sapiens</i> DNA cross-link repair 1C (DCLRE1C), transcript variant d, mRNA.	1672
DCLRE1C	NM_001033858	<i>Homo sapiens</i> DNA cross-link repair 1C (DCLRE1C), transcript variant c, mRNA.	1673
DCLRE1C	NM_022487	<i>Homo sapiens</i> DNA cross-link repair 1C (DCLRE1C), transcript variant b, mRNA.	1674
DDX1	NM_004939	<i>Homo sapiens</i> DEAD/H-box helicase 1 (DDX1), mRNA.	1675
DDX58	NM_014314	<i>Homo sapiens</i> DEAD (Asp-Glu-Ala-Asp) box polypeptide 58 (DDX58), mRNA.	1676
DHX58	NM_024119	<i>Homo sapiens</i> DEXH (Asp-Glu-X-His) box polypeptide 58 (DHX58), mRNA.	1677
DKC1	NM_001142463	<i>Homo sapiens</i> dyskerin pseudouridine synthase 1 (DKC1), transcript variant 2, mRNA.	1678
DKC1	NM_001363	<i>Homo sapiens</i> dyskerin pseudouridine synthase 1 (DKC1), transcript variant 1, mRNA.	1679
DNMT3B	NM_001207055	<i>Homo sapiens</i> DNA methyltransferase 3 beta (DNMT3B), transcript variant 7, mRNA.	1680
DNMT3B	NM_001207056	<i>Homo sapiens</i> DNA methyltransferase 3 beta (DNMT3B), transcript variant 8, mRNA.	1681
DNMT3B	NM_006892	<i>Homo sapiens</i> DNA methyltransferase 3 beta (DNMT3B), transcript variant 1, mRNA.	1682
DNMT3B	NM_175848	<i>Homo sapiens</i> DNA methyltransferase 3 beta (DNMT3B), transcript variant 2, mRNA.	1683
DNMT3B	NM_175849	<i>Homo sapiens</i> DNA methyltransferase 3 beta (DNMT3B), transcript variant 3, mRNA.	1684
DNMT3B	NM_175850	<i>Homo sapiens</i> DNA methyltransferase 3 beta (DNMT3B), transcript variant 6, mRNA.	1685
DOCK2	NM_004946	<i>Homo sapiens</i> dedicator of cytokinesis 2 (DOCK2), mRNA.	1686
DOCK8	NM_203447	<i>Homo sapiens</i> dedicator of cytokinesis 8 (DOCK8), transcript variant 1, mRNA.	1687
DOCK8	NM_001190458	<i>Homo sapiens</i> dedicator of cytokinesis 8 (DOCK8), transcript variant 2, mRNA.	1688
DOCK8	NM_001193536	<i>Homo sapiens</i> dedicator of cytokinesis 8 (DOCK8), transcript variant 3, mRNA.	1689
DSC1	NM_004948	<i>Homo sapiens</i> desmocollin 1 (DSC1), transcript variant Dsc1b, mRNA.	1690
DSC1	NM_024421	<i>Homo sapiens</i> desmocollin 1 (DSC1), transcript variant Dsc1a, mRNA.	1691
EGR1	NM_001964	<i>Homo sapiens</i> early growth response 1 (EGR1), mRNA.	1692
ELANE	NM_001972	<i>Homo sapiens</i> elastase, neutrophil expressed (ELANE), mRNA.	1693
EPG5	NM_020964	<i>Homo sapiens</i> ectopic P-granules autophagy protein 5 homolog (EPG5), mRNA.	1694
ETF1	NM_004730	<i>Homo sapiens</i> eukaryotic translation termination factor 1 (ETF1), transcript variant 1, mRNA.	1695
F9	NM_000133	<i>Homo sapiens</i> coagulation factor IX (F9), transcript variant 1, mRNA.	1696
FAS	NM_000043	<i>Homo sapiens</i> Fas cell surface death receptor (FAS), transcript variant 1, mRNA.	1697
FAS	NM_152871	<i>Homo sapiens</i> Fas cell surface death receptor (FAS), transcript variant 2, mRNA.	1698
FAS	NM_152872	<i>Homo sapiens</i> Fas cell surface death receptor (FAS), transcript variant 3, mRNA.	1699
FAS	NR_028033	<i>Homo sapiens</i> Fas cell surface death receptor (FAS), transcript variant 4, non-coding RNA.	1700
FAS	NR_028034	<i>Homo sapiens</i> Fas cell surface death receptor (FAS), transcript variant 5, non-coding RNA.	1701
FAS	NR_028035	<i>Homo sapiens</i> Fas cell surface death receptor (FAS), transcript variant 6, non-coding RNA.	1702
FAS	NR_028036	<i>Homo sapiens</i> Fas cell surface death receptor (FAS), transcript variant 7, non-coding RNA.	1703
FASLG	NM_000639	<i>Homo sapiens</i> Fas ligand (TNF superfamily, member 6) (FASLG), mRNA.	1704
FCGR2A	NM_001136219	<i>Homo sapiens</i> Fc fragment of IgG receptor IIa (FCGR2A), transcript variant 1, mRNA.	1705
FCGR2A	NM_021642	<i>Homo sapiens</i> Fc fragment of IgG receptor IIa (FCGR2A), transcript variant 2, mRNA.	1706
FCGR3A	NM_000569	<i>Homo sapiens</i> Fc fragment of IgG receptor IIIa (FCGR3A), transcript variant 1, mRNA.	1707
FCGR3A	NM_001127592	<i>Homo sapiens</i> Fc fragment of IgG receptor IIIa (FCGR3A), transcript variant 2, mRNA.	1708
FCGR3A	NM_001127593	<i>Homo sapiens</i> Fc fragment of IgG receptor IIIa (FCGR3A), transcript variant 3, mRNA.	1709
FCGR3A	NM_001127595	<i>Homo sapiens</i> Fc fragment of IgG receptor IIIa (FCGR3A), transcript variant 4, mRNA.	1710
FCGR3A	NM_001127596	<i>Homo sapiens</i> Fc fragment of IgG receptor IIIa (FCGR3A), transcript variant 5, mRNA.	1711
FCN3	NM_003665	<i>Homo sapiens</i> ficolin 3 (FCN3), transcript variant 1, mRNA.	1712
FCN3	NM_173452	<i>Homo sapiens</i> ficolin 3 (FCN3), transcript variant 2, mRNA.	1713
FEZ1	NM_005103	<i>Homo sapiens</i> fasciculation and elongation protein zeta 1 (FEZ1), transcript variant 1, mRNA.	1714

TABLE 12-continued

Non-redundant list of transcript variants that correspond to the set of genes that no CNV 'solutions' have been reported in the 71 PML cases			
RefSeq_Gene_ Symbol	RefSeq_Accession_ Number	mRNA_Description	SEQ ID
FEZ1	NM_022549	<i>Homo sapiens</i> fasciculation and elongation protein zeta 1 (FEZ1), transcript variant 2, mRNA.	1715
FOS	NM_005252	<i>Homo sapiens</i> Fos proto-oncogene, AP-1 transcription factor subunit (FOS), mRNA.	1716
FOXH1	NM_003923	<i>Homo sapiens</i> forkhead box H1 (FOXH1), mRNA.	1717
FOXN1	NM_003593	<i>Homo sapiens</i> forkhead box N1 (FOXN1), mRNA.	1718
FOXP3	NM_001114377	<i>Homo sapiens</i> forkhead box P3 (FOXP3), transcript variant 2, mRNA.	1719
FOXP3	NM_014009	<i>Homo sapiens</i> forkhead box P3 (FOXP3), transcript variant 1, mRNA.	1720
FPR1	NM_001193306	<i>Homo sapiens</i> formyl peptide receptor 1 (FPR1), transcript variant 1, mRNA.	1721
FPR1	NM_002029	<i>Homo sapiens</i> formyl peptide receptor 1 (FPR1), transcript variant 2, mRNA.	1722
G6PC3	NM_138387	<i>Homo sapiens</i> glucose 6 phosphatase, catalytic, 3 (G6PC3), transcript variant 1, mRNA.	1723
G6PC3	NR_028582	<i>Homo sapiens</i> glucose 6 phosphatase, catalytic, 3 (G6PC3), transcript variant 2, non-coding RNA.	1724
G6PC3	NR_028581	<i>Homo sapiens</i> glucose 6 phosphatase, catalytic, 3 (G6PC3), transcript variant 3, non-coding RNA.	1725
GATA2	NM_001145661	<i>Homo sapiens</i> GATA binding protein 2 (GATA2), transcript variant 1, mRNA.	1726
GATA2	NM_001145662	<i>Homo sapiens</i> GATA binding protein 2 (GATA2), transcript variant 3, mRNA.	1727
GATA2	NM_032638	<i>Homo sapiens</i> GATA binding protein 2 (GATA2), transcript variant 2, mRNA.	1728
GFI1	NM_005263	<i>Homo sapiens</i> growth factor independent 1 transcription repressor (GFI1), transcript variant 1, mRNA.	1729
GFI1	NM_001127215	<i>Homo sapiens</i> growth factor independent 1 transcription repressor (GFI1), transcript variant 2, mRNA.	1730
GFI1	NM_001127216	<i>Homo sapiens</i> growth factor independent 1 transcription repressor (GFI1), transcript variant 3, mRNA.	1731
GOLGB1	NM_001256486	<i>Homo sapiens</i> golgin B1 (GOLGB1), transcript variant 1, mRNA.	1732
GOLGB1	NM_001256487	<i>Homo sapiens</i> golgin B1 (GOLGB1), transcript variant 3, mRNA.	1733
GOLGB1	NM_001256488	<i>Homo sapiens</i> golgin B1 (GOLGB1), transcript variant 4, mRNA.	1734
GOLGB1	NM_004487	<i>Homo sapiens</i> golgin B1 (GOLGB1), transcript variant 2, mRNA.	1735
GPRC5A	NM_003979	<i>Homo sapiens</i> G protein-coupled receptor class C group 5 member A (GPRC5A), mRNA.	1736
GRAP2	NM_004810	<i>Homo sapiens</i> GRB2-related adaptor protein 2 (GRAP2), mRNA.	1737
HAX1	NM_006118	<i>Homo sapiens</i> HCLS1 associated protein X-1 (HAX1), transcript variant 1, mRNA.	1738
HAX1	NM_001018837	<i>Homo sapiens</i> HCLS1 associated protein X-1 (HAX1), transcript variant 2, mRNA.	1739
HELLS	NM_018063	<i>Homo sapiens</i> helicase, lymphoid-specific (HELLS), transcript variant 1, mRNA.	1740
HIVEP1	NM_002114	<i>Homo sapiens</i> human immunodeficiency virus type I enhancer binding protein 1 (HIVEP1), mRNA.	1741
HIVEP2	NM_006734	<i>Homo sapiens</i> human immunodeficiency virus type I enhancer binding protein 2 (HIVEP2), mRNA.	1742
HIVEP3	NM_001127714	<i>Homo sapiens</i> human immunodeficiency virus type I enhancer binding protein 3 (HIVEP3), transcript variant 2, mRNA.	1743
HIVEP3	NM_024503	<i>Homo sapiens</i> human immunodeficiency virus type I enhancer binding protein 3 (HIVEP3), transcript variant 1, mRNA.	1744
HIVEP3	NR_038260	<i>Homo sapiens</i> human immunodeficiency virus type I enhancer binding protein 3 (HIVEP3), transcript variant 3, non-coding RNA.	1745
HIVEP3	NR_038261	<i>Homo sapiens</i> human immunodeficiency virus type I enhancer binding protein 3 (HIVEP3), transcript variant 4, non-coding RNA.	1746
HP	NM_001126102	<i>Homo sapiens</i> haptoglobin (HP), transcript variant 2, mRNA.	1747
HP	NM_005143	<i>Homo sapiens</i> haptoglobin (HP), transcript variant 1, mRNA.	1748
HPCAL1	NM_002149	<i>Homo sapiens</i> hippocalcin like 1 (HPCAL1), transcript variant 1, mRNA.	1749
HPCAL1	NM_134421	<i>Homo sapiens</i> hippocalcin like 1 (HPCAL1), transcript variant 2, mRNA.	1750
HPCAL1	NM_001258357	<i>Homo sapiens</i> hippocalcin like 1 (HPCAL1), transcript variant 3, mRNA.	1751
HPCAL1	NM_001258358	<i>Homo sapiens</i> hippocalcin like 1 (HPCAL1), transcript variant 4, mRNA.	1752
HPCAL1	NM_001258359	<i>Homo sapiens</i> hippocalcin like 1 (HPCAL1), transcript variant 5, mRNA.	1753
HTR2A	NM_000621	<i>Homo sapiens</i> 5-hydroxytryptamine receptor 2A (HTR2A), transcript variant 1, mRNA.	1754
HTR2A	NM_001165947	<i>Homo sapiens</i> 5-hydroxytryptamine (serotonin) receptor 2A, G protein-coupled (HTR2A), transcript variant 2, mRNA.	1755
ICOS	NM_012092	<i>Homo sapiens</i> inducible T-cell costimulator (ICOS), mRNA.	1756
IDI1	NM_004508	<i>Homo sapiens</i> isopentenyl-diphosphate delta isomerase 1 (IDI1), transcript variant 1, mRNA.	1757
IFIH1	NM_022168	<i>Homo sapiens</i> interferon induced with helicase C domain 1 (IFIH1), mRNA.	1758
IFNAR1	NM_000629	<i>Homo sapiens</i> interferon (alpha, beta and omega) receptor 1 (IFNAR1), mRNA.	1759
IFNAR2	NM_207584	<i>Homo sapiens</i> interferon (alpha, beta and omega) receptor 2 (IFNAR2), transcript variant 3, mRNA.	1760
IFNAR2	NM_207585	<i>Homo sapiens</i> interferon (alpha, beta and omega) receptor 2 (IFNAR2), transcript variant 1, mRNA.	1761
IFNAR2	NM_000874	<i>Homo sapiens</i> interferon (alpha, beta and omega) receptor 2 (IFNAR2), transcript variant 2, mRNA.	1762
IFNG	NM_000619	<i>Homo sapiens</i> interferon gamma (IFNG), mRNA.	1763
IFNGR1	NM_000416	<i>Homo sapiens</i> interferon gamma receptor 1 (IFNGR1), mRNA.	1764
IFNGR2	NM_005534	<i>Homo sapiens</i> interferon gamma receptor 2 (interferon gamma transducer 1) (IFNGR2), transcript variant 2, mRNA.	1765
IGLL1	NM_020070	<i>Homo sapiens</i> immunoglobulin lambda like polypeptide 1 (IGLL1), transcript variant 1, mRNA.	1766

TABLE 12-continued

Non-redundant list of transcript variants that correspond to the set of genes that no CNV 'solutions' have been reported in the 71 PML cases			
RefSeq_Gene_ Symbol	RefSeq_Accession_ Number	mRNA_Description	SEQ ID
IGLL1	NM_152855	<i>Homo sapiens</i> immunoglobulin lambda like polypeptide 1 (IGLL1), transcript variant 2, mRNA.	1767
IKBKB	NM_001190720	<i>Homo sapiens</i> inhibitor of kappa light polypeptide gene enhancer in B-cells, kinase beta (IKBKB), transcript variant 2, mRNA.	1768
IKBKB	NM_001242778	<i>Homo sapiens</i> inhibitor of kappa light polypeptide gene enhancer in B-cells, kinase beta (IKBKB), transcript variant 7, mRNA.	1769
IKBKB	NM_001556	<i>Homo sapiens</i> inhibitor of kappa light polypeptide gene enhancer in B-cells, kinase beta (IKBKB), transcript variant 1, mRNA.	1770
IKBKB	NR_033818	<i>Homo sapiens</i> inhibitor of kappa light polypeptide gene enhancer in B-cells, kinase beta (IKBKB), transcript variant 5, non-coding RNA.	1771
IKBKB	NR_033819	<i>Homo sapiens</i> inhibitor of kappa light polypeptide gene enhancer in B-cells, kinase beta (IKBKB), transcript variant 6, non-coding RNA.	1772
IKBKB	NR_040009	<i>Homo sapiens</i> inhibitor of kappa light polypeptide gene enhancer in B-cells, kinase beta (IKBKB), transcript variant 8, non-coding RNA.	1773
IKBKG	NM_001099856	<i>Homo sapiens</i> inhibitor of kappa light polypeptide gene enhancer in B-cells, kinase gamma (IKBKG), transcript variant 2, mRNA.	1774
IKBKG	NM_001099857	<i>Homo sapiens</i> inhibitor of kappa light polypeptide gene enhancer in B-cells, kinase gamma (IKBKG), transcript variant 1, mRNA.	1775
IKBKG	NM_001145255	<i>Homo sapiens</i> inhibitor of kappa light polypeptide gene enhancer in B-cells, kinase gamma (IKBKG), transcript variant 4, mRNA.	1776
IKBKG	NM_003639	<i>Homo sapiens</i> inhibitor of kappa light polypeptide gene enhancer in B-cells, kinase gamma (IKBKG), transcript variant 3, mRNA.	1777
IKZF1	NM_001220765	<i>Homo sapiens</i> IKAROS family zinc finger 1 (IKZF1), transcript variant 2, mRNA.	1778
IKZF1	NM_001220767	<i>Homo sapiens</i> IKAROS family zinc finger 1 (IKZF1), transcript variant 4, mRNA.	1779
IKZF1	NM_001220768	<i>Homo sapiens</i> IKAROS family zinc finger 1 (IKZF1), transcript variant 5, mRNA.	1780
IKZF1	NM_001220770	<i>Homo sapiens</i> IKAROS family zinc finger 1 (IKZF1), transcript variant 7, mRNA.	1781
IKZF1	NM_001220771	<i>Homo sapiens</i> IKAROS family zinc finger 1 (IKZF1), transcript variant 8, mRNA.	1782
IKZF1	NM_006060	<i>Homo sapiens</i> IKAROS family zinc finger 1 (IKZF1), transcript variant 1, mRNA.	1783
IL10	NM_000572	<i>Homo sapiens</i> interleukin 10 (IL10), mRNA.	1784
IL10RA	NM_001558	<i>Homo sapiens</i> interleukin 10 receptor subunit alpha (IL10RA), transcript variant 1, mRNA.	1785
IL10RA	NR_026691	<i>Homo sapiens</i> interleukin 10 receptor subunit alpha (IL10RA), transcript variant 2, non-coding RNA.	1786
IL10RB	NM_000628	<i>Homo sapiens</i> interleukin 10 receptor subunit beta (IL10RB), mRNA.	1787
IL12B	NM_002187	<i>Homo sapiens</i> interleukin 12B (IL12B), mRNA.	1788
IL12RB1	NM_005535	<i>Homo sapiens</i> interleukin 12 receptor subunit beta 1 (IL12RB1), transcript variant 1, mRNA.	1789
IL12RB1	NM_153701	<i>Homo sapiens</i> interleukin 12 receptor subunit beta 1 (IL12RB1), transcript variant 2, mRNA.	1790
IL17F	NM_052872	<i>Homo sapiens</i> interleukin 17F (IL17F), mRNA.	1791
IL17RA	NM_014339	<i>Homo sapiens</i> interleukin 17 receptor A (IL17RA), transcript variant 1, mRNA.	1792
IL1B	NM_000576	<i>Homo sapiens</i> interleukin 1, beta (IL1B), mRNA.	1793
IL21	NM_001207006	<i>Homo sapiens</i> interleukin 21 (IL21), transcript variant 2, mRNA.	1794
IL21	NM_021803	<i>Homo sapiens</i> interleukin 21 (IL21), transcript variant 1, mRNA.	1795
IL21R	NM_181078	<i>Homo sapiens</i> interleukin 21 receptor (IL21R), transcript variant 2, mRNA.	1796
IL21R	NM_181079	<i>Homo sapiens</i> interleukin 21 receptor (IL21R), transcript variant 3, mRNA.	1797
IL21R	NM_021798	<i>Homo sapiens</i> interleukin 21 receptor (IL21R), transcript variant 1, mRNA.	1798
IL2RA	NM_000417	<i>Homo sapiens</i> interleukin 2 receptor, alpha (IL2RA), transcript variant 1, mRNA.	1799
IL2RG	NM_000206	<i>Homo sapiens</i> interleukin 2 receptor subunit gamma (IL2RG), mRNA.	1800
IL4R	NM_000418	<i>Homo sapiens</i> interleukin 4 receptor (IL4R), transcript variant 1, mRNA.	1801
IL4R	NM_001257406	<i>Homo sapiens</i> interleukin 4 receptor (IL4R), transcript variant 3, mRNA.	1802
IL4R	NM_001257407	<i>Homo sapiens</i> interleukin 4 receptor (IL4R), transcript variant 4, mRNA.	1803
IL4R	NM_001257997	<i>Homo sapiens</i> interleukin 4 receptor (IL4R), transcript variant 5, mRNA.	1804
IL7	NM_000880	<i>Homo sapiens</i> interleukin 7 (IL7), transcript variant 1, mRNA.	1805
IL7	NM_001199886	<i>Homo sapiens</i> interleukin 7 (IL7), transcript variant 2, mRNA.	1806
IL7	NM_001199887	<i>Homo sapiens</i> interleukin 7 (IL7), transcript variant 3, mRNA.	1807
IL7	NM_001199888	<i>Homo sapiens</i> interleukin 7 (IL7), transcript variant 4, mRNA.	1808
IL7R	NM_002185	<i>Homo sapiens</i> interleukin 7 receptor (IL7R), transcript variant 1, mRNA.	1809
IRAK4	NM_001114182	<i>Homo sapiens</i> interleukin 1 receptor associated kinase 4 (IRAK4), transcript variant 1, mRNA.	1810
IRAK4	NM_001145256	<i>Homo sapiens</i> interleukin 1 receptor associated kinase 4 (IRAK4), transcript variant 3, mRNA.	1811
IRAK4	NM_001145257	<i>Homo sapiens</i> interleukin 1 receptor associated kinase 4 (IRAK4), transcript variant 4, mRNA.	1812
IRAK4	NM_001145258	<i>Homo sapiens</i> interleukin 1 receptor associated kinase 4 (IRAK4), transcript variant 5, mRNA.	1813
IRAK4	NM_016123	<i>Homo sapiens</i> interleukin 1 receptor associated kinase 4 (IRAK4), transcript variant 2, mRNA.	1814
IRF3	NM_001197122	<i>Homo sapiens</i> interferon regulatory factor 3 (IRF3), transcript variant 2, mRNA.	1815
IRF3	NM_001197123	<i>Homo sapiens</i> interferon regulatory factor 3 (IRF3), transcript variant 3, mRNA.	1816
IRF3	NM_001197124	<i>Homo sapiens</i> interferon regulatory factor 3 (IRF3), transcript variant 4, mRNA.	1817
IRF3	NM_001197125	<i>Homo sapiens</i> interferon regulatory factor 3 (IRF3), transcript variant 5, mRNA.	1818
IRF3	NM_001197126	<i>Homo sapiens</i> interferon regulatory factor 3 (IRF3), transcript variant 6, mRNA.	1819

TABLE 12-continued

Non-redundant list of transcript variants that correspond to the set of genes that no CNV 'solutions' have been reported in the 71 PML cases			
RefSeq_Gene_ Symbol	RefSeq_Accession_ Number	mRNA_Description	SEQ ID
IRF3	NM_001197127	<i>Homo sapiens</i> interferon regulatory factor 3 (IRF3), transcript variant 7, mRNA.	1820
IRF3	NM_001197128	<i>Homo sapiens</i> interferon regulatory factor 3 (IRF3), transcript variant 8, mRNA.	1821
IRF3	NM_001571	<i>Homo sapiens</i> interferon regulatory factor 3 (IRF3), transcript variant 1, mRNA.	1822
IRF3	NR_045568	<i>Homo sapiens</i> interferon regulatory factor 3 (IRF3), transcript variant 9, non-coding RNA.	1823
IRF7	NM_001572	<i>Homo sapiens</i> interferon regulatory factor 7 (IRF7), transcript variant a, mRNA.	1824
IRF7	NM_004029	<i>Homo sapiens</i> interferon regulatory factor 7 (IRF7), transcript variant b, mRNA.	1825
IRF7	NM_004031	<i>Homo sapiens</i> interferon regulatory factor 7 (IRF7), transcript variant d, mRNA.	1826
IRF8	NM_002163	<i>Homo sapiens</i> interferon regulatory factor 8 (IRF8), mRNA.	1827
IRGM	NM_001145805	<i>Homo sapiens</i> immunity related GTPase M (IRGM), mRNA.	1828
ISG15	NM_005101	<i>Homo sapiens</i> ISG15 ubiquitin-like modifier (ISG15), mRNA.	1829
ITK	NM_005546	<i>Homo sapiens</i> IL2 inducible T-cell kinase (ITK), mRNA.	1830
ITSN2	NM_006277	<i>Homo sapiens</i> intersectin 2 (ITSN2), transcript variant 1, mRNA.	1831
ITSN2	NM_019595	<i>Homo sapiens</i> intersectin 2 (ITSN2), transcript variant 3, mRNA.	1832
ITSN2	NM_147152	<i>Homo sapiens</i> intersectin 2 (ITSN2), transcript variant 2, mRNA.	1833
JAGN1	NM_032492	<i>Homo sapiens</i> jagunal homolog 1 (<i>Drosophila</i>) (JAGN1), mRNA.	1834
JAK3	NM_000215	<i>Homo sapiens</i> Janus kinase 3 (JAK3), mRNA.	1835
JMY	NM_152405	<i>Homo sapiens</i> junction mediating and regulatory protein, p53 cofactor (JMY), mRNA.	1836
JUN	NM_002228	<i>Homo sapiens</i> Jun proto-oncogene, AP-1 transcription factor subunit (JUN), mRNA.	1837
KITLG	NM_000899	<i>Homo sapiens</i> KIT ligand (KITLG), transcript variant b, mRNA.	1838
KITLG	NM_003994	<i>Homo sapiens</i> KIT ligand (KITLG), transcript variant a, mRNA.	1839
LAMTOR2	NM_001145264	<i>Homo sapiens</i> late endosomal/lysosomal adaptor, MAPK and MTOR activator 2 (LAMTOR2), transcript variant 2, mRNA.	1840
LAMTOR2	NM_014017	<i>Homo sapiens</i> late endosomal/lysosomal adaptor, MAPK and MTOR activator 2 (LAMTOR2), transcript variant 1, mRNA.	1841
LCK	NM_005356	<i>Homo sapiens</i> LCK proto-oncogene, Src family tyrosine kinase (LCK), transcript variant 2, mRNA.	1842
LCK	NM_001042771	<i>Homo sapiens</i> LCK proto-oncogene, Src family tyrosine kinase (LCK), transcript variant 1, mRNA.	1843
LCP2	NM_005565	<i>Homo sapiens</i> lymphocyte cytosolic protein 2 (SH2 domain containing leukocyte protein of 76 kDa) (LCP2), mRNA.	1844
LIG1	NM_000234	<i>Homo sapiens</i> DNA ligase 1 (LIG1), transcript variant 1, mRNA.	1845
LIG4	NM_001098268	<i>Homo sapiens</i> DNA ligase 4 (LIG4), transcript variant 3, mRNA.	1846
LIG4	NM_002312	<i>Homo sapiens</i> DNA ligase 4 (LIG4), transcript variant 1, mRNA.	1847
LIG4	NM_0206937	<i>Homo sapiens</i> DNA ligase 4 (LIG4), transcript variant 2, mRNA.	1848
LRBA	NM_001199282	<i>Homo sapiens</i> LPS responsive beige-like anchor protein (LRBA), transcript variant 1, mRNA.	1849
LRBA	NM_006726	<i>Homo sapiens</i> LPS responsive beige-like anchor protein (LRBA), transcript variant 2, mRNA.	1850
LYST	NM_000081	<i>Homo sapiens</i> lysosomal trafficking regulator (LYST), transcript variant 1, mRNA.	1851
MAGEA9	NM_005365	<i>Homo sapiens</i> MAGE family member A9 (MAGEA9), mRNA.	1852
MAGEA9B	NM_001080790	<i>Homo sapiens</i> MAGE family member A9B (MAGEA9B), mRNA.	1853
MAGT1	NM_032121	<i>Homo sapiens</i> magnesium transporter 1 (MAGT1), mRNA.	1854
MALT1	NM_006785	<i>Homo sapiens</i> MALT1 paracaspase (MALT1), transcript variant 1, mRNA.	1855
MALT1	NM_173844	<i>Homo sapiens</i> MALT1 paracaspase (MALT1), transcript variant 2, mRNA.	1856
MAP3K2	NM_006609	<i>Homo sapiens</i> mitogen-activated protein kinase kinase kinase 2 (MAP3K2), mRNA.	1857
MAPK1	NM_002745	<i>Homo sapiens</i> mitogen-activated protein kinase 1 (MAPK1), transcript variant 1, mRNA.	1858
MAPK1	NM_138957	<i>Homo sapiens</i> mitogen-activated protein kinase 1 (MAPK1), transcript variant 2, mRNA.	1859
MAPK3	NM_001040056	<i>Homo sapiens</i> mitogen-activated protein kinase 3 (MAPK3), transcript variant 2, mRNA.	1860
MAPK3	NM_001109891	<i>Homo sapiens</i> mitogen-activated protein kinase 3 (MAPK3), transcript variant 3, mRNA.	1861
MAPK3	NM_002746	<i>Homo sapiens</i> mitogen-activated protein kinase 3 (MAPK3), transcript variant 1, mRNA.	1862
MAVS	NM_020746	<i>Homo sapiens</i> mitochondrial antiviral signaling protein (MAVS), transcript variant 1, mRNA.	1863
MAVS	NM_001206491	<i>Homo sapiens</i> mitochondrial antiviral signaling protein (MAVS), transcript variant 3, mRNA.	1864
MAVS	NR_037921	<i>Homo sapiens</i> mitochondrial antiviral signaling protein (MAVS), transcript variant 2, non-coding RNA.	1865
MECP2	NM_004992	<i>Homo sapiens</i> methyl-CpG binding protein 2 (MECP2), transcript variant 1, mRNA.	1866
MECP2	NM_001110792	<i>Homo sapiens</i> methyl-CpG binding protein 2 (MECP2), transcript variant 2, mRNA.	1867
MEX3C	NM_016626	<i>Homo sapiens</i> mex-3 RNA binding family member C (MEX3C), mRNA.	1868
MRE11A	NM_005590	<i>Homo sapiens</i> MRE11 homolog A, double strand break repair nuclease (MRE11A), transcript variant 2, mRNA.	1869
MRE11A	NM_005591	<i>Homo sapiens</i> MRE11 homolog A, double strand break repair nuclease (MRE11A), transcript variant 1, mRNA.	1870
MS4A1	NM_021950	<i>Homo sapiens</i> membrane spanning 4-domains A1 (MS4A1), transcript variant 3, mRNA.	1871
MS4A1	NM_152866	<i>Homo sapiens</i> membrane spanning 4-domains A1 (MS4A1), transcript variant 1, mRNA.	1872
MSN	NM_002444	<i>Homo sapiens</i> moesin (MSN), mRNA.	1873
MYD88	NM_001172566	<i>Homo sapiens</i> myeloid differentiation primary response 88 (MYD88), transcript variant 5, mRNA.	1874
MYD88	NM_001172567	<i>Homo sapiens</i> myeloid differentiation primary response 88 (MYD88), transcript variant 1, mRNA.	1875
MYD88	NM_001172568	<i>Homo sapiens</i> myeloid differentiation primary response 88 (MYD88), transcript variant 3, mRNA.	1876

TABLE 12-continued

Non-redundant list of transcript variants that correspond to the set of genes that no CNV 'solutions' have been reported in the 71 PML cases			
RefSeq_Gene_ Symbol	RefSeq_Accession_ Number	mRNA_Description	SEQ ID
MYD88	NM_001172569	<i>Homo sapiens</i> myeloid differentiation primary response 88 (MYD88), transcript variant 4, mRNA.	1877
MYD88	NM_002468	<i>Homo sapiens</i> myeloid differentiation primary response 88 (MYD88), transcript variant 2, mRNA.	1878
NBN	NM_002485	<i>Homo sapiens</i> nibrin (NBN), mRNA.	1879
NFIC	NM_001245005	<i>Homo sapiens</i> nuclear factor I C (NFIC), transcript variant 4, mRNA.	1880
NFIC	NM_0205843	<i>Homo sapiens</i> nuclear factor I C (NFIC), transcript variant 2, mRNA.	1881
NFIC	NM_001245002	<i>Homo sapiens</i> nuclear factor I C (NFIC), transcript variant 1, mRNA.	1882
NFIC	NM_001245004	<i>Homo sapiens</i> nuclear factor I C (NFIC), transcript variant 3, mRNA.	1883
NFIC	NM_005597	<i>Homo sapiens</i> nuclear factor I C (NFIC), transcript variant 5, mRNA.	1884
NFKB1	NM_003998	<i>Homo sapiens</i> nuclear factor of kappa light polypeptide gene enhancer in B-cells 1 (NFKB1), transcript variant 1, mRNA.	1885
NFKB1	NM_001165412	<i>Homo sapiens</i> nuclear factor of kappa light polypeptide gene enhancer in B-cells 1 (NFKB1), transcript variant 2, mRNA.	1886
NFKB2	NM_001077494	<i>Homo sapiens</i> nuclear factor of kappa light polypeptide gene enhancer in B-cells 2 (NFKB2), transcript variant 1, mRNA.	1887
NFKB2	NM_002502	<i>Homo sapiens</i> nuclear factor of kappa light polypeptide gene enhancer in B-cells 2 (NFKB2), transcript variant 2, mRNA.	1888
NFKB2	NM_001261403	<i>Homo sapiens</i> nuclear factor of kappa light polypeptide gene enhancer in B-cells 2 (NFKB2), transcript variant 4, mRNA.	1889
NFKBIA	NM_020529	<i>Homo sapiens</i> NFKB inhibitor alpha (NFKBIA), mRNA.	1890
NHEJ1	NM_024782	<i>Homo sapiens</i> non-homologous end joining factor 1 (NHEJ1), mRNA.	1891
NLRP3	NM_183395	<i>Homo sapiens</i> NLR family, pyrin domain containing 3 (NLRP3), transcript variant 2, mRNA.	1892
NLRP3	NM_004895	<i>Homo sapiens</i> NLR family, pyrin domain containing 3 (NLRP3), transcript variant 1, mRNA.	1893
NLRP3	NM_001127462	<i>Homo sapiens</i> NLR family, pyrin domain containing 3 (NLRP3), transcript variant 5, mRNA.	1894
NLRP3	NM_001127461	<i>Homo sapiens</i> NLR family, pyrin domain containing 3 (NLRP3), transcript variant 4, mRNA.	1895
NLRP3	NM_001079821	<i>Homo sapiens</i> NLR family, pyrin domain containing 3 (NLRP3), transcript variant 3, mRNA.	1896
NLRP3	NM_001243133	<i>Homo sapiens</i> NLR family, pyrin domain containing 3 (NLRP3), transcript variant 6, mRNA.	1897
NOD2	NM_022162	<i>Homo sapiens</i> nucleotide-binding oligomerization domain containing 2 (NOD2), mRNA.	1898
ORAI1	NM_032790	<i>Homo sapiens</i> ORAI calcium release-activated calcium modulator 1 (ORAI1), mRNA.	1899
OSTM1	NM_014028	<i>Homo sapiens</i> osteopetrosis associated transmembrane protein 1 (OSTM1), mRNA.	1900
PGM3	NM_001199917	<i>Homo sapiens</i> phosphoglucomutase 3 (PGM3), transcript variant 1, mRNA.	1901
PGM3	NM_001199918	<i>Homo sapiens</i> phosphoglucomutase 3 (PGM3), transcript variant 3, mRNA.	1902
PGM3	NM_015599	<i>Homo sapiens</i> phosphoglucomutase 3 (PGM3), transcript variant 2, mRNA.	1903
PGM3	NM_001199919	<i>Homo sapiens</i> phosphoglucomutase 3 (PGM3), transcript variant 4, mRNA.	1904
PIAS1	NM_016166	<i>Homo sapiens</i> protein inhibitor of activated STAT 1 (PIAS1), transcript variant 2, mRNA.	1905
PIK3R1	NM_181523	<i>Homo sapiens</i> phosphoinositide-3-kinase regulatory subunit 1 (PIK3R1), transcript variant 1, mRNA.	1906
PIK3R1	NM_181524	<i>Homo sapiens</i> phosphoinositide-3-kinase regulatory subunit 1 (PIK3R1), transcript variant 3, mRNA.	1907
PIK3R1	NM_181504	<i>Homo sapiens</i> phosphoinositide-3-kinase regulatory subunit 1 (PIK3R1), transcript variant 2, mRNA.	1908
PIK3R1	NM_001242466	<i>Homo sapiens</i> phosphoinositide-3-kinase regulatory subunit 1 (PIK3R1), transcript variant 4, mRNA.	1909
PLCG2	NM_002661	<i>Homo sapiens</i> phospholipase C gamma 2 (PLCG2), mRNA.	1910
PMS2	NM_000535	<i>Homo sapiens</i> PMS1 homolog 2, mismatch repair system component (PMS2), transcript variant 1, mRNA.	1911
PNP	NM_000270	<i>Homo sapiens</i> purine nucleoside phosphorylase (PNP), mRNA.	1912
POLA1	NM_016937	<i>Homo sapiens</i> polymerase (DNA directed), alpha 1, catalytic subunit (POLA1), mRNA.	1913
POLE	NM_006231	<i>Homo sapiens</i> DNA polymerase epsilon, catalytic subunit (POLE), mRNA.	1914
PRF1	NM_001083116	<i>Homo sapiens</i> perforin 1 (PRF1), transcript variant 2, mRNA.	1915
PRF1	NM_005041	<i>Homo sapiens</i> perforin 1 (PRF1), transcript variant 1, mRNA.	1916
PRKCD	NM_006254	<i>Homo sapiens</i> protein kinase C delta (PRKCD), transcript variant 1, mRNA.	1917
PRKCD	NM_212539	<i>Homo sapiens</i> protein kinase C delta (PRKCD), transcript variant 2, mRNA.	1918
PRKDC	NM_001081640	<i>Homo sapiens</i> protein kinase, DNA-activated, catalytic polypeptide (PRKDC), transcript variant 2, mRNA.	1919
PRKDC	NM_006904	<i>Homo sapiens</i> protein kinase, DNA-activated, catalytic polypeptide (PRKDC), transcript variant 1, mRNA.	1920
PROC	NM_000312	<i>Homo sapiens</i> protein C, inactivator of coagulation factors Va and VIIIa (PROC), mRNA.	1921
PSMB8	NM_004159	<i>Homo sapiens</i> proteasome (prosome, macropain) subunit, beta type, 8 (PSMB8), transcript variant 1, mRNA.	1922
PSMB8	NM_148919	<i>Homo sapiens</i> proteasome (prosome, macropain) subunit, beta type, 8 (PSMB8), transcript variant 2, mRNA.	1923
PTEN	NM_000314	<i>Homo sapiens</i> phosphatase and tensin homolog (PTEN), transcript variant 1, mRNA.	1924
PTPRC	NM_001267798	<i>Homo sapiens</i> protein tyrosine phosphatase, receptor type C (PTPRC), transcript variant 5, mRNA.	1925

TABLE 12-continued

Non-redundant list of transcript variants that correspond to the set of genes that no CNV 'solutions' have been reported in the 71 PML cases			
RefSeq_Gene_ Symbol	RefSeq_Accession_ Number	mRNA_Description	SEQ ID
PTPRC	NM_002838	<i>Homo sapiens</i> protein tyrosine phosphatase, receptor type C (PTPRC), transcript variant 1, mRNA.	1926
PTPRC	NM_080921	<i>Homo sapiens</i> protein tyrosine phosphatase, receptor type C (PTPRC), transcript variant 2, mRNA.	1927
PTPRC	NR_052021	<i>Homo sapiens</i> protein tyrosine phosphatase, receptor type C (PTPRC), transcript variant 4, non-coding RNA.	1928
PURA	NM_005859	<i>Homo sapiens</i> purine rich element binding protein A (PURA), mRNA.	1929
RAB27A	NM_183235	<i>Homo sapiens</i> RAB27A, member RAS oncogene family (RAB27A), transcript variant 3, mRNA.	1930
RAB27A	NM_183236	<i>Homo sapiens</i> RAB27A, member RAS oncogene family (RAB27A), transcript variant 4, mRNA.	1931
RAB27A	NM_004580	<i>Homo sapiens</i> RAB27A, member RAS oncogene family (RAB27A), transcript variant 1, mRNA.	1932
RAB27A	NM_183234	<i>Homo sapiens</i> RAB27A, member RAS oncogene family (RAB27A), transcript variant 2, mRNA.	1933
RAB7A	NM_004637	<i>Homo sapiens</i> RAB7A, member RAS oncogene family (RAB7A), mRNA.	1934
RABGEF1	NM_014504	<i>Homo sapiens</i> RAB guanine nucleotide exchange factor (GEF) 1 (RABGEF1), transcript variant 4, mRNA.	1935
RAC2	NM_002872	<i>Homo sapiens</i> ras-related C3 botulinum toxin substrate 2 (rho family, small GTP binding protein Rac2) (RAC2), mRNA.	1936
RAD51	NM_001164270	<i>Homo sapiens</i> RAD51 recombinase (RAD51), transcript variant 3, mRNA.	1937
RAD51	NM_002875	<i>Homo sapiens</i> RAD51 recombinase (RAD51), transcript variant 1, mRNA.	1938
RAD51	NM_133487	<i>Homo sapiens</i> RAD51 recombinase (RAD51), transcript variant 2, mRNA.	1939
RAD51	NM_001164269	<i>Homo sapiens</i> RAD51 recombinase (RAD51), transcript variant 4, mRNA.	1940
RAG1	NM_000448	<i>Homo sapiens</i> recombination activating gene 1 (RAG1), mRNA.	1941
RAG2	NM_000536	<i>Homo sapiens</i> recombination activating gene 2 (RAG2), transcript variant 1, mRNA.	1942
RAG2	NM_001243785	<i>Homo sapiens</i> recombination activating gene 2 (RAG2), transcript variant 3, mRNA.	1943
RAG2	NM_001243786	<i>Homo sapiens</i> recombination activating gene 2 (RAG2), transcript variant 4, mRNA.	1944
RBCK1	NM_006462	<i>Homo sapiens</i> RANBP2-type and C3HC4-type zinc finger containing 1 (RBCK1), transcript variant 1, mRNA.	1945
RBCK1	NM_031229	<i>Homo sapiens</i> RANBP2-type and C3HC4-type zinc finger containing 1 (RBCK1), transcript variant 2, mRNA.	1946
RFX5	NM_000449	<i>Homo sapiens</i> regulatory factor X5 (RFX5), transcript variant 1, mRNA.	1947
RFX5	NM_001025603	<i>Homo sapiens</i> regulatory factor X5 (RFX5), transcript variant 2, mRNA.	1948
RFXANK	NM_003721	<i>Homo sapiens</i> regulatory factor X associated ankyrin containing protein (RFXANK), transcript variant 1, mRNA.	1949
RFXANK	NM_134440	<i>Homo sapiens</i> regulatory factor X associated ankyrin containing protein (RFXANK), transcript variant 2, mRNA.	1950
RFXAP	NM_000538	<i>Homo sapiens</i> regulatory factor X associated protein (RFXAP), mRNA.	1951
RIPK1	NM_003804	<i>Homo sapiens</i> receptor (TNFRSF)-interacting serine-threonine kinase 1 (RIPK1), mRNA.	1952
RIPK3	NM_006871	<i>Homo sapiens</i> receptor-interacting serine-threonine kinase 3 (RIPK3), mRNA.	1953
RMRP	NR_003051	<i>Homo sapiens</i> RNA component of mitochondrial RNA processing endoribonuclease (RMRP), RNase MRP RNA.	1954
RNASEH2A	NM_006397	<i>Homo sapiens</i> ribonuclease H2, subunit A (RNASEH2A), mRNA.	1955
RNASEH2B	NM_001142279	<i>Homo sapiens</i> ribonuclease H2, subunit B (RNASEH2B), transcript variant 2, mRNA.	1956
RNASEH2B	NM_024570	<i>Homo sapiens</i> ribonuclease H2, subunit B (RNASEH2B), transcript variant 1, mRNA.	1957
RNASEH2C	NM_032193	<i>Homo sapiens</i> ribonuclease H2, subunit C (RNASEH2C), mRNA.	1958
RNASEL	NM_021133	<i>Homo sapiens</i> ribonuclease L (2',5'-oligoadenylate synthetase-dependent) (RNASEL), mRNA.	1959
RNF168	NM_152617	<i>Homo sapiens</i> ring finger protein 168 (RNF168), mRNA.	1960
RNF31	NM_017999	<i>Homo sapiens</i> ring finger protein 31 (RNF31), mRNA.	1961
RNU4ATAC	NR_023343	<i>Homo sapiens</i> RNA, U4atac small nuclear (U12-dependent splicing) (RNU4ATAC), small nuclear RNA.	1962
RTEL1	NM_016434	<i>Homo sapiens</i> regulator of telomere elongation helicase 1 (RTEL1), transcript variant 1, mRNA.	1963
RTEL1	NM_032957	<i>Homo sapiens</i> regulator of telomere elongation helicase 1 (RTEL1), transcript variant 2, mRNA.	1964
RTEL1-TNFRSF6B	NR_037882	<i>Homo sapiens</i> RTEL1-TNFRSF6B readthrough (NMD candidate) (RTEL1-TNFRSF6B), long non-coding RNA.	1965
SALL2	NM_005407	<i>Homo sapiens</i> spalt like transcription factor 2 (SALL2), transcript variant 1, mRNA.	1966
SAMHD1	NM_015474	<i>Homo sapiens</i> SAM domain and HD domain 1 (SAMHD1), mRNA.	1967
SBDS	NM_016038	<i>Homo sapiens</i> Shwachman-Bodian-Diamond syndrome (SBDS), mRNA.	1968
SH2D1A	NM_001114937	<i>Homo sapiens</i> SH2 domain containing 1A (SH2D1A), transcript variant 2, mRNA.	1969
SH2D1A	NM_002351	<i>Homo sapiens</i> SH2 domain containing 1A (SH2D1A), transcript variant 1, mRNA.	1970
SHARPIN	NM_030974	<i>Homo sapiens</i> SHANK-associated RH domain interactor (SHARPIN), transcript variant 1, mRNA.	1971
SHARPIN	NR_038270	<i>Homo sapiens</i> SHANK-associated RH domain interactor (SHARPIN), transcript variant 2, non-coding RNA.	1972
SKIV2L	NM_006929	<i>Homo sapiens</i> superkiller viralicidic activity 2-like (<i>S. cerevisiae</i>) (SKIV2L), mRNA.	1973
SLC37A4	NM_001164277	<i>Homo sapiens</i> solute carrier family 37 (glucose-6-phosphate transporter), member 4 (SLC37A4), transcript variant 1, mRNA.	1974

TABLE 12-continued

Non-redundant list of transcript variants that correspond to the set of genes that no CNV 'solutions' have been reported in the 71 PML cases			
RefSeq_Gene_ Symbol	RefSeq_Accession_ Number	mRNA_Description	SEQ ID
SLC37A4	NM_001164278	<i>Homo sapiens</i> solute carrier family 37 (glucose-6-phosphate transporter), member 4 (SLC37A4), transcript variant 2, mRNA.	1975
SLC37A4	NM_001164279	<i>Homo sapiens</i> solute carrier family 37 (glucose-6-phosphate transporter), member 4 (SLC37A4), transcript variant 3, mRNA.	1976
SLC37A4	NM_001467	<i>Homo sapiens</i> solute carrier family 37 (glucose-6-phosphate transporter), member 4 (SLC37A4), transcript variant 4, mRNA.	1977
SLC37A4	NM_001164280	<i>Homo sapiens</i> solute carrier family 37 (glucose-6-phosphate transporter), member 4 (SLC37A4), transcript variant 5, mRNA.	1978
SLC46A1	NM_001242366	<i>Homo sapiens</i> solute carrier family 46 member 1 (SLC46A1), transcript variant 2, mRNA.	1979
SLC46A1	NM_080669	<i>Homo sapiens</i> solute carrier family 46 member 1 (SLC46A1), transcript variant 1, mRNA.	1980
SLC8A1	NM_001112800	<i>Homo sapiens</i> solute carrier family 8 member A1 (SLC8A1), transcript variant B, mRNA.	1981
SLC8A1	NM_001112801	<i>Homo sapiens</i> solute carrier family 8 member A1 (SLC8A1), transcript variant C, mRNA.	1982
SLC8A1	NM_001112802	<i>Homo sapiens</i> solute carrier family 8 member A1 (SLC8A1), transcript variant D, mRNA.	1983
SLC8A1	NM_001252624	<i>Homo sapiens</i> solute carrier family 8 member A1 (SLC8A1), transcript variant E, mRNA.	1984
SLC8A1	NM_021097	<i>Homo sapiens</i> solute carrier family 8 member A1 (SLC8A1), transcript variant A, mRNA.	1985
SMAD2	NM_001003652	<i>Homo sapiens</i> SMAD family member 2 (SMAD2), transcript variant 2, mRNA.	1986
SMAD2	NM_001135937	<i>Homo sapiens</i> SMAD family member 2 (SMAD2), transcript variant 3, mRNA.	1987
SMAD2	NM_005901	<i>Homo sapiens</i> SMAD family member 2 (SMAD2), transcript variant 1, mRNA.	1988
SMAD3	NM_005902	<i>Homo sapiens</i> SMAD family member 3 (SMAD3), transcript variant 1, mRNA.	1989
SMAD3	NM_001145102	<i>Homo sapiens</i> SMAD family member 3 (SMAD3), transcript variant 2, mRNA.	1990
SMAD3	NM_001145103	<i>Homo sapiens</i> SMAD family member 3 (SMAD3), transcript variant 3, mRNA.	1991
SMAD3	NM_001145104	<i>Homo sapiens</i> SMAD family member 3 (SMAD3), transcript variant 4, mRNA.	1992
SMAD4	NM_005359	<i>Homo sapiens</i> SMAD family member 4 (SMAD4), mRNA.	1993
SNAP29	NM_004782	<i>Homo sapiens</i> synaptosomal-associated protein, 29 kDa (SNAP29), mRNA.	1994
SNAR-A1	NR_004435	<i>Homo sapiens</i> small ILF3/NF90-associated RNA A1 (SNAR-A1), small nuclear RNA.	1995
SNAR-A10	NR_024229	<i>Homo sapiens</i> small ILF3/NF90-associated RNA A10 (SNAR-A10), small nuclear RNA.	1996
SNAR-A11	NR_024225	<i>Homo sapiens</i> small ILF3/NF90-associated RNA A11 (SNAR-A11), small nuclear RNA.	1997
SNAR-A12	NR_004437	<i>Homo sapiens</i> small ILF3/NF90-associated RNA A12 (SNAR-A12), small nuclear RNA.	1998
SNAR-A13	NR_024216	<i>Homo sapiens</i> small ILF3/NF90-associated RNA A13 (SNAR-A13), small nuclear RNA.	1999
SNAR-A14	NR_024242	<i>Homo sapiens</i> small ILF3/NF90-associated RNA A14 (SNAR-A14), small nuclear RNA.	2000
SNAR-A2	NR_004436	<i>Homo sapiens</i> small ILF3/NF90-associated RNA A2 (SNAR-A2), small nuclear RNA.	2001
SNAR-A3	NR_024214	<i>Homo sapiens</i> small ILF3/NF90-associated RNA A3 (SNAR-A3), small nuclear RNA.	2002
SNAR-A4	NR_024215	<i>Homo sapiens</i> small ILF3/NF90-associated RNA A4 (SNAR-A4), small nuclear RNA.	2003
SNAR-A5	NR_024223	<i>Homo sapiens</i> small ILF3/NF90-associated RNA A5 (SNAR-A5), small nuclear RNA.	2004
SNAR-A6	NR_024227	<i>Homo sapiens</i> small ILF3/NF90-associated RNA A6 (SNAR-A6), small nuclear RNA.	2005
SNAR-A7	NR_024224	<i>Homo sapiens</i> small ILF3/NF90-associated RNA A7 (SNAR-A7), small nuclear RNA.	2006
SNAR-A8	NR_024228	<i>Homo sapiens</i> small ILF3/NF90-associated RNA A8 (SNAR-A8), small nuclear RNA.	2007
SNAR-A9	NR_024226	<i>Homo sapiens</i> small ILF3/NF90-associated RNA A9 (SNAR-A9), small nuclear RNA.	2008
SNAR-B1	NR_024231	<i>Homo sapiens</i> small ILF3/NF90-associated RNA B1 (SNAR-B1), small nuclear RNA.	2009
SNAR-B2	NR_024230	<i>Homo sapiens</i> small ILF3/NF90-associated RNA B2 (SNAR-B2), small nuclear RNA.	2010
SNAR-C1	NR_024220	<i>Homo sapiens</i> small ILF3/NF90-associated RNA C1 (SNAR-C1), small nuclear RNA.	2011
SNAR-C2	NR_024217	<i>Homo sapiens</i> small ILF3/NF90-associated RNA C2 (SNAR-C2), small nuclear RNA.	2012
SNAR-C3	NR_024221	<i>Homo sapiens</i> small ILF3/NF90-associated RNA C3 (SNAR-C3), small nuclear RNA.	2013
SNAR-C4	NR_024218	<i>Homo sapiens</i> small ILF3/NF90-associated RNA C4 (SNAR-C4), small nuclear RNA.	2014
SNAR-C5	NR_024219	<i>Homo sapiens</i> small ILF3/NF90-associated RNA C5 (SNAR-C5), small nuclear RNA.	2015
SNAR-D	NR_024243	<i>Homo sapiens</i> small ILF3/NF90-associated RNA D (SNAR-D), small nuclear RNA.	2016
SNAR-E	NR_024258	<i>Homo sapiens</i> small ILF3/NF90-associated RNA E (SNAR-E), small nuclear RNA.	2017
SNAR-F	NR_004384	<i>Homo sapiens</i> small ILF3/NF90-associated RNA F (SNAR-F), small nuclear RNA.	2018
SNAR-G1	NR_004383	<i>Homo sapiens</i> small ILF3/NF90-associated RNA G1 (SNAR-G1), small nuclear RNA.	2019
SNAR-G2	NR_024244	<i>Homo sapiens</i> small ILF3/NF90-associated RNA G2 (SNAR-G2), small nuclear RNA.	2020
SNAR-H	NR_024342	<i>Homo sapiens</i> small ILF3/NF90-associated RNA H (SNAR-H), small nuclear RNA.	2021
SNAR-I	NR_024343	<i>Homo sapiens</i> small ILF3/NF90-associated RNA I (SNAR-I), small nuclear RNA.	2022
SNCA	NM_000345	<i>Homo sapiens</i> synuclein, alpha (non A4 component of amyloid precursor) (SNCA), transcript variant 1, mRNA.	2023
SNCA	NM_001146054	<i>Homo sapiens</i> synuclein, alpha (non A4 component of amyloid precursor) (SNCA), transcript variant 2, mRNA.	2024
SNCA	NM_001146055	<i>Homo sapiens</i> synuclein, alpha (non A4 component of amyloid precursor) (SNCA), transcript variant 3, mRNA.	2025
SNCA	NM_007308	<i>Homo sapiens</i> synuclein, alpha (non A4 component of amyloid precursor) (SNCA), transcript variant 4, mRNA.	2026
SNX10	NM_013322	<i>Homo sapiens</i> sorting nexin 10 (SNX10), transcript variant 2, mRNA.	2027
SNX10	NM_001199835	<i>Homo sapiens</i> sorting nexin 10 (SNX10), transcript variant 1, mRNA.	2028
SNX10	NM_001199837	<i>Homo sapiens</i> sorting nexin 10 (SNX10), transcript variant 3, mRNA.	2029
SNX10	NM_001199838	<i>Homo sapiens</i> sorting nexin 10 (SNX10), transcript variant 4, mRNA.	2030
SNX10	NR_037670	<i>Homo sapiens</i> sorting nexin 10 (SNX10), transcript variant 5, non-coding RNA.	2031
SP110	NM_004509	<i>Homo sapiens</i> SP110 nuclear body protein (SP110), transcript variant a, mRNA.	2032
SP110	NM_080424	<i>Homo sapiens</i> SP110 nuclear body protein (SP110), transcript variant c, mRNA.	2033
SP110	NM_001185015	<i>Homo sapiens</i> SP110 nuclear body protein (SP110), transcript variant d, mRNA.	2034
SP110	NM_004510	<i>Homo sapiens</i> SP110 nuclear body protein (SP110), transcript variant b, mRNA.	2035
SP140	NM_001005176	<i>Homo sapiens</i> SP140 nuclear body protein (SP140), transcript variant 2, mRNA.	2036
SP140	NM_007237	<i>Homo sapiens</i> SP140 nuclear body protein (SP140), transcript variant 1, mRNA.	2037

TABLE 12-continued

Non-redundant list of transcript variants that correspond to the set of genes that no CNV 'solutions' have been reported in the 71 PML cases			
RefSeq_Gene_ Symbol	RefSeq_Accession_ Number	mRNA_Description	SEQ ID
SPINK5	NM_001127698	<i>Homo sapiens</i> serine peptidase inhibitor, Kazal type 5 (SPINK5), transcript variant 1, mRNA.	2038
SPINK5	NM_006846	<i>Homo sapiens</i> serine peptidase inhibitor, Kazal type 5 (SPINK5), transcript variant 2, mRNA.	2039
SPINK5	NM_001127699	<i>Homo sapiens</i> serine peptidase inhibitor, Kazal type 5 (SPINK5), transcript variant 3, mRNA.	2040
SQSTM1	NM_003900	<i>Homo sapiens</i> sequestosome 1 (SQSTM1), transcript variant 1, mRNA.	2041
SQSTM1	NM_001142298	<i>Homo sapiens</i> sequestosome 1 (SQSTM1), transcript variant 2, mRNA.	2042
SQSTM1	NM_001142299	<i>Homo sapiens</i> sequestosome 1 (SQSTM1), transcript variant 3, mRNA.	2043
SRSF1	NM_001078166	<i>Homo sapiens</i> serine and arginine rich splicing factor 1 (SRSF1), transcript variant 2, mRNA.	2044
SRSF1	NM_006924	<i>Homo sapiens</i> serine and arginine rich splicing factor 1 (SRSF1), transcript variant 1, mRNA.	2045
SRSF1	NR_034041	<i>Homo sapiens</i> serine and arginine rich splicing factor 1 (SRSF1), transcript variant 3, non-coding RNA.	2046
STAT1	NM_007315	<i>Homo sapiens</i> signal transducer and activator of transcription 1 (STAT1), transcript variant alpha, mRNA.	2047
STAT1	NM_139266	<i>Homo sapiens</i> signal transducer and activator of transcription 1 (STAT1), transcript variant beta, mRNA.	2048
STAT2	NM_005419	<i>Homo sapiens</i> signal transducer and activator of transcription 2, 113 kDa (STAT2), transcript variant 1, mRNA.	2049
STAT2	NM_198332	<i>Homo sapiens</i> signal transducer and activator of transcription 2, 113 kDa (STAT2), transcript variant 2, mRNA.	2050
STAT3	NM_003150	<i>Homo sapiens</i> signal transducer and activator of transcription 3 (STAT3), transcript variant 2, mRNA.	2051
STAT3	NM_139276	<i>Homo sapiens</i> signal transducer and activator of transcription 3 (STAT3), transcript variant 1, mRNA.	2052
STAT3	NM_213662	<i>Homo sapiens</i> signal transducer and activator of transcription 3 (STAT3), transcript variant 3, mRNA.	2053
STAT5B	NM_012448	<i>Homo sapiens</i> signal transducer and activator of transcription 5B (STAT5B), mRNA.	2054
STIM1	NM_003156	<i>Homo sapiens</i> stromal interaction molecule 1 (STIM1), transcript variant 2, mRNA.	2055
STK4	NM_006282	<i>Homo sapiens</i> serine/threonine kinase 4 (STK4), mRNA.	2056
STX11	NM_003764	<i>Homo sapiens</i> syntaxin 11 (STX11), mRNA.	2057
STXBP2	NM_001127396	<i>Homo sapiens</i> syntaxin binding protein 2 (STXBP2), transcript variant 2, mRNA.	2058
STXBP2	NM_001272034	<i>Homo sapiens</i> syntaxin binding protein 2 (STXBP2), transcript variant 3, mRNA.	2059
STXBP2	NM_006949	<i>Homo sapiens</i> syntaxin binding protein 2 (STXBP2), transcript variant 1, mRNA.	2060
STXBP2	NR_073560	<i>Homo sapiens</i> syntaxin binding protein 2 (STXBP2), transcript variant 4, non-coding RNA.	2061
SYNCRIP	NM_001159673	<i>Homo sapiens</i> synaptotagmin binding cytoplasmic RNA interacting protein (SYNCRIP), transcript variant 2, mRNA.	2062
SYNCRIP	NM_001159674	<i>Homo sapiens</i> synaptotagmin binding cytoplasmic RNA interacting protein (SYNCRIP), transcript variant 3, mRNA.	2063
SYNCRIP	NM_001159676	<i>Homo sapiens</i> synaptotagmin binding cytoplasmic RNA interacting protein (SYNCRIP), transcript variant 5, mRNA.	2064
SYNCRIP	NM_001159677	<i>Homo sapiens</i> synaptotagmin binding cytoplasmic RNA interacting protein (SYNCRIP), transcript variant 6, mRNA.	2065
SYNCRIP	NM_001253771	<i>Homo sapiens</i> synaptotagmin binding cytoplasmic RNA interacting protein (SYNCRIP), transcript variant 7, mRNA.	2066
SYNCRIP	NM_001159675	<i>Homo sapiens</i> synaptotagmin binding cytoplasmic RNA interacting protein (SYNCRIP), transcript variant 4, mRNA.	2067
SYNCRIP	NM_006372	<i>Homo sapiens</i> synaptotagmin binding cytoplasmic RNA interacting protein (SYNCRIP), transcript variant 1, mRNA.	2068
T	NM_001270484	<i>Homo sapiens</i> T brachyury transcription factor (T), transcript variant 2, mRNA.	2069
T	NM_003181	<i>Homo sapiens</i> T brachyury transcription factor (T), transcript variant 1, mRNA.	2070
TAP1	NM_000593	<i>Homo sapiens</i> transporter 1, ATP binding cassette subfamily B member (TAP1), transcript variant 1, mRNA.	2071
TAP2	NM_018833	<i>Homo sapiens</i> transporter 2, ATP binding cassette subfamily B member (TAP2), transcript variant 2, mRNA.	2072
TAP2	NM_000544	<i>Homo sapiens</i> transporter 2, ATP binding cassette subfamily B member (TAP2), transcript variant 1, B allele, mRNA.	2073
TAPBP	NM_003190	<i>Homo sapiens</i> TAP binding protein (tapasin) (TAPBP), transcript variant 1, mRNA.	2074
TAPBP	NM_172209	<i>Homo sapiens</i> TAP binding protein (tapasin) (TAPBP), transcript variant 3, mRNA.	2075
TAPBP	NM_172208	<i>Homo sapiens</i> TAP binding protein (tapasin) (TAPBP), transcript variant 2, mRNA.	2076
TAZ	NM_000116	<i>Homo sapiens</i> tafazzin (TAZ), transcript variant 1, mRNA.	2077
TAZ	NM_181312	<i>Homo sapiens</i> tafazzin (TAZ), transcript variant 3, mRNA.	2078
TAZ	NM_181311	<i>Homo sapiens</i> tafazzin (TAZ), transcript variant 2, mRNA.	2079
TAZ	NM_181313	<i>Homo sapiens</i> tafazzin (TAZ), transcript variant 4, mRNA.	2080
TAZ	NR_024048	<i>Homo sapiens</i> tafazzin (TAZ), transcript variant 5, non-coding RNA.	2081
TBK1	NM_013254	<i>Homo sapiens</i> TANK binding kinase 1 (TBK1), mRNA.	2082
TBX1	NM_005992	<i>Homo sapiens</i> T-box 1 (TBX1), transcript variant B, mRNA.	2083
TBX1	NM_080646	<i>Homo sapiens</i> T-box 1 (TBX1), transcript variant A, mRNA.	2084
TBX1	NM_080647	<i>Homo sapiens</i> T-box 1 (TBX1), transcript variant C, mRNA.	2085

TABLE 12-continued

Non-redundant list of transcript variants that correspond to the set of genes that no CNV 'solutions' have been reported in the 71 PML cases			
RefSeq_Gene_ Symbol	RefSeq_Accession_ Number	mRNA_Description	SEQ ID
TCIRG1	NM_006019	<i>Homo sapiens</i> T-cell immune regulator 1, ATPase H+ transporting V0 subunit a3 (TCIRG1), transcript variant 1, mRNA.	2086
TCIRG1	NM_006053	<i>Homo sapiens</i> T-cell immune regulator 1, ATPase H+ transporting V0 subunit a3 (TCIRG1), transcript variant 2, mRNA.	2087
TICAM1	NM_182919	<i>Homo sapiens</i> toll like receptor adaptor molecule 1 (TICAM1), mRNA.	2088
TLR3	NM_003265	<i>Homo sapiens</i> toll like receptor 3 (TLR3), mRNA.	2089
TLR4	NM_003266	<i>Homo sapiens</i> toll like receptor 4 (TLR4), transcript variant 3, mRNA.	2090
TLR4	NM_138554	<i>Homo sapiens</i> toll like receptor 4 (TLR4), transcript variant 1, mRNA.	2091
TLR4	NM_138557	<i>Homo sapiens</i> toll like receptor 4 (TLR4), transcript variant 4, mRNA.	2092
TMEM173	NM_198282	<i>Homo sapiens</i> transmembrane protein 173 (TMEM173), mRNA.	2093
TNF	NM_000594	<i>Homo sapiens</i> tumor necrosis factor (TNF), mRNA.	2094
TNFAIP3	NM_001270507	<i>Homo sapiens</i> TNF alpha induced protein 3 (TNFAIP3), transcript variant 2, mRNA.	2095
TNFAIP3	NM_001270508	<i>Homo sapiens</i> TNF alpha induced protein 3 (TNFAIP3), transcript variant 1, mRNA.	2096
TNFAIP3	NM_006290	<i>Homo sapiens</i> TNF alpha induced protein 3 (TNFAIP3), transcript variant 3, mRNA.	2097
TNFRSF11A	NM_003839	<i>Homo sapiens</i> tumor necrosis factor receptor superfamily, member 11a, NFKB activator (TNFRSF11A), transcript variant 1, mRNA.	2098
TNFRSF11A	NM_001270949	<i>Homo sapiens</i> tumor necrosis factor receptor superfamily, member 11a, NFKB activator (TNFRSF11A), transcript variant 2, mRNA.	2099
TNFRSF11A	NM_001270950	<i>Homo sapiens</i> tumor necrosis factor receptor superfamily, member 11a, NFKB activator (TNFRSF11A), transcript variant 3, mRNA.	2100
TNFRSF11A	NM_001270951	<i>Homo sapiens</i> tumor necrosis factor receptor superfamily, member 11a, NFKB activator (TNFRSF11A), transcript variant 4, mRNA.	2101
TNFRSF11B	NM_002546	<i>Homo sapiens</i> tumor necrosis factor receptor superfamily, member 11b (TNFRSF11B), mRNA.	2102
TNFRSF13B	NM_012452	<i>Homo sapiens</i> TNF receptor superfamily member 13B (TNFRSF13B), mRNA.	2103
TNFRSF4	NM_003327	<i>Homo sapiens</i> TNF receptor superfamily member 4 (TNFRSF4), mRNA.	2104
TNFRSF8	NM_001243	<i>Homo sapiens</i> TNF receptor superfamily member 8 (TNFRSF8), transcript variant 1, mRNA.	2105
TNFSF11	NM_003701	<i>Homo sapiens</i> tumor necrosis factor (ligand) superfamily, member 11 (TNFSF11), transcript variant 1, mRNA.	2106
TNFSF11	NM_033012	<i>Homo sapiens</i> tumor necrosis factor (ligand) superfamily, member 11 (TNFSF11), transcript variant 2, mRNA.	2107
TNFSF12	NM_003809	<i>Homo sapiens</i> tumor necrosis factor superfamily member 12 (TNFSF12), transcript variant 1, mRNA.	2108
TNFSF12	NR_037146	<i>Homo sapiens</i> tumor necrosis factor superfamily member 12 (TNFSF12), transcript variant 2, non-coding RNA.	2109
TP53	NM_000546	<i>Homo sapiens</i> tumor protein p53 (TP53), transcript variant 1, mRNA.	2110
TP53	NM_001126112	<i>Homo sapiens</i> tumor protein p53 (TP53), transcript variant 2, mRNA.	2111
TP53	NM_001126113	<i>Homo sapiens</i> tumor protein p53 (TP53), transcript variant 4, mRNA.	2112
TP53	NM_001126114	<i>Homo sapiens</i> tumor protein p53 (TP53), transcript variant 3, mRNA.	2113
TP53	NM_001126115	<i>Homo sapiens</i> tumor protein p53 (TP53), transcript variant 5, mRNA.	2114
TP53	NM_001126116	<i>Homo sapiens</i> tumor protein p53 (TP53), transcript variant 6, mRNA.	2115
TP53	NM_001126117	<i>Homo sapiens</i> tumor protein p53 (TP53), transcript variant 7, mRNA.	2116
TP53	NM_001126118	<i>Homo sapiens</i> tumor protein p53 (TP53), transcript variant 8, mRNA.	2117
TP53	NM_001276695	<i>Homo sapiens</i> tumor protein p53 (TP53), transcript variant 4, mRNA.	2118
TP53	NM_001276696	<i>Homo sapiens</i> tumor protein p53 (TP53), transcript variant 3, mRNA.	2119
TP53	NM_001276697	<i>Homo sapiens</i> tumor protein p53 (TP53), transcript variant 5, mRNA.	2120
TP53	NM_001276698	<i>Homo sapiens</i> tumor protein p53 (TP53), transcript variant 6, mRNA.	2121
TP53	NM_001276699	<i>Homo sapiens</i> tumor protein p53 (TP53), transcript variant 7, mRNA.	2122
TP53	NM_001276760	<i>Homo sapiens</i> tumor protein p53 (TP53), transcript variant 1, mRNA.	2123
TP53	NM_001276761	<i>Homo sapiens</i> tumor protein p53 (TP53), transcript variant 2, mRNA.	2124
TRAF3	NM_001199427	<i>Homo sapiens</i> TNF receptor associated factor 3 (TRAF3), transcript variant 4, mRNA.	2125
TRAF3	NM_003300	<i>Homo sapiens</i> TNF receptor associated factor 3 (TRAF3), transcript variant 3, mRNA.	2126
TRAF3	NM_145725	<i>Homo sapiens</i> TNF receptor associated factor 3 (TRAF3), transcript variant 1, mRNA.	2127
TRAF3	NM_145726	<i>Homo sapiens</i> TNF receptor associated factor 3 (TRAF3), transcript variant 2, mRNA.	2128
TRAF6	NM_004620	<i>Homo sapiens</i> TNF receptor-associated factor 6, E3 ubiquitin protein ligase (TRAF6), transcript variant 2, mRNA.	2129
TRAF6	NM_145803	<i>Homo sapiens</i> TNF receptor-associated factor 6, E3 ubiquitin protein ligase (TRAF6), transcript variant 1, mRNA.	2130
TREX1	NM_007248	<i>Homo sapiens</i> three prime repair exonuclease 1 (TREX1), transcript variant 5, mRNA.	2131
TREX1	NM_033629	<i>Homo sapiens</i> three prime repair exonuclease 1 (TREX1), transcript variant 4, mRNA.	2132
TREX1	NM_016381	<i>Homo sapiens</i> three prime repair exonuclease 1 (TREX1), transcript variant 1, mRNA.	2133
TRNT1	NM_182916	<i>Homo sapiens</i> tRNA nucleotidyl transferase 1 (TRNT1), transcript variant 1, mRNA.	2134
TTC7A	NM_020458	<i>Homo sapiens</i> tetratricopeptide repeat domain 7A (TTC7A), transcript variant 2, mRNA.	2135
TYK2	NM_003331	<i>Homo sapiens</i> tyrosine kinase 2 (TYK2), mRNA.	2136
UNC119	NM_005148	<i>Homo sapiens</i> unc-119 lipid binding chaperone (UNC119), transcript variant 1, mRNA.	2137
UNC119	NM_054035	<i>Homo sapiens</i> unc-119 lipid binding chaperone (UNC119), transcript variant 2, mRNA.	2138
UNC13D	NM_199242	<i>Homo sapiens</i> unc-13 homolog D (UNC13D), mRNA.	2139
UNC93B1	NM_030930	<i>Homo sapiens</i> unc-93 homolog B1 (<i>C. elegans</i>) (UNC93B1), mRNA.	2140
UNG	NM_080911	<i>Homo sapiens</i> uracil DNA glycosylase (UNG), transcript variant 2, mRNA.	2141
UNG	NM_003362	<i>Homo sapiens</i> uracil DNA glycosylase (UNG), transcript variant 1, mRNA.	2142
USP18	NM_017414	<i>Homo sapiens</i> ubiquitin specific peptidase 18 (USP18), mRNA.	2143

TABLE 12-continued

Non-redundant list of transcript variants that correspond to the set of genes that no CNV 'solutions' have been reported in the 71 PML cases			
RefSeq_Gene_ Symbol	RefSeq_Accession_ Number	mRNA_Description	SEQ ID
USP20	NM_006676	<i>Homo sapiens</i> ubiquitin specific peptidase 20 (USP20), transcript variant 1, mRNA.	2144
USP20	NM_001008563	<i>Homo sapiens</i> ubiquitin specific peptidase 20 (USP20), transcript variant 2, mRNA.	2145
USP20	NM_001110303	<i>Homo sapiens</i> ubiquitin specific peptidase 20 (USP20), transcript variant 3, mRNA.	2146
VAPA	NM_003574	<i>Homo sapiens</i> VAMP associated protein A (VAPA), transcript variant 1, mRNA.	2147
VAPA	NM_194434	<i>Homo sapiens</i> VAMP associated protein A (VAPA), transcript variant 2, mRNA.	2148
VCP	NM_007126	<i>Homo sapiens</i> valosin containing protein (VCP), mRNA.	2149
VDAC1	NM_003374	<i>Homo sapiens</i> voltage dependent anion channel 1 (VDAC1), transcript variant 1, mRNA.	2150
VDAC1	NR_036624	<i>Homo sapiens</i> voltage dependent anion channel 1 (VDAC1), transcript variant 3, non-coding RNA.	2151
VDAC1	NR_036625	<i>Homo sapiens</i> voltage dependent anion channel 1 (VDAC1), transcript variant 2, non-coding RNA.	2152
VPS13B	NM_017890	<i>Homo sapiens</i> vacuolar protein sorting 13 homolog B (yeast) (VPS13B), transcript variant 5, mRNA.	2153
VPS13B	NM_181661	<i>Homo sapiens</i> vacuolar protein sorting 13 homolog B (yeast) (VPS13B), transcript variant 4, mRNA.	2154
VPS13B	NM_015243	<i>Homo sapiens</i> vacuolar protein sorting 13 homolog B (yeast) (VPS13B), transcript variant 3, mRNA.	2155
VPS13B	NR_047582	<i>Homo sapiens</i> vacuolar protein sorting 13 homolog B (yeast) (VPS13B), transcript variant 6, non-coding RNA.	2156
VPS13B	NM_152564	<i>Homo sapiens</i> vacuolar protein sorting 13 homolog B (yeast) (VPS13B), transcript variant 1, mRNA.	2157
VPS45	NM_007259	<i>Homo sapiens</i> vacuolar protein sorting 45 homolog (VPS45), transcript variant 1, mRNA.	2158
WAS	NM_000377	<i>Homo sapiens</i> Wiskott-Aldrich syndrome (WAS), mRNA.	2159
WEE1	NM_003390	<i>Homo sapiens</i> WEE1 G2 checkpoint kinase (WEE1), transcript variant 1, mRNA.	2160
WEE1	NM_001143976	<i>Homo sapiens</i> WEE1 G2 checkpoint kinase (WEE1), transcript variant 2, mRNA.	2161
WIPF1	NM_001077269	<i>Homo sapiens</i> WAS/WASL interacting protein family member 1 (WIPF1), transcript variant 2, mRNA.	2162
WIPF1	NM_003387	<i>Homo sapiens</i> WAS/WASL interacting protein family member 1 (WIPF1), transcript variant 1, mRNA.	2163
XIAP	NM_001204401	<i>Homo sapiens</i> X-linked inhibitor of apoptosis, E3 ubiquitin protein ligase (XIAP), transcript variant 2, mRNA.	2164
XIAP	NM_001167	<i>Homo sapiens</i> X-linked inhibitor of apoptosis, E3 ubiquitin protein ligase (XIAP), transcript variant 1, mRNA.	2165
XIAP	NR_037916	<i>Homo sapiens</i> X-linked inhibitor of apoptosis, E3 ubiquitin protein ligase (XIAP), transcript variant 3, non-coding RNA.	2166
YBX1	NM_004559	<i>Homo sapiens</i> Y-box binding protein 1 (YBX1), transcript variant 1, mRNA.	2167
YWHAZ	NM_001135699	<i>Homo sapiens</i> tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein zeta (YWHAZ), transcript variant 3, mRNA.	2168
YWHAZ	NM_001135700	<i>Homo sapiens</i> tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein zeta (YWHAZ), transcript variant 4, mRNA.	2169
YWHAZ	NM_001135701	<i>Homo sapiens</i> tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein zeta (YWHAZ), transcript variant 5, mRNA.	2170
YWHAZ	NM_001135702	<i>Homo sapiens</i> tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein zeta (YWHAZ), transcript variant 6, mRNA.	2171
YWHAZ	NM_003406	<i>Homo sapiens</i> tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein zeta (YWHAZ), transcript variant 1, mRNA.	2172
YWHAZ	NM_145690	<i>Homo sapiens</i> tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein zeta (YWHAZ), transcript variant 2, mRNA.	2173
ZAP70	NM_001079	<i>Homo sapiens</i> zeta chain of T cell receptor associated protein kinase 70 (ZAP70), transcript variant 1, mRNA.	2174
ZAP70	NM_207519	<i>Homo sapiens</i> zeta chain of T cell receptor associated protein kinase 70 (ZAP70), transcript variant 2, mRNA.	2175
ZBTB24	NM_014797	<i>Homo sapiens</i> zinc finger and BTB domain containing 24 (ZBTB24), transcript variant 1, mRNA.	2176
ZBTB24	NM_001164313	<i>Homo sapiens</i> zinc finger and BTB domain containing 24 (ZBTB24), transcript variant 2, mRNA.	2177

[0626] Table 12 lists all transcript variants for genes in Table 6 that were not ‘discovered’ by PBio on the basis of aCGH (CNV identified genes). The SEQ ID NOS correspond to transcript variants (oftentimes more than one per gene).

TABLE 13

Genes for which the total burden of heterozygous, damaging variants was found to be statistically greater in PML cases versus ExAC controls								
GENE	Ave CASES	Ave EXAC CASES	Ave EXAC SAMPLES	Ave FET	Ave FET corr(419)	Ave OR	Ethnicity	Overlap
PLCG2	17	1,806	31,277	1.43E-10	6.21E-08	10.27	EUR	EUR + AFR
RBCK1	6	187	29,324	4.27E-07	1.86E-04	24.60	EUR	
EPG5	9	764	32,835	7.11E-07	3.09E-04	10.79	EUR	
IL17F	4	61	33,346	1.67E-06	7.28E-04	54.57	EUR	
SHARPIN	8	646	32,162	2.58E-06	1.12E-03	10.84	EUR	EUR + AFR
PRF1	8	715	33,027	4.44E-06	1.93E-03	10.04	EUR	
JAGN1	5	163	27,768	6.80E-06	2.96E-03	21.71	EUR	
TAP1	5	203	28,125	1.80E-05	7.82E-03	17.63	EUR	
POLE	11	1,660	29,108	2.84E-05	1.23E-02	5.51	EUR	EUR + AFR
LRBA	11	1,876	32,136	3.47E-05	1.51E-02	5.38	EUR	
EHF	3	49	32,588	4.83E-05	2.10E-02	48.59	EUR	
IL12B	3	58	33,112	7.44E-05	3.23E-02	41.70	EUR	
ATL2	8	31	5,041	4.03E-11	1.75E-08	90.11	AFR	AFR
NHEJ1	6	27	4,384	5.48E-09	2.39E-06	64.56	AFR	
LYST	11	291	4,748	1.09E-08	4.76E-06	16.85	AFR	
HIVEP1	9	150	4,432	7.41E-08	3.22E-05	23.83	AFR	
AP3B1	5	46	4,937	1.69E-06	7.36E-04	33.23	AFR	EUR + AFR
TNFRSF10A	7	149	4,626	3.28E-06	1.43E-03	15.03	AFR	
PIK3CD	7	148	4,549	3.52E-06	1.53E-03	14.87	AFR	
PLCG2	8	256	4,410	1.47E-05	6.41E-03	9.99	AFR	
PNP	3	11	5,189	2.00E-05	8.69E-03	78.45	AFR	EUR + AFR
POLE	8	297	4,752	2.48E-05	1.08E-02	9.23	AFR	
MCEE	3	13	5,164	3.10E-05	1.35E-02	66.04	AFR	
DOCK2	6	173	5,023	6.39E-05	2.78E-02	11.21	AFR	
ALG12	4	43	4,252	6.73E-05	2.93E-02	23.03	AFR	

[0627] Table 13 lists genes for which the total burden of heterozygous, damaging variants was found to be statistically greater in PML cases versus ExAC controls. Gene burden analysis was performed as described below at minor allele frequency (MAF) cutoffs of 0.01, 0.02, 0.03, 0.04 and 0.05. Not all genes survived statistical analysis at all MAF cutoffs. For each gene that survived at multiple MAF cutoffs, the averages of the Fishers Exact Test (FET), nominal and

corrected, were calculated, as were the other relevant metrics. Two genes overlapped between AFR and EUR analyses. FETs were corrected for multiple testing with the number of genes used in this study (419). Only genes for which FET_corr was <0.05 and in which variants affected >10% of cases within the given ethnicity (>2 for AFR, >4 for EUR) were considered for inclusion.

TABLE 14

Top tier of variants found to be significant on the basis of variant burden analysis							
Gene	Variant (hg19)	Genotype	PML EUR	PML AFR	PML LAT	ExAC EUR	ExAC AFR
			44	21	5		
PLCG2	chr16: 81942175, A > G	het	2	5	0	512/32281	88/4707
IFIH1	chr2: 163136505, C > G	het	6	1	0	611/33155	23/5182
TCIRG1	chr11: 67818269, G > A	het	0	4	0	103/33193	200/5170
IGLL1	chr22: 23917192, G > T	het	4	3	1	751/33348	603/5183
MAVS	chr20: 3846397, C > T	hom	4	4	0	800/32122	684/4982

TABLE 14-continued

Top tier of variants found to be significant on the basis of variant burden analysis							
SHARPIN	chr8: 145154222, G > A	het	8	4	0	2916/33177	59/4865
CHD7	chr8: 61654298, T > A	het	5	0	0	1103/33106	39/4840
CX3CR1	chr3: 39323163, A > C	hom	11	4	0	4723/31219	193/4376
LRBA	chr4: 151199080, G > A	hom	3	3	0	2260/33328	20/5195
HIVEP3	chr1:420 47208, C > G	het	5	3	1	3383/32494	123/5061
IFIH1	chr2: 163124051, C > T	hom	20	3	1	12107/33356	184/5199
RNASEL	chr1: 182554557, C > T	hom	7	2	0	4543/33356	78/5202
Gene	ExAC LAT	PML EUR OR	PML EUR FET	PML AFR OR	PML AFR FET	PML ALL OR	PML ALL FET
PLCG2	116/5548	2.95	0.154755	16.40	0.0000	6.49	0.0002
IFIH1	119/5671	8.41	0.000156	11.22	0.0927	6.38	0.0002
TCIRG1	60/5770	NA	NA	5.85	0.0082	7.31	0.0028
IGLL1	236/5782	4.34	0.017218	1.27	0.7286	3.47	0.0036
MAVS	52/5691	3.92	0.023868	1.48	0.5191	3.47	0.0036
SHARPIN	171/5780	2.31	0.053526	19.17	0.0001	2.68	0.0040
CHD7	93/5725	3.72	0.015268	NA	NA	2.64	0.0485
CX3CR1	1357/5491	1.87	0.088087	5.10	0.0128	1.51	0.1806
LRBA	54/5785	1.01	1.000000	43.13	0.0001	1.69	0.2736
HIVEP3	902/5756	1.10	0.803620	6.69	0.0143	1.30	0.4283
IFIH1	1076/5776	1.46	0.212471	4.54	0.0374	1.21	0.4372
RNASEL	167/5785	1.20	0.658473	6.91	0.0403	1.22	0.5616

[0628] Table 14 lists the top tier of variants that were found to be significant on the basis of variant burden analysis, as described below. For each variant (genome coordinates are based on UCSC hg19), detailed information is presented of the numbers of EUR and AFR cases that carry the variant, along with the ethnic-specific and aggregate statistical metrics.

TABLE 15

Second tier of variants found on the basis of variant burden analysis							
Gene	Variant (hg19)	Genotype	PML EUR 44	PML AFR 21	PML LAT 5	ExAC EUR	ExAC AFR
SHARPIN	chr8: 145154824, A > C	het	3	0	0	2/30,670	0/4,471
RTEL1	chr20: 62305450, C > T	het	0	2	0	1/32,552	0/4,838
IGLL1	chr22: 23915745, G > A	het	2	0	1	19/33,348	74/5,184
PGM3	chr6: 83884161, C > G	het	0	2	0	0/33,069	26/5,167
ATM	chr11: 108202772, G > T	het	3	0	0	170/32,707	3/5,099
TMEM173	chr5: 138856923, C > T	het	2	2	0	108/32,327	204/4,842

TABLE 15-continued

Second tier of variants found on the basis of variant burden analysis							
CLCN7	chr16: 1510535, C > T	het	0	2	0	1/32,898	66/5,119
MAVS	chr20: 3843027, C > A	hom	4	2	0	803/33,206	167/5,171
ORAI1	chr12: 122064788, G > GT	het	4	0	0	371/28,708	5/3,555
RBFOX1	chr16: 7714909, C > T	het	0	2	0	1/33,367	69/4,902
MALT1	chr18: 56401523, C > T	het	4	0	0	466/33,239	9/5,179
GFI1	chr1: 92946625, G > C	het	2	1	0	206/29,111	6/4,156
DOCK2	chr5: 169081453, G > C	het	0	2	0	48/33,350	43/5,201
ATM	chr11: 108117787, C > T	het	2	0	0	93/33,256	2/5,151
SNAP29	chr22: 21235389, A > G	het	3	0	0	283/32,917	21/5,149
TICAM1	chr19: 4817657, C > T	het	0	2	0	32/31,437	71/4,814
GTPBP4	chr10: 1060218, G > A	hom	3	0	0	334/33,367	20/5,202
BACH1	chr21: 30698953, T > G	het	2	0	0	134/33,122	4/5,103
DOCK8	chr9: 304628, G > A	het	2	0	0	149/33,298	2/5,161
STXBP2	chr19: 7712287, G > C	het	2	0	0	161/32,104	4/4,626
FAS	chr10: 90771767, G > A	het	2	0	0	175/33,304	3/5,182
GOLGB1	chr3: 121415370, T > C	het	3	2	0	1,111/33,349	26/5,179
FUK	chr16: 70503095, A > G	het	4	0	0	741/33,341	23/4,899
IL10	chr1: 206945738, C > T	het	2	0	0	206/33,343	2/5,198
ITK	chr5: 156593120, C > T	het	2	0	0	206/33,353	5/5,203
STIM2	chr4: 27019452, C > T	het	2	0	0	219/33,369	5/5,202
ASH1L	chr1: 155317682, C > T	het	2	0	0	218/33,367	2/5,203
TBC1D16	chr17: 77926526, C > T	het	3	0	0	496/31,905	15/4,845
LYST	chr1: 235840495, G > T	het	3	0	0	517/33,239	10/5,156
SALL2	chr14: 21993359, G > A	het	3	0	0	519/31,729	14/4,520
CHD7	chr8: 61757805, C > T	het	3	0	0	517/32,880	14/4,872

TABLE 15-continued

Second tier of variants found on the basis of variant burden analysis							
BLM	chr15: 91306241, G > A	het	2	0	0	266/33,277	10/5,061
NOD2	chr16: 50741791, C > T	het	0	2	0	285/33,369	21/5,203
IGLL1	chr22: 23915583, T > C	het	2	0	0	265/33,334	21/5,183
TTC7A	chr2: 47205921, C > T	het	3	0	0	589/33,202	13/5,173
KITLG	chr12: 88900891, C > A	het	4	0	0	1,023/33,226	31/5,158
ATR	chr3: 142281353, C > G	het	4	0	0	1,037/33,343	14/5,130
ATM	chr11: 108123551, C > T	het	0	2	0	217/29,921	66/4,955
CR2	chr1: 207641950, C > T	het	0	2	0	391/33,363	19/5,203
HIVEP2	chr6: 143092151, T > C	het	3	2	0	1,718/33,370	50/4,901
ITSN2	chr2: 24431184, C > T	hom	3	2	0	2,019/33,339	17/5,186
ITSN2	chr2: 24432937, C > T	hom	3	2	0	2,026/32,472	17/4,958
DOCK8	chr9: 312134, G > A	het	3	2	0	2,114/33,251	79/5,180
VPS13B	chr8: 100205255, G > A	het	0	2	0	811/33,345	19/5,192
NRIP1	chr21: 16339852, T > C	het	0	2	0	901/33,355	19/5,203
Gene	ExAC LAT	PML EUR OR	PML EUR FET	PML AFR OR	PML AFR FET	PML ALL OR	PML ALL FET
SHARPIN	0/5,302	1122.00	0.000000	NA	NA	905.40	0.0000
RTEL1	0/5,737	NA	NA	1240.64	0.0000	1268.41	0.0000
IGLL1	9/5,783	83.53	0.000351	NA	NA	19.41	0.0006
PGM3	3/5,748	NA	NA	20.81	0.0055	44.58	0.0011
ATM	7/5,713	14.00	0.001636	NA	NA	10.78	0.0032
TMEM173	58/5,770	14.21	0.009863	2.39	0.2226	6.97	0.0033
CLCN7	0/5,732	NA	NA	8.06	0.0308	19.18	0.0055
MAVS	46/5,779	4.04	0.021706	3.15	0.1480	3.98	0.0056
ORAI1	16/5,354	7.64	0.002562	NA	NA	5.76	0.0064
RBFOX1	4/5,782	NA	NA	7.37	0.0361	17.48	0.0066
MALT1	40/5,760	7.03	0.003411	NA	NA	5.14	0.0093
GFI1	39/5,114	6.68	0.039391	34.58	0.0347	6.80	0.0113
DOCK2	27/5,786	NA	NA	12.63	0.0137	11.02	0.0155
ATM	28/5,756	16.98	0.007047	NA	NA	10.53	0.0169
SNAP29	32/5,740	8.44	0.006584	NA	NA	5.79	0.0171
TICAM1	19/5,687	NA	NA	7.03	0.0392	10.08	0.0183
GTPBP4	21/5,786	7.24	0.009925	NA	NA	5.25	0.0220
BACH1	6/5,778	11.72	0.014110	NA	NA	8.96	0.0227
DOCK8	5/5,762	10.59	0.017020	NA	NA	8.31	0.0261
STXBP2	11/5,686	9.45	0.021028	NA	NA	7.06	0.0350
FAS	10/5,731	9.01	0.022902	NA	NA	6.89	0.0365
GOLGB1	84/5,779	2.12	0.180743	20.86	0.0055	2.71	0.0443
FUK	73/5,787	4.40	0.016488	NA	NA	3.13	0.0449
IL10	6/5,787	7.66	0.030787	NA	NA	6.06	0.0458
ITK	5/5,789	7.66	0.030770	NA	NA	6.01	0.0466
STIM2	9/5,789	7.21	0.034346	NA	NA	5.57	0.0532
ASH1L	17/5,789	7.24	0.034067	NA	NA	5.48	0.0548
TBC1D16	21/5,707	4.63	0.031269	NA	NA	3.53	0.0584

TABLE 15-continued

Second tier of variants found on the basis of variant burden analysis							
LYST	35/5,756	4.63	0.031299	NA	NA	3.47	0.0606
SALL2	17/5,718	4.40	0.035528	NA	NA	3.37	0.0650
CHD7	41/5,765	4.58	0.032169	NA	NA	3.36	0.0654
BLM	17/5,756	5.91	0.048875	NA	NA	4.40	0.0799
NOD2	2/5,789	NA	NA	25.97	0.0037	4.21	0.0860
IGLL1	26/5,787	5.94	0.048403	NA	NA	4.15	0.0881
TTC7A	61/5,759	4.05	0.043427	NA	NA	2.94	0.0891
KITLG	40/5,760	3.15	0.046242	NA	NA	2.38	0.0964
ATR	69/5,785	3.12	0.047671	NA	NA	2.33	0.1021
ATM	40/5,425	NA	NA	7.80	0.0327	3.64	0.1093
CR2	8/5,754	NA	NA	28.72	0.0031	3.09	0.1422
HIVEP2	209/5,788	1.35	0.494339	10.21	0.0202	1.64	0.2458
ITSN2	55/5,784	1.14	0.748301	32.01	0.0025	1.55	0.3862
ITSN2	56/5,672	1.10	0.753875	30.59	0.0028	1.50	0.3937
DOCK8	161/5,768	1.08	0.757661	6.80	0.0415	1.37	0.4238
VPS13B	100/5,778	NA	NA	28.66	0.0031	1.37	0.6600
NRIP1	64/5,780	NA	NA	28.72	0.0031	1.30	0.6698

[0629] Table 15 lists the second tier of variants that were found on the basis of variant burden analysis, as described below. For each variant (genome coordinates are UCSC hg19), detailed information is presented of the numbers of EUR and AFR cases that carry the variant, along with the ethnic-specific and aggregate statistical metrics.

TABLE 16

Potential testing scenario, based on top variant burden hits						
Gene/Variant	Cases		Proportion of Cohort (n = 70) Test Method	Patient information		
	solved			Primary disease	Ethnicity	Gender
All 4 SNVs	28	40%	genotyping	M, H, O	A, E	both
SHARPIN, IFIH1, PLCG2 SNVs	24	34%	genotyping	M, H, O	A, E	both
IFIH1, PLCG2 SNVs	13	19%	genotyping	M, H, O	A, E	both
SHARPIN SNV	13	19%	genotyping	M, H	A, E	both
IFIH1 SNV	7	10%	genotyping	M, H, O	A, E	both
PLCG2 SNV	7	10%	genotyping	M, H	A, E	both
CHD7 SNV	5	7%	genotyping	M, H, O	E	both

[0630] Table 16 lists a potential testing scenario, based on top variant burden hits (reported in Table 14). The analysis is for illustrative purposes only, it being acknowledged that greater diagnostic yields can be obtained by assaying for a larger number of variants, including those listed in Table 15. Examples are given for diagnostic yield using singleton

variants, as well as a variety of combinations, including the use of the top 4 variants. For this set of variants, the test method is described as genotyping, as opposed to whole gene sequencing (i.e., determination of the status at each of the bases, which yields a binary output, as opposed to identification of variants elsewhere in the relevant genes).

TABLE 17

Potential testing scenario using genes identified as having a greater burden of damaging, heterozygous variants in the PML cohort					
GENE	Ave		Ethnic-specific yield (%)	Overall yield (EUR + AFR) (%)	Test Method
	CASES	Ethnicity			
PLCG2	17/44	EUR	38	38	Gene sequencing
PLCG2	8/21	AFR	38		Gene sequencing
POLE	8/21	AFR	38		Gene sequencing
POLE	11/44	EUR	25	29	Gene sequencing
LRBA	11/44	EUR	25		Gene sequencing
EPG5	9/44	EUR	20		Gene sequencing
SHARPIN	8/44	EUR	18		Gene sequencing

[0631] Table 17 lists a potential testing scenario using genes identified as having a greater burden of damaging, heterozygous variants in the PML cohort (see Table 13). The nature of the testing method is ‘gene sequencing’ since the variants are not known in advance—any and all potentially damaging variants need to be considered in such an assay.

TABLE 18

Summary of genes that survive case-level, gene burden and/or variant burden analyses			
Gene	Case Level	Variant Burden	Gene Burden
PLCG2	Yes	Yes	Yes
CHD7	Yes	Yes	
IFIH1	Yes	Yes	
AP3B1	Yes		Yes
EPG5	Yes		Yes
PIK3CD	Yes		Yes
LRBA		Yes	Yes
SHARPIN		Yes	Yes

[0632] Table 18 represents a summary of genes that survive case-level (2 or more examples in Tables 7, 8), gene burden and/or variant burden analyses (based on Tables 13 and 14). Of note is that PLCG2 satisfies all 3 criteria (2 or more examples, in Table 8, presence in Tables 13, 14). This summary demonstrates that many genes have been identified as significant on the basis of independent analysis methods.

Example 11—Figures Referenced in this Study

[0633] FIGS. 1-12 represent example CNV data from the PML gene discovery study (71 PML cases, see Table 7 for patient information) using array CGH (methods described herein). In each figure/drawing: 1) genome coordinates are listed at the top (hg18 assembly, chromosome number and position depicted); 2) data track 1 (labeled ‘Genes’) depicts the location of the RefSeq genes (exons are dark gray portions of the bars, introns are light gray portions of the bars); 3) data track 2 (labeled ‘Normal Cohort’) depicts the size and location of CNVs found in the NVE cohort (PBio’s proprietary control database consisting of CNV findings in apparently healthy—i.e. normal—subjects, see methods herein) with the y-axis corresponding to the number of NVE subjects that have the CNV; and 4) remaining data tracks are CNV data found in individual PML patients wherein the y-axis corresponds to the log 2 ratio (see methods herein), points represent individual probes on the microarray, and line segments are shifted positive (copy number gain) or negative (copy number loss) based on the output of DNA-copy, the CNV calling algorithm. Typical log 2 ratios for gains and losses on the Agilent 1M microarray (see methods herein) and our experimental protocols are: 0.6 for duplications, 1.0 for triplications (or homozygous duplications), -1.0 for heterozygous deletions, and <-2 (often -4.0 to -6.0) for homozygous deletions. Relevant genes are labeled in the ‘Genes’ data track.

[0634] FIG. 1 represents an example of a gene impacted by germline and acquired CNVs. Germline CNVs that impact the PRKCB gene include patient PML50 with a 4.8 Kb intronic heterozygous loss (also found in 7 Normal subjects) and patient PML11 with a 7.3 Kb intronic gain (also found in 1 Normal subject). Acquired CNVs were found in 6 PML patients, a series of gains at -23.9 Mb with

varying log 2 ratios, suggestive of a mixed cell population (array CGH experiments were performed on blood-derived genomic DNA).

[0635] FIG. 2 represents an example of potentially PML-relevant genes (TNFRSF13C and CENPM) impacted by acquired CNVs. Acquired CNVs were found in 9 PML patients, a series of gains at -40.6 Mb with varying log 2 ratios, suggestive of a mixed cell population (array CGH experiments were performed on blood-derived genomic DNA). All 9 PML patients (see Table 7 for patient information) had a primary diagnosis of HIV and were mixed gender (3 females and 6 males) and ethnicity (4 African ancestry and 5 European ancestry).

[0636] FIG. 3 represents an example of a gene impacted by germline and acquired CNVs. A germline CNV, which is a 7.2 Kb intronic heterozygous loss (not found in Normal subjects, but an adjacent loss is found in 8 Normal subjects) that impacts the PKHD1 gene, was detected in patient PML26. Acquired CNVs were found in 3 PML patients, a series of gains at -51.9 Mb with varying log 2 ratios, suggestive of a mixed cell population (array CGH experiments were performed on blood-derived genomic DNA).

[0637] FIG. 4 represents an example of a gene impacted by a recurrent CNV loss. The 14.7 Kb intronic deletion impacts the BMPR2 gene. Heterozygous deletions were detected in patients PML58 and MVGS811-13a (also found in 2 Normal subjects), and a homozygous deletion was detected in patient PML29 (none found in Normal subjects). All three PML patients are males and their primary disease is HIV (see Table 7).

[0638] FIG. 5 represents an example of a gene impacted by a recurrent CNV gain. The 10.2 Kb exonic gain disrupts the COMMD6 gene. Two PML patients, PML29 and MVGS811-13a, have a homozygous duplication (log 2 ratio comparable to triplications) based on the observation that 1000 genomes subjects are reported to have this gain (see hg19 assembly DGV variant esv3632749, which reports 148 of 2504 subjects as having this gain; no Normals were found in PBio’s NVE db). Both PML patients are males and their primary disease is HIV (see Table 7).

[0639] FIG. 6 represents an example of a gene impacted by a recurrent CNV gain. The 27.4 Kb exonic gain disrupts the KCTD7 gene and the right breakpoint is 16-90 Kb upstream of RABGEF 1 transcript variants (RefSeq: NM_001287060, NR_104676, NM_014504, NM_001287062, NM_001287061). Patient PML29 has a homozygous duplication (log 2 ratio comparable to triplications) based on the observation that 1000 genomes subjects are reported to have this gain (see hg19 assembly DGV variant esv3613515, which reports 28 of 2504 subjects as having this gain; no Normals were found in PBio’s NVE db). Patient PML63 has a duplication. Both PML patients are males of African ancestry and their primary disease is HIV (see Table 7).

[0640] FIG. 7 represents an example of a gene impacted by a recurrent CNV gain. The 344 Kb exonic gain disrupts the FPR2 and ZNF616 genes (via left and right breakpoints) and additional genes fully encompassed by this CNV are: FPR3, ZNF350, ZNF350-AS1, ZNF432, ZNF577, ZNF613, ZNF614, ZNF615, ZNF649, ZNF649-AS1, ZNF841. Patient PML03 has a homozygous duplication (log 2 ratio comparable to triplications) based on the observation that 3 Normal subjects (PBio’s NVE db) are found to have a duplication of this region, along with patient PML10. Both

PML patients are females of European ancestry and their primary diseases are HIV and MS (see Table 7).

[0641] FIG. 8 represents an example of a gene impacted by a recurrent CNV loss. The 1.1 Kb exonic deletion impacts the PIK3CD and PIK3CD-AS1 (previous gene symbol was Clorf200) genes. A homozygous deletion was detected in patient MVGS811-13a and this loss (heterozygous or homozygous) was not found in Normal subjects or the DGV public CNV database. The PML patient is a male and his primary disease is HIV (see Table 7). He is presumed to be of EUR ancestry (ethnicities were not available for MVGS samples).

[0642] FIG. 9 represents an example of a gene impacted by an intergenic, recurrent CNV gain. The 16.7 Kb intergenic gain has a left breakpoint that is 105 Kb upstream of the CD180 gene (RefSeq transcript variant NM_005582). Patient MVGS995-4a has a homozygous duplication (log 2 ratio comparable to triplications) based on the observation that 1000 genomes subjects are reported to have this gain (see hg19 assembly DGV variant esv3605336, which reports 2 of 2504 subjects as having this gain; no Normals were found in PBio's NVE db). The PML patient is a male of European ancestry and his primary disease is MS (see Table 7).

[0643] FIG. 10 represents an example of a gene impacted by an intergenic, recurrent CNV loss. The 7.7 Kb intergenic homozygous deletion has a left breakpoint that is 3-4 Kb upstream of VDAC1 transcript variants (RefSeq: NM_003374, NR_036625, NR_036624). This loss (heterozygous or homozygous) was not found in Normal subjects or the DGV public CNV database. Patient PML30 is a male of European ancestry and his primary disease is HIV (see Table 7).

[0644] FIG. 11 represents an example of a gene impacted by an intergenic, recurrent CNV loss. The 6.8 Kb intergenic homozygous deletion has a left breakpoint that is 4 Kb downstream of EGR1 transcript variant (RefSeq: NM_001964) and 26 Kb downstream of ETF1 transcript variants (RefSeq: NM_001256302, NM_004730, NM_001282185, NM_001291975, NM_001291974). This loss was found to be homozygous in 1 Normal subject and the loss was also reported in the DGV public CNV database (see hg19 assembly DGV variant esv3606925, which reports 33 of 2504 subjects as having this loss, homozygous vs. heterozygous subjects are unknown). Patient PML69 is a male of European ancestry and his primary disease (condition) is kidney transplant (see Table 7, reported as 'Other'). Patient PML69 was treated with CTLA4-Ig (belatacept, a CD28-B7 costimulation blocker and T-cell anergy inducer). The CD28 pathway includes links to the patient's genetic finding (e.g., homozygous deletion adjacent to the EGR1 gene) and several other genes that may be related to immunodeficiency (e.g., CD40LG, ITK, LCK, LRBA, PIK3CD, PIK3R1, PLCG2, WAS, and ZAP70) (Dekeyser M et al. Open Forum Infect Diseases, 2016, Refractory T-Cell Anergy and Rapidly Fatal Progressive Multifocal Leukoencephalopathy following Prolonged CTLA4 Therapy).

[0645] FIG. 12 represents an example of a gene impacted by an intergenic, recurrent CNV loss. The 5.6 Kb intergenic homozygous deletion has a left breakpoint that is 20 Kb upstream of IITSN2 transcript variants (RefSeq: NM_019595, NM_006277, NM_147152). Heterozygous losses were found in 50 Normal subjects and the loss was also reported in the DGV public CNV database (see hg19

assembly DGV variant esv3590068, which reports 222 of 2504 subjects as having this loss, homozygous vs. heterozygous subjects are unknown). Patient PML65 is a male of African ancestry and his primary disease is HIV (see Table 7).

[0646] FIG. 13 represents an example of known and/or predicted protein interactions using the String database (string-db.org; see Szklarczyk et al., (2015) and references therein). A non-redundant list of all genes reported in Table 7 (43 genes, which included those whose expression was inferred to be impacted by a nearby intergenic CNV) as best solutions/explanations for 61 of 71 PML cases (11 PML cases are reported as 'unsolved', including 1 case for which only CGH data was obtained) was assessed using the String db. The 'minimum required interaction score' was set to 'high confidence (0.7)' and no additional 'interactors' were added. Of the 43 input genes, 21 were found to have high confidence interactions, as shown in the figure, along with annotation of the number of PML cases that had each of these genes as a solution/explanation (e.g., 3 PML cases in Table 7 were found to have a PLCG2 solution).

Example 12—Gene Burden Analysis

[0647] Gene burden analysis was performed as follows. Using a variety of in-house scripts, and data downloaded from ExAC (exac.broadinstitute.org), a count was performed for all variants occurring in each of the 419 genes listed in Table 6. Each variant was classified according to whether it was deemed damaging (on the basis of at least one of the prediction algorithms SIFT, PolyPhen2 or MutationTaster) or non-damaging, heterozygous or homozygous. This was performed in parallel for PML variants and those found in ExAC. ExAC data for which quality/coverage was <80% of expected was not used and gene burden analysis could not therefore be performed.

[0648] An ethnic-specific (EUR or AFR only, there were too few LAT cases for this type of analysis) comparison was then performed for each of 4 categories:

[0649] Homozygous damaging

[0650] Homozygous non-damaging

[0651] Heterozygous damaging

[0652] Heterozygous non-damaging

[0653] For all 4 categories, variants with minor allele frequency (MAF) cutoffs of 0.01, 0.02, 0.03, 0.04, 0.05 and 0.1 were considered.

[0654] For each comparison, odds ratios (OR) and Fisher's exact test (FET) were calculated for the comparison of numbers of PML cases with at least one variant of the type under consideration and those in ExAC. Correction for multiple testing was performed by multiplying the FET by the number of genes being considered (419). Only genes for whom the FET_corrected was <0.05 were included in Table 13, which contains data on the average values for a given gene at all MAFs that passed FET_correction. In practice, only the category of heterozygous damaging yielded significant genes.

Example 13—Variant Burden Analysis

[0655] For each variant identified in at least one PML case, a count was performed in order to obtain the frequency of the same variant in the cohort as a whole. This aggregate data was compared to counts for the same variant as reported

in ExAC. ExAC data was filtered for quality/coverage and variant burden analysis was not performed if ExAC coverage was <80% expected.

[0656] Variant burden analysis was performed separately for EUR (n=44 cases) and AFR (n=21 cases) cohorts (LAT cohort was too small) and the OR and FET values calculated. From this analysis, only variants with OR>1 (i.e., potentially indicative of increased risk for PML) for both ethnicities (AFR and EUR) and for which the ExAC frequency of the variant was <5% were considered. Furthermore, only those variants for which the frequency in the ethnic-specific cohort was >10% (5 or more EUR cases, 3 or more AFR cases) were considered top-tier (Table 14), although other variants have been tabulated in Table 15.

Example 14—Exemplary PML Risk Prediction Tests

[0657] Table 16 provides exemplary markers for creating a low-cost, simple (genotype specific SNVs) PML risk prediction test. Other embodiments could be similarly devised from other SNVs reported in Tables 14 and 15. Different combinations of SNVs from Tables 14, 15 could be utilized in tests of varying complexity, to develop a test that would yield higher diagnostic yields than the top example listed in Table 16 (i.e., 40%).

[0658] Table 17 provides exemplary genes that could be included in a gene panel sequencing test for PML risk prediction. Other embodiments could be similarly devised from genes reported in Table 13, or from other tables disclosed herein.

[0659] Table 9 contains ‘example’ variants that may be considered as ‘AD’ causes of immunodeficiency (i.e., presence of just 1 of the 2 reported het SNVs in a given patient may be causing immunodeficiency), which may increase the risk for PML. For example, this may be a more likely scenario for het SNVs that are ‘novel’ in the ExAC db (i.e., not found in the general population), and even more likely if such novel SNVs are found in ≥2 PML cases (irrespective of the invoked disease model). Examples of this include the following 3 genes:

[0660] AK2, 2 cases (Table 9)

[0661] chr1:33476435, C>A, novel in ExAC

[0662] PML20 and PML33, AFR and EUR, both HIV

[0663] EPG5, 2 cases (Table 9)

[0664] chr18: 43445601, T>G, novel in ExAC

[0665] PML25 and PML27, both EUR, both HIV

[0666] TNFRSF11A, 9 cases (Table 7)

[0667] chr18: 60052034, A>C, novel in ExAC

[0668] see Table 7 for case IDs, 2 AFR and 7 EUR, all HIV

[0669] It can be appreciated by those skilled in the art that immunodeficiency genes presently known to cause AR disease may potentially cause AD disease. Numerous examples have been reported in the literature, including several of the genes listed in Table 6 (e.g., Disease_model is indicated as AD_AR for 32 genes, such as ADAR and TICAM1).

Example 15—Exemplary 96-Genes Panel PML Risk Prediction Tests

[0670] Table 19 contains an exemplary 96-gene panel based on genes that were found to have at least one PML

case count from Tables 7 and 8. The “Genes” and “Case_level_solutions” columns showed genes and total number of PML cases (with at least one ‘case level’ solution) reported in Tables 7 and 8. In addition, the top 7 genes (CHD7, IFIH1, IGLL1, MAVS, PLCG2, SHARPIN, TCIRG1) from Table 14 with SNVs based on ‘PML_ALL_FET’ values <0.05 (column 0) were also included in Table 19. Among these 7 genes, 3 genes (IGLL1, MAVS, SHARPIN) with SNVs were based on ‘PML_ALL_FET’ values <0.05 (column 0) from Table 15.

TABLE 19

exemplary 96-gene panel	
Genes	Case_level_solutions
AP3B1	5
APOL1	1
ASH1L	1
ATM	1
ATR	3
BLM	1
CARD11	3
CDKN1B	1
CHD7	4
CLCN7	1
DCLRE1C	3
DDX58	1
DOCK8	8
EGR1	1
EPG5	3
ETF1	1
FPR2	1
GATA2	2
GFI1	4
HIVEP1	1
HIVEP2	2
HTR2A	1
IDO2	1
IFIH1	3
IFNGR2	1
IFNLR1	1
IGLL1	0
IKBKB	1
IL17F	1
IL1B	1
IL21R	1
IRAK4	2
ITSN2	2
JUN	2
KAT6B	1
KCTD7	1
LIG4	1
LRBA	1
MALL	1
MAPK3	2
MAVS	0
MCEE	1
MKL1	1
MYD88	1
NBN	1
NFKB1	3
NOD2	6
NRIP1	1
PIAS1	1
PIAS2	1
PIK3CD	4
PIK3CD-AS1	1
PIK3R1	1
PKHD1	3
PLCG2	5
PNPT1	1
POLA1	1
POLE	1
PRF1	1

TABLE 19-continued

exemplary 96-gene panel	
Genes	Case_level_solutions
PRKCB	1
PRKCD	1
PRKCH	1
PRKDC	4
PSTPIP1	1
PTEN	1
PTPRC	2
RABGEF1	1
RAD51	1
RAG1	4
RAG2	2
RIPK1	1
RIPK3	2
RNF168	2
RTEL1	2
SHARPIN	1
SKIV2L	1
SMAD4	1
STIM1	2
STIM2	1
STXBP2	3
TAP2	1
TBK1	2
TCIRG1	1
TICAM1	2
TLR3	2
TLR4	1
TNFRSF11A	10
TNFRSF13B	1
TNFRSF8	1
TP53	1
TRAF3	1
TRAFD1	1
TRPM2	1
VPS45	1
WEE1	2
ZAP70	3
TOTAL (96 genes)	172
Non-redundant cases	67
Dx yield for PML cohort (n = 70)	95.7%

[0671] The non-redundant number of PML cases and diagnostic yield are listed in the last 2 rows of Table 19. Specifically, a test including the 96 genes had a diagnostic yield of 95.7% based on the genetic findings from the 70 PML cases used in the present study.

Example 16—Exemplary 39-Gene Panel PML Risk Prediction Tests

[0672] Table 20 contains an exemplary 39-gene panel based on genes that were found to have multiple PML case count from Tables 7 and 8. The “Genes” and “Case_level_solutions” columns showed genes and total number of PML cases (with at least two ‘case level’ solutions) reported in Tables 7 and 8. In addition, the top 7 genes (CHD7, IFIH1, IGLL1, MAVS, PLCG2, SHARPIN, TCIRG1) from Table 14 with SNVs based on ‘PML_ALL_FET’ values <0.05 (column 0) were also included in Table 20. Among these 7 genes, 3 genes (IGLL1, MAVS, SHARPIN) with SNVs were based on ‘PML_ALL_FET’ values <0.05 (column 0) from Table 15.

TABLE 20

exemplary 39-gene panel	
Genes	Case_level_solutions
AP3B1	5
ATR	3
CARD11	3
CHD7	4
DCLRE1C	3
DOCK8	8
EPG5	3
GATA2	2
GFI1	4
HIVEP2	2
IFIH1	3
IGLL1	0
IRAK4	2
ITSN2	2
JUN	2
MAPK3	2
MAVS	0
NFKB1	3
NOD2	6
PIK3CD	4
PKHD1	3
PLCG2	5
PRKDC	4
PTPRC	2
RAG1	4
RAG2	2
RIPK3	2
RNF168	2
RTEL1	2
SHARPIN	1
STIM1	2
STXBP2	3
TBK1	2
TCIRG1	1
TICAM1	2
TLR3	2
TNFRSF11A	10
WEE1	2
ZAP70	3
TOTAL (39 genes)	115
Non-redundant cases	57
Dx yield for PML cohort (n = 70)	81.4%

[0673] The non-redundant number of PML cases and diagnostic yield are listed in the last 2 rows of Table 20. Specifically, a test including the 39 genes had a diagnostic yield of 81.4% based on the genetic findings from the 70 PML cases used in the present study.

Example 17—Exemplary 23-Gene Panel PML Risk Prediction Tests

[0674] Table 21 contains an exemplary 23-gene panel based on genes that were found to have multiple PML case count from Tables 7 and 8. The “Genes” and “Case_level_solutions” columns showed genes and total number of PML cases (with at least three ‘case level’ solutions) reported in Tables 7 and 8. In addition, the top 7 genes (CHD7, IFIH1, IGLL1, MAVS, PLCG2, SHARPIN, TCIRG1) from Table 14 with SNVs based on ‘PML_ALL_FET’ values <0.05 (column 0) were also included in Table 21. Among these 7 genes, 3 genes (IGLL1, MAVS, SHARPIN) with SNVs were based on ‘PML_ALL_FET’ values <0.05 (column 0) from Table 15.

TABLE 21

exemplary 23-gene panel	
Genes	Case_level_solutions
AP3B1	5
ATR	3
CARD11	3
CHD7	4
DCLRE1C	3
DOCK8	8
EPG5	3
GFI1	4
IFIH1	3
IGLL1	0
MAVS	0
NFKB1	3
NOD2	6
PIK3CD	4
PKHD1	3
PLCG2	5
PRKDC	4
RAG1	4
SHARPIN	1
STXBP2	3
TCIRG1	1
TNFRSF11A	10
ZAP70	3
TOTAL (23 genes)	83
Non-redundant cases	50
Dx yield for PML cohort (n = 70)	71.4%

[0675] The non-redundant number of PML cases and diagnostic yield are listed in the last 2 rows of Table 21. Specifically, a test including the 23 genes had a diagnostic yield of 71.4% based on the genetic findings from the 70 PML cases used in the present study.

Example 18—Exemplary 15-Gene Panel PML Risk Prediction Tests

[0676] Table 22 contains an exemplary 15-gene panel based on genes that were found to have multiple PML case count from Tables 7 and 8. The “Genes” and “Case_level_solutions” columns showed genes and total number of PML cases (with at least four ‘case level’ solutions) reported in Tables 7 and 8. In addition, the top 7 genes (CHD7, IFIH1, IGLL1, MAVS, PLCG2, SHARPIN, TCIRG1) from Table 14 with SNVs based on ‘PML_ALL_FET’ values <0.05 (column 0) were also included in Table 22. Among these 7 genes, 3 genes (IGLL1, MAVS, SHARPIN) with SNVs were based on ‘PML_ALL_FET’ values <0.05 (column 0) from Table 15.

TABLE 22

exemplary 15-gene panel	
Genes	Case_level_solutions
AP3B1	5
CHD7	4
DOCK8	8
GFI1	4
IFIH1	3
IGLL1	0
MAVS	0
NOD2	6
PIK3CD	4
PLCG2	5

TABLE 22-continued

exemplary 15-gene panel	
Genes	Case_level_solutions
PRKDC	4
RAG1	4
SHARPIN	1
TCIRG1	1
TNFRSF11A	10
TOTAL (15 genes)	59
Non-redundant cases	39
Dx yield for PML cohort (n = 70)	55.7%

[0677] The non-redundant number of PML cases and diagnostic yield are listed in the last 2 rows of Table 22. Specifically, a test including the 15 genes had a diagnostic yield of 55.7% based on the genetic findings from the 70 PML cases used in the present study.

Example 19—Exemplary 11-Gene Panel PML Risk Prediction Tests

[0678] Table 23 contains an exemplary 11-gene panel based on genes that were found to have multiple PML case count from Tables 7 and 8. The “Genes” and “Case_level_solutions” columns showed genes and total number of PML cases (with at least five ‘case level’ solutions) reported in Tables 7 and 8. In addition, the top 7 genes (CHD7, IFIH1, IGLL1, MAVS, PLCG2, SHARPIN, TCIRG1) from Table 14 with SNVs based on ‘PML_ALL_FET’ values <0.05 (column 0) were also included in Table 23. Among these 7 genes, 3 genes (IGLL1, MAVS, SHARPIN) with SNVs were based on ‘PML_ALL_FET’ values <0.05 (column 0) from Table 15.

TABLE 23

exemplary 11-gene panel	
Genes	Case_level_solutions
AP3B1	5
CHD7	4
DOCK8	8
IFIH1	3
IGLL1	0
MAVS	0
NOD2	6
PLCG2	5
SHARPIN	1
TCIRG1	1
TNFRSF11A	10
TOTAL (11 genes)	43
Non-redundant cases	33
Dx yield for PML cohort (n = 70)	47.1%

[0679] The non-redundant number of PML cases and diagnostic yield are listed in the last 2 rows of Table 23. Specifically, a test including the 11 genes had a diagnostic yield of 47.1% based on the genetic findings from the 70 PML cases used in the present study.

Example 20—Exemplary 10-Gene Panel PML Risk Prediction Tests

[0680] Table 24 contains an exemplary 10-SNV panel based on top 7 SNVs in Table 14 and 3 SNVs from Table 15 (based on overlapping genes between 14 and 15: IGLL1,

MAVS, SHARPIN). Specifically, Using the top 10 SNVs (7 from Table 14, along with 3 from Table 15, residing in genes already selected from Table 14), an additive count (column “Case total additive (non-redundant)”) was performed to determine how many PML cases had at least one of the variants when these were considered in order (e.g., column “Order (FET)”: ‘1’, first, followed by ‘1’+‘2’, followed by ‘1’+‘2’+‘3’, etc). Since some individuals harbor more than one variant, the additive count is not equal to the simple sum of PML case numbers for each variant (column “Case total per SNV”). All genome coordinates are based on hg19 build. **[0681]** An additive count was performed for ExAC subjects (column “ExAC subjects total additive (redundant)”), as follows: i) The average cohort size for ExAC for all variants was calculated; ii) Each total subject count (all ethnicities) was normalized to this average cohort size. The ExAC additive count represents a simple addition: labeled as “redundant” in column “ExAC subjects total additive (redundant)”, because information regarding the possible presence of multiple variants in the same individual is not available; iii) Odds Ratios (ORs) and Fisher’s Exact test (FET) values were calculated (columns “PML ALL OR additive” and “PML ALL FET additive”).

TABLE 24

exemplary 10-gene panel										
Order (FET) ¹	Table source	Gene	Variant (hg19)	Genotype	Case total per SNV	Case total additive (non-redundant) ²	Dx yield (non-redundant)	ExAC subjects total additive (redundant) ³	PML ALL OR additive	PML ALL FET additive
1	14	PLCG2	chr16: 81942175, A > G	het	7	7	10%	730	6.50	2.00E-04
2	14	IFIH1	chr2: 163136505, C > G	het	7	13	19%	1,473	6.49	6.37E-07
3	14	TCIRG1	chr11: 67818269, G > A	het	4	16	23%	1,830	6.73	2.94E-08
4	14	IGLL1	chr22: 23917192, G > T	het	8	22	31%	3,388	5.42	9.41E-09
5	14	MAVS	chr20: 3846397, C > T	hom	8	26	37%	4,947	4.60	2.13E-08
6	14	SHARPIN	chr8: 145154222, G > A	het	12	33	47%	8,064	3.91	5.10E-08
7	14	CHD7	chr8: 61654298, T > A	het	5	36	51%	9,292	3.89	3.26E-08
8	15	SHARPIN	chr8: 145154824, A > C	het	3	37	53%	9,294	4.12	8.10E-09
9	15	IGLL1	chr22: 23915745, G > A	het	3	38	54%	9,394	4.30	2.59E-09
10	15	MAVS	chr20: 3843027, C > A	hom	6	38	54%	10,393	3.77	5.26E-08

¹SNV order based on lowest FET value reported in Tables 14 and 15 for combined ethnicities

²PML case total = 70

³ExAC subject total = 43,419 (average for the 10 SNVs)

[0682] It can be appreciated by those skilled in the art that the above gene panels were selected based on the present genetic findings in 70 PML cases. Furthermore, a gene not presently selected for any of these exemplary gene panels may be added to the gene panel. For example, genes in which only 1 PML case was found to have variants fulfilling the criteria may be added to the gene panel if genetic validation in additional PML cases shows a ‘n=1 case’ gene is impacted by more than 1 PML case when the data are examined for a new set of PML cases. In some cases, additional genes (e.g., PML-linked genes such as DOCK8, BAG3, STAT1) may be added to the gene panel.

[0683] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

SEQUENCE LISTING

The patent application contains a lengthy sequence listing. A copy of the sequence listing is available in electronic form from the USPTO web site (<https://seqdata.uspto.gov/?pageRequest=docDetail&DocID=US20250277269A1>). An electronic copy of the sequence listing will also be available from the USPTO upon request and payment of the fee set forth in 37 CFR 1.19(b)(3).

What is claimed is:

1. A method of treating a condition in a subject in need of an immunosuppressive medication therapy, comprising: administering a therapeutically effective amount of an immunosuppressive medication to the subject, wherein the subject has a decreased risk of progressive multifocal leukoencephalopathy (PML) due to an infection of the brain by John Cunningham virus (JCV), wherein the subject's decreased risk is associated with an absence of one or more genetic variations in the subject, wherein the subject has been tested for a presence of the one or more genetic variations with a genetic assay and has been identified as not having the one or more genetic variations, wherein the one or more genetic variations has an odds ratio (OR) of 2 or more, and wherein the OR is:

$$[D_D/D_N]/[D/N_N],$$

wherein D_D is the number of subjects in a diseased cohort of subjects with the one or more genetic variations; D_N the number of subjects in the diseased cohort without the one or more genetic variations; N_D is the number of subjects in a non-diseased cohort of subjects with the one or more genetic variations; and N_N is the number of subjects in the non-diseased cohort without the one or more genetic variations, wherein the diseased cohort of subjects have PML, and wherein the non-diseased cohort of subjects do not have PML.

2. The method of claim 1, wherein the one or more genetic variations disrupt or modulate a gene selected from the group consisting of a TMEM173 gene, a TCIRG1 gene, a ORAI1 gene, a ITSN2 gene, a IFNGR2 gene, a IKBKB gene, a IL21R gene, a LIG4 gene, a PIK3R1 gene, a PNPT1 gene, a POLE gene, a PSTPIP1 gene, a TAP2 gene, a TNFRSF13B gene, and a TRPM2 gene.

3. The method of claim 2, further comprising testing the subject for the presence of the one or more genetic variations with the genetic assay prior to the administering.

4. The method of claim 2, wherein the genetic assay comprises microarray analysis, PCR, sequencing, nucleic acid hybridization, or any combination thereof.

5. The method of claim 2, wherein the genetic assay comprises whole exome sequencing or whole genome sequencing.

6. The method of claim 2, wherein the genetic assay comprises analyzing nucleic acid information of the subject obtained by whole exome sequencing or whole genome sequencing.

7. The method of claim 2, wherein the subject has been tested with a JCV antibody test, a CD62L test, or a CSF IgM oligoclonal bands test.

8. The method of claim 2, wherein the one or more genetic variations is a single nucleotide polymorphism (SNP) or a single nucleotide variation (SNV).

9. The method of claim 2, wherein the one or more genetic variations is a deletion.

10. The method of claim 2, wherein the one or more genetic variations comprise (i) a genetic variation in the TMEM173 gene that is chr5:138856923 C>T; a genetic variation in the TCIRG1 gene that is chr11:67818269 G>A; (iii) a genetic variation in the ORAI1 gene is chr12:122064788 G>GT; or (iv) any combination thereof, and wherein chromosome positions of the one or more genetic variations are defined with respect to UCSC hg19.

11. The method of claim 2, wherein the one or more genetic variations comprise (i) a genetic variation in the ITSN2 gene selected from the group consisting of chr2:24431184 C>T, chr2:24432937 C>T, and chr2:24435599 G>A; (ii) a genetic variation in the IFNGR2 gene that is chr21:34809232 C>T; (iii) a genetic variation in the IKBKB gene that is chr8:42176189 G>A; or (iv) any combination thereof, and wherein chromosome positions of the one or more genetic variations are defined with respect to UCSC hg19.

12. The method of claim 2, wherein the one or more genetic variations comprise (i) a genetic variation in the IL21R gene that is chr16:27460020 G>A; (ii) a genetic variation in the LIG4 gene selected from the group consisting of chr13:108861092 G>T and chr13:108863591 G>A; (iii) a genetic variation in the PNPT1 gene that is chr2:55910961 T>C; or (iv) any combination thereof, and wherein chromosome positions of the one or more genetic variations are defined with respect to UCSC hg19.

13. The method of claim 2, wherein the one or more genetic variations comprise (i) a genetic variation in the POLE gene selected from the group consisting of chr12:133201381 T>A, chr12:133202816 C>T, chr12:133209020 G>C, chr12:133220526 T>C, chr12:133220544 C>T, chr12:133237658 T>G, chr12:133245026 G>A, chr12:133252406 C>A, chr12:133253971 C>T, and chr12:133253995 G>A; (ii) a genetic variation in the PSTPIP1 gene that is chr15:77329479 C>T; (iii) a genetic variation in the TAP2 gene that is chr6:32797809 C>T; or (iv) any combination thereof, and wherein chromosome positions of the one or more genetic variations are defined with respect to UCSC hg19.

14. The method of claim 2, wherein the one or more genetic variations comprise (i) a genetic variation in the TNFRSF13B gene that is chr17:16852187 A>G; (ii) a genetic variation in the TRPM2 gene selected from the group consisting of chr21:45786650 C>T, chr21:45795833 G>T, chr21:45815307 T>C, chr21:45815331 G>A, chr21:45815425 C>G, chr21:45820196 C>T, chr21:45826486

G>A, chr21: 45845699 G>A, and 45855099 C>T; or (iii) any combination thereof, and wherein chromosome positions of the one or more genetic variations are defined with respect to UCSC hg19.

15. The method of claim 2, wherein the one or more genetic variations comprise two or more genetic variations.

16. The method of claim 2, wherein the one or more genetic variations comprise at least 5 genetic variations.

17. The method of claim 2, wherein prior to testing the subject for the presence of the one or more genetic variations with the genetic assay the method further comprises obtaining biological samples from subjects with PML and (a) confirming each biological sample is not a duplicate of any other biological sample based on nucleic acid information of the biological samples or (b) determining a sex genotype for each biological sample based on nucleic acid information of the biological samples, and confirming the sex genotype of each biological sample is the same as a sex phenotype of the subject with PML from which the biological sample was obtained.

18. The method of claim 2, wherein the one or more genetic variations comprise a copy number variation as set forth in any one of SEQ ID NOs: 20, 111-113, and 148-154.

19. The method of claim 2, wherein the condition is Crohn's disease.

20. The method of claim 2, wherein the condition is multiple sclerosis or a relapsing form of multiple sclerosis.

* * * * *