

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent	12399123
Kind Code	B1
Date of Patent	August 26, 2025
Inventor(s)	Shah; Preyas et al.

Spatial targeting of analytes

Abstract

Systems and methods are provided for targeting one or more features in a region of interest, and efficiently removing the features from an array of features. The systems and methods can remove one or more beads that contain analytes, intermediate agents, or a combination thereof, from the region of interest of a substrate by emitting light toward the beads on the substrate.

Inventors:	Shah; Preyas (Pleasanton, CA), Ramachandran Iyer; Eswar Prasad (Sunnyvale, CA)
Applicant:	10x Genomics, Inc. (Pleasanton, CA)
Family ID:	1000005553864
Assignee:	10x Genomics, Inc. (Pleasanton, CA)
Appl. No.:	17/172816
Filed:	February 10, 2021

Related U.S. Application Data

us-provisional-application US 62976783 20200214

Publication Classification

Int. Cl.: **G01N21/64** (20060101); **G01N1/02** (20060101); **G01N1/06** (20060101); **G01N1/28** (20060101); **G01N33/543** (20060101); **G02B21/00** (20060101); **G02B21/06** (20060101)

U.S. Cl.:

CPC **G01N21/6452** (20130101); **G01N21/6456** (20130101); **G01N33/543** (20130101); **G02B21/0016** (20130101); **G02B21/06** (20130101); **G01N2001/028** (20130101); **G01N2001/068** (20130101); **G01N2001/284** (20130101); **G01N2021/6419** (20130101); **G01N2021/6421** (20130101)

Field of Classification Search

CPC: G01N (21/6452); G01N (21/6456); G01N (33/543); G01N (2001/028); G01N (2001/068); G01N (2001/284); G01N (2021/6419); G01N (2021/6421); G02B (21/0016); G02B (21/06)

USPC: 506/3

References Cited

U.S. PATENT DOCUMENTS

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
4683195	12/1986	Mullis	N/A	N/A
4683202	12/1986	Mullis	N/A	N/A
4800159	12/1988	Mullis	N/A	N/A
4883867	12/1988	Lee	N/A	N/A
4965188	12/1989	Mullis	N/A	N/A
5002882	12/1990	Lunnen	N/A	N/A
5130238	12/1991	Malek	N/A	N/A
5308751	12/1993	Ohkawa	N/A	N/A
5321130	12/1993	Yue	N/A	N/A
5410030	12/1994	Yue	N/A	N/A
5436134	12/1994	Haugland	N/A	N/A
5455166	12/1994	Walker	N/A	N/A
5494810	12/1995	Barany et al.	N/A	N/A
5503980	12/1995	Cantor	N/A	N/A

5512439	12/1995	Hornes	N/A	N/A
5512462	12/1995	Cheng	N/A	N/A
5582977	12/1995	Yue	N/A	N/A
5599675	12/1996	Brenner	N/A	N/A
5641658	12/1996	Adams	N/A	N/A
5648245	12/1996	Fire et al.	N/A	N/A
5658751	12/1996	Yue	N/A	N/A
5750341	12/1997	Macevicz	N/A	N/A
5763175	12/1997	Brenner	N/A	N/A
5830711	12/1997	Barany et al.	N/A	N/A
5837832	12/1997	Chee et al.	N/A	N/A
5854033	12/1997	Lizardi	N/A	N/A
5863753	12/1998	Haugland	N/A	N/A
5871921	12/1998	Landegren et al.	N/A	N/A
5912148	12/1998	Eggerding	N/A	N/A
6013440	12/1999	Lipshutz	N/A	N/A
6027889	12/1999	Barany et al.	N/A	N/A
6060240	12/1999	Kamb et al.	N/A	N/A
6130073	12/1999	Eggerding	N/A	N/A
6143496	12/1999	Brown	N/A	N/A
6153389	12/1999	Haarer	N/A	N/A
6165714	12/1999	Lane et al.	N/A	N/A
6210891	12/2000	Nyren	N/A	N/A
6210894	12/2000	Brennan	N/A	N/A
6214587	12/2000	Dattagupta	N/A	N/A
6258568	12/2000	Nyren	N/A	N/A
6266459	12/2000	Walt	N/A	N/A
6274320	12/2000	Rothberg	N/A	N/A
6300063	12/2000	Lipshutz et al.	N/A	N/A
6309824	12/2000	Drmanac	N/A	N/A
6344316	12/2001	Lockhart	N/A	N/A
6355431	12/2001	Chee	N/A	N/A
6368801	12/2001	Faruqi	N/A	N/A
6401267	12/2001	Drmanac	N/A	N/A
6404907	12/2001	Gilchrist	N/A	N/A
6432360	12/2001	Church et al.	N/A	N/A
6503713	12/2002	Rana	N/A	N/A
6506561	12/2002	Cheval et al.	N/A	N/A
6544732	12/2002	Chee	N/A	N/A
6620584	12/2002	Chee	N/A	N/A
6632641	12/2002	Brennan	N/A	N/A
6737236	12/2003	Pieken et al.	N/A	N/A
6770441	12/2003	Dickinson	N/A	N/A
6773886	12/2003	Kaufman	N/A	N/A
6787308	12/2003	Balasubramanian	N/A	N/A
6800453	12/2003	Labauer	N/A	N/A
6812005	12/2003	Fan et al.	N/A	N/A
6828100	12/2003	Ronaghi	N/A	N/A
6833246	12/2003	Balasubramanian	N/A	N/A
6859570	12/2004	Walt	N/A	N/A
6864052	12/2004	Drmanac	N/A	N/A
6897023	12/2004	Fu	N/A	N/A
6942968	12/2004	Dickinson et al.	N/A	N/A
7057026	12/2005	Barnes	N/A	N/A
7115400	12/2005	Adessi	N/A	N/A
7118883	12/2005	Inoue	N/A	N/A
7128893	12/2005	Leamon et al.	N/A	N/A
7166431	12/2006	Chee et al.	N/A	N/A
7211414	12/2006	Hardin	N/A	N/A
7255994	12/2006	Lao	N/A	N/A
7258976	12/2006	Mitsuhashi	N/A	N/A
7297518	12/2006	Quake	N/A	N/A
7329492	12/2007	Hardin	N/A	N/A
7361488	12/2007	Fan et al.	N/A	N/A
7378242	12/2007	Hurt	N/A	N/A
7393665	12/2007	Brenner	N/A	N/A
7405281	12/2007	Xu	N/A	N/A
7407757	12/2007	Brenner	N/A	N/A
7537897	12/2008	Brenner	N/A	N/A
7563576	12/2008	Chee	N/A	N/A
7582420	12/2008	Oliphant et al.	N/A	N/A
7595883	12/2008	El Gamal	N/A	N/A
7601498	12/2008	Mao	N/A	N/A
7635566	12/2008	Brenner	N/A	N/A
7674752	12/2009	He	N/A	N/A

7709198	12/2009	Luo et al.	N/A	N/A
7776567	12/2009	Mao	N/A	N/A
7803943	12/2009	Mao	N/A	N/A
7910304	12/2010	Drmanac	N/A	N/A
7955794	12/2010	Shen et al.	N/A	N/A
7960119	12/2010	Chee	N/A	N/A
8003354	12/2010	Shen et al.	N/A	N/A
8148068	12/2011	Brenner	N/A	N/A
8206917	12/2011	Chee	N/A	N/A
8288103	12/2011	Oliphant	N/A	N/A
8460865	12/2012	Chee	N/A	N/A
8481257	12/2012	Van Eijk	N/A	N/A
8603743	12/2012	Liu et al.	N/A	N/A
8604182	12/2012	Luo et al.	N/A	N/A
8815512	12/2013	Van Eijk	N/A	N/A
8835358	12/2013	Fodor	N/A	N/A
8911945	12/2013	Van Eijk	N/A	N/A
8951726	12/2014	Luo et al.	N/A	N/A
9062348	12/2014	Van Eijk	N/A	N/A
9085798	12/2014	Chee	N/A	N/A
9194001	12/2014	Brenner	N/A	N/A
9290808	12/2015	Fodor	N/A	N/A
9290809	12/2015	Fodor	N/A	N/A
9328383	12/2015	Van Eijk	N/A	N/A
9334536	12/2015	Van Eijk	N/A	N/A
9371598	12/2015	Chee	N/A	N/A
9506061	12/2015	Brown et al.	N/A	N/A
9593365	12/2016	Frisen et al.	N/A	N/A
9598728	12/2016	Barany et al.	N/A	N/A
9644204	12/2016	Hindson et al.	N/A	N/A
9694361	12/2016	Bharadwaj	N/A	N/A
9702004	12/2016	Van Eijk	N/A	N/A
9727810	12/2016	Fodor et al.	N/A	N/A
9777324	12/2016	Van Eijk	N/A	N/A
9783841	12/2016	Nolan et al.	N/A	N/A
9834814	12/2016	Peter et al.	N/A	N/A
9850536	12/2016	Oliphant et al.	N/A	N/A
9868979	12/2017	Chee et al.	N/A	N/A
9879313	12/2017	Chee et al.	N/A	N/A
10002316	12/2017	Fodor et al.	N/A	N/A
10023907	12/2017	Van Eijk	N/A	N/A
10030261	12/2017	Frisen et al.	N/A	N/A
10041949	12/2017	Bendall et al.	N/A	N/A
10059990	12/2017	Boyden et al.	N/A	N/A
10208982	12/2018	Bannish et al.	N/A	N/A
10273541	12/2018	Hindson et al.	N/A	N/A
10308982	12/2018	Chee	N/A	N/A
10357771	12/2018	Bharadwaj	N/A	N/A
10415080	12/2018	Dunaway et al.	N/A	N/A
10472669	12/2018	Chee	N/A	N/A
10480022	12/2018	Chee	N/A	N/A
10480029	12/2018	Bent et al.	N/A	N/A
10494667	12/2018	Chee	N/A	N/A
10501777	12/2018	Beechem et al.	N/A	N/A
10550429	12/2019	Harada et al.	N/A	N/A
10590244	12/2019	Delaney et al.	N/A	N/A
10640816	12/2019	Beechem et al.	N/A	N/A
10662468	12/2019	Chee	N/A	N/A
10724078	12/2019	Van Driel et al.	N/A	N/A
10725027	12/2019	Bell	N/A	N/A
10774372	12/2019	Chee et al.	N/A	N/A
10774374	12/2019	Frisen et al.	N/A	N/A
10787701	12/2019	Chee	N/A	N/A
10858702	12/2019	Lucero et al.	N/A	N/A
10913975	12/2020	So et al.	N/A	N/A
10914730	12/2020	Chee et al.	N/A	N/A
10927403	12/2020	Chee et al.	N/A	N/A
10961566	12/2020	Chee	N/A	N/A
10995361	12/2020	Chen et al.	N/A	N/A
11001879	12/2020	Chee	N/A	N/A
11008607	12/2020	Chee	N/A	N/A
11008608	12/2020	Samusik et al.	N/A	N/A
11046996	12/2020	Chee et al.	N/A	N/A
11067567	12/2020	Chee	N/A	N/A
11156603	12/2020	Chee	N/A	N/A

11162132	12/2020	Frisen et al.	N/A	N/A
11168350	12/2020	Nolan et al.	N/A	N/A
11208684	12/2020	Chee	N/A	N/A
11214796	12/2021	Shirai et al.	N/A	N/A
11286515	12/2021	Chee et al.	N/A	N/A
11293917	12/2021	Chee	N/A	N/A
11299774	12/2021	Frisen et al.	N/A	N/A
11313856	12/2021	Chee	N/A	N/A
11332790	12/2021	Chell et al.	N/A	N/A
11352659	12/2021	Frisen et al.	N/A	N/A
11352667	12/2021	Hauling et al.	N/A	N/A
11359228	12/2021	Chee et al.	N/A	N/A
11365442	12/2021	Chee	N/A	N/A
11371086	12/2021	Chee	N/A	N/A
11384386	12/2021	Chee	N/A	N/A
11390912	12/2021	Frisen et al.	N/A	N/A
11401545	12/2021	Chee	N/A	N/A
11407992	12/2021	Dadhwal	N/A	N/A
11408029	12/2021	Katirae et al.	N/A	N/A
11434524	12/2021	Ramachandran Iyer et al.	N/A	N/A
11447807	12/2021	Church et al.	N/A	N/A
11459607	12/2021	Terry et al.	N/A	N/A
11479809	12/2021	Frisen et al.	N/A	N/A
11479810	12/2021	Chee	N/A	N/A
11492612	12/2021	Dadhwal	N/A	N/A
11501440	12/2021	Weisenfeld et al.	N/A	N/A
11505828	12/2021	Chell et al.	N/A	N/A
11512308	12/2021	Gallant et al.	N/A	N/A
11519022	12/2021	Chee	N/A	N/A
11519033	12/2021	Schnall-Levin et al.	N/A	N/A
11530438	12/2021	Persson et al.	N/A	N/A
11535887	12/2021	Gallant et al.	N/A	N/A
11542543	12/2022	Chee	N/A	N/A
11549138	12/2022	Chee	N/A	N/A
11560587	12/2022	Chee	N/A	N/A
11560592	12/2022	Chew et al.	N/A	N/A
11560593	12/2022	Chell et al.	N/A	N/A
11592447	12/2022	Uytingco et al.	N/A	N/A
11608498	12/2022	Gallant et al.	N/A	N/A
11608520	12/2022	Galonska et al.	N/A	N/A
11613773	12/2022	Frisen et al.	N/A	N/A
11618897	12/2022	Kim et al.	N/A	N/A
11618918	12/2022	Chee et al.	N/A	N/A
11624063	12/2022	Dadhwal	N/A	N/A
11624086	12/2022	Uytingco et al.	N/A	N/A
11634756	12/2022	Chee	N/A	N/A
11649485	12/2022	Yin et al.	N/A	N/A
11661626	12/2022	Katirae et al.	N/A	N/A
11680260	12/2022	Kim et al.	N/A	N/A
11692218	12/2022	Engblom et al.	N/A	N/A
11702693	12/2022	Bharadwaj	N/A	N/A
11702698	12/2022	Stoeckius	N/A	N/A
11713480	12/2022	Lee	N/A	N/A
11732292	12/2022	Chee	N/A	N/A
11732299	12/2022	Ramachandran Iyer	N/A	N/A
11732300	12/2022	Bava	N/A	N/A
11733238	12/2022	Chee	N/A	N/A
11739372	12/2022	Frisen et al.	N/A	N/A
11739381	12/2022	Chew et al.	N/A	N/A
11753673	12/2022	Chew et al.	N/A	N/A
11753674	12/2022	Chee et al.	N/A	N/A
11753675	12/2022	Ramachandran Iyer	N/A	N/A
11761030	12/2022	Chee	N/A	N/A
11761038	12/2022	Stoeckius	N/A	N/A
11767550	12/2022	Chee	N/A	N/A
11768175	12/2022	Kim et al.	N/A	N/A
11773433	12/2022	Gallant et al.	N/A	N/A
11781130	12/2022	Dadhwal	N/A	N/A
11788122	12/2022	Frisen et al.	N/A	N/A
11795498	12/2022	Frisen et al.	N/A	N/A
11795507	12/2022	Chell et al.	N/A	N/A
11808769	12/2022	Uytingco et al.	N/A	N/A
11821024	12/2022	Chee et al.	N/A	N/A
11821035	12/2022	Bent et al.	N/A	N/A
11827935	12/2022	Ramachandran Iyer et al.	N/A	N/A

11835462	12/2022	Bava	N/A	N/A
11840687	12/2022	Gallant et al.	N/A	N/A
11840724	12/2022	Chew et al.	N/A	N/A
11845979	12/2022	Engblom et al.	N/A	N/A
11859178	12/2023	Gallant et al.	N/A	N/A
11866767	12/2023	Uytingco et al.	N/A	N/A
11866770	12/2023	Chee	N/A	N/A
11873482	12/2023	Kim et al.	N/A	N/A
11891654	12/2023	Alvarado Martinez et al.	N/A	N/A
11898205	12/2023	Bava	N/A	N/A
11926822	12/2023	Gohil et al.	N/A	N/A
11926863	12/2023	Boutet	N/A	N/A
11926867	12/2023	Yin et al.	N/A	N/A
11933957	12/2023	Tentori et al.	N/A	N/A
11952627	12/2023	Stoeckius	N/A	N/A
11959076	12/2023	Kim et al.	N/A	N/A
11959130	12/2023	Galonska et al.	N/A	N/A
11965213	12/2023	Williams	N/A	N/A
11970739	12/2023	Chew et al.	N/A	N/A
11981958	12/2023	Galonska	N/A	N/A
11981960	12/2023	Lin et al.	N/A	N/A
11981965	12/2023	Chell et al.	N/A	N/A
RE50065	12/2023	Frisen et al.	N/A	N/A
12024741	12/2023	Tentori et al.	N/A	N/A
12031177	12/2023	Tentori et al.	N/A	N/A
12060604	12/2023	Katiraei et al.	N/A	N/A
12071655	12/2023	Sukovich et al.	N/A	N/A
12076701	12/2023	Bava	N/A	N/A
12098417	12/2023	Engblom et al.	N/A	N/A
12098985	12/2023	Cox et al.	N/A	N/A
12110541	12/2023	Bava	N/A	N/A
12117439	12/2023	Delaney et al.	N/A	N/A
12128403	12/2023	Kim et al.	N/A	N/A
12129516	12/2023	Tentori et al.	N/A	N/A
12157124	12/2023	Cox et al.	N/A	N/A
12180543	12/2023	Uytingco et al.	N/A	N/A
12195790	12/2024	Sukovich et al.	N/A	N/A
12203134	12/2024	Nagendran et al.	N/A	N/A
12209280	12/2024	Mignardi et al.	N/A	N/A
12223751	12/2024	Li et al.	N/A	N/A
12228544	12/2024	Kim et al.	N/A	N/A
12234505	12/2024	Chee	N/A	N/A
12241060	12/2024	Kim et al.	N/A	N/A
12241890	12/2024	Delaney et al.	N/A	N/A
12249085	12/2024	Tentori et al.	N/A	N/A
12265079	12/2024	Bent	N/A	N/A
12270077	12/2024	Schnall-Levin et al.	N/A	N/A
12275988	12/2024	Galonska et al.	N/A	N/A
12281357	12/2024	Tentori et al.	N/A	N/A
12286673	12/2024	Bava	N/A	N/A
12287264	12/2024	Cox et al.	N/A	N/A
2002/0040275	12/2001	Cravatt	N/A	N/A
2002/0164611	12/2001	Bamdad	N/A	N/A
2003/0017451	12/2002	Wang et al.	N/A	N/A
2003/0022207	12/2002	Balasubramanian	N/A	N/A
2003/0148335	12/2002	Shen et al.	N/A	N/A
2003/0162216	12/2002	Gold	N/A	N/A
2003/0211489	12/2002	Shen et al.	N/A	N/A
2003/0224419	12/2002	Corcoran	N/A	N/A
2003/0232348	12/2002	Jones et al.	N/A	N/A
2003/0232382	12/2002	Brennan	N/A	N/A
2004/0033499	12/2003	Ilseley et al.	N/A	N/A
2004/0067492	12/2003	Peng et al.	N/A	N/A
2004/0096853	12/2003	Mayer	N/A	N/A
2004/0106110	12/2003	Balasubramanian	N/A	N/A
2005/0037393	12/2004	Gunderson et al.	N/A	N/A
2005/0048580	12/2004	Labaer	N/A	N/A
2005/0100900	12/2004	Kawashima et al.	N/A	N/A
2005/0130173	12/2004	Leamon et al.	N/A	N/A
2005/0136414	12/2004	Gunderson et al.	N/A	N/A
2005/0170373	12/2004	Monforte	N/A	N/A
2005/0191656	12/2004	Drmanac et al.	N/A	N/A
2005/0191698	12/2004	Chee et al.	N/A	N/A
2005/0196786	12/2004	Levy	N/A	N/A
2005/0202433	12/2004	Van Beuningen	N/A	N/A

2005/0227271	12/2004	Kwon	N/A	N/A
2005/0260653	12/2004	LaBaer	N/A	N/A
2006/0041385	12/2005	Bauer et al.	N/A	N/A
2006/0211001	12/2005	Yu et al.	N/A	N/A
2006/0216775	12/2005	Burkart et al.	N/A	N/A
2006/0263789	12/2005	Kincaid	N/A	N/A
2007/0020640	12/2006	McCloskey et al.	N/A	N/A
2007/0054288	12/2006	Su et al.	N/A	N/A
2007/0099208	12/2006	Drmanac et al.	N/A	N/A
2007/0128624	12/2006	Gormley et al.	N/A	N/A
2007/0128656	12/2006	Agrawal	N/A	N/A
2007/0172873	12/2006	Brenner et al.	N/A	N/A
2007/0207482	12/2006	Church et al.	N/A	N/A
2007/0251824	12/2006	Patton	N/A	N/A
2007/0254305	12/2006	Paik et al.	N/A	N/A
2007/0269805	12/2006	Hogers	N/A	N/A
2007/0280517	12/2006	De La Torre-Bueno et al.	N/A	N/A
2008/0009420	12/2007	Schroth et al.	N/A	N/A
2008/0108804	12/2007	Hayashizaki et al.	N/A	N/A
2008/0160580	12/2007	Adessi et al.	N/A	N/A
2008/0199929	12/2007	Yeung et al.	N/A	N/A
2008/0220434	12/2007	Thomas	N/A	N/A
2008/0261204	12/2007	Lexow	N/A	N/A
2008/0286795	12/2007	Kawashima et al.	N/A	N/A
2009/0005252	12/2008	Drmanac et al.	N/A	N/A
2009/0006002	12/2008	Honisch et al.	N/A	N/A
2009/0018024	12/2008	Church et al.	N/A	N/A
2009/0026082	12/2008	Rothberg et al.	N/A	N/A
2009/0082212	12/2008	Williams	N/A	N/A
2009/0099041	12/2008	Church et al.	N/A	N/A
2009/0105959	12/2008	Braverman et al.	N/A	N/A
2009/0117573	12/2008	Fu et al.	N/A	N/A
2009/0127589	12/2008	Rothberg et al.	N/A	N/A
2009/0155781	12/2008	Drmanac et al.	N/A	N/A
2009/0197326	12/2008	El Gamal et al.	N/A	N/A
2009/0233802	12/2008	Bignell et al.	N/A	N/A
2009/0253581	12/2008	van Eijk et al.	N/A	N/A
2009/0270273	12/2008	Burns et al.	N/A	N/A
2009/0289184	12/2008	Deininger	N/A	N/A
2009/0291854	12/2008	Weisinger-Mayr et al.	N/A	N/A
2009/0305237	12/2008	Cantor et al.	N/A	N/A
2009/0312193	12/2008	Kim et al.	N/A	N/A
2010/0035249	12/2009	Hayashizaki et al.	N/A	N/A
2010/0120097	12/2009	Matz et al.	N/A	N/A
2010/0120098	12/2009	Grunenwald et al.	N/A	N/A
2010/0145037	12/2009	Brive et al.	N/A	N/A
2011/0028685	12/2010	Purkayastha et al.	N/A	N/A
2011/0059436	12/2010	Hardin et al.	N/A	N/A
2011/0152111	12/2010	Fan et al.	N/A	N/A
2011/0245111	12/2010	Chee	N/A	N/A
2011/0275077	12/2010	James	N/A	N/A
2012/0135871	12/2011	van Eijk et al.	N/A	N/A
2012/0202698	12/2011	van Eijk et al.	N/A	N/A
2012/0245053	12/2011	Shirai et al.	N/A	N/A
2013/0040842	12/2012	Lim	506/9	C12Q 1/6806
2013/0122516	12/2012	Meares	N/A	N/A
2013/0171621	12/2012	Luo et al.	N/A	N/A
2014/0066318	12/2013	Frisen et al.	N/A	N/A
2014/0155274	12/2013	Xie et al.	N/A	N/A
2014/0270435	12/2013	Dunn	N/A	N/A
2014/0274731	12/2013	Raymond et al.	N/A	N/A
2014/0323330	12/2013	Glezer et al.	N/A	N/A
2014/0378350	12/2013	Hindson et al.	N/A	N/A
2015/0000854	12/2014	Gann-Fetter et al.	N/A	N/A
2015/0072867	12/2014	Soldatov	N/A	N/A
2015/0148239	12/2014	Jon	N/A	N/A
2015/0292988	12/2014	Bharadwaj et al.	N/A	N/A
2015/0344942	12/2014	Frisen et al.	N/A	N/A
2016/0108458	12/2015	Frei et al.	N/A	N/A
2016/0138091	12/2015	Chee et al.	N/A	N/A
2016/0145677	12/2015	Chee et al.	N/A	N/A
2016/0253584	12/2015	Fodor et al.	N/A	N/A
2016/0289740	12/2015	Fu et al.	N/A	N/A
2016/0298180	12/2015	Chee	N/A	N/A
2017/0016053	12/2016	Beechem et al.	N/A	N/A

2017/0016909	12/2016	Beechem et al.	N/A	N/A
2017/0029875	12/2016	Zhang et al.	N/A	N/A
2017/0067096	12/2016	Wassie et al.	N/A	N/A
2017/0089811	12/2016	Tillberg et al.	N/A	N/A
2017/0159109	12/2016	Zheng et al.	N/A	N/A
2017/0220733	12/2016	Zhuang et al.	N/A	N/A
2017/0233722	12/2016	Seelig et al.	N/A	N/A
2017/0241911	12/2016	Rockel et al.	N/A	N/A
2017/0275669	12/2016	Weissleder et al.	N/A	N/A
2017/0342405	12/2016	Fu et al.	N/A	N/A
2018/0051322	12/2017	Church et al.	N/A	N/A
2018/0057873	12/2017	Zhou et al.	N/A	N/A
2018/0088112	12/2017	Fan et al.	N/A	N/A
2018/0112261	12/2017	Van Driel et al.	N/A	N/A
2018/0201980	12/2017	Chee et al.	N/A	N/A
2018/0216161	12/2017	Chen et al.	N/A	N/A
2018/0216162	12/2017	Belhocine et al.	N/A	N/A
2018/0245142	12/2017	So et al.	N/A	N/A
2018/0291439	12/2017	van Eijk et al.	N/A	N/A
2018/0305681	12/2017	Jovanovich et al.	N/A	N/A
2018/0334670	12/2017	Bharadwaj et al.	N/A	N/A
2019/0017106	12/2018	Frisen et al.	N/A	N/A
2019/0024153	12/2018	Frisen et al.	N/A	N/A
2019/0024154	12/2018	Frisen et al.	N/A	N/A
2019/0055594	12/2018	Samusik et al.	N/A	N/A
2019/0064173	12/2018	Bharadwaj et al.	N/A	N/A
2019/0085383	12/2018	Church et al.	N/A	N/A
2019/0161796	12/2018	Hauling et al.	N/A	N/A
2019/0177777	12/2018	Chee	N/A	N/A
2019/0177778	12/2018	Chee	N/A	N/A
2019/0177789	12/2018	Hindson et al.	N/A	N/A
2019/0177800	12/2018	Boutet et al.	N/A	N/A
2019/0194709	12/2018	Church et al.	N/A	N/A
2019/0203275	12/2018	Frisen et al.	N/A	N/A
2019/0233878	12/2018	Delaney et al.	N/A	N/A
2019/0249226	12/2018	Bent et al.	N/A	N/A
2019/0249248	12/2018	Beechem et al.	N/A	N/A
2019/0262831	12/2018	West et al.	N/A	N/A
2019/0264268	12/2018	Frisen et al.	N/A	N/A
2019/0271030	12/2018	Chee	N/A	N/A
2019/0271031	12/2018	Chee	N/A	N/A
2019/0300943	12/2018	Chee et al.	N/A	N/A
2019/0300944	12/2018	Chee et al.	N/A	N/A
2019/0300945	12/2018	Chee et al.	N/A	N/A
2019/0309353	12/2018	Chee	N/A	N/A
2019/0309354	12/2018	Chee	N/A	N/A
2019/0309355	12/2018	Chee	N/A	N/A
2019/0323071	12/2018	Chee	N/A	N/A
2019/0323088	12/2018	Boutet et al.	N/A	N/A
2019/0330617	12/2018	Church et al.	N/A	N/A
2019/0338353	12/2018	Belgrader et al.	N/A	N/A
2019/0367969	12/2018	Belhocine et al.	N/A	N/A
2019/0367982	12/2018	Belhocine et al.	N/A	N/A
2019/0367997	12/2018	Bent et al.	N/A	N/A
2020/0002763	12/2019	Belgrader et al.	N/A	N/A
2020/0024641	12/2019	Nolan et al.	N/A	N/A
2020/0047010	12/2019	Lee et al.	N/A	N/A
2020/0048690	12/2019	Chee	N/A	N/A
2020/0063191	12/2019	Kennedy-Darling et al.	N/A	N/A
2020/0063195	12/2019	Chee	N/A	N/A
2020/0063196	12/2019	Chee	N/A	N/A
2020/0071751	12/2019	Daugharthy et al.	N/A	N/A
2020/0080136	12/2019	Zhang et al.	N/A	N/A
2020/0109443	12/2019	Chee	N/A	N/A
2020/0173985	12/2019	Dong et al.	N/A	N/A
2020/0224244	12/2019	Nilsson et al.	N/A	N/A
2020/0239946	12/2019	Dewal	N/A	N/A
2020/0256867	12/2019	Hennek et al.	N/A	N/A
2020/0277663	12/2019	Iyer	N/A	N/A
2020/0277664	12/2019	Frenz	N/A	N/A
2020/0299757	12/2019	Chee et al.	N/A	N/A
2020/0325531	12/2019	Chee	N/A	N/A
2020/0370095	12/2019	Farmer et al.	N/A	N/A
2020/0399687	12/2019	Frisen et al.	N/A	N/A
2020/0407781	12/2019	Schnall-Levin	N/A	N/A

2021/0010068	12/2020	Chee et al.	N/A	N/A
2021/0010070	12/2020	Schnall-Levin et al.	N/A	N/A
2021/0095331	12/2020	Fan et al.	N/A	N/A
2021/0123040	12/2020	Macosko et al.	N/A	N/A
2021/0130881	12/2020	Cox	N/A	N/A
2021/0140982	12/2020	Uytingco et al.	N/A	N/A
2021/0150707	12/2020	Weisenfeld et al.	N/A	N/A
2021/0155982	12/2020	Yin et al.	N/A	N/A
2021/0158522	12/2020	Weisenfeld et al.	N/A	N/A
2021/0172007	12/2020	Chee et al.	N/A	N/A
2021/0189475	12/2020	Tentori et al.	N/A	N/A
2021/0190770	12/2020	Delaney et al.	N/A	N/A
2021/0198741	12/2020	Williams	N/A	N/A
2021/0199660	12/2020	Williams et al.	N/A	N/A
2021/0207202	12/2020	Chee	N/A	N/A
2021/0214785	12/2020	Stoeckius	N/A	N/A
2021/0222235	12/2020	Chee	N/A	N/A
2021/0222241	12/2020	Bharadwaj	N/A	N/A
2021/0222242	12/2020	Ramachandran Iyer	N/A	N/A
2021/0222253	12/2020	Uytingco	N/A	N/A
2021/0223227	12/2020	Stoeckius	N/A	N/A
2021/0230584	12/2020	Mikkelsen et al.	N/A	N/A
2021/0230681	12/2020	Patterson et al.	N/A	N/A
2021/0230692	12/2020	Daugharthy et al.	N/A	N/A
2021/0237022	12/2020	Bava	N/A	N/A
2021/0238581	12/2020	Mikkelsen et al.	N/A	N/A
2021/0238664	12/2020	Bava et al.	N/A	N/A
2021/0238675	12/2020	Bava	N/A	N/A
2021/0238680	12/2020	Bava	N/A	N/A
2021/0247316	12/2020	Bava	N/A	N/A
2021/0255175	12/2020	Chee et al.	N/A	N/A
2021/0262018	12/2020	Bava et al.	N/A	N/A
2021/0262019	12/2020	Alvarado Martinez et al.	N/A	N/A
2021/0269864	12/2020	Chee	N/A	N/A
2021/0270822	12/2020	Chee	N/A	N/A
2021/0285036	12/2020	Yin et al.	N/A	N/A
2021/0285046	12/2020	Chell et al.	N/A	N/A
2021/0292748	12/2020	Frisen et al.	N/A	N/A
2021/0292822	12/2020	Frisen et al.	N/A	N/A
2021/0317510	12/2020	Chee et al.	N/A	N/A
2021/0317524	12/2020	Lucero et al.	N/A	N/A
2021/0324457	12/2020	Ramachandran Iyer et al.	N/A	N/A
2021/0332424	12/2020	Schnall-Levin	N/A	N/A
2021/0332425	12/2020	Pfeiffer et al.	N/A	N/A
2021/0348221	12/2020	Chell et al.	N/A	N/A
2022/0002791	12/2021	Frisen et al.	N/A	N/A
2022/0003755	12/2021	Chee	N/A	N/A
2022/0010367	12/2021	Ramachandran Iyer et al.	N/A	N/A
2022/0017951	12/2021	Ramachandran Iyer et al.	N/A	N/A
2022/0025446	12/2021	Shah	N/A	N/A
2022/0025447	12/2021	Tentori et al.	N/A	N/A
2022/0033888	12/2021	Schnall-Levin et al.	N/A	N/A
2022/0049293	12/2021	Frenz et al.	N/A	N/A
2022/0049294	12/2021	Uytingco et al.	N/A	N/A
2022/0064630	12/2021	Bent et al.	N/A	N/A
2022/0081728	12/2021	Williams	N/A	N/A
2022/0090058	12/2021	Frisen et al.	N/A	N/A
2022/0090175	12/2021	Uytingco et al.	N/A	N/A
2022/0090181	12/2021	Gallant et al.	N/A	N/A
2022/0098576	12/2021	Dadhwaj	N/A	N/A
2022/0098661	12/2021	Chew et al.	N/A	N/A
2022/0106632	12/2021	Galonska et al.	N/A	N/A
2022/0106633	12/2021	Engblom et al.	N/A	N/A
2022/0112486	12/2021	Ramachandran Iyer et al.	N/A	N/A
2022/0112545	12/2021	Chee	N/A	N/A
2022/0119869	12/2021	Ramachandran Iyer et al.	N/A	N/A
2022/0127659	12/2021	Frisen et al.	N/A	N/A
2022/0127666	12/2021	Katirae et al.	N/A	N/A
2022/0127672	12/2021	Stoeckius	N/A	N/A
2022/0145361	12/2021	Frenz et al.	N/A	N/A
2022/0154255	12/2021	Chee et al.	N/A	N/A
2022/0170083	12/2021	Khaled et al.	N/A	N/A
2022/0195422	12/2021	Gallant et al.	N/A	N/A
2022/0195505	12/2021	Frisen et al.	N/A	N/A
2022/0196644	12/2021	Chee	N/A	N/A

2022/0213526	12/2021	Frisen et al.	N/A	N/A
2022/0220544	12/2021	Ach et al.	N/A	N/A
2022/0241780	12/2021	Tentori et al.	N/A	N/A
2022/0267844	12/2021	Ramachandran Iyer et al.	N/A	N/A
2022/0282329	12/2021	Chell et al.	N/A	N/A
2022/0290217	12/2021	Frenz et al.	N/A	N/A
2022/0290219	12/2021	Chee	N/A	N/A
2022/0298560	12/2021	Frisen et al.	N/A	N/A
2022/0325325	12/2021	Chee et al.	N/A	N/A
2022/0326251	12/2021	Uytingco et al.	N/A	N/A
2022/0333171	12/2021	Chee	N/A	N/A
2022/0333191	12/2021	Mikkelsen et al.	N/A	N/A
2022/0333192	12/2021	Uytingco	N/A	N/A
2022/0333195	12/2021	Schnall-Levin et al.	N/A	N/A
2022/0334031	12/2021	Delaney et al.	N/A	N/A
2022/0348905	12/2021	Dadhwal	N/A	N/A
2022/0348992	12/2021	Stoeckius et al.	N/A	N/A
2022/0356464	12/2021	Kim et al.	N/A	N/A
2022/0364163	12/2021	Stahl et al.	N/A	N/A
2022/0389491	12/2021	Chee	N/A	N/A
2022/0389503	12/2021	Mikkelsen et al.	N/A	N/A
2022/0389504	12/2021	Chew et al.	N/A	N/A
2022/0403455	12/2021	Ramachandran Iyer et al.	N/A	N/A
2022/0404245	12/2021	Chell et al.	N/A	N/A
2023/0002812	12/2022	Stoeckius et al.	N/A	N/A
2023/0014008	12/2022	Shastri	N/A	N/A
2023/0017773	12/2022	Kim et al.	N/A	N/A
2023/0031305	12/2022	Hernandez Neuta et al.	N/A	N/A
2023/0033960	12/2022	Gallant et al.	N/A	N/A
2023/0034039	12/2022	Shahjamali	N/A	N/A
2023/0034216	12/2022	Bava	N/A	N/A
2023/0040363	12/2022	Chee	N/A	N/A
2023/0042088	12/2022	Chee	N/A	N/A
2023/0042817	12/2022	Mignardi	N/A	N/A
2023/0047782	12/2022	Tentori et al.	N/A	N/A
2023/0056549	12/2022	Dadhwal	N/A	N/A
2023/0064372	12/2022	Chell et al.	N/A	N/A
2023/0069046	12/2022	Chew et al.	N/A	N/A
2023/0077364	12/2022	Patterson et al.	N/A	N/A
2023/0080543	12/2022	Katirae et al.	N/A	N/A
2023/0081381	12/2022	Chew et al.	N/A	N/A
2023/0100497	12/2022	Frisen et al.	N/A	N/A
2023/0107023	12/2022	Chee	N/A	N/A
2023/0111225	12/2022	Chew et al.	N/A	N/A
2023/0113230	12/2022	Kim et al.	N/A	N/A
2023/0126825	12/2022	Nagendran et al.	N/A	N/A
2023/0129552	12/2022	Ramachandran Iyer	N/A	N/A
2023/0135010	12/2022	Tentori et al.	N/A	N/A
2023/0143569	12/2022	Iyer et al.	N/A	N/A
2023/0145575	12/2022	Gallant et al.	N/A	N/A
2023/0147726	12/2022	Hadrup et al.	N/A	N/A
2023/0151412	12/2022	Chee	N/A	N/A
2023/0159994	12/2022	Chee	N/A	N/A
2023/0159995	12/2022	Iyer et al.	N/A	N/A
2023/0160008	12/2022	Chell et al.	N/A	N/A
2023/0175045	12/2022	Katsori et al.	N/A	N/A
2023/0183684	12/2022	Gallant et al.	N/A	N/A
2023/0183785	12/2022	Frisen et al.	N/A	N/A
2023/0194469	12/2022	Tentori et al.	N/A	N/A
2023/0194470	12/2022	Kim et al.	N/A	N/A
2023/0203478	12/2022	Kim et al.	N/A	N/A
2023/0212650	12/2022	Chew et al.	N/A	N/A
2023/0212655	12/2022	Chee	N/A	N/A
2023/0212656	12/2022	Chow et al.	N/A	N/A
2023/0220368	12/2022	Kim	N/A	N/A
2023/0220454	12/2022	Bent et al.	N/A	N/A
2023/0220455	12/2022	Galonska et al.	N/A	N/A
2023/0227811	12/2022	Dadhwal	N/A	N/A
2023/0228762	12/2022	Uytingco et al.	N/A	N/A
2023/0242973	12/2022	Frisen et al.	N/A	N/A
2023/0242976	12/2022	Tentori et al.	N/A	N/A
2023/0265488	12/2022	Gohil et al.	N/A	N/A
2023/0265489	12/2022	Uytingco et al.	N/A	N/A
2023/0265491	12/2022	Tentori et al.	N/A	N/A
2023/0267625	12/2022	Tentori et al.	N/A	N/A

2023/0279474	12/2022	Katirae	N/A	N/A
2023/0279477	12/2022	Kvastad et al.	N/A	N/A
2023/0279481	12/2022	Marrache et al.	N/A	N/A
2023/0287399	12/2022	Gallant et al.	N/A	N/A
2023/0287475	12/2022	Chell et al.	N/A	N/A
2023/0287481	12/2022	Katsori et al.	N/A	N/A
2023/0295699	12/2022	Sukovich et al.	N/A	N/A
2023/0295722	12/2022	Bharadwaj	N/A	N/A
2023/0304072	12/2022	Gohil et al.	N/A	N/A
2023/0304074	12/2022	Chee et al.	N/A	N/A
2023/0304078	12/2022	Frisen et al.	N/A	N/A
2023/0313279	12/2022	Giacomello et al.	N/A	N/A
2023/0323340	12/2022	Dadhwal	N/A	N/A
2023/0323434	12/2022	Yin et al.	N/A	N/A
2023/0323436	12/2022	Chee	N/A	N/A
2023/0323447	12/2022	Schnall-Levin et al.	N/A	N/A
2023/0323453	12/2022	Stoeckius	N/A	N/A
2023/0332138	12/2022	Kim et al.	N/A	N/A
2023/0332211	12/2022	Chee	N/A	N/A
2023/0332212	12/2022	Chew et al.	N/A	N/A
2023/0332227	12/2022	Ramachandran Iyer	N/A	N/A
2023/0332247	12/2022	Singh et al.	N/A	N/A
2023/0351619	12/2022	Tentori et al.	N/A	N/A
2023/0358733	12/2022	Chee	N/A	N/A
2023/0366008	12/2022	Chew et al.	N/A	N/A
2023/0383285	12/2022	Kim et al.	N/A	N/A
2023/0383344	12/2022	Stoeckius	N/A	N/A
2023/0392204	12/2022	Chell et al.	N/A	N/A
2023/0393071	12/2022	Bava	N/A	N/A
2023/0407404	12/2022	Baumgartner et al.	N/A	N/A
2023/0416807	12/2022	Chee	N/A	N/A
2023/0416808	12/2022	Sukovich et al.	N/A	N/A
2023/0416850	12/2022	Singh et al.	N/A	N/A
2024/0002931	12/2023	Bava	N/A	N/A
2024/0011081	12/2023	Chee	N/A	N/A
2024/0011090	12/2023	Chew et al.	N/A	N/A
2024/0018572	12/2023	Mignardi	N/A	N/A
2024/0018575	12/2023	Gallant et al.	N/A	N/A
2024/0018589	12/2023	Schnall-Levin et al.	N/A	N/A
2024/0026445	12/2023	Ramachandran Iyer et al.	N/A	N/A
2024/0033743	12/2023	Tentori et al.	N/A	N/A
2024/0035937	12/2023	Cox et al.	N/A	N/A
2024/0043908	12/2023	Chew et al.	N/A	N/A
2024/0043925	12/2023	Bent et al.	N/A	N/A
2024/0052343	12/2023	Gallant et al.	N/A	N/A
2024/0053351	12/2023	Uytingco et al.	N/A	N/A
2024/0060115	12/2023	Chee et al.	N/A	N/A
2024/0067953	12/2023	Mikkelsen et al.	N/A	N/A
2024/0068016	12/2023	Frisen et al.	N/A	N/A
2024/0068017	12/2023	Lundeberg et al.	N/A	N/A
2024/0076723	12/2023	Mignardi	N/A	N/A
2024/0080346	12/2023	Engblom et al.	N/A	N/A
2024/0084365	12/2023	Frisen et al.	N/A	N/A
2024/0084366	12/2023	Chee	N/A	N/A
2024/0084383	12/2023	Ramachandran Iyer et al.	N/A	N/A
2024/0093274	12/2023	Frisen et al.	N/A	N/A
2024/0093290	12/2023	Stahl et al.	N/A	N/A
2024/0110228	12/2023	Uytingco et al.	N/A	N/A
2024/0124933	12/2023	Chell et al.	N/A	N/A
2024/0125772	12/2023	Delaney et al.	N/A	N/A
2024/0141327	12/2023	Kim et al.	N/A	N/A
2024/0158838	12/2023	Alvarado Martinez et al.	N/A	N/A
2024/0175080	12/2023	Galonska et al.	N/A	N/A
2024/0182968	12/2023	Bava	N/A	N/A
2024/0191286	12/2023	Boutet et al.	N/A	N/A
2024/0200121	12/2023	Boutet	N/A	N/A
2024/0209425	12/2023	Yin et al.	N/A	N/A
2024/0218427	12/2023	Sukovich et al.	N/A	N/A
2024/0218432	12/2023	Mielinis	N/A	N/A
2024/0219701	12/2023	Tentori et al.	N/A	N/A
2024/0253036	12/2023	Kim et al.	N/A	N/A
2024/0263218	12/2023	Katirae et al.	N/A	N/A
2024/0271190	12/2023	Stoeckius et al.	N/A	N/A
2024/0271195	12/2023	Mikhael et al.	N/A	N/A
2024/0279747	12/2023	Williams	N/A	N/A

2024/0287600	12/2023	Iyer et al.	N/A	N/A
2024/0294971	12/2023	Galonska	N/A	N/A
2024/0294974	12/2023	Galonska et al.	N/A	N/A
2024/0294975	12/2023	Lin et al.	N/A	N/A
2024/0301488	12/2023	Stoeckius	N/A	N/A
2024/0301489	12/2023	Chew et al.	N/A	N/A
2024/0360494	12/2023	Costa et al.	N/A	N/A
2024/0368711	12/2023	Giacomello et al.	N/A	N/A
2024/0377297	12/2023	Cox et al.	N/A	N/A
2024/0385088	12/2023	Kim et al.	N/A	N/A
2024/0392349	12/2023	Frisen et al.	N/A	N/A
2024/0392351	12/2023	Chee	N/A	N/A
2024/0392352	12/2023	Stahl et al.	N/A	N/A
2024/0392353	12/2023	Engblom et al.	N/A	N/A
2024/0401109	12/2023	Kim et al.	N/A	N/A
2024/0401117	12/2023	Bava	N/A	N/A
2024/0401118	12/2023	Tentori et al.	N/A	N/A
2024/0404301	12/2023	Li et al.	N/A	N/A
2024/0408593	12/2023	Kim et al.	N/A	N/A
2024/0416315	12/2023	Bava	N/A	N/A
2024/0417783	12/2023	Chew et al.	N/A	N/A
2024/0417784	12/2023	Sukovich et al.	N/A	N/A
2025/0002980	12/2024	Tentori et al.	N/A	N/A
2025/0002982	12/2024	Stoeckius et al.	N/A	N/A
2025/0003956	12/2024	Delaney et al.	N/A	N/A
2025/0019689	12/2024	Galonska et al.	N/A	N/A
2025/0019749	12/2024	Katirae et al.	N/A	N/A
2025/0066762	12/2024	Man et al.	N/A	N/A
2025/0066770	12/2024	Costa	N/A	N/A
2025/0073719	12/2024	Cox et al.	N/A	N/A
2025/0075261	12/2024	Kim	N/A	N/A
2025/0101501	12/2024	Chee	N/A	N/A
2025/0101502	12/2024	Chee	N/A	N/A
2025/0101504	12/2024	Nagendran et al.	N/A	N/A
2025/0122564	12/2024	Mignardi et al.	N/A	N/A
2025/0122565	12/2024	Schnall-Levin et al.	N/A	N/A
2025/0129412	12/2024	Uyingco et al.	N/A	N/A
2025/0129421	12/2024	Schnall-Levin et al.	N/A	N/A
2025/0137043	12/2024	Tentori	N/A	N/A

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
1680604	12/2004	CN	N/A
1923471	12/2007	EP	N/A
2002017	12/2007	EP	N/A
2130913	12/2008	EP	N/A
2881465	12/2014	EP	N/A
2987792	12/2015	EP	A61K 49/0023
3013984	12/2015	EP	N/A
2350648	12/2016	EP	N/A
3511423	12/2018	EP	N/A
3541956	12/2018	EP	N/A
2007-014297	12/2006	JP	N/A
2009-036694	12/2008	JP	N/A
WO 1989/010977	12/1988	WO	N/A
WO 1991/006678	12/1990	WO	N/A
WO 1995/025116	12/1994	WO	N/A
WO 1995/035505	12/1994	WO	N/A
WO 2001/007915	12/2000	WO	N/A
WO 2002/059355	12/2001	WO	N/A
WO 2002/077283	12/2001	WO	N/A
WO 2003/002979	12/2002	WO	N/A
WO 2003/010176	12/2002	WO	N/A
WO 2005/007814	12/2004	WO	N/A
WO 2007/073171	12/2006	WO	N/A
WO 2007/076726	12/2006	WO	N/A
WO 2007/145612	12/2006	WO	N/A
WO 2008/075086	12/2007	WO	N/A
WO 2009/032167	12/2008	WO	N/A
WO 2009/152928	12/2008	WO	N/A
WO 2009/156725	12/2008	WO	N/A
WO 2010/088517	12/2009	WO	N/A
WO 2010/126614	12/2009	WO	N/A
WO 2011/068088	12/2010	WO	N/A
WO 2011/127099	12/2010	WO	N/A

WO 2012/140224	12/2011	WO	N/A
WO 2012/159089	12/2011	WO	N/A
WO 2012/168003	12/2011	WO	N/A
WO 2013/123442	12/2012	WO	N/A
WO 2013/131962	12/2012	WO	N/A
WO 2013/138510	12/2012	WO	N/A
WO 2013/150082	12/2012	WO	N/A
WO 2013/150083	12/2012	WO	N/A
WO 2014/060483	12/2013	WO	N/A
WO 2014/210223	12/2013	WO	N/A
WO 2014/210225	12/2013	WO	N/A
WO 2015/031691	12/2014	WO	N/A
WO 2016/138496	12/2015	WO	N/A
WO 2016/138500	12/2015	WO	N/A
WO 2016/166128	12/2015	WO	N/A
WO 2016/168825	12/2015	WO	N/A
WO 2017/019456	12/2016	WO	N/A
WO 2017/075293	12/2016	WO	N/A
WO 2017/096158	12/2016	WO	N/A
WO 2018/064640	12/2017	WO	N/A
WO 2018/091676	12/2017	WO	N/A
WO 2018/175779	12/2017	WO	N/A
WO 2019/213254	12/2018	WO	N/A
WO 2019/213294	12/2018	WO	N/A
WO 2020/028194	12/2019	WO	N/A
WO 2020/047002	12/2019	WO	N/A
WO 2020/047005	12/2019	WO	N/A
WO 2020/047010	12/2019	WO	N/A
WO 2020/053655	12/2019	WO	N/A
WO 2020/061064	12/2019	WO	N/A
WO 2020/061066	12/2019	WO	N/A
WO 2020/061108	12/2019	WO	N/A
WO 2020/076979	12/2019	WO	N/A
WO 2020/099640	12/2019	WO	N/A
WO 2020/123301	12/2019	WO	N/A
WO 2020/123305	12/2019	WO	N/A
WO 2020/123309	12/2019	WO	N/A
WO 2020/123311	12/2019	WO	N/A
WO 2020/123316	12/2019	WO	N/A
WO 2020/123317	12/2019	WO	N/A
WO 2020/123318	12/2019	WO	N/A
WO 2020/123319	12/2019	WO	N/A
WO 2020/123320	12/2019	WO	N/A
WO 2020/160044	12/2019	WO	N/A
WO 2020/167862	12/2019	WO	N/A
WO 2020/176788	12/2019	WO	N/A
WO 2020/176882	12/2019	WO	N/A
WO 2020/190509	12/2019	WO	N/A
WO 2020/198071	12/2019	WO	N/A
WO 2020/206285	12/2019	WO	N/A
WO 2020/219901	12/2019	WO	N/A
WO 2020/243579	12/2019	WO	N/A
WO 2021/041974	12/2020	WO	N/A
WO 2021/067246	12/2020	WO	N/A
WO 2021/067514	12/2020	WO	N/A
WO 2021/091611	12/2020	WO	N/A
WO 2021/092433	12/2020	WO	N/A
WO 2021/097255	12/2020	WO	N/A
WO 2021/102003	12/2020	WO	N/A
WO 2021/102005	12/2020	WO	N/A
WO 2021/102039	12/2020	WO	N/A
WO 2021/116715	12/2020	WO	N/A
WO 2021/133842	12/2020	WO	N/A
WO 2021/133845	12/2020	WO	N/A
WO 2021/133849	12/2020	WO	N/A
WO 2021/142233	12/2020	WO	N/A
WO 2021/168261	12/2020	WO	N/A
WO 2021/168278	12/2020	WO	N/A
WO 2021/207610	12/2020	WO	N/A
WO 2021/216708	12/2020	WO	N/A
WO 2021/225900	12/2020	WO	N/A
WO 2021/236625	12/2020	WO	N/A
WO 2021/236929	12/2020	WO	N/A
WO 2021/237056	12/2020	WO	N/A
WO 2021/237087	12/2020	WO	N/A

WO 2021/242834	12/2020	WO	N/A
WO 2021/247543	12/2020	WO	N/A
WO 2021/247568	12/2020	WO	N/A
WO 2021/247593	12/2020	WO	N/A
WO 2021/252499	12/2020	WO	N/A
WO 2021/252576	12/2020	WO	N/A
WO 2021/252591	12/2020	WO	N/A
WO 2021/252747	12/2020	WO	N/A
WO 2021/263111	12/2020	WO	N/A
WO 2022/025965	12/2021	WO	N/A
WO 2022/060798	12/2021	WO	N/A
WO 2022/060953	12/2021	WO	N/A
WO 2022/061150	12/2021	WO	N/A
WO 2022/061152	12/2021	WO	N/A
WO 2022/087273	12/2021	WO	N/A
WO 2022/099037	12/2021	WO	N/A
WO 2022/103712	12/2021	WO	N/A
WO 2022/109181	12/2021	WO	N/A
WO 2022/132645	12/2021	WO	N/A
WO 2022/140028	12/2021	WO	N/A
WO 2022/147005	12/2021	WO	N/A
WO 2022/147296	12/2021	WO	N/A
WO 2022/164615	12/2021	WO	N/A
WO 2022/178267	12/2021	WO	N/A
WO 2022/198068	12/2021	WO	N/A
WO 2022/221425	12/2021	WO	N/A
WO 2022/226057	12/2021	WO	N/A
WO 2022/236054	12/2021	WO	N/A
WO 2022/243303	12/2021	WO	N/A
WO 2022/226372	12/2021	WO	N/A
WO 2022/256503	12/2021	WO	N/A
WO 2022/271820	12/2021	WO	N/A
WO 2023/287765	12/2022	WO	N/A
WO 2023/018799	12/2022	WO	N/A
WO 2023/034489	12/2022	WO	N/A
WO 2023/044071	12/2022	WO	N/A
WO 2023/076345	12/2022	WO	N/A
WO 2023/086880	12/2022	WO	N/A
WO 2023/102118	12/2022	WO	N/A
WO 2023/122033	12/2022	WO	N/A
WO 2023/150098	12/2022	WO	N/A
WO 2023/150163	12/2022	WO	N/A
WO 2023/150171	12/2022	WO	N/A
WO 2023/215552	12/2022	WO	N/A
WO 2023/225519	12/2022	WO	N/A
WO 2023/229988	12/2022	WO	N/A
WO 2023/250077	12/2022	WO	N/A
WO 2024/015578	12/2023	WO	N/A
WO 2024/035844	12/2023	WO	N/A
WO 2024/081212	12/2023	WO	N/A
WO 2024/086167	12/2023	WO	N/A
WO 2024/086776	12/2023	WO	N/A
WO 2024/102809	12/2023	WO	N/A
WO 2024/137826	12/2023	WO	N/A
WO 2024/145224	12/2023	WO	N/A
WO 2024/145441	12/2023	WO	N/A
WO 2024/145445	12/2023	WO	N/A
WO 2024/145491	12/2023	WO	N/A
WO 2024/206603	12/2023	WO	N/A
WO 2024/220882	12/2023	WO	N/A
WO 2024/238900	12/2023	WO	N/A
WO 2024/254316	12/2023	WO	N/A
WO 2025/029605	12/2024	WO	N/A
WO 2025/029627	12/2024	WO	N/A
WO 2025/043076	12/2024	WO	N/A
WO 2025/072119	12/2024	WO	N/A

OTHER PUBLICATIONS

Borm et al., “High throughput Human embryo spatial transcriptome mapping by surface transfer of tissue RNA,” Abstracts Selected Talks, Single Cell Genom 2019, 1 pages (Abstract Only). cited by applicant

Chen et al., “Large field of view-spatially resolved transcriptomics at nanoscale resolution,” bioRxiv, Jan. 19, 2021, retrieved from URL <<https://www.biorxiv.org/node/1751045.abstract>>, 37 pages. cited by applicant

Cho et al., “Seq-Scope: Submicrometer-resolution spatial transcriptomics for single cell and subcellular studies,” bioRxiv, Jan. 27, 2021, retrieved from URL <<https://www.biorxiv.org/node/1754517.abstract>>, 50 pages. cited by applicant

Codeluppi et al., “Spatial organization of the somatosensory cortex revealed by osmFISH,” Nature Methods, Nov. 2018, 15:932-935. cited by applicant

Eng et al., "Microtranscriptome-scale super-resolved imaging in tissues using RNA seqFISH+," *Nature*, Apr. 2019, 568(7751):235-239, 37 pages. cited by applicant

Goh et al., "Highly Specific Multiplexed RNA Imaging in Tissues With Split-FISH," *Nat Methods*, Jun. 15, 2020, 17(7):689-693, 21 pages. cited by applicant

Liu et al., "High-Spatial-Resolution Multi-Omnics Sequencing via Deterministic Barcoding in Tissue," *Cell*, Nov. 13, 2020, 183(6):1665-1681, 36 pages. cited by applicant

Liu et al., "Spatial transcriptome sequencing of FFPE tissues at cellular level," *bioRxiv* 788992, Oct. 14, 2020, 39 pages. cited by applicant

Takei et al., "Integrated Spatial Genomics Reveals Global Architecture of Single Nuclei," *Nature*, Jan. 27, 2021, 590(7845):344-350, 53 pages. cited by applicant

Xia et al., "Spatial transcriptome profiling by MERFISH reveals subcellular RNA compartmentalization and cell cycle-dependent gene expression", *Proceedings of the National Academy of Sciences*, Sep. 2019, 116(39):19490-19499. cited by applicant

Burgess, "Spatial transcriptomics coming of age," *Nat Rev Genet.*, Jun. 2019, 20(6):317, 1 page. cited by applicant

Asp et al., "A spatiotemporal organ-wide gene expression and cell atlas of the developing human heart," *Cell*, Dec. 12, 2019, 179(7):1647-1660. cited by applicant

Kuhn et al., "A novel, high-performance random array platform for quantitative gene expression profiling," *Genome Res*, 2004, 14:2347-2356. cited by applicant

Ncbi.nlm.nih.gov, [online], "Molecular Inversion Probe Assay," available on or before Oct. 14, 2014, via Internet Archive: Wayback Machine URL<<https://web.archive.org/web/20141014124037/https://www.ncbi.nlm.nih.gov/probe/docs/techmip/>>, retrieved on Jun. 16, 2021, retrieved from URL<<https://www.ncbi.nlm.nih.gov/probe/docs/techmip/>>, 2 pages. cited by applicant

10xGenomics.com, [online], "Visium Spatial Gene Expression Reagent Kits—User Guide," Jan. 2022, retrieved on Jun. 27, 2024, retrieved from URL<<https://web.archive.org/web/20230326192142/https://www.10xgenomics.com/support/spatial-gene-expression-fresh-frozen/documentation/steps/library-spatial-gene-expression-reagent-kits-user-guide>>, 71 pages. cited by applicant

Dalma-Weiszhausz et al., "The affymetrix GeneChip platform: an overview," *Methods Enzymol.*, 2006, 410:3-28. cited by applicant

Miller et al., "Chapter 11—Solid and Suspension Microarrays for Microbial Diagnostics," *Methods in Microbiology*, 2015, 42:395-431. cited by applicant

Vickovic et al., "SM-Omics: An automated Platform for High-Throughput Spatial Multi-Omics," *bioRxiv*, Oct. 2020, 40 pages. cited by applicant

Holscher et al., "Application of Laser-Assisted Microdissection for Tissue and Cell-Specific Analysis of RNA," *Progress in Botany*, Jan. 2008, 69(3):141-167. cited by applicant

Sun et al., "Statistical Analysis of Spatial Expression Pattern for Spatially Resolved Transcriptomic Studies," *Nature Methods*, Jan. 27, 2020, 17(2): 193-200. cited by applicant

Svensson et al., "SpatialDE: identification of spatially variable genes," *Nature Methods*, May 2018, 15:343-346, 15 pages. cited by applicant

Asp et al., "Spatially Resolved Transcriptomes—Next Generation Tools for Tissue Exploration," *Bioessays*, Oct. 2020, 42(10):e1900221, 16 pages. cited by applicant

Bergenstr hle et al., "Seamless integration of image and molecular analysis for spatial transcriptomics workflows," *BMC Genomics*, Jul. 2020, 21(1):482, 7 pages. cited by applicant

Bolotin et al., "MiXCR: software for comprehensive adaptive immunity profiling," *Nat Methods.*, May 2015, 12(5):380-1. cited by applicant

Espina et al., "Laser-capture microdissection," *Nat Protoc*, 2006, 1(2):586-603. cited by applicant

[No Author Listed], "Chromium Next GEM Single Cell 3' Reagent Kits v3.1—User Guide," 10x Genomics, Document No. CG000204, Nov. 2019, 58 pages. cited by applicant

[No Author Listed], "Chromium Next GEM Single Cell 3' Reagent Kits v3.1 (Dual Index)—User Guide," 10x Genomics, Mar. 2021, Document No. CG000304, 58 pages. cited by applicant

[No Author Listed], "HuSNP Mapping Assay User's Manual," Affymetrix Part No. 90094 (Affymetrix, Santa Clara, Calif.), GeneChip, 2000, 104 pages. cited by applicant

[No Author Listed], "Microarray technologies have excellent possibilities in genomics-related researches," *Science Tools From Amersham Pharmacia Biotech* (English Translation). cited by applicant

10xGenomics.com, [online], "Visium Spatial Gene Expression Reagent Kits—Tissue Optimization—User Guide," Jul. 2020, retrieved on May 25, 2021, retrieved from URL<https://assets.ctfassets.net/an68im79xiti/5UJrN0cH17rEk0UXwdl9It/e54d99fb08a8f1500aba503005a04a56/CG000238_VisiumSpatialTissueOptimizationUserGuide.pdf>, 42 pages. cited by applicant

10xGenomics.com, [online], "Visium Spatial Gene Expression Reagent Kits—Tissue Optimization," Nov. 2019, retrieved on Jan. 25, 2022, retrieved from URL<https://assets.ctfassets.net/an68im79xiti/4q03w6959AJFxfS5lee9/6a2ac61cf6388a72564eeb96bc294967/CG000238_VisiumSpatialTissueOptimizationNov2019.pdf>, 46 pages. cited by applicant

10xGenomics.com, [online], "Visium Spatial Gene Expression Reagent Kits—Tissue Optimization," Oct. 2020, retrieved on Dec. 28, 2021, retrieved from URL<https://assets.ctfassets.net/an68im79xiti/5UJrN0cH17rEk0UXwdl9It/e54d99fb08a8f1500aba503005a04a56/CG000238_VisiumSpatialTissueOptimizationOct2020.pdf>, 43 pages. cited by applicant

10xGenomics.com, [online], "Visium Spatial Gene Expression Reagent Kits—User Guide," Jun. 2020, retrieved on May 25, 2021, retrieved from URL<https://assets.ctfassets.net/an68im79xiti/3GGIfH3RWpd1bFVhalpexR/8baa08d9007157592b65b2cdc7130990/CG000239_VisiumSpatialGeneExpressionUserGuide.pdf>, 69 pages. cited by applicant

10xGenomics.com, [online], "Visium Spatial Gene Expression Reagent Kits—User Guide," Oct. 2020, retrieved on Dec. 28, 2021, retrieved from URL<https://assets.ctfassets.net/an68im79xiti/3GGIfH3RWpd1bFVhalpexR/8baa08d9007157592665b2cdc7130990/CG000239_VisiumSpatialGeneExpressionOct2020.pdf>, 70 pages. cited by applicant

Adessi et al., "Solid phase DNA amplification: characterisation of primer attachment and amplification mechanisms," *Nucl. Acids Res.*, 2000, 28(20):E87, 8 pages. cited by applicant

Affymetrix, "GeneChip Human Genome U133 Set," retrieved from the Internet: on the World Wide Web at affymetrix.com/support/technical/datasheets/hgu133a.pdf, on Feb. 26, 2003, 2 pages. cited by applicant

Affymetrix, "Human Genome U95Av2," Internet Citation, retrieved from the internet: on the World Wide Web [affymetrix.com](http://affymetrix.com/support/technical/datasheets/hgu95a.pdf), retrieved on Oct. 2, 2002, 1 page. cited by applicant

Albretsen et al., "Applications of magnetic beads with covalently attached oligonucleotides in hybridization: Isolation and detection of specific measles virus RNA in cell lysate," *Anal. Biochem.*, 1990, 189(1):40-50. cited by applicant

Allawi et al., "Thermodynamics and NMR of Internal G T Mismatches in DNA," *Biochemistry*, 1996, 36(34):10581-10594. cited by applicant

Armani et al., "2D-PCR: a method of mapping DNA in tissue sections," *Lab Chip*, 2009, 9(24):3526-34. cited by applicant

Atkinson et al., "An Updated Protocol for High Throughput Plant Tissue Sectioning," *Front Plant Sci*, 2017, 8:1721, 8 pages. cited by applicant

Atkinson, "Overview of Translation: Lecture Manuscript," U of Texas, 2000, DD, pp. 6.1-6.8. cited by applicant

Bains et al., "A novel method for nucleic acid sequence determination," *Journal of Theoretical Biology*, 1988, 135(3), 303-7. cited by applicant

Barnes, "PCR amplification of up to 35-kb DNA with high fidelity and high yield from lambda bacteriophage templates," *Proc. Natl. Acad. Sci USA*, 1994, 91(12):3500-3504. cited by applicant

Beattie et al., "Advances in genosensor research," *Clin Chem.*, May 1995, 41(5):700-6. cited by applicant

Beechem et al., "High-Plex Spatially Resolved RNA and Protein Detection Using Digital Spatial Profiling: A Technology Designed for Immuno-oncology Biomedical Translational Research," *Methods Mol Biol*, 2020, Chapter 25, 2055:563-583. cited by applicant

Biosyntagma.com, [online], "Resolving Heterogeneity One Cell at a Time," available on or before Apr. 21, 2017, via Internet Archive: Wayback Machine URL<<https://web.archive.org/web/20170421212315/http://www.biosyntagma.com/>>, retrieved on Sep. 29, 2021, URL<<http://www.biosyntagma.com/>>, 3 pages. cited by applicant

Birney et al., "Identification and analysis of functional elements in 1% of the human genome by the ENCODE pilot project," *Nature*, 2007, 447(7146):799-816. cited by applicant

Blanchard et al., "High-density oligonucleotide arrays," *Biosensors & Bioelectronics*, 1996, 11(6-7):687-690. cited by applicant

Blokzijl et al., "Profiling protein expression and interactions: proximity ligation as a tool for personalized medicine," *J Intern. Med.*, 2010, 268(3):232-245. cited by applicant

Blow, "Tissue Issues," *Nature*, 2007, 448(7156):959-962. cited by applicant

Brandon et al., "Mitochondrial mutations in cancer," *Oncogene*, 2006, 25(34):4647-4662. cited by applicant

Brenner et al., "Gene expression analysis by massively parallel signature sequencing (MPSS) on microbead arrays," *Nat. Biotech.*, 2000, 18(6):630-634. cited by applicant

Brenner et al., "In vitro cloning of complex mixtures of DNA on microbeads: physical separation of differentially expressed cDNAs," *Proc. Natl. Acad. Sci. USA*, 1997, 94(14):7660-7665. cited by applicant

Brow, "35—The Cleavase I enzyme for mutation and polymorphism scanning," *PCR Applications Protocols for Functional Genomics*, 1999, pp. 537-550. cited by applicant

Brown et al., "Retroviral integration: structure of the initial covalent product and its precursor, and a role for the viral IN protein," *Proc Natl Acad Sci USA*, 1988, 85(12):4061-4066. cited by applicant

cited by applicant
 Buenrostro et al., "Transposition of native chromatin for multimodal regulatory analysis and personal epigenomics," *Nat Methods*, Dec. 2013, 10(12):1213-1218. cited by applicant
 Bullard et al., "Direct comparison of nick-joining activity of the nucleic acid ligases from bacteriophage T4," *Biochem. J.* 2006, 398(1):135-144. cited by applicant
 Burgess, "A space for transcriptomics," *Nature Reviews Genetics*, 2016, 17(8):436-7. cited by applicant
 Burgess, "Finding structure in gene expression," *Nature Reviews Genetics*, 2018, 19(5):249, 1 page. cited by applicant
 Ertsey et al., "Coverslip Mounted-Immersion Cycled in Situ RT-PCR for the Localization of mRNA in Tissue Sections," *Biotechniques*, 1998, 24(1):92-100. cited by applicant
 Cha et al., "Specificity, efficiency, and fidelity of PCR," *Genome Res.*, 1993, 3(3):S18-29. cited by applicant
 Chandra et al., "Cell-free synthesis-based protein microarrays and their applications," *Proteomics*, 2009, 5(6):717-30. cited by applicant
 Chatterjee et al., "Mitochondrial DNA mutations in human cancer. *Oncogene*," 2006, 25(34):4663-4674. cited by applicant
 Chen et al., "RNA imaging. Spatially resolved, highly multiplexed RNA profiling in single cells," *Science*, Apr. 2015, 348(6233):aaa6090, 21 pages. cited by applicant
 Chen et al., "Spatial Transcriptomics and In Situ Sequencing to Study Alzheimer's Disease," *Cell*, Aug. 2020, 182(4):976-991. cited by applicant
 Chen et al., "μCB-seq: microfluidic cell barcoding and sequencing for high-resolution imaging and sequencing of single cells," *Lab Chip*, Nov. 2020, 20(21):3955-3965. cited by applicant
 Constantine et al., "Use of genechip high-density oligonucleotide arrays for gene expression monitoring," *Life Science News*, Amersham Life Science, 1998, 1(1):1-4. cited by applicant
 Cornett et al., "MALDI imaging mass spectrometry: molecular snapshots of biochemical systems," *Nature Methods*, 2007, 4(10):828-833. cited by applicant
 Credle et al., "Multiplexed analysis of fixed tissue RNA using Ligation in situ Hybridization," *Nucleic Acids Research*, 2017, 45(14):e128, 9 pages. cited by applicant
 Crosetto et al., "Spatially resolved transcriptomics and beyond," *Nature Review Genetics*, 2015, 16(1):57-66. cited by applicant
 Czarnik, "Encoding methods for combinatorial chemistry," *Curr Opin Chem Biol.*, Jun. 1997, 1(1):60-6. cited by applicant
 Dabl et al., "Circle-to-circle amplification for precise and sensitive DNA analysis," *Proc. Natl. Acad. Sci.*, 2004, 101(13):4548-4553. cited by applicant
 Daubendiek et al., "Rolling-Circle RNA Synthesis: Circular Oligonucleotides as Efficient Substrates for T7 RNA Polymerase," *J. Am. Chem. Soc.*, 1995, 117(1):1-5. cited by applicant
 Davies et al., "How best to identify chromosomal interactions: a comparison of approaches," *Nat. Methods*, 2017, 14(2):125-134. cited by applicant
 Dean et al., "Comprehensive human genome amplification using multiple displacement amplification," *Proc Natl. Acad. Sci. USA*, 2002, 99(8):5261-66. cited by applicant
 Duncan et al., "Affinity chromatography of a sequence-specific DNA binding protein using Teflon-linked oligonucleotides," *Anal. Biochem.*, 1988, 169(1):10-14. cited by applicant
 Eguiluz et al., "Multitissue array review: a chronological description of tissue array techniques, applications and procedures," *Pathology Research and Practice*, 2004, 200(1):1-6. cited by applicant
 Eldridge et al., "An in vitro selection strategy for conferring protease resistance to ligand binding peptides," *Protein Eng Des Sel.*, 2009, 22(11):691-698. cited by applicant
 Ellington et al., "Antibody-based protein multiplex platforms: technical and operational challenges," *Clin Chem*, 2010, 56(2):186-193. cited by applicant
 Emmert-Buck et al., "Laser capture microdissection," *Science*, Nov. 1996, 274(5289):998-1001. cited by applicant
 Fire et al., "Rolling replication of short DNA circles," *Proc. Natl. Acad. Sci.*, 1995, 92(10):4641-4645. cited by applicant
 Fluidigm, "Equivalence of Imaging Mass Cytometry and Immunofluorescence on FFPE Tissue Sections," *White Paper*, 2017, 12 pages. cited by applicant
 Fodor et al., "Light-directed, spatially addressable parallel chemical synthesis," *Science*, 1995, 251(4995):767-773. cited by applicant
 Forster et al., "A human gut bacterial genome and culture collection for improved metagenomic analyses," *Nature Biotechnology*, 2019, 37(2):186-192. cited by applicant
 Frese et al., "Formylglycine aldehyde Tag—protein engineering through a novel post-translational modification," *ChemBioChem.*, 2009, 10(3):425-27. cited by applicant
 Fu et al., "Counting individual DNA molecules by the stochastic attachment of diverse labels," *PNAS*, 2011, 108(22):9026-9031. cited by applicant
 Fu et al., "Continuous Polony Gels for Tissue Mapping with High Resolution and RNA Capture Efficiency," *bioRxiv*, 2021, 20 pages. cited by applicant
 Fullwood et al., "Next-generation DNA sequencing of paired-end tags (PET) for transcriptome and genome analyses," *Genome Res.*, 2009, 19(4):521-532. cited by applicant
 Ganguli et al., "Pixelated spatial gene expression analysis from tissue," *Nat Commun.*, Jan. 2018, 9(1):202, 9 pages. cited by applicant
 Gao et al., "Q&A: Expansion microscopy," *BMC Biology*, 15:50, 9 pages, 2017. cited by applicant
 Gene@arrays[online], BeadArray Technology, available on or before Feb. 14, 2015, via Internet Archive: Wayback Machine URL <https://web.archive.org/web/20150214084616/http://genearrays.com/services/microarrays/illumina/beadarray-technology/>, [retrieved on Jan. 30, 2020], 3 pages. cited by applicant
 Giacomello et al., "Spatially resolved transcriptome profiling in model plant species," *Nature Plants* 3, 17061, 11 pages, 2017. cited by applicant
 Gnanaprasagam, "Unlocking the molecular archive: the emerging use of formalin-fixed paraffin-embedded tissue for biomarker research in urological cancer," *PLoS One*, 2015, 10(2):e0117478, 10 pages. cited by applicant
 Goldkorn et al., "A simple and efficient enzymatic method for covalent attachment of DNA to cellulose. Application for hybridization-restriction analysis and DNA probes," *Nucleic Acids Res.*, 1986, 14(22):9171-9191. cited by applicant
 Gracia Villacampa et al., "Genome-wide Spatial Expression Profiling in FFPE Tissues," *bioRxiv*, 2020, pp. 38 pages. cited by applicant
 Gunderson et al., "Decoding randomly ordered DNA arrays," *Genome Research*, 2004, 14(5):870-877. cited by applicant
 Guo et al., "Direct fluorescence analysis of genetic polymorphisms by hybridization with oligonucleotide arrays on glass supports," *Nucleic Acids Res.*, Dec. 2000, 28(24):E125-E128. cited by applicant
 Gupta et al., "Single-cell isoform RNA sequencing characterizes isoforms in thousands of cerebellar cells," *Nature Biotechnol.*, Oct. 2018, 36:1197-1202. cited by applicant
 Hadley et al., "Determining composition of micron-scale protein deposits in neurodegenerative disease by spatially targeted optical microproteomics," *Elife*, 2019, 8:e35555, 10 pages. cited by applicant
 Hamaguchi et al., "Direct reverse transcription-PCR on oligo(dT)-immobilized polypropylene microplates after capturing total mRNA from crude cell lysates," *Anal. Biochem.*, 2004, 344(1):2256-63. cited by applicant
 Hayes et al., "Electrophoresis of proteins and nucleic acids: I-Theory," *BMJ*, Sep. 1989, 299(6703):843-6. cited by applicant
 He et al., "In situ synthesis of protein arrays," *Current Opinion in Biotechnology*, 2008, 19(1):4-9. cited by applicant
 He, "Cell-free protein synthesis: applications in proteomics and biotechnology," *New Biotechnology*, 2008, 25(2-3):126-132. cited by applicant
 Hejatko et al., "In situ hybridization technique for mRNA detection in whole mount *Arabidopsis* samples," *Nature Protocols*, 2006, 1(4):1939-1946. cited by applicant
 Hiatt et al., "Parallel, tag-directed assembly of locally derived short sequence reads," *Nature Methods*, 2010, 7(2):119-25. cited by applicant
 Inoue and Wittbrodt, "One for All—A Highly Efficient and Versatile Method for Fluorescent Immunostaining in Fish Embryos," *PLoS One* 6, e19713, 2011. cited by applicant
 Jamur et al., "Permeabilization of cell membranes," *Method Mol. Biol.*, 2010, 588:63-66. cited by applicant
 Jemt et al., "An automated approach to prepare tissue-derived spatially barcoded RNA-sequencing libraries," *Scientific Reports*, 2016, 6:37137, 10 pages. cited by applicant
 Kapteyn et al., "Incorporation of non-natural nucleotides into template-switching oligonucleotides reduces background and improves cDNA synthesis from vesicles," *BMC Genomics*, Jul. 2010, 11:413, 9 pages. cited by applicant
 Kashyap et al., "Selective local lysis and sampling of live cells for nucleic acid analysis using a microfluidic probe," *Sci Rep.*, Jul. 2016, 6:29579, 10 pages. cited by applicant
 Korbel et al., "Paired-end mapping reveals extensive structural variation in the human genome," *Science*, 2007, 318(5849):420-426. cited by applicant
 Kozlov et al., "A highly scalable peptide-based assay system for proteomics," *PLoS One*, 2012, 7(6):e37441, 10 pages. cited by applicant
 Kristensen et al., "High-Throughput Methods for Detection of Genetic Variation," *Biotechniques*, Feb. 2001, 30(2):318-332. cited by applicant
 Kurz et al., "cDNA-protein fusions: covalent protein-gene conjugates for the in vitro selection of peptides and proteins," *ChemBioChem.*, 2001, 2(9):666-72. cited by applicant
 Kwok, "High-throughput genotyping assay approaches," *Pharmacogenomics*, Feb. 2000, 1(1):95-100. cited by applicant
 Lage et al., "Whole genome analysis of genetic alterations in small DNA samples using hyperbranched strand displacement amplification and array-CGH," *Genomics*, 2000, 13(2):294-307. cited by applicant
 Landegren et al., "Reading bits of genetic information: methods for single-nucleotide polymorphism analysis," *Genome Res.*, Aug. 1998, 8(8):769-76. cited by applicant

Langdale et al., "A rapid method of gene detection using DNA bound to Sephacryl," *Gene*, 1985, 36(3):201-210. cited by applicant

Lassmann et al., "A Novel Approach for Reliable Microarray Analysis of Microdissected Tumor Cells From Formalin-Fixed and Paraffin-Embedded Colorectal Specimens," *J Mol Med*, 87, 211-224, 2009. cited by applicant

Lee et al., "Fluorescent in situ sequencing (FISSEQ) of RNA for gene expression profiling in intact cells and tissues," *Nature Protocols*, 2015, 10(3):442-458.

Lein et al., "The promise of spatial transcriptomics for neuroscience in the era of molecular cell typing," *Science* 358, 64-69, 2017. cited by applicant

Leriche et al., "Cleavable linkers in chemical biology," *Bioorganic & Medicinal Chemistry*, 2012, 20:571-582. cited by applicant

Linnarsson, "Recent advances in DNA sequencing methods—general principles of sample preparation," *Experimental Cell Research*, 2010, 316(8):1339-1343.

Liu et al., "High-Spatial-Resolution Multi-Omics Atlas Sequencing of Mouse Embryos via Deterministic Barcoding in Tissue," *BioRxiv*, 2019, 55 pages. cited by applicant

Lizardi et al., "Mutation detection and single-molecule counting using isothermal rolling-circle amplification," *Nat. Genet.*, 1998, 19(3):225-232. cited by applicant

Lundberg et al., "Multiplexed homogeneous proximity ligation assays for high-throughput protein biomarker research in serological material," *Mol Cell Proteomics*, 2004, 3(12):M110.004978, 11 pages. cited by applicant

MacBeath et al., "Printing proteins as microarrays for high-throughput function determination," *Science*, Sep. 2000, 289(5485):1760-1763. cited by applicant

Marx, "Method of the Year: spatially resolved transcriptomics," *Nature Methods*, 2021, 18(1):9-14. cited by applicant

Merritt et al., "Multiplex digital spatial profiling of proteins and RNA in fixed tissue," *Nat Biotechnol*, May 2020, 38(5):586-599. cited by applicant

Metzker, "Sequencing technologies—the next generation," *Nature Reviews Genetics*, 2010, 11(1):31-46. cited by applicant

Miller et al., "Basic concepts of microarrays and potential applications in clinical microbiology," *Clinical Microbiology Reviews*, 2009, 22(4):611-633. cited by applicant

Mishra et al., "Three-dimensional genome architecture and emerging technologies: looping in disease," *Genome Medicine*, 2017, 9(1):87, 14 pages. cited by applicant

Mitra et al., "Digital genotyping and haplotyping with polymerase colonies," *Proc. Natl. Acad. Sci. USA*, May 2003, 100(10):5926-5931. cited by applicant

Mizusawa et al., "A bacteriophage lambda vector for cloning with BamHI and Sau3A," *Gene*, 1982, 20(3):317-322. cited by applicant

Nanostring, "GeoMx—nCounter DSP Instrument User Manual," MAN-10088-05 for software v2.1, Nov. 2020, 106 pages. cited by applicant

Nanostring, "GeoMx—NGS DSP Instrument User Manual," MAN-10116-03 for software v2.1, Dec. 2020, 104 pages. cited by applicant

Nikiforov et al., "The use of 96-well polystyrene plates for DNA hybridization-based assays: an evaluation of different approaches to oligonucleotide immobilization," *J. Microencapsul.*, May 1995, 22(1):201-9. cited by applicant

Nowak et al., "Entering the Postgenome Era," *Science*, 1995, 270(5235):368-71. cited by applicant

Oliveira et al., "Inverse electron demand Diels-Alder reactions in chemical biology," *Chemical Society Reviews*, Aug. 2017, 46(16):4895-4950. cited by applicant

Pemov et al., "DNA analysis with multiplex microarray-enhanced PCR," *Nucl. Acids Res.*, Jan. 2005, 33(2):e11, 9 pages. cited by applicant

Perler et al., "Intervening sequences in an Archaea DNA polymerase gen," *Proc Natl Acad Sci USA*, Jun. 1992, 89(12):5577-5581. cited by applicant

Petterson et al., "Generations of sequencing technologies," *Genomics*, 2009, 93(2):105-111. cited by applicant

Polsky-Cynkin et al., "Use of DNA immobilized on plastic and agarose supports to detect DNA by sandwich hybridization," *Clin. Chem.*, 1985, 31(9):1438-1441.

Ranki et al., "Sandwich hybridization as a convenient method for the detection of nucleic acids in crude samples," *Gene*, 1983, 21(1-2):77-85. cited by applicant

Reinartz et al., "Massively parallel signature sequencing (MPSS) as a tool for in-depth quantitative gene expression profiling in all organisms," *Brief Funct Genomics*, 2002, 1(1):95-104. cited by applicant

Rodrigues et al., "Slide-seq: A scalable technology for measuring genome-wide expression at high spatial resolution," *Science*, 2019, 363(6434):1463-1467. cited by applicant

Ronaghi et al., "A sequencing method based on real-time pyrophosphate," *Science*, Jul. 1998, 281(5375):363-365. cited by applicant

Ronaghi et al., "Real-time DNA sequencing using detection of pyrophosphate release," *Analytical Biochemistry*, Nov. 1996, 242(1):84-89. cited by applicant

Ronaghi, "Pyrosequencing sheds light on DNA sequencing," *Genome Res*, Jan. 2001, 11(1):3-11. cited by applicant

Salmén et al., "Barcoded solid-phase RNA capture for Spatial Transcriptomics profiling in mammalian tissue sections," *Nature Protocols*, Oct. 2018, 13(11):2018-2027. cited by applicant

Saxonov et al., "10x Genomics, Mastering Biology to Advance Human Health," PowerPoint, 10x, 2020, 41 pages. cited by applicant

Schena et al., "Quantitative monitoring of gene expression patterns with a complementary DNA microarray," *Science*, Oct. 1995, 270(5235):467-470. cited by applicant

Shalon et al., "A DNA microarray system for analyzing complex DNA samples using two-color fluorescent probe hybridization," *Genome Res.*, Jul. 1996, 6(7):1215-1222. cited by applicant

Stahl et al., "Visualization and analysis of gene expression in tissue sections by spatial transcriptomics," *Science*, Jun. 2016, 353(6294):78-82. cited by applicant

Stahl et al., "Visualization and analysis of gene expression in tissue sections by spatial transcriptomics," *Supplementary Materials, Science*, Jul. 2016, 353(6294):78-82. cited by applicant

Stimpson et al., "Real-time detection of DNA hybridization and melting on oligonucleotide arrays by using optical wave guides," *Proc Natl Acad Sci USA*, Jun. 2000, 97(12):6581-6586. cited by applicant

Strell et al., "Placing RNA in context and space—methods for spatially resolved transcriptomics," *The FEBS Journal*, 2019, 286(8):1468-1481. cited by applicant

Thiery et al., "Multiplex target protein imaging in tissue sections by mass spectrometry—TAMSIM," *Rapid Commun. Mass Spectrom.*, 2007, 21:823-829. cited by applicant

Tijssen et al., "Overview of principles of hybridization and the strategy of nucleic acid assays" in *Techniques in Biochemistry and Molecular Biology—Hybridization*, 1993, 24(Chapter 2), 65 pages. cited by applicant

Trejo et al., "Extraction-free whole transcriptome gene expression analysis of FFPE sections and histology-directed subareas of tissue," *PLoS One*, Feb. 2019, 14(2):e0199999. cited by applicant

Twyman et al., "Techniques Patents for SNP Genotyping," *Pharmacogenomics*, Jan. 2003, 4(1):67-79. cited by applicant

Van Gelder et al., "Amplified RNA synthesized from limited quantities of heterogeneous cDNA," *Proc. Natl. Acad. Sci. USA*, 1990, 87(5):1663-1667. cited by applicant

Vasiliskov et al., "Fabrication of microarray of gel-immobilized compounds on a chip by copolymerization," *Biotechniques*, Sep. 1999, 27(3):592-606. cited by applicant

Vickovic et al., "High-definition spatial transcriptomics for in situ tissue profiling," *Nat Methods*, Oct. 2019, 16(10):987-990. cited by applicant

Vogelstein et al., "Digital PCR," *Proceedings of the National Academy of Sciences*, Aug. 1999, 96(16):9236-9241. cited by applicant

Walker et al., "Strand displacement amplification—an isothermal, in vitro DNA amplification technique," *Nucleic Acids Research*, 1992, 20(7):1691-1696. cited by applicant

Wang et al., "Imaging-based pooled CRISPR screening reveals regulators of lncRNA localization," *Proc Natl Acad Sci USA*, May 2019, 116(22):10842-10850. cited by applicant

Wang et al., "High-fidelity mRNA amplification for gene profiling," *Nature Biotechnology*, Apr. 2000, 18(4):457-459. cited by applicant

Worthington et al., "Cloning of random oligonucleotides to create single-insert plasmid libraries," *Anal Biochem*, 2001, 294(2):169-175. cited by applicant

Yershov et al., "DNA analysis and diagnostics on oligonucleotide microchips," *Proc. Natl. Acad. Sci. USA*, May 1996, 93(10):4913-4918. cited by applicant

Yoda et al., "Site-specific gene expression analysis using an automated tissue micro-dissection punching system," *Sci Rep.*, Jun. 2017, 7(1):4325, 11 pages. cited by applicant

Zhu et al., "Reverse transcriptase template switching: a SMART approach for full-length cDNA library construction," *Biotechniques*, Apr. 2001, 30(4):892-899. cited by applicant

Zlobec et al., "Next-generation tissue microarray (ngTMA) increases the quality of biomarker studies: an example using CD3, CD8, and CD45RO in the tumor microenvironment of different solid tumor types," *Journal of Translational Medicine*, 2013 11(104):1-7. cited by applicant

Primary Examiner: Priest; Aaron A

Assistant Examiner: Beil; Randi Lynn

Attorney, Agent or Firm: Fish & Richardson P.C.

Background/Summary

CROSS-REFERENCE TO RELATED APPLICATION (1) This application claims the benefit of U.S. Patent Application Ser. No. 62/976,783 titled SPATIAL TARGETING OF ANALYTES, filed Feb. 14, 2020, the disclosure of which is hereby incorporated by reference in its entirety.

BACKGROUND

(1) Cells within a tissue have differences in cell morphology and/or function due to varied analyte levels (e.g., gene and/or protein expression) within the different cells. The specific position of a cell within a tissue (e.g., the cell's position relative to neighboring cells or the cell's position relative to the tissue microenvironment) can affect, e.g., the cell's morphology, differentiation, fate, viability, proliferation, behavior, signaling, and cross-talk with other cells in the tissue.

(2) Spatial heterogeneity has been previously studied using techniques that typically provide data for a handful of analytes in the context of intact tissue or a portion of a tissue (e.g., tissue section), or provide significant analyte data from individual, single cells, but fails to provide information regarding the position of the single cells from the originating biological sample (e.g., tissue).

(3) Users may be interested in focusing on particular regions of interest of a biological sample. Techniques are needed to identify a region of interest and collect analytes from the region of interest in an efficient and reliable manner.

SUMMARY

(4) This document generally relates to apparatuses, systems, and methods for preparing a sample, such as a biological sample. In particular, the present disclosure provides systems and methods for spatially targeting one or more analytes in a region of interest, and efficiently obtaining the analytes from a biological sample.

(5) Some implementations of the present disclosure provide systems and methods for removing one or more features (e.g., beads) that contain analytes of interest from a substrate by emitting light toward the features on the substrate. In some implementations, a feature capture device is provided. The feature capture device can include a capture surface configured to face at least the features of interest on the substrate. The capture surface can be activated to attach (e.g., adhesively attach) the features of interest when a beam of light is projected on an area of the capture surface that corresponds to the features of interest on the substrate. For example, the feature capture device can include a transparent body having a first surface and a second surface that, for example, may be arranged opposite to the first surface. A light beam can enter the first surface of the feature capture device, pass through the transparent body, and reach the second surface that includes the capture surface. In some implementations, a light source can emit a beam of light with a predetermined beam divergence that is suitable for a particular size of region of interest. For example, light can be a laser of a particular wavelength configured to activate a limited region on the substrate.

(6) Some implementations of the present disclosure can include an instrument for identifying a region of interest on an array of features affixed to the substrate. Examples of such an instrument include a microscope, a camera, or other suitable imaging devices configured to visualize the array of features on the substrate. A region of interest in the array of features can be manually determined by a user who examines the array through the instrument. Alternatively or in addition, the instrument can be configured to automatically identify a region of interest in the array of features based on one or more inputs or factors, such as particular types of analytes of interest (e.g., cell type, RNA, Protein, DNA, etc.), particular research purposes, etc.

(7) Once the feature capture device has captured the features of interest, it can be removed from the position where it captured the features from the substrate, and processed to retrieve the captured features from the capture surface of the feature capture device. Various methods can be used to retrieve the captured features from the feature capture device.

(8) In alternative implementations, the present disclosure provides a substrate that attaches an array of features with photo-sensitive bonds, such as cleavable linkers. A region of interest can be identified on the feature array using, for example, the instrument described herein, and a beam of light is projected onto the region of interest so that the photo-sensitive bonds affixing the features in the region of interest are cleaved, and the features in the region of interest are released from the substrate. For example, a light beam can be projected onto a bottom surface of the substrate that is opposite to a top surface on which the feature array is affixed with the photo-sensitive bonds. The light beam passes through the transparent body of the substrate and reaches the top surface to cleave the photo-sensitive bonds. In some implementations, a light source emits a beam of light with a predetermined beam divergence that is suitable for a particular size of region of interest. For example, light can be a laser of a particular wavelength (e.g., ultraviolet (UV) light) configured to be projected on a limited region on the substrate. In some implementations, the removed features can be collected within a reservoir containing a suitable liquid.

(9) Particular embodiments described herein include a system for preparing a biological sample. The system includes a substrate, a feature capture block, a light generator, and a controller. The substrate is configured to place a feature array thereon. The array of features may be configured to capture analytes from the biological sample. The feature capture block may include a capture surface configured to face the feature array on the substrate. The light generator may be configured to emit light. The controller may be configured to activate the light generator to emit light toward the capture surface of the feature capture block to permit the capture surface to capture at least one feature from the feature array.

(10) In some implementations, the system can optionally include one or more of the following features. The feature capture block may include a transparent body to permit light to pass through the body. The transparent body of the feature capture block may have a light entering surface onto which the light is emitted. The light may pass through the transparent body and be directed toward the capture surface. The light entering surface may be arranged to be opposite to the capture surface. The capture surface may include a feature capture material. The feature capture material may include a light-activated or heat-activated adhesive. The light-activated adhesive may include a polymer which can be triggered to be adhesive under selected conditions. The feature capture block may be configured to be arranged such that the capture surface contacts with the feature array on the substrate. The feature capture block may be configured to be arranged such that the capture surface is spaced apart from the feature array on the substrate. The light generator may be configured to generate a laser of a predetermined wavelength. The laser may include an ultraviolet (UV) laser. Other light sources can also be used, such as a Digital Micromirror Device (DMD). The system may include a microscope configured to visualize a region of interest on the feature array. The controller may be configured to activate the light generator to emit the light toward a portion of the capture surface corresponding to the region of interest on the feature array. The light can be applied to the portion of the capture surface directly from the light generator. Alternatively, the light can be deflected from the light generator and then interact with the portion of the capture surface. The light-emitted portion of the capture surface may be permitted to capture at least one feature in the region of interest on the feature array. The feature array may include a spatially-barcoded bead array. The feature array may include a plurality of hydrogel beads. Alternatively or in addition, the feature array may include a plurality of silica beads, which may have an average diameter of 50 microns or smaller (e.g., nanometer scale). The feature array may include a plurality of beads being uniform in size (e.g., diameter). The feature array may be affixed to the substrate in a monolayer.

(11) Particular embodiments described herein include a method of preparing a biological sample. The method may include providing a substrate affixing a feature array thereon, the feature array capturing analytes from the biological sample; arranging a feature capture block relative to the substrate such that a capture surface of the feature capture block faces the feature array on the substrate; identifying a region of interest on the feature array; emitting light (either directly or after deflected) toward a portion of the capture surface of the feature capture block, the portion of the capture surface corresponding to the region of interest on the feature array; permitting for the light-emitted portion of the capture surface to capture at least one feature within the region of interest on the feature array; removing the feature capture block from the substrate; and retrieving the at least one feature from the capture surface of the feature capture block.

(12) In some implementations, the method can optionally include one or more of the following attributes. The method may include placing the substrate on a substrate stage of a microscope to visualize the feature array on the substrate including the region of interest. The feature capture block may include a transparent body to permit the light to pass through the body. The transparent body of the feature capture block may have a light entering surface onto which the light is emitted. The light may pass through the transparent body and be directed toward the capture surface. The light entering surface may be arranged to be opposite to the capture surface. The capture surface may include a feature capture material. The feature capture material may include a light- or heat-activated adhesive. The light-activated adhesive may include a polymer which can be triggered to be adhesive under selected conditions. The feature capture block may be configured to be arranged such that the capture surface contacts with the feature array on the substrate. The feature capture block may be configured to be arranged such that the capture surface is spaced apart from the feature array on the substrate. The light may include a laser of a predetermined wavelength. The light may include an ultraviolet (UV) laser. Other light sources can also be used, such as a Digital Micromirror Device (DMD). The feature array may include a spatially-barcoded bead array. The feature array may include a plurality of beads being uniform in size (e.g., diameter) and arranged in a monolayer.

(13) Particular embodiments described herein include a system for preparing a biological sample. The system may include a substrate, a light generator, and a controller. The substrate may be configured to include photo-sensitive bonds and a feature array attached to the substrate via the photo-sensitive bonds. The feature array may be configured to capture analytes from the biological sample. The light generator may be configured to emit light. The controller may be configured to activate the light generator to emit the light toward a region of interest of the feature array to cleave the photo-sensitive bonds in the region of interest and permit for at least one feature within the region of interest of the feature array to be released from the substrate.

(14) In some implementations, the system can optionally include one or more of the following features. The substrate may include a transparent body having a first surface and a second surface. The first surface may be configured to include the photo-sensitive bonds and the feature array. The light generator may be configured to emit the light onto the second surface of the substrate such that the light passes through the transparent body and reaches the photo-sensitive bonds on the first surface of the substrate. The first surface may be arranged to be opposite to the second surface. The light generator may be configured to generate a laser of a predetermined wavelength. The laser may include an ultraviolet (UV) laser. Other light sources can also be used, such as a Digital Micromirror Device (DMD). The system may include a reservoir containing a fluid into which the at least one feature is released from the substrate. The photo-sensitive bonds may include photo-cleavable linkers. The photo-cleavable linkers may include at least one of photo-sensitive chemical bonds include-amino-3-(2-nitrophenyl) propionic acid (ANP), phenacyl ester derivatives, 8-quinoliny benzenesulfonate, dicoumarin, 6-bromo-7-alkixycoumarin-4-ylmethoxycarbonyl, a bimane-based linker, or a bis-arylhydrazone based linker. The system may include a microscope configured to visualize a region of interest on the feature array. The controller may be configured to activate the light generator to emit the light toward a portion of the substrate corresponding to the region of interest on the feature array. The feature array may include a spatially-barcoded bead array. The feature array may include a plurality of hydrogel beads. The feature array may include a plurality of beads being uniform in size (e.g., diameter). The feature array may be affixed to the substrate in a monolayer.

(15) Particular embodiments described herein include a method of preparing a biological sample. The method may include providing a substrate affixing a feature array using photo-sensitive bonds; identifying a region of interest on the feature array; emitting light toward a portion of the substrate, the portion of the capture surface corresponding to the region of interest on the feature array; permitting for the photo-sensitive bonds in the region of interest to be cleaved so that at least one feature within the region of interest is released from the substrate; and collecting the at least one feature.

(16) In some implementations, the method can optionally include one or more of the following features. The method may include placing the substrate on a substrate stage of a microscope to visualize the feature array on the substrate including the region of interest. The substrate may include a transparent body having a first surface and a second surface. The first surface configured to place the photo-sensitive bonds and the feature array. Emitting light may include emitting light onto second surface of the substrate such that the light passes through the transparent body and reaches the photo-sensitive bonds on the first surface of the substrate. The first surface may be arranged to be opposite to the second surface. The light may include a laser of a predetermined wavelength. The light may include an ultraviolet (UV) laser. Other light sources can also be used, such as a Digital Micromirror Device (DMD). Collecting the at least one feature may include collecting the at least one feature within a fluid contained in a reservoir. The photo-sensitive bonds may include photo-cleavable linkers. The photo-cleavable linkers may include at least one of photo-sensitive chemical bonds include 3-amino-3-(2-nitrophenyl) propionic acid (ANP), phenacyl ester derivatives, 8-quinoliny benzenesulfonate, dicoumarin, 6-bromo-7-alkixycoumarin-4-ylmethoxycarbonyl, a bimane-based linker, or a bis-arylhydrazone based linker. The feature array may include a spatially-barcoded bead array. The feature array may include a plurality of hydrogel beads. The feature array may include a plurality of beads being uniform in size (e.g., diameter). The feature array may be affixed to the substrate in a monolayer.

(17) As such, the present disclosure provides solutions for efficiently performing spatially resolved capture of analytes by physically dissecting a region of interest from captured analytes on the feature array.

(18) All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, patent application, or item of information was specifically and individually indicated to be incorporated by reference. To the extent publications, patents, patent applications, and items of information incorporated by reference contradict the disclosure contained in the specification, the specification is intended to supersede and/or take precedence over any such contradictory material.

(19) Where values are described in terms of ranges, it should be understood that the disclosure includes the disclosure of all possible sub-ranges within such ranges, as well as specific numerical values that fall within such ranges irrespective of whether a specific numerical value or specific sub-range is expressly stated.

(20) The term "each," when used in reference to a collection of items, is intended to identify an individual item in the collection but does not necessarily refer to every item in the collection, unless expressly stated otherwise, or unless the context of the usage clearly indicates otherwise.

(21) Various embodiments of the features of this disclosure are described herein. However, it should be understood that such embodiments are provided merely by way of example, and numerous variations, changes, and substitutions can occur to those skilled in the art without departing from the scope of this disclosure. It should also be understood that various alternatives to the specific embodiments described herein are also within the scope of this disclosure.

Description

DESCRIPTION OF DRAWINGS

(1) The following drawings illustrate certain embodiments of the features and advantages of this disclosure. These embodiments are not intended to limit the scope of the appended claims in any manner. Like reference symbols in the drawings indicate like elements.

(2) FIG. 1 illustrates an exemplary system for preparing a biological sample.

(3) FIGS. 2A-C illustrate an exemplary process for performing spatially-targeted feature capture.

(4) FIG. 3 is a flowchart of an exemplary process for performing spatially-targeted feature capture.

(5) FIG. 4 illustrates another exemplary system for preparing a biological sample.

(6) FIGS. 5A-C illustrate an exemplary process for performing spatially-targeted feature capture.

(7) FIG. 6 is a flowchart of an exemplary process for performing spatially-targeted feature capture.

(8) FIG. 7 is a block diagram of computing devices that may be used to implement the systems and methods described in this document, as either a

client or as a server or plurality of servers.

DETAILED DESCRIPTION

(9) In general, the present disclosure relates to devices, instruments, systems, and methods for preparing a biological sample. In particular, the present disclosure provide systems and methods for spatially targeting one or more analytes in a region of interest, and efficiently removing the analytes from a biological sample. Some implementations of the present disclosure provide systems and methods for removing one or more beads that contain analytes of interest or intermediate agents representative of analytes of interest (e.g., an analyte capture agent or a ligation product) from a substrate by emitting light toward the beads on the substrate.

(10) In some implementations, the present disclosure provides a bead capture block that includes a capture surface configured to face at least the beads of interest on the substrate. The capture surface can be activated to adhere to the beads of interest when a beam of light is projected on an area of the capture surface that corresponds to the beads of interest on the substrate. For example, the bead capture block can include a transparent body having a first surface and a second surface that, for example, may be arranged opposite to the first surface. A light beam can enter the first surface of the bead capture block, pass through the transparent body, and reach the second surface that includes the capture surface. Various types of light sources can be selected depending on various factors, such as the characteristics of the capture surface. In some implementations, a light source is configured to emit a beam of light with a predetermined beam divergence that is suitable for a particular size of region of interest. For example, light can be a laser of a particular wavelength configured to activate a limited region on the substrate. Examples of the wave length include 250-350 nm or 350-460 nm UV light.

(11) Some implementations of the present disclosure can include an instrument for identifying a region of interest on an array of beads affixed to the substrate. Examples of such an instrument include a microscope, a camera, or other suitable imaging devices configured to visualize the array of beads on the substrate. A region of interest in the bead array can be manually determined by a user who examines the bead array through the instrument. Alternatively or in addition, the instrument is configured to automatically identify a region of interest in the bead array based on one or more inputs or factors, such as particular types of analytes of interest (e.g., cell type such as immune cells or cancer cells), particular research purposes (e.g., diagnosis), etc.

(12) Once the bead capture block has captured the beads of interest, it can be removed from the position that it captured the beads from the substrate, and processed to retrieve the captured beads from the capture surface of the bead capture block. Various methods can be used to retrieve the captured beads from the bead capture block.

(13) In alternative implementations, the present disclosure provides a substrate that attaches an array of beads with photo-sensitive bonds, such as cleavable linkers. A region of interest can be identified on the bead array using, for example, the instrument described herein, and a beam of light is projected onto the region of interest so that the photo-sensitive bonds affixing the beads in the region of interest are cleaved, and the beads in the region of interest are released from the substrate. For example, a light beam can be projected onto a bottom surface of the substrate that is opposite to a top surface on which the bead array is affixed with the photo-sensitive bonds. The light beam passes through transparent body of the substrate and reaches the top surface to cleave the photo-sensitive bonds. Various types of light sources can be selected based on various factors, such as the characteristics of the substrate and the photo-sensitive bonds being used. In some implementations, a light source is configured to emit a beam of light with a predetermined beam divergence that is suitable for a particular size of region of interest. For example, light can be a laser of a particular wavelength (e.g., ultraviolet (UV) light) configured to activate a limited region on the substrate. In some implementations, the removed beads can be collected within a reservoir (e.g., containing a suitable liquid). In some implementations, the released beads can be attached to an additional substrate (e.g., a flow cell, a wafer, a bead, a slide) using a covalent bond, such as through photo-click chemistry. For example, the released beads and the additional substrate can include complementary reactive moieties (e.g., a terminal alkene and a tetrazine or a tetrazole) that can perform a cycloaddition reaction upon irradiation with an appropriate wavelength of light, thereby covalently linking the released beads to the additional substrate. Additional examples of photo-click chemistry are described in Oliveira, et al. *Chemical Society Reviews* 46.16 (2017): 4895-4950, incorporated herein by reference.

(14) The techniques described herein allow a user to obtain analytes in a targeted area for analysis before and after a biological sample is thoroughly analyzed. Further, the substrate that captures an array of features that capture analytes can be used for analysis of a particular region of interest, and can be used again for analysis of another region of interest simultaneously or at a later time. In some implementations, the feature array can include spatially-barcoded capture probes so that the analytes included in the target features removed from a targeted area can be accurately identified in a spatial analysis, as described herein.

(15) Referring to FIGS. 1-3, an example system for preparing a biological sample is described. In some implementations, the system is configured to obtain one or more analytes that are spatially targeted. For example, the system can be configured to remove one or more features that contain analytes of interest from a substrate by projecting light toward the features of interest on the substrate.

(16) FIG. 1 illustrates an exemplary system **100** for preparing a biological sample. The system **100** includes a substrate **102**, a targeted feature capture device **104**, and a targeted sample preparation system **106**.

(17) In some implementations, the substrate **102** is configured to position an array of features **110**. The feature array **110** can include spatially-barcoded capture probes, or an array of beads with such capture probes. The array of features **110** can be directly or indirectly attached or fixed to the substrate **102**, or disposed on the substrate **102** in various manners described herein. The array of features **110** can include spatially-barcoded features for array-based spatial analysis, as described herein.

(18) The features in the feature array **110** can include analytes captured from a biological sample, such as a cell or a tissue. Therefore, the substrate **102** does not include the biological sample (e.g., cell, tissue, etc.), but includes the feature array. Alternatively, the features in the feature array **110** can include intermediate agents (e.g., analyte capture agents and/or ligation products) obtained after interacting with analytes in the biological sample. Therefore, the substrate **102** does not include the biological sample (e.g., cell, tissue, etc.), but includes the feature array. Various systems and methods can be used to capture analytes from a sample using the feature array **110**. For example, the substrate **102** with the feature array **110** can be used for array-based spatial analysis, as described herein. For example, an array-based spatial analysis method described herein can be used to transfer one or more analytes from a sample to the feature array **110** on the substrate. Each feature may be associated with a unique spatial location on the array, and used to identify the analytes and the spatial location of each analyte based on the relative spatial location of the feature to which that the analyte is bound on the feature array **110**. Exemplary array-based spatial analysis methods are further described herein.

(19) The targeted feature capture device **104** is configured to selectively retrieve one or more of the features that are of interest from the feature array **110**. The targeted feature capture device **104** is configured to be positioned at a distance from the feature array **110** of the substrate **102**, or abutted to the feature array **110** of the substrate **102** so that one or more features of interest can be captured by the targeted feature capture device **104**.

(20) In some implementations, the feature capture device **104** includes a body **120** having a light receiving surface **122** and a capture surface **124**. The body **120** can be configured as a block that is transparent so that light can pass through. The body **120** can be made of various materials, such as polymer, glass, and other suitable material having desired light transmittance. As described herein, the light receiving surface **122** of the body **120** provides a surface that receives light emitted thereon. The capture surface **124** of the body **120** is configured to be disposed relative to the feature array **110** on the substrate **102**. For example, the capture surface **124** of the body **120** can be positioned at a distance from the feature array **110** on the substrate **102**. Alternatively, the capture surface **124** can be positioned to be abutted to the feature array **110** of the substrate **102**.

(21) The capture surface **124** is configured to capture one or more features from the feature array **110** on the substrate **102** when the light is projected on a region of the capture surface **124** that corresponds to the features in the feature array **110**. In some implementations, the capture surface **124** includes a selective adhesive layer **126** that can be triggered to be adhesive under predetermined conditions. For example, the selective adhesive layer

126 is not adhesive under normal conditions (e.g., when activated by light or heat), and becomes adhesive only when one or more predetermined conditions (e.g., when activated by light of a certain wavelength or heat of a certain temperature) are satisfied. In some implementations, the selective adhesive layer **126** includes a light-activated adhesive material that is activated to be adhesive when light of predetermined characteristic is projected onto the layer **126**. Examples of such a light-activated adhesive material include special polymer and other suitable materials. The light-activated adhesive material can be provided to the capture surface **124** in various manners. In one example, the light-activated adhesive material can be coated on the capture surface **124** of the body **120**.

(22) The targeted sample preparation system **106** is configured to identify a region of interest **204** on the feature array **110** and retrieve selected features from the feature array **110** using the targeted feature capture device **104**. In some implementations, the targeted sample preparation system **106** can include a region of interest (ROI) identifying device **130**, a light generator **132**, a feature retrieving device **134**, and a controller **136**.

(23) The ROI identifying device **130** is configured to identify a ROI **204** on the feature array **110** so that one or more features are selectively retrieved from the feature array **110**. In some implementations, the ROI identifying device **130** can take an image of the feature array **110** on the substrate **102** in real-time and provide the image to a user so that the user can visualize the feature array **110**. For example, the ROI identifying device **130** can include, or be connected to, a display device (e.g., a display screen), and stream the image of the feature array **110** to the display device that can output the image. Examples of the ROI identifying device **130** can include a microscope, a camera, or other suitable imaging devices configured to visualize the feature array **110** of the substrate **102**.

(24) In some implementations, the ROI identifying device **130** is configured to automatically identify a ROI on the feature array **110** based on one or more inputs or factors, such as particular types of analytes of interest, particular research purposes, etc. For example, the ROI identifying device **130** can employ a machine learning model that is trained for one or more analysis or research purposes, and configured to automatically determine a ROI on the feature array **110** to meet a particular analysis or research purpose.

(25) In some implementations, a biological sample can be stained to facilitate visualization. Example methods for staining a sample are described herein. The ROI identifying device **130** can capture an image of the stained sample. Optionally, the ROI identifying device **130** can interpret low pass sequencing data (e.g., via amplification of second strand synthesis on the sample removed from the substrate). The removed sample can be placed back on the substrate. The ROI identifying device **130** can visually identify a region of interest on the image based on the stain and/or the sequencing data that have been optionally obtained. The ROI identifying device **130** can be used to permit a user to select a ROI based on the stain image. In some implementation, the ROI identifying device **130**, which may employ a machine learning algorithm, can automatically identify and mark a ROI on the stained image. In some implementations, the user can control the light generator **132** to emit light toward the identified ROI on the feature array **110** through the feature capture device **104**. In other implementations, the light generator **132** can be automatically controlled and guided to be placed so that light is emitted toward the identified ROI on the feature array **110** through the feature capture device **104**.

(26) The light generator **132** can be configured to generate and emit light on the substrate **102** through the feature capture device **104**. In some implementations, the light generator **132** is manually controllable to be positioned relative to the substrate **102** and/or the feature capture device **104**, and also manually controllable to generate and emit light to a particular area (e.g., the identified ROI) on the feature array **110** of the substrate **102**. For example, the light generator **132** can include a user interface (e.g., physical or virtual buttons, switches, or other suitable input devices) for receiving user input of controlling the light generator **132**. Alternatively or in addition, the light generator **132** is configured to automatically adjust its position relative to the substrate **102** and/or the feature capture device **104** and generate and emit light toward a particular area (e.g., the identified ROI) on the feature array **110** of the substrate **102**.

(27) The feature retrieving device **134** is configured to retrieve features from the feature capture device **104**. For example, when the features are captured at the capture surface **124** of the feature capture device **104**, the feature retrieving device **134** is used to release the features from the capture surface **124** of the feature capture device **104**. The feature retrieving device **134** can use various configurations and methods to retrieve the features from the feature capture device **104**. For example, the feature retrieving device **134** can provide a dissolvent to dissolve a material (e.g., the selective adhesive layer **126**) coated on the capture surface **124** so that the features are released from the feature capture device **104**.

(28) The controller **136** can be configured to control the ROI identifying device **130**, the light generator **132**, and/or the feature retrieving device **134**. In some implementations, the controller **136** is configured to control the ROI identifying device **130** to identify the ROI on the feature array **110** and receive data indicative of the identified ROI. The controller **136** can control the light generator **132** to arrange or orient the light generator **132** in a desired position relative to the feature array **110** on the substrate **102**, and/or generate and project light onto the identified ROI on the feature array **110** through the feature capture device **104** so that the features on the identified ROI are captured to the capture surface **124** of the feature capture device **104**. The controller **136** can further control the feature retrieving device **134** to release the captured features from the capture surface **124** for further analysis.

(29) FIGS. 2A-C illustrate an exemplary process **200** for performing spatially-targeted feature capture. In this example, the process **200** is illustrated to be performed using the system **100** described in FIG. 1. Referring to FIG. 2A, the process **200** can be performed after the feature array **110** on the substrate **102** has captured analytes from a biological sample. The feature capture device **104** is arranged to a desired position relative to the substrate **102**. For example, the feature capture device **104** is arranged such that the capture surface **124** faces at least a portion of the feature array **110** on the substrate **102**. In some implementations, the feature capture device **104** can be arranged such that the capture surface **124** is spaced apart at a distance **202** from the feature array **110** on the substrate **102**. The distance **202** can be determined to be small enough for the material **126** at the capture surface **124** to extend toward and adhere to targeted features of the feature array **110**, when the material **126** is triggered by a projected light as described herein. The distance **202** can range between 0 to 100 microns in some examples. In another example, the distance **202** can range between 0 to 50 microns. A bigger distance **202** is possible depending on, for example, the characteristics (e.g., a degree of stretching when light-activated) of the material **126** at the capture surface **124**. In other implementations, the feature capture device **104** can be arranged such that the capture surface **124** contacts with at least a portion of the feature array **110**.

(30) In some implementations, the ROI identifying device **130** can be used to identify a region of interest (ROI) **204** in the feature array **110** on the substrate **102**. The ROI **204** is determined to include one or more target features **206** among the feature array **110**. As described herein, the ROI identifying device **130** can provide a visual image of the feature array **110** to a user so that the user can identify the ROI **204** from the visual image of the feature array **110**. In addition or alternatively, the ROI identifying device **130** can automatically identify the ROI **204**, and mark the ROI **204** on the image of the feature array **110**, so that light can be accurately projected onto the ROI **204**.

(31) Referring to FIG. 2B, the light generator **132** is used to emit light **208** onto the ROI **204** of the feature array **110**. In some implementations, the light generator **132** generates a beam of light **208** of suitable wavelength and projects the light beam **208** onto the ROI **204**. For example, the light beam **208** can be a laser of suitable wavelength.

(32) In some implementations, the light **208** is projected onto the light receiving surface **122**, and passes through the transparent body **120** to reach the capture surface **124**. For example, the light receiving surface **122** is arranged opposite to the capture surface **124** so that the light **208** travels a linear path through the body **120**. Alternatively, the light receiving surface **122** can be arranged a different side of the body **120** so that the light **208** is projected onto such a different side of the body **120** and travels along a non-linear path from the light receiving surface **122** to the capture surface **124**. In this configuration, one or more reflectors can be included in the body **120** to change the path of the light **208**, or the interior of the body **120** can be configured to form a non-linear light path toward the capture surface **124**.

(33) The light **208** can locally activate the layer **126** at the capture surface **124** so that the layer **126** becomes adhesive at the area (e.g., the ROI **204**) that receives the light **208**. For example, the layer **126** can absorb the energy from the light **208** and change its properties to become adhesive. The area of the layer **126** that becomes locally adhesive can attach to the light-targeted features **206** on the substrate **102**.

(34) Referring to FIG. 2C, the feature capture device **104** that attaches the light-targeted features **206** can be removed from the light-targeted features **206**. The light-targeted features **206** can be retrieved from the capture surface **124** of the feature capture device **104**.

(35) In some implementations, the feature retrieving device **134** can be used to release the light-targeted features **206** from the capture surface **124**. As described herein, for example, the feature retrieving device **134** can include a chamber that contains a dissolvent for dissolving a material (e.g., the selective adhesive layer **126**) coated on the capture surface **124**. The feature retrieving device **134** can be at least partially immersed in the chamber so that the features are released from the feature capture device **104**. Alternatively, the feature retrieving device **134** can provide a spray or other types of injectors for discharging a dissolvent toward the material (e.g., the selective adhesive layer **126**) coated on the capture surface **124** so that the features are released from the feature capture device **104**.

(36) FIG. 3 is a flowchart of an example process **300** for performing spatially-targeted feature capture. In some implementations, the process **300** can be at least partially performed using the system **100** described in FIG. 1. In addition or alternatively, the process **300** can be used to implement at least part of the process **200** described in FIGS. 2A-C.

(37) The process **300** can include placing a feature array on a substrate (**302**). The feature array can include spatially-barcoded capture probes, or an array of beads with such capture probes or functionality. For example, the feature array can be directly or indirectly attached or fixed to the substrate. Other methods can be used to place the feature array on the substrate.

(38) The process **300** can include capturing analytes from a biological sample on the feature array of the substrate (**304**). Various methods can be used for capturing analytes from a sample. In one example, a biological sample is permeabilized to allow analytes to migrate away from the sample and toward the feature array of the substrate. The analytes can interact (e.g., hybridize) with the features (e.g., capture probes therein) on the feature array.

(39) The process **300** can include arranging a feature capture cap relative to the substrate (**306**). The feature capture cap can be configured to be identical or similar to the feature capture device **104**. In some implementations, the feature capture cap is at least partially made of a transparent body and includes a capture surface that is light-activated to be adhesive. The capture surface of the cap can be arranged close to, or abutted to, the feature array on the substrate.

(40) The process **300** can include identifying a region of interest (ROI) on the feature array (**308**). An ROI is to determine one or more target features that contain analytes of interest on the feature array. In some implementations, an ROI can be identified by a user who can review an image generated for the feature array on the substrate. In alternative implementations, the ROI can be automatically identified on the feature array so that analytes of interest can be obtained by subsequent steps in the process **300**.

(41) The process **300** can include emitting light onto the substrate through the cap (**310**). For example, a beam of light (e.g., a laser beam of suitable wavelength) can be projected through the cap onto the ROI of the feature array on the substrate. When the light hits the capture surface of the cap, the capture surface can absorb the energy and become locally adhesive. Then, the adhesive area of the capture surface can attach target features in the ROI so that the target features in the ROI are removed from the feature array on the substrate.

(42) The process **300** can include removing the cap (**312**). The cap with the target features attached to the capture surface can be pulled away from the substrate. The process **300** can include removing the target features from the cap (**314**). Various methods can be used to retrieve the target features from the capture surface of the cap. For example, the target features can be removed from the cap by a dissolvent for dissolving the materials coated on the capture surface.

(43) Referring to FIGS. 4-6, another exemplary system for preparing a biological sample is described. In some implementations, the system is configured to obtain one or more analytes (or intermediate agents, such as analyte capture agents) that are spatially targeted. For example, the system can be configured to remove one or more beads that contain analytes of interest from a substrate by projecting light toward the beads of interest on the substrate.

(44) FIG. 4 illustrates an exemplary system **500** for preparing a biological sample. The system **500** includes a substrate **502** and a targeted sample preparation system **506**. In some implementation, the substrate **502** and the targeted sample preparation system **506** can be configured similarly to the substrate **102** and the targeted sample preparation system **106** described herein.

(45) In some implementations, the substrate **502** is configured to position an array of features **510**. The feature array **510** can include spatially-barcoded capture probes, or an array of beads with such capture probes. The array of features **510** can be directly or indirectly attached or fixed to the substrate **502**, or disposed on the substrate **502** in various manners described herein. The array of features **510** can include spatially-barcoded features for array-based spatial analysis, as described herein.

(46) In this example, the feature array **510** can be attached to the substrate **502** with photo-sensitive bonds **602** (FIG. 5A), such as cleavable linkers. An exemplary photo-sensitive bond (e.g., cleavable capture probe) is described herein, for example with reference to FIG. 7. For example, the photo-sensitive bonds **602** can be cleaved when exposed to light, such as UV light. Example linkers of the photo-sensitive bond include-amino-3-(2-nitrophenyl) propionic acid (ANP), phenacyl ester derivatives, 8-quinolinyln benzenesulfonate, dicoumarin, 6-bromo-7-alkoxycoumarin-4-ylmethoxycarbonyl, a bimeane-based linker, and a bis-arylhydrazone based linker.

(47) The features in the feature array **510** can include analytes captured from a biological sample, such as a cell or a tissue. Various systems and methods can be used to capture analytes from a sample using the feature array **510**. For example, the substrate **502** with the feature array **510** can be used for array-based spatial analysis, as described herein. For example, an array-based spatial analysis method described herein can be used to transfer one or more analytes from a sample to the feature array **510** on the substrate. Each feature may be associated with a unique spatial location on the array, and used to identify the analytes and the spatial location of each analyte based on the relative spatial location of the feature to which that the analyte is bound on the feature array **510**. Example array-based spatial analysis methods are further described herein.

(48) The targeted sample preparation system **506** is configured to identify a region of interest (ROI) **604** on the feature array **510** and retrieve selected features from the feature array **510** by cleaving the linkers of the features. In some implementations, the targeted sample preparation system **506** can include a ROI identifying device **530**, a light generator **532**, a feature collecting device **534**, and a controller **536**.

(49) The ROI identifying device **530** is configured to identify a ROI **604** on the feature array **510** so that one or more features are selectively retrieved from the feature array **510**. In some implementations, the ROI identifying device **530** can take an image of the feature array **510** on the substrate **502** in real-time and provide the image to a user so that the user can visualize the feature array **510**. For example, the ROI identifying device **530** can include, or be connected to, a display device (e.g., a display screen), and stream the image of the feature array **510** to the display device that can output the image. Examples of the ROI identifying device **530** can include a microscope, a camera, or other suitable imaging devices configured to visualize the feature array **510** of the substrate **502**.

(50) In some implementations, the ROI identifying device **530** is configured to automatically identify a ROI on the feature array **510** based on one or more inputs or factors, such as particular types of analytes of interest, particular research purposes, etc. For example, the ROI identifying device **530** can employ a machine learning model that is trained for one or more analysis or research purposes, and configured to automatically determine a ROI on the feature array **510** to meet a particular analysis or research purpose.

(51) In some implementations, a biological sample can be stained to facilitate visualization. Example methods for staining a sample are described herein. The ROI identifying device **530** can capture an image of the stained sample. Optionally, the ROI identifying device **530** can interpret low pass sequencing data (e.g., via amplification of second strand synthesis on the sample removed from the substrate). The removed sample can be placed back on the substrate. The ROI identifying device **530** can visually identify a region of interest on the image based on the stain and/or the sequencing data that have been optionally obtained. The ROI identifying device **530** can be used to permit a user to select a ROI based on the stain image. In some implementations, the ROI identifying device **530**, which may employ a machine learning algorithm, can automatically identify and mark a ROI

on the stained image. In some implementations, the user can control the light generator 532 to emit light toward the identified ROI on the feature array 510. In other implementations, the light generator 532 can be automatically controlled and guided to be placed so that light is emitted onto the identified ROI of the feature array 510.

(52) The light generator 532 is configured to generate and emit light onto the substrate 502. In some implementations, the light generator 532 is manually controllable to be positioned relative to the substrate 502, and also manually controllable to generate and emit light to a particular area (e.g., the identified ROI) on the feature array 510 of the substrate 502. For example, the light generator 532 includes a user interface (e.g., physical or virtual buttons, switches, or other suitable input devices) for receiving a user input of controlling the light generator 532. Alternatively or in addition, the light generator 532 is configured to automatically adjust its position relative to the substrate 502 and generate and emit light toward a particular area (e.g., the identified ROI) on the feature array 510 of the substrate 502.

(53) The feature collecting device 534 is configured to retrieve target features from the substrate 502. For example, when the linkers are cleaved and the features are released from the substrate 502, the feature collecting device 534 is used to collect the released features. The feature collecting device 534 can use various configurations and methods to retrieve the features removed from the substrate 502. In some implementations, the feature collecting device 534 can include one or more reservoirs that each contain a fluid and is used to collect the removed target features 606 therein. In some implementations, the feature collecting device 534 can include a plurality of reservoirs used to separately collect target features of different characteristics (e.g., physical properties, cell types, gene expression profiles, disease vs. normal, etc.).

(54) The controller 536 can be configured to control the ROI identifying device 530, the light generator 532, and/or the feature collecting device 534. In some implementations, the controller 536 is configured to control the ROI identifying device 530 to identify the ROI on the feature array 510 and receive data indicative of the identified ROI. The controller 536 can control the light generator 532 to arrange or orient the light generator 532 in a desired position relative to the feature array 510 on the substrate 502, and/or generate and project light onto the identified ROI on the feature array 510 of the substrate 502 so that the features on the identified ROI are removed from the substrate 502. The controller 536 can further control the feature collecting device 534 to collect the removed features for further analysis.

(55) FIGS. 5A-C illustrate an exemplary process 600 for performing spatially-targeted feature capture. In this example, the process 600 is illustrated to be performed using the system 500 described in FIG. 4. Referring to FIG. 5A, the process 600 can be performed after the feature array 510 on the substrate 502 has captured analytes from a biological sample. As described herein, the feature array 510 can be attached to the substrate 502 with photo-sensitive bonds 602, such as linkers that are cleavable when exposed to light.

(56) In some implementations, the ROI identifying device 530 can be used to identify a region of interest (ROI) 604 in the feature array 510 on the substrate 502. The ROI 604 is determined to include one or more target features 606 among the feature array 510. As described herein, the ROI identifying device 530 can provide a visual image of the feature array 510 to a user so that the user can identify the ROI 604 from the visual image of the feature array 510. In addition or alternatively, the ROI identifying device 530 can automatically identify the ROI 604, and mark the ROI 604 on the image of the feature array 510, so that light can be accurately projected onto the ROI 604.

(57) Referring to FIG. 5B, the light generator 532 is used to emit light 608 onto the ROI 604 of the feature array 510. In some implementations, the light generator 532 generates a beam of light 608 of suitable wavelength and projects the light beam 608 onto the ROI 604. For example, the light beam 608 can be an ultraviolet (UV) laser. Alternatively, the light generator 532 can include a digital micro-mirror device.

(58) In some implementations, the light 608 is projected onto a bottom surface 610 of the substrate 502 that is opposite to a top surface 612 of the substrate 502 on which the feature array 510 is placed. The light 608 that enters the bottom surface 610 of the substrate 502 passes through a transparent body 614 of the substrate 502 and reaches the top surface 612 of the substrate 502. The light 608 can locally cleave the photo-sensitive bonds 602 in the ROI 604 so that the target features 606 in the ROI 604 are released from the substrate 502.

(59) Referring to FIG. 5C, the target features 606 are released from the substrate 502 when the photo-sensitive bonds 602 are cleaved by the light 608 projected onto the ROI 604. In some implementations, the feature collecting device 534 can be used to collect the target features 606. As described herein, for example, the feature collecting device 534 can include a reservoir that contains a fluid, and the removed target features 606 can be collected in the reservoir. In some implementations, a plurality of reservoirs can be provided and used to separately collect target features 606 of different characteristics (e.g., physical properties, cell types, gene expression profiles, disease vs. normal, etc.). For example, a first reservoir can be used to collect target features 606 having a first characteristic while a second reservoir can be used to collect target features 606 having a second, different characteristic.

(60) FIG. 6 is a flowchart of an exemplary process 700 for performing spatially-targeted feature capture. In some implementations, the process 700 can be at least partially performed using the system 100 described in FIG. 4. In addition or alternatively, the process 700 can be used to implement at least part of the process 600 described in FIGS. 5A-5C.

(61) The process 700 can include placing a feature array on a substrate with cleavable linkers (702). The feature array can include spatially-barcoded capture probes, or an array of beads with such capture probes or functionality. Additional description of spatially-barcoded capture probes is provided herein; see also, for example, WO 2020/176788 (e.g., Section (II) (b), such as subsections (i)-(vi)), U.S. Patent Application Publication No. 2020/0277663 (e.g., Section (II) (b) (e.g., subsections (i)-(vi))), and/or U.S. Patent Application Publication No. 2019/0264268 (e.g., paragraphs [0088]-[0109]). For example, the feature array can be directly or indirectly attached or fixed to the substrate. Other methods can be used to place the feature array on the substrate. The feature array can be attached to the substrate using cleavable linkers or other types of photo-sensitive bonds as described herein. The linkers are cleavable when exposed to light of certain characteristics (e.g., wavelength). For example, the linkers can be selected to be cleaved when exposed to a UV light.

(62) The process 700 can include capturing analytes from a biological sample on the feature array of the substrate (704). Various methods can be used for capturing analytes from a sample. In one example, a biological sample is permeabilized to allow analytes to migrate away from a sample and toward the feature array of the substrate. The analytes can interact with the features (e.g., capture probes therein) on the feature array.

(63) The process 700 can include identifying a region of interest (ROI) on the feature array (706). An ROI is used to determine one or more target features that contain analytes of interest on the feature array. In some implementations, an ROI can be identified by a user who can review an image generated for the feature array on the substrate. In alternative implementations, the ROI can be automatically identified on the feature array so that the analytes of interest can be obtained by subsequent steps in the process 700.

(64) The process 700 can include emitting light onto the substrate through the cap (708). For example, a beam of light (e.g., a laser beam of suitable wavelength) can be projected onto the ROI of the feature array of the substrate. When the light hits the cleavable linkers in the ROI, the linkers are cleaved so that the features attached to the linkers are released from the substrate.

(65) The process 700 can include collecting the removed target features (710). Various methods can be used to retrieve the target features from the capture surface of the cap. For example, the target features removed from the substrate can be collected within a fluid of a particular type contained in a reservoir.

(66) FIG. 7 is a block diagram of computing devices 900 that may be used to implement the systems and methods described in this document, as either a client or as a server or plurality of servers. For example, at least some of the components described in FIG. 7 can be used to implement the targeted sample preparation system 106, 506 and other instruments, apparatuses, and devices described herein. Computing device 900 is intended to represent various forms of digital computers, such as laptops, desktops, workstations, personal digital assistants, servers, blade servers, mainframes, and other appropriate computers. The components shown here, their connections and relationships, and their functions, are meant to be examples only, and are not meant to limit implementations described and/or claimed in this document.

(67) Computing device 900 includes a processor 902, memory 904, a storage device 906, a high-speed interface 908 connecting to memory 904 and

high-speed expansion ports **910**, and a low speed interface **912** connecting to low speed bus **914** and storage device **906**. Each of the components **902**, **904**, **906**, **908**, **910**, and **912**, are interconnected using various busses, and may be mounted on a common motherboard or in other manners as appropriate. The processor **902** can process instructions for execution within the computing device **900**, including instructions stored in the memory **904** or on the storage device **906** to display graphical information for a GUI on an external input/output device, such as display **916** coupled to high-speed interface **908**. In other implementations, multiple processors and/or multiple buses may be used, as appropriate, along with multiple memories and types of memory. Also, multiple computing devices **900** may be connected, with each device providing portions of the necessary operations (e.g., as a server bank, a group of blade servers, or a multi-processor system).

(68) The memory **904** stores information within the computing device **900**. In one implementation, the memory **904** is a volatile memory unit or units. In another implementation, the memory **904** is a non-volatile memory unit or units. The memory **904** may also be another form of computer-readable medium, such as a magnetic or optical disk.

(69) The storage device **906** is capable of providing mass storage for the computing device **900**. In one implementation, the storage device **906** may be or contain a computer-readable medium, such as a floppy disk device, a hard disk device, an optical disk device, or a tape device, a flash memory or other similar solid state memory device, or an array of devices, including devices in a storage area network or other configurations. A computer program product can be tangibly embodied in an information carrier. The computer program product may also contain instructions that, when executed, perform one or more methods, such as those described above. The information carrier is a computer- or machine-readable medium, such as the memory **904**, the storage device **906**, or memory on processor **902**.

(70) The high-speed controller **908** manages bandwidth-intensive operations for the computing device **900**, while the low speed controller **912** manages lower bandwidth-intensive operations. Such allocation of functions is an example only. In one implementation, the high-speed controller **908** is coupled to memory **904**, display **916** (e.g., through a graphics processor or accelerator), and to high-speed expansion ports **910**, which may accept various expansion cards (not shown). In the implementation, low-speed controller **912** is coupled to storage device **906** and low-speed expansion port **914**. The low-speed expansion port, which may include various communication ports (e.g., USB, Bluetooth, Ethernet, wireless Ethernet) may be coupled to one or more input/output devices, such as a keyboard, a pointing device, a scanner, or a networking device such as a switch or router, e.g., through a network adapter.

(71) The computing device **900** may be implemented in a number of different forms, as shown in the figure. For example, it may be implemented as a standard server **920**, or multiple times in a group of such servers. It may also be implemented as part of a rack server system **924**. In addition, it may be implemented in a personal computer such as a laptop computer **922**. Alternatively, components from computing device **900** may be combined with other components in a mobile device. Each of such devices may contain one or more of computing device **900**, and an entire system may be made up of multiple computing devices **900** communicating with each other.

(72) Additionally computing device **900** can include Universal Serial Bus (USB) flash drives. The USB flash drives may store operating systems and other applications. The USB flash drives can include input/output components, such as a wireless transmitter or USB connector that may be inserted into a USB port of another computing device.

(73) Various implementations of the systems and techniques described here can be realized in digital electronic circuitry, integrated circuitry, specially designed ASICs (application specific integrated circuits), computer hardware, firmware, software, and/or combinations thereof. These various implementations can include implementation in one or more computer programs that are executable and/or interpretable on a programmable system including at least one programmable processor, which may be special or general purpose, coupled to receive data and instructions from, and to transmit data and instructions to, a storage system, at least one input device, and at least one output device.

(74) These computer programs (also known as programs, software, software applications or code) include machine instructions for a programmable processor, and can be implemented in a high-level procedural and/or object-oriented programming language, and/or in assembly/machine language. As used herein, the terms “machine-readable medium” “computer-readable medium” refers to any computer program product, apparatus and/or device (e.g., magnetic discs, optical disks, memory, Programmable Logic Devices (PLDs)) used to provide machine instructions and/or data to a programmable processor, including a machine-readable medium that receives machine instructions as a machine-readable signal. The term “machine-readable signal” refers to any signal used to provide machine instructions and/or data to a programmable processor.

(75) To provide for interaction with a user, the systems and techniques described here can be implemented on a computer having a display device (e.g., a CRT (cathode ray tube) or LCD (liquid crystal display) monitor) for displaying information to the user and a keyboard and a pointing device (e.g., a mouse or a trackball) by which the user can provide input to the computer. Other kinds of devices can be used to provide for interaction with a user as well; for example, feedback provided to the user can be any form of sensory feedback (e.g., visual feedback, auditory feedback, or tactile feedback); and input from the user can be received in any form, including acoustic, speech, or tactile input.

(76) The systems and techniques described here can be implemented in a computing system that includes a back end component (e.g., as a data server), or that includes a middleware component (e.g., an application server), or that includes a front end component (e.g., a client computer having a graphical user interface or a Web browser through which a user can interact with an implementation of the systems and techniques described here), or any combination of such back end, middleware, or front end components. The components of the system can be interconnected by any form or medium of digital data communication (e.g., a communication network). Examples of communication networks include a local area network (“LAN”), a wide area network (“WAN”), peer-to-peer networks (having ad-hoc or static members), grid computing infrastructures, and the Internet.

(77) The computing system can include clients and servers. A client and server are generally remote from each other and typically interact through a communication network. The relationship of client and server arises by virtue of computer programs running on the respective computers and having a client-server relationship to each other.

(78) Spatial analysis methodologies and compositions described herein can provide a vast amount of analyte and/or expression data for a variety of analytes within a biological sample at high spatial resolution, while retaining native spatial context. Spatial analysis methods and compositions can include, e.g., the use of a capture probe including a spatial barcode (e.g., a nucleic acid sequence that provides information as to the location or position of an analyte within a cell or a tissue sample (e.g., mammalian cell or a mammalian tissue sample) and a capture domain that is capable of binding to an analyte (e.g., a protein and/or a nucleic acid) produced by and/or present in a cell. Spatial analysis methods and compositions can also include the use of a capture probe having a capture domain that captures an intermediate agent for indirect detection of an analyte. For example, the intermediate agent can include a nucleic acid sequence (e.g., a barcode) associated with the intermediate agent. Detection of the intermediate agent is therefore indicative of the analyte in the cell or tissue sample.

(79) Non-limiting aspects of spatial analysis methodologies and compositions are described in U.S. Pat. Nos. 10,774,374, 10,724,078, 10,480,022, 10,059,990, 10,041,949, 10,002,316, 9,879,313, 9,783,841, 9,727,810, 9,593,365, 8,951,726, 8,604,182, 7,709,198, U.S. Patent Application Publication Nos. 2020/239946, 2020/080136, 2020/0277663, 2020/024641, 2019/330617, 2019/264268, 2020/256867, 2020/224244, 2019/194709, 2019/161796, 2019/085383, 2019/055594, 2018/216161, 2018/051322, 2018/0245142, 2017/241911, 2017/089811, 2017/067096, 2017/029875, 2017/0016053, 2016/108458, 2015/000854, 2013/171621, WO 2018/091676, WO 2020/176788, Rodriques et al., Science 363 (6434): 1463-1467, 2019; Lee et al., Nat. Protoc. 10 (3): 442-458, 2015; Trejo et al., PLOS ONE 14 (2): e0212031, 2019; Chen et al., Science 348 (6233): aaa6090, 2015; Gao et al., BMC Biol. 15:50, 2017; and Gupta et al., Nature Biotechnol. 36:1197-1202, 2018; the Visium Spatial Gene Expression Reagent Kits User Guide (e.g., Rev C, dated June 2020), and/or the Visium Spatial Tissue Optimization Reagent Kits User Guide (e.g., Rev C, dated July 2020), both of which are available at the 10× Genomics Support Documentation website, and can be used herein in any combination. Further non-limiting aspects of spatial analysis methodologies and compositions are described herein.

(80) Some general terminology that may be used in this disclosure can be found in Section (I)(b) of WO 2020/176788 and/or U.S. Patent Application

Publication No. 2020/0277663. Typically, a “barcode” is a label, or identifier, that conveys or is capable of conveying information (e.g., information about an analyte in a sample, a bead, and/or a capture probe). A barcode can be part of an analyte, or independent of an analyte. A barcode can be attached to an analyte. A particular barcode can be unique relative to other barcodes. For the purpose of this disclosure, an “analyte” can include any biological substance, structure, moiety, or component to be analyzed. The term “target” can similarly refer to an analyte of interest.

(81) Analytes can be broadly classified into one of two groups: nucleic acid analytes, and non-nucleic acid analytes. Examples of non-nucleic acid analytes include, but are not limited to, lipids, carbohydrates, peptides, proteins, glycoproteins (N-linked or O-linked), lipoproteins, phosphoproteins, specific phosphorylated or acetylated variants of proteins, amidation variants of proteins, hydroxylation variants of proteins, methylation variants of proteins, ubiquitylation variants of proteins, sulfation variants of proteins, viral proteins (e.g., viral capsid, viral envelope, viral coat, viral accessory, viral glycoproteins, viral spike, etc.), extracellular and intracellular proteins, antibodies, and antigen binding fragments. In some embodiments, the analyte(s) can be localized to subcellular location(s), including, for example, organelles, e.g., mitochondria, Golgi apparatus, endoplasmic reticulum, chloroplasts, endocytic vesicles, exocytic vesicles, vacuoles, lysosomes, etc. In some embodiments, analyte(s) can be peptides or proteins, including without limitation antibodies and enzymes. Additional examples of analytes can be found in Section (I)(c) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663. In some embodiments, an analyte can be detected indirectly, such as through detection of an intermediate agent, for example, a ligation product or an analyte capture agent (e.g., an oligonucleotide-conjugated antibody), such as those described herein.

(82) A “biological sample” is typically obtained from the subject for analysis using any of a variety of techniques including, but not limited to, biopsy, surgery, and laser capture microscopy (LCM), and generally includes cells and/or other biological material from the subject. In some embodiments, a biological sample can be a tissue section. In some embodiments, a biological sample can be a fixed and/or stained biological sample (e.g., a fixed and/or stained tissue section). Non-limiting examples of stains include histological stains (e.g., hematoxylin and/or eosin) and immunological stains (e.g., fluorescent stains). In some embodiments, a biological sample (e.g., a fixed and/or stained biological sample) can be imaged. Biological samples are also described in Section (I)(d) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663.

(83) In some embodiments, a biological sample is permeabilized with one or more permeabilization reagents. For example, permeabilization of a biological sample can facilitate analyte capture. Exemplary permeabilization agents and conditions are described in Section (I)(d)(ii)(13) or the Exemplary Embodiments Section of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663.

(84) Array-based spatial analysis methods involve the transfer of one or more analytes from a biological sample to an array of features on a substrate, where each feature is associated with a unique spatial location on the array. Subsequent analysis of the transferred analytes includes determining the identity of the analytes and the spatial location of the analytes within the biological sample. The spatial location of an analyte within the biological sample is determined based on the feature to which the analyte is bound (e.g., directly or indirectly) on the array, and the feature's relative spatial location within the array.

(85) A “capture probe” refers to any molecule capable of capturing (directly or indirectly) and/or labelling an analyte (e.g., an analyte of interest) in a biological sample. In some embodiments, the capture probe is a nucleic acid or a polypeptide. In some embodiments, the capture probe includes a barcode (e.g., a spatial barcode and/or a unique molecular identifier (UMI)) and a capture domain. In some embodiments, a capture probe can include a cleavage domain and/or a functional domain (e.g., a primer-binding site, such as for next-generation sequencing (NGS)). See, e.g., Section (II)(b) (e.g., subsections (i)-(vi)) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663. Generation of capture probes can be achieved by any appropriate method, including those described in Section (II)(d)(ii) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663.

(86) In some embodiments, more than one analyte type (e.g., nucleic acids and proteins) from a biological sample can be detected (e.g., simultaneously or sequentially) using any appropriate multiplexing technique, such as those described in Section (IV) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663.

(87) In some embodiments, detection of one or more analytes (e.g., proteins) can be performed using one or more analyte capture agents. As used herein, an “analyte capture agent” refers to an agent that interacts with an analyte (e.g., an analyte in a biological sample) and with a capture probe (e.g., a capture probe attached to a substrate or a feature) to identify the analyte. In some embodiments, the analyte capture agent includes: (i) an analyte binding moiety (e.g., that binds to an analyte), for example, an antibody or antigen-binding fragment thereof; (ii) analyte binding moiety barcode; and (iii) an analyte capture sequence. As used herein, the term “analyte binding moiety barcode” refers to a barcode that is associated with or otherwise identifies the analyte binding moiety. As used herein, the term “analyte capture sequence” refers to a region or moiety configured to hybridize to, bind to, couple to, or otherwise interact with a capture domain of a capture probe. In some cases, an analyte binding moiety barcode (or portion thereof) may be able to be removed (e.g., cleaved) from the analyte capture agent. Additional description of analyte capture agents can be found in Section (II)(b)(ix) of WO 2020/176788 and/or Section (II)(b)(viii) U.S. Patent Application Publication No. 2020/0277663.

(88) There are at least two methods to associate a spatial barcode with one or more neighboring cells, such that the spatial barcode identifies the one or more cells, and/or contents of the one or more cells, as associated with a particular spatial location. One method is to promote analytes or analyte proxies (e.g., intermediate agents) out of a cell and towards a spatially-barcode array (e.g., including spatially-barcode capture probes). Another method is to cleave spatially-barcode capture probes from an array and promote the spatially-barcode capture probes towards and/or into or onto the biological sample.

(89) In some cases, capture probes may be configured to prime, replicate, and consequently yield optionally barcoded extension products from a template (e.g., a DNA or RNA template, such as an analyte or an intermediate agent (e.g., a ligation product or an analyte capture agent), or a portion thereof), or derivatives thereof (see, e.g., Section (II)(b)(vii) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663 regarding extended capture probes). In some cases, capture probes may be configured to form ligation products with a template (e.g., a DNA or RNA template, such as an analyte or an intermediate agent, or portion thereof), thereby creating ligation products that serve as proxies for a template.

(90) As used herein, an “extended capture probe” refers to a capture probe having additional nucleotides added to the terminus (e.g., 3' or 5' end) of the capture probe thereby extending the overall length of the capture probe. For example, an “extended 3' end” indicates additional nucleotides were added to the most 3' nucleotide of the capture probe to extend the length of the capture probe, for example, by polymerization reactions used to extend nucleic acid molecules including templated polymerization catalyzed by a polymerase (e.g., a DNA polymerase or a reverse transcriptase). In some embodiments, extending the capture probe includes adding to a 3' end of a capture probe a nucleic acid sequence that is complementary to a nucleic acid sequence of an analyte or intermediate agent specifically bound to the capture domain of the capture probe. In some embodiments, the capture probe is extended using reverse transcription. In some embodiments, the capture probe is extended using one or more DNA polymerases. The extended capture probes include the sequence of the capture probe and the sequence of the spatial barcode of the capture probe.

(91) In some embodiments, extended capture probes are amplified (e.g., in bulk solution or on the array) to yield quantities that are sufficient for downstream analysis, e.g., via next-generation sequencing. In some embodiments, extended capture probes (e.g., DNA molecules) act as templates for an amplification reaction (e.g., a polymerase chain reaction).

(92) Additional variants of spatial analysis methods, including in some embodiments, an imaging step, are described in Section (II) (a) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663. Analysis of captured analytes (and/or intermediate agents or portions thereof), for example, including sample removal, extension of capture probes, sequencing (e.g., of a cleaved extended capture probe and/or a cDNA molecule complementary to an extended capture probe), sequencing on the array (e.g., using, for example, in situ hybridization or in situ ligation approaches), temporal analysis, and/or proximity capture, is described in Section (II) (g) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663. Some quality control measures are described in Section (II) (h) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663.

(93) Spatial information can provide information of biological importance. For example, the methods and compositions described herein can allow for: identification of one or more biomarkers (e.g., diagnostic, prognostic, and/or for determination of efficacy of a treatment) of a disease or disorder; identification of a candidate drug target for treatment of a disease or disorder; identification (e.g., diagnosis) of a subject as having a disease or disorder; identification of stage and/or prognosis of a disease or disorder in a subject; identification of a subject as having an increased likelihood of developing a disease or disorder; monitoring of progression of a disease or disorder in a subject; determination of efficacy of a treatment of a disease or disorder in a subject; identification of a patient subpopulation for which a treatment is effective for a disease or disorder; modification of a treatment of a subject with a disease or disorder; selection of a subject for participation in a clinical trial; and/or selection of a treatment for a subject with a disease or disorder.

(94) Spatial information can provide information of biological importance. For example, the methods and compositions described herein can allow for: identification of transcriptome and/or proteome expression profiles (e.g., in healthy and/or diseased tissue); identification of multiple analyte types in close proximity (e.g., nearest neighbor analysis); determination of up- and/or down-regulated genes and/or proteins in diseased tissue; characterization of tumor microenvironments (e.g., differentiate immune cells from cancer cells); characterization of tumor immune responses; characterization of cells types and their co-localization in tissue; and identification of genetic variants within tissues (e.g., based on gene and/or protein expression profiles associated with specific disease or disorder biomarkers).

(95) Typically, for spatial array-based methods, a substrate functions as a support for direct or indirect attachment of capture probes to features of the array. A “feature” is an entity that acts as a support or repository for various molecular entities used in spatial analysis. In some embodiments, some or all of the features in an array are functionalized for analyte capture. Exemplary substrates are described in Section (II)(c) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663. Exemplary features and geometric attributes of an array can be found in Sections (II)(d)(i), (II)(d)(iii), and (II)(d)(iv) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663.

(96) Generally, analytes and/or intermediate agents (or portions thereof) can be captured when contacting a biological sample with a substrate including capture probes (e.g., a substrate with capture probes embedded, spotted, printed, fabricated on the substrate, or a substrate with features (e.g., beads, wells) comprising capture probes). As used herein, “contact,” “contacted,” and/or “contacting,” a biological sample with a substrate refers to any contact (e.g., direct or indirect) such that capture probes can interact (e.g., bind covalently or non-covalently (e.g., hybridize)) with analytes from the biological sample. Capture can be achieved actively (e.g., using electrophoresis) or passively (e.g., using diffusion). Analyte capture is further described in Section (II) (e) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663.

(97) In some cases, spatial analysis can be performed by attaching and/or introducing a molecule (e.g., a peptide, a lipid, or a nucleic acid molecule) having a barcode (e.g., a spatial barcode) to a biological sample (e.g., to a cell in a biological sample). In some embodiments, a plurality of molecules (e.g., a plurality of nucleic acid molecules) having a plurality of barcodes (e.g., a plurality of spatial barcodes) are introduced to a biological sample (e.g., to a plurality of cells in a biological sample) for use in spatial analysis. In some embodiments, after attaching and/or introducing a molecule having a barcode to a biological sample, the biological sample can be physically separated (e.g., dissociated) into single cells or cell groups for analysis. Some such methods of spatial analysis are described in Section (III) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663.

(98) In some cases, spatial analysis can be performed by detecting multiple oligonucleotides that hybridize to an analyte. In some instances, for example, spatial analysis can be performed using RNA-templated ligation (RTL). Methods of RTL have been described previously. See, e.g., Credle et al., *Nucleic Acids Res.* 2017 Aug. 21; 45 (14): e128. Typically, RTL includes hybridization of two oligonucleotides to adjacent sequences on an analyte (e.g., an RNA molecule, such as an mRNA molecule). In some instances, the oligonucleotides are DNA molecules. In some instances, one of the oligonucleotides includes at least two ribonucleic acid bases at the 3' end and/or the other oligonucleotide includes a phosphorylated nucleotide at the 5' end. In some instances, one of the two oligonucleotides includes a capture domain (e.g., a poly(A) sequence, a non-homopolymeric sequence). After hybridization to the analyte, a ligase (e.g., SplintR ligase) ligates the two oligonucleotides together, creating a ligation product. In some instances, the two oligonucleotides hybridize to sequences that are not adjacent to one another. For example, hybridization of the two oligonucleotides creates a gap between the hybridized oligonucleotides. In some instances, a polymerase (e.g., a DNA polymerase) can extend one of the oligonucleotides prior to ligation. After ligation, the ligation product is released from the analyte. In some instances, the ligation product is released using an endonuclease (e.g., RNase H). The released ligation product can then be captured by capture probes (e.g., instead of direct capture of an analyte) on an array, optionally amplified, and sequenced, thus determining the location and optionally the abundance of the analyte in the biological sample.

(99) During analysis of spatial information, sequence information for a spatial barcode associated with an analyte is obtained, and the sequence information can be used to provide information about the spatial distribution of the analyte in the biological sample. Various methods can be used to obtain the spatial information. In some embodiments, specific capture probes and the analytes they capture are associated with specific locations in an array of features on a substrate. For example, specific spatial barcodes can be associated with specific array locations prior to array fabrication, and the sequences of the spatial barcodes can be stored (e.g., in a database) along with specific array location information, so that each spatial barcode uniquely maps to a particular array location.

(100) Alternatively, specific spatial barcodes can be deposited at predetermined locations in an array of features during fabrication such that at each location, only one type of spatial barcode is present so that spatial barcodes are uniquely associated with a single feature of the array. Where necessary, the arrays can be decoded using any of the methods described herein so that spatial barcodes are uniquely associated with array feature locations, and this mapping can be stored as described above.

(101) When sequence information is obtained for capture probes and/or analytes during analysis of spatial information, the locations of the capture probes and/or analytes can be determined by referring to the stored information that uniquely associates each spatial barcode with an array feature location. In this manner, specific capture probes and captured analytes are associated with specific locations in the array of features. Each array feature location represents a position relative to a coordinate reference point (e.g., an array location, a fiducial marker) for the array. Accordingly, each feature location has an “address” or location in the coordinate space of the array.

(102) Some exemplary spatial analysis workflows are described in the Exemplary Embodiments section of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663. See, for example, the Exemplary embodiment starting with “In some non-limiting examples of the workflows described herein, the sample can be immersed . . .” of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663. See also, e.g., the Visium Spatial Gene Expression Reagent Kits User Guide (e.g., Rev C, dated June 2020), and/or the Visium Spatial Tissue Optimization Reagent Kits User Guide (e.g., Rev C, dated July 2020).

(103) In some embodiments, spatial analysis can be performed using dedicated hardware and/or software, such as any of the systems described in Sections (II)(e)(ii) and/or (V) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663, or any of one or more of the devices or methods described in Sections Control Slide for Imaging, Methods of Using Control Slides and Substrates for, Systems of Using Control Slides and Substrates for Imaging, and/or Sample and Array Alignment Devices and Methods, Informational labels of WO 2020/123320.

(104) Suitable systems for performing spatial analysis can include components such as a chamber (e.g., a flow cell or sealable, fluid-tight chamber) for containing a biological sample. The biological sample can be mounted for example, in a biological sample holder. One or more fluid chambers can be connected to the chamber and/or the sample holder via fluid conduits, and fluids can be delivered into the chamber and/or sample holder via fluidic pumps, vacuum sources, or other devices coupled to the fluid conduits that create a pressure gradient to drive fluid flow. One or more valves can also be connected to fluid conduits to regulate the flow of reagents from reservoirs to the chamber and/or sample holder.

(105) The systems can optionally include a control unit that includes one or more electronic processors, an input interface, an output interface (such

as a display), and a storage unit (e.g., a solid state storage medium such as, but not limited to, a magnetic, optical, or other solid state, persistent, writeable and/or re-writeable storage medium). The control unit can optionally be connected to one or more remote devices via a network. The control unit (and components thereof) can generally perform any of the steps and functions described herein. Where the system is connected to a remote device, the remote device (or devices) can perform any of the steps or features described herein. The systems can optionally include one or more detectors (e.g., CCD, CMOS) used to capture images. The systems can also optionally include one or more light sources (e.g., LED-based, diode-based, lasers) for illuminating a sample, a substrate with features, analytes from a biological sample captured on a substrate, and various control and calibration media.

(106) The systems can optionally include software instructions encoded and/or implemented in one or more of tangible storage media and hardware components such as application specific integrated circuits. The software instructions, when executed by a control unit (and in particular, an electronic processor) or an integrated circuit, can cause the control unit, integrated circuit, or other component executing the software instructions to perform any of the method steps or functions described herein.

(107) In some cases, the systems described herein can detect (e.g., register an image) the biological sample on the array. Exemplary methods to detect the biological sample on an array are described in PCT Application No. 2020/061064 and/or U.S. patent application Ser. No. 16/951,854.

(108) Prior to transferring analytes from the biological sample to the array of features on the substrate, the biological sample can be aligned with the array. Alignment of a biological sample and an array of features including capture probes can facilitate spatial analysis, which can be used to detect differences in analyte presence and/or level within different positions in the biological sample, for example, to generate a three-dimensional map of the analyte presence and/or level. Exemplary methods to generate a two- and/or three-dimensional map of the analyte presence and/or level are described in PCT Application No. 2020/053655 and spatial analysis methods are generally described in WO 2020/061108 and/or U.S. patent application Ser. No. 16/951,864.

(109) In some cases, a map of analyte presence and/or level can be aligned to an image of a biological sample using one or more fiducial markers, e.g., objects placed in the field of view of an imaging system which appear in the image produced, as described in the Substrate Attributes Section, Control Slide for Imaging Section of WO 2020/123320, PCT Application No. 2020/061066, and/or U.S. patent application Ser. No. 16/951,843. Fiducial markers can be used as a point of reference or measurement scale for alignment (e.g., to align a sample and an array, to align two substrates, to determine a location of a sample or array on a substrate relative to a fiducial marker) and/or for quantitative measurements of sizes and/or distances.

Claims

1. A method comprising: providing a substrate comprising a feature array, wherein the feature array comprises a plurality of features and analytes, intermediate agents, or a combination thereof that were previously captured by the plurality of features from a biological sample; arranging a feature capture block relative to the substrate such that a capture surface of the feature capture block faces the feature array on the substrate; identifying a region of interest on the feature array; emitting light toward a portion of the capture surface of the feature capture block that corresponds to the region of interest on the feature array to expose the portion of the capture surface to the light such that a light-activated adhesive on the portion of the capture surface is activated; permitting the light-activated adhesive on the portion of the capture surface exposed to the light to capture at least one feature of the plurality of features within the region of interest on the feature array, the at least one feature comprising at least one element of the analytes, the intermediate agents, or the combination thereof; removing the feature capture block from the substrate; and retrieving the at least one feature from the capture surface of the feature capture block.
 2. The method of claim 1, further comprising: placing the substrate on a substrate stage of a microscope to visualize the feature array on the substrate including the region of interest.
 3. The method of claim 1, wherein the feature capture block includes a transparent body to permit the light to pass through the transparent body, wherein the transparent body of the feature capture block has a light entering surface onto which the light is emitted, the light passing through the transparent body and directed toward the capture surface, and wherein the light entering surface is arranged to be opposite to the capture surface.
 4. The method of claim 1, wherein the capture surface includes a feature capture material, wherein the feature capture material comprises a light-activated adhesive.
 5. The method of claim 1, wherein the feature capture block is arranged such that the capture surface contacts with the feature array on the substrate.
 6. The method of claim 1, wherein the feature capture block is arranged such that the capture surface is spaced apart from the feature array on the substrate.
 7. The method of claim 1, wherein the light comprises a laser of a predetermined wavelength, and wherein the light comprises an ultraviolet (UV) laser.
 8. The method of claim 1, wherein the feature array comprises a spatially-barcoded bead array having a plurality of beads being uniform in diameter and arranged in a monolayer, and wherein the plurality of beads corresponds to the plurality of features.
 9. The method of claim 1, wherein the feature array is attached to the substrate via cleavable linkers.
 10. The method of claim 9, wherein the cleavable linkers comprise one or more photo-sensitive chemical bonds.
 11. The method of claim 10, wherein the one or more photo-sensitive chemical bonds comprise at least one chemical bond selected from the group consisting of: 3-amino-3-(2-nitrophenyl) propionic acid (ANP), phenacyl ester derivatives, 8-quinolinyln benzenesulfonate, dicoumarin, 6-bromo-7-alkixycoumarin-4-ylmethoxycarbonyl, a bimanane-based linker, and a bis-arylhydrazone based linker.
 12. The method of claim 1, wherein identifying the region of interest on the feature array comprises determining that the one or more features of the plurality of features of the feature array contains the at least one element of the analytes, the intermediate agents, or the combination thereof.
 13. The method of claim 1, wherein identifying the region of interest on the feature array comprises identifying, using a machine learning algorithm, the region of interest on the feature array.
 14. The method of claim 1, wherein retrieving the at least one feature from the capture surface of the feature capture block comprises collecting the at least one feature within a reservoir containing a liquid.
 15. The method of claim 1, wherein identifying the region of interest on the feature array comprises: providing an image of the feature array to a user; and receiving a selection by the user indicative of the region of interest.
 16. The method of claim 1, wherein identifying the region of interest on the feature array comprises selecting the region of interest based on a stained image of the feature array.
 17. The method of claim 1, wherein emitting the light toward the portion of the capture surface of the feature capture block comprises emitting light onto the capture surface through the feature capture block.
-