



US 20250280833A1

(19) **United States**

(12) **Patent Application Publication**  
**McFatrigh et al.**

(10) **Pub. No.: US 2025/0280833 A1**

(43) **Pub. Date: Sep. 11, 2025**

(54) **METHODS AND COMPOSITIONS FOR  
IMPROVING SOYBEAN YIELD**

(60) Provisional application No. 62/884,423, filed on Aug.  
8, 2019.

(71) Applicant: **NewLeaf Symbiotics, Inc.**, St. Louis,  
MO (US)

**Publication Classification**

(72) Inventors: **Charles Michael McFatrigh**, St. Louis,  
MO (US); **Janne Kerovuo**, St. Louis,  
MO (US)

(51) **Int. Cl.**  
**A01N 63/20** (2020.01)  
**C12R 1/01** (2006.01)

(52) **U.S. Cl.**  
CPC ..... **A01N 63/20** (2020.01); **C12R 2001/01**  
(2021.05)

(21) Appl. No.: **19/028,044**

(22) Filed: **Jan. 17, 2025**

**Related U.S. Application Data**

(63) Continuation of application No. 17/633,600, filed on  
Feb. 8, 2022, filed as application No. PCT/US2020/  
034235 on May 22, 2020.

(57) **ABSTRACT**

The present disclosure provides yield enhancing composi-  
tions comprising *Methylobacterium*, methods for improving  
soybean yield, and methods of making the compositions.  
Also provided are isolated soybean yield enhancing *Meth-*  
*ylobacterium*.

**Specification includes a Sequence Listing.**

## METHODS AND COMPOSITIONS FOR IMPROVING SOYBEAN YIELD

### CROSS-REFERENCE TO RELATED APPLICATIONS

**[0001]** This patent application is a continuation application of U.S. patent application Ser. No. 17/633,600 filed Feb. 8, 2022, which is a 35 U.S.C. § 371 application of International Patent Application No. PCT/US2020/034235 filed May 22, 2020, which claims the benefit of U.S. Patent Application No. 62/884,423, filed Aug. 8, 2019, each of which is incorporated herein by reference in its entirety.

### SEQUENCE LISTING STATEMENT

**[0002]** A sequence listing containing the file named “NLSYM9971USN2 Sequence Listing.xml” which is 97,129 bytes (measured in MS-Windows®) and created on Mar. 31, 2025, contains 79 nucleotide sequences, is provided herewith via the USPTO’s EFS system, and is incorporated herein by reference in its entirety.

### BACKGROUND

**[0003]** One-carbon organic compounds such as methane and methanol are found extensively in nature, and are utilized as carbon sources by bacteria classified as methanotrophs and methylotrophs. Methanotrophic bacteria include species in the genera *Methylobacter*, *Methylomonas*, *Methylomicrobium*, *Methylococcus*, *Methylosinus*, *Methylocystis*, *Methylosphaera*, *Methylocaldum*, and *Methylocella* (Lidstrom, 2006). Methanotrophs possess the enzyme methane monooxygenase, that incorporates an atom of oxygen from O<sub>2</sub> into methane, forming methanol. All methanotrophs are obligate one-carbon utilizers that are unable to use compounds containing carbon-carbon bonds. Methylotrophs, on the other hand, can also utilize more complex organic compounds, such as organic acids, higher alcohols, sugars, and the like. Thus, methylotrophic bacteria are facultative methylotrophs. Methylotrophic bacteria include species in the genera *Methylobacterium*, *Hyphomicrobium*, *Methylophilus*, *Methylobacillus*, *Methylophaga*, *Aminobacter*, *Methylorhabdus*, *Methylopilula*, *Methylosulfonomonas*, *Marinosulfonomonas*, *Paracoccus*, *Xanthobacter*, *Ancylobacter* (also known as *Microcycylus*), *Thiobacillus*, *Rhodopseudomonas*, *Rhodobacter*, *Acetobacter*, *Bacillus*, *Mycobacterium*, *Arthobacter*, and *Nocardia* (Lidstrom, 2006).

**[0004]** Most methylotrophic bacteria of the genus *Methylobacterium* are pink-pigmented. They are conventionally referred to as PPFM bacteria, being pink-pigmented facultative methylotrophs. Green (2005, 2006) identified twelve validated species in the genus *Methylobacterium*, specifically *M. aminovorans*, *M. chloromethanicum*, *M. dichloromethanicum*, *M. extorquens*, *M. fujisawaense*, *M. mesophilicum*, *M. organophilum*, *M. radiotolerans*, *M. rhodesianum*, *M. rhodinum*, *M. thiocyanatum*, and *M. zemannii*. However, *M. nidulans* is a nitrogen-fixing *Methylobacterium* that is not a PPFM (Sy et al., 2001). *Methylobacterium* are found in soil, dust, fresh water, sediments, and leaf surfaces, as well as in industrial and clinical environments (Green, 2006).

### SUMMARY

**[0005]** Methods for improving soybean plant yield that comprise: (a) applying a composition to a soybean plant or a part thereof wherein the composition comprises (i) one or more of *Methylobacterium* NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), NLS0610 (ISO26), or variants thereof, or (ii) a combination of *Methylobacterium* isolate NLS0109 (NRRL B-67340) or a variant thereof, and *Methylobacterium* isolate NLS0017 (B-50931) or NLS0610 (ISO26), or variants thereof; wherein said composition further comprises at least one additional component selected from the group consisting of an additional active ingredient, an agriculturally acceptable adjuvant, and an agriculturally acceptable excipient; and, (b) growing the soybean plant to maturity, thereby improving yield of the soybean plant, are provided herein. In certain embodiments of the methods, the composition is applied to a soybean seed. In certain embodiments of the aforementioned methods, the composition comprises a solid substance with the *Methylobacterium* grown thereon and adhered thereto, or an emulsion having the *Methylobacterium* grown therein. In certain embodiments of any of the aforementioned methods, the composition comprises the *Methylobacterium* at a titer of about 1×10<sup>6</sup> CFU/gm to about 1×10<sup>14</sup> CFU/gm for a solid composition or at a titer of about 1×10<sup>6</sup> CFU/mL to about 1×10<sup>11</sup> CFU/mL for a liquid composition. In certain embodiments of the aforementioned methods, the composition comprises NLS0934 (NRRL B-67341), NLS1181 (ISO25), or a variant thereof. In certain embodiments of any of the aforementioned methods, the *Methylobacterium* variant is glyphosate resistant or glufosinate resistant. In certain embodiments of the aforementioned methods, the applied composition coats or partially coats the soybean plant or a part thereof. In certain embodiments of the aforementioned methods, the composition is applied to foliage of the soybean plant. In certain embodiments of the aforementioned methods, the composition further comprises a fungicide. In certain embodiments of the aforementioned methods, the composition is applied at about the VE to about the R6 stage of development. In certain embodiments of the aforementioned methods, the composition is applied at about the V2 to about the V3 stage of development, about the V3 to V4, or V3 stage of development. In certain embodiments of the aforementioned methods, the methods further comprise the step of harvesting seed from the mature soybean plant. In certain embodiments of the aforementioned methods, yield of harvested seed is increased in comparison to yield of harvested seed obtained from a control soybean plant that did not receive an application of the *Methylobacterium*. In certain embodiments of the aforementioned methods, the composition is applied by spraying, coating, partially coating, immersing, and/or imbibing the soybean plant or plant part with the composition. In certain embodiments of the aforementioned methods, the applied composition coats or partially coats the soybean plant or a part thereof, wherein partial coating includes coating at least 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 98%, 99%, or about 99.5% of the surface area of the soybean plant or a part thereof. In certain embodiments of the aforementioned methods, the soybean plant part is a seed. In certain embodiments of the aforementioned methods, the composition comprises one or more of *Methylobacterium* isolate NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693

(ISO23), NLS1179 (ISO24), NLS1181 (ISO25), NLS0610 (ISO26), or a combination of *Methylobacterium* isolates NLS0109 (NRRL B-67340), and NLS0017 (B-50931) or NLS0610 (ISO26). In certain embodiments of the aforementioned methods, the composition further comprises an additional active ingredient. In certain embodiments of the aforementioned methods, the additional active ingredient is selected from the group consisting of a fungicide, insecticide, nematocide, and biological. In certain embodiments of the aforementioned methods, the biological is a biocontrol agent. In certain embodiments of the aforementioned methods, the additional active ingredient is selected from the group consisting of clothianidin, *Bacillus firmus*, abamectin, thiamethoxam, imidacloprid, azoxystrobin, fluopyram, fluoxastrobin, ipconazole, mefenoxam, metalaxyl, penflufen, prothioconazole, pyraclostrobin, and sedaxane. In certain embodiments of the aforementioned methods, the composition comprises (i) a *Methylobacterium* wherein the chromosomal genomic DNA has at least 99%, 99.9, 99.8, 99.7, 99.6%, or 99.5% sequence identity to chromosomal genomic DNA of one or more of NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), or NLS0610 (ISO26), or (ii) a combination of *Methylobacterium* isolates wherein the chromosomal genomic DNA of a first isolate has at least 99%, 99.9, 99.8, 99.7, 99.6%, or 99.5% sequence identity to chromosomal genomic DNA of NLS0109 (NRRL B-67340), and chromosomal DNA of a second isolate has at least 99%, 99.9, 99.8, 99.7, 99.6%, or 99.5% sequence identity to chromosomal genomic DNA of NLS0017 (B-50931) or NLS0610 (ISO26). In certain embodiments of the aforementioned methods, the composition comprises (i) a *Methylobacterium* wherein the assembled genome DNA sequence of the *Methylobacterium* has an average nucleotide identity (ANI) score of at least 99.00 when compared to the assembled genome DNA sequence of NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), or NLS0610 (ISO26), or (ii) a combination of *Methylobacterium* isolates wherein the ANI score of the assembled genome of the first isolate has an average nucleotide identity (ANI) score of at least 99.00 when compared to the assembled genome DNA sequence of NLS0109 (NRRL B-67340), and assembled genome DNA sequence of a second isolate has an average nucleotide identity (ANI) score of at least 99.00 when compared to the assembled genome DNA sequence of NLS0017 (B-50931) or NLS0610 (ISO26). In certain embodiments of the aforementioned methods where the composition comprises one or more of NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), NLS0610 (ISO26), or variants thereof, the composition further comprises an additional *Methylobacterium* selected from the group consisting of ISO01 (NRRL B-50929), ISO02 (NRRL B-50930), ISO03 (NRRL B-50931), ISO04 (NRRL B-50932), ISO05 (NRRL B-50933), ISO06 (NRRL B-50934), ISO07 (NRRL B-50935), ISO08 (NRRL B-50936), ISO09 (NRRL B-50937), ISO10 (NRRL B-50938), ISO11 (NRRL B-50939), ISO12 (NRRL B-50940), ISO13 (NRRL B-50941), ISO14 (NRRL B-50942) and ISO16 (NRRL B-67340).

**[0006]** Also provided herein are soybean plants or soybean plant parts that are coated or partially coated with a composition comprising a *Methylobacterium*, wherein the *Methylobacterium* is one or more of (i) NLS0934 (NRRL

B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), NLS0610 (ISO26), or variants thereof, or (ii) a combination of *Methylobacterium* isolate NLS0109 (NRRL B-67340) or a variant thereof, and *Methylobacterium* isolate NLS0017 (B-50931) or a variant thereof. In certain embodiments, the composition further comprises at least one additional component selected from the group consisting of an additional active ingredient, an agriculturally acceptable adjuvant, and an agriculturally acceptable excipient. In certain aforementioned embodiments, the composition comprises the *Methylobacterium* at a titer of about  $1 \times 10^6$  CFU/gm to about  $1 \times 10^{14}$  CFU/gm for a solid composition or at a titer of about  $1 \times 10^6$  CFU/mL to about  $1 \times 10^{11}$  CFU/mL for a liquid composition. In certain aforementioned embodiments, the *Methylobacterium* is NLS0934 (NRRL B-67341) or NLS1181 (ISO25). In certain embodiments of any of the aforementioned compositions, the soybean plant part is selected from the group consisting of a seed, a stem, a flower, a leaf, a petiole, a pod, and an axillary bud. In certain aforementioned embodiments, the composition comprises one or more of *Methylobacterium* NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), NLS0610 (ISO26), or a combination of *Methylobacterium* isolates NLS0109 (NRRL B-67340), and NLS0017 (B-50931) or NLS0610. In certain aforementioned embodiments, the composition comprises an additional active ingredient. In certain aforementioned embodiments, the additional active ingredient is selected from the group consisting of a fungicide, insecticide, nematocide, and biological. In certain aforementioned embodiments, the biological is a biocontrol agent. In certain aforementioned embodiments, the additional active ingredient is selected from the group consisting of clothianidin, *Bacillus firmus*, abamectin, thiamethoxam, imidacloprid, azoxystrobin, fluopyram, fluoxastrobin, ipconazole, mefenoxam, metalaxyl, penflufen, prothioconazole, pyraclostrobin, and sedaxane. In certain aforementioned embodiments, the composition comprises one or more *Methylobacterium* wherein the chromosomal genomic DNA has at least 99%, 99.9, 99.8, 99.7, 99.6%, or 99.5% sequence identity to chromosomal genomic DNA of NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), or NLS0610 (ISO26), or (ii) a combination of *Methylobacterium* isolates wherein the chromosomal genomic DNA of a first isolate has at least 99%, 99.9, 99.8, 99.7, 99.6%, or 99.5% sequence identity to chromosomal genomic DNA of NLS0109 (NRRL B-67340), and wherein the chromosomal genomic DNA of a second isolate has at least 99%, 99.9, 99.8, 99.7, 99.6%, or 99.5% sequence identity to chromosomal genomic DNA of NLS0017 (B-50931) or NLS0610 (ISO26). In certain embodiments, the composition comprises (i) a *Methylobacterium* having genomic DNA comprising one or more polynucleotide marker fragments of at least 50, 60, 100, 120, 180, 200, 240, or 300 nucleotides of SEQ ID NOS: 1-15. In certain of the aforementioned embodiments, the composition further comprises an additional *Methylobacterium* selected from the group consisting of ISO01 (NRRL B-50929), ISO02 (NRRL B-50930), ISO03 (NRRL B-50931), ISO04 (NRRL B-50932), ISO05 (NRRL B-50933), ISO06 (NRRL B-50934), ISO07 (NRRL B-50935), ISO08 (NRRL B-50936), ISO09 (NRRL B-50937), ISO10 (NRRL B-50938), ISO11 (NRRL B-50939), ISO12 (NRRL B-50940), ISO13 (NRRL

B-50941), and ISO14 (NRRL B-50942); or a *Methylobacterium* having genomic DNA comprising one or more polynucleotide marker fragments of at least 50, 60, 100, 120, 180, 200, 240, or 300 nucleotides of SEQ ID NOS: 61-79. [0007] Also provided herein are isolated *Methylobacterium* strains NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), NLS0610 (ISO26), or variants thereof. Compositions comprising one or more of *Methylobacterium* NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), NLS0610 (ISO26), variants thereof are also provided. In certain embodiments, the *Methylobacterium* strains or compositions coat or partially coat a soybean plant or a part thereof. In certain embodiments, the composition coats or partially coats the soybean seed. In certain embodiments, compositions that coat or partially coat a soybean plant, soybean seed, or other soybean plant part, comprise one or more of NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), NLS0610 (ISO26), or variants thereof.

## DESCRIPTION

### Definitions

[0008] The term “and/or” where used herein is to be taken as specific disclosure of each of the two specified features or components with or without the other. Thus, the term “and/or” as used in a phrase such as “A and/or B” herein is intended to include “A and B,” “A or B,” “A” (alone), and “B” (alone). Likewise, the term “and/or” as used in a phrase such as “A, B, and/or C” is intended to encompass each of the following embodiments: A, B, and C; A, B, or C; A or C; A or B; B or C; A and C; A and B; B and C; A (alone); B (alone); and C (alone).

[0009] Where a term is provided in the singular, embodiments comprising the plural of that term are also provided.

[0010] As used herein, the terms “include,” “includes,” and “including” are to be construed as at least having the features or encompassing the items to which they refer while not excluding any additional unspecified features or unspecified items.

[0011] As used herein, the phrases “adhered thereto” and “adherent” refer to *Methylobacterium* that are associated with a solid substance by growing, or having been grown, on a solid substance.

[0012] As used herein, the phrase “active ingredient” refers to a biological or pesticide in a composition for treatment of plants and/or plant parts.

[0013] As used herein, the term “biological” refers to a component of a composition for treatment of plants or plant parts comprised of or derived from a microorganism. Biologicals include biocontrol agents, other beneficial microorganisms, microbial extracts, natural products, plant growth activators or plant defense agents. Non-limiting examples of biocontrol agents include bacteria, fungi, beneficial nematodes, and viruses. In certain compositions, a biological can comprise a mono-culture or co-culture of *Methylobacterium*, or a combination of *Methylobacterium* strains or isolates that have been separately cultured.

[0014] As used herein, the phrase “agriculturally acceptable adjuvant” refers to a substance that enhances the performance of a biological or pesticide in a composition for treatment of plants and/or plant parts. In certain compositions, a biological can comprise a mono-culture or co-culture of *Methylobacterium*.

[0015] As used herein, the phrase “agriculturally acceptable excipient” refers to an essentially inert substance that can be used as a diluent and/or carrier for a biological or pesticide in a composition for treatment of plants and/or plant parts. In certain compositions, a biological can comprise a mono-culture or co-culture of *Methylobacterium*.

[0016] As used herein, the term “strain” shall include all isolates of such strain.

[0017] As used herein, the phrase “control plant” refers to a plant that had not received treatment with a yield enhancing *Methylobacterium* or composition comprising the same at either the seed or any subsequent stage of the control plant’s development. In certain embodiments, a control plant can be a plant that was treated with an additional active ingredient or a yield neutral *Methylobacterium*.

[0018] As used herein, the phrase “co-culture of *Methylobacterium*” refers to a *Methylobacterium* culture comprising at least two strains of *Methylobacterium* or at least two species of *Methylobacterium*.

[0019] As used herein, the phrase “contaminating microorganism” refers to microorganisms in a culture, fermentation broth, fermentation broth product, or composition that were not identified prior to introduction into the culture, fermentation broth, fermentation broth product, or composition.

[0020] As used herein, “variant” when used in the context of a *Methylobacterium* isolate, refers to any isolate that has chromosomal genomic DNA with at least 99%, 99.9, 99.8, 99.7, 99.6%, or 99.5% sequence identity to chromosomal genomic DNA of a deposited *Methylobacterium* isolate provided herein. A variant of an isolate can be obtained from various sources including soil, plants or plant material and water, particularly water associated with plants and/or agriculture. Variants include derivatives obtained from deposited isolates. *Methylobacterium* isolates or strains can be sequenced (for example as taught by Sanger et al. (1977), Bentley et al. (2008) or Caporaso et al. (2012)) and genome-scale comparison of the sequences conducted (Konstantinidis et al. (2005)) using sequence analysis tools, such as BLAST, as taught by Altschul et al. (1990) or clustalw ([www.cbi.ac.uk/Tools/msa/clustalw2/](http://www.cbi.ac.uk/Tools/msa/clustalw2/)).

[0021] As used herein, “derivative” when used in the context of a *Methylobacterium* isolate, refers to any *Methylobacterium* that is obtained from a deposited *Methylobacterium* isolate provided herein. Derivatives of a *Methylobacterium* isolate include, but are not limited to, derivatives obtained by selection, derivatives obtained by mutagenesis and selection, and genetically transformed *Methylobacterium* obtained from the *Methylobacterium* isolate. A “derivative” can be identified, for example based on genetic identity to the strain or isolate from which it was obtained and will generally exhibit chromosomal genomic DNA with at least 99%, 99.9, 99.8, 99.7, 99.6%, or 99.5% sequence identity to chromosomal genomic DNA of the strain or isolate from which it was derived.

[0022] As used herein, “sequence identity” when used to evaluate whether a particular *Methylobacterium* strain is a variant or derivative of a *Methylobacterium* strain provided herein refers to a measure of nucleotide-level genomic similarity between the coding regions of two genomes. Sequence identity between the coding regions of bacterial genomes can be calculated, for example, by determining the Average Nucleotide Identity (ANI) score using FastANI (Jain et al. “High throughput ANI analysis of 90K prokary-

otic genomes reveals clear species boundaries”, Nat Communications 9, 5114 (2018)) and Han et al. (“ANI tools web: a web tool for fast genome comparison within multiple bacterial strains”; Database, 2016, 1-5).

**[0023]** As used herein, the term “emulsion” refers to a colloidal mixture of two immiscible liquids wherein one liquid is the continuous phase and the other liquid is the dispersed phase. In certain embodiments, the continuous phase is an aqueous liquid and the dispersed phase is liquid that is not miscible, or partially miscible, in the aqueous liquid.

**[0024]** As used herein, the phrase “essentially free of contaminating microorganisms” refers to a culture, fermentation broth, fermentation product, or composition where at least about 95% of the microorganisms present by amount or type in the culture, fermentation broth, fermentation product, or composition are the desired *Methylobacterium* or other desired microorganisms of pre-determined identity.

**[0025]** As used herein, the phrase “inanimate solid substance” refers to a substance which is insoluble or partially soluble in water or aqueous solutions and which is either non-living or which is not a part of a still-living organism from which it was derived.

**[0026]** As used herein, the phrase “mono-culture of *Methylobacterium*” refers to a *Methylobacterium* culture consisting of a single strain of *Methylobacterium*.

**[0027]** As used herein, a “pesticide” refers to an insecticide, fungicide, nematocide, bactericide, or any combination thereof.

**[0028]** As used herein, the phrase “bacteriostatic agent” refers to agents that inhibit growth of bacteria but do not kill the bacteria.

**[0029]** As used herein, the phrase “pesticide does not substantially inhibit growth of said *Methylobacterium*” refers to any pesticide that when provided in a composition comprising a fermentation product comprising a solid substance wherein a mono-culture or co-culture of *Methylobacterium* is adhered thereto, results in no more than a 50% inhibition of *Methylobacterium* growth when the composition is applied to a plant or plant part in comparison to a composition lacking the pesticide. In certain embodiments, the pesticide results in no more than a 40%, 20%, 10%, 5%, or 1% inhibition of *Methylobacterium* growth when the composition is applied to a plant or plant part in comparison to a composition lacking the pesticide.

**[0030]** As used herein, the term “*Methylobacterium*” refers to genera and species in the methylobacteriaceae family, including bacterial species in the *Methylobacterium* genus and proposed *Methylorubrum* genus (Green and Ardley (2018)). *Methylobacterium* includes pink-pigmented facultative methylotrophic bacteria (PPFM) and also encompasses the non-pink-pigmented *Methylobacterium nodulans*, as well as colorless mutants of *Methylobacterium* isolates. For example, and not by way of limitation, “*Methylobacterium*” refers to bacteria of the species listed below as well as any new *Methylobacterium* species that have not yet been reported or described that can be characterized as *Methylobacterium* or *Methylorubrum* based on phylogenetic analysis: *Methylobacterium adhaesivum*; *Methylobacterium oryzae*; *Methylobacterium aerolatum*; *Methylobacterium oxalidis*; *Methylobacterium aquaticum*; *Methylobacterium persicinum*; *Methylobacterium brachiatum*; *Methylobacterium phyllosphaerae*; *Methylobacterium brachythecii*; *Methylobacterium phyllostachyos*; *Methylobacterium bulla-*

*tum*; *Methylobacterium platani*; *Methylobacterium cerastii*; *Methylobacterium pseudosasicola*; *Methylobacterium currus*; *Methylobacterium radiotolerans*; *Methylobacterium dankookense*; *Methylobacterium soli*; *Methylobacterium frigidaeris*; *Methylobacterium specialis*; *Methylobacterium fujisawaense*; *Methylobacterium tardum*; *Methylobacterium gnaphalii*; *Methylobacterium tarhaniae*; *Methylobacterium goesingense*; *Methylobacterium thuringiense*; *Methylobacterium gossypicola*; *Methylobacterium trifolii*; *Methylobacterium gregans*; *Methylobacterium variabile*; *Methylobacterium haplocladii*; *Methylobacterium aminovorans* (*Methylorubrum aminovorans*); *Methylobacterium hispanicum*; *Methylobacterium extorquens* (*Methylorubrum extorquens*); *Methylobacterium indicum*; *Methylobacterium podarium* (*Methylorubrum podarium*); *Methylobacterium iners*; *Methylobacterium populi* (*Methylorubrum populi*); *Methylobacterium isbiliense*; *Methylobacterium pseudosasa* (*Methylorubrum pseudosasa*); *Methylobacterium jeotgali*; *Methylobacterium rhodesianum* (*Methylorubrum rhodesianum*); *Methylobacterium komagatae*; *Methylobacterium rhodinum* (*Methylorubrum rhodinum*); *Methylobacterium longum*; *Methylobacterium salsuginis* (*Methylorubrum salsuginis*); *Methylobacterium marchantiae*; *Methylobacterium suomiense* (*Methylorubrum suomiense*); *Methylobacterium mesophilicum*; *Methylobacterium thiocyanatum* (*Methylorubrum thiocyanatum*); *Methylobacterium nodulans*; *Methylobacterium zatmanii* (*Methylorubrum zatmanii*); *Methylobacterium organophilum*.

**[0031]** As used herein, the phrase “solid substance” refers to a substance which is insoluble or partially soluble in water or aqueous solutions.

**[0032]** As used herein, the phrase “solid phase that can be suspended therein” refers to a solid substance that can be distributed throughout a liquid by agitation.

**[0033]** As used herein, the term “non-regenerable” refers to either a plant part or processed plant product that cannot be regenerated into a whole plant.

**[0034]** As used herein, the phrase “substantially all of the solid phase is suspended in the liquid phase” refers to media wherein at least 95%, 98%, or 99% of solid substance(s) comprising the solid phase are distributed throughout the liquid by agitation.

**[0035]** As used herein, the phrase “substantially all of the solid phase is not suspended in the liquid phase” refers to media where less than 5%, 2%, or 1% of the solid is in a particulate form that is distributed throughout the media by agitation.

**[0036]** To the extent to which any of the preceding definitions is inconsistent with definitions provided in any patent or non-patent reference incorporated herein by reference, any patent or non-patent reference cited herein, or in any patent or non-patent reference found elsewhere, it is understood that the preceding definition will be used herein.

Yield Enhancing *Methylobacterium*, Compositions Comprising Yield Enhancing *Methylobacterium*, Methods of their Use, and Methods of Making

**[0037]** Various yield enhancing *Methylobacterium* isolates, compositions comprising these *Methylobacterium*, methods of using the compositions to improve soybean plant yield, and methods of making the compositions are provided herein. In certain embodiments, yield enhancing *Methylobacterium* isolates are selected from NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), NLS0610 (ISO26), and vari-

ants thereof, and a combination of *Methylobacterium* isolate NLS0109 (NRRL B-67340) or a variant thereof, and *Methylobacterium* isolate NLS0017 (B-50931) or NLS0610 (ISO26) or variants thereof. Amounts of the compositions that comprise yield enhancing *Methylobacterium* sufficient to provide for improved soybean plant yield can be determined by measuring any or all of changes in yield relative to untreated plants or plant parts. In certain embodiments, yield can be assessed by measuring output of seed on a per unit area basis (i.e. bushels per acre, kilograms per hectare, and the like), where the yield enhancing *Methylobacterium* treated plants or plants grown from *Methylobacterium* treated seed are grown at about the same density as the control plants. In certain embodiments, yield can be assessed by measuring output on a per plant or per plant part basis (grams of seed per plant, grams of seed per pod, pods per plant, seeds per pod and the like) of the yield enhancing *Methylobacterium* treated plants in comparison to untreated control plants.

**[0038]** Isolated yield enhancing *Methylobacterium* are provided herein. In certain embodiments, the *Methylobacterium* is selected from the group consisting of *M. gregans*, *M. dankookense*, *M. radiotolerans*, *M. komagatae*, and *M. bullatum*. In certain embodiments, the yield enhancing *Methylobacterium* isolate is selected from the group consisting of NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), NLS0610 (ISO26), and variants thereof, and combinations thereof. In certain embodiments, the yield enhancing *Methylobacterium* isolate is selected from the group consisting of NLS0934 (NRRL B-67341), NLS1181 (ISO25), and variants thereof. In certain embodiments, the yield enhancing *Methylobacterium* is a combination of *Methylobacterium* isolates NLS0109, and NLS0017 (NRRL B-50931) or NLS0610 (ISO26). In certain embodiments, the yield enhancing *Methylobacterium* isolate can enhance yield when applied to a soybean seed, when applied in vegetative stages of soybean development, or when applied during reproductive stages of soybean development. In certain embodiments, the yield enhancing *Methylobacterium* has a chromosomal genomic DNA with at least 99%, 99.9, 99.8, 99.7, 99.6%, or 99.5% sequence identity to chromosomal genomic DNA of NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), or NLS0610 (ISO26)); or the yield enhancing *Methylobacterium* is a combination of *Methylobacterium* variants of NLS0109 having chromosomal genomic DNA with at least 99%, 99.9, 99.8, 99.7, 99.6%, or 99.5% sequence identity to chromosomal genomic DNA of NLS0109 (NRRL B-67340), and variants of NLS0017 (NRRL B-50931) or NLS0610 (ISO26) variants having chromosomal genomic DNA with at least 99%, 99.9, 99.8, 99.7, 99.6%, or 99.5% sequence identity to chromosomal genomic DNA of NLS0017 (NRRL B-50931) or NLS0610 (ISO26). In certain embodiments, the yield enhancing *Methylobacterium* provides for at least about 2%, at least about 3%, at least about 4%, at least about 5%, at least about 6%, at least about 7%, at least about 8%, at least about 9%, at least about 10%, or at least about 15% increases in yield of a treated plant or a plant arising from a treated seed in comparison to untreated control plants or plants grown from untreated seeds. In certain embodiments, the yield enhancing *Methylobacterium* provides for at least about 2% or at least about 5% to at least about a 10% or at least about a 20%

increase in yield of a treated plant or a plant grown from a treated seed in comparison to untreated control plants or plants arising from untreated seeds.

**[0039]** In certain embodiments, the yield enhancing *Methylobacterium* provides for increased yield when applied to a seed. In certain embodiments, the yield enhancing *Methylobacterium* provides for increases in yield when applied during soybean vegetative stages of development. In certain embodiments, the yield enhancing *Methylobacterium* provides for increased yield when applied just prior to or during soybean reproductive stages of development. In some embodiments, a *Methylobacterium* is applied more than once, wherein such applications can be to a soybean seed and a soybean plant at one or more stages of development, or can be more than one application to a soybean plant at two or more developmental stages. In certain embodiments of any of the aforementioned compositions, the composition comprises a solid substance wherein a mono-culture or co-culture of *Methylobacterium* is adhered thereto. In certain embodiments where the *Methylobacterium* is adhered to a solid substance, the composition comprises a colloid formed by the solid substance wherein a mono-culture or co-culture of *Methylobacterium* is adhered thereto and a liquid. In certain embodiments, the colloid is a gel. In certain embodiments of certain aforementioned compositions, composition is an emulsion that does not contain a solid substance. In certain embodiments of any of the aforementioned compositions, the yield enhancing *Methylobacterium* is selected from the group consisting of NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), NLS0610 (ISO26), and variants thereof. In certain embodiments of any of the aforementioned compositions, the yield enhancing *Methylobacterium* is a combination of *Methylobacterium* NLS0109, and NLS0017 or NLS0610. In certain embodiments, the yield enhancing *Methylobacterium* has a chromosomal genomic DNA with at least 99%, 99.9, 99.8, 99.7, 99.6%, or 99.5% sequence identity to chromosomal genomic DNA of NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), or NLS0610 (ISO26); or the yield enhancing *Methylobacterium* is a combination of *Methylobacterium* variants of NLS0109, and NLS0017 or NLS0610 having chromosomal genomic DNA with at least 99%, 99.9, 99.8, 99.7, 99.6%, or 99.5% sequence identity to chromosomal genomic DNA of NLS0109 (NRRL B-67340), NLS0017 (NRRL B-50931) or NLS0610.

**[0040]** In certain embodiments, isolated yield enhancing *Methylobacterium* can be identified by treating a plant, a seed, soil in which the plant or a plant arising from the seed are grown, or other plant growth media in which the plant or a plant arising from the seed are grown and assaying for increased yield.

**[0041]** In certain embodiments, soybean seed or soybeans in the vegetative stages of development are treated with the yield enhancing *Methylobacterium*. The vegetative stages of soybean are as follows: VE (emergence), VC (cotyledon stage), V1 (first trifoliate leaf), V2 (second trifoliate leaf), V3 (third trifoliate leaf), V4 (fourth trifoliate leaf), up to V (n) (nth trifoliate leaf, where the final number of trifoliate leaves depend on the soybean variety and environmental conditions). A description of the soybean vegetative stages can be found on the world wide web (internet) at [extension.agron.iastate.edu/soybean/production\\_growth-](http://extension.agron.iastate.edu/soybean/production_growth-)

stages.html and in “Soybean Growth and Development”, Pedersen, P., Iowa State University Extension and Outreach publication PM 1945 December 2009). In certain embodiments, the yield enhancing *Methylobacterium* are applied at about the VE to about the V4, V5, V6, or Vn stage of development, where n is the number of trifoliolate leaves present immediately prior to entering the reproductive stages of development. In certain embodiments, the yield enhancing *Methylobacterium* are applied at about the VC, V1, V2, or V3 to about the V4, V5, V6, or Vn stage of development, where n is the number of trifoliolate leaves present immediately prior to entering the reproductive stages of development. In certain embodiments, the yield enhancing *Methylobacterium* that is applied to the seed or during the vegetative stage is selected from the group consisting of NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), NLS0610 (ISO26), and variants thereof.

**[0042]** In certain embodiments, the yield enhancing *Methylobacterium* are applied to a transgenic soybean plant that is herbicide, insect or disease tolerant. In certain embodiments, the yield enhancing *Methylobacterium* are applied before, during, or after the application of glyphosate to a transgenic soybean plant that is glyphosate tolerant. Commercially available glyphosate formulations that can be used include, but are not limited to, Roundup Original MAX®, Roundup PowerMAX®, Roundup UltraMax®, or Roundup WeatherMAX® (Monsanto Co., St. Louis, MO., USA); Touchdown IQ® or Touchdown Total® (Syngenta, Wilmington, Delaware, USA); Glyphomax®, Glyphomax Plus®, or Glyphomax XRT® (Dow Agrosciences LLC, Indianapolis, IN, USA). Soybean plants are typically sprayed with glyphosate at about the V3-V4 vegetative development stage. In certain embodiments, the yield enhancing *Methylobacterium* that is applied before, during, or after the application of glyphosate is a derivative of NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), or NLS0610 (ISO26)), that is selected for glyphosate resistance. Selections for glyphosate resistant bacteria that have been described (Comai et al., Science 221 (4608): 370-371) can be adapted for selection of yield enhancing *Methylobacterium* that are glyphosate resistant. The selection and use of glyphosate resistant yield enhancing *Methylobacterium* from mutagenized or other populations of *Methylobacterium* such as NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), or NLS0610 (ISO26), and variants thereof, is provided herein.

**[0043]** In certain embodiments, soybean seed or soybeans in the late vegetative stages to reproductive stages of development are treated with the yield enhancing *Methylobacterium*. The late vegetative stages of soybean are the V5 or V6, to V (n) (nth trifoliolate leaf, where the final number of trifoliolate leaves depend on the soybean variety and environmental conditions) stages. The reproductive stages of soybean development are: R1 (beginning flowering—at least one flower on any node); R2 (full flowering—an open flower at one of the two uppermost nodes); R3 (beginning pod-pods are 5 mm at one of the four uppermost nodes); R4 (full pod-pods at 2 cm at one of the four uppermost nodes); R5 (Beginning seed-seed is 3 mm long in the pod at one of the four uppermost nodes on the main stem); R6 (full seed-pod containing a green seed that fills the pod capacity at one of

the four uppermost nodes on the main stem); R7 (beginning maturity—one normal pod on the main stem has reached its mature pod color); and R8 (full maturity—95% of the pods have reached their full mature color. In certain embodiments, the yield enhancing *Methylobacterium* that is applied to late vegetative or reproductive stage is selected from the group consisting of NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), NLS0610 (ISO26), and variants thereof, and a combination of *Methylobacterium* NLS0109, and NLS0017 or NLS0610, and variants thereof.

**[0044]** Various *Methylobacterium* isolates provided herein are disclosed in Table 1.

TABLE 1

| <i>Methylobacterium</i> isolates |             |         |                                |
|----------------------------------|-------------|---------|--------------------------------|
| Deposit Identifier               | ISOLATE No. | NLS No. | USDA ARS NRRL No. <sup>1</sup> |
| <i>Methylobacterium</i> sp. #1   | ISO01       | NLS0046 | NRRL B-50929                   |
| <i>Methylobacterium</i> sp. #2   | ISO02       | NLS0020 | NRRL B-50930                   |
| <i>Methylobacterium</i> sp. #3   | ISO03       | NLS0017 | NRRL B-50931                   |
| <i>Methylobacterium</i> sp. #4   | ISO04       | NLS0042 | NRRL B-50932                   |
| <i>Methylobacterium</i> sp. #5   | ISO05       | NLS0089 | NRRL B-50933                   |
| <i>Methylobacterium</i> sp. #6   | ISO06       | NLS0068 | NRRL B-50934                   |
| <i>Methylobacterium</i> sp. #7   | ISO07       | NLS0065 | NRRL B-50935                   |
| <i>Methylobacterium</i> sp. #8   | ISO08       | NLS0069 | NRRL B-50936                   |
| <i>Methylobacterium</i> sp. #9   | ISO09       | NLS0062 | NRRL B-50937                   |
| <i>Methylobacterium</i> sp. #10  | ISO10       | NLS0064 | NRRL B-50938                   |
| <i>Methylobacterium</i> sp. #11  | ISO11       | NLS0021 | NRRL B-50939                   |
| <i>Methylobacterium</i> sp. #12  | ISO12       | NLS0066 | NRRL B-50940                   |
| <i>Methylobacterium</i> sp. #13  | ISO13       | NLS0037 | NRRL B-50941                   |
| <i>Methylobacterium</i> sp. #14  | ISO14       | NLS0038 | NRRL B-50942                   |
| <i>Methylobacterium</i> #16      | ISO16       | NLS0109 | NRRL B-67340                   |
| <i>Methylobacterium</i> #17      | ISO17       | NLS0934 | NRRL B-67341                   |
| <i>Methylobacterium</i> sp #22   | ISO22       | NLS0497 | NRRL B-67925                   |
| <i>Methylobacterium</i> sp #23   | ISO23       | NLS0693 | NRRL B-67926                   |
| <i>Methylobacterium</i> sp #24   | ISO24       | NLS1179 | NRRL B-67929                   |
| <i>Methylobacterium</i> sp #25   | ISO25       | NLS1181 | NRRL B-67927                   |
| <i>Methylobacterium</i> sp #26   | ISO26       | NLS0610 | NRRL B-67892                   |

<sup>1</sup>Deposit number for strain deposited with the AGRICULTURAL RESEARCH SERVICE CULTURE COLLECTION (NRRL) of the National Center for Agricultural Utilization Research, Agricultural Research Service, U.S. Department of Agriculture, 1815 North University Street, Peoria, Illinois 61604 U.S.A. under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure. Access to the deposited material will be available during pendency of the patent application making reference to it, to one determined by the Director to be entitled to access under 37 CFR § 1.14 and 35 USC § 122 (see In re Lundak, 227 USPQ 90, 94-95 (Fed. Cir. 1985)(citing 35 U.S.C. 114). Subject to 37 CFR § 1.808(b), all restrictions imposed by the depositor on the availability to the public of the deposited material will be irrevocably removed upon the granting of any patent from this patent application.

**[0045]** Samples of the following *Methylobacterium* sp. strains have been deposited with the AGRICULTURAL RESEARCH SERVICE CULTURE COLLECTION (NRRL) of the National Center for Agricultural Utilization Research, Agricultural Research Service, U.S. Department of Agriculture, 1815 North University Street, Peoria, Illinois 61604 U.S.A. under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure. *Methylobacterium* sp. NRRL B-50929, NRRL B-50930, NRRL B-50931, NRRL B-50932, NRRL B-50933, NRRL B-50934, NRRL B-50935, NRRL B-50936, NRRL B-50937, NRRL B-50938, NRRL B-50939, NRRL B-50940, NRRL B-50941 and NRRL B-50942 were deposited with NRRL on Mar. 12, 2014. *Methylobacterium* sp. NRRL B-67340 and NRRL B-67341 were deposited with NRRL on Nov. 18, 2016. *Methylobacterium* sp. *Methylobacterium* sp. NRRL B-67892 was deposited with NRRL on Nov. 26, 2019. *Methylobacterium* sp. NRRL B-67925, NRRL B-67926, and

NRRL B-67927 were deposited with NRRL on Feb. 21, 2020. *Methylobacterium* sp. NRRL B-67929 was deposited with NRRL on Mar. 3, 2020.

**[0046]** Variants of a *Methylobacterium* isolate listed in Table 1 include derivatives obtained therefrom by genetic transformation, mutagenesis and/or insertion of a heterologous sequence. In some embodiments, such variants are identified by the presence of chromosomal genomic DNA with at least 99%, 99.9, 99.8, 99.7, 99.6%, or 99.5% sequence identity to chromosomal genomic DNA of the strain from which it was derived. In certain embodiments, derivatives and other variants are distinguished by the presence of one or more unique DNA sequences that include: (i) a unique sequence of SEQ ID NOs: 1-15 and SEQ ID NOs: 61-79. Specific assays and primer sequences that use such sequences for specific detection of *Methylobacterium* strains from isolated DNA or various samples from treated plants are described in the examples herein, and in WO2020010264.

**[0047]** Co-assigned patents or patent applications that disclose additional specific uses of certain *Methylobacterium* strains of Table 1 such as: increasing corn yield (US20160295868); improving lettuce cultivation (U.S. Pat. No. 10,212,939); improving tomato growth (U.S. Pat. No. 10,368,547); improving soybean yield (US2016/0302423); improving fruit production (U.S. Pat. No. 10,111,438); controlling corn rootworm (US20170238553); controlling root lesion nematodes (U.S. Pat. No. 10,448,645); controlling root knot nematodes (U.S. Pat. No. 10,098,353); and controlling fungal disease (US20180295841 and US20190364905) are each incorporated herein by reference in their entireties.

**[0048]** Also provided herein are methods for improving soybean yield that comprise applying any of the aforementioned compositions provided herein to a plant or a plant part in an amount that provides for increased soybean yield in the plant, plant part, or a plant obtained therefrom relative to infection of a control plant, plant part, or plant obtained therefrom that had not received an application of the composition. In certain embodiments, the plant part is selected from the group consisting of a leaf, a stem, a flower, a root, a pod, a coleoptile and a seed. In certain embodiments, the method further comprises the step of harvesting at least one plant part selected from the group consisting of a leaf, a stem, a flower, a root, a pod, or a seed from the plant or plant part. In certain embodiments of any of the aforementioned methods, the methods further comprise obtaining a processed food or feed composition from the plant or plant part. In certain embodiments, the processed food or feed composition is a meal or a paste. In certain embodiments of any of the aforementioned methods, the yield enhancing *Methylobacterium* is selected from the group consisting of NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), NLS0610 (ISO26), and variants thereof. In certain embodiments of any of the aforementioned compositions, the yield enhancing *Methylobacterium* is a combination of *Methylobacterium* NLS0109, and NLS0017 or NLS0610, or variants thereof.

**[0049]** Also provided are methods of making the compositions useful for improving soybean yield that comprise combining a yield enhancing *Methylobacterium* with an agriculturally acceptable excipient and/or with an agriculturally acceptable adjuvant. In certain embodiments of the methods, the *Methylobacterium* is adhered to a solid sub-

stance. In certain embodiments of the methods, the *Methylobacterium* is adhered to a solid substance, where the solid substance is combined with a liquid to form a composition that is a colloid. In certain embodiments of the methods, the colloid is a gel. In certain embodiments of the methods, *Methylobacterium* adhered to a solid substance is provided by culturing the *Methylobacterium* in the presence of the solid substance. In certain embodiments of the methods, the composition comprises an emulsion. In certain embodiments of the methods, the *Methylobacterium* is provided by culturing the *Methylobacterium* in an emulsion. In certain embodiments of any of the aforementioned methods, the yield enhancing *Methylobacterium* is selected from the group consisting of NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), NLS0610 (ISO26), and variants thereof. In certain embodiments of any of the aforementioned compositions, the yield enhancing *Methylobacterium* is a combination of *Methylobacterium* NLS0109, and NLS0017 or NLS0610, or variants thereof.

**[0050]** Methods where *Methylobacterium* are cultured in biphasic media comprising a liquid phase and a solid substance have been found to significantly increase the resultant yield of *Methylobacterium* relative to methods where the *Methylobacterium* are cultured in liquid media alone. In certain embodiments, the methods can comprise growing the *Methylobacterium* in liquid media with a particulate solid substance that can be suspended in the liquid by agitation under conditions that provide for *Methylobacterium* growth. In certain embodiments where particulate solid substances are used, at least substantially all of the solid phase can thus be suspended in the liquid phase upon agitation. Such particulate solid substances can comprise materials that are about 1 millimeter or less in length or diameter. In certain embodiments, the degree of agitation is sufficient to provide for uniform distribution of the particulate solid substance in the liquid phase and/or optimal levels of culture aeration. However, in other embodiments provided herein, at least substantially all of the solid phase is not suspended in the liquid phase, or portions of the solid phase are suspended in the liquid phase and portions of the solid phase are not suspended in the liquid phase. Non-particulate solid substances can be used in certain biphasic media where the solid phase is not suspended in the liquid phase. Such non-particulate solid substances include, but are not limited to, materials that are greater than about 1 millimeter in length or diameter. Such particulate and non-particulate solid substances also include, but are not limited to, materials that are porous, fibrous, or otherwise configured to provide for increased surface areas for adherent growth of the *Methylobacterium*. Biphasic media where portions of the solid phase are suspended in the liquid phase and portions of the solid phase are not suspended in the liquid phase can comprise a mixture of particulate and non-particulate solid substances. Such particulate and non-particulate solid substances used in any of the aforementioned biphasic media also include, but are not limited to, materials that are porous, fibrous, or otherwise configured to provide for increased surface areas for adherent growth of the *Methylobacterium*. In certain embodiments, the media comprises a colloid formed by a solid and a liquid phase. A colloid comprising a solid and a liquid can be pre-formed and added to liquid media or can be formed in media containing a solid and a liquid. Colloids comprising a solid and a liquid can be formed by subjecting



certain solid substances to a chemical and/or thermal change. In certain embodiments, the colloid is a gel. In certain embodiments, the liquid phase of the media is an emulsion. In certain embodiments, the emulsion comprises an aqueous liquid and a liquid that is not miscible, or only partially miscible, in the aqueous liquid. Liquids that are not miscible, or only partially miscible, in water include, but are not limited to, any of the following: (1) liquids having a miscibility in water that is equal to or less than that of pentanol, hexanol, or heptanol at 25 degrees C.; (2) liquids comprising an alcohol, an aldehyde, a ketone, a fatty acid, a phospholipid, or any combination thereof; (3) alcohols selected from the group consisting of aliphatic alcohols containing at least 5 carbons and sterols; (4) an animal oil, microbial oil, synthetic oil, plant oil, or combination thereof; and/or, (5) a plant oil is selected from the group consisting of corn, soybean, cotton, peanut, sunflower, olive, flax, coconut, palm, rapeseed, sesame seed, safflower, and combinations thereof. In certain embodiments, the immiscible or partially immiscible liquid can comprise at least about 0.02% to about 20% of the liquid phase by mass. In certain embodiments, the methods can comprise obtaining a biphasic culture media comprising the liquid, the solid, and *Methylobacterium* and incubating the culture under conditions that provide for growth of the *Methylobacterium*. Biphasic culture medias comprising the liquid, the solid, and *Methylobacterium* can be obtained by a variety of methods that include, but are not limited to, any of: (a) inoculating a biphasic media comprising the liquid and the solid substance with *Methylobacterium*; (b) inoculating the solid substance with *Methylobacterium* and then introducing the solid substance comprising the *Methylobacterium* into the liquid media; (c) inoculating the solid substance with *Methylobacterium*, incubating the *Methylobacterium* on the solid substance, and then introducing the solid substance comprising the *Methylobacterium* into the liquid media; or (d) any combination of (a), (b), or (c). Methods and compositions for growing *Methylobacterium* in biphasic media comprising a liquid and a solid are disclosed in co-assigned U.S. Pat. Nos. 9,181,541 and 9,845,462, which are incorporated herein by reference in their entirety, and in co-assigned International Patent Publication WO2013181610, published Dec. 5, 2013, which is incorporated herein by reference in its entirety.

**[0051]** Methods where *Methylobacterium* are cultured in media comprising an emulsion have also been found to significantly increase the resultant yield of *Methylobacterium* relative to methods where the *Methylobacterium* are cultured in liquid media alone. In certain embodiments, the methods for making the compositions provided herein can comprise growing the yield enhancing *Methylobacterium* in an emulsion under conditions that provide for *Methylobacterium* growth. Medias comprising the emulsion and yield enhancing *Methylobacterium* can be obtained by a variety of methods that include, but are not limited to, any of: (a) inoculating a media comprising the emulsion with *Methylobacterium*; (b) inoculating the aqueous liquid with the *Methylobacterium*, introducing the non-aqueous liquid, and mixing to form an emulsion; (c) inoculating the aqueous liquid with the *Methylobacterium*, introducing the non-aqueous liquid, and mixing to form an emulsion; or (d) any combination of (a), (b), or (c). In certain embodiments, the emulsion comprises an aqueous liquid and a liquid that is not miscible, or only partially miscible, in the aqueous liquid.

Non-aqueous liquids that are not miscible, or only partially miscible, in water include, but are not limited to, any of the following: (1) liquids having a miscibility in water that is equal to or less than that of n-pentanol, n-hexanol, or n-heptanol at 25 degrees C.; (2) liquids comprising an alcohol, an aldehyde, a ketone, a fatty acid, a phospholipid, or any combination thereof; (3) alcohols is selected from the group consisting of aliphatic alcohols containing at least 5, 6, or 7 carbons and sterols; (4) an animal oil, microbial oil, synthetic oil, plant oil, or combination thereof; and/or, (5) a plant oil is selected from the group consisting of corn, soybean, cotton, peanut, sunflower, olive, flax, coconut, palm, rapeseed, sesame seed, safflower, and combinations thereof. In certain embodiments, the immiscible or partially immiscible non-aqueous liquid can comprise at least about 0.02% to about 20% of the emulsion by mass. In certain embodiments, the immiscible or partially immiscible non-aqueous liquid can comprise at least about any of about 0.05%, 0.1%, 0.5%, or 1% to about 3%, 5%, 10%, or 20% of the emulsion by mass. Methods and compositions for growing *Methylobacterium* in media comprising an emulsion are disclosed in co-assigned U.S. Pat. No. 10,287,544 and International Patent Publication WO2014194189, published Dec. 4, 2014, which are incorporated herein by reference in their entirety.

**[0052]** In some embodiments, the composition or method disclosed herein may comprise one or more additional components. In some embodiments an additional component can be an additional active ingredient, for example, a pesticide or a second biological. The pesticide may be, for example, an insecticide, a fungicide, an herbicide, or a nematocide. The second biological can be a biocontrol agent.

**[0053]** Non-limiting examples of insecticides and nematocides include carbamates, diamides, macrocyclic lactones, neonicotinoids, organophosphates, phenylpyrazoles, pyrethrins, spinosyns, synthetic pyrethroids, tetrone and tetramic acids. In particular embodiments insecticides and nematocides include abamectin, aldicarb, aldoxycarb, bifenthrin, carbofuran, chlorantraniliporle, chlothianidin, cyfluthrin, cyhalothrin, cypermethrin, deltamethrin, dinotefuran, emamectin, ethiprole, fenamiphos, fipronil, flubendiamide, fosthiazate, imidacloprid, ivermectin, lambda-cyhalothrin, milbemectin, nitenpyram, oxamyl, permethrin, tioxazafen, spinetoram, spinosad, spirotetramat, spirotetramat, tefluthrin, thiacloprid, thiamethoxam, and thiodicarb.

**[0054]** Non-limiting examples of useful fungicides include aromatic hydrocarbons, benzimidazoles, benzthiadiazole, carboxamides, carboxylic acid amides, morpholines, phenylamides, phosphonates, quinone outside inhibitors (e.g. strobilurins), thiazolidines, thiophanates, thiophene carboxamides, and triazoles. Particular examples of fungicides include acibenzolar-S-methyl, azoxystrobin, benalaxyl, bixafen, boscalid, carbendazim, cyproconazole, dimethomorph, epoxiconazole, fluopyram, fluoxastrobin, flutianil, flutolanil, fluxapyroxad, fosetyl-A1, ipconazole, isopyrazam, kresoxim-methyl, mefenoxam, metalaxyl, metconazole, myclobutanil, orysastrobin, penflufen, penthiopyrad, picoxystrobin, propiconazole, prothioconazole, pyraclostrobin, sedaxane, silthiofam, tebuconazole, thifluzamide, thiophanate, tolclofos-methyl, trifloxystrobin, and triticonazole.

**[0055]** Non-limiting examples of herbicides include ACCase inhibitors, acetanilides, AHAS inhibitors, carotenoid biosynthesis inhibitors, EPSPS inhibitors, glutamine

synthetase inhibitors, PPO inhibitors, PS II inhibitors, and synthetic auxins. Particular examples of herbicides include acetoachlor, clethodim, dicamba, flumioxazin, fomesafen, glyphosate, glufosinate, mesotrione, quizalofop, saflufenacil, sulcotrione, and 2,4-D.

**[0056]** In some embodiments, compositions or methods disclosed herein may comprise an additional active ingredient which may be an additional biological. The additional biological could be a biocontrol agent, other beneficial microorganisms, microbial extracts, natural products, plant growth activators or a plant defense agent. Non-limiting examples of biocontrol agents include bacteria, fungi, beneficial nematodes, and viruses.

**[0057]** In certain embodiments, an additional biological can be *Methylobacterium*. In certain embodiments, the additional biological is a *Methylobacterium* listed in Table 1, or variants thereof. Thus, compositions or methods comprised herein can comprise two or more of NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), NLS0610 (ISO26), or variants thereof. In certain embodiments, an additional biological is selected from the group consisting of ISO01 (NRRL B-50929), ISO02 (NRRL B-50930), ISO03 (NRRL B-50931), ISO04 (NRRL B-50932), ISO05 (NRRL B-50933), ISO06 (NRRL B-50934), ISO07 (NRRL B-50935), ISO08 (NRRL B-50936), ISO09 (NRRL B-50937), ISO10 (NRRL B-50938), ISO11 (NRRL B-50939), ISO12 (NRRL B-50940), ISO13 (NRRL B-50941), ISO14 (NRRL B-50942), ISO16 (NRRL B-67340), and variants thereof. In certain embodiments, the aforementioned variant thereof can be a *Methylobacterium* having chromosomal genomic DNA with at least 99%, 99.9, 99.8, 99.7, 99.6%, or 99.5% sequence identity to chromosomal genomic DNA of ISO01 (NRRL B-50929), ISO02 (NRRL B-50930), ISO03 (NRRL B-50931), ISO04 (NRRL B-50932), ISO05 (NRRL B-50933), ISO06 (NRRL B-50934), ISO07 (NRRL B-50935), ISO08 (NRRL B-50936), ISO09 (NRRL B-50937), ISO10 (NRRL B-50938), ISO11 (NRRL B-50939), ISO12 (NRRL B-50940), ISO13 (NRRL B-50941), ISO14 (NRRL B-50942) or ISO16 (NRRL B-67340). In certain embodiments, an additional biological can be a *Methylobacterium* selected from *M. gregans*, *M. radiotolerans*, *M. extorquens*, *M. populi*, *M. salsuginis*, *M. brachiatum*, and *M. komagatae*.

**[0058]** In certain embodiments, the additional biological can be a bacterium of the genus *Actinomyces*, *Agrobacterium*, *Arthrobacter*, *Alcaligenes*, *Aureobacterium*, *Azobacter*, *Beijerinckia*, *Bacillus*, *Brevibacillus*, *Burkholderia*, *Chromobacterium*, *Clostridium*, *Clavibacter*, *Comomonas*, *Corynebacterium*, *Curtobacterium*, *Enterobacter*, *Flavobacterium*, *Gluconobacter*, *Hydrogenophaga*, *Klebsiella*, *Methylobacterium*, *Paenibacillus*, *Pasteuria*, *Phingobacterium*, *Photobacterium*, *Phyllobacterium*, *Pseudomonas*, *Rhizobium*, *Bradyrhizobium*, *Serratia*, *Stenotrophomonas*, *Variovorax*, or *Xenorhabdus*. In particular embodiments the bacteria is selected from the group consisting of *Bacillus amyloliquefaciens*, *Bacillus cereus*, *Bacillus firmus*, *Bacillus licheniformis*, *Bacillus pumilus*, *Bacillus sphaericus*, *Bacillus subtilis*, *Bacillus thuringiensis*, *Chromobacterium suttsuga*, *Pasteuria penitrens*, *Pasteuria usage*, and *Pseudomona fluorescens*.

**[0059]** In certain embodiments the additional biological can be a fungus of the genus *Alternaria*, *Ampelomyces*, *Aspergillus*, *Aureobasidium*, *Beauveria*, *Colletotrichum*,

*Coniothyrium*, *Gliocladium*, *Metarhizium*, *Muscodora*, *Paecilomyces*, *Trichoderma*, *Typhula*, *Ulocladium*, or *Verticillium*. In particular embodiments the fungus is *Beauveria bassiana*, *Coniothyrium minitans*, *Gliocladium vixens*, *Muscodora albus*, *Paecilomyces lilacinus*, or *Trichoderma polysporum*.

**[0060]** In further embodiments the additional biological can be a nodulation related factor, plant growth activator or plant defense agent including, but not limited to harpin, *Reynoutria sachalinensis*, jasmonate, lipochitooligosaccharides (LCO), and isoflavones. In certain embodiments, additional biological components can include a combination of a bacterium and a nodulation related factor, plant growth activator or plant defense agent. One such combination includes, but not limited to, Optimize®, a combination of *Bradyrhizobium japonicum* and LCO.

**[0061]** In further embodiments, the additional biological can include, but is not limited to, various *Bacillus* sp., *Pseudomonas* sp., *Coniothyrium* sp., *Pantoea* sp., *Streptomyces* sp., and *Trichoderma* sp. Microbial biopesticides can be a bacterium, fungus, virus, or protozoan. Particularly useful biopesticidal microorganisms include various *Bacillus subtilis*, *Bacillus thuringiensis*, *Bacillus pumilis*, *Pseudomonas syringae*, *Trichoderma harzianum*, *Trichoderma virens*, and *Streptomyces lydicus* strains. Other microorganisms that are added can be genetically engineered or wild-type isolates that are available as pure cultures. In certain embodiments, it is anticipated that the biological or biocontrol agent can be provided in the fermentation broth, fermentation broth product, or composition in the form of a spore.

**[0062]** In certain embodiments, the fermentation broth, fermentation broth product, or compositions that comprise yield enhancing *Methylobacterium* can further comprise one or more introduced additional active ingredients or microorganisms of pre-determined identity other than *Methylobacterium*.

**[0063]** In certain embodiments, the liquid culture medium is prepared from inexpensive and readily available components, including, but not limited to, inorganic salts such as potassium phosphate, magnesium sulfate and the like, carbon sources such as glycerol, methanol, glutamic acid, aspartic acid, succinic acid and the like, and amino acid blends such as peptone, tryptone, and the like. Exemplary liquid media that can be used include, but are not limited to, ammonium mineral salts (AMS) medium (Whittenbury et al., 1970), Vogel-Bonner (VB) minimal culture medium (Vogel and Bonner, 1956), and LB broth ("Luria-Bertani Broth").

**[0064]** In general, a solid substance that can be used in the methods and compositions to provide for the efficient growth of *Methylobacterium* can be any suitable solid substance which is insoluble or only partially soluble in water or aqueous solutions. Such suitable solid substances are also non-bacteriocidal or non-bacteriostatic with respect to yield enhancing *Methylobacterium* when the solid substances are provided in the liquid culture media. In certain embodiments, such suitable solid substances are also solid substances that are readily obtained in sterile form or rendered sterile. Solid substances can be sterilized by any method that provides for removal of contaminating microorganisms and thus include, but are not limited to, methods such as autoclaving, irradiation, chemical treatment, and any combination thereof. These solid substances include natural

substances of animal, plant, microbial, fungal, or mineral origin, manmade substances, or combinations of natural and manmade substances. In certain embodiments, solid substances are inanimate solid substances. Inanimate solid substances of animal, plant, microbial, or fungal origin can be obtained from animals, plants, microbes, or fungi that are inviable (i.e. no longer living) or that have been rendered inviable. Diatom shells are thus inanimate solid substances when previously associated diatom algae have been removed or otherwise rendered inviable. Since diatom shells are inanimate solid substances, they are not considered to be photosynthetic organisms or photosynthetic microorganisms. In certain embodiments, solid substances include, but are not limited to, sand, silt, soil, clay, ash, charcoal, diatomaceous earth and other similar minerals, ground glass or glass beads, ground ceramic materials, ceramic beads, bentonite, kaolin, talc, perlite, mica, vermiculite, silicas, quartz powder, montmorillonite, and combinations thereof. In certain embodiments, a solid substance can be a polymer or polymeric beads. Polymers that can be used as a solid substance include, but are not limited to, various polysaccharides such as cellulosic polymers and chitinous polymers which are insoluble or only partially soluble in water or aqueous solutions, agar (i.e. galactans), and combinations thereof. In certain embodiments, a solid substance can be an insoluble or only partially soluble salt crystal. Salt crystals that can be used include, but are not limited to, insoluble or only partially soluble carbonates, chromates, sulfites, phosphates, hydroxides, oxides, and sulfides. In certain embodiments, a solid substance can be a microbial cell, fungal cell, microbial spore, or fungal spore. In certain embodiments, a solid substance can be a microbial cell or microbial spore wherein the microbial cell or microbial spore is not a photosynthetic microorganism. In certain embodiments, the microbial cell or microbial spore is not a photosynthetic microorganism, where the photosynthetic microorganism is selected from the group consisting of algae, cyanobacteria, diatoms, *Botryococcus braunii*, *Chlorella*, *Dunaliella tertiolecta*, *Gracilaria*, *Pleurochrysis carterae*, Sargassum, and Ulva. In still other embodiments, a solid substance can be an inactivated (i.e. inviable) microbial cell, fungal cell, microbial spore, or fungal spore. In still other embodiments, a solid substance can be a quiescent (i.e. viable but not actively dividing) microbial cell, fungal cell, microbial spore, or fungal spore. In still other embodiments, a solid substance can be cellular debris of microbial origin. In still other embodiments, a solid substance can be particulate matter from any part of a plant. Plant parts that can be used to obtain a solid substance include, but are not limited to, cobs, husks, hulls, leaves, roots, flowers, stems, barks, seeds, and combinations thereof. Products obtained from processed plant parts including, but not limited to, bagasse, wheat bran, soybean grits, crushed seed cake, stover, and the like can also be used. Such plant parts, processed plants, and/or processed plant parts can be milled to obtain a solid material in a particulate form that can be used. In certain embodiments, wood or a wood product including, but not limited to, wood pulp, sawdust, shavings, and the like can be used. In certain embodiments, a solid substance can be a particulate matter from an animal(s), including, but not limited to, bone meal, gelatin, ground or powdered shells, hair, macerated hide, and the like.

**[0065]** In certain embodiments, a solid substance is provided in a particulate form that provides for distribution of

the solid substance in the culture media. In certain embodiments, a solid substance is comprised of particle of about 2 microns to about 1000 microns in average length or average diameter. In certain embodiments, a solid substance is comprised of particle of about 1 micron to about 1000 microns in average length or average diameter. In certain embodiments, a solid substance is a particle of about 1, 2, 4, 10, 20, or 40 microns to any of about 100, 200, 500, 750, or 1000 microns in average length or average diameter. Desirable characteristics of particles used in the methods and compositions provided herein include suitable wettability such that the particles can be suspended throughout the media upon agitation.

**[0066]** In certain embodiments, a solid substance is provided in the media as a colloid wherein the continuous phase is a liquid and the dispersed phase is the solid. Suitable solids that can be used to form colloids in liquid media used to grow yield enhancing *Methylobacterium* include, but are not limited to, various solids that are referred to as hydrocolloids. Such hydrocolloids used in the media, methods and compositions provided herein can be hydrophilic polymers, of plant, animal, microbial, or synthetic origin. Hydrocolloid polymers used in the methods can contain many hydroxyl groups and/or can be polyelectrolytes. Hydrocolloid polymers used in the compositions and methods provided herein include, but are not limited to, agar, alginate, arabinoxylan, carrageenan, carboxymethylcellulose, cellulose, curdlan, gelatin, gellan,  $\beta$ -glucan, guar gum, gum arabic, locust bean gum, pectin, starch, xanthan gum, and mixtures thereof. In certain embodiments, a colloid used in the media, methods, and compositions provided herein can comprise a hydrocolloid polymer and one or more proteins.

**[0067]** In certain embodiments, a solid substance can provide for adherent growth of the yield enhancing *Methylobacterium* on the solid substance. Yield enhancing *Methylobacterium* that are adhered to a solid substance are *Methylobacterium* that cannot be substantially removed by simply washing the solid substance with the adherent yield enhancing *Methylobacterium* with growth media whereas non-adherent *Methylobacterium* can be substantially removed by washing the solid substance with liquid growth media. In this context, "substantially removed" means that at least about 30%, 40%, 50%, 60%, 70%, or 80% the *Methylobacterium* present are removed when the solid substance is washed with three volumes of liquid growth media. Such washing can be effected by a variety of methods including, but not limited to, decanting liquid from a washed solid phase or passing liquid through a solid phase on a filter that permits flow through of bacteria in the liquid. In certain embodiments, the adherent yield enhancing *Methylobacterium* that are associated with the solid can include both *Methylobacterium* that are directly attached to a solid and/or *Methylobacterium* that are indirectly attached to a solid substance. *Methylobacterium* that are indirectly attached to a solid substance include, but are not limited to, *Methylobacterium* that are attached to another *Methylobacterium* or to another microorganism that is attached to the solid substance, *Methylobacterium* that are attached to a solid substance by being attached to another substance that is attached to the solid substance, and the like. In certain embodiments, at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 98%, 99%, 99.5% or 99.9% of the *Methylobacterium* in the fermentation broth, fermentation broth product, or compositions are *Methylobacterium* that

are adhered to a solid substance. In certain embodiments, adherent yield enhancing *Methylobacterium* can be present on the surface of a solid substance in the fermentation broth, fermentation broth product, or composition at a density of at least about 1 *Methylobacterium*/20 square micrometers, of at least about 1 *Methylobacterium*/10 square micrometers, of at least about 1 *Methylobacterium*/10 square micrometers, of at least about 1 *Methylobacterium*/5 square micrometers, of at least about 1 *Methylobacterium*/2 square micrometers, or of at least about 1 *Methylobacterium*/square micrometer. In certain embodiments, adherent yield enhancing *Methylobacterium* can be present on the surface of a solid substance in the fermentation broth, fermentation broth product, or composition at a density of at least about 1 *Methylobacterium*/20 square micrometers to about 1 *Methylobacterium*/square micrometer, of at least about 1 *Methylobacterium*/10 square micrometers to about 1 *Methylobacterium*/square micrometer, of at least about 1 *Methylobacterium*/10 square micrometers to about 1 *Methylobacterium*/square micrometer, of at least about 1 *Methylobacterium*/5 square micrometers to about 1 *Methylobacterium*/square micrometer, or of at least about 1 *Methylobacterium*/2 square micrometers to about 1 *Methylobacterium*/square micrometer. In certain embodiments, adherent yield enhancing *Methylobacterium* can be present on the surface of a solid substance in the fermentation broth, fermentation broth product, or composition at a density of at least about 1 *Methylobacterium*/20 square micrometers to about 1 *Methylobacterium*/2 square micrometers, of at least about 1 *Methylobacterium*/10 square micrometers to about 1 *Methylobacterium*/2 square micrometers, of at least about 1 *Methylobacterium*/10 square micrometers to about 1 *Methylobacterium*/2 square micrometers, or of at least about 1 *Methylobacterium*/5 square micrometers to about 1 *Methylobacterium*/2 square micrometers. Biphasic fermentation broths provided herein can comprise a liquid phase that contains non-adherent *Methylobacterium*. In certain embodiments, titers of non-adherent *Methylobacterium* in the liquid phase can be less than about 100,000, 10,000, or 1,000 colony forming units (CFU) per ml. In certain embodiments of any of the aforementioned compositions, the yield enhancing *Methylobacterium* is selected from the group consisting of NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), NLS0610 (ISO26), and variants thereof. In certain embodiments of any of the aforementioned compositions, the yield enhancing *Methylobacterium* is selected from the group consisting of NLS0934 (NRRL B-67341), NLS1181 (ISO25), and variants thereof. In certain embodiments of any of the aforementioned compositions, the yield enhancing *Methylobacterium* is a combination of *Methylobacterium* NLS0109 and NLS0017, or variants thereof.

**[0068]** Fermentation broths and compositions with yield enhancing *Methylobacterium* at a titer of greater than about  $5 \times 10^8$  colony-forming units per milliliter, at a titer of greater than about  $1 \times 10^9$  colony-forming units per milliliter, at a titer of greater than about  $1 \times 10^{10}$  colony-forming units per milliliter, at a titer of at least about  $3 \times 10^{10}$  colony-forming units per milliliter are provided herein. In certain embodiments, fermentation broths and compositions provided herein can comprise yield enhancing *Methylobacterium* at a titer of at least about  $5 \times 10^8$  colony-forming units per milliliter to at least about  $3 \times 10^{10}$  colony-forming units per milliliter, at least about  $5 \times 10^8$  colony-forming units per

milliliter to at least about  $4 \times 10^{10}$  colony-forming units per milliliter, or at least about  $5 \times 10^8$  colony-forming units per milliliter to at least about  $6 \times 10^{10}$  colony-forming units per milliliter. In certain embodiments, fermentation broths provided herein can comprise yield enhancing *Methylobacterium* at a titer of at least about  $1 \times 10^9$  colony-forming units per milliliter to at least about  $3 \times 10^{10}$  colony-forming units per milliliter, at least about  $1 \times 10^9$  colony-forming units per milliliter to at least about  $4 \times 10^{10}$  colony-forming units per milliliter, or at least about  $1 \times 10^9$  colony-forming units per milliliter to at least about  $6 \times 10^{10}$  colony-forming units per milliliter. In certain embodiments, fermentation broths provided herein will comprise yield enhancing *Methylobacterium* at a titer of at least about  $1 \times 10^{10}$  colony-forming units per milliliter to at least about  $3 \times 10^{10}$  colony-forming units per milliliter, at least about  $1 \times 10^{10}$  colony-forming units per milliliter to at least about  $4 \times 10^{10}$  colony-forming units per milliliter, or at least about  $1 \times 10^{10}$  colony-forming units per milliliter to at least about  $6 \times 10^{10}$  colony-forming units per milliliter. In certain embodiments, fermentation broths provided herein will comprise yield enhancing *Methylobacterium* at a titer of, at least about  $3 \times 10^{10}$  colony-forming units per milliliter to at least about  $4 \times 10^{10}$  colony-forming units per milliliter, or at least about  $3 \times 10^{10}$  colony-forming units per milliliter to at least about  $6 \times 10^{10}$  colony-forming units per milliliter. In certain embodiments of any of the aforementioned compositions, the yield enhancing *Methylobacterium* is selected from the group consisting of NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), NLS0610 (ISO26), and variants thereof. In certain embodiments of any of the aforementioned compositions, the yield enhancing *Methylobacterium* is a combination of *Methylobacterium* NLS0109 and NLS0017, or variants thereof.

**[0069]** Yield enhancing *Methylobacterium* can be obtained as fermentation products and used to make various compositions useful for treating plants or plant parts to improve soybean yield. The *Methylobacterium* compositions can be applied to plants or plant parts in various forms, including for example as liquid compositions or as dried compositions, for example dried powders. Dried powders can be obtained by various methods, including for example, by spray-drying, freeze-drying, vacuum-drying, air-drying and fluidized bed-drying. Plants or plant parts that have been at least partially coated or coated with the fermentation broth products or compositions comprising yield enhancing *Methylobacterium* are thus provided. In certain embodiments, the plant part is a seed. Partial coating of a plant, a plant part, such as a seed includes, but is not limited to coating at least about 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 98%, 99%, or about 99.5% of the surface area of the plant, plant part, or plant seed. Also provided are processed plant products that contain the fermentation broth products or compositions with yield enhancing *Methylobacterium* or adherent yield enhancing *Methylobacterium*. In some embodiments, solid substances with adherent yield enhancing *Methylobacterium* can be used to make various compositions that are particularly useful for treating plant seeds. Seeds that have been at least partially coated with the fermentation broth products or compositions are thus provided. Partial coating of a seed includes, but is not limited to coating at least about 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 98%, 99%, or about 99.5% of the surface area of the seed. Also provided are processed

seed products, including, but not limited to, meal, flour, feed, and flakes that contain the fermentation broth products or compositions provided herein. In certain embodiments, the processed plant product will be non-regenerable (i.e. will be incapable of developing into a plant). In certain embodiments, the solid substance used in the fermentation product or composition that at least partially coats the plant or plant part or that is contained in the processed plant or plant part product comprises a solid substance and associated or adherent yield enhancing *Methylobacterium* that can be readily identified by comparing a treated and an untreated plant or plant part, or processed product thereof. In certain embodiments, the yield enhancing *Methylobacterium* is selected from the group consisting of NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), NLS0610 (ISO26), and variants thereof. In certain embodiments, the yield enhancing *Methylobacterium* is selected from the group consisting of NLS0934 (NRRL B-67341) and/or NLS1181 (ISO25). In certain embodiments of any of the aforementioned compositions, the yield enhancing *Methylobacterium* is a combination of *Methylobacterium* NLS0109, and NLS0017 or NLS0610, or variants thereof.

**[0070]** Compositions useful for treating plants or plant parts that comprise yield enhancing *Methylobacterium* can also further comprise additional components, including an active ingredient, an agriculturally acceptable adjuvant or an agriculturally acceptable excipient. An agriculturally acceptable adjuvant or an agriculturally acceptable excipient is typically an ingredient that does not cause undue phytotoxicity or other adverse effects when exposed to a plant or plant part. In certain embodiments, a solid substance used in a fermentation broth can itself be an agriculturally acceptable adjuvant or an agriculturally acceptable excipient so long as it is not bacteriocidal or bacteriostatic to the *Methylobacterium*. In some embodiments agriculturally acceptable adjuvants and/or excipients are added to *Methylobacterium* to increase stability and/or shelf life. In other embodiments, the composition further comprises at least one of an agriculturally acceptable adjuvant or an agriculturally acceptable excipient.

**[0071]** Any of the aforementioned compositions can also further comprise an additional active ingredient. In certain embodiments, the additional active ingredient is a pesticide used in the composition that does not substantially inhibit growth of the *Methylobacterium*. As *Methylobacterium* are gram negative bacteria, suitable bacteriocides used in the compositions can include, but are not limited to, bacteriocides that exhibit activity against gram positive bacteria but not gram negative bacteria. Compositions provided herein can also comprise a bacteriostatic agent that does not substantially inhibit growth of the *Methylobacterium*. Bacteriostatic agents suitable for use in compositions provided herein include, but are not limited to, those that exhibit activity against gram positive bacteria but not gram negative bacteria. Any of the aforementioned compositions can also be an essentially dry product (i.e. having about 5% or less water content), a mixture of the composition with an emulsion, or a suspension.

**[0072]** Agriculturally acceptable adjuvants used in the compositions that comprise yield enhancing *Methylobacterium* include, but are not limited to, components that enhance product efficacy and/or products that enhance ease of product application. Adjuvants that enhance product

efficacy can include various wetters/spreaders that promote adhesion to and spreading of the composition on plant parts, stickers that promote adhesion to the plant part, penetrants that can promote contact of the active agent with interior tissues, extenders that increase the half-life of the active agent by inhibiting environmental degradation, and humectants that increase the density or drying time of sprayed compositions. Wetters/spreaders used in the compositions can include, but are not limited to, non-ionic surfactants, anionic surfactants, cationic surfactants, amphoteric surfactants, organo-silicate surfactants, and/or acidified surfactants. Stickers used in the compositions can include, but are not limited to, latex-based substances, terpene/pinylene, and pyrrolidone-based substances. Penetrants can include mineral oil, vegetable oil, esterified vegetable oil, organo-silicate surfactants, and acidified surfactants. Extenders used in the compositions can include, but are not limited to, ammonium sulphate, or menthene-based substances. Humectants used in the compositions can include, but are not limited to, glycerol, propylene glycol, and diethyl glycol. Adjuvants that improve ease of product application include, but are not limited to, acidifying/buffering agents, anti-foaming/de-foaming agents, compatibility agents, drift-reducing agents, dyes, and water conditioners. Anti-foaming/de-foaming agents used in the compositions can include, but are not limited to, dimethopolysiloxane. Compatibility agents used in the compositions can include, but are not limited to, ammonium sulphate. Drift-reducing agents used in the compositions can include, but are not limited to, polyacrylamides, and polysaccharides. Water conditioners used in the compositions can include, but are not limited to, ammonium sulphate.

**[0073]** Methods of treating plants and/or plant parts with the fermentation broths, fermentation broth products, and compositions comprising yield enhancing *Methylobacterium* are also provided herein. Treated plants, and treated plant parts obtained therefrom, include, but are not limited to, soybean. As used herein, the term soybean includes, but is not limited to all varieties, subspecies, and cultivars of *Glycine max*. Soybean subspecies include, but are not limited to, *Glycine max* L. ssp. *max* and *Glycine max* ssp. *Formosana*. Plant parts that are treated include, but are not limited to, leaves, stems, flowers, roots, seeds, pods, and the like. Seeds or other propagules of any of the aforementioned plants can be treated with the fermentation broths, fermentation broth products, fermentation products, and/or compositions provided herein.

**[0074]** In certain embodiments, plants and/or plant parts are treated by applying the fermentation broths, fermentation broth products, fermentation products, and compositions that comprise yield enhancing *Methylobacterium* as a spray. Such spray applications include, but are not limited to, treatments of a single plant part or any combination of plant parts. Spraying can be achieved with any device that will distribute the fermentation broths, fermentation broth products, fermentation products, and compositions to the plant and/or plant part(s). Useful spray devices include a boom sprayer, a hand or backpack sprayer, crop dusters (i.e. aerial spraying), and the like. Spraying devices and or methods providing for application of the fermentation broths, fermentation broth products, fermentation products, and compositions to either one or both of the adaxial surface and/or abaxial surface can also be used. Plants and/or plant parts that are at least partially coated with any of a biphasic

fermentation broth, a fermentation broth product, fermentation product, or compositions that comprise a solid substance with yield enhancing *Methylobacterium* adhered thereto are also provided herein. In certain embodiments, the plant part is a seed. Partial coating of a plant or a plant part includes, but is not limited to coating at least about 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 98%, 99%, or about 99.5% of the surface area of the plant or plant part. In some embodiments, the plant part is a seed and partial coating includes, but is not limited to coating at least about 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 98%, 99%, or about 99.5% of the surface area of the seed. Also provided herein are processed plant products that comprise a solid substance with yield enhancing *Methylobacterium* adhered thereto.

[0075] In certain embodiments, seeds are treated by exposing the seeds to the fermentation broths, fermentation broth products, fermentation products, and compositions that comprise yield enhancing *Methylobacterium*. Seeds can be treated with the fermentation broths, fermentation broth products, and compositions provided herein by methods including, but not limited to, imbibition, coating, spraying, and the like. Seed treatments can be effected with both continuous and/or a batch seed treaters. In certain embodiments, the coated seeds can be prepared by slurring seeds with a coating composition containing a fermentation broth or fermentation broth product comprising yield enhancing *Methylobacterium* and air drying the resulting product. Air drying can be accomplished at any temperature that is not deleterious to the seed or the *Methylobacterium*. In certain embodiments, air drying is accomplished at a temperature not greater than 30 degrees Centigrade. The proportion of coating that comprises yield enhancing *Methylobacterium* includes, but is not limited to, a range of 0.1 to 25% by weight of the seed, 0.5 to 5% by weight of the seed, and 0.5 to 2.5% by weight of seed. In certain embodiments, a solid substance used in the seed coating or treatment will have yield enhancing *Methylobacterium* adhered thereon. In certain embodiments, a solid substance used in the seed coating or treatment will be associated with yield enhancing *Methylobacterium* and will be present in a fermentation broth, fermentation broth product, or composition obtained by the methods provided herein. Various seed treatment compositions and methods for seed treatment disclosed in U.S. Pat. Nos. 5,106,648, 5,512,069, and 8,181,388 are incorporated herein by reference in their entireties and can be adapted for use with an active agent comprising the fermentation broths, fermentation broth products, or compositions provided herein. In certain embodiments, the composition used to treat the seed can contain agriculturally acceptable excipients that include, but are not limited to, woodflours, clays, activated carbon, diatomaceous earth, fine-grain inorganic solids, calcium carbonate and the like. Clays and inorganic solids that can be used with the fermentation broths, fermentation broth products, or compositions provided herein include, but are not limited to, calcium bentonite, kaolin, china clay, talc, perlite, mica, vermiculite, silicas, quartz powder, montmorillonite and mixtures thereof. Agriculturally acceptable adjuvants that promote sticking to the seed that can be used include, but are not limited to, polyvinyl acetates, polyvinyl acetate copolymers, hydrolyzed polyvinyl acetates, polyvinylpyrrolidone-vinyl acetate copolymer, polyvinyl alcohols, polyvinyl alcohol copolymers, polyvinyl methyl ether, polyvinyl methyl ether-maleic anhydride

copolymer, waxes, latex polymers, celluloses including ethylcelluloses and methylcelluloses, hydroxy methylcelluloses, hydroxypropylcellulose, hydroxymethylpropylcelluloses, polyvinyl pyrrolidones, alginates, dextrans, maltodextrins, polysaccharides, fats, oils, proteins, karaya gum, jaguar gum, tragacanth gum, polysaccharide gums, mucilage, gum arabics, shellacs, vinylidene chloride polymers and copolymers, soybean-based protein polymers and copolymers, lignosulfonates, acrylic copolymers, starches, polyvinylacrylates, zeins, gelatin, carboxymethylcellulose, chitosan, polyethylene oxide, acrylamide polymers and copolymers, polyhydroxyethyl acrylate, methylacrylamide monomers, alginate, ethylcellulose, polychloroprene and syrups or mixtures thereof. Other useful agriculturally acceptable adjuvants that can promote coating include, but are not limited to, polymers and copolymers of vinyl acetate, polyvinylpyrrolidone-vinyl acetate copolymer and water-soluble waxes. Various surfactants, dispersants, anticaking agents, foam-control agents, and dyes disclosed herein and in U.S. Pat. No. 8,181,388 can be adapted for use with an active agent comprising the fermentation broths, fermentation broth products, or compositions provided herein.

[0076] Provided herein are compositions that comprise yield enhancing *Methylobacterium* that provide for increased yield of soybean plants relative to untreated plants, plant parts, and plants obtained therefrom that have not been exposed to the compositions. In certain embodiments, plant parts, including, but not limited to, a seed, a leaf, a flower, a stem, a root, a pod, or a coleoptile can be treated with the compositions provided herein to increase soybean plant yield. Treatments or applications can include, but are not limited to, spraying, coating, partially coating, immersing, and/or imbibing the plant or plant parts with the compositions provided herein. Partial coating of a soybean plant or a soybean plant part includes, but is not limited to coating at least about 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 98%, 99%, or about 99.5% of the surface area of the soybean plant or soybean plant part. In some embodiments, the soybean plant part that is partially coated is a soybean plant seed. In certain embodiments, a seed, a leaf, a flower, a stem, a root, a pod, or a coleoptile can be immersed and/or imbibed with a liquid, semi-liquid, emulsion, or slurry of a composition provided herein. Such seed immersion or imbibition can be sufficient to provide for increased yield in a treated soybean plant or soybean plant grown from a treated seed in comparison to an untreated soybean plant or soybean plant grown from an untreated seed. In certain embodiments, plant seeds can be immersed and/or imbibed for at least 1, 2, 3, 4, 5, or 6 hours. Such immersion and/or imbibition can, in certain embodiments, be conducted at temperatures that are not deleterious to the plant seed or the *Methylobacterium*. In certain embodiments, the seeds can be treated at about 15 to about 30 degrees Centigrade or at about 20 to about 25 degrees Centigrade. In certain embodiments, seed imbibition and/or immersion can be performed with gentle agitation. In certain embodiments, the soybean seed or VE stage soybean plant is exposed to the composition by providing the composition in furrow. Providing the composition in furrow represents one of several methods provided herein for applying a composition to a soybean seed or to a soybean plant at about the VE stage of soybean plant development.

[0077] Compositions provided herein comprising yield enhancing *Methylobacterium* and related methods are there-

fore expected to be useful in improving yield in a wide variety of soybean plants, including, but not limited to, all varieties, subspecies, and cultivars of *Glycine max*.

**[0078]** In certain embodiments, an amount of a composition provided herein that is sufficient to provide for increased soybean yield can be a composition with yield enhancing *Methylobacterium* at a titer of at least about  $1 \times 10^6$  colony-forming units per milliliter, at least about  $5 \times 10^6$  colony-forming units per milliliter, at least about  $1 \times 10^7$  colony-forming units per milliliter, at least about  $5 \times 10^8$  colony-forming units per milliliter, at least about  $1 \times 10^9$  colony-forming units per milliliter, at least about  $1 \times 10^{10}$  colony-forming units per milliliter, or at least about  $3 \times 10^{10}$  colony-forming units per milliliter. In certain embodiments, an amount of a composition provided herein that is sufficient to provide for increased soybean yield to a plant or plant part can be a composition with yield enhancing *Methylobacterium* at a titer of at least about  $1 \times 10^6$  colony-forming units per milliliter, at least about  $5 \times 10^6$  colony-forming units per milliliter, at least about  $1 \times 10^7$  colony-forming units per milliliter, at least about  $5 \times 10^8$  colony-forming units per milliliter, or at least about  $6 \times 10^{10}$  colony-forming units per milliliter of a liquid or an emulsion. In certain embodiments, an amount of a composition provided herein that is sufficient to provide for increased soybean yield can be a fermentation broth product with a yield enhancing *Methylobacterium* titer of a solid phase of that product is at least about  $1 \times 10^6$  colony-forming units per milliliter, at least about  $5 \times 10^6$  colony-forming units per milliliter, at least about  $1 \times 10^7$  colony-forming units per milliliter, or at least about  $5 \times 10^8$  colony-forming units per milliliter to at least about  $6 \times 10^{10}$  colony-forming units per milliliter of a liquid or an emulsion. In certain embodiments, an amount of a composition provided herein that is sufficient to provide for increased soybean yield can be a composition with a *Methylobacterium* titer of at least about  $1 \times 10^6$  colony-forming units per gram, at least about  $5 \times 10^6$  colony-forming units per gram, at least about  $1 \times 10^7$  colony-forming units per gram, or at least about  $5 \times 10^8$  colony-forming units per gram to at least about  $6 \times 10^{10}$  colony-forming units of *Methylobacterium* per gram of particles in the composition containing the particles that comprise a solid substance wherein a mono-culture or co-culture of yield enhancing *Methylobacterium* is adhered thereto. In certain embodiments, an amount of a composition provided herein that is sufficient to provide for increased soybean yield to a plant or plant part can be a composition with a *Methylobacterium* titer of at least about  $1 \times 10^6$  colony-forming units per mL, at least about  $5 \times 10^6$  colony-forming units per mL, at least about  $1 \times 10^7$  colony-forming units per mL, or at least about  $5 \times 10^8$  colony-forming units per mL to at least about  $6 \times 10^{10}$  colony-forming units of *Methylobacterium* per mL in a composition comprising an emulsion wherein a mono-culture or co-culture of a yield enhancing *Methylobacterium* adhered to a solid substance is provided therein or grown therein. In certain embodiments, an amount of a composition provided herein that is sufficient to provide for increased soybean yield to a plant or plant part can be a composition with a *Methylobacterium* titer of at least about  $1 \times 10^6$  colony-forming units per mL, at least about  $5 \times 10^6$  colony-forming units per mL, at least about  $1 \times 10^7$  colony-forming units per mL, or at least about  $5 \times 10^8$  colony-forming units per mL to at least about  $6 \times 10^{10}$  colony-forming units of *Methylobacterium* per mL of in a composition comprising an emulsion

wherein a mono-culture or co-culture of a yield enhancing *Methylobacterium* is provided therein or grown therein. In certain embodiments of any of the aforementioned compositions, the *Methylobacterium* is selected from the group consisting of (i) NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), NLS0610 (ISO26), and variants thereof; and (ii) a combination of *Methylobacterium* isolate NLS0109 (NRRL B-67340) or a variant thereof, and *Methylobacterium* isolate NLS0017 (B-50931) or NLS0610, or variants thereof. In certain embodiments of any of the aforementioned compositions, the composition is a lyophilized composition.

**[0079]** In certain embodiments, an amount of a composition provided herein that is sufficient to provide for increased soybean yield can be a composition with a *Methylobacterium* at a titer of at least about  $1 \times 10^4$  colony-forming units per milliliter, at least about  $1 \times 10^5$  colony-forming units per milliliter, at least about  $1 \times 10^6$  colony-forming units per milliliter, at least about  $5 \times 10^6$  colony-forming units per milliliter, at least about  $1 \times 10^7$  colony-forming units per milliliter, at least about  $5 \times 10^8$  colony-forming units per milliliter, at least about  $1 \times 10^9$  colony-forming units per milliliter, at least about  $1 \times 10^{10}$  colony-forming units per milliliter, or at least about  $3 \times 10^{10}$  colony-forming units per milliliter. In certain embodiments, an amount of a composition provided herein that is sufficient to provide for increased soybean yield can be a composition with *Methylobacterium* at a titer of at least about  $1 \times 10^4$  colony-forming units per milliliter, at least about  $1 \times 10^5$  colony-forming units per milliliter, at least about  $1 \times 10^6$  colony-forming units per milliliter, at least about  $5 \times 10^6$  colony-forming units per milliliter, at least about  $1 \times 10^7$  colony-forming units per milliliter, or at least about  $5 \times 10^8$  colony-forming units per milliliter to at least about  $6 \times 10^{10}$  colony-forming units per milliliter of a liquid or an emulsion. In certain embodiments, an amount of a composition provided herein that is sufficient to provide for increased soybean yield can be a fermentation broth product with a *Methylobacterium* titer of a solid phase of that product is at least about  $1 \times 10^4$  colony-forming units per gram, at least about  $1 \times 10^5$  colony-forming units per gram, at least about  $1 \times 10^6$  colony-forming units per gram, at least about  $5 \times 10^6$  colony-forming units per gram, at least about  $1 \times 10^7$  colony-forming units per gram, or at least about  $5 \times 10^8$  colony-forming units per gram to at least about  $6 \times 10^{10}$  colony-forming units of *Methylobacterium* per gram, at least about  $1 \times 10^{11}$  colony-forming units of *Methylobacterium* per gram, at least about  $1 \times 10^{12}$  colony-forming units of *Methylobacterium* per gram, at least about  $1 \times 10^{13}$  colony-forming units of *Methylobacterium* per gram, or at least about  $5 \times 10^{13}$  colony-forming units of *Methylobacterium* per gram of the solid phase. In certain embodiments, an amount of a composition provided herein that is sufficient to provide for increased soybean yield can be a composition with a *Methylobacterium* titer of at least about  $1 \times 10^6$  colony-forming units per gram, at least about  $5 \times 10^6$  colony-forming units per gram, at least about  $1 \times 10^7$  colony-forming units per gram, or at least about  $5 \times 10^8$  colony-forming units per gram to at least about  $6 \times 10^{10}$  colony-forming units of *Methylobacterium* per gram, at least about  $1 \times 10^{11}$  colony-forming units of *Methylobacterium* per gram, at least about  $1 \times 10^{12}$  colony-forming units of *Methylobacterium* per gram, at least about  $1 \times 10^{13}$  colony-forming units of *Methylobacterium* per gram, or at least about  $5 \times 10^{13}$  colony-forming units of *Methylobacterium* per gram.



gram of particles in the composition containing the particles that comprise a solid substance wherein a mono-culture or co-culture of *Methylobacterium* is adhered thereto. In certain embodiments, an amount of a composition provided herein that is sufficient to provide for increased soybean yield can be a composition with a *Methylobacterium* titer of at least about  $1 \times 10^6$  colony-forming units per mL, at least about  $5 \times 10^6$  colony-forming units per mL, at least about  $1 \times 10^7$  colony-forming units per mL, or at least about  $5 \times 10^8$  colony-forming units per mL to at least about  $6 \times 10^{10}$  colony-forming units of *Methylobacterium* per mL in a composition comprising an emulsion wherein a mono-culture or co-culture of a *Methylobacterium* adhered to a solid substance is provided therein or grown therein. In certain embodiments, an amount of a composition provided herein that is sufficient to provide for increased soybean yield can be a composition with a *Methylobacterium* titer of at least about  $1 \times 10^6$  colony-forming units per mL, at least about  $5 \times 10^6$  colony-forming units per mL, at least about  $1 \times 10^7$  colony-forming units per mL, or at least about  $5 \times 10^8$  colony-forming units per mL to at least about  $6 \times 10^{10}$  colony-forming units of *Methylobacterium* per mL of in a composition comprising an emulsion wherein a mono-culture or co-culture of a *Methylobacterium* is provided therein or grown therein. In certain embodiments of any of the aforementioned compositions, the *Methylobacterium* is selected from the group consisting of (i) NLS0934 (NRRL B-67341), NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), NLS0610 (ISO26), and variants thereof; and (ii) a combination of *Methylobacterium* isolate NLS0109 (NRRL B-67340) or a variant thereof, and *Methylobacterium* isolate NLS0017 (B-50931) or NLS0610 (ISO26), or variants thereof.

#### EXAMPLES

**[0080]** The following examples are included to demonstrate illustrative, non-limiting embodiments of the disclosure. It will be appreciated by those of skill in the art that the techniques disclosed in the following examples represent techniques discovered by the Applicants to function well in the practice of the invention. However, those of skill in the art should, in light of the instant disclosure, appreciate that many changes can be made in the specific embodiments that are disclosed, while still obtaining like or similar results, without departing from the scope of the disclosure.

##### Example 1

###### Preparation of *Methylobacterium*

**[0081]** In preparation for the field trials, *Methylobacterium* cultures were grown in ammonium mineral salts (AMS)+peptone media containing a carbon source (AMS+carbon (15 g/liter)+peptone (10 g/liter)) at 30° C. for 4 days. To prepare one liter of liquid AMS medium+carbon+peptone ("AMS media solution"), the following components were added to a total volume of one (1) liter of distilled water: 700 milligrams of dibasic potassium phosphate anhydrous, 540 milligrams of monobasic potassium phosphate anhydrous, one gram of magnesium sulfate heptahydrate, 500 milligrams of ammonium chloride anhydrous, 200 milligrams of calcium chloride dihydrate, 15 g of a carbon source, and 10 g of peptone.

**[0082]** The AMS media solution was sterilized by autoclaving. After growing for 4 days at 30° C., cultures were harvested by centrifugation at 5000 rpm for 10 minutes and then re-suspended in AMS media solution with 20% (v/v) glycerol as a cryoprotectant. The cultures were aliquoted and frozen at -80° C. until thawed for use. The liquid *Methylobacterium* preparations were applied to commercial soybean seeds that had been treated with conventional fungicides and insecticides. The liquid *Methylobacterium* preparations were applied to the seeds at a rate of about  $1 \times 10^5$  CFU/seed to about  $1 \times 10^6$  CFU/seed.

##### Example 2

###### Increased Soybean Yield by Application of *Methylobacterium* Compositions in 2017 Yield Trials

**[0083]** Soybean microplot field trials were established at four locations throughout the Midwest soybean growing region for the purpose of evaluating 67 *Methylobacterium* isolates applied as a seed treatment to commercial soybean seeds. Six controls that were not treated with a *Methylobacterium* isolate were included in the trials.

**[0084]** The microplot trials were conducted as randomized complete block design. There were six replications of each of the *Methylobacterium* isolates and controls at each location. The trials were established within existing farmer field sites and managed with local agronomic practices. All soybean varieties used were glyphosate-tolerant varieties. The trials were harvested for yield at physiological maturity with a commercial harvest combine.

#### Results

**[0085]** The analysis of yield results was conducted with JMP14 (SAS Institute), and the assumptions of normality and equal variance were tested prior to executing the Analysis of variance (ANOVA). ANOVA was conducted with the Analyze-Fit Model routine. Comparisons of *Methylobacterium* isolates to the average of the six untreated controls were performed with two-tailed t-tests applied to the pairwise differences between least-squares means estimated from the ANOVA model, under the null hypothesis that the difference in means was zero.

**[0086]** Across-locations analyses for the four locations were conducted according to the following model:  $[2] Y_{hijk} = M + I_i + S_j + IS_{ij} + L_h + R(L)_{k(h)} + LI_{hi} + IR(L)_{ik(h)} + LS_{hj} + LIS_{hij} + e_{hijk}$ , where  $Y_{hijk}$  is the yield at location h of isolate i at stage j in replicate k, M represents the overall mean,  $I_i$  is the fixed effect of isolate i,  $S_j$  is the fixed effect of stage j,  $IS_{ij}$  is the fixed effect of the interaction of isolate i and stage j,  $L_h$  is the random effect of location h,  $R(L)_{k(h)}$  is the random effect of replicate k nested within location h,  $LI_{hi}$  is the random effect of the interaction of location h and isolate i,  $IR(L)_{ik(h)}$  is the random effect of the interaction of isolate i and replicate k nested within location h,  $LS_{hj}$  is the random effect of the interaction of location h and stage j,  $LIS_{hij}$  is the random effect of the three-way interaction of location h with isolate i and stage j, and  $e_{hijk}$  is the random error.

**[0087]** Yield results for isolates selected for further analysis are shown in Table 2 below.



TABLE 2

| Yield results from 2017 soybean yield trials |                   |                                  |                               |                          |                                  |
|----------------------------------------------|-------------------|----------------------------------|-------------------------------|--------------------------|----------------------------------|
| Treatment                                    | Mean yield (Bu/A) | Yield difference from UTC (Bu/A) | Yield difference from UTC (%) | Con-trast p-value v. UTC | Win rate (# locations above UTC) |
| NLS0934                                      | 92.9              | +6.0                             | +6.9%                         | 0.0086                   | 4/4 (100%)                       |
| NLS1181                                      | 92.0              | +5.1                             | +5.9%                         | 0.025                    | 4/4 (100%)                       |
| NLS1179                                      | 91.6              | +4.7                             | +5.4%                         | 0.039                    | 4/4 (100%)                       |
| <i>Methylobacterium</i> S4                   | 91.2              | +4.3                             | +4.9%                         | 0.055                    | 4/4 (100%)                       |
| <i>Methylobacterium</i> S3                   | 91.1              | +4.2                             | +4.8%                         | 0.062                    | 4/4 (100%)                       |
| NLS0693                                      | 91.0              | +4.1                             | +4.7%                         | 0.075                    | 3/4 (75%)                        |
| NLS0610                                      | 90.0              | +3.1                             | +3.6%                         | 0.17                     | 4/4 (100%)                       |
| UTC                                          | 86.9              | —                                | —                             | —                        | —                                |
| <i>Methylobacterium</i> S5                   | 85.7              | -1.2                             | -1.4%                         | 0.59                     | 2/4 (50%)                        |
| NLS0497                                      | 85.7              | -1.2                             | -1.4%                         | 0.59                     | 1/4 (25%)                        |
| <i>Methylobacterium</i> S2                   | 85.7              | -1.2                             | -1.4%                         | 0.59                     | 2/4 (50%)                        |
| <i>Methylobacterium</i> S1                   | 82.4              | -4.5                             | -4.5%                         | 0.041                    | 1/4 (25%)                        |

\*P-values taken from a two-tailed t-test comparing each treatment to the Untreated Check (UTC).

### Example 3

#### Increased Soybean Yield by Application of *Methylobacterium* Compositions in 2018 Yield Trials

[0088] Strains for use in 2018 field trials were selected based on results from the 2017 field trials described above and cultures were prepared for inoculation as described in Example 1. Isolates demonstrating positive yield results in 2017 and several isolates that demonstrated a negative effect on yield were selected. Soybean field trials were established at four locations throughout the Midwest soybean growing region for the purpose of evaluating the *Methylobacterium* isolates applied as a seed treatment to commercial soybean seeds. Controls that were not treated with a *Methylobacterium* isolate were included in the trials.

[0089] The trials were conducted as randomized complete block design consisting of four 30-inch rows that were 40 feet long. There were four replications of each of the *Methylobacterium* isolates and controls at each location. The trials were established within existing farmer field sites and managed with local agronomic practices. Harvest was done per standard grower practices and plot yields were determined using conventional metered combine harvesters. Plot test weights were used to calculate bu/acre yield estimates for each plot.

### Results

[0090] The analysis of yield results was conducted as described above. Data for isolates showing an increase in soybean yield are provided in Table 3 below.

TABLE 3

| Yield results from 2018 soybean yield trials. |                   |                                  |                               |                          |                                  |
|-----------------------------------------------|-------------------|----------------------------------|-------------------------------|--------------------------|----------------------------------|
| Treatment                                     | Mean yield (Bu/A) | Yield difference from UTC (Bu/A) | Yield difference from UTC (%) | Con-trast p-value v. UTC | Win rate (# locations above UTC) |
| NLS0934                                       | 73.8              | +3.0                             | +4.2%                         | 0.0050                   | 4/4 (100%)                       |
| NLS1181                                       | 73.5              | +2.7                             | +3.8%                         | 0.0098                   | 4/4 (100%)                       |
| NLS0497                                       | 73.2              | +2.4                             | +3.4%                         | 0.025                    | 3/4 (75%)                        |
| NLS0693                                       | 72.6              | +1.8                             | +2.5%                         | 0.097                    | 3/4 (75%)                        |
| NLS1179                                       | 72.5              | +1.7                             | +2.4%                         | 0.10                     | 2/4 (50%)                        |
| NLS0610                                       | 70.9              | +0.1                             | +0.1%                         | 0.94                     | 2/4 (50%)                        |
| UTC                                           | 70.8              | —                                | —                             | —                        | —                                |
| <i>Methylobacterium</i> S1                    | 70.8              | 0                                | 0                             | 1.00                     | 2/4 (50%)                        |
| <i>Methylobacterium</i> S2                    | 70.7              | -0.1                             | -0.1%                         | 0.92                     | 2/4 (50%)                        |
| <i>Methylobacterium</i> S3                    | 70.7              | -0.1                             | -0.1%                         | 0.86                     | 2/4 (50%)                        |
| <i>Methylobacterium</i> S4                    | 70.1              | -0.7                             | -1.0%                         | 0.47                     | 2/4 (50%)                        |
| <i>Methylobacterium</i> S5                    | 70.0              | -0.8                             | -1.1%                         | 0.39                     | 2/4 (50%)                        |

\*P-values taken from a two-tailed t-test comparing each treatment to the Untreated Check (UTC).

### Example 4

#### Increased Soybean Yield by Application of Dry *Methylobacterium* Compositions

[0091] In preparation for the field trials, *Methylobacterium* cultures were grown in ammonium mineral salts (AMS)+peptone media containing a carbon source (AMS+carbon (15 g/liter)+peptone (10 g/liter)) at 30° C. for 4 days. To prepare one liter of liquid AMS medium+carbon+peptone (“AMS media solution”), the following components were added to a total volume of one (1) liter of distilled water: 700 milligrams of dibasic potassium phosphate anhydrous, 540 milligrams of monobasic potassium phosphate anhydrous, one gram of magnesium sulfate heptahydrate, 500 milligrams of ammonium chloride anhydrous, 200 milligrams of calcium chloride dihydrate, 15 g of a carbon source, and 10 g of peptone.

[0092] The AMS media solution was sterilized by autoclaving. After growing for 4 days at 30° C., cultures were harvested by centrifugation at 5000 rpm for 10 minutes and cell pellets from centrifugation were re-suspended in AMS media solution at one tenth the original volume (“concentrated slurry”).

[0093] The concentrated slurry and commercial formulation additives were mixed and the resulting mixture was frozen at -80° C. Frozen mixture was freeze-dried using standard industrial freeze-drying conditions and then ground into a powder. The powder was applied to commercial soybean seeds that had been treated with conventional fungicides and insecticides at a rate of about 1×10<sup>5</sup> CFU/seed to about 1×10<sup>6</sup> CFU/seed.

[0094] Soybean field trials were established at multiple locations throughout the Midwest soybean growing region (9 locations in 2017 and 11 locations in 2018). The trials were conducted as randomized complete block design consisting of four 30-inch rows that were 40 feet long. There were four replications of each of the *Methylobacterium* isolates and controls at each location. *Methylobacterium* S6

positive control contained a *Methylobacterium* isolate that was previously shown to increase soybean yield (US20160302423). The trials were established within existing farmer field sites and managed with local agronomic practices. Harvest was done per standard grower practices and plot yields were determined using conventional metered combine harvesters. Plot test weights were used to calculate bu/acre yield estimates for each plot. Results are reported in Table 4 below.

TABLE 4

| Treatment                  | 2018<br>Yield<br>bu/ac | 2017<br>Yield<br>bu/ac | 2017/18<br>average<br>yield | 2 yr average<br>increase vs<br>UTC (bu) |
|----------------------------|------------------------|------------------------|-----------------------------|-----------------------------------------|
| <i>Methylobacterium</i> S6 | 62.56                  | 58.10                  | 60.33                       | 4.47                                    |
| Control                    |                        |                        |                             |                                         |
| NLS0610/NLS0109            | 61.77                  | 58.51                  | 60.14                       | 4.28                                    |
| NLS0017/NLS0109            | 61.24                  | 60.02                  | 60.63                       | 4.77                                    |
| NLS0610                    | 62.09                  | 58.52                  | 60.31                       | 4.45                                    |
| Optimize ®                 | 60.08                  |                        |                             |                                         |
| UTC                        | 57.47                  | 54.24                  | 55.86                       |                                         |

#### Example 5 Detection and Identification of *Methylobacterium* Strains, Variants and Derivatives

**[0095]** Assays are disclosed for detection or identification of specific *Methylobacterium* strains and closely related derivatives. Genomic DNA fragments unique to a *Methylobacterium* strain are identified and qPCR Locked Nucleic Acid (LNA) based assays are developed.

**[0096]** Genomic DNA sequences of *Methylobacterium* strains are compared by BLAST analysis of approximately 300 bp fragments using a sliding window of from 1-25 nucleotides to whole genome sequences of over 1000 public and proprietary *Methylobacterium* isolates. Genomic DNA fragments are identified that have weak BLAST alignments, indicative of approximately 60-95% identity over the entire fragment, to corresponding fragments of a *Methylobacterium* of interest. Fragments from the target *Methylobacterium* strain genome corresponding to the identified weak alignment regions were selected for assay development.

**[0097]** Target fragment sequences that distinguish NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), and NLS0610 (ISO26), from related *Methylobacterium* isolates are provided in Table 5 below.

TABLE 5

| Unique Fragment Sequences of <i>Methylobacterium</i> Strains |             |              |                                                                                                                                                                                                                                                                                                                                               |
|--------------------------------------------------------------|-------------|--------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Strain                                                       | Fragment    | SEQ<br>ID NO | Sequence                                                                                                                                                                                                                                                                                                                                      |
| NLS0497                                                      | ref1_46464  | 1            | CTCAACGTTGCCGCTTAACCGAAGACTAACTA<br>ACATAAATTTCAGTTAGCCGCGAAACAGAATAT<br>GATACTTGGTCTTATACGACGCATAGAGGCGCA<br>TTCCGTC AAGTCGCGCATTG CAGAGTCCCCCGCA<br>AAATTTCGTTTTCTGGGAGCGAGCAGAAACGTGT<br>GCACGTGTTCTGTCACTCTATTTTCTGGTTTGA<br>GCAATCATCTCGCGGCTCAGATCCCGTCGTTCCA<br>TAGAGTGGTTTTCCGTTTCAGTATGAATTGTGAGC<br>AGCCCCGCGGCTATATCTACTTCGACGTATA |
| NLS0497                                                      | ref1_85227  | 2            | ACCAACAGGCAGAGAGGCTTACAAAATGCTC<br>TAGATATCTATAATCGCGCCGGTCTGGATAATA<br>GGCGGGAAAAGTCAAATATTCTTATCGGACTTG<br>GCGATGCACTAGCGCACTGAGCAAATTTATAG<br>ACGCGAAAATTTCTACAGTGAAGGTTTAGCAG<br>TCCGAGCGCGTCTATGACGGTGTGTCGGACTA<br>GTGCTGTCTGCAATTTCAGAGCAGCCGTTCCGGC<br>TGGTTCAGTTTGGGTATGCCCTTTCTGTACCAGC<br>CTGTGACTGGTCTGGTATGTAGCCCTTGACCAAT       |
| NLS0497                                                      | ref1_98103  | 3            | ATGAAGCTCCGTCGTGTTTATGTAAGAACGTTT<br>GCAGTTTCTACGATGCTGAAGAACTCATTCTGGA<br>CGGAGATATTTCTATTATTATAGGCCAAATGGT<br>GGAGGAAAAACAAATCTACTTGATGCGACGATA<br>CACTTGTGAGAAGACATTTATTGCAATCGTGGG<br>CTGTCGTTCCGCATACATACAGCCAATAAACTA<br>TTCCGATCACTTCCAACCAATGATCAATCACA<br>AACTATGCGCTGGAGAAGTACAGCGGTCGCGAA<br>AGCGAAGACCAAGTCGTTGAGTTGATATC         |
| NLS0693                                                      | ref1_622066 | 4            | CGGGGGCGGTCTCCTCATATATGGAGGGAATTA<br>ATCTGAAGGGTCGAGAGCCCCGCTACGGCACAC<br>ACACGGCGGGCGGGCGGTGGGCTCCTTCTCATA<br>TACGCTTCTGATAATTCGACCGTAATCGCCCCG<br>CCGGCCGAGCGGCCAGACCTTATTAGTCCAGCA<br>TCACACGGTGATCGTATCGTGGAGGGCGAACA<br>ACCGCCTAAGGTTGCGCACGTAGCCATCCACAT<br>CCGCCCTTGCCGCTCGCGCAAGGGAAGCAGAA<br>GCTTGATAGCGACTTCGCGGGCGCTTTTGGTAGC<br>C |
| NLS0693                                                      | ref1_2496   | 5            | TGTGGGCGCTCATGGAACGGGGGACGTGCTTT<br>TTTGATCTTAGCCATTGTCATCATCCATCGGTC                                                                                                                                                                                                                                                                         |

TABLE 5-continued

| Unique Fragment Sequences of Methylobacterium Strains |              |              |                                                                                                                                                                                                                                                                                                                                                 |
|-------------------------------------------------------|--------------|--------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Strain                                                | Fragment     | SEQ<br>ID NO | Sequence                                                                                                                                                                                                                                                                                                                                        |
|                                                       |              |              | CGGGTGACAGGCCTTTACGCACATAACAACGGT<br>TATTTTCGATCAGCCCCCTCCTGATCGCACAAAGTC<br>AGCTATACAGATGCTTCTGAAAGCCAACAAGG<br>CGTTACGTATCTGCGTGACCCACCGAGCTGCGC<br>TGCGGCATCGCTGACAGCTCTGTATTTTAATTG<br>AGGAGGTGGGGCAATTTAGCCTGGTCAAGCTCG<br>CCAACGTCTTCCTTCACATATTGAGACAGGAC                                                                                |
| NLS0693                                               | ref1_2640477 | 6            | AGAGTGTATTTCCCCAAGACTTTATCTCTTTGCT<br>CATCACAACAGCCCGCTCAAGATGATATGCG<br>CTGGGATTGTGCGCGACCCGTGTAAATTGTAT<br>TTACCTCTCGGCTCATGTCCCTGTGCGCGGAGC<br>GATTGGAGGTAAAGTAATCTCAACTTTGCGGCC<br>CATGGCTGATGCTAGCTCAGCTACGCGACCTAA<br>GGTAATATCTTTGTGTCGCGGAGTTCGCGATGA<br>ATGACCGAGCGATTGACGCCGATGGTGGGGCA<br>ATATCGGCTTGAGTTAGACCGTTCTTGCTTC            |
| NLS1179                                               | ref1_687571  | 7            | ACCTGTCTCGACATCGGGGCGAGCCACGAGGCG<br>TCGGCCGTGGTAGATCTCGCGGAAGCGGTAGTA<br>GTGGCGGAGGTCCCATCGGGATCGATGGGGGA<br>GTCCGGCGTACCTTCTCCCTCTTCTGCGATTAGC<br>GACAGGGCTCGCATGGCGCTGGGTGCGTCAGTG<br>ATGGCGAACCACTACTATCGGGGAACCAACCGC<br>CTGTGACGAGCTGGCGGTCCGGGTTTCTTTGA<br>AGACAGCGTCACCCAGCGCTTCGATCTTCTCAAT<br>CAAGGCTGCGTAGAACTGCCCGATGGTCGCGAA       |
| NLS1179                                               | ref1_695522  | 8            | GACGTGGTGACGGTGGCACCAGTCGCCAGCGA<br>AGCGGCTACTGTACTGAGCCGCGAGTCATGTCG<br>GCTGAAGAAATGGTCGATATGCCCAGGGAGGT<br>CGGGCGTAGGTCGTACCTCAAATTCACCGGCAG<br>GATCTTGTCAGCAAGGCGTCCGAATTCGCATC<br>GACCTGACTTAGGAAATTACCGAGGTAATCGCT<br>CGGTCCGTTTCGCGTCGGCGGCTGCATCGCCAG<br>CAGCGCCTTATGCTCGGTGCTCAACTCACGAGC<br>GAATATTGTCCAGGACAGGTTGTGACCTCGGC<br>GAC    |
| NLS1179                                               | ref1_705877  | 9            | GAGCATGGCAGGCCACCACGCATCGCTTCTTCA<br>TCACCTTGCTCGGACGACTTGCTGCTTTGTCGGT<br>CTCTCTCTCAAGGACGAGACGCTGCGACACTTG<br>CTTCGTGAGAGTGCCAGGATTAATCCCGTCAAT<br>ACCACTACTACGTGAGTGGCGTGCGCCGGCG<br>GCACGAGGACGTCGACGCGGAACGATTGATTC<br>GAGACGCTAACTTTGACGTGTACAATTTAGTCAC<br>GCTTTTCTTAGCGACGCGGAGATAGCCTCGTTG<br>GGCCGCTTGCTCACCGCGATGCAGCCGAGTT          |
| NLS1181                                               | ref1_54084   | 10           | AACGCGATGGGAGCAGCACGTAATGGGCGTCAT<br>ACAAGATATAAAGAAGGACTTGAAGGTAGAAT<br>AGATAACCTTAAGGAAAATATTGATAATATCAG<br>TTCCAGTATCACATGCACAATACAAGGAGTGCC<br>GAAGGTCGAGTCAGAAAGAAATTTGGGATGTGC<br>AAACGCTTCGTTTGACCAAAATACAGAAAATGTG<br>GATGAATGGTCTACGGATAGGGCTGCATTTCTC<br>GACGCCGTGAAAAGTTTGATAGACCGCAACGCA<br>AGTAATGGAAAATTCGGAGTTCCAGGATCATAT<br>TG |
| NLS1181                                               | ref1_4816166 | 11           | TTTTCTTTTCGCGCCTATGACCTGAGTGCGGCCG<br>AAGCTGTCCGCTTATTAGAGTTTACAACCAGAT<br>GCGAGCGGCTTAGCTCTTCTCCGTTATTAAAGCA<br>GGGCGCCTCTTCTTAGGGGCGCCCTTCATATTT<br>AATCTTGTCTGATGTCTGGCGCCATATCAGACAA<br>ACATCAGCGTTGACCTTGATTTGCATCTACATT<br>AGGGTTGCTCCAGAAATGGAGCCGCGATGTCC<br>GTCCCGTACCTCAAGCGAGAGATGTGGGGTGTG<br>TATTACATCCATTGGCGCGAGGGCGGCCG         |
| NLS1181                                               | ref1_2292077 | 12           | GTTGGAGATGGTGCGTTTGCAACCGTCAGATC<br>GCCTACTGTGAATCCGGTGACCTCCCGAGAGAA                                                                                                                                                                                                                                                                           |

TABLE 5-continued

| Unique Fragment Sequences of <i>Methylobacterium</i> Strains |              |           |                                                                                                                                                                                                                                                                                                                                          |
|--------------------------------------------------------------|--------------|-----------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Strain                                                       | Fragment     | SEQ ID NO | Sequence                                                                                                                                                                                                                                                                                                                                 |
|                                                              |              |           | CGTGAAGGTAACGTGTCGTAGTCTCGCCAACTGC<br>CAAAGAGGTGTCCGCAACGGCGATGGTCGCTGT<br>CGGGGCGCTTGTTTGGACATCAACGGTGAGACC<br>CGCCGAGGCAGGCCCGGTGTACCAGCGGTATC<br>GACGGCCCTAGCGGTGACCGTGTGCGATCCAGC<br>CGTTAGCGTCGAACTGGTGATGCTATAGGTCCC<br>GCCGATCGCCACTGCCGAGCCCAGAATGGTCGT<br>GCC                                                                   |
| NLS0610                                                      | ref1_2810264 | 13        | ACCGAAGGCGTCCCCGGACACGAAGGCCTGAAACAC<br>CATATCTGTGGCGATCAGGCCGACGTGGTCGCGGAC<br>TTCAACTGGCAGAGAATGCCAGGCCGCTTCGATTCA<br>GATGATACTGGTACGGACATAGGAGCGGCTTAGCTT<br>TCTCAGTGCAAATGTGATTGATTCGGGCTCAAAATG<br>ATCTTGATCGGACGAGACGTTTCAATCCATGTCGTG<br>TTGCCATCGCCGATCGGTGCGTCAAGAGACAGATGG<br>CGCCGACCGTAGATACGCGTTGGGTTGCCCGCACC<br>GCTTCTCCA |
| NLS0610                                                      | ref1_322980  | 14        | GGAGGTGTGATCTGATGATGTGCTGGATGAAAT<br>TGGCGGTGAGCACTTGTTCAAGTTGGCCAGCTC<br>GACGAGATCGGCGTGATGCTCGGCGTCGATCAG<br>GATGTTCAAGGAGACCGGACGTACGAGGACTT<br>GGTATTAGCGCCGTTGCGCATCAGCTTGACGCT<br>TGCTCTGCTTCTCAGCGTGCCGCGTCAGGATGAC<br>CCTGATGTAGCTGTTGAGGTTGATGCCGTAATAG<br>CCTGCGGACTCTGTGAGATCCCGCGAAGATCG<br>TCGGCGAGGGTCAGGCGGATGGTGCTGGTCGG |
| NLS0610                                                      | ref1_2785241 | 15        | AAGTAACCGCTCAACATGATCTTCAGCATGTTGT<br>CCAACAGCAGGAGAATACATGTAATTCACCATG<br>ACCGGCAAGCTGCGACTGGCCATTGCTTCCACC<br>GCTTGAATGTAGCGATCGAATTCGCAAAATCA<br>GGGTGGAATGAAAATATCGAACCAAACTGCGAG<br>CCTTGAATCCGTTCTGCAAAATTATCGAAAATT<br>TTCTTGGCCGACTGCCGTTGAAAACATTCTTAC<br>GTTTACATGCGGCCCGCTGAAACAAGACAGTC<br>TACCAGCTCTGGGAAATGGGGTGAAGGTCGG   |

**[0098]** Regions in SEQ ID NOS: 1-15 where corresponding regions in other *Methylobacterium* strains were identified as having one or more nucleotide mismatches from the target *Methylobacterium* strain sequence were selected, and qPCR primers designed using Primer3 software (Untergasser et al. (2012), Koressaar et al. (2007)) to flank the mismatch regions, have a melting temperature (T<sub>m</sub>) in the range of 55-60 degrees, and to generate a PCR DNA fragment of approximately 100 bp. The probe sequence was designed with a 5' FAM reporter dye, a 3' Iowa Black FQ quencher, and contains one to six LNA bases (Integrated

DNA Technologies, Coralville, Iowa). At least 1 of the LNA bases is in the position of a mismatch, while the other LNA bases are used to raise the T<sub>m</sub>. The T<sub>m</sub> of the probe sequence is targeted to be 10 degrees above the T<sub>m</sub> of the primers.

**[0099]** Primer and probe sequences for specific detection of NLS0497 (ISO22), NLS0693 (ISO23), NLS1179 (ISO24), NLS1181 (ISO25), NLS0610 (ISO26), and variants thereof, are provided as SEQ ID NOS: 16-51 in Table 6. Each of the probes contains a 5' FAM reporter dye and a 3' Iowa Black FQ quencher.

TABLE 6

| Primers and Probes for Specific Detection of Strains |           |                       |
|------------------------------------------------------|-----------|-----------------------|
| Primer/Probe                                         | SEQ ID NO | Sequence*             |
| NLS0497_ref011_46464_For                             | 16        | CACGTGTTCTGTCACTCTATT |
| NLS0497_ref011_46464_Rev                             | 17        | GAACCGGAAACCACCTCTATG |
| NLS0497_ref011_46464_Probe                           | 18        | AGCAATCATCTCGCGGCTC   |
| NLS0497_ref022_85227_For                             | 19        | GCGCACTGAGCAAATTTATAG |
| NLS0497_ref022_85227_Rev                             | 20        | CTGAATTGCGAGCAGACA    |

TABLE 6-continued

| Primers and Probes for Specific Detection of Strains |              |                          |
|------------------------------------------------------|--------------|--------------------------|
| Primer/Probe                                         | SEQ<br>ID NO | Sequence*                |
| NLS0497_ref022_85227_Probe                           | 21           | AGTGAAGGTTTAGCAGTCCG     |
| NLS0497_ref015_98103_For                             | 22           | CTACTTGATGCGACGATACAC    |
| NLS0497_ref015_98103_Rev                             | 23           | GGTTGGAAGTGATCGGAATAG    |
| NLS0497_ref015_98103_Probe                           | 24           | TCGTTGCGCCATACATACACG    |
| NLS0693_ref1_622066_For                              | 25           | GCTCCTTCCTCATATACGCTTC   |
| NLS0693_ref1_622066_Rev                              | 26           | CCGTGTGATGCTGGACTAATAA   |
| NLS0693_ref1_622066_Probe                            | 27           | TAATTTCGACCGTAATCGCC     |
| NLS0693_ref6_2496_For                                | 28           | TGATCGCACAAAGTCAGCTATAC  |
| NLS0693_ref6_2496_Rev                                | 29           | CGAGCTTGACCAGGCTAAAT     |
| NLS0693_ref6_2496_Probe                              | 30           | ACGCCTTTGTTGGCTTTCAG     |
| NLS0693_ref1_2640477_For                             | 31           | CGGAGCGATTGGAGGTAAAG     |
| NLS0693_ref1_2640477_Rev                             | 32           | CTCCGCGACACAAAGATATTA    |
| NLS0693_ref1_2640477_Probe                           | 33           | AATCTCAACTTTCGCGCCC      |
| NLS1179_ref1_687571_For                              | 34           | CTTCTCCCTCTTCTGCGATTAG   |
| NLS1179_ref1_687571_Rev                              | 35           | GACGCTGTCTTCAAAGGAAAC    |
| NLS1179_ref1_687571_Probe                            | 36           | TGATGGCGAACAACTACT       |
| NLS1179_ref1_695522_For                              | 37           | GCATCGACCTGACTTAGGAAA    |
| NLS1179_ref1_695522_Rev                              | 38           | GAGTTGACGACCGAGCATAA     |
| NLS1179_ref1_695522_Probe                            | 39           | AGGTAATCGCTCGGTCCGT      |
| NLS1179_ref1_705877_For                              | 40           | CTTTGTCGGTCTCTCTCTCAAG   |
| NLS1179_ref1_705877_Rev                              | 41           | CAGCTGACGTAGTAGTGGTATTG  |
| NLS1179_ref1_705877_Probe                            | 42           | TCGTCAGAGTGCCAGGATT      |
| NLS1181_ref1_54084_For                               | 43           | GTTCCAGTATCACATGCACAATAC |
| NLS1181_ref1_54084_Rev                               | 44           | CGTAGGACCATTTCATCCACAT   |
| NLS1181_ref1_54084_Probe                             | 45           | TTTGGGATGTGCAAACGCT      |
| NLS1181_ref1_4816166_For                             | 46           | GCTTAGCTCTTCCTCCGTTATT   |
| NLS1181_ref1_4816166_Rev                             | 47           | ATTCTGGGAGCAACCCTAATG    |
| NLS1181_ref1_4816166_Probe                           | 48           | ATCTTGTCGATGCTGGCG       |
| NLS1181_ref1_2292077_For                             | 49           | CGTGAAGGTAAGTGTCTAGTTC   |
| NLS1181_ref1_2292077_Rev                             | 50           | CACCGTTGATGTCCAACAAG     |
| NLS1181_ref1_2292077_Probe                           | 51           | AACTGCCAAAGAGGTGTCC      |
| NLS0610_ref1_2810264_For                             | 52           | CATAGGAGCGGCTTAGCTTTCT   |
| NLS0610_ref1_2810264_Rev                             | 53           | GATGGCAACACGACATGGATTG   |
| NLS0610_ref1_2810264_Probe                           | 54           | CAGTGCAAATGTGATTGATTC    |
| NLS0610_ref1_322980_For                              | 55           | GACGTACGCAGGACTTGGTATTAG |
| NLS0610_ref1_322980_Rev                              | 56           | CTCAACAGCTACATCAGGGTCATC |

TABLE 6-continued

| Primers and Probes for Specific Detection of Strains |           |                              |
|------------------------------------------------------|-----------|------------------------------|
| Primer/Probe                                         | SEQ ID NO | Sequence*                    |
| NLS0610_ref1_322980_Probe                            | 57        | TTGCTCT <u>T</u> GTTCACGCGTG |
| NLS0610_ref1_2785241_For                             | 58        | AACCAAACTGCGAGCCTTGA         |
| NLS0610_ref1_2785241_Rev                             | 59        | GGGCCGCATGTAACGTAAGA         |
| NLS0610_ref1_2785241_Probe                           | 60        | TTCTT <u>G</u> GCCGACTGCCG   |

\*Bold and underlined letters represent the position of an LNA base

Use of Primer/Probe Sets on Isolated DNA to Detect Specific *Methylobacterium* Isolates and Variants Thereof and Distinguish from Related *Methylobacterium* Isolates

**[0100]** Each 10 ul qPCR reaction contains 5 ul of Quantabio PerfeCTa qPCR ToughMix 2× Mastermix, Low ROX from VWR, 0.05 ul of 100 uM forward primer, 0.05 ul of 100 uM reverse primer, 0.05 ul of 50 uM probe, 2.85 ul nuclease free water and 2 ul of DNA template. Approximately 1 ng of DNA template is used per reaction. The reaction is conducted in a ThermoFisher QuantStudio™ 6

Flex Real-Time PCR System with the following program: 95° C. for 3 min, then 40 cycles of 95° C. for 15 sec and 60° C. for 45 seconds. The analysis software on the PCR instrument calculates a threshold and Ct value for each sample. Each sample is run in triplicate on the same qPCR plate. A positive result is indicated where the delta Ct between positive and negative controls is at least 5.

**[0101]** Other *Methylobacterium* isolates listed in Table 1 and variants thereof are identified by the presence of DNA fragments using assays as described above. Unique fragments for use in such methods are provided in Table 7.

TABLE 7

| Strain  | Fragment    | SEQ ID NO | Sequence                                                                                                                                                                                                                                                                                                                                      |
|---------|-------------|-----------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| NLS0017 | ref4_930    | 61        | GCAAAACGACCTAATAGTTCTACAGCGGCATGCG<br>CCAAGTCAGCGCGGTGAACAGTATACCTGGGAG<br>CAACTTGTCCTCCGAAACCCACATAAAACAAATT<br>ACTCCTGGCAGTGCCGAGTCCATCAAAATCGAAT<br>ACAATATTTCTCGAGGAGGCATCTGTAATAGCCT<br>GCCAAGCAACAAAGCTATGGCGCCGTTATGACT<br>TTCATTGCTTCTGGTAGACATAAAATAATATGCC<br>GATTTGTGATCCCAATGTAGAAATATGCCGCAT<br>CAATTGCGCCAAGTTTATTTCGGATCGAT    |
| NLS0017 | ref1_142021 | 62        | GGCGCCAACGGTATGATCGCATGATTTTCTCGCG<br>GCATAGCTTGCGGGAATGGCGTATTTGGCGCTCT<br>CCTCAGGAATTTCTAAGGGCATACGAGGAACCT<br>TACAGCACTTTTACTGGTATTTGTAGTGACAGC<br>GGAGGAGGCTGGTGCTCAGGTAATCGTGATGA<br>AGTGATCCGGGCCATTTCGGGGCGCGTTTCTAGTC<br>TTTCCAATCCGCGCCCTGTACCAGTATTACGCC<br>GGACCGGTCTGCGCCGCGCCGCTTGTACCGC<br>CCTAAATGTCTAAGAGCGTCTAACAAAGC        |
| NLS0017 | ref1_142636 | 63        | GACGATATCGCTCATCTTCACTGCGATTGAAGCTG<br>GTGCCGTACTGCATAGGGATGAAAAGTGATGC<br>GGATAGACGGCTGACGGGAAAGCGCCTGGTCGA<br>TCGAAGACTTTGCTGACGAGGTGTGGTAGCCCC<br>GGATATAGGCATCGAAGGCCGGGACGTTGATCCC<br>ATCCTTTGCCTTATCTTGACTGGCGTCGTCGCGTG<br>CCGTGAGAACGGGCACGTCGAGGTCATCGAGG<br>CCAGCACCTTGCGGAACACCTGCGTTCCGCGGTT<br>GGGATTATCGACGGGAACGCGGTGGCCGC     |
| NLS0020 | ref3_25009  | 64        | GCCCTTCTGTGTCAGGCGATATTGTATAATGGCGTT<br>GCCCCAATAGAAGCAGCCATTGTCGCGAGGGCA<br>GCAGCGACGCTAGGTCGAAAGAGCATCCTAATCT<br>CGATCAAGATGCGACTGAGATTTCTGATGAAAAAT<br>ATCTAGACACAAGCAAAGCTGGTGAAATTACAA<br>CGATCATGGCGACAATTGCGGCCAATTCGGCCGG<br>AACTTGAAGGAACATAAAATGAATATTACAAA<br>TATACCGCAAAGCATGTAGAGTTGCTACACCAAG<br>GGTCGGGACGTCCAAAAAACTCACTGAGGA |

TABLE 7-continued

| Strain  | Fragment     | SEQ ID NO | Sequence                                                                                                                                                                                                                                                                                                                                          |
|---------|--------------|-----------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| NLS0020 | ref3_25219   | 65        | GGAACATAAAAAATGAATATTACAAATATACCGC<br>AAAGCATGTAGAGTTGCTACACCAAGGGTCGGG<br>ACGTCCAAAAAACTCACTGAGGAAGTCGACTG<br>GAAGCACGAGGCGCCCCCCCCAGGAGCGGGGCG<br>ACCGGCAAGGGGGCCCGCAATTGTGCCATGATC<br>GACCAGCTTAGGTAGGATCCTCTTTCGACCTAAC<br>GAATGGCTGCTTCTATTGGGGCAACGCCATTATA<br>CAATATCGCCTGACCATCTGGAACGCGGCCCGGT<br>CCACCGGCAGGTTGGCGACGACAGCGTCGGAG       |
| NLS0020 | ref1_4361220 | 66        | CGGCGTCGACCAGCCGGGCGAACTGCTTGGGCAT<br>GCTCTCCCGCGACGCCGGCCACAGCCGCGTCCCC<br>GTCCCTCCGCACAGGATCATCGGGTGGATTTGAA<br>AGGCAAAACGGGACATCAGGATAGGCCGCTCAG<br>GCGTTGGCGCTGAGGCGCTTGATGTCGGCGTCGA<br>CCATCTCGGTGATCAGCGCCTCGAGGCTGGTCTC<br>GGCTCCCGAGCCGAGGTGCGCTTGGCTTGGCG<br>GGGTTGCCCAGCAGCACCTCGACCTCTGCCGGCC<br>GGAACAGCGCCGGGTCGACGATCAGGTGG        |
| NLS0020 | ref1_4602420 | 67        | CTGGACATGCGCCCCACCCGGCCAAGTCCGACCG<br>CACCGGCAACCGCTCCTGTAGTCGTCGTCATCGT<br>TCTCACCCCTGAGGCGGAGACCGTCCGCTAACGG<br>GGTGCTCAAGCAACCGTGGGCGGAGGAACAC<br>GCACGTAGTCGCGTTTCAAGGTTGCGACGAACGC<br>CTCGGCCATGCCGTTGCTCTGCGGGCTCTCCAGC<br>GGCGTCGTTTTTGGCACCAACCAAGGTCGCGGG<br>CGAAGCGGCGCGTGTGCGGGGACTGTGAGGAA<br>TTTCGTGTGGGGCGGCCATAGTGGATCCG           |
| NLS0089 | ref1_194299  | 68        | GGAAATCGGCTTCAAGTACGACGTACGCCGGCC<br>ATGCAGGTCACGGGTGCACTGTTCATCTCGAGC<br>GGGACAACCAGCCGTTCCCTCGAAGCTGGAGTC<br>CGGCCTCGTCTTGGCGCAGGTGAGACACGCCACC<br>CAGGGCGCGGAAATCGGCCTGGCCGGCTATCTAA<br>CCGATTGGTGGCAGGTCTTGGCGGCTACGCTTA<br>TACCGAGGCACGCGTACTCTCGCCACTGGAAGAC<br>GATGGAGACGTGATCGCAGCAGGTAATCTCGTCG<br>GCAACGTTCCGCTAAATACTTTTCAGTCT         |
| NLS0089 | ref1_194305  | 69        | CGGCCTGGCCGGCTATCTAACCGATTGGTGGCAG<br>GTCTTTGGCGGCTACGCTTATACCGAGGCACGCG<br>TACTCTCGCCACTGGAAGACGATGGAGACGTGAT<br>CGCAGCAGGTAATCTCGTCGGCAACGTTCCGCTA<br>AATACTTTTCAGTCTGTTCAACAAGTTCGATATCA<br>ACGAGAATTTCTCCGTTGCTCTGGGCTATTACTAT<br>CAGGATGCCAGCTTTGCCCTCCTCAGACAATGCAG<br>TGCGTTTGCCAAGTTATTCGCGGTTTCGATGGCGG<br>GTTGTTCTATCGATTTCGACGAGTTGAC  |
| NLS0089 | ref1_194310  | 70        | ACGTTCCGCTAAATACTTTTCAGTCTGTTCAACAA<br>GTTTCGATATCAACGAGAATTTCTCCGTTGCTCTG<br>GGCTATTACTATCAGGATGCCAGCTTTGCCTCCTC<br>AGACAATGCAGTGCGTTTGGCCAAGTTATTCGCGG<br>TTCGATGGCGGGTTGTTCTATCGATTTCGACGAGT<br>TGACACGCGTTTCAGCTTAGCGTCGAGAACATTTT<br>CGACAGGCGTTACATCATCAACTCCAACAACAAAC<br>AACAACCTCACGCTGGCGCGCCGAGAACAGTCC<br>GCGTGCAATTGATCGCTCGGTTCTAAA |
| NLS0042 | ref1_86157   | 71        | AGCCCAACAAGCCTGATGCACCTTAACATACCTC<br>TAATGTGCGGCCAATTGCTTGGCGGCAGGGGAT<br>GTTGTATCGTCATAGGCTTGTCTAACCGGAACCTT<br>GTTTGCCAATCTCTTGGCGATCGCAACGCCAT<br>CTCGTGTTCTGTCACCATGTGCGCGTTCTCTAAT<br>TGCACTCATGGTGCCACGTGCACCTCCGATCGTC<br>TCGTGTCTAGAATGAAGGTGGGAACAACCTTACA<br>CAGGCTTTTCGCGACGCGCAATTTCTGGTTTCTCC<br>GCCTCGGATGTGGGTTTGAGCGCTTC         |
| NLS0042 | ref1_142469  | 72        | CTTTTCATTTGTCATGATCTCGACCAAGGTATTCA<br>CGGCAAGCTCGGTCTGTTGCTTAGCAAGTGCCCTG<br>AACTTCGGAACGATCGGCTCTCGACCTTCGGG<br>TTCGAGACCTGTCCCTTTTGAAACACGTCGCC                                                                                                                                                                                                |

TABLE 7-continued

| Strain  | Fragment     | SEQ ID NO | Sequence                                                                                                                                                                                                                                                                                                                                     |
|---------|--------------|-----------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|         |              |           | TACACTTTTCGGGATCAAGGTGCGGGTTGGCTTT<br>GGTCAAAATTCTCTGGCGTCCCATTACACGCCCT<br>CCGCATCATCGTTCCCGCGAAGCATCTGACCCCC<br>GACTTCCGCGAGGAAGCGTGTGGCGTGATCCTCG<br>AAGCGGAATGCCACCTCGAAGTGTTC                                                                                                                                                           |
| NLS0042 | ref1_142321  | 73        | CAGCAGCAAGCAGATCGTTGAAAACCGCTTGAA<br>CCGCATCTTGATCGGGACCGGAACCAATCAGGTC<br>ATCTAGGTAAACCGAGACGTAAACTCGTTTGCGC<br>TCGGCATCTTTCAGAACGTCCGTGATGCCAGACC<br>GCATTAGTACCATCGTCGCCAAGGCGGGCGACTG<br>AACGAAGCCGATCGGCAGAGTAACGGGGACC<br>GCCCCAATCGGGTTGCGAACGCAAGACCACTTA<br>GCAAAGGTTTCGAGCAGCGCCGAAGTTCGCATGGT<br>GGAGAGCCGCGCAACACGGTTCCGTGATA    |
| NLS0064 | ref1_153668  | 74        | TAGACATTCCAACAAACCGGCAAGAGGCTCGTCC<br>TCACTCGAGGATTTGTTGGGACTTGCAATGATGTC<br>GAAGCGGAGCCGTTATGACCTGGGTGCGATCATG<br>CGCCGAGCATGGGAGATGGCTCGGAGGCGGCA<br>TTCGCGGTTGGCGAGCGGGACGGACTCACCTTG<br>CTGCCGCGATGCGCAGCGCGTGGGCCGAAGCCA<br>AGTTGGCACTCGCGCCACGAAGACGGAGCAGG<br>ATCGTCTCTCTCCGAGCGACATGATCGGACATGA<br>GGACGCCCTACCAAGGCCGGGTTCTAAAAATAT |
| NLS0064 | ref1_3842117 | 75        | AAGATGGATACGACAAGCGCGATTACATTATTTG<br>CGAAATAGATGGACAAATAAAGACAAAGGACT<br>GATGTATTTCTTAAATCTGGACAAGTTGACCTCT<br>TTCACATAGAAGTCACCACTCCCTTTGGGACAAAT<br>TTGGTGTCACGAAAACATAGAGGCCGAAGTCTTT<br>AGCTGAATTATCGCGCTCCGGTCTTATGCGGC<br>TGAGTGAAGCGCGGACAGCTTGCGAGCAGGGC<br>CGCCAAATGGCAGCCGGATGACACAATGCTCGGT<br>CTCCCGACGCTTCTTCAATCGGAGCGCT      |
| NLS0064 | ref1_3842278 | 76        | AGCTGAATTATCGCGCTCCGGTCTTTATGCGGC<br>TGAGTGAAGCGCGGACAGCTTGCGAGCAGGGC<br>CGCCAATGGCAGCCGGATGACACAATGCTCGGT<br>CTCCCGACGCTTCTTCAATCGGAGCGCTTCGCA<br>GCCCCGGGCGGCGCTCATGCGTCACGACCTGG<br>GCCCTGCGCACCTTCGCGGCCCGCCGTCCCGGC<br>AGATCCCTGATGCCCCAAGTGGGCGGCCACTCCA<br>TCAAAGAACC CGGCTGTGGCAGATCTCGTAGG<br>CATACCGAGGTTCCGCAGTGCCCCACCC          |
| NLS0109 | ref1_135566  | 77        | ACGGTCACCCACGGACTGGGCGAGTACCTCACC<br>GGTGTCTATCATAACGCCGAGTTAGTTTTCGAC<br>CGTCCCTTATGCGATGTACCACCGGTGTCGGCAG<br>CCGATTTCTGTCACCGGAGCTGGCGTTCGGGT<br>TCAGACCACCATCATCGGTACGATGCTCGGATT<br>GGACACGGGGCCTTCATCTCCCCGGCGTGACTA<br>TAGGAAACGGCGCATCTCGGGGCCAGGCGG<br>TCGTCAAGAGATGTCCACCTATGCGGTAGT<br>TGCTGGCGTCCCCGCGACCGTACGACGAT              |
| NLS0109 | ref1_135772  | 78        | CCAATAAAGCGTTGGCCGCTGGGCAACCCGAT<br>CCGAGCCTAAGACTCAAAGCGCAAGCGAACACT<br>TGGTAGAGACAGCCCGCCGACTACGGCGTTCAG<br>CACTCTCCGGCTTGATCGGATAGGCATTGGTCA<br>AGGTGCCGTTGGTGATGACCTCGCCGCGCAAG<br>CGGCGAATTACTCGGATCAGCGGCCAGCACCTCG<br>ACCAAGTGTGCGAGCGCGACCAAAGGGCCACGT<br>TCGAGGACGTTTGAGGCGCGACCACTCTCGATAG<br>TCTCATCGTCGCGGCGAAGCTGCACCTCGA       |
| NLS0109 | ref1_169470  | 79        | CGATGGCACCGACCTGCCATGCCCTCTGCCGTCCG<br>CGCCAGAATGGTAAGAGGACGAAGGGGTAAG<br>GATCGTCGCTGCAGTGTGAGCAGCGACCAGAGA<br>AGGGGGCCGAACATCGGCATCAAACCTCGATTGC<br>CACTCGGACGCGAAGCGCGTCTTGAAGGAGGGA<br>TGGAAGCGAAACGGCCGAGAGTAACCGCCGAC<br>GAAAGATTGCACCCCTCATCGAGCAGGATCGGA                                                                              |



TABLE 7-continued

| Strain | Fragment | SEQ ID |                                                                       |
|--------|----------|--------|-----------------------------------------------------------------------|
|        |          | NO     | Sequence                                                              |
|        |          |        | GGTGAAGGCAAGCGTGGTTATTGGTAAGTGCA<br>AAAAATATAATGGTAGCGTCAGATCTAGCGTTC |

## REFERENCES

- ## REFERENCES
- [10102] Altschul S. F., Gish W., Miller W., Myers E. W., Lipman D. J. (1990) Basic local alignment search tool. *J. Mol. Biol.* 215:403-410
- [10103] Bentley D. R., Balasubramanian S., Swerdlow H. P., Smith G. P., Milton J., Brown D. G., Hall K. P., Evers D. J., Barnes C. L., Bignell H. R. et al. (2008) Accurate whole human genome sequencing using reversible terminator chemistry. *Nature* 456:53-59.
- [10104] Caporaso J. G., Lauber C. L., Walters W. A., Berg-Lyons D., Huntley J., Fierer N., Owens S. M., Betley J., Fraser L., Bauer M., Gormley N., Gilbert J. A., Smith G., Knight R. (2012) Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. *ISME J* 6:1621-1624.
- [10105] Comai et al. (1983) An altered *aroA* gene product confers resistance to the herbicide glyphosate. *Science* 221 (4608): 370-371.
- [10106] Green, P. N. 2005. *Methylobacterium*. In Brenner, D. J., N. R. Krieg, and J. T. Staley (eds.). "Bergey's Manual of Systematic Bacteriology. Volume two, The Proteobacteria. Part C, The alpha-, beta-, delta-, and epsilonproteobacteria." Second edition. Springer, New York. Pages 567-571.
- [10107] Green, P. N. 2006. *Methylobacterium*. In Dworkin, M., S. Falkow, E. Rosenberg, K. H. Schleifer, and E. Stackebrandt (eds.). "The Prokaryotes. A Handbook on the Biology of Bacteria. Volume 5. Proteobacteria: Alpha and Beta Subclasses." Third edition. Springer, New York. Pages 257-265.
- [10108] Green and Ardley (2018) Review of the genus *Methylobacterium* and closely related organisms: a proposal that some *Methylobacterium* species be reclassified into a new genus, *Methylorubrum* gen. nov. *International Journal of Systematic and Evolutionary Microbiology* 68:2727-2748.
- [10109] Konstantinidis, K. T. and Tiedje J. M. (2005) *Proc. Nat. Acad. Sci. USA* 102:2567-2572.
- [10110] Lidstrom, M. E. 2006. Aerobic methylotrophic prokaryotes. In Dworkin, M., S. Falkow, E.
- [10111] Rosenberg, K. H. Schleifer, and E. Stackebrandt (eds.). "The Prokaryotes. A Handbook on the Biology of Bacteria. Volume 2. Ecophysiology and biochemistry." Third edition. Springer, New York. Pages 618-634.
- [10112] Sanger F, Nicklen S, Coulson AR (1977) DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci USA.* 74 (12): 5463-5467
- [10113] Sy, A., Giraud, E., Jourand, P., Garcia, N., Willems, A., De Lajudic, P., Prin, Y., Neyra, M., Gillis, M., Boivin-Masson, C., and Dreyfus, B. 2001. Methylotrophic *Methylobacterium* Bacteria Nodulate and Fix Nitrogen in Symbiosis with Legumes. *Jour. Bacteriol.* 183 (1): 214-220.
- [10114] Vogel, H. J., and D. M. Bonner. 1956. Acetylornithinase of *Escherichia coli*: Partial purification and some properties. *J. Biol. Chem.* 218:97-106.
- [10115] Whittenbury, R., S. L. Davies, and J. F. Wilkinson. 1970. Enrichment, isolation and some properties of methane-utilizing bacteria. *J. Gen. Microbiol.* 61:205-218.
- [10116] Having illustrated and described the principles of the present disclosure, it should be apparent to persons skilled in the art that the disclosure can be modified in arrangement and detail without departing from such principles.
- [10117] Although the materials and methods have been described in terms of various embodiments and illustrative examples, it will be apparent to those of skill in the art that variations can be applied to the materials and methods described herein without departing from the concept, spirit and scope of the disclosure. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the disclosure as defined by the appended claims.

## SEQUENCE LISTING

```

Sequence total quantity: 79
SEQ ID NO: 1          moltype = DNA  length = 300
FEATURE
source                1..300
                      mol_type = genomic DNA
                      organism = Methylobacterium sp.

SEQUENCE: 1
ctcaacggttgc cgcgccttaac cgaagactaa ctaacataaa ttctcagttag ccgcgaaaca 60
gaatatgata cttggtcttta tacgacgcat agaggcgcat tccgtcaagt cgcgcattgc 120
agagtccccc cgaaaaattcg ttttctggga gcgagcagaa acgtgtgcac gtgtttctgtc 180
actctatttt ctggttttaga gcaatcatct cgcggctcag atcccgtcgt tccatagagt 240
ggtttccggt tcagtatgaa ttgtgagcag cccctcgggc tatatctact tcgacgtata 300

SEQ ID NO: 2          moltype = DNA  length = 300
FEATURE
source                1..300
                      mol_type = genomic DNA
                      organism = Methylobacterium sp.

```

-continued

---

SEQUENCE: 2  
 accaacaggc agaagaggct tacaaaaatg ctctagatat ctataatcgc gccggtctgg 60  
 ataataaggcg ggaaaagtca aatattctta tccgacttgg cgatgcagct agcgcaactga 120  
 gcaaatttat agacgcgaaa aatttctaca gtgaagggtt agcagtcga gcgcggtcat 180  
 gacggtgctt gccgactagt gtctgctcgc aattcagagc agcggttcg gctgggttcag 240  
 tttgggtatg ccttttctgt accagcctgt gactgggtctg gtatgtagcc cttgaccaat 300

SEQ ID NO: 3                   moltype = DNA   length = 300  
 FEATURE                   Location/Qualifiers  
 source                   1..300  
                           mol\_type = genomic DNA  
                           organism = Methylobacterium sp.

SEQUENCE: 3  
 atgaagctcc gtcgtgttta tgtaaaagac gttcgcagtt tctacgatgc tgaagaactc 60  
 attctggacg gagatatttc tattattata ggcccaaatg gtggaggaaa aacaaatcta 120  
 ctgatgcga cgatacactt gttgagaaga catttattgc aatcgtggcg tgtcgttcgc 180  
 catacataca cgccaataaa ctattccgat cacttccaac caaatgatca aatcacaaac 240  
 tatgcgctgg agaagtacag cggtcgcgaa agcgaagacc aagtcgttga gttcgatatc 300

SEQ ID NO: 4                   moltype = DNA   length = 300  
 FEATURE                   Location/Qualifiers  
 source                   1..300  
                           mol\_type = genomic DNA  
                           organism = Methylobacterium sp.

SEQUENCE: 4  
 cggggggcgt ctctcatat atggagggaa ttaatctgaa gggtcggagc ccccgctacg 60  
 gcacacacac ggcggcgggc ggtgggctcc ttctcatat acgcttcctg ataatttcga 120  
 ccgtaatcgc ccgcggcgcc agcggccaga ccttattagt ccagcatcac acggtgcac 180  
 gtatcgtgga gggcgaaaca ccgcctaagg ttccgccagt agccatccac atccgccctt 240  
 gccgctcgcg caagggaagc agaatgcttg atagcgactt cgcgggcgctc tttgtagacc 300

SEQ ID NO: 5                   moltype = DNA   length = 300  
 FEATURE                   Location/Qualifiers  
 source                   1..300  
                           mol\_type = genomic DNA  
                           organism = Methylobacterium sp.

SEQUENCE: 5  
 tgtggggcgt catggaacg ggggacgtgc ttttttgatc tctagccatt gtcacatcc 60  
 atcgtgccg gtagacaggcc ttacgcaca taacaacggt tatttcgatc agcccccctc 120  
 tgatgcgaca agtcagctat acagatgctt ctgaaagcca acaaggcggt tacgtatctg 180  
 cgtgacccca ccgagctcgc ctgcgcgcat gctgacagct ctgtatttta attcgaggag 240  
 gtggggcaat ttgacctggt caagctcgcc aacgtcttcc ttcacatatt gagacaggac 300

SEQ ID NO: 6                   moltype = DNA   length = 300  
 FEATURE                   Location/Qualifiers  
 source                   1..300  
                           mol\_type = genomic DNA  
                           organism = Methylobacterium sp.

SEQUENCE: 6  
 agagtgtatt tccccaagac tttatctctt tgctcatcac aaaacagccc gctcaagatg 60  
 atatgcgctg ggattttgtgc cgcgacccgt tgtaattgta tttacctctc ggctcatgtt 120  
 ccctgtcgcc agcgacaggg gaggtaaagt aatctcaact ttgcggccca tggctgatgc 180  
 tagctcagct acgcgacctt aggtaatatc tttgtgtccg cggagttcgc gatgaatgac 240  
 cgagcgattg acgcgcatgg tgcgggcaat atcggttga gttagaccgt tcttggtctc 300

SEQ ID NO: 7                   moltype = DNA   length = 300  
 FEATURE                   Location/Qualifiers  
 source                   1..300  
                           mol\_type = genomic DNA  
                           organism = Methylobacterium sp.

SEQUENCE: 7  
 acctgtctcg acatcggggg cagccacgag gcgtcggccg tggtagatct cggcgaagcg 60  
 gtagtagtgg gcgaggtccc catcgggacg gatgggggag tccggcgtag cttctccctc 120  
 ttctgcgatt agcgacaggg ctgcgcatgg cgtgggtgcg tcagtgatgg cgaacaacct 180  
 actatcgggg aaccaccgcc tgtcgacgag ctgggggtcc gggtttcctt tgaagacagc 240  
 gtcacccagc gcttcgatct tctcaatcaa ggctgcgtag aactgcccga tggctcgcaa 300

SEQ ID NO: 8                   moltype = DNA   length = 300  
 FEATURE                   Location/Qualifiers  
 source                   1..300  
                           mol\_type = genomic DNA  
                           organism = Methylobacterium sp.

SEQUENCE: 8  
 gacgtggtga cgggtggcacc agtcgcccag cgaagcggct actgtactga gcccgagtc 60  
 atgtcggctg aagaaatggt cgatatgccc cagggaagtc gggcgtaggt cgtaacctca 120  
 attcaccggc aggatcttgt ccagcaagcg gtccgaattc gcacgcacct gacttaggaa 180  
 attaccgagg taatcgctcg gtcggttcgc gtcggcgccc tgcacgcgca gcagcgctt 240

-continued

---

atgctcggtc gtcaactcac gagegaatat tgtccaggac aggttgtcga cctcggcgac 300

SEQ ID NO: 9                   moltype = DNA   length = 300  
 FEATURE                    Location/Qualifiers  
 source                     1..300  
                           mol\_type = genomic DNA  
                           organism = Methylobacterium sp.

SEQUENCE: 9  
 gagcatggca ggccaccacg catcgcttct tcatcacctt gctcggacga cttgctgctt 60  
 tgtcgggtctc tctctcaagg acgagacgct gcgacacttg cttcgtcaga gtgccaggat 120  
 taatccccgt caataccact actacgtcag ctggcggtgcg ccgggcggca cgaaggacgt 180  
 cgacgcggaa cgattgattc gagacgctaa ctttgacgtg tacaatttag tcacgctttt 240  
 cttagcgac gccagatag cctcgttggg ccgcttgctc accgccgatg cagccgagtt 300

SEQ ID NO: 10               moltype = DNA   length = 300  
 FEATURE                    Location/Qualifiers  
 source                     1..300  
                           mol\_type = genomic DNA  
                           organism = Methylobacterium sp.

SEQUENCE: 10  
 aacgcgatgg gagcagcagc taatgggctg catacaagat ataaaagaag gacttgaagg 60  
 tagaatagat aaccttaagg aaaatattga taatacagc tccagtatca catgcacaat 120  
 acaaggagtg ccgaagggtc agtcagaaaag aattttggga tgtgcaaacg cttcgtttga 180  
 ccaaatacag aaaatgtgga tgaatggtcc tacggatagg gctgcatttc tcgacgccgt 240  
 gaaaagtgtg atagaccgca acgcaagtaa tggaaaattc ggagttccag gatcatattg 300

SEQ ID NO: 11               moltype = DNA   length = 300  
 FEATURE                    Location/Qualifiers  
 source                     1..300  
                           mol\_type = genomic DNA  
                           organism = Methylobacterium sp.

SEQUENCE: 11  
 ttttcctttc gcgcctatga cctgagtgctg gccgaagctg tccgcttatt agagttttac 60  
 aaccagatgc gagcggctta gctcttctc cgttattaaag cagggcgcct cttcttaggg 120  
 gcgccccctc atatttaac tctgtctgat tctggcgcca tatcagacaa acatcagcgt 180  
 tgaccttgat tttgcatcta cattaggggtt gctcccagaa tggagccgcc gatgtccgtc 240  
 ccgtacctca agcgagagat gtgggggtgtg tattacatcc attggcgcca gggcggccgc 300

SEQ ID NO: 12               moltype = DNA   length = 300  
 FEATURE                    Location/Qualifiers  
 source                     1..300  
                           mol\_type = genomic DNA  
                           organism = Methylobacterium sp.

SEQUENCE: 12  
 gttggagatg gtgcccgttg caaccgtcag atcgccact gtgaatccgg tgacctcccg 60  
 agagaacgtg aaggttaactg tcgtagtctc gccaaactgcc aaagaggtgt ccgcaacggc 120  
 gatggtcgct gtcggggcgc ttgtttggac atcaacggtg agaccgcgcg aggcaggccc 180  
 cgtgttacca gcggtatcga cggccctagc ggtgaccgtg tgcgatccag ccgttagcgt 240  
 cgaaactggtg atgctatagg tcccgccgat cgccactgcc gagcccagaa tggtcgtgcc 300

SEQ ID NO: 13               moltype = DNA   length = 300  
 FEATURE                    Location/Qualifiers  
 source                     1..300  
                           mol\_type = genomic DNA  
                           organism = Methylobacterium sp.

SEQUENCE: 13  
 accgaaggcg tccccggaca cgaaggcctg aaacaccata tctgtggcga tcaggccgac 60  
 gtggtcgcgg acttcaactg gcagagaatg ccaggccgct tcgatttcag atgatactgg 120  
 tacggacata ggagcggctt agctttctca gtgcaaatgt gattgattcc ggtcaaaaa 180  
 tgatcttgat cggacgagac gttttcaatc catgtcgtgt tgccatcgcc gatcggtgcg 240  
 tcaagagaca gatggcgccg accgtagata cgcgcttcggg ttgcccgcac cgcttctcca 300

SEQ ID NO: 14               moltype = DNA   length = 300  
 FEATURE                    Location/Qualifiers  
 source                     1..300  
                           mol\_type = genomic DNA  
                           organism = Methylobacterium sp.

SEQUENCE: 14  
 ggaggtgtga tctgatgatg tgctggatga aattggcggt cgagcacttg ttcagcttgg 60  
 ccagctcgac gagatcggcg tgatgctcgg cgtcgatcag gatgttcagc gagaccggac 120  
 gtacgcagga cttggatata gcgcggttgc gcatcagctt gcagccttgc tctgcttctc 180  
 agcgtgccgc gtcaggatga cctgatgta gctgttgagg ttgatgcgt aatagcctgc 240  
 ggactctgtg agatccccgc gaagatcgct gcgcagggtc aggcggatgg tgctggtcgg 300

SEQ ID NO: 15               moltype = DNA   length = 300  
 FEATURE                    Location/Qualifiers

-continued

---

```

source                1..300
                      mol_type = genomic DNA
                      organism = Methylobacterium sp.

SEQUENCE: 15
aagtaaccgc tcaacatgat cttcagcatg ttgtccaaca gcaggagaat acatgtaatt 60
caccatgacc ggcaagctgc gactggccat tgcttcacac gcttgaatgt agcgatcgaa 120
tttcgcaaaa tcagggtgga atgaaaatat cgaaccaaac tgcgagcctt gaatccgttc 180
tgcaaaatta tcgaaaaaatt ttcttggccg actgcccgtt gaaaacattc ttacgtttac 240
atgcggcccg cctgaaaaca gacagtctac cagctctggg aaatgggggt gaagggtcgg 300

SEQ ID NO: 16          moltype = DNA length = 21
FEATURE               Location/Qualifiers
misc_feature           1..21
                      note = Primer
source                1..21
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 16
cacgtgttct gtcactctat t                                     21

SEQ ID NO: 17          moltype = DNA length = 20
FEATURE               Location/Qualifiers
misc_feature           1..20
                      note = Primer
source                1..20
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 17
gaaccggaaa ccactctatg                                     20

SEQ ID NO: 18          moltype = DNA length = 19
FEATURE               Location/Qualifiers
misc_feature           1..19
                      note = Probe
source                1..19
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 18
agcaatcatc tcgcggtc                                     19

SEQ ID NO: 19          moltype = DNA length = 21
FEATURE               Location/Qualifiers
misc_feature           1..21
                      note = Primer
source                1..21
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 19
gcgcactgag caaatTTata g                                     21

SEQ ID NO: 20          moltype = DNA length = 18
FEATURE               Location/Qualifiers
misc_feature           1..18
                      note = Primer
source                1..18
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 20
ctgaattgcg agcagaca                                     18

SEQ ID NO: 21          moltype = DNA length = 20
FEATURE               Location/Qualifiers
misc_feature           1..20
                      note = Probe
source                1..20
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 21
agtgaagggt tagcagtcgg                                     20

SEQ ID NO: 22          moltype = DNA length = 21
FEATURE               Location/Qualifiers
misc_feature           1..21
                      note = Primer
source                1..21
                      mol_type = other DNA

```

-continued

---

|                          |                                |    |
|--------------------------|--------------------------------|----|
| SEQUENCE: 22             | organism = synthetic construct |    |
| ctacttgatg cgacgataca c  |                                | 21 |
| SEQ ID NO: 23            | moltype = DNA length = 21      |    |
| FEATURE                  | Location/Qualifiers            |    |
| misc_feature             | 1..21                          |    |
|                          | note = Primer                  |    |
| source                   | 1..21                          |    |
|                          | mol_type = other DNA           |    |
|                          | organism = synthetic construct |    |
| SEQUENCE: 23             |                                |    |
| ggttggaagt gatcggaata g  |                                | 21 |
| SEQ ID NO: 24            | moltype = DNA length = 20      |    |
| FEATURE                  | Location/Qualifiers            |    |
| misc_feature             | 1..20                          |    |
|                          | note = Probe                   |    |
| source                   | 1..20                          |    |
|                          | mol_type = other DNA           |    |
|                          | organism = synthetic construct |    |
| SEQUENCE: 24             |                                |    |
| tcgttcgcca tacatacacg    |                                | 20 |
| SEQ ID NO: 25            | moltype = DNA length = 22      |    |
| FEATURE                  | Location/Qualifiers            |    |
| misc_feature             | 1..22                          |    |
|                          | note = Primer                  |    |
| source                   | 1..22                          |    |
|                          | mol_type = other DNA           |    |
|                          | organism = synthetic construct |    |
| SEQUENCE: 25             |                                |    |
| gtccttcct catatacgct tc  |                                | 22 |
| SEQ ID NO: 26            | moltype = DNA length = 22      |    |
| FEATURE                  | Location/Qualifiers            |    |
| misc_feature             | 1..22                          |    |
|                          | note = Primer                  |    |
| source                   | 1..22                          |    |
|                          | mol_type = other DNA           |    |
|                          | organism = synthetic construct |    |
| SEQUENCE: 26             |                                |    |
| ccgtgtgatg ctggactaat aa |                                | 22 |
| SEQ ID NO: 27            | moltype = DNA length = 20      |    |
| FEATURE                  | Location/Qualifiers            |    |
| misc_feature             | 1..20                          |    |
|                          | note = Probe                   |    |
| source                   | 1..20                          |    |
|                          | mol_type = other DNA           |    |
|                          | organism = synthetic construct |    |
| SEQUENCE: 27             |                                |    |
| taatttcgac cgtaatcgcc    |                                | 20 |
| SEQ ID NO: 28            | moltype = DNA length = 22      |    |
| FEATURE                  | Location/Qualifiers            |    |
| misc_feature             | 1..22                          |    |
|                          | note = Primer                  |    |
| source                   | 1..22                          |    |
|                          | mol_type = other DNA           |    |
|                          | organism = synthetic construct |    |
| SEQUENCE: 28             |                                |    |
| tgatcgaca agtcagctat ac  |                                | 22 |
| SEQ ID NO: 29            | moltype = DNA length = 20      |    |
| FEATURE                  | Location/Qualifiers            |    |
| misc_feature             | 1..20                          |    |
|                          | note = Primer                  |    |
| source                   | 1..20                          |    |
|                          | mol_type = other DNA           |    |
|                          | organism = synthetic construct |    |
| SEQUENCE: 29             |                                |    |
| cgagcttgac caggctaaat    |                                | 20 |
| SEQ ID NO: 30            | moltype = DNA length = 20      |    |
| FEATURE                  | Location/Qualifiers            |    |

-continued

---

|                           |                                |    |
|---------------------------|--------------------------------|----|
| misc_feature              | 1..20                          |    |
|                           | note = Probe                   |    |
| source                    | 1..20                          |    |
|                           | mol_type = other DNA           |    |
|                           | organism = synthetic construct |    |
| SEQUENCE: 30              |                                |    |
| acgcctttgt ttgctttcag     |                                | 20 |
| SEQ ID NO: 31             | moltype = DNA length = 20      |    |
| FEATURE                   | Location/Qualifiers            |    |
| misc_feature              | 1..20                          |    |
|                           | note = Primer                  |    |
| source                    | 1..20                          |    |
|                           | mol_type = other DNA           |    |
|                           | organism = synthetic construct |    |
| SEQUENCE: 31              |                                |    |
| cggagcgcatt ggaggtaaag    |                                | 20 |
| SEQ ID NO: 32             | moltype = DNA length = 22      |    |
| FEATURE                   | Location/Qualifiers            |    |
| misc_feature              | 1..22                          |    |
|                           | note = Primer                  |    |
| source                    | 1..22                          |    |
|                           | mol_type = other DNA           |    |
|                           | organism = synthetic construct |    |
| SEQUENCE: 32              |                                |    |
| ctccgcgcac acaaagatat ta  |                                | 22 |
| SEQ ID NO: 33             | moltype = DNA length = 19      |    |
| FEATURE                   | Location/Qualifiers            |    |
| misc_feature              | 1..19                          |    |
|                           | note = Probe                   |    |
| source                    | 1..19                          |    |
|                           | mol_type = other DNA           |    |
|                           | organism = synthetic construct |    |
| SEQUENCE: 33              |                                |    |
| aatctcaact ttgcggccc      |                                | 19 |
| SEQ ID NO: 34             | moltype = DNA length = 22      |    |
| FEATURE                   | Location/Qualifiers            |    |
| misc_feature              | 1..22                          |    |
|                           | note = Primer                  |    |
| source                    | 1..22                          |    |
|                           | mol_type = other DNA           |    |
|                           | organism = synthetic construct |    |
| SEQUENCE: 34              |                                |    |
| cttctccctc ttctgcgcatt ag |                                | 22 |
| SEQ ID NO: 35             | moltype = DNA length = 21      |    |
| FEATURE                   | Location/Qualifiers            |    |
| misc_feature              | 1..21                          |    |
|                           | note = Primer                  |    |
| source                    | 1..21                          |    |
|                           | mol_type = other DNA           |    |
|                           | organism = synthetic construct |    |
| SEQUENCE: 35              |                                |    |
| gacgctgtct tcaaaggaaa c   |                                | 21 |
| SEQ ID NO: 36             | moltype = DNA length = 19      |    |
| FEATURE                   | Location/Qualifiers            |    |
| misc_feature              | 1..19                          |    |
|                           | note = Probe                   |    |
| source                    | 1..19                          |    |
|                           | mol_type = other DNA           |    |
|                           | organism = synthetic construct |    |
| SEQUENCE: 36              |                                |    |
| tgatggcgaa caacctact      |                                | 19 |
| SEQ ID NO: 37             | moltype = DNA length = 21      |    |
| FEATURE                   | Location/Qualifiers            |    |
| misc_feature              | 1..21                          |    |
|                           | note = Primer                  |    |
| source                    | 1..21                          |    |
|                           | mol_type = other DNA           |    |
|                           | organism = synthetic construct |    |
| SEQUENCE: 37              |                                |    |

-continued

---

|                            |                                |
|----------------------------|--------------------------------|
| gcacgcacct gacttaggaa a    | 21                             |
| SEQ ID NO: 38              | moltype = DNA length = 20      |
| FEATURE                    | Location/Qualifiers            |
| misc_feature               | 1..20                          |
|                            | note = Primer                  |
| source                     | 1..20                          |
|                            | mol_type = other DNA           |
|                            | organism = synthetic construct |
| SEQUENCE: 38               |                                |
| gagttgacga ccgagcataa      | 20                             |
| SEQ ID NO: 39              | moltype = DNA length = 19      |
| FEATURE                    | Location/Qualifiers            |
| misc_feature               | 1..19                          |
|                            | note = Probe                   |
| source                     | 1..19                          |
|                            | mol_type = other DNA           |
|                            | organism = synthetic construct |
| SEQUENCE: 39               |                                |
| aggtaatcgc tcggtccgt       | 19                             |
| SEQ ID NO: 40              | moltype = DNA length = 22      |
| FEATURE                    | Location/Qualifiers            |
| misc_feature               | 1..22                          |
|                            | note = Primer                  |
| source                     | 1..22                          |
|                            | mol_type = other DNA           |
|                            | organism = synthetic construct |
| SEQUENCE: 40               |                                |
| ctttgtcggc ctctctctca ag   | 22                             |
| SEQ ID NO: 41              | moltype = DNA length = 23      |
| FEATURE                    | Location/Qualifiers            |
| misc_feature               | 1..23                          |
|                            | note = Primer                  |
| source                     | 1..23                          |
|                            | mol_type = other DNA           |
|                            | organism = synthetic construct |
| SEQUENCE: 41               |                                |
| cagctgacgt agtagtggtta ttg | 23                             |
| SEQ ID NO: 42              | moltype = DNA length = 19      |
| FEATURE                    | Location/Qualifiers            |
| misc_feature               | 1..19                          |
|                            | note = Probe                   |
| source                     | 1..19                          |
|                            | mol_type = other DNA           |
|                            | organism = synthetic construct |
| SEQUENCE: 42               |                                |
| tcgtcagagt gccaggatt       | 19                             |
| SEQ ID NO: 43              | moltype = DNA length = 24      |
| FEATURE                    | Location/Qualifiers            |
| misc_feature               | 1..24                          |
|                            | note = Primer                  |
| source                     | 1..24                          |
|                            | mol_type = other DNA           |
|                            | organism = synthetic construct |
| SEQUENCE: 43               |                                |
| gttccagtat cacatgcaca atac | 24                             |
| SEQ ID NO: 44              | moltype = DNA length = 21      |
| FEATURE                    | Location/Qualifiers            |
| misc_feature               | 1..21                          |
|                            | note = Primer                  |
| source                     | 1..21                          |
|                            | mol_type = other DNA           |
|                            | organism = synthetic construct |
| SEQUENCE: 44               |                                |
| cgtaggacca ttcattccaca t   | 21                             |
| SEQ ID NO: 45              | moltype = DNA length = 19      |
| FEATURE                    | Location/Qualifiers            |
| misc_feature               | 1..19                          |
|                            | note = Probe                   |

-continued

---

|                          |                                |    |
|--------------------------|--------------------------------|----|
| source                   | 1..19                          |    |
|                          | mol_type = other DNA           |    |
|                          | organism = synthetic construct |    |
| SEQUENCE: 45             |                                |    |
| tttgggatgt gcaaacgct     |                                | 19 |
| SEQ ID NO: 46            | moltype = DNA length = 22      |    |
| FEATURE                  | Location/Qualifiers            |    |
| misc_feature             | 1..22                          |    |
|                          | note = Primer                  |    |
| source                   | 1..22                          |    |
|                          | mol_type = other DNA           |    |
|                          | organism = synthetic construct |    |
| SEQUENCE: 46             |                                |    |
| gcttagctct tcctccgta tt  |                                | 22 |
| SEQ ID NO: 47            | moltype = DNA length = 21      |    |
| FEATURE                  | Location/Qualifiers            |    |
| misc_feature             | 1..21                          |    |
|                          | note = Primer                  |    |
| source                   | 1..21                          |    |
|                          | mol_type = other DNA           |    |
|                          | organism = synthetic construct |    |
| SEQUENCE: 47             |                                |    |
| attctgggag caacccta g    |                                | 21 |
| SEQ ID NO: 48            | moltype = DNA length = 20      |    |
| FEATURE                  | Location/Qualifiers            |    |
| misc_feature             | 1..20                          |    |
|                          | note = Probe                   |    |
| source                   | 1..20                          |    |
|                          | mol_type = other DNA           |    |
|                          | organism = synthetic construct |    |
| SEQUENCE: 48             |                                |    |
| atctgtctg atgtctggcg     |                                | 20 |
| SEQ ID NO: 49            | moltype = DNA length = 22      |    |
| FEATURE                  | Location/Qualifiers            |    |
| misc_feature             | 1..22                          |    |
|                          | note = Primer                  |    |
| source                   | 1..22                          |    |
|                          | mol_type = other DNA           |    |
|                          | organism = synthetic construct |    |
| SEQUENCE: 49             |                                |    |
| cgtgaaggta actgtcgtag tc |                                | 22 |
| SEQ ID NO: 50            | moltype = DNA length = 21      |    |
| FEATURE                  | Location/Qualifiers            |    |
| misc_feature             | 1..21                          |    |
|                          | note = Primer                  |    |
| source                   | 1..21                          |    |
|                          | mol_type = other DNA           |    |
|                          | organism = synthetic construct |    |
| SEQUENCE: 50             |                                |    |
| caccgttgat gtccaaacaa g  |                                | 21 |
| SEQ ID NO: 51            | moltype = DNA length = 19      |    |
| FEATURE                  | Location/Qualifiers            |    |
| misc_feature             | 1..19                          |    |
|                          | note = Probe                   |    |
| source                   | 1..19                          |    |
|                          | mol_type = other DNA           |    |
|                          | organism = synthetic construct |    |
| SEQUENCE: 51             |                                |    |
| aactgcaaaa gaggtgtcc     |                                | 19 |
| SEQ ID NO: 52            | moltype = DNA length = 22      |    |
| FEATURE                  | Location/Qualifiers            |    |
| misc_feature             | 1..22                          |    |
|                          | note = Primer                  |    |
| source                   | 1..22                          |    |
|                          | mol_type = other DNA           |    |
|                          | organism = synthetic construct |    |
| SEQUENCE: 52             |                                |    |
| cataggagcg gcttagcttt ct |                                | 22 |



-continued

---

|                             |                                |    |
|-----------------------------|--------------------------------|----|
| SEQ ID NO: 53               | moltype = DNA length = 22      |    |
| FEATURE                     | Location/Qualifiers            |    |
| misc_feature                | 1..22                          |    |
|                             | note = Primer                  |    |
| source                      | 1..22                          |    |
|                             | mol_type = other DNA           |    |
|                             | organism = synthetic construct |    |
| SEQUENCE: 53                |                                |    |
| gatggcaaca cgacatggat tg    |                                | 22 |
| SEQ ID NO: 54               | moltype = DNA length = 21      |    |
| FEATURE                     | Location/Qualifiers            |    |
| misc_feature                | 1..21                          |    |
|                             | note = Probe                   |    |
| source                      | 1..21                          |    |
|                             | mol_type = other DNA           |    |
|                             | organism = synthetic construct |    |
| SEQUENCE: 54                |                                |    |
| cagtgcaaat gtgattgatt c     |                                | 21 |
| SEQ ID NO: 55               | moltype = DNA length = 24      |    |
| FEATURE                     | Location/Qualifiers            |    |
| misc_feature                | 1..24                          |    |
|                             | note = Primer                  |    |
| source                      | 1..24                          |    |
|                             | mol_type = other DNA           |    |
|                             | organism = synthetic construct |    |
| SEQUENCE: 55                |                                |    |
| gacgtacgca ggacttggtat ttag |                                | 24 |
| SEQ ID NO: 56               | moltype = DNA length = 24      |    |
| FEATURE                     | Location/Qualifiers            |    |
| misc_feature                | 1..24                          |    |
|                             | note = Primer                  |    |
| source                      | 1..24                          |    |
|                             | mol_type = other DNA           |    |
|                             | organism = synthetic construct |    |
| SEQUENCE: 56                |                                |    |
| ctcaacagct acatcagggt catc  |                                | 24 |
| SEQ ID NO: 57               | moltype = DNA length = 20      |    |
| FEATURE                     | Location/Qualifiers            |    |
| misc_feature                | 1..20                          |    |
|                             | note = Probe                   |    |
| source                      | 1..20                          |    |
|                             | mol_type = other DNA           |    |
|                             | organism = synthetic construct |    |
| SEQUENCE: 57                |                                |    |
| ttgctctgct tctcagcgtg       |                                | 20 |
| SEQ ID NO: 58               | moltype = DNA length = 20      |    |
| FEATURE                     | Location/Qualifiers            |    |
| misc_feature                | 1..20                          |    |
|                             | note = Primer                  |    |
| source                      | 1..20                          |    |
|                             | mol_type = other DNA           |    |
|                             | organism = synthetic construct |    |
| SEQUENCE: 58                |                                |    |
| aacaaaactg cgagccttga       |                                | 20 |
| SEQ ID NO: 59               | moltype = DNA length = 21      |    |
| FEATURE                     | Location/Qualifiers            |    |
| misc_feature                | 1..21                          |    |
|                             | note = Primer                  |    |
| source                      | 1..21                          |    |
|                             | mol_type = other DNA           |    |
|                             | organism = synthetic construct |    |
| SEQUENCE: 59                |                                |    |
| gggccgcatg taaacgtaag a     |                                | 21 |
| SEQ ID NO: 60               | moltype = DNA length = 17      |    |
| FEATURE                     | Location/Qualifiers            |    |
| misc_feature                | 1..17                          |    |
|                             | note = Probe                   |    |
| source                      | 1..17                          |    |
|                             | mol_type = other DNA           |    |

-continued

---

organism = synthetic construct

SEQUENCE: 60  
ttcttgccg actgccg 17

SEQ ID NO: 61                   moltype = DNA   length = 300  
FEATURE                       Location/Qualifiers  
source                         1..300  
                              mol\_type = genomic DNA  
                              organism = Methylobacterium sp.

SEQUENCE: 61  
gcaaaacgac ctaatagttc tacagcggca tgcgccaaagt cagcgcggtg aacagtatac 60  
ctgggagcaa cttgtcctcc gaaacccaca taaaacaaat tactcctggc agtgcccagt 120  
ccatcaaaat cgaatacaat atttctcgag gaggcacatg taatagcctg ccaaagcaac 180  
aaagctatgg cgcggttatg actttcattg cttctggtag acataaaata atatgccgat 240  
tgtgatcccc aaatgtagaa tattgccgca tcaattgcgc caagtttatt tcggatcgat 300

SEQ ID NO: 62                   moltype = DNA   length = 300  
FEATURE                       Location/Qualifiers  
source                         1..300  
                              mol\_type = genomic DNA  
                              organism = Methylobacterium sp.

SEQUENCE: 62  
ggcgccaacg gtatgatcgc atgattttcc tgcggcatag cttgcgggaa tggcgtattt 60  
ggcgtctctc tcaggaattt ctaagggcat acgcaggaac tctacagcac ttttactggt 120  
atttttagt gacagcggag gaggtggtg ctcaaggtaa tcgtgatgaa gtgatccggg 180  
ccattcgggg cgcgtttcta gtctttccaa tccgcgccct gtaccacgta ttacgccgga 240  
ccggtctcgc cgcgcgcccc ctcttgaccg ccctaaatgt ctaagagcgt ctaacaaagc 300

SEQ ID NO: 63                   moltype = DNA   length = 300  
FEATURE                       Location/Qualifiers  
source                         1..300  
                              mol\_type = genomic DNA  
                              organism = Methylobacterium sp.

SEQUENCE: 63  
gacgatatcg ctcattctta ctgcattgaa gctgggtgcg tactgcatag ggatgaaaaa 60  
gtgatgcgga tagacgctg acgggaaagc gcctgggtcga tcgaagactt tgctgacgag 120  
gttggtgtag aaccggatat aggcattcga gcccgggacg ttgatcccat cctttgcctt 180  
atcttgactg gcgtcgtcgc gtgcgcgtcag aacggggcacg tcgcaggtea tcgaggccag 240  
caccttgccg aacacctgcg ttccgcgctt gggattatcg acggcgaacg cgggtggccgc 300

SEQ ID NO: 64                   moltype = DNA   length = 300  
FEATURE                       Location/Qualifiers  
source                         1..300  
                              mol\_type = genomic DNA  
                              organism = Methylobacterium sp.

SEQUENCE: 64  
gcccttctgt caggcgatat tgtataatgg cgttgcccca atagaagcag ccattcgtgc 60  
gagggcagca gcgacgctag gtgcgaagag catcctaate tcgatcaaga tgcgactgag 120  
atttctgatg aaaatatcta gacacaagca aagctggtga aattacaacg atcatggcga 180  
caattgcccc caattcggcc ggaacttgaa ggaacataaa aatgaatatt acaaatatac 240  
cgcaagcat gtagagtgc tacaccaagg gtccgggacgt ccaaaaaaac tactgagga 300

SEQ ID NO: 65                   moltype = DNA   length = 300  
FEATURE                       Location/Qualifiers  
source                         1..300  
                              mol\_type = genomic DNA  
                              organism = Methylobacterium sp.

SEQUENCE: 65  
ggaacataaa aatgaatatt acaaatatac cgcaagcat gtagagttgc tacaccaagg 60  
gtcgggacgt ccaaaaaaac tcactgagga agtcgactgg aagcacgagg cgccccccc 120  
aggagcgggg cgaccggcaa gggggcccgcc aattgtcgcc atgatcgacc agcttaggta 180  
ggatcctctt tcgacctaac gaatggctgc ttctattggg gcaacgccat tatacaatat 240  
cgctgacca tctggaacgc ggcccgttcc accggcagggt tggcgacgac agcgtcgga 300

SEQ ID NO: 66                   moltype = DNA   length = 300  
FEATURE                       Location/Qualifiers  
source                         1..300  
                              mol\_type = genomic DNA  
                              organism = Methylobacterium sp.

SEQUENCE: 66  
cggcgtcgac cagccggggc aactgcttgg gcatgctctc ccgcgacgcc ggccacagcc 60  
gcgtccccgt cctcccgcac aggatcatcg ggtggatttg aaaggcaaaa cgggacatca 120  
ggataggccg ctcaggcggt ggcgctgagg cgcttgatgt cggcgtcgac catctcggtg 180  
atcagcgccg cgaggctggt ctgcgcctcc cagccgaagg tcgccttggc cttggcgggg 240  
ttgcccgaca gcaacctgac ctctgcggc cggaacacgc ccgggtcgac gatcaggtag 300

-continued

---

SEQ ID NO: 67                   moltype = DNA   length = 300  
 FEATURE                       Location/Qualifiers  
 source                        1..300  
                               mol\_type = genomic DNA  
                               organism = Methylobacterium sp.

SEQUENCE: 67

```

ctggacatgc gccacccecg gccaaagtcc accgcaccgg caaccgctcc tgtagtctgc 60
gtcatcgctc tcacccctga ggcggagacc gtccgctaac ggggtgtctc aagcaaccgt 120
ggggcggagg aacacgcacg tagtcgcgtt tcaaggttcg cacgaacgcc tcggccatgc 180
cgttgctctg cgggctctcc agcggcgctc tttttggcac caaaccaagg tcggggcgga 240
agcgcgcgct gtcgcgggga ctgtcaggaa ttctgtgtgg gggcgcccat agtggatccg 300

```

SEQ ID NO: 68                   moltype = DNA   length = 300  
 FEATURE                       Location/Qualifiers  
 source                        1..300  
                               mol\_type = genomic DNA  
                               organism = Methylobacterium sp.

SEQUENCE: 68

```

ggaaatcggc ttcaagtacg acgtcacgcc ggccatgcag gtcacgggtg cactgttcaa 60
tctcgagcgc gacaaccagc cgttcccttc gaacgtggag tccggcctcg tccttggcgc 120
aggtcagaca cgcaccacgg gcgcggaaat cggcctggcc ggctatctaa ccgattgggtg 180
gcaggtcttt ggcgggtacg cttataccga ggacgcgcta ctctgccac tggaaagaca 240
tggagacgtg atcgcagcag gtaatctcgt cggcaacgtt ccgctaaata ctttcagtct 300

```

SEQ ID NO: 69                   moltype = DNA   length = 300  
 FEATURE                       Location/Qualifiers  
 source                        1..300  
                               mol\_type = genomic DNA  
                               organism = Methylobacterium sp.

SEQUENCE: 69

```

cggcctggcc ggctatctaa ccgattgggt gcaggtcttt ggcggctacg cttataccga 60
ggcaccgcgt ctctgccacg tggaaagaca tggagacgtg atcgcagcag gtaatctcgt 120
cggcaacggt ccgctaaata ctttcagtct gttcaacaag ttcgatatca acgagaattt 180
ctccgttctg ctgggctatt actatcagga tgccagcttt gcctcctcag acaatgcagt 240
gcgtttgcca agttattcgc ggttcgatgg cgggttgttc tatcgattcg acgagttgac 300

```

SEQ ID NO: 70                   moltype = DNA   length = 300  
 FEATURE                       Location/Qualifiers  
 source                        1..300  
                               mol\_type = genomic DNA  
                               organism = Methylobacterium sp.

SEQUENCE: 70

```

acgttcgcgt aaatactttc agtctgttca acaagttcga tatcaacgag aattttctccg 60
ttgtctctgg ctattactat caggatgcca gctttgcctc ctcagacaat gcagtgcgtt 120
tgccaagtta ttccggttcc gatggcgggt tgttctatcg attcgacgag ttgacacgcg 180
ttcagcttag cgtcgagaaac attttcgaca ggcgtttacat catcaactcc aacaacaaca 240
acaacctcac gcctggcgcg ccgagaacag tccgcgtgca attgatcgct cggttctaaa 300

```

SEQ ID NO: 71                   moltype = DNA   length = 300  
 FEATURE                       Location/Qualifiers  
 source                        1..300  
                               mol\_type = genomic DNA  
                               organism = Methylobacterium sp.

SEQUENCE: 71

```

agccacacaag cctgatgcac ttaactacat cctctaattg cgcgccaat tgcttggcgg 60
caggggatgtg tgtatcgtea taggcttgtc taaccggaac ttgtttgcca atctctttgg 120
cgatcgcaac cgcctatctc tgttcgtcaa ccatgtgcgc gttcctctaa ttgacctcat 180
ggtgccacgt gcacctccga tcgtctcgtg tctagaatga aggtgggaac aaccttacac 240
aggctttcgc gacgcgcgaa tttctggttt ctccgcctcg gatgtgggtt tgagcgcttc 300

```

SEQ ID NO: 72                   moltype = DNA   length = 300  
 FEATURE                       Location/Qualifiers  
 source                        1..300  
                               mol\_type = genomic DNA  
                               organism = Methylobacterium sp.

SEQUENCE: 72

```

cttttcattt gtcatgatct cgaccaaggt attcacggca agctcggtct gttgcttagc 60
aagtgcctga acttcgcgaa cgatcggctc tcgacccttc gggttcgaga cctgtccctt 120
ttgaaaacca cgtgccctac acttttcggg atcaaggtgc gggttggctt tggtaaaat 180
tctctggcgt cccattacac gccctccgca tcactgttcc cgcgaacgat ctgacccccc 240
acttcgcgga ggaagcgtgt ggcgtgatcc tcgaagcgga atgccacctc gaactgttcc 300

```

SEQ ID NO: 73                   moltype = DNA   length = 300  
 FEATURE                       Location/Qualifiers  
 source                        1..300  
                               mol\_type = genomic DNA

-continued

---

```

                                organism = Methylobacterium sp.
SEQUENCE: 73
cagcagcaag cagatcgttg aaaaccgctt gaaccgcacg ttgatcgga cgggaaccaa 60
tcagggtcac taggtaaacc gagacgtaaa ctcggttgcg ctccgcatct ttcagaacgt 120
ccgtgatgcc agacgcgatt agtaccatcg tcgccaaggc gggcgactga acgaagccga 180
tcggcagaga gtaacgggga cggccctaa tcgggttgcg aacgcaagac cacttagcaa 240
aggttcgagc acggccgaac ttcgcatggt ggagagccgc ggcaacacgg ttccgtgata 300

SEQ ID NO: 74      moltype = DNA length = 300
FEATURE           Location/Qualifiers
source            1..300
                  mol_type = genomic DNA
                  organism = Methylobacterium sp.
SEQUENCE: 74
tagacattcc aacaaaccgg caagaggctc gtccctactc gaggatttgt tgggacttgc 60
atgatgtcga agcggagccg ttatgacctg ggtgcatca tgcgccgagc atgggagatg 120
gctcggggagg cggcattcgc ggttgccgag cgggcacgga ctacacctgc tgcgcgatg 180
cgacgctcgc gggccgaagc caagttggca ctccgcccc cgaagacgga gcaggatcgt 240
ctctctccga gcgacatgat cggacatgag gacgcctacc aaggccgggt tctaaaaat 300

SEQ ID NO: 75      moltype = DNA length = 300
FEATURE           Location/Qualifiers
source            1..300
                  mol_type = genomic DNA
                  organism = Methylobacterium sp.
SEQUENCE: 75
aagatggata cgacaagcgc gattacatta ttgcgaaat agatggacaa ataaaagaca 60
aaggactgat gtatttcctt aaatctggac aagttgacct cttcacata gaagtcacca 120
ctccctttgg gacaatttgg tgtcacgaaa acatagaggg cgaactctct agctgaatta 180
tcgcgctcgc ggttcttatg cggctgagtg aagcgcggga cagcttgcca gcaggccgcg 240
caatggcagc cgggatgaca caatgctcgg tctcccgacg cttcttcaat cgggagcgct 300

SEQ ID NO: 76      moltype = DNA length = 299
FEATURE           Location/Qualifiers
source            1..299
                  mol_type = genomic DNA
                  organism = Methylobacterium sp.
SEQUENCE: 76
agctgaatta tcgcgctcgc ggttcttatg cggctgagtg aagcgcggga cagcttgcca 60
gcaggccgcg caatggcagc cgggatgaca caatgctcgg tctcccgacg cttcttcaat 120
cgggagcgct tcgcagccgc gggcggcgcg ctcatgcgtc acgacctggg ccttgccgac 180
cttcgcggcc ccgcgcgtcc gccagatccc tgatgcccc aagtgggcgc cactccatca 240
aagaaccccg gctctgtggc gatctcgtag gcataccgag gttccgcagt gcccccacc 299

SEQ ID NO: 77      moltype = DNA length = 300
FEATURE           Location/Qualifiers
source            1..300
                  mol_type = genomic DNA
                  organism = Methylobacterium sp.
SEQUENCE: 77
acggtcaccc caccgactgg gcgagtacct caccgggtgt ctatcataac gccgagttag 60
ttttcgaccg tcccttatgc gatgtaccac cgggtgctcg agccgatttc gtcccaccg 120
gagctggcgt tccggttcag accaccatca tcggtcacga tgtctggatt ggacacgggg 180
ccttcacatc ccccgcgctg actataggaa acggcgcgat cgtcggggcc caggcggctg 240
tcacaagaga tgtcccacc tatgcggtag ttgctggcgt ccccgcgacc gtacgacgat 300

SEQ ID NO: 78      moltype = DNA length = 300
FEATURE           Location/Qualifiers
source            1..300
                  mol_type = genomic DNA
                  organism = Methylobacterium sp.
SEQUENCE: 78
ccaataaaag cggtggccgc ctgggcaacc cgatccgagc ctaagactca aagcgcaagc 60
gaacacttgg tagagacagc ccgccgacta cggcggtcca gcaactctcc gctttgatcg 120
gataggcatt ggtcaagggt cgggtggtga tgacctcgcc cggcgcaagc ggcgaattac 180
tcggatcagc ggccagcacc tcgaccaagt gtcggagcgc gaccaaaggg ccacgttcga 240
ggacgtttga ggcgcgacca gtctcgatag tctcatcgct cggcgcaagc tgcacctcga 300

SEQ ID NO: 79      moltype = DNA length = 300
FEATURE           Location/Qualifiers
source            1..300
                  mol_type = genomic DNA
                  organism = Methylobacterium sp.
SEQUENCE: 79
cgatggcacc gacctgccat gccctgcccg tccgcgccag aatggtaaag aggacgaagg 60
gggtaaggat cgtcgctgca gtgttgagca gcgaccagag aagggggccc aacatcggca 120

```

-continued

---

```

tcaaacctcg attgccactc ggaacgcgaag cgcgtcttga aggagggatg gaagcgaaac 180
ggccgcagag taacgcgcga cgaaagattg caccctcat cgagcaggat cggaggtgaa 240
ggcaagcgtg gggtattggt aagtgcacaaa aatataatgg tagcgtcaga tctagcgttc 300

```

---

**1.-22.** (canceled)

**23.** A soybean plant or soybean plant part that is coated or partially coated with a composition comprising a *Methylobacterium* selected from the group consisting of consisting of: NLS0497 (NRRL B-67925), NLS0693 (NRRL B-67926), NLS1179 (NRRL B-67929), NLS1181 (NRRL B-67927), or NLS0610 (NRRL B-67892), NLS0109 (NRRL B-67340) and NLS0610 (NRRL B-67892), and variants thereof.

**24.** The soybean plant or soybean plant part of claim **23**, wherein the composition further comprises at least one additional component selected from the group consisting of an additional active ingredient, an agriculturally acceptable adjuvant, and an agriculturally acceptable excipient.

**25.** The soybean plant or soybean plant part of claim **23**, wherein the composition comprises the *Methylobacterium* at a titer of about  $1 \times 10^6$  CFU/gm to about  $1 \times 10^{14}$  CFU/gm for a solid composition or at a titer of about  $1 \times 10^6$  CFU/mL to about  $1 \times 10^{11}$  CFU/mL for a liquid composition.

**26.** The soybean plant or soybean plant part of claim **23**, wherein the *Methylobacterium* is NLS0610 (NRRL B-67892), or a variant thereof.

**27.** The soybean plant or soybean plant part of claim **23**, wherein the soybean plant part is selected from the group consisting of a seed, a stem, a flower, a leaf, a petiole, a pod, and an axillary bud.

**28.** The soybean plant or soybean plant part of claim **23**, wherein the composition is applied by spraying, coating, partially coating, immersing, and/or imbibing the soybean plant or plant part with the composition.

**29.** The soybean plant or soybean plant part of claim **24**, wherein the additional component is an additional active ingredient.

**30.** The soybean plant or soybean plant part of claim **29**, wherein the additional active ingredient is selected from the group consisting of a fungicide, insecticide, nematocide, and an additional biological.

**31.** The soybean plant or soybean plant part of claim **29**, wherein the additional biological is a biocontrol agent.

**32.** The soybean plant or soybean plant part of claim **29**, wherein the additional active ingredient:

(a) is selected from the group consisting of clothianidin, *Bacillus firmus*, abamectin, thiamethoxam, imidacloprid, azoxystrobin, fluopyram, fluoxastrobin, ipconazole, mefenoxam, metalaxyl, penflufen, prothioconazole, pyraclostrobin, and sedaxane;

(b) is selected from the group consisting of NLS0046 (NRRL B-50929), NLS0020 (NRRL B-50930), NLS0017 (NRRL B-50931), NLS0042 (NRRL B-50932), NLS0089 (NRRL B-50933), NLS0068 (NRRL B-50934), NLS0065 (NRRL B-50935), NLS0069 (NRRL B-50936), NLS0062 (NRRL B-50937), NLS0064 (NRRL B-50938), NLS0021 (NRRL B-50939), NLS0066 (NRRL B-50940), NLS0037 (NRRL B-50941), NLS0038 (NRRL B-50942), NLS0109 (NRRL B-67340), and variants thereof; and/or

(c) is a bacterium of the genus *Actinomycetes*, *Agrobacterium*, *Arthrobacter*, *Alcaligenes*, *Aureobacterium*, *Azobacter*, *Bacillus*, *Beijerinckia*, *Brevibacillus*, *Burkholderia*, *Chromobacterium*, *Clostridium*, *Clavibacter*, *Comomonas*, *Corynebacterium*, *Curtobacterium*, *Enterobacter*, *Flavobacterium*, *Gluconobacter*, *Hydrogenophaga*, *Klebsiella*, *Paenibacillus*, *Pasteuria*, *Phingobacterium*, *Photorhabdus*, *Phyllobacterium*, *Pseudomonas*, *Rhizobium*, *Bradyrhizobium*, *Serratia*, *Stenotrophomonas*, *Variovorax*, or *Xenorhabdus*.

**33.-34.** (canceled)

\* \* \* \* \*