



US012404280B2

(12) **United States Patent**
Park et al.

(10) **Patent No.:** **US 12,404,280 B2**

(45) **Date of Patent:** **Sep. 2, 2025**

(54) **HETEROCYCLIC COMPOUND, AND
ORGANIC LIGHT-EMITTING ELEMENT
USING SAME**

(71) Applicant: **LT MATERIALS CO., LTD.**, Yongin
(KR)

(72) Inventors: **Seong-Jong Park**, Yongin-si (KR);
Hyun-Ju La, Yongin-si (KR);
Won-Jang Jeong, Yongin-si (KR);
Dong-Jun Kim, Yongin-si (KR)

(73) Assignee: **LT MATERIALS CO., LTD.**,
Yongin-si (KR)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 802 days.

(21) Appl. No.: **17/637,342**

(22) PCT Filed: **Aug. 19, 2020**

(86) PCT No.: **PCT/KR2020/011019**

§ 371 (c)(1),

(2) Date: **Feb. 22, 2022**

(87) PCT Pub. No.: **WO2021/040310**

PCT Pub. Date: **Mar. 4, 2021**

(65) **Prior Publication Data**

US 2022/0324877 A1 Oct. 13, 2022

(30) **Foreign Application Priority Data**

Aug. 23, 2019 (KR) 10-2019-0103655

(51) **Int. Cl.**

H01L 51/54 (2006.01)

C07D 498/04 (2006.01)

H10K 85/60 (2023.01)

H10K 50/18 (2023.01)

(52) **U.S. Cl.**

CPC **C07D 498/04** (2013.01); **H10K 85/621**
(2023.02); **H10K 85/654** (2023.02); **H10K**
85/6572 (2023.02); **H10K 85/6574** (2023.02);
H10K 85/6576 (2023.02); **H10K 50/18**
(2023.02)

(58) **Field of Classification Search**

None

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

6,551,723	B1	4/2003	Okada et al.
2015/0243895	A1	8/2015	Lim et al.
2016/0380208	A1	12/2016	La et al.
2018/0282295	A1	10/2018	Parham et al.
2019/0288218	A1	9/2019	La et al.
2020/0066996	A1	2/2020	La et al.
2020/0339513	A1	10/2020	Heo et al.

FOREIGN PATENT DOCUMENTS

CN	105358533	A1	2/2016
CN	109384786	A	2/2019
JP	2000-63818	A	2/2000
JP	6479770	B2	3/2019
KR	10-2017-0084190	A	7/2017
KR	10-2018-0062343	A	6/2018
KR	10-2018-0075398	A	7/2018
KR	10-2019-0078139	A	7/2019

OTHER PUBLICATIONS

International Search Report (PCT/ISA/210) issued in PCT/KR2020/011019 mailed on Dec. 1, 2020.

Primary Examiner — Andrew K Bohaty

(74) *Attorney, Agent, or Firm* — Birch, Stewart, Kolasch
& Birch, LLP


(57)

ABSTRACT


The present specification relates to a heterocyclic compound represented by Chemical Formula 1, and an organic light emitting device using the same.

15 Claims, 3 Drawing Sheets


【FIG. 1】




【FIG. 2】



【FIG. 3】



【FIG. 4】



【FIG. 5】

CATHODE
ELECTRON INJECTION LAYER
SECOND ELECTRON TRANSFER LAYER
SECOND HOLE BLOCKING LAYER
SECOND STACK LIGHT EMITTING LAYER
SECOND ELECTRON BLOCKING LAYER
SECOND HOLE TRANSFER LAYER
P-TYPE CHARGE GENERATION LAYER
N-TYPE CHARGE GENERATION LAYER
FIRST ELECTRON TRANSFER LAYER
FIRST HOLE BLOCKING LAYER
FIRST STACK LIGHT EMITTING LAYER
FIRST ELECTRON BLOCKING LAYER
FIRST HOLE TRANSFER LAYER
FIRST HOLE INJECTION LAYER
ANODE
SUBSTRATE

1

HETEROCYCLIC COMPOUND, AND ORGANIC LIGHT-EMITTING ELEMENT USING SAME

TECHNICAL FIELD

The present specification relates to a heterocyclic compound, and an organic light emitting device using the same.

This application claims priority to and the benefits of Korean Patent Application No. 10-2019-0103655, filed with the Korean Intellectual Property Office on Aug. 23, 2019, the entire contents of which are incorporated herein by reference.

BACKGROUND ART

An electroluminescent device is one type of self-emissive display devices, and has an advantage of having a wide viewing angle, and a high response speed as well as having an excellent contrast.

An organic light emitting device has a structure disposing an organic thin film between two electrodes. When a voltage is applied to an organic light emitting device having such a structure, electrons and holes injected from the two electrodes bind and pair in the organic thin film, and light emits as these annihilate. The organic thin film may be formed in a single layer or a multilayer as necessary.

A material of the organic thin film may have a light emitting function as necessary. For example, as a material of the organic thin film, compounds capable of forming a light emitting layer themselves alone may be used, or compounds capable of performing a role of a host or a dopant of a host-dopant-based light emitting layer may also be used. In addition thereto, compounds capable of performing roles of hole injection, hole transfer, electron blocking, hole blocking, electron transfer, electron injection and the like may also be used as a material of the organic thin film.

Development of an organic thin film material has been continuously required for enhancing performance, lifetime or efficiency of an organic light emitting device.

DISCLOSURE


Technical Problem

The present specification is directed to providing a heterocyclic compound, and an organic light emitting device using the same.

Technical Solution

One embodiment of the present specification provides a heterocyclic compound represented by the following Chemical Formula 1.

[Chemical Formula 1]



2

In Chemical Formula 1,

X is O; S; or NR,

L1 to L3 are the same as or different from each other, and each independently a direct bond; a substituted or unsubstituted arylene group having 6 to 60 carbon atoms; or a substituted or unsubstituted heteroarylene group having 2 to 60 carbon atoms,

Z1 to Z3 are the same as or different from each other, and each independently selected from the group consisting of hydrogen; deuterium; a halogen group; a cyano group; $\text{—P(=O)(R104)(R105)}$; —N(R106)(R107) ; a substituted or unsubstituted alkyl group having 1 to 60 carbon atoms; a substituted or unsubstituted alkenyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkynyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkoxy group having 1 to 60 carbon atoms; a substituted or unsubstituted cycloalkyl group having 3 to 60 carbon atoms; a substituted or unsubstituted heterocycloalkyl group having 2 to 60 carbon atoms; a substituted or unsubstituted aryl group having 6 to 60 carbon atoms; and a substituted or unsubstituted heteroaryl group having 2 to 60 carbon atoms,

R1 and R2 are the same as or different from each other, and each independently selected from the group consisting of hydrogen; deuterium; a halogen group; a cyano group; $\text{—P(=O)(R104')(R105')}$; —N(R106')(R107') ; a substituted or unsubstituted alkyl group having 1 to 60 carbon atoms; a substituted or unsubstituted alkenyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkynyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkoxy group having 1 to 60 carbon atoms; a substituted or unsubstituted cycloalkyl group having 3 to 60 carbon atoms; a substituted or unsubstituted heterocycloalkyl group having 2 to 60 carbon atoms; a substituted or unsubstituted aryl group having 6 to 60 carbon atoms; and a substituted or unsubstituted heteroaryl group having 2 to 60 carbon atoms, or adjacent two or more groups bond to each other to form a substituted or unsubstituted aliphatic hydrocarbon ring; a substituted or unsubstituted aromatic hydrocarbon ring; a substituted or unsubstituted aliphatic heteroring; or a substituted or unsubstituted aromatic heteroring,

R is hydrogen; deuterium; a halogen group; a cyano group; a substituted or unsubstituted alkyl group having 1 to 60 carbon atoms; a substituted or unsubstituted alkenyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkynyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkoxy group having 1 to 60 carbon atoms; a substituted or unsubstituted cycloalkyl group having 3 to 60 carbon atoms; a substituted or unsubstituted heterocycloalkyl group having 2 to 60 carbon atoms; a substituted or unsubstituted aryl group having 6 to 60 carbon atoms; or a substituted or unsubstituted heteroaryl group having 2 to 60 carbon atoms,

R104 to R107 and R104' to R107' are each independently hydrogen; deuterium; a halogen group; an alkyl group; an alkenyl group; an alkoxy group; a cycloalkyl group; an aryl group; or a heteroaryl group,

m, p, x, n, q and y are each an integer of 1 to 5,

a is an integer of 1 to 3, and

when m, p, x, n, q, y and a are each an integer of 2 or greater, substituents in the parentheses are the same as or different from each other.

Another embodiment of the present application provides an organic light emitting device including a first electrode; a second electrode provided opposite to the first electrode; and one or more organic material layers provided between the first electrode and the second electrode, wherein one or more layers of the organic material layers include the heterocyclic compound represented by Chemical Formula 1.

Advantageous Effects

A compound described in the present specification can be used as a material of an organic material layer of an organic light emitting device. In the organic light emitting device, the compound is capable of performing a role of a hole injection material, a hole transfer material, a light emitting material, an electron transfer material, an electron injection material or the like. Particularly, the compound can be used as an electron transfer layer material or a charge generation material of the organic light emitting device.

Specifically, when using the compound represented by Chemical Formula 1 in an organic material layer, a driving voltage of the device can be lowered, light efficiency can be enhanced, and lifetime properties of the device can be enhanced.

DESCRIPTION OF DRAWINGS

FIG. 1 to FIG. 5 are diagrams each illustrating a lamination structure of an organic light emitting device according to One embodiment of the present specification.

- 100: Substrate
- 200: Anode
- 300: Organic Material Layer
- 301: Hole Injection Layer
- 302: Hole Transfer Layer
- 303: Light Emitting Layer
- 304: Hole Blocking Layer
- 305: Electron Transfer Layer
- 306: Electron Injection Layer
- 400: Cathode

Mode for Disclosure

Hereinafter, the present specification will be described in more detail.

In the present specification, a certain part "including" certain constituents means capable of further including other constituents, and does not exclude other constituents unless particularly stated on the contrary.

In the present specification, "substituted or unsubstituted" means being substituted with one or more substituents selected from the group consisting of deuterium; a halogen group; a cyano group; a linear or branched alkyl group having 1 to 60 carbon atoms; a linear or branched alkenyl group having 2 to 60 carbon atoms; a linear or branched alkynyl group having 2 to 60 carbon atoms; a monocyclic or polycyclic cycloalkyl group having 3 to 60 carbon atoms; a monocyclic or polycyclic heterocycloalkyl group having 2 to 60 carbon atoms; a monocyclic or polycyclic aryl group having 6 to 60 carbon atoms; a monocyclic or polycyclic heteroaryl group having 2 to 60 carbon atoms; a silyl group; a phosphine oxide group; an alkylamine group having 1 to 20 carbon atoms; a monocyclic or polycyclic arylamine group having 6 to 60 carbon atoms; and a monocyclic or polycyclic heteroarylamine group having 2 to 60 carbon atoms, or being unsubstituted, or being substituted with a

substituent linking two or more substituents selected from among the substituents illustrated above, or being unsubstituted.

The term "substitution" means a hydrogen atom bonding to a carbon atom of a compound being changed to another substituent, and the position of substitution is not limited as long as it is a position at which the hydrogen atom is substituted, that is, a position at which a substituent can substitute, and when two or more substituents substitute, the two or more substituents may be the same as or different from each other.

In the present specification, a "case of a substituent being not indicated in a chemical formula or compound structure" means that a hydrogen atom bonds to a carbon atom. However, since deuterium (^2H) is an isotope of hydrogen, some hydrogen atoms may be deuterium.

In one embodiment of the present application, a "case of a substituent being not indicated in a chemical formula or compound structure" may mean that positions that may come as a substituent may all be hydrogen or deuterium. In other words, since deuterium is an isotope of hydrogen, some hydrogen atoms may be deuterium that is an isotope, and herein, a content of the deuterium may be from 0% to 100%.


In one embodiment of the present application, in a "case of a substituent being not indicated in a chemical formula or compound structure", hydrogen and deuterium may be mixed in compounds when deuterium is not explicitly excluded such as a deuterium content being 0%, a hydrogen content being 100% or substituents being all hydrogen.

In one embodiment of the present application, deuterium is one of isotopes of hydrogen, is an element having deuterium formed with one proton and one neutron as a nucleus, and may be expressed as hydrogen-2, and the elemental symbol may also be written as D or ^2H .

In one embodiment of the present application, an isotope means an atom with the same atomic number (Z) but with a different mass number (A), and may also be interpreted as an element with the same number of protons but with a different number of neutrons.

In one embodiment of the present application, a meaning of a content T % of a specific substituent may be defined as $T2/T1 \times 100 = T\%$ when the total number of substituents that a basic compound may have is defined as T1, and the number of specific substituents among these is defined as T2.

In other words, in one example, having a deuterium content of 20% in a phenyl group represented by



means that the total number of substituents that the phenyl group may have is 5 (T1 in the formula), and the number of deuterium among these is 1 (T2 in the formula). In other words, having a deuterium content of 20% in a phenyl group may be represented by the following structural formulae.


In the present specification, the silyl group is a substituent including Si, having the Si atom directly linked as a radical,

7

and is represented by $-\text{Si}(\text{R101})(\text{R102})(\text{R103})$. R101 to R103 are the same as or different from each other, and may be each independently a substituent formed with at least one of hydrogen; deuterium; a halogen group; an alkyl group; an alkenyl group; an alkoxy group; a cycloalkyl group; an aryl group; and a heteroaryl group. Specific examples of the silyl group may include a trimethylsilyl group, a triethylsilyl group, a t-butyltrimethylsilyl group, a vinyltrimethylsilyl group, a propyltrimethylsilyl group, a triphenylsilyl group, a diphenylsilyl group, a phenylsilyl group and the like, but are not limited thereto.

In the present specification, the fluorenyl group may be substituted, and adjacent substituents may bond to each other to form a ring.

When the fluorenyl group is substituted,



and the like may be included, however, the structure is not limited thereto.

In the present specification, the heteroaryl group includes O, S, Se, N or Si as a heteroatom, includes monocyclic or polycyclic having 2 to 60 carbon atoms, and may be further substituted with other substituents. Herein, the polycyclic means a group in which the heteroaryl group is directly linked to or fused with other cyclic groups. Herein, the other cyclic groups may be a heteroaryl group, but may also be different types of cyclic groups such as a cycloalkyl group,

8

a heterocycloalkyl group and an aryl group. The number of carbon atoms of the heteroaryl group may be from 2 to 60, specifically from 2 to 40 and more specifically from 3 to 25. Specific examples of the heteroaryl group may include a pyridyl group, a pyrrolyl group, a pyrimidyl group, a pyridazinyl group, a furanyl group, a thiophene group, an imidazolyl group, a pyrazolyl group, an oxazolyl group, an isoxazolyl group, a thiazolyl group, an isothiazolyl group, a triazolyl group, a furazanyl group, an oxadiazolyl group, a thiadiazolyl group, a dithiazolyl group, a tetrazolyl group, a pyranlyl group, a thiopyranlyl group, a diazinyl group, an oxazinyl group, a thiazinyl group, a dioxynyl group, a triazinyl group, a tetrazinyl group, a quinolyl group, an isoquinolyl group, a quinazolinyl group, an isoquinazolinyl group, a quinoxalinyl group, a naphthyridyl group, an acridinyl group, a phenanthridinyl group, an imidazopyridinyl group, a diazanaphthalenyl group, a triazaindene group, an indolyl group, an indolizinyl group, a benzothiazolyl group, a benzoxazolyl group, a benzimidazolyl group, a benzothiophene group, a benzofuran group, a dibenzothiophene group, a dibenzofuran group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a phenazinyl group, a dibenzosilole group, spirobi(dibenzosilole), a dihydrophenazinyl group, a phenoxazinyl group, a phenanthridyl group, an imidazopyridinyl group, a thienyl group, an indolo[2,3-a]carbazolyl group, an indolo[2,3-b]carbazolyl group, an indolinyl group, a 10,11-dihydro-dibenzo[b,f]azepine group, a 9,10-dihydroacridinyl group, a phenanthrazinyl group, a phenothiazinyl group, a phthalazinyl group, a naphthylidinyll group, a phenanthrolinyl group, a benzo[c][1,2,5]thiadiazolyl group, a 5,10-dihydrobenzo[b,e][1,4]azasilinyl group, a pyrazolo[1,5-c]quinazolinyl group, a pyrido[1,2-b]indazolyl group, a pyrido[1,2-a]imidazo[1,2-e]indolinyl group, a 5,11-dihydroindeno[1,2-b]carbazolyl group and the like, but are not limited thereto.

In the present specification, examples of the aryl group and the heteroaryl group described above may be respectively applied to the arylene group and the heteroarylene group except that they are a divalent group.

In the present specification, the phosphine oxide group is represented by $-\text{P}(=\text{O})(\text{R104})(\text{R105})$, and R104 and R105 are the same as or different from each other and may be each independently a substituent formed with at least one of hydrogen; deuterium; a halogen group; an alkyl group; an alkenyl group; an alkoxy group; a cycloalkyl group; an aryl group; and a heteroaryl group. Specifically, the phosphine oxide group may be substituted with an aryl group, and as the aryl group, the examples described above may be applied. Examples of the phosphine oxide group may include a diphenylphosphine oxide group, a dinaphthylphosphine oxide group and the like, but are not limited thereto.

In the present specification, the amine group is represented by $-\text{N}(\text{R106})(\text{R107})$, and R106 and R107 are the same as or different from each other and may be each independently a substituent formed with at least one of hydrogen; deuterium; a halogen group; an alkyl group; an alkenyl group; an alkoxy group; a cycloalkyl group; an aryl group; and a heteroaryl group. The amine group may be selected from the group consisting of $-\text{NH}_2$; a monoalkylamine group; a monoarylamine group; a monoheteroarylamine group; a dialkylamine group; a diarylamine group; a diheteroarylamine group; an alkylarylamine group; an alkylheteroarylamine group; and an arylheteroarylamine group, and although not particularly limited thereto, the number of carbon atoms is preferably from 1 to 30. Specific examples of the amine group may include a methylamine group, a dimethylamine group, an ethylamine group, a

diethylamine group, a phenylamine group, a naphthylamine group, a biphenylamine group, a dibiphenylamine group, an anthracenylamine group, a 9-methyl-anthracenylamine group, a diphenylamine group, a phenylnaphthylamine group, a ditolylamine group, a phenyltolylamine group, a triphenylamine group, a biphenylnaphthylamine group, a phenylbiphenylamine group, a biphenylfluorenylamine group, a phenyltriphenylenylamine group, a biphenyltriphenylenylamine group and the like, but are not limited thereto.

In the present specification, an “adjacent” group may mean a substituent substituting an atom directly linked to an atom substituted by the corresponding substituent, a substituent sterically most closely positioned to the corresponding substituent, or another substituent substituting an atom substituted by the corresponding substituent. For example, two substituents substituting ortho positions in a benzene ring, and two substituents substituting the same carbon in an aliphatic ring may be interpreted as groups “adjacent” to each other.

As the aliphatic hydrocarbon ring, the aromatic hydrocarbon ring, the aliphatic heteroring or the aromatic heteroring that the adjacent groups may form, the structures illustrated as the cycloalkyl group, the aryl group, the cycloheteroalkyl group and the heteroaryl group described above may be applied except for those that are not a monovalent group.

One embodiment of the present specification provides a compound represented by Chemical Formula 1.

Chemical Formula 1 has a structure in which thiadiazole, oxadiazole or triazole is fused to phenanthridine, and has a wide band gap and a high T1 value by disconnecting electron distribution of HOMO (Highest Occupied Molecular Orbital) and LUMO (Lowest Unoccupied Molecular Orbital) by the thiadiazole, the oxadiazole or the triazole.

When having a low T1, excitons generated in a light emitting layer pass over to an electron transfer layer and resultantly emit light in the electron transfer layer or at an interface of the electron transfer layer by causing charge unbalance in the light emitting layer, and as a result, an organic electric device has a decrease in color purity, and decreases in efficiency and lifetime. By having a wide band gap, holes passing from the light emitting layer may be prevented due to a low HOMO level, and a high LUMO level may readily take over electrons in an electron injection layer and readily transfer the electrons to the light emitting layer.

In one embodiment of the present specification, X may be O; S; or NR.

In another embodiment, X may be O.

In another embodiment, X may be S.

In another embodiment, X may be NR.

In one embodiment of the present specification, R may be hydrogen; deuterium; a halogen group; a cyano group; a substituted or unsubstituted alkyl group having 1 to 60 carbon atoms; a substituted or unsubstituted alkenyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkynyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkoxy group having 1 to 60 carbon atoms; a substituted or unsubstituted cycloalkyl group having 3 to 60 carbon atoms; a substituted or unsubstituted heterocycloalkyl group having 2 to 60 carbon atoms; a substituted or unsubstituted aryl group having 6 to 60 carbon atoms; or a substituted or unsubstituted heteroaryl group having 2 to 60 carbon atoms.

In another embodiment, R may be a substituted or unsubstituted aryl group having 6 to 60 carbon atoms.

In another embodiment, R may be a substituted or unsubstituted aryl group having 6 to 40 carbon atoms.

In another embodiment, R may be a substituted or unsubstituted aryl group having 6 to 20 carbon atoms.

In another embodiment, R may be a phenyl group.

In one embodiment of the present specification, L1 to L3 are the same as or different from each other, and may be each independently a direct bond; a substituted or unsubstituted arylene group having 6 to 60 carbon atoms; or a substituted or unsubstituted heteroarylene group having 2 to 60 carbon atoms.

In another embodiment, L1 to L3 are the same as or different from each other, and may be each independently a direct bond; a substituted or unsubstituted arylene group having 6 to 40 carbon atoms; or a substituted or unsubstituted heteroarylene group having 2 to 40 carbon atoms.

In another embodiment, L1 to L3 are the same as or different from each other, and may be each independently a direct bond; or a substituted or unsubstituted arylene group having 6 to 20 carbon atoms.

In another embodiment, L1 to L3 are the same as or different from each other, and may be each independently a direct bond; or an arylene group having 6 to 20 carbon atoms unsubstituted or substituted with a halogen group.

In another embodiment, L1 to L3 are the same as or different from each other, and may be each independently a direct bond; a phenylene group unsubstituted or substituted with a halogen group; a biphenylene group; or a terphenylene group.

In one embodiment of the present specification, Z1 to Z3 are the same as or different from each other, and may be each independently selected from the group consisting of hydrogen; deuterium; a halogen group; a cyano group; —P(=O) (R104)(R105); —N(R106)(R107) ; a substituted or unsubstituted alkyl group having 1 to 60 carbon atoms; a substituted or unsubstituted alkenyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkynyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkoxy group having 1 to 60 carbon atoms; a substituted or unsubstituted cycloalkyl group having 3 to 60 carbon atoms; a substituted or unsubstituted heterocycloalkyl group having 2 to 60 carbon atoms; a substituted or unsubstituted aryl group having 6 to 60 carbon atoms; and a substituted or unsubstituted heteroaryl group having 2 to 60 carbon atoms.

In another embodiment, Z1 to Z3 are the same as or different from each other, and may be each independently hydrogen; deuterium; a cyano group; —P(=O) (R104)(R105); a substituted or unsubstituted aryl group having 6 to 60 carbon atoms; or a substituted or unsubstituted heteroaryl group having 2 to 60 carbon atoms.

In another embodiment, Z1 to Z3 are the same as or different from each other, and may be each independently hydrogen; deuterium; a cyano group; —P(=O) (R104)(R105); a substituted or unsubstituted aryl group having 6 to 40 carbon atoms; or a substituted or unsubstituted heteroaryl group having 2 to 40 carbon atoms. R104 and R105 may be each independently an alkyl group or an aryl group.

In another embodiment, Z1 to Z3 are the same as or different from each other, and may be each independently hydrogen; deuterium; a cyano group; —P(=O) (R104)(R105); a substituted or unsubstituted phenyl group; a biphenyl group; a naphthyl group; a substituted or unsubstituted pyridine group; a substituted or unsubstituted pyrimidine group; a substituted or unsubstituted triazine group; a substituted or unsubstituted quinoline group; a substituted or unsubstituted benzimidazolyl group; a benzoxazolyl group; a benzothiazolyl group; a carbazolyl group;

11

a dibenzofuran group; a dibenzothiophene group; a phenanthrenyl group; a phenanthrolinyl group; a 9,9-dimethyl-9H-fluorenyl group; or a benzo[4,5]thieno[3,2-d]pyrimidine group. R104 and R105 may be each independently an alkyl group or an aryl group.

In another embodiment, Z1 to Z3 are the same as or different from each other, and may be each independently hydrogen; deuterium; a cyano group; $\text{—P(=O)(R104)(R105)}$; a phenyl group unsubstituted or substituted with one or more substituents selected from the group consisting of a halogen group, a cyano group, a phenyl group, a methyl group and a carbazole group; a biphenyl group; a naphthyl group; a pyridine group unsubstituted or substituted with a phenyl group or a pyridine group; a pyrimidine group unsubstituted or substituted with one or more substituents selected from the group consisting of a phenyl group, a biphenyl group, a naphthyl group, a phenanthrenyl group and toluene; a triazine group unsubstituted or substituted with one or more substituents selected from the group consisting of a phenyl group, a biphenyl group, a naphthyl group, a phenanthrenyl group and toluene; a quinoline group unsubstituted or substituted with a phenyl group; a benzimidazolyl group unsubstituted or substituted with one or more substituents selected from the group consisting of a phenyl group and an ethyl group; a benzoxazolyl group; a benzothiazolyl group; a carbazolyl group; a dibenzofuran group; a dibenzothiophene group; a phenanthrenyl group; a phenanthrolinyl group; a 9,9-dimethyl-9H-fluorenyl group; or a benzo[4,5]thieno[3,2-d]pyrimidine group, and R104 and R105 may be each independently a methyl group or a phenyl group.


In one embodiment of the present specification, R1 and R2 are the same as or different from each other, and may be each independently selected from the group consisting of hydrogen; deuterium; a halogen group; a cyano group; $\text{—P(=O)(R104')(R105')}$; —N(R106')(R107') ; a substituted or unsubstituted alkyl group having 1 to 60 carbon atoms; a substituted or unsubstituted alkenyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkynyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkoxy group having 1 to 60 carbon atoms; a substituted or unsubstituted cycloalkyl group having 3 to 60 carbon atoms; a substituted or unsubstituted heterocycloalkyl group having 2 to 60 carbon atoms; a substituted or unsubstituted aryl group having 6 to 60 carbon atoms; and a substituted or unsubstituted heteroaryl group having 2 to 60 carbon atoms, or adjacent two or more groups may bond to each other to form a substituted or unsubstituted aliphatic hydrocarbon ring; a substituted or unsubstituted aromatic hydrocarbon ring; a substituted or unsubstituted aliphatic heteroring; or a substituted or unsubstituted aromatic heteroring.

In another embodiment, R1 and R2 may each be hydrogen; or deuterium.


In the heterocyclic compound provided in one embodiment of the present specification, Chemical Formula 1 is represented by any one of the following Chemical Formulae 2 to 5.

12


[Chemical Formula 2]




[Chemical Formula 3]



[Chemical Formula 4]



[Chemical Formula 5]



In Chemical Formula 2 to Chemical Formula 5, X, m, p, x, n, q, y and a have the same definitions as in Chemical Formula 1,

L11 to L13 are the same as or different from each other, and each independently a direct bond; a substituted or unsubstituted arylene group having 6 to 60 carbon atoms; or a substituted or unsubstituted heteroarylene group having 2 to 60 carbon atoms,

Z11 to Z13 are the same as or different from each other, and each independently selected from the group consisting of a cyano group; $\text{—P(=O)(R104)(R105)}$; a substituted or unsubstituted aryl group having 6 to 60 carbon atoms; and a substituted or unsubstituted heteroaryl group having 2 to 60 carbon atoms,

R11 and R12 are the same as or different from each other, and each independently selected from the group con-

13

sisting of hydrogen; deuterium; a halogen group; a cyano group; $-\text{P}(=\text{O})(\text{R104})(\text{R105})$; $-\text{N}(\text{R106})(\text{R107})$; a substituted or unsubstituted alkyl group having 1 to 60 carbon atoms; a substituted or unsubstituted alkenyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkynyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkoxy group having 1 to 60 carbon atoms; a substituted or unsubstituted cycloalkyl group having 3 to 60 carbon atoms; and a substituted or unsubstituted heterocycloalkyl group having 2 to 60 carbon atoms.

R104, R105 and R104' to R107' are each independently hydrogen; deuterium; a halogen group; an alkyl group; an alkenyl group; an alkoxy group; a cycloalkyl group; an aryl group; or a heteroaryl group,

b is 1 or 2,

c is an integer of 1 to 4,

when b is 2, two R11s are the same as or different from each other, and

when c is 2 or greater, a plurality of R12s are the same as or different from each other.

In another embodiment, L11 to L13 are the same as or different from each other, and may be each independently a direct bond; a substituted or unsubstituted arylene group having 6 to 60 carbon atoms; or a substituted or unsubstituted heteroarylene group having 2 to 60 carbon atoms.

In another embodiment, L11 to L13 are the same as or different from each other, and may be each independently a direct bond; a substituted or unsubstituted arylene group having 6 to 40 carbon atoms; or a substituted or unsubstituted heteroarylene group having 2 to 40 carbon atoms.

In another embodiment, L11 to L13 are the same as or different from each other, and may be each independently a direct bond; or a substituted or unsubstituted arylene group having 6 to 20 carbon atoms.

In another embodiment, L11 to L13 are the same as or different from each other, and may be each independently a direct bond; or an arylene group having 6 to 20 carbon atoms unsubstituted or substituted with a halogen group.

In another embodiment, L11 to L13 are the same as or different from each other, and may be each independently a direct bond; a phenylene group unsubstituted or substituted with a halogen group; a biphenylene group; or a terphenylene group.

In another embodiment, Z11 to Z13 are the same as or different from each other, and may be each independently a cyano group; $-\text{P}(=\text{O})(\text{R104})(\text{R105})$; a substituted or unsubstituted aryl group having 6 to 60 carbon atoms; or a substituted or unsubstituted heteroaryl group having 2 to 60 carbon atoms.

In another embodiment, Z11 to Z13 are the same as or different from each other, and may be each independently a cyano group; $-\text{P}(=\text{O})(\text{R104})(\text{R105})$; a substituted or unsubstituted aryl group having 6 to 40 carbon atoms; or a substituted or unsubstituted heteroaryl group having 2 to 40 carbon atoms. R104 and R105 may be each independently an alkyl group or an aryl group.

In another embodiment, Z11 to Z13 are the same as or different from each other, and may be each independently a cyano group; $-\text{P}(=\text{O})(\text{R104})(\text{R105})$; a substituted or

14

unsubstituted phenyl group; a biphenyl group; a naphthyl group; a substituted or unsubstituted pyridine group; a substituted or unsubstituted pyrimidine group; a substituted or unsubstituted triazine group; a substituted or unsubstituted quinoline group; a substituted or unsubstituted benzimidazolyl group; a benzoxazolyl group; a benzothiazolyl group; a carbazolyl group; a dibenzofuran group; a dibenzothiophene group; a phenanthrenyl group; a phenanthrolinyl group; a 9,9-dimethyl-9H-fluorenyl group; or a benzo [4,5]thieno[3,2-d]pyrimidine group. R104 and R105 may be each independently an alkyl group or an aryl group.

In another embodiment, Z11 to Z13 are the same as or different from each other, and may be each independently a cyano group; $-\text{P}(=\text{O})(\text{R104})(\text{R105})$; a phenyl group unsubstituted or substituted with one or more substituents selected from the group consisting of a halogen group, a cyano group, a phenyl group, a methyl group and a carbazole group; a biphenyl group; a naphthyl group; a pyridine group unsubstituted or substituted with a phenyl group or a pyridine group; a pyrimidine group unsubstituted or substituted with one or more substituents selected from the group consisting of a phenyl group, a biphenyl group, a naphthyl group, a phenanthrenyl group and toluene; a triazine group unsubstituted or substituted with one or more substituents selected from the group consisting of a phenyl group, a biphenyl group, a naphthyl group, a phenanthrenyl group and toluene; a quinoline group unsubstituted or substituted with a phenyl group; a benzimidazolyl group unsubstituted or substituted with one or more substituents selected from the group consisting of a phenyl group and an ethyl group; a benzoxazolyl group; a benzothiazolyl group; a carbazolyl group; a dibenzofuran group; a dibenzothiophene group; a phenanthrenyl group; a phenanthrolinyl group; a 9,9-dimethyl-9H-fluorenyl group; or a benzo[4,5]thieno[3,2-d]pyrimidine group, and R104 and R105 may be each independently a methyl group or a phenyl group.

In one embodiment of the present specification, R11 and R12 are the same as or different from each other, and may be each independently selected from the group consisting of hydrogen; deuterium; a halogen group; a cyano group; $-\text{P}(=\text{O})(\text{R104})(\text{R105})$; $-\text{N}(\text{R106})(\text{R107})$; a substituted or unsubstituted alkyl group having 1 to 60 carbon atoms; a substituted or unsubstituted alkenyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkynyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkoxy group having 1 to 60 carbon atoms; a substituted or unsubstituted cycloalkyl group having 3 to 60 carbon atoms; and a substituted or unsubstituted heterocycloalkyl group having 2 to 60 carbon atoms.




In one embodiment of the present specification, R11 and R12 are the same as or different from each other, and may be each independently hydrogen; deuterium; or a substituted or unsubstituted alkyl group having 1 to 60 carbon atoms.

In one embodiment of the present specification, R11 and R12 are the same as or different from each other, and may be each independently hydrogen; or deuterium.

In one embodiment of the present specification, R11 and R12 may be hydrogen.





In the heterocyclic compound provided in one embodiment of the present specification, Chemical Formula 1 is represented by any one of the following compounds.

15






16

-continued



17

-continued

**18**

-continued

8


11

5

10

15

20




12

9 25

30

35

40



45

13


10

50

55




60

65



19

-continued

**20**

-continued

14

5

10

15

20

25

15

30

35

40

45

50

16

55




60

65

17




18

19



21

-continued

**22**

-continued

20

5

10

15

20

21

25

30

35

40

22 45


50

55


60

65


23



24




25



23

-continued



26

5


10

15


20

25


27



28

**24**

-continued



29

25

27

30

35

40

45


28

50

55

60


65



31

25

-continued

**26**

-continued

32

5

10

15

20

25

30

35

33 40

45


50

55



60

65

34





35



27


-continued

**28**

-continued


36

38




39

20



37


40



65

29

-continued



41

5

10

15


20

25

30

35


42 40



65

30

-continued



43

5

10

15

20

25

30

35

42 40


45

50

55

60


65



44

31

-continued



45

5

10

15

20

25

46

30

35

40

47

45

50


55

60

65

32

-continued





48

49

33

-continued

**34**

-continued

50

52


5

10

15

20

25




53

30

35

40

51



54


45

50

55



60

65



35

-continued

**36**

-continued

55

57

5

10

15

20

25

30

35

40

56

45



50

55

60




65

58






37

-continued




**38**

-continued



39


-continued

**40**

-continued

65


68



20


66

69




67

70



41

-continued



71

5


10

15

20

42

-continued



74

72 25

30

35

40


73 45

50


55

60



65



75






76






43

-continued


**44**

-continued



45

-continued



83

5

10

15

20

84

25


30

35

40

85


45



65


46

-continued




86

87






88







47

-continued





**48**

-continued







49

-continued


**50**

-continued









51

-continued


**52**

-continued



53

-continued

**54**

-continued

111

113

5

10

15

20

25

30

35

40

112

114



45

50

55


60

65



55

-continued

**56**

-continued

115

5

10

15

20

25

30

35

40

116

45

50



55

60

65


117

118



57

-continued

**58**

-continued

119

121

5

10

15

20

25

30

35

120 40

122



45

50

55



60

65



59

-continued

**60**

-continued

123

125

5

10

15

20

25

30

35

40

124


45

50


55

60


65



126





127



61

-continued

**62**

-continued

128

5

10

15

20

25

30

35

40

129


45

50


55

60

65





130



131

63

-continued

**64**

-continued

132

134

5

10

15

20

25

30

35

133

40

135



45

50

55



60

65



65

-continued

**66**

-continued

136

5

10

15

20

25

30

35

40

137


45

50


55

60

65




138



139

67

-continued

**68**

-continued

140

5

10

15

20

25

30

35

40

141

45


50

55



60

65

142




143





69

-continued

**70**

-continued





150

5

10

15

20

25

30

35

40

151

45


50

55

60


65

72
-continued



152



153



151

45


154



73

-continued

155




5

10

15

20

156




30

35

40

157




60

65


74

-continued


158



159






160





75

-continued


**76**

-continued



77


-continued




166

78


-continued




168



169




167




170

79

-continued

**80**

-continued



30

35

40

172


45

50


55

60


65



174



-continued



25

30

35

40

176

45

50


55

60


65

-continued

177




178



83

-continued



179

5

10

15

20

25

30

35

40

180

45

50


55

60

65



84




-continued




181

182







187

5

10

15

20

25

30

35

40

188

45


50

55

60



65

88
-continued




189

190



89

-continued

**90**

-continued

191

5

10

15

20

25

30

35

40

192

45


50

55


60


65

193



194





195

5

10

15

20


25

30

35

40


196



60

65

92
-continued



197

5

10

15

20

25

30

35

40

45


50






55

60

65




198





95

-continued

**96**

-continued

204

5

10

15

20

205

25

30

35

40

206

45


50

55


60

65


207



208




209



97

-continued

**98**

-continued

210

212

5

10

15

20

25

30

35

40

211

213



45

50

55




60

65



99

-continued

**100**

-continued

214

5

10

15

20

215

25

30

35

40

216 45


50

55


60

65


217



218






219



101

-continued

**102**

-continued

220

223

5

10

15

20

221

25

224

30

35

40

222

45




225

50

55




60

65






103

-continued




**104**

-continued






105

-continued




**106**

-continued



107

-continued

**108**

-continued

238

241

5

10

15

20

25

239

242

30

35

40

45

240




243

50

55




60

65





109

-continued


**110**

-continued



111

-continued



249

5

10

15

20

25

30

35

40

250

45

50


55

60

65

112

-continued




251

252

113

-continued

**114**

-continued

253

5

10

15

20

25

30

35

40

254

45


50

55



60

65

255




256



115

-continued

**116**

-continued

257

259

5

10

15

20

25

30

35

40

258

260



45

50

55


60

65



117

-continued

**118**

-continued

261

5

10

15

20

25

30

35

40

262

45

50



55

60

65



263

264



119

-continued

**120**

-continued

265

267

5

10

15

20

25

30

35

40

266

268



45

50

55


60

65



121

-continued

**122**

-continued

269

271

5

10

15

20

25

30

35

270 40

45



50

55

60


65

272



123


-continued

**124**

-continued

273

275




30

35


40

274

276




65



125

-continued

**126**

-continued

277

279

5

10

15

20

25

30

35

40

278

45



50

55

60


65

280



127

-continued

**128**

-continued

281

283

5

10

15

20

25

30

35

40

282

284



45

50

55

60


65



129

-continued

285



5

10

15

20


25

30

35

40

286




65

130

-continued

287



25

30

35

40

286

45


50

55

60


65

288



131

-continued



289

5

10

15

20


25

30

35

40


290



65

132

-continued



291

5

10

15

20


25

30

35

40

290



292

45

50


55

60

65

133

-continued



293

5

10

15

20

25

30

35

40

294

45

50


55

60

65

134

-continued




295

296

135

-continued

**136**

-continued

297

299

5

10

15

20

25

30

35

40

298

45



50

55

60


65

300



137

-continued



301

5

10

15

20

25

30

35

40

302


45

50


55

60


65

**138**

-continued





303





304

139

-continued


**140**

-continued



141

-continued

**142**

-continued


310

5

10

15

20




313

25
311

30

35

40



314


45
312

50

55

60




65



315

143

-continued

**144**

-continued

316

5

10

15

20

317

25

30

35

40

318


45

50


55

60


65



319






320



321

145

-continued

**146**

-continued


322

5

10

15

20



325

25


323

30

35

40

45



326


324

50

55

60




65






327

147

-continued




**148**

-continued



149

-continued

**150**

-continued

334

5

10

15

20

335 25

30

35

40

45



336

50

55

60

65




337

338

151

-continued

**152**

-continued

339

5

10

15

20

25

30

35

40

340 45


50

55



60

65

341





342



153

-continued

**154**

-continued

343

345

5

10

15

20

25

30

35

40

344

346



45

50

55


60

65



155

-continued

**156**

-continued

347

5

10

15

20

25

30

35

40

348

45

50



55

60

65



349

350



157

-continued

**158**

-continued

351

5

10

15

20

25

30

35

40

352

45


50

55


60

65

353





354



159

-continued

**160**

-continued

355

5

10

15

20

25

30

35

40

356



45

50

55

60

65





357

358

161

-continued

**162**

-continued

359

5

10

15

20

25

30

35

40

360 45


50

55


60

65

361





362



163

-continued

**164**

-continued

363

5

10

15

20

25

30

35

40

364



45

50

55

60

65





365

366

165

-continued

**166**

-continued

367

5

10

15

20

25

30

35

40

368

45

50



55

60

65


369

370



167

-continued

**168**

-continued

371

5

10

15

20

25

30

35

40

372

45

50



55

60

65



373

374



169

-continued

**170**

-continued

375

5

10

15

20

25

30

35

40

376

45

50



55

60

65




377

378






171

-continued

**172**

-continued



379

5

10

15

20

380 25

30

35

40

381 45

50

55



60

65

382

383

384

173**174**

-continued

385

5

10

15

20

25

386


45

50

55

60


65



387




388

389



175


-continued

**176**

-continued


390

393




391

394




392

395




177

-continued



**178**

-continued



179

-continued

**180**

-continued

403

405

5

10

15

20

25

30

35

40

45

404



50

55

60



65

406



181

-continued

**182**

-continued

407

5

10

15

20

25

30

35

40

408

45

50



55

60

65



409

410



183

-continued

**184**

-continued

411

5

10

15

20

25

30

35

40

412



45

50

55

60

65





413

414

185

-continued

**186**

-continued

415

5

10

15

20

25

30

35

40



416 45

50

55

60

65




417

418

187

-continued



419

5

10

15

20

25

30

35

40


419 45

50


55

60

65


**188**

-continued




421

422



189

-continued



423

5

10

15

20

25

30

35

40

45


424

50


55

60

65


**190**

-continued





425

426



191

-continued

**192**

-continued

427

5

10

15

20

25

30

35

40

428

45

50



55

60

65



429

430



193

-continued

**194**

-continued

431

433

5

10

15

20

25

30

35

40

432

45

434




50

55

60




65

435



195

-continued

**196**

-continued

436

439

5

10

15

437

20

25

30

35

40

438

440



45

50

55




60

65



197

-continued

**198**

-continued

441

5

10

15

20

442

25

30

35

40

443


45

50

55

60



65



444



445

446



199

-continued

**200**

-continued

447

5

10

15

20

25

30

35

40

448




45

50

55

60

65






449

450

451

201

-continued



452

5

10

15

453

20

25

30

35

40

454

45

50



55

60

65

202

-continued






455

456

203

-continued

**204**

-continued

457

460

5

10

15

20

458

25

30

35

40

459


45

50


55

60


65



461





462



205

-continued

**206**

-continued

463

5

10

15

20

25

30

35

40

464

45



50

55

60

65



465



466

207

-continued

**208**

-continued

467

469

5

10

15

20

25

30

35

40

468

45

50



470

55

60


65

471










209

-continued





**210**

-continued







211

-continued





**212**

-continued






213

-continued




**214**

-continued






215

-continued



**216**

-continued



217

-continued

**218**

-continued

501

5

10

15

20

25

30

35

40

502

45



50

55

60

65




503



504

219

-continued

**220**

-continued

505

5

10

15

20

506

25

30

35

40

507




45

50

55

60

65





508

509

510

221

-continued

**222**

-continued

511

5

10

15

20

25

30

35

40

512




45

50

55

60

65







513

514



515

223

-continued

**224**

-continued



517

30

35

518

45

50

519

55

60

65

520





521

522




523

225

-continued


**226**

-continued



227


-continued




531

15

532 20




533



40

45



534


55

60

65

228

-continued



535

10

15

25

30

35

40

45

50

55

60

65

70

75

80

85

90

95

100

105

110

115

120

125

130

135

140

145

150

155

160

165

170

175

180

185

190

195

200

205

210

215

220

225

230

235

240

245

250

255

260

265

270

275

280

285

290

295

300

305

310

315

320

325

330

335

340

345

350

355

360

365

370

375

380

385

390

395

400

405

410

415

420

425

430

435

440

445

450

455

460

465

470

475

480

485

490

495

500

505

510

515

520

525

530

535

540

545

550

555

560

565

570

575

580

585

590

595

600

605

610

615

620

625

630

635

640

645

650

655

660

665

670

675

680

685

690

695

700

705

710

715

720

725

730

735

740

745

750

755

760

765

770

775

780

785

790

795

800

805

810

815

820

825

830

835

840

845

850

855

860

865

870

875

880

885

890

895

900

905

910

915

920

925

930

935

940

945

950

955

960

965

970

975

980

985

990

995

1000

1005

1010

1015

1020

1025

1030

1035

1040

1045

1050

1055

1060

1065

1070

1075

1080

1085

1090

1095

1100

1105

1110

1115

1120

1125

1130

1135

1140

1145

1150

1155

1160

1165

1170

1175

1180

1185

1190

1195

1200

1205

1210

1215

1220

1225

1230

1235

1240

1245

1250

1255

1260

1265

1270

1275

1280

1285

1290

1295

1300

1305

1310

1315

1320

1325

1330

1335

1340

1345

1350

1355

1360

1365

1370

1375

1380

1385

1390

1395

1400

1405

1410

1415

1420

1425

1430

1435

1440

1445

1450

1455

1460

1465

1470

1475

1480

1485

1490

1495

1500

1505

1510

1515

1520

1525

1530

1535

1540

1545

1550

1555

1560

1565

1570

1575

1580

1585

1590

1595

1600

1605

1610

1615

1620

1625




1630

1635




1640

229

-continued





**230**

-continued







231

-continued


**232**

-continued



233

-continued




552

5

10

15

20




553

25

30

35

40

45
554

50


55

60


65

234


-continued



555




556



557

235

-continued



558

5

10

15

20

25


559

30

35

40

45




560

50


55

60

65


**236**

-continued




561

562








563








237

-continued

**238**






-continued



239






-continued

574

**240**


-continued

579




241

-continued





**242**

-continued







243

-continued




**244**

-continued






245

-continued


**246**

-continued



247

-continued




607

5

10

15

20



608


25

30

35

40

45



609

50


55

60


65

248


-continued



610




611



612

249

-continued



613

5


10

15

20


250

-continued



616

614 25




30

35

40

45



617



615

50

55

60





65







618

251

-continued





**252**

-continued






253

-continued




**254**

-continued







255

-continued


**256**

-continued



257

-continued




641

5

10

15

20




642

30

35

40

45



643

50


55

60

65


258

-continued




644


645



646







647






259

-continued




**260**

-continued




261

-continued

**262**


-continued

655




658

656 25



659


657



660

263

-continued



In addition, by introducing various substituents to the structure of Chemical Formula 1, compounds having unique properties of the introduced substituents may be synthesized. For example, by introducing substituents normally used as hole injection layer materials, hole transfer layer materials, light emitting layer materials, electron transfer

264

layer materials and charge generation layer materials used for manufacturing an organic light emitting device to the core structure, materials satisfying conditions required for each organic material layer may be synthesized.

5 In addition, by introducing various substituents to the structure of Chemical Formula 1, the energy band gap may be finely controlled, and meanwhile, properties at interfaces between organic materials are enhanced, and material applications may become diverse.

10 One embodiment of the present specification provides an organic light emitting device including a first electrode; a second electrode; and one or more organic material layers provided between the first electrode and the second electrode, wherein one or more layers of the organic material layers include the heterocyclic compound represented by Chemical Formula 1.

In one embodiment of the present specification, the first electrode may be an anode, and the second electrode may be a cathode.

20 In another embodiment of the present specification, the first electrode may be a cathode, and the second electrode may be an anode.

In one embodiment of the present specification, the organic light emitting device may be a blue organic light emitting device, and the heterocyclic compound according to Chemical Formula 1 may be used as a material of the blue organic light emitting device.

In another embodiment of the present specification, the organic light emitting device may be a green organic light emitting device, and the heterocyclic compound according to Chemical Formula 1 may be used as a material of the green organic light emitting device.

In another embodiment of the present specification, the organic light emitting device may be a red organic light emitting device, and the heterocyclic compound according to Chemical Formula 1 may be used as a material of the red organic light emitting device.

Specific descriptions on the heterocyclic compound represented by Chemical Formula 1 are the same as the descriptions provided above.

The organic light emitting device of the present specification may be manufactured using common organic light emitting device manufacturing methods and materials except that one or more of the organic material layers are formed using the heterocyclic compound described above.

45 The heterocyclic compound may be formed into an organic material layer through a solution coating method as well as a vacuum deposition method when manufacturing the organic light emitting device. Herein, the solution coating method means spin coating, dip coating, inkjet printing, screen printing, a spray method, roll coating and the like, but is not limited thereto.

The organic material layer of the organic light emitting device of the present specification may be formed in a single layer structure, but may be formed in a multilayer structure in which two or more organic material layers are laminated. For example, the organic light emitting device of the present disclosure may have a structure including a hole injection layer, a hole transfer layer, a light emitting layer, an electron transfer layer, an electron injection layer and the like as the organic material layer. However, the structure of the organic light emitting device is not limited thereto, and may include a smaller number of organic material layers.

In the organic light emitting device of the present specification, the organic material layer includes an electron transfer layer, and the electron transfer layer may include the heterocyclic compound.

In the organic light emitting device of the present specification, the organic material layer includes a hole blocking layer, and the hole blocking layer may include the heterocyclic compound.

The organic light emitting device of the present disclosure may further include one, two or more layers selected from the group consisting of a light emitting layer, a hole injection layer, a hole transfer layer, an electron injection layer, an electron transfer layer, an electron blocking layer and a hole blocking layer.

FIG. 1 to FIG. 4 illustrate a lamination order of electrodes and organic material layers of an organic light emitting device according to one embodiment of the present specification. However, the scope of the present application is not limited to these diagrams, and structures of organic light emitting devices known in the art may also be used in the present application.

FIG. 1 illustrates an organic light emitting device in which an anode (200), an organic material layer (300) and a cathode (400) are consecutively laminated on a substrate (100). However, the structure is not limited to such a structure, and as illustrated in FIG. 2, an organic light emitting device in which a cathode, an organic material layer and an anode are consecutively laminated on a substrate may also be obtained.

FIG. 3 and FIG. 4 illustrate cases of the organic material layer being a multilayer. The organic light emitting device according to FIG. 3 includes a hole injection layer (301), a hole transfer layer (302), a light emitting layer (303), an electron transfer layer (305) and an electron injection layer (306), and the organic light emitting device according to FIG. 4 includes a hole injection layer (301), a hole transfer layer (302), a light emitting layer (303), a hole blocking layer (304), an electron transfer layer (305) and an electron injection layer (306). However, the scope of the present application is not limited to such a lamination structure, and as necessary, layers other than the light emitting layer may not be included, and other necessary functional layers may be further added.

The organic material layer including Chemical Formula 1 may further include other materials as necessary.

In addition, the organic light emitting device according to one embodiment of the present specification includes an anode, a cathode, and two or more stacks provided between the anode and the cathode, the two or more stacks each independently include a light emitting layer, a charge generation layer is included between the two or more stacks, and the charge generation layer includes the heterocyclic compound represented by Chemical Formula 1.

In addition, the organic light emitting device according to one embodiment of the present specification includes an anode, a first stack provided on the anode and including a first light emitting layer, a charge generation layer provided on the first stack, a second stack provided on the charge generation layer and including a second light emitting layer, and a cathode provided on the second stack. Herein, the charge generation layer may include the heterocyclic compound represented by Chemical Formula 1. In addition, the first stack and the second stack may each independently further include one or more types of the hole injection layer, the hole transfer layer, the hole blocking layer, the electron transfer layer, the electron injection layer and the like described above.

The organic light emitting device according to one embodiment of the present specification includes a first electrode; a first stack provided on the first electrode and including a first light emitting layer; a charge generation

layer provided on the first stack; a second stack provided on the charge generation layer and including a second light emitting layer; and a second electrode provided on the second stack, wherein the charge generation layer may include the heterocyclic compound represented by Chemical Formula 1.

The organic light emitting device according to one embodiment of the present specification includes a first electrode; a second electrode; and an organic material layer provided between the first electrode and the second electrode, wherein the organic material layer includes two or more stacks, and the two or more stacks each independently include a light emitting layer, a charge generation layer is included between the two or more stacks, and the charge generation layer may include the heterocyclic compound represented by Chemical Formula 1.

The organic light emitting device according to one embodiment of the present specification includes a first electrode; a second electrode; and an organic material layer provided between the first electrode and the second electrode, wherein the organic material layer includes a first stack including a first light emitting layer; a charge generation layer provided on the first stack; and a second stack provided on the charge generation layer and including a second light emitting layer, and the charge generation layer may include the heterocyclic compound represented by Chemical Formula 1.

The charge generation layer may be an N-type charge generation layer or a P-type charge generation layer, and the charge generation layer may further include a dopant known in the art in addition to the heterocyclic compound represented by Chemical Formula 1.

The charge generation layer may be an N-type charge generation layer, and the N-type charge generation layer may include the heterocyclic compound represented by Chemical Formula 1.

In the organic light emitting device according to one embodiment of the present specification, the second stack further includes a hole injection layer, and the hole injection layer includes the heterocyclic compound represented by Chemical Formula 1.

As the organic light emitting device according to one embodiment of the present specification, an organic light emitting device having a 2-stack tandem structure is illustrated in FIG. 5.

Herein, the first electron blocking layer, the first hole blocking layer, the second hole blocking layer and the like described in FIG. 5 may not be included in some cases.

In the organic light emitting device according to one embodiment of the present specification, materials other than the compound of Chemical Formula 1 are illustrated below, however, these are for illustrative purposes only and not for limiting the scope of the present application, and may be replaced by materials known in the art.

As the anode material, materials having relatively large work function may be used, and transparent conductive oxides, metals, conductive polymers or the like may be used. Specific examples of the anode material include metals such as vanadium, chromium, copper, zinc and gold, or alloys thereof; metal oxides such as zinc oxide, indium oxide, indium tin oxide (ITO) and indium zinc oxide (IZO); combinations of metals and oxides such as ZnO:Al or SnO₂:Sb; conductive polymers such as poly(3-methylthiophene), poly[3,4-(ethylene-1,2-dioxy)thiophene](PEDOT), polypyrrole and polyaniline, and the like, but are not limited thereto.

As the cathode material, materials having relatively small work function may be used, and metals, metal oxides,

267

conductive polymers or the like may be used. Specific examples of the cathode material include metals such as magnesium, calcium, sodium, potassium, titanium, indium, yttrium, lithium, gadolinium, aluminum, silver, tin and lead, or alloys thereof; multilayer structure materials such as LiF/Al or LiO₂/Al, and the like, but are not limited thereto.

As the hole injection material, known hole injection materials may be used, and for example, phthalocyanine compounds such as copper phthalocyanine disclosed in U.S. Pat. No. 4,356,429, or starburst-type amine derivatives such as tris(4-carbazoyl-9-ylphenyl)amine (TCTA), 4,4',4"-tri[phenyl(m-tolyl)amino]triphenylamine (m-MTDATA) or 1,3,5-tris[4-(3-methylphenyl)phenylamino]phenyl]benzene (m-MTDAPB) described in the literature [Advanced Material, 6, p. 677 (1994)], polyaniline/dodecylbenzene sulfonic acid, poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate), polyaniline/camphor sulfonic acid or polyaniline/poly(4-styrenesulfonate) that are conductive polymers having solubility, and the like, may be used.

As the hole transfer material, pyrazoline derivatives, arylamine-based derivatives, stilbene derivatives, triphenyldiamine derivatives and the like may be used, and low molecular or high molecular materials may also be used.

As the electron transfer material, metal complexes of oxadiazole derivatives, anthraquinodimethane and derivatives thereof, benzoquinone and derivatives thereof, naphthoquinone and derivatives thereof, anthraquinone and derivatives thereof, tetracyanoanthraquinodimethane and derivatives thereof, fluorenone derivatives, diphenyldicyanoethylene and derivatives thereof, diphenoquinone derivatives, 8-hydroxyquinoline and derivatives thereof, and the like, may be used, and high molecular materials may also be used as well as low molecular materials.

As examples of the electron injection material, LiF is typically used in the art, however, the present application is not limited thereto.

As the light emitting material, red, green or blue light emitting materials may be used, and as necessary, two or more light emitting materials may be mixed and used. Herein, two or more light emitting materials may be used by being deposited as individual sources of supply or by being premixed and deposited as one source of supply. In addition, fluorescent materials may also be used as the light emitting material, however, phosphorescent materials may also be used. As the light emitting material, materials emitting light by bonding electrons and holes injected from an anode and a cathode, respectively, may be used alone, however, materials having a host material and a dopant material involving in light emission together may also be used.

When mixing light emitting material hosts, same series hosts may be mixed, or different series hosts may be mixed. For example, any two or more types of materials among n-type host materials or p-type host materials may be selected and used as a host material of a light emitting layer.

The organic light emitting device according to one embodiment of the present specification may be a top-emission type, a bottom-emission type or a dual-emission type depending on the materials used.

The heterocyclic compound according to one embodiment of the present specification may also be used in an organic electronic device including an organic solar cell, an organic photo conductor, an organic transistor and the like under a similar principle used in the organic light emitting device.


Hereinafter, the present specification will be described in more detail with reference to examples, however, these are

268


for illustrative purposes only, and the scope of the present application is not limited thereto.

PREPARATION EXAMPLE

[Preparation Example 1] Preparation of Compound 1



269



1) Preparation of Compound 1-1

After dissolving 4,7-dibromobenzo[c][1,2,5]thiadiazole (49 g, 166.6 mmol) and 2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (30 g, 136.9 mmol) in a mixed solution of toluene (1000 mL), ethanol (200 mL) and H₂O (200 mL), Pd(PPh₃)₄ (tetrakis(triphenylphosphine)palladium(0)) (7.9 g, 6.845 mmol) and K₃PO₄ (87.1 g, 410.7 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. Subsequently, the solvent was removed using a rotary evaporator, and then the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 1-1 (39 g, yield 90%).

2) Preparation of Compound 1-2

After dissolving Compound 1-1 (39 g, 127.3 mmol) in methylene chloride (500 mL), triethylamine (18 mL, 127.3 mmol) was introduced thereto. After lowering the temperature from room temperature to 0° C., benzoyl chloride (19.8 g, 137.74 mmol) dissolved in methylene chloride was slowly added dropwise thereto. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 1-2 (45 g, yield 86%).

3) Preparation of Compound 1-3

After dissolving Compound 1-2 (45 g, 109.6 mmol) in nitrobenzene (500 mL), POCl₃ (25 g, 164.4 mmol) was slowly added dropwise thereto, and then the result was stirred for 4 hours at 150° C. After the reaction was completed, the result was neutralized with an aqueous NaHCO₃ solution and then extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 1-3 (40 g, yield 92%).

4) Preparation of Compound 1-4

After dissolving Compound 1-3 (40 g, 101.9 mmol) and bis(pinacolato)diboron (32.63 g, 128.48 mmol) in 1,4-dioxane (400 mL), Pd(dppf)Cl₂ ([1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium (II)) (2.51 g, 3.43 mmol) and KOAc (potassium acetate) (25.22 g, 256.96 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with ethyl acetate and H₂O. The organic layer


270

was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was passed through silica gel to obtain Compound 1-4 (40 g, yield 89%).

5) Preparation of Compound 1


After adding 2-chloro-4,6-diphenyl-1,3,5-triazine (6.1 g, 22.76 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol) and toluene/ethanol/H₂O to Compound 1-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 1 (9.3 g, yield 76%).

[Preparation Example 2] Preparation of Compound 16




271

-continued





1-4



16

After adding 4-chloro-2,6-diphenylpyrimidine (6.1 g, 22.76 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol) and toluene/ethanol/H₂O to Compound 1-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 16 (9.4 g, yield 76%).


[Preparation Example 3] Preparation of Compound 31




1-1

272


-continued




1-2



1-3



1-4




31

After adding 2-(4-bromophenyl)-4,6-diphenyl-1,3,5-triazine (8.8 g, 22.76 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol) and toluene/ethanol/H₂O to Compound 1-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by


273

column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 31 (10.1 g, yield 72%).

[Preparation Example 4] Preparation of Compound 46


**274**

-continued




After adding 4-(4-bromophenyl)-2,6-diphenylpyrimidine (8.8 g, 22.76 mmol), $\text{Pd}(\text{PPh}_3)_4$ (1.25 g, 1.09 mmol), K_2CO_3 (25.0 g, 65.4 mmol) and toluene/ethanol/ H_2O to Compound 1-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110°C . After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO_4 , and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 46 (10.2 g, yield 72%).

[Preparation Example 5] Preparation of Compound 61




275


-continued



1-3



1-4




61


After adding 2-(3-bromophenyl)-4,6-diphenyl-1,3,5-triazine (8.8 g, 22.76 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol) and toluene/ethanol/H₂O to Compound 1-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 61 (10.9 g, yield 77%).

[Preparation Example 6] Preparation of Compound


76

**276**


-continued




1-1




1-2



1-3



1-4




76



277

After adding 4-(3-bromophenyl)-2,6-diphenylpyrimidine (8.8 g, 22.76 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol) and toluene/ethanol/H₂O to Compound 1-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 76 (10.5 g, yield 74%).

[Preparation Example 7] Preparation of Compound 91


**278**

-continued




After adding 4'-(4-bromophenyl)-2,2':6',2''-terpyridine (8.8 g, 22.76 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol) and toluene/ethanol/H₂O to Compound 1-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 91 (10.7 g, yield 76%).

[Preparation Example 8] Preparation of Compound 107



279

-continued




After adding 2-bromo-9-phenyl-1,10-phenanthroline (7.6 g, 22.76 mmol), $\text{Pd}(\text{PPh}_3)_4$ (1.25 g, 1.09 mmol), K_2CO_3 (25.0 g, 65.4 mmol) and toluene/ethanol/ H_2O to Compound 1-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110°C . After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO_4 , and after removing the solvent using a rotary evaporator, the result was purified by column

280

chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 107 (10.0 g, yield 77%).


5

[Preparation Example 9] Preparation of Compound 109



281

-continued



109

1) Preparation of Compound 2-1

After dissolving 4-bromobenzo[c][1,2,5]thiadiazole (50 g, 232.5 mmol) and 2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (50.9 g, 232.5 mmol) in toluene (1000 mL), ethanol (200 mL) and H₂O (200 mL), Pd(PPh₃)₄ (13.4 g, 11.62 mmol) and K₃PO₄ (148.1 g, 697.5 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. Subsequently, the solvent was removed using a rotary evaporator, and then the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 2-1 (45 g, yield 85%).

2) Preparation of Compound 2-2

After dissolving Compound 2-1 (45 g, 197.9 mmol) in methylene chloride (450 mL), triethylamine (27 mL, 197.9 mmol) was introduced thereto. After lowering the temperature from room temperature to 0° C., 4-bromobenzoyl chloride (47.7 g, 217.6 mmol) dissolved in methylene chloride was slowly added dropwise thereto. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 2-2 (70 g, yield 85%).

3) Preparation of Compound 2-3

After dissolving Compound 2-2 (70 g, 170.6 mmol) in nitrobenzene (700 mL), POCl₃ (39.2 g, 255.9 mmol) was slowly added dropwise thereto, and then the result was stirred for 4 hours at 150° C. After the reaction was completed, the result was neutralized with an aqueous NaHCO₃ solution and then extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 2-3 (60 g, yield 90%).

4) Preparation of Compound 2-4

After dissolving Compound 2-3 (60 g, 152.9 mmol) and bis(pinacolato)diboron (48.95 g, 192.7 mmol) in 1,4-dioxane (600 mL), Pd(dppf)Cl₂ (3.76 g, 5.15 mmol) and


282

KOAc (37.8 g, 385.4 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with ethyl acetate and H₂O. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was passed through silica gel to obtain Compound 2-4 (60 g, yield 89%).


5) Preparation of Compound 109

After adding 2-chloro-4,6-diphenyl-1,3,5-triazine (6.1 g, 22.76 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol) and toluene/ethanol/H₂O to Compound 2-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 109 (9.3 g, yield 76%).


[Preparation Example 10] Preparation of Compound 124



2-1





2-2



2-3

283





-continued



After adding 4-chloro-2,6-diphenylpyrimidine (6.1 g, 22.76 mmol), $\text{Pd}(\text{PPh}_3)_4$ (1.25 g, 1.09 mmol), K_2CO_3 (25.0 g, 65.4 mmol), and toluene/ethanol/ H_2O to Compound 2-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110°C . After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO_4 , and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 124 (9.7 g, yield 78%).


284

[Preparation Example 11] Preparation of Compound 139




285

-continued






After adding 2-(4-bromophenyl)-4,6-diphenyl-1,3,5-triazine (8.8 g, 22.76 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol) and toluene/ethanol/H₂O to Compound 2-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 139 (11.0 g, yield 78%).

[Preparation Example 12] Preparation of Compound 144


**286**

-continued




287

-continued






After adding 9-(4-(4-(4-bromophenyl)-6-phenyl-1,3,5-triazin-2-yl)phenyl)-9H-carbazole (12.6 g, 22.76 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol) and toluene/ethanol/H₂O to Compound 2-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 144 (14.0 g, yield 78%).

[Preparation Example 13] Preparation of Compound 154


**288**

-continued



289

-continued




154


After adding 4-(4-bromophenyl)-2,6-diphenylpyrimidine (8.8 g, 22.76 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol) and toluene/ethanol/H₂O to Compound 2-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 154 (11.2 g, yield 79%).

290


-continued



2-2




2-3




2-4

[Preparation Example 14] Preparation of Compound 174



2-1






174

After adding 9-(4-(4-(3-bromophenyl)-6-phenyl-1,3,5-triazin-2-yl)phenyl)-9H-carbazole (12.6 g, 22.76 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol)

291


and toluene/ethanol/H₂O to Compound 2-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 174 (14.0 g, 78%).

[Preparation Example 15] Preparation of Compound 199



292





-continued




After adding 4'-(4-bromophenyl)-2,2':6',2''-terpyridine (8.8 g, 22.76 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol) and toluene/ethanol/H₂O to Compound 2-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 199 (10.7 g, yield 76%).

293

[Preparation Example 16] Preparation of Compound 213



**294**

-continued




After adding (4-bromophenyl)diphenylphosphine oxide (8.1 g, 22.76 mmol), $\text{Pd}(\text{PPh}_3)_4$ (1.25 g, 1.09 mmol), K_2CO_3 (25.0 g, 65.4 mmol) and toluene/ethanol/ H_2O to Compound 2-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110°C . After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO_4 , and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 213 (10.7 g, yield 79%).

[Preparation Example 17] Preparation of Compound 215



295

-continued




After adding 2-bromo-9-phenyl-1,10-phenanthroline (7.6 g, 22.76 mmol), $\text{Pd}(\text{PPh}_3)_4$ (1.25 g, 1.09 mmol), K_2CO_3 (25.0 g, 65.4 mmol) and toluene/ethanol/ H_2O to Compound 2-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110°C . After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO_4 , and after removing the solvent using a rotary evaporator, the result was purified by column

296


chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 215 (10.5 g, yield 81%).

[Preparation Example 18] Preparation of Compound 217



297

-continued



217

298

5) Preparation of Compound 217

After adding 2-chloro-4,6-diphenyl-1,3,5-triazine (6.1 g, 22.76 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol) and toluene/ethanol/H₂O to Compound 3-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 217 (9.3 g, yield 76%).

[Preparation Example 19] Preparation of Compound 232

1) Preparation of Compound 3-1

After dissolving 4-bromobenzo[c][1,2,5]thiadiazole (50 g, 232.5 mmol) and 5-chloro-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl) aniline (58.9 g, 232.5 mmol) in toluene (1000 mL), ethanol (200 mL) and H₂O (200 mL), Pd(PPh₃)₄ (13.4 g, 11.63 mmol) and K₃PO₄ (148 g, 97.5 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 3-1 (50 g, yield 82%).

2) Preparation of Compound 3-2



After dissolving Compound 3-1 (50 g, 191 mmol) in methylene chloride (500 mL), triethylamine (26.6 mL, 191 mmol) was introduced thereto. After lowering the temperature from room temperature to 0° C., benzoyl chloride (29.5 g, 210.1 mmol) dissolved in methylene chloride was slowly added dropwise thereto. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 3-2 (60 g, yield 86%).

3) Preparation of Compound 3-3


After dissolving Compound 3-2 (60 g, 164 mmol) in nitrobenzene (600 mL), POCl₃ (37.7 g, 246 mmol) was slowly added dropwise thereto, and then the result was stirred for 4 hours at 150° C. After the reaction was completed, the result was neutralized with an aqueous NaHCO₃ solution and then extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 3-3 (50 g, yield 87%).

4) Preparation of Compound 3-4


After dissolving Compound 3-3 (50 g, 143.7 mmol) and bis(pinacolato)diboron (54.7 g, 215.6 mmol) in 1,4-dioxane (500 mL), Pd(dppf)Cl₂ (5.02 g, 6.86 mmol) and KOAc (42.3 g, 431.1 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with ethyl acetate and H₂O. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was passed through silica gel to obtain Compound 3-4 (50 g, yield 79%).



3-1




3-2




3-3

299

-continued





3-4



232

After adding 4-chloro-2,6-diphenylpyrimidine (6.1 g, 22.76 mmol), $\text{Pd}(\text{PPh}_3)_4$ (1.25 g, 1.09 mmol), K_2CO_3 (25.0 g, 65.4 mmol) and toluene/ethanol/ H_2O to Compound 3-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO_4 , and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 232 (9.7 g, 78%).


[Preparation Example 20] Preparation of Compound 247




3-1

300


-continued




3-2



3-3



3-4








247

After adding 2-(4-bromophenyl)-4,6-diphenyl-1,3,5-triazine (8.8 g, 22.76 mmol), $\text{Pd}(\text{PPh}_3)_4$ (1.25 g, 1.09 mmol), K_2CO_3 (25.0 g, 65.4 mmol) and toluene/ethanol/ H_2O to Compound 3-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO_4 , and after removing the solvent using a rotary evaporator, the result was purified by

301


column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 247 (11.1 g, yield 78%).

[Preparation Example 21] Preparation of Compound 253





302

-continued







After adding 9-(3-(4-(4-bromophenyl)-6-phenyl-1,3,5-triazin-2-yl)phenyl)-9H-carbazole (12.6 g, 22.76 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol) and toluene/ethanol/H₂O to Compound 3-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 253 (13.0 g, yield 73%).

[Preparation Example 22] Preparation of Compound 262



303

-continued








After adding 4-(4-bromophenyl)-2,6-diphenylpyrimidine (8.8 g, 22.76 mmol), $\text{Pd}(\text{PPh}_3)_4$ (1.25 g, 1.09 mmol), K_2CO_3 (25.0 g, 65.4 mmol) and toluene/ethanol/ H_2O to Compound 3-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110°C . After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO_4 , and after removing the solvent using a rotary evaporator, the result was purified by column

304


chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 262 (10.9 g, yield 77%).

[Preparation Example 23] Preparation of Compound 268





305

-continued







After adding 9-(3-(4-(4-bromophenyl)-6-phenylpyrimidin-2-yl)phenyl)-9H-carbazole (12.6 g, 22.76 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol) and toluene/ethanol/H₂O to Compound 3-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 268 (13.5 g, yield 75%).

[Preparation Example 24] Preparation of Compound 283

**306**

-continued







After adding 9-(3-(4-(3-bromophenyl)-6-phenyl-1,3,5-triazin-2-yl)phenyl)-9H-carbazole (12.6 g, 22.76 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol)

307



and toluene/ethanol/H₂O to Compound 3-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 283 (13.1 g, yield 73%).

[Preparation Example 25] Preparation of Compound 298




308

-continued








After adding 9-(3-(4-(3-bromophenyl)-6-phenylpyrimidin-2-yl)phenyl)-9H-carbazole (12.6 g, 22.76 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol) and toluene/ethanol/H₂O to Compound 3-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 298 (13.7 g, yield 73%).

[Preparation Example 26] Preparation of Compound 307



309

-continued







After adding 4'-(4-bromophenyl)-2,2':6',2''-terpyridine (8.8 g, 22.76 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃

310


(25.0 g, 65.4 mmol) and toluene/ethanol/H₂O to Compound 3-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 307 (10.9 g, yield 77%).

[Preparation Example 27] Preparation of Compound 323




311

-continued




3-4



323

After adding 2-bromo-9-phenyl-1,10-phenanthroline (7.6 g, 22.76 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol) and toluene/ethanol/H₂O to Compound 3-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 323 (9.9 g, yield 76%).


[Preparation Example 28] Preparation of Compound 325




4-1

312


-continued




4-2



4-3



4-4



325

1) Preparation of Compound 4-1

After dissolving 4,7-dibromobenzo[c][1,2,5]oxadiazole (50 g, 179.9 mmol) and 2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (39.4 g, 179.9 mmol) in toluene (1000 mL), ethanol (200 mL) and H₂O (200 mL), Pd(PPh₃)₄ (10.4 g, 8.995 mmol) and K₃PO₄ (114.5 g, 539.7 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the

313

result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 4-1 (45 g, yield 86%).

2) Preparation of Compound 4-2

After dissolving Compound 4-1 (45 g, 155.1 mmol) in methylene chloride (500 mL), triethylamine (21.6 mL, 155.1 mmol) was introduced thereto. After lowering the temperature from room temperature to 0° C., benzoyl chloride (23.9 g, 170.61 mmol) dissolved in methylene chloride was slowly added dropwise thereto. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 4-2 (50 g, yield 82%).

3) Preparation of Compound 4-3

After dissolving Compound 4-2 (50 g, 126.8 mmol) in nitrobenzene (500 mL), POCl₃ (25 g, 164.4 mmol) was slowly added dropwise thereto, and then the result was stirred for 4 hours at 150° C. After the reaction was completed, the result was neutralized with an aqueous NaHCO₃ solution and then extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 4-3 (40 g, yield 83%).


4) Preparation of Compound 4-4

After dissolving Compound 4-3 (40 g, 106.3 mmol) and bis(pinacolato)diboron (40.4 g, 159.4 mmol) in 1,4-dioxane (400 mL), Pd(dppf)Cl₂ (4.3 g, 5.31 mmol) and KOAc (31.2 g, 318.9 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with ethyl acetate and H₂O. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was passed through silica gel to obtain Compound 4-4 (40 g, yield 88%).

5) Preparation of Compound 325






After adding 2-(4-bromophenyl)-4,6-diphenyl-1,3,5-triazine (9.1 g, 23.62 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol) and toluene/ethanol/H₂O to Compound 4-4 (10.0 g, 23.62 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 325 (11 g, yield 76%).

[Preparation Example 29] Preparation of Compound 326



314

-continued







326

After adding 4-(4-bromophenyl)-2,6-diphenylpyrimidine (9.1 g, 23.62 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃

315


(25.0 g, 65.4 mmol) and toluene/ethanol/H₂O to Compound 4-4 (10.0 g, 23.62 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 326 (11.5 g, yield 80%).

[Preparation Example 30] Preparation of Compound 337



316

-continued



1) Preparation of Compound 5-1

After dissolving 4-bromobenzo[c][1,2,5]oxadiazole (50 g, 251.2 mmol) and 2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (55 g, 251.2 mmol) in toluene (1000 mL), ethanol (200 mL) and H₂O (200 mL), Pd(PPh₃)₄ (14.5 g, 12.56 mmol) and K₃PO₄ (159.9 g, 753.6 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 5-1 (45 g, yield 85%).

2) Preparation of Compound 5-2

After dissolving Compound 5-1 (45 g, 213.0 mmol) in methylene chloride (500 mL), triethylamine (29.7 mL, 213.0 mmol) was introduced thereto. After lowering the temperature from room temperature to 0° C., 4-bromobenzoyl chloride (51.4 g, 234.3 mmol) dissolved in methylene chloride was slowly added dropwise thereto. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 5-2 (70 g, yield 83%).

3) Preparation of Compound 5-3

After dissolving Compound 5-2 (50 g, 126.8 mmol) in nitrobenzene (500 mL), POCl₃ (25 g, 164.4 mmol) was slowly added dropwise thereto, and then the result was stirred for 4 hours at 150° C. After the reaction was completed, the result was neutralized with an aqueous NaHCO₃ solution and then extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 5-3 (42 g, yield 87%).

4) Preparation of Compound 5-4

After dissolving Compound 5-3 (40 g, 106.3 mmol) and bis(pinacolato)diboron (40.4 g, 159.4 mmol) in 1,4-dioxane (400 mL), Pd(dppf)Cl₂ (4.3 g, 5.31 mmol) and KOAc (31.2




317

g, 318.9 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with ethyl acetate and H₂O. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was passed through silica gel to obtain Compound 5-4 (41 g, yield 90%).

5) Preparation of Compound 337




After adding 2-(4-bromophenyl)-4,6-diphenyl-1,3,5-triazine (9.1 g, 23.62 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol) and toluene/ethanol/H₂O to Compound 5-4 (10.0 g, 23.62 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 337 (10 g, yield 70%).

[Preparation Example 31] Preparation of Compound 338




318

-continued








After adding 4-(4-bromophenyl)-2,6-diphenylpyrimidine (9.1 g, 23.62 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol) and toluene/ethanol/H₂O to Compound 5-4 (10.0 g, 23.62 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 338 (10.4 g, yield 72%).

[Preparation Example 32] Preparation of Compound 353




319

-continued



320

-continued



After dissolving 4,7-dibromobenzo[c][1,2,5]thiadiazole (50 g, 170.1 mmol) and phenylboronic acid (20.7 g, 170.1 mmol) in toluene (1000 mL), ethanol (200 mL) and H₂O (200 mL), Pd(PPh₃)₄ (7.9 g, 6.845 mmol) and K₃PO₄ (87.1 g, 410.7 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 6-1 (45 g, yield 91%).

2) Preparation of Compound 6-2

After dissolving Compound 6-1 (45 g, 154.5 mmol) and 2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl) aniline (33.8 g, 154.5 mmol) in toluene (900 mL), ethanol (180 mL) and H₂O (180 mL), Pd(PPh₃)₄ (8.9 g, 7.72 mmol) and K₃PO₄ (63.9 g, 463.5 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 6-2 (38.7 g, yield 83%).

3) Preparation of Compound 6-3

After dissolving Compound 6-2 (38.7 g, 127.3 mmol) in methylene chloride (387 mL), triethylamine (18 mL, 127.3 mmol) was introduced thereto. After lowering the temperature from room temperature to 0° C., 4-bromobenzoyl chloride (27.9 g, 137.74 mmol) dissolved in methylene chloride was slowly added dropwise thereto. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 6-3 (52 g, yield 84%).

321

4) Preparation of Compound 6-4

After dissolving Compound 6-3 (52 g, 106.9 mmol) in nitrobenzene (520 mL), POCl_3 (24.5 g, 160.3 mmol) was slowly added dropwise thereto, and then the result was stirred for 4 hours at 150°C . After the reaction was completed, the result was neutralized with an aqueous NaHCO_3 solution and then extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO_4 . After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 6-4 (45 g, yield 90%).



5) Preparation of Compound 6-5

After dissolving Compound 6-4 (40 g, 85.4 mmol) and bis(pinacolato)diboron (32.63 g, 128.48 mmol) in 1,4-dioxane (400 mL), Pd(dppf)Cl_2 (3.48 g, 4.27 mmol) and KOAc (25.22 g, 256.96 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with ethyl acetate and H_2O . The organic layer was dried with anhydrous MgSO_4 , and after removing the solvent using a rotary evaporator, the result was passed through silica gel to obtain Compound 6-5 (40 g, yield 90%).


6) Preparation of Compound 353

After adding 2-(4-bromophenyl)-4,6-diphenyl-1,3,5-triazine (7.5 g, 19.4 mmol), $\text{Pd(PPh}_3)_4$ (1.25 g, 1.09 mmol), K_2CO_3 (25.0 g, 65.4 mmol) and toluene/ethanol/ H_2O to Compound 6-5 (10.0 g, 19.4 mmol), the result was stirred for 6 hours at 110°C . After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO_4 , and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 353 (10 g, yield 74%).

[Preparation Example 33] Preparation of Compound 354




6-1




6-2

322


-continued



6-3




6-4



6-5

323




-continued



354

After adding 4-(4-bromophenyl)-2,6-diphenylpyrimidine (7.5 g, 19.4 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol) and toluene/ethanol/H₂O to Compound 6-5 (10.0 g, 19.4 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 354 (10.5 g, yield 77%).


[Preparation Example 34] Preparation of Compound 361




7-2

324


-continued



7-3




7-4



7-5

325



1) Preparation of Compound 7-1

After dissolving 4,7-dibromobenzo[c][1,2,5]oxadiazole (55.5 g, 200 mmol) and phenylboronic acid (24.4 g, 200 mmol) in toluene (1000 mL), ethanol (200 mL) and H₂O (200 mL), Pd(PPh₃)₄ (11.5 g, 10 mmol) and K₃PO₄ (127.3 g, 600 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 7-1 (49 g, yield 89%).

2) Preparation of Compound 7-2

After dissolving Compound 7-1 (42.5 g, 154.5 mmol) and 2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (33.8 g, 154.5 mmol) in toluene (1000 mL), ethanol (200 mL) and H₂O (200 mL), Pd(PPh₃)₄ (8.9 g, 7.72 mmol) and K₃PO₄ (63.9 g, 463.5 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 7-2 (37.7 g, yield 83%).

3) Preparation of Compound 7-3

After dissolving Compound 7-2 (36.6 g, 127.3 mmol) in methylene chloride (450 mL), triethylamine (18 mL, 127.3 mmol) was introduced thereto. After lowering the temperature from room temperature to 0° C., 4-bromobenzoyl chloride (27.9 g, 137.74 mmol) dissolved in methylene chloride was slowly added dropwise thereto. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 7-3 (50.3 g, yield 84%).

326

4) Preparation of Compound 7-4

After dissolving Compound 7-3 (50.2 g, 106.9 mmol) in nitrobenzene (502 mL), POCl₃ (24.5 g, 160.3 mmol) was slowly added dropwise thereto, and then the result was stirred for 4 hours at 150° C. After the reaction was completed, the result was neutralized with an aqueous NaHCO₃ solution and then extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 7-4 (40 g, yield 83%).


5) Preparation of Compound 7-5

After dissolving Compound 7-4 (38.6 g, 85.4 mmol) and bis(pinacolato)diboron (32.63 g, 128.48 mmol) in 1,4-dioxane (400 mL), Pd(dppf)Cl₂ (3.48 g, 4.27 mmol) and KOAc (25.22 g, 256.96 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with ethyl acetate and H₂O. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was passed through silica gel to obtain Compound 7-5 (39 g, yield 91%).

6) Preparation of Compound 361


After adding 2-(4-bromophenyl)-4,6-diphenyl-1,3,5-triazine (7.76 g, 20.0 mmol), Pd(PPh₃)₄ (1.15 g, 1.00 mmol), K₂CO₃ (8.29 g, 60.0 mmol) and toluene/ethanol/H₂O to Compound 7-5 (10.0 g, 20.0 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 361 (10.4 g, yield 76%).

[Preparation Example 35] Preparation of Compound 362





327

-continued



**328**

-continued






After adding 4-(4-bromophenyl)-2,6-diphenylpyrimidine (7.71 g, 20.0 mmol), Pd(PPh₃)₄ (1.15 g, 1.00 mmol), K₂CO₃ (8.29 g, 60.0 mmol) and toluene/ethanol/H₂O to Compound 7-5 (10.0 g, 20.0 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 362 (10.7 g, yield 78%).

[Preparation Example 36] Preparation of Compound 409




329

-continued

**330**

-continued



1) Preparation of Compound 8-1

After dissolving 4,7-dibromobenzo[c][1,2,5]oxadiazole (55.5 g, 200 mmol) and phenylboronic acid (24.4 g, 200 mmol) in toluene (1000 mL), ethanol (200 mL) and H₂O (200 mL), Pd(PPh₃)₄ (11.5 g, 10 mmol) and K₃PO₄ (127.3 g, 600 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 8-1 (49 g, yield 89%).

2) Preparation of Compound 8-2

After dissolving Compound 8-1 (42.5 g, 154.5 mmol) and 4-chloro-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl) aniline (39.2 g, 154.5 mmol) in toluene (1000 mL), ethanol (200 mL) and H₂O (200 mL), Pd(PPh₃)₄ (8.9 g, 7.72 mmol) and K₃PO₄ (63.9 g, 463.5 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 8-2 (41.2 g, yield 83%).

3) Preparation of Compound 8-3

After dissolving Compound 8-2 (40.9 g, 127.3 mmol) in methylene chloride (480 mL), triethylamine (18 mL, 127.3 mmol) was introduced thereto. After lowering the temperature from room temperature to 0° C., 4-bromobenzoyl chloride (27.9 g, 137.74 mmol) dissolved in methylene chloride was slowly added dropwise thereto. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column

331

chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 8-3 (49 g, yield 90%).

4) Preparation of Compound 8-4

After dissolving Compound 8-3 (45.5 g, 106.9 mmol) in nitrobenzene (455 mL), POCl_3 (24.5 g, 160.3 mmol) was slowly added dropwise thereto, and then the result was stirred for 4 hours at 150°C . After the reaction was completed, the result was neutralized with an aqueous NaHCO_3 solution and then extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO_4 . After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 8-4 (38 g, yield 87%).




5) Preparation of Compound 8-5

After dissolving Compound 8-4 (34.8 g, 85.4 mmol) and bis(pinacolato)diboron (32.63 g, 128.48 mmol) in 1,4-dioxane (400 mL), Pd(dppf)Cl_2 (3.48 g, 4.27 mmol) and KOAc (25.22 g, 256.96 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with ethyl acetate and H_2O . The organic layer was dried with anhydrous MgSO_4 , and after removing the solvent using a rotary evaporator, the result was passed through silica gel to obtain Compound 8-5 (36 g, yield 84%).

6) Preparation of Compound 409

After adding 2-(4-bromophenyl)-4,6-diphenyl-1,3,5-triazine (7.76 g, 20.0 mmol), $\text{Pd(PPh}_3)_4$ (1.15 g, 1.00 mmol), K_2CO_3 (8.29 g, 60.0 mmol) and toluene/ethanol/ H_2O to Compound 8-5 (10.0 g, 20.0 mmol), the result was stirred for 6 hours at 110°C . After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO_4 , and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 409 (9.9 g, yield 72%).


[Preparation Example 37] Preparation of Compound 411




8-2

332


-continued



8-3




8-4



8-5



333

-continued







After adding 4'-(4-bromophenyl)-2,2':6',2''-terpyridine (7.76 g, 20.0 mmol), Pd(PPh₃)₄ (1.15 g, 1.00 mmol), K₂CO₃ (8.29 g, 60.0 mmol) and toluene/ethanol/H₂O to Compound 8-5 (10.0 g, 20.0 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 411 (10.2 g, yield 75%).

[Preparation Example 38] Preparation of Compound 412


**334**

-continued





335

-continued






After adding 2-bromo-9-phenyl-1,10-phenanthroline (6.7 g, 20.0 mmol), Pd(PPh₃)₄ (1.15 g, 1.00 mmol), K₂CO₃ (8.29 g, 60.0 mmol) and toluene/ethanol/H₂O to Compound 8-5 (10.0 g, 20.0 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain red Compound 412 (10.1 g, yield 80%).

[Preparation Example 39] Preparation of Compound 417


**336**

-continued



337

-continued



1) Preparation of Compound 9-1

After dissolving 4,7-dibromo-2-phenyl-2H-benzo[d][1,2,3]triazole (35.3 g, 100 mmol) and phenylboronic acid (12.2 g, 100 mmol) in toluene (700 mL), ethanol (14 mL) and H₂O (14 mL), Pd(PPh₃)₄ (5.75 g, 5 mmol) and K₃PO₄ (65.7 g, 300 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 9-1 (30 g, yield 85%).

2) Preparation of Compound 9-2

After dissolving Compound 9-1 (27.05 g, 77.25 mmol) and 4-chloro-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl) aniline (39.2 g, 77.25 mmol) in toluene (770 mL), ethanol (154 mL) and H₂O (154 mL), Pd(PPh₃)₄ (4.45 g, 3.86 mmol) and K₃PO₄ (63.9 g, 231.75 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was

338

extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 9-2 (26 g, yield 84%).

3) Preparation of Compound 9-3

After dissolving Compound 9-2 (25.3 g, 63.65 mmol) in methylene chloride (300 mL), triethylamine (9 mL, 63.65 mmol) was introduced thereto. After lowering the temperature from room temperature to 0° C., 4-bromobenzoyl chloride (13.95 g, 68.87 mmol) dissolved in methylene chloride was slowly added dropwise thereto. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 9-3 (49 g, yield 90%).

4) Preparation of Compound 9-4

After dissolving Compound 9-3 (26.7 g, 53.45 mmol) in nitrobenzene (267 mL), POCl₃ (12.25 g, 80.15 mmol) was slowly added dropwise thereto, and then the result was stirred for 4 hours at 150° C. After the reaction was completed, the result was neutralized with an aqueous NaHCO₃ solution and then extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 9-4 (22.8 g, yield 88%).


5) Preparation of Compound 9-5

After dissolving Compound 9-4 (20.6 g, 42.7 mmol) and bis(pinacolato)diboron (16.3 g, 64.24 mmol) in 1,4-dioxane (210 mL), Pd(dppf)Cl₂ (1.74 g, 2.13 mmol) and KOAc (12.56 g, 128.4 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with ethyl acetate and H₂O. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was passed through silica gel to obtain Compound 9-5 (20.1 g, yield 82%).

6) Preparation of Compound 417






After adding 2-(4-bromophenyl)-4,6-diphenyl-1,3,5-triazine (6.75 g, 17.4 mmol), Pd(PPh₃)₄ (1.00 g, 0.87 mmol), K₂CO₃ (7.2 g, 52.2 mmol) and toluene/ethanol/H₂O to Compound 9-5 (10.0 g, 17.4 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 417 (10 g, yield 76%).

[Preparation Example 40] Preparation of Compound 425




339

-continued

**340**

-continued



1) Preparation of Compound 10-1

After dissolving 4,7-dibromobenzo[c][1,2,5]thiadiazole (58.8 g, 200 mmol) and phenylboronic acid (24.4 g, 200 mmol) in toluene (1000 mL), ethanol (200 mL) and H₂O (200 mL), Pd(PPh₃)₄ (11.5 g, 10 mmol) and K₃PO₄ (127.3 g, 600 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 10-1 (52 g, yield 89%).

2) Preparation of Compound 10-2

After dissolving Compound 10-1 (45 g, 154.5 mmol) and 5-chloro-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (39.2 g, 154.5 mmol) in toluene (1000 mL), ethanol (200 mL) and H₂O (200 mL), Pd(PPh₃)₄ (8.9 g, 7.72 mmol) and K₃PO₄ (63.9 g, 463.5 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 10-2 (43.4 g, yield 83%).

3) Preparation of Compound 10-3

After dissolving Compound 10-2 (43.0 g, 127.3 mmol) in methylene chloride (430 mL), triethylamine (18 mL, 127.3 mmol) was introduced thereto. After lowering the temperature from room temperature to 0° C., 4-bromobenzoyl chloride (27.9 g, 137.74 mmol) dissolved in methylene chloride was slowly added dropwise thereto. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO₄. After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 10-3 (50.58 g, 90%).

341

4) Preparation of Compound 10-4

After dissolving Compound 10-3 (47.2 g, 106.9 mmol) in nitrobenzene (455 mL), POCl_3 (24.5 g, 160.3 mmol) was slowly added dropwise thereto, and then the result was stirred for 4 hours at 150°C . After the reaction was completed, the result was neutralized with an aqueous NaHCO_3 solution and then extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO_4 . After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 10-4 (39.5 g, yield 87%).


5) Preparation of Compound 10-5

After dissolving Compound 10-4 (36.2 g, 85.4 mmol) and bis(pinacolato)diboron (32.63 g, 128.48 mmol) in 1,4-dioxane (362 mL), Pd(dppf)Cl_2 (3.48 g, 4.27 mmol) and KOAc (25.22 g, 256.96 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with ethyl acetate and H_2O . The organic layer was dried with anhydrous MgSO_4 , and after removing the solvent using a rotary evaporator, the result was passed through silica gel to obtain Compound 10-5 (38 g, yield 86%).

6) Preparation of Compound 425


After adding 2-(4-bromophenyl)-4,6-diphenyl-1,3,5-triazine (7.52 g, 19.4 mmol), $\text{Pd(PPh}_3)_4$ (1.12 g, 0.97 mmol), K_2CO_3 (8.04 g, 58.2 mmol) and toluene/ethanol/ H_2O to Compound 10-5 (10.0 g, 19.4 mmol), the result was stirred for 6 hours at 110°C . After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO_4 , and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 425 (10.2 g, yield 75%).

[Preparation Example 41] Preparation of Compound 426



342


-continued



343


After adding 4-(4-bromophenyl)-2,6-diphenylpyrimidine (7.51 g, 19.4 mmol), $\text{Pd}(\text{PPh}_3)_4$ (1.12 g, 0.97 mmol), K_2CO_3 (8.04 g, 58.2 mmol) and toluene/ethanol/ H_2O to Compound 10-5 (10.0 g, 19.4 mmol), the result was stirred for 6 hours at 110°C . After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO_4 , and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 426 (10.4 g, yield 77%).

[Preparation Example 42] Preparation of Compound 449



344

-continued



1) Preparation of Compound 11-1

After dissolving 4-bromobenzo[c][1,2,5]thiadiazole (45 g, 154.5 mmol) and 2-chloro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl) aniline (39.2 g, 154.5 mmol) in toluene (1000 mL), ethanol (200 mL) and H_2O (200 mL), $\text{Pd}(\text{PPh}_3)_4$ (8.9 g, 7.72 mmol) and K_3PO_4 (63.9 g, 463.5 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO_4 . After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 11-1 (43.4 g, yield 83%).

2) Preparation of Compound 11-2

After dissolving Compound 11-1 (43.0 g, 127.3 mmol) in methylene chloride (430 mL), triethylamine (18 mL, 127.3 mmol) was introduced thereto. After lowering the temperature from room temperature to 0°C , 4-bromobenzoyl chloride (27.9 g, 137.74 mmol) dissolved in methylene chloride was slowly added dropwise thereto. After the reaction was completed, the result was extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO_4 . After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 11-2 (50.58 g, yield 90%).

3) Preparation of Compound 11-3

After dissolving Compound 11-2 (47.2 g, 106.9 mmol) in nitrobenzene (455 mL), POCl_3 (24.5 g, 160.3 mmol) was slowly added dropwise thereto, and then the result was stirred for 4 hours at 150°C . After the reaction was completed, the result was neutralized with an aqueous NaHCO_3 solution and then extracted with methylene chloride and distilled water, and the organic layer was dried with anhydrous MgSO_4 . After removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 11-3 (39.5 g, yield 87%).

4) Preparation of Compound 11-4

After dissolving Compound 11-3 (36.2 g, 85.4 mmol) and bis(pinacolato)diboron (32.63 g, 128.48 mmol) in 1,4-dioxane (362 mL), $\text{Pd}(\text{dppf})\text{Cl}_2$ (3.48 g, 4.27 mmol) and




345

KOAc (25.22 g, 256.96 mmol) were introduced thereto, and the result was stirred for 5 hours under reflux. After the reaction was completed, the result was extracted with ethyl acetate and H₂O. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was passed through silica gel to obtain Compound 11-4 (38 g, yield 86%).

5) Preparation of Compound 449



After adding 2-(4-bromophenyl)-4,6-diphenyl-1,3,5-triazine (7.52 g, 19.4 mmol), Pd(PPh₃)₄ (1.12 g, 0.97 mmol), K₂CO₃ (8.04 g, 58.2 mmol) and toluene/ethanol/H₂O to Compound 11-4 (10.0 g, 19.4 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 449 (10 g, yield 74%).

[Preparation Example 43] Preparation of Compound 450




346

-continued




After adding 4-(4-bromophenyl)-2,6-diphenylpyrimidine (7.51 g, 19.4 mmol), Pd(PPh₃)₄ (1.12 g, 0.97 mmol), K₂CO₃ (8.04 g, 58.2 mmol) and toluene/ethanol/H₂O to Compound 11-4 (10.0 g, 19.4 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 450 (10.4 g, yield 77%).

[Preparation Example 44] Preparation of Compound 473




347


-continued




1-1




1-2



1-3



1-4



473


After adding 4'-bromo-2,4,6-trifluoro-[1,1'-biphenyl]-3, 65
5-dicarbonitrile (7.67 g, 22.76 mmol), Pd(PPh₃)₄ (1.25 g,
1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol) and toluene/

348

ethanol/H₂O to Compound 1-4 (10.0 g, 22.76 mmol), the
result was stirred for 6 hours at 110° C. After the reaction
was completed, the result was cooled to room temperature,
and extracted with distilled water and dichloromethane. The
organic layer was dried with anhydrous MgSO₄, and after
removing the solvent using a rotary evaporator, the result
was purified by column chromatography using dichlo-
romethane and hexane as a developing solvent to obtain
Compound 473 (9.7 g, yield 74%).

[Preparation Example 45] Preparation of
Compound 485


15



20

25

30



2-1

35


40

45


50

55

60




2-2



2-3


349

-continued




After adding 4'-bromo-2,3,5,6-tetrafluoro-[1,1'-biphenyl]-4-carbonitrile (7.5 g, 22.76 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol) and toluene/ethanol/H₂O to Compound 2-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 485 (10.1 g, yield 79%).

[Preparation Example 46] Preparation of Compound 505

**350**

-continued







After adding 4'-bromo-2,4,6-trifluoro-[1,1'-biphenyl]-3,5-dicarbonitrile (7.6 g, 22.76 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol) and toluene/ethanol/

351



H₂O to Compound 3-4 (10.0 g, 22.76 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 505 (9.5 g, yield 73%).

[Preparation Example 47] Preparation of Compound 527



352

-continued



After adding 4'-bromo-2,3,5,6-tetrafluoro-[1,1'-biphenyl]-4-carbonitrile (7.5 g, 23.62 mmol), Pd(PPh₃)₄ (1.25 g, 1.09 mmol), K₂CO₃ (25.0 g, 65.4 mmol), and toluene/ethanol/H₂O to Compound 3-4 (10.0 g, 23.61 mmol), the result was stirred for 6 hours at 110° C. After the reaction was completed, the result was cooled to room temperature, and extracted with distilled water and dichloromethane. The organic layer was dried with anhydrous MgSO₄, and after removing the solvent using a rotary evaporator, the result was purified by column chromatography using dichloromethane and hexane as a developing solvent to obtain Compound 527 (9.4 g, yield 75%).

Compounds were prepared in the same manner as in the preparation examples, and the synthesis identification results are shown in the following Tables 1 and 2.

TABLE 1

NO	¹ H NMR (CDCl ₃ , 300 Mz)
1	8.36 (4H, d), 8.20-8.19 (4H, m), 7.94 (1H, d), 7.85 (1H, m), 7.70-7.65 (3H, m), 7.50-7.49 (7H, m)
16	8.35 (2H, m), 8.23-8.19 (5H, m), 7.94-7.85 (4H, m), 7.70-7.49 (10H, m)
31	8.36 (4H, d), 8.20-8.19 (3H, m), 8.07 (1H, s), 7.96-7.94 (3H, m), 7.85 (1H, m), 7.70-7.65 (3H, m), 7.50-7.49 (7H, m), 7.25 (2H, d)
46	8.35-8.30 (4H, m), 8.23-8.19 (4H, m), 8.07 (1H, s), 7.94-7.85 (4H, m), 7.70-7.49 (10H, m), 7.25 (2H, d)
61	8.38-8.36 (5H, m), 8.19 (2H, m), 8.07 (1H, s), 7.94 (2H, m), 7.85 (1H, m), 7.73-7.61 (5H, m), 7.50-7.49 (7H, m)
76	8.35 (2H, d), 8.23-8.19 (4H, m), 8.07 (1H, s), 7.94-7.85 (6H, m), 7.73-7.49 (12H, m)
91	9.18-9.14 (4H, m), 8.55 (1H, d), 8.20-8.19 (3H, m), 8.07 (1H, s), 7.94 (1H, m), 7.85 (1H, m), 7.74-7.65 (5H, m), 7.49 (1H, m), 7.25-7.23 (6H, m)
107	8.71 (2H, d), 8.60 (1H, s), 8.33 (2H, m), 8.20-8.19 (4H, m), 7.94-7.85 (3H, m), 7.70-7.49 (7H, m), 7.29 (2H, d)

353

TABLE 1-continued

NO	¹ H NMR (CDCl ₃ , 300 Mz)
109	8.69 (2H, d), 8.36 (4H, d), 8.20 (1H, d), 7.99-7.94 (4H, m), 7.85 (1H, m), 7.70 (1H, m), 7.50 (7H, m)
139	8.69 (2H, d), 8.36 (4H, m), 8.20 (1H, d), 7.99-7.94 (4H, m), 7.85 (3H, m), 7.70 (1H, m), 7.50 (7H, m), 7.25 (2H, d)
144	8.69 (2H, d), 8.55 (1H, d), 8.36 (2H, d), 8.20-8.19 (2H, m), 7.96-7.85 (12H, m), 7.70 (1H, m), 7.58-7.50 (6H, m), 7.35 (1H, m), 7.25-7.16 (4H, m)
174	8.69 (2H, d), 8.55 (1H, d), 8.38-8.36 (3H, m), 8.20-8.19 (2H, m), 7.99-7.85 (11H, m), 7.73-7.70 (2H, m), 7.61-7.50 (7H, m), 7.35 (1H, m), 7.20-7.16 (2H, m)
199	9.18-9.14 (4H, m), 8.69 (2H, d), 8.55 (2H, d), 8.20 (1H, d), 7.99-7.94 (2H, m), 7.85 (3H, m), 7.74-7.70 (3H, m), 7.50 (1H, d), 7.25-7.23 (6H, m)
213	8.69 (2H, d), 8.20 (1H, d), 7.99-7.94 (6H, m), 7.85-7.70 (8H, m), 7.51-7.50 (7H, m)
215	8.71-8.69 (6H, m), 8.33 (2H, d), 8.20 (2H, m), 7.99-7.85 (4H, m), 7.70 (1H, m) 7.55-7.49 (4H, m), 7.29 (2H, d)
217	8.39-8.36 (6H, m), 8.24-8.19 (3H, m), 7.99 (1H, d), 7.65 (2H, m), 7.50-7.49 (8H, m)
232	8.39-8.35 (4H, m), 8.24-8.19 (4H, m), 7.99-7.94 (3H, m), 7.65-7.49 (10H, m)
247	8.45 (1H, d), 8.36 (4H, m), 8.23-8.19 (3H, m), 7.99-7.90 (4H, m), 7.65 (2H, m), 7.50-7.49 (8H, m), 7.25 (2H, d)
253	8.55 (1H, d), 8.45 (1H, d), 8.36 (2H, m), 8.24-8.19 (6H, m), 7.99-7.90 (5H, m), 7.68-7.49 (11H, m), 7.35 (1H, m), 7.25-7.16 (4H, m)
262	8.45 (1H, m), 8.35-8.19 (8H, m), 8.19-7.85 (6H, m), 7.65-7.49 (10H, m)
268	8.55 (1H, d), 8.45 (1H, d), 8.35-8.19 (10H, m), 8.19-7.80 (6H, m), 7.68-7.49 (11H, m), 7.35 (1H, m), 7.20-7.16 (2H, m)
283	8.55 (1H, d), 8.45-8.36 (4H, m), 8.24-8.19 (6H, m), 7.99-7.90 (4H, m), 7.73-7.49 (13H, m), 7.35 (1H, m), 7.20-7.16 (2H, m)
298	8.55 (1H, d), 8.45 (1H, d), 8.35 (2H, d), 8.23-8.19 (6H, m), 7.99-7.90 (5H, m), 7.80-7.49 (14H, m), 7.35 (1H, m), 7.20-7.16 (2H, m)
307	9.18-9.14 (4H, m), 8.55 (2H, d), 8.45 (1H, d), 8.23-8.19 (3H, m) 7.99 (1H, d), 7.90 (1H, d), 7.74-7.65 (4H, m), 7.50-7.49 (2H, m), 7.25-7.23 (6H, m)
323	8.76-8.63 (4H, m), 8.39-8.33 (3H, m), 8.19 (2H, d), 7.99 (1H, d), 7.90 (1H, d), 7.65-7.49 (7H, m), 7.29 (2H, d)
325	8.36 (4H, m), 8.20-8.19 (3H, m), 8.07 (1H, s), 7.96-7.94 (3H, m), 7.85 (1H, m), 7.70-7.65 (3H, m), 7.50-7.49 (7H, m), 7.25 (2H, m)

354

TABLE 1-continued

NO	¹ H NMR (CDCl ₃ , 300 Mz)
326	8.35-8.30 (4H, m), 8.23-8.19 (4H, m), 8.07 (1H, s), 7.94-7.85 (4H, m), 7.70-7.49 (10H, m), 7.25 (2H, m)
337	8.69 (2H, d), 8.36 (4H, m), 8.20 (1H, d), 7.99-7.96 (4H, m), 7.85 (3H, m), 7.70 (1H, m), 7.50 (7H, m), 7.25 (2H, d)
338	8.35 (2H, d), 8.23-8.19 (4H, m), 7.99-7.85 (7H, m), 7.70-7.49 (14H, m)
353	8.69 (2H, d), 8.36 (4H, m), 8.20 (1H, d), 8.07 (1H, s), 7.96-7.94 (3H, m), 7.85 (3H, m), 7.70 (1H, m), 7.50-7.41 (7H, m), 7.41-7.19 (6H, m)
354	8.69 (2H, d), 8.35-8.30 (4H, m), 8.23-8.20 (2H, m), 8.07 (1H, s), 7.94-7.85 (8H, m), 7.70 (1H, m), 7.55-7.41 (7H, m)
361	8.69 (2H, d), 8.36 (4H, d), 8.20 (1H, d), 8.07 (1H, s), 7.96-7.94 (3H, m), 7.85 (3H, m), 7.70 (1H, m), 7.50-7.41 (7H, m), 7.25-7.19 (6H, m)
362	8.69 (2H, d), 8.35-8.30 (4H, m), 8.23-8.20 (2H, m), 8.07 (1H, s), 7.94-7.85 (8H, m), 7.70 (1H, m), 7.55-7.41 (7H, m), 7.19 (4H, m)
409	8.36-8.31 (5H, m), 8.19 (2H, d), 8.09-8.07 (2H, m), 7.96-7.94 (3H, m), 7.65 (2H, m), 7.50-7.41 (8H, m), 7.25-7.19 (6H, m)
411	9.18-9.14 (4H, m), 8.55 (2H, d), 8.31 (1H, m), 8.19 (2H, m), 8.09-8.07 (2H, m), 7.94 (1H, s), 7.74-7.65 (4H, m), 7.49-7.41 (2H, m), 7.25-7.19 (10H, m)
412	8.82 (1H, d), 8.71 (2H, d), 8.47 (1H, s), 8.33-8.19 (6H, m), 8.07 (1H, s), 7.90 (1H, d), 7.65-7.41 (7H, m), 7.29 (2H, d), 7.19 (4H, m)
417	8.36-8.31 (5H, m), 8.19 (2H, m), 8.09-8.06 (4H, m), 7.96-7.94 (3H, m), 7.65-7.41 (13H, m), 7.25-7.19 (6H, m)
425	8.45 (1H, d), 8.36 (4H, d), 8.23-8.19 (3H, m), 8.07 (1H, s), 7.96-7.90 (3H, m), 7.65 (2H, m), 7.50-7.41 (8H, m), 7.25-7.19 (6H, m)
426	8.45 (1H, d), 8.35-8.19 (8H, m), 8.07 (1H, s), 7.94-7.85 (5H, m), 7.65-7.41 (10H, m)
449	8.36 (4H, d), 8.19-8.14 (3H, m), 7.99-7.96 (4H, m), 7.65 (3H, m), 7.50-7.49 (8H, m), 7.25 (2H, d)
450	8.35-8.30 (4H, m), 8.23-8.14 (4H, m), 7.99-7.94 (4H, m), 7.65-7.49 (11H, m), 7.25 (2H, d)
473	8.20-8.19 (3H, m), 8.07 (1H, s), 7.94 (1H, d), 7.85 (1H, m), 7.70-7.65 (3H, m), 7.49 (1H, m), 7.25 (4H, s)
485	8.69 (2H, d), 8.20 (1H, d) 7.99-7.94 (2H, m), 7.85 (2H, d), 7.70 (1H, t), 7.50 (1H, d), 7.25 (4H, s)
505	8.45 (1H, d), 8.23-8.19 (3H, m), 7.99 (1H, d), 7.65 (2H, m) 7.50-7.49 (2H, m)
527	8.69 (2H, d), 8.20 (1H, d), 7.99-7.94 (2H, m), 7.85 (3H, m), 7.70 (1H, t), 7.50 (1H, d), 7.25 (4H, s)

TABLE 2

Compound	FD-MS	Compound	FD-MS
1	m/z = 544.14 (C34H20N6S = 544.63)	2	m/z = 620.17 (C40H24N6S = 620.73)
3	m/z = 696.20 (C46H28N6S = 696.83)	4	m/z = 620.17 (C40H24N6S = 620.73)
5	m/z = 696.20 (C46H28N6S = 696.83)	6	m/z = 709.20 (C46H27N7S = 709.83)
7	m/z = 709.20 (C46H27N7S = 709.83)	8	m/z = 572.17 (C36H24N6S = 572.17)
9	m/z = 644.17 (C42H24N6S = 644.17)	10	m/z = 696.20 (C46H28N6S = 696.83)
11	m/z = 710.18 (C46H26N6OS = 710.81)	12	m/z = 726.16 (C46H26N6S2 = 726.87)
13	m/z = 744.18 (C46H29N6OPS = 744.81)	14	m/z = 736.24 (C49H32N6S = 736.89)
15	m/z = 744.20 (C50H28N6S = 744.87)	16	m/z = 543.15 (C35H21N5S = 543.64)
17	m/z = 619.18 (C41H25N5S = 619.74)	18	m/z = 695.21 (C47H29N5SS = 695.84)
19	m/z = 619.18 (C41H25N5S = 619.74)	20	m/z = 695.21 (C47H29N5S = 695.84)
21	m/z = 708.20 (C47H28N6S = 708.84)	22	m/z = 708.20 (C47H28N6S = 708.84)
23	m/z = 571.18 (C37H25N5S = 571.70)	24	m/z = 643.1S (C43H25N5S = 643.76)
25	m/z = 708.20 (C47H28N6S = 708.84)	26	m/z = 709.19 (C47H27N5OS = 709.82)
27	m/z = 725.17 (C47H27N5S2 = 725.88)	28	m/z = 743.19 (C47H30N5OPS = 743.82)
29	m/z = 735.24 (C50H33N5S = 735.90)	30	m/z = 743.21 (C51H29N5S = 743.88)
31	m/z = 620.17 (C40H24N6S = 620.73)	32	m/z = 696.20 (C46H28N6S = 696.83)
33	m/z = 772.24 (C52H32N6S = 772.93)	34	m/z = 696.20 (C46H28N6S = 696.83)
35	m/z = 772.24 (C52H32N6S = 772.93)	36	m/z = 785.23 (C52H31N7S = 785.92)
37	m/z = 785.23 (C52H31N7S = 785.92)	38	m/z = 648.20 (C42H28N6S = 648.78)
39	m/z = 720.20 (C48H28N6S = 720.85)	40	m/z = 772.24 (C52H32N6S = 772.93)
41	m/z = 786.22 (C52H30N6OS = 786.91)	42	m/z = 802.19 (C52H30N6S2 = 802.97)
43	m/z = 820.21 (C52H33N6OPS = 820.91)	44	m/z = 812.27 (C55H36N6S = 812.99)
45	m/z = 820.24 (C56H32N6S = 820.97)	46	m/z = 619.18 (C41H25N5S = 619.74)
47	m/z = 695.21 (C47H29N5S = 695.84)	48	m/z = 771.24 (C53H33N5S = 771.94)
49	m/z = 695.21 (C47H29N5S = 695.84)	50	m/z = 771.24 (C53H33N5S = 771.94)
51	m/z = 784.24 (C53H32N6S = 784.94)	52	m/z = 784.24 (C53H32N6S = 784.94)
53	m/z = 647.21 (C43H29N5S = 647.80)	54	m/z = 719.21 (C49H29N5S = 719.86)

TABLE 2-continued

Compound	FD-MS	Compound	FD-MS
55	m/z = 771.24 (C53H33N5S = 771.94)	56	m/z = 785.22 (C53H31N5OS = 785.92)
57	m/z = 801.20 (C53H31N5S2 = 801.98)	58	m/z = 819.22 (C53H34N5OPS = 819.92)
59	m/z = 811.27 (56H37N5S = 812.00)	60	m/z = 819.24 (C57H33N5S = 819.98)
61	m/z = 620.17 (C40H24N6S = 620.73)	62	m/z = 696.20 (C46H28N6S = 696.83)
63	m/z = 772.24 (C52H32N6S = 772.93)	64	m/z = 696.20 (C46H28N6S = 696.83)
65	m/z = 772.24 (C52H32N6S = 772.93)	66	m/z = 785.23 (C52H31N7S = 785.92)
67	m/z = 785.23 (C52H31N7S = 785.92)	68	m/z = 648.20 (C42H28N6S = 648.78)
69	m/z = 720.20 (C48H28N6S = 720.85)	70	m/z = 772.24 (C52H32N6S = 772.93)
71	m/z = 786.22 (C52H30N6OS = 786.91)	72	m/z = 802.19 (C52H30N6S2 = 802.97)
73	m/z = 820.21 (C52H33N6OPS = 820.91)	74	m/z = 812.27 (C55H36N6S = 812.99)
75	m/z = 820.24 (C56H32N6S = 820.97)	76	m/z = 619.18 (C41H25N5S = 619.74)
77	m/z = 695.21 (C47H29N5S = 695.84)	78	m/z = 771.24 (C53H33N5S = 771.94)
79	m/z = 695.21 (C47H29N5S = 695.84)	80	m/z = 771.24 (C53H33N5S = 771.94)
81	m/z = 784.24 (C53H32N6S = 784.94)	82	m/z = 784.24 (C53H32N6S = 784.94)
83	m/z = 647.21 (C43H29N5S = 647.80)	84	m/z = 719.21 (C49H29N5S = 719.86)
85	m/z = 771.24 (C53H33N5S = 771.94)	86	m/z = 785.22 (C53H31N5OS = 785.92)
87	m/z = 801.20 (C53H31N5S2 = 801.98)	88	m/z = 819.22 (C53H34N5OPS = 819.92)
89	m/z = 811.27 (56H37N5S = 812.00)	90	m/z = 819.24 (C57H33N5S = 819.98)
91	m/z = 620.17 (C40H24N6S = 620.73)	92	m/z = 719.21 (C49H29N5S = 719.86)
93	m/z = 573.10 (C35H19N5S2 = 573.69)	94	m/z = 467.12 (C29H17N5S = 467.55)
95	m/z = 533.16 (C34H23N5S = 533.65)	96	m/z = 619.18 (C41H25N5S = 619.74)
97	m/z = 465.10 (C27H20N3OPS = 465.51)	98	m/z = 517.13 (C33H19N5S = 517.61)
99	m/z = 575.12 (C35H21N5S2 = 575.70)	100	m/z = 517.13 (C33H19N5S = 517.61)
101	m/z = 522.09 (C32H18N4S2 = 522.64)	102	m/z = 506.12 (C32H18N4OS = 506.58)
103	m/z = 581.16 (C38H23N5S = 581.69)	104	m/z = 581.16 (C38H23N5S = 581.69)
105	m/z = 589.13 (C37H24N3OPS = 589.65)	106	m/z = 589.13 (C37H24N3OPS = 589.65)
107	m/z = 567.15 (C37H21N5S = 567.67)	108	m/z = 491.12 (C31H17N5S = 491.57)
109	m/z = 544.14 (C34H20N6S = 544.63)	110	m/z = 620.17 (C40H24N6S = 620.73)
111	m/z = 696.20 (C46H28N6S = 696.83)	112	m/z = 620.17 (C40H24N6S = 620.73)
113	m/z = 696.20 (C46H28N6S = 696.83)	114	m/z = 709.20 (C46H27N7S = 709.83)
115	m/z = 709.20 (C46H27N7S = 709.83)	116	m/z = 572.17 (C36H24N6S = 572.17)
117	m/z = 644.17 (C42H24N6S = 644.17)	118	m/z = 696.20 (C46H28N6S = 696.83)
119	m/z = 710.18 (C46H26N6OS = 710.81)	120	m/z = 726.16 (C46H26N6S2 = 726.87)
121	m/z = 744.18 (C46H29N6OPS = 744.81)	122	m/z = 736.24 (C49H32N6S = 736.89)
123	m/z = 744.20 (C50H28N6S = 744.87)	124	m/z = 543.15 (C35H21N5S = 543.64)
125	m/z = 619.18 (C41H25N5S = 619.74)	126	m/z = 695.21 (C47H29N5S = 695.84)
127	m/z = 619.18 (C41H25N5S = 619.74)	128	m/z = 695.21 (C47H29N5S = 695.84)
129	m/z = 708.20 (C47H28N6S = 708.84)	130	m/z = 708.20 (C47H28N6S = 708.84)
131	m/z = 571.18 (C37H25N5S = 571.70)	132	m/z = 643.18 (C43H25N5S = 643.76)
133	m/z = 708.20 (C47H28N6S = 708.84)	134	m/z = 709.19 (C47H27N5OS = 709.82)
135	m/z = 725.17 (C47H27N5S2 = 725.88)	136	m/z = 743.19 (C47H30N5OPS = 743.82)
137	m/z = 735.24 (C50H33N5S = 735.90)	138	m/z = 743.21 (C51H29N5S = 743.88)
139	m/z = 620.17 (C40H24N6S = 620.73)	140	m/z = 696.20 (C46H28N6S = 696.83)
141	m/z = 772.24 (C52H32N6S = 772.93)	142	m/z = 696.20 (C46H28N6S = 696.83)
143	m/z = 772.24 (C52H32N6S = 772.93)	144	m/z = 785.23 (C52H31N7S = 785.92)
145	m/z = 785.23 (C52H31N7S = 785.92)	146	m/z = 648.20 (C42H28N6S = 648.78)
147	m/z = 720.20 (C48H28N6S = 720.85)	148	m/z = 772.24 (C52H32N6S = 772.93)
149	m/z = 786.22 (C52H30N6OS = 786.91)	150	m/z = 802.19 (C52H30N6S2 = 802.97)
151	m/z = 820.21 (C52H33N6OPS = 820.91)	152	m/z = 812.27 (C55H36N6S = 812.99)
153	m/z = 820.24 (C56H32N6S = 820.97)	154	m/z = 619.18 (C41H25N5S = 619.74)
155	m/z = 695.21 (C47H29N5S = 695.84)	156	m/z = 771.24 (C53H33N5S = 771.94)
157	m/z = 695.21 (C47H29N5S = 695.84)	158	m/z = 771.24 (C53H33N5S = 771.94)
159	m/z = 784.24 (C53H32N6S = 784.94)	160	m/z = 784.24 (C53H32N6S = 784.94)
161	m/z = 647.21 (C43H29N5S = 647.80)	162	m/z = 719.21 (C49H29N5S = 719.86)
163	m/z = 771.24 (C53H33N5S = 771.94)	164	m/z = 785.22 (C53H31N5OS = 785.92)
165	m/z = 801.20 (C53H31N5S2 = 801.98)	166	m/z = 819.22 (C53H34N5OPS = 819.92)
167	m/z = 811.27 (56H37N5S = 812.00)	168	m/z = 819.24 (C57H33N5S = 819.98)
169	m/z = 620.17 (C40H24N6S = 620.73)	170	m/z = 696.20 (C46H28N6S = 696.83)
171	m/z = 772.24 (C52H32N6S = 772.93)	172	m/z = 696.20 (C46H28N6S = 696.83)
173	m/z = 772.24 (C52H32N6S = 772.93)	174	m/z = 785.23 (C52H31N7S = 785.92)
175	m/z = 785.23 (C52H31N7S = 785.92)	176	m/z = 648.20 (C42H28N6S = 648.78)
177	m/z = 720.20 (C48H28N6S = 720.85)	178	m/z = 772.24 (C52H32N6S = 772.93)
179	m/z = 786.22 (C52H30N6OS = 786.91)	180	m/z = 802.19 (C52H30N6S2 = 802.97)
181	m/z = 820.21 (C52H33N6OPS = 820.91)	182	m/z = 812.27 (C55H36N6S = 812.99)
183	m/z = 820.24 (C56H32N6S = 820.97)	184	m/z = 619.18 (C41H25N5S = 619.74)
185	m/z = 695.21 (C47H29N5S = 695.84)	186	m/z = 771.24 (C53H33N5S = 771.94)
187	m/z = 695.21 (C47H29N5S = 695.84)	188	m/z = 771.24 (C53H33N5S = 771.94)
189	m/z = 784.24 (C53H32N6S = 784.94)	190	m/z = 784.24 (C53H32N6S = 784.94)
191	m/z = 647.21 (C43H29N5S = 647.80)	192	m/z = 719.21 (C49H29N5S = 719.86)
193	m/z = 771.24 (C53H33N5S = 771.94)	194	m/z = 785.22 (C53H31N5OS = 785.92)
195	m/z = 801.20 (C53H31N5S2 = 801.98)	196	m/z = 819.22 (C53H34N5OPS = 819.92)
197	m/z = 811.27 (56H37N5S = 812.00)	198	m/z = 819.24 (C57H33N5S = 819.98)
199	m/z = 620.17 (C40H24N6S = 620.73)	200	m/z = 719.21 (C49H29N5S = 719.86)
201	m/z = 573.10 (C35H19N5S2 = 573.69)	202	m/z = 467.12 (C29H17N5S = 467.55)
203	m/z = 533.16 (C34H23N5S = 533.65)	204	m/z = 619.18 (C41H25N5S = 619.74)
205	m/z = 465.10 (C27H20N3OPS = 465.51)	206	m/z = 517.13 (C33H19N5S = 517.61)
207	m/z = 575.12 (C35H21N5S2 = 575.70)	208	m/z = 517.13 (C33H19N5S = 517.61)
209	m/z = 522.09 (C32H18N4S2 = 522.64)	210	m/z = 506.12 (C32H18N4OS = 506.58)

TABLE 2-continued

Compound	FD-MS	Compound	FD-MS
211	m/z = 581.16 (C38H23N5S = 581.69)	212	m/z = 581.16 (C38H23N5S = 581.69)
213	m/z = 589.13 (C37H24N3OPS = 589.65)	214	m/z = 589.13 (C37H24N3OPS = 589.65)
215	m/z = 567.15 (C37H21N5S = 567.67)	216	m/z = 491.12 (C31H17N5S = 491.57)
217	m/z = 544.14 (C34H20N6S = 544.63)	218	m/z = 620.17 (C40H24N6S = 620.73)
219	m/z = 696.20 (C46H28N6S = 696.83)	220	m/z = 620.17 (C40H24N6S = 620.73)
221	m/z = 696.20 (C46H28N6S = 696.83)	222	m/z = 709.20 (C46H27N7S = 709.83)
223	m/z = 709.20 (C46H27N7S = 709.83)	224	m/z = 572.17 (C36H24N6S = 572.17)
225	m/z = 644.17 (C42H24N6S = 644.17)	226	m/z = 696.20 (C46H28N6S = 696.83)
227	m/z = 710.18 (C46H26N6OS = 710.81)	228	m/z = 726.16 (C46H26N6S2 = 726.87)
229	m/z = 744.18 (C46H29N6OPS = 744.81)	230	m/z = 736.24 (C49H32N6S = 736.89)
231	m/z = 744.20 (C50H28N6S = 744.87)	232	m/z = 543.15 (C35H21N5S = 543.64)
233	m/z = 619.18 (C41H25N5S = 619.74)	234	m/z = 695.21 (C47H29N5SS = 695.84)
235	m/z = 619.18 (C41H25N5S = 619.74)	236	m/z = 695.21 (C47H29N5S = 695.84)
237	m/z = 708.20 (C47H28N6S = 708.84)	238	m/z = 708.20 (C47H28N6S = 708.84)
239	m/z = 571.18 (C37H25N5S = 571.70)	240	m/z = 643.18 (C43H25N5S = 643.76)
241	m/z = 708.20 (C47H28N6S = 708.84)	242	m/z = 709.19 (C47H27N5OS = 709.82)
243	m/z = 725.17 (C47H27N5S2 = 725.88)	244	m/z = 743.19 (C47H30N5OPS = 743.82)
245	m/z = 735.24 (C50H33N5S = 735.90)	246	m/z = 743.21 (C51H29N5S = 743.88)
247	m/z = 620.17 (C40H24N6S = 620.73)	248	m/z = 696.20 (C46H28N6S = 696.83)
249	m/z = 772.24 (C52H32N6S = 772.93)	250	m/z = 696.20 (C46H28N6S = 696.83)
251	m/z = 772.24 (C52H32N6S = 772.93)	252	m/z = 785.23 (C52H31N7S = 785.92)
253	m/z = 785.23 (C52H31N7S = 785.92)	254	m/z = 648.20 (C42H28N6S = 648.78)
255	m/z = 720.20 (C48H28N6S = 720.85)	256	m/z = 772.24 (C52H32N6S = 772.93)
257	m/z = 786.22 (C52H30N6OS = 786.91)	258	m/z = 802.19 (C52H30N6S2 = 802.97)
259	m/z = 820.21 (C52H33N6OPS = 820.91)	260	m/z = 812.27 (C55H36N6S = 812.99)
261	m/z = 820.24 (C56H32N6S = 820.97)	262	m/z = 619.18 (C41H25N5S = 619.74)
263	m/z = 695.21 (C47H29N5S = 695.84)	264	m/z = 771.24 (C53H33N5S = 771.94)
265	m/z = 695.21 (C47H29N5S = 695.84)	266	m/z = 771.24 (C53H33N5S = 771.94)
267	m/z = 784.24 (C53H32N6S = 784.94)	268	m/z = 784.24 (C53H32N6S = 784.94)
269	m/z = 647.21 (C43H29N5S = 647.80)	270	m/z = 719.21 (C49H29N5S = 719.86)
271	m/z = 771.24 (C53H33N5S = 771.94)	272	m/z = 785.22 (C53H31N5OS = 785.92)
273	m/z = 801.20 (C53H31N5S2 = 801.98)	274	m/z = 819.22 (C53H34N5OPS = 819.92)
275	m/z = 811.27 (56H37N5S = 812.00)	276	m/z = 819.24 (C57H33N5S = 819.98)
277	m/z = 620.17 (C40H24N6S = 620.73)	278	m/z = 696.20 (C46H28N6S = 696.83)
279	m/z = 772.24 (C52H32N6S = 772.93)	280	m/z = 696.20 (C46H28N6S = 696.83)
281	m/z = 772.24 (C52H32N6S = 772.93)	282	m/z = 785.23 (C52H31N7S = 785.92)
283	m/z = 785.23 (C52H31N7S = 785.92)	284	m/z = 648.20 (C42H28N6S = 648.78)
285	m/z = 720.20 (C48H28N6S = 720.85)	286	m/z = 772.24 (C52H32N6S = 772.93)
287	m/z = 786.22 (C52H30N6OS = 786.91)	288	m/z = 802.19 (C52H30N6S2 = 802.97)
289	m/z = 820.21 (C52H33N6OPS = 820.91)	290	m/z = 812.27 (C55H36N6S = 812.99)
291	m/z = 820.24 (C56H32N6S = 820.97)	292	m/z = 619.18 (C41H25N5S = 619.74)
293	m/z = 695.21 (C47H29N5S = 695.84)	294	m/z = 771.24 (C53H33N5S = 771.94)
295	m/z = 695.21 (C47H29N5S = 695.84)	296	m/z = 771.24 (C53H33N5S = 771.94)
297	m/z = 784.24 (C53H32N6S = 784.94)	298	m/z = 784.24 (C53H32N6S = 784.94)
299	m/z = 647.21 (C43H29N5S = 647.80)	300	m/z = 719.21 (C49H29N5S = 719.86)
301	m/z = 771.24 (C53H33N5S = 771.94)	302	m/z = 785.22 (C53H31N5OS = 785.92)
303	m/z = 801.20 (C53H31N5S2 = 801.98)	304	m/z = 819.22 (C53H34N5OPS = 819.92)
305	m/z = 811.27 (56H37N5S = 812.00)	306	m/z = 819.24 (C57H33N5S = 819.98)
307	m/z = 620.17 (C40H24N6S = 620.73)	308	m/z = 719.21 (C49H29N5S = 719.86)
309	m/z = 573.10 (C35H19N5S2 = 573.69)	310	m/z = 467.12 (C29H17N5S = 467.55)
311	m/z = 533.16 (C34H23N5S = 533.65)	312	m/z = 619.18 (C41H25N5S = 619.74)
313	m/z = 465.10 (C27H20N3OPS = 465.51)	314	m/z = 517.13 (C33H19N5S = 517.61)
315	m/z = 575.12 (C35H21N5S2 = 575.70)	316	m/z = 517.13 (C33H19N5S = 517.61)
317	m/z = 522.09 (C32H18N4S2 = 522.64)	318	m/z = 506.12 (C32H18N4OS = 506.58)
319	m/z = 581.16 (C38H23N5S = 581.69)	320	m/z = 581.16 (C38H23N5S = 581.69)
321	m/z = 589.13 (C37H24N3OPS = 589.65)	322	m/z = 589.13 (C37H24N3OPS = 589.65)
323	m/z = 567.15 (C37H21N5S = 567.67)	324	m/z = 491.12 (C31H17N5S = 491.57)
325	m/z = 604.20 (C40H24N6O = 604.67)	326	m/z = 603.20 (C41H25N5O = 603.68)
327	m/z = 604.20 (C40H24N6O = 604.67)	328	m/z = 551.17 (C37H21N5O = 551.60)
329	m/z = 679.24 (C46H29N7 = 679.78)	330	m/z = 678.25 (C47H30N6 = 678.79)
331	m/z = 679.24 (C46H29N7 = 679.78)	332	m/z = 626.22 (C43H26N6 = 626.72)
333	m/z = 620.17 (C40H24N6S = 620.73)	334	m/z = 619.18 (C41H25N5S = 619.74)
335	m/z = 620.17 (C40H24N6S = 620.73)	336	m/z = 567.15 (C37H21N5S = 567.67)
337	m/z = 604.20 (C40H24N6O = 604.67)	338	m/z = 603.20 (C41H25N5O = 603.68)
339	m/z = 604.20 (C40H24N6O = 604.67)	340	m/z = 551.17 (C37H21N5O = 551.60)
341	m/z = 604.20 (C40H24N6O = 604.67)	342	m/z = 603.20 (C41H25N5O = 603.68)
343	m/z = 604.20 (C40H24N6O = 604.67)	344	m/z = 551.17 (C37H21N5O = 551.60)
345	m/z = 679.24 (C46H29N7 = 679.78)	346	m/z = 678.25 (C47H30N6 = 678.79)
347	m/z = 679.24 (C46H29N7 = 679.78)	348	m/z = 626.22 (C43H26N6 = 626.72)
349	m/z = 679.24 (C46H29N7 = 679.78)	350	m/z = 678.25 (C47H30N6 = 678.79)
351	m/z = 679.24 (C46H29N7 = 679.78)	352	m/z = 626.22 (C43H26N6 = 626.72)
353	m/z = 696.20 (C46H28N6S = 696.83)	354	m/z = 695.21 (C47H29N5S = 695.84)
355	m/z = 696.20 (C46H28N6S = 696.83)	356	m/z = 643.18 (C43H25N5S = 643.76)
357	m/z = 696.20 (C46H28N6S = 696.83)	358	m/z = 695.21 (C47H29N5S = 695.84)
359	m/z = 696.20 (C46H28N6S = 696.83)	360	m/z = 643.18 (C43H25N5S = 643.76)
361	m/z = 680.23 (C46H28N6O = 680.77)	362	m/z = 679.23 (C47H29N5O = 679.78)
363	m/z = 680.23 (C46H28N6O = 680.77)	364	m/z = 627.20 (C43H25N5O = 627.70)
365	m/z = 680.23 (C46H28N6O = 680.77)	366	m/z = 679.23 (C47H29N5O = 679.78)

TABLE 2-continued

Compound	FD-MS	Compound	FD-MS
367	m/z = 680.23 (C46H28N6O = 680.77)	368	m/z = 627.20 (C43H25N5O = 627.70)
369	m/z = 755.27 (C52H33N7 = 755.88)	370	m/z = 754.28 (C53H34N6 = 754.89)
371	m/z = 755.27 (C52H33N7 = 755.88)	372	m/z = 702.25 (C49H30N6 = 702.82)
373	m/z = 755.27 (C52H33N7 = 755.88)	374	m/z = 754.28 (C53H34N6 = 754.89)
375	m/z = 755.27 (C52H33N7 = 755.88)	376	m/z = 702.25 (C49H30N6 = 702.82)
377	m/z = 696.20 (C46H28N6S = 696.83)	378	m/z = 695.21 (C47H29N5S = 695.84)
379	m/z = 696.20 (C46H28N6S = 696.83)	380	m/z = 643.18 (C43H25N5S = 643.76)
381	m/z = 620.17 (C40H24N6S = 620.73)	382	m/z = 619.18 (C41H25N5S = 619.74)
383	m/z = 620.17 (C40H24N6S = 620.73)	384	m/z = 567.15 (C37H21N5S = 567.67)
385	m/z = 680.23 (C46H28N6O = 680.77)	386	m/z = 679.23 (C47H29N5O = 679.78)
387	m/z = 680.23 (C46H28N6O = 680.77)	388	m/z = 627.20 (C43H25N5O = 627.70)
389	m/z = 604.20 (C40H24N6O = 604.67)	390	m/z = 603.20 (C41H25N5O = 603.68)
391	m/z = 604.20 (C40H24N6O = 604.67)	392	m/z = 551.17 (C37H21N5O = 551.60)
393	m/z = 755.27 (C52H33N7 = 755.88)	394	m/z = 754.28 (C53H34N6 = 754.89)
395	m/z = 755.27 (C52H33N7 = 755.88)	396	m/z = 702.25 (C49H30N6 = 702.82)
397	m/z = 679.24 (C46H29N7 = 679.78)	398	m/z = 678.25 (C47H30N6 = 678.79)
399	m/z = 679.24 (C46H29N7 = 679.78)	400	m/z = 626.22 (C43H26N6 = 626.72)
401	m/z = 696.20 (C46H28N6S = 696.83)	402	m/z = 695.21 (C47H29N5S = 695.84)
403	m/z = 696.20 (C46H28N6S = 696.83)	404	m/z = 643.18 (C43H25N5S = 643.76)
405	m/z = 620.17 (C40H24N6S = 620.73)	406	m/z = 619.18 (C41H25N5S = 619.74)
407	m/z = 620.17 (C40H24N6S = 620.73)	408	m/z = 567.15 (C37H21N5S = 567.67)
409	m/z = 680.23 (C46H28N6O = 680.77)	410	m/z = 679.23 (C47H29N5O = 679.78)
411	m/z = 680.23 (C46H28N6O = 680.77)	412	m/z = 627.20 (C43H25N5O = 627.70)
413	m/z = 604.20 (C40H24N6O = 604.67)	414	m/z = 603.20 (C41H25N5O = 603.68)
415	m/z = 604.20 (C40H24N6O = 604.67)	416	m/z = 551.17 (C37H21N5O = 551.60)
417	m/z = 755.27 (C52H33N7 = 755.88)	418	m/z = 754.28 (C53H34N6 = 754.89)
419	m/z = 755.27 (C52H33N7 = 755.88)	420	m/z = 702.25 (C49H30N6 = 702.82)
421	m/z = 679.24 (C46H29N7 = 679.78)	422	m/z = 678.25 (C47H30N6 = 678.79)
423	m/z = 679.24 (C46H29N7 = 679.78)	424	m/z = 626.22 (C43H26N6 = 626.72)
425	m/z = 696.20 (C46H28N6S = 696.83)	426	m/z = 695.21 (C47H29N5S = 695.84)
427	m/z = 696.20 (C46H28N6S = 696.83)	428	m/z = 643.18 (C43H25N5S = 643.76)
429	m/z = 680.23 (C46H28N6O = 680.77)	430	m/z = 679.23 (C47H29N5O = 679.78)
431	m/z = 680.23 (C46H28N6O = 680.77)	432	m/z = 627.20 (C43H25N5O = 627.70)
433	m/z = 604.20 (C40H24N6O = 604.67)	434	m/z = 603.20 (C41H25N5O = 603.68)
435	m/z = 604.20 (C40H24N6O = 604.67)	436	m/z = 551.17 (C37H21N5O = 551.60)
437	m/z = 755.27 (C52H33N7 = 755.88)	438	m/z = 754.28 (C53H34N6 = 754.89)
439	m/z = 755.27 (C52H33N7 = 755.88)	440	m/z = 702.25 (C49H30N6 = 702.82)
441	m/z = 679.24 (C46H29N7 = 679.78)	442	m/z = 678.25 (C47H30N6 = 678.79)
443	m/z = 679.24 (C46H29N7 = 679.78)	444	m/z = 626.22 (C43H26N6 = 626.72)
445	m/z = 696.20 (C46H28N6S = 696.83)	446	m/z = 695.21 (C47H29N5S = 695.84)
447	m/z = 696.20 (C46H28N6S = 696.83)	448	m/z = 643.18 (C43H25N5S = 643.76)
449	m/z = 620.17 (C40H24N6S = 620.73)	450	m/z = 619.18 (C41H25N5S = 619.74)
451	m/z = 620.17 (C40H24N6S = 620.73)	452	m/z = 567.15 (C37H21N5S = 567.67)
453	m/z = 680.23 (C46H28N6O = 680.77)	454	m/z = 679.23 (C47H29N5O = 679.78)
455	m/z = 680.23 (C46H28N6O = 680.77)	456	m/z = 627.20 (C43H25N5O = 627.70)
457	m/z = 604.20 (C40H24N6O = 604.67)	458	m/z = 603.20 (C41H25N5O = 603.68)
459	m/z = 604.20 (C40H24N6O = 604.67)	460	m/z = 551.17 (C37H21N5O = 551.60)
461	m/z = 755.27 (C52H33N7 = 755.88)	462	m/z = 754.28 (C53H34N6 = 754.89)
463	m/z = 755.27 (C52H33N7 = 755.88)	464	m/z = 702.25 (C49H30N6 = 702.82)
465	m/z = 679.24 (C46H29N7 = 679.78)	466	m/z = 678.25 (C47H30N6 = 678.79)
467	m/z = 679.24 (C46H29N7 = 679.78)	468	m/z = 626.22 (C43H26N6 = 626.72)
469	m/z = 562.08 (C32H14F4N4S = 562.54)	470	m/z = 544.09 (C32H15F3N4S = 544.55)
471	m/z = 544.09 (C32H15F3N4S = 544.55)	472	m/z = 526.10 (C32H16F2N4S = 526.56)
473	m/z = 569.09 (C33H14F3N5S = 569.56)	474	m/z = 551.10 (C33H15F2N5S = 551.57)
475	m/z = 544.09 (C32H15F3N4S = 544.55)	476	m/z = 551.10 (C33H15F2N5S = 551.57)
477	m/z = 555.08 (C31H14F5N3S = 555.52)	478	m/z = 562.08 (C32H14F4N4S = 562.54)
479	m/z = 526.10 (C32H16F2N4S = 526.56)	480	m/z = 544.09 (C32H15F3N4S = 544.55)
481	m/z = 540.11 (C34H16N6S = 540.60)	482	m/z = 515.12 (C33H7N5S = 515.59)
483	m/z = 515.12 (C33H7N5S = 515.59)	484	m/z = 490.12 (C32H18N4S = 490.58)
485	m/z = 562.08 (C32H14F4N4S = 562.54)	486	m/z = 544.09 (C32H15F3N4S = 544.55)
487	m/z = 544.09 (C32H15F3N4S = 544.55)	488	m/z = 526.10 (C32H16F2N4S = 526.56)
489	m/z = 569.09 (C33H14F3N5S = 569.56)	490	m/z = 551.10 (C33H15F2N5S = 551.57)
491	m/z = 544.09 (C32H15F3N4S = 544.55)	492	m/z = 551.10 (C33H15F2N5S = 551.57)
493	m/z = 555.08 (C31H14F5N3S = 555.52)	494	m/z = 562.08 (C32H14F4N4S = 562.54)
495	m/z = 526.10 (C32H16F2N4S = 526.56)	496	m/z = 544.09 (C32H15F3N4S = 544.55)
497	m/z = 540.11 (C34H16N6S = 540.60)	498	m/z = 515.12 (C33H7N5S = 515.59)
499	m/z = 515.12 (C33H7N5S = 515.59)	500	m/z = 490.12 (C32H18N4S = 490.58)
501	m/z = 562.08 (C32H14F4N4S = 562.54)	502	m/z = 544.09 (C32H15F3N4S = 544.55)
503	m/z = 544.09 (C32H15F3N4S = 544.55)	504	m/z = 526.10 (C32H16F2N4S = 526.56)
505	m/z = 569.09 (C33H14F3N5S = 569.56)	506	m/z = 551.10 (C33H15F2N5S = 551.57)
507	m/z = 544.09 (C32H15F3N4S = 544.55)	508	m/z = 551.10 (C33H15F2N5S = 551.57)
509	m/z = 555.08 (C31H14F5N3S = 555.52)	510	m/z = 562.08 (C32H14F4N4S = 562.54)
511	m/z = 526.10 (C32H16F2N4S = 526.56)	512	m/z = 544.09 (C32H15F3N4S = 544.55)
513	m/z = 540.11 (C34H16N6S = 540.60)	514	m/z = 515.12 (C33H7N5S = 515.59)
515	m/z = 515.12 (C33H7N5S = 515.59)	516	m/z = 490.12 (C32H18N4S = 490.58)
517	m/z = 524.13 (C34H16N6O = 524.54)	518	m/z = 553.11 (C33H14F3N5O = 553.50)
519	m/z = 546.11 (C32H14F4N4O = 546.48)	520	m/z = 510.12 (C32H16F2N4O = 510.50)
521	m/z = 524.13 (C34H16N6O = 524.54)	522	m/z = 553.11 (C33H14F3N5O = 553.50)

TABLE 2-continued

Compound	FD-MS	Compound	FD-MS
523	m/z = 546.11 (C32H4F4N4O = 546.48)	524	m/z = 510.12 (C32H16F2N4O = 510.50)
525	m/z = 524.13 (C34H16N6O = 524.54)	526	m/z = 553.11 (C33H14F3N5O = 553.50)
527	m/z = 546.11 (C32H4F4N4O = 546.48)	528	m/z = 510.12 (C32H16F2N4O = 510.50)
529	m/z = 599.18 (C40H21N7 = 599.65)	530	m/z = 628.16 (C39H19F3N6 = 628.61)
531	m/z = 621.15 (C38H19F4N5 = 621.59)	532	m/z = 585.17 (C38H21F2N5 = 585.61)
533	m/z = 599.18 (C40H21N7 = 599.65)	534	m/z = 628.16 (C39H19F3N6 = 628.61)
535	m/z = 621.15 (C38H19F4N5 = 621.59)	536	m/z = 585.17 (C38H21F2N5 = 585.61)
537	m/z = 599.18 (C40H21N7 = 599.65)	538	m/z = 628.16 (C39H19F3N6 = 628.61)
539	m/z = 621.15 (C38H19F4N5 = 621.59)	540	m/z = 585.17 (C38H21F2N5 = 585.61)
541	m/z = 565.11 (C35H15N7S = 565.61)	542	m/z = 594.08 (C34H13F3N6S = 594.57)
543	m/z = 587.08 (C33H13F4N5S = 587.55)	544	m/z = 551.10 (C33H15F2N5S = 551.57)
545	m/z = 565.11 (C35H15N7S = 565.61)	546	m/z = 594.08 (C34H13F3N6S = 594.57)
547	m/z = 587.08 (C33H13F4N5S = 587.55)	548	m/z = 551.10 (C33H15F2N5S = 551.57)
549	m/z = 565.11 (C35H15N7S = 565.61)	550	m/z = 594.08 (C34H13F3N6S = 594.57)
551	m/z = 587.08 (C33H13F4N5S = 587.55)	552	m/z = 551.10 (C33H15F2N5S = 551.57)
553	m/z = 565.11 (C35H15N7S = 565.61)	554	m/z = 594.08 (C34H13F3N6S = 594.57)
555	m/z = 587.08 (C33H13F4N5S = 587.55)	556	m/z = 551.10 (C33H15F2N5S = 551.57)
557	m/z = 565.11 (C35H15N7S = 565.61)	558	m/z = 594.08 (C34H13F3N6S = 594.57)
559	m/z = 587.08 (C33H13F4N5S = 587.55)	560	m/z = 551.10 (C33H15F2N5S = 551.57)
561	m/z = 565.11 (C35H15N7S = 565.61)	562	m/z = 594.08 (C34H13F3N6S = 594.57)
563	m/z = 587.08 (C33H13F4N5S = 587.55)	564	m/z = 551.10 (C33H15F2N5S = 551.57)
565	m/z = 565.11 (C35H15N7S = 565.61)	566	m/z = 594.08 (C34H13F3N6S = 594.57)
567	m/z = 587.08 (C33H13F4N5S = 587.55)	568	m/z = 551.10 (C33H15F2N5S = 551.57)
569	m/z = 565.11 (C35H15N7S = 565.61)	570	m/z = 594.08 (C34H13F3N6S = 594.57)
571	m/z = 587.08 (C33H13F4N5S = 587.55)	572	m/z = 551.10 (C33H15F2N5S = 551.57)
573	m/z = 565.11 (C35H15N7S = 565.61)	574	m/z = 594.08 (C34H13F3N6S = 594.57)
575	m/z = 587.08 (C33H13F4N5S = 587.55)	576	m/z = 551.10 (C33H15F2N5S = 551.57)
577	m/z = 565.11 (C35H15N7S = 565.61)	578	m/z = 594.08 (C34H13F3N6S = 594.57)
579	m/z = 587.08 (C33H13F4N5S = 587.55)	580	m/z = 551.10 (C33H15F2N5S = 551.57)
581	m/z = 565.11 (C35H15N7S = 565.61)	582	m/z = 594.08 (C34H13F3N6S = 594.57)
583	m/z = 587.08 (C33H13F4N5S = 587.55)	584	m/z = 551.10 (C33H15F2N5S = 551.57)
585	m/z = 565.11 (C35H15N7S = 565.61)	586	m/z = 594.08 (C34H13F3N6S = 594.57)
587	m/z = 587.08 (C33H13F4N5S = 587.55)	588	m/z = 551.10 (C33H15F2N5S = 551.57)
589	m/z = 565.11 (C35H15N7S = 565.61)	590	m/z = 594.08 (C34H13F3N6S = 594.57)
591	m/z = 587.08 (C33H13F4N5S = 587.55)	592	m/z = 551.10 (C33H15F2N5S = 551.57)
593	m/z = 565.11 (C35H15N7S = 565.61)	594	m/z = 594.08 (C34H13F3N6S = 594.57)
595	m/z = 587.08 (C33H13F4N5S = 587.55)	596	m/z = 551.10 (C33H15F2N5S = 551.57)
597	m/z = 565.11 (C35H15N7S = 565.61)	598	m/z = 594.08 (C34H13F3N6S = 594.57)
599	m/z = 587.08 (C33H13F4N5S = 587.55)	600	m/z = 551.10 (C33H15F2N5S = 551.57)
601	m/z = 565.11 (C35H15N7S = 565.61)	602	m/z = 594.08 (C34H13F3N6S = 594.57)
603	m/z = 587.08 (C33H13F4N5S = 587.55)	604	m/z = 551.10 (C33H15F2N5S = 551.57)
605	m/z = 565.11 (C35H15N7S = 565.61)	606	m/z = 594.08 (C34H13F3N6S = 594.57)
607	m/z = 587.08 (C33H13F4N5S = 587.55)	608	m/z = 551.10 (C33H15F2N5S = 551.57)
609	m/z = 565.11 (C35H15N7S = 565.61)	610	m/z = 594.08 (C34H13F3N6S = 594.57)
611	m/z = 587.08 (C33H13F4N5S = 587.55)	612	m/z = 551.10 (C33H15F2N5S = 551.57)
613	m/z = 549.13 (C35H15N7O = 549.55)	614	m/z = 578.11 (C34H13F3N6O = 578.51)
615	m/z = 571.10 (C33H13F4N5O = 571.49)	616	m/z = 535.12 (C33H15F2N5O = 535.51)
617	m/z = 549.13 (C35H15N7O = 549.55)	618	m/z = 578.11 (C34H13F3N6O = 578.51)
619	m/z = 571.10 (C33H13F4N5O = 571.49)	620	m/z = 535.12 (C33H15F2N5O = 535.51)
621	m/z = 549.13 (C35H15N7O = 549.55)	622	m/z = 578.11 (C34H13F3N6O = 578.51)
623	m/z = 571.10 (C33H13F4N5O = 571.49)	624	m/z = 535.12 (C33H15F2N5O = 535.51)
625	m/z = 549.13 (C35H15N7O = 549.55)	626	m/z = 578.11 (C34H13F3N6O = 578.51)
627	m/z = 571.10 (C33H13F4N5O = 571.49)	628	m/z = 535.12 (C33H15F2N5O = 535.51)
629	m/z = 549.13 (C35H15N7O = 549.55)	630	m/z = 578.11 (C34H13F3N6O = 578.51)
631	m/z = 571.10 (C33H13F4N5O = 571.49)	632	m/z = 535.12 (C33H15F2N5O = 535.51)
633	m/z = 549.13 (C35H15N7O = 549.55)	634	m/z = 578.11 (C34H13F3N6O = 578.51)
635	m/z = 571.10 (C33H13F4N5O = 571.49)	636	m/z = 535.12 (C33H15F2N5O = 535.51)
637	m/z = 624.18 (C41H20N8 = 624.66)	638	m/z = 653.15 (C40H18F3N7 = 353.62)
639	m/z = 646.15 (C39H18F4N6 = 646.60)	640	m/z = 610.17 (C39H20F2N6 = 610.62)
641	m/z = 624.18 (C41H20N8 = 624.66)	642	m/z = 653.15 (C40H18F3N7 = 353.62)
643	m/z = 646.15 (C39H18F4N6 = 646.60)	644	m/z = 610.17 (C39H20F2N6 = 610.62)
645	m/z = 624.18 (C41H20N8 = 624.66)	646	m/z = 653.15 (C40H18F3N7 = 353.62)
647	m/z = 646.15 (C39H18F4N6 = 646.60)	648	m/z = 610.17 (C39H20F2N6 = 610.62)
649	m/z = 624.18 (C41H20N8 = 624.66)	650	m/z = 653.15 (C40H18F3N7 = 353.62)
651	m/z = 646.15 (C39H18F4N6 = 646.60)	652	m/z = 610.17 (C39H20F2N6 = 610.62)
653	m/z = 548.14 (C35H16N8 = 548.56)	654	m/z = 577.12 (C34H14F3N7 = 577.53)
655	m/z = 570.12 (C33H14F4N6 = 570.51)	656	m/z = 534.14 (C33H16F2N6 = 534.52)
657	m/z = 624.18 (C41H20N8 = 624.66)	658	m/z = 653.15 (C40H18F3N7 = 353.62)
659	m/z = 646.15 (C39H18F4N6 = 646.60)	660	m/z = 610.17 (C39H20F2N6 = 610.62)
661	m/z = 619.08 (C35H12F3N7S = 619.58)	662	m/z = 619.08 (C35H12F3N7S = 619.58)
663	m/z = 619.08 (C35H12F3N7S = 619.58)	664	m/z = 619.08 (C35H12F3N7S = 619.58)


363

<Experimental Example 1-1> Manufacture of Organic Light Emitting Device

Comparative Example 1-1

A transparent ITO electrode thin film obtained from glass for an OLED (manufactured by Samsung-Corning Co., Ltd.) was ultrasonic cleaned using trichloroethylene, acetone, ethanol and distilled water consecutively for 5 minutes each, stored in isopropanol, and used.


Next, an ITO substrate was installed in a substrate folder of a vacuum deposition apparatus, and the following 4,4', 4''-tris(N,N-(2-naphthyl)-phenylamino)triphenylamine (2-TNATA) was introduced to a cell in the vacuum deposition apparatus.



2-TNATA

Subsequently, the chamber was evacuated until the degree of vacuum therein reached 10^{-6} torr, and then 2-TNATA was evaporated by applying a current to the cell to deposit a hole injection layer having a thickness of 600 Å on the ITO substrate.

To another cell in the vacuum deposition apparatus, the following N,N'-bis(α-naphthyl)-N,N'-diphenyl-4,4'-diamine (NPB) was introduced, and evaporated by applying a current to the cell to deposit a hole transfer layer having a thickness of 300 Å on the hole injection layer.




NPB


After forming the hole injection layer and the hole transfer layer as above, a blue light emitting material having a

364

structure as below was deposited thereon as a light emitting layer. Specifically, in one side cell in the vacuum deposition apparatus, H1, a blue light emitting host material, was vacuum deposited to a thickness of 200 Å, and D1, a blue light emitting dopant material, was vacuum deposited thereon by 5% with respect to the host material.




H1



D1



Subsequently, a compound of the following Structural Formula E1 was deposited to a thickness of 300 Å on the light emitting layer to form an electron transfer layer. As an electron injection layer, lithium fluoride (LiF) was deposited to a thickness of 10 Å, and an Al cathode was employed to a thickness of 1,000 Å, and as a result, an OLED device was manufactured.




E1

365

-continued



BBQB



TBQB

Meanwhile, all the organic compounds required to manufacture the OLED device were vacuum sublimation purified under 10^{-8} torr to 10^{-6} torr by each material to be used in the OLED manufacture.

Example 1-1 to Example 1-41 and Comparative Examples 1-2 to 1-4

Organic electroluminescent devices were manufactured in the same manner as in Comparative Example 1-1 except that, in Example 1-1 to Example 1-41 and Comparative Examples 1-2 to 1-4, compounds of the following Table 3

366

prepared in the preparation examples were used instead of E1 used when forming the electron transfer layer in Comparative Example 1-1.

<Experimental Example 1-2> Evaluation on Organic Electroluminescent Device

For each of the organic light emitting devices manufactured in Comparative Example 1-1 to Comparative Example 1-4 and Example 1-1 to Example 1-41, electroluminescent (EL) properties were measured using M7000 manufactured by McScience Inc., and with the measurement results, T_{95} was measured when standard luminance was 700 cd/m^2 through a lifetime measurement system (M6000) manufactured by McScience Inc. Results of measuring driving voltage, light emission efficiency, color coordinate (CIE) and lifetime of the blue organic light emitting devices manufactured according to the present disclosure are as shown in Table 3.

TABLE 3

		Com- pound	Driving Voltage (V)	Light Emission Efficiency (cd/A)	CIE (x, y)	Life- time (T ₉₅)
25	Example 1-1	1	5.03	6.68	(0.134, 0.101)	65
	Example 1-2	16	4.92	6.83	(0.134, 0.102)	43
	Example 1-3	31	4.90	6.76	(0.134, 0.101)	42
	Example 1-4	46	4.84	6.98	(0.134, 0.103)	43
30	Example 1-5	61	5.05	6.32	(0.134, 0.102)	37
	Example 1-6	76	4.81	6.97	(0.134, 0.101)	42
	Example 1-7	91	4.79	7.10	(0.134, 0.102)	42
	Example 1-8	107	4.78	7.11	(0.134, 0.101)	42
	Example 1-9	109	4.81	7.01	(0.134, 0.101)	41
	Example 1-10	139	4.90	6.88	(0.134, 0.100)	44
	Example 1-11	144	4.85	6.89	(0.134, 0.101)	54
	Example 1-12	174	4.78	7.05	(0.134, 0.100)	50
35	Example 1-13	199	5.01	6.67	(0.134, 0.100)	41
	Example 1-14	213	4.78	7.17	(0.134, 0.100)	43
	Example 1-15	215	4.85	6.81	(0.134, 0.100)	44
	Example 1-16	217	4.80	6.90	(0.134, 0.100)	44
40	Example 1-17	232	4.81	6.96	(0.134, 0.102)	43
	Example 1-18	247	4.76	7.00	(0.134, 0.101)	39
	Example 1-19	253	4.74	7.05	(0.134, 0.102)	40
	Example 1-20	262	4.77	7.02	(0.134, 0.100)	39
	Example 1-21	268	4.85	6.83	(0.134, 0.103)	43
	Example 1-22	283	4.85	6.86	(0.134, 0.100)	55
	Example 1-23	298	4.79	6.88	(0.134, 0.102)	41
	Example 1-24	307	4.79	6.99	(0.134, 0.101)	56
45	Example 1-25	323	4.85	6.89	(0.134, 0.100)	37
	Example 1-26	325	4.82	6.90	(0.134, 0.102)	40
	Example 1-27	326	4.83	6.98	(0.134, 0.103)	44
	Example 1-28	337	4.83	6.91	(0.134, 0.100)	39
50	Example 1-29	338	4.77	7.00	(0.134, 0.103)	54
	Example 1-30	353	4.98	6.62	(0.134, 0.102)	43
	Example 1-31	354	4.92	6.71	(0.134, 0.100)	45
	Example 1-32	361	4.99	6.71	(0.134, 0.099)	44
	Example 1-33	362	4.85	6.88	(0.134, 0.102)	45
	Example 1-34	409	4.85	6.91	(0.134, 0.100)	46
	Example 1-35	411	4.85	6.91	(0.134, 0.103)	43
	Example 1-36	412	4.92	6.65	(0.134, 0.102)	45
55	Example 1-37	417	4.79	6.72	(0.134, 0.101)	40
	Example 1-38	425	4.85	6.62	(0.134, 0.102)	43
	Example 1-39	426	4.70	6.92	(0.134, 0.102)	45
	Example 1-40	449	4.98	6.70	(0.134, 0.099)	43
	Example 1-41	450	4.96	6.71	(0.134, 0.100)	42
	Comparative Example 1-1	E1	5.56	5.91	(0.134, 0.100)	28
	Comparative Example 1-2	E2	5.52	6.09	(0.134, 0.101)	28
	Comparative Example 1-3	BBQB	5.50	6.10	(0.134, 0.101)	30
65	Comparative Example 1-4	TBQB	5.51	6.15	(0.134, 0.102)	29

367


As seen from the results of Table 3, the organic light emitting device using the electron transfer layer material of the blue organic light emitting device of the present disclosure had lower driving voltage and significantly improved light emission efficiency and lifetime compared to the comparative examples. Particularly, it was identified that Compounds 91, 107, 213, 253, 262 and 338 were superior in all aspects of driving, efficiency and lifetime.

Such a result is considered to be due to the fact that, when using the disclosed compound having proper length and strength, and flatness as an electron transfer layer, a compound in an excited state is made by receiving electrons under a specific condition, and particularly when an excited state is formed in the hetero-skeleton site of the compound, excited energy moves to a stable state before the excited hetero-skeleton site goes through other reactions, and as a result, the relatively stabilized compound is capable of efficiently transferring electrons without the compound being decomposed or destroyed. For reference, those that are stable when excited are considered to be aryl or acene-based compounds or polycyclic hetero-compounds. Accordingly, it is considered that excellent results in all aspects of driving, efficiency and lifetime were obtained by the compound of the present disclosure enhancing enhanced electron-transfer properties or improved stability.

<Experimental Example 2-1> Manufacture of Organic Light Emitting Device

Comparative Example 2-1

A transparent ITO electrode thin film obtained from glass for an OLED (manufactured by Samsung-Corning Co., Ltd.) was ultrasonic cleaned using trichloroethylene, acetone, ethanol and distilled water consecutively for 5 minutes each, stored in isopropanol, and used. Next, an ITO substrate was installed in a substrate folder of a vacuum deposition apparatus, and the following 4,4',4''-tris(N,N-(2-naphthyl)-phenylamino)triphenylamine (2-TNATA) was introduced to a cell in the vacuum deposition apparatus.




2-TNATA

Subsequently, the chamber was evacuated until the degree of vacuum therein reached 10^{-6} torr, and then 2-TNATA was


368

evaporated by applying a current to the cell to deposit a hole injection layer having a thickness of 600 Å on the ITO substrate. To another cell in the vacuum deposition apparatus, the following N,N'-bis(α-naphthyl)-N, N'-diphenyl-4, 4'-diamine (NPB) was introduced, and evaporated by applying a current to the cell to deposit a hole transfer layer having a thickness of 300 Å on the hole injection layer.




NPB

After forming the hole injection layer and the hole transfer layer as above, a blue light emitting material having a structure as below was deposited thereon as a light emitting layer. Specifically, in one side cell in the vacuum deposition apparatus, H1, a blue light emitting host material, was vacuum deposited to a thickness of 200 Å, and D1, a blue light emitting dopant material, was vacuum deposited thereon by 5% with respect to the host material.



H1




D1

Subsequently, as a hole blocking layer, a compound of the following Structural Formula E2 was deposited to a thickness of 50 Å.


Subsequently, as an electron transfer layer, a compound of the following Structural Formula E1 was deposited to a thickness of 300 Å.

369



370

-continued



E1

5

10


15

20

TBQB

E2

30



35

40

Example 2-1 to Example 2-41, Comparative
Example 2-2 and Comparative Example 2-3


Organic electroluminescent devices were manufactured in the same manner as in Comparative Example 2-1 except that, in Example 2-1 to Example 2-41, Comparative Example 2-2 and Comparative Example 2-3, compounds of the following Table 4 prepared in the preparation examples were used instead of E2 used when forming the hole blocking layer in Comparative Example 2-1.

<Experimental Example 2-2> Evaluation on
Organic Electroluminescent Device

For each of the organic light emitting devices manufactured in Comparative Example 2-1 to Comparative Example 2-3 and Example 2-1 to Example 2-41, electroluminescent (EL) properties were measured using M7000 manufactured by McScience Inc., and with the measurement results, T_{95} was measured when standard luminance was 700 cd/m² through a lifetime measurement system (M6000) manufactured by McScience Inc. Results of measuring driving voltage, light emission efficiency, color coordinate (CIE) and lifetime of the blue organic light emitting devices manufactured according to the present disclosure are as shown in Table 4.

TABLE 4

60		Com- pound	Driving Voltage (V)	Light	CIE (x, y)	Life- time (T ₉₅)
				Emission Efficiency (cd/A)		
65	Example 2-1	1	4.96	6.77	(0.134, 0.101)	56
	Example 2-2	16	5.07	6.83	(0.134, 0.102)	55
	Example 2-3	31	5.05	6.81	(0.134, 0.101)	70
	Example 2-4	46	4.98	6.75	(0.134, 0.103)	55



BBQB

371

TABLE 4-continued

	Com- pound	Driving Voltage (V)	Light Emission Efficiency (cd/A)	CIE (x, y)	Life- time (T ₉₅)
Example 2-5	61	4.95	6.95	(0.134, 0.102)	51
Example 2-6	76	4.89	6.94	(0.134, 0.101)	52
Example 2-7	91	4.93	6.88	(0.134, 0.102)	58
Example 2-8	107	5.05	6.65	(0.134, 0.101)	57
Example 2-9	109	5.04	6.75	(0.134, 0.101)	58
Example 2-10	139	5.10	6.58	(0.134, 0.100)	73
Example 2-11	144	4.85	7.14	(0.134, 0.101)	49
Example 2-12	174	5.08	7.02	(0.134, 0.100)	43
Example 2-13	199	4.90	6.71	(0.134, 0.100)	48
Example 2-14	213	4.93	6.71	(0.134, 0.100)	45
Example 2-15	215	5.01	6.95	(0.134, 0.100)	53
Example 2-16	217	4.86	6.86	(0.134, 0.100)	56
Example 2-17	232	4.85	6.89	(0.134, 0.102)	46
Example 2-18	247	5.04	7.06	(0.134, 0.101)	50
Example 2-19	253	5.10	6.97	(0.134, 0.102)	51
Example 2-20	262	4.84	7.14	(0.134, 0.100)	44
Example 2-21	268	4.88	7.02	(0.134, 0.103)	42
Example 2-22	283	5.06	6.65	(0.134, 0.100)	51
Example 2-23	298	5.05	6.78	(0.134, 0.102)	41
Example 2-24	307	5.05	6.77	(0.134, 0.101)	48
Example 2-25	323	4.90	6.95	(0.134, 0.100)	58
Example 2-26	325	5.11	6.46	(0.134, 0.102)	73
Example 2-27	326	4.78	7.20	(0.134, 0.103)	57
Example 2-28	337	5.07	6.53	(0.134, 0.100)	71
Example 2-29	338	4.85	7.19	(0.134, 0.103)	41
Example 2-30	353	4.91	6.65	(0.134, 0.102)	47
Example 2-31	354	5.05	6.71	(0.134, 0.100)	55
Example 2-32	361	5.00	6.77	(0.134, 0.099)	58
Example 2-33	362	4.96	6.87	(0.134, 0.102)	53
Example 2-34	409	4.99	6.74	(0.134, 0.100)	46
Example 2-35	411	4.96	6.80	(0.134, 0.103)	43
Example 2-36	412	5.03	6.69	(0.134, 0.102)	50
Example 2-37	417	4.84	7.13	(0.134, 0.101)	55
Example 2-38	425	4.80	7.14	(0.134, 0.102)	52
Example 2-39	426	4.82	7.14	(0.134, 0.102)	50
Example 2-40	449	4.79	7.15	(0.134, 0.099)	48
Example 2-41	450	5.04	6.85	(0.134, 0.100)	49
Comparative Example 2-1	E2	5.52	6.09	(0.134, 0.101)	28
Comparative Example 2-2	BBQB	5.50	6.10	(0.134, 0.101)	30
Comparative Example 2-3	TBQB	5.51	6.15	(0.134, 0.102)	29

As seen from the results of Table 4, the organic light emitting device using the hole blocking layer material of the blue organic light emitting device of the present disclosure had lower driving voltage and significantly improved light emission efficiency and lifetime compared to the comparative examples. Particularly, it was identified that Compounds 144, 262, 268, 326, 338, 417, 425, 426 and 449 were significantly superior in all aspects of driving, efficiency and lifetime.

Such results are due to the fact that the compound is a bipolar type having both a p-type and an n-type, and therefore, hole leakage is prevented and excitons are effectively trapped in the light emitting layer.

<Experimental Example 3-1> Manufacture of Organic Light Emitting Device

Comparative Example 3-1

A glass substrate on which indium tin oxide (ITO) was coated as a thin film to a thickness of 1,500 Å was cleaned with distilled water ultrasonic waves. After the cleaning with distilled water was finished, the substrate was ultrasonic


372

cleaned with solvents such as acetone, methanol and isopropyl alcohol, then dried, and UVO treatment was conducted for 5 minutes using UV in a UV cleaner. After that, the substrate was transferred to a plasma cleaner (PT), and after conducting plasma treatment under vacuum for ITO work function and residual film removal, the substrate was transferred to a thermal deposition apparatus for organic deposition.

On the transparent ITO electrode (anode), organic materials were formed in a 2-stack white organic light emitting device (WOLED) structure. As for the first stack, the following TAPC was thermal vacuum deposited first to a thickness of 300 Å to form a hole transfer layer. After forming the hole transfer layer, a light emitting layer was thermal vacuum deposited thereon as follows. As the light emitting layer, the following TCz1, a host, was 8% doped with the following Flrpic, a blue phosphorescent dopant, and deposited to 300 Å. After forming an electron transfer layer to 400 Å using the following TmPyPB, the following BBQB compound was 20% doped with Cs₂CO₃ to form as a charge generation layer to 100 Å.

As for the second stack, MoO₃ was thermal vacuum deposited first to a thickness of 50 Å to form a hole injection layer. A hole transfer layer, a common layer, was formed to 100 Å by 20% doping MoO₃ to the following TAPC, and then depositing the following TAPC to 300 Å. A light emitting layer was formed thereon by 8% doping the following Ir(ppy)₃, a green phosphorescent dopant, to TCz1, a host, and depositing the result to 300 Å, and then an electron transfer layer was formed to 600 Å using the following TmPyPB. Lastly, an electron injection layer was formed on the electron transfer layer by depositing lithium fluoride (LiF) to a thickness of 10 Å, and then a cathode was formed on the electron injection layer by depositing an aluminum (Al) cathode to a thickness of 1,200 Å, and as a result, an organic light emitting device was manufactured.


Meanwhile, all the organic compounds required to manufacture the organic light emitting diode (OLED) device were vacuum sublimation purified under 10⁻⁸ torr to 10⁻⁶ torr for each material to be used in the OLED manufacture.




TAPC

373



-continued



TCz1




FIrpic

Ir(ppy)₃


TmPyPB

374

-continued




BBQB



TBQB

E2



Example 3-1 to Example 3-41, Comparative
Example 3-2 and Comparative Example 3-3

Organic electroluminescent devices were manufactured in the same manner as in Comparative Example 3-1 except that, in Example 3-1 to Example 3-41, Comparative Example 3-2 and Comparative Example 3-3, compounds of the following Table 5 were used instead of BBQB used when forming the charge generation layer in Comparative Example 3-1.

<Experimental Example 3-2> Evaluation an
Organic Electroluminescent Device

For each of the organic light emitting devices manufactured as above, electroluminescent (EL) properties were measured using M7000 manufactured by McScience Inc., and with the measurement results, T_{95} was measured when standard luminance was 3,500 cd/m² through a lifetime measurement system (M6000) manufactured by McScience Inc. Results of measuring driving voltage, light emission efficiency, color coordinate (CIE) and lifetime of the white organic light emitting devices manufactured according to the present disclosure are as shown in Table 5.

TABLE 5

	Com- pound	Driving Voltage (V)	Light Emission Efficiency (cd/A)	CIE (x, y)	Life- time (T_{95})
Example 3-1	1	7.78	63.95	(0.218, 0.427)	61
Example 3-2	16	7.60	65.45	(0.220, 0.431)	40
Example 3-3	31	7.57	64.75	(0.220, 0.431)	39
Example 3-4	46	7.48	66.88	(0.200, 0.421)	40
Example 3-5	61	7.81	60.55	(0.228, 0.436)	35
Example 3-6	76	7.44	66.74	(0.243, 0.442)	39
Example 3-7	91	7.41	67.99	(0.221, 0.433)	39
Example 3-8	107	7.39	68.12	(0.208, 0.415)	39
Example 3-9	109	7.44	67.13	(0.233, 0.433)	38
Example 3-10	139	7.57	65.89	(0.238, 0.438)	41
Example 3-11	144	7.50	66.01	(0.225, 0.429)	50
Example 3-12	174	7.39	67.56	(0.209, 0.415)	47
Example 3-13	199	7.74	63.89	(0.231, 0.440)	38
Example 3-14	213	7.39	68.71	(0.211, 0.419)	40
Example 3-15	215	7.49	65.26	(0.209, 0.419)	41
Example 3-16	217	7.42	66.11	(0.207, 0.409)	41
Example 3-17	232	7.43	66.66	(0.208, 0.415)	40
Example 3-18	247	7.35	67.06	(0.214, 0.420)	36
Example 3-19	253	7.33	67.56	(0.224, 0.429)	37
Example 3-20	262	7.37	67.21	(0.221, 0.434)	36
Example 3-21	268	7.50	65.45	(0.212, 0.422)	40
Example 3-22	283	7.49	65.70	(0.228, 0.418)	51
Example 3-23	298	7.40	65.89	(0.231, 0.420)	38
Example 3-24	307	7.41	66.98	(0.219, 0.411)	52
Example 3-25	323	7.50	65.98	(0.219, 0.411)	35
Example 3-26	325	7.45	66.11	(0.210, 0.412)	37
Example 3-27	326	7.47	66.84	(0.218, 0.421)	41
Example 3-28	337	7.46	66.21	(0.209, 0.432)	36
Example 3-29	338	7.38	67.04	(0.231, 0.418)	50
Example 3-30	353	7.69	63.38	(0.243, 0.442)	40
Example 3-31	354	7.61	64.23	(0.205, 0.411)	42
Example 3-32	361	7.72	64.22	(0.243, 0.442)	41
Example 3-33	362	7.50	65.88	(0.209, 0.419)	42
Example 3-34	409	7.49	66.18	(0.210, 0.420)	43
Example 3-35	411	7.50	66.20	(0.231, 0.419)	40
Example 3-36	412	7.49	66.24	(0.229, 0.420)	40
Example 3-37	417	7.31	66.80	(0.224, 0.423)	35
Example 3-38	425	7.45	66.27	(0.220, 0.424)	37
Example 3-39	426	7.29	67.57	(0.221, 0.430)	40
Example 3-40	449	7.56	65.46	(0.205, 0.411)	39
Example 3-41	450	7.54	65.64	(0.231, 0.418)	39
Comparative Example 3-1	BBQB	8.43	58.11	(0.220, 0.429)	27
Comparative Example 3-2	TBQB	8.47	58.90	(0.222, 0.430)	28
Comparative Example 3-3	E2	8.45	58.05	(0.221, 0.431)	25

As seen from the results of Table 5, the organic light emitting device using the charge generation layer material of the 2-stack white organic light emitting device of the present disclosure had lower driving voltage and improved light emission efficiency compared to the comparative examples. Particularly, it was identified that Compounds 91, 107, 213, 253, 262 and 338 were significantly superior in all aspects of driving, efficiency and lifetime.

Such a result is considered to be due to the fact that the compound of the present disclosure used as an N-type charge generation layer formed with the disclosed skeleton having proper length and strength, and flatness and a proper hetero-compound capable of binding to metals forms a gap state in the N-type charge generation layer by doping an alkali metal or an alkaline earth metal thereto, and electrons produced from a P-type charge generation layer are readily injected into an electron transfer layer through the gap state produced in the N-type charge generation layer. Accordingly, the P-type charge generation layer may favorably inject and transfer electrons to the N-type charge generation layer, and as a result, driving voltage was lowered, and efficiency and lifetime were improved in the organic light emitting device.

In addition, by the core structure of the present application having a substituent at two places and thereby combining an electron-deficient substituent and an aryl or acene-based substituent, the electron-deficient substituent readily receives electrons from the electron injection layer, and the aryl or acene-based substituent stabilizes the molecule itself and transfers the supplied electrons to the light emitting layer, which may enhance device properties. Accordingly, higher molecular stability and device properties may be obtained compared to the materials in which phenanthridine is single substituted as in the comparative examples.




<Experimental Example 4-1> Manufacture of
Organic Light Emitting Device

Comparative Example 4-1

A glass substrate on which ITO was coated as a thin film to a thickness of 1,500 Å was cleaned with distilled water ultrasonic waves. After the cleaning with distilled water was finished, the substrate was ultrasonic cleaned with solvents such as acetone, methanol and isopropyl alcohol, then dried, and UVO treatment was conducted for 5 minutes using UV in a UV cleaner. After that, the substrate was transferred to a plasma cleaner (PT), and after conducting plasma treatment under vacuum for ITO work function and residual film removal, the substrate was transferred to a thermal deposition apparatus for organic deposition. On the transparent ITO electrode (anode), organic materials were formed in a 2-stack white organic light emitting device (WOLED) structure. As for the first stack, the following TAPC was thermal vacuum deposited first to a thickness of 300 Å to form a hole transfer layer. After forming the hole transfer layer, a light emitting layer was thermal vacuum deposited thereon as follows. As the light emitting layer, the following TCz1, a host, was 8% doped with the following Flrpic, a blue phosphorescent dopant, and deposited to 300 Å. After forming an electron transfer layer to 400 Å using the following TmPyPB, the following Bphen was 20% doped with Cs₂CO₃ to form as a charge generation layer to 100 Å. As for the second stack, MoO₃ was thermal vacuum deposited first to a thickness of 50 Å to form a hole injection layer. A hole transfer layer, a common layer, was formed to 100 Å by 20% doping MoO₃ to the following TAPC, and then depositing the following TAPC to 300 Å. A light emitting layer was formed thereon by 8% doping the following Ir(ppy)₃, a green phosphorescent dopant, to the following TCz1, a host, and depositing the result to 300 Å, and then an electron transfer layer was formed to 600 Å using the following TmPyPB. Lastly, an electron injection layer was formed on the electron transfer layer by depositing lithium fluoride (LiF) to a thickness of 10 Å, and then a cathode was formed


377

on the electron injection layer by depositing an aluminum (Al) cathode to a thickness of 1,200 Å, and as a result, an organic electroluminescent device was manufactured. Meanwhile, all the organic compounds required to manufacture the OLED device were vacuum sublimation purified under 10^{-8} torr to 10^{-6} torr for each material to be used in the OLED manufacture.





378

-continued



TmPyPB



Example 4-1 to Example 4-4

Organic electroluminescent devices were manufactured in the same manner as in Comparative Example 4-1 except that, in Example 4-1 to Example 4-4, compounds presented in Table 6 were used instead of MoO₃ used when forming the hole injection layer in Comparative Example 4-1.

<Experimental Example 4-2> Evaluation on Organic Electroluminescent Device

For each of the organic electroluminescent devices manufactured as above, electroluminescent (EL) properties were measured using M7000 manufactured by McScience Inc., and with the measurement results, T₉₅ was measured when standard luminance was 3,500 cd/m² through a lifetime measurement system (M6000) manufactured by McScience Inc. Results of measuring driving voltage, light emission efficiency, color coordinate (CIE) and lifetime of the white organic electroluminescent device manufactured according to the present disclosure are as shown in Table 6.

379

TABLE 6


	Com- pound	Driving Voltage (V)	Light Emission Efficiency (cd/A)	CIE (x, y)	Life- time (T ₉₅)
Example 4-1	473	5.12	6.61	(0.134, 0.100)	52
Example 4-2	485	5.14	6.65	(0.134, 0.101)	53
Example 4-3	505	5.17	6.72	(0.134, 0.100)	55
Example 4-4	527	5.24	6.71	(0.134, 0.101)	55
Comparative Example 4-1	MoO ₃	5.45	6.06	(0.134, 0.101)	44

As seen from the results of Table 6, the organic electroluminescent device using the hole injection layer material of the 2-stack white organic electroluminescent device of the present disclosure had lower driving voltage and improved light emission efficiency compared to Comparative Example 4-1.

Such a result is considered to be due to the fact that holes may be smoothly injected by being formed with materials having an energy level similar to the energy level of the hole transfer layer receiving holes from a P-type charge generation layer, and, by having a low LUMO energy level, electrons produced from the P-type charge generation layer, which are in a stable state due to anionization, are readily injected through a gap state produced in an N-type charge generation layer. Accordingly, the P-type charge generation layer may favorably inject and transfer electrons to the N-type charge generation layer, and as a result, driving voltage was lowered, and efficiency and lifetime were improved in the organic light emitting device.

The invention claimed is:

1. A heterocyclic compound represented by the following Chemical Formula 1:



wherein, in Chemical Formula 1,

X is O; S; or NR;

L1 to L3 are the same as or different from each other, and each independently a direct bond; a substituted or unsubstituted arylene group having 6 to 60 carbon atoms; or a substituted or unsubstituted heteroarylene group having 2 to 60 carbon atoms;

Z1 to Z3 are the same as or different from each other, and each independently selected from the group consisting of hydrogen; deuterium; a halogen group; a cyano group; —P(=O)(R104)(R105); —N(R106)(R107); a substituted or unsubstituted alkyl group having 1 to 60 carbon atoms; a substituted or unsubstituted alkenyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkynyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkoxy group having 1 to 60 carbon atoms; a substituted or unsubstituted cycloalkyl group having 3 to 60 carbon atoms;

380

a substituted or unsubstituted heterocycloalkyl group having 2 to 60 carbon atoms; a substituted or unsubstituted aryl group having 6 to 60 carbon atoms; and a substituted or unsubstituted heteroaryl group having 2 to 60 carbon atoms;

R1 and R2 are the same as or different from each other, and each independently selected from the group consisting of hydrogen; deuterium; a halogen group; a cyano group; —P(=O)(R104')(R105'); —N(R106')(R107'); a substituted or unsubstituted alkyl group having 1 to 60 carbon atoms; a substituted or unsubstituted alkenyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkynyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkoxy group having 1 to 60 carbon atoms; a substituted or unsubstituted cycloalkyl group having 3 to 60 carbon atoms; a substituted or unsubstituted heterocycloalkyl group having 2 to 60 carbon atoms; a substituted or unsubstituted aryl group having 6 to 60 carbon atoms; and a substituted or unsubstituted heteroaryl group having 2 to 60 carbon atoms, or adjacent two or more groups bond to each other to form a substituted or unsubstituted aliphatic hydrocarbon ring; a substituted or unsubstituted aromatic hydrocarbon ring; a substituted or unsubstituted aliphatic heteroring; or a substituted or unsubstituted aromatic heteroring;

R is hydrogen; deuterium; a halogen group; a cyano group; a substituted or unsubstituted alkyl group having 1 to 60 carbon atoms; a substituted or unsubstituted alkenyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkynyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkoxy group having 1 to 60 carbon atoms; a substituted or unsubstituted cycloalkyl group having 3 to 60 carbon atoms; a substituted or unsubstituted heterocycloalkyl group having 2 to 60 carbon atoms; a substituted or unsubstituted aryl group having 6 to 60 carbon atoms; or a substituted or unsubstituted heteroaryl group having 2 to 60 carbon atoms;


R104 to R107 and R104' to R107' are each independently hydrogen; deuterium; a halogen group; an alkyl group; an alkenyl group; an alkoxy group; a cycloalkyl group; an aryl group; or a heteroaryl group;

m, p, x, n, q and y are each an integer of 1 to 5;

a is an integer of 1 to 3; and

when m, p, x, n, q, y and a are each an integer of 2 or greater, substituents in the parentheses are the same as or different from each other.


2. The heterocyclic compound of claim 1, wherein Chemical Formula 1 is represented by one of the following Chemical Formulae 2 to 5:




381

-continued


[Chemical Formula 3]



[Chemical Formula 4]



[Chemical Formula 5]



in Chemical Formula 2 to Chemical Formula 5,
X, m, p, x, n, q, y, a and R have the same definitions as
in Chemical Formula 1,

L11 to L13 are the same as or different from each other,
and each independently a direct bond; a substituted or
unsubstituted arylene group having 6 to 60 carbon
atoms; or a substituted or unsubstituted heteroarylene
group having 2 to 60 carbon atoms,

Z11 to Z13 are the same as or different from each other,
and each independently selected from the group consisting of
a cyano group; $\text{—P(=O)(R104)(R105)}$; a substituted or unsubstituted aryl group having 6 to 60
carbon atoms; and a substituted or unsubstituted heteroaryl group having 2 to 60 carbon atoms,

R11 and R12 are the same as or different from each other,
and each independently selected from the group consisting of
hydrogen; deuterium; a halogen group; a cyano group; $\text{—P(=O)(R104')(R105')}$; —N(R106')(R107') ; a substituted or unsubstituted alkyl group
having 1 to 60 carbon atoms; a substituted or unsubstituted alkenyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkynyl group having 2 to 60 carbon atoms; a substituted or unsubstituted alkoxy
group having 1 to 60 carbon atoms; a substituted or unsubstituted cycloalkyl group having 3 to 60 carbon atoms; and a substituted or unsubstituted heterocycloalkyl group having 2 to 60 carbon atoms,

382

R104, R105 and R104' to R107' are each independently
hydrogen; deuterium; a halogen group; an alkyl group;
an alkenyl group; an alkoxy group; a cycloalkyl group;
an aryl group; or a heteroaryl group,

b is 1 or 2,

c is an integer of 1 to 4,

when b is 2, two R11s are the same as or different from
each other, and

when c is 2 or greater, a plurality of R12s are the same as
or different from each other.



3. The heterocyclic compound of claim 1, wherein R1 and
R2 are hydrogen.

4. The heterocyclic compound of claim 1, wherein L1 to
L3 are the same as or different from each other, and each
independently a direct bond; a phenylene group unsubstituted
or substituted with a halogen group; a biphenylene
group; or a terphenylene group.

5. The heterocyclic compound of claim 1, wherein R is a
substituted or unsubstituted aryl group having 6 to 60 carbon
atoms.




6. The heterocyclic compound of claim 1, wherein Z1 to
Z3 are the same as or different from each other, and each
independently hydrogen; deuterium; a cyano group;
 $\text{—P(=O)(R104)(R105)}$; a substituted or unsubstituted aryl
group having 6 to 40 carbon atoms; or a substituted or
unsubstituted heteroaryl group having 2 to 40 carbon atoms.

7. The heterocyclic compound of claim 1, wherein
Chemical Formula 1 is represented by any one of the
following compounds:



383

-continued

**384**

-continued

3

6

5

10

15

20

4

35

40

45

5

55





60

65

7


8

9



385

-continued



10

5


10

15

20

11

25




30

35

40

45

12



50


55

60

65


386

-continued




13

14






15



387

-continued

**388**

-continued

16

19

5

10

15

20

17

20

25

30

35

40

45

21




18

50

55




60

65



389

-continued

**390**

-continued

22

25

5

10

15

20

25

23

30

35

40

24




45

50




55

60

65






-continued





-continued

31



393

-continued

**394**

-continued

34

5

10

15

20

25

30

35

35 40

45



50

55

60

65


36



37

395


-continued



38

396

-continued



40

5

10

15

20


25

30

35

40

39




45

50

55

60



65



41

397

-continued

**398**

-continued

42

44

5

10

15

20

25

45

30

35

40

43

45


50

46

55

60

65



40

43



45

50

55



60

65



399

-continued

**400**

-continued

47

5

10

15

20

25

30

35

48

40



45

50

55

60

65





49

50

401


-continued

**402**

-continued

51

53



20

25

30

35

52

40

54


45

50

55



60

65



403

-continued

**404**

-continued

55

57

5

10

15

20

25

30

35

56 40

58



45

50

55



60

65






405

-continued




**406**

-continued






407

-continued






408

-continued






409

-continued




**410**

-continued






411

-continued




**412**

-continued



413

-continued

**414**

-continued

82

85

5

10

15

20

25

83

30

35

40

84 45

50




55

60




65

86

87







-continued








-continued

91







417

-continued


**418**

-continued



419

-continued



104


5

10

15


420

-continued



108

105 20



25

30


106

35

40

45

107 50




55

60

65


109

110



421


-continued



111

422

-continued



113


25

30

35

40

112




45

50

55

60


65



114

423

-continued

**424**

-continued

115

5

10

15

20

25

30

35

40

116

45


50

55



60

65


117




118



-continued




120



-continued

121



25


30

35

40




120

122





427

-continued



**428**

-continued



429

-continued

**430**

-continued

128

5

10

15

20

25

30

35

40

129


45

50

55


60

65





130

131



431

-continued

**432**

-continued

132

134

5

10

15

20

25

30

35

40

133

135



45

50

55



60

65





433


-continued

**434**


-continued



-continued



-continued



30

35

40

141

45

50



55

60

65




142

143







437

-continued

**438**

-continued





150

5

10

15

20

25

30

35

40

151

45


50

55

60


65

-continued



152



153



151


45

154



441

-continued




155

5

10

15

20




156

25

30

35

40



157

50


55

60


65

442

-continued




158




159

160



443

-continued



161

5

10

15

20

25

162

30

35

40

163



45

50


55

60


65

**444**

-continued





164






165

445

-continued


**446**

-continued






447

-continued

**448**

-continued



450

175

5

10

15

20



25

30

35

40

176



177

5

10

15

20

25

30

35

40

176


45

50

55

60

65




178

451

-continued

179



5

10

15


20

25

30

35

180 40



45

50

55


60

65

452

-continued

181




30

35

180 40

182



45

50



55

60



65

453

-continued


**454**

-continued



455

-continued



187

5

10

15

20


25

30

35

40


188



65

456

-continued



189

5

10

15

20


25

30

35

40

188




190

65

457

-continued

**458**

-continued

191

5

10

15

20

25

30

35

40

192

45


50

55



60

65

193




194



459

-continued



195

5

10

15

20


25

30

35

460

-continued



197

196 40


45

50


55

60



65



198



-continued



-continued

201

199

5

10

15

20

25

30

35

40

200



45

50

55

60

65



202

203


45

50

55




60

65



463

-continued

**464**

-continued

204

5

10

15

20

205

25

30

35

40

206 45

50

55




60

65

207



208

209



465

-continued

**466**

-continued

210

5

10

15

20

25

30

35

40

211

45



50

55

60

65




212



213

467

-continued

**468**

-continued

214

5

10

15

20

215

25

30

35

40

216


45

50

55

60



65



217




218

219



469

-continued

**470**

-continued

220


223

5

10

15

20



221


25

30

35

40

224



222 45


50

55

60


65

225



471


-continued



226

472

-continued



229

5


10

15

20


25

227



45

228



65


30

35

40

45

230







231

50

55

60



473**474**

-continued

232

235

5

10

15

20

233

25

30

35

40

45




234

50

55




60

65



475

-continued

**476**

-continued

238

241

5

10

15

20

25

239

242

30

35

40

45

240




243

50

55


60

65



477

-continued



244

5


10

15

20

25


245



40

45

246



50


55

60


65

478

-continued




247



248

479

-continued

**480**

-continued

249

5

10

15

20

25

30

35

40

250

45


50

55


60

65

251




252



481

-continued

**482**

-continued

253

5

10

15

20

25

30

35

40

254

45


50

55



60

65

255




256



483

-continued

**484**

-continued

257

5

10

15

20

25

30

35

40

258

45


50

55



60

65

259




260



485

-continued

**486**

-continued

261

5

10

15

20

25

30

35

40

262

45

50



55

60

65



263

264



487

-continued

**488**

-continued

265

267

5

10

15

20

25

30

35

40

266

268



45

50

55


60

65



489

-continued

**490**

-continued

269

271

5

10

15

20

25

30

35

40

270




45

50

55


60

65



491


-continued




273

492


-continued



275



274




276

65

493

-continued

**494**

-continued

277

279

5

10

15

20

25

30

35

40

278




45

50

55



60

65



495

-continued

**496**

-continued

281

283

5

10

15

20

25

30

35

40

282

284



45

50

55

60


65



497

-continued

285



5

10

15

20


25

30

35

40

286




65

498

-continued

287



25

30

35

40


45

50

55

60


288



65

499

-continued



289

5

10

15

20


25

30

35

40


290



65

500

-continued



291

5

10

15

20


25

30

35

40

290



292

45

50


55

60

65

501

-continued



293

5

10

15

20

25

30

35

40

294

45

50


55

60

65

502

-continued





295

296

503

-continued

**504**

-continued

297

299

5

10

15

20

25

30

35

40

298

300



45

50

55


60

65



505

-continued



301

5

10

15

20


25

30

35

40


302



65

506

-continued



303

5

10

15

20


25

30

35

40

302



304

45

50



55

60

65

507

-continued

**508**

-continued

305

5

10

15

20

25

30

35

40

306



45

50

55

60

65






307

308

309

509

-continued

**510**

-continued


310

5

10

15

20



313


311

25

30

35

40



314

312


45

50

55

60




65



315

511

-continued

**512**

-continued

316

5

10

15

20

317

25

30

35

40




318 45

50

55

60

65






319

320

321

513

-continued

**514**

322


325

5

10

15

20



25


323

30

35

40

45




324

50

55

60

65






326

327

515

-continued

**516**

-continued

328


5

10

15

20

329




331

25

30

35

40



332

45


330

50

55

60




65



333

517

-continued

**518**

-continued

334

5

10

15

20

335 25

30

35

40

45



336

50

55

60

65





337

338

519

-continued

**520**

-continued

339

5

10

15

20

25

30

35

40



340 45

50

55

60




65



341

342

-continued



-continued

343

346

5

10

15

20

344 25

30

35

40

345

45



50

55

60


65

347



523

-continued

**524**

-continued

348

5

10

15

20

25

30

35

40

349


45

50


55

60


65



350




351



525

-continued

**526**

-continued

352

5

10

15

20

25

30

35

40

353

45


50

55



60

65


354




355



-continued




-continued




358

357 40




359



529

-continued

**530**

-continued

360

5

10

15

20

25

30

35

361 40

45


50

55



60

65

362




363



531

-continued

**532**

-continued

364

5

10

15

20

25

30

35

365 40


45

50


55

60


65



366





367



533

-continued

**534**

-continued

368

370

5

10

15

20

25

30

35

369 40

371



45

50

55


60

65



535

-continued



372

5

10

15

20

25

30

35

40

373

45

50


55

60

65

536

-continued





374

375

537

-continued

**538**

-continued

376

5

10

15

20

25

30

35

40

377




45

50

55

60

65





378

379

380

539

-continued

**540**

-continued

381

5

10

15

20

25

30

35

40

382


45

50

55

60



65



383




384

385



541

-continued

**542**

-continued

386

5

10

15

20

387 25

30

35

40

45



388

50

55

60

65






389

390

543


-continued

**544**

-continued

391

394



20

25

395

392

30

35

40

45

393


396

50

55


60

65



545

-continued




397

5

10

15




398

20

25

30




35

399

40

45



400

50


55

60


65

546

-continued




401



402

547

-continued

**548**

-continued

403

5

10

15

20

25

30

35

40

404

45

50



55

60

65



405

406



549

-continued

**550**

-continued

407

5

10

15

20

25

30

35

40

408

45

50



55

60

65


409

410



551

-continued

**552**

-continued

411

5

10

15

20

25

30

35

40

412

45

50



55

60

65



413

414



553

-continued

**554**

-continued

415

5

10

15

20

25

30

35

40

416



45

50

55

60

65




417

418

555

-continued



419

5

10

15

20

25

30

35

40


420 45

50


55

60


65

**556**

421




422



557

-continued

**558**

-continued

423

5

10

15

20

25

30

35

40

45

424


50

55



60

65

425




426






559

-continued





560

-continued



561

-continued

**562**

-continued

431

433

5

10

15

20

25

30

35

40

432

45

434




50

55

60




65

435



563

-continued

**564**

-continued

436

439

5

10

15

437

20

25

30

35

40

438

45



50

55

60




65

440



565

-continued

**566**

-continued

441

5

10

15

20

442

25

30

35

40

443


45

50

55

60



65



444



445

446



567

-continued

**568**

-continued

447

449

5

10

15

20

25

30

35

40

448

45

450




451

50

55




60

65



569

-continued

**570**

-continued

452

455

5

10

15

453

20

25

30

35

40

454



45

50

55

60




65



456

571

-continued

**572**

-continued

457

5

10

15

20

458

25

30

35

40

459

45


50

55


60

65


460



461





462



573

-continued

**574**

-continued

463

5

10

15

20

25

30

35

40

464

45



50

55

60

65



465



466

575

-continued

**576**

-continued

467

5

10

15

20

25

30

35

40

468

45

50

55



60

65

469


470

471










577

-continued


**578**

-continued




579

-continued





**580**

-continued






581

-continued




**582**

-continued






583

-continued



**584**

-continued



585

-continued

**586**

-continued

501

5

10

15

20

25

30

35

40

502




45

50

55

60

65






503

504

505

587

-continued

**588**

-continued

506

5

10

15

20

507

25

30

35

40

508

45

50

55




60

65

509




510

511



589

-continued

**590**

-continued

512

5

10

15

20

513

25

30

35

40

45




514

50

55

60

65




515

516

517


591

-continued



593

-continued



526

5


10

15

20

25

527




30

35

40

45

528




55

60

65


594

-continued




529

530







531






595

-continued

**596**


-continued



597

-continued

539




5

10

15

20

540




30

35

40

45

541 50



55


60

65


598

-continued


542




543



544







545







599

-continued


**600**

-continued



601

-continued



554

5

10

15

20

25


555

30

35

40

45




556

50


55

60

65


**602**

-continued




557

558




559









603

-continued


**604**

-continued




605

-continued






**606**

-continued








607

-continued


**608**

-continued



609

-continued



586


5

10

15

587

20




25

30

35

588




40

45

50

589

55




60

65


610

-continued




590


591



592




593



611


-continued

594




20

595




40

596




597

55


**612**

-continued


598




599



600






601






613

-continued


**614**

-continued



615

-continued



608

5


10

15

20

616

-continued



611

609

25

30

35

40

610


45

50


55

60

65




613



617

-continued



614

5


10

15

20

618

-continued



617



615 25

30

35

40

45



618



616

50

55

60


65



619

619

-continued

**620**


-continued

620

5

10

15



624

625

20



621

25

30

35

622





626

40

45

50

623





627

55





60

65






621

-continued




**622**

-continued







623

-continued




**624**

-continued






625

-continued





**626**

-continued






627

-continued




**628**

-continued






629

-continued


**630**

-continued



631

-continued



661

632


-continued

664


5

10

15



662



25

30

35

8. An organic light emitting device comprising:
a first electrode;

20

a second electrode; and

one or more organic material layers provided between the first electrode and the second electrode,

wherein one or more layers of the organic material layers include the heterocyclic compound of claim 1.

25

9. The organic light emitting device of claim 8, wherein the organic material layer includes an electron transfer layer, and the electron transfer layer includes the heterocyclic compound.

10. The organic light emitting device of claim 8, wherein the organic material layer includes a hole blocking layer, and the hole blocking layer includes the heterocyclic compound.

30

11. The organic light emitting device of claim 8, further comprising one, two or more layers selected from the group consisting of a light emitting layer, a hole injection layer, a hole transfer layer, an electron injection layer, an electron transfer layer, an electron blocking layer and a hole blocking layer.

35

12. The organic light emitting device of claim 8 comprising:

663

a first stack provided on the first electrode and including a first light emitting layer;

40

a charge generation layer provided on the first stack;

a second stack provided on the charge generation layer and including a second light emitting layer; and the second electrode provided on the second stack.

45

13. The organic light emitting device of claim 12, wherein the charge generation layer includes the heterocyclic compound.

14. The organic light emitting device of claim 12, wherein the charge generation layer includes an N-type charge generation layer, and the N-type charge generation layer includes the heterocyclic compound.

50

15. The organic light emitting device of claim 12, wherein the second stack further includes a hole injection layer, and the hole injection layer includes the heterocyclic compound.

* * * * *