

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent	12404333
Kind Code	B2
Date of Patent	September 02, 2025
Inventor(s)	Fang; Jianmin et al.

Anti PD-L1 antibody and use thereof

Abstract

Provided in the present invention is a PD-L1 immunosuppressant, and specifically provided are a monoclonal antibody capable of targeting PD-L1, and nucleotides, combinations of polynucleotides, expression vectors and combinations of expression vectors encoding the antibody. Also provided in the present invention is a conjugate or pharmaceutical composition containing the above-mentioned anti PD-L1 antibody. Further provided in the present invention is the use of the above-mentioned anti PD-L1 antibody, nucleotides, combinations of polynucleotides, expression vectors, combinations of expression vectors, conjugate or pharmaceutical composition in the preparation of a medicament for treating or preventing cancers.

Inventors:	Fang; Jianmin (Shandong, CN), Jiang; Jing (Shandong, CN), Li; Shenjun (Shandong, CN), Zhao; Guorui (Shandong, CN)
Applicant:	REMEGEN CO., LTD. (Shandong, CN)
Family ID:	74684171
Assignee:	REMEGEN CO., LTD. (Shandong, CN)
Appl. No.:	17/312884
Filed (or PCT Filed):	August 25, 2020
PCT No.:	PCT/CN2020/110935
PCT Pub. No.:	WO2021/037007
PCT Pub. Date:	March 04, 2021

Prior Publication Data

Document Identifier	Publication Date
US 20220177590 A1	Jun. 09, 2022

Foreign Application Priority Data

CN

201910805440.7

Aug. 29, 2019

Publication Classification

Int. Cl.: C12N15/63 (20060101); A61K39/00 (20060101); A61P35/00 (20060101); C07K16/28 (20060101); C12N15/62 (20060101)

U.S. Cl.:

CPC C07K16/2827 (20130101); A61P35/00 (20180101); A61K2039/505 (20130101); C07K2317/24 (20130101); C07K2317/76 (20130101); C07K2317/92 (20130101); C12N15/62 (20130101); C12N15/63 (20130101); C12N2510/00 (20130101)

Field of Classification Search

USPC: None

References Cited

U.S. PATENT DOCUMENTS

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
9334331	12/2015	Igawa	N/A	C07K 16/36
10421807	12/2018	Gonzales	N/A	A61P 17/08
2019/0083644	12/2018	Yoo et al.	N/A	N/A

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
3137211	12/2020	CA	N/A
104470949	12/2014	CN	N/A
109195991	12/2018	CN	N/A
109232740	12/2018	CN	N/A
109627338	12/2018	CN	N/A
3378871	12/2017	EP	N/A
3470429	12/2018	EP	N/A
3858862	12/2020	EP	N/A
2018-536401	12/2017	JP	N/A
2523894	12/2013	RU	N/A
2580045	12/2015	RU	N/A
201620936	12/2015	TW	N/A
WO 2007/005874	12/2006	WO	N/A
WO 2017/020858	12/2016	WO	N/A
WO 2017/058859	12/2016	WO	N/A
WO 2017/132562	12/2016	WO	N/A
WO 2017/220990	12/2016	WO	N/A
WO 2018/052818	12/2017	WO	N/A
WO 2019/129211	12/2018	WO	N/A

OTHER PUBLICATIONS

Al Qaraghuli et al. (2020, Nature Scientific Reports 10:13969). cited by examiner

Edwards et al. (2003, JMB 334:103-118). cited by examiner

Lloyd et al. (2009, Protein Engineering, Eng. Design & Selection 22(3): 159-168). cited by examiner

Goel et al. (2004, J. Immunol. 173: 7358-7367). cited by examiner

Khan et al. (2014, J. Immunol. 192: 5398-5405). cited by examiner

Poosarla et al. (2017, Biotechn. Bioeng. 114(6): 1331 -1342). cited by examiner

Rabia et al. (2018, Biochemical Engineering Journal 137:365-374). cited by examiner

Gura (Science, v278, 1997, pp. 1041-1042). cited by examiner

Freshney (Culture of Animal Cells, A Manual of Basic Technique, Alan R. Liss, Inc., 1983, New York, p. 4). cited by examiner

Zips et al. (2005, In Vivo 19:1-8). cited by examiner

Editorial, (2013, Nature Biotechnology 31:85). cited by examiner

International Search Report in PCT/CN2020/110935 mailed Dec. 2, 2020. cited by applicant

Chen, L and Flies, D B., “Molecular mechanisms of T cell co-stimulation and co-inhibition”, Nature Reviews Immunology, 2013, 13(4): 227-242. cited by applicant

Ohaegbulam, K C et al., “Human cancer immunotherapy with antibodies to the PD-1 and PD-L1 pathway,” Trends in Molecular Medicine, 2015, 21(1): 24-33. cited by applicant

Ishida, M et al., “Differential expression of PD-L1 and PD-L2, ligands for an inhibitory receptor PD-1, in the cells of lymphohematopoietic tissues,” Immunology Letters, 2002, 84(1): 57-62. cited by applicant

Taube, J M et al., “Colocalization of inflammatory response with B7-h1 expression in human melanocytic lesions supports an adaptive resistance mechanism of immune escape”, Science Translational Medicine, 2012, 4(127): 127ra37. cited by applicant

Tseng, S Y et al., “B7-DC, a new dendritic cell molecule with potent costimulatory properties for T cells,” The Journal of Experimental Medicine, 2001, 193(7): 839-845. cited by applicant

Butte, M J et al., “Programmed death-1 ligand 1 interacts specifically with the B7-1 costimulatory molecule to inhibit T cell responses,” Immunity, 2007, 27(1): 111-122. cited by applicant

Cheng X et al., “Structure and interactions of the human programmed cell death 1 receptor,” The Journal of Biological Chemistry, 2013, 288(17): 11771-11785. cited by applicant

Dirix, Luc Y., “Avelumab, an anti-PD-L1 antibody, in patients with locally advanced or metastatic breast cancer: a phase 1b JAVELIN Solid Tumor Study”, Breast Cancer Res Treat (2018) 167:671-686. cited by applicant

Boyerinas, Benjamin, et al., “Antibody-dependent cellular cytotoxicity (ADCC) activity of a novel anti-PD-L1 antibody avelumab (MSB0010718C) on human tumor cells,” Cancer Immunol Res. Oct. 2015; 3(10): 1148-1157. cited by applicant

Liu, An, “Discovery on small molecular inhibitors of PD-1/PD-L1 pathway”, Jilin: Jilin University School of Pharmaceutical Sciences, 2016, (in 77 pages). cited by applicant

European Search Report received in European Application No. 20859087.7 dated Jun. 13, 2022 in 9 pages. cited by applicant

Japanese Office Action received in Japanese Application No. 2021-549665 dated Aug. 18, 2022 in 14 pages. cited by applicant

Russian Office Action in Russian Patent Application No. 2021108266 issued Jan. 21, 2022. cited by applicant

Pakula, Andrew A., et al., “Genetic Analysis of Protein Stability and Function,” Annual Rev. Genet. 1989, 23:289-310. cited by applicant

Background/Summary

FIELD

(1) The present invention relates to the field of biomedicine, specifically to an anti-PD-L1 antibody or antigen-binding fragment thereof, and medical use thereof.

REFERENCE TO SEQUENCE LISTING

(2) A Sequence Listing submitted as an ASCII text file via EFS-Web is hereby incorporated by reference in accordance with 35 U.S.C. § 1.52(e). The name of the ASCII text file for the Sequence Listing is 35124494_1.TXT, the date of creation of the ASCII text file is Jun. 10, 2021, and the size of the ASCII text file is 14,066 bytes.

BACKGROUND

(3) Recently, in the field of tumor therapy, increasing efforts have been dedicated to utilizing the body's immune system to defense tumors. This method of inhibition and killing of tumor cells by mobilizing the body's immune system is called as tumor immunotherapy. Tumor immunotherapy, including cell immunotherapy, tumor vaccines, passive immunotherapy targeting tumors, and immune checkpoint inhibitors, is currently the most promising research direction in the field of tumor therapy and has yielded a number of prospective research results.

(4) Immune checkpoint refers to a signaling pathway that controls the intensity of T cell immune response by balancing costimulatory and co-suppressive signals (Reference 1). Immune checkpoints can maintain immune tolerance by regulating the intensity of the autoimmune response under normal circumstances. When the body is invaded by tumors, however, the activation of immune checkpoints can inhibit autoimmunity, which favors the growth and escape of tumor cells. Immune checkpoints such as CTLA-4, programmed death receptor-1 (PD-1)/programmed death ligand-1 (PD-L1) and TIM-3 are the key negatively regulatory molecules, playing an important role in tumors' immune evasion. Blocking the negatively regulatory pathway of immune checkpoints with a specific antibody and rebuilding the ability of body's immune system to recognize and kill tumor cells have achieved good therapeutic outcomes in tumor immunotherapy.

(5) Among the known immune checkpoints, PD-1/PD-L1 has drawn high attention in tumor immune the anti-PD-L1 antibody or antigen-binding fragment to be expressed, and recovering the antibody or fragment.

(6) The present invention further provides a method for enhancing T cell function, comprising administering an effective amount of the above-mentioned pharmaceutical composition of the present invention to dysfunctional T cells.

(7) In another aspect, the present invention provides a method for the treatment or prevention of cancer, comprising administering a therapeutically effective amount of the antibody, polynucleotide, polynucleotide combination, expression vector, conjugate and/or pharmaceutical composition of the present invention to a subject in need thereof.

(8) In yet another aspect, the present invention provides use of the anti-PD-L1 antibody or antigen-binding fragment thereof, the polynucleotide, the polynucleotide combination, the corresponding nucleic acid construct or vector encoding the antibody or antigen-binding fragment thereof, the antibody-drug conjugate or the pharmaceutical composition of the present invention for the manufacture of a medicament for use in the treatment or prevention of cancer.

(9) In yet another aspect, the present invention provides the antibody, polynucleotide, polynucleotide combination, expression vector, conjugate and/or pharmaceutical composition of the present invention, for use in the treatment or prevention of cancer.

(10) Further, the cancer is a solid tumor.

- (11) Further, the solid tumor is lung cancer, colorectal cancer, breast cancer, ovarian cancer, melanoma, bladder cancer, urothelial cancer, kidney cancer, liver cancer, salivary cancer, stomach cancer, gliomas, thyroid cancer, thymic cancer, epithelial cancer, head and neck cancer, gastric and pancreatic cancer.
- (12) Further, the lung cancer is non-small cell lung cancer.
- (13) Further, the ovarian cancer is triple negative breast cancer.
- (14) Yet another aspect of the present invention provides a method for the treatment of T cell dysfunction, comprising administering a therapeutically effective amount of the above-mentioned pharmaceutical composition of the present invention to a patient with T cell dysfunction. The T cell dysfunction comprises infection and tumor immunization. The tumor immunization is caused by a cancer selected from breast cancer, lung cancer, colon cancer, ovarian cancer, melanoma, bladder cancer, kidney cancer, liver cancer, salivary cancer, stomach cancer, glioma, thyroid cancer, thymic cancer, epithelial cancer, head and neck cancer, and gastric and pancreatic cancer.
- (15) The present invention further provides use of the anti-PD-L1 antibody or antigen-binding fragment thereof, the polynucleotide, the polynucleotide combination, the corresponding nucleic acid construct or vector encoding the antibody or antigen-binding fragment thereof, the antibody-drug conjugate or the pharmaceutical composition of the present invention for the manufacture of a medicament for use in treatment or prevention of cancer.
- (16) The anti-PD-L1 monoclonal antibody provided by the present invention is a novel monoclonal antibody targeting PD-L1 with a new CDR sequence and amino acid sequence. The anti-PD-L1 monoclonal antibody provided by the present invention has a surprising affinity and specificity to human PD-L1, and has significant advantages in competing with human PD-1 to bind to human PD-L1. The anti-PD-L1 monoclonal antibody provided by the present invention exhibits unexpected tumor inhibitory effects in in vitro animal models of both non-small cell lung cancer and colorectal cancer, and provides a better option for inhibiting tumor progression.
-

Description

BRIEF DESCRIPTION OF DRAWINGS

- (1) FIG. 1 shows an affinity binding and dissociation curve of hAAG5D8 to human PD-L1, where the curve 1 indicates 100 nM hAAG5D8, the curve 2 indicates 50 nM hAAG5D8, and the curve 3 indicates 25 nM hAAG5D8.
- (2) FIG. 2 shows a curve of competing between hAAG5D8 and human PD-1 for binding.
- (3) FIG. 3 shows the binding ability of hAAG5D8 to various B7 family proteins (PD-L1, PD-L2, B7-H3, PD-1 and CD80).
- (4) FIG. 4 shows the growth curve of tumor after administration of hAAG5D8 (5 mg/kg), MPDL3820A (5 mg/kg, positive control) or human IgG1 (5 mg/kg, negative control) in a subcutaneous xenograft tumor model of human non-small cell lung cancer.
- (5) FIG. 5 shows the change in tumor volume after administration of hAAG5D8 (1 mg/kg, 3 mg/kg, or 9 mg/kg), MPDL3820A (10 mg/kg, positive control) or PBS (negative control) in a subcutaneous xenograft tumor model of human colorectal cancer.

DETAILED DESCRIPTION

(6) Definitions

(7) Unless defined otherwise, all technical and scientific terms used herein have the same meaning as understood by those ordinarily skilled in the art. With regard to the definitions and terms in the art, reference may be made to Current Protocols in Molecular Biology (Ausubel). The standard three- and/or one-letter code used for expressing one of 20 common L-amino acids in the art is adopted as the abbreviation of an amino acid residue.

(8) In the present invention, a method for determining or numbering the complementarity

determining region (CDR) of an antibody's variable domain includes IMGT, Kabat, Chothia, AbM and Contact, which are well known in the art.

(9) For the purposes of the present invention, the “consistency”, “identity” or “similarity” between two nucleic acid or amino acid sequences refers to the percentage of identical nucleotides or identical amino acid residues between the two sequences to be compared after optimal alignment. The percentage is purely statistical and the differences between the two sequences are randomly distributed and cover their full length. Sequence comparison between two nucleic acid or amino acid sequences are usually performed by comparing these sequences after they have been optimally matched, and the comparison can be performed on a segment or on a “comparison window”. In addition to manual implementation, the optimal alignment for comparing sequences can also be performed by the local homology algorithm of Smith and Waterman (1981) [Ad. App. Math. 2: 482], the local homology algorithm of Needleman and Wunsch (1970) [J. Mol. Biol. 48: 443], the similarity search method of Pearson and Lipman (1988) [Proc. Natl. Acad. Sci. USA 85: 2444], or a computer software using these algorithms (GAP, BESTFIT, FASTA and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI, or BLAST N or BLAST P comparison software).

(10) As used herein, “antibody” is used in a broadest sense and encompasses various antibodies including, but not limited to, a monoclonal antibody, a polyclonal antibody, and a multispecific antibody (e.g., a bispecific antibody). As used herein, “antigen-binding fragment” refers to an antibody fragment consisting of or comprising a partial sequence of a heavy or light variable chain of an antibody from which it is derived, wherein the partial sequence is capable of retaining the same binding specificity as the antibody from which it is derived and a sufficient affinity, preferably equal to at least 1/100, more preferably at least 1/10 of the affinity of the antibody from which it is derived. Such a functional fragment comprises a minimum of 5 amino acids, preferably 10, 15, 25, 50 or 100 contiguous amino acids of the antibody sequence from which it is derived, including (particularly) Fab, F(ab'), F(ab').sub.2, Fv, dAb, Fd, a complementarity determining region (CDR) fragment, a single chain antibody (scFv), and a bivalent single chain antibody, that contains at least an immunoglobulin fragment enough to allow a specific antigen to bind to the polypeptide. The above fragments can be prepared by a synthetic or enzymatic method, or by chemical cleavage of an intact immunoglobulin, or can be genetically engineered by recombinant DNA technology. The preparation methods thereof are well known in the art. A heavy chain contains a heavy chain variable region (abbreviated as VH) and a heavy chain constant region. The heavy chain constant region contains three domains, CH1, CH2 and CH3. A light chain contains a light chain variable region (abbreviated as VL) and a light chain constant region. The light chain constant region contains a domain, CL. VH and VL regions can be further subdivided into multiple regions with high variability, called as complementarity determining regions (CDRs), interspersed with more conservative regions called as framework regions (FRs). Each VH and VL is composed of three CDRs and four FRs, which are arranged from the amino terminal to the carboxy terminal in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. These variable regions of the heavy and light chains contain a binding domain that interacts with an antigen. The constant region of an antibody can mediate binding of an immunoglobulin to a host tissue or factor, including various cells in the immune system (such as effector cells) and the first component of the classical complement system (C1q). Chimeric or humanized antibodies are also encompassed by the antibodies according to the present invention.

(11) The term “humanized antibody” refers to an antibody that contains a CDR region derived from a non-human antibody, with the rest deriving from one (or several) human antibody. Moreover, in order to retain binding affinity, some residues at the backbone (called FR) segment can be modified (Jones et al., Nature, 321: 522-525, 1986; Verhoeyen et al., Science, 239: 1534-1536, 1988; Riechmann et al., Nature, 332: 323-327, 1988). Humanized antibodies or fragments thereof according to the present invention can be prepared by techniques known to those skilled in the art

(e.g., described in the document Singer et al., J. Immun. 150: 2844-2857, 1992; Mountain et al., Biotechnol. Genet. Eng. Rev., 10: 1-142, 1992; or Bebbington et al., Bio/Technology, 10: 169-175, 1992).

(12) The term “chimeric antibody” refers to an antibody in which the variable region sequence is from one species while the constant region sequence is from another species, for example, an antibody in which the variable region sequence is from a mouse antibody while the constant region sequence is from a human antibody. A chimeric antibody or a fragment thereof according to the present invention can be prepared by using genetic recombination technology. For example, the chimeric antibody can be produced by cloning a recombinant DNA comprising a promoter and a sequence encoding a variable region of a non-human, especially a murine monoclonal antibody according to the present invention, and a sequence encoding a constant region of a human antibody. The chimeric antibody of the present invention encoded by such a recombinant gene will be, for example, a murine-human chimera whose specificity is determined by the variable region derived from murine DNA, and the isotype is determined by the constant region derived from human DNA. For methods for preparing a chimeric antibody, for example, reference can be made to the document Verhoeven et al. (BioEssays, 8:74, 1988).

(13) The term “monoclonal antibody” refers to a preparation of an antibody molecule consisting of a single molecule. Monoclonal antibody compositions display a single binding specificity and affinity for a particular epitope.

(14) The term an “isolated” nucleic acid molecule refers to a nucleic acid molecule identified and separated from at least one contaminant nucleic acid molecules, and is generally associated with the contaminant nucleic acid molecule in the natural source of an antibody nucleic acid. An isolated nucleic acid molecule is different in form or environment from when it is found in nature, and therefore different from that existing in natural cells. However, an isolated nucleic acid molecule comprises a nucleic acid molecule contained in cells where an antibody is usually expressed, and where for example, it is located on a different chromosomal position from that in a natural cell.

(15) Generally, in order to prepare a monoclonal antibody or functional fragment thereof, especially a murine-derived monoclonal antibody or functional fragment thereof, reference can be made to the technology especially described in the manual “Antibodies” (Harlow and Lane, Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor NY, pp. 726, 1988) or the technique for preparation from hybridoma cells described by Kohler and Milstein (Nature, 256: 495-497, 1975).

EXAMPLES

(16) The embodiments of the present invention will be described in detail below in conjunction with examples. However, it will be understood by those skilled in the art that the following examples are only used to illustrate the present invention and should not be regarded as limiting the scope of the present invention.

Example 1 Production of Anti-PD-L1 Antibody

(17) Through extensive screening of anti-PD-L1 antibodies obtained after immunizing mice, a candidate murine antibody mAAG5D8 was determined. After sequence alignment in the antibody variable region database, a human IgG1 framework region with a high homology to the murine PD-L1 antibody mAAG5D8 was determined. For mAAG5D8, a variety of humanized antibodies was further designed and compared for their affinity, and finally a candidate humanized anti-PD-L1 antibody hAAG5D8 was determined.

(18) TABLE-US-00001 TABLE 6 CDRs 1-3 amino acid sequences of heavy chain and light chain variable regions of anti-PD-L1 humanized antibody hAAG5D8 (determined by IMGT method) Heavy chain CDR1 SEQ ID NO: 1 GFSLSRYS CDR2 SEQ ID NO: 2 IWGVGTT CDR3 SEQ ID NO: 3 ARNWGTADYFDY Light chain CDR1 SEQ ID NO: 4 KSVHTSGYSY CDR2 SEQ ID NO: 5 LAS CDR3 SEQ ID NO: 6 QHSGELPYT

(19) The heavy chain variable region of anti-PD-L1 humanized antibody hAAG5D8 comprises an amino acid sequence of (SEQ ID NO: 31):

(20) TABLE-US-00002 QVQLQESGPG LVKPSETLSL TCTVSGESLS RYSVHWIRQP
PGKGLEWLGM IWGVGITDYN 60 SALKSRLTIS KDTSKNQFSL KLSSVTAADT
AVYYCARNWG TADYFDYWGQ GTIVTVSSAS 120

(21) The light chain variable region of anti-PD-L1 humanized antibody hAAG5D8 comprises an amino acid sequence of (SEQ ID NO: 32):

(22) TABLE-US-00003 DIVLTQSPAS LAVSPGQRAT ITCRLSKEVH TSGYSIMHWY
QQKPGQPPKI LIYLASNLES 60 GVPARFSGSG SGTDFTLTIN PVEANDTANY
YCQHSGLPEY TFGGGTKVEI KRT 113

(23) The amino acid sequences of the heavy chain and light chain of anti-PD-L1 humanized antibody hAAG5D8 are shown in SEQ ID NO: 33 and SEQ ID NO: 34, respectively.

Example 2 Comparison of Affinity of hAAG5D8 to Human PD-L1

(24) The affinity of hAAG5D8 to human PD-L1 was detected using biolayer interferometry (BLI). The wells in A-D columns 1, 3, and 5 of a black 96-well plate were added with a PBS solution as baseline 1, baseline 2 and a dissociation solution, respectively. The wells in column 2 were added with a PD-L1 solution (R&D company), in column 4 with a hAAG5D8 solution (The concentrations were as follows: 100 nM, 50 nM, 25 nM, and 0 nM), in column 10 with an imidazole solution, in column 11 with water, and in column 12 with a nickel sulfate solution. Ni-NTA probe (Fortebio) was employed in the experiment. First, the probe was dipped in A-D column 1 for 180 seconds to stabilize the baseline. The Ni-NTA probe was then dipped in A-D column 2 for 300 seconds to immobilize PD-L1 on the probe. The Ni-NTA probe was then dipped in A-D column 3 for 120 seconds to stabilize the baseline. The Ni-NTA probe was then dipped in A-D column 4 for 600 seconds to allow the binding of hAAG5D8 with a different concentration to the immobilized PD-L1 protein on the probe. The Ni-NTA probe was then dipped in A-D column 5 for 600 seconds to allow the antibody to dissociate spontaneously. Finally, the Ni-NTA probe was dipped in A-D columns 10, 11, and 12 successively to force the immobilized PD-L1 dissociating from the probe. Data analysis was performed by Data analysis 7.0 to obtain the binding and dissociation equilibrium constant $K_{sub.D}$.

(25) Results and conclusions: The binding and dissociation equilibrium constant $K_{sub.D}$ value and affinity binding and dissociation curve of hAAG5D8 to human PD-L1 are shown in Table 7 and FIG. 1, respectively. The experimental results show that the affinity of hAAG5D8 to human PD-L1 is much higher than the affinity of PD-1 to PD-L1 of 8.2 M (Molecular Interactions of Antibody Drugs Targeting PD-1, PD-L1, and CTLA-4 in Immuno-Oncology, Hyun Tae Lee et al., Molecules 2019, 24, 1190; doi:10.3390/molecules24061190, Mar. 26, 2019, last paragraph on page 4).

(26) TABLE-US-00004 TABLE 7 Affinity kinetic constant of anti-PD-L1 antibody binding to PD-L1 Main parameter $K_{sub.D}(M)$ Affinity kinetic constant $7.62E-11 \pm 3.32E-11$

Example 3 Competing Between hAAG5D8 and Human PD-1 for Binding

(27) The ability of hAAG5D8 to compete with PD-1 to bind to PD-L1 was tested by ELISA. PD-L1 (Fc Tag) was diluted to 1 $\mu\text{g/mL}$ with a coating buffer (6 mM Na_2CO_3 , 14 mM NaHCO_3), and then added 100 μL to each well followed by overnight incubation at 4° C. The plate was then washed with PBST, and added with 250 μL of a blocking solution (3% BSA/PBST) to each well to block for 2 h at 25° C. Afterwards, the plate was washed with PBST and added with 50 μL of human PD-1 at a concentration of 6 $\mu\text{g/mL}$ as well as 50 μL of a gradiently diluted hAAG5D8 solution (at a concentration of 8000 ng/mL, 2000 ng/mL, 500 ng/mL, 250 ng/mL, 125 ng/mL, 62.5 ng/mL, 31.25 ng/mL, 6.25 ng/mL, or 1.25 ng/mL) to each well followed by mixing well and incubation at 25° C. for 2 h. Next, the plate was washed with PBST and added with 100 μL of an enzyme-labeled secondary antibody (goat anti-human IgG-Fc-HRP diluted 1:5000) to each well, followed by incubation at 25° C. for 1 h. Finally, color development was performed and values were read at 450 nm. A four-parameter equation was used for curve fitting to calculate the

EC.sub.50 value.

(28) Results and conclusions: EC.sub.50 value and the competitive binding curve of hAAG5D8 competitively binding to human PD-L1 are shown in Table 8 and FIG. 2, respectively.

Experimental results show that hAAG5D8 can effectively block the binding of PD-1 to PD-L1.

(29) TABLE-US-00005 TABLE 8 Competing between hAAG5D8 and human PD-1 for binding (pM, Mean $\pm$ SD) Main parameter Anti-PD-L1 antibody EC.sub.50 of competing with human PD-1 715.56 ± 61.58

Example 4 Affinity of hAAG5D8 to a Variety of Other B7 Family Proteins

(30) The affinity of hAAG5D8 to a variety of other B7 family proteins (PD-L1, PD-L2, B7-H3, PD-1 and CD80) was tested by ELISA. hAAG5D8 was diluted to 100 μ g/mL with a coating buffer (6 mM Na.sub.2CO.sub.3, 14 mM NaHCO.sub.3), and then added 100 μ L to each well followed by overnight incubation at 4° C. The plate was then washed with PBST, and added with 250 μ L of a blocking solution (3% BSA/PBST) to each well to block for 2 h at 25° C. Afterwards, the plate was washed with PBST and added with 100 μ L of a protein sample (rhPD-L1, rhPD-L2, rhB7-H3, rhPD-1 or rhCD80) at a concentration of 5 μ g/mL to each well followed by incubation at 25° C. for 2 h. Next, the plate was washed with PBST and added with 100 μ L of an enzyme-labeled secondary antibody (the secondary antibody anti-His-tag diluted 1:5000) to each well followed by incubation at 25° C. for 1 h. Finally, the plate was washed with PBST and then color development is performed. Values were read at 450 nm.

(31) Results and conclusions: The binding of hAAG5D8 to a B7 family protein (PD-L1, PD-L2, B7-H3, PD-1 or CD80) is shown in FIG. 3. The results show that hAAG5D8 can specifically bind to PD-L1 but not PD-L2, and hAAG5D8 does not bind to other proteins of the same family with related functions (B7-H3, PD-1, and CD80). Accordingly, hAAG5D8 binds to PD-L1 with extremely high specificity.

Example 5 Study on the Therapeutic Efficacy of hAAG5D8 on Subcutaneously Transplanted Tumor in Human Non-Small Cell Lung Cancer HCC827 NCG Mice

(32) Twelve highly immunodeficient male NCG mice aged 4-5 weeks, weighing 20-26 g (purchased from Model Animal Research Center of Nanjing University) were injected with 100 μ L of human peripheral blood mononuclear cells PBMC (1×10^7 cells/100 μ L, isolated from blood of healthy donors) via tail vein injection. Three days after tail vein injection, mice were inoculated subcutaneously with human non-small cell lung cancer HCC827 cells (ATCC) on the right flank. Three days after inoculation (an average tumor volume of 57 mm³), the mice were randomly divided into 3 groups (A-1 group, A-2 group, and A-3 group) with 4 mice in each group, and administered according to the dosage regimen shown in Table 9. Both positive control and negative control were set in this experiment, wherein the positive control was MPDL3820A (i.e. Atezolizumab) and the negative control was Human IgG1 (Crown Bioscience (Taicang) Co., Ltd., lot no. AB160083). On the day of the fourth administration (that is, the 13th day after inoculation of tumor cells), the experimental mice were euthanized under CO₂ and then tumors were taken. The relative tumor inhibition (TGI.sub.RTV) and the tumor growth inhibition (TGI.sub. Δ TV) were calculated for evaluation of drug efficacy.

(33) The equation for calculating the relative tumor inhibition:

$TGI_{sub.RTV} = 1 - T_{sub.RTV} / C_{sub.RTV}$ (%). (T.sub.RTV is the relative tumor volume (TV) of the positive control group or treatment group at a specific time point, C.sub.RTV is the relative tumor volume of the negative control group at a specific time point, T.sub.RTV/C.sub.RTV is the percentage value of the relative tumor volume of the positive control group or treatment group and the relative tumor volume of the negative control group, $RTV = V_{sub.t} / V_{sub.0}$, V.sub.0 is the tumor volume of the animal at the time of grouping, and V.sub.t is the tumor volume of the animal after treatment.)

(34) The equation for calculating the tumor growth inhibition: $TGI_{sub.\Delta TV} \% = (1 - \Delta T / \Delta C) \times 100\%$. (ΔT =the average tumor volume of the positive control group or treatment group at a specific time

point—the average tumor volume of the positive control group or the treatment group at the beginning of administration, and AC=the average tumor volume of the negative control group at a specific time point—the average tumor volume of the negative control group at the beginning of administration.)

(35) Results and conclusions: The tumor inhibitory effects of hAAG5D8 on human non-small cell lung cancer models are shown in Table 10 and FIG. 4, respectively. Table 10 shows the relative tumor inhibition and tumor growth inhibition of hAAG5D8 on human non-small cell lung cancer cells, and FIG. 4 shows tumor growth curves after administration. Experimental results show that hAAG5D8 has a significant inhibitory effect on human non-small cell lung cancer.

(36) TABLE-US-00006 TABLE 9 Dosage regimen for studying the tumor inhibitory activity of hAAG5D8 on human non-small cell lung cancer Group Subject Drug Dose (mg/kg) Route Volume Period 1 A-1 group Human IgG1 5 Intravenous injection 10 μ L/g BIW $\times$ 4 times 2 A-2 group MPDL3820A 5 Intravenous injection 10 μ L/g BIW $\times$ 4 times 3 A-3 group hAAG5D8 5 Intravenous injection 10 μ L/g BIW $\times$ 4 times Note: Group 1 is the negative control group, Group 2 is the positive control group, and Group 3 is the treatment group.

(37) TABLE-US-00007 TABLE 10 Relative tumor inhibition and tumor growth inhibition of hAAG5D8 on human non-small cell lung cancer tumor cells Drug TGI.sub.RTV(%) TGI.sub. Δ .sub.TV(%) Human IgG1 (5 mg/kg), — — MPDL3820A (5 mg/kg) 25.3 49.5 hAAG5D8 (5 mg/kg) 35.9 63.3

Example 6 Study on the Therapeutic Efficacy of hAAG5D8 on Subcutaneously Transplanted Tumor in Mouse Colorectal Cancer MC38 Transgenic Mice

(38) Each of 30 human PD-1 transgenic female C57 mice (Animal Experiment Center of Tongji University) was implanted subcutaneously 3 $\times$ 10⁶ mouse colorectal cancer cells expressing human PD-L1 (Tongji University, code: MC38-hPD-L1) on the left flank. When the tumor volume reached about 100 mm³, the mice were randomly divided into 5 groups: C-1 group (negative control group, PBS, 6 mice), C-2 group (positive control group, Atezolizumab, 3 mg/kg, 6 mice), C-3 group (administration group, hAAG5D8, 1 mg/kg, 6 mice), C-4 group (administration group, hAAG5D8, 3 mg/kg, 6 mice) and C-5 group (administration group, hAAG5D8, 9 mg/kg, 6 mice). The mice were administered intraperitoneally 2 times a week, 5 times in total. The tumor volume was measured twice a week until the 16th day after the beginning of administration.

(39) Calculation Equations:

(40) Tumor volume: TV=D1 $\times$ D2²/2, wherein D1 and D2 represent tumor long diameter and tumor short diameter, respectively.

(41) Relative tumor volume: RTV=T.sub.VT/T.sub.V1, where T.sub.V1 is the tumor volume before administration, and T.sub.VT is the tumor volume on each measurement.

(42) Relative tumor proliferation: T/C(%)=T.sub.RTV/C.sub.RTV $\times$ 100%, wherein T.sub.RTV represents the RTV of the treatment group or positive control group, and C.sub.RTV represents the RTV of the negative control group.

(43) Relative tumor inhibition: TGI.sub.RTV(%)=(1-T.sub.RTV/C.sub.RTV) $\times$ 100%.

(44) Results and conclusions: The tumor inhibitory effects of hAAG5D8 on mouse colorectal cancer are shown in Table 11 and FIG. 5. The results show that hAAG5D8 at a concentration of 3 mg/kg or above has a significant tumor inhibitory effect on subcutaneously transplanted tumors of mouse colorectal cancer. At the dose of 3 mg/kg, the tumor inhibitory effect of the anti-PD-L1 antibody has been significantly superior to that of Atezolizumab.

(45) TABLE-US-00008 TABLE 11 Tumor volume changes in tumor-bearing mice after administration Dose Tumor volume TV (mm³) T.sub.RTV/ TGI.sub.RTV Group (mg/kg) D0 D16 C.sub.RTV(%) (%) PBS — 99.9 $\pm$ 13.2 1974.2 $\pm$ 413.2 — — Atezolizumab 3 95.9 $\pm$ 16.8 1111.6 $\pm$ 435.2* 53 47 hAAG5D8 1 95.5 $\pm$ 9.8 1138.8 $\pm$ 144.9* 64 36 hAAG5D8 3 90.7 $\pm$ 12.4 664.8 $\pm$ 92.0** 39 61 hAAG5D8 9 91.2 $\pm$ 15.9 319.1 $\pm$ 0190.4** 14 86 Note: The day of administration was defined as D0; *P < 0.05, **P < 0.01 vs PBS

REFERENCES

(46) 1. CHEN L, FLIES D B. Molecular mechanisms of T cell co-stimulation and co-inhibition [J]. Nature reviews Immunology, 2013, 13(4): 227-42. 2. OHAEBULAM K C, ASSALA, LAZAR-MOLNAR E, et al. Human cancer immunotherapy with antibodies to the PD-1 and PD-L1 pathway [J]. Trends in molecular medicine, 2015, 21(1): 24-33. 3. ISHIDA M, IWAI Y, TANAKA Y, et al. Differential expression of PD-L1 and PD-L2, ligands for an inhibitory receptor PD-1, in the cells of lymphohematopoietic tissues [J]. Immunology letters, 2002, 84(1): 57-62. 4. TAUBE J M, ANDERS R A, YOUNG G D, et al. Colocalization of inflammatory response with B7-h1 expression in human melanocytic lesions supports an adaptive resistance mechanism of immune escape [J]. Science translational medicine, 2012, 4(127): 127ra37. 5. TSENG S Y, OTSUJI M, GORSKI K, et al. B7-DC, a new dendritic cell molecule with potent costimulatory properties for T cells [J]. The Journal of experimental medicine, 2001, 193(7): 839-46. 6. BUTTE M J, KEIR M E, PHAMDUY T B, et al. Programmed death-1 ligand 1 interacts specifically with the B7-1 costimulatory molecule to inhibit T cell responses [J]. Immunity, 2007, 27(1): 111-22. 7. CHENG X, VEVERKA V, RADHAKRISHNAN A, et al. Structure and interactions of the human programmed cell death 1 receptor [J]. The Journal of biological chemistry, 2013, 288(17): 11771-85. 8. An Liu, Discovery on small molecular inhibitors of PD-1/PD-L1 pathway [D], Jilin: Jilin University School of Pharmaceutical Sciences, 2016, 15.

Claims

1. An isolated anti-PD-L1 antibody or antigen-binding fragment thereof comprising a heavy chain variable region and a light chain variable region, wherein the heavy chain variable region and the light chain variable region comprise the following CDRs: (1) CDRs 1-3 of the heavy chain variable region comprising the amino acid sequences of SEQ ID NOs: 1-3 respectively, and the CDRs 1-3 of the light chain variable region comprising the amino acid sequences of SEQ ID NOs: 4-6 respectively; or (2) CDRs 1-3 of the heavy chain variable region comprising the amino acid sequences of SEQ ID NOs: 7-9 respectively, and CDRs 1-3 of the light chain variable region comprising the amino acid sequences of SEQ ID NOs: 10-12 respectively; or (3) CDRs 1-3 of the heavy chain variable region comprising the amino acid sequences of SEQ ID NOs: 13-15 respectively, and CDRs 1-3 of the light chain variable region comprising the amino acid sequences of SEQ ID NOs: 16-18 respectively; or (4) CDRs 1-3 of the heavy chain variable region comprising the amino acid sequences of SEQ ID NOs: 19-21 respectively, and CDRs 1-3 of the light chain variable region comprising the amino acid sequences of SEQ ID NOs: 22-24 respectively; or (5) CDRs 1-3 of the heavy chain variable region comprising the amino acid sequences of SEQ ID NOs: 25-27 respectively, and CDRs 1-3 of the light chain variable region comprising the amino acid sequences of SEQ ID NOs: 28-30 respectively.
2. The antibody or antigen-binding fragment thereof according to claim 1, wherein: (1) the heavy chain variable region comprises the same CDRs 1-3 as in SEQ ID NO: 31 and are more than 95% identical to SEQ ID NO: 31; and (2) the light chain variable region comprises the same CDRs 1-3 as in SEQ ID NO: 32 and are more than 95% identical to SEQ ID NO: 32.
3. The antibody or antigen-binding fragment thereof according to claim 1, wherein: (1) the heavy chain variable region comprises the amino acid sequence as shown in SEQ ID NO: 31; and (2) the light chain variable region comprises the amino acid sequence as shown in SEQ ID NO: 32.
4. The antibody or antigen-binding fragment thereof according to claim 1, wherein the antibody is a monoclonal antibody, chimeric antibody, humanized antibody, bispecific antibody, multispecific antibody or Fab Fragment, F(ab') fragment, F(ab').sub.2 fragment, Fv fragment, dAb, Fd, or single chain antibody (scFv).
5. An isolated polynucleotide encoding the antibody or antigen-binding fragment thereof according to claim 1.

6. A combination of an isolated polynucleotide encoding the heavy chain of the antibody or antigen-binding fragment thereof according to claim 1 and an isolated polynucleotide encoding the light chain of the antibody or antigen-binding fragment thereof according to claim 1.
 7. A nucleic acid construct comprising the polynucleotide according to claim 5.
 8. A host cell comprising the nucleic acid construct of claim 7.
 9. An antibody-drug conjugate comprising the antibody or antigen-binding fragment thereof according to claim 1 and a drug toxin.
 10. A composition comprising the antibody or antigen-binding fragment thereof according to claim 1, and a pharmaceutically acceptable carrier.
 11. A method for the manufacture of an anti-PD-L1 antibody, comprising culturing the host cell according to claim 8 under a condition suitable for a vector encoding the anti-PD-L1 antibody or antigen-binding fragment to be expressed, and recovering the antibody or fragment.
 12. The antibody or antigen-binding fragment thereof according to claim 1, wherein: (1) the heavy chain comprises the amino acid sequence as shown in SEQ ID NO: 33; and (2) the light chain comprises the amino acid sequence as shown in SEQ ID NO: 34.
-