

(19) **United States**

(12) **Patent Application Publication**
Ben Khaled et al.

(10) **Pub. No.: US 2025/0283103 A1**
(43) **Pub. Date: Sep. 11, 2025**

(54) **METHOD OF MODULATING THE
ALKALOID CONTENT OF TOBACCO**

Publication Classification

(71) Applicant: **Nicoventures Trading Limited,**
London (GB)

(51) **Int. Cl.**
C12N 15/82 (2006.01)
A24B 3/12 (2006.01)
A24B 15/00 (2006.01)
C12N 9/12 (2006.01)

(72) Inventors: **Sara Ben Khaled**, London (GB);
Javier Galdon-Armero, London (GB);
Francisco Anastacio De Abreu E
Lima, London (GB)

(52) **U.S. Cl.**
CPC *C12N 15/8243* (2013.01); *A24B 3/12*
(2013.01); *A24B 15/00* (2013.01); *C12N*
9/1241 (2013.01); *C12Y 207/07012* (2013.01)

(73) Assignee: **Nicoventures Trading Limited,**
London (GB)

(57) **ABSTRACT**

(21) Appl. No.: **18/859,895**

The present invention provides a method for modulating the alkaloid content of a plant (e.g. a tobacco plant), the method comprising modifying said plant by modulating the activity or expression of Galactose-1-phosphate uridyl transferase. The present invention also provides for the use of Galactose-1-phosphate uridyl transferase for modulating the alkaloid content of a plant, as well as tobacco cells, plants, plant propagation materials, harvested leaves, processed tobaccos, or tobacco products obtainable in accordance with the invention.

(22) PCT Filed: **Apr. 26, 2023**

(86) PCT No.: **PCT/GB2023/051108**

§ 371 (c)(1),

(2) Date: **Oct. 24, 2024**

(30) **Foreign Application Priority Data**

Apr. 27, 2022 (GB) 2206109.7

Specification includes a Sequence Listing.




Figure 1




Figure 2




Figure 3

Gene: Nitab4.5_0002825g0060.2 genomic sequence (SEQ ID No. 1):

```
ACATAGATTATTATCTAGAATATGGGTACTCTTGCACTTACACAAAACTAGAGGAGCAAAGTTGATC
AATTTCCCTCACGFTTTTATATATTCCACGAAACCATGTTCTTAACTTCCGAGATTCTCAGTGAAGCATC
GAAGTGAAGTGGAGGAGCACCAAATTCGGAGCCCCGACACAAAGGCAAATGGCGGAGGCAGAAGCACC
AAATCGGAGCCCCGAAATTCGGAAAGACAGAATCCACAACAGGTGGGTCCCTATTCTCGCCGGCGAGAT
CACGGCGGGCCGTCTGACTTCAAGGCAAATCGAACCACAAACCGAACAATAAAACAGAATGCCCATTT
TGCGCCGGCCACGAGCACGAGTGTGCGCCGGAGATTTTTCGTGTGCCGGCAAATTCACCCGCCGATTG
GAAAATCAGAGTTATCCAAAATCTCTTCCGGCAGTCAGCAGAGAATTGGACTTCCAAAATCCGGTTT
CTGCCCTCGGGGACGTAGCTGTTAGTGGGTTCCGGTTTCACGATGTGGTTATTGAATCGCCGGTTCAT
TCAGTAAATATGAGCGATCTTCTCCGTCAAAAGTTGGGGAAGTTATGCTGGCTTTTAAGAAGCGGGT
CGAGCAGCTCCAGAATTATGATAATGTCAAATATGTCAGGTTTATTCTGATCCCGTATAAAAATGC
ATTGGTTTTTCGTATCAATTTGTCTATATTTGTAGTTGATACTGTATATTCATTCTTTGATTTTATAT
GAATTTAAAATAAATGCATTTAAAAGCTACATAAAAATAGTAGAGACCATGACTATTTAAAAGCTAAAA
GTATTTAACAAAGTATATGATGAAGGCTTGCTGCAATGGTGAATTTTTTGTGTTGCAGTAAAGGACTTT
GAAGATTTACCTGGCTTTAATCTTTTGGTGTAATTGCTTATATGGGTTTCATTTCCATTAACCTGAAAG
TGTTTTGTGTTTTGTACAAATAAAATATAAGAGTAAACGTTAATGGTAGAGCCCATCAATGTTGTACA
AAGTGACCAACCAATTATAGCTTCTTTTATGTTATATTTAGCAAACCTATTTTGTCAATTTAGTTAATT
TTATCTTTAGCTTCTTTTTATTTAGCTGCTTATCATTTGTCTGCTTTTCATTTTGTGGCTTTTAAATT
TGGCAGACTGAATGATTTTTCTGTGTAACATTGGTGGAGAAAAGGCTTAGGCTGACAACATGTTTGTC
TTTAGGTTTTAATTGAACAAATATAACAGCATTTGACGTTTGTTACTGGAAGAGTAGCTCTTTACTTA
GACCAACTTGCTAAAATACTGAACCAATTATGAGAGGCTTTTGTTGGAACTCATAGACTGTCAAGT
GTCACCTGCTTGAGGGCTTATCATTAAGGAGAATAATTTTAGAGGAAGCTTTTCCATTAATCTGCATA
GGAGCTTGATATGAATCTCCATCCATTTATCTGACATGGTTTGATTTTCATGGTAGATTATTTGA
GTTTGATTTTTTAGTGACAGGAGAAAATCTGGTAGCATTTAAACGAAGTTGGTTTGATATGGTAAATA
TCAACTTGACTGTCAATTTGAACTATTTAACAGAACTGCTCTTAACTTGTAAGTCATGTAGCAGTTT
ATCCAAAAAATAAATACTGTAAATCATGTAGCAAGTAGCATATTTGTGGTATCAAGCCAAACCTT
GTTTGATATCCCAGATTACAAACCATTTTGTAGAGGTATTGTTAGCTGCACATTCTTGTTACTTGTTT
CCTGTTCCATTGCAGAATACTCCAAAATATGCACTTCTATCTACTCATACCAAATATATGTAAGAGAA
CTTGAGAAGTGATTAGTTGTGATGAAATTGGAGATGAATGAGCACAATATCTGATCTCATGCCTTTTG
TGCGTGCAAGGAATTTGCAGTACGGAACAGCATGTGTTGTTTTTTCCACTTTCTTGTTGAGTATAATG
ATTTGTACTGTACTGAGCAAGATTGTCCCATCAAAGGCTACTGGACTATATAACACTAATTCATTCAA
TGGCTTATAAAGGTGTTCAAAAACCATGGAGCCTCTGCTGGTGCTTCAATGAGCCACTCTCACAGCCA
GATGATTGCTCTTCCCATTTGTTCTCCAACAGTTTCTGCTCGTCTTGACAGTATGAGGGAATACTACA
AGCAGACTGGGAAATGCAGCCTGTGTGATATTC AACCAAATGAATTGTTAATTGATCAATCAGCCCAT
TTCATATCATTTGGTTCCCTTTGCTGCAACTTTTGCTTTTGAGATTTGGATAATTCCTCGTGATCCTC
CTCTCATTTTTCATGAATCGATAATGAAAAGGTAATCGATGCTTTTCTTGTCATAAAGATTGTTTTCA
TGAGAAAATTATGTGAAGTTACACAGTTTTAACTTTGCAAATATCAGTTATCTCCAAATTCTACGTGA
GAAAAGTCACAACGGTAGACCATGAGGGCCTTGACTTATCTCAAGTAGAATTTGCTGATTAGTGAAGA
GAAAAGGAAAACTCATCAAAAACTGATATTCAGGAAAAAACGTACGTTGTTTCAGGCTAAACTAATT
TTTTGATTGTTTCTCCATCTCAACAACTACTTAGTTGAAAAGATTGTGTATGTTTTACCAATGAAATA
TGCAGGCAAAATGATCTTGGAGGGTTGTTGAAACTCATGCTTTTGAAGATGTCTTTCAGTTGAACAAT
CCACCATTCAACCTCCTTATTCATACCTCCCCATTTCAAGTAGATCCCTCATATGCTCCTTCCACACA
CTGGTTTTTACAAATAGCTCCCCATTTGAGTGGAGTTGGTGGATTGAGATAGCAACCGGATGTCACA
TAAATCCTGTTTTCCAGAGGATGCTGCCAAAATCTGAGGGATGTGAGAATCCTTAATTAACAGTAT
TTAGGGGGAATCTATTAAATCTAATTATTCATATCTGTAGACACAGTAACTAAGGAGATAAAGAGGAT
GTGGATACAGGGAAGATGTACCTTTTTATGCGGAGTTATGTGTAGATTCGATTTTGTGATCCATTAC
AAATGTTATCTTATGTTACACGGTCATTTTCACGCAAAAAATAATTCTCCCTCCATTTCAATTTACCT
ATTTTCTTTTGTAGTCTATGCCAACAAAAATGACCACTTCTTTATTTTGA
```

Figure 4

Nitab4.5_0002825g0060.2 coding sequence (SEQ ID No: 2):

ACATAGATTATTATCTAGAATATGGGTACTCTTGCACTTACACAAAACTAGAGGAGCAAAGTTGATC
AATTTCCCTCACGTTTTATATATATCCACGAAACCATGTTCTTAACCTCCGAGATTCTCAGTGAAGCATC
GAAGTGAAGTGGAGGAGCACCAAATTCGGAGCCCCGACACAAAGGCAAATGGCGGAGGCAGAAGCACC
AAATCGGAGCCCCGAAATTCGGAAAGACAGAATCCACAACAGGTGGGTCTTATTCTCGCCGGCGAGAT
CACGGCCGGCCGTCTGACTTCAAGGCAAATCGAACCACAAACCGAACAATAAAACAGAATGCCCATTT
TGCGCCGGCCACGAGCACGAGTGTGCGCCGGAGATTTTTCGTGTGCCGGCAAATTCACCCGGCGATTG
GAAAATCAGAGTTATCCAAAATCTCTTCCGGCAGTCAGCAGAGAATTGGACTTCCAAAATCCGGTTT
CTGCCCTCGGGGACGTAGCTGTTAGTGGGTTCCGGGTTTACGATGTGGTTATTGAATCGCCGGTTCAT
TCAGTAAATATGAGCGATCTTCTCCGTCAAGTTGGGGAAGTTATGCTGGCTTTTAAGAAGCGGGT
CGAGCAGCTCCAGAATTATGATAATGTCAAATATGTTTCAAGGTGTTCAAAAACCATGGAGCCTCTGCTG
GTGCTTCAATGAGCCACTCTCACAGCCAGATGATTGCTCTTCCATTGTTTCCCAACAGTTTCTGCT
CGTCTTGACAGTATGAGGGAATACTACAAGCAGACTGGGAAATGCAGCCTGTGTGATATTCAACCAA
TGAATTGTTAATTGATCAATCAGCCCAATTCATATCATTTGGTTCCCTTTGCTGCAACTTTTGCTTTTG
AGATTTGGATAAATTCCTCGTGATCACTCCTCTCATTTTCATGAACCTCGATAATGAAAAGGCAAATGAT
CTTGAGGGTTGTTGAAACTCATGCTTTTGAAGATGTCTTTGCAGTTGAACAATCCACCATTCAACCT
CCTTATTCATACCTCCCCATTTCAAGTAGATCCCTCATATGCTCCTTCCACACACTGGTTTTTACAAA
TAGCTCCCCATTTGAGTGGAGTTGGTGGATTTCGAGATAGCAACCGGATGTCACATAAATCCTGTTTTT
CCAGAGGATGCTGCCAAAATTCAGAGGATGTGAGAATCCTTAATTAACAGTATTTAGGGGGAATCTA
TTAAATCTAATTATTCATATCTGTAGACACAGTAACTAAGGAGATAAAGAGGATGTGGATACAGGGAA
GATGTACCCTTTTTATGCGGAGTTATGTGTAGATTTCGATTTTGTGATCCATTACAAATGTTATCTTAT
GTTACACGGTCATTTTACGCAAAAAATAATTCTCCCTCCATTTCAATTTACCTATTTTCTTTTGTAGT
CTATGCCAACAAAATGACCACTTTCTTTATTTGGA

Figure 5

Nitab4.5_0002825g0060.2 amino acid sequence: (SEQ ID No: 3)

MAEAEAPNRSPEIRKDRIHNRWVLFSPARSRRPSDFKAKSNPQPNKTECPFCAGHEHECAPEIFRVP
ANSTADWKIRVIQNLFPAVSRELDQNPVSALGDVAVSGFGFHDVVIESPVHSVNMSDLSPSQVGEVM
LAFKKRVEQLQNYDNVYVQVFNHGSAGASMSHSHSQMIALPIVPPTVSARLDSMREYYKQTKCS
LCDIQPNELLIDQSAHFISLVPFAATFAFEIWIIPRDHSSHFEHLDNEKANDLGGLLKLMLLKMSLQL
NNPPFNLLIHTSPFQVDPSYAPSTHWFLQIAPHLSGVGGFEIATGCHINPVFPEDAANKILRDVRIIN

Figure 6

Identifier	Nic 1 ERF - Gene
Nitab4.5_0003090g0020.1	<i>ERF17L3ΔN</i>
Nitab4.5_0003090g0030.1	<i>ERF199</i>
Nitab4.5_0003665g0040.1	<i>JRE5L2</i>
Nitab4.5_0004620g0010.1	<i>ERF210</i>
Nitab4.5_0004620g0030.1	<i>ERF91</i>
Nitab4.5_0004620g0080.1	<i>ERF29</i>
Nitab4.5_0004620g0090.3	<i>ERF130</i>
Nitab4.5_0004620g0095.1	<i>ERF16</i>
Nitab4.5_0006382g0040.1	<i>ERF110</i>

Figure 7

Identifier	Nic 2 ERF - Gene
Nitab4.5_0002924g0010.1	ERF17LI
Nitab4.5_0002924g0020.2	ERF179
Nitab4.5_0002924g0040.2	ERF17
Nitab4.5_0002924g0045.1	ERF168
Nitab4.5_0002924g0050.2	ERF115
Nitab4.5_0006499g0010.1	ERF104
Nitab4.5_0006499g0020.2	ERF221
Nitab4.5_0012667g0020.2	ERF91L1
Nitab4.5_0015055g0010.2	ERF189

Figure 8**Nitab4.5_0003090g0030.1 (ERF199)**

SEQ ID No. 7. >Nitab4.5_0003090g0030.1 720 residues [genomic]
ATGGCAATGGAAATGAATCCAGCTGACGAAACCTTGTTTTCTCCGACTCTCATCTCCTTGAATCGATAAAGCAA
CATCTTCTTGACGATTCAGATTTTTCTGAAATTTTTTCGTCGATGAATTCTAGCAACGAAATATTGCCTAACAGT
CCTAGCTCAAGTTTTAGCAGCTTCGACTTCGACTGCAGCTTCCTTAATTGGGATGAAAACCTCTGAGGAAACATTA
ATACCAACTGATCAGAATCCTTCACATGAATCCCATGAAAAGTACTCCGAGTCCGAGGAGAAAACCCAGGGCCCT
GGGGTGGCGCGTGAGAAAAACGCGCCGCGAGATTGGACGCGGTACATAGGAGTGAAACGGCGACCGTGGGGGACG
TTTTCGGCGGAGACAAGAGACCCAAGTAGGAAAGGTGAAGGTGCAAGGCTGTGGTTAGGAACTTACGAGACCGCA
GAGGATGCAGCGTTAGCTTACGATCAAGCCGCTTTCAAATCCGCGGCTCGAGAGCTCGGCTCAATTTTCCTCAC
TTAATCGGCTCAAACATGCCTAAGCCGGCTAGAGTTACAGCGAGGCGTAGTCGTACGCGCTCACCCGAGCCATCG
TCTTCTTCATCCACCTCATCATCAGAAAATGTACCAAGGAAAAGGAATATAGATGTGATAAATTCATAGCCAAA
GCCAAATTCCTTTGTCATAGCTTGAATTTACAGAGATTAGCTTAA

Figure 9

SEQ ID No. 8. >Nitab4.5_0003090g0030.1 720 residues [cdna]
ATGGCAATGGAAATGAATCCAGCTGACGAAACCTTGTTTTCTCCGACTCTCATCTCCTTGAATCGATAAAGCAA
CATCTTCTTGACGATTCAGATTTTTCTGAAATTTTTTCGTCGATGAATTCTAGCAACGAAATATTGCCTAACAGT
CCTAGCTCAAGTTTTAGCAGCTTCGACTTCGACTGCAGCTTCCTTAATTGGGATGAAAACCTCTGAGGAAACATTA
ATACCAACTGATCAGAATCCTTCACATGAATCCCATGAAAAGTACTCCGAGTCCGAGGAGAAAACCCAGGGCCCT
GGGGTGGCGCGTGAGAAAAACGCGCCGCGAGATTGGACGCGGTACATAGGAGTGAAACGGCGACCGTGGGGGACG
TTTTCGGCGGAGACAAGAGACCCAAGTAGGAAAGGTGAAGGTGCAAGGCTGTGGTTAGGAACTTACGAGACCGCA
GAGGATGCAGCGTTAGCTTACGATCAAGCCGCTTTCAAATCCGCGGCTCGAGAGCTCGGCTCAATTTTCCTCAC
TTAATCGGCTCAAACATGCCTAAGCCGGCTAGAGTTACAGCGAGGCGTAGTCGTACGCGCTCACCCGAGCCATCG
TCTTCTTCATCCACCTCATCATCAGAAAATGTACCAAGGAAAAGGAATATAGATGTGATAAATTCATAGCCAAA
GCCAAATTCCTTTGTCATAGCTTGAATTTACAGAGATTAGCTTAA

Figure 10

SEQ ID No. 9. >Nitab4.5_0003090g0030.1 720 residues [cds]
ATGGCAATGGAAATGAATCCAGCTGACGAAACCTTGTTTTCTCCGACTCTCATCTCCTTGAATCGATAAAGCAA
CATCTTCTTGACGATTCAGATTTTTCTGAAATTTTTTCGTCGATGAATTCTAGCAACGAAATATTGCCTAACAGT
CCTAGCTCAAGTTTTAGCAGCTTCGACTTCGACTGCAGCTTCCTTAATTGGGATGAAAACCTCTGAGGAAACATTA
ATACCAACTGATCAGAATCCTTCACATGAATCCCATGAAAAGTACTCCGAGTCCGAGGAGAAAACCCAGGGCCCT
GGGGTGGCGCGTGAGAAAAACGCGCCGCGAGATTGGACGCGGTACATAGGAGTGAAACGGCGACCGTGGGGGACG
TTTTCGGCGGAGACAAGAGACCCAAGTAGGAAAGGTGAAGGTGCAAGGCTGTGGTTAGGAACTTACGAGACCGCA
GAGGATGCAGCGTTAGCTTACGATCAAGCCGCTTTCAAATCCGCGGCTCGAGAGCTCGGCTCAATTTTCCTCAC
TTAATCGGCTCAAACATGCCTAAGCCGGCTAGAGTTACAGCGAGGCGTAGTCGTACGCGCTCACCCGAGCCATCG
TCTTCTTCATCCACCTCATCATCAGAAAATGTACCAAGGAAAAGGAATATAGATGTGATAAATTCATAGCCAAA
GCCAAATTCCTTTGTCATAGCTTGAATTTACAGAGATTAGCTTAA

Figure 11

SEQ ID No. 10. >Nitab4.5_0003090g0030.1 239 residues [peptide]
MAMEMNPADETLFFSDSHLLESIKQHLLDDSDFSEIFSSMNSSNEILPNSPSSSFSSSFDSDCSFLNWDENSEETL
IPTDQNPESHESHEKYSEEEKTQGPVAREKNAPRDWTRYIGVKRRPWGTFSAETRDPSRKGEARLWLGTYETA
EDAALAYDQAFAKIRGSRARLNFPHLIGSNMPKPARVTARRSRTSRPEPSSSSSTSSSENVPKRKNIDVINSIAK
AKFLCHSLNLQRLA

Figure 12**Nitab4.5_0015055g0010.2 (ERF189)**

SEQ ID No. 11. >Nitab4.5_0015055g0010.2 699 residues [genomic]
ATGGAAATGAATCTAGCTGACGAAACCTTGTTTTCTCTGAGTCTCATCTCCTTGAATCGATAAAGCAACATCTT
CTTGATGATTAGATTTTTCTGAAATTTTTTCGCCGATGAGTTCAAGCAACGAAATATTGCCTAACAGTCTTAGC
TCAAGTTTTAGCAGCTTCGACTGCAGCTTCCTCAATTGGGATGAAAACCTTGAGGAAACATTAATACCAACTGAT
CAAAATCCTTCACATGAGAAGTGCTCCGAGTCCGAGGAGCAAACCCAGGGCCCAGCGGTGGTGCGTGAGAAAAAC
GCGCCGCGAGATTGGACGCGGTATATAGGAGTGAAACGGCGGCCGTGGGGGACGTTTTTCGGCGGAGACAAGAGAC
CCAAGTAGGAAAGGTGAAGGTGCAAGGCTGTGGTTAGGAACTTACGAGACCGCAGAGGATGCAGCGTTGGCTTAC
GATCAAGCCGCTTTCAAATCCGCGGCTCGAGAGCTCGGCTCAATTTTCCCACTTAATTGGCTCAAACATGCCT
AAGCCGGCTAGAGTAACAGCGAGGCGTAGTCGTACGCGCTCACCCGAGCCATCCTCTTCTTCATCCACTTCATCA
TCAGAAAATGTGCCAAGAAAAAGGAATATAGATGTGATAAATTCATAGCCAAAGCCAAATTCCTTTGTCATAGC
T

Figure 13

SEQ ID No. 12. >Nitab4.5_0015055g0010.2 699 residues [cdna]
 ATGGAAATGAATCTAGCTGACGAAACCTTGTTTTCTCTGAGTCTCATCTCCTTGAATCGATAAAGCAACATCTT
 CTTGATGATTCAGATTTTTCTGAAATTTTTTCGCCGATGAGTTCAAGCAACGAAATATTGCCTAACAGTCCTAGC
 TCAAGTTTTAGCAGCTTCGACTGCAGCTTCCTCAATTGGGATGAAAACCTTGAGGAAACATTAATACCAACTGAT
 CAAAATCCTTCACATGAGAAGTGCTCCGAGTCCGAGGAGCAAACCCAGGGCCCAGCGGTGGTGCGTGAGAAAAAC
 GCGCCGCGAGATTGGACGCGGTATATAGGAGTGAAACGGCGGCCGTGGGGGACGTTTTTCGGCGGAGACAAGAGAC
 CCAAGTAGGAAAGGTGAAGGTGCAAGGCTGTGGTTAGGAACCTTACGAGACCGCAGAGGATGCAGCGTTGGCTTAC
 GATCAAGCCGCTTTCAAAATCCGCGGCTCGAGAGCTCGGCTCAATTTTCCCCACTTAATTGGCTCAAACATGCCT
 AAGCCGGCTAGAGTAACAGCGAGGCGTAGTCGTACGCGCTCACCAGGCCATCCTCTTCTTCATCCACTTCATCA
 TCAGAAAATGTGCCAAGAAAAAGGAATATAGATGTGATAAATTCATAGCCAAAGCCAAATTCCTTTGTCATAGC
 TTAAATTTACAGAGATTAGCTTAA

Figure 14

SEQ ID No. 13. >Nitab4.5_0015055g0010.2 699 residues [cds]
 ATGGAAATGAATCTAGCTGACGAAACCTTGTTTTCTCTGAGTCTCATCTCCTTGAATCGATAAAGCAACATCTT
 CTTGATGATTCAGATTTTTCTGAAATTTTTTCGCCGATGAGTTCAAGCAACGAAATATTGCCTAACAGTCCTAGC
 TCAAGTTTTAGCAGCTTCGACTGCAGCTTCCTCAATTGGGATGAAAACCTTGAGGAAACATTAATACCAACTGAT
 CAAAATCCTTCACATGAGAAGTGCTCCGAGTCCGAGGAGCAAACCCAGGGCCCAGCGGTGGTGCGTGAGAAAAAC
 GCGCCGCGAGATTGGACGCGGTATATAGGAGTGAAACGGCGGCCGTGGGGGACGTTTTTCGGCGGAGACAAGAGAC
 CCAAGTAGGAAAGGTGAAGGTGCAAGGCTGTGGTTAGGAACCTTACGAGACCGCAGAGGATGCAGCGTTGGCTTAC
 GATCAAGCCGCTTTCAAAATCCGCGGCTCGAGAGCTCGGCTCAATTTTCCCCACTTAATTGGCTCAAACATGCCT
 AAGCCGGCTAGAGTAACAGCGAGGCGTAGTCGTACGCGCTCACCAGGCCATCCTCTTCTTCATCCACTTCATCA
 TCAGAAAATGTGCCAAGAAAAAGGAATATAGATGTGATAAATTCATAGCCAAAGCCAAATTCCTTTGTCATAGC
 TTAAATTTACAGAGATTAGCTTAA

Figure 15

SEQ ID No. 14.
 >Nitab4.5_0015055g0010.2 232 residues [peptide]
 MEMNLADETLFFSESHLLESIKQHLLDDSDSEIFSPMSSSNEILNPSPPSSSFSSFDCSFLNWDENFEETLIPTD
 QNPSHEKCESEEEQTQGPVAVREKNAPRDWTRYIGVKRPPWGTFSAETRDPSRKGEARLWLGTYETAEDAALAY
 DQAAFKIRGSRARLNFPHLIGSNMPKPARVTARRSRTRSPEPSSSSSTSSSENVPKRKNIDVINSIAKAKFLCHS
 LNLQRL

Figure 16

TRV1: SEQ ID No. 15
 GATGGTAAGAAGAAAGGCGCGCAGTATGCGATAGCTCTTCACAGCCTGTATGACTTCAAGTTGAAAGACTTGATG
 GCTACTATGGTTGAGAAGAAAACCTAAAGTGGTTTCATGCTGCTATGCTTTTGGCTCCTGAAAGTATGTTAGTGGAC
 GAAGGTCCATTACCTTCTGTTGACGGTTACTACATGAAGAAGAACGGGAAGATCTATTTCCGGTTTTGAGAAAGAT
 CCTTCCTTTTCTTACATTTCATGACTGGGAAGAGTACAAGAAGTATCTACTGGGGAAGCCAGTGAGTTACCAAGGG
 AATGTGTCTACTTCGAACCGTGGCAGGTGAGAGGAGACACAATGCTTTTTTCGATCTACAGGATAGCTGGAGTT
 CCGAGGAGGTCTCTATCATCGCAAGAGTACTACCGAAGAATATATATCAGTAGATGGGAAAGCATGGTTGTTGTC
 CCAATTTTTCGATCTGGTTCGAATCAACGCGAGAGTTGGTCAAGAAAGACCTGTTTGTAGAGAAACAATTCATGGAC
 AAGTGTTTGGATTACATAGCTAGGTTATCTGACCAGCAGCTGACCATAAGCAATGTTAAATCATACTTGAGTTCA
 AATAATTGGGTCTTATTCATAAACGGGGCGGCCGTGAAGAACAAGCAAAGTGATAGATTCTCGAGATTTACAGTTG
 TTGGCTCAAACCTTTGCTAGTGAAGGAACAAGTGGCGAGACCTGTCTGAGGGAGTTGCGTGAAGCAATTCGACT
 GAGACGAAACCTATCACGTCATTGACTGATGTGCTGGGTTTAAATATCAAGAAAACCTGTGGAAGCAGTTTGCTAAC
 AAGATCGCAGTCGGCGGATTTCGTTGGCATGGTTGGTACTCTAATTGGATTCTATCCAAAGAAGGTACTAACCTGG
 GCGAAGGACACACCAAATGGTCCAGAACTATGTTACGAGAACCTCGCACAAAACCAAGGTGATAGTATTTCTGAGT
 GTTGTGTATGCCATTGGAGGAATCACGCTTATGCGTCGAGACATCCGAGATGGACTGGTGAAAAAACTATGTGAT
 ATGTTTGATATCAAACGGGGGGGCCATGTCTTAGACGTTGAGAATCCGTGCCGCTATTATGAAATCAACGATTTT
 TTTAGCAGTCTGTATTCGGCATCTGAGTCCGGTGAGACCGTTTTTACCAGATTTATCCGAGGTAAGGCAAGTCT
 GATAAGCTATTGCAGCAGAAGAAAGAAATCGCTGACGAGTTTCTAAGTGCAAAATTCCTCTAACTATTCTGGCAGT
 TCGGTGAGAACTTCTCCACCATCGGTGGTTCGTTTCATCTCGAAGCGGACTGGGTCTGTTGTTGGAAGACAGTAAC
 GTGCTGACCCAAGCTAGAGTTGGAGTTTCAAGAAAGGTAGACGATGAGGAGATCATGGAGCAGTTTCTGAGTGGT
 CTTATTGACACTGAAGCAGAAATTGACGAGGTTGTTTTCAGCTGAAATGTGAAAGAGGGGAAACAAGC
 GTACAAAAGGTGTTGTGTAACCTTTAACGCCACAGGATTTGAGAACGTGTTGCCAGCTTGCAAAACCTTTGGTC
 AGCAAAGGAAAAACGGTCAACCGTTCGATTACTTCAAGTGATGGGAGGTGAGAGATTACCAAAAAGGCCGPTT
 GTCAGTGGAGACGATTCTGTGGACGCTAGAAGAGAGTTTCTGTACTACTTAGATGCGGAGAGAGTTCGCTCAAAT

GATGAAATTATGTCTCTGTATCGTGACTATTCGAGAGGAGTTATTCGAACTGGAGGTCAGAATTACCCGCACGGA
CTGGGAGTGTGGGATGTGGAGATGAAGAACTGGTGCATACGTCCAGTGGTCACTGAACATGCTTATGTGTTCCAA
CCAGACAAACGTATGGATGATTGGTCGGGATACTTAGAAGTGGCTGTTTGGGAACGAGGTATGTTGGTCAACGAC
TTCGCGGTGCGAAAGGATGAGTGATTATGTCATAGTTTGGCATCAGACGTATCTTTGCAATAACAGGTAATAATCC
TCTCTCTTGATATTTTTAAATTATAGAATTAATTAGTTTACTTTATTTCTTTACTATATGATTTAAATAGTTTAAAT
CTTGTTTTTGGAGTAACTATTCGATTTTGATATTTGTATTCGTCTTACAAAGTTGGAAATACATGATGATATTTTC
TTTTGAACGTGATACCTACCAATACTAATCTTACGGAATCTTTTAAATAGAGCACTAATCAACATGGAACTAAAGA
CCAATTCCTTAAGTGTCTCTGTGTACAGTTTCATTTTAGTAGTGGCTTAAAGTATTATTATCTCCCTTCATGCGGG
GCAATTATGTAGATTAAAATCGAAATTATATAAAATTTACATAAGTCTAAGTCTAGGGTCTCCAGCTAATTGTTA
TTTTTTTAAACGATGTTGACTAAAGCAATAACGACGTTGACTTGTGTAAACAGGTTGATCTTGGACAATTTAAGT
GCCCTGGATCTAGGACCAGTTAACTGTTCTTTTGAATTAGTTGACGGTGTACCTGGTTGTGGTAAGTCGACAATG
ATTGTCAACTCAGCTAATCCTTGTGTGCGATGTGGTTCTCTCTACTGGGAGAGCAGCAACCGACGACTTGATCGAG
AGATTGCGCAGCAAAGGTTTTCCATGCAAATTGAAAAGGAGAGTGAAGACGGTTGATTCTTTTTTGTGTCATTGT
GTCGATGGTTCTTTAACCGGAGACGTGTTGCATTTTCGACGAAGCTCTCATGGCCCATGCTGGTATGGTGACTTT
TGCGCTCAGATAGCTGGTGCTAAACGATGTATCTGTCAAGGAGATCAGAATCAAATTTCTTTCAAGCCTAGGGTA
TCTCAAGTTGATTTGAGGTTTTCTAGTCTGGTCGGAAAGTTTGACATTGTTACAGAAAAAGAGAACTTACAGA
AGTCCAGCAGATGTGGCTGCCGATTTGAACAAGTACTTACTGAGATGTGAGAACACATAACCGGACTGCTAAT
TCGATGACGGTGAGGAAGATTGTGTCTAAAGAACAGGTTTCTTTGAAGCCTGGTGCTCAGTACATAACTTTCCCTT
CAGTCTGAGAAGAAGGAGTTGGTAAATTTGTTGGCATTGAGGAAAGTGGCAGCTAAAGTGAGTACAGTACACGAG
TCGCAAGGAGAGACATTCAAAGATGTAGTCTTAGTCAGGACGAAACCTACGGATGACTCAATCGCTAGAGGTCGG
GAGTACTTAATCGTGGCATTGTGCGCTCACACACAATCACTTGTGTATGAACTGTGAAAGAGGACGATGTAAGC
AAAGAGATCAGGGAAAGTGCCGCGCTTACGAAGGCGGCTTTGGCAAGATTTTTTGTACTGAGACCGTCTTATGA
CGGTTTTCGGTCTAGGTTTGTATGCTTTAGACATCATGAAGGGCTTGCCTGGTCCAGATTGAGTACGATTACG
GACTTGGAGATGTGGTACGACGCTTGTGTTCCGGGAAATTCGTTAAGAGACTCAAGCCTAGACGGGTATTTGGTG
GCAACGACTGATTGCAATTTGCGATTAGACAATGTTACGATCAAAAGTGGAACTGGAAAGACAAGTTTGCTGAA
AAAGAAACGTTTTCTGAAACCGGTTATTCGTACTGCTATGCCTGACAAAAGGAAGACTACTCAGTTGGAGAGTTTG
TTAGCATTGCGAAAAAGGAACCAAGCGGCACCCGATCTACAAGAAAATGTGCACGCGACAGTTCTAATCGAAGAG
ACGATGAAGAAGCTGAAATCTGTTGTCTACGATGTGGGAAAAATTCGGGCTGATCCTATTGTCAATAGAGCTCAA
ATGGAGAGATGGTGGAGAAAATCAAAGCACAGCGGTACAGGCTAAGGTAGTAGCAGATGTGAGAGAGTTACATGAA
ATAGACTATTCTGCTTTACATGTATATGATCAAATCTGACGTGAAACCTAAGACTGATTTAACACCGCAATTTGAA
TACTCAGCTCTACAGACTGTTGTGTATCACGAGAAGTTGATCAACTCGTTGTTGGTCCAATTTTCAAAGAAATTT
AATGAACGCAAGTTGGATGCTATGCAACCACATTTTGTGTTCAACACGAGAATGACATCGAGTGATTTAAACGAT
CGAGTGAAGTTCTTAAATACGGAAGCGGCTTACGACTTTGTTGAGATAGACATGTCTAAATTCGACAAGTCGGCA
AATCGCTTCCATTTACAACCTGCAGCTGGAGATTTACAGGTTATTTGGGCTGGATGAGTGGGCGGCTTCCTTTGG
GAGGTGTGCGCACTCAAACCTACTGTGAGAGATATTCAAATGGTATGATGGCGCATATTTGGTACCAACAAAAG
AGTGGAGATGCTGATACCTTATAATGCAAATTCAGATAGAACACTGTGTGCACTCTTGTCTGAATTACCATTGGAG
AAAGCAGTCATGGTTACATATGGAGGAGATGACTCACTGATTGCGTTTCTTAGAGGAACGCAGTTTGTGATCCG
TGTCCAAAGTTGGCTACTAAGTGAATTTTCGAGTGCAAGATTTTAAAGTACGATGTCCCAATGTTTTGTGGGAAG
TTCTTGTCTTAAGACGTCATCGTGTACGAGTTCTGTCGGATCCGGTAAAGTTCTGACGAAGTTGGGGAAAAAG
AGTATAAAGGATGTGCAACATTTAGCCGAGATCTACATCTCGCTGAATGATTTCAATAGAGCTCTGGGAACTAC
ATGGTGGTATCCAACTGTCCGAGTCTGTTTCAGACCGGTATTTGTACAAAGGTGATTCTGTTTCATGCGCTTTGT
GCGCTATGGAAGCATATTAAGAGTTTTACAGCTCTGTGTACATTATTCGAGACGAAAACGATAAGGAATGAAC
CCGGCTAAGGTTGATTGGAAGAAGGCACAGAGAGCTGTGTCAAACCTTTACGACTGGTAATATGGAAGACAAGTC
ATTGGTCACCTTGAAGAAGAAGACTTTCGAAGTCTCAAATTTCTCAAATCTAGGGGCCATTGAATTGTTGTGGA
CGGTAGGAGGAAGAGACCGAAGTATTTTACAGAAGAAGAGAACTGTCTAAATCATGTTGGTGGGAAGAAGAG
TGAACACAAGTTAGACGTTTTTGGACAAAGGGATTACAAAATGATTAATCTTACGCGTTTTCTAAAGGTAGTAGG
TGTACAACTAGTTGTAACATCACATCTACCTGCAGATACGCTGGGTTCAATCAAATCGATCTGTTGGATTTCGAG
ACTTACTGAGAAAAGAAAGAGAGGAAAGACTATTTCAGAGATTCAAAGCTCGAGCTTGCGATAACTGTTTCAGTTGC
GCAGTACAAGTTGAATACAGTATTTCCACACAGGAGAACCTACTTGTATGTCTGGAAGGTGGGTGTATTTCTGA
GGGCGTTCGGGTCTGTGACGGTACATACCTTTTCAGTATCGAAGTGTGCTGCTAATATGGGTGCTACTGATTTCGAC
TAGGCGCCTCAATGTGGAAGAAGCTGAACAGTTTCGGATTACATTGAAGGCGATTTTACCGATCAAGAGGTTTTCGG
TGAGTTCATGTCTTTGAAACAAGTGGAGATGAAGACGATTGAGGCGAAGTACGATGGTCTTACAGACCAGCTAC
TACTAGACCTAAGTCATTATTGTCAAGTGAAGATGTTAAGAGAGCGTCTAATAAGAAAACTCGTCTTAATGCAT
AAAGAAATTTATTGTCAATATGACGTGTGACTCAAGGGTTGTGTGAATGAAGTCACTGTTCTTGGTCACGAGAC
GTGTAGTATCGGTTCATGCTAACAAATTCGGAAGCAAGTTGCTGACATGGTGGTGTACACGTAGGTGTGCGGA
AAATAATTGTGGATGGTTTTGTGTGTGTTATCAATGATTTTACTTTTGATGTGTATAATTGTTGTGCGCGTAG
TCACCTTGAAAAGTGTGCTAAACGTGTTGAAACAAGAAATCGAGAAATTTGGAAACAAATTCGACGAAATCAAGC
TGAAAACATGTCTGCGACAGCTAAAAAGTCTCATAATTCAAGACCTCTAAGAAGAAATTCAAAGAGGACAGAGA
ATTTGGGACACCAAAAAGATTTTTAAGAGATGATGTTCTTTTCGGGATTGATCGTTTGTGTTTTGATTTTA
TTTTATATTGTTATCTGTTTCTGTGTATAGACTGTTTGAGATTGGCGCTTGGCCGACTCATTGCTTACCATAGG
GGAACGGACTTTGTTTGTGTTGTTATTTTATTTGATTTTATTTAAATTTCTCAATGATCTGAAAAGGCCCTCGAGG

CTAAGAGATTATTGGGGGGTGAGTAAGTACTTTTAAAGTGATGATGGTTACAAAGGCAAAAGGGGTAAAACCCCT
CGCCTACGTAAGCGTTATTACGCCCCGATCCCCCGGGAGCTCGAATTCGCTGAAATCACCAGTCTCTCTCTACA
AATCTATCTCTCTCTATTTTTTCCATAAATAATGTGTGAGTAGTTTCCCGATAAGGGAAATTAGGGTTCTTATAG
GGTTTCGCTCATGTGTTGAGCATATAAGAAACCCCTTAGTATGTATTTGTATTTGTAAAATACTTCTATTATCAAT
AAAATTTCTAATTCCTAAACCAAAATCCAGTACTAAAATCCAGATCTCCTAAAGTCCCTATAGATCTTTGTCTGT
GAATATAAACACGACACGAGACGACTAAACCTGGAGCCCAGACGCCGTTTCAAGCTAGAAGTACCCTTAGGCAG
GAGGCCGTTAGGGAAAAGATGCTAAGGCAGGGTTGGTTACGTTGACTCCCCCGTAGGTTTGGTTTAAATATGATG
AAGTGGACGGAAGGAAGGAGACAAGGAAGGATAAGGTTGCAGGCCCTGTGCAAGGTAAGAAGATGGAAATTT
TGATAGAGGTACGCTACTATACTTATACTATACTAAGGGAATGCTTGTATTTATACCCTATACCCCCTAATAA
CCCCCTTATCAATTTAAGAAATAATCCGCATAAGCCCCCGCTTAAAAATTTGGTATCAGAGCCATGAATAGGTCTAT
GACCAAACTCAAGAGGATAAAACCTCACCAAAATACGAAAGAGTTCTTAACTCTAAAGATAAAAGATCTTTCAA
GATCAAACTAGTTCCCTCACACCGGAGCATGCGATATCCTCGACCTGCAGGCATGCAAGCTTGGCGTAATCATG
GTCATAGCTGTTTCTGTGTGAAATTGTTATCCGCTCACAATTCACACAACATACGAGCCGGAAGCATAAAGTG
TAAAGCCTGGGGTGCCATATGAGTGAGCTAACTCACATTAATTGCGTTGCGCTCACTGCCCCGCTTCCAGTCGGG
AAACCTGTCTGTGCCAGCTGCATTAATGAATCGGCCAACGCGCGGGGAGAGGCGGTTTGCCTATTGGGCCAAAGAC
AAAAGGGCGACATTCAACCGATTGAGGGAGGGAAGGTAATATTGACGGAAATTATTATTAAAGGTGAATTATC
ACCGTCACCGACTTGAGCCATTGCGAATTAGAGCCAGCAAAATCACCAGTAGCACCATTACCATTAGCAAGGCC
GGAAACGTCACCAATGAAACCATCGATAGCAGCACCCTAATCAGTAGCGACAGAATCAAGTTTGCCTTTAGCGTCT
AGACTGTAGCGCGTTTTTCATCGGCATTTTTCGGTTCATAGCCCCCTTATTAGCGTTTGGCATCTTTTTCATAACAAA
ATCACCGGAACAGAGCCACCACCGGAACCGCCTCCCTCAGAGCCGCCACCCTCAGAACCGCCACCCTCAGAGCC
ACCACCCTCAGAGCCGCCACCAGAACCACCAGAGCCGCCGCCAGCATTGACAGGAGGCCCGATCTAGTAACA
TAGATGACACCGCGCGCGATAATTTATCCTAGTTTTCGCGCTATATTTTGTTCCTATCGCGTATTAAATGTATA
ATTGCGGGACTCTAATCATAAAAACCCATCTCATAAATAACGTCATGCATTACATGTTAATTATTACATGCTTAA
CGTAATTCAACAGAAATTATATGATAATCATCGCAAGACCGGCAACAGGATTCAATCTTAAGAACTTTATTGCC
AAATGTTTTGAACGATCGGGGATCATCCGGGTCTGTGGCGGGAATCCACGAAAATATCCGAACGCAGCAAGATAT
CGCGGTGCATCTCGGTCTTGCTTGGCAGTCGCCGCCGACGCCGTTGATGTGGACGCCGGGCCGATCATATTGT
CGCTCAGGATCGTGGCGTTGTGCTTGTTCGGCCGTTGCTGTCTGAATGATATCGGCACCTTCGACCGCCTGTTCGG
CAGAGATCCCCGTGGCGAAGAACTCCAGCATGAGATCCCCGCGCTGGAGGATCATCCAGCCGCGCTCCCGAAAA
CGATTCCGAAGCCCAACCTTTTCATAGAAGGCGCGGTGGAATCGAAATCTCGTGATGGCAGGTTGGGCGCTCGCTT
GGTCCGTCATTTTCAACCCAGAGTCCCGCTCAGAAGAACTCGTCAAGAAGGCGATAGAAGGCGATGCGCTGCGA
ATCGGGAGCGCGATACCGTAAAGCACGAGGAAGCGGTACGCCCATTCGCCGCCAAGCTCTTCAGCAATATCACG
GGTAGCCAACGCTATGTCTGATAGCGGTCCGCCACACCCAGCCGCCACAGTCGATGAATCCAGAAAAGCGGCC
ATTTTCCACCATGATATTCGGAAGCAGGCATCGCCATGGGTACGACGAGATCATCGCCGTCGGGCATGCGCGC
CTTGAGCCTGGCGAACAGTTTCGGCTGGCGCGAGCCCTGATGCTCTTCGTCCAGATCATCCTGATCGACAAGACC
GGCTTCCATCCGAGTACGTGCTCGCTCGATGCGATGTTTCGTTGGTGGTCAATGGGCAGGTAGCCGATCAAG
CGTATGCAGCCCGCGCATTGCATCAGCCATGATGGATACTTTCTCGGCAGGAGCAAGGTGAGATGACAGGAGATC
CTGCCCCGGCACTTCGCCCCAATAGCAGCCAGTCCCTTCCCGCTTCAGTGACAACGTCGAGCACAGCTGCGCAAGG
AACGCCCGTCTGTGGCCAGCCACGATAGCCCGCTGCTCGTCTGCTGAGTTCATTTCAGGGCACCAGGAGGTGCGT
CTTGACAAAAGAACCAGCGCGCCCTGCGCTGACAGCCGGAACACGCGCGCATCAGAGCAGCCGATGTCTGTTG
TGCCAGATCATAGCCGAATAGCCTTCCACCAAGCGCGGAGAACCTGCGTGCAATCCATCTTGTTCATCAT
GCGAAACGATCCAGATCCGGTGCAGATTATTTGGATTGAGAGTGAATATGAGACTCTAATTGGATACCGAGGGGA
ATTTATGGAACGTCAGTGGAGCATTTTTGACAAGAAATATTTGCTAGCTGATAGTGACCTTAGGCGACTTTTGAA
CGCGCAATAATGGTTTTCTGACGTATGTGCTTAGCTCATTAACCTCCAGAAACCCGCGGCTGAGTGGCTCCTTCAA
CGTTGCGGTTCTGTGCTTCCAAACGTAACCGGCTTGTCGCCGCTCATCGCGGGGGTTCATAACGTGACTCCCT
TAATTCTCCGCTCATGATCAGATTGTCGTTTCCCGCCTTCAGTTTAAACTATCAGTGTTTGACAGGATATATTGG
CGGGTAAACCTAAGAGAAAAGAGCGTTTATTAGAATAATCGGATATTTAAAGGGCGTGAAAAGGTTTATCCGTT
CGTCCATTTGTATGTGCATGCCAACACAGGGTTCCCCAGATCTGGCGCCGCCAGCGAGACGAGCAAGATTGGC
CGCCGCCGAAACGATCCGACAGCGCGCCAGCACAGGTGCGCAGGCAAAATTGCACCAACGCATACAGCGCCAGC
AGAATGCCATAGTGGCGGTGACGTCTGAGTGAACACAGATCCGCGCAGGAGGCCCGGACACCCGCGCATAATC
AGGCCGATGCCGACAGCTCGAGCGGACAGTGCTCAGATCAGGGGTATGTTGGTTTACGCTTACGCTGGC
CTCCGGACAGCCTCCGCTGGTCCGATTGAACGCGCGGATTCTTTATCACTGATAAGTTGGTGGACATATTATGT
TTATCAGTGATAAAGTGTCAAGCATGACAAAGTTGCAGCCGAATACAGTGATCCGTGCCGCCCTGGACCTGTTGA
ACGAGGTGCGCGTAGACGGTCTGACGACACGCAAACTGGCGGAACGGTTGGGGGTTTCAGCAGCCGGCGCTTTACT
GGCACTTCAGGAACAAGCGGGCGCTGCTCGACGCACTGGCCGAAGCCATGCTGGCGGAGAATCATACGCATTCCG
TGCCGAGAGCCGACGACTGGCGCTCATTTCTGATCGGGAATGCCCGCAGCTTCAGGCAGGCGCTGCTCGCCT
ACCGCGATGGCGCGCGCATCCATGCCGGCACGCGACCGGGCGCACCGCAGATGGAAACGGCCGACGCGCAGCTTC
GCTTCTCTGCGAGGCGGGTTTTTCGGCCGGGGACGCCGTCAATGCGCTGATGACAATCAGCTACTTCACTGTTG
GGGCCGTGCTTGAGGAGCAGGCCGGCGACAGCGATGCCGGCGAGCGCGGCGGCACCGTTGAACAGGCTCCGCTCT
CGCCGCTGTTGCGGGCCGCGATAGACGCCCTTCGACGAAGCCGGTCCGGACGACGCTTCGAGCAGGACTCGCGG
TGATTGTCGATGGATTGGCGAAAAGGAGGCTCGTTGTGAGGAACGTTGAAGGACCGAGAAAGGTTGACGATTGAT
CAGGACCGCTGCCGAGCGCAACCCACTCACTACAGCAGAGCCATGTAGACAACATCCCCCTCCCCCTTCCACCG

CGTCAGACGCCCGTAGCAGCCCGCTACGGGCTTTTTTCATGCCCTGCCCTAGCGTCCAAGCCTCACGGCCGCGCTC
GGCCTCTCTGGCGGCCCTTCTGGCGCTCTTCCGCTTCCTCGCTCACTGACTCGCTGCGCTCGGTGCTTCCGCTGCG
GCGAGCGGTATCAGCTCACTCAAAGGCGGTAATACGGTTATCCACAGAATCAGGGGATAACGCAGGAAAGAACAT
GTGAGCAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCC
CCCTGACGAGCATCACAAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATACCAGGC
GTTTTCCCCCTGGAAGCTCCCTCGTGCGCTCTCTGTTCCGACCTGCGCTTACCGGATACCTGTCCGCCTTTCT
CCCTTCGGGAAGCGTGGCGCTTTTCCGCTGCATAACCCCTGCTTCGGGGTCATTATAGCGATTTTTTCGGTATATC
CATCCTTTTTTCGCACGATATACAGGATTTTGCCAAAGGGTTCGTGTAGACTTTCCTTGGTGTATCCAAACGGCGTC
AGCCGGGCAGGATAGGTGAAGTAGGCCACCCGCGAGCGGGTGTTCCTTCTTCACTGTCCCTTATTTCGCACCTGG
CGGTGCTCAACGGGAATCTGCTCTGCGAGGCTGGCCGGCTACCGCCGGCGTAACAGATGAGGGCAAGCGGATGG
CTGATGAAACCAAGCCAACCAGGAAGGGCAGCCACCTATCAAGGTGTACTGCCTTCCAGACGAACGAAGAGCGA
TTGAGGAAAAGGCGGCGGCGGCCGCGCATGAGCCTGTGCGCTACCTGCTGGCCGTGCGCCAGGGCTACAAAATCA
CGGGCGTCTGAGTATGAGCACGTCCGCGAGCTGGCCCGCATCAATGGCGACCTGGGCGCGCTGGGCGGCTGC
TGAAACTCTGGCTCACCGACGACCCGCGCACGGCGCGGTTGCGGTGATGCCACGATCCTCGCCCTGCTGGCGAAGA
TCGAAGAGAAGCAGGACGAGCTTGGCAAGGTGATGATGGGCGTGGTCCGCCCCAGGGCAGAGCCATGACTTTTTT
AGCCGCTAAAACGGCCGGGGGTGCGCGTGATTGCCAAGCACGTCCCATGCGCTCCATCAAGAAGAGCGACTTC
GCGGAGCTGGTGAAGTACATCACCGACGAGCAAGGCAAGACCGAGCGCCTTTGCGACGCTCACCGGGCTGGTTC
CCTCGCGCTGGGCTGGCGGCGCTCTATGGCCCTGCAACCGCGCAGAAACGCGCTCGAAGCCGTGTGCGAGACA
CCGCGGCGCGCGCGCTTGTGGATACCTCGCGGAAAACCTTGGCCCTCACTGACAGATGAGGGGCGGACCTTGACAC
TTGAGGGGCGGACTCACCGGCGCGCGCTTGACAGATGAGGGGCGGCTCGATTTTGGCCGGCGGACGTGGAGCTG
GCCAGCCTCGCAATCGGCGAAAACGCCTGATTTTACGCGAGTPTCCACAGATGATGTGGACAAGCCTGGGGAT
AAGTGCCCTGCGGTATTGACACTTGAGGGGCGGACTACTGACAGATGAGGGGCGCGATCCTTGACACTTGAGGG
GCAGAGTGCTGACAGATGAGGGGCGCACCTATTGACATTTGAGGGGCTGTCCACAGGCAGAAAATCCAGCATTTG
CAAGGGTTTTCCGCGCGTTTTTCGGCCACCGCTAACCTGTCTTTTAACTGCTTTTAAACCAATATTTATAAACCT
TGTTTTTAACAGGGCTGCGCCCTGTGCGCGTGACCGCGCACGCCGAAGGGGGGTGCCCCCCTTCTCGAACCCCT
CCCGGCCGCTAACGCGGGCTCCCATCCCCCAGGGGTGCGCCCTCGGCCGGAACGGCTCACCCAAAAA
TGGCAGCGCTGGCAGTCTTGCCATTGCGGGGATCGGGGCGTAACGGGATGGGCGATCAGCCGAGCGCGACGC
CCGGAAGCATTTGACGTGCGCGAGGTGCTGGCATCGACATTCAGCGACAGGTGCGGGGAGTGAGGGCGCGCGCC
TGGGTGGCGGCTGCCCTTCACTTCGGCCGTCGGGGCATTCACGGACTTCATGGCGGGGCGGCAATTTTACCT
TGGGCATTCTTGGCATAGTGGTTCGCGGGTGCCGTGCTCGTGTTCGGGGGTGCGATAAACCCAGCGAACCATTTGA
GGTGATAGGTAAGATTATACCGAGGTATGAAAACGAGAATTGGACCTTTACAGAATTACTCTATGAAGCGCCATA
TTTAAAAAGCTACCAAGACGAAGAGGATGAAGAGGATGAGGAGGCAGATTGCCTTGAATATATTGACAATACTGA
TAAGATAATATATCTTTTATATAGAAGATATCGCCGTATGTAAGGATTTTCAAGGGGCAAGGCATAGGCAGCGCGC
TTATCAATATATCTATAGAATGGGCAAAAGCATAAAAACTTGATGGAATAATGCTTGAACCCAGGACAATAACC
TTATAGCTTGTAAATCTATCATAATTGGGTAATGACTCCAACCTTATTGATAGTGTATTTATGTTACAGATAATGCC
CGATGACTTTGTGATGACGCTCCACCGATTTTGAGAACGACAGCGACTTCCGTCCCAGCGGTGCCAGGTGCTGCC
TCAGATTACAGTTATGCCGCTCAATTCGCTGCGTATATCGCTTGCTGATTACGTGCAGCTTTCCCTTCAGGCGGG
ATTCATACAGCGGCCAGCCATCCGTCATCCATATCACCACGTCAAAGGGTGACAGCAGGCTCATAAGACGCCCCA
GCGTCGCCATGATGCGTTACCGAATACGTGCGCAACACCGCTCTTCCGGAGACTGTACACGCTAAACAGCC
AGCGCTGGCGCATTTAGCCCGACATAGCCCACTGTTCGTCCATTTCCGCGCAGACGATGAGCTGCTGCCG
GCTGTATGCGCGAGGTTACCGACTGCGGCCTGAGTTTTTAAAGTGACGTAAAATCGTGTGAGGCCAACGCCCAT
AATGCGGGCTGTTGCCCGCATCCAACGCCATTGATGGCCATATCAATGATTTTCTGGTGCGTACCGGGTTGAGA
AGCGGTGTAAGTGAAGTGCAGTTGCCATGTTTTACGGCAGTGAGAGCAGAGATAGCGCTGATGTCCGGCGGTGCT
TTTGCCGTTACGCACCACCCGTCAGTAGCTGAACAGGAGGACAGCTGATAGACACAGAAGCCACTGGAGCACC
TCAAAAACACCATCATACACTAAATCAGTAAGTTGGCAGCATCACCCATAATTGTGGTTTTCAAAATCGGCTCCGT
CGATACTATGTTATACGCCAACTTTGAAAACAACCTTTGAAAAGCTGTTTTCTGGTATTTAAGGTTTTAGAATGC
AAGGAACAGTGAATTGGAGTTCGTCTTGTTATAATTAGCTTCTTGGGGTATCTTTAAATACTGTAGAAAAGAGGA
AGGAAATAATAAATGGCTAAAATGAGAATATCACCGGAATTGAAAAAACTGATCGAAAAATACCGCTGCGTAAAA
GATACGGAAGGAATGTCTCCTGCTAAGGTATATAAGCTGCTGGGAGAAAATGAAAACCTATATTTAAAAATGACG
GACAGCCGGTATAAAGGACCACTATGATGTGGAACGGGAGACATGATGCTATGGCTGGAAGGAAAGCTG
CCTGTTCCAAAGGTCCTGCACTTTGAACGGCATGATGGCTGGAGCAATCTGCTCATGAGTGAGGCGGATGGCGTC
CTTTGCTCGGAAGAGTATGAAGATGAACAAAGCCCTGAAAAGATTATCGAGCTGTATGCGGAGTGATCAGGCTC
TTTCACTCCATCGACATATCGGATTGTCCCTATACGAATAGCTTAGACAGCCGCTTAGCCGAATTGGATTACTTA
CTGAATAACGATCTGGCCGATGTGGATTGCGAAAACCTGGGAAGAAGACACTCCATTTAAAGATCCGCGCGAGCTG
TATGATTTTTTTAAAGACGGAAAAGCCCCGAAGAGGAACCTTGTCTTTTCCACGGCGACCTGGGAGACAGCAACATC
TTTGTGAAAGATGGCAAAGTAAGTGGCTTTATTGATCTTGGGAGAAGCGGCAGGGCGGACAAGTGGTATGACATT
GCCTTCTGCGTCCGGTCGATCAGGGAGGATATCGGGGAAGAACAGTATGTGAGCTATTTTTTGACTTACTGGGG
ATCAAGCCTGATTGGGAGAAAATAAAATATTATATTTTACTGGATGAATTGTTTTAGTACCTAGATGTGGCGCAA
CGATGCCGGCGACAAGCAGGAGCGCACCGACTTCTCCGCATCAAGTGTTTTGGCTCTCAGGCCGAGGCCACGG
CAAGTATTTGGGCAAGGGTCCGTGGTATTCTGTCAGGGCAAGATTGGAATACCAAGTACGAGAAGGACGGCCA
GACGGTCTACGGGACCGACTTCATTGCCGATAAGGTGGATTATCTGGACACCAAGGCACCAGGCGGTCAAATCA

GGAATAAGGGGCACATTGCCCGGCGTGAGTCGGGGCAATCCCGCAAGGAGGGTGAATGAATCGGACGTTTGACCG
GAAGGCATACAGGCAAGAACTGATCGACGCGGGGTTTTCCGCGGAGGATGCCGAAACCATCGCAAGCCGACCGT
CATGCGTGCGCCCCGCGAAACCTTCCAGTCCGTCGGCTCGATGGTCCAGCAAGCTACGGCCAAGATCGAGCGCGA
CAGCGTGCAACTGGCTCCCCCTGCCCTGCCCGCGCCATCGGCCGCCGTGGAGCGTTTCGCGTCGCTCGAACAGGA
GGCGGCAGGTTTGGCGAAGTCGATGACCATCGACACGCGAGGAACTATGACGACCAAGAAGCGAAAAACCGCCGG
CGAGGACCTGGCAAAACAGGTCAGCGAGGCCAAGCAGGCCCGGTTGCTGAAACACACGAAGCAGCAGATCAAGGA
AATGACGCTTTCCCTTGTTCGATATTGCGCCGTGGCCGGACACGATGCGAGCGATGCCAAACGACACGCCCCGCTC
TGCCCTGTTTACCACGCGCAACAAGAAAAATCCCGCGCGAGGCGCTGCAAAACAAGGTCATTTTCCACGTCACAA
GGACGTGAAGATCACCTACACCGGCGTCGAGCTGCGGGCCGACGATGACGAACTGGTGTGGCAGCAGGTGTTGGA
GTACGCGAAGCGCACCCCTATCGGCGAGCCGATCACCTTACGTTTACGAGCTTTGCCAGGACCTGGGCTGGTC
GATCAATGGCCGGTATTACACGAAGGCCGAGGAATGCTGTGCGCGCTACAGGCGACGGCGATGGGCTTACGTC
CGACCGCGTTGGGCACCTGGAATCGGTGTGCTGCTGACCGCTTCCGCGTCTGGACCGTGGCAAGAAAACGTC
CCGTTGCCAGGTCTGATCGACGAGGAAATCGTCTGCTGTTTGTGTCGCGACCACTACACGAAATTCATATGGGA
GAAGTACCGCAAGCTGTGCGCGACGGCCCCGACGGATGTTGACTATTTTTCAGCTCGCACCGGGAGCCGTACCCGCT
CAAGCTGGAAACCTTCCGCTCATGTGCGGATCGGATTCACCCGCGTGAAGAAGTGGCGCGAGCAGGTGCGCGA
AGCCTGCGAAGAGTTGCGAGGCGAGCGGCTGGTGGAAACGCGCTGGGTCAATGATGACCTGGTGCATTGCAAACG
CTAGGGCCTTTGTGGGTGAGTTCCGCTGGGGGTTTACGACCGCAGCGCTTACTGGCATTTCAGGAACAAGCGGG
CACTGCTCGAGCAGCTTGTCTGCTCAGTATCGCTCGGGACGACGCGCGCTCTACGAACCTGCCGATAAACAGA
GGATTAAAATTGACAATTTGTGATTAAGGCTCAGATTTCAGCGGCTGGAGCGGCCGACGTGCAGGATTTCCGCGAG
ATCCGATTGTGCGCCCTGAAGAAAGCTCCAGAGATGTTGCGGTCCGTTTACGAGCACGAGGAGAAAAAGCCCATG
GAGGCGTTGCTGAACGGTTGCGAGATGCCGTGGCATTGCGCGCTTACATCGACGGCGAGATCATTTGGGCTGTG
GTCTTCAAACAGGAGGACGGCCCAAGGACGCTACAAGGCGCATCTGTCCGGCGTTTTCTGGAGCCCGAACAG
CGAGGCCGAGGGGTGCGCGGTATGCTGCTGCGGGCGTTGCCGGCGGTTTATTGCTCGTGATGATCGTCCGACAG
ATTCCAACGGGAATCTGGTGGATGCGCATCTTCATCTCGGCGCACTTAATATTTTCGCTATTTCTGGAGCTTGTTG
TTTATTTTCGGTCTACCGCTGCGGGGCGGGGTGCGGGCGACGGTAGGCGCTGTGCAGCCGCTGATGGTCTGTTT
ATCTCTGCCGCTCTGCTAGGTAGCCCGATACGATTGATGGCGGTCTTGGGGGCTATTTGCGGAACGCGGGCGTG
GCGCTGTTGGTGTGACACCAACGCGAGCGCTAGATCTGTGCGGCTCGCAGCGGGGCTGGCGGGGGCGGTTTCC
ATGGCGTTTCGGAACCGTGCTGACCCGCAAGTGGCAACCTCCCGTGCCCTCTGCTACCTTTACCGCTTGCAACTG
GCGGCGGAGGACTTCTGCTCGTTCCAGTAGCTTTAGTGTGTTGATCCGCAATCCCGATGCTTACAGGAACCAAT
GTTCTCGGCGCTGGCGTGGCTCGGCTGATCGGAGCGGGTTTAACTTACTTCTTTTGGTTCCGGGGGATCTCGCGA
CTCGAACCTACAGTTGTTTCTTACTGGGCTTTCTCAGCCCCAGATCTGGGGTGCATCAGCCGGGGATGCATCAG
GCCGACAGTCGGAACCTTCCGGTCCCCGACCTGTACCATTCGGTGAGCAATGGATAGGGGAGTTGATATCGTCAAC
GTTCACTTCTAAAGAAATAGCGCCACTCAGCTTCTCAGCGGCTTATCCAGCGATTTCTATTATGTGCGCATA
GTTCTCAAGATCGACAGCCTGTACGGTTAAGCGAGAAATGAATAAGAAGGCTGATAATTCGGATCTCTGCGAGG
GAGATGATATTTGATCACAGGCGCAACGCTCTGTCTATCGTTACAATCAACATGCTACCCTCCGCGAGATCATCC
GTGTTTCAAACCCGGCAGCTTAGTTGCCGTTCTTCCGAATAGCATCGGTAACATGAGCAAAGTCTGCCGCTTAC
AACGGCTCTCCCGCTGACGCCGTCCCGGACTGATGGGCTGCCGTGATCGAGTGGTGATTTTGTGCCGAGCTGCCG
GTCGGGGAGCTGTTGGCTGGCTGGTGGCAGGATATATTGTGGTGTAACAAATTGACGCTTAGACAACTTAATAA
CACATTGCGGACGTTTTTAATGTACTGGGGTGGTTTTCTTTTACCAGTGAGACGGGCAACAGCTGATTGCCCT
TCACCGCTTGGCCCTGAGAGAGTTGACAGCAAGCGGTTCCACGTTGGTTTGGCCCCAGCAGGCGAAATCTGTGTTGA
TGGTGGTTCCGAAATCGGCAAAATCCCTTATAAATCAAAGAATAGCCCCGAGATAGGGTTGAGTGTGTTCCAGT
TTGGAACAAGAGTCCACTATTAAAGAACGTGGACTCCAACGTCAAAGGGCGAAAAACCGTCTATCAGGGCGATGG
CCCACTACGTGAACCATCACCCAAATCAAGTTTTTTGGGGTCGAGGTGCCGTAAAGCACTAAATCGGAACCCCTAA
AGGGAGCCCCCGATTTAGAGCTTGACGGGGAAGCCGGCGAACGTGGCGAGAAAGGAAGGGAAGAAAGCGAAAGG
AGCGGGCGCCATTACGGCTGCGCAACTGTTGGGAAGGGCGATCGGTGCGGGGCTCTTCGCTATTACGCCAGCTGG
CGAAAGGGGGATGTGCTGCAAGCGGATTAAGTTGGGTAACGCCAGGGTTTTTCCAGTACAGACGTTGTAAAACGA
CGGCCAGTGAATTCGAGCTCGGTACCCCCCTACTCCAAAAATGTCAAAGATACAGTCTCAGAAGACCAAGGGCT
ATTGAGACTTTTCAACAAAGGGTAATTTCCGGAAACCTCCTCGGATTCATTGCCCAGCTATCTGTCACTTCATC
GAAAGGACAGTAGAAAGGAAGGTGGCTCCTACAAATGCCATCATTTGCGATAAAGGAAAGGCTATCATTTCAAGAT
GCCTTGCGCAGAGTGGTCCCAAAGATGGACCCCAACACGAGGAGCATCGTGGAAGAAAGAGACGTCCCAACC
ACGTCTTCAAAGCAAGTGGATTGATGTGACATCTCCACTGACGTAAGGGATGACGCACAATCCCACTATCCCTTCG
CAAGACCCCTTCTTCTATATAAGGAAGTTTCAATTTTCAATTTGGAGAGGACAGCCCCAAGCTTTCTAGAGGATCCATAAA
ACATTTCAATCCTTTGAACGCGGTAGAACGTGCTAATTGGATTTTGGTGAGAACGCGGTAGAACGTACTTATCAC
CTACAGTTTTATTTTGTGTTTTCTTTTGGTTAATCTATCCAGCTTAGTACCGAGTGGGGGAAAGTGACTGGTGT
GCCTAAAACCTTTTCTTTTGATACTTTGTAAAAATACATACAGATACAATGGCGAACGGTAACCTCAAGTTGTCTC
AATTGCTCAATGTGGACGAGATGTCTGCTGAGCAGAGGAGTCAATTTCTTTGACTTGATGCTGACTAAACCTGATT
GTGAGATCGGGCAAATGATGCAAAGAGTTGTTGTTGATAAAGTCGATGACATGATTAGAGAAAGAAAGACTAAAG
ATCCAGTGATTGTTTCATGAAGTTCTTTCTCAGAAGGAACAGAACAGTTGATGGAAATTTATCTGAATTCATA
TCGTGTTTAAAGACGACAAAAACATGGTTCATGGGTTTGGCGCTGCTGAGCGAAACTACAAGCTTTATTGCTTT
TAGATAGAGTTCTGCTCTGCAAGAGGTGGATGACATCGGTGGTCAATGGTCTGTTTTGGGTAACCTAGAGGTGAGA
AAAGGATTCAATCCTGTTGTCCAAATCTAGATATTCGGGATGATCAGAGAGAAATTTCTCGACAGATATTTCTTA

CTGCTATTGGTGATCAAGCTAGAAGTGGTAAGAGACAGATGTCGGAGAATGAGCTGTGGATGTATGACCAATTTT
GTGAAAATATTGCTGCGCCTAACGCGGTTAGGTGCAATAATACATATCAGGGTTGTACATGTAGGGGTTTT

METHOD OF MODULATING THE ALKALOID CONTENT OF TOBACCO

FIELD OF THE INVENTION

[0001] The present invention relates to methods of modulating the alkaloid content a plant or part thereof or cell or cell culture. The invention also extends to methods of modulating the expression and/or activity of polypeptides which modulate alkaloid content within plants. Alternatively, the invention provides methods of modulating the expression and/or activity of genes which encode polypeptides which modulate alkaloid content within plants. The invention also extends to constructs, which can be used to modulate the polypeptides. The invention further relates to plant cells and plants modified to achieve a modulation in alkaloid content. The invention also relates to a processed and harvested leaf from such modulated plants and use thereof in a tobacco industry product, including combustible smoking articles.

BACKGROUND

[0002] Alkaloids are a group of naturally occurring compounds which mostly contain basic nitrogen atoms and are produced by a large variety of organisms including bacteria, fungi, plants and animals.

[0003] Alkaloids may be classified according to the similarity of the carbon skeleton e.g. indole-, isoquinoline- and pyridine-like. Pyridine derivatives are one class of monomeric alkaloids; this class includes simple derivatives of pyridine, polycyclic condensed and noncondensing pyridine derivatives and sesquiterpene pyridine derivatives. Examples are nicotine, normicotine, pseudooxynicotine, anabasine, myosmine and anatabine.

[0004] Most of the known biological functions of alkaloids are related to protection. Neuroactive molecules, such as caffeine, cocaine, morphine, and nicotine, act as defence compounds against invading predators. The accumulation of these alkaloids is the result of signal transduction cascades that monitor gene expression, enzyme activities, and alkaloid concentrations. The fine-tuning of alkaloid content in the plant involves negative feedback loops and degradative pathways.

[0005] Nicotine occurs naturally in several varieties of plants but is found at the highest level in the tobacco plant. Cultivated tobacco produces 2-4% alkaloids of total dry weight. Nicotine is produced in wild and cultivated *Nicotiana* species and plays an important role in plant defence against herbivores and insects (Voelckel et al. (2001) *Oecologia* 127(2): 274-280, incorporated herein by reference). It accounts for ~90% of the total alkaloid content. The remaining 10% of the alkaloid pool is mostly constituted by the structurally related compounds normicotine, anatabine, anabasine and pseudooxynicotine (PON).

[0006] The regulation of alkaloid content in tobacco is complex. Several factors including genotype, environment, fertilization and agronomic practices (e.g. topping) affect alkaloid levels in tobacco plants. Some key regulators of nicotine biosynthesis are well characterized, for example putrescine N-methyltransferase (PMT), which plays a pivotal role in this pathway, is activated by members of the ethylene responsive factor (ERF) superfamily, the largest

transcription factor family in the tobacco genome (Rushton et al. (2008) *Plant Physiol.* 147(1): 280-295 incorporated herein by reference).

[0007] Tobacco pyridine alkaloids are precursors of tobacco-specific nitrosamines (TSNAs) that form during the post-harvest leaf curing. The four primary TSNAs found in cured tobacco leaves are N'-nitrosonornicotine (NNN), N'-nitrosoanatabine (NAT), N'-nitrosoanabasine (NAB) and 4-(methyl nitrosamino)-1-(3-pyridyl)-1-butanone (NNK). During the post-harvest leaf curing, reactions between pyridine alkaloids and nitrosating species leads to the formation of TSNAs.

[0008] PON is likely to function as the direct precursor in the synthesis of the TSNA NNK (Bush et al., 2001 incorporated herein by reference). Reducing the production and accumulation of TSNAs is of high importance. The CYP82E family of nicotine demethylase genes is one of the primary regulators of nicotine to normicotine conversion, and altering their activity or accumulation may result in a decrease in NNN levels.

[0009] As described in the Examples, the inventors sought to investigate genes responsible for alkaloid and/or TSNA precursor synthesis, with the aim of modulating alkaloid content in plants, e.g. decreasing TSNA content in tobacco.

SUMMARY OF THE INVENTION

[0010] It has been surprisingly found that by modulating the activity or expression of Galactose-1-phosphate uridyl transferase, the alkaloid content and/or TSNA content or precursor of TSNA content of plants can be modulated. The gene(s) as taught herein, for example Nitab4.5_0002825g0060.2, are regulators of alkaloid and TSNA precursor content in cultivated tobacco. In particular, the gene (s) as taught herein, for example Nitab4.5_0002825g0060.2, is a regulator of alkaloid content in cultivated tobacco. Nitab4.5_0002825g0060.2 encodes Galactose-1-phosphate uridyl transferase according to the present invention. Homologues of Nitab4.5_0002825g0060.2 are provided in Table 1.

[0011] According to the present invention, tobacco industry products with modulated alkaloid content and commercially desirable traits sought after by consumers of tobacco industry products can be produced. In some instances, consumers may desire a product with low levels of alkaloid content. In some instances, consumers may desire a product with low levels of TSNA precursors.

[0012] The present invention may be particularly useful in the field of plant molecular farming, where plants (such as tobacco and other *Nicotiana* spp.) are used for the production of proteins, peptides, and metabolites e.g. for the production of therapeutics and pharmaceuticals such as antibiotics, virus like particles, or neutraceuticals or small molecules. Tobacco has been used for the development of an HIV-neutralising antibody in an EU-funded project called PharmPlant and Medicago Inc., Canada have worked on a tobacco-based platform for the production of virus-like particles for flu vaccine manufacture.

[0013] Thus, a plant according to the present invention may be used for molecular farming to reduce or eliminate the presence of nicotinic alkaloids. The use of a low nicotine plant or rootstock is beneficial in molecular farming and would reduce downstream processing costs associated with purification.

[0014] The present inventors have surprisingly determined a method for modulating (e.g. decreasing) the alkaloid content, of a plant (e.g. a tobacco plant) by modulating (e.g. decreasing) the activity or expression of Galactose-1-phosphate uridyl transferase. The alkaloid content (e.g. the content of one or more of nicotine, nornicotine, PON, anabasine, anatabine or myosmine, suitably the content of one or more of nicotine, nornicotine, PON, anabasine or anatabine) of a plant (e.g. tobacco plant) may be decreased by decreasing the activity or expression of Galactose-1-phosphate uridyl transferase or may be increased by increasing the activity or expression of Galactose-1-phosphate uridyl transferase.

[0015] Prior to the present invention it had not been known that modulation of the activity or expression of Galactose-1-phosphate uridyl transferase as described herein could be used to modulate alkaloid content or modulate TSNA precursor content in particular, nornicotine, PON, anabasine and/or anatabine content.

[0016] The present invention provides a method of modulating (e.g. decreasing) the alkaloid content of a tobacco plant or a part thereof or tobacco plant cell, the method comprising modifying said plant or plant cell by modulating (e.g. decreasing) the activity or expression of Galactose-1-phosphate uridyl transferase.

[0017] In another aspect, the present invention provides a method of modulating (e.g. decreasing) the content of a tobacco specific nitrosamine (TSNA) or a precursor of a TSNA in a tobacco plant or plant part thereof or tobacco plant cell, the method comprising modifying said plant or plant cell by modulating (e.g. decreasing) the activity or expression of Galactose-1-phosphate uridyl transferase.

[0018] In another aspect, the present invention provides a method for producing a plant or part thereof, a cell or cell culture, a plant propagation material, a leaf, a cut harvested leaf, a processed leaf or a cut and processed leaf which has modulated (e.g. decreased) alkaloid content, the method comprising modifying said plant or cell culture to modulate the activity or expression of Galactose-1-phosphate uridyl transferase.

[0019] In another aspect, the present invention provides the use of at least one gene encoding a Galactose-1-phosphate uridyl transferase for modulating alkaloid content of a tobacco cell or tobacco plant or part thereof.

[0020] Suitably the Galactose-1-phosphate uridyl transferase may:

[0021] a) comprise an amino acid sequence as set out in SEQ ID No. 3; or a functional variant or functional fragment or orthologue of SEQ ID No. 3; or a sequence which has at least 80% identity to SEQ ID No. 3; or a homologue of SEQ ID No. 3; or

[0022] b) is encoded by a nucleotide sequence as set out in SEQ ID No. 1 or 2; or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2; or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2.

[0023] Suitably, the alkaloid content is modulated (e.g. decreased) in comparison to a plant or cell culture which has not been modified to modulate the activity or expression of Galactose-1-phosphate uridyl transferase.

[0024] In another aspect, the present invention provides a tobacco plant or part thereof or a tobacco cell or cell culture which has been modified to modulate (e.g. decrease) the

activity or expression of Galactose-1-phosphate uridyl transferase and the tobacco plant or part thereof or tobacco cell or cell culture has decreased alkaloid and/or TSNA precursor content in comparison to an unmodified plant or unmodified cell or cell culture.

[0025] In a further aspect, the present invention provides a plant propagation material obtainable (e.g. obtained) from a plant according to the present invention or from a plant or cell or cell culture produced by the method or use according to the present invention.

[0026] Suitably, the alkaloid content of the plant may be decreased in comparison to a plant or cell culture which has not been modified to modulate the activity or expression of Galactose-1-phosphate uridyl transferase.

[0027] Suitably, the content of one or more alkaloids selected from nicotine, nornicotine, PON, anabasine, myosmine and anatabine may be modulated (e.g. decreased), preferably the content of nicotine, nornicotine and/or PON is modulated (e.g. decreased).

[0028] Suitably, the nicotine content may be decreased.

[0029] In one aspect, the present invention provides the use of a tobacco plant or part thereof or tobacco cell or cell culture according to the present invention, or of a plant produced according to the present invention to breed a plant.

[0030] In another aspect, the present invention provides the use of a tobacco plant or part thereof or a tobacco cell or cell culture according to the present invention, or of a plant produced by a method according to the present invention for production of a product.

[0031] In another aspect, the present invention provides the use of a tobacco plant or part thereof according to the present invention, or of a plant produced by a method according to the present invention to grow a crop.

[0032] In another aspect, the present invention provides the use of a tobacco plant or part thereof according to the present invention, or of a plant produced by a method according to the present invention to produce a leaf.

[0033] In one aspect, the present invention provides a harvested leaf of a plant according to the present invention, or obtainable from a plant propagated from a propagation material according to the present invention, or obtainable from a plant obtained by a use according to the present invention, or obtainable from a plant produced by a method according to the present invention.

[0034] Suitably, the harvested leaf of a plant may be a cut harvested leaf.

[0035] In one aspect, the present invention provides a processed leaf, preferably a processed tobacco leaf, preferably a non-viable processed tobacco leaf:

[0036] obtainable (e.g. obtained) from a plant obtainable from a use according to the present invention;

[0037] obtainable (e.g. obtained) by processing a plant according to the present invention; obtainable (e.g. obtained) from a plant propagated from a plant propagation material according to the present invention; or

[0038] obtainable (e.g. obtained) by processing a harvested leaf of a plant according to the present invention; or

[0039] obtainable (e.g. obtained) from a plant produced by a method according to the present invention.

[0040] Suitably, the processed leaf may be processed by curing, fermenting, pasteurising or a combination thereof.

[0041] Suitably, the processed leaf may be a cut processed leaf.

[0042] In one aspect, the present invention provides a cured tobacco material made from a plant or a part thereof according to the present invention, or a harvested leaf according to the present invention, or a processed leaf according to the present invention.

[0043] In another aspect, the present invention provides a tobacco blend comprising said cured tobacco material according to the present invention.

[0044] In one aspect, the present invention provides a tobacco industry product prepared from: a tobacco plant or part thereof or tobacco cell or cell culture according to the present invention;

[0045] a tobacco plant or part thereof propagated from a tobacco plant propagation material according to the present invention;

[0046] a harvested leaf of a plant according to the present invention;

[0047] a processed leaf according to the present invention.

[0048] Suitably, the tobacco product may be a combustible smoking article.

[0049] Suitably, the tobacco product may be a smokeless tobacco product.

[0050] Suitably, the tobacco product may be a non-combustible aerosol provision system such as a tobacco heating device or an aerosol-generating device.

[0051] In another aspect, the present invention provides a combustible smoking article, non-combustible aerosol provisioning system, smokeless tobacco product or tobacco heating device comprising a plant or a part thereof according to the present invention or an extract (e.g. a tobacco extract) thereof or a tobacco cell culture according to the present invention; or a cured tobacco material according to the present invention; or a tobacco blend according to the present invention.

[0052] In one aspect, the present invention provides the use of a nucleotide sequence encoding Galactose-1-phosphate uridyl transferase protein which:

[0053] a) encodes an amino acid sequence as set out in SEQ ID No. 3; or a functional variant or functional fragment or orthologue of SEQ ID No. 3; or a sequence which has at least 80% identity to SEQ ID No. 3; or a homologue of SEQ ID No. 3; or

[0054] b) comprises a sequence as set out in SEQ ID No. 1 or 2; or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2; or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2;

[0055] to select a plant having modulated (e.g. reduced) alkaloid content and/or modulated (e.g. reduced) content of TSNA or a precursor of a TSNA.

[0056] In another aspect, the present invention provides a mutant of a plant carrying a heritable mutation in a nucleotide sequence which:

[0057] a) encodes an amino acid sequence as set out in SEQ ID No. 3; or a functional variant or functional fragment or orthologue of SEQ ID No. 3; or a sequence which has at least 80% identity to SEQ ID No. 3; or a homologue of SEQ ID No. 3; or

[0058] b) comprises a sequence as set out in SEQ ID No. 1 or 2; or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2; or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2;

[0059] wherein said heritable mutation modulates (e.g. decreases) the activity or expression of Galactose-1-phosphate uridyl transferase and wherein the mutant plant has modulated (e.g. decreased) alkaloid content and/or modulated content of a TSNA or a precursor of a TSNA relative to a comparable plant which does not carry said heritable mutation.

[0060] In another aspect, the present invention provides progeny or seed of a mutant plant which carries the heritable mutation according to the present invention.

[0061] In one aspect, the present invention provides a harvested leaf, a processed leaf or cured tobacco material produced from a plant comprising a modification in a nucleotide sequence which:

[0062] a) encodes an amino acid sequence as set out in SEQ ID No. 3; or a functional variant or functional fragment or orthologue of SEQ ID No. 3; or a sequence which has at least 80% identity to SEQ ID No. 3; or a homologue of SEQ ID No. 3; or

[0063] b) comprises a sequence as set out in SEQ ID No. 1 or 2; or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2; or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2;

[0064] wherein said modification modulates (e.g. decreases) the activity or expression of Galactose-1-phosphate uridyl transferase and wherein said plant has modulated (e.g. decreased) alkaloid content and/or modulated content of a TSNA or a precursor of a TSNA relative to a comparable plant which does not carry said modification in said Galactose-1-phosphate uridyl transferase.

[0065] In one aspect, the activity or expression of at least one Nic1 ERF gene (such as any one or more of those in FIG. 6; SEQ ID No 7-10) and/or at least one Nic2 ERF gene (such as any one or more of those in FIG. 7; SEQ ID No 11-14) is modulated in addition to the at least one Galactose-1-phosphate uridyl transferase. Suitably, said at least one Nic1 ERF and/or Nic2 ERF gene may comprise a mutation which decreases its expression and/or activity. In one aspect, the activity of ERF199 (e.g. at least one of SEQ ID NO. 7-10) is modulated (e.g. decreased) in addition to the modulation (e.g. decrease in activity and or expression) of the at least one Galactose-1-phosphate uridyl transferase. In one aspect, the activity of ERF189 (e.g. at least one of SEQ ID No.11-14) is modulated (e.g. decreased) in addition to the modulation (e.g. decrease in activity and or expression) of the at least one Galactose-1-phosphate uridyl transferase. In one aspect, the activity of ERF199 (e.g. at least one of SEQ ID NO. 7-10) and ERF189 (e.g. at least one of SEQ ID No.11-14) is modulated (e.g. decreased) in addition to the modulation (e.g. decrease in activity and or expression) of the at least one Galactose-1-phosphate uridyl transferase. Exemplary sequences of ERF 199 and ERF189 are provided in FIGS. 8-45 (SEQ ID Nos 7-14). See also WO2018237107, which is incorporated herein in by reference in its entirety.

BRIEF DESCRIPTION OF THE DRAWINGS

[0066] Embodiments of the invention will now be described, by way of example only, with reference to the accompanying drawings, in which:

[0067] FIG. 1 shows the alkaloid content of plants targeted with VIGS construct for Nitab4.5_0002825g0060.2. Graph

is representative of 2 biological replicates (N=12) analysed by one-way ANOVA and Tukey's multiple-comparison post-test. Values are shown as means $\pm$ SEM. Asterisks indicate statistical significance of P value 0.05. Pyridine alkaloids: nicotine, normicotine and pseudonycotine (PON).

[0068] FIG. 2 shows the measured endpoint absorbance values (340 nm) for the production of B-NADPH over a 60 minute reaction by 1.7 pg of Nitab4.5_0002825g0060.2 enzyme or 0.5 pg of positive control enzyme (recombinant human Galactose-1-phosphate uridyl transferase, MyBio-source, MBS143204) in the presence of 0.67 mM uridine 5'-diphosphoglucose and 1.3 mM galactose 1-Phosphate. The results for the control reactions are also shown on the same graph and indicate that the reaction is dependent on uridine 5'-diphosphoglucose and galactose 1-Phosphate. MBP and the no enzyme control (-ve) both have limited activity. The commercial positive control displays >5-fold higher activity per microgram of protein in this assay but the Nitab4.5_0002825g0060.2 enzyme (labelled T4372) demonstrates reliable, substrate-dependent activity. Error bars are standard deviation from four replicates.

[0069] FIG. 3 shows the genomic sequence of Nitab4.5_0002825g0060.2 (SEQ ID No. 1).

[0070] FIG. 4 shows the coding sequence of Nitab4.5_0002825g0060.2 (SEQ ID No. 2).

[0071] FIG. 5 shows the amino acid sequence of Nitab4.5_0002825g0060.2 (SEQ ID No. 3).

[0072] FIG. 6 provides a table of Nic1 ERFs.

[0073] FIG. 7 provides a table of Nic2 ERFs.

[0074] FIGS. 8-15 show the sequences of ERF199 (SEQ ID Nos 7-10) and ERF189 (SEQ ID Nos 11-14).

[0075] FIG. 16 shows the sequence of TRV1 (SEQ ID No 15) as used in Example 1.

[0076] Some sequences disclosed herein contain "X" or "N" in nucleotide sequences. "X" or "N" can be any nucleotide or a deletion or insertion of one or more nucleotides. For example, in some cases a string of "X"s or "N"s are shown. The number of "X"s or "N"s does not necessarily correlate with the actual number of nucleotides at that position. There may be more or fewer nucleotides than shown as "X" or "N" in the sequence.

DETAILED DESCRIPTION

[0077] For the first time the present inventors have shown that by modulating the activity or expression of at least one Galactose-1-phosphate uridyl transferase in a plant (e.g. a tobacco plant) or a cell (e.g. tobacco cell), the alkaloid and/or TSNA precursor content of the plant (or processed plant) or cell can be modulated.

[0078] Galactose-1-phosphate uridylyltransferase or GalT plays an important role in galactose metabolism. GalT catalyses the transfer of an uridyldiphosphate group on galactose (or glucose) 1-phosphate. During the reaction, the uridyl moiety links to a histidine residue. In the *Escherichia coli* enzyme, it has been shown that two histidine residues separated by a single proline residue are essential for enzyme activity. The closest homologue of Nitab4.5_0002825g0060.2 in *Arabidopsis* (At5g18200, 62% identity) catalyses the conversion of ADP-glucose and inorganic phosphate (Pi) into glucose-1-phosphate and ADP. The role of GalT in the regulation of pyridine alkaloid content in tobacco was until now unknown.

[0079] The present invention provides a method of modulating (e.g. decreasing) the alkaloid content of a plant or a

part thereof, the method comprising modifying said plant by modulating (e.g. decreasing) the activity or expression of at least one Galactose-1-phosphate uridyl transferase.

[0080] Also provided is a method of modulating (e.g. decreasing) the content of a tobacco specific nitrosamine (TSNA) precursor in a tobacco plant or plant part thereof, the method comprising modifying said plant by modulating (e.g. decreasing) the activity or expression of at least one Galactose-1-phosphate uridyl transferase.

[0081] The at least one Galactose-1-phosphate uridyl transferase may be selected from an amino acid sequence as set out in SEQ ID No. 3, or a functional variant or functional fragment or orthologue thereof, or a sequence which has at least 80% identity to SEQ ID No. 3, or a homologue of SEQ ID No. 3; or said at least one Galactose-1-phosphate uridyl transferase may be encoded by a which comprises a sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2, or a homologue of SEQ ID No. 1 or 2.

[0082] In one embodiment at least two genes encoding Galactose-1-phosphate uridyl transferase are modified selected from the group of: genes which encode polypeptides comprising an amino acid sequence as set out in SEQ ID No. 3 or a functional variant or functional fragment or orthologue thereof, or a sequence which has at least 80% identity to SEQ ID No. 3, or a homologue of SEQ ID No. 3; or genes encoding Galactose-1-phosphate uridyl transferase comprising a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2, or a homologue of SEQ ID No. 1 or 2.

[0083] In one embodiment, at least three, such as at least four, such as at least five, such as at least six, such as at least seven, such as at least eight, such as at least nine, such as ten Galactose-1-phosphate uridyl transferases are modulated, wherein the Galactose-1-phosphate uridyl transferases comprise an amino acid sequence as set out in SEQ ID No. 3 or a functional variant or functional fragment or orthologue thereof, or a sequence which has at least 80% identity to SEQ ID No. 3, or a homologue of SEQ ID No. 3; or comprise a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2, or a homologue of SEQ ID No.1 or 2.

[0084] In one aspect, the at least one Galactose-1-phosphate uridyl transferase comprises or consists of an amino acid sequence as set out in: SEQ ID No. 3 or a functional variant or functional fragment or orthologue thereof, or a sequence which has at least 80% identity to SEQ ID No. 3; or wherein the Galactose-1-phosphate uridyl transferase comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2 or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2; or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2. In one aspect, the activity or expression of at least one further gene is modulated. Suitably, at least two (or at least three or at least four or at least five or at least six or at least seven or at least eight or at least nine) additional genes selected from Table 1 or a sequence having at least 80% sequence identity thereto may also be modulated.

[0085] The “expression” of Galactose-1-phosphate uridyl transferase may refer to the level of transcription, translation i.e. protein expression.

[0086] Measurement of the level or amount of a gene product may be carried out by any suitable method, for example comparison of mRNA transcript levels, protein or peptide levels, and/or phenotype of a plant, between a modified plant and comparable plant which has not been modified according to the present invention.

[0087] The term “a comparable product” as defined herein would be one derived from a plant (e.g. a tobacco plant) which had not been modified according to the present invention, but in which all other relevant features were the same (e.g. plant species, growing conditions, method of processing the plant, e.g. tobacco, etc.). The comparable product according to the present invention may mean a plant (e.g. a tobacco plant) or a part thereof, such as a leaf (e.g. a tobacco leaf), a harvested leaf (e.g. a harvested tobacco leaf), a cut harvested leaf (e.g. a cut harvested tobacco leaf), a processed leaf (e.g. a processed tobacco leaf) or plant propagation material (e.g. tobacco plant propagation material), or a product comprising said plant or part therefore, e.g. a tobacco industry product or combinations thereof obtainable or obtained from a plant which has not been modified in accordance with the present invention, e.g. to modulate the activity or expression of a gene encoding a Galactose-1-phosphate uridyl transferase. In one embodiment a comparable product is one which does not comprise a gene encoding a Galactose-1-phosphate uridyl transferase whose activity or expression has been modulated.

[0088] The term “modifying” or “modified” as used herein means a plant (e.g. a tobacco plant) or nucleic acid sequence that has been altered or changed. The present invention comprises the modification of plants using techniques for genetic modification of plants or non-genetic modification of plants. Such methods are well known in the art and examples of genetic modification techniques include transformation, transgenics, cisgenics, and gene editing methods. Examples of non-genetic modification techniques include fast-neutron mutagenesis, chemical mutagenesis e.g. ethyl methane-sulfonate (EMS) mutagenesis and modern population analysis approaches.

[0089] In one embodiment a natural variant which has a modified gene encoding a Galactose-1-phosphate uridyl transferase is selected and that trait or gene is bred into a second plant which may have commercially desirable traits.

[0090] In one embodiment the plant according to the present invention is a transgenic plant. In one embodiment the plant according to the invention is a non-transgenic plant.

[0091] The term “unmodified plant” as defined herein would be a plant (e.g. a tobacco plant) which had not been modified according to the present invention, e.g. to modulate the activity or expression of Galactose-1-phosphate uridyl transferase or to modify the nucleic acid sequence of at least one gene encoding a Galactose-1-phosphate uridyl transferase; and in which all other relevant features were the same (e.g. plant species, growing conditions, method of processing tobacco, etc.). In one embodiment an unmodified plant is one which does not comprise a gene encoding a Galactose-1-phosphate uridyl transferase whose activity or expression has been modulated. In one embodiment, an unmodified plant is one which does not comprise a modified

nucleic acid sequence which encodes at least one gene encoding a Galactose-1-phosphate uridyl transferase.

Galactose-1-Phosphate Uridyl Transferase

[0092] A “Galactose-1-phosphate uridyl transferase” as used herein refers to a protein which catalyses facilitates the simultaneous conversion of uridine diphosphoglucose (UDP-glucose) and galactose-1-phosphate (gal-1P) to uridine diphosphogalactose (UDP-galactose) and glucose-1-phosphate.

[0093] Suitably, a Galactose-1-phosphate uridyl transferase may be identified using the assay described in Example 2 herein. For example, a Galactose-1-phosphate uridyl transferase may be identified using the assay described in Mayes, J. S. and Hanson R. G. (1966) *Methods in Enzymology* IX, 708-713.

[0094] An illustrative sequence of Galactose-1-phosphate uridyl transferase from tobacco is shown in SEQ ID No. 3.

[0095] A Galactose-1-phosphate uridyl transferase may be identified by comparing the amino acid sequence of the protein in question to the amino acid sequence of SEQ ID No. 3.

[0096] In one aspect, a Galactose-1-phosphate uridyl transferase in accordance with the present invention comprises or consists of an amino acid sequence set forth in SEQ ID No. 3 or a sequence with at least 80% identity thereto; or a functional fragment thereof which has Galactose-1-phosphate uridyl transferase activity.

[0097] Domains within the amino acid sequence of a protein may be identified using domain prediction software known in the art. Domains are also described in protein databases such as UniprotKB. Without wishing to be bound by theory, it is hypothesized that modulating content of Galactose-1-phosphate uridyl transferase in a plant cell or modulating activity, of Galactose-1-phosphate uridyl transferase in a plant would alter the metabolic pathways producing alkaloids and TSNA precursors, resulting in modulated alkaloid and/or TSNA precursor content.

[0098] In one embodiment, Galactose-1-phosphate uridyl transferase comprises an amino acid sequence shown as SEQ ID No. 3, or a sequence which has at least 80% identity thereto, or a homologue thereof. Suitably, a homologue of SEQ ID No. 3 may be selected from the group comprising the amino acid sequences provided in Table 1 or a sequence which has at least 80% identity thereto.

[0099] In one embodiment, Galactose-1-phosphate uridyl transferase comprises an amino acid sequence shown as SEQ ID No. 3, or a sequence which has at least 80% identity thereto (preferably at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identity thereto). In one embodiment Galactose-1-phosphate uridyl transferase comprises an amino acid sequence shown in Table 1, or a sequence which has at least 80% identity thereto (preferably at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identity thereto).

[0100] In one embodiment, the Galactose-1-phosphate uridyl transferase according to the present invention comprises or consists of an amino acid shown as SEQ ID No. 3. In one embodiment, the Galactose-1-phosphate uridyl transferase according to the present invention comprises or consists of an amino acid shown in Table 1.

[0101] Suitably, the protein may be from *Nicotiana tabacum*.

[0102] In one embodiment, Galactose-1-phosphate uridyl transferase is encoded by a polynucleotide sequence wherein the gene (prior to mutation) comprises a polynucleotide sequence shown as SEQ ID No. 1, or a sequence which has at least 80% identity thereto, or a homologue thereof.

[0103] Suitably, a homologue of SEQ ID No. 1 may be selected from the group comprising the polynucleotide sequences provided in Table 1 or a sequence which has at least 80% identity thereto.

[0104] In one embodiment, Galactose-1-phosphate uridyl transferase is encoded by a polynucleotide sequence shown as SEQ ID No. 1, or a sequence which has at least 80% identity thereto (preferably at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identity thereto). In one embodiment Galactose-1-phosphate uridyl transferase is encoded by a polynucleotide sequence shown in Table 1, or a sequence which has at least 80% identity thereto (preferably at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identity thereto).

[0105] In one embodiment, the Galactose-1-phosphate uridyl transferase is encoded by a polynucleotide sequence which comprises or consists of a polynucleotide sequence shown as SEQ ID No. 1. In one embodiment, the Galactose-1-phosphate uridyl transferase is encoded by a polynucleotide sequence which comprises or consists of a polynucleotide sequence shown in Table 1.

[0106] In one embodiment, Galactose-1-phosphate uridyl transferase is encoded by a polynucleotide sequence wherein the gene (prior to mutation) comprises a polynucleotide sequence shown as SEQ ID No. 2, or a sequence which has at least 80% identity thereto, or a homologue thereof.

[0107] Suitably, a homologue of SEQ ID No. 2 may be selected from the group comprising the polynucleotide sequences provided in Table 1 or a sequence which has at least 80% identity thereto.

[0108] In one embodiment, Galactose-1-phosphate uridyl transferase is encoded by a polynucleotide sequence shown as SEQ ID No. 2, or a sequence which has at least 80% identity thereto (preferably at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identity thereto). In one embodiment Galactose-1-phosphate uridyl transferase is encoded by a polynucleotide sequence shown in Table 1, or a sequence which has at least 80% identity thereto (preferably at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identity thereto).

[0109] In one embodiment, the Galactose-1-phosphate uridyl transferase is encoded by a polynucleotide sequence which comprises or consists of a polynucleotide sequence shown as SEQ ID No. 2. In one embodiment, the Galactose-1-phosphate uridyl transferase is encoded by a polynucleotide sequence which comprises or consists of a polynucleotide sequence shown in Table 1.

TABLE 1

Homologues of SEQ ID No. 1, 2 and 3
Homologue genomic sequence (SEQ ID NO: 4): ACCTCAGCTTTCATATATTATACCTTTCCACGAAACGCTGTTCTTAACCTCCAGATTTT CAGTGAAACATCGAAGTGAAGTGGAGGAGCACCAGTTCGGAGCCCC GACACAAAGGCAATGGCGGAGGCAGAACGACCAATCGGAGCCCCGAAATCCGGAAGAGC AGAATCCACAAACAGGTGGGTCCATTCTCGCCGGCGAGATCAGGGCG GCCGTCTGACTTCAAGGCAAAATCGAACCCACACCGAACAATAAACAGAAATGCCCATTT TGCGCCGGCCACGAGCACGAGTGTGCGCCGGAGATTTTCGTGTGCC GGCAGATTCCACCGCTGATTGGAAAAATCAGAGTTATCCAAATCTCTTTCCGGCAGTCAGC AGAGAATTGGAGTTCCAAATTCGGTTTCTGCCCTCGGGGACGTAGC TGCTAGTGGGTTTCGGGTTTCACGATGTGGTTATTGAATCGCCGTTCAATCAATAAATATG AGCGATCTTTCCGCGTCTCAAGTTGGGGAAGTTATGCTGGCTTTTAA GAAGCGGGTCGAGCAGCTTCAGAAATGTGATAATGTCAAATATGTTAGGTTTTATCTGA TCCCGTCTAAAAATGCATTGGTTTTCGTATCAATTTGCTATATTTT GTACTGCATATTCATTCTTTGATTTTATATGAATTTTAAATAAATGCATTTAAAGCTAC ATAAAATAGTAGAGACCATGACTGTTAAAGCTAAAGTATTTAAC AAAGTATATATGAAGCTTACTGCAATGGTGAATTTTTGTTTGGCGTAAAGTACTTTGA AGATTACCTGGCTTTAATCTTTTGGTGCGATTGCTTATATGGGTTT ATTTCCATTAACCTGAAAGTGTGTTTTGTTTTGTACAAATAAATAAAGAGTAAACGGTAA TGGTAGAGCCCATCAATGTTTGTACAAAGTGGCCAAACCAATCTAGC TTCTTTTATGTTATATTTAGCAACCTATTTGTGCTAGGTTAAGTTTATATTTAGGTTT TTTTTATTTAGCTGCTTATCATTGTCTGCTTTGGTTTTGTGGCTT TTTAATTTGGCAGACTGAATGATTTTTCTGTGTAACATTGGTGGAGAGAAGGCTGACAACA TGTTTGTCTTTAGGTTTTTAATAGAACAAATATAAGCTAGTTCCAAAT ACTTTAGATCTTGCTGCTTTGAAGTTTGTACTGGAAGAGTAGCTCTTTAATCAGACCAAC TTGCTAAATACTGAACCAATATGAGAGGTTTTTGTGGAACTC ATAGACTGTCAAGTGTCAACTGCTTGAGGGCTTATCATTAAAGAGATAATTTTAGAGGAA GCTTTTCCATTAACTGCATAGGAGCTTGATATGAACCTCCATC CATTTATCTGATATGGTTTGATTTTCATTGTAGATTATTTGAGTTTGTATTTTAGTGACA ACTGACAGGAGAAATCTGGTAGCATTAAATGAAGTTGGTTTGATA TGGTAAATATCAATTTGACTGTAAATTTGAACATTTAAACAGAACTACTCTAAACTTGTA AGTCAATGTAGCAGTTTATCCAAAAAATAAAAAACCTTGTAAT CATGTAGCATATTTGTAGTATCAAGCAAACTTGTGGATATCCAGATTACAAACCAT TTTGAGAGGTATTTAGCTGACATCTCTGTTACTTGGTTCTGTT CCATTGCAGAACTCTAAATATGCACTTCTATCTACTATACCAATATATGTAAGAGA ACTTGAGAAGTGATTAGTTCTCATAAATGGAGATGAATGAGCACAA TATCTGATCTCATGCCCTTTGTGCGTGCAAGGAATTTGCAGTACGGAACAGCATGTATTGT TTTTTCCACTTCTGTTGAGCGCTAATGATTTTACTGTACTCAGC AAGATTGTCCCATCAAAGGCTACTGGACTATATAACACTAATTCATTCAATGGCTTATAA AGGTGTTCAAAACCATGGAGCCTCTGCTGGTGCTCAATGAGCCAT

TABLE 1-continued

Homologues of SEQ ID No. 1, 2 and 3
TCTCACAGCCAGATGATTGCTCTTCCCATTGTTCTCCAGCAGTTTCTGCTCGTCTTGACA GTATGAGGGAATACTACAAGCAGACTGGGAAGTGACGCTTTGTGAT ATTCAACCAATGAATTGTTAATTGATCAATCAGCCATTTTCATATCATTGGTTCCCTTTG CTGCAACTTTTGTCTTTGAGATTTGGATAATTCCTCGTGATCACTCC TCTCATTTTCATGAACCTGATAATGAAAAGGTAATCGATGCTTTTCTTGCCATAAGTTG TTTTTCATGAGAAAATATGTGAAGTTACACAGTTTAACTTTGCAAA TATCAGTTATCCCCAAATCTACGTGAGAAAAGTCAACCGTAGACCATGAGGGACTTTG ACTTTTCTCAAGTAGAATTGCTGATTAGTGAAGAGAAAAGGAAAAC TCATAAAAAAATGATATTTCTCTGTTTCAGGCTAACTAATTTTAGATTGTTTCTCTATC TCAACAACCACTTAGTTGAAAAGGTTTGTGATGTTTACCAATCAA ATATGCAGGCAATGATCTTGGAGGGTTGTTGAACTCGTGCTTTTGAAGATGTCTTTGCA GTTGAACCAATCCACCATTCACCTCCTTATTCATACCTCCCATTTT AAGTAGATCCCTCATATGCTCCTTCCACACACTGGTTTTTACAAATAGCTCCCATTTGAG TGGAGTTGGTGGATTGAGATAGCAACCGGATGTACATAAATCCTG TTTTCCAGAGGATGCTGCCAAATTTCTAAGGGATGTGAGATCCTTAATTAACAGTATTT AGGGGAATCTATTAATCTAATATGTCATATCTTTAGACACAGTAA CTAAGGAGATAAAGAGGATGTGGCTACAGGGAAGTTGTACCCCTTTTATGCGGAGTTATGC GTAGATTCGATTTTGTATCCATTACAAATGTTTCTTATGTTACAC GGTCATTTTCACGTAAAAAATAATACATGAGAT Homologue coding sequence (SEQ ID NO: 5): ATGGCGGAGGCAGAAGCACCAATCGGAGCCCCGAAATCCGGAAGACAGAATCCACAACAG GTGGGTCTTATTCTCGCCGGCAGATCAGCGCGGCGTCTGACTTC AAGGCAAAATCGAACCCCAACCGAACCAATAAACAAGATGCCATTTTGCGCCGCCACGA GCACAGTGTGCGCCGGAGATTTTTCGTGTGCCCGCAGATCCACC GCTGATTGGAATAACAGAGTTATCCAAATCTCTTTCGGCAGTCAGCAGAGAATTGGAGTT CCAAATTCGGTTTCTGCCCTCGGGGACGTAGCTGCTAGTGGGTTT GGGTTTCACGATGTGGTTATTGAATCGCCGTTTCAATCAATAAATATGAGCGATCTTTCGCC GTCTCAAGTTGGGAAGTTATGCTGGCTTTTAAGAAGCGGGTCGAG CAGCTTCAGAATTGTGATAATGTCAAATATGTTCAAGTGTTCAAAACCATGGAGCCTCTGC TGGTGCTTCAATGAGCCATTCTCACAGCCAGATGATTGCTCTTCCC ATTGTTCTCCAGCAGTTTCTGCTCGTCTTGACAGTATGAGGGAATACTACAAGCAGACTGG GAAGTGACGCCTTTGTATATTCAACCAATGAATTGTTAATTGAT CAATCAGCCCATTTTCATATCATTGGTTCCCTTTGCTGCAACTTTTGTCTTTGAGATTTGGAT AATTCTCTGATCACTCCTCTCATTTTCATGAACCTGATAATGAA AAGGCAATGATCTTGGAGGGTTGTGAACTCGTGCTTTTGAAGATGTCTTTCAGTGTGAA CAATCCACCATTCACCTCCTTATTCATACCTCCCATTTTCAAGTA GATCCCTCATATGCTCCTTCCACACACTGGTTTTTACAAATAGCTCCCATTTGAGTGGAGT TGGTGGATTCGAGATAGCAACCGGATGTACATAAATCCTGTTTTT CCAGAGGATGCTGCCAAATTTCTAAGGGATGTGAGAATCCTTAATTAA Homologue amino acid sequence (SEQ ID NO: 6): MAEAEAPNRSPEIRKDIRHNWVLFSPARSRPSPDFKAKSNPQPNKTECPFCAGHEHECAP EIFRPVADSTADWKIRVIQNLFPVSRLEFQNSVSLGDDVAASGF GFHDVVIESPVHSINMSDLSPSQVGEVMLAFKKRVEQLQNCNDVKYVQVFNKHGASAGASMS HSHSQMIALPIVPPAVSARLDSMREYYKQTKGKSLCDIQPNELLID QSAHFISLVPPAATPAFEIWIIPRDHSHFHELDNEKANDLGGLLKLVLKMSLQLNNPPFN LLIHTSPFQVDPSPYAPSTHWFLQIAPHLSGVGGFELATGCHINPVF PEDAAKILRDVRILN

[0110] Suitably, the protein for use according to the present invention may be encoded by a polynucleotide sequence from *Nicotiana tabacum*.

[0111] In one aspect the present invention provides a method of decreasing the alkaloid content of a plant or part thereof or cell (e.g. plant cell), the method comprising modifying said plant by decreasing or inhibiting the activity or expression of at least one Galactose-1-phosphate uridyl transferase.

[0112] In one aspect the present invention provides a method of decreasing the alkaloid content of a plant or part thereof or plant cell, the method comprising modifying said plant by decreasing or inhibiting the activity or expression of at least one Galactose-1-phosphate uridyl transferase comprising the amino acid sequence shown as SEQ ID No. 3, or a sequence which has at least 80% identity thereto or wherein the at least one gene encoding a Galactose-1-phosphate uridyl transferase comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional

variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2.

[0113] In one aspect the present invention provides a method of decreasing the content of a tobacco specific nitrosamine (TSNA) precursor in a plant or part thereof (e.g. leaf), the method comprising modifying said plant by decreasing or inhibiting the activity or expression of at least one Galactose-1-phosphate uridyl transferase. Suitably, the method comprising modifying said plant by decreasing or inhibiting the activity or expression of at least one Galactose-1-phosphate uridyl transferase comprising the amino acid sequence shown as SEQ ID No. 3, or a sequence which has at least 80% identity thereto or wherein the at least one gene encoding a Galactose-1-phosphate uridyl transferase comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2.

[0114] In one aspect the present invention provides a method of decreasing the content of a tobacco specific nitrosamine (TSNA) precursor in a plant or part thereof (e.g. leaf), the method comprising modifying said plant by decreasing or inhibiting the activity or expression of at least one Galactose-1-phosphate uridyl transferase.

[0115] In one aspect the present invention provides a method of decreasing the content of a TSNA in a processed leaf, such as a cured leaf, the method comprising:

[0116] modifying a plant by decreasing or inhibiting the activity or expression of at least one Galactose-1-phosphate uridyl transferase;

[0117] harvesting a leaf from said plant;

[0118] and processing e.g. curing said harvested leaf.

[0119] Suitably, the method of decreasing the content of a TSNA in a processed leaf may comprise: modifying said plant by decreasing or inhibiting the activity or expression of at least one Galactose-1-phosphate uridyl transferase comprising the amino acid sequence shown as SEQ ID No. 3, or a sequence which has at least 80% identity thereto or wherein the at least one gene encoding a Galactose-1-phosphate uridyl transferase comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2. The term “decreasing” or “inhibiting” (e.g. inhibiting the activity or expression of Galactose-1-phosphate uridyl transferase) as used herein means that the activity or expression of the gene encoding the Galactose-1-phosphate uridyl transferase is lower or decreased compared with the activity or expression of the gene in a comparable product.

[0120] In one aspect, the present invention provides a method of increasing the alkaloid content of a plant or part thereof or cell (e.g. plant cell), the method comprising modifying said plant by increasing or enhancing the activity or expression of at least one gene encoding a Galactose-1-phosphate uridyl transferase.

[0121] In one aspect, the present invention provides a method of increasing the alkaloid content of a plant or part thereof or plant cell, the method comprising modifying said plant by increasing or enhancing the activity or expression of at least one Galactose-1-phosphate uridyl transferase comprising the amino acid sequence shown as SEQ ID No. 3, or a sequence which has at least 80% identity thereto or wherein the at least one gene encoding a Galactose-1-phosphate uridyl transferase comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2. In one aspect the present invention provides a method of increasing the content of a tobacco specific nitrosamine (TSNA) precursor in a plant or part thereof (e.g. leaf), the method comprising modifying said plant by increasing or enhancing the activity or expression of at least one Galactose-1-phosphate uridyl transferase.

[0122] In one aspect the present invention provides a method of increasing the content of a tobacco specific nitrosamine (TSNA) precursor in a plant or part thereof (e.g. leaf), the method comprising modifying said plant by increasing or enhancing the activity or expression of at least one Galactose-1-phosphate uridyl transferase comprising the amino acid sequence shown as SEQ ID No. 3, or a

sequence which has at least 80% identity thereto, or wherein the at least one gene encoding an Galactose-1-phosphate uridyl transferase comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2.

[0123] The term “increasing” or “enhancing” (e.g. increasing the activity or expression of at least one gene encoding a Galactose-1-phosphate uridyl transferase) as used herein means that the activity or expression of the gene encoding the Galactose-1-phosphate uridyl transferase is higher or increased compared with the activity or expression of the gene in a comparable product. According to the present invention, the activity or expression of Galactose-1-phosphate uridyl transferase is modulated.

[0124] In one aspect the present invention provides a method of modulating (i.e. increasing or decreasing) the alkaloid content of a plant or part thereof or cell (e.g. plant cell), the method comprising modifying said plant by modulating (i.e. increasing or decreasing) the activity of at least one Galactose-1-phosphate uridyl transferase.

[0125] The term “activity” refers to any functionality of the Galactose-1-phosphate uridyl transferase. For example the activity of the Galactose-1-phosphate uridyl transferase may refer to its ability to simultaneously catalyse conversion of uridine diphosphoglucose (UDP-glucose) and galactose-1-phosphate (gal-1P) to uridine diphosphogalactose (UDP-galactose) and glucose-1-phosphate. Methods for measuring such activity are known in the art.

[0126] Modulation of the activity of Galactose-1-phosphate uridyl transferase may entail increasing or decreasing the activity of the Galactose-1-phosphate uridyl transferase.

[0127] Increasing the activity of Galactose-1-phosphate uridyl transferase refers to enhancing or improving the ability of the Galactose-1-phosphate uridyl transferase to carry out a particular function in comparison to Galactose-1-phosphate uridyl transferase in a plant that has not been modified in accordance with the invention.

[0128] Decreasing the activity of Galactose-1-phosphate uridyl transferase refers to reducing, inhibiting or disrupting the ability of the Galactose-1-phosphate uridyl transferase to carry out a particular function in comparison to Galactose-1-phosphate uridyl transferase in a plant that has not been modified in accordance with the invention. The activity of Galactose-1-phosphate uridyl transferase may be reduced to such an extent that the activity is prevented or eliminated.

[0129] In some embodiments the activity of Galactose-1-phosphate uridyl transferase be modulated (i.e. increased or decreased) by at least about 10% 20% 30%, or 40%, suitably at least about 50%, 60%, 70%, more suitably at least about 80%, 90%, 95% or 100% in comparison to the activity of a gene encoding a Galactose-1-phosphate uridyl transferase in a plant (e.g. a tobacco plant) which has not been modified in accordance with the present invention. Suitably, the activity may be decreased.

[0130] In some embodiments the modulated Galactose-1-phosphate uridyl transferase exhibits increased or decreased activity compared to an unmodified Galactose-1-phosphate uridyl transferase. The modulated Galactose-1-phosphate uridyl transferase may exhibit at least about 1%, at least about 3%, at least about 5%, at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least

about 80% or at least about 90% increased or decreased activity compared to an unmodified Galactose-1-phosphate uridyl transferase.

[0131] Techniques are known in the art for measuring protein activities, for example methods for measuring Galactose-1-phosphate uridyl transferase activity are described in Mayes and Hanson *supra*.

[0132] In one aspect the present invention provides a method of modulating (i.e. increasing or decreasing) the alkaloid content of a plant or part thereof or cell (e.g. plant cell), the method comprising modifying said plant by modulating (i.e. increasing or decreasing) the expression of at least one Galactose-1-phosphate uridyl transferase.

[0133] The “expression” of a gene refers to the degree to which the information encoded in the gene is converted to a functionality. The level of expression of a gene may be equated with the amount of the product of that gene present in a cell or organism. A modification that modulates (i.e. increases or decreases) the expression of a gene is one that increases the amount of the product of that gene in a plant or cell in comparison to an unmodified plant or cell.

[0134] In some embodiments the expression of Galactose-1-phosphate uridyl transferase gene is modulated (i.e. increased or decreased) in comparison to the expression of a gene encoding a Galactose-1-phosphate uridyl transferase in a plant (e.g. a tobacco plant) which has not been modified in accordance with the present invention.

[0135] In some embodiments the expression of Galactose-1-phosphate uridyl transferase gene may be modulated (i.e. increased or decreased) by at least about 10% 20% 30%, or 40%, suitably at least about 50%, 60%, 70%, more suitably at least about 80%, 90%, 95% or 100% in comparison to the expression of a gene encoding a Galactose-1-phosphate uridyl transferase in a plant (e.g. a tobacco plant) which has not been modified in accordance with the present invention. Suitably, expression may be decreased.

[0136] In some embodiments, the modulated Galactose-1-phosphate uridyl transferase exhibits increased or decreased expression compared to an unmodified Galactose-1-phosphate uridyl transferase. The modulated Galactose-1-phosphate uridyl transferase may exhibit at least about 1%, at least about 3%, at least about 5%, at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80% or at least about 90% increased or decreased expression compared to an unmodified Galactose-1-phosphate uridyl transferase. Suitably, expression may be decreased.

[0137] Typically, genes are transcribed to mRNA, which is translated to protein, the final gene product.

[0138] Proteins may be sequestered in cellular stores and/or degraded. The expression of a gene may be modulated by modulating any or all of these steps. Accordingly, in some embodiments the modification modulates (e.g. decreases) expression of at least one gene encoding a Galactose-1-phosphate uridyl transferase in one of the following ways:

[0139] modulating (e.g. decreasing) transcription from the at least one gene encoding a Galactose-1-phosphate uridyl transferase;

[0140] modulating (e.g. decreasing) translation of the mRNA from the at least one gene encoding a Galactose-1-phosphate uridyl transferase;

[0141] modulating (e.g. decreasing) release of the Galactose-1-phosphate uridyl transferase from intracellular stores;

[0142] modulating (e.g. increasing) the rate of degradation of the Galactose-1-phosphate uridyl transferase and/or

[0143] introducing a mutation which modifies the amino acid sequence of the Galactose-1-phosphate uridyl transferase to decrease or increase its activity e.g. to decrease Galactose-1-phosphate uridyl transferase activity.

[0144] The expression of specific genes encoding Galactose-1-phosphate uridyl transferase can be measured by measuring transcription and/or translation of the gene. Methods for measuring transcription are well known in the art and include, amongst others, northern blot, RNA-Seq, in situ hybridization, DNA microarrays and RT-PCR. Alternatively, the expression of a gene may be measured indirectly by measuring the level of the gene product for example the protein encoded by said gene. For example, the expression of Galactose-1-phosphate uridyl transferase may be determined by measuring the presence of the protein using an antibody specific for the Galactose-1-phosphate uridyl transferase by western blot.

Modifying

[0145] The plant or cell may be modified in any way that modulates activity or expression of at least one Galactose-1-phosphate uridyl transferase. Types of modifications to plants and cells that modulate activity or expression of genes, as well as techniques to achieve those modifications, are known in the art.

[0146] In some embodiments the present invention provides a method of decreasing the alkaloid content of a plant or part thereof or cell (e.g. plant cell), the method comprising modifying said plant by decreasing or inhibiting the activity or expression of at least one Galactose-1-phosphate uridyl transferase as described herein.

[0147] In some embodiments the present invention provides a method of decreasing the content of a tobacco specific nitrosamine (TSNA) or a precursor of a TSNA in a tobacco plant or plant part thereof, the method comprising modifying said plant or a cell culture by decreasing the activity or expression of at least one Galactose-1-phosphate uridyl transferase as described herein.

[0148] Any method known in the art for decreasing or inhibiting the activity or expression of a gene or protein may be used in the methods according to the present invention.

[0149] Suitably, the activity or expression of the gene encoding a Galactose-1-phosphate uridyl transferase may be reduced, partly inactivated, inhibited, eliminated, knocked out or lost such that the protein activity, expression or function of the gene encoding a Galactose-1-phosphate uridyl transferase may not be detectable.

[0150] In one aspect, the at least one gene encoding a Galactose-1-phosphate uridyl transferase is knocked out. In other words, the gene encoding a Galactose-1-phosphate uridyl transferase has been rendered completely inoperative.

[0151] By way of example, the present method may comprise:

[0152] providing a mutation in a nucleic acid sequence which encodes a protein comprising the amino acid

sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto;

[0153] providing a mutation in a regulatory region (e.g. a promoter or an enhancer) which contributes to controlling the expression of a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto;

[0154] providing an antisense RNA, siRNA or miRNA which reduces the level of nucleic acid sequence encoding a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto.

[0155] Each of the above approaches results in the reduction or prevention of activity or expression of a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto or wherein the at least one gene encoding a Galactose-1-phosphate uridyl transferase comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2.

[0156] As used herein, the term “mutation” encompasses a natural genetic variant or an engineered variant. In particular, the term “mutation” refers to a variation in the nucleotide sequence encoding the amino acid sequence or in the amino acid sequence compared to the sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% (preferably at least 85%, preferably at least 90%, preferably at least 93%, preferably at least 95%, preferably at least 98%, preferably at least 99%) sequence identity thereto.

[0157] In one embodiment the mutation decreases the alkaloid content of a plant. In another embodiment, the mutation decreases the content of at least one TSNA precursor in a plant or part thereof, or leaf such as a harvested or processed leaf. In one embodiment the mutation decreases the content of one or more TSNA selected from NNN, NNK, NAT, NAB, preferably NNN and/or NNK content is decreased in a processed leaf. Suitably, the TSNA content is reduced in relation to a comparable product.

[0158] In one embodiment, a method according to the present invention may comprise providing a nucleic acid sequence to a plant or part thereof or plant cell, wherein said nucleic acid results in the reduction or elimination of the activity or expression of at least one Galactose-1-phosphate uridyl transferase.

[0159] In one embodiment, a method according to the present invention may comprise providing a nucleic acid sequence to a plant or part thereof or plant cell, wherein said nucleic acid results in the modification of the nucleic acid sequence of at least one Galactose-1-phosphate uridyl transferase.

[0160] Suitably said nucleic acid sequence may be introduced to the plant or part thereof or cell. Suitably an endogenous nucleic acid sequence in the plant or part thereof or cell may be modified to encode the polypeptide according to the present invention (e.g. by gene editing). For example, an endogenous nucleotide sequence may be modified to decrease the activity or expression of at least one Galactose-1-phosphate uridyl transferase.

[0161] In a preferred embodiment, each copy of a nucleic acid sequence encoding a protein comprising a sequence shown as SEQ ID No. 3, or a sequence which has at least 80% sequence identity thereto or wherein the at least one gene encoding a Galactose-1-phosphate uridyl transferase comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2, which is present in the plant is modified e.g. inhibited or mutated. Suitably, some or all of the homologues listed in Table 1, or corresponding sequences which have at least 80% sequence identity thereto are modified e.g. inhibited or mutated.

[0162] In a preferred embodiment, some or all of the homologues of the Galactose-1-phosphate uridyl transferase as described herein are modified e.g. inhibited or mutated. Suitably, some or all of the homologues listed in Table 1, or corresponding sequences which have at least 80% sequence identity thereto are modified e.g. inhibited or mutated.

[0163] In some embodiments the plant or plant cell according to the present invention is homozygous. Suitably, the plant or plant cell may be homozygous for the modification e.g. inhibition or mutation.

[0164] In some embodiments the plant or plant cell according to the present invention expresses only the modified e.g. mutated nucleic acid encoding at least one Galactose-1-phosphate uridyl transferase. In other words, in some embodiments no endogenous (or endogenous and functional protein) is present in the plant according to the present invention. In other words, if any endogenous protein is present it is preferably in an inactive form.

[0165] In one embodiment the present method may comprise providing a mutation in the nucleic acid sequence shown as SEQ ID No. 1, or 2, or a nucleic acid sequence which has at least 80% identity thereto, or a homologue of SEQ ID No. 1 or SEQ ID No. 2.

[0166] The mutation may alter the plant genome such that a nucleic acid sequence encoding a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto, or a homologue of SEQ ID No. 3, is completely or partially deleted or otherwise modified to inhibit or eliminate the activity of the Galactose-1-phosphate uridyl transferase. In some embodiments the mutation does not alter the level or expression of the protein but reduces inhibits or eliminates the activity of the Galactose-1-phosphate uridyl transferase.

[0167] Suitably, at least one mutation may be in a non-cytoplasmic domain, a cytoplasmic domain or a transmembrane domain of the Galactose-1-phosphate uridyl transferase.

[0168] The mutation may interrupt the nucleic acid sequence which encodes a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto, or a homologue of SEQ ID No. 3.

[0169] The interruption may cause the nucleic acid sequence to not be transcribed and/or translated. The nucleic acid sequence may be interrupted, for example, by deleting or otherwise modifying the ATG start codon of the nucleic acid sequence such that translation of the protein is reduced or prevented.

[0170] The nucleic acid sequence may comprise one or more nucleotide change(s) that reduce or prevent expression of the protein or affect protein trafficking. For example, expression of the protein may be reduced or prevented by introduction of one or more pre-mature stop codons, a frame shift, a splice mutation or a non-tolerated amino acid substitution in the open reading frame. A premature stop codon refers to a mutation which introduces a stop codon into the open reading frame and prevents translation of the entire amino acid sequence. The premature stop codon may be a TAG (“amber”), TAA (“ochre”), or TGA (“opal” or “umber”) codon.

[0171] A frame-shift mutation (also called a framing error or a reading frame shift) is a mutation caused by indels (insertions or deletions) of a number of nucleotides in a nucleic acid sequence that is not divisible by three. Due to the triplet nature of gene expression by codons, the insertion or deletion can change the reading frame, resulting in a completely different translation from the original. A frame-shift mutation will often cause the reading of the codons after the mutation to code for different amino acids. The frameshift mutation will commonly result in the introduction of a premature stop codon.

[0172] A splice mutation inserts, deletes or changes a number of nucleotides in the specific site at which splicing takes place during the processing of precursor messenger RNA into mature messenger RNA. The deletion of the splicing site results in one or more introns remaining in mature mRNA and may lead to the production of abnormal proteins.

[0173] A non-tolerated amino acid substitution refers to a mutation which causes a non-synonymous amino acid substitution in the protein which results in reduced or ablated function of the protein.

[0174] Any method known in the art for providing a mutation in a nucleic acid sequence may be used in the method according to the present invention. For example, homologous recombination may be used, in which a vector is created in which the relevant nucleic acid sequence(s) are mutated and used to transform plants or plant cells. Recombinant plants or plant cells expressing the mutated sequence may then be selected.

[0175] In one embodiment the mutation introduces a non-tolerated amino acid substitution in a protein comprising an amino acid sequence shown as SEQ ID No. 3, or a sequence which has at least 80% sequence identity thereto, or a homologue of SEQ ID No. 3.

[0176] In some embodiments, the Galactose-1-phosphate uridyl transferase may contain a mutation which decreases the expression or activity of the Galactose-1-phosphate uridyl transferase. The mutation may be a deletion, a splice mutant or codon encoding a non-tolerated amino acid substitution.

[0177] In one embodiment, the nucleic acid sequence encoding the Galactose-1-phosphate uridyl transferase may be wholly or partially deleted. The deletion may be continuous, or may comprise a plurality of sections of sequence. The deletion preferably removes a sufficient amount of nucleotide sequence such that the nucleic acid sequence no longer encodes a functional Galactose-1-phosphate uridyl transferase. The deletion may be total, in which case 100% of the coding portion of the nucleic acid sequence is absent, when compared to the corresponding genome of a comparable unmodified plant. The deletion may, for example,

remove at least 50, 60, 70, 80 or 90% of the coding portion of the nucleic acid sequence. Suitably, at least part of the protein may be deleted. The deletion may, for example, remove at least 10, 20, 30, 40, 50, 60, 70, 80 or 90% of the coding portion of the protein.

[0178] The deletion may remove at least 10 amino acids (such as at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90 amino acids) from the Galactose-1-phosphate uridyl transferase. Suitably, the deletion may remove at least 10 amino acids (such as at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90 amino acids) of the Galactose-1-phosphate uridyl transferase wherein the sequence of the Galactose-1-phosphate uridyl transferase is aligned with SEQ ID No. 3. Suitably, the deletion may remove at least 10 amino acids (such as at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90 amino acids) from the Galactose-1-phosphate uridyl transferase wherein the Galactose-1-phosphate uridyl transferase prior to deletion comprises an amino acid sequence set forth in SEQ ID No. 3, or a sequence which has at least 80% sequence identity thereto, or a homologue of SEQ ID No. 3. Suitably, the Galactose-1-phosphate uridyl transferase prior to deletion comprises an amino acid sequence set forth in SEQ ID No. 3.

[0179] Suitably, the protein for use according to the present invention may comprise a truncated Galactose-1-phosphate uridyl transferase. Suitably, the truncated protein may be a truncated version of an amino acid sequence set forth in SEQ ID No. 3, or a sequence which has at least 80% sequence identity thereto, or a homologue of SEQ ID No. 3. Suitably, the truncated protein lacks at least 10 amino acids (such as at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90 amino acids, at least 100 amino acids, at least 110 amino acids) from the Galactose-1-phosphate uridyl transferase.

[0180] The deletion may remove at least part of a domain. The deletion may for example, remove at least 10, 20, 30, 40, 50, 60, 70, 80 or 90% of a domain. Suitably, the deletion may remove at least 5 amino acids, at least 10 amino acids, at least 15, at least 20, at least 25, at least 30 amino acids, at least 40 amino acids, at least 50 amino acids, at least 60 amino acids, at least 70 amino acids, at least 80 amino acids of a domain. Suitably, the deletion may remove 5 amino acids, 10 amino acids, 15, 20 amino acids, 25 amino acids, 30 amino acids, 40 amino acids, 50 amino acids, 60 amino acids, 70 amino acids, 80 amino acids of a domain.

[0181] The deletion may remove at least part of a domain. The deletion may, for example, remove at least one or at least two amino acids from a domain. Suitably, a domain may be completely deleted.

[0182] Methods for deletion of nucleic acid sequences in plants are known in the art. For example, homologous recombination may be used, in which a vector is created in which the relevant nucleic acid sequence(s) are missing and used to transform plants or plant cells. Recombinant plants or plant cells expressing the new portion of sequence may then be selected.

[0183] Plant cells transformed with a vector as described herein may be grown and maintained in accordance with well-known tissue culturing methods such as by culturing the cells in a suitable culture medium supplied with the necessary growth factors such as amino acids, plant hormones, vitamins, etc.

[0184] Modification of the nucleic acid sequence may be performed using targeted mutagenesis methods (also referred to as targeted nucleotide exchange (TNE) or oligo-directed mutagenesis (ODM)). Targeted mutagenesis methods include, without limitation, those employing zinc finger nucleases, TALENs (see WO2011/072246 and WO2010/079430), Cas9-like, Cas9/crRNA/tracrRNA, Cas9/gRNA, or other CRISPR systems (see WO 2014/071006 and WO2014/093622), meganucleases (see WO2007/047859 and WO2009/059195), or targeted mutagenesis methods employing mutagenic oligonucleotides, possibly containing chemically modified nucleotides for enhancing mutagenesis with sequence complementarity to the gene, into plant protoplasts (e.g., KeyBase® or TALENs).

[0185] Alternatively, mutagenesis systems such as TILLING (Targeting Induced Local Lesions IN Genomics; McCallum et al. (2000) *Nat. Biotech.* 18:455, and McCallum et al. (2000) *Plant Physiol.* 123, 439-442, both incorporated herein by reference) may be used to generate plant lines which comprise a gene encoding a protein having a mutation. TILLING uses traditional chemical mutagenesis (e.g. ethyl methanesulfonate (EMS) mutagenesis, which produces random mutations) followed by high-throughput screening for mutations. Thus, plants, seeds, cells and tissues comprising a gene having the desired mutation may be obtained.

[0186] The method may comprise the steps of mutagenizing plant seeds (e.g. EMS mutagenesis), pooling of plant individuals or DNA, PCR amplification of a region of interest, heteroduplex formation and high-throughput detection, identification of the mutant plant, sequencing of the mutant PCR product. It is understood that other mutagenesis and selection methods may equally be used to generate such modified plants. Seeds may, for example, be radiated or chemically treated and the plants may be screened for a modified phenotype.

[0187] Fast neutron deletion mutagenesis may be used in a reverse genetics sense (i.e. with PCR) to identify plant lines carrying a deletion in the endogenous gene. See for example Ohshima et al. (1998) *Virology* 213:472-481; Okubara et al. (1994) *Genetics* 137:867-874; and Quesada et al. (2000) *Genetics* 154:421-4315 which are incorporated herein by reference.

[0188] In another approach, dominant mutants may be used to trigger RNA silencing due to gene inversion and recombination of a duplicated gene locus. See for example Kusaba et al. (2003) *Plant Cell* 15:1455-1467 (incorporated herein by reference).

[0189] Modified plants may be distinguished from non-modified plants, i.e., wild type plants, by molecular methods, such as the mutation(s) present in the DNA, and by the modified phenotypic characteristics. The modified plants may be homozygous or heterozygous for the modification. Preferably modified plants are homozygous for the modification.

[0190] In one embodiment the method of reducing or preventing the activity or expression of a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto; or a homologue of SEQ ID No. 3 does not comprise treating the plant with a chemical (e.g. an agrochemical).

[0191] Other ways of reducing or preventing the expression will be apparent to one skilled in the art and include the

use of virus-induced gene silencing (VIGs), micro RNA silencing, RNAi, antisense, tDNA insertions, or dominant negative constructs (or antimorphic mutations).

[0192] In one embodiment the expression of Galactose-1-phosphate uridyl transferase, such as a gene encoding a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto, may be reduced or eliminated by virus-induced gene silencing.

[0193] In one embodiment the expression of Galactose-1-phosphate uridyl transferase, such as a gene encoding a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto, may be reduced or eliminated by microRNAs.

[0194] In one embodiment the expression of Galactose-1-phosphate uridyl transferase, such as a gene encoding a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto, may be reduced or eliminated by RNAi.

[0195] In one embodiment the expression of Galactose-1-phosphate uridyl transferase, such as a gene encoding a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto, may be reduced or eliminated by antisense suppression.

[0196] In one embodiment the expression of Galactose-1-phosphate uridyl transferase, such as a gene encoding a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto, may be reduced or eliminated by sense suppression.

[0197] In one embodiment the expression of Galactose-1-phosphate uridyl transferase, such as a gene encoding a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto, may be reduced or eliminated by tDNA insertions.

[0198] In one embodiment the expression of Galactose-1-phosphate uridyl transferase, such as a gene encoding a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto, may be reduced or eliminated by dominant negative constructs (or antimorphic mutations).

[0199] In one embodiment the expression of Galactose-1-phosphate uridyl transferase, such as a gene encoding a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto may be reduced or eliminated by a targeted mutagenesis based system.

[0200] In one embodiment the expression of Galactose-1-phosphate uridyl transferase, such as a gene encoding a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto, may be reduced or eliminated by a gene editing e.g. CRISPR based system.

[0201] In one embodiment the expression of a gene encoding a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% sequence identity thereto may be reduced or eliminated by zinc finger nuclease, TALENs, meganucleases, mutagenic oligonucleotides or TILLING.

[0202] In some embodiments the present invention provides a method of increasing the alkaloid content of a plant or part thereof or cell (e.g. plant cell), the method comprising modifying said plant by increasing or enhancing the activity or expression of at least one Galactose-1-phosphate uridyl transferase.

[0203] Any method known in the art for increasing or enhancing the activity or expression of a gene may be used in the methods according to the present invention.

[0204] In some embodiments the method may comprise overexpressing at least one gene encoding a Galactose-1-phosphate uridyl transferase. Suitably the method may comprise expressing one or more additional copies of the at least one gene encoding a Galactose-1-phosphate uridyl transferase in the plant or cell. Suitably the method may comprise modifying the endogenous copy of the at least one gene encoding a Galactose-1-phosphate uridyl transferase such that its expression is increased. The method may comprise mutating the coding sequence of the at least one gene encoding a Galactose-1-phosphate uridyl transferase. The method may comprise mutating a regulatory sequence that regulates expression of the at least one gene encoding a Galactose-1-phosphate uridyl transferase.

[0205] Suitably the method may comprise transforming a cell of a plant (e.g. a tobacco plant) with a genetic construct which encodes at least one Galactose-1-phosphate uridyl transferase comprising an amino acid sequence as set out in SEQ ID No. 3, or a functional variant or functional fragment or orthologue thereof, or a sequence which has at least 80% identity to SEQ ID No. 3; or a homologue of SEQ ID No. 3; or wherein the at least one gene encoding a Galactose-1-phosphate uridyl transferase comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2; or which comprises a nucleotide sequence which encodes a protein which is capable of promoting or augmenting at least one endogenous Galactose-1-phosphate uridyl transferase. It will be appreciated that each of these options would result in an increased activity and expression of the polypeptide encoded by the at least one Galactose-1-phosphate uridyl transferase. The method may comprise regenerating the plant from the transformed cell. There is provided use of genetic construct which is capable of increasing the activity and/or expression of a polypeptide encoded by at least one gene encoding a Galactose-1-phosphate uridyl transferase for increasing the alkaloid content (e.g. nicotine content) in a plant or part thereof or cell transformed with the construct.

[0206] The genetic construct may encode a polypeptide comprising the amino acid SEQ ID No. 3, or a functional variant or functional fragment or orthologue thereof, or a sequence which has at least 80% identity to SEQ ID No. 3; or comprise a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2.

[0207] In another embodiment, the invention relates to a method of increasing the alkaloid content of a plant or part thereof or a cell, comprising modifying said plant or cell by increasing the activity of at least one Galactose-1-phosphate uridyl transferase.

[0208] In one embodiment the activity of at least one gene encoding a Galactose-1-phosphate uridyl transferase may be increased by introducing (or providing) a mutation to at least one gene encoding a Galactose-1-phosphate uridyl transferase.

[0209] Suitably, the activity of at least one gene encoding a Galactose-1-phosphate uridyl transferase may be increased by introducing a mutation to at least one gene encoding a Galactose-1-phosphate uridyl transferase which comprises an amino acid sequence as set out in SEQ ID No. 3; or a functional variant or functional fragment or orthologue thereof, or a sequence which has at least 80% identity to SEQ ID No. 3; or wherein the at least one gene encoding a Galactose-1-phosphate uridyl transferase comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2.

[0210] In some embodiments a modification which increases the activity or expression of at least one Galactose-1-phosphate uridyl transferase and thereby increases alkaloid content by one of the following:

[0211] modulating (e.g. increasing) transcription from the at least one gene encoding a Galactose-1-phosphate uridyl transferase;

[0212] modulating (e.g. increasing) translation of the mRNA from the at least one gene encoding a Galactose-1-phosphate uridyl transferase;

[0213] modulating (e.g. increasing) release of the Galactose-1-phosphate uridyl transferase from intracellular stores; and/or

[0214] modulating (e.g. decreasing) the rate of degradation of the Galactose-1-phosphate uridyl transferase.

Alkaloid Content

[0215] In one embodiment the present invention provides a method of modulating the alkaloid content of a plant (e.g. a tobacco plant) or a part thereof, the method comprising modifying said plant by modulating the activity or expression of at least one Galactose-1-phosphate uridyl transferase. The term “modulating” is used herein to mean either increasing or decreasing.

[0216] The term “increasing alkaloid content” is used herein to mean that the alkaloid content in the product of the present invention (e.g. plant, part thereof (e.g. leaf), processed leaf or a product made from the plant (e.g. a tobacco industry product)) is higher compared with a comparable product which has not been modified in accordance with the present invention.

[0217] The term “decreasing alkaloid content” is used herein to mean that alkaloid content in the product of the present invention (e.g. plant, part thereof (e.g. leaf), processed leaf or a product made from the plant (e.g. a tobacco industry product)) is lower compared with a comparable product which has not be modified in accordance with the present invention.

[0218] In some embodiments, the modulation of alkaloid content refers to an increase in alkaloid content wherein the activity or expression of at least one Galactose-1-phosphate uridyl transferase is increased (or example the protein is overexpressed).

[0219] In some embodiments, the modulation of alkaloid content refers to a decrease in alkaloid content wherein the

expression of at least one Galactose-1-phosphate uridyl transferase is decreased or inhibited or eliminated.

[0220] In a further aspect, the alkaloid content is measured from leaves. In one aspect the alkaloid content is measured from green leaves. In a further aspect, the alkaloid content is measured from cured leaves, e.g. air-cured, flue-cured, fire-cured or sun-cured leaves. In a further aspect, the alkaloid content is measured from flue-cured leaves. In a further aspect, the alkaloid content is measured from air-cured leaves.

[0221] The term “alkaloid content” is used herein to mean the concentration and/or total amount of the entire group of compounds classified as alkaloids or the concentration and/or total amount of one or more compounds classified as alkaloids. Alkaloids typically present in tobacco include nornicotine, PON, anatabine, anabasine, nicotine, and myosmine. In some embodiments the content of one or more alkaloids, such as two or more alkaloids, such as three or more alkaloids, such as four or more alkaloids, such as five or more alkaloids, such as all six alkaloids, selected from nicotine, nornicotine, PON, anatabine, anabasine and myosmine is modulated. In some embodiments the content of one or more alkaloids, such as two or more alkaloids, such as three or more alkaloids, such as four or more alkaloids, such as five or more alkaloids, such as all six alkaloids, selected from nicotine, nornicotine, PON, anatabine, anabasine and myosmine is increased. In some embodiments the content of one or more alkaloids, such as two or more alkaloids, such as three or more alkaloids, such as four or more alkaloids, such as five or more alkaloids, such as all six alkaloids, selected from nicotine, nornicotine, PON, anatabine, anabasine and myosmine is decreased. In some embodiments the total alkaloid content of the plant or cell is modulated. In some embodiments the total alkaloid content is increased. In some embodiments the total alkaloid content is increased.

[0222] Any method known in the art for determining the concentration and/or total content of alkaloids may be used. One preferred method for analysing alkaloid content involves the analysis by gas chromatography-flame ionization detection method (GC-FID) or by reversed phase high performance liquid chromatography with tandem mass spectrometry (LC-MS/MS).

[0223] In one embodiment there is provided a method for producing a plant (e.g. a tobacco plant) or part thereof, a plant propagation material (e.g. a tobacco plant propagation material), a cell (e.g. a tobacco cell), a leaf (e.g. a tobacco leaf), a harvested leaf (e.g. a harvested tobacco leaf), a cut harvested leaf (e.g. a cut harvested tobacco leaf), a processed leaf (e.g. a processed tobacco leaf), a cut and processed leaf (e.g. a cut and processed tobacco leaf), a product comprising said plant or part thereof (e.g. a tobacco industry product) or combinations thereof obtainable or obtained by a plant of the invention which has modulated alkaloid content, the method comprising modifying said plant to modulate the activity or expression of Galactose-1-phosphate uridyl transferase. The modulated alkaloid content may be determined by comparing the alkaloid content in the plant (e.g. tobacco plant) or part thereof, plant propagation material (e.g. tobacco plant propagation material), a cell (e.g. a tobacco cell), leaf (e.g. tobacco leaf), harvested leaf (e.g. a harvested tobacco leaf), cut harvested leaf (e.g. a cut harvested tobacco leaf), processed leaf (e.g. processed tobacco leaf), cut and processed leaf (e.g. cut and processed tobacco leaf), a product com-

prising a plant or part thereof of the present invention, e.g. a tobacco industry product, or combinations thereof with a comparable product.

[0224] Suitably the alkaloid content may be modulated in a plant, e.g. a tobacco plant e.g. modified tobacco plant. Suitably the alkaloid content may be modulated in a leaf (e.g. a tobacco leaf e.g. a tobacco leaf from a modified tobacco plant). Suitably the alkaloid content may be modulated in a harvested leaf (e.g. a harvested tobacco leaf from a modified tobacco plant). Suitably the alkaloid content may be modulated in a cut harvested leaf (e.g. a cut harvested tobacco leaf from a modified tobacco plant). Suitably the alkaloid content may be modulated in a processed leaf (e.g. a processed tobacco leaf e.g. a processed tobacco leaf from a modified tobacco plant). Suitably the alkaloid content may be modulated in a cut and processed leaf (e.g. a cut and processed tobacco leaf e.g. a cut and processed tobacco leaf from a modified tobacco plant). Suitably the alkaloid content may be modulated in a cured leaf (e.g. cured a tobacco leaf from a modified tobacco plant). Suitably the alkaloid content may be modulated in an extract of a green leaf (e.g. a green tobacco leaf from a modified tobacco plant). Suitably the alkaloid content may be modulated in a product comprising the plant of the present invention or part thereof (e.g. a tobacco industry product, for example a tobacco industry product produced from a modified tobacco plant or part thereof). Suitably the alkaloid content may be modulated in any one of the above products or combinations thereof. Suitably the modulation of alkaloid content described above may be an increase in alkaloid content. Suitably the modulation of alkaloid content described above may be a decrease in alkaloid content (e.g. a decrease in nornicotine and/or PON content).

[0225] In one embodiment the content of one or more alkaloids selected from nornicotine, PON, anatabine and anabasine is decreased. In one embodiment the content of nornicotine is decreased. In one embodiment the content of PON is decreased. In one embodiment the content of anatabine is decreased. In one embodiment the content of anabasine is decreased.

[0226] In one embodiment the nicotine content of a modified plant (e.g. tobacco plant), plant propagation material (e.g. tobacco plant propagation material), leaf (e.g. tobacco leaf), harvested leaf (e.g. harvested tobacco leaf), cut harvested leaf (e.g. cut harvested tobacco leaf), processed leaf (e.g. processed tobacco leaf), cut and processed leaf (e.g. cut and processed tobacco leaf) or tobacco industry product from a modified tobacco plant is not substantially decreased. Suitably, the nicotine content is at least 85% (such as at least 90%, such as at least 95%, such as at least 98%, such as at least 99%) of the nicotine content of a comparable product.

[0227] In one embodiment the alkaloid content of a plant (e.g. tobacco plant) or part thereof may be modulated by at least 0.5, 1.5, 2, 3 or 4 fold when compared to the alkaloid content of a plant (e.g. tobacco plant) or part thereof, respectively, which has not been modified to modulate the activity or expression of at least one gene encoding a Galactose-1-phosphate uridyl transferase and which has been grown under similar growth conditions. Suitably the alkaloid content may be modulated by about 0.5 fold to about 4 fold. Suitably the alkaloid content may be modulated by about 4 fold. Suitably the modification may be an increase or a decrease in alkaloid content.

[0228] Suitably the modulation may be of one or more alkaloids selected from nicotine, nornicotine, PON, anatabine, anabasine and myosmine. Suitably the modulation may be of one or more alkaloids selected from nicotine, nornicotine, PON, anatabine and anabasine. Suitably, the nornicotine content may be reduced. Suitably, the PON content may be reduced. Suitably, the anatabine content may be reduced. Suitably, the anabasine content may be reduced.

[0229] In one embodiment of the invention the alkaloid content of a plant (e.g. a tobacco plant) or part thereof may be modulated by at least 1%, 2%, 5%, 8%, 10%, 12%, 15%, 20%, 25%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 100% in comparison to a plant (e.g. a tobacco plant) or part thereof which has not been modified according to the present invention. In one embodiment the alkaloid content may be modulated by at least 30% in comparison to an unmodified plant or part thereof.

[0230] In one embodiment the alkaloid content may be modulated by at least 40% in comparison to an unmodified plant or part thereof. In one embodiment the alkaloid content may be modulated by at least 50% in comparison to an unmodified plant or part thereof. In one embodiment the alkaloid content may be modulated by at least 60% in comparison to an unmodified plant or part thereof. The modulation may be an increase or a decrease in alkaloid content when compared to an unmodified plant (e.g. a tobacco plant) or part thereof.

[0231] Suitably the modulation may be of total alkaloid content. Suitably the modulation may be of one or more alkaloids selected from nicotine, nornicotine, PON, anatabine, anabasine and myosmine. Suitably the modulation may be of one or more alkaloids selected from nornicotine, nicotine, PON, anatabine and anabasine. Suitably the modulation may be of nornicotine content, such as decrease in nornicotine content. Suitably the modulation may be of anabasine content, such as decrease in anabasine content. Suitably the modulation may be of PON content, such as decrease in PON content. Suitably the modulation may be of anatabine content, such as decrease in anatabine content.

[0232] Suitably the modulation may be of more than one alkaloid, such as two or more alkaloids, such as three or more alkaloids, such as four or more alkaloids, such as five or more alkaloids, such as all six alkaloids, selected from nicotine, nornicotine, PON, anatabine, anabasine and myosmine.

[0233] In some embodiments the alkaloid content of the plant may be modulated by between about 5% and about 100%, by between about 10% and about 90%, by between about 20% and about 80%, by between about 30% and about 70%, by between about 40% and 60%, by between about 40% and 50%, or by between about 50% and 60%.

Tobacco-Specific Nitrosamine (TSNA) Content

[0234] In one embodiment the present invention provides a method of decreasing the content of at least one tobacco-specific nitrosamine (TSNA) precursor in a plant (e.g. a tobacco plant) or a part thereof or in a tobacco cell. Suitably, the method may comprise modifying said plant by modulating the activity or expression of at least one Galactose-1-phosphate uridyl transferase. In one embodiment, the present invention provides a method of producing a processed leaf with decreased TSNA content (e.g. relative to a comparable product). The method of producing a processed leaf with decreased TSNA content may comprise:

[0235] modifying a plant by decreasing or inhibiting the activity or expression of at least one Galactose-1-phosphate uridyl transferase;

[0236] harvesting a leaf from said plant;

[0237] and processing e.g. curing said harvested leaf.

[0238] The TSNA may be measured in a processed tobacco, e.g. cured tobacco or reconstituted tobacco. In one embodiment the TSNA content is measured and/or modified (e.g. reduced) in a cured tobacco plant or part thereof (e.g. in cured tobacco leaf).

[0239] The term “tobacco-specific nitrosamine” or “TSNA” as used herein has its usual meaning in the art, namely a nitrosamine which is found only in tobacco industry products or other nicotine-containing products. Suitably the at least one tobacco-specific nitrosamine may be N'-nitrosonornicotine (NNN), 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK), N'-nitrosoanatabine (NAT) or N'-nitrosoanabasine (NAB).

[0240] The term “precursor thereto” when used in relation to at least one tobacco-specific nitrosamine refers to one or more chemicals or compounds of a tobacco plant that give rise to the formation of a tobacco-specific nitrosamine or are involved in the nitrosation reaction leading to tobacco-specific nitrosamine production.

[0241] In one embodiment the TSNA may be one or more of group selected from: N'-nitrosonornicotine (NNN), 4-(methyl nitrosamino)-1-(3-pyridyl)-1-butanone (NNK), N'-nitrosoanatabine (NAT) and N'-nitrosoanabasine (NAB). Suitably the at least one tobacco-specific nitrosamine may be NNK or NNN. In one embodiment the tobacco-specific nitrosamine is NNN. In another embodiment the tobacco-specific nitrosamine is NNK.

[0242] In one embodiment the precursor of the TSNA is one or more of the group selected from nornicotine, anabasine, anatabine, and an oxidised derivative of nicotine such as pseudooxynicotine (PON).

[0243] In one embodiment the TSNA is N'-nitrosonornicotine (NNN) and/or the precursor is nornicotine. In one embodiment the content of NNN is decreased. In one embodiment the content of nornicotine is decreased. In one embodiment the content of NNN and nornicotine is decreased. In one embodiment the TSNA is 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) and/or the precursor is PON. In one embodiment the content of NNK is decreased. In one embodiment the content of PON is decreased. In one embodiment the content of NNK and PON is decreased. In one embodiment the TSNA is N'-nitrosoanatabine (NAT) and/or the precursor is anatabine. In one embodiment the content of NAT is decreased. In one embodiment the content of anatabine is decreased. In one embodiment the content of NAT and anatabine is decreased.

[0244] In one embodiment the TSNA is N'-nitrosoanabasine (NAB) and/or the precursor is anabasine. In one embodiment the content of NAB is decreased. In one embodiment the content of nornicotine is decreased. In one embodiment the content of NAB and anabasine is decreased. The precursor of the TSNA (e.g. NNK, NNN, NAB and/or NAT) may be measured in green tobacco leaf, e.g. prior to processing, e.g. prior to curing. In one embodiment the precursor of the TSNA (e.g. NNK, NNN, NAB and/or NAT) is measured and/or modified (e.g. reduced) in a green tobacco leaf, e.g. prior to processing, e.g. prior to curing.

[0245] In one embodiment carrying out a method and/or use of the invention results in a reduction of at least one

TSNA or a precursor thereto in the modified tobacco plant (or part thereof) when compared to a tobacco plant (or part thereof) which has not been modified in accordance with the present invention.

[0246] The terms “reducing at least one TSNA or precursor thereto” or “reduction of at least one TSNA or precursor thereto” are used herein to mean that the concentration and/or total content of the at least one TSNA or precursor thereto in the product, method or use of the invention is lower in relation to a comparable product, method or use. For example, a comparable tobacco industry product would be derived from a tobacco plant which had not been modified according to the present invention, but in which all other relevant features were the same (e.g. plant species, growing conditions, method of processing tobacco, etc.).

[0247] Any method known in the art for determining the concentration and/or levels of at least one TSNA or precursor thereto may be used. In particular a method such may comprise the addition of deuterium labelled internal standard, an aqueous extraction and filtration, followed by analysis using reversed phase high performance liquid chromatography with tandem mass spectrometry (LC-MS/MS) may be used. Other examples for determining the concentration and/or level of a precursor to a tobacco-specific nitrosamine include a method such as the one detailed in CORESTA recommended method CRM-72: Determination of Tobacco Specific Nitrosamines in Tobacco and Tobacco Products by LC-MS/MS; CRM being developed into ISO/DIS 21766 or Wagner et al. (2005) Analytical Chemistry 77(4), 1001-1006 all of which are incorporated herein by reference.

[0248] Suitably the concentration and/or total content of the at least one tobacco-specific nitrosamine or precursor thereto may be reduced by carrying out a method and/or use of the present invention.

[0249] Suitably the concentration and/or level of the at least one tobacco-specific nitrosamine or precursor thereto may be reduced in a tobacco plant of the invention (e.g. obtainable or obtained by a method and/or use of the invention) when compared to the concentration and/or level of the at least one tobacco-specific nitrosamine(s) or precursor thereto in a tobacco plant which has not been modified in accordance with present invention.

[0250] The concentration and/or total content of the at least one tobacco-specific nitrosamine(s) or precursor thereto may be reduced in a tobacco leaf, harvested leaf, processed tobacco leaf, tobacco industry product or combinations thereof obtainable or obtained from a tobacco plant (or part of a tobacco plant or a tobacco cell culture) of the invention when compared with a tobacco leaf, harvested leaf, processed tobacco leaf, tobacco industry product or combinations thereof obtainable or obtained from a tobacco plant (or part of a tobacco plant or a tobacco cell culture) which has not been modified in accordance with the present invention.

[0251] Suitably the concentration and/or total content of the at least one tobacco-specific nitrosamine or precursor thereto may be reduced in a processed tobacco leaf.

[0252] Suitably the concentration and/or level of the at least one tobacco-specific nitrosamine or precursor thereto may be reduced in a tobacco industry product.

[0253] In one embodiment the at least one tobacco-specific nitrosamine or precursor thereto may be reduced by at least about 1%, at least about 3%, at least about 5%, at least

about 10%, at least about 20%, at least about 30%, at least about 40% or at least about 50%. In some embodiments the at least one tobacco-specific nitrosamine or precursor thereto may be reduced by between about 5% and about 50%, by between about 10% and about 50%, by between about 20% and about 50%, by between about 30% and about 50%, or by between about 40% and 50%.

[0254] In relation to processed (e.g. cured) tobacco leaf (e.g. cured or reconstituted), the at least one tobacco-specific nitrosamine or precursor thereto may be reduced by between about 5000 ng/g and about 50 ng/g, by between about 4000 ng/g and about 100 ng/g, by between about 3000 ng/g and 500 ng/g or by between 2000 ng/g and 1000 ng/g. In some embodiments the at least one tobacco-specific nitrosamine or precursor thereto may be reduced by at least about 5000 ng/g, at least about 4000 ng/g, at least about 3000 ng/g, at least about 2000 ng/g, at least about 1000 ng/g, at least about 500 ng/g, at least about 100 ng/g or at least about 50 ng/g.

Biomass Production

[0255] In some instances, it may be desirable to produce plants or biomass with high alkaloid levels e.g. high levels of nicotine content so that nicotine may be purified to produce a pure nicotine product for example for use in devices which utilize liquid containing nicotine (e.g. e-cigarettes) or within tobacco heating devices. For example, the production of nicotine in this way could reduce costs of nicotine extraction for the production of e-liquids for e-cigarettes.

[0256] In one aspect, the present invention provides a method of producing a biomass comprising: growing a cell which has been engineered to modulate (e.g. increase) the activity or expression of a gene encoding a Galactose-1-phosphate uridyl transferase under conditions to produce a biomass. Suitably, the activity or expression of Galactose-1-phosphate uridyl transferase may be increased in order to increase the concentration and/or total nicotine content.

[0257] In one embodiment, the present invention provides a method of producing a biomass having modified (e.g. increased) concentration and/or total content of nicotine, comprising growing a cell which has been engineered to increase the activity or expression of at least one Galactose-1-phosphate uridyl transferase comprising an amino acid sequence as set out in SEQ ID No. 3, or a functional variant or functional fragment or orthologue thereof, or a sequence which has at least 80% identity to SEQ ID No. 3; or a homologue of SEQ ID No. 3; or wherein the at least one Galactose-1-phosphate uridyl transferase comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2.

[0258] The cell may be engineered by any method known in the art to modify the activity or expression of at least one Galactose-1-phosphate uridyl transferase. Suitably, the cell may be engineered to express an exogenous gene encoding a Galactose-1-phosphate uridyl transferase. Suitably, the cell may be engineered to overexpress a gene encoding a Galactose-1-phosphate uridyl transferase.

[0259] Suitably, the biomass may contain a higher concentration and/or total content of nicotine compared with the biomass produced by a comparable cell which has not been modified in accordance with the present invention.

[0260] Suitably the cell for use in biomass production may be a plant cell, such as a tobacco cell.

[0261] Suitably the cell for use in biomass production may be a yeast cell.

[0262] In one embodiment the cell (e.g. yeast cell) may be further modified to comprise one or more sequences that increases nicotinic alkaloid biosynthesis. Suitably these one or more sequences may be incorporated into a nucleic acid construct that is suitable for cell (e.g. yeast cell) transformation. The one or more sequences may be overexpressed in the cell (e.g. yeast cell). The sequences may be selected from one or more of the following genes: MPO (or Methylputrescine Oxidase or MPO1 or MPO2); A622 (or Isoflavone reductase-like protein or Isoflavone reductase homolog or Isoflavone reductase-like protein); BBL (or Berberine bridge enzyme or Berberine bridge enzyme-like or BBE or NBB1); PMT (or Putrescine N-Methyltransferase or putrescine methyltransferase or S-adenosyl-L-methionine:putrescine N-methyltransferase or PMT or PMT1 or PMT2 or PMT3 or PMT4) and QPT (or quinolinate phosphoribosyltransferase). In one embodiment, the sequences may be selected from one or more of the following genes: BBL, A622, PMT and MPO (MPO1 or MPO2). Genes suitable for modification in this way may be taught in US2016032299 for example, which is incorporated herein by reference.

Commercially Desirable Traits

[0263] In one embodiment the plants of the present invention have modified (i.e. increased or decreased) total alkaloid content and/or modified (i.e. increased or decreased) content of one or more alkaloids, whilst the flavour characteristics and/or other commercially desirable traits are at least maintained. Suitably, the plants of the present invention may have decreased total alkaloid content and/or decreased content of one or more alkaloids, whilst the flavour characteristics and/or other commercially desirable traits are at least maintained.

[0264] In one embodiment, the plants of the present invention produce leaves of a similar grade and/or quality to plants which have not been modified according to the invention.

[0265] In one embodiment, the plants of the present invention have reduced nornicotine and/or PON and/or anabasine and/or anatabine content without a significant change in the flavour characteristics of the plant (e.g. compared with the same plant which has not been modified in accordance with the present invention).

[0266] In one embodiment the plants of the present invention have decreased TSNA precursor content without a significant change (e.g. decrease) in other commercially desirable traits of the plant (e.g. compared with the same plant which has not been modified in accordance with the present invention). In particular, the yield of the modified plant is preferably not reduced compared with the same plant which has not been modified in accordance with the present invention.

[0267] Therefore in one embodiment the methods and uses of the present invention relate to decreasing TSNA precursor content whilst maintaining the flavour characteristics and/or other commercially desirable traits (e.g. yield).

[0268] The term “commercially desirable traits” as used herein will include traits such as yield, mature plant height, harvestable leaf number, average node length, cutter leaf length, cutter leaf width, quality (e.g. leaf quality, suitably

cured leaf quality), abiotic (for instance drought) stress tolerance, herbicide tolerance and/or biotic (for instance insect, bacteria or fungus) stress tolerance.

[0269] Leaf quality may be measured based on colour, texture and aroma of the cured leaf, for example according to United States Department of Agriculture (USDA) grades and standards.

[0270] Tobacco grades are evaluated based on factors including, but not limited to, the leaf stalk position, leaf size, leaf colour, leaf uniformity and integrity, ripeness, texture, elasticity, sheen (related with the intensity and the depth of coloration of the leaf as well as the shine), hygroscopicity (the faculty of the tobacco leaves to absorb and to retain the ambient moisture), and green nuance or cast. Leaf grade can be determined using standard methods known in the art, for example, using an Official Standard Grade published by the Agricultural Marketing Service of the US Department of Agriculture (7 U.S.C. § 511). See, e.g., Official Standard Grades for Burley Tobacco (U.S. Type 31 and Foreign Type 93), effective Nov. 5, 1990 (55 F.R. 40645); Official Standard Grades for Flue-Cured Tobacco (U.S. Types 11, 12, 13, 14 and Foreign Type 92), effective Mar. 27, 1989 (54 F.R. 7925); Official Standard Grades for Pennsylvania Seedleaf Tobacco (U.S. Type 41), effective Jan. 8, 1965 (29 F.R. 16854); Official Standard Grades for Ohio Cigar-Leaf Tobacco (U.S. Types 42, 43, and 44), effective Dec. 8, 1963 (28 F.R. 11719 and 28 F.R. 11926); Official Standard Grades for Wisconsin Cigar-Binder Tobacco (U.S. Types 54 and 55), effective Nov. 20, 1969 (34 F.R. 17061); Official Standard Grades for Wisconsin Cigar-Binder Tobacco (U.S. Types 54 and 55), effective Nov. 20, 1969 (34 F.R. 17061); Official Standard Grades for Georgia and Florida Shade-Grown Cigar-Wrapper Tobacco (U.S. Type 62), Effective April 1971. A USDA grade index value can be determined according to an industry accepted grade index. See e.g. Bowman et al. (1988) Tobacco Science, 32:39-40; Legacy Tobacco Document Library (Bates Document #523267826-523267833, Jul. 1, 1988, Memorandum on the Proposed Burley Tobacco Grade Index); and Miller et al. (1990) Tobacco Intern., 192:55-57 (all foregoing references are incorporated herein in their entirety).

[0271] In one aspect, a USDA grade index is a 0-100 numerical representation of federal grade received and is a weighted average of all stalk positions. A higher grade index indicates higher quality. Alternatively, leaf grade may be determined via hyper-spectral imaging. See e.g. WO 2011/027315 (which is incorporated herein by reference).

[0272] In one embodiment, a tobacco plant of the present invention provides tobacco of commercially acceptable grade.

[0273] Suitably, the tobacco plant of the present invention provides cured tobacco of commercially acceptable grade.

[0274] In one embodiment, a tobacco plant of the present invention is capable of producing leaves having a USDA grade index value of at least about 70% of the USDA grade index value of leaves of a comparable plant when grown in similar growth conditions. Suitably, tobacco plants disclosed herein may be capable of producing leaves having a USDA grade index value of at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 98% of the USDA grade index value of a control plant when grown in similar growth conditions. Suitably, tobacco plants disclosed herein may be capable of producing leaves having

a USDA grade index value of between 65% and 130%, between 70% and 130%, between 75% and 130%, between 80% and 130%, between 85% and 130%, between 90% and 130%, between 95% and 130%, between 100% and 130%, between 105% and 130%, between 110% and 130%, between 115% and 130%, or between 120% and 130% of the USDA grade index value of a comparable plant.

[0275] In one aspect, the tobacco plant of the present invention is capable of producing leaves having a USDA grade index value of at least 50. Suitably, tobacco plants disclosed herein may be capable of producing leaves having a USDA grade index value of 55 or more, 60 or more, 65 or more, 70 or more, 75 or more, 80 or more, 85 or more, 90 or more, and 95 or more.

[0276] Unless specified otherwise, used herein, tobacco yield refers to cured leaf yield which is calculated based on the weight of cured tobacco leaves per acre under standard field conditions following standard agronomic and curing practice.

[0277] In one aspect, a plant (e.g. a tobacco plant) of the present invention has a yield between 50% and 150%, between 55% and 145%, between 60% and 140%, between 65% and 135%, between 70% and 130%, between 75% and 125%, between 80% and 120%, between 85% and 115%, between 90% and 110%, between 95% and 105%, between 50% and 100%, between 55% and 100%, between 60% and 100%, between 65% and 100%, between 70% and 100%, between 75% and 100%, between 80% and 100%, between 85% and 100%, between 90% and 100%, between 95% and 100%, between 100% and 150%, between 105% and 150%, between 110% and 150%, between 115% and 150%, between 120% and 150%, between 125% and 150%, between 130% and 150%, between 135% and 150%, between 140% and 150%, or between 145% and 150% of the yield of a comparable plant when grown in similar field conditions.

[0278] In another aspect, the plant (e.g. a tobacco plant) yield of the present invention is approximately 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, or 3.0 times of the yield of a comparable plant when grown in similar field conditions.

[0279] In another aspect, the yield of a tobacco plant of the present invention is comparable to the yield of the flue cured comparable plant when grown in similar field conditions.

[0280] In one aspect, a tobacco plant of the present invention provides a yield selected from the group consisting of about between 1200 and 3500, between 1300 and 3400, between 1400 and 3300, between 1500 and 3200, between 1600 and 3100, between 1700 and 3000, between 1800 and 2900, between 1900 and 2800, between 2000 and 2700, between 2100 and 2600, between 2200 and 2500, and between 2300 and 2400 lbs/acre.

[0281] In another aspect, a tobacco plant of the present invention provides a yield selected from the group consisting of about between 1200 and 3500, between 1300 and 3500, between 1400 and 3500, between 1500 and 3500, between 1600 and 3500, between 1700 and 3500, between 1800 and 3500, between 1900 and 3500, between 2000 and 3500, between 2100 and 3500, between 2200 and 3500, between 2300 and 3500, between 2400 and 3500, between 2500 and 3500, between 2600 and 3500, between 2700 and 3500, between 2800 and 3500, between 2900 and 3500, between 3000 and 3500, and between 3100 and 3500 lbs/acre.

[0282] In a further aspect, a tobacco plant of the present invention provides a yield selected from the group consisting of about between 1200 and 3500, between 1200 and 3400, between 1200 and 3300, between 1200 and 3200, between 1200 and 3100, between 1200 and 3000, between 1200 and 2900, between 1200 and 2800, between 1200 and 2700, between 1200 and 2600, between 1200 and 2500, between 1200 and 2400, between 1200 and 2300, between 1200 and 2200, between 1200 and 2100, between 1200 and 2000, between 1200 and 1900, between 1200 and 1800, between 1200 and 1700, between 1200 and 1600, between 1200 and 1500, and between 1200 and 1400 lbs/acre.

Plant Breeding

[0283] In one embodiment the present invention provides a method of producing a plant having a modified alkaloid content and/or modified content of a tobacco specific nitrosamine (TSNA) precursor comprising:

[0284] a. crossing a donor plant having modified (e.g. decreased) nicotine content and/or modified (e.g. decreased) content of a tobacco specific nitrosamine (TSNA) precursor and wherein the activity or expression of at least one Galactose-1-phosphate uridyl transferase according to the present invention has been modulated (e.g. decreased) in the donor plant in accordance with the present invention with a recipient tobacco plant that does not have modified nicotine content or modified content of a tobacco specific nitrosamine (TSNA) precursor and possesses commercially desirable traits;

[0285] b. isolating genetic material from a progeny of said donor plant crossed with said recipient plant; and

[0286] c. performing molecular marker-assisted selection with a molecular marker comprising:

[0287] i. identifying an introgressed region comprising a mutation in a polynucleotide sequence encoding a protein defined in a.

[0288] Suitably, the activity or expression of a protein comprising an amino acid sequence as set out in SEQ ID No. 3, or a functional variant or functional fragment or orthologue thereof, or a sequence which has at least 80% identity to SEQ ID No. 3; or a homologue of SEQ ID No. 3; or a protein encoded by a nucleotide sequence as set out in SEQ ID No. 1 or 2 or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2 or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2; is modulated in the donor plant when compared to a comparable plant. Suitably, the alkaloid content and/or TSNA precursor content is decreased by said method. Suitably, the alkaloid content and/or TSNA precursor content is decreased and the activity or expression of said Galactose-1-phosphate uridyl transferase is decreased or inhibited.

[0289] The molecular marker assisted selection may comprise performing PCR to identify an introgressed nucleic acid sequence comprising a mutation which modulates the activity or expression of a protein comprising the amino acid sequence shown as SEQ ID No. 3, or an amino acid sequence which has at least 80% identity thereto.

Tobacco Plants

[0290] The present invention provides methods, uses directed to plants (e.g. tobacco plants) as well as a cell (e.g.

a tobacco cell), a cell culture, a plant (e.g. a tobacco plant) and a plant propagation material.

[0291] The term “tobacco plant” as used herein refers to a plant in the genus *Nicotiana* that is used in the production of tobacco industry products. Non-limiting examples of suitable “tobacco” plants include *N. tabacum* and *N. rustica* (for example, *N. tabacum* L., LA B21, LN KY171, TI 1406, Basma, Galpao, Perique, Beinhart 1000-1, and Petico).

[0292] The tobacco material can be derived or obtained from varieties of *Nicotiana tabacum* types, commonly known as Burley varieties, flue or bright varieties and dark varieties. In some embodiments, the tobacco material is derived from a Burley, Virginia or a dark tobacco plant. The tobacco plant may be selected from Burley tobacco, rare tobacco, speciality tobacco, expanded tobacco or the like.

[0293] The use of tobacco cultivars and elite tobacco cultivars is also contemplated herein. The tobacco plant for use herein may therefore be a tobacco variety or elite tobacco cultivar. Particularly useful *Nicotiana tabacum* varieties include Flue-cured Virginia type, Burley type, and Oriental type.

[0294] In some embodiments, the tobacco plant may be, for example, selected from one or more of the following varieties: L. cultivar T.I. 1068, AA 37-1, B 13P, Xanthi (Mitchell-Mor), KT D #3 Hybrid 107, Bel-W3, 79-615, Samsun Holmes NN, F4 from cross BU21xHoja Parado, line 97, KTRDC #2 Hybrid 49, KTRDC #4 Hybrid 1 10, Burley 21, PM016, KTRDC #5 KY 160 SI, KTRDC #7 FCA, KTRDC #6 TN 86 SI, PMO21, K 149, K 326, K 346, K 358, K 394, K 399, K 730, KY 10, KY 14, KY 160, KY 17, KY 8959, KY 9, KY 907, MD 609, McNair 373, NC 2000, PG 01, PG 04, P01, P02, P03, RG 11, RG 17, RG 8, Speight G-28, TN 86, TN 90, VA 509, AS44, Banket AI, Basma Drama B84/31, Basma I Zichna ZP4/B, BasmaXanthi BX 2A, Batek, Besuki Jember, C104, Coker 319, Coker 347, Criollo Misionero, PM092, Delcrest, Djebel 81, DVH 405, Galpao Comum, HB04P, Hicks Broadleaf, Kabakulak Ellassona, PM102, Kutsage E1, KY 14xL8, KY 171, LA BU 21, McNair 944, NC 2326, NC 71, NC 297, NC 3, PVH 03, PVH 09, PVH 19, PVH 2110, Red Russian, Samsun, Saplak, Simmaba, Talgar 28, PM132, Wislica, Yayaldag, NC 4, TR Madole, Prilep HC-72, Prilep P23, Prilep PB 156/1, Prilep P12-2/1, Yaka JK-48, Yaka JB 125/3, TI-1068, KDH-960, TI-1070, TW136, PM204, PM205, Basma, TKF 4028, L8, TKF 2002, TN 90, GR141, Basma xanthi, GR149, GR153, and Petit Havana.

[0295] Non-limiting examples of varieties or cultivars are: BD 64, CC 101, CC 200, CC 27, CC 301, CC 400, CC 500, CC 600, CC 700, CC 800, CC 900, Coker 176, Coker 319, Coker 371 Gold, Coker 48, CD 263, DF91 1, DT 538 LC, Galpao tobacco, GL 26H, GL 350, GL 600, GL 737, GL 939, GL 973, HB 04P, HB 04P LC, HB3307PLC, Hybrid 403LC, Hybrid 404LC, Hybrid 501 LC, K 149, K 326, K 346, K 358, K394, K 399, K 730, KDH 959, KT 200, KT204LC, KY10, KY14, KY 160, KY 17, KY 171, KY 907, KY907LC, KTY14xL8 LC, Little Crittenden, McNair 373, McNair 944, msKY 14xL8, Narrow Leaf Madole, Narrow Leaf Madole LC, NBH 98, N-126, N-777LC, N-7371 LC, NC 100, NC 102, NC 2000, NC 291, NC 297, NC 299, NC 3, NC 4, NC 5, NC 6, NC7, NC 606, NC 71, NC 72, NC 810, NC BH 129, NC 2002, Neal Smith Madole, OXFORD 207, PD 7302 LC, PD 7309 LC, PD 7312 LC ‘Periq’e’ tobacco, PVH03, PVH09, PVH19, PVH50, PVH51, R 610, R 630, R 7-1 1, R 7-12, RG 17, RG 81, RG H51, RG H 4, RG H 51,

RS 1410, Speight 168, Speight 172, Speight 179, Speight 210, Speight 220, Speight 225, Speight 227, Speight 234, Speight G-28, Speight G-70, Speight H-6, Speight H20, Speight NF3, TI 1406, TI 1269, TN 86, TN86LC, TN 90, TN 97, TN97LC, TN D94, TN D950, TR (Tom Rosson) Madole, VA 309, VA359, AA 37-1, B 13P, Xanthi (Mitchell-Mor), Bel-W3, 79-615, Samsun Holmes NN, KTRDC number 2 Hybrid 49, Burley 21, KY 8959, KY 9, MD 609, PG 01, PG 04, P01, P02, P03, RG 1 1, RG 8, VA 509, AS44, Banket AI, Basma Drama B84/31, Basma I Zichna ZP4/B, Basma Xanthi BX 2A, Batek, Besuki Jember, C104, Coker 347, Criollo Misionero, Delcrest, Djebel 81, DVH 405, Galpao Comum, HB04P, Hicks Broadleaf, Kabakulak Ellassona, Kutsage E1, LA BU 21, NC 2326, NC 297, PVH 21 10, Red Russian, Samsun, Saplak, Simmaba, Talgar 28, Wislica, Yayaldag, Prilep HC-72, Prilep P23, Prilep PB 156/1, Prilep P12-2/1, Yaka JK-48, Yaka JB 125/3, TI-1068, KDH-960, TI-1070, TW136, Basma, TKF 4028, L8, TKF 2002, GR141, Basma xanthi, GR149, GR153, Petit Havana. Low converter subvarieties of the above, even if not specifically identified herein, are also contemplated.

[0296] The tobacco plant may be a Burley, Flue-cured Virginia, or Oriental.

[0297] In one embodiment the plant propagation material may be obtainable from a plant (e.g. a tobacco plant) of the invention.

[0298] A “plant propagation material” as used herein refers to any plant matter taken from a plant from which further plants may be produced. Suitably, a plant propagation material may be selected from a seed, plant calli and plant clumps. Suitably the plant propagation material may be a seed. Suitably, the plant propagation material may be plant calli. Suitably the plant propagation material may be plant clumps.

[0299] In one embodiment the cell (e.g. tobacco cell), cell culture, tobacco plant and/or plant propagation material may be obtainable (e.g. obtained) by a method according to the invention.

[0300] Suitably a tobacco plant according to the present invention may have modulated (e.g. decreased) nicotine content when compared to an unmodified tobacco plant, wherein the tobacco plant has been modified to modulate (e.g. decrease) the activity or expression of at least one Galactose-1-phosphate uridyl transferase. Suitably a tobacco plant according to the present invention may have decreased nicotine content when compared to an unmodified tobacco plant, wherein the tobacco plant has been modified to decrease or inhibit the activity or expression of at least one Galactose-1-phosphate uridyl transferase.

[0301] Suitably a tobacco plant according to the present invention may have modulated (e.g. reduced) content of a tobacco specific nitrosamine (TSNA) precursor when compared to an unmodified tobacco plant, wherein the tobacco plant has been modified to modulate (e.g. decrease) the activity or expression of at least one Galactose-1-phosphate uridyl transferase.

[0302] Suitably a tobacco plant according to the present invention may have decreased tobacco specific nitrosamine (TSNA) precursor content when compared to an unmodified tobacco plant, wherein the tobacco plant has been modified to decrease or inhibit the activity or expression of at least one Galactose-1-phosphate uridyl transferase.

[0303] In one embodiment the tobacco plant in accordance with the present invention comprises a tobacco cell of the invention.

[0304] In another embodiment the plant propagation material may be obtainable (e.g. obtained) from a tobacco plant of the invention.

[0305] In one embodiment there is provided the use of a tobacco plant as described herein to breed a tobacco plant.

[0306] The present invention also provides in another embodiment the use of a tobacco plant of the foregoing embodiments for the production of a tobacco industry product.

[0307] In another embodiment there is provided the use of a tobacco plant of the invention to grow a crop.

[0308] In one embodiment there is provided the use of a cell as provided for in the foregoing embodiments for production of a tobacco industry product.

[0309] In one embodiment the present invention provides a cell culture (e.g. in in vitro culture).

[0310] The tobacco cell culture may be a cell suspension culture. These cells cultured in vitro may be incorporated into a tobacco industry product, e.g. as a substitute for conventional tobacco particles, shreds, fine cut or long cut tobacco lamina, as an additive ingredient or as both a substitute and an additive. Suitably, the cell culture may produce nicotine.

[0311] In one embodiment there is provided the use of a cell culture, e.g. a harvested and/or processed cell culture according to the present invention for the production of a tobacco industry product.

[0312] The tobacco cells harvested from an in vitro culture may be dried, e.g. freeze-dried, for example to produce a powder.

[0313] In one embodiment, the cell culture is a tobacco cell culture. The skilled person will be aware of known methods for establishing in vitro cultures of tobacco cells. By way of example only, the following method may be used: collecting seeds from a tobacco plant of interest and sterilising their exterior to eliminate unwanted organisms, planting said seeds to grow a tobacco plant of interest, removing tissue from the tobacco plant (for example, from the tobacco stem) for use as an explant, establishing a callus culture from the tobacco explant, establishing a cell suspension culture from the callus culture, and harvesting culture material (e.g. including tobacco cells) to produce a tobacco cell culture.

[0314] The tobacco cells can be harvested by various methods, including filtration, e.g. vacuum filtration. The sample may be washed in the filter by adding water and the remaining liquid removed with the filtration, e.g. vacuum filtration.

[0315] The harvested tobacco cell culture may be further processed, e.g. dried, such as air-dried and/or freeze-dried. The harvested tobacco cell culture or dried harvested tobacco cell culture or an extract therefrom may be incorporated into tobacco industry products according to the present invention.

[0316] In one embodiment, the present invention provides a plant (e.g. tobacco plant) or part thereof for use in molecular farming. Suitably, a plant or part thereof modified in accordance with the present invention may be used in the manufacture of proteins such as therapeutics e.g. antibiotics, virus like particles, nutraceuticals or small molecules.

[0317] In one embodiment, the present invention provides a method for the production of proteins (e.g. therapeutic

proteins), the method comprising modifying a plant or part thereof capable of producing said protein (e.g. therapeutic protein) by modulating (e.g. decreasing) the activity or expression of at least one Galactose-1-phosphate uridyl transferase having an amino acid sequence as set out in SEQ ID No.3, or a functional variant or functional fragment or orthologue thereof, or a sequence which has at least 80% identity to SEQ ID No. 3; or wherein the at least one Galactose-1-phosphate uridyl transferase comprises a nucleotide sequence as set out in SEQ ID No. 1 or 2, or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2, or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2; and culturing the plant under conditions sufficient to allow the production of said protein (e.g. therapeutic protein).

Products

[0318] The present invention also provides for products obtainable or obtained from plants according to the present invention. Products are provided which are obtainable or obtained from a plant in which the activity or expression of Galactose-1-phosphate uridyl transferase has been modulated. In one embodiment, the product may comprise a construct of the invention which modulates the activity or expression of at least one Galactose-1-phosphate uridyl transferase as defined herein. In one embodiment, the product may comprise a construct of the invention which modifies the nucleic acid sequence of at least one Galactose-1-phosphate uridyl transferase as defined herein. The present invention also provides for products obtainable or obtained from tobacco according to the present invention.

[0319] In one embodiment there is provided the use of a tobacco plant of the invention to produce a tobacco leaf.

[0320] Suitably the tobacco leaf may be subjected to downstream applications such as processing.

[0321] Thus in one embodiment the use of the foregoing embodiment may provide a processed tobacco leaf. Suitably the tobacco leaf may be subjected to curing, fermenting, pasteurising or combinations thereof. In another embodiment the tobacco leaf may be cut. In some embodiments the tobacco leaf may be cut before or after being subjected to curing, fermenting, pasteurising or combinations thereof.

[0322] In one embodiment the present invention provides a harvested leaf of a tobacco plant of the invention.

[0323] In a further embodiment the harvested leaf may be obtainable (e.g. obtained) from a tobacco plant propagated from a propagation material of the present invention.

[0324] In another embodiment there is provided a harvest leaf obtainable from a method or use of the present invention.

[0325] Suitably the harvested leaf may be a cut harvested leaf.

[0326] In some embodiments the harvested leaf may comprise viable tobacco cells. In other embodiments the harvested leaf may be subjected to further processing.

[0327] There is also provided a processed tobacco leaf.

[0328] The processed tobacco leaf may be obtainable from a tobacco plant of the invention. Suitably the processed tobacco leaf may be obtainable from a tobacco plant obtained in accordance with any of the methods and/or uses of the present invention.

[0329] Suitably, the processed leaf may comprise reduced content of one or more TSNAs selected from NNN, NNK,

NAT and NAB. Suitably, the content of NNN may be reduced. Suitably, the content of NNK may be reduced. Suitably, the content of NAT may be reduced. Suitably, the content of NAB may be reduced. Suitably, the reduction in TSNA content is in relation to a comparable product which has not been modified according to the present invention.

[0330] In another embodiment the processed tobacco leaf may be obtainable (e.g. obtained) from a tobacco plant propagated from a tobacco plant propagation material according to the present invention.

[0331] The processed tobacco leaf of the present invention may be obtainable (e.g. obtained) by processing a harvested leaf of the invention.

[0332] The term “processed tobacco leaf” as used herein refers to a tobacco leaf that has undergone one or more processing steps to which tobacco is subjected to in the art. A “processed tobacco leaf” comprises no or substantially no viable cells.

[0333] The term “viable cells” refers to cells which are able to grow and/or are metabolically active. Thus, if a cell is said to not be viable, also referred to as “non-viable” then a cell does not display the characteristics of a viable cell.

[0334] The term “substantially no viable cells” means that less than about 5% of the total cells are viable. Preferably, less than about 3%, more preferably less than about 1%, even more preferably less than about 0.1% of the total cells are viable.

[0335] In one embodiment the processed tobacco leaf may be processed by one or more of: curing, fermenting and/or pasteurising.

[0336] Suitably the processed tobacco leaf may be processed by curing.

[0337] Tobacco leaf may be cured by any method known in the art. In one embodiment tobacco leaf may be cured by one or more of the curing methods selected from the group consisting of: air curing, fire curing, flue curing and sun curing.

[0338] Suitably the tobacco leaf may be air cured.

[0339] Typically air curing is achieved by hanging tobacco leaf in well-ventilated barns and allowing to dry. This is usually carried out over a period of four to eight weeks. Air curing is especially suitable for burley tobacco.

[0340] Suitably the tobacco leaf may be fire cured. Fire curing is typically achieved by hanging tobacco leaf in large barns where fires of hardwoods are kept on continuous or intermittent low smoulder and usually takes between three days and ten weeks, depending on the process and the tobacco. In another embodiment the tobacco leaf may be flue cured. Flue curing may comprise stringing tobacco leaves onto tobacco sticks and hanging them from tier-poles in curing barns. The barns usually have a flue which runs from externally fed fire boxes. Typically this results in tobacco that has been heat-cured without being exposed to smoke. Usually the temperature will be raised slowly over the course of the curing with the whole process taking approximately 1 week.

[0341] Suitably the tobacco leaf may be sun cured. This method typically involves exposure of uncovered tobacco to the sun.

[0342] Suitably the processed tobacco leaf may be processed by fermenting.

[0343] Fermentation can be carried out in any manner known in the art. Typically during fermentation, the tobacco leaves are piled into stacks (a bulk) of cured tobacco covered

in e.g. burlap to retain moisture. The combination of the remaining water inside the leaf and the weight of the tobacco generates a natural heat which ripens the tobacco. The temperature in the centre of the bulk is monitored daily. In some methods every week, the entire bulk is opened. The leaves are then removed to be shaken and moistened and the bulk is rotated so that the inside leaves go outside and the bottom leaves are placed on the top of the bulk. This ensures even fermentation throughout the bulk. The additional moisture on the leaves, plus the actual rotation of the leaves themselves, generates heat, releasing the tobacco’s natural ammonia and reducing nicotine, while also deepening the colour and improving the tobacco’s aroma. Typically the fermentation process continues for up to 6 months, depending on the variety of tobacco, stalk position on the leaf, thickness and intended use of leaf.

[0344] Suitably the processed tobacco leaf may be processed by pasteurising. Pasteurising may be particularly preferred when the tobacco leaf will be used to make a smokeless tobacco industry product, most preferably snus.

[0345] Tobacco leaf pasteurisation may be carried out by any method known in the art. For example pasteurisation may be carried out as detailed in J Foulds, L Ramstrom, M Burke, K Fagerstrom. Effect of smokeless tobacco (snus) on smoking and public health in Sweden Tobacco Control (2003) 12: 349-359, the teaching of which is incorporated herein by reference.

[0346] During the production of snus, pasteurisation is typically carried out by a process in which the tobacco is heat treated with steam for 24-36 hours (reaching temperatures of approximately 100° C.). This results in an almost sterile product and without wishing to be bound by theory one of the consequences of this is believed to be a limitation of further TSNA formation.

[0347] In one embodiment the pasteurisation may be steam pasteurisation.

[0348] In some embodiments the processed tobacco leaf may be cut. The processed tobacco leaf may be cut before or after processing. Suitably, the processed tobacco leaf may be cut after processing.

[0349] In one embodiment, the use of the foregoing embodiment may provide reconstituted tobacco.

[0350] In one embodiment, there is provided reconstituted tobacco.

[0351] “Reconstituted” as used herein may also be referred to as recon, recycled or homogenized sheet tobacco and refers to tobacco material generated from remnants of tobacco leaf after processing.

[0352] Reconstituted tobacco allows the production of a consistent, high quality blend and allows the adjustment of the ratio of individual components.

[0353] Reconstituted tobacco may be nano fibre recon (nanofibers can be extracted in solid or liquid form), paper making recon (which uses stems, scraps, and midribs, etc. as the raw material) or slurry type recon (which uses a mixture of fines and tobacco stems, ground to power, mixed with water and vegetable binding agent; the soluble residue is formed to sheets by extracting the water).

[0354] Any method known in the art may be used for making reconstituted tobacco, for example see CORESTA Congress, Sapporo, 2012, Smoke Science/Product Technology Groups, SSPT 12 (incorporated herein by reference).

[0355] In some embodiments the tobacco plant, harvested leaf of a tobacco plant and/or processed tobacco leaf may be

used to extract nicotine. The extraction of nicotine can be achieved using any method known in the art. For example a method for extracting nicotine from tobacco is taught in U.S. Pat. No. 2,162,738 which is incorporated herein by reference.

[0356] In one aspect, the present invention provides cured tobacco material made from a tobacco plant or part thereof according to the invention.

[0357] Suitably, the cured tobacco may comprise a reduced content of one or more TSNA's selected from NNK, NNN, NAT and NAB. Suitably, the content of NNN may be reduced. Suitably, the content of NNK may be reduced. Suitably, the content of NAT may be reduced. Suitably, the content of NAB may be reduced. Suitably, the reduction in TSNA content is in relation to a comparable product which has not been modified according to the present invention.

[0358] In another aspect, the present invention provides a tobacco blend comprising tobacco material made from a tobacco plant or part thereof according to the present invention, or from a tobacco cell culture according to the present invention. In one aspect, the present invention provides a tobacco blend comprising cured tobacco material according to the present invention.

[0359] Suitably, the tobacco blend according to the present invention may comprise approximately 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80% or 90% tobacco from a tobacco plant or part thereof according to the present invention, or from a tobacco cell culture according to the present invention. Suitably, the tobacco blend may comprise approximately 10% tobacco from a tobacco plant or part thereof according to the present invention, or from a tobacco cell culture according to the present invention. Suitably, the tobacco blend may comprise approximately 20% tobacco from a tobacco plant or part thereof according to the present invention, or from a tobacco cell culture according to the present invention. Suitably, the tobacco blend may comprise approximately 30% tobacco from a tobacco plant or part thereof according to the present invention, or from a tobacco cell culture according to the present invention. Suitably, the tobacco blend may comprise approximately 40% tobacco from a tobacco plant or part thereof according to the present invention, or from a tobacco cell culture according to the present invention. Suitably, the tobacco blend may comprise approximately 50% tobacco from a tobacco plant or part thereof according to the present invention, or from a tobacco cell culture according to the present invention. Suitably, the tobacco blend may comprise approximately 60% tobacco from a tobacco plant or part thereof according to the present invention, or from a tobacco cell culture according to the present invention. Suitably, the tobacco blend may comprise approximately 70% tobacco from a tobacco plant or part thereof according to the present invention, or from a tobacco cell culture according to the present invention. Suitably, the tobacco blend may comprise approximately 80% tobacco from a tobacco plant or part thereof according to the present invention, or from a tobacco cell culture according to the present invention. Suitably, the tobacco blend may comprise approximately 90% tobacco from a tobacco plant or part thereof according to the present invention, or from a tobacco cell culture according to the present invention.

[0360] In one aspect, a tobacco blend product of the present invention comprises at least about 5, 10, 20, 30, 40, 50, 60, 70, 80, 90, or 95 percent by dry weight of tobacco

cured from a tobacco plant or part thereof according to the present invention, or a tobacco cell culture according to the present invention.

[0361] Suitably, the cured tobacco material may be air cured. Suitably, the cured tobacco material may be flue cured. Suitably, the cured tobacco material may be sun cured. Suitably, the cured tobacco material may be fire cured.

[0362] A tobacco industry product or smoking article according to the present invention may comprise the tobacco material (e.g. cured tobacco material or reconstituted tobacco material) according to the present invention.

[0363] In another aspect the present invention provides a tobacco industry product.

[0364] In one embodiment the tobacco industry product according to the present invention may be a blended tobacco industry product. Suitably, the tobacco blend may comprise cured tobacco material according to the present invention.

[0365] In one embodiment the tobacco industry product may be prepared from a tobacco plant of the invention or a part thereof.

[0366] Suitably the tobacco plant or part thereof may be propagated from a tobacco plant propagation material according to the present invention.

[0367] The term "part thereof" as used herein in the context of a tobacco plant refers to a portion of the tobacco plant. Suitably, the "part thereof" may be a leaf, root or stem of a tobacco plant or the flowers. Suitably, the "part thereof" may be a leaf, root or stem of a tobacco plant.

Tobacco Industry Product

[0368] As used herein, the term "tobacco industry product" is intended to include combustible smoking articles such as cigarettes, cigarillos, cigars, tobacco for pipes or for roll-your-own cigarettes, (whether based on tobacco, tobacco derivatives, expanded tobacco, reconstituted tobacco, tobacco substitutes or other smokable material), non-combustible aerosol provision systems such as heating products that release compounds from substrate materials without burning such as electronic cigarettes, tobacco heating products, and hybrid systems to generate aerosol from a combination of substrate materials, for example hybrid systems containing a liquid or gel or solid substrate, as well as aerosolizable substrate materials used within these aerosol provision systems; and aerosol-free delivery articles such as lozenges, gums, patches, articles comprising breathable powders and smokeless tobacco industry products such as snus and snuff, which aerosol-free delivery articles may or may not deliver nicotine.

[0369] In one embodiment the tobacco industry product may be prepared from (e.g. may comprise) a tobacco plant of the invention or a part thereof.

[0370] Suitably the tobacco plant or part thereof may be propagated from a tobacco plant propagation material according to the present invention.

[0371] The term "part thereof" as used herein in the context of a tobacco plant refers to a portion of the tobacco plant. Preferably the "part thereof" is a leaf of a tobacco plant.

[0372] In another embodiment the tobacco industry product may be prepared from a harvested leaf of the invention.

[0373] In a further embodiment the tobacco industry product may be prepared from a processed tobacco leaf of the invention.

[0374] Suitably the tobacco industry product may be prepared from a tobacco leaf processed by one or more of: curing, fermenting and/or pasteurising.

[0375] Suitably the tobacco industry product may comprise a cut tobacco leaf, optionally processed as per the foregoing embodiment.

[0376] In another embodiment, the tobacco industry product may be prepared from a tobacco cell culture according to the present invention.

[0377] In another embodiment, the tobacco industry product may be prepared from (e.g. may comprise) a cured tobacco material according to the present invention.

[0378] In another embodiment, the tobacco industry product may be prepared from (e.g. may comprise) a tobacco blend according to the present invention.

[0379] In one embodiment the tobacco industry product may be a smoking article.

[0380] As used herein, the term “smoking article” can include smokeable products, such as rolling tobacco, cigarettes, cigars and cigarillos whether based on tobacco, tobacco derivatives, expanded tobacco, reconstituted tobacco or tobacco substitutes.

[0381] In another embodiment the tobacco industry product may be a smokeless tobacco industry product.

[0382] The term “smokeless tobacco industry product” as used herein refers to a tobacco industry product that is not intended to be smoked and/or subjected to combustion.

[0383] Smokeless tobacco industry products (including heat-not-burn materials) may contain tobacco in any form, including dried particles, shreds, granules, powders, or slurry, deposited on, mixed in, surrounded by, or combined with other ingredients in any format, such as flakes, films, tabs, foams, or beads.

[0384] In one embodiment a smokeless tobacco industry product may include snus, snuff, chewing tobacco or the like.

[0385] In one embodiment, the tobacco industry product is a combustible smoking article, selected from the group consisting of a cigarette, a cigarillo and a cigar.

[0386] In one embodiment, the tobacco industry product comprises one or more components of a combustible smoking article, such as a filter, a filter rod, a filter rod segments, tobacco, a tobacco rod, a tobacco rod segment, a spill, an additive release component such as a capsule, a thread, beads, a paper such as a plug wrap, a tipping paper or a cigarette paper.

[0387] In one embodiment, the tobacco industry product is a non-combustible aerosol provision system.

[0388] In one embodiment, the tobacco industry product comprises one or more components of a non-combustible aerosol provision system, such as a heater and an aerosolizable substrate.

[0389] In one embodiment, the aerosol provision system is an electronic cigarette also known as a vaping device.

[0390] In one embodiment the electronic cigarette comprises a heater, a power supply capable of supplying power to the heater, an aerosolizable substrate such as a liquid or gel, a housing and optionally a mouthpiece.

[0391] In one embodiment the aerosolizable substrate is contained in a substrate container. In one embodiment the substrate container is combined with or comprises the heater.

[0392] In one embodiment, the tobacco industry product is a heating product which releases one or more compounds by

heating, but not burning, a substrate material. The substrate material is an aerosolizable material which may be for example tobacco or other non-tobacco products, which may or may not contain nicotine. In one embodiment, the heating product is a tobacco heating product.

[0393] In one embodiment, the heating product is an electronic device.

[0394] In one embodiment, the tobacco heating product comprises a heater, a power supply capable of supplying power to the heater, an aerosolizable substrate such as a solid or gel material.

[0395] In one embodiment the heating product is a non-electronic article.

[0396] In one embodiment the heating product comprises an aerosolizable substrate such as a solid or gel material and a heat source which is capable of supplying heat energy to the aerosolizable substrate without any electronic means, such as by burning a combustion material, such as charcoal.

[0397] In one embodiment the heating product also comprises a filter capable of filtering the aerosol generated by heating the aerosolizable substrate.

[0398] In some embodiments the aerosolizable substrate material may comprise a vapour or aerosol generating agent or a humectant, such as glycerol, propylene glycol, triacetin or diethylene glycol.

[0399] In one embodiment, the tobacco industry product is a hybrid system to generate aerosol by heating, but not burning, a combination of substrate materials. The substrate materials may comprise for example solid, liquid or gel which may or may not contain nicotine. In one embodiment, the hybrid system comprises a liquid or gel substrate and a solid substrate. The solid substrate may be for example tobacco or other non-tobacco products, which may or may not contain nicotine. In one embodiment, the hybrid system comprises a liquid or gel substrate and tobacco.

[0400] In a further embodiment the tobacco industry product may be a tobacco heating device or hybrid device or e-cigarette or the like.

[0401] Typically in tobacco heating devices or hybrid devices, an aerosol is generated by the transfer of heat from a heat source to a physically separate aerosol-forming substrate or material, which may be located within, around or downstream of the heat source. During smoking, volatile compounds are released from the aerosol-forming substrate by heat transfer from the heat source and entrained in air drawn through the smoking article. As the released compounds cool, they condense to form an aerosol that is inhaled by the user.

[0402] Aerosol-generating articles and devices for consuming or smoking tobacco heating devices are known in the art. They can include, for example, electrically heated aerosol-generating devices in which an aerosol is generated by the transfer of heat from one or more electrical heating elements of the aerosol-generating device to the aerosol-forming substrate of a tobacco heating device.

[0403] Suitably the tobacco heating device may be an aerosol-generating device.

[0404] Preferably the tobacco heating device may be a heat-not-burn device. Heat-not-burn devices are known in the art and release compounds by heating, but not burning, tobacco.

[0405] An example of a suitable, heat-not-burn device may be one taught in WO2013/034459 or GB2515502 which are incorporated herein by reference.

[0406] In one embodiment the aerosol-forming substrate of a tobacco heating device may be a tobacco industry product in accordance with the present invention.

[0407] In one embodiment the tobacco heating device may be a hybrid device.

Polynucleotides/Polypeptides/Constructs

[0408] In certain embodiments of the present invention, constructs which modulate activity or expression of at least one Galactose-1-phosphate uridyl transferase as described herein may be transformed into plant cells, suitably under the direction of a promoter.

[0409] In certain embodiments of the present invention, constructs which decrease (i.e. inhibit) the activity or expression of at least one Galactose-1-phosphate uridyl transferase as described herein may be transformed into plant cells under the direction of a promoter. For example, the genetic construct may be a gene editing construct or may comprise an RNAi molecule, which may comprise a small interfering RNA (siRNA) molecule, or a short hairpin loop (shRNA) molecule.

[0410] In certain embodiments of the present invention, constructs which increase activity or expression of a gene encoding a Galactose-1-phosphate uridyl transferase as described herein may be transformed into plant cells, suitably under the direction of a promoter e.g. constructs which encode a gene encoding a Galactose-1-phosphate uridyl transferase such as an endogenous Galactose-1-phosphate uridyl transferase.

[0411] Constructs may be introduced into plants according to the present invention by means of suitable vector, e.g. plant transformation vectors. A plant transformation vector may comprise an expression cassette comprising 5'-3' in the direction of transcription, a promoter sequence, a construct sequence targeting a gene encoding a Galactose-1-phosphate uridyl transferase as described herein and, optionally a 3' untranslated, terminator sequence including a stop signal for RNA polymerase and a polyadenylation signal for polyadenylase. The promoter sequence may be present in one or more copies, and such copies may be identical or variants of a promoter sequence as described above. The terminator sequence may be obtained from plant, bacterial or viral genes. Suitable terminator sequences are the pea *rbcS* E9 terminator sequence, the *nos* terminator sequence derived from the nopaline synthase gene of *Agrobacterium tumefaciens* and the 35S terminator sequence from cauliflower mosaic virus, for example. A person skilled in the art will be readily aware of other suitable terminator sequences.

[0412] The construct of the present invention may also comprise a gene expression enhancing mechanism to increase the strength of the promoter. An example of such an enhancer element is one derived from a portion of the promoter of the pea plastocyanin gene, and which is the subject of International Patent Application No. WO 97/20056 which is incorporated herein by reference. Suitable enhancer elements may be the *nos* enhancer element derived from the nopaline synthase gene of *Agrobacterium tumefaciens* and the 35S enhancer element from cauliflower mosaic virus, for example.

[0413] These regulatory regions may be derived from the same gene as the promoter DNA sequence or may be derived from different genes, from *Nicotiana tabacum* or other organisms, for example from a plant of the family Solan-

aceae, or from the subfamily Cestroideae. All of the regulatory regions should be capable of operating in cells of the tissue to be transformed.

[0414] The promoter DNA sequence may be derived from the same gene as the gene of interest, e.g. the gene the promoter is going to direct, for instance a gene encoding a Galactose-1-phosphate uridyl transferase according to the invention, a coding sequence used in the present invention or may be derived from a different gene, from *Nicotiana tabacum*, or another organism, for example from a plant of the family Solanaceae, or from the subfamily Cestroideae.

[0415] The expression cassette may be incorporated into a basic plant transformation vector, such as pBIN 19 Plus, pBI 101, pKYLX71:35S2, pCambia2300 or other suitable plant transformation vectors known in the art. In addition to the expression cassette, the plant transformation vector will contain such sequences as are necessary for the transformation process. These may include the *Agrobacterium* vir genes, one or more T-DNA border sequences, and a selectable marker or other means of identifying transgenic plant cells.

[0416] The term "expression vector or plant transformation vector" means a construct capable of in vivo or in vitro expression. Preferably, the expression vector is incorporated in the genome of the organism. In one embodiment the vector of the present invention expresses a protein e.g. Galactose-1-phosphate uridyl transferase as described herein. The term "incorporated" preferably relates to stable incorporation into the genome.

[0417] Techniques for transforming plants are well known within the art and include *Agrobacterium*-mediated transformation, for example. The basic principle in the construction of genetically modified plants is to insert genetic information in the plant genome so as to obtain a stable maintenance of the inserted genetic material. A review of the general techniques may be found in articles by Potrykus (*Annu Rev Plant Physiol Plant Mol Biol* [1991]42:205-225) and Christon (*AgroFood-Industry Hi-Tech* March/April 1994 17-27), which are incorporated herein by reference. Typically, in *Agrobacterium*-mediated transformation a binary vector carrying a foreign DNA of interest, i.e. a construct according to the present invention, is transferred from an appropriate *Agrobacterium* strain to a target plant by the co-cultivation of the *Agrobacterium* with explants from the target plant. Transformed plant tissue is then regenerated on selection media, which selection media comprises a selectable marker and plant growth hormones. An alternative is the floral dip method (Clough & Bent, 1998 *Plant J.* 1998 December; 16(6):735-43, which is incorporated herein by reference) whereby floral buds of an intact plant are brought into contact with a suspension of the *Agrobacterium* strain containing the chimeric gene, and following seed set, transformed individuals are germinated and identified by growth on selective media. Direct infection of plant tissues by *Agrobacterium* is a simple technique which has been widely employed and which is described in Butcher et al. (1980) *Tissue Culture Methods for Plant Pathologists*, eds.: D. S. Ingrams and J. P. Helgeson, 203-208 which is incorporated herein by reference.

[0418] Further suitable transformation methods include direct gene transfer into protoplasts using polyethylene glycol or electroporation techniques, particle bombardment, micro-injection and the use of silicon carbide fibres for example. Transforming plants using ballistic transformation

and production of fertile transgenic maize plants by silicon carbide whisker-mediated transformation is taught in Frame et al. (1994) *The Plant Journal* 6(6): 941-948, which is incorporated herein by reference, and viral transformation techniques is taught in, for example, Meyer et al. (1992) *Mol. Gen. Genet.* 231(3): 345-352, which is incorporated herein by reference. The use of cassava mosaic virus as a vector system for plants is taught in Meyer et al. (1992) *Gene* 110: 213-217, which is incorporated herein by reference. Further teachings on plant transformation may be found in EP-A-0449375, incorporated herein by reference.

[0419] In a further aspect, the present invention relates to a vector system which carries a construct and introducing it into the genome of an organism, such as a plant, suitably a tobacco plant. The vector system may comprise one vector, but it may comprise two vectors. In the case of two vectors, the vector system is normally referred to as a binary vector system. Binary vector systems are described in further detail in Gynheung et al. (1980) *Binary Vectors*, Plant Molecular Biology Manual A3, 1-19, which is incorporated herein by reference.

[0420] One extensively employed system for transformation of plant cells uses the Ti plasmid from *Agrobacterium tumefaciens* or a Ri plasmid from *Agrobacterium rhizogenes* described by An et al. (1986) *Plant Physiol.* 81, 301-305 and Butcher et al. (1980) *Tissue Culture Methods for Plant Pathologists* eds.: D. S. Ingrams and J. P. Helgeson, 203-208 which are incorporated herein by reference. After each introduction method of the desired exogenous gene according to the present invention in the plants, the presence and/or insertion of further DNA sequences may be necessary. The use of T-DNA for the transformation of plant cells has been intensively studied and is described in EP-A-120516; Hoekema (1985) *The Binary Plant Vector System*, Offset-drukkerij Kanters B. B., Amsterdam Chapter V; Fraley et al. *Crit. Rev. Plant Sci.* 4:1-46; and An et al. (1985) *EMBO J* 4: 277-284, all incorporated herein by reference.

[0421] Plant cells transformed with construct(s) which modulate the activity or expression of at least one Galactose-1-phosphate uridyl transferase may be grown and maintained in accordance with well-known tissue culturing methods such as by culturing the cells in a suitable culture medium supplied with the necessary growth factors such as amino acids, plant hormones, vitamins, etc. The term "transgenic plant" in relation to the present invention includes any plant that comprises a construct which modulates the activity or expression of at least one Galactose-1-phosphate uridyl transferase according to the invention. Accordingly a transgenic plant is a plant which has been transformed with a construct according to the invention. Preferably the transgenic plant exhibits modulated (e.g. reduced) alkaloid content and/or modulated (e.g. reduced) TSNA precursor content according to the present invention. The term "transgenic plant" does not cover native nucleotide coding sequences in their natural environment when they are under the control of their native promoter which is also in its natural environment.

[0422] In one aspect, a gene encoding a Galactose-1-phosphate uridyl transferase, a construct, a plant transformation vector or a plant cell according to the present invention is in an isolated form. The term "isolated" means that the sequence is at least substantially free from at least one other component with which the sequence is naturally associated in nature and as found in nature.

[0423] In one aspect, a gene encoding a Galactose-1-phosphate uridyl transferase, a construct, plant transformation vector or a plant cell according to the invention is in a purified form. The term "purified" means in a relatively pure state, e.g. at least about 90% pure, or at least about 95% pure or at least about 98% pure.

[0424] The term "nucleotide sequence" as used herein refers to an oligonucleotide sequence or polynucleotide sequence, and variant, homologues, fragments and derivatives thereof (such as portions thereof). The nucleotide sequence may be of genomic or synthetic or recombinant origin, which may be double-stranded or single-stranded whether representing the sense or anti-sense strand.

[0425] The term "nucleotide sequence" in relation to the present invention includes genomic DNA, cDNA, synthetic DNA, and RNA. Preferably it means DNA, more preferably cDNA sequence coding for the present invention.

[0426] In a preferred embodiment, the nucleotide sequence when relating to and when encompassed by the present invention, i.e. the gene encoding a Galactose-1-phosphate uridyl transferase according to the present invention, includes the native nucleotide sequence when in its natural environment and when it is linked to its naturally associated sequence(s) that is/are also in its/their natural environment. For ease of reference, we shall call this preferred embodiment the "native nucleotide sequence". In this regard, the term "native nucleotide sequence" means an entire nucleotide sequence that is in its native environment and when operatively linked to an entire promoter with which it is naturally associated, which promoter is also in its native environment.

[0427] The nucleotide sequence for use in the present invention may be present in a vector in which the nucleotide sequence is operably linked to regulatory sequences capable of providing for the expression of the nucleotide sequence by a suitable host organism. The constructs for use in the present invention may be transformed into a suitable host cell as described herein to provide for expression of a polypeptide of the present invention. The choice of vector e.g. a plasmid, cosmid, or phage vector will often depend on the host cell into which it is to be introduced. Vectors may be used in vitro, for example for the production of RNA or used to transfect, transform, transduce or infect a host cell.

[0428] In some applications, the nucleotide sequence for use in the present invention is operably linked to a regulatory sequence which is capable of providing for the expression of the nucleotide sequence, such as by the chosen host cell. By way of example, the present invention covers a vector comprising the nucleotide sequence of gene encoding a Galactose-1-phosphate uridyl transferase as described herein operably linked to such a regulatory sequence, i.e. the vector is an expression vector.

[0429] The term "operably linked" refers to a juxtaposition wherein the components described are in a relationship permitting them to function in their intended manner. A regulatory sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences.

[0430] The term "regulatory sequences" includes promoters and enhancers and other expression regulation signals. The term "promoter" is used in the normal sense of the art, e.g. an RNA polymerase binding site. The nucleotide

sequence within a construct which encodes Galactose-1-phosphate uridyl transferase may be operably linked to at least a promoter.

[0431] The term “construct”—which is synonymous with terms such as “cassette” or “vector”—includes a nucleotide sequence for use according to the present invention directly or indirectly attached to a promoter.

[0432] An example of an indirect attachment is the provision of a suitable spacer group such as an intron sequence, such as the Sh1-intron or the ADH intron, intermediate the promoter and the nucleotide sequence of the present invention. The same is true for the term “fused” in relation to the present invention which includes direct or indirect attachment. In some cases, the terms do not cover the natural combination of the nucleotide sequence coding for the protein ordinarily associated with the wild type gene promoter and when they are both in their natural environment. The construct may even contain or express a marker, which allows for the selection of the genetic construct.

[0433] In some embodiments, a promoter may be operably linked to nucleotide sequence in a construct or vector which is used to modulate the concentration and/or total content of nicotine in a cell or cell culture or tobacco plant or part thereof.

[0434] In some embodiments the promoter may be selected from the group consisting of: a constitutive promoter, a tissue-specific promoter, a developmentally-regulated promoter and an inducible promoter.

[0435] In one embodiment the promoter may be a constitutive promoter.

[0436] A constitutive promoter directs the expression of a gene throughout the various parts of a plant continuously during plant development, although the gene may not be expressed at the same level in all cell types. Examples of known constitutive promoters include those associated with the cauliflower mosaic virus 35S transcript (Odell J T, Nagy F, Chua N H. (1985). Identification of DNA sequences required for activity of the cauliflower mosaic virus 35S promoter. *Nature*. 313 810-2), the rice actin 1 gene (Zhang W, McElroy D, Wu R. (1991). Analysis of rice Act1 5' region activity in transgenic rice plants. *Plant Cell* 3 1155-65) and the maize ubiquitin 1 gene (Cormejo M J, Luth D, Blankenship K M, Anderson O D, Blechl A E. (1993). Activity of a maize ubiquitin promoter in transgenic rice. *Plant Molec. Biol.* 23 567-81). Constitutive promoters such as the Carnation Etched Ring Virus (CERV) promoter (Hull R, Sadler J, Longstaff M (1986) (CaMV/35S), figwort mosaic virus 35S promoter. The sequence of carnation etched ring virus DNA: comparison with cauliflower mosaic virus and retroviruses. *EMBO Journal*, 5(2):3083-3090).

[0437] The constitutive promoter may be selected from a: a carnation etched ring virus (CERV) promoter, a cauliflower mosaic virus (CaMV 35S promoter), a promoter from the rice actin 1 gene or the maize ubiquitin 1 gene.

[0438] The promoter may be a tissue specific promoter. A tissue-specific promoter is one which directs the expression of a gene in one (or a few) parts of a plant, usually throughout the lifetime of those plant parts. The category of tissue-specific promoter commonly also includes promoters whose specificity is not absolute, i.e. they may also direct expression at a lower level in tissues other than the preferred tissue. Tissue specific promoters include the phaseolin-promoter, legumin b4-promoter, usp-promoter, sbp-promoter, ST-LS1 promoter, B33 (patatin class I promoter).

[0439] In another embodiment the promoter may be a developmentally-regulated promoter.

[0440] A developmentally-regulated promoter directs a change in the expression of a gene in one or more parts of a plant at a specific time during plant development. The gene may be expressed in that plant part at other times at a different (usually lower) level, and may also be expressed in other plant parts.

[0441] In one embodiment the promoter may be an inducible promoter.

[0442] An inducible promoter is capable of directing the expression of a gene in response to an inducer. In the absence of the inducer the gene will not be expressed. The inducer may act directly upon the promoter sequence, or may act by counteracting the effect of a repressor molecule. The inducer may be a chemical agent such as a metabolite, a protein, a growth regulator (such as auxin and salicylic acid which activate the OCS promoter), or a toxic element, a physiological stress such as heat, light (such as the soybean SSU promoter), wounding (e.g. the nos, nopaline synthase promoter), or osmotic pressure, or an indirect consequence of the action of a pathogen or pest. A developmentally-regulated promoter might be described as a specific type of inducible promoter responding to an endogenous inducer produced by the plant or to an environmental stimulus at a particular point in the life cycle of the plant. Examples of known inducible promoters include those associated with wound response, such as described by Warner S A, Scott R, Draper J. ((1993) *Plant J.* 3 191-201), temperature response as disclosed by Benfey & Chua (1989) (Benfey, P. N., and Chua, N-H. ((1989) *Science* 244 174-181), and chemically induced, as described by Gatz ((1995) *Methods in Cell Biol.* 50 411-424).

[0443] A nucleotide sequence encoding either a protein which has the specific properties of a gene encoding a Galactose-1-phosphate uridyl transferase as defined herein or a protein which is suitable for modification may be identified and/or isolated and/or purified from any cell or organism producing said protein. Various methods are well known within the art for the identification and/or isolation and/or purification of nucleotide sequences. By way of example, PCR amplification techniques to prepare more of a sequence may be used once a suitable sequence has been identified and/or isolated and/or purified.

[0444] In a yet further alternative, the nucleotide sequence encoding the Galactose-1-phosphate uridyl transferase may be prepared synthetically by established standard methods, e.g. the phosphoroamidite method described by Beaucage et al. (1981) *Tetrahedron Letters* 22, 1859-1869 which is incorporated herein by reference, or the method described by Matthes et al. (1984) *EMBO J.* 3, 801-805 which is incorporated herein by reference. In the phosphoroamidite method, oligonucleotides are synthesised, e.g. in an automatic DNA synthesiser, purified, annealed, ligated and cloned in appropriate vectors.

[0445] As used herein, the term “amino acid sequence” is synonymous with the term “polypeptide” and/or the term “protein”. In some instances, the term “amino acid sequence” is synonymous with the term “peptide”. In some instances, the term “amino acid sequence” is synonymous with the term “enzyme”.

[0446] The present invention also encompasses the use of sequences having a degree of sequence identity or sequence homology with amino acid sequence(s) of a polypeptide

having the specific properties defined herein or of any nucleotide sequence i.e. a gene encoding a Galactose-1-phosphate uridyl transferase (hereinafter referred to as a “homologous sequence(s)”). Here, the term “homologue” means an entity having a certain homology with the subject amino acid sequences and the subject nucleotide sequences. Here, the term “homology” can be equated with “identity”.

[0447] The homologous amino acid sequence and/or nucleotide sequence and/or fragments should provide and/or encode a polypeptide which retains the functional activity and/or enhances the activity of the Galactose-1-phosphate uridyl transferase. Typically, the homologous sequences will comprise the same catalytic sites etc. as the subject amino acid sequence for instance or will encode the same catalytic sites. Although homology can also be considered in terms of similarity (i.e. amino acid residues having similar chemical properties/functions), in the context of the present invention it is preferred to express homology in terms of sequence identity. Homologous sequences typically retain functional domains or motifs.

[0448] In one embodiment, a homologous sequence is taken to include an amino acid sequence or nucleotide sequence which has one, two or several additions, deletions and/or substitutions compared with the subject sequence.

[0449] In one embodiment, homologues of Galactose-1-phosphate uridyl transferase SEQ ID No. 3 are provided in Table 1. The present invention extends to the utilisation of homologues listed in Table 1 and to sequences having at least 80% sequence identity thereto.

Sequence Identity

[0450] Sequence identity comparisons can be conducted by eye, or more usually, with the aid of readily available sequence comparison programs. These commercially available computer programs can calculate % homology between two or more sequences. % homology or % identity may be calculated over contiguous sequences, i.e. one sequence is aligned with the other sequence and each amino acid in one sequence is directly compared with the corresponding amino acid in the other sequence, one residue at a time. This is called an “ungapped” alignment. Typically, such ungapped alignments are performed only over a relatively short number of residues.

[0451] Although this is a very simple and consistent method, it fails to take into consideration that, for example, in an otherwise identical pair of sequences, one insertion or deletion will cause the following amino acid residues to be put out of alignment, thus potentially resulting in a large reduction in % homology when a global alignment is performed. Consequently, most sequence comparison methods are designed to produce optimal alignments that take into consideration possible insertions and deletions without penalising unduly the overall homology score. This is achieved by inserting “gaps” in the sequence alignment to try to maximise local homology.

[0452] However, these more complex methods assign “gap penalties” to each gap that occurs in the alignment so that, for the same number of identical amino acids, a sequence alignment with as few gaps as possible—reflecting higher relatedness between the two compared sequences—will achieve a higher score than one with many gaps. “Affine gap costs” are typically used that charge a relatively high cost for the existence of a gap and a smaller penalty for each subsequent residue in the gap. This is the most commonly

used gap scoring system. High gap penalties will of course produce optimised alignments with fewer gaps. Most alignment programs allow the gap penalties to be modified. However, it is preferred to use the default values when using such software for sequence comparisons.

[0453] Calculation of maximum % homology therefore firstly requires the production of an optimal alignment, taking into consideration gap penalties. A suitable computer program for carrying out such an alignment is the Vector NTI (Invitrogen Corp.). Examples of software that can perform sequence comparisons include, but are not limited to, the BLAST package (see Ausubel et al. (1999) Short Protocols in Molecular Biology, 4th Ed—Chapter 18), BLAST 2 (see FEMS Microbiol Lett 1999 174(2): 247-50; FEMS Microbiol Lett 1999 177(1): 187-8 and tatiana(a)ncbi.nlm.nih.gov), FASTA (Altschul et al. 1990 J. Mol. Biol. 403-410) and AlignX for example. At least BLAST, BLAST 2 and FASTA are available for offline and online searching (see Ausubel et al. 1999, pages 7-58 to 7-60).

[0454] Although the final % homology can be measured in terms of identity, the alignment process itself is typically not based on an all-or-nothing pair comparison. Instead, a scaled similarity score matrix is generally used that assigns scores to each pairwise comparison based on chemical similarity or evolutionary distance. An example of such a matrix commonly used is the BLOSUM62 matrix—the default matrix for the BLAST suite of programs. Vector NTI programs generally use either the public default values or a custom symbol comparison table if supplied (see user manual for further details). For some applications, it is preferred to use the default values for the Vector NTI package.

[0455] Alternatively, percentage homologies may be calculated using the multiple alignment feature in Vector NTI (Invitrogen Corp.), based on an algorithm, analogous to CLUSTAL (Higgins DG & Sharp PM (1988), Gene 73(1), 237-244). Once the software has produced an optimal alignment, it is possible to calculate % homology, preferably % sequence identity. The software typically does this as part of the sequence comparison and generates a numerical result.

[0456] Should gap penalties be used when determining sequence identity, then preferably the following parameters are used for pairwise alignment:

FOR BLAST			
GAP OPEN			0
GAP EXTENSION			0

FOR CLUSTAL		DNA	PROTEIN
WORD SIZE	2		1
GAP PENALTY	15		10
GAP EXTENSION	6.66		0.1

[0457] In one embodiment, CLUSTAL may be used with the gap penalty and gap extension set as defined above. In some embodiments the gap penalties used for BLAST or CLUSTAL alignment may be different to those detailed above. The skilled person will appreciate that the standard parameters for performing BLAST and CLUSTAL alignments may change periodically and will be able to select

appropriate parameters based on the standard parameters detailed for BLAST or CLUSTAL alignment algorithms at the time.

[0458] Suitably, the degree of identity with regard to a nucleotide sequence is determined over at least 50 contiguous nucleotides, preferably over at least 60 contiguous nucleotides, preferably over at least 70 contiguous nucleotides, preferably over at least 80 contiguous nucleotides, preferably over at least 90 contiguous nucleotides, preferably over at least 100 contiguous nucleotides, preferably over at least 150 contiguous nucleotides, preferably over at least 200 contiguous nucleotides, preferably over at least 250 contiguous nucleotides, preferably over at least 300 contiguous nucleotides, preferably over at least 350 contiguous nucleotides, preferably over at least 400 contiguous nucleotides, preferably over at least 450 contiguous nucleotides, preferably over at least 500 contiguous nucleotides, preferably over at least 550 contiguous nucleotides, preferably over at least 600 contiguous nucleotides, preferably over at least 650 contiguous nucleotides, or preferably over at least 700 contiguous nucleotides.

[0459] Suitably, the degree of identity with regard to a nucleotide, cDNA, cds or amino acid sequence may be determined over the whole sequence.

[0460] The sequences may also have deletions, insertions or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent substance. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues as long as the secondary binding activity of the substance is retained. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values include leucine, isoleucine, valine, glycine, alanine, asparagine, glutamine, serine, threonine, phenylalanine, and tyrosine.

[0461] Conservative substitutions may be made, for example according to the Table below. Amino acids in the same block in the second column and preferably in the same line in the third column may be substituted for each other:

ALIPHATIC	Non-polar	G A P I L V
	Polar - uncharged	C S T M N Q
	Polar - charged	D E K R
	AROMATIC	H F W Y

[0462] The present invention also encompasses homologous substitution (substitution and replacement are both used herein to mean the interchange of an existing amino acid residue, with an alternative residue) that may occur i.e. like-for-like substitution such as basic for basic, acidic for acidic, polar for polar etc. Non-homologous substitution may also occur i.e. from one class of residue to another or alternatively involving the inclusion of unnatural amino acids such as ornithine (hereinafter referred to as Z), diaminobutyric acid ornithine (hereinafter referred to as B), norleucine ornithine (hereinafter referred to as O), pyrilylalanine, thienylalanine, naphthylalanine and phenylglycine.

[0463] Replacements may also be made by unnatural amino acids include; alpha* and alpha-disubstituted* amino acids, N-alkyl amino acids*, lactic acid*, halide derivatives of natural amino acids such as trifluorotyrosine*, p-Cl-phenylalanine*, p-Br-phenylalanine*, p-1-phenylalanine*, L-allyl-glycine*, beta-alanine*, L-alpha-amino butyric acid*, L-gamma-amino butyric acid*, L-alpha-amino isobutyric acid*, L-e-amino caproic acid#, 7-amino heptanoic acid*, L-methionine sulfone#, L-norleucine*, L-norvaline*, p-nitro-L-phenylalanine*, L-hydroxyproline#, L-thiopropine*, methyl derivatives of phenylalanine (Phe) such as 4-methyl-Phe*, pentamethyl-Phe*, L-Phe (4-amino)#, L-Tyr (methyl)*, L-Phe (4-isopropyl)*, L-Tic (1,2,3,4-tetrahydroisoquinoline-3-carboxyl acid)*, L-diaminopropionic acid# and L-Phe (4-benzyl)*. The notation * has been utilised for the purpose of the discussion above (relating to homologous or non-homologous substitution), to indicate the hydrophobic nature of the derivative whereas # has been utilised to indicate the hydrophilic nature of the derivative, ## indicates amphipathic characteristics.

[0464] Variant amino acid sequences may include suitable spacer groups that may be inserted between any two amino acid residues of the sequence including alkyl groups such as methyl, ethyl or propyl groups in addition to amino acid spacers such as glycine or beta-alanine residues. A further form of variation, involves the presence of one or more amino acid residues in peptoid form, which will be well understood by those skilled in the art. For the avoidance of doubt, "the peptoid form" is used to refer to variant amino acid residues wherein the alpha-carbon substituent group is on the residue's nitrogen atom rather than the alpha-carbon. Processes for preparing peptides in the peptoid form are known in the art, for example Simon et al. (1992) PNAS 89(20), 9367-9371 and Horwell (1995) Trends Biotechnol. 13(4), 132-134.

[0465] The nucleotide sequences for use in the present invention may include within them synthetic or modified nucleotides. A number of different types of modification to oligonucleotides are known in the art. These include methylphosphonate and phosphorothioate backbones and/or the addition of acridine or polylysine chains at the 3' and/or 5' ends of the molecule. For the purposes of the present invention, it is to be understood that the nucleotide sequences described herein may be modified by any method available in the art. Such modifications may be carried out in order to enhance the in vivo activity or life span of nucleotide sequences of the present invention. The present invention also encompasses sequences that are complementary to the nucleic acid sequences of the present invention or sequences that are capable of hybridising either to the sequences of the present invention or to sequences that are complementary thereto. The term "hybridisation" as used herein shall include "the process by which a strand of nucleic acid joins with a complementary strand through base pairing" as well as the process of amplification as carried out in polymerase chain reaction (PCR) technologies.

[0466] The present invention also relates to nucleotide sequences that can hybridise to the nucleotide sequences of the present invention (including complementary sequences of those presented herein). Preferably, hybridisation is determined under stringency conditions (e.g. 50° C. and 0.2×SSC {1×SSC=0.15 M NaCl, 0.015 M Na₃citrate pH 7.0}). More preferably, hybridisation is determined under high stringency conditions (e.g. 65° C. and 0.1×SSC {1×SSC=0.15 M NaCl, 0.015 M Na₃citrate pH 7.0}).

[0467] A review of the general techniques used for transforming plants may be found in articles such as Potrykus et al. (1991) *Annu Rev Plant Physiol. Plant Mol. Biol.* 42:205-225 and Christou et al. (1994) *Agro-Food-Industry Hi-Tech March/April* 17-27, which are incorporated herein by reference. Further teachings on plant transformation may be found in EP-A-0449375, incorporated herein by reference.

[0468] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. Singleton, et al., *DICTIONARY OF MICROBIOLOGY AND MOLECULAR BIOLOGY*, 20 ED., John Wiley and Sons, New York (1994), and Hale & Marham, *THE HARPER COLLINS DICTIONARY OF BIOLOGY*, Harper Perennial, NY (1991) provide one of skill with a general dictionary of many of the terms used in this disclosure.

[0469] This disclosure is not limited by the exemplary methods and materials disclosed herein, and any methods and materials similar or equivalent to those described herein can be used in the practice or testing of embodiments of this disclosure. Numeric ranges are inclusive of the numbers defining the range. Unless otherwise indicated, any nucleic acid sequences are written left to right in 5' to 3' orientation; amino acid sequences are written left to right in amino to carboxy orientation, respectively.

[0470] The headings provided herein are not limitations of the various aspects or embodiments of this disclosure which can be had by reference to the specification as a whole. Accordingly, the terms defined immediately below are more fully defined by reference to the specification as a whole.

[0471] Amino acids are referred to herein using the name of the amino acid, the three letter abbreviation or the single letter abbreviation. The term "protein", as used herein, includes proteins, polypeptides, and peptides. As used herein, the term "amino acid sequence" is synonymous with the term "polypeptide" and/or the term "protein". In some instances, the term "amino acid sequence" is synonymous with the term "peptide". In some instances, the term "amino acid sequence" is synonymous with the term "enzyme".

[0472] In the present disclosure and claims, the conventional one-letter and three-letter codes for amino acid residues may be used. The 3-letter code for amino acids as defined in conformity with the IUPACIUB Joint Commission on Biochemical Nomenclature (JCBN). It is also understood that a polypeptide may be coded for by more than one nucleotide sequence due to the degeneracy of the genetic code.

[0473] Other definitions of terms may appear throughout the specification. Before the exemplary embodiments are described in more detail, it is to be understood that this disclosure is not limited to particular embodiments described, as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present disclosure will be limited only by the appended claims.

[0474] Where a range of values is provided, it is understood that each intervening value, to the tenth of the unit of the lower limit unless the context clearly dictates otherwise, between the upper and lower limits of that range is also specifically disclosed. Each smaller range between any stated value or intervening value in a stated range and any other stated or intervening value in that stated range is

encompassed within this disclosure. The upper and lower limits of these smaller ranges may independently be included or excluded in the range, and each range where either, neither or both limits are included in the smaller ranges is also encompassed within this disclosure, subject to any specifically excluded limit in the stated range. Where the stated range includes one or both of the limits, ranges excluding either or both of those included limits are also included in this disclosure.

[0475] It must be noted that as used herein and in the appended claims, the singular forms "a", "an", and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "an enzyme" or "Galactose-1-phosphate uridyl transferase" includes a plurality of such candidate agents and equivalents thereof known to those skilled in the art, and so forth.

Advantages

[0476] It has been surprisingly found that by modulating the activity or expression of Galactose-1-phosphate uridyl transferase as taught herein which acts as a positive regulator of alkaloid content in tobacco, the alkaloid and/or TSNA precursor content of plants can be modulated. Thereby tobacco industry products with modulated (e.g. reduced) alkaloid content) and/or reduced TSNA precursor content and commercially desirable traits sought after by consumers of tobacco industry products can be produced.

[0477] The present inventors have surprisingly determined a method for modulating the alkaloid content and/or TSNA precursor content of a plant (e.g. tobacco plant) by modulating the activity or expression of Galactose-1-phosphate uridyl transferase. Alkaloid or TSNA precursor content of a plant (e.g. tobacco plant) may be decreased by decreasing or inhibiting the activity or expression of Galactose-1-phosphate uridyl transferase as described herein. Prior to the present invention it had not been known that modulation of the activity or expression of Galactose-1-phosphate uridyl transferase as described herein could be used to modulate alkaloid (and/or TSNA precursor content of a plant (e.g. a tobacco plant).

[0478] The publications discussed herein are provided solely for their disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that such publications constitute prior art to the claims appended hereto.

EXAMPLES

Example 1—Virus-Induced Gene Silencing (VIGS) of Nitab4.5_0002825g0060.2 Decreases Alkaloid Content in Leaves

Virus-Induced Gene Silencing (VIGS)

[0479] For virus induced gene silencing, a 300-nucleotide cDNA fragment of Nitab4.5_0002825g0060.2:

(TRV2: SEQ ID No. 16)

TGTTGATATTCAACCAATGAATTGTTAATTGATCAATCAGCCCATTTCA

TATCATTGGTTCCCTTTGCTGCACTTTTGCTTTTGAGATTGGATAAT

TCCTCGTGATCACTCCTCTCATTTTCATGAACGATAATGAAAAGGCA

-continued

AATGATCTTGGAGGGTTGTTGAAACTCATGCTTTTGAAGATGTCTTTGC
 AGTTGAACAATCCACCATTCAACCTCCTTATTCATACCTCCCCATTTCA
 AGTAGATCCCTCATATGCTCCTTCCACACACTGGTTTTTACAAATAGCT
 CCCCAT

was synthesized and cloned with In-Fusion cloning kit into pTV00 (between EcoRI and XhoI sites). The plasmid was then transformed into *A. tumefaciens* GV3101.

[0480] The TRV vector comprising both (TRV RNA1 SEQ ID No. 15) and (TRV RNA2) comprising the targeted nucleotide sequence were separately propagated in *A. tumefaciens*. These cultures were mixed (1:1) and syringe-infiltration into 2-week-old TN90 plants. The silencing effect was assessed five weeks post-virus infection by assessing the expression level of the target gene.

Silencing

[0481] VIGS assays were performed as previously described (Ratcliff et al., 2001) Ratcliff, F et al., (2001), The Plant Journal, 25: 237-245 (incorporated herein by reference). Briefly, independent cultures of *A. tumefaciens* GV3101 carrying TRV2 and TRV1 plasmids were propagated overnight in LB medium supplemented with appropriate antibiotics. Cultures were resuspended in VIGS buffer (10 mM morpholineethanesulfonic acid pH 5.6, 10 mM MgCl₂, and 100 μM acetocyringone) adjusting optical density to OD₆₀₀=1, and incubated overnight at room temperature in the dark. These cultures were mixed (1:1) and syringe-infiltrated into 2-week-old TN90 plants. The silencing effect was assessed two weeks post-virus infection by assessing the expression level of the target gene. TRV-Luciferase was used as a negative control and TRV-PDS (reduced chlorophyll content of the silenced leaves) was used as a phenotypic silencing control.

Alkaloid Measurement

[0482] Relative content of pyridine alkaloids was determined by reversed phase high performance liquid chromatography with tandem mass spectrometry (LC-MS/MS). Chromatographic separation was achieved using a Gemini-NX column (100 mm×3.0 mm, particle size 3 µm, Phenomenex) and gradient chromatographic separation using 6.5 mM ammonium acetate buffer (aq) (pH10) and Methanol.

[0483] Mass Spectrometer operates in electrospray (ESI) positive mode using scheduled MRM data acquisition. Two MRM transitions were monitored for each analyte and one for the isotope labelled internal standard.

Analyte	Precursor Ion	Daughter Ion (quant/confirm)
Nicotine	163.1	130/106
Nicotine d4	167.1	134.1
Anabasine	163.1	80/120
Anatabine	161.1	144/80
Normicotine	149.1	80/130
Normicotine d4	153.1	84.1
PON	176.1	106.0/148
PON d4	183.1	110.0

Results

[0484] Alkaloid content of plants targeted with VIGS construct for Nitab4.5_0002825g0060.2. Graph is representative of 2 biological replicates (N=12) analysed by one-way ANOVA and Tukey's multiple-comparison post-test. Values are shown as means $\pm$ SEM. Asterisks indicate statistical significance of P value $\leq$ 0.05.

[0485] VIGS of Nitab4.5_0002825g0060.2 leads to a decrease in alkaloid content in leaves, in particular a decrease in nicotine, nor nicotine and PON content.

CONCLUSIONS

[0486] Nitab4.5_0002825g0060.2 is a positive regulator of alkaloid content, in particular alkaloid content in leaves and is a regulator of pyridine alkaloids in tobacco.


Example 2—Nitab4.5 0002825q0060.2 has
Galactose-1-Phosphate Uridyl Transferase
Functional Activity

[0487] The present inventors characterised Nita4.5_0002825g0060.2 as a gene encoding for Galactose-1-phosphate uridyl transferase. This enzyme facilitates the simultaneous conversion of uridine diphosphoglucose (UDP-glucose) and galactose-1-phosphate (gal-1P) to uridine diphosphogalactose (UDP-galactose) and glucose-1-phosphate.

[0488] To confirm the categorisation a functional assay was performed to determine the activity of Nitab4.5_0002825g0060.2.

[0489] The galactose-1-phosphate uridyl transferase coupled assay protocol described in Mayes, and Hanson supra was used to functionally validate and measure the activity of Nitab4.5_0002825g0060.2. This coupled assay results in the formation of B-NADPH which is monitored by an increase in absorbance at 340 nm.

Coupled Enzyme Reaction:



Results

[0490] FIG. 2 shows the measured endpoint absorbance values (340 nm) for the production of B-NADPH over a 60 minute reaction by 1.7 pg of Nitab4.5_0002825g0060.2 enzyme or 0.5 pg of positive control enzyme (recombinant human Galactose-1-phosphate uridyl transferase, MyBioSource, MBS143204) in the presence of 0.67 mM uridine 5'-diphosphoglucose and 1.3 mM galactose 1-Phosphate. The results for the control reactions are also shown on the same graph and indicate that the reaction is dependent on uridine 5'-diphosphoglucose and galactose 1-Phosphate.

MBP and the no enzyme control both have limited activity. The commercial positive control displays >5-fold higher activity per microgram of protein in this assay but the Nitab4.5_0002825g0060.2 enzyme does demonstrate reliable, substrate-dependant activity. Error bars are standard deviation from four replicates.

CONCLUSIONS

[0491] Nitab4.5_0002825g0060.2 is capable of functioning as a Galactose-1-phosphate uridyl transferase and as

shown in Example 1, is a positive regulator of alkaloid content, in particular alkaloid content in leaves and is a regulator of pyridine alkaloids in tobacco.

Example 3—Homologue Testing

[0492] The effects of the homologues of SEQ ID No. 3, namely those listed in Table 1, are tested in assays as described in the above examples.

SEQUENCE LISTING

```
Sequence total quantity: 16
SEQ ID NO: 1          moltype = DNA length = 3178
FEATURE              Location/Qualifiers
source                1..3178
                     mol_type = genomic DNA
                     organism = Nicotiana tabacum

SEQUENCE: 1
acatagatta  ttatctagaa  tatgggtact  cttgcactta  cacaaaaact  agaggagcaa  60
agttgatcaa  ttctctcaag  ttttatatat  tccacgaaac  catgttctta  acttccgaga  120
ttctcagtga  agcatcgaag  tgaagtggag  gagcaccaaa  ttcggagccc  cgacacaaag  180
gcaaatggcg  gaggcagaag  caccaaatcg  gagccccgaa  attcggaaaag  acagaatcca  240
caacaggtgg  gtcctattct  cgccggcgag  atcacggcgg  cegtctgact  tcaaggcaaa  300
atcgaaccca  caaccgaaca  ataaaacaga  atgcccattt  tgcgccggcc  acgagcacga  360
gtgtgcgcgg  gagatttttc  gtgtgcgcgg  aaattccacc  gccgattgga  aaatcagagt  420
tatccaaaat  ctctttccgg  agtcagcag  agaattggac  ttccaaaatc  cggttttctg  480
cctcggggac  gtatgtgta  gtgggttcgg  gtttcacgat  gtggttattg  aatcgccggg  540
tcattcagta  aatgatgagc  atctttctcc  gtcacaagtt  ggggaagtta  tgctggcttt  600
taagaagcgg  gtcgagcagc  tccagaatta  tgataatgtc  aaatatgttc  aggttttatt  660
ctgatcccg  ataaaaatgc  attggttttc  gtatcaattt  gtctatattt  tgtagttgat  720
actgtatatt  cattctttga  ttttatatga  atttaaaata  aatgcattta  aaagctacat  780
aaaaatagta  gagaccatga  ctattaaaag  ctaaaagtat  ttaacaaagt  atatgatgaa  840
ggcttgctgc  aatggtgaat  tttttgtttg  cagtaaaagg  ctttgaagat  ttacctggct  900
ttaatctttt  ggtgttaattg  ctatatggg  ttcatttcca  ttaacttgaa  agtgttttgt  960
ttttgtacaa  ataaaataa  agagttaaag  ttaattgtag  agcccatcaa  tgtttgtaca  1020
aagtgaacaa  ccaattatag  ctctttttat  gttatattta  gcaaacctat  tttgtcattt  1080
agttaatttt  atcttttagc  tctttttatt  tcagctgctt  atcatttgtc  tgcttttcat  1140
ttgtggcttt  ttaatttggc  agactgaatg  atttttctgt  gtaacattgg  tggagaaaaa  1200
gcttaggctg  acaacatggt  tgtctttagg  ttttaattga  acaaatataa  cagcatttga  1260
cgtttggta  tggaagagta  gctctttact  tagaccaact  tgctaaaata  ctgaacccaa  1320
ttatgagagg  cttttggtag  aaactcatag  actgtcaagt  gtcacctgct  tgagggtcta  1380
tcattaagga  gaataatttt  agaggaaagt  tttccattaa  tctgcatagg  agcttgtagt  1440
atgaactctc  catccattta  tctgacatgg  tttgattttc  atggtagatt  atttgagttt  1500
gtatttttga  tgacaggaga  aaatctggta  gcattttaa  gaagtgggtt  tgataggtta  1560
aatatcaact  tgactgtcaa  tttgaactat  ttaacagaac  tgctcttaaa  cttgtaagtc  1620
atgtagcagt  ttatccaaaa  aaaaaaaat  actgtaaatc  atgtagcaag  tagcatattt  1680
gtggtatcaa  gccaaacctt  gtttgatata  ccagattaca  aaccttttt  gagaggtaat  1740
gttagctgca  cattcttggt  acttggttcc  tgttccattg  cagaatactc  caaaaatatgc  1800
acttctatct  actcatacca  aatatatgta  agagaacttg  agaagtgatt  agttgtcatg  1860
aaattggaga  tgaatgagca  caatatctga  tctcatgccc  tttgtgcgtg  caaggaaatt  1920
gcagtacgga  acagcatgtg  ttgttttttc  cactttcttg  ttgagtataa  tgattttgtac  1980
tgtactgagc  aagattgtcc  catcaaaagg  tactggacta  tataacacta  attcattcaa  2040
tggcttataa  aggtgttcaa  aaacctgga  gccctgctg  gtgcttcaat  gagccactcc  2100
cacagccaga  tgattgctct  tccattggt  cctccaacag  tttctgctcg  tcttgacagt  2160
atgagggaa  actacaagca  gactgggaaa  tgcagcctgt  gtgatattca  accaaatgaa  2220
ttgttaattg  atcaatcagc  ccatttcata  tcatttggtt  cctttgctgc  aacttttgct  2280
tttgagattt  ggataattcc  tcgtgatcac  tcctctcatt  ttcatgaact  cgataatgaa  2340
aaggtaatcg  atgcttttct  tgtcaataag  attgttttca  tgagaaaatt  atgtgaagtt  2400
acacagtttt  aactttgcaa  atatcagtta  tctccaaatt  ctacgtgaga  aaagtccaaa  2460
cggtagacca  tgaggccctt  gacttatctc  aagtagaatt  tgctgattag  tgaagagaaa  2520
aggaaaactc  atcaaaaaac  tgatattcag  gaaaaaacgt  acgttgtttc  aggctaaact  2580
aattttttga  ttgtttctcc  atctcaacaa  ctacttagtt  gaaaagattt  gtgatgtttt  2640
accaatgaaa  tatgcaggca  aatgatcttg  gaggttggtt  gaaactcatg  cttttgaaga  2700
tgtctttgca  gttgaacaat  ccaccattca  acctccttat  tcataacctc  ccattttcaag  2760
tagatccctc  atatgctcct  tccacacact  ggtttttaca  aatagctccc  catttgagtg  2820
gagttggtgg  attcgagata  gcaaccggat  gtcacataaa  tcctgttttc  ccagaggatg  2880
ctgccccaa  tctgagggat  gtgagatacc  ttaattaaca  gtatttaggg  ggaatctatt  2940
aaatcctaatt  attcatatct  gtgacacag  taactaagga  gataaagagg  atgtggatag  3000
agggaagatg  tacccttttt  atgaggagtt  atgtgtagat  tcgattttgt  gatccattac  3060
aaatgttatc  ttatgttaca  cggtcatttt  cagcagaaaa  ataattctcc  ctccatttca  3120
atttacctat  tttcttttta  gtctatgcca  acaaaaatga  ccactttctt  tatttgga  3178
```

SEQ ID NO: 2 moltype = DNA length = 1464

-continued

FEATURE
source

Location/Qualifiers
1..1464
mol_type = genomic DNA
organism = *Nicotiana tabacum*

SEQUENCE: 2

acatagatta	ttatctagaa	tatgggtact	cttgcactta	cacaaaaact	agaggagcaa	60
agttgatcaa	tttcctcacg	ttttatatat	tccacgaaac	catgtttctta	acttccgaga	120
ttctcagtga	agcatcgaag	tgaagtggag	gagcaccaaa	ttcggagccc	cgacacaaag	180
gcaaatggcg	gaggcagaag	caccaaactcg	gagcccccga	attcggaag	acagaatcca	240
caacaggtgg	gtcctattct	cgccggcgag	atcacggcgg	ccgtctgact	tcaaggcaaa	300
atcgaaccca	caacogaaca	ataaaacaga	atgcccaatt	tgcgccggcc	acgagcacga	360
gtgtgcccgc	gagatttttc	gtgtgccggc	aaattccacc	gccgattgga	aaatcagagt	420
tatccaaaat	ctctttccgg	cagtcagcag	agaattggac	ttccaaaatc	cggttttctgc	480
cctcggggac	gtagctgtta	gtgggttcgg	gtttcacgat	gtggttattg	aatcgccggt	540
tcattcagta	aatatgagcg	atctttctcc	gtcacaaagt	ggggaagtta	tgctggcttt	600
taagaagcgg	gtcgcagcag	tccagaatta	tgataatgtc	aaatatgttc	aggtgttcaa	660
aaacctatga	gcctctgctg	gtgcttcaat	gagccactct	cacagccaga	tgattgctct	720
tcccattgtt	cctccaacag	ttctgctcgc	tcttgacagt	atgagggaa	actacaagca	780
gactgggaaa	tgcagcctgt	gtgatatcca	accaaataga	ttgttaattg	atcaatcagc	840
ccatttcata	tcatttggtc	cctttgctgc	aacttttgct	tttgagattt	ggataattcc	900
tcgtgatcac	tcctctcatt	ttcatgaact	cgataatgaa	aaggcaaatg	atcttggagg	960
gttgttga	ctcatgcttt	tgaagatgtc	tttgcagttg	aacaatccac	cattcaacct	1020
ccttattcat	acctccccat	ttcaagtaga	tcctccatat	gctccttcca	cacactgggt	1080
tttacaata	gctccccatt	tgagtggagt	tggtggattc	gagatagcaa	ccgcatgtca	1140
cataaatcct	gttttccag	tcgatgtcgc	caaaattctg	agggatgtga	gaatccttaa	1200
ttaacagtat	ttagggggaa	gctattaaat	ctaattatct	atatctgtag	acacagtaac	1260
taaggagata	aagaggatgt	ggatacaggg	aagatgtacc	ctttttatgc	ggagttagt	1320
gtagattcga	ttttgtgatc	cattacaaat	gttatcttat	gttacacggg	cattttcacg	1380
caaaaaataa	tttccctccc	atttcaattt	acctattttc	tttttagtct	atgccacaaa	1440
aatgaccac	tttctttatt	tgga				1464

SEQ ID NO: 3

FEATURE
source

moltype = AA length = 339
Location/Qualifiers
1..339
mol_type = protein
organism = *Nicotiana tabacum*

SEQUENCE: 3

MAEAEAPNRS	PEIRKDRHNN	RWVLFSPARS	RRPSDFKAKS	NPQPNKTEC	PFCAGHEHEC	60
APFIFRPVAN	STADWKIRVI	QNLFPVAVSRE	LDFQNPVSAL	GDVAVSGFGF	HDVVIESPVH	120
SVNMSDLSPS	QVGEVMLAFK	KRVEQLQNYD	NVKYVQVFKN	HGASAGASMS	HSQSQMIALP	180
IVPPTVSARL	DSMREYYQOT	GKCSLQDIQP	NELLIDQSAH	FISLVPPFAAT	FAFEIWIIPR	240
DHSSHFHELD	NEKANDLGGI	LKLMLLKMMSL	QLNNPPFNLL	IHTSPFQVDP	SYAPSTHWFL	300
QIAPHLSGVG	GFEIATGCHI	NPVFPEDAAS	ILRDVRILN			339

SEQ ID NO: 4

FEATURE
source

moltype = DNA length = 3057
Location/Qualifiers
1..3057
mol_type = genomic DNA
organism = unidentified

SEQUENCE: 4

acctcacgtt	tcatatatta	taccttttcc	acgaaacgct	gttcttaacc	tcccagattt	60
tcagtgaaac	atcgaagtga	agtggaggag	cacaaaattc	ggagcccccga	cacaaaggca	120
aatggcggag	gcagaagcac	caaatcggag	ccccgaaatc	cggaagagaca	gaatccacaa	180
caggtgggtc	ctatttctgc	cgccgagatc	acggcggccg	tctgacttca	aggcaaaatc	240
gaacccacaa	ccgaacaata	aaacagaatg	cccattttgc	gccggccacg	agcacgagtg	300
tgcccgccgag	atttttcgtg	tgccggcaga	ttccaccgct	gattggaaaa	tcagagttaa	360
ccaaaatctc	tttccggcag	tcagcagaga	attggagttc	caaaaatcgg	tttctgccct	420
cggggacgta	gctgctagt	ggttcggggt	tcacgatgtg	gttattgaat	cgccggttca	480
ttcaataaat	atgagcgatc	tttcgccgtc	tcaagttggg	gaagttagtc	tggtctttta	540
gaagcgggtc	gagcagcttc	agaattgtga	taatgtcaaa	tatgttcagg	ttttattctg	600
atcccgctca	aaaatgcatt	ggttttcgta	tcaattttgc	tatatattgt	actgcatatt	660
cattctttga	ttttatatga	attttaaaat	aaatgcattt	aaaagctaca	taaaaatagt	720
agagaccatg	actgttaaaa	gctaaaagta	tttaacaaag	tatatattga	aggcttactg	780
caatgggtgaa	ttttttgttt	gcggtaaaat	actttgaaga	tttaacctgc	tttaattctt	840
tggtgcgatt	gcttatatgg	gttcattttc	attaaactga	aagtggtttg	tttttgtaca	900
aataaaat	aagagtaaac	ggtaaatggta	gagcccatca	atgtttgtac	aaagtggcca	960
accaattcta	gctctcttta	tggttatatt	agcaaaccta	ttttgtcatt	gagtttaagt	1020
tatatattag	ttctttttat	ttcagctgct	tatcatttgt	ctgctttggg	tttgtggctt	1080
tttaattttg	cagactgaat	gattttttct	tgtaacattg	gtggagagaa	ggctgacaac	1140
atgtttgtct	ttagggttta	atagaacaaa	tataagctag	ttccaaatac	tttagatctt	1200
gctgctttga	agtttggttac	tggaagagta	gctctttta	cagaccaaact	tgctaaaata	1260
ctgaacacaaa	ttatgagagg	tttttggttg	aaactcatag	actgtcaagt	gtcaactgct	1320
tgagggtcta	tcattaagga	gaataatttt	agagggaagt	tttccattaa	tctgcataag	1380
agcttggtgat	atgaactctc	catccattta	tctgatattg	tttgattttc	attgtagatt	1440
atttgagttt	gtattttttag	tgacaactga	caggagaaaa	tctggtagca	tttaaatgaa	1500
gttggtttga	tatggtaaat	atcaatttga	ctgtaaat	gaactattta	acagaactac	1560

-continued

```

tcttaaaactt gtaagtcacg tagcagttta tcccaaaaaa aaataaaaaa ccttgtaaat 1620
catgtagcat atttgtagta tcaagcaaaa ccttgttgga tatccagat tacaaaacat 1680
ttttgagagg tattgttagc tgcacattct tgttacttgg ttccctgtcc attgcagaat 1740
actctaaaaat atgcacttct atctactcat accaaatata tgtaagagaa cttgagaagt 1800
gattagttct cataaatttg agatgaatga gcacaatata tgatctcatg ccttttgtgc 1860
gtgcaaggaa tttgcagtag ggaacagcat gtattgtttt ttccactttc ttgttgagcg 1920
ctaagtattt atactgtact cagcaagatt gtcccatcaa agggctactg gactatataa 1980
cactaattca ttcaatggct tataaagggtg ttcaaaaacc atggagcctc tgctggtgct 2040
tcaatgagcc attctcagag ccagatgatt gctcttccca ttgttccctc agcagtttct 2100
gctcgtcttg acagtagtag ggaatactac aagcagactg ggaagtgcag cctttgtgat 2160
attcaaccac atgaattggt aattgatcaa tcagcccatc tcatactatt ggttcccttt 2220
gctgcaactt ttgcttttga gatttggata attcctcgtg atcactcctc tcattttcat 2280
gaactcgata atgaaaaggc aatcgatgct ttctctgtcc ataagtttgt ttcatgaga 2340
aaattatgtg aagttacaca gttttaactt tgcaaatatc agttatcccc aaattctacg 2400
tgagaaaagt cacaaaggta gaccatgagg gactttgact ttctcgaagt agaatttgc 2460
gattagttaa gagaaaagga aactcataa aaaaactgat attctctgtt tcaggctaaa 2520
ctaattttta gattgtttct ctatctcaac aaccacttag ttgaaaaggc ttgtgtatgt 2580
tttaccatcc aaatatgcag gcaaatgata ttggagggtt gttgaaactc gtgcttttga 2640
agatgtcttt gcagttgaac aatccaccat tcaacctcct tttcatacc tccccatttc 2700
aagtagatcc ctcatatgct ccttccacac actggttttt acaaatagct cccccattga 2760
gtggagtttg tggattcgag atagcaaccg gatgtcacat aaatcctggt ttcccagagg 2820
atgctgccaa aattctaaag gatgtgagaa tccttaatta acagtattta gggggaatct 2880
attaaatcta attatgcata tctttagaca cagtaactaa ggagataaag aggatgtggc 2940
tacaggggag ttgtaccctt tttatgcgga gttatgcgta gattcgattt tgtaatccat 3000
tacaatgtt ttcttatgtt acacggtcat tttcacgtaa aaaaataatc atgagat 3057

```

SEQ ID NO: 5 moltype = DNA length = 1020

FEATURE Location/Qualifiers

source 1..1020
mol_type = genomic DNA
organism = unidentified

SEQUENCE: 5

```

atggcggagg cagaagcacc aaatcggagc cccgaaatcc ggaaagacag aatccacaac 60
aggtgggtcc tattctcgcc ggccagatca cggcggccgt ctgacttcaa ggcaaaatcg 120
aaccacaac cgaacaataa aacagaatgc ccatttttgc ccggccacga gcacgagtgt 180
cgccgggaga ttttctggtg gccggcagat tccaccgctg attggaaaat cagagttatc 240
caaaatctct ttccggcagt cagcagagaa ttggagttcc aaaattcggc ttctgcccctc 300
ggggacgtag ctgctagtgg gttcgggttt cagcatgtgg ttattgaatc gccgggttcat 360
tcaataaata tgagcgatct ttccgctgct caagttgggg aagttatgct ggcttttaag 420
aagcgggtcg agcagcttca gaattgtgat aatgtcaaat atgttcaggc gttcaaaaac 480
catggagcct ctgctgtgtg ttcaatgagc cattctcaca gccagatgat tgctcttccc 540
attgttctct cagcagtttc tgctcgtctt gacagtagta gggaatacta caagcagact 600
gggaagtcca gcttttgtga tattcaacca aatgaattgt taattgatca atcagcccat 660
ttcatatcat tggttccctt tgctgcaact tttgcttttg agatttggat aattcctcgt 720
gatcactcct ctcattttca tgaactcgat aatgaaaagg caaatgatct tggaggggtg 780
ttgaaaactc tgccttttga gatgtctttg cagttgaaca atccaccatt caacctcctt 840
attcatacct ccccatttca agtagatccc tcatatgctc cttccacaca ctgggtttta 900
caaatagctc cccatttgag tggagtttgt ggattcgaga tagcaaccgg atgtcacata 960
aatcctgttt tcccagagga tgctgccaac attctaaggc atgtgagaat ccttaattaa 1020

```

SEQ ID NO: 6 moltype = AA length = 339

FEATURE Location/Qualifiers

source 1..339
mol_type = protein
organism = unidentified

SEQUENCE: 6

```

MAEAEAPNRS PEIRKDRHNN RWVLFSPARS RRPSPDFKAKS NPQPNKTEC PFCAGHEHEC 60
APEIFRPVAD STADWKIRVI QNLFPVRSRE LEFQNSVSAL GDVAASGFGF HDVVIESPVI 120
SINMSDLSPS QVGEVMLAFK KRVEQLQNCD NVKYVQVFKN HGASAGASMS HSHSQMIALP 180
IVPPAVSARL DSMREYYKQT GKCSLCDIQP NELLIDQSAH FISLVPPAAT FAFEIWIIPR 240
DHSSHFHELD NEKANDLGGI LKLVLLKMSL QLNNPPFNLL IHTSPFQVDP SYAPSTHFWL 300
QIAPHLSGVG GFEIATGCHI NPVPEDAAK ILRDVRILN 339

```

SEQ ID NO: 7 moltype = DNA length = 720

FEATURE Location/Qualifiers

source 1..720
mol_type = genomic DNA
organism = Nicotiana tabacum

SEQUENCE: 7

```

atggcaatgg aaatgaatcc agctgacgaa acctgtgttt tctccgactc tcattctcct 60
gaatcgataa agcaacatct tcttgacgat tcagattttt ctgaaatttt ttctgcgatg 120
aattctagca acgaaatatt gcctaacagt cctagctcaa gtttttagcag cttcgaactc 180
gactgcagct tccttaattg ggaatgaaac tctgaggaaa catataatcc aactgatcag 240
aatccttcac atgaatccca tgaagagtag tccgagtcgg aggagaaaac ccaggggcct 300
ggggtggcgc gtgagaaaaa cgcgcgcgca gattggacgc ggtacatagg agtgaacagg 360
cgaccgtggg ggacgttttc ggcggagaca agagacccaa gtaggaaagg tgaaggtgca 420

```

-continued

```

aggetgtggt taggaactta cgagaccgca gaggatgcag cgtagctta cgatcaagcc 480
gctttcaaaa tccgcggctc gagagctcgg ctcaattttc ctcacttaat cggctcaaac 540
atgcctaagc cggctagagt tacagcgagg cgtagtcgta cgcgctcacc cgagccatcg 600
tcttcttcat ccacctcatc atcagaaaat gtaccaagga aaaggaatat agatgtgata 660
aattccatag ccaagcccaa attcctttgt catagcttga atttacagag attagcttaa 720

```

```

SEQ ID NO: 8      moltype = DNA length = 720
FEATURE          Location/Qualifiers
source           1..720
                 mol_type = other DNA
                 organism = synthetic construct

```

```

SEQUENCE: 8
atggcaatgg aaatgaatcc agctgacgaa acctgtgttt tctccgactc tcattctcctt 60
gaatcgataa agcaacatct tcttgacgat tcagattttt ctgaaatttt ttcgtcgatg 120
aattctagca acgaaatatt gcctaacagt cctagctcaa gttttagcag cttcgacttc 180
gactgcagct tccttaattg gtaggaaaac tctgaggaaa cattaatacc aactgatcag 240
aatccttcac atgaatccca tgaagagtac tccgagtcgg aggagaaaac ccagggccct 300
ggggtggcgc gtgagaaaaa cgcgcgcgca gattggacgc ggtacatagg agtgaacagg 360
cgaccgtggg ggacgttttc ggcggagaca agagacccaa gtaggaaagg tgaaggtgca 420
aggetgtggt taggaactta cgagaccgca gaggatgcag cgtagctta cgatcaagcc 480
gctttcaaaa tccgcggctc gagagctcgg ctcaattttc ctcacttaat cggctcaaac 540
atgcctaagc cggctagagt tacagcgagg cgtagtcgta cgcgctcacc cgagccatcg 600
tcttcttcat ccacctcatc atcagaaaat gtaccaagga aaaggaatat agatgtgata 660
aattccatag ccaagcccaa attcctttgt catagcttga atttacagag attagcttaa 720

```

```

SEQ ID NO: 9      moltype = DNA length = 720
FEATURE          Location/Qualifiers
source           1..720
                 mol_type = genomic DNA
                 organism = Nicotiana tabacum

```

```

SEQUENCE: 9
atggcaatgg aaatgaatcc agctgacgaa acctgtgttt tctccgactc tcattctcctt 60
gaatcgataa agcaacatct tcttgacgat tcagattttt ctgaaatttt ttcgtcgatg 120
aattctagca acgaaatatt gcctaacagt cctagctcaa gttttagcag cttcgacttc 180
gactgcagct tccttaattg gtaggaaaac tctgaggaaa cattaatacc aactgatcag 240
aatccttcac atgaatccca tgaagagtac tccgagtcgg aggagaaaac ccagggccct 300
ggggtggcgc gtgagaaaaa cgcgcgcgca gattggacgc ggtacatagg agtgaacagg 360
cgaccgtggg ggacgttttc ggcggagaca agagacccaa gtaggaaagg tgaaggtgca 420
aggetgtggt taggaactta cgagaccgca gaggatgcag cgtagctta cgatcaagcc 480
gctttcaaaa tccgcggctc gagagctcgg ctcaattttc ctcacttaat cggctcaaac 540
atgcctaagc cggctagagt tacagcgagg cgtagtcgta cgcgctcacc cgagccatcg 600
tcttcttcat ccacctcatc atcagaaaat gtaccaagga aaaggaatat agatgtgata 660
aattccatag ccaagcccaa attcctttgt catagcttga atttacagag attagcttaa 720

```

```

SEQ ID NO: 10     moltype = AA length = 239
FEATURE          Location/Qualifiers
source           1..239
                 mol_type = protein
                 organism = Nicotiana tabacum

```

```

SEQUENCE: 10
MAMEMNPADE TLFFSDSHLL ESIKQHLLDD SDFSEIFSSM NSSNEILPNS PSSFSSSFD 60
DCSFLNWDEN SEETLIPTDQ NPSHESHEKY SESEKQTQGP GVAREKNAPR DWTRYIGVKR 120
RPWGTFSAPT RDPSPKGEA RLWLGTYETA EDAALAYDQA AFKIRGSRAR LNFPHLIGSN 180
MPKPARVTAR RSRTRSPEPS SSSSTSSSEN VPRKRNI DVI NSIAKAKFLC HSLNLQRLA 239

```

```

SEQ ID NO: 11     moltype = DNA length = 699
FEATURE          Location/Qualifiers
source           1..699
                 mol_type = genomic DNA
                 organism = Nicotiana tabacum

```

```

SEQUENCE: 11
atggaaatga atctagctga cgaacacctg tttttctctg agtctcatct ccttgaatcg 60
ataaagcaac atcttcttga tgattcagat ttttttgaaa ttttttcgcc gatgagtcca 120
agcaacgaaa tattgcctaa cagtcttagc tcaagtttta gcagcttcga ctgcagcttc 180
ctcaattggg atgaaaactt tgaggaaaac ttaataccaa ctgatcaaaa tcttccatc 240
gagaagtgtc ccgagtcgca ggagcaaac ccaggcccgag cgggtggtgcg tgagaaaaac 300
gcccgcgagc attggacgag gtatatagga gtgaaacggc ggccgtgggg gacgttttcg 360
gcggagacaa gagaccgaag taggaaaggc gaaggtgcaa ggctgtggtt aggaacttac 420
gagaccgcagc aggatgcagc gttggcttac gatcaagccg ctttcaaaat ccgcggctcg 480
agagctcggc tcaattttcc ccacttaatt ggctcaaaac tgcctaagcc ggctagagta 540
acagcgaggc gtagtcgtac gcgctcacc gagccatcct cttcttcate cacttcatca 600
tcagaaaatg tgccaagaaa aaggaatata gatgtgataa attccatagc caaagccaaa 660
ttcctttgtc atagcttaaa tttacagaga ttagcttaa 699

```

```

SEQ ID NO: 12     moltype = DNA length = 699
FEATURE          Location/Qualifiers

```

-continued

```

source                1..699
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 12
atggaatga atctagctga cgaacacctg tttttctctg agtctcatct ccttgaatcg 60
ataaagcaac atcttcttga tgattcagat ttttctgaaa ttttttcgcc gatgagttca 120
agcaacgaaa tattgcctaa cagtcctagc tcaagtttta gcagcttcga ctgcagcttc 180
ctcaattggg atgaaaactt tgaggaaaca ttaataccaa ctgatcaaaa tccttcacat 240
gagaagtgtc ccgagtcgga ggagcaaac ccaggcccgag cgggtggtgcg tgagaaaaac 300
gcgcgcgcgag attggacgcg gtatatagga gtgaaacggc ggccgtgggg gacgttttcg 360
gcggagacaa gagacccaag taggaaaggt gaaggtgcaa ggctgtggtt aggaacttac 420
gagaccgcag aggatgcagc gttggcttac gatcaagccg ctttcaaaat ccgcggctcg 480
agagctcggc tcaatttttc ccacttaatt ggctcaaaac tgccaaagcc ggctagagta 540
acagcgaggc gtagtcgtac gcgctcacc gagccatcct cttcttcac cacttcacatca 600
tcagaaaatg tgccaagaaa aaggaatata gatgtgataa attccatagc caaagccaaa 660
ttcctttgtc atagcttaaa ttacagaga ttagcttaa 699

SEQ ID NO: 13        moltype = DNA length = 699
FEATURE              Location/Qualifiers
source                1..699
                      mol_type = genomic DNA
                      organism = Nicotiana tabacum

SEQUENCE: 13
atggaatga atctagctga cgaacacctg tttttctctg agtctcatct ccttgaatcg 60
ataaagcaac atcttcttga tgattcagat ttttctgaaa ttttttcgcc gatgagttca 120
agcaacgaaa tattgcctaa cagtcctagc tcaagtttta gcagcttcga ctgcagcttc 180
ctcaattggg atgaaaactt tgaggaaaca ttaataccaa ctgatcaaaa tccttcacat 240
gagaagtgtc ccgagtcgga ggagcaaac ccaggcccgag cgggtggtgcg tgagaaaaac 300
gcgcgcgcgag attggacgcg gtatatagga gtgaaacggc ggccgtgggg gacgttttcg 360
gcggagacaa gagacccaag taggaaaggt gaaggtgcaa ggctgtggtt aggaacttac 420
gagaccgcag aggatgcagc gttggcttac gatcaagccg ctttcaaaat ccgcggctcg 480
agagctcggc tcaatttttc ccacttaatt ggctcaaaac tgccaaagcc ggctagagta 540
acagcgaggc gtagtcgtac gcgctcacc gagccatcct cttcttcac cacttcacatca 600
tcagaaaatg tgccaagaaa aaggaatata gatgtgataa attccatagc caaagccaaa 660
ttcctttgtc atagcttaaa ttacagaga ttagcttaa 699

SEQ ID NO: 14        moltype = AA length = 231
FEATURE              Location/Qualifiers
source                1..231
                      mol_type = protein
                      organism = Nicotiana tabacum

SEQUENCE: 14
MEMNLADETL FFSHLLLES IKQHLLDDSD FSEIFSPMSS SNEILPNSPS SSFSSFDCSF 60
LNWDENFEET LIPTDQNPSE EKSESEEEQT QGPAVVREKN APRDWTRYIG VKRRPWGTF 120
AETRDPSRK EGARLWLGTY ETAEADALAY DQAFKIRGS RARLNFPHLI GSNMPKPARV 180
TARRSRTRSP BPSSSSSTSS SENVPRKRNI DVINSIAKAK FLCHSLNLQR L 231

SEQ ID NO: 15        moltype = DNA length = 20171
FEATURE              Location/Qualifiers
source                1..20171
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 15
gatggtaaga agaaaggcgc gcagtatgcy atagctcttc acagcctgta tgacttcaag 60
ttgaaagact tgatggctac tatgggttag aagaaaacta aagtgggtca tgctgctatg 120
ctttttgtct ctgaaagtat gttagtggac gaaggtccat taccttctgt tgacggttac 180
tacatgaaga agaacgggaa gatctatctt ggttttgaga aagatccttc cttttcttac 240
attcatgact gggaagagta caagaagtat ctactgggga agccagtgcg ttaccaaggg 300
aatgtgttct acttcgaacc gtggcaggtg agaggagaca caatgctttt ttcatctac 360
aggatagctg gagttccgag gaggctctta tcatcgcaag agtactaccg aagaatataat 420
atcagtgaat gggaaagcat gttgtgtgtc ccaatttttc atctgggtcg atcaacgcga 480
gagttggtca agaaagacct gttttagtag aaacaattca tggacaagtg ttggattac 540
atagctaggt tatctgacca gcagctgacc ataagcaatg ttaaatcata cttgagttca 600
aataattggg tcttattcat aaacggggcg gccgtgaaga acaagcaag tgtagattct 660
cgagatttac agttgtgtgc tcaaaccttg ctagtgaagg aacaagtggc gagacctgtc 720
atgagggagt tgcgtgaagc aattctgact gagacgaaac ctatcacgct attgactgat 780
gtgctgggtt taatatcaag aaaactgtgg aagcagtttg ctaacaagat cgcagtcggc 840
ggattcgttg gcatggttgg tactctaatt ggattctatc caaagaaggt actaacctgg 900
gcgaaggaca caccaaatgg tccagaacta tgttacgaga actcgcacaa aaccaaggtg 960
atagtatttc tgagtgttgt gtagtccatt ggaggaatca cgcttatgcy tcgagacatc 1020
cgagatggac aggtgaaaaa acctatgtgat atgtttgata tcaaacgggg ggcccatgct 1080
ttagacgttg agaatccgtg ccgctattat gaaatcaacg atttcttttag cagtctgtat 1140
tcggcatctg agtccgtgga gaccgtttta ccagatttat ccgaggtaaa agccaagtct 1200
gataagctat tgcagcagaa gaaagaaatc gctgacgagt ttctaagtgc aaaaattctct 1260
aactattctg gcagttcggg gaagaacttc ccaccatcgg tggtcgggtt atctcgaagc 1320
ggactgggtc tgttgttgga agacagtaac gtgctgaccg aagctagagt tggagtttca 1380

```

-continued

agaaaggtag	acgatgagga	gatcatggag	cagtttctga	gtggtcttat	tgacactgaa	1440
gcagaaattg	acgagggttg	ttcagccttt	tcagctgaat	gtgaaagagg	ggaaacaagc	1500
ggtacaaagg	tggtgtgtaa	acottttaacg	ccaccaggat	ttgagaacgt	gttgccagct	1560
gtcaaacctt	tggtcagcaa	aggaaaaacg	gtcaaacgtg	tcgattactt	ccaagtgatg	1620
ggaggtgaga	gattaccaaa	aaggccggtt	gtcagtgagg	acgattctgt	ggacgctaga	1680
agagagtttc	tgtactactt	agatgcggag	agagtgcctc	aaaatgatga	aattatgtct	1740
ctgtatcgtg	actattcgag	aggagttatt	cgaactggag	gtcagaatta	cccgcaaggga	1800
ctgggagtg	gggatgtgga	gatgaagaac	tggtgcatac	gtccagtggt	cactgaacat	1860
gcttatgtgt	tccaaccaga	caaacgtatg	gatgattggt	cgggataactt	agaagtggct	1920
gtttgggaac	gaggtatggt	ggtcaacgac	ttcgcggctg	aaaggatgag	tgattatgtc	1980
atagttttcg	atcagacgta	tctttgcaat	aacaggtaat	aatcctctct	cttgatattt	2040
ttaaattata	gaatttaatta	gtttacttta	ttctttacta	tatgatttaa	atagtttaat	2100
cttggttttg	agtaaacctat	tcgattttga	tatttgtatt	cgtcctacaa	agtggaatat	2160
actgatgata	ttttcttttg	aacgtgatac	ctaccaatag	taatcttacy	gaatctttta	2220
atagagcact	aatacaactat	gaactaaaga	ccaattctta	agtgctctcg	ttgtacagtt	2280
catttttagta	gtgcgtttta	gtattattat	ctcccttcat	gcggggcaat	tatgtagatt	2340
aaaatcgaaa	ttatataaaa	tttataatag	tctaagctta	gggtctccag	ctaattgtta	2400
tttttttaac	gatgttgact	aaagcaataa	cgacgttgac	ttgtgtttaa	cagggtgatc	2460
ttggacaatt	taagtgccct	ggatctagga	ccagtttaact	gttcttttga	attagttgac	2520
gggtgacctg	gttgtgtgta	gtcgacaatg	attgtcaact	cagctaactc	ttgtgtcgat	2580
gtgggttctc	ctaactggag	agcagcaacc	gacgacttga	tcgagagatt	cgcgagcaaa	2640
gggtttccat	gcaaatagaa	aggagagtg	aagacgggtg	attctttttt	gatgcattgt	2700
gtcgatggtt	ctttaaccgg	agacgtgtgt	catttcgacg	aagctctcat	ggcccatgct	2760
ggtatggtgt	acttttgcgc	tcagatagct	gggtgctaac	gatgtatctg	tcaaggagat	2820
cagaatcaaa	tttctttcaa	gcctagggtta	tctcaagttg	atttgaggtt	ttctagtctg	2880
gtcggaaagt	ttgacattgt	tacagaaaaa	agagaaaact	acagaagtcc	agcagatgtg	2940
gctgcgctat	tgaacaagta	ctatactgga	gatgtcagaa	cacataacgc	gactgctaatt	3000
tcgatgacgg	tgaaggaagt	tgtgtctaaa	gaacagggtt	ctttgaagcc	tggtgctcag	3060
tacataaact	tccttcagtc	tgagaagaag	gagttggtaa	atttgttgcc	attgaggaaa	3120
gtggcagcta	aagtgtgtac	agtacacgag	tcgcaaggag	agacattcaa	agatgtagt	3180
ctagtacgga	gcaaacctca	ggatgactca	atcgctagag	gtcgggagta	cttaactcgt	3240
gcattgtcgc	gtcacacaca	atcactttgt	tatgaaactg	tgaaagagga	cgatgtaagc	3300
aaagagatca	gggaaagtgc	cgcgcttacg	aaggcggctt	tggcaagatt	ttttgttact	3360
gagaccgtct	tatgacggtt	tcggtctagg	tttgatgtct	ttagacatca	tgaaggccct	3420
tcgcgcgttc	cagattcagg	tacgattacg	gacttgagga	tgtggtacga	cgctttgttt	3480
ccgggaaatt	cgtaagagaa	ctcaagccta	gacgggtatt	tggtggcaac	gactgatgtc	3540
aatttgcgat	tagacaatgt	tacgatcaaa	agtggaacct	ggaaagacaa	gtttgctgaa	3600
aaagaaacgt	ttctgaacc	gggtattcgt	actgctatgc	ctgacaaaag	gaagactact	3660
cagttggaga	gtttgttagc	attgcagaaa	aggaaaccaag	cggcacccca	tctacaagaa	3720
aatgtgcacg	cgacacatgt	aatcgaaagag	acgatgaaga	agctgaaatc	tgttgtctac	3780
gatgtgggaa	aaattcgggc	tgatcctatt	gtcaatagag	ctcaaatgga	gagatgggtg	3840
agaaatcaaa	gcacagcggt	acaggctaag	gtagtagcac	atgtgagaga	gttacatgaa	3900
atagactatt	tgctttacat	gtatatgac	aaatctgacg	tgaacctcaa	gactgattta	3960
acaccgcaat	ttgaatactc	agctctacag	actgttgtgt	atcacgagaa	gttgatcaac	4020
tcgttgttcg	gtccaatttt	caaagaaatt	aatgaacgca	agttggatgc	tatgcaacca	4080
cattttgtgt	tcaacacgag	aatgacatcg	agtgatttaa	acgatcgagt	gaagttctta	4140
aatacggaag	cggtctacga	ctttgttgag	atagacatgt	ctaaattcga	caagtccggca	4200
aatcgcttcc	atttacaact	gcagctggag	atttacaggt	tatttgggct	ggatgagttg	4260
gcggccttcc	tttgggaggt	gtcgacacat	caaaactactg	tgagagatat	tcaaaatgggt	4320
atgatggcgc	atatttggta	ccaacaaaag	agtgagagtg	ctgatactta	taatgcataa	4380
tcagatagaa	cactgtgtgc	actctgtctc	gaattaccat	tggagaaagc	agtcagtggt	4440
acatatggag	gagatgactc	actgatttgc	tttcttagag	gaacgcagtt	tgttgatccg	4500
tgtccaaagt	tggtacttaa	gtggaatttc	gagtgcaaga	tttttaagta	cgatgtccca	4560
atgttttgtg	ggaagttctt	gcttaagacg	tcatcgtgtt	acgagttcgt	gccagatccg	4620
gtaaaagttc	tgacgaagtt	ggggaaaaag	agtataaagg	atgtgcaaca	tttagccgag	4680
atctacatct	cgctgaatga	ttccaataga	gctcttggga	actacatggt	ggtatccaaa	4740
ctgtccgagt	ctgtttcaga	cgggtatttg	tacaaagggt	attctgttca	tgcgctttgt	4800
gcgctatgga	agcatattaa	gagtttttaca	gctctgtgta	cattattccg	agacgaaaac	4860
gataaggaat	tgaacccggc	taagggttgat	tggaaagaag	cacagagagc	tgtgtcaaac	4920
ttttacgact	ggtaaatatg	aagacaagtc	attggtcacc	ttgaagaaga	agactttcga	4980
agttctcaaa	ttctcaaatc	taggggccat	tgaattgttt	gtggacggta	ggaggaagag	5040
accgaagtat	tttcacagaa	gaagagaaac	tgtcctaata	catgtttggg	ggaagaagag	5100
tgaacacaa	ttagacgttt	ttgaccaaag	ggattacaaa	atgattaaat	cttacgcggt	5160
tctaaaggta	gtagggtgat	aactagttgt	aacatcacat	ctacctgcag	atacgccctg	5220
gttcatctaa	atcgatctgt	tggtatcgag	acttactgag	aaaagaaaag	gaggaagagc	5280
tattcagaga	ttcaaaagctc	gagcttgcca	taactgttca	gttcgcaggt	acaaggttga	5340
atcaggtatt	tccacacagg	agaacgtact	tgatgtctgg	aaggtgggtt	gtatttctga	5400
gggcgttccg	gtctgtgaag	gtacataccc	tttcagtatc	gaagtgtcgc	taatatgggt	5460
tgctactgat	tcgactaggc	gcctcaatgt	ggaagaactg	aacagttccg	attacattga	5520
aggcgatttt	accgatcaag	aggttttccg	tgaagttcat	tctttgaaac	aagtgagagat	5580
gaagacgatt	gaggcgaaag	acgatgggtcc	ttacagacca	gctactacta	gacctaaagt	5640
attattgtca	agtgaagatg	ttaaagagag	gtctaataag	aaaaactcgt	cttaatgcat	5700
aaagaaattt	attgtcaata	tgacgtgtgt	actcaagggt	tgtgtgaatg	aagtcaactgt	5760
tcttgggtcac	gagacgtgta	gtatcgggtca	tgctaacaaa	ttgcgaaagc	aagttgctga	5820
catgggtgtg	gtcacacgta	gggtgtcgga	aaataattgt	ggatgggtttg	tctgtgttgt	5880
tatcaatgat	tttacttttg	atgtgtataa	ttgtgtgggc	cgtagtccac	ttgaaaagtg	5940

-continued

tcgtaaacgt	gttgaacaa	gaaatcgaga	aatttggaaa	caaattcgac	gaaatcaagc	6000
tgaaaacatg	tctgcgacag	ctaaaaagtc	tcataattcg	aagacctcta	agaagaaatt	6060
caaaaggagc	agagaatttg	ggacacccaaa	aagattttta	agagatgatg	ttcctttcgg	6120
gattgatcgt	ttgtttgctt	tttgatttta	ttttatattg	ttatctgttt	ctgtgtatag	6180
actgtttgag	attggcgctt	ggcgcactca	ttgtcttacc	ataggggaac	ggactttgtt	6240
tggtgtgtta	ttttatttgt	attttattaa	aattctcaat	gatctgaaaa	ggcctcgagg	6300
ctaagagatt	attggggggt	gagtaagtac	ttttaaagtg	atgatggtta	caaaggcaaa	6360
aggggtaaaa	ccctcgcct	acgtaagcgt	tattacgccc	ggatcccccg	gggagctcga	6420
attcgcgtga	atcaccagtc	tctctctaca	aatctatctc	tctctatttt	ttccataaat	6480
aatgtgtgag	tagtttcccg	ataagggaaa	ttagggttct	tataggggtt	cgctcatgtg	6540
ttgagcctat	aagaaacccct	tagtatgtat	ttgtatttgt	aaaatacttc	tattatcaat	6600
aaaattttcta	attcctaaaa	ccaaaatcca	gtactaaaa	ccagatctcc	taaagtcctc	6660
atagatcttt	gtcgtgaata	taaaccagac	acgagacgac	taaacctgga	gcccagacgc	6720
cggttcgaagc	tagaagtacc	gcttaggcag	gaggccgtta	gggaaaagat	gctaaggcag	6780
ggttgggttac	gttgactccc	ccgtagggtt	ggtttaataa	tgatgaagtg	gacggaagga	6840
aggaggaaga	caaggaagga	taagggtgca	ggcctgtgc	aaggtaaaga	gatggaaatt	6900
tgatagaggt	acgctactat	acttatacta	tacgctaagg	gaatgcttgt	atttataccc	6960
tataccccct	aataacccct	tatcaattta	agaaataatc	cgcataagcc	cccgttataa	7020
aattgggtatc	agagccatga	ataggtctat	gacccaaact	caagaggata	aaacctcacc	7080
aaaaatcga	agagttctta	actctaaaga	taaaagatct	ttcaagatca	aaactagtcc	7140
cctcacaccg	gagcatcgca	tatcctcgac	ctgcaggcat	gcaagcttgg	cgtaatcatg	7200
gtcatagctg	tttctgtgtg	gaaattgtta	tccgctcaca	attccacaca	acatacgagc	7260
cgggaagcata	aagtgtaaa	ctcgggggtg	ctaattgagt	agctaaactc	cattaatgtc	7320
gttcgcgtca	ctgcccgtt	tccagtcggg	aaacctgtcg	tgccagctgc	attaatgaat	7380
cggccaacgc	ggggggagag	gcggtttgcg	tattgggcca	aagacaaaag	ggcgacattc	7440
aaccgatgga	gggaggggaag	gtaaatattg	acggaaatta	ttcattaaag	gtgaattatc	7500
accgtcacccg	acttgagcca	tttggaatt	agagccagca	aaatcaccag	tagcaccatt	7560
accattagca	aggccggaaa	cgtcaccaat	gaaaccatcg	atagcagcac	cgtaatcagt	7620
agcgacagaa	tcaagtttgc	ctttagcgtc	agaactgtag	gcgttttcat	cggcattttc	7680
ggtcataacc	cccttattag	cgtttgccat	cttttcataa	tcaaaatcac	cggaaaccaga	7740
gccaccaccg	gaaccgcctc	cctcagagcc	gccaccctca	gaaccgccac	cctcagagcc	7800
accaccctca	gagccgcac	cagaaccacc	accagagccg	ccgccagcat	tgacaggagg	7860
cccgatctag	taacatagat	gacaccgcgc	gcgataattt	atcctagttt	gcgcgctata	7920
ttttgttttc	tatcgcgtgt	taaatgtata	attgcgggac	tctaatcata	aaaaccocat	7980
tcataaataa	cgtcatgcat	tacatgttaa	ttattacatg	cttaacgtaa	ttcaacagaa	8040
atttatatgat	aatcatcgca	agaccggcaa	caggattcaa	tcttaagaaa	ctttatttgc	8100
aaatgtttga	acgatccggg	atcatccggg	tctgtggcgg	gaactccacg	aaaatatccg	8160
aacgcagcaa	gatatccggg	tgcactcctg	tcttgccctg	gcagtcgccc	ccgacgccgt	8220
tgatgtggac	gccggggccc	atcatattgt	cgctcaggat	cggtggcgtg	tgcttgctcg	8280
ccgttgctgt	cgtaatgata	tccgcacctt	cgaccgcctg	ttccgcagag	atcccgtggg	8340
cgaagaactc	cagcatgaga	tccccgcgct	ggaggatcat	ccagccggcg	ttccggaaaa	8400
cgattccgaa	gcccacacct	tcatagaagg	cggcggtgga	atcgaaatct	cgatgatggc	8460
ggttgggcgt	gtcttggtcg	gtcatttcga	acccagagat	cccgtccaga	agaactcgtc	8520
aagaaggcga	tagaaggcga	tgccgtcgca	atcggggagc	gcgataccgt	aaagcacgag	8580
gaagcgggtca	gcccattcgc	cgccaaagtc	ttcagcaata	tcacgggtag	ccaacgctat	8640
gtcctgatag	cggtccgcga	caccacagcc	gccacagtcg	atgaatccag	aaaagcggcc	8700
attttccacc	atgatattcg	gcaagcaggc	atcgccatgg	gtcacgacga	gatcatcgcc	8760
tcggggcatg	cgcccttga	gcttgccgaa	cagttcggct	ggcgcgagcc	cctgatgctc	8820
ttcgtccaga	tcactcgtga	gcacaagacc	ggcttccatc	cgagtacgtg	ctcgctcgat	8880
gcgatgtttc	gcttggtggt	cgaatgggca	ggtagccgga	tcaagcgtat	gcagccgccc	8940
cattgcatca	gccatgatgc	atactttctc	ggcaggagca	aggtgagatg	acaggagatc	9000
ctgccccggc	acttcgccca	atagcagcca	gtcccctccc	gcttcagtga	caacgtcgag	9060
cacagctcgc	caagggaacc	ccgtcgtggc	cagccacgat	agccgcgctg	cctcgtcctg	9120
cagttcattc	agggcaccgg	acaggtcggt	cttgacaaaa	agaaccgggc	gcccctgcgc	9180
tgacagccgg	aacacggcgg	catcagagca	gccgattgtc	tgtgttgccc	agtcataagg	9240
gaatagcctc	tccaccacaag	cggccgggga	acctgcgtgc	aatccatctt	gttcaatcat	9300
gcgaacagat	ccagatccgg	tcagatttat	ttggattgag	agtgaatatg	agactcta	9360
tggtatccga	ggggaattta	tggaacgtca	gtggagcatt	tttgacaaga	aatatttgct	9420
agctgatagt	gaccttaggc	gacttttgaa	cgcgcaataa	tggtttctga	cgatgtgtgt	9480
tagctcatta	aactccagaa	accgcgggct	gagtggtccc	ttcaacgttg	cggttctgtc	9540
agttccaaac	gtaaaaaggc	ttgtcccgcg	tcactggcgg	gggtccataa	gtgactccct	9600
taattctccg	ctcatgatca	gattgtcggt	tcccgccttc	agttttaaact	atcagtgttt	9660
gacaggat	attggcggtt	aaacctaa	gaaaagagcg	tttattagaa	taatcgggata	9720
tttaaaagg	cgtgaaaagg	tttatccgtt	cgtccatttg	tatgtgcacg	ccaaccacag	9780
ggttccccag	atctggcgcc	ggccagcgag	acgagcaaga	ttggcccgcc	cccgaacaga	9840
tcgacagcgc	cgccacgac	aggtgcgcag	gcaaattgca	ccaacgcata	cagcgccagc	9900
agaatgccat	agtggcggtt	gacgtcgttc	gagtgaaaca	gatcgcgag	gaggcccgcc	9960
agcacccgca	taactcaggcc	gatgccgaca	gcgtcgagcg	cgacagtgtc	cagaattacg	10020
atcaggggta	tgttggtgtt	cacgtctggc	ctccggacca	gcctccgctg	gtccgattga	10080
acgcgcggat	tctttatcac	tgataagttg	gtggacatat	tatgtttatc	agtgataaag	10140
tgtcaagcat	gcataagttg	cagccgaata	cagtgatccg	tgccgcctcg	gacctgttga	10200
acgaggtcgg	cgtagacggt	ctgacgacac	gcaaaactgg	ggaacgggtt	gggggttcagc	10260
agccggcgct	ttactggcac	ttcaggaaca	agccggcgct	gctcgacgca	ctggccgaag	10320
ccatgctggc	ggagaatcat	acgcattcgg	tgccgagagc	cgacgacgac	tgccgctcat	10380
ttctgatcgg	gaatggccgc	agcttcaggc	agggcgtgct	cgccctaccg	gatggcgcg	10440
gcatccatgc	cggcacgcga	ccggggcgac	cgcagatgga	aacggccgac	gcgcagcttc	10500

-continued

gcttctctctg	cgaggcggggt	ttttcgggcg	gggacgcctg	caatgcgctg	atgacaatca	10560
gctactttcac	tggtggggccc	gtgcttgagg	agcaggcccg	cgacagcgat	gcccggcgagc	10620
gcgggcggcac	cggtgaacag	gctccgctct	cgccgctgtt	gcccggccgag	atagacgcct	10680
tcgacgaagc	cggtccggac	gcagcggttc	agcagggact	cgcggtgatt	gtcgatggat	10740
tggcgaaaaag	gaggtcgtgt	gtcaggaaac	ttgaaggacc	gagaaagggt	gacgattgat	10800
caggaccgct	gcccggagcgc	aacccactca	ctacagcaga	gccatgtaga	caacatcccc	10860
tccccctttc	caccgcgtca	gacgcccgtg	gcagcccgtc	acgggctttt	tcatgccctg	10920
ccttagcgctc	caagcctcac	ggcgcgcgtc	ggcctctctg	gcccgccttc	ggcgctcttc	10980
cgcttctctcg	ctcactgact	cgctgcgtc	ggctgttccg	ctgcggcgag	cggtatcagc	11040
tcactcaaaag	gcggtaatat	ggttatccac	agaatcaggg	gataacgcag	gaaagaacat	11100
gtgagcaaaa	ggccagcaaa	aggccaggaa	cctgaaaaag	gcccgcgtgc	tggcgttttt	11160
ccataggctc	cgccccctcg	acgagcatca	caaaaatcga	cgctcaagtc	agagggtggcg	11220
aaacccgaca	ggactataaa	gataccaggc	gtttccccct	ggaagctccc	tctgtcgctc	11280
tctgtttccg	accctgcgcg	ttaccggata	cctgtccgct	tttctccctt	cggaagcggt	11340
ggcgcttttc	cgctgcataa	cctgtcttcg	gggtcattat	agcgattttt	tccgtatatc	11400
catccttttt	cgcaogatat	acaggatttt	gccaaagggt	tctgtatagc	tttcttgggt	11460
gtatccaacg	gcgtcagccg	ggcaggatag	gtgaagtagg	cccaccccg	agcgggtggt	11520
cctttcttcac	tgctcccttat	tcgcacctgg	cggtgctcaa	cggaatcct	gctctgcgag	11580
gctggccggc	taccgcccgc	gtaacagatg	agggcaagcg	gatggctgat	gaaaccaagc	11640
caaccagga	gggagcccca	cctatcaagg	tgtactgcct	tccagacgaa	cgaagacgga	11700
ttgaggaaaa	ggcgggcgcg	gcccgcagta	gcctgtcgcc	ctacctgctg	gcccgtcgcc	11760
agggctacaa	aatcacgggc	gtcgtggact	atgagcacgt	ccgcgagctg	gcccgcacat	11820
atggcgacct	gggcccgcctg	ggcgccctgc	tgaacctctg	gctcacgcag	gaccccgcga	11880
cgcgccggtt	cggtgatgcc	acgatcctcg	cctgtctggc	gaagatcgaa	gagaagcagg	11940
acgagcttgg	caaggtcatg	atggcgctgg	tccgcccag	ggcagagcca	tgactttttt	12000
agccgctaaa	agccgcgggg	ggtgcgcgtg	attgccaaag	acgtcccttc	gcgctccatc	12060
aagaagagcg	acttcgcgga	gctggtgaag	tacatcaccc	acgagcaagg	caagaccgag	12120
cgcttttgcg	acgctcacgc	ggctggttgc	cctcgccgct	gggctggcgg	ccgtctatgg	12180
ccctgcaaac	gcccagaaaa	cgccgtcgaa	gcctgtgctg	agacacccgc	gcccgcggcg	12240
ttgtggatag	ctcgccgaaa	acttgccctc	cactgacaga	tgagggcgcg	acgttgacac	12300
ttgagggggc	gactcacccc	gcgcggcggt	gacagatgag	gggcaggctc	gatttcgggc	12360
ggcgacgtgg	agtggtccag	cctcgcaaat	ggcgaaaaac	gcctgatttt	acgcgagttt	12420
cccacagatg	atgtggacaa	gcttggggat	aagtgcctct	cggtattgac	acttgagggg	12480
cgcgactact	gacagatgag	ggcgccgcat	cttgacactt	gaggggcaga	gtgctgacag	12540
atgagggggc	cacctattga	catttgaggg	gctgtccaca	ggcagaaaa	ccagcatttg	12600
caaggggttc	cgcccggttt	tcggccaccg	ctaacctgtc	ttttaacctg	cttttaaac	12660
aatatttata	aaccttggtt	ttaacagggg	ctgcgcccgt	tgccggtgac	cgccgcagcc	12720
gaaggggggt	gccccccctt	ctcgaaacct	cccgcccgcc	taacgcgggc	ctcccatccc	12780
cccaggggct	gcgcccctcg	gcccgcgaac	gcctcacccc	aaaaatggca	gcgctggcag	12840
tccttgccat	gtgcgggcat	ggggcagtaa	cgggatgggc	gatcagcccg	agcgcgacgc	12900
ccggaagcat	tgacgtgcgc	cagggtgctg	catcgacatt	cagcgaccag	gtgcccggga	12960
gtgagggcg	cgccctgggt	ggcgccctgc	ccttcacttc	ggccgtcggg	gcattcacgg	13020
acttcattgc	ggggccggga	atttttacct	tgggcattct	tgccatagtg	gtcgcgggtg	13080
ccgtgctcgt	gttcgggggt	gcgataaaac	cagcgaaaca	tttgagggtg	taggtgaagt	13140
tataccgagg	tatgaaaacg	agaattggac	ctttacagaa	ttactctatg	aagcgcata	13200
tttaaaaagc	taccaagacg	aagaggatga	agaggatgag	gaggcagatt	gccttgaata	13260
tattgacaat	actgataaga	taatatatct	tttatataga	agatatcgcc	gtatgtaagg	13320
atctcagggg	gcaaggcata	ggcagcgccg	ttatcaatat	atctatagaa	tgggcgaagc	13380
ataaaaactt	gcattggacta	atgcttgaaa	cccaggacaa	taaccttata	gcttgttaat	13440
tctatcataa	ttgggtaatg	actccaactt	attgatagtg	ttttatgttc	agataatgcc	13500
cgatgacttt	gtcatgcagc	tcacccgatt	ttgagaacga	cagcgacttc	cgtcccagcc	13560
gtgcagggtg	ctgcctcaga	ttcaggttat	gcccgtcaat	tcgctgcgta	tatcgcttgc	13620
tgattacgtg	cagctttccc	ttcaggcggg	attcatacag	ggccagccca	tcctgcatcc	13680
atatcaccac	gtcaaaagggt	gacagcagcc	tcataagacg	cccagcgctc	gccatagtgc	13740
gttcaccgaa	tacgtgcgca	acaaccgtct	tcggagagct	gtcatacgcg	taaaacagcc	13800
agcgtggggc	cgattttagc	ccgacatagc	cccactgttc	gtccatttcc	gcgcagacga	13860
tgacgtcact	gcccggctgt	atgcgcgagg	ttaccgactg	cgccctgagt	tttttaagtg	13920
acgtaaaaatc	gtgttgaggc	caacgcccct	aatgcgggct	gttgcccggc	atccaaagcc	13980
attcatggcc	atatcaatga	ttttctgggt	cgtaccgggt	tgagaagcgg	tgtaagtga	14040
ctgcagttgc	catgtttttac	ggcagtgaga	gcagagatag	cgctgatgtc	cgccggtgct	14100
tttgccgtta	cgcaccaccc	cgtcagtagc	tgaacaggag	ggacagctga	tagacacaga	14160
agccactgga	gcacctcaaa	aacaccatca	tacactaaat	cagtaagttg	gcagcatcac	14220
ccataaattgt	ggtttcaaaa	tcgggtccgt	cgatactatg	ttatacgcca	actttgaaaa	14280
caactttgaa	aaagctgttt	tctggtatgt	aagggttttag	aatgcaaggga	acagtgaatt	14340
ggagttcgtc	ttgtttataat	tagcttcttg	gggtatcttt	aaatactgta	gaaaagagga	14400
aggaaaataat	aaatggctaa	aatgagaata	tcaccggaat	tgaaaaaact	gatcgaaaaa	14460
taccgctcgc	taaaagatac	ggaagggaat	tctcctgcta	aggtatataa	gctgggtggga	14520
gaaaatgaaa	acctatattt	aaaaatgacg	gacagccggt	ataaagggag	cacctatgat	14580
gtggaacggg	aaaaggacat	gatgctatgg	ctggaaggaa	agctgcctgt	tccaaagggtc	14640
ctgcactttg	aacggcatga	tggtctggagc	aatctgctca	tgagtgaagg	cgatggcgctc	14700
ctttgctcgg	aagagtatga	agctgaacaa	agccctgaaa	agattatcga	gctgtatgcg	14760
gagtgcatca	ggctctttca	ctccatcgac	atatcggtat	gtccctatac	gaatagctta	14820
gacagccgct	tagccgaatt	ggattactta	ctgaataacg	atctggccga	tgtggattgc	14880
gaaaactggg	aagaagacac	tccattttaa	gatccgcgcg	agctgtatga	ttttttaaag	14940
acggaaaaagc	ccgaagagga	acttgtcttt	tcccacggcg	acctgggaga	cagcaacatc	15000
tttgtgaaag	atggcaaaat	aagtggcttt	attgatcttg	ggagaagcgg	caggggcgac	15060

-continued

aagtggatg	acattgcctt	ctgcgtccgg	tcgatcaggg	aggatatcgg	ggaagaacag	15120
tatgtcgagc	tattttttga	cttactgggg	atcaagcctg	attggggagaa	aataaaatat	15180
tatatattac	tggatgaatt	gttttagtac	ctagatgtgg	cgcaacgatg	ccggcgacaa	15240
gcaggagcgc	accgacttct	tccgcacaa	gtgttttggc	tctcaggccg	aggcccacgg	15300
caagtatttg	ggcaaggvgg	cgctggatt	cgtagcaggg	aagattcgga	ataccaagta	15360
cgagaaggac	ggcgagacgg	tctacgggg	cgacttcatt	cccagataag	tggattatct	15420
ggacaccaag	gcaccaggcg	gggtcaaatca	ggaataaggg	cacattgccc	cgccgtgagt	15480
cggggcaatc	ccgcaaggag	gggtgaatgaa	tccgacgttt	gaccggaagg	catacaggca	15540
agaactgato	gacgcgggg	tttccgccc	ggatgccgaa	accatcgcaa	gccgcaccgt	15600
catgcgtgcg	ccccgcgaaa	ccttccagtc	cgtagcgtcg	atgggtccagc	aagctacggc	15660
caagatcgag	cgcgacagcg	tgcactggc	tccccctgcc	ctgcccgcgc	catcgccgcg	15720
cgtaggagcgt	tcgcgtcgtc	tcgaacagga	ggcggcaggt	ttggcggaag	cgatgaccat	15780
cgacacgcga	ggaactatga	cgaccaagaa	gcgaaaaacc	gccggcgagg	acctggcaaa	15840
acaggtcagc	gaggccaagc	aggcccggtt	gctgaaacac	acgaagcagc	agatcaagga	15900
aatgcagcgt	tccttggtcg	atattgcgcc	gtggccggac	acgatgcgag	cgatgccaaa	15960
cgacacggcc	cgctctgccc	tggtcaccac	gcgcaacaag	aaaatccgcg	gcgaggcgct	16020
gcaaaacaag	gtcattttcc	acgtcaacaa	ggacgtgaag	atcacctaca	ccggcgctcg	16080
gctgcggggc	gacgatgaag	aactggtgtg	gcagcaggtg	ttggagtacg	cgaaagcgac	16140
ccctatcgcc	gagcogacta	ccttcacgtt	ctacgagcgt	tgccaggacc	tgggctggtc	16200
gatcaatggc	gggtattaca	cgaaaggcga	ggaatgcctg	tcgcgcctac	aggcgacggc	16260
gatgggcttc	acgtccgacc	gcgtttgggca	cctggaatcg	gtgtcgctgc	tgcaaccgct	16320
ccgcgtcctg	gaccgtggca	agaaaacgtc	ccgttgccag	gtcctgatcg	acgaggaat	16380
cgctgtgctg	tttgcctggc	accactacac	gaaattcata	tgggagaagt	accgcaagct	16440
gtcgcgcagc	gcccgcagga	tgttcgacta	tttcagctcg	caccgggagc	cgtaccgcgt	16500
caagctggaa	accctccgcc	tcattgtcgg	atccgattcc	accgcgtga	agaagtggcg	16560
cgagcaggtc	ggcgaaagcct	gcgaagaggt	gcgagggcagc	ggcctgggtg	aacacgcctg	16620
ggatcaatgat	gacgtggtgc	attgcaaacg	ctagggccctt	gtggggctag	ttccggtcgt	16680
gggttcagca	cgcagcgtct	tactggcatt	tcaggaacaa	gcgggcactg	ctcgacgcac	16740
ttgcttcgct	cagtagcgtc	cgggacgcac	ggcgcgctct	acgaactgcc	gataaaacaga	16800
ggattaaaa	tgacaattgt	gattaaagct	cagattcgac	ggcttggagc	ggccgacgtg	16860
caaggattcc	gcgagatcct	attgtcggcc	ctgaagaaag	ctccagagat	gttcgggtcc	16920
gtttacgagc	acgaggagaa	aaagcccatg	gaggcgctcg	ctgaacgggt	gcgagatgcc	16980
gtggcattcg	gcgcctacat	cgacggcgag	atcatttggc	tgtcggtctt	caaacaggag	17040
gacggcccca	aggacgccta	caaggcgcat	ctgtccggcg	ttttcgtgga	gcccgaacag	17100
cgaggccgag	gggtcgcggg	tatgctgctg	cgggcggtgc	cgggcggttt	attgctcgtg	17160
atgatcgtcc	gacagattcc	aacgggaatc	tgggtgagtc	gcattctcat	cctcggcgca	17220
cttaatatct	cgctattctg	gagctgtgtg	tttatttcgg	tctaccgect	gcccggcggg	17280
gtcgcggcga	cggtaggcgc	tgtgcagccg	ctgatggctg	tggtcatctc	tgccgctcgt	17340
ctaggtagcc	cgatagcatt	gatggcggtc	ctgggggcta	tttcgggaac	tgccggcgctg	17400
gcgctgttgg	tttgacacac	aaacgcagcg	ctagatcctg	tcggcgtcgc	agcgggcctg	17460
gcggggcgcg	gttccatggc	gttcggaacc	gtgtgacccc	gcaagtggca	acctcccctg	17520
cctctgctca	cctttaccgc	ctggcaactg	gcggccggag	gacttctgct	cgttccagta	17580
gctttagtg	ttgatccgca	aatcccgatg	cctacaggaa	ccaatgttct	cgccctggcg	17640
tggtcgggcc	tgatcggagc	gggtttaacc	tacttccctt	gggtccgggg	gatctcgcca	17700
ctcgaaacct	cagttgtttc	cttactgggc	tttctcagcc	ccagatctgg	ggtagatcag	17760
ccgggggatgc	atcaggccga	cagtccgaa	ttcgggtccc	cgacctgtac	cattcgggtga	17820
gcaatggata	ggggagttga	tatcgtcaac	gttcacttct	aaagaaatag	cgccactcag	17880
cttctcagc	ggctttatcc	agcgatttcc	tattatgtcg	gcattagtct	caagatcgac	17940
agcctgtcac	gggttaagcga	gaaatgaata	agaaggctga	taattcggat	ctctgcgagg	18000
gagatgatat	ttgatcacag	gcagcaacgc	tctgtcatcg	ttacaatcaa	catgctaccc	18060
tcgcgcagat	catccgtgtt	tcaaaccccg	cagcttagtt	gcggttcttc	cgaatagcat	18120
cggtaacatg	acgaagctct	gcgcgcttac	aacggctctc	ccgctgacgc	cgtccgggac	18180
tgatgggctg	cctgtatcga	gtgggtgatt	tgtgcgcagc	tgccggtcgg	ggagctggtg	18240
ctgggctggg	ggcaggatat	attgtgggtg	aaacaaattg	acgcttagac	aacttaataa	18300
cacattgcgg	acgtttttta	tgtactgggg	tggtttttct	tttcaccagt	gagacgggca	18360
acagctgatt	gcccttcacc	gcctggccct	gagagaggtg	cagcaagcgg	tcacagctgg	18420
tttggccag	caggcgaaaa	tctgttttga	tgggtggttc	gaaatcggca	aaatccctta	18480
taaatcaaaa	gaatagcccg	agatagggtt	gagtggtgtt	ccagtttggg	acaagagtc	18540
actattaaag	aacgtggact	ccaacgtcaa	agggcgaaaa	accgtctatc	agggcgatgg	18600
cccactacgt	gaaccatcac	ccaaatcaag	ttttttgggg	tcgaggtgcc	gtaaaagcact	18660
aaatcggaac	cctaaaggga	gccccgatt	tagagcttga	cggggaaagg	cgccgaacgt	18720
ggcgagaaa	gaaggggaaga	aagcgaaagg	agcggggcgc	attcaggctg	cgcaactggt	18780
gggaaggcg	atcggtgcgg	gcctcttcgc	tattacgcca	gctggcgaaa	gggggatgtg	18840
ctgcaaggcg	attaagttgg	gtaacgccag	ggttttccca	gtcacgacgt	tgtaaaaacga	18900
cgccagtgga	attcgagctc	ggtacccccc	tactccaaaa	atgtcaaaaga	tacagtctca	18960
gaagacccaa	gggctattga	gactttttcaa	caaagggtaa	tttcgggaaa	cctcctcgga	19020
ttccattgcc	cagctatctg	tcacttcac	gaaaggacag	tagaaaaagg	aggtggctcc	19080
tacaaatgcc	atcattgcga	taaggaaaag	gctatcattc	aagatgcctc	tgccgacagt	19140
ggtcccaagg	atggaccccc	accacagagg	agcatcgtgg	aaaaagaaga	cgtcccaacc	19200
acgtcttcaa	agcaagtgga	ttgatgtgac	atctccactg	acgtaaggga	tgacgcacaa	19260
ttccactatc	ccttcgaaga	ccctcttct	atataaggaa	gttcatttca	ttggagagg	19320
acagcccaag	ctttctagag	gatccataaa	acatttcaat	cccttgaacg	cggtagaacg	19380
tgctaattgg	attttgttga	gaacgcggta	gaacgtactt	atcacctaca	gttttatttt	19440
gtttttcttt	ttggtttaat	ctatccagct	tagtaccgag	tgggggaaag	tgactgggtg	19500
gcctaaaaac	tttcttttga	tactttgtaa	aaatacatac	agatacaatg	gcgaacggta	19560
acttcaagtt	gtctcaattg	ctcaatgtgg	acgagatgtc	tgctgagcag	aggagtcatt	19620

-continued

```

tctttgactt gatgctgact aaacctgatt gtgagatcgg gcaaatgatg caaagagttg 19680
ttgttgataa agtcgatgac atgattagag aaagaaagac taaagatcca gtgattgttc 19740
atgaagtctt ttctcagaag gaacagaaca agttgatgga aatttatcct gaattcaata 19800
tcgtgtttaa agacgacaaa aacatgggtc atgggtttgc ggctgctgag cgaaaactac 19860
aagctttatt gcttttagat agagttcctg ctctgcaaga ggtggatgac atcggtgggtc 19920
aatggtcggt ttgggttaact agaggtgaga aaaggattca ttctgtgtg ccaaatctag 19980
atattcggga tgatcagaga gaaatttctc gacagatatt tcttactgct attggtgac 20040
aagctagaag tggtaagaga cagatgtcgg agaatgagct gtggatgtat gaccaatttc 20100
gtgaaaatat tgctgcgcct aacgcgggta ggtgcaataa tacatatcag ggttgtacat 20160
gtagggggtt t 20171

```

```

SEQ ID NO: 16      moltype = DNA length = 300
FEATURE
source             Location/Qualifiers
                   1..300
                   mol_type = other DNA
                   organism = synthetic construct

```

```

SEQUENCE: 16
tgtgatattc aaccaaatga attgttaatt gatcaatcag cccatttcat atcattgggt 60
ccctttgctg caacttttgc ttttgagatt tggataatc ctcgtgatca ctctctcat 120
tttcatgaac tcgataatga aaaggcaaat gatcttgagg ggttggtgaa actcatgctt 180
ttgaagatgt ctttgcagtt gaacaatcca ccatccaacc tccttattca tacctcccca 240
tttcaagtag atccctcata tgctccttcc acacactggt ttttacaat agctcccat 300

```

1. A method of modulating (e.g. decreasing) the alkaloid content of a tobacco plant or a part thereof or tobacco plant cell, the method comprising modifying said plant or plant cell by modulating (e.g. decreasing) the activity or expression of Galactose-1-phosphate uridyl transferase.

2. A method of modulating (e.g. decreasing) the content of a tobacco specific nitrosamine (TSNA) or a precursor of a TSNA in a tobacco plant or plant part thereof or tobacco plant cell, the method comprising modifying said plant or plant cell by modulating (e.g. decreasing) the activity or expression of Galactose-1-phosphate uridyl transferase.

3. A method for producing a plant or part thereof, a cell or cell culture, a plant propagation material, a leaf, a cut harvested leaf, a processed leaf or a cut and processed leaf which has modulated (e.g. decreased) alkaloid content, the method comprising modifying said plant or cell culture to modulate the activity or expression of Galactose-1-phosphate uridyl transferase.

4. Use of at least one gene encoding a Galactose-1-phosphate uridyl transferase for modulating alkaloid content of a tobacco cell or tobacco plant or part thereof.

5. A method or use according to any preceding claim, wherein the Galactose-1-phosphate uridyl transferase:

- a) comprises an amino acid sequence as set out in SEQ ID No. 3; or a functional variant or functional fragment or orthologue of SEQ ID No. 3; or a sequence which has at least 80% identity to SEQ ID No. 3; or a homologue of SEQ ID No. 3; or
- b) is encoded by a nucleotide sequence as set out in SEQ ID No. 1 or 2; or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2; or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2.

6. A method or use according to any one of the preceding claims, wherein the alkaloid content is modulated (e.g. decreased) in comparison to a plant or cell culture which has not been modified to modulate the activity or expression of Galactose-1-phosphate uridyl transferase.

7. A tobacco plant or part thereof or a tobacco cell or cell culture which has been modified to modulate (e.g. decrease) the activity or expression of Galactose-1-phosphate uridyl

transferase and the tobacco plant or part thereof or tobacco cell or cell culture has decreased alkaloid and or TSNA precursor content in comparison to an unmodified plant or unmodified cell or cell culture.

8. A plant propagation material obtainable (e.g. obtained) from a plant according to claim 7 or from a plant or cell or cell culture produced by the method or use of any one of claims 1 to 6.

9. A method or use according to any one of claims 1-6, or a plant or part thereof or cell or cell culture according to claim 7, or a plant propagation material according to claim 8, wherein the alkaloid content of the plant is decreased in comparison to a plant or cell culture which has not been modified to modulate the activity or expression of Galactose-1-phosphate uridyl transferase.

10. A method or use according to any one of claims 1-6 or 9, a tobacco plant or part thereof or tobacco cell or cell culture according to claim 7 or 9, or a plant propagation material according to claim 8, wherein the content of one or more alkaloids selected from nicotine, nornicotine, PON, anabasine, myosmine and anatabine is modulated (e.g. decreased), preferably the content of nicotine, nornicotine and/or PON is modulated (e.g. decreased).

11. A method or use according to claim 10, a tobacco plant or part thereof or tobacco cell or tobacco cell culture according to claim 10, or a plant propagation material according to claim 10 wherein the nicotine content is decreased.

12. Use of a tobacco plant or part thereof or tobacco cell or cell culture according to any one of claims 7 or 9-11, or of a plant produced by the method of any one of claims 1 to 6, or 9 to 11 to breed a plant.

13. Use of a tobacco plant or part thereof or a tobacco cell or cell culture according to any one of claims 7 or 9-11, or of a plant produced by the method of any one of claims 1 to 6, or 9 to 11 for production of a product.

14. Use of a tobacco plant or part thereof according to any one of claims 7 or 9-11, or of a plant produced by the method of any one of claims 1 to 6, or 9 to 11 to grow a crop.

15. Use of a tobacco plant or part thereof according to any one of claims 7 or 9-11, or of a plant produced by the method of any one of claims 1 to 6, or 9 to 11 to produce a leaf.

16. A harvested leaf of a plant according to any one of claims **7** or **9-11**, or obtainable from a plant propagated from a propagation material according to any one of claims **8-11**, or obtainable from a plant obtained by a use according to any one of claims **4-6** or **9-11**, or obtainable from a plant produced by the method of any one of claims **1-6**, or **9-11**.

17. A harvested leaf of a plant according to claim **16**, wherein the harvested leaf of a plant is a cut harvested leaf.

18. A processed leaf, preferably a processed tobacco leaf, preferably a non-viable processed tobacco leaf:

obtainable (e.g. obtained) from a plant obtainable from a use according to any one of claims **4-6** or **9-11**;

obtainable (e.g. obtained) by processing a plant according to any one of claims **7** or **9-11**;

obtainable (e.g. obtained) from a plant propagated from a plant propagation material according to any one of claims **8-11**; or

obtainable (e.g. obtained) by processing a harvested leaf of a plant according to claim **16** or **17**; or

obtainable (e.g. obtained) from a plant produced by the method of any one of claims **1** to **6**, or **9** to **11**.

19. A processed leaf according to claim **18**, wherein the leaf is processed by curing, fermenting, pasteurising or a combination thereof.

20. A processed leaf according to claim **18** or **19**, wherein the processed leaf is a cut processed leaf.

21. Cured tobacco material made from a plant or a part thereof according to any one of claims **7** or **9-11**, or a harvested leaf according to claim **16** or **17**, or a processed leaf according to any of claims **18-20**.

22. A tobacco blend comprising said cured tobacco material of claim **21**.

23. A tobacco industry product prepared from:

a tobacco plant or part thereof or tobacco cell or cell culture according to any one of claims **7** or **9-11**;

a tobacco plant or part thereof propagated from a tobacco plant propagation material according to claim **8**;

a harvested leaf of a plant according to claim **16** or **17**;

a processed leaf according to any one of claims **18-20**.

24. A tobacco industry product according to claim **23**, wherein the tobacco product is a combustible smoking article.

25. A tobacco industry product according to claim **23**, wherein the tobacco product is a smokeless tobacco product.

26. A tobacco product according to claim **23**, wherein the tobacco product is a non-combustible aerosol provision system such as a tobacco heating device or an aerosol-generating device.

27. A combustible smoking article, non-combustible aerosol provisioning system, smokeless tobacco product or tobacco heating device comprising a plant or a part thereof according to any one of claims **7** or **9-11** or an extract (e.g. a tobacco extract) thereof or a tobacco cell culture according to any one of claims **7** or **9-11**; or a cured tobacco material according to claim **21**; or a tobacco blend according to claim **22**.

28. Use of a nucleotide sequence encoding Galactose-1-phosphate uridyl transferase which:

a) encodes an amino acid sequence as set out in SEQ ID No. 3; or a functional variant or functional fragment or orthologue of SEQ ID No. 3; or a sequence which has at least 80% identity to SEQ ID No. 3; or a homologue of SEQ ID No. 3; or

b) comprises a sequence as set out in SEQ ID No. 1 or 2; or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2; or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2;

to select a plant having modulated (e.g. reduced) alkaloid content and/or modulated (e.g. reduced) content of TSNA or a precursor of a TSNA.

29. A mutant of a plant carrying a heritable mutation in a nucleotide sequence which:

a) encodes an amino acid sequence as set out in SEQ ID No. 3; or a functional variant or functional fragment or orthologue of SEQ ID No. 3; or a sequence which has at least 80% identity to SEQ ID No. 3; or a homologue of SEQ ID No. 3; or

b) comprises a sequence as set out in SEQ ID No. 1 or 2; or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2; or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2;

wherein said heritable mutation modulates (e.g. decreases) the activity or expression of Galactose-1-phosphate uridyl transferase and wherein the mutant plant has modulated (e.g. decreased) alkaloid content and/or modulated content of a tobacco specific nitrosamine (TSNA) or a precursor of a TSNA relative to a comparable plant which does not carry said heritable mutation.

30. Progeny or seed of a mutant plant which carries the heritable mutation according to claim **29**.

31. A harvested leaf, a processed leaf or cured tobacco material produced from a plant comprising a modification in a nucleotide sequence which:

a) encodes an amino acid sequence as set out in SEQ ID No. 3; or a functional variant or functional fragment or orthologue of SEQ ID No. 3; or a sequence which has at least 80% identity to SEQ ID No. 3; or a homologue of SEQ ID No. 3; or

b) comprises a sequence as set out in SEQ ID No. 1 or 2; or a functional variant or functional fragment or orthologue of SEQ ID No. 1 or 2; or a nucleic acid sequence which has at least 80% identity to SEQ ID No. 1 or 2; or a homologue of SEQ ID No. 1 or 2;

wherein said modification modulates (e.g. decreases) the activity or expression of Galactose-1-phosphate uridyl transferase and wherein said plant has modulated (e.g. decreased) alkaloid content and/or modulated content of a tobacco specific nitrosamine (TSNA) or a precursor of a TSNA relative to a comparable plant which does not carry said modification in said Galactose-1-phosphate uridyl transferase.

* * * * *