

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent Application Publication

20250276063

Kind Code

A1

Publication Date

September 04, 2025

Inventor(s)

van der TOUW; William et al.

PLACENTA-DERIVED NK CELLS AS A SENOLYTIC FOR THERAPEUTIC AND OTHER USES

Abstract

Provided herein are methods of killing senescent cells, e.g., killing senescent cells in a human subject. Also provided herein are methods of treating a disease or disorder associated with cellular senescence in a subject in need thereof comprising administering to the subject an effective amount of placenta-derived NK cells to the subject. The present invention also provides compositions comprising NK, e.g., CYNK cells, for killing senescent cells and for the treatment of a disease or disorder associated with cellular senescence.

Inventors: van der TOUW; William (Florham Park, NJ), HARIRI; Robert J (Florham Park, NJ), MAHLAKÖIV; Tanel (Florham Park, NJ)

Applicant: Celularity Inc. (Florham Park, NJ)

Family ID: 83189022

Assignee: Celularity Inc. (Florham Park, NJ)

Appl. No.: 18/293243

Filed (or PCT Filed): July 29, 2022

PCT No.: PCT/US2022/074326

Related U.S. Application Data

us-provisional-application US 63227328 20210729

Publication Classification

Int. Cl.: A61K40/15 (20250101); A61K35/17 (20250101); A61K40/42 (20250101); A61P35/00 (20060101); C12N5/0783 (20100101)

U.S. Cl.:

CPC A61K40/15 (20250101); A61K35/17 (20130101); A61K40/4224 (20250101); A61P35/00 (20180101); C12N5/0646 (20130101);

Background/Summary

[0001] This application claims priority to U.S. Provisional Patent Application No. 63/227,328, filed Jul. 29, 2021, the disclosure of which is incorporated herein by reference in its entirety.

1. FIELD

[0002] Provided herein are methods of producing populations of natural killer (NK) cells and/or ILC3 cells from a population of hematopoietic stem or progenitor cells in media comprising stem cell mobilizing factors, e.g., three-stage methods of producing NK cells and/or ILC3 cells in media comprising stem cell mobilizing factors starting with hematopoietic stem or progenitor cells from cells of the placenta, for example, from placental perfusate (e.g., human placental perfusate) or other tissues, for example, umbilical cord blood or peripheral blood. Further provided herein are methods of using the placental perfusate, the NK cells and/or ILC3 cells and/or NK progenitor cells described herein, to, e.g., kill or suppress the proliferation of senescent cells.

2. BACKGROUND

[0003] Natural killer (NK) cells play a key role in immune surveillance against stressed cell states including infection, tumor transformation, and cell senescence. Using a broad repertoire of surface receptors, NK cells sense the environment to differentiate between normal and malignant cells (cancerous or infected). Broadly, the two major classes of signals triggering specific cytotoxicity of infected cells by NK cells are: 1) downregulated MHC I expression, and 2) induced expression of NK cell receptor ligands. The molecular ligands for NK cell activating receptors are endogenous—induced by infection and consequent cell stress, or exogenous—molecules expressed by the pathogen. In the case of cell senescence, ageing is associated with the accumulation of senescent cells which secrete inflammatory signals and mediate a senescent-associated phenotype. NK cells are able to recognize senescent cells specifically through the recognition of stress antigens like MICA/B (Krizhanovsky, V. et al. Cell 2008)(Sagiv A. Aging. 2016). Senescent cells have been shown to escape NK cell targeted killing by specifically overexpressing MHC-I inhibitory ligand HLA-E to counteract the presentation of stress ligands (Pereira B I et al. Nat. Comm. 2019). We present the feasibility of using exogenous NK cells for the elimination of senescent cells in order to reverse the deleterious effects of senescence-associated ageing. We also explore the feasibility of genetically knocking-out the expression of NKG2A so that manufactured cells are not sensitive to HLA-E driven escape of senescent cells by tumor targeting.

[0004] Senescence is associated with physiological decline of an organism and with a range of age-related diseases. Cellular senescence occurs when cells reach their replicative capacity and irreversible growth arrest, resulting in accumulation of dysfunctional senescent cells, decreased tissue repair capacity and a failure to maintain tissue homeostasis. Increased cell senescence in an aging organism is related to increased inflammation and tissue fibrosis (ref: PMID: 23140366). It has been demonstrated that reduction of senescent cells in an organism can improve organ function and healthy lifespan (ref: PMID: 32555459, PMID: 22048312, PMID: 26840489).

[0005] Senescent cells are characterized by a range of phenotypical features, including the expression of senescence-associated molecules on the cell surface (ref: PMID: 34359910). Ligands

of an activating natural killer (NK) cell receptor have been shown to mediate immunosurveillance of senescent cells (ref: PMID: 26878797, PMID: 31160572, PMID: 30575733).

[0006] NK cells are innate lymphoid cells that play a key role in immune surveillance against aberrant cells. NK cells circulate the body probing the environment using an array of activating and inhibitory cell surface receptors that distinguish between healthy and stressed cells. NK cell activating receptors, including natural cytotoxicity receptors (NCRs) NKp30, NKp44, NKp46 and NKG2D, recognize cell surface stress ligands that are upregulated on transformed, senescent or infected cells (PMID: 18425107). Upon target cell recognition, NK cells release cytotoxic molecules perforins and granzymes to kill target cells (ref: PMID: 17124515, PMID: 30575733). In addition to direct elimination of target (senescent cells), NK cells play a role in reducing tissue fibrosis and promote tissue repair (ref: PMID: 16515819, PMID: 16472598, PMID: 28373874).

[0007] Celularity Inc. has developed CYNK-001, an off-the-shelf allogeneic NK cell population derived from human placental hematopoietic stem cells, that efficiently kills various tumor cell lines in vitro and in vivo (ref: PMID: 33741730, PMID: 23641243). While adoptive NK cell therapies, including CYNK-001, are being investigated in the clinical setting for the treatment of hematological and solid tumors, NK cell therapies and their potential benefit in aging has not been investigated.

3. SUMMARY

[0008] The present invention provides methods of killing senescent cells comprising contacting the senescent cells with placenta-derived natural killer (NK) cells.

[0009] The present invention provides methods of treating a disease or disorder associated with cellular senescence in a subject in need thereof comprising administering to the subject an effective amount of placenta-derived NK cells to the subject so as thereby to provide an effective treatment of the disease or disorder in the subject.

[0010] The present invention provides compositions comprising placenta-derived NK cells for use in killing senescent cells in a subject.

[0011] The present invention provides uses of a composition comprising placenta-derived NK cells for the manufacture of a medicament for killing senescent cells in a subject.

[0012] The present invention provides compositions comprising placenta-derived NK cells for use treating a disease or disorder associated with cellular senescence in a subject. The present invention provides uses of a composition comprising placenta-derived NK cells for use treating a disease or disorder associated with cellular senescence in a subject.

[0013] In some embodiments of the invention the placenta-derived NK cells comprise at least 20% CD56+CD3⁻ natural killer cells.

[0014] In some embodiments of the invention the placenta-derived NK cells comprise at least 40% CD56+CD3⁻ natural killer cells.

[0015] In some embodiments of the invention the placenta-derived NK cells comprise at least 60% CD56+CD3⁻ natural killer cells.

[0016] In some embodiments of the invention the placenta-derived NK cells comprise at least 80% CD56+CD3⁻ natural killer cells.

[0017] In some embodiments of the invention said placenta derived natural killer cells are human placenta derived natural killer cells.

[0018] In some embodiments of the invention said placenta derived natural killer cells are hematopoietic stem cell-derived natural killer cells.

[0019] In some embodiments of the invention said placenta derived natural killer cells are CD34⁺ hematopoietic stem cell-derived natural killer cells.

[0020] In some embodiments of the invention said placenta derived natural killer cells are CYNK cells.

[0021] The method of any one of claims **1-10**, wherein the NK cells are characterized by expression of one or more markers selected from the group consisting of FGFBP2, GZMH,

CCL3L3, GZMM, CXCR4, ZEB2, KLF2, LITAF, RORA, LYAR, CNOT1, IFNG, DUSP2, ATG2A, CD7, PMAIP1, PPP2R5C, NR4A2, ZFP36L2, PIK3R1, KLRF1, SNHG9, MT2A, RGS2, CHD1, DUSP1, EML4, ZFP36, ZC3H12A, DNAJB6, SBDS, IRF1, TSC22D3, TSPYL2, PNRC1, ISCA1, JUNB, WHAMM, RICTOR, TNFAIP3, EPC1, MVD, CLK1, ARL4C, REL, KMT2E, YPEL5, AMD1, BTG2, and IDS which is lower than expression of said markers in peripheral blood natural killer cells and/or expression of one or more markers selected from the group consisting of NDFIP2, LINC00996, MAL, CCL1, MB, SPINK2, C15orf48, CAMK1, KLRC1, TNFSF10, TNFRSF18, IL32, CAPG, AC092580.4, S100A11, TNFRSF4, ENO1, FCER1G, CCND2, KRT81, MRPS6, ANXA2, PTGER2, GLO1, HAVCR2, PYCARD, LAT2, SLC16A3, COTL1, PKM, TALDO1, CD96, NCR3, KRT86, STMN1, LTB, ARPC1B, ARPC5, FKBP1A, TIMP1, GZMK, CD59, PGK1, RGS10, EVL, RAC2, LGALS1, ITGB7, TUBB, PGAM1, PRF1, GZMB, IL2RB, KLRC2, and KLRB1 which is higher than expression of said markers in peripheral blood natural killer cells.

[0022] In some embodiments of the invention the NK cells are characterized by expression of one or more markers selected from the group consisting of FGFBP2, GZMH, CCL3L3, GZMM, CXCR4, ZEB2, KLF2, LITAF, RORA, LYAR, CNOT1, IFNG, DUSP2, ATG2A, CD7, PMAIP1, PPP2R5C, NR4A2, ZFP36L2, PIK3R1, KLRF1, SNHG9, MT2A, RGS2, CHD1, DUSP1, EML4, ZFP36, ZC3H12A, DNAJB6, SBDS, IRF1, TSC22D3, TSPYL2, PNRC1, ISCA1, JUNB, WHAMM, RICTOR, TNFAIP3, EPC1, MVD, CLK1, ARL4C, REL, KMT2E, YPEL5, AMD1, BTG2, and IDS which is lower than expression of said markers in peripheral blood natural killer cells.

[0023] In some embodiments of the invention expression of 2, 3, 4, 5, 6, 7, 8, 9, 10, or more markers selected from the group consisting of FGFBP2, GZMH, CCL3L3, GZMM, CXCR4, ZEB2, KLF2, LITAF, RORA, LYAR, CNOT1, IFNG, DUSP2, ATG2A, CD7, PMAIP1, PPP2R5C, NR4A2, ZFP36L2, PIK3R1, KLRF1, SNHG9, MT2A, RGS2, CHD1, DUSP1, EML4, ZFP36, ZC3H12A, DNAJB6, SBDS, IRF1, TSC22D3, TSPYL2, PNRC1, ISCA1, JUNB, WHAMM, RICTOR, TNFAIP3, EPC1, MVD, CLK1, ARL4C, REL, KMT2E, YPEL5, AMD1, BTG2, and IDS is lower than expression of said markers in peripheral blood natural killer cells.

[0024] In some embodiments of the invention the NK cells are characterized by expression of one or more markers selected from the group consisting of NDFIP2, LINC00996, MAL, CCL1, MB, SPINK2, C15orf48, CAMK1, KLRC1, TNFSF10, TNFRSF18, IL32, CAPG, AC092580.4, S100A11, TNFRSF4, ENO1, FCER1G, CCND2, KRT81, MRPS6, ANXA2, PTGER2, GLO1, HAVCR2, PYCARD, LAT2, SLC16A3, COTL1, PKM, TALDO1, CD96, NCR3, KRT86, STMN1, LTB, ARPC1B, ARPC5, FKBP1A, TIMP1, GZMK, CD59, PGK1, RGS10, EVL, RAC2, LGALS1, ITGB7, TUBB, PGAM1, PRF1, GZMB, IL2RB, KLRC2, and KLRB1 which is higher than expression of said markers in peripheral blood natural killer cells.

[0025] In some embodiments of the invention expression of 2, 3, 4, 5, 6, 7, 8, 9, 10, or more markers selected from the group consisting of NDFIP2, LINC00996, MAL, CCL1, MB, SPINK2, C15orf48, CAMK1, KLRC1, TNFSF10, TNFRSF18, IL32, CAPG, AC092580.4, S100A11, TNFRSF4, ENO1, FCER1G, CCND2, KRT81, MRPS6, ANXA2, PTGER2, GLO1, HAVCR2, PYCARD, LAT2, SLC16A3, COTL1, PKM, TALDO1, CD96, NCR3, KRT86, STMN1, LTB, ARPC1B, ARPC5, FKBP1A, TIMP1, GZMK, CD59, PGK1, RGS10, EVL, RAC2, LGALS1, ITGB7, TUBB, PGAM1, PRF1, GZMB, IL2RB, KLRC2, and KLRB1 is higher than expression of said markers in peripheral blood natural killer cells.

[0026] In some embodiments of the invention the NK cells are prepared by the methods presented herein.

[0027] In some embodiments of the invention the disease or disorder is a fibrotic disease or disorder.

[0028] In some embodiments of the invention the disease or disorder is an inflammatory disease or disorder.

[0029] In some embodiments of the invention the disease or disorder is a metabolic disease or disorder.

[0030] In some embodiments of the invention the disease or disorder is a liver disease or disorder.

[0031] In some embodiments of the invention the disease or disorder is a kidney disease or disorder.

[0032] In some embodiments of the invention the disease or disorder is a lung disease or disorder.

[0033] In some embodiments of the invention the disease or disorder is a eye disease or disorder.

[0034] In some embodiments of the invention the disease or disorder is a skeletal disease or disorder.

[0035] In some embodiments of the invention the disease or disorder is a skin disease or disorder.

[0036] In some embodiments of the invention the disease or disorder is a gastrointestinal disease or disorder.

[0037] In some embodiments of the invention the disease or disorder is a bone marrow disease or disorder.

[0038] In some embodiments of the invention the disease or disorder is a cardiovascular disease or disorder.

[0039] In some embodiments of the invention the disease or disorder is a circulatory disease or disorder.

[0040] In some embodiments of the invention the disease or disorder is a neuronal disease or disorder.

[0041] In some embodiments of the invention the disease or disorder is an age-related disease or disorder.

[0042] In some embodiments of the invention the disease or disorder is osteoarthritis.

[0043] In some embodiments of the invention the disease or disorder is intervertebral disc degeneration.

[0044] In some embodiments of the invention the disease or disorder is atherosclerosis.

[0045] In some embodiments of the invention the disease or disorder is hardening of the arteries.

[0046] In some embodiments of the invention the disease or disorder is heart disease.

[0047] In some embodiments of the invention the disease or disorder is neurodegeneration.

[0048] In some embodiments of the invention the disease or disorder is neuroinflammation.

[0049] In some embodiments of the invention the disease or disorder is Alzheimer's disease.

[0050] In some embodiments of the invention the disease or disorder is myeloproliferative neoplasm (MPN).

[0051] In some embodiments of the invention the disease or disorder is myelodysplastic syndrome (MDS).

[0052] In some embodiments of the invention the disease or disorder is osteoporosis.

[0053] In some embodiments of the invention the disease or disorder is age-induced frailty.

[0054] In some embodiments of the invention the effective treatment comprises a reduction in fibrosis.

[0055] In some embodiments of the invention the effective treatment comprises a reduction in inflammation.

[0056] In some embodiments of the invention the effective treatment comprises a reduction in senescent cell numbers.

[0057] In some embodiments of the invention the effective treatment comprises a reduction in relative senescent cell numbers.

[0058] In some embodiments of the invention the effective treatment comprises an increase in organ or tissue function.

[0059] In some embodiments of the invention the senescent cell is selected from the group consisting of a senescent fibroblast, a senescent pre-adipocyte, a senescent epithelial cell, a senescent chondrocyte, a senescent intervertebral disc cell, a senescent satellite cell, a senescent

muscle cell, a senescent liver cell, a senescent kidney cell, a senescent liver cell, a senescent kidney cell, a senescent bone marrow cell, a senescent vascular cell, a senescent immune cell, a senescent heart cell, a senescent arterial cell, a senescent venous cell, a senescent gastrointestinal cell, a senescent neuron, and a senescent endothelial cell.

[0060] In some embodiments of the invention the CYNK cells are prepared by the methods presented herein.

Terminology

[0061] As used herein, the term CYNK are CD34⁺ cell-derived NK cells produced by the methods described herein. In specific embodiments, CYNK cells are placental-derived NK cells. In other specific embodiments, CYNK-001 is a specific formulation of CYNK cells.

[0062] As used herein, the terms “immunomodulatory compound” and “IMiDTM” do not encompass thalidomide.

[0063] As used herein, “lenalidomide” means 3-(4'-aminoisoindoline-1'-one)-1-piperidine-2,6-dione (Chemical Abstracts Service name) or 2,6-Piperidinedione,3-(4-amino-1,3-dihydro-1-oxo-2H-isoindol-2-yl)-(International Union of Pure and Applied Chemistry (IUPAC) name). As used herein, “pomalidomide” means 4-amino-2-(2,6-dioxopiperidin-3-yl)isoindole-1,3-dione.

[0064] As used herein, “multipotent,” when referring to a cell, means that the cell has the capacity to differentiate into a cell of another cell type. In certain embodiments, “a multipotent cell” is a cell that has the capacity to grow into a subset of the mammalian body's approximately 260 cell types. Unlike a pluripotent cell, a multipotent cell does not have the capacity to form all of the cell types.

[0065] As used herein, “feeder cells” refers to cells of one type that are co-cultured with cells of a second type, to provide an environment in which the cells of the second type can be maintained, and perhaps proliferate. Without being bound by any theory, feeder cells can provide, for example, peptides, polypeptides, electrical signals, organic molecules (e.g., steroids), nucleic acid molecules, growth factors (e.g., bFGF), other factors (e.g., cytokines), and metabolic nutrients to target cells. In certain embodiments, feeder cells grow in a mono-layer.

[0066] As used herein, the “natural killer cells” or “NK cells” produced using the methods described herein, without further modification, include natural killer cells from any tissue source.

[0067] As used herein, the “ILC3 cells” produced using the methods described herein, without further modification, include ILC3 cells from any tissue source.

[0068] As used herein, “placental perfusate” means perfusion solution that has been passed through at least part of a placenta, e.g., a human placenta, e.g., through the placental vasculature, and includes a plurality of cells collected by the perfusion solution during passage through the placenta.

[0069] As used herein, “placental perfusate cells” means nucleated cells, e.g., total nucleated cells, isolated from, or isolatable from, placental perfusate.

[0070] As used herein, “tumor cell suppression,” “suppression of tumor cell proliferation,” and the like, includes slowing the growth of a population of tumor cells, e.g., by killing one or more of the tumor cells in said population of tumor cells, for example, by contacting or bringing, e.g., NK cells or an NK cell population produced using a three-stage method described herein into proximity with the population of tumor cells, e.g., contacting the population of tumor cells with NK cells or an NK cell population produced using a three-stage method described herein. In certain embodiments, said contacting takes place in vitro or ex vivo. In other embodiments, said contacting takes place in vivo.

[0071] As used herein, the term “hematopoietic cells” includes hematopoietic stem cells and hematopoietic progenitor cells.

[0072] As used herein, the “undefined component” is a term of art in the culture medium field that refers to components whose constituents are not generally provided or quantified. Examples of an “undefined component” include, without limitation, serum, for example, human serum (e.g., human serum AB) and fetal serum (e.g., fetal bovine serum or fetal calf serum).

[0073] As used herein, “+”, when used to indicate the presence of a particular cellular marker,

means that the cellular marker is detectably present in fluorescence activated cell sorting over an isotype control; or is detectable above background in quantitative or semi-quantitative RT-PCR. [0074] As used herein, “–”, when used to indicate the presence of a particular cellular marker, means that the cellular marker is not detectably present in fluorescence activated cell sorting over an isotype control; or is not detectable above background in quantitative or semi-quantitative RT-PCR.

Description

4. BRIEF DESCRIPTION OF THE FIGURES

[0075] FIG. 1 shows expansion of NK cells for compounds CRL1-CRL11.

[0076] FIG. 2 shows expansion of NK cells for compounds CRL12-CRL22.

[0077] FIG. 3 shows expansion of NK cells relative to SR1 positive control.

[0078] FIG. 4 shows expansion of CD34+ cells from which the NK cells were derived.

[0079] FIG. 5 shows cytotoxicity of the expanded NK cultures.

[0080] FIGS. 6A-6C show that PNK cells highly express genes encoding the cytotoxic machinery. FIG. 6A CYNK cells were combined with peripheral blood derived NK cells (PB-NK) at 1:1 ratio and gene expression analyzed on single cell level using 10× Genomics Chromium platform and Illumina sequencing. Bioinformatics analysis utilized 10× Genomics Cell Ranger analysis pipeline. Transcript analysis was restricted to Granzyme B (GZMB) expressing cells. FIG. 6B A representative tSNE plot depicting PNK and PB-NK cells as distinct populations. FIG. 6C tSNE plots of selected NK cell-associated genes. The data is representative of two donors.

[0081] FIG. 7 shows that PNK and PB-NK cells differentially express genes encoding NK cell receptors. The expression of selected NK cell receptor genes analyzed by real-time quantitative PCR in peripheral blood NK cells (PB-NK) and CD11a+-bead-purified PNK cells. An alternative name indicated above the histogram for selected markers. The data represents mean±SD of three donors for CYNK and PBNK cells (n=3). *p<0.05, **p<0.005, ***p<0.001.

[0082] FIG. 8 shows the gating strategy for PB-NK and CYNK cells. CYNK and PBMC cells were thawed and stained with fluorophore-coupled antibodies targeting NK cell receptors. The figure demonstrates representative dot plots and the gating strategy for the identification of CYNK and PB-NK cells. See FIG. 9 for further characterization of the populations.

[0083] FIG. 9 shows differential expression of surface proteins on CYNK and PB-NK cells. CYNK and PB-NK cells were pre-gated as indicated in FIG. 8.

[0084] FIG. 10 shows that CYNK cells form a distinct cell population from PB-NK cells based on surface protein expression. tSNE plots demonstrating differential clustering of CYNK and PB-NK cells based on their surface markers. tSNE plots were generated of flow cytometry data using FlowJo software.

[0085] FIGS. 11A-11C show that CYNK-001 cells express NK cell activating receptors involved in senescent cell recognition. FIG. 11A: Representative flow cytometry dot plots demonstrating the gating strategy for the analysis of CYNK-001 cells. Thawed CYNK-001 cells were stained with fluorophore-conjugated antibodies recognizing indicated NK cell markers and analyzed by flow cytometry. Lineage included CD3, CD14 and CD19. CYNK-001 are defined as live CD3– CD14– CD19– CD56+ cells. FIG. 11B: Proportion of indicated marker expression on CYNK-001 cells. FIG. 11C: Representative histograms of the expression of indicated NK activating cell receptors on CYNK cells. mean±SD, n=6 (CYNK-001 developed from 6 donors), FMO—fluorescence minus one.

[0086] FIG. 12 shows that tumor cell lines express NKG2D ligands. Representative histograms demonstrating the binding of Fc-coupled recombinant NKG2D (NKG2D-Fc) or antibodies against major NKG2D ligands ULBP2/5/6 and MICA/B to tumor cell lines. Indicated tumor cells were

cultured for 24 h on tissue culture plates and trypsinized using TrypLE reagent for antibody staining and flow cytometric analysis. Dotted vertical line indicates the binding of the fluorophore-coupled IgG control antibody.

[0087] FIG. 13 shows that NKG2D mediates recognition and killing of target cells by CYNK-001. Indicated tumor cell lines were seeded on 96-well format RTCA E-plates for the evaluation of cytolysis by CYNK-001 using an impedance-based RTCA platform. After 24 h of tumor cell expansion, NKG2D blocking antibody (aNKG2D) or a non-specific antibody control (IgG) were added to tumor cells at a final concentration of 10 μ g/ml together with CYNK-001 cells. CYNK-001 were added at an effector-to-target ration of 10:1 for U251, 2.5:1 for LN229, 1.25:1 for DLD-1, 5:1 for HCT116 and 2.5:1 for HT29. Cytolysis of target cells was evaluated over indicated time (h). mean $\pm$ SD, n=2 (technical replicates).

5. DETAILED DESCRIPTION

[0088] Provided herein are novel methods of producing and expanding NK cells and/or ILC3 cells from hematopoietic cells, e.g., hematopoietic stem cells or progenitor cells. Also provided herein are methods, e.g., three-stage methods, of producing NK cell populations and/or ILC3 cell populations from hematopoietic cells, e.g., hematopoietic stem cells or progenitor cells. The hematopoietic cells (e.g., CD34+ hematopoietic stem cells) used to produce the NK cells and/or ILC3 cells, and NK cell populations and/or ILC3 cell populations, may be obtained from any source, for example, without limitation, placenta, umbilical cord blood, placental blood, peripheral blood, spleen or liver. In certain embodiments, the NK cells and/or ILC3 cells or NK cell populations and/or ILC3 cell populations are produced from expanded hematopoietic cells, e.g., hematopoietic stem cells and/or hematopoietic progenitor cells. In one embodiment, hematopoietic cells are collected from a source of such cells, e.g., placenta, for example from placental perfusate, umbilical cord blood, placental blood, peripheral blood, spleen, liver (e.g., fetal liver) and/or bone marrow.

[0089] The hematopoietic cells used to produce the NK cells and/or ILC3 cells, and NK cell populations and/or ILC3 cell populations, may be obtained from any animal species. In certain embodiments, the hematopoietic stem or progenitor cells are mammalian cells. In specific embodiments, said hematopoietic stem or progenitor cells are human cells. In specific embodiments, said hematopoietic stem or progenitor cells are primate cells. In specific embodiments, said hematopoietic stem or progenitor cells are canine cells. In specific embodiments, said hematopoietic stem or progenitor cells are rodent cells.

5.1. Hematopoietic Cells

[0090] Hematopoietic cells useful in the methods disclosed herein can be any hematopoietic cells able to differentiate into NK cells and/or ILC3 cells, e.g., precursor cells, hematopoietic progenitor cells, hematopoietic stem cells, or the like. Hematopoietic cells can be obtained from tissue sources such as, e.g., bone marrow, cord blood, placental blood, peripheral blood, liver or the like, or combinations thereof. Hematopoietic cells can be obtained from placenta. In a specific embodiment, the hematopoietic cells are obtained from placental perfusate. In one embodiment, the hematopoietic cells are not obtained from umbilical cord blood. In one embodiment, the hematopoietic cells are not obtained from peripheral blood. Hematopoietic cells from placental perfusate can comprise a mixture of fetal and maternal hematopoietic cells, e.g., a mixture in which maternal cells comprise greater than 5% of the total number of hematopoietic cells. In certain embodiments, hematopoietic cells from placental perfusate comprise at least about 90%, 95%, 98%, 99% or 99.5% fetal cells.

[0091] In another specific embodiment, the hematopoietic cells, e.g., hematopoietic stem cells or progenitor cells, from which the NK cell populations and/or ILC3 cell populations produced using a three-stage method described herein are produced, are obtained from placental perfusate, umbilical cord blood, fetal liver, mobilized peripheral blood, or bone marrow. In another specific embodiment, the hematopoietic cells, e.g., hematopoietic stem cells or progenitor cells, from which

the NK cell populations and/or ILC3 cell populations produced using a three-stage method described herein are produced, are combined cells from placental perfusate and cord blood, e.g., cord blood from the same placenta as the perfusate. In another specific embodiment, said umbilical cord blood is isolated from a placenta other than the placenta from which said placental perfusate is obtained. In certain embodiments, the combined cells can be obtained by pooling or combining the cord blood and placental perfusate. In certain embodiments, the cord blood and placental perfusate are combined at a ratio of 100:1, 95:5, 90:10, 85:15, 80:20, 75:25, 70:30, 65:35, 60:40, 55:45: 50:50, 45:55, 40:60, 35:65, 30:70, 25:75, 20:80, 15:85, 10:90, 5:95, 100:1, 95:1, 90:1, 85:1, 80:1, 75:1, 70:1, 65:1, 60:1, 55:1, 50:1, 45:1, 40:1, 35:1, 30:1, 25:1, 20:1, 15:1, 10:1, 5:1, 1:1, 1:5, 1:10, 1:15, 1:20, 1:25, 1:30, 1:35, 1:40, 1:45, 1:50, 1:55, 1:60, 1:65, 1:70, 1:75, 1:80, 1:85, 1:90, 1:95, 1:100, or the like by volume to obtain the combined cells. In a specific embodiment, the cord blood and placental perfusate are combined at a ratio of from 10:1 to 1:10, from 5:1 to 1:5, or from 3:1 to 1:3. In another specific embodiment, the cord blood and placental perfusate are combined at a ratio of 10:1, 5:1, 3:1, 1:1, 1:3, 1:5 or 1:10. In a more specific embodiment, the cord blood and placental perfusate are combined at a ratio of 8.5:1.5 (85%:15%).

[0092] In certain embodiments, the cord blood and placental perfusate are combined at a ratio of 100:1, 95:5, 90:10, 85:15, 80:20, 75:25, 70:30, 65:35, 60:40, 55:45: 50:50, 45:55, 40:60, 35:65, 30:70, 25:75, 20:80, 15:85, 10:90, 5:95, 100:1, 95:1, 90:1, 85:1, 80:1, 75:1, 70:1, 65:1, 60:1, 55:1, 50:1, 45:1, 40:1, 35:1, 30:1, 25:1, 20:1, 15:1, 10:1, 5:1, 1:1, 1:5, 1:10, 1:15, 1:20, 1:25, 1:30, 1:35, 1:40, 1:45, 1:50, 1:55, 1:60, 1:65, 1:70, 1:75, 1:80, 1:85, 1:90, 1:95, 1:100, or the like by total nucleated cells (TNC) content to obtain the combined cells. In a specific embodiment, the cord blood and placental perfusate are combined at a ratio of from 10:1 to 10:1, from 5:1 to 1:5, or from 3:1 to 1:3. In another specific embodiment, the cord blood and placental perfusate are combined at a ratio of 10:1, 5:1, 3:1, 1:1, 1:3, 1:5 or 1:10.

[0093] In another specific embodiment, the hematopoietic cells, e.g., hematopoietic stem cells or progenitor cells from which said NK cell populations and/or ILC3 cell populations produced using a three-stage method described herein are produced, are from both umbilical cord blood and placental perfusate, but wherein said umbilical cord blood is isolated from a placenta other than the placenta from which said placental perfusate is obtained.

[0094] In certain embodiments, the hematopoietic cells are CD34⁺ cells. In specific embodiments, the hematopoietic cells useful in the methods disclosed herein are CD34^{sup.}+CD38^{sup.}+ or CD34^{sup.}+CD38^{sup.}-. In a more specific embodiment, the hematopoietic cells are CD34^{sup.}+CD38^{sup.}-Lin^{sup.}-. In another specific embodiment, the hematopoietic cells are one or more of CD2^{sup.}-, CD3^{sup.}-, CD11b^{sup.}-, CD11c^{sup.}-, CD14^{sup.}-, CD16^{sup.}-, CD19^{sup.}-, CD24^{sup.}-, CD56^{sup.}-, CD66b^{sup.}- and/or glycophorin A^{sup.}-. In another specific embodiment, the hematopoietic cells are CD2^{sup.}-, CD3^{sup.}-, CD11b^{sup.}-, CD11c^{sup.}-, CD14^{sup.}-, CD16^{sup.}-, CD19^{sup.}-, CD24^{sup.}-, CD56^{sup.}-, CD66b^{sup.}- and glycophorin A^{sup.}-. In another more specific embodiment, the hematopoietic cells are CD34^{sup.}+CD38^{sup.}-CD33^{sup.}-CD117^{sup.}-. In another more specific embodiment, the hematopoietic cells are CD34^{sup.}+CD38^{sup.}-CD33^{sup.}-CD117^{sup.}-CD235^{sup.}-CD36^{sup.}-.

[0095] In another embodiment, the hematopoietic cells are CD45^{sup.}+. In another specific embodiment, the hematopoietic cells are CD34^{sup.}+CD45^{sup.}+. In another embodiment, the hematopoietic cell is Thy-1. In a specific embodiment, the hematopoietic cell is CD34^{sup.}+Thy-1^{sup.}+. In another embodiment, the hematopoietic cells are CD133^{sup.}+. In specific embodiments, the hematopoietic cells are CD34^{sup.}+CD133^{sup.}+ or CD133^{sup.}+Thy-1^{sup.}+. In another specific embodiment, the CD34^{sup.}+ hematopoietic cells are CXCR4^{sup.}+. In another specific embodiment, the CD34^{sup.}+ hematopoietic cells are CXCR4^{sup.}-. In another embodiment, the hematopoietic cells are positive for KDR (vascular growth factor receptor 2). In specific embodiments, the hematopoietic cells are CD34^{sup.}+KDR^{sup.}+, CD133^{sup.}+KDR^{sup.}+ or Thy-1^{sup.}+KDR^{sup.}+. In certain other embodiments, the

hematopoietic cells are positive for aldehyde dehydrogenase (ALDH^{sup.}+), e.g., the cells are CD34^{sup.}+ALDH^{sup.}+

[0096] In certain other embodiments, the CD34^{sup.}+ cells are CD45^{sup.}-. In specific embodiments, the CD34^{sup.}+ cells, e.g., CD34^{sup.}+, CD45^{sup.}- cells express one or more, or all, of the miRNAs hsa-miR-380, hsa-miR-512, hsa-miR-517, hsa-miR-518c, hsa-miR-519b, hsa-miR-520a, hsa-miR-337, hsa-miR-422a, hsa-miR-549, and/or hsa-miR-618.

[0097] In certain embodiments, the hematopoietic cells are CD34^{sup.}-.

[0098] The hematopoietic cells can also lack certain markers that indicate lineage commitment, or a lack of developmental naiveté. For example, in another embodiment, the hematopoietic cells are HLA-DR^{sup.}-. In specific embodiments, the hematopoietic cells are CD34^{sup.}+HLA-DR^{sup.}-, CD133^{sup.}+HLA-DR^{sup.}-, Thy-1^{sup.}+HLA-DR^{sup.}- or ALDH^{sup.}+HLA-DR^{sup.}- In another embodiment, the hematopoietic cells are negative for one or more, or all, of lineage markers CD2, CD3, CD11b, CD11c, CD14, CD16, CD19, CD24, CD56, CD66b and glycophorin A.

[0099] Thus, hematopoietic cells can be selected for use in the methods disclosed herein on the basis of the presence of markers that indicate an undifferentiated state, or on the basis of the absence of lineage markers indicating that at least some lineage differentiation has taken place. Methods of isolating cells, including hematopoietic cells, on the basis of the presence or absence of specific markers is discussed in detail below.

[0100] Hematopoietic cells used in the methods provided herein can be a substantially homogeneous population, e.g., a population comprising at least about 95%, at least about 98% or at least about 99% hematopoietic cells from a single tissue source, or a population comprising hematopoietic cells exhibiting the same hematopoietic cell-associated cellular markers. For example, in various embodiments, the hematopoietic cells can comprise at least about 95%, 98% or 99% hematopoietic cells from bone marrow, cord blood, placental blood, peripheral blood, or placenta, e.g., placenta perfusate.

[0101] Hematopoietic cells used in the methods provided herein can be obtained from a single individual, e.g., from a single placenta, or from a plurality of individuals, e.g., can be pooled. Where the hematopoietic cells are obtained from a plurality of individuals and pooled, the hematopoietic cells may be obtained from the same tissue source. Thus, in various embodiments, the pooled hematopoietic cells are all from placenta, e.g., placental perfusate, all from placental blood, all from umbilical cord blood, all from peripheral blood, and the like.

[0102] Hematopoietic cells used in the methods disclosed herein can, in certain embodiments, comprise hematopoietic cells from two or more tissue sources. For example, in certain embodiments, when hematopoietic cells from two or more sources are combined for use in the methods herein, a plurality of the hematopoietic cells used to produce natural killer cells using a three-stage method described herein comprise hematopoietic cells from placenta, e.g., placenta perfusate. In various embodiments, the hematopoietic cells used to produce NK cell populations and/or ILC3 cell populations produced using a three-stage method described herein, comprise hematopoietic cells from placenta and from cord blood; from placenta and peripheral blood; from placenta and placental blood, or placenta and bone marrow. In one embodiment, the hematopoietic cells comprise hematopoietic cells from placental perfusate in combination with hematopoietic cells from cord blood, wherein the cord blood and placenta are from the same individual, i.e., wherein the perfusate and cord blood are matched. In embodiments in which the hematopoietic cells comprise hematopoietic cells from two tissue sources, the hematopoietic cells from the sources can be combined in a ratio of, for example, 1:10, 2:9, 3:8, 4:7, 5:6, 6:5, 7:4, 8:3, 9:2, 1:10, 1:9, 1:8, 1:7, 1:6, 1:5, 1:4, 1:3, 1:2, 1:1, 2:1, 3:1, 4:1, 5:1, 6:1, 7:1, 8:1 or 9:1.

5.1.1. Placental Hematopoietic Stem Cells

[0103] In certain embodiments, the hematopoietic cells used in the methods provided herein are placental hematopoietic cells. In one embodiment, placental hematopoietic cells are CD34. In a

specific embodiment, the placental hematopoietic cells are predominantly (e.g., at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 98%) CD34.sup.+CD38.sup.+ cells. In another specific embodiment, the placental hematopoietic cells are predominantly (e.g., at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 98%) CD34.sup.+CD38+ cells. Placental hematopoietic cells can be obtained from a post-partum mammalian (e.g., human) placenta by any means known to those of skill in the art, e.g., by perfusion.

[0104] In another embodiment, the placental hematopoietic cell is CD45.sup.-. In a specific embodiment, the hematopoietic cell is CD34.sup.+CD45.sup.-. In another specific embodiment, the placental hematopoietic cells are CD34.sup.+CD45.sup.+.

5.2. Production of Natural Killer and/or ILC3 Cells and Natural Killer Cell and/or ILC3 Cell Populations

[0105] Production of NK cells and/or ILC3 cells and NK cell and/or ILC3 cell populations by the present methods comprises expanding a population of hematopoietic cells. During cell expansion, a plurality of hematopoietic cells within the hematopoietic cell population differentiate into NK cells and/or ILC3 cells. In one aspect, provided herein is a method of producing NK cells comprising culturing hematopoietic stem cells or progenitor cells, e.g., CD34.sup.+ stem cells or progenitor cells, in a first medium comprising a stem cell mobilizing agent and thrombopoietin (Tpo) to produce a first population of cells, subsequently culturing said first population of cells in a second medium comprising a stem cell mobilizing agent and interleukin-15 (IL-15), and lacking Tpo, to produce a second population of cells, and subsequently culturing said second population of cells in a third medium comprising IL-2 and IL-15, and lacking a stem cell mobilizing agent and LMWH, to produce a third population of cells, wherein the third population of cells comprises natural killer cells that are CD56+, CD3-, and wherein at least 70%, for example at least 80%, of the natural killer cells are viable. In certain embodiments, such natural killer cells comprise natural killer cells that are CD16-. In certain embodiments, such natural killer cells comprise natural killer cells that are CD94+. In certain embodiments, such natural killer cells comprise natural killer cells that are CD94+ or CD16+. In certain embodiments, such natural killer cells comprise natural killer cells that are CD94- or CD16-. In certain embodiments, such natural killer cells comprise natural killer cells that are CD94+ and CD16+. In certain embodiments, such natural killer cells comprise natural killer cells that are CD94- and CD16-. In certain embodiments, said first medium and/or said second medium lack leukemia inhibiting factor (LIF) and/or macrophage inflammatory protein-1 alpha (MIP-1 α). In certain embodiments, said third medium lacks LIF, MIP-1 α , and FMS-like tyrosine kinase-3 ligand (Flt-3L). In specific embodiments, said first medium and said second medium lack LIF and MIP-1 α , and said third medium lacks LIF, MIP-1 α , and Flt3L. In certain embodiments, none of the first medium, second medium or third medium comprises heparin, e.g., low-molecular weight heparin.

[0106] In one aspect, provided herein is a method of producing NK cells comprising (a) culturing hematopoietic stem or progenitor cells in a first medium comprising a stem cell mobilizing agent and thrombopoietin (Tpo) to produce a first population of cells; (b) culturing the first population of cells in a second medium comprising a stem cell mobilizing agent and interleukin-15 (IL-15), and lacking Tpo, to produce a second population of cells; and (c) culturing the second population of cells in a third medium comprising IL-2 and IL-15, and lacking LMWH, to produce a third population of cells; wherein the third population of cells comprises natural killer cells that are CD56+, CD3-, and CD11a+. In certain embodiments, said first medium and/or said second medium lack leukemia inhibiting factor (LIF) and/or macrophage inflammatory protein-1 alpha (MIP-1 α). In certain embodiments, said third medium lacks LIF, MIP-1 α , and FMS-like tyrosine kinase-3 ligand (Flt-3L). In specific embodiments, said first medium and said second medium lack LIF and MIP-1 α , and said third medium lacks LIF, MIP-1 α , and Flt3L. In certain embodiments, none of the first medium, second medium or third medium comprises heparin, e.g., low-molecular weight heparin.

[0107] In one aspect, provided herein is a method of producing NK cells comprising (a) culturing hematopoietic stem or progenitor cells in a first medium comprising a stem cell mobilizing agent and thrombopoietin (Tpo) to produce a first population of cells; (b) culturing the first population of cells in a second medium comprising a stem cell mobilizing agent and interleukin-15 (IL-15), and lacking Tpo, to produce a second population of cells; and (c) culturing the second population of cells in a third medium comprising IL-2 and IL-15, and lacking each of stem cell factor (SCF) and LMWH, to produce a third population of cells; wherein the third population of cells comprises natural killer cells that are CD56+, CD3-, and CD11a+. In certain embodiments, said first medium and/or said second medium lack leukemia inhibiting factor (LIF) and/or macrophage inflammatory protein-1 alpha (MIP-1 α). In certain embodiments, said third medium lacks LIF, MIP-1 α , and FMS-like tyrosine kinase-3 ligand (Flt-3L). In specific embodiments, said first medium and said second medium lack LIF and MIP-1 α , and said third medium lacks LIF, MIP-1 α , and Flt3L. In certain embodiments, none of the first medium, second medium or third medium comprises heparin, e.g., low-molecular weight heparin.

[0108] In one aspect, provided herein is a method of producing NK cells comprising (a) culturing hematopoietic stem or progenitor cells in a first medium comprising a stem cell mobilizing agent and thrombopoietin (Tpo) to produce a first population of cells; (b) culturing the first population of cells in a second medium comprising a stem cell mobilizing agent and interleukin-15 (IL-15), and lacking Tpo, to produce a second population of cells; and (c) culturing the second population of cells in a third medium comprising IL-2 and IL-15, and lacking each of SCF, a stem cell mobilizing agent, and LMWH, to produce a third population of cells; wherein the third population of cells comprises natural killer cells that are CD56+, CD3-, and CD11a+. In certain embodiments, said first medium and/or said second medium lack leukemia inhibiting factor (LIF) and/or macrophage inflammatory protein-1 alpha (MIP-1 α). In certain embodiments, said third medium lacks LIF, MIP-1 α , and FMS-like tyrosine kinase-3 ligand (Flt-3L). In specific embodiments, said first medium and said second medium lack LIF and MIP-1 α , and said third medium lacks LIF, MIP-1 α , and Flt3L. In certain embodiments, none of the first medium, second medium or third medium comprises heparin, e.g., low-molecular weight heparin.

[0109] In one aspect, provided herein is a method of producing NK cells comprising (a) culturing hematopoietic stem or progenitor cells in a first medium comprising a stem cell mobilizing agent and thrombopoietin (Tpo) to produce a first population of cells; (b) culturing the first population of cells in a second medium comprising a stem cell mobilizing agent and interleukin-15 (IL-15), and lacking Tpo, to produce a second population of cells; (c) culturing the second population of cells in a third medium comprising IL-2 and IL-15, and lacking each of a stem cell mobilizing agent and LMWH, to produce a third population of cells; and (d) isolating CD11a+ cells from the third population of cells to produce a fourth population of cells; wherein the fourth population of cells comprises natural killer cells that are CD56+, CD3-, and CD11a+. In certain embodiments, said first medium and/or said second medium lack leukemia inhibiting factor (LIF) and/or macrophage inflammatory protein-1 alpha (MIP-1 α). In certain embodiments, said third medium lacks LIF, MIP-1 α , and FMS-like tyrosine kinase-3 ligand (Flt-3L). In specific embodiments, said first medium and said second medium lack LIF and MIP-1 α , and said third medium lacks LIF, MIP-1 α , and Flt3L. In certain embodiments, none of the first medium, second medium or third medium comprises heparin, e.g., low-molecular weight heparin.

[0110] In certain embodiments, of any of the above embodiments, said natural killer cells express perforin and EOMES. In certain embodiments, said natural killer cells do not express either ROR γ t or IL1R1.

[0111] In one aspect, provided herein is a method of producing ILC3 cells comprising (a) culturing hematopoietic stem or progenitor cells in a first medium comprising a stem cell mobilizing agent and thrombopoietin (Tpo) to produce a first population of cells; (b) culturing the first population of cells in a second medium comprising a stem cell mobilizing agent and interleukin-15 (IL-15), and

lacking Tpo, to produce a second population of cells; and (c) culturing the second population of cells in a third medium comprising IL-2 and IL-15, and lacking LMWH, to produce a third population of cells; wherein the third population of cells comprises ILC3 cells that are CD56+, CD3-, and CD11a-. In certain embodiments, said first medium and/or said second medium lack leukemia inhibiting factor (LIF) and/or macrophage inflammatory protein-1 alpha (MIP-1 α). In certain embodiments, said third medium lacks LIF, MIP-1 α , and FMS-like tyrosine kinase-3 ligand (Flt-3L). In specific embodiments, said first medium and said second medium lack LIF and MIP-1 α , and said third medium lacks LIF, MIP-1 α , and Flt3L. In certain embodiments, none of the first medium, second medium or third medium comprises heparin, e.g., low-molecular weight heparin.

[0112] In one aspect, provided herein is a method of producing ILC3 cells comprising (a) culturing hematopoietic stem or progenitor cells in a first medium comprising a stem cell mobilizing agent and thrombopoietin (Tpo) to produce a first population of cells; (b) culturing the first population of cells in a second medium comprising a stem cell mobilizing agent and interleukin-15 (IL-15), and lacking Tpo, to produce a second population of cells; and (c) culturing the second population of cells in a third medium comprising a stem cell mobilizing agent, IL-2 and IL-15, and lacking LMWH, to produce a third population of cells; wherein the third population of cells comprises ILC3 cells that are CD56+, CD3-, and CD11a-. In certain embodiments, said first medium and/or said second medium lack leukemia inhibiting factor (LIF) and/or macrophage inflammatory protein-1 alpha (MIP-1 α). In certain embodiments, said third medium lacks LIF, MIP-1 α , and FMS-like tyrosine kinase-3 ligand (Flt-3L). In specific embodiments, said first medium and said second medium lack LIF and MIP-1 α , and said third medium lacks LIF, MIP-1 α , and Flt3L. In certain embodiments, none of the first medium, second medium or third medium comprises heparin, e.g., low-molecular weight heparin.

[0113] In one aspect, provided herein is a method of producing ILC3 cells comprising (a) culturing hematopoietic stem or progenitor cells in a first medium comprising a stem cell mobilizing agent and thrombopoietin (Tpo) to produce a first population of cells; (b) culturing the first population of cells in a second medium comprising a stem cell mobilizing agent and interleukin-15 (IL-15), and lacking Tpo, to produce a second population of cells; and (c) culturing the second population of cells in a third medium comprising SCF, IL-2 and IL-15, and lacking LMWH, to produce a third population of cells; wherein the third population of cells comprises ILC3 cells that are CD56+, CD3-, and CD11a-. In certain embodiments, said first medium and/or said second medium lack leukemia inhibiting factor (LIF) and/or macrophage inflammatory protein-1 alpha (MIP-1 α). In certain embodiments, said third medium lacks LIF, MIP-1 α , and FMS-like tyrosine kinase-3 ligand (Flt-3L). In specific embodiments, said first medium and said second medium lack LIF and MIP-1 α , and said third medium lacks LIF, MIP-1 α , and Flt3L. In certain embodiments, none of the first medium, second medium or third medium comprises heparin, e.g., low-molecular weight heparin.

[0114] In one aspect, provided herein is a method of producing ILC3 cells comprising (a) culturing hematopoietic stem or progenitor cells in a first medium comprising a stem cell mobilizing agent and thrombopoietin (Tpo) to produce a first population of cells; (b) culturing the first population of cells in a second medium comprising a stem cell mobilizing agent and interleukin-15 (IL-15), and lacking Tpo, to produce a second population of cells; and (c) culturing the second population of cells in a third medium comprising a stem cell mobilizing agent, SCF, IL-2 and IL-15, and lacking LMWH, to produce a third population of cells; wherein the third population of cells comprises ILC3 cells that are CD56+, CD3-, and CD11a-. In certain embodiments, said first medium and/or said second medium lack leukemia inhibiting factor (LIF) and/or macrophage inflammatory protein-1 alpha (MIP-1 α). In certain embodiments, said third medium lacks LIF, MIP-1 α , and FMS-like tyrosine kinase-3 ligand (Flt-3L). In specific embodiments, said first medium and said second medium lack LIF and MIP-1 α , and said third medium lacks LIF, MIP-1 α , and Flt3L. In certain embodiments, none of the first medium, second medium or third medium comprises heparin, e.g., low-molecular weight heparin.

[0115] In one aspect, provided herein is a method of producing ILC3 cells comprising (a) culturing hematopoietic stem or progenitor cells in a first medium comprising a stem cell mobilizing agent and thrombopoietin (Tpo) to produce a first population of cells; (b) culturing the first population of cells in a second medium comprising a stem cell mobilizing agent and interleukin-15 (IL-15), and lacking Tpo, to produce a second population of cells; (c) culturing the second population of cells in a third medium comprising IL-2 and IL-15, and lacking each of a stem cell mobilizing agent and LMWH, to produce a third population of cells; and (d) isolating CD11a⁻ cells, or removing CD11a⁺ cells, from the third population of cells to produce a fourth population of cells; wherein the fourth population of cells comprises ILC3 cells that are CD56⁺, CD3⁻, and CD11a⁻. In certain embodiments, said first medium and/or said second medium lack leukemia inhibiting factor (LIF) and/or macrophage inflammatory protein-1 alpha (MIP-1 α). In certain embodiments, said third medium lacks LIF, MIP-1 α , and FMS-like tyrosine kinase-3 ligand (Flt-3L). In specific embodiments, said first medium and said second medium lack LIF and MIP-1 α , and said third medium lacks LIF, MIP-1 α , and Flt3L. In certain embodiments, none of the first medium, second medium or third medium comprises heparin, e.g., low-molecular weight heparin.

[0116] In certain embodiments, said ILC3 cells express ROR γ t and IL1R1. In certain embodiments, said ILC3 cells do not express either perforin or EOMES.

5.2.1. Production of NK Cell and/or ILC3 Cell Populations Using a Three-Stage Method

[0117] In one embodiment, provided herein is a three-stage method of producing NK cell and/or ILC3 cell populations. In certain embodiments, the method of expansion and differentiation of the hematopoietic cells, as described herein, to produce NK cell and/or ILC3 cell populations according to a three-stage method described herein comprises maintaining the cell population comprising said hematopoietic cells at between about 2×10^4 and about 6×10^6 cells per milliliter. In certain aspects, said hematopoietic stem or progenitor cells are initially inoculated into said first medium from 1×10^4 to 1×10^5 cells/mL. In a specific aspect, said hematopoietic stem or progenitor cells are initially inoculated into said first medium at about 3×10^4 cells/mL.

[0118] In certain aspects, said first population of cells are initially inoculated into said second medium from 5×10^4 to 5×10^5 cells/mL. In a specific aspect, said first population of cells is initially inoculated into said second medium at about 1×10^5 cells/mL.

[0119] In certain aspects said second population of cells is initially inoculated into said third medium from 1×10^5 to 5×10^6 cells/mL. In certain aspects, said second population of cells is initially inoculated into said third medium from 1×10^5 to 1×10^6 cells/mL. In a specific aspect, said second population of cells is initially inoculated into said third medium at about 5×10^5 cells/mL. In a more specific aspect, said second population of cells is initially inoculated into said third medium at about 5×10^5 cells/mL in a spinner flask. In a specific aspect, said second population of cells is initially inoculated into said third medium at about 3×10^5 cells/mL. In a more specific aspect, said second population of cells is initially inoculated into said third medium at about 3×10^5 cells/mL in a static culture.

[0120] In a certain embodiment, the three-stage method comprises a first stage ("stage 1") comprising culturing hematopoietic stem cells or progenitor cells, e.g., CD34⁺ stem cells or progenitor cells, in a first medium for a specified time period, e.g., as described herein, to produce a first population of cells. In certain embodiments, the first medium comprises a stem cell mobilizing agent and thrombopoietin (Tpo). In certain embodiments, the first medium comprises in addition to a stem cell mobilizing agent and Tpo, one or more of LMWH, Flt-3L, SCF, IL-6, IL-7, G-CSF, and GM-CSF. In a specific embodiment, the first medium comprises in addition to a stem cell mobilizing agent and Tpo, each of LMWH, Flt-3L, SCF, IL-6, IL-7, G-CSF, and GM-CSF. In a specific embodiment, the first medium lacks added LMWH. In a specific embodiment, the first medium lacks added desulphated glycosaminoglycans. In a specific embodiment, the first medium lacks LMWH. In a specific embodiment, the first medium lacks desulphated glycosaminoglycans. In a specific embodiment, in addition to a stem cell mobilizing agent and Tpo, each of Flt-3L, SCF,

IL-6, IL-7, G-CSF, and GM-CSF. In specific embodiments, the first medium lacks leukemia inhibiting factor (LIF), macrophage inhibitory protein-1alpha (MIP-1 α) or both.

[0121] In certain embodiments, subsequently, in “stage 2” said cells are cultured in a second medium for a specified time period, e.g., as described herein, to produce a second population of cells. In certain embodiments, the second medium comprises a stem cell mobilizing agent and interleukin-15 (IL-15) and lacks Tpo. In certain embodiments, the second medium comprises, in addition to a stem cell mobilizing agent and IL-15, one or more of LMWH, Flt-3, SCF, IL-6, IL-7, G-CSF, and GM-CSF. In certain embodiments, the second medium comprises, in addition to a stem cell mobilizing agent and IL-15, each of LMWH, Flt-3, SCF, IL-6, IL-7, G-CSF, and GM-CSF. In a specific embodiment, the second medium lacks added LMWH. In a specific embodiment, the second medium lacks added desulphated glycosaminoglycans. In a specific embodiment, the second medium lacks heparin, e.g., LMWH. In a specific embodiment, the second medium lacks desulphated glycosaminoglycans. In certain embodiments, the second medium comprises, in addition to a stem cell mobilizing agent and IL-15, each of Flt-3, SCF, IL-6, IL-7, G-CSF, and GM-CSF. In specific embodiments, the second medium lacks leukemia inhibiting factor (LIF), macrophage inhibitory protein-1alpha (MIP-1 α) or both.

[0122] In certain embodiments, subsequently, in “stage 3” said cells are cultured in a third medium for a specified time period, e.g., as described herein, to produce a third population of cell, e.g., natural killer cells. In certain embodiments, the third medium comprises IL-2 and IL-15, and lacks a stem cell mobilizing agent and LMWH. In certain embodiments, the third medium comprises in addition to IL-2 and IL-15, one or more of SCF, IL-6, IL-7, G-CSF, and GM-CSF. In certain embodiments, the third medium comprises, in addition to IL-2 and IL-15, each of SCF, IL-6, IL-7, G-CSF, and GM-CSF. In specific embodiments, the first medium lacks one, two, or all three of LIF, MIP-1 α , and Flt3L. In specific embodiments, the third medium lacks added desulphated glycosaminoglycans. In specific embodiments, the third medium lacks desulphated glycosaminoglycans. In specific embodiments, the third medium lacks heparin, e.g., LMWH.

[0123] In a specific embodiment, the three-stage method is used to produce NK cell and/or ILC3 cell populations. In certain embodiments, the three-stage method is conducted in the absence of stromal feeder cell support. In certain embodiments, the three-stage method is conducted in the absence of exogenously added steroids (e.g., cortisone, hydrocortisone, or derivatives thereof).

[0124] In certain aspects, said first medium used in the three-stage method comprises a stem cell mobilizing agent and thrombopoietin (Tpo). In certain aspects, the first medium used in the three-stage method comprises, in addition to a stem cell mobilizing agent and Tpo, one or more of Low Molecular Weight Heparin (LMWH), Flt-3 Ligand (Flt-3L), stem cell factor (SCF), IL-6, IL-7, granulocyte colony-stimulating factor (G-CSF), or granulocyte-macrophage-stimulating factor (GM-CSF). In certain aspects, the first medium used in the three-stage method comprises, in addition to a stem cell mobilizing agent and Tpo, each of LMWH, Flt-3L, SCF, IL-6, IL-7, G-CSF, and GM-CSF. In certain aspects, the first medium used in the three-stage method comprises, in addition to a stem cell mobilizing agent and Tpo, each of Flt-3L, SCF, IL-6, IL-7, G-CSF, and GM-CSF. In a specific aspect, the first medium lacks added LMWH. In a specific aspect, the first medium lacks added desulphated glycosaminoglycans. In a specific aspect, the first medium lacks LMWH. In a specific aspect, the first medium lacks desulphated glycosaminoglycans. In certain aspects, said Tpo is present in the first medium at a concentration of from 1 ng/mL to 100 ng/mL, from 1 ng/mL to 50 ng/mL, from 20 ng/mL to 30 ng/mL, or about 25 ng/mL. In other aspects, said Tpo is present in the first medium at a concentration of from 100 ng/mL to 500 ng/mL, from 200 ng/mL to 300 ng/mL, or about 250 ng/mL. In certain aspects, when LMWH is present in the first medium, the LMWH is present at a concentration of from 1 U/mL to 10 U/mL; the Flt-3L is present at a concentration of from 1 ng/mL to 50 ng/mL; the SCF is present at a concentration of from 1 ng/mL to 50 ng/mL; the IL-6 is present at a concentration of from 0.01 ng/mL to 0.1 ng/mL; the IL-7 is present at a concentration of from 1 ng/mL to 50 ng/mL; the G-CSF is present at a

concentration of from 0.01 ng/mL to 0.50 ng/mL; and the GM-CSF is present at a concentration of from 0.005 ng/mL to 0.1 ng/mL. In certain aspects, in the first medium, the Flt-3L is present at a concentration of from 1 ng/mL to 50 ng/mL; the SCF is present at a concentration of from 1 ng/mL to 50 ng/mL; the IL-6 is present at a concentration of from 0.01 ng/mL to 0.1 ng/mL; the IL-7 is present at a concentration of from 1 ng/mL to 50 ng/mL; the G-CSF is present at a concentration of from 0.01 ng/mL to 0.50 ng/mL; and the GM-CSF is present at a concentration of from 0.005 ng/mL to 0.1 ng/mL. In certain aspects, when LMWH is present in the first medium, the LMWH is present at a concentration of from 4 U/mL to 5 U/mL; the Flt-3L is present at a concentration of from 20 ng/mL to 30 ng/mL; the SCF is present at a concentration of from 20 ng/mL to 30 ng/mL; the IL-6 is present at a concentration of from 0.04 ng/mL to 0.06 ng/mL; the IL-7 is present at a concentration of from 20 ng/mL to 30 ng/mL; the G-CSF is present at a concentration of from 0.20 ng/mL to 0.30 ng/mL; and the GM-CSF is present at a concentration of from 0.005 ng/mL to 0.5 ng/mL. In certain aspects, in the first medium, the Flt-3L is present at a concentration of from 20 ng/mL to 30 ng/mL; the SCF is present at a concentration of from 20 ng/mL to 30 ng/mL; the IL-6 is present at a concentration of from 0.04 ng/mL to 0.06 ng/mL; the IL-7 is present at a concentration of from 20 ng/mL to 30 ng/mL; the G-CSF is present at a concentration of from 0.20 ng/mL to 0.30 ng/mL; and the GM-CSF is present at a concentration of from 0.005 ng/mL to 0.5 ng/mL. In certain aspects, when LMWH is present in the first medium, the LMWH is present at a concentration of about 4.5 U/mL; the Flt-3L is present at a concentration of about 25 ng/mL; the SCF is present at a concentration of about 27 ng/mL; the IL-6 is present at a concentration of about 0.05 ng/mL; the IL-7 is present at a concentration of about 25 ng/mL; the G-CSF is present at a concentration of about 0.25 ng/mL; and the GM-CSF is present at a concentration of about 0.01 ng/mL. In certain aspects, in the first medium, the Flt-3L is present at a concentration of about 25 ng/mL; the SCF is present at a concentration of about 27 ng/mL; the IL-6 is present at a concentration of about 0.05 ng/mL; the IL-7 is present at a concentration of about 25 ng/mL; the G-CSF is present at a concentration of about 0.25 ng/mL; and the GM-CSF is present at a concentration of about 0.01 ng/mL. In certain embodiments, said first medium additionally comprises one or more of the following: antibiotics such as gentamycin; antioxidants such as transferrin, insulin, and/or beta-mercaptoethanol; sodium selenite; ascorbic acid; ethanolamine; and glutathione. In certain embodiments, the medium that provides the base for the first medium is a cell/tissue culture medium known to those of skill in the art, e.g., a commercially available cell/tissue culture medium such as SCGM™, STEMMACT™, GBGM®, AIM-V®, X-VIVO™ 10, X-VIVO™ 15, OPTIMIZER, STEMSPAN® H3000, CELLGRO COMPLETE™, DMEM:Ham's F12 ("F12") (e.g., 2:1 ratio, or high glucose or low glucose DMEM), Advanced DMEM (Gibco), EL08-1D2, Myelocult™ H5100, IMDM, and/or RPMI-1640; or is a medium that comprises components generally included in known cell/tissue culture media, such as the components included in GBGM®, AIM-V®, X-VIVO™ 10, X-VIVO™ 15, OPTIMIZER, STEMSPAN® H3000, CELLGRO COMPLETE™, DMEM:Ham's F12 ("F12") (e.g., 2:1 ratio, or high glucose or low glucose DMEM), Advanced DMEM (Gibco), EL08-1D2, Myelocult™ H5100, IMDM, and/or RPMI-1640. In certain embodiments, said first medium is not GBGM®. In specific embodiments of any of the above embodiments, the first medium lacks LIF, MIP-1 α , or both.

[0125] In certain aspects, said second medium used in the three-stage method comprises a stem cell mobilizing agent and interleukin-15 (IL-15), and lacks Tpo. In certain aspects, the second medium used in the three-stage method comprises, in addition to a stem cell mobilizing agent and IL-15, one or more of LMWH, Flt-3, SCF, IL-6, IL-7, G-CSF, and GM-CSF. In certain aspects, the second medium used in the three-stage method comprises, in addition to a stem cell mobilizing agent and IL-15, each of LMWH, Flt-3, SCF, IL-6, IL-7, G-CSF, and GM-CSF. In certain aspects, the second medium used in the three-stage method comprises, in addition to a stem cell mobilizing agent and IL-15, each of Flt-3, SCF, IL-6, IL-7, G-CSF, and GM-CSF. In a specific aspect, the second medium lacks added LMWH. In a specific aspect, the second medium lacks added desulphated

glycosaminoglycans. In a specific aspect, the second medium lacks LMWH. In a specific aspect, the second medium lacks desulphated glycosaminoglycans. In certain aspects, said IL-15 is present in said second medium at a concentration of from 1 ng/mL to 50 ng/mL, from 10 ng/mL to 30 ng/mL, or about 20 ng/mL. In certain aspects, when LMWH is present in said second medium, the LMWH is present at a concentration of from 1 U/mL to 10 U/mL; the Flt-3L is present at a concentration of from 1 ng/mL to 50 ng/mL; the SCF is present at a concentration of from 1 ng/mL to 50 ng/mL; the IL-6 is present at a concentration of from 0.01 ng/mL to 0.1 ng/mL; the IL-7 is present at a concentration of from 1 ng/mL to 50 ng/mL; the G-CSF is present at a concentration of from 0.01 ng/mL to 0.50 ng/mL; and the GM-CSF is present at a concentration of from 0.005 ng/mL to 0.1 ng/mL. In certain aspects, in said second medium, the Flt-3L is present at a concentration of from 1 ng/mL to 50 ng/mL; the SCF is present at a concentration of from 1 ng/mL to 50 ng/mL; the IL-6 is present at a concentration of from 0.01 ng/mL to 0.1 ng/mL; the IL-7 is present at a concentration of from 1 ng/mL to 50 ng/mL; the G-CSF is present at a concentration of from 0.01 ng/mL to 0.50 ng/mL; and the GM-CSF is present at a concentration of from 0.005 ng/mL to 0.1 ng/mL. In certain aspects, when LMWH is present in the second medium, the LMWH is present in the second medium at a concentration of from 4 U/mL to 5 U/mL; the Flt-3L is present at a concentration of from 20 ng/mL to 30 ng/mL; the SCF is present at a concentration of from 20 ng/mL to 30 ng/mL; the IL-6 is present at a concentration of from 0.04 ng/mL to 0.06 ng/mL; the IL-7 is present at a concentration of from 20 ng/mL to 30 ng/mL; the G-CSF is present at a concentration of from 0.20 ng/mL to 0.30 ng/mL; and the GM-CSF is present at a concentration of from 0.005 ng/mL to 0.5 ng/mL. In certain aspects, in the second medium, the Flt-3L is present at a concentration of from 20 ng/mL to 30 ng/mL; the SCF is present at a concentration of from 20 ng/mL to 30 ng/mL; the IL-6 is present at a concentration of from 0.04 ng/mL to 0.06 ng/mL; the IL-7 is present at a concentration of from 20 ng/mL to 30 ng/mL; the G-CSF is present at a concentration of from 0.20 ng/mL to 0.30 ng/mL; and the GM-CSF is present at a concentration of from 0.005 ng/mL to 0.5 ng/mL. In certain aspects, when LMWH is present in the second medium, the LMWH is present in the second medium at a concentration of from 4 U/mL to 5 U/mL; the Flt-3L is present at a concentration of from 20 ng/mL to 30 ng/mL; the SCF is present at a concentration of from 20 ng/mL to 30 ng/mL; the IL-6 is present at a concentration of from 0.04 ng/mL to 0.06 ng/mL; the IL-7 is present at a concentration of from 20 ng/mL to 30 ng/mL; the G-CSF is present at a concentration of from 0.20 ng/mL to 0.30 ng/mL; and the GM-CSF is present at a concentration of from 0.005 ng/mL to 0.5 ng/mL. In certain aspects, in the second medium, the Flt-3L is present at a concentration of from 20 ng/mL to 30 ng/mL; the SCF is present at a concentration of from 20 ng/mL to 30 ng/mL; the IL-6 is present at a concentration of from 0.04 ng/mL to 0.06 ng/mL; the IL-7 is present at a concentration of from 20 ng/mL to 30 ng/mL; the G-CSF is present at a concentration of from 0.20 ng/mL to 0.30 ng/mL; and the GM-CSF is present at a concentration of from 0.005 ng/mL to 0.5 ng/mL. In certain aspects, when LMWH is present in the second medium, the LMWH is present in the second medium at a concentration of about 4.5 U/mL; the Flt-3L is present at a concentration of about 25 ng/mL; the SCF is present at a concentration of about 27 ng/mL; the IL-6 is present at a concentration of about 0.05 ng/mL; the IL-7 is present at a concentration of about 25 ng/mL; the G-CSF is present at a concentration of about 0.25 ng/mL; and the GM-CSF is present at a concentration of about 0.01 ng/mL. In certain aspects, in the second medium, the Flt-3L is present at a concentration of about 25 ng/mL; the SCF is present at a concentration of about 27 ng/mL; the IL-6 is present at a concentration of about 0.05 ng/mL; the IL-7 is present at a concentration of about 25 ng/mL; the G-CSF is present at a concentration of about 0.25 ng/mL; and the GM-CSF is present at a concentration of about 0.01 ng/mL. In certain embodiments, said second medium additionally comprises one or more of the following: antibiotics such as gentamycin; antioxidants such as transferrin, insulin, and/or beta-mercaptoethanol; sodium selenite; ascorbic acid; ethanolamine; and glutathione. In certain embodiments, the medium that provides the base for the second medium is a cell/tissue culture

medium known to in the art, e.g., a commercially available cell/tissue culture medium such as SCGM™, STEMMACS™, GBGM®, AIM-V®, X-VIVO™ 10, X-VIVO™ 15, OPTMIZER, STEMSPAN® H3000, CELLGRO COMPLETE™, DMEM:Ham's F12 ("F12") (e.g., 2:1 ratio, or high glucose or low glucose DMEM), Advanced DMEM (Gibco), EL08-1D2, Myelocult™ H5100, IMDM, and/or RPMI-1640; or is a medium that comprises components generally included in known cell/tissue culture media, such as the components included in GBGM®, AIM-V®, X-VIVO™ 10, X-VIVO™ 15, OPTMIZER, STEMSPAN® H3000, CELLGRO COMPLETE™, DMEM:Ham's F12 ("F12") (e.g., 2:1 ratio, or high glucose or low glucose DMEM), Advanced DMEM (Gibco), EL08-1D2, Myelocult™ H5100, IMDM, and/or RPMI-1640. In certain embodiments, said second medium is not GBGM®. In specific embodiments of any of the above embodiments, the first medium lacks LIF, MIP-1 α , or both.

[0126] In certain aspects, said third medium used in the three-stage method comprises IL-2 and IL-15, and lacks a stem cell mobilizing agent and LMWH. In certain aspects, said third medium used in the three-stage method comprises IL-2 and IL-15, and lacks LMWH. In certain aspects, said third medium used in the three-stage method comprises IL-2 and IL-15, and lacks SCF and LMWH. In certain aspects, said third medium used in the three-stage method comprises IL-2 and IL-15, and lacks SCF, a stem cell mobilizing agent and LMWH. In certain aspects, said third medium used in the three-stage method comprises a stem cell mobilizing agent, IL-2 and IL-15, and lacks LMWH. In certain aspects, said third medium used in the three-stage method comprises SCF, IL-2 and IL-15, and lacks LMWH. In certain aspects, said third medium used in the three-stage method comprises a stem cell mobilizing agent, SCF, IL-2 and IL-15, and lacks LMWH. In certain aspects, said third medium used in the three-stage method comprises IL-2 and IL-15, and lacks a stem cell mobilizing agent and LMWH. In certain aspects, the third medium used in the three-stage method comprises, in addition to IL-2 and IL-15, one or more of SCF, IL-6, IL-7, G-CSF, or GM-CSF. In certain aspects, the third medium used in the three-stage method comprises, in addition to IL-2 and IL-15, each of SCF, IL-6, IL-7, G-CSF, and GM-CSF. In certain aspects, said IL-2 is present in said third medium at a concentration of from 10 U/mL to 10,000 U/mL and said IL-15 is present in said third medium at a concentration of from 1 ng/mL to 50 ng/mL. In certain aspects, said IL-2 is present in said third medium at a concentration of from 100 U/mL to 10,000 U/mL and said IL-15 is present in said third medium at a concentration of from 1 ng/mL to 50 ng/mL. In certain aspects, said IL-2 is present in said third medium at a concentration of from 300 U/mL to 3,000 U/mL and said IL-15 is present in said third medium at a concentration of from 10 ng/mL to 30 ng/mL. In certain aspects, said IL-2 is present in said third medium at a concentration of about 1,000 U/mL and said IL-15 is present in said third medium at a concentration of about 20 ng/mL. In certain aspects, in said third medium, the SCF is present at a concentration of from 1 ng/mL to 50 ng/mL; the IL-6 is present at a concentration of from 0.01 ng/mL to 0.1 ng/mL; the IL-7 is present at a concentration of from 1 ng/mL to 50 ng/mL; the G-CSF is present at a concentration of from 0.01 ng/mL to 0.50 ng/mL; and the GM-CSF is present at a concentration of from 0.005 ng/mL to 0.1 ng/mL. In certain aspects, in said third medium, the SCF is present at a concentration of from 20 ng/mL to 30 ng/mL; the IL-6 is present at a concentration of from 0.04 ng/mL to 0.06 ng/mL; the IL-7 is present at a concentration of from 20 ng/mL to 30 ng/mL; the G-CSF is present at a concentration of from 0.20 ng/mL to 0.30 ng/mL; and the GM-CSF is present at a concentration of from 0.005 ng/mL to 0.5 ng/mL. In certain aspects, in said third medium, the SCF is present at a concentration of about 22 ng/mL; the IL-6 is present at a concentration of about 0.05 ng/mL; the IL-7 is present at a concentration of about 20 ng/mL; the G-CSF is present at a concentration of about 0.25 ng/mL; and the GM-CSF is present at a concentration of about 0.01 ng/mL. In certain aspects, the third medium comprises 100 ng/mL IL-7, 1000 ng/mL IL-2, 20 ng/mL IL-15, and 10 stem cell mobilizing agent and lacks SCF. In certain aspects, the third medium comprises 20 ng/mL IL-7, 1000 ng/mL IL-2, 20 ng/mL IL-15, and stem cell mobilizing agent and lacks SCF. In certain aspects, the third medium comprises 20 ng/mL IL-7, 20 ng/mL IL-

15, and stem cell mobilizing agent and lacks SCF. In certain aspects, the third medium comprises 100 ng/mL IL-7, 22 ng/mL SCF, 1000 ng/mL IL-2, and 20 ng/mL IL-15 and lacks stem cell mobilizing agent. In certain aspects, the third medium comprises 22 ng/mL SCF, 1000 ng/mL IL-2, and 20 ng/mL IL-15 and lacks stem cell mobilizing agent. In certain aspects, the third medium comprises 20 ng/mL IL-7, 22 ng/mL SCF, 1000 ng/mL IL-2, and 20 ng/mL IL-15 and lacks stem cell mobilizing agent. In certain aspects, the third medium comprises 20 ng/mL IL-7, 22 ng/mL SCF, and 1000 ng/mL IL-2 and lacks stem cell mobilizing agent. In specific embodiments of any of the above embodiments, the first medium lacks one, two, or all three of LIF, MIP-1 α , Flt-3L. [0127] In certain embodiments, said third medium additionally comprises one or more of the following: antibiotics such as gentamycin; antioxidants such as transferrin, insulin, and/or beta-mercaptoethanol; sodium selenite; ascorbic acid; ethanolamine; and glutathione. In certain embodiments, the medium that provides the base for the third medium is a cell/tissue culture medium known to those of skill in the art, e.g., a commercially available cell/tissue culture medium such as SCGMTM, STEMMACSTM, GBGM[®], AIM-V[®], X-VIVOTM 10, X-VIVOTM 15, OPTIMIZER, STEMSPAN[®] H3000, CELLGRO COMPLETETM, DMEM:Ham's F12 ("F12") (e.g., 2:1 ratio, or high glucose or low glucose DMEM), Advanced DMEM (Gibco), EL08-1D2, MyelocultTM H5100, IMDM, and/or RPMI-1640; or is a medium that comprises components generally included in known cell/tissue culture media, such as the components included in GBGM[®], AIM-V[®], X-VIVOTM 10, X-VIVOTM 15, OPTIMIZER, STEMSPAN[®] H3000, CELLGRO COMPLETETM, DMEM:Ham's F12 ("F12") (e.g., 2:1 ratio, or high glucose or low glucose DMEM), Advanced DMEM (Gibco), EL08-1D2, MyelocultTM H5100, IMDM, and/or RPMI-1640. In certain embodiments, said third medium is not GBGM[®].

[0128] Generally, the particularly recited medium components do not refer to possible constituents in an undefined component of said medium. For example, said Tpo, IL-2, and IL-15 are not comprised within an undefined component of the first medium, second medium or third medium, e.g., said Tpo, IL-2, and IL-15 are not comprised within serum. Further, said LMWH, Flt-3, SCF, IL-6, IL-7, G-CSF, and/or GM-CSF are not comprised within an undefined component of the first medium, second medium or third medium, e.g., said LMWH, Flt-3, SCF, IL-6, IL-7, G-CSF, and/or GM-CSF are not comprised within serum.

[0129] In certain aspects, said first medium, second medium or third medium comprises human serum-AB. In certain aspects, any of said first medium, second medium or third medium comprises 1% to 20% human serum-AB, 5% to 15% human serum-AB, or about 2, 5, or 10% human serum-AB.

[0130] In certain embodiments, in the three-stage methods described herein, said hematopoietic stem or progenitor cells are cultured in said first medium for 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 days. In certain embodiments, in the three-stage methods described herein, cells are cultured in said second medium for 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 days. In certain embodiments, in the three-stage methods described herein, cells are cultured in said third medium for 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 days, or for more than 30 days.

[0131] In a specific embodiment, in the three-stage methods described herein, said hematopoietic stem or progenitor cells are cultured in said first medium for 7-13 days to produce a first population of cells, before said culturing in said second medium; said first population of cells are cultured in said second medium for 2-6 days to produce a second population of cells before said culturing in said third medium; and said second population of cells are cultured in said third medium for 10-30 days, i.e., the cells are cultured a total of 19-49 days.

[0132] In a specific embodiment, in the three-stage methods described herein, in the three-stage methods described herein, said hematopoietic stem or progenitor cells are cultured in said first medium for 8-12 days to produce a first population of cells, before said culturing in said second medium; said first population of cells are cultured in said second medium for 3-5 days to produce a

second population of cells before said culturing in said third medium; and said second population of cells are cultured in said third medium for 15-25 days, i.e., the cells are cultured a total of 26-42 days.

[0133] In a specific embodiment, in the three-stage methods described herein, said hematopoietic stem or progenitor cells are cultured in said first medium for about 10 days to produce a first population of cells, before said culturing in said second medium; said first population of cells are cultured in said second medium for about 4 days to produce a second population of cells before said culturing in said third medium; and said second population of cells are cultured in said third medium for about 21 days, i.e., the cells are cultured a total of about 35 days.

[0134] In certain aspects, the three-stage method disclosed herein produces at least 5000-fold more natural killer cells as compared to the number of hematopoietic stem cells initially inoculated into said first medium. In certain aspects, said three-stage method produces at least 10,000-fold more natural killer cells as compared to the number of hematopoietic stem cells initially inoculated into said first medium. In certain aspects, said three-stage method produces at least 50,000-fold more natural killer cells as compared to the number of hematopoietic stem cells initially inoculated into said first medium. In certain aspects, said three-stage method produces at least 75,000-fold more natural killer cells as compared to the number of hematopoietic stem cells initially inoculated into said first medium. In certain aspects, the viability of said natural killer cells is determined by 7-aminoactinomycin D (7AAD) staining. In certain aspects, the viability of said natural killer cells is determined by annexin-V staining. In specific aspects, the viability of said natural killer cells is determined by both 7-AAD staining and annexin-V staining. In certain aspects, the viability of said natural killer cells is determined by trypan blue staining.

[0135] In certain aspects, the three-stage method disclosed herein produces at least 5000-fold more ILC3 cells as compared to the number of hematopoietic stem cells initially inoculated into said first medium. In certain aspects, said three-stage method produces at least 10,000-fold more ILC3 cells as compared to the number of hematopoietic stem cells initially inoculated into said first medium. In certain aspects, said three-stage method produces at least 50,000-fold more ILC3 cells as compared to the number of hematopoietic stem cells initially inoculated into said first medium. In certain aspects, said three-stage method produces at least 75,000-fold more ILC3 cells as compared to the number of hematopoietic stem cells initially inoculated into said first medium.

[0136] In certain aspects, the three-stage method produces natural killer cells that comprise at least 20% CD56+CD3⁻ natural killer cells. In certain aspects, the three-stage method produces natural killer cells that comprise at least 40% CD56+CD3⁻ natural killer cells. In certain aspects, the three-stage method produces natural killer cells that comprise at least 60% CD56+CD3⁻ natural killer cells. In certain aspects, the three-stage method produces natural killer cells that comprise at least 70% CD56+CD3⁻ natural killer cells. In certain aspects, the three-stage method produces natural killer cells that comprise at least 80% CD56+CD3⁻ natural killer cells.

[0137] In certain aspects, the three-stage method disclosed herein produces natural killer cells that comprise at least 20% CD56+CD3⁻CD11a⁺ natural killer cells. In certain aspects, the three-stage method disclosed herein produces natural killer cells that comprise at least 40% CD56+CD3⁻CD11a⁺ natural killer cells. In certain aspects, the three-stage method disclosed herein produces natural killer cells that comprise at least 60% CD56+CD3⁻CD11a⁺ natural killer cells. In certain aspects, the three-stage method disclosed herein produces natural killer cells that comprise at least 80% CD56+CD3⁻CD11a⁺ natural killer cells.

[0138] In certain aspects, the three-stage method disclosed herein produces ILC3 cells that comprise at least 20% CD56+CD3⁻CD11a⁻ ILC3 cells. In certain aspects, the three-stage method disclosed herein produces ILC3 cells that comprise at least 40% CD56+CD3⁻CD11a⁻ ILC3 cells. In certain aspects, the three-stage method disclosed herein produces ILC3 cells that comprise at least 60% CD56+CD3⁻CD11a⁻ ILC3 cells. In certain aspects, the three-stage method disclosed herein produces natural killer cells that comprise at least 80% CD56+CD3⁻CD11a⁻ ILC3 cells.

[0139] In certain aspects, the three-stage method produces natural killer cells that exhibit at least 20% cytotoxicity against K562 cells when said natural killer cells and said K562 cells are co-cultured in vitro or ex vivo at a ratio of 10:1. In certain aspects, the three-stage method produces natural killer cells that exhibit at least 35% cytotoxicity against the K562 cells when said natural killer cells and said K562 cells are co-cultured in vitro or ex vivo at a ratio of 10:1. In certain aspects, the three-stage method produces natural killer cells that exhibit at least 45% cytotoxicity against the K562 cells when said natural killer cells and said K562 cells are co-cultured in vitro or ex vivo at a ratio of 10:1. In certain aspects, the three-stage method produces natural killer cells that exhibit at least 60% cytotoxicity against the K562 cells when said natural killer cells and said K562 cells are co-cultured in vitro or ex vivo at a ratio of 10:1. In certain aspects, the three-stage method produces natural killer cells that exhibit at least 75% cytotoxicity against the K562 cells when said natural killer cells and said K562 cells are co-cultured in vitro or ex vivo at a ratio of 10:1.

[0140] In certain aspects, the three-stage method produces ILC3 cells that exhibit at least 20% cytotoxicity against K562 cells when said ILC3 cells and said K562 cells are co-cultured in vitro or ex vivo at a ratio of 10:1. In certain aspects, the three-stage method produces ILC3 cells that exhibit at least 35% cytotoxicity against the K562 cells when said ILC3 cells and said K562 cells are co-cultured in vitro or ex vivo at a ratio of 10:1. In certain aspects, the three-stage method produces ILC3 cells that exhibit at least 45% cytotoxicity against the K562 cells when said ILC3 cells and said K562 cells are co-cultured in vitro or ex vivo at a ratio of 10:1. In certain aspects, the three-stage method produces ILC3 cells that exhibit at least 60% cytotoxicity against the K562 cells when said ILC3 cells and said K562 cells are co-cultured in vitro or ex vivo at a ratio of 10:1. In certain aspects, the three-stage method produces ILC3 cells that exhibit at least 75% cytotoxicity against the K562 cells when said ILC3 cells and said K562 cells are co-cultured in vitro or ex vivo at a ratio of 10:1.

[0141] In certain aspects, after said third culturing step, said third population of cells, e.g., said population of natural killer cells and/or ILC3 cells, is cryopreserved. In certain aspects, after said fourth step, said fourth population of cells, e.g., said population of natural killer cells and/or ILC3 cells, is cryopreserved.

[0142] In certain aspects, provided herein are populations of cells comprising natural killer cells, i.e., natural killers cells produced by a three-stage method described herein. Accordingly, provided herein is an isolated natural killer cell population produced by a three-stage method described herein. In a specific embodiment, said natural killer cell population comprises at least 20% CD56+CD3⁻ natural killer cells. In a specific embodiment, said natural killer cell population comprises at least 40% CD56+CD3⁻ natural killer cells. In a specific embodiment, said natural killer cell population comprises at least 60% CD56+CD3⁻ natural killer cells. In a specific embodiment, said natural killer cell population comprises at least 80% CD56+CD3⁻ natural killer cells. In a specific embodiment, said natural killer cell population comprises at least 60% CD16⁻ cells. In a specific embodiment, said natural killer cell population comprises at least 80% CD16⁻ cells. In a specific embodiment, said natural killer cell population comprises at least 20% CD94⁺ cells. In a specific embodiment, said natural killer cell population comprises at least 40% CD94⁺ cells.

[0143] In certain aspects, provided herein is a population of natural killer cells that is CD56+CD3⁻CD117+CD11a⁺, wherein said natural killer cells express perforin and/or EOMES, and do not express one or more of ROR γ t, aryl hydrocarbon receptor (AHR), and IL1R1. In certain aspects, said natural killer cells express perforin and EOMES, and do not express any of ROR γ t, aryl hydrocarbon receptor, or IL1R1. In certain aspects, said natural killer cells additionally express T-bet, GZMB, NKP46, NKP30, and NKG2D. In certain aspects, said natural killer cells express CD94. In certain aspects, said natural killer cells do not express CD94.

[0144] In certain aspects, provided herein is a population of ILC3 cells that is CD56+CD3⁻

CD117+CD11a⁻, wherein said ILC3 cells express one or more of ROR γ t, aryl hydrocarbon receptor, and IL1R1, and do not express one or more of CD94, perforin, and EOMES. In certain aspects, said ILC3 cells express ROR γ t, aryl hydrocarbon receptor, and IL1R1, and do not express any of CD94, perforin, or EOMES. In certain aspects, said ILC3 cells additionally express CD226 and/or 2B4. In certain aspects, said ILC3 cells additionally express one or more of IL-22, TNF α , and DNAM-1. In certain aspects, said ILC3 cells express CD226, 2B4, IL-22, TNF α , and DNAM-1.

[0145] In certain aspects, provided herein is a method of producing a cell population comprising natural killer cells and ILC3 cells, comprising (a) culturing hematopoietic stem or progenitor cells in a first medium comprising a stem cell mobilizing agent and thrombopoietin (Tpo) to produce a first population of cells; (b) culturing the first population of cells in a second medium comprising a stem cell mobilizing agent and interleukin-15 (IL-15), and lacking Tpo, to produce a second population of cells; (c) culturing the second population of cells in a third medium comprising IL-2 and IL-15, and lacking each of a stem cell mobilizing agent and LMWH, to produce a third population of cells; and (d) separating CD11a⁺ cells and CD11a⁻ cells from the third population of cells; and (e) combining the CD11a⁺ cells with the CD11a⁻ cells in a ratio of 50:1, 40:1, 30:1, 20:1, 10:1, 5:1, 4:1, 3:1, 2:1, 1:1, 1:2, 1:3, 1:4, 1:5, 1:10, 1:20, 1:30, 1:40, or 1:50 to produce a fourth population of cells. In certain embodiments, said first medium and/or said second medium lack leukemia inhibiting factor (LIF) and/or macrophage inflammatory protein-1 alpha (MIP-1 α). In certain embodiments, said third medium lacks LIF, MIP-1 α , and FMS-like tyrosine kinase-3 ligand (Flt-3L). In specific embodiments, said first medium and said second medium lack LIF and MIP-1 α , and said third medium lacks LIF, MIP-1 α , and Flt3L. In certain embodiments, none of the first medium, second medium or third medium comprises heparin, e.g., low-molecular weight heparin. In certain aspects, in the fourth population of cells, the CD11a⁺ cells and CD11a⁻ cells are combined in a ratio of 50:1. In certain aspects, in the fourth population of cells, the CD11a⁺ cells and CD11a⁻ cells are combined in a ratio of 20:1. In certain aspects, in the fourth population of cells, the CD11a⁺ cells and CD11a⁻ cells are combined in a ratio of 10:1. In certain aspects, in the fourth population of cells, the CD11a⁺ cells and CD11a⁻ cells are combined in a ratio of 5:1. In certain aspects, in the fourth population of cells, the CD11a⁺ cells and CD11a⁻ cells are combined in a ratio of 1:1. In certain aspects, in the fourth population of cells, the CD11a⁺ cells and CD11a⁻ cells are combined in a ratio of 1:5. In certain aspects, in the fourth population of cells, the CD11a⁺ cells and CD11a⁻ cells are combined in a ratio of 1:10. In certain aspects, in the fourth population of cells, the CD11a⁺ cells and CD11a⁻ cells are combined in a ratio of 1:20. In certain aspects, in the fourth population of cells, the CD11a⁺ cells and CD11a⁻ cells are combined in a ratio of 1:50.

5.3. Stem Cell Mobilizing Factors

5.3.1. Chemistry Definitions

[0146] To facilitate understanding of the disclosure of stem cell mobilizing factors set forth herein, a number of terms are defined below.

[0147] Generally, the nomenclature used herein and the laboratory procedures in biology, cellular biology, biochemistry, organic chemistry, medicinal chemistry, and pharmacology described herein are those well known and commonly employed in the art. Unless defined otherwise, all technical and scientific terms used herein generally have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs.

[0148] The term “about” or “approximately” means an acceptable error for a particular value as determined by one of ordinary skill in the art, which depends in part on how the value is measured or determined. In certain embodiments, the term “about” or “approximately” means within 1, 2, 3, or 4 standard deviations. In certain embodiments, the term “about” or “approximately” means within 50%, 20%, 15%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, 0.5%, or 0.05% of a given value or range.

[0149] As used herein, any “R” group(s) such as, without limitation, R^{sup.a}, R^{sup.b}, R^{sup.c},

R.sup.d, R.sup.e, R.sup.f, R.sup.g, R.sup.h, R.sup.m, R.sup.G, R.sup.J, R.sup.K, R.sup.U, R.sup.V, R.sup.Y, and R.sup.Z represent substituents that can be attached to the indicated atom. An R group may be substituted or unsubstituted. If two “R” groups are described as being “taken together” the R groups and the atoms they are attached to can form a cycloalkyl, cycloalkenyl, aryl, heteroaryl or heterocycle. For example, without limitation, if R.sup.a and R.sup.b of an NR.sup.aR.sup.b group are indicated to be “taken together,” it means that they are covalently bonded to one another to form a ring:

##STR00001##

[0150] In addition, if two “R” groups are described as being “taken together” with the atom(s) to which they are attached to form a ring as an alternative, the R groups are not limited to the variables or substituents defined previously.

[0151] Whenever a group is described as being “optionally substituted” that group may be unsubstituted or substituted with one or more of the indicated substituents. Likewise, when a group is described as being “unsubstituted or substituted” if substituted, the substituent(s) may be selected from one or more the indicated substituents. If no substituents are indicated, it is meant that the indicated “optionally substituted” or “substituted” group may be substituted with one or more group(s) individually and independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, acylalkyl, hydroxy, alkoxy, alkoxyalkyl, aminoalkyl, amino acid, aryl, heteroaryl, heterocyclyl, aryl(alkyl), heteroaryl(alkyl), heterocyclyl(alkyl), hydroxyalkyl, acyl, cyano, halogen, thiocarbonyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, S-sulfonamido, N-sulfonamido, C-carboxy, O-carboxy, isocyanato, thiocyanato, isothiocyanato, azido, nitro, silyl, sulfenyl, sulfinyl, sulfonyl, haloalkyl, haloalkoxy, trihalomethanesulfonyl, trihalomethanesulfonamido, an amino, a mono-substituted amino group and a di-substituted amino group.

[0152] As used herein, “C.sub.a to C.sub.b” in which “a” and “b” are integers refer to the number of carbon atoms in an alkyl, alkenyl or alkynyl group, or the number of carbon atoms in the ring of a cycloalkyl, cycloalkenyl, aryl, heteroaryl or heteroalicyclyl group. That is, the alkyl, alkenyl, alkynyl, ring(s) of the cycloalkyl, ring(s) of the cycloalkenyl, ring(s) of the aryl, ring(s) of the heteroaryl or ring(s) of the heteroalicyclyl can contain from “a” to “b”, inclusive, carbon atoms. Thus, for example, a “C.sub.1 to C.sub.4 alkyl” group refers to all alkyl groups having from 1 to 4 carbons, that is, CH.sub.3—, CH.sub.3CH.sub.2—, CH.sub.3CH.sub.2CH.sub.2—, (CH.sub.3).sub.2CH—, CH.sub.3CH.sub.2CH.sub.2CH.sub.2—, CH.sub.3CH.sub.2CH(CH.sub.3)— and (CH.sub.3).sub.3C—. If no “a” and “b” are designated with regard to an alkyl, alkenyl, alkynyl, cycloalkyl cycloalkenyl, aryl, heteroaryl or heteroalicyclyl group, the broadest range described in these definitions is to be assumed.

[0153] As used herein, “alkyl” refers to a straight or branched hydrocarbon chain that comprises a fully saturated (no double or triple bonds) hydrocarbon group. The alkyl group may have 1 to 20 carbon atoms (whenever it appears herein, a numerical range such as “1 to 20” refers to each integer in the given range; e.g., “1 to 20 carbon atoms” means that the alkyl group may consist of 1 carbon atom, 2 carbon atoms, 3 carbon atoms, etc., up to and including 20 carbon atoms, although the present definition also covers the occurrence of the term “alkyl” where no numerical range is designated). The alkyl group may also be a medium size alkyl having 1 to 10 carbon atoms. The alkyl group could also be a lower alkyl having 1 to 6 carbon atoms. The alkyl group of the compounds may be designated as “C.sub.1-C.sub.4 alkyl” or similar designations. By way of example only, “C.sub.1-C.sub.4 alkyl” indicates that there are one to four carbon atoms in the alkyl chain, i.e., the alkyl chain is selected from methyl, ethyl, propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, and t-butyl. Typical alkyl groups include, but are in no way limited to, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tertiary butyl, pentyl and hexyl. The alkyl group may be substituted or unsubstituted.

[0154] As used herein, “alkenyl” refers to an alkyl group that contains in the straight or branched

hydrocarbon chain one or more double bonds. Examples of alkenyl groups include allenyl, vinylmethyl and ethenyl. An alkenyl group may be unsubstituted or substituted.

[0155] As used herein, “alkynyl” refers to an alkyl group that contains in the straight or branched hydrocarbon chain one or more triple bonds. Examples of alkynyls include ethynyl and propynyl. An alkynyl group may be unsubstituted or substituted.

[0156] As used herein, “cycloalkyl” refers to a completely saturated (no double or triple bonds) mono- or multi-cyclic hydrocarbon ring system. When composed of two or more rings, the rings may be joined together in a fused fashion. Cycloalkyl groups can contain 3 to 10 atoms in the ring(s) or 3 to 8 atoms in the ring(s). A cycloalkyl group may be unsubstituted or substituted. Typical cycloalkyl groups include, but are in no way limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl.

[0157] As used herein, “cycloalkenyl” refers to a mono- or multi-cyclic hydrocarbon ring system that contains one or more double bonds in at least one ring; although, if there is more than one, the double bonds cannot form a fully delocalized pi-electron system throughout all the rings (otherwise the group would be “aryl,” as defined herein). Cycloalkenyl groups can contain 3 to 10 atoms in the ring(s) or 3 to 8 atoms in the ring(s). When composed of two or more rings, the rings may be connected together in a fused fashion. A cycloalkenyl group may be unsubstituted or substituted.

[0158] As used herein, “aryl” refers to a carbocyclic (all carbon) monocyclic or multicyclic aromatic ring system (including fused ring systems where two carbocyclic rings share a chemical bond) that has a fully delocalized pi-electron system throughout all the rings. The number of carbon atoms in an aryl group can vary. For example, the aryl group can be a C.sub.6-C.sub.14 aryl group, a C.sub.6-C.sub.10 aryl group, or a C.sub.6 aryl group. Examples of aryl groups include, but are not limited to, benzene, naphthalene and azulene. An aryl group may be substituted or unsubstituted.

[0159] As used herein, “heteroaryl” refers to a monocyclic or multicyclic aromatic ring system (a ring system with fully delocalized pi-electron system) that contain(s) one, two, three or more heteroatoms, that is, an element other than carbon, including but not limited to, nitrogen, oxygen and sulfur. The number of atoms in the ring(s) of a heteroaryl group can vary. For example, the heteroaryl group can contain 4 to 14 atoms in the ring(s), 5 to 10 atoms in the ring(s) or 5 to 6 atoms in the ring(s). Furthermore, the term “heteroaryl” includes fused ring systems where two rings, such as at least one aryl ring and at least one heteroaryl ring, or at least two heteroaryl rings, share at least one chemical bond. Examples of heteroaryl rings include, but are not limited to, those described herein and the following: furan, furazan, thiophene, benzothiophene, phthalazine, pyrrole, oxazole, benzoxazole, 1,2,3-oxadiazole, 1,2,4-oxadiazole, thiazole, 1,2,3-thiadiazole, 1,2,4-thiadiazole, benzothiazole, imidazole, benzimidazole, indole, indazole, pyrazole, benzopyrazole, isoxazole, benzoisoxazole, isothiazole, triazole, benzotriazole, thiadiazole, tetrazole, pyridine, pyridazine, pyrimidine, pyrazine, purine, pteridine, quinoline, isoquinoline, quinazoline, quinoxaline, cinnoline and triazine. A heteroaryl group may be substituted or unsubstituted.

[0160] As used herein, “heterocyclyl” or “heteroalicyclyl” refers to three-, four-, five-, six-, seven-, eight-, nine-, ten-, up to 18-membered monocyclic, bicyclic, and tricyclic ring system wherein carbon atoms together with from 1 to 5 heteroatoms constitute said ring system. A heterocycle may optionally contain one or more unsaturated bonds situated in such a way, however, that a fully delocalized pi-electron system does not occur throughout all the rings. The heteroatom(s) is an element other than carbon including, but not limited to, oxygen, sulfur, and nitrogen. A heterocycle may further contain one or more carbonyl or thiocarbonyl functionalities, so as to make the definition include oxo-systems and thio-systems such as lactams, lactones, cyclic imides, cyclic thioimides and cyclic carbamates. When composed of two or more rings, the rings may be joined together in a fused fashion. Additionally, any nitrogens in a heterocyclyl may be quaternized. Heterocyclyl or heteroalicyclic groups may be unsubstituted or substituted. Examples of such

“heterocyclyl” or “heteroalicycyl” groups include, but are not limited to, those described herein and the following: 1,3-dioxin, 1,3-dioxane, 1,4-dioxane, 1,2-dioxolane, 1,3-dioxolane, 1,4-dioxolane, 1,3-oxathiane, 1,4-oxathiin, 1,3-oxathiolane, 1,3-dithiole, 1,3-dithiolane, 1,4-oxathiane, tetrahydro-1,4-thiazine, 1,3-thiazinane, 2H-1,2-oxazine, maleimide, succinimide, barbituric acid, thiobarbituric acid, dioxopiperazine, hydantoin, dihydrouacil, trioxane, hexahydro-1,3,5-triazine, imidazoline, imidazolidine, isoxazoline, isoxazolidine, oxazoline, oxazolidine, oxazolidinone, thiazoline, thiazolidine, morpholine, oxirane, piperidine N-Oxide, piperidine, piperazine, pyrrolidine, pyrrolidone, pyrrolidione, 4-piperidone, pyrazoline, pyrazolidine, 2-oxopyrrolidine, tetrahydropyran, 4H-pyran, tetrahydrothiopyran, thiamorpholine, thiamorpholine sulfoxide, thiamorpholine sulfone, and their benzo-fused analogs (e.g., benzimidazolidinone, tetrahydroquinoline, and 3,4-methylenedioxyphenyl).

[0161] As used herein, “aralkyl” and “aryl(alkyl)” refer to an aryl group connected, as a substituent, via a lower alkylene group. The lower alkylene and aryl group of an aralkyl may be substituted or unsubstituted. Examples include but are not limited to benzyl, 2-phenylalkyl, 3-phenylalkyl and naphthylalkyl.

[0162] As used herein, “heteroaralkyl” and “heteroaryl(alkyl)” refer to a heteroaryl group connected, as a substituent, via a lower alkylene group. The lower alkylene and heteroaryl group of heteroaralkyl may be substituted or unsubstituted. Examples include but are not limited to 2-thienylalkyl, 3-thienylalkyl, furylalkyl, thienylalkyl, pyrrolylalkyl, pyridylalkyl, isoxazolylalkyl, imidazolylalkyl and their benzo-fused analogs.

[0163] A “heteroalicycyl(alkyl)” and “heterocyclyl(alkyl)” refer to a heterocyclic or a heteroalicyclic group connected, as a substituent, via a lower alkylene group. The lower alkylene and heterocyclyl of a heteroalicycyl(alkyl) may be substituted or unsubstituted. Examples include but are not limited tetrahydro-2H-pyran-4-yl(methyl), piperidin-4-yl(ethyl), piperidin-4-yl(propyl), tetrahydro-2H-thiopyran-4-yl(methyl), and 1,3-thiazinan-4-yl(methyl).

[0164] “Lower alkylene groups” are straight-chained —CH.sub.2— tethering groups, forming bonds to connect molecular fragments via their terminal carbon atoms. Examples include but are not limited to methylene (—CH.sub.2—), ethylene (—CH.sub.2CH.sub.2—), propylene (—CH.sub.2CH.sub.2CH.sub.2—), and butylene (—CH.sub.2CH.sub.2CH.sub.2CH.sub.2—). A lower alkylene group can be substituted by replacing one or more hydrogen of the lower alkylene group with a substituent(s) listed under the definition of “substituted.”

[0165] As used herein, “alkoxy” refers to the formula —OR wherein R is an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl) is defined herein. A non-limiting list of alkoxys are methoxy, ethoxy, n-propoxy, 1-methylethoxy (isopropoxy), n-butoxy, iso-butoxy, sec-butoxy, tert-butoxy, phenoxy and benzoxy. An alkoxy may be substituted or unsubstituted.

[0166] As used herein, “acyl” refers to a hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl) connected, as substituents, via a carbonyl group. Examples include formyl, acetyl, propanoyl, benzoyl and acryl. An acyl may be substituted or unsubstituted.

[0167] As used herein, “acylalkyl” refers to an acyl connected, as a substituent, via a lower alkylene group. Examples include aryl-C(=O)—(CH.sub.2).sub.n— and heteroaryl-C(=O)—(CH.sub.2).sub.n—, where n is an integer in the range of 1 to 6.

[0168] As used herein, “alkoxyalkyl” refers to an alkoxy group connected, as a substituent, via a lower alkylene group. Examples include C.sub.1-4 alkyl-O—(CH.sub.2).sub.n—, wherein n is an integer in the range of 1 to 6.

[0169] As used herein, “aminoalkyl” refers to an optionally substituted amino group connected, as a substituent, via a lower alkylene group. Examples include H.sub.2N(CH.sub.2).sub.n—, wherein n is an integer in the range of 1 to 6.

[0170] As used herein, “hydroxyalkyl” refers to an alkyl group in which one or more of the

hydrogen atoms are replaced by a hydroxy group. Exemplary hydroxyalkyl groups include but are not limited to, 2-hydroxyethyl, 3-hydroxypropyl, 2-hydroxypropyl, and 2,2-dihydroxyethyl. A hydroxyalkyl may be substituted or unsubstituted.

[0171] As used herein, “haloalkyl” refers to an alkyl group in which one or more of the hydrogen atoms are replaced by a halogen (e.g., mono-haloalkyl, di-haloalkyl and tri-haloalkyl). Such groups include but are not limited to, chloromethyl, fluoromethyl, difluoromethyl, trifluoromethyl, chloro-fluoroalkyl, chloro-difluoroalkyl and 2-fluoroisobutyl. A haloalkyl may be substituted or unsubstituted.

[0172] As used herein, “haloalkoxy” refers to an alkoxy group in which one or more of the hydrogen atoms are replaced by a halogen (e.g., mono-haloalkoxy, di-haloalkoxy and tri-haloalkoxy). Such groups include but are not limited to, chloromethoxy, fluoromethoxy, difluoromethoxy, trifluoromethoxy, chloro-fluoroalkyl, chloro-difluoroalkoxy and 2-fluoroisobutoxy. A haloalkoxy may be substituted or unsubstituted.

[0173] A “sulfenyl” group refers to an “—SR” group in which R can be hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). A sulfenyl may be substituted or unsubstituted.

[0174] A “sulfinyl” group refers to an “—S(=O)—R” group in which R can be the same as defined with respect to sulfenyl. A sulfinyl may be substituted or unsubstituted.

[0175] A “sulfonyl” group refers to an “SO.sub.2R” group in which R can be the same as defined with respect to sulfenyl. A sulfonyl may be substituted or unsubstituted.

[0176] An “O-carboxy” group refers to a “RC(=O)O—” group in which R can be hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl), as defined herein. An O-carboxy may be substituted or unsubstituted.

[0177] The terms “ester” and “C-carboxy” refer to a “—C(=O)OR” group in which R can be the same as defined with respect to O-carboxy. An ester and C-carboxy may be substituted or unsubstituted.

[0178] A “thiocarbonyl” group refers to a “—C(=S)R” group in which R can be the same as defined with respect to O-carboxy. A thiocarbonyl may be substituted or unsubstituted.

[0179] A “trihalomethanesulfonyl” group refers to an “X.sub.3CSO.sub.2—” group wherein each X is a halogen.

[0180] A “trihalomethanesulfonamido” group refers to an “X.sub.3CS(O).sub.2N(R.sub.A)—” group wherein each X is a halogen, and R.sub.A hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl).

[0181] The term “amino” as used herein refers to a —NH.sub.2 group.

[0182] As used herein, the term “hydroxy” refers to a —OH group.

[0183] A “cyano” group refers to a “—CN” group.

[0184] The term “azido” as used herein refers to a —N.sub.3 group.

[0185] An “isocyanato” group refers to a “—NCO” group.

[0186] A “thiocyanato” group refers to a “—CNS” group.

[0187] An “isothiocyanato” group refers to an “—NCS” group.

[0188] A “carbonyl” group refers to a C=O group.

[0189] An “S-sulfonamido” group refers to a “—SO.sub.2N(R.sub.AR.sub.B)” group in which R.sub.A and R.sub.B can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An S-sulfonamido may be substituted or unsubstituted.

[0190] An “N-sulfonamido” group refers to a “RSO.sub.2N(R.sub.A)—” group in which R and R.sub.A can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or

heterocyclyl(alkyl). An N-sulfonamido may be substituted or unsubstituted.

[0191] An “O-carbamyl” group refers to a “—OC(=O)N(R.sub.AR.sub.B)” group in which R.sub.A and R.sub.B can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An O-carbamyl may be substituted or unsubstituted.

[0192] An “N-carbamyl” group refers to an “ROC(=O)N(R.sub.A)—” group in which R and R.sub.A can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An N-carbamyl may be substituted or unsubstituted.

[0193] An “O-thiocarbamyl” group refers to a “—OC(=S)—N(R.sub.AR.sub.B)” group in which R.sub.A and R.sub.B can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An O-thiocarbamyl may be substituted or unsubstituted.

[0194] An “N-thiocarbamyl” group refers to an “ROC(=S)N(R.sub.A)—” group in which R and R.sub.A can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An N-thiocarbamyl may be substituted or unsubstituted.

[0195] A “C-amido” group refers to a “—C(=O)N(R.sub.AR.sub.B)” group in which R.sub.A and R.sub.B can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). A C-amido may be substituted or unsubstituted.

[0196] An “N-amido” group refers to a “RC(=O)N(R.sub.A)—” group in which R and R.sub.A can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An N-amido may be substituted or unsubstituted.

[0197] A “urea” group refers to “N(R)—C(=O)—NR.sub.AR.sub.B group in which R can be hydrogen or an alkyl, and R.sub.A and R.sub.B can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). A urea may be substituted or unsubstituted.

[0198] The term “halogen atom” or “halogen” as used herein, means any one of the radio-stable atoms of column 7 of the Periodic Table of the Elements, such as, fluorine, chlorine, bromine and iodine.

[0199] As used herein,  custom-character indicates a single or double bond, unless stated otherwise.

[0200] Where the numbers of substituents is not specified (e.g. haloalkyl), there may be one or more substituents present. For example “haloalkyl” may include one or more of the same or different halogens. As another example, “C.sub.1-C.sub.3 alkoxyphenyl” may include one or more of the same or different alkoxy groups containing one, two or three atoms.

[0201] As used herein, the abbreviations for any protective groups, amino acids and other compounds, are, unless indicated otherwise, in accord with their common usage, recognized abbreviations, or the IUPAC-IUB Commission on Biochemical Nomenclature (See, Biochem. 11:942-944 (1972)).

[0202] In certain embodiments, “optically active” and “enantiomerically active” refer to a collection of molecules, which has an enantiomeric excess of no less than about 50%, no less than about 70%, no less than about 80%, no less than about 90%, no less than about 91%, no less than about 92%, no less than about 93%, no less than about 94%, no less than about 95%, no less than about 96%, no less than about 97%, no less than about 98%, no less than about 99%, no less than about 99.5%, or no less than about 99.8%. In certain embodiments, the compound comprises about 95% or more of the desired enantiomer and about 5% or less of the less preferred enantiomer based on the total weight of the two enantiomers in question.

[0203] In describing an optically active compound, the prefixes R and S are used to denote the absolute configuration of the optically active compound about its chiral center(s). The (+) and (−) are used to denote the optical rotation of an optically active compound, that is, the direction in which a plane of polarized light is rotated by the optically active compound. The (−) prefix indicates that an optically active compound is levorotatory, that is, the compound rotates the plane of polarized light to the left or counterclockwise. The (+) prefix indicates that an optically active compound is dextrorotatory, that is, the compound rotates the plane of polarized light to the right or clockwise. However, the sign of optical rotation, (+) and (−), is not related to the absolute configuration of a compound, R and S.

[0204] The term “isotopic variant” refers to a compound that contains an unnatural proportion of an isotope at one or more of the atoms that constitute such a compound. In certain embodiments, an “isotopic variant” of a compound contains unnatural proportions of one or more isotopes, including, but not limited to, hydrogen (¹H), deuterium (²H), tritium (³H), carbon-11 (¹¹C), carbon-12 (¹²C), carbon-13 (¹³C), carbon-14 (¹⁴C), nitrogen-13 (¹³N), nitrogen-14 (¹⁴N), nitrogen-15 (¹⁵N), oxygen-14 (¹⁴O), oxygen-15 (¹⁵O), oxygen-16 (¹⁶O), oxygen-17 (¹⁷O), oxygen-18 (¹⁸O), fluorine-17 (¹⁷F), fluorine-18 (¹⁸F), phosphorus-31 (³¹P), phosphorus-32 (³²P), phosphorus-33 (³³P), sulfur-32 (³²S), sulfur-33 (³³S), sulfur-34 (³⁴S), sulfur-35 (³⁵S), sulfur-36 (MS), chlorine-35 (³⁵Cl), chlorine-36 (³⁶Cl), chlorine-37 (³⁷Cl), bromine-79 (⁷⁹Br), bromine-81 (⁸¹Br), iodine-123 (¹²³I), iodine-125 (¹²⁵I), iodine-127 (¹²⁷I), iodine-129 (¹²⁹I), and iodine-131 (¹³¹I). In certain embodiments, an “isotopic variant” of a compound is in a stable form, that is, non-radioactive. In certain embodiments, an “isotopic variant” of a compound contains unnatural proportions of one or more isotopes, including, but not limited to, hydrogen (¹H), deuterium (²H), carbon-12 (¹²C), carbon-13 (¹³C), nitrogen-14 (¹⁴N), nitrogen-15 (¹⁵N), oxygen-16 (¹⁶O), oxygen-17 (¹⁷O), oxygen-18 (¹⁸O), fluorine-17 (¹⁷F), phosphorus-31 (³¹P), sulfur-32 (³²S), sulfur-33 (³³S), sulfur-34 (³⁴S), sulfur-36 (³⁶S), chlorine-35 (³⁵Cl), chlorine-37 (³⁷Cl), bromine-79 (⁷⁹Br), bromine-81 (⁸¹Br), and iodine-127 (¹²⁷I). In certain embodiments, an “isotopic variant” of a compound is in an unstable form, that is, radioactive. In certain embodiments, an “isotopic variant” of a compound contains unnatural proportions of one or more isotopes, including, but not limited to, tritium (³H), carbon-11 (¹¹C), carbon-14 (¹⁴C), nitrogen-13 (¹³N), oxygen-14 (¹⁴O), oxygen-15 (¹⁵O), fluorine-18 (¹⁸F), phosphorus-32 (³²P), phosphorus-33 (³³P), sulfur-35 (³⁵S), chlorine-36 (³⁶Cl), iodine-123 (¹²³I), iodine-125 (¹²⁵I), iodine-129 (¹²⁹I), and iodine-131 (¹³¹I). It will be understood that, in a compound as provided herein, any hydrogen can be ²H, for example, or any carbon can be ¹³C, for example, or any nitrogen can be ¹⁵N, for example, or any oxygen can be ¹⁸O, for example, where feasible according to the judgment of one of skill. In certain embodiments, an “isotopic variant” of a compound contains unnatural proportions of deuterium (D).

[0205] The term “solvate” refers to a complex or aggregate formed by one or more molecules of a solute, e.g., a compound provided herein, and one or more molecules of a solvent, which present in a stoichiometric or non-stoichiometric amount. Suitable solvents include, but are not limited to, water, methanol, ethanol, n-propanol, isopropanol, and acetic acid. In certain embodiments, the solvent is pharmaceutically acceptable. In one embodiment, the complex or aggregate is in a crystalline form. In another embodiment, the complex or aggregate is in a noncrystalline form. Where the solvent is water, the solvate is a hydrate. Examples of hydrates include, but are not limited to, a hemihydrate, monohydrate, dihydrate, trihydrate, tetrahydrate, and pentahydrate.

[0206] The phrase “an enantiomer, a mixture of enantiomers, a mixture of two or more diastereomers, or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate,

hydrate, or prodrug thereof” has the same meaning as the phrase “(i) an enantiomer, a mixture of enantiomers, a mixture of two or more diastereomers, or an isotopic variant of the compound referenced therein; (ii) a pharmaceutically acceptable salt, solvate, hydrate, or prodrug of the compound referenced therein; or (iii) a pharmaceutically acceptable salt, solvate, hydrate, or prodrug of an enantiomer, a mixture of enantiomers, a mixture of two or more diastereomers, or an isotopic variant of the compound referenced therein.”

5.3.2. Stem Cell Mobilizing Compounds

[0207] In certain aspects, the stem cell mobilizing factor is a compound having Formula (I), (I-A), (I-B), (I-C), or (I-D), as described below.

Formula (I)

[0208] Some embodiments disclosed herein relate to a compound of Formula (I), or a pharmaceutically acceptable salt thereof, having the structure:

##STR00002##

wherein: each  custom-character can independently represent a single bond or a double bond; R^{sup.J} can be selected from the group consisting of —NR^{sup.a}R^{sup.b}, —OR^{sup.b}, and =O; wherein if R^{sup.J} is =O, then  custom-character joining G and J represents a single bond and G is N and the N is substituted with R^{sup.G}; otherwise  custom-character joining G and J represents a double bond and G is N; R^{sup.a} can be hydrogen or C_{sub.1-4} alkyl; R^{sup.b} can be R^{sup.c} or —(C_{sub.1-4} alkyl)-R^{sup.c}; R^{sup.c} can be selected from the group consisting of: —OH, —O(C_{sub.1-4} alkyl), —O(C_{sub.1-4} haloalkyl); —C(=O)NH_{sub.2}; unsubstituted C_{sub.6-10} aryl; substituted C_{sub.6-10} aryl; unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R^{sup.c} moiety indicated as substituted can be substituted with one or more substituents E, wherein each E can be independently selected from the group consisting of: —OH, C_{sub.1-4} alkyl, C_{sub.1-4} haloalkyl, —O(C_{sub.1-4} alkyl), and —O(C_{sub.1-4} haloalkyl); R^{sup.K} can be selected from the group consisting of: hydrogen, unsubstituted C_{sub.1-6} alkyl; substituted C_{sub.1-6} alkyl; —NH(C_{sub.1-4} alkyl); —N(C_{sub.1-4} alkyl)_{sub.2}, unsubstituted C_{sub.6-10} aryl; substituted C_{sub.6-10} aryl; unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R^{sup.K} moiety indicated as substituted can be substituted with one or more substituents Q, wherein each Q is independently selected from the group consisting of: —OH, C_{sub.1-4} alkyl, C_{sub.1-4} haloalkyl, halo, cyano, —O—(C_{sub.1-4} alkyl), and —O—(C_{sub.1-4} haloalkyl); R^{sup.G} can be selected from the group consisting of hydrogen, C_{sub.1-4} alkyl, and —(C_{sub.1-4} alkyl)-C(=O)NH_{sub.2}; R^{sup.Y} and R^{sub.Z} can each independently be absent or be selected from the group consisting of: hydrogen, halo, C_{sub.1-6} alkyl, —OH, —O—(C_{sub.1-4} alkyl), —NH(C_{sub.1-4} alkyl), and —N(C_{sub.1-4} alkyl)_{sub.2}; or R^{sup.Y} and R^{sup.Z} taken together with the atoms to which they are attached can joined together to form a ring selected from:

##STR00003##

wherein said ring can be optionally substituted with one, two, or three groups independently selected from C_{sub.1-4} alkyl, C_{sub.1-4} haloalkyl, halo, cyano, —OH, —O—(C_{sub.1-4} alkyl), —N(C_{sub.1-4} alkyl)_{sub.2}, unsubstituted C_{sub.6-10} aryl, C_{sub.6-10} aryl substituted with 1-5 halo atoms, and —O—(C_{sub.1-4} haloalkyl); and wherein if R^{sup.Y} and R^{sup.Z} taken together forms

##STR00004##

then R^{sup.J} can be —OR^{sup.b} or =O; R^{sup.d} can be hydrogen or C_{sub.1-4} alkyl; R^{sup.m} can be selected from the group consisting of C_{sub.1-4} alkyl, halo, and cyano; J can be C; and X, Y, and Z can each be independently N or C, wherein the valency of any carbon atom is filled as

needed with hydrogen atoms.

[0209] In some embodiments,  custom-character can represent a single bond. In other embodiments,  custom-character can represent a double bond. In some embodiments,  custom-character joining Y and Z can represent a single bond. In other embodiments,  custom-character joining Y and Z can represent a double bond. In some embodiments, when  custom-character joining G and J represents a single bond, G can be N and the N is substituted with R.sup.G. In other embodiments, when  custom-character joining G and J represents a double bond, G can be N. In some embodiments, when  custom-character joining G and J represents a double bond, then  custom-character joining J and R.sup.J can be a single bond. In some embodiments, when  custom-character joining G and J represents a double bond, then  custom-character joining J and R.sup.J can not be a double bond. In some embodiments, when  custom-character joining J and R.sup.J represents a double bond, then  custom-character joining G and J can be a single bond. In some embodiments, when  custom-character joining J and R.sup.J represents a double bond, then  custom-character joining G and J can not be a double bond.

[0210] In some embodiments, R.sup.J can be —NR.sup.aR.sup.b. In other embodiments, R.sup.J can be —OR.sup.b. In still other embodiments, R.sup.J can be =O. In some embodiments, when R.sup.J is =O, then  custom-character joining G and J represents a single bond and G is N and the N is substituted with R.sup.G. In some embodiments, R.sup.G is —CH.sub.2CH.sub.2—C(=O)NH.sub.2.

[0211] In some embodiments, R.sup.a can be hydrogen. In some embodiments, R.sup.a can be C.sub.1-C.sub.4 alkyl. For example, R.sup.a can be methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl or tert-butyl.

[0212] In some embodiments, R.sup.b can be R.sup.c. In some embodiments, R.sup.b can be —(C.sub.1-C.sub.4 alkyl)-R.sup.c. For example, R.sup.b can be —CH.sub.2—R.sup.c, —CH.sub.2CH.sub.2—R.sup.c, —CH.sub.2CH.sub.2CH.sub.2—R.sup.c, or —CH.sub.2CH.sub.2CH.sub.2CH.sub.2—R.sup.c. In some embodiments, when R.sup.b is —CH.sub.2CH.sub.2—R.sup.c, R.sup.c can be —O(C.sub.1-C.sub.4 alkyl). In other embodiments, when R.sup.b is —CH.sub.2CH.sub.2—R.sup.c, R.sup.c can be —O(C.sub.1-C.sub.4 haloalkyl). In still other embodiments, when R.sup.b is —CH.sub.2CH.sub.2—R.sup.c, R can be —C(=O)NH.sub.2.

[0213] In some embodiments, R.sup.c can be —OH. In some embodiments, R.sup.c can be —O(C.sub.1-C.sub.4 alkyl). In some embodiments, R.sup.c can be —O(C.sub.1-C.sub.4 haloalkyl). In some embodiments, R.sup.c can be —C(=O)NH.sub.2. In some embodiments, R.sup.c can be unsubstituted C.sub.6-10 aryl. In some embodiments, R.sup.c can be substituted C.sub.6-10 aryl. In some embodiments, R.sup.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S. In some embodiments, R.sup.c can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S. In some embodiments, when a R.sup.c moiety is indicated as substituted, the moiety can be substituted with one or more, for example, one, two, three, or four substituents E. In some embodiments, E can be —OH. In some embodiments, E can be C.sub.1-C.sub.4 alkyl. In some embodiments, E can be C.sub.1-C.sub.4 haloalkyl. In some embodiments, E can be —O(C.sub.1-C.sub.4 alkyl). In some embodiments, E can be —O(C.sub.1-C.sub.4 haloalkyl).

[0214] In some embodiments, when R.sup.b is —CH.sub.2CH.sub.2—R.sup.c, R.sup.c can be unsubstituted C.sub.6-10 aryl. In other embodiments, when R.sup.b is —CH.sub.2CH.sub.2—R.sup.c, R.sup.c can be substituted C.sub.6-10 aryl. In still other embodiments, when R.sup.b is —CH.sub.2CH.sub.2—R.sup.c, R.sup.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S. In yet still other embodiments, R.sup.b can be —(C.sub.1-C.sub.4 alkyl)-R.sup.c and R.sup.c can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S. When a R.sup.c moiety is indicated as substituted, the moiety can be substituted with one or more, for

example, one, two, three, or four substituents E. In some embodiments, E can be —OH. In other embodiments, E can be C.sub.1-C.sub.4 alkyl. In still other embodiments, E can be C.sub.1-C.sub.4 haloalkyl. In still other embodiments, E can be —O(C.sub.1-C.sub.4 alkyl). In still other embodiments, E can be —O(C.sub.1-C.sub.4 haloalkyl).

[0215] In some embodiments, when R^{sup.b} is —CH₂CH₂—R^{sup.c}, R^{sup.c} can be phenyl. In other embodiments, when R^{sup.b} is —CH₂CH₂—R^{sup.c}, R^{sup.c} can be naphthyl. In still other embodiments, when R^{sup.b} is —CH₂CH₂—R^{sup.c}, R^{sup.c} can be hydroxyphenyl. In still other embodiments, when R^{sup.b} is —CH₂CH₂—R^{sup.c}, R^{sup.c} can be indolyl.

[0216] In some embodiments, R^{sup.K} can be hydrogen. In other embodiments, R^{sup.K} can be unsubstituted C.sub.1-6 alkyl. For example, in some embodiments, R^{sup.K} can be methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, pentyl (branched and straight-chained), or hexyl (branched and straight-chained). In other embodiments, R^{sup.K} can be substituted C.sub.1-6 alkyl. In other embodiments, R^{sup.K} can be —NH(C.sub.1-4 alkyl). For example, in some embodiments, R^{sup.K} can be —NH(CH₃), —NH(CH₂CH₃), —NH(isopropyl), or —NH(sec-butyl). In other embodiments, R^{sup.K} can be —N(C.sub.1-4 alkyl).sub.2.

[0217] In some embodiments, R^{sup.K} can be unsubstituted C.sub.6-10 aryl. In other embodiments, R^{sup.K} can be substituted C.sub.6-10 aryl. In other embodiments, R^{sup.K} can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S. In other embodiments, R^{sup.K} can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S. When a R^{sup.K} moiety is indicated as substituted, the moiety can be substituted with one or more, for example, one, two, three, or four substituents Q. In some embodiments, Q can be —OH. In other embodiments, Q can be C.sub.1-4 alkyl. In still other embodiments, Q can be C.sub.1-4 haloalkyl. In still other embodiments, Q can be halo. In still other embodiments, Q can be cyano. In still other embodiments, Q can be —O—(C.sub.1-4 alkyl). In still other embodiments, Q can be —O—(C.sub.1-4 haloalkyl).

[0218] In some embodiments, R^{sup.K} can be phenyl or naphthyl. In other embodiments, R^{sup.K} can be benzothiophenyl. In other embodiments, R^{sup.K} can be benzothiophenyl. In other embodiments, R^{sup.K} can be benzothiophenyl. In still other embodiments, R^{sup.K} can be pyridinyl. In yet still other embodiments, R^{sup.K} can be pyridinyl substituted with one or more substituents Q. For example, R^{sup.K} can be methylpyridinyl, ethylpyridinyl cyanopyridinyl, chloropyridinyl, fluoropyridinyl, or bromopyridinyl.

[0219] In some embodiments, R^{sup.G} can be hydrogen. In some embodiments, R^{sup.G} can be C.sub.1-4 alkyl. In some embodiments, R^{sup.G} can be —(C.sub.1-4 alkyl)-C(=O)NH₂.

[0220] In some embodiments, R^{sup.Y} and R^{sup.Z} can independently be absent. In other embodiments, R^{sup.Y} and R^{sup.Z} can independently be hydrogen. In other embodiments, R^{sup.Y} and R^{sup.Z} can independently be halo. In other embodiments, R^{sup.Y} and R^{sup.Z} can independently be C.sub.1-6 alkyl. In other embodiments, R^{sup.Y} and R^{sup.Z} can independently be —OH. In still other embodiments, R^{sup.Y} and R^{sup.Z} can independently be —O—(C.sub.1-4 alkyl). In other embodiments, R^{sup.Y} and R^{sup.Z} can independently be —NH(C.sub.1-4 alkyl). For example, R^{sup.Y} and R^{sup.Z} can independently be —NH(CH₃), —NH(CH₂CH₃), —NH(isopropyl), or —NH(sec-butyl). In other embodiments, R^{sup.Y} and R^{sup.Z} can independently be —N(C.sub.1-4 alkyl).sub.2.

[0221] In some embodiments, R^{sup.Y} and R^{sup.Z} taken together with the atoms to which they are attached can be joined together to form a ring. In some embodiments, R^{sup.Y} and R^{sup.Z} taken together with the atoms to which they are attached can be joined together to form

##STR00005##

In other embodiments, R^{sup.Y} and R^{sup.Z} taken together with the atoms to which they are attached can be joined together to form

##STR00006##

In other embodiments, R.sup.Y and R.sup.Z taken together with the atoms to which they are attached can be joined together to form

##STR00007##

In still other embodiments, R.sup.Y and R.sup.Z taken together with the atoms to which they are attached can be joined together to form

##STR00008##

In yet still other embodiments, R.sup.Y and R.sup.Z taken together with the atoms to which they are attached can be joined together to form

##STR00009##

In other embodiments, R.sup.Y and R.sup.Z taken together with the atoms to which they are attached can be joined together to form

##STR00010##

In yet other embodiments, R.sup.Y and R.sup.Z taken together with the atoms to which they are attached can be joined together to form

##STR00011##

In yet still other embodiments, R.sup.Y and R.sup.Z taken together with the atoms to which they are attached can be joined together to form

##STR00012##

In other embodiments, R.sup.Y and R.sup.Z taken together with the atoms to which they are attached can be joined together to form

##STR00013##

In still other embodiments, R.sup.Y and R.sup.Z taken together with the atoms to which they are attached can be joined together to form and

##STR00014##

In some embodiments, when R.sup.Y and R.sup.Z taken together with the atoms to which they are attached can be joined together to form a ring, the ring can be substituted with one, two, or three groups independently selected from C.sub.1-C.sub.4 alkyl, —N(C.sub.1-C.sub.4 alkyl).sub.2, cyano, unsubstituted phenyl, and phenyl substituted with 1-5 halo atoms.

[0222] In some embodiments, when R.sup.Y and R.sup.Z taken together forms

##STR00015##

then R.sup.J can be —OR.sup.b or =O.

[0223] In some embodiments, R.sup.Y and R.sup.Z taken together with the atoms to which they are attached can be joined together to form

##STR00016##

In other embodiments, R.sup.Y and R.sup.Z taken together with the atoms to which they are attached can be joined together to form

##STR00017##

In other embodiments, R.sup.Y and R.sup.Z taken together with the atoms to which they are attached can be joined together to form

##STR00018##

In other embodiments, R.sup.Y and R.sup.Z taken together with the atoms to which they are attached can be joined together to form

##STR00019##

In other embodiments, R.sup.Y and R.sup.Z taken together with the atoms to which they are attached can be joined together to form

##STR00020##

In other embodiments, R.sup.Y and R.sup.Z taken together with the atoms to which they are attached can be joined together to form

##STR00021##

In other embodiments, R.sup.Y and R.sup.Z taken together with the atoms to which they are

attached can be joined together to form

##STR00022##

In other embodiments, R^{sup}.Y and R^{sup}.Z taken together with the atoms to which they are attached can be joined together to form

##STR00023##

In other embodiments, R^{sup}.Y and R^{sup}.Z taken together with the atoms to which they are attached can be joined together to form

##STR00024##

In other embodiments, R^{sup}.Y and R^{sup}.Z taken together with the atoms to which they are attached can be joined together to form

##STR00025##

In some embodiments, when R^{sup}.Y and R^{sup}.Z taken together with the atoms to which they are attached can be joined together to form a ring, the ring can be substituted with one, two, or three groups independently selected from C_{sub.1}-C_{sub.4} alkyl, —N(C_{sub.1}-C_{sub.4} alkyl)_{sub.2}, cyano, unsubstituted phenyl, and phenyl substituted with 1-5 halo atoms. In some embodiments, R^{sup}.Y and R^{sup}.Z taken together with the atoms to which they are attached can be

##STR00026##

In other embodiments, R^{sup}.Y and R^{sup}.Z taken together with the atoms to which they are attached can be

##STR00027##

In still other embodiments, R^{sup}.Y and R^{sup}.Z taken together with the atoms to which they are attached can be

##STR00028##

In yet still other embodiments, R^{sup}.Y and R^{sup}.Z taken together with the atoms to which they are attached can be

##STR00029##

In other embodiments, R^{sup}.Y and R^{sup}.Z taken together with the atoms to which they are attached can be

##STR00030##

[0224] In some embodiments, R^{sup}.d can be hydrogen. In other embodiments, R^{sup}.d can be C_{sub.1}-C_{sub.4} alkyl. For example R^{sup}.d can be methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl or tert-butyl. In still other embodiments, R^{sup}.d can be halo. In other embodiments, R^{sup}.d can be cyano.

[0225] In some embodiments, R^{sup}.m can be hydrogen. In other embodiments, R^{sup}.m can be C_{sub.1}-C_{sub.4} alkyl. For example R^{sup}.m can be methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl or tert-butyl. In still other embodiments, R^{sup}.m can be halo. For example, R^{sup}.m can be fluoro, chloro, bromo, or iodo. In other embodiments, R^{sup}.m can be cyano.

[0226] In some embodiments, X, Y, and Z can each be independently N or C, wherein the valency of any carbon atom is filled as needed with hydrogen atoms. In some embodiments, X can be N, Y can be N, and Z can be N. In other embodiments, X can be N, Y can be N, and Z can be CH. In some embodiments, X can be N, Y can be CH, and Z can be N. In still other embodiments, X can be CH, Y can be N, and Z can be N. In yet still other embodiments, X can be CH, Y can be CH, and Z can be N. In other embodiments, X can be CH, Y can be N, and Z can be CH. In yet other embodiments, X can be N, Y can be CH, and Z can be CH. In other embodiments, X can be CH, Y can be CH, and Z can be CH.

[0227] In some embodiments, R^{sup}.a can be hydrogen; R^{sup}.b can be —(C_{sub.1}-C_{sub.4} alkyl)-R^{sup}.c; R^{sup}.c can be selected from the group consisting of: —C(=O)NH_{sub.2}; unsubstituted C_{sub.6-10} aryl; substituted C_{sub.6-10} aryl; unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R^{sup}.c

moiety indicated as substituted is substituted with one or more substituents E, wherein each E can be independently selected from the group consisting of: —OH, C.sub.1-C.sub.4 alkyl, C.sub.1-C.sub.4 haloalkyl, —O(C.sub.1-C.sub.4 alkyl), and —O(C.sub.1-C.sub.4 haloalkyl); R.sup.K can be selected from the group consisting of: hydrogen, unsubstituted C.sub.1-6 alkyl; —NH(C.sub.1-4 alkyl); —N(C.sub.1-4 alkyl).sub.2, unsubstituted C.sub.6-10 aryl; substituted C.sub.6-10 aryl; unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.K moiety indicated as substituted is substituted with one or more substituents Q, wherein each Q can be independently selected from the group consisting of: —OH, C.sub.1-4 alkyl, C.sub.1-4 haloalkyl, halo, cyano, —O—(C.sub.1-4 alkyl), and —O—(C.sub.1-4 haloalkyl); R.sup.G can be —(C.sub.1-4 alkyl)-C(=O)NH.sub.2; R.sup.Y and R.sup.Z can each be independently absent or be selected from the group consisting of: hydrogen, C.sub.1-6 alkyl, and —NH(C.sub.1-4 alkyl); or R.sup.Y and R.sup.Z taken together with the atoms to which they are attached can be joined together to form a ring selected from:

##STR00031##

wherein said ring can be optionally substituted with one, two, or three groups independently selected from C.sub.1-4 alkyl, C.sub.1-4 haloalkyl, halo, cyano, —OH, —O—(C.sub.1-4 alkyl), —N(C.sub.1-4 alkyl).sub.2, unsubstituted C.sub.6-C.sub.10 aryl, C.sub.6-C.sub.10 aryl substituted with 1-5 halo atoms, and —O—(C.sub.1-4 haloalkyl); R.sup.d can be C.sub.1-C.sub.4 alkyl; R.sup.m can be cyano; and X, Y, and Z can each be independently N or C, wherein the valency of any carbon atom is filled as needed with hydrogen atoms.

[0228] In some embodiments, R.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be selected from the group consisting of: unsubstituted phenyl, substituted phenyl, indolyl, and —C(=O)NH.sub.2; R.sup.K can be selected from the group consisting of: hydrogen, methyl, substituted pyridinyl, unsubstituted benzothiophenyl, and —NH(C.sub.1-C.sub.4 alkyl); R.sup.G can be —CH.sub.2CH.sub.2—C(=O)NH.sub.2; R.sup.Y can be —NH(C.sub.1-C.sub.4 alkyl); R.sup.Z can be absent or hydrogen; or R.sup.Y and R.sup.Z taken together with the atoms to which they are attached can be joined together to form a ring selected from:

##STR00032##

wherein said ring can be optionally substituted with one, two, or three groups independently selected from C.sub.1-C.sub.4 alkyl, —N(C.sub.1-C.sub.4 alkyl).sub.2, cyano, unsubstituted phenyl, and phenyl substituted with 1-5 halo atoms; R.sup.d can be C.sub.1-C.sub.4 alkyl; R.sup.m can be cyano; and X can be N or CH.

[0229] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N;  joining G and J can be a double bond; R.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; or R.sup.c can be substituted C.sub.6-10 aryl, substituted with one or more E, wherein E is —OH; R.sup.K can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; or R.sup.K can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; substituted with one or more Q, wherein Q can be selected from cyano, halo, or C.sub.1-C.sub.4 alkyl; R.sup.Y and R.sup.Z taken together can be

##STR00033##

[0230] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N;  joining G and J can be a double bond; R.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; or R.sup.c can be substituted C.sub.6-10 aryl, substituted with one or more E, wherein E is —OH; R.sup.K can be hydrogen, C.sub.1-4 alkyl, or

unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and R.sup.Y and R.sup.Z taken together can be

##STR00034##

[0231] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N;  joining G and J can be a double bond; R.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; or R.sup.c can be substituted C.sub.6-10 aryl, substituted with one or more E, wherein E is —OH; R.sup.K can be hydrogen, C.sub.1-4 alkyl, or unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and R.sup.Y and R.sup.Z taken together can be

##STR00035##

[0232] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N;  joining G and J can be a double bond, R.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be substituted C.sub.6-10 aryl; substituted with one or more E, wherein E can be

—OH; R.sup.K can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.Y can be —NH(C.sub.1-4 alkyl); R.sup.Z can be hydrogen; J can be C; X can be N; Y can be C; Z can be C; and  joining Y and Z can be a double bond. In some embodiments, the compound of Formula (I) can be 4-(2-((2-(benzo[b]thiophen-3-yl)-6-(isopropylamino)pyrimidin-4-yl)amino)ethyl)phenol.

[0233] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N;  joining G and J can be a double bond; R.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c, R.sup.c can be substituted C.sub.6-10 aryl, substituted with one or more E, wherein E can be

—OH; R.sup.K can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.Y and R.sup.Z taken together is

##STR00036##

wherein the ring is substituted with C.sub.1-C.sub.4 alkyl; J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I) can be 4-(2-((2-(benzo[b]thiophen-3-yl)-7-isopropylthieno[3,2-d]pyrimidin-4-yl)amino)ethyl)phenol.

[0234] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N;  joining G and J can be a double bond; R.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c, R.sup.c can be substituted C.sub.6-10 aryl, substituted with one or more E, wherein E can be —OH; R.sup.K can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.Y and R.sup.Z taken together is

##STR00037##

R.sup.d; can be C.sub.1-C.sub.4 alkyl; J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I) can be 4-(2-((2-(benzo[b]thiophen-3-yl)-7-isopropyl-6,7-dihydro-5H-pyrrolo[2,3-d]pyrimidin-4-yl)amino)ethyl)phenol.

[0235] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N;  joining G and J can be a double bond; R.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c, R.sup.c can be substituted C.sub.6-10 aryl, substituted with one or more E, wherein E can be —OH; R.sup.K can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.Y and R.sup.Z taken together is

##STR00038##

R.sup.d can be C.sub.1-C.sub.4 alkyl; J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I) can be 2-(benzo[b]thiophen-3-yl)-4-((4-hydroxyphenethyl)amino)-7-isopropyl-5,7-dihydro-6H-pyrrolo[2,3-d]pyrimidin-6-one.

[0236] In some embodiments, when R^{sup}.J is —OR^{sup}.b; G can be N;  joining G and J can be a double bond; R^{sup}.b can be —CH₂CH₂—R^{sup}.c; R^{sup}.c can be —C(=O)NH₂; R^{sup}.K can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R^{sup}.Y and R^{sup}.Z taken together can be

##STR00039##

R^{sup}.d can be C₁₋₄ alkyl; J can be C; X can be N; Y can be C; and Z is C. In some embodiments, the compound of Formula (I) can be 3-((2-(benzo[b]thiophen-3-yl)-9-isopropyl-9H-purin-6-yl)oxy)propanamide.

[0237] In some embodiments, when R^{sup}.J is —NR^{sup}.aR^{sup}.b; G can be N;  joining G and J can be a double bond; R^{sup}.b can be —CH₂CH₂—R^{sup}.c; R^{sup}.c can be substituted C₆₋₁₀ aryl, substituted with one or more E, wherein E is —OH; R^{sup}.K is unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R^{sup}.Y and R^{sup}.Z taken together can be

##STR00040##

wherein said ring is substituted with —N(C₁₋₄ alkyl)₂; J can be C; X can be N; Y can be C; and Z is C. In some embodiments, the compound of Formula (I) can be 4-(2-((2-(benzo[b]thiophen-3-yl)-8-(dimethylamino)pyrimido[5,4-d]pyrimidin-4-yl)amino)ethyl)phenol.

[0238] In some embodiments, when R^{sup}.J is —NR^{sup}.aR^{sup}.b; G can be N;  joining G and J can be a double bond; R^{sup}.a can be hydrogen; R^{sup}.b can be —CH₂CH₂—R^{sup}.c; R^{sup}.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R^{sup}.K can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R^{sup}.K moiety indicated as substituted is substituted with one or more Q, wherein Q is cyano; R^{sup}.Y can be —NH(C₁₋₄ alkyl); R^{sup}.Z can be absent; J can be C; X can be C; Y can be C; Z can be N; and  joining Y and Z can be a double bond. In some embodiments, the compound of Formula (I) can be 5-(2-((2-(1H-indol-3-yl)ethyl)amino)-6-(sec-butylamino)pyrimidin-4-yl)nicotinonitrile.

[0239] In some embodiments, when R^{sup}.J is —NR^{sup}.aR^{sup}.b; G can be N;  joining G and J can be a double bond; R^{sup}.a can be hydrogen; R^{sup}.b can be —CH₂CH₂—R^{sup}.c; R^{sup}.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R^{sup}.K can be unsubstituted C₁₋₆ alkyl; R^{sup}.Y and R^{sup}.Z taken together can

##STR00041##

wherein the ring is substituted with unsubstituted C₆₋₁₀ aryl; J can be C; X can be N; Y can be C; Z can be C. In some embodiments, the compound of Formula (I) can be N-(2-(1H-indol-3-yl)ethyl)-2-methyl-6-phenylthieno[2,3-d]pyrimidin-4-amine

[0240] In some embodiments, when R^{sup}.J can be —NR^{sup}.aR^{sup}.b; G can be N;  joining G and J can be a double bond; R^{sup}.a can be hydrogen; R^{sup}.b can be —CH₂CH₂—R^{sup}.c; R^{sup}.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R^{sup}.K can be hydrogen; R^{sup}.Y and R^{sup}.Z taken together can be

##STR00042##

wherein the ring is substituted with substituted C₆₋₁₀ aryl; J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I) can be N-(2-(1H-indol-3-yl)ethyl)-6-(4-fluorophenyl)thieno[2,3-d]pyrimidin-4-amine

[0241] In some embodiments, when R^{sup}.J is =O; G can be N substituted with R^{sup}.G;  joining G and J can be a single bond; R^{sup}.G can be —(C₁₋₄ alkyl)-C(=O)NH₂; R^{sup}.K can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R^{sup}.Y and R^{sup}.Z taken together can be

##STR00043##

R.sup.d can be C.sub.1-C.sub.4 alkyl; J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I) can be 3-(2-(benzo[b]thiophen-3-yl)-9-isopropyl-6-oxo-6,9-dihydro-1H-purin-1-yl)propanamide.

[0242] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N;  custom-character joining G and J can be a double bond R.sup.a can be hydrogen R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.K can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.K moiety indicated as substituted is substituted with one or more Q, wherein Q can be halo; R.sup.Y and R.sup.Z taken together can be

##STR00044##

J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I) can be N-(2-(1H-indol-3-yl)ethyl)-2-(5-fluoropyridin-3-yl)quinazolin-4-amine.

[0243] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G is N;  custom-character joining G and J can be a double bond; R.sup.a can be hydrogen R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.K can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.K moiety indicated as substituted is substituted with one or more Q, wherein Q can be cyano; R.sup.Y and R.sup.Z taken together is

##STR00045##

J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I) can be 5-(4-((2-(1H-indol-3-yl)ethyl)amino)quinazolin-2-yl)nicotinonitrile.

[0244] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N;  custom-character joining G and J can be a double bond; R.sup.a can be hydrogen R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.K can be —NH(C.sub.1-4 alkyl); R.sup.Y and R.sup.Z taken together can be

##STR00046##

J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I) can be N.sup.4-(2-(1H-indol-3-yl)ethyl)-N.sup.2-(sec-butyl)quinazoline-2,4-diamine.

[0245] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N;  custom-character joining G and J can be a double bond; R.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be substituted C.sub.6-10 aryl, substituted with one or more E, wherein E is —OH; R.sup.K can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.Y and R.sup.Z taken together can be

##STR00047##

wherein the ring is substituted with cyano; R.sup.d can be C.sub.1-C.sub.4 alkyl; J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I) can be 2-(benzo[b]thiophen-3-yl)-4-((4-hydroxyphenethyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-5-carbonitrile.

[0246] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N;  custom-character joining G and J can be a double bond; R.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.K can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.Y and R.sup.Z taken together can be

##STR00048##

wherein the ring is substituted with C.sub.1-4 alkyl; J can be C; X can be C; Y can be N; and Z can be C; wherein the valency of any carbon atom is filled as needed with hydrogen atoms. In some embodiments, the compound of Formula (I) can be N-(2-(1H-indol-3-yl)ethyl)-6-(benzo[b]thiophen-3-yl)-3-isopropylimidazo[1,5-a]pyrazin-8-amine

[0247] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N;  joining G and J can be a double bond; R.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be substituted C.sub.6-10 aryl, substituted with one or more E, wherein E is —OH; R.sup.K can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R and R.sup.Z taken together can be

##STR00049## wherein the ring can be substituted with C.sub.1-4 alkyl; J can be C; X can be C; Y can be N; and Z can be C; wherein the valency of any carbon atom is filled as needed with hydrogen atoms. In some embodiments, the compound of Formula (I) can be 4-(2-((6-(benzo[b]thiophen-3-yl)-3-isopropylimidazo[1,5-a]pyrazin-8-yl)amino)ethyl)phenol.

[0248] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N;  joining G and J represents a double bond; R.sup.a can be hydrogen R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.K can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.K moiety indicated as substituted is substituted with one or more Q, wherein Q is cyano; R.sup.Y and R.sup.Z taken together is

##STR00050##

wherein the ring is substituted with C.sub.1-C.sub.4 alkyl; J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I) can be 5-(4-((2-(1H-indol-3-yl)ethyl)amino)-7-isopropylthieno[3,2-d]pyrimidin-2-yl)nicotinonitrile.

[0249] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N;  joining G and J represents a double bond; R.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.K can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.K moiety indicated as substituted is substituted with one or more Q, wherein Q is halo; R.sup.Y and R.sup.Z taken together can be

##STR00051##

wherein the ring is substituted with C.sub.1-C.sub.4 alkyl; J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I) can be N-(2-(1H-indol-3-yl)ethyl)-2-(5-fluoropyridin-3-yl)-7-isopropylthieno[3,2-d]pyrimidin-4-amine.

[0250] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N;  joining G and J can be a double bond; R.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.K can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.K moiety indicated as substituted is substituted with one or more Q, wherein Q is halo; R.sup.Y and R.sup.Z taken together can be

##STR00052##

J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I) can be N-(2-(1H-indol-3-yl)ethyl)-2-(5-fluoropyridin-3-yl)furo[3,2-d]pyrimidin-4-amine.

[0251] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N;  joining G and J can be a double bond; R.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be unsubstituted five- to ten-membered heteroaryl

having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.K can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.K moiety indicated as substituted is substituted with one or more Q, wherein Q is C.sub.1-C.sub.4 alkyl; R.sup.Y and R.sup.Z taken together can be

##STR00053##

J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I) can be N-(2-(1H-indol-3-yl)ethyl)-2-(5-methylpyridin-3-yl)furo[3,2-d]pyrimidin-4-amine. PGP-ART

[0252] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N;  custom-character joining G and J can be a double bond; R.sup.a can be hydrogen; R.sup.b can be —

CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.K can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.K moiety indicated as substituted is substituted with one or more Q, wherein Q is C.sub.1-C.sub.4 alkyl; R.sup.Y and R.sup.Z taken together can be

##STR00054##

wherein the ring is substituted with C.sub.1-C.sub.4 alkyl J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I) can be N-(2-(1H-indol-3-yl)ethyl)-7-isopropyl-2-(5-methylpyridin-3-yl)thieno[3,2-d]pyrimidin-4-amine.

[0253] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G is N;  custom-character joining G and J can be a double bond; R.sup.a can be hydrogen; R.sup.b can be —

CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.K can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.K moiety indicated as substituted is substituted with one or more Q, wherein Q is cyano; R.sup.Y and R.sup.Z taken together can be

##STR00055##

J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I) can be 5-(4-((2-(1H-indol-3-yl)ethyl)amino)furo[3,2-d]pyrimidin-2-yl)nicotinonitrile.

[0254] In some embodiments, provided herein is compound of Formula (I), wherein the compound

can be selected from: [0255] 4-(2-((2-(benzo[b]thiophen-3-yl)-6-(isopropylamino)pyrimidin-4-yl)amino)ethyl)phenol; [0256] 4-(2-((2-(benzo[b]thiophen-3-yl)-7-isopropylthieno[3,2-d]pyrimidin-4-yl)amino)ethyl)phenol; [0257] 4-(2-((2-(benzo[b]thiophen-3-yl)-7-isopropyl-6,7-dihydro-5H-pyrrolo[2,3-d]pyrimidin-4-yl)amino)ethyl)phenol; [0258] 2-(benzo[b]thiophen-3-yl)-4-((4-hydroxyphenethyl)amino)-7-isopropyl-5,7-dihydro-6H-pyrrolo[2,3-d]pyrimidin-6-one; [0259] 3-((2-(benzo[b]thiophen-3-yl)-9-isopropyl-9H-purin-6-yl)oxy)propanamide; [0260] 4-(2-((2-(benzo[b]thiophen-3-yl)-8-(dimethylamino)pyrimido[5,4-d]pyrimidin-4-yl)amino)ethyl)phenol; [0261] 5-(2-((2-(1H-indol-3-yl)ethyl)amino)-6-(sec-butylamino)pyrimidin-4-yl)nicotinonitrile; [0262] N-(2-(1H-indol-3-yl)ethyl)-2-methyl-6-phenylthieno[2,3-d]pyrimidin-4-amine; [0263] N-(2-(1H-indol-3-yl)ethyl)-6-(4-fluorophenyl)thieno[2,3-d]pyrimidin-4-amine; [0264] 3-(2-(benzo[b]thiophen-3-yl)-9-isopropyl-6-oxo-6,9-dihydro-1H-purin-1-yl)propanamide; [0265] N-(2-(1H-indol-3-yl)ethyl)-2-(5-fluoropyridin-3-yl)quinazolin-4-amine; [0266] 5-(4-((2-(1H-indol-3-yl)ethyl)amino)quinazolin-2-yl)nicotinonitrile; [0267] N.sup.4-(2-(1H-indol-3-yl)ethyl)-N.sup.2-(sec-butyl)quinazoline-2,4-diamine; [0268] 2-(benzo[b]thiophen-3-yl)-4-((4-hydroxyphenethyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-5-carbonitrile; [0269] N-(2-(1H-indol-3-yl)ethyl)-6-(benzo[b]thiophen-3-yl)-3-isopropylimidazo[1,5-a]pyrazin-8-amine; [0270] 4-(2-((6-(benzo[b]thiophen-3-yl)-3-isopropylimidazo[1,5-a]pyrazin-8-yl)amino)ethyl)phenol; [0271] 5-(4-((2-(1H-indol-3-yl)ethyl)amino)-7-isopropylthieno[3,2-d]pyrimidin-2-yl)nicotinonitrile; [0272] N-(2-(1H-indol-3-yl)ethyl)-2-(5-fluoropyridin-3-yl)-7-isopropylthieno[3,2-d]pyrimidin-4-amine; [0273] N-(2-(1H-indol-3-yl)ethyl)-2-(5-fluoropyridin-3-

yl)furo[3,2-d]pyrimidin-4-amine; [0274] N-(2-(1H-indol-3-yl)ethyl)-2-(5-methylpyridin-3-yl)furo[3,2-d]pyrimidin-4-amine; [0275] N-(2-(1H-indol-3-yl)ethyl)-7-isopropyl-2-(5-methylpyridin-3-yl)thieno[3,2-d]pyrimidin-4-amine; [0276] 5-(4-((2-(1H-indol-3-yl)ethyl)amino)furo[3,2-d]pyrimidin-2-yl)nicotinonitrile; and [0277] pharmaceutically acceptable salts thereof.

Formula (I-A)

[0278] In some embodiments provided herein, the compound of Formula (I) can have the structure of Formula (I-A):

##STR00056##

including pharmaceutically acceptable salts thereof, wherein: R^{sup.J} can be —NR^{sup.a}R^{sup.b}; R^{sup.a} can be hydrogen or C_{sub.1}-C_{sub.4} alkyl; R^{sup.b} can be R^{sup.c} or —(C_{sub.1}-C_{sub.4} alkyl)-R^{sup.c}; R^{sup.c} can be selected from the group consisting of: unsubstituted C_{sub.6-10} aryl; substituted C_{sub.6-10} aryl; unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R^{sup.c} moiety indicated as substituted is substituted with one or more substituents E, wherein each E can be independently selected from the group consisting of: —OH, C_{sub.1}-C_{sub.4} alkyl, C_{sub.1}-C_{sub.4} haloalkyl, —O(C_{sub.1}-C_{sub.4} alkyl), and —O(C_{sub.1}-C_{sub.4} haloalkyl); R^{sup.K} can be selected from the group consisting of: hydrogen, unsubstituted C_{sub.1-6} alkyl; [0279] —NH(C_{sub.1-4} alkyl); —N(C_{sub.1-4}alkyl).sub.2, unsubstituted C_{sub.6-10} aryl; substituted C_{sub.6-10} aryl; unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R^{sup.K} moiety indicated as substituted is substituted with one or more substituents Q, wherein each Q can be independently selected from the group consisting of: —OH, C_{sub.1-4} alkyl, C_{sub.1-4} haloalkyl, halo, cyano, —O—(C_{sub.1-4} alkyl), and —O—(C_{sub.1-4} haloalkyl); Y and Z can each be C; X can be N or CH; W can be O or S; and R^{sup.a} can be hydrogen or C_{sub.1}-C_{sub.4} alkyl.

[0280] In some embodiments, R^{sup.a} can be hydrogen. In other embodiments, R^{sup.a} can be C_{sub.1}-C_{sub.4} alkyl.

[0281] In some embodiments, R^{sup.b} can be —(C_{sub.1}-C_{sub.4} alkyl)-R^{sup.c}. For example, R^{sup.b} can be —CH_{sub.2}—R^{sup.c}, —CH_{sub.2}CH_{sub.2}—R^{sup.c}, —CH_{sub.2}CH_{sub.2}CH_{sub.2}—R^{sup.c}, or —CH_{sub.2}CH_{sub.2}CH_{sub.2}CH_{sub.2}—R^{sup.c}.

[0282] In some embodiments, R^{sup.c} can be —OH. In some embodiments, R^{sup.c} can be —O(C_{sub.1}-C_{sub.4} alkyl). In some embodiments, R^{sup.c} can be —O(C_{sub.1}-C_{sub.4} haloalkyl). In some embodiments, R^{sup.c} can be —C(=O)NH_{sub.2}. In some embodiments, R^{sup.c} can be unsubstituted C_{sub.6-10} aryl. In some embodiments, R^{sup.c} can be substituted C_{sub.6-10} aryl. In some embodiments, R^{sup.c} can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S. In some embodiments, R^{sup.c} can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S. In some embodiments, when a R^{sup.c} moiety is indicated as substituted, the moiety can be substituted with one or more, for example, one, two, three, or four substituents E. In some embodiments, E can be —OH. In some embodiments, E can be C_{sub.1}-C_{sub.4} alkyl. In some embodiments, E can be C_{sub.1}-C_{sub.4} haloalkyl. In some embodiments, E can be —O(C_{sub.1}-C_{sub.4} alkyl). In some embodiments, E can be —O(C_{sub.1}-C_{sub.4} haloalkyl). In some embodiments R^{sup.c} can be phenyl. In other embodiments, R^{sup.c} can be hydroxyphenyl. In still other embodiments, R^{sup.c} can be indolyl.

[0283] In some embodiments, R^{sup.K} can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S. In some embodiments, R^{sup.K} can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the

group consisting of O, N, and S; wherein the substituted heteroaryl can be substituted with one or more substituents Q, wherein each Q can be independently selected from the group consisting of: —OH, C.sub.1-4 alkyl, C.sub.1-4 haloalkyl, halo, cyano, —O—(C.sub.1-4 alkyl), and —O—(C.sub.1-4 haloalkyl). In some embodiments, R.sup.K can be pyridinyl. In other embodiments, R.sup.K can be pyridinyl substituted with one or more substituents Q. For example, R.sup.K can be methylpyridinyl, ethylpyridinyl, cyanopyridinyl, chloropyridinyl, fluoropyridinyl, or bromopyridinyl.

[0284] In some embodiments, R.sup.e can be hydrogen. In some embodiments, R.sup.e can be C.sub.1-C.sub.4 alkyl. For example, R.sup.e can be methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl or tert-butyl.

[0285] In some embodiments, R.sup.a can be hydrogen; R.sup.b can be —(C.sub.1-C.sub.4 alkyl)—R.sup.c; R.sup.c can be selected from the group consisting of: unsubstituted C.sub.6-10 aryl; substituted C.sub.6-10 aryl; unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.c moiety indicated as substituted is substituted with one or more substituents E, wherein each E can be independently selected from the group consisting of: —OH, C.sub.1-C.sub.4 alkyl, C.sub.1-C.sub.4 haloalkyl, —O(C.sub.1-C.sub.4 alkyl), and —O(C.sub.1-C.sub.4 haloalkyl); R.sup.K can be selected from the group consisting of: unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein the substituted heteroaryl is substituted with one or more substituents Q, wherein each Q can be independently selected from the group consisting of: —OH, C.sub.1-4 alkyl, C.sub.1-4 haloalkyl, halo, cyano, —O—(C.sub.1-4 alkyl), and —O—(C.sub.1-4 haloalkyl); and R.sup.e can be C.sub.1-C.sub.4 alkyl.

[0286] In some embodiments, R.sup.a can be hydrogen; R.sup.b can be —(CH.sub.2—CH.sub.2)—R.sup.c; R.sup.c can be selected from the group consisting of: substituted phenyl and unsubstituted indolyl; wherein the substituted phenyl is substituted with one substituent E, wherein E can be —OH; R.sup.K can be selected from the group consisting of: unsubstituted benzothiophenyl and substituted pyridinyl; wherein the substituted pyridinyl is substituted with one substituent Q, wherein Q can be selected from the group consisting of: C.sub.1-4 alkyl, halo, and cyano; and R.sup.e can be isopropyl.

[0287] In some embodiments, when W is O, R.sup.J can be —NR.sup.aR.sup.b; R.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be selected from the group consisting of: unsubstituted C.sub.6-10 aryl; substituted C.sub.6-10 aryl; unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.c moiety indicated as substituted is substituted with one or more substituents E, wherein each E can be independently selected from the group consisting of: —OH, C.sub.1-C.sub.4 alkyl, and —O(C.sub.1-C.sub.4 alkyl); R.sup.K can be selected from the group consisting of unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.K moiety indicated as substituted is substituted with one or more substituents Q, wherein each Q can be independently selected from the group consisting of: —C.sub.1-4 alkyl, halo, cyano, and —O—(C.sub.1-4 alkyl); Y and Z can each be C; X can be N or CH; and R.sup.e can be hydrogen or C.sub.1-C.sub.4 alkyl.

[0288] In some embodiments, when W is S, R.sup.J can be —NR.sup.aR.sup.b; R.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be selected from the group consisting of: unsubstituted C.sub.6-10 aryl; substituted C.sub.6-10 aryl; unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and

substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.c moiety indicated as substituted is substituted with one or more substituents E, wherein each E can be independently selected from the group consisting of: —OH, C.sub.1-C.sub.4 alkyl, and —O(C.sub.1-C.sub.4 alkyl); R.sup.K can be selected from the group consisting of unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.K moiety indicated as substituted is substituted with one or more substituents Q, wherein each Q can be independently selected from the group consisting of: —C.sub.1-4 alkyl, halo, cyano, and —O—(C.sub.1-4 alkyl); Y and Z can each be C; X can be N or CH; and R.sup.e can be hydrogen or C.sub.1-C.sub.4 alkyl. [0289] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N; R.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.K can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.K moiety indicated as substituted is substituted with one or more Q, wherein Q is C.sub.1-C.sub.4 alkyl; W can be S; R.sup.e can be C.sub.1-C.sub.4 alkyl; J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I-A) can be N-(2-(1H-indol-3-yl)ethyl)-7-isopropyl-2-(5-methylpyridin-3-yl)thieno[3,2-d]pyrimidin-4-amine.

[0290] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N; R.sup.a can be hydrogen R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.K can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.K moiety indicated as substituted is substituted with one or more Q, wherein Q is cyano; W can be S; R.sup.e can be C.sub.1-C.sub.4 alkyl; J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I-A) can be 5-(4-((2-(1H-indol-3-yl)ethyl)amino)-7-isopropylthieno[3,2-d]pyrimidin-2-yl)nicotinonitrile.

[0291] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N; R.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.K can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.K moiety indicated as substituted is substituted with one or more Q, wherein Q is halo; W can be S; R.sup.e can be C.sub.1-C.sub.4 alkyl; J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I-A) can be N-(2-(1H-indol-3-yl)ethyl)-2-(5-fluoropyridin-3-yl)-7-isopropylthieno[3,2-d]pyrimidin-4-amine.

[0292] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N; R.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c, R.sup.c can be substituted C.sub.6-10 aryl, substituted with one or more E, wherein E can be —OH; R.sup.K can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; W can be S; R.sup.e can be C.sub.1-C.sub.4 alkyl; J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I-A) can be 4-(2-((2-(benzo[b]thiophen-3-yl)-7-isopropylthieno[3,2-d]pyrimidin-4-yl)amino)ethyl)phenol.

[0293] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N; R.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.K can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.K moiety indicated as substituted is substituted with one or more Q, wherein Q is halo; W can be O; R.sup.e can be hydrogen; J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I-A) can be N-(2-(1H-indol-3-yl)ethyl)-2-(5-fluoropyridin-3-yl)furo[3,2-d]pyrimidin-4-amine.

[0294] In some embodiments, when R.sup.J is —R.sup.aR.sup.b; G can be N; custom-character joining G and J can be a double bond; R.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.K can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.K moiety indicated as substituted is substituted with one or more Q, wherein Q is C.sub.1-C.sub.4 alkyl; W can be 0; R.sup.e can be hydrogen; J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I-A) can be N-(2-(1H-indol-3-yl)ethyl)-2-(5-methylpyridin-3-yl)furo[3,2-d]pyrimidin-4-amine.

[0295] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G is NR.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.K can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.K moiety indicated as substituted is substituted with one or more Q, wherein Q is cyano; W can be O; R.sup.e can be hydrogen; J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I-A) can be 5-(4-((2-(1H-indol-3-yl)ethyl)amino)furo[3,2-d]pyrimidin-2-yl)nicotinonitrile.

[0296] In some embodiments, the compound of Formula (I-A), or a pharmaceutically acceptable salt thereof, can be selected from the group consisting of: [0297] N-(2-(1H-indol-3-yl)ethyl)-7-isopropyl-2-(5-methylpyridin-3-yl)thieno[3,2-d]pyrimidin-4-amine; [0298] 5-(4-((2-(1H-indol-3-yl)ethyl)amino)-7-isopropylthieno[3,2-d]pyrimidin-2-yl)nicotinonitrile; N-(2-(1H-indol-3-yl)ethyl)-2-(5-fluoropyridin-3-yl)-7-isopropylthieno[3,2-d]pyrimidin-4-amine; [0299] 4-(2-((2-(benzo[b]thiophen-3-yl)-7-isopropylthieno[3,2-d]pyrimidin-4-yl)amino)ethyl)phenol; [0300] N-(2-(1H-indol-3-yl)ethyl)-2-(5-fluoropyridin-3-yl)furo[3,2-d]pyrimidin-4-amine; [0301] N-(2-(1H-indol-3-yl)ethyl)-2-(5-methylpyridin-3-yl)furo[3,2-d]pyrimidin-4-amine; and [0302] 5-(4-((2-(1H-indol-3-yl)ethyl)amino)furo[3,2-d]pyrimidin-2-yl)nicotinonitrile.

Formula (I-B)

[0303] In other embodiments provided herein, the compound of Formula (I) can have the structure of Formula (I-B):

##STR00057##

including pharmaceutically acceptable salts thereof, wherein: R.sup.a can be hydrogen or C.sub.1-C.sub.4 alkyl; R.sup.b can be R.sup.c or —(C.sub.1-4 alkyl)-R.sup.c; R.sup.c can be selected from the group consisting of: —OH, —O(C.sub.1-C.sub.4 alkyl), —O(C.sub.1-C.sub.4 haloalkyl); —C(=O)NH.sub.2; unsubstituted C.sub.6-10 aryl; substituted C.sub.6-10 aryl; unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.c moiety indicated as substituted is substituted with one or more substituents E, wherein each E can be independently selected from the group consisting of: —OH, C.sub.1-C.sub.4 alkyl, C.sub.1-C.sub.4 haloalkyl, —O(C.sub.1-C.sub.4 alkyl), and —O(C.sub.1-C.sub.4 haloalkyl); R.sup.K can be selected from the group consisting of: hydrogen, unsubstituted C.sub.1-6 alkyl; substituted C.sub.1-6 alkyl; —NH(C.sub.1-4 alkyl); —N(C.sub.1-4 alkyl).sub.2, unsubstituted C.sub.6-10 aryl; substituted C.sub.6-10 aryl; unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.K moiety indicated as substituted is substituted with one or more substituents Q, wherein each Q can be independently selected from the group consisting of: —OH, C.sub.1-4 alkyl, C.sub.1-4 haloalkyl, halo, cyano, —O—(C.sub.1-4 alkyl), and —O—(C.sub.1-4 haloalkyl); R.sup.G can be selected from the group consisting of hydrogen, C.sub.1-4 alkyl, and —(C.sub.1-4 alkyl)-C(=O)NH.sub.2; R.sup.f can be selected from the group consisting of hydrogen, C.sub.1-4 alkyl, unsubstituted C.sub.6-C.sub.10 aryl, and C.sub.6-C.sub.10 aryl substituted with 1-5 halo

atoms; U can be N or CR.sup.U; V can be S or NR.sup.V; R.sup.U can be selected from the group consisting of hydrogen, C.sub.1-4 alkyl, halo, and cyano; R.sup.V can be hydrogen or C.sub.1-C.sub.4 alkyl; wherein when U is CR.sup.U and V is NR.sup.V, R.sup.U is selected from the group consisting of C.sub.1-4 alkyl, halo, and cyano; Y and Z can each be C; and X can be N or CH.
[0304] In some embodiments, R.sup.a can be hydrogen. In other embodiments, R.sup.a can be C.sub.1-C.sub.4 alkyl.

[0305] In some embodiments, R.sup.b can be —(C.sub.1-C.sub.4 alkyl)-R.sup.c. For example, R.sup.b can be —CH.sub.2—R.sup.c, —CH.sub.2CH.sub.2—R.sup.c, —CH.sub.2CH.sub.2CH.sub.2—R.sup.c, or —CH.sub.2CH.sub.2CH.sub.2CH.sub.2—R.sup.c. In certain embodiments, R.sup.b can be —(CH.sub.2CH.sub.2)—R.sup.c. In certain embodiments, R.sup.b can be —(CH.sub.2CH.sub.2)—C(=O)NH.sub.2. In certain embodiments, R.sup.b can be —(CH.sub.2CH.sub.2)-(indolyl). In certain embodiments, R.sup.b can be —(CH.sub.2CH.sub.2)-(hydroxyphenyl).

[0306] In some embodiments, R.sup.c can be —OH. In some embodiments, R.sup.c can be —O(C.sub.1-C.sub.4 alkyl). In some embodiments, R.sup.c can be —O(C.sub.1-C.sub.4 haloalkyl). In some embodiments, R.sup.c can be —C(=O)NH.sub.2. In some embodiments, R.sup.c can be unsubstituted C.sub.6-10 aryl. In some embodiments, R.sup.c can be substituted C.sub.6-10 aryl. In some embodiments, R.sup.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S. In some embodiments, R.sup.c can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S. In some embodiments, when a R.sup.c moiety is indicated as substituted, the moiety can be substituted with one or more, for example, one, two, three, or four substituents E. In some embodiments, E can be

—OH. In some embodiments, E can be C.sub.1-C.sub.4 alkyl. In some embodiments, E can be C.sub.1-C.sub.4 haloalkyl. In some embodiments, E can be —O(C.sub.1-C.sub.4 alkyl). In some embodiments, E can be —O(C.sub.1-C.sub.4 haloalkyl).

[0307] In some embodiments, R.sup.K can be hydrogen. In other embodiments, R.sup.K can be C.sub.1-C.sub.4 alkyl. For example, R.sup.K can be methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl or tert-butyl. In some embodiments, R.sup.K can be selected from the group consisting of: unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein the substituted heteroaryl can be substituted with one or more substituents Q, wherein each Q can be independently selected from the group consisting of: —OH, C.sub.1-4 alkyl, C.sub.1-4 haloalkyl, halo, cyano, —O—(C.sub.1-4 alkyl), and —O—(C.sub.1-4 haloalkyl). In certain embodiments, R.sup.K can be benzothiophenyl. In other embodiments, R.sup.K can be pyridinyl substituted with one or more substituents Q. For example, R.sup.K can be methylpyridinyl, ethylpyridinyl, cyanopyridinyl, chloropyridinyl, fluoropyridinyl, or bromopyridinyl.

[0308] In some embodiments, R.sup.G can be selected from the group consisting of hydrogen, C.sub.1-4 alkyl, and —(C.sub.1-4 alkyl)-C(=O)NH.sub.2. In certain embodiments, R.sup.G can be —(CH.sub.2CH.sub.2)—C(=O)NH.sub.2.

[0309] In some embodiments, R.sup.f can be hydrogen. In other embodiments, R.sup.f can be C.sub.1-4 alkyl. For example, R.sup.f can be methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl or tert-butyl. In some embodiments, R.sup.f can be unsubstituted C.sub.6-C.sub.10 aryl. In other embodiments, R.sup.f can be C.sub.6-C.sub.10 aryl substituted with 1-5 halo atoms. In certain embodiments, R.sup.f can be phenyl substituted with 1-5 halo atoms. In certain embodiments, R.sup.f can be fluorophenyl.

[0310] In some embodiments, U can be N. In other embodiments, U can be CR.sup.U.

[0311] In some embodiments, V can be S. In other embodiments, V can be NR.sup.V.

[0312] In some embodiments, R.sup.U can be hydrogen. In some embodiments, R.sup.U can be

C.sub.1-4 alkyl. In other embodiments R.sup.U can be halo. For example, R.sup.U can be fluoro, chloro, bromo, or iodo. In still other embodiments, R.sup.U can be cyano.

[0313] In some embodiments, R.sup.V can be hydrogen. In other embodiments, R.sup.V can be C.sub.1-4 alkyl. For example, R.sup.V can be methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl or tert-butyl. In some embodiments, Y and Z can each be C and X can be N. In other embodiments, Y and Z can each be C and X can be CH.

[0314] In some embodiments, R.sup.a can be hydrogen; R.sup.b can be —(C.sub.1-4 alkyl)-R.sup.c; R.sup.c can be selected from the group consisting of: —C(=O)NH.sub.2, unsubstituted C.sub.6-10 aryl; substituted C.sub.6-10 aryl; unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.c moiety indicated as substituted can be substituted with one or more substituents E, wherein each E can be independently selected from the group consisting of: —OH, C.sub.1-C.sub.4 alkyl, C.sub.1-C.sub.4 haloalkyl, —O(C.sub.1-C.sub.4 alkyl), and —O(C.sub.1-C.sub.4 haloalkyl); R.sup.K can be selected from the group consisting of: unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein the substituted heteroaryl is substituted with one or more substituents Q, wherein each Q can be independently selected from the group consisting of: —OH, C.sub.1-4 alkyl, C.sub.1-4 haloalkyl, halo, cyano, —O—(C.sub.1-4 alkyl), and —O—(C.sub.1-4 haloalkyl); R.sup.G is C.sub.1-4 alkyl or

—(C.sub.1-4 alkyl)-C(=O)NH.sub.2; R.sup.f can be selected from the group consisting of hydrogen, unsubstituted phenyl, and phenyl substituted with 1-5 halo atoms; Y and Z each can be C; and X can be CH.

[0315] In some embodiments, R.sup.a can be hydrogen; R.sup.b can be —(CH.sub.2—CH.sub.2)—R.sup.c; R.sup.c can be selected from the group consisting of: —C(=O)NH.sub.2, substituted phenyl and unsubstituted indolyl; wherein the substituted phenyl is substituted with one substituent E, wherein E can be —OH; R.sup.K can be selected from the group consisting of: unsubstituted benzothiohenyl and substituted pyridinyl; wherein the substituted pyridinyl is substituted with one substituent Q, wherein Q can be selected from the group consisting of: C.sub.1-4 alkyl, halo, and cyano; R.sup.G can be —(CH.sub.2CH.sub.2)—C(=O)NH.sub.2; R.sup.f can be selected from the group consisting of hydrogen, phenyl, and fluorophenyl; Y and Z each can be C; and X can be CH.

[0316] In some embodiments, when V is S, R.sup.a can be hydrogen or C.sub.1-C.sub.4 alkyl; R.sup.b can be R.sup.c or —(CH.sub.2—CH.sub.2)—R.sup.c; R.sup.c can be selected from the group consisting of: —C(=O)NH.sub.2; unsubstituted C.sub.6-10 aryl; substituted C.sub.6-10 aryl; unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.c moiety indicated as substituted is substituted with one or more substituents E, wherein each E can be independently selected from the group consisting of: —OH, C.sub.1-C.sub.4 alkyl, and —O(C.sub.1-C.sub.4 alkyl); R.sup.K can be selected from the group consisting of: hydrogen, unsubstituted C.sub.1-6 alkyl; substituted C.sub.1-6 alkyl; —NH(C.sub.1-4 alkyl); and —N(C.sub.1-4 alkyl).sub.2; wherein a R.sup.K moiety indicated as substituted is substituted with one or more substituents Q, wherein each Q can be independently selected from the group consisting of: —OH, C.sub.1-4 alkyl, halo, cyano, and —O—(C.sub.1-4 alkyl); R.sup.G can be selected from the group consisting of hydrogen, C.sub.1-4 alkyl, and —(C.sub.1-4 alkyl)-C(=O)NH.sub.2; R.sup.f can be selected from the group consisting of hydrogen, C.sub.1-4 alkyl, unsubstituted C.sub.6-C.sub.10 aryl, and C.sub.6-C.sub.10 aryl substituted with 1-5 halo atoms; U can be CR.sup.U; R.sup.U can be selected from the group consisting of hydrogen, C.sub.1-4 alkyl, halo, and cyano; Y and Z can each be C; and X can be N.

[0317] In some embodiments, when V is NR.sup.V, R.sup.a can be hydrogen or C.sub.1-C.sub.4

alkyl; R^{sup.b} can be R^{sup.c} or —(CH_{sub.2}—CH_{sub.2})—R^{sup.c}; R^{sup.c} can be selected from the group consisting of: —C(=O)NH_{sub.2}; unsubstituted C_{sub.6-10} aryl; substituted C_{sub.6-10} aryl; unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R^{sup.c} moiety indicated as substituted is substituted with one or more substituents E, wherein each E can be independently selected from the group consisting of: —OH, C_{sub.1-4} alkyl, C_{sub.1-4}, and —O(C_{sub.1-4} alkyl); R^{sup.K} can be selected from the group consisting of: unsubstituted C_{sub.6-10} aryl; substituted C_{sub.6-10} aryl; unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R^{sup.K} moiety indicated as substituted is substituted with one or more substituents Q, wherein each Q can be independently selected from the group consisting of: —OH, C_{sub.1-4} alkyl, halo, cyano, and —O—(C_{sub.1-4} alkyl); R^{sup.G} can be selected from the group consisting of hydrogen, C_{sub.1-4} alkyl, and —(C_{sub.1-4} alkyl)-C(=O)NH_{sub.2}; R^{sup.f} can be hydrogen; U can be N or CR^{sup.U}; R^{sup.U} can be selected from the group consisting of C_{sub.1-4} alkyl, halo, and cyano; R^{sup.V} can be hydrogen or C_{sub.1-4} alkyl; Y and Z can each be C; and X can be N or CH.

[0318] In some embodiments, when R^{sup.J} is —OR^{sup.b}; G can be N;  custom-character joining G and J can be a double bond; R^{sup.b} can be —CH_{sub.2}CH_{sub.2}—R^{sup.c}; R^{sup.c} can be

—C(=O)NH_{sub.2}; R^{sup.K} can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; U can be N; V can be NR^{sup.v}; R^{sup.v} can be C_{sub.1-4} alkyl; R^{sup.f} can be hydrogen; J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I-B) can be 3-((2-(benzo[b]thiophen-3-yl)-9-isopropyl-9H-purin-6-yl)oxy)propanamide.

[0319] In some embodiments, when R^{sup.J} is =O; G can be N substituted with R^{sup.G};  custom-character joining G and J can be a single bond; R^{sup.G} can be —(C_{sub.1-4} alkyl)-C(=O)NH_{sub.2}; R^{sup.K} can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; U can be N; V can be NR^{sup.v}; R^{sup.v} can be C_{sub.1-4} alkyl; R^{sup.f} can be hydrogen; J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I-B) can be 3-(2-(benzo[b]thiophen-3-yl)-9-isopropyl-6-oxo-6,9-dihydro-1H-purin-1-yl)propanamide.

[0320] In some embodiments, when R^{sup.J} is —NR^{sup.a}R^{sup.b}; G can be N;  custom-character joining G and J can be a double bond; R^{sup.a} can be hydrogen; R^{sup.b} can be —CH_{sub.2}CH_{sub.2}—R^{sup.c}; R^{sup.c} can be substituted C_{sub.6-10} aryl, substituted with one or more E, wherein E is —OH; R^{sup.K} can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; U can be CR^{sup.u}; R^{sup.u} can be cyano; V can be NR^{sup.v}; R^{sup.v} can be C_{sub.1-4} alkyl; R^{sup.f} can be hydrogen; J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I-B) can be 2-(benzo[b]thiophen-3-yl)-4-((4-hydroxyphenethyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-5-carbonitrile.

[0321] In some embodiments, when R^{sup.J} is —NR^{sup.a}R^{sup.b}; G can be N;  custom-character joining G and J can be a double bond; R^{sup.a} can be hydrogen; R^{sup.b} can be —CH_{sub.2}CH_{sub.2}—R^{sup.c}; R^{sup.c} can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R^{sup.K} can be unsubstituted C_{sub.1-6} alkyl; U can be CR^{sup.u}; R^{sup.u} can be hydrogen; V can be S; R^{sup.f} can be phenyl; J can be C; X can be N; Y can be C; Z can be C. In some embodiments, the compound of Formula (I-B) can be N-(2-(1H-indol-3-yl)ethyl)-2-methyl-6-phenylthieno[2,3-d]pyrimidin-4-amine.

[0322] In some embodiments, when R^{sup.J} can be —NR^{sup.a}R^{sup.b}; G can be N;  custom-character joining G and J can be a double bond; R^{sup.a} can be hydrogen; R^{sup.b} can be

—CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.K can be hydrogen; U can be CR.sup.u; R.sup.u can be hydrogen; V can be S; R.sup.f can be fluorophenyl; J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I-B) can be N-(2-(1H-indol-3-yl)ethyl)-6-(4-fluorophenyl)thieno[2,3-d]pyrimidin-4-amine.

[0323] In some embodiments, the compound of Formula (I-B), or a pharmaceutically acceptable salt thereof, can be selected from the group consisting of: [0324] 3-((2-(benzo[b]thiophen-3-yl)-9-isopropyl-9H-purin-6-yl)oxy)propanamide; [0325] 3-(2-(benzo[b]thiophen-3-yl)-9-isopropyl-6-oxo-6,9-dihydro-1H-purin-1-yl)propanamide; [0326] 2-(benzo[b]thiophen-3-yl)-4-((4-hydroxyphenethyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-5-carbonitrile; [0327] N-(2-(1H-indol-3-yl)ethyl)-2-methyl-6-phenylthieno[2,3-d]pyrimidin-4-amine; and [0328] N-(2-(1H-indol-3-yl)ethyl)-6-(4-fluorophenyl)thieno[2,3-d]pyrimidin-4-amine.

Formula (I-C)

[0329] In still other embodiments provided herein, the compound of Formula (I) can have the structure of Formula (I-C):

##STR00058##

including pharmaceutically acceptable salts thereof, wherein: R.sup.J can be —NR.sup.aR.sup.b; R.sup.a can be hydrogen or C.sub.1-C.sub.4 alkyl; R.sup.b can be R.sup.c or —(C.sub.1-C.sub.4 alkyl)-R.sup.c; R.sup.c can be selected from the group consisting of: unsubstituted C.sub.6-10 aryl; substituted C.sub.6-10 aryl; unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.c moiety indicated as substituted is substituted with one or more substituents E, wherein each E can be independently selected from the group consisting of: —OH, C.sub.1-C.sub.4 alkyl, C.sub.1-C.sub.4 haloalkyl, —O(C.sub.1-C.sub.4 alkyl), and —O(C.sub.1-C.sub.4 haloalkyl); R.sup.K can be selected from the group consisting of: hydrogen, unsubstituted C.sub.1-6 alkyl; —NH(C.sub.1-4 alkyl); —N(C.sub.1-4 alkyl).sub.2, unsubstituted C.sub.6-10 aryl; substituted C.sub.6-10 aryl; unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.K moiety indicated as substituted is substituted with one or more substituents Q, wherein each Q can be independently selected from the group consisting of: —OH, C.sub.1-4 alkyl, C.sub.1-4 haloalkyl, halo, cyano, —O—(C.sub.1-4 alkyl), and —O—(C.sub.1-4 haloalkyl); A can be N or CH; B can be N or CH; R.sup.g can be selected from the group consisting of hydrogen, C.sub.1-4 alkyl, and —N(C.sub.1-4 alkyl).sub.2; Y and Z can each be C; and X can be N or CH.

[0330] In some embodiments, R.sup.K can be —NH(C.sub.1-4 alkyl). For example, in some embodiments, R.sup.K can be —NH(CH.sub.3), —NH(CH.sub.2CH.sub.3), —NH(isopropyl), or —NH(sec-butyl). In some embodiments, R.sup.K can be unsubstituted benzothiophenyl. In other embodiments, R.sup.K can be substituted pyridinyl. For example, R.sup.K can be methylpyridinyl, ethylpyridinyl, cyanopyridinyl, chloropyridinyl, fluoropyridinyl, or bromopyridinyl.

[0331] In some embodiments, A can be N and B can be N. In other embodiments, A can be N and B can be CH. In still other embodiments, A can be CH and B can be N. In yet still other embodiments, A can be CH and B can be CH.

[0332] In some embodiments, R.sup.g can be hydrogen. In other embodiments, R.sup.g can be —N(C.sub.1-4 alkyl).sub.2. In certain embodiments, R.sup.g can be —N(CH.sub.3).sub.2.

[0333] In some embodiments, R.sup.a can be hydrogen; R.sup.b can be —(C.sub.1-C.sub.4 alkyl)-R.sup.c; R.sup.c can be selected from the group consisting of: unsubstituted C.sub.6-10 aryl; substituted C.sub.6-10 aryl; unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.c moiety

indicated as substituted is substituted with one or more substituents E, wherein each E can be independently selected from the group consisting of: —OH, C.sub.1-C.sub.4 alkyl, C.sub.1-C.sub.4 haloalkyl, —O(C.sub.1-C.sub.4 alkyl), and —O(C.sub.1-C.sub.4 haloalkyl); R.sup.K can be selected from the group consisting of: —NH(C.sub.1-4 alkyl); unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein the substituted heteroaryl is substituted with one or more substituents Q, wherein each Q can be independently selected from the group consisting of: —OH, C.sub.1-4 alkyl, C.sub.1-4 haloalkyl, halo, cyano, —O—(C.sub.1-4 alkyl), and —O—(C.sub.1-4 haloalkyl); and R.sup.g can be hydrogen or —N(C.sub.1-4 alkyl).sub.2.

[0334] In some embodiments, R.sup.a can be hydrogen; R.sup.b can be —(C.sub.1-C.sub.4 alkyl)-R.sup.c; R.sup.c can be selected from the group consisting of: substituted phenyl and unsubstituted indolyl; wherein the substituted phenyl is substituted with one or more substituents E, wherein each E can be independently selected from the group consisting of: —OH, C.sub.1-C.sub.4 alkyl, C.sub.1-C.sub.4 haloalkyl, —O(C.sub.1-C.sub.4 alkyl), and —O(C.sub.1-C.sub.4 haloalkyl); R.sup.K can be selected from the group consisting of: —NH(C.sub.1-4 alkyl); unsubstituted benzothiophenyl; and substituted pyridinyl; wherein the substituted pyridinyl is substituted with one or more substituents Q, wherein each Q can be independently selected from the group consisting of: —OH, C.sub.1-4 alkyl, C.sub.1-4 haloalkyl, halo, cyano, —O—(C.sub.1-4 alkyl), and —O—(C.sub.1-4 haloalkyl); and R.sup.g can be hydrogen or —N(C.sub.1-4 alkyl).sub.2.

[0335] In some embodiments, R.sup.a can be hydrogen; R.sup.b can be —(CH.sub.2CH.sub.2)—R.sup.c; R.sup.c can be selected from the group consisting of: substituted phenyl and unsubstituted indolyl; wherein the substituted phenyl is substituted with one substituent E, wherein E can be —OH; R.sup.K can be selected from the group consisting of: —NH(sec-butyl); unsubstituted benzothiophenyl, and substituted pyridinyl; wherein the substituted pyridinyl is substituted with one or more substituents Q, wherein each Q can be independently selected from the group consisting of: C.sub.1-4 alkyl, halo, and cyano; and R.sup.g can be hydrogen or —N(CH.sub.3).sub.2.

[0336] In some embodiments, when A is C and B is C, R.sup.J can be —NR.sup.aR.sup.b; G can be N; R.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be substituted C.sub.6-10 aryl, substituted with one or more E, wherein E is —OH; or unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.K can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.g can be hydrogen; J can be C; X can be N; Y can be C; and Z is C.

[0337] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N; R.sup.a can be hydrogen; R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be substituted C.sub.6-10 aryl, substituted with one or more E, wherein E is —OH; R.sup.K is unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; A can be N; B can be N; R.sup.g can be —N(C.sub.1-4 alkyl).sub.2; J can be C; X can be N; Y can be C; and Z is C. In some embodiments, the compound of Formula (I-C) can be 4-(2-((2-(benzo[b]thiophen-3-yl)-8-(dimethylamino)pyrimido[5,4-d]pyrimidin-4-yl)amino)ethyl)phenol.

[0338] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G can be N; R.sup.a can be hydrogen R.sup.b can be —CH.sub.2CH.sub.2—R.sup.c; R.sup.c can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R.sup.K can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R.sup.K moiety indicated as substituted is substituted with one or more Q, wherein Q can be halo; A can be CH; B can be CH; R.sup.g can be hydrogen; J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I-C) can be N-(2-(1H-indol-3-yl)ethyl)-2-(5-fluoropyridin-3-yl)quinazolin-4-amine.

[0339] In some embodiments, when R.sup.J is —NR.sup.aR.sup.b; G is N;  custom-character

joining G and J can be a double bond; R^{sup.a} can be hydrogen R^{sup.b} can be —CH_{sub.2}CH_{sub.2}—R^{sup.c}; R^{sup.c} can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R^{sup.K} can be substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R^{sup.K} moiety indicated as substituted is substituted with one or more Q, wherein Q can be cyano; A can be CH; B can be CH; R⁹ can be hydrogen; J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I-C) can be 5-(4-((2-(1H-indol-3-yl)ethyl)amino)quinazolin-2-yl)nicotinonitrile.

[0340] In some embodiments, when R^{sup.J} is —NR^{sup.a}R^{sup.b}; G can be N;  joining G and J can be a double bond; R^{sup.a} can be hydrogen R^{sup.b} can be —CH_{sub.2}CH_{sub.2}—R^{sup.c}; R^{sup.c} can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R^{sup.K} can be —NH(C_{sub.1-4} alkyl); A can be CH; B can be CH; R^{sup.g} can be hydrogen; J can be C; X can be N; Y can be C; and Z can be C. In some embodiments, the compound of Formula (I-C) can be N^{sup.4}-(2-(1H-indol-3-yl)ethyl)-N^{sup.2}-(sec-butyl)quinazoline-2,4-diamine.

[0341] In some embodiments, the compound of Formula (I-C), or a pharmaceutically acceptable salt thereof, can be selected from the group consisting of: [0342] 4-(2-((2-(benzo[b]thiophen-3-yl)-8-(dimethylamino)pyrimido[5,4-d]pyrimidin-4-yl)amino)ethyl)phenol; [0343] N-(2-(1H-indol-3-yl)ethyl)-2-(5-fluoropyridin-3-yl)quinazolin-4-amine; [0344] 5-(4-((2-(1H-indol-3-yl)ethyl)amino)quinazolin-2-yl)nicotinonitrile; and [0345] N^{sup.4}-(2-(1H-indol-3-yl)ethyl)-N-(sec-butyl)quinazoline-2,4-diamine.

Formula (I-D)

[0346] In yet still other embodiments provided herein, the compound of Formula (I) can have the structure of Formula (I-D):

##STR00059##

including pharmaceutically acceptable salts thereof, wherein: R^{sup.J} can be —NR^{sup.a}R^{sup.b}; R^{sup.a} can be hydrogen or C_{sub.1-4} alkyl; R^{sup.b} can be R^{sup.c} or —(C_{sub.1-4} alkyl)-R^{sup.c}; R^{sup.c} can be selected from the group consisting of: unsubstituted C_{sub.6-10} aryl; substituted C_{sub.6-10} aryl; unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R^{sup.c} moiety indicated as substituted is substituted with one or more substituents E, wherein each E can be independently selected from the group consisting of: —OH, C_{sub.1-4} alkyl, C_{sub.1-4} haloalkyl, —O(C_{sub.1-4} alkyl), and —O(C_{sub.1-4} haloalkyl); R^{sup.K} can be selected from the group consisting of: unsubstituted C_{sub.6-10} aryl; substituted C_{sub.6-10} aryl; unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R^{sup.K} moiety indicated as substituted is substituted with one or more substituents Q, wherein each Q can be independently selected from the group consisting of: —OH, C_{sub.1-4} alkyl, C_{sub.1-4} haloalkyl, halo, cyano, —O—(C_{sub.1-4} alkyl), and —O—(C_{sub.1-4} haloalkyl); R^{sup.h} can be hydrogen or C_{sub.1-4} alkyl; D can be N or CH; Y can be N; Z can be C; and X can be N or CH.

[0347] In some embodiments, R^{sup.h} can be hydrogen. In other embodiments, R^{sup.h} can be C_{sub.1-4} alkyl. For example, R^{sup.h} can be methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl or tert-butyl.

[0348] In some embodiments, D can be N. In other embodiments, D can be CH.

[0349] In some embodiments, when D is N, Y can be N, Z can be C, and X can be N. In other embodiments, when D is N, Y can be N, Z can be C, and X can be CH. In some embodiments, when D is CH, Y can be N, Z can be C, and X can be N. In other embodiments, when D is CH, Y can be N, Z can be C, and X can be CH.

[0350] In some embodiments, R^{sup.a} can be hydrogen; R^{sup.b} can be —(C_{sub.1-4} alkyl)-R^{sup.c}; R^{sup.c} can be selected from the group consisting of: unsubstituted C_{sub.6-10} aryl; substituted C_{sub.6-10} aryl; unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R^{sup.c} moiety indicated as substituted is substituted with one or more substituents E, wherein each E can be independently selected from the group consisting of: —OH, C_{sub.1-4} alkyl, C_{sub.1-4} haloalkyl, —O(C_{sub.1-4} alkyl), and —O(C_{sub.1-4} haloalkyl); R^{sup.K} can be selected from the group consisting of: unsubstituted C_{sub.6-10} aryl; substituted C_{sub.6-10} aryl; unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; and substituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; wherein a R^{sup.K} moiety indicated as substituted is substituted with one or more substituents Q, wherein each Q can be independently selected from the group consisting of: —OH, C_{sub.1-4} alkyl, C_{sub.1-4} haloalkyl, halo, cyano, —O—(C_{sub.1-4} alkyl), and —O—(C_{sub.1-4} haloalkyl); and R^{sup.h} can be hydrogen or C_{sub.1-4} alkyl.

[0351] In some embodiments, R^{sup.a} can be hydrogen; R^{sup.b} can be —(C_{sub.1-4} alkyl)-R^{sup.c}; R^{sup.c} can be selected from the group consisting of: substituted phenyl and unsubstituted indolyl; wherein the substituted phenyl is substituted with one or more substituents E, wherein each E can be independently selected from the group consisting of: —OH, C_{sub.1-4} alkyl, C_{sub.1-4} haloalkyl, —O(C_{sub.1-4} alkyl), and —O(C_{sub.1-4} haloalkyl); R^{sup.K} can be unsubstituted benzothiophenyl; and R^{sup.h} can be hydrogen or C_{sub.1-4} alkyl.

[0352] In some embodiments, R^{sup.a} can be hydrogen; R^{sup.b} can be —(CH_{sub.2}—CH_{sub.2})—R^{sup.c}; R^{sup.c} can be selected from the group consisting of: substituted phenyl and unsubstituted indolyl; wherein the substituted phenyl is substituted with one substituent E, wherein E can be —OH; R^{sup.K} can be unsubstituted benzothiophenyl; and R^{sup.h} can be hydrogen or C_{sub.1-4} alkyl.

[0353] In some embodiments, when D is N; R^{sup.J} is —NR^{sup.a}R^{sup.b}; G can be N; R^{sup.a} can be hydrogen; R^{sup.b} can be —CH_{sub.2}CH_{sub.2}—R^{sup.c}; R^{sup.c} can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; or substituted C_{sub.6-10} aryl, substituted with one or more E, wherein E is —OH; R^{sup.K} can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; R^{sup.h} can be C_{sub.1-4} alkyl; J can be C; X can be C; Y can be N; and Z can be C; wherein the valency of any carbon atom is filled as needed with hydrogen atoms.

[0354] In some embodiments, when R^{sup.J} is —NR^{sup.a}R^{sup.b}; G can be N; R^{sup.a} can be hydrogen; R^{sup.b} can be —CH_{sub.2}CH_{sub.2}—R^{sup.c}; R^{sup.c} can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S or substituted C_{sub.6-10} aryl, substituted with one or more E, wherein E is —OH; R^{sup.K} can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; D can be N; R^{sup.h} can be C_{sub.1-4} alkyl; J can be C; X can be C; Y can be N; and Z can be C; wherein the valency of any carbon atom is filled as needed with hydrogen atoms. In some embodiments, the compound of Formula (I-D) can be N-(2-(1H-indol-3-yl)ethyl)-6-(benzo[b]thiophen-3-yl)-3-isopropylimidazo[1,5-a]pyrazin-8-amine.

[0355] In some embodiments, when R^{sup.J} is —NR^{sup.a}R^{sup.b}; G can be N;  joining G and J can be a double bond; R^{sup.a} can be hydrogen; R^{sup.b} can be —CH_{sub.2}CH_{sub.2}—R^{sup.c}; R^{sup.c} can be substituted C_{sub.6-10} aryl, substituted with one or more E, wherein E is —OH; R^{sup.K} can be unsubstituted five- to ten-membered heteroaryl having 1-4 atoms selected from the group consisting of O, N, and S; D can be N; R^{sup.h} can be C_{sub.1-4} alkyl; J can be C; X can be C; Y can be N; and Z can be C; wherein the valency of any carbon atom is filled as needed with hydrogen atoms. In some embodiments, the compound of Formula (I-D)

can be 4-(2-((6-(benzo[b]thiophen-3-yl)-3-isopropylimidazo[1,5-a]pyrazin-8-yl)amino)ethyl)phenol.

[0356] In some embodiments, the compound of Formula (I-D), or a pharmaceutically acceptable salt thereof, can be selected from the group consisting of: [0357] N-(2-(1H-indol-3-yl)ethyl)-6-(benzo[b]thiophen-3-yl)-3-isopropylimidazo[1,5-a]pyrazin-8-amine; and 4-(2-((6-(benzo[b]thiophen-3-yl)-3-isopropylimidazo[1,5-a]pyrazin-8-yl)amino)ethyl)phenol.

[0358] The compounds provided herein may be enantiomerically pure, such as a single enantiomer or a single diastereomer, or be stereoisomeric mixtures, such as a mixture of enantiomers, e.g., a racemic mixture of two enantiomers; or a mixture of two or more diastereomers. As such, one of skill in the art will recognize that administration of a compound in its (R) form is equivalent, for compounds that undergo epimerization in vivo, to administration of the compound in its (S) form. Conventional techniques for the preparation/isolation of individual enantiomers include synthesis from a suitable optically pure precursor, asymmetric synthesis from achiral starting materials, or resolution of an enantiomeric mixture, for example, chiral chromatography, recrystallization, resolution, diastereomeric salt formation, or derivatization into diastereomeric adducts followed by separation.

5.4. Isolation of NK Cells

[0359] Methods of isolating natural killer cells are known in the art and can be used to isolate the natural killer cells, e.g., NK cells produced using the three-stage method, described herein. For example, NK cells can be isolated or enriched, for example, by staining cells, in one embodiment, with antibodies to CD56 and CD3, and selecting for CD56^{sup.}+CD3^{sup.}- cells. In certain embodiments, the NK cells are enriched for CD56^{sup.}+CD3^{sup.}- cells in comparison with total cells produced using the three-stage method, described herein. NK cells, e.g., cells produced using the three-stage method, described herein, can be isolated using a commercially available kit, for example, the NK Cell Isolation Kit (Miltenyi Biotec). NK cells, e.g., cells produced using the three-stage method, described herein, can also be isolated or enriched by removal of cells other than NK cells in a population of cells that comprise the NK cells, e.g., cells produced using the three-stage method, described herein. For example, NK cells, e.g., cells produced using the three-stage method, described herein, may be isolated or enriched by depletion of cells displaying non-NK cell markers using, e.g., antibodies to one or more of CD3, CD4, CD14, CD19, CD20, CD36, CD66b, CD123, HLA DR and/or CD235a (glycophorin A). Negative isolation can be carried out using a commercially available kit, e.g., the NK Cell Negative Isolation Kit (DynaL Biotech). Cells isolated by these methods may be additionally sorted, e.g., to separate CD11a⁺ and CD11a⁻ cells, and/or CD117⁺ and CD117⁻ cells, and/or CD16^{sup.}+ and CD16^{sup.}- cells, and/or CD94^{sup.}- and CD94^{sup.}-. In certain embodiments, cells, e.g., cells produced by the three-step methods described herein, are sorted to separate CD11a⁺ and CD11a⁻ cells. In specific embodiments, CD11a⁺ cells are isolated. In certain embodiments, the cells are enriched for CD11a^{sup.}+ cells in comparison with total cells produced using the three-stage method, described herein. In specific embodiments, CD11a⁻ cells are isolated. In certain embodiments, the cells are enriched for CD11a⁻ cells in comparison with total cells produced using the three-stage method, described herein. In certain embodiments, cells are sorted to separate CD117⁺ and CD117⁻ cells. In specific embodiments, CD117⁺ cells are isolated. In certain embodiments, the cells are enriched for CD117^{sup.}+ cells in comparison with total cells produced using the three-stage method, described herein. In specific embodiments, CD117⁻ cells are isolated. In certain embodiments, the cells are enriched for CD117⁻ cells in comparison with total cells produced using the three-stage method, described herein. In certain embodiments, cells are sorted to separate CD16^{sup.}+ and CD16^{sup.}- cells. In specific embodiments, CD16^{sup.}+ cells are isolated. In certain embodiments, the cells are enriched for CD16^{sup.}+ cells in comparison with total cells produced using the three-stage method, described herein. In specific embodiments, CD16^{sup.}- cells are isolated. In certain embodiments, the cells are enriched for CD16⁻ cells in comparison with total cells produced using

the three-stage method, described herein. In certain embodiments, cells are sorted to separate CD94.sup.+ and CD94.sup.- cells. In specific embodiments, CD94.sup.+ cells are isolated. In certain embodiments, the cells are enriched for CD94.sup.+ cells in comparison with total cells produced using the three-stage method, described herein. In specific embodiments, CD94.sup.- cells are isolated. In certain embodiments, the cells are enriched for CD94- cells in comparison with total cells produced using the three-stage method, described herein. In certain embodiments, isolation is performed using magnetic separation. In certain embodiments, isolation is performed using flow cytometry.

[0360] Methods of isolating ILC3 cells are known in the art and can be used to isolate the ILC3 cells, e.g., ILC3 cells produced using the three-stage method, described herein. For example, ILC3 cells can be isolated or enriched, for example, by staining cells, in one embodiment, with antibodies to CD56, CD3, and CD11a, and selecting for CD56.sup.+CD3.sup.- CD11a.sup.- cells. ILC3 cells, e.g., cells produced using the three-stage method, described herein, can also be isolated or enriched by removal of cells other than ILC3 cells in a population of cells that comprise the ILC3 cells, e.g., cells produced using the three-stage method, described herein. For example, ILC3 cells, e.g., cells produced using the three-stage method, described herein, may be isolated or enriched by depletion of cells displaying non-ILC3 cell markers using, e.g., antibodies to one or more of CD3, CD4, CD11a, CD14, CD19, CD20, CD36, CD66b, CD94, CD123, HLA DR and/or CD235a (glycophorin A). Cells isolated by these methods may be additionally sorted, e.g., to separate CD117.sup.+ and CD117.sup.- cells. NK cells can be isolated or enriched, for example, by staining cells, in one embodiment, with antibodies to CD56, CD3, CD94, and CD11a, and selecting for CD56.sup.+CD3.sup.-CD94.sup.+CD11a.sup.+ cells. NK cells, e.g., cells produced using the three-stage method, described herein, can also be isolated or enriched by removal of cells other than NK cells in a population of cells that comprise the NK cells, e.g., cells produced using the three-stage method, described herein. In certain embodiments, the NK cells are enriched for CD56.sup.+CD3.sup.-CD94.sup.+CD11a.sup.+ cells in comparison with total cells produced using the three-stage method, described herein.

[0361] In one embodiment, ILC3 cells are isolated or enriched by selecting for CD56.sup.+CD3.sup.-CD11a.sup.- cells. In certain embodiments, the ILC3 cells are enriched for CD56.sup.+CD3.sup.-CD11a.sup.- cells in comparison with total cells produced using the three-stage method, described herein. In one embodiment, ILC3 cells are isolated or enriched by selecting for CD56.sup.+CD3.sup.-CD11a.sup.-CD117+ cells. In certain embodiments, the ILC3 cells are enriched for CD56.sup.+CD3.sup.-CD11a.sup.-CD117+ cells in comparison with total cells produced using the three-stage method, described herein. In one embodiment, ILC3 cells are isolated or enriched by selecting for CD56.sup.+CD3.sup.-CD11a.sup.-CD117CDIL1R1.sup.+ cells. In certain embodiments, the ILC3 cells are enriched for CD56.sup.+CD3.sup.-CD11a.sup.-CD117CDIL1R1.sup.+ cells in comparison with total cells produced using the three-stage method, described herein.

[0362] In one embodiment, NK cells are isolated or enriched by selecting for CD56.sup.+CD3.sup.-CD94.sup.+CD11a.sup.+ cells. In certain embodiments, the NK cells are enriched for CD56.sup.+CD3.sup.-CD94.sup.+CD11a.sup.+ cells in comparison with total cells produced using the three-stage method, described herein. In one embodiment, NK cells are isolated or enriched by selecting for CD56.sup.+CD3.sup.-CD94.sup.+CD11a.sup.+CD117.sup.- cells. In certain embodiments, the NK cells are enriched for CD56.sup.+CD3.sup.-CD94.sup.+CD11a.sup.+CD117.sup.- cells in comparison with total cells produced using the three-stage method, described herein.

[0363] Cell separation can be accomplished by, e.g., flow cytometry, fluorescence-activated cell sorting (FACS), or, in one embodiment, magnetic cell sorting using microbeads conjugated with specific antibodies. The cells may be isolated, e.g., using a magnetic activated cell sorting (MACS) technique, a method for separating particles based on their ability to bind magnetic beads (e.g.,

about 0.5-100 μm diameter) that comprise one or more specific antibodies, e.g., anti-CD56 antibodies. Magnetic cell separation can be performed and automated using, e.g., an AUTOMACS™ Separator (Miltenyi). A variety of useful modifications can be performed on the magnetic microspheres, including covalent addition of antibody that specifically recognizes a particular cell surface molecule or hapten. The beads are then mixed with the cells to allow binding. Cells are then passed through a magnetic field to separate out cells having the specific cell surface marker. In one embodiment, these cells can then be isolated and re-mixed with magnetic beads coupled to an antibody against additional cell surface markers. The cells are again passed through a magnetic field, isolating cells that bound both the antibodies. Such cells can then be diluted into separate dishes, such as microtiter dishes for clonal isolation.

5.5. Placental Perfusate

[0364] NK cells and/or ILC3 cells, e.g., NK cell and/or ILC3 cell populations produced according to the three-stage method described herein may be produced from hematopoietic cells, e.g., hematopoietic stem or progenitors from any source, e.g., placental tissue, placental perfusate, umbilical cord blood, placental blood, peripheral blood, spleen, liver, or the like. In certain embodiments, the hematopoietic stem cells are combined hematopoietic stem cells from placental perfusate and from cord blood from the same placenta used to generate the placental perfusate. Placental perfusate comprising placental perfusate cells that can be obtained, for example, by the methods disclosed in U.S. Pat. Nos. 7,045,148 and 7,468,276 and U.S. Patent Application Publication No. 2009/0104164, the disclosures of which are hereby incorporated in their entireties.

5.5.1. Cell Collection Composition

[0365] The placental perfusate and perfusate cells, from which hematopoietic stem or progenitors may be isolated, or useful in tumor suppression or the treatment of an individual having tumor cells, cancer or a viral infection, e.g., in combination with the NK cells and/or ILC3 cells, e.g., NK cell and/or ILC3 cell populations produced according to the three-stage method provided herein, can be collected by perfusion of a mammalian, e.g., human post-partum placenta using a placental cell collection composition. Perfusate can be collected from the placenta by perfusion of the placenta with any physiologically-acceptable solution, e.g., a saline solution, culture medium, or a more complex cell collection composition. A cell collection composition suitable for perfusing a placenta, and for the collection and preservation of perfusate cells is described in detail in related U.S. Application Publication No. 2007/0190042, which is incorporated herein by reference in its entirety.

[0366] The cell collection composition can comprise any physiologically-acceptable solution suitable for the collection and/or culture of stem cells, for example, a saline solution (e.g., phosphate-buffered saline, Krebs's solution, modified Krebs's solution, Eagle's solution, 0.9% NaCl, etc.), a culture medium (e.g., DMEM, H.DMEM, etc.), and the like.

[0367] The cell collection composition can comprise one or more components that tend to preserve placental cells, that is, prevent the placental cells from dying, or delay the death of the placental cells, reduce the number of placental cells in a population of cells that die, or the like, from the time of collection to the time of culturing. Such components can be, e.g., an apoptosis inhibitor (e.g., a caspase inhibitor or JNK inhibitor); a vasodilator (e.g., magnesium sulfate, an antihypertensive drug, atrial natriuretic peptide (ANP), adrenocorticotropin, corticotropin-releasing hormone, sodium nitroprusside, hydralazine, adenosine triphosphate, adenosine, indomethacin or magnesium sulfate, a phosphodiesterase inhibitor, etc.); a necrosis inhibitor (e.g., 2-(1H-Indol-3-yl)-3-pentylamino-maleimide, pyrrolidine dithiocarbamate, or clonazepam); a TNF- α inhibitor; and/or an oxygen-carrying perfluorocarbon (e.g., perfluorooctyl bromide, perfluorodecyl bromide, etc.).

[0368] The cell collection composition can comprise one or more tissue-degrading enzymes, e.g., a metalloprotease, a serine protease, a neutral protease, a hyaluronidase, an RNase, or a DNase, or the like. Such enzymes include, but are not limited to, collagenases (e.g., collagenase I, II, III or IV, a collagenase from *Clostridium histolyticum*, etc.); dispase, thermolysin, elastase, trypsin,

LIBERASE, hyaluronidase, and the like.

[0369] The cell collection composition can comprise a bacteriocidally or bacteriostatically effective amount of an antibiotic. In certain non-limiting embodiments, the antibiotic is a macrolide (e.g., tobramycin), a cephalosporin (e.g., cephalexin, cephadrine, cefuroxime, cefprozil, cefaclor, cefixime or cefadroxil), a clarithromycin, an erythromycin, a penicillin (e.g., penicillin V) or a quinolone (e.g., ofloxacin, ciprofloxacin or norfloxacin), a tetracycline, a streptomycin, etc. In a particular embodiment, the antibiotic is active against Gram(+) and/or Gram(-) bacteria, e.g., *Pseudomonas aeruginosa*, *Staphylococcus aureus*, and the like.

[0370] The cell collection composition can also comprise one or more of the following compounds: adenosine (about 1 mM to about 50 mM); D-glucose (about 20 mM to about 100 mM); magnesium ions (about 1 mM to about 50 mM); a macromolecule of molecular weight greater than 20,000 daltons, in one embodiment, present in an amount sufficient to maintain endothelial integrity and cellular viability (e.g., a synthetic or naturally occurring colloid, a polysaccharide such as dextran or a polyethylene glycol present at about 25 g/l to about 100 g/l, or about 40 g/l to about 60 g/l); an antioxidant (e.g., butylated hydroxyanisole, butylated hydroxytoluene, glutathione, vitamin C or vitamin E present at about 25 μ M to about 100 μ M); a reducing agent (e.g., N-acetylcysteine present at about 0.1 mM to about 5 mM); an agent that prevents calcium entry into cells (e.g., verapamil present at about 2 μ M to about 25 μ M); nitroglycerin (e.g., about 0.05 g/L to about 0.2 g/L); an anticoagulant, in one embodiment, present in an amount sufficient to help prevent clotting of residual blood (e.g., heparin or hirudin present at a concentration of about 1000 units/l to about 100,000 units/l); or an amiloride containing compound (e.g., amiloride, ethyl isopropyl amiloride, hexamethylene amiloride, dimethyl amiloride or isobutyl amiloride present at about 1.0 μ M to about 5 μ M).

5.5.2. Collection and Handling of Placenta

[0371] Generally, a human placenta is recovered shortly after its expulsion after birth. In one embodiment, the placenta is recovered from a patient after informed consent and after a complete medical history of the patient is taken and is associated with the placenta. In one embodiment, the medical history continues after delivery.

[0372] Prior to recovery of perfusate, the umbilical cord blood and placental blood are removed. In certain embodiments, after delivery, the cord blood in the placenta is recovered. The placenta can be subjected to a conventional cord blood recovery process. Typically a needle or cannula is used, with the aid of gravity, to exsanguinate the placenta (see, e.g., Anderson, U.S. Pat. No. 5,372,581; Hessel et al., U.S. Pat. No. 5,415,665). The needle or cannula is usually placed in the umbilical vein and the placenta can be gently massaged to aid in draining cord blood from the placenta. Such cord blood recovery may be performed commercially, e.g., LifeBank Inc., Cedar Knolls, N.J., ViaCord, Cord Blood Registry and CryoCell. In one embodiment, the placenta is gravity drained without further manipulation so as to minimize tissue disruption during cord blood recovery.

[0373] Typically, a placenta is transported from the delivery or birthing room to another location, e.g., a laboratory, for recovery of cord blood and collection of perfusate. The placenta can be transported in a sterile, thermally insulated transport device (maintaining the temperature of the placenta between 20-28° C.), for example, by placing the placenta, with clamped proximal umbilical cord, in a sterile zip-lock plastic bag, which is then placed in an insulated container. In another embodiment, the placenta is transported in a cord blood collection kit substantially as described in U.S. Pat. No. 7,147,626. In one embodiment, the placenta is delivered to the laboratory four to twenty-four hours following delivery. In certain embodiments, the proximal umbilical cord is clamped, for example within 4-5 cm (centimeter) of the insertion into the placental disc prior to cord blood recovery. In other embodiments, the proximal umbilical cord is clamped after cord blood recovery but prior to further processing of the placenta.

[0374] The placenta, prior to collection of the perfusate, can be stored under sterile conditions and at either room temperature or at a temperature of 5 to 25° C. (centigrade). The placenta may be

stored for a period of longer than forty eight hours, or for a period of four to twenty-four hours prior to perfusing the placenta to remove any residual cord blood. The placenta can be stored in an anticoagulant solution at a temperature of 5° C. to 25° C. (centigrade). Suitable anticoagulant solutions are well known in the art. For example, a solution of heparin or warfarin sodium can be used. In one embodiment, the anticoagulant solution comprises a solution of heparin (e.g., 1% w/w in 1:1000 solution). In some embodiments, the exsanguinated placenta is stored for no more than 36 hours before placental perfusate is collected.

5.5.3. Placental Perfusion

[0375] Methods of perfusing mammalian placentae and obtaining placental perfusate are disclosed, e.g., in Hariri, U.S. Pat. Nos. 7,045,148 and 7,255,879, and in U.S. Application Publication Nos. 2009/0104164, 2007/0190042 and 20070275362, issued as U.S. Pat. No. 8,057,788, the disclosures of which are hereby incorporated by reference herein in their entireties.

[0376] Perfusate can be obtained by passage of perfusion solution, e.g., saline solution, culture medium or cell collection compositions described above, through the placental vasculature. In one embodiment, a mammalian placenta is perfused by passage of perfusion solution through either or both of the umbilical artery and umbilical vein. The flow of perfusion solution through the placenta may be accomplished using, e.g., gravity flow into the placenta. For example, the perfusion solution is forced through the placenta using a pump, e.g., a peristaltic pump. The umbilical vein can be, e.g., cannulated with a cannula, e.g., a TEFLON® or plastic cannula, that is connected to a sterile connection apparatus, such as sterile tubing. The sterile connection apparatus is connected to a perfusion manifold.

[0377] In preparation for perfusion, the placenta can be oriented in such a manner that the umbilical artery and umbilical vein are located at the highest point of the placenta. The placenta can be perfused by passage of a perfusion solution through the placental vasculature, or through the placental vasculature and surrounding tissue. In one embodiment, the umbilical artery and the umbilical vein are connected simultaneously to a pipette that is connected via a flexible connector to a reservoir of the perfusion solution. The perfusion solution is passed into the umbilical vein and artery. The perfusion solution exudes from and/or passes through the walls of the blood vessels into the surrounding tissues of the placenta, and is collected in a suitable open vessel from the surface of the placenta that was attached to the uterus of the mother during gestation. The perfusion solution may also be introduced through the umbilical cord opening and allowed to flow or percolate out of openings in the wall of the placenta which interfaced with the maternal uterine wall. In another embodiment, the perfusion solution is passed through the umbilical veins and collected from the umbilical artery, or is passed through the umbilical artery and collected from the umbilical veins, that is, is passed through only the placental vasculature (fetal tissue).

[0378] In one embodiment, for example, the umbilical artery and the umbilical vein are connected simultaneously, e.g., to a pipette that is connected via a flexible connector to a reservoir of the perfusion solution. The perfusion solution is passed into the umbilical vein and artery. The perfusion solution exudes from and/or passes through the walls of the blood vessels into the surrounding tissues of the placenta, and is collected in a suitable open vessel from the surface of the placenta that was attached to the uterus of the mother during gestation. The perfusion solution may also be introduced through the umbilical cord opening and allowed to flow or percolate out of openings in the wall of the placenta which interfaced with the maternal uterine wall. Placental cells that are collected by this method, which can be referred to as a “pan” method, are typically a mixture of fetal and maternal cells.

[0379] In another embodiment, the perfusion solution is passed through the umbilical veins and collected from the umbilical artery, or is passed through the umbilical artery and collected from the umbilical veins. Placental cells collected by this method, which can be referred to as a “closed circuit” method, are typically almost exclusively fetal.

[0380] The closed circuit perfusion method can, in one embodiment, be performed as follows. A

post-partum placenta is obtained within about 48 hours after birth. The umbilical cord is clamped and cut above the clamp. The umbilical cord can be discarded, or can be processed to recover, e.g., umbilical cord stem cells, and/or to process the umbilical cord membrane for the production of a biomaterial. The amniotic membrane can be retained during perfusion, or can be separated from the chorion, e.g., using blunt dissection with the fingers. If the amniotic membrane is separated from the chorion prior to perfusion, it can be, e.g., discarded, or processed, e.g., to obtain stem cells by enzymatic digestion, or to produce, e.g., an amniotic membrane biomaterial, e.g., the biomaterial described in U.S. Application Publication No. 2004/0048796. After cleaning the placenta of all visible blood clots and residual blood, e.g., using sterile gauze, the umbilical cord vessels are exposed, e.g., by partially cutting the umbilical cord membrane to expose a cross-section of the cord. The vessels are identified, and opened, e.g., by advancing a closed alligator clamp through the cut end of each vessel. The apparatus, e.g., plastic tubing connected to a perfusion device or peristaltic pump, is then inserted into each of the placental arteries. The pump can be any pump suitable for the purpose, e.g., a peristaltic pump. Plastic tubing, connected to a sterile collection reservoir, e.g., a blood bag such as a 250 mL collection bag, is then inserted into the placental vein. Alternatively, the tubing connected to the pump is inserted into the placental vein, and tubes to a collection reservoir(s) are inserted into one or both of the placental arteries. The placenta is then perfused with a volume of perfusion solution, e.g., about 750 ml of perfusion solution. Cells in the perfusate are then collected, e.g., by centrifugation.

[0381] In one embodiment, the proximal umbilical cord is clamped during perfusion, and, more specifically, can be clamped within 4-5 cm (centimeter) of the cord's insertion into the placental disc.

[0382] The first collection of perfusion fluid from a mammalian placenta during the exsanguination process is generally colored with residual red blood cells of the cord blood and/or placental blood. The perfusion fluid becomes more colorless as perfusion proceeds and the residual cord blood cells are washed out of the placenta. Generally from 30 to 100 mL of perfusion fluid is adequate to initially flush blood from the placenta, but more or less perfusion fluid may be used depending on the observed results.

[0383] In certain embodiments, cord blood is removed from the placenta prior to perfusion (e.g., by gravity drainage), but the placenta is not flushed (e.g., perfused) with solution to remove residual blood. In certain embodiments, cord blood is removed from the placenta prior to perfusion (e.g., by gravity drainage), and the placenta is flushed (e.g., perfused) with solution to remove residual blood.

[0384] The volume of perfusion liquid used to perfuse the placenta may vary depending upon the number of placental cells to be collected, the size of the placenta, the number of collections to be made from a single placenta, etc. In various embodiments, the volume of perfusion liquid may be from 50 mL to 5000 mL, 50 mL to 4000 mL, 50 mL to 3000 mL, 100 mL to 2000 mL, 250 mL to 2000 mL, 500 mL to 2000 mL, or 750 mL to 2000 mL. Typically, the placenta is perfused with 700-800 mL of perfusion liquid following exsanguination.

[0385] The placenta can be perfused a plurality of times over the course of several hours or several days. Where the placenta is to be perfused a plurality of times, it may be maintained or cultured under aseptic conditions in a container or other suitable vessel, and perfused with a cell collection composition, or a standard perfusion solution (e.g., a normal saline solution such as phosphate buffered saline ("PBS") with or without an anticoagulant (e.g., heparin, warfarin sodium, coumarin, bishydroxycoumarin), and/or with or without an antimicrobial agent (e.g., β -mercaptoethanol (0.1 mM); antibiotics such as streptomycin (e.g., at 40-100 μ g/ml), penicillin (e.g., at 40 U/ml), amphotericin B (e.g., at 0.5 μ g/ml). In one embodiment, an isolated placenta is maintained or cultured for a period of time without collecting the perfusate, such that the placenta is maintained or cultured for 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or 24 hours, or 2 or 3 or more days before perfusion and collection of perfusate. The perfused placenta

can be maintained for one or more additional time(s), e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24 or more hours, and perfused a second time with, e.g., 700-800 mL perfusion fluid. The placenta can be perfused 1, 2, 3, 4, 5 or more times, for example, once every 1, 2, 3, 4, 5 or 6 hours. In one embodiment, perfusion of the placenta and collection of perfusion solution, e.g., placental cell collection composition, is repeated until the number of recovered nucleated cells falls below 100 cells/mL. The perfusates at different time points can be further processed individually to recover time-dependent populations of cells, e.g., total nucleated cells. Perfusates from different time points can also be pooled.

5.5.4. Placental Perfusate and Placental Perfusate Cells

[0386] Typically, placental perfusate from a single placental perfusion comprises about 100 million to about 500 million nucleated cells, including hematopoietic cells from which NK cells and/or ILC3 cells, e.g., NK cells and/or ILC3 cells produced according to the three-stage method described herein, may be produced by the method disclosed herein. In certain embodiments, the placental perfusate or perfusate cells comprise CD34.sup.+ cells, e.g., hematopoietic stem or progenitor cells. Such cells can, in a more specific embodiment, comprise CD34.sup.+CD45.sup.- stem or progenitor cells, CD34.sup.+CD45.sup.+ stem or progenitor cells, or the like. In certain embodiments, the perfusate or perfusate cells are cryopreserved prior to isolation of hematopoietic cells therefrom. In certain other embodiments, the placental perfusate comprises, or the perfusate cells comprise, only fetal cells, or a combination of fetal cells and maternal cells.

5.6. NK Cells

5.6.1. NK Cells Produced by Three-Stage Method

[0387] In another embodiment, provided herein is an isolated NK cell population, wherein said NK cells are produced according to the three-stage method described above.

[0388] In one embodiment, provided herein is an isolated NK cell population produced by a three-stage method described herein, wherein said NK cell population comprises a greater percentage of CD3-CD56+ cells than an NK progenitor cell population produced by a three-stage method described herein, e.g., an NK progenitor cell population produced by the same three-stage method with the exception that the third culture step used to produce the NK progenitor cell population was of shorter duration than the third culture step used to produce the NK cell population. In a specific embodiment, said NK cell population comprises about 70% or more, in some embodiments, 75%, 80%, 85%, 90%, 95%, 98%, or 99% CD3-CD56+ cells. In another specific embodiment, said NK cell population comprises no less than 80%, 85%, 90%, 95%, 98%, or 99% CD3-CD56+ cells. In another specific embodiment, said NK cell population comprises between 70%-75%, 75%-80%, 80%-85%, 85%-90%, 90%-95%, or 95%-99% CD3-CD56+ cells.

[0389] In certain embodiments, said CD3.sup.-CD56.sup.- cells in said NK cell population comprises CD3.sup.-CD56.sup.+ cells that are additionally NKp46.sup.+. In certain embodiments, said CD3.sup.-CD56.sup.+ cells in said NK cell population comprises CD3.sup.-CD56.sup.+ cells that are additionally CD16-. In certain embodiments, said CD3.sup.-CD56.sup.+ cells in said NK cell population comprises CD3.sup.-CD56.sup.- cells that are additionally CD16+. In certain embodiments, said CD3.sup.-CD56.sup.+ cells in said NK cell population comprises CD3.sup.-CD56.sup.+ cells that are additionally CD94-. In certain embodiments, said CD3.sup.-CD56.sup.+ cells in said NK cell population comprises CD3.sup.-CD56.sup.+ cells that are additionally CD94+. In certain embodiments, said CD3.sup.-CD56.sup.+ cells in said NK cell population comprises CD3.sup.-CD56.sup.+ cells that are additionally CD11a.sup.+. In certain embodiments, said CD3.sup.-CD56.sup.+ cells in said NK cell population comprises CD3.sup.-CD56.sup.+ cells that are additionally NKp30.sup.+. In certain embodiments, said CD3.sup.-CD56.sup.+ cells in said NK cell population comprises CD3.sup.-CD56.sup.+ cells that are additionally CD161.sup.+. In certain embodiments, said CD3.sup.-CD56.sup.+ cells in said NK cell population comprises CD3.sup.-CD56.sup.+ cells that are additionally DNAM-1.sup.+. In certain embodiments, said CD3.sup.-CD56.sup.+ cells in said NK cell population comprises

CD3.sup.-CD56.sup.+ cells that are additionally T-bet.sup.+.

[0390] In one embodiment, an NK cell population produced by a three-stage method described herein comprises cells which are CD117+. In one embodiment, an NK cell population produced by a three-stage method described herein comprises cells which are NKG2D+. In one embodiment, an NK cell population produced by a three-stage method described herein comprises cells which are NKp44+. In one embodiment, an NK cell population produced by a three-stage method described herein comprises cells which are CD244+. In one embodiment, an NK cell population produced by a three-stage method described herein comprises cells which express perforin. In one embodiment, an NK cell population produced by a three-stage method described herein comprises cells which express EOMES. In one embodiment, an NK cell population produced by a three-stage method described herein comprises cells which express granzyme B. In one embodiment, an NK cell population produced by a three-stage method described herein comprises cells which secrete IFN γ , GM-CSF and/or TNF α .

5.7. ILC3 Cells

5.7.1. ILC3 Cells Produced by Three-Stage Method

[0391] In another embodiment, provided herein is an isolated ILC3 cell population, wherein said ILC3 cells are produced according to the three-stage method described above.

[0392] In one embodiment, provided herein is an isolated ILC3 cell population produced by a three-stage method described herein, wherein said ILC3 cell population comprises a greater percentage of CD3-CD56+ cells than an ILC3 progenitor cell population produced by a three-stage method described herein, e.g., an ILC3 progenitor cell population produced by the same three-stage method with the exception that the third culture step used to produce the ILC3 progenitor cell population was of shorter duration than the third culture step used to produce the ILC3 cell population. In a specific embodiment, said ILC3 cell population comprises about 70% or more, in some embodiments, 75%, 80%, 85%, 90%, 95%, 98%, or 99% CD3-CD56+ cells. In another specific embodiment, said ILC3 cell population comprises no less than 80%, 85%, 90%, 95%, 98%, or 99% CD3-CD56+ cells. In another specific embodiment, said ILC3 cell population comprises between 70%-75%, 75%-80%, 80%-85%, 85%-90%, 90%-95%, or 95%-99% CD3-CD56+ cells.

[0393] In certain embodiments, said CD3.sup.-CD56.sup.+ cells in said ILC3 cell population comprises CD3.sup.-CD56.sup.+ cells that are additionally NKp46.sup.-. In certain embodiments, said CD3.sup.-CD56.sup.+ cells in said ILC3 cell population comprises CD3.sup.-CD56.sup.+ cells that are additionally CD16-. In certain embodiments, said CD3.sup.-CD56.sup.+ cells in said ILC3 cell population comprises CD3.sup.-CD56.sup.+ cells that are additionally IL1R1+. In certain embodiments, said CD3.sup.-CD56.sup.+ cells in said ILC3 cell population comprises CD3.sup.-CD56.sup.+ cells that are additionally CD94-. In certain embodiments, said CD3.sup.-CD56.sup.+ cells in said ILC3 cell population comprises CD3.sup.-CD56.sup.+ cells that are additionally ROR γ t+. In certain embodiments, said CD3.sup.-CD56.sup.+ cells in said ILC3 cell population comprises CD3.sup.-CD56.sup.+ cells that are additionally CD11a.sup.-. In certain embodiments, said CD3.sup.-CD56.sup.+ cells in said ILC3 cell population comprises CD3.sup.-CD56.sup.+ cells that are additionally T-bet+.

[0394] In one embodiment, an ILC3 cell population produced by a three-stage method described herein comprises cells which are CD117+. In one embodiment, an ILC3 cell population produced by a three-stage method described herein comprises cells which are NKG2D.sup.-. In one embodiment, an ILC3 cell population produced by a three-stage method described herein comprises cells which are NKp30.sup.-. In one embodiment, an ILC3 cell population produced by a three-stage method described herein comprises cells which are CD244+. In one embodiment, an ILC3 cell population produced by a three-stage method described herein comprises cells which are DNAM-1+. In one embodiment, an ILC3 cell population produced by a three-stage method described herein comprises cells which express AHR. In one embodiment, an ILC3 cell population

produced by a three-stage method described herein comprises cells which do not express perforin. In one embodiment, an ILC3 cell population produced by a three-stage method described herein comprises cells which do not express EOMES. In one embodiment, an ILC3 cell population produced by a three-stage method described herein comprises cells which do not express granzyme B. In one embodiment, an ILC3 cell population produced by a three-stage method described herein comprises cells which secrete IL-22 and/or IL-8.

[0395] In certain aspects, cell populations produced by the three-stage method described herein comprise CD11a⁺ cells and CD11a⁻ cells in a ratio of 50:1, 40:1, 30:1, 20:1, 10:1, 5:1, 4:1, 3:1, 2:1, 1:1, 1:2, 1:3, 1:4, 1:5, 1:10, 1:20, 1:30, 1:40, or 1:50. In certain aspects, a population of cells described herein comprises CD11a⁺ cells and CD11a^{sup.-} cells in a ratio of 50:1. In certain aspects, a population of cells described herein comprises CD11a⁺ cells and CD11a⁻ cells in a ratio of 20:1. In certain aspects, a population of cells described herein comprises CD11a⁺ cells and CD11a⁻ cells in a ratio of 10:1. In certain aspects, a population of cells described herein comprises CD11a⁺ cells and CD11a⁻ cells in a ratio of 5:1. In certain aspects, a population of cells described herein comprises CD11a⁺ cells and CD11a⁻ cells in a ratio of 1:1. In certain aspects, a population of cells described herein comprises CD11a⁺ cells and CD11a⁻ cells in a ratio of 1:5. In certain aspects, a population of cells described herein comprises CD11a⁺ cells and CD11a⁻ cells in a ratio of 1:10. In certain aspects, a population of cells described herein comprises CD11a⁺ cells and CD11a⁻ cells in a ratio of 1:20. In certain aspects, a population of cells described herein comprises CD11a⁺ cells and CD11a⁻ cells in a ratio of 1:50.

[0396] In certain aspects, cell populations described herein are produced by combining the CD11a⁺ cells with the CD11a⁻ cells in a ratio of 50:1, 40:1, 30:1, 20:1, 10:1, 5:1, 4:1, 3:1, 2:1, 1:1, 1:2, 1:3, 1:4, 1:5, 1:10, 1:20, 1:30, 1:40, or 1:50 to produce a combined population of cells. In certain aspects, a combined population of cells described herein comprises CD11a⁺ cells and CD11a⁻ cells combined in a ratio of 50:1. In certain aspects, a combined population of cells described herein comprises CD11a⁺ cells and CD11a⁻ cells combined in a ratio of 20:1. In certain aspects, a combined population of cells described herein comprises CD11a⁺ cells and CD11a⁻ cells combined in a ratio of 10:1. In certain aspects, a combined population of cells described herein comprises CD11a⁺ cells and CD11a⁻ cells combined in a ratio of 5:1. In certain aspects, a combined population of cells described herein comprises CD11a⁺ cells and CD11a⁻ cells combined in a ratio of 1:1. In certain aspects, a combined population of cells described herein comprises CD11a⁺ cells and CD11a⁻ cells combined in a ratio of 1:5. In certain aspects, a combined population of cells described herein comprises CD11a⁺ cells and CD11a⁻ cells combined in a ratio of 1:10. In certain aspects, a combined population of cells described herein comprises CD11a⁺ cells and CD11a⁻ cells combined in a ratio of 1:20. In certain aspects, a combined population of cells described herein comprises CD11a⁺ cells and CD11a⁻ cells combined in a ratio of 1:50.

[0397] In certain aspects, cell populations produced by the three-stage method described herein comprise NK cells and ILC3 cells in a ratio of 50:1, 40:1, 30:1, 20:1, 10:1, 5:1, 4:1, 3:1, 2:1, 1:1, 1:2, 1:3, 1:4, 1:5, 1:10, 1:20, 1:30, 1:40, or 1:50. In certain aspects, a population of cells described herein comprises NK cells and ILC3 cells in a ratio of 50:1. In certain aspects, a population of cells described herein comprises NK cells and ILC3 cells in a ratio of 20:1. In certain aspects, a population of cells described herein comprises NK cells and ILC3 cells in a ratio of 10:1. In certain aspects, a population of cells described herein comprises NK cells and ILC3 cells in a ratio of 5:1. In certain aspects, a population of cells described herein comprises NK cells and ILC3 cells in a ratio of 1:1. In certain aspects, a population of cells described herein comprises NK cells and ILC3 cells in a ratio of 1:5. In certain aspects, a population of cells described herein comprises NK cells and ILC3 cells in a ratio of 1:10. In certain aspects, a population of cells described herein comprises NK cells and ILC3 cells in a ratio of 1:20. In certain aspects, a population of cells described herein comprises NK cells and ILC3 cells in a ratio of 1:50.

[0398] In certain aspects, cell populations described herein are produced by combining the NK cells with the ILC3 cells in a ratio of 50:1, 40:1, 30:1, 20:1, 10:1, 5:1, 4:1, 3:1, 2:1, 1:1, 1:2, 1:3, 1:4, 1:5, 1:10, 1:20, 1:30, 1:40, or 1:50 to produce a combined population of cells. In certain aspects, a combined population of cells described herein comprises NK cells and ILC3 cells combined in a ratio of 50:1. In certain aspects, a combined population of cells described herein comprises NK cells and ILC3 cells combined in a ratio of 20:1. In certain aspects, a combined population of cells described herein comprises NK cells and ILC3 cells combined in a ratio of 10:1. In certain aspects, a combined population of cells described herein comprises NK cells and ILC3 cells combined in a ratio of 5:1. In certain aspects, a combined population of cells described herein comprises NK cells and ILC3 cells combined in a ratio of 1:1. In certain aspects, a combined population of cells described herein comprises NK cells and ILC3 cells combined in a ratio of 1:5. In certain aspects, a combined population of cells described herein comprises NK cells and ILC3 cells combined in a ratio of 1:10. In certain aspects, a combined population of cells described herein comprises NK cells and ILC3 cells combined in a ratio of 1:20. In certain aspects, a combined population of cells described herein comprises NK cells and ILC3 cells combined in a ratio of 1:50.

5.8. Compositions Comprising NK Cells and/or ILC3 Cells

5.8.1. NK Cells and/or ILC3 Cells Produced Using the Three-Stage Method

[0399] In some embodiments, provided herein is a composition, e.g., a pharmaceutical composition, comprising an isolated NK cell and/or ILC3 cell population produced using the three-stage method described herein. In a specific embodiment, said isolated NK cell and/or ILC3 cell population is produced from hematopoietic cells, e.g., hematopoietic stem or progenitor cells isolated from placental perfusate, umbilical cord blood, and/or peripheral blood. In another specific embodiment, said isolated NK cell and/or ILC3 cell population comprises at least 50% of cells in the composition. In another specific embodiment, said isolated NK cell and/or ILC3 cell population, e.g., CD3^{sup}-CD56^{sup}+ cells, comprises at least 80%, 85%, 90%, 95%, 98% or 99% of cells in the composition. In certain embodiments, no more than 5%, 10%, 15%, 20%, 25%, 30%, 35%, or 40% of the cells in said isolated NK cell and/or ILC3 cell population are CD3^{sup}-CD56^{sup}+ cells. In certain embodiments, said CD3^{sup}-CD56^{sup}+ cells are CD16^{sup}-.

[0400] NK cell and/or ILC3 cell populations produced using the three-stage method described herein, can be formulated into pharmaceutical compositions for use in vivo. Such pharmaceutical compositions comprise a population of NK cells and/or ILC3 cells in a pharmaceutically-acceptable carrier, e.g., a saline solution or other accepted physiologically-acceptable solution for in vivo administration. Pharmaceutical compositions of the invention can comprise any of the NK cell and/or ILC3 cell populations described elsewhere herein.

[0401] The pharmaceutical compositions of the invention comprise populations of cells that comprise 50% viable cells or more (that is, at least 50% of the cells in the population are functional or living). Preferably, at least 60% of the cells in the population are viable. More preferably, at least 70%, 80%, 90%, 95%, or 99% of the cells in the population in the pharmaceutical composition are viable.

[0402] The pharmaceutical compositions of the invention can comprise one or more compounds that, e.g., facilitate engraftment; stabilizers such as albumin, dextran 40, gelatin, hydroxyethyl starch, and the like.

[0403] When formulated as an injectable solution, in one embodiment, the pharmaceutical composition of the invention comprises about 1.25% HSA and about 2.5% dextran. Other injectable formulations, suitable for the administration of cellular products, may be used.

[0404] In one embodiment, the compositions, e.g., pharmaceutical compositions, provided herein are suitable for systemic or local administration. In specific embodiments, the compositions, e.g., pharmaceutical compositions, provided herein are suitable for parenteral administration. In specific embodiments, the compositions, e.g., pharmaceutical compositions, provided herein are suitable for

injection, infusion, intravenous (IV) administration, intratumoral administration, or intratumor administration. In specific embodiments, the compositions, e.g., pharmaceutical compositions, provided herein are suitable for administration via a device, a matrix, or a scaffold. In specific embodiments, the compositions, e.g., pharmaceutical compositions provided herein are suitable for injection. In specific embodiments, the compositions, e.g., pharmaceutical compositions, provided herein are suitable for administration via a catheter. In specific embodiments, the compositions, e.g., pharmaceutical compositions, provided herein are suitable for local injection. In more specific embodiments, the compositions, e.g., pharmaceutical compositions, provided herein are suitable for local injection directly into a solid tumor (e.g., a sarcoma). In specific embodiments, the compositions, e.g., pharmaceutical compositions, provided herein are suitable for injection by syringe. In specific embodiments, the compositions, e.g., pharmaceutical compositions, provided herein are suitable for administration via guided delivery. In specific embodiments, the compositions, e.g., pharmaceutical compositions, provided herein are suitable for injection aided by laparoscopy, endoscopy, ultrasound, computed tomography, magnetic resonance, or radiology.

[0405] In certain embodiments, the compositions, e.g., pharmaceutical compositions provided herein, comprising NK cells and/or ILC3 cells produced using the methods described herein, are provided as pharmaceutical grade administrable units. Such units can be provided in discrete volumes, e.g., 15 mL, 20 mL, 25 mL, 30 mL, 35 mL, 40 mL, 45 mL, 50 mL, 55 mL, 60 mL, 65 mL, 70 mL, 75 mL, 80 mL, 85 mL, 90 mL, 95 mL, 100 mL, 150 mL, 200 mL, 250 mL, 300 mL, 350 mL, 400 mL, 450 mL, 500 mL, or the like. Such units can be provided so as to contain a specified number of cells, e.g., NK cells and/or ILC3 cells, e.g., 1×10^4 , 5×10^4 , 1×10^5 , 5×10^5 , 1×10^6 , 5×10^6 , 1×10^7 , 5×10^7 , 1×10^8 , 5×10^8 or more cells per milliliter, or 1×10^4 , 5×10^4 , 1×10^5 , 5×10^5 , 1×10^6 , 5×10^6 , 1×10^7 , 5×10^7 , 1×10^8 , 5×10^8 , 1×10^9 , 5×10^9 , 1×10^{10} , 5×10^{10} , 1×10^{11} or more cells per unit. In specific embodiments, the units can comprise about, at least about, or at most about 1×10^4 , 5×10^4 , 1×10^5 , 5×10^5 , 1×10^6 , 5×10^6 or more NK cells and/or ILC3 cells per milliliter, or 1×10^4 , 5×10^4 , 1×10^5 , 5×10^5 , 1×10^6 , 5×10^6 , 1×10^7 , 5×10^7 , 1×10^8 , 5×10^8 , 1×10^9 , 5×10^9 , 1×10^{10} , 5×10^{10} , 1×10^{11} or more cells per unit. Such units can be provided to contain specified numbers of NK cells and/or ILC3 cells or NK cell and/or ILC3 cell populations and/or any of the other cells. In specific embodiments, the NK cells and ILC3 cells are present in ratios provided herein.

[0406] In another specific embodiment, said isolated NK cells and/or ILC3 cells in said composition are from a single individual. In a more specific embodiment, said isolated NK cells and/or ILC3 cells comprise NK cells and/or ILC3 cells from at least two different individuals. In another specific embodiment, said isolated NK cells and/or ILC3 cells in said composition are from a different individual than the individual for whom treatment with the NK cells and/or ILC3 cells is intended. In another specific embodiment, said NK cells have been contacted or brought into proximity with an immunomodulatory compound or thalidomide in an amount and for a time sufficient for said NK cells to express detectably more granzyme B or perforin than an equivalent number of natural killer cells, i.e. NK cells not contacted or brought into proximity with said immunomodulatory compound or thalidomide. In another specific embodiment, said composition additionally comprises an immunomodulatory compound or thalidomide. In certain embodiments, the immunomodulatory compound is a compound described below. See, e.g., U.S. Pat. No. 7,498,171, the disclosure of which is hereby incorporated by reference in its entirety. In certain embodiments, the immunomodulatory compound is an amino-substituted isoindoline. In one embodiment, the immunomodulatory compound is 3-(4-amino-1-oxo-1,3-dihydroisoindol-2-yl)-piperidine-2,6-dione; 3-(4'-aminoisoindoline-1'-one)-1-piperidine-2,6-dione; 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione; or 4-Amino-2-(2,6-dioxopiperidin-3-yl)isoindole-1,3-dione. In another embodiment, the immunomodulatory compound is pomalidomide, or

lenalidomide. In another embodiment, said immunomodulatory compound is a compound having the structure

##STR00060##

wherein one of X and Y is C=O, the other of X and Y is C=O or CH₂, and R² is hydrogen or lower alkyl, or a pharmaceutically acceptable salt, hydrate, solvate, clathrate, enantiomer, diastereomer, racemate, or mixture of stereoisomers thereof. In another embodiment, said immunomodulatory compound is a compound having the structure

##STR00061## [0407] wherein one of X and Y is C=O and the other is CH₂ or C=O; [0408] R¹ is H, (C₁-C₈)alkyl, (C₃-C₇)cycloalkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, benzyl, aryl, (C₀-C₄)alkyl-(C₁-C₆)heterocycloalkyl, (C₀-C₄)alkyl-(C₂-C₅)heteroaryl, C(O)R³, C(S)R³, C(O)OR⁴, (C₁-C₈)alkyl-N(R⁶)₂, (C₁-C₈)alkyl-OR⁵, (C₁-C₈)alkyl-C(O)OR⁵, C(O)NHR³, C(S)NHR³, C(O)NR³R^{3'}, C(S)NR³R^{3'} or (C₁-C₈)alkyl-O(CO)R⁵; [0409] R² is H, F, benzyl, (C₁-C₈)alkyl, (C₂-C₈)alkenyl, or (C₂-C₈)alkynyl; [0410] R³ and R^{3'} are independently (C₁-C₈)alkyl, (C₃-C₇)cycloalkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, benzyl, aryl, (C₀-C₄)alkyl-(C₁-C₆)heterocycloalkyl, (C₀-C₄)alkyl-(C₂-C₅)heteroaryl, (C₀-C₈)alkyl-N(R⁶)₂, (C₁-C₈)alkyl-OR⁵, (C₁-C₈)alkyl-C(O)OR⁵, (C₁-C₈)alkyl-O(CO)R⁵, or C(O)OR⁵; [0411] R⁴ is (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, (C₁-C₄)alkyl-OR⁵, benzyl, aryl, (C₀-C₄)alkyl-(C₁-C₆)heterocycloalkyl, or (C₀-C₄)alkyl-(C₂-C₅)heteroaryl; [0412] R⁵ is (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, benzyl, aryl, or (C₂-C₅)heteroaryl; [0413] each occurrence of R⁶ is independently H, (C₁-C₈)alkyl, (C₂-C₅)alkenyl, (C₂-C₈)alkynyl, benzyl, aryl, (C₂-C₅)heteroaryl, or (C₀-C₈)alkyl-C(O)O—R⁵ or the R⁶ groups can join to form a heterocycloalkyl group; [0414] n is 0 or 1; and [0415] * represents a chiral-carbon center; [0416] or a pharmaceutically acceptable salt, hydrate, solvate, clathrate, enantiomer, diastereomer, racemate, or mixture of stereoisomers thereof. In another embodiment, said immunomodulatory compound is a compound having the structure

##STR00062## [0417] wherein: [0418] one of X and Y is C=O and the other is CH₂ or C=O; [0419] R is H or CH₂OCOR'; [0420] (i) each of R¹, R², R³, or R⁴, independently of the others, is halo, alkyl of 1 to 4 carbon atoms, or alkoxy of 1 to 4 carbon atoms or (ii) one of R¹, R², R³, or R⁴ is nitro or —NHR⁵ and the remaining of R¹, R², R³, or R⁴ are hydrogen; [0421] R⁵ is hydrogen or alkyl of 1 to 8 carbons [0422] R⁶ hydrogen, alkyl of 1 to 8 carbon atoms, benzo, chloro, or fluoro; [0423] R' is R⁷—CHR¹⁰—N(R⁸R⁹); [0424] R⁷ is m-phenylene or p-phenylene or —(C_nH_{2n})— in which n has a value of 0 to 4; [0425] each of R⁸ and R⁹ taken independently of the other is hydrogen or alkyl of 1 to 8 carbon atoms, or R⁸ and R⁹ taken together are tetramethylene, pentamethylene, hexamethylene, or —CH₂CH₂XCH₂CH₂— in which X₁ is —O—, —S—, or —NH—; [0426] R¹⁰ is hydrogen, alkyl of to 8 carbon atoms, or phenyl; and [0427] * represents a chiral-carbon center;

or a pharmaceutically acceptable salt, hydrate, solvate, clathrate, enantiomer, diastereomer, racemate, or mixture of stereoisomers thereof.

[0428] In another specific embodiment, the composition additionally comprises one or more anticancer compounds, e.g., one or more of the anticancer compounds described below.

[0429] In a more specific embodiment, the composition comprises NK cells and/or ILC3 cells from another source, or made by another method. In a specific embodiment, said other source is

placental blood and/or umbilical cord blood. In another specific embodiment, said other source is peripheral blood. In more specific embodiments, the NK cell and/or ILC3 cell population in said composition is combined with NK cells and/or ILC3 cells from another source, or made by another method in a ratio of about 100:1, 95:5, 90:10, 85:15, 80:20, 75:25, 70:30, 65:35, 60:40, 55:45: 50:50, 45:55, 40:60, 35:65, 30:70, 25:75, 20:80, 15:85, 10:90, 5:95, 100:1, 95:1, 90:1, 85:1, 80:1, 75:1, 70:1, 65:1, 60:1, 55:1, 50:1, 45:1, 40:1, 35:1, 30:1, 25:1, 20:1, 15:1, 10:1, 5:1, 1:1, 1:5, 1:10, 1:15, 1:20, 1:25, 1:30, 1:35, 1:40, 1:45, 1:50, 1:55, 1:60, 1:65, 1:70, 1:75, 1:80, 1:85, 1:90, 1:95, 1:100, or the like.

[0430] In another specific embodiment, the composition comprises an NK cell and/or ILC3 cell population produced using the three-stage method described herein and either isolated placental perfusate or isolated placental perfusate cells. In a more specific embodiment, said placental perfusate is from the same individual as said NK cell and/or ILC3 cell population. In another more specific embodiment, said placental perfusate comprises placental perfusate from a different individual than said NK cell and/or ILC3 cell population. In another specific embodiment, all, or substantially all (e.g., greater than 90%, 95%, 98% or 99%) of cells in said placental perfusate are fetal cells. In another specific embodiment, the placental perfusate or placental perfusate cells, comprise fetal and maternal cells. In a more specific embodiment, the fetal cells in said placental perfusate comprise less than about 90%, 80%, 70%, 60% or 50% of the cells in said perfusate. In another specific embodiment, said perfusate is obtained by passage of a 0.9% NaCl solution through the placental vasculature. In another specific embodiment, said perfusate comprises a culture medium. In another specific embodiment, said perfusate has been treated to remove erythrocytes. In another specific embodiment, said composition comprises an immunomodulatory compound, e.g., an immunomodulatory compound described below, e.g., an amino-substituted isoindoline compound. In another specific embodiment, the composition additionally comprises one or more anticancer compounds, e.g., one or more of the anticancer compounds described below.

[0431] In another specific embodiment, the composition comprises an NK cell and/or ILC3 cell population and placental perfusate cells. In a more specific embodiment, said placental perfusate cells are from the same individual as said NK cell and/or ILC3 cell population. In another more specific embodiment, said placental perfusate cells are from a different individual than said NK cell and/or ILC3 cell population. In another specific embodiment, the composition comprises isolated placental perfusate and isolated placental perfusate cells, wherein said isolated perfusate and said isolated placental perfusate cells are from different individuals. In another more specific embodiment of any of the above embodiments comprising placental perfusate, said placental perfusate comprises placental perfusate from at least two individuals. In another more specific embodiment of any of the above embodiments comprising placental perfusate cells, said isolated placental perfusate cells are from at least two individuals. In another specific embodiment, said composition comprises an immunomodulatory compound. In another specific embodiment, the composition additionally comprises one or more anticancer compounds, e.g., one or more of the anticancer compounds described below.

6. KITS

[0432] Provided herein is a pharmaceutical pack or kit comprising one or more containers filled with one or more of the compositions described herein, e.g., a composition comprising NK cells and/or ILC3 cells produced by a method described herein, e.g., NK cell and/or ILC3 cell populations produced using the three-stage method described herein. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

[0433] The kits encompassed herein can be used in accordance with the methods described herein, e.g., methods of suppressing the growth of tumor cells and/or methods of treating cancer, e.g.,

hematologic cancer, and/or methods of treating viral infection. In one embodiment, a kit comprises NK cells and/or ILC3 cells produced by a method described herein or a composition thereof, in one or more containers. In a specific embodiment, provided herein is a kit comprising an NK cell and/or ILC3 cell population produced by a three-stage method described herein, or a composition thereof.

7. EXAMPLES

7.1. Example 1: Three-Stage Method of Producing Natural Killer Cells from Hematopoietic Stem or Progenitor Cells

[0434] CD34^{sup}.+ cells are cultured in the following medium formulations for the indicated number of days, and aliquots of cells are taken for assessment of cell count, cell viability, characterization of natural killer cell differentiation and functional evaluation.

[0435] Stage 1 medium: 90% Stem Cell Growth Medium (SCGM) (CellGro®), 10% Human Serum-AB, supplemented with 25 ng/mL or 250 ng/mL recombinant human thrombopoietin (TPO), 25 ng/mL recombinant human Flt3L, 27 ng/mL recombinant human stem cell factor (SCF), 25 ng/mL recombinant human IL-7, 0.05 ng/mL or 0.025 ng/mL recombinant human IL-6, 0.25 ng/mL or 0.125 ng/mL recombinant human granulocyte colony-stimulating factor (G-CSF), 0.01 ng/mL or 0.025 ng/mL recombinant human granulocyte-macrophage colony-stimulating factor (GM-CSF), 0.10% gentamicin, and 1 to 10 μ m StemRegenin-1 (SR-1) or other stem cell mobilizing agent.

[0436] Stage 2 medium: 90% SCGM, 10% Human Serum-AB, supplemented with 25 ng/mL recombinant human Flt3L, 27 ng/mL recombinant human SCF, 25 ng/mL recombinant human IL-7, 20 ng/mL recombinant human IL-15, 0.05 ng/mL or 0.025 ng/mL recombinant human IL-6, 0.25 ng/mL or 0.125 ng/mL recombinant human granulocyte colony-stimulating factor (G-CSF), 0.01 ng/mL or 0.025 ng/mL recombinant human granulocyte-macrophage colony-stimulating factor (GM-CSF), 0.10% gentamicin, and 1 to 10 μ m SR1 or other stem cell mobilizing agent.

[0437] Stage 3 medium: 90% STEMMACTTM, 10% Human Serum-AB, 0.025 mM 2-mercaptoethanol (55 mM), supplemented with 22 ng/mL recombinant human SCF, 1000 U/mL recombinant human IL-2, 20 ng/mL recombinant human IL-7, 20 ng/mL recombinant human IL-15, 0.05 ng/mL or 0.025 ng/mL recombinant human IL-6, 0.25 ng/mL or 0.125 ng/mL recombinant human granulocyte colony-stimulating factor (G-CSF), 0.01 ng/mL or 0.025 ng/mL recombinant human granulocyte-macrophage colony-stimulating factor (GM-CSF), and 0.10% gentamicin.

[0438] Cells are seeded at Day 0 at 3×10^4 cells/mL in Stage 1 media, and cells are tested for purity by a CD34⁺ and CD45⁺ count and viability by 7AAD staining. At Day 5 cells are counted and seeded to a concentration of 1×10^5 cells/mL with Stage 1 medium. At Day 7 cells are counted and seeded to a concentration of 1×10^5 cells/mL with Stage 1 medium.

[0439] At Day 10, cells are counted and seeded to a concentration of 1×10^5 cells/mL in Stage 2 medium. At Day 12, cells are counted and seeded to a concentration of 3×10^5 cells/mL in Stage 2 medium. At Day 14, cells are counted and seeded in Stage 3 medium. Cells are maintained in Stage 3 media until day 35.

[0440] Alternatively, the following protocol is used through Day 14: Cells seeded at Day 0 at 7.5×10^5 cells/mL in Stage 1 media, and cells are tested for purity by a CD34⁺ and CD45⁺ count and viability by 7AAD staining. At Day 7 cells are counted and seeded to a concentration of 3×10^5 cells/mL with Stage 1 medium. At Day 9 cells are counted and seeded to a concentration of 3×10^5 cells/mL with Stage 2 medium. At Day 12, cells are counted and seeded to a concentration of 3×10^5 cells/mL in Stage 2 medium. At Day 14, cells are counted and seeded to a concentration of 3×10^5 cells/mL in Stage 2 medium.

[0441] Seeding of cells into at passage is performed either by dilution of the culture with fresh media or by centrifugation of cells and resuspension/addition of fresh media.

[0442] For harvest, cells are spun at 400 $\times$ g for seven minutes, followed by suspension of the pellet in an equal volume of Plasmalyte A. The suspension is spun at 400 $\times$ g for seven minutes, and the

resulting pellet is suspended in 10% HSA (w/v), 60% Plasmalyte A (v/v) at the target cell concentration. The cells are then strained through a 70 µm mesh, the final container is filled, an aliquot of the cells are tested for viability, cytotoxicity, purity, and cell count, and the remainder is packaged.

7.2. Example 2: Selection of Stem Cell Mobilizing Agents for the Expansion of NK Cells

[0443] The following compounds were investigated for their ability to promote the expansion of NK cell populations in vitro:

[0444] 4-(2-((2-(benzo[b]thiophen-3-yl)-6-(isopropylamino)pyrimidin-4-yl)amino)ethyl)phenol (“CRL1”)

##STR00063## [0445] 4-(2-((2-(benzo[b]thiophen-3-yl)-7-isopropylthieno[3,2-d]pyrimidin-4-yl)amino)ethyl)phenol)) (“CRL2”)

##STR00064## [0446] 4-(2-((2-(benzo[b]thiophen-3-yl)-7-isopropyl-6,7-dihydro-5H-pyrrolo[2,3-d]pyrimidin-4-yl)amino)ethyl)phenol (“CRL3”)

##STR00065## [0447] 2-(benzo[b]thiophen-3-yl)-4-((4-hydroxyphenethyl)amino)-7-isopropyl-5,7-dihydro-6H-pyrrolo[2,3-d]pyrimidin-6-one (“CRL4”)

##STR00066## [0448] 3-((2-(benzo[b]thiophen-3-yl)-9-isopropyl-9H-purin-6-yl)oxy)propanamide (“CRL5”)

##STR00067## [0449] 4-(2-((2-(benzo[b]thiophen-3-yl)-8-(dimethylamino)pyrimido[5,4-d]pyrimidin-4-yl)amino)ethyl)phenol (“CRL6”)

##STR00068## [0450] 5-(2-((2-(1H-indol-3-yl)ethyl)amino)-6-(sec-butylamino)pyrimidin-4-yl)nicotinonitrile (“CRL7”)

##STR00069## [0451] N-(2-(1H-indol-3-yl)ethyl)-2-methyl-6-phenylthieno[2,3-d]pyrimidin-4-amine (“CRL8”)

##STR00070## [0452] N-(2-(1H-indol-3-yl)ethyl)-6-(4-fluorophenyl)thieno[2,3-d]pyrimidin-4-amine (“CRL9”)

##STR00071## [0453] 3-(2-(benzo[b]thiophen-3-yl)-9-isopropyl-6-oxo-6,9-dihydro-1H-purin-1-yl)propanamide (“CRL10”)

##STR00072## [0454] N-(2-(1H-indol-3-yl)ethyl)-2-(5-fluoropyridin-3-yl)quinazolin-4-amine (“CRL11”)

##STR00073## [0455] 5-(4-((2-(1H-indol-3-yl)ethyl)amino)quinazolin-2-yl)nicotinonitrile (“CRL12”)

##STR00074## [0456] N.sup.4-(2-(H-indol-3-yl)ethyl)-N.sup.2-(sec-butyl)quinazoline-2,4-diamine (“CRL13”)

##STR00075## [0457] 2-(benzo[b]thiophen-3-yl)-4-((4-hydroxyphenethyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-5-carbonitrile (“CRL14”)

##STR00076## [0458] N-(2-(1H-indol-3-yl)ethyl)-6-(benzo[b]thiophen-3-yl)-3-isopropylimidazo[1,5-a]pyrazin-8-amine (“CRL15”)

##STR00077## [0459] 4-(2-((6-(benzo[b]thiophen-3-yl)-3-isopropylimidazo[1,5-a]pyrazin-8-yl)amino)ethyl)phenol (“CRL16”)

##STR00078## [0460] 5-(4-((2-(1H-indol-3-yl)ethyl)amino)-7-isopropylthieno[3,2-d]pyrimidin-2-yl)nicotinonitrile (“CRL17”)

##STR00079## [0461] N-(2-(1H-indol-3-yl)ethyl)-2-(5-fluoropyridin-3-yl)-7-isopropylthieno[3,2-d]pyrimidin-4-amine (“CRL18”)

##STR00080## [0462] N-(2-(1H-indol-3-yl)ethyl)-2-(5-fluoropyridin-3-yl)furo[3,2-d]pyrimidin-4-amine (“CRL19”)

##STR00081## [0463] N-(2-(1H-indol-3-yl)ethyl)-2-(5-methylpyridin-3-yl)furo[3,2-d]pyrimidin-4-amine (“CRL20”)

##STR00082## [0464] N-(2-(1H-indol-3-yl)ethyl)-7-isopropyl-2-(5-methylpyridin-3-yl)thieno[3,2-d]pyrimidin-4-amine (“CRL21”)

##STR00083##

and [0465] 5-(4-((2-(1H-indol-3-yl)ethyl)amino)furo[3,2-d]pyrimidin-2-yl)nicotinonitrile

("CRL22")

##STR00084##

7.3. Example 3: Characterization of Three-Stage NK Cells

Methods

[0466] UCB CD34+ cells were cultivated in presence of cytokines including thrombopoietin, SCF, Flt3 ligand, IL-7, IL-15 and IL-2 for 35 days to produce three-stage NK cells, as described in Example 1. Multi-color flow cytometry was used to determine the phenotypic characteristics of three-stage NK cells.

[0467] For biological testing, the compounds were provided to culture to evaluate their effects on NK cell expansion and differentiation. Specifically, donors of CD34+ cells (StemCell Technology) were thawed and expanded in vitro following NK culture protocol. During the first 14 days of the culture, each CRL compounds was dissolved in DMSO and added to the culture at 10 μ M concentration. SR1 (at 10 μ M) served as a positive control compound, while DMSO alone without any compound served as a negative control. At the end of the culture on Day 35, cell expansion, natural killer (NK) cell differentiation and cytotoxicity of the cells against K562 tumor cell line were characterized. Due to the large number of the compounds, the testing was performed in two experiments, CRL1-11 and CRL 12-22. The same donors were used for each experiment. Positive and negative controls were also included in both experiments.

Results

[0468] Cell expansion data showed that 20 out of the 22 compounds supported NK expansion at 10 μ M concentration. Except for CRL7 and CRL13, the rest of the compounds all resulted in a NK expansion of 2,000~15,000 fold over 35 days (FIG. 1 and FIG. 2). Among all the compounds, CRL19, 20 and 22 supported cell expansion the best, and they demonstrated a similar level of expansion compared to SR1 at Day 35 (FIG. 3). CD34 cell expansion at Day 14 of the culture showed a similar trend that most of the compounds supported CD34 cells expansion, and CRL19, 20 and 22 achieved the highest CD34 cell expansion at Day 14 (FIG. 4).

[0469] Cytotoxicity assay was run using compound cultured cells against K562 tumor cells at 10:1 effector to target ratio (FIG. 5) to evaluate cell functions. The results showed that the cells cultured with compounds killed 30~60% of K562 cells at 10:1 E:T ratio, indicating that the cells present NK functions. For both donors, cells cultured with CRL17, 18, 19 and 21 demonstrated similar or greater killing activities compared to those cultured with SR1.

CONCLUSIONS

[0470] In summary, we found that all the compounds except CRL7 and CRL13 supported PNK-007 expansion and differentiation. Expansion with the compounds ranged from 2,000~15, 000 fold over 35 days, and the culture achieved more than 70% of NK cells. Among these compounds, CRL 19, 20 and 22 demonstrated very similar expansion, differentiation and cytotoxicity profiles as SR1 for PNK-007 culture. CRL 17, 18, and 21 resulted in slightly less expansion compared to SR1 but increased CD56+/CD11a+ subpopulation, and also increased killing activities of the cells.

7.4 Example 4: Further Characterization of Three-Stage NK Cells

Methods

[0471] Cells: Frozen PBMC were acquired from Stem Cell Technologies. Peripheral blood derived NKs (PB-NK) cells were isolated from fresh blood of healthy donors using the Human NK Cell Enrichment Kit (Stem Cell Technologies) according to manufacturer's instructions. CYNK cells were generated from umbilical cord blood-derived CD34+ stem cells (Ref: Zhang et al. J Immunother Cancer. 2015). Briefly, the CD34+ cells were cultivated in the presence of cytokines including thrombopoietin, SCF, Flt3 ligand, IL-7, IL-15 and IL-2 for 35 days. PBNK and CYNK cells were cryopreserved until analysis.

[0472] Magnetic-activated cell sorting: PNK cells were stained with PE Mouse Anti-Human CD11a (BD) and CD11a+ PNK cells concentrated using anti-PE MicroBeads according to manufacturer's instructions (Miltenyi Biotec).

[0473] Single cell RNA sequencing: CYNK cells were combined with PB-NK at 1:1 ratio and gene expression analyzed on single cell level using 10× Genomics Chromium platform and Illumina sequencing. Bioinformatics analysis utilized 10× Genomics Cell Ranger analysis pipeline.

[0474] Flow Cytometry: Cryopreserved cells were rapidly thawed in a 37° C. water bath and washed once in RPMI1640+10% hiFBS (heat inactivated Fetal Bovine Serum, Gibco), followed by LIVE/DEAD™ Fixable Aqua Stain in PBS. Cells were washed with FACS buffer (PBS+2% FBS) followed by incubation in blocking solution (Brilliant Stain buffer, Mouse IgG2a isotype k control and Human BD Fc Block (all from BD)). Cells were washed with FACS buffer and incubated with fluorophore-coupled antibodies in FACS buffer for 25 min on ice. Cells were washed with FACS buffer before analysis on Fortessa X20 flow cytometer (BD).

[0475] qRT-PCR: RNA was isolated from cells using Quick-RNA Miniprep kit (Qiagen) according to the manufacturer's instructions. cDNA was synthesized using SuperScript IV Reverse Transcriptase (Thermo Fisher Scientific) in a standard reaction. RT-PCR was performed using Taqman Gene expression assays (Applied Biosystems). Expression levels were calculated relative to GAPDH (Hs02758991) using the $\Delta\Delta C_t$ method.

Results

[0476] CYNK cells efficiently kill various tumor cell lines in vitro, however, the mechanisms CYNK cells use to induce cell death remains poorly understood (ref). To elucidate on the activating NK cell receptors, the intracellular signaling pathways and molecular mechanisms CYNK cells employ to carry out their functional roles, we used single-cell RNA sequencing (scRNAseq) as an unbiased approach to compare CYNK cells to peripheral blood NK cells (PB-NK) (FIG. 6A). Unbiased transcriptional clustering revealed two distinct signatures differentiating between CYNK and PB-NK cells (FIG. 6B). Tables 1 and 2 list top 50 upregulated genes per cluster in PB-NK and CYNK cells, respectively. The gene set expressed higher in PB-NK cells included genes associated with NK cell functional roles, including FGFBP2, granzymes (GZMH, GZMM), CXCR4, KLRF1, KLF2, IFNG (Table 1).

[0477] FGFBP2, encoding fibroblast growth factor-binding protein, is known to be secreted by cytotoxic lymphocytes.

[0478] Granzymes are a group of serine proteases which are stored in the cytotoxic granules of NK cells and cytotoxic T lymphocytes (ref). While GzmA and GzmB induce target cell death upon release to their cytoplasm and have been extensively studied, less is known about the functional role of GzmH, GzmK and GzmM.

[0479] CXCR4 regulates NK cell homing to bone marrow.

[0480] KLRF1 encodes NKp80, an activating C-type lectin-like immunoreceptor that is activated upon binding to activation-induced C-type lectin (AICL), inducing NK cell cytotoxicity and cytokine secretion.

[0481] Transcription factor KLF2 that regulates both NK cell proliferation and survival.

[0482] NK cell-derived IFN-7 (IFNG gene) is a key immunoregulatory factor secreted from activated NK cells that promotes adaptive immune response by modulating dendritic cell and T cell responses.

TABLE-US-00001 TABLE 1 Top 50 upregulated genes per PB-NK cluster. Feature CYNK PB-NK PB-NK Log2 PB-NK P- Feature ID Name Average Average Fold Change Value 1

ENSG00000137441	FGFBP2	0.099352	2.935962	4.88363	4.09E-78	2	ENSG00000100450	GZMH	0.136708	2.484828	4.182845	2.49E-58	3	ENSG00000276085	CCL3L3	0.072152	1.251852	4.115143	2.13E-49	4	ENSG00000197540	GZMM	0.134235	1.982728	3.883559	1.40E-50	5	ENSG00000121966	CXCR4	0.403236	5.935725	3.879087	9.19E-51	6	ENSG00000169554	ZEB2	0.127877	1.860789	3.861967	7.03E-50	7	ENSG00000127528	KLF2	0.172475	1.92761	3.481483	1.86E-40	8	ENSG00000189067	LITAF	0.297791	3.231559	3.439184	1.06E-39	9	ENSG00000069667	RORA	0.101913	1.055542	3.371425	3.26E-37	10	ENSG00000145220	LYAR	0.142448	1.306592	3.196402	2.39E-33	11	ENSG00000125107	CNOT1	0.208595	1.809824			
-----------------	--------	----------	----------	---------	----------	---	-----------------	------	----------	----------	----------	----------	---	-----------------	--------	----------	----------	----------	----------	---	-----------------	------	----------	----------	----------	----------	---	-----------------	-------	----------	----------	----------	----------	---	-----------------	------	----------	----------	----------	----------	---	-----------------	------	----------	---------	----------	----------	---	-----------------	-------	----------	----------	----------	----------	---	-----------------	------	----------	----------	----------	----------	----	-----------------	------	----------	----------	----------	----------	----	-----------------	-------	----------	----------	--	--	--

3.116348 3.39E-32 12 ENSG00000115317 IFNG 0.193317 1.639941 3.083863 1.11E-29 13
 ENSG00000158050 DUSP2 0.40774 3.322164 3.025836 4.12E-30 14 ENSG00000110046
 ATG2A 0.190226 1.508942 2.987028 3.39E-29 15 ENSG00000173762 CD7 0.492697 3.641922
 2.885402 1.77E-27 16 ENSG00000141682 PMAIP1 0.252398 1.820017 2.849558 6.51E-26 17
 ENSG00000078304 PPP2R5C 0.381864 2.591665 2.762207 6.15E-25 18 ENSG00000153234
 NR4A2 0.399174 2.622622 2.715393 5.59E-24 19 ENSG00000152518 ZFP36L2 0.856899
 5.585388 2.703993 4.72E-24 20 ENSG00000145675 PIK3R1 0.325168 2.078618 2.675822
 2.70E-23 21 ENSG00000150045 KLRF1 0.191285 1.177103 2.620822 4.78E-22 22
 ENSG00000255198 SNHG9 0.516983 2.951818 2.512937 1.34E-20 23 ENSG00000125148
 MT2A 0.51504 2.913311 2.499426 9.06E-20 24 ENSG00000116741 RGS2 0.203737 1.147279
 2.492865 1.51E-19 25 ENSG00000153922 CHD1 0.252574 1.350762 2.418474 9.42E-19 26
 ENSG00000120129 DUSP1 2.078529 9.865317 2.24638 2.58E-16 27 ENSG00000143924 EML4
 0.256284 1.150299 2.165756 7.80E-15 28 ENSG00000128016 ZFP36 2.22866 9.777355
 2.132849 1.32E-14 29 ENSG00000163874 ZC3H12A 0.261759 1.120475 2.097382 7.47E-14 30
 ENSG00000105993 DNAJB6 0.6506 2.667169 2.035058 2.98E-13 31 ENSG00000126524 SBDS
 0.534822 2.185078 2.030148 3.57E-13 32 ENSG00000125347 IRF1 1.450448 5.812277 2.002193
 7.32E-13 33 ENSG00000157514 TSC22D3 1.103379 4.30409 1.963373 2.57E-12 34
 ENSG00000184205 TSPYL2 0.592137 2.247746 1.924086 1.14E-11 35 ENSG00000146278
 PNRC1 1.362312 5.156149 1.919832 7.77E-12 36 ENSG00000135070 ISCA1 0.27898 1.043084
 1.90227 2.06E-11 37 ENSG00000171223 JUNB 4.09462 15.11622 1.883884 2.20E-11 38
 ENSG00000156232 WHAMM 0.316425 1.146147 1.856513 7.14E-11 39 ENSG00000164327
 RICTOR 0.318279 1.101977 1.791406 3.85E-10 40 ENSG00000118503 TNFAIP3 0.550807
 1.902316 1.787777 3.93E-10 41 ENSG00000120616 EPC1 0.562199 1.846066 1.714953
 2.17E-09 42 ENSG00000167508 MVD 0.309448 1.00722 1.702322 4.11E-09 43
 ENSG00000013441 CLK1 0.690164 2.216412 1.682859 4.62E-09 44 ENSG00000188042
 ARL4C 0.437325 1.388136 1.666056 8.18E-09 45 ENSG00000162924 REL 0.553809 1.736208
 1.648145 1.14E-08 46 ENSG00000005483 KMT2E 0.79402 2.460289 1.631225 1.47E-08 47
 ENSG00000119801 YPEL5 0.966141 2.98202 1.625617 1.70E-08 48 ENSG00000123505 AMD1
 0.558578 1.664102 1.574595 6.03E-08 49 ENSG00000159388 BTG2 0.751541 2.22132 1.563151
 7.55E-08 50 ENSG00000010404 IDS 0.723193 2.128073 1.556757 8.48E-08

[0483] Top differentially expressed genes in CYNK cluster that are encode factors associated with
 NK cell functional role include surface receptors and co-receptors (CD96, NCR3, CD59, KLRC1),
 TNFSF10, immune checkpoint genes (TNFRSF18, TNFRSF4, HAVCR2), NK cell receptor
 adaptor molecule genes (FCER1G and LAT2) (Table 2).

TABLE-US-00002 TABLE 2 Top 50 upregulated genes per CYNK cluster. CYNK Log2 Feature
 PBNK CYNK Fold CYNK P- Feature ID Name Average Average Change Value 1
 ENSG00000102471 NDFIP2 0.077391 1.45981 4.230949 1.69E-22 2 ENSG00000242258
 LINC00996 0.063046 1.183921 4.222944 5.04E-22 3 ENSG00000172005 MAL 0.057005
 1.03529 4.173813 1.35E-21 4 ENSG00000108702 CCL1 0.078524 1.334494 4.080611 5.11E-09
 5 ENSG00000198125 MB 0.10193 1.683947 4.041355 1.45E-20 6 ENSG00000128040 SPINK2
 0.087962 1.233641 3.804242 7.88E-19 7 ENSG00000166920 C15orf48 0.078901 1.018246
 3.683547 6.40E-18 8 ENSG00000134072 CAMK1 0.151762 1.932724 3.667647 2.13E-18 9
 ENSG00000134545 KLRC1 0.509273 4.740451 3.217889 9.47E-16 10 ENSG00000121858
 TNFSF10 0.295975 2.682764 3.178801 6.44E-15 11 ENSG00000186891 TNFRSF18 1.182011
 10.09017 3.093605 6.96E-15 12 ENSG00000008517 IL32 4.345617 37.08234 3.093395 6.60E-15
 13 ENSG00000042493 CAPG 0.369213 3.112494 3.074529 9.91E-15 14 ENSG00000235576
 AC092580.4 0.44736 3.660475 3.031759 2.23E-14 15 ENSG00000163191 S100A11 0.41527
 3.364804 3.017543 2.42E-14 16 ENSG00000186827 TNFRSF4 0.135529 1.097816 3.01448
 1.91E-13 17 ENSG00000074800 ENO1 2.166202 16.05066 2.889567 1.86E-13 18
 ENSG00000158869 FCER1G 0.734274 5.393877 2.876632 2.43E-13 19 ENSG00000118971

CCND2 0.457175 3.324621 2.861636 3.21E-13 20 ENSG00000205426 KRT81 0.169883
1.187806 2.803005 3.69E-12 21 ENSG00000243927 MRPS6 0.358643 2.29304 2.675597
6.10E-12 22 ENSG00000182718 ANXA2 0.206125 1.282389 2.635118 3.48E-11 23
ENSG00000125384 PTGER2 0.175546 1.08713 2.628037 4.29E-11 24 ENSG00000124767
GLO1 0.214053 1.289543 2.588793 6.50E-11 25 ENSG00000135077 HAVCR2 0.175924
1.031051 2.548543 1.51E-10 26 ENSG00000103490 PYCARD 0.183097 1.070527 2.545209
1.34E-10 27 ENSG00000086730 LAT2 0.178566 1.04156 2.541707 1.53E-10 28
ENSG00000141526 SLC16A3 0.282006 1.622835 2.523282 1.73E-10 29 ENSG00000103187
COTL1 0.894342 5.013779 2.486834 1.45E-10 30 ENSG00000067225 PKM 1.099712 6.145949
2.482453 1.11E-10 31 ENSG00000177156 TALDO1 0.196687 1.084745 2.46115 4.23E-10 32
ENSG00000153283 CD96 0.368458 2.029162 2.460314 1.66E-10 33 ENSG00000204475 NCR3
0.640272 3.472457 2.438804 2.31E-10 34 ENSG00000170442 KRT86 0.257845 1.372733
2.410873 1.02E-09 35 ENSG00000117632 STMN1 0.468878 2.413499 2.36315 1.22E-09 36
ENSG00000227507 LTB 3.831437 19.41653 2.341609 1.09E-09 37 ENSG00000130429 ARPC1B
0.570053 2.846585 2.31957 1.27E-09 38 ENSG00000162704 ARPC5 0.347317 1.717418 2.30484
1.66E-09 39 ENSG00000088832 FKBP1A 0.40017 1.978205 2.304629 1.60E-09 40
ENSG00000102265 TIMP1 0.385447 1.902345 2.302248 1.96E-09 41 ENSG00000113088
GZMK 0.290312 1.403201 2.27168 1.37E-08 42 ENSG00000085063 CD59 0.215186 1.035997
2.265377 7.12E-09 43 ENSG00000102144 PGK1 1.405879 6.735348 2.260328 2.92E-09 44
ENSG00000148908 RGS10 0.217451 1.014713 2.220352 1.33E-08 45 ENSG00000196405 EVL
1.186164 5.50471 2.214345 5.41E-09 46 ENSG00000128340 RAC2 1.063092 4.917253 2.209516
5.72E-09 47 ENSG00000100097 LGALS1 4.427539 20.46621 2.208968 6.05E-09 48
ENSG00000139626 ITGB7 0.50059 2.285445 2.19016 8.54E-09 49 ENSG00000196230 TUBB
1.062715 4.838214 2.186651 1.22E-08 50 ENSG00000171314 PGAM1 0.670096 3.046436
2.18433 8.56E-09

[0484] To better understand how the cytotoxic response is initiated in CYNK cells, we specifically analyzed the expression of manually chosen genes encoding well characterized proteins leading from target detection to a cytolytic response, with main focus on NK cell receptors and adaptor molecule (Table 3). Differential gene expression analysis showed high expression of the two key cytotoxic molecules perforin (PRF1) and granzyme B (GZMB) in CYNK cells. Similarly, most receptors that were differentially expressed between CYNK and PB-NK cells, with the exception of KLRF1 (encoding NKp80), were higher expressed on CYNK cells. Expression of selected NK cell effector and receptor genes is visualized on tSNE plots in FIG. 6C. Elevated expression of genes encoding components of the NK cell cytotoxic machinery correlate well with the high cytotoxic activity of CYNK cells against a broad range of target cells.

TABLE-US-00003 TABLE 3 Top differentially expressed genes encoding factors regulating NK cell cytolytic function. Genes that had <1 count per cell across the entire cluster were excluded.

Log2 Feature Value	CYNK	PBNK	Fold	CYNK	P- Feature ID	Name	Alias	Average	Average Change
1	ENSG00000134545	KLRC1	NKG2A	4.740451	0.509273	3.217889	9.47E-16	CD159a	2
	ENSG00000121858	TNFSF10	TRAIL	2.682764	0.295975	3.178801	6.44E-15	3	
	ENSG00000186891	TNFRSF18	GITR	10.09017	1.182011	3.093605	6.96E-15	4	
	ENSG00000186827	TNFRSF4	CD134	1.097816	0.135529	3.014481	1.91E-13	OX40	5
	ENSG00000135077	HAVCR2	TIM-3	1.031051	0.175924	2.548543	1.51E-10	6	
	ENSG00000153283	CD96	Tactile	2.029162	0.368458	2.460314	1.66E-10	7	ENSG00000204475
	NCR3	CD337	3.472457	0.640272	2.438804	2.31E-10	NKp30	8	MAC-IP, ENSG00000085063
	CD59	MIRL	1.035997	0.215186	2.265377	7.12E-09	protectin	9	ENSG00000139626
	ITGB7	2.285445	0.50059	2.19016	8.54E-09	10	ENSG00000180644	PRF1	3.589295
	0.887169	2.016259	8.95E-08	11	ENSG00000100453	GZMB	11.6194	3.515453	1.725026
	4.27E-06	12	ENSG00000100385	IL2RB	2.568753	0.956632	1.424929	0.000126	13
	ENSG00000205809	KLRC2	NKG2C	1.419451	0.784861	0.854636	0.026587	CD159c	14
	ENSG00000111796	KLRB1							

CD161 18.74844 10.45953 0.842324 0.027995 15 15NSG00000150045 KLRF1 NKp80 0.191285
1.177103 -2.62082 4.78E-22

[0485] We next analyzed the transcriptional profile of CYNK and PB-NK cells by quantitative real-time PCR (qRT-PCR) focusing on selected NK cell-associated genes that were highly and/or differentially expressed in the scRNAseq dataset (FIG. 7). RNA was extracted from freshly thawed naïve cells post isolation or culture. qRT-PCR demonstrated high expression of CD69, KLRK1 and KLRB1 relative to the housekeeping gene GAPDH in both CYNK and PB-NK cells, whereas, KLRK1 and KLRB1, encoding for NKG2D and CD161/KLRB1, respectively, were significantly higher expressed in PB-NK cells. Significant differential expression of NKp80, encoded by KLRF1 gene, earlier seen by scRNAseq (Table 3), was confirmed by qRT-PCR. Similarly, KLRD1 was higher expressed on PB-NK compared to CYNK cells. Together, the data show higher expression of the inhibitory killer cell lectin-like receptor (KLRB1, KLRD1, KLRF1) expression on PB-NK cells when compared to CYNK cells. The two C-type lectin receptor genes KLRC1 and KLRC2, encoding the inhibitory NKG2A and the activating NKG2C, were higher expressed in CYNK cells. Of the natural cytotoxicity receptors (NCRs), only NCR2 (encoding NKp44) was differentially expressed with high expression in CYNK cells and almost no expression in PB-NK cells. Two co-activating NK cell receptor genes CD244 (2B4) and CD226 (DNAM-1) were slightly higher expressed in PB-NK compared to CYNK cells. Alongside the typical ligand-activated NK cell receptor genes, we also analyzed the expression of FCGR3A encoding an Fc receptor CD16 that is required for antibody-dependent cell-mediated cytotoxicity. Whereas scRNAseq data demonstrated no significant differential expression of FCGR3A, by qRT-PCR it was highly expressed in the PB-NK cells and at a very low level in CYNK cells. The expression of two genes TNFRSF18 and TNFSF10 that were highly differentially expressed by scRNAseq and elevated in the CYNK cluster, were also analyzed by qRT-PCR. The PCR data confirms high expression of these genes encoding for GITR and TRAIL, respectively, on CYNK cells relative to low level expression in PB-NK cells.

[0486] Lastly, we characterized CYNK cells relative to PB-NK by surface protein expression using flow cytometry. Antibodies targeting various NK cell receptors were chosen based on the transcriptional characterization by scRNAseq and qRT-PCR (Tables 1-3, GIG. 6 and FIG. 7). NK cells express high level of the NK cell marker CD56 and lack the expression of T cell, B cell and myeloid cell markers CD3, CD19 and CD14, respectively (FIG. 8). Whereas a majority of PB-NK cells express CD56 at a low level, a small subset of PB-NK cells express CD56 at a level seen in CYNK cells (FIG. 9). NCR analysis demonstrated a high expression of NKp44 in CYNK cells, whereas, NKp44 was expressed at a low level in PB-NK, corresponding well to our transcriptional analysis (FIG. 7). NKp80, on the other hand, was expressed on PB-NK cell and little on CYNK, also confirming the transcriptional data of KLRF1 expression (Table 1 and FIG. 7). CD16 was virtually not expressed on CYNK cells, whereas the majority of PB-NK cells expressed CD16 at a high level. CD16 protein expression, therefore, also corresponds well to transcriptional analysis (Table 1 and FIG. 7). The expression of killer cell lectin-like receptors was comparable between CYNK and PB-NK cells, with CYNK cells demonstrating higher mean fluorescence intensity compared to PB-NK cells for NKG2D, NKG2C, CD94 (NKG2C) and NKG2A. GITR, a checkpoint inhibitor molecule, encoded by TNFRSF18, was not expressed on PB-NK cells but highly on all CYNK cells, correlating well to qRT-PCR data.

[0487] We used the flow cytometry dataset (FIG. 8 and FIG. 9) to perform an unbiased analysis of the surface marker expression on CYNK and PB-NK cell populations (FIG. 10). Antibody-stained CYNK and PBMC cells were mixed for acquisition and analyzed by flow cytometry. It is evident from the tSNE plots that CYNK and PB-NK cells cluster separately from each other and other peripheral blood cells when looking at the localization of CD56- and CD3/CD14/CD19-positive cells on the plot. High expression of NKp44 (CD336) and GITR (CD357) enable the identification of CYNK cells as GITR is virtually not expressed in any cell type in the PBMC subsets. PB-NK

cells on the other hand, highly express CD16 and NKp80 that are not expressed on CYNK cells. Altogether, we have identified cell surface markers that allow to distinguish CYNK cells from PB-NK with high confidence.

7.5 Example 5: Senescent Cell Killing

[0488] Celularity has established a procedure to generate clinical-grade placental CD34+ stem cell-derived NK (CYNK) cells. CYNK-001 cells are currently being evaluated in the clinic as an anti-cancer therapeutic. CYNK-001 cells highly express (mRNA and protein level) an array of typical NK cell activating receptors, including NKp30, NKp44, NKp46 and NKG2D. Celularity has also demonstrated the ability to knockout NKG2A during our CYNK manufacturing process, resulting in an NK cell which is not sensitive to HLA-E dependent inhibition as demonstrated by cytotoxicity experiments using U251 glioblastoma cell line. Based on these mechanistic studies, we anticipate similar sensitivity of senescent cells to CYNK cell targeted cytotoxicity which can additionally be enhanced by NKG2A gene knockout.

[0489] Here, we demonstrate that CYNK-001 express functional NKG2D, an NK cell receptor implicated in clearing senescent cells from the body (ref: PMID: 26878797, PMID: 31160572).

[0490] Results: NK cells recognize transformed, stressed or infected cells via specific receptor-ligand interactions. To explore the ability of CYNK-001 to recognize target cells, we first analyzed the cell surface expression of NK cell activating receptors that are known to mediate the anti-tumor activity of NK cells. CYNK-001 cells present with the nominal NK cell phenotype: a lack of expression of lineage markers CD3, CD14 and CD19, but high expression of CD56 (FIG. 11A) and NK cell activating receptors NKp30, NKp44, NKp46, NKG2D and DNAM-1 (FIGS. 11B and 11C).

[0491] Ligands for the NK cell activating receptor NKG2D are upregulated on senescent cells (ref: PMID: 34359910). To analyze whether NKG2D receptor on CYNK-001 cells is functional, we screened different tumor cell lines for the expression of NKG2D receptor ligands. First, we analyzed the binding of recombinant NKG2D protein to its ligands on tumor cell lines.

Recombinant Fc-coupled NKG2D bound to tumor cell lines at different efficiency, suggesting different level of expression of NKG2D ligands on tumor cells (FIG. 12). We also used antibodies specific for major known NKG2D ligands MHC class I chain-related protein A and B (MICA/B) and members of the UL16 binding protein (ULBP) family. ULBP2/5/6 were highly expressed on most analyzed tumor cell lines. MICA/B was only detected on K562 cells. Together, the data indicates, that tumor cell lines express different NKG2D ligands and could be recognized by CYNK-001 cells.

[0492] To specifically analyze the involvement of NKG2D on CYNK-001 in target cell recognition, we performed a cytotoxicity assay on NKG2D ligand-expressing cell lines. Co-culture of CYNK-001 cells with tumor cell lines induced rapid tumor cell death (FIG. 13). Addition of an NKG2D blocking antibody in the reaction reduced cytolysis against all analyzed tumor cell lines compared to a non-specific isotype control antibody.

[0493] Mean killing was reduced from 50.91% (IgG) to 28.91% with NKG2D blocking antibody on LN229 cell line. Mean killing was reduced from 50.55% (IgG) to 35.05% with NKG2D blocking antibody on U251 cell line. Mean killing was reduced from 57.40% (IgG) to 41.57% with NKG2D blocking antibody on DLD-1 cell line. Mean killing was reduced from 54.11% (IgG) to 40.88% with NKG2D blocking antibody on HCT116 cell line. Mean killing was reduced from 51.73% (IgG) to 29.33% with NKG2D blocking antibody on HT29 cell line. The data indicates that CYNK-001 uses NKG2D receptor for the recognition and cytotoxic elimination of target cells.

EQUIVALENTS

[0494] The present invention is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the invention in addition to those described will become apparent to those skilled in the art from the foregoing description and accompanying figures. Such modifications are intended to fall within the scope of the appended claims.

[0495] All references cited herein are incorporated herein by reference in their entirety and for all purposes to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference in its entirety for all purposes. The citation of any publication is for its disclosure prior to the filing date and should not be construed as an admission that the present invention is not entitled to antedate such publication by virtue of prior invention.

Claims

1. A method of killing senescent cells comprising contacting the senescent cells with placenta-derived natural killer (NK) cells.
2. A method of treating a disease or disorder associated with cellular senescence in a subject in need thereof comprising administering to the subject an effective amount of placenta-derived NK cells to the subject so as thereby to provide an effective treatment of the disease or disorder in the subject.
3. The method of claim 1 or claim 2, wherein the placenta-derived NK cells comprise at least 20% CD56+CD3⁻ natural killer cells.
4. The method of claim 1 or claim 2, wherein the placenta-derived NK cells comprise at least 40% CD56+CD3⁻ natural killer cells.
5. The method of claim 1 or claim 2, wherein the placenta-derived NK cells comprise at least 60% CD56+CD3⁻ natural killer cells.
6. The method of claim 1 or claim 2, wherein the placenta-derived NK cells comprise at least 80% CD56+CD3⁻ natural killer cells.
7. The method of any of claims 1-6, wherein said placenta derived natural killer cells are human placenta derived natural killer cells.
8. The method of any of claims 1-6, wherein said placenta derived natural killer cells are hematopoietic stem cell-derived natural killer cells.
9. The method of any of claims 1-6, wherein said placenta derived natural killer cells are CD34⁺ hematopoietic stem cell-derived natural killer cells.
10. The method of any of claims 1-6, wherein said placenta derived natural killer cells are CYNK cells.
11. The method of any one of claims 1-10, wherein the NK cells are characterized by expression of one or more markers selected from the group consisting of FGFBP2, GZMH, CCL3L3, GZMM, CXCR4, ZEB2, KLF2, LITAF, RORA, LYAR, CNOT1, IFNG, DUSP2, ATG2A, CD7, PMAIP1, PPP2R5C, NR4A2, ZFP36L2, PIK3R1, KLRF1, SNHG9, MT2A, RGS2, CHD1, DUSP1, EML4, ZFP36, ZC3H12A, DNAJB6, SBDS, IRF1, TSC22D3, TSPYL2, PNRC1, ISCA1, JUNB, WHAMM, RICTOR, TNFAIP3, EPC1, MVD, CLK1, ARL4C, REL, KMT2E, YPEL5, AMD1, BTG2, and IDS which is lower than expression of said markers in peripheral blood natural killer cells and/or expression of one or more markers selected from the group consisting of NDFIP2, LINC00996, MAL, CCL1, MB, SPINK2, C15orf48, CAMK1, KLRC1, TNFSF10, TNFRSF18, IL32, CAPG, AC092580.4, S100A11, TNFRSF4, ENO1, FCER1G, CCND2, KRT81, MRPS6, ANXA2, PTGER2, GLO1, HAVCR2, PYCARD, LAT2, SLC16A3, COTL1, PKM, TALDO1, CD96, NCR3, KRT86, STMN1, LTB, ARPC1B, ARPC5, FKBP1A, TIMP1, GZMK, CD59, PGK1, RGS10, EVL, RAC2, LGALS1, ITGB7, TUBB, PGAM1, PRF1, GZMB, IL2RB, KLRC2, and KLRB1 which is higher than expression of said markers in peripheral blood natural killer cells.
12. The method of any one of claims 1-11, wherein the NK cells are characterized by expression of one or more markers selected from the group consisting of FGFBP2, GZMH, CCL3L3, GZMM, CXCR4, ZEB2, KLF2, LITAF, RORA, LYAR, CNOT1, IFNG, DUSP2, ATG2A, CD7, PMAIP1, PPP2R5C, NR4A2, ZFP36L2, PIK3R1, KLRF1, SNHG9, MT2A, RGS2, CHD1, DUSP1, EML4, ZFP36, ZC3H12A, DNAJB6, SBDS, IRF1, TSC22D3, TSPYL2, PNRC1, ISCA1, JUNB,

WHAMM, RICTOR, TNFAIP3, EPC1, MVD, CLK1, ARL4C, REL, KMT2E, YPEL5, AMD1, BTG2, and IDS which is lower than expression of said markers in peripheral blood natural killer cells.

13. The method of claim 12, wherein expression of 2, 3, 4, 5, 6, 7, 8, 9, 10, or more markers selected from the group consisting of FGFBP2, GZMH, CCL3L3, GZMM, CXCR4, ZEB2, KLF2, LITAF, RORA, LYAR, CNOT1, IFNG, DUSP2, ATG2A, CD7, PMAIP1, PPP2R5C, NR4A2, ZFP36L2, PIK3R1, KLRF1, SNHG9, MT2A, RGS2, CHD1, DUSP1, EML4, ZFP36, ZC3H12A, DNAJB6, SBDS, IRF1, TSC22D3, TSPYL2, PNRC1, ISCA1, JUNB, WHAMM, RICTOR, TNFAIP3, EPC1, MVD, CLK1, ARL4C, REL, KMT2E, YPEL5, AMD1, BTG2, and IDS is lower than expression of said markers in peripheral blood natural killer cells.

14. The method of any one of claims 1-13, wherein the NK cells are characterized by expression of one or more markers selected from the group consisting of NDFIP2, LINC00996, MAL, CCL1, MB, SPINK2, C15orf48, CAMK1, KLRC1, TNFSF10, TNFRSF18, IL32, CAPG, AC092580.4, S100A11, TNFRSF4, ENO1, FCER1G, CCND2, KRT81, MRPS6, ANXA2, PTGER2, GLO1, HAVCR2, PYCARD, LAT2, SLC16A3, COTL1, PKM, TALDO1, CD96, NCR3, KRT86, STMN1, LTB, ARPC1B, ARPC5, FKBP1A, TIMP1, GZMK, CD59, PGK1, RGS10, EVL, RAC2, LGALS1, ITGB7, TUBB, PGAM1, PRF1, GZMB, IL2RB, KLRC2, and KLRB1 which is higher than expression of said markers in peripheral blood natural killer cells.

15. The method of claim 14, wherein expression of 2, 3, 4, 5, 6, 7, 8, 9, 10, or more markers selected from the group consisting of NDFIP2, LINC00996, MAL, CCL1, MB, SPINK2, C15orf48, CAMK1, KLRC1, TNFSF10, TNFRSF18, IL32, CAPG, AC092580.4, S100A11, TNFRSF4, ENO1, FCER1G, CCND2, KRT81, MRPS6, ANXA2, PTGER2, GLO1, HAVCR2, PYCARD, LAT2, SLC16A3, COTL1, PKM, TALDO1, CD96, NCR3, KRT86, STMN1, LTB, ARPC1B, ARPC5, FKBP1A, TIMP1, GZMK, CD59, PGK1, RGS10, EVL, RAC2, LGALS1, ITGB7, TUBB, PGAM1, PRF1, GZMB, IL2RB, KLRC2, and KLRB1 is higher than expression of said markers in peripheral blood natural killer cells.

16. The method of any one of claims 1-15, wherein the NK cells are prepared by the methods presented herein.

17. The method of any one of claims 2-16, wherein the disease or disorder is a fibrotic disease or disorder.

18. The method of any one of claims 2-16, wherein the disease or disorder is a inflammatory disease or disorder.

19. The method of any one of claims 2-16, wherein the disease or disorder is a metabolic disease or disorder.

20. The method of any one of claims 2-19, wherein the disease or disorder is a liver disease or disorder.

21. The method of any one of claims 2-19, wherein the disease or disorder is a kidney disease or disorder.

22. The method of any one of claims 2-19, wherein the disease or disorder is a lung disease or disorder.

23. The method of any one of claims 2-19, wherein the disease or disorder is a eye disease or disorder.

24. The method of any one of claims 2-19, wherein the disease or disorder is a skeletal disease or disorder.

25. The method of any one of claims 2-19, wherein the disease or disorder is a skin disease or disorder.

26. The method of any one of claims 2-19, wherein the disease or disorder is a gastrointestinal disease or disorder.

27. The method of any one of claims 2-19, wherein the disease or disorder is a bone marrow disease or disorder.

28. The method of any one of claims 2-19, wherein the disease or disorder is a cardiovascular disease or disorder.
 29. The method of any one of claims 2-19, wherein the disease or disorder is a circulatory disease or disorder.
 30. The method of any one of claims 2-19, wherein the disease or disorder is a neuronal disease or disorder.
 31. The method of any one of claims 2-19, wherein the disease or disorder is an age-related disease or disorder.
 32. The method of any one of claims 2-19, wherein the disease or disorder is osteoarthritis.
 33. The method of any one of claims 2-19, wherein the disease or disorder is intervertebral disc degeneration.
 34. The method of any one of claims 2-19, wherein the disease or disorder is atherosclerosis.
 35. The method of any one of claims 2-19, wherein the disease or disorder is hardening of the arteries.
 36. The method of any one of claims 2-19, wherein the disease or disorder is heart disease.
 37. The method of any one of claims 2-19, wherein the disease or disorder is neurodegeneration.
 38. The method of any one of claims 2-19, wherein the disease or disorder is neuroinflammation.
 39. The method of any one of claims 2-19, wherein the disease or disorder is Alzheimer's disease.
 40. The method of any one of claims 2-19, wherein the disease or disorder is myeloproliferative neoplasm (MPN).
 41. The method of any one of claims 2-19, wherein the disease or disorder is myelodysplastic syndrome (MDS).
 42. The method of any one of claims 2-19, wherein the disease or disorder is osteoporosis.
 43. The method of any one of claims 2-19, wherein the disease or disorder is age-induced frailty.
 44. The method of any one of claims 2-43, wherein the effective treatment comprises a reduction in fibrosis.
 45. The method of any one of claims 2-43, wherein the effective treatment comprises a reduction in inflammation.
 46. The method of any one of claims 2-43, wherein the effective treatment comprises a reduction in senescent cell numbers.
 47. The method of any one of claims 2-43, wherein the effective treatment comprises a reduction in relative senescent cell numbers.
 48. The method of any one of claims 2-43, wherein the effective treatment comprises an increase in organ or tissue function.
 49. The method of claim 1, wherein the senescent cell is selected from the group consisting of a senescent fibroblast, a senescent pre-adipocyte, a senescent epithelial cell, a senescent chondrocyte, a senescent intervertebral disc cell, a senescent satellite cell, a senescent muscle cell, a senescent liver cell, a senescent kidney cell, a senescent liver cell, a senescent kidney cell, a senescent bone marrow cell, a senescent vascular cell, a senescent immune cell, a senescent heart cell, a senescent arterial cell, a senescent venous cell, a senescent gastrointestinal cell, a senescent neuron, and a senescent endothelial cell.
 50. A composition comprising placenta-derived NK cells for use in killing senescent cells in a subject.
 51. Use of composition comprising placenta-derived NK cells for the manufacture of a medicament for killing senescent cells in a subject.
 52. A composition comprising placenta-derived NK cells for use treating a disease or disorder associated with cellular senescence in a subject. Use of a composition comprising placenta-derived NK cells for use treating a disease or disorder associated with cellular senescence in a subject.
-