

(19) **United States**

(12) **Patent Application Publication**
KUROSAKI et al.

(10) **Pub. No.: US 2025/0277003 A1**
(43) **Pub. Date: Sep. 4, 2025**

(54) **GLYCOSYLATED RBD AND USE THEREOF**

(71) Applicant: **OSAKA UNIVERSITY**, Suita-shi,
Osaka (JP)
(72) Inventors: **Tomohiro KUROSAKI**, Suita-shi (JP);
Ryo SHINNAKASU, Suita-shi (JP);
Shuhei SAKAKIBARA, Suita-shi (JP)
(73) Assignee: **OSAKA UNIVERSITY**, Suita-shi,
Osaka (JP)

(21) Appl. No.: **18/287,773**
(22) PCT Filed: **Apr. 22, 2022**
(86) PCT No.: **PCT/JP2022/018507**
§ 371 (c)(1),
(2) Date: **Oct. 20, 2023**

Related U.S. Application Data

(60) Provisional application No. 63/178,346, filed on Apr. 22, 2021.

Publication Classification

(51) **Int. Cl.**
C07K 14/005 (2006.01)
A61K 39/00 (2006.01)
A61K 39/215 (2006.01)
A61P 31/14 (2006.01)
(52) **U.S. Cl.**
CPC **C07K 14/005** (2013.01); **A61K 39/215**
(2013.01); **A61P 31/14** (2018.01); **A61K**
2039/575 (2013.01)

(57) **ABSTRACT**

As a technique for efficiently producing anti-coronavirus antibodies with cross-reactivity, provided is a polypeptide comprising an amino acid sequence having at least 60% or more identity to the amino acid sequence of SEQ ID NO: 2, wherein the amino acid sequence is modified so as to allow sugar chain attachment to positions corresponding to positions 85, 164, and 172 of the amino acid sequence of SEQ ID NO: 2.

Specification includes a Sequence Listing.





Fig. 1-1

A



B




Fig. 1-2




Fig. 2-1




Fig. 2-2




Figure 2 is a scatter plot showing the concentration of anti-CoV-2 IgG in serum (ng/ml) for different groups. The y-axis is logarithmic, ranging from 10^2 to 10^7 ng/ml. The x-axis categories are Pre-imm., CoV-2 wt, GM9, GM14, and CoV-1 wt. Data points are represented by different symbols: squares for CoV-2 wt, triangles for GM9, inverted triangles for GM14, and circles for CoV-1 wt. A horizontal dashed line at $10^{2.5}$ ng/ml indicates the detection limit. Statistical significance is indicated by asterisks (*, **, ***) above the data points.

Fig. 3-2




Fig. 4




Fig. 5-1




Fig. 5-2





Fig. 6

A



B




Fig. 7




Fig. 8




Fig. 9-1




Fig. 9-2





Fig. 9-3



GM14-2

	RBD Binding								Neutralization								
	Clonal lineage	VH mutation	VK mutation	CoV-1 (Clade 1a)	WIV1 (Clade 1a)	SHC014 (Clade 1a)	CoV-2 (Clade 1b)	PaGX (Clade 1b)	AL-103 (Clade 2)	Rs4081 (Clade 2)	Rf1/2004 (Clade 2)	BM48-31 (Clade 3)	CoV-1 (Clade 1a)	WIV1 (Clade 1a)	SHC014 (Clade 1a)	CoV-2 (Clade 1b)	PaGX (Clade 1b)
Clonal lineage:	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]
VH/V _K mutations:	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]
Binding (gMFI):	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]
Neutralization (IC ₅₀ , μg/ml):	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]	[Pattern]

Clonal lineage: [Color key] Colored (clonal lineages) [Box] No color (singletons)


VH/V_K mutations: [Scale 0-16]

Binding (gMFI): [Scale 0-30K]

Neutralization (IC₅₀, μg/ml): [Scale 10-0.1] [Box] >10 ND: Not determined

Fig. 9-5

B



C

	Affinity (K_D [M])									Neutralization (IC50 [μ g/ml])				
	Clade 1a			1b		2		3	Clade 1a			1b		
	CoV-1	WIV1	SHC014	CoV-2	PaGX	AL-103	Rs4048	RT1/2004	BM48-31	CoV-1	WIV1	SHC014	CoV-2	PaGX
Ab 15	1.97E-06	1.89E-06	2.21E-06	2.21E-06	1.43E-06	4.43E-06	7.84E-06	7.85E-06	7.31E-06	0.183	0.334	0.471	0.865	1.40
Ab 25	8.21E-06	7.59E-06	5.15E-06	4.54E-06	5.26E-06	5.01E-06	1.47E-06	8.21E-07	1.29E-06	0.135	0.135	0.191	0.701	0.262
Ab 47	5.55E-06	4.62E-06	3.47E-06	2.10E-06	1.90E-06	8.95E-06	1.62E-07	nb	nb	0.367	2.83	1.47	4.36	3.77
Ab 129	1.46E-06	2.07E-06	1.56E-06	2.79E-06	2.94E-06	nb	nb	nb	nb	1.07	1.27	>10	5.85	1.41
Ab 130	2.61E-06	1.59E-06	5.09E-07	1.90E-06	1.43E-06	nb	nb	nb	nb	2.58	4.69	>10	>10	>10
Ab 283	2.99E-06	7.20E-06	3.07E-06	2.92E-06	3.75E-06	nb	nb	nb	nb	1.68	1.12	>10	>10	>10
Ab 132	3.37E-06	3.36E-06	5.30E-07	2.52E-07	2.40E-06	nb	nb	nb	nb	0.942	2.86	>10	2.63	>10
Ab 137	1.94E-06	2.03E-06	4.51E-07	2.89E-06	1.31E-06	nb	nb	nb	nb	6.96	1.49	>10	>10	>10
Ab 185	2.76E-06	1.69E-06	5.10E-07	4.92E-06	2.48E-06	nb	nb	nb	nb	2.51	3.68	>10	>10	>10

Fig. 9-6




Fig. 10




Fig. 11




Fig. 12-2

mAb ID	HC					LC					Based on flow cytometry as-sey					K562 (test target)							
	IGHV	IGHD	IGHJ	CDR3	SEQ ID NO	SHM	AA num	NRKV	NRCL	CDR3	SEQ ID NO	SHM	AA num	Usage	ConV gMR	WCV gMR	ConV gMR	WCV gMR	4.1% (class3) gMR	15.4% (class3) gMR	ConV	WCV	ConV
60-2	IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	30	14	9	IGHV12-44	IGHJ2	GHVH33PPT	41	5	3	4	18122	17968	12487	71104	31541	>10	>10	>10	>10
	IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	30	14	7	IGHV12-44	IGHJ2	GHVH33PPT	41	5	3	4	12771	17556	11688	18156	5010	>10	>10	>10	>10
	IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	30	15	10	IGHV12-44	IGHJ2	GHVH33PPT	41	6	7	4	14321	11556	9210	13800	5239	>10	>10	>10	>10
	IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	31	17	12	IGHV12-44	IGHJ2	GHVH33PPT	42	7	5	4	15673	8716	6832	12444	3488	>10	>10	>10	>10
	IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	31	18	12	IGHV12-44	IGHJ2	GHVH33PPT	42	10	5	4	11643	8079	6850	7240	1898	>10	>10	>10	>10
	IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	32	17	13	IGHV12-44	IGHJ2	GHVH33PPT	42	6	3	4	12144	7547	7136	10375	2421	>10	>10	>10	>10
	IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	31	13	16	IGHV12-44	IGHJ2	GHVH33PPT	42	11	6	4	14888	14638	10863	16308	8667	>10	>10	>10	>10
	IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	30	13	8	IGHV12-44	IGHJ2	GHVH33PPT	41	4	3	4	18241	13834	14610	24079	11680	>10	>10	>10	>10
	IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	30	12	7	IGHV12-44	IGHJ2	GHVH33PPT	41	4	3	4	18080	20688	13162	23838	7859	>10	>10	>10	>10
	IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	30	14	14	IGHV12-44	IGHJ2	GHVH33PPT	41	11	6	4	14038	11610	11438	16315	5864	>10	>10	>10	>10
	IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	31	21	18	IGHV12-44	IGHJ2	GHVH33PPT	42	13	8	4	14194	3731	33172	13564	4381	>10	>10	>10	>10
	IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	30	11	6	IGHV12-44	IGHJ2	GHVH33PPT	41	0	5	4	14941	17502	15912	16956	5446	>10	>10	>10	>10
	IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	30	17	10	IGHV12-44	IGHJ2	GHVH33PPT	42	5	3	4	13861	7613	5790	3724	4531	>10	>10	>10	>10
	IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	30	11	7	IGHV12-44	IGHJ2	GHVH33PPT	41	10	8	4	13686	8503	2811	11642	3578	>10	>10	>10	>10
	IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	30	11	7	IGHV12-44	IGHJ2	GHVH33PPT	41	7	3	4	13875	10167	10872	14486	3945	>10	>10	>10	>10
	IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10
	IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10
	IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10
	IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10
	IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2	TCVGVGVGVGV	34	16	5	IGHV12-44	IGHJ2	GHVH33PPT	43	7	5	4	12347	15272	16638	4109	878	>10	>10	>10	>10	
IGHV1516	IGHD3-3	IGHJ2																					

12-3

[illegible]

Fig. 13

[illegible]

Fig. 14




Fig. 15

MFVFLVLLPLVSSQCVNLTTTRTQLPPAYTNSFTRGVYYPDKVFRSSVLHSTQDLFLPFFSN
 VTWFHAIHVSGTNGTKRFDNPVLPFNDGVYFASTEKSNIIRGWIFGTTLDSTQSLIVNN
 ATNVVIKVCEFQFCNDPFLGVYYHKNNKSWMESEFRVYSSANNCTFEYVSQPFLMDLE
 GKQGNFKNLREFVFKNIDGYFKIYSKHTPINLVRDLPQGFSALEPLVDLPIGINITRFQTLA
 LHRSYLTPGDSSSGWTAGAAAYYVGYLQPRTFLLKYNENGITITDAVDCALDPLSETKCTLK
 SFTVEKGIYQTSNFRVQPTESIVRFP**NITNLCPFGEVFNATRFASVYAWNRRKRISNCVA**
DYSVLYNSASFSTFKCYGVSPTKLNDLCFTNVYADSFVIRGDEVQRQIAPGQTGKIAD
YNYKLPPDDFTGCVIAWNSNNLDSKVGGNYNLYRLFRKSNLKPFERDISTEIQAG
STPCNGVEGFNCYFPLOSYGFQPTNGVGYQPYRVVLSFELLHAPATVCGPKKSTN
 LVKNKCVNFNFNGLTGTGVLTESNKKFLPFQQFGRDIADTTDAVRDPQTLEILDITPCSF
 GVSVITPGTNTSNQVAVLYQDVNCTEVPVAIHADQLTPTWRVYSTGSNVFQTRAGCLIG
 AEHVNNSEYCDIPGAGICASYQTQTNPRRARSVASQSIIAYTMSLGAENSVAYSNNNSIA
 IPTNFTISVTTEILPVSMTKTSVDCTMYICGDSTECSNLLLQYGSFCTQLNRALTGIAVEQD
 KNTQEVFAQVKQIYKTPPIKDFGGFNFSQILPDPSKPSKRSFIEDLLFNKVTADAGFIKQY
 GDCLGDIAARDLICAQKFNGLTVLPPLTDEMIAQYTSALLAGTITSGWTFGAGAALQIPF
 AMQMAYRFNGIGVTQNVLYENQKLIANQFNSAIGKIQDSLSTASALGKLQDVVNQNA
 QALNTLVKQLSSNFGAISSVLNDILSRDKVEAEVQIDRLITGRQLQSLQTYVTQQLIRAAEI
 RASANLAATKMSECVLGQSKRVDFCGKGYHLSFPQSAPHGVFLHVTYVPAQEKNT
 TAPAICHGKAHFPREGVFVSNGTHWFVTQRNFYEPQIITDNTFVSGNCDVVIGIVNN
 TVYDPLQPELDSFKEELDKYFKNHTSPDVLGDISGINASVVNIQKEIDRLNEVAKNLNES
 LIDLQELGKYEQYIKWPWYIWLGFIAGLIAIVMVTIMLCCMTSCCCLKGCCSCGSCCKFD
 EDDSEPVLKGVKLHYT

GLYCOSYLATED RBD AND USE THEREOF**CROSS REFERENCE TO RELATED APPLICATIONS**

[0001] This application is a National Stage of International Application No. PCT/JP2022/018507 filed Apr. 22, 2022, claiming priority/benefit based on U.S. Provisional Patent Application No. 63/178,346 filed Apr. 22, 2021.

INCORPORATION BY REFERENCE OF SEQUENCE LISTING

[0002] The content of the electronically submitted sequence listing, file name: Q292734_substitute sequence listing as filed; size: 45,290 bytes; and date of creation: Mar. 28, 2024, filed herewith, is incorporated herein by reference in its entirety.

TECHNICAL FIELD

[0003] Disclosed are techniques relating to sugar chain modification of the receptor-binding domain (RBD) of SARS-CoV-2 spike protein.

BACKGROUND ART

[0004] The COVID-19 pandemic, caused by the β -coronavirus severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), is a global health crisis. Coronavirus entry into host cells is mediated by the virus S protein, which forms trimeric spikes on the viral surface. The entry receptor for SARS-CoV-2 and SARS-CoV (herein called SARS-CoV-1) is the human cell-surface angiotensin converting enzyme 2 (ACE2), and the receptor-binding domain (RBD) of the spike from both of these viruses binds ACE2 with high affinity. Hence, it is believed that the RBD can become a primary target for neutralizing antibodies.

[0005] However, given that prior coronavirus epidemics (e.g., SARS-CoV-1 and MERS-CoV) have occurred due to zoonotic coronaviruses crossing the species barrier (Non-patent Literature (NPL) 1), the potential for the emergence of similar viruses in the future poses a significant threat to global public health, even in the face of effective vaccines for current viruses. For instance, one pre-emergent bat coronavirus, WIV1-CoV, is highly homologous to SARS-CoV-1 and SARS-CoV-2 and has been shown to be able to infect human ACE2-expressing cells (NPL 2). However, sera from SARS-CoV-2 infected individuals have shown very limited cross-neutralization of WIV1-CoV, except for rare individuals with low-levels of neutralizing antibodies (NPL 3).

CITATION LIST**Non-Patent Literature**

- [0006]** NPL 1: Wacharapluesadee, S., Tan, C. W., Maneecorn, P., Duengkae, P., Zhu, F., Joyjinda, Y., Kaewpom, T., Chia, W. N., Ampoot, W., Lim, B. L., et al. (2021). Evidence for SARS-CoV-2 related coronaviruses circulating in bats and pangolins in Southeast Asia. *Nat Commun* 12, 972.
- [0007]** NPL 2: Menachery, V. D., B. L. Yount Jr., K. Debbink, S. Agnihothram, L. E. Gralinski, J. A. Plante, R. L. Graham, T. Scobey, X. Y. Ge, E. F. Donaldson, et

al. 2015. A SARS-like cluster of circulating bat coronaviruses shows potential for human emergence. *Nat. Med.* 21: 1508-1513.

- [0008]** NPL 3: Garcia-Beltran, W. F., Lam, E. C., Astudillo, M. G., Yang, D., Miller, T. E., Feldman, J., Hauser, B. M., Caradonna, T. M., Clayton, K. L., Nitido, A. D., et al. (2021). COVID-19-neutralizing antibodies predict disease severity and survival. *Cell* 184, 476-488 e411.

SUMMARY OF INVENTION**Technical Problem**

[0009] An object is to provide a technique for obtaining an anti-coronavirus antibody with cross-reactivity.

Solution to Problem

[0010] The SARS-CoV-2 RBD is composed of two subdomains (FIG. 1A). The core-RBD subdomain consists of a central B sheet flanked by α -helices and presents a stable folded scaffold for the receptor-binding motif (RBM, residues 437-508) (Shang et al., 2020, Structural basis of receptor recognition by SARS-CoV-2. *Nature* 581, 221-224), which encodes ACE2 binding and receptor specificity (herein, the RBM-encoded region is called the head-RBD subdomain) (Shang et al., 2020, the same as above). Although initial analysis of convalescent sera from SARS-CoV-2-infected individuals demonstrated that many neutralizing antibodies (Class 1 and 2) recognize the head-RBD subdomain (FIG. 1B), cross-neutralization of known antibodies is very limited, as stated above. The present inventors thus conducted extensive research and found, in terms of broadly protective responses, that the core-RBD subdomain could be a more promising target.

[0011] One of the potential obstacles to targeting the core-RBD subdomain is that the epitopes in this subdomain are immuno-subdominant. Possible approaches to circumvent this problem include at least the following: i) altering the immunogen surface through targeted point mutations and deletions (Jardine et al., 2013, Rational HIV immunogen design to target specific germline B cell receptors. *Science* 340, 711-716; Impagliazzo et al., 2015, A stable trimeric influenza hemagglutinin stem as a broadly protective immunogen. *Science* 349, 1301-1306; Valkenburg et al., 2016, Stalking influenza by vaccination with pre-fusion headless HA mini-stem. *Sci Rep* 6, 22666); and ii) introducing N-linked glycans believed to shield the neighboring epitopes by means of the NxS/T sequons (Duan et al., 2018, Glycan Masking Focuses Immune Responses to the HIV-1 CD4-Binding Site and Enhances Elicitation of VRC01-Class Precursor Antibodies. *Immunity* 49, 301-311 e305; Eggink et al., 2014, Guiding the immune response against influenza virus hemagglutinin toward the conserved stalk domain by hyperglycosylation of the globular head domain. *J Virol* 88, 699-704). To facilitate the immune responses to the core-RBD subdomain, the present inventors introduced glycans into specific sites in the SARS-CoV-2 head-RBD subdomain, demonstrating that the glycan engineering facilitated the elicitation of potent cross-neutralizing antibodies towards SARS-CoV-1 and WIV1-CoV (clade 1 sarbecoviruses). The inventors conducted further study and research based on these findings to thus provide the subject matter represented by the following.

Item 1.

[0012] A polypeptide comprising an amino acid sequence having at least 60% or more identity to the amino acid sequence of SEQ ID NO: 2, wherein the amino acid sequence is modified so as to allow sugar chain attachment to positions corresponding to positions 85, 164, and 172 of the amino acid sequence of SEQ ID NO: 2.

Item 2.

[0013] The polypeptide according to Item 1, wherein the modified amino acid sequence is further modified so as to allow sugar chain attachment to a position corresponding to position 118 and/or a position corresponding to position 147 of the amino acid sequence of SEQ ID NO: 2.

Item 3.

[0014] The polypeptide according to Item 1 or 2, wherein the sugar chain is an N-type sugar chain.

Item 4.

[0015] The polypeptide according to any one of Items 1 to 3, wherein the sugar chain is attached to each of positions corresponding to positions 85, 164, and 172 of the amino acid sequence of SEQ ID NO: 2.

Item 5.

[0016] The polypeptide according to Item 4, wherein a sugar chain is further attached to a position corresponding to position 118 and/or a position corresponding to position 147.

Item 6.

[0017] The polypeptide according to Item 4 or 5, wherein the sugar chain is an N-type sugar chain.

Item 7.

[0018] A polynucleotide comprising a base sequence encoding the polypeptide of Item 1 or 2.

Item 8.

[0019] A composition comprising the polypeptide of any one of Items 1 to 6 or the polynucleotide of Item 7.

Item 9.

[0020] The composition according to Item 8, which is a vaccine.

Item 10.

[0021] A method for producing an anti-coronavirus antibody, comprising immunizing an animal with the polypeptide of any one of Items 1 to 6.

Item 11.

[0022] An antibody obtained by the method of Item 10.

Advantageous Effects of Invention

[0023] Provided are techniques for efficiently producing anti-coronavirus antibodies with cross-reactivity.

BRIEF DESCRIPTION OF DRAWINGS

[0024] FIG. 1 shows design of SARS-CoV-2 RBD glycan-masked mutants. (A) Structure of RBD of CoV-2 S (PDB: 6YZ5) that engages the ACE-2 ectodomain (PDB: 6M0J). (B) Epitopes of representative anti-CoV-2 mAbs. (C) (left) Glycosylation sites in RBD WT, GM9, and GM14. (C) (right) Ribbon models of RBD GM9 and GM14. The spheres are the introduced N-type glycan sites. (D) Glycan occupancy at N-linked glycosylation sites of RBD WT, GM9, and GM14 determined by LC-MS. (E) ELISA binding of previously reported mAbs against SARS-CoV-2 RBD (CB6, C002, S309, CR3022, and EY6A) to WT, GM9 and GM14. Anti-C. albicans hIgG1 mAb 23B12 was used for a negative control. The heatmap shows the percentage of binding (the value of the area under the curve [AUC] in ELISA for RBD WT is set at 100% for each mAb). Representative results from three independent experiments are shown.

[0025] FIG. 2 shows design and expression of CoV-2 RBD glycan-masked mutants. (A) Structure of RBD-ACE2 complex. (B) Parental amino acid sequences and introduced NXT sequons. (C) SDS-PAGE of RBD WT, GM9, and GM14.

[0026] FIG. 3 shows that glycan mutants of CoV-2 spike RBD immunogen effectively elicited CoV-1 S RBD specific antibodies. (A) Schematic overview of the experimental design. (B) CoV-2 RBD (left) or CoV-1 RBD (right) specific serum IgG1 antibody levels in serum from pre-immune mice, or CoV-2 RBD, GM9, GM14, or CoV-1 RBD WT immunized mice were measured by ELISA. Samples were pooled from three independent experiments (3-4 mice each group). (C) Class 3/4 type serum antibodies from CoV-2 RBD, GM9, GM14, or CoV-1 RBD WT immunized mice were measured by Epitope-Blocking ELISA with a mixture of CR3022/EY6A/5309 human IgG1 mAbs (left). Affinity measurements were determined by ELISA, expressed as the binding ratio to low density/high density of plate bound CoV-1 RBD protein (right). (D) WIV1 RBD-specific serum IgG1 antibody levels (left) and affinity (right) were measured as in (B) and (C, right), respectively. Horizontal lines indicate mean values; each symbol indicates one mouse. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; unpaired Student's t-test.

[0027] FIG. 4 shows that immunization with a nanoparticle antigen leads to the effective induction of CoV-2 S RBD specific antibodies. ELISA plots for CoV-2 RBD wt probe recognition of serum from respective biotin (+) RBD/St-avi nanoparticles, biotin (−) RBD/St-avi nanoparticles or only biotin (+) RBD immunized mice 3 weeks post primary immunization. Horizontal lines indicate mean values; each symbol indicates one mouse. The CR3022 mouse IgG1 monoclonal antibody (InvivoGen) was used as a standard. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; unpaired Student's t-test.

[0028] FIG. 5 shows that glycan engineering of the SARS-CoV-2 head-RBD elicited cross-neutralizing activities against SARS-CoV-1 and WIV1-CoV. (A) BALB/c mice were prime-boost-immunized with CoV-2 RBD WT, GM9, GM14 or CoV-1 RBD as shown in FIG. 3. Sera were collected 7 days after boost and pre-incubated with CoV-2 S, CoV-1 S, or WIV-1 CoV S-pseudotyped VSVΔG-luc for 1 h. The mixture was incubated with VeroE6 TMPRESS2 cells overnight. The NT50s of each group of mice are shown in the graphs. NT50s of the sample below the detection limit (1:20) was set as 20. * $P < 0.05$, Wilcoxon test. RBD WT (n=11); GM9 (n=11); GM14 (n=11); CoV-1 RBD (n=8). (B)

Neutralization activities of plasma samples from pre-pandemic healthy donors (HC, n=8) or convalescent COVID-19 patients (n=24) against CoV-2, CoV-1 and WIV-1 CoV S-pseudotyped VSVΔG-luc. NT50s of the sample below the detection limit (1:20) was set as 20. (C) Donor information. (D) ADE of infection by DENV1 single-round infectious particles (SRIP) by anti-DENV1 E mAb 3H12. DENV1 SRIP were pre-incubated with different concentrations of 3H12 or control mouse IgG1 and then incubated with RAW264.7 cells. The luciferase activity was measured 72 h post infection. (E) Sera of immunized mice were serially diluted and mixed with CoV-2 S pseudotyped VSVΔG-luc for 1 h and incubated with RAW264.7 cells. The luciferase activity was measured 24 h post infection. Representative results from two to three independent experiments are shown. Data are mean±SEM. Dotted lines in the graphs (NT50=20) represent the lower limit of detection (A and B). *, p<0.05; **, p<0.01; ***, p<0.001; unpaired Student's t-test.

[0029] FIG. 6 shows cross-reactivity of sera and GC B cells in mice immunized with GM9 or GM14. (A) ELISA plots for CoV-2 RBD wt probe recognition of serum of mice after prime/boost with GM9 or GM14. Sera were collected at 3 weeks post primary immunization. (B) Representative FACS plots of GC B from dLNs after immunization. Cells were gated for antigen-binding IgG+GC B cells (CD138-B220+IgG+GL7+Fas+). The graph (right) shows the percentage of positive cells for RBD wt binding among GM9 or GM14-binding cells (n=4 each).

[0030] FIG. 7 shows that immunization with CoV-2 GM antigens leads to the activation of CoV-1 and CoV-2 cross-reactive GC B cells. (A) Representative flow cytometry plots analyzing GC B cells (CD138-B220+IgG+GL7+Fas+) from lymph nodes of mice 3 weeks post primary immunization with each antigen. The plots are representative of findings from 2 independent experiments (left). Quantification of absolute numbers of CD138-B220+IgG+GL7+Fas+ GC B cells (right) from multiple mice in one experiment. Results are representative of 2 independent experiments. Horizontal lines indicate mean values; each symbol indicates one mouse. (B) Representative analytical results by flow cytometry of CoV-2 RBD-binding MLN GC B cells (CD138-B220+IgG+GL7+Fas+CoV-2 RBD(APC)+CoV-2 RBD). Plots are representative of findings from 2 independent experiments (left). Absolute numbers of GC B cells (right) from multiple mice in one experiment are shown. Results are representative of 2 independent experiments. (C) Cross-reactivity of CoV-2 RBD-binding MLN GC B cells (CD138-B220+IgG+GL7+Fas+CoV-2 RBD(APC)+CoV-2 RBD(PE-Cy7+)) with CoV-1 RBD assessed by flow cytometry. Plots are representative of findings from 2 independent experiments (left). The frequency of CoV-1 and CoV-2 RBD cross-reactive GC B cells (right) from multiple mice in one experiment. Results are representative of 2 independent experiments. Horizontal lines indicate mean values; each symbol indicates one mouse. *, P<0.05; **, P<0.01; ***, P<0.001; unpaired Student's t-test.

[0031] FIG. 8 shows authentic virus neutralizing assay. A mixture of 100 TCID50 virus and serially diluted, heat-inactivated plasma samples were incubated for 1 h before being placed on VeroE6-TMPRSS2 cells seeded in 96-well plates. After culturing for 4 days, cells were fixed with formalin and stained with crystal violet solution. Cut-off dilution index with >50% cytopathic effect was presented as

microneutralization titer. Microneutralization titer of the sample below the detection limit (1:40 dilution) was set as 20. *, P<0.05; **, P<0.01; unpaired Student's t-test.

[0032] FIG. 9 shows that GC-derived mAbs exhibited breadth of RBD binding and neutralization against clade 1 SARS-related CoVs. (A) Clonality, VH and VK mutations and antibody class, binding to S RBDs of CoV-1, WIV1-CoV, CoV-2, BatCoV Anlong-103 (AL-103) and BtCoV BM48-31, neutralizing activity against CoV-1, WIV1-CoV and CoV-2 pseudoviruses of mAbs derived from GC B cells of mice immunized with GM9 (GM9-1 and GM9-2) or GM14 (GM14-1 and GM14-2). (B) Correlation between mAb binding to CoV-2 RBD and CoV-1 or WIV1-CoV RBD. Binding of mAbs with CoV-1, WIV1 and CoV-2 RBDs were measured by bead-based flow cytometric assays. The binding signals to respective RBDs (gMFI) are plotted in the graph. Spearman's rank correlation coefficients and p values are shown. (C) Binding affinity of mAbs with RBD proteins. The equilibrium dissociation constants (M) and IC₅₀s (μg/ml) of individual clones are shown. (D) Neutralizing activity (IC₅₀ [μg/ml]) of representative cross-reactive mAbs and mouse IgG2c recombinant antibodies carrying the variable regions of previously isolated S304 (class 3) and CR3022 (class 4) human anti-RBD antibodies.

[0033] FIG. 10 shows correlation between gMFI from bead-based flow-cytometry assay and RMax from biolayer interferometry (Octet) assay. gMFI and KD(M), RMax, kon (1/Ms) and koff (1/s) of bivalent mAb-CoV-2 RBD interactions are shown. Spearman's rank test.

[0034] FIG. 11 shows generation of cross-reactive memory B cells and long-lived plasma cells in mice immunized with GM9 or GM14. (A) Schematic overview of the experimental design. (B) Cross-reactivity of CoV-2 RBD-specific memory B cells (CD138-B220+IgG1+CD38+GL7-CoV-2 RBD(APC)+CoV-2 RBD(PE-Cy7+)) with CoV-1 RBD from spleen of mice one month post boost immunization with each antigen. Frequency of CoV-1 and CoV-2 RBD cross-reactive memory B cells (upper) and quantification of absolute numbers of total CoV-2-specific memory B cells (lower) from multiple mice in one experiment. Results are representative of 2 independent experiments. (C) ELISPOT analysis of CoV-1- (left) or CoV-2-(right) specific IgG1 ASC responses of pre-immune mice, or one month post boost immunizations with each antigen. Results are representative of 2 independent experiments. Horizontal lines indicate mean values; each symbol indicates one mouse. *, P<0.05; **, P<0.01; ***, P<0.001; unpaired Student's t-test.

[0035] FIG. 12 shows characteristics regarding mAb generated from GC B cells of mice immunized with GM9 or GM14.

[0036] FIG. 13 shows comparisons among the RBD sequences of representative sarbecoviruses of the genus β-coronavirus. The regions enclosed by squares are receptor binding motifs (RBD head regions). The star symbols represent sites into which the sugar chain modification was inserted. The RBD amino acid sequences used for the alignment in FIG. 13 are SEQ ID NOs: 2, 107, 108, 109, 110, 111, 112, 113, and 114 in this order from the top.

[0037] FIG. 14 shows a phylogenetic tree of RBD sequences of representative sarbecoviruses of the genus β-coronavirus. The numerical values in parentheses indicate homology to SARS-Cov-2 RBD.

[0038] FIG. 15 shows the amino acid sequence (SEQ ID NO: 1) of the SARS-CoV-2 spike protein. The region shown in bold is the receptor binding domain (SEQ ID NO: 2).

DESCRIPTION OF EMBODIMENTS

1. Definition

[0039] In the present specification, the terms “comprising,” “containing,” and “having” include the concepts of containing, including, consisting essentially of, and consisting of.

[0040] In the present specification, the “identity” of amino acid sequences refers to the degree of identicalness of two or more amino acid sequences that can be compared with each other. Thus, the higher the identicalness of two amino acid sequences, the higher the identity or similarity of these sequences. The level of amino acid sequence identity is determined, for example, using FASTA, which is a tool for sequence analysis, with default parameters. The level of amino acid sequence identity can otherwise be determined using the algorithm BLAST by Karlin and Altschul. A program called “BLASTX,” which is based on such an algorithm of BLAST, has been developed. Specific procedures of these analysis methods are known, and reference may be made to the website (<http://www.ncbi.nlm.nih.gov/>) of the National Center of Biotechnology Information (NCBI). The “identity” of base sequences is also defined accordingly.

[0041] In the present specification, the term “corresponding positions” in terms of amino acid sequences and nucleotide sequences can be determined by aligning (alignment of) a target sequence and a reference sequence (e.g., the amino acid sequence of SEQ ID NO: 2) so as to achieve maximum homology between the residues present in each sequence. Alignment can be conducted by using known algorithms, and the procedures thereof are known to those skilled in the art. For example, alignment can be performed by using the Clustal W multiple alignment program (Nucleic Acids Res, 1994, 22:4673-4680) with default settings. Alternatively, Clustal W2 or Clustal omega, revised versions of Clustal W, can also be used. Clustal W, Clustal W2, and Clustal omega are available, for example, on the websites of the European Bioinformatics Institute (EBI [www.ebi.ac.uk/index.html]) and the DNA Data Bank of Japan operated by the National Institute of Genetics (DDBJ [www.ddbj.nig.ac.jp/Welcome-j.html])).

[0042] In the present specification, “conservative substitution” means a substitution of an amino acid residue with an amino acid residue having a similar side chain. For example, a substitution between amino acid residues having a basic side chain, such as lysine, arginine, and histidine, is considered to be a conservative substitution. Other examples that are considered to be a conservative substitution include a substitution between amino acid residues having an acidic side chain, such as aspartic acid and glutamic acid; a substitution between amino acid residues having an uncharged polar side chain, such as glycine, asparagine, glutamine, serine, threonine, tyrosine, and cysteine; a substitution between amino acid residues having a nonpolar side chain, such as alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, and tryptophan; a substitution between amino acid residues having a R-branched side chain, such as threonine, valine, and isoleucine; and a

substitution between amino acid residues having an aromatic side chain, such as tyrosine, phenylalanine, tryptophan, and histidine.

2. Polypeptide

[0043] The polypeptide preferably has a certain degree or more of identity to the amino acid sequence of SEQ ID NO: 2. The certain degree or more means, for example, 60% or more, 65% or more, 70% or more, 75% or more, 80% or more, 85% or more, 90% or more, 91% or more, 92% or more, 93% or more, 94% or more, 95% or more, 96% or more, 97% or more, 98% or more, or 99% or more.

[0044] SEQ ID NO: 2 is the amino acid sequence of RBD composed of 199 amino acid residues at positions 331 to 529 of the amino acid sequence of SARS-CoV-2 S protein (SEQ ID NO: 1). In one embodiment, the polypeptide may be shorter than 199 residues in length or longer than 199 amino acid residues in length. In one embodiment, the polypeptide is 1 residue, 2 residues, 3 residues, 4 residues, 5 residues, 6 residues, 7 residues, 8 residues, 9 residues, or 10 residues longer than 199 residues in length, or 1 residue, 2 residues, 3 residues, 4 residues, 5 residues, 6 residues, 7 residues, 8 residues, 9 residues, or 10 residues shorter than 199 residues in length. When the polypeptide is longer than 199 residues in length, the longer sequence portion preferably has a certain degree or more of identity to the corresponding amino acid sequence of SEQ ID NO: 1. The meaning of the phrase “certain degree or more” used here is as described above. In one embodiment, the polypeptide may comprise, for example, a Tag sequence at the C- or N-terminus.

[0045] In one embodiment, if the amino acid sequence of the polypeptide does not exactly match the amino acid sequence of SEQ ID NO: 1 or 2, the mismatched amino acid residues are preferably based on conservative amino acid substitutions in the corresponding amino acid residues in the amino acid sequence of SEQ ID NO: 1 or 2.

[0046] The polypeptide preferably has an amino acid sequence that is modified so as to allow sugar chain attachment to positions corresponding to positions 85, 164 and 172 of the amino acid sequence of SEQ ID NO: 2. The type of sugar chain is not limited and may be either an N-type sugar chain or O-type sugar chain. In one embodiment, the sugar chain is preferably an N-type sugar chain.

[0047] Techniques for sugar chain attachment to specific positions of a polypeptide are known. For example, for N-type sugar chain attachment to a position corresponding to position 85 of the amino acid sequence of SEQ ID NO: 2, the amino acid residue corresponding to position 85 of the amino acid sequence of SEQ ID NO: 2 is preferably an aspartic acid residue while the amino acid residue corresponding to position 87 is preferably a serine residue or a threonine residue (preferably a threonine residue). Similarly, for N-type sugar chain attachment to a position corresponding to position 164 of the amino acid sequence of SEQ ID NO: 2, the amino acid residue corresponding to position 164 of the amino acid sequence of SEQ ID NO: 2 is preferably an aspartic acid residue while the amino acid residue corresponding to position 166 is preferably a serine residue or a threonine residue (preferably a threonine residue). Similarly, for N-type sugar chain attachment to a position corresponding to position 172 of the amino acid sequence of SEQ ID NO: 2, the amino acid residue corresponding to position 172 of the amino acid sequence of SEQ ID NO: 2 is preferably an aspartic acid residue while the amino acid

residue corresponding to position 174 is preferably a serine residue or a threonine residue.

[0048] In one embodiment, the polypeptide preferably has an amino acid sequence that is modified so as to allow sugar chain attachment to a position corresponding to position 118 and/or a position corresponding to position 147, in addition to positions corresponding to positions 85, 164, and 172 of the amino acid sequence in SEQ ID NO: 2. In one embodiment, the polypeptide preferably has an amino acid sequence that is modified so as to allow sugar chain attachment to positions corresponding to positions 118 and 147, in addition to positions corresponding to positions 85, 164, and 172 of the amino acid sequence of SEQ ID NO: 2. In one embodiment, the polypeptide preferably has an amino acid sequence that is modified so as to allow sugar chain attachment to positions corresponding to positions 85, 164, and 172 of the amino acid sequence of SEQ ID NO: 2, wherein the amino acid sequence is not modified so as to allow sugar chain attachment to a position corresponding to position 118 and/or a position corresponding to position 147 of the amino acid sequence of SEQ ID NO: 2. In one embodiment, the polypeptide preferably has an amino acid sequence that is not modified so as to allow sugar chain attachment to at least one position corresponding to at least one position selected from the group consisting of positions 87, 126, 141, 152, 154, and 157 of the amino acid sequence of SEQ ID NO: 2, from the viewpoint of allowing stable expression thereof. The phrase “at least one” used here can mean at least two, at least three, at least four, at least five, or at least six. The type of sugar chain is not limited and may be either an N-type sugar chain or O-type sugar chain. In one embodiment, the sugar chain is preferably an N-type sugar chain.

[0049] For N-type sugar chain attachment to a position corresponding to position 118 of the amino acid sequence of SEQ ID NO: 2, the amino acid residue corresponding to position 118 of the amino acid sequence of SEQ ID NO: 2 is preferably an aspartic acid residue, and the amino acid residue corresponding to position 120 is preferably a serine residue or a threonine residue (preferably a threonine residue). Similarly, for N-type sugar chain attachment to a position corresponding to position 147 of the amino acid sequence of SEQ ID NO: 2, the amino acid residue corresponding to position 147 of the amino acid sequence of SEQ ID NO: 2 is preferably an aspartic acid residue, and the amino acid residue corresponding to position 149 is preferably a serine residue or a threonine residue (preferably a threonine residue).

[0050] In one embodiment of the polypeptide, in the amino acid sequence of SEQ ID NO: 2, the amino acid residues at positions 85, 164, and 172 are preferably substituted with an aspartic acid residue, and the amino acid residues at positions 87, 166, and 174 are preferably substituted with a serine residue or a threonine residue (preferably a threonine residue). In one embodiment of the polypeptide, in the amino acid sequence of SEQ ID NO: 2, the amino acid residue at positions 85, 118, 147, 164, and 172 are preferably substituted with an aspartic acid residue, and the amino acid residues at positions 87, 120, 149, 166, and 174 are preferably substituted with a serine residue or a threonine residue (preferably, a threonine residue).

[0051] In one embodiment, the polypeptide preferably has a certain degree or more of glycan occupancy at one or more positions corresponding to one or more positions selected from the group consisting of positions 85, 118, 147, 164, and

172 of the amino acid sequence of SEQ ID NO: 2. The glycan occupancy can be measured by the method described in the Examples below. The phrase “certain degree or more” means, for example, 30% or more, 40% or more, 50% or more, 60% or more, 70% or more, 80% or more, or 90% or more. In one embodiment, the phrase “one or more” includes two or more, three or more, or four or more.

[0052] The polypeptide may be in the form of a pharmaceutically acceptable salt formed with an acid or base. The salt is not limited as long as it is a pharmaceutically acceptable salt, and may be either an acid salt or a basic salt. Examples of acid salts include inorganic acid salts, such as hydrochloride, hydrobromide, sulfate, nitrate, and phosphate; organic acid salts, such as acetate, propionate, tartrate, fumarate, maleate, malate, citrate, methanesulfonate, and para-toluenesulfonate; and amino acid salts, such as aspartate and glutamate. Examples of basic salts include alkali metal salts, such as sodium salts and potassium salts; and alkaline earth metal salts, such as calcium salts and magnesium salts.

[0053] The polypeptide may otherwise be in the form of a solvate. The solvent is not limited as long as it is pharmaceutically acceptable, and may be, for example, water, ethanol, glycerol, or acetic acid.

[0054] The polypeptide can be easily prepared according to a known genetic engineering method, according to its amino acid sequence. For example, the polypeptide can be prepared using PCR, restriction enzyme cleavage, a DNA ligation technique, an in vitro transcription/translation technique, a recombinant protein production technique, etc. In one embodiment, the polypeptide can be obtained by expressing a polynucleotide encoding the polypeptide in an appropriate host.

[0055] The polypeptide may be purified after synthesis. For example, the polypeptide may be obtained by extraction from cells harvested and collected by centrifugation, filtration, etc. from a culture product. Specifically, at the first step of extraction, the cells can be destroyed by means of digestion with an enzyme, destruction caused by osmotic pressure, a sudden increase or decrease in pressure, sonication, various homogenizers, etc. The destroyed cells can then be fractionated by physical means such as low-speed centrifugation, ultra-centrifugation, filtration, molecular sieving, and membrane concentration, chemical means such as a precipitant, a solubilizer, an adsorption/desorption agent, and a dispersant, or physicochemical means such as electrophoresis, column chromatography, a support, dialysis, and salting-out. These techniques may be used in combination. In applying these techniques, physicochemical conditions, such as temperature, pressure, pH, and ion strength, can be suitably set.

[0056] The polypeptide is useful for efficiently producing anti-coronavirus antibodies with cross-reactivity. In one embodiment, cross-reactivity means reactivity to SARS-CoV-2 and coronaviruses other than SARS-CoV-2. In one embodiment, the coronaviruses other than SARS-CoV-2 are those belonging to the genus β -coronavirus, and preferably those belonging to the subgenus sarbecovirus of the genus β -coronavirus. In one embodiment, the coronaviruses other than SARS-CoV-2 are SARS-related coronaviruses. In one embodiment, examples of the coronaviruses other than SARS-CoV-2 include SARS-CoV-1, WIV1-CoV, MERS-CoV, and Pangolin-CoV. In one embodiment, the anti-coronavirus antibodies with cross-reactivity are preferably

neutralizing antibodies, and preferably broadly neutralizing antibodies. Such antibodies are useful in preventing, treating, or alleviating diseases or conditions caused by coronaviruses. Accordingly, the polypeptide can be used as a vaccine or therapeutic agent for these purposes.

3. Polynucleotide

[0057] The polynucleotide preferably has a base sequence encoding the polypeptide described above. In one embodiment, the polynucleotide is preferably in the form of an expression cassette of the polypeptide. The expression cassette is not limited as long as it is a polynucleotide that is capable of expressing a polypeptide in cells. Typical example of the expression cassette is a polynucleotide comprising a promoter and a base sequence that encodes a polypeptide under the control of the promoter.

[0058] The promoter contained in the expression cassette is not limited and can be suitably selected according to, for example, the type of cells into which the expression cassette is introduced. Examples include pol II promoters. Specific examples include a CMV promoter, EF1 promoter, SV40 promoter, MSCV promoter, hTERT promoter, β -actin promoter, and CAG promoter. Other examples include a tryptophan promoter, lac promoter, T7 promoter, T5 promoter, T3 promoter, SP6 promoter, arabinose-induced promoter, cold-shock promoter, and tetracycline-induced promoter.

[0059] The cell type (origin) used to introduce the polynucleotide and produce the polypeptide is not limited. Examples include animal cells, plant cells, and fungi, such as yeast. Particular examples of animals include various mammals, such as humans, monkeys, mice, rats, dogs, cats, rabbits, pigs, horses, cows, sheep, goats, and deer, non-mammalian vertebrates, and invertebrates. In one embodiment, mammal cells are preferred. The type of cells is not limited, and cells from various tissues or cells having various properties can be used. Examples include blood cells, hematopoietic stem cells/progenitor cells, gametes (sperm and ovum), fibroblast cells, epithelial cells, vascular endothelial cells, nerve cells, liver cells, keratinocytes, muscle cells, epidermal cells, endocrine cells, ES cells, iPS cells, tissue stem cells, and cancer cells.

[0060] The expression cassette can comprise other elements (e.g., multiple cloning sites (MCS), drug resistance genes, replication origins, enhancer sequences, repressor sequences, insulator sequences, reporter proteins (e.g., fluorescent protein etc.)—coding sequences and drug resistance gene-coding sequences), if necessary. The MCS is not limited as long as it contains multiple (e.g., 2 to 50, preferably 2 to 20, and more preferably 2 to 10) restriction enzyme sites.

[0061] Examples of drug resistance genes include chloramphenicol resistance genes, tetracycline resistance genes, neomycin resistance genes, erythromycin resistance genes, spectinomycin resistance genes, kanamycin resistance genes, hygromycin resistance genes, and puromycin resistance genes.

[0062] The reporter protein is not limited as long as it is a light-emitting (color-developing) protein, which emits light (develops a color) by reacting with a specific substrate, or it is a fluorescent protein, which emits fluorescence from excitation light. Examples of light-emitting (color-developing) proteins include luciferase, β -galactosidase, chloramphenicol acetyltransferase, and R-glucuronidase. Examples of fluorescent proteins include GFP, Azami-Green, ZsGreen,

GFP2, HyPer, Sirius, BFP, CFP, Turquoise, Cyan, TFP1, YFP, Venus, ZsYellow, Banana, Kusabira-Orange, RFP, DsRed, AsRed, Strawberry, Jred, KillerRed, Cherry, HcRed, and mPlum.

[0063] The expression cassette can constitute an expression vector, together with another sequence, if necessary. The type of vector is not limited, and examples include plasmid vectors of animal cell expression plasmids etc.; and virus vectors of retroviruses, lentiviruses, adenoviruses, adeno-associated viruses, herpes viruses, Sendai viruses, etc.

[0064] The polynucleotide can be easily prepared according to a known genetic engineering method. For example, the polynucleotide can be prepared using PCR, restriction enzyme cleavage, a DNA ligation technique, etc.

[0065] The polynucleotide enables the production of the polypeptide described above. Thus, the polynucleotide can be used for the same purposes as in the polypeptide described above (e.g., a vaccine for diseases or symptoms caused by coronaviruses).

4. Composition

[0066] The composition preferably comprises the polypeptide or polynucleotide described above. The composition comprising the polypeptide or polynucleotide described above is useful as a vaccine or a pharmaceutical composition.

[0067] The compositions may comprise other components according to the purpose. Examples of other components include a stabilizer for increasing the heat resistance of the vaccine and an adjuvant as an auxiliary for enhancing immunogenicity. Examples of stabilizers include sugars and amino acids. Examples of adjuvants include aluminum compounds (e.g., aluminum hydroxide gel), CpG oligodeoxynucleotide, mineral oil, vegetable oil, alum, bentonite, silica, muramyl dipeptide derivatives, thymosin, and interleukin. The adjuvants may be used alone or in a combination of two or more.

[0068] The composition can comprise bases, carriers, solvents, dispersants, emulsifiers, buffers, stabilizers, excipients, binders, disintegrates, lubricants, thickeners, humectants, colorants, fragrances, chelating agents, and the like, if necessary.

[0069] Examples of the dosage form of the vaccine or pharmaceutical composition include, but are not limited to, injections, such as aqueous injections, non-aqueous injections, suspension injections, and solid injections; oral preparations, such as tablets, capsules, granules, powders, fine granules, syrups, enteric agents, sustained-release capsules, chewable tablets, drops, pills, liquids for internal use, confectionery tablets, sustained-release preparations, and sustained-release granules; and preparations for external use, such as nasal drops, inhalants, rectal suppositories, inserts, enemas, and jellies.

[0070] The content of the active ingredient in the composition can be set based on, for example, the administration subject, the administration route, the dosage form, the conditions of the patient, and the determination of the doctor. For example, the content of the polypeptide or polynucleotide in the composition may be 0.0001 to 95 wt %, and preferably 0.001 to 50 wt %.

[0071] The amount of the vaccine or pharmaceutical composition for use can be set in consideration of various factors, such as the administration route, the health condi-

tions of the subject, the age, gender, body weight of the subject, pharmacological findings such as pharmacokinetics and toxicological characteristics, use or non-use of a drug delivery system, whether it is administered as part of a combinational drug with other drugs, acquisition of immunity, and acquisition of boosting effects. For example, the vaccine or pharmaceutical composition can be used such that the administration amount of the active ingredient per single administration is 1 µg to 100 mg/kg (body weight) or 50 µg to 10 mg/kg (body weight). The interval and frequency of administration is not limited, and the administration is preferably performed, for example, at an interval of about 1 to 30 weeks.

5. Antibody and Production Method Therefor

[0072] An anti-coronavirus antibody with cross-reactivity can be produced by using the polypeptide described above as an antigen.

[0073] Examples of the type of the antibody include polyclonal antibodies, monoclonal antibodies, human chimeric antibodies, humanized antibodies, fully human antibodies, single-stranded antibodies, and antigen-binding fragments (Fab, F(ab')₂, minibody, scFv-Fc, Fv, scFv, diabody, triabody, and tetrabody.) A human chimeric antibody is an antibody in which the sequence of the variable region of the antibody is from an animal other than a human (e.g. a mouse or cow) while the sequence of the constant region of the antibody is from a human.

[0074] Methods for producing these antibodies are known, and ordinary methods can be used for the production (Current Protocols in Molecular Biology, Chapter 11.12 to 11.13 (2000)). Specifically, for a polyclonal antibody, an antigen can be immunized into a non-human animal, such as a domestic rabbit, to obtain the antibody from the serum of the immunized animal according to an ordinary method. A monoclonal antibody can be obtained from hybridoma cells prepared by immunizing a non-human animal, such as a mouse or a cow, with an antigen, followed by cell fusion of the resulting spleen cells with myeloma cells (Current Protocols in Molecular Biology, edit. Ausubel et al. (1987), publish. John Wiley and Sons, Section 11.4 to 11.11). Immunization of an animal can be conducted by administering the polypeptide described above to an animal or by administering the polynucleotide to the animal and allowing it to express in the animal's body.

[0075] The antibody can also be produced by enhancing immunological reactions by using various adjuvants according to the host type. Examples of such adjuvants include mineral gels, such as Freund's adjuvant and aluminum hydroxide; surface-active substances, such as lysolecithin, pluronic polyol, polyanion, peptide, oil emulsion, keyhole limpet hemocyanin, and dinitrophenol; and human adjuvants, such as BCG (Bacille de Calmette et Guérin) and Corynebacterium parvum.

[0076] In one embodiment, cross-reactivity of an antibody means reactivity to SARS-CoV-2 and coronaviruses other than SARS-CoV-2. In one embodiment, the coronaviruses other than SARS-CoV-2 are those belonging to the genus β-coronavirus, and preferably those belonging to the sub-genus Sarbecovirus of the genus β-coronavirus. In one embodiment, the coronaviruses other than SARS-CoV-2 are SARS-related coronaviruses. In one embodiment, examples of the coronaviruses other than SARS-CoV-2 include SARS-CoV-1, WIV1-CoV, MERS-CoV, and Pangolin-CoV.

In one embodiment, the antibody preferably recognizes the core-RBD subdomain of the SARS-CoV-2 S protein. In one embodiment, the anti-coronavirus antibodies with cross-reactivity are preferably neutralizing antibodies, and preferably broadly neutralizing antibodies. Such antibodies are useful in preventing, treating, or alleviating diseases or conditions caused by coronaviruses.

EXAMPLES

Materials and Test Methods

Mice

[0077] BALB/c mice were purchased from SLC Japan and maintained under specific-pathogen-free conditions. Sex-matched 7- to 8-week-old mice were used for all experiments. All animal experiments were conducted in accordance with the animal experiment guidelines of Osaka University.

Human Subjects

[0078] Human blood samples were collected at Tokyo Shinagawa Hospital and plasma was isolated using Vacutainer CPT Tubes (Becton, Dickinson). The study protocol was approved by the National Institute of Infectious Diseases Ethic Review Board for Human Subjects, the ethics committees of Tokyo Shinagawa Hospital, and Osaka University. All participants provided written informed consent in accordance with the Declaration of Helsinki.

Vaccine Design

[0079] The mammalian expression construct for parental rRBD (pCAGGS-CoV-2 RBD-His-Avi) encodes aa 331-530 of SARSCoV-2 S which has MERS S protein-derived signal peptide (MIHSVFLLMFLLTPTESYVD) at the N terminus and a 6×His-Avi-tag (HHHHHHGLNDIFEAQKIEWHE) at the C terminus. For glycan masking, NxT sequons were introduced in the RBD surface residues shown in FIG. 2. Mutations expected to disrupt the structure of RBD or with the low glycosylation score (<0.5) by NetNGlyc (<http://www.cbs.dtu.dk/services/NetNGlyc/>) were avoided. The candidate constructs were transfected into Expi293F cells to examine whether added glycosylation prevented RBD expression. Antigenicity of the GM mutants was evaluated by ELISA using known anti-RBD antibodies.

Recombinant Protein Expression and Purification

[0080] The plasmids for the recombinant SARS-CoV RBD and GM mutants were transiently transfected into Expi293F cells (Thermo Fisher Scientific) with the ExpiFectamine 293 Transfection Kit (Thermo Fisher Scientific) according to the manufacturer's protocol. After 3-4 days, supernatants were collected and passed through a 0.45-micrometer filter. The recombinant proteins were purified from supernatants by Talon metal affinity resin (Takara) according to the manufacturer's protocol. The elute from the resin was concentrated using an Amicon Ultra 4 10,000 NWML, in which the elution buffer was exchanged to PBS. For nanoparticle antigens and probes, post-translational biotinylation of rRBD and GM mutants was performed in culture via co-expression of the BirA enzyme by co-transfection of a BirA-Flag expression vector (Addgene). Cells

were maintained in culture supplemented with 100 μ M biotin (Sigma-Aldrich) after transfection as described above. **[0081]** For preparation of ELISA or ELISPOT antigens, mammalian expression constructs of CoV-2 RBD WT, GM9, GM14 and SARS-related CoV RBDs containing a thrombin cleavage site (LVPRGS) in front of the C-terminal 6 $\times$ His-Avi-tag were transiently transfected into Expi293F cells. Purified proteins were treated with thrombin using a Thrombin kit (Merck) according to the manufacturer's protocol. The cleaved 6 $\times$ His-AviTag and thrombin were

3.1 (Thermo Fisher Scientific). The search parameters were as follows: a mass tolerance of ± 5 ppm, confidence score of > 80 and ID type of MS2. Carboxymethylation (+58.005 Da) was set as a static modification of Cys residues. A human glycan database stored in the software was used for glycopeptide identifications. The integrated peak area of the multiple precursor ions from each glycopeptide was calculated, and the relative peak area and glycan occupancy (%) were calculated by the following formula:

$$\begin{aligned} \text{Relative peak area} &= (\text{The peak area of glycopeptide}) / \\ &\quad (\text{The total peak area of glycopeptides and nonglycosylated peptide}) \\ \text{Glycan occupancy (\%)} &= (\text{The total relative peak areas of glycopeptides}) / \\ &\quad (\text{The total relative peak areas of glycopeptides and nonglycosylated peptide}) \times 100 \end{aligned}$$

removed by Talon metal affinity resin (Takara) and Benzamidine Sepharose (GE), respectively. Purity or biotinylation efficiency of recombinant proteins was confirmed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE).

Anti-CoV-2 S RBD Monoclonal Antibodies

[0082] DNA fragments encoding heavy and kappa or lambda chain variable regions from known anti-CoV-2 mAbs (Pinto et al., 2020, Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV antibody. *Nature* 583, 290-295; Yuan et al., 2020, Structural basis of a shared antibody response to SARS-CoV-2. *Science* 369, 1119-1123; Zhou et al., 2020, Structural basis for the neutralization of SARS-CoV-2 by an antibody from a convalescent patient. *Nat Struct Mol Biol* 27, 950-958) were synthesized (Eurofins) and cloned into human IgG1, Igkappa and Iglambda expression vectors, respectively. The heavy and light chain expression vectors were co-transfected into Expi293F. Respective mAbs were purified with Protein G Sepharose (GE) according to the manufacturer's protocol.

N-Linked Glycan Occupancy Analysis by Liquid Chromatography/Mass Spectrometry (LC/MS)

[0083] Sample preparation was performed by referring to Tajiri-Tsukada et al., 2020 (Establishment of a highly precise multi-attribute method for the characterization and quality control of therapeutic monoclonal antibodies. *Bioengineered* 11, 984-1000). The sample protein (10 μ g) was treated using MPEX PTS Reagents (GL Sciences). The reduced and carboxymethylated protein was digested with 2 μ g of chymotrypsin (1 μ g/ml, Promega) at 37° C. for 3 days. The resulting peptides were desalted using an Oasis HLB μ Elution plate (Waters), and dried and dissolved in 50 μ l of 0.1% FA solution. The sample solution was analyzed by liquid chromatography/mass spectrometry (LC/MS) using the parallel acquisition mode using higher-energy collisional dissociation (HCD) and electron-transfer/higher-energy collisional dissociation (EThcD). The EThcD-MS/MS condition was as follows: 23% of supplemental activation collision energy, m/z 2 of isolation window and 100 ms of maximum injection time. The peptide identifications were performed by database searching using BioPharma Finder™

Nanoparticle Coating

[0084] Streptavidin-coated 0.11- μ m nanoparticles (Bangs Laboratories) were washed twice with PBS and 5.5 μ g nanoparticles were incubated with 50 μ g biotinylated proteins in 80 μ l PBS per one mouse for 5 hours at 4° C. Coating efficiency was measured by flow cytometry.

Immunization

[0085] At 7 to 8 weeks of age, each antigen group was vaccinated with a prime immunization, and three weeks later mice were boosted with a second vaccination. Prior to inoculation, immunogen suspensions were gently mixed 2:1 vol/vol with AddaVax adjuvant (InvivoGen) to reach a final concentration of 0.4 mg/mL antigen. Mice were injected intramuscularly using a 29G $\times$ 1/2 needle syringe (Terumo) with 60 μ L per injection site (120 μ L total) of immunogen under isoflurane anesthesia. For prime immunization antigens were used as nanoparticles and for boost immunization, antigens were used as monomeric proteins.

Flow Cytometry

[0086] Single-cell suspensions were prepared from inguinal and iliac lymph nodes or spleen. Inguinal and iliac lymph nodes were collected and pooled for individual mice. Detection of CoV-1 or CoV-2 RBD-specific B cells was carried out using biotinylated rRBD pre-labeled with fluorophore-conjugated streptavidin. To exclude induced 6 $\times$ His-, Avi-Tag-, biotin- and streptavidin-specific B cells, samples were pre-stained with decoy probe. Cell samples were analyzed using an Attune NxT flow cytometer (Thermo Fisher Scientific) and sorted using a FACS Aria II cell sorter (BD Bioscience). Data were analyzed using FlowJo software V10 (Tree Star).

BCR Cloning and Antibody Expression

[0087] CoV-1 and CoV-2-S RBD-specific B cells were single-cell sorted from a lymph node of immunized mice at day 7 after boost as described above. Cloning and expression of antibodies were performed according to a known method (von Boehmer et al., 2016, Sequencing and cloning of antigen-specific antibodies from mouse memory B cells. *Nat Protoc* 11, 1908-1923; Inoue et al., 2020, Exit from germinal center to become quiescent memory B cells depends on

metabolic reprogramming and provision of a survival signal. *J Exp Med* 218). Monoclonal antibodies were expressed using the Expi293 Expression System (Thermo Fisher Scientific) and purified from the culture supernatants of Expi293F cells by Protein G Sepharose (GE).

Flow Cytometry Analysis of mAb Binding to the Antigen

[0088] Anti-mouse IgK microparticles and negative control particles (BD CompBeads) were mixed at 1:1 ratio. The beads were then mixed with 250 ng of purified mAbs (mIgG2c/mIgK) for 20 min, washed with PBS containing 2% FBS, followed by labeling with PE-conjugated antigen for 20 min. Binding capacity of mAbs to antigen was assessed by flow cytometry (BD FACSCanto II and FACS Aria II).

ELISA

[0089] Nunc Maxisorp Immuno plates (Thermo Fisher Scientific) were coated with each rRBD (100 ng/50 μ l). Blocking was carried out using BlockingOne solution (Nacalai Tesque). Plates were sequentially incubated with serially diluted serum samples or monoclonal antibodies cloned from single GC B cells and IgG1-HRP detection antibodies (Southern Biotech). Detection was carried out using KPL SureBlue TMB Microwell Peroxidase Substrate (SeraCare). The CR3022 mouse IgG1 monoclonal antibody (InvivoGen) was used as a standard.

[0090] For affinity ELISAs, serially diluted sera were incubated on plates coated with 1 μ g/ml CoV-1 RBD protein (low density) or 8 μ g/ml CoV-1 RBD (high density). The affinity of CoV-1 RBD-specific IgG1 was expressed as a ratio of binding to low-density:high-density CoV-1 RBD-coated plates as in the disclosure of Wang et al., 2015 (Anti-HA Glycoforms Drive B Cell Affinity Selection and Determine Influenza Vaccine Efficacy. *Cell* 162, 160-169).

[0091] For mAb-based epitope blocking ELISA, CoV-1 RBD or CoV-2 RBD coated plates for serum samples or cloned monoclonal antibodies respectively were prepared. After blocking was carried out, CR3022 (class 4), EY6A (class 4), and/or S309 (class 3) human IgG1 mAbs (150 ng 50 μ l-1 each) in a mixture or alone was added. After incubation at RT for 3 h, the plate was washed with PBST, then serum samples or cloned monoclonal antibodies were added and incubated at 37° C. for 1 h. After three washes, HRP conjugated goat anti-mouse IgG1 or IgG2c, respectively, were added and incubated for 1 h at RT. Occupancy of Class 3/4 type antibodies in whole anti-CoV-1 RBD antibodies was calculated by subtraction of titers measured by epitope blocking ELISA and conventional serum titer ELISA and then showed as a percentage.

ELISPOT Assay

[0092] Plates with a cellulose membrane bottom were coated with CoV-1 RBD or CoV-2 RBD (100 ng 50 μ l-1). Bone marrow cells were added to the wells and incubated in RPMI1640 medium supplemented with 10% FBS, 50 μ M 2-ME and 2 mM sodium pyruvate for 5 h at 37° C. under 5% CO₂. After washing with PBS with 0.05% Tween 20, goat anti-mouse IgG1 Ab was added to the wells followed by addition of alkaline phosphatase-labeled anti-goat IgG Ab. Spots were visualized by BCIP/NBT substrate (Promega) and counted.

Pseudovirus Assay

[0093] Preparation of SARS CoV S protein-pseudotyped VSVΔG-luc was performed according to the disclosures of Tani et al., 2010 (Involvement of ceramide in the propagation of Japanese encephalitis virus. *J Virol* 84, 2798-2807) and Yoshida et al., 2021 (SARS-CoV-2-induced humoral immunity through B cell epitope analysis in COVID-19 infected individuals. *Sci Rep* 11, 5934). HEK293T cells were transfected with expression plasmids for respective CoV S proteins (SARS-CoV-2 Wuhan strain S, SARS-CoV-1 S, and SARS-related CoV WIV1 S) by using TransIT-LT1 (Mirus) according to the manufacturer's instructions. At 24 hours after transfection, VSVΔG-luc virus (MOI=0.1) was inoculated onto the transfectants. After 2-hour incubation, cells were washed with DMEM and were further cultivated for additional 24-48 hours. Cell-free supernatant was harvested and used for the neutralizing assay according to the disclosure of Nie et al., 2020 (Quantification of SARS-CoV-2 neutralizing antibody by a pseudotyped virus-based assay. *Nat Protoc* 15, 3699-3715). Mouse sera and human plasma samples were incubated at 56° C. for 30 min and then serially diluted from 1/20 in culture medium. SARS-CoV S-pseudotyped VSVΔG-luc was incubated with different dilution of mouse sera, human plasma or recombinant antibodies for 1 hour at 4° C., and then inoculated onto a monolayer culture of VeroE6-TM-PRSS2 in a 96-well plate. At 16-hour post inoculation, cells were washed with PBS and then lysed with luciferase cell culture lysis reagent (Promega). After centrifugation, cleared cell lysates were incubated with firefly luciferase assay substrate (Promega) in 96-well white polystyrene plates (Corning). Luciferase activity was measured by Glo-Max Discover luminometer (Promega). NT50 or IC₅₀ was calculated by Prism software (GraphPad).

Antibody-Dependent Enhancement (ADE) Assay

[0094] DENV1 single-round infectious particles (SRIP) were prepared according to the disclosure in Matsuda et al., 2018 (High-throughput neutralization assay for multiple flaviviruses based on single-round infectious particles using dengue virus type 1 reporter replicon. *Sci Rep* 8, 16624). For evaluation of the ADE assay system using RAW264.7 cells, anti-E mouse monoclonal antibody 3H12 (Yamanaka et al., 2013, A mouse monoclonal antibody against dengue virus type 1 Mochizuki strain targeting envelope protein domain II and displaying strongly neutralizing but not enhancing activity. *J Virol* 87, 12828-12837) was pre-incubated with DENV1 SRIP (multiplicity of infection [MOI]=0.1) for 1 h at 4° C., and then inoculated onto RAW264.7 cells in 96-well plate. After 2 hours, cells were washed with PBS and cultivated in fresh medium for 3 days. Infectivity was measured using a NanoLuc assay kit (Promega). To examine whether antisera had ADE of CoV-2 infection, SARS-CoV-2 S pseudotyped VSVΔG-luc was incubated with sera from immunized mice for 1 h at 4° C., and then inoculated onto RAW264.7 cells in a 96-well plate. After 2 hours, cells were washed with PBS and cultivated in fresh medium for 16-20 hours. Luciferase activity of infected cells was measured as above.

Virus Neutralization Assay

[0095] A mixture of 100 TCID₅₀ virus and serially diluted, heat-inactivated plasma samples (2-fold serial dilu-

tions starting from 1:40 dilution) were incubated at 37° C. for 1 h before being placed on VeroE6-TMPRSS2 cells seeded in 96-well flat-bottom plates (TPP). VeroE6-TMPRSS2 cells were maintained in low glucose DMEM (Fujifilm) containing 10% heat-inactivated fetal bovine serum, 1 mg/mL Geneticin (Thermo Fisher Scientific), and 100 U/mL penicillin/streptomycin (Thermo Fisher Scientific) at 37° C. supplied with 5% CO₂. After culturing for 4 days, cells were fixed with 20% formalin (Fujifilm), and stained with crystal violet solution (Sigma-Aldrich). Cut-off dilution index with >50% cytopathic effect was presented as microneutralization titer. Microneutralization titer of the sample below the detection limit (1:40 dilution) was set as 20.

Biolayer Interferometry Assay

[0096] The kinetics of monoclonal antibody binding to antigen was determined with the OctetRED96e system (ForteBio) at 30° C. with shaking at 1,000 rpm. Biotinylated-RBD proteins from CoV-1, CoV-2, WIV1-CoV, Bat SARS-related AL-103 and BM48-31 were loaded at 6 µg/ml in 1× kinetics buffer (0.1% BSA, 0.02% Tween-20 in PBS) for 900 s onto streptavidin biosensors (ForteBio) and incubated with serially diluted Fab antibodies (100, 33.3, 11.1, and 0 nM) for 120 s, followed by immersion in 1× kinetics buffer for 300 s of dissociation time. The binding curves were fit in a 1:1 binding model and the dissociation constant (K_D) values were calculated by Octet Data Analysis software (ForteBio).

Phylogenetic Analysis of Antibody Clones

[0097] For each cell, V, D, J gene and CDR3 assignment was conducted on the full length BCR sequences using IgBlast and the IMGT mouse references. Clones were defined separately for each experiment, on the basis of their heavy chain only, using the DefineClones function of the Change-O package, and 38/38 cells of GM14 #1 and 26/31 cells of GM14 #2 were considered the same clone. Clonal tree reconstruction was performed for these two major clones using Alakazam (both Change-O and Alakazam are part of the Immcantation analysis framework (Gupta et al., 2015, Change-O: a toolkit for analyzing large-scale B cell immunoglobulin repertoire sequencing data. *Bioinformatics* 31, 3356-3358)). Reconstructed clonal trees were finally plotted using Cytoscape.

Statistics

[0098] Statistical analyses were performed by a two-tailed unpaired and paired Student's t-test using GraphPad Prism software. P values <0.05 were considered significant. Error bars denote ±SD.

Design of SARS-CoV-2 RBD Glycan Mutants

[0099] To engineer the SARS-CoV-2 head-RBD subdomain (FIG. 1A), N-linked glycosylation sites (N_xT motif) were introduced in this subdomain (Barnes et al., 2020, Structures of Human Antibodies Bound to SARS-CoV-2 Spike Reveal Common Epitopes and Recurrent Features of Antibodies. *Cell* 182, 828-842 e816), which contains essential residues for ACE-2 engagement (FIG. 1A and FIG. 1C). The already established monoclonal antibodies (mAbs) recognizing SARS-CoV-2 RBD are classified into four types; as shown in FIG. 1B, class 1/2 and 3/4 recognize epitopes in the head-RBD and core-RBD subdomain, respectively. Five

potential sites for the introduction of N_xT sequons (GM14) were identified based on the following criteria; (1) surface residues on the class 1 or 2 epitope; (2) residues outside the class 3 and 4 epitopes in the core-RBD subdomain (Barnes et al., 2020, the same as above) (FIG. 1C); (3) residues within, or conformationally close to the non-conserved patches of the head-RBD subdomain (A in FIG. 2); (4) N_xT mutations not expected to disrupt the overall structure of RBD; (5) not expected to reduce RBD protein expression or stability (Starr et al., 2020, Deep Mutational Scanning of SARS-CoV-2 Receptor Binding Domain Reveals Constraints on Folding and ACE2 Binding. *Cell* 182, 1295-1310 e1220). Although the N415 site is somewhat distal from the head-RBD subdomain (FIG. 1C), this site was already reported to be a part of the epitope recognized by the class 1 mAb CB6 (Shi et al., 2020, A human neutralizing antibody targets the receptor-binding site of SARS-CoV-2. *Nature* 584, 120-124). Two glycan mutants GM9 and GM14 that have 3 and 5 additional glycosylation sites, respectively, were designed. Structural modeling predicted that these additional glycans would prevent antigen recognition by class 1 and 2 antibodies (Barnes et al., 2020, the same as above; Yuan et al., 2020, the same as above).

Results

[0100] The glycan mutants were coupled with 6×His-Avi-tag at the C terminus and expressed in mammalian Expi293 cells. They exhibited a higher molecular weight than wild-type RBD in SDS-PAGE (FIG. 2B). The glycan occupancy of each introduced glycosylation sites was determined by liquid chromatography-mass spectrometry (LC-MS). All added glycosylation sites had occupancies greater than 75% except the added N494 and N502 glycan from GM9, which showed 70% and 57%, respectively (FIG. 1D). The process of N-glycosylation in the endoplasmic reticulum (ER) is affected by conformational constraints (Breitling and Aebi, 2013, N-linked protein glycosylation in the endoplasmic reticulum. *Cold Spring Harb Perspect Biol* 5, a013359). The somewhat lower glycan occupancy at N494 and N502 could be due to limited accessibility of oligosaccharyl transferases to these asparagine residues, which are positioned in close proximity to each other.

[0101] It was then investigated whether the glycan modifications prevent binding of class 1/2 antibodies but maintain antigenic epitopes for class 3/4 antibodies. In ELISA, all tested antibodies were strong binders to wildtype RBD (FIG. 1E). mAb CB6, a stereotypic class 1 neutralizing antibody, completely failed to bind to either of the glycan mutants (GM9 and GM14). Although the binding of mAb C002, a class 2 antibody (Robbiani et al., 2020, Convergent antibody responses to SARS-CoV-2 in convalescent individuals. *Nature* 584, 437-442), to RBD was abolished by GM14, this mAb still bound weakly to GM9, indicating that the five mutations are required for complete inhibition of class 2 mAb C002. In contrast, core-RBD subdomain targeting mAb S309 (class 3), mAb CR3022 (class 4), and mAb EY6A (class 4) (Pinto et al., 2020, the same as above; Yuan et al., 2020, the same as above) (Zhou et al., 2020, the same as above) bound to wildtype RBD, GM9, or GM14 at a similar level, indicating that the introduced mutations for additional glycosylation do not affect overall structure of the core-RBD subdomain (FIG. 1E).

Glycan Engineering of the SARS-CoV-2 Head-RBD Generated More Antibodies Towards the Core-RBD with Higher Affinity

[0102] To overcome the limited immunogenicity of the relatively small SARS-CoV-2 RBD, it was multivalently displayed on streptavidin polystyrene nanoparticles. Consistent with the disclosure in Walls et al., 2020 (Elicitation of Potent Neutralizing Antibody Responses by Designed Protein Nanoparticle Vaccines for SARS-CoV-2. *Cell* 183, 1367-1382 e1317), in contrast to soluble monomeric RBD, the particulate RBD gave rise to more robust primary antibody responses (FIG. 4). These particulate antigens (wildtype RBD, GM9, or GM14) were injected intramuscularly into BALB/c mice with AddaVax adjuvant. The mice were boosted with respective non-particulate antigens 3 weeks after priming (FIG. 3A). Sera taken at 7 days after boosting were tested for reactivity to SARS-CoV-2 RBD. Since, like alum, AddaVax dominantly induces IgG1 type antibody in the mouse (Sato et al., 2020, Repurposing the psoriasis drug Oxarol to an ointment adjuvant for the influenza vaccine. *Int Immunol* 32, 499-507), the IgG1 response was measured. To first address the possibility that the mice may have mounted an immune response towards neo-epitopes generated by glycan engineering that the parental SARS-CoV-2 RBD probe would not detect, serum and GC B cell responses were examined from the respective immunization protocols using immunogen-matched glycosylation mutant and parental probes. ELISA assay analysis indicated similar RBD-specific serum titers between the probe sets; by flow cytometry analysis, about 10 or 15% of GC B cells bound to only immunogen-matched glycosylation mutant probes (FIGS. 6A and B). Hence, to some extent, B cells recognizing neo-epitopes appear to emerge upon GM9 or GM14 immunization. Nevertheless, more than 85% of the responding cells were reactive to the parental CoV-2 RBD probe, thus, the parental SARS-CoV-2 RBD probe were usually used in these assays.

[0103] The IgG1 from mice immunized with glycan mutants displayed similar reactivity to CoV-2, assessed by ELISA, to that by wildtype RBD immunization (FIG. 3B, left). In regard to the reactivity to CoV-1, GM9- or GM14-immunized mice exhibited about 8-fold higher than wildtype RBD immunization (FIG. 3B, right). Given the sequence conservation of the RBD-core subdomain between SARS-CoV-2 and CoV-1 (FIG. 2A), it is most likely that the enhanced cross-reactivity to CoV-1 by GM9 or GM14 immunization is due to generation of more IgG1 towards the conserved RBD-core subdomain of CoV-2. Supporting this possibility, CoV-1 reactive IgG1 elicited by GM9 or GM14 immunization was about 90 or 85% inhibited by pre-treatment of verified class 3/4 anti-CoV-2 core-RBD antibodies (mixtures of mAb S309, mAb CR3022, and mAb EY6A), respectively, like wildtype RBD immunization (FIG. 3C, left). Next, the affinity of cross-reactive IgG1 elicited by GM9 or GM14 immunization was characterized. The affinity of IgG1 for CoV-1 RBD was measured using an affinity ELISA assay that was modified from the nitrophenyl system; this assay measures the ratio of high-affinity to all-affinity binding IgG1. As shown in FIG. 3C, right, in contrast to wildtype RBD immunization, cross-reactive IgG1 elicited by GM9 or GM14 immunization had significantly higher affinity. When the WIV1-CoV was used instead of the SARS-CoV-1 RBD probe, both glycan mutants induced about 8-fold higher serum titers (FIG. 3D, left). In regard to

affinity, although two among 11 wildtype RBD-immunized mice showed high ratio of high-affinity to all-affinity binding, sera from the remaining nine mice had relatively lower affinity compared with GM9- or GM14-immunized mice (FIG. 3D, right). Together, glycan engineering of the SARS-CoV-2 head-RBD subdomain facilitates the generation of core-RBD subdomain specific antibodies with higher affinity.

Glycan Engineering of the SARS-CoV-2 Head-RBD Elicited Cross-Neutralizing Antibodies Against SARS-CoV-1 and Other Related Viruses

[0104] To determine neutralizing activity, a vesicular stomatitis virus (VSV)-based pseudovirus method (Nie et al., 2020, the same as above) was mainly employed, and half-maximal neutralization titers (NT50s) were calculated. CoV-2 wildtype RBD, GM9, or GM14 immunization gave rise to similar neutralizing activity to SARS-CoV-2. GM9 or GM14 immunization manifested somewhat increased or decreased activity, respectively, compared with CoV-2 wildtype RBD immunization (FIG. 5A). By using virus neutralizing assay, this tendency by GM9 or GM14 immunization was also observed (FIG. 8).

[0105] To measure the neutralizing activity against CoV-2 variants, VSV-based viruses with K417N/E484K/N501Y mutation from the B.1.351 variant were used. The results indicated that even CoV-2 wildtype RBD immunization elicited somewhat higher neutralizing activity than wildtype CoV-2. Given that human sera by mRNA vaccines showed low neutralizing activity against viruses with the E484K mutation (Chen et al., 2021, Resistance of SARS-CoV-2 variants to neutralization by monoclonal and serum-derived polyclonal antibodies. *Nat. Med.* 27:717-726), the immunization technique of the present invention can produce more resistant sera.

[0106] Towards SARS-CoV-1, antisera elicited by GM9 or GM14 immunization possessed neutralizing activity with about 15- or 10-fold higher potency, respectively, than CoV-2 wildtype RBD immunization, assessed by the VSV-based pseudovirus method (FIG. 5A). Similarly, these sera exhibited substantial neutralizing activity against WIV1-CoV, SHC014, and PaGX (FIG. 5A). Importantly, the levels of neutralization for SARS-CoV-1 and WIV1-CoV observed after GM9 or GM14 immunization were similar to convalescent plasma samples from SARS-CoV-2-infected individuals for their neutralizing activity of the SARS-CoV-2, analyzed by the same laboratory in the same assay format (FIG. 5B and FIG. 5C). Given that the major epitopes of the CoV-1 reactive antibodies elicited by GM9 or GM14 immunization are located in the core-RBD subdomain (FIG. 3C, left), it is most likely that these anti-core-RBD antibodies contribute to cross-neutralization of SARS-CoV-1, WIV1-CoV, SHC014, and PaGX.

[0107] Previous studies of SARS-CoV-1 and other viruses have suggested that antibody-dependent enhancement (ADE) of infection might take place after vaccination (Smatti et al., 2018, Viral-Induced Enhanced Disease Illness. *Front Microbiol* 9, 2991). To examine this possibility, VSV-based pseudovirus was incubated with serially diluted sera from immunized mice and then inoculated on Raji, a human B lymphoma cell line frequently used to evaluate the ADE activity of SARS-CoV (Verma et al., 2009, Analysis of the Fc gamma receptor-dependent component of neutralization measured by anthrax toxin neutralization assays. *Clin Vac*

cine Immunol 16, 1405-1412). As a positive control for this assay, it was shown that infectivity of CoV-2 S pseudovirus was significantly enhanced by addition of anti-CoV-2 SRBD MW05 chimeric human/mouse IgG1 antibody (FIG. 5D, center). In this experimental setting, like sera from mice immunized with SARS-CoV-2 wildtype RBD, sera from mice immunized with GM9 or GM14 showed no significant ADE activity (FIG. 5D, right).

Reactivity Profiles of Cross-Reactive GC B Cells

[0108] To complement the above serological characterization, the reactivity of GC B cells isolated post priming was assessed. As demonstrated by flow analysis, wildtype RBD, GM9, or GM14 immunization gave rise to similar levels of overall GC B cells (FIG. 7A). Among the CoV-2+ cells, the frequency of GC B cells positive for both CoV-2 and CoV-1 probes were increased by immunization of GM9 or GM14, compared with wildtype RBD immunization (FIGS. 7B and C).

[0109] To examine cross-reactivity profiles for each CoV-2+/CoV-1+GC B cells, mAbs from single cell sorted GC B cells from four individual mouse (GM9-1, GM9-2, GM14-1, or GM14-2) were characterized. Sequence analysis revealed that, in each mouse, more than 85% of the isolated mAbs were members of expanded clonal lineages. Particularly in GM14-1 or GM14-2 mouse, almost all the mAbs were encoded by the same combination of VH/VK with different DH and JK segments: GM14-1 (VH14-3-DH3-2-JH4/VK4-111-JK2), and GM14-2 (VH14-3-DH2-3-JH4/VK4-111-JK1) (FIG. 12).

[0110] Among all the analyzed mAbs, many of them showed blockade of their binding to CoV-2 RBD by authentic class 4 mAbs, except that mAbs derived from VH14-1/VK8-27 gene pairing were blocked by authentic class 3 mAb (FIG. 12). For assessment of the breadth of recognition to sarbecovirus RBDs by the above mAbs, a relative binding index was assigned to each antibody by employing flow cytometry-based measurement (FIG. 9A) (Bajic et al., 2019, Influenza Antigen Engineering Focuses Immune Responses to a Subdominant but Broadly Protective Viral Epitope. Cell Host Microbe 25, 827-835 e826; Leach et al., 2019, Requirement for memory B-cell activation in protection from heterologous influenza virus reinfection. Int Immunol 31, 771-779); the relative binding indexes represented well RMax of bivalent IgG antibodies measured by Octet kinetics assay (FIG. 10). RBDs were selected from distantly related clade 2 (Anlong-103 [AL-103]) and clade 3 (BM48-31) (Lau et al., 2020, Possible Bat Origin of Severe Acute Respiratory Syndrome Coronavirus 2. Emerg Infect Dis 26,

1542-1547) (Letko et al., 2020, Functional assessment of cell entry and receptor usage for SARS-CoV-2 and other lineage B betacoronaviruses. Nat Microbiol 5, 562-569), in addition to clade 1 (SARS-CoV-1 and WIV1-CoV as clade 1a, and SARS-CoV-2 and PaGX as clade 1b). As expected, almost all the isolated mAbs from GM9-1, GM9-2, GM14-1, or GM14-2 mouse displayed binding reactivity to clade 1a and clade 1b RBDs. The relative binding indexes were well correlated between CoV-1/WIV1/SHC014/PaGX and CoV-2 in GM9- or GM14-immunized mice (FIG. 9B). In regard to clade 2 and 3RBD, some of the isolated mAbs from GM9-1 or GM9-2, but only a few from GM14-1 mouse, exhibited their significant binding reactivity (FIG. 9A).

[0111] Then, the neutralization activities of each mAb towards CoV-1, WIV1, SHC014, PaGX, or CoV-2 were evaluated by the aforementioned VSV-based pseudotype assays (FIG. 9A and FIG. 12). Because of the large number of antibodies, screening was performed and nine mAbs were focused using the following two criteria (FIG. 9C). First, the neutralization activities are below 10 µg/ml IC₅₀ (half-maximal inhibitory concentration) towards at least two types of viruses among three. The highest potency towards these viruses is below 3 µg/ml IC₅₀. As shown in FIG. 9D, like authentic mAb S309 or CR3022, these mAbs possessed higher IC₅₀s for CoV-2 than those observed for CoV-1 or WIV1. In VH14-3-DH3-2-JH4/VK4-111-JK2 (GM14-1) or VH14-3-DH2-3-JH4/VK4-111-JK1 (GM14-2) lineage, it was demonstrated that the mAb with the highest binding index for CoV-2 does not possess the highest potency of neutralization activity towards CoV-2 (FIG. 9C). Thus, antibody neutralization potency is likely governed at least in part by factors beyond binding affinity.

Elicitation of Cross-Reactive Long-Lived Plasma Cells and Memory B Cells

[0112] Most successful vaccine approaches rely on the generation of memory B cells and long-lived plasma cells (LLPCs). As shown in FIG. 11A, analysis was carried out at seven weeks. At this time, class-switched IgG1 type memory B cells were similarly generated by wildtype RBD, GM9, or GM14 immunization, although GM9 and GM14 gave rise to more CoV-1+CoV-2+ cells (FIG. 11B). Furthermore, as assessed by SARS-CoV-1-specific antibody secreting cells (ASC) in the bone marrow, GM9 or GM14 immunization induced more CoV-1+ ASC (FIG. 11C), together indicating that both cross-reactive memory compartments are more efficiently generated by this immunization protocol using GM9 or GM14, compared with wildtype RBD immunization.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 114

<210> SEQ ID NO 1

<211> LENGTH: 1273

<212> TYPE: PRT

<213> ORGANISM: SARS-CoV-2

<400> SEQUENCE: 1

Met Phe Val Phe Leu Val Leu Pro Leu Val Ser Ser Gln Cys Val
1 5 10 15

-continued

Asn	Leu	Thr	Thr	Arg	Thr	Gln	Leu	Pro	Pro	Ala	Tyr	Thr	Asn	Ser	Phe
			20					25					30		
Thr	Arg	Gly	Val	Tyr	Tyr	Pro	Asp	Lys	Val	Phe	Arg	Ser	Ser	Val	Leu
		35					40					45			
His	Ser	Thr	Gln	Asp	Leu	Phe	Leu	Pro	Phe	Phe	Ser	Asn	Val	Thr	Trp
	50					55					60				
Phe	His	Ala	Ile	His	Val	Ser	Gly	Thr	Asn	Gly	Thr	Lys	Arg	Phe	Asp
65					70					75					80
Asn	Pro	Val	Leu	Pro	Phe	Asn	Asp	Gly	Val	Tyr	Phe	Ala	Ser	Thr	Glu
				85					90					95	
Lys	Ser	Asn	Ile	Ile	Arg	Gly	Trp	Ile	Phe	Gly	Thr	Thr	Leu	Asp	Ser
			100					105					110		
Lys	Thr	Gln	Ser	Leu	Leu	Ile	Val	Asn	Asn	Ala	Thr	Asn	Val	Val	Ile
		115					120					125			
Lys	Val	Cys	Glu	Phe	Gln	Phe	Cys	Asn	Asp	Pro	Phe	Leu	Gly	Val	Tyr
	130					135					140				
Tyr	His	Lys	Asn	Asn	Lys	Ser	Trp	Met	Glu	Ser	Glu	Phe	Arg	Val	Tyr
145					150					155					160
Ser	Ser	Ala	Asn	Asn	Cys	Thr	Phe	Glu	Tyr	Val	Ser	Gln	Pro	Phe	Leu
				165					170					175	
Met	Asp	Leu	Glu	Gly	Lys	Gln	Gly	Asn	Phe	Lys	Asn	Leu	Arg	Glu	Phe
			180					185					190		
Val	Phe	Lys	Asn	Ile	Asp	Gly	Tyr	Phe	Lys	Ile	Tyr	Ser	Lys	His	Thr
		195					200					205			
Pro	Ile	Asn	Leu	Val	Arg	Asp	Leu	Pro	Gln	Gly	Phe	Ser	Ala	Leu	Glu
	210					215					220				
Pro	Leu	Val	Asp	Leu	Pro	Ile	Gly	Ile	Asn	Ile	Thr	Arg	Phe	Gln	Thr
225					230					235					240
Leu	Leu	Ala	Leu	His	Arg	Ser	Tyr	Leu	Thr	Pro	Gly	Asp	Ser	Ser	Ser
				245					250					255	
Gly	Trp	Thr	Ala	Gly	Ala	Ala	Ala	Tyr	Tyr	Val	Gly	Tyr	Leu	Gln	Pro
			260					265					270		
Arg	Thr	Phe	Leu	Leu	Lys	Tyr	Asn	Glu	Asn	Gly	Thr	Ile	Thr	Asp	Ala
		275					280					285			
Val	Asp	Cys	Ala	Leu	Asp	Pro	Leu	Ser	Glu	Thr	Lys	Cys	Thr	Leu	Lys
	290					295					300				
Ser	Phe	Thr	Val	Glu	Lys	Gly	Ile	Tyr	Gln	Thr	Ser	Asn	Phe	Arg	Val
305					310					315					320
Gln	Pro	Thr	Glu	Ser	Ile	Val	Arg	Phe	Pro	Asn	Ile	Thr	Asn	Leu	Cys
				325					330					335	
Pro	Phe	Gly	Glu	Val	Phe	Asn	Ala	Thr	Arg	Phe	Ala	Ser	Val	Tyr	Ala
		340						345					350		
Trp	Asn	Arg	Lys	Arg	Ile	Ser	Asn	Cys	Val	Ala	Asp	Tyr	Ser	Val	Leu
		355					360					365			
Tyr	Asn	Ser	Ala	Ser	Phe	Ser	Thr	Phe	Lys	Cys	Tyr	Gly	Val	Ser	Pro
	370					375					380				
Thr	Lys	Leu	Asn	Asp	Leu	Cys	Phe	Thr	Asn	Val	Tyr	Ala	Asp	Ser	Phe
385					390					395					400
Val	Ile	Arg	Gly	Asp	Glu	Val	Arg	Gln	Ile	Ala	Pro	Gly	Gln	Thr	Gly
			405					410					415		
Lys	Ile	Ala	Asp	Tyr	Asn	Tyr	Lys	Leu	Pro	Asp	Asp	Phe	Thr	Gly	Cys

420																425								430							
Val	Ile	Ala	Trp	Asn	Ser	Asn	Asn	Leu	Asp	Ser	Lys	Val	Gly	Gly	Asn																
435				440				445																							
Tyr	Asn	Tyr	Leu	Tyr	Arg	Leu	Phe	Arg	Lys	Ser	Asn	Leu	Lys	Pro	Phe																
450				455				460																							
Glu	Arg	Asp	Ile	Ser	Thr	Glu	Ile	Tyr	Gln	Ala	Gly	Ser	Thr	Pro	Cys																
465				470				475				480																			
Asn	Gly	Val	Glu	Gly	Phe	Asn	Cys	Tyr	Phe	Pro	Leu	Gln	Ser	Tyr	Gly																
485				490				495																							
Phe	Gln	Pro	Thr	Asn	Gly	Val	Gly	Tyr	Gln	Pro	Tyr	Arg	Val	Val	Val																
500				505				510																							
Leu	Ser	Phe	Glu	Leu	Leu	His	Ala	Pro	Ala	Thr	Val	Cys	Gly	Pro	Lys																
515				520				525																							
Lys	Ser	Thr	Asn	Leu	Val	Lys	Asn	Lys	Cys	Val	Asn	Phe	Asn	Phe	Asn																
530				535				540																							
Gly	Leu	Thr	Gly	Thr	Gly	Val	Leu	Thr	Glu	Ser	Asn	Lys	Lys	Phe	Leu																
545				550				555				560																			
Pro	Phe	Gln	Gln	Phe	Gly	Arg	Asp	Ile	Ala	Asp	Thr	Thr	Asp	Ala	Val																
565				570				575																							
Arg	Asp	Pro	Gln	Thr	Leu	Glu	Ile	Leu	Asp	Ile	Thr	Pro	Cys	Ser	Phe																
580				585				590																							
Gly	Gly	Val	Ser	Val	Ile	Thr	Pro	Gly	Thr	Asn	Thr	Ser	Asn	Gln	Val																
595				600				605																							
Ala	Val	Leu	Tyr	Gln	Asp	Val	Asn	Cys	Thr	Glu	Val	Pro	Val	Ala	Ile																
610				615				620																							
His	Ala	Asp	Gln	Leu	Thr	Pro	Thr	Trp	Arg	Val	Tyr	Ser	Thr	Gly	Ser																
625				630				635				640																			
Asn	Val	Phe	Gln	Thr	Arg	Ala	Gly	Cys	Leu	Ile	Gly	Ala	Glu	His	Val																
645				650				655																							
Asn	Asn	Ser	Tyr	Glu	Cys	Asp	Ile	Pro	Ile	Gly	Ala	Gly	Ile	Cys	Ala																
660				665				670																							
Ser	Tyr	Gln	Thr	Gln	Thr	Asn	Ser	Pro	Arg	Arg	Ala	Arg	Ser	Val	Ala																
675				680				685																							
Ser	Gln	Ser	Ile	Ile	Ala	Tyr	Thr	Met	Ser	Leu	Gly	Ala	Glu	Asn	Ser																
690				695				700																							
Val	Ala	Tyr	Ser	Asn	Asn	Ser	Ile	Ala	Ile	Pro	Thr	Asn	Phe	Thr	Ile																
705				710				715				720																			
Ser	Val	Thr	Thr	Glu	Ile	Leu	Pro	Val	Ser	Met	Thr	Lys	Thr	Ser	Val																
725				730				735																							
Asp	Cys	Thr	Met	Tyr	Ile	Cys	Gly	Asp	Ser	Thr	Glu	Cys	Ser	Asn	Leu																
740				745				750																							
Leu	Leu	Gln	Tyr	Gly	Ser	Phe	Cys	Thr	Gln	Leu	Asn	Arg	Ala	Leu	Thr																
755				760				765																							
Gly	Ile	Ala	Val	Glu	Gln	Asp	Lys	Asn	Thr	Gln	Glu	Val	Phe	Ala	Gln																
770				775				780																							
Val	Lys	Gln	Ile	Tyr	Lys	Thr	Pro	Pro	Ile	Lys	Asp	Phe	Gly	Gly	Phe																
785				790				795				800																			
Asn	Phe	Ser	Gln	Ile	Leu	Pro	Asp	Pro	Ser	Lys	Pro	Ser	Lys	Arg	Ser																
805				810				815																							
Phe	Ile	Glu	Asp	Leu	Leu	Phe	Asn	Lys	Val	Thr	Leu	Ala	Asp	Ala	Gly																
820				825				830																							

Phe	Ile	Lys	Gln	Tyr	Gly	Asp	Cys	Leu	Gly	Asp	Ile	Ala	Ala	Arg	Asp
835															
Leu	Ile	Cys	Ala	Gln	Lys	Phe	Asn	Gly	Leu	Thr	Val	Leu	Pro	Pro	Leu
850															
Leu	Thr	Asp	Glu	Met	Ile	Ala	Gln	Tyr	Thr	Ser	Ala	Leu	Leu	Ala	Gly
865															
Thr	Ile	Thr	Ser	Gly	Trp	Thr	Phe	Gly	Ala	Gly	Ala	Ala	Leu	Gln	Ile
885															
Pro	Phe	Ala	Met	Gln	Met	Ala	Tyr	Arg	Phe	Asn	Gly	Ile	Gly	Val	Thr
900															
Gln	Asn	Val	Leu	Tyr	Glu	Asn	Gln	Lys	Leu	Ile	Ala	Asn	Gln	Phe	Asn
915															
Ser	Ala	Ile	Gly	Lys	Ile	Gln	Asp	Ser	Leu	Ser	Ser	Thr	Ala	Ser	Ala
930															
Leu	Gly	Lys	Leu	Gln	Asp	Val	Val	Asn	Gln	Asn	Ala	Gln	Ala	Leu	Asn
945															
Thr	Leu	Val	Lys	Gln	Leu	Ser	Ser	Asn	Phe	Gly	Ala	Ile	Ser	Ser	Val
965															
Leu	Asn	Asp	Ile	Leu	Ser	Arg	Leu	Asp	Lys	Val	Glu	Ala	Glu	Val	Gln
980															
Ile	Asp	Arg	Leu	Ile	Thr	Gly	Arg	Leu	Gln	Ser	Leu	Gln	Thr	Tyr	Val
995															
Thr	Gln	Gln	Leu	Ile	Arg	Ala	Ala	Glu	Ile	Arg	Ala	Ser	Ala	Asn	
1010															
Leu	Ala	Ala	Thr	Lys	Met	Ser	Glu	Cys	Val	Leu	Gly	Gln	Ser	Lys	
1025															
Arg	Val	Asp	Phe	Cys	Gly	Lys	Gly	Tyr	His	Leu	Met	Ser	Phe	Pro	
1040															
Gln	Ser	Ala	Pro	His	Gly	Val	Val	Phe	Leu	His	Val	Thr	Tyr	Val	
1055															
Pro	Ala	Gln	Glu	Lys	Asn	Phe	Thr	Thr	Ala	Pro	Ala	Ile	Cys	His	
1070															
Asp	Gly	Lys	Ala	His	Phe	Pro	Arg	Glu	Gly	Val	Phe	Val	Ser	Asn	
1085															
Gly	Thr	His	Trp	Phe	Val	Thr	Gln	Arg	Asn	Phe	Tyr	Glu	Pro	Gln	
1100															
Ile	Ile	Thr	Thr	Asp	Asn	Thr	Phe	Val	Ser	Gly	Asn	Cys	Asp	Val	
1115															
Val	Ile	Gly	Ile	Val	Asn	Asn	Thr	Val	Tyr	Asp	Pro	Leu	Gln	Pro	
1130															
Glu	Leu	Asp	Ser	Phe	Lys	Glu	Glu	Leu	Asp	Lys	Tyr	Phe	Lys	Asn	
1145															
His	Thr	Ser	Pro	Asp	Val	Asp	Leu	Gly	Asp	Ile	Ser	Gly	Ile	Asn	
1160															
Ala	Ser	Val	Val	Asn	Ile	Gln	Lys	Glu	Ile	Asp	Arg	Leu	Asn	Glu	
1175															
Val	Ala	Lys	Asn	Leu	Asn	Glu	Ser	Leu	Ile	Asp	Leu	Gln	Glu	Leu	
1190															
Gly	Lys	Tyr	Glu	Gln	Tyr	Ile	Lys	Trp	Pro	Trp	Tyr	Ile	Trp	Leu	
1205															

-continued

Gly	Phe	Ile	Ala	Gly	Leu	Ile	Ala	Ile	Val	Met	Val	Thr	Ile	Met
1220						1225					1230			
Leu	Cys	Cys	Met	Thr	Ser	Cys	Cys	Ser	Cys	Leu	Lys	Gly	Cys	Cys
1235						1240					1245			
Ser	Cys	Gly	Ser	Cys	Cys	Lys	Phe	Asp	Glu	Asp	Asp	Ser	Glu	Pro
1250						1255					1260			
Val	Leu	Lys	Gly	Val	Lys	Leu	His	Tyr	Thr					
1265						1270								

<210> SEQ ID NO 2
 <211> LENGTH: 199
 <212> TYPE: PRT
 <213> ORGANISM: SARS-CoV-2

<400> SEQUENCE: 2

Asn	Ile	Thr	Asn	Leu	Cys	Pro	Phe	Gly	Glu	Val	Phe	Asn	Ala	Thr	Arg
1				5					10					15	
Phe	Ala	Ser	Val	Tyr	Ala	Trp	Asn	Arg	Lys	Arg	Ile	Ser	Asn	Cys	Val
			20					25					30		
Ala	Asp	Tyr	Ser	Val	Leu	Tyr	Asn	Ser	Ala	Ser	Phe	Ser	Thr	Phe	Lys
		35					40					45			
Cys	Tyr	Gly	Val	Ser	Pro	Thr	Lys	Leu	Asn	Asp	Leu	Cys	Phe	Thr	Asn
	50					55					60				
Val	Tyr	Ala	Asp	Ser	Phe	Val	Ile	Arg	Gly	Asp	Glu	Val	Arg	Gln	Ile
65					70					75				80	
Ala	Pro	Gly	Gln	Thr	Gly	Lys	Ile	Ala	Asp	Tyr	Asn	Tyr	Lys	Leu	Pro
			85						90					95	
Asp	Asp	Phe	Thr	Gly	Cys	Val	Ile	Ala	Trp	Asn	Ser	Asn	Asn	Leu	Asp
		100						105						110	
Ser	Lys	Val	Gly	Gly	Asn	Tyr	Asn	Tyr	Leu	Tyr	Arg	Leu	Phe	Arg	Lys
		115					120					125			
Ser	Asn	Leu	Lys	Pro	Phe	Glu	Arg	Asp	Ile	Ser	Thr	Glu	Ile	Tyr	Gln
	130					135					140				
Ala	Gly	Ser	Thr	Pro	Cys	Asn	Gly	Val	Glu	Gly	Phe	Asn	Cys	Tyr	Phe
145					150					155					160
Pro	Leu	Gln	Ser	Tyr	Gly	Phe	Gln	Pro	Thr	Asn	Gly	Val	Gly	Tyr	Gln
			165						170					175	
Pro	Tyr	Arg	Val	Val	Val	Leu	Ser	Phe	Glu	Leu	Leu	His	Ala	Pro	Ala
		180						185					190		
Thr	Val	Cys	Gly	Pro	Lys	Lys									
		195													

<210> SEQ ID NO 3
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 3

Val	Arg	Asp	Tyr	Gly	Tyr	Gly	Tyr	Phe	Asp	Val
1				5					10	

<210> SEQ ID NO 4
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

-continued

<400> SEQUENCE: 4

Val Arg Gly Tyr Gly Tyr Gly Tyr Phe Asp Val
1 5 10

<210> SEQ ID NO 5

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 5

Val Arg Asp Tyr Gly Tyr Gly Tyr Phe Asp Asp
1 5 10

<210> SEQ ID NO 6

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 6

Ala Ile Ile Thr Thr Val Leu Pro Trp Phe Ala Tyr
1 5 10

<210> SEQ ID NO 7

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 7

Ala Ile Met Thr Thr Val Leu Pro Trp Phe Ala Tyr
1 5 10

<210> SEQ ID NO 8

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 8

Ala Ile Ile Thr Thr Val Ala Pro Phe Phe Ala Tyr
1 5 10

<210> SEQ ID NO 9

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 9

Ala Ile Ile Thr Thr Val Ala Pro Tyr Phe Ala Tyr
1 5 10

<210> SEQ ID NO 10

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 10

Ala Leu Ile Thr Thr Val Leu Pro Trp Phe Ala Phe
1 5 10

<210> SEQ ID NO 11

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

-continued

<400> SEQUENCE: 11

Ala Leu Ile Thr Thr Val Leu Pro Trp Phe Thr Phe
1 5 10

<210> SEQ ID NO 12

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 12

Thr Arg Glu Ser Asp Tyr Gly Phe Phe Asp Tyr
1 5 10

<210> SEQ ID NO 13

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 13

Ala Arg Glu Ser Asp Tyr Gly Phe Phe Asp Cys
1 5 10

<210> SEQ ID NO 14

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 14

Val Arg Glu Ser Asp Tyr Gly Tyr Phe Ala Tyr
1 5 10

<210> SEQ ID NO 15

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 15

Ala Arg Glu Ser Asp Tyr Gly Phe Phe Asp Tyr
1 5 10

<210> SEQ ID NO 16

<211> LENGTH: 18

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 16

Ala Arg His Gly Gly Thr Tyr Tyr Tyr Gly Ile Ser Pro Tyr Ala Met
1 5 10 15

Asp Tyr

<210> SEQ ID NO 17

<211> LENGTH: 18

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 17

Ala Arg His Ser Gly Thr Tyr Tyr Tyr Gly Ser Ser Pro Tyr Ala Met
1 5 10 15

Asp Tyr

-continued

<210> SEQ ID NO 18
<211> LENGTH: 18
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 18

Ala Arg His Ser Gly Thr Tyr Tyr Tyr Gly Ser Ser Pro Tyr Gly Met
1 5 10 15

Asp Tyr

<210> SEQ ID NO 19
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 19

Ala Arg Gly Trp Leu Gly His Tyr Phe Asp Tyr
1 5 10

<210> SEQ ID NO 20
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 20

Gln His His Tyr Gly Thr Pro Leu Thr
1 5

<210> SEQ ID NO 21
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 21

Gln His His Tyr Gly Asn Pro Leu Thr
1 5

<210> SEQ ID NO 22
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 22

Gln His His Tyr Gly Ser Pro Leu Thr
1 5

<210> SEQ ID NO 23
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 23

Gln His His Tyr Asp Ala Pro Leu Thr
1 5

<210> SEQ ID NO 24
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 24

-continued

Gln His His Tyr Gly Ala Pro Leu Thr
1 5

<210> SEQ ID NO 25
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 25

His Gln Tyr Leu Ser Ser Trp Thr
1 5

<210> SEQ ID NO 26
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 26

Gln His His Tyr Gly Ser Pro Pro Trp Thr
1 5 10

<210> SEQ ID NO 27
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 27

Gln His His Tyr Gly Thr Pro Pro Trp Thr
1 5 10

<210> SEQ ID NO 28
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 28

Gln His Phe Trp Asp Thr Pro Tyr Thr
1 5

<210> SEQ ID NO 29
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 29

Gln Gln His Tyr Ser Thr Pro Arg Thr
1 5

<210> SEQ ID NO 30
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 30

Thr Ile Gly Tyr Gly Tyr Gly Tyr Phe Asp Tyr
1 5 10

<210> SEQ ID NO 31
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 31

-continued

Thr Lys Gly Tyr Gly Tyr Gly Tyr Phe Asp Tyr
1 5 10

<210> SEQ ID NO 32
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 32

Thr Lys Gly Tyr Gly Tyr Gly Tyr Phe Asp Gln
1 5 10

<210> SEQ ID NO 33
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 33

Ser Ile Gly Tyr Gly Tyr Gly Tyr Phe Asp Tyr
1 5 10

<210> SEQ ID NO 34
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 34

Ala Arg Asn Phe Asp Val
1 5

<210> SEQ ID NO 35
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 35

Ala Arg Tyr Phe Asp Val
1 5

<210> SEQ ID NO 36
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 36

Val Arg Asp Tyr Gly Tyr Gly Tyr Phe Gly Val
1 5 10

<210> SEQ ID NO 37
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 37

Thr Ile Gly Phe Gly Tyr Gly Tyr Phe Asp Tyr
1 5 10

<210> SEQ ID NO 38
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

-continued

<400> SEQUENCE: 38

Ala Arg His Tyr Asp Tyr Asp Asp Gly Met Asp Tyr
1 5 10

<210> SEQ ID NO 39

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 39

Ala Arg Tyr Gly Asn Asp Trp Tyr Phe Asp Val
1 5 10

<210> SEQ ID NO 40

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 40

Thr Thr Pro Val Tyr Asp Gly Tyr Tyr Asp Tyr Ala Met Asp Cys
1 5 10 15

<210> SEQ ID NO 41

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 41

Gln His His Tyr Gly Thr Pro Tyr Thr
1 5

<210> SEQ ID NO 42

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 42

Gln His His Tyr Gly Ser Pro Tyr Thr
1 5

<210> SEQ ID NO 43

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 43

Gln Gln Trp Asn Asn Tyr Pro Tyr Thr
1 5

<210> SEQ ID NO 44

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 44

Gln Gln Trp Ser Ser Tyr Pro Tyr Thr
1 5

<210> SEQ ID NO 45

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

-continued

<400> SEQUENCE: 45

Gln His His Tyr Gly Pro Pro Tyr Thr
1 5

<210> SEQ ID NO 46

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 46

Gln His His Tyr Gly Pro Pro Phe Thr
1 5

<210> SEQ ID NO 47

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (9)..(9)

<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

<400> SEQUENCE: 47

Phe Gln Gly Ser His Val Pro Arg Xaa
1 5

<210> SEQ ID NO 48

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 48

Gln Gln Gly Gly Glu Asp Pro Trp Thr
1 5

<210> SEQ ID NO 49

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 49

Ala Arg Leu Asp Ser Ser Tyr Tyr Val Asp Tyr Pro Met Asp Tyr
1 5 10 15

<210> SEQ ID NO 50

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 50

Ala Arg Leu Asp Ser Ser Tyr Tyr Val Asp Tyr Pro Leu Asp Tyr
1 5 10 15

<210> SEQ ID NO 51

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 51

Ala Arg Leu Asp Ser Ser Ala Tyr Val Asp Tyr Pro Met Asp Tyr
1 5 10 15

-continued

<210> SEQ ID NO 52
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 52

Val	Arg	Leu	Asp	Ser	Ser	Ala	Tyr	Val	Gly	Tyr	Pro	Met	Asp	Tyr
1				5					10					15

<210> SEQ ID NO 53
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 53

Ala	Arg	Leu	Asp	Thr	Ser	Ser	Tyr	Val	Asp	Tyr	Ala	Met	Asp	Tyr
1				5					10					15

<210> SEQ ID NO 54
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 54

Ala	Arg	Leu	Asp	Ser	Ser	Ala	Tyr	Val	Glu	Tyr	Pro	Met	Asp	Tyr
1				5					10					15

<210> SEQ ID NO 55
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 55

Ala	Arg	Leu	Asp	Ser	Ser	Thr	Tyr	Val	Asp	Tyr	Ala	Met	Asp	Tyr
1				5					10					15

<210> SEQ ID NO 56
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 56

Ala	Arg	Leu	Asp	Ser	Ser	Ala	Tyr	Val	Gly	Tyr	Ala	Met	Asp	Tyr
1				5					10					15

<210> SEQ ID NO 57
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 57

Ala	Arg	Leu	Asp	Ser	Ser	Ala	Tyr	Val	Asp	Tyr	Pro	Met	Asp	Phe
1				5					10					15

<210> SEQ ID NO 58
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 58

Ala	Arg	Leu	Asp	Ser	Ser	Ala	Tyr	Leu	Gly	Tyr	Pro	Leu	Asp	Tyr
1				5					10					15

-continued

<210> SEQ ID NO 59
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 59

Ala Arg Leu Asp Ser Ser Gly Tyr Val Gly Tyr Ala Met Asp Tyr
1 5 10 15

<210> SEQ ID NO 60
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 60

Ala Arg Leu Asp Ser Ala Ala Tyr Val Gly Tyr Ala Met Asp Tyr
1 5 10 15

<210> SEQ ID NO 61
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 61

Ile Arg Leu Asp Ser Ser Tyr Tyr Val Asp Tyr Pro Val Asp Tyr
1 5 10 15

<210> SEQ ID NO 62
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 62

Ala Arg Leu Asp Ser Ser Tyr Tyr Val Asp Tyr Pro Val Asp Tyr
1 5 10 15

<210> SEQ ID NO 63
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 63

Ala Arg Leu Asp Ile Ser Asp Tyr Val Asp Tyr Pro Met Asp His
1 5 10 15

<210> SEQ ID NO 64
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 64

Ala Arg Leu Asp Ser Ser Ser Tyr Val Asp Tyr Pro Met Asp Tyr
1 5 10 15

<210> SEQ ID NO 65
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 65

Ala Arg Leu Asp Ser Ser Ala Tyr Val Gly Tyr Pro Met Asp Tyr

-continued

1	5	10	15
---	---	----	----

<210> SEQ ID NO 66
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 66

Ala	Arg	Leu	Asp	Ser	Ser	Cys	Tyr	Val	Asp	Tyr	Pro	Val	Asp	Tyr
1				5				10					15	

<210> SEQ ID NO 67
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 67

Ala	Arg	Leu	Asp	Ser	Ser	Tyr	Phe	Val	Asp	Tyr	Pro	Leu	Asp	Ser
1				5				10					15	

<210> SEQ ID NO 68
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 68

Val	Arg	Leu	Asp	Ser	Ser	Tyr	Tyr	Val	Asp	Tyr	Pro	Val	Asp	Tyr
1				5				10					15	

<210> SEQ ID NO 69
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 69

Ala	Arg	Gln	Arg	Gly	Ser	Tyr	Tyr	Arg	Tyr	Asp	Asp	Tyr	Ala	Met	Asp
1				5				10					15		

Tyr

<210> SEQ ID NO 70
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 70

Ala	Arg	Trp	Gly	Asn	Tyr	Val	Ser	Tyr	Tyr	Thr	Met	Asp	Tyr
1				5				10					

<210> SEQ ID NO 71
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 71

Leu	Gln	Tyr	Asp	Glu	Phe	Pro	Tyr	Thr
1				5				

<210> SEQ ID NO 72
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

-continued

<400> SEQUENCE: 72

Leu Gln Tyr Asp Glu Phe Pro Phe Thr
1 5

<210> SEQ ID NO 73

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 73

Gln Gln Tyr Asp Glu Phe Pro Tyr Thr
1 5

<210> SEQ ID NO 74

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 74

Leu Gln Tyr Asp Asp Phe Pro Tyr Thr
1 5

<210> SEQ ID NO 75

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 75

Leu Gln Tyr Ala Glu Phe Pro Tyr Thr
1 5

<210> SEQ ID NO 76

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 76

Leu His Tyr Asp Glu Phe Pro Tyr Thr
1 5

<210> SEQ ID NO 77

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 77

Phe Gln Gly Ser His Val Pro Trp Thr
1 5

<210> SEQ ID NO 78

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 78

Gln His Tyr His Ser Tyr Pro Pro Met Trp Thr
1 5 10

<210> SEQ ID NO 79

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

-continued

<400> SEQUENCE: 79

Ala Arg Leu Asp Tyr Asp Gly Tyr Tyr Glu Tyr Ala Met Asp Tyr
1 5 10 15

<210> SEQ ID NO 80

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 80

Ala Arg Leu Asp Tyr Asp Thr Tyr Tyr Glu Tyr Ala Met Asp Tyr
1 5 10 15

<210> SEQ ID NO 81

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 81

Ala Arg Leu Asp Phe Asp Gly Tyr Tyr Glu Tyr Pro Met Asp Tyr
1 5 10 15

<210> SEQ ID NO 82

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 82

Ser Arg Leu Asp Phe Asp Gly Tyr Tyr Glu Tyr Ala Met Asp Tyr
1 5 10 15

<210> SEQ ID NO 83

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 83

Thr Arg Leu Asp Tyr Asp Gly Tyr Tyr Asp Tyr Ala Val Asp Tyr
1 5 10 15

<210> SEQ ID NO 84

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 84

Thr Arg Leu Asp Tyr Asp Gly Tyr Tyr Asp Tyr Ala Met Asp Tyr
1 5 10 15

<210> SEQ ID NO 85

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 85

Ala Arg Leu Asp Tyr Asp Gly Tyr Tyr Asp Phe Ala Met Asp Tyr
1 5 10 15

<210> SEQ ID NO 86

<211> LENGTH: 15

<212> TYPE: PRT

-continued

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 86

Ala Arg Leu Asp Tyr Asp Gly Tyr Tyr Glu Tyr Pro Met Asp Tyr
1 5 10 15

<210> SEQ ID NO 87

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 87

Ala Arg Leu Asp Tyr Asp Gly Tyr Tyr Glu Asn Ala Met Asp Tyr
1 5 10 15

<210> SEQ ID NO 88

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 88

Ala Arg Leu Asp Tyr Asp Ala Tyr Tyr Glu Tyr Ala Met Asp Tyr
1 5 10 15

<210> SEQ ID NO 89

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 89

Ala Arg Leu Asp Tyr Ala Gly Tyr Tyr Glu Tyr Ala Met Asp Tyr
1 5 10 15

<210> SEQ ID NO 90

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 90

Thr Arg Leu Asp Tyr Glu Gly Tyr Tyr Glu Tyr Ala Met Asp Tyr
1 5 10 15

<210> SEQ ID NO 91

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 91

Thr Arg Leu Asp Tyr Asp Ser Tyr Tyr Glu Tyr Ala Leu Asp Tyr
1 5 10 15

<210> SEQ ID NO 92

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 92

Ser Arg Leu Asp Tyr Asp Gly Tyr Tyr Glu Tyr Ala Met Asp Tyr
1 5 10 15

<210> SEQ ID NO 93

<211> LENGTH: 15

-continued

<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 93

Ser Arg Leu Asp Tyr Asp Gly Tyr Tyr Glu Tyr Thr Met Asp Tyr
1 5 10 15

<210> SEQ ID NO 94
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 94

Ala Arg Leu Asp Tyr Asp Gly Tyr Tyr Asp Tyr Ala Leu Asp Tyr
1 5 10 15

<210> SEQ ID NO 95
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 95

Ala Arg Leu Asp Tyr Asp Gly Tyr Tyr Glu Tyr Pro Leu Asp Tyr
1 5 10 15

<210> SEQ ID NO 96
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 96

Ala Arg Leu Asp Tyr Asp Gly Tyr Tyr Asp Tyr Ala Met Asp Tyr
1 5 10 15

<210> SEQ ID NO 97
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 97

Ala Arg Leu Asp Tyr Asp Gly Tyr Tyr Glu Tyr Ala Val Asp Tyr
1 5 10 15

<210> SEQ ID NO 98
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 98

Ala Arg Leu Asp Tyr Asp Gly Tyr Tyr Glu Tyr Ala Leu Asp Tyr
1 5 10 15

<210> SEQ ID NO 99
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 99

Ala Arg Leu Asp Tyr Asp Ala Tyr Tyr Glu Asn Ala Met Asp Tyr
1 5 10 15

<210> SEQ ID NO 100

-continued

<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 100

Gly Arg Leu Asp Ser Leu His Tyr Tyr Asp Tyr Gly Phe Asp Tyr
1 5 10 15

<210> SEQ ID NO 101
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 101

Val Arg Leu Asp Tyr Asp Gly Tyr Tyr Glu Tyr Ser Met Asp Tyr
1 5 10 15

<210> SEQ ID NO 102
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 102

Thr Arg Leu Asp Tyr Asp Gly Tyr Tyr Glu Tyr Ala Leu Asp Tyr
1 5 10 15

<210> SEQ ID NO 103
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 103

Ala Arg Asp Gly Gly Asp Tyr Asp Trp Phe Ala Tyr
1 5 10

<210> SEQ ID NO 104
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 104

Leu Gln Tyr Asp Glu Phe Pro Arg Thr
1 5

<210> SEQ ID NO 105
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 105

Leu Gln Tyr Asp Asp Phe Pro Arg Thr
1 5

<210> SEQ ID NO 106
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 106

Gln His His Tyr Gly Ser Ser Thr
1 5

-continued

<210> SEQ ID NO 107

<211> LENGTH: 199

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 107

Asn Ile Thr Asn Leu Cys Pro Phe Gly Glu Val Phe Asn Ala Ser Lys
1 5 10 15

Phe Ala Ser Val Tyr Ala Trp Asn Arg Lys Arg Ile Ser Asn Cys Val
20 25 30

Ala Asp Tyr Ser Val Leu Tyr Asn Ser Thr Ser Phe Ser Thr Phe Lys
35 40 45

Cys Tyr Gly Val Ser Pro Thr Lys Leu Asn Asp Leu Cys Phe Thr Asn
50 55 60

Val Tyr Ala Asp Ser Phe Val Val Lys Gly Asp Glu Val Arg Gln Ile
65 70 75 80

Ala Pro Gly Gln Thr Gly Val Ile Ala Asp Tyr Asn Tyr Lys Leu Pro
85 90 95

Asp Asp Phe Thr Gly Cys Val Ile Ala Trp Asn Ser Val Lys Gln Asp
100 105 110

Ala Leu Thr Gly Gly Asn Tyr Gly Tyr Leu Tyr Arg Leu Phe Arg Lys
115 120 125

Ser Lys Leu Lys Pro Phe Glu Arg Asp Ile Ser Thr Glu Ile Tyr Gln
130 135 140

Ala Gly Ser Thr Pro Cys Asn Gly Gln Val Gly Leu Asn Cys Tyr Tyr
145 150 155 160

Pro Leu Glu Arg Tyr Gly Phe His Pro Thr Thr Gly Val Asn Tyr Gln
165 170 175

Pro Phe Arg Val Val Val Leu Ser Phe Glu Leu Leu Asn Gly Pro Ala
180 185 190

Thr Val Cys Gly Pro Lys Leu
195

<210> SEQ ID NO 108

<211> LENGTH: 198

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 108

Asn Ile Thr Asn Leu Cys Pro Phe Gly Glu Val Phe Asn Ala Thr Lys
1 5 10 15

Phe Pro Ser Val Tyr Ala Trp Glu Arg Lys Lys Ile Ser Asn Cys Val
20 25 30

Ala Asp Tyr Ser Val Leu Tyr Asn Ser Thr Phe Phe Ser Thr Phe Lys
35 40 45

Cys Tyr Gly Val Ser Ala Thr Lys Leu Asn Asp Leu Cys Phe Ser Asn
50 55 60

Val Tyr Ala Asp Ser Phe Val Val Lys Gly Asp Asp Val Arg Gln Ile
65 70 75 80

Ala Pro Gly Gln Thr Gly Val Ile Ala Asp Tyr Asn Tyr Lys Leu Pro
85 90 95

Asp Asp Phe Met Gly Cys Val Leu Ala Trp Asn Thr Arg Asn Ile Asp
100 105 110

Ala Thr Ser Thr Gly Asn Tyr Asn Tyr Lys Tyr Arg Tyr Leu Arg His

-continued

115	120	125
Gly Lys Leu Arg Pro Phe	Glu Arg Asp Ile Ser Asn Val Pro Phe Ser	
130	135	140
Pro Asp Gly Lys Pro Cys Thr Pro Pro Ala Leu Asn Cys Tyr Trp Pro		
145	150	155
Leu Asn Asp Tyr Gly Phe Tyr Thr Thr Thr Gly Ile Gly Tyr Gln Pro		
165	170	175
Tyr Arg Val Val Val Leu Ser Phe Glu Leu Leu Asn Ala Pro Ala Thr		
180	185	190
Val Cys Gly Pro Lys Leu		
195		

<210> SEQ ID NO 109
 <211> LENGTH: 198
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 109

Asn Ile Thr Asn Leu Cys Pro Phe Gly Glu Val Phe Asn Ala Thr Thr	
1	5
Phe Pro Ser Val Tyr Ala Trp Glu Arg Lys Arg Ile Ser Asn Cys Val	
20	25
Ala Asp Tyr Ser Val Leu Tyr Asn Ser Thr Ser Phe Ser Thr Phe Lys	
35	40
Cys Tyr Gly Val Ser Ala Thr Lys Leu Asn Asp Leu Cys Phe Ser Asn	
50	55
Val Tyr Ala Asp Ser Phe Val Val Lys Gly Asp Asp Val Arg Gln Ile	
65	70
Ala Pro Gly Gln Thr Gly Val Ile Ala Asp Tyr Asn Tyr Lys Leu Pro	
85	90
Asp Asp Phe Thr Gly Cys Val Leu Ala Trp Asn Thr Arg Asn Ile Asp	
100	105
Ala Thr Gln Thr Gly Asn Tyr Asn Tyr Lys Tyr Arg Ser Leu Arg His	
115	120
Gly Lys Leu Arg Pro Phe Glu Arg Asp Ile Ser Asn Val Pro Phe Ser	
130	135
Pro Asp Gly Lys Pro Cys Thr Pro Pro Ala Phe Asn Cys Tyr Trp Pro	
145	150
Leu Asn Asp Tyr Gly Phe Tyr Ile Thr Asn Gly Ile Gly Tyr Gln Pro	
165	170
Tyr Arg Val Val Val Leu Ser Phe Glu Leu Leu Asn Ala Pro Ala Thr	
180	185
Val Cys Gly Pro Lys Leu	
195	

<210> SEQ ID NO 110
 <211> LENGTH: 198
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 110

Asn Ile Thr Asn Leu Cys Pro Phe Gly Glu Val Phe Asn Ala Thr Thr	
1	5
Phe Pro Ser Val Tyr Ala Trp Glu Arg Lys Arg Ile Ser Asn Cys Val	

-continued

20					25					30					
Ala	Asp	Tyr	Ser	Val	Leu	Tyr	Asn	Ser	Thr	Ser	Phe	Ser	Thr	Phe	Lys
	35						40					45			
Cys	Tyr	Gly	Val	Ser	Ala	Thr	Lys	Leu	Asn	Asp	Leu	Cys	Phe	Ser	Asn
	50					55					60				
Val	Tyr	Ala	Asp	Ser	Phe	Val	Val	Lys	Gly	Asp	Asp	Val	Arg	Gln	Ile
	65					70					75				80
Ala	Pro	Gly	Gln	Thr	Gly	Val	Ile	Ala	Asp	Tyr	Asn	Tyr	Lys	Leu	Pro
				85					90					95	
Asp	Asp	Phe	Leu	Gly	Cys	Val	Leu	Ala	Trp	Asn	Thr	Asn	Ser	Lys	Asp
			100					105					110		
Ser	Ser	Thr	Ser	Gly	Asn	Tyr	Asn	Tyr	Leu	Tyr	Arg	Trp	Val	Arg	Arg
		115					120					125			
Ser	Lys	Leu	Asn	Pro	Tyr	Glu	Arg	Asp	Leu	Ser	Asn	Asp	Ile	Tyr	Ser
	130					135					140				
Pro	Gly	Gly	Gln	Ser	Cys	Ser	Ala	Val	Gly	Pro	Asn	Cys	Tyr	Asn	Pro
	145					150					155				160
Leu	Arg	Pro	Tyr	Gly	Phe	Phe	Thr	Thr	Ala	Gly	Val	Gly	His	Gln	Pro
				165					170					175	
Tyr	Arg	Val	Val	Val	Leu	Ser	Phe	Glu	Leu	Leu	Asn	Ala	Pro	Ala	Thr
			180					185					190		
Val	Cys	Gly	Pro	Lys	Leu										
		195													

<210> SEQ ID NO 111

<211> LENGTH: 180

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 111

Asn	Ile	Thr	Asn	Arg	Cys	Pro	Phe	Asp	Ser	Ile	Phe	Asn	Ala	Ser	Arg
1				5					10					15	
Phe	Pro	Ser	Val	Tyr	Ala	Trp	Glu	Arg	Thr	Lys	Ile	Ser	Asp	Cys	Val
			20					25					30		
Ala	Asp	Tyr	Thr	Val	Leu	Tyr	Asn	Ser	Thr	Ser	Phe	Ser	Thr	Phe	Lys
		35					40					45			
Cys	Tyr	Gly	Val	Ser	Pro	Ser	Lys	Leu	Ile	Asp	Leu	Cys	Phe	Thr	Ser
	50					55					60				
Val	Tyr	Ala	Asp	Thr	Phe	Leu	Ile	Arg	Phe	Ser	Glu	Val	Arg	Gln	Val
	65				70				75					80	
Ala	Pro	Gly	Glu	Thr	Gly	Val	Ile	Ala	Asp	Tyr	Asn	Tyr	Arg	Leu	Pro
			85					90					95		
Asp	Asp	Phe	Thr	Gly	Cys	Val	Ile	Ala	Trp	Asn	Thr	Ala	Asn	Gln	Asp
			100					105					110		
Val	Gly	Ser	Tyr	Phe	Tyr	Arg	Ser	His	Arg	Ser	Thr	Lys	Leu	Lys	Pro
		115						120				125			
Phe	Glu	Arg	Asp	Leu	Ser	Ser	Asp	Glu	Asn	Gly	Val	Arg	Thr	Leu	Ser
	130					135					140				
Thr	Tyr	Asp	Phe	Asn	Pro	Tyr	Val	Pro	Leu	Asp	Tyr	Gln	Ala	Thr	Arg
	145				150						155				160
Val	Val	Val	Leu	Ser	Phe	Glu	Leu	Leu	Asn	Ala	Pro	Ala	Thr	Val	Cys
			165						170					175	

-continued

Gly Pro Lys Leu
180

<210> SEQ ID NO 112
 <211> LENGTH: 180
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 112

Asn Ile Thr Asn Leu Cys Pro Phe Asp Lys Val Phe Asn Ala Thr Arg
 1 5 10 15
 Phe Pro Ser Val Tyr Ala Trp Glu Arg Thr Lys Ile Ser Asp Cys Val
 20 25 30
 Ala Asp Tyr Thr Val Phe Tyr Asn Ser Thr Ser Phe Ser Thr Phe Asn
 35 40 45
 Cys Tyr Gly Val Ser Pro Ser Lys Leu Ile Asp Leu Cys Phe Thr Ser
 50 55 60
 Val Tyr Ala Asp Thr Phe Leu Ile Arg Phe Ser Glu Val Arg Gln Val
 65 70 75 80
 Ala Pro Gly Gln Thr Gly Val Ile Ala Asp Tyr Asn Tyr Lys Leu Pro
 85 90 95
 Asp Asp Phe Thr Gly Cys Val Ile Ala Trp Asn Thr Ala Lys Gln Asp
 100 105 110
 Val Gly Ser Tyr Phe Tyr Arg Ser His Arg Ser Ser Lys Leu Lys Pro
 115 120 125
 Phe Glu Arg Asp Leu Ser Ser Glu Glu Asn Gly Val Arg Thr Leu Ser
 130 135 140
 Thr Tyr Asp Phe Asn Gln Asn Val Pro Leu Glu Tyr Gln Ala Thr Arg
 145 150 155 160
 Val Val Val Leu Ser Phe Glu Leu Leu Asn Ala Pro Ala Thr Val Cys
 165 170 175

Gly Pro Lys Leu
180

<210> SEQ ID NO 113
 <211> LENGTH: 180
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 113

Asn Ile Thr Asn Arg Cys Pro Phe Asp Lys Val Phe Asn Ala Ser Arg
 1 5 10 15
 Phe Pro Asn Val Tyr Ala Trp Glu Arg Thr Lys Ile Ser Asp Cys Val
 20 25 30
 Ala Asp Tyr Thr Val Leu Tyr Asn Ser Thr Ser Phe Ser Thr Phe Lys
 35 40 45
 Cys Tyr Gly Val Ser Pro Ser Lys Leu Ile Asp Leu Cys Phe Thr Ser
 50 55 60
 Val Tyr Ala Asp Thr Phe Leu Ile Arg Ser Ser Glu Val Arg Gln Val
 65 70 75 80
 Ala Pro Gly Glu Thr Gly Val Ile Ala Asp Tyr Asn Tyr Lys Leu Pro
 85 90 95
 Asp Asp Phe Thr Gly Cys Val Ile Ala Trp Asn Thr Ala Lys Gln Asp
 100 105 110

```
<210> SEQ ID NO 114
<211> LENGTH: 195
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 114
```

Asn	Ile	Thr	Gln	Leu	Cys	Pro	Phe	Asn	Glu	Val	Phe	Asn	Ile	Thr	Ser	1		
				5					10								15	
Phe	Pro	Ser	Val	Tyr	Ala	Trp	Glu	Arg	Met	Arg	Ile	Thr	Asn	Cys	Val			
				20					25								30	
Ala	Asp	Tyr	Ser	Val	Leu	Tyr	Asn	Ser	Ser	Ala	Ser	Phe	Ser	Thr	Phe			
				35					40								45	
Gln	Cys	Tyr	Gly	Val	Ser	Pro	Thr	Lys	Leu	Asn	Asp	Leu	Cys	Phe	Ser			
				50					55								60	
Ser	Val	Tyr	Ala	Asp	Tyr	Phe	Val	Val	Lys	Gly	Asp	Asp	Val	Arg	Gln			
				65					70								75	
Ile	Ala	Pro	Ala	Gln	Thr	Gly	Val	Ile	Ala	Asp	Tyr	Asn	Tyr	Lys	Leu			
				85					90								95	
Pro	Asp	Asp	Phe	Thr	Gly	Cys	Val	Ile	Ala	Trp	Asn	Thr	Asn	Ser	Leu			
				100					105								110	
Asp	Ser	Ser	Asn	Glu	Phe	Phe	Tyr	Arg	Arg	Phe	Arg	His	Gly	Lys	Ile			
				115					120								125	
Lys	Pro	Tyr	Gly	Arg	Asp	Leu	Ser	Asn	Val	Leu	Phe	Asn	Pro	Ser	Gly			
				130					135								140	
Gly	Thr	Cys	Ser	Ala	Glu	Gly	Leu	Asn	Cys	Tyr	Lys	Pro	Leu	Ala	Ser			
				145					150								155	
Tyr	Gly	Phe	Thr	Gln	Ser	Ser	Gly	Ile	Gly	Phe	Gln	Pro	Tyr	Arg	Val			
				165					170								175	
Val	Val	Leu	Ser	Phe	Glu	Leu	Leu	Asn	Ala	Pro	Ala	Thr	Val	Cys	Gly			
				180					185								190	
Pro	Lys	Gln																
				195														

1-11. (canceled)

12. A polypeptide comprising an amino acid sequence having at least 90% or more identity to the amino acid sequence of SEQ ID NO: 2, wherein the amino acid sequence is modified so as to allow sugar chain attachment to positions corresponding to any one or more of: positions 85, 118, 147, 164, and 172 of the amino acid sequence of SEQ ID NO: 2.

13. The polypeptide according to claim **12**, wherein the polypeptide comprises an amino acid sequence having at least 95% or more identity to the amino acid sequence of SEQ ID NO: 2.

14. The polypeptide according to claim **12**, wherein the polypeptide comprises an amino acid sequence having at least 97% or more identity to the amino acid sequence of SEQ ID NO: 2.

15. The polypeptide according to claim **12**, wherein the polypeptide comprises an amino acid sequence having at least 98% or more identity to the amino acid sequence of SEQ ID NO: 2.

16. The polypeptide according to claim **12**, wherein the sugar chain is an N-type sugar chain.

17. The polypeptide according to claim **12**, wherein a sugar chain is attached to each of positions corresponding to positions 85, 164, and 172 of the amino acid sequence of SEQ ID NO: 2.

18. The polypeptide according to claim **17**, wherein a sugar chain is further attached to a position corresponding to position 118 and/or a position corresponding to position 147.

19. A polynucleotide comprising a base sequence encoding the polypeptide of claim **12**.

20. A vaccine or pharmaceutical composition comprising the polypeptide of claim **12**.

21. A vaccine or pharmaceutical composition comprising the polynucleotide of claim **19**.

22. A method for producing an anti-coronavirus antibody, comprising immunizing an animal with the polypeptide of claim **12**.

23. A method for producing an anti-coronavirus antibody, comprising immunizing an animal with the polynucleotide of claim **19**.

24. An antibody obtained by the method of claim **22**.

* * * * *