



US 20250277012A1

(19) **United States**(12) **Patent Application Publication**
BHANDARI et al.(10) **Pub. No.: US 2025/0277012 A1**(43) **Pub. Date: Sep. 4, 2025**(54) **CONJUGATED HEPCIDIN MIMETICS**

filed on Apr. 1, 2021, provisional application No. 63/169,533, filed on Apr. 1, 2021, provisional application No. 63/169,527, filed on Apr. 1, 2021.

(71) Applicant: **Protagonist Therapeutics, Inc.**,
Newark, CA (US)(72) Inventors: **Ashok BHANDARI**, Pleasanton, CA (US); **Jie ZHANG**, Salisbury (AU); **Gregory Thomas BOURNE**, Moggill, Brisbane (AU); **Mark Leslie SMYTHE**, Bardon (AU); **Tran Trung TRAN**, Newark, CA (US); **Roopa TARANATH**, Cupertino, CA (US)**Publication Classification**(51) **Int. Cl.****C07K 14/575** (2006.01)**C07K 1/107** (2006.01)(52) **U.S. Cl.**CPC **C07K 14/575** (2013.01); **C07K 1/1077** (2013.01)(21) Appl. No.: **18/284,835**(22) PCT Filed: **Mar. 31, 2022**(86) PCT No.: **PCT/US2022/022809**

§ 371 (c)(1),

(2) Date: **Sep. 28, 2023****Related U.S. Application Data**

(60) Provisional application No. 63/325,323, filed on Mar. 30, 2022, provisional application No. 63/169,538,

(57) **ABSTRACT**

The present invention provides hepcidin analogues with improved in vivo half lives, and related pharmaceutical compositions and methods of use thereof. The present invention provides novel peptide analogues, including both peptide monomer analogues and peptide dimer analogues, having hepcidin activity, and also having other beneficial properties making the peptides of the present invention suitable alternatives to hepcidin.

Specification includes a Sequence Listing.

CONJUGATED HEPCIDIN MIMETICS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of and priority to U.S. Provisional Application No. 63/169,527, filed Apr. 1, 2021; U.S. Provisional Application No. 63/169,533, filed Apr. 1, 2021; U.S. Provisional Application No. 63/169,538, filed Apr. 1, 2021; and U.S. Provisional Application No. 63/325,323, filed Mar. 30, 2022, each of which is hereby incorporated by reference in its entirety.

FIELD OF THE INVENTION

[0002] The present invention relates, inter alia, to certain hepcidin peptide analogues, including both peptide monomers and peptide dimers, and conjugates and derivatives thereof, as well as compositions comprising the peptide analogues, and to the use of the peptide analogues in the treatment and/or prevention of a variety of diseases, conditions or disorders, including treatment and/or prevention of erythrocytoses, such as polycythemia vera, iron overload diseases such as hereditary hemochromatosis, iron-loading anemias, and other conditions and disorders described herein.

BACKGROUND

[0003] Hepcidin (also referred to as LEAP-1), a peptide hormone produced by the liver, is a regulator of iron homeostasis in humans and other mammals. Hepcidin acts by binding to its receptor, the iron export channel ferroportin, causing its internalization and degradation. Human hepcidin is a 25-amino acid peptide (Hep25). See Krause et al. (2000) FEBS Lett 480:147-150, and Park et al. (2001) J Biol Chem 276:7806-7810. The structure of the bioactive 25-amino acid form of hepcidin is a simple hairpin with 8 cysteines that form 4 disulfide bonds as described by Jordan et al. J Biol Chem 284:24155-67. The N terminal region is required for iron-regulatory function, and deletion of 5 N-terminal amino acid residues results in a loss of iron-regulatory function. See Nemeth et al. (2006) Blood 107: 328-33.

[0004] Abnormal hepcidin activity is associated with iron overload diseases, including hereditary hemochromatosis (HH) and iron-loading anemias. Hereditary hemochromatosis is a genetic iron overload disease that is mainly caused by hepcidin deficiency or in some cases by hepcidin resistance. This allows excessive absorption of iron from the diet and development of iron overload. Clinical manifestations of HH may include liver disease (e.g., hepatic cirrhosis NASH, and hepatocellular carcinoma), diabetes, and heart failure. Currently, the only treatment for HH is regular phlebotomy, which is very burdensome for the patients. Iron-loading anemias are hereditary anemias with ineffective erythropoiesis such as β -thalassemia, which are accompanied by severe iron overload. Complications from iron overload are the main causes of morbidity and mortality for these patients. Hepcidin deficiency is the main cause of iron overload in non-transfused patients, and contributes to iron overload in transfused patients. The current treatment for iron overload in these patients is iron chelation, which is very burdensome, sometimes ineffective, and accompanied by frequent side effects.


[0005] Hepcidin has several limitations that restrict its use as a drug, including a difficult synthetic process due in part to aggregation and precipitation of the protein during folding, which in turn leads to low bioavailability, injection site reactions, immunogenicity, and high cost of goods. What are needed in the art are compounds having hepcidin activity and also possessing other beneficial physical properties such as improved solubility, stability, and/or potency, so that hepcidin-like compounds might be produced affordably and used to treat hepcidin-related diseases and disorders such as, e.g., those described herein.

[0006] The present invention addresses such needs, providing novel peptide analogues, including both peptide monomer analogues and peptide dimer analogues, having hepcidin activity, and also having other beneficial properties making the peptides of the present invention suitable alternatives to hepcidin.

BRIEF SUMMARY OF THE INVENTION

[0007] The present invention generally relates to peptide analogues, including both monomer and dimers, exhibiting hepcidin activity and methods of using the same.

[0008] In one aspect, the present invention provides a hepcidin analogue comprising a peptide comprising an amino acid sequence of Formula (I'):



or a pharmaceutically acceptable salt or solvate thereof, wherein:

[0009] R^1 is hydrogen, $\text{C}_1\text{--C}_6$ alkyl, $\text{C}_6\text{--C}_{12}$ aryl, $\text{C}_6\text{--C}_{12}$ aryl- $\text{C}_1\text{--C}_6$ alkyl, $\text{C}_1\text{--C}_{20}$ alkanoyl, $\text{C}_1\text{--C}_{20}$ cycloalkanoyl, $\text{C}_6\text{--C}_{10}$ aryl- $\text{C}_1\text{--C}_6$ alkylene- $\text{C}(\text{O})\text{---}$, 5- or 6-membered heteroaryl- $\text{C}(\text{O})\text{---}$ or 5- or 6-membered heterocycloalkyl- $\text{C}(\text{O})\text{---}$, wherein the $\text{C}_1\text{--C}_{20}$ alkanoyl, $\text{C}_6\text{--C}_{10}$ aryl- $\text{C}_1\text{--C}_6$ alkylene- $\text{C}(\text{O})\text{---}$, 5- or 6-membered heteroaryl- $\text{C}(\text{O})\text{---}$ and 5- or 6-membered heterocycloalkyl- $\text{C}(\text{O})\text{---}$ of R^1 are each optionally substituted with 1, 2 or 3 substituents independently selected from halo, OH, CN, $\text{C}_1\text{--C}_6$ alkoxy, $\text{C}_1\text{--C}_6$ alkyl, ---COOH , $\text{---COO}(\text{C}_1\text{--C}_6)$, $\text{C}_1\text{--C}_6$ haloalkyl and $\text{C}_1\text{--C}_6$ haloalkoxy;

[0010] R^2 is NH_2 , substituted amino, OH, substituted hydroxy, $\text{C}_1\text{--C}_{20}$ alkylamino, phenyl- $\text{C}_1\text{--C}_8$ alkylene- NH--- , wherein $\text{C}_1\text{--C}_{20}$ alkylamino and phenyl- $\text{C}_1\text{--C}_8$ alkylene- NH--- are each optionally substituted with a substituent independently selected from NH_2 , ---COOH and phenyl;


[0011] X1 is absent, or is Asp, isoAsp, Asp(OMe), Glu, bhGlu, bGlu, Gly, N-substituted Gly, Ala, Gln, Arg, Dab, Leu, Lys, Dap, Orn, (D)Asp, (D)Arg, Tet1, Tet2, Lys, substituted Lys, (D)Lys, substituted (D)Lys, or Cys, (D)Cys, aMeCys, hCys, Pen, (D)Pen, NMe-Glu, Aad, Dap or isoGlu;

[0012] X2 is Ala, Thr, Gly, N-substituted Gly, Ser, Cys, 4S_Mcp, 4R_Mcp, NMe-Thr, aMePro, Hyp_3R, HyP_3S, hSer or Thr_Me;

[0013] X3 is Ala, Gly, N-substituted Gly, His, substituted His, Cys, (D)Cys, aMeCys, hCys, Pen, 3Pal, 4Pal, BIP, Ala 3Quin, Trp_50H or His_1Me;


[0014] X4 is Ala, Phe, Dpa, Gly, N-substituted Gly, bhPhe, a-MePhe, NMe-Phe, D-Phe, 2Pal, or Cys, (D)Cys, aMeCys, hCys, Pen or DIP;

- [0015] X5 is Ala, Pro, D-Pro, bhPro, D-bhPro, NPC, D-NPC, Gaba, 2-Pyrrolidinepropanoic acid (Ppa), 2-Pyrrolidinebutanoic acid (Pba), Glu, Lys, substituted Lys, (D)Lys, substituted (D)Lys, or Cys, (D)Cys, aMeCys, hCys, Pen, NMe-Cys, 4S_Mcp, 4R_Mcp, Morph or Tic;
- [0016] X6 is Cys, (D)Cys, aMeCys, hCys, Pen, dPen, NMe-Cys, Ala, Lys, (D)Lys, substituted Lys or substituted (D)Lys;
- [0017] X7 is absent, or is Ala, Gly, N-substituted Gly, Ile, Val, Leu, NLeu, Lys, substituted Lys, (D)Lys, substituted (D)Lys, Cys, (D)Cys, aMeCys, hCys, Pen, NMe-Cys, Phe or DIP;
- [0018] X8 is absent or is Ala, (D)Ala, Ile, Gly, N-substituted Gly, Glu, Val, Leu, NLeu, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys, aMeLys, 123Triazole, Cys, (D)Cys, aMeCys, hCys, Pen, Lys_Me3;
- [0019] X9 is absent, or is Ala, Ile, Gly, N-substituted Gly, Val, Leu, NLeu, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys, Cys, (D)Cys, aMeCys, hCys, Pen, NMe-Phe, aMePhe, Dip, dDIP, BIP, aMePhe or substituted Phe;
- [0020] X10 is absent, or Ala, Gly, N-substituted Gly, Ile, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys, Cys, (D)Cys, aMeCys, hCys, Pen, Tle, substituted (D)Lys, NMe-Lys, NMe-dLys, substituted NMe-Lys, substituted NMe-dLys or Mor_propanoic_acid;
- [0021] X11 is absent, or is Ala, Pro, bhPhe, Lys, substituted Lys, (D)Lys, NMe-Phe, NMe-dSer, NMe-dGln, NMe-dLeu, NMe-dTyr, 1Nal or NMe-Ile;
- [0022] each of X12-X14 is absent, or is independently an amino acid;
- [0023] the peptide of formula (I') is optionally further conjugated with any amino acid;
- [0024] any of the amino acids of the peptide of formula (I') is optionally replaced with the corresponding (D)-amino acid or optionally N-substituted;
- [0025] (i) at least one of X1, X3-X5, X7-X10 is Cys, (D)Cys, aMeCys, hCys, Pen, (D)Pen, NMe-Cys, 4S_Mcp or 4R_Mcp, wherein the peptide is cyclized by taking the mercapto group on the side chain of one of X1, X3-X5 and X7-X10, the mercapto group on the side chain of X6 and L^x to form a —S-L^x— linkage; or the mercapto group on the side chain of X1, the mercapto group on the side chain of X8 and L^x taken together form a —S-L^x— linkage; or the mercapto group on the side chain of X2, the mercapto group on the side chain of X5 and L^x taken together form a —S-L^x— linkage, wherein each L^x is independently a bond or a linking group; and
- [0026] (ii) the functional groups on the side chains of X1 and X10 are optionally taken together to form an amide linkage; or the functional group on the side chain of X1 and the functional group of R² are optionally taken together to form an amide linkage.
- and wherein
- [0027] Dapa is diaminopropanoic acid; Dpa or DIP is 3,3-diphenylalanine or b,b-diphenylalanine; bhPhe is b-homophenylalanine; Bip is biphenylalanine; bhPro is b-homoproline; Tic is L-1,2,3,4,-tetrahydro-isoquinoline-3-carboxylic acid; NPC is L-nipecotic acid; bhTrp is b-homoTryptophane; 1-Nal is 1-naphthylalanine; 2-Nal is 2-naphthylalanine; Orn is ornithine; Nleu is norleucine; 2Pal is 2-pyridylalanine; Ppa is 2-(R)-Pyrrolidinepropanoic acid, Pba is 2-(R)-Pyrrolidinebutanoic acid; substituted Phe is phenylalanine wherein phenyl is substituted with F, Cl, Br, I, OH, methoxy, dimethoxy, dichloro, dimethyl, difluoro, pentafluoro, allyloxy, azido, nitro, 4-carbamoyl-2,6-dimethyl, trifluoromethoxy, trifluoromethyl, phenoxy, benzyloxy, carbamoyl, t-Bu, carboxyl, CN, or guanidine;
- [0028] substituted bhPhe is b-homophenylalanine wherein phenyl is substituted with F, Cl, Br, I, OH, methoxy, dimethoxy, dichloro, dimethyl, difluoro, pentafluoro, allyloxy, azido, nitro, 4-carbamoyl-2,6-dimethyl, trifluoromethoxy, trifluoromethyl, phenoxy, benzyloxy, carbamoyl, t-Bu, carboxyl, CN, or guanidine;
- [0029] substituted Trp is N-methyl-L-tryptophan, a-methyltryptophan, or tryptophan substituted with F, Cl, OH, or t-Bu;
- [0030] substituted bhTrp is N-methyl-L-b-homotryptophan, a-methyl-b-homotryptophan, or b-homotryptophan substituted with F, Cl, OH, or t-Bu;
- [0031] substituted Lys, substituted dLys or substituted NMe-Lys or substituted NMe-dLys is Lys, dLys, NMe-Lys or NMe-dLys, wherein the F-amino group on the side chain of Lys, dLys, NMe-Lys or NMe-dLys is covalently bound to C₁-C₆ alkanoyl, a half-life extension moiety or a linking moiety covalently linked to a half-life extension moiety, wherein the linking moiety comprises one or more linker moieties covalently bound to each other;
- [0032] Tet1 is (S)-(2-amino)-3-(2H-tetrazol-5-yl)propanoic acid; and Tet2 is (S)-(2-amino)-4-(1H-tetrazol-5-yl)butanoic acid;
- [0033] 123Triazole is




and

[0034] Dab is



[0035] In certain embodiments, the present invention includes a hepcidin analogue comprising a peptide of Formula (I):



or a pharmaceutically acceptable salt or solvate thereof, wherein:

[0036] R¹ is hydrogen, C₁-C₆ alkyl, C₆-C₁₂ aryl, C₆-C₁₂ aryl-C₁-C₆ alkyl, C₁-C₂₀ alkanoyl, C₁-C₂₀ cycloalkanoyl;

[0037] R^2 is NH_2 , substituted amino, OH, or substituted hydroxy;

[0038] X0 is absent, or is Cys, (D)Cys, aMeCys, hCys, or Pen;

[0039] X1 is absent, or is Asp, isoAsp, Asp(OMe), Glu, bhGlu, bGlu, Gly, N-substituted Gly, Gla, Glp, Ala, Arg, Dab, Leu, Lys, Dap, Orn, (D)Asp, (D)Arg, Tet1, or Tet2, Lys, substituted Lys, (D)Lys, substituted (D)Lys, or Cys, (D)Cys, aMeCys, hCys, or Pen;

[0040] X2 is Ala, Thr, Gly, N-substituted Gly, or Ser;

[0041] X3 is Ala, Gly, N-substituted Gly, His, substituted His, or Cys, (D)Cys, aMeCys, hCys, or Pen;

[0042] X4 is Ala, Phe, Dpa, Gly, N-substituted Gly, bhPhe, a-MePhe, NMe-Phe, D-Phe, 2Pal, or Cys, (D)Cys, aMeCys, hCys, or Pen;

[0043] X5 is Ala, Pro, D-Pro, bhPro, D-bhPro, NPC, D-NPC, Gaba, 2-Pyrrolidinepropanoic acid (Ppa), 2-Pyrrolidinebutanoic acid (Pba), Glu, Lys, substituted Lys, (D)Lys, substituted (D)Lys, or Cys, (D)Cys, aMeCys, hCys, or Pen;

[0044] X6 is Cys, (D)Cys, aMeCys, hCys, or Pen;

[0045] X7 is absent, or is Ala, Gly, N-substituted Gly, Ile, Val, Leu, NLeu, Lys, substituted Lys, (D)Lys, substituted (D)Lys, or Cys, (D)Cys, aMeCys, hCys, or Pen;

[0046] X8 is absent or is Ala, (D)Ala, Ile, Gly, N-substituted Gly, Glu, Val, Leu, NLeu, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys, aMeLys, 123Triazole, or Cys, (D)Cys, aMeCys, hCys, or Pen;

[0047] X9 is absent, or is Ala, Ile, Gly, N-substituted Gly, Val, Leu, NLeu, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys, or Cys, (D)Cys, aMeCys, hCys, or Pen;

[0048] X10 is absent, or is Ala, Gly, N-substituted Gly, Ile, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys, or Cys, (D)Cys, aMeCys, hCys, or Pen;

[0049] X11 is absent, or is Ala, Pro, bhPhe, Lys, substituted Lys, or (D)Lys;

and

each of X12-X14 is absent, or is independently any amino acid;

provided that:

[0050] i) the peptide may further be conjugated at any amino acid;

[0051] ii) any of the amino acids of the peptide may be the corresponding (D)-amino acid of the amino acid or may be N-substituted; and

[0052] iii) at least one of X0, X1, X3-X5, X7-X10 is Cys, (D)Cys, aMeCys, hCys, or Pen and Cys, (D)Cys, aMeCys, hCys, or Pen form a disulfide bond with X6;

and

[0053] Dapa is diaminopropanoic acid; Dpa or DIP is 3,3-diphenylalanine or b,b-diphenylalanine; bhPhe is b-homophenylalanine; Bip is biphenylalanine; bhPro is b-homoproline; Tic is L-1,2,3,4-tetrahydro-isoquinoline-3-carboxylic acid; NPC is L-nipecotic acid; bhTrp is b-homoTryptophane; 1-Nal is 1-naphthylalanine; 2-Nal is 2-naphthylalanine; Orn is orinithine; Nleu is norleucine; 2Pal is 2-pyridylalanine; Ppa is 2-(R)-Pyrrolidinepropanoic acid, Pba is 2-(R)-Pyrrolidinebutanoic acid; substituted Phe is phenylalanine wherein phenyl is substituted with F, Cl, Br, I, OH, methoxy, dimethoxy, dichloro, dimethyl, difluoro, pentafluoro, allyloxy, azido, nitro, 4-carbamoyl-2,6-dimethyl, trifluoromethoxy, trifluoromethyl, phenoxy, benzyloxy, carbamoyl, t-Bu, carboxyl, CN, or guanidine;


[0054] substituted bhPhe is b-homophenylalanine wherein phenyl is substituted with F, Cl, Br, I, OH, methoxy, dimethoxy, dichloro, dimethyl, difluoro, pentafluoro, allyloxy, azido, nitro, 4-carbamoyl-2,6-dimethyl, trifluoromethoxy, trifluoromethyl, phenoxy, benzyloxy, carbamoyl, t-Bu, carboxyl, CN, or guanidine;

[0055] substituted Trp is N-methyl-L-tryptophan, a-methyltryptophan, or tryptophan substituted with F, Cl, OH, or t-Bu;

[0056] substituted bhTrp is N-methyl-L-b-homotryptophan, a-methyl-b-homotryptophan, or b-homotryptophan substituted with F, Cl, OH, or t-Bu;


[0057] Tet1 is (S)-(2-amino)-3-(2H-tetrazol-5-yl)propanoic acid; and Tet2 is (S)-(2-amino)-4-(1H-tetrazol-5-yl)butanoic acid;

[0058] 123Triazole is



and

[0059] Dab is



[0060] In one embodiment, X0 is Cys, (D)Cys, aMeCys, hCys, or Pen; and X0 and X6 are linked via a disulfide bond.

[0061] In one embodiment, X1 is Cys, (D)Cys, aMeCys, hCys, or Pen; and X1 and X6 are linked via a disulfide bond.

[0062] In one embodiment, X3 is Cys, (D)Cys, aMeCys, hCys, or Pen; and X3 and X6 are linked via a disulfide bond.

[0063] In one embodiment, X4 is Cys, (D)Cys, aMeCys, hCys, or Pen; and X4 and X6 are linked via a disulfide bond.

[0064] In one embodiment, X5 is Cys, (D)Cys, aMeCys, hCys, or Pen; and X5 and X6 are linked via a disulfide bond.

[0065] In one embodiment, X7 is Cys, (D)Cys, aMeCys, hCys, or Pen; and X7 and X6 are linked via a disulfide bond.

[0066] In one embodiment, X8 is Cys, (D)Cys, aMeCys, hCys, or Pen; and X8 and X6 are linked via a disulfide bond.

[0067] In one embodiment, X9 is Cys, (D)Cys, aMeCys, hCys, or Pen; and X9 and X6 are linked via a disulfide bond.

[0068] In one embodiment, X10 is Cys, (D)Cys, aMeCys, hCys, or Pen; and X10 and X6 are linked via a disulfide bond.

[0069] In one embodiment, R^1 is IVA or isovaleric acid.

[0070] In one embodiment, R^2 is NH_2 . In one embodiment, R^2 is OH.

[0071] In particular embodiments of any of the hepcidin analogues of the present invention, the substituted Lys or substituted (D)Lys is Lys or (D)Lys substituted directly or via a linker with an acid selected from C12 (Lauric acid), C14 (Myristic acid), C16 (Palmitic acid), C18 (Stearic

acid), C20, C12 diacid, C14 diacid, C16 diacid, C18 diacid, C20 diacid, biotin, and isovaleric acid, or a residue thereof. In one embodiment, the linker is Ahx, PEG, or PEG-Ahx.

[0072] In particular embodiments of any of the hepcidin analogues of the present invention, X8 or X10 is Lys or (D)Lys substituted with LIZ; wherein L1 is absent, Dapa, D-Dapa, or isoGlu, PEG, Ahx, isoGlu-PEG, PEG-isoGlu, PEG-Ahx, isoGlu-Ahx, or isoGlu-PEG-Ahx; Ahx is an aminohexanoic acid moiety; PEG is $—[C(O)—CH_2-(Peg)_n-N(H)]_m—$, or $—[C(O)—CH_2—CH_2-(Peg)_n-N(H)]_m—$; and Peg is $—OCH_2CH_2—$, m is 1, 2, or 3; and n is an integer between 1-100K; and Z is a half-life extension moiety. In one embodiment, the half-life extension moiety is C₁₀-C₂₁ alkanoyl.

[0073] In certain embodiments, a peptide analogue or dimer of the present invention comprises an isovaleric acid moiety conjugated to an N-terminal X1 residue. In certain embodiments, a peptide analogue or dimer of the present invention comprises an isovaleric acid moiety conjugated to an N-terminal Asp residue. In certain embodiments, a peptide analogue or dimer of the present invention comprises an isovaleric acid moiety conjugated to an N-terminal Glu residue.

[0074] In certain embodiments, a peptide analogue of the present invention comprises an amidated C-terminal residue.

[0075] In certain embodiments, a peptide analogue of the present invention comprises or consists of an amino acid sequence or structure shown in Tables 6A-6B, or a variant thereof.

[0076] In a related embodiment, the present invention includes a polynucleotide that encodes a peptide of a hepcidin analogue or dimer (or monomer subunit of a dimer) of the present invention.

[0077] In a further related embodiment, the present invention includes a vector comprising a polynucleotide of the invention. In particular embodiments, the vector is an expression vector comprising a promoter operably linked to the polynucleotide, e.g., in a manner that promotes expression of the polynucleotide.

[0078] In another embodiment, the present invention includes a pharmaceutical composition, comprising a hepcidin analogue, dimer, polynucleotide, or vector of the present invention, and a pharmaceutically acceptable carrier, excipient or vehicle.

[0079] In another embodiment, the present invention provides a method of binding a ferroportin or inducing ferroportin internalization and degradation, comprising contacting the ferroportin with at least one hepcidin analogue, dimer or composition of the present invention.

[0080] In a further embodiment, the present invention includes a method for treating a disease of iron metabolism in a subject in need thereof comprising providing to the subject an effective amount of a pharmaceutical composition of the present invention. In certain embodiments, the pharmaceutical composition is provided to the subject by an oral, intravenous, peritoneal, intradermal, subcutaneous, intramuscular, intrathecal, inhalation, vaporization, nebulization, sublingual, buccal, parenteral, rectal, vaginal, or topical route of administration. In certain embodiments, the pharmaceutical composition is provided to the subject by an oral or subcutaneous route of administration. In certain embodiments, the disease of iron metabolism is an iron overload disease. In certain embodiments, the pharmaceutical composition is provided to the subject at most or about twice

daily, at most or about once daily, at most or about once every two days, at most or about once a week, or at most or about once a month.

[0081] In particular embodiments, the hepcidin analogue is provided to the subject at a dosage of about 1 mg to about 100 mg or about 1 mg to about 5 mg.

[0082] In another embodiment, the present invention provides a device comprising pharmaceutical composition of the present invention, for delivery of a hepcidin analogue or dimer of the invention to a subject, optionally orally or subcutaneously.

[0083] In yet another embodiment, the present invention includes a kit comprising a pharmaceutical composition of the invention, packaged with a reagent, a device, or an instructional material, or a combination thereof.

DETAILED DESCRIPTION OF THE INVENTION

[0084] The present invention relates generally to hepcidin analogue peptides and methods of making and using the same. In certain embodiments, the hepcidin analogues exhibit one or more hepcidin activity. In certain embodiments, the present invention relates to hepcidin peptide analogues comprising one or more peptide subunit that forms a cyclized structures through an intramolecular bond, e.g., an intramolecular disulfide bond. In particular embodiments, the cyclized structure has increased potency and selectivity as compared to non-cyclized hepcidin peptides and analogies thereof. In particular embodiments, hepcidin analogue peptides of the present invention exhibit increased half-lives, e.g., when delivered orally, as compared to hepcidin or previous hepcidin analogues.

Definitions and Nomenclature

[0085] Unless otherwise defined herein, scientific and technical terms used in this application shall have the meanings that are commonly understood by those of ordinary skill in the art. Generally, nomenclature used in connection with, and techniques of, chemistry, molecular biology, cell and cancer biology, immunology, microbiology, pharmacology, and protein and nucleic acid chemistry, described herein, are those well-known and commonly used in the art.

[0086] As used herein, the following terms have the meanings ascribed to them unless specified otherwise.

[0087] Throughout this specification, the word “comprise” or variations such as “comprises” or “comprising” will be understood to imply the inclusion of a stated integer (or components) or group of integers (or components), but not the exclusion of any other integer (or components) or group of integers (or components).

[0088] The singular forms “a,” “an,” and “the” include the plurals unless the context clearly dictates otherwise.

[0089] The term “including” is used to mean “including but not limited to.” “Including” and “including but not limited to” are used interchangeably.

[0090] The terms “patient,” “subject,” and “individual” may be used interchangeably and refer to either a human or a non-human animal. These terms include mammals such as humans, primates, livestock animals (e.g., bovines, porcines), companion animals (e.g., canines, felines) and rodents (e.g., mice and rats). The term “mammal” refers to

any mammalian species such as a human, mouse, rat, dog, cat, hamster, guinea pig, rabbit, livestock, and the like.

[0091] The term “peptide,” as used herein, refers broadly to a sequence of two or more amino acids joined together by peptide bonds. It should be understood that this term does not connote a specific length of a polymer of amino acids, nor is it intended to imply or distinguish whether the polypeptide is produced using recombinant techniques, chemical or enzymatic synthesis, or is naturally occurring.

[0092] The term “peptide analogue” or “hepcidin analogue” as used herein, refers broadly to peptide monomers and peptide dimers comprising one or more structural features and/or functional activities in common with hepcidin, or a functional region thereof. In certain embodiments, a peptide analogue includes peptides sharing substantial amino acid sequence identity with hepcidin, e.g., peptides that comprise one or more amino acid insertions, deletions, or substitutions as compared to a wild-type hepcidin, e.g., human hepcidin, amino acid sequence. In certain embodiments, a peptide analogue comprises one or more additional modification, such as, e.g., conjugation to another compound. Encompassed by the term “peptide analogue” is any peptide monomer or peptide dimer of the present invention. In certain instances, a “peptide analog” may also or alternatively be referred to herein as a “hepcidin analogue,” “hepcidin peptide analogue,” or a “hepcidin analogue peptide.”

[0093] The recitations “sequence identity,” “percent identity,” “percent homology,” or, for example, comprising a “sequence 50% identical to,” as used herein, refer to the extent that sequences are identical on a nucleotide-by-nucleotide basis or an amino acid-by-amino acid basis over a window of comparison. Thus, a “percentage of sequence identity” may be calculated by comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical nucleic acid base (e.g., A, T, C, G, I) or the identical amino acid residue (e.g., Ala, Pro, Ser, Thr, Gly, Val, Leu, Ile, Phe, Tyr, Trp, Lys, Arg, His, Asp, Glu, Asn, Gln, Cys and Met) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison (i.e., the window size), and multiplying the result by 100 to yield the percentage of sequence identity.

[0094] Calculations of sequence similarity or sequence identity between sequences (the terms are used interchangeably herein) can be performed as follows. To determine the percent identity of two amino acid sequences, or of two nucleic acid sequences, the sequences can be aligned for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). In certain embodiments, the length of a reference sequence aligned for comparison purposes is at least 30%, preferably at least 40%, more preferably at least 50%, 60%, and even more preferably at least 70%, 80%, 90%, 100% of the length of the reference sequence. The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position.

[0095] The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

[0096] The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. In some embodiments, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch, (1970, *J. Mol. Biol.* 48: 444-453) algorithm which has been incorporated into the GAP program in the GCG software package, using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another preferred embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package, using an NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. Another exemplary set of parameters includes a Blossum 62 scoring matrix with a gap penalty of 12, a gap extend penalty of 4, and a frameshift gap penalty of 5. The percent identity between two amino acid or nucleotide sequences can also be determined using the algorithm of E. Meyers and W. Miller (1989, *Cabios*, 4: 11-17) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

[0097] The peptide sequences described herein can be used as a “query sequence” to perform a search against public databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, et al., (1990, *J. Mol. Biol.* 215: 403-10). BLAST nucleotide searches can be performed with the NBLAST program, score=100, wordlength=12 to obtain nucleotide sequences homologous to nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score=50, wordlength=3 to obtain amino acid sequences homologous to protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al. (*Nucleic Acids Res.* 25:3389-3402, 1997). When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used.

[0098] The term “conservative substitution” as used herein denotes that one or more amino acids are replaced by another, biologically similar residue. Examples include substitution of amino acid residues with similar characteristics, e.g., small amino acids, acidic amino acids, polar amino acids, basic amino acids, hydrophobic amino acids and aromatic amino acids. See, for example, the table below. In some embodiments of the invention, one or more Met residues are substituted with norleucine (Nle) which is a bioisostere for Met, but which, as opposed to Met, is not readily oxidized. In some embodiments, one or more Trp residues are substituted with Phe, or one or more Phe residues are substituted with Trp, while in some embodiments, one or more Pro residues are substituted with Npc, or one or more Npc residues are substituted with Pro. Another example of a conservative substitution with a residue normally not found in endogenous, mammalian peptides and proteins is the conservative substitution of Arg or Lys with,

for example, ornithine, canavanine, aminoethylcysteine or another basic amino acid. In some embodiments, another conservative substitution is the substitution of one or more Pro residues with bhPro or Leu or D-Npc (isonipecotic acid). For further information concerning phenotypically silent substitutions in peptides and proteins, see, for example, Bowie et. al. Science 247, 1306-1310, 1990. In the scheme below, conservative substitutions of amino acids are grouped by physicochemical properties. I: neutral, hydrophilic, II: acids and amides, III: basic, IV: hydrophobic, V: aromatic, bulky amino acids.

I	II	III	IV	V
A	N	H	M	F
S	D	R	L	Y
T	E	K	I	W
P	Q		V	
G			C	

[0099] In the scheme below, conservative substitutions of amino acids are grouped by physicochemical properties. VI: neutral or hydrophobic, VII: acidic, VIII: basic, IX: polar, X: aromatic.

VI	VII	VIII	IX	X
A	E	H	M	F
L	D	R	S	Y
I		K	T	W
P			C	
G			N	
V			Q	

[0100] The term “amino acid” or “any amino acid” as used here refers to any and all amino acids, including naturally occurring amino acids (e.g., α -amino acids), unnatural amino acids, modified amino acids, and non-natural amino acids. It includes both D- and L-amino acids. Natural amino acids include those found in nature, such as, e.g., the 23 amino acids that combine into peptide chains to form the building-blocks of a vast array of proteins. These are primarily L stereoisomers, although a few D-amino acids occur in bacterial envelopes and some antibiotics. The 20 “standard,” natural amino acids are listed in the above tables. The “non-standard,” natural amino acids are pyrrolysine (found in methanogenic organisms and other eukaryotes), selenocysteine (present in many noneukaryotes as well as most eukaryotes), and N-formylmethionine (encoded by the start codon AUG in bacteria, mitochondria and chloroplasts). “Unnatural” or “non-natural” amino acids are non-proteinogenic amino acids (i.e., those not naturally encoded or found in the genetic code) that either occur naturally or are chemically synthesized. Over 140 natural amino acids are known and thousands of more combinations are possible.

Examples of “unnatural” amino acids include β -amino acids (β^3 and β^2), homo-amino acids, proline and pyruvic acid derivatives, 3-substituted alanine derivatives, glycine derivatives, ring-substituted phenylalanine and tyrosine derivatives, linear core amino acids, diamino acids, D-amino acids, and N-methyl amino acids. Unnatural or non-natural amino acids also include modified amino acids. “Modified” amino acids include amino acids (e.g., natural amino acids) that have been chemically modified to include a group, groups, or chemical moiety not naturally present on the amino acid.

[0101] As is clear to the skilled artisan, the peptide sequences disclosed herein are shown proceeding from left to right, with the left end of the sequence being the N-terminus of the peptide and the right end of the sequence being the C-terminus of the peptide. Among sequences disclosed herein are sequences incorporating a “Hy-” moiety at the amino terminus (N-terminus) of the sequence, and either an “—OH” moiety or an “—NH₂” moiety at the carboxy terminus (C-terminus) of the sequence. In such cases, and unless otherwise indicated, a “Hy-” moiety at the N-terminus of the sequence in question indicates a hydrogen atom, corresponding to the presence of a free primary or secondary amino group at the N-terminus, while an “—OH” or an “—NH₂” moiety at the C-terminus of the sequence indicates a hydroxy group or an amino group, corresponding to the presence of an amido (CONH₂) group at the C-terminus, respectively. In each sequence of the invention, a C-terminal “—OH” moiety may be substituted for a C-terminal “—NH₂” moiety, and vice-versa. It is further understood that the moiety at the amino terminus or carboxy terminus may be a bond, e.g., a covalent bond, particularly in situations where the amino terminus or carboxy terminus is bound to a linker or to another chemical moiety, e.g., a PEG moiety.

[0102] The term “NH₂,” as used herein, refers to the free amino group present at the amino terminus of a polypeptide. The term “OH,” as used herein, refers to the free carboxy group present at the carboxy terminus of a peptide. Further, the term “Ac,” as used herein, refers to Acetyl protection through acylation of the C- or N-terminus of a polypeptide.

[0103] The term “carboxy,” as used herein, refers to —CO₂H.

[0104] For the most part, the names of naturally occurring and non-naturally occurring aminoacyl residues used herein follow the naming conventions suggested by the IUPAC Commission on the Nomenclature of Organic Chemistry and the IUPAC-IUB Commission on Biochemical Nomenclature as set out in “Nomenclature of α -Amino Acids (Recommendations, 1974)” Biochemistry, 14(2), (1975). To the extent that the names and abbreviations of amino acids and aminoacyl residues employed in this specification and appended claims differ from those suggestions, they will be made clear to the reader. Some abbreviations useful in describing the invention are defined below in the following Table 1.

TABLE 1

Abbreviations of Non-Natural Amino Acids and Chemical Moieties	
Abbreviation	Definition
bh, b-h, bhomo, or b-homo	β -homo
Aad	(S)-2-aminoadipic acid
DIG	Diglycolic acid

TABLE 1-continued





Abbreviations of Non-Natural Amino Acids and Chemical Moieties	
Abbreviation	Definition
Dapa or Dap	Diaminopropionic acid
Daba or Dab	Diaminobutyric acid
Pen	Penicillamine
dPen	D-Penicillamine
dPhe, dF or (D)Phe	D-phenylalanine
Hexadecane_Amine	$\text{—NH—(CH}_2\text{)}_{15}\text{CH}_3$
Dodecyl_Amine	$\text{—NH—(CH}_2\text{)}_{11}\text{CH}_3$
Sarc or Sar	Sarcosine
Cit	Citrulline
Cav	Cavanine
dGln	D-glutamine
NMe_Ile	N-methyl-isoleucine
Nva	Norvaline
Nva_Morphlino	(S)-2-amino-5-morpholinopentanoic acid
	
Morph	(S)-morpholine-3-carboxylic acid
	
Mor_propanoic_acid	(S)-2-amino-3-morpholinopropanoic acid
	
NMe-Arg	N-Methyl-Arginine
NMe-Trp	N-Methyl-Tryptophan
NMe-Phe or NMe_Phe	N-Methyl-Phenylalanine
Ac-	Acetyl
2-Nal	2-Naphthylalanine
1-Nal	1-Naphthylalanine
Bip	Biphenylalanine
2Pal	2-Pyridylalanine
βAla or bAla	beta-Alanine
Aib	2-aminoisobutyric acid
Azt	azetidine-2-carboxylic acid
Tic	L-1,2,3,4-Tetrahydroisoquinoline-3-carboxylic acid
Phe(OMe) or Tyr(Me)	Tyrosine (4-Methyl)
N-MeLys or (Me)Lys	N-Methyl-Lysine
Lys_Me3 or Lys(Me3)	

TABLE 1-continued




Abbreviations of Non-Natural Amino Acids and Chemical Moieties	
Abbreviation	Definition
12-amino-dodecanoic acid	
Dpa or DIP dDIP or dPpa	β,β -diphenylalanine or (S)-2-amino-3,3-diphenylpropanoic acid D- β,β -diphenylalanine or (R)-2-amino-3,3-diphenylpropanoic acid
N_Ethylamine	
NMe_Glu or N-MeGlu	N-methylglutamic acid
p-Xylene	
NH ₂ CONH ₂ COOH Phe(4-F), Phe(4F), (4-F)Phe or (4F)Phe Phe(4-CF ₃), Phe(4 CF ₃), (4-CF ₃)Phe or (4CF ₃)Phe Phe(2,3,5-triF), or (2,3,5-triF)Phe Palm (Peg)n Peg2 Peg4 Peg8 Peg11 Peg12 1Peg2 or 1PEG2 1Peg2-1Peg2 or 1PEG2-1PEG2 2Peg2 or PEG2 2Peg4 or PEG4 1Peg8 or 1PEG8 2Peg8 or PEG8 1Peg11 or 1PEG11 2Peg11 or PEG11 2Peg11' or 2Peg12 or PEG12 2Peg11'_Palm or 2Peg12'_Palm or PEG12'_Palm 2Peg11'_C18_ Diacid or 2Peg12_ C18_Diacid or PEG12_C18_Diacid 2Peg11'_Ahx_ Palm or 2Peg12'_Ahx_ Palm or PEG12'_Ahx_Palm	Free Amine Amide Acid 4-Fluoro-L-Phenylalanine (4-Trifluoromethyl)-L-Phenylalanine (2,3,5-Trifluoro)-L-Phenylalanine Palmitic or Palmitoyl or C(O)-(CH ₂) ₁₄ CH ₃ -(OCH ₂ CH ₂) _n - n is 1, 2, 3, 4, etc -(OCH ₂ CH ₂) ₂ - -(OCH ₂ CH ₂) ₄ - -(OCH ₂ CH ₂) ₈ - -(OCH ₂ CH ₂) ₁₁ - -(OCH ₂ CH ₂) ₁₂ - -[C(O)-CH ₂ -(Peg) ₂ -NH]- or -[C(O)-CH ₂ -(OCH ₂ CH ₂) ₂ -NH]- -[C(O)-CH ₂ -(OCH ₂ CH ₂) ₂ -NH-C(O)-CH ₂ -(OCH ₂ CH ₂) ₂ -NH]- -[C(O)-CH ₂ -CH ₂ -(Peg) ₂ -NH]- or -[C(O)-CH ₂ -CH ₂ -(OCH ₂ CH ₂) ₂ -NH]- -[C(O)-CH ₂ -CH ₂ -(Peg) ₄ -NH]- or -[C(O)-CH ₂ -CH ₂ -(OCH ₂ CH ₂) ₄ -NH]- -[C(O)-CH ₂ -(Peg) ₈ -NH]- or -[C(O)-CH ₂ -(OCH ₂ CH ₂) ₈ -NH]- -[C(O)-CH ₂ -CH ₂ -(Peg) ₈ -NH]- or -[C(O)-CH ₂ -CH ₂ -(OCH ₂ CH ₂) ₈ -NH]- -[C(O)-CH ₂ -(Peg) ₁₁ -NH]- or -[C(O)-CH ₂ -(OCH ₂ CH ₂) ₁₁ -NH]- -[C(O)-CH ₂ -CH ₂ -(Peg) ₁₁ -NH]- or -[C(O)-CH ₂ -CH ₂ -(OCH ₂ CH ₂) ₁₁ -NH]- -[C(O)-CH ₂ -CH ₂ -(Peg) ₁₂ -NH]- or -[C(O)-CH ₂ -CH ₂ -(OCH ₂ CH ₂) ₁₂ -NH]- -[C(O)-CH ₂ -CH ₂ -(Peg) ₁₂ -NH]-C(O)-(CH ₂) ₁₄ CH ₃ or -[C(O)-CH ₂ -CH ₂ -(OCH ₂ CH ₂) ₁₂ -NH]-C(O)-(CH ₂) ₁₄ CH ₃ -[C(O)-CH ₂ -CH ₂ -(Peg) ₁₂ -NH]-C(O)-(CH ₂) ₁₆ C(O)OH or -[C(O)-CH ₂ -CH ₂ -(OCH ₂ CH ₂) ₁₂ -NH]-C(O)-(CH ₂) ₁₆ C(O)OH -[C(O)-CH ₂ -CH ₂ -(Peg) ₁₂ -NH]-C(O)-(CH ₂) ₅ -N(H)-C(O)-(CH ₂) ₁₄ CH ₃ or -[C(O)-CH ₂ -CH ₂ -(OCH ₂ CH ₂) ₁₂ -NH]-C(O)-(CH ₂) ₅ -N(H)-C(O)-(CH ₂) ₁₄ CH ₃

TABLE 1-continued

Abbreviation	Definition
Lys(2Peg11'_Palm) or Lys(PEG11'_Palm) or Lys(PEG12_Palm)	
Lys(2Peg11'_C18_Diacid) or Lys(PEG12_C18_Diacid)	
Lys(2Peg11'_Ahx_Palm) or Lys(PEG12_Ahx_Palm)	
Lys(2Peg11'_IsoGlu_Palm) or Lys(PEG12_IsoGlu_Palm)	
Lys(2Peg11'_Ahx_C18_Diacid) or Lys(PEG12_Ahx_C18_Diacid)	
Lys(2Peg11'_Ahx_IsoGlu_C18_Diacid) or Lys(PEG12_Ahx_IsoGlu_C18_Diacid)	
Lys(2Peg11'_Ahx_IsoGlu_Behenic_acid) or Lys(PEG12_Ahx_IsoGlu_Behenic_acid)	
IsoGlu_Palm	

TABLE 1-continued

Abbreviation	Definition
Lys(IsoGlu_Palm) or Z_minus	
Lys(Ahx_Palm)	
Lys(1Peg2_1Peg2_Ahx_C18_Diacid) or Lys(1PEG2_1PEG2_Ahx_C18_Diacid)	
Lys(1Peg2_1Peg2_IsoGlu_C18_Diacid) or Lys(1PEG2_1PEG2_IsoGlu_C18_Diacid)	
(D)Lys(Peg11_OMe) or (D)Lys(PEG11_OMe)	
Peg11_OMe or PEG11_OMe	$-[C(O)-CH_2-CH_2-(OCH_2CH_2)_{11}-OMe]$
Lys(Ahx_Ahx_C18_Diacid)	
Lys(Ado_C18_Diacid)	

TABLE 1-continued

Abbreviations of Non-Natural Amino Acids and Chemical Moieties	
Abbreviation	Definition
Lys(Ado_Palm)	
Lys(Ado_IsoGlu_C18_Diacid)	
Lys(2Peg11'_2Peg11'_Palm) or Lys(PEG12_PEG12_Palm)	
Lys(2Peg4_Palm) or Lys(PEG4_Palm)	
Lys(2Peg4_Ahx_Palm) or Lys(PEG4_Ahx_Palm)	
Lys(Ac)	
Lys(Ahx)	

TABLE 1-continued








Abbreviation	Definition
Lys(Ahx_PEG20K)	
Lys([Lys (2Peg11'_Palm) or Lys(Lys (PEG12_Palm) (PEG12_Palm)] ₂)	
Lys(1Peg2_Ahx_Palm) or Lys(1PEG2_Ahx_Palm)	
Lys(1Peg2_Ahx_C18_Diacid) or Lys(1PEG2_Ahx_C18_Diacid)	
Lys(2Peg8_Ahx_Palm) or Lys(PEG8_Ahx_Palm)	
Lys(2Peg8_Ahx_C18_Diacid) or Lys(PEG8_Ahx_C18_Diacid)	
Lys(1Peg2_1Peg2_Ahx_Palm) or Lys(1PEG2_1PEG2_Ahx_Palm)	

TABLE 1-continued








Abbreviations of Non-Natural Amino Acids and Chemical Moieties	
Abbreviation	Definition
Lys(2Peg11'_AlbuTag) or Lys(PEG12'_AlbuTag)	
(D)Lys_IVA	
Lys(2Peg11'_IsoGlu_C14_Diacid) or Lys(PEG12'_IsoGlu_C14_Diacid)	
Lys(2Peg11'_IsoGlu_C16_Diacid) or Lys(PEG12'_IsoGlu_C16_Diacid)	
Lys(2Peg11'_IsoGlu_C18_Diacid) or Lys(PEG12'_IsoGlu_C18_Diacid)	
Lys(2Peg11'_IsoGlu_C20_Diacid) or Lys(PEG12'_IsoGlu_C20_Diacid)	
Peg13	Bifunctional PEG linker with 13 PolyEthylene Glycol units
Peg25	Bifunctional PEG linker with 25 PolyEthylene Glycol units
Peg1K	Bifunctional PEG linker with PolyEthylene Glycol Mol wt of 1000 Da
Peg2K	Bifunctional PEG linker with PolyEthylene Glycol Mol wt of 2000 Da
Peg3.4K	Bifunctional PEG linker with PolyEthylene Glycol Mol wt of 3400 Da
Peg5K	Bifunctional PEG linker with PolyEthylene Glycol Mol wt of 5000 Da
IDA or Ida	Iminodiacetic acid
IDA-Palm	(Palmityl)-Iminodiacetic acid
hPhe	homoPhenylalanine
Ahx	Aminohexanoic acid
Isovaleric Acid	
DIG-OH	Glycolic monoacid
Triazine	Amino propyl Triazine di-acid
Boc-Triazine	Boc-Triazine di-acid
Trifluorobutyric acid	4,4,4-Trifluorobutyric acid
2-Methyltrifluorobutyric acid	2-methyl-4,4,4-Butyric acid

TABLE 1-continued





Abbreviations of Non-Natural Amino Acids and Chemical Moieties	
Abbreviation	Definition
Trifluoropentanoic acid	5,5,5-Trifluoropentanoic acid
1,4-Phenylenediacetic acid	para-Phenylenediacetic acid
1,3-Phenylenediacetic acid	meta-Phenylenediacetic acid
DTT	Dithiothreitol
βhTrp or bhTrp	β-homoTryptophane
βhPhe or bhPhe	β-homophenylalanine
Phe(4-CF ₃)	4-TrifluoromethylPhenylalanine
βGlu or bGlu	β-Glutamic acid
Asp_OMe or (OMe)Asp	L-Aspartic acid β-methyl ester
Glu_OMe or (OMe)Glu	L-Glutamic acid gamma-methyl ester
	
βhGlu or bhGlu	β-homoglutamic acid
2-2-Indane	2-Aminoindane-2-carboxylic acid
1-1-Indane	1-Aminoindane-1-carboxylic acid
hCha	homocyclohexylalanine
Cyclobutyl	Cyclobutylalanine
hLeu	Homoleucine
Gla	γ-Carboxy-glutamic acid
Glp	Pyroglutamic acid
Aep	3-(2-aminoethoxy)propanoic acid
Aea	(2-aminoethoxy)acetic acid
IsoGlu-octanoic acid	octanoyl-γ-Glu
K-octanoic acid	octanoyl-ε-Lys
Dapa(Palm)	Hexadecanoyl-β-Diaminopropionic acid
IsoGlu-Palm	hexadecanoyl-γ-Glu
C-StBu	S-tert-butylthio-cysteine
C-tBu	S-tert-butyl-cysteine
N-MeCys, (Me)Cys or NMeCys	N-methyl-cysteine
a-MeCys, aMeCys, or α-MeCys	α-methyl-cysteine
hCys or Hcy	homo-cysteine
dCys, (D)Cys or dC	D-cysteine
4R_Mcp	(2S,4R)-4-mercaptopyrrolidine-2-carboxylic acid
	
4S_Mcp	(2S,4S)-4-mercaptopyrrolidine-2-carboxylic acid
	
MOMHA	2-(3-methoxy-3-oxopropyl)-6-methylheptanoic acid
	

TABLE 1-continued





Abbreviations of Non-Natural Amino Acids and Chemical Moieties	
Abbreviation	Definition
Dapa(AcBr)	NY-(bromoacetyl)-2,3-diaminopropionic acid
Tle	tert-Leucine
Phg	phenylglycine
Oic	octahydroindole-2-carboxylic acid
Chg	α -cyclohexylglycine
GP-(Hyp)	Gly-Pro-HydroxyPro
Inp	isonipecotic acid or 
Amc	4-(aminomethyl)cyclohexane carboxylic acid
Betaine	(CH ₃) ₃ NCH ₂ CH ₂ CO ₂ H
D-Npc or D-NPC	(D)-nipecotic acid
Npc or NPC	Nipecotic acid
(D)Lys, D-Lys,	D-Lysine
dLys, or dK	
dLys_Carnitine_	
Alkyl	
Orn	Ornithine
Homoserine or hSer	homoserine
dSer or (D)Ser	D-serine
Nleu or Nle	Norleucine
α -MePro, aMePro,	α -methylproline
or α -MePro	
Hyp_3R	(2S,3R)-3-hydroxypyrrolidine-2-carboxylic acid
Hyp_3S	(2S,3S)-3-hydroxypyrrolidine-2-carboxylic acid
bhPro or bhP	beta-homoproline
1-MeHis, His_1Me,	1-Methyl-histidine
His(1-Me), or	
MeHis	
DiIsoAmylAmine_	
CH ₂ _Acid	
(Me)Glu or Glu_Me	N-Me-glutamic acid
3Pal or 3-Pal	3-pyridylalanine 
3Quin or 3-Quin	3-Quinolinyllalanine 
aMeF or a-MePhe	Alpha-methylphenylalanine
or (a-Me)Phe	
Me_Thr or	N-Me-threonine
NMe_Thr	

TABLE 1-continued








Abbreviations of Non-Natural Amino Acids and Chemical Moieties	
Abbreviation	Definition
Hyp	 hydroxyproline (all isomers)
Tet1	 (S)-2-amino-3-(2H-tetrazol-5-yl)propanoic acid
Tet2	(S)-2-amino-4-(1H-tetrazol-5-yl)butanoic acid
Z_minus	Lys(isoglu_Palm)
(Me)Ile or	N-methyl-isoleucine
(N-Me)Ile	
C18_Diacid_	
isoGlu_1Peg2_	
1Peg2 or C18_	
Diacid_isoGlu_	
1PEG2_1PEG2	
PropanoicP,	γ-aminobutyric acid (NH ₂ CH ₂ CH ₂ CH ₂ CO ₂ H)
ProtanoicPro	
or Ppa	—C(O)-alkanyl
ButanoicP,	
ButanoicPro,	—C(O)-alkenyl
or Pba	
Gaba	
or GABA	
alkanoyl	
alkenoyl	
iso Asp	
Lys(Gal) or	
Lys_Gal	
dLys_Gal	
Lys_1PEG2_	
1PEG2_Dap_	
C18_Diacid	

TABLE 1-continued










Abbreviations of Non-Natural Amino Acids and Chemical Moieties	
Abbreviation	Definition
Lys_Acrylamide	
dK_Acrylamide	
Lys_PEG11_OMe	
dLys_PEG11_OMe	
dLys_PEG8'_OMe or dLYs_PEG7_OMe)	
dLys_PEG4'_OMe or dLYs_PEG3_OMe	
PEG20K	 Compound prepared using the above reagent from SUNBRIGHT® ME-200 HS (MW~20,000)  Compound prepared using the above reagent from SUNBRIGHT® ME-300 CS (MW~30,000)
PEG40KB	 Compound prepared using the above reagent from SUNBRIGHT® GL2400GS2 (MW~40,000)

TABLE 1-continued

Abbreviations of Non-Natural Amino Acids and Chemical Moieties	
Abbreviation	Definition
NMe_Lys	N-methyl-lysine
NMe_dLys	N-methyl-D-lysine
NMe_dSer	N-methyl-D-serine
NMe_Ser	N-methyl-serine
NMe_Tyr	N-methyl-tyrosine
NMe_dTyr	N-methyl-D-tyrosine
NMe_dLeu	N-methyl-D-leucine
NMe_dPhe	N-methyl-D-phenylalanine
NMe_Gln	N-methyl-glutamine
NMe_dGln	N-methyl-D-glutamine
Trp_5OH	(S)-2-amino-3-(6-methoxy-1H-indol-3-yl)propanoic acid
Trp_6OMe	(S)-2-amino-3-(6-methoxy-1H-indol-3-yl)propanoic acid
Nicotinic_acid	Nicotinic acid
4_Fluoro-phenylacetic_acid	2-(4-fluorophenyl)acetic acid
Thr_Me or Thr(Me)	O-methyl-L-threonine
N_Butyl_Phe	
diphenylethylamine	2,2-diphenylethylamino
Amylamine	pentylamino

[0105] Throughout the present specification, unless naturally occurring amino acids are referred to by their full name (e.g. alanine, arginine, etc.), they are designated by their conventional three-letter or single-letter abbreviations (e.g. Ala or A for alanine, Arg or R for arginine, etc.). In the case of less common or non-naturally occurring amino acids, unless they are referred to by their full name (e.g., sarcosine, ornithine, etc.), frequently employed three- or four-character codes are employed for residues thereof, including, Sar or Sarc (sarcosine, i.e. N-methylglycine), Aib (α -aminoisobutyric acid), Daba (2,4-diaminobutanoic acid), Dapa (2,3-diaminopropanoic acid), γ -Glu (γ -glutamic acid), pGlu (pyroglutamic acid), Gaba (γ -aminobutanoic acid), β -Pro (pyrrolidine-3-carboxylic acid), 8Ado (8-amino-3,6-dioxaoctanoic acid), Abu (4-aminobutyric acid), bhPro (β -homo-proline), bhPhe (β -homo-L-phenylalanine), bhAsp (β -homo-aspartic acid), Dpa (β,β diphenylalanine), Ida (Iminodiacetic acid), hCys (homocysteine), bhDpa (β -homo- β,β -diphenylalanine).

[0106] Furthermore, R¹ can in all sequences be substituted with isovaleric acid or equivalent. In some embodiments, wherein a peptide of the present invention is conjugated to an acidic compound such as, e.g., isovaleric acid, isobutyric acid, valeric acid, and the like, the presence of such a conjugation is referenced in the acid form. So, for example, but not to be limited in any way, instead of indicating a conjugation of isovaleric acid to a peptide by referencing isovaleroyl, in some embodiments, the present application may reference such a conjugation as isovaleric acid.

[0107] It is understood that for each of the hepcidin analogue formulas provided herein, bonds may be indicated by a “—” or implied based on the formula and constituent (s). For example, “B7(L1Z)” is understood to include a bond between B7 and L1 if L1 is present, or between B7 and Z if L1 is absent. Similarly, “B5(L1Z)” is understood to include a bond between B5 and L1 if L1 is present, or between B5 and Z if L1 is absent. In addition, it is understood that a bond exists between L1 and Z when both are present. Accordingly, definitions of certain substituents, such as e.g., B7, L1 and J, may include “—” before and/or after the defined substituent, but in each instance, it is understood that the substituent is bonded to other substituents via a single bond. For

example, where “J” is defined as Lys, D-Lys, Arg, Pro, -Pro-Arg-, etc., it is understood that J is bound to Xaa2 and Y1 via single bonds. Thus, definitions of substituents may include or may not include “—”, but are still understood to be bonded to adjacent substituents.

[0108] The term “L-amino acid,” as used herein, refers to the “L” isomeric form of a peptide, and conversely the term “D-amino acid” refers to the “D” isomeric form of a peptide. In certain embodiments, the amino acid residues described herein are in the “L” isomeric form, however, residues in the “D” isomeric form can be substituted for any L-amino acid residue, as long as the desired functional is retained by the peptide.

[0109] Unless otherwise indicated, reference is made to the L-isomeric forms of the natural and unnatural amino acids in question possessing a chiral center. Where appropriate, the D-isomeric form of an amino acid is indicated in the conventional manner by the prefix “D” before the conventional three-letter code (e.g. Dasp, (D)Asp or D-Asp; Dphe, (D)Phe or D-Phe).

[0110] As used herein, a “lower homolog of Lys” refers to an amino acid having the structure of Lysine but with one or more fewer carbons in its side chain as compared to Lysine.

[0111] As used herein, a “higher homolog of Lys” refers to an amino acid having the structure of Lysine but with one or more additional carbon atoms in its side chain as compared to Lysine.

[0112] The term “DRP,” as used herein, refers to disulfide rich peptides.

[0113] The term “dimer,” as used herein, refers broadly to a peptide comprising two or more monomer subunits. Certain dimers comprise two DRPs. Dimers of the present invention include homodimers and heterodimers. A monomer subunit of a dimer may be linked at its C- or N-terminus, or it may be linked via internal amino acid residues. Each monomer subunit of a dimer may be linked through the same site, or each may be linked through a different site (e.g., C-terminus, N-terminus, or internal site).

[0114] The term “isostere replacement” or “isostere substitution” are used interchangeably herein to refer to any amino acid or other analog moiety having chemical and/or structural properties similar to a specified amino acid. In

certain embodiments, an isostere replacement is a conservative substitution with a natural or unnatural amino acid.

[0115] The term “cyclized,” as used herein, refers to a reaction in which one part of a polypeptide molecule becomes linked to another part of the polypeptide molecule to form a closed ring, such as by forming a disulfide bridge or other similar bond.

[0116] The term “subunit,” as used herein, refers to one of a pair of polypeptide monomers that are joined to form a dimer peptide composition.

[0117] The term “linker moiety,” as used herein, refers broadly to a chemical structure that is capable of linking or joining together two peptide monomer subunits to form a dimer.

[0118] The term “linking group,” as used herein, refers broadly to a chemical structure that is capable of covalently linking or joining together two other molecular fragments.

[0119] The term “solvate” in the context of the present invention refers to a complex of defined stoichiometry formed between a solute (e.g., a hepcidin analogue or pharmaceutically acceptable salt thereof according to the invention) and a solvent. The solvent in this connection may, for example, be water, ethanol or another pharmaceutically acceptable, typically small-molecular organic species, such as, but not limited to, acetic acid or lactic acid. When the solvent in question is water, such a solvate is normally referred to as a hydrate.

[0120] As used herein, a “disease of iron metabolism” includes diseases where aberrant iron metabolism directly causes the disease, or where iron blood levels are dysregulated causing disease, or where iron dysregulation is a consequence of another disease, or where diseases can be treated by modulating iron levels, and the like. More specifically, a disease of iron metabolism according to this disclosure includes iron overload diseases, iron deficiency disorders, disorders of iron biodistribution, other disorders of iron metabolism and other disorders potentially related to iron metabolism, etc. Diseases of iron metabolism include hemochromatosis, HFE mutation hemochromatosis, ferroportin mutation hemochromatosis, transferrin receptor 2 mutation hemochromatosis, hemojuvelin mutation hemochromatosis, hepcidin mutation hemochromatosis, juvenile hemochromatosis, neonatal hemochromatosis, hepcidin deficiency, transfusional iron overload, thalassemia, thalassemia intermedia, alpha thalassemia, sideroblastic anemia, porphyria, porphyria cutanea tarda, African iron overload, hyperferritinemia, ceruloplasmin deficiency, atransferrinemia, congenital dyserythropoietic anemia, hypochromic microcytic anemia, sickle cell disease, polycythemia vera (primary and secondary), secondary erythrocytoses, such as Chronic obstructive pulmonary disease (COPD), post-renal transplant, Chuvash, HIF and PHD mutations, and idiopathic, myelodysplasia, pyruvate kinase deficiency, iron deficiency of obesity, other anemias, benign or malignant tumors that overproduce hepcidin or induce its overproduction, conditions with hepcidin excess, Friedreich ataxia, gracile syndrome, Hallervorden-Spatz disease, Wilson’s disease, pulmonary hemosiderosis, hepatocellular carcinoma, cancer, hepatitis, cirrhosis of liver, pica, chronic renal failure, insulin resistance, diabetes, atherosclerosis, neurodegenerative disorders, multiple sclerosis, Parkinson’s disease, Huntington’s disease, and Alzheimer’s disease.

[0121] In some embodiments, the disease and disorders are related to iron overload diseases such as iron hemochro-

matosis, HFE mutation hemochromatosis, ferroportin mutation hemochromatosis, transferrin receptor 2 mutation hemochromatosis, hemojuvelin mutation hemochromatosis, hepcidin mutation hemochromatosis, juvenile hemochromatosis, neonatal hemochromatosis, hepcidin deficiency, transfusional iron overload, thalassemia, thalassemia intermedia, alpha thalassemia, sickle cell disease, myelodysplasia, sideroblastic infections, diabetic retinopathy, and pyruvate kinase deficiency.

[0122] In some embodiments, the hepcidin analogues of the invention are used to treat diseases and disorders that are not typically identified as being iron related. For example, hepcidin is highly expressed in the murine pancreas suggesting that diabetes (Type I or Type II), insulin resistance, glucose intolerance and other disorders may be ameliorated by treating underlying iron metabolism disorders. See Ilyin, G. et al. (2003) FEBS Lett. 542 22-26, which is herein incorporated by reference. As such, peptides of the invention may be used to treat these diseases and conditions. Those skilled in the art are readily able to determine whether a given disease can be treated with a peptide according to the present invention using methods known in the art, including the assays of WO 2004092405, which is herein incorporated by reference, and assays which monitor hepcidin, hemojuvelin, or iron levels and expression, which are known in the art such as those described in U.S. Pat. No. 7,534,764, which is herein incorporated by reference.

[0123] In certain embodiments of the present invention, the diseases of iron metabolism are iron overload diseases, which include hereditary hemochromatosis, iron-loading anemias, alcoholic liver diseases and chronic hepatitis C.

[0124] The term “pharmaceutically acceptable salt,” as used herein, represents salts or zwitterionic forms of the peptides or compounds of the present invention which are water or oil-soluble or dispersible, which are suitable for treatment of diseases without undue toxicity, irritation, and allergic response; which are commensurate with a reasonable benefit/risk ratio, and which are effective for their intended use. The salts can be prepared during the final isolation and purification of the compounds or separately by reacting an amino group with a suitable acid. Representative acid addition salts include acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, glycerophosphate, hemisulfate, heptanoate, hexanoate, formate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethansulfonate (isethionate), lactate, maleate, mesitylenesulfonate, methanesulfonate, naphthylenesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, trichloroacetate, trifluoroacetate, phosphate, glutamate, bicarbonate, para-toluenesulfonate, and undecanoate. Also, amino groups in the compounds of the present invention can be quaternized with methyl, ethyl, propyl, and butyl chlorides, bromides, and iodides; dimethyl, diethyl, dibutyl, and diamyl sulfates; decyl, lauryl, myristyl, and steryl chlorides, bromides, and iodides; and benzyl and phenethyl bromides. Examples of acids which can be employed to form therapeutically acceptable addition salts include inorganic acids such as hydrochloric, hydrobromic, sulfuric, and phosphoric, and organic acids such as oxalic, maleic, succinic, and citric. A pharmaceutically acceptable salt may suitably be a salt chosen, e.g., among acid addition salts and basic salts. Examples of acid addition

salts include chloride salts, citrate salts and acetate salts. Examples of basic salts include salts where the cation is selected among alkali metal cations, such as sodium or potassium ions, alkaline earth metal cations, such as calcium or magnesium ions, as well as substituted ammonium ions, such as ions of the type $N(R_1)(R_2)(R_3)(R_4)^+$, where R_1 , R_2 , R_3 and R_4 independently will typically designate hydrogen, optionally substituted C1-6-alkyl or optionally substituted C2-6-alkenyl. Examples of relevant C1-6-alkyl groups include methyl, ethyl, 1-propyl and 2-propyl groups. Examples of C2-6-alkenyl groups of possible relevance include ethenyl, 1-propenyl and 2-propenyl. Other examples of pharmaceutically acceptable salts are described in "Remington's Pharmaceutical Sciences", 17th edition, Alfonso R. Gennaro (Ed.), Mark Publishing Company, Easton, PA, USA, 1985 (and more recent editions thereof), in the "Encyclopaedia of Pharmaceutical Technology", 3rd edition, James Swarbrick (Ed.), Informa Healthcare USA (Inc.), NY, USA, 2007, and in J. Pharm. Sci. 66: 2 (1977). Also, for a review on suitable salts, see *Handbook of Pharmaceutical Salts: Properties, Selection, and Use* by Stahl and Wermuth (Wiley-VCH, 2002). Other suitable base salts are formed from bases which form non-toxic salts. Representative examples include the aluminum, arginine, benzathine, calcium, choline, diethylamine, diolamine, glycine, lysine, magnesium, meglumine, olamine, potassium, sodium, tromethamine, and zinc salts. Hemisalts of acids and bases may also be formed, e.g., hemisulphate and hemicalcium salts.

[0125] The term "N(alpha)Methylation", as used herein, describes the methylation of the alpha amine of an amino acid, also generally termed as an N-methylation.

[0126] The term "sym methylation" or "Arg-Me-sym", as used herein, describes the symmetrical methylation of the two nitrogens of the guanidine group of arginine. Further, the term "asym methylation" or "Arg-Me-asym" describes the methylation of a single nitrogen of the guanidine group of arginine.

[0127] The term "acylating organic compounds", as used herein refers to various compounds with carboxylic acid functionality that are used to acylate the N-terminus of an amino acid subunit prior to forming a C-terminal dimer. Non-limiting examples of acylating organic compounds include cyclopropylacetic acid, 4-Fluorobenzoic acid, 4-fluorophenylacetic acid, 3-Phenylpropionic acid, Succinic acid, Glutaric acid, Cyclopentane carboxylic acid, 3,3,3-trifluoropropionic acid, 3-Fluoromethylbutyric acid, Tetrahydro-2H-Pyran-4-carboxylic acid.

[0128] The term " C_{n-m} " or " C_n-C_m " indicates a range which includes the endpoints, wherein n and m are integers and indicate the number of carbons. Examples include C_{1-4} , C_{1-6} , C_{1-20} , C_1-C_6 , C_1-C_8 and the like

[0129] The term "alkyl" includes a straight chain or branched, noncyclic or cyclic, saturated aliphatic hydrocarbon containing from 1 to 24 carbon atoms. Representative saturated straight chain alkyls include, but are not limited to, methyl, ethyl, n-propyl, n-butyl, n-pentyl, n-hexyl, and the like, while saturated branched alkyls include, without limitation, isopropyl, sec-butyl, isobutyl, tert-butyl, isopentyl, and the like. Representative saturated cyclic alkyls include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and the like, while unsaturated cyclic alkyls include, without limitation, cyclopentenyl, cyclohexenyl, and the like.

[0130] The term "alkylene" refers to a divalent alkyl group, particularly having from 1 to 24 carbon atoms. The term is exemplified by groups such as methylene ($-\text{CH}_2-$), ethylene ($-\text{CH}_2\text{CH}_2-$), the propylene isomers (e.g., $-\text{CH}_2\text{CH}_2\text{CH}_2-$ and $-\text{CH}(\text{CH}_3)\text{CH}_2-$) and the like.

[0131] The term "alkoxy" refers to the group "alkyl-O—". Examples of alkoxy groups include, e.g., methoxy, ethoxy, n-propoxy, iso-propoxy, n-butoxy, tert-butoxy, sec-butoxy, n-pentoxy, n-hexoxy and 1,2-dimethylbutoxy.

[0132] The term "halo" refers to atoms occupying group VIIA of the periodic table, such as fluoro, chloro, bromo or iodo.

[0133] The term "haloalkyl" refers to an unbranched or branched alkyl group as defined above, wherein one or more (e.g., 1 to 6 or 1 to 3) hydrogen atoms are replaced by a halogen. For example, where a residue is substituted with more than one halogen, it may be referred to by using a prefix corresponding to the number of halogen moieties attached. Dihaloalkyl and trihaloalkyl refer to alkyl substituted with two ("di") or three ("tri") halo groups, which may be, but are not necessarily, the same halogen. Examples of haloalkyl include, e.g., trifluoromethyl, difluoromethyl, fluoromethyl, trichloromethyl, 2,2,2-trifluoroethyl, 1,2-difluoroethyl, 3-bromo-2-fluoropropyl, 1,2-dibromoethyl and the like.

[0134] The term "haloalkoxy" refers to a radical $-\text{OR}$ where R is haloalkyl group as defined above e.g., trifluoromethoxy, 2,2,2-trifluoroethoxy, difluoromethoxy, and the like.

[0135] The term "alkanoyl" means an alkyl-C(O)— group, wherein the alkyl group is as defined herein. Representative alkanoyl groups include methanoyl, ethanoyl, 3-methylbutanoyl, $\text{CH}_3(\text{CH}_2)_{14}\text{C(O)}-$, and the like.

[0136] As used herein, "Amide" means $-\text{CONH}-$ or $-\text{NHC(O)}-$ linkage.

[0137] As used herein, the term "thio" or "mercapto" means an $-\text{SH}$ group.

[0138] As used herein, the term "N-substituted" means the a-amino group of the amino acid is substituted. In some embodiments, the N-substituted means the a-amino group of the amino acid is substituted with 1 or 2 C_{1-6} alkyl groups, such as methyl, ethyl or propyl, and the like.

[0139] As used herein "Aryl" by itself or as part of another substituent refers to a polyunsaturated, aromatic, hydrocarbon group containing from 6 to 14 carbon atoms, or 6 to 10 carbon atoms, which can be a single ring or multiple rings (up to three rings) which are fused together or linked covalently. Thus the phrase includes, but is not limited to, groups such as phenyl, biphenyl, anthracenyl, naphthyl by way of example. Non-limiting examples of unsubstituted aryl groups include phenyl, 1-naphthyl, 2-naphthyl and 4-biphenyl.

[0140] As used herein, "functional group" on the side chain of an amino acid means $-\text{COOH}$, $-\text{NH}_2$, $-\text{NH}-$, $-\text{SH}$, $-\text{SCH}_3$, $-\text{OH}$, $-\text{C(O)NH}_2$, guanidiny, imidazoyl, pyrrolidiny, phenyl, indolyl, and the like. In some embodiments, the functional group include $-\text{COOH}$, $-\text{NH}_2$, OH , SH , guanidiny, and the like.

[0141] As used herein, a "therapeutically effective amount" of the peptide agonists of the invention is meant to describe a sufficient amount of the peptide agonist to treat an hepcidin-related disease, including but not limited to any of the diseases and disorders described herein (for example, a

disease of iron metabolism). In particular embodiments, the therapeutically effective amount will achieve a desired benefit/risk ratio applicable to any medical treatment.

Peptide Analogues of Hepcidin

[0142] The present invention provides peptide analogues of hepcidin, which may be monomers or dimers (collectively “hepcidin analogues”).

[0143] In some embodiments, a hepcidin analogue of the present invention binds ferroportin, e.g., human ferroportin. In certain embodiments, hepcidin analogues of the present invention specifically bind human ferroportin. As used herein, “specifically binds” refers to a specific binding agent’s preferential interaction with a given ligand over other agents in a sample. For example, a specific binding agent that specifically binds a given ligand, binds the given ligand, under suitable conditions, in an amount or a degree that is observable over that of any nonspecific interaction with other components in the sample. Suitable conditions are those that allow interaction between a given specific binding agent and a given ligand. These conditions include pH, temperature, concentration, solvent, time of incubation, and the like, and may differ among given specific binding agent and ligand pairs, but may be readily determined by those skilled in the art. In some embodiments, a hepcidin analogue of the present invention binds ferroportin with greater specificity than a hepcidin reference compound (e.g., any one of the hepcidin reference compounds provided herein). In some embodiments, a hepcidin analogue of the present invention exhibits ferroportin specificity that is at least about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 200%, 300%, 400%, 500%, 700%, 1000%, or 10,000% higher than a hepcidin reference compound (e.g., any one of the hepcidin reference compounds provided herein). In some embodiments, a hepcidin analogue of the present invention exhibits ferroportin specificity that is at least about 5 fold, or at least about 10, 20, 50, or 100 fold higher than a hepcidin reference compound (e.g., any one of the hepcidin reference compounds provided herein).

[0144] In certain embodiments, a hepcidin analogue of the present invention exhibits a hepcidin activity. In some embodiments, the activity is an in vitro or an in vivo activity, e.g., an in vivo or an in vitro activity described herein. In some embodiments, a hepcidin analogue of the present invention exhibits at least about 50%, 60%, 70%, 80%, 90%, 95%, 97%, 98%, 99%, or greater than 99% of the activity exhibited by a hepcidin reference compound (e.g., any one of the hepcidin reference compounds provided herein).

[0145] In some embodiments, a hepcidin analogue of the present invention exhibits at least about 50%, 60%, 70%, 80%, 90%, 95%, 97%, 98%, 99%, or greater than 99% of the ferroportin binding ability that is exhibited by a hepcidin reference compound. In some embodiments, a hepcidin analogue of the present invention has a lower EC₅₀ or IC₅₀ (i.e., higher binding affinity) for binding to ferroportin, (e.g., human ferroportin) compared to a hepcidin reference compound. In some embodiments, a hepcidin analogue of the present invention has an EC₅₀ or IC₅₀ in a ferroportin competitive binding assay that is at least about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 200%, 300%, 400%, 500%, 700%, or 1000% lower than a hepcidin reference compound.

[0146] In certain embodiments, a hepcidin analogue of the present invention exhibits increased hepcidin activity as

compared to a hepcidin reference compound. In some embodiments, the activity is an in vitro or an in vivo activity, e.g., an in vivo or an in vitro activity described herein. In certain embodiments, the hepcidin analogue of the present invention exhibits 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 60, 70, 80, 90, 100, 120, 140, 160, 180, or 200-fold greater hepcidin activity than a hepcidin reference compound. In certain embodiments, the hepcidin analogue of the present invention exhibits at least about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 97%, 98%, 99% or greater than 99%, 100%, 200% 300%, 400%, 500%, 700%, or 1000% greater activity than a hepcidin reference compound.

[0147] In some embodiments, a peptide analogue of the present invention exhibits at least about 50%, 60%, 70%, 80%, 90%, 95%, 97%, 98%, 99%, or greater than 99%, 100%, 200% 300%, 400%, 500%, 700%, or 1000% greater in vitro activity for inducing the degradation of human ferroportin protein as that of a hepcidin reference compound, wherein the activity is measured according to a method described herein.

[0148] In some embodiments, a peptide or a peptide dimer of the present invention exhibits at least about 50%, 60%, 70%, 80%, 90%, 95%, 97%, 98%, 99%, or greater than 99%, 100%, 200% 300%, 400%, 500%, 700%, or 1000% greater in vivo activity for inducing the reduction of free plasma iron in an individual as does a hepcidin reference compound, wherein the activity is measured according to a method described herein.

[0149] In some embodiments, the activity is an in vitro or an in vivo activity, e.g., an in vivo or an in vitro activity described herein. In certain embodiments, a hepcidin analogue of the present invention exhibits 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 60, 70, 80, 90, 100, 120, 140, 160, 180, or 200-fold greater or at least about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 200%, 300%, 400%, 500%, 700%, or 1000% greater activity than a hepcidin reference compound, wherein the activity is an in vitro activity for inducing the degradation of ferroportin, e.g., as measured according to the Examples herein; or wherein the activity is an in vivo activity for reducing free plasma iron, e.g., as measured according to the Examples herein.

[0150] In some embodiments, the hepcidin analogues of the present invention mimic the hepcidin activity of Hep25, the bioactive human 25-amino acid form, are herein referred to as “mini-hepcidins”. As used herein, in certain embodiments, a compound (e.g., a hepcidin analogue) having a “hepcidin activity” means that the compound has the ability to lower plasma iron concentrations in subjects (e.g. mice or humans), when administered thereto (e.g. parenterally injected or orally administered), in a dose-dependent and time-dependent manner. See e.g. as demonstrated in Rivera et al. (2005), Blood 106:2196-9. In some embodiments, the peptides of the present invention lower the plasma iron concentration in a subject by at least about 1.2, 1.5, 2, 3, 4, 5, 6, 7, 8, 9, or 10-fold, or at least about 5%, 10%, 20%, 25%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or about 99%.

[0151] In some embodiments, the hepcidin analogues of the present invention have in vitro activity as assayed by the ability to cause the internalization and degradation of ferroportin in a ferroportin-expressing cell line as taught in Nemeth et al. (2006) Blood 107:328-33. In some embodiments, in vitro activity is measured by the dose-dependent

loss of fluorescence of cells engineered to display ferroportin fused to green fluorescent protein as in Nemeth et al. (2006) Blood 107:328-33. Aliquots of cells are incubated for 24 hours with graded concentrations of a reference preparation of Hep25 or a mini-hepcidin. As provided herein, the EC_{50} values are provided as the concentration of a given compound (e.g. a hepcidin analogue peptide or peptide dimer of the present invention) that elicits 50% of the maximal loss of fluorescence generated by a reference compound. The EC_{50} of the Hep25 preparations in this assay range from 5 to 15 nM and in certain embodiments, preferred hepcidin analogues of the present invention have EC_{50} values in in vitro activity assays of about 1,000 nM or less. In certain embodiments, a hepcidin analogue of the present invention has an EC_{50} in an in vitro activity assay (e.g., as described in Nemeth et al. (2006) Blood 107:328-33 or the Example herein) of less than about any one of 0.01, 0.05, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, 200 or 500 nM. In some embodiments, a hepcidin analogue or biotherapeutic composition (e.g., any one of the pharmaceutical compositions described herein) has an EC_{50} or IC_{50} value of about 1 nM or less.

[0152] Other methods known in the art for calculating the hepcidin activity and in vitro activity of the hepcidin analogues according to the present invention may be used. For example, in certain embodiments, the in vitro activity of the hepcidin analogues or the reference peptides is measured by their ability to internalize cellular ferroportin, which is determined by immunohistochemistry or flow cytometry using antibodies which recognizes extracellular epitopes of ferroportin. Alternatively, in certain embodiments, the in vitro activity of the hepcidin analogues or the reference peptides is measured by their dose-dependent ability to inhibit the efflux of iron from ferroportin-expressing cells that are preloaded with radioisotopes or stable isotopes of iron, as in Nemeth et al. (2006) Blood 107:328-33.

[0153] In some embodiments, the hepcidin analogues of the present invention exhibit increased stability (e.g., as measured by half-life, rate of protein degradation) as compared to a hepcidin reference compound. In certain embodiments, the stability of a hepcidin analogue of the present invention is increased at least about 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 60, 70, 80, 90, 100, 120, 140, 160, 180, or 200-fold greater or at least about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 200%, 300%, 400%, or 500% greater than a hepcidin reference compound. In some embodiments, the stability is a stability that is described herein. In some embodiments, the stability is a plasma stability, e.g., as optionally measured according to the method described herein. In some embodiments, the stability is stability when delivered orally.

[0154] In particular embodiments, a hepcidin analogue of the present invention exhibits a longer half-life than a hepcidin reference compound. In particular embodiments, a hepcidin analogue of the present invention has a half-life under a given set of conditions (e.g., temperature, pH) of at least about 5 minutes, at least about 10 minutes, at least about 20 minutes, at least about 30 minutes, at least about 45 minutes, at least about 1 hour, at least about 2 hour, at least about 3 hours, at least about 4 hours, at least about 5 hours, at least about 6 hours, at least about 12 hours, at least about 18 hours, at least about 1 day, at least about 2 days, at least

about 4 days, at least about 7 days, at least about 10 days, at least about two weeks, at least about three weeks, at least about 1 month, at least about 2 months, at least about 3 months, or more, or any intervening half-life or range in between, about 5 minutes, about 10 minutes, about 20 minutes, about 30 minutes, about 45 minutes, about 1 hour, about 2 hour, about 3 hours, about 4 hours, about 5 hours, about 6 hours, about 12 hours, about 18 hours, about 1 day, about 2 days, about 4 days, about 7 days, about 10 days, about two weeks, about three weeks, about 1 month, about 2 months, about 3 months, or more, or any intervening half-life or range in between. In some embodiments, the half-life of a hepcidin analogue of the present invention is extended due to its conjugation to one or more lipophilic substituent or half-life extension moiety, e.g., any of the lipophilic substituents or half-life extension moieties disclosed herein. In some embodiments, the half-life of a hepcidin analogue of the present invention is extended due to its conjugation to one or more polymeric moieties, e.g., any of the polymeric moieties or half-life extension moieties disclosed herein. In certain embodiments, a hepcidin analogue of the present invention has a half-life as described above under the given set of conditions wherein the temperature is about 25° C., about 4° C., or about 37° C., and the pH is a physiological pH, or a pH about 7.4.

[0155] In certain embodiments, a hepcidin analogue of the present invention, comprising a conjugated half-life extension moiety, has an increased serum half-life following oral, intravenous or subcutaneous administration as compared to the same analogue but lacking the conjugated half-life extension moiety. In particular embodiments, the serum half-life of a hepcidin analogue of the present invention following any of oral, intravenous or subcutaneous administration is at least 12 hours, at least 24 hours, at least 30 hours, at least 36 hours, at least 48 hours, at least 72 hours or at least 168 h. In particular embodiments, it is between 12 and 168 hours, between 24 and 168 hours, between 36 and 168 hours, or between 48 and 168 hours.

[0156] In certain embodiments, a hepcidin analogue of the present invention, e.g., a hepcidin analogue comprising a conjugated half-life extension moiety, results in decreased concentration of serum iron following oral, intravenous or subcutaneous administration to a subject. In particular embodiments, the subject's serum iron concentration is decreased to less than 10%, less than 20%, less than 25%, less than 30%, less than 40%, less than 50%, less than 60%, less than 70%, less than 80%, or less than 90% of the serum iron concentration in the absence of administration of the hepcidin analogue to the subject. In particular embodiments, the decreased serum iron concentration remains for at least 1 hour, at least 4 hours, at least 10 hours, at least 12 hours, at least 24 hours, at least 36 hours, at least 48 hours, or at least 72 hours following administration to the subject. In particular embodiments, it remains for between 12 and 168 hours, between 24 and 168 hours, between 36 and 168 hours, or between 48 and 168 hours. In one embodiment, the serum iron concentration of the subject is reduced to less than 20% at about 4 hours or about 10 hours following administration to the subject, e.g., intravenously, orally, or subcutaneously. In one embodiment, the serum iron concentration of the subject is reduced to less than 50% or less than 60% for about 24 to about 30 hours following administration, e.g., intravenously, orally, or subcutaneously.

[0157] In some embodiments, the half-life is measured in vitro using any suitable method known in the art, e.g., in some embodiments, the stability of a hepcidin analogue of the present invention is determined by incubating the hepcidin analogue with pre-warmed human serum (Sigma) at 37° C. Samples are taken at various time points, typically up to 24 hours, and the stability of the sample is analyzed by separating the hepcidin analogue from the serum proteins and then analyzing for the presence of the hepcidin analogue of interest using LC-MS.

[0158] In some embodiments, the stability of the hepcidin analogue is measured in vivo using any suitable method known in the art, e.g., in some embodiments, the stability of a hepcidin analogue is determined in vivo by administering the peptide or peptide dimer to a subject such as a human or any mammal (e.g., mouse) and then samples are taken from the subject via blood draw at various time points, typically up to 24 hours. Samples are then analyzed as described above in regard to the in vitro method of measuring half-life. In some embodiments, in vivo stability of a hepcidin analogue of the present invention is determined via the method disclosed in the Examples herein.

[0159] In some embodiments, the present invention provides a hepcidin analogue as described herein, wherein the hepcidin analogue exhibits improved solubility or improved aggregation characteristics as compared to a hepcidin reference compound. Solubility may be determined via any suitable method known in the art. In some embodiments, suitable methods known in the art for determining solubility include incubating peptides (e.g., a hepcidin analogue of the present invention) in various buffers (Acetate pH4.0, Acetate pH5.0, Phos/Citrate pH5.0, Phos Citrate pH6.0, Phos pH 6.0, Phos pH 7.0, Phos pH7.5, Strong PBS pH 7.5, Tris pH7.5, Tris pH 8.0, Glycine pH 9.0, Water, Acetic acid (pH 5.0 and other known in the art) and testing for aggregation or solubility using standard techniques. These include, but are not limited to, visual precipitation, dynamic light scattering, Circular Dichroism and fluorescent dyes to measure surface hydrophobicity, and detect aggregation or fibrillation, for example. In some embodiments, improved solubility means the peptide (e.g., the hepcidin analogue of the present invention) is more soluble in a given liquid than is a hepcidin reference compound.

[0160] In certain embodiments, the present invention provides a hepcidin analogue as described herein, wherein the hepcidin analogue exhibits a solubility that is increased at least about 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 60, 70, 80, 90, 100, 120, 140, 160, 180, or 200-fold greater or at least about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 200%, 300%, 400%, or 500% greater than a hepcidin reference compound in a particular solution or buffer, e.g., in water or in a buffer known in the art or disclosed herein.

[0161] In certain embodiments, the present invention provides a hepcidin analogue as described herein, wherein the hepcidin analogue exhibits decreased aggregation, wherein the aggregation of the peptide in a solution is at least about 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 60, 70, 80, 90, 100, 120, 140, 160, 180, or 200-fold less or at least about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 200%, 300%, 400%, or 500% less than a hepcidin reference compound in a particular solution or buffer, e.g., in water or in a buffer known in the art or disclosed herein.

[0162] In some embodiments, the present invention provides a hepcidin analogue, as described herein, wherein the hepcidin analogue exhibits less degradation (i.e., more degradation stability), e.g., greater than or about 10% less, greater than or about 20% less, greater than or about 30% less, greater than or about 40 less, or greater than or about 50% less than a hepcidin reference compound. In some embodiments, degradation stability is determined via any suitable method known in the art. In some embodiments, suitable methods known in the art for determining degradation stability include the method described in Hawe et al J Pharm Sci, VOL. 101, NO. 3, 2012, p 895-913, incorporated herein in its entirety. Such methods are in some embodiments used to select potent sequences with enhanced shelf lives.

[0163] In some embodiments, the hepcidin analogue of the present invention is synthetically manufactured. In other embodiments, the hepcidin analogue of the present invention is recombinantly manufactured.

[0164] The various hepcidin analogue monomer and dimer peptides of the invention may be constructed solely of natural amino acids. Alternatively, these hepcidin analogues may include unnatural or non-natural amino acids including, but not limited to, modified amino acids. In certain embodiments, modified amino acids include natural amino acids that have been chemically modified to include a group, groups, or chemical moiety not naturally present on the amino acid. The hepcidin analogues of the invention may additionally include D-amino acids. Still further, the hepcidin analogue peptide monomers and dimers of the invention may include amino acid analogs. In particular embodiments, a peptide analogue of the present invention comprises any of those described herein, wherein one or more natural amino acid residues of the peptide analogue is substituted with an unnatural or non-natural amino acid, or a D-amino acid.

[0165] In certain embodiments, the hepcidin analogues of the present invention include one or more modified or unnatural amino acids. For example, in certain embodiments, a hepcidin analogue includes one or more of Daba, Dapa, Pen, Sar, Cit, Pba, Cav, HLeu, 2-Nal, 1-Nal, d-1-Nal, d-2-Nal, Bip, Phe(4-OMe), Tyr(4-OMe), β hTrp, β hPhe, Phe(4-CF₃), 2-2-Indane, 1-1-Indane, Cyclobutyl, β hPhe, hLeu, Glu, Phe(4-NH₂), hPhe, 1-Nal, Nle, 3-3-diPhe, cyclobutyl-Ala, Cha, Bip, β -Glu, Phe(4-Guan), homo amino acids, D-amino acids, and various N-methylated amino acids. One having skill in the art will appreciate that other modified or unnatural amino acids, and various other substitutions of natural amino acids with modified or unnatural amino acids, may be made to achieve similar desired results, and that such substitutions are within the teaching and spirit of the present invention.

[0166] The present invention includes any of the hepcidin analogues described herein, e.g., in a free or a salt form.

[0167] Compounds described herein include isotopically-labeled compounds, which are identical to those recited in the various formulas and structures presented herein, but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into the present compounds include isotopes of hydrogen, carbon, nitrogen, oxygen, fluorine and chlorine, such as ²H, ³H, ¹³C, ¹⁴C, ¹⁵N, ¹⁸O, ¹⁷O, ³⁵S, ¹⁸F, ³⁶Cl, respectively. Certain isotopically-labeled compounds described herein, for

example those into which radioactive isotopes such as ^3H and ^{14}C are incorporated, are useful in drug and/or substrate tissue distribution assays. Further, substitution with isotopes such as deuterium, i.e., ^2H , can afford certain therapeutic advantages resulting from greater metabolic stability, for example increased in vivo half-life or reduced dosage requirements. In particular embodiments, the compounds are isotopically substituted with deuterium. In more particular embodiments, the most labile hydrogens are substituted with deuterium.

[0168] The hepcidin analogues of the present invention include any of the peptide monomers or dimers described herein linked to a linker moiety, including any of the specific linker moieties described herein.

[0169] The hepcidin analogues of the present invention include peptides, e.g., monomers or dimers, comprising a peptide monomer subunit having at least 85%, at least 90%, at least 92%, at least 94%, at least 95%, at least 98%, or at least 99% amino acid sequence identity to a hepcidin analogue peptide sequence described herein (e.g., any one of the peptides disclosed herein), including but not limited to any of the amino acid sequences shown in Tables 2 and 3.


[0170] In certain embodiments, a peptide analogue of the present invention, or a monomer subunit of a dimer peptide analogue of the present invention, comprises or consists of 7 to 35 amino acid residues, 8 to 35 amino acid residues, 9 to 35 amino acid residues, 10 to 35 amino acid residues, 7 to 25 amino acid residues, 8 to 25 amino acid residues, 9 to 25 amino acid residues, 10 to 25 amino acid residues, 7 to 18 amino acid residues, 8 to 18 amino acid residues, 9 to 18 amino acid residues, or 10 to 18 amino acid residues, and, optionally, one or more additional non-amino acid moieties, such as a conjugated chemical moiety, e.g., a half-life extension moiety, a PEG, or a linker moiety. In particular embodiments, a monomer subunit of a hepcidin analogue comprises or consists of 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, or 35 amino acid residues. In particular embodiments, a monomer subunit of a hepcidin analogue of the present invention comprises or consists of 10 to 18 amino acid residues and, optionally, one or more additional non-amino acid moieties, such as a conjugated chemical moiety, e.g., a PEG or linker moiety. In various embodiments, the monomer subunit comprises or consists of 7 to 35 amino acid residues, 9 to 18 amino acid residues, or 10 to 18 amino acid residues. In particular embodiments of any of the various Formulas described herein, X comprises or consists of 7 to 35 amino acid residues, 8 to 35 amino acid residues, 9 to 35 amino acid residues, 10 to 35 amino acid residues, 7 to 25 amino acid residues, 8 to 25 amino acid residues, 9 to 25 amino acid residues, 10 to 25 amino acid residues, 7 to 18 amino acid residues, 8 to 18 amino acid residues, 9 to 18 amino acid residues, or 10 to 18 amino acid residues.

[0171] In particular embodiments, a hepcidin analogue or dimer of the present invention does not include any of the compounds described in PCT/US2014/030352 or PCT/US2015/038370.

Peptide Hepcidin Analogues

[0172] In certain embodiments, hepcidin analogues of the present invention comprise a single peptide subunit, optionally conjugated to an acid moiety. In certain embodiments, the acid moiety is conjugated directly or via a linker.

[0173] In one aspect, the present invention provides a hepcidin analogue comprising a peptide comprising an amino acid sequence of Formula (I'):



or a pharmaceutically acceptable salt or solvate thereof, wherein:

[0174] R^1 is hydrogen, $\text{C}_1\text{—C}_6$ alkyl, $\text{C}_6\text{—C}_{12}$ aryl, $\text{C}_6\text{—C}_{12}$ aryl- $\text{C}_1\text{—C}_6$ alkyl, $\text{C}_1\text{—C}_{20}$ alkanoyl, $\text{C}_1\text{—C}_{20}$ cycloalkanoyl, $\text{C}_6\text{—C}_{10}$ aryl- $\text{C}_1\text{—C}_6$ alkylene- $\text{C}(\text{O})\text{—}$, 5- or 6-membered heteroaryl- $\text{C}(\text{O})\text{—}$ or 5- or 6-membered heterocycloalkyl- $\text{C}(\text{O})\text{—}$, wherein the $\text{C}_1\text{—C}_{20}$ alkanoyl, $\text{C}_6\text{—C}_{10}$ aryl- $\text{C}_1\text{—C}_6$ alkylene- $\text{C}(\text{O})\text{—}$, 5- or 6-membered heteroaryl- $\text{C}(\text{O})\text{—}$ and 5- or 6-membered heterocycloalkyl- $\text{C}(\text{O})\text{—}$ of R^1 are each optionally substituted with 1, 2 or 3 substituents independently selected from halo, OH, CN, $\text{C}_1\text{—C}_6$ alkoxy, $\text{C}_1\text{—C}_6$ alkyl, —COOH , $\text{—COO}(\text{C}_1\text{—C}_6)$, $\text{C}_1\text{—C}_6$ haloalkyl and $\text{C}_1\text{—C}_6$ haloalkoxy;

[0175] R^2 is NH_2 , substituted amino, OH, substituted hydroxy, $\text{C}_1\text{—C}_{20}$ alkylamino, phenyl- $\text{C}_1\text{—C}_8$ alkylene- NH— , wherein $\text{C}_1\text{—C}_{20}$ alkylamino and phenyl- $\text{C}_1\text{—C}_8$ alkylene- NH— are each optionally substituted with a substituent independently selected from NH_2 , —COOH and phenyl;

[0176] X1 is absent, or is Asp, isoAsp, Asp(OMe), Glu, bhGlu, bGlu, Gly, N-substituted Gly, Gla, Glp, Ala, Arg, Dab, Leu, Lys, Dap, Orn, (D)Asp, (D)Arg, Tet1, Tet2, Lys, substituted Lys, (D)Lys, substituted (D)Lys, or Cys, (D)Cys, aMeCys, hCys, Pen, (D)Pen, NMe-Glu, Aad, Dap or isoGlu;

[0177] X2 is Ala, Thr, Gly, N-substituted Gly, Ser, Cys, 4S_Mcp, 4R_Mcp, NMe-Thr, aMePro, Hyp_3R, HyP_3S, hSer or Thr_Me;

[0178] X3 is Ala, Gly, N-substituted Gly, His, substituted His, Cys, (D)Cys, aMeCys, hCys, Pen, 3Pal, 4Pal, BIP, Ala 3Quin, Trp_50H or His_1Me;

[0179] X4 is Ala, Phe, Dpa, Gly, N-substituted Gly, bhPhe, a-MePhe, NMe-Phe, D-Phe, 2Pal, or Cys, (D)Cys, aMeCys, hCys, Pen or DIP;

[0180] X5 is Ala, Pro, D-Pro, bhPro, D-bhPro, NPC, D-NPC, Gaba, 2-Pyrrolidinepropanoic acid (Ppa), 2-Pyrrolidinebutanoic acid (Pba), Glu, Lys, substituted Lys, (D)Lys, substituted (D)Lys, or Cys, (D)Cys, aMeCys, hCys, Pen, NMe-Cys, 4S_Mcp, 4R_Mcp, Morph or Tic;

[0181] X6 is Cys, (D)Cys, aMeCys, hCys, Pen, dPen, NMe-Cys, Ala, Lys, (D)Lys, substituted Lys or substituted (D)Lys;

[0182] X7 is absent, or is Ala, Gly, N-substituted Gly, Ile, Val, Leu, NLeu, Lys, substituted Lys, (D)Lys, substituted (D)Lys, Cys, (D)Cys, aMeCys, hCys, Pen, NMe-Cys, Phe or DIP;

[0183] X8 is absent or is Ala, (D)Ala, Ile, Gly, N-substituted Gly, Glu, Val, Leu, NLeu, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys, aMeLys, 123Triazole, Cys, (D)Cys, aMeCys, hCys, Pen, Lys_Me3;

[0184] X9 is absent, or is Ala, Ile, Gly, N-substituted Gly, Val, Leu, NLeu, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys, Cys, (D)Cys, aMeCys, hCys, Pen, NMe-Phe, aMePhe, Dip, dDIP, BIP, aMePhe or substituted Phe;

[0185] X10 is absent, or Ala, Gly, N-substituted Gly, Ile, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys, Cys, (D)Cys, aMeCys, hCys, Pen, Tle, substituted (D)Lys, NMe-Lys, NMe-dLys, substituted NMe-Lys, substituted NMe-dLys or Mor_propanoic acid;

[0186] X11 is absent, or is Ala, Pro, bhPhe, Lys, substituted Lys, (D)Lys, NMe-Phe, NMe-dSer, NMe-dGln, NMe-dLeu, NMe-dTyr, INal or NMe-Ile;

[0187] each of X12-X14 is absent, or is independently an amino acid;

[0188] the peptide of formula (I') is optionally further conjugated with any amino acid;

[0189] any of the amino acids of the peptide of formula (I') is optionally replaced with the corresponding (D)-amino acid or optionally N-substituted;

[0190] (i) at least one of X1, X3-X5, X7-X10 is Cys, (D)Cys, aMeCys, hCys, Pen, (D)Pen, NMe-Cys, 4S_Mcp or 4R_Mcp, wherein the peptide is cyclized by taking the mercapto group on the side chain of one of X1, X3-X5 and X7-X10, the mercapto group on the side chain of X6 and L^x to form a —S-L^x-S— linkage; or the mercapto group on the side chain of X1, the mercapto group on the side chain of X8 and L^x taken together form a —S-L^x-S— linkage; or the mercapto group on the side chain of X2, the mercapto group on the side chain of X5 and L^x taken together form a —S-L^x-S— linkage, wherein each L^x is independently a bond or a linking group; and

[0191] (ii) the functional groups on the side chains of X1 and X10 are optionally taken together to form an amide linkage; or the functional group on the side chain of X1 and the functional group of R² are optionally taken together to form an amide linkage;

and wherein

[0192] Dapa is diaminopropanoic acid; Dpa or DIP is 3,3-diphenylalanine or b,b-diphenylalanine; bhPhe is b-homophenylalanine; Bip is biphenylalanine; bhPro is b-homoproline; Tic is L-1,2,3,4-tetrahydro-isoquinoline-3-carboxylic acid; NPC is L-nipecotic acid; bhTrp is b-homoTryptophane; 1-Nal is 1-naphthylalanine; 2-Nal is 2-naphthylalanine; Om is orinithine; Nleu is norleucine; 2Pal is 2-pyridylalanine; Ppa is 2-(R)-Pyrrolidinepropanoic acid, Pba is 2-(R)-Pyrrolidinebutanoic acid; substituted Phe is phenylalanine wherein phenyl is substituted with F, Cl, Br, I, OH, methoxy, dimethoxy, dichloro, dimethyl, difluoro, pentafluoro, allyloxy, azido, nitro, 4-carbamoyl-2,6-dimethyl, trifluoromethoxy, trifluoromethyl, phenoxy, benzyloxy, carbamoyl, t-Bu, carboxyl, CN, or guanidine;

[0193] substituted bhPhe is b-homophenylalanine wherein phenyl is substituted with F, Cl, Br, I, OH, methoxy, dimethoxy, dichloro, dimethyl, difluoro, pentafluoro, allyloxy, azido, nitro, 4-carbamoyl-2,6-dimethyl, trifluoromethoxy, trifluoromethyl, phenoxy, benzyloxy, carbamoyl, t-Bu, carboxyl, CN, or guanidine;


[0194] substituted Trp is N-methyl-L-tryptophan, a-methyltryptophan, or tryptophan substituted with F, Cl, OH, or t-Bu;

[0195] substituted bhTrp is N-methyl-L-b-homotryptophan, a-methyl-b-homotryptophan, or b-homotryptophan substituted with F, Cl, OH, or t-Bu;

[0196] substituted Lys, substituted dLys or substituted NMe-Lys or substituted NMe-dLys is Lys, dLys, NMe-


Lys or NMe-dLys, wherein the 8-amino group on the side chain of Lys, dLys, NMe-Lys or NMe-dLys is covalently bound to C₁-C₆ alkanoyl, a half-life extension moiety or a linking moiety covalently linked to a half-life extension moiety, wherein the linking moiety comprises one or more linker moieties covalently bound to each other; Tet1 is (S)-(2-amino)-3-(2H-tetrazol-5-yl)propanoic acid; and Tet2 is (S)-(2-amino)-4-(1H-tetrazol-5-yl)butanoic acid;

[0197] 123Triazole is




and

[0198] Dab is



[0199] In some embodiments of hepcidin analogues of Formula (I') or a pharmaceutically acceptable salt or solvate thereof, the invention provides a hepcidin analogue comprising a peptide comprising an amino acid sequence of formula (X):



wherein:

[0200] (i) one of X1, X3-X5, X7-X10 is Cys, (D)Cys, aMeCys, hCys, Pen, (D)Pen, NMe-Cys, 4S_Mcp or 4R_Mcp, wherein the peptide is cyclized by taking the mercapto group on the side chain of one of X1, X3-X5 and X7-X10, the mercapto group on the side chain of X6 and L^x to form a —S-L^x-S— linkage; or the mercapto group on the side chain of X1, the mercapto group on the side chain of X8 and L^x taken together form a —S-L^x-S— linkage; or the mercapto group on the side chain of X2, the mercapto group on the side chain of X5 and L^x taken together form a —S-L^x-S— linkage, wherein each L^x is independently a bond or a linking group; and

[0201] (ii) the functional groups on the side chains of X1 and X10 are optionally taken together to form an amide linkage; or the functional group on the side chain of X1 and the functional group of R² are optionally taken together to form an amide linkage.

[0202] In some embodiments of peptides of formula (I') or (X), or a pharmaceutically acceptable salt or solvate thereof, X1 is Asp, isoAsp, Asp(OMe), Glu, bhGlu, bGlu, Gly, N-substituted Gly, Gla, Glp, Ala, Arg, Dab, Leu, Lys, Dap, Orn, (D)Asp, (D)Arg, Tet1, Tet2, Lys, substituted Lys, (D)Lys, substituted (D)Lys, NMe-Glu, Aad, Dap or isoGlu;

[0203] X2 is Ala, Thr, Gly, N-substituted Gly, Ser, NMe-Thr, aMePro, Hyp_3R, HyP_3S, hSer or Thr_Me;

[0204] X3 is Ala, Gly, N-substituted Gly, His, substituted His, 3Pal, 4Pal, BIP, Ala_3Quin, Trp_50H or His_1Me;

[0205] X4 is Ala, Phe, Dpa, Gly, N-substituted Gly, bhPhe, a-MePhe, NMe-Phe, D-Phe, 2Pal or DIP;

[0206] X5 is Ala, Pro, D-Pro, bhPro, D-bhPro, NPC, D-NPC, Gaba, 2-Pyrrolidinepropanoic acid (Ppa), 2-Pyrrolidinebutanoic acid (Pba), Glu, Lys, substituted Lys, (D)Lys, substituted (D)Lys, Morph or Tic;

[0207] X6 is Cys, (D)Cys, aMeCys, hCys, Pen, dPen or NMe-Cys;

[0208] X7 is absent, Cys, (D)Cys, aMeCys, hCys, Pen, NMe-Cys or Phe;


[0209] X8 is absent, or is Ala, (D)Ala, Ile, Gly, N-substituted Gly, Glu, Val, Leu, NLeu, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys, 123Triazole or Lys_Me3;

[0210] X9 is absent, or is Ala, Ile, Gly, N-substituted Gly, Val, Leu, NLeu, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys NMe-Phe, aMePhe, Dip, dDIP, BIP, aMePhe or substituted Phe;

[0211] X10 is absent, or Ala, Gly, N-substituted Gly, Ile, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys, Tle, substituted (D)Lys, NMe-Lys, NMe-dLys, substituted NMe-Lys, substituted NMe-dLys or Mor_propanoic_acid;

[0212] X11 is absent, or is Ala, Pro, bhPhe, Lys, substituted Lys, (D)Lys, NMe-Phe, NMe-dSer, NMe-dGln, NMe-dLeu, NMe-dTyr, 1Nal or NMe-Ile;


[0213] the mercapto group on the side chain of X6, the mercapto group on the side chain of X7 and L^x taken together form a $-S-L^x-S-$ linkage, wherein L^x is a bond or a linking group selected from C_{1-8} alkylene, phenylene,




wherein the alkylene and phenylene are each optionally substituted with 1 or 2 substituents independently selected from C_{1-6} alkyl, halo, CN, OH, $-COOH$, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy or NH_2 ; and L^a and L^b are each independently C_{1-4} alkylene; and

[0214] the functional groups on the side chains of X1 and X10 are optionally taken together to form an amide linkage; or the functional group on the side chain of X1 and the functional group of R^2 are optionally taken together to form an amide linkage. In one embodiment, X6 and X7 form a disulfide bond, wherein X6 is Cys and X7 is Cys. In another embodiment, X6 and X7 form a disulfide bond, wherein X6 is Cys and X7 is NMe-Cys. In another embodiment, X6 and X7 form a disulfide bond, wherein X6 is hCys and X7 is hCys. In another embodiment, X6 and X7 form a disulfide bond, wherein X6 is Cys and X7 is hCys. In another embodiment, X6 and X7 form a disulfide bond, wherein X6 is Pen and X7 is Cys. In another embodiment, X6 and X7 form a disulfide bond, wherein X6 is Cys and X7 is Pen. In another embodiment, X6 and X7 form a disulfide bond, wherein X6 is Pen and X7 is Pen.


[0215] In some embodiments of peptides of formula (I') or (X), or a pharmaceutically acceptable salt or solvate thereof, L^x is a bond or



wherein L^a and L^b are each independently C_{1-6} alkyl. In one embodiment, L^x is a bond. In another embodiment, L^x is



wherein L^a and L^b are each independently C_{1-6} alkyl, for example, L^a and L^b are each CH_3 . In some embodiments, L^x is



wherein L^a and L^b are each independently C_{1-6} alkyl, for example methyl. In one embodiment, L^x is p-Xylene.

[0216] In some embodiments of peptides of formula (I') or (X), or a pharmaceutically acceptable salt or solvate thereof, X1 is Cys, (D)Cys, aMeCys, hCys, Pen or (D)Pen;

[0217] X2 is Ala, Thr, Gly, N-substituted Gly, Ser, NMe-Thr, aMePro, Hyp_3R, HyP_3S, hSer or Thr_Me;

[0218] X3 is Ala, Gly, N-substituted Gly, His, substituted His, 3Pal, 4Pal, BIP, Ala_3Quin, Trp_50H or His_1Me;

[0219] X4 is Ala, Phe, Dpa, Gly, N-substituted Gly, bhPhe, a-MePhe, NMe-Phe, D-Phe, 2Pal or DIP;

[0220] X5 is Ala, Pro, D-Pro, bhPro, D-bhPro, NPC, D-NPC, Gaba, 2-Pyrrolidinepropanoic acid (Ppa), 2-Pyrrolidinebutanoic acid (Pba), Glu, Lys, substituted Lys, (D)Lys, substituted (D)Lys, Morph or Tic;

[0221] X6 is Ala, Lys, (D)Lys, substituted Lys or substituted (D)Lys;

[0222] X7 is absent, or is Ala, Gly, N-substituted Gly, Ile, Val, Leu, NLeu, Lys, substituted Lys, (D)Lys, substituted (D)Lys, Phe or DIP;

- [0223] X8 is Cys, (D)Cys, aMeCys, hCys or Pen;
 [0224] X9 is absent, or is Ala, Ile, Gly, N-substituted Gly, Val, Leu, NLeu, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys NMe-Phe, aMePhe, Dip, dDIP, BIP, aMePhe or substituted Phe;
 [0225] X10 is absent, or Ala, Gly, N-substituted Gly, Ile, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys, Tle, substituted (D)Lys, NMe-Lys, NMe-dLys, substituted NMe-Lys, substituted NMe-dLys or Mor-propanoic acid;
 [0226] X11 is absent, or is Ala, Pro, bhPhe, Lys, substituted Lys, (D)Lys, NMe-Phe, NMe-dSer, NMe-dGln, NMe-dLeu, NMe-dTyr, 1Nal or NMe-Ile;
 [0227] the mercapto group on the side chain of X1 and the mercapto group on the side chain of X8 taken together form a disulfide bond; and

the functional groups on the side chains of X1 and X10 are optionally taken together to form an amide linkage; or the functional group on the side chain of X1 and the functional group of R² are optionally taken together to form an amide linkage.

In one embodiment, X1 and X8 form a disulfide bond, wherein X1 is hCy and X8 is Cys. In another embodiment, X1 and X8 form a disulfide bond, wherein X1 is Pen and X8 is Cys. In another embodiment, X1 and X8 form a disulfide bond, wherein X1 is Cys and X8 is dPen. In another embodiment, X1 and X8 form a disulfide bond, wherein X1 is dCys and X8 is dCys. In another embodiment, X1 and X8 form a disulfide bond, wherein X1 is dCys and X8 is Cys. In another embodiment, X1 and X8 form a disulfide bond, wherein X1 is Cys and X8 is hCys. In another embodiment, X1 and X8 form a disulfide bond, wherein X1 is hcy and X8 is Pen. In another embodiment, X1 and X8 form a disulfide bond, wherein X1 is hCys and X8 is hCys. In another embodiment, X1 and X8 form a disulfide bond, wherein X1 is Cys and X8 is Cys.

[0228] In some embodiments peptides of formula (I') or (X), or a pharmaceutically acceptable salt or solvate thereof, X1 is Asp, isoAsp, Asp(OMe), Glu, bhGlu, bGlu, Gly, N-substituted Gly, Glu, Glp, Ala, Arg, Dab, Leu, Lys, Dap, Orn, (D)Asp, (D)Arg, Tet1, Tet2, Lys, substituted Lys, (D)Lys, substituted (D)Lys, NMe-Glu, Aad, Dap or isoGlu;

- [0229] X2 is Cys, 4S_Mcp or 4R_Mcp;
 [0230] X3 is Ala, Gly, N-substituted Gly, His, substituted His, 3Pal, 4Pal, BIP, Ala 3Quin, Trp_50H or His_1Me;
 [0231] X4 is Ala, Phe, Dpa, Gly, N-substituted Gly, bhPhe, a-MePhe, NMe-Phe, D-Phe, 2Pal or DIP;
 [0232] X5 is Cys, (D)Cys, aMeCys, hCys, Pen, NMe-Cys, 4S_Mcp or 4R_Mcp;
 [0233] X6 is Cys, (D)Cys, aMeCys, hCys, Pen, dPen or NMe-Cys;
 [0234] X7 is absent, Cys, (D)Cys, aMeCys, hCys, Pen, NMe-Cys or Phe;
 [0235] X8 is absent, or is Ala, (D)Ala, Ile, Gly, N-substituted Gly, Glu, Val, Leu, NLeu, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys, 123Triazole or Lys_Me3;
 [0236] X9 is absent, or is Ala, Ile, Gly, N-substituted Gly, Val, Leu, NLeu, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys NMe-Phe, aMePhe, Dip, dDIP, BIP, aMePhe or substituted Phe;
 [0237] X10 is absent, or Ala, Gly, N-substituted Gly, Ile, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted

(D)Lys, Tle, substituted (D)Lys, NMe-Lys, NMe-dLys, substituted NMe-Lys, substituted NMe-dLys or Mor-propanoic acid;

- [0238] X11 is absent, or is Ala, Pro, bhPhe, Lys, substituted Lys, (D)Lys, NMe-Phe, NMe-dSer, NMe-dGln, NMe-dLeu, NMe-dTyr, 1Nal or NMe-Ile;
 [0239] the mercapto group on the side chain of X2 and the mercapto group on the side chain of X5 taken together form a disulfide bond; and


the functional groups on the side chains of X1 and X10 are optionally taken together to form an amide linkage; or the functional group on the side chain of X1 and the functional group of R² are optionally taken together to form an amide linkage.

In one embodiment, X2 and X5 form a disulfide bond, wherein X2 is Cys and X5 is Cys. In another embodiment, X2 and X5 form a disulfide bond, wherein X2 is Cys and X5 is NMe-Cys. In another embodiment, X2 and X5 form a disulfide bond, wherein X2 is Cys and X5 is hCys. In another embodiment, X2 and X5 form a disulfide bond, wherein X2 is Cys and X5 is 4S_Mcp. In another embodiment, X2 and X5 form a disulfide bond, wherein X2 is Cys and X5 is 4R_Mcp. In another embodiment, X2 and X5 form a disulfide bond, wherein X2 is 4S_Mcp and X5 is 4R_Mcp. In another embodiment, X2 and X5 form a disulfide bond, wherein X2 is 4S_Mcp and X5 is 4S_Mcp. In another embodiment, X2 and X5 form a disulfide bond, wherein X2 is 4R_Mcp and X5 is 4R_Mcp.

[0240] In some embodiments peptides of formula (I') or (X), or a pharmaceutically acceptable salt or solvate thereof, wherein the peptide is cyclized by taking the functional groups the side chains of X1 and X10 together to form an amide linkage, i.e., —C(O)NH— or —NHC(O)— linkage. In certain instances, the peptide is cyclized by taking the amino or carboxy group on the side chain of X1 and carboxy or amino group on the said chain of X10 together to form an amide linkage. In some embodiments, X1 and X10 form an amide linkage and X6 and X7 form a disulfide bond. In one embodiment, X1 and X10 form an amide linkage and X6 and X7 form a disulfide bond, wherein X1 is Glu and X10 is dK.

[0241] In some embodiments peptides of formula (I') or (X), or a pharmaceutically acceptable salt or solvate thereof, wherein the peptide is cyclized by taking the functional groups on the side chains of X1 and functional group of R² together to form an amide linkage, i.e., —C(O)NH— or —NHC(O)— linkage. In certain instances, the peptide is cyclized by taking the amino or carboxy group on the side chain of X1 and the carboxy or amino group of R² together to form an amide linkage. In some embodiments, X1 and R² form an amide linkage and X6 and X7 form a disulfide bond. In one embodiment, X1 and R² form an amide linkage and X6 and X7 form a disulfide bond, wherein X1 is Glu and R² is C₁-C₂₀ alkylamino or phenyl-C₁-C₈ alkylene-NH—, each of which is substituted with a substituent independently selected from NH₂ and —COOH. In one embodiment, R² is C₁₋₈ alkylamino substituted with NH₂ or COOH. In one embodiment, R² is 2-aminoethylamino.

[0242] In certain embodiments, the present invention includes a hepcidin analogue comprising a peptide of Formula (I):



or a pharmaceutically acceptable salt or solvate thereof,

wherein:

- [0243] R¹ is hydrogen, C₁-C₆ alkyl, C₆-C₁₂ aryl, C₆-C₁₂ aryl-C₁-C₆ alkyl, C₁-C₂₀ alkanoyl, C₁-C₂₀ cycloalkanoyl;
- [0244] R² is NH₂, substituted amino, OH, or substituted hydroxy;
- [0245] X0 is absent, or is Cys, (D)Cys, aMeCys, hCys, or Pen;
- [0246] X1 is absent, or is Asp, isoAsp, Asp(OMe), Glu, bhGlu, bGlu, Gly, N-substituted Gly, Gla, Glp, Ala, Arg, Dab, Leu, Lys, Dap, Orn, (D)Asp, (D)Arg, Tet1, or Tet2, Lys, substituted Lys, (D)Lys, substituted (D)Lys, or Cys, (D)Cys, aMeCys, hCys, or Pen;
- [0247] X2 is Ala, Thr, Gly, N-substituted Gly, or Ser;
- [0248] X3 is Ala, Gly, N-substituted Gly, His, substituted His, or Cys, (D)Cys, aMeCys, hCys, or Pen;
- [0249] X4 is Ala, Phe, Dpa, Gly, N-substituted Gly, bhPhe, a-MePhe, NMe-Phe, D-Phe, 2Pal, or Cys, (D)Cys, aMeCys, hCys, or Pen;
- [0250] X5 is Ala, Pro, D-Pro, bhPro, D-bhPro, NPC, D-NPC, Gaba, 2-Pyrrolidinepropanoic acid (Ppa), 2-Pyrrolidinebutanoic acid (Pba), Glu, Lys, substituted Lys, (D)Lys, substituted (D)Lys, or Cys, (D)Cys, aMeCys, hCys, or Pen;
- [0251] X6 is Cys, (D)Cys, aMeCys, hCys, or Pen;
- [0252] X7 is absent, or is Ala, Gly, N-substituted Gly, Ile, Val, Leu, NLeu, Lys, substituted Lys, (D)Lys, substituted (D)Lys, or Cys, (D)Cys, aMeCys, hCys, or Pen;
- [0253] X8 is absent or is Ala, (D)Ala, Ile, Gly, N-substituted Gly, Glu, Val, Leu, NLeu, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys, aMeLys, 123Triazole, or Cys, (D)Cys, aMeCys, hCys, or Pen;
- [0254] X9 is absent, or is Ala, Ile, Gly, N-substituted Gly, Val, Leu, NLeu, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys, or Cys, (D)Cys, aMeCys, hCys, or Pen;
- [0255] X10 is absent, or is Ala, Gly, N-substituted Gly, Ile, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys, or Cys, (D)Cys, aMeCys, hCys, or Pen;
- [0256] X11 is absent, or is Ala, Pro, bhPhe, Lys, substituted Lys, or (D)Lys;

and

- [0257] each of X12-X14 is absent, or is independently any amino acid;

provided that:

- [0258] i) the peptide may further be conjugated at any amino acid;
- [0259] ii) any of the amino acids of the peptide may be the corresponding (D)-amino acid of the amino acid or may be N-substituted; and
- [0260] iii) at least one of X0, X1, X3-X5, X7-X10 is Cys, (D)Cys, aMeCys, hCys, or Pen and Cys, (D)Cys, aMeCys, hCys, or Pen form a disulfide bond with X6;

and

- [0261] Dapa is diaminopropanoic acid; Dpa or DIP is 3,3-diphenylalanine or b,b-diphenylalanine; bhPhe is b-homophenylalanine; Bip is biphenylalanine; bhPro is b-homoproline; Tic is L-1,2,3,4,-tetrahydro-isoquinoline-3-carboxylic acid; NPC is L-nipecotic acid; bhTrp is b-homoTryptophane; 1-Nal is 1-naphthylalanine; 2-Nal is 2-naphthylalanine; Orn is orinithine; Nleu is norleucine; 2Pal is 2-pyridylalanine; Ppa is 2-(R)-Pyrrolidinepropanoic acid, Pba is 2-(R)-Pyrrolidinebu-

tanoic acid; substituted Phe is phenylalanine wherein phenyl is substituted with F, Cl, Br, I, OH, methoxy, dimethoxy, dichloro, dimethyl, difluoro, pentafluoro, allyloxy, azido, nitro, 4-carbamoyl-2,6-dimethyl, trifluoromethoxy, trifluoromethyl, phenoxy, benzyloxy, carbamoyl, t-Bu, carboxyl, CN, or guanidine;


- [0262] substituted bhPhe is b-homophenylalanine wherein phenyl is substituted with F, Cl, Br, I, OH, methoxy, dimethoxy, dichloro, dimethyl, difluoro, pentafluoro, allyloxy, azido, nitro, 4-carbamoyl-2,6-dimethyl, trifluoromethoxy, trifluoromethyl, phenoxy, benzyloxy, carbamoyl, t-Bu, carboxyl, CN, or guanidine;

- [0263] substituted Trp is N-methyl-L-tryptophan, a-methyltryptophan, or tryptophan substituted with F, Cl, OH, or t-Bu;

- [0264] substituted bhTrp is N-methyl-L-b-homotryptophan, a-methyl-b-homotryptophan, or b-homotryptophan substituted with F, Cl, OH, or t-Bu;


- [0265] Tet1 is (S)-(2-amino)-3-(2H-tetrazol-5-yl)propanoic acid; and Tet2 is (S)-(2-amino)-4-(1H-tetrazol-5-yl)butanoic acid;

- [0266] 123Triazole is



and

- [0267] Dab is



- [0268] In one embodiment, X0 is Cys, (D)Cys, aMeCys, hCys, or Pen; and X0 and X6 are linked via a disulfide bond.

- [0269] In one embodiment, X1 is Cys, (D)Cys, aMeCys, hCys, or Pen; and X1 and X6 are linked via a disulfide bond.

- [0270] In one embodiment, X3 is Cys, (D)Cys, aMeCys, hCys, or Pen; and X3 and X6 are linked via a disulfide bond.

- [0271] In one embodiment, X4 is Cys, (D)Cys, aMeCys, hCys, or Pen; and X4 and X6 are linked via a disulfide bond.

- [0272] In one embodiment, X5 is Cys, (D)Cys, aMeCys, hCys, or Pen; and X5 and X6 are linked via a disulfide bond.

- [0273] In one embodiment, X7 is Cys, (D)Cys, aMeCys, hCys, or Pen; and X7 and X6 are linked via a disulfide bond.

- [0274] In one embodiment, X8 is Cys, (D)Cys, aMeCys, hCys, or Pen; and X8 and X6 are linked via a disulfide bond.

- [0275] In one embodiment, X9 is Cys, (D)Cys, aMeCys, hCys, or Pen; and X9 and X6 are linked via a disulfide bond.

- [0276] In one embodiment, X10 is Cys, (D)Cys, aMeCys, hCys, or Pen; and X10 and X6 are linked via a disulfide bond.


- [0277] In one embodiment,

- [0278] X1 is Glu, Dab, Dap, Orn, Lys, or Tet1;

- [0279] X2 is Thr;


- [0280] X3 is His or 1MeHis;

- [0281] X4 is Dpa;
 [0282] X5 is Ala or Pro;
 [0283] X6 is absent, Ala, Glu, or substituted Lys;
 [0284] X7 is absent, or is Ala, Ile, Lys, substituted Lys, (D)Lys, or substituted (D)Lys;
 [0285] X8 is absent, or is Ala, Ile, Glu, Asp, 123Triazole, Lys, substituted Lys, (D)Lys, substituted (D)Lys, or aMeLys;
 [0286] X9 is absent, or is bhPhe;
 [0287] X10 is absent, or is Ala, Ile, Phe, bhPhe, Lys, substituted Lys, (D)Lys, or substituted (D)Lys; and
 [0288] X11 is absent, or is Pro, bhPhe, Lys, substituted Lys, or (D)Lys.
 [0289] In one embodiment, X1 is Glu.
 [0290] In one embodiment, X2 is Thr.
 [0291] In one embodiment, X3 is His.
 [0292] In one embodiment, X4 is Phe or Dpa.
 [0293] In one embodiment, X5 is Pro.
 [0294] In one embodiment, X7 is Ile.
 [0295] In one embodiment, X8 is Lys, substituted Lys, (D)Lys, or substituted (D)Lys.
 [0296] In one embodiment, X9 is Phe or bhF.
 [0297] In one embodiment, X10 is Lys, substituted Lys, (D)Lys, or substituted (D)Lys.
 [0298] In one embodiment, X11 is absent, Arg, Lys, substituted Lys, (D)Lys, or substituted (D)Lys.
 [0299] In one embodiment, the peptide is according to Formula II:




wherein R^1 , R^2 , X6, X8, and X10-X14 are as in claim 1; and wherein Cys and X6 linked via a disulfide bond.

[0300] In one embodiment, the peptide is according to Formula III:




wherein R^1 , R^2 , X6, X8, and X10-X14 are as in claim 1; and wherein Cys and X6 linked via a disulfide bond.

[0301] In one embodiment, the peptide is according to Formula IV:




wherein R^1 , R^2 , X6, X8, and X10-X14 are as in claim 1; and wherein Cys and X6 linked via a disulfide bond.

[0302] In one embodiment, the peptide is according to Formula V:




wherein R^1 , R^2 , X6, X8, and X10-X14 are as in claim 1; and wherein Cys and X6 linked via a disulfide bond.

[0303] In one embodiment, the peptide is according to Formula VI:





wherein R^1 , R^2 , X6, and X10-X14 are as in claim 1; and wherein Cys and X6 linked via a disulfide bond.

[0304] In one embodiment, the peptide is according to Formula VII:



wherein R^1 , R^2 , X6, X8, X10, and X11-X14 are as in claim 1; and wherein Cys and X6 linked via a disulfide bond.

[0305] In one embodiment, the peptide is according to Formula VIIIa or VIIIb:



wherein R^1 , R^2 , X6, X8, X10, and X11-X14 are as in claim 1; and wherein Cys and X6 linked via a disulfide bond.

- [0306] In one embodiment, X6 is Pen.
 [0307] In one embodiment, X6 is hCys.
 [0308] In one embodiment, X6 is aMeCys.
 [0309] In one embodiment, X6 is (D)Cys.
 [0310] In one embodiment, X6 is Cys.
 [0311] In one embodiment, X8 is Lys substituted with Ahx-Palm.
 [0312] In one embodiment, X8 is absent, Lys, substituted Lys, (D)Lys, or substituted (D)Lys.
 [0313] In one embodiment, X8 is absent.
 [0314] In one embodiment, X8 is (D)Lys.
 [0315] In one embodiment, X8 is Lys.
 [0316] In one embodiment, X8 is Lys substituted with Ahx-Palm.
 [0317] In one embodiment, X8 is Lys(Ahx_Palm).
 [0318] In one embodiment, X8 is a conjugated amino acid.
 [0319] In one embodiment, X8 is conjugated Lys or (D)Lys.
 [0320] In one embodiment, X8 is Lys(L1Z) or (D)Lys(L1Z), wherein L1 is a linker, and wherein Z is a half-life extension moiety.
 [0321] In one embodiment, X10 is Glu.
 [0322] In one embodiment, X10 is absent, Lys, substituted Lys, (D)Lys, or substituted (D)Lys.
 [0323] In one embodiment, X10 is absent.
 [0324] In one embodiment, X10 is (D)Lys.
 [0325] In one embodiment, X10 is Lys.
 [0326] In one embodiment, X10 is Lys substituted with Ahx-Palm.
 [0327] In one embodiment, X10 is Lys(Ahx_Palm).
 [0328] In one embodiment, X10 is a conjugated amino acid.
 [0329] In one embodiment, X10 is conjugated Lys or (D)Lys.
 [0330] In one embodiment, X10 is Lys(L1Z) or (D)Lys(L1Z), wherein L1 is a linker, and wherein Z is a half-life extension moiety.
 [0331] In one embodiment, X8 is (D)Lys; and X10 is Lys(L1Z) or (D)Lys(L1Z). In one embodiment, X10 is Lys(L1Z).
 [0332] In one embodiment, X8 is Lys(L1Z) or (D)Lys(L1Z); and X10 is absent, Glu, Lys, or (D)Lys. In one embodiment, X8 is Lys(L1Z). In another embodiment, X10 is absent. In yet another embodiment, X10 is (D)Lys. In a particular embodiment, X10 is Glu.
 [0333] In one embodiment, L1 is a single bond.
 [0334] In one embodiment, L1 is iso-Glu.
 [0335] In one embodiment, L1 is Ahx.
 [0336] In one embodiment, L1 is iso-Glu-Ahx.
 [0337] In one embodiment, L1 is PEG.
 [0338] In one embodiment, L1 is PEG-Ahx.
 [0339] In one embodiment, L1 is iso-Glu-PEG-Ahx.

[0340] In one embodiment, PEG is $-\text{C}(\text{O})-\text{CH}_2-(\text{Peg})_n-\text{N}(\text{H})\text{m}-$, or $-\text{C}(\text{O})-\text{CH}_2-\text{CH}_2-(\text{Peg})_n-\text{N}(\text{H})\text{m}-$; and Peg is $-\text{OCH}_2\text{CH}_2-$, m is 1, 2, or 3; and n is an integer between 1-100, or is 10K, 20K, or 30K.

[0341] In one embodiment, m is 1.

[0342] In one embodiment, m is 2.

[0343] In one embodiment, n is 2.

[0344] In one embodiment, n is 4.

[0345] In one embodiment, n is 8.

[0346] In one embodiment, n is 11.

[0347] In one embodiment, n is 12.

[0348] In one embodiment, n is 20K.

[0349] In one embodiment, PEG is 1Peg2; and 1Peg2 is $-\text{C}(\text{O})-\text{CH}_2-(\text{Peg})_2-\text{N}(\text{H})-$.

[0350] In one embodiment, PEG is 2Peg2; and 2Peg2 is $-\text{C}(\text{O})-\text{CH}_2-\text{CH}_2-(\text{Peg})_2-\text{N}(\text{H})-$.

[0351] In one embodiment, PEG is 1Peg2-1Peg2; and each 1Peg2 is $-\text{C}(\text{O})-\text{CH}_2-\text{CH}_2-(\text{Peg})_2-\text{N}(\text{H})-$.

[0352] In one embodiment, PEG is 1Peg2-1Peg2; and 1Peg2-1Peg2 is $-\text{C}(\text{O})-\text{CH}_2-(\text{OCH}_2\text{CH}_2)_2-\text{NH}-\text{C}(\text{O})-\text{CH}_2-(\text{OCH}_2\text{CH}_2)_2-\text{NH}-$.

[0353] In one embodiment, PEG is 2Peg4; and 2Peg4 is $-\text{C}(\text{O})-\text{CH}_2-\text{CH}_2-(\text{Peg})_4-\text{N}(\text{H})-$, or $-\text{C}(\text{O})-\text{CH}_2-\text{CH}_2-(\text{OCH}_2\text{CH}_2)_4-\text{NH}-$.

[0354] In one embodiment, PEG is 1Peg8; and 1Peg8 is $-\text{C}(\text{O})-\text{CH}_2-(\text{Peg})_8-\text{N}(\text{H})-$, or $-\text{C}(\text{O})-\text{CH}_2-(\text{OCH}_2\text{CH}_2)_8-\text{NH}-$.

[0355] In one embodiment, PEG is 2Peg8; and 2Peg8 is $-\text{C}(\text{O})-\text{CH}_2-\text{CH}_2-(\text{Peg})_8-\text{N}(\text{H})-$, or $-\text{C}(\text{O})-\text{CH}_2-\text{CH}_2-(\text{OCH}_2\text{CH}_2)_8-\text{NH}-$.

[0356] In one embodiment, PEG is 1Peg11; and 1Peg11 is $-\text{C}(\text{O})-\text{CH}_2-(\text{Peg})_{11}-\text{N}(\text{H})-$, or $-\text{C}(\text{O})-\text{CH}_2-(\text{OCH}_2\text{CH}_2)_{11}-\text{NH}-$.

[0357] In one embodiment, PEG is 2Peg11; and 2Peg11 is $-\text{C}(\text{O})-\text{CH}_2-\text{CH}_2-(\text{Peg})_{11}-\text{N}(\text{H})-$, or $-\text{C}(\text{O})-\text{CH}_2-\text{CH}_2-(\text{OCH}_2\text{CH}_2)_{11}-\text{NH}-$.

[0358] In one embodiment, PEG is 2Peg11' or 2Peg12; and 2Peg11' or 2Peg12 is $-\text{C}(\text{O})-\text{CH}_2-\text{CH}_2-(\text{Peg})_{12}-\text{N}(\text{H})-$, or $-\text{C}(\text{O})-\text{CH}_2-\text{CH}_2-(\text{OCH}_2\text{CH}_2)_{12}-\text{NH}-$.

[0359] In one embodiment, when PEG is attached to Lys, the $-\text{C}(\text{O})-$ of PEG is attached to Ne of Lys.

[0360] In one embodiment, when PEG is attached to isoGlu, the $-\text{N}(\text{H})-$ of PEG is attached to $-\text{C}(\text{O})-$ of isoGlu.

[0361] In one embodiment, when PEG is attached to Ahx, the $-\text{N}(\text{H})-$ of PEG is attached to $-\text{C}(\text{O})-$ of Ahx.

[0362] In one embodiment, when PEG is attached to Palm, the $-\text{N}(\text{H})-$ of PEG is attached to $-\text{C}(\text{O})-$ of Palm.

[0363] In one embodiment, Z is Palm.

[0364] In one embodiment, -L1Z is:

[0365] PEG11_OMe;

[0366] PEG12_C18 acid;

[0367] 1PEG2_1PEG2_Ahx_Palm;

[0368] 1PEG2_Ahx_Palm;

[0369] Ado_Palm;

[0370] Ahx_Palm;

[0371] Ahx_PEG20K;

[0372] PEG12_Ahx_IsoGlu_Behenic;

[0373] PEG12 Ahx Palm;

[0374] PEG12_DEKHKHS_Palm;

[0375] PEG12_IsoGlu_C18 acid;

[0376] PEG12_Ahx_C18 acid;

[0377] PEG12_IsoGlu_Palm;

[0378] PEG12_KKK_Palm;

[0379] PEG12_KKKG_Palm;

[0380] PEG12_DEKHKHS_Palm;

[0381] PEG12 Palm;

[0382] PEG12_PEG12_Palm;

[0383] PEG20K;

[0384] PEG4_Ahx_Palm;

[0385] PEG4 Palm;

[0386] PEG8_Ahx_Palm; or

[0387] IsoGlu_Palm;

[0388] -1PEG2_1PEG2_Dap_C18_Diacid;

[0389] -1PEG2_1PEG2_IsoGlu_C10_Diacid;

[0390] -1PEG2_1PEG2_IsoGlu_C12_Diacid;

[0391] -1PEG2_1PEG2_IsoGlu_C14_Diacid;

[0392] -1PEG2_1PEG2_IsoGlu_C16_Diacid;

[0393] -1PEG2_1PEG2_IsoGlu_C18_Diacid;

[0394] -1PEG2_1PEG2_IsoGlu_C22_Diacid;

[0395] -1PEG2_1PEG2_Ahx_C18_Diacid;

[0396] -1PEG2_1PEG2_C18_Diacid;

[0397] -1PEG8_IsoGlu_C18_Diacid;

[0398] -IsoGlu_C18_Diacid;

[0399] -PEG12_Ahx_C18_Diacid;

[0400] -PEG12_C16_Diacid;

[0401] -PEG12_C18_Diacid;

[0402] -1PEG2_1PEG2_1PEG2_C18_Diacid;

[0403] -1PEG2_1PEG2_1PEG2_IsoGlu_C18_Diacid;

[0404] -PEG12_IsoGlu_C18_Diacid;

[0405] -PEG4_IsoGlu_C18_Diacid; or

[0406] -PEG4_PEG4_IsoGlu_C18_Diacid;

[0407] wherein

[0408] PEG11_OMe is $-\text{C}(\text{O})-\text{CH}_2-\text{CH}_2-(\text{OCH}_2\text{CH}_2)_{11}-\text{OMe}$;

[0409] 1PEG2 is $-\text{C}(\text{O})-\text{CH}_2-(\text{OCH}_2\text{CH}_2)_2-\text{NH}-$;

[0410] PEG4 is $-\text{C}(\text{O})-\text{CH}_2-\text{CH}_2-(\text{OCH}_2\text{CH}_2)_4-\text{NH}-$;

[0411] PEG8 is $-\text{C}(\text{O})-\text{CH}_2-\text{CH}_2-(\text{OCH}_2\text{CH}_2)_8-\text{NH}-$;

[0412] 1PEG8 is $-\text{C}(\text{O})-\text{CH}_2-(\text{OCH}_2\text{CH}_2)_8-\text{NH}-$;

[0413] PEG12 is $-\text{C}(\text{O})-\text{CH}_2-\text{CH}_2-(\text{OCH}_2\text{CH}_2)_{12}-\text{NH}-$;


[0414] Ado is $-\text{C}(\text{O})-(\text{CH}_2)_{11}-\text{NH}-$;

[0415] Cn acid is $-\text{C}(\text{O})(\text{CH}_2)_n-2-\text{CH}_3$; C18 acid is $-\text{C}(\text{O})-(\text{CH}_2)_{16}-\text{Me}$;

[0416] Palm is $-\text{C}(\text{O})-(\text{CH}_2)_{14}-\text{Me}$;

[0417] isoGlu is isoglutamic acid;


[0418] isoGlu_Palm is



[0419] Ahx is $-\text{C}(\text{O})-(\text{CH}_2)_5-\text{NH}-$;


[0420] Cn_Diacid is $-\text{C}(\text{O})-(\text{CH}_2)_{n-2}-\text{COOH}$; wherein n is 10, 12, 14, 16, 18, or 22.

[0421] In one embodiment, X6 or X10 is Lys(1PEG2_1PEG2_IsoGlu_C_n_Diacid); and Lys(1PEG2_1PEG2_IsoGlu_C_n_Diacid) is




and n is 10, 12, 14, 16, or 18.

[0422] In one embodiment, X6 or X10 is (D)Lys(1PEG2_1PEG2_IsoGlu_C_n_Diacid); and (D)Lys(1PEG2_1PEG2_IsoGlu_C_n_Diacid) is




[0423] and n is 10, 12, 14, 16, or 18.

[0424] In one embodiment, X6 or X10 is Lys(1PEG8_IsoGlu_C_n_Diacid); and Lys(1PEG8IsoG_C_n_Diacid) is




and n is 10, 12, 14, 16, or 18.

[0425] In one embodiment, X6 or X10 is (D)Lys(1PEG8_IsoGlu_C_n_Diacid); and (D)Lys(1PEG8_IsoGlu_C_n_Diacid) is




and n is 10, 12, 14, 16, or 18.

[0426] In one embodiment, X6 or X10 is Lys(1PEG2_1PEG2_Dap_C_n_Diacid); and Lys(1PEG2_1PEG2_Dap_C_n_Diacid) is




[0427] and n is 10, 12, 14, 16, or 18.

[0428] In one embodiment, X6 or X10 is Lys(IsoGlu_C_n_Diacid); and Lys(IsoGlu_C_n_Diacid) is




[0430] In one embodiment, X6 or X10 is (D)Lys(IsoGlu_C_n_Diacid); and (D)Lys(IsoGlu_C_n_Diacid) is



[0431] and n is 10, 12, 14, 16, or 18.


[0429] and n is 10, 12, 14, 16, or 18.

[0432] In one embodiment, X6 or X10 is Lys(PEG12_IsoGlu_C_n_Diacid); and Lys(PEG12_IsoGlu_C_n_Diacid) is




and n is 10, 12, 14, 16, or 18.

[0433] In one embodiment, X6 or X10 is (D)Lys(PEG12_IsoGlu_C_n_Diacid); and (D)Lys(PEG12_IsoGlu_C_n_Diacid) is




[0434] and n is 10, 12, 14, 16, or 18.

[0435] In one embodiment, X6 or X10 is Lys(PEG4_IsoGlu_C_n_Diacid); and Lys(PEG4_IsoGlu_C_n_Diacid) is



[0436] and n is 10, 12, 14, 16, or 18.


[0437] In one embodiment, X6 or X10 is (D)Lys(PEG4_IsoGlu_C_n_Diacid); and (D)Lys(PEG4_IsoGlu_C_n_Diacid) is




[0438] and n is 10, 12, 14, 16, or 18.

[0439] In one embodiment, X6 or X10 is Lys(PEG4_PEG4_IsoGlu_C_n_Diacid); and Lys(PEG4_PEG4_IsoGlu_C_n_Diacid) is

[0440] and n is 10, 12, 14, 16, or 18.




[0441] In one embodiment, X6 or X10 is (D)Lys(PEG4_PEG4_IsoGlu_C_n_Diacid); and (D)Lys(PEG4_PEG4_IsoGlu_C_n_Diacid) is




[0442] and n is 10, 12, 14, 16, or 18.

[0443] In one embodiment, X6 or X10 is Lys(IsoGlu_C_n_Diacid); and Lys(IsoGlu_C_n_Diacid) is




[0445] In one embodiment, X6 or X10 is (D)Lys(IsoGlu_C_n_Diacid); and (D)Lys(IsoGlu_C_n_Diacid) is



[0446] and n is 10, 12, 14, 16, or 18


[0447] In one embodiment, X6 or X10 is Lys(PEG12_Ahx_C_n_Diacid); and Lys(PEG12_Ahx_C_n_Diacid) is

[0444] and n is 10, 12, 14, 16, or 18.




[0448] and n is 10, 12, 14, 16, or 18.

[0449] In one embodiment, X6 or X10 is Lys(PEG12_Ahx_C_n_Diacid); and Lys(PEG12_Ahx_C_n_Diacid) is




[0450] and n is 10, 12, 14, 16, or 18.

[0451] In one embodiment, X6 or X10 is (D)Lys(PEG12_Ahx_C_n_Diacid); and (D)Lys(PEG12_Ahx_C_n_Diacid) is



[0452] and n is 10, 12, 14, 16, or 18.


[0453] In one embodiment, X6 or X10 is Lys(PEG12_C_n_Diacid); and Lys(PEG12_C_n_Diacid) is



[0454] and n is 10, 12, 14, 16, or 18.


[0455] In one embodiment, X6 or X10 is (D)Lys(PEG12_C_n_Diacid); and


[0456] (D)Lys(PEG12_C_n_Diacid) is




[0457] and n is 10, 12, 14, 16, or 18.

- [0458] In one embodiment, R^2 is NH_2 .
- [0459] In one embodiment, R^2 is substituted amino.
- [0460] In one embodiment, R^2 is N-alkylamino.
- [0461] In one embodiment, R^2 is N-alkylamino, wherein alkyl is further substituted or unsubstituted.
- [0462] In one embodiment, R^2 is N-alkylamino, wherein alkyl is further substituted aryl or heteroaryl.
- [0463] In one embodiment, R^2 is alkylamino, wherein alkyl is unsubstituted or substituted with aryl; and alkyl is ethyl, propyl, butyl, or pentyl.
- [0464] In one embodiment, R^2 is alkylamino, wherein alkyl is unsubstituted or substituted with phenyl; and alkyl is ethyl, propyl, butyl, or pentyl.
- [0465] In one embodiment, R^2 is OH.
- [0466] In one embodiment, R^1 is C1-C20 alkanoyl.
- [0467] In one embodiment, R^1 is IVA or isovaleric acid.
- [0468] In one embodiment, the peptide is a linear peptide.
- [0469] In one embodiment, the peptide is a lactam.
- [0470] In one embodiment, the peptide is a lactam, wherein any free $-NH_2$ is cyclized with any free $-C(O)_2H$.
- [0471] The disclosure includes a hepcidin analogue or pharmaceutically acceptable salt or solvate thereof comprising or consisting of a peptide, wherein the peptide is any one of the peptides listed in Tables 6A-B.
- [0472] The disclosure includes a hepcidin analogue or pharmaceutically acceptable salt or solvate thereof comprising or consisting of a peptide, wherein the peptide is:







-continued




-continued




-continued




-continued

Isovaleric Acid-E-T-H-[Dpa]-[Cys]-[Cys]-[Lys(1PEG2_1PEG2_Ahx_C18_Diacid)]-[bhPhe]-[(D)Lys]-NH₂

ID# 21

Isovaleric Acid-E-T-H-[Dpa]-P-[Cys]-[Cys]-[Lys(1PEG2_1PEG2_Ahx_C18_Diacid)]-[bhPhe]-[(D)Lys]-NH₂


-continued



Isovaleric Acid-E-T-H-[Dpa]-P-[Cys]-I-[Lys(IPEG2_1PEG2_Ahx_C18_Diacid)]-[bzlPhe]-[Cys]-NH2

-continued

ID# 27



-continued

ID# 36




TABLE 7

Exemplary peptides of the invention	
Seq ID No.	Peptide
110	




TABLE 7-continued


Exemplary peptides of the invention	
Seq ID No.	Peptide
111	 Isovaleric_Acid-[Hcy]-T-H-[DIP]-P-A-I-[Hcy]-[bhF]-[Lys_Ahx_Palm]-[dK]-NH₂

TABLE 7-continued



Exemplary peptides of the invention	
Seq ID No.	Peptide
122	 Isovaleric_Acid-E-C-H-[DIP]-C-[Lys_1PEG2_1PEG2_DMG_N_2ae_Palm]-[N_Butyl_Phe]
127	 Isovaleric_Acid-E-[4S_Mcp]-H-[DIP]-[4R_Mcp]-[Lys_1PEG2_1PEG2_Dap_Palm]-[N_Butyl_Phe]

TABLE 7-continued

Exemplary peptides of the invention	
Seq ID No.	Peptide
142	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_Palm]-[DIP]-[dK]-NH₂

TABLE 7-continued


Exemplary peptides of the invention	
Seq ID No.	Peptide
143	 Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_DMG_N_2ae_Palm]-[DIP]-[dK]-NH₂

TABLE 7-continued


Exemplary peptides of the invention	
Seq ID No.	Peptide
150	 Isovaleric_Acid-E-[Hyp_3R]-H-[DIP]-P-C-C-[NMe_Lys_DMG_N_2ae_Palm]-[DIP]-NH₂

TABLE 7-continued


Exemplary peptides of the invention	
Seq ID No.	Peptide
160	 Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_DMG_N_2ae_C18_Diacid_18_OMe]-[DIP]-[dK]-NH₂

TABLE 7-continued


Exemplary peptides of the invention	
Seq ID No.	Peptide
161	 Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_Alix_Palm]-[DIP]-N_Ethylamine

TABLE 7-continued

Exemplary peptides of the invention	
Seq ID No.	Peptide
163	Isovaleric Acid-E-T-H-[DIP]-P-C-[NMe_Lys_Ahx_Palm]-Diphenylethylamine

TABLE 7-continued


Exemplary peptides of the invention	
Seq ID No.	Peptide
164	 Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_Palm]-[DIP]-N_Ethylamine

TABLE 7-continued


Exemplary peptides of the invention	
Seq ID No.	Peptide
165	 Isovaleric Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_Palm]-[DIP]-[dDap]-NH₂

TABLE 7-continued


Exemplary peptides of the invention	
Seq ID No.	Peptide
168	 Isovaleric_Acid-[IleSer]-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_Palm]-[DIP]-[dK]-NH₂

TABLE 7-continued


Exemplary peptides of the invention	
Seq ID No.	Peptide
169	 Isovaleric_Acid-E-[NMe_Thr]-H-[DIP]-P-C-C-[NMe_Lys_DMKG_N_2ac_Palm]-[DIP]-NH₂

TABLE 7-continued

Exemplary peptides of the invention	
Seq ID No.	Peptide
170	Isovaleric_Acid-E-[NMe_Thr]-H-[DIP]-P-C-C-NMe_Lys_Alx_DMG_N_2ae_Palm-[DIP]-[dK]-NH₂

TABLE 7-continued

Exemplary peptides of the invention	
Seq ID No.	Peptide
172	Isovaleric_Acid-E-[NMe_Thr]-H-[DIP]-P-C-C-[NMe_Lys_Alix_Palm]-[DIP]-NH₂

TABLE 7-continued

Exemplary peptides of the invention	
Seq ID No.	Peptide
173	Isovaleric_Acid-E-[NMe_Thr]-H-[DIP]-P-C-C-[NMe_Lys_Ahx_Palm]-[DIP]-[dK]-NH₂

TABLE 7-continued


Exemplary peptides of the invention	
Seq ID No.	Peptide
175	 Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[NMe_Lys_DMG_N_2ae_Palm]-[DIP]-[dK]-NH₂

TABLE 7-continued

Exemplary peptides of the invention	
Seq ID No.	Peptide
193	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_Alix_DMG_N_2ae_C18_Diacid]-[DIP]-[dK]-NH₂

TABLE 7-continued


Exemplary peptides of the invention	
Seq ID No.	Peptide
194	 Isovaleric_Acid-E-T-H-[DIP]-P—C—C—[NMe_Lys_DMG_N_2ae_Palm]-[DIP]-[dK]-NH₂

TABLE 7-continued


Exemplary peptides of the invention	
Seq ID No.	Peptide
195	 Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_DMG_N_2ae_IsoGlu_Palm]-[DIP]-[dK]-NH₂

TABLE 7-continued

Exemplary peptides of the invention	
Seq ID No.	Peptide
196	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_Palm]-[DIP]-[dK]-NH₂

TABLE 7-continued


Exemplary peptides of the invention	
Seq ID No.	Peptide
197	

TABLE 7-continued


Exemplary peptides of the invention	
Seq ID No.	Peptide
198	 Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_Alix_DMG_N_2ac_C18_Diacid]-[DIP]-[dK]-NH₂

TABLE 7-continued

Exemplary peptides of the invention	
Seq ID No.	Peptide
210	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[Lys-Alrx_DMG_N_2ae_C18_Diacid]-[DIP]-[dK]-NH₂

TABLE 7-continued


Exemplary peptides of the invention	
Seq ID No.	Peptide
218	 Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH₂

TABLE 7-continued

Exemplary peptides of the invention	
Seq ID No.	Peptide
223	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_IsoGlu_Palm]-[hhF]-[dK]-NH₂

TABLE 7-continued

Exemplary peptides of the invention	
Seq ID No.	Peptide
242	Isovaleric_Acid-Aad-T-[3Pal]-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[DIP]-[dK]-NH₂

TABLE 7-continued


Exemplary peptides of the invention	
Seq ID No.	Peptide
246	 Isovaleric_Acid-[Aad]-T-[3Pal]-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhf]-[dK]-NH₂

TABLE 7-continued


Exemplary peptides of the invention	
Seq ID No.	Peptide
247	 Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-[NMe_dSer]-NH₂

TABLE 7-continued


Exemplary peptides of the invention	
Seq ID No.	Peptide
249	 Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe Lys 1PEG2 1PEG2 Dap C18 Diacid]-[bifF]-[dK]-[NMe dLeu]-NH₂

TABLE 7-continued


Exemplary peptides of the invention	
Seq ID No.	Peptide
250	 Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-[NMe_dPhe]-NH₂

TABLE 7-continued


Exemplary peptides of the invention	
Seq ID No.	Peptide
275	 Isovaleric_Acid-E-TH-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[DIP]-[NMe_Lys]-NH₂

TABLE 7-continued


Exemplary peptides of the invention	
Seq ID No.	Peptide
288	 Isovaleric_Acid-[Tet1]-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bbF]-[dK]-NH₂

TABLE 7-continued

Exemplary peptides of the invention	
Seq ID No.	Peptide
307	Isoleucine_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_Palm]-[bhF]-[dK]-NH₂

TABLE 7-continued


Exemplary peptides of the invention	
Seq ID No.	Peptide
308	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_Palm]-[bif]-[dK]-NH₂

[0473] In some embodiments, the invention provides a peptide set forth in Table 7, or a pharmaceutical acceptable salt and solvate thereof.

[0474] In some embodiments, the invention provides a peptide set forth in Table 6C, or a pharmaceutical acceptable salt and solvate thereof, wherein Cys and Cys; dCys and dCys; Cys and dCys; Cys and dPen; Cys and hCys; hcys and hCys; or hCys and Pen are cyclized to form a disulfide bond. In certain embodiments, the mercapto groups on the side chains of an amino acid pair, for example, Cys and Cys; dCys and dCys; Cys and dCys; Cys and dPen; Cys and hCys; hcys and hCys; or hCys and Pen in a peptide set forth in Table 6C are taken together to form a disulfide bond.

[0475] In some embodiments, the invention provides a peptide set forth in Table 6D, or a pharmaceutical acceptable salt and solvate thereof, wherein Cys and Cys; Cys and NMe-Cys; Cys and hCys; Cys and dCys; Cys and 4S_Mcp; Cys and 4R_Mcp; or 4S_Mcp and 4R_Mcp are cyclized to form a disulfide bond. In certain embodiments, the mercapto groups on the side chains of an amino acid pair, for example, Cys and Cys; Cys and NMe-Cys; Cys and hCys; Cys and dCys; Cys and 4S_Mcp; Cys and 4R_Mcp; or 4S_Mcp and 4R_Mcp in a peptide set forth in Table 6D are taken together to form a disulfide bond.

[0476] In some embodiments, the invention provides a peptide set forth in Table 6E, or a pharmaceutical acceptable salt and solvate thereof, wherein Cys and Cys; Cys and Pen; Cys and hCys; hcys and hCys; Cys and NMe-Cys; or Pen and Pen in the peptide are cyclized to form a disulfide bond. In certain embodiments, the mercapto groups on the side chains of an amino acid pair, for example, Cys and Cys; Cys and Pen; Cys and hCys; hcys and hCys; Cys and NMe-Cys; or Pen and Pen in a peptide set forth in Table 6D are taken together to form a disulfide bond. In some embodiments, the invention provides a peptide set forth in Table 6E, wherein (i) Cys and Cys in the peptide are cyclized to form a disulfide bond and (ii) the carboxy group on the side chain of Glu and the amino group on the side chain of dLys in the peptide are taken together to form an amide linkage, i.e., $-C(O)NH-$ linkage. In certain embodiments, the invention provides a peptide set forth in Table 6E, wherein (i) Cys and Cys in the peptide are cyclized to form a disulfide bond and (ii) the carboxy group on the side chain of Glu and the amino group of N-Ethylamine are taken together to form an amide linkage, i.e., $-C(O)NH-$ linkage. In other embodiments, the invention provides a peptide set forth in Table 6E, wherein Cys and Cys in the peptide are cyclized through a p-xylene linker to form a



linkage.

[0477] In one embodiment, R^2 is NH_2 . In another embodiment, R^2 is substituted amino. In another embodiment, R^2 is alkylamino or (substituted alkyl)amino. In another embodiment, R^2 is methylamino, ethylamino, propylamino, benzylamino, or phenethylamino.

[0478] In one embodiment, R^2 is OH.

[0479] In one embodiment, R^1 is C_1 - C_{20} alkanoyl.

[0480] In one embodiment, R^1 is IVA or isovaleric acid.

[0481] In certain embodiments of any of the peptide analogues having any of the various Formulae set forth herein, R^1 is selected from methyl, acetyl, formyl, benzoyl, trifluoroacetyl, isovaleryl, isobutyryl, octanoyl, and conjugated amides of lauric acid, hexadecanoic acid, and γ -Glu-hexadecanoic acid.

[0482] In certain embodiments, substituted Lys is Lys substituted with Ac, PEG, Ahx, isoGlu, C_{10} - C_{20} alkanoyl, PEG-Ahx, PEG-isoGlu, Ahx- C_{10} - C_{20} alkanoyl, isoGlu- C_{10} - C_{20} alkanoyl, PEG-Ahx- C_{10} - C_{20} alkanoyl, PEG-isoGlu- C_{10} - C_{20} alkanoyl, or any of the others described herein. In one embodiment, Lys is substituted to N^ϵ of Lys.

[0483] In certain embodiments, substituted (D)Lys is (D)Lys substituted with Ac, PEG, Ahx, isoGlu, C_{10} - C_{20} alkanoyl, PEG-Ahx, PEG-isoGlu, Ahx- C_{10} - C_{20} alkanoyl, isoGlu- C_{10} - C_{20} alkanoyl, PEG-Ahx- C_{10} - C_{20} alkanoyl, PEG-isoGlu- C_{10} - C_{20} alkanoyl, or any of the others described herein. In one embodiment, (D)Lys is substituted to N^ϵ of (D)Lys.

[0484] In certain embodiment, C_{10} - C_{20} alkanoyl is Palm.

[0485] In certain embodiment, the present invention includes a polypeptide comprising an amino acid sequence set forth in Tables 6A-B, or having any amino acid sequence with at least 85%, at least 90%, at least 92%, at least 94%, or at least 95% identity to any of these amino acid sequences.

[0486] In certain embodiment, the present invention includes a hepcidin analogue having a structure or comprising an amino acid sequence set forth below:

[0487] In certain embodiment, the present invention includes a hepcidin analogue having a structure or comprising an amino acid sequence set forth below:

[0488] In a particular embodiment, the peptide is any one of peptides wherein the FPN activity is <100 nM. In another particular embodiment, the peptide is any one of peptides wherein the FPN activity is <50 nM. In another particular embodiment, the peptide is any one of peptides wherein the FPN activity is <20 nM. In another particular embodiment, the peptide is any one of peptides wherein the FPN activity is <10 nM. In more particular embodiment, the peptide is any one of peptides wherein the FPN activity is <5 nM.

Peptide Analogue Conjugates

[0489] In certain embodiments, hepcidin analogues of the present invention, including both monomers and dimers, comprise one or more conjugated chemical substituents, such as lipophilic substituents and polymeric moieties, collectively referred to herein as half-life extension moieties. Without wishing to be bound by any particular theory, it is believed that the lipophilic substituent binds to albumin in the bloodstream, thereby shielding the hepcidin analogue from enzymatic degradation, and thus enhancing its half-life. In addition, it is believed that polymeric moieties enhance half-life and reduce clearance in the bloodstream, and in some cases enhance permeability through the epithelium and retention in the lamina propria. Moreover, it is also surmised that these substituents in some cases may enhance permeability through the epithelium and retention in the lamina propria. The skilled person will be well aware of suitable techniques for preparing the compounds employed in the context of the invention. For examples of non-limiting suitable chemistry, see, e.g., WO98/08871, WO00/55184,

WO00/55119, Madsen et al. (J. Med. Chem. 2007, 50, 6126-32), and Knudsen et al. 2000 (J. Med. Chem. 43, 1664-1669).

[0490] In one embodiment, the side chains of one or more amino acid residues (e.g., Lys residues) in a hepcidin analogue of the invention is further conjugated (e.g., covalently attached) to a lipophilic substituent or other half-life extension moiety. The lipophilic substituent may be covalently bonded to an atom in the amino acid side chain, or alternatively may be conjugated to the amino acid side chain via one or more spacers or linker moieties. The spacer or linker moiety, when present, may provide spacing between the hepcidin analogue and the lipophilic substituent.

[0491] In certain embodiments, the lipophilic substituent or half-life extension moiety comprises a hydrocarbon chain having from 4 to 30 C atoms, for example at least 8 or 12 C atoms, and preferably 24 C atoms or fewer, or 20 C atoms or fewer. The hydrocarbon chain may be linear or branched and may be saturated or unsaturated. In certain embodiments, the hydrocarbon chain is substituted with a moiety which forms part of the attachment to the amino acid side chain or the spacer, for example an acyl group, a sulfonyl group, an N atom, an O atom or an S atom. In some embodiments, the hydrocarbon chain is substituted with an acyl group, and accordingly the hydrocarbon chain may form part of an alkanoyl group, for example palmitoyl, caproyl, lauroyl, myristoyl or stearoyl.

[0492] A lipophilic substituent may be conjugated to any amino acid side chain in a hepcidin analogue of the invention. In certain embodiment, the amino acid side chain includes a carboxy, hydroxyl, thiol, amide or amine group, for forming an ester, a sulphonyl ester, a thioester, an amide or a sulphonamide with the spacer or lipophilic substituent. For example, the lipophilic substituent may be conjugated to Asn, Asp, Glu, Gln, His, Lys, Arg, Ser, Thr, Tyr, Trp, Cys or Dbu, Dpr or Orn. In certain embodiments, the lipophilic substituent is conjugated to Lys. An amino acid shown as Lys in any of the formula provided herein may be replaced by, e.g., Dbu, Dpr or Orn where a lipophilic substituent is added.

[0493] In further embodiments of the present invention, alternatively or additionally, the side-chains of one or more amino acid residues in a hepcidin analogue of the invention may be conjugated to a polymeric moiety or other half-life extension moiety, for example, in order to increase solubility and/or half-life in vivo (e.g., in plasma) and/or bioavailability. Such modifications are also known to reduce clearance (e.g. renal clearance) of therapeutic proteins and peptides.

[0494] As used herein, "Polyethylene glycol" or "PEG" is a polyether compound of general formula $H-(O-CH_2-CH_2)_n-OH$. PEGs are also known as polyethylene oxides (PEOs) or polyoxyethylenes (POEs), depending on their molecular weight PEO, PEE, or POG, as used herein, refers to an oligomer or polymer of ethylene oxide. The three names are chemically synonymous, but PEG has tended to refer to oligomers and polymers with a molecular mass below 20,000 g/mol, PEO to polymers with a molecular mass above 20,000 g/mol, and POE to a polymer of any molecular mass. PEG and PEO are liquids or low-melting solids, depending on their molecular weights. Throughout this disclosure, the 3 names are used indistinguishably. PEGs are prepared by polymerization of ethylene oxide and are commercially available over a wide range of molecular weights from 300 g/mol to 10,000,000 g/mol. While PEG

and PEO with different molecular weights find use in different applications, and have different physical properties (e.g., viscosity) due to chain length effects, their chemical properties are nearly identical. The polymeric moiety is preferably water-soluble (amphiphilic or hydrophilic), non-toxic, and pharmaceutically inert. Suitable polymeric moieties include polyethylene glycols (PEG), homo- or copolymers of PEG, a monomethyl-substituted polymer of PEG (mPEG), or polyoxyethylene glycerol (POG). See, for example, Int. J. Hematology 68:1 (1998); Bioconjugate Chem. 6:150 (1995); and Crit. Rev. Therap. Drug Carrier Sys. 9:249 (1992). Also encompassed are PEGs that are prepared for purpose of half-life extension, for example, mono-activated, alkoxy-terminated polyalkylene oxides (POA's) such as mono-methoxy-terminated polyethylene glycols (mPEG's); bis activated polyethylene oxides (glycols) or other PEG derivatives are also contemplated. Suitable polymers will vary substantially by weights ranging from about 200 to about 40,000 are usually selected for the purposes of the present invention. In certain embodiments, PEGs having molecular weights from 200 to 2,000 daltons or from 200 to 500 daltons are used. Different forms of PEG may also be used, depending on the initiator used for the polymerization process, e.g., a common initiator is a mono-functional methyl ether PEG, or methoxypoly(ethylene glycol), abbreviated mPEG. Other suitable initiators are known in the art and are suitable for use in the present invention.

[0495] Lower-molecular-weight PEGs are also available as pure oligomers, referred to as monodisperse, uniform, or discrete. These are used in certain embodiments of the present invention.

[0496] PEGs are also available with different geometries: branched PEGs have three to ten PEG chains emanating from a central core group; star PEGs have 10 to 100 PEG chains emanating from a central core group; and comb PEGs have multiple PEG chains normally grafted onto a polymer backbone. PEGs can also be linear. The numbers that are often included in the names of PEGs indicate their average molecular weights (e.g. a PEG with $n=9$ would have an average molecular weight of approximately 400 daltons, and would be labeled PEG 400).

[0497] As used herein, "PEGylation" is the act of coupling (e.g., covalently) a PEG structure to the hepcidin analogue of the invention, which is in certain embodiments referred to as a "PEGylated hepcidin analogue". In certain embodiments, the PEG of the PEGylated side chain is a PEG with a molecular weight from about 200 to about 40,000. In certain embodiments, the PEG portion of the conjugated half-life extension moiety is PEG3, PEG4, PEG5, PEG6, PEG7, PEG8, PEG9, PEG10, or PEG11. In particular embodiments, it is PEG11. In certain embodiments, the PEG of a PEGylated spacer is PEG3 or PEG8. In some embodiments, a spacer is PEGylated. In certain embodiments, the PEG of a PEGylated spacer is PEG3, PEG4, PEG5, PEG6, PEG7, PEG8, PEG9, PEG10, or PEG11. In certain embodiments, the PEG of a PEGylated spacer is PEG3 or PEG8.

[0498] In some embodiments, the present invention includes a hepcidin analogue peptide (or a dimer thereof) conjugated with a PEG that is attached covalently, e.g., through an amide, a thiol, via click chemistry, or via any other suitable means known in the art. In particular embodiments PEG is attached through an amide bond and, as such, certain PEG derivatives used will be appropriately functionalized. For example, in certain embodiments, PEG11, which

is O-(2-aminoethyl)-O'-(2-carboxyethyl)-undecaethyl-ene glycol, has both an amine and carboxylic acid for attachment to a peptide of the present invention. In certain embodiments, PEG25 contains a diacid and 25 glycol moieties.

[0499] Other suitable polymeric moieties include poly-amino acids such as poly-lysine, poly-aspartic acid and poly-glutamic acid (see for example Gombotz, et al. (1995), Bioconjugate Chem., vol. 6: 332-351; Hudecz, et al. (1992), Bioconjugate Chem., vol. 3, 49-57 and Tsukada, et al. (1984), J. Natl. Cancer Inst., vol. 73: 721-729. The polymeric moiety may be straight-chain or branched. In some embodiments, it has a molecular weight of 500-40,000 Da, for example 500-10,000 Da, 1000-5000 Da, 10,000-20,000 Da, or 20,000-40,000 Da.

[0500] In some embodiments, a hepcidin analogue of the invention may comprise two or more such polymeric moieties, in which case the total molecular weight of all such moieties will generally fall within the ranges provided above.

[0501] In some embodiments, the polymeric moiety may be coupled (by covalent linkage) to an amino, carboxyl or thiol group of an amino acid side chain. Certain examples are the thiol group of Cys residues and the epsilon amino group of Lys residues, and the carboxyl groups of Asp and Glu residues may also be involved.

[0502] The skilled worker will be well aware of suitable techniques which can be used to perform the coupling reaction. For example, a PEG moiety bearing a methoxy group can be coupled to a Cys thiol group by a maleimido linkage using reagents commercially available from Nektar Therapeutics AL. See also WO 2008/101017, and the references cited above, for details of suitable chemistry. A maleimide-functionalised PEG may also be conjugated to the side-chain sulfhydryl group of a Cys residue.


[0503] As used herein, disulfide bond oxidation can occur within a single step or is a two-step process. As used herein, for a single oxidation step, the trityl protecting group is often

employed during assembly, allowing deprotection during cleavage, followed by solution oxidation. When a second disulfide bond is required, one has the option of native or selective oxidation. For selective oxidation requiring orthogonal protecting groups, Acn and Trityl is used as the protecting groups for cysteine. Cleavage results in the removal of one protecting pair of cysteine allowing oxidation of this pair. The second oxidative deprotection step of the cysteine protected Acn group is then performed. For native oxidation, the trityl protecting group is used for all cysteines, allowing for natural folding of the peptide.

[0504] A skilled worker will be well aware of suitable techniques which can be used to perform the oxidation step.

[0505] In particular embodiments, a hepcidin analogue of the present invention comprises a half-life extension moiety, which may be selected from but is not limited to the following: Ahx-Palm, PEG2-Palm, PEG11-Palm, isoGlu-Palm, dapa-Palm, isoGlu-Lauric acid, isoGlu-Mysteric acid, and isoGlu-Isovaleric acid.

[0506] In particular embodiments, a hepcidin analogue comprises a half-life extension moiety having the structure shown below, wherein $n=0$ to 24 or $n=14$ to 24:



[0507] In certain embodiments, a hepcidin analogue of the present invention comprises a conjugated half-life extension moiety shown in Table 2.

TABLE 2

Illustrative Half-Life Extension Moieties	
#	Conjugates
C1	
	C12 (Lauric acid)
C2	
	C14 (Mysteric acid)
C3	
	C16 (Palm or Palmitic acid)

TABLE 2-continued










Illustrative Half-Life Extension Moieties	
#	Conjugates
C4	 C18 (Stearic acid)
C5	 C20
C6	 C12 diacid
C7	 C14 diacid
C8	 C16 diacid
C9	 C18 diacid
C10	 C20 diacid
C11	 Biotin

TABLE 2-continued

Illustrative Half-Life Extension Moieties	
#	Conjugates
C12	 Isovaleric acid

[0508] In certain embodiments, a half-life extension moiety is conjugated directly to a hepcidin analogue, while in other embodiments, a half-life extension moiety is conjugated to a hepcidin analogue peptide via a linker moiety, e.g., any of those depicted in Table 3.

TABLE 3






Illustrative Linker Moieties*	
#	Linker Moiety
L1	IsoGlu
L2	 Dapa
L3	 Ahx
L4	Lipidic based linkers:  n = 1 to 24
L5	 n = 1 to 25 —[C(O)CH₂CH₂(OCH₂CH₂)_nN(H)]— PEG based linkers (n=5-25) PEG based linkers
L6	 IsoGlu-Ahx

TABLE 3-continued

Illustrative Linker Moieties*	
#	Linker Moiety
L7	—[C(O)—CH ₂ -(Peg)2-NH]— or —[C(O)—CH ₂ —(OCH ₂ CH ₂) ₂ —NH]— (1Peg2)
L8	—[(C(O)—CH ₂ —(OCH ₂ CH ₂) ₂ —NH—C(O)—CH ₂ —(OCH ₂ CH ₂) ₂ —NH]— (1Peg2-1Peg2)
L9	—[C(O)—CH ₂ —CH ₂ -(Peg)2-NH]— or —[C(O)—CH ₂ —CH ₂ —(OCH ₂ CH ₂) ₂ —NH]— (2Peg2)
L10	—[C(O)—CH ₂ —CH ₂ -(Peg)4-NH]— or —[C(O)—CH ₂ —CH ₂ —(OCH ₂ CH ₂) ₄ —NH]— (2Peg4)
L11	—[C(O)—CH ₂ -(Peg)8-NH]— or —[C(O)—CH ₂ —(OCH ₂ CH ₂) ₈ —NH]— (1Peg8)
L12	—[C(O)—CH ₂ —CH ₂ -(Peg)8-NH]— or —[C(O)—CH ₂ —CH ₂ —(OCH ₂ CH ₂) ₈ —NH]— (2Peg8)
L17	—[C(O)—CH ₂ -(Peg)11-NH]— or —[C(O)—CH ₂ —(OCH ₂ CH ₂) ₁₁ —NH]— (1Peg11)
L18	—[C(O)—CH ₂ —CH ₂ -(Peg)11-NH]— or —[C(O)—CH ₂ —CH ₂ —(OCH ₂ CH ₂) ₁₁ —NH]— (2Peg11)
L19	—[C(O)—CH ₂ —CH ₂ -(Peg)12-NH]— or —[C(O)—CH ₂ —CH ₂ —(OCH ₂ CH ₂) ₁₂ —NH]— (2Peg11' or 2Peg12)

*(Peg) is —(OCH₂CH₂)—

[0509] With reference to linker structures shown in Table 3, reference to n=1 to 24 or n=1 to 25, or the like, (e.g., in L4, or L5) indicates that n may be any integer within the recited range. Additional linker moieties can be used are shown in “Abbreviation” table.

[0510] In particular embodiments, a hepcidin analogue of the present invention comprises any of the linker moieties shown in Table 3 and any of the half-life extension moieties shown in Table 2, including any of the following combinations shown in Table 4.

TABLE 4

Illustrative Combinations of Linkers and Half-Life Extension Moieties in Hepcidin Analogues					
Linker	Half-Life Extension Moiety	Linker	Half-Life Extension Moiety	Linker	Half-Life Extension Moiety
L1	C1	L1	C2	L1	C3
L2	C1	L2	C2	L2	C3
L3	C1	L3	C2	L3	C3
L4	C1	L4	C2	L4	C3
L5	C1	L5	C2	L5	C3
L6	C1	L6	C2	L6	C3
L7	C1	L7	C2	L7	C3
L8	C1	L8	C2	L8	C3
L9	C1	L9	C2	L9	C3
L10	C1	L10	C2	L10	C3
L11	C1	L11	C2	L11	C3
L12	C1	L12	C2	L12	C3
L13	C1	L13	C2	L13	C3
L14	C1	L14	C2	L14	C3
L15	C1	L15	C2	L15	C3
L16	C1	L16	C2	L16	C3
L17	C1	L17	C2	L17	C3
L18	C1	L18	C2	L18	C3
L19	C1	L19	C2	L18	C3
L1	C4	L1	C5	L1	C6
L2	C4	L2	C5	L2	C6
L3	C4	L3	C5	L3	C6

TABLE 4-continued

Illustrative Combinations of Linkers and Half-Life Extension Moieties in Hepcidin Analogues					
Linker	Half-Life Extension Moiety	Linker	Half-Life Extension Moiety	Linker	Half-Life Extension Moiety
L4	C4	L4	C5	L4	C6
L5	C4	L5	C5	L5	C6
L6	C4	L6	C5	L6	C6
L7	C4	L7	C5	L7	C6
L8	C4	L8	C5	L8	C6
L9	C4	L9	C5	L9	C6
L10	C4	L10	C5	L10	C6
L11	C4	L11	C5	L11	C6
L12	C4	L12	C5	L12	C6
L13	C4	L13	C5	L13	C6
L14	C4	L14	C5	L14	C6
L15	C4	L15	C5	L15	C6
L16	C4	L16	C5	L16	C6
L17	C4	L17	C5	L17	C6
L18	C4	L18	C5	L18	C6
L19	C4	L19	C5	L18	C6
L1	C7	L1	C8	L1	C9
L2	C7	L2	C8	L2	C9
L3	C7	L3	C8	L3	C9
L4	C7	L4	C8	L4	C9
L5	C7	L5	C8	L5	C9
L6	C7	L6	C8	L6	C9
L7	C7	L7	C8	L7	C9
L8	C7	L8	C8	L8	C9
L9	C7	L9	C8	L9	C9
L10	C7	L10	C8	L10	C9
L11	C7	L11	C8	L11	C9
L12	C7	L12	C8	L12	C9
L13	C7	L13	C8	L13	C9
L14	C7	L14	C8	L14	C9
L15	C7	L15	C8	L15	C9
L16	C7	L16	C8	L16	C9
L17	C7	L17	C8	L17	C9
L18	C7	L18	C8	L18	C9
L19	C7	L19	C8	L18	C9

TABLE 4-continued

Illustrative Combinations of Linkers and Half-Life Extension Moieties in Hepcidin Analogues					
Linker	Half-Life Extension Moieties	Linker	Half-Life Extension Moieties	Linker	Half-Life Extension Moieties
L1	C10	L1	C11	L1	C12
L2	C10	L2	C11	L2	C12
L3	C10	L3	C11	L3	C12
L4	C10	L4	C11	L4	C12
L5	C10	L5	C11	L5	C12
L6	C10	L6	C11	L6	C12
L7	C10	L7	C11	L7	C12
L8	C10	L8	C11	L8	C12
L9	C10	L9	C11	L9	C12
L10	C10	L10	C11	L10	C12
L11	C10	L11	C11	L11	C12
L12	C10	L12	C11	L12	C12
L13	C10	L13	C11	L13	C12
L14	C10	L14	C11	L14	C12
L15	C10	L15	C11	L15	C12
L16	C10	L16	C11	L16	C12
L17	C10	L17	C11	L17	C12
L18	C10	L18	C11	L18	C12
L19	C10	L19	C11	L18	C12

[0511] In certain embodiments, a hepcidin analogue comprises two or more linkers. In particular embodiments, the two or more linkers are concatamerized, i.e., bound to each other.

[0512] In related embodiments, the present invention includes polynucleotides that encode a polypeptide having a peptide sequence present in any of the hepcidin analogues described herein.

[0513] In addition, the present invention includes vectors, e.g., expression vectors, comprising a polynucleotide of the present invention.

Methods of Treatment

[0514] In some embodiments, the present invention provides methods for treating a subject afflicted with a disease or disorder associated with dysregulated hepcidin signaling, wherein the method comprises administering to the subject a hepcidin analogue of the present invention. In some embodiments, the hepcidin analogue that is administered to the subject is present in a composition (e.g., a pharmaceutical composition). In one embodiment, a method is provided for treating a subject afflicted with a disease or disorder characterized by increased activity or expression of ferroportin, wherein the method comprises administering to the individual a hepcidin analogue or composition of the present invention in an amount sufficient to (partially or fully) bind to and agonize ferroportin or mimic hepcidin in the subject. In one embodiment, a method is provided for treating a subject afflicted with a disease or disorder characterized by dysregulated iron metabolism, wherein the method comprises administering to the subject a hepcidin analogue or composition of the present invention.

[0515] In some embodiments, methods of the present invention comprise providing a hepcidin analogue or a composition of the present invention to a subject in need thereof. In particular embodiments, the subject in need thereof has been diagnosed with or has been determined to be at risk of developing a disease or disorder characterized by dysregulated iron levels (e.g., diseases or disorders of iron metabolism; diseases or disorders related to iron over-

load; and diseases or disorders related to abnormal hepcidin activity or expression). In particular embodiments, the subject is a mammal (e.g., a human).

[0516] In certain embodiments, the disease or disorder is a disease of iron metabolism, such as, e.g., an iron overload disease, iron deficiency disorder, disorder of iron biodistribution, or another disorder of iron metabolism and other disorder potentially related to iron metabolism, etc. In particular embodiments, the disease of iron metabolism is hemochromatosis, HFE mutation hemochromatosis, ferroportin mutation hemochromatosis, transferrin receptor 2 mutation hemochromatosis, hemojuvelin mutation hemochromatosis, hepcidin mutation hemochromatosis, juvenile hemochromatosis, neonatal hemochromatosis, hepcidin deficiency, transfusional iron overload, thalassemia, thalassemia intermedia, alpha thalassemia, beta thalassemia, sideroblastic anemia, porphyria, porphyria cutanea tarda, African iron overload, hyperferritinemia, ceruloplasmin deficiency, atransferrinemia, congenital dyserythropoietic anemia, hypochromic microcytic anemia, sickle cell disease, polycythemia vera (primary and secondary), secondary erythrocytoses, such as Chronic obstructive pulmonary disease (COPD), post-renal transplant, Chuvash, HIF and PHD mutations, and idiopathic, myelodysplasia, pyruvate kinase deficiency, hypochromic microcytic anemia, transfusion-dependent anemia, hemolytic anemia, iron deficiency of obesity, other anemias, benign or malignant tumors that overproduce hepcidin or induce its overproduction, conditions with hepcidin excess, Friedreich ataxia, gracile syndrome, Hallervorden-Spatz disease, Wilson's disease, pulmonary hemosiderosis, hepatocellular carcinoma, cancer (e.g., liver cancer), hepatitis, cirrhosis of liver, pica, chronic renal failure, insulin resistance, diabetes, atherosclerosis, neurodegenerative disorders, dementia, multiple sclerosis, Parkinson's disease, Huntington's disease, or Alzheimer's disease.

[0517] In certain embodiments, the disease or disorder is related to iron overload diseases such as iron hemochromatosis, HFE mutation hemochromatosis, ferroportin mutation hemochromatosis, transferrin receptor 2 mutation hemochromatosis, hemojuvelin mutation hemochromatosis, hepcidin mutation hemochromatosis, juvenile hemochromatosis, neonatal hemochromatosis, hepcidin deficiency, transfusional iron overload, thalassemia, thalassemia intermedia, alpha thalassemia, sickle cell disease, myelodysplasia, sideroblastic infections, diabetic retinopathy, and pyruvate kinase deficiency.

[0518] In certain embodiments, the disease or disorder is one that is not typically identified as being iron related. For example, hepcidin is highly expressed in the murine pancreas suggesting that diabetes (Type I or Type II), insulin resistance, glucose intolerance and other disorders may be ameliorated by treating underlying iron metabolism disorders. See Ilyin, G. et al. (2003) FEBS Lett. 542 22-26, which is herein incorporated by reference. As such, peptides of the invention may be used to treat these diseases and conditions. Those skilled in the art are readily able to determine whether a given disease can be treated with a peptide according to the present invention using methods known in the art, including the assays of WO 2004092405, which is herein incorporated by reference, and assays which monitor hepcidin, hemojuvelin, or iron levels and expression, which are known in the art such as those described in U.S. Pat. No. 7,534,764, which is herein incorporated by reference.

[0519] In certain embodiments, the disease or disorder is postmenopausal osteoporosis.

[0520] In certain embodiments of the present invention, the diseases of iron metabolism are iron overload diseases, which include hereditary hemochromatosis, iron-loading anemias, alcoholic liver diseases, heart disease and/or failure, cardiomyopathy, and chronic hepatitis C.

[0521] In particular embodiments, any of these diseases, disorders, or indications are caused by or associated with a deficiency of hepcidin or iron overload.

[0522] In some embodiments, methods of the present invention comprise providing a hepcidin analogue of the present invention (i.e., a first therapeutic agent) to a subject in need thereof in combination with a second therapeutic agent. In certain embodiments, the second therapeutic agent is provided to the subject before and/or simultaneously with and/or after the pharmaceutical composition is administered to the subject. In particular embodiments, the second therapeutic agent is iron chelator. In certain embodiments, the second therapeutic agent is selected from the iron chelators Deferoxamine and Deferasirox (Exjade™). In another embodiment, the method comprises administering to the subject a third therapeutic agent.

[0523] The present invention provides compositions (for example pharmaceutical compositions) comprising one or more hepcidin analogues of the present invention and a pharmaceutically acceptable carrier, excipient or diluent. A pharmaceutically acceptable carrier, diluent or excipient refers to a non-toxic solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. Prevention of the action of microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents such as sugars, sodium chloride, and the like.

[0524] The term “pharmaceutically acceptable carrier” includes any of the standard pharmaceutical carriers. Pharmaceutically acceptable carriers for therapeutic use are well known in the pharmaceutical art and are described, for example, in “Remington’s Pharmaceutical Sciences”, 17th edition, Alfonso R. Gennaro (Ed.), Mark Publishing Company, Easton, PA, USA, 1985. For example, sterile saline and phosphate-buffered saline at slightly acidic or physiological pH may be used. Suitable pH-buffering agents may, e.g., be phosphate, citrate, acetate, tris(hydroxymethyl)aminomethane (TRIS), N-tris(hydroxymethyl)methyl-3-aminopropanesulfonic acid (TAPS), ammonium bicarbonate, diethanolamine, histidine, arginine, lysine or acetate (e.g. as sodium acetate), or mixtures thereof. The term further encompasses any carrier agents listed in the US Pharmacopeia for use in animals, including humans.

[0525] In certain embodiments, the compositions comprise two or more hepcidin analogues disclosed herein. In certain embodiments, the combination is selected from one of the following: (i) any two or more of the hepcidin analogue peptide monomers shown therein; (ii) any two or more of the hepcidin analogue peptide dimers disclosed herein; (iii) any one or more of the hepcidin analogue peptide monomers disclosed herein, and any one or more of the hepcidin analogue peptide dimers disclosed herein.

[0526] It is to be understood that the inclusion of a hepcidin analogue of the invention (i.e., one or more hepcidin analogue peptide monomers of the invention or one or more hepcidin analogue peptide dimers of the present inven-

tion) in a pharmaceutical composition also encompasses inclusion of a pharmaceutically acceptable salt or solvate of a hepcidin analogue of the invention. In particular embodiments, the pharmaceutical compositions further comprise one or more pharmaceutically acceptable carrier, excipient, or vehicle.

[0527] In certain embodiments, the invention provides a pharmaceutical composition comprising a hepcidin analogue, or a pharmaceutically acceptable salt or solvate thereof, for treating a variety of conditions, diseases, or disorders as disclosed herein or elsewhere (see, e.g., Methods of Treatment, herein). In particular embodiments, the invention provides a pharmaceutical composition comprising a hepcidin analogue peptide monomer, or a pharmaceutically acceptable salt or solvate thereof, for treating a variety of conditions, diseases, or disorders as disclosed herein elsewhere (see, e.g., Methods of Treatment, herein). In particular embodiments, the invention provides a pharmaceutical composition comprising a hepcidin analogue peptide dimer, or a pharmaceutically acceptable salt or solvate thereof, for treating a variety of conditions, diseases, or disorders as disclosed herein.

[0528] The hepcidin analogues of the present invention may be formulated as pharmaceutical compositions which are suited for administration with or without storage, and which typically comprise a therapeutically effective amount of at least one hepcidin analogue of the invention, together with a pharmaceutically acceptable carrier, excipient or vehicle.

[0529] In some embodiments, the hepcidin analogue pharmaceutical compositions of the invention are in unit dosage form. In such forms, the composition is divided into unit doses containing appropriate quantities of the active component or components. The unit dosage form may be presented as a packaged preparation, the package containing discrete quantities of the preparation, for example, packaged tablets, capsules or powders in vials or ampoules. The unit dosage form may also be, e.g., a capsule, cachet or tablet in itself, or it may be an appropriate number of any of these packaged forms. A unit dosage form may also be provided in single-dose injectable form, for example in the form of a pen device containing a liquid-phase (typically aqueous) composition. Compositions may be formulated for any suitable route and means of administration, e.g., any one of the routes and means of administration disclosed herein.

[0530] In particular embodiments, the hepcidin analogue, or the pharmaceutical composition comprising a hepcidin analogue, is suspended in a sustained-release matrix. A sustained-release matrix, as used herein, is a matrix made of materials, usually polymers, which are degradable by enzymatic or acid-base hydrolysis or by dissolution. Once inserted into the body, the matrix is acted upon by enzymes and body fluids. A sustained-release matrix desirably is chosen from biocompatible materials such as liposomes, polylactides (polylactic acid), polyglycolide (polymer of glycolic acid), polylactide co-glycolide (copolymers of lactic acid and glycolic acid) polyanhydrides, poly(ortho)esters, polypeptides, hyaluronic acid, collagen, chondroitin sulfate, carboxylic acids, fatty acids, phospholipids, polysaccharides, nucleic acids, polyamino acids, amino acids such as phenylalanine, tyrosine, isoleucine, polynucleotides, polyvinyl propylene, polyvinylpyrrolidone and silicone. One embodiment of a biodegradable matrix is a matrix of

one of either polylactide, polyglycolide, or polylactide co-glycolide (co-polymers of lactic acid and glycolic acid).

[0531] In certain embodiments, the compositions are administered parenterally, subcutaneously or orally. In particular embodiments, the compositions are administered orally, intracisternally, intravaginally, intraperitoneally, intrarectally, topically (as by powders, ointments, drops, suppository, or transdermal patch, including delivery intravitreally, intranasally, and via inhalation) or buccally. The term “parenteral” as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous, intradermal and intra-articular injection and infusion. Accordingly, in certain embodiments, the compositions are formulated for delivery by any of these routes of administration.

[0532] In certain embodiments, pharmaceutical compositions for parenteral injection comprise pharmaceutically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, or sterile powders, for reconstitution into sterile injectable solutions or dispersions just prior to use. Examples of suitable aqueous and non-aqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), carboxymethylcellulose and suitable mixtures thereof, beta-cyclodextrin, vegetable oils (such as olive oil), and injectable organic esters such as ethyl oleate. Proper fluidity may be maintained, for example, by the use of coating materials such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants. These compositions may also contain adjuvants such as preservative, wetting agents, emulsifying agents, and dispersing agents. Prolonged absorption of an injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption, such as aluminum monostearate and gelatin.

[0533] Injectable depot forms include those made by forming microcapsule matrices of the hepcidin analogue in one or more biodegradable polymers such as polylactide-polyglycolide, poly(orthoesters), poly(anhydrides), and (poly)glycols, such as PEG. Depending upon the ratio of peptide to polymer and the nature of the particular polymer employed, the rate of release of the hepcidin analogue can be controlled. Depot injectable formulations are also prepared by entrapping the hepcidin analogue in liposomes or microemulsions compatible with body tissues.

[0534] The injectable formulations may be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium just prior to use.

[0535] Hepcidin analogues of the present invention may also be administered in liposomes or other lipid-based carriers. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically acceptable and metabolizable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to a hepcidin analogue of the present invention, stabilizers, preservatives, excipients, and the like. In certain embodiments, the lipids comprise phospholipids, including the phosphatidyl cholines (lecithins) and serines, both natural and synthetic. Methods to form liposomes are known in the art.

[0536] Pharmaceutical compositions to be used in the invention suitable for parenteral administration may comprise sterile aqueous solutions and/or suspensions of the peptide inhibitors made isotonic with the blood of the recipient, generally using sodium chloride, glycerin, glucose, mannitol, sorbitol, and the like.

[0537] In some aspects, the invention provides a pharmaceutical composition for oral delivery. Compositions and hepcidin analogues of the instant invention may be prepared for oral administration according to any of the methods, techniques, and/or delivery vehicles described herein. Further, one having skill in the art will appreciate that the hepcidin analogues of the instant invention may be modified or integrated into a system or delivery vehicle that is not disclosed herein, yet is well known in the art and compatible for use in oral delivery of peptides.

[0538] In certain embodiments, formulations for oral administration may comprise adjuvants (e.g. resorcinols and/or nonionic surfactants such as polyoxyethylene oleyl ether and n-hexadecylpolyethylene ether) to artificially increase the permeability of the intestinal walls, and/or enzymatic inhibitors (e.g. pancreatic trypsin inhibitors, diisopropylfluorophosphate (DFF) or trasylol) to inhibit enzymatic degradation. In certain embodiments, the hepcidin analogue of a solid-type dosage form for oral administration can be mixed with at least one additive, such as sucrose, lactose, cellulose, mannitol, trehalose, raffinose, maltitol, dextran, starches, agar, alginates, chitins, chitosans, pectins, gum tragacanth, gum arabic, gelatin, collagen, casein, albumin, synthetic or semisynthetic polymer, or glyceride. These dosage forms can also contain other type(s) of additives, e.g., inactive diluting agent, lubricant such as magnesium stearate, paraben, preserving agent such as sorbic acid, ascorbic acid, alpha-tocopherol, antioxidants such as cysteine, disintegrators, binders, thickeners, buffering agents, pH adjusting agents, sweetening agents, flavoring agents or perfuming agents.

[0539] In particular embodiments, oral dosage forms or unit doses compatible for use with the hepcidin analogues of the present invention may include a mixture of hepcidin analogue and nondrug components or excipients, as well as other non-reusable materials that may be considered either as an ingredient or packaging. Oral compositions may include at least one of a liquid, a solid, and a semi-solid dosage forms. In some embodiments, an oral dosage form is provided comprising an effective amount of hepcidin analogue, wherein the dosage form comprises at least one of a pill, a tablet, a capsule, a gel, a paste, a drink, a syrup, ointment, and suppository. In some instances, an oral dosage form is provided that is designed and configured to achieve delayed release of the hepcidin analogue in the subject's small intestine and/or colon.

[0540] In one embodiment, an oral pharmaceutical composition comprising a hepcidin analogue of the present invention comprises an enteric coating that is designed to delay release of the hepcidin analogue in the small intestine. In at least some embodiments, a pharmaceutical composition is provided which comprises a hepcidin analogue of the present invention and a protease inhibitor, such as aprotinin, in a delayed release pharmaceutical formulation. In some instances, pharmaceutical compositions of the instant invention comprise an enteric coat that is soluble in gastric juice at a pH of about 5.0 or higher. In at least one embodiment, a pharmaceutical composition is provided comprising an

enteric coating comprising a polymer having dissociable carboxylic groups, such as derivatives of cellulose, including hydroxypropylmethyl cellulose phthalate, cellulose acetate phthalate and cellulose acetate trimellitate and similar derivatives of cellulose and other carbohydrate polymers.

[0541] In one embodiment, a pharmaceutical composition comprising a hepcidin analogue of the present invention is provided in an enteric coating, the enteric coating being designed to protect and release the pharmaceutical composition in a controlled manner within the subject's lower gastrointestinal system, and to avoid systemic side effects. In addition to enteric coatings, the hepcidin analogues of the instant invention may be encapsulated, coated, engaged or otherwise associated within any compatible oral drug delivery system or component. For example, in some embodiments a hepcidin analogue of the present invention is provided in a lipid carrier system comprising at least one of polymeric hydrogels, nanoparticles, microspheres, micelles, and other lipid systems.

[0542] To overcome peptide degradation in the small intestine, some embodiments of the present invention comprise a hydrogel polymer carrier system in which a hepcidin analogue of the present invention is contained, whereby the hydrogel polymer protects the hepcidin analogue from proteolysis in the small intestine and/or colon. The hepcidin analogues of the present invention may further be formulated for compatible use with a carrier system that is designed to increase the dissolution kinetics and enhance intestinal absorption of the peptide. These methods include the use of liposomes, micelles and nanoparticles to increase GI tract permeation of peptides.

[0543] Various bioresponsive systems may also be combined with one or more hepcidin analogue of the present invention to provide a pharmaceutical agent for oral delivery. In some embodiments, a hepcidin analogue of the instant invention is used in combination with a bioresponsive system, such as hydrogels and mucoadhesive polymers with hydrogen bonding groups (e.g., PEG, poly(methacrylic) acid [PMAA], cellulose, Eudragit®, chitosan and alginate) to provide a therapeutic agent for oral administration. Other embodiments include a method for optimizing or prolonging drug residence time for a hepcidin analogue disclosed herein, wherein the surface of the hepcidin analogue surface is modified to comprise mucoadhesive properties through hydrogen bonds, polymers with linked mucins or/and hydrophobic interactions. These modified peptide molecules may demonstrate increase drug residence time within the subject, in accordance with a desired feature of the invention. Moreover, targeted mucoadhesive systems may specifically bind to receptors at the enterocytes and M-cell surfaces, thereby further increasing the uptake of particles containing the hepcidin analogue.

[0544] Other embodiments comprise a method for oral delivery of a hepcidin analogue of the present invention, wherein the hepcidin analogue is provided to a subject in combination with permeation enhancers that promote the transport of the peptides across the intestinal mucosa by increasing paracellular or transcellular permeation. For example, in one embodiment, a permeation enhancer is combined with a hepcidin analogue, wherein the permeation enhancer comprises at least one of a long-chain fatty acid, a bile salt, an amphiphilic surfactant, and a chelating agent. In one embodiment, a permeation enhancer comprising sodium N-[hydroxybenzoyl]amino]caprylate is used to form a weak

noncovalent association with the hepcidin analogue of the instant invention, wherein the permeation enhancer favors membrane transport and further dissociation once reaching the blood circulation. In another embodiment, a hepcidin analogue of the present invention is conjugated to oligoarginine, thereby increasing cellular penetration of the peptide into various cell types. Further, in at least one embodiment a noncovalent bond is provided between a peptide inhibitor of the present invention and a permeation enhancer selected from the group consisting of a cyclodextrin (CD) and a dendrimers, wherein the permeation enhancer reduces peptide aggregation and increasing stability and solubility for the hepcidin analogue molecule.

[0545] Other embodiments of the invention provide a method for treating a subject with a hepcidin analogue of the present invention having an increased half-life. In one aspect, the present invention provides a hepcidin analogue having a half-life of at least several hours to one day in vitro or in vivo (e.g., when administered to a human subject) sufficient for daily (q.d.) or twice daily (b.i.d.) dosing of a therapeutically effective amount. In another embodiment, the hepcidin analogue has a half-life of three days or longer sufficient for weekly (q.w.) dosing of a therapeutically effective amount. Further, in another embodiment, the hepcidin analogue has a half-life of eight days or longer sufficient for bi-weekly (b.i.w.) or monthly dosing of a therapeutically effective amount. In another embodiment, the hepcidin analogue is derivatized or modified such that it has a longer half-life as compared to the underivatized or unmodified hepcidin analogue. In another embodiment, the hepcidin analogue contains one or more chemical modifications to increase serum half-life.

[0546] When used in at least one of the treatments or delivery systems described herein, a hepcidin analogue of the present invention may be employed in pure form or, where such forms exist, in pharmaceutically acceptable salt form.

Dosages

[0547] The total daily usage of the hepcidin analogues and compositions of the present invention can be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular subject will depend upon a variety of factors including: a) the disorder being treated and the severity of the disorder; b) activity of the specific compound employed; c) the specific composition employed, the age, body weight, general health, sex and diet of the patient; d) the time of administration, route of administration, and rate of excretion of the specific hepcidin analogue employed; e) the duration of the treatment; f) drugs used in combination or coincidental with the specific hepcidin analogue employed, and like factors well known in the medical arts.

[0548] In particular embodiments, the total daily dose of the hepcidin analogues of the invention to be administered to a human or other mammal host in single or divided doses may be in amounts, for example, from 0.0001 to 300 mg/kg body weight daily or 1 to 300 mg/kg body weight daily. In certain embodiments, a dosage of a hepcidin analogue of the present invention is in the range from about 0.0001 to about 100 mg/kg body weight per day, such as from about 0.0005 to about 50 mg/kg body weight per day, such as from about 0.001 to about 10 mg/kg body weight per day, e.g. from about 0.01 to about 1 mg/kg body weight per day, admin-

istered in one or more doses, such as from one to three doses. In particular embodiments, a total dosage is about 1 mg, about 2 mg, about 3 mg, about 4 mg, about 5 mg, about 6 mg, about 7 mg, about 8 mg, about 9 mg, or about 10 mg about once or twice weekly, e.g., for a human patient. In particular embodiments, the total dosage is in the range of about 1 mg to about 5 mg, or about 1 mg to about 3 mg, or about 2 mg to about 3 mg per human patient, e.g., about once weekly.

[0549] In various embodiments, a hepcidin analogue of the invention may be administered continuously (e.g. by intravenous administration or another continuous drug administration method), or may be administered to a subject at intervals, typically at regular time intervals, depending on the desired dosage and the pharmaceutical composition selected by the skilled practitioner for the particular subject. Regular administration dosing intervals include, e.g., once daily, twice daily, once every two, three, four, five or six days, once or twice weekly, once or twice monthly, and the like.

[0550] Such regular hepcidin analogue administration regimens of the invention may, in certain circumstances such as, e.g., during chronic long-term administration, be advantageously interrupted for a period of time so that the medicated subject reduces the level of or stops taking the medication, often referred to as taking a “drug holiday.” Drug holidays are useful for, e.g., maintaining or regaining sensitivity to a drug especially during long-term chronic treatment, or to reduce unwanted side-effects of long-term chronic treatment of the subject with the drug. The timing of a drug holiday depends on the timing of the regular dosing regimen and the purpose for taking the drug holiday (e.g., to regain drug sensitivity and/or to reduce unwanted side effects of continuous, long-term administration). In some embodiments, the drug holiday may be a reduction in the dosage of the drug (e.g. to below the therapeutically effective amount for a certain interval of time). In other embodiments, administration of the drug is stopped for a certain interval of time before administration is started again using the same or a different dosing regimen (e.g. at a lower or higher dose and/or frequency of administration). A drug holiday of the invention may thus be selected from a wide range of time-periods and dosage regimens. An exemplary drug holiday is two or more days, one or more weeks, or one or more months, up to about 24 months of drug holiday. So, for example, a regular daily dosing regimen with a peptide, a peptide analogue, or a dimer of the invention may, for example, be interrupted by a drug holiday of a week, or two weeks, or four weeks, after which time the preceding, regular dosage regimen (e.g. a daily or a weekly dosing regimen) is resumed. A variety of other drug holiday regimens are envisioned to be useful for administering the hepcidin analogues of the invention.

[0551] Thus, the hepcidin analogues may be delivered via an administration regime which comprises two or more administration phases separated by respective drug holiday phases.

[0552] During each administration phase, the hepcidin analogue is administered to the recipient subject in a therapeutically effective amount according to a pre-determined administration pattern. The administration pattern may comprise continuous administration of the drug to the recipient subject over the duration of the administration phase. Alternatively, the administration pattern may comprise adminis-

tration of a plurality of doses of the hepcidin analogue to the recipient subject, wherein said doses are spaced by dosing intervals.

[0553] A dosing pattern may comprise at least two doses per administration phase, at least five doses per administration phase, at least 10 doses per administration phase, at least 20 doses per administration phase, at least 30 doses per administration phase, or more.

[0554] Said dosing intervals may be regular dosing intervals, which may be as set out above, including once daily, twice daily, once every two, three, four, five or six days, once or twice weekly, once or twice monthly, or a regular and even less frequent dosing interval, depending on the particular dosage formulation, bioavailability, and pharmacokinetic profile of the hepcidin analogue of the present invention.

[0555] An administration phase may have a duration of at least two days, at least a week, at least 2 weeks, at least 4 weeks, at least a month, at least 2 months, at least 3 months, at least 6 months, or more.

[0556] Where an administration pattern comprises a plurality of doses, the duration of the following drug holiday phase is longer than the dosing interval used in that administration pattern. Where the dosing interval is irregular, the duration of the drug holiday phase may be greater than the mean interval between doses over the course of the administration phase. Alternatively the duration of the drug holiday may be longer than the longest interval between consecutive doses during the administration phase.

[0557] The duration of the drug holiday phase may be at least twice that of the relevant dosing interval (or mean thereof), at least 3 times, at least 4 times, at least 5 times, at least 10 times, or at least 20 times that of the relevant dosing interval or mean thereof.

[0558] Within these constraints, a drug holiday phase may have a duration of at least two days, at least a week, at least 2 weeks, at least 4 weeks, at least a month, at least 2 months, at least 3 months, at least 6 months, or more, depending on the administration pattern during the previous administration phase.

[0559] An administration regime comprises at least 2 administration phases. Consecutive administration phases are separated by respective drug holiday phases. Thus the administration regime may comprise at least 3, at least 4, at least 5, at least 10, at least 15, at least 20, at least 25, or at least 30 administration phases, or more, each separated by respective drug holiday phases.

[0560] Consecutive administration phases may utilise the same administration pattern, although this may not always be desirable or necessary. However, if other drugs or active agents are administered in combination with a hepcidin analogue of the invention, then typically the same combination of drugs or active agents is given in consecutive administration phases. In certain embodiments, the recipient subject is human.

[0561] In some embodiments, the present invention provides compositions and medicaments comprising at least one hepcidin analogue as disclosed herein. In some embodiments, the present invention provides a method of manufacturing medicaments comprising at least one hepcidin analogue as disclosed herein for the treatment of diseases of iron metabolism, such as iron overload diseases. In some embodiments, the present invention provides a method of manufacturing medicaments comprising at least one hepci-

din analogue as disclosed herein for the treatment of diabetes (Type I or Type II), insulin resistance, or glucose intolerance. Also provided are methods of treating a disease of iron metabolism in a subject, such as a mammalian subject, and preferably a human subject, comprising administering at least one hepcidin analogue, or composition as disclosed herein to the subject. In some embodiments, the hepcidin analogue or the composition is administered in a therapeutically effective amount. Also provided are methods of treating diabetes (Type I or Type II), insulin resistance, or glucose intolerance in a subject, such as a mammalian subject, and preferably a human subject, comprising administering at least one hepcidin analogue or composition as disclosed herein to the subject. In some embodiments, the hepcidin analogue or composition is administered in a therapeutically effective amount.

[0562] In some embodiments, the invention provides a process for manufacturing a hepcidin analogue or a hepcidin analogue composition (e.g., a pharmaceutical composition), as disclosed herein.

[0563] In some embodiments, the invention provides a device comprising at least one hepcidin analogue of the present invention, or pharmaceutically acceptable salt or solvate thereof for delivery of the hepcidin analogue to a subject.

[0564] In some embodiments, the present invention provides methods of binding a ferroportin or inducing ferroportin internalization and degradation which comprises contacting the ferroportin with at least one hepcidin analogue, or hepcidin analogue composition as disclosed herein.

[0565] In some embodiments, the present invention provides methods of binding a ferroportin to block the pore and exporter function without causing ferroportin internalization. Such methods comprise contacting the ferroportin with at least one hepcidin analogue, or hepcidin analogue composition as disclosed herein.

[0566] In some embodiments, the present invention provides kits comprising at least one hepcidin analogue, or hepcidin analogue composition (e.g., pharmaceutical composition) as disclosed herein packaged together with a reagent, a device, instructional material, or a combination thereof.

[0567] In some embodiments, the present invention provides a method of administering a hepcidin analogue or hepcidin analogue composition (e.g., pharmaceutical composition) of the present invention to a subject via implant or osmotic pump, by cartridge or micro pump, or by other means appreciated by the skilled artisan, as well-known in the art. In some embodiments, the present invention provides complexes which comprise at least one hepcidin analogue as disclosed herein bound to a ferroportin, preferably a human ferroportin, or an antibody, such as an antibody which specifically binds a hepcidin analogue as disclosed herein, Hep25, or a combination thereof.

[0568] In some embodiments, the hepcidin analogue of the present invention has a measurement (e.g., an EC_{50}) of less than 500 nM within the FPN internalization assay. As a skilled person will realize, the function of the hepcidin analogue is dependent on the tertiary structure of the hepcidin analogue and the binding surface presented. It is therefore possible to make minor changes to the sequence encoding the hepcidin analogue that do not affect the fold or are not on the binding surface and maintain function. In other embodiments, the present invention provides a hepci-

din analogue having 85% or higher (e.g., 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.5%) identity or homology to an amino acid sequence of any hepcidin analogue described herein that exhibits an activity (e.g., hepcidin activity), or lessens a symptom of a disease or indication for which hepcidin is involved.

[0569] In other embodiments, the present invention provides a hepcidin analogue having 85% or higher (e.g., 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.5%) identity or homology to an amino acid sequence of any hepcidin analogue presented herein, or a peptide according to any one of the formulae or hepcidin analogues described herein.

[0570] In some embodiments, a hepcidin analogue of the present invention may comprise functional fragments or variants thereof that have at most 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid substitutions compared to one or more of the specific peptide analogue sequences recited herein.

[0571] In addition to the methods described in the Examples herein, the hepcidin analogues of the present invention may be produced using methods known in the art including chemical synthesis, biosynthesis or in vitro synthesis using recombinant DNA methods, and solid phase synthesis. See e.g. Kelly & Winkler (1990) *Genetic Engineering Principles and Methods*, vol. 12, J. K. Setlow ed., Plenum Press, NY, pp. 1-19; Merrifield (1964) *J Amer Chem Soc* 85:2149; Houghten (1985) *PNAS USA* 82:5131-5135; and Stewart & Young (1984) *Solid Phase Peptide Synthesis*, 2ed. Pierce, Rockford, IL, which are herein incorporated by reference. The hepcidin analogues of the present invention may be purified using protein purification techniques known in the art such as reverse phase high-performance liquid chromatography (HPLC), ion-exchange or immunoaffinity chromatography, filtration or size exclusion, or electrophoresis. See Olsnes, S. and A. Pihl (1973) *Biochem.* 12(16): 3121-3126; and Scopes (1982) *Protein Purification*, Springer-Verlag, NY, which are herein incorporated by reference. Alternatively, the hepcidin analogues of the present invention may be made by recombinant DNA techniques known in the art. Thus, polynucleotides that encode the polypeptides of the present invention are contemplated herein. In certain preferred embodiments, the polynucleotides are isolated. As used herein "isolated polynucleotides" refers to polynucleotides that are in an environment different from that in which the polynucleotide naturally occurs.

EXAMPLES

[0572] The following examples demonstrate certain specific embodiments of the present invention. The following examples were carried out using standard techniques that are well known and routine to those of skill in the art, except where otherwise described in detail. It is to be understood that these examples are for illustrative purposes only and do not purport to be wholly definitive as to conditions or scope of the invention. As such, they should not be construed in any way as limiting the scope of the present invention.

Abbreviations

- [0573]** DCM: dichloromethane
- [0574]** DMF: N,N-dimethylformamide
- [0575]** NMP: N-methylpyrrolidone

- [0576] HBTU: O-(Benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate
- [0577] HATU: 2-(7-aza-1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium
- [0578] hexafluorophosphate
- [0579] DCC: Dicyclohexylcarbodiimide
- [0580] NHS: N-hydroxysuccinimide
- [0581] DIPEA: diisopropylethylamine
- [0582] EtOH: ethanol
- [0583] Et₂O: diethyl ether
- [0584] H₂: hydrogen
- [0585] TFA: trifluoroacetic acid
- [0586] TIS: triisopropylsilane
- [0587] ACN: acetonitrile
- [0588] HPLC: high performance liquid chromatography
- [0589] ESI-MS: electron spray ionization mass spectrometry
- [0590] PBS: phosphate-buffered saline
- [0591] Boc: t-butoxycarbonyl
- [0592] Fmoc: Fluorenylmethyloxycarbonyl
- [0593] Acn: acetamidomethyl
- [0594] IVA: Isovaleric acid (or Isovaleryl)
- [0595] K(): In the peptide sequences provided herein, wherein a compound or chemical group is presented in parentheses directly after a Lysine residue, it is to be understood that the compound or chemical group in the parentheses is a side chain conjugated to the Lysine residue. So, e.g., but not to be limited in any way, K—[(PEG8)] indicates that a PEG8 moiety is conjugated to a side chain of this Lysine.
- [0596] Palm: Indicates conjugation of a palmitic acid (palmitoyl).

Synthetic Protocol—1

Synthesis of Peptide Monomers

[0597] Peptide monomers of the present invention were synthesized using the Merrifield solid phase synthesis techniques on Protein Technology's Symphony multiple channel synthesizer. The peptides were assembled using HBTU (O-Benzotriazole-N,N,N',N'-tetramethyl-uronium-hexafluoro-phosphate), Diisopropylethylamine (DIEA) coupling conditions. For some amino acid couplings PyAOP (7-Azabenzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate) and DIEA conditions were used. Rink Amide MBHA resin (100-200 mesh, 0.57 mmol/g) was used for peptide with C-terminal amides and pre-loaded Wang Resin with N- α -Fmoc protected amino acid was used for peptide with C-terminal acids. The coupling reagents (HBTU and DIEA premixed) were prepared at 100 mmol concentration. Similarly, amino acids solutions were prepared at 100 mmol concentration. Peptide inhibitors of the present invention were identified based on medical chemistry optimization and/or phage display and screened to identify those having superior binding and/or inhibitory properties.

Assembly

[0598] The peptides were assembled using standard Symphony protocols. The peptide sequences were assembled as follows: Resin (250 mg, 0.14 mmol) in each reaction vial was washed twice with 4 ml of DMF followed by treatment

with 2.5 ml of 20% 4-methyl piperidine (Fmoc de-protection) for 10 min. The resin was then filtered and washed two times with DMF (4 ml) and re-treated with Piperidine for additional 30 minute. The resin was again washed three times with DMF (4 ml) followed by addition 2.5 ml of amino acid and 2.5 ml of HBTU-DIEA mixture. After 45 min of frequent agitations, the resin was filtered and washed three times with DMF (4 ml each). For a typical peptide of the present invention, double couplings were performed. After completing the coupling reaction, the resin was washed three times with DMF (4 ml each) before proceeding to the next amino acid coupling.

Cleavage

[0599] Following completion of the peptide assembly, the peptide was cleaved from the resin by treatment with cleavage reagent, such as reagent K (82.5% trifluoroacetic acid, 5% water, 5% thioanisole, 5% phenol, 2.5% 1,2-ethanedithiol). The cleavage reagent was able to successfully cleave the peptide from the resin, as well as all remaining side chain protecting groups.

[0600] The cleaved peptides were precipitated in cold diethyl ether followed by two washings with ethyl ether. The filtrate was poured off and a second aliquot of cold ether was added, and the procedure repeated. The crude peptide was dissolved in a solution of acetonitrile water (7:3 with 1% TFA) and filtered. The quality of linear peptide was then verified using electrospray ionization mass spectrometry (ESI-MS) (Micromass/Waters ZQ) before being purified.

Purification

[0601] Analytical reverse-phase, high performance liquid chromatography (HPLC) was performed on a Gemini C18 column (4.6 mm×250 mm) (Phenomenex). Semi-Preparative reverse phase HPLC was performed on a Gemini 10 μ m C18 column (22 mm×250 mm) (Phenomenex). Separations were achieved using linear gradients of buffer B in A (Mobile phase A: water containing 0.15% TFA, mobile phase B: Acetonitrile (ACN) containing 0.1% TFA), at a flow rate of 1 mL/min (analytical) and 20 mL/min (preparative). Separations were achieved using linear gradients of buffer B in A (Mobile phase A: water containing 0.15% TFA, mobile phase B: Acetonitrile (ACN) containing 0.1% TFA), at a flow rate of 1 mL/min (analytical) and 15 mL/min (preparative).

Synthetic Protocol—2

Synthesis of Peptide Monomers

[0602] Peptide monomers of the present invention were synthesized using standard Fmoc solid phase synthesis techniques on a CEM Liberty Bluem microwave peptide synthesizer. The peptides were assembled using Oxyma/DIC (ethyl cyanohydroxyiminoacetate/diisopropylcarbodiimide) with microwave heating. Rink Amide-MBHA resin (100-200 mesh, 0.66 mmol/g) was used for peptides with C-terminal amides and pre-loaded Wang Resin with N- α -Fmoc protected amino acid was used for peptide with C-terminal acids. Oxyma was prepared as a 1M solution in DMF with 0.1M DIEA. DIC was prepared as 0.5M solution in DMF. The Amino acids were prepared at 200 mM. Peptide inhibitors of the present invention were identified based on

medicinal chemistry optimization and/or phage display and screened to identify those having superior binding and/or inhibitory properties.

Assembly

[0603] The peptides were made using standard CEM Liberty Blue protocols. The peptide sequences were assembled as follows: Resin (400 mg, 0.25 mmol) was suspended in 10 ml of 50/50 DMF/DCM. The resin was then transferred to the reaction vessel in the microwave cavity. The peptide was assembled using repeated Fmoc deprotection and Oxyma/DIC coupling cycles. For deprotection, 20% 4-methylpiperidine in DMF was added to the reaction vessel and heated to 90° C. for 65 seconds. The deprotection solution was drained and the resin washed three times with DMF. For most amino acids, 5 equivalents of amino acid, Oxyma and DIC were then added to the reaction vessel and microwave irradiation rapidly heated the mixing reaction to 90° C. for 4 min. For Arginine and Histidine residues, milder conditions using respective temperatures of 75 and 50° C. for 10 min were used to prevent racemization. Rare and expensive amino acids were often coupled manually overnight at room temperature using only 1.5-2 eq of reagents. Difficult couplings were often double coupled 2x4 min at 90° C. After coupling the resin was washed with DMF and the whole cycle was repeated until the desired peptide assembly was completed.

Cleavage

[0604] Following completion of the peptide assembly, the peptide was then cleaved from the resin by treatment with a standard cleavage cocktail of 91:5:2:2 TFA/H₂O/TIPS/DODT for 2 hrs. If more than one Arg(Pbf) residue was present the cleavage was allowed to go for an additional hour.

[0605] The cleaved peptides were precipitated in cold diethyl ether. The filtrate was decanted off and a second aliquot of cold ether was added, and the procedure was repeated. The quality of linear peptide was then verified using electrospray ionization mass spectrometry (ESI-MS) (Waters® Micromass® ZQ™) before being purified.

Purification

[0606] Analytical reverse-phase, high performance liquid chromatography (HPLC) was performed on a Gemini® C18 column (4.6 mm×250 mm) (Phenomenex). Semi-Preparative reverse phase HPLC was performed on a Gemini® 10 µm C18 column (22 mm×250 mm) (Phenomenex) or Jupiter® 10 µm, 300 Å C18 column (21.2 mm×250 mm) (Phenomenex). Separations were achieved using linear gradients of buffer B in A (Mobile phase A: water containing 0.15% TFA, mobile phase B: Acetonitrile (ACN) containing 0.1% TFA), at a flow rate of 1 mL/min (analytical) and 20 mL/min (preparative).

Example 1A

Synthesis of Peptide Analogues

[0607] Unless otherwise specified, reagents and solvents employed in the following were available commercially in standard laboratory reagent or analytical grade, and were used without further purification.

Procedure for Solid-Phase Synthesis of Peptides

Method A

[0608] Peptide analogues of the invention were chemically synthesized using optimized 9-fluorenylmethoxy carbonyl (Fmoc) solid phase peptide synthesis protocols. For C-terminal amides, rink-amide resin was used, although wang and trityl resins were also used to produce C-terminal acids. The side chain protecting groups were as follows: Glu, Thr and Tyr: O-tButyl; Trp and Lys: t-Boc (t-butyloxycarbonyl); Arg: N-gamma-2,2,4,6,7-pentamethyldihydrobenzofuran-5-sulfonyl; His, Gln, Asn, Cys: Trityl. For selective disulfide bridge formation, AcM (acetamidomethyl) was also used as a Cys protecting group. For coupling, a four to ten-fold excess of a solution containing Fmoc amino acid, HBTU and DIEA (1:1:1.1) in DMF was added to swelled resin [HBTU: 0-(Benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate; DIEA: diisopropylethylamine; DMF: dimethylformamide]. HATU (0-(7-azabenzotriazol-1-yl)-1,1,3,3,-tetramethyluronium hexafluorophosphate) was used instead of HBTU to improve coupling efficiency in difficult regions. Fmoc protecting group removal was achieved by treatment with a DMF, piperidine (2:1) solution.

Method B

[0609] Alternatively, peptides were synthesized utilizing the CEM liberty Blue Microwave assisted peptide synthesizer. Using the Liberty Blue, Fmoc deprotection was carried out by addition of 20% 4-methylpiperidine in DMF with 0.1M Oxyma in DMF and then heating to 90° C. using microwave irradiation for 4 min. After DMF washes the Fmoc-amino acids were coupled by addition of 0.2M amino acid (4-6 eq), 0.5M DIC (4-6 eq) and 1M Oxyma (with 0.1M DIEA) 4-6 eq (all in DMF). The coupling solution is heated using microwave radiation to 90° C. for 4 min. A second coupling is employed when coupling Arg or other sterically hindered amino acids. When coupling with histidine, the reaction is heated to 50° C. for 10 min. The cycles are repeated until the full length peptide is obtained.

Procedure for Cleavage of Peptides Off Resin

[0610] Side chain deprotection and cleavage of the peptide analogues of the invention (e.g., Compound No. 2) was achieved by stirring dry resin in a solution containing trifluoroacetic acid, water, ethanedithiol and tri-isopropylsilane (90:5:2.5:2.5) for 2 to 4 hours. Following TFA removal, peptide was precipitated using ice-cold diethyl ether. The solution was centrifuged and the ether was decanted, followed by a second diethyl ether wash. The peptide was dissolved in an acetonitrile, water solution (1:1) containing 0.1% TFA (trifluoroacetic acid) and the resulting solution was filtered. The linear peptide quality was assessed using electrospray ionization mass spectrometry (ESI-MS).

Procedure for Purification of Peptides

[0611] Purification of the peptides of the invention (e.g., Compound No. 2) was achieved using reverse-phase high performance liquid chromatography (RP-HPLC). Analysis was performed using a C18 column (3 µm, 50×2 mm) with a flow rate of 1 mL/min. Purification of the linear peptides was achieved using preparative RP-HPLC with a C18 column (5 µm, 250×21.2 mm) with a flow rate of 20 mL/min.

Separation was achieved using linear gradients of buffer B in A (Buffer A: Aqueous 0.05% TFA; Buffer B: 0.043% TFA, 90% acetonitrile in water).

[0612] One of skill in the art will appreciate that standard methods of peptide synthesis may be used to generate the compounds of the invention.

Conjugation of Half-Life Extension Moieties

[0613] Conjugation of peptides were performed on resin. Lys(ivDde) was used as the key amino acid. After assembly of the peptide on resin, selective deprotection of the ivDde group occurred using 3×5 min 2% hydrazine in DMF for 5 min. Activation and acylation of the linker using HBTU, DIEA 1-2 equivalents for 3 h, and Fmoc removal followed by a second acylation with the lipidic acid gave the conjugated peptide.

Example 1B

Synthesis of Peptide Id #31

Isovaleric Acid-E-T-H-[Dpa]-P-[Cys]-[Cys]-[Lys
(1PEG2_1PEG2_IsoGlu_C18_Diacid)]-[bhPhe]-
[(D)Lys]-NH₂

[0614] The TFA salt of Peptide #31 was synthesized on Rink amide resin. Upon completion, 50 mg of >95% pure Peptide #31 was isolated as a white powder. Peptide #31 was constructed on Rink Amide MBHA (100-200 mesh, 0.66 mmol/g) resin using standard Fmoc protection synthesis conditions. The constructed peptide was isolated from the resin and protecting groups by cleavage with strong acid followed by precipitation. The crude precipitate was then purified by RP-HPLC. Lyophilization of pure fractions gave the final product Peptide #31.

Peptide Assembly

[0615] Swell Resin: 200 mg of Rink Amide MBHA solid phase resin (0.66 mmol/g loading) was transferred to a 250 mL reaction vessel. The resin was swelled with 60 mL of DMF (2 hrs).

[0616] Step 1: Coupling of Fmoc-(D)Lys(Boc)-OH: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL of amino acid Fmoc-(D)Lys(Boc)-OH in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0617] Step 2: Coupling of Fmoc-βhomo-L-Phe-OH: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL of amino Fmoc-βhomo-L-Phe-OH in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After complet-

ing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0618] Step 3 Coupling of Fmoc-L-Lys(IvDde)-OH: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL of amino acid Fmoc-L-Lys(IvDde)-OH in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0619] Step 4 Coupling of Fmoc-L-Cys(Trt)-OH: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL of amino acid Fmoc-L-Cys(Trt)-OH in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0620] Step 5 Coupling of Fmoc-L-Cys(Trt)-OH: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL of amino acid Fmoc-L-Cys(Trt)-OH in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0621] Step 6: Coupling of Fmoc-Pro-OH: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL of amino acid Fmoc-Pro-OH in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0622] Step 7: Coupling of Fmoc-L-DIP-OH: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL of amino acid Fmoc-L-DIP-OH in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the

coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0623] Step 8: Coupling of Fmoc-L-His(Trt)-OH: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL of amino acid Fmoc-L-His(Trt)-OH in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0624] Step 9: Coupling of Fmoc-L-Thr(tBu)-OH: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL of amino acid Fmoc-L-Thr(tBu)-OH in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0625] Step 9': Coupling of Fmoc-L-Glu(tBu)-OH: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL of amino acid (2S)-2-[[[9H-Fmoc-L-Glu(tBu)-OH: in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0626] Step 10: Coupling of Isovaleric acid: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL Isovaleric acid in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0627] Step 11: IvDde removal and Coupling of Fmoc-1PEG2-OH: The IvDde was removed from the Lys C-terminus of the resin bound peptide using 2-5% hydrazine in DMF (4×30 min), followed by a DMF wash. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL of amino acid Fmoc-1PEG2-OH in DMF (200 mM) and 2.0 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the cou-

pling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0628] Step 12: Coupling of Fmoc-1PEG2-OH: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL Fmoc-1PEG2-OH in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0629] Step 13: Coupling of Fmoc-L-Glu-OtBu: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL Fmoc-Glu-OtBu in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0630] Step 14: Coupling of tBuO₂C(CH₂)₁₆CO₂H: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL tBuO₂C(CH₂)₁₆CO₂H in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0631] Step 15: TFA Cleavage and Ether precipitation: 10 ml of the cleavage cocktail [TFA cleavage cocktail (90/5/2.5/2.5 TFA/water/Tips/DODT) was added to the protected resin bound peptide and shaken for two hours. Cold Diethyl Ether was added forming a white precipitate that was then centrifuged. The ether was decanted to waste and 2 more ether washes of the precipitate were performed. The resulting white precipitate cake was dissolved in acetonitrile/water (7:3) and filtered before oxidation and purification.

[0632] Step 16: The peptide in acetonitrile/water (7:3) was diluted to 250 mL. 12 in MeOH was added dropwise until a sustained brown color resulted. The mixture was stirred for 2-5 mins before ascorbic acid was added portion wise until the solution became clear. The solution was filtered and ready for purification by RP-HPLC.

[0633] Step 17: RP-HPLC purification: Semi-Preparative reverse phase HPLC was performed on a Gemini® 10 µm C18 column (22 mm×250 mm) (Phenomenex). Separations were achieved using linear gradients of buffer B in A (Mobile phase A: water containing 0.15% TFA, mobile phase B: Acetonitrile (ACN) containing 0.1% TFA), at a flow rate of 20 mL/min (preparative).

[06334] Step 18: Final Lyophilization and Analysis: The collected fractions were analyzed by analytical RP-HPLC, and all fractions >95% purity were combined. Lyophilization of the combined fractions gave Peptide #31 as a white powder with a purity of >95%. Low resolution LC/MS of purified #31 gave one major charged state of the peptide, $[M+2H]^{2+}$ of 1064.3. The experimental mass agrees with the theoretical molecular weight of 2126.6 Da.

Example 1C

Synthesis of Peptide #42

Isovaleric Acid-[Glu_OMe]-T-H-[Dpa]-P-[Cys]-
[Cys]-[Lys(1PEG2_1PEG2_Dap_C18_Diacid)]-
[bhPhe]-[(D)Lys]-NH₂

[06335] The TFA salt of Peptide #42 was synthesized on Rink amide resin. Upon completion, 52 mg of >95% pure Peptide #42 was isolated as a white powder.

[06336] Peptide #42 was constructed on Rink Amide MBHA (100-200 mesh, 0.66 mmol/g) resin using standard Fmoc protection synthesis conditions. The constructed peptide was isolated from the resin and protecting groups by cleavage with strong acid followed by precipitation. The crude precipitate was then purified by RP-HPLC. Lyophilization of pure fractions gave the final product Peptide #42.

Peptide Assembly

[06337] Swell Resin: 200 mg of Rink Amide MBHA solid phase resin (0.66 mmol/g loading) was transferred to a 250 mL reaction vessel. The resin was swelled with 60 mL of DMF (2 hrs).

[06338] Step 1: Coupling of Fmoc-(D)Lys(Boc)-OH: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL of amino acid Fmoc-(D)Lys(Boc)-OH in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[06339] Step 2: Coupling of Fmoc-β-homo-L-Phe-OH: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL of amino Fmoc-β-homo-L-Phe-OH in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0640] Step 3: Coupling of Fmoc-L-Lys(IvDde)-OH: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of

DMF (3×0.1 min) and followed by addition of 2.5 mL of amino acid Fmoc-L-Lys(IvDde)-OH in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0641] Step 4: Coupling of Fmoc-L-Cys(Trt)-OH: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL of amino acid Fmoc-L-Cys(Trt)-OH in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0642] Step 5: Coupling of Fmoc-L-Cys(Trt)-OH: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL of amino acid Fmoc-L-Cys(Trt)-OH in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0643] Step 6: Coupling of Fmoc-Pro-OH: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL of amino acid Fmoc-Pro-OH in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0644] Step 7: Coupling of Fmoc-L-DIP-OH: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL of amino acid Fmoc-L-DIP-OH in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0645] Step 8: Coupling of Fmoc-L-His(Trt)-OH: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of

DMF (3×0.1 min) and followed by addition of 2.5 mL of amino acid FMOC-L-His(Trt)-OH in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0646] Step 9: Coupling of FMOC-L-Thr(tBu)-OH: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL of amino acid FMOC-L-Thr(tBu)-OH in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0647] Step 9': Coupling of FMOC-L-Glu(OMe)-OH: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL of amino acid (2S)-2-[(9H-FMOC-L-Glu(OMe)-OH): in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0648] Step 10: Coupling of Isovaleric acid: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL Isovaleric acid in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0649] Step 11: IvDde removal and Coupling of Fmoc-1PEG2-OH: The IvDde was removed from the Lys C-terminus of the resin bound peptide using 2-5% hydrazine in DMF (4×30 min), followed by a DMF wash. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL of amino acid Fmoc-1PEG2-OH in DMF (200 mM) and 2.0 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0650] Step 12: Coupling of Fmoc-1PEG2-OH: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF

(3×0.1 min) and followed by addition of 2.5 mL Fmoc-1PEG2-OH in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0651] Step 13: Coupling of Fmoc-L-Dap(Boc)-OH: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL Fmoc-Dap(Boc)-OH in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0652] Step 14: Coupling of tBuO₂C(CH₂)₁₆CO₂H: Deprotection of the Fmoc group was accomplished by two treatments with 2.5 ml of 20% piperidine in DMF twice to the swollen Rink Amide resin for 5 and 10 min respectively. After deprotection the resin was washed with 3.75 mL of DMF (3×0.1 min) and followed by addition of 2.5 mL tBuO₂C(CH₂)₁₆CO₂H in DMF (200 mM) and 2.5 mL of coupling reagent HBTU-DIEA mixture in DMF (200 and 220 mM). The coupling reaction was mixed for 1 hr, filtered and repeated once (double coupling). After completing the coupling reaction, the resin was washed with 6.25 mL of DMF (3×0.1 min) prior to starting the next deprotection/coupling cycle.

[0653] Step 15: TFA Cleavage and Ether precipitation: 10 ml of the cleavage cocktail [TFA cleavage cocktail (90/5/2.5/2.5 TFA/water/Tips/DODT)] was added to the protected resin bound peptide and shaken for two hours. Cold Diethyl Ether was added forming a white precipitate that was then centrifuged. The ether was decanted to waste and 2 more ether washes of the precipitate were performed. The resulting white precipitate cake was dissolved in acetonitrile/water (7:3) and filtered before oxidation and purification.

[0654] Step 16: The peptide in acetonitrile/water (7:3) was diluted to 250 mL. 12 in MeOH was added dropwise until a sustained brown color resulted. The mixture was stirred for 2-5 mins before ascorbic acid was added portion wise until the solution became clear. The solution was filtered and ready for purification by RP-HPLC.

[0655] Step 17: RP-HPLC purification: Semi-Preparative reverse phase HPLC was performed on a Gemini® 10 µm C18 column (22 mm×250 mm) (Phenomenex). Separations were achieved using linear gradients of buffer B in A (Mobile phase A: water containing 0.15% TFA, mobile phase B: Acetonitrile (ACN) containing 0.1% TFA), at a flow rate of 20 mL/min (preparative).

[0656] Step 18: Final Lyophilization and Analysis: The collected fractions were analyzed by analytical RP-HPLC, and all fractions >95% purity were combined. Lyophilization of the combined fractions gave Peptide #42 as a white powder with a purity of >95%. Low resolution LC/MS of purified #42 gave 2 charged states of the peptide, [M+3H]³⁺

of 700.0 and the molecular ion [M+2H]²⁺ of 1049.4. The experimental mass agrees with the theoretical molecular weight of 2097.6 Da.

Example 2

Activity of Peptide Analogues

[0657] Peptide analogues were tested in vitro for induction of internalization of the human ferroportin protein. Following internalization, the ferroportin protein is degraded. The assay used (FPN activity assay) measures a decrease in fluorescence of the receptor.

[0658] The cDNA encoding the human ferroportin (SLC40A1) was cloned from a cDNA clone from Origene (NM_014585). The DNA encoding the ferroportin was amplified by PCR using primers also encoding terminal restriction sites for subcloning, but without the termination codon. The ferroportin receptor was subcloned into a mammalian GFP expression vector containing a neomycin (G418) resistance marker in such that the reading frame of the ferroportin was fused in frame with the GFP protein. The fidelity of the DNA encoding the protein was confirmed by DNA sequencing. HEK293 cells were used for transfection of the ferroportin-GFP receptor expression plasmid. The cells were grown according to standard protocol in growth medium and transfected with the plasmids using Lipofectamine (manufacturer's protocol, Invitrogen). The cells stably expressing ferroportin-GFP were selected using G418 in the growth medium (in that only cells that have taken up and incorporated the cDNA expression plasmid survive) and sorted several times on a Cytomation MoFlo™ cell sorter to obtain the GFP-positive cells (488 nm/530 nm). The cells were propagated and frozen in aliquots.

[0659] To determine activity of the hepcidin analogues (compounds) on the human ferroportin, the cells were incubated in 96 well plates in standard media, without phenol red. Compound was added to desired final concentration for at least 18 hours in the incubator. Following incubation, the remaining GFP-fluorescence was determined either by whole cell GFP fluorescence (Envision plate reader, 485/535 filter pair), or by Beckman Coulter Quanta™ flow cytometer (express as Geometric mean of fluorescence intensity at 485 nm/525 nm). Compound was added to desired final concentration for at least 18 hours but no more than 24 hours in the incubator.

[0660] In certain experiments, reference compounds included native Hepcidin, Mini-Hepcidin, and R1-Mini-Hepcidin, which is an analog of mini-hepcidin. The "RI" in

RI-Mini-Hepcidin refers to Retro Inverse. A retro inverse peptide is a peptide with a reversed sequence in all D amino acids. An example is that Hy-Glu-Thr-His-NH₂ becomes Hy-DHis-DThr-DGlu-NH₂. The EC₅₀ of these reference compounds for ferroportin internalization/degradation was determined according to the FPN activity assay described above. These peptides served as control standards.

TABLE 5

Reference compounds		
Name	Sequence	Potency EC50 (nM)
Hepcidin	Hy-DTHFPIC(1) IFC(2) C(3) GC(2) C(4) HRSKC(3) GMC(4) C(1) KT-OH (SEQ ID NO: 501)	34
Mini-Hepcidin 1-9	Hy-DTHFPICIF-NH ₂ (SEQ ID NO: 502)	712
RI-Mini-Hepcidin	Hy-DPhe-DIle-DCys-DIle-DPro-DPhe-DHis-DThr-DAsp-NH ₂ (SEQ ID NO: 503)	>10 μM
Ref. Compd 1	Isovaleric acid-DTHFPC IKF-Lys[PEG11-Palm]-PRSKGCK-NH ₂ (SEQ ID NO: 601)	30
Ref. Compd. 2	Isovaleric acid-DTHFPCKF-Lys[PEG11-Palm]-PRSK-[SAR]-CK-NH ₂ (SEQ ID NO: 602)	13

The potency IC₅₀ or EC₅₀ values (nM) determined for various peptide analogues of the present invention are provided in Tables 6A-6E. These values were determined as described herein. Compound ID numbers are indicated by "Compd ID," and reference compounds are indicated by "Ref Compd." FPN IC₅₀ or EC₅₀ values determined from these data are shown in Tables 6A-6E. Where not shown, data was not yet determined.

TABLE 6A

Illustrative Hepcidin Analogues				
Compd ID	SEQ ID No.	Peptide	Cyclized@	FPN IC50 (nM) T47D MSA IC50 (nM)
1	1	Isovaleric Acid-[Cys]-T-H-[Dpa]-P-[Cys]-I-[(D)Lys]-[bhPhe]-[Lys(Ahx_Palm)]-[(D)Lys]-NH2;	Cys-Cys form a disulfide bond	**
2	2	Isovaleric Acid-E-T-[Cys]-[Dpa]-P-[Cys]-I-[(D)Lys]-[bhPhe]-[Lys(Ahx_Palm)]-[(D)Lys]-NH2;	Cys-Cys form a disulfide bond	**

TABLE 6A-continued

Illustrative Hepcidin Analogues					
Compd ID	SEQ ID No.	Peptide	Cyclized [@]	FPN IC ₅₀ (nM)	T47D MSA IC ₅₀ (nM)
3	3	Isovaleric Acid-E-T-H-[Cys]-P-[Cys]-I-[(D)Lys]-[bhPhe]-[Lys(Ahx_Palm)]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	***	
4	4	Isovaleric Acid-E-T-H-[Dpa]-[Cys]-[Cys]-I-[(D)Lys]-[bhPhe]-[Lys(Ahx_Palm)]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	***	
5	5	Isovaleric Acid-E-T-H-[Dpa]-[Cys]-[Cys]-I-[(D)Lys]-[bhPhe]-[Lys(Ahx_Palm)]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	***	
6	6	Isovaleric Acid-E-T-H-[Dpa]-P-[Cys]-[Cys]-[(D)Lys]-[bhPhe]-[Lys(Ahx_Palm)]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	***	
7	7	Isovaleric Acid-E-T-H-[Dpa]-P-[Cys]-I-[Cys]-[bhPhe]-[Lys(Ahx_Palm)]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	***	
8	8	Isovaleric Acid-E-T-H-[Dpa]-P-[Cys]-I-[Cys]-[bhPhe]-[Lys(Ahx_Palm)]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	**	
9	9	Ac-[Cys]-E-T-H-[Dpa]-P-[Cys]-I-[Lys(Ahx_Palm)]-[bhPhe]-NH ₂ ;	Cys-Cys form a disulfide bond	**	
10	10	Isovaleric Acid-[Cys]-E-T-H-[Dpa]-P-[Cys]-I-[Lys(Ahx_Palm)]-[bhPhe]-NH ₂ ;	Cys-Cys form a disulfide bond	**	
11	11	Pen-E-T-H-[Dpa]-P-[Cys]-I-[Lys(Ahx_Palm)]-[bhPhe]-NH ₂ ;	Pen-Cys form a disulfide bond	*	
12	12	Ac-[hCys]-E-T-H-[Dpa]-P-[Cys]-I-[Lys(Ahx_Palm)]-[bhPhe]-NH ₂ ;	hCys-Cys form a disulfide bond	**	
13	13	Isovaleric Acid-[hCys]-E-T-H-[Dpa]-P-[Cys]-I-[Lys(Ahx_Palm)]-[bhPhe]-NH ₂ ;	hCys-Cys form a disulfide bond	**	
14	14	Ac-[hCys]-E-T-H-[Dpa]-P-[hCys]-I-[Lys(Ahx_Palm)]-[bhPhe]-NH ₂ ;	hCys-hCys form a disulfide bond	**	
15	15	Isovaleric Acid-[hCys]-E-T-H-[Dpa]-P-[hCys]-I-[Lys(Ahx_Palm)]-[bhPhe]-NH ₂ ;	hCys-hCys form a disulfide bond	*	
16	16	hCys-V-E-T-H-[Dpa]-P-[hCys]-I-[Lys(Ahx_Palm)]-[bhPhe]-NH ₂ ;	hCys-hCys form a disulfide bond	*	
17	17	[Cys]-V-E-T-H-[Dpa]-P-[hCys]-I-[Lys(Ahx_Palm)]-[bhPhe]-NH ₂ ;	Cys-hCys form a disulfide bond	**	
18	18	Isovaleric Acid-E-T-H-F-P-[Pen]-I-[Lys(1PEG2_1PEG2_IsoGlu_C18_Diacid)]-F-E-P-R-S-K-G-[Pen]-K-NH ₂ ;	Pen-Pen form a disulfide bond	*	*
19	19	Isovaleric Acid-E-T-H-F-P-[Pen]-I-[Lys(1PEG2_1PEG2_IsoGlu_C18_Diacid)]-F-E-P-R-S-K-G-[Cys]-K-NH ₂ ;	Cys-Cys form a disulfide bond	*	

TABLE 6A-continued

Illustrative Hepcidin Analogues					
Compd ID	SEQ ID No.	Peptide	Cyclized [@]	FPN IC ₅₀ (nM)	T47D MSA IC ₅₀ (nM)
20	20	Isovaleric Acid-E-T-H-[Cys]-P-[Cys]-I-[Lys(1PEG2_1PEG2_Ahx_C18_Diacid)]-[bhPhe]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	**	
21	21	Isovaleric Acid-E-T-H-[Dpa]-[Cys]-[Cys]-I-[Lys(1PEG2_1PEG2_Ahx_C18_Diacid)]-[bhPhe]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	***	***
22	22	Isovaleric Acid-E-T-H-[Dpa]-P-[Cys]-[Cys]-[Lys(1PEG2_1PEG2_Ahx_C18_Diacid)]-[bhPhe]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	***	***
23	23	Isovaleric Acid-E-T-H-[Dpa]-P-[Cys]-I-[Lys(1PEG2_1PEG2_Ahx_C18_Diacid)]-[bhPhe]-[Cys]-NH ₂ ;	Cys-Cys form a disulfide bond	***	***
24	24	Isovaleric Acid-E-T-H-[Cys]-P-[Cys]-I-[(D)Lys]-[bhPhe]-[Lys(1PEG2_1PEG2_Ahx_C18_Diacid)]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	**	
25	25	Isovaleric Acid-E-T-H-[Dpa]-[Cys]-[Cys]-I-[(D)Lys]-[bhPhe]-[Lys(1PEG2_1PEG2_Ahx_C18_Diacid)]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	**	
26	26	Isovaleric Acid-E-T-H-[Dpa]-P-[Cys]-[Cys]-[(D)Lys]-[bhPhe]-[Lys(1PEG2_1PEG2_Ahx_C18_Diacid)]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	***	
27	27	Isovaleric Acid-E-T-H-[Dpa]-P-[Cys]-I-[Cys]-[bhPhe]-[Lys(1PEG2_1PEG2_Ahx_C18_Diacid)]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	***	***
28	28	Isovaleric Acid-E-T-H-[Dpa]-P-[Cys]-[Cys]-[(D)Lys]-[bhPhe]-[Lys(1PEG2_1PEG2_IsoGlu_C18_Diacid)]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	**	
29	29	Isovaleric Acid-E-T-H-[Cys]-P-[Cys]-I-[Lys(1PEG2_1PEG2_IsoGlu_C18_Diacid)]-[bhPhe]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	**	
30	30	Isovaleric Acid-E-T-H-[Dpa]-[Cys]-[Cys]-I-[Lys(1PEG2_1PEG2_IsoGlu_C18_Diacid)]-[bhPhe]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	***	***
31	31	Isovaleric Acid-E-T-H-[Dpa]-P-[Cys]-[Cys]-[Lys(1PEG2_1PEG2_IsoGlu_C18_Diacid)]-[bhPhe]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	***	***
32	32	Isovaleric Acid-E-T-H-[Dpa]-P-[Cys]-I-[Lys(1PEG2_1PEG2_IsoGlu_C18_Diacid)]-[bhPhe]-[Cys]-NH ₂ ;	Cys-Cys form a disulfide bond	***	**
33	33	Isovaleric Acid-E-T-H-[Cys]-P-[Cys]-I-[(D)Lys]-[bhPhe]-[Lys(1PEG2_1PEG2_IsoGlu_C18_Diacid)]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	*	
34	34	Isovaleric Acid-E-T-H-[Dpa]-[Cys]-[Cys]-I-[(D)Lys]-[bhPhe]-[Lys(1PEG2_1PEG2_IsoGlu_C18_Diacid)]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	**	
36	36	Isovaleric Acid-E-T-H-[Dpa]-P-[Cys]-I-[Cys]-[bhPhe]-[Lys(1PEG2_1PEG2_IsoGlu_C18_Diacid)]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	***	***
39	39	Isovaleric Acid-E-T-H-[Dpa]-P-[Cys]-[Cys]-[Lys(Ahx_Palm)]-[bhPhe]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	***	***

TABLE 6A-continued

Illustrative Hepcidin Analogues					
Compd ID	SEQ ID No.	Peptide	Cyclized [@]	FPN IC ₅₀ (nM)	T47D MSA IC ₅₀ (nM)
40	40	Isovaleric Acid-[Glu_OMe]-T-H-[Dpa]-P-[Cys]-[Cys]-[Lys(Ahx_Palm)]-[bhPhe]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	***	****
41	41	Isovaleric Acid-E-T-H-[Dpa]-P-[Cys]-[Cys]-[Lys(1PEG2_1PEG2_Dap_C18_Diacid)]-[bhPhe]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	***	***
42	42	Isovaleric Acid-[Glu_OMe]-T-H-[Dpa]-P-[Cys]-[Cys]-[Lys(1PEG2_1PEG2_Dap_C18_Diacid)]-[bhPhe]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	***	***
43	43	Isovaleric Acid-[Cys]-T-H-[Dpa]-P-A-I-[Cys]-[bhPhe]-[Lys(Ahx_Palm)]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	****	*
44	44	Isovaleric Acid-E-T-H-[Dpa]-P-[Cys]-[MeCys]-[Lys(Ahx_Palm)]-[bhPhe]-[(D)Lys]-NH ₂ ;	Cys-Cys form a disulfide bond	****	****
45	45	Isovaleric Acid-E-[Cys]-H-[Dpa]-[Cys]-A-I-[(D)Lys]-[bhPhe]-[Lys(Ahx_Palm)]-[(D)Lys]-A-NH ₂ ;	Cys-Cys form a disulfide bond	***	****
46	46	Isovaleric Acid-E-[Cys]-H-[Dpa]-[(D)Cys]-A-I-[(D)Lys]-[bhPhe]-[Lys(Ahx_Palm)]-[(D)Lys]-A-NH ₂ ;	Cys-Cys form a disulfide bond	****	

TABLE 6B

Illustrative Hepcidin Analogues					
Compd ID	SEQ ID No.	Peptide	Cyclized [@]	FPN IC ₅₀ (nM)	T47D MSA IC ₅₀ (nM)
101	101	Isovaleric Acid-D-T-H-F-P-[Cys]-[Cys]-I-[Lys(IsoGlu_Palm)]-F-E-P-R-OH;	Cys-Cys form a disulfide bond	***	

[0661] In Tables 6A-6E, for FPN and T47D internalization assays, the symbols representing the IC₅₀ values have the following meanings: ****=1 nM≤IC₅₀≤10 nM; ***=10 nM<IC₅₀≤100 nM; **=100 nM<IC₅₀≤500 nM; *=500 nM. Where not shown, data has not become available yet.

TABLE 6C

Illustrative Hepcidin Analogues					
SEQ ID No.	Peptide	Cyclized [@]	FPN IC ₅₀ (nM)	T47D MSA IC ₅₀ (nM)	
102	Isovaleric_Acid-[Hcy]-T-H-[DIP]-P-A-I-C-[bhF]-[Lys_Ahx_Palm]-dK-OH				
103	Isovaleric_Acid-[dPen]-T-H-DIP-P-A-I-C-[bhF]-[Lys_Ahx_Palm]-dK-NH ₂	Cys-dPen form a disulfide bond	**		

TABLE 6C-continued

Illustrative Hecpudin Analogues				
SEQ ID No.	Peptide	Cyclized [@]	FPN IC ₅₀ (nM)	T47D MSA IC ₅₀ (nM)
104	Isovaleric__Acid-C-T-H-[DIP]-P-A-I-[dPen]-[bhF]-[Lys__Ahx__Palm]-dK-NH ₂	Cys-dPen form a disulfide bond	***	
105	Isovaleric__Acid-[dC]-T-H-[DIP]-P-A-I-[dC]-[bhF]-[Lys__Ahx__Palm]-[dK]-NH ₂	dCys-dCys form a disulfide bond	***	
106	Isovaleric__Acid-C-T-H-[DIP]-P-A-I-[dC]-[bhF]-[Lys__Ahx__Palm]-dK-NH ₂	Cys-dCys form a disulfide bond	**	
107	Isovaleric__Acid-[dC]-T-H-[DIP]-P-A-I-C-[bhF]-[Lys__Ahx__Palm]-[dK]-NH ₂	dCys-Cys form a disulfide bond	***	
108	Isovaleric__Acid-C-T-H-[DIP]-P-A-I-[Hcy]-[bhF]-[Lys__Ahx__Palm]-[dK]-NH ₂	Cys-hCys form a disulfide bond	***	
109	Isovaleric__Acid-[Hcy]-T-H-[DIP]-P-A-I-[Pen]-[bhF]-[Lys__Ahx__Palm]-[dK]-NH ₂	hCys-Pen form a disulfide bond	***	
110	Isovaleric__Acid-[Hcy]-T-H-[DIP]-P-A-I-C-[bhF]-[Lys__Ahx__Palm]-[dK]-NH ₂	hCys-Cys form a disulfide bond	****	
111	Isovaleric__Acid-[Hcy]-T-H-[DIP]-P-A-I-[Hcy]-[bhF]-[Lys__Ahx__Palm]-[dK]-NH ₂	hCys-hCys form a disulfide bond	***	

TABLE 6D

Illustrative Hecpudin Analogues				
SEQ ID No.	Peptide	Cyclized [@]	FPN IC ₅₀ (nM)	T47D MSA IC ₅₀ (nM)
112	Isovaleric__Acid-E-C-H-[DIP]-C-[Lys__1PEG2__1PEG2__Dap__C18__Diacid]-[N__Butyl__Phe]	Cys-Cys form a disulfide bond		
113	Isovaleric__Acid-E-C-H-[DIP]-C-[Lys__Me3]-[N__Butyl__Phe]	Cys-Cys form a disulfide bond		
114	Isovaleric__Acid-E-C-H-[DIP]-C-[Lys__Ahx__Palm]-[N__Butyl__Phe]	Cys-Cys form a disulfide bond		
115	Isovaleric__Acid-E-C-H-[DIP]-C-[Lys__DMG__N__2ae__Palm]-[N__Butyl__Phe]	Cys-Cys form a disulfide bond		
116	Isovaleric__Acid-E-C-H-[DIP]-C-[Lys__Dap__Palm]-[N__Butyl__Phe]	Cys-Cys form a disulfide bond		
117	Isovaleric__Acid-E-C-H-[DI]P-C-[Lys__Ahx__Dap__Palm]-[N__Butyl__Phe]	Cys-Cys form a disulfide bond		

TABLE 6D-continued

Illustrative Hepcidin Analogues			
SEQ ID No.	Peptide	Cyclized [@]	FPN IC ₅₀ (nM)
			T47D MSA IC ₅₀ (nM)
118	Isovaleric_Acid-E-C-H-[DIP]-C-[Lys_1PEG2_1PEG2_Dap_Palm]-[N_Butyl_Phe]	Cys-Cys form a disulfide bond	
119	Isovaleric_Acid-E-C-H-[DIP]-[NMe-Cys]-[Lys_1PEG2_1PEG2_DMKG_N_2ae_Palm]-[N_Butyl_Phe]	Cys-NMe-Cys form a disulfide bond	
120	Isovaleric_Acid-E-C-H-[DIP]-[Hcy]-[Lys_1PEG2_1PEG2_DMKG_N_2ae_Palm]-[N_Butyl_Phe]	Cys-Hcy form a disulfide bond	
121	Isovaleric_Acid-E-C-H-[DIP]-C-[Lys_1PEG2_1PEG2_DMKG_N_2ae_Palm]-Diphenylethylamine	Cys-Cys form a disulfide bond	***
122	Isovaleric_Acid-E-C-H-[DIP]-C-[Lys_1PEG2_1PEG2_DMKG_N_2ae_Palm]-[N_Butyl_Phe]	Cys-Cys form a disulfide bond	****
123	Isovaleric_Acid-[Glu_OME]-C-H-[DIP]-C-[Lys_DMKG_N_2ae_Palm]-Amylamine	Cys-Cys form a disulfide bond	*
124	Isovaleric_Acid-E-C-H-[DIP]-C-[Lys_DMKG_N_2ae_Palm]-Amylamine	Cys-Cys form a disulfide bond	**
125	Isovaleric_Acid-E-C-H-[DIP]-[4S_Mcp]-[Lys_Ahx_DMKG_N_2ae_Palm]-[N_Butyl_Phe]	Cys-[4S_Mcp] form a disulfide bond	*** *
126	Isovaleric_Acid-E-C-H-[DIP]-[4R_Mcp]-[Lys_Ahx_DMKG_N_2ae_Palm]-[N_Butyl_Phe]	Cys-4R_Mcp form a disulfide bond	*** ***
127	Isovaleric_Acid-E-[4S_Mcp]-H-[DIP]-[4R_Mcp]-[Lys_1PEG2_1PEG2_Dap_Palm]-[N_Butyl_Phe]	[4S_Mcp]-[4R_Mcp] form a disulfide bond	*** ****
128	Isovaleric_Acid-E-C-H-[DIP]-dC-[Lys_Ahx_Palm]-[DIP]-NH ₂	Cys-dCys form a disulfide bond	*** **
129	Isovaleric_Acid-E-C-H-[DIP]-C-[Lys_Ahx_Palm]-[DIP]-NH ₂	Cys-Cys form a disulfide bond	*
130	Isovaleric_Acid-E-C-H-[DIP]-C-[Hexadecane_Amine]	Cys-Cys form a disulfide bond	
131	Isovaleric_Acid-E-C-H-[DIP]-C-[12-aminododecanoic acid]	Cys-Cys form a disulfide bond	
132	Isovaleric_Acid-E-C-H-[DIP]-C-[Dodecyl_Amine]	Cys-Cys form a disulfide bond	
133	Isovaleric_Acid-E-C-H-[DIP]-C-[Lys_Ahx_Palm]-OH	Cys-Cys form a disulfide bond	
134	Isovaleric_Acid-E-C-H-[DIP]-P-C-I-[dK]-[bhF]-[Lys_Ahx_Palm]-[dK]-NH ₂	Cys-Cys form a disulfide bond	***

TABLE 6E

Illustrative Hepcidin Analogues			
SEQ ID No.	Peptide	Cyclized@	FPN IC ₅₀ (nM)
			T47D MSA IC ₅₀ (nM)
135	[MOMHA]-T-H-[DIP]-P-C-C-[dK]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	**
136	[MOMHA]-T-H-[DIP]-P-C-C-[Lys_Me3]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	
137	[MOMHA]-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_DMG_N_2ae_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
138	[MOMHA]-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_DMG_N_2ae_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
139	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[Lys_Me3]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	
140	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[Lys_Me3]-[dDIP]-NH ₂	Cys-Cys form a disulfide bond	***
141	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[Lys_Me3]-[DIP]-NH ₂	Cys-Cys form a disulfide bond	***
142	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_Palm]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond; and Glu and dLys form an amide linkage	**** **
143	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_DMG_N_2ae_Palm]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond and E and dLys form an amide linkage	*** ****
144	Isovaleric_Acid-E-[NMe_Thr]-H-[DIP]-P-C-C-[NMe_Lys_DMG_N_2ae_C18_Diacid]-[DIP]-NH ₂	Cys-Cys form a disulfide bond	
145	Isovaleric_Acid-[Glu_OMe]-[aMe_Pro]-H-[DIP]-P-C-C-[NMe_Lys_Ahx_DMG_N_2ae_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	***
146	Isovaleric_Acid-[Glu_OMe]-[Thr_Me]-H-[DIP]-P-C-C-[NMe_Lys_DMG_N_2ae_C18_Diacid_OMe]-[DIP]-NH ₂	Cys-Cys form a disulfide bond	**
147	Isovaleric_Acid-[Glu_OMe]-[Thr_Me]-H-[DIP]-P-C-C-[NMe_Lys_Ahx_DMG_N_2ae_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
148	Isovaleric_Acid-[Glu_OMe]-S-H-[NMe_Phe]-P-C-C-[NMe_Lys_Ahx_DMG_N_2ae_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	* *
149	Isovaleric_Acid-[Glu_OMe]-[NMe_Thr]-H-[DIP]-P-C-C-[NMe_Lys_DMG_N_2ae_Palm]-[DIP]-[NMe_Lys]-NH ₂	Cys-Cys form a disulfide bond	***
150	Isovaleric_Acid-E-[Hyp_3R]-H-[DIP]-P-C-C-[NMe_Lys_DMG_N_2ae_Palm]-[DIP]-NH ₂	Cys-Cys form a	****

TABLE 6E-continued

Illustrative Hepcidin Analogues			
SEQ ID No.	Peptide	Cyclized@	FPN IC ₅₀ (nM)
			T47D MSA IC ₅₀ (nM)
151	Isovaleric_Acid-[Glu_OMe]-[Hyp_3R]-H-[DIP]-P-C-C-[NMe_Lys_DMG_N_2ae_Palm]-[DIP]-[dK]-NH ₂	disulfide bond Cys-Cys form a disulfide bond	****
152	Isovaleric_Acid-E-[Hyp_3R]-H-[DIP]-P-C-C-[NMe_Lys_DMG_N_2ae_Palm]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	****
153	Isovaleric_Acid-[Glu_OMe]-[NMe_Thr]-H-[DIP]-P-C-C-[NMe_Lys_Ahx_DMG_N_2ae_C18_Diacid]-[DIP]-[NMe_Lys]-NH ₂	Cys-Cys form a disulfide bond	**
154	Isovaleric_Acid-E-[Hyp_3R]-H-[DIP]-P-C-C-[NMe_Lys_Ahx_DMG_N_2ae_C18_Diacid]-[DIP]-NH ₂	Cys-Cys form a disulfide bond	****
155	Isovaleric_Acid-[Glu_OMe]-[Hyp_3R]-H-[DIP]-P-C-C-[NMe_Lys_Ahx_DMG_N_2ae_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	***
156	Isovaleric_Acid-E-[Hyp_3R]-H-[DIP]-P-C-C-[NMe_Lys_Ahx_DMG_N_2ae_C18_Diacid]-[DIP]-[NMe_Lys]-NH ₂	Cys-Cys form a disulfide bond	***
157	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_Dap_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	***
158	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_DMG_N_2ae_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	***
159	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_DMG_N_2ae_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	***
160	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_DMG_N_2ae_C18_Diacid_18_OMe]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond; and E and dK form an amide linkage	***
161	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_Palm]-[DIP]-N_Ethylamine	Cys-Cys form a disulfide bond; and E and N_Ethylamine form an amide linkage	**** **
162	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[hC_CH ₂ CH ₂ CH ₂ NH_Palm]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** ****
163	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_Palm]-Diphenylethylamine	Cys-Cys form a disulfide bond	*** ****
164	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_Palm]-[DIP]-N_Ethylamine	Cys-Cys form a disulfide bond	****
165	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_Palm]-[DIP]-[dDap]-NH ₂	Cys-Cys form a disulfide bond	****

TABLE 6E-continued

Illustrative Hepcidin Analogues			
SEQ ID No.	Peptide	Cyclized@	FPN IC ₅₀ (nM)
			T47D MSA IC ₅₀ (nM)
166	Isovaleric_Acid-[NMe_Glu]-T-H-[DIP]-P-C- C-[NMe_Lys_Ahx_Palm]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	****
167	Isovaleric_Acid-[NMe_Glu]-T-H-[DIP]-P-C- C-[NMe_Lys_Ahx_DMGMG_N_2ae_Palm]- [DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	****
168	Isovaleric_Acid-[hSer]-T-H-[DIP]-P-C-C- [NMe_Lys_Ahx_Palm]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	****
169	Isovaleric_Acid-E-[NMe_Thr]-H-[DIP]-P-C- C-[NMe_Lys_DMGMG_N_2ae_Palm]-[DIP]-NH ₂	Cys-Cys form a disulfide bond	****
170	Isovaleric_Acid-E-[NMe_Thr]-H-[DIP]-P-C- C-[NMe_Lys_Ahx_DMGMG_N_2ae_Palm]- [DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	****
171	Isovaleric_Acid-E-P-H-[DIP]-P-C-C- [NMe_Lys_Ahx_Palm]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	****
172	Isovaleric_Acid-E-[NMe_Thr]-H-[DIP]-P-C- C-[NMe_Lys_Ahx_Palm]-[DIP]-NH ₂	Cys-Cys form a disulfide bond	****
173	Isovaleric_Acid-E-[NMe_Thr]-H-[DIP]-P-C- C-[NMe_Lys_Ahx_Palm]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	****
174	Isovaleric_Acid-E-[Hyp_3R]-H-[DIP]-P-C-C- [NMe_Lys_Ahx_Palm]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	***
175	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C- C-[NMe_Lys_DMGMG_N_2ae_Palm]-[DIP]- [dK]-NH ₂	Cys-Cys form a disulfide bond	****
176	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C- C-[NMe_Lys_DMGMG_N_2ae_IsoGlu_Palm]- [DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	
177	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C- C-[NMe_Lys_Ahx_Palm]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	
178	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C- C-[NMe_Lys_Ahx_DMGMG_N_2ae_Palm]- [DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	
179	Isovaleric_Acid-E-T-H-[DIP]-P-C-C- [NMe_Lys_Ahx_Palm]-[DIP]-NH ₂	Cys-Cys form a disulfide bond	
180	Isovaleric_Acid-E-T-[3Pal]-[DIP]-P-C-C- [NMe_Lys_Ahx_Palm]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	
181	Isovaleric_Acid-E-T-H-DIP-P-C-C- [NMe_Lys_Ahx_Palm]-[DIP]-[dK]- [NMe_Phe]-NH ₂	Cys-Cys form a disulfide bond	
182	Isovaleric_Acid-E-T-H-[DIP]-P-C-C- [NMe_Lys_Ahx_Palm]-[DIP]-Tle-NH ₂	Cys-Cys form a disulfide bond	

TABLE 6E-continued

Illustrative Hepcidin Analogues			
SEQ ID No.	Peptide	Cyclized@	FPN IC ₅₀ (nM)
			T47D MSA IC ₅₀ (nM)
183	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_DMG_N_2ae_C18_Diacid]-[Bip]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
184	Isovaleric_Acid-[Glu_OMe]-[NMe_Thr]-H-[DIP]-P-C-C-[NMe_Phe]-NH ₂	Cys-Cys form a disulfide bond	* **
185	Isovaleric_Acid-[Glu_OMe]-[NMe_Thr]-H-[DIP]-P-C-C-[NMe_Lys_DMG_N_2ae_C18_Diacid]-[NMe_Phe]-NH ₂	Cys-Cys form a disulfide bond	* **
186	Isovaleric_Acid-[Glu_OMe]-[NMe_Thr]-H-[DIP]-P-C-C-[NMe_Lys_DMG_N_2ae_C18_Diacid]-[DIP]-NH ₂	Cys-Cys form a disulfide bond	*** **
187	Isovaleric_Acid-E-[NMe_Thr]-H-[DIP]-P-C-C-[NMe_Lys_DMG_N_2ae_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
188	Isovaleric_Acid-E-P-H-[DIP]-P-C-C-[NMe_Lys_Ahx_DMG_N_2ae_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	* **
189	Isovaleric_Acid-E-[NMe_Thr]-H-[DIP]-P-C-C-[NMe_Lys_Ahx_DMG_N_2ae_C18_Diacid]-[DIP]-NH ₂	Cys-Cys form a disulfide bond	*** **
190	Isovaleric_Acid-E-[NMe_Thr]-H-[DIP]-P-C-C-[NMe_Lys_Ahx_DMG_N_2ae_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
191	Isovaleric_Acid-E-[Hyp_3R]-H-[DIP]-P-C-C-[NMe_Lys_Ahx_DMG_N_2ae_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
192	Isovaleric_Acid-E-[Hyp_3S]-H-[DIP]-P-C-C-[NMe_Lys_Ahx_DMG_N_2ae_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
193	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_DMG_N_2ae_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
194	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_DMG_N_2ae_Palm]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
195	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_DMG_N_2ae_IsoGlu_Palm]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
196	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_Palm]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
197	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_DMG_N_2ae_Palm]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
198	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_DMG_N_2ae_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
199	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_DMG_N_2ae_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
200	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[Lys_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **

TABLE 6E-continued

Illustrative Hepcidin Analogues			
SEQ ID No.	Peptide	Cyclized@	FPN IC ₅₀ (nM)
			T47D MSA IC ₅₀ (nM)
201	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[Lys_Ahx_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	
202	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[Lys_Ahx_DMG_N_2ae_C18_Diacid]-[DIP]-[dLys_PEG11_OMe]-NH ₂	Cys-Cys form a disulfide bond	
203	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[Lys_1PEG2_1PEG2_DMG_N_2ae_C18_Diacid]-[DIP]-[dLys_PEG11_OMe]-NH ₂	Cys-Cys form a disulfide bond	***
204	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[Lys_1PEG2_1PEG2_DMG_N_2ae_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	
205	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[Lys_Ahx_DMG_N_2ae_C18_Diacid]-[bhF]-[dLys_PEG11_OMe]-NH ₂	Cys-Cys form a disulfide bond	
206	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[NMe_Lys_Dap_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
207	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_Dap_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	**** **
208	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[dLys_Ahx_DMG_N_2ae_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	***
209	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[Lys_DMG_N_2ae_DMG_N_2ae_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	***
210	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[Lys_Ahx_DMG_N_2ae_C18_Diacid]-[DIP]-Tle-NH ₂	Cys-Cys form a disulfide bond	***
211	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[Lys_Ahx_DMG_N_2ae_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	***
212	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[dLys_Ahx_DMG_N_2ae_C18_Diacid]-[DIP]-Tle-NH ₂	Cys-Cys form a disulfide bond	***
213	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[dLys_Ahx_DMG_N_2ae_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	***
214	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[DIP]-[dLys_PEG11_OMe]-NH ₂	Cys-Cys form a disulfide bond	*** **
215	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[Bip]-[dLys_PEG11_OMe]-NH ₂	Cys-Cys form a disulfide bond	**** **
216	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[Bip]-[dK]-NH ₂	Cys-Cys form a disulfide bond	**** ****
217	Isovaleric_Acid-E-T-H-[DIP]-[Morph]-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	***
218	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys and Cys cyclized through a p-	***

TABLE 6E-continued

Illustrative Hepcidin Analogues			
SEQ ID No.	Peptide	Cyclized@	T47D MSA IC ₅₀ (nM)
		Xylene linker	
219	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dLys_Carnitine_Alkyl]-NH ₂	Cys-Cys form a disulfide bond	***
220	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[NMe_Lys_IsoGlu_Palm]-[bhF]-[dLys_PEG11_OMe]-NH ₂	Cys-Cys form a disulfide bond	****
221	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[NMe_Lys_IsoGlu_Palm]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	****
222	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_IsoGlu_Palm]-[bhF]-[dLys_PEG11_OMe]-NH ₂	Cys-Cys form a disulfide bond	*** **
223	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_IsoGlu_Palm]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	****
224	Isovaleric_Acid-E-T-H-[DIP]-[Tic]-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	**
225	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[NMe_Phe]-[NMe_dLys]-NH ₂	Cys-Cys form a disulfide bond	*** **
226	Isovaleric_Acid-L-T-[3Pal]-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[aMe_Phe]-[dK]-NH ₂	Cys-Cys form a disulfide bond	**
227	Isovaleric_Acid-L-T-[3Pal]-[DIP]-[bhP]-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	**
228	Isovaleric_Acid-L-T-H-[DIP]-[bhP]-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	***
229	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_DMG_N_2ae_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	
230	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_DMG_N_2ae_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	***
231	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[NMeLys_1PEG2_1PEG2_DMG_N_2ae_C18Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	***
232	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_DMG_N_2ae_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
233	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
234	Isovaleric_Acid-E-[hSer]-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	**
235	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dLys_PEG11_OMe]-NH ₂	Cys-Cys form a disulfide bond	

TABLE 6E-continued

Illustrative Hepcidin Analogues			
SEQ ID No.	Peptide	Cyclized@	FPN IC ₅₀ (nM)
			T47D MSA IC ₅₀ (nM)
236	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dLys_PEG11_OMe]-NH ₂	Cys-Cys form a disulfide bond	
237	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[Mor_propanoic_acid]-NH ₂	Cys-Cys form a disulfide bond	
238	Isovaleric_Acid-L-T-[3Pal]-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[Phe_4COOH]-[dK]-NH ₂	Cys-Cys form a disulfide bond	
239	Isovaleric_Acid-L-T-[3Pal]-[DIP]-[Npc]-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	
240	Isovaleric_Acid-L-T-H-[DIP]-[Npc]-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	
241	Isovaleric_Acid-L-T-[3Pal]-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	**
242	Isovaleric_Acid-Aad-T-[3Pal]-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
243	Isovaleric_Acid-E-T-[3Pal]-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[DIP]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
244	Isovaleric_Acid-[Aad]-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	***
245	Isovaleric_Acid-[Aad]-T-H-[DIP]-[Npc]-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	***
246	Isovaleric_Acid-[Aad]-T-[3Pal]-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	***
247	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-[NMe_dSer]-NH ₂	Cys-Cys form a disulfide bond	*** ****
248	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-[NMe_dGln]-NH ₂	Cys-Cys form a disulfide bond	*** **
249	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-[NMe_dLeu]-NH ₂	Cys-Cys form a disulfide bond	*** **
250	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-[NMe_dPhe]-NH ₂	Cys-Cys form a disulfide bond	*** ****
251	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-[NMe_Gln]-NH ₂	Cys-Cys form a disulfide bond	***
252	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-[NMe_Leu]-NH ₂	Cys-Cys form a disulfide bond	

TABLE 6E-continued

Illustrative Hepcidin Analogues			
SEQ ID No.	Peptide	Cyclized@	FPN IC ₅₀ (nM)
			T47D MSA IC ₅₀ (nM)
253	Isovaleric_Acid-E-[Hyp_3S]-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	***
254	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dLys_PEG11_OMe]-NH ₂	Cys-Cys form a disulfide bond	*** **
255	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dLys_PEG11_OMe]-NH ₂	Cys-Cys form a disulfide bond	*** **
256	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-NH ₂	Cys-Cys form a disulfide bond	**** **
257	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-[NMe_dTyr]-NH ₂	Cys-Cys form a disulfide bond	*** **
258	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-[1Na]-NH ₂	Cys-Cys form a disulfide bond	*** **
259	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-[NMe_Tyr]-NH ₂	Cys-Cys form a disulfide bond	*** **
260	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-[NMe_Ser]-NH ₂	Cys-Cys form a disulfide bond	*** **
261	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-[Na]-NH ₂	Cys-Cys form a disulfide bond	*** **
262	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-[NMe_Ile]-NH ₂	Cys-Cys form a disulfide bond	*** **
263	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-[NMe_Phe]-NH ₂	Cys-Cys form a disulfide bond	*** **
264	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[Dab]-NH ₂	Cys-Cys form a disulfide bond	*** **
265	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[Nva_Morphino]-NH ₂	Cys-Cys form a disulfide bond	*** **
266	Isovaleric_Acid-E-T-H-[DIP]-P-C-[NMe_Cys]-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[NMe_Phe]-[dK]-NH ₂	Cys-NMe_Cys form a disulfide bond	***
267	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[NMe_dLys_Ac]-NH ₂	Cys-Cys form a disulfide bond	**
268	Isovaleric_Acid-E-T-H-[DIP]-P-[Hcy]-[Hcy]-[Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	homoCys-homoCys form a disulfide bond	*** **
269	Isovaleric_Acid-E-T-H-[DIP]-P-[Hcy]-[Hcy]-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	homoCys-homoCys form a disulfide bond	*** **

TABLE 6E-continued

Illustrative Hepcidin Analogues				T47D	
SEQ ID No.	Peptide	Cyclized@	FPN IC ₅₀ (nM)	MSA IC ₅₀ (nM)	
270	Isovaleric_Acid-E-T-H-[DIP]-P-C-[Hcy]-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-homoCys form a disulfide bond	***	***	
271	Isovaleric_Acid-E-T-H-[DIP]-P-[Pen]-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Pen-Cys form a disulfide bond	***		
272	Isovaleric_Acid-E-T-H-[DIP]-P-C-[Pen]-[Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Pen-Cys form a disulfide bond	***	***	
273	Isovaleric_Acid-E-T-H-[DIP]-P-[Pen]-[Pen]-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Pen-Pen form a disulfide bond	**		
274	Isovaleric_Acid-E-T-H-[DIP]-P-[Pen]-[Pen]-[Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Pen-Pen form a disulfide bond	***		
275	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[DIP]-[NMe_Lys]-NH ₂	Cys-Cys form a disulfide bond	***	***	
276	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[NMe_dLys]-NH ₂	Cys-Cys form a disulfide bond	***		
277	Isovaleric_Acid-E-T-H-[DIP]-P-C-[NMe_Cys]-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[NMe_Phe]-[NMe_Lys]-NH ₂	Cys-NMe_Cys form a disulfide bond	***		
278	Isovaleric_Acid-E-T-H-[DIP]-P-C-[NMe_Cys]-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-NMe_Cys form a disulfide bond	***		
279	Isovaleric_Acid-E-T-[His_1Me]-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[Me_Phe]-[dK]-NH ₂	Cys-Cys form a disulfide bond	***		
280	Isovaleric_Acid-E-T-[His_1Me]-[DIP]-[Npc]-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*		
281	Isovaleric_Acid-E-T-H-[DIP]-P-C-[NMe_Cys]-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[NMe_Phe]-[dK]-NH ₂	Cys-NMe_Cys form a disulfide bond	**		
282	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_DMG_N_2ae_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	***	***	
283	Isovaleric_Acid-E-T-H-DIP-P-C-C-NMe_Lys_1PEG2_1PEG2_DMG_N_2ae_C18_Diacid-bhF-dK-NH ₂	Cys-Cys form a disulfide bond	***	***	
284	Isovaleric_Acid-E-T-[Trp_5OH]-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*		
285	Isovaleric_Acid-E-T-[His_1Me]-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	***	***	
286	Isovaleric_Acid-E-T-[Phe_4CF3]-[DIP]-P-C-C-	Cys-Cys form a	**		

TABLE 6E-continued

Illustrative Hepcidin Analogues			
SEQ ID No.	Peptide	Cyclized@	FPN IC ₅₀ (nM)
			T47D MSA IC ₅₀ (nM)
287	[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂ Isovaleric_Acid-E-T-[Trp_6OMe]-[DIP]-P-C-C-	disulfide bond Cys-Cys form a	*
288	[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂ Isovaleric_Acid-[Tet1]-T-H-[DIP]-P-C-C-	disulfide bond Cys-Cys form a	*** **
289	[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂ Isovaleric_Acid-[Dap]-T-H-[DIP]-P-C-C-	disulfide bond Cys-Cys form a	**
290	[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[Tle]-NH ₂ Isovaleric_Acid-E-T-H-[DIP]-P-C-C-	disulfide bond Cys-Cys form a	*** **
291	[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[NMe_Phe]-[dK]-NH ₂ Isovaleric_Acid-E-T-H-[DIP]-P-C-C-	disulfide bond Cys-Cys form a	*** **
292	[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[NMe_Lys]-NH ₂ Isovaleric_Acid-E-T-H-[DIP]-P-C-C-	disulfide bond Cys-Cys form a	*** **
293	[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[DIP]-[dK]-NH ₂ Isovaleric_Acid-E-T-H-[DIP]-P-C-C-	disulfide bond Cys-Cys form a	*** **
294	[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-NH ₂ Isovaleric_Acid-E-T-H-[DIP]-P-C-C-	disulfide bond Cys-Cys form a	*** **
295	[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-NH ₂ Isovaleric_Acid-E-T-H-[DIP]-P-C-C-	disulfide bond Cys-Cys form a	*** **
296	[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂ Isovaleric_Acid-E-T-[Ala_3Quin]-[DIP]-P-C-C-	disulfide bond Cys-Cys form a	* *
297	[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂ Isovaleric_Acid-E-T-[Bip]-[DIP]-P-C-C-	disulfide bond Cys-Cys form a	*
298	[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂ 4_Fluorophenylacetic_acid-E-T-H-[DIP]-P-C-C-	disulfide bond Cys-Cys form a	*** **
299	[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂ Nicotinic_acid-E-T-H-[DIP]-P-C-C-	disulfide bond Cys-Cys form a	*** **
300	[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂ Inp-E-T-H-[DIP]-P-C-C-	disulfide bond Cys-Cys form a	** **
301	[aMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂ Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-	disulfide bond Cys-Cys form a	*** **
302	[aMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂ Isovaleric_Acid-E-T-H-[DIP]-P-C-C-	disulfide bond Cys-Cys form a	*** **
303	[NMe_Lys_PEG12_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂ Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-	disulfide bond Cys-Cys form a	*** **

TABLE 6E-continued

Illustrative Hepcidin Analogues			
SEQ ID No.	Peptide	Cyclized@	FPN IC ₅₀ (nM)
			T47D MSA IC ₅₀ (nM)
304	Isovaleric_Acid-E-T-H-DIP-P-C-C-[NMe_Lys_PEG12_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
305	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
306	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_1PEG2_1PEG2_Dap_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
307	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_Palm]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
308	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[NMe_Lys_Ahx_Palm]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
309	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[Lys_DMG_N_2ae_DMG_N_2ae_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
310	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[Lys_DMG_N_2ae_DMG_N_2ae_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
311	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[Lys_Ahx_DMG_N_2ae_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
312	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[Lys_Ahx_DMG_N_2ae_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
313	Isovaleric_Acid-[Glu_OMe]-H-[DIP]-P-C-C-[Lys_DMG_N_2ae_DMG_N_2ae_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
314	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[Lys_DMG_N_2ae_DMG_N_2ae_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
315	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[Lys_Ahx_DMG_N_2ae_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
316	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[Lys_Ahx_DMG_N_2ae_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
317	Isovaleric_Acid-[Glu_OMe]-T-H-[DIP]-P-C-C-[Lys_DMG_N_2ae_DMG_N_2ae_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **
318	Isovaleric_Acid-E-T-H-[DIP]-P-C-C-[Lys_DMG_N_2ae_DMG_N_2ae_C18_Diacid]-[bhF]-[dK]-NH ₂	Cys-Cys form a disulfide bond	*** **

Example 2C

Activity of Peptide Analogues

[0662] The potency of the peptides in causing ferroportin internalization was evaluated in a T47D cell-based assay. T47D cell line (HTB 133, ATCC) is a human breast carci-

noma adherent cell line which endogenously expresses ferroportin. In this internalization assay, the potency of the test peptides was evaluated in presence of serum albumin which is the main protein component in the blood. T47D cells were maintained in RPMI media (containing required amount of fetal bovine serum) and regularly sub-cultured. In preparation for the assay, the cells were seeded in 96-well

plates at a density of 80-100 k cells per well in 100 μ l volume and allowed to rest overnight. On the next day, test peptides were first prepared in dilution series (10-point series, starting concentration of $\sim 5 \mu$ M, typically 3-4 $\times$ fold dilution steps), all with 0.5% mouse serum albumin (MSA purified from mouse serum; Sigma, A3139). The test peptide dilution series were allowed to incubate at room temperature for 30 min. Then the media was aspirated from the 96-well cell plate and test peptide dilution series were added. After 1 hr incubation, the media with test peptides was aspirated out and AF647-conjugated detection peptide was added at fixed concentration of 200 nM. The AF647-conjugated detection peptide was previously verified to bind to ferroportin and cause its internalization. The cells were washed again after a 2 hr incubation in preparation for flow cytometry analysis. The Median Fluorescence Intensity (MFI) of the AF647-positive population was measured (after removing dead cells and non-singlets from the analysis). The MFI values were used to generate a dose-response curve and obtain IC₅₀ potencies for the test peptides. The IC₅₀ potencies were calculated by using 4-parameter non-linear fitting function in Graphpad Prism (Tables 6A-B).

Example 2D

LAD2 Activity of Peptide Analogues

[0663] In anaphylactoid reactions, the main mechanism involves the direct stimulation of mast cells or basophils leading to the release of anaphylactic mediators such as histamine and β -hexosaminidase. A recent study by McNeil et al. (McNeil B D et al., 2015) reported that MrgprX2, a specific membrane receptor on human mast cells, induces anaphylactoid reactions. The LAD2 (Laboratory of Allergic Diseases 2) human mast cell line derived from human mast cell sarcoma/leukemia (Kirshenbaum et al., 2003), is commonly employed to study anaphylactoid reactions because its biological properties are identical to those of primary human mast cells including the overexpression of the MrgprX2 receptor and sensitivity towards degranulating peptides (Kulka et al., 2008). The release of anaphylactic mediators such as (3-hexosaminidase, is assessed by fluorometric quantification.

[0664] The degranulation potential of hepcidin mimetics were evaluated in the LAD2 cells. On the day of the assay, serial dilutions of compounds were added to LAD2 cells plated at 20000 cells/well in a 96-well plate. After incubation for 30 minutes, the amount of β -hexosaminidase released into the supernatants and in cell lysates was quantified using the fluorogenic substrate 4-methylumbelliferyl-N-acetyl-b-D-glucosaminide. Dose-response curves were generated by plotting the % of β -hexosaminidase release (y-axis) against the concentrations of peptides tested (x-axis). The EC₅₀ values and standard errors were calculated using XLfit 5.5.0.5 based on the following equation: 4 Parameter Sigmoidal Model: $f = (A + ((B - A) / (1 + ((C/x)^D))))$ where A=Emin, B=Emax, C=EC₅₀ and D=slope.

[0665] References: McNeil B D et al., Nature, 12, 519 (2015); Kirshenbaum et al. Leukemia Res. 27, 677 (2003); Kulka et al. Immunology 123, 398 (2008).

Example 3

In Vivo Validation of Peptide Analogues

[0666] Hepcidin analogues of the present invention were tested for in vivo activity, to determine their ability to decrease free Fe²⁺ in serum.

[0667] A hepcidin analogue or vehicle control were administered to mice (n=3/group) at 1000 nmol/kg either intravenously or subcutaneously. Serum samples were taken from groups of mice administered with the hepcidin analog at 30 min, 1 h, 2 h, 4 h, 10 h, 24 h, 30 h, 36 h, and 48 h post-administration. Iron content in plasma/serum was measured using a colorimetric assay on the Cobas c 111 according to instructions from the manufacturer of the assay (assay: IRON2: ACN 661).

[0668] In another experiment, various hepcidin analogues or vehicle control were administered to mice (n=3/group) at 1000 nmol/kg subcutaneously. Serum samples were taken from groups of mice administered with vehicle or hepcidin analog at 30 h and 36 h post-administration. Iron content in plasma/serum was measured using a colorimetric assay on the Cobas c 111 according to instructions from the manufacturer of the assay (assay: IRON2: ACN 661).

[0669] These studies demonstrate that hepcidin analogues of the present invention reduce serum iron levels for at least 30 hours, thus demonstrating their increased serum stability.

Example 4

In Vitro Validation of Peptide Analogues

[0670] Based in part on the structure activity relationships (SAR) determined from the results of the experiments described herein, a variety of Hepcidin-like peptides of the present invention were synthesized using the method described in Example 1, and in vitro activity was tested as described in Example 2. Reference compounds included native Hepcidin, Mini-Hepcidin, R1-Mini-Hepcidin, Reference Compound 1 and Reference Compound 2. IC₅₀ or EC₅₀ values of the peptides are shown in summary Tables 6A-6E.

Example 5

Plasma Stability

[0671] Plasma stability experiments were undertaken to complement the in vivo results and assist in the design of potent, stable Ferroportin agonists. In order to predict the stability in rat and mouse plasma, ex vivo stability studies were initially performed in these matrices.

[0672] Peptides of interest (20 μ M) were incubated with pre-warmed plasma (BioreclamationIVT) at 37° C. Aliquots were taken at various time points up to 24 hours (e.g. 0, 0.25, 1, 3, 6 and 24 hr), and immediately quenched with 4 volumes of organic solvent (acetonitrile/methanol (1:1) and 0.1% formic acid, containing 1 μ M internal standard). Quenched samples were stored at 4° C. until the end of the experiment and centrifuged at 17,000 g for 15 minutes. The supernatant were diluted 1:1 with deionized water and analyzed using LC-MS. Percentage remaining at each time point was calculated based on the peak area ratio (analyte over internal standard) relative to the initial level at time zero. Half-lives were calculated by fitting to a first-order exponential decay equation using GraphPad.

Example 6

Reduction of Serum Iron in Mice

[0673] Hepcidin mimetic compounds, designed for oral stability, were tested for systemic absorption by PO dosing in a wild type mouse model C₅₇BL/6. The animals were

acclimatized in normal rodent diet for 4-5 days prior to study start and fasted overnight prior to study start. Groups of 4 animals each received either Vehicle or the Compounds. The compounds were formulated in Saline at a concentration of 5 mg/mL. The mice received dosing solution via oral gavage at volume of 200 μ L per animal of body weight 20 g. Each group received 1 dose of compounds at 50 mg/kg/dose. The group marked for vehicle received only the formulation. Blood was drawn at 4 hours post-dose and serum was prepared for PK and PD measurements. The compound concentration was measured by mass spectrometry method and iron concentration in the samples was measured using the colorimetric method on Roche cobas c system.

Example 7

Reduction of Serum Iron in Mice

[0674] In another experiment, a new set of compounds were tested for systemic absorption by PO dosing in a wild type mouse model C57BL/6. The animals were acclimatized in normal rodent diet for 4-5 days prior to study start. Over the night prior to the first dose, the mice were switched to a low iron diet (with 2 ppm iron) and this diet was maintained during the rest of the study. Groups of 5 animals each received either Vehicle or the Compounds. The concentration of compounds was at 30 mg/mL, formulated in 0.7% NaCl+10 mM NaAcetate buffer. Food was withdrawn around 2 hours prior to each dose to ensure that the stomach was clear of any food particles prior to PO dosing. The mice received dosing solution via oral gavage at volume of 200 μ L per animal of body weight 20 g. Each group received 2 doses of compound at 300 mg/kg/dose, on successive days. The group marked for vehicle received only the formulation. Blood was drawn at 4.5 hours post-last-dose and serum was prepared for PD measurements. Serum iron concentration was measured using the colorimetric method on Roche cobas c system.

Example 8

Pharmacodynamic Effects for the Serum Iron Reducing Abilities of a Representative Compound in Mice

[0675] In a second in vivo study, the representative compound was tested for pharmacodynamic effect with a single dose of 300 mg/kg/dose vs. 2 doses of 300 mg/kg over two days QD (once per day). C57BL/6 mice were acclimatized in normal rodent diet for 4-5 days prior to study start. Over the night prior to the first dose, the mice were switched to a low iron diet (with 2 ppm iron) and this diet was maintained during the rest of the study. Groups of 5 animals each received either Vehicle or the Compounds. The compound was formulated in 0.7% NaCl+10 mM NaAcetate buffer at 30 mg/mL concentration. Food was withdrawn around 2 hours prior to each dose to ensure that the stomach was clear of any food particles prior to PO dosing. The mice received dosing solution via oral gavage at volume of 200 μ L per animal of body weight 20 g.

Example 9

PK/PD Effects of Oral Dosing of a Representative Compound of the Present Invention in Mice

[0676] In another in vivo study with healthy Wild Type mouse model C57BL/6, representative Compound was

tested for PK and PD effect with multiple dosing over three days. The mice were maintained under normal rodent feed during the acclimatization and switched to iron-deficient diet (with ~2 ppm iron) one night prior to the first dose. Groups of 5 mice each received a total of 6 doses of either vehicle or a representative compound of the present invention at different dose strengths, in a BID format over three days. Mice were dosed via oral gavage with the representative compound formulated in 0.7% saline and 10 mM Sodium Acetate. The different groups received either vehicle, 150 mg/kg/dose BID, 75 mg/kg/dose BID, 37.5 mg/kg/dose BID, or 18.75 mg/kg/dose BID. An additional group received 100 mg/kg/dose BID in addition to a total of 100 mg/kg/day of compound in drinking water (DW), thereby receiving a total dose of 300 g/kg/day. At 3 hours post-last-dose the vehicle group marked for iron-challenge and all the compound dosed groups received iron solution via oral gavage at 4 mg/kg iron per animal. Blood was collected at 90 min post-iron-challenge to prepare serum for PK and PD measurements. The compound concentration was measured by mass spectrometry method and iron concentration in the samples was measured using the colorimetric method on Roche cobas c system.

Example 10

Reduction of Serum Iron in Mice

[0677] In a separate triage, a new set of compounds were tested for their pharmacodynamic effect when dosed orally in the wild type mouse model C57BL/6. The animals were acclimatized in normal rodent diet for 4-5 days prior to study start. The group of 5 animals designated to receive two doses of a representative compound received an iron-deficient diet (with 2-ppm iron) on the night prior to the first dose and all the other groups designated for single dose of different compounds were treated with iron-deficient diet for two nights prior to the compound dosing. The concentration of compounds in the dosing solution was at 30 mg/mL, formulated in 0.7% NaCl+10 mM NaAcetate buffer. Food was withdrawn around 2 hours prior to any dosing to ensure that the stomach was clear of any food particles prior to PO dosing. The mice received dosing solution via oral gavage at volume of 200 μ L per animal of body weight 20 g. The group marked for vehicle received only the formulation. Blood was drawn at 4.5 hours post-last-dose and serum was prepared for PD measurements. Serum iron concentration was measured using the colorimetric method on Roche cobas c system.

Example 11

Stability in Simulated Gastric Fluid

[0678] Blank SGF was prepared by adding 2 g sodium chloride, 7 mL hydrochloric acid (37%) in a final volume of 1 L water, and adjusted pH to 1.2.

[0679] SGF was prepared by dissolving 320 mg Pepsin (Sigma®, P6887, from Porcine Stomach Mucosa) in 100 mL Blank SGF and stirred at room temperature for 30 minutes. The solution was filtered through 0.45 μ m membrane and aliquot and stored at -20° C.

[0680] Experimental compounds of interest (at a concentration of 20 μ M) were incubated with pre-warmed SGF at 37° C. Aliquots were taken at various time points up to 24 hours (e.g., 0, 0.25, 1, 3, 6 and 24 hr), and immediately

quenched with 4 volumes of organic solvent (acetonitrile/methanol (1:1) and 0.1% formic acid, containing 1 μM internal standard). Quenched samples were stored at 4° C. until the end of the experiment and centrifuged at 4,000 rpm for 10 minutes. The supernatant were diluted 1:1 with deionized water and analyzed using LC-MS. Percentage remaining at each time point was calculated based on the peak area ratio (analyte over internal standard) relative to the initial level at time zero. Half-lives were calculated by fitting to a first-order exponential decay equation using GraphPad.

Example 12

Stability in Simulated Intestinal Fluids

[0681] Blank FaSSIF was prepared by dissolving 0.348 g NaOH, 3.954 g sodium phosphate monobasic monohydrate and 6.186 g NaCl in a final volume of 1 liter water (pH adjusted to 6.5).

[0682] FaSSIF was prepared by dissolving 1.2 g porcine pancreatin (Chem-supply, PL378) in 100 mL Blank FaSSIF and stirred at room temperature for 30 minutes. The solution was filtered through 0.45 μm membrane and aliquot and stored at -20° C.

[0683] Experimental compounds of interest (20 μM) were incubated with pre-warmed FaSSIF (1% pancreatin in final incubation mixture) at 37° C. Aliquots were taken at various time points up to 24 hours (e.g. 0, 0.25, 1, 3, 6 and 24 hr), and immediately quenched with 4 volumes of organic solvent (acetonitrile/methanol (1:1) and 0.1% formic acid, containing 1 μM internal standard). Quenched samples were stored at 4° C. until the end of the experiment and centrifuged at 4,000 rpm for 10 minutes. The supernatant were diluted 1:1 with deionized water and analyzed using LC-MS. Percentage remaining at each time point was calculated based on the peak area ratio (analyte over internal standard) relative to the initial level at time zero. Half-lives were calculated by fitting to a first-order exponential decay equation using GraphPad.

Example 13

Modified Experimental for Peptides Prone to “Non-Specific Binding”

[0684] Compounds of interest (at concentration of 20 μM) were mixed with pre-warmed FaSSIF (1% pancreatin in final working solution). The solution mixture was aliquoted and incubated at 37° C. The number of aliquots required was equivalent to the number of time points (e.g. 0, 0.25, 1, 3, 6 and 24 hr). At each time point, one aliquot was taken and immediately quenched with 4 volumes of organic solvent (acetonitrile/methanol (1:1) and 0.10% formic acid, containing 1 μM internal standard). The remaining steps were the same as the generic experimental.

[0685] All of the above U.S. patents, U.S. patent application publications, U.S. patent applications, foreign patents, foreign patent applications and non-patent publications referred to in this specification and/or listed in the Application Data Sheet, are incorporated herein by reference, in their entirety.

[0686] At least some of the chemical names or sequences of compounds of the invention as given and set forth in this application, may have been generated on an automated basis by use of a commercially available chemical naming software program, and have not been independently verified. In the instance where the indicated chemical name or sequence and the depicted structure differ, the depicted structure will control. In the chemical structures where a chiral center exists in a structure but no specific stereochemistry is shown for the chiral center, both enantiomers associated with the chiral structure are encompassed by the structure. Similarly, for the peptides where E/Z isomers exist but are not specifically mentioned, both isomers are specifically disclosed and covered.

[0687] From the foregoing it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention.

SEQUENCE LISTING

```
<160> NUMBER OF SEQ ID NOS: 323

<210> SEQ ID NO 1
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: D from lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Aminohexanoic acid aplmityl conjugation
<220> FEATURE:
```

-continued

<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 1

Cys Thr His Xaa Pro Cys Ile Lys Xaa Lys Lys
1 5 10

<210> SEQ ID NO 2
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: D from lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Aminohexanoic acid aplmityl conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 2

Glu Thr Cys Xaa Pro Cys Ile Lys Xaa Lys Lys
1 5 10

<210> SEQ ID NO 3
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: D from lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Aminohexanoic acid aplmityl conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 3

Glu Thr His Cys Pro Cys Ile Lys Xaa Lys Lys
1 5 10

<210> SEQ ID NO 4
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: D from lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Aminohexanoic acid aplmityl conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 4

Glu Thr His Xaa Cys Cys Ile Lys Xaa Lys Lys
1 5 10

<210> SEQ ID NO 5
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: D from lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Aminohexanoic acid aplmityl conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 5

Glu Thr His Xaa Cys Cys Ile Lys Xaa Lys Lys
1 5 10

<210> SEQ ID NO 6
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: D from lysine

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Aminohexanoic acid aplmityl conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 6

Glu Thr His Xaa Pro Cys Cys Lys Xaa Lys Lys
1 5 10

<210> SEQ ID NO 7
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Aminohexanoic acid aplmityl conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 7

Glu Thr His Xaa Pro Cys Ile Cys Xaa Lys Lys
1 5 10

<210> SEQ ID NO 8
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Aminohexanoic acid aplmityl conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 8

Glu Thr His Xaa Pro Cys Ile Cys Xaa Lys Lys
1 5 10

-continued

<210> SEQ ID NO 9
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: ACETYLATION
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Aminohexanoic acid aplmityl conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine

<400> SEQUENCE: 9

Cys Glu Thr His Xaa Pro Cys Ile Lys Xaa
1 5 10

<210> SEQ ID NO 10
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid N-terminal conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Aminohexanoic acid aplmityl conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine

<400> SEQUENCE: 10

Cys Glu Thr His Xaa Pro Cys Ile Lys Xaa
1 5 10

<210> SEQ ID NO 11
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is penicillamine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)

-continued

<223> OTHER INFORMATION: Aminohexanoic acid aplmityl conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine

<400> SEQUENCE: 11

Xaa Glu Thr His Xaa Pro Cys Ile Lys Xaa
1 5 10

<210> SEQ ID NO 12
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is homo-cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: ACETYLTATION
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Aminohexanoic acid aplmityl conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine

<400> SEQUENCE: 12

Xaa Glu Thr His Xaa Pro Cys Ile Lys Xaa
1 5 10

<210> SEQ ID NO 13
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is homo-cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Aminohexanoic acid aplmityl conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine

<400> SEQUENCE: 13

Xaa Glu Thr His Xaa Pro Cys Ile Lys Xaa
1 5 10

-continued

<210> SEQ ID NO 14
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is homo-cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: ACETYLATION
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is homo-cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Aminohexanoic acid aplmityl conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine

<400> SEQUENCE: 14

Xaa Glu Thr His Xaa Pro Xaa Ile Lys Xaa
1 5 10

<210> SEQ ID NO 15
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is homo-cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is homo-cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Aminohexanoic acid aplmityl conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine

<400> SEQUENCE: 15

Xaa Glu Thr His Xaa Pro Xaa Ile Lys Xaa
1 5 10

-continued

<210> SEQ ID NO 16
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is homo-cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is homo-cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Aminohexanoic acid aplmityl conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine

<400> SEQUENCE: 16

Xaa Val Glu Thr His Xaa Pro Xaa Ile Lys Xaa
1 5 10

<210> SEQ ID NO 17
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is homo-cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Aminohexanoic acid aplmityl conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine

<400> SEQUENCE: 17

Cys Val Glu Thr His Xaa Pro Xaa Ile Lys Xaa
1 5 10

<210> SEQ ID NO 18
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Xaa is penicillamine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (14)..(14)
<223> OTHER INFORMATION: 1PEG2_1PEG2_IsoGlu_C18_Diacid conjugation

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (16)..(16)
<223> OTHER INFORMATION: Xaa is penicillamine

<400> SEQUENCE: 18

Glu Thr His Phe Pro Xaa Ile Lys Phe Glu Pro Arg Ser Lys Gly Xaa
1 5 10 15

Lys

<210> SEQ ID NO 19
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Xaa is penicillamine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (14)..(14)
<223> OTHER INFORMATION: 1PEG2_1PEG2_IsoGlu_C18_Diacid conjugation

<400> SEQUENCE: 19

Glu Thr His Phe Pro Xaa Ile Lys Phe Glu Pro Arg Ser Lys Gly Cys
1 5 10 15

Lys

<210> SEQ ID NO 20
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Ahx_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 20

Glu Thr His Cys Pro Cys Ile Lys Xaa Lys
1 5 10

<210> SEQ ID NO 21
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: isovaleric acid conjugation
<220> FEATURE:

-continued

<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Ahx_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 21

Glu Thr His Xaa Cys Cys Ile Lys Xaa Lys
1 5 10

<210> SEQ ID NO 22
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Ahx_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 22

Glu Thr His Xaa Pro Cys Cys Lys Xaa Lys
1 5 10

<210> SEQ ID NO 23
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Ahx_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)

-continued

<223> OTHER INFORMATION: Xaa is beta-homophenylalanine

<400> SEQUENCE: 23

Glu Thr His Xaa Pro Cys Ile Lys Xaa Cys
1 5 10

<210> SEQ ID NO 24

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized hepcidin analogues

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (1)..(1)

<223> OTHER INFORMATION: isovaleric acid conjugation

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (8)..(8)

<223> OTHER INFORMATION: D form lysine

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (9)..(9)

<223> OTHER INFORMATION: Xaa is beta-homophenylalanine

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (10)..(10)

<223> OTHER INFORMATION: 1PEG2_1PEG2_Ahx_C18_Diacid conjugation

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (11)..(11)

<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 24

Glu Thr His Cys Pro Cys Ile Lys Xaa Lys Lys
1 5 10

<210> SEQ ID NO 25

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized hepcidin analogues

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (1)..(1)

<223> OTHER INFORMATION: isovaleric acid conjugation

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (4)..(4)

<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (8)..(8)

<223> OTHER INFORMATION: D form lysine

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (9)..(9)

<223> OTHER INFORMATION: Xaa is beta-homophenylalanine

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (10)..(10)

<223> OTHER INFORMATION: 1PEG2_1PEG2_Ahx_C18_Diacid conjugation

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (11)..(11)

<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 25

Glu Thr His Xaa Cys Cys Ile Lys Xaa Lys Lys
1 5 10

-continued

<210> SEQ ID NO 26
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: D form lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Ahx_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 26

Glu Thr His Xaa Pro Cys Cys Lys Xaa Lys Lys
1 5 10

<210> SEQ ID NO 27
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Ahx_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 27

Glu Thr His Xaa Pro Cys Ile Cys Xaa Lys Lys
1 5 10

<210> SEQ ID NO 28
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: D form lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: 1PEG2_1PEG2_IsoGlu_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 28

Glu Thr His Xaa Pro Cys Cys Lys Xaa Lys Lys
1 5 10

<210> SEQ ID NO 29
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_IsoGlu_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 29

Glu Thr His Cys Pro Cys Ile Lys Xaa Lys
1 5 10

<210> SEQ ID NO 30
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_IsoGlu_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 30

Glu Thr His Xaa Cys Cys Ile Lys Xaa Lys
1 5 10

<210> SEQ ID NO 31
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_IsoGlu_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 31

Glu Thr His Xaa Pro Cys Cys Lys Xaa Lys
1 5 10

<210> SEQ ID NO 32
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_IsoGlu_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine

<400> SEQUENCE: 32

-continued

Glu Thr His Xaa Pro Cys Ile Lys Xaa Cys
1 5 10

<210> SEQ ID NO 33
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: D form lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: 1PEG2_1PEG2_IsoGlu_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 33

Glu Thr His Cys Pro Cys Ile Lys Xaa Lys Lys
1 5 10

<210> SEQ ID NO 34
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: D form lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: 1PEG2_1PEG2_IsoGlu_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 34

Glu Thr His Xaa Cys Cys Ile Lys Xaa Lys Lys
1 5 10

<210> SEQ ID NO 35

-continued

<400> SEQUENCE: 35

000

<210> SEQ ID NO 36
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: 1PEG2_1PEG2_IsoGlu_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 36

Glu Thr His Xaa Pro Cys Ile Cys Xaa Lys Lys
1 5 10

<210> SEQ ID NO 37

<400> SEQUENCE: 37

000

<210> SEQ ID NO 38

<400> SEQUENCE: 38

000

<210> SEQ ID NO 39
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:

-continued

<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 39

Glu Thr His Xaa Pro Cys Cys Lys Xaa Lys
1 5 10

<210> SEQ ID NO 40
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 40

Xaa Thr His Xaa Pro Cys Cys Lys Xaa Lys
1 5 10

<210> SEQ ID NO 41
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 41

-continued

Glu	Thr	His	Xaa	Pro	Cys	Cys	Lys	Xaa	Lys
1				5					10

<210> SEQ ID NO 42
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 42

Xaa	Thr	His	Xaa	Pro	Cys	Cys	Lys	Xaa	Lys
1				5					10

<210> SEQ ID NO 43
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 43

Cys	Thr	His	Xaa	Pro	Ala	Ile	Cys	Xaa	Lys
1				5					10

<210> SEQ ID NO 44
<211> LENGTH: 10

-continued

<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is N-methyl-cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 44

Glu Thr His Xaa Pro Cys Xaa Lys Xaa Lys
1 5 10

<210> SEQ ID NO 45
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: D form lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 45

Glu Cys His Xaa Cys Ala Ile Lys Xaa Lys Lys
1 5 10

<210> SEQ ID NO 46
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta,beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: D form cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: D form lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 46

Glu Cys His Xaa Cys Ala Ile Lys Xaa Lys Lys
1 5 10

<210> SEQ ID NO 47

<400> SEQUENCE: 47

000

<210> SEQ ID NO 48

<400> SEQUENCE: 48

000

<210> SEQ ID NO 49

<400> SEQUENCE: 49

000

<210> SEQ ID NO 50

<400> SEQUENCE: 50

000

<210> SEQ ID NO 51

<400> SEQUENCE: 51

000

<210> SEQ ID NO 52

<400> SEQUENCE: 52

-continued

000

<210> SEQ ID NO 53

<400> SEQUENCE: 53

000

<210> SEQ ID NO 54

<400> SEQUENCE: 54

000

<210> SEQ ID NO 55

<400> SEQUENCE: 55

000

<210> SEQ ID NO 56

<400> SEQUENCE: 56

000

<210> SEQ ID NO 57

<400> SEQUENCE: 57

000

<210> SEQ ID NO 58

<400> SEQUENCE: 58

000

<210> SEQ ID NO 59

<400> SEQUENCE: 59

000

<210> SEQ ID NO 60

<400> SEQUENCE: 60

000

<210> SEQ ID NO 61

<400> SEQUENCE: 61

000

<210> SEQ ID NO 62

<400> SEQUENCE: 62

000

<210> SEQ ID NO 63

<400> SEQUENCE: 63

-continued

000

<210> SEQ ID NO 64

<400> SEQUENCE: 64

000

<210> SEQ ID NO 65

<400> SEQUENCE: 65

000

<210> SEQ ID NO 66

<400> SEQUENCE: 66

000

<210> SEQ ID NO 67

<400> SEQUENCE: 67

000

<210> SEQ ID NO 68

<400> SEQUENCE: 68

000

<210> SEQ ID NO 69

<400> SEQUENCE: 69

000

<210> SEQ ID NO 70

<400> SEQUENCE: 70

000

<210> SEQ ID NO 71

<400> SEQUENCE: 71

000

<210> SEQ ID NO 72

<400> SEQUENCE: 72

000

<210> SEQ ID NO 73

<400> SEQUENCE: 73

000

<210> SEQ ID NO 74

-continued

<400> SEQUENCE: 74

000

<210> SEQ ID NO 75

<400> SEQUENCE: 75

000

<210> SEQ ID NO 76

<400> SEQUENCE: 76

000

<210> SEQ ID NO 77

<400> SEQUENCE: 77

000

<210> SEQ ID NO 78

<400> SEQUENCE: 78

000

<210> SEQ ID NO 79

<400> SEQUENCE: 79

000

<210> SEQ ID NO 80

<400> SEQUENCE: 80

000

<210> SEQ ID NO 81

<400> SEQUENCE: 81

000

<210> SEQ ID NO 82

<400> SEQUENCE: 82

000

<210> SEQ ID NO 83

<400> SEQUENCE: 83

000

<210> SEQ ID NO 84

<400> SEQUENCE: 84

000

<210> SEQ ID NO 85

-continued

<400> SEQUENCE: 85

000

<210> SEQ ID NO 86

<400> SEQUENCE: 86

000

<210> SEQ ID NO 87

<400> SEQUENCE: 87

000

<210> SEQ ID NO 88

<400> SEQUENCE: 88

000

<210> SEQ ID NO 89

<400> SEQUENCE: 89

000

<210> SEQ ID NO 90

<400> SEQUENCE: 90

000

<210> SEQ ID NO 91

<400> SEQUENCE: 91

000

<210> SEQ ID NO 92

<400> SEQUENCE: 92

000

<210> SEQ ID NO 93

<400> SEQUENCE: 93

000

<210> SEQ ID NO 94

<400> SEQUENCE: 94

000

<210> SEQ ID NO 95

<400> SEQUENCE: 95

000

-continued

<210> SEQ ID NO 96

<400> SEQUENCE: 96

000

<210> SEQ ID NO 97

<400> SEQUENCE: 97

000

<210> SEQ ID NO 98

<400> SEQUENCE: 98

000

<210> SEQ ID NO 99

<400> SEQUENCE: 99

000

<210> SEQ ID NO 100

<400> SEQUENCE: 100

000

<210> SEQ ID NO 101

<211> LENGTH: 13

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized hepcidin analogues

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (1)..(1)

<223> OTHER INFORMATION: Isovaleric acid conjugation

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (9)..(9)

<223> OTHER INFORMATION: IsoGlu_Palm conjugation

<400> SEQUENCE: 101

Asp Thr His Phe Pro Cys Cys Ile Lys Phe Glu Pro Arg

1

5

10

<210> SEQ ID NO 102

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized hepcidin analogues

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (1)..(1)

<223> OTHER INFORMATION: Xaa is homo-cysteine

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (1)..(1)

<223> OTHER INFORMATION: Isovaleric acid conjugation

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (4)..(4)

<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine

<220> FEATURE:

<221> NAME/KEY: MOD_RES

-continued

<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 102

Xaa Thr His Xaa Pro Ala Ile Cys Xaa Lys Lys
1 5 10

<210> SEQ ID NO 103
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is D form penicillamine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 103

Xaa Thr His Xaa Pro Ala Ile Cys Xaa Lys Lys
1 5 10

<210> SEQ ID NO 104
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is D form penicillamine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 104

Cys Thr His Xaa Pro Ala Ile Xaa Xaa Lys Lys
1 5 10

<210> SEQ ID NO 105
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: D form cystein
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: D form cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 105

Cys Thr His Xaa Pro Ala Ile Cys Xaa Lys Lys
1 5 10

<210> SEQ ID NO 106
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: D form cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES

-continued

<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 106

Cys Thr His Xaa Pro Ala Ile Cys Xaa Lys Lys
1 5 10

<210> SEQ ID NO 107
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: D form cystein
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 107

Cys Thr His Xaa Pro Ala Ile Cys Xaa Lys Lys
1 5 10

<210> SEQ ID NO 108
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is homo-cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 108

Cys Thr His Xaa Pro Ala Ile Xaa Xaa Lys Lys
1 5 10

<210> SEQ ID NO 109
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is homo-cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is penicillamine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 109

Xaa Thr His Xaa Pro Ala Ile Xaa Xaa Lys Lys
1 5 10

<210> SEQ ID NO 110
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is homo-cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES

-continued

<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 110

Xaa Thr His Xaa Pro Ala Ile Cys Xaa Lys Lys
1 5 10

<210> SEQ ID NO 111
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is homo-cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is homo-cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 111

Xaa Thr His Xaa Pro Ala Ile Xaa Xaa Lys Lys
1 5 10

<210> SEQ ID NO 112
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is N-Butyl Phenylalanine

<400> SEQUENCE: 112

Glu Cys His Xaa Cys Lys Xaa
1 5

<210> SEQ ID NO 113
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Xaa is trimethyl-L-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is N-Butyl Phenylalanine

<400> SEQUENCE: 113

Glu Cys His Xaa Cys Xaa Xaa
1 5

<210> SEQ ID NO 114
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is N-Butyl Phenylalanine

<400> SEQUENCE: 114

Glu Cys His Xaa Cys Lys Xaa
1 5

<210> SEQ ID NO 115
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:

-continued

<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: DMG_N_2ae_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is N-Butyl Phenylalanine

<400> SEQUENCE: 115

Glu Cys His Xaa Cys Lys Xaa
1 5

<210> SEQ ID NO 116
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Dap_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is N-Butyl Phenylalanine

<400> SEQUENCE: 116

Glu Cys His Xaa Cys Lys Xaa
1 5

<210> SEQ ID NO 117
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Ahx_Dap_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is N-Butyl Phenylalanine

<400> SEQUENCE: 117

-continued

Glu Cys His Xaa Cys Lys Xaa
1 5

<210> SEQ ID NO 118
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is N-Butyl Phenylalanine

<400> SEQUENCE: 118

Glu Cys His Xaa Cys Lys Xaa
1 5

<210> SEQ ID NO 119
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is N-methyl cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: 1PEG2_1PEG2_DMG_N_2ae_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is N-Butyl Phenylalanine

<400> SEQUENCE: 119

Glu Cys His Xaa Xaa Lys Xaa
1 5

<210> SEQ ID NO 120
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is homo-cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: 1PEG2_1PEG2_DMG_N_2ae_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is N-Butyl Phenylalanine

<400> SEQUENCE: 120

Glu Cys His Xaa Xaa Lys Xaa
1 5

<210> SEQ ID NO 121
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: 1PEG2_1PEG2_DMG_N_2ae_Palm conjugation

<400> SEQUENCE: 121

Glu Cys His Xaa Cys Lys
1 5

<210> SEQ ID NO 122
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: 1PEG2_1PEG2_DMG_N_2ae_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is N-Butyl Phenylalanine

<400> SEQUENCE: 122

Glu Cys His Xaa Cys Lys Xaa
1 5

-continued

<210> SEQ ID NO 123
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: DMG_N_2ae_Palm conjugation

<400> SEQUENCE: 123

Xaa Cys His Xaa Cys Lys
1 5

<210> SEQ ID NO 124
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: DMG_N_2ae_Palm conjugation

<400> SEQUENCE: 124

Glu Cys His Xaa Cys Lys
1 5

<210> SEQ ID NO 125
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is (2S,4S)-4-mercaptopyrrolidine-2-carboxylic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES

-continued

<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is N-Butyl Phenylalanine

<400> SEQUENCE: 125

Glu Cys His Xaa Xaa Lys Xaa
1 5

<210> SEQ ID NO 126
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is (2S,4R)-4-mercaptopyrrolidine-2-carboxylic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is N-Butyl Phenylalanine

<400> SEQUENCE: 126

Glu Cys His Xaa Xaa Lys Xaa
1 5

<210> SEQ ID NO 127
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is (2S,4S)-4-mercaptopyrrolidine-2-carboxylic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is (2S,4R)-4-mercaptopyrrolidine-2-carboxylic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_Palm conjugation
<220> FEATURE:

-continued

<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is N-Butyl Phenylalanine

<400> SEQUENCE: 127

Glu Xaa His Xaa Xaa Lys Xaa
1 5

<210> SEQ ID NO 128
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: D form cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine

<400> SEQUENCE: 128

Glu Cys His Xaa Cys Lys Xaa
1 5

<210> SEQ ID NO 129
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine

<400> SEQUENCE: 129

Glu Cys His Xaa Cys Lys Xaa
1 5

<210> SEQ ID NO 130
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: C terminal Hexadecane_Amine conjugation

<400> SEQUENCE: 130

Glu Cys His Xaa Cys
1 5

<210> SEQ ID NO 131
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Xaa is 12-aminododecanoic acid

<400> SEQUENCE: 131

Glu Cys His Xaa Cys Xaa
1 5

<210> SEQ ID NO 132
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: C terminal Dodecyl_Amine conjugation

<400> SEQUENCE: 132

Glu Cys His Xaa Cys
1 5

<210> SEQ ID NO 133
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Ahx_Palm conjugation

<400> SEQUENCE: 133

Glu Cys His Xaa Cys Lys
1 5

<210> SEQ ID NO 134
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: D form lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 134

Glu Cys His Xaa Pro Cys Ile Lys Xaa Lys Lys
1 5 10

<210> SEQ ID NO 135
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: 2-(3-methoxy-3-oxopropyl)-6-methylheptanoic
acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: D form lysine

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 135

Thr His Xaa Pro Cys Cys Lys Xaa Lys
1 5

<210> SEQ ID NO 136
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: 2-(3-methoxy-3-oxopropyl)-6-methylheptanoic
acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is trimethyl-L-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 136

Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5

<210> SEQ ID NO 137
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: 2-(3-methoxy-3-oxopropyl)-6-methylheptanoic
acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 137

Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5

<210> SEQ ID NO 138
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: 2-(3-methoxy-3-oxopropyl)-6-methylheptanoic
acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 138

Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5

<210> SEQ ID NO 139
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is trimethyl-L-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

-continued

<400> SEQUENCE: 139

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 140
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is trimethyl-L-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is D form beta, beta-diphenylalanine

<400> SEQUENCE: 140

Glu Thr His Xaa Pro Cys Cys Xaa Xaa
1 5

<210> SEQ ID NO 141
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is trimethyl-L-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine

<400> SEQUENCE: 141

Glu Thr His Xaa Pro Cys Cys Xaa Xaa
1 5

<210> SEQ ID NO 142
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES

-continued

<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 142

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 143
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: DMG_N_2ae_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 143

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 144
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa in N-methyl-threonine

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: DMG_N_2ae_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine

<400> SEQUENCE: 144

Glu Xaa His Xaa Pro Cys Cys Xaa Xaa
1 5

<210> SEQ ID NO 145
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is alpha-methylproline
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 145

Xaa Xaa His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 146
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES

-continued

<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is O-methyl-L-threonine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: DMG_N_2ae_C18_Diacid_OMe conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine

<400> SEQUENCE: 146

Xaa Xaa His Xaa Pro Cys Cys Xaa Xaa
1 5

<210> SEQ ID NO 147
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is O-methyl-L-threonine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 147

Xaa Xaa His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

-continued

<210> SEQ ID NO 148
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is N-methyl-phenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 148

Xaa Ser His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 149
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is N-methyl-threonine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: DMG_N_2ae_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)

-continued

<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine

<400> SEQUENCE: 149

Xaa Xaa His Xaa Pro Cys Cys Xaa Xaa Xaa
1 5 10

<210> SEQ ID NO 150
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is (2S,3R)-3-hydroxypyrrolidine-2-carboxylic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: DMG_N_2ae_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine

<400> SEQUENCE: 150

Glu Xaa His Xaa Pro Cys Cys Xaa Xaa
1 5

<210> SEQ ID NO 151
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is (2S,3R)-3-hydroxypyrrolidine-2-carboxylic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)

-continued

<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: DMG_N_2ae_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysing

<400> SEQUENCE: 151

Xaa Xaa His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 152
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is (2S,3R)-3-hydroxypyrrolidine-2-carboxylic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: DMG_N_2ae_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 152

Glu Xaa His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 153
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is N-methyl-threonine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine

<400> SEQUENCE: 153

Xaa Xaa His Xaa Pro Cys Cys Xaa Xaa Xaa
1 5 10

<210> SEQ ID NO 154
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is (2S,3R)-3-hydroxypyrrolidine-2-carboxylic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine

<400> SEQUENCE: 154

Glu Xaa His Xaa Pro Cys Cys Xaa Xaa
1 5

<210> SEQ ID NO 155
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:

-continued

<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is (2S,3R)-3-hydroxypyrrolidine-2-carboxylic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysing

<400> SEQUENCE: 155

Xaa Xaa His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 156
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is (2S,3R)-3-hydroxypyrrolidine-2-carboxylic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine

<400> SEQUENCE: 156

-continued

Glu Xaa His Xaa Pro Cys Cys Xaa Xaa Xaa
1 5 10

<210> SEQ ID NO 157
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysing

<400> SEQUENCE: 157

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 158
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 158

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys

-continued

1	5	10
---	---	----

<210> SEQ ID NO 159
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 159

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 160
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid_18_OMe conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 160

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

-continued

<210> SEQ ID NO 161
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid_18_OMe conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: C terminal N_Ethylamine conjugation

<400> SEQUENCE: 161

Glu Thr His Xaa Pro Cys Cys Xaa Xaa
1 5

<210> SEQ ID NO 162
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is homo-cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: CH₂CH₂CH₂NH_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 162

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 163

-continued

<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: C terminal Diphenylethylamine conjugation

<400> SEQUENCE: 163

Glu Thr His Xaa Pro Cys Cys Xaa
1 5

<210> SEQ ID NO 164
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: C terminal Diphenylethylamine conjugation

<400> SEQUENCE: 164

Glu Thr His Xaa Pro Cys Cys Xaa
1 5

<210> SEQ ID NO 165
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:

-continued

<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Diaminopropionic acid

<400> SEQUENCE: 165

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Xaa
1 5 10

<210> SEQ ID NO 166
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is N-methylglutamic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 166

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 167
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)

-continued

<223> OTHER INFORMATION: Xaa is N-methylglutamic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 167

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 168
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is homo-serine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 168

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 169
<211> LENGTH: 9
<212> TYPE: PRT

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is N-methyl-threonine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: DMG_N_2ae_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine

<400> SEQUENCE: 169

Glu Xaa His Xaa Pro Cys Cys Xaa Xaa
1 5

<210> SEQ ID NO 170
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is N-methyl-threonine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: DMG_N_2ae_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 170

Glu Xaa His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 171

-continued

<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 171

Glu Pro His Xaa Pro Cys Cys Xaa Xaa Xaa
1 5 10

<210> SEQ ID NO 172
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is N-methyl-threonine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine

<400> SEQUENCE: 172

Glu Xaa His Xaa Pro Cys Cys Xaa Xaa
1 5

<210> SEQ ID NO 173
<211> LENGTH: 10
<212> TYPE: PRT

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is N-methyl-threonine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 173

Glu Xaa His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 174
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is (2S,3R)-3-hydroxypyrrolidine-2-carboxylic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 174

-continued

Glu	Xaa	His	Xaa	Pro	Cys	Cys	Xaa	Xaa	Lys
1				5					10

<210> SEQ ID NO 175
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: DMG_N_2ae_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 175

Xaa	Thr	His	Xaa	Pro	Cys	Cys	Xaa	Xaa	Lys
1				5					10

<210> SEQ ID NO 176
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: DMG_N_2ae_IsoGlu_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:

-continued

<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 176

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 177
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 177

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 178
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)

-continued

<223> OTHER INFORMATION: Ahx_DMg_N_2ae_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 178

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 179
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine

<400> SEQUENCE: 179

Glu Thr His Xaa Pro Cys Cys Xaa Xaa
1 5

<210> SEQ ID NO 180
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is 3-pyridylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:

-continued

<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 180

Glu Thr Xaa Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 181
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: Xaa is N-methyl-phenylalanine

<400> SEQUENCE: 181

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys Xaa
1 5 10

<210> SEQ ID NO 182
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)

-continued

<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa is tert-Leucine

<400> SEQUENCE: 182

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Xaa
1 5 10

<210> SEQ ID NO 183
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is Biphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 183

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 184
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is N-methyl-threonine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:

-continued

<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-phenylalanine

<400> SEQUENCE: 184

Xaa Xaa His Xaa Pro Cys Cys Xaa
1 5

<210> SEQ ID NO 185
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is N-methyl-threonine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is N-methyl-phenylalanine

<400> SEQUENCE: 185

Xaa Xaa His Xaa Pro Cys Cys Xaa Xaa
1 5

<210> SEQ ID NO 186
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is N-methyl-threonine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)

-continued

<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine

<400> SEQUENCE: 186

Xaa Xaa His Xaa Pro Cys Cys Xaa Xaa
1 5

<210> SEQ ID NO 187
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is N-methyl-threonine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 187

Glu Xaa His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 188
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:

-continued

<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 188

Glu Pro His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 189
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is N-methyl-threonine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine

<400> SEQUENCE: 189

Glu Xaa His Xaa Pro Cys Cys Xaa Xaa
1 5

<210> SEQ ID NO 190
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is N-methyl-threonine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)

-continued

<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 190

Glu Xaa His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 191
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is (2S,3R)-3-hydroxypyrrolidine-2-carboxylic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 191

Glu Xaa His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 192
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is (2S,3S)-3-hydroxypyrrolidine-2-

-continued

carboxylic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 192

Glu Xaa His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 193
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 193

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 194
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:

-continued

<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: DMG_N_2ae_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 194

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 195
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: DMG_N_2ae_IsoGlu_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 195

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 196
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)

-continued

<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 196

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 197
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 197

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 198
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:

-continued

<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 198

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 199
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 199

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 200
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)

-continued

<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 200

Xaa	Thr	His	Xaa	Pro	Cys	Cys	Lys	Xaa	Lys
1			5					10	

<210> SEQ ID NO 201
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 201

Xaa	Thr	His	Xaa	Pro	Cys	Cys	Lys	Xaa	Lys
1			5					10	

<210> SEQ ID NO 202
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:

-continued

<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: PEG11_OMe conjugation

<400> SEQUENCE: 202

Xaa Thr His Xaa Pro Cys Cys Lys Xaa Lys
1 5 10

<210> SEQ ID NO 203
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: PEG11_OMe conjugation

<400> SEQUENCE: 203

Glu Thr His Xaa Pro Cys Cys Lys Xaa Lys
1 5 10

<210> SEQ ID NO 204
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)

-continued

<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_DMGM_N2aeC18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 204

Glu Thr His Xaa Pro Cys Cys Lys Xaa Lys
1 5 10

<210> SEQ ID NO 205
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMGM_N2aeC18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: PEG11_OME conjugation

<400> SEQUENCE: 205

Xaa Thr His Xaa Pro Cys Cys Lys Xaa Lys
1 5 10

<210> SEQ ID NO 206
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:

-continued

<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 206

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 207
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 207

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 208
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)

-continued

<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: D form lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 208

Xaa Thr His Xaa Pro Cys Cys Lys Xaa Lys
1 5 10

<210> SEQ ID NO 209
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: DMG_N_2ae_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 209

Xaa Thr His Xaa Pro Cys Cys Lys Xaa Lys
1 5 10

<210> SEQ ID NO 210
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:

-continued

<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa is tert-leucine

<400> SEQUENCE: 210

Xaa Thr His Xaa Pro Cys Cys Lys Xaa Xaa
1 5 10

<210> SEQ ID NO 211
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 211

Xaa Thr His Xaa Pro Cys Cys Lys Xaa Lys
1 5 10

<210> SEQ ID NO 212
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)

-continued

<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: D form lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa is tert-leucine

<400> SEQUENCE: 212

Xaa Thr His Xaa Pro Cys Cys Lys Xaa Xaa
1 5 10

<210> SEQ ID NO 213
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: D form lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 213

Xaa Thr His Xaa Pro Cys Cys Lys Xaa Lys
1 5 10

<210> SEQ ID NO 214
<211> LENGTH: 10
<212> TYPE: PRT

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: PEG11_OMe conjugation

<400> SEQUENCE: 214

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 215
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is Biphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: PEG11_OMe conjugation

<400> SEQUENCE: 215

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 216
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is Biphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 216

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 217
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is (S)-morpholine-3-carboxylic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES

-continued

<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 217

Glu Thr His Xaa Xaa Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 218
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 218

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 219
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Carnitine_Alkyl conjugation

<400> SEQUENCE: 219

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 220
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: IsoGlu_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: PEG11_OMe conjugation

<400> SEQUENCE: 220

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 221
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES

-continued

```

<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: IsoGlu_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

```

```

<400> SEQUENCE: 221

```

```

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1           5           10

```

```

<210> SEQ ID NO 222
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: IsoGlu_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: PEG11_OMe conjugation

```

```

<400> SEQUENCE: 222

```

```

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1           5           10

```

```

<210> SEQ ID NO 223
<211> LENGTH: 10

```

-continued

<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: IsoGlu_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 223

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 224
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is L-1,2,3,4-Tetrahydroisoquinoline-3-carboxylic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 224

Glu Thr His Xaa Xaa Cys Cys Xaa Xaa Lys
1 5 10

-continued

<210> SEQ ID NO 225
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is N-Methyl-Phenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa is D form N-Methyl-Lysine

<400> SEQUENCE: 225

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Xaa
1 5 10

<210> SEQ ID NO 226
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is 3-pyridylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is Alpha-methylphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 226

Leu Thr Xaa Xaa Pro Cys Cys Xaa Xaa Lys

-continued

1	5	10
---	---	----

```

<210> SEQ ID NO 227
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is 3-pyridylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is beta-homoproline
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

```

<400> SEQUENCE: 227

```

Leu Thr Xaa Xaa Xaa Cys Cys Xaa Xaa Lys
1           5           10

```

```

<210> SEQ ID NO 228
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is beta-homoproline
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES

```

-continued

<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 228

Leu Thr His Xaa Xaa Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 229
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 229

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 230
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_DMGM_N2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 230

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 231
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_DMGM_N2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 231

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 232
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES

-continued

<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 232

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 233
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 233

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 234
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is homo-serine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 234

Glu Xaa His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 235
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: PEG11_OMe conjugation

<400> SEQUENCE: 235

Xaa Thr His Xaa Pro Cys Cys Lys Xaa Lys
1 5 10

<210> SEQ ID NO 236
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: PEG11_OMe conjugation

<400> SEQUENCE: 236

Glu Thr His Xaa Pro Cys Cys Lys Xaa Lys
1 5 10

<210> SEQ ID NO 237
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-Methyl-Lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa is (S)-2-amino-3-morpholinopropanoic acid

<400> SEQUENCE: 237

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Xaa
1 5 10

<210> SEQ ID NO 238
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is 3-pyridylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is Phe(4-COOH)
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 238

Leu Thr Xaa Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 239
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is 3-pyridylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is Nipicotic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 239

-continued

Leu Thr Xaa Xaa Xaa Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 240
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is Nipecotic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 240

Leu Thr His Xaa Xaa Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 241
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is 3-pyridylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:

-continued

<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 241

Leu Thr Xaa Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 242
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is (S)-2-aminoadipic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is 3-pyridylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 242

Xaa Thr Xaa Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 243
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is 3-pyridylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)

-continued

<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 243

Glu Thr Xaa Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 244
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is (S)-2-aminoadipic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 244

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 245
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is (S)-2-aminoadipic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:

-continued

<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is Xaa is Nipecotic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 245

Xaa Thr His Xaa Xaa Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 246
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is (S)-2-aminoadipic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is 3-pyridylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 246

Xaa Thr Xaa Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 247

-continued

<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: Xaa is D form N-Methyl-Serine

<400> SEQUENCE: 247

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys Xaa
1 5 10

<210> SEQ ID NO 248
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: Xaa is D form N-Methyl-Glutamine

<400> SEQUENCE: 248

-continued

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys Xaa
1 5 10

<210> SEQ ID NO 249
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: Xaa is D form N-Methyl-Leucine

<400> SEQUENCE: 249

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys Xaa
1 5 10

<210> SEQ ID NO 250
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: Xaa is D form N-Methyl-Phenylalanine

<400> SEQUENCE: 250

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys Xaa
1 5 10

<210> SEQ ID NO 251
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: Xaa is N-Methyl-Glutamine

<400> SEQUENCE: 251

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys Xaa
1 5 10

<210> SEQ ID NO 252
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES

-continued

<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: Xaa is N-Methyl-Leucine

<400> SEQUENCE: 252

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys Xaa
1 5 10

<210> SEQ ID NO 253
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is (2S,3S)-3-hydroxypyrrolidine-2-carboxylic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 253

Glu Xaa His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 254
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)

-continued

<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: PEG11_OME conjugation

<400> SEQUENCE: 254

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 255
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: PEG11_OME conjugation

<400> SEQUENCE: 255

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 256
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:

-continued

<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine

<400> SEQUENCE: 256

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa
1 5

<210> SEQ ID NO 257
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: Xaa is D form N-Methyl-Tyrosine

<400> SEQUENCE: 257

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys Xaa
1 5 10

<210> SEQ ID NO 258
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: Xaa is 1-Napthylalanine

<400> SEQUENCE: 258

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys Xaa
1 5 10

<210> SEQ ID NO 259
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: Xaa is N-Methyl-Tyrosine

<400> SEQUENCE: 259

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys Xaa
1 5 10

-continued

<210> SEQ ID NO 260
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: Xaa is N-Methyl-Serine

<400> SEQUENCE: 260

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys Xaa
1 5 10

<210> SEQ ID NO 261
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: Xaa is Napthylalanine

-continued

<400> SEQUENCE: 261

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys Xaa
1 5 10

<210> SEQ ID NO 262
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: Xaa is N-Methyl-Isoleucine

<400> SEQUENCE: 262

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys Xaa
1 5 10

<210> SEQ ID NO 263
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES

-continued

<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: Xaa is N-Methyl-Phenylalanine

<400> SEQUENCE: 263

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys Xaa
1 5 10

<210> SEQ ID NO 264
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa is Diaminobutyric acid

<400> SEQUENCE: 264

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Xaa
1 5 10

<210> SEQ ID NO 265
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa is (S)-2-amino-5-morpholinopentanoic acid

<400> SEQUENCE: 265

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Xaa
1 5 10

<210> SEQ ID NO 266
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is N-methyl-cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is N-methyl-phenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 266

Glu Thr His Xaa Pro Cys Xaa Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 267
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES

-continued

<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa is D form N-methyl-Lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Acetyl conjugation

<400> SEQUENCE: 267

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Xaa
1 5 10

<210> SEQ ID NO 268
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(7)
<223> OTHER INFORMATION: Xaa is homocysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 268

Glu Thr His Xaa Pro Xaa Xaa Lys Xaa Lys
1 5 10

<210> SEQ ID NO 269
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(7)
<223> OTHER INFORMATION: Xaa is homocysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 269

Glu Thr His Xaa Pro Xaa Xaa Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 270
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is homocysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 270

Glu Thr His Xaa Pro Cys Xaa Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 271
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES

-continued

<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Xaa is penicillamine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 271

Glu Thr His Xaa Pro Xaa Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 272
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is penicillamine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 272

Glu Thr His Xaa Pro Cys Xaa Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 273
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(7)
<223> OTHER INFORMATION: Xaa is penicillamine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 273

Glu Thr His Xaa Pro Xaa Xaa Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 274
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(7)
<223> OTHER INFORMATION: Xaa is penicillamine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 274

Glu Thr His Xaa Pro Xaa Xaa Lys Xaa Lys
1 5 10

<210> SEQ ID NO 275
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES

-continued

<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa is N-methyl-Lysine

<400> SEQUENCE: 275

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Xaa
1 5 10

<210> SEQ ID NO 276
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa is D form N-methyl-Lysine

<400> SEQUENCE: 276

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Xaa
1 5 10

<210> SEQ ID NO 277
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is N-methyl-cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is N-Methyl-Phenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa is N-methyl-Lysine

<400> SEQUENCE: 277

Glu Thr His Xaa Pro Cys Xaa Xaa Xaa Xaa
1 5 10

<210> SEQ ID NO 278
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is N-methyl-cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 278

Glu Thr His Xaa Pro Cys Xaa Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 279
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is 1-Methyl-histidine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is N-methyl-phenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 279

Glu Thr Xaa Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 280
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is 1-Methyl-histidine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is Nipicotic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

-continued

<400> SEQUENCE: 280

Glu Thr Xaa Xaa Xaa Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 281
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is N-methyl-cysteine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is N-methyl-phenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 281

Glu Thr His Xaa Pro Cys Xaa Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 282
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_DMg_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)

-continued

<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 282

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 283
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 283

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 284
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is (S)-2-amino-3-(6-methoxy-1H-indol-3-yl)propanoic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 284

Glu Thr Xaa Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 285
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is 1-Methyl-histidine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 285

Glu Thr Xaa Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 286
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is (4-Trifluoromethyl)-L-Phenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES

-continued

<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 286

Glu Thr Xaa Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 287
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is (S)-2-amino-3-(6-methoxy-1H-indol-3-yl)propanoic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 287

Glu Thr Xaa Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 288
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is (S)-2-amino-3-(2H-tetrazol-5-yl)propanoic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES

-continued

<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 288

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 289
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is Diaminopropionic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 289

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 290
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa is tert-leucine

<400> SEQUENCE: 290

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Xaa
1 5 10

<210> SEQ ID NO 291
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is N-Methyl-Phenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 291

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 292
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES

-continued

<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa is N-Methyl-Lysine

<400> SEQUENCE: 292

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Xaa
1 5 10

<210> SEQ ID NO 293
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form lysine

<400> SEQUENCE: 293

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 294
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation

<400> SEQUENCE: 294

Glu Thr His Xaa Pro Cys Cys Xaa
1 5

<210> SEQ ID NO 295
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine

<400> SEQUENCE: 295

Glu Thr His Xaa Pro Cys Cys Xaa Xaa
1 5

<210> SEQ ID NO 296
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is 3-Quinolinyllalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES

-continued

<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 296

Glu Thr Xaa Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 297
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is Biphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 297

Glu Thr Xaa Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 298
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: 4_Fluorophenylacetic_acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 298

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 299
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Nicotinic_acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 299

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 300
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: isonipecotic acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES

-continued

<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 300

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 301
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric Acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is alpha-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 301

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 302
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric Acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is alpha-methyl-lysine

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 302

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 303
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric Acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: PEG12_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 303

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 304
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric Acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES

-continued

<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: PEG12_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 304

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 305
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric Acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 305

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 306
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric Acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: 1PEG2_1PEG2_Dap_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 306

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 307
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric Acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 307

Xaa Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 308
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric Acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES

-continued

<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is N-methyl-lysine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 308

Glu Thr His Xaa Pro Cys Cys Xaa Xaa Lys
1 5 10

<210> SEQ ID NO 309
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric Acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: DMG_N_2ae_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 309

Xaa Thr His Xaa Pro Cys Cys Lys Xaa Lys
1 5 10

<210> SEQ ID NO 310
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric Acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine

-continued

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: DMG_N_2ae_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 310

Glu Thr His Xaa Pro Cys Cys Lys Xaa Lys
1 5 10

<210> SEQ ID NO 311
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric Acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 311

Xaa Thr His Xaa Pro Cys Cys Lys Xaa Lys
1 5 10

<210> SEQ ID NO 312
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric Acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES

-continued

<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 312

Glu Thr His Xaa Pro Cys Cys Lys Xaa Lys
1 5 10

<210> SEQ ID NO 313
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric Acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: DMG_N_2ae_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 313

Xaa His Xaa Pro Cys Cys Lys Xaa Lys
1 5

<210> SEQ ID NO 314
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric Acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: DMG_N_2ae_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: D form Lysine

-continued

<400> SEQUENCE: 314

Glu His Xaa Pro Cys Cys Lys Xaa Lys
1 5

<210> SEQ ID NO 315
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric Acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 315

Xaa Thr His Xaa Pro Cys Cys Lys Xaa Lys
1 5 10

<210> SEQ ID NO 316
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric Acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ahx_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 316

Glu Thr His Xaa Pro Cys Cys Lys Xaa Lys
1 5 10

-continued

<210> SEQ ID NO 317
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is L-Glutamic acid gamma-methyl ester
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric Acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: DMG_N_2ae_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 317

Xaa Thr His Xaa Pro Cys Cys Lys Xaa Lys
1 5 10

<210> SEQ ID NO 318
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized hepcidin analogues
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric Acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is beta, beta-diphenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: DMG_N_2ae_DMG_N_2ae_C18_Diacid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa is beta-homophenylalanine
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: D form Lysine

<400> SEQUENCE: 318

Glu Thr His Xaa Pro Cys Cys Lys Xaa Lys
1 5 10

<210> SEQ ID NO 319
<211> LENGTH: 25
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 319

Asp Thr His Phe Pro Ile Cys Ile Phe Cys Cys Gly Cys Cys His Arg
1 5 10 15

Ser Lys Cys Gly Met Cys Cys Lys Thr
 20 25

<210> SEQ ID NO 320

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 320

Asp Thr His Phe Pro Ile Cys Ile Phe
1 5

<210> SEQ ID NO 321

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Made in Lab - Hepcidin Analogue

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (1)..(1)

<223> OTHER INFORMATION: D form phenylalanine

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (2)..(2)

<223> OTHER INFORMATION: D form isoleucine

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (3)..(3)

<223> OTHER INFORMATION: D form cysteine

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (4)..(4)

<223> OTHER INFORMATION: D form isoleucine

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (5)..(5)

<223> OTHER INFORMATION: D form proline

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (6)..(6)

<223> OTHER INFORMATION: D form phenylalanine

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (7)..(7)

<223> OTHER INFORMATION: D form histidine

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (8)..(8)

<223> OTHER INFORMATION: D form threonine

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (9)..(9)

<223> OTHER INFORMATION: D form aspartic acid

<400> SEQUENCE: 321

Phe Ile Cys Ile Pro Phe His Thr Asp
1 5

<210> SEQ ID NO 322

<211> LENGTH: 17

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Made in Lab - Hepcidin Analogue

<220> FEATURE:

-continued

```

<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: 2Peg11'-Palm conjugation

<400> SEQUENCE: 322

Asp Thr His Phe Pro Cys Ile Lys Phe Lys Pro Arg Ser Lys Gly Cys
1           5           10          15

Lys

<210> SEQ ID NO 323
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Made in Lab - Hepcidin Analogue
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Isovaleric acid conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: 2Peg11'-Palm conjugation
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (15)..(15)
<223> OTHER INFORMATION: Xaa is Sarcosine


<400> SEQUENCE: 323

Asp Thr His Phe Pro Cys Ile Lys Phe Lys Pro Arg Ser Lys Xaa Cys
1           5           10          15

Lys

```

1. A hepcidin analogue comprising a peptide comprising an amino acid sequence of Formula (I'):



or a pharmaceutically acceptable salt or solvate thereof, wherein:

R¹ is hydrogen, C₁–C₆ alkyl, C₆–C₁₂ aryl, C₆–C₁₂ aryl-C₁–C₆ alkyl, C₁–C₂₀ alkanoyl, C₁–C₂₀ cycloalkanoyl, C₆–C₁₀ aryl-C₁–C₆ alkylene-C(O)—, 5- or 6-membered heteroaryl-C(O)— or 5- or 6-membered heterocycloalkyl-C(O)—, wherein the C₁–C₂₀ alkanoyl, C₆–C₁₀ aryl-C₁–C₆ alkylene-C(O)—, 5- or 6-membered heteroaryl-C(O)— and 5- or 6-membered heterocycloalkyl-C(O)— of R¹ are each optionally substituted with 1, 2 or 3 substituents independently selected from halo, OH, CN, C₁–C₆ alkoxy, C₁–C₆ alkyl, —COOH, —COO(C₁–C₆), C₁–C₆ haloalkyl and C₁–C₆ haloalkoxy;

R² is NH₂, substituted amino, OH, substituted hydroxy, C₁–C₂₀ alkylamino, phenyl-C₁–C₈ alkylene-NH—, wherein C₁–C₂₀ alkylamino and phenyl-C₁–C₈ alkylene-NH— are each optionally substituted with a substituent independently selected from NH₂, —COOH and phenyl;

X1 is absent, or is Asp, isoAsp, Asp(OMe), Glu, bhGlu, bGlu, Gly, N-substituted Gly, Gla, Glp, Ala, Arg, Dab,

Leu, Lys, Dap, Orn, (D)Asp, (D)Arg, Tet1, Tet2, Lys, substituted Lys, (D)Lys, substituted (D)Lys, or Cys, (D)Cys, aMeCys, hCys, Pen, (D)Pen, NMe-Glu, Aad, Dap or isoGlu;

X2 is Ala, Thr, Gly, N-substituted Gly, Ser, Cys, 4S_Mcp, 4R_Mcp, NMe-Thr, aMePro, Hyp_3R, HyP_3S, hSer or Thr_Me;

X3 is Ala, Gly, N-substituted Gly, His, substituted His, Cys, (D)Cys, aMeCys, hCys, Pen, 3Pal, 4Pal, BIP, Ala_3Quin, Trp_50H or His_1Me;

X4 is Ala, Phe, Dpa, Gly, N-substituted Gly, bhPhe, a-MePhe, NMe-Phe, D-Phe, 2Pal, or Cys, (D)Cys, aMeCys, hCys, Pen or DIP;

X5 is Ala, Pro, D-Pro, bhPro, D-bhPro, NPC, D-NPC, Gaba, 2-Pyrrolidinepropanoic acid (Ppa), 2-Pyrrolidinebutanoic acid (Pba), Glu, Lys, substituted Lys, (D)Lys, substituted (D)Lys, or Cys, (D)Cys, aMeCys, hCys, Pen, NMe-Cys, 4S_Mcp, 4R_Mcp, Morph or Tic;

X6 is Cys, (D)Cys, aMeCys, hCys, Pen, dPen, NMe-Cys, Ala, Lys, (D)Lys, substituted Lys or substituted (D)Lys;

X7 is absent, or is Ala, Gly, N-substituted Gly, Ile, Val, Leu, NLeu, Lys, substituted Lys, (D)Lys, substituted (D)Lys, Cys, (D)Cys, aMeCys, hCys, Pen, NMe-Cys, Phe or DIP;

X8 is absent or is Ala, (D)Ala, Ile, Gly, N-substituted Gly, Glu, Val, Leu, NLeu, Phe, bhPhe, Lys, substituted Lys,

(D)Lys, substituted (D)Lys, aMeLys, 123Triazole, Cys, (D)Cys, aMeCys, hCys, Pen, Lys_Me3;

X9 is absent, or is Ala, Ile, Gly, N-substituted Gly, Val, Leu, NLeu, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys, Cys, (D)Cys, aMeCys, hCys, Pen, NMe-Phe, aMePhe, Dip, dDIP, BIP, aMePhe or substituted Phe;

X10 is absent, or is Ala, Gly, N-substituted Gly, Ile, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys, Cys, (D)Cys, aMeCys, hCys, Pen, Tle, substituted (D)Lys, NMe-Lys, NMe-dLys, substituted NMe-Lys, substituted NMe-dLys or Mor_propanoic_acid;

X11 is absent, or is Ala, Pro, bhPhe, Lys, substituted Lys, (D)Lys, NMe-Phe, NMe-dSer, NMe-dGln, NMe-dLeu, NMe-dTyr, 1Nal or NMe-Ile;

each of X12-X14 is absent, or is independently an amino acid;

the peptide of formula (I') is optionally further conjugated with any amino acid;

any of the amino acids of the peptide of formula (I') is optionally replaced with the corresponding (D)-amino acid or optionally N-substituted;

(i) optionally, at least one of X1, X3-X5, X7-X10 is Cys, (D)Cys, aMeCys, hCys, Pen, (D)Pen, NMe-Cys, 4S_Mcp or 4R_Mcp, wherein the peptide is cyclized by taking the mercapto group on the side chain of one of X1, X3-X5 and X7-X10, the mercapto group on the side chain of X6 and L^x to form a —S-L^x— linkage; or the mercapto group on the side chain of X1, the mercapto group on the side chain of X8 and L^x taken together form a —S-L^x— linkage; or the mercapto group on the side chain of X2, the mercapto group on the side chain of X5 and L^x taken together form a —S-L^x— linkage, wherein each L^x is independently a bond or a linking group; and

(ii) the functional groups on the side chains of X1 and X10 are optionally taken together to form an amide linkage; or the functional group on the side chain of X1 and the functional group of R² are optionally taken together to form an amide linkage;

and wherein

Dapa is diaminopropanoic acid; Dpa or DIP is 3,3-diphenylalanine or b,b-diphenylalanine; bhPhe is b-homophenylalanine; Bip is biphenylalanine; bhPro is b-homoproline; Tic is L-1,2,3,4,-tetrahydro-isoquinoline-3-carboxylic acid; NPC is L-nipecotic acid; bhTrp is b-homoTryptophane; 1-Nal is 1-naphthylalanine; 2-Nal is 2-naphthylalanine; Orn is ornithine; Nleu is norleucine; 2Pal is 2-pyridylalanine; Ppa is 2-I-Pyrrolidinepropanoic acid, Pba is 2-I-Pyrrolidinebutanoic acid; substituted Phe is phenylalanine wherein phenyl is substituted with F, Cl, Br, I, OH, methoxy, dimethoxy, dichloro, dimethyl, difluoro, pentafluoro, allyloxy, azido, nitro, 4-carbamoyl-2,6-dimethyl, trifluoromethoxy, trifluoromethyl, phenoxy, benzyloxy, carbamoyl, t-Bu, carboxyl, CN, or guanidine;

substituted bhPhe is b-homophenylalanine wherein phenyl is substituted with F, Cl, Br, I, OH, methoxy, dimethoxy, dichloro, dimethyl, difluoro, pentafluoro, allyloxy, azido, nitro, 4-carbamoyl-2,6-dimethyl, trifluoromethoxy, trifluoromethyl, phenoxy, benzyloxy, carbamoyl, t-Bu, carboxyl, CN, or guanidine;


substituted Trp is N-methyl-L-tryptophan, a-methyltryptophan, or tryptophan substituted with F, Cl, OH, or t-Bu;

substituted bhTrp is N-methyl-L-b-homotryptophan, a-methyl-b-homotryptophan, or b-homotryptophan substituted with F, Cl, OH, or t-Bu;


substituted Lys, substituted dLys or substituted Nme-Lys or substituted Nme-dLys is Lys, dLys, Nme-Lys or Nme-dLys, wherein the F-amino group on the side chain of Lys, dLys, Nme-Lys or Nme-dLys is covalently bound to C₁-C₆ alkanoyl, a half-life extension moiety or a linking moiety covalently linked to a half-life extension moiety, wherein the linking moiety comprises one or more linker moieties covalently bound to each other;

Tet1 is (S)-(2-amino)-3-(2H-tetrazol-5-yl)propanoic acid; and Tet2 is (S)-(2-amino)-4-(1H-tetrazol-5-yl)butanoic acid;


123Triazole is



and
Dab is



2. The hepcidin analogue of claim 1 comprising a peptide comprising an amino acid sequence of Formula (X):




or a pharmaceutically acceptable salt or solvate thereof, wherein:

(i) one of X1, X3-X5, X7-X10 is Cys, (D)Cys, aMeCys, hCys, Pen, (D)Pen, Nme-Cys, 4S_Mcp or 4R_Mcp, wherein the peptide is cyclized by taking the mercapto group on the side chain of one of X1, X3-X5 and X7-X10, the mercapto group on the side chain of X6 and L^x to form a —S-L^x— linkage; or the mercapto group on the side chain of X1, the mercapto group on the side chain of X8 and L^x taken together form a —S-L^x— linkage; or the mercapto group on the side chain of X2, the mercapto group on the side chain of X5 and L^x taken together form a —S-L^x— linkage, wherein each L^x is independently a bond or a linking group; and

(ii) the functional groups on the side chains of X1 and X10 are optionally taken together to form an amide linkage; or the functional group on the side chain of X1 and the functional group of R² are optionally taken together to form an amide linkage.

3. The hepcidin analogue of claim 1, comprising a peptide according to Formula I:



or a pharmaceutically acceptable salt or solvate thereof, wherein:

R¹ is hydrogen, C₁-C₆ alkyl, C₆-C₁₂ aryl, C₆-C₁₂ aryl-C₁-C₆ alkyl, C₁-C₂₀ alkanoyl, C₁-C₂₀ cycloalkanoyl;

R² is NH₂, substituted amino, OH, or substituted hydroxy;

X0 is absent, or is Cys, (D)Cys, aMeCys, hCys, or Pen;

X1 is absent, or is Asp, isoAsp, Asp(OMe), Glu, bhGlu, bGlu, Gly, N-substituted Gly, Gla, Glp, Ala, Arg, Dab, Leu, Lys, Dap, Orn, (D)Asp, (D)Arg, Tet1, or Tet2, Lys, substituted Lys, (D)Lys, substituted (D)Lys, or Cys, (D)Cys, aMeCys, hCys, or Pen;

X2 is Ala, Thr, Gly, N-substituted Gly, or Ser;

X3 is Ala, Gly, N-substituted Gly, His, substituted His, or Cys, (D)Cys, aMeCys, hCys, or Pen;

X4 is Ala, Phe, Dpa, Gly, N-substituted Gly, bhPhe, a-MePhe, Nme-Phe, D-Phe, 2Pal, or Cys, (D)Cys, aMeCys, hCys, or Pen;

X5 is Ala, Pro, D-Pro, bhPro, D-bhPro, NPC, D-NPC, Gaba, 2-Pyrrolidinepropanoic acid (Ppa), 2-Pyrrolidinebutanoic acid (Pba), Glu, Lys, substituted Lys, (D)Lys, substituted (D)Lys, or Cys, (D)Cys, aMeCys, hCys, or Pen;

X6 is Cys, (D)Cys, aMeCys, hCys, or Pen;

X7 is absent, or is Ala, Gly, N-substituted Gly, Ile, Val, Leu, Nleu, Lys, substituted Lys, (D)Lys, substituted (D)Lys, or Cys, (D)Cys, aMeCys, hCys, or Pen;

X8 is absent or is Ala, (D)Ala, Ile, Gly, N-substituted Gly, Glu, Val, Leu, Nleu, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys, aMeLys, 123Triazole, or Cys, (D)Cys, aMeCys, hCys, or Pen;

X9 is absent, or is Ala, Ile, Gly, N-substituted Gly, Val, Leu, Nleu, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys, or Cys, (D)Cys, aMeCys, hCys, or Pen;

X10 is absent, or is Ala, Gly, N-substituted Gly, Ile, Phe, bhPhe, Lys, substituted Lys, (D)Lys, substituted (D)Lys, or Cys, (D)Cys, aMeCys, hCys, or Pen;

X11 is absent, or is Ala, Pro, bhPhe, Lys, substituted Lys, or (D)Lys; and

each of X12-X14 is absent, or is independently any amino acid;

provided that:

i) the peptide may further be conjugated at any amino acid;

ii) any of the amino acids of the peptide may be the corresponding (D)-amino acid of the amino acid or may be N-substituted; and

iii) at least one of X0, X1, X3-X5, X7-X10 is Cys, (D)Cys, aMeCys, hCys, or Pen and Cys, (D)Cys, aMeCys, hCys, or Pen form a disulfide bond with X6; and

Dapa is diaminopropanoic acid; Dpa or DIP is 3,3-diphenylalanine or b,b-diphenylalanine; bhPhe is b-homophenylalanine; Bip is biphenylalanine; bhPro is b-homoproline; Tic is L-1,2,3,4-tetrahydro-isoquinoline-3-carboxylic acid; NPC is L-nipecotic acid; bhTrp is b-homoTryptophane; 1-Nal is 1-naphthylalanine; 2-Nal is 2-naphthylalanine; Orn is ornithine; Nleu is norleucine; 2Pal is 2-pyridylalanine; Ppa is 2-I-Pyrrolidinepropanoic acid, Pba is 2-I-Pyrrolidinebutanoic acid; substituted Phe is phenylalanine wherein phenyl is substituted with F, Cl, Br, I, OH, methoxy, dime-

thoxy, dichloro, dimethyl, difluoro, pentafluoro, allyloxy, azido, nitro, 4-carbamoyl-2,6-dimethyl, trifluoromethoxy, trifluoromethyl, phenoxy, benzyloxy, carbamoyl, t-Bu, carboxyl, CN, or guanidine;


substituted bhPhe is b-homophenylalanine wherein phenyl is substituted with F, Cl, Br, I, OH, methoxy, dimethoxy, dichloro, dimethyl, difluoro, pentafluoro, allyloxy, azido, nitro, 4-carbamoyl-2,6-dimethyl, trifluoromethoxy, trifluoromethyl, phenoxy, benzyloxy, carbamoyl, t-Bu, carboxyl, CN, or guanidine;

substituted Trp is N-methyl-L-tryptophan, a-methyltryptophan, or tryptophan substituted with F, Cl, OH, or t-Bu;


substituted bhTrp is N-methyl-L-b-homotryptophan, a-methyl-b-homotryptophan, or b-homotryptophan substituted with F, Cl, OH, or t-Bu;

Tet1 is (S)-(2-amino)-3-(2H-tetrazol-5-yl)propanoic acid; and Tet2 is (S)-(2-amino)-4-(1H-tetrazol-5-yl)butanoic acid;

123Triazole is



and
Dab is



4.-12. (canceled)

13. The hepcidin analogue or pharmaceutically acceptable salt or solvate thereof according to claim 1, wherein X1 is Glu, X2 is Thr, X3 is His, X4 is Dpa, and X5 is Pro.

14.-17. (canceled)

18. The hepcidin analogue or pharmaceutically acceptable salt or solvate thereof according to claim 13, wherein X7 is Ile.


19. (canceled)

20. The hepcidin analogue or pharmaceutically acceptable salt or solvate thereof according to claim 13, wherein X9 is Phe or bhF.

21. The hepcidin analogue or pharmaceutically acceptable salt or solvate thereof according to claim 13, wherein X10 is Lys, substituted Lys, (D)Lys, or substituted (D)Lys.

22.-25. (canceled)

26. The hepcidin analogue or pharmaceutically acceptable salt or solvate thereof according to claim 1, wherein the peptide is according to Formula V:



wherein Cys and X6 are linked via a disulfide bond.

27.-33. (canceled)

34. The hepcidin analogue of claim 1, or pharmaceutically acceptable salt or solvate thereof, wherein

X6 is Cys and X7 is Cys; or
 X6 is Cys and X7 is NMe-Cys; or
 X6 is hCys and X7 is hCys; or
 X6 is Cys and X7 is hCys;
 X6 is Pen and X7 is Cys; or
 X6 is Cys and X7 is Pen; or
 X6 is Pen and X7 is Pen.

35.-38. (canceled)

39. The hepcidin analogue of claim 1, or pharmaceutically acceptable salt or solvate thereof, wherein X1 is Glu and X10 is (D)Lys.

40.-49. (canceled)

50. The hepcidin analogue or pharmaceutically acceptable salt or solvate thereof according to claim 1, wherein X8 is absent.

51.-56. (canceled)

57. The hepcidin analogue or pharmaceutically acceptable salt or solvate thereof according to claim 1, wherein X8 is Lys(L1Z) or (D)Lys(L1Z), wherein L1 is a linker, and wherein Z is a half-life extension moiety.

58. The hepcidin analogue or pharmaceutically acceptable salt or solvate thereof according to claim 1, wherein X10 is Glu.

59. (canceled)

60. The hepcidin analogue or pharmaceutically acceptable salt or solvate thereof according to claim 1, wherein X10 is absent, Lys, substituted Lys, (D)Lys, or substituted (D)Lys.

61. (canceled)

62. The hepcidin analogue or pharmaceutically acceptable salt or solvate thereof according to claim 1, wherein X10 is (D)Lys.

63.-66. (canceled)

67. The hepcidin analogue or pharmaceutically acceptable salt or solvate thereof according to claim 1, wherein X10 is conjugated Lys or (D)Lys.

68. The hepcidin analogue or pharmaceutically acceptable salt or solvate thereof according to claim 1, wherein X10 is Lys(L1Z) or (D)Lys(L1Z), wherein L1 is a linker, and wherein Z is a half-life extension moiety.

69.-99. (canceled)

100. The hepcidin analogue or pharmaceutically acceptable salt or solvate thereof according to claim 68, wherein -L1Z is:

PEG11 OMe;
 PEG12_C18 acid;
 1PEG2_1PEG2_Ahx_Palm;
 1PEG2_Ahx_Palm;
 Ado_Palm;
 Ahx_Palm;
 Ahx_PEG20K;
 PEG12_Ahx_IsoGlu_Behenic;
 PEG12_Ahx_Palm;
 PEG12 DEKHKS_Palm;
 PEG12_IsoGlu_C18 acid;
 PEG12_Ahx_C18 acid;
 PEG12_IsoGlu_Palm;
 PEG12_KKK_Palm;
 PEG12_KKKG_Palm;
 PEG12_DEKHKS_Palm;
 PEG12_Palm;
 PEG12_PEG12 Palm;
 PEG20K;
 PEG4_Ahx_Palm;
 PEG4_Palm;

PEG8 Ahx Palm; or

IsoGlu_Palm;

-1PEG2_1PEG2_Dap_C18_Diacid;

-1PEG2_1PEG2_IsoGlu_C10_Diacid;

-1PEG2_1PEG2_IsoGlu_C12_Diacid;

-1PEG2_1PEG2_IsoGlu_C14_Diacid;

-1PEG2_1PEG2_IsoGlu_C16_Diacid;

-1PEG2_1PEG2_IsoGlu_C18_Diacid;

-1PEG2_1PEG2_IsoGlu_C22_Diacid;

-1PEG2_1PEG2_Ahx_C18_Diacid;

-1PEG2_1PEG2_C18_Diacid;

-1PEG8_IsoGlu_C18_Diacid;

-IsoGlu_C18_Diacid;

-PEG12_Ahx_C18_Diacid;

-PEG12_C16_Diacid;

-PEG12_C18_Diacid;

-1PEG2_1PEG2_1PEG2_C18_Diacid;

-1PEG2_1PEG2_1PEG2_IsoGlu_C18_Diacid;

-PEG12_IsoGlu_C18_Diacid;

-PEG4_IsoGlu_C18_Diacid; or

-PEG4_PEG4_IsoGlu_C18_Diacid;

wherein

PEG11_OME is $-\text{[C(O)—CH}_2\text{—CH}_2\text{—(OCH}_2\text{CH}_2\text{)]}_{11}\text{—OMe}$;

1PEG2 is $-\text{C(O)—CH}_2\text{—(OCH}_2\text{CH}_2\text{)}_2\text{—NH—}$;

PEG4 is $-\text{C(O)—CH}_2\text{—CH}_2\text{—(OCH}_2\text{CH}_2\text{)}_4\text{—NH—}$;

PEG8 is $-\text{[C(O)—CH}_2\text{—CH}_2\text{—(OCH}_2\text{CH}_2\text{)}_8\text{—NH—}$;

1PEG8 is $-\text{[C(O)—CH}_2\text{—(OCH}_2\text{CH}_2\text{)}_8\text{—NH—}$;

PEG12 is $-\text{[C(O)—CH}_2\text{—CH}_2\text{—(OCH}_2\text{CH}_2\text{)}_{12}\text{—NH—}$;


Ado is $-\text{[C(O)—(CH}_2\text{)}_{11}\text{—NH—}$;

Cn acid is $-\text{C(O)(CH}_2\text{)}_n\text{—2-CH}_3$; C18 acid is $-\text{C(O)—(CH}_2\text{)}_{16}\text{—Me}$;

Palm is $-\text{C(O)—(CH}_2\text{)}_{14}\text{—Me}$;

isoGlu is isoglutamic acid;

isoGlu_Palm is



Ahx is $-\text{[C(O)—(CH}_2\text{)}_5\text{—NH—}$;

Cn_Diacid is $-\text{C(O)—(CH}_2\text{)}_{n-2}\text{—COOH}$; wherein n is 10, 12, 14, 16, 18, or 22.

101.-129. (canceled)

130. The hepcidin analogue or pharmaceutically acceptable salt or solvate thereof according to claim 1, wherein R¹ is IVA or isovaleric acid.

131. The hepcidin analogue or pharmaceutically acceptable salt or solvate thereof according to claim 1, wherein the peptide is a linear peptide.


132.-133. (canceled)

134. The hepcidin analogue of claim 1, or pharmaceutically acceptable salt or solvate thereof comprising or consisting of a peptide, wherein the peptide is any one of the peptides listed in Tables 6A-6B.


135. The hepcidin analogue of claim 1, or pharmaceutically acceptable salt or solvate thereof, wherein the hepcidin analogue comprises or consists of a peptide set forth in Tables 6C-6E and Table 7.

136. The hepcidin analogue of claim 1, or pharmaceutically acceptable salt or solvate thereof comprising or consisting of a peptide, wherein the peptide is

ID# 3




ID# 4




-continued

ID# 5




ID# 6





-continued


ID# 7

Isovaleric Acid—E—T—H—[Dpa]—P—[Cys]—I—[Cys]—[bhPhe]—[Lys(Ahx_Palm)]—[(D)Lys]—NH₂



ID# 21

Isovaleric Acid—E—T—H—[Dpa]—[Cys]—[Cys]—I—[Lys(1PEG2_1PEG2_Ahx_C18_Diacid)]—[bhPhe]—[(D)Lys]—NH₂


ID# 22



-continued


Isoleucine—E—T—H—[Dpa]—P—[Cys]—[Cys]—[Lys(1PEG2_1PEG2_Ahx_C18_Diacid)]—[bhPhe]—[(D)Lys]—NH₂

ID# 23


Isoleucine—E—T—H—[Dpa]—P—[Cys]—I—[Lys(1PEG2_1PEG2_Ahx_C18_Diacid)]—[bhPhe]—[(D)Lys]—NH₂

-continued

ID# 27



ID# 36



137.-138. (canceled)

139. A pharmaceutical composition comprising the hepcidin analogue or pharmaceutically acceptable salt or solvate thereof of claim 1, and a pharmaceutically acceptable carrier, excipient or vehicle.

140. A method of binding a ferroportin or inducing ferroportin internalization and degradation, comprising contacting the ferroportin with at least one hepcidin analogue or pharmaceutically acceptable salt or solvate thereof of claim 1, or the pharmaceutical composition of claim 139.

141. A method for treating a disease of iron metabolism or a disease or disorder associated with dysregulated hepcidin signaling in a subject in need thereof comprising providing to the subject an effective amount of the hepcidin analogue or pharmaceutically acceptable salt or solvate thereof of claim 1 or the pharmaceutical composition of claim 131.

142. (canceled)

143. The method of claim 141, wherein the pharmaceutical composition is provided to the subject by an oral, intravenous, peritoneal, intradermal, subcutaneous, intramuscular, intrathecal, inhalation, vaporization, nebulization, sublingual, buccal, parenteral, rectal, vaginal, or topical route of administration.

144. The method of claim 143, wherein the pharmaceutical composition is provided to the subject by an oral or subcutaneous route of administration.

145. The method of claim 141, wherein the disease or disorder is a disease of iron metabolism.

146. The method of claim 145, wherein the disease of iron metabolism is an iron overload disease.

147. The method of claim 141, wherein the disease or disorder is a hemochromatosis, a thalassemia, or a polycythemia vera.

148. The method of claim 141, wherein the hepcidin analogue or pharmaceutically acceptable salt or solvate thereof or the pharmaceutical composition is provided to the subject at most twice daily, at most once daily, at most once every two days, at most once a week, or at most once a month.

149. The method of claim 141, wherein the hepcidin analogue or pharmaceutically acceptable salt or solvate thereof or the pharmaceutical composition is provided to the subject at a dosage of about 1 mg to about 100 mg.

150. A device comprising the pharmaceutical composition of claim 139, for delivery of the hepcidin analogue or pharmaceutically acceptable salt or solvate thereof to a subject, optionally orally or subcutaneously.

151. A kit comprising the pharmaceutical composition of claim 139, packaged with a reagent, a device, or an instructional material, or a combination thereof.

* * * * *