



US 20250277038A1

(19) **United States**

(12) **Patent Application Publication**
WANG et al.

(10) **Pub. No.: US 2025/0277038 A1**

(43) **Pub. Date: Sep. 4, 2025**

(54) **ANTI-PD-L1 ANTIBODY AND USE THEREOF**

(30) **Foreign Application Priority Data**

(71) Applicants: **SHANGHAI JEMINCARE
PHARMACEUTICAL CO., LTD.,**
Shanghai (CN); **JIANGXI
JEMINCARE GROUP CO., LTD.,**
Nanchang (CN)

Feb. 10, 2021 (CN) 202110183554.X

(72) Inventors: **Zongda WANG**, Shanghai (CN);
Jianqiu SONG, Shanghai (CN);
Xiaowu LIU, Shanghai (CN); **Xiaodan
CAO**, Shanghai (CN); **Sujun DENG**,
Shanghai (CN); **Zhongzong PAN**,
Shanghai (CN); **Xueping WANG**,
Shanghai (CN)

Publication Classification

(51) **Int. Cl.**
C07K 16/28 (2006.01)
A61K 47/68 (2017.01)
A61P 37/04 (2006.01)
(52) **U.S. Cl.**
CPC **C07K 16/2827** (2013.01); **A61K 47/6849**
(2017.08); **A61P 37/04** (2018.01); **C07K**
2317/24 (2013.01); **C07K 2317/33** (2013.01);
C07K 2317/52 (2013.01); **C07K 2317/565**
(2013.01); **C07K 2317/567** (2013.01); **C07K**
2317/76 (2013.01); **C07K 2317/92** (2013.01)

(73) Assignees: **SHANGHAI JEMINCARE
PHARMACEUTICAL CO., LTD.,**
Shanghai (CN); **JIANGXI
JEMINCARE GROUP CO., LTD.,**
Nanchang (CN)

(57) **ABSTRACT**

(21) Appl. No.: **18/264,819**

(22) PCT Filed: **Feb. 9, 2022**


(86) PCT No.: **PCT/CN2022/075599**

§ 371 (c)(1),

(2) Date: **Aug. 9, 2023**

Provided is an isolated antigen binding protein capable of binding to a PD-L1 protein, capable of blocking binding between a PD-1 protein and the PD-L1 protein, and/or capable of stimulating immune cells to secrete cytokines. Also provided is use of the isolated antigen binding protein in the preparation of a drug.

Specification includes a Sequence Listing.



µg/mL	1910G2H2L2H2	1910G2H2L3H3	1910G2H2L4H4	1910G2H2L7H7	1910G2H2L9H9	Atazoximab	IgG1 N297A
EC50	0.02547	0.01965	0.01285	0.02185	0.01757	0.03887	NA




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5

ANTI-PD-L1 ANTIBODY AND USE THEREOF**CROSS-REFERENCE TO RELATED APPLICATIONS**

[0001] This application is a national stage entry of International application PCT/CN2022/075599 filed on Feb. 9, 2022 that claims the priority of Chinese Patent Application 202110183554.X filed on Feb. 10, 2021. These application are incorporated herein by reference in their entirety.

FIELD OF THE INVENTION

[0002] The present application relates to the field of biomedicine, in particular to an anti-PD-L1 antibody and its use in the manufacture of a medicament.

REFERENCE TO SEQUENCE LISTING SUBMITTED VIA EFS-WEB

[0003] This application includes an electronically submitted sequence listing in _ST25.text format. The _ST25.text file contains a sequence listing entitled "U 021435-8_ST25.text". The sequence listing contained in this ST25. text file is part of the specification and is hereby incorporated by reference herein in its entirety.

BACKGROUND OF THE INVENTION

[0004] Malignant tumor is a disease that seriously threaten human health worldwide at present, and it is the main type of disease that causes human death. With the aging of population in China, the incidence of cancer is increasing, and effective therapeutic drugs need to be developed urgently, among which the development of immune checkpoint drugs has become a research hotspot in recent years.

[0005] The mechanism of PD-1 or PD-L1 immunotherapy is to design specific monoclonal antibodies against PD-1 or PD-L1, prevent the recognition of PD-1 and PD-L1, and restore the normal function of T cells, so that T cells can effectively kill tumor cells. Currently, many therapeutic antibodies have been developed against this signaling pathway, such as Durvalumab and Atezolizumab. However, during treating, there are widely existing phenomena such as low response rate and susceptible drug resistance. Therefore, an anti-tumor PD-L1 antibody with stable structure, excellent therapeutic effect and suitable for large-scale industrial production is urgently needed.

SUMMARY OF THE INVENTION

[0006] The present application provides an isolated antigen-binding protein having one or more of the following properties: (1) capable of binding to PD-L1 protein derived from a primate with a K_D value of 1×10^9 M or less; (2) capable of blocking the binding of PD-L1 protein to PD-1 protein; and (3) capable of stimulating the secretion of cytokines in immune cells.

[0007] In some embodiments, the primate includes a human and/or a monkey.

[0008] In some embodiments, the isolated antigen-binding protein includes an antibody or an antigen-binding fragment thereof.

[0009] In some embodiments, the antigen-binding fragment includes Fab, Fab', F(ab)₂, Fv fragment, F(ab')₂, scFv, di-scFv, VHH and/or dAb.

[0010] In some embodiments, the antibody is selected from the group consisting of monoclonal antibody, chimeric antibody, humanized antibody and fully human antibody.

[0011] In some embodiments, the isolated antigen-binding protein competes with a reference antibody for binding to PD-L1 protein, wherein the reference antibody comprises a heavy chain variable region VH and a light chain variable region VL, the VH of the reference antibody comprises HCDR1, HCDR2 and HCDR3, the VL of the reference antibody comprises LCDR1, LCDR2 and LCDR3, and the HCDR1 of the reference antibody comprises an amino acid sequence as set forth in SEQ ID NO: 1, the HCDR2 of the reference antibody comprises an amino acid sequence as set forth in SEQ ID NO: 2, the HCDR3 of the reference antibody comprises an amino acid sequence as set forth in SEQ ID NO: 3, the LCDR1 of the reference antibody comprises an amino acid sequence as set forth in SEQ ID NO: 4, the LCDR2 of the reference antibody comprises an amino acid sequence as set forth in SEQ ID NO: 5, and the LCDR3 of the reference antibody comprises an amino acid sequence as set forth in SEQ ID NO: 6.

[0012] In some embodiments, the isolated antigen-binding protein comprises HCDR3, and the HCDR3 comprises an amino acid sequence as set forth in SEQ ID NO: 3.

[0013] In some embodiments, the isolated antigen-binding protein comprises HCDR2, and the HCDR2 comprises an amino acid sequence as set forth in SEQ ID NO: 2.

[0014] In some embodiments, the isolated antigen-binding protein comprises HCDR1, and the HCDR1 comprises an amino acid sequence as set forth in SEQ ID NO: 1.

[0015] In some embodiments, the isolated antigen-binding protein comprises HCDR1, HCDR2 and HCDR3, wherein the HCDR1 comprises an amino acid sequence as set forth in SEQ ID NO: 1, the HCDR2 comprises an amino acid sequence as set forth in SEQ ID NO: 2, and the HCDR3 comprises an amino acid sequence as set forth in SEQ ID NO: 3.

[0016] In some embodiments, the isolated antigen-binding protein comprises a heavy chain variable region VH, wherein the VH comprises a framework region H-FR1, and the C-terminus of the H-FR1 is directly or indirectly linked to the N-terminus of the HCDR1, and the H-FR1 comprises an amino acid sequence as set forth in SEQ ID NO: 50.

[0017] In some embodiments, the H-FR1 comprises an amino acid sequence as set forth in any one of SEQ ID NO: 7 or 8.

[0018] In some embodiments, the VH comprises a framework region H-FR2, the H-FR2 is located between the HCDR1 and the HCDR2, and the H-FR2 comprises an amino acid sequence as set forth in SEQ ID NO: 51.

[0019] In some embodiments, the H-FR2 comprises an amino acid sequence as set forth in any one of SEQ ID NO: 9 to 11.

[0020] In some embodiments, the VH comprises a framework region H-FR3, the H-FR3 is located between the HCDR2 and the HCDR3, and the H-FR3 comprises an amino acid sequence as set forth in SEQ ID NO: 52.

[0021] In some embodiments, the H-FR3 comprises an amino acid sequence as set forth in any one of SEQ ID NO: 12 to 17.

[0022] In some embodiments, the VH comprises a framework region H-FR4, the N-terminus of the H-FR4 is linked

to the C-terminus of the HCDR3, and the H-FR4 comprises an amino acid sequence as set forth in SEQ ID NO: 53.

[0023] In some embodiments, the H-FR4 comprises an amino acid sequence as set forth in SEQ ID NO: 18 or 19.

[0024] In some embodiments, the isolated antigen-binding protein comprises H-FR1, H-FR2, H-FR3 and H-FR4, wherein the H-FR1 comprises an amino acid sequence as set forth in SEQ ID NO: 7 or 8, the H-FR2 comprises an amino acid sequence as set forth in any one of SEQ ID NO: 9 to 11, the H-FR3 comprises an amino acid sequence as set forth in any one of SEQ ID NO: 12 to 17 and the H-FR4 comprises an amino acid sequence as set forth in SEQ ID NO: 18 or 19.

[0025] In some embodiments, the isolated antigen-binding protein comprises H-FR1, H-FR2, H-FR3 and H-FR4, and the H-FR1, H-FR2, H-FR3 and H-FR4 are selected from any one of groups of amino acid sequences as follows:

[0026] (1) H-FR1: SEQ ID NO: 7, H-FR2: SEQ ID NO: 9, H-FR3: SEQ ID NO: 12 and H-FR4: SEQ ID NO: 18;

[0027] (2) H-FR1: SEQ ID NO: 8, H-FR2: SEQ ID NO: 10, H-FR3: SEQ ID NO: 13 and H-FR4: SEQ ID NO: 19;

[0028] (3) H-FR1: SEQ ID NO: 8, H-FR2: SEQ ID NO: 11, H-FR3: SEQ ID NO: 14 and H-FR4: SEQ ID NO: 18;

[0029] (4) H-FR1: SEQ ID NO: 8, H-FR2: SEQ ID NO: 10, H-FR3: SEQ ID NO: 15 and H-FR4: SEQ ID NO: 19;

[0030] (5) H-FR1: SEQ ID NO: 8, H-FR2: SEQ ID NO: 10, H-FR3: SEQ ID NO: 16 and H-FR4: SEQ ID NO: 19; and

[0031] (6) H-FR1: SEQ ID NO: 8, H-FR2: SEQ ID NO: 11, H-FR3: SEQ ID NO: 17 and H-FR4: SEQ ID NO: 19.

[0032] In some embodiments, the isolated antigen-binding protein comprises a heavy chain variable region VH, and the VH comprises an amino acid sequence as set forth in SEQ ID NO: 54.

[0033] In some embodiments, the VH comprises an amino acid sequence as set forth in any one of SEQ ID NO: 20 to 25.

[0034] In some embodiments, the isolated antigen-binding protein comprises a heavy chain constant region, and the heavy chain constant region is derived from a human IgG constant region.

[0035] In some embodiments, the heavy chain constant region is derived from a human IgG1 constant region, and the human IgG1 constant region comprises an amino acid sequence as set forth in SEQ ID NO: 41.

[0036] In some embodiments, the isolated antigen-binding protein comprises an antibody heavy chain HC, and the HC comprises an amino acid sequence as set forth in any one of SEQ ID NO: 42 to 46.

[0037] In some embodiments, the isolated antigen-binding protein comprises LCDR3, and the LCDR3 comprises an amino acid sequence as set forth in SEQ ID NO: 6.

[0038] In some embodiments, the isolated antigen-binding protein comprises LCDR2, and the LCDR2 comprises an amino acid sequence as set forth in SEQ ID NO: 5.

[0039] In some embodiments, the isolated antigen-binding protein comprises LCDR1, and the LCDR1 comprises an amino acid sequence as set forth in SEQ ID NO: 4.

[0040] In some embodiments, the isolated antigen-binding protein comprises LCDR1, LCDR2 and LCDR3, wherein

the LCDR1 comprises an amino acid sequence as set forth in SEQ ID NO: 4, the LCDR2 comprises an amino acid sequence as set forth in SEQ ID NO: 5, and the LCDR3 comprises an amino acid sequence as set forth in SEQ ID NO: 6.

[0041] In some embodiments, the isolated antigen-binding protein comprises a light chain variable region VL, wherein the VL comprises a framework region L-FR1, the C-terminus of the L-FR1 is directly or indirectly linked to the N-terminus of the LCDR1, and the L-FR1 comprises an amino acid sequence as set forth in SEQ ID NO: 55.

[0042] In some embodiments, the L-FR1 comprises an amino acid sequence as set forth in SEQ ID NO: 26 or 27.

[0043] In some embodiments, the VL comprises a framework region L-FR2, the L-FR2 is located between the LCDR1 and the LCDR2, and the L-FR2 comprises an amino acid sequence as set forth in SEQ ID NO: 56.

[0044] In some embodiments, the L-FR2 comprises an amino acid sequence as set forth in SEQ ID NO: 28 or 29.

[0045] In some embodiments, the VL comprises a framework region L-FR3, the L-FR3 is located between the LCDR2 and the LCDR3, and the L-FR3 comprises an amino acid sequence as set forth in SEQ ID NO: 57.

[0046] In some embodiments, the L-FR3 comprises an amino acid sequence as set forth in any one of SEQ ID NO: 30 to 33.

[0047] In some embodiments, the VL comprises a framework region L-FR4, the N-terminus of the L-FR4 is linked to the C-terminus of the LCDR3, and the L-FR4 comprises an amino acid sequence as set forth in SEQ ID NO: 58.

[0048] In some embodiments, the L-FR4 comprises an amino acid sequence as set forth in SEQ ID NO: 34 or 35.

[0049] In some embodiments, the isolated antigen-binding protein comprises L-FR1, L-FR2, L-FR3 and L-FR4, wherein the L-FR1 comprises an amino acid sequence as set forth in SEQ ID NO: 26 or 27, the L-FR2 comprises an amino acid sequence as set forth in SEQ ID NO: 28 or 29, the L-FR3 comprises an amino acid sequence as set forth in any one of SEQ ID NO: 30 to 33, and the L-FR4 comprises an amino acid sequence as set forth in SEQ ID NO: 34 or 35.

[0050] In some embodiments, the isolated antigen-binding protein comprises L-FR1, L-FR2, L-FR3 and L-FR4, and the L-FR1, L-FR2, L-FR3 and L-FR4 are selected from any one of groups of amino acid sequences as follows:

[0051] (1) L-FR1: SEQ ID NO: 26, L-FR2: SEQ ID NO: 28, L-FR3: SEQ ID NO: 30 and L-FR4: SEQ ID NO: 34;

[0052] (2) L-FR1: SEQ ID NO: 27, L-FR2: SEQ ID NO: 29, L-FR3: SEQ ID NO: 31 and L-FR4: SEQ ID NO: 35;

[0053] (3) L-FR1: SEQ ID NO: 27, L-FR2: SEQ ID NO: 29, L-FR3: SEQ ID NO: 32 and L-FR4: SEQ ID NO: 35; and

[0054] (4) L-FR1: SEQ ID NO: 27, L-FR2: SEQ ID NO: 29, L-FR3: SEQ ID NO: 33 and L-FR4: SEQ ID NO: 35.

[0055] In some embodiments, the isolated antigen-binding protein comprises a light chain variable region VL, and the VL comprises an amino acid sequence as set forth in SEQ ID NO: 59.

[0056] In some embodiments, the VL comprises an amino acid sequence as set forth in any one of SEQ ID NO: 36 to 39.

[0057] In some embodiments, the isolated antigen-binding protein comprises an antibody light chain constant region, and the antibody light chain constant region comprises an amino acid sequence as set forth in SEQ ID NO: 40.

[0058] In some embodiments, the isolated antigen-binding protein comprises an antibody light chain LC, and the LC comprises an amino acid sequence as set forth in any one of SEQ ID NO: 47 to 49.

[0059] In another aspect, the present application provides a polypeptide comprising the isolated antigen-binding protein.

[0060] In another aspect, the present application provides an immunoconjugate comprising the isolated antigen-binding protein or the polypeptide.

[0061] In another aspect, the present application provides an isolated nucleic acid molecule(s) encoding the isolated antigen-binding protein.

[0062] In another aspect, the present application provides a vector comprising the nucleic acid molecule.

[0063] In another aspect, the present application provides a cell comprising the nucleic acid molecule or the vector.

[0064] In another aspect, the present application provides a method for preparing the isolated antigen-binding protein, comprising culturing the cell under conditions that the isolated antigen-binding protein can be expressed.

[0065] In another aspect, the present application provides a pharmaceutical composition comprising the isolated antigen-binding protein, the nucleic acid molecule, the vector and/or the cell, and optionally a pharmaceutically acceptable carrier.

[0066] In another aspect, the present application provides the use of the isolated antigen-binding protein, the nucleic acid molecule, the vector, the cell and/or the pharmaceutical composition in the manufacture of a medicament for preventing, alleviating and/or treating a tumor.

[0067] In another aspect, the present application provides a method for preventing, alleviating or treating a tumor, comprising administering the isolated antigen-binding protein and/or the polypeptide to a subject in need.

[0068] In another aspect, the present application provides the isolated antigen-binding protein, for use in preventing, alleviating or treating a tumor.

[0069] In some embodiments, the tumor comprises a tumor with a high expression of PD-1 or PD-L1.

[0070] In some embodiments, the tumor comprises a solid tumor.

[0071] In some embodiments, the tumor comprises breast cancer, lung cancer, gastric cancer and urothelial carcinoma.

[0072] In another aspect, the present application provides a method for inhibiting the binding of PD-1 protein to PD-L1 protein, comprising administering the isolated antigen-binding protein and/or the polypeptide.

[0073] In another aspect, the present application provides a method for stimulating the secretion of cytokines in immune cells, comprising administering the isolated antigen-binding protein and/or the polypeptide.

[0074] In another aspect, the present application provides a method for detecting the presence and/or content of PD-L1 protein, comprising administering the isolated antigen-binding protein and/or the polypeptide.

[0075] In another aspect, the present application provides a kit comprising the isolated antigen-binding protein, the

polypeptide, the immunoconjugate, the nucleic acid molecule, the vector, the cell and/or the pharmaceutical composition.

[0076] Other aspects and advantages of the present invention will be readily apparent to those skilled in the art from the following detailed description. In the following detailed description, only exemplary embodiments of the present invention are shown and described. As will be recognized by those skilled in the art, the contents of the present application enable those skilled in the art to modify the disclosed specific embodiments without departing from the spirit and scope of the present application to which the present application relates. Accordingly, the descriptions in the drawings and specification of the present application are only exemplary but not restrictive.

BRIEF DESCRIPTION OF DRAWINGS

[0077] The specific features of the present application to which the present application relates are shown in the appended claims. The features and advantages of the present application to which the present application relates will be better understood by referring to the exemplary embodiments and drawings described in detail below. A brief description of the attached drawings is as follows:

[0078] FIG. 1 shows that the antigen-binding protein described in the present application binds to human PD-L1 protein on HEK293 cell.

[0079] FIG. 2 shows that the antigen-binding protein described in the present application binds to monkey PD-L1 protein on CHOK1 cell.

[0080] FIG. 3 shows that the antigen-binding protein described in the present application blocks the binding of human PD-1 to human PD-L1 protein on HEK293 cell.

[0081] FIG. 4 shows that the antigen-binding protein described in the present application blocks the binding of human PD-1 to monkey PD-L1 protein on CHOK1 cell.

[0082] FIG. 5 shows that the antigen-binding protein described in the present application stimulates immune cells to secrete cytokine IL-2.

DETAILED DESCRIPTION OF THE INVENTION

[0083] The embodiments of the present application will be described in the following by specific examples, and those skilled in the art can easily understand other advantages and effects of the present application from the disclosure in this specification.

Definition of Terms

[0084] In the present application, the term “PD-1” generally refers to the programmed death 1 receptor, which is also be called “programmed death 1”, “CD279”, “differentiation cluster 279”, “PD1”, “PDCD1” or “CD297”. PD-1 protein usually comprises an extracellular IgV domain, a transmembrane domain and an intracellular tail. PD-1 is usually expressed on T cells, B cells, natural killer T cells, activated monocytes and dendritic cells (DC). PD-1 can bind to its ligands PD-L1 and PD-L2. The term “PD-1” encompasses any natural PD-1 or modified PD-1 derived from any vertebrates, including mammals, such as primates (for example, human or monkeys) and rodents (for example, mice or rats). The term encompasses “full-length”, unprocessed PD-1 and any form of PD-1 produced by processing

in cells. PD-1 can exist as a transmembrane protein or as a soluble protein. “PD-1” comprises complete PD-1 and fragments thereof, as well as functional variants, isoforms, homologues, derivatives and analogues of PD-1, and analogues with at least one common epitope with PD-1. PD-1 sequences are known in the art. For example, an exemplary full-length human PD-1 protein sequence can be found by NCBI Accession Number NP_005009.2, and an exemplary full-length cynomolgus monkey PD-1 protein sequence can be found by NCBI Accession Number NP_001271065 or Uniprot Accession Number B0LAJ3.

[0085] In the present application, the term “PD-L1” generally refers to the programmed death ligand 1 protein. PD-L1 is also called differentiation cluster 274 (CD274) or B7 homologue 1 (B7-H1), and it is a protein encoded by (human) CD274 gene. PD-L1 can bind to its receptor, such as programmed death 1 (PD-1). The complexation of PD-L1 and PD-1 plays an immunosuppressive role by inhibiting T cell proliferation and producing cytokines IL-2 and IFN- γ . The term “PD-L1” encompasses any natural PD-L1 or modified PD-1 derived from any vertebrates, including mammals, such as primates (for example, human or monkeys) and rodents (for example, mice or rats). The term encompasses “full-length”, unprocessed PD-L1 and any form of PD-L1 produced by processing in cells. PD-L1 can exist as a transmembrane protein or as a soluble protein. The term also encompasses naturally occurring variants of PD-L1, such as splicing variants or allelic variants. The basic structure of PD-L1 comprises four domains: extracellular Ig-like V domain and Ig-like C2 domain, transmembrane domain and cytoplasmic domain. PD-L1 sequences are known in the art. For example, information about human PD-L1 gene (including genomic DNA sequence) can be found by NCBI Gene ID No.29126. The amino acid sequence of an exemplary full-length human PD-L1 protein can be found by NCBI Accession Number NP_054862 or UniProt Accession Number Q9NZQ7.

[0086] In the present application, the term “antigen-binding protein” generally refers to a protein containing a portion that binds to an antigen, and optionally a scaffold or skeleton part portion that allows the portion that binds to an antigen with a conformation that promotes the binding of the antigen-binding protein to an antigen. An antigen-binding protein may typically comprise an antibody light chain variable region (VL), an antibody heavy chain variable region (VH) or both of them, and functional fragments thereof. The variable regions of heavy and light chains contain binding domains that interact with antigens. Examples of antigen-binding proteins include but not limited to antibodies, antigen-binding fragments, immunoconjugates, multispecific antibodies (e.g., bispecific antibodies), antibody fragments, antibody derivatives, antibody analogs or fusion proteins, or the like, as long as they show the required antigen-binding activity.

[0087] In the present application, the term “antibody” generally refers to an immunoglobulin that is reactive to a specified protein or peptide or fragment thereof. Antibodies can be antibodies from any class, including but not limited to IgG, IgA, IgM, IgD and IgE, and antibodies from any subclass (e.g. IgG1, IgG2, IgG3, and IgG4). The antibody may comprise a heavy chain constant region selected from IgG1, IgG2, IgG3, or IgG4, for example. The antibody may also comprise a light chain selected from kappa (κ) or

lambda (λ), for example. The antibodies of the present application can be derived from any species.

[0088] In the present application, the term “antigen-binding fragment” generally refers to a portion of an antibody molecule, which contains an amino acid residue that interacts with an antigen and gives the antibody specificity and affinity to an antigen. Examples of antigen-binding fragments may include but not limited to Fab, Fab', F(ab)₂, a Fv fragment, F(ab')₂, scFv, di-scFv and/or dAb. In the present application, the term “Fab” generally refers to a fragment containing a heavy chain variable domain and a light chain variable domain, and also contains a constant domain of the light chain and a first constant domain (CH1) of the heavy chain; the term “Fab'” generally refers to a fragment different from Fab by adding a small amount of residues (including one or more cysteine from the antibody hinge region) to the carboxyl end of the CH1 domain of the heavy chain; the term “F(ab')₂” generally refers to the dimer of Fab', which contains antibody fragments of two Fab fragments connected by a disulfide bridge on the hinge region. The term “Fv” generally refers to a smallest antibody fragment containing complete antigen recognition and binding sites. In some embodiments, the fragment can be composed of a dimer with a heavy chain variable region and a light chain variable region in closely non-covalent bonding; the term “dsFv” generally refers to disulfide-stabilized Fv fragments, in which the bond between a single light chain variable region and a single heavy chain variable region is disulfide bond. The term “dAb fragment” generally refers to an antibody fragment composed of a VH domain. In the present application, the term “scFv” generally refers to a monovalent molecule formed by covalent connection and pairing of a heavy chain variable domain and a light chain variable domain of an antibody through a flexible peptide linker; such scFv molecule can comprise the general structure: NH₂—VL-linker-VH—COOH or NH₂—VH-linker-VL—COOH.

[0089] In the present application, the term “variable region” or “variable domain” generally refers to the domain of antibody heavy chain or light chain involved in the binding of antibody to antigen. In the present application, the term “variable” generally means that some portions of the sequence of the variable domain of an antibody change strongly, forming the binding and specificity of various specific antibodies to their specific antigens. Variability is distributed in the whole variable region of antibody unevenly. It mainly exists in three segments in the light chain variable region and the heavy chain variable region, which are called complementarity determining region (CDR) or high variable region (HVR), namely LCDR1, LCDR2, LCDR3, HCDR1, HCDR2 and HCDR3 respectively. The more highly conservative portion of the variable domain is called a framework region (FR). The variable domains of natural heavy chain and light chain comprise four FR regions (H-FR1, H-FR2, H-FR3, H-FR4, L-FR1, L-FR2, L-FR3, L-FR4) respectively, and most of them have B-folding configuration and are connected by three CDR structural loop regions. CDRs in each chain are close together through FR regions, and together with the CDRs from the other chain form an antigen binding site of the antibody.

[0090] In this field, variable regions of antibodies or CDRs of antibodies can be numbered by various methods, such as Kabat numbering scheme and definition rules based on

sequence variability (see Kabat et al., *protein sequence in immunology*, 5th Edition, National Institutes of Health, Bethesda, Maryland (1991)), Chothia numbering scheme and definition rules based on the position of structural loop regions (see, A1-Lazikani et al., *J Mol Biol* 273:927-48, 1997), IMGT numbering scheme and definition rules based on amino acid sequence alignment of germline V gene by efranc et al., and Honneger's numbering scheme (AHO's), Martin numbering scheme, Gelfand numbering scheme, etc. See Mathieu Dondelinger et al., Understanding the Significance and Implications of Antibody Numbering and Antigen-Binding Surface/Residue Definition, *Front. Immunol.*, 16 Oct. 2018.

[0091] In the present application, the term “monoclonal antibody” generally refers to an antibody obtained from a group of basically homogeneous antibodies, that is, each antibody constituting the group is the same, except for possible naturally occurring mutations and/or post-translational modifications (e.g., isomerization, amidation) that may exist in a very small amount. Monoclonal antibodies are highly specific and for a single antigenic site.

[0092] In the present application, the term “chimeric antibody” generally refers to an antibody in which the variable region is derived from one species and the constant region is derived from another species. Generally, the variable region is derived from an antibody of an experimental animal such as a rodent (“parent antibody”), and the constant region is derived from a human antibody, so that the obtained chimeric antibody is less likely to cause an adverse immune response in a human individual, compared with a parent (e.g., mouse-derived) antibody.

[0093] In the present application, the term “humanized antibody” generally refers to an antibody in which some or all of the amino acids other than the CDR region of a non-human antibody (e.g. a mouse antibody) are replaced by corresponding amino acids derived from human immunoglobulin. Addition, deletion, insertion, substitution or modification of amino acids in the CDR region can also be allowed, as long as they still retain the ability of antibodies to bind to specific antigens. A humanized antibody may optionally comprise at least a portion of the constant region of human immunoglobulin. A humanized antibody retains the antigen specificity similar to that of the original antibody. A “humanized” form of a non-human (e.g., murine) antibody may comprise a chimeric antibody derived from a sequence of a non-human immunoglobulin at a minimum. In some cases, the residues of CDR region in human immunoglobulin (receptor antibody) can be replaced by the residues of CDR region of a non-human species (donor antibody) (such as mice, rats, rabbits or non-human primates) with desired properties, affinity and/or ability. In some cases, the residues in the FR region of human immunoglobulin can be replaced by corresponding non-human residues. In addition, the humanized antibody may comprise amino acid modifications that are not present in the recipient antibody or in the donor antibody.

[0094] In the present application, the term “fully human antibody” generally refers to an antibody in which all portions (including the variable region and the constant region of the antibody) are encoded by genes of human origin. The methods to obtain fully human antibodies in this field may comprise phage display technology, transgenic mouse technology, ribosome display technology and RNA-polypeptide technology.

[0095] In the present application, the terms “binding”, “specifically binding” or “specific for” generally refer to measurable and reproducible interactions, such as the binding of antigens with antibodies, which can determine the presence of a target in the presence of heterogeneous populations of molecules, including biological molecules. For example, an antibody binds to an epitope through its antigen-binding domain, and this binding requires some complementarity between the antigen-binding domain and the epitope. For example, an antibody that specifically binds to a target (which may be an epitope) is an antibody that binds to this target with greater affinity, binding affinity, and easier and/or longer duration than binding of the antibody to other targets. An antibody is called to “specifically bind” to an antigen when it is more likely to bind to an epitope through its antigen-binding domain than binding of the antibody to a random, unrelated epitope.

[0096] In the present application, the terms “KD” and “ K_D ” can be used interchangeably and usually refer to the equilibrium dissociation constant, and “KD” is a ratio of the dissociation rate constant (k_{dis} , also known as “off-rate (k_{off})” or “kd”) to the association rate constant (k_{on} , also known as “bonding rate (k_{on})” or “ka”). Association rate constant (k_{on}), dissociation rate constant (k_{dis}) and equilibrium dissociation constant (KD) can be used to express the binding affinity of antigen-binding proteins (e.g. antibodies) to antigens. Methods for determining the association and dissociation rate constants are well known in the art, including but not limited to Biolayer interferometry (BLI), Radioimmunoassay (RIA), Equilibrium Dialysis, Surface Plasmon Resonance (SPR), Fluorescence Resonance Energy Transfer (FRET), co-immunoprecipitation (Co-IP) and protein Chip Technology. If measured under different conditions (e.g. salt concentration and pH), the measured affinity of a specific protein-protein interaction may be different.

[0097] In the present application, the term “primate” usually refers to monkey and ape species, and includes monkey species, such as the monkeys from the *Macaca* (such as, cynomolgus (*Macaca fascicularis*) and/or rhesus monkey (*Macaca mulatta*)) and baboon (*Papio ursinus*), as well as marmoset (a species from *Callithrix*), squirrel monkey (a species from *Saimiri*), and tamarin (a species from *Saguinus*), and ape species, such as chimpanzee (*Pan troglodytes*), and also includes *Homo sapiens*.

[0098] In the present application, the terms “polypeptide” or “protein” are used interchangeably and generally refer to polymers of amino acid residues. The term also applies to amino acid polymers in which one or more amino acid residues are analogs or mimetics of corresponding naturally occurring amino acids, as well as naturally occurring amino acid polymers. The term may also comprise modified amino acid polymers, for example, by adding sugar residues to form glycoproteins or modification of phosphorylation. Polypeptides and proteins can be produced by naturally occurring and non-recombinant cells or genetically engineered or recombinant cells, and may comprise molecules with amino acid sequences of natural protein, or molecules with deletion, addition and/or substitution of one or more amino acids of natural sequences. The terms “polypeptide” and “protein” particularly comprise deletion, addition and/or substitution of one or more amino acids of a sequence of the antigen-binding protein described in the present application.

[0099] In the present application, the term “homologue” generally refers to an amino acid sequence or nucleotide

sequence that has certain homology with the wild-type amino acid sequence and the wild-type nucleotide sequence. The term “homology” can be equivalent to sequence “identity”. Homologous sequences may comprise amino acid sequences that may be at least 80%, 85%, 90%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8% or 99.9% identical to the subject sequence. Typically, the homologue will comprise the same active site or the like as the subject amino acid sequence. Homology may be considered in terms of similarity (i.e., amino acid residues having similar chemical properties/functions), or may be expressed in terms of sequence identity. In the present application, a sequence having percentage identity in any one of the mentioned SEQ ID NO of a mentioned amino acid sequence or nucleotide sequence refers to a sequence having the percentage identity over the entire length of the mentioned SEQ ID NO.

[0100] In the present application, the term “isolated” generally refers to biological materials (e.g. viruses, nucleic acids, or protein) that are substantially free of components normally associated with or interacting with their naturally occurring environment. The isolated biological materials optionally includes additional materials that are not found in its natural environment (e.g., nucleic acids or proteins). In the present application, when it comes to protein, “isolation” usually means that the molecule is isolated and separated from the whole organism in which the molecule is found to exist naturally, or there are basically no other biomacromolecules of the same type. When a nucleic acid molecule is referred to, it is completely or partially separated from the sequence that naturally binds to it, or the nucleic acid has a heterologous sequence that binds to it, or the nucleic acid is separated from the chromosome.

[0101] In the present application, the term “immunoconjugate” generally refers to the substance formed by the connection of antigen-binding protein with other active agents, which may be a small molecular active agent, such as a chemotherapeutic agent, a toxin, an immunotherapy agent, an imaging probe or a fluorescent probe.

[0102] In the present application, the term “nucleic acid” molecule generally refers to any length of isolated form of nucleotide, deoxynucleotide or ribonucleotide or its analogues isolated from its natural environment or artificially synthesized.

[0103] In the present application, the term “vector” generally refers to a nucleic acid molecule capable of self-replication in a suitable host, which transfers the inserted nucleic acid molecule into and/or between host cells. The vector may comprise a vector mainly used for inserting DNA or RNA into cells, a vector mainly used for replicating DNA or RNA, and a vector mainly used for transcription and/or expression of translation of DNA or RNA. The vector also comprises a vector with a plurality of the above functions. The vector can be a polynucleotide that can be transcribed and translated into a polypeptide when introduced into a suitable host cell. Generally, the vector can produce a desired expression product by culturing a suitable host cell comprising the vector.

[0104] In the present application, the term “cell” generally refers to an individual cell, cell line or cell culture that may comprise or already has comprised a plasmid or vector including a nucleic acid molecule described in the present application, or that can express an antigen-binding protein described in the present application. The cells may comprise

progeny of a single host cell. Because of natural, unexpected or intentional mutation, the progeny cells may not be exactly the same as the original parent cells in morphology or genome, but they can express the antibody or antigen-binding fragment thereof described in the present application. The cells can be obtained by transfecting cells in vitro using the vector described in the present application. The cells can be prokaryotic cells (e.g. *Escherichia coli*) or eukaryotic cells (e.g. yeast cells, e.g. COS cells, China hamster ovary (CHO) cells, HeLa cells, HEK293 cells, COS-1 cells, NSO cells or myeloma cells). In some cases, the cells may be mammalian cells. For example, the mammalian cells may be CHO-K1 cells.

[0105] In the present application, the term “pharmaceutical composition” generally refers to a preparation that exists in a form that allows the biological activity of an active ingredient to be effective, and does not contain additional ingredients with unacceptable toxicity to a subject to whom the composition is to be administered.

[0106] In the present application, the term “treatment” generally refers to the desire to change the natural course of the individual being treated, and can be clinical intervention to achieve prevention and treatment or in the process of clinical lesions. Desirable therapeutic effects comprise, but not limited to, preventing the occurrence or recurrence of the disease, relieving symptoms, weakening any direct or indirect pathological consequences of the disease, preventing metastasis, reducing the rate of disease progression, improving or alleviating the disease state and alleviating or improving the prognosis. In some cases, the antigen-binding protein (e.g. anti-PD-L1 antibody) of the present application can be used to delay disease development or slow disease progression.

[0107] In the present application, the term “administration” generally refers to a method of administering a certain dose of a compound (e.g., an anticancer therapeutic agent) or a pharmaceutical composition (e.g., a pharmaceutical composition comprising an anticancer therapeutic agent) to a subject (e.g., a patient). Administration can be carried out by any suitable means, including parenteral, intrapulmonary and intranasal administration, and (if necessary for local treatment) intralesional administration. Parenteral infusion comprises, for example, intramuscular, intravenous, intra-arterial, intraperitoneal or subcutaneous administration.

[0108] In the present application, the term “tumor” generally refers to the growth and proliferation of all neoplastic cell (either malignant or benign) and all precancerous and cancerous cells and tissues. In the present application, the tumor can be a tumor with a high expression of PD-1 or PD-L1 in cells and tissues. Tumors may comprise solid tumors and/or non-solid tumors (e.g., hematomas, lymphomas).

[0109] In the present application, the term “between” generally means that the C-terminus of an amino acid fragment is directly or indirectly linked to the N-terminus of a first amino acid fragment, and its N-terminus is directly or indirectly linked to the C-terminus of a second amino acid fragment. In the light chain, for example, the N-terminus of the L-FR2 is directly or indirectly linked to the C-terminus of the LCDR1, and the C-terminus of the L-FR2 is directly or indirectly linked to the N-terminus of the LCDR2. For another example, the N-terminus of the L-FR3 is directly or indirectly linked to the C-terminus of the LCDR2, and the C-terminus of the L-FR3 is directly or indirectly linked to

the N-terminus of the LCDR3. In the heavy chain, for example, the N-terminus of the H-FR2 is directly or indirectly linked to the C-terminus of the HCDR1, and the C-terminus of the H-FR2 is directly or indirectly linked to the N-terminus of the HCDR2. For another example, the N-terminus of the H-FR3 is directly or indirectly linked to the C-terminus of the HCDR2, and the C-terminus of the H-FR3 is directly or indirectly linked to the N-terminus of the HCDR3. In the present application, the “first amino acid fragment” and the “second amino acid fragment” can be any one of the same or different amino acid fragments.

[0110] In the present application, the term “includes/include” generally refers to the meaning of comprising, inclusive, containing, or encompassing. In some cases, it also represents the meanings of “is”, and “consist of”.

[0111] In the present application, the term “about” generally refers to a change in the range of 0.5%-10% above or below a specified value, such as a change in the range of 0.5%, 1%, 1.5%, 2%, 2.5%, 3%, 3.5%, 4%, 4.5%, 5%, 5.5%, 6%, 6.5%, 7%, 7.5%, 8%, 8.5%, 9%, 9.5%, or 10% above or below a specified value.

DETAILED DESCRIPTION OF THE PRESENT APPLICATION

Antigen Binding Protein

[0112] In an aspect, the present application provides an isolated antigen-binding protein, which can bind to PD-L1 derived from primates (e.g. humans or monkeys) with a KD value of 1×10^{-9} M or less. The binding affinity of primate PD-L1 antigen-binding protein to PD-L1 can be determined by any method known in the art. In some cases, the binding affinity can be determined by surface plasmon resonance (SPR), enzyme-linked immunosorbent assay (ELISA), binding antigen precipitation method, equilibrium dialysis method and biolayer interference (BLI). In some cases, the binding affinity and KD value of PD-L1 antigen-binding protein to PD-L1 can be determined by biolayer interference (BLI). For example, an ForteBio Octet molecular interaction analyzer can be used to analyze the binding kinetics between antigens and antibodies.

[0113] In the present application, the isolated antigen-binding protein can bind to PD-L1 derived from primates with a KD value of 1×10^{-9} M or less. For example, the KD value can be a value of about 1×10^{-9} M or less, about 9×10^{-10} M or less, about 8×10^{-10} M or less, about 7×10^{-10} M or less, about 6×10^{-10} M or less, about 5×10^{-10} M or less, about 4×10^{-10} M or less, about 3×10^{-10} M or less, about 2×10^{-10} M or less, and about 1×10^{-10} M or less binding to PD-L1 derived from humans, for example, detected using a ForteBio Octet molecular interaction analyzer.

[0114] In another case, the binding activity of PD-L1 antigen-binding protein described in the present application to PD-L1 can be detected by flow cytometry or enzyme-linked immunosorbent assay.

[0115] For example, in FACS test, host cells stably expressing human PD-L1 (such as HEK293 cells) are used, and the EC50 value of the PD-L1 antigen-binding protein binding to PD-L1 is between about 0.0001 $\mu\text{g/mL}$ and about 100 $\mu\text{g/mL}$, for example, between about 0.001 $\mu\text{g/mL}$ and about 10 $\mu\text{g/mL}$, between about 0.001 $\mu\text{g/mL}$ and about 5 $\mu\text{g/mL}$, between about 0.001 $\mu\text{g/mL}$ and about 1 $\mu\text{g/mL}$, between about 0.01 $\mu\text{g/mL}$ and about 0.5 $\mu\text{g/mL}$, between

about 0.01 $\mu\text{g/mL}$ and about 0.1 $\mu\text{g/mL}$, or between about 0.01 $\mu\text{g/mL}$ and about 0.05 $\mu\text{g/mL}$.

[0116] For example, in FACS test, host cells stably expressing monkey PD-L1 (such as CHOK1 cells) are used, and the EC50 value of PD-L1 antigen-binding protein binding to PD-L1 is between about 0.0001 $\mu\text{g/mL}$ and about 100 $\mu\text{g/mL}$, for example, between about 0.001 $\mu\text{g/mL}$ and about 10 $\mu\text{g/mL}$, between about 0.001 $\mu\text{g/mL}$ and about 5 $\mu\text{g/mL}$, between about 0.01 $\mu\text{g/mL}$ and about 1 $\mu\text{g/mL}$, between about 0.02 $\mu\text{g/mL}$ and about 0.5 $\mu\text{g/mL}$, and between about 0.03 $\mu\text{g/mL}$ and about 0.1 $\mu\text{g/mL}$.

[0117] In another aspect, the antigen-binding protein described in the present application can block the binding of PD-1 to PD-L1. In some cases, the blocking of the binding of PD-1 to PD-L1 by the antigen-binding protein can be determined by flow cytometer FACS or enzyme-linked immunosorbent assay (ELISA).

[0118] For example, host cells stably expressing human PD-L1 (such as HEK293 cells) are first incubated with a decreasing amount of unlabeled antigen-binding protein, and then incubated with biotin-labeled PD-1 protein. Then, cells were analyzed using FACS to confirm that the antigen-binding protein blocks the binding of PD-1 to PD-L1. For example, between about 0.001 $\mu\text{g/mL}$ and about 10 $\mu\text{g/mL}$, between about 0.001 $\mu\text{g/mL}$ and about 5 $\mu\text{g/mL}$, between about 0.01 $\mu\text{g/mL}$ and about 1 $\mu\text{g/mL}$, between about 0.02 $\mu\text{g/mL}$ and about 0.5 $\mu\text{g/mL}$, and between about 0.02 $\mu\text{g/mL}$ and about 0.1 $\mu\text{g/mL}$.

[0119] For example, host cells stably expressing monkey PD-L1 (such as CHOK1 cells) are first incubated with a decreasing amount of unlabeled antigen-binding protein, and then incubated with biotin-labeled PD-1 protein. Then, cells were analyzed using FACS to confirm that the antigen-binding protein blocks the binding of PD-1 to PD-L1. For example, between about 0.001 $\mu\text{g/mL}$ to about 10 $\mu\text{g/mL}$, about 0.001 $\mu\text{g/mL}$ to about 5 $\mu\text{g/mL}$, about 0.01 $\mu\text{g/mL}$ to about 1 $\mu\text{g/mL}$, and about 0.1 $\mu\text{g/mL}$ to about 0.7 $\mu\text{g/mL}$.

[0120] In the present application, the isolated antigen-binding protein may comprise CDR3 from a heavy chain variable region VH, and the VH may comprise an amino acid sequence as set forth in SEQ ID NO: 54.

[0121] In the present application, the isolated antigen-binding protein may comprise HCDR3, and the HCDR3 may comprise CDR3 of VH with an amino acid sequence as set forth in any one of SEQ ID NO: 20 to 25. For example, the isolated antigen-binding protein may comprise HCDR3, and the HCDR3 may comprise an amino acid sequence as set forth in SEQ ID NO: 3.

[0122] In the present application, the isolated antigen-binding protein may comprise CDR2 from a heavy chain variable region VH, and the VH may comprise an amino acid sequence as set forth in SEQ ID NO: 54.

[0123] In the present application, the isolated antigen-binding protein may comprise HCDR2, and the HCDR2 may comprise CDR2 of VH with an amino acid sequence as set forth in any one of SEQ ID NO: 20 to 25. For example, the isolated antigen-binding protein may comprise HCDR2, and the HCDR2 may comprise an amino acid sequence as set forth in SEQ ID NO: 2.

[0124] In the present application, the isolated antigen-binding protein may comprise CDR1 from a heavy chain variable region VH, and the VH may comprise an amino acid sequence as set forth in SEQ ID NO: 54.

[0125] In the present application, the isolated antigen-binding protein may comprise HCDR1, and the HCDR1 may comprise CDR1 of VH with an amino acid sequence as set forth in any one of SEQ ID NO: 20 to 25. For example, the isolated antigen-binding protein may comprise HCDR1, and the HCDR1 may comprise an amino acid sequence as set forth in SEQ ID NO: 1.

[0126] In the present application, the isolated antigen-binding protein may comprise HCDR1, HCDR2 and HCDR3, the HCDR1 may comprise CDR1 of VH with an amino acid sequence as set forth in SEQ ID NO: 54, the HCDR2 may comprise CDR2 of VH with an amino acid sequence as set forth in SEQ ID NO: 54, and the HCDR3 may comprise CDR3 of VH with an amino acid sequence as set forth in SEQ ID NO: 54.

[0127] In the present application, the isolated antigen-binding protein may comprise HCDR1, HCDR2 and HCDR3, the HCDR1 may comprise CDR1 of VH with an amino acid sequence as set forth in any one of SEQ ID NO: 20 to 25, the HCDR2 may comprise CDR2 of VH with an amino acid sequence as set forth in any one of SEQ ID NO: 20 to 25, and the HCDR3 may comprise CDR3 of VH with an amino acid sequence as set forth in any one of SEQ ID NO: 20 to 25.

[0128] For example, the isolated antigen-binding protein may comprise CDR1 of VH with an amino acid sequence as set forth in SEQ ID NO: 20, may comprise CDR2 of VH with an amino acid sequence as set forth in SEQ ID NO: 20, and may comprise CDR3 of VH with an amino acid sequence as set forth in SEQ ID NO: 20.

[0129] For example, the isolated antigen-binding protein may comprise CDR1 of VH with an amino acid sequence as set forth in SEQ ID NO: 21, may comprise CDR2 of VH with an amino acid sequence as set forth in SEQ ID NO: 21, and may comprise CDR3 of VH with an amino acid sequence as set forth in SEQ ID NO: 21.

[0130] For example, the isolated antigen-binding protein may comprise CDR1 of VH with an amino acid sequence as set forth in SEQ ID NO: 22, may comprise CDR2 of VH with an amino acid sequence as set forth in SEQ ID NO: 22, and may comprise CDR3 of VH with an amino acid sequence as set forth in SEQ ID NO: 22.

[0131] For example, the isolated antigen-binding protein may comprise CDR1 of VH with an amino acid sequence as set forth in SEQ ID NO: 23, may comprise CDR2 of VH with an amino acid sequence as set forth in SEQ ID NO: 23, and may comprise CDR3 of VH with an amino acid sequence as set forth in SEQ ID NO: 23.

[0132] For example, the isolated antigen-binding protein may comprise CDR1 of VH with an amino acid sequence as set forth in SEQ ID NO: 24, may comprise CDR2 of VH with an amino acid sequence as set forth in SEQ ID NO: 24, and may comprise CDR3 of VH with an amino acid sequence as set forth in SEQ ID NO: 24.

[0133] For example, the isolated antigen-binding protein may comprise CDR1 of VH with an amino acid sequence as set forth in SEQ ID NO: 25, may comprise CDR2 of VH with an amino acid sequence as set forth in SEQ ID NO: 25, and may comprise CDR3 of VH with an amino acid sequence as set forth in SEQ ID NO: 25.

[0134] For example, the isolated antigen-binding protein described in the present application may comprise HCDR1, HCDR2 and HCDR3, the HCDR1 may comprise an amino acid sequence as set forth in SEQ ID NO: 1, the HCDR2

may comprise an amino acid sequence as set forth in SEQ ID NO: 2, and the HCDR3 may comprise an amino acid sequence as set forth in SEQ ID NO: 3.

[0135] In the present application, the isolated antigen-binding protein may comprise H-FR1, and the H-FR1 may comprise an amino acid sequence as set forth in SEQ ID NO: 50: EVQLVESGGGLVQPGX₁₆SLX₁₉LSCX₂₃AS (SEQ ID NO: 50), where X₁₆ is G or R, X₁₉ is K or R, and X₂₃ is A or V. For example, it may be divided according to Chothia rules.

[0136] In some cases, compared with an amino acid sequence as set forth in SEQ ID NO: 7, the H-FR1 may comprise at least amino acid substitutions at positions selected from the group consisting of amino acid substitutions at X₁₆, X₁₉ and/or X₂₃.

[0137] For example, the H-FR1 may comprise an amino acid sequence as set forth in SEQ ID NO: 7 or 8.

[0138] In the present application, the isolated antigen-binding protein may comprise H-FR2, and the H-FR2 may comprise an amino acid sequence as set forth in SEQ ID NO: 51: WMX₃WX₅RQAPGKGLEWVASI (SEQ ID NO: 51), where X₃ is S or T, and X₅ is I or V. For example, it may be divided according to Chothia rules.

[0139] In some cases, compared with an amino acid sequence as set forth in SEQ ID NO: 9, the H-FR2 may comprise at least amino acid substitutions at positions selected from the group consisting of amino acid substitutions at X₃ and/or X₅.

[0140] For example, the H-FR2 may comprise an amino acid sequence as set forth in any one of SEQ ID NO: 9 to 11.

[0141] In the present application, the isolated antigen-binding protein may comprise H-FR3, and the H-FR3 may comprise an amino acid sequence as set forth in SEQ ID NO: 52:

TTYX₄DSVKGRFTISRDX₁₇X₁₈KX₂₀X₂₁LYLQMNSLRX₃₁EDTAX₃₆YYCSR (SEQ ID NO: 52), where X₄ is A, P or V, X₁₇ is D or N, X₁₈ is A or S, X₂₀ is N or S, X₂₁ is S or T, X₃₁ is A or S, and X₃₆ is T or V. For example, it may be divided according to Chothia rules.

[0142] In some cases, compared with an amino acid sequence as set forth in SEQ ID NO: 12, the H-FR3 may comprise at least amino acid substitutions at positions selected from the group consisting of amino acid substitutions at X₄, X₁₇, X₁₈, X₂₀, X₂₁, X₃₁ and/or X₃₆.

[0143] For example, the H-FR3 may comprise an amino acid sequence as set forth in any one of SEQ ID NO: 12 to 17.

[0144] In the present application, the isolated antigen-binding protein may comprise H-FR4, and the H-FR4 may comprise an amino acid sequence as set forth in SEQ ID NO: 53: WGQGX₅X₆VTVSS (SEQ ID NO: 53), where X₅ is T or V, and X₆ is L or M. For example, it may be divided according to Chothia rules.

[0145] In some cases, compared with an amino acid sequence as set forth in SEQ ID NO: 18, the H-FR4 may comprise at least amino acid substitution at X₆ and/or X₅.

[0146] For example, the H-FR4 may comprise an amino acid sequence as set forth in SEQ ID NO: 18 or 19.

[0147] In the present application, the isolated antigen-binding protein may comprise a heavy chain variable region VH, and the VH may comprise an amino acid sequence as set forth in SEQ ID NO: 54: EVQLVESGGGLVQPGX₁₆SLX₁₉LSCX₂₃ASGFTFSNFWMX₃₅WX₃₇RQAPGKGLEWVASITHSGGITYYX₆₁DSVKGRFTISRDX₇₄X₇₅KX₇₇X₇₈

LYLQMNSLRX₈₈EDTAX₉₃YYCS

RDPTAPFDYWGQGX₁₁₂X₁₁₃VTVSS (SEQ ID NO: 54), where X₁₆ is G or R, X₁₉ is K or R, X₂₃ is A or V, X₃₅ is S or T, X₃₇ is I or V, X₆₁ is A, P or V, X₇₄ is D or N, X₇₅ is A or S, X₇₇ is N or S, X₇₈ is S or T, X₈₈ is A or S, X₉₃ is T or V, X₁₁₂ is T or V, and X₁₁₃ is L or M. For example, it may be divided according to Chothia rules.

[0148] In some cases, compared with an amino acid sequence as set forth in SEQ ID NO: 20, the VH may comprise at least amino acid substitutions at positions selected from the group consisting of amino acid substitutions at X₁₆, X₁₉, X₂₃, X₃₅, X₃₇, X₆₁, X₇₄, X₇₅, X₇₇, X₇₈, X₈₈, X₉₃, X₁₁₂ and/or X₁₁₃.

[0149] For example, the VH may comprise an amino acid sequence as set forth in any one of SEQ ID NO: 20 to 25.

[0150] In the present application, the isolated antigen-binding protein may comprise CDR3 from a light chain variable region VL, and the VL may comprise an amino acid sequence as set forth in SEQ ID NO: 59.

[0151] In the present application, the isolated antigen-binding protein may comprise LCDR3, and the LCDR3 may comprise CDR3 of VL with an amino acid sequence as set forth in any one of SEQ ID NO: 36 to 39. For example, the isolated antigen-binding protein may comprise LCDR3, and the LCDR3 may comprise an amino acid sequence as set forth in SEQ ID NO: 6.

[0152] In the present application, the isolated antigen-binding protein may comprise CDR2 from a light chain variable region VL, and the VL may comprise an amino acid sequence as set forth in SEQ ID NO: 59.

[0153] In the present application, the isolated antigen-binding protein may comprise LCDR2, and the LCDR2 may comprise CDR2 of VL with an amino acid sequence as set forth in any one of SEQ ID NO: 36 to 39. For example, the isolated antigen-binding protein may comprise LCDR2, and the LCDR2 may comprise an amino acid sequence as set forth in SEQ ID NO: 5.

[0154] In the present application, the isolated antigen-binding protein may comprise CDR1 from a light chain variable region VL, and the VL may comprise an amino acid sequence as set forth in SEQ ID NO: 59.

[0155] In the present application, the isolated antigen-binding protein may comprise LCDR1, and the LCDR1 may comprise CDR1 of VL with an amino acid sequence as set forth in any one of SEQ ID NO: 36 to 39. For example, the isolated antigen-binding protein may comprise LCDR1, and the LCDR1 may comprise an amino acid sequence as set forth in SEQ ID NO: 4.

[0156] In the present application, the isolated antigen-binding proteins may comprise LCDR1, LCDR2 and LCDR3, the LCDR1 may comprise CDR1 of VL with an amino acid sequence as set forth in SEQ ID NO: 59, the LCDR2 may comprise CDR2 of VL with an amino acid sequence as set forth in SEQ ID NO: 59, and the LCDR3 may comprise CDR3 of VL with an amino acid sequence as set forth in SEQ ID NO: 59.

[0157] In the present application, the isolated antigen-binding proteins may comprise LCDR1, LCDR2 and LCDR3, the LCDR1 may comprise CDR1 of VL with an amino acid sequence as set forth in any one of SEQ ID NO: 36 to 39, the LCDR2 may comprise CDR2 of VL with an amino acid sequence as set forth in any one of SEQ ID NO:

36 to 39, and the LCDR3 may comprise CDR3 of VL with an amino acid sequence as set forth in any one of SEQ ID NO: 36 to 39.

[0158] For example, the isolated antigen-binding protein may comprise CDR1 of VL with an amino acid sequence as set forth in SEQ ID NO: 36, may comprise CDR2 of VL with an amino acid sequence as set forth in SEQ ID NO: 36, and may comprise CDR3 of VL with an amino acid sequence as set forth in SEQ ID NO: 36.

[0159] For example, the isolated antigen-binding protein may comprise CDR1 of VL with an amino acid sequence as set forth in SEQ ID NO: 37, may comprise CDR2 of VL with an amino acid sequence as set forth in SEQ ID NO: 37, and may comprise CDR3 of VL with an amino acid sequence as set forth in SEQ ID NO: 37.

[0160] For example, the isolated antigen-binding protein may comprise CDR1 of VL with an amino acid sequence as set forth in SEQ ID NO: 38, may comprise CDR2 of VL with an amino acid sequence as set forth in SEQ ID NO: 38, and may comprise CDR3 of VL with an amino acid sequence as set forth in SEQ ID NO: 38.

[0161] For example, the isolated antigen-binding protein may comprise CDR1 of VL with an amino acid sequence as set forth in SEQ ID NO: 39, may comprise CDR2 of VL with an amino acid sequence as set forth in SEQ ID NO: 39, and may comprise CDR3 of VL with an amino acid sequence as set forth in SEQ ID NO: 38.

[0162] For example, the isolated antigen-binding proteins described in the present application may comprise LCDR1, LCDR2 and LCDR3, the LCDR1 may comprise an amino acid sequence as set forth in SEQ ID NO: 4, the LCDR2 may comprise an amino acid sequence as set forth in SEQ ID NO: 5, and the LCDR3 may comprise an amino acid sequence as set forth in SEQ ID NO: 6.

[0163] In the present application, the isolated antigen-binding protein may comprise L-FR1, and the L-FR1 may comprise an amino acid sequence as set forth in SEQ ID NO: 55: DIQMTQSPX₅SLSASX₁₅GDX₁₈VTITC (SEQ ID NO: 55), where X₅ is P or S, X₁₅ is L or V, and X₁₈ is Q or R. For example, it may be divided according to Chothia rules.

[0164] In some cases, compared with an amino acid sequence as set forth in SEQ ID NO: 26, the L-FR1 may comprise at least amino acid substitutions at positions selected from the group consisting of amino acid substitutions at X₉, X₁₅ and/or X₁₈.

[0165] For example, the L-FR1 may comprise an amino acid sequence as set forth in SEQ ID NO: 26 or 27.

[0166] In the present application, the isolated antigen-binding protein may comprise L-FR2, and the L-FR2 may comprise an amino acid sequence as set forth in SEQ ID NO: 56: WYQQKPGKAPX₁₁QLIR (SEQ ID NO: 56), where X₁₁ is K or R. For example, it may be divided according to Chothia rules.

[0167] In some cases, compared with an amino acid sequence as set forth in SEQ ID NO: 28, the H-FR2 may comprise at least amino acid substitutions at positions selected from the group consisting of amino acid substitutions at X₁₁.

[0168] For example, the L-FR2 may comprise an amino acid sequence as set forth in SEQ ID NO: 28 or 29.

[0169] In the present application, the isolated antigen-binding protein may comprise L-FR3, and the L-FR3 may comprise an amino acid sequence as set forth in SEQ ID NO: 57: GX₂PSRFSGSGSGKDX₁₅X₁₆FX₁₈ISX₂₁X₂₂X₂₃X₂₄

EDIAX₂₉YYC (SEQ ID NO: 57), where X₂ is T or V, X₁₅ is F or Y, X₁₆ is S or T, X₁₈ is S or T, X₂₁ is N or S, X₂₂ is L or V, X₂₃ is E or Q, X₂₄ is P or S, and X₂₉ is S or T. For example, it may be divided according to Chothia rules.

[0170] In some cases, compared with an amino acid sequence as set forth in SEQ ID NO: 30, the L-FR3 may comprise at least amino acid substitutions at positions selected from the group consisting of amino acid substitutions at X₂, X₁₅, X₁₆, X₁₈, X₂₁, X₂₂, X₂₃, X₂₄ and/or X₂₉.

[0171] For example, the L-FR3 may comprise an amino acid sequence as set forth in any one of SEQ ID NO: 30 to 33.

[0172] In the present application, the isolated antigen-binding protein may comprise L-FR4, and the L-FR4 may comprise an amino acid sequence as set forth in SEQ ID NO: 58: FGX₃GTKLEX₉K (SEQ ID NO: 58), where X₃ is A or Q and X₉ is I or L. For example, it may be divided according to Chothia rules.

[0173] In some cases, compared with an amino acid sequence as set forth in SEQ ID NO: 34, the L-FR4 may comprise at least amino acid substitutions at positions selected from the group consisting of amino acid substitutions at X₃ and/or X₉.

[0174] For example, the L-FR4 may comprise an amino acid sequence as set forth in SEQ ID NO: 34 or 35.

[0175] In the present application, the isolated antigen-binding protein may comprise a light chain variable region VL, and the VL may comprise an amino acid sequence as set forth in SEQ ID NO: 59: DIQMTQSPX₆SLSASX₁₅GDX₁₈VTITCQASQNNINYIAWYQQKPGKAPX₄₅QLIRYTS TLVSGX₅₈PSRFSGSGSGKDX₇₁X₇₂FX₇₄ISX₇₇X₇₈X₇₉X₈₀EDIAX₈₅YYCLQYDNPNTF GX₁₀₀GTKLEX₁₀₆K (SEQ ID NO: 59), where X₉ is P or S, X₁₅ is L or V, X₁₈ is Q or R, X₄₅ is K or R, X₅₈ is T or V, X₇₁ is F or Y, X₇₂ is S or T, X₇₄ is T or S, X₇₇ is N or S, X₇₈ is L or V, X₇₉ is E or Q, X₈₀ is P or S, X₈₅ is S or T, X₁₀₀ is A or Q, and X₁₀₆ is L or I. For example, it may be divided according to Chothia rules.

[0176] In some cases, compared with an amino acid sequence as set forth in SEQ ID NO: 36, and the VL may comprise at least amino acid substitutions at positions selected from the group consisting of X₉, X₁₅, X₁₈, X₄₅, X₅₈, X₇₁, X₇₂, X₇₄, X₇₇, X₇₈, X₇₉, X₈₀, X₈₅, X₁₀₀ and/or X₁₀₆.

[0177] For example, the VL may comprise an amino acid sequence as set forth in any one of SEQ ID NO: 36 to 39.

[0178] In the present application, the isolated antigen-binding protein may comprise HCDR1, HCDR2, HCDR3, LCDR1, LCDR2 and LCDR3, and the HCDR1 may comprise an amino acid sequence as set forth in SEQ ID NO: 1, and the HCDR2 may comprise an amino acid sequence as set forth in SEQ ID NO: 2, the HCDR3 may comprise an amino acid sequence as set forth in SEQ ID NO: 3, the LCDR1 may comprise an amino acid sequence as set forth in SEQ ID NO: 4, the LCDR2 may comprise an amino acid sequence as set forth in SEQ ID NO: 5, and the LCDR3 may comprise an amino acid sequence as set forth in SEQ ID NO: 6.

[0179] In the present application, the isolated antigen-binding protein may comprise VH and VL, the VH may comprise an amino acid sequence as set forth in SEQ ID NO: 54, and the VL may comprise an amino acid sequence as set forth in SEQ ID NO: 59.

[0180] In the present application, the isolated antigen-binding protein may comprise VH and VL, the VH may comprise an amino acid sequence as set forth in any one of

SEQ ID NO: 20 to 25, and the VL may comprise an amino acid sequence as set forth in any one of SEQ ID NO: 36 to 39.

[0181] For example, the isolated antigen-binding protein may comprise VH and VL, the VH may comprise an amino acid sequence as set forth in SEQ ID NO: 20, and the VL may comprise an amino acid sequence as set forth in SEQ ID NO: 36.

[0182] For example, the isolated antigen-binding protein may comprise VH and VL, the VH may comprise an amino acid sequence as set forth in SEQ ID NO: 21, and the VL may comprise an amino acid sequence as set forth in SEQ ID NO: 37.

[0183] For example, the isolated antigen-binding protein may comprise VH and VL, the VH may comprise an amino acid sequence as set forth in SEQ ID NO: 22, and the VL may comprise an amino acid sequence as set forth in SEQ ID NO: 38.

[0184] For example, the isolated antigen-binding protein may comprise VH and VL, the VH may comprise an amino acid sequence as set forth in SEQ ID NO: 23, and the VL may comprise an amino acid sequence as set forth in SEQ ID NO: 37.

[0185] For example, the isolated antigen-binding protein may comprise VH and VL, the VH may comprise an amino acid sequence as set forth in SEQ ID NO: 24, and the VL may comprise an amino acid sequence as set forth in SEQ ID NO: 37.

[0186] For example, the isolated antigen-binding protein may comprise VH and VL, the VH may comprise an amino acid sequence as set forth in SEQ ID NO: 25, and the VL may comprise an amino acid sequence as set forth in SEQ ID NO: 39.

[0187] In the present application, the isolated antigen-binding protein may comprise a heavy chain constant region, which may be derived from IgG. For example, the heavy chain constant region may be derived from human IgG. the heavy chain constant region may be derived from IgG1, IgG2, IgG3 and IgG4. For example, the heavy chain constant region may be derived from human IgG1. Compared with natural human IgG1, the heavy chain constant region may be amino acid mutated. For example, the heavy chain constant region may be a sequence obtained by performing N297A amino acid point mutation on IgG1. For example, the heavy chain constant region may comprise an amino acid sequence as set forth in SEQ ID NO: 41.

[0188] In the present application, the isolated antigen-binding protein may comprise a heavy chain, and the heavy chain may comprise an amino acid sequence as set forth in any one of SEQ ID NO: 42 to 46.

[0189] In the present application, the isolated antigen-binding protein may comprise a light chain constant region, and the light chain constant region may be derived from a light chain λ and a light chain κ . For example, the light chain constant region may comprise an amino acid sequence as set forth in SEQ ID NO: 40.

[0190] In the present application, the isolated antigen-binding protein may comprise a light chain, and the light chain may comprise an amino acid sequence as set forth in any one of SEQ ID NO: 47 to 49.

[0191] For example, the isolated antigen-binding protein may comprise a heavy chain and a light chain, the heavy chain may comprise an amino acid sequence as set forth in

SEQ ID NO: 42, and the light chain may comprise an amino acid sequence as set forth in SEQ ID NO: 47.

[0192] For example, the isolated antigen-binding protein may comprise a heavy chain and a light chain, the heavy chain may comprise an amino acid sequence as set forth in SEQ ID NO: 43, and the light chain may comprise an amino acid sequence as set forth in SEQ ID NO: 48.

[0193] For example, the isolated antigen-binding protein may comprise a heavy chain and a light chain, the heavy chain may comprise an amino acid sequence as set forth in SEQ ID NO: 44, and the light chain may comprise an amino acid sequence as set forth in SEQ ID NO: 47.

[0194] For example, the isolated antigen-binding protein may comprise a heavy chain and a light chain, the heavy chain may comprise an amino acid sequence as set forth in SEQ ID NO: 45, and the light chain may comprise an amino acid sequence as set forth in SEQ ID NO: 47.

[0195] For example, the isolated antigen-binding protein may comprise a heavy chain and a light chain, the heavy chain may comprise an amino acid sequence as set forth in SEQ ID NO: 46, and the light chain may comprise an amino acid sequence as set forth in SEQ ID NO: 49.

[0196] In the present application, a part of the amino acid sequence of each heavy chain or light chain of the antigen-binding protein is homologous to the corresponding amino acid sequence in an antibody from a specific species, or belongs to a specific category. For example, the variable region and constant portion of light chain and heavy chain are both from the variable region and constant region of antibody of one animal species (such as human). In the present application, the homologue may be a protein and/or a polypeptide having at least about 85% (for example, at least about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or more) identity with the amino acid sequence of the protein and/or the polypeptide (for example, an antibody or fragment thereof that specifically binds to PD-L1 protein).

[0197] In the present application, the homology generally refers to similarity, likeness or association between two or more sequences. The alignment for determining the percentage of sequence homology can be realized in various ways known in the art, for example, using publicly available computer softwares such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) softwares. Those skilled in the art may determine appropriate parameters for alignment sequences, including any algorithm needed to achieve maximum alignment within the full-length sequence being compared or within the target sequence area. The homology may also be determined by the following methods: FASTA and BLAST. For the description of FASTA algorithm, please refer to W. R. Pearson and D. J. Lipman, "Improved Tools for Biological Sequence Comparison", *Proc. Natl. Acad. Sci.*, 85:2444-2448, 1988; and D. J. Lipman and W. R. Pearson, "Rapid and sensitive protein similarity searches", *Science*, 227:1435-1441, 1989. Descriptions of BLAST algorithm may be found in S. Altschul, W. Gish, W. Miller, E. W. Myers and D. Lipman, "Basic Local Alignment Search Tool", *Journal of Molecular Biology*, 215:403-410, 1990.

Test Method

[0198] The physical/chemical properties and/or biological activities of the PD-1 antigen-binding protein described in

the present application may be identified, screened or characterized by various assays known in the art.

[0199] In one aspect, for example, the antigen binding activity of the antigen-binding protein of the present application may be tested by known methods such as enzyme-linked immunosorbent assay (ELISA), western blot (e.g., protein blot), flow cytometry (e.g., FACS), immunohistochemistry, immunofluorescence or the like.

Nucleic Acid, Vector, Host Cell and Preparation Method

[0200] In another aspect, the present application also provides one or more isolated nucleic acid molecules. The one or more nucleic acid molecules may encode the antigen-binding protein described herein. For example, each nucleic acid molecule in the one or more nucleic acid molecules may encode the complete antigen-binding protein, or may encode a portion thereof (for example, one or more of HCDR1-3, LCDR1-3, VL, VH, a light chain or a heavy chain).

[0201] The nucleic acid molecule described in the present application may be isolated. For example, it may be produced or synthesized by the following methods: (i) amplification in vitro, for example, polymerase chain reaction (PCR) amplification, (ii) cloning and recombination, (iii) purification, for example, fractionation by enzyme digestion and gel electrophoresis, or (iv) synthesis, for example, chemical synthesis. In some embodiments, the isolated nucleic acid is a nucleic acid molecule prepared by recombinant DNA technology.

[0202] In the present application, the nucleic acid encoding the antibody and antigen-binding fragment thereof may be prepared by various methods known in the art, including but not limited to restriction fragment manipulation or overlapping extension PCR using synthetic oligonucleotides.

[0203] In another aspect, the present application provides one or more vectors comprising one or more nucleic acid molecules described herein. Each vector may comprise one or more of the nucleic acid molecules. In addition, the vector may also comprise other genes, such as marker genes that allow selection of the vector in appropriate host cells and under appropriate conditions. In addition, the vector may also comprise an expression control element that allows the coding region to be correctly expressed in an appropriate host. For example, the vector is an expression vector.

[0204] In another aspect, the present application provides a host cell, and the host cell may comprise one or more nucleic acid molecules and/or one or more vectors described herein. In some embodiments, each or every host cell may comprise one or more nucleic acid molecules or vectors described herein. In some embodiments, each or every host cell may comprise a plurality (e.g., two or more) or a variety (e.g., two or more kinds) of nucleic acid molecules or vectors described herein.

[0205] In another aspect, the present application provides a method for preparing the antibody or antigen-binding fragment thereof. The method may comprise culturing the host cell described in the present application under conditions such that the antibody or antigen-binding fragment thereof is expressed. For example, with suitable culture medium, suitable temperature and culture time, etc., these methods are known to those skilled in the art.

Pharmaceutical Composition, Method and Use

[0206] In another aspect, the present application provides a pharmaceutical composition. The pharmaceutical composition may comprise the antigen-binding protein, the polypeptide, the nucleic acid molecule, the vector, the host cell described herein, and optionally a pharmaceutically acceptable carrier. The pharmaceutically acceptable adjuvant is not toxic to the recipient at the dosage and concentration adopted, and the pharmaceutical composition in the present application may also comprise more than one active compounds, which are those active compounds with complementary activities and usually do not adversely affect each other. The form and effective amount of such pharmaceuticals depend on, for example, the amount and form of antagonists present in the preparation, and the clinical parameters of the subject.

[0207] The pharmaceutical composition may be used for inhibiting tumor growth. For example, the pharmaceutical composition of the present application may inhibit or delay the development or progression of a disease, reduce the tumor size (even basically eliminate the tumor), and/or alleviate and/or stabilize the disease conditions.

[0208] The pharmaceutical composition of the present application may comprise a prophylactically and/or therapeutically effective amount of the antibody, antigen-binding fragment thereof. The prophylactically and/or therapeutically effective amount is the dosage that is required to prevent and/or treat (at least partially treat) a disease or disorder and/or any complications thereof in a subject having or at risk of developing the disease or disorder.

[0209] In another aspect, the present application provides a use of the antigen-binding protein and/or the fusion protein in the manufacture of a medicament. The medicament is for use in treating cancer, inhibiting tumor growth and/or inhibiting tumor cell proliferation. In some embodiments, the tumor or cancer is a tumor or cancer with abnormal expression of PD-L1. In the present application, the tumor may include a tumor with a high expression of PD-1 or PD-L1. In the present application, the tumor may include a tumor with a high expression of PD-L1. In the present application, the tumor may include a solid tumor. In the present application, the tumor may include breast cancer, lung cancer, gastric cancer and urothelial carcinoma.

[0210] In another aspect, the present application provides a method for inhibiting the binding of PD-L1 to PD-1, which comprises administering the antigen-binding protein and/or the polypeptide described herein. For example, the method may be ex vivo or in vitro. For example, the method may be a method for non-therapeutic purposes. In some cases, the method may comprise contacting a biological sample with the antigen-binding protein and/or PD-1 described herein under conditions that allow the antigen-binding protein and/or PD-1 to bind to PD-L1, detecting whether a complex is formed between the antigen-binding protein and PD-L1, and detecting whether a complex is formed between PD-1 and PD-L1.

[0211] In another aspect, the present application provides a method for stimulating the secretion of cytokines in immune cells, which comprises administering the isolated antigen-binding protein and/or the polypeptide. The cytokine may be IL-2. The immune cells may be lymphocytes. For example, T lymphocytes. For example, the method may be ex vivo or in vitro. For example, the method may be a method for non-therapeutic purpose.

[0212] In another aspect, the present application provides a method for detecting the presence and/or content of PD-L1 protein, which comprises administering the isolated antigen-binding protein and/or the polypeptide. For example, the method can be ex vivo or in vitro. For example, the method may be a method for non-therapeutic purposes.

[0213] The present application also provides the use of antigen-binding protein in a method for diagnosing a subject suffering from a tumor or cancer, the method comprises: determining the presence or expression level of PD-L1 in a sample obtained from the subject by contacting the sample with the antigen-binding protein of the present application and detecting the presence of the bound antibody.

[0214] In another aspect, the present application provides a chimeric antigen receptor (CAR), which may comprise the nucleic acid molecule or antigen-binding protein described in the present application.

[0215] In another aspect, the present application provides an antibody drug conjugate, which may include the antigen-binding fragment.

[0216] In another aspect, the present application provides a kit, which may comprise the antigen-binding protein, the chimeric antigen receptor, the genetically modified cell, the antibody drug conjugate described in the present application, and/or the pharmaceutical composition described in the present application. It may comprise the antigen-binding protein, the chimeric antigen receptor, the genetically modified cell described in the present application, and/or the antibody drug conjugate described in the present application in a single common container, and may also be optionally in combination with one or more therapeutic agents, and optionally formulated together in a pharmaceutical compositions

[0217] In another aspect, the present application provides a drug delivery device, which may be used for administering the antigen-binding protein or a pharmaceutical composition thereof.

[0218] Without being bound by any theory, the following examples are only for explaining various embodiments of the invention, and they are not used to limit the scope of the invention.

EXAMPLE

Example 1. Preparation of Murine Anti-Human PD-L1 Monoclonal Antibody

[0219] Emulsification was carried out with Human PD-L1/B7-H1 Protein, Fc Tag (Acro Biosystem PD1-H5258) and PADRE, and then 6 to 8-week-old female SD rats were injected intraperitoneally with 50 µg Human PD-L1/B7-H1 Protein, Fc Tag/rat, 25 µg PADRE/rat; after that, booster immunization was carried out at intervals of two to three weeks. The rats were immunized three times in total, and two weeks after the last immunization, the tail blood of rats was taken to determine the titer of anti-human PDL1 antibody in serum. After 50 µg protein without Freund's adjuvant was injected intraperitoneally into the rats for ictus immunisatorius, the spleen cells of the rats were taken and fused with myeloma (Sp2/0) cells on the third day.

[0220] Clones that specifically bind to HEK293-PD-L1 cells and can simultaneously block the binding of receptor PD-1 to HEK293-hPD-L1 cells were screened from the fusion plate by flow cytometer and subcloning was carried out by limited dilution method, and finally a monoclonal that

can secrete specific antibodies was screened. Monoclonal antibodies were obtained by small-scale production and purification in serum-free medium for subsequent identification, including the detection of the binding ability of antibodies to HEK293-PD-L1, the function of blocking binding of PD-1 to HEK293-hPD-L1 cell, and the mixed lymphocyte experiment to finally obtain candidate anti-human PD-L1 monoclonal antibody 50G2-3. The sequences of the variable regions of murine hybridoma clone 50G2-3 obtained by monoclonal antibody sequencing is as follows:

```
>50G2-3 light chain variable region sequence
                                     (SEQ ID NO: 36)
DIQMTQSPPSLSASLGDQVTITCQASQNINNYIAWYQQKPGKAPRQLIRY
TSTLVSGTSPSRFSGSGSKDYFSISNVESEDIASYYCLQYDNPNTFGA
GTKLELK

>50G2-3 heavy chain variable region sequence
                                     (SEQ ID NO: 20)
EVQLVESGGGLVQPGRSLKSCVASGFTFSNFWMTWIRQAPGKLEWVAS
ITHSGGITYYPDSVKGRFTISRDDAKSTLYLQMNSLRSEDTATYYCSRDP
TEAPFDYWGGQVMVTVSS
```

TABLE 1

Heavy Chain and Light Chain CDR Sequences of Murine Hybridoma Clone 50G2-3			
Heavy Chain		Light Chain	
HCDR1	GFTFSNF (SEQ ID NO: 1)	LCDR1	QASQNINNYIA (SEQ ID NO: 4)
HCDR2	THSGGI (SEQ ID NO: 2)	LCDR2	YTSTLVS (SEQ ID NO: 5)
HCDR3	DPTEAPFDY (SEQ ID NO: 3)	LCDR3	LQYDNPNT (SEQ ID NO: 6)

Example 2. Humanization of Monoclonal Antibodies

[0221] The most homologous human Germline antibody (data source: IMGT) was selected as the humanized design framework (the framework of the light chain was based on IGKV1-33*01, IGKJ2*01, and the framework of the heavy chain was based on IGHV3-7*01, IGHJ4*01), and the variable regions of the light and heavy chains of the antibody were numbered by Chothia [Chothia & Lesk, 1987], and CDR regions of the antibody were defined as: CDRL1 (L24-L34), CDRL2 (L50-L56), CDRL3 (L89-L97), CDRH1 (H26-H32), CDRH2 (H52-H56) and CDRH3 (H95-H97), and humanized mutations of amino acids of light and heavy chain variable regions of antibodies were carried out according to sequence alignment and structural information of variable regions; expression vectors were designed, genes were synthesized, recombinant antibodies were expressed by mammalian cells and purified, the differences in activity and physical and chemical properties between humanized antibodies and chimeric antibodies were compared, and 1 to 2 rounds of humanization optimization were carried out.

[0222] Germline antibody sequence information:

```
IGKV1-33*01
                                     (SEQ ID NO: 60)
DIQMTQSPSSLSASVGDRTTITCQASQDISNYLNWYQQKPKAPKLLIYD
ASNLETGVPSPSRFSGSGSGTDFTFTISSLPEDIATYYCQQYDNLFP
IGKJ2*01
                                     (SEQ ID NO: 61)
YTFGGQGTKLEIK
IGHV3-7*01
                                     (SEQ ID NO: 62)
EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYWMWVRQAPGKLEWVAN
IKQDGSSEKYYVDSVKGRFTISRDNAKNSLYLQMNSLRADTAIVYCAR
IGHJ4*01
                                     (SEQ ID NO: 63)
YFDYWGGQTLVTVSS
```

[0223] The following optimized designs of light and heavy chains were humanized sequences transplanted based on the sequence of the above Germanline antibody as framework CDRs (1910G2HzL0 and 1910G2HzH0 are the light and heavy chains of the chimeric antibody, and 1910G2HzL2, 1910G2HzH2, 1910G2HzL3, 1910G2HzH3, 1910G2HzL4, 1910G2HzH4, 1910G2HzL7, 1910G2HzH7, 1910G2HzL9 and 1910G2HzH9 are the light and heavy chains of the humanized antibody).

```
1910G2HzL0 light chain variable region sequence
                                     (SEQ ID NO: 36)
DIQMTQSPPSLSASLGDQVTITCQASQNINNYIAWYQQKPGKAPRQLIRY
TSTLVSGTSPSRFSGSGSKDYFSISNVESEDIASYYCLQYDNPNTFGA
GTKLELK

1910G2HzL2 light chain variable region sequence
                                     (SEQ ID NO: 37)
DIQMTQSPSSLSASVGDRTTITCQASQNINNYIAWYQQKPGKAPQLIRY
TSTLVSGVPSRFRSGSGSKDYFTFTISSLPEDIATYYCLQYDNPNTFGQ
GTKLEIK

1910G2HzL3 light chain variable region sequence
                                     (SEQ ID NO: 38)
DIQMTQSPSSLSASVGDRTTITCQASQNINNYIAWYQQKPGKAPQLIRY
TSTLVSGTSPSRFSGSGSKDYFTFTISNLQPEDATYYCLQYDNPNTFGQ
GTKLEIK

1910G2HzL4 light chain variable region sequence
                                     (SEQ ID NO: 37)
Same as 1910G2HzL2

1910G2HzL7 light chain variable region sequence
                                     (SEQ ID NO: 37)
Same as 1910G2HzL2

1910G2HzL9 light chain variable region sequence
                                     (SEQ ID NO: 39)
DIQMTQSPSSLSASVGDRTTITCQASQNINNYIAWYQQKPGKAPQLIRY
TSTLVSGVPSRFRSGSGSKDYFTFTISSLPEDIATYYCLQYDNPNTFGQ
GTKLEIK
```

-continued

1910G2HzH0 heavy chain variable region sequence
(SEQ ID NO: 20)
EVQLVESGGGLVQPGRSLKSLCVASGFTFSNFWMTWIRQAPGKGLEWVAS

ITHSGGITYYPDSVKGRFTISRDDAKSTLYLQMNSLRSEDATYYCSRDP

TEAPFDYWQGVMVTSS

1910G2HzH2 heavy chain variable region sequence
(SEQ ID NO: 21)
EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFWMTWVRQAPGKGLEWVAS

ITHSGGITYYVDSVKGRFTISRDDAKSSLYLQMNSLRAEDTAVYYCSRDP

TEAPFDYWQGTLVTSS

1910G2HzH3 heavy chain variable region sequence
(SEQ ID NO: 22)
EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFWMSWVRQAPGKGLEWVAS

ITHSGGITYYVDSVKGRFTISRDNKSSLYLQMNSLRAEDTAVYYCSRDP

TEAPFDYWQGVMVTSS

1910G2HzH4 heavy chain variable region sequence
(SEQ ID NO: 23)
EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFWMTWVRQAPGKGLEWVAS

ITHSGGITYYPDSVKGRFTISRDDSKNTLYLQMNSLRAEDTAVYYCSRDP

TEAPFDYWQGTLVTSS

1910G2HzH7 heavy chain variable region sequence
(SEQ ID NO: 24)
EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFWMTWVRQAPGKGLEWVAS

ITHSGGITYYADSVKGRFTISRDDSKSTLYLQMNSLRAEDTAVYYCSRDP

TEAPFDYWQGTLVTSS

1910G2HzH9 heavy chain variable region sequence
(SEQ ID NO: 25)
EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFWMSWVRQAPGKGLEWVAS

ITHSGGITYYADSVKGRFTISRDDSKNTLYLQMNSLRAEDTAVYYCSRDP

TEAPFDYWQGTLVTSS

[0224] The light chains and heavy chains of the antibodies described above are combined to obtain antigen-binding proteins 1910G2HzL2H2, 1910G2HzL3H3, 1910G2HzL4H4, 1910G2HzL7H7 and 1910G2HzL9H9.

Example 3. Determination of Kinetic Parameters of the Antigen-Binding Protein of the Present Application

[0225] (1) The affinities of humanized antibodies 1910G2HzL2H2, 1910G2HzL3H3, 1910G2HzL4H4, 1910G2HzL7H7 and 1910G2HzL9H9 to human PD-L1 (Acro, Catalog Number: H5229) were determined by Octet RED96e (Fortebio), both antigens and antibodies were diluted by 1xPBST (1xPBS: Sangon, B548117-0500; 0.02% Tween 20: Sigma-Alorich, P1379), and the concentration of antigen for use was 100 nM, and the concentration of antibody for use was 50 nM.

[0226] (2) Detection of samples on machines (Octet Data Acquisition 11.1.0.11): First, the sample was added into a 96-well plate (Greiner bio-one, 655209) with a system of 200 μ L/well. Then the software parameters were set, the plate temperature was set to 30° C., and the frequency of collecting standard dynamic signals was 5.0 HZ. Next, the AHC sensor (Fortebio, Catalog Number: 18-0015) was

pre-wetted with 1xPBST for 10 minutes, and then the sample was detected on the instrument. Each cycle includes the following steps: 1) immersing in buffer for 60 seconds; 2) detecting whether the antigen nonspecifically binding to the sensor; 3) regeneration of 10 mM glycine solution with pH 1.7; 4) immersing in the buffer for 60 seconds; 5) immobilizing the antibody on the sensor for 20 seconds; 6) immersing the sensor in the buffer for 180 seconds; 7) the antigen binding to the antibody for 180 seconds; 8) dissociating the antigen from the antibody for 10 minutes; 9) regenerating the sensor.

(3) Data Analysis

[0227] The association rate (Ka) and dissociation rate (Kd) of the antigen-antibody at a ratio of 1:1 were measured by Data Analysis 11.0 software of Fortebio, and based on these, the equilibrium dissociation constant (KD) of antibodies were calculated. The results are as shown in the Table below, indicating that the affinities of humanized antibodies 1910G2HzL2H2, 1910G2HzL3H3, 1910G2HzL4H4, 1910G2HzL7H7 and 1910G2HzL9H9 to human PD-L1 are comparable to Atezolimumab, and even with lower KD values. Theses indicated that the antigen-binding proteins of the present application had higher binding affinities.

TABLE 2			
Affinity of Humanized Antibodies to Human PD-L1			
Humanized Antibody	KD (M)	kon (1/Ms)	kdis (1/s)
1910G2HzL2H2	2.435E-10	1.03E+06	2.50E-04
1910G2HzL3H3	4.522E-10	1.06E+06	4.77E-04
1910G2HzL4H4	1.601E-10	1.03E+06	1.65E-04
1910G2HzL7H7	3.019E-10	9.58E+05	2.89E-04
1910G2HzL9H9	2.137E-10	9.80E+05	2.10E-04
Atezolimumab	4.837E-10	6.44E+05	3.11E-04

Example 4. Binding Activity of the Antigen-Binding Protein of the Present Application to Antigen PD-L1 on the Cell Surface

[0228] CHOK1-cynoPD-L1 or HEK293-hPD-L1 were collected with 5E6 cells in each 96-well plate, centrifuged at 300 g for 5 minutes, resuspended with pre-cooled FACS buffer, evenly spread on a 3799 U-shaped plate by 100 μ L/well, and blocked at room temperature for 15 minutes. Humanized antibodies were taken, started at an initial concentration of 3.33 μ g/mL, and then diluted three-fold to establish 12 gradients. After blocking and centrifugation at 2000 rpm for 5 minutes, and the supernatant was removed. Diluted antibodies were added by 100 μ L/well, and the mixture was mixed evenly, and incubated at 4° C. for 1 hour. The mixture was centrifuged at 2000 rpm for 5 minutes, the supernatant was removed, and the residue was washed with FACS Buffer for 1-2 times. A secondary antibody goat-anti-human488, 1:1000 was added by 100 μ L/well, and incubated at 4° C. for 1 hour, centrifuged at 2000 rpm for 5 minutes, then the supernatant was removed, and the residue was washed with FACS Buffer for 1-2 times, and then FACS buffer was added by 30 μ L/well for resuspension of the cells and detection on the instrument.

[0229] FIG. 1 showed that the antigen-binding protein described in the present application binds to the human PD-L1 protein on the HEK293 cell. FIG. 2 showed that the

antigen-binding protein described in the present application binds to the monkey PD-L1 protein on the CHOK1 cell. Results were as shown in FIG. 1 and FIG. 2, and the binding activities of antigen-binding proteins 1910G2HzL2H2, 1910G2HzL3H3, 1910G2HzL4H4, 1910G2HzL7H7 and 1910G2HzL9H9 of the present application to antigen PD-L1 on the cell surface were comparable to or even higher than Atezolimumab. It was indicated that the binding activities of the antigen-binding proteins of the present application to PD-L1 were excellent.

Example 5. The Antigen-Binding Protein of the Present Application Blocks the Binding Activity of Antigen PD-L1 on the Cell Surface to PD-1

[0230] HEK293-hPD-L1 or CHOK1-cynoPD-L1 were collected by 5E6 cells in each 96-well plate, centrifuged at 300 g for 5 minutes, resuspended with pre-cooled FACS buffer, evenly spread on a 3799 U-shaped plate by 100 μ L/well, and blocked at room temperature for 15 minutes. Humanized antibodies were taken and started at an initial concentration of 10 μ g/mL, and then diluted three-fold to obtain 12 gradients. Biotin-PD-1-mFc of 0.224 mg/mL was prepared into 2 μ g/mL. After blocking and centrifugation at 2000 rpm for 5 minutes, and the supernatant was removed. Diluted antibodies were added by 50 μ L/well, the mixture was mixed evenly, and 2 μ g/mL of Biotin-PD-1-mFc was added by 50 μ L/well, mixed evenly, and incubated at 4° C. for 1 hour. After a centrifugation at 2000 rpm for 5 minutes, the supernatant was removed, and the residue was washed with FACS Buffer for 1-2 times. A secondary antibody SA488,1:1000 was added by 100 μ L/well, and incubated at 4° C. for 1 hour, centrifuged at 2000 rpm for 5 minutes, then the supernatant was removed. The residue was washed with FACS Buffer for 1-2 times, and then FACS buffer was added by 30 μ L/well for resuspension of the cells and detection on the instrument. FIG. 3 showed that the antigen-binding protein described in the present application blocked the binding of PD-1 to the human PD-L1 protein on the HEK293 cell. FIG. 4 showed that the antigen-binding protein described in the present application blocked the binding of PD-1 to the monkey PD-L1 protein on the CHOK1 cell. Results were as shown in FIG. 3 and FIG. 4, and the activities of antigen binding 3 proteins 1910G2HzL2H2, 1910G2HzL3H3, 1910G2HzL4H4, 1910G2HzL7H7 and 1910G2HzL9H9 of the present application to block the binding of antigen PD-L1 on the cell surface to PD-1 were comparable to Atezolimumab. It was indicated that the

activities of the antigen-binding proteins of the present application to block the binding of PD-L1 to PD-1 were excellent.

Example 6. Mixed Lymphocyte Reaction: Secretion of Cytokine IL-2

[0231] Human dendritic DC cells were resuscitated with cell culture medium (1640+2% FBS), the density of DC cells was adjusted to 1×10^5 to 1×10^7 cells/mL, then mitomycin C was added with a final concentration of 50 μ g/mL, treating at 37 degrees in the dark for 30 minutes, then 10 mL culture medium was added for termination; the mixture was centrifuged at 400 g for 10 minutes, and then washed with 10 mL culture medium. Gradient dilution of anti-PD1 antibody: the greatest final concentration of antibody was 2.5 μ g/mL (the preparation concentration was 10 μ g/mL), and the gradient dilution was ten-fold (5 concentration points+ one 0 concentration), and then 50 μ L of prepared anti-PD1 antibody was added into the corresponding cell culture plate (Corning, Catalog Number: 3599). DC cells treated with human peripheral blood lymphocytes PBMC and mitomycin C were collected, the density of DC cells was adjusted to 2×10^5 cells/mL, and then the cells were added into the culture plate by 50 μ L/well, i.e., the number of DC cells per well was 1×10^4 cells/well; the density of PBMC cells was adjusted to 2×10^6 cells/mL, and then the cells were added into the culture plate by 100 μ L/well, i.e., the number of PBMC cells per well was 2×10^5 cells/well. The cell culture plate was placed in a cell incubator with 5% carbon dioxide at 37° C. for 5 days. After 5 days, the supernatant was collected following a centrifugation with 300 g for 5 minutes, and the content of IL-2 was detected with Human IL-2 ELISA kit (BD, Catalog Number: 550611). The detection method was strictly performed in accordance with the kit instructions, and the data was processed with GraphPad Prism software. The results were as shown in FIG. 5, indicating all of the antigen-binding proteins 1910G2HzL2H2, 1910G2HzL3H3, 1910G2HzL4H4, 1910G2HzL7H7 and 1910G2HzL9H9 of the present application can stimulate the lymphocytes to secrete cytokines, with better effects than Atezolimumab. It was indicated that the antigen-binding proteins of the present application had high ability to stimulate immune cells to secrete cytokines. [0232] The foregoing detailed description is provided by way of explanation and example, and is not intended to limit the scope of the appended claims. Many variations of the embodiments enumerated in the present invention are obvious to those skilled in the art, and they are within the scope of the appended claims and their equivalents.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 63

<210> SEQ ID NO 1

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: HCDR1

<400> SEQUENCE: 1

Gly Phe Thr Phe Ser Asn Phe

1

5

-continued

<210> SEQ ID NO 2
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: HCDR2

<400> SEQUENCE: 2

Thr His Ser Gly Gly Ile
1 5

<210> SEQ ID NO 3
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: HCDR3

<400> SEQUENCE: 3

Asp Pro Thr Glu Ala Pro Phe Asp Tyr
1 5

<210> SEQ ID NO 4
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: LCDR1

<400> SEQUENCE: 4

Gln Ala Ser Gln Asn Ile Asn Asn Tyr Ile Ala
1 5 10

<210> SEQ ID NO 5
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: LCDR2

<400> SEQUENCE: 5

Tyr Thr Ser Thr Leu Val Ser
1 5

<210> SEQ ID NO 6
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: LCDR3

<400> SEQUENCE: 6

Leu Gln Tyr Asp Asn Val Pro Asn Thr
1 5

<210> SEQ ID NO 7
<211> LENGTH: 25
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 50G2-3 H-FR1

<400> SEQUENCE: 7

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Arg

-continued

1	5	10	15
---	---	----	----

Ser Leu Lys Leu Ser Cys Val Ala Ser
20 25

<210> SEQ ID NO 8
<211> LENGTH: 25
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 1910G2HzH2, 1910G2HzH3, 1910G2HzH4, 1910G2HzH7
and 1910G2HzH9 H-FR1

<400> SEQUENCE: 8

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser
20 25

<210> SEQ ID NO 9
<211> LENGTH: 19
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 50G2-3 H-FR2

<400> SEQUENCE: 9

Trp Met Thr Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
1 5 10 15

Ala Ser Ile

<210> SEQ ID NO 10
<211> LENGTH: 19
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 1910G2HzH2, 1910G2HzH4 and 1910G2HzH7 H-FR2

<400> SEQUENCE: 10

Trp Met Thr Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
1 5 10 15

Ala Ser Ile

<210> SEQ ID NO 11
<211> LENGTH: 19
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 1910G2HzH3 and 1910G2HzH9 H-FR2

<400> SEQUENCE: 11

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
1 5 10 15

Ala Ser Ile

<210> SEQ ID NO 12
<211> LENGTH: 41
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 50G2-3 H-FR3

<400> SEQUENCE: 12

-continued

Thr Tyr Tyr Pro Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp
1 5 10 15

Asp Ala Lys Ser Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ser Glu
20 25 30

Asp Thr Ala Thr Tyr Tyr Cys Ser Arg
35 40

<210> SEQ ID NO 13
<211> LENGTH: 41
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 1910G2HzH2 H-FR3

<400> SEQUENCE: 13

Thr Tyr Tyr Val Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp
1 5 10 15

Asp Ala Lys Ser Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu
20 25 30

Asp Thr Ala Val Tyr Tyr Cys Ser Arg
35 40

<210> SEQ ID NO 14
<211> LENGTH: 41
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 1910G2HzH3 H-FR3

<400> SEQUENCE: 14

Thr Tyr Tyr Val Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp
1 5 10 15

Asn Ala Lys Ser Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu
20 25 30

Asp Thr Ala Val Tyr Tyr Cys Ser Arg
35 40

<210> SEQ ID NO 15
<211> LENGTH: 41
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 1910G2HzH4 H-FR3

<400> SEQUENCE: 15

Thr Tyr Tyr Pro Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp
1 5 10 15

Asp Ser Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu
20 25 30

Asp Thr Ala Val Tyr Tyr Cys Ser Arg
35 40

<210> SEQ ID NO 16
<211> LENGTH: 41
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 1910G2HzH7 H-FR3

<400> SEQUENCE: 16

-continued

```

Thr Tyr Tyr Ala Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp
1           5           10           15
Asp Ser Lys Ser Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu
           20           25           30
Asp Thr Ala Val Tyr Tyr Cys Ser Arg
           35           40

```

```

<210> SEQ ID NO 17
<211> LENGTH: 41
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 1910G2HzH9 H-FR3

```

```

<400> SEQUENCE: 17

```

```

Thr Tyr Tyr Ala Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp
1           5           10           15
Asp Ser Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu
           20           25           30
Asp Thr Ala Val Tyr Tyr Cys Ser Arg
           35           40

```

```

<210> SEQ ID NO 18
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 50G2-3 and 1910G2HzH3 H-FR4

```

```

<400> SEQUENCE: 18

```

```

Trp Gly Gln Gly Val Met Val Thr Val Ser Ser
1           5           10

```

```

<210> SEQ ID NO 19
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 1910G2HzH2, 1910G2HzH4, 1910G2HzH7 and
1910G2HzH9 H-FR4

```

```

<400> SEQUENCE: 19

```

```

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
1           5           10

```

```

<210> SEQ ID NO 20
<211> LENGTH: 118
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 50G2-3 VH

```

```

<400> SEQUENCE: 20

```

```

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Arg
1           5           10           15
Ser Leu Lys Leu Ser Cys Val Ala Ser Gly Phe Thr Phe Ser Asn Phe
           20           25           30
Trp Met Thr Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
           35           40           45
Ala Ser Ile Thr His Ser Gly Gly Ile Thr Tyr Tyr Pro Asp Ser Val

```

-continued

50	55	60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ala Lys Ser Thr Leu Tyr		
65	70	75 80
Leu Gln Met Asn Ser Leu Arg Ser Glu Asp Thr Ala Thr Tyr Tyr Cys		
	85	90 95
Ser Arg Asp Pro Thr Glu Ala Pro Phe Asp Tyr Trp Gly Gln Gly Val		
	100	105 110
Met Val Thr Val Ser Ser		
	115	

<210> SEQ ID NO 21
 <211> LENGTH: 118
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: 1910G2HzH2 VH

<400> SEQUENCE: 21

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly	
1	5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Phe	
	20 25 30
Trp Met Thr Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val	
	35 40 45
Ala Ser Ile Thr His Ser Gly Gly Ile Thr Tyr Tyr Val Asp Ser Val	
	50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ala Lys Ser Ser Leu Tyr	
65	70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys	
	85 90 95
Ser Arg Asp Pro Thr Glu Ala Pro Phe Asp Tyr Trp Gly Gln Gly Thr	
	100 105 110
Leu Val Thr Val Ser Ser	
	115

<210> SEQ ID NO 22
 <211> LENGTH: 118
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: 1910G2HzH3 VH

<400> SEQUENCE: 22

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly	
1	5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Phe	
	20 25 30
Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val	
	35 40 45
Ala Ser Ile Thr His Ser Gly Gly Ile Thr Tyr Tyr Val Asp Ser Val	
	50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Ser Ser Leu Tyr	
65	70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys	
	85 90 95

-continued

Ser	Arg	Asp	Pro	Thr	Glu	Ala	Pro	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Val
			100					105					110		

Met	Val	Thr	Val	Ser	Ser
					115

<210> SEQ ID NO 23
 <211> LENGTH: 118
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: 1910G2HzH4 VH

<400> SEQUENCE: 23

Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly
1				5					10					15	

Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Asn	Phe
			20					25					30		

Trp	Met	Thr	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val
		35					40					45			

Ala	Ser	Ile	Thr	His	Ser	Gly	Gly	Ile	Thr	Tyr	Tyr	Pro	Asp	Ser	Val
	50					55					60				

Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asp	Ser	Lys	Asn	Thr	Leu	Tyr
65					70					75				80	

Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
			85						90					95	

Ser	Arg	Asp	Pro	Thr	Glu	Ala	Pro	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Thr
			100					105					110		

Leu	Val	Thr	Val	Ser	Ser
					115

<210> SEQ ID NO 24
 <211> LENGTH: 118
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: 1910G2HzH7 VH

<400> SEQUENCE: 24

Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly
1				5					10					15	

Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Asn	Phe
			20					25					30		

Trp	Met	Thr	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val
		35					40					45			

Ala	Ser	Ile	Thr	His	Ser	Gly	Gly	Ile	Thr	Tyr	Tyr	Ala	Asp	Ser	Val
	50					55					60				

Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asp	Ser	Lys	Ser	Thr	Leu	Tyr
65					70					75				80	

Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
			85						90					95	

Ser	Arg	Asp	Pro	Thr	Glu	Ala	Pro	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Thr
			100					105					110		

Leu	Val	Thr	Val	Ser	Ser
					115

<210> SEQ ID NO 25

-continued

<211> LENGTH: 118
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 1910G2HzH9 VH

<400> SEQUENCE: 25

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Phe
20 25 30
Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45
Ala Ser Ile Thr His Ser Gly Gly Ile Thr Tyr Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Thr Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ser Arg Asp Pro Thr Glu Ala Pro Phe Asp Tyr Trp Gly Gln Gly Thr
100 105 110
Leu Val Thr Val Ser Ser
115

<210> SEQ ID NO 26
<211> LENGTH: 23
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 50G2-3 L-FR1

<400> SEQUENCE: 26

Asp Ile Gln Met Thr Gln Ser Pro Pro Ser Leu Ser Ala Ser Leu Gly
1 5 10 15
Asp Gln Val Thr Ile Thr Cys
20

<210> SEQ ID NO 27
<211> LENGTH: 23
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 1910G2HzL2, 1910G2HzL3 and 1910G2HzL9 L-FR1

<400> SEQUENCE: 27

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15
Asp Arg Val Thr Ile Thr Cys
20

<210> SEQ ID NO 28
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 50G2-3 L-FR2

<400> SEQUENCE: 28

Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Arg Gln Leu Ile Arg
1 5 10 15

-continued

<210> SEQ ID NO 29
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 1910G2HzL2, 1910G2HzL3 and 1910G2HzL9 L-FR2

<400> SEQUENCE: 29

Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Gln Leu Ile Arg
1 5 10 15

<210> SEQ ID NO 30
<211> LENGTH: 32
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 50G2-3 L-FR3

<400> SEQUENCE: 30

Gly Thr Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Lys Asp Tyr Ser
1 5 10 15

Phe Ser Ile Ser Asn Val Glu Ser Glu Asp Ile Ala Ser Tyr Tyr Cys
20 25 30

<210> SEQ ID NO 31
<211> LENGTH: 32
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 1910G2HzL2 L-FR3

<400> SEQUENCE: 31

Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Lys Asp Tyr Thr
1 5 10 15

Phe Thr Ile Ser Ser Leu Gln Pro Glu Asp Ile Ala Thr Tyr Tyr Cys
20 25 30

<210> SEQ ID NO 32
<211> LENGTH: 32
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 1910G2HzL3 L-FR3

<400> SEQUENCE: 32

Gly Thr Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Lys Asp Tyr Thr
1 5 10 15

Phe Thr Ile Ser Ser Leu Gln Pro Glu Asp Ile Ala Thr Tyr Tyr Cys
20 25 30

<210> SEQ ID NO 33
<211> LENGTH: 32
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 1910G2HzL9 L-FR3

<400> SEQUENCE: 33

Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Lys Asp Phe Thr
1 5 10 15

Phe Thr Ile Ser Ser Leu Gln Pro Glu Asp Ile Ala Thr Tyr Tyr Cys

-continued

20	25	30
----	----	----

<210> SEQ ID NO 34
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 50G2-3 L-FR4

<400> SEQUENCE: 34

Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
1 5 10

<210> SEQ ID NO 35
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 1910G2HzL2, 1910G2HzL3 and 1910G2HzL9 L-FR4

<400> SEQUENCE: 35

Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
1 5 10

<210> SEQ ID NO 36
<211> LENGTH: 107
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 50G2-3 VL

<400> SEQUENCE: 36

Asp Ile Gln Met Thr Gln Ser Pro Pro Ser Leu Ser Ala Ser Leu Gly
1 5 10 15

Asp Gln Val Thr Ile Thr Cys Gln Ala Ser Gln Asn Ile Asn Asn Tyr
20 25 30

Ile Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Arg Gln Leu Ile
35 40 45

Arg Tyr Thr Ser Thr Leu Val Ser Gly Thr Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Lys Asp Tyr Ser Phe Ser Ile Ser Asn Val Glu Ser
65 70 75 80

Glu Asp Ile Ala Ser Tyr Tyr Cys Leu Gln Tyr Asp Asn Val Pro Asn
85 90 95

Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
100 105

<210> SEQ ID NO 37
<211> LENGTH: 107
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 1910G2HzL2 VL

<400> SEQUENCE: 37

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Gln Ala Ser Gln Asn Ile Asn Asn Tyr
20 25 30

Ile Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Gln Leu Ile

-continued

35	40	45
Arg Tyr Thr Ser Thr Leu Val Ser Gly Val Pro Ser Arg Phe Ser Gly		
50	55	60
Ser Gly Ser Gly Lys Asp Tyr Thr Phe Thr Ile Ser Ser Leu Gln Pro		
65	70	75 80
Glu Asp Ile Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp Asn Val Pro Asn		
85	90	95
Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys		
100	105	

<210> SEQ ID NO 38
 <211> LENGTH: 107
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: 1910G2HzL3 VL

<400> SEQUENCE: 38

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15
Asp Arg Val Thr Ile Thr Cys Gln Ala Ser Gln Asn Ile Asn Asn Tyr
20 25 30
Ile Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Gln Leu Ile
35 40 45
Arg Tyr Thr Ser Thr Leu Val Ser Gly Thr Pro Ser Arg Phe Ser Gly
50 55 60
Ser Gly Ser Gly Lys Asp Tyr Thr Phe Thr Ile Ser Asn Leu Gln Pro
65 70 75 80
Glu Asp Ile Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp Asn Val Pro Asn
85 90 95
Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
100 105

<210> SEQ ID NO 39
 <211> LENGTH: 107
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: 1910G2HzL9 VL

<400> SEQUENCE: 39

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15
Asp Arg Val Thr Ile Thr Cys Gln Ala Ser Gln Asn Ile Asn Asn Tyr
20 25 30
Ile Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Gln Leu Ile
35 40 45
Arg Tyr Thr Ser Thr Leu Val Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60
Ser Gly Ser Gly Lys Asp Phe Thr Phe Thr Ile Ser Ser Leu Gln Pro
65 70 75 80
Glu Asp Ile Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp Asn Val Pro Asn
85 90 95
Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
100 105

-continued

<210> SEQ ID NO 40
 <211> LENGTH: 107
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Light Chain Constant Region

<400> SEQUENCE: 40

Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
 1 5 10 15
 Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
 20 25 30
 Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
 35 40 45
 Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
 50 55 60
 Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
 65 70 75 80
 Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
 85 90 95
 Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 100 105

<210> SEQ ID NO 41
 <211> LENGTH: 330
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Heavy Chain Constant Region

<400> SEQUENCE: 41

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
 1 5 10 15
 Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 20 25 30
 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 35 40 45
 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 50 55 60
 Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
 65 70 75 80
 Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
 85 90 95
 Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
 100 105 110
 Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
 115 120 125
 Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
 130 135 140
 Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
 145 150 155 160
 Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
 165 170 175
 Glu Gln Tyr Ala Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
 180 185 190

-continued

His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
 195 200 205
 Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
 210 215 220
 Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu
 225 230 235 240
 Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
 245 250 255
 Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
 260 265 270
 Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
 275 280 285
 Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
 290 295 300
 Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
 305 310 315 320
 Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 325 330

<210> SEQ ID NO 42
 <211> LENGTH: 448
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: 1910G2HzH2 Heavy Chain

<400> SEQUENCE: 42

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Phe
 20 25 30
 Trp Met Thr Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ala Ser Ile Thr His Ser Gly Gly Ile Thr Tyr Tyr Val Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ala Lys Ser Ser Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ser Arg Asp Pro Thr Glu Ala Pro Phe Asp Tyr Trp Gly Gln Gly Thr
 100 105 110
 Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro
 115 120 125
 Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly
 130 135 140
 Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn
 145 150 155 160
 Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln
 165 170 175
 Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser
 180 185 190
 Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser
 195 200 205

-continued

```

Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr
210                215                220

His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser
225                230                235                240

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg
245                250                255

Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro
260                265                270

Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala
275                280                285

Lys Thr Lys Pro Arg Glu Glu Gln Tyr Ala Ser Thr Tyr Arg Val Val
290                295                300

Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr
305                310                315                320

Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr
325                330                335

Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu
340                345                350

Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys
355                360                365

Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser
370                375                380

Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp
385                390                395                400

Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser
405                410                415

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala
420                425                430

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
435                440                445

```

```

<210> SEQ ID NO 43
<211> LENGTH: 448
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 1910G2HzH3 Heavy Chain

```

```

<400> SEQUENCE: 43

```

```

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1          5          10          15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Phe
20        25        30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35        40        45

Ala Ser Ile Thr His Ser Gly Gly Ile Thr Tyr Tyr Val Asp Ser Val
50        55        60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Ser Ser Leu Tyr
65        70        75        80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85        90        95

Ser Arg Asp Pro Thr Glu Ala Pro Phe Asp Tyr Trp Gly Gln Gly Val
100       105       110

```


<400> SEQUENCE: 44

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

-continued

Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Asn	Phe	
			20				25						30			
Trp	Met	Thr	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	
			35				40				45					
Ala	Ser	Ile	Thr	His	Ser	Gly	Gly	Ile	Thr	Tyr	Tyr	Pro	Asp	Ser	Val	
			50				55				60					
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asp	Ser	Lys	Asn	Thr	Leu	Tyr	
			65				70				75			80		
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	
			85						90			95				
Ser	Arg	Asp	Pro	Thr	Glu	Ala	Pro	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	
			100						105			110				
Leu	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	
			115						120			125				
Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	
			130						135			140				
Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	
			145						150			155			160	
Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	
			165						170			175				
Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	
			180						185			190				
Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	
			195						200			205				
Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	
			210						215			220				
His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	
			225						230			235			240	
Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	
			245						250			255				
Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	
			260						265			270				
Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	
			275						280			285				
Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Ala	Ser	Thr	Tyr	Arg	Val	Val	
			290						295			300				
Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	
			305						310			315			320	
Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	
			325						330			335				
Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	
			340						345			350				
Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	
			355						360			365				
Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	
			370						375			380				
Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	
			385						390			395			400	
Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	
			405						410			415				

-continued

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala
 420 425 430

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 435 440 445

<210> SEQ ID NO 45
 <211> LENGTH: 448
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: 1910G2HzH7 Heavy Chain

<400> SEQUENCE: 45

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Phe
 20 25 30

Trp Met Thr Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ala Ser Ile Thr His Ser Gly Gly Ile Thr Tyr Tyr Ala Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Ser Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ser Arg Asp Pro Thr Glu Ala Pro Phe Asp Tyr Trp Gly Gln Gly Thr
 100 105 110

Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro
 115 120 125

Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly
 130 135 140

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn
 145 150 155 160

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln
 165 170 175

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser
 180 185 190

Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser
 195 200 205

Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr
 210 215 220

His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser
 225 230 235 240

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg
 245 250 255

Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro
 260 265 270

Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala
 275 280 285

Lys Thr Lys Pro Arg Glu Glu Gln Tyr Ala Ser Thr Tyr Arg Val Val
 290 295 300

Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr
 305 310 315 320

-continued

Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr
 325 330 335
 Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu
 340 345 350
 Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys
 355 360 365
 Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser
 370 375 380
 Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp
 385 390 395 400
 Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser
 405 410 415
 Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala
 420 425 430
 Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 435 440 445

 <210> SEQ ID NO 46
 <211> LENGTH: 448
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: 1910G2HzH9 Heavy Chain

 <400> SEQUENCE: 46

 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Phe
 20 25 30
 Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ala Ser Ile Thr His Ser Gly Gly Ile Thr Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ser Arg Asp Pro Thr Glu Ala Pro Phe Asp Tyr Trp Gly Gln Gly Thr
 100 105 110
 Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro
 115 120 125
 Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly
 130 135 140
 Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn
 145 150 155 160
 Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln
 165 170 175
 Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser
 180 185 190
 Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser
 195 200 205
 Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr
 210 215 220

-continued

His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	225	230	235	240
Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	245	250	255	
Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	260	265	270	
Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	275	280	285	
Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Ala	Ser	Thr	Tyr	Arg	Val	Val	290	295	300	
Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	305	310	315	320
Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	325	330	335	
Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	340	345	350	
Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	355	360	365	
Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	370	375	380	
Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	385	390	395	400
Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	405	410	415	
Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	420	425	430	
Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys	435	440	445	

<210> SEQ ID NO 47
 <211> LENGTH: 214
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: 1910G2HzL2 Light Chain

<400> SEQUENCE: 47

Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly	1	5	10	15
Asp	Arg	Val	Thr	Ile	Thr	Cys	Gln	Ala	Ser	Gln	Asn	Ile	Asn	Asn	Tyr	20	25	30	
Ile	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Gln	Leu	Ile	35	40	45	
Arg	Tyr	Thr	Ser	Thr	Leu	Val	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	50	55	60	
Ser	Gly	Ser	Gly	Lys	Asp	Tyr	Thr	Phe	Thr	Ile	Ser	Ser	Leu	Gln	Pro	65	70	75	80
Glu	Asp	Ile	Ala	Thr	Tyr	Tyr	Cys	Leu	Gln	Tyr	Asp	Asn	Val	Pro	Asn	85	90	95	
Thr	Phe	Gly	Gln	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Arg	Thr	Val	Ala	Ala	100	105	110	
Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	Asp	Glu	Gln	Leu	Lys	Ser	Gly	115	120	125	

-continued

Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
 130 135 140

Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
 145 150 155 160

Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
 165 170 175

Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
 180 185 190

Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
 195 200 205

Phe Asn Arg Gly Glu Cys
 210

<210> SEQ ID NO 48
 <211> LENGTH: 214
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: 1910G2HzL3 Light Chain

<400> SEQUENCE: 48

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15

Asp Arg Val Thr Ile Thr Cys Gln Ala Ser Gln Asn Ile Asn Asn Tyr
 20 25 30

Ile Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Gln Leu Ile
 35 40 45

Arg Tyr Thr Ser Thr Leu Val Ser Gly Thr Pro Ser Arg Phe Ser Gly
 50 55 60

Ser Gly Ser Gly Lys Asp Tyr Thr Phe Thr Ile Ser Asn Leu Gln Pro
 65 70 75 80

Glu Asp Ile Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp Asn Val Pro Asn
 85 90 95

Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala
 100 105 110

Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
 115 120 125

Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
 130 135 140

Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
 145 150 155 160

Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
 165 170 175

Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
 180 185 190

Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
 195 200 205

Phe Asn Arg Gly Glu Cys
 210

<210> SEQ ID NO 49
 <211> LENGTH: 214
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:

-continued

<223> OTHER INFORMATION: 1910G2HzL9 Light Chain

<400> SEQUENCE: 49

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15

Asp Arg Val Thr Ile Thr Cys Gln Ala Ser Gln Asn Ile Asn Asn Tyr
 20 25 30

Ile Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Gln Leu Ile
 35 40 45

Arg Tyr Thr Ser Thr Leu Val Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60

Ser Gly Ser Gly Lys Asp Phe Thr Phe Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80

Glu Asp Ile Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp Asn Val Pro Asn
 85 90 95

Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala
 100 105 110

Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
 115 120 125

Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
 130 135 140

Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
 145 150 155 160

Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
 165 170 175

Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
 180 185 190

Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
 195 200 205

Phe Asn Arg Gly Glu Cys
 210

<210> SEQ ID NO 50

<211> LENGTH: 25

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: 50G2-3 H-FR1; 1910G2HzH2, 1910G2HzH3,
 1910G2HzH4, 1910G2HzH7 and 1910G2HzH9 H-FR1

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (16)..(16)

<223> OTHER INFORMATION: Xaa = Gly or Arg

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (19)..(19)

<223> OTHER INFORMATION: Xaa = Lys or Arg

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (23)..(23)

<223> OTHER INFORMATION: Xaa = Ala or Val

<400> SEQUENCE: 50

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Xaa
 1 5 10 15

Ser Leu Xaa Leu Ser Cys Xaa Ala Ser
 20 25

-continued

<210> SEQ ID NO 51
<211> LENGTH: 19
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 50G2-3 H-FR2; 1910G2HzH2, 1910G2HzH4 and
1910G2HzH7 H-FR2; 1910G2HzH3 and 1910G2HzH9 H-FR2
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa = Ser or Thr
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa = Ile or Val

<400> SEQUENCE: 51

Trp Met Xaa Trp Xaa Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
1 5 10 15

Ala Ser Ile

<210> SEQ ID NO 52
<211> LENGTH: 41
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 50G2-3 H-FR3; 1910G2HzH2 H-FR3; 1910G2HzH3
H-FR3; 1910G2HzH4 H-FR3; 1910G2HzH7 H-FR3; 1910G2HzH9 H-FR3
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa = Ala, Pro or Val
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (17)..(17)
<223> OTHER INFORMATION: Xaa = Asp or Asn
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (18)..(18)
<223> OTHER INFORMATION: Xaa = Ala or Ser
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: Xaa = Asn or Ser
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (21)..(21)
<223> OTHER INFORMATION: Xaa = Ser or Thr
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (31)..(31)
<223> OTHER INFORMATION: Xaa = Ala or Ser
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (36)..(36)
<223> OTHER INFORMATION: Xaa = Thr or Val

<400> SEQUENCE: 52

Thr Tyr Tyr Xaa Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp
1 5 10 15

Xaa Xaa Lys Xaa Xaa Leu Tyr Leu Gln Met Asn Ser Leu Arg Xaa Glu
20 25 30

Asp Thr Ala Xaa Tyr Tyr Cys Ser Arg
35 40

<210> SEQ ID NO 53
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: 50G2-3 and 1910G2HzH3 H-FR4; 1910G2HzH2,
1910G2HzH4, 1910G2HzH7 and 1910G2HzH9 H-FR4
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa = Thr or Val
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Xaa = Leu or Met

<400> SEQUENCE: 53

Trp Gly Gln Gly Xaa Xaa Val Thr Val Ser Ser
1 5 10

<210> SEQ ID NO 54
<211> LENGTH: 118
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 50G2-3 VH; 1910G2HzH2 VH; 1910G2HzH3 VH;
1910G2HzH4 VH; 1910G2HzH7 VH; 1910G2HzH9 VH
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (16)..(16)
<223> OTHER INFORMATION: Xaa = Gly or Arg
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (19)..(19)
<223> OTHER INFORMATION: Xaa = Lys or Arg
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (23)..(23)
<223> OTHER INFORMATION: Xaa = Ala or Val
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (35)..(35)
<223> OTHER INFORMATION: Xaa = Ser or Thr
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (37)..(37)
<223> OTHER INFORMATION: Xaa = Ile or Val
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (61)..(61)
<223> OTHER INFORMATION: Xaa = Ala, Pro or Val
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (74)..(74)
<223> OTHER INFORMATION: Xaa = Asp or Asn
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (75)..(75)
<223> OTHER INFORMATION: Xaa = Ala or Ser
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (77)..(77)
<223> OTHER INFORMATION: Xaa = Asn or Ser
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (78)..(78)
<223> OTHER INFORMATION: Xaa = Ser or Thr
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (88)..(88)
<223> OTHER INFORMATION: Xaa = Ala or Ser
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (93)..(93)
<223> OTHER INFORMATION: Xaa = Thr or Val
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (112)..(112)

-continued

<223> OTHER INFORMATION: Xaa = Thr or Val
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (113)..(113)
<223> OTHER INFORMATION: Xaa = Leu or Met

<400> SEQUENCE: 54

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Xaa
1 5 10 15
Ser Leu Xaa Leu Ser Cys Xaa Ala Ser Gly Phe Thr Phe Ser Asn Phe
 20 25 30
Trp Met Xaa Trp Xaa Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
Ala Ser Ile Thr His Ser Gly Gly Ile Thr Tyr Tyr Xaa Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Xaa Xaa Lys Xaa Xaa Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Xaa Glu Asp Thr Ala Xaa Tyr Tyr Cys
 85 90 95
Ser Arg Asp Pro Thr Glu Ala Pro Phe Asp Tyr Trp Gly Gln Gly Xaa
100 105 110
Xaa Val Thr Val Ser Ser
 115

<210> SEQ ID NO 55
<211> LENGTH: 23
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 50G2-3 L-FR1; 1910G2HzL2, 1910G2HzL3 and
1910G2HzL9 L-FR1
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa = Pro or Ser
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (15)..(15)
<223> OTHER INFORMATION: Xaa = Leu or Val
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (18)..(18)
<223> OTHER INFORMATION: Xaa = Gln or Arg

<400> SEQUENCE: 55

Asp Ile Gln Met Thr Gln Ser Pro Xaa Ser Leu Ser Ala Ser Xaa Gly
1 5 10 15
Asp Xaa Val Thr Ile Thr Cys
 20

<210> SEQ ID NO 56
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 50G2-3 L-FR2; 1910G2HzL2, 1910G2HzL3 and
1910G2HzL9 L-FR2
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: Xaa = Lys or Arg

<400> SEQUENCE: 56

-continued

Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Xaa Gln Leu Ile Arg
 1 5 10 15

<210> SEQ ID NO 57
 <211> LENGTH: 32
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: 50G2-3 L-FR3; 1910G2HzL2 L-FR3; 1910G2HzL3
 L-FR3; 1910G2HzL9 L-FR3
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (2)..(2)
 <223> OTHER INFORMATION: Xaa = Thr or Val
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (15)..(15)
 <223> OTHER INFORMATION: Xaa = Phe or Tyr
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (16)..(16)
 <223> OTHER INFORMATION: Xaa = Ser or Thr
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (18)..(18)
 <223> OTHER INFORMATION: Xaa = Ser or Thr
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (21)..(21)
 <223> OTHER INFORMATION: Xaa = Asn or Ser
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (22)..(22)
 <223> OTHER INFORMATION: Xaa = Leu or Val
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (23)..(23)
 <223> OTHER INFORMATION: Xaa = Glu or Gln
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (24)..(24)
 <223> OTHER INFORMATION: Xaa = Pro or Ser
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (29)..(29)
 <223> OTHER INFORMATION: Xaa = Ser or Thr
 <400> SEQUENCE: 57

Gly Xaa Pro Ser Arg Phe Ser Gly Ser Gly Lys Asp Xaa Xaa
 1 5 10 15

Phe Xaa Ile Ser Xaa Xaa Xaa Xaa Glu Asp Ile Ala Xaa Tyr Tyr Cys
 20 25 30

<210> SEQ ID NO 58
 <211> LENGTH: 10
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: 50G2-3 L-FR4; 1910G2HzL2, 1910G2HzL3 and
 1910G2HzL9 L-FR4
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (3)..(3)
 <223> OTHER INFORMATION: Xaa = Ala or Gln
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (9)..(9)
 <223> OTHER INFORMATION: Xaa = Ile or Leu
 <400> SEQUENCE: 58

Phe Gly Xaa Gly Thr Lys Leu Glu Xaa Lys

-continued

1	5	10
---	---	----

```
<210> SEQ ID NO 59
<211> LENGTH: 107
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 50G2-3 VL; 1910G2HzL2 VL; 1910G2HzL3 VL;
1910G2HzL9 VL
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa = Pro or Ser
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (15)..(15)
<223> OTHER INFORMATION: Xaa = Leu or Val
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (18)..(18)
<223> OTHER INFORMATION: Xaa = Gln or Arg
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (45)..(45)
<223> OTHER INFORMATION: Xaa = Lys or Arg
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (58)..(58)
<223> OTHER INFORMATION: Xaa = Thr or Val
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (71)..(71)
<223> OTHER INFORMATION: Xaa = Phe or Tyr
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (72)..(72)
<223> OTHER INFORMATION: Xaa = Ser or Thr
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (74)..(74)
<223> OTHER INFORMATION: Xaa = Ser or Thr
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (77)..(77)
<223> OTHER INFORMATION: Xaa = Asn or Ser
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (78)..(78)
<223> OTHER INFORMATION: Xaa = Leu or Val
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (79)..(79)
<223> OTHER INFORMATION: Xaa = Glu or Gln
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (80)..(80)
<223> OTHER INFORMATION: Xaa = Pro or Ser
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (85)..(85)
<223> OTHER INFORMATION: Xaa = Ser or Thr
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (100)..(100)
<223> OTHER INFORMATION: Xaa = Ala or Gln
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (106)..(106)
<223> OTHER INFORMATION: Xaa = Ile or Leu

<400> SEQUENCE: 59

Asp Ile Gln Met Thr Gln Ser Pro Xaa Ser Leu Ser Ala Ser Xaa Gly
1 5 10 15
```

-continued

```

Asp Xaa Val Thr Ile Thr Cys Gln Ala Ser Gln Asn Ile Asn Asn Tyr
      20                25                30

Ile Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Xaa Gln Leu Ile
      35                40                45

Arg Tyr Thr Ser Thr Leu Val Ser Gly Xaa Pro Ser Arg Phe Ser Gly
      50                55                60

Ser Gly Ser Gly Lys Asp Xaa Xaa Phe Xaa Ile Ser Xaa Xaa Xaa Xaa
      65                70                75                80

Glu Asp Ile Ala Xaa Tyr Tyr Cys Leu Gln Tyr Asp Asn Val Pro Asn
      85                90                95

Thr Phe Gly Xaa Gly Thr Lys Leu Glu Xaa Lys
      100                105

```

```

<210> SEQ ID NO 60
<211> LENGTH: 95
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 60

```

```

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1      5                10                15

Asp Arg Val Thr Ile Thr Cys Gln Ala Ser Gln Asp Ile Ser Asn Tyr
      20                25                30

Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
      35                40                45

Tyr Asp Ala Ser Asn Leu Glu Thr Gly Val Pro Ser Arg Phe Ser Gly
      50                55                60

Ser Gly Ser Gly Thr Asp Phe Thr Phe Thr Ile Ser Ser Leu Gln Pro
      65                70                75                80

Glu Asp Ile Ala Thr Tyr Tyr Cys Gln Gln Tyr Asp Asn Leu Pro
      85                90                95

```

```

<210> SEQ ID NO 61
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 61

```

```

Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
1      5                10

```

```

<210> SEQ ID NO 62
<211> LENGTH: 98
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 62

```

```

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1      5                10                15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
      20                25                30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
      35                40                45

Ala Asn Ile Lys Gln Asp Gly Ser Glu Lys Tyr Tyr Val Asp Ser Val
      50                55                60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr

```

-continued

65	70	75	80
Leu Gln Met Asn Ser	Leu Arg Ala Glu Asp	Thr Ala Val Tyr	Tyr Cys
	85	90	95
Ala Arg			
<210> SEQ ID NO 63			
<211> LENGTH: 15			
<212> TYPE: PRT			
<213> ORGANISM: Homo sapiens			
<400> SEQUENCE: 63			
Tyr Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser			
1	5	10	15

1-66. (canceled)

67. An antibody or an antigen-binding fragment thereof binding to a human and/or a monkey PD-L1 protein, comprising HCDR1, HCDR2 and HCDR3 and LCDR1, LCDR2 and LCDR3, wherein the HCDR1 comprises or consists of an amino acid sequence as set forth in SEQ ID NO: 1, the HCDR2 comprises or consists of an amino acid sequence as set forth in SEQ ID NO: 2, and the HCDR3 comprises or consists of an amino acid sequence as set forth in SEQ ID NO: 3, the LCDR1 comprises or consists of an amino acid sequence as set forth in SEQ ID NO: 4, the LCDR2 comprises or consists of an amino acid sequence as set forth in SEQ ID NO: 5, and the LCDR3 comprises or consists of an amino acid sequence as set forth in SEQ ID NO: 6.

68. The antibody or antigen binding fragment according to claim **67**, wherein the antigen-binding fragment includes a Fab, a Fab', a F(ab)₂, a Fv fragment, a F(ab')₂, a scFv, a di-scFv, a VHH and/or a dAb.

69. The antibody or antigen binding fragment according to claim **67**, wherein the antibody is selected from the group consisting of a monoclonal antibody, a chimeric antibody, a humanized antibody and a fully human antibody.

70. The antibody or antigen binding fragment according to claim **67**, comprising H-FR1, H-FR2, H-FR3 and H-FR4, and the H-FR1, H-FR2, H-FR3 and H-FR4 are selected from any one of groups of amino acid sequences as follows:

- (1) H-FR1: SEQ ID NO: 7, H-FR2: SEQ ID NO: 9, H-FR3: SEQ ID NO: 12 and H-FR4: SEQ ID NO: 18;
- (2) H-FR1: SEQ ID NO: 8, H-FR2: SEQ ID NO: 10, H-FR3: SEQ ID NO: 13 and H-FR4: SEQ ID NO: 19;
- (3) H-FR1: SEQ ID NO: 8, H-FR2: SEQ ID NO: 11, H-FR3: SEQ ID NO: 14 and H-FR4: SEQ ID NO: 18;
- (4) H-FR1: SEQ ID NO: 8, H-FR2: SEQ ID NO: 10, H-FR3: SEQ ID NO: 15 and H-FR4: SEQ ID NO: 19;
- (5) H-FR1: SEQ ID NO: 8, H-FR2: SEQ ID NO: 10, H-FR3: SEQ ID NO: 16 and H-FR4: SEQ ID NO: 19; and
- (6) H-FR1: SEQ ID NO: 8, H-FR2: SEQ ID NO: 11, H-FR3: SEQ ID NO: 17 and H-FR4: SEQ ID NO: 19;

or

comprising L-FR1, L-FR2, L-FR3 and L-FR4, and the L-FR1, L-FR2, L-FR3 and L-FR4 are selected from any one of groups of amino acid sequences as follows:

- (1) L-FR1: SEQ ID NO: 26, L-FR2: SEQ ID NO: 28, L-FR3: SEQ ID NO: 30 and L-FR4: SEQ ID NO: 34;

- (2) L-FR1: SEQ ID NO: 27, L-FR2: SEQ ID NO: 29, L-FR3: SEQ ID NO: 31 and L-FR4: SEQ ID NO: 35;
- (3) L-FR1: SEQ ID NO: 27, L-FR2: SEQ ID NO: 29, L-FR3: SEQ ID NO: 32 and L-FR4: SEQ ID NO: 35; and
- (4) L-FR1: SEQ ID NO: 27, L-FR2: SEQ ID NO: 29, L-FR3: SEQ ID NO: 33 and L-FR4: SEQ ID NO: 35.

71. The antibody or antigen binding fragment according to claim **67**, wherein the antibody or antigen binding fragment comprises a heavy chain variable region VH or a light chain variable region VL,

wherein the VH comprises or consists of an amino acid sequence as set forth in any one of SEQ ID NO: 20 to 25, or wherein the VL comprises or consists of an amino acid sequence as set forth in any one of SEQ ID NO: 36 to 39.

72. The antibody or antigen binding fragment according to claim **67**, wherein the antibody or antigen binding fragment comprises a heavy chain constant region, and the heavy chain constant region is derived from a human IgG constant region, optionally, the heavy chain constant region is derived from a human IgG1 constant region, and the human IgG1 constant region comprises an amino acid sequence as set forth in SEQ ID NO: 41,

or the antibody or antigen binding fragment comprises an antibody light chain constant region, and the antibody light chain constant region comprises an amino acid sequence as set forth in SEQ ID NO: 40.

73. The antibody or antigen binding fragment according to claim **67**, wherein the antibody or antigen binding fragment comprises an antibody heavy chain HC or an antibody light chain LC, and the HC comprises or consists of an amino acid sequence as set forth in any one of SEQ ID NO: 42 to 46, and the LC comprises an amino acid sequence as set forth in any one of SEQ ID NO: 47 to 49.

74. An immunoconjugate comprising the antibody or antigen binding fragment of claim **67**.

75. An isolated nucleic acid molecule(s) encoding the antibody or antigen binding fragment of claim **67**.

76. A vector comprising the nucleic acid molecule according to claim **75**.

77. A cell comprising the nucleic acid molecule according to claim **75** or the vector comprising said nucleic acid molecule.

78. A method for preparing the antibody or antigen binding fragment of claim **67**, comprising culturing the cell

comprising the nucleic acid molecule encoding the antibody or antigen binding fragment of claim 67 or the vector comprising said nucleic acid molecule under conditions that the antibody or antigen binding fragment of claim 67 can be expressed.

79. A pharmaceutical composition comprising the antibody or antigen binding fragment of claim 67, the nucleic acid molecule encoding said antibody or antigen binding fragment, the vector comprising said nucleic acid molecule and/or the cell of comprising said nucleic acid molecule or said vector, and optionally a pharmaceutically acceptable carrier.

80. A method for preventing, alleviating or treating a tumor, comprising administering the antibody or antigen binding fragment of claim 67 to a subject in need, optionally the tumor includes a tumor with a high expression of PD-1 or PD-L1.

81. The method according to claim 80, wherein the tumor includes a solid tumor.

82. The method according to claim 80, wherein the tumor includes breast cancer, lung cancer, gastric cancer and urothelial carcinoma.

83. A method for inhibiting the binding of PD-1 protein to PD-L1 protein or stimulating the secretion of cytokines in immune cells or detecting the presence and/or content of PD-L1 protein, comprising administering the antibody or antigen binding fragment of claim 67, the immunoconjugate and/or the pharmaceutical composition comprising the antibody or antigen binding fragment of claim 67.

84. A kit comprising the antibody or antigen binding fragment of claim 67, the immunoconjugate and/or the pharmaceutical composition comprising the antibody or antigen binding fragment of claim 67.

85. The antibody or antigen binding fragment according to claim 67, comprising a heavy chain variable region VH and a light chain variable region VL, wherein

VH comprises or consists of an amino acid sequence as set forth in SEQ ID NO: 20, and the VL comprises or consists of an amino acid sequence as set forth in SEQ ID NO:36;

VH comprises or consists of an amino acid sequence as set forth in SEQ ID NO: 21, and the VL comprises or consists of an amino acid sequence as set forth in SEQ ID NO:37;

VH comprises or consists of an amino acid sequence as set forth in SEQ ID NO: 22, and the VL comprises or consists of an amino acid sequence as set forth in SEQ ID NO:38;

VH comprises or consists of an amino acid sequence as set forth in SEQ ID NO: 23, and the VL comprises or consists of an amino acid sequence as set forth in SEQ ID NO:39;

VH comprises or consists of an amino acid sequence as set forth in SEQ ID NO: 24, and the VL comprises or consists of an amino acid sequence as set forth in SEQ ID NO:37; or

VH comprises or consists of an amino acid sequence as set forth in SEQ ID NO: 25, and the VL comprises or consists of an amino acid sequence as set forth in SEQ ID NO:39.

86. The antibody or antigen binding fragment according to claim 67, comprising a heavy chain HC and a light chain LC, wherein

HC comprises or consists of an amino acid sequence as set forth in SEQ ID NO: 42, and the LC comprises or consists of an amino acid sequence as set forth in SEQ ID NO:47;

HC comprises or consists of an amino acid sequence as set forth in SEQ ID NO: 43, and the LC comprises or consists of an amino acid sequence as set forth in SEQ ID NO:48;

HC comprises or consists of an amino acid sequence as set forth in SEQ ID NO: 44, and the LC comprises or consists of an amino acid sequence as set forth in SEQ ID NO:47;

HC comprises or consists of an amino acid sequence as set forth in SEQ ID NO: 45, and the LC comprises or consists of an amino acid sequence as set forth in SEQ ID NO:47; or

HC comprises or consists of an amino acid sequence as set forth in SEQ ID NO: 46, and the LC comprises or consists of an amino acid sequence as set forth in SEQ ID NO:49.

* * * * *