



US012409177B2

(12) **United States Patent**
Cao et al.

(10) **Patent No.:** **US 12,409,177 B2**
(45) **Date of Patent:** **Sep. 9, 2025**

(54) **FGFR INHIBITORS FOR THE TREATMENT OF CANCER**

(71) Applicant: **BridGene Biosciences, Inc.**, Sunnyvale, CA (US)

(72) Inventors: **Ping Cao**, Sunnyvale, CA (US); **Chao Zhang**, Sunnyvale, CA (US); **Michael J. Bishop**, Sunnyvale, CA (US)

(73) Assignee: **BridGene Biosciences, Inc.**, Sunnyvale, CA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 949 days.

(21) Appl. No.: **17/430,621**

(22) PCT Filed: **Feb. 14, 2020**

(86) PCT No.: **PCT/US2020/018356**

§ 371 (c)(1),

(2) Date: **Aug. 12, 2021**

(87) PCT Pub. No.: **WO2020/168237**

PCT Pub. Date: **Aug. 20, 2020**

(65) **Prior Publication Data**

US 2022/0218703 A1 Jul. 14, 2022

Related U.S. Application Data

(60) Provisional application No. 62/805,854, filed on Feb. 14, 2019.

(51) **Int. Cl.**

A61K 31/506 (2006.01)

A61K 45/06 (2006.01)

(Continued)

(52) **U.S. Cl.**

CPC **A61K 31/506** (2013.01); **A61K 45/06** (2013.01); **C07D 401/04** (2013.01); **C07D 401/14** (2013.01); **C07D 405/14** (2013.01)

(58) **Field of Classification Search**

CPC **A61K 31/506**; **A61K 45/06**; **C07D 401/04**; **C07D 401/14**; **C07D 405/14**; **A61P 35/00**

See application file for complete search history.

(56)

References Cited

U.S. PATENT DOCUMENTS

4,118,571 A 10/1978 Leshner et al.
7,880,000 B2 2/2011 Geuns-Meyer et al.

(Continued)

FOREIGN PATENT DOCUMENTS

CN 103596951 A 2/2014
GB 0121825 10/2001

(Continued)

OTHER PUBLICATIONS

Barillari et al., "Classical Bioisosteres", Bioisosteres in Medicinal Chemistry Ch. 2, Ed. Brown, Wiley-VCH Verlag GmbH & Co., pp. 15-29 (Year: 2012).*

CN Office Action in Chinese Application No. 2020800146914, dated May 5, 2023, 19 pages (with English translation).

Unknown, "Virtual Screening of Targeted FGFR4 Inhibitors and Theoretical Computational Study of Their Selective Mechanisms," China Master's Thesis Full Text Database Database (Master) Medicine and Health Science and Technology Series, May 2018, 64 pages (with English abstract).

(Continued)

Primary Examiner — Jeffrey S Lundgren

Assistant Examiner — Alexander K. Showalter

(74) *Attorney, Agent, or Firm* — Clark & Elbing LLP


(57)

ABSTRACT

The disclosure provides novel FGFR inhibitors based on the pyridinylpyrimidine. The disclosure includes inhibitors with broad inhibitory activity against all FGFR isoforms, and inhibitors with selective inhibition against FGFR4. These novel pyridinylpyrimidine-based FGFR inhibitors, or their derivatives, have strong potential to be used to treat cancer.

12 Claims, 3 Drawing Sheets

Specification includes a Sequence Listing.



(51) **Int. Cl.**

A61P 35/00 (2006.01)
C07D 401/04 (2006.01)
C07D 401/14 (2006.01)
C07D 405/14 (2006.01)

WO WO 2011075620 A1 6/2011

OTHER PUBLICATIONS

Wei, "Exploration of the Mechanism by Which Fisetin Induces Apoptosis in Breast Cancer Cells and Inhibits Embryonic Development in Zebrafish by Acting on the FGFR4 Pathway", China Master's Thesis Full Text Database Database (Master) Medicine and Health Science and Technology Series, Jan. 2015, 77 pages (with English abstract).

EP Extended Search Report in European Application No. 20755723, dated Oct. 11, 2022, 9 pages.

Hari et al., "Conformation-Selective Inhibitors Reveal Differences in the Activation and Phosphate-Binding Loops of the Tyrosine Kinases Abl and Src", ACS Chemical Biology, Oct. 2013, 8(12):2734-2743.

Ranjitkar et al., "Affinity Reagents that Target a Specific Inactive Form of Protein Kinases", Chemistry & Biology, Feb. 2010, 17(2):195-206.

Ranjitkar et al. "Affinity-Based Probes Based on Type II Kinase Inhibitors," J Am Chem Soc. 134(46):19017-19025 (Nov. 2012) (21 pages).

* cited by examiner

(56)


References Cited

U.S. PATENT DOCUMENTS

8,476,434	B2	7/2013	Geuns-Meyer et al.	
2006/0009453	A1	1/2006	Geuns-Meyer et al.	
2007/0213325	A1	9/2007	Cee et al.	
2009/0069327	A1 *	3/2009	Ding	C07D 239/42 544/122
2011/0201602	A1	8/2011	Geuns-Meyer et al.	
2012/0035171	A1	2/2012	Saxty et al.	
2022/0048883	A1 *	2/2022	Zhang	A61P 35/00

FOREIGN PATENT DOCUMENTS

GB	2 369 359	A	5/2002
JP	2009525978	A	7/2009
JP	2014512357	A	5/2014
WO	WO 2007/092531	A2	8/2007



Subdomain V

FGFR1	-IVEYASKGNLR-	SEQ ID NO: 57
FGFR2	-IVEYASKGNLR-	SEQ ID NO: 58
FGFR3	-LVEYAAKGNLR-	SEQ ID NO: 59
FGFR4	-IVEYAAKGNLR-	SEQ ID NO: 60

FIGURE 1




FIGURE 2A




FIGURE 2B




FIGURE 3A




FIGURE 3B

1

FGFR INHIBITORS FOR THE TREATMENT OF CANCER

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a 35 USC § 371 National Stage application of International Application No. PCT/US2020/018356 filed Feb. 14, 2020, now pending; which claims the benefit under 35 USC § 119 (e) to U.S. Application Ser. No. 62/805,854 filed Feb. 14, 2019. The disclosure of each of the prior applications is considered part of and is incorporated by reference in the disclosure of this application.

INCORPORATION BY REFERENCE OF SEQUENCE LISTING

The material in the accompanying sequence listing is hereby incorporated by reference into this application. The accompanying sequence listing text file, named BRIDGE1100-1_ST25.txt, was created on Aug. 5, 2021 and is 1,084 kB in size. The file can be accessed using Microsoft Word on a computer that uses Windows OS.

FIELD

The present disclosure relates generally to novel fibroblast growth factor receptor (FGFR) inhibitors, and more specifically to the use of pyridinylpyrimidine-based FGFR4 specific inhibitors for the treatment of cancer.

BACKGROUND

The human family of fibroblast growth factor receptors (FGFRs) is composed of four receptor tyrosine kinases that bind 18 ligands called fibroblast growth factors (FGFs). The four members (FGFR1, FGFR2, FGFR3, and FGFR4) are highly conserved among each other and consist of extracellular ligand-binding domains, a transmembrane segment, and a cytoplasmic tyrosine kinase domain. Upon binding of ligands to the extracellular domains of FGFRs, the kinase domains are activated by autophosphorylation and then phosphorylate cytoplasmic substrates, triggering downstream signaling cascades that control cell growth and differentiation.

The FGFR signaling pathway is an important and validated target for cancer therapeutics since it plays a crucial role in tumor proliferation, angiogenesis, migration, and survival. Mutations and overexpression of FGFRs and their ligands have been reported in several cancers, such as breast, lung, bladder, prostate, and gastric. For instance, amplification of FGFR1 has been found in about 10% of breast cancers (predominantly in estrogen receptor positive diseases), in 10-20% of squamous non-small-cell lung cancer (NSCLC), ovarian cancer (~5%), and bladder cancer (3%). FGFR2 amplification has been detected in gastric (5-10%) and breast cancers (4% of triple negative cases), and mutations in FGFR2 occur in 12% of endometrial carcinomas. FGFR3 mutations were identified in about 70% of non-muscle-invasive bladder cancers and 10-20% of invasive high-grade bladder cancers. Amplification and activating mutations in FGFR4 have been described in 8% of rhabdomyosarcoma patients. In addition, many preclinical studies have reported FGFR4 overexpression in prostate, colon, and liver cancers.

A number of FGFR small-molecule inhibitors have been developed and evaluated in clinical trials for the treatment of

2


cancers, but most of them are pan-FGFR inhibitors with promiscuous kinome activity, such as BGJ398 and LY-2874455. It has been found, from sequence analysis, that FGFR4 contains a cysteine (Cys552) located near the ATP-binding site, in the hinge region of the receptor, which is unique within the FGFR family and rare among other kinases. In fact, the first selective FGFR4 inhibitor, BLU9931, was discovered recently targeting this unique cysteine and exhibited very good specificity and significant antitumor activity against hepatocellular carcinoma in vivo. However, the potency and bioavailability of BLU9931 is suboptimal for clinical applications.

SUMMARY

Embodiments of the present disclosure include novel pyridinylpyrimidine-based compounds that are potent FGFR4 specific inhibitors. The FGFR4 specific inhibitors can be used as targeted therapies to treat cancer.

In some embodiments, the present disclosure provides a compound of Formula (I)


Formula (I)



or an optically pure stereoisomer, solvate, or pharmaceutically acceptable salt thereof.


In some embodiments, each R₁, R₂, R₃, R₄, and R₅ can be independently H, F, Cl, Br, C₁₋₄ alkyl, cyclopropyl, N₃, NH₂, NO₂, CF₃, OCF₃, OCHF₂, or OC₁₋₄alkyl. In some embodiments, R₆ can be (CH₂)₀₋₅CH=CH₂, (CH₂)₀₋₅C≡CH, NHCO(CH₂)₀₋₅CH=CH₂, NH(CH₂)₀₋₅CH=CH₂, OCO(CH₂)₀₋₅CH=CH₂, O(CH₂)₀₋₅CH=CH₂, NHCO(CH₂)₀₋₅C≡CH, or OCO(CH₂)₀₋₅C≡CH.

In some embodiments, Linker can be selected from the group consisting of C₁₋₂₀alkyl, COC₁₋₂₀alkyl, CO₂C₁₋₂₀alkyl,




3

-continued




phenyl, naphthyl, anthracene,



4


-continued



optionally substituted with one or more substituents selected from the group consisting of F, Cl, Br, C₁₋₄ alkyl, CF₃, CHF₂, NO₂, —NR₇—(CH₂)_n—R₈, —CO NR₇—(CH₂)_n—R₈, —CO—(CH₂)_n—R₈, —OCO—(CH₂)_n—R₈, and —O—(CH₂)_n—R₈.

In some embodiments, n can be an integer selected from 0 to 5. In some embodiments, R₇ can be selected from the group consisting of H, C₁₋₄ alkyl, and COC₁₋₄alkyl.


In some embodiments, R₈ can be selected from the group consisting of C₁₋₂₀ alkyl COC₁₋₂₀alkyl, CO₂C₁₋₂₀alkyl,



5

optionally substituted with one or more substituents selected from the group consisting of F, Cl, Br, C₁₋₄ alkyl, CF₃, CHF₂, and NO₂.

The present disclosure also provides a compound of Formula (II)




Formula (II)

or an optically pure stereoisomer, solvate, or pharmaceutically acceptable salt thereof.


In some embodiments, R₁ can be (CH₂)₀₋₅CH=CH₂, (CH₂)₀₋₅C≡CH, NHCO(CH₂)₀₋₅CH=CH₂, NH(CH₂)₀₋₅CH=CH₂, OCO(CH₂)₀₋₅CH=CH₂, O(CH₂)₀₋₅CH=CH₂, NHCO(CH₂)₀₋₅C≡CH, or OCO(CH₂)₀₋₅C≡CH.

Linker can be selected from the group consisting of C₁₋₂₀alkyl, COC₁₋₂₀alkyl, CO₂C₁₋₂₀alkyl,



phenyl, naphthyl, anthracene,


6




optionally substituted with one or more substituents selected from the group consisting of F, Cl, Br, C₁₋₄ alkyl, CF₃, CHF₂, NO₂, —NR₂—(CH₂)_n—R₃, —CO NR₂—(CH₂)_n—R₃, —CO—(CH₂)_n—R₃, —OCO—(CH₂)_n—R₃, and —O—(CH₂)_n—R₃.

In some embodiments, R₂ can be selected from the group consisting of H, C₁₋₄ alkyl, and COC₁₋₄alkyl. n is an integer selected from 0 to 5.


In some embodiments, R₃ can be selected from the group consisting of C₁₋₂₀ alkyl, COC₁₋₂₀alkyl, CO₂C₁₋₂₀alkyl,







9





-continued



In some embodiments, the compound can be selected from the group consisting of


**10**

-continued



11

-continued



The present disclosure also provides a pharmaceutical formulation including the compound according to Formula (I), Formula (II), Formula (III), or an optically pure stereoisomer or pharmaceutically acceptable salt thereof. The present disclosure further provides a method for treating cancer

12

in a subject including administering a compound with the structure of Formula (I), Formula (II), Formula (III), or an optically pure stereoisomer or pharmaceutically acceptable salt thereof to the subject. In some embodiments, the compound targets amino acid residue 484 of SEQ ID NO: 52, amino acid residue 512 of SEQ ID NO: 56, or amino acid residue 552 of SEQ ID NO: 50 or 54. The present disclosure also provides a method of inhibiting a kinase activity including contacting a cell with a compound with the structure of Formula (I), Formula (II), Formula (III), or an optically pure stereoisomer or pharmaceutically acceptable salt thereof. In some embodiments, the kinase can be Anaplastic lymphoma kinase (ALK), Epidermal growth factor receptor (EGFR), Ephrin type-3 receptor 3 (EPH-B3), Focal adhesion kinase (FAK), Fibroblast growth factor receptor 1 (FGFR1), Fibroblast growth factor receptor 2 (FGFR2), Fibroblast growth factor receptor 3 (FGFR3), Fibroblast growth factor receptor 4 (FGFR4), Mast/stem cell growth factor receptor (SCFR or KIT), Mitogen-activated protein kinase kinase 1 (MAP2K1 or MEK1), Hepatocyte growth factor receptor (HGFR or MET), Platelet-derived growth factor receptor alpha (PDGFRA), Platelet-derived growth factor receptor beta (PDGFRB), Proto-oncogene tyrosine kinase receptor (RET), Proto-oncogene tyrosine-protein kinase (ROS) or Tyrosine-protein kinase receptor (TYRO 3). The cell can be a cancer cell. The cancer cell can be a breast, lung, bladder, prostate, ovarian, endometrial, rhabdomyosarcoma, liver or gastric cancer cell.

Other features and advantages can become apparent from the following detailed drawings and description.

BRIEF DESCRIPTION OF THE DRAWINGS

In order to facilitate a full understanding of the present disclosure, reference is now made to the accompanying drawings, in which like elements are referenced with like numerals. These drawings should not be construed as limiting the present disclosure, but are intended to be illustrative only. The drawings are not necessarily to scale, or inclusive of all elements of a system, emphasis instead generally being placed upon illustrating the concepts, structures, and techniques sought to be protected herein.

FIG. 1 is a schematic representation of FGFR kinases. A. Schematic representation of the selective inhibition of FGFRs by electrophilic inhibitors by covalently targeting a thiol group (SH) in the cysteine residue. B. Partial sequence alignment of FGFR kinases within the subdomain V highlighting the unique cysteine residue that FGFR4 contains near the ATP-binding site;

FIG. 2A is the pharmacokinetic data of compound 1 administered by intraperitoneal injection (IP) into mice at 10 mg/kg, according to some embodiments of the present disclosure;

FIG. 2B is the pharmacokinetic data of compound 1 administered by intravenous injection (IV) into mice at 1 mg/kg and per os (PO) at 5 mg/kg, according to some embodiments of the present disclosure;

FIG. 3A is the mean plasma concentration time profile of compound 5 by IP administration in female CD-1 mice at 10 mg/kg, according to some embodiments of the present disclosure; and

FIG. 3B is the pharmacokinetic data of compound 5 by IV administration at 5 mg/kg and PO administration at 100 mg/kg, according to some embodiments of the present disclosure.

13

DESCRIPTION

Disclosed herein is a class of chemical compounds as FGFR4-selective inhibitors. These small molecule inhibitors can be used as targeted therapies to treat cancer.

Fibroblast growth factor receptors (FGFRs) are highly conserved receptors consisting of extracellular ligand-binding domain, a transmembrane segment, and a cytoplasmic tyrosine kinase domain. The human FRFR family includes four members, FGFR1, FGFR2, FGFR3, and FGFR4, which can be bound by 18 different ligands called fibroblast growth factors (FGFs). Each receptor is composed of an extracellular domain, consisting of three immunoglobulin-like domain (IgI IgII, and IgIII) and an acid box, the IgII and IgIII domains constituting the FGF ligand-binding site; a transmembrane domain; and a tyrosine kinase cytoplasmic domain. FGFRs also contain hinge region (subdomain V), located near the ATP-binding site (SEQ ID NOs 57-60). FGFRs encoding mRNA are subjected to alternative splicing, giving rise to several protein-coding splice variants or isoforms (SEQ ID NOs: 1; 3; 5; 7; 9; 11; 13; 15; 17; 19; 21; 23; 25; 27; 29; 31; 33; 35; 37; 39; 41; 43; 45; 47; 49; 51; 53; and 55). As shown in Table 3, the human FGFR1 gene encodes 9 protein coding splice variants (SEQ ID Nos: 2; 4; 6; 8; 10; 12; 14; 16; and 18), the human FGFR2 gene encodes 11 protein coding splice variants (SEQ ID Nos: 20; 22; 24; 26; 28; 30; 32; 34; 36; 38; and 40), the human FGFR3 gene encodes four protein coding splice variants (SEQ ID Nos: 42; 44; 46; and 48), and the human FGFR4 gene encodes four protein coding splice variants (SEQ ID Nos: 50; 52; 54; and 56).

There are 18 members in the FGF family of ligands (FGF1-FGF10 and FGF16-FGF23). The binding of a ligand to the extracellular domain of a FGFR leads to receptor dimerization resulting in the activation of the tyrosine-kinase domain by auto-phosphorylation. Subsequently, an activated FGFR phosphorylates cytoplasmic substrates, such as FGFR substrate 2 (FRS2) and phosphatase C γ (PLC γ) triggering downstream signaling cascades. Activated FRS2 promotes the RAS-mitogen-activated protein kinase (MAPK) or the phosphoinositide 3-kinase (PI3K)-AKT pathways that regulate cell proliferation, differentiation and survival. On the other hand, the activation of PLC γ lead to calcium release and regulates events that mediate cell motility.

Deregulation of FGFR signaling has been linked to oncogenesis through several mechanisms including activating mutations, gene amplification or changes in post-transcriptional processing, and translocation, leading to constitutive activation of the receptor.

Specifically, FGFR4 amplification and activating mutations have been described in patients with rhabdomyosarcoma, and FGFR4 overexpression have been linked to prostate, colon, breast and liver cancers. FGFR4 differs from the other FGFRs by the presence of a cysteine in the hinge region, which is unique within the FGFR family and rare among other kinases. Depending on the isoform, the cysteine is located at different positions: Cys484 of SEQ ID NO: 52, Cys512 of SEQ ID NO: 56, or Cys552 of SEQ ID NO: 50 or 54. This unique cysteine can be targeted for the design of FGFR4 specific inhibitors exhibiting very good specificity.

As used herein, the term "FGFR inhibitor" or "FGFRi" refers to any compound capable of inhibiting the enzymatic activity of FGFR, including its own auto-phosphorylation and the kinase activity. Such inhibitors efficiently inhibit FGFRs, and are said to "inhibit", "decrease", or "reduce" the bio-


14

logical activity of FGFRs. The FGFR inhibitors of the disclosure can be "pan-inhibitor" and present a broad efficiency at inhibiting one or more of FGFR1-FGFR4, or present a specific efficiency at inhibiting only one FGFR, FGFR4 for example.

The efficiency of a compound can be referred to by its IC₅₀ value. The "IC₅₀" is the half-maximal inhibitory concentration (IC₅₀) of a compound. As used herein, the IC₅₀ of a FGFRi refers to the concentration of inhibitor which is sufficient to induce the inhibition of the enzymatic activity of FGFR halfway between the baseline and maximum after a specified exposure time. The IC₅₀ value of the present disclosure specifically refers to the concentration of FGFR inhibitor which is sufficient to induce the inhibition of one or more FGFRs, i.e. FGFR1, FGFR2, FGFR3 and/or FGFR4.

In some embodiments, the present disclosure provides a compound of Formula (I)


Formula (I)



or an optically pure stereoisomer, solvate, or pharmaceutically acceptable salt thereof.


In some embodiments, each R₁, R₂, R₃, R₄, and R₅ can be independently H, F, Cl, Br, C₁₋₄ alkyl, cyclopropyl, N₃, NH₂, NO₂, CF₃, OCF₃, OCHF₂, or OC₁₋₄alkyl. In some embodiments, R₆ can be (CH₂)₀₋₅CH=CH₂, (CH₂)₀₋₅C≡CH, NHCO(CH₂)₀₋₅CH=CH₂, NH(CH₂)₀₋₅CH=CH₂, OCO(CH₂)₀₋₅CH=CH₂, O(CH₂)₀₋₅CH=CH₂, NHCO(CH₂)₀₋₅C≡CH, or OCO(CH₂)₀₋₅C≡CH.

In some embodiments, Linker can be selected from the group consisting of C₁₋₂₀ alkyl, COC₁₋₂₀alkyl, CO₂C₁₋₂₀alkyl,




15


-continued



phenyl, naphthyl, anthracene,

**16**


-continued



optionally substituted with one or more substituents selected from the group consisting of F, Cl, Br, C_{1-4} alkyl, CF_3 , CHF $_2$, NO_2 , $-NR_7-(CH_2)_n-R_8$, $-CO-NR_7-(CH_2)_n-R_8$, $-CO-(CH_2)_n-R_8$, $-OCO-(CH_2)_n-R_8$, and $-O-(CH_2)_n-R_8$. The number of the optional substituents can be an integer selected from 0 to 4.

In some embodiments, n can be an integer selected from 0 to 5. In some embodiments, R_7 can be selected from the group consisting of H, C_{1-4} alkyl, and COC_{1-4} alkyl.

In some embodiments, R_8 can be selected from the group consisting of C_{1-20} alkyl, COC_{1-20} alkyl, CO_2C_{1-20} alkyl,




optionally substituted with one or more substituents selected from the group consisting of F, Cl, Br, C_{1-4} alkyl, CF_3 , CHF $_2$, and NO_2 .

In some embodiments, R_1 and R_5 are Cl, R_2 and R_4 are OMe, and R_3 is H.

The present disclosure also provides a compound of Formula (III)


Formula (III)



or an optically pure stereoisomer, solvate or pharmaceutically acceptable salt thereof.

In some embodiments, each R₁, R₂, R₃, R₄, and R₅ can be independently selected from the group consisting of H, F, Cl, Br, C₁₋₄ alkyl, cyclopropyl, N₃, NH₂, NO₂, CF₃, OCF₃, OCHF₂, and OC₁₋₄alkyl. n can be an integer selected from 0 to 5.

In some embodiments, R₆ can be selected from the group consisting of C₁₋₂₀ alkyl, COC₁₋₂₀alkyl, CO₂C₁₋₂₀alkyl,




optionally substituted with one or more substituents selected from the group consisting of F, Cl, Br, C₁₋₄ alkyl, CF₃, CHF₂, and NO₂. R₇ can be selected from the group consisting of H, C₁₋₄ alkyl, and COC₁₋₄alkyl. The number of the optional substituents can be an integer selected from 0 to 4.

In some embodiments, R₁ and R₅ are Cl, R₂ and R₄ are OMe, and R₃ is H.

In some embodiments, the present disclosure provides a compound of Formula (IV)

Formula (IV)




or an optically pure stereoisomer, solvate, or pharmaceutically acceptable salt thereof.

In some embodiments, each R_1 , R_2 , R_3 , R_4 , and R_5 can be independently H, F, Cl, Br, C_{1-4} alkyl, cyclopropyl, N_3 , NH_2 , NO_2 , CF_3 , OCF_3 , $OCHF_2$, or OC_{1-4} alkyl. In some embodiments, R_6 can be $(CH_2)_{0.5}CH=CH_2$, $(CH_2)_{0.5}C\equiv CH$, $NHCO(CH_2)_{0.5}CH=CH_2$, $NH(CH_2)_{0.5}CH=CH_2$, $OCO(CH_2)_{0.5}CH=CH_2$, $O(CH_2)_{0.5}CH=CH_2$, $NHCO(CH_2)_{0.5}C\equiv CH$, or $OCO(CH_2)_{0.5}C\equiv CH$.


In some embodiments, each A and B can be independently NR₇, CONR₇, O, CO, or OCO.

In some embodiments, Core can be selected from the group consisting of




Each D can be independently N or CH.

In some embodiments, Linker can be selected from the group consisting of C₁₋₂₀ alkyl, COC₁₋₂₀alkyl, CO₂C₁₋₂₀alkyl,




21


-continued



phenyl, naphthyl, anthracene,

**22**


-continued



optionally substituted with one or more substituents selected from the group consisting of F, Cl, Br, C₁₋₄ alkyl, CF₃, CHF₂, NO₂, —NR₇—(CH₂)_n—R₈, —CO NR₇—(CH₂)_n—R₈, —CO—(CH₂)_n—R₈, —OCO—(CH₂)_n—R₈, and —O—(CH₂)_n—R₈. The number of the optional substituents can be an integer selected from 0 to 4.


In some embodiments, n can be an integer selected from 0 to 5. In some embodiments, R₇ can be selected from the group consisting of H, C₁₋₄ alkyl, and COC₁₋₄alkyl.

In some embodiments, R₈ can be selected from the group consisting of C₁₋₂₀alkyl, COC₁₋₂₀alkyl, CO₂C₁₋₂₀alkyl,




23

-continued

**24**

-continued



optionally substituted with one or more substituents selected from the group consisting of F, Cl, Br, C₁₋₄ alkyl, CF₃, CHF₂, and NO₂.

In some embodiments, the compound of the present disclosure is selected from the group consisting of compound 1-4 and 6-16 as shown in Table 1.


TABLE 1

The FGFR inhibitor compounds in the present disclosure.	
Compound Number	Chemical Structure
compound 1	
compound 2	
compound 3	


The FGFR inhibitor compounds in the present disclosure.

Compound Number Chemical Structure


compound 4



compound 5




compound 6




The FGFR inhibitor compounds in the present disclosure.

Compound Number Chemical Structure


compound 7



compound 8




compound 9




The FGFR inhibitor compounds in the present disclosure.

Compound Number Chemical Structure


compound 10



compound 11




compound 12




The FGFR inhibitor compounds in the present disclosure.

Compound Number Chemical Structure

compound 13



compound 14



compound 15






TABLE 1-continued

The FGFR inhibitor compounds in the present disclosure.	
Compound Number	Chemical Structure
compound 16	

The present disclosure also provides a pharmaceutical formulation including the compound according to Formula (I), Formula (II), Formula (III), or an optically pure stereoisomer or pharmaceutically acceptable salt thereof. The present disclosure further provides a method for treating cancer in a subject including administering a compound with the structure of Formula (I), Formula (II), Formula (III), or an optically pure stereoisomer or pharmaceutically acceptable salt thereof to the subject. The present disclosure also provides a method of inhibiting a kinase activity including contacting a cell with a compound with the structure of Formula (I), Formula (II), Formula (III), or an optically pure stereoisomer or pharmaceutically acceptable salt thereof. In some embodiments, the kinase can be Anaplastic lymphoma kinase (ALK), Epidermal growth factor receptor (EGFR), Ephrin type-3 receptor 3 (EPH-B3), Focal adhesion kinase (FAK), Fibroblast growth factor receptor 1 (FGFR1), Fibroblast growth factor receptor 2 (FGFR2), Fibroblast growth factor receptor 3 (FGFR3), Fibroblast growth factor receptor 4 (FGFR4), Mast/stem cell growth factor receptor (SCFR or KIT), Mitogen-activated protein kinase kinase 1 (MAP2K1 or MEK1), Hepatocyte growth factor receptor (HGFR or MET), Platelet-derived growth factor receptor alpha (PDG-FRA), Platelet-derived growth factor receptor beta (PDG-FRB), Proto-oncogene tyrosine kinase receptor (RET), Proto-oncogene tyrosine-protein kinase (ROS) or Tyrosine-protein kinase receptor (TYRO 3). The cell can be a cancer cell. The cancer cell can be a breast, lung, bladder, prostate, ovarian, endometrial, rhabdomyosarcoma, liver or gastric cancer cell.

The present disclosure specifically provides compounds with non-obvious property improvement compared to control structures such as compound 5 with the structure of



The IUPAC name of the compound is N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)amino)pyridin-3-yl)pyrimidin-4-yl)amino)-3-methylphenyl)acrylamide.

The term “about” will be understood by persons of ordinary skill in the art. Whether the term “about” is used explicitly or not, every quantity given herein refers to the actual given value, and it is also meant to refer to the approximation to such given value that would be reasonably inferred based on the ordinary skill in the art.

Alkyl groups refer to univalent groups derived from alkanes by removal of a hydrogen atom from any carbon atom, which include straight chain and branched chain with from 1 to 12 carbon atoms, and typically from 1 to about 10 carbons or in some embodiments, from 1 to about 6 carbon atoms, or in other embodiments having 1, 2, 3 or 4 carbon atoms. Examples of straight chain alkyl groups include, but are not limited to, methyl, ethyl, n-propyl, n-butyl, n-pentyl, and n-hexyl groups. Examples of branched chain alkyl groups include, but are not limited to isopropyl, isobutyl, sec-butyl and tert-butyl groups. Alkyl groups may be substituted or unsubstituted. Representative substituted alkyl groups may be mono-substituted or substituted more than once, such as, but not limited to, mono-, di-, or tri-substituted. As used herein, the term alkyl, unless otherwise stated, refers to both cyclic and noncyclic groups.

35

The terms "cyclic alkyl" or "cycloalkyl" refer to univalent groups derived from cycloalkanes by removal of a hydrogen atom from a ring carbon atom. Cycloalkyl groups are saturated or partially saturated non-aromatic structures with a single ring or multiple rings including isolated, fused, bridged, and spiro ring systems, having 3 to 14 carbon atoms, or in some embodiments, from 3 to 12, or 3 to 10, or 3 to 8, or 3, 4, 5, 6 or 7 carbon atoms. Cycloalkyl groups may be substituted or unsubstituted. Representative substituted cycloalkyl groups may be mono-substituted or substituted more than once, such as, but not limited to, mono-, di-, or tri-substituted. Examples of monocyclic cycloalkyl groups include, but are not limited to cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl groups. Examples of multi-cyclic ring systems include, but are not limited to, bicycle [4.4.0]decane, bicycle[2.2.1]heptane, spiro[2.2]pentane, and the like.

Alkenyl groups refer to straight and branched chain and cycloalkyl groups as defined above, with one or more double bonds between two carbon atoms. Alkenyl groups may have 2 to about 12 carbon atoms, or in some embodiment from 1 to about 10 carbons or in other embodiments, from 1 to about 6 carbon atoms, or 1, 2, 3 or 4 carbon atoms in other embodiments. Alkenyl groups may be substituted or unsubstituted. Representative substituted alkenyl groups may be mono-substituted or substituted more than once, such as, but not limited to, mono-, di-, or tri-substituted. Examples of alkenyl groups include, but are not limited to, vinyl, allyl, $-\text{CH}=\text{CH}(\text{CH}_3)$, $-\text{CH}=\text{C}(\text{CH}_3)_2$, $-\text{C}(\text{CH}_3)=\text{CH}_2$, cyclopentenyl, cyclohexenyl, butadienyl, pentadienyl, and hexadienyl, among others.

Alkynyl groups refer to straight and branched chain and cycloalkyl groups as defined above, with one or more triple bonds between two carbon atoms. Alkynyl groups may have 2 to about 12 carbon atoms, or in some embodiment from 1 to about 10 carbons or in other embodiments, from 1 to about 6 carbon atoms, or 1, 2, 3 or 4 carbon atoms in other embodiments. Alkynyl groups may be substituted or unsubstituted. Representative substituted alkynyl groups may be mono-substituted or substituted more than once, such as, but not limited to, mono-, di-, or tri-substituted. Exemplary alkynyl groups include, but are not limited to, ethynyl, propargyl, and $-\text{C}\equiv\text{C}(\text{CH}_3)$, among others.

Aryl groups are cyclic aromatic hydrocarbons that include single and multiple ring compounds, including multiple ring compounds that contain separate and/or fused aryl groups. Aryl groups may contain from 6 to about 18 ring carbons, or in some embodiments from 6 to 14 ring carbons or even 6 to 10 ring carbons in other embodiments. Aryl group also includes heteroaryl groups, which are aromatic ring compounds containing 5 or more ring members, one or more ring carbon atoms of which are replaced with heteroatom such as, but not limited to, N, O, and S. Aryl groups may be substituted or unsubstituted. Representative substituted aryl groups may be mono-substituted or substituted more than once, such as, but not limited to, mono-, di-, or tri-substituted. Aryl groups include, but are not limited to, phenyl, biphenylenyl, triphenylenyl, naphthyl, anthryl, and pyrenyl groups.

Suitable heterocyclyl groups include cyclic groups with atoms of at least two different elements as members of its rings, of which one or more is a heteroatom such as, but not limited to, N, O, or S. Heterocyclyl groups may include 3 to about 20 ring members, or 3 to 18 in some embodiments, or about 3 to 15, 3 to 12, 3 to 10, or 3 to 6 ring members. The ring systems in heterocyclyl groups may be unsaturated, partially saturated, and/or saturated. Heterocyclyl groups

36

may be substituted or unsubstituted. Representative substituted heterocyclyl groups may be mono-substituted or substituted more than once, such as, but not limited to, mono-, di-, or tri-substituted. Exemplary heterocyclyl groups include, but are not limited to, pyrrolidinyl, tetrahydrofuryl, dihydrofuryl, tetrahydrothienyl, tetrahydrothiopyranyl, piperidyl, morpholinyl, thiomorpholinyl, thioxanyl, piperazinyl, azetidiny, aziridinyl, imidazolidinyl, pyrazolidinyl, thiazolidinyl, tetrahydrothiophenyl, tetrahydrofuranly, dioxolyl, furanyl, thiophenyl, pyrrolyl, imidazolyl, pyrazolyl, pyrazolinyl, triazolyl, tetrazolyl, oxazolyl, isoxazolyl, thiazolyl, thiazolinyl, oxetanyl, thietanyl, homopiperidyl, oxepanyl, thiepanyl, oxazepinyl, diazepinyl, thiazepinyl, 1,2,3,6-tetrahydropyridyl, indolinyl, 2H-pyranly, 4H-pyranly, dioxolanyl, dioxanyl, purinyl, quinoliziny, cinnolinyl, phthalazinyl, pteridinyl, and benzothiazolyl groups.

Polycyclic or polycyclyl groups refer to two or more rings in which two or more carbons are common to the two adjoining rings, wherein the rings are "fused rings"; if the rings are joined by one common carbon atom, these are "spiro" ring systems. Rings that are joined through non-adjacent atoms are "bridged" rings. Polycyclic groups may be substituted or unsubstituted. Representative polycyclic groups may be substituted one or more times.

Halogen groups include F, Cl, Br, and I; nitro group refers to $-\text{NO}_2$; cyano group refers to $-\text{CN}$; isocyano group refers to $-\text{N}=\text{C}$; epoxy groups encompass structures in which an oxygen atom is directly attached to two adjacent or non-adjacent carbon atoms of a carbon chain or ring system, which is essentially a cyclic ether structure. An epoxide is a cyclic ether with a three-atom ring.

An alkoxy group is a substituted or unsubstituted alkyl group, as defined above, singular bonded to oxygen. Alkoxy groups may be substituted or unsubstituted. Representative substituted alkoxy groups may be substituted one or more times. Exemplary alkoxy groups include, but are not limited to, methoxy, ethoxy, propoxy, butoxy, pentoxy, hexoxy, isopropoxy, sec-butoxy, tert-butoxy, cyclopropyloxy, cyclobutyloxy, cyclopentyloxy, and cyclohexyloxy groups.

The terms "amine" and "amino" refer to derivatives of ammonia, wherein one of more hydrogen atoms have been replaced by a substituent which include, but are not limited to alkyl, alkenyl, aryl, and heterocyclyl groups. Carbamate groups refers to $-\text{O}(\text{C}=\text{O})\text{NR}_1\text{R}_2$, where R_1 and R_2 are independently hydrogen, aliphatic groups, aryl groups, or heterocyclyl groups.

The term "optionally substituted" means the anteceding group may be substituted or unsubstituted. When substituted, the substituents of an "optionally substituted" group may include, without limitation, one or more substituents independently selected from the following groups or a particular designated set of groups, alone or in combination: lower alkyl, lower alkenyl, lower alkynyl, lower alkanoyl, lower heteroalkyl, lower heterocycloalkyl, lower haloalkyl, lower haloalkenyl, lower haloalkynyl, lower perhaloalkyl, lower perhaloalkoxy, lower cycloalkyl, phenyl, aryl, aryloxy, lower alkoxy, lower haloalkoxy, oxo, lower acyloxy, carbonyl, carboxyl, lower alkylcarbonyl, lower carboxyester, lower carboxamido, cyano, hydrogen, halogen, hydroxy, amino, lower alkylamino, arylamino, amido, nitro, thiol, lower alkylthio, lower haloalkylthio, lower perhaloalkylthio, arylthio, sulfonate, sulfonic acid, trisubstituted silyl, N_3 , SH, SCH_3 , $\text{C}(\text{O})\text{CH}_3$, CO_2CH_3 , CO_2H , pyridinyl, thiophene, furanyl, lower carbamate, and lower urea. Two substituents may be joined together to form a fused five-, six-, or seven-membered carbocyclic or heterocyclic ring consisting of zero to three heteroatoms, for example forming methyl-

enedioxy or ethylenedioxy. An optionally substituted group may be unsubstituted (e.g., $-\text{CH}_2\text{CH}_3$), fully substituted (e.g., $-\text{CF}_2\text{CF}_3$), monosubstituted (e.g., $-\text{CH}_2\text{CH}_2\text{F}$) or substituted at a level anywhere in-between fully substituted and monosubstituted (e.g., $-\text{CH}_2\text{CF}_3$). Where substituents are recited without qualification as to substitution, both substituted and unsubstituted forms are encompassed. Where a substituent is qualified as "substituted," the substituted form is specifically intended. Additionally, different sets of optional substituents to a particular moiety may be defined as needed; in these cases, the optional substitution will be as defined, often immediately following the phrase, "optionally substituted with."

Pharmaceutically acceptable salts of compounds described herein include conventional nontoxic salts or quaternary ammonium salts of a compound, e.g., from non-toxic organic or inorganic acids. For example, such conventional nontoxic salts include those derived from inorganic acids such as hydrochloride, hydrobromic, sulfuric, sulfamic, phosphoric, nitric, and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, palmitic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isothionic, and the like. In other cases, described compounds may contain one or more acidic functional groups and, thus, are capable of forming pharmaceutically acceptable salts with pharmaceutically acceptable bases. These salts can likewise be prepared in situ in the administration vehicle or the dosage form manufacturing process, or by separately reacting the purified compound in its free acid form with a suitable base, such as the hydroxide, carbonate or bicarbonate of a pharmaceutically acceptable metal cation, with ammonia, or with a pharmaceutically acceptable organic primary, secondary or tertiary amine. Representative alkali or alkaline earth salts include the lithium, sodium, potassium, calcium, magnesium, and aluminum salts and the like. Representative organic amines useful for the formation of base addition salts include ethylamine, diethylamine, ethylenediamine, ethanolamine, diethanolamine, piperazine and the like.

The term "treatment" is used interchangeably herein with the term "therapeutic method" and refers to both 1) therapeutic treatments or measures that cure, slow down, lessen symptoms of, and/or halt progression of a diagnosed pathologic conditions, disease or disorder, and 2) and prophylactic/preventative measures. Those in need of treatment may include individuals already having a particular medical disease or disorder as well as those who may ultimately acquire the disorder (i.e., those needing preventive measures).

The term "subject" as used herein refers to any individual or patient to which the subject methods are performed. Generally, the subject is human, although as will be appreciated by those in the art, the subject may be an animal.

The terms "therapeutically effective amount", "effective dose", "therapeutically effective dose", "effective amount," or the like refer to the amount of a subject compound that will elicit the biological or medical response in a tissue, system, animal or human that is being sought by administering said compound. Generally, the response is either amelioration of symptoms in a patient or a desired biological outcome. Such amount should be sufficient to inhibit FGFR enzymatic activity.

Also disclosed herein are pharmaceutical compositions including compounds with the structures of Formula (I),

Formula (II), or Formula (III). The term "pharmaceutically acceptable carrier" refers to a non-toxic carrier that may be administered to a patient, together with a compound of this disclosure, and which does not destroy the pharmacological activity thereof. Pharmaceutically acceptable carriers that may be used in these compositions include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

Pharmaceutically acceptable carriers that may be used in the pharmaceutical compositions of this disclosure include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, wool fat and self-emulsifying drug delivery systems (SEDDS) such as α -tocopherol, polyethyleneglycol 1000 succinate, or other similar polymeric delivery matrices.

In pharmaceutical composition comprising only the compounds described herein as the active component, methods for administering these compositions may additionally comprise the step of administering to the subject an additional agent or therapy. Such therapies include, but are not limited to, an anemia therapy, a diabetes therapy, a hypertension therapy, a cholesterol therapy, neuropharmacologic drugs, drugs modulating cardiovascular function, drugs modulating inflammation, immune function, production of blood cells; hormones and antagonists, drugs affecting gastrointestinal function, chemotherapeutics of microbial diseases, and/or chemotherapeutics of neoplastic disease. Other pharmacological therapies can include any other drug or biologic found in any drug class. For example, other drug classes can comprise allergy/cold/ENT therapies, analgesics, anesthetics, anti-inflammatories, antimicrobials, antivirals, asthma/pulmonary therapies, cardiovascular therapies, dermatology therapies, endocrine/metabolic therapies, gastrointestinal therapies, cancer therapies, immunology therapies, neurologic therapies, ophthalmic therapies, psychiatric therapies or rheumatologic therapies. Other examples of agents or therapies that can be administered with the compounds described herein include a matrix metalloprotease inhibitor, a lipoxigenase inhibitor, a cytokine antagonist, an immunosuppressant, a cytokine, a growth factor, an immunomodulator, a prostaglandin or an anti-vascular hyperproliferation compound.

The term "therapeutically effective amount" as used herein refers to the amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue, system, animal, individual or human that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes one or more of the

following: (1) Preventing the disease; for example, preventing a disease, condition or disorder in an individual that may be predisposed to the disease, condition or disorder but does not yet experience or display the pathology or symptomatology of the disease, (2) Inhibiting the disease; for example, inhibiting a disease, condition or disorder in an individual that is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (i.e., arresting further development of the pathology and/or symptomatology), and (3) Ameliorating the disease; for example, ameliorating a disease, condition or disorder in an individual that is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (i.e., reversing the pathology and/or symptomatology).

The compounds of this disclosure may be employed in a conventional manner for controlling the disease described herein, including, but not limited to, cancer. Such methods of treatment, their dosage levels and requirements may be selected by those of ordinary skill in the art from available methods and techniques. For example, the compounds of this disclosure may be combined with a pharmaceutically acceptable adjuvant for administration to a patient suffering from cancer in a pharmaceutically acceptable manner and in an amount effective to treat cancer.

Alternatively, the compounds of this disclosure may be used in compositions and methods for treating or protecting individuals against the diseases described herein, including but not limited to a cancer, over extended periods of time. The compounds may be employed in such compositions either alone or together with other compounds of this disclosure in a manner consistent with the conventional utilization of such compounds in pharmaceutical compositions. For example, a compound of this disclosure may be combined with pharmaceutically acceptable adjuvants conventionally employed in vaccines and administered in prophylactically effective amounts to protect individuals over an extended period of time against the diseases described herein, including, but not limited to, cancer.

As used herein, the terms "combination," "combined," and related terms refer to the simultaneous or sequential administration of therapeutic agents in accordance with this disclosure. For example, a described compound may be administered with another therapeutic agent simultaneously or sequentially in separate unit dosage forms or together in a single unit dosage form. Accordingly, the present disclosure provides a single unit dosage form comprising a described compound, an additional therapeutic agent, and a pharmaceutically acceptable carrier, adjuvant, or vehicle. Two or more agents are typically considered to be administered "in combination" when a patient or individual is simultaneously exposed to both agents. In many embodiments, two or more agents are considered to be administered "in combination" when a patient or individual simultaneously shows therapeutically relevant levels of the agents in a particular target tissue or sample (e.g., in brain, in serum, etc.).

When the compounds of this disclosure are administered in combination therapies with other agents, they may be administered sequentially or concurrently to the patient. Alternatively, pharmaceutical or prophylactic compositions according to this disclosure comprise a combination of ivermectin, or any other compound described herein, and another therapeutic or prophylactic agent. Additional therapeutic agents that are normally administered to treat a particular disease or condition may be referred to as "agents appropriate for the disease, or condition, being treated."

The compounds utilized in the compositions and methods of this disclosure may also be modified by appending appropriate functionalities to enhance selective biological properties. Such modifications are known in the art and include those, which increase biological penetration into a given biological system (e.g., blood, lymphatic system, or central nervous system), increase oral availability, increase solubility to allow administration by injection, alter metabolism and/or alter rate of excretion.

According to a preferred embodiment, the compositions of this disclosure are formulated for pharmaceutical administration to a subject or patient, e.g., a mammal, preferably a human being. Such pharmaceutical compositions are used to ameliorate, treat or prevent any of the diseases described herein including but not limited to cancer in a subject.

Agents of the disclosure are often administered as pharmaceutical compositions comprising an active therapeutic agent, i.e., and a variety of other pharmaceutically acceptable components. See Remington's Pharmaceutical Science (15th ed., Mack Publishing Company, Easton, Pa., 1980). The preferred form depends on the intended mode of administration and therapeutic application. The compositions can also include, depending on the formulation desired, pharmaceutically acceptable, non-toxic carriers or diluents, which are defined as vehicles commonly used to formulate pharmaceutical compositions for animal or human administration. The diluent is selected so as not to affect the biological activity of the combination. Examples of such diluents are distilled water, physiological phosphate-buffered saline, Ringer's solutions, dextrose solution, and Hank's solution. In addition, the pharmaceutical composition or formulation may also include other carriers, adjuvants, or nontoxic, nontherapeutic, nonimmunogenic stabilizers and the like.

In some embodiments, the present disclosure provides pharmaceutically acceptable compositions comprising a therapeutically effective amount of one or more of a described compound, formulated together with one or more pharmaceutically acceptable carriers (additives) and/or diluents for use in treating the diseases described herein, including, but not limited to cancer. While it is possible for a described compound to be administered alone, it is preferable to administer a described compound as a pharmaceutical formulation (composition) as described herein. Described compounds may be formulated for administration in any convenient way for use in human or veterinary medicine, by analogy with other pharmaceuticals.

As described in detail, pharmaceutical compositions of the present disclosure may be specially formulated for administration in solid or liquid form, including those adapted for the following: oral administration, for example, drenches (aqueous or non-aqueous solutions or suspensions), tablets, e.g., those targeted for buccal, sublingual, and systemic absorption, boluses, powders, granules, pastes for application to the tongue; parenteral administration, for example, by subcutaneous, intramuscular, intravenous or epidural injection as, for example, a sterile solution or suspension, or sustained-release formulation; topical application, for example, as a cream, ointment, or a controlled-release patch or spray applied to the skin, lungs, or oral cavity; intravaginally or intrarectally, for example, as a pessary, cream or foam; sublingually; ocularly; transdermally; or nasally, pulmonary and to other mucosal surfaces.

Wetting agents, emulsifiers and lubricants, such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, release agents, coating agents, sweetening, flavoring

and perfuming agents, preservatives and antioxidants can also be present in the compositions.

Examples of pharmaceutically acceptable antioxidants include: water soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite and the like; oil-soluble antioxidants, such as ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, alpha-tocopherol, and the like; and metal chelating agents, such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid, phosphoric acid, and the like.

Formulations for use in accordance with the present disclosure include those suitable for oral, nasal, topical (including buccal and sublingual), rectal, vaginal and/or parenteral administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. The amount of active ingredient, which can be combined with a carrier material, to produce a single dosage form will vary depending upon the host being treated, and the particular mode of administration. The amount of active ingredient that can be combined with a carrier material to produce a single dosage form will generally be that amount of the compound, which produces a therapeutic effect. Generally, this amount will range from about 1% to about 99% of active ingredient. In some embodiments, this amount will range from about 5% to about 70%, from about 10% to about 50%, or from about 20% to about 40%.

In certain embodiments, a formulation as described herein comprises an excipient selected from the group consisting of cyclodextrins, liposomes, micelle forming agents, e.g., bile acids, and polymeric carriers, e.g., polyesters and polyanhydrides; and a compound of the present disclosure. In certain embodiments, an aforementioned formulation renders orally bioavailable a described compound of the present disclosure.

Methods of preparing formulations or compositions comprising described compounds include a step of bringing into association a compound of the present disclosure with the carrier and, optionally, one or more accessory ingredients. In general, formulations may be prepared by uniformly and intimately bringing into association a compound of the present disclosure with liquid carriers, or finely divided solid carriers, or both, and then, if necessary, shaping the product.

The pharmaceutical compositions may be in the form of a sterile injectable preparation, for example, as a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to techniques known in the art using suitable dispersing or wetting agents (such as, for example, Tween 80) and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are mannitol, water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including synthetic mono- or diglycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as those described in Pharmacopeia Helvetica, or a similar alcohol. Other commonly used surfactants, such as Tweens, Spans and other

emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation.

In some cases, in order to prolong the effect of a drug, it may be desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material having poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution, which in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

Injectable depot forms are made by forming microcapsule matrices of the described compounds in biodegradable polymers such as polylactide-polyglycolide. Depending on the ratio of drug to polymer, and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions, which are compatible with body tissue.

The pharmaceutical compositions of this disclosure may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, and aqueous suspensions and solutions. In the case of tablets for oral use, carriers, which are commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried cornstarch. When aqueous suspensions and solutions and propylene glycol are administered orally, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening and/or flavoring and/or coloring agents may be added.

Formulations described herein suitable for oral administration may be in the form of capsules, cachets, pills, tablets, lozenges (using a flavored basis, usually sucrose and acacia or tragacanth), powders, granules, or as a solution or a suspension in an aqueous or non-aqueous liquid, or as an oil-in-water or water-in-oil liquid emulsion, or as an elixir or syrup, or as pastilles (using an inert base, such as gelatin and glycerin, or sucrose and acacia) and/or as mouth washes and the like, each containing a predetermined amount of a compound of the present disclosure as an active ingredient. Compounds described herein may also be administered as a bolus, electuary or paste.

In solid dosage forms for oral administration (capsules, tablets, pills, dragees, powders, granules and the like), an active ingredient is mixed with one or more pharmaceutically-acceptable carriers, such as sodium citrate or dicalcium phosphate, and/or any of the following: fillers or extenders, such as starches, lactose, sucrose, glucose, mannitol, and/or silicic acid; binders, such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinyl pyrrolidone, sucrose and/or acacia; humectants, such as glycerol; disintegrating agents, such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate; solution retarding agents, such as paraffin; absorption accelerators, such as quaternary ammonium compounds; wetting agents, such as, for example, cetyl alcohol, glycerol monostearate, and non-ionic surfactants; absorbents, such as kaolin and bentonite clay; lubricants, such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate,

and mixtures thereof; and coloring agents. In the case of capsules, tablets and pills, the pharmaceutical compositions may also comprise buffering agents. Solid compositions of a similar type may also be employed as fillers in soft and hard-shelled gelatin capsules using such excipients as lactose or milk sugars, as well as high molecular weight polyethylene glycols and the like.

Tablets may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared using binder (for example, gelatin or hydroxypropylmethyl cellulose), lubricant, inert diluent, preservative, disintegrant (for example, sodium starch glycolate or cross-linked sodium carboxymethyl cellulose), surface-active or dispersing agent. Molded tablets may be made in a suitable machine in which a mixture of the powdered compound is moistened with an inert liquid diluent. If a solid carrier is used, the preparation can be in tablet form, placed in a hard gelatin capsule in powder or pellet form, or in the form of a troche or lozenge. The amount of solid carrier will vary, e.g., from about 25 to 800 mg, preferably about 25 mg to 400 mg. When a liquid carrier is used, the preparation can be, e.g., in the form of a syrup, emulsion, soft gelatin capsule, sterile injectable liquid such as an ampule or nonaqueous liquid suspension. Where the composition is in the form of a capsule, any routine encapsulation is suitable, for example, using the aforementioned carriers in a hard gelatin capsule shell.

Tablets and other solid dosage forms, such as dragees, capsules, pills and granules, may optionally be scored or prepared with coatings and shells, such as enteric coatings and other coatings well known in the pharmaceutical-formulating art. They may alternatively or additionally be formulated so as to provide slow or controlled release of the active ingredient therein using, for example, hydroxypropylmethyl cellulose in varying proportions to provide the desired release profile, other polymer matrices, liposomes and/or microspheres. They may be formulated for rapid release, e.g., freeze-dried. They may be sterilized by, for example, filtration through a bacteria-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions that can be dissolved in sterile water, or some other sterile injectable medium immediately before use. These compositions may also optionally contain opacifying agents and may be of a composition that they release the active ingredient(s) only, or preferentially, in a certain portion of the gastrointestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes. The active ingredient can also be in micro-encapsulated form, if appropriate, with one or more of the above-described excipients.

Liquid dosage forms for oral administration of compounds of the disclosure include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active ingredient, the liquid dosage forms may contain inert diluents commonly used in the art, such as, for example, water or other solvents, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor and sesame oils), glycerol, tetrahydrofuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

Besides inert diluents, oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, coloring, perfuming and preservative agents.

Suspensions, in addition to active compounds, may contain suspending agents as, for example, ethoxylated isosteryl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, and mixtures thereof.

The pharmaceutical compositions of this disclosure may also be administered in the form of suppositories for rectal administration. These compositions can be prepared by mixing a compound of this disclosure with a suitable non-irritating excipient, which is solid at room temperature but liquid at the rectal temperature and therefore will melt in the rectum to release the active components. Such materials include, but are not limited to, cocoa butter, beeswax and polyethylene glycols.

Topical administration of the pharmaceutical compositions of this disclosure is especially useful when the desired treatment involves areas or organs readily accessible by topical application. For application topically to the skin, the pharmaceutical composition should be formulated with a suitable ointment containing the active components suspended or dissolved in a carrier. Carriers for topical administration of the compounds of this disclosure include, but are not limited to, mineral oil, liquid petroleum, white petroleum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, the pharmaceutical composition can be formulated with a suitable lotion or cream containing the active compound suspended or dissolved in a carrier. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water. The pharmaceutical compositions of this disclosure may also be topically applied to the lower intestinal tract by rectal suppository formulation or in a suitable enema formulation. Topically-administered transdermal patches are also included in this disclosure.

The pharmaceutical compositions of this disclosure may be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other solubilizing or dispersing agents known in the art.

For ophthalmic use, the pharmaceutical compositions may be formulated as micronized suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative such as benzylalkonium chloride. Alternatively, for ophthalmic uses, the pharmaceutical compositions may be formulated in an ointment such as petrolatum.

Transdermal patches have the added advantage of providing controlled delivery of a compound of the present disclosure to the body. Dissolving or dispersing the compound in the proper medium can make such dosage forms. Absorption enhancers can also be used to increase the flux of the compound across the skin. Either providing a rate controlling membrane or dispersing the compound in a polymer matrix or gel can control the rate of such flux.

Examples of suitable aqueous and nonaqueous carriers, which may be employed in the pharmaceutical compositions of the disclosure, include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of

coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

Such compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Inclusion of one or more antibacterial and/or antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like, may be desirable in certain embodiments. It may alternatively or additionally be desirable to include isotonic agents, such as sugars, sodium chloride, and the like into the compositions. In addition, prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents, which delay absorption such as aluminum monostearate and gelatin.

In certain embodiments, a described compound or pharmaceutical preparation is administered orally. In other embodiments, a described compound or pharmaceutical preparation is administered intravenously. Alternative routes of administration include sublingual, intramuscular, and transdermal administrations.

When compounds described herein are administered as pharmaceuticals, to humans and animals, they can be given per se or as a pharmaceutical composition containing, for example, 0.1% to 99.5% (more preferably, 0.5% to 90%) of active ingredient in combination with a pharmaceutically acceptable carrier.

Preparations described herein may be given orally, parenterally, topically, or rectally. They are of course given in forms suitable for the relevant administration route. For example, they are administered in tablets or capsule form, by injection, inhalation, eye lotion, ointment, suppository, etc. administration by injection, infusion or inhalation; topical by lotion or ointment; and rectal by suppositories. Oral administrations are preferred.

Such compounds may be administered to humans and other animals for therapy by any suitable route of administration, including orally, nasally, as by, for example, a spray, rectally, intravaginally, parenterally, intracisternally and topically, as by powders, ointments or drops, including buccally and sublingually.

Regardless of the route of administration selected, compounds described herein which may be used in a suitable hydrated form, and/or the pharmaceutical compositions of the present disclosure, are formulated into pharmaceutically-acceptable dosage forms by conventional methods known to those of skill in the art.

Actual dosage levels of the active ingredients in the pharmaceutical compositions of the disclosure may be varied so as to obtain an amount of the active ingredient that is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient.

The terms "administration of" and or "administering" should be understood to mean providing a pharmaceutical composition in a therapeutically effective amount to the subject in need of treatment. Administration routes can be enteral, topical or parenteral. As such, administration routes include but are not limited to intracutaneous, subcutaneous, intravenous, intraperitoneal, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, transdermal, transtracheal, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal and intrasternal, oral, sublingual buccal, rectal, vaginal, nasal ocular administrations, as well infusion, inhalation, and nebulization.

The term "cancer" refers to a group diseases characterized by abnormal and uncontrolled cell proliferation starting at

one site (primary site) with the potential to invade and to spread to others sites (secondary sites, metastases) which differentiate cancer (malignant tumor) from benign tumor. Virtually all the organs can be affected, leading to more than 100 types of cancer that can affect humans. Cancers can result from many causes including genetic predisposition, viral infection, exposure to ionizing radiation, exposure environmental pollutant, tobacco and or alcohol use, obesity, poor diet, lack of physical activity or any combination thereof.

Exemplary cancers described by the national cancer institute include: Acute Lymphoblastic Leukemia, Adult; Acute Lymphoblastic Leukemia, Childhood; Acute Myeloid Leukemia, Adult; Adrenocortical Carcinoma; Adrenocortical Carcinoma, Childhood; AIDS-Related Lymphoma; AIDS-Related Malignancies; Anal Cancer; Astrocytoma, Childhood Cerebellar; Astrocytoma, Childhood Cerebral; Bile Duct Cancer, Extrahepatic; Bladder Cancer; Bladder Cancer, Childhood; Bone Cancer, Osteosarcoma/Malignant Fibrous Histiocytoma; Brain Stem Glioma, Childhood; Brain Tumor, Adult; Brain Tumor, Brain Stem Glioma, Childhood; Brain Tumor, Cerebellar Astrocytoma, Childhood; Brain Tumor, Cerebral Astrocytoma/Malignant Glioma, Childhood; Brain Tumor, Ependymoma, Childhood; Brain Tumor, Medulloblastoma, Childhood; Brain Tumor, Supratentorial Primitive Neuroectodermal Tumors, Childhood; Brain Tumor, Visual Pathway and Hypothalamic Glioma, Childhood; Brain Tumor, Childhood (Other); Breast Cancer; Breast Cancer and Pregnancy; Breast Cancer, Childhood; Breast Cancer, Male; Bronchial Adenomas/Carcinoids, Childhood; Carcinoid Tumor, Childhood; Carcinoid Tumor, Gastrointestinal; Carcinoma, Adrenocortical; Carcinoma, Islet Cell; Carcinoma of Unknown Primary; Central Nervous System Lymphoma, Primary; Cerebellar Astrocytoma, Childhood; Cerebral Astrocytoma/Malignant Glioma, Childhood; Cervical Cancer; Childhood Cancers; Chronic Lymphocytic Leukemia; Chronic Myelogenous Leukemia; Chronic Myeloproliferative Disorders; Clear Cell Sarcoma of Tendon Sheaths; Colon Cancer; Colorectal Cancer, Childhood; Cutaneous T-Cell Lymphoma; Endometrial Cancer; Ependymoma, Childhood; Epithelial Cancer, Ovarian; Esophageal Cancer; Esophageal Cancer, Childhood; Ewing's Family of Tumors; Extracranial Germ Cell Tumor, Childhood; Extragonadal Germ Cell Tumor; Extrahepatic Bile Duct Cancer; Eye Cancer, Intraocular Melanoma; Eye Cancer, Retinoblastoma; Gallbladder Cancer; Gastric (Stomach) Cancer; Gastric (Stomach) Cancer, Childhood; Gastrointestinal Carcinoid Tumor; Germ Cell Tumor, Extracranial, Childhood; Germ Cell Tumor, Extragonadal; Germ Cell Tumor, Ovarian; Gestational Trophoblastic Tumor; Glioma, Childhood Brain Stem; Glioma, Childhood Visual Pathway and Hypothalamic; Hairy Cell Leukemia; Head and Neck Cancer; Hepatocellular (Liver) Cancer, Adult (Primary); Hepatocellular (Liver) Cancer, Childhood (Primary); Hodgkin's Lymphoma, Adult; Hodgkin's Lymphoma, Childhood; Hodgkin's Lymphoma During Pregnancy; Hypopharyngeal Cancer; Hypothalamic and Visual Pathway Glioma, Childhood; Intraocular Melanoma; Islet Cell Carcinoma (Endocrine Pancreas); Kaposi's Sarcoma; Kidney Cancer; Laryngeal Cancer; Laryngeal Cancer, Childhood; Leukemia, Acute Lymphoblastic, Adult; Leukemia, Acute Lymphoblastic, Childhood; Leukemia, Acute Myeloid, Adult; Leukemia, Acute Myeloid, Childhood; Leukemia, Chronic Lymphocytic; Leukemia, Chronic Myelogenous; Leukemia, Hairy Cell; Lip and Oral Cavity Cancer; Liver Cancer, Adult (Primary); Liver Cancer, Childhood (Primary); Lung Cancer, Non-Small Cell; Lung Cancer, Small Cell; Lymphoblas-

tic Leukemia, Adult Acute; Lymphoblastic Leukemia, Childhood Acute; Lymphocytic Leukemia, Chronic; Lymphoma, AIDS—Related; Lymphoma, Central Nervous System (Primary); Lymphoma, Cutaneous T-Cell; Lymphoma, Hodgkin's, Adult; Lymphoma, Hodgkin's; Childhood; Lymphoma, Hodgkin's During Pregnancy; Lymphoma, Non-Hodgkin's, Adult; Lymphoma, Non-Hodgkin's, Childhood; Lymphoma, Non-Hodgkin's During Pregnancy; Lymphoma, Primary Central Nervous System; Macroglobulinemia, Waldenström's; Male Breast Cancer; Malignant Mesothelioma, Adult; Malignant Mesothelioma, Childhood; Malignant Thymoma; Medulloblastoma, Childhood; Melanoma; Melanoma, Intraocular; Merkel Cell Carcinoma; Mesothelioma, Malignant; Metastatic Squamous Neck Cancer with Occult Primary; Multiple Endocrine Neoplasia Syndrome, Childhood; Multiple Myeloma/Plasma Cell Neoplasm; Mycosis Fungoides; Myelodysplasia Syndromes; Myelogenous Leukemia, Chronic; Myeloid Leukemia, Childhood Acute; Myeloma, Multiple; Myeloproliferative Disorders, Chronic; Nasal Cavity and Paranasal Sinus Cancer; Nasopharyngeal Cancer; Nasopharyngeal Cancer, Childhood; Neuroblastoma; Non-Hodgkin's Lymphoma, Adult; Non-Hodgkin's Lymphoma, Childhood; Non-Hodgkin's Lymphoma During Pregnancy; Non-Small Cell Lung Cancer; Oral Cancer, Childhood; Oral Cavity and Lip Cancer; Oropharyngeal Cancer; Osteosarcoma/Malignant Fibrous Histiocytoma of Bone; Ovarian Cancer, Childhood; Ovarian Epithelial Cancer; Ovarian Germ Cell Tumor; Ovarian Low Malignant Potential Tumor; Pancreatic Cancer; Pancreatic Cancer, Childhood; Pancreatic Cancer, Islet Cell; Paranasal Sinus and Nasal Cavity Cancer; Parathyroid Cancer; Penile Cancer; Pheochromocytoma; Pineal and Supratentorial Primitive Neuroectodermal Tumors, Childhood; Pituitary Tumor; Plasma Cell Neoplasm/Multiple Myeloma; Pleuropulmonary Blastoma; Pregnancy and Breast Cancer; Pregnancy and Hodgkin's Lymphoma; Pregnancy and Non-Hodgkin's Lymphoma; Primary Central Nervous System Lymphoma; Primary Liver Cancer, Adult; Primary Liver Cancer, Childhood; Prostate Cancer; Rectal Cancer; Renal Cell (Kidney) Cancer; Renal Cell Cancer, Childhood; Renal Pelvis and Ureter, Transitional Cell Cancer; Retinoblastoma; Rhabdomyosarcoma, Childhood; Salivary Gland Cancer; Salivary Gland Cancer, Childhood; Sarcoma, Ewing's Family of Tumors; Sarcoma, Kaposi's; Sarcoma (Osteosarcoma/Malignant Fibrous Histiocytoma of Bone; Sarcoma, Rhabdomyosarcoma, Childhood; Sarcoma, Soft Tissue, Adult; Sarcoma, Soft Tissue, Childhood; Sezary Syndrome; Skin Cancer; Skin Cancer, Childhood; Skin Cancer (Melanoma); Skin Carcinoma, Merkel Cell; Small Cell Lung Cancer; Small Intestine Cancer; Soft Tissue Sarcoma, Adult; Soft Tissue Sarcoma, Childhood; Squamous Neck Cancer with Occult Primary, Metastatic; Stomach (Gastric) Cancer; Stomach (Gastric) Cancer, Childhood; Supratentorial Primitive Neuroectodermal Tumors, Childhood; T-Cell Lymphoma, Cutaneous; Testicular Cancer; Thymoma, Childhood; Thymoma, Malignant; Thyroid Cancer; Thyroid Cancer, Childhood; Transitional Cell Cancer of the Renal Pelvis and Ureter; Trophoblastic Tumor, Gestational; Unknown Primary Site, Cancer of, Childhood; Unusual Cancers of Childhood; Ureter and Renal Pelvis, Transitional Cell Cancer; Urethral Cancer; Uterine Sarcoma; Vaginal Cancer; Visual Pathway and Hypothalamic Glioma, Childhood; Vulvar Cancer; Waldenström's Macroglobulinemia; and Wilms' Tumor.

In certain aspects, cancer include Lung cancer, Breast

cancer, Non-Hodgkin lymphoma, Leukemia, Pancreatic cancer, Kidney cancer, endometrial cancer, Head and neck cancer, Lip cancer, oral cancer, Thyroid cancer, Brain cancer, Ovary cancer, Melanoma, Gallbladder cancer, Laryngeal cancer, Multiple myeloma, Nasopharyngeal cancer, Hodgkin lymphoma, Testis cancer and Kaposi sarcoma.

In certain aspects, the method further includes administering a chemotherapeutic agent. The compounds of the disclosure can be administered in combination with one or more additional therapeutic agents. The phrases “combination therapy”, “combined with” and the like refer to the use of more than one medication or treatment simultaneously to increase the response. The FGFR inhibitor of the present disclosure might for example be used in combination with other drugs or treatment in use to treat cancer. In various aspect, the compound is administered prior to, simultaneously with or following the administration of the chemotherapeutic agent.

The term “anti-cancer therapy” refers to any therapy or treatment that can be used for the treatment of a cancer. Anti-cancer therapies include, but are not limited to, surgery, radiotherapy, chemotherapy, immune therapy and targeted therapies.

Examples of chemotherapeutic agents or anti-cancer agents include, but are not limited to, Actinomycin, Azacitidine, Azathioprine, Bleomycin, Bortezomib, Carboplatin, Capecitabine, Cisplatin, Chlorambucil, Cyclophosphamide, Cytarabine, Daunorubicin, Docetaxel, Doxifluridine, Doxorubicin, Epirubicin, Epothilone, Etoposide, Fluorouracil, Gemcitabine, Hydroxyurea, Idarubicin, Imatinib, Irinotecan, Mechlorethamine, Mercaptopurine, Methotrexate, Mitoxantrone, Oxaliplatin, Paclitaxel, Pemetrexed, Teniposide, Tioguanine, Topotecan, Valrubicin, Vinblastine, Vincristine, Vindesine, Vinorelbine, panitumumab, Erbitux (cetuximab), matuzumab, IMC-IIF 8, TheraCIM hR3, denosumab, Avastin (bevacizumab), Humira (adalimumab), Herceptin (trastuzumab), Remicade (infliximab), rituximab, Synagis (palivizumab), Mylotarg (gemtuzumab oxogamicin), Raptiva (efalizumab), Tysabri (natalizumab), Zenapax (daclizumab), NeutroSpec (Technetium 99mTc) fanolesomab, tocilizumab, ProstaScint (Indium-111 labeled Capromab Pentetide), Bexxar (tositumomab), Zevalin (ibritumomab tiuxetan (IDEC-Y2B8) conjugated to yttrium 90), Xolair (omalizumab), MabThera (Rituximab), ReoPro (abciximab), MabCampath (alemtuzumab), Simulect (basiliximab), LeukoScan (sulesomab), CEA-Scan (arcitumomab), Verluma (nofetumomab), Panorex (Edrecolomab), alemtuzumab, CDP 870, natalizumab Gilotrif (afatinib), Lynparza (olaparib), Perjeta (pertuzumab), Otdivo (nivolumab), Bosulif (bosutinib), Cabometyx (cabozantinib), Ogivri (trastuzumab-dkst), Sutent (sunitinib malate), Adcetris (brentuximab vedotin), Alecensa (alectinib), Calquence (acalabrutinib), Yescarta (ciloleucel), Verzenio (abemaciclib), Keytruda (pembrolizumab), Aliqopa (copanlisib), Nerlynx (neratinib), Imfinzi (durvalumab), Darzalex (daratumumab), Tecentriq (atezolizumab), and Tarceva (erlotinib). Examples of immunotherapeutic agent include, but are not limited to, interleukins (IL-2, IL-12), cytokines (Interferons, G-CSF, imiquimod), chemokines (CCL3, CC126, CXCL7), immunomodulatory imide drugs (thalidomide and its analogues).

In a further embodiment, the disclosure provides a method of inhibiting a kinase activity including contacting a cell with a compound of Formula (I), Formula (II), or Formula (III). In one aspect, the kinase is selected from the group consisting of ALK, EGFR, EPH-B3, FAK, FGFR1, FGFR2, FGFR3, FGFR4, KIT, MEK1, MET, PDGFR-AL-

49

PHA, PDGFR-BETA, RET, ROS and TYRO 3. In certain aspects, the kinase is FGFR1, FGFR2, FGFR3 and/or FGFR4. In another aspect, the kinase is FGFR4. In various aspects, the cell is a cancer cell. In many aspects, the cancer cell is a breast, lung, bladder, prostate, ovarian, endometrial, rhabdomyosarcoma, liver or gastric cancer cell.

Referring to FIG. 1, the FGFR4 kinase contains a cysteine (Cys552) located near the ATP-binding site, in the hinge region of the receptor, which is unique within the FGFR family and rare among other kinases. Covalent inhibitors of FGFR4 kinase afford potent and selective inhibition of FGFRs by covalently targeting a thiol group (SH) in the cysteine residue.

The screening method employed in the current disclosure according to some embodiments is described as following. Screen probes against live cells or live animals can include the step of (1) treatment of live cells with an electrophilic probe leads to the covalent linkage between target proteins and the probe; (2) lyse cells; (3) add biotin-azide to conjugate biotin to probe-modified proteins via the Copper-Catalyzed Azide-Alkyne Cycloaddition (CuAAC) click reaction; (4) use streptavidin resin to pull down and enrich biotin-conjugated proteins; (5) digest resin-bound proteins with trypsin; (6) analyze resulting tryptic peptides with liquid chromatography-mass spectrometry (LC-MS) or MS; (7) obtain information on identity of target proteins as well as strength of probe-target interaction in cells. The target confirmation and selection can include the steps of (1) use orthogonal methods such as western blots to confirm that the identified proteins indeed bind to the probe; and (2) select the most disease-relevant target-lead pairs for further pre-clinical and clinical studies.


Presented below are examples discussing the design and evaluation of efficacy of new pyridinylpyrimidine-based FGFR inhibitors, contemplated for the discussed applications. The following examples are provided to further illustrate the embodiments of the present disclosure, but are not intended to limit the scope of the disclosure. While they are typical of those that might be used, other procedures, methodologies, or techniques known to those skilled in the art may alternatively be used.

EXAMPLES

Example 1


Synthesis and Characterization of the Pyridinylpyrimidine-Based FGFR Inhibitors (Compounds 1-4)

Scheme 1. Synthesis of compound 1.




50

-continued



51


-continued




compound 1

Compound 1 was synthesized based on synthetic scheme 1.

Scheme 2. Synthesis of compound 2.

**52**


-continued



compound 2





Compound 2 was synthesized based on synthetic scheme 2.

Scheme 3. Synthesis of compound 3.



53


-continued








compound 3

Compound 3 was synthesized based on synthetic scheme 3.

Scheme 4. Synthesis of compound 4.


**54**

-continued



55

-continued



compound 4

56


Compound 4 was synthesized based on synthetic scheme 4.

Example 2

Synthesis of Compound 6


(N-[2-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]-5-(1-methylazetidin-3-yl)oxy-phenyl]prop-2-enamide)

Scheme 5. Synthesis of compound 6.




57

58




Compound 6 was synthesized based on synthetic scheme 5. The synthetic process for each of the step is described as below.

55



-continued



60


Step 1 to synthesize compound 6: A solution of 4-amino-3-nitro-phenol (3 g, 19.46 mmol), tert-butyl 3-iodoazetidine-1-carboxylate (6.06 g, 21.41 mmol), Cs_2CO_3 (9.51 g, 29.20 mmol) in DMF (60 mL) was stirred at 90° C. for 16 hours under N_2 atmosphere. The mixture was diluted with water (5

59


mL), extracted with EtOAc (10 mL×3), then the combined organic layers were washed with sat. aq. NaCl (5 mL×7), dried over Na₂SO₄, filtered and concentrated. The residue was purified by silica gel chromatography (PE/EtOAc=1/0 to 3/1) to afford tert-butyl 3-(4-amino-3-nitro-phenoxy)aze-

60

tidine-1-carboxylate (5.4 g, 17.46 mmol, 90% yield) as a red solid. ¹H NMR (400 MHz, DMSO-d₆): δ=7.28 (d, J=4.0 Hz, 1H), 7.18-7.14 (m, 2H), 7.02 (d, J=9.2 Hz, 1H), 4.97-4.92 (m, 1H), 4.26-4.23 (m, 2H), 3.79-3.76 (m, 2H), 1.38 (s, 9H).




Step 2 to synthesize compound 6: To a solution of N-(2,6-dichloro-3,5-dimethoxy-phenyl)-3-(6-methylsulfonylpyrimidin-4-yl)pyridin-2-amine (800 mg, 1.76 mmol), tert-butyl 3-(4-amino-3-nitro-phenoxy)azetidine-1-carboxylate (652 mg, 2.11 mmol) in DMAc (15 mL) was added t-BuOK (595 mg, 5.31 mmol). The mixture was stirred at 45° C. for 2 hours. The mixture was diluted with water (20 mL), extracted with EtOAc (30 mL×3), then the combined organic layers were washed with sat. aq. NaCl (20 mL×7), dried over Na₂SO₄, filtered and concentrated. The residue was purified by silica gel chromatography (PE/EtOAc=1/0 to 3/1) to afford tert-butyl 3-[4-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]-3-nitro-phenoxy]azetidine-1-carboxylate (648 mg, 0.947 mmol, 54% yield) as a yellow solid.



61

62




6-5


15

Step 3 to synthesize compound 6: To a solution of tert-butyl 3-[4-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]-3-nitro-phenoxy]azetidine-1-carboxylate (648 mg, 0.947 mmol) in dioxane (10 mL) was added HCl/dioxane (4 M, 2.37 mL). The mixture was stirred at 25° C. for 2 hours. The mixture was concentrated to afford N-[4-(azetidin-3-yloxy)-2-nitro-phenyl]-6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-amine (550 mg, 0.886 mmol, 93% yield, HCl salt) as a yellow solid.

20



6-5



6-6

Step 4 to synthesize compound 6: To a solution of N-[4-(azetidin-3-yloxy)-2-nitro-phenyl]-6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-amine (550 mg, 0.886 mmol, HCl salt) in MeOH (15 mL) was added aqueous HCHO (2.88 g, 35.43 mmol, 2.64 mL, 37% purity), NaBH(OAc)₃ (376 mg, 1.77 mmol), CH₃COOH (213 mg, 3.54 mmol). The mixture was stirred at 25° C. for 4 hours. The mixture was concentrated and the residue was purified by silica gel chromatography (DCM/MeOH=1/0 to 10/1) to afford 6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]-N-[4-(1-methylazetidin-3-yloxy)-2-nitro-phenyl]pyrimidin-4-amine (450 mg, 0.752 mmol, 85% yield) as a red solid.



55

60

65

63


64




6-7

Step 5 to synthesize compound 6: To a solution of 6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]-N-[4-

zene-1,2-diamine (400 mg, 0.704 mmol, 93% yield) as a yellow solid.



6-7



compound 6

60

(1-methylazetidin-3-yl)oxy-2-nitro-phenyl]pyrimidin-4-amine (450 mg, 0.752 mmol) in THE (15 mL) and H₂O (15 mL) was added NH₄Cl (201 mg, 3.76 mmol), Fe (210 mg, 3.76 mmol). The mixture was stirred at 65° C. for 4 hours. The mixture was filtered and the filtrate was concentrated to afford N₁-[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]-4-(1-methylazetidin-3-yl)oxy-ben-

65

Step 6 to synthesize compound 6: To a solution of N₁-[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]-4-(1-methylazetidin-3-yl)oxy-benzene-1,2-diamine (200 mg, 0.352 mmol) in DCM (10 mL) was added Et₃N (36 mg, 0.352 mmol) and prop-2-enoyl chloride (32 mg, 0.352 mmol). The mixture was stirred at 25° C. for 0.5 hour. The mixture was diluted with water (10 mL), extracted

65

with DCM (10 mL×3), then the combined organic layers were dried over Na₂SO₄, filtered and concentrated. The residue was purified by preparative HPLC (water (0.225% FA)-ACN) to afford N-[2-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]-5-(1-methylazetidin-3-yl)oxy-phenyl]prop-2-enamide (8.5 mg, 4% yield, 96% purity) as a white solid. LCMS: t_R=2.928 min in 0-60CD_4 min_Pos_220&254_Shimadzu.lcm, MS (ESI) m/z=622.2 [M+H]⁺. ¹H NMR (400 MHz, DMSO-d₆): δ=10.93 (s, 1H), 9.68 (s, 1H), 9.02 (s, 1H), 8.61 (s, 1H), 8.39 (s, 1H), 8.06-7.95 (m, 2H), 7.41-7.31 (m, 2H), 6.86-6.83 (m, 2H), 6.69-6.66 (m, 2H), 6.25 (d, J=2.0 Hz, 1H), 5.73-5.70

66


(m, 1H), 4.76-4.70 (m, 1H), 3.93 (s, 6H), 3.76-3.72 (m, 2H), 3.01-2.98 (m, 2H), 2.30 (s, 3H).

Example 3

Synthesis of Compound 7


N-[2-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]-5-(1-methylazetidin-3-yl)methoxy]phenyl]prop-2-enamide

Scheme 6. Synthesis of compound 7.




67

68



Compound 7 was synthesized based on synthetic scheme 6. The synthetic process for each of the step is described as below.



Step 1 to synthesize compound 7: To a solution of tert-butyl 3-(hydroxymethyl)azetidine-1-carboxylate (2 g, 10.68 mmol) in DCM (5 mL) was added DABCO (2.40 g, 21.36 mmol) and TsCl (2.44 g, 12.82 mmol). The mixture was stirred at 0° C. for 0.5 hr. The mixture was filtered and the filtrate was diluted with H₂O (50 mL) and extracted with DCM (50 mL×3). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford tert-butyl 3-(p-tolylsulfonyloxymethyl)azetidine-1-carboxylate (3 g, 8.79 mmol, 82% yield) as a colorless oil, which was used directly in the next step.


Step 2 to synthesize compound 7: To a solution of tert-butyl 3-(p-tolylsulfonyloxymethyl)azetidine-1-carboxylate (2 g, 5.86 mmol) in DMF (20 mL) was added Cs₂CO₃ (3.82 g, 11.72 mmol) and 4-amino-3-nitrophenol (1.35 g, 8.79 mmol). The mixture was stirred at 80° C. for 12 hr. The mixture was quenched by addition of H₂O (50 mL) at 25° C.,

69

and then diluted with H₂O (50 mL) and extracted with EtOAc (50 mL×3). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. After that the residue was purified by flash silica gel chromatography (ISCO®; 12 g SepaFlash® Silica Flash Column, Eluent of 0~50% Ethyl acetate/Petroleum ether gradient @ 40 mL/min) to afford tert-butyl 3-[(4-amino-3-nitro-phenoxy)methyl]azetidine-1-carboxylate (1.4 g, 4.33 mmol, 74% yield) as a brown solid.


70

pyridin-2-amine (1.97 g, 4.33 mmol). The mixture was stirred at 40° C. for 2 hr. The mixture was quenched by addition of H₂O (50 mL) at 25° C., and then diluted with H₂O (50 mL) and extracted with EtOAc (50 mL×3). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. After that the residue was purified by flash silica gel chromatography (ISCO®; 12 g SepaFlash® Silica Flash Column, Eluent of 0~70% Ethyl acetate/




Step 3 to synthesize compound 7: To a solution of tert-butyl 3-[(4-amino-3-nitro-phenoxy)methyl]azetidine-1-carboxylate (1.4 g, 4.33 mmol) in DMF (5 mL) was added t-BuOK (583 mg, 5.20 mmol) and N-(2,6-dichloro-3,5-dimethoxy-phenyl)-3-(6-methylsulfonylpyrimidin-4-yl)

Petroleum ether gradient @ 40 mL/min) to afford tert-butyl 3-[[4-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]-3-nitro-phenoxy]methyl]azetidine-1-carboxylate (1 g, 1.29 mmol, 30% yield, 90% purity) as a yellow solid.





71

72



7-6

Step 4 to synthesize compound 7: A solution of tert-butyl 3-[[4-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]-3-nitro-phenoxymethyl]azetidine-1-carboxylate (0.95 g, 1.36 mmol) in HCl/dioxane (4 M, 10 mL) was stirred at 25° C. for 0.5 hr. The mixture was concentrated under reduced pressure to afford N-[4-(azetidin-3-ylmethoxy)-2-nitro-phenyl]-6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-amine (1.17 g, 1.35 mmol, 99% yield, 69% purity) as a yellow solid, which was used directly in the next step.




7-7

Step 5 to synthesize compound 7: To a solution of N-[4-(azetidin-3-ylmethoxy)-2-nitro-phenyl]-6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-amine (1 g, 1.22 mmol) in MeOH (10 mL) was added aqueous HCHO (3.96 g, 48.79 mmol, 3.63 mL, 37% purity), CH₃COOH (293 mg, 4.88 mmol) and NaBH(OAc)₃ (517 mg, 2.44 mmol). The mixture was stirred at 25° C. for 4 hr. The mixture was concentrated under reduced pressure to give a residue. After that the residue was purified by flash silica gel chromatography (ISCO®; 12 g SepaFlash® Silica Flash Column, Eluent of 0~80% Ethyl acetate/Petroleum ether gradient @ 60 mL/min) to afford 6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]-N-[4-(1-methylazetidin-3-ylmethoxy)-2-nitro-phenyl]pyrimidin-4-amine (0.5 g, 0.735 mmol, 60% yield, 90% purity) as a yellow solid.


73

74



Step 6 to synthesize compound 7: To a solution of 6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]-N-[4-[(1-methylazetidin-3-yl)methoxy]-2-nitro-phenyl]pyrimidin-4-amine (0.5 g, 0.816 mmol) in THE (1 mL) and H₂O (1 mL) was added Fe (228 mg, 4.08 mmol) and NH₄C₁ (218 mg, 4.08 mmol). The mixture was stirred at 60° C. for 5 hr. The mixture was filtered and the filtrate was concentrated under reduced pressure to give a residue. After that the

30 mixture was quenched by addition of H₂O (50 mL) at 25° C., and then diluted with H₂O (50 mL) and extracted with EtOAc (50 mL×3). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford N₁-[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]-4-[(1-methylazetidin-3-yl)methoxy]benzene-1,2-diamine (350 mg, 0.60 mmol, 74% yield) as a brown solid.



75

Step 7 to synthesize compound 7: To a solution of N₁-[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]-4-[(1-methylazetidin-3-yl)methoxy]benzene-1,2-diamine (0.2 g, 0.343 mmol) in DCM (5 mL) was added DIEA (133 mg, 1.03 mmol) and prop-2-enoyl chloride (34 mg, 0.378 mmol). The mixture was stirred at 0° C. for 0.5 hr. The mixture was quenched by addition of H₂O (50 mL) at 25° C., and then diluted with H₂O (50 mL) and extracted with DCM (50 mL×3). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. After that the residue was purified by preparative HPLC (water (0.225% FA)-ACN) to afford N-[2-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]-5-[(1-methylazetidin-3-yl)methoxy]phenyl]prop-2-enamide (4.8 mg, 2% yield, 92.8% purity) as a white solid.

76


LCMS: t_R =1.425 min in 10-80AB_4 min_220&254_Shimadzu.lcm, MS (ESI) m/z =636.3 [M+H]⁺. ¹H NMR (400 MHz, Methanol): δ =8.62 (s, 1H), 8.50 (s, 1H), 8.00-7.98 (m, 2H), 7.58-7.44 (m, 2H), 7.02-6.78 (m, 4H), 6.45-6.33 (m, 2H), 5.76 (d, J=9.6 Hz, 1H), 4.35-4.14 (m, 6H), 3.96 (s, 6H), 3.24, 2.98 (s, 3H), 2.61-2.57 (m, 1H).

Example 4

Synthesis of Compound 8


N-[2-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]-5-[(1-methyl-4-piperidyl)oxy]phenyl]prop-2-enamide

Scheme 7. Synthesis of compound 8.




77

78






8-6




compound 8

Compound 8 was synthesized based on synthetic scheme 7. The synthetic process for each of the step is described as below.



8-2




8-4

Step 1 to synthesize compound 8: To a solution of 1-methylpiperidin-4-ol (1.3 g, 11.29 mmol) in THF (10 mL) was added KOH (2.53 g, 45.15 mmol) and TsCl (3.23 g, 16.93 mmol) at 0° C. The mixture was stirred at 25° C. for 16 hr. The mixture was poured into water (30 ml) and filtered; the filter cake was collected, dried to afford (1-methyl-4-piperidyl) 4-methylbenzenesulfonate (1.8 g, 6.68 mmol, 59% yield) as a yellow oil.

Step 2 to synthesize compound 8: To a solution of 4-amino-3-nitro-phenol (400 mg, 2.60 mmol) and (1-methyl-4-piperidyl) 4-methylbenzenesulfonate (1.05 g, 3.89 mmol) in DMF (20 mL) was added Cs₂CO₃ (1.69 g, 5.19 mmol). The mixture was stirred at 80° C. for 8 hr. The residue was diluted with EtOAc (20 mL) and extracted with H₂O (60 mL). The combined organic layers were washed with H₂O (30 mL×2), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by flash silica gel chromatography (ISCO®; 12 g SepaFlash® Silica Flash Column, Eluent of 0-10% methanol/dichloromethane @ 40 mL/min) to afford 5-[(1-methyl-4-piperidyl)oxy]-2-nitro-aniline (230 mg, 0.915 mmol, 35% yield) as a brown solid.


79

80



Step 3 to synthesize compound 8: To a solution of 4-[(1-methyl-4-piperidyl)oxy]-2-nitro-aniline (138 mg, 0.549 mmol) and N-(2,6-dichloro-3,5-dimethoxy-phenyl)-3-(6-methylsulfonylpyrimidin-4-yl)pyridin-2-amine (300 mg, 0.659 mmol) in DMAc (10 mL) was added tBuOK (123 mg, 1.10 mmol). The mixture was stirred at 50° C. for 2 hr. The residue was diluted with H₂O (20 mL) and extracted with EtOAc (20 mL×3). The combined organic layers were


washed with H₂O (30 mL×2), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by flash silica gel chromatography (ISCO®; 12 g SepaFlash® Silica Flash Column, Eluent of 0~10% methanol/dichloromethane @ 40 mL/min) to afford 6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]-N-[4-[(1-methyl-4-piperidyl)oxy]-2-nitro-phenyl]pyrimidin-4-amine (200 mg, 0.16 mmol, 29% yield) as a brown solid.




81

82

Step 4 to synthesize compound 8: To a solution of 6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]-N-[4-[(1-methyl-4-piperidyl)oxy]-2-nitro-phenyl]pyrimidin-4-amine (200 mg, 0.16 mmol) in THF (10 mL) and H₂O (10 mL) was added NH₄C_l (43 mg, 0.8 mmol) and Fe (45 mg, 0.8 mmol). The mixture was stirred at 65° C. for 3 hr. The mixture was filtered with diatomite, and the filtrate was concentrated under reduced pressure to give a residue. The residue was purified by flash silica gel chromatography (ISCO®; 12 g SepaFlash® Silica Flash Column, Eluent of 0~10% Methanol/Dichloromethane @ 40 mL/min) to afford N₁-[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]-4-[(1-methyl-4-piperidyl)oxy]benzene-1,2-diamine (38 mg, 0.064 mmol, 40% yield) as a yellow solid.



8-6




compound 8

Step 5 to synthesize compound 8: To a solution of N₁-[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]-4-[(1-methyl-4-piperidyl)oxy]benzene-1,2-diamine (38 mg, 0.064 mmol) in DCM (10 mL) and DIEA (16 mg, 0.127 mmol) was added prop-2-enoyl chloride (7 mg, 0.076 mmol). The mixture was stirred at 0° C. for 0.5 hr. The mixture was concentrated under reduced pressure to give the residue. The residue was purified by preparative HPLC (water (0.225% FA)-ACN) to afford N-[2-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]-5-[(1-methyl-4-piperidyl)oxy]phenyl]prop-2-enamide (11.1 mg, 0.017 mmol, 27% yield, 99.30% purity) as a yellow solid.

Synthesis of Compound 9


N-[2-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]-5-(1-methylpyrrolidin-3-yl)oxy-phenyl]prop-2-enamide

Scheme 8. Synthesis of compound 9.




85

-continued





compound 9

Compound 9 was synthesized based on synthetic scheme 8. The synthetic process for each of the step is described as below.




Step 1 to synthesize compound 9: To a solution of 1-methylpyrrolidin-3-ol (1.0 g, 9.89 mmol) and KOH (2.22 g, 39.55 mmol) in THE (30 mL) was added TsCl (2.83 g, 14.83 mmol) at 0° C. The mixture was stirred at 25° C. for 16 hours. The mixture was filtered and the filtrate was concentrated to afford (1-methylpyrrolidin-3-yl) 4-methylbenzenesulfonate (1.1 g, 4.31 mmol, 44% yield) as yellow oil.




86

-continued




Step 2 to synthesize compound 9: To a solution of 4-amino-3-nitro-phenol (600 mg, 3.89 mmol) in DMF (20 mL) was added Cs₂CO₃ (2.54 g, 7.79 mmol). The mixture was stirred at 25° C. for 0.5 hour, then (1-methylpyrrolidin-3-yl) 4-methylbenzenesulfonate (1.09 g, 4.28 mmol) was added thereto. The mixture was stirred at 80° C. for 6 hours. The mixture was diluted with water (20 mL), extracted with EtOAc (50 mL×3), then the combined organic layers were washed with sat. aq. NaCl (20 mL×7), dried over Na₂SO₄, filtered and concentrated. The residue was purified by silica gel chromatography (DCM/MeOH=1/0 to 10/1) to afford 4-(1-methylpyrrolidin-3-yl)oxy-2-nitro-aniline (600 mg, 2.53 mmol, 65% yield) as a red solid.



tBuOK, DMAc

87


88




9-5

Step 3 to synthesize compound 9: To a solution of 4-(1-methylpyrrolidin-3-yl)oxy-2-nitro-aniline (200 mg, 0.843 mmol), N-(2,6-dichloro-3,5-dimethoxy-phenyl)-3-(6-methylsulfonylpyrimidin-4-yl)pyridin-2-amine (422 mg, 0.927 mmol) in DMAC (10 mL) was added t-BuOK (286 mg, 2.55 mmol). The mixture was stirred at 40° C. for 2 hours. The mixture was diluted with water (20 mL), extracted with EtOAc (50 mL×3), then the combined organic layers were dried over Na₂SO₄, filtered and concentrated to afford 6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]-N-[4-(1-methylpyrrolidin-3-yl)oxy-2-nitro-phenyl]pyrimidin-4-amine (500 mg, crude) as yellow solid.

Step 4 to synthesize compound 9: To a solution of 6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]-N-[4-(1-methylpyrrolidin-3-yl)oxy-2-nitro-phenyl]pyrimidin-4-amine (300 mg, 0.49 mmol) in THF (10 mL) and H₂O (10 mL) was added Fe (137 mg, 2.45 mmol) and NH₄Cl (131 mg, 2.45 mmol). The mixture was stirred at 65° C. for 4 hours. The mixture was filtered and the filtrate was extracted with EtOAc (30 mL×3), then the combined organic layers were dried over Na₂SO₄, filtered and concentrated to afford N₁-[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]-4-(1-methylpyrrolidin-3-yl)oxy-benzene-1,2-diamine (150 mg, 0.258 mmol, 52% yield) as a yellow solid.




9-5




9-6

89

90



9-6



compound 9

40

Step 5 to synthesize compound 9: To a solution of N₁-[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl] 45
 pyrimidin-4-yl]-4-(1-methylpyrrolidin-3-yl)oxy-benzene-1, 2-diamine (120 mg, 0.206 mmol) in DCM (10 mL) was
 added prop-2-enoyl chloride (19 mg, 0.206 mmol) and DIEA (27 mg, 0.206 mmol). The mixture was stirred at 25°
 C. for 1 hour. The mixture was diluted with water (5 mL), 50
 extracted with DCM (10 mL×3), then the combined organic
 layers were dried over Na₂SO₄, filtered and concentrated.
 The residue was purified by preparative HPLC (water
 (0.225% FA)-ACN) to afford N-[2-[[6-[2-(2,6-dichloro-3,5-
 dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]-5-(1- 55
 methylpyrrolidin-3-yl)oxy-phenyl]prop-2-enamide (50 mg,
 0.0778 mmol, 38% yield, 99% purity) as yellow solid.
 LCMS: t_R =1.129 min in 10-80AB_4
 min_220&254_Shimadzu.lcm, MS (ESI) m/z =636.2 60
 [M+H]⁺. ¹H NMR (400 MHz, DMSO-d₆): δ=10.88 (s, 1H),
 10.15 (s, 1H), 9.65 (s, 1H), 9.10 (s, 1H), 8.64 (s, 1H), 8.06
 (d, J=4.4 Hz, 1H), 7.96 (d, J=7.6 Hz, 1H), 7.54 (s, 1H), 7.46
 (d, J=8.4 Hz, 1H), 7.01-6.98 (m, 1H), 6.88-6.83 (m, 3H),
 6.52-6.49 (m, 1H), 6.26-6.21 (m, 1H), 5.74-5.71 (m, 1H), 65
 5.17-5.12 (m, 1H), 3.93 (s, 6H), 3.81-3.61 (m, 1H), 3.41-
 2.89 (m, 6H), 2.61-2.54 (m, 1H), 2.30-2.10 (m, 1H).

91

Example 6


Synthesis of Compound 10

N-[2-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]-5-tetrahydrofuran-3-yloxy-phenyl]prop-2-enamide

5


92

Scheme 9. Synthesis of compound 10.




93

Compound 10 was synthesized based on synthetic scheme 9. The synthetic process for each of the step is described as below.



94

Step 1 to synthesize compound 10: To a solution of 4-amino-3-nitro-phenol (4 g, 25.95 mmol), tetrahydrofuran-3-ol (2.29 g, 25.95 mmol) and PPh₃ (8.17 g, 31.14 mmol) in THE (25 mL) was added a solution of DTBAD (8.96 g, 38.93 mmol) in THE (25 mL) dropwise at 0° C. under N₂. The mixture was stirred at 0° C. for 0.5 h then warmed to 25° C. for 15.5 h. The mixture was concentrated. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate=5/1 to 3/1) and the residue was recrystallized from CH₃CN to afford 2-nitro-4-tetrahydrofuran-3-yloxy-aniline (1.1 g, 4.78 mmol, 18% yield, 97.5% purity) as a red solid. LCMS: t_R=0.824 min in 0-30AB_4 min_220&254_Shimadzu.lcm, MS (ESI) m/z=225.2 [M+H]⁺. ¹H NMR (400 MHz, CDCl₃): δ=7.44 (s, 1H), 7.00-6.95 (m, 1H), 6.72-6.66 (m, 1H), 5.84 (s, 2H), 4.84-4.80 (m, 1H), 3.93-3.83 (m, 4H), 2.19-2.03 (m, 2H).





55

Step 2 to synthesize compound 10: To a solution of N-(2,6-dichloro-3,5-dimethoxy-phenyl)-3-(6-methylsulfonylpyrimidin-4-yl)pyridin-2-amine (300 mg, 0.66 mmol) and 2-nitro-4-tetrahydrofuran-3-yloxy-aniline (296 mg, 1.32 mmol) in DMAc (30 mL) was added tBuOK (148 mg, 1.32 mmol). The mixture was stirred at 40° C. for 2 hr. The mixture was poured into water (60 mL) and filtered, the filter cake was collected in vacuo to afford 6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]-N-(2-nitro-4-tetrahydrofuran-3-yloxy-phenyl)pyrimidin-4-amine (240 mg, 0.40 mmol, 61% yield) as a yellow solid.

95


96




10-5

Step 3 to synthesize compound 10: To a solution of 6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]-N-(2-nitro-4-tetrahydrofuran-3-yloxy-phenyl)pyrimidin-4-amine (240 mg, 0.40 mmol) in THF (10 mL) and H₂O (10 mL) was added Fe (112 mg, 2.00 mmol) and NH₄Cl (107 mg, 2.00 mmol). The mixture was stirred at 65° C. for 3 hr. The mixture was filtered with diatomite, and the filtrate was

concentrated under reduced pressure to afford N₁-[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]-4-tetrahydrofuran-3-yloxy-benzene-1,2-diamine (300 mg, 0.368 mmol, 92% yield, 90% purity) as a red solid. LCMS: t_R =0.922 min in 0-60AB_2 min_220&254_Shimadzu.lcm, MS (ESI) m/z =569.1 [M+H]⁺.



10-5



compound 10

97

Step 4 to synthesize compound 10: To a solution of N₁-[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]-4-tetrahydrofuran-3-yloxy-benzene-1,2-diamine (220 mg, 0.386 mmol) in DCM (10 mL) and DIEA (100 mg, 0.772 mmol) and prop-2-enoyl chloride (31 mg, 0.348 mmol) at 0° C. The mixture was stirred at 0° C. for 0.5 hr. The mixture was concentrated under reduced pressure. The residue was purified by preparative HPLC (water (0.225% FA)-ACN) to afford N-[2-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]-5-tetrahydrofuran-3-yloxy-phenyl]prop-2-enamide (100 mg, 0.159 mmol, 41% yield, 99.2% purity) as a yellow solid. LCMS: t_R =2.148 min in 0-60AB_4 min_220&254_Shimadzu.lcm, MS (ESI) m/z =623.1

98


[M+H]⁺. ¹H NMR (400 MHz, DMSO-d₆): δ=9.66 (s, 1H), 8.67 (s, 1H), 8.07-7.93 (m, 2H), 7.44-7.39 (m, 2H), 6.92 (s, 1H), 6.88-6.77 (m, 3H), 6.53-6.49 (m, 1H), 6.26-6.21 (m, 1H), 5.71 (dd, J=10.0 and 2.0 Hz, 1H), 5.00 (t, J=5.6 Hz, 1H), 3.94 (s, 6H), 3.89-3.76 (m, 4H), 2.26-2.21 (m, 1H), 2.04-1.98 (m, 1H).

Example 7

Synthesis of Compound 11



N-[2-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]-5-(oxetan-3-yloxy)phenyl]prop-2-enamide

Scheme 10. Synthesis of compound 11.




99

100





Compound 11 was synthesized based on synthetic scheme 30 10. The synthetic process for each of the step is described as below.



Step 1 to synthesize compound 11: To a solution of 4-amino-3-nitro-phenol (200 mg, 1.30 mmol) in DMF (5 mL) was added Cs₂CO₃ (846 mg, 2.60 mmol). The mixture was stirred at 25° C. for 0.5 hour, then oxetan-3-yl 4-methylbenzenesulfonate (444 mg, 1.95 mmol) was added. The mixture was heated to 80° C. for 6 hours. The mixture was diluted with water (10 mL), extracted with EtOAc (20 mL×3), then the combined organic layers were dried over Na₂SO₄, filtered and concentrated. The residue was purified by silica gel chromatography (PE/EtOAc=1/0 to 3/1) to afford 2-nitro-4-(oxetan-3-yloxy)aniline (100 mg, 0.476 mmol, 37% yield) as red solid. ¹H NMR (400 MHz, DMSO-d₆): δ=7.30 (s, 2H), 7.16-7.09 (m, 2H), 7.20 (d, J=9.2 Hz, 1H), 5.27-5.22 (m, 1H), 4.91-4.87 (m, 2H), 4.54-4.51 (m, 2H).

101


102




11-4

Step 2 to synthesize compound 11: To a solution of N-(2,6-dichloro-3,5-dimethoxy-phenyl)-3-(6-methylsulfonylpyrimidin-4-yl)pyridin-2-amine (300 mg, 0.659 mmol), 2-nitro-4-(oxetan-3-yloxy)aniline (138 mg, 0.659 mmol) in DMAc (10 mL) was added t-BuOK (223 mg, 1.99 mmol) at 0° C. The mixture was then heated to 40° C. for 2 hours. The mixture was diluted with water (10 mL), extracted with EtOAc (20 mL×3). Then the combined organic layers were washed with sat. aq. NaCl (10 mL×7), dried over Na₂SO₄, filtered and concentrated. The residue was triturated with CH₃CN (5 mL) at 25° C. for 5 min. The precipitation was filtered, dried to afford 6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]-N-[2-nitro-4-(oxetan-3-yloxy)phenyl]pyrimidin-4-amine (300 mg, 0.512 mmol, 78% yield) as a yellow solid.

-continued



11-5



11-4

55


60

65


Step 3 to synthesize compound 11: To a solution of 6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]-N-[2-nitro-4-(oxetan-3-yloxy)phenyl]pyrimidin-4-amine (200 mg, 0.34 mmol) in THE (10 mL) and H₂O (10 mL) was added Fe (95 mg, 1.71 mmol) and NH₄Cl (91 mg, 1.71 mmol). The mixture was stirred at 65° C. for 4 hours. The mixture was cooled to room temperature, filtered, and the filtrate was concentrated to afford N₁-[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]-4-(oxetan-3-yloxy)benzene-1,2-diamine (180 mg, crude) as yellow solid.

103

104



11-5



compound 11


Step 4 to synthesize compound 11: To a solution of N₁-[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]-4-(oxetan-3-yloxy)benzene-1,2-diamine (150 mg, 0.27 mmol) in DCM (15 mL) was added prop-2-enoyl chloride (19 mg, 0.216 mmol) at 0° C. The mixture was stirred at 25° C. for 0.5 hour. The mixture was quenched by water (10 mL), then extracted with DCM (50 mL×3), dried over Na₂SO₄, filtered and concentrated. The residue was purified preparative HPLC (Colum: Xtimate C18 100*30 mm*3 um, water (0.04% NH₃·H₂O+10 mM NH₄HCO₃)-ACN; B % from 63 to 83; Gradient Time: 30 min; Flow rate: 25 mL/min) then further purified by chiral SFC (Instrument: CAS-QD-ANA-SFC-SD (Agilent 1260 with DAD detector), Column: Chiralcel OJ-3 100×4.6 mm I.D., 3 um. Mobile phase: A: CO₂ B: ethanol (0.05% DEA), Gradient: from 5% to 40% of B in 4.5 min and hold 40% for 2.5 min, then 5% of B for 1 min Flow rate: 2.8 mL/min, Column temperature: 40° C.) to afford N-[2-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]-5-(oxetan-3-yloxy)phenyl]prop-2-enamide (30 mg, 0.049 mmol, 18% yield, 100% purity) as yellow solid. LCMS: t_R=2.023 min in:10-80CD₄ min_Pos_220&254_Shimadzu.lcm, MS (ESI) m/z=609.2 [M+H]⁺. ¹H NMR (400 MHz, DMSO-d₆): δ=10.93 (s, 1H), 9.66 (s, 1H), 8.98 (s, 1H), 8.61 (s, 1H), 8.07-7.96 (m, 2H), 7.42-7.34 (m, 2H), 6.92 (s, 1H), 6.86-6.84 (m, 2H), 6.66-6.55 (m, 2H), 6.26-6.21 (m, 1H), 5.74-5.71 (m, 1H), 5.29-5.26 (m, 1H), 4.94-4.91 (m, 2H), 4.59-4.56 (m, 2H), 3.37 (s, 6H).

Example 8

Synthesis of Compound 12




N-[2-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]-5-(2-pyrrolidin-1-ylethoxy)phenyl]prop-2-enamide

Scheme 11. Synthesis of compound 12.




105


-continued

**106**


-continued




Compound 12 was synthesized based on synthetic scheme 11. The synthetic process for each of the step is described as below.




Step 1 to synthesize compound 12: 4-amino-3-nitrophenol (1.5 g, 9.73 mmol), Boc₂O (3.19 g, 14.60 mmol) and Et₃N (984 mg, 9.73 mmol) dissolved in DCM (10 mL). Then a solution of DMAP (118 mg, 0.973 mmol) in DCM (10 mL) was slowly added dropwise to the above solution. Then the mixture was stirred at 25° C. for 2 hours. The mixture was concentrated under reduced pressure to remove DCM. The residue was purified by silica gel chromatography. (ISCO®; 20 g SepaFlash® Silica Flash Column, Eluent of 0-30% Ethyl acetate/Petroleum ether gradient @ 40 mL/min) to afford tert-butyl N-(4-hydroxy-2-nitrophenyl)carbamate (2.35 g, 95% yield) as a yellow solid. LCMS: t_R=1.169 min in 10-80AB_2 min_220&254_Shimadzu.lcm, MS (ESI) m/z=255.1 [M+H]⁺.




107



Step 2 to synthesize compound 12: A solution of 1-(2-chloroethyl)pyrrolidine (2.25 g, 13.22 mmol, HCl salt), tert-butyl N-(4-hydroxy-2-nitro-phenyl)carbamate (2.8 g, 11.01 mmol), Cs₂CO₃ (7.18 g, 22.03 mmol), in DMF (100 mL) was stirred at 50° C. for 8 hours. The mixture was extracted with EtOAc (200 mL×5). The combined organic layers were washed with H₂O (50 mL×6), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by flash silica gel chromatography (ISCO®; 20 g SepaFlash® Silica Flash Column, Eluent of 0-35% Ethyl acetate/Petroleum ether gradient @ 40 mL/min) to afford tert-butyl N-[2-nitro-4-(2-pyrrolidin-1-ylethoxy)phenyl]carbamate (1.2 g, 31% yield) as a black solid. LCMS: t_R=0.980 min in 10-80 AB₂ min_{220&254}_Shimadzu.lcm, MS (ESI) m/z=352.1 [M+H]⁺.




HCl/dioxane





Step 3 to synthesize compound 12: A mixture of tert-butyl N-[2-nitro-4-(2-pyrrolidin-1-ylethoxy)phenyl]carbamate (1.2 g, 3.41 mmol), HCl/dioxane (4 M, 0.85 mL) in MeOH (20 mL) was stirred at 25° C. for 1 hr. The mixture was adjusted to pH=8 by addition of sat. aq. NaHCO₃ and the resulting mixture was concentrated to afford 2-nitro-4-(2-pyrrolidin-1-ylethoxy)aniline (1.0 g, crude) as a yellow solid. LCMS: t_R=0.882 min in 0-60 AB₂ min_{220&254}_Shimadzu.lcm, MS (ESI) m/z=252.1 [M+H]⁺.

108





Step 4 to synthesize compound 12: To a solution of N-(2,6-dichloro-3,5-dimethoxy-phenyl)-3-(6-methylsulfonylpyrimidin-4-yl)pyridin-2-amine (352 mg, 0.659 mmol), 2-nitro-4-(2-pyrrolidin-1-ylethoxy)aniline (500 mg, 1.99 mmol) in DMF (30 mL) was added t-BuOK (223 mg, 1.99 mmol) at 0° C. Then the mixture was stirred at 40° C. for 2 hours under N₂ atmosphere. The mixture was extracted with EtOAc (150 mL×5). The combined organic layers were washed with H₂O (30 mL×6), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by flash silica gel chromatography (ISCO®; 12 g SepaFlash® Silica Flash Column, Eluent of 0-15% MeOH/DCM gradient @ 20 mL/min) to afford 6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]-N-[2-nitro-4-(2-pyrrolidin-1-ylethoxy)phenyl]pyrimidin-4-amine (539 mg, 69% yield, 53% purity) as a yellow solid. LCMS: t_R=1.004 min in 10-80 AB₂ min_{220&254}_Shimadzu.lcm, MS (ESI) m/z=626.1 [M+H]⁺.

109



Step 5 to synthesize compound 12: A mixture of 6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-amine (400 mg, 0.638 mmol) and Pd/C (500 mg, 10% purity) in MeOH (30 mL) was stirred at 25° C. for 2 hours under H₂ atmosphere. The mixture was filtered and concentrated under reduced pressure to give a residue. The mixture was concentrated to afford N-[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]-4-(2-pyrrolidin-1-ylethoxy)benzene-1,2-diamine (200 mg, crude) as a black solid. LCMS: *t_R*=1.218 min in 10-80 AB_4 min_220&254_Shimadzu.lcm, MS (ESI) *m/z*=596.3 [M+H]⁺.

110



Step 6 to synthesize compound 12: To a solution of N-[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]-4-(2-pyrrolidin-1-ylethoxy)benzene-1,2-diamine (150 mg, 0.251 mmol) in DCM (4 mL) and Et₃N (28 mg, 0.277 mmol) was added a solution of prop-2-enoyl chloride (25 mg, 0.277 mmol) in DCM (3 mL) dropwise at 0° C. The mixture was stirred at 25° C. for 0.5 hour. The mixture was concentrated under reduced pressure to remove DCM. The residue was purified by preparative HPLC (Column: Xtimate C18 100*30 mm*3 um water (0.04% NH₃·H₂O+10 mM NH₄HCO₃)-ACN; B % from 63 to 83; Gradient Time: 30 min; Flow rate: 25 mL/min) to afford N-[2-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]-5-(2-pyrrolidin-1-ylethoxy)phenyl]prop-2-enamide (9.1 mg, 5% yield, 98% purity) as a white solid. LCMS: *t_R*=1.550 min in 10-80CD_4 min_220&254_Shimadzu.lcm, MS (ESI) *m/z*=650.2 [M+H]⁺. ¹H NMR (400 MHz, DMSO-*d*₆): δ=10.95 (s, 1H), 9.64 (s, 1H), 8.94 (s, 1H), 8.61 (s, 1H), 8.05-7.95 (m, 2H), 7.45-7.38 (m, 2H), 6.94-6.81 (m, 4H), 6.55-6.48 (m, 1H), 6.25-6.21 (m, 1H), 5.72 (d, J=11.2 Hz, 1H), 4.07 (t, J=5.6 Hz, 2H), 3.92 (s, 6H), 2.89-2.73 (m, 6H), 1.69 (s, 4H).


111

Example 9


Synthesis of Compound 13

N-[5-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]-1-methyl-imidazol-4-yl]prop-2-enamide


Scheme 12. Synthesis of compound 13.

**112**


-continued




Compound 13 was synthesized based on synthetic scheme 12. The synthetic process for each of the step is described as below.





Step 1 to synthesize compound 13: A mixture of 5-chloro-1-methyl-4-nitroimidazole (2 g, 12.38 mmol) in NH_3/MeOH (10 mL, 7 M) was stirred at 100°C . for 16 hr in an autoclave. The mixture was cooled down to room temperature, and the precipitation was filtered, washed with MeOH (10 mL $\times$ 2), dried to afford 3-methyl-5-nitroimidazol-4-amine (1.6 g, 11.26 mmol, 91% yield) as a yellow solid. ^1H NMR (400 MHz, DMSO-d_6): $\delta=7.55$ (s, 2H), 7.25 (s, 1H), 3.47 (s, 3H).



113




Step 2 to synthesize compound 13: To a solution of N-(2,6-dichloro-3,5-dimethoxy-phenyl)-3-(6-methylsulfonylpyrimidin-4-yl)pyridin-2-amine (500 mg, 1.10 mmol) and 3-methyl-5-nitro-imidazol-4-amine (312 mg, 2.20 mmol) in DMF (20 mL) was added tBuOK (246 mg, 2.20 mmol). The mixture was stirred at 40° C. for 12 hr under N₂ atmosphere. The mixture was diluted with H₂O (20 mL) and extracted with EtOAc (10 mL×2). The combined organic layers were washed with H₂O (10 mL×3), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by flash silica gel chromatography (ISCO®; 12 g SepaFlash® Silica Flash Column, Eluent of 0~10% Ethyl acetate/Petroleum ether gradient @ 50 mL/min) to afford 6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]-N-(3-methyl-5-nitro-imidazol-4-yl)pyrimidin-4-amine (320 mg, 0.619 mmol, 56% yield) as a yellow solid.



Step 3 to synthesize compound 13: To a solution of 6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]-N-(3-

114

methyl-5-nitro-imidazol-4-yl)pyrimidin-4-amine (300 mg, 0.58 mmol) in MeOH (5 mL) was added Pd/C (5 mg, 10% purity) under N₂ atmosphere. The suspension was degassed and purged with H₂ for 3 times. The mixture was stirred under H₂ (15 psi) at 25° C. for 2 hr. The liquid was filtered with diatomite, and the filtrate was concentrated under reduced pressure to afford N₅-[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]-1-methyl-imidazole-4,5-diamine (200 mg, 0.41 mmol, 71% yield) as a brown solid.




Step 4 to synthesize compound 13: To a solution of N₅-[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]-1-methyl-imidazole-4,5-diamine (100 mg, 0.205 mmol) in DCM (10 mL) and DIEA (53 mg, 0.41 mmol) was added prop-2-enoyl chloride (15 mg, 0.164 mmol) at 0° C. The mixture was stirred at 0° C. for 0.5 hr. The mixture was concentrated under reduced pressure and the residue was purified by preparative HPLC (Column: Phenomenex Luna C18 100*40 mm*3 um; water (0.225% FA)-ACN; B % from 20 to 40; Gradient time: 9 min; Flow rate: 25 mL/min) to afford N₅-[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]-1-methyl-imidazol-4-yl]prop-2-enamide (13.6 mg, 0.024 mmol, 12% yield, 95.34% purity) as a yellow solid. LCMS: t_R=0.968 min in 10-80AB_4 min_220&254_Shimadzu.lcm, MS (ESI) m/z=541.2 [M+H]⁺. ¹H NMR (400 MHz, DMSO-d₆): δ=10.99 (s, 1H), 10.45 (s, 1H), 9.26 (s, 1H), 8.72 (s, 1H), 8.13-8.09 (m, 2H), 7.21-6.86 (m, 3H), 6.43-6.23 (m, 2H), 5.79-5.76 (m, 1H), 4.04 (s, 6H), 3.55 (s, 3H).

115

Example 10


Synthesis of Compound 14

Scheme 13. Synthesis of compound 14.




116


-continued



Compound 14 was synthesized based on synthetic scheme 13. The synthetic process for each of the step is described as below.





Step 1 to synthesize compound 14: To the mixture of compound 14-2 (30.0 g, 194.8 mmol, 1.0 eq) in ethyl acetate (600 mL) was added compound 14-1 (19 g, 214 mmol, 1.1 eq) and triphenylphosphine (56.0 g, 214.1 mmol, 1.1 eq). Then diethyl azodicarboxylate (33.5 mL, 214.1 mmol, 1.1 eq) was added dropwise. The mixture was stirred at room temperature overnight. The reaction was monitored by TLC. Then the mixture was filtered through a pad of Celite and sintered funnel. The result mixture was extracted with ethyl acetate (3×600 mL) and added 1N HCl and water. The aqueous phase was extracted with ethyl acetate and ammonium hydroxide. The combined organic phase was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by column chromatography on a silica gel (dichloromethane:methanol, 100:1-50:1-30:1) to give compound 14-3 (4.6 g, 11%).




117

-continued




Step 2 to synthesize compound 14: To the mixture of compound 14-3 (10.0 g, 44.4 mmol, 1.1 eq) and compound 14-4 (18.4 g, 40.4 mmol, 1.0 eq) in dimethylformamide (60 mL) was added potassium tert-butoxide (13.6 g, 121.2 mmol, 3.0 eq) under nitrogen atmosphere. Then the mixture was stirred at 40° C. for 2 h. TLC analysis of the reaction mixture showed partial conversion to the desired product. The mixture was cooled to room temperature and extracted with ethyl acetate and water. The combined organics were washed with brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by column chromatography on a silica gel (dichloromethane:methanol, 100:1-50:1) to afford compound 14-5 (5 g, 21%).





118

-continued



Step 3 to synthesize compound 14: To the mixture of compound 14-5 (5.0 g, 8.3 mmol, 1.0 eq) in ethanol/water (100 mL/20 mL) was added Fe (2.33 g, 41.7 mmol, 5.0 eq) and ammonium chloride (4.46 g, 83.4 mmol, 10.0 eq). Then the mixture was stirred at 85° C. for 2 h. TLC analysis of the reaction mixture showed partial conversion to the desired product. The mixture was filtered and extracted with ethyl acetate (120 mL×3). The combined organics were washed with brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by column chromatography on a silica gel (dichloromethane:methanol, 10:1) to afford compound 14-6 (4 g, 85%).



Step 4 to synthesize compound 14: To the mixture of compound 14-6 (4.0 g, 7.03 mmol, 1.0 eq) in dichloromethane (40 mL) was added compound 14-7 (632.6 mg, 7.03 mmol, 1.0 eq) at 0° C. The mixture was stirred at room temperature for 2 h. TLC analysis of the reaction mixture showed full conversion to the desired product. Then the mixture was quenched with water (200 mL) and extracted


119

with dichloromethane (40 mL×2). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by column chromatography on a silica gel (dichloromethane: methanol, 10:1) to afford the crude product (purity: 90%). The crude product was dissolved in dichloromethane (50 mL). Then the mixture was added 20% sodium hydroxide solution (10 mL). The mixture was stirred at room temperature for 30 min. Then the mixture was extracted with dichloromethane and washed with brine. The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give compound 14 (2.0 g, 46%) as yellow solid. LCMS: $[M+1]^+$ 624. ^1H NMR (400 MHz, DMSO): δ 10.94 (s, 1H), 8.58 (s, 1H), 8.02-8.01 (m, 1H), 7.92 (m, 1H), 7.43 (m, 1H), 7.39-7.37 (m, 1H), 6.96 (m, 1H), 6.80-6.78 (m, 3H), 6.76 (m, 1H), 6.46 (m, 1H), 6.17 (m, 1H), 5.73 (s, 1H), 5.69-5.66 (m, 1H), 4.03-4.00 (m, 3H), 3.90 (s, 8H), 2.61-2.59 (m, 3H) and 2.19 (s, 8H).

Example 11


Synthesis of Compound 15

Scheme 14. Synthesis of compound 15.




120

-continued




Compound 15 was synthesized based on synthetic scheme 14. The synthetic process for each of the step is described as below.




Step 1 to synthesize compound 15: To the mixture of compound 15-2 (30.0 g, 194.6 mmol, 1.0 eq) in ethyl acetate (600 mL) was added compound 15-1 (25.1 g, 214 mmol, 1.1 eq) and triphenylphosphine (56.0 g, 214.1 mmol, 1.1 eq). Then diethyl azodicarboxylate (33.5 mL, 214.1 mmol, 1.1 eq) was added dropwise. The mixture was stirred at room temperature overnight. The reaction was monitored by TLC. Then the mixture was filtered through a pad of celite and sintered funnel. The result mixture was extracted with ethyl acetate (3×600 mL) and added 1N HCl and water. The aqueous phase was extracted with ethyl acetate and ammonium hydroxide. The combined organic phase was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by column

121

chromatography on a silica gel (dichloromethane:methanol, 100:1~50:1~30:1) to give compound 15-3 (17.5 g, 36%).




Step 2 to synthesize compound 15: To the mixture of compound 15-3 (2.2 g, 8.8 mmol, 2.0 eq) and compound 15-4 (2.0 g, 4.4 mmol, 1.0 eq) in dimethylformamide (30 mL) was added potassium tert-butoxide (1.48 g, 13.2 mmol, 3.0 eq) under nitrogen atmosphere. Then the mixture was stirred at 40°C for 2 h. TLC analysis of the reaction mixture showed partial conversion to the desired product. The mixture was cooled to room temperature and extracted with ethyl acetate and water. The combined organics were washed with brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by column chromatography on a silica gel (dichloromethane:methanol, 100:1-50:1) to afford compound 15-5 (550 mg, 20%).




122

-continued



Step 3 to synthesize compound 15: To the mixture of compound 15-5 (1.0 g, 1.6 mmol, 1.0 eq) in ethanol/water (20 mL/4 mL) was added Fe (446 mg, 8.0 mmol, 5.0 eq) and ammonium chloride (880 mg, 16 mmol, 10.0 eq). Then the mixture was stirred at 80°C for 2 h. TLC analysis of the reaction mixture showed partial conversion to the desired product. The mixture was filtered and extracted with ethyl acetate (20 mL $\times$ 3). The combined organics were washed with brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by column chromatography on a silica gel (dichloromethane:methanol, 10:1) to afford compound 15-6 (280 mg, 29%).



Step 4 to synthesize compound 15: To the mixture of compound 15-6 (280 mg, 0.47 mmol, 1.0 eq) in dichloromethane (5 mL) was added compound 15-7 (42.5 mg, 0.47 mmol, 1.0 eq) at 0°C . The mixture was stirred at room temperature for 2 h. TLC analysis of the reaction mixture showed full conversion to the desired product. Then the


123

mixture was quenched with water (20 mL) and extracted with dichloromethane (5 mL $\times$ 2). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by column chromatography on a silica gel (dichloromethane: methanol, 10:1) to afford compound 15 (81 mg, 29%) as yellow solid. LCMS: $[M+1]^+$ 652. ^1H NMR (400 MHz, CDCl_3): δ 11.12 (s, 1H), 9.30 (s, 1H), 8.62 (s, 1H), 8.40 (m, 1H), 8.12 (m, 1H), 7.85 (m, 1H), 7.38 (m, 1H), 7.25 (m, 1H), 7.00 (s, 1H), 6.66 (m, 1H), 6.45 (m, 2H), 6.38-6.35 (m, 4H), 5.71-5.69 (m, 1H), 3.34 (s, 2H), 3.89 (s, 8H), 3.46 (s, 1H), 3.34 (m, 2H), 3.17 (m, 5H) and 1.38-1.35 (m, 8H).

Example 12


Synthesis of Compound 16

Scheme 15. Synthesis of compound 16.




124

-continued




125

-continued




compound 16

126




Step 2 to synthesize compound 16: To a solution of trans-2-aminocyclopentanol (22 g, 217.51 mmol) in THE (220 mL) was added TEA (26.41 g, 261.01 mmol) and Boc_2O (52.22 g, 239.26 mmol, 54.97 mL, 1.1 eq). The mixture was stirred at 15° C. for 16 hr. The mixture was concentrated under reduced pressure to remove THF. The residue was diluted with EtOAc (20 mL) and extracted with H_2O (20 mL $\times$ 3), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by flash silica gel chromatography (ISCO®; 80 g SepaFlash® Silica Flash Column, Eluent of 0–30% Ethyl acetate/Petroleum ether gradient @ 60 mL/min) to afford tert-butyl N-[trans-2-hydroxycyclopentyl]carbamate (14 g, 69.56 mmol, 32% yield) as a white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ =6.72 (d, J =6.4 Hz, 1H), 4.60 (s, 1H), 3.80–3.75 (m, 1H), 3.49 (br, 1H), 1.86–1.54 (m, 4H), 1.38 (s, 9H), 1.32–1.29 (m, 2H).



16-4


Compound 16 was synthesized based on synthetic scheme 15. The synthetic process for each of the step is described as below.




Step 1 to synthesize compound 16: To a solution of 6-oxabicyclo[3.1.0]hexane (7 g, 83.22 mmol) in i-PrOH (40 mL) was added $\text{NH}_3\cdot\text{H}_2\text{O}$ (35.00 g, 249.65 mmol 25% purity). The mixture was stirred at 60° C. for 16 hr in a sealed tube. The mixture was concentrated under reduced pressure to afford trans-2-aminocyclopentanol (26 g, crude) as a brown oil. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ =3.68–3.53 (m, 1H), 2.88–2.77 (m, 1H), 1.82–1.77 (m, 2H), 1.58–1.53 (m, 2H), 1.44–1.40 (m, 1H), 1.22–1.16 (m, 1H).

Step 3 to synthesize compound 16: A mixture of tert-butyl N-[trans-2-hydroxycyclopentyl]carbamate (14 g, 69.56 mmol), isoindoline-1,3-dione (11.26 g, 76.52 mmol), PPh_3 (20.07 g, 76.52 mmol) and DIAD (16.18 g, 80.00 mmol) in THE (150 mL) was degassed and purged with N_2 for 3 times, and then the mixture was stirred at 0° C. for 1 hr. The mixture was concentrated under reduced pressure to remove THF. The residue was diluted with H_2O (150 mL), and extracted with EtOAc (150 mL $\times$ 3), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by flash silica gel chromatography (ISCO®; 80 g SepaFlash® Silica Flash Column, Eluent of 0–30% Ethyl acetate/Petroleum ether gradient @ 80 mL/min). The crude product was further re-crystallization from EtOH (100 mL) at 80° C., then the precipitation was filtered, dried to afford tert-butyl N-[cis-2-(1,3-dioxoisindolin-2-yl)cyclopentyl]carbamate (15 g, 45.40 mmol, 65% yield) as a white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ =7.83–7.8 (m, 5H), 6.67 (d, J =7.6 Hz, 1H), 4.49–4.45 (m, 1H), 1.92–1.41 (m, 6H), 1.01 (s, 9H).

127




Step 4 to synthesize compound 16: To a solution of tert-butyl N-[cis-2-(1,3-dioxoisindolin-2-yl)cyclopentyl] carbamate (5 g, 15.13 mmol) in MeOH (50 mL) was added NH_2NH_2 (709 mg, 16.65 mmol, 80% purity). The mixture was stirred at 50° C. for 1.5 hr. The mixture was concentrated under reduced pressure to remove MeOH. The residue was diluted with H_2O (50 mL), then extracted with EtOAc (50 mL $\times$ 3), dried over Na_2SO_4 , filtered and concentrated to afford tert-butyl N-[cis-2-aminocyclopentyl]carbamate (1.1 g, 5.49 mmol, 36% yield) as a yellow oil.




128

-continued




Step 5 to synthesize compound 16: To a solution of N-(2,6-dichloro-3,5-dimethoxy-phenyl)-3-[(6-methylsulfonyl)pyrimidin-4-yl]pyridin-2-amine (758 mg, 1.66 mmol) and tert-butyl N-[cis-2-aminocyclopentyl]carbamate (1 g, 4.99 mmol) in DMAc (10 mL) was added DIEA (430 mg, 3.33 mmol). The mixture was stirred at 130° C. for 12 hr. The mixture was diluted with H_2O (20 mL) and extracted with EtOAc (20 mL $\times$ 3), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by flash silica gel chromatography (ISCO®; 12 g SepaFlash® Silica Flash Column, Eluent of 0-30% Ethyl acetate/Petroleum ether gradient @ 40 mL/min) to afford tert-butyl N-[2-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]cyclopentyl]carbamate (cis mixture, 500 mg, 0.869 mmol, 52% yield) as a yellow solid. LCMS: t_R =1.412 min in 10-80AB_2 min_220&254_Shimadzu.lcm, MS (ESI) m/z =575.2 $[\text{M}+\text{H}]^+$.




129

-continued



**130**

-continued



Chiral SFC

Step 6 to synthesize compound 16: A solution of tert-butyl N-[2-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]cyclopentyl]carbamate (500 mg, 0.868 mmol, cis mixture) in HCl/dioxane (4 M, 10 mL) was stirred at 25° C. for 0.5 hr. The mixture was concentrated to afford N₁-[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]cyclopentane-1,2-diamine (800 mg, crude) as a white solid.



Step 7 to synthesize compound 16: To a solution of N₁-[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]cyclopentane-1,2-diamine (700 mg, 1.47 mmol, cis mixture) in DCM (10 mL) was added DIEA (190 mg, 1.47 mmol) and prop-2-enoyl chloride (67 mg, 0.736 mmol). The mixture was stirred at 0° C. for 0.5 hr. The residue was diluted with H₂O (20 mL) and extracted with EtOAc (20 mL×3), dried over Na₂SO₄, filtered and concn-

131

trated under reduced pressure to give a residue. The residue was purified by flash silica gel chromatography (ISCO®; 12 g SepaFlash® Silica Flash Column, Eluent of 0~10% dichloromethane/Petroleum ether gradient @ 40 mL/min) to give desired product (170 mg, 94% purity) as a white solid, which was further separated by SFC (column: DAICEL CHIRALPAK AD (250 mm*30 mm, 10 μ m); mobile phase: [0.1% NH₃·H₂O ETOH]; B %: 35%-35%, 30 min) to afford (+) N-[cis-2-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]cyclopentyl]prop-2-enamide (43.7 mg, 0.081 mmol, 98.85% purity, single cis isomer) and (–) N-[cis-2-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]cyclopentyl]prop-2-enamide (46.5 mg, 0.087 mmol, 99.10% purity, single cis isomer) as a white solid.


(+)N-[cis-2-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]cyclopentyl]prop-2-enamide (the first peak in chiral SFC). LCMS: t_R =1.476 min in 10-80AB_4 min_220&254_Shimadzu.lcm, MS (ESI) m/z =529.2 [M+H]⁺. ¹H NMR (400 MHz, DMSO-d₆): δ =11.09 (s, 1H), 8.48 (s, 1H), 8.04-8.02 (m, 3H), 7.27 (d, J=8.0 Hz, 1H), 6.97 (s, 1H), 6.87-6.83 (m, 2H), 6.17 (s, 1H), 6.05-5.95 (m, 1H), 5.48 (s, 1H), 4.41-4.25 (m, 2H), 3.93 (s, 6H), 2.02-1.57 (m, 6H). Chiral SFC: t_R =5.118 min (Instrument column: Chiralpak AD-3 100×4.6 mm I.D., 3 μ m, Mobile phase: A: CO₂ B:ethanol (0.05% DEA), Gradient: from 5% to 40% of B in 4.5 min and hold 40% for 2.5 min, then 5% of B for 1 min; Flow rate: 2.8 mL/min Column temperature: 40° C.; UV detection: 220 nm), ee %=99%. [α]_D²⁰=+13 (c=0.10, MeOH).

(–)N-[cis-2-[[6-[2-(2,6-dichloro-3,5-dimethoxy-anilino)-3-pyridyl]pyrimidin-4-yl]amino]cyclopentyl]prop-2-enamide (the second peak in chiral SFC). LCMS: t_R =1.481 min in 10-80AB_4 min_220&254_Shimadzu.lcm, MS (ESI) m/z =529.2 [M+H]⁺. ¹H NMR (400 MHz, DMSO-d₆): δ =11.09 (s, 1H), 8.48 (s, 1H), 8.04-8.02 (m, 3H), 7.27 (d, J=8.0 Hz, 1H), 6.97 (s, 1H), 6.87-6.83 (m, 2H), 6.16 (s, 1H), 6.05-5.95 (m, 1H), 5.48 (s, 1H), 4.41-4.25 (m, 2H), 3.93 (s, 6H), 2.02-1.57 (m, 6H). Chiral SFC: t_R =6.002 min (Instrument column: Chiralpak AD-3 100×4.6 mm I.D., 3 μ m, Mobile phase: A: CO₂ B:ethanol (0.05% DEA), Gradient: from 5% to 40% of B in 4.5 min and hold 40% for 2.5 min, then 5% of B for 1 min; Flow rate: 2.8 mL/min Column temperature: 40° C.; UV detection: 220 nm), ee %=99.1%. [α]_D²⁰=–13 (c=0.10, MeOH).

Example 13

Evaluation of the In Vitro Efficacy of FGFR Inhibitors

inventive compound 1



132

-continued

control compound 5




Table 2 below compares the properties of inventive compound 1 and control compound 5, which indicates that compound 1 exhibits superior properties in solubility, permeability, mouse plasma stability, mouse liver microsome stability, in vivo exposure, and bioavailability than control compound 5. The inventive compound 1 improves the FGFR4-selective inhibitory effect. More importantly, the plasma stability of the inventive compound is significantly improved. Also, control compound 5 has poor solubility in the solution of 5% DMSO and 95% of (20% HPBC in PBS). The inventive compound 1, on the other hand, exhibits a much-improved solubility which facilitates its administration.

TABLE 2

Comparison of the properties between inventive compound 1 and control compound 5.

Compound	Compound 1	Compound 5
Molecular weight	530.12	550.13
s + LogP	3.13	5.08
s + LogD 7.4	3.12	5.08
MDCK Permeability Papp, A-B (1X10 ⁻⁶ cm/s)	25.8	11.4
Protein binding (%)	92	—
T_PSA Å ²	119.52	110.29
IC ₅₀ towards FGFR1 (nM)	1280	>30000
IC ₅₀ towards FGFR2 (nM)	1870	>30000
IC ₅₀ towards FGFR3 (nM)	4440	>30000
IC ₅₀ towards FGFR4 (nM)	20	4.1
Hepatocyte Clin Ms (mL/min/10 ⁶ cells) (SE)	43.63 ± 1.30	41.70 ± 1.73
Ms Plasma Stability t _{1/2} (min)(SE)	1147.5 ± 347.0	221.2 ± 55.7
CL (L/h/kg)/V _{ss} (L/kg)	1.6/0.88/0.52/20.9	0.84/1.53/115(?)/18.0
t _{1/2} (h)/% F		

Referring to FIGS. 2 and 3, the inventive compound 5 also exhibits an non-obvious improvement of pharmacokinetic data when administered into mice. Table 3 below summarizes the critical pharmacokinetic data from these experiments.

133

TABLE 3

Pharmacokinetic data of inventive compound 1 and control compound 5.						
Route	Compound					
	Compound 1			Compound 5		
	IP	IV	PO	IP	IV	PO
Dose (mg/kg)	10	1	5	10	5	100
FIGS.	FIG. 2A	FIG. 2B	FIG. 2B	FIG. 3A	FIG. 3B	FIG. 3B
T _{max} (hr)	0.25	0.083	3.17	0.25	0.083	6.67
C _{max} (ng/mL)	14867	849	119	865	470	180
T _{1/2} (hr)	2.81	0.522	2.73	1.23	15.1	—
MRT _{last} (hr)	0.67	0.515	3.42	1.57	7.71	6.1
MRT _{inf} (hr)	0.67	0.537	4.02	1.66	18.5	—

134

TABLE 3-continued

Pharmacokinetic data of inventive compound 1 and control compound 5.						
Route	Compound					
	Compound 1			Compound 5		
	IP	IV	PO	IP	IV	PO
10 AUC _{last} (hr*ng/mL)	9608	618	465	1270	896	1450
AUC _{inf} (hr*ng/mL)	9609	621	648	1280	1240	—
15 Cl (mL/hr/kg)	—	1632	—	—	837	8.09
V _{ss} (mL/kg)	—	876	—	—	15300	—
F (%)	—	—	20.9	—	—	8.09

TABLE 4



FGFR inhibition data of the disclosed compounds.					
Compound No.	Molecular Structure	FGFR1 Inhibition	FGFR2 Inhibition	FGFR3 Inhibition	FGFR4 inhibition
Compound 1		—	—	—	++
Compound 2		++	+	+	+++

TABLE 4-continued




FGFR inhibition data of the disclosed compounds.					
Compound No.	Molecular Structure	FGFR1 Inhibition	FGFR2 Inhibition	FGFR3 Inhibition	FGFR4 inhibition
Compound 3		ND	ND	ND	+
Compound 4		ND	ND	ND	++
Compound 5 (Control)		—	—	—	+++

TABLE 4-continued




FGFR inhibition data of the disclosed compounds.					
Compound No.	Molecular Structure	FGFR1 Inhibition	FGFR2 Inhibition	FGFR3 Inhibition	FGFR4 inhibition
Compound 6		+	+	—	+++
Compound 7		+	+	—	+++
Compound 8		—	+	—	+++

TABLE 4-continued




FGFR inhibition data of the disclosed compounds.					
Compound No.	Molecular Structure	FGFR1 Inhibition	FGFR2 Inhibition	FGFR3 Inhibition	FGFR4 inhibition
Compound 9		+	+	—	+++
Compound 10		—	—	—	+++
Compound 11		—	—	—	+++

TABLE 4-continued






FGFR inhibition data of the disclosed compounds.					
Compound No.	Molecular Structure	FGFR1 Inhibition	FGFR2 Inhibition	FGFR3 Inhibition	FGFR4 inhibition
Compound 12		+	+	—	+++
Compound 13		ND	ND	ND	—
Compound 14		—	—	—	+++

TABLE 4-continued

FGFR inhibition data of the disclosed compounds.					
Compound No.	Molecular Structure	FGFR1 Inhibition	FGFR2 Inhibition	FGFR3 Inhibition	FGFR4 inhibition
Compound 15		—	—	—	+++
Compound 16		ND	ND	ND	+

The evaluation standards of IC₅₀ values against the kinases in the table are shown below.

+++ below or equal to 10 nM

++ between 10 and 100 nM

+ between 100 and 1000 nM

— above or equal 1000 nM

ND Not determined

45

The FGFR inhibition data of the disclosed compounds are shown in Table 4. Compound 5 is used as a control.

TABLE 5

FGFR sequences.		
SEQ ID	FGFR1	AGATGCAGGGGCGCAAACGCCAAAGGAGACCAGGCTGTAGGAAGAGAAGGGCAGAGC
NO: 1	isoform 1	GCCGGACAGCTCGGCCGCTCCCCGTCTTTGGGGCCGCGGCTGGGGAACACAAGG
	Nucleic acid	CCCAGCAGGCAGCTGCAGGGGGCGGAGGCGGAGGAGGGACCAGCGCGGTGGGAGTG
	sequence	AGAGAGCGAGCCCTCGCGCCCCCGCGCGCATAGCGCTCGGAGCGCTCTTGGGCCA
		CAGGCGCGGCTCTCGCGGGCGGGCGGAGCTAGCGGGAGCCGGACGCCGTGCA
		GCCGCAGCGCGGAGGAACCCGGGTGTGCCGGAGCTGGGCGGCACGTCCGGACG
		GGACCGAGACCCCTCGTAGCGCATTCGCGCGACCTCGCCTTCCCCGGCGCGAGCGC
		GCCGCTGCTTGAAAAGCCGCGGAACCAAGGACTTTTCTCCGCTCCGAGCTCGGGG
		GCCCCGAGGGCGCACGGTACCCGTGCTGCAGTCGGGACGCGCGGCGCCGGGGCC
		TCCGACAGGCGATGGAGCCCGGTCTGCAAGGAAAGTGAGGCGCCGCGCTGCGTTCT
		GGAGGAGGGGGCACAAAGTCTGGAGACCCGGGTGGCGGACGGAGCCCTCCCCC
		GCCCCGCTCCGGGGCACCAGCTCCGGCTCCATTGTTCCCGCCCGGGCTGGAGGCGC
		CGAGCACCGAGCGCCGCGGGAGTCGAGCGCGGCGCGGAGCTCTTGCAGCCCGC
		CAGGACCCGAACAGAGCCCGGGGGCGGCGGGCCGGAGCCGGGACGCGGGCACACGC
		CCGTCGCACAAGCCACGGCGGACTCTCCGAGGCGGAACCTCCACGCCGAGCGAGG
		GTCACTTTGAAAAGGAGGATCGAGCTCACTGTGGAGTATCCATGGAGATGTGGAGCC
		TTGTCAACCACTCTAACTGCAGAACTGGGATGTGGAGCTGGAAGTGCCCTCTTC
		TGGGCTGTGCTGTCACAGCCACACTCTGCACCGCTAGGCCGTCCCGACCTTGCTT

TABLE 5-continued

FGFR sequences.

GAACAAGCCAGCCCTGGGAGCCCTGTGGAAGTGGAGTCCTTCTGGTCCACCCC
 GGTGACCTGCTGCAGCTTCGCTGTCGGCTGCGGGACGATGTGCAGAGCATCAACTGG
 CTGCGGGACGGGGTGACGTGGCGGAAGCAACCGCACCCGCATCAGGGGAGGAG
 GTGGAGGTGCAGGACTCCGTGCCCGCAGACTCCGGCCTCTATGCTTGCCTAACCCAGC
 AGCCCCTCGGGCAGTGACACCACCTACTTCTCCGTCAATGTTTCAGATGCTCTCCCC
 TCCTCGAGGATGATGATGATGATGATGACTCCTCTTCAGAGGAGAAGAAACAGAT
 AACACCAAAACCAACCGTATGCCCGTAGCTCCATATTGGACATCCCCAGAAAAGATG
 GAAAAGAAATTGCATGCAGTGCCGGCTGCCAAGACAGTGAAGTTCAAATGCCCTTCC
 AGTGGGACCCCAAACCCCACTGCGCTGGTTGAAAATGGCAAAGAAATCAAACCT
 GACCACAGAATTGGAGGCTACAAGGTCGGTTATGCCACCTGGAGCATCATAATGGAC
 TCTGTGGTGCCTCTGACAAGGGCAACTACACCTGCATTGTGGAGAAATGAGTACGGC
 AGCATCAACCACACATACCAGCTGGATGTCGTGGAGCGGTCCCTCACCGGCCCATC
 CTGCAAGCAGGGTGGCCCGCAACAAACAGTGGCCCTGGGTAGCAACGTGGAGTTC
 ATGTGTAAGGTGTACAGTGACCCGCAGCGCACATCCAGTGGCTAAGACATCGAG
 GTGAATGGGAGCAAGATTGGCCCAAGACAACCTGCCTTATGTCCAGATCTTTGAGACT
 GCTGGAGTTAATACCACCGACAAGAGATGGAGGTGCTTCACTTAAGAAATGTCTCC
 TTTGAGGACGCGGGGAGTATACGTGCTTGGCGGGTAACCTATCGGACTCTCCCAT
 CACTCTGCATGGTTGACCGTCTTGAAGCCCTGGAAGAGAGGGCCGGCAGTGATGACC
 TCGCCCCGTGACTGGAGATCATCATCTATTGCACAGGGGCCCTTCTCATCTCCTGC
 ATGGTGGGGTCCGTTCATCGTCTACAAGATGAAGAGTGGTACCAAGAGAGTGACTTC
 CACAGCCAGATGGCTGTGCACAAGCTGGCCAAAGAGCATCCCTCTGCGACAGCAGGTA
 ACAGTGTCTGCTGACTCCAGTGCATCCATGAACCTCTGGGGTCTTCTGTGTTCCGCCA
 TCACGGCTCTCTCCAGTGGGACTCCCATGCTAGCAGGGGTCTCTGAGTATGAGCTT
 CCCGAAGACCCCTCGCTGGGAGCTGCCTCGGGACAGACTGGTCTTAGGCAAAACCCCTG
 GGAGAGGGCTGCTTTGGGCAGGTGGTGTGGCAGAGGCTATCGGGCTGGACAAGGAC
 AAACCAACCGTGTGACCAAGTGGCTGTGAAGATGTTGAAGTCGGACGCAACAGAG
 AAAGACTTGTGAGACTGATCTCAGAAATGGAGATGATGAAGATGATCGGGAAGCAT
 AAGAATATCATCAACCTGCTGGGGGCTGCACGCAGGATGGTCCCTTGTATGTCATC
 GTGGAGTATGCCCTCAAGGGCAACCTGCGGGAGTACCTGCAGGCCCGGAGGCCCCCA
 GGGCTGGAATACTGCTACAACCCAGCCACAACCCAGAGGAGCAGCTCTCTCCAAG
 GACCTGGTGTCTTCCGCTACCAAGGTGGCCCGAGGCATGGAGTATCTGGCCTCCAAG
 AAGTGCATACACGAGACCTGGCAGCCAGGAATGTCTGGTGACAGAGGCAATGTG
 ATGAAGATAGCAGACTTGGCCTCGCACGGGACATTACCAACATCGACTACTATAAA
 AAGACAACCAACGGCCGAGTGCCTGTGAAGTGGATGGCACCAGGCGATTATTGAC
 CGGATCTACACCCACAGAGTGTGTGTGCTTTTCGGGGTGTCTCTGTGGGAGATC
 TTCACTCTGGGCGGCTCCCCATACCCCGGTGTGCTGTGGAGGAACCTTTCAAGCTG
 CTGAAGGAGGGTCACCGCATGGACAAGCCAGTAACCTGCACCAACGAGCTGTACATG
 ATGATCGGGGACTGCTGGCATGCAGTGCCCTCACAGAGACCCACCTTCAAGCAGCTG
 GTGGAAGACCTGGACCGCATCGTGGCCTTGACCTCCAAACAGGAGTACCTGGACCTG
 TCCATGCCCCCTGGACAGTACTCCCCAGCTTTCCCGACACCCGAGCTCTACGTGC
 TCCTCAGGGGAGGATTCCGTCTTCTCTCATGAGCCGCTGCCCGAGGAGCCCTGCCTG
 CCCCAGACCCAGCCAGCTTGCCCAATGGCGGACTCAAACGCCGCTGACTGCCACCC
 ACACGCCCTCCCCAGACTCCACCGTCAGCTGTAACCTCACCCACAGCCCTGCTGG
 GCCACCACTGTCCGTCCTGTCCCTTTCTGCTGGCAGGAGCCGGCTGCCTACC
 AGGGGCTTCTGTGTGGCTGCTTCAACCCACTCAGCTCACCTCTCTCTCCACCT
 CCTCTCCACCTGCTGGTGAGAGGTGCAAGAGGCAGATCTTTGCTGCCAGCCACTTC
 ATCCCCCTCCAGATGTTTGGACCAACACCCCTCTCTGCCACAGGCACTGCTGGAGG
 GCAGGGAGTGGGAGCCAATGAACAGGCATGCAAGTGAAGCTTCTGAGCTTTCTCC
 TGTGCGTTTGGTCTGTTTTGCCCTTACCCATAAGCCCTCGCACTCTGGTGGCAGGT
 GCCTTGTCTCAGGGCTACAGCAGTAGGGAGGTGAGTGTCTGCTGCTCGATTGAAG
 GTGACCTCTGCCCCAGATAGGTGGTGCAGTGGCTTATTAATCCGATACATAGTTT
 CTTTGTGACCAATGCTCGGTACAGAGGATGGTGAAGGCAAGGCCAGGTTGGGG
 CAGTGTGTGGCCCTGGGGCCAGGCCCAAACTGGGGGCTCTGTATATAGCTATGAA
 GAAAACACAAAGTGTATAAATCTGAGTATATATTACATGTCTTTTAAAGGGTCTG
 TTACCAGAGATTTACCCATCGGTAAGATGCTCCTGGTGGCTGGGAGGCATCAGTTG
 CTATATATTAAAAACAAAAAAGAAAAAAGGAAAAATGTTTTTAAAAAGGTCATATA
 TTTTTTGCTACTTTTGTGTTTTATTTTTTAAATTATGTTCTAAACCTATTTTCAG
 TTTAGGTCCCTCAATAAAAAATGCTGCTGCTTCATTATCTATGGGCTGTATGAAA
 GGGTGGGAATGTCCACTGGAAAGAAGGGACACCCACGGGCCCTGGGGCTAGGTCTGT
 CCCGAGGGCACCGCATGCTCCCGCGCAGGTTCCTTGTAACCTCTTCTCTTAGGTCT
 CTGCACCCAGACCTCACGACGCACCTCCTGCCTCTCCGTGCTTTTGAAAGTCAGA
 AAAAGAAGATGTCTGCTTCGAGGGCAGGAACCCCATCATGCAGTAGAGGCGCTGGG
 CAGAGAGTCAAGGCCAGCAGCATCGACCATGGATGGTTCTCTCAAGGAAACCGG
 TGGGGTTGGGCTGGGGAGGGGGCACCTACCTAGGAATAGCCACGGGTAGAGCTACA
 GTGATTAAAGAGAAAGCAAGGGCGCGGTGTGCACGCCTGTAATCCAGCACTTTGG
 GACACCGAGGTGGGCAGATCACTCAGGTGAGGAGTTTGAGACCAGCCTGGCCAAC
 TAGTGAAACCCCATCTCTACTAAAAATGCAAAAATATCTCAGGCATGGTGGCACACG
 CCTGTAATCCAGCTCCACAGGAGGCTGAGGCAGAAATCCCTGAAGCTGGGAGGCGG
 AGGTGTCAGTGAGCCGAGATTGCGCCATTGCACCTCAGCCTGGGCAACAGAGAAAAAC
 AAAAAGGAAAAACAAATGATGAAGTCTGCAGAAACTGAAACCCAGACATGTGTCTGC
 CCCCTCTATGTGGGCATGGTTTTGCCAGTGTCTTAAGTGCAGGAGAACATGTCAAC
 TGAGGCTAGTTTTGCATTCAGGTCCCTGGCTTCGTTCTTGTGGTATGCCTCCCCA
 GATCGTCCTTCTGTATCCATGTGACAGACTGTATTTGTTGGGACTGTGCGAGATC
 TTGGCTTCTACAGTTCTTCTGTCCAACTCCATCCTGTCCTCAGGAACGGGGG
 AAAATTCTCGAATGTTTTTGGTTTTTGGTGTCTGGAATTTACTTCTGCCACCTG
 CTGGTATCACTGTCTCACTAAGTGGATTCTGGCTCCCCGTACCTCATGGCTCAA
 ACTACCACTCCTCAGTCGCTATATTAAGCTTATATTTGTCTGGATTACTGCTAAAT
 ACAAAAGAAAGTTCAATATGTTTTTCTGTAGGAAAAATGGGATTGTGCTTTA

TABLE 5-continued

FGFR sequences.		
SEQ ID NO: 2	FGFR1 isoform 1 Amino acid sequence	AATTTCTGAGCTAGGGATTTTTTGGCAGCTGCAGTGTGGCGACTATTGTAAAATTC TCTTTGTTTCTCTCTGTAAATAGCACCTGCTAACATTACAATTGTATTTATGTTTA AAGAAGGCATCATTGGTGAACAGAACTAGGAAATGAATTTTAGCTCTTAAAGCA TTTGCTTTGAGACCGCACAGGAGTGTCTTTCCTTGTAACACAGTGATGATAATTTCT GCCTTGCCCTACCTTGAAGCAATGTTGTGTGAAGGATGAAGAATCTAAAGTCTT CATAAGTCCCTGGGAGAGGTGCTAGAAAAATATAAGGCATATCATAATTACAGTGA TGTCCTTGCTGTACTACTCAAATCACCCAAAATTTCCCCAAGACTGCGCTAGCT GTCAAATAAAGACAGTGAAATTGACCTG
SEQ ID NO: 3	FGFR1 isoform 2 Nucleic acid sequence	MWSWKLLFWAVLVATLCTARPSPTLPEQAQPWGAPVEVESFLVHPGDLQLRCL RDDVQSINWLRDGVQLAESNRTRI TGEVEVQDSVPADSLYACVTSPPSGSDTYF SVNVSDALPSSDDDDDDSSSEKETDNTKPNRMPVAPYWTSPKMEKKLHAVPAA KTVKFKCPSGTPNPTLRWLKNGKEFKPDHRIGGYKVRYATWSIIMDSVVPDKGNY TCIVENEYGSINHTYQLDVVERSPHRPI LQAGLPANKTVALGSNVEFMCKVYSDPQP HIQWLKHIEVNGSKI GPDNLPYVQILKTAGVNTTDKEMEVLHLRNVSFEDAGYTCCL AGNSIGLSHSHAWLTVLEALEERPAVMTSPLYLEII IYCTGAF LISCMVGSVIVYKM KSGTKKSDPHSQMAVHKLAKSI PLRRQVTVSADSSASMN SGVLLVRPSRLSSSGTPM LAGVSEYELPEDPRWELPRDRLVLGKPLGEGCFQVVLAEAI GLDKDKPNRVTKVAV KMLKSDATEKDLSDLISEMEMMKMIGKHKNI INLLGACTQDGPLYVIVEYASKGNLR EYLQARRPPGLECYCYNPSHNPEEQLS SKDLVSCAYQVARGMEYLA SKKCIHRDLAAR NVLVTEDNVMKIADFLGLARDIHHIDYKKTNGRLPVKWMAPALFDR IYTHQSDVW SFGVLLWEI FTLGGSPYPGPVPEELFKLLKEGHRMDKPSNCTNELYMMMRDCWHAVP SQRPTFKQLVEDLDR IVALTSNQEYLDLSMPLDQYSPSPDTRSSSTCSSGSDSVFSH EPLPEEPCLPRHPAQLANGGLKRR GCCGGCGCATAGCGCTCGGAGCGCTCTTGGCGGCACAGGCGCGGCGTCTCGGCGGC GGGCGGCAGCTAGCGGAGCGGCGACCGCGGTGCAGCCGAGCGCGCGGAGGAAACC GGGTGTGCCGGGAGCTGGGCGGCACGCTCCGGACGGGACCGAGACCCCTCGTAGCGC ATTGCGCGACCTCGCCTTCCCGGCGCGAGCGCGCGCTGCTTGAAAAGCCGCGG AACCCAAGGACTTTTCTCCGGTCCGAGCTCGGGGCGCCCGCAGGGCGCACGCTACC CGTGCTGCAGTCGGGCACGCGCGGCGCGCGGCGCTCCGCGAGGCGATGGAGCCCGG TCTGCAAGGAAAGTAGGCGCGCGCGCTGCGTTCTGGAGGAGGGGGGCAAGGTCT GGAGACCCCGGTGGCGGACGGGAGCCTCCCCCGCCCGCCTCCGGGGCACCGAGC TCCGGCTCCATGTTCCCGCCCGGGCTGGAGGCGCGAGCACCGAGCGCGCGCGGGA GTCGAGCGCGCGCGCGGAGCTCTTGCAGCCCGCGCAGGACCGGAACAGAGCCCGG GGCGGCGGGCCGGAGCCGGGACGCGGGCACACGCCGCTCGCACAGCCACGGCGG ACTCTCCCGAGGCGGAACCTCCACGCCGAGCGAGGGTCAGTTTGAAAAGGAGGATCG AGCTCACTGTGGAGTATCCATGGAGATGTGGAGCCTTGTACCAACCTCTAACTGCA GAACCTGGGATGTGGAGCTGGAAGTGCTCCTCTTCTGGGCTGTGCTGGTACAGCCA CACTCTGCACCGCTAGGCCCTCCCGACCTTGCTGAACAAGCCAGCCCTGGGGAG CCCGTGTGGAAGTGGAGTCTTCTTGGTCCACCCCGGTGACCTGCTGCAGCTTCGCT GTCGGCTGCGGGACGATGTGCAGAGCATCAACTGGCTGCGGGACGGGGTGCAGCTGG CGGAAAGCAACCGCACCCGCATCACAGGGGAGGAGGTGGAGGTGCAGGACTCCGTGC CCGAGACTCCGGCTCTATGCTTGCCTAACAGCAGCCCTCGGGCAGTGACACCA CCTACTTCTCCGTCAATGTTTCAGATGCTCTCCCTCCTCGGAGGATGATGATGATG ATGATGACTCCTCTTCAGAGGAGAAAGAAACAGATAACACCAAAACCAACCCCGTAG CTCCATATTGGACATCCCCAGAAAAGATGGAAAAGAAATTGCATGCAGTGCCCGCTG CCAAGACAGTGAAGTTCAAATGCCCTTCAGTGGGACCCCAAACCCACACTGCGCT GGTTGAAAAATGGCAAAGATTCAAACCTGACCACAGAATTGGAGGCTACAAGGTCC GTTATGCCACCTGGAGCATCATAATGGACTCTGTGGTGCCTCTGACAAGGGCAACT ACACCTGCATTGTGGAGAATGAGTACGGCAGCATCAACCACATACAGCTGGATG TCGTGGAGCGGTCCCTCACCGGCCATCTGCAAGCAGGGTTGCCCGCAACAAA CAGTGGCCCTGGGTAGCAACGTGGAGTTCATGTGTAAGGTGTACAGTGACCCGACG CGCACATCCAGTGGCTAAAGCACATCGAGGTGAATGGGAGCAAGATTGGCCAGACA ACCTGCCTTATGTCCAGATCTTGAAGACTGCTGGAGTTAATACCACCGACAAAGAGA TGGAGGTGCTTCACTTAAGAAATGCTCCTTTGAGGACGCGAGGGAGTATACGTGCT TGGCGGGTAACCTATCGGACTCTCCATCACTCTGCATGGTTGACCGTTCTGGAAG CCCTGGAAGAGAGGCCCGGAGTGATGACCTCGCCCCGTGACCTGGAGATCATCATCT ATTGCACAGGGGCTTCTCATCTCCTGCATGGTGGGTGGGTGATCATCGTCTACAAGA TGAAGAGTGGTACCAAGAAGAGTGACTTCCACAGCAGATGGCTGTGCACAAGCTGG CCAAGAGCATCCCTCTGCGCAGACAGGTAACAGTGTCTGTGACTCCAGTGATCCCA TGAACCTCTGGGTTCTTCTGGTTCGGCCATCAGGCTCTCTCCAGTGGGACTCCCA TGCTAGCAGGGGTCTCTGAGTATGAGCTTCCGAAGACCTCGCTGGGAGCTGCCTC GGGACAGACTGGTCTTAGGCAACCCCTGGGAGAGGGCTGCTTTGGGCAGGTGGTGT TGGCAGAGGCTATCGGGCTGGACAAGGACAAACCAACCGTGTGACCAAGTGGCTG TGAAGATGTTGAAGTCGAGCGCAACAGAGAAAGACTTGTGAGACTGATCTCAGAAA TGGAGATGATGAAGATGATCGGAAGCATAAGAATATCATCAACCTGCTGGGGGCT GCACGAGGATGGTCCCTTGATGTGTCATCGTGGAGTATGCCTCAAGGGCAACCTGC GGGAGTACCTGCAGGCCCGGAGGCCCCAGGGCTGGAATACTGCTACAACCCAGCC ACAACCCAGAGGAGCAGCTCTCTCCAAGGACCTGGTGTCTGCGCTACCAAGGTGG CCCGAGGCATGGAGTATCTGGCTCCAAGAAGTGATACACCGAGACTGGCAGCCA GGAATGCTCTGGTGACAGAGGACAATGTGATGAAGATAGCAGACTTGGCTCGCAC GGGACATTACACCATCGACTACTATAAAAAGACAACCAACGGCCGACTGCCTGTGA AGTGGATGGACCCGAGGCATTATTTGACCGGATCTACACCCACAGAGTGTGTGT GGTCTTTTCGGGTGCTCCTGTGGGAGATCTTCACTCTGGGCGGCTCCCATACCCG GTGTGCTGTGGAGGAACCTTTCAAGTGTCTGAAGGAGGTACCCGATGGACAAGC CCAGTAACCTGCACCAACGAGCTGTACATGATGATGCGGGACTGCTGCATGCACTGC CCTCACAGAGACCCACTTCAAGCAGTGTGTGAAGACCTGGACCGCATCGTGGCT

TABLE 5-continued

FGFR sequences.	
SEQ ID NO: 4	FGFR1 isoform 2 Amino acid sequence TGACCTCCAACCCAGGAGTACCTGGACCTGTCCATGCCCTGGACCACTACTCCCCCA GCTTTCGCCGACACCCGGAGCTCTACGTGCTCCTCAGGGGAGGATTCCGCTTCTCTC ATGAGCCGCTGCCCGAGGAGCCCTGCCTGCCCCGACACCCAGCCAGCTTGCCAATG GCGGACTCAAACGCCGCTGACTGCCACCCACACGCCCTCCCCAGACTCCACCGTCTAG CTGTAAACCTCACCCACAGCCCTGCTGGGCCACACCTGTCCGTCCTGTCCCCCT TTCTGTCTGGCAGGAGCGGCTGCCTACCAGGGGCTTCTGTGTGGCTGCCTTCA CCCCACCTCAGCTCACCTCTCCCTCCACCTCCTCTCCACCTGTGGTGAGAGGTGCAA AGAGGCAGATCTTTGCTGCCAGCCACTTCATCCCTCCAGATGTTGGACCAACACC CCTCCTGCCACCAGGCACTGCCTGGAGGGCAGGGAGTGGGAGCCAATGAACAGGCA TGCAAGTGAGAGCTTCCTGAGCTTTCTCCTGTGCGTTTGGTCTGTTTTGCCTTCAAC CATAAGCCCTCGCACTCTGGTGGCAGGTGCCTGTCTCAGGGCTACAGCAGTAGG GAGGTCACTGCTTCGTGCTCGATTGAAGGTGACCTCTGCCCCAGATAGGTGGTGCC AGTGGCTTATTAATTCCGATACTAGTTTGTCTTGTGACCAATGCTGGTACCAGA GGATGGTGAGGCGAAGGCCAGGTGGGGGCAAGTGTGTGGCCCTGGGGCCAGCCCC AAACCTGGGGGCTCTGTATATAGCTATGAAGAAAACACAAAGTGATATAATCTGAGTA TATATTTACATGTCTTTTAAAGGGTCGTTACCAGAGATTACCCATCGGGTAAGA TGCTCCTGGTGGCTGGGAGGCATCAGTTGCTATATATTAAGAAAAGAAAAAGAAAA AAGGAAATGTTTTTAAAGGTCATATATTTTGTCTACTTTGTCTGTTTTATTTT TTTAAATATGTTCTAAACCTATTTTTCAGTTTAGTCCCTCAATAAAAAATGTCTGCT GCTTCATTATCTATGGGCTGTATGAAAAGGGTGGGAATGTCCACTGGAAAGAGGG ACACCCACGGGCCCTGGGGCTAGGTCTGTCCCGAGGGCACCAGTGTCCCGCGCA GGTTCCCTGTAACTCTTCTTCTAGGTCTGCACCCAGACCTCACGACGACCTCC TGCTCTCCGCTGCTTTTGGAAAGTCAGAAAAGAGATGTCTGCTTCGAGGGCAGG AACCCCATCCATGCAGTAGAGGCGCTGGGCAGAGAGTCAAGGCCAGCAGCCATCGA CCATGGATGGTTTCTTCCAAGGAAACCGTGGGGTGGGGTGGGGAGGGGGCACCTA CCTAGGAATAGCCACGGGGTAGAGCTACAGTGATTAAAGAGAAAGCAAGGCGCGGT TGCTCACGCTGTAAATCCAGCACTTTGGGACACCGAGGTGGGCAGATCACTTCAGG TCAGGAGTTTGAGACCAGCCTGGCCAACCTAGTGAACCCCATCTCTACTAAAAATG CAAAAATTATCCAGGCATGGTGGCACACGCTGTAAATCCAGCTCCACAGGAGGCTG AGGCAGAAATCCCTGAAGCTGGGAGGCGGAGGTGTCAGTGAGCCGAGATTGCGCCAT TGCACTCCAGCCTGGGCAACAGAGAAAAACAAAAGGAAAAACAAATGATGAAGGTCTG CAGAACTGAAACCCAGACATGTGTCTGCCCTCTATGTGGGCATGTTTTCAGG TGCTTCTAAGTGCAGGAGACATGTACCTGAGGCTAGTTTGTGATTCAGGTCCCTG GCTTCGTTTCTGTTGGTATGCCTCCCGAGATCGTCTTCTGTATCCATGTGACCA GACTGTATTTGTTGGGACTGTCGAGATCTTGGCTTCTTACAGTTCTTCTGTCCAA ACTCCATCCTGTCTCCTCAGGAACGGGGGAAAAATCTCCGAATGTTTTTGGTTTTT GGCTGCTTGGAAATTAATCTCTGCCACCTGCTGGTCACTGCTCTCACTAAGTGGG TTCTGGCTCCCCGTACCTCATGGCTCAAACTACCACTCCTCAGTCGCTATATTA GCTTATATTTTGTGGATTACTGCTAAATACAAAAGAAAGTTCAATATGTTTTTCAAT TCTGTAGGAAAAAGGAGTTGCTGCTTTAAATTTCTGAGCTAGGAGTTTTTGGCAG CTGCACTGTTGGGCACTATTGTAATAATCTCTTGTCTCTCTGTAATAGCACCT GCTAACATTACAATTTGTATTTATGTTTAAAGAGGCATCATTTGGTGAACAGAACT AGGAAATGAATTTTGTAGCTTTAAAGCATTTGCTTTGAGACCGCACAGGAGTGTCT TTCTTGTAAACAGTGATGATAATTTCTGCTTGGCCCTACCTTGAAGCAATGTTG TGTGAAGGATGAAGAACTTAAAGTCTTCAAGTCTTGGGAGAGGTGCTAGAAA AATATAAGGCACTATCATAATTACAGTGATGCTCTGTCTGTACTACTCAAATCACC CACAAATTTCCCCAAGACTGCGCTAGCTGTCAAATAAAGACAGTGAATTTGACCT GA
SEQ ID NO: 5	FGFR1 isoform 3 Nucleic acid sequence AGCGCTCTTGCGGCCACAGGCGCGGCTCCTCGGCGGCGGGCGGAGCTAGCGGGAG CCGGGACGCGCGGTGCAGCCGAGCGCGCGGAGGAACCCGGGTGTGCGGGAGCTGGG CGGCCACGTCGGACGGGACCGAGACCCCTCGTAGCGCATTCGGGCGACCTCGCCTT CCCCAGGCGCGAGCGCGCCCTGCTTGAAAGCCGCGGAACCAAGGACTTTTCTCC GGTCCGAGCTCGGGGCGCCCGCAGGGCGCACGGTACCCGTGCTGCACTCGGGCAGC CCGCGGCGCGGGGCTCCGACGGGCGATGGAGCCCGGTCTGCAAGGAAAGTGAAGG GCCGCCCTGCGTTCTGGAGGAGGGGGGCACAAGGTCTGGAGACCCCGGGTGGCGGA CGGGAGCCCTCCCCCGCCCCGCTCCGGGGCACAGCTCCGGCTCCATTGTTCCCG CCGGGGCTGGAGGCGCCGAGCACCGAGCGCCCGGGAGTCAGCGCGCGGCGCGGA GCTCTTGCAGCCCGCGCAGGACCGAACAGAGCCCGGGGCGGGCGGGCGGAGCCGG GGACGCGGGCACACGCCCGCTCGCACAAAGCCACGGCGGACTCTCCGAGGCGGAACC TCCACGCCGAGCGAGGTCAGTTTGAAAGGAGGATCGAGCTCACTGTGGAGTATCC ATGGAGATGTGGAGCCTGTGACCAACCTCTAACTGCAGAACTGGGATGTGGAGCTG

TABLE 5-continued

FGFR sequences.

GAAGTGCCTCCTCTTCTGGGCTGTGCTGGTCACAGCCACACTCTGCACCGCTAGGCC
 GTCCCCGACCTTGCCCTGAACAAGCCAGCCCTGGGGAGCCCTGTGGAAGTGGAGTC
 CTTCTCTGGTCCACCCCGGTGACCTGCTGCAGCTTCGCTGTGCGGTGCGGGACGATGT
 GCAGAGCATCAACTGGCTGCGGGACGGGGTGACGCTGGCGGAAAGCAACCGCACCCG
 CATCACAGGGGAGGAGGTGGAGGTGCAGGACTCCGTGCCCGCAGACTCCGGCCTCTA
 TGCTTGCGTAACAGCAGCCCTCGGGCAGTGACACCACCTACTTCTCCGTCAATGT
 TTCAGATGCTCTCCCCCTCGGAGGATGATGATGATGATGACTCTCTTTCAGA
 GGAGAAAGAAACAGATAACACCAACCAACCGTATGCCCGTAGCTCCATATTGGAC
 ATCCCCAGAAAAGATGGAAGAAATTCATGTCAGTGCCGGCTGCCAAGACAGTGAA
 GTTCAAATGCCCTTCCAGTGGGACCCCAACCCACACTGCGCTGGTTGAAAAATGG
 CAAAGAATTCAAACCTGACCACAGAATTGGAGGTACAAGGTCCGTATGCGACCTG
 GAGCATCATAATGGACTCTGTGTGCCCTCTGACAAGGGCAACTACACCTGCAATTGT
 GGAGAATGAGTACGGCAGCATCAACCACATACAGCTGGATGTCGTGGAGCGGTG
 CCCTCACCGGCCCATCCTGCAAGCAGGGTTGCCGCCAACAAACAGTGGCCCTGGG
 TAGCAACGTGGAGTTCATGTGTAAGGTGTACAGTGACCCCGCAGCCGCACATCCAGTG
 GCTAAAGCACATCGAGGTGAATGGGAGCAAGATTGGCCAGACAACCTGCCTTATGT
 CCAGATCTTGAAGACTGCTGGAGTTAATACACCGACAAAGAGATGGAGGTGCTTCA
 CTTAAGAAATGCTCTCTTTGAGGACGAGGGGAGTATACGTGCTTGGCGGGTAATC
 TATCGGACTCTCCCATCACTCTGCATGGTTGACCGTTCTGGAAGCCCTGGAAGAGAG
 GCCGGCAGTGATGACCTCGCCCCCTGTACCTGGAGATCATCATCTATTGCACAGGGGC
 CTTCTCATCTCTGTCATGTTGGGGTGGTTCATCGTCTACAAGATGAAGAGTGGTAC
 CAAGAAGAGTGACTTCCACAGCCAGATGGCTGTGCACAAGCTGGCCAAGAGCATCCC
 TCTGCGCAGACAGGTGTCTGCTGACTCCAGTGTCATCCATGAACCTCGGGGTTCTTCT
 GGTTTCGGCCATCACGGCTCTCTCCAGTGGGACTCCCATGCTAGCAGGGGTCTCTGA
 GTATGAGCTTCCCGAAGACCTCGCTGGGAGCTGCCCTCGGGACAGACTGGTCTTAGG
 CAAACCCCTGGGAGAGGGCTGCTTTGGGAGGTGGTGTGGCAGAGGCTATCGGGCT
 GGACAAGGACAAACCCCAACCGTGTGACCAAGTGGCTGTGAAGATGTTGAAGTCGGA
 CGCAACAGAGAAAGACTTGTGACAGCTGATCTCAGAAATGGAGATGATGAAGATGAT
 CGGGAAGCATAGAAATATCATCAACCTGCTGGGGGCTGCACGCAGGATGGTCCCTT
 GTATGTGTCGTGGAGTATGCTTCAAGGGCAACCTGCGGGAGTACCTGCAGGCCCG
 GAGGCCCCAGGGCTGGAATACTGCTACAACCCAGCCACAACCCAGAGGAGCAGCT
 CTCTTCAAGGACCTGGTGTCTTGCCTTACAGGTGGCCGAGGCTAGGAGTATCT
 GGCTTCAAGAGTGATACACCGAGACTTGGCAGCCAGGAATGCTCTGGTGACAGA
 GGACAATGTGATGAAGATAGCAGACTTTGGCCCTCGCACGGGACATTACCCACATCGA
 CTACTATAAAAGACAAACCAACGGCCGACTGCCGTGTAAGTGGATGGCACCCGAGGC
 ATTTATTTGACCGGATCTACACCCACAGAGTGTGTGTGCTTTTCGGGGTGCTCCT
 GTGGGAGATCTTCACTCTGGGCGGCTCCCATACCCGGTGTGCCCTGTGGAGGAAT
 TTTCAAGCTGCTGAAGGAGGGTCAACGCATGGACAAGCCAGTAACCTGCACCAACGA
 GCTGTACATGATGATGCGGGACTGCTGGCATGCAGTGCCCTCACAGAGACCCACCTT
 CAAGCAGCTGGTGAAGACCTGGACCGCATCGTGGCTTGAACCTCAACCCAGGAGTA
 CCTGGACCTGTCCATGCCCTGGACAGTACTCCCGAGCTTTCCCGACACCCGGAG
 CTCTACGTGCTCCTCAGGGAGGATTCCGTCTTCTCTCATGAGCCGTGCCCGAGGA
 GCCCTGCCTGCCCGACACCCAGCCAGCTTGGCAATGGCGGACTCAAACGCCGCTG
 ACTGCCACCCACACGCCCTCCCGAGACTCCACCGTCAGCTGAACCTCAACCCACAG
 CCCCTGCTGGGCCACCACTGTCCGTCCCTGTCCCTTTCTGCTGGCAGGAGCCG
 GCTGCCCTACAGGGGCTTCTGTGTGGCTGCCTTCAACCCACTCAGCTCACCTCT
 CCCTCCACCTCTCTCCACCTGCTGGTGAGAGGTGCAAGAGGCAGATCTTTGTCTG
 CAGCCACTTTCATCCCTCCAGATGTTGGACCAACACCCCTCTCTGCCACCCAGGCAC
 TGCTTGAGGGCAGGGAGTGGGAGCCAATGAACAGGCATGCAAGTGAAGAGCTTCTCG
 AGCTTTCTCCTGTGCGTTTGGTCTGTTTGCCTTCAACCCATAAGCCCTCGCACTCT
 GGTGGCAGGTGCCCTTGTCTCAGGGCTACAGCAGTAGGGAGGTGAGTGTCTCGTGCC
 TCGATTGAAGGTGACCTCTGCCCCAGATAGGTGGTGGCAGTGCTTATTAATTCGCA
 TACTAGTTTGTCTTGTGACCAATGCTGGTACCAGAGGATGGTGAGGCGAAGGCC
 AGGTGGGGGAGTGTGTGGCCCTGGGGCCAGCCCAAACTGGGGGCTCTGTATA
 TAGCTATGAAGAAAAACAAAGTGTATAAATCTGAGTATATATTTACATGCTTTTTT
 AAAAGGGTCTGTACCAGAGATTACCCATCGGGTAAGATGCTCTGTGTGGCTGGGAG
 GCATCAGTTGCTATATATTAATAACAAAAAGAAAAAAGGAAAAATGTTTTAAAAA
 AGGTCATATATTTTTTGCTACTTTTGCTGTTTTATTTTTTAAATATGTTCTAAAC
 CTATTTTCAGTTTAGGTCCCTCAATAAAATTTGCTGCTGCTTCAATTATCTATGGGC
 TGTATGAAAAGGGTGGGAATGTCCACTGGAAGAAGGGACACCCACGGGCCCTGGGG
 CTAGGTCTGTCCCGAGGGCACCGCATGCTCCCGGGCAGGTTCCCTGTAACTCTTC
 TTCTTAGTCTTGACCCAGACCTCACGACGCACCTCTGCTCTCCGCTGCTTTTG
 GAAAGTCAGAAAAAGAGATGTCTGCTTCAGGGCAGGAACCCCATCCATGCAGTAG
 AGGCGCTGGGCAGAGAGTCAAGGCCAGCAGCCATCGACCATGGATGGTTTCTCCCA
 AGGAAACCGGTGGGTTGGGCTGGGGAGGGGGACCTACCTAGGAATAGCACGGGG
 TAGAGCTACAGTGATTAAGAGGAAAGCAAGGGCGCGGTGCTCACGCTGTAATCCC
 AGCACTTTGGGACACCGAGGTGGGCAGATCACTTCAGGTGAGGATTTGAGACCAG
 CTGGCCAACTTAGTGAACCCCATCTCTACTAAAAATGCAAAATATCCAGGCATG
 GTGGCACACGCCTGTAATCCAGCTCCACAGGAGGCTGAGGCAGAAATCCCTTGAAGC
 TGGGAGGCGGAGGTTGCAAGTGAGCCGAGATTGCCCATTCGACTCCAGCTGGGCAA
 CAGAGAAAAAAGGAAACAAATGATGAAGGTCTGCAGAACTGAAACCCAGAC
 ATGTGTCTGCCCCCTCTATGTGGGCATGGTTTTGCCAGTGCTTCTAAGTCAGGAGA
 ACATGTCACTGAGGCTAGTTTTGCATTCAAGTCCCTGGCTTCGTTCTTGTGGTA
 TGCTTCCAGATCGTCTTCTGTATCCATGTGACAGACTGTATTGTTGGGACT
 GTCGAGATCTTGCTTCTTACAGTTCTTCTGTCCAACTCCATCTGTCCCTCAG
 GAACGGGGGAAAAATCTCCGAATGTTTTTGGTTTTTGGCTGCTTGAATTTACTT
 CTGCCACCTGCTGGTCACTCTGCTCACTAAGTGGATTCTGGCTCCCCCGTACCT
 CATGGCTCAAACTACCACTCTCAGTCGTATATTAAGCTTATATTTTGTGGATT

TABLE 5-continued

FGFR sequences.		
SEQ ID NO: 6	FGFR1 isoform 3 Amino acid sequence	ACTGCTAAATACAAAAGAAAGTTCAATATGTTTTTCATTTCTGTAGGGAAAAATGGGAT TGCTGCTTTAAATTTCTGAGCTAGGGATTTTTTGGCAGCTGCAGTGTGGCGACTAT TGTAATAATCTCTTTGTTTCTCTCTGTAAATAGCACCTGCTAACATTACAAATTTGTA TTTATGTTTAAAGAAGGCATCATTGGTGAACAGAACTAGGAAATGAATTTTAGCT CTTAAAAGCATTTGCTTTGAGACCGCACAGGAGTGTCTTTCCTGTAAAAACAGTGAT GATAATTTCTGCCTTGCCCTACCTTGAAGCAATGTTGTGTGAAGGATGAAGAATC TAAAAGTCTTCATAAGTCCTTGGGAGAGGTGCTAGAAAAATATAAGGCATATCATA ATTA MWSWKCLLFWAVLVATLCTARPSPTLPEQAQPWGAPVEVESFLVHPGDLQLRCL RDDVQSINWLRDGVQLAESNRTRI TGEVEVQDSVPADSLYACVTSPPSGSDTTYF SVNVSDALPSSDDDDDDSSSEKETDNTKPNRMPVAPYWTSPKMEKKLHAVPAA KTVKFKCPSGTPNPTLRWLKNGKEFKPDHRIGGYKVRYATWSIIMDSVVPSPDKNY TCIVENEYGSINHTYQLDVVERSPHRPI LQAGLPANKTVALGSNVEFMCKVYSDPQP HIQWLKHIEVNGSKI GPDNLPIVQILKTAGVNTTDKEMEVLHLRNVSFEDAGYTCCL AGNSIGLSHSHSAWLTVLEALEERPAMVTSPLYLEII IYCTGAF LISCMVGSVIVYKM KSGTKKSDPHSQMAVHKLAKSI PLRRQVSADSSASMNSGVLLVRPRLSSSGTPMLA GVSEYELPEDPRWELPRDRLVLGKPLGEGCFGQVVLAEI GLDKDKPNRVTKVAVKM LKSDATEKDLSDLISEMEMMMKIGKHKNI INLLGACTQDGPLYVIVEYASKGNLREY LQARRPGLLEYCYNPSHNPEEQLSKDLVSCAYQVARGMEYLASKKCIHRDLAARNV LVTEDNMVKIADFLARDIHHIDYKKTTNGRLPVKWMPEALFDRIYTHQSDVWSF GVLLWEIFTLGGSPYPGVPEELFKLLKEGHRMDKPSNCTNELYMMMRDCWHAVPSQ RPTFKQLVEDLDRIVALTSNQEYLDLSMPLDQYSPSPFDPTRSSSTCSSGSDSVFSHEP LPEEPCLPRHPAQLANGGLKRR
SEQ ID NO: 7	FGFR1 isoform 4 Nucleic acid sequence	CCCTTTCACCTCCTGGCTCCTCCCGGGCGATCCGCGCCCTTGGGTCTCCCCTCCC TTCCTCCGTCGCGTCTCCTGCGCCCTCCCTGCGCTCGTCCCGCCGCTCTTCCC GCCGCCCACTTTTCTCCAACTCGCGCTCGGGAGCTGGCGAGGCGGCGCGGCTCC TCAAAGTGGGAGAGCTTCAAGGTCACGTGGTCCGTCCAGCCCTGCTATCTCACCAG ACACTGTCCACCTGTATGTTGGATCAGTACTCCAGTGAGAAGACAGCAGGCCTTT CACCCATGCAGCCATTCACTCTTTCATAACCACTGTGATGGAGGCAAGGGTCAGTT TGAAAAGGAGGATCGAGCTCACTGTGGAGTATCCATGGAGATGTGGAGCCTTGTAC CAACTCTAACTGCAGAACTGGGATGTGGAGCTGGAAGTGCCTCCTCTCTGGGCTG TGCTGGTACAGCCCACTCTGCACCGCTAGGCCGTCCCGACCTTGCTGAACAAG CCCAGCCCTGGGGAGCCCTGTGGAAGTGGAGTCTTCTGGTCCACCCCGGTGACC TGCTGCAGCTTCGCTGTGCGCTGCGGACGATGTGCAGAGCATCACTGGCTGCGGG ACGGGGTGCAGCTGGCGGAAAGCAACCGCACCCGCATCACAGGGGAGGAGGTGGAGG TGCAAGGACTCCGTGCCCGCAGACTCCGGCCTCTATGCTTGCCTAACAGCAGCCCT CGGGCAGTGACACCACCTACTTCTCCGTCAATGTTTCAGATGCTCTCCCCTCCTCGG AGGATGATGATGATGATGATGACTCCTCTTCAGAGGAGAAAGAAACAGATAACACCA AACCAAACCCGTAGCTCCATATTGGACATCCCAGAAAAGATGGAAGAAATATGC ATGCAGTGCCGGCTGCGCAAGCAGTGAAGTCAAATGCCCTTCCAGTGGGACCCCAA ACCCCACTGCGCTGGTTGAAAATGGCAAAGAAATCAAACCTGACCACAGAATTG GAGGCTACAAGTCCGTTATGCCACCTGGAGCATCATAATGGACTCTGTGGTGCCT CTGACAAGGGCACTACACCTGCATTGTGAGAAATGAGTACGGCAGCATCAACCA CATACCAGCTGGATGTCGTGGAGCGGTCCCTCACCGGCCCATCTCTCAAGCAGGGT TGCCCGCAACAAAACAGTGGCCCTGGGTAGCAACGTGGAGTTCATGTGAAGTGT ACAGTGACCCGCGCAGCCGACATCCAGTGGCTAAAGCACATCGAGGTGAATGGGAGCA AGATTGGCCCAAGCAACCTGCCTTATGTCCAGATCTTGAAGACTGCTGGAGTTAATA CCACCGACAAGAGATGGAGGTGCTTCACTTAAGAAATGTCCTCTTGGAGACGCGAG GGGAGTATACGTGCTTGGCGGGTAACCTATCGGACTCTCCATCACTCTGCATGGT TGACCGTTCTGGAAGCCCTGGAAGAGAGGCGGCGAGTGATGACCTCGCCCTGTACC TGGAGATCATCATCTATTGCACAGGGCCCTTCTCATCTCCTGCATGTGGGGTGG TCATCGTCTACAAGATGAAGAGTGGTACCAAGAGAGTGACTTCCACAGCCAGATGG CTGTGCACAAGCTGGCCAAGAGCATCCCTCTGCGCAGACAGGTAACAGTGTCTGTG ACTCCAGTGATCATGAACCTTGGGGTCTTCTGGTTTCGGCCATCACGGCTCTCCT CCAGTGGGACTCCCATGCTAGCAGGGGTCTCTGAGTATGAGCTTCCCGAAGACCTC GCTGGGAGCTGCCCTCGGGACAGACTGGTCTTAGGCAAAACCTGGGAGAGGGCTGT TTGGGAGGTGGTGTGGCAGAGGCTATCGGGCTGGACAAGGACAAACCAACCGTG TGACCAAAGTGGCTGTGAAGATGTTGAAGTCGGACGCAACAGAGAAAGACTTGTGAG ACCTGATCTCAGAAATGGAGATGATGAAGATGATCGGAAGCATAAGAATATCATCA ACCTGCTGGGGGCTCGACGCAGGATGGTCCCTTGATGTGATCGTGGAGTATGCTC CCAAGGGCAACTGCGGGAGTACCTGCAGGCCCGGAGGCCCCAGGGCTGGAATACT GCTACAACCCAGCCACAACCCAGAGGAGCAGCTCTCTCCAAGGACCTGGTGTCT GCGCTACAGGTGGCCGAGGATGGAGTATCTGGCTCCAAGAGTGCATACACC GAGACCTGGCAGCCAGGAATGTCTGGTGACAGAGGACAATGTGATGAAGATAGCAG ACTTTGGCCTCGCACGGGACATTCACCATCGACTACTATAAAAGACAACCAACG GCCGACTGCTGTGAAGTGGATGGCACCAGAGGCAATTTTGAACCGATCTACACCC ACCAGAGTGTGTGTGGTCTTTCGGGGTGTCTGTGGGAGATCTTCACTCTGGGCG GCTCCCCATACCCCGTGTGCTGTGGAGGAACTTTCAAGCTGCTGAAGGAGGGTC ACCGCATGGACAAGCCAGTAACCTGCACCAACGAGCTGTACATGATGATGCGGGACT GCTGGCATGCACTGCCCTCACAGAGACCCACCTTCAAGCAGCTGGTGAAGACCTGG ACCGCATCGTGGCTTGACCTCCAACAGGAGTACCTGGACCTGTCCATGCCCTGG ACCGTACTCCCCAGCTTTCCGACACCCGAGCTCTACGTGCTCTCAGGGGAGG ATTCCGTCTTCTCTCATGAGCCGCTGCCCGAGGAGCCCTGCCTGCCCCGACACCCAG CCCAGCTTGGCAATGGCGGACTCAAACGCGCTGACTGCCACCCACACGCCCCCTCC AGACTCCACCGTCAGCTGTAACCTCACCCACAGCCCTGCTGGGCCCAACACCTGT CCGTCCCTGTCCCTTTCTGCTGGCAGGAGCCGGCTGCTACAGGGGCTTCTCTG

TABLE 5-continued

FGFR sequences.	
SEQ ID NO: 8	TGTGGCCTGCCTTCACCCCACTCAGCTCACCTCTCCCTCCACCTCTCTCCACCTGC TGGTGAGAGGTGCAAGAGGCAGATCTTTGCTGCCAGCCACTTCATCCCTCCCAGA TGTTGGACCAACACCCCTCCCTGCCACCAGGCACTGCCTGGAGGGCAGGAGTGGGA GCCAATGAACAGGCATGCAAGTGAGAGCTTCCCTGAGCTTTCTCCTGTCGGTTTGGTC TGTTTTGCCCTTCACCCATAAGCCCTCGCACTCTGGTGGCAGGTGCCTTGTCTCAG GGCTACAGCAGTAGGGAGGTGAGTGTCTCGTGCCTCGATTGAAGGTGACCTCTGCCC CAGATAGGTGGTGGCAGTGGCTTATTAATTCGATACTAGTTTGGCTTTGCTGACCAA ATGCTCTGGTACAGAGGATGGTGAGGCGAAGGCCAGGTTGGGGGCAGTGTGTGGCC CTGGGGCCAGCCCCAACTGGGGGCTCTGTATATAGCTATGAAGAAACACAAAGT GTATAAATCTGAGTATATATTTACATGTCTTTTAAAGGGTCGTTACAGAGATTT ACCCATCGGGTAAGATGCTCCTGGTGGCTGGAGGCATCAGTTGCTATATATTAATA ACAAAAAGAAAAAAGGAAAAATGTTTTTAAAGGTCATATATTTTTTGTACTT TTGCTGTTTTATTTTTTAAATTATGTTCTAAACCTATTTTCAGTTTAGGTCCCTCA ATAAAAATTGCTGCTGCTTCATTTATCTATGGGCTGTATGAAAAGGGTGGGAATGTC CACTGGAAGAAGGGACACCCACGGGCCCTGGGGCTAGGTCTGTCCGAGGGCACCG CATGCTCCCGGCGCAGGTTCTTGTAACTCTCTCTCTCTAGGTCTGCAACCCAGACC TCACGACGCACTCTGCTCTCCGCTGCTTTTGGAAAGTCAGAAAAAGAGATGTC TGCTTCGAGGGCAGGAACCCCATCCATGCAGTAGAGGCGCTGGGCAGAGAGTCAAGG CCCAGCAGCCATCGACCATGGATGGTTTCTCCAAGGAAACCGGTGGGGTTGGGCTG GGGAGGGGGCACTTACCTAGGAATAGCCACGGGTAGAGCTACAGTGATTAAGAGGA AAGCAAGGGCGCGGTGCTCACGCTGTAATCCAGCACTTTGGGACACCGAGGTGG GCAGATCACTTCAGGTGAGGATTTGAGACCAGCCTGGCCAACCTAGTGAACCCCA TCTCTACTAAAAATGCAAAATATCCAGGCATGGTGGCACACGCTGTAATCCAG CTCCACAGGAGGCTGAGGCAGAAATCCCTTGAAGCTGGGAGGCGAGGTTCAGTGTAG CCGAGATTGCGCCATTGCACCTCCAGCTGGGCAACAGAGAAAAACAAAAGGAAAAACA AATGATGAAGTCTGCAGAACTGAAACCCAGACATGTGCTGCCCTCTATGTGG GCATGGTTTTGCCAGTCTCTAAGTGCAGAGAACATGTCACCTGAGGCTAGTTTT GCATTCAAGTCCCTGGCTTCGTTTCTTGTGGTATGCCTCCCGAGATCGTCTCTCCT GTATCCATGTGACCACTGTATTTGTTGGGACTGTGCGAGATCTTGGCTCTTACA GTTCTTCTGTCCAACTCCATCTGTCCCTCAGGAACGGGGGAAAAATCTCCGAA TGTTTTTGGTTTTTTGGCTGCTTGAATTTACTTCTGCCACCTGCTGGTCATCACTG TCCTCACTAAGTGGATTCTGGCTCCCCGTACCTCATGGCTCAAATACCACTCCTC AGTCGCTATATTAAGCTTATATTTTGTGGATTACTGCTAAATACAAAAGAAAGT CAATATGTTTTCACTTCTGTAGGAAAAATGGGATTGCTGCTTAAATTTCTGAGCTA GGGATTTTTTGGCAGCTGCAGTGTGGCGACTATTGTAAATTTCTTTGTTTCTCT CTGTAAATAGCACCTGCTAACATTACAATTGTATTATGTTTAAAGAGGCATCAT TTGGTGAACAGAACTAGGAAATGAATTTTAGCTCTTAAAGCATTTGCTTTGAGAC CGCACAGGAGTGTCTTCTTGTGTAACAGTGTATGATAATTCTGCTCTGGCCCTAC CTTGAAGCAATGTTGTGTAAGGATGAAGAATCTAAAGTCTTCATAAGTCTTGG GAGAGGTGCTAGAAAAATATAAGGCATATCATAATTACAGTGTATGCTTGTCTGT ACTACTCAAATCACCAAAATTTCCCAAGACTGGCTAGCTGTCAAATAAAGA CAGTGAAATTGACCTGA
SEQ ID NO: 9	MEARVSLKRRIELTVEYPWRCGALSPTSNCRTGMWSWKCLLFWAVLVATLCTARPS PTLPEQAQPWGAPVEVESFLVHPGDLQLRLRDDVQSINWLRDGVQLAESNRTRI TGEEVEVQDSVPADSGLYACVTTSSPSGSDTTYFSVNVSDALPSEDDDDDDSSSEE KETDNTKPNVPVAPYWTSPKMEKKLHAVPAKTVKFKCPSSGTPNPTLRWLKNGKEF KPDHRIIGYKVRATWSIIMDSVVP SDKNYTCIVENEYGSINHTYQLDVVERSPHR P I L Q A G L P A N K T V A L G S N V E F M C K V Y S D P O P H I Q W L K H I E V N G S K I G P D N L P Y V Q I L K T A G V N T D K E M E V L H L R N V S F E D A G E Y T C L A G N S I G L S H H S A W L T V L E A L E E R P A V M T S P L Y L E I I I Y C T G A F L I S C M V G S V I V Y K M K S G T K K S D F H S Q M A V H K L A K S I P L R R Q V T V S A D S S A S M N S G V L L V R P S R L S S S G T P M L A G V S E Y E L P E D P R W E L P R D R L V L G K P L G E G C F G Q V V L A E A I G L D K D K P N R V T K V A V K M L K S D A T E K D L S D L I S E M E M M K M I K H K N I I N L L G A C T Q D G P L Y V I V E Y A S K N L R E Y L Q A R R P P G L E Y C Y N P S H N P E E Q L S S K D L V S C A Y Q V A R G M E Y L A S K K C I H R D L A A R N V L V T E D N V M K I A D F L A R D I H H I D Y Y K K T T N G R L P V K W M A P E A L F D R I Y T H Q S D V W S F G V L L W E I F T L G G S P Y P G V P V E E L F K L L K E G H R M D K P S N C T N E L Y M M R D C W H A V P S Q R P T F K Q L V E D L D R I V A L T S N Q E Y L D L S M P L D Q Y S P S F P D T R S S T C S S G E D S V F S H E P L P E E P C L P R H P A Q L A N G G L K R R
SEQ ID NO: 9	CGCCTCTTGGCGCCACAGGCGCGCGCTCCTCGGCGGCGGGCGGCAGCTAGCGGGAGC CGGGACGCCGGTGTCAGCCGCGAGCGCGCGGAGGAACCCGGGTGTGCCGGGAGCTGGGC GGCCAGCTCGGACCGGACCGAGACCCCTCGTAGCGCATTCGCGCGACCTCGCCTTC CCCGGCCGCGAGCGCGCGCTGTCTGAAAAGCCGCGGAACCCAGGACTTTCTCCG GTCCGAGCTCGGGCGCCCGCAGGGCGCACCGTACCCGTGCTGCAGTCCGGCACGC CGCGCGCCGGGGCTCCGCAAGGCGATGGAGCCCGGTCTGCAAGGAAAGTGAGGCG CCGCCGCTGCGTTCTGAGGAGGGGGGCACAAGGTCTGGAGACCCCGGTGGCGGAC GGGAGCCCTCCCCCGCCCGCTCCGGGGCACAGCTCCGGCTCCATTGTTCCCGC CCGGCTGGAGGCGCGAGCACGAGCGCCCGGGAGTCGAGCGCCGCGCGCGGAG CTCTTGCAACCCCGCAGGACCCGAACAGAGCCCGGGGGCGGGCGGGCGGAGCCGGG GACGCGGGCACACGCGCCGCTCGCACAGCCACGGCGGACTCTCCGAGGGCGGAACCT CCACGCGAGCGAGGGTCAGTTGAAAAGGAGGATCGAGCTCACTGTGGAGTATCCA TGGAGATGTGAGCCTTGTACCAACCTCTAATGCAGAACTGGGATGTGGAGCTGG AAGTGCCTCTCTTCTGGGCTGTGCTGGTCACAGCCCACTCTGCACCCGTAGGCCG TCCCCGACCTTGCTCTGAACAAGGATGGCAGCTGTGACCCGGGATTTCCGGTGAGATG TTCTGCACTCTGGCCGGGTCTGTCAGCGCAAGCCAGCCCTGGGGAGCCCCGTGTGG AAGTGGAGTCTCTTCTGGTCCACCCCGGTGACCTGTGTCAGCTTCGCTGTGCGCTGC GGGACGATGTGCAGAGCATCACTGGCTGCGGGACGGGGTGACGTGGCGGAAGCA ACCGCACCCGCATCACAGGGGAGGAGTGGAGTGCAGGACTCCGTGCCCGCAGACT

TABLE 5-continued

		FGFR sequences.
SEQ ID NO: 10	FGFR1	CCGGCCTCTATGCTTGCCTAACAGCAGCCCCCTCGGGCAGTGACACCACCTACTTCT CCGTCAATGTTTCAGATGCTCTCCCCCTCCTCGGAGGATGATGATGATGATGACT CCTCTTCAGAGGAGAAAGAAACAGATAACACCAAACCAAACCGTATGCCCGTAGCTC CATATTGGACATCCCCAGAAAAGATGGAAAAGAAATTGCATGCAGTGCCCGGTGCCA AGACAGTGAAGTTCAAATGCCCTTCCAGTGGGACCCCAAACCCACACTGCGCTGGT TGAAAAATGGCAAAGAATTCAAACCTGACCACAGAATTGGAGGCTACAAGGTCCGTT ATGCCACCTGGAGCATCATAATGGACTCTGTGGTGCCCTCTGACAAGGGCAACTACA CCTGCATTGTGGAGAATGAGTACGGCAGCATCAACCACACATACCAGCTGGATGTCG TGGAGCGGTCCCCCTACCGGCCCATCTGCAAGCAGGGTTGCCCGCCACAAAACAG TGGCCCTGGGTAGCAACGTGGAGTTCATGTGTAAGGTGTACAGTGACCCGCAGCCGC ACATCCAGTGGCTAAAGCACATCGAGGTGAATGGGAGCAAGATTGGCCAGACAACC TGCCTTATGTCCAGATCTTGAAGACTGCTGGAGTTAATACCACCGACAAAGATGG AGGTGCTTCACTTAAGAAATGTCTCCTTTGAGGACGCAGGGGAGTATACGTGCTTGG CGGGTAACCTCTATCGGACTCTCCCATCACTCTGCATGGTTGACCGTCTTGGAGGCC TGGAAGAGAGGCCCGCAGTGATGACCTCGCCCCGTGACCTGGAGATCATCATCTATT GCACAGGGGCCCTTCTCATCTCTGCATGGTGGGTCCGTCATCGTCTACAAGATGA AGAGTGGTACCAAGAAGAGTGACTTCCACAGCCAGATGGCTGTGCACAAGCTGGCCA AGAGCATCCCTCTGCGCAGACAGGTGTCTGCTGACTCCAGTGATCATGAACTCTG GGGTCTTCTGGTTCGGCCATCAGGGCTCTCCTCCAGTGGGACTCCCATGCTAGCAG GGGTCTCTGAGTATGAGCTTCCGAAAGACCTCGCTGGGAGCTGCCCGGGACAGAC TGGTCTTAGGCAAACCCCTGGGAGAGGGTGTCTTGGGCAGGTGGTGTGGCAGAGG CTATCGGGCTGGACAAGGACAAACCCAAACCGTGTGACCAAAGTGGCTGTGAAGATGT TGAAGTCGAGCCACACAGAGAAAGACTTGTGAGACTGATCTCAGAAATGGAGATGA TGAAGATGATCGGGAAGCATAGAATATCATCAACCTGCTGGGGGCTGCACGCAGG ATGGTCCCTTGATGTGTCATCGTGGAGTATGCCCTCAAGGGCAACCTGCGGGAGTACC TGCAGGCCCGAGGCCCGCCAGGGCTGGAATACTGCTACAACCCAGCCACAACCCAG AGGAGCAGCTCTCCTCCAAGGACCTGGTGTCTGCGCCTACAGGTGGCCCGAGGCA TGGAGTATCTGGCCTCCAAGAAGTGCATACACCGAGACCTGGCAGCCAGGAATGTCC TGGTGACAGAGGACAAATGTGATGAAGATAGCAGACTTGGCCTCGCACGGGACATTC ACCACATCGACTACTATAAAAAGACAACCAACGGCCGACTGCCCTGTGAAGTGGATGG CACCCGAGGCATTATTTGACCGGATCTACACCCACAGAGTGATGTGTGGTCTTTCG GGGTCTCTGTGGGAGATCTTCACTCTGGGCGGCTCCCCATACCCCGGTGTGCTCG TGGAGGAACCTTTCAAGCTGCTGAAGGAGGGTCACCGCATGGACAAGCCAGTAACT GCACCAACGAGCTGTACATGATGATGCGGGACTGCTGGCATGCAGTGCCCTCACAGA GACCCACCTTCAAGCAGCTGGTGGAGACCTGGACCGCATCGTGGCCTTGACCTCCA ACCAGGAGTACCTGGACCTGTCCATGCCCTGGACAGTACTCCCCCAGCTTTCCCG ACACCCGAGGCTCTACGTGCTCTCAGGGGAGGATTCCGCTCTTCTCTCATGAGCCGC TGCCCGAGGAGCCCTGCCTGCCCGACACCCAGCCAGCTTGCCAAATGGCGGATCA AACGCCGCTGACTGCCACCCACACGCCCTCCCCAGACTCCACCGTCAGCTGTAAACC TCACCCACAGCCCTGCTGGGCCACACACCTGTCCGTCCCTGTCCCTTTCTGTGCTG GCAGGAGCCGCTGCCTACCAGGGGCTTCTGTGTGGCCTGCCTTCAACCCACTCA GCTCACCTCTCCCTCCACCTCCTCTCCACCTGTGTTGAGAGGTGCAAGAGGCGAGA TCTTTGTGTCAGCCACTTATCCCCCTCCAGATGTTGGACCAACACCCCTCCCTGC CACAGGCACTGCTCGAGGGCAGGGAGTGGAGCCCAATGAACAGGCATGCAAGTGA GAGCTTCTGAGCTTTCTCTGTGCTGGTTGGTCTGTTTGCCTTACCCATAAGCCC CTCGCACTCTGGTGGCAGGTGCCTTGTCTCAGGGCTACAGCAGTAGGGAGGTGAGT GCTTCGTGCTCGATTGAAGGTGACCTCTGCCCCAGATAGTGGTGGCAGTGGCTTA TTAATTCGATACTAGTTTGTCTTGTGACCAATGCCTGGTACCAGAGGATGGTGA GGCGAAGGCCAGGTTGGGGCAGTGTGTGGCCCTGGGGCCAGCCCCAAACTGGGG GCTCTGTATATAGCTATGAAGAAAACACAAAGTGTATAAATCTGAGTATATATTAC ATGTCTTTTAAAGGGTCGTTACAGAGATTTACCATCGGTAAGATGCTCCTGG TGGCTGGGAGGCATCAGTTGCTATATATATAAAACAAAAAGAAAAAGGAAAAAT GTTTTTAAAGGTCATATATTTTTTGTACTTTTGTCTGTTTATTTTTTAAATTA TGTTCTAAACCTATTTTCAGTTTAGGTCCCTCAATAAAATTTGCTGCTGCTTC
	isoform 5 Amino acid sequence	
SEQ ID NO: 11	FGFR1	MAAVTRDFGEMLLHSGRVLPAAEQPWGAPVEVESFLVHPGDLQLRLRDDVQSIN WLRDGVQLAESNRTRITGEEVEVQDSVPADSGLYACVTSPPSGSDTTYFSVNVSDAL PSSDDDDDDSSSEKETDNTKPNRMPVAPYWTSPKMEKKLHAPPAKTVKFKCP SSGTPNPTLRWLKNGKEFKPDHRIGGYKVRATWSIIMDSVVPDCKGNYTCIVENEY GSINHTYQLDVVERSPhRPILQAGLPANKTVALGSNVEFMCKVYSDPQPHIQWLKHI EVNGSKIGPNDLPVYQILKTAGVNTTDKEMEVLHLRNVSFEDAGEYTCLAGNSIGLS HHSAWLTVLEALERPAVMTSPYLEII IYCTGAFLISCMVGSVIVYKMSGTKKSD FHSQMAVHKLAKSIPLRRQVSADSSASMNSGVLLVRPSRLSSSGTPMLAGVSEYELP EDPRWELPRDLRLVLGKPLGEGCFGQVVLAEAIGLDKDPNVRVTKVAVKMLKSDATEK LDSLISEMEMMKMI GKHKNI IINLLACTQDGPLYVIVEYASKNLREYLQARRPPG LEYCYNPSHNFEEQLSSKDLVSCAYQVARGMEYLASKKCIHRDLAARNVLVTEDNVM KIADFLGARDIHIDYKKTTNGLRFPVKWMAPEALFDRIYTHQSDVWSFGVLLWEIF TLGSPYPGPVPEELFKLLKEGHRMDKPSNCTNELYNNMRDCWHAVPSQRPTFKQLV EDLDRIVALTSNQEYLDLSMLDQYSPSPDTRSSSTCSSGEDSVFSHEPLPEEPCLP RHPAQLANGGLKRR
	isoform 6 Nucleic acid sequence	
		CTTGCGGCCACAGGCGCGGCGTCTCGGCGGCGGGCGGCAGCTAGCGGGAGCCGGGA CGCCCGGTGCAGCCGACGCGCGGAGGAACCGGGTGTGCCGGGAGCTGGGCGGCCA CGTCCGGACGGGACCGAGACCCCTCGTAGCGATTTGCGGCGACCTCGCCTTCCCGG CCGCGAGCGCGCCGCTGCTTGAAGCCCGCGGAACCCAAGGACTTTTCTCCGGTCCG AGCTCGGGGCGCCCGCAGGGCGCACGGTACCCGTGCTGCAGTCGGGCACGCCGCGG CGCCGGGCGCTCCGAGGGCGATGGAGCCGGTCTGCAAGGAAGTAGGCGCGCCGCG GCTCGGTCTGGAGGAGGGGGGCAAGGTCTGGAGACCCCGGTGGCGGACGGGAG

TABLE 5-continued

FGFR sequences.		
SEQ ID	FGFR1	MWSWKCLLFWAVLVLTATLCTARPSPTLPEQDALPSSSEDDDDDDSSSEKETDNTKP
NO: 12	isoform 6	NRMPVAPYWTSPEKMEKKLHAVPAKTVKPKPSSSGTPNPTRLRWLKNKGEFKPDHRI
	Amino acid	GGYKVRVATWSIIMDSVVPDKNYTCIVENEYGSINHLYQLDVVERSPHRPLQAG
	sequence	LPANKTVALGSNVFEMCKVYSDPQPHIQWLKHIEVNGSKIGPDNLPYVQILKTAGVN
		TTDKEMEVLHLRNVSFEDAGEYTCLAGNSIGLSHHSAWLTVLEALERPAVMTSPLY
		LEIIYCTGAFLISCMVGSVIVYKMKSGTKKSDPHSQMAVHKLAKSIPLRRQVTVSA
		DSSASMSGVLLVRPSRLSSSGTPMLAGVSEYELPEDPRWELPRDLVLGKPLGEGC
		FGQVVLAEIIGLDKDKPNRVTKVAVKMLKSDATEKDLSDLI SEMEMMKMIGKHKNII
		NLLGACTQDGLFYVIVEYASKNLREYLQARRPPGLEYYCNPNSHPNPEQLSSKDLVS
		CAYQVARGMEYLASKKCIHRDLAARNVLVTEEDNVMKIADFLGLARDIHHIDYKKTTN
		GRLPVKWMAPEALFDRIYTHQSDVWSFGVLLWEIFTLGGSPYPGVPEELFKLLKEG
		HRMDKPSNCTNELYMMMRDCWHAVPSQRPTFKQLVEDLDRI VALTSNQEYLDLSMPL
		DQYSPSPDTRSSSTCSSGEDSVFSHEPLPEEPCLPRHPAQLANGGLKRR

TABLE 5-continued

FGFR sequences.		
SEQ ID NO: 13	FGFR1 isoform 7 Nucleic acid sequence	CTTGCGGCCACAGGCGCGCGTCTCTGGCGGGCGGGCGGAGCTAGCGGGAGCCGGGA CGCCGGTGCAGCCGCGAGCGCGCGGAGGAACCCGGGTGTGCCGGGAGTGGGCGGCCA CGTCCGGACGGGACCGAGACCCCTCGTAGCGCATTTGCGGCGACCTCGCTTCCCCGG CCGCGAGCGCGCCGCTGCTTGAAAAGCCGCGGAACCCAGGACTTTTCTCCGGTCCG AGCTCGGGGCGCCCCGAGGGCGCACGGTACCCGTGCTGCAGTCGGGCACGCGCGG CGCCGGGGCTCCGCGAGGGGATGGAGCCCGGTCTGCAAGGAAAGTGAAGCGCCGCC GCTGCGTTCTGGAGGAGGGGGGCACAAGGTCTGGAGACCCCGGGTGGCGGACGGGAG CCCTCCCCCGCCCCGCTCCGGGGCACAGCTCCGGTCCATTGTTCCCGCCCGG CTGGAGCGCCGAGCACCGAGCGCCCGGGAGTCGAGCGCCGGCCGCGAGCTCTT GCGACCCCGCCAGACCCGAACAGAGCCCGGGGGCGGGCGGAGCCGGGAGCGCGGAGCG GGGCACACGCCCGCTCGCACAAGCCACGCGGACTCTCCGAGGCGGAACCTCCACG CCGAGCGAGGGTCAGTTTGAAAAGGAGGATCGAGCTCACTGTGGAGTATCCATGGAG ATGTGGAGCCTGTGCACCAACCTTAAGTGCAGAACTGGGATGTGGAGCTGGAAGTG CCTCTCTTCTGGGCTGTGCTGGTCAAGCCACACTCTGCACCGCTAGGCCGCTCCC GACCTTGCTGAACAAGATGCTCTCCCTCTCTCGAGGATGATGATGATGATGATGA CTCTCTTTCAGAGGAGAAAGAAACAGATAACACCAAAACAAACCCCTAGCTCCATA TTGGACATCCCGAAAAGATGGAAGAAATTCATGTCAGTGCCGGCTGCCAAGAC AGTGAAGTTCAAATGCCCTTCCAGTGGGACCCAAACCCCACTGCGCTGGTTGAA AAATGGCAAAGAAATCAAACCTGACCACAGAATGGAGGCTACAAGTCCGTTATGC CACCTGGAGCATCATAATGGACTCTGTGGTGCCTCTGCAAGGGCAACTACACCTG CATTGTGGAGATGAGTACGGCAGCATCAACCCACATACAGCTGGATGCTCGTGA GCGGTCCCTTACCGGCCATCTGCAAGCAGGTTGCCCGCAACAAACAGTGGC CCTGGGTAGCAACGTGGAGTTCATGTGAAGGTGTACAGTGACCCGAGCGCGACAT CCAGTGGCTAAAGCACATCGAGGTGAATGGAGCAAGATTGGCCGACACAACCTGCC TTATGTCCAGATCTGAAGACTGCTGGAGTTAATACCCGACAAAGAGATGGAGGT GCTTCACTTAAGAAATGTCTCTTTGAGGACGCGGGGAGTATACGTGCTTGGCGGG TAACCTTATCGGACTCTCCATCACTCTGCATGGTTGACCGTTCTGGAAGCCCTGGA AGAGAGGCCGGCAGTGATGACCTCGCCCTGTACCTGGAGATCATCATCTATTGCAC AGGGGCTTCTCTCATCTCCTGCATGGTGGGTCGGTTCATCGCTACAAGATGAAGAG TGGTACCAAGAAGAGTGAATCCACAGCCAGATGGTGTGCACAAGCTGGCCAAAGAG CATCCCTCTGCGCAGACAGGTAACAGTGTCTGTGACTCCAGTGATCCATGAATC TGGGTTTCTTCTGGTTCGGCCATCACGGCTCTCTCCAGTGGGACTCCCATGCTAGC AGGGGTCTCTGAGTATGAGCTTCCCGAAGACCTCGCTGGGAGCTGCCTCGGGACAG ACTGGTCTTAGGCAAAACCCCTGGGAGAGGGCTGCTTTGGGCGAGTGGTGTGGCAGA GGCTATCGGGCTGGACAAGGACAAACCCAAACCGTGTGACCAAGTGGCTGTGAAGAT GTTGAAGTTCGACGCAACAGAGAAAGACTTGTGAGACTGATCTCAGAAATGGAGAT GATGAAGATGATCGGGAAGCATAAGAATATCATCAACCTGCTGGGGCTGCACGCA GGATGGTCCCTGTATGTATCATCTGGAGTATGCTTCAAGGGCAACCTGCGGGAGTA CCTGCAGGCCCGGAGGCCCGCAGGGCTGGAATACTGCTACAACCCAGCCACAACCC AGAGGAGCAGCTCTCTCCAAAGGACCTGGTGTCTGCGCCTACAGGTGGCCCGAGG CATGGAGTATCTGGCTTCAAGAGTGCATACCCGAGACTGGCAGCCAGGAATGT CCTGGTGACAGAGGACAATGTATGAAGATAGCAGACTTTGGCTCTGCACGGGACAT TCACCACATCGACTACTATAAAAGACAACCAACGGCCGACTGCCTGTGAAGTGGAT GGCACCCGAGGCATTATTTGACCGGATCTACACCCAGAGTGAATGTGTGGCTTT CGGGGTCTCTGTGGGAGATCTTCACTCTGGGCGGCTCCCATACCCCGGTGTGCC TGTGGAGGAACCTTTCAAGCTGCTGAAGGAGGGTCAACCGCATGGACAAGCCAGTAA CTGCACCAACGAGCTGTACATGATGATGCGGGACTGCTGGCATGAGTGCCTCACA GAGACCCACCTTCAAGCAGCTGGTGAAGACCTGGACCGCATCGTGGCTTGACCTC CAACCAGGAGTACCTGGACCTGTCCATGCCCCCTGGACCACTCTCCCGAGCTTTCC CGACACCCGAGGCTTACGTGCTCTCAGGGGAGGATTCGCTCTTCTCTCATGAGCC GCTGCCCGAGGAGCCCTGCTGCCCCGACACCCAGCCAGCTTGCCAATGGCGGACT CAAACGCGCTGACTGCCACCCACACGCCCTCCCGAGACTCCACCGTCAGCTGTAAC CTCACCCACAGCCCTGCTGGGCCCCACACCTGTCGTCCTGTCCTCTTCTCTGCTG TGGCAGGAGCCGGCTGCTACAGGGGCTTCTGTGTGGCTGCTTCAACCCACT CAGCTCACCTCTCCCTCCACCTCTCTCCACCTGCTGGTGGAGGTGCAAGAGGCA GATCTTTGCTGCCAGCCACTTCATCCCTCCAGATGTTGGACCAACACCCCTCCCT GCCACAGGCACTGCTGGAGGGCAGGGAGTGGGAGCAATGAACAGGCATGCAAGT GAGAGCTTCTGAGCTTTCTCTCTGTCGGTGTGGTCTGTTTTGCTTCAACCATAGC CCCTCGCACTCTGGTGGCAGGTGCTTGTCTCAGGGCTACAGCAGTAGGGAGGTCA GTGCTTCTGCTCTGATTGAAGGTGACCTCTGCCCCAGATAGGTGGTGGCAGTGGCT TATTAATTCGATACTAGTTTGTCTTGTGACCAATGCTGGTACAGAGGATGGT GAGGCGAAGGCACAGTTGGGGGAGTGTGTGGCCCTGGGGCCAGCCCAAACTGG GGGCTCTGTATATAGCTATGAAGAAAACAAAGTGTATAAATCTGAGTATATATTT ACATGTCTTTTAAAAGGGTCGTTACAGAGATTTACCCATCGGGTAAGATGCTCCT GGTGGCTGGGAGGCATCAGTTGCTATATATATAAAACAAAAAGAAAAAGGAAA ATGTTTTTAAAAGGTATATATTTTTTGCTACTTTTGTGTTTTATTTTTTAAAT TATGTTCTAAACCTATTTTTCAGTTTAGGTCCCTCAATAAAATTTGCTGCTGCTT
SEQ ID NO: 14	FGFR1 isoform 7 Amino acid sequence	MWSWKCLLFWAVLVTATLCTARPSPTLPEQDALPSSSEDDDDDDSSSEKETDNTPK NPVAPYWTSPEKMEKKLHAVPAKTKVFKCPSSGTPNPTRLWLKNGEFPKPDHRIIG YKRYATWSIIMDSVVPDSKGNITCI VENEYGSINHTYQLDVVERSHPRLILQAGLP ANKTVALGSNVEFMCKVYSDPQPHIQWLKHIEVNGSKI GPDNLPYVQILKTAGVNTT DKEMEVLHLRNVSFEDAGEYTC LAGNSIGLSHSAWLTVLEALEERPAMVTSPLYLE II IYCTGAFLISCMVGSVIVYKMSGTTKSDPHSQMAVHKLAKSIP LRRQVTVSADS SASMNSGVLLVRPSRLSSSGTPMLAGVSEYELPEDPRWELPRDLRLVGLKPLGEGCFG QVVLAEAI GLDKDPNRVTKVAVKMLKSDATEKDLSDLI SEMEMMKMIGKHKNI INL LGACTQDGPLYVIVEYASKGNLREYLQARRPPGLECYNPSHNPEQLSSKDLVSCA YQVARGMEYLASKKCIHRDLAARNVLTEDNVMKIADFGDLARDIHHIDYKKTNTNGR

TABLE 5-continued

FGFR sequences.		
SEQ ID	FGFR1	LPVKWMAPEALFDRIYTHQSDVWSFGVLLWEIFTLGGSPYPGVPVEELFKLLKEGHR
NO: 15	isoform 8	MDKPSNCTNELYMMMRDCWHAVPSQRPTFKQLVEDLDRI VALTSNQEYLDLSMPLDQ
	Nucleic acid	YSPSPFDTRSSTCSSGEDSVFSHEPLPEEPCLPRHPAQLANGGLKRR
	sequence	
		GCTTTGCCCGCCGACGCCAGCCGGGGCCGGCGCCTCCCTCCGCTCGCCGCCGCC
		CTTTCACCTCCTGGCTCCCTCCCGGGCGATCCGCGCCCCCTGGGTCTCCCTCCCTT
		CCCTCCGTCGCGCTCTCCTGCGCCCCCTCCCTGCGCTCGTCCCGCCGCTCTTCCCGC
		CGCCCAACTTTCTCTCAACTCGCGCTCGGGAGCTGGCGAGGCGGGCGGCTCCTC
		AGGTCAAGTTGAAAAGGAGGATCGAGCTCACTGTGGAGTATCCATGGAGATGTGGAG
		CCTTGTACCAACCTCTAAGTGCAGAACTGGGATGTGGAGCTGGAAGTGCCTCCTCT
		TCTGGGCTGTGCTGGTCACAGCCACACTCTGCACCGCTAGGCCGTCCTCCGACCTTG
		CTGAACAAGCCAGCCCTGGGGAGCCCTGTGGAAGTGGAGTCTTCTTGGTCCACC
		CCGGTGACCTGTGTCAGCTTCTGTGTGGGTGCGGGACGATGTGCAGAGCATCAACT
		GGCTGCGGGACGGGGTGCAGCTGGCGGAAAGCAACCGCACCCGCATCACAGGGGAGG
		AGGTGGAGGTGCAGGACTCCGTGCCCGCAGACTCCGGCCTCTATGCTTGCCTAACA
		GCAGCCCTCGGGCAGTGACACCACCTACTTCTCCGTCAATGTTTCAGATGCTCTCC
		CCTCTCGGAGGATGATGATGATGATGATGACTCCTCTTCAGAGGAGAAAGAACAG
		ATAACACCAAAACCAACCCCGTAGCTCCATATTGGACATCCCGAGAAAGATGGAAA
		AGAAATGTCATGCAGTGCCGGCTGCCAAGACAGTGAAGTTCAAATGCCCTTCCAGTG
		GGACCCCAACCCACACTGCGCTGGTTGAAAATGGCAAAGAAATCAAACCTGACC
		ACAGAAATGGAGGTCACAAAGTCCGTTATGCCACCTGGAGCATCATATGGACTCTG
		TGGTGCCCTCTGACAAGGGCAACTACACCTGCATTGTGGAGAATGAGTACGGCAGCA
		TCAACCACACATACCAGCTGGATGCTGTGGAGCGGTCCCTCACCGGCCCATCTGCG
		AAGCAGGGTTGCCCGCAACAAACAGTGGCCCTGGGTAGCAACGTGGAGTTCATGT
		GTAAGGTGTACAGTGACCCGACGCGCACATCCAGTGGCTAAAGCACATCGAGGTGA
		ATGGGAGCAAGATTGGCCAGACAACCTGCCTTATGTCCAGATCTTGAAGACTGCTG
		GAGTTAATACACCGACAAAGAGATGGAGGTGCTTCACTTAAGAAATGTCTCCTTG
		AGGACGCGGGGAGTATACGTGCTTGGCGGGTAACCTATCCGAGCTCTCCATCACT
		CTGCATGGTTGACCGTTCTGGAAGCCCTGGAAGAGAGGCCGGCAGTATGACCTCGC
		CCCTGTACCTGGAGATCATCATCTATTGCACAGGGGCCCTTCTCATCTCTGCATGG
		TGGGGTCCGTCTCGTCTACAAGATGAAGAGTGGTACCAAGAAGAGTGACTTCCACA
		GCCAGATGGCTGTGCACAAGCTGGCCAAGAGCATCCCTCTGCGCAGACAGGTAAACG
		TGCTGTGACTCCAGTGCATCCATGAACCTCTGGGGTCTTCTGGTTCGGCCATCAC
		GGCTCTCCTCCAGTGGGACTCCCATGTAGCAGGGGTCTCTGAGTATGAGCTTCCCG
		AAGACCTCGCTGGGAGCTGCCCTGGGACAGACTGGTCTTAGGCAAAACCCCTGGGAG
		AGGGCTGCTTTGGGCAGGTGGTGTGGCAGAGGCTATCGGGCTGGACAAGGACAAAC
		CCAACCGTGTGACCAAAGTGGCTGTGAAGATGTTGAAGTCGGACGCAACAGAGAAAG
		ACTTGTGACACCTGATCTCAGAAATGGAGATGATGAAGATGATCGGGAAGCATAAGA
		ATATCATCAACCTGCTGGGGGCTGCACGACAGGATGGTCCCTTGTATGTCTCGTGG
		AGTATGCCTCCAAGGGCAACCTGCGGGAGTACCTGCAGGCCCCGAGGCCCCAGGGC
		TGGAATACTGCTACAACCCAGCCACAAACCCAGAGGAGCAGCTCTCCTCCAAGGACC
		TGGTGTCTGCGCCTACAGGTGGCCCGAGGCATGGAGTATCTGGCCTCCAAGAGT
		GCATACACCGAGACCTGGCAGCCAGGAATGTCTGGTGACAGAGGCAATGTGATGA
		AGATAGCAGACTTTGGCCTCGCACGGGACATTCAACCATCGACTACTATAAAAAGA
		CAACCAACGGCCGACTGCCTGTGAAGTGGATGGCACCAGGAGCATTATTTGACCGGA
		TCTACACCCACAGAGTATGTGTGGTCTTTCCGGGTGCTCTGTGGGAGATCTTCA
		CTCTGGGCGGCTCCCATACCCCGGTGTGCTGTGGAGGAACCTTTCAAGCTGCTGA
		AGGAGGGTACCCGATGGACAAGCCAGTAACCTGCACCAACGAGCTGTACATGATGA
		TGCGGGACTGCTGGCATGCAGTGCCCTCACAGAGACCCACCTTCAAGCAGCTGGTGG
		AAGACCTGGACCGCATCGTGGCCTTGACCTCCAACAGGAGTACCTGGACCTGTCCA
		TGCCCCGAGACAGTACTCCCCAGCTTTCCCGACACCCGGAGCTCTACGTGCTCCT
		CAGGGGAGGATTCGCTCTTCTCATGAGCCGCTGCCCGAGGAGCCTGCCTGCCCC
		GACACCCAGCCAGCTTGCCAATGGCGGACTCAAACGCCGCTGACTGCCACCCACAC
		GCCCTCCCCAGACTCCACCGTCAGCTGTAACCTCACCCACAGCCCTGCTGGGCCC
		ACCACCTGTCCGTCCTGTCCCTTTCTGTGCTGGCAGGAGCCGGCTGCCACAGGG
		GCCTTCTGTGTGGCTGGCTTCAACCCACTCAGCTCACCTCTCCCTCCACCTCCTC
		TCCACCTGTGTGGTGGAGGTGCAAGAGGCGAGATCTTGTGTCAGCCACTTCATCC
		CCTCCCAGATGTTGGACCAACACCCCTCCCTGCCACCCAGGCACTGCCTGGAGGGCAG
		GGAGTGGGAGCCAATGAACAGGCATGCAAGTGAAGCTTCTGAGCTTTCTCCTGTC
		GGTTTGGTCTGTTTTGCCTTCAACCATAAAGCCCTCGCACTCTGGTGGCAGGTGCCT
		TGTCCTCAGGGCTACAGCAGTAGGGAGGTGAGTCTTCTGTCCTCGATTGAAGGTGA
		CCTCTGCCCCAGATAGGTGGTGGCAGTGGCTTATTAATCCGATACTAGTTTGTCTT
		GCTGACCAAATGCTGTGACAGAGGATGGTGAGGCGAAGGCCAGGTTGGGGGAGT
		GTGTGTGGCCCTGGGGCCAGCCCCAACTGGGGGCTCTGTATATAGCTATGAAGAAA
		ACACAAAGTGTATAAATCTGAGTATATATTTACATGCTTTTAAAGGGTCGTTAC
		CAGAGATTTACCCATCGGGTAAGATGCTCCTGGTGGCTGGGAGGCATCAGTTGCTAT
		ATATTAACCAACAAAAAGAAAAAAGGAAATGTTTTAAAGAGTCATATATTTT
		TTGCTACTTTTGTGTTTTATTTTTTAAATATGTTCTAAACCTATTTTCAGTTTA
		GGTCCCTCAATAAAAAATGTGTGCTGCTTCATT
SEQ ID	FGFR1	MWSWKLLFWAVLVATLCTARPSPTLPEQAQPGWAPVEVESFLVHPGDLQLRCL
NO: 16	isoform 8	RDDVQSINWLRDGVQLAESNRTRI TGEVEVQDSVPADSLYACVTS SPSGSDTTYF
	Amino acid	SVNVSDALPSSDDDDDDSSSEKETDNTKPNVPVAPYWTSPEKMEKKLHAVPAAKT
	sequence	VKFKCPSSGTFNPTLRWLKNGKEFKPDHRIGGYKVRATWSIIMDSVVPDCKGNYTC
		IVENEYGSINHTYQLDVVERSPHRPILQAGLPANKTVALGNSVEFMCKVYSDPQPHI
		QWLKHIEVNGSKI GPDNL PYVQILKTAGVNTDKEMEVLHLRNVSFEDAGEYTCLAG
		NSIGLSHSAWLTVLEALEERPAMVTSPLYLEI I IYCTGAFLISCMVGSVIVYKMS
		GTKKSDPHSQMAVHKLAKSIPLRQVTVSADSSASMNSGVLLVRPSRLSSSGT PMLA

TABLE 5-continued

FGFR sequences.		
SEQ ID NO: 17	FGFR1 isoform 9 Nucleic acid sequence	GVSEYELPEDPRWELPRDLVLGKPLGEGCFGQVVLAEI GLDKDKPNRVTKVAVKM LKSDATEKDLSDLISEMEMMKMIGKHKNIIINLLGACTQDGPLYVIVEYASKGNLREY LQARRPPGLECYNPSHNPEEQLSKDLVSCAYQVARGMEYLASKKCIHRDLAARNV LVTEDNVMKIADPGLARDIHIIIDYYKKTNGRLPVKWMAPALFDRIYTHQSDVWSF GVLLWEIFTLGGSPYPGPVVEELFKLLKEGHRMDKPSNCTNELYMMMRDCWHAVPSQ RPTFKQLVEDLDRIVALTSNQEYLDLSMPLDQYSPSPDTRSSSTCSSGSDSVFSHEP LPEEPCLPRHPAQLANGGLKRR AACTTTTCTCCAAGTTCGCGCTCGGGAGCTGGCGAGGCGGCGGGCTCCTCAAAGT GGGAGAGCTTCAAGGTACGTGGTCCGTCCAGCCCCGTGCTATCTCACCAGACACTGT CCACCTGTATGTTGGATCAGTACTCCAGTGAGAAGACAGCAGGCACCTTTCACCCAT GCAGCCATTTCAGTCTTCATAACCACTGTGATGGAGGCAAGGGTCAAGTTGAAAAG GAGGATCGAGCTCACTGTGGAGTATCCATGGAGATGTGGAGCCTTGTCACCAACCTC TAAGTCAGAACTGGGATGTGGAGCTGGAAGTGCCCTCCTCTCTGGGCTGTGCTGGT CACAGCCACACTCTGCACCGCTAGGCCGTCCCCGACCTTGCCTGAAACAAGCCAGCC CTGGGGAGCCCCGTGGAAGTGGAGTCTTCTCGGTCCACCCCGGTGACCTGTGTGCA GCTTCGCTGTCCGGTGCAGGACGATGTGCAGAGCATCAACTGGCTGCAGGACGGGGT GCAGCTGGCGAAAGCAACCGCACCCGCATCACAGGGGAGGAGGTGGAGGTGCAGGA CTCCGTGCCCGCAGACTCCGGCTCTATGCTTGCGTAACAGCAGCCCCCTCGGGCAG TGACACCACTACTTCTCCGTCAATGTTTCAGATGCTCTCCCTCCTCGGAGGATGA TGATGATGATGATGACTCCTCTTCAGAGGAGAAAGAAACAGATAACCAACCAACAAA CCCCGTAGCTCCATATTGGACATCCCAGAAAAGATGGAAGAAAGAAATGCATGCAGT GCCGGCTGCCAAGACAGTGAAGTTCAAATGCCCTTCCAGTGGGACCCCAACCCAC ACTGCGCTGGTTGAAAATGGCAAGAATTCAAACCTGACCACAGAATTGGAGGCTA CAAGGTCGTTATGCCACCTGGAGCATCATAATGGACTCTGTGGTGGCCCTCTGACAA GGGCAACTACACCTGCATTGTGGAAGATGAGTACGGCAGCATCAACCACACATACCA GCTGGATGTGCTGGAGCGGTCCCCCTACCGGCCCATCTGCAAGCAGGGTGGCCCGC CAACAAAACAGTGGCCCTGGGTAGCAACCTGGAGTTCATGTGTAAGGTGTACAGTGA CCCGCAGCCGCACATCCAGTGGCTAAAGCACATCGAGGTGAATGGGAGCAAGATTGG CCCAGACAACCTGCCTTATGTCCAGATCTTGAAGACTGTCTGGAGTTAATACCACCGA CAAAGAGATGGAGGTGCTTCACTTAAGAAATGTCTCCTTTGAGGACGCAGGGGAGTA TACGTGCTTGGCGGGTAACTCTATCGGACTCTCCATCACTCTGCATGGTTGACCGT TCTGGAAGCCCTGGAAGAGAGGCCCGGAGTGTGACCTCGCCCTGTACCTGGAGAT CATCATCTATTGCACAGGGGCCCTTCTCATCTCCTGCATGGTGGGTGCGTCACTGT CTACAAGATGAAGAGTGGTACCAAGAAGAGTGAAGTTCACAGCCAGATGGCTGTGCA CAAGCTGGCCAGAGCATCCCTCTGCGCAGACAGGTAAACAGTGTCTGCTGACTCCAG TGCATCCATGAACCTCTGGGGTCTTCTGTGTTGGCCATCACGGCTCTCTCCAGTGG GACTCCCATGCTAGCAGGGGTCTCTGAGTATGAGCTTCCCGAAGACCTCGCTGGGA GCTGCCCTCGGACAGACTGGTCTTAGGCAAAACCCCTGGGAGAGGGCTGCTTTGGGCA GGTGGTGTGGCAGAGGCTATCGGGCTGGACAAGGACAAACCAACCGTGTGACCAA AGTGGCTGTGAAGATGTTGAAGTCGGACGCAACAGAGAAAGACTTGTGACAGCTGAT CTCAGAAATGGAGATGATGAAGATGATCGGGAAGCATAAGAAATATCATCAACCTGCT GGGGGCTGTCACGCAGGATGGTCCCTTGATGTGTCATCGTGGAGTATGCCTCCAAGGG CAACCTGCGGGAGTACCTGCAGGCCCGGAGGCCCCAGGGCTGGAATACTGCTACAA CCCCAGCCACAACCCAGAGGAGCAGCTCTCTCCAAGGACCTGGTGTCTGCGCCTA CCAGGTGGCCCGAGGCATGGAGTATCTGGCCCTCAAGAAGTGCATACCCGAGACCT GGCAGCCAGGAATGTCTGTGACAGAGGACAAATGTGATGAAGATAGCAGACTTTGG CCTCGCACGGGACATTACCCACATCGACTACTATAAAAAGACAACCAACGGCCGACT GCCTGTGAAGTGGATGGCACCCGAGGCATTATTGACCGGATCTACACCCACACAGAG TGATGTGTGGTCTTTTGGGGTGTCTCTGTGGGAGATCTTCACTCTGGGCGGCTCCCC ATACCCCGGTGTGCTCTGTGGAGGAACTTTCAAGCTGCTGAAGGAGGGTCAACGCAT GGACAAGCCAGTAAGTGCACCAACGAGCTGTACATGATGATGCGGGACTGTCTGGCA TGCAGTGCCCTCAGAGAGACCACTTCAAGCAGCTGGTGAAGACCTGGACCGCAT CGTGGCCTTGACCTCCAACAGGAGTACCTGGACCTGTCCATGCCCTGGACCAT CTCCCCAGCTTTCCCGACACCCGAGCTCTACGTGCTCTCAGGGGAGGATTCCGT CTTCTCTCATGAGCCGCTGCCCGAGGAGCCCTGCTGCCCGACACCCAGCCAGCT TGCCAATGGCGGACTCAAACGCGCTGACTGCCACCCACACGCCCCCTCCCAGACTCC ACCGTCAGTGTAAACCTCACCCACAGCCCTGTGGGCCACACACCTGTCCGTCC TGTCCTCTTCTGCTGGCA
SEQ ID NO: 18	FGFR1 isoform 9 Amino acid sequence	MWSWKCLLFWAVLVTATLCTARPSPTLPEQAQPWGAPVEVESFLVHPGDLQLRCRL RDDVQSINWLRDGVQLAESNRTRITGEEVEVQDSVPADSGLYACVTSSPSGSDTTYF SVNVSDALPSSSEDDDDDDSSSEKETDNTKPNVPAPYWTSPKMKKLLHAVPAKT VKFKCPSSGTPNPTLRWLKNGKEFKPDHRIIGYKVRATWSIIMDSVVPDKGNVTC IVENEYGSINHTYQLDVVERSPHRPIIQAQGLPANKTVALGSNVEFMCKVYSDPQPHI QWLKHI EVNGSKI GPDNLPIVQILKTAGVNTTDKEMEVLHLRNVSFEDAGEYTCLAG NSIGLSHSAWLTVLEALEERPAVMTSPLEYLIIYCTGAFLISCMVGSVIVYKMS GTKKSDPHSQMAVHKLAKSIPLRRQVTVSADSSASMSGVLLVLRPSRLSSSGTPMLA GVSEYELPEDPRWELPRDLVLGKPLGEGCFGQVVLAEI GLDKDKPNRVTKVAVKM LKSDATEKDLSDLISEMEMMKMIGKHKNIIINLLGACTQDGPLYVIVEYASKGNLREY LQARRPPGLECYNPSHNPEEQLSKDLVSCAYQVARGMEYLASKKCIHRDLAARNV LVTEDNVMKIADPGLARDIHIIIDYYKKTNGRLPVKWMAPALFDRIYTHQSDVWSF GVLLWEIFTLGGSPYPGPVVEELFKLLKEGHRMDKPSNCTNELYMMMRDCWHAVPSQ RPTFKQLVEDLDRIVALTSNQEYLDLSMPLDQYSPSPDTRSSSTCSSGSDSVFSHEP LPEEPCLPRHPAQLANGGLKRR

TABLE 5-continued

FGFR sequences.		
SEQ ID NO: 19	FGFR2 isoform 1 Nucleic acid sequence	GGCGGCGGCTGGAGGAGAGCGCGGTGGAGAGCCGAGCGGGCGGGCGGGTGC GGA GCGGGCGAGGAGCGCGCGCGGCCGCCACAAAGCTCGGGCGCGCGGGGTGCATGC GGCGTACCTGGCCCGCGCGCGGCTGCTCTCCGGGCTGGCGGGGGCCGCCGCGAG CCCCGGGGGCCCCGAGGCCGCGAGCTTGCCTGCGCGCTCTGAGCCTTCGCAACTCGCG AGCAAAAGTTTGGTGGAGGCAACGCCAAGCCTGAGTCCTTTCTCTCTCGTTCCCA AATCCGAGGGCAGCCCGCGGGGCTCATGCCCGCGCTCCTCCGAGCCTGGGGTACGC GTGAAGCCCCGGGAGGCTTGGCGCGCGGCGAAGACCCAGGACCACTCTTGCCTTTG GAGTTGCTCCCCGCAACCCCGGGCTCGTCGCTTCTCCATCCGAGCCACGCGGGG GCGGGGACACACAGGTGCGGAGGAGCGTTGCCATTCAAGTGACTGCAGCAGCAGC GGCAGCGCCTCGGTTCTGAGCCACCGCAGGCTGAAGGCATTGCGCGTAGTCCATG CCCGTAGAGGAAGTGTGCAGATGGGATTAACGTCCACATGGAGATATGGAAGAGGAC CGGGGATTGGTACCGTAACCATGGTCAGCTGGGGTCTTTTCATCTGCCTGGTCTG TCACCATGGCAACCTTGTCCTCGGCCCGGCCCTCTTCAAGTTAGTTGAGGATACCA CATTAGAGCCAGAAGATGCCATCTCATCCGAGATGATGAGGATGACACCGATGGTG CGGAAGATTGTGTAGTGAAGACAGTAACCAAGAGAGACCATATCGGACCAACA CAGAAAAGATGGAAGAGCGGCTCCATGCTGTGCTCGGGCCAACTGTCAAGTTTC GCTGCCAGCCGGGGGAACCAATGCCAACCATGCGGTGGCTGAAAAACGGGAAGG AGTTTAAGCAGGAGCATCGCATTGGAGGCTACAAGGTACGAAACCGACACTGGAGCC TCATTATGGAAGTGTGGTCCCCTCTGACAAGGGAATTATACCTGTGTAGTGGAGA ATGAATACGGTCCATCAATCACAGTACCACCTGGATGTTGTGAGCGATCGCCTC ACCGGCCCATCTCCAAGCCGAGCTGCCGCAATGCCTCCACAGTGGTGGAGGAG ACGTAGAGTTTGTCTGCAAGGTTTACAGTGATGCCAGCCCCACATCCAGTGGATCA AGCACGTGGAAGAAACGGCAGTAATAACGGGCCCGAGGGCTGCCCTACCTCAAGG TTCTCAAGGCCGCGCGTGTAAACACCACGACAAAGAGATTGAGGTTCTCTATATTC GGAATGTAACTTTGGAGGACGCTGGGGAATATACGTGCTTGGCGGTAATTTCTATTG GGATATCTTTCACTCTGCATGGTTGACAGTTCTGCCAGCGCTGGAGAGAGAAAGG AGATTACAGCTTCCCAGACTACCTGGAGATAGCCATTTACTGCATAGGGGTCTTCT TAATCGCCTGTATGGTGGTAACAGTCATCTGTGCCGAATGAAGAACACGACCAAGA AGCCAGACTTCAGCAGCCAGCCGGCTGTGCACAGCTGACCAACGTATCCCCCTGC GGAGACAGGTAACAGTTTCGGCTGAGTCCAGCTCCTCCATGAACCTCCAACCCCCGC TGGTGAGGATAACAACACGCTCTCTTCAACGGCAGACACCCCCATGCTGGCAGGGG TCTCCAGTATGAACCTCCAGAGGACCAAAATGGGAGTTTCCAAGAGATAAGCTGA CACTGGGCAAGCCCTGGGAGAAGGTTGCTTGGGCAAGTGGTCATGGCGAAGCAG TGGGAATTGACAAGACAAGCCCAAGGAGGCGGTCACCGTGGCCGTGAAGATGTTGA AAGATGATGCCACAGAGAAAGACCTTTCTGATCTGGTGTGAGAGATGGAGATGATGA AGATGATTGGGAAACACAAGAAATATCATAAATCTTCTTGGAGCTGCACACAGGATG GGCCTCTCTATGTATAGTTGAGTATGCCCTCTAAAGGCAACCTCCGAGAATACCTCC GAGCCCGGAGGCCACCCGGGATGGAGTACTCCTATGACATTAACCGTGTTCTCGAGG AGCAGATGACCTTCAAGGACTTGGTGTATGCACCTACCAGCTGGCCAGAGGCATGG AGTACTTGGCTTCCCCAAATGTATTATCTGAGATTAGCAGCCAGAAATGTTTGG TAACAGAAACAAATGTGATGAAATAGCAGACTTTGAGCTCGCCAGAGATATCAACA ATATAGACTATTACAAAAGACCACCAATGGGCGGCTTCCAGTCAAGTGGATGGCTC CAGAAGCCCTGTTTGATAGAGTATACACTCATCAGAGTGTCTGGTCTCTCGGGG TGTTAATGTGGGAGATCTTCACTTATGGGGGCTCGCCCTACCCAGGATTCCTCGTG AGGAACCTTTTAAGCTGCTGAAGGAAGGACACAGAATGGATAAGCCAGCCAACTGCA CCAACGAACTGTACATGATGATGAGGGACTGTTGGCATGCAGTGCCCTCCAGAGAC CAACGTTCAAGCAGTTGGTAGAAGACTTGGATCGAATTTCACTCTCACAAACATG AGGAATACTTGGACCTCAGCCAACCTCTCGAACAGTATTACCTAGTTACCTTGACA CAAGAAGTTCTGTTCTTTCAGGAGATGATTCTGTTTTTCTCCAGACCCCATGCCTT ACGAACCATGCCTTCTCAGTATCCACACATAAACGGCAGTGTTAAACATGAAATGA CTGTGTCTGCTGTCCCCAACAGGACAGCACTGGGAACCTAGCTACACTGAGCAGG GAGACCATGCCTCCAGAGCTTGTGTCTCCACTTGTATATAGGATCAGAGGAGTA AATAATTGGAAGAAATAGTATCAGCATATGTGTAAGATTATACAGTTGAAAACCTGTA ATCTTCCCAGGAGGAGAAGAGGTTTCTGGAGCAGTGGACTGCCACAAGCCACCAT GTAACCCCTCTACCTGCCGTGCGTACTGGCTGTGGACCAGTAGGACTCAAGTGGA CGTGCCTTCTGCCCTTCTTGTAAATTTGTAATAATTGGAGAAGATTATGTGAGCA CACACTTACAGAGCACAAATGCAGTATATAGGTGCTGGATGTATGTAATATATTTCA AATTATGTATAAATATATATATATTTTACAAGGAGTTATTTTTTGTATGATTTT AAATGGATGTCCTCAATGCACCTAGAAAATTGGTCTCTCTTTTTTAAATAGCTATTTG CTAAATGCTGTCTTACACATAATTTCTAATTTTCAACGAGCAGAGGTGGAAGAAAT ACTTTTGCTTTTCAGGGAAGATGATATACGTTAATTTATTAATAAATTTGGTAATATA CAAAACAAATTAATCATTATAGTTTTTTTGTAAATTAAGTGGCATTTCTATGCAGG CAGCACAGCAGACTAGTTAATCTATTGCTTGGACTTAACAGTTATCAGATCCTTTG AAAAGAGAATATTACAAATATATGACTAATTTGGGGAAATGAAGTTTGTATTTATT TGTGTTAAATGCTGCTGTGACAGGATGTTCTTAGACCTCCTAAAGTCCCATATT AAAAGAACTCATTATAGGAAGGTGTTTCAATTTTGGTGTGCAACCCCTGTATTACGT CAACGCAACGCTTAACCTGAGCTTCCCAAGATAAATGGTACCAGCGTCTCTTAAAG ATGCTTAAATCCATTCTTGGAGCAGACCTTAGTTGAAATGATAGCAGAAATGTGCT TCTCTCTGGCAGCTGGCCTTCTGCTTCTGAGTTGCACATTAATCAGATTAGCCTGTA TTCTCTTCAAGTAAATTTGATAATGGCTTCCAGACTCTTTGGCGTTGGAGACGCTG TTAGGATCTCAAGTCCCATCATAGAAAATTGAAACACAGAGTTGTTCTGTCTGATAG TTTTGGGGATACGTCCATCTTTTAAGGGATTGCTTTCATCTAATTTCTGGCAGGACC TCACCAAAAGATCCAGCCTCATACCTACATCAGACAAAATATCGCCGTGTTCTCTTC TGTAATAAGTATTGTTGTTTTGCTTTGGAAACACCCACTCACTTTGCAATAGCCGTG CAAGATGAATGCAGATTACACTGATCTTATGTGTTACAAAATTTGGAGAAAGTATTTA ATAAACCTGTAAATTTTTTACTGACAATAAAAATGTTTCTACAGATATTAATGTT AACAGACAAAATAAGTGCACGCACTTATTTTTT

TABLE 5-continued

FGFR sequences.		
SEQ ID NO: 20	FGFR2 isoform 1 Amino acid sequence	MVSWGRFICLVVVTMATLSLARPSFSLVEDTTLEPEDAISSGDDEDDTDGAEDFVSE NSNNKRAPYWTNTEKMEKRLHAVPAANTVKFRCPAGGNPMPMTMRWLKNGKEFKQEH IGGYKVRNQHWSLIMESVVPDSDKGNVTCVVENEYGSINHTYHLVDVVERSPHRPILQA GLPANASTVVGGDVEFVKVYSDAQPHIQWIKHVEKNKSKYGPDGLPYLKVLKAAGV NTTDKEIEVLYIRNVTFEDAGEYTCLAGNSIGISFHSAWLTVLPAPGREKEITASPD YLEIAIYICIGVFLIACMVVTVILCRMKNNTTKPDFSSQPAVHKLTKRIPLRRQVTVS AESSSSMNSNTPLVRITRLSSTADTTPMLAGVSEYELPEDPKWEFPRDKLTGLKPLG EGCFGQVVMABEAVGIDKDKPKEAVTVAVKMLKDDATEKDLSDLVSEMEMMMKMGKH NI INLLGACTQDGLPYVIVEYASKGNLREYLRARRPPGMEYSYDINRVPEEQMTFKD LVSCYQLARGMEYLASQKCIHRDLAARNVLVTENNVMKIADPGLARDINNIDYKK TTNGRLPVKWMABEALFDRVYTHQSDVWSFGVLMWEIFTLGGSPYPGIPVEELFKLL KEGHRMDKPANCTNELYMMMRDCWHAVPSQRPTFKQLVEDLDRILTLTTNEEYLDLS QPLEQYSPSPYDTRSSSCSSGDSDVFSFDPMPYEPCLPQYPHINGSVKT
SEQ ID NO: 21	FGFR2 isoform 2 Nucleic acid sequence	CCCAAGGACCACTCTTCTGCGTTTGGAGTTGCTCCCGCAACCCCGGGCTCGTCGCT TTCTCCATCCCGACCCACGCGGGGCGCGGGGACAACACAGGTCGCGGAGGAGCGTTG CCATTCAAGTGACTGCAGCAGCAGCGGCGAGCGCTCGGTTCCCTGAGCCACCCGAGG CTGAAGGCATTGCGCGTAGTCCATGCCCGTAGAGGAAGTGTGCAGATGGGATTAACG TCCACATGGAGATATGGAAGAGGACCGGGGATTGGTACCCTAACCATGGTCAGCTGG GGTCGTTTCATCTGCTGGTCTGGTACCATGGCAACCTTGTCCCTGGCCCGGCCC TCCTTCAGTTTATGTTGAGGATACACATTAGAGCCAGAAGAGCCACCAACCAATAC CAATCTCTCAACCAGAAGTGACGTGGCTGCGCCAGGGGAGTCGCTAGAGGTGCGG TGCTGTGTAAGAGATGCGCGCGTGATCAGTTGGACTAAGGATGGGGTGCACTTGGGG CCCAACAATAGGACAGTGCTTATTGGGGAGTACTTGACAGATAAAGGGCGCCACGCCT AGAGACTCCGGCTCTATGCTTGTACTGCGCAGTAGGACTGTAGACAGTGAAACTTGG TACTTCATGGTGAATGTACAGATGCCATCTCATCCGAGATGATGAGGATGACACC GATGGTGGGGAAGATTTGTGTCAGTGAGAACAGTAACAACAAGAGAGCACCATACTGG ACCAACACAGAAAAGATGGAAGCGGCTCCATGCTGTGCTGCGGCCAACACTGTC AAGTTTCGCTGCGCCAGCGGGGGGAACCAATGCCAACCATGCGGTGGCTGAAAAAC GGGAAGGAGTTTAAGCAGGAGCATCGCATTGGAGGCTACAGGTACGAAACAGCAC TGGAGCCTCATTAAGGAAAGTGTGGTCCCCTGACAGGGAATATACCTGTGTA GTGGAGAATGAATACGGGTCCATCAATCACAGTACCACCTGGATGTTGTGGAGCGA TCGCCTCACCAGCCATCCTCCAGCGGACTGCGGCAAAATGCCTCCACAGTGGTC GGAGGAGACGTAGAGTTTGTCTGCAAGGTTTACAGTGATGCCAGCCCCACATCCAG TGGATCAAGCACGTGGAAGAACCGGCAGTAATAACGGGCCCGAGCGGCTGCGCTAC CTCAAGGTTCTCAAGGCGCGCGGTGTTAACACACGACAGAAAGAGATTGAGGTTCTC TATATTGGAATGTAACTTTTGGAGACGCTGGGGAATATACGTGCTTGGCGGGTAAT TCTATTGGGATATCCTTCTACTCTGCATGGTTGACAGTTCTGCCAGCGCCTGGAAGA GAAAGAGAGATTACAGCTTCCCGAGCTACCTGGAGATAGCCATTTACTGCATAGGG GTCTTCTTAATCGCTGTATGGTGGTAACAGTCATCCTGTGCGGAATGAAGAACAG ACCAAGAAGCCAGACTTCAGCAGCCAGCGGCTGTGCACAAGCTGACCAACAGTATC CCCTGCGGAGACAGGTAACAGTTTCGGCTGAGTCCAGCTCCTCCATGAACCTCAAC ACCCCGCTGGTGAGGATAACAACACGCTCTCTTCAACGGCAGACACCCCATGTCTG GCAGGGGTCTCCGAGTATGAACCTCCAGAGGACCAAAATGGGAGTTTCCAAGAGAT AAGCTGACACTGGGCAAGCCCTGGGAGAAGGTGTCTTGGGCAAGTGGTCATGGCG GAAGCAGTGGGAATTGACAAAGACAAGCCCAAGGAGCGGTCACCGTGGCCGTGAAG ATGTTGAAAGATGATGCCACAGAGAAAGACCTTCTGATCTGGTGTGAGAGTGGAG ATGATGAAGATGATTGGGAAACACAAGAAATATCATAAATCTTCTTGGAGCTGCACA CAGGATGGGCTCTCTATGTATAGTTGAGTATGCCCTCTAAAGGCAACCTCCGAGAA TACCTCCGAGCCCGAGGCCACCCGGGATGGAGTACTCTATGACATTAACCGTGT CCTGAGGAGCAGATGACCTTCAAGGACTTGGTGTATGCACCTACCAGCTGGCCAGA GGCATGGAGTACTTGGCTTCCCAAAATGTATTCATCGAGATTTAGCAGCCAGAAAT GTTTTGTTAAGAGAAACAAATGTGATGAAATAGCAGACTTTGGACTCGCCAGAGAT ATCAACAATATAGACTATTACAAAAGACCAACCAATGGGCGGCTTCCAGTCAAGTGG ATGGCTCCAGAGCCCTGTTTGTATAGAGTATACACTCATCAGAGTGTCTGGTCTC TTCGGGGTGTAAATGTGGGAGATCTTCACTTAGGGGGCTCGCCCTACCAGGGGATT CCCGTGGAGGAACTTTTAAGCTGCTGAAGGAAGGACACAGAATGGATAAGCCAGCC AACTGCACCAACGAACCTGTATCATGATGATGAGGACTGTTGGCATGCAGTGCCCTCC CAGAGACCAACGTTCAAGCAGTTGGTAGAAGACTTGGATCGAATTTCACTCTCACA ACCAATGAGGAATACTTGGACCTCAGCCAACCTCTCGAACAGTATTACCTAGTTAC CCTGACACAAGAGTTCTTGTCTTCCAGGAGATGATCTGTTTTTCTCCAGACCCC ATGCCTTACGAACCATGCCTTCTCAGTATCCACACATAAACCGCAGTGTTAAACA TGAATGACTGTGTCTGCTGTCCCAAACAGGACAGCACTGGGAACCTAGCTACACT GAGCAGGAGAGACCATGCCTCCAGAGCTTGTGTCTCCACTTGTATATATGGATCAG AGGAGTAAATAATGGAAAAGTAATCAGCATATGTGTAAGATTATACAGTTGAAA ACTTGTAATCTTCCCGAGGAGAGAAGAGGTTTCTGGAGCAGTGGACTGCCACAAG CCACCATGTAAACCCCTCTCACCTGCCGTGCGTACTGGCTGTGGACCAGTAGGACTCA AGGTGGACGTGCGTTCTGCTTCTTGTAAATTTGTAATAATTGGAGAAGATTAT GTCAGCACACACTTACAGAGCACAATGCGAGTATATAGGTGCTGGATGTATGTAAT ATATTCAAATATGTATAAATATATATATATATTTACAAGGAGTTATTTTTGTAT TGATTTTAAATGGATGCCCAATGCACCTAGAAAATTGGTCTCTCTTTTTTAAATAG CTATTTGCTAAATGCTGTTCTTACACATAATTTCTTAATTTTACCAGCAGAGGTG GAAAAATACTTTTGCTTTTCCAGGAAAATGGTATAACGTTAATTTATTAATAAATGG TAATATACAAAACAAATTAATCATTATAGTTTTTTTTGTAATTAAGTGGCATTCT ATGCAGGCAGCACAGCAGACTAGTTAATCTATTGCTTGGACTTAACAGTTATCAGA TCCTTTGAAAAGAGAATATTTACAATATATGACTAATTTGGGGAAAATGAAGTTTG ATTTATTTGTGTTTAAATGCTGCTGTGAGACGATTGTTCTTAGACCTCCTAAATGCC CCATATTTAAAGAACTCATTATAGGAAGTGTTCATTTTGTGTGCAACCTGTG

TABLE 5-continued

FGFR sequences.		
SEQ ID	FGFR2	ATTACGTCAACGCAACGTCTAACTGGACTTCCCAAGATAAATGGTACCAGCGTCCTC TAAAAAGATGCCTTAATCCATTCCCTTGAGGACAGACCTTAGTTGAAATGATAGCAGA ATGTGCTTCTCTCGCAGCTGGCCTTCTGCTTCTGAGTTGCACATTAATCAGATTA GCCTGTATTCTCTTCAGTGAATTTTGATAATGGCTTCCAGACTCTTTGGCGTTGGAG ACGCCTGTAGGATCTTCAAGTCCCATCATAGAAAATTGAAACACAGAGTTGTTCTG CTGATAGTTTGGGGATACGTCCATCTTTTAAGGGATTGCTTTCATCTAATTTCTGG CAGGACCTCACCAAAAGATCCAGCCTCATACCTACATCAGACAAAATATCGCCGTTG TTCCTTCTGTACTAAAGTATTGTGTTTTGCTTTGGAAACACCCACTCACTTTGCAAT AGCCGTGCAAGATGAATGCAGATTACACTGATCTTATGTGTTACAAAATTGGAGAAA GTATTTAATAAAACCTGTTAATTTTATACTGACAATAAAAATGTTTCTACAGATAT TAATGTTAACAGACAAAATAAATGTACGCAACTTA
NO: 22	isoform 2 Amino acid sequence	MVSWGRFICLVVVTMATLSLARPSFSLVEDTTLEPEEPPTKYQISQPEVYVAAPGES LEVRCLLKDAAVISWTKDGVHLGPNNRTVLIGEYLQIKGATPRDSGLYACTASRTVD SETWYFVMVNTDAISSGDDEDDTDGAEDFVSENSNNKRAPYWTNTEKMEKRLHAVPA ANTVKFRCPAGGNPMPTMRWLKNGKEFKQEHRIGGYKVRNQHWSLIMESVVP SDKGN YTCVENEYGSINHTYHLDVVERS PHRP ILQAGLPANASTVVGDDVEFVKVYSDAQ PHIQWIKHVEKNGSKYGPDLPLYLKVLAAGVNTTDKEIEVLYIRNVT FEDAGEYTC LAGNSIGISFHSAWLTVLPAPGREKEITASPDYLEIAIYICIGVFLIACMVVTVILCR MKNTTKKPDFSQPAVHKLTKRIPLRQVTVSAESSSSMNSNTPLVRIITRLSSAD TPMLAGVSEYELPEDPKWEPFRDKLTGKPLGEGCFGQVVMAEAVGIDKDKPKKEAVT VAVKMLKDDATEKDLSDLVSEMMEKMI GKHKNI INLLGACTQDGPLYVIVEYASKG NLREYLRRRRP PGMEYSYDINRVPEEQMTFKDLVSCYQLARGMEYLASQKCIHRDL AARNVLVTENNVMKIADFLGLARDINNIDYKKTNGRLPVKWMPEALFDRVYTHQS DVWSFGVLMEWIFTLGGSPYPGIPVEELFKLLKEGHRMDKPANCTNELYMMMRDCWH AVPSQRPTFKQLVEDLDRILTLTNEEYLDLSQPLEQYSPSYDTRSSCSSGDDSVF SPDPMPYEPCLPQYPHINGSVKT
SEQ ID	FGFR2	ACAGACTCTCCCGCAGAAGTACCCAGCAAGAAGCCTTTGGGAGCAGTAGAGATGG AGTTTCACTATGTTGCCCAGGCTAGCCTTGAACCTCCTGACCTCAGATGATCTGCCCG CGCAGGCCTCCCGAAGTGCTGGGATTACAGGCATGAGCCACCGCCTGGCCTGCCA ACTCTTGTAAAGATCTCGAAGGAAACATTTTCTCCCTGAAGGAAACCCAGCTATG CAGACACAGCTGATAATCTTGCAATCTCTGAAAGATGTTGCACCCCTATGGCAAGTG GCGGCTGCTGAGGCTCTGACGTGACTCCAGGCATGAACGCTCTCAGCTGTGTTTAC CTCAGCTCCTCGGAGGGAGCCTGGGAGACTGACGCCTGAGTTTACATCAGTGTCAT AAACCCAGCACAACCTAGGAGGGGACCTCCTGCCTAGTGTGTGTGGGTGAGGAGAT AGAAAAGCTCTCACTGAGTAACTGGACAAGGTCAATATACCTCGCTGATTGAGAAG ACTTCACTCTCTGCAAAAGAGACGTGTGTGTTTAGAGGAAGTGGGAGCCCCAGCC GATTCTGCAAGACTTCCGAGAGTCAGATATCCAGACAGAAGATGCGGACACCTGGGT GACCAGACAGCGAAGAGGAAAGAAACAAAACGAGCATGTGCCAAGCCTGTGAGGGAGA AAGGGCAACAAACAGTGACCTTCCACAGAAATGTGTTTAAACAAAACAAAACAGCT CTTTGGCGTTGTCTAAGAGACTGCCATTTTGAGGAAAGAGCGATCGCCTCACCGGCC CATCCTCCAAGCCGACTGCCCGCAATGCCTCCACAGTGGTCGGAGGAGACGTAGA GTTTGTCTGCAAGTTTACAGTGATGCCAGCCCCACATCCAGTGGATCAAGCAGCT GGAAAAGAACGGCAGTAAATACGGGCCCGACGGGCTGCCCTACCTCAAGGTTCTCAA GGCCGCCGCTGTTAAACACCACGGACAAAGAGATTGAGGTTCTCTATATTGGAAATGT AAGTTTGGAGGAGCTGGGGAATATACGTGCTTGGCGGGTAATTCTATTGGGATATC CTTTCACTCTGCAATGGTTGACAGTTCTGCCAGCGCTGGAAGAGAAAAGGAGATTAC AGCTTCCCAGACTACCTGGAGATAGCCATTTACTGCATAGGGGTCTTCTTAATCGC CTGTATGGTGGTAACAGTCATCTGTGCCGAATGAAGAACACGACCAAGAAGCCAGA CTTCAGCAGCCAGCGGCTGTGCACAAGCTGACCAAACGTATCCCCCTGCGGAGACA GGTAACAGTTTTCGGCTGAGTCCAGCTCCTCCATGAACCTCAACACCCCGCTGGTGAG GATAACAAACAGCCCTCTCTCAACGGCAGACACCCCATGCTGGCAGGGGTCTCCGA GTATGAACCTTCAGAGGACCCAAAATGGGAGTTTCCAAGAGATAAGCTGACACTGGG CAAGCCCTGGGAGAAGGTTGCTTTGGGCAAGTGGTCATGGCGGAAGCAGTGGGAAT TGACAAAGACAAGCCCAAGGAGCGGTACCCGTGGCGCTGAAGATGTTGAAAGATGA TGCCACAGAGAAAGACCTTTCTGATCTGGTGTGAGAGATGGAGATGATGAAGATGAT TGGGAAACACAAGATATCATAAATCTTTTGGAGCCTGCACACAGGATGGGCTCT CTATGTCATAGTTGAGTATGCCCTTAAAGGCAACCTCCGAGAATACCTCCGAGCCCG GAGGCCACCCGGGATGGAGTACTCTATGACATTAACCGTGTTCCTGAGGAGCAGAT GACCTTCAAGGACTTGGTGTATGCACCTACCAGCTGGCCAGAGGCATGGAGTACTT GGCTTCCCAAAAATGTAATTCATCGAGATTTAGCAGCCAGAAATGTTTGGTAACAGA AAACAATGTGATGAAAATAGCAGACTTTGGAAGTCCGAGAGATATCAACAATATAGA CTATTACAAAAGACCACCAATGGGCGGCTTCCAGTCAAGTGGATGGCTCCAGAAGC CCTGTTTGTAGATATACACTATCAGAGTGATGTCTGGTCTTCCGGGTGTTAAT GTGGGAGATCTTCACTTTAGGGGGCTCGCCCTACCCAGGGATTCCCGTGGAGGAAC TTTAAAGCTGCTGAAGGAAGGACACAGAATGGATAAGCCAGCCAACTGCACCAACGA ACTGTACATGATGATGAGGAGTGTGGCATGCAAGTCCCTCCAGAGACCAACGTT CAAGCAGTTGGTAGAAGACTTGGATCGAATTTCTCACTCTCACAAACCAATGAGGAATA CTTGGACCTCAGCCAACTCTCGAACAGTATTCACCTAGTTACCTTGACACAAGAAG TTCTTGTCTTTCAGGAGATGATCTGTGTTTTTCTCCAGACCCCATGCTTACGAACC ATGCCTTCTCAGTATCCACACATAAACGGCAGTGTAAACATGAATGACTGTGTC TGCTGTCCCCAACAGGACAGCACTGGGAACCTAGCTACACTGAGCAGGAGAGACCA TGCTTCCAGAGCTTGTGTCTCCACTGTATATATGGATCAGAGGAGTAAATAAT GGAAAAGTAATCAGCATATGTGTAAGATTATACAGTTGAAAACCTGTAAATCTTCC CCAGGAGGAGAAGAAGTTTCTGGAGCAGTGGACTGCCACAAGCCACCATGTAAACCC CTCTCACTGCCGTGCTACTGGCTGTGGACCAAGTAGGACTCAAGGTGGACGTGCGT TCTGCCCTTCTTGTAAATTTTGTAAATATGGAGAAGATTTATGTCAGCACACACTT
NO: 23	isoform 3 Nucleic acid sequence	

TABLE 5-continued

FGFR sequences.		
SEQ ID NO: 24	FGFR2 isoform 3 Amino acid sequence	ACAGAGCACAAATGCAGTATATAGGTGCTGGATGTATGTAATATATATTCAAATTATG TATAAATATATATTATATATTACAAGGAGTTATTTTTTGTATTGATTTTAAATGGA TGTCCTCAATGCACCTAGAAAATTGGTCTCTCTTTTTTAATAGCTATTTGCTAAATG CTGTTCTTACACATAATTTCTTAATTTTACCCGAGCAGAGGTGGAAAAATACTTTTG CTTTTCAGGGAAAAATGGTATAACGTTAATTTTATTAATAAATGGTAATATACAAAACA ATTAATCATTATAGTTTTTTTTGTAAATTAAGTGGCATTCTATGCAGGCAGCACACA GCAGACTAGTTAATCTATTGCTTGGACTTAAGTAGTTATCAGATCCTTTGAAAAGAG AATATTTACAATATATGACTAATTTGGGGAAAAATGAAGTTTGTATTATTGTGTTT AAATGCTGCTGTCAGACGATTGTTCTTAGACCTCCTAAATGCCCATATTAAAGAA CTCATTATAGGAAGGTGTTTCATTTTGGTGTGCAACCTGTCTATTACGTCAACGCA ACGTCTAAGTGGACTTCCCAAGATAAATGGTACCAGCGTCTCTTAAAGATGCCTT AATCCATTCTTGGAGCAGACCTTAGTTGAAATGATAGCAGAAATGTGCTCTCTCT GGCAGCTGGCCTTCTGCTTCTGAGTTGCACATTAATCAGATTAGCCTGTATTCTCTT CAGTGAATTTTGATAATGGCTTCCAGACTCTTTGGCGTTGGAGACGCTGTTAGGAT CTTCAAGTCCCATCATAGAAAATTGAAACACAGAGTTGTTCTGCTGATAGTTTTGGG GATACGTCCTATCTTTTAAAGGATGCTTTTCATCTAATTTCTGGCAGGACCTCACCAA AAGATCCAGCTCATACCTACATCAGACAAAATATCGCCGTGTCTCTCTGTACTA AAGTATTGTGTTTTGCTTTGGAAACCCCACTCACTTGAATAGCCGTGCAAGATG AATGCAGATTACACTGATCTTATGTGTTACAAAATGGAGAAAGTATTAAATAAAC CTGTTAATTTTTATCTAGACAATAAAAATGTTTCTACAGATATTAAATGTTAAACAAGA
SEQ ID NO: 25	FGFR2 isoform 4 Nucleic acid sequence	TGACTGCAGCAGCAGCGCGCAGCGCTCGGTTCTGAGCCACCGCAGGCTGAAGGCA TTGCGCGTAGTCCATGCCCCGTAGAGGAAGTGTGCAGATGGGATTAACGCTCCACATGG AGATATGGAAGAGGACCGGGGATTGGTACCGTAACCATGGTCAGCTGGGTCGCTTTC ATCTGCTGGTCTGTTGGTCAACATGGCAACCTTGTCCCTGGCCCGGCCCTCCTTCAGT TTAGTTGAGGATACACATTAGAGCCAGAAGAGCCACCAACCAATACCAATCTCT CAACCAGAAGTGTACGTGGCTGCGCCAGGGGAGTTCGCTAGAGGTGCGCTGCCTGTTG AAAGATGCCCGCTGATCAGTTGGACTAAGGATGGGTGCACTTGGGGCCCAACAAT AGGACAGTGCTTATTGGGGAGTACTTGCAGATAAAGGGCCGACGCTAGAGACTCC GGCCTCTATGCTTGTACTGCCAGTAGGACTGTAGACAGTGAAACTTGGTACTTCATG GTGAATGTCACAGATGCCATCTCATCCGAGATGATGAGGATGACACCGATGGTGCG GAAGATTTGTGAGTGAAGACAGTAACAACAAGAGAGCACCATACTGGACCAACACA GAAAAGATGAAAAGCGGCTCCATGCTGTGCTGCGGCCAACACTGTCAAGTTTCGC TGCCAGCGCGGGGGAACCAATGCCAACATGCGGTGGCTGAAAACCGGGAAGGAG TTTAAGCAGGAGCATCGCATTTGGAGGCTACAAGGTACGAAACAGCACTGGAGCCTC ATTATGGAAGTGTGGTCCCATCTGACAAAGGAAATTATACCTGTGTAGTGGAGAAT GAATACGGGTCCATCAATCACACGTACCACCTGGATGTTGTGGCGCTGGAAGAGAA AAGGAGATTACAGCTTCCCCAGACTACCTGGAGATAGCCATTTACTGCATAGGGGTC TTCTTAATCGCCTGTATGGTGGTAACAGTCATCCTGTGCCGAATGAAGAACACGACC AAGAAGCCAGACTTCAGCAGCCAGCCGGCTGTGCACAAGCTGACCAACGATATCCCC CTGCGGAGACAGGTAACAGTTTCGGCTGAGTCCAGCTCCTCCATGAACCTCAACACC CCGCTGGTGAGGATAACAACACGCTCTCTTCAACGGCAGACACCCCATGCTGGCA GGGTCTCCGAGTATGAACCTTCCAGAGGACCAAAATGGGAGTTTCCAAGAGATAAG CTGACACTGGGCAAGCCCTGGGAGAAGGTTGCTTTGGGCAAGTGGTCATGGCGGAA GCAGTGGGAATTGACAAAGACAAGCCCAAGGAGGCGGTACCGTGGCCGTGAAGATG TTGAAAGATGATGCCACAGAGAAAGACCTTTCTGATCTGGTGTGAGAGATGGAGATG ATGAAGATGATTGGGAAACACAAGAAATATCAATAATCTTCTGGAGCCTGCACACAG GATGGGCTCTCTATGTATAGTTGAGTATGCTCTAAAGGCAACCTCCGAGAATAC CTCCGAGCCCGGAGGCCACCCGGGATGGAGTACTCTATGACATTAAACGTTGTTCT GAGGAGCAGATGACCTTCAAGGACTTGGTGTGATGCACTTACCAGTGGCCAGAGGC ATGGAGTACTTGGCTTCCCAAAATGTATTATCATGAGATTAGCAGCCAGAAATGTT TTGGTAACAGAAAACAAATGTATGATAAAATAGCAGACTTTGGACTCGCCAGAGATATC AACAAATATAGACTATTACAAAAGACCACCAATGGGCGGCTTCCAGTCAAGTGGATG GCTCCAGAAGCCCTGTTTGATAGAGTATACACTCATCAGAGTATGCTGGTCTCTC GGGTGTTAATGTGGGAGATCTTCACTTTAGGGGGCTCGCCCTACCAGGGATTCCC GTGAGGAAGCTTTTAAAGTGTGTAAGGAAGGACACAGAATGGATAAGCCAGCCAAC TGCACCAACGAACGTATCATGATGATGAGGGACTGTTGGCATGCAGTGCCTCCCAG AGACCAACGTTCAAGCAGTTGGTAGAAGACTTGGATCGAATTCTCACTCTCACAACC AATGAGGAATACTTGGACCTCAGCAACCTCTGAACAGTATTACCTAGTTTACCTT GACACAAGAAGTCTTGTCTTTCAGGAGATGATTCTGTTTTTCTCAGACCCCATG CCTTACGAACCATGCCTTCTCAGTATCCACACATAAACGGCAGTGTTAAACATGA ATGACTGTGTGCTGTCTTCCCAACAGGACAGCACTGGGAACCTAGCTACACTGAG CAGGGAGACCATGCCTCCCAGAGCTTGTGTCTCCACTTGTATATATGGATCAGAGG AGTAAATAATTGGAAGTAATCAGCATATGTGTAAGATTATACAGTTGAAAACCT TGTAATCTTCCCAGGAGGAGAAGAGGTTTCTGGAGCAGTGGACTGCCACAAGCCA CCATGTAACCCCTCTCACCTGCCGTGCGTACTGGCTGTGAGACCAGTAGGACTCAAG

TABLE 5-continued

FGFR sequences.		
SEQ ID NO: 26	FGFR2 isoform 4 Amino acid sequence	TGGACGTGCGTTCTGCGCTTCTGTTAATTTTGTAAATTTGGAGAAGATTTATGTC AGCACACACTTACAGAGCACAAATGCAGTATATAGGTGCTGGATGTATGTAATATA TTCAAATTATGTATAAATATATATATATATTTTACAAGGAGTTATTTTTTGTATTGA TTTTAAATGGATGTCCTCAATGCACCTAGAAAATGGTCTCTCTTTTTTAAATAGCTA TTTGCTAAATGCTGTTCTTACACATAATTTCTTAATTTTACCAGAGCAGAGGTGGAA AAATACTTTTGCTTTTCAAGGAAAATGGTATAACGTTAATTTATTAATAAATGGTAA TATACAAAACAATTAATCATTATAGTTTTTTTGTAAATTAAGTGGCATTCTATG CAGGCAGCACAGCAGACTAGTTAATCTATTGCTTGGACTTAAGTATATCAGATCC TTTGAAAGAGAAATATTACAATATATGACTAATTTGGGAAAATGAAGTTTGTATT TATTTGTGTTTAAATGCTGCTGTCAGACGATTGTTCTTAGACCTCTAAATGCCCCA TATTAAAGAACTCATTATAGGAAGGTGTTTCAATTTGGTGTGCAACCTGTCATT ACGTCAACGCAACGCTTAACCTGGACTTCCCAAGATAAATGGTACCAGCGTCTCTTA AAAGATGCTTAAATCCATTCTTGGAGACAGACCTTAGTTGAAATGATAGCAGAATG TGCTTCTCTGCGCAGCTGGCCTTCTGCTTCTGAGTTGCACATTAATCAGATTAGCC TGATTCTCTTCAAGTCCCATCATAGAAAATGAAACACAGAGTTGTTCTGCTG CTGTTAGGATCTTCAAGTCCCATCATAGAAAATGAAACACAGAGTTGTTCTGCTG ATAGTTTGGGGATACGTCCTATCTTTTAAAGGATGCTTTTCACTAATCTGCGCAG GACCTCACCAAAAGATCCAGCCTCATACCTACATCAGACAAAATATCGCCGTTGTTT CTTCTGTACTAAAGTATTGTGTTTGTCTTTGAAAACACCCACTCACTTTGCAATAGC CGTGCAAGATGAATGCAGATTACACTGATCTTATGTGTTACAAAATGGAGAAAAGTA TTTAATAAAACCTGTTAATTTTATACTGACAATAAAAATGTTTCTACAGATATTAA TGTTAAACAAGACAAAATAATGTACGCAACTTATTTTTTT MVSWGRFICLVVVTMATLSLARPSFSLVEDTTLEPEEPPTYKQISQPEVYVAAPGES LEVRCCLKDAAVISWTKDGVHLGPNRNTVLIGEYLQIKGATPRDSGLYACTASRTVD SETWYFMVNVTDIAISSGDEDDTDGAEDFVSENSNNKRAPYWTNTEKMEKRLHAVPA ANTVKFRCPAGGNPMPTMRWLKNGKEFKQEHRI GGYKVRNQHWSLIMESVVPDCKGN YTCVVENEYGSINHTYHLDVVPAGREKEITASPDYLEIAIYICIGVFLIACMVVTIL CRMKNNTKKPDFSSQPAVHKLTKRIPLRRQVTVSAESSSSMNSNTPLVRITRRLSST ADTPMLAGVSEYELPEDPKWEFPRDKLTGKPLGEGCFGQVMAEAVGIDKDKPKKEA VTVAVKMLKDDATEKDLSDLVSEMEMMKMIGKHNIINLLGACTQDGPLYVIVEYAS KGNLREYLRARRPPGMEYSYDINRVP EEQMTFKDLVSCYQLARGMEYLASQKCIHR DLAARNVLVTENNVMKIADFGGLARDINNIDYKKTNGRLPVKWMPEALFDRVYTH QSDVWSFGVLMWEIFTLGGSPYPGIPVEELFKLLKEGHRMDKPA NCTNELYMMMRDC WHAVPSQRPTFKQLVEDLDRILTLTTNEEYLDLSQPLEQYSPSYDPTRSSCSSGDDS VFSPDMPYEPCLPQYPHINGSVKT
SEQ ID NO: 27	FGFR2 isoform 5 Nucleic acid sequence	AATTTGTTGAGGAATTTCCCCCTAGCCTTGACCCCTTGACAGCTCCCGCTCCTACTC AGTGCTGGGGAGAAGTAGGGAGGCCTTAAGCGAAGAGATGGGCTGCACTTTGGAGG AGCCGGACACTGTTGACTTTTCTGATGTGAAATCTACCCAGGAACAAAACACCACTG ACTGCAGCAGCAGCGCAGCGCTCGGTTCTGAGCCACCCGAGGCTGAAGGCATT GCGCGTAGTCCATGCCGTAGAGGAAGTGTGCAGATGGGATTACGCTCCACATGGAG ATATGGAAGAGGACCGGGGATTGGTACCGTAACCATGGTCACTGGGTCGTTTCAT CTGCTCGTGTGTTTACCATGGCAACCTTGTCCTTGGCCCGGCCCTCCTTCAGTTT AGTTGAGGATACACATTAGAGCCAGAAGATGCCATCTCATCCGGAGATGATGAGGA TGACACCGATGGTGCGGAAGATTTTGTCACTGAGAACAGTAACAACAAGAGAGCACC ATACTGGACCAACACAGAAAAGATGGAAAAGCGGCTCCATGCTGTGCTTGCAGCCAA CACTGTCAAGTTTCTGCTGCCAGCAGCGGGGGAACCAATGCCAACATGCGGTGGCT GAAAAACGGGAAGGAGTTTAAAGCAGGAGCATCGCATTTGGAGGCTACAAGGTACGAAA CCAGCACTGGAGCTCATTATGGAAGTGTGGTCCCATCTGACAAGGGAATATATAC CTGTGTAGTGAGGAATGAATACGGGTCCATCAATCACACGTACCACCTGGATGTTGT GGAGCGATCGCCTCACCGGCCATCCTCCAAAGCCGACTGCCGGCAATGCCCTCCAC AGTGGTCGGAGGAGACGTAGAGTTTGTCTGCAAGGTTTACAGTGATGCCAGCCCA CATCCAGTGATCAAGCACGTGGAAGAAGACGGCAGTAAATACGGGCCCGACGGGCT GCCCTACCTCAAGTTTCTCAAGGCCCGCGGTGTTAACACCACGGACAAAAGAGATTGA GGTTCTCTATATTCGGAATGTAACTTTGGAGACGCTGGGGAATATACGTGCTTGGC GGGTAATTCATTGGGATATCCTTCACTCTGCATGGTTGACAGTTCTGCCAGCGCC TGGAAGAGAAAAGGAGATTACAGCTTCCCGAGACTACCTGGAGATAGCCATTTACTG CATAGGGGTCTTCTTAATCGCCTGTATGGTGGTAACAGTCATCCTGTGCCGAATGAA GAACACGACCAAGAAGCCAGACTTCAGCAGCCAGCCGGCTGTGCACAAGCTGACCAA ACGTATCCCCCTGCGGAGACAGGTAACAGTTTCCGGTGAAGTCCAGCTCCTCCATGAA CTCCAACACCCCGCTGGTGAAGATAACAACACGCTCTCTTCAACGGCAGACACCCC CATGCTGGCAGGGGTCTCCGAGTATGAACCTCCAGAGGACCCAAAATGGGAGTTTCC AAGAGATAAGCTGACACTGGGCAAGCCCTGGGAGAAGGTTGCTTTGGGCAAGTGGT CATGGCGGAAGCAGTGGGAATTGACAAAGACAAGCCCAAGGAGCGGTCACCGTGGC CGTGAAGATGTTGAAAGATGATGCCACAGAGAAGACCTTCTGATCTGGTGTGAGA GATGGAGATGATGAAGATGATTGGGAAACACAAGAATATATAAATCTTCTTGGAGC CTGCACACAGGATGGGCTCTCTATGTCATAGTTGAGTATGCTTAAAGGCAACCT CCGAGAATACCTCCGAGCCCGGAGGCCACCCGGGATGGAGTACTCTATGACATTAA CCGTGTTCTCTGAGGAGCAGATGACCTTCAAGGACTTGGTGTGATGCACCTACCAGCT GGCCAGAGGATGAGTACTTGGCTTCCCAAAAATGTATTTCATCAGAGATTAGCAGC CAGAAATGTTTGGTAACAGAAAACAATGTGATGAAAATAGCAGACTTTGGACTCGC CAGAGATATCAACAATATAGACTATTACAAAAGACCAACAATGGGCGGCTTCCAGT CAAGTGGATGGCTCCAGAAGCCCTGTTTGTATAGAGTATACACTCATCAGAGTATGT CTGGTCTTTCGGGGTGTAAATGTGGGAGATCTTCACTTTAGGGGGCTCGCCCTACCC AGGGATTCCTGAGGGAACCTTTTAAAGTGTCTGAAGGAAGGACACAGAATGGATAA GCCAGCCAAGTGCACCAACGAAGTGTACATGATGATGAGGAGCTGTTGGCATGCAGT GCCCTCCAGAGACCAACGTTCAAGCAGTTGGTAGAAGACTTGGATCGAATTCAC

TABLE 5-continued

FGFR sequences.		
SEQ ID NO. 28	FGFR2 isoform 5 Amino acid sequence	TCTCACAACCAATGAGGAGGAGAAGAAGGTTTCTGGAGCAGTGGACTGCCACAAGCC ACCATGTAACCCCTCTCACCTGCCGTGCGTACTGGCTGTGGACCAGTAGGACTCAAG GTGGACGTGCGTTCTGCCTTCTTGTAAATTTTGTAAATTTGGAGAAGATTTATGT CAGCACACACTTACAGAGCACAATGTCAGTATATAGGTGCTGGATGTATGTAAATAT ATTCAAATTATGTATAAATATATATTATATATTACAAAGGAGTTATTTTTGTATTG ATTTTAAATGGATGCCAATGCACCTAGAAAATTTGGTCTCTCTTTTTTAAATAGCT ATTTGCTAAATGCTGTTCTTACACATAATTTCTTAATTTTCACCGAGCAGAGGTGGA AAAATACTTTTGTCTTTCAGGGAATGGTATAACGTTAATTTATTAATAAATGGTA ATATACAAAACAATTAATCATTATAGTTTTTTTTGTAAATTTAAGTGGCATTCTAT GCAGGCAGCAGCAGACTAGTTAATCTATTGCTTGGACTTAACTAGTTATCAGATC CTTTGAAAAGAGAATATTACAAATATATGACTAATTTGGGGAATGAAGTTTTGAT TTATTTGTGTTTAAATGCTGCTGTGAGACGATTGTTCTTAGACCTCCTAAATGCCCC ATATTAAAGAACTCATTTCATAGGAAGGTGTTTCATTTTGGTGTCAACCCCTGTCT TACGTCAACGCAACGCTCAACTGGACTTCCCAAGATAAATGGTACCAGCGTCTCTT AAAAGATGCTTAATCCATTCTTGGAGCAGACCTTAGTTGAAATGATAGCAGAAT GTGCTTCTCTTGGCAGCTGGCCTTCTGCTTCTGAGTTGCACATTAATCAGATTAGC CTGTATTCTCTCAGTGAATTTTGATAATGGCTTCCAGACTCTTGGCGTTGGAGAC GCCTGTTAGGATCTTCAAGTCCATCATAGAAAATTGAAACACAGAGTTGTTCTGCT GATAGTTTTGGGGATACGTCCATCTTTTTAAGGGATTGCTTTCATCTAATTTCTGGCA GGACCTCACAAAAGATCCAGCCTCATACCTACATCAGACAAAATACGCGGTGTT CCTTCTGTACTAAAGTATTGTGTTTGTCTTGGAAACACCCACTCATTGCAATAG CCGTGCAAGATGAATGCAGATTACACTGATCTTATGTGTTACAAAATGGAGAAAGT ATTTAATAAACCTGTTAATTTTATACTGACAATAAAATGTTTCTACAGATATTA ATGTTAACAAGACAAAATAAATGTACGCAACTTATTTTTTT
SEQ ID NO. 29	FGFR2 isoform 6 Nucleic acid sequence	MVSWGRFICLVVVTMATLSLARPSFSLVEDTTLEPEDAISSGDEDDTDGAEDFVSE NSNNKRAPYWTNTEKMEKRLHAVPAANTVKFRCPAGGNPMTMRWLKNGKEFKQEH IGGYKVRNQHWLSIMESVVPDKNYTCVVENEYGSINHTYHLDVVERSHPRPILQA GLPANASTVVGDDVEFVKVYDAQPHIQWIKHVEKNKSGYGPDLPLKVLKAAGV NTTDKEIEVLYIRNVTFEDAGEYTCLAGNSIGISFHSAWLTVLPAPGREKEITASPD YLEIAIYICGVFLIACMVVTVILCRMKNNTTKPDFSSQPAVHKLTKRIPLRRQVTVS AESSSSMNSNTPLVRIITRLSSADT PMLAGVSEYELPEDPKWFFPRDKLTGLKPLG EGCFGQVMAEAVGIDKDKPKEAVTVAVKMLKDDATEKDLSDLVSEMEMMKMIGKHK NIINLLGACTQDGPLYVIVEYASKGNLREYLRARRPPGMEYSYDINRVPEEQMTFKD LVSCYQLARGMEYLASQKCIHRDLAARNVLVTENNVMKIADFLGARDINNIIDYYKK TTNGRLPVKWMPEALFDRVYTHQSDVWSPGLMWEIFTLGGSPYPGIPVEELFKLL KEGRHMDKPNANTNELYMMMRDCHWAVPSQRPTFKQLVEDLDRILTLTNEEEKKVS GAVDCHKPPCNPSPHLPCVLAVDQ GGCGGCGGCTGGAGGAGAGCGCGGTGGAGAGCCGAGCGGGCGGGCGGGTGC GCGGCGAGGGAGCGCGCGCGGCCGCCACAAAGCTCGGGCGCGCGGGGTGCATGC GGCGTACCTGGCCCGCGCGCGGCGACTGCTCTCCGGGTGGCGGGGGCGCGCCGCGAG CCCCGGGGGCCCCGAGGCCGAGCTTGCCCTGCGCGCTCTGAGCCTTCGCAACTCGCG AGCAAAAGTTTGGTGGAGGCAACGCCAAGCCTGAGTCCTTTCTCTCTCGTTCCCA AATCCGAGGGCAGCCCGCGGGCGTCATGCCCGCGCTCCTCCGAGCCTGGGGTACGC GTGAAGCCCGGAGGCTTGGCGCGCGGCAAGACCCAGGACCACTCTTCTGCGTTTG GAGTTGCTCCCGCAACCCCGGGCTCGTCGCTTCTCCATCCCGACCCACGCGGGG GCGGGGACACACAGGTGCGGAGGAGCGTTGCCATTCAAGTGACTGCAGCAGCAGC GGCAGCGCCTCGGTTCTGAGCCACCGCAGGCTGAAGGCATTGCGCGTAGTCCATG CCCGTAGGAAGTGTGCAGATGGGATTAACGTCCACATGGAGATATGGAAGAGGAC CGGGGATTGTTACGTAACCATGGTCAGCTGGGGTCGTTTCATCTGCCTGGTCTG TCACCATGGCAACCTTGTCCTGGCCCGGCCCTCTTCAGTTTAGTTGAGGATACCA CATTAGAGCCAGAAGGAGCACCATCTGAGCAACACAGAAAAGATGGAAGAGCGG TCCATGCTGTGCTGCGGCCAACACTGTCAAGTTTCGCTGCCAGCCGGGGGGAACC CAATGCCAACCATGCGGTGGCTGAAAAACGGGAAGGAGTTAAGCAGGAGCATCGCA TTGGAGGCTACAAGGTACGAAACACGACTGGAGCCTCATTATGGAAGTGTGGTCC CATCTGACAAGGGAAATATACCTGTGTAGTGGAGAATGAATACGGGTCCATCAATC ACACGTACCACCTGGATGTTGTGGAGCGATCGCTCACCGGCCATCTCCAAGCCG GACTGCCGGCAATGCCTCCACAGTGGTGGAGGAGACGTAGAGTTGTCTGCAAGG TTTACAGTGATGCCAGCCACATCCAGTGGATCAAGCAGCTGGAAGAAAGAACGGCA GTAATACGGGCCCGACGGGCTGCCCTACCTCAAGGTTCTCAAGGCCCGCGGTGTTA ACACACGGACAAAGAGATTGAGGTTCTCTATATTGGAATGTAACTTTGAGGAGC CTGGGAATATACGTGCTTGGCGGGTAATCTATTGGGATATCTTTCACTCTGCAT GGTTGACAGTTCTGCCAGCGCCTGGAAGAGAAAAGGAGATTACAGCTTCCCAGACT ACCTGGAGATAGCCATTACTGCATAGGGGCTTCTTAACTCGCTGTATGGTGGTAA CAGTCATCTGTGCGGAATGAAGAACACGACCAAGAACCCAGACTTTCAGCAGCCAGC CGGCTGTGCACAAGCTGACCAACGATATCCCCGCGGAGACAGGTTTCCGGTGTAGT CCAGCTCTCTCAACTCAACACCCCGCTGGTGGAGATAACAACACCGCTCTCTT CAACGGCAGACACCCCATGCTGGCAGGGGTCTCCGAGTATGAACCTTCAGAGGACC CAAAATGGGAGTTTCAAGAGATAAGCTGACACTGGGCAAGCCCTGGGAGAAGGTT GCTTTGGCAAGTGGTCATGGCGAAGCAGTGGGAATTGACAAGACAAGCCCAAGG AGGCGGTACCGTGGCGTGAAGATGTTGAAAGATGATGCCACAGAGAAAGACCTTT CTGATCTGGTGTGAGAGATGAGATGATGAAGATGATTGGGAAACACAAGAATATCA TAAATCTTCTGGAGCCTGCACACAGGATGGGCTCTCTATGTCATAGTTGAGTATG CCTCTAAAGGCAACCTCCGAGAATACCTCCGAGCCCGGAGGCCACCGGATGGAGT ACTCCTATGACATTAACCGTGTCTGAGGAGCAGATGACCTTCAAGGACTTGGTGT CATGACCTACAGCTGGCCAGAGCATGGAGTACTTGGCTTCCAAAAATGTATTC ATCGAGATTTAGCAGCCAGAAATGTTTGGTAACAGAAAACATGTGATGAAAATG

TABLE 5-continued

FGFR sequences.		
SEQ ID NO: 30	FGFR2 isoform 6 Amino acid sequence	CAGACTTTGGAATCGCCAGAGATATCAACAATATAGACTATTACAAAAGACCACCA ATGGGCGGCTTCCAGTCAAGTGGATGGCTCCAGAAGCCCTGTTTGATAGAGTATACA CTCATCAGAGTGTGTCTGGTCTTCCGGGTGTTAATGTGGGAGATCTTCACTTTAG GGGGCTCGCCCTACCCAGGGATTCCCGTGGAGGAACCTTTTAAAGCTGCTGAAGGAAG GACACAGAATGGATAAGCCAGCCAACTGCACCAACGAAGTGTACATGATGATGAGGG ACTGTTGGCATGCAGTGCCCTCCAGAGACCAACGTTCAAGCAGTTGGTAGAAGACT TGGATCGAATTCTCACTCTCACAACCAATGAGGAATACTTGGACCTCAGCCAACCTC TCGAACAGTATTCACCTAGTTACCCTGACACAAGAAGTTCTTGTCTTCAGGAGATG ATTCTGTTTTTCTCCAGACCCATGCCTTACGAACCATGCCTTCCTCAGTATCCAC ACATAAACCGCAGTGTAAAAACATGAATGACTGTGTCTGCCTGTCCCAACAGGAC AGCACTGGGAACCTAGCTACACTGAGCAGGGAGACCATGCCTCCAGAGCTTGTGT CTCCACTTGTATATATGGATCAGAGGAGTAAATAATTGGAAAAGTAATCAGCATATG TGTAAGATTATACAGTTGAAAACCTTGTAATCTTCCCAAGGAGGAGAAGAAGGTTT CTGGAGCAGTGGACTGCCACAAGCCACCATGTAACCCCTCTCACCTGCCGTGCGTAC TGGCTGTGGACCAGTAGGACTCAAGGTGGACGTGCGTCTCGCTTCCTTGTTAATTT TGTAATAATTGGAGAAGATTATGTGACACACACTTACAGAGCACAAATGCAGTAT ATAGGTGCTGGATGTATGTAATATATTTCAAATATGTATAAATATATATATATAT TTACAAGGAGTATTTTTTGTATTGATTTAAATGGATGTCCCAATGCACCTAGAAA ATTGGTCTCTCTTTTTTAATAGCTATTGTCTAAATGCTGTTCTTACACATAATTT TTAATTTTCCAGCAGCAGAGGTGAAAAAATACTTTGCTTTCCAGGAAAAATGGTATA ACGTTAATTATTAATAAATTTGGTAATATACAAACAATTAATCATTATAGTTTTT TTTGTAAATTTAAGTGGCATTCTATGACGAGCAGACAGCAGACTAGTTAATCTATTG CTTGGACTTAAGTAGTTATCAGATCCTTTGAAAAGAGAATATTTACAATATATGACT AATTTGGGAAAA
SEQ ID NO: 30	FGFR2 isoform 6 Amino acid sequence	MVSWGRFICLVVMTATLSLARPSFSLVEDTTLEPEGAPYWTNTEKMEKRLHAVPAA NTVKFRCPAGGNPMTMRWLKNGKEFKQEHRIIGYKVRNQHWSLIMESVVPKDNV TCVVENEYGSINHTYHLVDVVERSPHRPILOAGLPANASTVVGDDVEFVKVYSDAQ HIQWIKHVEKNGSKYGPDLPLKVLKAAVNTTDEIEVLIRNVTFEDAGEYTL AGNSIGISFHSAWLTVLPAPGREKEITASPDYLEIAIYICIGVFLIAMVVTVILCRM KNTTKKPDFSSQPAVHKLTKRIPLRRQVSAESSSSMNSNTPLVRITTRLSSADTPM LAGVSEYELPEDPKWEFPRDKLTGLKPLGEGCFQGVMAEAVGIDKDKPEAVTVAV KMLKDDATEKDLSDLVSEMEMMKMIGKHNIINLLGACTQDGPLYVIVEYASKGNLR EYLRARRPPGMEYSYDINRVPEEQMTFKDLVSCYQLARGMEYLASQKCIHRDLAAR NVLVTENNVMKIADFLARDINNIDYKKTNGRLPVKWMAPALFDRVYTHQSDVW SFGVLMWEIFTLGGSPYPGIPVEELFKLLKEGHRMDKPANCTNELYMMMRDCWHAVP SQRPTFKQLVEDLDRILTLTNEEYLDLSQPLEQYSPSYDPDRSSCSSGDDSVFSPD PMPYEPCLPQYPHINGSVKT
SEQ ID NO: 31	FGFR2 isoform 7 Nucleic acid sequence	CCGGCCGCGAGCCCGGGGGCCCCGAGGCCGCGAGCTTGCCTGCGCGCTCTGAGCCTT CGCAACTCGCGAGCAAAGTTTGGTGGAGGCCAACGCCAAGCCTGAGTCCTTCTTCTCT CTCGTTCCCAAAATCCGAGGGCAGCCCGCGGGCGCTCATGCCCGCGCTCTCCCGCAGC CTGGGGTACGCGTGAAGCCCGGGAGGCTTGGCGCCGGCGAAGACCAAGGACCACTC TTCTGCGTTTGGAGTTGCTCCCGCAACCCCGGGCTCGTCGCTTTCTCCATCCCGAC CCACGCGGGGCGCGGGGACAACACAGGTGCGGGAGGAGCGTGCCTATTCAAGTGACT GCAGCAGCAGCGGCGAGCGCTCGGTTCTTGAGCCCAACCGCAGGCTGAAGGCATTGCG CGTAGTCCATGCGCGTAGAGGAAGTGTGCAGATGGGATTAACGTCCACATGGAGATA TGGAAGAGGACCGGGGATTGGTACCGTAACCATGGTCACTGGGGTGGTTTCATCTG CCTGGTCTGGTCAACATGGCAACCTTGTCCTTGGCCCGGCCCTCTTCACTTTAGT TGAGGATACCAATTAGAGCCAGAAGGCCAACCAAAATACCAATCTCTCAACC AGAAAGTGACGTGGCTGCGCCAGGGGAGTGCCTAGAGGTGCGCTGCTGTGAAAGA TGCCCGCGTGATCAGTTGGACTAAGGATGGGTGCACCTTGGGGCCCAACAATAGGAC AGTGCTTATGGGGAGTACTTGCAAGATAAAGGGCGCCACGCTTAGAGACTCCGGCT CTATGCTTGTAAGTGCAGTAGGACTGTAGACAGTGAAACTTGGTACTTCATGGTGAA TGTCACAGATGCCATCTCATCCGAGATGATGAGGATGACACCGATGGTGGCGAAGA TTTTGTGAGTGAGAACAGTAACAACAAGAGAGCACCATAGTGAGCCAAACAGAAAA GATGGAAAAGCGGCTCCATGCTGTGCTGCGGCCAACACTGTCAAGTTTCGCTGCCCC AGCCGGGGGAACCAATGCCAACCATGCGGTGGCTGAAAAACGGGAAGGAGTTTAA GCAGGAGCATCGCATTTGGAGGCTACAAGGTACGAAACAGCACTGGAGCCTCATTAT GGAAAGTGTGGTCCCATCTGACAGGGAAATTATACCTGTGTAGTGGAGAATGAATA CGGGTCCATCAATCACACGTACCACTGGATGTTGTGGAGCGATCGCTCACCAGGCC CATCTCCAAGCGGACTGCGGCAAAATGCCTCCACAGTGGTGGAGGAGAGCTAGA GTTTGTCTGCAAGGTTTACAGTGATGCCAGCCCCACATCCAGTGGATCAAGCACGT GGAAAAGAACGGCAGTAAATACGGGCCCGACGGGCTGCCCTACCTCAAGGTTCTCAA GGTTTCGGCTGAGTCCAGCTCCTCATGAACCTCAACACCCCGCTGGTGAGGATAAC AACACGCTCTCTTCAACGGCAGACACCCCATGCTGGCAGGGGTCTCCGAGTATGA ACTTCCAGAGGACCAAAATGGGAGTTTCAAGAGATAAGCTGACACTGGGCAAGCC CCTGGGAGAAGGTTGCTTTGGGCAAGTGGTCATGGCGAAGCAGTGGGAATTGACAA AGACAAGCCCAAGGAGGCGGTACCGTGGCCGTGAAGATGTGAAAGATGATGCCAC AGAGAAAGACCTTTCTGATCTGGTGTGAGAGATGGAGATGATGAAGATGATTGGGAA ACACAAGAATATCATAAATCTTCTTGAGCCTGCACACAGGATGGGCTCTCTATGT CATAGTTGAGTATGCCTCTAAAGGCAACCTCCGAGAATACCTCCGAGCCCGGAGGCC ACCCGGGATGGAGTACTCTATGACATTAAACCGTGTCTCTGAGGAGCAGATGACCTT CAAGGACTTGGTGTATGCACCTTACCAGCTGGCCAGAGGATGGAGTACTTGGCTTC CCAAAATGTATTATCGAGATTAGCAGCCAGAAATGTTTGGTAACAGAAAACAA TGATGATGAAAATAGCAGACTTTGGACTCGCCAGAGATATCAACAATATAGACTATTA CAAAAAGACCACCAATGGGCGGCTTCAGTCAAGTGGATGGCTCCAGAAGCCCTGTT TGATAGAGTATACACTCATCAGAGTGTCTGGTCTTCGGGTGTTAATGTGGGA

TABLE 5-continued

FGFR sequences.		
SEQ ID NO: 32	FGFR2 isoform 7 Amino acid sequence	GATCTTCACCTTTAGGGGGCTCGCCCTACCCAGGGATTCCCGTGGAGGAACCTTTTAA GCTGCTGAAGGAAGGACACAGAATGGATAAGCCAGCCAACTGCACCAACGAACGTGA CATGATGATGAGGACTGTTGGCATGCAGTGCCCTCCAGAGACCAACGTTCAGCA GTTGGTAGAAGACTTGGATCGAATTCCTACTCTCACAACCAATGAGGAATACTTGGGA CCTCAGCCAACCTCTCGAACAGTATTACCTAGTTACCTTGACACAGAAGTTCTTG TTCTTCAGGAGATGATTCTGTTTTTCTCCAGACCCCATGCCTTACGAACCATGCCT TCCTCAGTATCCACACATAAACCGCAGTGTTAAACATGAATGACTGTGTCTGCCTG TCCCAACAGGACAGCACTGGGAACCTAGCTACACTGAGCAGGAGACCATGCCTC CCAGAGCTTGTGTCTCCACTTGTATATATGGATCAGAGGAGTAAATAATTGGAAAA GTAATCAGCATATGTGTAAGATTATACAGTTGAAAACCTGTAATCTTCCCCAGGA GGAGAAGAAGTTTCTGGAGCAGTGGACTGCCACAAGCCACCATGTAACCCCTCTCA CCTGCCGTGCGTACTGGCTGTGGACCAGTAGGACTCAAGGTGGACGTGCGTTCTGCC TTCTTGTGAATTTTGAATAATTGGAGAAGATTATGTGAGCACACACTTACAGAG CACAAATGCAGTATATAGGTGCTGGATGTATGTAATATATTCAAATTATGTATAAA TATATATTATATATTACAAGGAGTTATTTTTTGTATTGATTTAAATGGATGTCCC AATGCACCTAGAAAAATGGTCTCTCTTTTTTAAATAGCTATTGCTAAATGCTGTTC TTACACATAATTTCTTAATTTTACCGAGCAGAGGTGAAAAATACTTTTGGCTTTCA GGAAAAATGTTATAACGTTAATTTATTAATAAATTGGTAATATACAAAAACA
SEQ ID NO: 33	FGFR2 isoform 8 Nucleic acid sequence	MVSWGRFICLVVMTATLSLARPSFSLVEDTTLEPEEPPTKYQISQPEVYVAAPGES LEVRCLLKDAAVISWTKDGVHLGPNNRTVLIGEYLQIKGATPRDSGLYACTASRTVD SETWYFMVNVTDAISSGDEDDTDGAEDFVSSENSNNKRAPYWTNTEKMEKRLHAVPA ANTVKFRCPAGGNPMPMTMRWLKNGKEFKQEHRIGGYKVRNQHWSLINESVVPDKN YTCVENEYGSINHTYHLDVVERSPhRPI LQAGLPANASTVVGDDVEFVCKVYSDAQ PHIQWIKHVEKNGSKYGPDLPLKVLKVAESSSSMSNSNTPLVRIITRSLSTADTP MLAGVSEYELFEDPKWEFPRDKLTGKPLGEGCFQVVMABEAVGIDKDKPEAVTVA VKMLKDDATEKDLSDLVSEMEMMKMI GKHKNI INLLGACTQDGPLYVIVEYASKGNL REYLRRRRPPGMEYSYDINRVPEEQMTFKDLVSCTYQLARGMEYLASQKCIHRDLAA RNVLVTENNVMKIADFGIARDINNIDYKKTNTNRLPVKWMABEALFDRVYTHQSDV WSFGVLMWEIFTLGGSPYPGIPVEELFKLLKEGHRMDKPANCTNELYMMMRDCWHAV PSQRPTFKQLVEDLDRLITLTITNEEYLDLSQPLEQYSPSPYDTRSSCSSGDDSVFSP DPMPEPCLPQYPHINGSVKT GAGCGGGCGAGGGAGCGCGCGCGGCCCAAGCTCGGGCGCGCGGGGCTGCAT CGCGCGTACCTGGCCCGCGCGCGCACTGCTCTCCGGGCTGGCGGGGCGCGCGCG AGCCCCGGGGGCCCCGAGGCGCGAGCTTGCTGCGCGCTCTGAGCCTTCGCAACTCG CGAGCAAAGTTTGGTGGAGGCAACGCCAAGCCTGAGTCTCTTCTCTCTCGTTCC CAAAATCCGAGGCGAGCCCGCGGGCGTATGCCCCGCGCTCCTCCGAGCCTGGGGTAC GCGTGAAGCCCGGAGGCTTGGCGCCGGCGAAGACCAAGGACCACTCTTCTGCGTT TGGAGTTGCTCCCGCAACCCCGGGCTCGTCGCTTTCTCCATCCCGACCCACGCGGG GCGCGGGGCAACACAGGTGCGCGAGGAGCGTTGCCATTCAAGTGACTGCAGCAGCA GCGGCAGCGCCTCGGTTCTGAGCCACCGCAGGCTGAAGGCATTGCGCGTAGTCCA TGCCCGTAGAGGAAGTGTGCAGATGGGATTAACGTCCACATGGAGATATGGAAGAGG ACCGGGATTTGGTACCGTAACCATGGTCAGCTGGGGTCTGTTTATCTGCGCTGGTCTG GGTCACCATGGCAACCTTGTCTTGGCCCGGCCCTCTTCAAGTTAGTTGAGGATAC CACATTAGAGCCAGAAGAGCCACCAACCAATACCAAAATCTCTCAACCAGAAGTGTA CGTGGCTGCGCCAGGGGAGTGTGATAGAGTGGCTGCGTCTGTTGAAAGATGCCCGGT GATCAGTTGGACTAAGGATGGGGTGCCTTGGGGCCCAACAAAGGACAGTGTCTTAT TGGGGGAGTACTTGAGATAAAGGGCGCCACGCCCTAGAGACTCCGGCTCTATGCTTG TACTGCCAGTAGGACTGTAGACAGTGAACCTTGGTACTTCAAGTGATGTGACAGCA TGCCATCTCATCCGAGATGATGAGGATGACACCGATGGTGGCGAAGATTTGTGAG TGAGAACAGTAACAAAGAGAGCACCATACTGGACCAACACAGAAAGATGGAAAA GCGGCTCCATGCTGTGCTGCGGCCAACACTGTCAAGTTTTCGCTGCCAGCCGGGGG GAACCAATGCCAACCATGCGGTGGCTGAAAAACGGGAAGGAGTTTAAAGCAGGAGCA TCGCATTGGAGGCTACAAGGTACGAAACAGCAGCTGGAGCCTCATTATGGAAGTGT GGTCCCATCTGACAAAGGAAATATACCTGTGTAGTGGAGAATGAATACGGGTCCAT CAATCACACGTACACCTGGATGTTGTGGAGCGATCGCCTCACCGGCCCATCTCCA AGCCGGACTGCCGCAAAATGCCCTCCACAGTGGTGGAGGAGAGCTAGAGTTTGTCTG CAAGGTTTACAGTGATGCCAGCCCCACATCCAGTGGATCAAGCACGTGGAAAAAGAA CGGCAGTAAATACGGGCCCGACGGGCTGCCCTACCTCAAGGTTCTCAAGCACTCGGG GATAAATAGTTCCAATGCAGAAGTGCTGGCTCTGTTCAATGTGACCGAGGCGGATGC TGGGGAATATATATGTAAGGTCTCCAATTATATAGGGCAGGCCAACAGCTCTGCTG GCTCACTGTCTTGCCAAAACAGCAAGCGCTGGAAGAGAAAAGGAGATTACAGCTTC CCCAGACTACCTGGAGATAGCCATTACTGCATAGGGGTCTTCTTAATCGCTGTAT GGTGGTAACAGTCACTCTGTGCGGAATGAAGAACACGACCAAGAAGCAGACTTCAG CAGCCAGCCGGCTGTGCACAGCTGACCAACGTATCCCCCTGCGGAGACAGGTAAC AGTTTCGGCTGAGTCCAGCTCCTCCATGAACCTCAACACCCCGCTGGTGGGATAAC AACACGCTCTCTTCAACGGCAGACACCCCATGCTGGCAGGGGTCTCCGAGTATGA ACTTCCAGAGGACCAAAATGGGAGTTTCCAAGAGATAAGCTGACACTGGGCAAGCC CCTGGGAGAAGGTTGCTTTGGGCAAGTGGTCATGGCGGAAGCAGTGGGAATTGACAA AGACAAGCCAAAGGAGGCGGTACCGTGGCCGTGAAGATGTTGAAAGATGATGCCAC AGAGAAAGACCTTTCTGATCTGGTGTGAGAGTGGAGATGATGAAGATGATTGGGAA ACACAAGAATATCATAAATCTTCTTGGAGCCTGCACACAGGATGGGCTCTCTATGT CATAGTTGAGTATGCTCTAAAGGCAACCTCCGAGAATACCTCCGAGCCCGGAGGCC ACCCGGGATGGAGTACTCTATGACATTAAACCGTGTCTCTGAGGAGCAGATGACCTT CAAGGACTTGGTGTATGCACCTACCAGCTGGCCAGAGGATGGAGTACTTGGCTTC CCAAAAATGTATTCATCGAGATTAGCAGCCAGAAATGTTTTGGTAACAGAAAAACA TGTGATGAAAATAGCAGACTTTGGACTCGCCAGAGATATCAACATATAGACTATTA

TABLE 5-continued

FGFR sequences.	
SEQ ID NO: 34	FGFR2 isoform 8 Amino acid sequence CAAAAAGACCACCAATGGGCGGCTTCAGTCAAGTGGATGGCTCCAGAAGCCCTGTT TGATAGAGTATACACTCATCAGAGTGATGCTGGTCCCTCGGGGTGTTAATGTGGGA GATCTTCACTTTAGGGGGCTCGCCCTACCCAGGGATTCCCGTGGAGGAACCTTTTAA GCTGCTGAAGGAAGGACACAGAATGGATAAGCCAGCCAACTGCACCAACGAAGTGA CATGATGATGAGGGACTGTTGGCATGCAGTGCCCTCCAGAGACCAACGTTCAAGCA GTTGGTAGAAGACTTGGATCGAATCTCACTCTCACAAACCAATGAGGAATACTTGGGA CCTCAGCCAACCTCTCGAACAGTATTACCTAGTTACCTTGACACAAGAAGTTCTTG TTCTTCAGGAGATGATTCTGTTTTTCTCCAGACCCCATGCCTTACGAACCATGCCT TCCTCAGTATCCACACATAAACGGCAGTGTTAAACATGA MVSWGRFICLVVVTMATLSLARPSFSLVEDTTLEPEEPPTYKYQISQPEVYVAAPGES LEVRCLLKDAAVISWTKDGVHLGPNNRTVLIGEYLQIKGATPRDSGLYACTASRTVD SETWYFMVNVDAISSGDEDDTDGAEDFVSENSNNKRAPYWTNTEKMEKRLHAVPA ANTVKFRCPAGGNPMPMTMRWLKNGKEFKQEHRI GGYKVRNQHWSLIMESVVPD KGN YTCVVENEYGS INHTYHLDVVERS PHRPILQAGLPANASTVVGGDVEFVCKVYSDAQ PHIQWIKHVEKNGSKYGPDLPLYLKV LKHS GINSSNAEVLALFNVTEADAGEYI CKV SNYIGQANQSAWLT VLPKQQAPGREKEI TASPDYLEIAI YCIGVFLIACMVTVILC RMKNTTKKPDFSSQPAVHKLTKRIPLRRQVTVSAESSSSMNSNTPLVRITRSLSTA DTPMLAGVSEYELPEDPKWEFPRDKLTLGKPLGEGCFGQVVM AEAVGIDKDKPKEAV TVAVKMLKDDATEKDLSDLVSEMEMMKMIGKHKNII NLLGACTQDGPLYV IVEYASK GNLREYLRARRPP GMEYSYD INRVPEEQMTFKDLVSC TYQLARGMEYLA SQKCIHRD LAARNVLVTENNVMKIADFG LARD INNI DYYKKTNGRLPVKWM APEALFDRVYTHQ SDVWSFGVLWMEI FTLGGSPYPGIPVEELFKLLKEGHRMDK PANCTNELYMMMRDCW HAVPSQRPTFKQLVEDLDRILTLTNEEYLDLSQPLEQYSPSPYDTRSSCSSGDDSV FSPDPMPIEPLCPQYPHINGSVKT
SEQ ID NO: 35	FGFR2 isoform 9 Nucleic acid sequence GAGAGCCGAGCGGGCGGGCGGGTGCAGAGCGGGCGAGGGAGCGCGCGCGGCGCG CACAAAGCTCGGGCGCGCGGGGCTGCATGCGCGGTACCTGGCCCGGCGCGGCGACT GCTCTCGGGCTGGCGGGGCGCGGCCGAGCCCGGGGGCCCCGAGGCCGCGAGCTT GCCTGCGCGCTCTGAGCCTTCGCAACTCGCGAGCAAAGTTTGGTGGAGGCAACGCCA AGCCTGAGTCTCTTCTCTCTCGTTCCTCCAAATCCGAGGGCAGCCCGCGGGCGTCA TGCCCGCGCTCTCTCCGAGCCTGGGGTACGCGTGAAGCCCGGAGGCTTGGCGCCGG CGAAGACCCAAAGGACCACTCTTCTGCGTTTGGAGTTGCTCCCGCAACCCCGGCTC GTCGCTTCTCCATCCCGACCCACGCGGGGCGCGGGGACAAACAGGTGCGGGAGGA CGCTTGCCATTCAAGTGA CTG CAGCAGCAGCGCAGCGCCTCGGTTCTGAGGCCAC CGCAGGCTGAAGGCATTGCGCGTAGTCCATGCCCGTAGAGGAAGTGTGCAGATGGGA TTAACGTCCACATGGAGATATGGAAGAGGACCGGGGATTGGTACCGTAACCATGGTC AGCTGGGGTCTTTTCATCTGCTTGGTCTGCTGCTCACCATGGCAACCTTGTCCTGGCC CGGCCCTCTTCAGTTTGTAGTGAGGATACCACTTAGAGCCAGAAGATGCCATCTCA TCCGGAGATGATGAGGATGACCCGATGGTGGGGAAGATTTTGTAGTGAGAACAGT AACAAAGAGAGACCACTACTGGACCAACACAGAAAAGATGGAAAAGCGGCTCCAT GCTGTGCTGCGGCAACACTGTCAAGTTTCTGCTGCCAGCCGGGGGGAACCCAAATG CCAACCATGCGGTGGCTGAAAACCGGAAGGAGTTTAAGCAGGAGCATCGCATTGGA GGCTACAAAGGTACGAAACCGCACTGGAGCCTCATTATGGAAAGTGTGGTCCCATCT GACAAGGGAATTATACCTGTGTAGTGAGGAATGAATACGGGTCCATCAATCACACG TACCACCTGGATGTTGTGAGCGATCGCCTCACCGGCCCATCTCCAAGCCGGA CTG CCGGCAAATGCTCCACAGTGGTCGGAGGAGACGTAGAGTTTGTCTGCAAGGTTTAC AGTGATGCCAGCCCACTCCAGTGGATCAAGCACGTGAAAAGAACCGGCAGTAAA TACGGGCCCCGACGGGCTGCCTACCTCAAGGTTCTCAAGCACTCGGGGATAAATAGT TCCAATGCAGAAGTGTGGCTCTGTTCAATGTGACCGAGGCGGATGCTGGGGAATAT ATATGTAAGGTCTCCAATTATATAGGGCAGGCCAACCAAGTCTGCCTGGCTCACTGTC CTGCCAAAACAGCAAGCGCCTGGAAGAGAAAAGGAGATTACAGCTTCCCCAGACTAC CTGGAGATAGCACTTTACTGCATAGGGTCTTCTTAATCGCCTGTATGGTGGTAACA GTCACTCTGTGCCAATGAAGAACACGACCAAGAAGCCAGACTTCAGCAGCCAGCCG GCTGTGCACAAGCTGACCAACGTATCCCCCTGCGGAGACAGGTAACAGTTTCGGCT GAGTCCAGCTCTCCATGAACCTCAACACCCCGCTGGTGAGGATAACCAACACGCCTC TCTTCAACGGCAGACACCCCATGCTGGCAGGGGTCTCCGAGTATGAACCTCCAGAG GACCCAAAATGGGAGTTTCCAAGAGATAAGCTGACACTGGGCAAGCCCCCTGGGAGAA GGTTGCTTTGGCAAGTGGTCATGGCGGAAGCAGTGGGAATTGACAAAGCAAGCCC AAGGAGGCGGTCAACGTGGCCGTGAAGATGTTGAAAGATGATGCCACAGAGAAAGAC CTTCTGATCTGGTGTGAGAGATGAGATGAAGATGATTGGGAAACACAAGAAT ATCATAAATCTTCTGGAGCCTGCACACAGGATGGGCCTCTCTATGTCAATAGTTGAG TATGCCCTCTAAAGGCAACCTCCGAGAATACCTCCGAGCCCGGAGGCCACCCGGGATG GAGTACTCTATGACATTAAACCGTGTCTCTGAGGAGCAGATGACCTTCAAGGACTTG GTGTCATGCACCTACCACTGGCCAGAGGATGGAGTACTTGGCTTCCCCAAAATAGT ATTCATCGAGATTAGCAGCCAGAAATGTTTTGGTAACAGAAACCAATGTGATGAAA ATAGCAGACTTTGGA CTG CCGCAGAGATATCAACAATATAGACTATTACAAAAAGACC ACCAATGGGCGGCTTCCAGTCAAGTGGATGGCTCCAGAAGCCCTGTTTGATAGAGTA TACACTCATCAGAGTGATGTCTGGTCTCTCGGGGTGTTAATGTGGGAGATCTTCACT TTAGGGGGCTCGCCCTACCCAGGATTCCTCGTGAGGAACCTTTTAAAGCTGCTGAAG GAAGGACACAGAAATGGATAAGCCAGCCAACTGCACCAACGAAGTGTACATGATGATG AGGGACTGTTGGCATGCAGTGCCCTCCAGAGACCAACGTTCAAGCAGTTGGTAGAA GACTTGGATCGAATCTCACTCTCACAAACCAATGAGATCTGAAAGTTTATGGCTTCA TTGAGAACTGGGAAAAGTTGGTCAGGCGCAGTGGCTCATGCCTGTAATCCAGCAC TTTGGGAGGCCGAGGCAGGCGGATCATGAGGTGAGGAGTTCCAGACCAGCCTGGCCA ACATGGTGAAACCTGTCTCTACTAAAGATACAAAAAATTAGCCGGGCGTGTGGTG TGCACCTGTAATCCAGCTACTCCGGAGGCTGAGGCAGGAGAGTCACTTGAACCGG GGAGGCGGAGGTTGAGTGAGCCGAGATCATGCCATTGCATTCCAGCCTTGGCGACA

TABLE 5-continued

FGFR sequences.		
SEQ ID NO: 36	FGFR2 isoform 9 Amino acid sequence	GAGCGAGACTCCGTCTCAAAAAAAAAATAAAAA MVSWGRFICLVVVTMATLSLARPSFSLVEDTTLEPEDAISSGDEDDTDGAEDFVSE NSNNKRAPYWTNTEKMEKRLHAVPAANTVKFRCPAGGNPMPMTMRWLKNGKEFKQEHR IGGYKVRNQHWSLIMESVVPDSDKGNVTCVENEYGSINHTYHLDVVERSPHRPILQA GLPANASTVVGDDVEFVKVYSDAQPHIQWIKHVEKNKSKYGPDGLPYLKV LKHSGI NSSNAEVLALFNVTEADAGEYICKVSNYIGQANQSAWLTVL PKQQA PGREKEITASP DYLEIAIYICIGVFLIACMVTVILCRMKNNTKKPDFSSQPAVHKLT KRIPLRRQVT SAESSSSMNSNTPLVRITRLSSTADTPMLAGVSEYELPEDPKWEFPRDKLT LKGL GEGCFGQVVM AEAVGIDKDKPKEAVTVAVKMLKDDATEKDLSDLVSEMEMMKMIGKH KNIIINLLGACTQDGPLYVIVEYASKGNLREYLRARRPPGMEYSYDINRVPEEQMTFK DLVSTYQLARGMEYLASQKCIHRDLAARNVLVTENNVMKIADFLGLARDINNIDYK KTTNGRLPVKWM APEALFDRVYTHQSDVWSFGVLMWEIFTLGGSPYPGIPVEELFKL LKEGHRMDKPANCTNELYMMMRDCWHAVPSQRPTFKQLVEDLDRILTLTTNEI
SEQ ID NO: 37	FGFR2 isoform 10 Nucleic acid sequence	GGACCGGGGATGGTACCGTAACCATGGTCAGCTGGGGTCGTTTCATCTGCCTGGTC GTGGTCACCATGGCAACCTTGTCCCTGGCCCGGCCCTCCTTCAGTTAGTTGAGGAT ACCATAGTAGGCCAGAGAGCCACCAACCAATACCAATCTCTCAACAGAGAGTG TACGTGGCTGCGCCAGGGGAGTCGCTAGAGGTGCGCTGCCTGTTGAAGATGCCGCC GTGATCAGTTGGACTAAGGATGGGGTGCACTTGGGGCCCAACATAGGACAGTGCTT ATTGGGGAGTACTTGACAGATAAGGGCGCCACGCCTAGAGACTCCGCGCTCTATGCT TGTAAGTCCAGTAGGACTGTAGACAGTGAAACTTGGTACTTCATGGTGAATGTCACA GATGCCATCTCATCCGAGATGATGAGGATGACACCGATGGTGCGGAAGATTTTGTG AGTGAGAACAGTAACAACAGAGAGCACCATACTGGACCAACACAGAAAAGATGGAA AAGCGGCTCCATGCTGTGCCTGCGGCCCAACACTGTCAAGTTTCGCTGCCAGCCGGG GGGAACCAATGCCAACCATGCGGTGGCTGAAAAACGGGAAGGAGTTTAAGCAGGAG CATCGCATTTGGAGGCTACAAGGTACGAAACAGCACTGGAGCCTCATTTGGAAGT GTGGTCCCATCTGACAAGGGAAATATACCTGTGTAGTGAGGAATGAATACGGGTCC ATCAATCACACGTACACCTGGATGTTGTGGAGCGATCGCCTCACCGGCCATCCTC CAAGCCGACTGCGCGCAATGCTCCACAGTGGTCGGAGGAGACGTAGAGTTTGTG TGCAAGGTTTACAGTGATGCCAGCCACATCCAGTGGATCAAGCACGTGGAAAAAG AACGGCAGTAAATACGGGCCGACGGGCTGCCCTACCTCAAGGTTCTCAAGCACTCG GGGATAAATAGTTCCAATGCGAAGTGTCTGGCTCTGTTCAATGTGACCGAGGCGGAT GCTGGGGAATATATATGTAAGGCTCCCAATATATAGGGCAGGCCAACAGTCTGCC TGGCTCACTGTCTGCCAAACAGCAAGCGCCTGGAAGAGAAAGGAGATTACAGCT TCCCAGACTACCTGGAGATAGCCATTACTGCATAGGGGCTCTCTTAATCGCTGT ATGGTGGTAACAGTCATCCTGTGCGCAATGAAGAACACGACCAAGAGCCAGACTTC AGCAGCCAGCCGGCTGTGCACAAGCTGACCAACGATATCCCCCTGCGGAGACAGGTA ACAGTTTCGGCTGAGTCCAGCTCCTCCATGAATCCCAACACCCCGCTGGTGAGGATA ACAACACGCCTCTCTTCAACGGCAGACACCCCCATGCTGGCAGGGGCTCTCCGAGTAT GAACCTCCAGAGGACCAAAATGGGAGTTTCCAAGAGATAAGCTGACACTGGGCAAG CCCCCTGGGAGAAGGTTGCTTTGGGCAAGTGGTCATGGCGGAAGCAGTGGGAATTGAC AAAGACAAGCCCAAGGAGGCGGTACCGTGGCCGTGAAGATGTTGAAGATGATGCC ACAGAGAAAGACCTTTCTGATCTGGTGTGAGAGATGGAGATGATGAAGATGATTGGG AAACACAAGAAATATCATAAATCTTCTTGGAGCCTGCACAGGATGGGCTCTCTAT GTACATAGTTGAGTATGCCTCTAAGGCAACCTCCGAGAATACCTCCGAGCCCGGAGG CCACCCGGGATGGAGTACTCTATGACATTAACCGTGTCTCTGAGGAGCAGATGACC TTCAAGGACTTTGGTGTATGCACCTACCAGCTGGCCAGAGGCATGGAGTACTTGGCT TCCCAAAAATGTATTATCGAGATTTAGCAGCCAGAAATGTTTGGTAACAGAAAAC AATGTGATGAAAATAGCAGACTTTGGACTCGCCAGAGATATCAACATATAGACTAT TACAAAAGACCACCAATGGGCGGCTTCCAGTCAAGTGGATGGCTCCAGAGCCCTG TTTGATAGAGTATACACTCATCAGAGTGTGCTGGTCTTCCGGGGTGTAAATGTGG GAGATCTTCACTTTAGGGGGCTCGCCCTACCCAGGGATTCCCGTGGAGGAACCTTT AAGCTGTGAAGGAAGGACACAGAAATGGATAAGCCAGCCAACTGCACCAACGAACG TACATGATGATGAGGGACTGTTGGCATGCAGTGCCTCCAGAGACCAACGTTCAAG CAGTTGGTAGAAGACTTGGATCGAATTCCTACTCTCACAACCAATGAGATCTGAAAG TTTATGGCTTCATTGAGAACTGGGAAAAGTTGGTCAGGCGCAGTGGCTCATGCCTG TAATCCAGCACTTTGGGAGGCGGAGCAGGCGGATCATGAGGTCAGGAGTTCCAGA CCAGCCTGGCCAACATGGTGAAACCTGTCTCTACTAAAGATACAAAAAATTAGCCG GGCGTGTGGTGTGCACCTGTAATCCAGCTACTCCGGGAGGCTGAGGCAGGAGAGT CACTTGAAACCGGGGAGGCGGAGGTGTCAGTGAGCCGAGATCATGCCATTGCATTCCA GCCTTGGCGACAGAGCGAGACTCCGTCT
SEQ ID NO: 38	FGFR2 isoform 10 Amino acid sequence	MVSWGRFICLVVVTMATLSLARPSFSLVEDTTLEPEEPPTKYQISQPEVYVAAPGES LEVRCLLKDAAVISWTKDGVHLGPNNRTVLI GEYLQIKGATPRDSGLYACTASRTVD SETWYFMVNVTDAISSGDEDDTDGAEDFVSENSNNKRAPYWTNTEKMEKRLHAVPA ANTVKFRCPAGGNPMPMTMRWLKNGKEFKQEHRIGGYKVRNQHWSLIMESVVPDSDKN YTCVENEYGSINHTYHLDVVERSPHRPILQAGLPANASTVVGDDVEFVKVYSDAQ PHIQWIKHVEKNKSKYGPDGLPYLKV LKHSGINSSNAEVLALFNVTEADAGEYICKV SNYIGQANQSAWLTVL PKQQA PGREKEITASPDYLEIAIYICIGVFLIACMVTVILC RMKNNTKKPDFSSQPAVHKLT KRIPLRRQVTSAESSSSMNSNTPLVRITRLSSTA DTPMLAGVSEYELPEDPKWEFPRDKLT LKGLGEGCFGQVVM AEAVGIDKDKPKEAV TVAVKMLKDDATEKDLSDLVSEMEMMKMIGKHKNIIINLLGACTQDGPLYVIVEYASK GNLREYLRARRPPGMEYSYDINRVPEEQMTFKDLVSTYQLARGMEYLASQKCIHRD LAARNVLVTENNVMKIADFLGLARDINNIDYKTTNGRLPVKWM APEALFDRVYTHQ SDVWSFGVLMWEIFTLGGSPYPGIPVEELFKLKEGHRMDKPANCTNELYMMMRDCW HAVPSQRPTFKQLVEDLDRILTLTTNEI

TABLE 5-continued

FGFR sequences.		
SEQ ID NO: 39	FGFR2 isoform 11 Nucleic acid sequence	GGTACCGTAACCATGGTCAGCTGGGGTCGTTTCATCTGCCTGGTCGTGGTCACCATG GCAACCTTGTCCTTGGCCCGGCCCTCCTTCAGTTTAGTTGAGGATACACATTAGAG CCAGAAGAGCCACCAACCAATACCAATCTCTCAACCAGAAGTGTACGTGGCTGCG CCAGGGGAGTCGCTAGAGGTGCGCTGCGCTGTTGAAAGATGCCGCCGTGATCAGTTGG ACTAAGGATGGGGTGCACCTTGGGGCCCAACAATAGGACAGTGTCTATTGGGGAGTAC TTGCAGATAAAGGGGCCACGCCCTAGAGACTCCGGCCTCTATGCTTGTACTGCCAGT AGGACTGTAGACAGTGAACCTTGGTACTTCATGGTGAATGTACAGATGCCATCTCA TCCGGAGATGATGAGGATGACACCGATGGTGCAGGAAGATTTTGTCAAGTGAACAGT AACCAACAGAGAGCACCATACTGGACCAACAGAGAAAGATGGAAAAGCGGCTCCAT GCTGTGCTGCGGCCAACACTGTCAAGTTTCGCTGCCCCAGCCGGGGGGAACCCCAATG CCAACCATGCGGTGGCTGAAAAACGGGAAGGAGTTTAAAGCAGGAGCATCGCATTTGA GGCTACAGGTACGAAACAGCACTGGAGCCTCATTATGAAAGTGTGGTCCCCTCT GACAAGGGAAATTATACCTGTGTAGTGAGGAATGAATACGGGTCATCAATCACACG TACCACCTGGATGTTGTGGAGCGATCGCCTCACCAGGCCATCCTCCAAGCCGAGTGC CCGCAATGCCTCCACAGTGGTCGGAGGAGACGTAGAGTTGTCTGCAAGGTTTAC AGTGATGCCAGCCCCACATCCAGTGGATCAAGCACGTGGAAAAGAACCGGAGTAAA TACGGGCCCGAGGGCTGCCCTACCTCAAGTTCTCAAGGCCCGCCGTTTAAACACC ACGGACAAAGAGATTGAGGTCTCTATATTGGAATGTAACCTTTGAGGACGCTGGG GAATATACGTGCTTGGCGGGTAATTTCTATTGGGATATCCTTTCACTCTGCATGGTTG ACAGTTCTGCCAGCGCCTGGAAGAGAAAAGGAGATTACAGCTTCCCAGACTACCTG GAGATAGCCATTTACTGCATAGGGGTCTTCTTAATCGCCTGTATGGTGGTAAACAGTC ATCCTGTGCGGAATGAAGAACACGACCAAGAAGCCAGACTTCAGCAGCCAGCCGGCT GTGCACAAGCTGACCAACGTATCCCCCTGCGGAGACAGGTTTCCGGCTGAGTCCAGC TCCTCCATGAACTCCAACACCCCGCTGGTGAGGATAACAACACGCCCTCTCTTCAACG GCAGACACCCCCATGTGCGCAGGGGTCTCCGAGTATGAACTTCCAGAGGACCCAAAA TGGGAGTTTCCAAGAGATAAGCTGACACTGGGCAAGCCCTGGGAGAAGGTTGCTTT GGGCAAGTGGTCTATGGCGGAAGCAGTGGGAATTGACAAAGACAGCCCAAGGAGGCG GTCAACCGTGGCCGTGAAGATGTTGAAAGATGATGCCACAGAGAAAGACCTTTCTGAT CTGGTGTACAGATGGAGATGATGAAGATGATTGGGAAACACAAGATATCATAAAT CTTCTTGGAGCTGCACACAGGATGGGCCTCTCTATGTCATAGTTGAGTATGCCTCT AAAGGCAACCTCCGAGAATACCTCCGAGCCCGGAGGCCACCCGGGATGGAGTACTCC TATGACATTAAACCGTGTCTGAGGAGCAGATGACCTTCAAGGACTTGGTGTATGCG ACCTACAGCTGGCCAGAGGCATGGAGTACTTGGCTTCCCAAAATGTATTCATCGA GATTTAGCAGCCAGAAATGTTTGGTAACAGAAAACAATGTGATGAAAATAGCAGAC TTTGGACTCGCCAGAGATATCAACAAATAGACTATTACAAAAGACCAACCAATGGG CGGCTTCCAGTCAAGTGGATGGCTCCAGAAGCCCTGTTGATAGAGTATACATCAT CAGAGTGTCTGCTGCTCGGGGTGTTAATGTGGGAGATCTTCACTTTAGGGGGC TCGCCCTACCCAGGGATTCCCGTGGAGGAACCTTTTAAGCTGCTGAAGGAAGGACAC AGAATGGATAAGCCAGCCAACCTGCACCAACGAAGTGTACATGATGATGAGGGACTGT TGGCATGCAGTGCCTCCAGAGACCAACGTTCAAGCAGTTGGTAGAAGACTTGGAT CGAATTCTACTCTCAACAACCAATGAGGAATACTTGGACCTCAGCCCAACCTCTCGAA CAGTATTACCTAGTTACCTGCACACAAGAGTCTTGTCTTTCAGGAGATGATTCT GTTTTTTCTCCAGACCCCATGCCTTACGAACCATGCCTTCTCAGTATCCACACATA AACGCGAGTGTAAACATGA
SEQ ID NO: 40	FGFR2 isoform 11 Amino acid sequence	MVSWGRFICLVVVTMATLSLARPSFSLVEDTTLEPEEPPTKYQISQPEVYVAAPGES LEVRCLLKDAAVISWTKDGVHLGPNNRVTLIGELYLQIKGATPRDSGLYACTASRTVD SETWYFMVNVTDIAISGGDEDDTDGAEDFVSENSNNKRAPYWTNTEKMEKRLHAVPA ANTVKFRCPAGGNPMPMTMRWLKNGKEFKQEHRIIGYKVRNQHWSLIMESVVPDKN YTCVENEYGSINHTYHLDVVERSPhRPILQAGLPANASTVVGDDVEFVKVYSDAQ PHIQWIKHVEKNGSKYGPDLPLYLVKLVKAAGVNTDKEIEVLYIRNVPFDAGEYTC LAGNSIGISFHSAWLTVLPAPGREKEITASPDYLEIAIYICGVFLIACMVVTVILCR MKNTTKKPDFSSQPAVHKLTKRIPLRQVSAESSSSMNSNTPLVIRITRLSSADTP MLAGVSEYELFEDPKWEFPRDKLTGKPLGEGCFGQVMAEAVGIDKDKPKEAVTVA VKMLKDDATEKDLSDLVSEMEMMKMIGKHKNIINLLGACTQDGPLYVIVEYASKGNL REYLRARRPPGMEYSYDINRVPEEQMTFKDLVSTCYQLARGMEYLASQKCIHRDLAA RNVLVTENNVMKIADFLGARDINNIDYKKTTNGRLPVKWMAPALFDRVYTHQSDV WSPFVLMWEIFTLGGSPYPGIPVEELFKLLKEGHRMDKPANCTNELYMMMRDCWHAV PSQRPTFKQLVEDLDRILTLTTNEEYLDLSQPLEQYSPSPYDTRSSCSSGDDSVFSP DPMPYEPCLPQYPHINGSVKT
SEQ ID NO: 41	FGFR3 isoform 1 Nucleic acid sequence	AGTGC CGGTGGCGGCGGCTCGCGGCGAGCTGGCGCGCGCGGTCTCTGCTCGCCG GTCGCACGGACGCACCGCGCGGCGCGCGCGCGGAGGAGCGGGCGGGAGCTGGGCCC GCGGACAGCGAGCGGAGCGGGAGCCGCGGTAGCGAGCGGGCTCCGGCGCTCGCC AGTCTCCGAGCGCGCGCGGCTCCCGCGGTGCGCGCGCGGGCGGTGGGGGCGAG CATGCCGCGCGCTGCTGAGGACGCGCGGCGCGCGCGCGCGCGCGCATGGGCGCCC CTGCTGCGCCCTCGCGCTCTGCGTGGCGGTGGCCATCGTGGCGCGCGCTCTCTCGG AGTCTTGGGAGCGAGCAGCGCGTCTGGGGCGAGCGGAGAGTCCCGGGCCAG AGCCCGGCGAGCAGGAGCAGTTGGTCTTGGCAGCGGGGATGCTGTGGAGCTGAGCT GTCCCCCGCCCGGGGGTGGTCCCATGGGGCCCACTGTCTGGGTCAAGGATGGCACAG GGCTGGTGCCCTCGGAGCGTGTCTGGTGGGGCCCGAGCGGTGCAGGTGTCTGAATG CCTCCACAGGAGCTCCGGGGCTACAGCTGCCGCGAGCGGCTCACGAGCGCGTAC TGTGCCACTTCAGTGTGCGGGTGACAGACGCTCCATCTCGGGAGATGACGAAGACG GGGAGGACGAGGCTGAGGACACAGGTGTGGACACAGGGGCCCTTACTGGACACGGC CCGAGCGGATGGACAAGAAGCTGTGGCGGTGCGCGCGCGCAACACCGTCCGCTTCC GCTGCCAGCCGCTGGCAACCCCACTCCCTCCATCTCTGGCTGAAGAACCGGAGGG AGTTCCCGCGCGAGCACCGCATTGGAGGCATCAAGCTGCGGCATCAGCAGTGGAGCC TGGTCATGGAAGCGTGGTGCCCTCGGACCGGCGCACTACACCTGCGCTGCTGGAGA

TABLE 5-continued

FGFR sequences.		
SEQ ID NO: 42	FGFR3 isoform 1 Amino acid sequence	ACAAGTTTGGCAGCATCCGGCAGACGTACACGCTGGACGTGCTGGAGCGCTCCCCGC ACCGGCCCATCCTGCAGGCGGGGCTGCCGGCCAACCAGACGGCGGTGCTGGGCAGCG ACGTGGAGTTCCACTGCAAGGTGTACAGTGACGCACAGCCCCACATCCAGTGGCTCA AGCACGTGGAGGTGAATGGCAGCAAGGTGGGCCCGGACGGCACACCTACGTTACCG TGCTCAAGACGGCGGGCGCTAACACCACCGACAAGGAGCTAGAGGTTCTCTCCTTGC ACAACTGCACCTTTGAGGACGCCGGGAGTACACCTGCCTGGCGGGCAATCTTATTG GGTTTTCTCATCACTCTGCGTGGCTGGTGGTGTGCTGCCAGCCGAGGAGGAGCTGGTGG AGGCTGACGAGGCGGGCAGTGTGTATGCAGGCATCCTCAGCTACGGGGTGGGCTTCT TCCTGTTTCATCCTGGTGGTGGCGGTGTGACGCTCTGCCGCTGCGCAGCCCCCCCA AGAAAGGCTGGGCTCCCCCACCCTGTCACAAAGATCTCCCGCTTCCCGCTCAAGCGAC AGGTGTCCTGGAGTCCAACGCTCCATGAGCTCCAACACACCACTGGTGCATCG CAAGGCTGTCTCAGGGGAGGGCCCCACGCTGGCCAATGTCTCCGAGCTCGAGCTGC CTGCCGACCCCAATGGGAGCTGTCTCGGGCCCGGTGACCTGGGCAAGCCCCCTTG GGGAGGGCTGCTTCGGCCAGGTGGTTCATGGCGAGGCCATCGGCATTGACAAGGACC GGGCGGCCAAGCCTGTACCGTAGCCGTGAAGATGCTGAAAGACGATGCCACTTGACA AGGACCTGTGCGACCTGGTGTCTGAGATGGAGATGATGAAGATGATCGGGAACACA AAAAATCATCAACCTGCTGGGCGCCTGCACGCAAGGCGGGCCCCCTGACGTGCTGG TGGAGTACGCGGCCAAGGGTAACCTGCGGGAGTTTCTGCGGGCGCGCGGCCCGCG GCCTGGACTACTCTTCGACACCTGCAAGCCGCCGAGGAGCAGCTCACCTTCAAGG ACCTGGTGTCTGTGCTTACCAGGTGGCCCGGGGCATGGAGTACTTGGCTCCCGA AGTGATCCACAGGACCTGGCTGCCCGCAATGTGCTGGTGACCGAGGACCAAGTGA TGAAGATCGCAGACTTCGGGCTGGCCCGGACGTGCACAACTCGACTACTACAAGA AGACGACCAACGGCCGGCTGCCGCTGAAGTGGATGGCGCCTGAGGCTTGTGTTGACC GAGTCTACACTCACCAGAGTGACGTCTGGTCTTGGGGTCTGCTCTGGGAGATCT TCACGCTGGGGGGCTCCCCGTACCCCGGCATCCCTGTGGAGGAGCTCTTCAAGCTGC TGAAGGAGGGCCACCGCATGGACAAGCCCGCCAAGTGCACACACGACCTGTACATGA TCATGCGGGAGTGTGTCATGCCGCGCCCTCCAGAGGGCCACCTTCAAGCAGCTGG TGGAGGACCTGGACCGGTGCTTACCCTGACGCTCCACCGACGAGTACCTGGACCTGT CGGCGCCTTTCGAGCAGTACTCCCGGGTGGCCAGGACACCCCAAGCTCCAGCTCCT CAGGGGACGACTCCGTGTTTGCACGACCTGCTGCCCCCGGCCCCACCCAGCAGTG GGGGCTCGCGGACGTGAAGGGCCACTGGTCCCCAACAATGTGAGGGGTCCCTAGCAG CCCACCTGCTGCTGGTGCACAGCCTCCCGGCATGAGACTCAGTGACAGATGGAG AGACAGCTACACAGAGCTTTGGTCTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGT GTGCACATCCGCGTGTGCTGTGTGCGCATCTTGCCTCCAGGTGCAGAGGTAC CCTGGGTGTCCCCGCTGCTGTGCAACGCTCTCCTGACTGGTGTGCTGACGACCCGAGG GCCTTTGTTCTGGGGGACCCAGTGCAGAAATGTAAGTGGGCCACCCGCTGGGACCC CCGTGGGGCAGGGAGCTGGGCCGACATGGCTCCGGCTCTGCTTTGCACACCGGG ACATCACAGGTTGGGCTCGGCCCTCCACACCCAAAGCTGAGCCTGCAGGGAAGC CCACATGTCCAGCACCTTGTGCTGGGGTGTAGTGGCACCGCCTCCCACTCCA GGCTTTCCCACTTCCCACTTGCCTTCAGAGACTGAAATTACGGGTACCTGAAGAT GGGAGCCTTTACCTTTTATGCAAAAGGTTTATTCGGAAACTAGTGATACATTTCTAT AAATAGATGTGTGTATATGATATATATACATATATATATAACATATATGGAAGA GGAAAGGCTGGTACAACGGAGGCTGCGACCTGGGGGCACAGGAGGACGGCATGG CCCTGGGCGGGGCTGGGGGGCGTGGAGGAGGCCACAGGGGTCTCACCCATGCA AGCAGAGGACACAGGGCCTTTCTGGCACCGCAGTTTGTGTTTAAACTGGACCTGTA TATTTGTAAAGCTATTTATGGGCCCTTGGCACTCTTGTTCACACCCCAACACTTC CAGCATTTAGCTGGCCACATGGCGGAGAGTTTAAATTTTAACTTATGACAACCGA GAAGGTTTATCCGCGCATAGAGGGACGGCCAAGAAATGTACGTCAGCCTGCCCGG AGCTGGAGGATCCCTCCAAGCCTAAAGGTTGTTAATAGTTGGAGGTGATTCAGT GAAGATATTTATTTCTTTGTCTTTTTCAGGAGAATTAGATTTCTATAGGATTTT TCTTTAGGAGATTTATTTTGGACTTCAAGCAAGCTGGTATTTTCATACAAATTC TTCTAATTGCTGTGTGTCCCAGGCAGGGAGACGGTTTCCAGGGAGGGGCCCGCCCTG TGTGACGTTCCGATGTTATAGATGTTACAAGTTTATATATATCTATATATATATA TTATTGAGTTTATTACAAGATGATTTGTGTAGACTTAACACTTCTTACGCAATGCT TCTAGAGTTTATAGCTGGAGTGTACCTTTCAAAGCTTGGAGGGAAGCCGTGAAT TCAGTTGTTGCTTCTGTACTGTTACTGGGCCCTGAGCTGGGCAGCTGTCCCTTGC TTGCTGACGGGCATGGCTCAGGGTGGTCTCTTCTTGGGGCCAGTGCATGTTGGC CAGAGGTGTCACCCAAACCGGCAGGTGCGATTTTGTAAACCGACGACGAACCTTCC GAAAAATAAGACACCTGGTGTCTAACCTGG
SEQ ID NO: 43	FGFR3 isoform 2	MGAPACALALCVAIVAGASSESLGTEQRVVGRAAEVPGPEPGQEQVLVFGSGDAV ELSCPPPGGPMGPTVWVKDGTGLVPSERVLVGPQRLQVLNASHEDSGAYSCRQLT QRVLCHFSSVRVTDAPSSGDDDEGDEAEDTGVDTGAPYWRPERMDKKLLAVPAANT VRFRCPAAGNPTPSISWLKNGREFRGEHRIGGIKLRHQQWSLVMSVVPVSDRGNITC VVENKFGSI RQTYTLDVLEERSPHRPI LQAGLPANQTAVLGSDFEFHKVYSDAQPHI QWLKHVEVNGSKVGPDPYVTVLKTAGANTTDKELEVLSLHNVTTFEDAGEYTCLAG NSIGFSHHSANLVLPAEEELVEADEAGSVYAGILSYGVGFLLFILVVAAVTLCRLR SPPKGLGSPVHKI SRFP LKRQVLSLESNASMSNTPLVRIARLSSGEGPTLANVSE LELPADPKWELSRARLT LKPLGEGCFQGVMAEAI GIDKDRAAKPVTVAVKMLKDD ATDKLSDLVSEMEMMKMI GKHKNI INLLGACTQGGPLYVLVEYAAKGNLREFLRAR RPPGLDYSFDTCKPPEEQLT FKLVS CAYQVARGMEYLAQKCIHRDLAARNVLVTE DNVMKIADPGLARDVHNLDYKKTTNGRLPVKWMAPALFDRVYTHQSDVWSFGVLL WEIFTLGGSPYPGIPVEELFKLKEGHRMDKPA NCTHDL YMIMRECWHAPSQRPTF KQLVEDLDRVLT VTS TDEYLDLSAPFEQYSPGGQDTPSSSSSGDSDVFAHDL LPPAP PSSGGSRT
SEQ ID NO: 43	FGFR3 isoform 2	GTCCGCGGCAGCTGGCGCCGCGCGGTCTGCTCTGCCGGTCCGACGACGACCCGGC GGGCGCCGGCCGGAGGACGGGCGGGAGCTGGGCCCGCGGACAGCAGCAGCCGGAGC

TABLE 5-continued

FGFR sequences.		
Nucleic acid sequence	GGGAGCCGCGCTAGCGAGCCGGGCTCCGGCGCTCGCCAGTCTCCCGAGCGGCGCCC GCCTCCCGCGGTGCCCGCGCCGGGCGGTGGGGGGCAGCATGCCCGCGCGCTGCC TGAGGACGCGCGGCCCGCCCCCGCCATGGGCGCCCCCTGCCCTGCGCCCTCGCGCT CTGCGTGGCGGTGCCCATCGTGGCCGGCGCCTCCTCGGAGTCTTGGGGACGGAGCA GCGCGTCTGGGGCGAGCGGCAGAAGTCCCGGGCCAGAGCCCGGCCAGCAGGAGCA GTTGGTCTTCGGCAGCGGGATGCTGTGGAGCTGAGCTGTCCCCCGCCCGGGGGTGG TCCCATGGGGCCACTGTCTGGGTCAAGGATGGCACAGGGCTGGTGGCCCTCGGAGCG TGTCCTGGTGGGGCCAGCGGCTGCGAGGTGCTGAATGCCTCCACGAGGACTCCGG GGCCTACAGTGC CGG CAGCGGCTCAGCAGCGCGTACTGTGCCACTTCAGTGTGCG GGTGACAGACGCTCCATCCTCGGGAGATGACGAAGACGGGGAGGACGAGGCTGAGGA CACAGGTGTGGACACAGGGGCCCTTACTGGACACGGCCGAGCGGATGGACAAGAA GCTGCTGGCGGTGCCGCGCCCAACACCGTCCGCTTCCGCTGCCAGCCGCTGGCAA CCCCACTCCCTCCATCTCCTGGCTGAAGAACGGCAGGGAGTTCGCGGGCAGCACCG CATTGGAGGCATCAAGCTGCGGCATCAGCAGTGGAGCCTGGTCATGGAAAGCGTGGT GCCCTCGGACCGCGCAACTACACCTGCGCTCGTGGAGAACAAAGTTTGGCAGCATCCG GCAGACGTACACGCTGGACGTGCTGGAGCGCTCCCGCACCGGCCCATCTGCAAGC GGGCGTGGCGCCCAACGAGCGCGGTGCTGGGCAGCGACGTGGAGTTCCACTGCAA GGTGTAAGTACGACGACAGCCCCACATCCAGTGGCTCAAGCAGTGGAGGTGAATGG CAGCAAGGTGGGCCCCGACGGCACACCTACGTTACCGTGTCAAGTCTCGGATCAG TGAGAGTGTGGAGGCGACGTGCGCCTCCGCTGGCCAAATGTGTGGAGCGGGACGG GGGCGAGTACCTCTGTGAGCCACCAATTTATAGCGTGGCCGAGAAAGGCTTTTG GCTGAGCGTTACGGGCCCCGAGCAGCCGAGGAGGAGCTGGTGGAGGCTGACGAGGC GGGCGTGTGTATGACGGCATCTCAGCTACGGGTGGGCTTCTCTCTGTTCATCCT GGTGGTGGCGGCTGTGACGCTCTGCCGCTGCGCGAGCCCCCAAGAAAGGCTGGG CTCCCCCAGCGTGACAAAGATCTCCCGCTTCCCGCTCAAGCGACAGGTGTCCCTGGA GTCCAACGCGTCCATGAGCTCCAACACACCACTGGTGCATCGCAAGGCTGTCCCT AGGGGAGGGCCCCAGCTGGCCATGTCTCCGAGCTCGAGCTGCCTGCCGACCCCAA ATGGGAGCTGTCTCGGGCCCGGTGACCTTGGGCAAGCCCTTGGGAGGGGTGCTT CGGCCAGTGGTCTATGGCGGAGGCCATCGGCATTGACAAGGACCGGCCGCCAAGCC TGTCACCGTAGCCGTGAAGATGTGAAAGACGATGCCACTGACAAGGACCTGTCCGA CCTGGTGTCTGAGATGGAGATGATGAAGATGATCGGAAACACAAAAACATCATCAA CCTGTGGGCGCCTGACCGCAGGGCGGGCCCTGTACGTGTGGTGGAGTACCGCGC CAAGGGTAACCTGCGGGAGTTTCTGCGGGCGCGCGGCCCGCGGCTGGACTACTC CTTCGACACCTGCAAGCCGCCCCGAGGAGCAGCTCACCTTCAAGGACCTGGTGTCTTG TGCTACCAAGTGGCCCGGGCATGGAGTACTTGGCCTCCAGAAAGTGATCCACAG GGACTTGGCTGCCCGCAATGTGTGTTGACCGAGGACAAAGTATGAAGATCGCAGA CTTCGGGCTGGCCCGGACGTGCACAACCTCGACTACTACAAGAAGACGACCAACGG CCGGCTGCCCGTGAAGTGGATGGCGCCTGAGGCCTTGTTTGACCGAGTCTACACTCA CCAGAGTACGCTGTGTCCTTTGGGGTCTGCTCTGGGAGATCTTACGCTGGGGGG CTCCCCGTACCCCGGCATCCTGTGGAGGAGCTCTTCAAGCTGCTGAAGGAGGGCCA CCGATGGACAAGCCCGCCAACTGCACACAGCCTGTACATGATCATCGGGAGTG CTGGCATGCCGCGCCCTCCAGAGGGCCACCTTCAAGCAGCTGGTGGAGGACCTGGA CCGTGTCTTACCGTGACGTCCACCGACGAGTACCTGGACCTGTGCGCGCCTTTGCA GCAGTACTCCCGGGTGGCCAGGACACCCCGAGCTCCAGCTCCTCAGGGGACGACT CGTGTGTGCGCACGACCTGTGCCCCCGGCCCCACCGAGTGGGGGCTCGCGGAC GTGAAGGGCCACTGGTCCCCAACATGTGAGGGGTCCCTAGCAGCCACCCCTGCTGC TGGTGACAGCCTCCTCCCGCATGAGACTCAGTGACAGATGGAGAGACAGCTACACA GAGCTTTGGTCTGTGTGTGTGTGTGCGTGTGTGTGTGTGTGTGTGCACATCCGCG TGTGCTGTGTGCGTGGCATCTTGCTCCAGGTGACAGGTACCTGGGTGTCCCG GCTGTGTGCAACGCTCTCCTGACTGGTGTGACGACCGAGGGGCTTTGTTCTGG GGGACCCAGTGCAGAAATGTAAGTGGGCCACCCGTTGGGACCCCGTGGGGCAGGG AGCTGGGCCCCGACATGGCTCCGGCCTCTGCTTTGACCCAGGGACATCACAGGGT GGCCTCGGCCCTCCACACCCAAAGCTGAGCCTGCAGGGAAGCCCCACATGTCAG CACCTTGTGCTGGGGTGTAGTGGCACCGCCTCCCGACCTCCAGGCTTTCCACTT CCCACCTGCCCCTCAGAGACTGAAATTACGGGTACCTGAAGATGGGAGCCTTTACC TTTTATGCAAAAGGTTTATTCGGAAACTAGTGTACATTTCTATAAATAGATGCTGT GTATATGGTATATATACATATATATATAACATATATGAAGAGGAAAAGGCTGGT ACAACGAGGCGCTGCGACCTTGGGGGCACAGGAGGACAGGATGGCCCTGGGCGGGC GTGGGGGGCGTGGAGGAGGCCCGAGGGGTCTACCCATGCAAGCAGAGGACAG GGCCTTTTGGCACCGCAGTTTGTGTTTAAACTGGACCTGTATATTGTAAAGCT ATTTATGGGCCCCGGCACTCTGTTTCCACACCCCAACACTTCCAGCATTTAGCTG GCCACATGGCGGAGAGTTTAAATTTTAACTTATGACAACCGAGAGGTTTATCCC GCCGATAGAGGGACGGCCAAGAAATGTACGTCCAGCTGCCCCGAGCTGGAGGATCC CCTCCAAGCCTAAAGGTTGTTAATAGTTGGAGGTGATTCAGTGAAGATATTTTAT TTCCTTTGCTCTTTTTCAGGAGAATTAGATTTCTATAGGATTTTCTTTAGGAGAT TATTTTTTGGACTTCAAAGCAAGCTGGTATTTTCATACAAATCTTCTAATTGCTGT GTGTCCAGGCAGGAGACGGTTTCCAGGAGGGGGCGGCCCTGTGTGACGGTTCCG ATGTTATAGATGTTACAAGTTTATATATATCTATATATATAATTTATGAGTTTTT ACAAGATGTATTTGTGTAGACTTAACACTTCTTACGCAATGCTTCTAGAGTTTAT AGCCTGGACTGCTACCTTTCAAAGCTTGGAGGGAAGCCGTGAATTGAGTTGGTTGCT TCTGTACTGTTACTGGGCCCTGAGTCTGGGCAGCTGTCCCTGTCTGCTGCGAGGGC CATGGCTCAGGGTGGTCTCTTCTTGGGGCCAGTGATGGTGGCCAGAGGTGTACC CAAACCGGACAGGTGCGATTTGTTAACCAGCGACGAACTTCCGAAAAATAAAGAC ACCTGGTTGCTAACCTGG	MGAPACALALCVAVAIVAGASSESLGTEQRVVGRAAEVPGPEPGQQEQLVFGSGDAV ELSCPPPGGPMGPTVWVKDGTGLVPSERVLVGPQRLQVLNASHEDSGAYSCRQLT QRVLCHFSVRVTDAPSSGDEDEGEDEEDTGVDTGAPYWTRPERMDKKLLAVPAANT

SEQ ID NO: 44
FGFR3
isoform 2
Amino acid

TABLE 5-continued

FGFR sequences.		
sequence		VRFRCPAAGNPTPSISWLKNGREFRGEHRIGGIKLRHQQWSLVMESVVPSPDRGNYTC VVENKFGSIRQTYTLDVLEERSPHRPILQAGLPANQTAVLGSDVEFHCKVYSDAQPHI QWLKHVEVNGSKVGPDPYVTVLKSWSSESVEADVRLRLANVSERDGGEYLCRATN FIGVAEKAFWLSVHGPRAAEEELVEADEAGSVYAGILSYGVGFLLVVAAVTLCL LRSPPKKGLGSPVTHKISRFPPLKRQVLSLESNASMSNTPLVRIARLSSGEGPTLANV SELELPADPKWELSRARLTLGKPLGEGCFQVVMAEAIGIDKDRAAKPVTVAVKMLK DDATDKDLSDLVSEMEMMKMIGKHKNIIINLLGACTQGGPLYVLVEYAAKGNLREFLR ARRPPGLDYSFDTCKPPEQLTFKDLVSCAYQVARGMEYLASQKCIHRDLAARNVLV TEDNVMKIADFGFLARDVHNLDYKKTNGRLPVKWMAPALFDRVYTHQSDVWSFGV LLWEIFTLGGSYPGPVPEELFKLLKEGHRMDKPANCTHDLYMIMRECWAHAPSQRP TFKQLVEDLDRVLTVSTDEYLDLSAPFEQYSPGGQDTPSSSSSGDSDVFAHDL LPP APPSSGGSRT
SEQ ID NO: 45	FGFR3 isoform 3 Nucleic acid sequence	GTCGCGGGCAGCTGGCGCCGCGCGGTCTGCTCTGCCGGTCGCACGGACGACCCGGC GGGCCGCGCGCCGAGGGACGGGGCGGGAGCTGGGCCCCGCGGACAGCGAGCCGGAGC GGGAGCCGCGCTAGCGAGCCGGGCTCCGGCGCTCGCCAGTCTCCCGAGCGGCGCCCC GCCTCCCGCGGTGCCCCGCGCGGGCGTGGGGGGCAGCATGCCCGCGCGCTGCC TGAGGACGCGCGGCCCGCCCGCCCATGGGCGCCCCCTGCCCTGCGCCCTCGCGCT CTGCGTGGCGTGGCCATCGTGGCCGCGCCTCCTCGGAGTCTTGGGGACGGAGCA CGCGCTCGTGGGGCGAGCGGCAGAAGTCCCGGGCCAGAGCCCGCCAGCAGGAGCA GTTGGTCTTGGGACGCGGGATGCTGTGGAGCTGAGCTGTCCCCCGCCCGGGGGTGG TCCCATGGGCCCCACTGTCTGGGTCAAGGATGGCACAGGGCTGGTGGCCCTCGGAGCG TGCTCTGGTGGGGCCCCAGCGGTGTCAGGTGCTGAATGCCTCCACGAGGACTCCGG GGCTACAGCTGCCGCGCAGGGCTCAGCGACGCGTACTGTGCCACTTCAGTGTGCG GGTGACAGACGCTCCATCCTCGGAGATGACGAAGACGGGGAGGACGAGGCTGAGGA CACAGGTGGACACAGGGGCCCTTACTGGACACGGCCGAGCGGATGGACAAGAA GCTGCTGGCGTGGCGCGGCCAACCCGTCGCTTCCGCTGCCAGCGCTGGCAA CCCCACTCCCTCCATCTCCTGGCTGAAGAAGCGGAGGAGTCCGCGCGGAGCACCCG CATTGGAGGCATCAAGCTGCGGCATCAGCAGTGGAGCCTGGTCATGGAAGCGTGGT GCCCTCGGACCGCGGCAACTACACTGCGCTCGTGGAGAACAAAGTTTGGCAGCATCCG GCAGACGTACACGCTGGACGTGCTGGAGCGCTCCCGCACCGGCCCATCTGCAAGC GGGGCTGCCGGCCCAACGACGCGCGGTGCTGGGCGAGCAGTGGAGTTCCACTGCAA GGTGTACAGTGACGCACAGCCCCACATCCAGTGGCTCAAGCAGTGGAGGTGAATGG CAGCAAGGTGGGCCCGGACGGCACACCTACGTTACCGTGTCTAAGTGTCTCCCTGGA GTCCAAACGCTCCATGAGCTCCAACACCACTGGTGGCATCGCAAGGCTGTCTC AGGGGAGGGCCCCACGCTGGCCATGTCTCCGAGCTCGAGCTGCCTGCCAGCCCCAA ATGGGAGCTGTCTCGGGCCCGGTGACCTGGGCAAGCCCCCTGGGGAGGGCTGTCTT CGGCCAGTGGTCTATGGCGGAGGCCATCGGCATTGACAAGGACCGGGCGCCCAAGCC TGTACCGTAGCCGTGAAGATGCTGAAAGACGATGCCACTGACAAGGACCTGTCCGA CCTGGTGTCTGAGATGGAGATGATGAAGATGATCGGAAACACAAAAACATCATCAA CCTGTGGGCGCCTGCACGCAAGGCGGGCCCCGTGACGTGCTGGTGGAGTACGCGGC CAAGGGTAACCTGCGGGAGTTTCTGCGGGCGCGCGGCCCGCGGCTGGAGTACTC CTTGCAACCTGCAAGCCGCGCCGAGGAGCAGCTCACCTTCAAGGACCTGGTGTCTCTG TGCTTACAGGTGGCCGCGGCATGGAGTACTTGGCCTCCAGAAAGTGATCCACAG GGACTTGGCTGCCCGCAATGTGTGGTGACCGAGGACAAAGTGTGAAGATCCGAGA CTTCCGGCTGGCCCGGACGTGCACAACCTCGACTACTACAAGAAGACGACCAACGG CCGCTGCCCGTGAAGTGGATGGCGCCTGAGGCCTTGTGACCGAGTCTACACTCA CCAGAGTGACGTCTGGTCTTTGGGGTCTGCTCTGGGAGATCTTACGCTGGGGGG CTCCTCCGTACCCCGGCATCCTGTGGAGGAGCTCTTCAAGCTGCTGAAGGAGGGCCA CCGCATGGACAAGCCCGCAACTGCACACGACCTGTACATGATCATCGGGAGTG CTGGCATGCCGCGCCCTCCAGAGGCCACCTTCAAGCAGCTGGTGGAGGACCTGGA CCGTGTCTTACCGTGACGTCCACCGACGAGTACCTGGACCTGTGCGCGCTTTCGA CGAGTACTCCCGGGTGGCCAGGACACCCCGAGCTCCAGCTCCTCAGGGGACGACTC CGTGTGTTGCCACGACCTGTGCCCCCGGCCACCCAGCAGTGGGGGCTCGCGGAC GTGAAGGGCCACTGGTCCCCAACATGTGAGGGGTCCCTAGCAGCCACCCCTGCTGC TGGTGACAGCCTCCTCCCGCATGAGACTCAGTGACAGATGGAGAGACGCTACACA GAGCTTTGGTCTGTGTGTGTGTGTGTGCGTGTGTGTGTGTGTGTGTGCACATCCGCG TGTGCTGTGTGCGTGGCATCTTGCCTCCAGGTGACAGGATACCTGGGTGTCCCC GCTGCTGTGCAACGCTCCTGACTGGTGTGTCAGCACCGAGGGGCTTTGTTCTGG GGGACCCAGTGCAAGTGAAGTGGGCCACCCGCTGGGACCCCGTGGGGCAGGG AGCTGGGCCCCGACATGGCTCCGGCCTCTGCCTTTGCACCCAGGGGACATCACAGGGT GGCTCGGCCCTCCACACCCAAAGCTGAGCCTGCAGGGAAGCCCCACATGTCCAG CACCTTGTGCTGGGGTGTAGTGGCACCGCCTCCACACCTCCAGGCTTTCCACTT CCCACCTGCCCCCTCAGAGACTGAAATTACGGGTACCTGAAGATGGGAGCCTTTACC TTTTATGCAAAAGTTTATTCCGAAACTAGTGATACATTTCTATAAATAGATGCTGT GTATATGGTATATATACATATATATATAACATATATGGAAGAGGAAAAGGCTGGT ACAACGAGGCGCTGCGACCTGGGGGCACAGGAGGCGAGGATGGCCCTGGGCGGGC GTGGGGGGCGTGGAGGGAGGCCACAGGGGTCTACCCATGCAAGCAGAGGACGAG GGCCTTTCTGGCACCGCAGTTTGTGTTTAAACTGGACCTGTATATTGTAAAGCT ATTTATGGGCCCCGCGACTCTTGTTCACACCCCAACACTTCCAGCATTTAGCTG GCCACATGGCGGAGAGTTTAAATTTTAACTTATGACAACCGAGAGGTTTATCCC GCCGATAGAGGGACGGCCAAGATGTACGTCCAGCTGCCCGGAGCTGGAGGATCC CCTCCAAGCCTAAAGGTTGTTAATAGTTGGAGGTGATCCAGTGAAGATATTTTAT TTCCTTTGCTCTTTTCAGGAGAATTAGATTTCTATAGGATTTTCTTTAGGAGAT TATTTTTTGGACTTCAAAGCAAGCTGGTATTTTCATACAAATCTTCTAATTGCTGT GTGTCCAGGCGAGGAGAGCGTTTCCAGGAGGGGGCGGCCCTGTGTGACGGTTCCG ATGTTATAGATGTTACAAGTTTATATATATCTATATATATAATTTATTGAGTTTTT ACAAGATGATTTTGTGTAGACTTAACACTTCTTACGCAATGCTTCTAGAGTTTTAT

TABLE 5-continued

FGFR sequences.		
SEQ ID NO: 46	FGFR3 isoform 3 Amino acid sequence	AGCCTGGACTGCTACCTTTCAAAGCTTGGAGGGAAGCCGTGAATTCAGTTGGTTCGT TCTGTACTGTACTGGGCCCTGAGTCTGGGCAGCTGTCCCTTGCTTGCCTGCAGGGC CATGGCTCAGGGTGGTCTCTTCTTGGGGCCAGTGCATGGTGGCCAGAGGTGTACC CAAACCGGCAGGTGCGATTTTGTTAACCCAGCGACGAACCTTCCGAAAAATAAAGAC ACCTGGTTGCTAACCTGG MGAPACALALCVAIVAGASSESLGTEQRVVGRAAEVPGPEPGQQEQLVFGSGDAV ELSCPPPGGGPMGPTVWVKDGTGLVPSERVLVGPQRLQVLNASHEDSGAYSCRQLT QRVLCHF SVRVTDAPSSGDEDEGEDEAEDTGVDTGAPYWTRPERMDKKLLAVPAANT VRFRCPAAGNPTPSISWLKNGREFRGEHRIGGIKLRHQQWSLVMSVVPVSDRGNYTC VVENKFGSIRQTYTLDVLEERSPHRPILQAGLPANQTAVLGSDFEFHCKVYSDAQPHI QWLKHVEVNGSKVGPDPYVTVLKVLSLESNASMSNTPLVRIARLSSGEGPTLANV SELELPADPKWELSRARLTGLKPLGEGCFGQVVMABEIGIDKDRAAKPVTVAVKMLK DDATDKDLSDLVSEMEMMKMIGKHNIINLLGACTQGGPLYVLVEYAAKGNLREFLR ARRPPGLDYSFDTCKPPEQLTFKDLVSCAYQVARGMEYLASQKCIHRDLAARNVLV TEDNVMKIADFLGARDVHNLDDYKKTNGRLPVKWMPEALFDRVYTHQSDVWSFGV LLWEIFTLGGSPYPGIPVEELFKLLKEGHRMDKPANCTHDLYMIMRECWHAAAPSQR TFKQLVEDLDRVLTVTSTDEYLDLSAPFEQYSPGGQDTPSSSSSGDSDVFAHDLPLP APPSSGGSRT
SEQ ID NO: 47	FGFR3 isoform 4 Nucleic acid sequence	CGCGCGCTGCCTGAGGACGCGCGGCCCGCCCGCCCGCCCATGGGCGCCCTGCCTGC GCCCTCGCGCTCTGCGTGGCCGTGGCCATCGTGGCCGGCGCCCTCTCGGAGTCCTTG GGGACGGAGCAGCGCGTCGTGGGGCGAGCGGCAGAAAGTCCCGGGCCAGAGCCCGGC CAGCAGGAGCAGTTGGTCTTCGGCAGCGGGGATGCTGTGGAGCTGAGCTGTCCCCCG CCCGGGGGTGGTCCCATGGGGCCACTGTCTGGGTCAAGGATGGCACAGGGCTGGTG CCCTCGAGCGTGTCTGGTGGGGCCCGAGCGGCTGCAGGTGCTGAATGCCTCCAC GAGGACTCCGGGGCTACAGCTGCCGCGCAGCGGCTCACGCGCGCGTACTGTGCCAC TTCAGTGTGCGGGTGACAGACGCTCCATCCTCGGGAGATGACGAAGACGGGGAGGAC GAGGCTGAGGACACAGGTGTGGACACAGGGGCCCTTACTGGACACGGCCCGAGCGG ATGGACAAGAAGCTGTGGCGGTGCCGGCCGCCAACACCGTCCGCTTCGCTGCCCA GCCGCTGGCAACCCACTCCCTCCATCTCCTGGCTGAAGAACGGCAGGGAGTTCGCG GGCGAGCACCGCATTTGGAGGCATCAAGCTGCGGCATCAGCAGTGGAGCTGGTCATG GAAAGCTGGTGCCCTCGGACCGCGGCAACTACACCTGCGCTCGTGGAGAACAAGTTT GGCAGCATCCGGCAGACGTACACGCTGGACGTGCTGGAGCGCTCCCGCACCGGCC ATCCTGCAAGCGGGCTGCCGGCCAACAGACGGCGGTGCTGGGCAGCGACGTGGAG TTCCACTGCAAGGTGTACAGTGACGCACAGCCCCACATCCAGTGGCTCAAGCACGTG GAGGTGAATGGCAGCAAGGTGGGCCCGGACGGCACACCTTACGTTACCGTGCTCAAG GTGTCCCTGGAGTCCAACGCGTCCATGAGCTCCAACACACCACTGGTGCGCATCGCA AGGTGTCTCAGGGGAGGGCCCCACGCTGGCCAATGTCTCCGAGCTCGAGCTGCCT GCCGACCCCAATGGGAGCTGTCTCGGGCCCGCTGACCTGGGCAAGCCCCCTGGG GAGGGCTGCTTCGCGCAGGTGGTCATGGCGGAGGCCATCGGCATTGACAAGGACCG GCCGCCAAGCTGTACCGTAGCCGTGAAGATGCTGAAAGACGATGCCACTGACAAG GACCTGTGCGGACCTGGTGTCTGAGATGGAGATGATGAAGATGATCGGGAACACAAA AACATCATCAACCTGCTGGGCGCCTGCACGACGGCGGGCCCTGTACGTGCTGGTG GAGTACGCGGCCAAGGGTAACCTGCGGGAGTTTCTGCGGGCGCGGCGGCCCGGGC CTGGACTACTCCTTCGACACCTGCAAGCGCCCGGAGGAGCAGCTCACCTTCAAGGAC CTGTGTCTCTGCTTACCAGGTGGCCCGGGGATGGAGTACTTGGCTCCAGAAAG TGCATCCACAGGGACCTGGCTGCCCGCAATGTGCTGGTGACCGAGGACAACGTGATG AAGATCGCAGACTTCGGGCTGGCCCGGACGTGCACAACCTCGACTACTACAAGAAG ACGACCAACGGCCGGCTGCCCGTGAAGTGGATGGCGCTGAGGCTTGTTTGACCGA GTCTACACTCACCAGAGTGACGTCTGGTCTTTGGGGTCTGCTCTGGGAGATCTTC ACGCTGGGGGGCTCCCGGTACCCCGGCATCCCTGTGGAGGAGCTCTTCAAGCTGCTG AAGGAGGGCCACGCATGGACAAGCCCGCCAAGTGCACACAGACCTGTACATGATC ATGCGGGAGTGCTGGCATGCCGCGCCTCCAGAGGGCCACCTTCAAGCAGCTGGTG GAGGACCTGGACCGTGTCTTACCGTGACGTCCACCGACGAGTACCTGGACCTGTGCG GCGCTTTCGAGCAGTACTCCCGGGTGGCCAGGACACCCCGAGCTCCAGCTCTCA GGGGACGACTCCGTGTTTGCCACGACCTGCTGCCCGGGCCCGCCAGCAGTGGG GGCTCGCGGACGTGAAGGGCCACTGGTCCCAACAATGTGAGGGGTCCCTAGCAGCC CACCTGCTGCTGGTGA
SEQ ID NO: 48	FGFR3 isoform 4 Amino acid sequence	MGAPACALALCVAIVAGASSESLGTEQRVVGRAAEVPGPEPGQQEQLVFGSGDAV ELSCPPPGGGPMGPTVWVKDGTGLVPSERVLVGPQRLQVLNASHEDSGAYSCRQLT QRVLCHF SVRVTDAPSSGDEDEGEDEAEDTGVDTGAPYWTRPERMDKKLLAVPAANT VRFRCPAAGNPTPSISWLKNGREFRGEHRIGGIKLRHQQWSLVMSVVPVSDRGNYTC VVENKFGSIRQTYTLDVLEERSPHRPILQAGLPANQTAVLGSDFEFHCKVYSDAQPHI QWLKHVEVNGSKVGPDPYVTVLKVLSLESNASMSNTPLVRIARLSSGEGPTLANV SELELPADPKWELSRARLTGLKPLGEGCFGQVVMABEIGIDKDRAAKPVTVAVKMLK DDATDKDLSDLVSEMEMMKMIGKHNIINLLGACTQGGPLYVLVEYAAKGNLREFLR ARRPPGLDYSFDTCKPPEQLTFKDLVSCAYQVARGMEYLASQKCIHRDLAARNVLV TEDNVMKIADFLGARDVHNLDDYKKTNGRLPVKWMPEALFDRVYTHQSDVWSFGV LLWEIFTLGGSPYPGIPVEELFKLLKEGHRMDKPANCTHDLYMIMRECWHAAAPSQR TFKQLVEDLDRVLTVTSTDEYLDLSAPFEQYSPGGQDTPSSSSSGDSDVFAHDLPLP APPSSGGSRT
SEQ ID NO: 49	FGFR4 isoform 1 Nucleic acid sequence	CGCTCGCGGCCACGCGCCGCTCGCGGGTACATTCTCGCTCCCGGCCGAGGAGCGCT CGGGCTGTCTGCGGACCTCGCCGCTGCAGGGTTCGCGCCGGCTGGAGCTGGGAGT GAGCGCGGAGGAGCCAGGTGAGGAGGAGCCAGGTGAGCAGGACCTGTGCTGGGC CGGAGTACGCGAGGCTCGAGGAAGGAGTGGTGGGAAGTCCAGCTTGGGTCCCTG

TABLE 5-continued

FGFR sequences.	
SEQ ID NO: 50	FGFRA isoform 1 Amino acid sequence MRLLLALLGVLLSVPGPPVLSLEASEEVELEPCLAPSLQEQELTVALGQPVRLCC GRAERGGHWYKEGSR LAPAGVRVGRGRLEIASFLPEDAGRYLCLARGSIMIVLQNL LTITGDSLTSNDDPKSHRDPNHRHSYPQAPYWTHPQRMKKLHVPAGNTVKFR CPAAGNPTPIRWLKDQAFHGENRIGGIRLRHQHWSLVMSVVPDRGTYTCLVEN AVGSIRYNYLLDVLERSPHRPI LQAGLPANTTAVVGSDELCKVYSDAQPHIQWLK HIVINGSSFGADGPPYVQVLKTADINSSEVEVLYLRNVSAEDAGEYTCLAGNSIGLS YQSAWLTVLPEEDPTWTAAPPEARITDI ILYASGSLALAVLLLLAGLYRQALHGRH PRPPATVQKLSRFPLARQFSLESSSGKSSSLVRGVRLSSSGPALLAGLVSLDLPL DPLWEFPRDRVLVKGKPLGEGCFQGVVRAEAFGMDPARPDQASTVAVKMLKDNASDKD LADLVSEMEVMKIGRHKNI INLLGVCTQEGPLYV IVECAAAGKNLREFLRARRPPGP DLSPDGPRSEGLSFPVLVSCAYQVARGMQYLESRKITHRDLAARNVLVTEDNVMK IADFGLARGVHHIDYKKTSGRLPVKWMAPALFDRVYTHQSDVVSFILLWEIFT LGGSPYPGI PVEELFSLREGHRMDRPPHCPPELYGLMRECWAAPSRPTFKQLVE ALDKVLLAVSEEYLDLRLTFGPYSPSGGDASSTCSSSDSVFSDHPLPLGSSSPFPGS GVQT
SEQ ID NO: 51	FGFRA isoform 2 Nucleic acid sequence GTCGCGGGTACATTCTCGCTCCCGGCCGAGGAGCGCTCGGGCTGTCTGCGGACCCCT GCCCGCTGCAGGGGTGCGCGCCGGCTGGAGCTGGGAGTGAGGCGCGGAGGAGCCAG GTGAGGAGGAGCCAGGAAGGCAGTTGGTGGGAAGTCCAGCTTGGGTCCCTGAGAGCT GTGAGAAGGAGATGCGGCTGCTGCTGGCCCTGTTGGGGGTCCTGCTGAGTGTGCTG GGCTCCAGTCTTGTCCCTGGAGGCCCTCTGAGGAAGTGGAGCTTGAGCCCTGCTTGG CTCCAGCTGGAGCAGCAAGAGCAGGAGCTGACAGTAGCCCTTGGGACGCTGTGCTG GTCTGTGCTGTGGGCGGGCTGAGCGTGGTGGCCACTGGTACAAGGAGGGCAGTCGCC TGGCACCTGCTGGCCGTGTACGGGGCTGGAGGGGCCCGCTAGAGATTGCCAGCTTCC TACCTGAGGATGCTGGCCGTACCTCTGCTGGCAGCAGGCTCCATGCTGCTGCTGCTG AGAATCTCACCTTGATTACAGGTGACTCCTTGACCTCCAGCAACGATGATGAGGACC

TABLE 5-continued

FGFR sequences.		
		CCAAGTCCCATAGGGACCCCTCGAATAGGCACAGTTACCCCCAGCAAGCACCCCTACT GGACACACCCCCAGCGCATGGAGAAGAACTGCATGCAGTACCTGCGGGGAACACCG TCAAGTTCCGCTGTCCAGCTGCAGGCAACCCACGCCCACCATCCGCTGGCTTAAGG ATGGACAGGCCTTTCATGGGGAGAACCGCATTGGAGGCATTCCGCTGCGCCATCAGC ACTGGAGTCTCGTGATGGAGAGCGTGGTGCCCTCGGACCGCGGCACATACACCTGCC TGGTAGAGAACGCTGTGGGCAGCATCCGCTATAACTACCTGCTAGATGTGCTGGAGC GGTCCCCGACCCGGCCCATCTGCAGGCGGGCTCCCGGCCAACACCACAGCCGTGG TGGGCAGCGACCTGGAGCTGCTGTGCAAGGTGTACAGCGATGCCAGCCCCACATCC AGTGGCTGAAGCACATCGTCATCAACGGCAGCAGCTTCGGAGCCGACGGTTTCCCT ATGTGCAAGTCTTAAAGATGCAGACATCAATAGCTCAGAGGTGGAGGTCTGTACC TCGGAAACGTGTGTCAGCGAGGACGCGAGGCGAGTACACCTGCCTCGAGGCAATTCCA TCGCCTCTCTTACCAGTCTGCTTGGCTCAGGTGCTGCCAGAGGAGGACCCACAT GGACCGCAGCAGCGCCGAGGCGAGTTCTCCCTGGAGTCAAGGCTCTTCCGGCAAGTC AAGCTCATCCTGGTACGAGCGTGGCTCTCTCTCCAGCGGCCCGCCTTGTCTCGC CGGCTCGCTGGTGTGGGAAGCCCCTAGGCGAGGGCTGCTTGGCCAGGTAGTAC GTGCAGAGGCTTTGGCATGGACCTGCCCGGCTGACCAAGCCAGCACTGTGGCCG TCAAGATGCTCAAAGACAACGCTCTGACAAGGACCTGGCCGACCTGGTCTCGGAGA TGGAGGTGATGAAGCTGATCGGCCGACACAAGAACATCATCAACCTGCTTGGTGTCT GCACCCAGGAAGGGCCCTGTACGTGATCGTGGAGTGCAGCGCCCAAGGGAAACCTGC GGGAGTTCTTGGGGCCCGCGCCCCAGGCCCGACCTCAGCCCCGACGGTCTCT GGAGCAGTGAGGGGCGCTCTCTTCCAGTCTGTGCTCTGCGCTACCAAGGTGG CCCGAGGCATGCAGTATCTGGAGTCCCGGAAGTGTATCCACCGGACCTGGCTGCC GCAATGTGCTGGTGACTGAGGACAATGTGATGAAGATTGCTGACTTGGGCTGGCC GCGCGTCCACCATGACTACTATAAGAAAACAGCAACGGCCGCTGCTGTGTA AGTGGATGGCGCCGAGGCTTGTGTTGACCGGGTGACACACCCAGAGTGACGTGT GGTCTTTGGGATCTGCTATGGAGATCTTACCCCTCGGGGCTCCCGTATCCTG GCATCCCGTGGAGAGCTGTTCTGCTGCTGCGGGAGGACATCGGATGGACCGAC CCCCACACTGCCCCCAGAGCTGTACGGGCTGATGCGTGAGTGTGGCACGACGCG CCTCCCAGAGGCTTACCTTCAAGCAGCTGGTGGAGGCGCTGGACAAGGTCTGCTGG CCGTCTCTGAGGAGTACCTCGACCTCCGCTGACCTTCGGACCTATTCCTCTCTG GTGGGGACGCCAGCAGCACCTGCTCTCCAGCGATTCTGTCTTCAGCCACGACCCCC TGCCATTGGGATCCAGCTCTTCCCTTCCGGTCTGGGGTGACAGATGAGCAAGGC TCAAGGTGTGACGGCACATAGGCTGGTGGCTTGGGCCTTGGGGCTCAGCCACAGC CTGACACAGTGTCTGACCTTGATAGCATGGGGCCCCCTGGCCAGAGTTGCTGTGCCG TGTCCAAGGGCCGTGCCCTTGCCTTGGAGCTGCCGTGCTGTCTGATGGCCCA AATGTACAGGTTCTGCTCGCTTCTTGGACCTTGGCGCTTAGTCCCATCCCGGTT TGGCTGAGCCTGGCTGGAGAGCTGCTATGCTAAACCTCTGCTCCCAATACACGCA GGAGTTCTGGGCTCTGAACCCCTTCCCCACACCTCCCTGCTGCTGCTG
SEQ ID NO: 52	FGFR4 isoform 2 Amino acid sequence	MRLLALLGVLLSVPGPPVLSLEASEEVELEPCLAPSLQQEQELTVALGQPVRLCC GRAERGGHWYKEGSR LAPAGVRVGRRLLEIASFLPEDAGRYLCARGSMIVLQNL LITGDSLTSNDDDEPKSHRDPNHRHSYPQAPYWTHPQRMKKLHAPAGNTVKFR CPAAGNPTPTIRWLKDGAFHGENRIGGIRLRHQHWSLVMSVVPSPDRGTYTCLVEN AVGSIRYNYLLDLVLESPHRPI LQAGLPANTTAVVGSDELLECKVYSDAQPHIQWLK HIVINGSFSGADGFPYVQVLKTADINSSEVEVLYLRNVSAEDAGEYTCLAGNSIGLS YQSAWLTVLPEEDPTWTAAPAEASSPWSQALPASQAHWPYEAACVSPPAAPPCSPASL VLGKPLGEGCFQVVRAEAFGMDPARPDQASTVAVKMLKDNASDKDLADLVSEMEVM KLIGRHKNI INLLGVCTQEGPLYVIVECAAKGNLREFLRARRPPGPDLPSPDGRSSE GPLSFPLVLS CAYQVARGMQYLESRKITHRDLAARNVLVTEDNVMKIADPGLARGVH HIDYKKTSNGRLPVKWMAPALFDRVYTHQSDVWSFGILLWEIFTLGGSPYPGPIV EELFSLLRGHRMDRPPHCPPELYGLMRECWAAPSQRPTFKQLVEALDKVLLAVSE EYDLRLTLFGPYSPSGDASSTCSSSDSVFSDHPLPLGSSSPFPGSGVQT
SEQ ID NO: 53	FGFR4 isoform 3 Nucleic acid sequence	ACATTCTCGCTCCCGCCGAGGAGCGCTCGGGCTGTCTGCGGACCTGCCGCGTGC AGGGGTGCGGCCCGCTGGAGCTGGGAGTGAGGCGCGGAGGAGCCAGGTGAGGAGG AGCCAGGAAGCAGTTGGTGGGAAGTCCAGCTTGGGTCCCTGAGAGCTGTGAGAAGG AGATGCGGCTGTGCTGGCCCTGTTGGGGGTCTGTGAGTGTGCTGGGCTCCAG TCTTGTCCCTGGAGGCTCTGAGGAAGTGAGCTTGAGCCCTGCCTGGCTCCAGCC TGGAGCAGCAAGAGCAGGAGCTGACAGTAGCCCTTGGGCAGCCTGTGCGTCTGTGCT GTGGGCGGGCTGAGCGTGGTGGCCACTGGTACAAGGAGGGCAGTCGCTGGCAGCTG CTGGCCGTGTACGGGGCTGGAGGGGCGCCTAGAGATTGCCAGCTTCTTACCTGAGG ATGCTGGCCGCTACCTGTGCTGGCAGGAGCTCCATGATCGTCTGAGAAATCTCA CCTTGATTACAGGTGACTCCTTGACCTCCAGCAACGATGATGAGGACCCCAAGTCCC ATAGGGACCCCTCGAATAGGCACAGTTACCCCCAGCAAGCACCTACTGGACACACC CCCAGCGCATGGAGAAGAACTGCATGCAGTACCTGCGGGGAACACCGCTCAAGTTCC GCTGTCCAGTGCAGGCAACCCACGCCCACCATCCGCTGGCTTAAGGATGGACAGG CCTTTCATGGGGAGAACCAGCATGGAGGCATTCCGGTGCGCCATCAGCACTGGAGTC TCGTGATGGAGAGCGTGGTGCCCTCGGACCGCGGCACATACCTGCTGCTGAGAGA ACGCTGTGGGCAGCATCCGCTATAACTACCTGCTAGATGTGCTGGAGCGGTCCCCGC ACCGGCCCATCTGACAGGCGGGCTCCCGGCCAACACCACAGCCGTGGTGGGCAGCG ACGTGGAGCTGCTGTGCAAGGTGTACAGCGATGCCAGCCCCACATCCAGTGGCTGA AGCACATCGTCATCAACGGCAGCAGCTTCGGAGCCGACGGTTTCCCTATGTGCAAG TCCTAAAGACTGCAGACATCAATAGCTCAGAGGTGGAGGTCTGTACCTGCGGAACG TGTACGCCGAGGACGCGAGGCGAGTACCTGCTCGAGGCAATTTCATCGGCTCT CCTACCAGTCTGCTGGCTCAGGTGCTGCCAGAGGAGGACCCACATGGACCGCAG CAGCGCCGAGGCGAGGTATACGGACATCATCTGTACGCGTGGGCTCCCTGGCT TGGCTGTGCTCTGCTGCTGGCGGGCTGTATCGAGGGCAGGCGCTCCAGCGCCGC ACCCCGCCCCCGCCACTGTGCGAGAAGTCTCCCGCTTCCCTCTGGCCCGACAGT

TABLE 5-continued

FGFR sequences.		
SEQ ID NO: 54	FGFR4 isoform 3 Amino acid sequence	TCTCCCTGGAGTCAGGCTCTTCCGGCAAGTCAAGCTCATCCCTGGTACGAGGCGTGC GTCTCTCCTCCAGCGGCCCCGCTTGTCTCGCCGGCTCGTGAGTCTAGATCTACCTC TCGACCCACTATGGGAGTTCCCCCGGACAGGCTGGTGTGGGAAGCCCTAGGCG AGGGCTGCTTTGGCCAGGTAGTACGTGCAGAGGCCCTTGGCATGGACCTGCCCCGGC CTGACCAAGCCAGCACTGTGGCCGTCAAGATGCTCAAAGACAACGCCTCTGACAAGG ACCTGGCCGACCTGGTCTCGGAGATGGAGGTGATGAAGCTGATCGGCCGACACAAGA ACATCATCAACCTGCTTGGTGTCTGCACCCAGGAAGGGCCCCGTGACGTGATCGTGG AGTGCGCCGCCAAGGAAACCTGCGGGAGTTCTGCGGGCCCGGCGCCCCCAGGCG CCGACCTCAGCCCCGACGGTCTCTCGGAGCAGTGAGGGCCGCTCTCCTTCCAGTCC TGGTCTCCTGCGCTACCAGGTGGCCCGAGGCATGCAGTATCTGAGTCCCGAAGT GTATCCACCGGACCTGGCTGCCCGCAATGTGCTGGTGAAGTGAAGACAATGTGATGA AGATTGTGACTTTGGGCTGGCCCGCGGCTCCACCACATTGACTACTATAAGAAAA CCAGCAACGGCCGCTGCCTGTGAAGTGGATGGCGCCCGAGGCCCTTGTGACCGGG TGTAACACACACAGAGTGACGTGTGGTCTTTGGGATCCTGCTATGGGAGATCTTCA CCCTCGGGGGCTCCCGTATCTGGCATCCCGGTGGAGGAGCTGTTCTCGCTCTGTC GGGAGGACATCGGATGGACCGACCCCACTGCCCCCAGAGCTGTACGGGTGA TGCGTGAGTGCTGGCAGCAGCGCCCTCCAGAGGCCCTACCTTCAAGCAGCTGGTGG AGGCGCTGGACAAGGTCTGCTGGCCGTCTCTGAGGAGTACCTCGACCTCCGCTGA CCTTCGAGCCCTATTCCTCTGCTGGTGGGACGCCAGCAGCAGCTGCTCTCCAGCG ATTCTGTCTCAGCCACGACCCCTGCCATTGGGATCCAGCTCTCTCCCTTCGGGT CTGGGTGTCAGACATGAGCAAGGCTCAAGGCTGTGCAGGCACATAGGCTGGTGGCT TGGGCTTGGGGCTCA
SEQ ID NO: 54	FGFR4 isoform 3 Amino acid sequence	MRLLLALLGVLLSVPGPPVLSLEASEEVELEPCLAPSLQQEQELTVALGQPVRLLC GRAERGGHWYKEGSRLLAPAGVRVGRRLRIASFLPEDAGRYLCLARGSIMIVLQNL LITGDSLTSNDDPKSHRDPNRSYPPQAPYWTHPQRMKKLHAVPAGNTVKFR CPAAGNPTPTIRWLKDGQAFHGENRIGIRLRHQHWSLVMSVVPDRGTYTCLVEN AVGSIRYNYLLDVLERSPHRPIQLQAGLPANTTAVVGSVDVLLCKVYSDAPHIQWLK HIVINGSSFGADGPPYVQVLKTADINSSEVEVLYLRNVSAEDAGEYTCLAGNSIGLS YQSAWLTVLPEEDPTWTAAPARYTDIILYASGSLALAVLLLAGLYRGQALHGRH PRPPATVQKLSRFPLARQFSLESGSSGKSSSLVRGVRLSSSGPALLAGLVSLDLPL DPLWEFPRDRLLVLGKPLGEGCFGQVVRAEAFGMDPARPDQASTVAVKMLKDNASDKD LADLVSEMEVMKLIGRHKNIINLLGVCTQEGPLYVIVECAAAGNKLREFLRARRPPGP LSPDGPRESSEGPLSPFVLVSCAYQVARGMQYLESRKCIRHDLAARNVLVTEDNVMK IADPGLARGVHHIDYKKTNGRLPVKWMPEALFDRVYTHQSDVWSFGILLWEIFT LGGSPYPGIPEEELFSLREGHRMDRPPHCPPELYGLMRECWAAPSORPTFKQLVE ALDKVLLAVSEYLDLRLTFGPYSPSGGDASSTCSSDSVFSHDPLPLGSSSPFPFGS GVQT
SEQ ID NO: 55	FGFR4 isoform 4 Nucleic acid sequence	AGTCCAGCTTGGGTCCCTGAGAGCTGTGAGAAGGAGATGCGGCTGCTGCTGGCCCTG TTGGGGTCTCTGAGTGTGCTGGGCTCCAGTCTGTCCCTGGAGGCTCTGAG GAAGTGGAGCTTGAGCCCTGCTGGCTCCAGCTGGAGCAGCAAGAGCAGGAGCTG ACAGTAGCCCTTGGGCAGCCTGTGCGTCTGTGCTGTGGCGGGCTGAGCGTGGTGGC CACTGGTACAAGGAGGCGAGTCGCTGGCACCTGCTGGCGGTGTACGGGCTGGAGG GGCCGCTAGAGATTGCCAGCTTCTACCTGAGGATGCTGGCCGCTACCTCTGCCTG GCACGAGGCTCCATGATCGTCTGCAGAATCTACCTTGATTACAGTGACTCCTTG ACCTCCAGCAACGATGATGAGGACCCCAAGTCCCATAGGACCCCTCGAATAGGCAC AGTTACCCCCAGCAAGCACCTACTGACACACCCAGCGCATGGAGAAGAACTG CATGCAGTACCTGCGGGAAACACCGTCAAGTCCGCTGTCCAGCTGCAGGCAACCC ACGCCACCATCCGCTGGCTTAAGGATGGACAGGCCTTTCATGGGGAGAACCGCATT GGAGGCATTGCGCTGCGCCATCAGCACTGGAGTCTCGTGATGGAGAGCGTGGTGGCC TCGACCGCGGCACATACACCTGCCTGGTAGAGAACGCTGTGGGCAGCATCCGCTAT AACTACCTGCTAGATGTGCTGGAGCGGTCCCCGACCGGCCATCCTGAGGCGGG CTCCCGCCAAACACACAGCGTGGTGGGACGACGTGGAGCTGCTGTGCAAGGTG TACAGCGATGCCAGCCCCACATCCAGTGGCTGAAGCACATCGTCATCAACGGCAGC AGCTTCGAGAGCCGACGGTTTCCCTATGTGCAAGTCTTAAAGACTGCAGACATCAAT AGCTCAGAGGTGGAGTCTGTACCTGCGGAACGTGTACGCGAGGACGACGAGCGAG TACACCTGCTCGAGGCAATTCATCGGCCTCTCTACAGTCTGCTGGCTCAGC GTGCTGCCAGGTACTGGGCGCATCCCCACCTCACATGTGACAGCCTGACTCCAGCA GGCAGAACCAGTCTCCCACTTTCAGTCTCTCTGGAGTCAAGGCTCTTCCGGCAAG TCAAGCTCATCCCTGGTACGAGGCGTGTCTCTCTCCAGCGGCCCGCTTGTGCT GCCGCGCTCGTGTGCTAGATCTACCTCTCGACCCACTATGGGAGTTCCTCCGGGAC AGGCTGGTGTCTGGGAAGCCCTAGGCGAGGCTGCTTTGGCCAGGTAGTACGTGCA GAGGCTTTGGCATGGACCTGCCCCGCTGACCAAGCCAGCACTGTGGCCGTCAAG ATGCTCAAAGACAACGCCTCTGACAAGGACCTGGCCGACCTGGTCTCGGAGATGGAG GTGATGAAGCTGATCGGCCGACACAAGAACATCATCAACCTGCTTGGTGTCTGCACC CAGGAAGGGCCCCGTGACGTGATCGTGGAGTGCGCCGCCAAGGGAACCTGCGGGAG TTCTGCGGGCCCGCGCCCCCAGGCCCCGACCTCAGCCCCGACGCTCTCGGAGC AGTGAGGGCCGCTCTCTTCCAGTCTGGTCTCTGCGCTACAGGTGGCCCGA GGCATGCAGTATCGGAGTCCCGAAGTGTATCCACCGGACCTGGCTGCCCGCAAT GTGCTGGTGACTGAGGACAATGTGATGAAGATTGCTGACTTTGGGCTGGCCCCGCGC

TABLE 5-continued

FGFR sequences.		
GTCCACCACATTGACTACTATAAGAAAACAGCAACGGCCGCCTGCCTGTGAAGTGG ATGGCGCCCGAGGCCTTGTGTTGACCGGGGTACACACACCAGAGTGACGTGTGGTCT TTTGGGATCCTGCTATGGGAGATCTTACCTCGGGGGCTCCCGTATCTGGCATC CCGGTGGAGGAGCTGTTCTCGCTGCTGCGGGAGGGACATCGGATGGACCGACCCCA CACTGCCCCCAGAGCTGTACGGGCTGATGCGTGAGTGCTGGCACGACGCGCCCTCC CAGAGGCCTACCTTCAAGCAGCTGGTGGAGGCGCTGGACAAGGTCCTGCTGGCCGTC TCTGAGGAGTACCTCGACCTCCGCCTGACCTTCGGACCCCTATTCCCCCTCTGGTGGG GACGCCAGCAGCACCTGCTCCTCCAGCGATTCTGTCTCAGCCACGACCCCTGCCA TTGGGATCCAGCTCCTTCCCTTCGGGCTCTGGGGTGACACATGAGCAAGGCTCAAG GCTGTGCAGGCACATAGGCTGGTGGCCTTGGGCCTTGGGGCTCAGCCACAGCCTGAC ACAGTGCTCGACCTTGATAGCATG		
SEQ ID NO: 56	FGFRA isoform 4 Amino acid sequence	MRLLALLGVLLSVPGPPVLSLEASEEVELEPCCLAPSLEQQEQELTVALGQPVRLCC GRAERGGHWYKEGSR LAPAGRVRGWRGRLEIASFLPEDAGRYLCLARGSMIVLQNL LITGDSLTSNDDDEPKSHRDPNRRHSYPQAPYVWTHPQRMKKLHAVPAGNTVKFR CPAAGNPTPTIRWLKDGQAFHGENRIGGIRLRHQHWSLVMSVVPVSDRGTYTCLVEN AVGSIRYNYLLDVLESPHRPI LQAGLPANTTAVVGSDELCKVYSDAQPHIQWLK HIVINGSSFGADGFPYVQVLKADINSSEVEVLVLRNVS AEDAGEYTCLAGNSIGLS YQSAWLTVLPGTGRI PHLTCDSLTPAGRTKSPTLQPSLES GSSGKSSSSSLVRGVRLS SSGPALLAGLVSLDLPLDPLWEFPRDRLVLGKPLGEGCFGQVVR AEFAGMDPARPDQ ASTVAVKMLKDNASDKDLADLVSEMEVMKLIGRHKNI INLLGVCTQEGPLYVIVECA AKGNLREFLRARRPPGDLSPDGPRSSEGPLSPFVLVSCAYQVARGMQYLESRKCTH RDLAARNVLVTEDNVMKIADFG LARGVHHIDYKKT SNGRLPVKWM APEALFDRVYT HQSDVWSFGILLWEI FTLGGSYPYGP IVEELFSL LREGHRMDRPPHCPPELYGLMRE CWAAPSQRPTFFKQLVEALDKVLLAVSE EYLDLRLTFGPYPSGGDASSTCSSSDSV FSDHPLPLGSSSPFPGSGVQT
SEQ ID NO: 57	FGFR1 Subdomain V Amino acid sequence	IVEYASKGNLR
SEQ ID NO: 58	FGFR2 Subdomain V Amino acid sequence	IVEYASKGNLR
SEQ ID NO: 59	FGFR3 Subdomain V Amino acid sequence	LVEYAAKGNLR
SEQ ID NO: 60	FGFR4 Subdomain V Amino acid sequence	IVECAAKGNLR

Bold: FGFR subdomain V

Underlined: Cysteine in FGFR4 subdomain V

Although the invention has been described with reference ⁴⁵ to the above examples, it will be understood that modifications and variations are encompassed within the spirit and scope of the invention. Accordingly, the invention is limited only by the following claims.


SEQUENCE LISTING

The patent contains a lengthy sequence listing. A copy of the sequence listing is available in electronic form from the USPTO web site (<https://seqdata.uspto.gov/?pageRequest=docDetail&DocID=US12409177B2>). An electronic copy of the sequence listing will also be available from the USPTO upon request and payment of the fee set forth in 37 CFR 1.19(b)(3).

205

What is claimed is:

1. A compound according to Formula (I)






or an optically pure stereoisomer, solvate, or pharmaceutically acceptable salt thereof,


wherein each of R_1 and R_5 is Cl, each of R_2 and R_4 is OMe, and R_3 is H;

R_6 is $\text{NHCOCH}=\text{CH}_2$

Linker is selected from the group consisting of




substituted phenyl, and



wherein said substituted phenyl comprises one or more substituents of the formula $-\text{O}-(\text{CH}_2)_n-\text{R}_8$; n is an integer selected from 0 to 5;

206


R_8 is selected from the group consisting of



optionally substituted with one or more substituents selected from the group consisting of F, Cl, Br, C_{1-4} alkyl, CF_3 , CHF_2 , and NO_2 ; and

R_7 is selected from the group consisting of H, C_{1-4} alkyl, and COC_{1-4} alkyl.


2. The compound of claim 1, wherein Linker is selected from the group consisting of



and substituted phenyl, wherein said substituted phenyl comprises one or more substituents of the formula $-\text{O}-(\text{CH}_2)_n-\text{R}_8$;

n is an integer selected from 0 to 2;


R_8 is selected from the group consisting of




and R_7 is selected from the group consisting of H and C_{1-4} alkyl.

207

3. A compound selected from the group consisting of





**208**

-continued






209

-continued

**210**

-continued



5. A pharmaceutical formulation comprising a compound according to any one of claims 1, 2, 3, and 4.

6. A method for treating cancer in a subject comprising administering a compound of claim 1 to the subject.

7. The method of claim 6, wherein the cancer is selected from the group consisting of breast, lung, bladder, prostate, ovarian, endometrial, rhabdomyosarcoma, liver, and gastric cancer.

8. The method of claim 7, further comprising administering a chemotherapeutic agent to the subject.

9. The method of claim 8, wherein the compound is administered prior to, simultaneously with, or following the administration of the chemotherapeutic agent.

10. A method for treating cancer in a subject comprising administering a compound of claim 2 to the subject.

11. A method for treating cancer in a subject comprising administering a compound of claim 3 to the subject.

12. A method for treating cancer in a subject comprising administering a compound of claim 4 to the subject.

* * * * *