



US 20250281405A1

(19) **United States**(12) **Patent Application Publication**
NALLANI et al.(10) **Pub. No.: US 2025/0281405 A1**(43) **Pub. Date: Sep. 11, 2025**(54) **POLYMERSOMES COMPRISING A SOLUBLE ENCAPSULATED POLYNUCLEOTIDE AND AN IONIZABLE LIPID AS WELL AS METHODS OF MAKING AND USES THEREOF**(71) Applicant: **ACM BIOLABS PTE LTD**, Singapore (SG)(72) Inventors: **Madhavan NALLANI**, Singapore (SG); **Jian Hang LAM**, Singapore (SG); **Liam MARTIN**, Singapore (SG); **Shrinivas VENKATARAMAN**, Singapore (SG); **Teck Wan CHIA**, Singapore (SG)(73) Assignee: **ACM BIOLABS PTE LTD**, Singapore (SG)(21) Appl. No.: **18/555,094**(22) PCT Filed: **Apr. 12, 2022**(86) PCT No.: **PCT/EP2022/059701**

§ 371 (c)(1),

(2) Date: **Oct. 12, 2023**(30) **Foreign Application Priority Data**

Apr. 12, 2021 (EP) 21167911.3

Publication Classification(51) **Int. Cl.***A61K 9/1273* (2025.01)*A61K 39/00* (2006.01)*A61K 39/145* (2006.01)*A61K 39/215* (2006.01)*A61P 37/04* (2006.01)(52) **U.S. Cl.**CPC *A61K 9/1273* (2013.01); *A61K 39/0005* (2013.01); *A61K 39/0011* (2013.01); *A61K 39/145* (2013.01); *A61K 39/215* (2013.01); *A61P 37/04* (2018.01); *A61K 2039/53* (2013.01)

(57)

ABSTRACT

The present invention relates to polymersomes comprising a soluble encapsulated antigen, wherein said soluble encapsulated antigen is a polynucleotide selected from a DNA molecule or a mRNA molecule, and wherein the polymersome further comprises an ionizable lipid.

Specification includes a Sequence Listing.

IgA titers from swine faecal swabs after oral immunisation with PEDv Spike protein encapsulated polymersomes




Figure 1




Figure 2

SOP Name:	Psomestfast.sop	Dispersant Name:	PBS
File Name:	260417.dts	Dispersant RI:	1.330
Record Number:	1	Viscosity (cP):	0.8984
Material RI:	1.45	Measurement Date und Time:	26 April, 2017 5:12:28 PM
Material Absorbtion:	0.001		
Temperature (°C)	25.0	Duration Used (s):	70
Count Rate (kcps):	174.8	Measurement	3.00
Cell Description:	Disposable micro cuvette (40µl)	Position (mm):	
		Attenuator:	7
Z-Average size (d.nm):	173.1	Size (d.nm):	
Pdi:	0.108	Peak 1:	196.2
Intercept:	0.927	Peak 2:	0.000
Result quality:	Good	Peak 3:	0.000
		% Instensity:	100.0
		St Dev (d.nm):	67.49
			0.000
			0.000




Fig.2A

Figure 2

Formulation	Antigen/Peptide/DNA	Size (d. nm)	PDI ¹
BD21	OVA	173.1	0.108
PDMS-PEG	PEDv S Protein	127.3	0.206
PDMS-PEG/DSPE-PEG	PEDv S Protein	126.4	0.183
PLA-PEG/Asolectin lipids	PEDv S Protein	131.2	0.185
BD21-NH ₂	eGFP-DNA	155.8	0.11

¹ PDI, polydispersity index from dynamic light scattering (DLS)

Fig. 2B

Figure 3




Figure 4




Figure 5




Fig. 5A

Figure 5




Fig. 5B

Figure 5




Fig. 5C

Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12





Figure 13



A.

Figure 13 (cont.)





Figure 14



**POLYMERSOMES COMPRISING A
SOLUBLE ENCAPSULATED
POLYNUCLEOTIDE AND AN IONIZABLE
LIPID AS WELL AS METHODS OF MAKING
AND USES THEREOF**

**CROSS-REFERENCE TO RELATED
APPLICATIONS**

[0001] The present invention is filed under 35 U.S.C. § 371 as the U.S. national phase of International Application No. PCT/EP2022/059701, filed 12 Apr. 2022, which designated the U.S. and claims the right of priority of European patent application 21167911.3 filed with the European Patent Office on 12 Apr. 2021, the entire content of each of which is incorporated herein for all purposes.

SEQUENCE LISTING

[0002] This application includes an electronically submitted sequence listing in .txt format. The .txt file contains a sequence listing entitled "SCH-15000-US1_ST25.txt" created on Apr. 11, 2024 and is 52,530 bytes in size. The sequence listing contained in this .txt file is part of the specification and is hereby incorporated by reference herein in its entirety.

TECHNICAL FIELD

[0003] The present invention relates to polymersomes including oxidation-stable polymersomes comprising a soluble or solubilized encapsulated antigen (e.g., a polynucleotide, e.g., a DNA molecule, a RNA molecule and/or combination/s thereof) and a ionizable lipid.

BACKGROUND OF THE INVENTION

[0004] Although immunization is a well-established process, there are differences in the response level elicited between different immunogens or antigens. For example, membrane proteins form a class of antigens that produce a low response level, which in turn means that large amounts of membrane proteins are required to generate or elicit an immune response to the desired level. Membrane proteins are notoriously difficult to synthesize and are insoluble in water without the presence of a detergent. This makes it expensive and difficult to obtain membrane proteins in sufficient quantity for immunization. Furthermore, membrane proteins require proper folding to function correctly. The immunogenicity of correctly folded native membrane proteins is typically much better than that of their solubilized forms, which may not be folded in a physiologically relevant manner. Thus, even though adjuvants may be used to boost the immunogenicity of such solubilized antigens, it is an inefficient method that does not provide too much of an advantage (e.g., WO2014/077781A1).

[0005] Although transfected cells and lipid-based systems have been used to present membrane protein antigens to increase the chances of isolating antibodies that may be efficient in vivo, these systems are often unstable (e.g., oxidation sensitive), tedious and costly. Moreover, the current state of the art for such membrane protein antigens is to use inactive virus-like particles for immunization.

[0006] On the other hand, vaccines are the most efficient way to prevent diseases, mainly infectious diseases [e.g., Liu et al., 2016]. As of today, most of the licensed vaccines are made of either live or killed viruses. Despite their effective-

ness in generating a humoral response (an antibody mediated response) to prevent viral propagation and entry into cells, safety of such vaccines remains a concern. In the past few decades, scientific advances have helped to overcome such issues by engineering vaccine vectors that are non-replicating recombinant viruses. In parallel, protein based antigens or subunit antigens are explored as safer alternatives. However, such protein based vaccines typically illicit poor immune (both humoral and cellular response). To improve immunogenic properties of antigens, several approaches have been used. For example, microencapsulation of antigens into polymers have been investigated extensively, although it did enhance the immunogenicity, aggregation and denaturing of antigens remain unsolved [e.g., Hilbert et al., 1999]. Furthermore, adjuvants (e.g., oil in water emulsions or polymer emulsions) [e.g., U.S. Pat. No. 9,636,397B2, US2015/0044242 A1] are used together with antigens to elicit a more pronounced humoral and cellular response. Despite these advances, they are less efficient in uptake and cross-presentation. To promote cross-presentation, based on the available information of the immune system during infection by viruses, viral like particles that mimics such properties have been exploited. Synthetic architectures such as liposomes with encapsulated antigens are particularly attractive. Liposomes are unilamellar self-assembling structures made of lipids and, cationic liposomes are more attractive and promising as delivery vehicles because of their efficient uptake by Antigen Presenting Cells (APCs) [e.g., Maji et al., 2016]. Furthermore, it allows to integrate immunomodulators such as Monophosphoryl Lipid A (MPL), CpG oligodeoxynucleotide, that are toll-like receptor (TLR) agonists which stimulate immune cells through receptors. Despite these opportunities of such delivery vehicles, one of the limiting factors is stability of liposomes in the presence of serum components. By PEGylations, loading with high melting temperature lipids, stability issues of liposomes are somewhat reduced with and one such well characterized example being interbilayered-crosslinked multilamellar vesicles (ICMVs), formed by stabilizing multilamellar vesicles with short covalent crosslinks linking lipids [e.g., Moon et al., 2011]. Other nanoparticle architectures have led to successful immunisations using nanodiscs [e.g., Kuai et al., 2017] or pH sensitive particles [e.g., Luo et al., 2017]. But such strategies either still requires adjuvants or are not as efficient outside the prototypical Ovalbumin (OVA) models.

[0007] In addition, polymersomes, offer as a stable alternative for liposomes and they have been used to integrate membrane proteins to elicit immune response [e.g., Quer et al., 2011, WO2014/077781A1]. Protein antigens were also encapsulated in a chemically altered membrane of the polymersome (however oxidation-sensitive membranes) to release antigens and the adjuvants to dendritic cells [e.g., Stano et al., 2013].

[0008] Despite this progress made by the use of polymers, there remains a need to provide for efficient uptake and stable cross-presentation delivery vehicles and methods that overcome, or at least alleviate, the above problems as well as possess an improved functionality inter alia in that they are also capable of eliciting a CD8⁽⁺⁾ T cell-mediated immune response, which is particularly important in treatment and/or prevention of infectious diseases, cancers and autoimmune diseases.

SUMMARY OF THE INVENTION

[0009] The present invention relates to polymersomes comprising a soluble encapsulated antigen, wherein said soluble encapsulated antigen is a polynucleotide (e.g., single or double stranded) selected from a RNA (e.g., mRNA) molecule or a DNA molecule and wherein the polymersome further comprises an ionizable lipid.

[0010] The invention also relates to the use of a polymersome having a diameter of about 120 nm or 140 nm or more comprising a soluble encapsulated antigen, wherein said soluble encapsulated antigen is selected from the group consisting of:

[0011] i) a polypeptide;

[0012] ii) a carbohydrate;

[0013] iii) a polynucleotide, preferably said polynucleotide is not an antisense oligonucleotide, further preferably said polynucleotide is a DNA or mRNA molecule, or

[0014] iv) a combination of i) and/or ii) and/or iii).

[0015] for eliciting an immune response.

[0016] The use of a collection of polymersomes having a mean diameter of about 120 nm, or 140 nm or more, the polymersomes of the collection comprising a soluble encapsulated antigen, wherein said soluble encapsulated antigen is selected from the group consisting of:

[0017] i) a polypeptide;

[0018] ii) a carbohydrate;

[0019] iii) a polynucleotide, preferably said polynucleotide is not an antisense oligonucleotide, further preferably said polynucleotide is a DNA or mRNA molecule, or

[0020] iv) a combination of i) and/or ii) and/or iii).

[0021] for eliciting an immune response.

[0022] Furthermore, in the course of the present invention it was found that providing the polymersomes of the present invention allows soluble (or solubilized) encapsulated (in said polymersomes) antigens to produce a stronger humoral immune response (compared to free antigens with or without adjuvants) as well as elicit a CD8⁽⁺⁾ T cell-mediated immune response. Consequently, an increase in the efficiency of antibody production in a subject is achieved. The increase in the efficiency can be attained with or without the use of adjuvants. Furthermore, the ability of the polymersomes of the present invention to elicit a CD8⁽⁺⁾ T cell-mediated immune response dramatically increases their potential as an immunotherapeutic antigen delivery and presentation system.

[0023] Because soluble (e.g., solubilized) encapsulated antigens presented by polymersomes, the antibodies produced by the use of polymersomes and methods of the present invention would not only have a higher production success rate and higher affinity for their corresponding in vitro or in vivo targets and accordingly improved sensitivity when used in various solution-based antibody applications, but also would make possible to easily raise antibodies to difficult antigens not capable of triggering antibody production by conventional methods using free antigen injections and/or decrease the amount of antigen required for such antibody production procedure thus decreasing the cost of such a production. Furthermore, soluble (e.g., solubilized) encapsulated antigens presented by polymersomes of the present invention are also capable of eliciting a CD8⁽⁺⁾ T cell-mediated immune response, which extends the use of corresponding polymersomes to cell-mediated immunity

and therefore improves their immunotherapeutic- and antigen delivery and presentation potential.

[0024] Therefore, the present application satisfies this demand by provision of oxidation-stable polymersomes that improve immunogenic properties of antigens, methods for their production and compositions comprising such polymersomes, described herein below, characterized in the claims and illustrated by the appended Examples and Figures.

Overview of the Sequence Listing

[0025] As described herein references are made to UniProtKB Accession Numbers (<http://www.uniprot.org/e.g.>, as available in UniProt release 2021_01, published Feb. 10, 2021).

[0026] As described herein references are made to NCBI GenBank Accession Numbers (<https://www.ncbi.nlm.nih.gov/genbank/release/current/e.g.>, as available in Release 242.0, published on Feb. 15, 2021).

[0027] SEQ ID NO: 1 is the amino acid sequence of the tumor neoantigen polypeptide Reps1 P45A derived from the colon cancer MC-38 mouse model.

[0028] SEQ ID NO: 2 is the amino acid sequence of the tumor neoantigen peptide Adpgk R304M derived from the colon cancer MC-38 mouse model.

[0029] SEQ ID NO: 3 is the amino acid sequence of the tumor neoantigen peptide Dpagt1 V213L derived from the colon cancer MC-38 mouse model.

[0030] SEQ ID NO: 4 is the amino acid sequence of the chicken Ovalbumin (OVA), UniProtKB Accession Number: P01012.

[0031] SEQ ID NO: 5 is the amino acid sequence of the influenza A virus (A/New York/38/2016 (H1N1)) hemagglutinin, UniProtKB Accession Number: A0A192ZYK0.

[0032] SEQ ID NO: 6 is the amino acid sequence of the influenza A virus (A/swine/4/Mexico/2009 (H1N1)) hemagglutinin, UniProtKB Accession Number: D2CE65.

[0033] SEQ ID NO: 7 is the amino acid sequence of the influenza A virus (A/Puerto rico/8/1934 (H1N1)) hemagglutinin.

[0034] SEQ ID NO: 8 is the amino acid sequence of the influenza A virus (A/California/07/2009 (H1N1)) hemagglutinin.

[0035] SEQ ID NO: 9 is the amino acid sequence of the tumor neoantigen polypeptide CD8 Trp2 173-196 derived from the melanoma B16-F10 mouse model.

[0036] SEQ ID NO: 10 is the amino acid sequence of the tumor neoantigen polypeptide CD4 M30 Kif18b K739N derived from the melanoma B16-F10 mouse model.

[0037] SEQ ID NO: 11 is the amino acid sequence of the tumor neoantigen polypeptide CD4 M44 Cpsf31 D314N derived from the melanoma B16-F10 mouse model.

[0038] SEQ ID NO: 12 is the amino acid sequence of the soluble portion (amino acid residues 19 to 1327) of the Porcine Epidemic Diarrhea virus (PEDv) Spike protein (S Protein) (UniProtKB Accession number: V5TA78)

[0039] SEQ ID NO: 13 is the amino acid sequence of the S1 region (amino acid residues 19 to 739) of the PEDv Spike protein (S Protein) and

[0040] SEQ ID NO: 14 is the amino acid sequence of the S2 region (amino acid residues 739 to 1327) of the PEDv Spike protein (S Protein)

[0041] SEQ ID NO: 15 is the amino sequence of the enhanced Green Fluorescent Protein (eGFP).

[0042] SEQ ID NO: 16 is the OVA mRNA nucleotide sequence.

BRIEF DESCRIPTION OF THE DRAWINGS

[0043] FIG. 1 shows a schematic view of the immunization with a polymersome of the present invention encapsulating antigens and measuring the humoral and cellular responses.

[0044] FIG. 2 shows the results of dynamic light scattering results for polymersome of the invention. FIG. 2A shows dynamic light scattering plot of OVA encapsulating polymersomes with a monodisperse population of 173.1 nm (diameter). FIG. 2B shows a table of mean diameter (Z average) measured by DLS for different polymersomes encapsulated with different antigens.

[0045] FIG. 3 shows an elution profile of OVA encapsulating polymersome in a size exclusion chromatography.

[0046] FIG. 4 shows sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE) of OVA encapsulating polymersomes.

[0047] FIG. 5 shows the results of encapsulation of a nucleic acid (here the coding gene of enhanced Green Fluorescent Protein (eGFP) in polymersomes of the invention and uptake of the polymers with the encapsulated nucleic acid in cells. FIG. 5A shows fluorescence intensity uptake of different polymersomes inside the cells and eGFP expression based on the DNA encapsulated in the polymersomes, while FIG. 5B and FIG. 5C show fluorescence images of cells that are transfected with DNA encapsulated polymersomes.

[0048] FIG. 6 shows antibody titers from the mice sera that were immunized with PBS, OVA alone, OVA with SAS adjuvant, OVA encapsulating polymersomes without adjuvants. Only ACM encapsulated OVA (herein after “ACM” refers to a polymersome of the present invention) was able to induce an IgG titer.

[0049] FIG. 7 shows antibody titers from the mice sera that were immunized with PBS, HA alone and HA encapsulating polymersomes without adjuvants. Only ACM encapsulated HA (polymersome of the present invention) was able to induce an IgG titer.

[0050] FIG. 8 shows results for a MC-38 mouse tumor model. Tumor volume was monitored in mice immunized with free peptides (open circle), ACM encapsulated peptides (closed square, polymersomes of the present invention) or with ACM encapsulated peptides together with an anti-PD1 antibody treatment (closed triangle). Tumor development was altered by ACM encapsulated peptides (polymersomes of the present invention) over free peptides, which is further potentiated by addition of the anti-PD1 antibody. No adjuvant was added in any of the groups.

[0051] FIG. 9 shows IgG Antibody titres and virus neutralisation (against the strain PEDv USA/Colorado/2013 (CO/13)) from mice sera that were immunised with PBS and with a soluble fragment of the PEDv S Protein that has been encapsulated in a polymersome used as herein (“Polymersomes encapsulated with SPIKE protein”) and in comparison, with killed PED virus (“Killed PEDv”) and ACM polymersomes only (i.e., without any antigen, “polymer-

somes only”). From the IgG Titre of FIG. 9, it is evident that both the ACM encapsulated fragment of the PEDv S Protein and the killed virus induce IgG titres. The virus neutralisation data shows that only the ACM encapsulated PEDv S protein results in a significant neutralising titre while the with a negative control (ACM Polymersomes without any antigen) and killed PED virus show negligible neutralisation.

[0052] FIG. 10 shows virus neutralization data (against the strain PEDv USA/Colorado/2013 (CO/13)) from sera generated from mice after immunization with PBS and different polymersomes (e.g., BD21 (as defined later), PDMS₄₆-PEO₃₇ (marked in the figure just as “PDMS”), PDMS₄₆-PEO₃₇ with DSPE-PEG (distearoylphosphatidylethanolamine [DSPE]polyethylene glycol) as added lipid, polyethylene glycol-polylactic acid (PLA-PEG) with added Asolectin lipids (commercially available phospholipids from soybean) encapsulating either full length soluble PED spike protein (in the case of “BD21 with soluble S protein”) or a S1 or S2 fragment thereof (in all other cases). From FIG. 10, it is evident that the groups of mice immunized with PBS sample do not show any virus neutralization, whereas all polymersome formulations show varying degree of virus neutralization regardless of whether they encapsulate the full length protein or a fragment thereof.

[0053] FIG. 11 shows IgA Antibody titers from swine immunised orally with ACM encapsulated PEDv S protein without the use of adjuvants. Titres are from faecal swabs. As seen in FIG. 11 the titres raises over time, showing that the orally administered polymersomes of the invention with PEDv S protein encapsulated therein, are able to elicit an immune response in the swine.

[0054] FIG. 12 shows a schematic representation of the Porcine Epidemic Diarrhea virus (PEDv) Spike protein (S Protein) (UniProtKB Accession number: V5TA78) and the soluble fragments of SEQ ID NO: 12 (amino acid residues 19 to 1327), SEQ ID NO: 13 (amino acid residues 19 to 739) and SEQ ID NO: 14 (amino acid residues 739 to 1327) that have been used for the encapsulation of soluble S Protein in polymersomes and subsequent immunization/vaccination of mice and pigs as described herein.

[0055] FIG. 13 shows DLS profiles demonstrating superior stability of OVA-mRNA loaded into ACM-polymersomes (BD/DOTAP as reference example) over OVA-mRNA loaded into LNPs (Onpattro composition) at -80°C . A.—LNPs; B.—ACM-polymersomes.

[0056] FIG. 14 shows of DLS profiles of empty polymersomes made with BD/DLIn-MC3-DMA (A.) ionizable lipid compared to empty BD/DOTAP (B.) polymersomes as reference examples demonstrating that both are formed in the same way.

DETAILED DESCRIPTION OF THE INVENTION

[0057] The following detailed description refers to the accompanying Examples and Figures that show, by way of illustration, specific details and embodiments, in which the invention may be practised. These embodiments are described in sufficient detail to enable those skilled in the art to practice the invention. Other embodiments may be utilized such that structural, logical, and eclectic changes may be made without departing from the scope of the invention. Various aspects of the present invention described herein are not necessarily mutually exclusive, as aspects of the present

invention can be combined with one or more other aspects to form new embodiments of the present invention.

[0058] In the present context, polymersomes are vesicles with a polymeric membrane, which are typically, but not necessarily, formed from the self-assembly of dilute solutions of one or more amphiphilic block copolymers, which can be of different types such as diblock and triblock (A-B-A or A-B-C). Polymersomes of the present invention may also be formed of tetra-block or penta-block copolymers. For tri-block copolymers, the central block is often shielded from the environment by its flanking blocks, while di-block copolymers self-assemble into bilayers, placing two hydrophobic blocks tail-to-tail, much to the same effect. In most cases, the vesicular membrane has an insoluble middle layer and soluble outer layers. The driving force for polymersome formation by self-assembly is considered to be the micro-phase separation of the insoluble blocks, which tend to associate in order to shield themselves from contact with water. Polymersomes of the present invention possess remarkable properties due to the large molecular weight of the constituent copolymers. Vesicle formation is favored upon an increase in total molecular weight of the block copolymers. As a consequence, diffusion of the (polymeric) amphiphiles in these vesicles is very low compared to vesicles formed by lipids and surfactants. Owing to this less mobility of polymer chains aggregated in vesicle structure, it is possible to obtain stable polymersome morphologies. Unless expressly stated otherwise, the term “polymersome” and “vesicle”, as used herein, are taken to be analogous and may be used interchangeably. Importantly, a polymersome of the invention can be formed from either one kind of block copolymers or from two or more kinds of block copolymers, meaning a polymersome can also be formed from a mixture of polymersomes and thus can contain two or more block copolymers. In some aspects, the polymersome of the present invention is oxidation-stable.

[0059] In some aspects, the present invention relates to a method for eliciting an immune response to a soluble (e.g., solubilized) encapsulated antigen in a subject. The method is suitable for injecting the subject with a composition comprising a polymersome (e.g., carrier or vehicle) having a membrane (e.g., circumferential membrane) of an amphiphilic polymer. The composition comprises a soluble (e.g., solubilized) antigen encapsulated by the membrane (e.g., circumferential membrane) of the amphiphilic polymer of the polymersome of the present invention. The antigen may be one or more of the following: i) a polypeptide; ii) a carbohydrate; iii) a polynucleotide (e.g., said polynucleotide is not an antisense oligonucleotide, preferably said polynucleotide is a DNA or messenger RNA (mRNA) molecule) or a combination of i) and/or ii) and/or iii).

[0060] In some further aspects, the present invention relates to polymersomes capable of eliciting a CD8(+) T cell-mediated immune response.

[0061] In some aspects, the present invention relates to polymersomes capable of targeting of lymph node-resident macrophages and/or B cells. Exemplary non-limiting targeting mechanisms envisaged by the present invention include: i) delivery of encapsulated antigens (e.g., polypeptides, etc.) to dendritic cells (DCs) for T cell activation (CD4 and/or CD8). Another one is: ii) delivery of whole folded antigens (e.g., proteins, etc.) that will be route to DC and will also trigger a titer (B cells).

[0062] In some aspects, the present invention relates to polymersomes encapsulating an antigen selected from a group consisting of: i) a self-antigen, ii) a non-self antigen, iii) a non-self immunogen and iv) a self-immunogen. Accordingly, the products and methods of the present invention are suitable for uses in settings (e.g., clinical settings) of induced tolerance, e.g., when targeting an autoimmune disease.

[0063] In some aspects, the present invention relates to polymersomes of the present invention comprising a lipid polymer.

[0064] The polymersomes of the present invention can also have co-encapsulated (i.e. encapsulated in addition to the antigen) one or more adjuvants. Examples of adjuvants include synthetic oligodeoxynucleotides (ODNs) containing unmethylated CpG motifs which can trigger cells that express Toll-like receptor 9 (including human plasmacytoid dendritic cells and B cells) to mount an innate immune response characterized by the production of Th1 and proinflammatory cytokines, cytokines such as Interleukin-1, Interleukin-2 or Interleukin-12, keyhole limpet hemocyanin (KLH), serum albumin, bovine thyroglobulin, or soybean trypsin inhibitor, too name only a few illustrative examples.

[0065] The polymersomes of the present invention can be of any size as long as the polymersomes are able to elicit an immune response. For example, the polymersomes may have a diameter of greater than 70 nm. The diameter of the polymersomes may range from about 100 nm to about 1 μ m, or from about 100 nm to about 750 nm, or from about 100 nm to about 500 nm. The diameter of the polymersome may further range from about 125 nm to about 175 nm or, from about 125 nm to about 250 nm, from about 140 nm to about 240 nm, from about 150 nm to about 235 nm, from about 170 nm to about 230 nm, or from about 220 nm to about 180 nm, or from about 190 nm to about 210 nm. The diameter of the polymersomes may, for example, about 200 nm; about 205 nm or about 210 nm. When used as a collection to elicit an immune response, the collection of polymersomes is typically a monodisperse population. The mean diameter of the used collection/population of polymersomes is typically above 70 nm, or above 120 nm, or above 125 nm, or above 130 nm, or above 140 nm, or above 150 nm, or above 160 nm, or for above 170 nm, or above 180 nm, or above 190 nm (cf. also FIG. 2 in this respect). The mean diameter of the collection of polymersomes may, for example, also in range of the individual polymersomes mentioned above, meaning the mean diameter of the collection of polymersomes may be in the range of 100 nm to about 1 μ m, or in the range of about 100 nm to about 750 nm, or in the range of about 100 nm to about 500 nm, or in the range from about 125 nm to about 250 nm, from about 140 nm to about 240 nm, from about 150 nm to about 235 nm, from about 170 nm to about 230 nm, or from about 220 nm to about 180 nm, or from about 190 nm to about 210 nm. The mean diameter of the collection of polymersomes may, for example, also be about 200 nm; about 205 nm or about 210 nm. The diameter can, for example, be determined by a dynamic light scattering (DLS) instrument using Z-average (d, nm), a preferred DLS parameter. Z-average size is the intensity weighted harmonic mean particle diameter (cf. Examples 1 and 2). In this context, it is noted that according to U.S. Pat. No. 8,323,696 of Hubbel et al, a collection of polymersomes should have a mean diameter of less than 70 nm to be able to elicit immune response. Similarly, Stano et al, supra, 2013, while

wanting to use smaller polymersome, used, due to technical constraints, polymersomes having a diameter of 125 nm+/-15 nm to elicit an immune response. Thus, it is surprising that a population/collection of polymersomes of the present invention with a mean diameter of, for example, than more 150 nm are able to induce both a cellular and a humoral immune response (cf. Example section). Such a collection of polymersomes may be in a form suitable for eliciting an immune response, for example, by injection or oral administration.

[0066] In some aspects, the present invention relates to compositions of the present invention suitable for intradermal, intraperitoneal, subcutaneous, intravenous, or intramuscular injection, or non-invasive administration of an antigen of the present invention, for example, oral administration or nasal administration. The composition may include a polymersome (e.g., carrier) of the present invention having a membrane (e.g., circumferential membrane) of an amphiphilic polymer. The composition further includes a soluble (e.g., solubilized) antigen encapsulated by the membrane of the amphiphilic polymer of the polymersome. The compositions of the present invention may be used for therapeutic purposes (for example, treatment of a subject suffering from a disease or for preventing from suffering from a disease, for example, by means of vaccination) or be used in antibody discovery, vaccine discovery, or targeted delivery.

[0067] In some aspects, polymersomes of the present invention have hydroxyl groups on their surface. In some further aspects, polymersomes of the present invention do not have hydroxyl groups on their surface.

[0068] In the present context, the term “encapsulated” means enclosed by a membrane (e.g., membrane of the polymersome of the present invention, e.g., embodied inside the lumen of said polymersome). With reference to an antigen the term “encapsulated” further means that said antigen is neither integrated into-nor covalently bound to-nor conjugated to said membrane (e.g., of a polymersome of the present invention). With reference to compartmentalization of the vesicular structure of polymersome as described herein the term “encapsulated” means that the inner vesicle is completely contained inside the outer vesicle and is surrounded by the vesicular membrane of the outer vesicle. The confined space surrounded by the vesicular membrane of the outer vesicle forms one compartment. The confined space surrounded by the vesicular membrane of the inner vesicle forms another compartment.

[0069] In the present context, the term “antigen” means any substance that may be specifically bound by components of the immune system. Only antigens that are capable of eliciting (or evoking or inducing) an immune response are considered immunogenic and are called “immunogens”. Exemplary non-limiting antigens are polypeptides derived from a soluble portion of proteins, hydrophobic polypeptides rendered soluble for encapsulation as well as aggregated polypeptides that are soluble as aggregates. The antigen may originate from within the body (“self-antigen”) or from the external environment (“non-self”).

[0070] Membrane proteins form a class of antigens that typically produce a low immune response level. Of specific interest, soluble (e.g., solubilized) membrane proteins (MPs) and membrane-associated peptides (MAPs) and fragments (i.e., portions) thereof (e.g., the antigens mentioned herein) are encapsulated by a polymersome, which may allow them

to be folded in a physiologically relevant manner. This greatly boosts the immunogenicity of such antigens so that when compared to free antigens, a smaller amount of the corresponding antigen can be used to produce the same level of the immune response. Furthermore, the larger size of the polymersomes (compared to free membrane proteins) allows them to be detected by the immune system more easily.

[0071] In the present context, the term “B16 peptide” refers to any neoantigen polypeptide derived from the spontaneous C57BL/6-derived B16 melanoma model (e.g., melanoma B16-F10 mouse model). Non-limiting examples thereof include the peptides of SEQ ID NO: 9, 10 and 11.

[0072] In the present context, the term “MC38 peptide” refers to any neoantigen polypeptide derived from the colon cancer MC38 mouse model. Non-limiting examples thereof include the peptides of SEQ ID NO: 1, 2 and 3.

[0073] In the present context, the term “Influenza hemagglutinin (HA)” refers to a glycoprotein found on the surface of influenza viruses. HA has at least 18 different antigens, which are all within the scope of the present invention. These subtypes are named H1 through H18. Non-limiting examples of “Influenza hemagglutinin (HA)” subtype H1 include the polypeptides of SEQ ID NOs: 5, 6, 7 and 8.

[0074] In the present context, the term “Swine Influenza hemagglutinin (HA)” refers to a glycoprotein found on the surface of swine influenza viruses, which is a family of influenza viruses endemic in pigs. Non-limiting examples of “Swine Influenza hemagglutinin (HA)” include subtype H1 of SEQ ID NO: 6.

[0075] In the present context, the term “PEDv S Protein” refers to SPIKE glycoprotein present on the surface of Porcine epidemic diarrhea virus (PEDV), which is a family of coronavirus in pigs. Non-limiting examples of soluble “PEDv S Protein” as may be used in the present invention include the entire soluble fragment consisting of the S1 and S2 region having the amino acid sequence of SEQ ID NO: 12, the soluble fragment of the S1 region of SEQ ID NO: 13, or the soluble fragment of the S2 region of SEQ ID NO: 14, of the Porcine Epidemic Diarrhea virus (PEDV) Spike protein (S Protein) (UniProtKB Accession number: V5TA78). It is of course also possible to use shorter fragments of the entire soluble fragment of the S1 and the S2 region or of either of the S1 or S2 regions alone (cf. FIG. 12 in this respect) It is of course also possible to use in polymersomes of the present invention a fragment that contains part of the S1 and part of the S2, say for example, amino acids 500 to 939 of the deposited sequence of the Spike protein. It is also noted here that a polymersome of the present invention may have encapsulated one or more different soluble fragments of the Spike protein, for example, the S1 region, the S2 region and/or the entire S1 and S2 region. In illustrative embodiments of a polymersomes of the invention, it has encapsulated therein one type of soluble fragments (for example, only the S1 region), two different types of soluble fragments (for example, the S1 and S2 region), three different types of soluble fragments (the S1 region, the S2 region and the entire soluble fragment of S1 and S2 of SEQ ID NO: 12 (amino acid residues 19 to 1327)) or even four different types of fragments (for example, the S1 region, the S2 region, the entire soluble fragment of S1 and S2 of SEQ ID NO: 12 (amino acid residues 19 to 1327) and as fourth type, the above-mentioned fragment that contains part of the S1 and part of the S2, say for example, amino acids 500 to

939 of the Spike protein sequence). It is also noted here that a polymersome of the present invention having encapsulated one or more different soluble fragments of the Spike protein are used in one preferred embodiment as oral vaccine against the Porcine Epidemic Diarrhea virus.

[0076] In the present context, the term “oxidation-stable” refers to a measure of polymersomes (or the corresponding polymers or membranes) resistance to oxidation, for example, using the method described by Scott et al., 2012. In this method a polymersome with an encapsulated antigen is incubated in a 0.5% solution of hydrogen peroxide and the amount of free (released) antigen can be quantified with UV/fluorescence HPLC. Polymersomes which release a substantial or all of the encapsulated antigen under these oxidizing conditions are considered to be oxidation sensitive. Another method of determining whether a block-copolymer and thus the resulting polymersome is oxidation stable or oxidation-sensitive is described in column 16 of U.S. Pat. No. 8,323,696. According to this method, polymers with functional groups that are oxidation-sensitive will be chemically altered by mild oxidizing agents, with a test for the same being enhanced solubility to 10% hydrogen peroxide for 20 h in vitro. As, for example, poly(propylene sulfide) (PPS) is an oxidation-sensitive polymer (see, for example, Scott et al 2012, supra and U.S. Pat. No. 8,323,696) PPS can serve as a reference to determine whether a polymer of interest and the respective polymersome of interest is oxidation-sensitive or oxidation stable. If, for example, the same or a higher amount of antigen, or about 90% or more of the amount, or about 80% or more, or about 70% or more, or about 60% or more is released from polymersomes of interest as it is from a PPS polymersome that has encapsulated therein the same antigen, then the polymersome is considered oxidation sensitive. If about only 0.5% or less, or about only 1.0% or less, or about 2% or less, or about 5% of less, or about 10% or less, or about 20% or less, or about 30% or less, or about 40% or less or about 50% or less of antigen is released from polymersomes of interest as it is from a PPS polymersome that has encapsulated therein the same antigen, then the polymersome is considered oxidation-stable. Thus, in line with this, PPS polymersomes as described in U.S. Pat. No. 8,323,696 or PPS-bl-PEG polymersomes, e.g., made from poly(propylene sulfide) (PPS) and poly(ethylene glycol) (PEG) as components as described in Stano et al, are not oxidation-stable polymersomes within the meaning of the present invention. Similarly, PPS30-PEG17 polymersomes are not oxidation-stable polymersomes within the meaning of the present invention. Other non-limiting examples of measuring oxidation stability include measurement of stability in the presence of serum components (e.g., mammalian serum, e.g., human serum components) or stability inside an endosome, for example.

[0077] In the present context, the term “reduction-stable” refers to a measure of polymersome resistance to reduction in a reducing environment.

[0078] In the present context, the term “serum” refers to blood plasma from which the clotting proteins have been removed.

[0079] In the present context, the term “oxidation-independent release” refers to a release of the polymersome content without or essentially without oxidation of the polymers forming the polymersomes.

[0080] The term “polypeptide” is equally used herein with the term “protein”. Proteins (including fragments thereof, preferably biologically active fragments, and peptides, usually having less than 30 amino acids) comprise one or more amino acids coupled to each other via a covalent peptide bond (resulting in a chain of amino acids). The term “polypeptide” as used herein describes a group of molecules, which, for example, consist of more than 30 amino acids. Polypeptides may further form multimers such as dimers, trimers and higher oligomers, i.e. consisting of more than one polypeptide molecule. Polypeptide molecules forming such dimers, trimers etc. may be identical or non-identical. The corresponding higher order structures of such multimers are, consequently, termed homo- or heterodimers, homo- or heterotrimers etc. An example for a heteromultimer is an antibody molecule, which, in its naturally occurring form, consists of two identical light polypeptide chains and two identical heavy polypeptide chains. The terms “polypeptide” and “protein” also refer to naturally modified polypeptides/proteins wherein the modification is effected e.g. by post-translational modifications like glycosylation, acetylation, phosphorylation and the like. Such modifications are well known in the art.

[0081] In the present context, the term “carbohydrates” refers to compounds such as aldoses and ketoses having the stoichiometric formula $C_n(H_2O)_n$ (e.g., hence “hydrates of carbon”). The generic term “carbohydrate” includes, but is not limited to, monosaccharides, oligosaccharides and polysaccharides as well as substances derived from monosaccharides by reduction of the carbonyl group (alditols), by oxidation of one or more terminal groups to carboxylic acids, or by replacement of one or more hydroxy group(s) by a hydrogen atom, an amino group, thiol group or similar groups. It also includes derivatives of these compounds.

[0082] In the present context, the term “polynucleotide” (also “nucleic acid”, which can be used interchangeably with the term “polynucleotide”) refers to macromolecules made up of nucleotide units which e.g., can be hydrolysable into certain pyrimidine or purine bases (usually adenine, cytosine, guanine, thymine, uracil), d-ribose or 2-deoxy-d-ribose and phosphoric acid. Non-limiting examples of “polynucleotide” include DNA molecules (e.g. cDNA or genomic DNA), RNA (mRNA), combinations thereof or hybrid molecules comprised of DNA and RNA. The nucleic acids can be double- or single-stranded and may contain double- and single-stranded fragments at the same time. Most preferred are double stranded DNA molecules and mRNA molecules.

[0083] In the present context, the term “antisense oligonucleotide” refers to a nucleic acid polymer, at least a portion of which is complementary to a nucleic acid which is present in a normal cell or in an affected cell. Exemplary “antisense oligonucleotide” include antisense RNA, siRNA, RNAi.

[0084] In the present context, the term “CD8(+) T cell-mediated immune response” refers to the immune response mediated by cytotoxic T cells (also known as TC, cytotoxic T lymphocyte, CTL, T-killer cells, cytolytic T cells, CD8(+) T-cells or killer T cells). Example of cytotoxic T cells include, but are not limited to antigen-specific effector CD8(+) T cells. In order for the T-cell receptors (TCR) to bind to the class I MHC molecule, the former must be accompanied by a glycoprotein called CD8, which binds to the constant portion of the class I MHC molecule. Therefore, these T cells are called CD8(+) T cells. Once activated, the

TC cell undergoes “clonal expansion” with the help of the cytokine Interleukin-2 (IL-2), which is a growth and differentiation factor for T cells. This increases the number of cells specific for the target antigen that can then travel throughout the body in search of antigen-positive somatic cells.

[0085] In the present context, the term “clonal expansion of antigen-specific CD8(+) T cells” refers to an increase in the number of CD8(+) T cells specific for the target antigen.

[0086] In the present context, the term “cellular immune response” refers to an immune response that does not involve antibodies, but rather involves the activation of phagocytes, antigen-specific cytotoxic T-lymphocytes, and the release of various cytokines in response to an antigen.

[0087] In the present context, the term “cytotoxic phenotype of antigen-specific CD8(+) T cells” refers to the set of observable characteristics of antigen-specific CD8(+) T cells related to their cytotoxic function.

[0088] In the present context, the term “lymph node-resident macrophages” refers to macrophages, which are large white blood cell that is an integral part of our immune system that use the process of phagocytosis to engulf particles and then digest them, present in lymph nodes that are small, bean-shaped glands throughout the body.

[0089] In the present context, the term “humoral immune response” refers to an immune response mediated by macromolecules found in extracellular fluids such as secreted antibodies, complement proteins, and certain antimicrobial peptides. Its aspects involving antibodies are often called antibody-mediated immunity.

[0090] In the present context, the term “B cells”, also known as B lymphocytes, are a type of white blood cell of the lymphocyte subtype. They function in the humoral immunity component of the adaptive immune system by secreting antibodies.

[0091] An “antibody” when used herein is a protein comprising one or more polypeptides (comprising one or more binding domains, preferably antigen binding domains) substantially or partially encoded by immunoglobulin genes or fragments of immunoglobulin genes. The term “immunoglobulin” (Ig) is used interchangeably with “antibody” herein. The recognized immunoglobulin genes include the kappa, lambda, alpha, gamma, delta, epsilon and mu constant region genes, as well as myriad immunoglobulin variable region genes. In particular, an “antibody” when used herein, is typically tetrameric glycosylated proteins composed of two light (L) chains of approximately 25 kDa each and two heavy (H) chains of approximately 50 kDa each. Two types of light chain, termed lambda and kappa, may be found in antibodies. Depending on the amino acid sequence of the constant domain of heavy chains, immunoglobulins can be assigned to five major classes: A, D, E, G, and M, and several of these may be further divided into subclasses (isotypes), e.g., IgG1, IgG2, IgG3, IgG4, IgA1, and IgA2, with IgG being preferred in the context of the present invention. An antibody relating to the present invention is also envisaged which has an IgE constant domain or portion thereof that is bound by the Fc epsilon receptor 1. An IgM antibody consists of 5 of the basic heterotetramer unit along with an additional polypeptide called a J chain, and contains 10 antigen binding sites, while IgA antibodies comprise from 2-5 of the basic 4-chain units which can polymerize to form polyvalent assemblages in combination with the J chain. In the case of IgGs, the 4-chain unit is

generally about 150,000 daltons. Each light chain includes an N-terminal variable (V) domain (VL) and a constant (C) domain (CL). Each heavy chain includes an N-terminal V domain (VH), three or four C domains (CHs), and a hinge region. The constant domains are not involved directly in binding an antibody to an antigen, but can exhibit various effector functions, such as participation of the antibody dependent cellular cytotoxicity (ADCC). If an antibody should exert ADCC, it is preferably of the IgG1 subtype, while the IgG4 subtype would not have the capability to exert ADCC.

[0092] The term “antibody” also includes, but is not limited to, but encompasses monoclonal, monospecific, poly- or multi-specific antibodies such as bispecific antibodies, humanized, camelized, human, single-chain, chimeric, synthetic, recombinant, hybrid, mutated, grafted, and in vitro generated antibodies, with chimeric or humanized antibodies being preferred. The term “humanized antibody” is commonly defined for an antibody in which the specificity encoding CDRs of HC and LC have been transferred to an appropriate human variable frameworks (“CDR grafting”). The term “antibody” also includes scFvs, single chain antibodies, diabodies or tetrabodies, domain antibodies (dAbs) and nanobodies. In terms of the present invention, the term “antibody” shall also comprise bi-, tri- or multimeric or bi-, tri- or multifunctional antibodies having several antigen binding sites.

[0093] Furthermore, the term “antibody” as employed in the invention also relates to derivatives of the antibodies (including fragments) described herein. A “derivative” of an antibody comprises an amino acid sequence which has been altered by the introduction of amino acid residue substitutions, deletions or additions. Additionally, a derivative encompasses antibodies which have been modified by a covalent attachment of a molecule of any type to the antibody or protein. Examples of such molecules include sugars, PEG, hydroxyl-, ethoxy-, carboxy- or amine-groups but are not limited to these. In effect the covalent modifications of the antibodies lead to the glycosylation, pegylation, acetylation, phosphorylation, amidation, without being limited to these.

[0094] The antibody relating to the present invention is preferably an “isolated” antibody. “Isolated” when used to describe antibodies disclosed herein, means an antibody that has been identified, separated and/or recovered from a component of its production environment. Preferably, the isolated antibody is free of association with all other components from its production environment. Contaminant components of its production environment, such as that resulting from recombinant transfected cells, are materials that would typically interfere with diagnostic or therapeutic uses for the polypeptide, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the antibody will be purified (1) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (2) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Ordinarily, however, an isolated antibody will be prepared by at least one purification step.

[0095] The term “essentially non-immunogenic” means that the block copolymer or amphiphilic polymer of the present invention does not elicit an adaptive immune

response, i.e., in comparison to an encapsulated immunogen, the block copolymer or amphiphilic polymer shows an immune response of less than 30%, preferably 20%, more preferably 10%, particularly preferably less than 9, 8, 7, 6 or 5%.

[0096] The term “essentially non-antigenic” means that the block copolymer or amphiphilic polymer of the present invention does not bind specifically with a group of certain products that have adaptive immunity (e.g., T cell receptors or antibodies), i.e., in comparison to an encapsulated antigen the block copolymer or amphiphilic polymer shows binding of less than 30%, preferably 20%, more preferably 10%, particularly preferably less than 9, 8, 7, 6 or 5%.

[0097] Typically, binding is considered specific when the binding affinity is higher than 10^{-6} M. Preferably, binding is considered specific when binding affinity is about 10^{-11} to 10^{-8} M (KD), preferably of about 10^{-11} to 10^{-9} M. If necessary, nonspecific binding can be reduced without substantially affecting specific binding by varying the binding conditions.

[0098] The term “amino acid” or “amino acid residue” typically refers to an amino acid having its art recognized definition such as an amino acid selected from the group consisting of: alanine (Ala or A); arginine (Arg or R); asparagine (Asn or N); aspartic acid (Asp or D); cysteine (Cys or C); glutamine (Gln or Q); glutamic acid (Glu or E); glycine (Gly or G); histidine (His or H); isoleucine (Ile or I); leucine (Leu or L); lysine (Lys or K); methionine (Met or M); phenylalanine (Phe or F); pro line (Pro or P); serine (Ser or S); threonine (Thr or T); tryptophan (Trp or W); tyrosine (Tyr or Y); and valine (Val or V), although modified, synthetic, or rare amino acids may be used as desired. Generally, amino acids can be grouped as having a nonpolar side chain (e.g., Ala, Cys, Ile, Leu, Met, Phe, Pro, Val); a negatively charged side chain (e.g., Asp, Glu); a positively charged sidechain (e.g., Arg, His, Lys); or an uncharged polar side chain (e.g., Asn, Cys, Gln, Gly, His, Met, Phe, Ser, Thr, Trp, and Tyr).

[0099] “Polyclonal antibodies” or “polyclonal antisera” refer to immune serum containing a mixture of antibodies specific for one (monovalent or specific antisera) or more (polyvalent antisera) antigens which may be prepared from the blood of animals immunized with the antigen or antigens.

[0100] Furthermore, the term “antibody” as employed in the invention also relates to derivatives or variants of the antibodies described herein which display the same specificity as the described antibodies. Examples of “antibody variants” include humanized variants of non-human antibodies, “affinity matured” antibodies (see, e.g. Hawkins et al. *J. Mol. Biol.* 254, 889-896 (1992) and Lowman et al., *Biochemistry* 30, 10832-10837 (1991)) and antibody mutants with altered effector function(s) (see, e.g., U.S. Pat. No. 5,648,260).

[0101] The term “monoclonal antibody” as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations and/or post-translation modifications (e.g., isomerizations, amidations) that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to conventional (polyclonal) antibody preparations which typically include different antibodies

directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they are synthesized by the hybridoma culture, uncontaminated by other immunoglobulins. The modifier “monoclonal” indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler et al., *Nature*, 256:495 (1975), or may be made by recombinant DNA methods (see, e.g., U.S. Pat. No. 4,816,567). The “monoclonal antibodies” may also be isolated from phage antibody libraries using the techniques described in Clackson et al., *Nature*, 352:624-628 (1991) and Marks et al., *J. Mol. Biol.*, 222:581-597 (1991), for example.

[0102] The monoclonal antibodies herein specifically include “chimeric” antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is (are) identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Pat. No. 4,816,567; Morrison et al., *Proc. Natl. Acad. Sci. USA*, 81:6851-6855 (1984)). Chimeric antibodies of interest herein include “primitized” antibodies comprising variable domain antigen-binding sequences derived from a non-human primate (e.g., Old World Monkey, Ape etc.) and human constant region sequences.

[0103] “Humanized” forms of non-human (e.g., murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F (ab')₂ or other antigen-binding subsequences of antibodies) of mostly human sequences, which contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a hypervariable region (also CDR) of the recipient are replaced by residues from a hypervariable region of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity, and capacity. In some instances, Fv framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, “humanized antibodies” as used herein may also comprise residues which are found neither in the recipient antibody nor the donor antibody. These modifications are made to further refine and optimize antibody performance. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones et al., *Nature*, 321:522-525 (1986); Reichmann et al., *Nature*, 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.*, 2:593-596 (1992).

[0104] The term “human antibody” includes antibodies having variable and constant regions corresponding substantially to human germline immunoglobulin sequences known in the art, including, for example, those described by Kabat et al. (See Kabat, et al. (1991) loc. cit.). The human

antibodies of the invention may include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or site-specific mutagenesis *in vitro* or by somatic mutation *in vivo*), for example in the CDRs, and in particular, CDR3. The human antibody can have at least one, two, three, four, five, or more positions replaced with an amino acid residue that is not encoded by the human germline immunoglobulin sequence.

[0105] As used herein, “*in vitro* generated antibody” refers to an antibody where all or part of the variable region (e.g., at least one CDR) is generated in a non-immune cell selection (e.g., an *in vitro* phage display, protein chip or any other method in which candidate sequences can be tested for their ability to bind to an antigen). This term thus preferably excludes sequences generated by genomic rearrangement in an immune cell.

[0106] A “bispecific” or “bifunctional antibody” is an artificial hybrid antibody having two different heavy/light chain pairs and two different binding sites. Bispecific antibodies can be produced by a variety of methods including fusion of hybridomas or linking of Fab' fragments. See, e.g., Songsivilai & Lachmann, *Clin. Exp. Immunol.* 79:315-321 (1990); Kostelny et al., *J. Immunol.* 148, 1547-1553 (1992). In one embodiment, the bispecific antibody comprises a first binding domain polypeptide, such as a Fab' fragment, linked via an immunoglobulin constant region to a second binding domain polypeptide.

[0107] Numerous methods known to those skilled in the art are available for obtaining antibodies or antigen-binding fragments thereof. For example, antibodies can be produced using recombinant DNA methods (U.S. Pat. No. 4,816,567). Monoclonal antibodies may also be produced by generation of hybridomas (see e.g., Kohler and Milstein (1975) *Nature*, 256:495-499) in accordance with known methods. Hybridomas formed in this manner are then screened using standard methods, such as enzyme-linked immunosorbent assay (ELISA) and surface plasmon resonance (BIACORE™) analysis, to identify one or more hybridomas that produce an antibody that specifically binds with a specified antigen. Any form of the specified antigen may be used as the immunogen, e.g., recombinant antigen, naturally occurring forms, any variants or fragments thereof, as well as antigenic peptide thereof.

[0108] In addition to the use of display libraries, the specified antigen can be used to immunize a non-human animal, e.g., a rodent, e.g., a mouse, hamster, or rat. In one embodiment, the non-human animal includes at least a part of a human immunoglobulin gene. For example, it is possible to engineer mouse strains deficient in mouse antibody production with large fragments of the human Ig loci. Using the hybridoma technology, antigen-specific monoclonal antibodies derived from the genes with the desired specificity may be produced and selected. See, e.g., XENOMOUSE™, Green et al. (1994) *Nature Genetics* 7:13-21, US 2003-0070185, WO 96/34096, and WO96/33735.

[0109] In another embodiment, a monoclonal antibody is obtained from the non-human animal, and then modified, e.g., humanized, deimmunized, chimeric, may be produced using recombinant DNA techniques known in the art. A variety of approaches for making chimeric antibodies have been described. See e.g., Morrison et al., *Proc. Natl. Acad. Sci. U.S.A.* 81:6851, 1985; Takeda et al., *Nature* 314:452, 1985; Cabilly et al., U.S. Pat. No. 4,816,567; Boss et al., U.S. Pat. No. 4,816,397; Tanaguchi et al., EP 171496; EP

173494, GB 2177096. Humanized antibodies may also be produced, for example, using transgenic mice that express human heavy and light chain genes, but are incapable of expressing the endogenous mouse immunoglobulin heavy and light chain genes. Winter describes an exemplary CDR-grafting method that may be used to prepare the humanized antibodies described herein (U.S. Pat. No. 5,225,539). All of the CDRs of a particular human antibody may be replaced with at least a portion of a non-human CDR, or only some of the CDRs may be replaced with non-human CDRs. It is only necessary to replace the number of CDRs required for binding of the humanized antibody to a predetermined antigen.

[0110] Humanized antibodies or fragments thereof can be generated by replacing sequences of the Fv variable domain that are not directly involved in antigen binding with equivalent sequences from human Fv variable domains. Exemplary methods for generating humanized antibodies or fragments thereof are provided by Morrison (1985) *Science* 229:1202-1207; by Oi et al. (1986) *BioTechniques* 4:214; and by U.S. Pat. Nos. 5,585,089; 5,693,761; 5,693,762; 5,859,205; and 6,407,213. Those methods include isolating, manipulating, and expressing the nucleic acid sequences that encode all or part of immunoglobulin Fv variable domains from at least one of a heavy or light chain. Such nucleic acids may be obtained from a hybridoma producing an antibody against a predetermined target, as described above, as well as from other sources. The recombinant DNA encoding the humanized antibody molecule can then be cloned into an appropriate expression vector.

[0111] In certain embodiments, a humanized antibody is optimized by the introduction of conservative substitutions, consensus sequence substitutions, germline substitutions and/or backmutations. Such altered immunoglobulin molecules can be made by any of several techniques known in the art, (e.g., Teng et al., *Proc. Natl. Acad. Sci. U.S.A.*, 80:7308-7312, 1983; Kozbor et al., *Immunology Today*, 4:7279, 1983; Olsson et al., *Meth. Enzymol.*, 92:3-16, 1982), and may be made according to the teachings of WO 92/06193 or EP 239400).

[0112] An antibody or fragment thereof may also be modified by specific deletion of human T cell epitopes or “deimmunization” by the methods disclosed in WO 98/52976 and WO 00/34317. Briefly, the heavy and light chain variable domains of an antibody can be analyzed for peptides that bind to MHC Class II; these peptides represent potential T-cell epitopes (as defined in WO 98/52976 and WO 00/34317). For detection of potential T-cell epitopes, a computer modeling approach termed “peptide threading” can be applied, and in addition a database of human MHC class II binding peptides can be searched for motifs present in the VH and VL sequences, as described in WO 98/52976 and WO 00/34317. These motifs bind to any of the 18 major MHC class II DR allotypes, and thus constitute potential T cell epitopes. Potential T-cell epitopes detected can be eliminated by substituting small numbers of amino acid residues in the variable domains, or preferably, by single amino acid substitutions. Typically, conservative substitutions are made. Often, but not exclusively, an amino acid common to a position in human germline antibody sequences may be used. Human germline sequences, e.g., are disclosed in Tomlinson, et al. (1992) *J. Mol. Biol.* 227:776-798; Cook, G. P. et al. (1995) *Immunol. Today* Vol. 16 (5): 237-242; Chothia, et al. (1992) *J. Mol. Biol.* 227:

799-817; and Tomlinson et al. (1995) EMBO J. 14:4628-4638. The V BASE directory provides a comprehensive directory of human immunoglobulin variable region sequences (compiled by Tomlinson, L. A. et al. MRC Centre for Protein Engineering, Cambridge, UK). These sequences can be used as a source of human sequence, e.g., for framework regions and CDRs. Consensus human framework regions can also be used, e.g., as described in U.S. Pat. No. 6,300,064.

[0113] “Effector cells”, preferably human effector cells are leukocytes which express one or more FcRs and perform effector functions. Preferably, the cells express at least FcγRm and perform ADCC effector function. Examples of human leukocytes which mediate ADCC include peripheral blood mononuclear cells (PBMC), natural killer (NK) cells, monocytes, cytotoxic T cells and neutrophils. The effector cells may be isolated from a native source, e.g., blood.

[0114] Techniques for production of antibodies, including polyclonal, monoclonal, humanized, bispecific and hetero-conjugate antibodies are known in the art, some of which are exemplified below.

1) Polyclonal Antibodies.

[0115] Polyclonal antibodies are generally raised in animals by multiple subcutaneous (sc) or intraperitoneal (ip) injections of the relevant antigen (e.g., encapsulated in a polymersome) and an adjuvant. It may be useful to conjugate the relevant antigen to a protein that is immunogenic in the species to be immunized, e.g., keyhole limpet hemocyanin (KLH), serum albumin, bovine thyroglobulin, or soybean trypsin inhibitor, using a bifunctional or derivatizing agent, e.g., maleimidobenzoyl sulfo succinimide ester (conjugation through cysteine residues), N-hydroxysuccinimide (through lysine residues), glutaraldehyde, succinic anhydride. Examples of adjuvants which may be employed include Freund’s complete adjuvant and MPL-TDM adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate). The immunization protocol may be selected by one skilled in the art without undue experimentation.

[0116] For example, the animals are immunized against the antigen, immunogenic conjugates, or derivatives by combining the protein or conjugate (for rabbits or mice, respectively) with 3 volumes of Freund’s complete adjuvant and injecting the solution intradermally at multiple sites. One month later, the animals are boosted with 1/5 to 1/10 the original amount of peptide or conjugate in Freund’s complete adjuvant by subcutaneous injection at multiple sites. Seven to fourteen days later, the animals are bled and the serum is assayed for antibody titer. Animals are boosted until the titer plateaus. Conjugates also can be made in recombinant cell culture as protein fusions. Also, aggregating agents such as alum are suitable used to enhance the immune response.

[0117] The term “immunizing” refers to the step or steps of administering one or more antigens to a non-human animal so that antibodies can be raised in the animal.

[0118] Specifically, the non-human animal is preferably immunized at least two, more preferably three times with said polypeptide (antigen), optionally in admixture with an adjuvant. An “adjuvant” is a nonspecific stimulant of the immune response. The adjuvant may be in the form of a composition comprising either or both of the following components: (a) a substance designed to form a deposit protecting the antigen(s) from rapid catabolism (e.g. mineral

oil, alum, aluminium hydroxide, liposome or surfactant (e.g. pluronic polyol) and (b) a substance that nonspecifically stimulates the immune response of the immunized host animal (e.g. by increasing lymphokine levels therein).

[0119] Exemplary molecules for increasing lymphokine levels include lipopolysaccharide (LPS) or a Lipid A portion thereof; *Bordetella pertussis*; pertussis toxin; *Mycobacterium tuberculosis*; and muramyl dipeptide (MDP). Examples of adjuvants include Freund’s adjuvant (optionally comprising killed *M. tuberculosis*; complete Freund’s adjuvant); aluminium hydroxide adjuvant; and monophosphoryl Lipid A-synthetic trehalose dicorynomycolate (MPL-TDM).

[0120] The “non-human animal” to be immunized herein is preferably a rodent. A “rodent” is an animal belonging to the rodentia order of placental mammals. Exemplary rodents include mice, rats, guinea pigs, squirrels, hamsters, ferrets etc, with mice being the preferred rodent for immunizing according to the method herein. Other non-human animals which can be immunized herein include non-human primates such as Old World monkey (e.g. baboon or macaque, including Rhesus monkey and cynomolgus monkey; see U.S. Pat. No. 5,658,570); birds (e.g. chickens); rabbits; goats; sheep; cows; horses; pigs; donkeys; dogs etc.

[0121] By “screening” is meant subjecting one or more monoclonal antibodies (e.g., purified antibody and/or hybridoma culture supernatant comprising the antibody) to one or more assays which determine qualitatively and/or quantitatively the ability of an antibody to bind to an antigen of interest.

[0122] By “immuno-assay” is meant an assay that determines binding of an antibody to an antigen, wherein either the antibody or antigen, or both, are optionally adsorbed on a solid phase (i.e., an “immunoabsorbent” assay) at some stage of the assay. Exemplary such assays include ELISAs, radioimmunoassays (RIAs), and FACS assays. Given the above, the present invention provides thus a monoclonal or polyclonal antibody obtainable by the aforescribed methods for the generation of an antibody, i.e., by immunizing a non-human animal as described before.

2) Monoclonal Antibodies.

[0123] Monoclonal antibodies are obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations and/or post-translational modifications (e.g., isomerizations, amidations) that may be present in minor amounts. Thus, the modifier “monoclonal” indicates the character of the antibody as not being a mixture of discrete antibodies. For example, the monoclonal antibodies may be made using the hybridoma method first described by Kohler et al., Nature, 256:495 (1975), or may be made by recombinant DNA methods (U.S. Pat. No. 4,816,567).

[0124] In the hybridoma method, a mouse or other appropriate host animal, such as a hamster, is immunized as hereinabove described to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the protein used for immunization.

[0125] Alternatively, lymphocytes may be immunized in vitro. Lymphocytes then are fused with myeloma cells using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, Monoclonal Antibodies: Principles and Practice, pp. 59-103 (Academic Press, 1986).

[0126] The immunizing agent will typically include the antigenic protein or a fusion variant thereof. Generally either peripheral blood lymphocytes ("PBLs") are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell. Goding, *Monoclonal Antibodies: Principles and Practice*, Academic Press (1986), pp. 59-103.

[0127] Immortalized cell lines are usually transformed mammalian cell, particularly myeloma cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells thus prepared are seeded and grown in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, parental myeloma cells. For example, if the parental myeloma cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine (HAT medium), which substances prevent the growth of HGPRT-deficient cells.

[0128] Preferred immortalized myeloma cells are those that fuse efficiently, support stable high-level production of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. Among these, preferred are murine myeloma lines, such as those derived from MOPC-21 and MPC-11 mouse tumors available from the Salk Institute Cell Distribution Center, San Diego, California USA, and SP-2 cells (and derivatives thereof, e.g., X63-Ag8-653) available from the American Type Culture Collection, Manassas, Virginia USA. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies (Kozbor, *J. Immunol.*, 133:3001 (1984); Brodeur et al., *Monoclonal Antibody Production Techniques and Applications*, pp. 51-63 (Marcel Dekker, Inc., New York, 1987)).

[0129] Culture medium in which hybridoma cells are growing is assayed for production of monoclonal antibodies directed against the antigen. Preferably, the binding specificity of monoclonal antibodies produced by hybridoma cells is determined by immunoprecipitation or by an in vitro binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA).

[0130] The culture medium in which the hybridoma cells are cultured can be assayed for the presence of monoclonal antibodies directed against desired antigen. Preferably, the binding affinity and specificity of the monoclonal antibody can be determined by immunoprecipitation or by an in vitro binding assay, such as radioimmunoassay (RIA) or enzyme-linked assay (ELISA). Such techniques and assays are known in the art. For example, binding affinity may be determined by the Scatchard analysis of Munson et al., *Anal. Biochem.*, 107:220 (1980).

[0131] After hybridoma cells are identified that produce antibodies of the desired specificity, affinity, and/or activity, the clones may be subcloned by limiting dilution procedures and grown by standard methods (Goding, *supra*). Suitable culture media for this purpose include, for example, D-MEM or RPMI-1640 medium. In addition, the hybridoma cells may be grown in vivo as ascites tumors in a mammal.

[0132] The monoclonal antibodies secreted by the subclones are suitably separated from the culture medium, ascites fluid, or serum by conventional immunoglobulin

purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

[0133] Monoclonal antibodies may also be made by recombinant DNA methods, such as those described in U.S. Pat. No. 4,816,567, and as described above. DNA encoding the monoclonal antibodies is readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as *E. coli* cells, simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, in order to synthesize monoclonal antibodies in such recombinant host cells. Review articles on recombinant expression in bacteria of DNA encoding the antibody include Skerra et al., *Curr. Opinion in Immunol.*, 5:256-262 (1993) and Pluckthun, *Immunol. Revs.* 130:151-188 (1992).

3) Humanized Antibodies

[0134] The antibodies of the invention may further comprise humanized or human antibodies. Humanized forms of non-human (e.g., murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F (ab')₂ or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. Humanized antibodies include human immunoglobulins (recipient antibody) in which residues from a complementarity determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues.

[0135] Humanized antibodies may also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domain, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. Jones et al., *Nature* 321:522-525 (1986); Riechmann et al., *Nature* 332:323-329 (1988) and Presta, *Curr. Opin. Struct. Biol.* 2:593-596 (1992).

[0136] Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers, Jones et al., *Nature* 321:522-525 (1986); Riechmann et al., *Nature* 332:323-327 (1988); Verhoeyen et al., *Science* 239:1534-1536 (1988), or through substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such

“humanized” antibodies are chimeric antibodies (U.S. Pat. No. 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

[0137] The choice of human variable domains, both light and heavy, to be used in making the humanized antibodies is very important to reduce antigenicity. According to the so-called “best-fit” method, the sequence of the variable domain of a rodent antibody is screened against the entire library of known human variable-domain sequences. The human sequence which is closest to that of the rodent is then accepted as the human framework (FR) for the humanized antibody. Sims et al., *J. Immunol.* 151:2296 (1993); Chothia et al., *J. Mol. Biol.*, 196:901 (1987).

[0138] Another method uses a particular framework derived from the consensus sequence of all human antibodies of a particular subgroup of light or heavy chains. The same framework may be used for several different humanized antibodies. Carter et al., *Proc. Natl. Acad. Sci. USA*, 89:4285 (1992); Presta et al., *J. Immunol.*, 151:2623 (1993).

[0139] It is further important that antibodies be humanized with retention of high affinity for the antigen and other favorable biological properties. To achieve this goal, according to a preferred method, humanized antibodies are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three-dimensional models of the parental and humanized sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, i.e., the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the recipient and import sequences so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved. In general, the CDR residues are directly and most substantially involved in influencing antigen binding.

[0140] Various forms of the humanized antibody are contemplated. For example, the humanized antibody may be an antibody fragment, such as an Fab, which is optionally conjugated with one or more cytotoxic agent(s) in order to generate an immunoconjugate.

[0141] Alternatively, the humanized antibody may be an intact antibody, such as an intact IgG1 antibody.

4) Human Antibodies

[0142] As an alternative to humanization, human antibodies can be generated. For example, it is now possible to produce transgenic animals (e.g., mice) that are capable, upon immunization, of producing a full repertoire of human antibodies in the absence of endogenous immunoglobulin production. For example, it has been described that the homozygous deletion of the antibody heavy-chain joining region (JH) gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody pro-

duction. Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice will result in the production of human antibodies upon antigen challenge. See, e.g., Jakobovits et al., *Proc. Natl. Acad. Sci. USA*, 90:2551 (1993); Jakobovits et al., *Nature*, 362:255-258 (1993); Bruggermann et al., *Year in Immun.*, 7:33 (1993); U.S. Pat. No. 5,591,669 and WO 97/17852.

[0143] Alternatively, phage display technology can be used to produce human antibodies and antibody fragments in vitro, from immunoglobulin variable (V) domain gene repertoires from unimmunized donors. McCafferty et al., *Nature* 348:552-553 (1990); Hoogenboom and Winter, *J. Mol. Biol.* 227:381 (1991). According to this technique, antibody V domain genes are cloned in-frame into either a major or minor coat protein gene of a filamentous bacteriophage, such as M13, and displayed as functional antibody fragments on the surface of the phage particle. Because the filamentous particle contains a single-stranded DNA copy of the phage genome, selections based on the functional properties of the antibody also result in selection of the gene encoding the antibody exhibiting those properties. Thus, the phage mimics some of the properties of the B-cell. Phage display can be performed in a variety of formats, reviewed in, e.g., Johnson, Kevin S. and Chiswell, David J., *Curr. Opin. Struct. Biol.* 3:564-571 (1993). Several sources of V-gene segments can be used for phage display. Clackson et al., *Nature* 352:624-628 (1991) isolated a diverse array of anti-oxazolone antibodies from a small random combinatorial library of V genes derived from the spleens of immunized mice. A repertoire of V genes from unimmunized human donors can be constructed and antibodies to a diverse array of antigens (including self-antigens) can be isolated essentially following the techniques described by Marks et al., *J. Mol. Biol.* 222:581-597 (1991), or Griffith et al., *EMBO J.* 12:725-734 (1993). See also, U.S. Pat. Nos. 5,565,332 and 5,573,905.

[0144] The techniques of Cole et al., and Boerner et al., are also available for the preparation of human monoclonal antibodies (Cole et al., *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, p. 77 (1985) and Boerner et al., *J. Immunol.* 147 (1): 86-95 (1991). Similarly, human antibodies can be made by introducing human immunoglobulin loci into transgenic animals, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resemble that seen in human in all respects, including gene rearrangement, assembly and antibody repertoire. This approach is described, for example, in U.S. Pat. Nos. 5,545,807; 5,545,806, 5,569,825, 5,625,126, 5,633,425, 5,661,016 and in the following scientific publications: Marks et al., *Bio/Technology* 10:779-783 (1992); Lonberg et al., *Nature* 368:856-859 (1994); Morrison, *Nature* 368:812-13 (1994), Fishwild et al., *Nature Biotechnology* 14:845-51 (1996), Neuberger, *Nature Biotechnology* 14:826 (1996) and Lonberg and Huszar, *Intern. Rev. Immunol.* 13:65-93 (1995). Finally, human antibodies may also be generated in vitro by activated B cells (see U.S. Pat. Nos. 5,567,610 and 5,229,275).

5) Bispecific and Polyspecific Antibodies

[0145] Bispecific antibodies (BsAbs) are antibodies that have binding specificities for at least two different epitopes, including those on the same or another protein. Alternatively, one arm can be armed to bind to the target antigen,

and another arm can be combined with an arm that binds to a triggering molecule on a leukocyte such as a T-cell receptor molecule (e.g., CD3), or Fc receptors for IgG (FcγR) such as FcγRI (CD64), FcγRII (CD32) and FcγRIII (CD16), so as to focus and localize cellular defense mechanisms to the target antigen-expressing cell. Such antibodies can be derived from full length antibodies or antibody fragments (e.g. F(ab')₂ bispecific antibodies).

[0146] Bispecific antibodies may also be used to localize cytotoxic agents to cells which express the target antigen. Such antibodies possess one arm that binds the desired antigen and another arm that binds the cytotoxic agent (e.g., methotrexate).

[0147] Methods for making bispecific antibodies are known in the art. Traditional production of full length bispecific antibodies is based on the coexpression of two immunoglobulin heavy chain-light chain pairs, where the two chains have different specificities. Millstein et al., *Nature*, 305:537-539 (1983). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of 10 different antibody molecules, of which only one has the correct bispecific structure. Purification of the correct molecule, which is usually done by affinity chromatography steps, is rather cumbersome, and the product yields are low. Similar procedures are disclosed in WO 93/08829 and in Trautnecker et al., *EMBO J.*, 10:3655-3659 (1991).

[0148] According to a different approach, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light chain binding, present in at least one of the fusions. DNAs encoding the immunoglobulin heavy chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. This provides for great flexibility in adjusting the mutual proportions of the three polypeptide fragments in embodiments when unequal ratios of the three polypeptide chains used in the construction provide the optimum yields. It is, however, possible to insert the coding sequences for two or all three polypeptide chains in one expression vector when the expression of at least two polypeptide chains in equal ratios results in high yields or when the ratios are of no particular significance.

[0149] In a preferred embodiment of this approach, the bispecific antibodies are composed of a hybrid immunoglobulin heavy chain with a first binding specificity in one arm, and a hybrid immunoglobulin heavy chain-light chain pair (providing a second binding specificity) in the other arm. It was found that this asymmetric structure facilitates the separation of the desired bispecific compound from unwanted immunoglobulin chain combinations, as the presence of an immunoglobulin light chain in only one half of the bispecific molecules provides for an easy way of separation. This approach is disclosed in WO 94/04690. For further details of generating bispecific antibodies, see, for example, Suresh et al., *Methods in Enzymology* 121:210 (1986).

[0150] According to another approach described in WO 96/27011 or U.S. Pat. No. 5,731,168, the interface between

a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the CH3 region of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g., tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chains(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g., alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

[0151] Techniques for generating bispecific antibodies from antibody fragments have been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan et al., *Science* 229:81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

[0152] Fab' fragments may be directly recovered from *E. coli* and chemically coupled to form bispecific antibodies. Shalaby et al., *J. Exp. Med.* 175:217-225 (1992) describes the production of fully humanized bispecific antibody F(ab')₂ molecules. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling in vitro to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the ErbB2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

[0153] Various techniques for making and isolating bivalent antibody fragments directly from recombinant cell culture have also been described. For example, bivalent heterodimers have been produced using leucine zippers. Kostelny et al., *J. Immunol.*, 148 (5): 1547-1553 (1992).

[0154] The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. The "diabody" technology described by Hollinger et al., *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific/bivalent antibody fragments. The fragments comprise a heavy-chain variable domain (VH) connected to a light-chain variable domain (VL) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the VH and VL domains of one fragment are forced to pair with the complementary VL and VH domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific/bivalent antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See Gruber et al., *J. Immunol.*, 152:5368 (1994).

[0155] Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt et al., *J. Immunol.* 147:60 (1991).

[0156] Exemplary bispecific antibodies may bind to two different epitopes on a given molecule. Alternatively, an anti-protein arm may be combined with an arm which binds to a triggering molecule on a leukocyte such as a T-cell receptor molecule (e.g., CD2, CD3, CD28 or B7), or Fc receptors for IgG (FcγR), such as FcγRI (CD64), FcγRII (CD32) and FcγRIII (CD16) so as to focus cellular defense mechanisms to the cell expressing the particular protein.

[0157] Another bispecific antibody of interest binds the protein of interest and further binds Human Serum Albumin.

[0158] The “diabody” technology described by Hollinger et al., *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a VH connected to a VL by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the VH and VL domains of one fragment are forced to pair with the complementary VL and VH domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (scFv) dimers has also been reported. See Gruber et al., *J. Immunol.*, 152:5368 (1994).

[0159] Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt et al. *J. Immunol.* 147:60 (1991).

[0160] A multivalent antibody may be internalized (and/or catabolized) faster than a bivalent antibody by a cell expressing an antigen to which the antibodies bind. The antibodies of the present invention can be multivalent antibodies (which are other than of the IgM class) with three or more antigen binding sites (e.g. tetravalent antibodies), which can be readily produced by recombinant expression of nucleic acid encoding the polypeptide chains of the antibody. The multivalent antibody can comprise a dimerization domain and three or more antigen binding sites. The preferred dimerization domain comprises (or consists of) an Fc region or a hinge region. In this scenario, the antibody will comprise an Fc region and three or more antigen binding sites amino-terminal to the Fc region. The preferred multivalent antibody herein comprises (or consists of) three to about eight, but preferably four, antigen binding sites. The multivalent antibody comprises at least one polypeptide chain (and preferably two polypeptide chains), wherein the polypeptide chain(s) comprise two or more variable domains. For instance, the polypeptide chain(s) may comprise VDI (X1_n-VD2-(X2)_n-Fc, wherein VDI is a first variable domain, VD2 is a second variable domain, Fc is one polypeptide chain of an Fc region, X1 and X2 represent an amino acid or polypeptide, and n is 0 or 1. For instance, the polypeptide chain(s) may comprise: VH-CHI-flexible linker-VH-CHI-Fc region chain; or VH-CHI-VH-CHI-Fc region chain. The multivalent antibody herein preferably further comprises at least two (and preferably four) light chain variable domain polypeptides. The multivalent antibody herein may, for instance, comprise from about two to about eight light chain variable domain polypeptides. The light chain variable domain polypeptides contemplated here comprise a light chain variable domain and, optionally, further comprise a CL domain.

[0161] Heteroconjugate antibodies are also within the scope of the present invention.

[0162] Heteroconjugate antibodies are composed of two covalently joined antibodies. For example, one of the antibodies in the heteroconjugate can be coupled to avidin, the other to biotin. It is contemplated that the antibodies may be prepared in vitro using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins may be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate and those disclosed, for example, in U.S. Pat. No. 4,676,980. Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable cross-linking agents are well known in the art, and are disclosed in U.S. Pat. No. 4,676,980, along with a number of cross-linking techniques.

[0163] For additional antibody production techniques, see *Antibodies: A Laboratory Manual*, eds. Harlow et al., Cold Spring Harbor Laboratory, 1988. The present invention is not necessarily limited to any particular source, method of production, or other special characteristics of an antibody.

[0164] The antibody relating to the present invention is preferably an “isolated” antibody. “Isolated” when used to describe antibodies disclosed herein, means an antibody that has been identified, separated and/or recovered from a component of its production environment. Preferably, the isolated antibody is free of association with all other components from its production environment. Contaminant components of its production environment, such as that resulting from recombinant transfected cells, are materials that would typically interfere with diagnostic or therapeutic uses for the polypeptide, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the antibody will be purified (1) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (2) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Ordinarily, however, an isolated antibody will be prepared by at least one purification step.

[0165] As used herein, “cancer” refers a broad group of diseases characterized by the uncontrolled growth of abnormal cells in the body. Unregulated cell division may result in the formation of malignant tumors or cells that invade neighboring tissues and may metastasize to distant parts of the body through the lymphatic system or bloodstream.

[0166] Non-limiting examples of cancers include squamous cell carcinoma, small-cell lung cancer, non-small cell lung cancer, squamous non-small cell lung cancer (NSCLC), non NSCLC, glioma, gastrointestinal cancer, renal cancer (e.g. clear cell carcinoma), ovarian cancer, liver cancer, colorectal cancer, endometrial cancer, kidney cancer (e.g., renal cell carcinoma (RCC)), prostate cancer (e.g. hormone refractory prostate adenocarcinoma), thyroid cancer, neuroblastoma, pancreatic cancer, glioblastoma (glioblastoma multiforme), cervical cancer, stomach cancer, bladder cancer, hepatoma, breast cancer, colon carcinoma, and head and neck cancer (or carcinoma), gastric cancer, germ cell tumor, pediatric sarcoma, sinonasal natural killer, melanoma (e.g., metastatic malignant melanoma, such as cutaneous or intraocular malignant melanoma), bone cancer, skin cancer, uterine cancer, cancer of the anal region, testicular cancer, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina,

carcinoma of the vulva, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, solid tumors of childhood, cancer of the ureter, carcinoma of the renal pelvis, neoplasm of the central nervous system (CNS), primary CNS lymphoma, tumor angiogenesis, spinal axis tumor, brain stem glioma, pituitary adenoma, Kaposi's sarcoma, epidermoid cancer, squamous cell cancer, T-cell lymphoma, environmentally-induced cancers including those induced by asbestos, virus-related cancers (e.g., human papilloma virus (HPV)-related tumor), and hematologic malignancies derived from either of the two major blood cell lineages, i.e., the myeloid cell line (which produces granulocytes, erythrocytes, thrombocytes, macrophages and mast cells) or lymphoid cell line (which produces B, T, NK and plasma cells), such as all types of leukemias, lymphomas, and myelomas, e.g., acute, chronic, lymphocytic and/or myelogenous leukemias, such as acute leukemia (ALL), acute myelogenous leukemia (AML), chronic lymphocytic leukemia (CLL), and chronic myelogenous leukemia (CML), undifferentiated AML (MO), myeloblastic leukemia (MI), myeloblastic leukemia (M2; with cell maturation), promyelocytic leukemia (M3 or M3 variant [M3V]), myelomonocytic leukemia (M4 or M4 variant with eosinophilia [M4E]), monocytic leukemia (M5), erythroleukemia (M6), megakaryoblastic leukemia (M7), isolated granulocytic sarcoma, and chloroma; lymphomas, such as Hodgkin's lymphoma (HL), non-Hodgkin's lymphoma (NHL), B-cell lymphomas, T-cell lymphomas, lymphoplasmacytoid lymphoma, monocytoid B-cell lymphoma, mucosa-associated lymphoid tissue (MALT) lymphoma, anaplastic (e.g., Ki 1+) large-cell lymphoma, adult T-cell lymphoma/leukemia, mantle cell lymphoma, angio immunoblastic T-cell lymphoma, angiocentric lymphoma, intestinal T-cell lymphoma, primary mediastinal B-cell lymphoma, precursor T-lymphoblastic lymphoma, T-lymphoblastic; and lymphoma/leukaemia (T-Lbly/T-ALL), peripheral T-cell lymphoma, lymphoblastic lymphoma, post-transplantation, lymphoproliferative disorder, true histiocytic lymphoma, primary central nervous system lymphoma, primary effusion lymphoma, lymphoblastic lymphoma (LBL), hematopoietic tumors of lymphoid lineage, acute lymphoblastic leukemia, diffuse large B-cell lymphoma, Burkitt's lymphoma, follicular lymphoma, diffuse histiocytic lymphoma (DHL), immunoblastic large cell lymphoma, precursor B-lymphoblastic lymphoma, cutaneous T-cell lymphoma (CTLC) (also called mycosis fungoides or Sezary syndrome), and lymphoplasmacytoid lymphoma (LPL) with Waldenstrom's macroglobulinemia; myelomas, such as IgG myeloma, light chain myeloma, nonsecretory myeloma, smoldering myeloma (also called indolent myeloma), solitary, plasmocytoma, and multiple myelomas, chronic lymphocytic leukemia (CLL), hairy cell lymphoma; hematopoietic tumors of myeloid lineage, tumors of mesenchymal origin, including fibrosarcoma and rhabdomyosarcoma; seminoma, teratocarcinoma, tumors of the central and peripheral nervous, including astrocytoma, schwannomas; tumors of mesenchymal origin, including fibrosarcoma, rhabdomyosarcoma, and osteosarcoma; and other tumors, including melanoma, xeroderma pigmentosum, keratoacanthoma, seminoma, thyroid follicular cancer and teratocarcinoma, hematopoietic tumors of lymphoid lineage, for example T-cell and B-cell tumors, including but not limited to T-cell disorders such as T-pro-

lymphocytic leukemia (T-PLL), including of the small cell and cerebriform cell type; large granular lymphocyte leukemia (LGL) preferably of the T-cell type; a/d T-NHL hepatosplenic lymphoma; peripheral/post-thymic T cell lymphoma (pleomorphic and immunoblastic subtypes); angiocentric (nasal) T-cell lymphoma; cancer of the head or neck, renal cancer, rectal cancer, cancer of the thyroid gland; acute myeloid lymphoma, as well as any combinations of said cancers. The methods described herein may also be used for treatment of metastatic cancers, refractory cancers (e.g., cancers refractory to previous immunotherapy, e.g., with a blocking CTLA-4 or PD-1 or PD-L1 antibody), and recurrent cancers.

[0167] The term "subject" is intended to include living organisms. Examples of subjects include mammals, e.g., humans, dogs, cows, horses, pigs, sheep, goats, cats, mice, rabbits, rats, and transgenic non-human animals. The subject (animal) can however be a non-mammalian animal such as a bird or a fish. In some preferred embodiments of the invention, the subject is a human, while in other some other preferred embodiments, the subject might be a farm animal, wherein the farm animal can be either a mammal or a non-mammalian animal. Examples of such non-mammalian animals are birds (e.g. poultry such as chicken, duck, goose or turkey), fishes (for example, fishes cultivated in aquaculture such as salmon, trout, or tilapia) or crustacean (such as shrimps or prawns). Examples of mammalian (life stock) animals includes goats; sheep; cows; horses; pigs; or donkeys. Other mammals include cats, dogs, mice and rabbits, for example. In illustrative embodiments the polymersomes of the present invention are used for the vaccination or immunization of the above-mentioned farm animals, both mammalian farm animals and non-mammalian farm animals (a bird, a fish, a crustacean) against virus infections (cf. the Example section in this regard). Accordingly, in such cases, polymersomes of the invention may have encapsulated therein soluble viral full length proteins or soluble fragments of viral full-length proteins.

[0168] When used for vaccinations of both humans and non-humans animals, polymersomes or compositions comprising polymersomes of the invention may be administered orally to the respective subject (cf. also the Example Section) dissolved only in a suitable (pharmaceutically acceptable) buffer such as phosphate-buffered saline (PBS) or 0.9% saline solution (an isotonic solution of 0.90% w/v of NaCl, with an osmolality of 308 mOsm/L). Alternatively, the polymersomes can be modified, for example, by a coating with natural polymers or can be formulated in particles of natural polymers such as alginate or chitosan or of synthetic polymers such as poly(d,l-lactide-co-glycolide) (PLG), poly(d,l-lactic-coglycolic acid) (PLGA), poly(g-glutamic acid) (g-PGA) [31,32] or poly(ethylene glycol) (PEG). These particles can either be particles in the micrometer range ("macrobeads") or nanoparticles, or nanoparticles incorporated into macrobeads all of which are well known in the art. See, for example, Hari et al, "Chitosan/calcium-alginate beads for oral delivery of insulin", Applied Polymer Science, Volume 59, Issue 11, 14 Mar. 1996, 1795-1801, the review of Sosnik "Alginate Particles as Platform for Drug Delivery by the Oral Route: State-of-the-Art" ISRN Pharmaceuticals Volume 2014, Article ID 926157, Machado et al, Encapsulation of DNA in Macroscopic and Nanosized Calcium Alginate Gel Particles", Langmuir 2013, 29, 15926-15935, International Patent Application WO 2015/110656,

the review “Nanoparticle vaccines” of Liang Zhao et al. Vaccine 32 (2014) 327-337) or Li et al “Chitosan-Alginate Nanoparticles as a Novel Drug Delivery System for Nifedipine” Int J Biomed Sci vol. 4 no. 3 Sep. 2008, 221-228. In illustrative embodiments of these polymersomes and oral formulations, the polymersomes that are used for vaccination have encapsulated therein a viral antigen that comprises a soluble portion of Influenza hemagglutinin, Swine Influenza hemagglutinin, Foot and Mouth Disease (FMD) virus protein such as the VP1, VP2 or VP3 coat protein (the VP1 coat protein contains the main antigenic determinants of the FMD virion, and hence changes in its sequence should be responsible for the high antigenic variability of the virus), Ovalbumin (OVA) or of the Porcine epidemic diarrhea (PED) virus SPIKE protein. As evident from the use of polymersomes comprising a soluble portion of the influenza hemagglutinin or a Foot and Mouth Disease (FMD) virus protein such as the VP1, VP2 or VP3 coat protein, the viral disease can affect any animal including birds and mammals, wherein a mammal can also be a human.

[0169] The term “effective dose” or “effective dosage” is defined as an amount sufficient to achieve or at least partially achieve the desired effect. The term “therapeutically effective dose” is defined as an amount sufficient to cure or at least partially arrest the disease and its complications in a patient already suffering from the disease. Amounts effective for this use will depend upon the severity of the infection and the general state of the subject’s own immune system. The term “patient” includes human and other mammalian subjects that receive either prophylactic or therapeutic treatment.

[0170] The appropriate dosage, or therapeutically effective amount, of the antibody or antigen binding portion thereof will depend on the condition to be treated, the severity of the condition, prior therapy, and the patient’s clinical history and response to the therapeutic agent. The proper dose can be adjusted according to the judgment of the attending physician such that it can be administered to the patient one time or over a series of administrations. The pharmaceutical composition can be administered as a sole therapeutic or in combination with additional therapies as needed.

[0171] If the pharmaceutical composition has been lyophilized, the lyophilized material is first reconstituted in an appropriate liquid prior to administration. The lyophilized material may be reconstituted in, e.g., bacteriostatic water for injection (BWI), physiological saline, phosphate buffered saline (PBS), or the same formulation the protein had been in prior to lyophilization.

[0172] Pharmaceutical compositions for injection may be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. In addition, a number of recent drug delivery approaches have been developed and the pharmaceutical compositions of the present invention are suitable for administration using these new methods, e.g., Inject-ease, Genject, injector pens such as Genen, and needleless devices such as MediJector and BioJector. The present pharmaceutical composition can also be adapted for yet to be discovered administration methods. See also Langer, 1990, Science, 249:1527-1533.

[0173] The pharmaceutical composition can also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously, into the ligament or tendon, subsynovially

or intramuscularly), by subsynovial injection or by intramuscular injection. Thus, for example, the formulations may be modified with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

[0174] The pharmaceutical compositions may also be in a variety of conventional depot forms employed for administration to provide reactive compositions. These include, for example, solid, semi-solid and liquid dosage forms, such as liquid solutions or suspensions, slurries, gels, creams, balms, emulsions, lotions, powders, sprays, foams, pastes, ointments, salves, balms and drops.

[0175] The pharmaceutical compositions may, if desired, be presented in a vial, pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. In one embodiment, the dispenser device may comprise a syringe having a single dose of the liquid formulation ready for injection. The syringe can be accompanied by instructions for administration.

[0176] The pharmaceutical composition may further comprise additional pharmaceutically acceptable components. Other pharmaceutically acceptable carriers, excipients, or stabilizers, such as those described in Remington’s Pharmaceutical Sciences 16th edition, Osol, A. Ed. (1980) may also be included in a protein formulation described herein, provided that they do not adversely affect the desired characteristics of the formulation. As used herein, “pharmaceutically acceptable carrier” means any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed and include: additional buffering agents; preservatives; co-solvents; antioxidants, including ascorbic acid and methionine; chelating agents such as EDTA; metal complexes (e.g., Zn-protein complexes); biodegradable polymers, such as polyesters; salt-forming counterions, such as sodium, polyhydric sugar alcohols; amino acids, such as alanine, glycine, asparagine, 2-phenylalanine, and threonine; sugars or sugar alcohols, such as lactitol, stachyose, mannose, sorbose, xylose, ribose, ribitol, myoinositol, galactose, galactitol, glycerol, cyclitols (e.g., inositol), polyethylene glycol; sulfur containing reducing agents, such as glutathione, thioctic acid, sodium thioglycolate, thioglycerol, [alpha]-monothioglycerol, and sodium thio sulfate; low molecular weight proteins, such as human serum albumin, bovine serum albumin, gelatin, or other immunoglobulins; and hydrophilic polymers, such as polyvinylpyrrolidone.

[0177] The formulations described herein are useful as pharmaceutical compositions in the treatment and/or prevention of the pathological medical condition as described herein in a patient in need thereof. The term “treatment” refers to both therapeutic treatment and prophylactic or preventative measures. Treatment includes the application or administration of the formulation to the body, an isolated tissue, or cell from a patient who has a disease/disorder, a symptom of a disease/disorder, or a predisposition toward a disease/disorder, with the purpose to cure, heal, alleviate,

relieve, alter, remedy, ameliorate, improve, or affect the disease, the symptom of the disease, or the predisposition toward the disease.

[0178] As used herein, the term “treating” and “treatment” refers to administering to a subject a therapeutically effective amount of a pharmaceutical composition according to the invention. A “therapeutically effective amount” refers to an amount of the pharmaceutical composition or the antibody which is sufficient to treat or ameliorate a disease or disorder, to delay the onset of a disease or to provide any therapeutic benefit in the treatment or management of a disease.

[0179] As used herein, the term “prophylaxis” refers to the use of an agent for the prevention of the onset of a disease or disorder. A “prophylactically effective amount” defines an amount of the active component or pharmaceutical agent sufficient to prevent the onset or recurrence of a disease.

[0180] As used herein, the terms “disorder” and “disease” are used interchangeably to refer to a condition in a subject. In particular, the term “cancer” is used interchangeably with the term “tumor”.

[0181] The kit of the invention will typically comprise the container described above and one or more other containers comprising materials desirable from a commercial and user standpoint, including buffers, diluents, filters, needles, syringes, and package inserts with instructions for use.

[0182] In the present context, the term “liposome” refers to a spherical vesicle having at least one lipid bilayer.

[0183] In the present context, the term “endosome” refers to a membrane-bound compartment (i.e., a vacuole) inside eukaryotic cells to which materials ingested by endocytosis are delivered.

[0184] In the present context, the term “late-endosome” refers to a pre-lysosomal endocytic organelle differentiated from early endosomes by lower luminal pH and different protein composition. Late endosomes are more spherical than early endosomes and are mostly juxtanuclear, being concentrated near the microtubule organizing center.

[0185] In the present context, the term “T helper cells” (also called TH cells or “effector CD4 (+) T cells”) refers to T lymphocytes that assist other white blood cells in immunologic processes, including maturation of B cells into plasma cells and memory B cells, and activation of cytotoxic T cells and macrophages. These cells are also known as “CD4 (+) T cells” because they express the CD4 glycoprotein on their surfaces. Helper T cells become activated when they are presented with e.g., peptide antigens, by MHC class II molecules, which are expressed on the surface of antigen-presenting cells (APCs).

[0186] As used herein, the term “self-antigen” refers to any molecule or chemical group of an organism which acts as an antigen in inducing antibody formation in another organism but to which the healthy immune system of the parent organism is tolerant.

[0187] As used herein, the term “% identity” refers to the percentage of identical amino acid residues at the corresponding position within the sequence when comparing two amino acid sequences with an optimal sequence alignment as exemplified by the ClustalW or X techniques as available from www.clustal.org, or equivalent techniques. Accordingly, both sequences (reference sequence and sequence of interest) are aligned, identical amino acid residues between both sequences are identified and the total number of identical amino acids is divided by the total number of amino

acids (amino acid length). The result of this division is a percent value, i.e. percent identity value/degree.

[0188] An immunization method of the present invention can be carried out using a either a full length soluble encapsulated antigen (e.g., protein) or fragment of the protein in a synthetic environment that allows its proper folding, and therefore the probability of isolating antibodies capable of detecting corresponding antigens (e.g., a membrane protein) in vivo would be higher. Moreover, the immunization and antibody generation can be carried out without any prior knowledge of the membrane protein structure, which may otherwise be necessary when using a peptide-based immunization approach.

[0189] Further, when compared to other techniques, the method of the present invention allows for a rapid and cost-effective production of membrane protein encapsulated in an oxidation-stable membrane environment.

[0190] In some aspects, the present invention relates to a method for eliciting an immune response to an antigen (e.g., an immunogen) in a subject. The method may include administering to the subject a composition including a polymersome of the present invention having a membrane (e.g., circumferential) of an amphiphilic polymer. The composition further includes a soluble antigen encapsulated by the membrane of the amphiphilic polymer of the polymersome of the present invention. The immunogen may be a membrane-associated protein. In some further aspects, the polymersome of the present invention comprises a lipid polymer. The administration may be carried out in any suitable fashion, for example, by oral administration, topical administration or injection.

[0191] The frequency of the administration (e.g. oral administration or injection) may be determined and adjusted by a person skilled in the art, dependent on the level of response desired. For example, weekly or bi-weekly administration (e.g. orally or by injection) of polymersomes of the present invention may be given to the subject, which may include a mammalian animal. The immune response can be measured by quantifying the blood concentration level of antibodies (titres) in the mammalian animal against the initial amount of antigen encapsulated by the polymersome of the present invention (cf., the Example Section).

[0192] The structure of the polymersomes may include amphiphilic block copolymers self-assembled into a vesicular format and encapsulating various antigens (e.g., soluble proteins, etc.), that are encapsulated by methods of solvent re-hydration, direct dispersion or by spontaneous self-assembly (e.g., Example 1 as described herein).

[0193] In the present context, the term “soluble antigen” as used herein means an antigen capable of being dissolved or liquefied. The term “soluble antigen” includes antigens that were “solubilized”, i.e., rendered soluble or more soluble, especially in water, by the action of a detergent or other agent. Exemplary non-limiting soluble antigens of the present invention include: polypeptides derived from a non-soluble portion of proteins, hydrophobic polypeptides rendered soluble for encapsulation as well as aggregated polypeptides that are soluble as aggregates.

[0194] In some aspects, the antigens (e.g., membrane proteins) of the present invention are solubilized with the aid of detergents, surfactants, temperature change or pH change. The vesicular structure provided by the amphiphilic block copolymers allows the antigens (e.g., membrane protein) to be folded in a physiologically correct and functional manner,

allowing the immune system of the target mammalian animal to detect said antigens, thereby producing a strong immune response.

[0195] In some aspects, the injection of the composition of the present invention may include intraperitoneal, subcutaneous, or intravenous, intramuscular injection, or non-invasive administration. In some other aspects, the injection of the composition of the present invention may include intradermal injection.

[0196] In some other aspects, the immune response level may be further heightened or boosted by including an adjuvant in the composition including the polymersome of the present invention. In such aspects, the polymersome and the adjuvant can be administered simultaneously to the subject.

[0197] In some aspects, a block copolymer or an amphiphilic polymer of the polymersome of the present invention is neither immunostimulant nor adjuvant.

[0198] In some other aspects, a block copolymer or an amphiphilic polymer of the polymersome of the present invention is immunostimulant and/or adjuvant.

[0199] In some further aspects, a polymersome of the present invention is immunogenic.

[0200] In some further aspects, a polymersome of the present invention is non-immunogenic.

[0201] In some aspects, the adjuvant may be administered separately from the administration of the composition of the present invention including the polymersome of the present invention. The adjuvant may be administered before, simultaneously, or after the administration of the composition including the polymersome encapsulating an antigen of the present invention. For example, the adjuvant may be injected to the subject after injecting the composition including the polymersome encapsulating an antigen of the present invention. In some aspects, the adjuvant can be encapsulated together with the antigen in the polymersomes.

[0202] A person skilled in the art would readily recognize and appreciate that the types of adjuvant to be injected depend on the types of antigen to be injected. The antigen may be an antigen of bacterial, viral, or fungi origin. For example, in the case where the antigen is OVA, the adjuvant may be Sigma Adjuvant System (SAS). Other antigen-adjuvant pairs are also suitable for use in the methods of the present invention. In certain aspects, the use of adjuvants is not needed. In yet other aspects, the present method works better, i.e., stronger immune response being evoked, without the use of adjuvants.

[0203] In some aspects, a membrane protein may be a transmembrane protein, G protein-coupled receptor, neurotransmitter receptor, kinase, porin, ABC transporter, ion transporter, acetylcholine receptor and cell adhesion receptor. The membrane proteins may also be fused to or coupled with a tag or may be tag-free. If the membrane proteins are tagged, then the tag may, for example, be selected from well-known affinity tags such as VSV, His-tag, Strep-tag®, Flag-tag, Intein-tag or GST-tag or a partner of a high affinity binding pair such as biotin or avidin or from a label such as a fluorescent label, an enzyme label, NMR label or isotope label.

[0204] In some aspects, the membrane proteins of fragments (or portions) thereof may be presented prior to encapsulation, or encapsulated simultaneously with the production of the protein through a cell-free expression system.

The cell-free expression system may be an in vitro transcription and translation system.

[0205] The cell-free expression system may also be an eukaryotic cell-free expression system such as the TNT system based on rabbit reticulocytes, wheat germ extract or insect extract, a prokaryotic cell-free expression system or an archaic cell-free expression system.


[0206] As mentioned above, the polymersomes may be formed of amphiphilic di-block or tri-block copolymers. In various aspects, the amphiphilic polymer may include at least one monomer unit of a carboxylic acid, an amide, an amine, an alkylene, a dialkylsiloxane, an ether or an alkylene sulphide.

[0207] In some aspects, the amphiphilic polymer may be a polyether block selected from the group consisting of an oligo(oxyethylene) block, a poly(oxyethylene) block, an oligo(oxypropylene) block, a poly(oxypropylene) block, an oligo(oxybutylene) block and a poly(oxybutylene) block. Further examples of blocks that may be included in the polymer include, but are not limited to, poly(acrylic acid), poly(methyl acrylate), polystyrene, poly(butadiene), poly(2-methyloxazoline), poly(dimethyl siloxane), poly(ϵ -caprolactone), poly(propylene sulphide), poly(N-isopropylacrylamide), poly(2-vinylpyridine), poly(2-(diethylamino)ethyl methacrylate), poly(2-diisopropylamino)ethylmethacrylate), poly(2-methacryloyloxy)ethylphosphorylcholine, poly(isoprene), poly(isobutylene), poly(ethylene-co-butylene) and poly(lactic acid). Examples of a suitable amphiphilic polymer include, but are not limited to, poly(ethyl ethylene)-b-poly(ethylene oxide) (PEE-b-PEO), poly(butadiene)-b-poly(ethylene oxide) (PBD-b-PEO), poly(styrene)-b-poly(acrylic acid) (PS-PAA), poly(dimethylsiloxane)-poly(ethylene oxide) (herein called PDMS-PEO) also known as poly(dimethylsiloxane-b-ethylene oxide), poly(2-methyloxazoline)-b-poly(dimethylsiloxane)-b-poly(2-methyloxazoline) (PMOXA-bPDMS-bPMOXA) including for example, triblock copolymers such as PMOXA₂₀-PDMS₅₄-PMOXA₂₀ (ABA) employed by May et al., 2013, poly(2-methyloxazoline)-b-poly(dimethylsiloxane)-b-poly(ethylene oxide) (PMOXA-b-PDMS-b-PEO), poly(ethylene oxide)-b-poly(propylene sulfide)-b-poly(ethylene oxide) (PEO-b-PPS-b-PEO) and a poly(ethylene oxide)-poly(butylene oxide) block copolymer. A block copolymer can be further specified by the average block length of the respective blocks included in a copolymer. Thus, PBMPEON indicates the presence of polybutadiene blocks (PB) with a length of M and polyethyleneoxide (PEO) blocks with a length of N. M and N are independently selected integers, which may for example be selected in the range from about 6 to about 60. Thus, PB₃₅PEO₁₈ indicates the presence of polybutadiene blocks with an average length of 35 and of polyethyleneoxide blocks with an average length of 18. In certain aspects, the PB-PEO diblock copolymer comprises 5-50 blocks PB and 5-50 blocks PEO. Likewise, PB₁₀PEO₂₄ indicates the presence of polybutadiene blocks with an average length of 10 and of polyethyleneoxide blocks with an average length of 24. Illustrative examples of suitable PB-PEO diblock copolymers that can be used in the present invention include the diblock copolymers PBD₂₁-PEO₁₄ (that is also commercially available) and [PBD]21-[PEO]12, (cf, WO2014/077781A1 and Nallani et al., 2011). As a further example E₀B_p indicates the presence of ethylene oxide blocks (E) with a length of 0 and butadiene blocks (B) with a length of P. Thus, O and P are independently selected


integers, e.g. in the range from about 10 to about 120. Thus, E16E22 indicates the presence of ethylene oxide blocks with an average length of 16 and of butadiene blocks with an average length of 22.

[0208] Turning to another preferred block copolymer that is used to form polymersome of the invention, poly(dimethylsiloxane-*b*-ethyleneoxide) (PDMS-PEO), it is noted that both linear and comb-type PDMS-PEO can be used herein (cf. Gaspard et al, “Mechanical Characterization of Hybrid Vesicles Based on Linear Poly(Dimethylsiloxane-*b*-Ethylene Oxide) and Poly(Butadiene-*b*-Ethylene Oxide) Block Copolymers” *Sensors* 2016, 16 (3), 390 which describes polymersomes formed from PDMS-PEO).

[0209] The structure of linear PDMS-PEO is shown in the following as formula (I)



while the structure of comb-type PDMS-PEO is shown in the following formula (II):



[0210] In line with the structural formula (I), the terminology PDMS_{*n*}-PEO_{*m*} indicates the presence of polydimethylsiloxane (PDMS) blocks with a length of *n* and polyethyleneoxide (PEO) blocks with a length of *m*. *m* and *n* are independently selected integers, each of which may, for example, be selected in the range from about 5 or about 6 to about 100, from about 5 to about 60 or from about 6 to about 60 or from about 5 to 50. For example, linear PDMS-PEO such as PDMS₁₂-PEO₄₆ or PDMS₄₇-PEO₃₆ are commercially available from Polymer Source Inc., Dorval (Montreal) Quebec, Canada. Accordingly, the PDMS-PEO block copolymer may comprise 5-100 blocks PDMS and 5-100 blocks PEO, 6-100 blocks PDMS and 6-100 blocks PEO, 5-100 blocks PDMS and 5-60 blocks PEO, or 5-60 blocks PDMS and 5-60 blocks PEO.

[0211] In accordance with the above, the present invention relates in one aspect to the method of eliciting an immune response in a subject, comprising administering to the subject a polymersome formed from PDMS-PEO carrying an antigen. The antigen can be associated/physically linked with the PDMS-PEO polymersome in any suitable way. For example, the PDMS-PEO polymersome may have a soluble antigen encapsulated therein as described in the present invention. Alternatively or in addition, the polymersome may have an antigen integrated/incorporated into the circumferential membrane of the polymersome as described in WO2014/077781A1. In this case, antigen is a membrane protein that is integrated with its (one or more) transmem-

brane domain into the circumferential membrane of the PDMS-PEO-polymersome. The integration can be achieved as described in WO2014/077781A1 or Nallani et al, “Proteopolymersomes: in vitro production of a membrane protein in polymersome membranes”, *Biointerphases*, 1 Dec. 2011, page 153. In case, the antigen is encapsulated in the PDMS-PEO polymersome, it may be a soluble antigen selected from the group consisting of a polypeptide, a carbohydrate, a polynucleotide and combinations thereof. The present invention further relates to a method for production of such encapsulated antigens in a polymersome formed from PDMS-PEO as well as to polymersomes produced by said method.

[0212] The present invention further relates to compositions comprising PDMS-PEO polymersomes carrying an

antigen. Also, in these compositions, the antigen can be associated/physically linked with the PDMS-PEO polymersome in any suitable way. For example, the PDMS-PEO polymersome may have a soluble antigen encapsulated therein as described in the present invention. Alternatively or in addition, the polymersome may have an antigen integrated/incorporated into the circumferential membrane of the polymersome as described in WO2014/077781A1. The present invention also relates to vaccines comprising such PDMS-PEO polymersomes carrying an antigen, methods of eliciting an immune response or methods for treatment, amelioration, prophylaxis or diagnostics of cancers, autoimmune or infectious diseases, such methods comprising providing PDMS-PEO polymersomes carrying an antigen to subject in need thereof.

[0213] In accordance with the above, the present invention also relates to the in vitro and in vivo use of a PDMS-PEO polymersomes carrying (or transporting) an antigen in a manner suitable for eliciting an immune response. The antigen can either be encapsulated in the PDMS-PEO polymersome or, for example, incorporated into the circumferential membrane of the polymersome as described in WO2014/077781A1.

[0214] In certain aspects, the polymersome of the present invention may contain one or more compartments (or otherwise termed “multicompartment”). Compartmentalization of the vesicular structure of polymersome allows for the co-existence of complex reaction pathways in living cell and helps to provide a spatial and temporal separation of many activities inside a cell. Accordingly, more than one type of antigens may be encapsulated by the polymersome of the present invention. The different antigens may have the same or different isoforms. Each compartment may also be formed of a same or a different amphiphilic polymer. In various aspects, two or more different antigens are integrated into the circumferential membrane of the amphiphilic polymer. Each compartment may encapsulate at least one of peptide, protein, and nucleic acid. The peptide, protein, polynucleotide or carbohydrate may be immunogenic.

[0215] Further details of suitable multicompartimentalized polymersomes can be found in WO 20121018306, the contents of which being hereby incorporated by reference in its entirety for all purposes.

[0216] The polymersomes may also be free-standing or immobilized on a surface, such as those described in WO 2010/1123462, the contents of which being hereby incorporated by reference in its entirety for all purposes.

[0217] In the case where the polymersome carrier contains more than one compartment, the compartments may comprise an outer block copolymer vesicle and at least one inner block copolymer vesicle, wherein the at least one inner block copolymer vesicle is encapsulated inside the outer block copolymer vesicle. In some aspects, each of the block copolymer of the outer vesicle and the inner vesicle includes a polyether block such as a poly(oxyethylene) block, a poly(oxypropylene) block, and a poly(oxybutylene) block. Further examples of blocks—that may be included in the copolymer include, but are not limited to, poly(acrylic acid), poly(methyl acrylate), polystyrene, poly(butadiene), poly(2-methyloxazoline), poly(dimethyl siloxane), poly(L-isocyanoalanine (2-thiophen-3-yl-ethyl)amide), poly(ϵ -caprolactone), poly(propylene sulphide), poly(N-isopropylacrylamide), poly(2-vinylpyridine), poly(2-(diethylamino)ethyl methacrylate), poly(2-(diisopropylamino)ethylmethacrylate), poly(2-(methacryloyloxy)ethylphosphorylcholine) and poly(lactic acid). Examples of suitable outer vesicles and inner vesicles include, but are not limited to, poly(ethyl ethylene)-b-poly(ethylene oxide) (PEE-b-PEO), poly(butadiene)-b-poly(ethylene oxide) (PBD-b-PEO), poly(styrene)-b-poly(acrylic acid) (PS-b-PAA), poly(ethylene oxide)-poly(caprolactone) (PEO-b-PCL), poly(ethylene oxide)-poly(lactic acid) (PEO-b-PLA), poly(isoprene)-poly(ethylene oxide) (PI-b-PEO), poly(2-vinylpyridine)-poly(ethylene oxide) (P2VP-b-PEO), poly(ethylene oxide)-poly(N-isopropylacrylamide) (PEO-b-PNIPAm), poly(ethylene glycol)-poly(propylene sulfide) (PEG-b-PPS), poly(methylphenylsilane)-poly(ethylene oxide) (PMPS-b-PEO-b-PMPS-b-PEO-b-PMPS), poly(2-methyloxazoline)-b-poly-(dimethylsiloxane)-b-poly(2-methyloxazoline) (PMOXA-b-PDMS-b-PMOXA), poly(2-methyloxazoline)-b-poly(dimethylsiloxane)-b-poly(ethylene oxide) (PMOXA-b-PDMS-b-PEO), poly[styrene-b-poly(L-isocyanoalanine (2-thiophen-3-yl-ethyl)amide)] (PS-b-PIAT), poly(ethylene oxide)-b-poly(propylene sulfide)-b-poly(ethylene oxide) (PEO-b-PPS-b-PEO) and a poly(ethylene oxide)-poly(butylene oxide) (PEO-b-PBO) block copolymer. A block copolymer can be further specified by the average number of the respective blocks included in a copolymer. Thus $PS_M\text{-PIAT}_N$ indicates the presence of polystyrene blocks (PS) with M repeating units and poly(L-isocyanoalanine (2-thiophen-3-yl-ethyl)amide) (PIAT) blocks with N repeating units. Thus, M and N are independently selected integers, which may for example be selected in the range from about 5 to about 95. Thus, $PS_{40}\text{-PIAT}_{50}$ indicates the presence of PS blocks with an average of 40 repeating units and of PIAT blocks with an average of 50 repeating units.

[0218] In some aspects, the invention relates to a method for production of an encapsulated antigen in polymersome, said method comprising: i) dissolving an amphiphilic polymer of the present invention in chloroform, preferably said amphiphilic polymer is polybutadiene-polyethylene oxide (BD); ii) drying said dissolved amphiphilic polymer to form

a polymer film; iii) adding a solubilized antigen to said dried amphiphilic polymer film from step ii), wherein said antigen is selected from the group consisting of: (a) a polypeptide; preferably said polypeptide is an antigen according to the present invention; (b) a carbohydrate; (c) a combination of a) and/or b) and/or c); iv) rehydrating said polymer film from step iii) to form polymer vesicles; v) optionally, filtering polymer vesicles from step iv) to purify polymer vesicles monodisperse vesicles; and/or vi) optionally, isolating said polymer vesicles from step iv) or v) from the non-encapsulated antigen.

[0219] In some other aspects, the invention relates to other methods for production of an encapsulated antigen in polymersome including methods based on mixing a non-aqueous solution of polymers in aqueous solution of antigens, sonication of corresponding mixed solutions of polymers and antigens, or extrusion of corresponding mixed solutions of polymers and antigens. Exemplary methods include those described in Rameez et al, Langmuir 2009, and in Neil et al Langmuir 2009, 25 (16), 9025-9029. In some other aspects, the invention relates to a polymersome (or plurality or collection of polymersomes) comprising a soluble encapsulated antigen, wherein said soluble encapsulated antigen is a polynucleotide (e.g., single or double stranded) selected from a RNA molecule (e.g., mRNA, self-amplifying mRNA or siRNA) or a DNA molecule or DNA/RNA hybrid and wherein the polymersome further comprises an ionizable lipid.

[0220] In some other aspects of the invention the ionizable lipid is a cationic ionizable lipid.

[0221] In some other aspects of the invention the ionizable lipid: (i) is either a tertiary amine (e.g., is not a quaternary amine) and/or comprising one or more tertiary amine moieties (e.g., not comprising a cationic quaternized ammonium moiety); and/or (ii) comprising a free electron pair at the nitrogen atom (e.g., ionizable nitrogen atom).

[0222] In some other aspects of the invention the ionizable lipid comprises or consists of: ionizable lipid DLin-MC3-DMA (i.e., (6Z,9Z,28Z,31Z)-Heptatriaconta-6,9,28,31-tetraen-19-yl 4-(dimethylamino) butanoate) having Formula I; ionizable lipid 319 having Formula II; ionizable lipid C12-200 having Formula III; ionizable lipid 5A2-SC8 having Formula IV; ionizable lipid 306Oi10 having Formula; ionizable lipid 5 having Formula VI; ionizable lipid SM-102 (i.e., heptadecan-9-yl 8-((2-hydroxyethyl) (6-oxo-6-(undecyloxy) hexyl)amino) octanoate) having Formula VII; ionizable lipid A9 having Formula VIII; ionizable lipid ALC-0315 (i.e., [(4-hydroxybutyl) azanediy]di(hexane-6,1-diyl)bis(2-hexyldecanoate) having Formula IX; ionizable lipid Arcturus 2.2 (8.8) 4 C CH₃ having Formula X; ionizable lipid Genevant CL1 having Formula XI; ionizable lipid ALC-0159 (i.e. 2-[(polyethylene glycol)-2000]-N, N-ditetradecylacetamide) having Formula XII; any one of above, wherein one or more alkyl chains thereof are further esterified.

[0223] In some preferred aspects of the invention the ionizable lipid is a cationic ionizable lipid comprising or consisting of: DLin-MC3-DMA ((6Z,9Z,28Z,31Z)-Heptatriaconta-6,9,28,31-tetraen-19-yl 4-(dimethylamino) butanoate).

[0224] In some other aspects of the invention the polymersome is capable of increasing thermostability and/or storage stability and/or immunogenicity of said polynucleotide (e.g., mRNA) and/or one or more polypeptides

encoded by said polynucleotide (e.g., mRNA) within said polymersome, preferably compared to that of the same polynucleotide (e.g., mRNA) and/or one or more polypeptides encoded by said same polynucleotide (e.g., mRNA) within another polymersome without ionizable lipid or within an ionizable lipid nanoparticle (LNP) comprising cholesterol under the same conditions, further preferably said thermostability and/or storage stability is increased in the temperature range from about -80°C . to about 4°C . (e.g., at -80°C ., -20°C . or at 4°C ., preferably at -80°C .), most preferably said immunogenicity is increased in the temperature range from about 36.5°C . to about 37.5°C .; further most preferably said increase is of about at least 5% (e.g., at least 10%, at least 20% or at least 30%).

[0225] In some other aspects of the invention the polymersome comprising: polybutadiene-polyethylene oxide (BD) and Dlin-MC3-DMA ((6Z,9Z,28Z,31Z)-Heptatriaconta-6,9,28,31-tetraen-19-yl 4-(dimethylamino) butanoate); and/or PBD-PEO (poly(butadiene)-poly(ethylene oxide)) and Dlin-MC3-DMA ((6Z,9Z,28Z,31Z)-Heptatriaconta-6,9,28,31-tetraen-19-yl 4-(dimethylamino) butanoate), preferably comprising 15% or 30% Dlin-MC3-DMA; and/or polybutadiene-polyethylene oxide (BD) and ionizable lipid 319 having Formula II; and/or polybutadiene-polyethylene oxide (BD) and ionizable lipid C12-200 having Formula III; and/or polybutadiene-polyethylene oxide (BD) and ionizable lipid 5A2-SC8 having Formula IV; and/or polybutadiene-polyethylene oxide (BD) and ionizable lipid 306Oi10 having Formula V; and/or polybutadiene-polyethylene oxide (BD) and ionizable lipid 5 having Formula VI; and/or polybutadiene-polyethylene oxide (BD) and ionizable lipid SM-102 having Formula VII; and/or polybutadiene-polyethylene oxide (BD) and ionizable lipid A9 having Formula VIII; and/or polybutadiene-polyethylene oxide (BD) and ionizable lipid ALC-0315 (i.e., [(4-hydroxybutyl) azanediyl]di(hexane-6,1-diyl)bis(2-hexyldecanoate) having Formula IX; and/or polybutadiene-polyethylene oxide (BD) and ionizable lipid Arcturus 2.2 (8.8) 4 C CH₃ having Formula X; and/or polybutadiene-polyethylene oxide (BD) and ionizable lipid Genevant CL1 having Formula XI; and/or polybutadiene-polyethylene oxide (BD) and ionizable lipid ALC-0159 (i.e. 2-[(polyethylene glycol)-2000]-N,N-ditetradecylacetamide) having Formula XII.

[0226] Compared to existing uptake and cross-presentation vehicles and methods based thereon the polymersomes of present invention inter alia offer the following advantages that are also aspects of the present invention:

[0227] The polymersomes are very efficient in uptake and cross-presentation to the immune system;

[0228] The immune response comprises a CD8⁽⁺⁾ T cell-mediated immune response;

[0229] The polymersomes are oxidation-stable;

[0230] The humoral response is stronger compared to that produced by free antigen-based techniques with or without adjuvants;

[0231] The immune response induced by polymersomes of the present invention could still be even further boosted using adjuvants;

[0232] The polymersomes of the present invention are inherently robust and can be tailored or functionalized to increase their circulation time in the body;

[0233] The polymersomes of the present invention are stable in the presence of serum components;

[0234] The polymersomes are inexpensive and quick to synthesize;

[0235] The amount of an antigen required to elicit an immune response by the methods of the present invention using polymersomes of the present invention is less compared to free antigen-based techniques with or without adjuvants.

[0236] The invention is also characterized by the following items:

[0237] 1. A polymersome (e.g., an oxidation-stable polymersome) comprising a soluble encapsulated antigen, wherein said soluble encapsulated antigen is selected from the group consisting of:

[0238] i) a polypeptide;

[0239] ii) a carbohydrate;

[0240] iii) a polynucleotide, preferably said polynucleotide is not an antisense oligonucleotide, further preferably said polynucleotide is a DNA or mRNA molecule.

[0241] iv) a combination of i) and/or ii) and/or iii).

[0242] 2. The polymersome according to any one of preceding items, wherein said polymersome is capable of eliciting a CD8⁽⁺⁾ T cell-mediated immune response, preferably said eliciting is an in vivo, ex vivo or in vitro eliciting.

[0243] 3. The polymersome according to any one of preceding items, wherein said antigen comprises a soluble portion of a membrane protein (MP) or a membrane-associated peptide (MAP), preferably said antigen comprises a soluble portion of Influenza hemagglutinin, Swine Influenza hemagglutinin, Ovalbumin (OVA), Porcine epidemic diarrhea virus SPIKE protein, B16 peptide or MC38 peptide, further preferably said antigen comprises a polypeptide which is at least 60% or more (e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to a polypeptide sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, and SEQ ID NOs: 12-14.

[0244] 4. The polymersome according to any one of preceding items, wherein said polymersome is stable in the presence of serum components, preferably said stability is an in vivo, ex vivo or in vitro stability.

[0245] 5. The polymersome according to any one of preceding items, wherein said polymersome is stable inside an endosome, preferably said stability is an in vivo, ex vivo or in vitro stability.

[0246] 6. The polymersome according to any one of preceding items, wherein said polymersome has an improved oxidation stability compared to corresponding oxidation stability of a liposome, preferably said improved stability is an in vivo, ex vivo or in vitro improved stability.

[0247] 7. The polymersome according to any one of preceding items, wherein said polymersome is capable of releasing its content comprising said soluble encapsulated antigen in an oxidation-independent manner and triggering CD8⁽⁺⁾ T cell-mediated immune response, preferably said releasing is an in vivo, ex vivo or in vitro releasing.

- [0248] 8. The polymersome according to any one of preceding items, wherein said polymersome is capable of eliciting a cellular immune response, wherein said cellular immune response comprises a CD8⁽⁺⁾ T cell-mediated immune response, preferably said immune response is an in vivo, ex vivo or in vitro immune response.
- [0249] 9. The polymersome according to any one of preceding items, wherein said polymersome is capable of eliciting a cellular and/or humoral immune response, wherein said cellular immune response comprises a CD8⁽⁺⁾ T cell-mediated immune response, preferably immune response is an in vivo, ex vivo or in vitro immune response.
- [0250] 10. The polymersome according to any one of preceding items, wherein said humoral immune response comprises production of specific antibodies, further preferably said immune response is an in vivo, ex vivo or in vitro immune response.
- [0251] 11. The polymersome according to any one of preceding items, wherein said polymersome is capable of enhancing the frequency of effector CD4⁽⁺⁾ T cells, preferably said enhancing is an in vivo, ex vivo or in vitro enhancing.
- [0252] 12. The polymersome according to any one of preceding items, wherein said cellular immune response comprises a T-cell mediated immune response, preferably said immune response is an in vivo, ex vivo or in vitro immune response.
- [0253] 13. The polymersome according to any one of preceding items, wherein said polymersome is capable of enhancing clonal expansion of antigen-specific CD8⁽⁺⁾ T cells compared to a free antigen, preferably said expansion is an in vivo, ex vivo or in vitro expansion.
- [0254] 14. The polymersome according to any one of preceding items, wherein said polymersome is capable of inducing antigen-specific effector CD8⁽⁺⁾ T cells, preferably said inducing is an in vivo, ex vivo or in vitro inducing.
- [0255] 15. The polymersome according to any one of preceding items, wherein said polymersome is capable of enhancing a cytotoxic phenotype of antigen-specific CD8⁽⁺⁾ T cells, preferably said enhancing is an in vivo, ex vivo or in vitro enhancing.
- [0256] 16. The polymersome according to any one of preceding items, wherein said polymersome is capable of targeting of lymph node-resident macrophages and/or B cells, preferably said targeting is an in vivo, ex vivo or in vitro targeting.
- [0257] 17. The polymersome according to any one of preceding items, wherein said polymersome is reduction-stable, preferably said polymersome is reduction-stable in the presence of serum components, further preferably said reduction-stability is an in vivo, ex vivo or in vitro reduction-stability.
- [0258] 18. The polymersome according to any one of preceding items, wherein said polymersome has reduced permeability, preferably said reduced permeability is compared to a corresponding permeability of a liposome, further preferably said permeability is an in vivo, ex vivo or in vitro permeability.
- [0259] 19. The polymersome according to any one of preceding items, wherein said polymersome is capable of releasing its content inside an endosome, preferably said endosome is a late-endosome, further preferably said releasing is an in vivo, ex vivo or in vitro releasing.
- [0260] 20. The polymersome according to any one of preceding items, wherein said polymersome is capable of one or more of the following:
- [0261] i) eliciting a cellular immune response; preferably said cellular immune response comprises a CD8⁽⁺⁾ T cell-mediated immune response; further preferably said cellular immune response is a CD8⁽⁺⁾ T cell-mediated immune response; most preferably said cellular immune response is against a soluble portion of Influenza hemagglutinin, Swine Influenza hemagglutinin, Ovalbumin (OVA), Porcine epidemic diarrhea virus SPIKE protein, B16 peptide or MC38 peptide, further most preferably said cellular immune response is against a polypeptide which is at least 60% or more (e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to a polypeptide sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11 SEQ ID NOS: 12-14;
- [0262] ii) releasing polymersome content inside an endosome, preferably said endosome is a late endosome; further preferably said content comprises a soluble portion of Influenza hemagglutinin, Swine Influenza hemagglutinin, Ovalbumin (OVA), Porcine epidemic diarrhea virus SPIKE protein, B16 peptide or MC38 peptide, most preferably said content comprises a polypeptide which is at least 60% or more (e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to a polypeptide sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NOS: 12-14;
- [0263] iii) releasing polymersome content in an oxidation-independent manner and triggering CD8⁽⁺⁾ T cell-mediated immune response; preferably said content comprises a soluble portion of Influenza hemagglutinin, Swine Influenza hemagglutinin, Ovalbumin (OVA), Porcine epidemic diarrhea virus SPIKE protein, B16 peptide or MC38 peptide, further preferably said content comprises a polypeptide which is at least 60% or more (e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to a polypeptide sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NOS: 12-14;
- [0264] iv) stimulating an immune response to said antigen; preferably said antigen comprises a soluble portion of Influenza hemagglutinin, Swine Influenza hemagglutinin, Ovalbumin (OVA), Porcine epidemic diarrhea virus SPIKE protein, B16 peptide or MC38

peptide, further preferably said antigen comprises a polypeptide which is at least 60% or more (e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to a polypeptide sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NOS: 12-14;

[0265] v) triggering a cross-protection induced by a CD8⁽⁺⁾ T cell-mediated immune response; preferably said response is against a soluble portion of Influenza hemagglutinin, Swine Influenza hemagglutinin, Porcine epidemic diarrhea virus SPIKE protein, Ovalbumin (OVA), B16 peptide or MC38 peptide, further preferably said response is against a polypeptide which is at least 60% or more (e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to a polypeptide sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, and SEQ ID NOS: 12-14;

[0266] vi) delivering a peptide or protein to an antigen-presenting cell (APC); preferably said peptide or protein comprises or is derived from a soluble portion of Influenza hemagglutinin, Swine Influenza hemagglutinin, Ovalbumin (OVA), Porcine epidemic diarrhea virus SPIKE protein, B16 peptide or MC38 peptide, further preferably said peptide or protein comprises or is derived from a polypeptide which is at least 60% or more (e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to a polypeptide sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, and SEQ ID NOS: 12-14;

[0267] vii) triggering an immune response comprising a CD8⁽⁺⁾ T cell-mediated immune response and/or CD4⁽⁺⁾ T cell-mediated immune response; preferably said response is against a soluble portion of Influenza hemagglutinin, Swine Influenza hemagglutinin, Ovalbumin (OVA), Porcine epidemic diarrhea virus SPIKE protein, B16 peptide or MC38 peptide, further preferably said response is against a polypeptide which is at least 60% or more (e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to a polypeptide sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, and SEQ ID NOS: 12-14;

[0268] viii) stimulating an immune response in a subject; preferably said response is against a soluble portion of Influenza hemagglutinin, Swine Influenza hemagglutinin, Ovalbumin (OVA), Porcine epidemic diarrhea virus SPIKE protein, B16 peptide or MC38 peptide, further preferably said response is against a polypeptide which is at least 60% or more (e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to a polypeptide sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, and SEQ ID NOS: 12-14;

[0269] ix) immunizing a non-human animal; preferably said immunizing is against a soluble portion of Influenza hemagglutinin, Swine Influenza hemagglutinin, Ovalbumin (OVA), Porcine epidemic diarrhea virus SPIKE protein, B16 peptide or MC38 peptide, further preferably said immunizing is against a polypeptide which is at least 60% or more (e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to a polypeptide sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, and SEQ ID NOS: 12-14;


[0270] x) said polymersome has an altered antigenicity compared to corresponding antigenicity of said antigen without said polymersome; preferably said antigen is a soluble portion of Influenza hemagglutinin, Swine Influenza hemagglutinin, Ovalbumin (OVA), Porcine epidemic diarrhea virus SPIKE protein, B16 peptide or MC38 peptide, further preferably said antigen is a polypeptide which is at least 60% or more (e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to a polypeptide sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, and SEQ ID NOS: 12-14;

[0271] xi) said polymersome has an altered immunogenicity compared to corresponding immunogenicity of said antigen without said polymersome, preferably said immunogen is a soluble portion of Influenza hemagglutinin, Swine Influenza hemagglutinin, Ovalbumin (OVA), Porcine epidemic diarrhea virus SPIKE protein, B16 peptide or MC38 peptide, further preferably said immunogen is a polypeptide which is at least 60% or more (e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to a polypeptide sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ


- ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, and SEQ ID NOs: 12-14.
- [0272] 21. The polymersome according to any one of preceding items, wherein said polymersome has one or more of the following properties:
- [0273] i) said polymersome comprises an oxidation-stable membrane; and/or
- [0274] ii) said polymersome is synthetic; and/or
- [0275] iii) said polymersome is free from non-encapsulated antigens or in a mixture with free non-encapsulated antigens; and/or
- [0276] iv) said polymersome comprises a membrane of an amphiphilic polymer; and/or
- [0277] and/or
- [0278] v) said polymersome comprises amphiphilic synthetic block copolymers forming a vesicle membrane; and/or
- [0279] vi) said polymersome has a diameter greater than 70 nm, wherein preferably the diameter is a range of about 100 nm to about 1 μ m, or in the range from about 120 nm to about 250 nm, or from about 125 nm to about 250 nm, from about 140 nm to about 240 nm, from about 150 nm to about 235 nm, from about 170 nm to about 230 nm, or from about 220 nm to about 180 nm, or from about 190 nm to about 210 nm and/or
- [0280] vii) said polymersome has a vesicular morphology;
- [0281] viii) said polymersome is self-assembling.
- [0282] 22. The polymersome of item 21, wherein the polymersome is in the form of a collection of polymersomes, wherein the mean diameter of the collection of polymersomes is in the range of about 100 nm to about 1 μ m, or in the range from about 100 nm to about 750 nm, or from about 100 nm to about 500 nm, or from about 120 nm to about 250 nm, from about 125 nm to about 250 nm, from about 140 nm to about 240 nm, from about 150 nm to about 235 nm, from about 170 nm to about 230 nm, or from about 220 nm to about 180 nm, or from about 190 nm to about 210 nm.
- [0283] 23. The polymersome according to any one of preceding items, wherein said antigen is an immunogen.
- [0284] 24. The polymersome according to any one of preceding items, wherein said antigen is selected from a group consisting of: i) a self-antigen, ii) a non-self antigen, iii) a non-self immunogen and iv) a self-immunogen.
- [0285] 25. The polymersome according to any one of preceding items, wherein said antigen is selected from the group consisting of:
- [0286] i) a polypeptide which is at least 80% or more (e.g., at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to a viral polypeptide sequence; preferably said viral polypeptide sequence is Influenza hemagglutinin or Swine Influenza hemagglutinin, further preferably said viral polypeptide sequence is selected from the group consisting of: SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7 and SEQ ID NO: 8;
- [0287] ii) a polypeptide which is at least 80% or more (e.g., at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to a bacterial polypeptide sequence;
- [0288] iii) a polypeptide which is at least 80% or more (e.g., at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to a mammalian or avian polypeptide sequence, preferably said mammalian or avian polypeptide sequence is Ovalbumin (OVA), a polypeptide which is at least 80% or more (e.g., at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to a SPIKE protein, B16 peptide or MC38 peptide, further preferably said mammalian or avian polypeptide sequence is selected from the group consisting of: SEQ ID NO: 4, SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11 and SEQ ID NOs: 12-14.
- [0289] 26. The polymersome according to any one of preceding items, wherein said mammalian polypeptide sequence is selected from the group consisting of: human, rodent, rabbit and horse polypeptide sequence.
- [0290] 27. The polymersome according to any one of preceding items, wherein said antigen is an antibody or a fragment thereof.
- [0291] 28. The polymersome according to any one of preceding items, wherein said antigen is selected from the group consisting of:
- [0292] i) Influenza hemagglutinin (HA), preferably selected from the group consisting of: SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7 and SEQ ID NO: 8;
- [0293] ii) Swine Influenza hemagglutinin (HA), preferably SEQ ID NO: 6;
- [0294] iii) Ovalbumin (OVA), preferably SEQ ID NO: 4;
- [0295] iv) Porcine epidemic diarrhea virus (PED), Spike Protein, preferably SEQ ID NOs: 12-14;
- [0296] v) B16 peptide, preferably selected from the group consisting of: SEQ ID NO: 9, SEQ ID NO: 10 and SEQ ID NO: 11;
- [0297] vi) MC38 peptide, preferably selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2 and SEQ ID NO: 3;
- [0298] vii) B16 and MC38 peptides, preferably said peptides are independently selected the groups: i) SEQ ID NOs: 1-3 and ii) SEQ ID NOs: 9-11.
- [0299] 29. The polymersome according to any one of preceding items, wherein said polymersome is selected from the group consisting of: cationic, anionic and nonionic polymersome and mixtures thereof.
- [0300] 30. The polymersome according to any one of preceding items, wherein said block copolymer or amphiphilic polymer is essentially non-immunogenic or essentially non-antigenic, preferably said block copolymer or amphiphilic polymer is non-immunogenic or non-antigenic.
- [0301] 31. The polymersome according to any one of preceding items, wherein said block copolymer or amphiphilic polymer is oxidation-stable.
- [0302] 32. The polymersome according to any one of preceding items, wherein said block copolymer or said amphiphilic polymer is neither immunostimulant nor adjuvant.

- [0303] 33. The polymersome according to any one of preceding items, wherein said amphiphilic polymer comprises a diblock or a triblock (A-B-A or A-B-C) copolymer.
- [0304] 34. The polymersome according to any one of preceding items, wherein said amphiphilic polymer comprises a copolymer poly(N-vinylpyrrolidone)-b-PLA.
- [0305] 35. The polymersome according to any one of preceding items, wherein said amphiphilic polymer comprises at least one monomer unit of a carboxylic acid, an amide, an amine, an alkylene, a dialkylsiloxane, an ether or an alkylene sulphide.
- [0306] 36. The polymersome according to any one of preceding items, wherein the amphiphilic polymer is a polyether block selected from the group consisting of an oligo(oxyethylene) block, a poly(oxyethylene) block, an oligo(oxypropylene) block, a poly(oxypropylene) block, an oligo(oxybutylene) block and a poly(oxybutylene) block.
- [0307] 37. The polymersome according to any one of preceding items, wherein said amphiphilic polymer is a poly(butadiene)-poly(ethylene oxide) (PB-PEO) diblock copolymer.
- [0308] 38. The polymersome according to any one of preceding items, wherein said PB-PEO diblock copolymer comprises 5-50 blocks PB and 5-50 blocks PEO.
- [0309] 39. The polymersome according to any one of preceding items, wherein said amphiphilic polymer is a poly(dimethylsiloxane)-poly(ethylene oxide) (PDMS-PEO) diblock copolymer, wherein preferably said PB-PEO diblock copolymer preferably comprises 5-100 blocks PDMS and 5-100 blocks PEO.
- [0310] 40. The polymersome according to any one of preceding items, wherein said polymersomes may comprises of block copolymers or amphiphilic polymers only or mixed with lipids.
- [0311] 41. The polymersome according to anyone of preceding items, wherein said the lipids comprises of synthetic or natural lipids or lipid mixtures or combination of synthetic and natural lipids.
- [0312] 42. The polymersome according to any one of preceding items, wherein said amphiphilic polymer is a poly(lactide)-poly(ethylene oxide)/1-palmitoyl-2-oleoyl-sn-glycero-3-phospho-L-serine (PLA-PEO/POPC) copolymer, preferably said PLA-PEO/POPC has a ratio of 50:50 and above (e.g., 50/50 or 75/25 or 90/10) of PLA-PEO to POPC (e.g., PLA-PEO/POPC).
- [0313] 43. The polymersome according to any one of preceding items, wherein said amphiphilic polymer is a poly(caprolactone)-poly(ethylene oxide)/1-palmitoyl-2-oleoyl-sn-glycero-3-phospho-L-serine (PCL-PEO/POPC) copolymer, preferably said PCL-PEO/POPC has a ratio of 50:50 and above (e.g., 50/50 or 75/25 or 90/10) of PCL-PEO to POPC (e.g., PCL-PEO/POPC).
- [0314] 44. The polymersome according to any one of preceding items, wherein said amphiphilic polymer is polybutadiene-polyethylene oxide (BD) or a poly (dimethylsiloxane)-poly(ethylene oxide) (PDMS-PEO) diblock copolymer.
- [0315] 45. The polymersome according to any one of preceding items, wherein said polymersome comprises diblock copolymers PBD₂₁-PEO₁₄ (herein referred to as “BD21”), PDMS₄₇-PEO₃₆ (PDMS-PEO) or the triblock copolymer PMOXA₁₂-PDMS₅₅-PMOXA₁₂.
- [0316] 46. The polymersome according to any one of preceding items, wherein said polymersome comprises one or more compartments.
- [0317] 47. The polymersome according to any one of preceding items, wherein said polymersome comprises one or more compartments, wherein each one of said one or more compartments encapsulates at least one peptide, protein, and nucleic acid, preferably said at least one of said peptide, protein, and nucleic acid is immunogenic or antigenic, further preferably said each one of the one or more compartments is comprised of a same or different amphiphilic polymer.
- [0318] 48. The polymersome according to any one of preceding items, wherein said polymersome comprises more than one compartment, wherein said compartments comprise an outer block copolymer vesicle and at least one inner block copolymer vesicle, wherein said at least one inner block copolymer vesicle is encapsulated inside the outer block copolymer vesicle, preferably said outer block copolymer vesicle is a polymersome formed of a copolymer independently selected from the group consisting of:
- [0319] i) poly [styrene-b-poly(L-isocyanooalanine (2-thiophen-3-yl-ethyl)amide)](PS-PIAT),
- [0320] ii) poly(butadiene)-poly(ethylene oxide) (PBD-PEO),
- [0321] iii) poly(ethylene oxide)-poly(caprolactone) (PEO-PCL),
- [0322] iv) poly(ethyl ethylene)-poly(ethylene oxide) (PEE-PEO),
- [0323] v) poly(ethylene oxide)-poly(lactic acid) (PEO-PLA),
- [0324] vi) poly(isoprene)-poly(ethylene oxide) (PI-PEO),
- [0325] vii) poly(2-vinylpyridine)-poly(ethylene oxide) (P2VP-PEO),
- [0326] viii) poly(ethylene oxide)-poly(N-isopropylacrylamide) (PEO-PNIPAm),
- [0327] ix) poly(styrene)-poly(acrylic acid) (PS-PAA),
- [0328] x) poly(ethylene glycol)-polypropylene sulfide) (PEG-PPS),
- [0329] xi) poly(2-methyloxazoline)-poly(dimethylsiloxane)-poly(2-methyloxazoline) (PMOXA-PDMS-PMOXA),
- [0330] xii) poly(ethylene oxide)-poly(dimethylsiloxane)-poly(2-methyloxazoline) (PEO-PDMS-PMOXA), and
- [0331] xiii) poly(methylphenylsilane)-poly(ethylene oxide) (PMPS-PEO-PMPS-PEO-PMPS);
- [0332] further preferably said at least one inner block copolymer vesicle is a polymersome formed of a copolymer independently selected from the group consisting of:
- [0333] xiv) poly [styrene-b-poly(L-isocyanooalanine (2-thiophen-3-yl-ethyl)amide)](PS-PIAT),
- [0334] xv) poly(butadiene)-poly(ethylene oxide) (PBD-PEO),
- [0335] xvi) poly(ethylene oxide)-poly(caprolactone) (PEO-PCL),


- [0336] xvii) poly(ethyl ethylene)-poly(ethylene oxide) (PEE-PEO), poly(ethylene oxide)-poly(lactic acid) (PEO-PLA), xviii)
- [0337] xix) poly(isoprene)-poly(ethylene oxide) (PI-PEO),
- [0338] xx) poly(2-vinylpyridine)-poly(ethylene oxide) (P2VP-PEO),
- [0339] xxi) poly(ethylene oxide)-poly(N-isopropylacrylamide) (PEO-PNIPAm),
- [0340] xxii) poly(styrene)-poly(acrylic acid) (PS-PAA),
- [0341] xxiii) poly(ethylene glycol)-polypropylene sulfide) (PEG-PPS),
- [0342] xxiv) poly(2-methyloxazoline)-poly(dimethylsiloxane)-poly(2-methyloxazoline) (PMOXA-PDMS-PMOXA),
- [0343] xxv) poly(ethylene oxide)-poly(dimethylsiloxane)-poly(2-methyloxazoline) (PEO-PDMS-PMOXA), and
- [0344] xxvi) poly(methylphenylsilane)-poly(ethylene oxide) (PMPS-PEO-PMPS-PEO-PMPS).
- [0345] 49. The polymersome according to any one of the preceding items, wherein said polymersome comprises a lipid polymer.
- [0346] 50. The polymersome according to any one of the preceding items, wherein the polymersome further comprises encapsulated adjuvant.
- [0347] 51. A method for production of encapsulated antigen in polymersome, said method comprising:
- [0348] i) dissolving an amphiphilic polymer in chloroform, preferably said amphiphilic polymer is Polybutadiene-Polyethylene oxide (BD);
- [0349] ii) drying said dissolved amphiphilic polymer to form a polymer film;
- [0350] iii) adding a solubilized antigen to said dried amphiphilic polymer film from step ii), wherein said antigen is selected from the group consisting of:
- [0351] a) a polypeptide; preferably said polypeptide antigen is according any one of preceding items, further preferably said polypeptide antigen comprises a soluble portion of Influenza hemagglutinin, Swine Influenza hemagglutinin, Ovalbumin (OVA), Porcine epidemic diarrhea virus SPIKE protein, B16 peptide or MC38 peptide, most preferably said polypeptide antigen is at least 60% or more (e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to a polypeptide sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, and SEQ ID NOs: 12-14;
- [0352] b) a carbohydrate;
- [0353] c) a polynucleotide, wherein said polynucleotide is not an antisense oligonucleotide, preferably said polynucleotide is a DNA or mRNA molecule;
- [0354] d) a combination of (a) and/or (b) and/or (c);
- [0355] iv) rehydrating said polymer film from step iii) to form polymer vesicles;
- [0356] v) optionally, filtering polymer vesicles from step iv) to purify polymer vesicles monodisperse vesicles; and/or
- [0357] vi) optionally, isolating said polymer vesicles from step iv) or v) from the non-encapsulated antigen.
- [0358] 52. The method for production of encapsulated antigen in polymersome according to any one of preceding items, wherein said polymersome is the polymersome according to any one of preceding items.
- [0359] 53. A polymersome produced by a method for production of encapsulated antigen in polymersome according to any one of preceding items.
- [0360] 54. A polymersome comprising a soluble encapsulated antigen, wherein said soluble encapsulated antigen is a polynucleotide (e.g., single or double stranded) selected from a RNA molecule (e.g., mRNA, self-amplifying mRNA or siRNA, preferably mRNA) or a DNA molecule or combination/s thereof (e.g., DNA/RNA hybrid) and wherein the polymersome further comprises an ionizable lipid, preferably said polymersome is the polymersome according to any one of preceding items.
- [0361] 55. The polymersome according to any one of preceding items, wherein said ionizable lipid is a cationic ionizable lipid.
- [0362] 56. The polymersome according to any one of preceding items, wherein said ionizable lipid:
- [0363] i) is a tertiary amine (e.g., is not a quaternary amine) and/or comprising one or more tertiary amine moieties (e.g., not comprising a cationic quaternized ammonium moiety, i.e., ionizable lipid is not DOTAP); and/or
- [0364] ii) comprising a free electron pair at the nitrogen atom (e.g., ionizable nitrogen atom); and/or
- [0365] iii) can or cannot be protonated to become quaternary amine.
- [0366] 57. The polymersome according to any one of preceding items, wherein said ionizable lipid comprising or consisting of:
- [0367] i) ionizable lipid DLin-MC3-DMA (i.e., (6Z, 9Z, 28Z, 31Z)-Heptatriaconta-6,9,28,31-tetraen-19-yl 4-(dimethylamino) butanoate) having Formula I:




[0368] ii) ionizable lipid 319 having Formula II:




[0369] iii) ionizable lipid C12-200 having Formula III:




[0370] iv) ionizable lipid 5A2-SC8 having Formula IV:




-continued




[0371] v) ionizable lipid 306Oi10 having Formula V:




[0372] vi) ionizable lipid 5 having Formula VI:




[0373] vii) ionizable lipid SM-102 (i.e., heptadecan-9-yl 8-((2-hydroxyethyl) (6-oxo-6-(undecyloxy) hexyl)amino) octanoate) having Formula VII:




[0374] viii) ionizable lipid A9 having Formula VIII:




[0375] ix) ionizable lipid ALC-0315 (i.e., [(4-hydroxybutyl) azanediy]di(hexane-6,1-diyl)bis(2-hexyldecanoate) having Formula IX:




[0376] x) ionizable lipid Arcturus 2.2 (8.8) 4 C CH3 having Formula X:



[0377] xi) ionizable lipid Genevant CL1 having Formula XI:



[0378] xii) ionizable lipid ALC-0159 (i.e. 2-[(polyethylene glycol)-2000]-N,N-ditetradecylacetamide) having Formula XII:



- [0379] xiii) any one of (i)-(xii), wherein one or more alkyl chains thereof are further esterified.
- [0380] 58. The polymersome according to any one of preceding items, wherein said polynucleotide encoding one or more polypeptides, preferably said one or more polypeptides are viral polypeptides.
- [0381] 59. The polymersome according to any one of preceding items, wherein said polymersome is capable of increasing thermostability and/or storage stability and/or immunogenicity of said polynucleotide (e.g., mRNA) and/or one or more polypeptides encoded by said polynucleotide (e.g., mRNA) within said polymersome, preferably compared to that of the same polynucleotide (e.g., mRNA) and/or one or more polypeptides encoded by said same polynucleotide (e.g., mRNA) within another polymersome without ionizable lipid or within an ionizable lipid nanoparticle (LNP) comprising cholesterol under the same conditions, further preferably said thermostability and/or storage stability is increased in the temperature range from about -80°C . to about 4°C . (e.g., at -80°C ., -20°C . or at 4°C ., preferably at -80°C .), most preferably said immunogenicity is increased in the temperature range from about 36.5°C . to about 37.5°C .; further most preferably said increase is of about at least 5% (e.g., at least 10%, at least 20% or at least 30%).
- [0382] 60. The polymersome according to any one of preceding items, wherein said polymersome is an oxidation-stable polymersome.
- [0383] 61. The polymersome according to any one of preceding items, wherein said polymersome is capable of eliciting a $\text{CD8}^{(+)}$ T cell-mediated immune response, preferably said eliciting is an in vivo, ex vivo or in vitro eliciting, further preferably said $\text{CD8}^{(+)}$ T cell-mediated immune response is against one or more polypeptides encoded by said polynucleotide.
- [0384] 62. The polymersome according to any one of preceding items, wherein said polymersome is capable of eliciting a cellular immune response, wherein said cellular immune response comprises a $\text{CD8}^{(+)}$ T cell-mediated immune response, preferably said immune response is an in vivo, ex vivo or in vitro immune response, further preferably said $\text{CD8}^{(+)}$ T cell-mediated immune response is against one or more polypeptides encoded by said polynucleotide.
- [0385] 63. The polymersome according to any one of preceding items, wherein said polymersome is capable of eliciting a cellular and/or humoral immune response, wherein said cellular immune response comprises a $\text{CD8}^{(+)}$ T cell-mediated immune response, preferably immune response is an in vivo, ex vivo or in vitro immune response, further preferably said cellular and/or humoral immune is against one or more polypeptides encoded by said polynucleotide, most preferably said $\text{CD8}^{(+)}$ T cell-mediated immune response is against one or more polypeptides encoded by said polynucleotide.
- [0386] 64. The polymersome according to any one of preceding items, wherein said humoral immune response comprises production of specific antibodies, further preferably said immune response is an in vivo, ex vivo or in vitro immune response, most preferably said humoral immune is against one or more polypeptides encoded by said polynucleotide.
- [0387] 65. The polymersome according to any one of preceding items, wherein said polymersome is capable of enhancing the frequency of effector $\text{CD4}^{(+)}$ T cells, preferably said enhancing is an in vivo, ex vivo or in vitro enhancing, further preferably said frequency of effector $\text{CD4}^{(+)}$ T cells is against one or more polypeptides encoded by said polynucleotide.
- [0388] 66. The polymersome according to any one of preceding items, wherein said cellular immune response comprises a T-cell mediated immune response, preferably said immune response is an in vivo, ex vivo or in vitro immune response, further preferably said cellular immune response comprising a T-cell mediated immune response is against one or more polypeptides encoded by said polynucleotide.
- [0389] 67. The polymersome according to any one of preceding items, wherein said polymersome is capable of enhancing clonal expansion of antigen-specific $\text{CD8}^{(+)}$ T cells compared to a free antigen, preferably said expansion is an in vivo, ex vivo or in vitro expansion, further preferably said clonal expansion of antigen-specific $\text{CD8}^{(+)}$ T cells is against one or more polypeptides encoded by said polynucleotide.
- [0390] 68. The polymersome according to any one of preceding items, wherein said polymersome is capable of inducing antigen-specific effector $\text{CD8}^{(+)}$ T cells, preferably said inducing is an in vivo, ex vivo or in vitro inducing, further preferably said inducing antigen-specific effector $\text{CD8}^{(+)}$ T cells is against one or more polypeptides encoded by said polynucleotide.
- [0391] 69. The polymersome according to any one of preceding items, wherein said polymersome is capable of enhancing a cytotoxic phenotype of antigen-specific $\text{CD8}^{(+)}$ T cells, preferably said enhancing is an in vivo, ex vivo or in vitro enhancing, further preferably said enhancing a cytotoxic phenotype of antigen-specific $\text{CD8}^{(+)}$ T cells is against one or more polypeptides encoded by said polynucleotide.
- [0392] 70. The polymersome according to any one of preceding items, wherein said polymersome is capable of targeting of lymph node-resident macrophages and/or B cells, preferably said targeting is an in vivo, ex vivo or in vitro targeting, further preferably said targeting of lymph node-resident macrophages and/or B cells is against one or more polypeptides encoded by said polynucleotide.
- [0393] 71. The polymersome according to any one of preceding items, wherein said polymersome is capable of one or more of the following:
- [0394] i) eliciting a cellular immune response; preferably said cellular immune response comprises a $\text{CD8}^{(+)}$ T cell-mediated immune response; further preferably said cellular immune response is a $\text{CD8}^{(+)}$ T cell-mediated immune response; most preferably said immune response is against one or more polypeptides encoded by said polynucleotide;
- [0395] ii) releasing polymersome content in an oxidation-independent manner and triggering $\text{CD8}^{(+)}$ T cell-mediated immune response; preferably said immune response is against one or more polypeptides encoded by said polynucleotide;
- [0396] iii) stimulating an immune response against one or more polypeptides encoded by said polynucleotide;

- [0397] iv) triggering a cross-protection induced by a CD8⁽⁺⁾ T cell-mediated immune response; preferably said immune response is against one or more polypeptides encoded by said polynucleotide;
- [0398] v) delivering one or more polypeptides encoded by said polynucleotide to an antigen-presenting cell (APC);
- [0399] vi) triggering an immune response comprising a CD8⁽⁺⁾ T cell-mediated immune response and/or CD4⁽⁺⁾ T cell-mediated immune response; preferably said immune response is against one or more polypeptides encoded by said polynucleotide;
- [0400] vii) stimulating an immune response in a subject; preferably said immune response is against one or more polypeptides encoded by said polynucleotide;
- [0401] viii) immunizing a non-human animal; preferably said immunizing is against one or more polypeptides encoded by said polynucleotide;
- [0402] ix) said polymersome has an altered antigenicity compared to corresponding antigenicity of said antigen without said polymersome;
- [0403] x) said polymersome has an altered immunogenicity compared to corresponding immunogenicity of said antigen without said polymersome.
- [0404] 72. The polymersome according to any one of preceding items, wherein said polymersome has one or more of the following properties:
- [0405] i) said polymersome comprises an oxidation-stable membrane; and/or
- [0406] ii) said polymersome is synthetic; and/or
- [0407] iii) said polymersome is free from non-encapsulated antigens or in a mixture with non-encapsulated antigens; and/or
- [0408] iv) said polymersome comprises a membrane of an amphiphilic polymer;
- [0409] and/or
- [0410] v) said polymersome comprises amphiphilic synthetic block copolymers forming a vesicle membrane; and/or
- [0411] vi) said polymersome has a diameter greater than 70 nm, preferably said diameter ranging from about 100 nm to about 1 μ m, or from about 100 nm to about 750 nm, or from about 100 nm to about 500 nm, or from about 125 nm to about 250 nm, from about 140 nm to about 240 nm, from about 150 nm to about 235 nm, from about 170 nm to about 230 nm, or from about 220 nm to about 180 nm, or from about 190 nm to about 210 nm, most preferably said diameter is of about 200 nm; and/or
- [0412] vii) said polymersome has a vesicular morphology;
- [0413] viii) said polymersome is self-assembling;
- [0414] ix) said polymersome comprises an amphiphilic block copolymer, wherein said amphiphilic block copolymer: (a) is capable of sensing pH (e.g., is capable of undergoing protonation or de-protonation depending on the pH); and/or (b) is modified and/or substituted to become a pH sensitive moiety (e.g., is capable of undergoing protonation or de-protonation depending on the pH), preferably said modification and/or substitution comprising modifying and/or substituting a hydroxyl group (e.g., C—OH) into/with —NH₃ and carboxyl group (e.g., —COOH) (e.g., modifying and/or substituting PBD-PEO—OH functional group into PBD-PEO—NH₃ and PBD-PEO—COOH).
- [0415] 73. The polymersome according to any one of preceding items, wherein the polymersome is in the form of a collection of polymersomes, wherein the mean diameter of the collection of polymersomes is in the range of about 100 nm to about 1 μ m, or from about 100 nm to about 750 nm, or from about 100 nm to about 500 nm, or from about 125 nm to about 250 nm, from about 140 nm to about 240 nm, from about 150 nm to about 235 nm, from about 170 nm to about 230 nm, or from about 220 nm to about 180 nm, or from about 190 nm to about 210 nm.
- [0416] 74. The polymersome according to any one of the preceding items, wherein said polynucleotide encodes an immunogen.
- [0417] 75. The polymersome according to any one of the preceding items, wherein said polynucleotide is a polynucleotide encoding one or more viral polypeptides, preferably encoding an Influenza hemagglutinin, Swine Influenza hemagglutinin and/or Porcine epidemic diarrhea virus SPIKE protein, further preferably said polynucleotide encoding a polypeptide selected from the group consisting of: SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7 and SEQ ID NO: 8 and SEQ ID NOs: 12, 13 and 14.
- [0418] 76. The polymersome according to any one of preceding items, wherein said polynucleotide encoding one or more of the following:
- [0419] i) Influenza hemagglutinin (HA), preferably selected from the group consisting of: SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7 and SEQ ID NO: 8;
- [0420] ii) Swine Influenza hemagglutinin (HA), preferably SEQ ID NO: 6;
- [0421] iii) Ovalbumin (OVA), preferably SEQ ID NO: 4;
- [0422] iv) B16 peptide, preferably selected from the group consisting of: SEQ ID NO: 9, SEQ ID NO: 10 and SEQ ID NO: 11;
- [0423] v) MC38 peptide, preferably selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2 and SEQ ID NO: 3;
- [0424] vi) B16 and MC38 peptides, preferably said peptides are independently selected the groups: i) SEQ ID NOs: 1-3 and ii) SEQ ID NOs: 9-11; and/or
- [0425] vii) Porcine epidemic diarrhea virus SPIKE protein and a soluble fragment thereof, preferably a fragment of SEQ ID NO: 12, 13 or 14.
- [0426] 77. The polymersome according to any one of preceding items, wherein said block copolymer or amphiphilic polymer is essentially non-immunogenic or essentially non-antigenic, preferably said block copolymer or amphiphilic polymer is non-immunogenic or non-antigenic.
- [0427] 78. The polymersome according to any one of preceding items, wherein said block copolymer or said amphiphilic polymer is neither immunostimulant nor adjuvant.
- [0428] 79. The polymersome according to any one of preceding items, wherein said amphiphilic polymer comprises a diblock or a triblock (A-B-A or A-B-C) copolymer.

- [0429] 80. The polymersome according to any one of preceding items, wherein said amphiphilic polymer comprises a copolymer poly(N-vinylpyrrolidone)-b-PLA.
- [0430] 81. The polymersome according to any one of preceding items, wherein said amphiphilic polymer comprises at least one monomer unit of a carboxylic acid, an amide, an amine, an alkylene, a dialkylsiloxane, an ether or an alkylene sulphide, preferably said amine is capable of protonation or de-protonation depending on the pH.
- [0431] 82. The polymersome according to any one of preceding items, wherein the amphiphilic polymer is a polyether block selected from the group consisting of an oligo(oxyethylene) block, a poly(oxyethylene) block, an oligo(oxypropylene) block, a poly(oxypropylene) block, an oligo(oxybutylene) block and a poly(oxybutylene) block.
- [0432] 83. The polymersome according to any one of preceding items, wherein said amphiphilic polymer is a poly(butadiene)-poly(ethylene oxide) (PB-PEO) diblock copolymer, or wherein said amphiphilic polymer is a poly(dimethylsiloxane)-poly(ethylene oxide) (PDMS-PEO) diblock copolymer.
- [0433] 84. The polymersome according to item 36, wherein said PB-PEO diblock copolymer comprises 5-50 blocks PB and 5-50 blocks PEO or wherein said PB-PEO diblock copolymer preferably comprises 5-100 blocks PDMS and 5-100 blocks PEO.
- [0434] 85. The polymersome according to any one of preceding items, wherein said amphiphilic polymer is a poly(lactide)-poly(ethylene oxide)/1-palmitoyl-2-oleoyl-sn-glycero-3-phospho-L-serine (PLA-PEO/POPC) copolymer, preferably said PLA-PEO/POPC has a ratio of 75 to 25 (e.g., 75/25) of PLA-PEO to POPC (e.g., PLA-PEO/POPC).
- [0435] 86. The polymersome according to any one of preceding items, wherein said amphiphilic polymer is a poly(caprolactone)-poly(ethylene oxide)/1-palmitoyl-2-oleoyl-sn-glycero-3-phospho-L-serine (PCL-PEO/POPC) copolymer, preferably said PCL-PEO/POPC has a ratio of 75 to 25 (e.g., 75/25) of PCL-PEO to POPC (e.g., PCL-PEO/POPC).
- [0436] 87. The polymersome according to any one of preceding items, wherein said amphiphilic polymer is polybutadiene-polyethylene oxide (BD).
- [0437] 88. The polymersome according to any one of preceding items, wherein said polymersome comprises diblock copolymer PBD₂₁-PEO₁₄ (BD21) and the triblock copolymer PMOXA 12-PDMS₅₅-PMOXA₁₂.
- [0438] 89. The polymersome according to any one of the preceding items, wherein said polymersome comprises a lipid polymer.
- [0439] 90. The polymersome according to any one of the preceding items, wherein said polymersome comprising:
- [0440] i) polybutadiene-polyethylene oxide (BD) and Dlin-MC3-DMA ((6Z,9Z,28Z,31Z)-Heptatriaconta-6,9,28,31-tetraen-19-yl 4-(dimethylamino) butanoate) having Formula I;
 - [0441] ii) PBD-PEO (poly(butadiene)-poly(ethylene oxide)) and Dlin-MC3-DMA ((6Z,9Z,28Z,31Z)-Heptatriaconta-6,9,28,31-tetraen-19-yl 4-(dimethylamino) butanoate), preferably comprising 15% or 30% Dlin-MC3-DMA;
 - [0442] iii) polybutadiene-polyethylene oxide (BD) and ionizable lipid 319 having Formula II;
 - [0443] iv) polybutadiene-polyethylene oxide (BD) and ionizable lipid C12-200 having Formula III;
 - [0444] v) polybutadiene-polyethylene oxide (BD) and ionizable lipid 5A2-SC8 having Formula IV;
 - [0445] vi) polybutadiene-polyethylene oxide (BD) and ionizable lipid 306Oi10 having Formula V;
 - [0446] vii) polybutadiene-polyethylene oxide (BD) and ionizable lipid 5 having Formula VI;
 - [0447] viii) polybutadiene-polyethylene oxide (BD) and ionizable lipid SM-102 having Formula VII;
 - [0448] ix) polybutadiene-polyethylene oxide (BD) and ionizable lipid A9 having Formula VIII;
 - [0449] x) polybutadiene-polyethylene oxide (BD) and ionizable lipid ALC-0315 (i.e., [(4-hydroxybutyl) azanediy]di(hexane-6,1-diyl)bis(2-hexyldecanoate) having Formula IX;
 - [0450] xi) polybutadiene-polyethylene oxide (BD) and ionizable lipid Arcturus 2.2 (8.8) 4 C CH3 having Formula X;
 - [0451] xii) polybutadiene-polyethylene oxide (BD) and ionizable lipid Genevant CL1 having Formula XI;
 - [0452] xiii) polybutadiene-polyethylene oxide (BD) and ionizable lipid ALC-0159 (i.e. 2-[(polyethylene glycol)-2000]-N,N-ditetradecylacetamide) having Formula XII.
- [0453] 91. A composition comprising a polymersome according to any one of preceding items.
- [0454] 92. The composition according to any one of preceding items, wherein said composition is a pharmaceutical or diagnostic composition.
- [0455] 93. The composition according to any one of preceding items, wherein said composition is an immunogenic, antigenic or immunotherapeutic composition.
- [0456] 94. The composition according to any one of preceding items, further comprising one or more immunostimulants and/or one or more adjuvants.
- [0457] 95. The composition according to any one of preceding items, wherein said composition is a vaccine.
- [0458] 96. The composition according to any one of preceding items, formulated for intradermal, intraperitoneal, intramuscular, subcutaneous, intravenous injection, or non-invasive administration to a mucosal surface.
- [0459] 97. Isolated antigen presenting cells or a hybridoma cell exposed to the polymersome or composition according to any one of preceding items.
- [0460] 98. The antigen presenting cells according to any one of preceding items, wherein said antigen presenting cells comprise a dendritic cell.
- [0461] 99. The antigen presenting cells according to any one of preceding items, wherein said antigen presenting cells comprise macrophages.
- [0462] 100. The antigen presenting cells according to any one of preceding items, wherein said antigen presenting cells comprise B-cells.
- [0463] 101. A vaccine comprising the polymersome, composition, antigen presenting cells or hybridoma

according to any one of preceding items, and further comprising a pharmaceutically accepted excipient or carrier.

[0464] 102. The vaccine according to any one of preceding items, wherein:

[0465] i) said antigen comprises Influenza hemagglutinin (HA), wherein said vaccine is an Influenza vaccine, preferably said Influenza hemagglutinin (HA) is at least 60% or more (e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to polypeptide selected from the group consisting of: SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7 and SEQ ID NO: 8;

[0466] ii) said antigen comprises Swine Influenza hemagglutinin (HA), wherein said vaccine is Swine Influenza vaccine, preferably said Swine Influenza hemagglutinin (HA) is at least 60% or more (e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to SEQ ID NO: 6;

[0467] iii) said antigen comprises Ovalbumin (OVA), wherein said vaccine is a cancer vaccine, preferably said Ovalbumin (OVA) is at least 60% or more (e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to SEQ ID NO: 4;

[0468] iv) said antigen comprises SPIKE protein (PEDv S), wherein said vaccine is as PED Vaccine, preferably said Porcine epidemic diarrhea virus SPIKE protein (S protein) which is at least 80% or more (e.g., at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical of SEQ ID NOs: 12-14,

[0469] v) said antigen comprises B16 peptide, wherein said vaccine is a cancer vaccine, preferably said peptide is selected from the group consisting of: SEQ ID NO: 9-11;

[0470] vi) said antigen comprises MC38 peptide, wherein said vaccine is a cancer vaccine preferably said peptide is selected from the group consisting of: SEQ ID NO: 1-3;

[0471] vii) said antigen comprises B16 and MC38 peptides, wherein said vaccine is a cancer vaccine, preferably said peptides are independently selected the groups: i) SEQ ID NOs: 1-3 and ii) SEQ ID NOs: 9-11;

[0472] viii) said antigen is at least 60% or more (e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to a polypeptide sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11; wherein said vaccine is a cancer vaccine.

[0473] 103. A kit comprising the polymersome, composition, antigen presenting cells, hybridoma or vaccine according to any one of preceding items.

[0474] 104. A method of eliciting an immune response in a subject (e.g. human), comprising:

[0475] i) providing the polymersome, composition, antigen presenting cells, hybridoma or vaccine according to any one of preceding items to said subject,

[0476] ii) administering said polymersome, composition, antigen presenting cells, hybridoma or vaccine to said subject, preferably said administering is intradermal, intraperitoneal, intramuscular, subcutaneous, intravenous injection, or non-invasive administration to a mucosal surface.

[0477] 105. The method of eliciting an immune response according to any one of preceding items, wherein said immune response is a broad immune response.

[0478] 106. The method of eliciting an immune response according to any one of preceding items, wherein said immune response comprises a CD8⁽⁺⁾ T cell-mediated immune response and/or CD4⁽⁺⁾ T cell-mediated immune response.

[0479] 107. A method for the treatment or prevention of an infectious disease, a cancer or autoimmune disease in a subject in need thereof (e.g. human) comprising administering to said subject a therapeutically effective amount of the polymersome, composition, antigen presenting cells, hybridoma or vaccine according to any one of preceding items, preferably said infectious disease is a viral or bacterial infectious disease.

[0480] 108. A method for immunizing a non-human animal, said method comprising the following steps:

[0481] i) providing the polymersome, composition, antigen presenting cells, hybridoma or vaccine according to any one of preceding items;

[0482] ii) immunizing said non-human animal with said polymersome, composition, antigen presenting cells, hybridoma or vaccine.

[0483] 109. A method for preparation of an antibody, comprising:

[0484] i) immunizing a non-human mammal with the polymersome, composition, antigen presenting cells, hybridoma or vaccine according to any one of preceding items;

[0485] ii) isolating an antibody obtained in step (i).

[0486] 110. The method according to any one of preceding items, wherein said antibody is a monoclonal antibody (mAb).

[0487] 111. The polymersome, composition, antigen presenting cells, hybridoma or vaccine according to any one of preceding items, for use as a medicament.

[0488] 112. The polymersome, composition, antigen presenting cells, hybridoma or vaccine according to any one of preceding items for use in one or more of the following methods:

[0489] i) in a method of antibody discovery and/or screening and/or preparation;

[0490] ii) in a method of vaccine discovery and/or screening and/or preparation;

[0491] iii) in a method of production or preparation of an immunogenic or immunostimulant composition;

[0492] iv) in a method of targeted delivery of a protein and/or peptide, preferably said targeted delivery is a targeted delivery of an antigenic protein and/or peptide according to any one of preceding items; further preferably said antigenic protein and/

or peptide comprises a soluble portion of a membrane protein (MP) or a membrane-associated peptide (MAP), most preferably said antigen comprises a soluble portion of Influenza hemagglutinin, Swine Influenza hemagglutinin, Ovalbumin (OVA), Porcine epidemic diarrhea virus SPIKE protein, B16 peptide or MC38 peptide, further most preferably said antigen is at least 60% or more (e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to a polypeptide sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, and SEQ ID NOs: 12-14; further most preferably said targeted delivery is carried out in a subject;

[0493] v) in a method of stimulating an immune response to an antigen, preferably said antigen is according to any one of preceding items, further preferably said antigen comprises a soluble portion of Influenza hemagglutinin, Swine Influenza hemagglutinin, Ovalbumin (OVA), Porcine epidemic diarrhea virus SPIKE protein, B16 peptide or MC38 peptide; further most preferably said antigen is at least 60% or more (e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to a polypeptide sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, and SEQ ID NOs: 12-14; further most preferably for use in stimulating an immune response to said antigen in a subject;

[0494] vi) in a method of triggering cross-protection induced by CD8⁽⁺⁾ T cell-mediated immune response, preferably in a method of triggering cross-protection induced by CD8⁽⁺⁾ T cell-mediated immune response against an antigen is according to any one of preceding items according to any one of preceding items; further preferably said antigen comprises a soluble portion of Influenza hemagglutinin, Swine Influenza hemagglutinin, Ovalbumin (OVA), Porcine epidemic diarrhea virus SPIKE protein, B16 peptide or MC38 peptide; most preferably said antigen is at least 60% or more (e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to a polypeptide sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, and SEQ ID NOs: 12-14;

[0495] vii) in a method of delivering a peptide and/or protein to an antigen-presenting cells (APCs) according to any one of preceding items; preferably said peptide and/or protein is an antigen according to any one of preceding items; further preferably said antigen comprises a soluble portion of Influenza

hemagglutinin, Swine Influenza hemagglutinin, Ovalbumin (OVA), Porcine epidemic diarrhea virus SPIKE protein, B16 peptide or MC38 peptide; most preferably said antigen is at least 60% or more (e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to a polypeptide sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, and SEQ ID NOs: 12-14;

[0496] viii) in a method of triggering an immune response comprising a CD8⁽⁺⁾ T cell-mediated immune response and/or CD4⁽⁺⁾ T cell-mediated immune response; preferably said response is against an antigen according to any one of preceding items; further preferably said antigen comprises a soluble portion of Influenza hemagglutinin, Swine Influenza hemagglutinin, Ovalbumin (OVA), Porcine epidemic diarrhea virus SPIKE protein, B16 peptide or MC38 peptide; further preferably said response is against an antigen which is at least 60% or more (e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100%) identical to a polypeptide sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, and SEQ ID NOs: 12-14;

[0497] ix) in a method for treatment, amelioration, prophylaxis or diagnostics of an infectious disease, preferably said infectious disease is a viral or bacterial infectious disease; further preferably said viral infectious disease is selected from a group consisting of: influenza infection, respiratory syncytial virus infection, herpes virus infection.

[0498] x) in a method for treatment, amelioration, prophylaxis or diagnostics of a cancer or an autoimmune disease;

[0499] xi) in a method for sensitizing cancer cells to chemotherapy;

[0500] xii) in a method for induction of apoptosis in cancer cells;

[0501] xiii) in a method for stimulating an immune response in a subject;

[0502] xiv) in a method for immunizing a non-human animal;

[0503] xv) in a method for preparation of hybridoma;

[0504] xvi) in a method according to any one of preceding items;

[0505] xvii) in a method according to any one of preceding i)-xvi), wherein said method is in vivo and/or ex vivo and/or in vitro method;

[0506] xviii) in a method according to any one of preceding i)-xvii), wherein said antigen is heterologous to the environment in which said antigen is used.

[0507] 113. Use of the polymersome, composition, antigen presenting cells, hybridoma or vaccine according to any one of preceding items for one or more of the following:

[0508] i) for antibody discovery and/or screening and/or preparation;

[0509] ii) for vaccine discovery and/or screening and/or preparation;

[0510] iii) for production or preparation of an immunogenic or immunostimulant composition;

[0511] iv) for targeted delivery of proteins and/or peptides, preferably said targeted delivery is a targeted delivery of antigenic proteins and/or peptides; further preferably said targeted delivery is carried out in a subject;

[0512] v) for stimulating an immune response to an antigen, preferably for use in stimulating an immune response to an antigen in a subject;

[0513] vi) for triggering cross-protection induced by a CD8⁽⁺⁾ T cell-mediated immune response;

[0514] vii) for delivering a peptide or protein to an antigen-presenting cell (APC); preferably said peptide or protein is an antigen, further preferably said peptide or protein is immunogenic or immunotherapeutic;

[0515] viii) for triggering an immune response comprising a CD8⁽⁺⁾ T cell-mediated immune response and/or CD4⁽⁺⁾ T cell-mediated immune response;

[0516] ix) in a method for treatment, amelioration, prophylaxis or diagnostics of an infectious disease, preferably said infectious disease is a viral or bacterial infectious disease; further preferably said viral infectious disease is selected from a group consisting of: influenza infection, respiratory syncytial virus infection; herpes virus infection;

[0517] x) for treatment, amelioration, prophylaxis or diagnostics of a cancer or an autoimmune disease;

[0518] xi) for sensitizing cancer cells to chemotherapy;

[0519] xii) for induction of apoptosis in cancer cells;

[0520] xiii) for stimulating an immune response in a subject;

[0521] xiv) for immunizing a non-human animal;

[0522] xv) for preparation of hybridoma;

[0523] xvi) in a method according to any one of preceding items;

[0524] xvii) for use according to any one of preceding i)-xvi), wherein said use is in vivo and/or ex vivo and/or in vitro use;

[0525] xviii) for use according to any one of preceding i)-xvii), wherein said antigen is heterologous to the environment in which said antigen is used.

[0526] 114. A method of eliciting an immune response in a subject, comprising administering to a subject a polymersome formed from PDMS-PEO carrying an antigen.

[0527] 115. The method of item 114, wherein the antigen is encapsulated within the PDMS-PEO polymersome.

[0528] 116. The method of item 115, wherein the antigen encapsulated in the PDMS-PEO polymersome is a soluble antigen.

[0529] 117. The method of item 116, wherein the antigen is selected from the group consisting of a polypeptide, a carbohydrate, a polynucleotide and combinations thereof.

[0530] 118. The method of item 114, wherein the antigen is integrated into the circumferential membrane of the PDMS-PEO polymersome.

[0531] 119. A PDMS-PEO polymersomes carrying an antigen.

[0532] 120. A polymersome of item 119, wherein the antigen is encapsulated within the PDMS-PEO polymersome.

[0533] 121. The polymersome of item 120, wherein the antigen encapsulated in the PDMS-PEO polymersome is a soluble antigen.

[0534] 122. The polymersome of item 121, wherein the antigen is selected from the group consisting of a polypeptide, a carbohydrate, a polynucleotide and combinations thereof.

[0535] 123. The polymersome of item 122, wherein the antigen is integrated into the circumferential membrane of the PDMS-PEO polymersome

[0536] 124. The polymersome of item 123, wherein the antigen is a membrane-associated protein or lipid antigen.

[0537] 125. The polymersome of item 124, wherein the membrane-associated protein is selected from the group consisting of a transmembrane protein, G protein-coupled receptor, neurotransmitter receptor, kinase, porin, ABC transporter, ion transporter, acetylcholine receptor, and a cell adhesion receptor.

[0538] 126. A pharmaceutical composition comprising a polymersome of any of items 119 to 125.

[0539] 127. The in vitro and in vivo use of a PDMS-PEO as defined in any of items 119 to 125 or a pharmaceutical composition of item 126 for eliciting an immune response.

EXAMPLES OF THE INVENTION

[0540] In order that the invention may be readily understood and put into practical effect, some aspects of the invention are described by way of the following non-limiting examples.

Materials and Methods

Example 1: Encapsulation of Ovalbumin, Peptides, Soluble HA, PEDv SPIKE Protein and eGFP DNA in Polymersomes

[0541] A 100 mg/ml stock of Polybutadiene-Polyethylene oxide (herein referred to as "BD21") is dissolved in chloroform. 100 μ L of the 100 mg/ml BD21 stock is then deposited into a borosilicate (12 $\times$ 75 mm) culture tube and slowly dried under a stream of nitrogen gas to form a thin polymer film. The film was further dried under vacuum for 6 hours in a desiccator. A 1 mL solution of 1-5 mg/ml solubilized Ovalbumin (OVA) protein in 1 $\times$ PBS buffer was then added to the culture tube. The mixture was stirred at 600 rpm, 4 $^{\circ}$ C. for at least 18 hours to rehydrate the film and to allow the formation of polymer vesicles. The turbid suspension was extruded through a 200-nm pore size Whatman Nucleopore membrane with an extruder (Avanti 1 mL liposome extruder, 21 strokes) to obtain monodisperse

vesicles [e.g., Fu et al., 2011, Lim. S. K, et al., 2017]. The protein containing BD21 polymer vesicles were purified from the non-encapsulated proteins by dialyzing the mixture against 1 L of 1× PBS using a dialysis membrane (300 kDa MWCO, cellulose ester membrane).

[0542] The final vesicle mixture was analysed for non-encapsulated protein using size-exclusion chromatography. Fractions of the vesicle peak from SEC were used to quantify the amount of protein encapsulation via SDS-PAGE. Vesicle size and mono-dispersity was characterized by dynamic light scattering instrument (Malvern, United Kingdom) (100× dilution with 1× PBS). For quantification of OVA encapsulated in polymersomes, samples were pre-treated with 20% DMSO followed by sample buffer, after which they were loaded on to the SDS-PAGE analysis.

[0543] For peptides encapsulation (exemplified by MC 38 neo-antigen peptides of SEQ ID NO: 1, 2 and 3), a similar protocol was followed. Peptides concentration was 0.5-0.3 mg/ml dissolved in PBS for encapsulation. After dialysis, an amount of encapsulated peptides was determined using Phenylalanine fluorescence (ex 270 nm/em 310 nm) using a Cary Eclipse Spectrophotometer (Agilent). Encapsulation of all 3 peptides was performed individually and concentration was determined to be 20-30 ug/ml for all peptides. An equivalent volume of each of 3 encapsulated peptides was mixed together just before injection into mice.

[0544] For HA encapsulation, a similar protocol was followed. Recombinant HA (H1N1/A/Puerto Rico/8/1934 strain) at a concentration of 10 µg/ml was dissolved in PBS for encapsulation. After dialysis, an amount of encapsulated peptides was determined by western blot. HA concentration after encapsulation was determined to be around 1 ug/ml. 100 ul were injected in mice.

[0545] For PEDv SPIKE protein encapsulation in BD21 polymersomes, a similar protocol was followed as described above. PEDv SPIKE protein (different constructs, SEQ ID Nos: 12-14) were expressed using Baculovirus expression system. Proteins isolated from the insect cells were added for encapsulation. Whereas, for encapsulation of PEDv SPIKE protein in polymersomes made of poly(dimethyl siloxane)-poly(ethylene oxide) (PDMS₄₆-PEO₃₇ obtained from Polymer Source, Quebec, Canada), or a mixture of block copolymers and lipids such as PDMS₄₆-PEO₃₇ (DSPE-PEG, PLA-PEG/POPC, PLA-PEG/Asolectin, a different protocol was followed in order to show the generality of the methods. Polymer and or polymer lipid mixture were dissolved in ethanol or any water miscible solvent and added dropwise to a protein solution to self-assemble and the proteins are encapsulated into polymersomes during self-assembly. Non-encapsulated proteins were removed by dialysis with PBS. After dialysis, amount of each polymersome sample encapsulated proteins was determined by densitometry. The concentration of proteins after encapsulation was determined to be around 1 µg/ml for each of these polymersome formulations. Polymersomes were encapsulated either with soluble SPIKE protein (SEQ 12) or S1 region of SPIKE protein (SEQ 13) and S2 region of SPIKE protein (SEQ 14). 100-200 µl of polymersomes (either only with soluble SPIKE protein or with mixture of polymersomes with S1 and S2 region of SPIKE proteins) were injected in mice and 1 ml of such polymersomes was orally administered to pigs.

[0546] For eGFR DNA encapsulation, a similar protocol as OVA encapsulation was followed. Briefly, block co-

polymers such as poly(butadiene)-poly(ethyleneoxide) (BD21), poly(butadiene)-poly(ethyleneoxide) modified with functional groups (e.g., NH₂, COOH) at the end of poly(ethylene oxide) chain (BD21-NH₂), mixture of block copolymers and lipids such as PLA-PEG/POPC, PLA-PEG/Asolectin, Dimethylaminoethane-carbamoyl (DC)-Cholesterol, 1,2-dioleoyl-3-trimethylammonium-propane (DOTAP) were dissolved in chloroform and transferred to a glass tube and slowly dried under a stream of nitrogen gas to form a thin film. The film was further dried under vacuum for 6 hours in a desiccator. 1 µg of eGFP DNA was added to the film and rehydrated overnight. Afterwards, the samples were extruded with 0.2 µm polycarbonate filter and dialyzed in HEPES Buffer.

Example 2: Transfection of eGFP DNA Encapsulated Polymersomes with HEK293T Cells

[0547] HEK293T cells were seeded with a density of 50,000 cells/well into a 48-well plate. For transfections (the Lipofectamine 2000 transfection), 1,000 µL of Opti-MEM I (Invitrogen), 2 µL of Lipofectamine 2000 (Invitrogen), and 1 µg of SF-GFP PC DNA (or polymersomes formulation containing 1 µg of SF-GFP PC DNA) were mixed. The transfection complexes were formed during 20 min incubation at RT. For transfection, the lipofectamine complex was added to the cells and incubated for 24 hr to 72 hr at 37 C and 5% CO₂. The efficiency of transfection was measured by GFP fluorescence, Ex 485 nm, Em 520 nm). For cellular uptake fluorescence measured at Ex 530 nm Em 560 nm. For imaging, aspirated the cell media followed by washed the cells with DPBS (with Ca²⁺/Mg²⁺) and fixed with 4% p-formaldehyde. Then, the glass cover-slip was removed and flipped into a glass slide containing a drop of 20 ul mounting media with DAPI. Finally, sealed the cover-slip with nail polish and stored at 4 C for future imaging. Fluorescence microscopy was used for imaging.

Example 3: Immunization of OVA Encapsulated Polymersomes for Antibody Titers

[0548] C57bl/6 mice were immunized using free OVA with or without Sigma Adjuvant System (SAS) and OVA encapsulated ACMs (polymersomes) by doing a prime and a boost 21 days later. All immunizations were performed with a final amount of OVA: 5-10 ug OVA/injection/mouse. Final bleeds were collected 42 days after prime. ELISA was then performed to assess titers: OVA was coated onto MaxiSorp plates (1 ug/ml) overnight. Plates were blocked using 3% BSA in PBS for 1 h at RT. All sera were diluted at 1:100 and incubated on plates for 1 h at RT. After 3 washes with PBS+0.05% Tween 20, secondary antibody anti-mouse IgG HRP coupled was incubated at 1:10,000 dilution for 1 h, RT (room temperature). After 3 washes with PBS/Tween 20 buffer, TMB substrate was added and reaction was stopped using 1M HCl. Optical densities were quantified at 450 nm.

Example 4: Immunization of HA Encapsulated Polymersomes for Antibody Titers

[0549] Similarly, Balb/c mice were immunized with free HA proteins (SEQ ID NO: 7), ACM encapsulated HA (polymersomes) in PBS or PBS control. All immunizations were performed with a same final amount of HA: 100 ng

HA/injection/mouse. Final bleeds were collected 42 days after prime and ELISA were performed as above using 1 ug/ml HA for plate coating.

Example 5: Immunization of MC 38 Neo-Antigen Peptides Encapsulated Polymersomes for Cellular Response

[0550] To observe a specific CD8 T cell response after immunization we used a MC-38 syngeneic tumour model. C57bl/6 mice were inoculated with subcutaneously at the right flank with MC-38 tumour cells (3×10^5) in 0.1 ml of PBS for tumour development. The inoculation day is defined as Day 0. The animals were randomized based on the bodyweights and immunizations were started at day 4 after the inoculation. Immunizations consisted of: free peptides, ACM encapsulated peptides (polymersomes) with and without co-treatment with a commercially available anti-PD-1 antibody. Peptides were: Reps1 P45A (SEQ ID NO: 1), Adpgk R304M (SEQ ID NO: 2) and Dpagt1 V213L (SEQ ID NO: 3) and were obtained from Genscript. 200 ul of peptides and peptides in ACMs were immunized subcutaneously on day 4, 11 and 18. The concentration of peptides in ACMs was determined to be 20-30 ug/ml, whereas for peptides alone 10 ug per injection per mice was used. The anti-PD1 antibody was injected intraperitoneally on day 5, 8, 12, 15, 19 and 22 at 5 mg/kg dosage. Animals were checked for any effects of tumour growth and treatments on normal behaviour such as mobility, food and water consumption, body weight gain/loss (body weights will be measured 3 times per week). Tumour sizes were measured 3 times per week in two dimensions using a caliper, and the volume was expressed in mm^3 using the formula: $V = 0.5 \times a \times b^2$ where a and b are the long and short diameters of the tumour, respectively.

Example 6: Immunization of Mice and Pigs with PEDv Spike Protein Encapsulated Polymersomes

[0551] Mice were immunised with ACM encapsulated PEDv spike protein and boosted with a second dose after 21 days, 150 ul-200 ul of polymersomes encapsulated with PEDv Spike protein were immunized. Sera was collected from the final bleed and was used for ELISA. Moreover, these sera were tested for their ability to neutralise the PEDV strain USA/Colorado/2013 (CO/13) through a conventional virus neutralisation. Furthermore, weaned pigs were orally vaccinated with 1 ml of polymersome encapsulated with PED SPIKE protein (after a prime on day 1 and a boost on day 14). A simple physiological solution was used for the oral vaccination.

Results

Encapsulation of Proteins and DNA of Example 1

[0552] OVA encapsulated polymersomes were purified by dialysis and size exclusion column (SEC) to remove the non-encapsulated proteins and analysed by dynamic light scattering. As shown in FIGS. 2A, FIGS. 3 and 4, an elution profile of OVA encapsulated polymersomes from SEC and a monodisperse population was observed.

[0553] Dynamic light scattering (DLS) data is presented in FIG. 2B for various of different polymersomes having encapsulated therein OVA, PEDv SPIKE protein or eGFP DNA. They are all measuring a mean diameter of 120

nm-180 nm using Z-average (d, nm), a preferred DLS parameter. Z-average size is the intensity weighted harmonic mean particle diameter, the values are in good agreement with earlier data [Fu et al., 2011, Lim. S. K, et al., 2017] of polymersomes.

Encapsulated eGFP DNA and Transfection Data of Example 2

[0554] eGFP DNA encapsulated polymersomes were transfected with HEK293T cells and after transfection, the uptake of ACM polymersomes were measured by fluorescence plate reader at Ex 530 nm and Em 560 nm and the transfection efficiency was measured by the GFP fluorescence (Ex 485 nm, Em 520 nm). As shown in FIG. 5, it is clear that polymersomes with DNA are able to penetrate into the cells and releasing the DNA to express the DNA to protein, all the polymersome formulations are taken up by the cells and are able to release the DNA, whereas the ratio of the polymersomes release versus the protein expression correlates well with its stability and biodegradability (FIG. 5A). Non-biodegradable polymersomes such as BD21 are taken up in smaller amount and the expression levels were lower comparing to the biodegradable polymersomes. Similar results were observed from the fluorescence images of cells as well (FIG. 5B & FIG. 5C).

Encapsulated OVA and Titers of Example 3

[0555] OVA encapsulated polymersomes were immunized in C57bl/6 mice by doing a prime and a boost 21 days later. Final bleeds were used for performing the ELISA. As shown in FIG. 6, it is clear that OVA encapsulated polymersomes is the only formulation able to trigger a titer in comparison to free OVA, OVA with adjuvants or control samples (PBS alone). The reason why OVA with SAS did not produce a titer may be due to the small amount of OVA used in the trial (around 5 ug per injection). Hence ACM encapsulated OVA was able to trigger a B cell response toward OVA in the form of an IgG serum titer specific for Ovalbumin.

Encapsulated HA and Titers of Example 4

[0556] HA (H1N1/A/Puerto Rico/8/1934 strain, SEQ ID NO: 7) encapsulated polymersomes were immunized in Balb/c mice by doing a prime and a boost 21 days later. Final bleeds were used for performing the ELISA. As shown in FIG. 7, it is clear that HA encapsulated polymersomes is the only formulation able to trigger a titer in comparison to free HA or control samples (PBS alone). The reason why free HA did not produce a titer may be due to the small amount of HA used in the trial (around 100 ng per injection). Hence ACM encapsulated HA was able to trigger a B cell response toward HA in the form of an IgG serum titer specific for HA.

Encapsulated MC-38 Neo Antigen Peptides and CD8 T Cell Response of Example 5

[0557] In order to show that ACM encapsulated antigen are able to trigger a CD8 T cell response we used a well-defined MC-38 syngeneic mouse tumour model which relies on the delivery of known CD8 antigenic peptides. High quantities of these peptides combined with adjuvants have been shown to trigger tumour control in therapeutic mouse models (e.g., Kuai et al., 2017, Luo et al., 2017). In addition, these effects were clearly correlated to the presence of peptide-specific CD8 T cells in the mouse blood. Hence

any tumour development difference between groups would be directly attributed to the presence of a peptide-specific pool of CD8 T cells. 4 days after inoculation with MC-38 cell lines, mice were immunized with either free peptides, ACM encapsulated peptides (polymersomes) with and without anti-PD1 antibody treatment as described in the section Materials and Methods herein. As shown in FIG. 8, immunization with encapsulated peptides was able to trigger an inhibitory effect in tumour development compared to free peptides. This effect was dramatically potentiated whenever anti-PD1 antibody injections were added. This data demonstrated that ACM encapsulated peptides (polymersomes) were able to trigger a peptide-specific CD8 T cell response most likely via the delivery of these peptides to dendritic cells, which resulted in tumour control. This effect was increased by addition of a checkpoint inhibitor such as the anti-PD1 antibody. Indeed, MC-38 has been shown to express PD-L1 molecule at their cell surface which is known to inhibit T cells killing activity inside tumours. Hence inhibition of such interaction by an antibody blocking PD1/PD-L1 interaction is known to reveal the presence of tumour specific T cells even further. Altogether this data clearly demonstrates that ACM encapsulated antigens (polymersomes) were able to trigger an antigen specific CD8 T cell response without addition of adjuvant.

Encapsulated PEDv Spike Protein and IgG, IgA
and Virus Neutralisation Response of Example 6

[0558] Mice were immunised with ACM encapsulated PEDv spike protein and boosted with a second dose after 21 days. Sera was collected from the final bleed and was used for ELISA. As can be seen in FIG. 8, antibodies that bind to SPIKE Protein coated on ELISA PLATE and the titers are of similar level to the animals vaccinated with killed virus in comparison with ACM vaccinated mice. Moreover, the sera were tested for their ability to neutralise the PEDV strain USA/Colorado/2013 (CO/13) through a conventional virus neutralisation experiment (FIG. 9). In here, it was observed that the virus neutralization occurs only for the sera from the mice immunized with ACM vaccine (i.e., ACM encapsulated PED Spike protein) while no neutralization was observed for the sera from the mice vaccinated with killed virus. Furthermore, different polymersomes (e.g., BD21, PDMS₄₆-PEO₃₇ (marked in FIG. 10 only as “PDMS”), PDMS₄₆-PEO₃₇/DSPE-PEG, PLA-PEG/Asolectin lipids) encapsulated with full length soluble protein and polymersomes mixture containing S1 and S2 region were immunized and the sera were tested for virus neutralization (FIG. 10). From FIG. 10, it is evident that the groups of mice immunized with PBS sample doesn't show any virus neutralization, whereas all other polymersome formulation shows varying degree of virus neutralization. Furthermore, when weaned pigs were orally vaccinated with ACM encapsulated PED SPIKE protein (after a prime on day 1 and a boost on day 14, an increase in specific IgA antibodies against the virus was observed from the faecal swabs collected and measured via ELISA, (see FIG. 11).

Example 7: Improved Stability (e.g., Storage) of
Polymersomes Comprising a Soluble Encapsulated
mRNA and an Ionizable Lipid

[0559] OVA mRNA (SEQ ID NO: 16) were prepared with ACM-polymersomes according to the following protocol/conditions/compositions:

[0560] 10 ug/ml OVA mRNA was prepared in 20 mM Tris, 4.5 mM acetate, 5% sucrose (e.g., based on Moderna LNP formulation).

[0561] BD/DOTAP (85/15) 400 mg/ml ACM-polymersomes were prepared in THF. ACM-polymersomes prepared with BD/DOTAP are used as reference example in Example 7. Accordingly, ACM-polymersomes with ionizable lipids can be prepared in the same way as BD/DOTAP polymersomes.

[0562] 11 ml (9 ml for homogenization, 1 ml for extrusion, 1 ml non-extruded).

[0563] Homogenization was carried out as follows: 5 strokes final pressure 75-80 Mpa-153 nm Pdl 0.18.

[0564] Homogenised batch split—4 ml dialysed 1000 kDa membrane, 4 ml split between 4° C., -20° C., -80° C.

[0565] Extruded batch split—0.5 ml dialysed, 0.5 ml split between 4° C., -20° C., -80° C.

[0566] Co-solvent batch split—0.5 ml dialysed, remainder split between 4° C., -20° C.

[0567] Subsequently, mRNA concentration was measured after homogenisation and mRNA encapsulation into ACM-polymersomes (Table 1 and Table 2).

TABLE 1

mRNA concentration measurement after homogenisation and OVA-mRNA encapsulation into ACM-polymersomes in the presence of detergent (Triton):			
With Triton X-100 (detergent)	Dilution	Concentration (µg/ml)	Average (µg/ml)
OVA mRNA 10 µg/ml	1x	7.6	8.9
stock	2x	10.3	
After co-solvent	1x	7.5	7.5
After homogenisation	1x	7.1	8.0
	2x	8.8	

TABLE 2

mRNA concentration measurement after homogenisation and OVA-mRNA encapsulation into ACM-polymersomes in the absence of detergent (Triton):			
Without Triton X-100 (no detergent)	Dilution	Concentration (µg/ml)	Average (µg/ml)
OVA mRNA 10 µg/ml	1x	5.5	9.5
stock	2x	9.5	
After co-solvent	1x	0.1	0.1
After homogenisation	1x	-0.2	-1.7
	2x	-1.7	

[0568] The results in Table 2 demonstrate successful mRNA encapsulation into ACM-polymersomes after homogenization in the absence of detergent.

[0569] Furthermore, the storage stability of OVA-mRNA encapsulated in ACM-polymersomes has been studied over the weekend at 4° C. (Table 3).

TABLE 3

Storage stability of OVA-mRNA encapsulated in ACM-polymersomes at 4° C.:			
ACM-mRNA (stored over weekend 4° C.)		µg/ml OVA mRNA concentration	
		Triton	No Triton
Co-solvent	Crude	7.4	2.2
	Dialysed	7.6	1.7
Extrusion	Crude	4.0	1.5
	Dialysed	3.2	1.0
Homogenisation	Crude	7.3	1.4
	Dialysed	8.8	1.7

[0570] Consequently, DLS profiles of OVA-mRNA loaded ACM BD/DOTAP polymersomes (as above) and LNPs (Onpattro composition, Tables 4-5 below) were compared in order to assess mRNA sample stability.

[0571] For that the following protocol/conditions/compositions were used:

[0572] Buffer: 20 mM Tris, 5% sucrose, pH 7.4.

[0573] ACM/LNP molar content: 5 µM.

[0574] OVA-mRNA (SEQ ID NO: 16) content: 25 µg/ml.

[0575] Storage temperatures: at -80° C.

[0576] DLS results are shown in FIG. 13 demonstrating superior stability of OVA-mRNA loaded into ACM-polymersomes over OVA-mRNA loaded into LNPs (Onpattro composition) at -80° C.

TABLE 4

Onpattro composition: Onpattro					
Component	MW	mg (0.5 mL)	mg/mL	mM	mol %
Dlin-MC3-DMA	642.1	13.0	13.0	20.25	49.3%
PEG2000-DMG	2509.2	1.6	1.6	0.64	1.6%
DSPC	790.1	3.3	3.3	4.18	10.2%
Cholesterol	386.7	6.2	6.2	0.04	39.0%
Total		24.1	24.1	41.1	100%

TABLE 5

Onpattro composition (cont.): Onpattro			
Component	MW	mg (0.3 mL)	mM
Potassium phosphate monobasic	136.1	0.2	1.5
Sodium phosphate dibasic	268.1	2.3	8.6
Sodium chloride	58.4	8.8	151

[0577] Further, the following empty ACM-polymersomes compositions were prepared:

[0578] Empty ACM-polymersomes with DOTAP (prepared as described above, BD/DOTAP polymersomes are used as a reference example): ACM concentration: 5 µM (7.7-8.6 mg/ml); Polymer (mol %): 85 or 70; DOTAP (mol %): 15 or 30; Buffer: 20 mM Tris 4.5 mM acetate 5% sucrose pH 7.4.

[0579] Empty ACM-polymersomes with DLin-MC3-DMA (BD/DLin-MC3-DMA): ACM concentration: 5 µM (7.6-8.6 mg/ml); Polymer (mol %): 85 or 70; MC3 (mol %):

15 or 30; Starting buffer: 20 mM sodium acetate pH 5 or 20 mM Tris 4.5 mM acetate 5% sucrose pH 7.4; Final buffer: 20 mM Tris 4.5 mM acetate 5% sucrose pH 7.4.

[0580] Subsequently, empty ACM polymersomes prepared with DLin-MC3-DMA ionizable lipid (BD/DLin-MC3-DMA) were compared to the reference DOTAP polymersomes (empty) by DLS demonstrating that both are formed in the same way (FIG. 14).

[0581] Further ACM-polymersomes comprising PBD-PEO and DLin-MC3-DMA were also prepared (Table 6):

TABLE 6

ACM-polymersomes with PBD-PEO and DLin-MC3-DMA:						
Materials		Mw (g/mol)	Concen- tration (mM)	Concen- tration (mM)	per 1 ml of formulation	
					mmoles	Mass (mg)
15%	PBD-PEO	1900	5.00	4.25	4.25E-03	8.08
MC3	DLin-MC3-DMA	642.1		0.75	7.50E-04	0.48
				Total	5.00E-03	8.56
30%	PBD-PEO	1900	5.00	3.50	3.50E-03	6.65
MC3	DLin-MC3-DMA	642.1		1.50	1.50E-03	0.96
				Total	5.00E-03	7.61

[0582] One skilled in the art would readily appreciate that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those inherent therein. Further, it will be readily apparent to one skilled in the art that varying substitutions and modifications may be made to the invention disclosed herein without departing from the scope and spirit of the invention. The compositions, methods, procedures, treatments, molecules and specific compounds described herein are presently representative of certain embodiments are exemplary and are not intended as limitations on the scope of the invention. Changes therein and other uses will occur to those skilled in the art which are encompassed within the spirit of the invention are defined by the scope of the claims. The listing or discussion of a previously published document in this specification should not necessarily be taken as an acknowledgement that the document is part of the state of the art or is common general knowledge.

[0583] The invention illustratively described herein may suitably be practiced in the absence of any element or elements, limitation or limitations, not specifically disclosed herein. Thus, for example, the terms “comprising”, “including”, “containing”, etc. shall be read expansively and without limitation. Additionally, the terms and expressions employed herein have been used as terms of description and not of limitation, and there is no intention in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof, but it is recognized that various modifications are possible within the scope of the invention claimed. Thus, it should be understood that although the present invention has been specifically disclosed by exemplary embodiments and optional features, modification and variation of the inventions embodied herein may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within the scope of this invention.

[0584] The invention has been described broadly and generically herein. Each of the narrower species and sub-generic groupings falling within the generic disclosure also form part of the invention. This includes the generic description of the invention with a proviso or negative limitation removing any subject matter from the genus, regardless of whether or not the excised material is specifically recited herein.

[0585] Other embodiments are within the following claims. In addition, where features or aspects of the invention are described in terms of Markush groups, those skilled in the art will recognize that the invention is also thereby described in terms of any individual member or subgroup of members of the Markush group.

REFERENCES

- [0586] Liu et al., Immune responses to vaccines delivered by encapsulation into and/or adsorption onto cationic lipid-PLGA hybrid nanoparticles, *Journal of Controlled Release*, 225 (2016): 230-239.
- [0587] Hilbert et al., Biodegradable microspheres containing influenza A vaccine: immune response to mice, *Vaccine* 17 (1999): 1065-1073.
- [0588] Miller et al., Adjuvant compositions and related methods, U.S. Pat. No. 9,636,397B2, Gerber et al., Adjuvant and vaccine compositions, US2015/0044242 A1.
- [0589] Maji et al., A lipid based antigen delivery system efficiently facilitates MHC class-I Antigen Presentation in Dendritic Cells to Stimulate CD8+ T cells, *Scientific Reports* 6 (2016): 27206.
- [0590] Moon et al., Interlayer-crosslinked multilamellar vesicles as synthetic vaccines for potent humoral and cellular responses, *Nature Materials* 10 (2011): 243-251.
- [0591] May et al., "In Vitro Expressed GPCR Inserted in Polymersome Membranes for Ligand-Binding Studies" (2013) *Angew. Chem. Int. Ed.*, 52, pages 749-753.
- Kuai et al., Designer vaccine nanodiscs for personalized cancer immunotherapy, *Nat Mater.* (2017) 16 (4): 489-496.
- [0592] Luo et al., A STING-activating nanovaccine for cancer immunotherapy, *Nat Nanotechnol.* (2017) 12 (7): 648-654.
- [0593] Quer et al., "Polymersomes enhance the immunogenicity of influenza subunit vaccine", *POLYMER CHEMISTRY*, GB, (2012), vol. 2, no. 7, page 1482.
- [0594] Nallani et al., Method for eliciting an immune response to an immunogen, WO2014/077781A1.
- [0595] Nallani et al., Proteopolymersomes: In vitro production of a membrane protein in polymersome membranes (2011), *Biointerphases*, 6 (4), pages 153-157.
- [0596] Neil et al., A Novel Method for the Encapsulation of Biomolecules into Polymersomes via Direct Hydration, *Langmuir* (2009), 25 (16), 9025-9029.
- [0597] Rameez et al. Large Scale Production of Vesicles by Hollow Fiber Extrusion: A Novel Method for Generating Polymersome Encapsulated Hemoglobin Dispersions *Langmuir* (2010) 26 (7), pp 5279-5285.
- Stano et al., Tunable T cell immunity towards a protein antigen using polymersomes vs. solid-core nanoparticles, *Biomaterials* 34 (2013): 4339-4346.
- [0598] Fu et al., Multicompartmentalized polymersomes for selective encapsulation of biomacromolecules, *Chemical Communication*, 2011, 47, 2862-2864.
- [0599] Lim. S. K., et al., Spontaneous formation of nanometer scale tubular vesicles in aqueous mixtures of lipid and block copolymer amphiphiles, *Soft Matter* 2017, 1107-1115.
- [0600] Scott et al., Dendritic cell activation and T cell priming with adjuvant- and antigen-loaded oxidation-sensitive polymersomes, *Biomaterials* 33 (2012) 6211e6

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 16

<210> SEQ ID NO 1

<211> LENGTH: 28

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Reps1 P45A peptide

<400> SEQUENCE: 1

Gly Arg Val Leu Glu Leu Phe Arg Ala Ala Gln Leu Ala Asn Asp Val
1 5 10 15

Val Leu Gln Ile Met Glu Leu Cys Gly Ala Thr Arg
20 25

<210> SEQ ID NO 2

<211> LENGTH: 28

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Adpgk R304M peptide

<400> SEQUENCE: 2

Gly Ile Pro Val His Leu Glu Leu Ala Ser Met Thr Asn Met Glu Leu
1 5 10 15

-continued

Met Ser Ser Ile Val His Gln Gln Val Phe Pro Thr
20 25

<210> SEQ ID NO 3
 <211> LENGTH: 25
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Dpagt1 V213L peptide

<400> SEQUENCE: 3

Glu Ala Gly Gln Ser Leu Val Ile Ser Ala Ser Ile Ile Val Phe Asn
1 5 10 15

Leu Leu Glu Leu Glu Gly Asp Tyr Arg
20 25

<210> SEQ ID NO 4
 <211> LENGTH: 386
 <212> TYPE: PRT
 <213> ORGANISM: Gallus gallus
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(386)
 <223> OTHER INFORMATION: Ovalbumin, UniProtKB Accession Number P01012

<400> SEQUENCE: 4

Met Gly Ser Ile Gly Ala Ala Ser Met Glu Phe Cys Phe Asp Val Phe
1 5 10 15

Lys Glu Leu Lys Val His His Ala Asn Glu Asn Ile Phe Tyr Cys Pro
20 25 30

Ile Ala Ile Met Ser Ala Leu Ala Met Val Tyr Leu Gly Ala Lys Asp
35 40 45

Ser Thr Arg Thr Gln Ile Asn Lys Val Val Arg Phe Asp Lys Leu Pro
50 55 60

Gly Phe Gly Asp Ser Ile Glu Ala Gln Cys Gly Thr Ser Val Asn Val
65 70 75 80

His Ser Ser Leu Arg Asp Ile Leu Asn Gln Ile Thr Lys Pro Asn Asp
85 90 95

Val Tyr Ser Phe Ser Leu Ala Ser Arg Leu Tyr Ala Glu Glu Arg Tyr
100 105 110

Pro Ile Leu Pro Glu Tyr Leu Gln Cys Val Lys Glu Leu Tyr Arg Gly
115 120 125

Gly Leu Glu Pro Ile Asn Phe Gln Thr Ala Ala Asp Gln Ala Arg Glu
130 135 140

Leu Ile Asn Ser Trp Val Glu Ser Gln Thr Asn Gly Ile Ile Arg Asn
145 150 155 160

Val Leu Gln Pro Ser Ser Val Asp Ser Gln Thr Ala Met Val Leu Val
165 170 175

Asn Ala Ile Val Phe Lys Gly Leu Trp Glu Lys Ala Phe Lys Asp Glu
180 185 190

Asp Thr Gln Ala Met Pro Phe Arg Val Thr Glu Gln Glu Ser Lys Pro
195 200 205

Val Gln Met Met Tyr Gln Ile Gly Leu Phe Arg Val Ala Ser Met Ala
210 215 220

Ser Glu Lys Met Lys Ile Leu Glu Leu Pro Phe Ala Ser Gly Thr Met
225 230 235 240

-continued

Ser Met Leu Val Leu Leu Pro Asp Glu Val Ser Gly Leu Glu Gln Leu
 245 250 255
 Glu Ser Ile Ile Asn Phe Glu Lys Leu Thr Glu Trp Thr Ser Ser Asn
 260 265 270
 Val Met Glu Glu Arg Lys Ile Lys Val Tyr Leu Pro Arg Met Lys Met
 275 280 285
 Glu Glu Lys Tyr Asn Leu Thr Ser Val Leu Met Ala Met Gly Ile Thr
 290 295 300
 Asp Val Phe Ser Ser Ser Ala Asn Leu Ser Gly Ile Ser Ser Ala Glu
 305 310 315 320
 Ser Leu Lys Ile Ser Gln Ala Val His Ala Ala His Ala Glu Ile Asn
 325 330 335
 Glu Ala Gly Arg Glu Val Val Gly Ser Ala Glu Ala Gly Val Asp Ala
 340 345 350
 Ala Ser Val Ser Glu Glu Phe Arg Ala Asp His Pro Phe Leu Phe Cys
 355 360 365
 Ile Lys His Ile Ala Thr Asn Ala Val Leu Phe Phe Gly Arg Cys Val
 370 375 380
 Ser Pro
 385

<210> SEQ ID NO 5
 <211> LENGTH: 566
 <212> TYPE: PRT
 <213> ORGANISM: Influenza A virus
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(566)
 <223> OTHER INFORMATION: Influenza A virus (A/New York/38/2016(H1N1))
 Hemagglutinin, UniProtKB Accession Number: A0A192ZYK0

<400> SEQUENCE: 5

Met Lys Ala Ile Leu Val Val Leu Leu Tyr Thr Phe Thr Thr Ala Asn
 1 5 10 15
 Ala Asp Thr Leu Cys Ile Gly Tyr His Ala Asn Asn Ser Thr Asp Thr
 20 25 30
 Val Asp Thr Val Leu Glu Lys Asn Val Thr Val Thr His Ser Val Asn
 35 40 45
 Leu Leu Glu Asp Lys His Asn Gly Lys Leu Cys Lys Leu Arg Gly Val
 50 55 60
 Ala Pro Leu His Leu Gly Lys Cys Asn Ile Ala Gly Trp Ile Leu Gly
 65 70 75 80
 Asn Pro Glu Cys Glu Ser Leu Ser Thr Ala Ser Ser Trp Ser Tyr Ile
 85 90 95
 Val Glu Thr Ser Asn Ser Asp Asn Gly Thr Cys Tyr Pro Gly Asp Phe
 100 105 110
 Ile Asn Tyr Glu Glu Leu Arg Glu Gln Leu Ser Ser Val Ser Ser Phe
 115 120 125
 Glu Arg Phe Glu Ile Phe Pro Lys Thr Ser Ser Trp Pro Asn His Asp
 130 135 140
 Ser Asn Lys Gly Val Thr Ala Ala Cys Pro His Ala Gly Ala Lys Ser
 145 150 155 160
 Phe Tyr Lys Asn Leu Ile Trp Leu Val Lys Lys Gly Asn Ser Tyr Pro
 165 170 175

-continued

Lys	Leu	Asn	Gln	Ser	Tyr	Ile	Asn	Asp	Lys	Gly	Lys	Glu	Val	Leu	Val	180	185	190
Leu	Trp	Gly	Ile	His	His	Pro	Ser	Thr	Thr	Ala	Asp	Gln	Gln	Ser	Leu	195	200	205
Tyr	Gln	Asn	Ala	Asp	Ala	Tyr	Val	Phe	Val	Gly	Thr	Ser	Arg	Tyr	Ser	210	215	220
Lys	Lys	Phe	Lys	Pro	Glu	Ile	Ala	Thr	Arg	Pro	Lys	Val	Arg	Asp	Gln	225	230	235
Glu	Gly	Arg	Met	Asn	Tyr	Tyr	Trp	Thr	Leu	Val	Glu	Pro	Gly	Asp	Lys	245	250	255
Ile	Thr	Phe	Glu	Ala	Thr	Gly	Asn	Leu	Val	Val	Pro	Arg	Tyr	Ala	Phe	260	265	270
Thr	Met	Glu	Arg	Asn	Ala	Gly	Ser	Gly	Ile	Ile	Ile	Ser	Asp	Thr	Pro	275	280	285
Val	His	Asp	Cys	Asn	Thr	Thr	Cys	Gln	Thr	Pro	Glu	Gly	Ala	Ile	Asn	290	295	300
Thr	Ser	Leu	Pro	Phe	Gln	Asn	Ile	His	Pro	Ile	Thr	Ile	Gly	Lys	Cys	305	310	315
Pro	Lys	Tyr	Val	Lys	Ser	Thr	Lys	Leu	Arg	Leu	Ala	Thr	Gly	Leu	Arg	325	330	335
Asn	Val	Pro	Ser	Ile	Gln	Ser	Arg	Gly	Leu	Phe	Gly	Ala	Ile	Ala	Gly	340	345	350
Phe	Ile	Glu	Gly	Gly	Trp	Thr	Gly	Met	Val	Asp	Gly	Trp	Tyr	Gly	Tyr	355	360	365
His	His	Gln	Asn	Glu	Gln	Gly	Ser	Gly	Tyr	Ala	Ala	Asp	Leu	Lys	Ser	370	375	380
Thr	Gln	Asn	Ala	Ile	Asp	Lys	Ile	Thr	Asn	Lys	Val	Asn	Ser	Val	Ile	385	390	395
Glu	Lys	Met	Asn	Thr	Gln	Phe	Thr	Ala	Val	Gly	Lys	Glu	Phe	Asn	His	405	410	415
Leu	Glu	Lys	Arg	Ile	Glu	Asn	Leu	Asn	Lys	Lys	Val	Asp	Asp	Gly	Phe	420	425	430
Leu	Asp	Ile	Trp	Thr	Tyr	Asn	Ala	Glu	Leu	Leu	Val	Leu	Leu	Glu	Asn	435	440	445
Glu	Arg	Thr	Leu	Asp	Tyr	His	Asp	Ser	Asn	Val	Lys	Asn	Leu	Tyr	Glu	450	455	460
Lys	Val	Arg	Asn	Gln	Leu	Lys	Asn	Asn	Ala	Lys	Glu	Ile	Gly	Asn	Gly	465	470	475
Cys	Phe	Glu	Phe	Tyr	His	Lys	Cys	Asp	Asn	Thr	Cys	Met	Glu	Ser	Val	485	490	495
Lys	Asn	Gly	Thr	Tyr	Asp	Tyr	Pro	Lys	Tyr	Ser	Glu	Glu	Ala	Lys	Leu	500	505	510
Asn	Arg	Glu	Lys	Ile	Asp	Gly	Val	Lys	Leu	Glu	Ser	Thr	Arg	Ile	Tyr	515	520	525
Gln	Ile	Leu	Ala	Ile	Tyr	Ser	Thr	Val	Ala	Ser	Ser	Leu	Val	Leu	Val	530	535	540
Val	Ser	Leu	Gly	Ala	Ile	Ser	Phe	Trp	Met	Cys	Ser	Asn	Gly	Ser	Leu	545	550	555
Gln	Cys	Arg	Ile	Cys	Ile											565		

-continued

<210> SEQ ID NO 6
<211> LENGTH: 566
<212> TYPE: PRT
<213> ORGANISM: Influenza A virus
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(566)
<223> OTHER INFORMATION: Influenza A virus (A/swine/4/Mexico/2009(H1N1))
Hemagglutinin, UniProtKB Accession Number: D2CE65

<400> SEQUENCE: 6

Met Lys Ala Ile Leu Val Val Leu Leu Tyr Thr Phe Ala Thr Ala Asn
1 5 10 15
Ala Asp Thr Leu Cys Ile Gly Tyr His Ala Asn Asn Ser Thr Asp Thr
20 25 30
Val Asp Thr Val Leu Glu Lys Asn Val Thr Val Thr His Ser Val Asn
35 40 45
Leu Leu Glu Asp Lys His Asn Gly Lys Leu Cys Lys Leu Arg Gly Val
50 55 60
Ala Pro Leu His Leu Gly Lys Cys Asn Ile Ala Gly Trp Ile Leu Gly
65 70 75 80
Asn Pro Glu Cys Glu Ser Leu Ser Thr Ala Ser Ser Trp Ser Tyr Ile
85 90 95
Val Glu Thr Ser Ser Ser Asp Asn Gly Thr Cys Tyr Pro Gly Asp Phe
100 105 110
Ile Asp Tyr Glu Glu Leu Arg Glu Gln Leu Ser Ser Val Ser Ser Phe
115 120 125
Glu Arg Phe Glu Ile Phe Pro Lys Thr Ser Ser Trp Pro Asn His Asp
130 135 140
Ser Asn Lys Gly Val Thr Ala Ala Cys Pro His Ala Gly Ala Lys Ser
145 150 155 160
Phe Tyr Lys Asn Leu Ile Trp Leu Val Lys Lys Gly Asn Ser Tyr Pro
165 170 175
Lys Leu Ser Lys Ser Tyr Ile Asn Asp Lys Gly Lys Glu Val Leu Val
180 185 190
Leu Trp Gly Ile His His Pro Pro Thr Ser Ala Asp Gln Gln Ser Leu
195 200 205
Tyr Gln Asn Ala Asp Thr Tyr Val Phe Val Gly Ser Ser Arg Tyr Ser
210 215 220
Lys Lys Phe Lys Pro Glu Ile Ala Ile Arg Pro Lys Val Arg Xaa Xaa
225 230 235 240
Glu Gly Arg Met Asn Tyr Tyr Trp Thr Leu Val Glu Pro Gly Asp Lys
245 250 255
Ile Thr Phe Glu Ala Thr Gly Asn Leu Val Val Pro Arg Tyr Ala Phe
260 265 270
Ala Met Glu Arg Asn Ala Gly Ser Gly Ile Ile Ile Ser Asp Thr Pro
275 280 285
Val His Asp Cys Asn Thr Thr Cys Gln Thr Pro Lys Gly Ala Ile Asn
290 295 300
Thr Ser Leu Pro Phe Gln Asn Ile His Pro Ile Thr Ile Gly Lys Cys
305 310 315 320
Pro Lys Tyr Val Lys Ser Thr Lys Leu Arg Leu Ala Thr Gly Leu Arg
325 330 335
Asn Val Pro Ser Ile Gln Ser Arg Gly Leu Phe Gly Ala Ile Ala Gly

-continued

340					345					350					
Phe	Ile	Glu	Gly	Gly	Trp	Thr	Gly	Met	Val	Asp	Gly	Trp	Tyr	Gly	Tyr
		355					360					365			
His	His	Gln	Asn	Glu	Gln	Gly	Ser	Gly	Tyr	Ala	Ala	Asp	Leu	Lys	Ser
		370				375						380			
Thr	Gln	Asn	Ala	Ile	Asp	Glu	Ile	Thr	Asn	Lys	Val	Asn	Ser	Val	Ile
385					390					395					400
Glu	Lys	Met	Asn	Thr	Gln	Phe	Thr	Ala	Val	Gly	Lys	Glu	Phe	Asn	His
			405						410					415	
Leu	Glu	Lys	Arg	Ile	Glu	Asn	Leu	Asn	Lys	Lys	Val	Asp	Asp	Gly	Phe
			420					425					430		
Leu	Asp	Ile	Trp	Thr	Tyr	Asn	Ala	Glu	Leu	Leu	Val	Leu	Leu	Glu	Asn
	435						440						445		
Glu	Arg	Thr	Leu	Asp	Tyr	His	Asp	Ser	Asn	Val	Lys	Asn	Leu	Tyr	Glu
	450					455					460				
Lys	Val	Arg	Ser	Gln	Leu	Lys	Asn	Asn	Ala	Lys	Glu	Ile	Gly	Asn	Gly
465					470					475					480
Cys	Phe	Glu	Phe	Tyr	His	Lys	Cys	Asp	Asn	Thr	Cys	Met	Glu	Ser	Val
			485						490					495	
Lys	Asn	Gly	Thr	Tyr	Asp	Tyr	Pro	Lys	Tyr	Ser	Glu	Glu	Ala	Lys	Leu
			500					505					510		
Asn	Arg	Glu	Glu	Ile	Asp	Gly	Val	Lys	Leu	Glu	Ser	Thr	Arg	Ile	Tyr
		515					520						525		
Gln	Ile	Leu	Ala	Ile	Tyr	Ser	Thr	Val	Ala	Ser	Ser	Leu	Val	Leu	Val
	530					535						540			
Val	Ser	Leu	Gly	Ala	Ile	Ser	Phe	Trp	Met	Cys	Ser	Asn	Gly	Ser	Leu
545					550					555					560
Gln	Cys	Arg	Ile	Cys	Ile										
				565											

<210> SEQ ID NO 7

<211> LENGTH: 565

<212> TYPE: PRT

<213> ORGANISM: Influenza A virus

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (1)..(565)

<223> OTHER INFORMATION: Influenza A virus (H1N1/A/Puerto rico/8/1934)
Hemagglutinin

<400> SEQUENCE: 7

Met	Lys	Ala	Asn	Leu	Leu	Val	Leu	Leu	Cys	Ala	Leu	Ala	Ala	Ala	Asp
1				5					10					15	
Ala	Asp	Thr	Ile	Cys	Ile	Gly	Tyr	His	Ala	Asn	Asn	Ser	Thr	Asp	Thr
			20					25					30		
Val	Asp	Thr	Val	Leu	Glu	Lys	Asn	Val	Thr	Val	Thr	His	Ser	Val	Asn
		35					40					45			
Leu	Leu	Glu	Asp	Ser	His	Asn	Gly	Lys	Leu	Cys	Arg	Leu	Lys	Gly	Ile
	50					55					60				
Ala	Pro	Leu	Gln	Leu	Gly	Lys	Cys	Asn	Ile	Ala	Gly	Trp	Leu	Leu	Gly
65				70						75					80
Asn	Pro	Glu	Cys	Asp	Pro	Leu	Leu	Pro	Val	Arg	Ser	Trp	Ser	Tyr	Ile
			85					90						95	
Val	Glu	Thr	Pro	Asn	Ser	Glu	Asn	Gly	Ile	Cys	Tyr	Pro	Gly	Asp	Phe

-continued

100							105					110				
Ile	Asp	Tyr	Glu	Glu	Leu	Arg	Glu	Gln	Leu	Ser	Ser	Val	Ser	Ser	Phe	
115							120					125				
Glu	Arg	Phe	Glu	Ile	Phe	Pro	Lys	Glu	Ser	Ser	Trp	Pro	Asn	His	Asn	
130							135					140				
Thr	Asn	Gly	Val	Thr	Ala	Ala	Cys	Ser	His	Glu	Gly	Lys	Ser	Ser	Phe	
145							150					155				
Tyr	Arg	Asn	Leu	Leu	Trp	Leu	Thr	Glu	Lys	Glu	Gly	Ser	Tyr	Pro	Lys	
165							170					175				
Leu	Lys	Asn	Ser	Tyr	Val	Asn	Lys	Lys	Gly	Lys	Glu	Val	Leu	Val	Leu	
180							185					190				
Trp	Gly	Ile	His	His	Pro	Pro	Asn	Ser	Lys	Glu	Gln	Gln	Asn	Leu	Tyr	
195							200					205				
Gln	Asn	Glu	Asn	Ala	Tyr	Val	Ser	Val	Val	Thr	Ser	Asn	Tyr	Asn	Arg	
210							215					220				
Arg	Phe	Thr	Pro	Glu	Ile	Ala	Glu	Arg	Pro	Lys	Val	Arg	Asp	Gln	Ala	
225							230					235				
Gly	Arg	Met	Asn	Tyr	Tyr	Trp	Thr	Leu	Leu	Lys	Pro	Gly	Asp	Thr	Ile	
245							250					255				
Ile	Phe	Glu	Ala	Asn	Gly	Asn	Leu	Ile	Ala	Pro	Met	Tyr	Ala	Phe	Ala	
260							265					270				
Leu	Ser	Arg	Gly	Phe	Gly	Ser	Gly	Ile	Ile	Thr	Ser	Asn	Ala	Ser	Met	
275							280					285				
His	Glu	Cys	Asn	Thr	Lys	Cys	Gln	Thr	Pro	Leu	Gly	Ala	Ile	Asn	Ser	
290							295					300				
Ser	Leu	Pro	Tyr	Gln	Asn	Ile	His	Pro	Val	Thr	Ile	Gly	Glu	Cys	Pro	
305							310					315				
Lys	Tyr	Val	Arg	Ser	Ala	Lys	Leu	Arg	Met	Val	Thr	Gly	Leu	Arg	Asn	
325							330					335				
Ile	Pro	Ser	Ile	Gln	Ser	Arg	Gly	Leu	Phe	Gly	Ala	Ile	Ala	Gly	Phe	
340							345					350				
Ile	Glu	Gly	Gly	Trp	Thr	Gly	Met	Ile	Asp	Gly	Trp	Tyr	Gly	Tyr	His	
355							360					365				
His	Gln	Asn	Glu	Gln	Gly	Ser	Gly	Tyr	Ala	Ala	Asp	Gln	Lys	Ser	Thr	
370							375					380				
Gln	Asn	Ala	Ile	Asn	Gly	Ile	Thr	Asn	Lys	Val	Asn	Thr	Val	Ile	Glu	
385							390					395				
Lys	Met	Asn	Ile	Gln	Phe	Thr	Ala	Val	Gly	Lys	Glu	Phe	Asn	Lys	Leu	
405							410					415				
Glu	Lys	Arg	Met	Glu	Asn	Leu	Asn	Lys	Lys	Val	Asp	Asp	Gly	Phe	Leu	
420							425					430				
Asp	Ile	Trp	Thr	Tyr	Asn	Ala	Glu	Leu	Leu	Val	Leu	Leu	Glu	Asn	Glu	
435							440					445				
Arg	Thr	Leu	Asp	Phe	His	Asp	Ser	Asn	Val	Lys	Asn	Leu	Tyr	Glu	Lys	
450							455					460				
Val	Lys	Ser	Gln	Leu	Lys	Asn	Asn	Ala	Lys	Glu	Ile	Gly	Asn	Gly	Cys	
465							470					475				
Phe	Glu	Phe	Tyr	His	Lys	Cys	Asp	Asn	Glu	Cys	Met	Glu	Ser	Val	Arg	
485							490					495				
Asn	Gly	Thr	Tyr	Asp	Tyr	Pro	Lys	Tyr	Ser	Glu	Glu	Ser	Lys	Leu	Asn	
500							505					510				

-continued

Arg Glu Lys Val Asp Gly Val Lys Leu Glu Ser Met Gly Ile Tyr Gln
515 520 525

Ile Leu Ala Ile Tyr Ser Thr Val Ala Ser Ser Leu Val Leu Leu Val
530 535 540

Ser Leu Gly Ala Ile Ser Phe Trp Met Cys Ser Asn Gly Ser Leu Gln
545 550 555 560

Cys Arg Ile Cys Ile
565

<210> SEQ ID NO 8
<211> LENGTH: 566
<212> TYPE: PRT
<213> ORGANISM: Influenza A virus
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(566)
<223> OTHER INFORMATION: Influenza A virus (A/California/07/2009(H1N1))
Hemagglutinin

<400> SEQUENCE: 8

Met Lys Ala Ile Leu Val Val Leu Leu Tyr Thr Phe Ala Thr Ala Asn
1 5 10 15

Ala Asp Thr Leu Cys Ile Gly Tyr His Ala Asn Asn Ser Thr Asp Thr
20 25 30

Val Asp Thr Val Leu Glu Lys Asn Val Thr Val Thr His Ser Val Asn
35 40 45

Leu Leu Glu Asp Lys His Asn Gly Lys Leu Cys Lys Leu Arg Gly Val
50 55 60

Ala Pro Leu His Leu Gly Lys Cys Asn Ile Ala Gly Trp Ile Leu Gly
65 70 75 80

Asn Pro Glu Cys Glu Ser Leu Ser Thr Ala Ser Ser Trp Ser Tyr Ile
85 90 95

Val Glu Thr Pro Ser Ser Asp Asn Gly Thr Cys Tyr Pro Gly Asp Phe
100 105 110

Ile Asp Tyr Glu Glu Leu Arg Glu Gln Leu Ser Ser Val Ser Ser Phe
115 120 125

Glu Arg Phe Glu Ile Phe Pro Lys Thr Ser Ser Trp Pro Asn His Asp
130 135 140

Ser Asn Lys Gly Val Thr Ala Ala Cys Pro His Ala Gly Ala Lys Ser
145 150 155 160

Phe Tyr Lys Asn Leu Ile Trp Leu Val Lys Lys Gly Asn Ser Tyr Pro
165 170 175

Lys Leu Ser Lys Ser Tyr Ile Asn Asp Lys Gly Lys Glu Val Leu Val
180 185 190

Leu Trp Gly Ile His His Pro Ser Thr Ser Ala Asp Gln Gln Ser Leu
195 200 205

Tyr Gln Asn Ala Asp Ala Tyr Val Phe Val Gly Ser Ser Arg Tyr Ser
210 215 220

Lys Lys Phe Lys Pro Glu Ile Ala Ile Arg Pro Lys Val Arg Asp Gln
225 230 235 240

Glu Gly Arg Met Asn Tyr Tyr Trp Thr Leu Val Glu Pro Gly Asp Lys
245 250 255

Ile Thr Phe Glu Ala Thr Gly Asn Leu Val Val Pro Arg Tyr Ala Phe
260 265 270

-continued

Ala Met	Glu Arg	Asn Ala	Gly Ser	Gly Ile	Ile Ile	Ile Ser	Asp Thr	Pro	
	275		280			285			
Val His	Asp Cys	Asn Thr	Thr Cys	Gln Thr	Pro Lys	Gly Ala	Ile Asn		
	290		295		300				
Thr Ser	Leu Pro	Phe Gln	Asn Ile	His Pro	Ile Thr	Ile Gly	Lys Cys		
305		310			315		320		
Pro Lys	Tyr Val	Lys Ser	Thr Lys	Leu Arg	Leu Ala	Thr Gly	Leu Arg		
		325		330			335		
Asn Ile	Pro Ser	Ile Gln	Ser Arg	Gly Leu	Phe Gly	Ala Ile	Ala Gly		
	340			345		350			
Phe Ile	Glu Gly	Gly Trp	Thr Gly	Met Val	Asp Gly	Trp Tyr	Gly Tyr		
	355		360		365				
His His	Gln Asn	Glu Gln	Gly Ser	Gly Tyr	Ala Ala	Asp Leu	Lys Ser		
	370		375		380				
Thr Gln	Asn Ala	Ile Asp	Glu Ile	Thr Asn	Lys Val	Asn Ser	Val Ile		
385		390			395		400		
Glu Lys	Met Asn	Thr Gln	Phe Thr	Ala Val	Gly Lys	Glu Phe	Asn His		
	405			410			415		
Leu Glu	Lys Arg	Ile Glu	Asn Leu	Asn Lys	Lys Val	Asp Asp	Gly Phe		
	420		425			430			
Leu Asp	Ile Trp	Thr Tyr	Asn Ala	Glu Leu	Leu Val	Leu Leu	Glu Asn		
	435		440		445				
Glu Arg	Thr Leu	Asp Tyr	His Asp	Ser Asn	Val Lys	Asn Leu	Tyr Glu		
	450		455		460				
Lys Val	Arg Ser	Gln Leu	Lys Asn	Asn Ala	Lys Glu	Ile Gly	Asn Gly		
465		470			475		480		
Cys Phe	Glu Phe	Tyr His	Lys Cys	Asp Asn	Thr Cys	Met Glu	Ser Val		
	485			490			495		
Lys Asn	Gly Thr	Tyr Asp	Tyr Pro	Lys Tyr	Ser Glu	Glu Ala	Lys Leu		
	500		505			510			
Asn Arg	Glu Glu	Ile Asp	Gly Val	Lys Leu	Glu Ser	Thr Arg	Ile Tyr		
	515		520			525			
Gln Ile	Leu Ala	Ile Tyr	Ser Thr	Val Ala	Ser Ser	Leu Val	Leu Val		
	530		535		540				
Val Ser	Leu Gly	Ala Ile	Ser Phe	Trp Met	Cys Ser	Asn Gly	Ser Leu		
545		550		555			560		
Gln Cys	Arg Ile	Cys Ile							
		565							

<210> SEQ ID NO 9

<211> LENGTH: 24

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: CD8 Trp2 173-196 peptide

<400> SEQUENCE: 9

Gln Pro	Gln Ile	Ala Asn	Cys Ser	Val Tyr	Asp Phe	Phe Val	Trp Leu		
1		5		10			15		

His Tyr	Tyr Ser	Val Arg	Asp Thr						
		20							

<210> SEQ ID NO 10

-continued

```

<211> LENGTH: 27
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: CD4 M30 Kif18b K739N peptide

```

```

<400> SEQUENCE: 10

```

```

Pro Ser Lys Pro Ser Phe Gln Glu Phe Val Asp Trp Glu Asn Val Ser
1           5           10           15

```

```

Pro Glu Leu Asn Ser Thr Asp Gln Pro Phe Leu
           20           25

```

```

<210> SEQ ID NO 11
<211> LENGTH: 27
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: CD4 M44 Cpsf31 D314N peptide

```

```

<400> SEQUENCE: 11

```

```

Glu Phe Lys His Ile Lys Ala Phe Asp Arg Thr Phe Ala Asn Asn Pro
1           5           10           15

```

```

Gly Pro Met Val Val Phe Ala Thr Pro Gly Met
           20           25

```

```

<210> SEQ ID NO 12
<211> LENGTH: 1309
<212> TYPE: PRT
<213> ORGANISM: Porcine epidemic diarrhea virus

```

```

<400> SEQUENCE: 12

```

```

Leu Pro Gln Asp Val Thr Arg Cys Ser Ala Asn Thr Asn Phe Arg Arg
1           5           10           15

```

```

Phe Phe Ser Lys Phe Asn Val Gln Ala Pro Ala Val Val Val Leu Gly
           20           25           30

```

```

Gly Tyr Leu Pro Ile Gly Glu Asn Gln Gly Val Asn Ser Thr Trp Tyr
           35           40           45

```

```

Cys Ala Gly Gln His Pro Thr Ala Ser Gly Val His Gly Ile Phe Leu
           50           55           60

```

```

Ser His Ile Arg Gly Gly His Gly Phe Glu Ile Gly Ile Ser Gln Glu
           65           70           75           80

```

```

Pro Phe Asp Pro Ser Gly Tyr Gln Leu Tyr Leu His Lys Ala Thr Asn
           85           90           95

```

```

Gly Asn Thr Asn Ala Thr Ala Arg Leu Arg Ile Cys Gln Phe Pro Ser
           100          105          110

```

```

Ile Lys Thr Leu Gly Pro Thr Ala Asn Asn Asp Val Thr Thr Gly Arg
           115          120          125

```

```

Asn Cys Leu Phe Asn Lys Ala Ile Pro Ala His Met Ser Glu His Ser
           130          135          140

```

```

Val Val Gly Ile Thr Trp Asp Asn Asp Arg Val Thr Val Phe Ser Asp
           145          150          155          160

```

```

Lys Ile Tyr Tyr Phe Tyr Phe Lys Asn Asp Trp Ser Arg Val Ala Thr
           165          170          175

```

```

Lys Cys Tyr Asn Ser Gly Gly Cys Ala Met Gln Tyr Val Tyr Glu Pro
           180          185          190

```

```

Thr Tyr Tyr Met Leu Asn Val Thr Ser Ala Gly Glu Asp Gly Ile Ser
           195          200          205

```

-continued

Tyr	Gln	Pro	Cys	Thr	Ala	Asn	Cys	Ile	Gly	Tyr	Ala	Ala	Asn	Val	Phe
210						215					220				
Ala	Thr	Glu	Pro	Asn	Gly	His	Ile	Pro	Glu	Gly	Phe	Ser	Phe	Asn	Asn
225					230					235					240
Trp	Phe	Leu	Leu	Ser	Asn	Asp	Ser	Thr	Leu	Val	His	Gly	Lys	Val	Val
				245					250					255	
Ser	Asn	Gln	Pro	Leu	Leu	Val	Asn	Cys	Leu	Leu	Ala	Ile	Pro	Lys	Ile
			260					265					270		
Tyr	Gly	Leu	Gly	Gln	Phe	Phe	Ser	Phe	Asn	Gln	Thr	Ile	Asp	Gly	Val
	275						280					285			
Cys	Asn	Gly	Ala	Ala	Val	Gln	Arg	Ala	Pro	Glu	Ala	Leu	Arg	Phe	Asn
290						295					300				
Val	Asp	Asp	Thr	Ser	Val	Ile	Leu	Ala	Glu	Gly	Ser	Ile	Val	Leu	His
305					310					315					320
Thr	Ala	Leu	Gly	Thr	Asn	Phe	Ser	Phe	Val	Cys	Ser	Asn	Ser	Ser	Asp
				325					330					335	
Pro	His	Leu	Ala	Thr	Phe	Ala	Ile	Pro	Leu	Gly	Ala	Thr	Gln	Val	Pro
			340					345					350		
Tyr	Tyr	Cys	Phe	Leu	Lys	Val	Asp	Thr	Tyr	Asn	Ser	Thr	Val	Tyr	Lys
		355					360					365			
Phe	Leu	Ala	Val	Leu	His	Pro	Thr	Val	Arg	Glu	Ile	Val	Ile	Thr	Lys
370						375				380					
Tyr	Gly	Asp	Val	Tyr	Val	Asn	Gly	Phe	Gly	Tyr	Leu	His	Leu	Gly	Leu
385					390					395					400
Leu	Asp	Ala	Val	Thr	Ile	Asn	Phe	Thr	Gly	His	Gly	Thr	Asp	Asp	Asp
				405					410					415	
Val	Pro	Gly	Phe	Trp	Thr	Ile	Ala	Ser	Thr	Asn	Phe	Val	Asp	Ala	Leu
			420					425					430		
Ile	Glu	Val	Gln	Gly	Thr	Ala	Ile	Gln	Arg	Ile	Leu	Tyr	Cys	Asp	Asp
	435					440						445			
Pro	Val	Ser	Gln	Leu	Lys	Cys	Ser	Gln	Val	Ala	Phe	Asp	Leu	Asp	Asp
	450					455					460				
Gly	Phe	Tyr	Pro	Ile	Ser	Ser	Arg	Asn	Leu	Leu	Ser	His	Glu	Gln	Pro
465				470						475					480
Ile	Ser	Phe	Val	Thr	Leu	Pro	Ser	Phe	Asn	Asp	His	Ser	Phe	Val	Asn
				485					490					495	
Ile	Thr	Val	Ser	Ala	Ser	Phe	Gly	Gly	His	Ser	Gly	Ala	Asn	Leu	Ile
			500					505					510		
Ala	Ser	Asp	Thr	Thr	Ile	Asn	Gly	Phe	Ser	Ser	Phe	Cys	Val	Asp	Thr
		515					520					525			
Arg	Gln	Phe	Thr	Ile	Ser	Leu	Phe	Tyr	Asn	Val	Thr	Asn	Ser	Tyr	Gly
	530					535					540				
Tyr	Val	Ser	Lys	Ser	Gln	Asp	Ser	Asn	Cys	Pro	Phe	Thr	Leu	Gln	Ser
545					550					555					560
Val	Asn	Asp	Tyr	Leu	Ser	Phe	Ser	Lys	Phe	Cys	Val	Ser	Thr	Ser	Leu
				565					570					575	
Leu	Ala	Ser	Ala	Cys	Thr	Ile	Asp	Leu	Phe	Gly	Tyr	Pro	Glu	Phe	Gly
			580					585					590		
Ser	Gly	Val	Lys	Phe	Thr	Ser	Leu	Tyr	Phe	Gln	Phe	Thr	Lys	Gly	Glu
	595						600					605			

-continued

Leu	Ile	Thr	Gly	Thr	Pro	Lys	Pro	Leu	Glu	Gly	Val	Thr	Asp	Val	Ser
610						615					620				
Phe	Met	Thr	Leu	Asp	Val	Cys	Thr	Lys	Tyr	Thr	Ile	Tyr	Gly	Phe	Lys
625					630					635					640
Gly	Glu	Gly	Ile	Ile	Thr	Leu	Thr	Asn	Ser	Ser	Phe	Leu	Ala	Gly	Val
			645						650					655	
Tyr	Tyr	Thr	Ser	Asp	Ser	Gly	Gln	Leu	Leu	Ala	Phe	Lys	Asn	Val	Thr
			660					665					670		
Ser	Gly	Ala	Val	Tyr	Ser	Val	Thr	Pro	Cys	Ser	Phe	Ser	Glu	Gln	Ala
		675						680				685			
Ala	Tyr	Val	Asp	Asp	Asp	Ile	Val	Gly	Val	Ile	Ser	Ser	Leu	Ser	Asn
	690					695					700				
Ser	Thr	Phe	Asn	Ser	Thr	Arg	Glu	Leu	Pro	Gly	Phe	Phe	Tyr	His	Ser
705					710					715					720
Asn	Asp	Gly	Ser	Asn	Cys	Thr	Glu	Pro	Val	Leu	Val	Tyr	Ser	Asn	Ile
				725					730					735	
Gly	Val	Cys	Lys	Ser	Gly	Ser	Ile	Gly	Tyr	Val	Pro	Ser	Gln	Ser	Gly
			740					745					750		
Gln	Val	Lys	Ile	Ala	Pro	Thr	Val	Ile	Gly	Asn	Ile	Ser	Ile	Pro	Thr
		755					760					765			
Asn	Phe	Ser	Met	Ser	Ile	Arg	Thr	Glu	Tyr	Leu	Gln	Leu	Tyr	Asn	Thr
	770					775					780				
Pro	Val	Ser	Val	Asp	Cys	Ala	Thr	Tyr	Val	Cys	Asn	Gly	Asn	Ser	Arg
785					790					795					800
Cys	Lys	Gln	Leu	Leu	Thr	Gln	Tyr	Thr	Ala	Ala	Cys	Lys	Thr	Ile	Glu
				805					810					815	
Ser	Ala	Leu	Gln	Leu	Ser	Ala	Arg	Leu	Glu	Ser	Val	Glu	Val	Asn	Ser
				820				825					830		
Met	Leu	Thr	Ile	Ser	Glu	Glu	Ala	Leu	Gln	Leu	Ala	Thr	Ile	Ser	Ser
		835					840					845			
Phe	Asn	Gly	Asp	Gly	Tyr	Asn	Phe	Thr	Asn	Val	Leu	Gly	Val	Ser	Val
	850					855					860				
Tyr	Asp	Pro	Ala	Ser	Gly	Arg	Val	Val	Gln	Lys	Arg	Ser	Phe	Ile	Glu
865					870					875					880
Asp	Leu	Leu	Phe	Asn	Lys	Val	Val	Thr	Asn	Gly	Leu	Gly	Thr	Val	Asp
				885					890					895	
Glu	Asp	Tyr	Lys	Arg	Cys	Ser	Asn	Gly	Arg	Ala	Val	Ala	Asp	Leu	Val
			900					905					910		
Cys	Ala	Gln	Tyr	Tyr	Ser	Gly	Val	Met	Val	Leu	Pro	Gly	Val	Val	Asp
		915					920					925			
Ala	Glu	Lys	Leu	His	Met	Tyr	Ser	Ala	Ser	Leu	Ile	Gly	Gly	Met	Val
	930					935					940				
Phe	Gly	Gly	Phe	Thr	Ala	Ala	Ala	Ala	Leu	Pro	Phe	Ser	Tyr	Ala	Val
945					950					955					960
Gln	Ala	Arg	Leu	Asn	Tyr	Leu	Ala	Leu	Gln	Thr	Asp	Val	Leu	Gln	Arg
				965					970					975	
Asn	Gln	Gln	Leu	Leu	Ala	Glu	Ser	Phe	Asn	Ser	Ala	Ile	Gly	Asn	Ile
			980					985					990		
Thr	Ser	Ala	Phe	Glu	Ser	Val	Lys	Glu	Ala	Ile	Ser	Gln	Thr	Ser	Lys
		995					1000					1005			
Gly	Leu	Asn	Thr	Val	Ala	His	Ala	Leu	Thr	Lys	Val	Gln	Glu	Phe	

Pro

<400> SEQUENCE: 13

Leu Pro Gln Asp Val Thr Arg Cys Ser Ala Asn Thr Asn Phe Arg Arg
1 5 10 15

Phe Phe Ser Lys Phe Asn Val Gln Ala Pro Ala Val Val Val Leu Gly
20 25 30

Gly Tyr Leu Pro Ile Gly Glu Asn Gln Gly Val Asn Ser Thr Trp Tyr

-continued

35	40	45
Cys Ala Gly Gln His Pro Thr Ala Ser Gly Val His Gly Ile Phe Leu		
50	55	60
Ser His Ile Arg Gly Gly His Gly Phe Glu Ile Gly Ile Ser Gln Glu		
65	70	75
Pro Phe Asp Pro Ser Gly Tyr Gln Leu Tyr Leu His Lys Ala Thr Asn		
	85	90
Gly Asn Thr Asn Ala Thr Ala Arg Leu Arg Ile Cys Gln Phe Pro Ser		
	100	105
Ile Lys Thr Leu Gly Pro Thr Ala Asn Asn Asp Val Thr Thr Gly Arg		
	115	120
Asn Cys Leu Phe Asn Lys Ala Ile Pro Ala His Met Ser Glu His Ser		
	130	135
Val Val Gly Ile Thr Trp Asp Asn Asp Arg Val Thr Val Phe Ser Asp		
145	150	155
Lys Ile Tyr Tyr Phe Tyr Phe Lys Asn Asp Trp Ser Arg Val Ala Thr		
	165	170
Lys Cys Tyr Asn Ser Gly Gly Cys Ala Met Gln Tyr Val Tyr Glu Pro		
	180	185
Thr Tyr Tyr Met Leu Asn Val Thr Ser Ala Gly Glu Asp Gly Ile Ser		
	195	200
Tyr Gln Pro Cys Thr Ala Asn Cys Ile Gly Tyr Ala Ala Asn Val Phe		
	210	215
Ala Thr Glu Pro Asn Gly His Ile Pro Glu Gly Phe Ser Phe Asn Asn		
225	230	235
Trp Phe Leu Leu Ser Asn Asp Ser Thr Leu Val His Gly Lys Val Val		
	245	250
Ser Asn Gln Pro Leu Leu Val Asn Cys Leu Leu Ala Ile Pro Lys Ile		
	260	265
Tyr Gly Leu Gly Gln Phe Phe Ser Phe Asn Gln Thr Ile Asp Gly Val		
	275	280
Cys Asn Gly Ala Ala Val Gln Arg Ala Pro Glu Ala Leu Arg Phe Asn		
	290	295
Val Asp Asp Thr Ser Val Ile Leu Ala Glu Gly Ser Ile Val Leu His		
305	310	315
Thr Ala Leu Gly Thr Asn Phe Ser Phe Val Cys Ser Asn Ser Ser Asp		
	325	330
Pro His Leu Ala Thr Phe Ala Ile Pro Leu Gly Ala Thr Gln Val Pro		
	340	345
Tyr Tyr Cys Phe Leu Lys Val Asp Thr Tyr Asn Ser Thr Val Tyr Lys		
	355	360
Phe Leu Ala Val Leu His Pro Thr Val Arg Glu Ile Val Ile Thr Lys		
	370	375
Tyr Gly Asp Val Tyr Val Asn Gly Phe Gly Tyr Leu His Leu Gly Leu		
385	390	395
Leu Asp Ala Val Thr Ile Asn Phe Thr Gly His Gly Thr Asp Asp Asp		
	405	410
Val Pro Gly Phe Trp Thr Ile Ala Ser Thr Asn Phe Val Asp Ala Leu		
	420	425
Ile Glu Val Gln Gly Thr Ala Ile Gln Arg Ile Leu Tyr Cys Asp Asp		
	435	440

-continued

```

Pro Val Ser Gln Leu Lys Cys Ser Gln Val Ala Phe Asp Leu Asp Asp
 450                      455                      460

Gly Phe Tyr Pro Ile Ser Ser Arg Asn Leu Leu Ser His Glu Gln Pro
465                      470                      475                      480

Ile Ser Phe Val Thr Leu Pro Ser Phe Asn Asp His Ser Phe Val Asn
                      485                      490                      495

Ile Thr Val Ser Ala Ser Phe Gly Gly His Ser Gly Ala Asn Leu Ile
                      500                      505                      510

Ala Ser Asp Thr Thr Ile Asn Gly Phe Ser Ser Phe Cys Val Asp Thr
 515                      520                      525

Arg Gln Phe Thr Ile Ser Leu Phe Tyr Asn Val Thr Asn Ser Tyr Gly
 530                      535                      540

Tyr Val Ser Lys Ser Gln Asp Ser Asn Cys Pro Phe Thr Leu Gln Ser
545                      550                      555                      560

Val Asn Asp Tyr Leu Ser Phe Ser Lys Phe Cys Val Ser Thr Ser Leu
 565                      570                      575

Leu Ala Ser Ala Cys Thr Ile Asp Leu Phe Gly Tyr Pro Glu Phe Gly
 580                      585                      590

Ser Gly Val Lys Phe Thr Ser Leu Tyr Phe Gln Phe Thr Lys Gly Glu
 595                      600                      605

Leu Ile Thr Gly Thr Pro Lys Pro Leu Glu Gly Val Thr Asp Val Ser
 610                      615                      620

Phe Met Thr Leu Asp Val Cys Thr Lys Tyr Thr Ile Tyr Gly Phe Lys
625                      630                      635                      640

Gly Glu Gly Ile Ile Thr Leu Thr Asn Ser Ser Phe Leu Ala Gly Val
 645                      650                      655

Tyr Tyr Thr Ser Asp Ser Gly Gln Leu Leu Ala Phe Lys Asn Val Thr
 660                      665                      670

Ser Gly Ala Val Tyr Ser Val Thr Pro Cys Ser Phe Ser Glu Gln Ala
 675                      680                      685

Ala Tyr Val Asp Asp Asp Ile Val Gly Val Ile Ser Ser Leu Ser Asn
 690                      695                      700

Ser Thr Phe Asn Ser Thr Arg Glu Leu Pro Gly Phe Phe Tyr His
705                      710                      715

```

<210> SEQ ID NO 14

<211> LENGTH: 590

<212> TYPE: PRT

<213> ORGANISM: Porcine epidemic diarrhea virus

<400> SEQUENCE: 14

```

Ser Asn Asp Gly Ser Asn Cys Thr Glu Pro Val Leu Val Tyr Ser Asn
 1                      5                      10                      15

Ile Gly Val Cys Lys Ser Gly Ser Ile Gly Tyr Val Pro Ser Gln Ser
 20                      25                      30

Gly Gln Val Lys Ile Ala Pro Thr Val Ile Gly Asn Ile Ser Ile Pro
 35                      40                      45

Thr Asn Phe Ser Met Ser Ile Arg Thr Glu Tyr Leu Gln Leu Tyr Asn
 50                      55                      60

Thr Pro Val Ser Val Asp Cys Ala Thr Tyr Val Cys Asn Gly Asn Ser
 65                      70                      75                      80

Arg Cys Lys Gln Leu Leu Thr Gln Tyr Thr Ala Ala Cys Lys Thr Ile

```

-continued

85								90				95				
Glu	Ser	Ala	Leu	Gln	Leu	Ser	Ala	Arg	Leu	Glu	Ser	Val	Glu	Val	Asn	
			100				105						110			
Ser	Met	Leu	Thr	Ile	Ser	Glu	Glu	Ala	Leu	Gln	Leu	Ala	Thr	Ile	Ser	
			115				120						125			
Ser	Phe	Asn	Gly	Asp	Gly	Tyr	Asn	Phe	Thr	Asn	Val	Leu	Gly	Val	Ser	
		130				135						140				
Val	Tyr	Asp	Pro	Ala	Ser	Gly	Arg	Val	Val	Gln	Lys	Arg	Ser	Phe	Ile	
145					150						155			160		
Glu	Asp	Leu	Leu	Phe	Asn	Lys	Val	Val	Thr	Asn	Gly	Leu	Gly	Thr	Val	
			165						170						175	
Asp	Glu	Asp	Tyr	Lys	Arg	Cys	Ser	Asn	Gly	Arg	Ala	Val	Ala	Asp	Leu	
			180						185						190	
Val	Cys	Ala	Gln	Tyr	Tyr	Ser	Gly	Val	Met	Val	Leu	Pro	Gly	Val	Val	
		195						200						205		
Asp	Ala	Glu	Lys	Leu	His	Met	Tyr	Ser	Ala	Ser	Leu	Ile	Gly	Gly	Met	
210					215						220					
Val	Phe	Gly	Gly	Phe	Thr	Ala	Ala	Ala	Ala	Leu	Pro	Phe	Ser	Tyr	Ala	
225					230						235			240		
Val	Gln	Ala	Arg	Leu	Asn	Tyr	Leu	Ala	Leu	Gln	Thr	Asp	Val	Leu	Gln	
			245						250						255	
Arg	Asn	Gln	Gln	Leu	Leu	Ala	Glu	Ser	Phe	Asn	Ser	Ala	Ile	Gly	Asn	
			260						265						270	
Ile	Thr	Ser	Ala	Phe	Glu	Ser	Val	Lys	Glu	Ala	Ile	Ser	Gln	Thr	Ser	
		275						280						285		
Lys	Gly	Leu	Asn	Thr	Val	Ala	His	Ala	Leu	Thr	Lys	Val	Gln	Glu	Phe	
290					295						300					
Val	Asn	Ser	Gln	Gly	Ala	Ala	Leu	Thr	Gln	Leu	Thr	Val	Gln	Pro	Gln	
305					310						315			320		
His	Asn	Phe	Gln	Ala	Ile	Ser	Ser	Ser	Ile	Asp	Asp	Ile	Tyr	Ser	Arg	
			325						330						335	
Leu	Asp	Ile	Leu	Ser	Ala	Asp	Val	Gln	Val	Asp	Arg	Leu	Ile	Thr	Gly	
			340						345						350	
Arg	Leu	Ser	Ala	Leu	Tyr	Cys	Phe	Val	Ala	Gln	Thr	Leu	Thr	Lys	Tyr	
		355						360						365		
Thr	Glu	Val	Gln	Ala	Ser	Arg	Lys	Leu	Ala	Gln	Gln	Lys	Val	Asn	Glu	
370					375						380					
Cys	Val	Lys	Ser	Gln	Ser	Gln	Arg	Tyr	Gly	Phe	Cys	Gly	Gly	Asp	Gly	
385					390						395			400		
Glu	His	Ile	Phe	Ser	Leu	Val	Gln	Ala	Ala	Pro	Gln	Gly	Leu	Leu	Phe	
			405						410						415	
Leu	His	Thr	Val	Leu	Val	Pro	Gly	Asp	Phe	Val	Asn	Val	Ile	Ala	Ile	
			420						425						430	
Ala	Gly	Leu	Cys	Val	Asn	Asp	Glu	Ile	Ala	Leu	Thr	Leu	Arg	Glu	Pro	
		435						440						445		
Gly	Leu	Val	Leu	Phe	Thr	His	Glu	Leu	Gln	Asn	Tyr	Asn	Ala	Thr	Glu	
450					455						460					
Tyr	Phe	Val	Ser	Ser	Arg	Arg	Met	Phe	Glu	Pro	Arg	Lys	Pro	Thr	Ala	
465					470						475			480		
Ser	Asp	Phe	Val	Gln	Ile	Glu	Ser	Cys	Val	Val	Thr	Tyr	Val	Asn	Leu	
			485						490						495	

-continued

Thr Arg Asp Gln Leu Pro Asp Val Ile Pro Asp Tyr Ile Asp Val Asn
500 505 510

Lys Thr Leu Asp Glu Ile Leu Ala Ser Leu Pro Asn Arg Thr Gly Pro
515 520 525

Ser Leu Pro Leu Asp Val Phe Asn Ala Thr Tyr Phe Asn Leu Thr Gly
530 535 540

Glu Met Ala Asp Leu Glu Leu Arg Ser Glu Ser Leu Arg Asn Asn Thr
545 550 555 560

Glu Glu Leu Gln Ser Leu Ile Tyr Asn Ile Asn Asn Thr Leu Val Asp
565 570 575

Leu Glu Trp Leu Asn Arg Val Glu Thr Tyr Ile Lys Trp Pro
580 585 590

<210> SEQ ID NO 15
 <211> LENGTH: 237
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: enhanced GFP

<400> SEQUENCE: 15

Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu Val Glu
1 5 10 15

Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Arg Gly Glu Gly
20 25 30

Glu Gly Asp Ala Thr Asn Gly Lys Leu Thr Leu Lys Phe Ile Cys Thr
35 40 45

Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr Leu Thr
50 55 60

Tyr Gly Val Leu Cys Phe Ser Arg Tyr Pro Asp His Met Lys Arg His
65 70 75 80

Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu Arg Thr
85 90 95

Ile Ser Phe Lys Asp Asp Gly Thr Tyr Lys Thr Arg Ala Glu Val Lys
100 105 110

Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly Ile Asp
115 120 125

Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr Asn Phe
130 135 140

Asn Ser His Asn Val Tyr Ile Thr Ala Asp Lys Gln Lys Asn Gly Ile
145 150 155 160

Lys Ala Tyr Phe Lys Ile Arg His Asn Val Glu Asp Gly Ser Val Gln
165 170 175

Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly Pro Val
180 185 190

Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Val Leu Ser Lys
195 200 205

Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Asp Val Thr
210 215 220

Ala Ala Gly Ile Thr His Gly Met Asp Glu Leu Tyr Lys
225 230 235

-continued

<210> SEQ ID NO 16
 <211> LENGTH: 1161
 <212> TYPE: RNA
 <213> ORGANISM: Gallus gallus

<400> SEQUENCE: 16

```

augggcagca ucggcgccgc cagcauggag uucugcuucg acguguucaa ggagcugaag    60
gugcaccacg ccaacgagaa caucuucac ugccccaucg ccaucaugag cgcccuggcc    120
augguguacc ugggcgccaa ggacagcacc cggacccaga ucaacaaggu ggugcgguuc    180
gacaagcugc cggguucgg cgacagcauc gagggccagu gcggcaccag cgugaacgug    240
cacagcagcc ugccggacau ccugaaccag aucaccaagc ccaacgacgu guacagcuuc    300
agccuggcca gccggcugua cgccgaggag cgguaaccca uccugcccga guaccugcag    360
ugcgugaagg agcuguaccg gggcggccug gagcccauca acuuccagac cgccgcccag    420
caggcccggg agcugaucaa cagcugggug gagagccaga ccaacggcau cauccggaac    480
gugcugcagc ccagcagcgu ggacagccag accgccaugg ugcuuggugaa cgccaucgug    540
uucaagggcc ugugggagaa gaccuucaag gacgaggaca cccaggccau gcccuuccgg    600
gugaccgagc aggagagcaa gcccgugcag augauguacc agaucggccu guuccgggug    660
gccagcaugg ccagcgagaa gaugaagauc cuggagcugc ccuucgccag cggcaccaug    720
agcaugcugg ugcugcugcc cgacgaggug agcggccugg agcagcugga gagcaucauc    780
aacuucgaga agcugaccga guggaccagc agcaacguga uggaggagcg gaagaucaag    840
guguaccugc cccggaugaa gauggaggag aaguacaacc ugaccagcgu gcugauggcc    900
augggcauca ccgacguguu cagcagcagc gccaacuga gcggaucag cagcgccgag    960
agccugaaga ucagccaggc cgugcacgcc gcccacgccg agaucaacga ggccggccgg    1020
gagguggugg gcagcgccga ggcggcgug gacgcccga gcgugagcga ggaguuccgg    1080
gccgaccacc ccuuccuguu cugcaucaag cacaucgcca ccaacgccgu gcuguucuuu    1140
ggccggugcg ugagccccug a                                     1161

```

1. A polymersome comprising a soluble encapsulated antigen, wherein said soluble encapsulated antigen is a polynucleotide selected from a RNA (e.g., mRNA) molecule or a DNA molecule and wherein the polymersome further comprises an ionizable lipid, wherein said ionizable lipid is: (i) a tertiary amine and/or comprising one or more tertiary amine moieties; and/or (ii) comprising a free electron pair at the nitrogen atom.

2. The polymersome according to claim 1, wherein said ionizable lipid is a cationic ionizable lipid.


3. The polymersome according to claim 1, wherein said ionizable lipid:

i) is a tertiary amine, wherein said ionizable lipid is not a quaternary amine, and/or comprising one or more tertiary amine moieties, wherein said ionizable lipid not comprising a cationic quaternized ammonium moiety; and/or


ii) comprising a free electron pair at the nitrogen atom, wherein said nitrogen atom is an ionizable nitrogen atom.

4. The polymersome according to claim 1, wherein said ionizable lipid comprises or consists of:


i) ionizable lipid DLin-MC3-DMA (i.e., (6Z,9Z,28Z,31Z)-Heptatriaconta-6,9,28,31-tetraen-19-yl 4-(dimethylamino) butanoate) having Formula I:




ii) ionizable lipid 319 having Formula II:



iii) ionizable lipid C12-200 having Formula III:




iv) ionizable lipid 5A2-SC8 having Formula IV:




CC(C)CCCCCCCCOC(=O)C1CCN(CC1)CCN(C)CCN2C(=O)C(=O)OCCCC(C)CCCCCCCCCCCCCCCCC(=O)OCCN(CCCCCCCCCCCC)CCCCCCCCCCCCCCCC(=O)OCC12CCCCCCCCC1CCCCCCCC2[illegible]


viii) ionizable lipid A9 having Formula VIII:




ix) ionizable lipid ALC-0315 (i.e., [(4-hydroxybutyl) azanediyl]di(hexane-6,1-diyl)bis(2-hexyldecanoate) having Formula IX:




x) ionizable lipid Arcturus 2.2 (8.8) 4 C CH₃ having Formula X:



xi) ionizable lipid Genevant CL1 having Formula XI:



xii) ionizable lipid ALC-0159 (i.e. 2-[(polyethylene glycol)-2000]-N, N-ditetradecylacetamide) having Formula XII:



xiii) any one of (i)-(xii), wherein one or more alkyl chains thereof are further esterified.

5. The polymersome according to claim 1, wherein said polynucleotide (e.g., SEQ ID NO: 16) encoding one or more polypeptides, preferably said one or more polypeptides are viral polypeptides.

6. The polymersome according to claim 1, wherein said polymersome is capable of increasing thermostability and/or storage stability and/or immunogenicity of said polynucleotide (e.g., mRNA) and/or one or more polypeptides encoded by said polynucleotide (e.g., mRNA) within said polymersome, preferably compared to that of the same polynucleotide (e.g., mRNA) and/or one or more polypeptides encoded by said same polynucleotide (e.g., mRNA) within another polymersome without ionizable lipid or within an ionizable lipid nanoparticle (LNP) comprising cholesterol under the same conditions, further preferably said thermostability and/or storage stability is increased in the temperature range from about -80°C . to about 4°C . (e.g., at -80°C ., -20°C . or at 4°C ., preferably at -80°C .), most preferably said immunogenicity is increased in the temperature range from about 36.5°C . to about 37.5°C .; further most preferably said increase is of about at least 5% (e.g., at least 10%, at least 20% or at least 30%).

7. The polymersome according to claim 1, wherein said polymersome is an oxidation-stable polymersome.

8.-18. (canceled)

19. The polymersome according to claim 1, wherein said polymersome has one or more of the following properties:

- i) said polymersome comprises an oxidation-stable membrane; and/or
- ii) said polymersome is synthetic; and/or
- iii) said polymersome is free from non-encapsulated antigens or in a mixture with non-encapsulated antigens; and/or
- iv) said polymersome comprises a membrane of an amphiphilic polymer; and/or
- v) said polymersome comprises amphiphilic synthetic block copolymers forming a vesicle membrane; and/or
- vi) said polymersome has a diameter greater than 70 nm, preferably said diameter ranging from about 100 nm to about 1 μm , or from about 100 nm to about 750 nm, or from about 100 nm to about 500 nm, or from about 125 nm to about 250 nm, from about 140 nm to about 240 nm, from about 150 nm to about 235 nm, from about 170 nm to about 230 nm, or from about 220 nm to about 180 nm, or from about 190 nm to about 210 nm, most preferably said diameter is of about 200 nm; and/or
- vii) said polymersome has a vesicular morphology;
- viii) said polymersome is self-assembling;
- ix) said polymersome comprises an amphiphilic block copolymer, wherein said amphiphilic block copolymer: (a) is capable of sensing pH (e.g., is capable of undergoing protonation or de-protonation depending on the pH); and/or (b) is modified and/or substituted to become a pH sensitive moiety (e.g., is capable of undergoing protonation or de-protonation depending on the pH), preferably said modification and/or substitution comprising modifying and/or substituting a hydroxyl group (e.g., $\text{C}-\text{OH}$) into/with $-\text{NH}_3$ and carboxyl group (e.g., $-\text{COOH}$) (e.g., modifying and/or substituting PBD-PEO-OH functional group into PBD-PEO- NH_3 and PBD-PEO- COOH).

20. (canceled)

21. The polymersome according to claim 1, wherein said polynucleotide (e.g., SEQ ID NO: 16) encodes an immunogen.

22. (canceled)

23. The polymersome according to claim 1, wherein said polynucleotide (e.g., SEQ ID NO: 16) encoding one or more of the following:

- i) Influenza hemagglutinin (HA), preferably selected from the group consisting of: SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7 and SEQ ID NO: 8;
- ii) Swine Influenza hemagglutinin (HA), preferably SEQ ID NO: 6;
- iii) Ovalbumin (OVA), preferably SEQ ID NO: 4;
- iv) B16 peptide, preferably selected from the group consisting of: SEQ ID NO: 9, SEQ ID NO: 10 and SEQ ID NO: 11;
- v) MC38 peptide, preferably selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2 and SEQ ID NO: 3;
- vi) B16 and MC38 peptides, preferably said peptides are independently selected the groups: i) SEQ ID NOs: 1-3 and ii) SEQ ID NOs: 9-11; and/or
- vii) Porcine epidemic diarrhea virus SPIKE protein and a soluble fragment thereof, preferably a fragment of SEQ ID NO: 12, 13 or 14.

24. (canceled)

25. (canceled)

26. The polymersome according to claim 19, wherein said amphiphilic polymer comprises a diblock or a triblock (A-B-A or A-B-C) copolymer.

27.-29. (canceled)

30. The polymersome according to claim 19, wherein said amphiphilic polymer is a poly(butadiene)-poly(ethylene oxide) (PB-PEO) diblock copolymer, or wherein said amphiphilic polymer is a poly(dimethylsiloxane)-poly(ethylene oxide) (PDMS-PEO) diblock copolymer.

31. (canceled)

32. The polymersome according to claim 19, wherein said amphiphilic polymer is a poly(lactide)-poly(ethylene oxide)/1-palmitoyl-2-oleoyl-sn-glycero-3-phospho-L-serine (PLA-PEO/POPC) copolymer, preferably said PLA-PEO/POPC has a ratio of 75 to 25 (e.g., 75/25) of PLA-PEO to POPC (e.g., PLA-PEO/POPC).

33. The polymersome according to claim 19, wherein said amphiphilic polymer is a poly(caprolactone)-poly(ethylene oxide)/1-palmitoyl-2-oleoyl-sn-glycero-3-phospho-L-serine (PCL-PEO/POPC) copolymer, preferably said PCL-PEO/POPC has a ratio of 75 to 25 (e.g., 75/25) of PCL-PEO to POPC (e.g., PCL-PEO/POPC).

34. The polymersome according to claim 19, wherein said amphiphilic polymer is polybutadiene-polyethylene oxide (BD).

35. The polymersome according to claim 1, wherein said polymersome comprises diblock copolymer PBD₂₁-PEO₁₄ (BD21) and the triblock copolymer PMOXA₁₂-PDMS₅₅-PMOXA₁₂.

36. The polymersome according to claim 1, wherein said polymersome comprises a lipid polymer.

37. The polymersome according to claim 1, wherein said polymersome comprising one or more of the following:

- i) polybutadiene-polyethylene oxide (BD) and ionizable lipid Dlin-MC3-DMA (i.e., (6Z,9Z,28Z,31Z)-Heptatriaconta-6,9,28,31-tetraen-19-yl 4-(dimethylamino) butanoate);

- ii) PBD-PEO (poly(butadiene)-poly(ethylene oxide)) and ionizable lipid DLin-MC3-DMA (i.e., (6Z,9Z,28Z,31Z)-Heptatriaconta-6,9,28,31-tetraen-19-yl 4-(dimethylamino) butanoate), preferably comprising 15% or 30% DLin-MC3-DMA;
- iii) polybutadiene-polyethylene oxide (BD) and ionizable lipid 319 having Formula II;
- iv) polybutadiene-polyethylene oxide (BD) and ionizable lipid C12-200 having Formula III;
- v) polybutadiene-polyethylene oxide (BD) and ionizable lipid 5A2-SC8 having Formula IV;
- vi) polybutadiene-polyethylene oxide (BD) and ionizable lipid 306Oi10 having Formula V;
- vii) polybutadiene-polyethylene oxide (BD) and ionizable lipid 5 having Formula VI;
- viii) polybutadiene-polyethylene oxide (BD) and ionizable lipid SM-102 having Formula VII;
- ix) polybutadiene-polyethylene oxide (BD) and ionizable lipid A9 having Formula VIII;
- x) polybutadiene-polyethylene oxide (BD) and ionizable lipid ALC-0315 (i.e., [(4-hydroxybutyl) azanediyl]di(hexane-6,1-diyl)bis(2-hexyldecanoate) having Formula IX;

- xi) polybutadiene-polyethylene oxide (BD) and ionizable lipid Arcturus 2.2 (8.8) 4 C CH₃ having Formula X;
- xii) polybutadiene-polyethylene oxide (BD) and ionizable lipid Genevant CL1 having Formula XI;
- xiii) polybutadiene-polyethylene oxide (BD) and ionizable lipid ALC-0159 (i.e. 2-[(polyethylene glycol)-2000]-N,N-ditetradecylacetamide) having Formula XII.

38. A composition comprising a polymersome comprising a soluble encapsulated antigen, wherein said soluble encapsulated antigen is a polynucleotide selected from a RNA (e.g., mRNA) molecule or a DNA molecule and wherein the polymersome further comprises an ionizable lipid, wherein said ionizable lipid is: (i) a tertiary amine and/or comprising one or more tertiary amine moieties; and/or (ii) comprising a free electron pair at the nitrogen atom.

39. The composition according to claim **38**, wherein said composition is a pharmaceutical or diagnostic composition.

40.-43. (canceled)

* * * * *