



US 20250270499A1

(19) **United States**

(12) **Patent Application Publication**
Troutman et al.

(10) **Pub. No.: US 2025/0270499 A1**

(43) **Pub. Date: Aug. 28, 2025**

(54) **RECOMBINANT PRODUCTION OF
BACTEROIDES FRAGILIS CAPSULAR
POLYSACCHARIDE A IN ESCHERICHIA
COLI**

C12N 9/90 (2013.01); *C12N 9/93* (2013.01);
C12R 2001/19 (2021.05); *C12Y 101/01095*
(2013.01); *C12Y 204/01001* (2013.01); *C12Y*
206/01021 (2013.01); *C12Y 501/03007*
(2013.01); *C12Y 504/99004* (2013.01); *C12Y*
604/01001 (2013.01)

(71) Applicant: **The University of North Carolina at
Charlotte, Charlotte, NC (US)**

(72) Inventors: **Jerry M. Troutman, Charlotte, NC
(US); Beth A. Scarbrough, Charlotte,
NC (US)**

(57)

ABSTRACT

Provided are methods for producing polysaccharides in bacteria by expressing in a bacterium one or more coding sequences selected from the group consisting of a *pglF* dehydrogenase coding sequence, a *wbpP* UDP-N-acetyl-d-glucosamine C4 epimerase coding sequence, a *wcfR* aminotransferase coding sequence, and a *wcfS* phosphoglycosyltransferase coding sequence, a *wcfQ* glycosyltransferase coding sequence, a *wcfO* pyruvyltransferase coding sequence, a *wcfP* glycosyltransferase coding sequence, a *wcfM* UDP-galactopyranose mutase coding sequence, a *wcfN* glycosyltransferase coding sequence, a *wza* polysaccharide export protein coding sequence, a *wzx* fippase coding sequence, a *wzy* polymerase coding sequence, and a *wzz* coding sequence, wherein at least one of the coding sequences is heterologous to the bacterium. Also provided are expression cassettes with one or more of the disclosed coding sequences, recombinant bacteria that harbor one or more of the expression cassettes, and methods for producing immunogenic compositions using the polysaccharides produced by the recombinant bacteria.

Specification includes a Sequence Listing.

Related U.S. Application Data

(60) Provisional application No. 63/557,286, filed on Feb. 23, 2024.


Publication Classification


(51) **Int. Cl.**

C12N 1/20 (2006.01)
C08B 37/08 (2006.01)
C12N 9/00 (2006.01)
C12N 9/04 (2006.01)
C12N 9/10 (2006.01)
C12N 9/90 (2006.01)
C12R 1/19 (2006.01)

(52) **U.S. Cl.**

CPC *C12N 1/20* (2013.01); *C08B 37/0072*
(2013.01); *C12N 9/0006* (2013.01); *C12N*
9/1051 (2013.01); *C12N 9/1096* (2013.01);





4520




Fig. 1




FIG. 2A




Fig. 2B




FIG. 2C




FIG. 3




FIG. 4




FIG. 5




FIG. 6A

FIG. 6B

FIG. 6C




FIG. 7




FIG. 8

Plasmid	Predicted CPISA Intermediate	Monoisotopic mass (Da)	[M-H] ⁻	[M-2H] ²⁻
1. Empty Vector	BP	846.67	846	
2. pBAS8	BPP-AATGal	1112.73	1112	555
3. pBAS9	BPP-AATGalGal	1274.76	1274	636
4. pBAS10	BPP-AATGalPyrGal	1344.79	1344	671
5. pBAS15	BPP-AATGalPyrGal-GalNAc	1547.87	1547	773
6. pBAS18	BPP-AATGalPyrGal-GalNAc-Gal	1709.92	1709	854




FIG. 9




FIG. 10





FIG. 11

**FIG. 12**




FIG. 13







FIG. 15

RECOMBINANT PRODUCTION OF BACTEROIDES FRAGILIS CAPSULAR POLYSACCHARIDE A IN ESCHERICHIA COLI

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Patent Application Ser. No. 63/557,286, filed Feb. 23, 2024, the disclosure of which is incorporated herein by reference in its entirety.

GOVERNMENT INTEREST

[0002] This invention was made with government support under R01 GM123251 awarded by the National Institutes of Health (NIH). The government has certain rights in the invention.

REFERENCE TO SEQUENCE LISTING XML

[0003] The Sequence Listing XML associated with the instant disclosure has been electronically submitted to the United States Patent and Trademark Office via the Patent Center as a 125,559 byte UTF-8-encoded XML file created on Feb. 24, 2025 and entitled "1276_13_2_ST26.xml". The Sequence Listing submitted via Patent Center is hereby incorporated by reference in its entirety.

TECHNICAL FIELD

[0004] The presently disclosed subject matter relates to compositions and methods for producing polysaccharides in bacteria. In some embodiments, the compositions and methods are based on expression of a plurality of coding sequences in the bacteria, which in some embodiments are coding sequences at least one of which is heterologous to the bacteria.

BACKGROUND

[0005] Polysaccharides have drawn considerable attention as potent and renewable sources of biological polymers. These polymers are ubiquitous among prokaryotes, and many possess unique chemical structures that bestow distinct physical and chemical attributes. Diverse physicochemical properties present advantages of applying these polymers in various settings such as wound healing, pharmaceuticals, cosmetics, and food industries.

[0006] A major hindrance to the application of many bacterial polysaccharides is the inability to isolate or synthesize them in large quantities. This is due to complex growth requirements of bacterial hosts, complicated and laborious isolation procedures, and difficulty of genetic manipulation of hosts to aid in isolation procedures and improve yield. Thus, recombinant polysaccharide production in model strains of *E. coli* have been underway, including multiple *Streptococcus pneumoniae* capsule serotypes, and the N-linked glycan from *Campylobacter jejuni* (Feldman et al., 2005; Kay et al., 2016; Nothaft et al., 2016; Kay et al., 2022). In both instances, these recombinant polysaccharides were employed for the development of glycoconjugate vaccines.

SUMMARY

[0007] This summary lists several embodiments of the presently disclosed subject matter, and in many cases lists variations and permutations of these embodiments. This summary is merely exemplary of the numerous and varied embodiments. Mention of one or more representative features of a given embodiment is likewise exemplary. Such an embodiment can typically exist with or without the feature(s) mentioned; likewise, those features can be applied to other embodiments of the presently disclosed subject matter, whether listed in this summary or not. To avoid excessive repetition, this summary does not list or suggest all possible combinations of such features.

[0008] In some embodiments, the presently disclosed subject matter relates to methods for producing polysaccharides in bacteria. In some embodiments, the methods comprise expressing in the bacterium a plurality of coding sequences, the plurality of coding sequences comprising nucleotide sequences that encode a pglF dehydrogenase gene product, a wbpP UDP-N-acetyl-d-glucosamine C4 epimerase gene product, a wcfR aminotransferase gene product, and a wcfS phospho-glycosyltransferase gene product, and optionally further comprise nucleotide sequences encoding one or more of a wcfQ glycosyltransferase gene product, a wcfO pyruvyltransferase gene product, a wcfP glycosyltransferase gene product, a wcfM UDP-galactopyranose mutase gene product, a wcfN glycosyltransferase gene product, a wza polysaccharide export protein gene product, a wzx fippase gene product, a wzy polymerase gene product, and a wzz gene product, and further wherein at least one of the coding sequences is heterologous to the bacterium. In some embodiments, the polysaccharide is a capsular polysaccharide A (CPSA) and the bacterium expresses each of the PglF gene product, the wbpP gene product, the WcfR gene product, the WcfS gene product, the WcfQ gene product, the WcfO gene product, the WcfP gene product, the WcfM gene product, the WcfN gene product, the wzz gene product, the wzx fippase gene product, and the wzy polymerase gene product, at least one of which is heterologous to the bacterium. In some embodiments, the bacterium is an *Escherichia coli* (*E. coli*) bacterium. In some embodiments, the *E. coli* bacterium has reduced or absent waal biological activity. In some embodiments, the pglF gene product is a *Campylobacter jejuni* gene product, the wbpP gene product is a *Vibrio vulnificus* gene product, the wcfR gene product is a *Bacteroides fragilis* gene product, the wcfS gene product is a *Bacteroides fragilis* gene product, the wcfQ gene product is a *Bacteroides fragilis* gene product, the wcfO gene product is a *Bacteroides fragilis* gene product, the wcfP gene product is a *Bacteroides fragilis* gene product, the wcfM gene product is a *Bacteroides fragilis* gene product, the wcfN gene product is a *Bacteroides fragilis* gene product, the wza gene product is an *Escherichia coli* gene product, the wzx fippase gene product is a *Bacteroides fragilis* gene product, the wzy polymerase gene product is a *Bacteroides fragilis* gene product, the wzz gene product is a *Bacteroides fragilis* gene product.

[0009] In some embodiments, the polysaccharide is present in a lipid bilayer of the bacterium, optionally wherein the lipid bilayer is part of the bacterium's cell wall.

[0010] In some embodiments, the polysaccharide is a CPSA, optionally an aggregate of two or more CPSA tetrasaccharide monomers.

[0011] In some embodiments, the plurality of coding sequences are transcribed to produce a polycistronic messenger RNA (mRNA) molecule in the bacterium. In some embodiments, the polycistronic mRNA molecule comprises a ribosome-binding site (RBS) operatively linked to a translation initiation codon of each of at least two of the plurality of coding sequences, optionally wherein the polycistronic mRNA molecule comprises a plurality of ribosome-binding sites (RBSs), with an RBS present 5' to each of at least two, optionally 5' to each, translation initiation codon present in the polycistronic mRNA, wherein at least one RBS is located 3' to a translation termination codon of a first coding sequence and 5' to a translation initiation codon of a second coding sequence present on the polycistronic mRNA molecule.

[0012] In some embodiments, the bacterium does not express a functional *waal* O-antigen ligase gene product.

[0013] The presently disclosed subject matter also relates in some embodiments to bacteria comprising a plurality of coding sequences, the plurality of coding sequences comprising nucleotide sequences that encode two or more of a *pglF* dehydrogenase gene product, a *wbpP* UDP-GlcNAc C4 epimerase gene product, a *wcfR* aminotransferase gene product, a *wcfS* phospho-glycosyltransferase gene product, a *wcfQ* glycosyltransferase gene product, a *wcfO* pyruvyltransferase gene product, a *wcfP* glycosyltransferase gene product, a *wcfM* UDP-galactopyranose mutase gene product, a *wcfN* glycosyltransferase gene product, a *wza* polysaccharide export protein gene product, a *wzx* fippase gene product, a *wzy* polymerase gene product, and a *wzz* gene product, wherein at least one of the plurality of coding sequences is heterologous to the bacterium. In some embodiments, the bacterium is an *E. coli* bacterium. In some embodiments, the *pglF* gene product is a *Campylobacter jejuni* gene product; and/or the *wbpP* gene product is a *Vibrio vulnificus* gene product; and/or the *wcfR* gene product is a *Bacteroides fragilis* gene product; and/or the *wcfS* gene product is a *Bacteroides fragilis* gene product; and/or the *wcfQ* gene product is a *Bacteroides fragilis* gene product; and/or the *wcfO* gene product is a *Bacteroides fragilis* gene product; and/or the *wcfP* gene product is a *Bacteroides fragilis* gene product; and/or the *wcfM* gene product is a *Bacteroides fragilis* gene product; and/or the *wcfN* gene product is a *Bacteroides fragilis* gene product; and/or the *wza* gene product is an *Escherichia coli* gene product; and/or the *wzx* fippase gene product is a *Bacteroides fragilis* gene product; and/or the *wzy* polymerase gene product is a *Bacteroides fragilis* gene product; and/or the *wzz* gene product is a *Bacteroides fragilis* gene product. In some embodiments, the plurality of coding sequences are transcribed in the bacterium to produce a polycistronic messenger RNA (mRNA) molecule in the bacterium. In some embodiments, the polycistronic mRNA molecule comprises a ribosome-binding site (RBS) operatively linked to a translation initiation codon of each of at least two of the plurality of coding sequences, optionally wherein the polycistronic mRNA molecule comprises a plurality of ribosome-binding sites (RBSs), with an RBS present 5' to each of at least two, optionally 5' to each, translation initiation codon present in the polycistronic mRNA, wherein at least one RBS is located 3' to a translation termination codon of a first coding sequence and 5' to a translation initiation codon of a second coding sequence present on the polycistronic mRNA molecule. In some embodiments, translation of the

polycistronic mRNA molecule in the bacterium results in production of a polysaccharide in the bacterium, optionally wherein the polysaccharide is present in a lipid bilayer of the bacterium, further optionally wherein the lipid bilayer is part of the bacterium's cell wall. In some embodiments, the polysaccharide is a CPSA, optionally an aggregate of two or more CPSA tetrasaccharide monomers.

[0014] In some embodiments, the presently disclosed subject matter relates to methods for producing immunogenic compositions. In some embodiments, the methods comprise mixing an antigenic molecule with a capsular polysaccharide A (CPSA), wherein the CPSA is produced by a bacterium as disclosed herein.

[0015] In some embodiments, the presently disclosed subject matter also relates to expression cassettes. In some embodiments, the expression cassettes comprise a plurality of coding sequences derived from at least two different bacterial species, wherein the plurality of coding sequences are selected from the group consisting of a *pglF* dehydrogenase gene coding sequence, a *wbpP* UDP-N-acetyl-d-glucosamine C4 epimerase gene coding sequence, a *wcfR* aminotransferase gene coding sequence, a *wcfS* phospho-glycosyltransferase gene coding sequence, a *wcfQ* glycosyltransferase gene coding sequence, a *wcfO* pyruvyltransferase gene coding sequence, a *wcfP* glycosyltransferase gene coding sequence, a *wcfM* UDP-galactopyranose mutase gene coding sequence, a *wcfN* glycosyltransferase gene coding sequence, a *wza* polysaccharide export protein gene coding sequence, a *wzx* fippase gene coding sequence, a *wzy* polymerase gene coding sequence, and a *wzz* gene coding sequence.

[0016] In some embodiments, the expression cassette comprises a combination of coding sequences selected from the group consisting of:

[0017] *pglF*, *wcfR*, and *wcfS*;

[0018] *pglF*, *wcfR*, *wcfS*, and *wcfQ*;

[0019] *pglF*, *wcfR*, *wcfS*, *wcfQ*, and *wcfO*;

[0020] *pglF*, *wcfR*, *wcfS*, *wcfQ*, *wcfO*, *wbpP*, and *wcfP*;

[0021] *pglF*, *wcfR*, *wcfS*, *wcfQ*, *wcfO*, *wbpP*, *wcfP*, *wcfM*, and *wcfN*; and

[0022] *pglF*, *wcfR*, *wcfS*, *wcfQ*, *wcfO*, *wbpP*, *wcfP*, *wcfM*, *wcfN*, *wzx*, and *wzy*.

[0023] In some embodiments, the expression cassette is operatively linked to a promoter such that transcription from the promoter results in the production of a polycistronic mRNA that is translatable to produce polypeptides encoded by the coding sequences. In some embodiments, the polycistronic mRNA molecule comprises a ribosome-binding site (RBS) operatively linked to a translation initiation codon of one or more of, optionally each of, the plurality of coding sequences, wherein at least one RBS is located 3' to a translation termination codon of a first coding sequence and 5' to a translation initiation codon of a second coding sequence, wherein the first coding sequence and the second coding sequence are adjacent to each other on the polycistronic mRNA molecule, the first coding sequence being present 5' to the second coding sequence. In some embodiments, the polycistronic mRNA molecule is operably linked to a promoter that is active in the bacterium, optionally wherein the promoter is an inducible promoter, optionally an isopropyl- β -d-thiogalactopyranoside (IPTG)-inducible promoter, further optionally wherein the promoter is a T5 promoter.

[0024] Accordingly, it is an object of the presently disclosed subject matter to provide recombinant nucleic acids that include coding sequences from different organisms that can be employed for producing polysaccharides including but not limited to CPSA in bacteria.

[0025] This and other objects are achieved in whole or in part by the presently disclosed subject matter. Further, an object of the presently disclosed subject matter having been stated above, other objects and advantages of the presently disclosed subject matter will become apparent to those skilled in the art after a study of the following Description, Figures, and EXAMPLES.

BRIEF DESCRIPTION OF THE FIGURES

[0026] FIG. 1 is a representation of the CPSA biosynthesis pathway and structure in *Bacteroides fragilis*, and a reconstitution of its biosynthesis in *E. coli*. The previously determined structure of the CPSA unit oligosaccharide is highlighted in yellow.

[0027] FIGS. 2A-2C illustrate Recombinant UDP-AAT-Gal and UDP-GalNAc biosynthesis in *E. coli*. (FIG. 2A) UDP-AATGal is produced from UDP-GlcNAc by PglF from *Campylobacter jejuni* and WcfR from *B. fragilis*. (FIG. 2B) UDP-GalNAc is produced from UDP-GlcNAc by WbpP from *Vibrio vulnificus*. (FIG. 2C) Schematic of CPSA biosynthesis in *E. coli* utilizing PglF and WbpP.

[0028] FIG. 3 depicts the LCMS identification of individual BPP-linked CPSA intermediates recombinantly produced in *E. coli*. Each BPP-linked CPSA intermediate was identified in lysates of *E. coli* DH5a cells recombinantly expressing genes from the CPSA biosynthesis operon using ESI-LCMS and SIM. Descriptions of plasmids used to generate individual intermediates in *E. coli* and the exact mass of each intermediate can be found in Table 4. Only chromatograms generated from SIM of the anticipated intermediates were overlaid in FIG. 3 (see also FIG. 11). 1: BP; 2: BPP-mono; 3: BPP-Di; 4: BPP-PyrDi; 5: BPP-Tri; 6: BPP-tetra.

[0029] FIG. 4 shows Both flippase (Wzy) and polymerase (Wzx) from *B. fragilis* may be required for polymerization of CPSA oligomers in *E. coli*. SDS-PAGE (left panel) and anti-CPSA western blot (right panel) of *E. coli* DH5a cell lysates expressing an empty vector control (pQE-80L), pBAS16 (-wzx, -wzy_{BF}), pBAS17 (+wzx, +wzy_{BF}), pBAS18 (+wzx_{BF}), or pBAS19 (+wzy_{BF}). The left panel is a Ponceau stained SDS-PAGE gel of normalized cell lysates. The right panel is an anti-CPSA western blot of cell lysates. Genes assembled in plasmids are listed in Tables 4 and 5. CPSA positive (+) indicates that all genes are present to produce CPSA oligosaccharide.

[0030] FIG. 5 shows confocal microscopy of cells expressing CPSA polymers (pBAS17; right panel) compared to an empty vector control (pQE-80L; left panel).

[0031] FIGS. 6A-6C show that CPSA polymers were detected in both aqueous and organic fractions of *E. coli* pBAS17 lysates. Wild-type (FIG. 6A) and a Δ waaL mutant (FIG. 6B) of *E. coli* MG1655 pBAS17 were evaluated for the presence of CPSA polymers in aqueous, organic, and insoluble cell fractions. Cell fractions were separated using SDS-PAGE and anti-CPSA antiserum. Lanes designated with an "A" are aqueous fractions, lanes designated with "O" are organic fractions, and lanes designated with "I" are insoluble cell fractions. (FIG. 6C) SDS-PAGE and anti-CPSA western blot of whole cell lysates.

[0032] FIG. 7 shows the whole cell western blot of *E. coli* MG1655 expressing pBAS16 or an empty vector control. The left panel shows a whole-cell dot blot of wild-type (WT) and mutant MG1655 expressing pBAS16 (-wzx, -wzy). The right panel shows PRO-Q™ Emerald staining of LPS in normalized, pronase-treated lysates. The lower (blue) arrow indicates lipid A core. The upper (white) arrow likely indicates CPSA oligomer(s) ligated to LOS.

[0033] FIG. 8 shows CPSA surface expression was not abolished in a Δ waaL or Δ wza mutant expressing pBAS17. The left panel is a whole cell dot blot of *E. coli* MG1655 expressing pBAS17 (+wzx, +wzy) or an empty vector control (pQE-80L; EV). The right panel shows PRO-Q™ Emerald 300 stain of LPS from pronase-treated cell lysates. The lower (blue) arrow indicates lipid A core. The upper (white) arrow likely indicates CPSA oligomer(s) ligated to lipid A core.

[0034] FIG. 9 is a series of spectra of recombinantly produced BPP-linked CPSA intermediates identified within *E. coli* cell lysates. Plasmids, CPSA intermediate, and [M-H]⁻ ion of each intermediate described in the Table above the spectra and correspond to each numbered spectrum.

[0035] FIG. 10 shows that CPSA intermediates were not detected in *E. coli* expressing an empty vector control. Lysates of *E. coli* expressing an empty vector control were evaluated with Liquid Chromatography-Electrospray Ionization-Mass Spectrometry (ESI-LC-MS) and selected ion monitoring (SIM) for all BPP-linked CPSA intermediates. A contaminant with a m/z identical to the first CPSA intermediate (BPP-mono) is observed at 2.84 min which is inconsistent with the retention time for a BPP-linked monosaccharide. The panel at the upper right is a zoomed in region of the chromatogram at the lower left, showing less abundant contaminants.

[0036] FIG. 11 is a series of graphs the LC-MS with SIM of recombinant *E. coli* cell lysates with overexpressed CPSA plasmids. Each *E. coli* lysate was evaluated for the presence of (1) BP, (2) the expected CPSA intermediate, and (3) the subsequent intermediate, with the exception of pBAS11. Cells expressing pBAS11, lacking the C4 epimerase, wbpP, did not produce the CPSA BPP-linked trisaccharide. Only BPP-linked pyruvylated disaccharide (BPP-pyrDi) could be detected in cell expressing pBAS11. pBAS15 does include wbpP, and BPP-linked trisaccharide was detected in these lysates. Plasmids are described in Table 4.

[0037] FIG. 12 is a whole-cell dot blot of *E. coli* DH5a expressing CPSA plasmids and showing that *E. coli* DH5a expressing pBAS16 and pBAS17 reacted with anti-CPSA antiserum. The plasmids employed for this dot blot are described in Table 4. A positive signal for CPSA was only observed with cells expressing the pBAS16 or the pBAS17 plasmid.

[0038] FIG. 13 graphs the LC-MS analysis of BPP-linked CPSA tetrasaccharide in *E. coli* cell lysates. Normalized cell lysates expressing pBAS16 (-wzx, -wzy; left panel) or pBAS17 (+wzx, +wzy; right panel) were evaluated for the presence of BP and BPP-linked CPSA tetrasaccharide using ESI-LC-MS with SIM. BPP-tetrasaccharide was still present in pBAS17 cell lysates but appeared to be much less abundant compared to cells expressing pBAS16.

[0039] FIGS. 14A and 14B are schematic diagrams of an exemplary pathway for producing CPSA in bacteria (FIG. 14A) and a representative arrangement of various coding

sequences in the expression cassettes and vectors of the presently disclosed subject matter (FIG. 14B).

[0040] FIG. 15 is a map of plasmid pBAS17, which includes the coding sequences shown in FIG. 14B.

DETAILED DESCRIPTION

[0041] The presently disclosed subject matter now will be described more fully hereinafter in the following detailed description of the presently disclosed subject matter, in which some, but not all embodiments of the presently disclosed subject matter are described. Indeed, the presently disclosed subject matter may be embodied in many different forms and should not be construed as limited to the embodiments set forth herein; rather, these embodiments are provided so that this disclosure will satisfy applicable legal requirements.

I. Definitions

[0042] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the presently described subject matter belongs.

[0043] Throughout the specification and claims, a given chemical formula or name shall encompass all active optical and stereoisomers, as well as racemic mixtures where such isomers and mixtures exist.

[0044] The terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting of the presently disclosed subject matter.

[0045] While the following terms are believed to be well understood by one of ordinary skill in the art, the following definitions are set forth to facilitate explanation of the presently disclosed subject matter.

[0046] References to techniques employed herein are intended to refer to the techniques as commonly understood in the art, including variations on those techniques or substitutions of equivalent techniques that would be apparent to one of skill in the art.

[0047] In describing the presently disclosed subject matter, it will be understood that a number of techniques and steps are disclosed. Each of these has individual benefit and each can also be used in conjunction with one or more, or in some cases all, of the other disclosed techniques.

[0048] Accordingly, for the sake of clarity, this description will refrain from repeating every possible combination of the individual steps in an unnecessary fashion. Nevertheless, the specification and claims should be read with the understanding that such combinations are entirely within the scope of the presently disclosed and claimed subject matter.

[0049] Following long-standing patent law convention, the terms “a”, “an”, and “the” refer to “one or more” when used in this application, including in the claims. For example, the phrase “a protein” refers to one or more proteins, including a plurality of the same protein. Similarly, the phrase “at least one”, when employed herein to refer to an entity, refers to, for example, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 75, 100, or more of that entity, including but not limited to whole number values between 1 and 100 and greater than 100.

[0050] Unless otherwise indicated, all numbers expressing quantities of ingredients, reaction conditions, and so forth used in the specification and claims are to be understood as being modified in all instances by the term “about”. The term

“about”, as used herein when referring to a measurable value such as an amount of mass, weight, time, volume, concentration, or percentage, is meant to encompass variations of in some embodiments +20%, in some embodiments +10%, in some embodiments +5%, in some embodiments +1%, in some embodiments +0.5%, and in some embodiments +0.1% from the specified amount, as such variations are appropriate to perform the disclosed methods and/or employ the disclosed compositions. Accordingly, unless indicated to the contrary, the numerical parameters set forth in this specification and attached claims are approximations that can vary depending upon the desired properties sought to be obtained by the presently disclosed subject matter.

[0051] As used herein, the term “and/or” when used in the context of a list of entities, refers to the entities being present singly or in combination. Thus, for example, the phrase “A, B, C, and/or D” includes A, B, C, and D individually, but also includes any and all combinations and subcombinations of A, B, C, and D.

[0052] As used herein, the term “adjuvant” refers to a substance that elicits an enhanced immune response when used in combination with a specific antigen.

[0053] As used herein, the terms “administration of” and/or “administering” a compound should be understood to refer to providing a compound of the presently disclosed subject matter to a subject in need of treatment.

[0054] The term “comprising”, which is synonymous with “including”, “containing”, or “characterized by”, is inclusive or open-ended and does not exclude additional, unrecited elements and/or method steps. “Comprising” is a term of art that means that the named elements and/or steps are present, but that other elements and/or steps can be added and still fall within the scope of the relevant subject matter.

[0055] As used herein, the phrase “consisting essentially of” limits the scope of the related disclosure or claim to the specified materials and/or steps, plus those that do not materially affect the basic and novel characteristic(s) of the disclosed and/or claimed subject matter. For example, a pharmaceutical composition can “consist essentially of” a pharmaceutically active agent or a plurality of pharmaceutically active agents, which means that the recited pharmaceutically active agent(s) is/are the only pharmaceutically active agent(s) present in the pharmaceutical composition. It is noted, however, that carriers, excipients, and/or other inactive agents can and likely would be present in such a pharmaceutical composition and are encompassed within the nature of the phrase “consisting essentially of”.

[0056] As used herein, the phrase “consisting of” excludes any element, step, or ingredient not specifically recited. It is noted that, when the phrase “consists of” appears in a clause of the body of a claim, rather than immediately following the preamble, it limits only the element set forth in that clause; other elements are not excluded from the claim as a whole. With respect to the terms “comprising”, “consisting of”, and “consisting essentially of”, where one of these three terms is used herein, the presently disclosed and claimed subject matter can include the use of either of the other two terms. For example, a composition that in some embodiments comprises a given active agent also in some embodiments can consist essentially of that same active agent, and indeed can in some embodiments consist of that same active agent.

[0057] The term “biocompatible”, as used herein, refers to a material that does not elicit a substantial detrimental response in the host.

[0058] As used herein, the terms “biologically active fragment” and “bioactive fragment” of a peptide encompass natural and synthetic portions of a longer peptide or protein that are capable of specific binding to their natural ligand and/or of performing a desired function of a protein, for example, a fragment of a protein of larger peptide which still contains the epitope of interest and is immunogenic.

[0059] The term “biological sample”, as used herein, refers to samples obtained from a subject, including but not limited to skin, hair, tissue, blood, plasma, cells, sweat, and urine.

[0060] A “coding region” of a gene comprises the nucleotide residues of the coding strand of the gene and the nucleotides of the non-coding strand of the gene which are homologous with or complementary to, respectively, the coding region of an mRNA molecule which is produced by transcription of the gene.

[0061] “Complementary” as used herein refers to the broad concept of subunit sequence complementarity between two nucleic acids (e.g., two DNA molecules). When a nucleotide position in both of the molecules is occupied by nucleotides normally capable of base pairing with each other at a given position, the nucleic acids are considered to be complementary to each other at this position. Thus, two nucleic acids are complementary to each other when a substantial number (in some embodiments at least 50%) of corresponding positions in each of the molecules are occupied by nucleotides that can base pair with each other (e.g., A:T and G:C nucleotide pairs). Thus, it is known that an adenine residue of a first nucleic acid region is capable of forming specific hydrogen bonds (“base pairing”) with a residue of a second nucleic acid region which is antiparallel to the first region if the residue is thymine or uracil.

[0062] Similarly, it is known that a cytosine residue of a first nucleic acid strand is capable of base pairing with a residue of a second nucleic acid strand which is antiparallel to the first strand if the residue is guanine. A first region of a nucleic acid is complementary to a second region of the same or a different nucleic acid if, when the two regions are arranged in an antiparallel fashion, at least one nucleotide residue of the first region is capable of base pairing with a residue of the second region. By way of example and not limitation, the first region comprises a first portion and the second region comprises a second portion, whereby, when the first and second portions are arranged in an antiparallel fashion, in some embodiments at least about 50%, in some embodiments at least about 75%, in some embodiments at least about 90%, and in some embodiments at least about 95% of the nucleotide residues of the first portion are capable of base pairing with nucleotide residues in the second portion. In some embodiments, all nucleotide residues of the first portion are capable of base pairing with nucleotide residues in the second portion.

[0063] As used herein, the terms “condition”, “disease condition”, “disease”, “disease state”, and “disorder” refer to physiological states in which diseased cells or cells of interest can be targeted with the compositions of the presently disclosed subject matter.

[0064] As used herein, the term “diagnosis” refers to detecting a risk or propensity to a condition, disease, or disorder. In any method of diagnosis exist false positives and false negatives. Any one method of diagnosis does not provide 100% accuracy.

[0065] A “disease” is a state of health of an animal wherein the animal cannot maintain homeostasis, and wherein if the disease is not ameliorated then the animal’s health continues to deteriorate.

[0066] In contrast, a “disorder” in an animal is a state of health in which the animal is able to maintain homeostasis, but in which the animal’s state of health is less favorable than it would be in the absence of the disorder. Left untreated, a disorder does not necessarily cause a further decrease in the animal’s state of health.

[0067] As used herein, an “effective amount” or “therapeutically effective amount” refers to an amount of a compound or composition sufficient to produce a selected effect, such as but not limited to alleviating symptoms of a condition, disease, or disorder. In the context of administering compounds in the form of a combination, such as multiple compounds, the amount of each compound, when administered in combination with one or more other compounds, may be different from when that compound is administered alone. Thus, an effective amount of a combination of compounds refers collectively to the combination as a whole, although the actual amounts of each compound may vary. The term “more effective” means that the selected effect occurs to a greater extent by one treatment relative to the second treatment to which it is being compared.

[0068] “Encoding” refers to the inherent property of specific sequences of nucleotides in a polynucleotide, such as a gene, a cDNA, or an mRNA, to serve as templates for synthesis of other polymers and macromolecules in biological processes having either a defined sequence of nucleotides (e.g., rRNA, tRNA, and mRNA) or a defined sequence of amino acids and the biological properties resulting therefrom. Thus, a gene encodes a protein if transcription and translation of an mRNA corresponding to or derived from that gene produces the protein in a cell or other biological system and/or an in vitro or ex vivo system. Both the coding strand, the nucleotide sequence of which is identical to the mRNA sequence (with the exception of uracil bases presented in the latter) and is usually provided in Sequence Listing, and the non-coding strand, used as the template for transcription of a gene or cDNA, can be referred to as encoding the protein or other product of that gene or cDNA.

[0069] In some embodiments, the terms “fragment”, “segment”, or “subsequence” as used herein refers to a portion of an amino acid sequence, comprising at least one amino acid, or a portion of a nucleic acid sequence comprising at least one nucleotide. Thus, in some embodiments, the terms “fragment”, “segment”, and “subsequence” are used interchangeably herein. In some embodiments, the term “fragment” refers to a compound (e.g., a small molecule compound, such as a small molecule comprising a purine scaffold) that can react with a reactive amino acid residue (e.g., a reactive cysteine) to form an adduct comprising a modified amino acid residue. Thus, in some embodiments, the terms “fragment” and “ligand” are used interchangeably. In some embodiments, the term “fragment” refers to that portion of a ligand that remains covalently attached to the reactive amino acid residue.

[0070] As used herein, a “ligand” is a compound (e.g., a purine-based compound) that specifically binds to a target compound or molecule, such as a reactive nucleophilic amino acid residue in a protein. In some embodiments, the ligand can bind to the target covalently. A ligand “specifically binds to” or “is specifically reactive with” a compound

(e.g., a reactive amino acid residue) when the ligand functions in a binding reaction which is determinative of the presence of the compound in a sample of heterogeneous compounds.

[0071] As used herein, a “functional” biological molecule is a biological molecule in a form in which it exhibits a property by which it can be characterized. A functional enzyme, for example, is one that exhibits the characteristic catalytic activity by which the enzyme can be characterized.

[0072] As used herein, the term “linker” refers to a molecule that joins two other molecules either covalently or noncovalently, such as but not limited to through ionic or hydrogen bonds or van der Waals interactions.

[0073] The term “pharmaceutical composition” refers to a composition comprising at least one active ingredient, whereby the composition is amenable to investigation for a specified, efficacious outcome in a mammal (for example, without limitation, a human). Those of ordinary skill in the art will understand and appreciate the techniques appropriate for determining whether an active ingredient has a desired efficacious outcome based upon the needs of the artisan.

[0074] “Pharmaceutically acceptable” means physiologically tolerable, for either human or veterinary application. Similarly, “pharmaceutical compositions” include formulations for human and veterinary use.

[0075] As used herein, the term “pharmaceutically acceptable carrier” means a chemical composition with which an appropriate compound or derivative can be combined and which, following the combination, can be used to administer the appropriate compound to a subject.

[0076] “Plurality” means at least two.

[0077] “Polypeptide” refers to a polymer composed of amino acid residues, related naturally occurring structural variants, and synthetic non-naturally occurring analogs thereof linked via peptide bonds, related naturally occurring structural variants, and synthetic non-naturally occurring analogs thereof.

[0078] As used herein, the term “mass spectrometry” (MS) refers to a technique for the identification and/or quantitation of molecules in a sample. MS includes ionizing the molecules in a sample, forming charged molecules; separating the charged molecules according to their mass-to-charge ratio; and detecting the charged molecules. MS allows for both the qualitative and quantitative detection of molecules in a sample. The molecules can be ionized and detected by any suitable means known to one of skill in the art. Some examples of mass spectrometry are “tandem mass spectrometry” or “MS/MS,” which are the techniques wherein multiple rounds of mass spectrometry occur, either simultaneously using more than one mass analyzer or sequentially using a single mass analyzer. The term “mass spectrometry” can refer to the application of mass spectrometry to protein analysis. In some embodiments, electrospray ionization (ESI) and matrix-assisted laser desorption/ionization (MALDI) can be used in this context. In some embodiments, intact protein molecules can be ionized by the above techniques, and then introduced to a mass analyzer. Alternatively, protein molecules can be broken down into smaller peptides, for example, by enzymatic digestion by a protease, such as trypsin. Subsequently, the peptides are introduced into the mass spectrometer and identified by peptide mass fingerprinting or tandem mass spectrometry.

[0079] As used herein, the term “mass spectrometer” is used to refer an apparatus for performing mass spectrometry

that includes a component for ionizing molecules and detecting charged molecules. Various types of mass spectrometers can be employed in the methods of the presently disclosed subject matter. For example, whole protein mass spectroscopy analysis can be conducted using time-of-flight (TOF) or Fourier transform ion cyclotron resonance (FT-ICR) instruments. For peptide mass analysis, MALDI time-of-flight instruments can be employed, as they permit the acquisition of peptide mass fingerprints (PMFs) at high pace. Multiple stage quadrupole-time-of-flight and the quadrupole ion trap instruments can also be used.

[0080] As used herein, the term “Western blot,” which can be also referred to as “immunoblot”, and related terms refer to an analytical technique used to detect specific proteins in a sample. The technique uses gel electrophoresis to separate the proteins, which are then transferred from the gel to a membrane (typically nitrocellulose or PVDF) and stained, in membrane, with antibodies specific to the target protein.

[0081] The expression “stable isotope labeling by amino acids in cell culture” (SILAC) is used herein to refer to an approach for incorporation of a label into proteins for mass spectrometry (MS)-based quantitative proteomics. SILAC comprises metabolic incorporation of a given “light” or “heavy” form of the amino acid into the proteins. For example, SILAC comprises the incorporation of amino acids with substituted stable isotopic nuclei (e.g. deuterium, ^{13}C , ^{15}N). In an illustrative SILAC experiment, two cell populations are grown in culture media that are identical, except that one of them contains a “light” and the other a “heavy” form of a particular amino acid (for example, ^{12}C and ^{13}C labeled L-lysine, respectively). When the labeled analog of an amino acid is supplied to cells in culture instead of the natural amino acid, it is incorporated into all newly synthesized proteins. After a number of cell divisions, each instance of the amino acid is replaced by its isotope-labeled analog. Since there is little chemical difference between the labeled amino acid and the natural amino acid isotopes, the cells behave substantially similar to the control cell population grown in the presence of a normal amino acid.

[0082] The term “protein” typically refers to large polypeptides. Conventional notation is used herein to portray polypeptide sequences: the left-hand end of a polypeptide sequence is the amino-terminus; the right-hand end of a polypeptide sequence is the carboxyl-terminus.

[0083] As used herein, the term “purified” and like terms relate to an enrichment of a molecule or compound relative to other components normally associated with the molecule or compound in a native environment. The term “purified” does not necessarily indicate that complete purity of the particular molecule has been achieved during the process.

[0084] A “highly purified” compound as used herein refers to a compound that is in some embodiments greater than 90% pure, that is in some embodiments greater than 95% pure, and that is in some embodiments greater than 98% pure.

[0085] The term “subject” as used herein refers to a member of species for which treatment and/or prevention of a disease or disorder using the compositions and methods of the presently disclosed subject matter might be desirable. Accordingly, the term “subject” is intended to encompass in some embodiments any member of the Kingdom Animalia including, but not limited to the phylum Chordata (e.g., members of Classes Osteichthyes (bony fish), Amphibia

(amphibians), Reptilia (reptiles), Aves (birds), and Mammalia (mammals), and all Orders and Families encompassed therein.

[0086] The compositions and methods of the presently disclosed subject matter are particularly useful for warm-blooded vertebrates. Thus, in some embodiments the presently disclosed subject matter concerns mammals and birds. More particularly provided are compositions and methods derived from and/or for use in mammals such as humans and other primates, as well as those mammals of importance due to being endangered (such as Siberian tigers), of economic importance (animals raised on farms for consumption by humans) and/or social importance (animals kept as pets or in zoos) to humans, for instance, carnivores other than humans (such as cats and dogs), swine (pigs, hogs, and wild boars), ruminants (such as cattle, oxen, sheep, giraffes, deer, goats, bison, and camels), rodents (such as mice, rats, and rabbits), marsupials, and horses. Also provided is the use of the disclosed methods and compositions on birds, including those kinds of birds that are endangered, kept in zoos, as well as fowl, and more particularly domesticated fowl, e.g., poultry, such as turkeys, chickens, ducks, geese, guinea fowl, and the like, as they are also of economic importance to humans. Thus, also provided is the use of the disclosed methods and compositions on livestock, including but not limited to domesticated swine (pigs and hogs), ruminants, horses, poultry, and the like.

[0087] A “sample”, as used herein, refers in some embodiments to a biological sample from a subject, including, but not limited to, normal tissue samples, diseased tissue samples, biopsies, blood, saliva, feces, semen, tears, and urine. A sample can also be any other source of material obtained from a subject which contains proteins, cells, tissues, or fluid of interest. A sample can also be obtained from cell or tissue culture.

[0088] The term “standard”, as used herein, refers to something used for comparison. For example, it can be a known standard agent or compound which is administered and used for comparing results when administering a test compound, or it can be a standard parameter or function which is measured to obtain a control value when measuring an effect of an agent or compound on a parameter or function. Standard can also refer to an “internal standard”, such as an agent or compound which is added at known amounts to a sample and is useful in determining such things as purification or recovery rates when a sample is processed or subjected to purification or extraction procedures before a marker of interest is measured. Internal standards are often a purified marker of interest which has been labeled, such as with a radioactive isotope, allowing it to be distinguished from an endogenous marker.

[0089] A “subject” of analysis, diagnosis, or treatment is an animal. Such animals include mammals, in some embodiments, humans.

[0090] As used herein, a “subject in need thereof” is a patient, animal, mammal, or human, who will benefit from the method of this presently disclosed subject matter.

[0091] The term “substantially pure” describes a compound, e.g., a protein or polypeptide, which has been separated from components which naturally accompany it. Typically, a compound is substantially pure when in some embodiments at least 10%, in some embodiments at least 20%, in some embodiments at least 50%, in some embodiments at least 60%, in some embodiments at least 75%, in

some embodiments at least 90%, and in some embodiments at least 99% of the total material (by volume, by wet or dry weight, or by mole percent or mole fraction) in a sample is the compound of interest. Purity can be measured by any appropriate method, e.g., in the case of polypeptides by column chromatography, gel electrophoresis, or HPLC analysis. A compound, e.g., a protein, is also substantially purified when it is essentially free of naturally associated components or when it is separated from the native contaminants which accompany it in its natural state.

[0092] The term “symptom”, as used herein, refers to any morbid phenomenon or departure from the normal in structure, function, or sensation, experienced by the patient and indicative of disease. In contrast, a “sign” is objective evidence of disease. For example, a bloody nose is a sign. It is evident to the patient, doctor, nurse, and other observers.

[0093] A “therapeutic” treatment is a treatment administered to a subject who exhibits signs of pathology for the purpose of diminishing or eliminating those signs.

[0094] A “therapeutically effective amount” of a compound is that amount of compound which is sufficient to provide a beneficial effect to the subject to which the compound is administered.

[0095] As used herein, the phrase “therapeutic agent” refers to an agent that is used to, for example, treat, inhibit, prevent, mitigate the effects of, reduce the severity of, reduce the likelihood of developing, slow the progression of, and/or cure, a disease or disorder.

[0096] The terms “treatment” and “treating” as used herein refer to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) the targeted pathologic condition, prevent the pathologic condition, pursue or obtain beneficial results, and/or lower the chances of the individual developing a condition, disease, or disorder, even if the treatment is ultimately unsuccessful. Those in need of treatment include those already with the condition as well as those prone to have or predisposed to having a condition, disease, or disorder, or those in whom the condition is to be prevented.

[0097] As used herein, the terms “vector”, “cloning vector”, and “expression vector” refer to a vehicle by which a polynucleotide sequence (e.g., a foreign gene) can be introduced into a host cell, so as to transduce and/or transform the host cell in order to promote expression (e.g., transcription and translation) of the introduced sequence. Vectors include plasmids, phages, viruses, etc.

[0098] All genes, gene names, and gene products disclosed herein are intended to correspond to homologs and/or orthologs from any species for which the compositions and methods disclosed herein are applicable. Thus, the terms include, but are not limited to genes and gene products from humans and mice. It is understood that when a gene or gene product from a particular species is disclosed, this disclosure is intended to be exemplary only, and is not to be interpreted as a limitation unless the context in which it appears clearly indicates.

II. Exemplary Compositions and Methods of Use Thereof

II.A. Introduction

[0099] In describing the presently disclosed subject matter, it will be understood that a number of techniques and steps are disclosed. Each of these has individual benefit and each can also be used in conjunction with one or more, or in

some cases all, of the other disclosed techniques. Accordingly, for the sake of clarity, this description will refrain from repeating every possible combination of the individual steps in an unnecessary fashion. Nevertheless, the specification and claims should be read with the understanding that such combinations are entirely within the scope of the presently disclosed subject matter and the claims.

[0100] A goal of the presently disclosed subject matter was to devise a recombinant system for the overproduction of capsular polysaccharide A (CPSA) from *Bacteroides fragilis* NTCT 9343. CPSA belongs to a unique class of zwitterionic polysaccharides that possess a distinct charge motif of alternating positive and negative charges. In contrast to neutral carbohydrates, zwitterionic polysaccharides elicit a potent, T-cell dependent immune response. 5-6 These distinctive and potent immunomodulatory properties of CPSA and other zwitterionic polysaccharides are modulated by both positively and negatively charged substituents, as well as their helical structure (see Tzianabos et al., 2000; Wang et al., 2000; Choi et al., 2002; Tzianabos et al., 2001). Akin to an immune response elicited by a proteinaceous antigen, CPSA is processed by antigen presenting cells and presented to CD4+ T-cells via the major histocompatibility complex II (MHCII). See e.g., Herget et al. (2008) Statistical analysis of the Bacterial Carbohydrate Structure Data Base (BCSDB): characteristics and diversity of bacterial carbohydrates in comparison with mammalian glycans. BMC Struct Biol August 11:8:35; Surana & Kasper (2012) The yin yang of bacterial polysaccharides: lessons learned from *B. fragilis* PSA. Immunol Rev 245(1):13-26; Porter & Martens. (2017) The Critical

[0101] Roles of Polysaccharides in Gut Microbial Ecology and Physiology. Annu Rev Microbiol 71:349-369. As such, CPSA has been employed in the development of completely carbohydrate-based vaccines (Troy & Kasper (2010) Beneficial effects of *Bacteroides fragilis* polysaccharides on the immune system. Front Biosci (Landmark Ed) 15:25-34). Paradoxically, CPSA also activates innate immune responses by stimulating the production of anti-inflammatory cytokines, such as IL-10 (Eribo et al. (2021) The intestinal commensal, *Bacteroides fragilis*, modulates host responses to viral infection and therapy: Lessons for exploration during *Mycobacterium tuberculosis* infection. Infect Immun 90(1):e0032121; Tolonen et al. (2022) Synthetic glycans control gut microbiome structure and mitigate colitis in mice. Nature Communications 13(1):1244). The activation of anti-inflammatory cytokines has advanced CPSA as a potential therapeutic for inflammation-associated diseases such as multiple sclerosis, ulcerative colitis, viral encephalitis, and irritable bowel syndrome (Tolonen et al. (2022) Synthetic glycans control gut microbiome structure and mitigate colitis in mice. Nature Communications 13 (1): 1244; Zheng et al. (2020) Capsular Polysaccharide From *Bacteroides fragilis* Protects Against Ulcerative Colitis in an Undegraded Form. Frontiers in Pharmacology 11:570476; Ramakrishna et al. (2019) *Bacteroides fragilis* polysaccharide A induces IL-10 secreting B and T cells that prevent viral encephalitis. Nature Communications 10(1):2153; Nothaft et al. (2016) Engineering the *Campylobacter jejuni* N-glycan to create an effective chicken vaccine. Scientific Reports 6(1):26511.

[0102] CPSA, a zwitterionic polymer, is composed of a repeating tetrasaccharide consisting of [$\rightarrow 3$]- β -d-pyrGalp-(1 $\rightarrow 3$)- α -d-AATGalp-(1 $\rightarrow 4$)[β -d-Galf-(1 $\rightarrow 3$)]- α -d-

GalpNAc-(1 $\rightarrow$], where AAT-Gal is 2-acetamido-4-amino-2,4,6-trideoxygalactopyranose, and pyrGal is 4,6-O-pyruvate-galactopyranose (FIG. 1). See Baumann et al., 1992. It is assembled on a common lipid anchor, bactoprenyl phosphate (BP; also known as undecaprenyl phosphate), at the cytoplasmic face of the inner membrane. Biosynthesis begins with the addition phospho-AATGal to BP by the initiating phospho-glycosyltransferase WcfS to produce BPP-AATGal (FIG. 1). See Mostafavi & Troutman, 2013. The remaining sugars, galactose (Gal), N-acetylgalactosamine (GalNAc), and galactofuranose (Galf) are sequentially appended to BPP-AATGal by the glycosyltransferases WcfQ, WcfP, and WcfN, respectively (Sharma et al., 2017). The pyruvyltransferase, WcfO, pyruvylates galactose after the formation of BPP-AATGal-Gal, and before the additional of GalNAc (Sharma et al., 2017). WcfM is a UDP-galactopyranose mutase that produces UDP-galactofuranose, the final sugar in CPSA repeat unit biosynthesis.

[0103] As disclosed herein, a recombinant system for CPSA production in *E. coli* was developed by implementing a stepwise method and strategic incorporation of unrelated, robust glycan modification enzymes. In doing so, the production of each lipid linked intermediate was achieved, and CPSA polymers of biologically relevant length were produced. Ultimately, an effective approach was modeled to evaluate recombinant polysaccharide production, which may be applied to other glycan biosynthesis systems that are not confined within a single operon. These results could aid in identification of potential targets to increase recombinant polymer quantity and increase availability of biologically relevant glycans.

II.B. Compositions: Nucleic Acids

[0104] In some embodiments, the presently disclosed subject matter provides nucleic acids that encode one or more biosynthetic enzymes that are involved in polysaccharide production in bacteria. By way of example and not limitation, the presently disclosed subject matter provides nucleic acids that encode one or more polypeptides that can be employed to produce a capsular polysaccharide A (CPSA) in a bacterium. CPSA can be produced via the actions of various genes from *Bacteroides fragilis* (optionally *Bacteroides fragilis* NTCT 9343), and in some embodiments the presently disclosed nucleic acid molecules employ one or more of the genes found in *Bacteroides fragilis*. These genes include the wcfR aminotransferase, the wcfS phospho-glycosyltransferase, the wcfQ glycosyltransferase, the wcfO pyruvyltransferase, the wcfP glycosyltransferase, the wcfM UDP-galactopyranose mutase, and the wcfN glycosyltransferase gene products, the nucleic acid and amino acid sequences of which are known. Exemplary such sequences include SEQ ID NOs: 79 (nucleic acid) and 80 (amino acid for wcfR, SEQ ID NOs: 81 and 82 for wcfS, SEQ ID NOs: 83 and 84 for wcfQ, SEQ ID NOs: 85 and 86 for wcfO, SEQ ID NOs: 87 and 88 for wcfP, SEQ ID NOs: 89 and 90 for wcfM, and SEQ ID NOs: 91 and 92 for wcfN).

[0105] In some embodiments, coding sequences for one or more of these gene products is provided to a bacterium, in some embodiments as part of an expression cassette. In some embodiments, at least one of the coding sequences is derived from an organism other than the species of the bacterium in which the coding sequences are introduced (i.e., in some embodiments, at least one coding sequence is heterologous to the bacterium).

[0106] As disclosed herein, additional gene products can also be employed to increase the efficiency and/or yield of the polysaccharide of interest. Such additional gene products can include, but are not limited to a *pglF* dehydrogenase gene product (optionally, a *Campylobacter jejuni* *pglF* gene product), a *wbpP* UDP-N-acetyl-d-glucosamine C4 epimerase gene product (optionally, a *Vibrio vulnificus* *wbpP* gene product), a *wza* polysaccharide export protein gene product (optionally an *E. coli* gene product, further optionally encoded by SEQ ID NO: 93 and/or having the amino acid sequence set forth in SEQ ID NO: 94), a *wzx* fippase gene product (optionally a *Bacteroides fragilis* gene product, further optionally encoded by SEQ ID NO: 95 and/or having the amino acid sequence set forth in SEQ ID NO: 96), and/or a *wzy* polymerase gene product (optionally a *Bacteroides fragilis* gene product, further optionally encoded by SEQ ID NO: 97 and/or having the amino acid sequence set forth in SEQ ID NO: 98).

[0107] As such, in some embodiments the presently disclosed subject matter provides expression cassettes comprising a plurality of coding sequences derived from at least two different bacterial species, wherein the plurality of coding sequences are selected from the group consisting of a *pglF* dehydrogenase gene coding sequence, a *wbpP* UDP-N-acetyl-d-glucosamine C4 epimerase gene coding sequence, a *wcfR* aminotransferase gene coding sequence, a *wcfS* phospho-glycosyltransferase gene coding sequence, a *wcfQ* glycosyltransferase gene coding sequence, a *wcfO* pyruvyltransferase gene coding sequence, a *wcfP* glycosyltransferase gene coding sequence, a *wcfM* UDP-galactopyranose mutase gene coding sequence, a *wcfN* glycosyltransferase gene coding sequence, a *wza* polysaccharide export protein gene coding sequence, a *wzx* fippase gene coding sequence, a *wzy* polymerase gene coding sequence, and a *wzz* gene coding sequence.

[0108] It is understood that within the scope of the presently disclosed subject matter are expression cassettes that comprise fewer than the full complement of coding sequences listed above. By way of example and not limitation, in some embodiments the expression cassette comprises a combination of coding sequences selected from the group consisting of:

[0109] *pglF*, *wcfR*, and *wcfS*;

[0110] *pglF*, *wcfR*, *wcfS*, and *wcfQ*;

[0111] *pglF*, *wcfR*, *wcfS*, *wcfQ*, and *wcfO*;

[0112] *pglF*, *wcfR*, *wcfS*, *wcfQ*, *wcfO*, *wbpP*, and *wcfP*;

[0113] *pglF*, *wcfR*, *wcfS*, *wcfQ*, *wcfO*, *wbpP*, *wcfP*, *wcfM*, and *wcfN*; and

[0114] *pglF*, *wcfR*, *wcfS*, *wcfQ*, *wcfO*, *wbpP*, *wcfP*, *wcfM*, *wcfN*, *wzx*, and *wzy*.

[0115] In order to express the plurality of coding sequences, in some embodiments the expression cassette of the presently disclosed subject matter is operatively linked to a promoter such that transcription from the promoter results in the production of a polycistronic mRNA that is translatable to produce polypeptides encoded by the coding sequences. In some embodiments, the polycistronic mRNA molecule comprises a ribosome-binding site (RBS) operatively linked to a translation initiation codon of one or more of, optionally each of, the plurality of coding sequences, wherein at least one RBS is located 3' to a translation termination codon of a first coding sequence and 5' to a translation initiation codon of a second coding sequence, wherein the first coding sequence and the second coding

sequence are adjacent to each other on the polycistronic mRNA molecule, the first coding sequence being present 5' to the second coding sequence. In some embodiments, the RBS comprises the nucleic acid sequence 5'-AGGAGG'-3'. In some embodiments, the RBS is attached to a linker sequence to form an RBS domain, wherein the RBS domain is just 5' of the translation initiator sequence of a coding sequence present in the expression cassette. In some embodiments the RBS/linker sequence comprises, consists essentially of, or consists of the nucleotide sequence 5'-AGGAGGATATACAT (SEQ ID NO: 100), wherein nucleotides 1-6 are the RBS and nucleotides 7-14 are the linker sequence.

[0116] In some embodiments, the polycistronic mRNA includes a plurality of RBSs, one operatively linked to each coding sequence. The RBSs that are present in the expression cassette (and hence the mRNA transcribed therefrom), can be the same or different. In some embodiments, the RBS cassette of SEQ ID NO: 100 is present just upstream of the initiator codon of each of the coding sequences present in the expression cassette.

[0117] In some embodiments, the expression cassette is transcribed in a bacterium in order to express the one or more coding sequences present thereon. In some embodiments, the expression cassette is transcribed to produce a polycistronic mRNA. In some embodiments, the polycistronic mRNA molecule is operably linked to a promoter that is active in the bacterium, optionally wherein the promoter is an inducible promoter, optionally an isopropyl- β -D-thiogalactopyranoside (IPTG)-inducible promoter, further optionally wherein the promoter is a T5 promoter. An exemplary T5 promoter comprises the nucleotide sequence presented in SEQ ID NO: 99.

[0118] As disclosed herein, the expression cassettes of the presently disclosed subject matter can encode some or all of the proteins required to produce a capsular polysaccharide A (CPSA) in a bacterium. For example, if CPSA production is desired, the expression cassette can provide a full complement of genes for CPSA production, which in some embodiments can be each of *pglF*, *wcfR*, *wcfS*, *wcfQ*, *wcfO*, *wbpP*, *wcfP*, *wcfM*, *wcfN*, *wzx*, and *wzy*. Expressing each of—*pglF*, *wcfR*, *wcfS*, *wcfQ*, *wcfO*, *wbpP*, *wcfP*, *wcfM*, *wcfN*, *wzx*, and *wzy* in a bacterium can result in the bacterium producing CPSA. However, as shown in FIG. 1, an expression cassette that lacks a *wcfN* coding sequence would be expected to produce a trisaccharide of AT-Gal/pyruvylated Gal/GalNAc, whereas an expression cassette that lacks a *wcfN* coding sequence and a *wcfP* coding sequence would be expected to produce a disaccharide of AT-Gal/pyruvylated Gal. As such, the coding sequences that are present in the expression cassette can determine which polysaccharides are produced.

[0119] In some embodiments, an expression cassette of the presently disclosed subject matter is present in a vector, optionally an expression vector. Methods for generating vectors that comprise recombinant nucleic acids are known. See e.g., Sambrook & Russell (2001) *Molecular Cloning: a Laboratory Manual*, 3rd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, United States of America.

II.C. Recombinant Bacteria

[0120] In some embodiments, the presently disclosed subject matter relates to bacteria comprising the expression

cassettes disclosed herein. As such, in some embodiments the presently disclosed subject matter relates to bacteria that have been transformed to carry an expression cassette comprising a plurality of coding sequences, the plurality of coding sequences comprising nucleotide sequences that encode two or more of a pglF dehydrogenase gene product, a wbpP UDP-GlcNAc C4 epimerase gene product, a wcfR aminotransferase gene product, a wcfS phospho-glycosyltransferase gene product, a wcfQ glycosyltransferase gene product, a wcfO pyruvyltransferase gene product, a wcfP glycosyltransferase gene product, a wcfM UDP-galactopyranose mutase gene product, a wcfN glycosyltransferase gene product, a wza polysaccharide export protein gene product, a wxz flippase gene product, a wzy polymerase gene product, and a wzz gene product, wherein at least one of the plurality of coding sequences is heterologous to the bacterium.

[0121] Any bacterium for which production of recombinant polysaccharides would be of interest can be employed in the compositions and methods of the presently disclosed subject matter. In some embodiments, the bacterium is an *E. coli* bacterium.

[0122] The bacteria of the presently disclosed subject matter carry at least one gene that is heterologous to (i.e., derived from a bacterium of a different strain, species, or genus) than the bacterium itself. As such, the bacteria are designed to carry at least one coding sequence that is not naturally found in that bacterium. Thus, in some embodiments the bacterium of the presently disclosed subject matter is an *E. coli* bacterium, and at least one of the pglF gene product, the wbpP gene product, the wcfR gene product, the wcfS gene product, the wcfQ gene product, the wcfO gene product, the wcfP gene product, the wcfM gene product, the wcfN gene product, the wza gene product, the wxz flippase gene product, the wzy polymerase gene product, and/or the wzz gene product is not naturally occurring in the *E. coli* bacterium. By way of example and not limitation, in some embodiments, the pglF gene product is a *Campylobacter jejuni* gene product, the wbpP gene product is a *Vibrio vulnificus* gene product, the wcfR gene product is a *Bacteroides fragilis* gene product, the wcfS gene product is a *Bacteroides fragilis* gene product, the wcfQ gene product is a *Bacteroides fragilis* gene product, the wcfO gene product is a *Bacteroides fragilis* gene product, the wcfP gene product is a *Bacteroides fragilis* gene product, the wcfM gene product is a *Bacteroides fragilis* gene product, the wcfN gene product is a *Bacteroides fragilis* gene product, the wza gene product is an *Escherichia coli* gene product, the wxz flippase gene product is a *Bacteroides fragilis* gene product, the wzy polymerase gene product is a *Bacteroides fragilis* gene product, and/or the wzz gene product is a *Bacteroides fragilis* gene product.

[0123] In some embodiments, the plurality of coding sequences are transcribed in the bacterium to produce a polycistronic messenger RNA (mRNA) molecule in the bacterium. In some embodiments, the polycistronic mRNA molecule comprises a ribosome-binding site (RBS) operatively linked to a translation initiation codon of each of at least two of the plurality of coding sequences present therein. In some embodiments, the polycistronic mRNA molecule comprises a plurality of ribosome-binding sites (RBSs), with an RBS present 5' to each of at least two, optionally 5' to each, translation initiation codon present in the polycistronic mRNA, wherein at least one RBS is

located 3' to a translation termination codon of a first coding sequence and 5' to a translation initiation codon of a second coding sequence present on the polycistronic mRNA molecule.

[0124] In some embodiments, translation of the polycistronic mRNA molecule in the bacterium results in production of a polysaccharide in the bacterium, optionally wherein the polysaccharide is present in a lipid bilayer of the bacterium, further optionally wherein the lipid bilayer is part of the bacterium's cell wall. In some embodiments, the polysaccharide is a CPSA, optionally an aggregate of two or more CPSA tetrasaccharide monomers.

[0125] It is noted that the various biological activities disclosed herein can be provided recombinantly using the nucleic acids and expression cassettes of the presently disclosed subject matter. However, it is also understood that for those biological activities for which the recombinant bacterium is modified to produce but that the recombinant bacteria already possesses such an activity, the nucleic acids of the presently disclosed subject matter can optionally exclude the relevant coding sequences and instead rely on the endogenous expression of those activities by the bacterium. As such, for those activities that the bacterium would be expected to express in the absence of the compositions of the presently disclosed subject matter, it is not required that such an activity would need to be provided.

[0126] Thus, for example, if the bacterium already expresses a wza gene product, a wxz flippase gene product, a wzy polymerase, and/or a wzz gene product, the biological activity can be provided endogenously.

II.C. Methods of Producing Polysaccharides Using the Nucleic Acids of the Presently Disclosed Subject Matter

[0127] In some embodiments, the presently disclosed subject matter employs the nucleic acids, optionally the expression cassettes disclosed herein, in methods for producing polysaccharides in bacteris. In some embodiments, the methods comprise expressing in the bacterium a plurality of coding sequences, the plurality of coding sequences comprising nucleotide sequences that encode a pglF dehydrogenase gene product, a wbpP UDP-N-acetyl-d-glucosamine C4 epimerase gene product, a wcfR aminotransferase gene product, and a wcfS phospho-glycosyltransferase gene product, and optionally further comprise nucleotide sequences encoding one or more of a wcfQ glycosyltransferase gene product, a wcfO pyruvyltransferase gene product, a wcfP glycosyltransferase gene product, a wcfM UDP-galactopyranose mutase gene product, a wcfN glycosyltransferase gene product, a wza polysaccharide export protein gene product, a wxz flippase gene product, a wzy polymerase gene product, and a wzz gene product, and further wherein at least one of the coding sequences is heterologous to the bacterium. In some embodiments, the polysaccharide is a capsular polysaccharide A (CPSA) and the bacterium expresses each of the PglF gene product, the wbpP gene product, the WcfR gene product, the WcfS gene product, the WcfQ gene product, the WcfO gene product, the WcfP gene product, the WcfM gene product, the WcfN gene product, the wzz gene product, the wxz flippase gene product, and the wzy polymerase gene product, at least one of which is heterologous to the bacterium. In some embodiments, the bacterium is an *Escherichia coli* bacterium.

[0128] In some embodiments, the pglF gene product is a *Campylobacter jejuni* gene product; the wbpP gene product

is a *Vibrio vulnificus* gene product; the wcfR gene product is a *Bacteroides fragilis* gene product; the wcfS gene product is a *Bacteroides fragilis* gene product; the wcfQ gene product is a *Bacteroides fragilis* gene product; the wcfO gene product is a *Bacteroides fragilis* gene product; the wcfP gene product is a *Bacteroides fragilis* gene product; the wcfM gene product is a *Bacteroides fragilis* gene product; the wcfN gene product is a *Bacteroides fragilis* gene product; the wza gene product is an *Escherichia coli* gene product; the wzx fippase gene product is a *Bacteroides fragilis* gene product; the wzy polymerase gene product is a *Bacteroides fragilis* gene product; and/or the wzz gene product is a *Bacteroides fragilis* gene product.

[0129] In some embodiments, action of the coding sequences disclosed herein in the bacterium results in the presence of the polysaccharide is present in a lipid bilayer of the bacterium, optionally wherein the lipid bilayer is part of the bacterium's cell wall. In some embodiments, the bacterium does not express a functional waal O-antigen ligase gene product.

EXAMPLES

[0130] The following EXAMPLES provide illustrative embodiments. In light of the present disclosure and the general level of skill in the art, those of skill will appreciate that the following EXAMPLES are intended to be exemplary only and that numerous changes, modifications, and alterations can be employed without departing from the scope of the presently disclosed subject matter.

[0131] Without further description, it is believed that one of ordinary skill in the art can, using the preceding description and the following illustrative EXAMPLES, make and utilize the compounds of the presently disclosed subject matter and practice the methods of the presently disclosed subject matter. The following EXAMPLES therefore particularly point out embodiments of the presently disclosed

subject matter and are not to be construed as limiting in any way the remainder of the disclosure.

Materials and Methods for the EXAMPLES

[0132] General. Polyclonal anti-CPSA antibodies were generously provided by Dr. Laurie Comstock. All bacterial cultures were grown in Miller Lysogeny Broth unless otherwise specified. *E. coli* mutants and pQE-80L plasmid were generously provided by Dr. Matthew Jorgensen of the University of Arkansas Medical Center. LC-MS analysis was performed on an Agilent 1260 Infinity II equipped with a quaternary pump and MSD. All mobile phases were prepared from LC-MS grade reagents.

[0133] Construction of CPSA plasmids. CPSA plasmids were constructed using the primers listed in Table 1. Briefly, pQE80L vector was isolated from *E. coli* DH5a cells. The vector was digested with restriction enzymes BamHI and HindII and purified with gel electrophoresis. Gene inserts with amplified with Phusion DNA polymerase (New England Biolabs). Amplicons were purified using the WIZ-ARD® SV Gel and PCR Clean-Up System (Promega). Purified, digested vector and inserts were incubated with NEBuilder HiFi DNA Assembly Mix (New England Biolabs) according to the manufacturer's instructions and incubated at 50° C., or overnight for constructs with more than 4 fragments. *E. coli* DH5a cells were then chemically transformed with 5 µL of assembly reaction mixture and plated on selective media (LB/Carb100). Successful transformants were confirmed by colony PCR.

[0134] An artificial SacI site was incorporated into pBAS8 immediately following wcfS. For construction of pBAS9-19, SacI digested pBAS8 was used as the vector backbone. For construction of pBAS16, wcfO and wcfQ were amplified as a single insert using pBAS10 as a template. For construction of pBAS 17-19, six genes (wcfOQP_{Bf} wbpP_{Vv}, wcfMN) were amplified as a single fragment using pBAS16 as a template.

TABLE 1

Primers Employed to Recreate CPSA Biosynthesis in <i>E. coli</i>		
Plasmid	Genes (Source)	Primers*
pBAS1	pglF _{Cj} (<i>V. vulnificus</i> M06-24)	CATCACCATCACCATCAGACATGATTTTTATATAAGCAAAAGATT AGC (SEQ ID NO: 1)
		GGATTTGTTGGTATTTGGTCATATGTATATCCTCTTATACACCTTCT TTATTGTGTTTAAATTC (SEQ ID NO: 2)
		ATGACCAAATACCAACAAATCC (SEQ ID NO: 3)
	wbpP _{Vv} (<i>V. vulnificus</i> M06-24)	GTTATCTCTTTATTGATTGTCCCATATGTATATCCTCTCTATTTTATAA ATCGCACATACC (SEQ ID NO: 4)
		ATGGGACAATCAATAAGAATAAC (SEQ ID NO: 5)
		GAGTCCAAGCTCAGCTAATTACTAACGAATGATGCTCCAAAATG (SEQ ID NO: 6)
pBAS8	pglF _{Cj} (<i>C. jejuni</i> ATCC)	GGATCGCATCACCATCACCATCAGgaATGATTTTTATATAAGCAAA AGATTAGC (SEQ ID NO: 7)
		GGTATTTTCATATGTATATCCTCTTATACACCTTCTTTATTGTG (SEQ ID NO: 8)
		AGGAGGATATACATATGAAAATACCTTTTTCACCACC (SEQ ID NO: 9)
	wcfR _{Bf} (<i>B. fragilis</i> ATCC 9343)	CAAATGTATATCCTCTCTAGTGGTGGTGGTGGTGTCTCGTTTCT GCAATTACG (SEQ ID NO: 10)
		CCACTAGAGGAGGATATACATTTGATCCGTTTTTTTGATATCG (SEQ ID NO: 11)
		GAGTCCAAGCTCAGCTAATTAGAGCTCTAACGAATGATGCTCC (SEQ ID NO: 12)

TABLE 1-continued

Primers Employed to Recreate CPSA Biosynthesis in <i>E. coli</i>		
Plasmid	Genes (Source)	Primers*
pBAS9*	wcfQ _{Bf} (<i>B. fragilis</i> ATCC 9343)	GCATCATTCGTTAGGAGCTCAGGAGGATATACATATGAAATTGGCTG TCATTTC (SEQ ID NO: 13) CAAGCTCAGCTAATTAGAGCTTCACTTGTCGTCATCGTCTTTGTAGTC ACGTACTGTTTTATAGATTATG (SEQ ID NO: 14)
pBAS10*	wcfQ _{Bf} (<i>B. fragilis</i> ATCC 9343) wcfO _{Bf} (<i>B. fragilis</i> ATCC 9343)	GCATCATTCGTTAGGAGCTCAGGAGGATATACATATGAAATTGGCTG TCATTTC (SEQ ID NO: 5) CCTTCAACGTACTGTTTTATAGATT (SEQ ID NO: 16) AATCTATAAAACAGTACGTTGAAGGAGGATATACATATGAGGAAGA TATTATTAACATATGG (SEQ ID NO: 17) CAAGCTCAGCTAATTAGAGCTTCACTTGTCGTCATCGTCTTTGTAGTC TGACATCATAAATTTATTACATATATTAATTAATC (SEQ ID NO: 18)
pBAS11*	wcfQ _{Bf} (<i>B. fragilis</i> ATCC 9343) wcfO _{Bf} (<i>B. fragilis</i> ATCC 9343) wcfP _{Bf} (<i>B. fragilis</i> ATCC 9343)	GCATCATTCGTTAGGAGCTCAGGAGGATATACATATGAAATTGGCTG TCATTTC (SEQ ID NO: 19) CCTTCAACGTACTGTTTTATAGATT (SEQ ID NO: 20) AATCTATAAAACAGTACGTTGAAGGAGGATATACATATGAGGAAGA TATTATTAACATATGG (SEQ ID NO: 21) CTCATATGTATATCCTCCTTCATGACATCATAAATTTATTACATATAT TAATTAATC (SEQ ID NO: 22) CATGAAGGAGGATATACATATGAGGGATGGAAGGCC (SEQ ID NO: 23) CAAGCTCAGCTAATTAGAGCTTCACTTGTCGTCATCGTCTTTGTAGTC TTTATGTAAAAGATTTAAATATATTCTTTC (SEQ ID NO: 24)
pBAS12*	wcfQ _{Bf} (<i>B. fragilis</i> ATCC 9343) wcfO _{Bf} (<i>B. fragilis</i> ATCC 9343) wcfP _{Bf} (<i>B. fragilis</i> ATCC 9343) wcfM _{Bf} (<i>B. fragilis</i> ATCC 9343) wcfN _{Bf} (<i>B. fragilis</i> ATCC 9343)	GCATCATTCGTTAGGAGCTCAGGAGGATATACATATGAAATTGGCTG TCATTTC (SEQ ID NO: 25) CCTTCAACGTACTGTTTTATAGATT (SEQ ID NO: 26) AATCTATAAAACAGTACGTTGAAGGAGGATATACATATGAGGAAGA TATTATTAACATATGG (SEQ ID NO: 27) CTCATATGTATATCCTCCTTCATGACATCATAAATTTATTACATATAT TAATTAATC (SEQ ID NO: 28) CATGAAGGAGGATATACATATGAGGGATGGAAGGCC (SEQ ID NO: 29) CCTCCTTCATTTATGTAAAAGATTTAAATATATTCTTTCGAG (SEQ ID NO: 30) AAATCTTTTACATAAATGAAGGAGGATATACATATGAAAAAATAAT ATGACTATCTAATTG (SEQ ID NO: 31) CAACTACAGCAAATATCTTCATATGTATATCCTCCTTCATAAGTCACT ATTTATAACTTTTCCAC (SEQ ID NO: 32) ATGAAGATATTGCTGTAGTTG (SEQ ID NO: 33) CAAGCTCAGCTAATTAGAGCTTTACTTGTCGTCATCGTCTTTGTAGTC ATAGGTAGCTCCATTTTTTTTACG (SEQ ID NO: 34)
pBAS15*	wcfQ _{Bf} (<i>B. fragilis</i> ATCC 9343) wcfO _{Bf} (<i>B. fragilis</i> ATCC 9343) wcfP _{Bf} (<i>B. fragilis</i> ATCC 9343) wbpP _{Vv} (<i>V. vulnificus</i> M06-24)	GCATCATTCGTTAGGAGCTCAGGAGGATATACATATGAAATTGGCTG TCATTTC (SEQ ID NO: 35) CCTTCAACGTACTGTTTTATAGATT (SEQ ID NO: 36) AATCTATAAAACAGTACGTTGAAGGAGGATATACATATGAGGAAGA TATTATTAACATATGG (SEQ ID NO: 37) CTCATATGTATATCCTCCTTCATGACATCATAAATTTATTACATATAT TAATTAATC (SEQ ID NO: 38) CATGAAGGAGGATATACATATGAGGGATGGAAGGCC (SEQ ID NO: 39) GTAAAAGATTTAAATATATTCTTTCGAG (SEQ ID NO: 40) CTCGAAAGAATATATTTAAATCTTTTACATAAATGAAGGAGGATATA <u>CATATGACCAAATACGAAAAATCC</u> (SEQ ID NO: 41) CAAGCTCAGCTAATTAGAGCTTTATTTTTTATCATTATATAAGCTTAT ATACC (SEQ ID NO: 42)
pBAS16*	wcfQ _{O_{Bf}} (from pBAS10) wcfP _{Bf} (<i>B. fragilis</i> ATCC 9343) wbpP _{Vv} (<i>V. vulnificus</i> M06-24)	GCATCATTCGTTAGGAGCTCAGGAGGATATACATATGAAATTGGCTG TCATTTC (SEQ ID NO: 43) CTCATATGTATATCCTCCTTCATGACATCATAAATTTATTACATATAT TAATTAATC (SEQ ID NO: 44) CATGAAGGAGGATATACATATGAGGGATGGAAGGCC (SEQ ID NO: 45) GTAAAAGATTTAAATATATTCTTTCGAG (SEQ ID NO: 46) CTCGAAAGAATATATTTAAATCTTTTACATAAATGAAGGAGGATATA <u>CATATGACCAAATACGAAAAATCC</u> (SEQ ID NO: 47) CAAGCTCAGCTAATTAGAGCTTTATTTTTTATCATTATATAAGCTTAT ATACC (SEQ ID NO: 48)

TABLE 1-continued

Primers Employed to Recreate CPSA Biosynthesis in <i>E. coli</i>		
Plasmid	Genes (Source)	Primers*
	wcfM _{Bf} (<i>B. fragilis</i> ATCC 9343)	GCTTTATAAATGATAAAAAATAAGGAGGATATACATATGAAAAA AAATATGACTATCTAATG (SEQ ID NO: 49) CAACTACAGCAATATCTTCATATGTATATCCTCCTTCATAAGTCACT ATTTATAACTTTTCCAC (SEQ ID NO: 50) ATGAAGATATTGCTGTAGTTG (SEQ ID NO: 51) CAAGCTCAGCTAATTAGAGCTTTAATAGGTAGCTCCATTTTTTTTACG (SEQ ID NO: 52)
pBAS17*	wcfQOP_wbpP_ wcfMN (from pBAS16) wzx _{Bf} (<i>B. fragilis</i> ATCC 9343) wzy _{Bf} (<i>B. fragilis</i> ATCC 9343)	GCATCATTCGTTAGGAGCTCAGGAGGATATACATATGAAATTGGCTG TCATTTC (SEQ ID NO: 53) CCTCCTTAAATAGGTAGCTCCATTTTTTTTACG (SEQ ID NO: 54) TGGAGCTACCTATTAAAGGAGGATATACATATGGGACAATCAATAAA GAATAAC (SEQ ID NO: 55) CCTCCTTCATTGTTTAAATATTGAAAGTAATAATTTC (SEQ ID NO: 56) CAATATTTAAACAATGAAGGAGGATATACATATGACTAGTACTTCTT TCTTTATTATTAAG (SEQ ID NO: 57) AAGCTCAGCTAATTAGAGCTTTAAATTTGATTTTGGCATTAAC (SEQ ID NO: 58)
pBAS18*	wcfQOP_wbpP_ wcfMN (from pBAS16) wzx _{Bf} (<i>B. fragilis</i> ATCC 9343)	GCATCATTCGTTAGGAGCTCAGGAGGATATACATATGAAATTGGCTG TCATTTC (SEQ ID NO: 59) CCTCCTTAAATAGGTAGCTCCATTTTTTTTACG (SEQ ID NO: 60) TGGAGCTACCTATTAAAGGAGGATATACATATGGGACAATCAATAAA GAATAAC (SEQ ID NO: 61) CAAGCTCAGCTAATTAGAGCTTCATTGTTTAAATATTGAAAGTAATA ATTTC (SEQ ID NO: 62)
pBAS19*	wcfQOP_wbpP_ wcfMN (from pBAS16) wzy _{Bf} (<i>B. fragilis</i> ATCC 9343)	GCATCATTCGTTAGGAGCTCAGGAGGATATACATATGAAATTGGCTG TCATTTC (SEQ ID NO: 63) CCTCCTTAAATAGGTAGCTCCATTTTTTTTACG (SEQ ID NO: 64) TGGAGCTACCTATTAAAGGAGGATATACATATGGGACAATCAATAAA GAATAAC (SEQ ID NO: 65) CAAGCTCAGCTAATTAGAGCTTCATTGTTTAAATATTGAAAGTAATA ATTTC (SEQ ID NO: 66)

*Plasmids constructed using SacI-digested pBAS8 as the vector backbone. See Materials and Methods.

**Sequences are presented in the 5' to 3' direction.

[0135] Extraction of BPP-linked CPSA intermediates. Cultures for glycolipid extraction were prepared by inoculating a single colony into 5 mL of LB/Carb¹⁰⁰ and 2% glucose. Cultures were grown overnight for 16 hour at 37° C., 220 rpm. Cell cultures were then diluted 1:1000 into 5 mL of LB/Carb¹⁰⁰ and grown at 37° C., 220 rpm until reaching an OD₆₀₀ of approximately 0.6 before induction with 0.1 mM IPTG. After overnight induction, liquid cultures were transferred to falcon tubes and centrifuged at 5,000×g for 15 minutes at 4° C.

[0136] The supernatant was discarded, and the cell pellet was resuspended in 10 mM PBS. The cell suspension was then transferred to a 15 mL glass centrifuge tube. To the cell suspension, 1 mL of chloroform and 2 mL of methanol were added to create a single-phase solution of water: chloroform: methanol (0.8:1:2.0). Each sample was vortexed and incubated at room temperature for 20 minutes to ensure cell lysis. Resulting insoluble materials were clarified from the cell lysate solution by centrifugation at 2,500×g for 20 minutes at room temperature. The supernatant was then transferred to a clean glass culture tube, frozen at -80° C., then dried under vacuum. Dried samples were resuspended in a solution of 1 mM ammonium hydroxide: n-propanol (1:1).

[0137] LC-MS analysis of BPP-linked CPSA intermediates. To evaluate the presence of CPSA intermediates in *E. coli*, 10 µL of cell extract was injected and separated using

Waters XBridge BEH C18 column (5 µm, 4.6×100 mm, 300 Å). For LC-MS analysis, mobile phase A consisted of 0.1% ammonium hydroxide pH 11, and mobile phase B was n-propanol. BPP-linked intermediates were separated using a 5-75% gradient of mobile phase B over 12 minutes, then 75-95% B for 5 minutes, and an isocratic hold at 95% B for 5 minutes at 1mL/min for a total run time of 22 minutes. The column was equilibrated for 5 min at 5% B prior to the next injection. Intermediates were detected using selected ion monitoring (SIM) of the predicted [M-H]⁻ and/or [M-H]⁻² ion of each intermediate. Total ion chromatograms were collected for each injection. Controls were prepared from parent strains expressing empty plasmids and evaluated for CPSA intermediates or isobaric compounds (FIG. 10).

[0138] Whole-cell dot blots. Cultures were prepared from single colonies, induced with 0.1 mM IPTG at OD₆₀₀ nm, and grown overnight at 37° C. at 220 rpm. Whole-cell dot blots were prepared using 5 µL of liquid culture dried on nitrocellulose. After drying, the nitrocellulose was blocked with a 5% milk solution for 30 min. The nitrocellulose was then rinsed gently with deionized water (×3) and incubated at room temperature for 1 hour with 1 mL of an adsorbed CPSA antiserum diluted 1:30 in sterile 10 mM PBS. CPSA antiserum was provided by Dr. Laurie Comstock (see Comstock et al., 1999). After incubation with anti-CPSA, the nitrocellulose was washed twice for 5 min in 0.3% PBST, then placed in alkaline phosphatase conjugated anti-rabbit

goat secondary antibody (1:10,000) for 1 hour at room temperature. The nitrocellulose was then washed 3× for 1 minute with 0.3% PBST and developed with NBT-BCIP.

[0139] SDS-PAGE and LPS staining. To evaluate LPS and CPS profiles of *E. coli* expressing either pBAS16 or pBAS17, overnight cultures were diluted 1:100 in LB and incubated at 37° C. with shaking until reaching an OD of ~0.6. Cell cultures were then induced with 0.01 mM IPTG and incubated overnight at 37° C. Cell cultures were normalized to an OD₆₀₀ of 1.0, and a 0.5 mL volume of each sample was pelleted and resuspended in 100 µL of 1× SDS dye. Cell samples were then lysed at 95° C. for 10 min and cooled to room temperature before 20 µg of pronase (Sigma) was added. The samples were incubated at room temperature for 20 min prior to SDS-PAGE. A 5 µL sample of each lysate

TABLE 2

<i>E. coli</i> Strains Employed	
Strain	Genotype
DH5α	F- ϕ 80lacZΔM15 Δ(lacZYA-argF)U169 recA1 endA1 hsdR17(rK-, mK+) phoA supE44 λ-thi-1 gyrA96 relA1
MG1655	K-12 F ⁻ λ ⁻ ilvG ⁻ rfb-50 rph-1
DR35	Reference
BAS5	MG1655 ΔwzxB::frit
BAS13	MG1655 Δwza::frit
MAJ975	MG1655 ΔwaaL::frit
BAS24	MG1655 Δwza::fritΔwaaL::cam

TABLE 3

Primers Employed for <i>E. coli</i> Mutants			
Strain Designation	Primers	Template	Check Primers
BAS5	GAAAGGCTCTTTACGTTAGATGAGCTTAT CAGATTAAATTAATTGCATGACATTACA CGTCTTGAGCGAT (SEQ ID NO: 67) GCACAAACGGCACCAACAAACCAGAAC CAACAATGATATAATCGTACATATGAATA TCCTCCTTAGTTCC (SEQ ID NO: 68)	pKD4	TGGCTGCTATTGGGC GAA (SEQ ID NO: 71) TCCACCGATATGATT TCTTTTC (SEQ ID NO: 72)
BAS24	TCAACAGTCAAGCAGTTTGGAAAAGTTA TCATCATTATAAAGGTAAACTGAATATC CTCCTTAGTTCC (SEQ ID NO: 69) TTGTATAGATAAGAAGTGAGTTTAACTC ACTTCTTAACTTGTTTATTCATTGTGTAG GCTGGAGC (SEQ ID NO: 70)	pKD3	TATCCCAATGGCATC G (SEQ ID NO: 73) ACCTAATTCACGTA CTCC (SEQ ID NO: 74)

was added to individual wells and separated at 40 mA for 60 min. Gels were then stained with the PRO-Q™ Emerald 300 kit (Thermo Fisher Scientific) according to the manufacturer's instructions. Generation of *E. coli* mutants. *E. coli* mutants (Table 2) were either generously provided by the Jorgenson laboratory or prepared using the lambda red recombinase method of Datsenko & Wanner, 2000. Briefly, parent strains of *E. coli* were transformed with pKD46. Single colonies of transformants were used to inoculate 5 mL cultures containing LB/Carb¹⁰⁰ and incubated at 30° C. Cultures were induced with 100 mM L-arabinose and returned to the incubator for 1 hour at 30° C. Cells were then pelleted at 5,000 RCF at 4° C. and washed six times with ice-cold 10% glycerol. These cells were used immediately for transformation or stored as aliquots at -80° C. For transformation, approximately 1-2 µg of purified amplicon was added to cell suspensions. Cells were then electroporated at 2,500 V and immediately recovered in LB at 30° C. overnight. Recovered cells were plated on LB/Cam²⁰ to select for waaL mutants. Colony PCR was used to confirm the correct insertion of antibiotic resistance cassettes. Primers used to generate and confirm mutants are listed in Table 3. BAS13 was prepared by transforming DR35, a strain provided by the Young Laboratory, with pCP20 to remove the Kan resistance cassette (Ranjit & Young, 2016). To ensure that the Kan resistance cassette was successfully removed, cells were tested for sensitivity against Kan⁵⁰. To cure the cells of the pCP20 plasmid, cells were cultured on LB agar and incubated at 37° C. and colony purified twice more at 37° C. Plasmid curing was confirmed by testing cells for sensitivity to Carb¹⁰⁰.

[0140] Confocal Microscopy. Cultures of *E. coli* MG1655 pQE-80L and MG1655 pBAS17 were prepared from single colonies. Once cultures reached an OD_{600nm} of 0.6, they were induced with 0.1 mM IPTG for 2 hours. Cells were pelleted at 5,000×g for 15 minutes at 4° C. and washed twice with 5 mL of 10 mM PBS. Cells were then resuspended 100 µL of 10 mM PBS. Primary anti-CPSA antiserum was added to cell suspensions at a final concentration of 1:25, to make a final cell suspension of 100 µL. Cells were incubated with CPSA-antiserum at room temperature with rocking for 30 minutes. Cells were then pelleted at 10,000×g for 5 minutes and washed twice with 1 mL of PBS. Cells were resuspended in a 1:2,500 dilution of FITC conjugated secondary antibody and incubated at room temperature with rocking for 30 minutes. Cells were again pelleted at 10,000×g for 5 minutes and washed once with 10 mM PBS. Cells were resuspended in 100 µL of 10 mM PBS and 10 µL were dropped onto 0.7% agar pads prepared on a clean glass slide. Once cells were dried, a cover slip was added. Confocal microscopy was performed by Dr. Shreya Goyal and Dr. James Grissom.

Example 1

Construction of a CPSA Expression Plasmid

[0141] To produce CPSA in *E. coli*, we first sought to assemble the CPSA locus from *B. fragilis* into a plasmid. However, the CPSA locus does not contain all genes required for the assembly of UDP-AATGal and UDP-Gal-

NAC, which are also not native to our *E. coli* host strains. In *B. fragilis*, UDP-AATGal is formed through the activity of a dehydrogenase that produces a 4-keto sugar, and an aminotransferase that adds an amine to the 4-position to produce a sugar with a 4-aminogalactose configuration. UDP-GalNAC could be readily formed through the epimerization of the *E. coli* abundant NDP-sugar UDP-GlcNAC. To ensure all NDP-sugars required for CPSA biosynthesis were available, Gibson assembly was used to incorporate pglF, a dehydrogenase from *Campylobacter jejuni*, which catalyzes dehydration of the 6-position of UDP-GlcNAC and oxidation of the 4-position to give UDP-2-acetamido-2,6-dideoxy- α -D-xylo-hexos-4-ulose, a precursor of UDP-AATGal (Olivier et al., 2006; Mostafavi & Troutman, 2013; Riegert et al., 2017). UDP-AATGal is formed from the 4-keto sugar by the aminotransferase WcfR, a gene product from the CPSA locus (FIG. 2A; see also Mostafavi & Troutman, 2013). We also incorporated wbpP from *Vibrio vulnificus* M0-624, a UDP-GlcNAC C4 epimerase that produces UDP-GalNAC (FIG. 2B; see also Park et al., 2006; Cunneen et al., 2013). Previous work from our laboratory and others have shown that PglF and WbpP are readily overproduced and functional after purification from *E. coli* (Schoenhofen et al., 2006; Reid et al., 2023).

[0142] Implementation of recombinant glycan production in *E. coli* has been demonstrated through the complete transfer of the biosynthesis operon into a plasmid for expression (Wacker et al., 2002; Ma & Wang, 2019). While this can be robust, it does not allow for evaluation of individual steps in the biosynthetic pathway, and bottlenecks in production can be challenging to delineate. To evaluate potential bottlenecks that may hinder CPSA biosynthesis in *E. coli*, we generated an incremental set of plasmids designed to produce each individual BPP-linked CPSA intermediate in a stepwise manner (FIG. 1, Table 4). It is important to note that each gene in Gibson assembled plasmids was preceded by a strong *E. coli* ribosomal binding site (RBS) often used for over production of gene products. Assembly of each plasmid was confirmed by PCR.

TABLE 4

Description of Plasmids used to Generate Individual CPSA Intermediates in <i>E. coli</i>		
Plasmid designation	Genes incorporated	Predicted CPSA Intermediate
pQE	[none]	BP
pBAS8	pglF _{Cj} , wcfRS	BPP-AATGal
pBAS9	pglF _{Cj} , wcfRSQ	BPP-AATGal-Gal
pBAS10	pglF _{Cj} , wcfRSQO	BPP-AATGal-PyrGal
pBAS11	pglF _{Cj} , wcfRSQOP	BPP-AATGal-PyrGal
pBAS12	pglF _{Cj} , wcfRSQOPMN	BPP-AATGal-PyrGal
pBAS15	pglF _{Cj} , wbpP _{Vv} , wcfRSQOP	BPP-AATGal-PyrGal-GalNAC
pBAS16	pglF _{Cj} , wbpP _{Vv} , wcfRSQOPMN	BPP-AATGal-PyrGal-GalNAC-GalF

Example 2

Stepwise Analysis of CPSA Repeat Unit
Biosynthesis by ESI-LC-MS

[0143] To investigate production of lipid-linked CPSA intermediates, we utilized an established ESI-LC-MS based method for tracking glycan intermediates in *E. coli* (Reid et

al., 2021). Previously, we showed the precise order and function of enzymes involved in CPSA production in vitro, which enabled us to predict the m/z of BPP-linked CPSA intermediates based on which genes were expressed (Table 4; see Sharma et al., 2017). To ensure that each intermediate corresponded to the inclusion of the appropriate Wcf glycosyltransferase(s), we used selected ion monitoring (SIM) to analyze each cell lysate for the [M-H]⁻ and/or [M-H]²⁻ ion of (1) BP, (2) the expected BPP-linked intermediate, and (3) the subsequent predicted intermediate (FIG. 3). Spectra were collected from the total ion count (TIC) of each injection (FIG. 1). We chose to monitor for later intermediates in the pathway to confirm that CPSA intermediates were not produced without the incorporation of the appropriate genes. Intermediates only corresponding to what was predicted were identified in cell lysates (i.e., cells did not produce later intermediates until the appropriate genes were expressed). Additionally, we did not observe isobaric compounds with identical or similar retention times of CPSA intermediates in cells harboring an empty vector (FIG. 10). BPP-linked pyruvylated trisaccharide was not produced until wbpP_{Vv} was co-expressed with wcfP and preceding glycosyltransferases (FIG. 11), confirming that the C4 epimerase was critical for CPSA oligosaccharide production in *E. coli* (Park et al., 2006).

Example 3

[0144] Co-expression of wxz and wzy from *B. fragilis* Produces CPSA Polymers in *E. coli* Once we confirmed production of the CPSA repeat unit by our recombinant *E. coli* strain, we evaluated whether an *E. coli* flippase or polymerase could catalyze polymer formation or whether the CPSA specific flippase or polymerase from *B. fragilis* was required. To determine if both Wxz and Wzy were necessary for CPSA polymerization, we constructed plasmids containing only *B. fragilis* flippase wxz (pBAS18) or polymerase wzy (pBAS19), and *B. fragilis* wxz and wzy together (pBAS17), along with all other genes required for oligosaccharide biosynthesis (Table 2A). To determine whether CPSA polymers were formed, we evaluated cell lysates using SDS-PAGE and Western blotting with an anti-CPSA antibody serum (FIG. 4). See also Comstock et al., 1999.

TABLE 5

Plasmids used to Determine if Co-expression of wxz _{Bf} and wzy _{Bf} is Needed for Recombinant Production of CPSA Polymers in <i>E. coli</i>	
Plasmid designation	Genes included
pQE-80L (Empty Vector)	[none]
pBAS16	pglF _{Cj} , wbpP _{Vv} , wcfRSQOPMN
pBAS17	pglF _{Cj} , wbpP _{Vv} , wcfRSQOPMN, wxz, wzy
pBAS18	pglF _{Cj} , wbpP _{Vv} , wcfRSQOPMN, wxz
pBAS19	pglF _{Cj} , wbpP _{Vv} , wcfRSQOPMN, wzy

[0145] Polymerized CPSA with a molecular weight around 30 kDa was only observed in cells expressing pBAS17 (+wxz_{Bf} +wzy_{Bf}), and not in cells expressing pBAS16 (-wxz_{Bf} -wzy_{Bf}), pBAS18 (+wxz_{Bf}), or pBAS19 (+wzy_{Bf}). See FIG. 3. We conclude that CPSA is polymerized in *E. coli* only when wxz_{Bf} and wzy_{Bf} are co-expressed, or that polymers produced in other strains were below the detection limits of this assay.

Example 4

CPSA Oligo- and Polymers are Likely Ligated to Lipid A Core

[0146] We next sought to determine if CPSA could be detected on the surface of *E. coli* expressing pBAS17 (+wzx_{Bf} +wzy_{Bf}). See Table 4. For this and remaining experiments, we transitioned to *E. coli* MG1655 as the host due to the availability of a small library of relevant mutants. Expression of CPSA in *E. coli* MG1655 did not qualitatively differ from CPSA expression in DH5 α (FIG. 12). Using confocal microscopy, we confirmed that intact cells expressing pBAS17 reacted with anti-CPSA antiserum and FITC conjugated secondary antibodies, while cells expressing an empty plasmid did not (FIG. 5). Using a whole-cell dot blot, we confirmed that pBAS8-15 did not react with anti-CPSA antiserum either but were surprised to find that cells expressing pBAS16 did (FIG. 12, Table 4).

[0147] To better understand CPSA glycoforms present in wild-type MG1655, cell lysates were partitioned into organic, aqueous, and insoluble fractions using the Bligh and Dyer method of lipid extraction (Bligh & Dyer, 1959; Henderson et al., 2013). For cells harboring pBAS17, the aqueous fraction was expected to contain water-soluble polysaccharides, potentially including lipid-linked CPSA intermediates (Reid et al., 2021). The organic fraction, typically used for phospholipid isolation, might also contain early BPP-linked CPSA intermediates (Reid et al., 2021). The insoluble phase, largely consisting of LPS would contain LPS-CPSA conjugates (Henderson et al., 2013). Each of these fractions were analyzed by SDS-PAGE and western blotting (FIG. 6). Intriguingly, polymerized CPSA was observed in both the aqueous and insoluble cell fractions in wild-type MG1655 pBAS17, with lower molecular weight polymers observed in the aqueous fraction (~30 kDa) than the organic fraction (~40 kDa). Combined with the previous results of surface expressed CPSA (FIG. 5; see also FIG. 12), this strongly suggested that CPSA can be ligated to LPS.

[0148] In *E. coli*, it has been shown that oligo- and polysaccharides other than O-antigens (O-Ag) can be ligated to the lipid A core of LPS, including colanic acid, enterobacterial common antigen (ECA), and peptidoglycan repeat units (Marolda et al., 2006; Meredith et al., 2007; Maciejewska et al., 2020). Additionally, recent developments in recombinant bacterial glycan production demonstrate that non-native glycans may be ligated to the lipid A core (Nothhaft et al., 2016; Kay et al., 2022). These glycan modifications of LPS are primarily attributed to the promiscuity of the *E. coli* O-antigen ligase, WaaL, which appends O-antigen to the heptose residue of the lipid A core (Abeyrathne et al., 2005; Ruan et al., 2012). To evaluate whether WaaL contributes to the production of higher molecular weight CPSA in the insoluble fraction of cell lysates, pBAS17 was overexpressed in an MG1655 Δ waaL mutant. Analysis of MG1655 Δ waaL pBAS17 cell lysate fractions revealed CPSA only in the aqueous fraction of cell lysates (FIG. 6) indicating that a Δ waaL mutant is needed to procure solely aqueous CPSA polymer.

[0149] In consideration of these previous findings, we thought it possible that CPSA oligomers and polymers may be ligated to the lipid A core of *E. coli* cells producing the CPSA repeat unit (pBAS16; -wzx_{Bf} -wzy_{Bf}) or polymer (pBAS17; +wzx, +wzy_{Bf}), respectively. As demonstrated by a whole-cell dot blot, CPSA surface expression appears to be

disrupted in waaL mutant expressing pBAS16 (FIG. 7). Based on this finding, we proposed that CPSA may be ligated in oligomeric form to LPS and requires WaaL for ligation to the lipid A core. To confirm this, LPS profiles of pronase treated cell lysates were evaluated using SDS-PAGE and the PRO-Q™ Emerald 300 Lipopolysaccharide Gel Stain Kit stain (Pierce). In MG1655, O-antigen production is disrupted by an IS5 insertion into the wbbL region of the O-antigen operon (Liu & Reeves, 1994). Therefore, LPS profiles of this strain would contain only lipid A core. Fluorescence imaging of PRO-Q™ Emerald 300 stained gels revealed an additional band of a higher molecular weight than lipid A core in pronase treated lysates of cells expressing pBAS16, but not in a waaL mutant or those expressing a plasmid control (FIG. 7). We were unable to detect these species with anti-PSA antiserum, which may be due levels below our detection limit. We also tested whether inactivation of wzxB, the O-antigen flippase in *E. coli* disrupted surface expression of CPSA or altered LPS profiles in cells expressing pBAS16. Western blot signals for Δ wzxB MG1655 pBAS16 were attenuated compared to wild type cells, but still present, indicating that WzxB may, in part, have a role in CPSA oligomer transport and surface expression (FIG. 7). Deletion of *E. coli* polysaccharide export protein, wza, had no effect on surface expression or LPS profiles of cells expressing pBAS16 compared to wild type.

Example 5

Evaluation of Potential Mechanisms of CPSA Polymer Export

[0150] Polysaccharide assembly and export typically occurs via a Wzx/Wzy dependent pathway and its cognate Wza transporter, or via an ATP-binding cassette (ABC) transporter such as observed with LPS transport (Whitfield, 2006). Since our evidence suggests that CPSA may be ligated to LPS and that *B. fragilis* Wza was not included in our constructs, we considered both of these pathways in *E. coli* as possible means of CPSA polymer export. We also considered that CPSA oligomers are still present in cells expressing pBAS17 (FIG. 13), even though CPSA polymers are produced. Like cells expressing pBAS16, if CPSA polymers or oligomers are ligated to Lipid A core and transported across the membrane via the well-defined Lpt transport pathway, a waaL mutant would abolish CPSA surface expression. Conversely, if CPSA expression was dependent on Wza_{Ec} for export across the outer membrane in cells expressing pBAS17, a wza_{Ec} mutant would no longer exhibit surface expression of CPSA.

[0151] To test whether CPSA polymer export is dependent on *E. coli* Wza and/or WaaL, we transformed pBAS17 into MG1655 lacking either wza or waaL and evaluated CPSA surface expression using a whole-cell dot blot. Surprisingly, a positive, but not as robust signal for CPSA was observed in a Δ wza and a Δ waaL mutant, indicating that neither may be essential for CPSA export (FIG. 8). To assess whether CPSA polymers are appended to Lipid A core in MG1655 pBAS17, we performed an analysis of LPS profiles (FIG. 8). SDS-PAGE of pronase treated lysates and staining with PRO-Q™ Emerald 300 revealed the presence of glycoforms of much higher molecular weights (~30 kDa), like those observed with anti-CPSA western blotting, and a ladder-like banding pattern in MG1655 pBAS17 (FIGS. 4 and 6).

Polymers of slightly lower average molecular weight were observed in the *waaL* mutant when compared to the *wza* mutant or wild type, suggesting that the polymer is no longer associated with lipid A core when *waaL* is inactivated (FIG. 8). Consistent with our findings in cells expressing pBAS16 (–*wzx*, –*wzy*), a low molecular weight band consistent with an oligomer of CPSA appended to lipid A core is no longer apparent in the *waaL* mutant.

Discussion of the EXAMPLES

[0152] A thorough understanding of recombinant polysaccharide biosynthesis in *E. coli* and hindrances in production at early stages could bolster production and isolation of rare polysaccharides. Our method demonstrates a systematic approach for evaluating individual steps within recombinant *Wzx/Wzy* dependent polysaccharide systems (Wacker et al., 2002; Ma & Wang, 2019). The efficacy of this method was demonstrated by production of complete polysaccharide of biologically relevant length, and used as a model to address missing or poorly expressed requisites, such as the UDP-GlcNAc C4 epimerase and 4,6-dehydratase not present in the CPSA operon (Kreisman et al., 2007). Additionally, we show that we were able to overcome poorly expressed components (*B. fragilis* UngD2) in *E. coli* by utilizing a well-expressed homolog (Coyne et al., 2008; Mostafavi & Troutman, 2013). Identification of bottlenecks in recombinant systems such as these will provide targeted opportunities to increase polymer quantity.

[0153] Moreover, our efforts provided the opportunity to evaluate whether co-expression of *wzx_{Bf}* and *wzy_{Bf}* was necessary for the translocation and concomitant polymerization of the CPSA unit oligosaccharide in *E. coli*. We were surprised to find CPSA polymers could not be detected without the expression of both *wzx_{Bf}* and *wzy_{Bf}* (Merino et al., 2016). This supports a previously proposed model that *Wzx*, *Wzy*, and *Wza* form a multi-protein complex wherein the translocase flips the unit oligosaccharide into the periplasm where it is then immediately “passed off” to the cognate polymerase and transporter or ligase (e.g., *WaaL*; see Feldman et al., 1999; Marolda et al., 2006). Countering this idea, was the expression and purification of a *Wzy* from *E. coli* without its associated translocase, and the reconstitution of bacterial polysaccharide biosynthesis in vitro (Woodward et al., 2010). It would be interesting to determine whether *Wzx_{Bf}* and *Wzy_{Bf}* colocalize and function as a unit in *E. coli*.

[0154] In the same regard, we found that deletion of *wzx_B* resulted in a significant attenuation of surface expressed CPSA in cells expressing pBAS16 (–*wzx*, –*wzy*), with a simultaneous decrease in a signal in SDS-PAGE gels that we propose is lipid A core-linked CPSA oligomer (FIG. 7). In future studies, it would be interesting to see if introduction of *wzx_{Bf}* in this strain (pBAS19, +*wzx_{Bf}* only) reverses this effect. The fact that a Δ *wzx_B* mutant did not appear to affect LPS/CPSA profiles in size distribution or signal intensity in cells expressing pBAS17 (+*wzx_{Bf}*, +*wzy_{Bf}*) suggests both *Wzx_{Bf}* functionality in *E. coli* and its ability to compensate for the lack of *wzx_B* in the *WaaL*-CPSA pathway (FIG. 8). This also implies promiscuous specificity of *E. coli* *Wzx_B* since it can accept CPSA oligomers, which could prove useful in the design of complex polysaccharide production systems.

[0155] Noting the potential activity of both flippases, *Wzx_{B_{Ec}}* and *Wzx_{B_{Bf}}* we were surprised to find that BPP-

tetrasaccharide could still be detected in cells expressing pBAS17, albeit at a much lower intensity (+*wzx*, +*wzy*; FIG. 5). This points toward a potential inefficiency in the activity of *Wzy_{Bf}* polymerase in *E. coli* and will be a future point of focus for improving CPSA output. Perhaps, this is due to a lack of the *B. fragilis* CPS transporter proteins in our construct, *Wza*, *Wzz*, and *Wzc*, which are suspected to form an active transporting complex (reviewed by Cuthbertson et al., 2009). Interplay of translocase and chain length regulator (Marolda et al., 2006).

[0156] Irregular cell shapes were also noted in strains expressing CPSA plasmids (FIG. 5), as well as anomalous growth curves upon plasmid induction. Future work should be dedicated to incorporating assembled genes into the *E. coli* genome as a recombinant operon for stability. This could also include the use of various genotypes to determine which gene products in *E. coli* hinder production and isolation of recombinant polysaccharides, specifically those that utilize the same nucleotide sugars and lipid precursor, BP. We show here, that a Δ *waaL* mutant is likely required to procure aqueous, non-lipid A associated CPSA, and has been demonstrated elsewhere in recombinant polysaccharide production systems (Kay et al., 2016). Current studies in our laboratory have implemented a Δ *waaL* strain for CPSA isolation and quantification of CPSA production in *E. coli*. Inclusion of the native CPSA chain length regulator, *Wzz* which is not included in the *B. fragilis* CPSA operon, to increase the length and yield of CPSA should also be considered for future studies. Systematic evaluation of the use of different transcription promoters to enhance polysaccharide quantity, recently demonstrated by Kay et al., 2016 will also be considered (Kay et al., 2022).

[0157] As such, disclosed herein are recombinant systems for producing polysaccharides including but not limited to CPSA production in bacteria including, but not limited to *E. coli*, by implementing a stepwise method and strategic incorporation of unrelated, robust glycan modification enzymes. In doing so, the production of each lipid-linked intermediate was confirmed along with CPSA polymers of biologically relevant length. Ultimately, provided herein are effective approaches to evaluating recombinant polysaccharide production, which may be applied to other glycan biosynthesis systems that are not confined within a single operon. These results could aid in the identification of potential targets to increase recombinant polymer quantity and increase the availability of biologically relevant glycans.

[0158] Summarily, disclosed herein are recombinant systems for production of CPSA and other polysaccharides in *E. coli* by implementing a stepwise method and strategic incorporation of unrelated, robust glycan modification enzymes. In doing so, it was confirmed the production of each lipid-linked intermediate and finally, CPSA polymers of biologically relevant length. Ultimately, disclosed herein is an effective approach to evaluate recombinant polysaccharide production, which can be applied to other glycan biosynthesis systems that are not confined within a single operon. These results could aid in the identification of potential targets to increase recombinant polymer quantity and increase the availability of biologically relevant glycans.

REFERENCES

[0159] All references cited in the instant disclosure (indicated by bracketed numbers) as well as those listed below, including but not limited to all patents, patent applications

and publications thereof, scientific journal articles, and database entries (such as but not limited to UniProt, EMBL, and GENBANK® biosequence database entries and including all annotations available therein) are incorporated herein by reference in their entireties to the extent that they supplement, explain, provide a background for, and/or teach methodology, techniques, and/or compositions employed herein. The discussion of the references is intended merely to summarize the assertions made by their authors. No admission is made that any reference (or a portion of any reference) is relevant prior art. Applicants reserve the right to challenge the accuracy and pertinence of any cited reference.

[0160] Abeyrathne et al. (2005) Functional characterization of WaaL, a ligase associated with linking O-antigen polysaccharide to the core of *Pseudomonas aeruginosa* lipopolysaccharide. *Journal of Bacteriology* 187(9):3002-3012.

[0161] Baumann et al. (1992) Structural elucidation of two capsular polysaccharides from one strain of *Bacteroides fragilis* using high-resolution NMR spectroscopy. *Biochemistry* 31(16):4081-4089.

[0162] Bligh & Dyer (1959) A rapid method of total lipid extraction and purification. *Canadian Journal of Biochemistry and Physiology* 37(8):911-917.

[0163] Brubaker et al. (1999) Mitogenic Activity of Purified Capsular Polysaccharide A from *Bacteroides fragilis*: Differential Stimulatory Effect on Mouse and Rat Lymphocytes In Vitro. *The Journal of Immunology* 162(4):2235-2242.

[0164] Choi et al. (2002) A Unique Structural Pattern Shared by T-Cell-Activating and Abscess-Regulating Zwitterionic Polysaccharides. *Biochemistry* 41(51):15144-15151.

[0165] Cobb et al. (2004) Polysaccharide Processing and Presentation by the MHCII Pathway. *Cell* 117(5):677-687.

[0166] Comstock et al. (1999) Analysis of a capsular polysaccharide biosynthesis locus of *Bacteroides fragilis*. *Infection and Immunity* 67(7):3525-3532.

[0167] Coyne et al. (2008) Role of glycan synthesis in colonization of the mammalian gut by the bacterial symbiont *Bacteroides fragilis*. *Proceedings of the National Academy of Sciences USA* 105(35):13099-13104.

[0168] Cunneen et al. (2013) Biosynthesis of UDP-GlcNAc, UndPP-GlcNAc and UDP-GlcNAcA Involves Three Easily Distinguished 4-Epimerase Enzymes, Gne, Gnu and GnaB. *PLOS One* 8(6):e67646.

[0169] Cuthbertson et al. (2009) Pivotal roles of the outer membrane polysaccharide export and polysaccharide copolymerase protein families in export of extracellular polysaccharides in gram-negative bacteria. *Microbiology and Molecular Biology Reviews* 73(1):155-177.

[0170] Datsenko & Wanner (2000) One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proceedings of the National Academy of Sciences USA* 97(12):6640-6645.

[0171] De Silva et al. (2009) Immunological Response from an Entirely Carbohydrate Antigen: Design of Synthetic Vaccines Based on Tn-PS A1 Conjugates. *Journal of the American Chemical Society* 131(28):9622-9623.

[0172] Feldman et al. (1999) The activity of a putative polysaccharide-linked sugar translocase (Wzx) involved in

Escherichia coli O antigen assembly is independent of the chemical structure of the O repeat. *J Biol Chem* 274(49):35129-38.

[0173] Feldman et al. (2005) Engineering N-linked protein glycosylation with diverse O antigen lipopolysaccharide structures in *Escherichia coli*. *Proceedings of the National Academy of Sciences USA* 102(8):3016-3021.

[0174] Henderson et al. (2013) Isolation and chemical characterization of lipid A from gram-negative bacteria. *Journal of Visualized Experiments* (79):e50623.

[0175] Kalka-Moll et al. (2002) Zwitterionic Polysaccharides Stimulate T Cells by MHC Class II-Dependent Interactions. *The Journal of Immunology* 169(11):6149-6153.

[0176] Kay et al. (2016) Recombinant expression of *Streptococcus pneumoniae* capsular polysaccharides in *Escherichia coli*. *Open Biology* 6(4):150243-150243.

[0177] Kay et al. (2022) Engineering a suite of *E. coli* strains for enhanced expression of bacterial polysaccharides and glycoconjugate vaccines. *Microb Cell Fact* 21(1) 66.

[0178] Kreisman et al. (2007) Structure and function relations with a T-cell-activating polysaccharide antigen using circular dichroism. *Glycobiology* 17(1):46-55.

[0179] Liu & Reeves (1994) *Escherichia coli* K12 regains its O antigen. *Microbiology* 140(1):49-57.

[0180] Ma & Wang (2019) RecET Direct Cloning of Polysaccharide Gene Cluster from Gram-Negative Bacteria. In *Bacterial Polysaccharides: Methods and Protocols* (Brockhausen (ed)). Springer Science+Business Media, New York, New York. pages 15-23.

[0181] Maciejewska et al. (2020) Lipopolysaccharide-Linked Enterobacterial Common Antigen (ECA (LPS)) Occurs in Rough Strains of *Escherichia coli* R1, R2, and R4. *International Journal of Molecular Sciences* 21(17):6038.

[0182] Marolda et al. (2006) Interplay of the Wzx translocase and the corresponding polymerase and chain length regulator proteins in the translocation and periplasmic assembly of lipopolysaccharide O antigen. *Journal of Bacteriology* 188(14):5124-5135.

[0183] Meredith et al. (2007) Modification of Lipopolysaccharide with Colanic Acid (M-antigen) Repeats in *Escherichia coli*. *Journal of Biological Chemistry* 282(11):7790-7798.

[0184] Merino et al. (2016) The first sugar of the repeat units is essential for the Wzy polymerase activity and elongation of the O-antigen lipopolysaccharide. *Future Microbiology* 11(7):903-918.

[0185] Mostafavi & Troutman (2013) Biosynthetic Assembly of the *Bacteroides fragilis* Capsular Polysaccharide A Precursor Bactoprenyl Diphosphate-Linked Acetamido-4-amino-6-deoxygalactopyranose. *Biochemistry* 52(11):1939-1949.

[0186] Nothaft et al. (2016) Engineering the *Campylobacter jejuni* N-glycan to create an effective chicken vaccine. *Scientific Reports* 6(1):26511.

[0187] Ochoa-Reparaz et al. (2010a) A polysaccharide from the human commensal *Bacteroides fragilis* protects against CNS demyelinating disease. *Mucosal Immunol* 3(5):487-495.

[0188] Ochoa-Repáraz et al. (2010b) Central nervous system demyelinating disease protection by the human commensal *Bacteroides fragilis* depends on polysaccharide A expression. *J Immunol* 2010, 185(7):4101-4108.

- [0189] Olivier et al. (2006) In vitro biosynthesis of UDP-N,N'-diacetyl bacillosamine by enzymes of the *Campylobacter jejuni* general protein glycosylation system. *Biochemistry* 45(45):13659-69.
- [0190] Park et al. (2006) Identification of the *Vibrio vulnificus* wbpP gene and evaluation of its role in virulence. *Infection and Immunity* 74(1) 721-728.
- [0191] Ramakrishna et al. (2019) *Bacteroides fragilis* polysaccharide A induces IL-10 secreting B and T cells that prevent viral encephalitis. *Nature Communications* 10(1): 2153.
- [0192] Ranjit & Young (2016) Colanic Acid Intermediates Prevent De Novo Shape Recovery of *Escherichia coli* Spheroplasts, Calling into Question Biological Roles Previously Attributed to Colanic Acid. *J Bacteriol* 198(8):1230-1240.
- [0193] Reid et al. (2021) Tracking Colanic Acid Repeat Unit Formation from Stepwise Biosynthesis Inactivation in *Escherichia coli*. *Biochemistry* 60(27):2221-2230.
- [0194] Reid et al. (2023) Chemoenzymatic Preparation of a *Campylobacter jejuni* Lipid-Linked Heptasaccharide on an Azide-Linked Polyisoprenoid. *ACS Omega* 8(17):15790-15798.
- [0195] Riegert et al. (2017) Structural and Biochemical Investigation of PglF from *Campylobacter jejuni* Reveals a New Mechanism for a Member of the Short Chain Dehydrogenase/Reductase Superfamily. *Biochemistry* 56(45): 6030-6040.
- [0196] Ruan et al. (2012) The WaaL O-antigen lipopolysaccharide ligase has features in common with metal ion-independent inverting glycosyltransferases. *Glycobiology* 22(2):288-299.
- [0197] Schoenhofen et al. (2006) Functional characterization of dehydratase/aminotransferase pairs from *Helicobacter* and *Campylobacter*: enzymes distinguishing the pseudaminic acid and bacillosamine biosynthetic pathways. *Journal of Biological Chemistry* 281(2):723-732.
- [0198] Sharma et al. (2017) Complete Tetrasaccharide Repeat Unit Biosynthesis of the Immunomodulatory *Bacteroides fragilis* Capsular Polysaccharide A. *ACS Chem Biol* 12(1):92-101.
- [0199] Tzianabos et al. (2000) T Cells Activated by Zwitterionic Molecules Prevent Abscesses Induced by Pathogenic Bacteria. *Journal of Biological Chemistry* 275(10): 6733-6740.
- [0200] Tzianabos et al. (2001) Structural rationale for the modulation of abscess formation by *Staphylococcus aureus* capsular polysaccharides. *Proceedings of the National Academy of Sciences USA* 98(16):9365-9370.
- [0201] Wacker et al. (2002) N-linked glycosylation in *Campylobacter jejuni* and its functional transfer into *E. coli*. *Science* 298(5599):1790-1793.
- [0202] Wang et al. (2000) Structural basis of the abscess-modulating polysaccharide A2 from *Bacteroides fragilis*. *Proceedings of the National Academy of Sciences USA* 97(25):13478.
- [0203] Whitfield (2006) Biosynthesis and assembly of capsular polysaccharides in *Escherichia coli*. *Annu Rev Biochem* 75:39-68.
- [0204] Woodward et al. (2010) In vitro bacterial polysaccharide biosynthesis: defining the functions of Wzy and Wzz. *Nature Chemical Biology* 6(6):418-423.
- [0205] Zheng et al. (2020) Capsular Polysaccharide From *Bacteroides fragilis* Protects Against Ulcerative Colitis in an Undegraded Form. *Frontiers in Pharmacology* 11:570476.
- [0206] Zhou et al. (2022) Antigen presenting cell response to polysaccharide A is characterized by the generation of anti-inflammatory macrophages. *Glycobiology* 32(2):136-147.
- [0207] It will be understood that various details of the presently disclosed subject matter can be changed without departing from the scope of the presently disclosed subject matter. Furthermore, the foregoing description is for the purpose of illustration only, and not for the purpose of limitation.

SEQUENCE LISTING

```

Sequence total quantity: 101
SEQ ID NO: 1          moltype = DNA length = 50
FEATURE              Location/Qualifiers
source                1..50
                     mol_type = other DNA
                     organism = synthetic construct

SEQUENCE: 1
catcaccatc accatcacga catgattttt tataaaagca aaagattagc          50

SEQ ID NO: 2          moltype = DNA length = 66
FEATURE              Location/Qualifiers
source                1..66
                     mol_type = other DNA
                     organism = synthetic construct

SEQUENCE: 2
ggatttggtg gtatttggtc atatgtatat cctcctttat acaccttctt tattgtgttt 60
aaattc                                                         66

SEQ ID NO: 3          moltype = DNA length = 22
FEATURE              Location/Qualifiers
source                1..22
                     mol_type = other DNA
                     organism = synthetic construct

SEQUENCE: 3
atgaccaa at accaacaat cc          22

SEQ ID NO: 4          moltype = DNA length = 61

```

-continued

FEATURE	Location/Qualifiers	
source	1..61	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 4		
gttattcttt attgattgtc ccatatgtat atcctcctct attttataaa tcgcacatac		60
c		61
SEQ ID NO: 5	moltype = DNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 5		
atgggacaat caataaagaa taac		24
SEQ ID NO: 6	moltype = DNA length = 44	
FEATURE	Location/Qualifiers	
source	1..44	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 6		
gagtccaagc tcagctaatt actaacgaat gatgctccaa aatg		44
SEQ ID NO: 7	moltype = DNA length = 56	
FEATURE	Location/Qualifiers	
source	1..56	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 7		
ggatcgcatc accatcacca tcacggaatg atttttata aaagcaaaag attagc		56
SEQ ID NO: 8	moltype = DNA length = 46	
FEATURE	Location/Qualifiers	
source	1..46	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 8		
ggtattttca tatgtatatc ctctttata cactttcttt attgtg		46
SEQ ID NO: 9	moltype = DNA length = 37	
FEATURE	Location/Qualifiers	
source	1..37	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 9		
aggaggatat acatatgaaa ataccttttt caccacc		37
SEQ ID NO: 10	moltype = DNA length = 57	
FEATURE	Location/Qualifiers	
source	1..57	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 10		
caaatgtata tcctctctta gtggtgggtg tggtgggtgc ggttttctgc aattacg		57
SEQ ID NO: 11	moltype = DNA length = 43	
FEATURE	Location/Qualifiers	
source	1..43	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 11		
ccactagagg aggatatata tttgatccgt ttttttgata tcg		43
SEQ ID NO: 12	moltype = DNA length = 44	
FEATURE	Location/Qualifiers	
source	1..44	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 12		
gagtccaagc tcagctaatt agagctccta acgaatgatg ctcc		44
SEQ ID NO: 13	moltype = DNA length = 54	
FEATURE	Location/Qualifiers	
source	1..54	
	mol_type = other DNA	

-continued

SEQUENCE: 13	organism = synthetic construct	
gcacattcgc ttaggagctc aggaggatat acatatgaaa ttggctgtca ttcc		54
SEQ ID NO: 14	moltype = DNA length = 70	
FEATURE	Location/Qualifiers	
source	1..70	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 14		
caagctcagc taattagagc ttcaattgtc gtcacgtctt ttgtagtcac gtactgtttt		60
atagattatg		70
SEQ ID NO: 15	moltype = DNA length = 54	
FEATURE	Location/Qualifiers	
source	1..54	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 15		
gcacattcgc ttaggagctc aggaggatat acatatgaaa ttggctgtca ttcc		54
SEQ ID NO: 16	moltype = DNA length = 25	
FEATURE	Location/Qualifiers	
source	1..25	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 16		
ccttcaacgt actgttttat agatt		25
SEQ ID NO: 17	moltype = DNA length = 62	
FEATURE	Location/Qualifiers	
source	1..62	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 17		
aatctataaa acagtcaggt gaaggaggat atacatatga ggaagatatt attaacatat		60
gg		62
SEQ ID NO: 18	moltype = DNA length = 83	
FEATURE	Location/Qualifiers	
source	1..83	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 18		
caagctcagc taattagagc ttcaattgtc gtcacgtctt ttgtagtcac acatcataaa		60
tttattacat atattaatta atc		83
SEQ ID NO: 19	moltype = DNA length = 54	
FEATURE	Location/Qualifiers	
source	1..54	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 19		
gcacattcgc ttaggagctc aggaggatat acatatgaaa ttggctgtca ttcc		54
SEQ ID NO: 20	moltype = DNA length = 25	
FEATURE	Location/Qualifiers	
source	1..25	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 20		
ccttcaacgt actgttttat agatt		25
SEQ ID NO: 21	moltype = DNA length = 62	
FEATURE	Location/Qualifiers	
source	1..62	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 21		
aatctataaa acagtcaggt gaaggaggat atacatatga ggaagatatt attaacatat		60
gg		62
SEQ ID NO: 22	moltype = DNA length = 57	
FEATURE	Location/Qualifiers	
source	1..57	
	mol_type = other DNA	

-continued

```

                                organism = synthetic construct
SEQUENCE: 22
ctcatatgta tatcctcctt catgacatca taaatttatt acatatatta attaatac 57

SEQ ID NO: 23      moltype = DNA length = 36
FEATURE           Location/Qualifiers
source            1..36
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 23
catgaaggag gatatacata tgagggatgg aaagcc 36

SEQ ID NO: 24      moltype = DNA length = 78
FEATURE           Location/Qualifiers
source            1..78
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 24
caagctcagc taattagagc ttcacttgtc gtcacgtctt ttgtagtctt tatgtaaaag 60
atttaaatat attctttc 78

SEQ ID NO: 25      moltype = DNA length = 54
FEATURE           Location/Qualifiers
source            1..54
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 25
gcatcattcg ttaggagctc aggaggatat acatatgaaa ttggtgtca tttc 54

SEQ ID NO: 26      moltype = DNA length = 25
FEATURE           Location/Qualifiers
source            1..25
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 26
ccttcaacgt actgttttat agatt 25

SEQ ID NO: 27      moltype = DNA length = 62
FEATURE           Location/Qualifiers
source            1..62
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 27
aatctataaa acagtacgtt gaaggaggat atacatatga ggaagatatt attaacatat 60
gg 62

SEQ ID NO: 28      moltype = DNA length = 57
FEATURE           Location/Qualifiers
source            1..57
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 28
ctcatatgta tatcctcctt catgacatca taaatttatt acatatatta attaatac 57

SEQ ID NO: 29      moltype = DNA length = 36
FEATURE           Location/Qualifiers
source            1..36
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 29
catgaaggag gatatacata tgagggatgg aaagcc 36

SEQ ID NO: 30      moltype = DNA length = 42
FEATURE           Location/Qualifiers
source            1..42
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 30
cttccttcac ttatgtaaaa gatttaaata tattctttcg ag 42

SEQ ID NO: 31      moltype = DNA length = 61
FEATURE           Location/Qualifiers
source            1..61
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 31

```

-continued

```

aaatctttta cataaatgaa ggaggatata catatgaaaa aaaaatatga ctatctaatt 60
g                                                                    61

SEQ ID NO: 32      moltype = DNA length = 66
FEATURE           Location/Qualifiers
source            1..66
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 32
caactacagc aaatatcttc atatgtatat cctccttcac aagtcactat ttataacttt 60
ttccac                                                                    66

SEQ ID NO: 33      moltype = DNA length = 22
FEATURE           Location/Qualifiers
source            1..22
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 33
atgaagatat ttgctgtagt tg                                                                    22

SEQ ID NO: 34      moltype = DNA length = 72
FEATURE           Location/Qualifiers
source            1..72
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 34
caagctcagc taattagagc tttacttgtc gtcacgtctt ttgtagtcat aggtagctcc 60
atttttttta cg                                                                    72

SEQ ID NO: 35      moltype = DNA length = 54
FEATURE           Location/Qualifiers
source            1..54
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 35
gcatcattcg ttaggagctc aggaggatat acatatgaaa ttggctgtca tttc      54

SEQ ID NO: 36      moltype = DNA length = 25
FEATURE           Location/Qualifiers
source            1..25
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 36
ccttcaacgt actgttttat agatt                                                                    25

SEQ ID NO: 37      moltype = DNA length = 62
FEATURE           Location/Qualifiers
source            1..62
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 37
aatctataaa acagtacgtt gaaggaggat atacatatga ggaagatatt attaacatat 60
gg                                                                    62

SEQ ID NO: 38      moltype = DNA length = 57
FEATURE           Location/Qualifiers
source            1..57
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 38
ctcatatgta tatcctcctt catgacatca taaatttatt acatatatta attaatac 57

SEQ ID NO: 39      moltype = DNA length = 36
FEATURE           Location/Qualifiers
source            1..36
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 39
catgaaggag gatatacata tgagggatgg aaagcc                                                                    36

SEQ ID NO: 40      moltype = DNA length = 28
FEATURE           Location/Qualifiers
source            1..28
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 40

```

-continued

gtaaaagatt taaatatatt ctttcgag	28
SEQ ID NO: 41	moltype = DNA length = 72
FEATURE	Location/Qualifiers
source	1..72
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 41	
ctcgaaagaa tatatttaaa tcttttacat aaatgaagga ggatatacat atgaccaa	60
acgaaaaaat cc	72
SEQ ID NO: 42	moltype = DNA length = 53
FEATURE	Location/Qualifiers
source	1..53
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 42	
caagctcagc taattagagc tttatTTTT atcatttata aagcttatat acc	53
SEQ ID NO: 43	moltype = DNA length = 54
FEATURE	Location/Qualifiers
source	1..54
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 43	
gcatcattcg ttaggagctc agggagatat acatatgaaa ttggctgtca ttcc	54
SEQ ID NO: 44	moltype = DNA length = 57
FEATURE	Location/Qualifiers
source	1..57
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 44	
ctcatatgta tatcctcctt catgacatca taaatttatt acatatatta attaatc	57
SEQ ID NO: 45	moltype = DNA length = 36
FEATURE	Location/Qualifiers
source	1..36
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 45	
catgaaggag gatatacata tgagggatgg aaagcc	36
SEQ ID NO: 46	moltype = DNA length = 28
FEATURE	Location/Qualifiers
source	1..28
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 46	
gtaaaagatt taaatatatt ctttcgag	28
SEQ ID NO: 47	moltype = DNA length = 72
FEATURE	Location/Qualifiers
source	1..72
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 47	
ctcgaaagaa tatatttaaa tcttttacat aaatgaagga ggatatacat atgaccaa	60
acgaaaaaat cc	72
SEQ ID NO: 48	moltype = DNA length = 53
FEATURE	Location/Qualifiers
source	1..53
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 48	
caagctcagc taattagagc tttatTTTT atcatttata aagcttatat acc	53
SEQ ID NO: 49	moltype = DNA length = 65
FEATURE	Location/Qualifiers
source	1..65
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 49	
gctttataaa tgataaaaaa taaaggagga tatacatatg aaaaaaaaa atgactatct	60
aattg	65

-continued

SEQ ID NO: 50 moltype = DNA length = 66
FEATURE Location/Qualifiers
source 1..66
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 50
caactacagc aaatatcttc atatgtatat cctccttcat aagtcactat ttataacttt 60
ttccac 66

SEQ ID NO: 51 moltype = DNA length = 22
FEATURE Location/Qualifiers
source 1..22
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 51
atgaagatat ttgctgtagt tg 22

SEQ ID NO: 52 moltype = DNA length = 48
FEATURE Location/Qualifiers
source 1..48
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 52
caagctcagc taattagagc tttaataggt agctccattt tttttacg 48

SEQ ID NO: 53 moltype = DNA length = 54
FEATURE Location/Qualifiers
source 1..54
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 53
gcatcattcg ttaggagctc aggaggatat acatatgaaa ttggctgtca tttc 54

SEQ ID NO: 54 moltype = DNA length = 33
FEATURE Location/Qualifiers
source 1..33
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 54
cctcctttaa taggtagctc catttttttt acg 33

SEQ ID NO: 55 moltype = DNA length = 54
FEATURE Location/Qualifiers
source 1..54
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 55
tggagctacc tattaaagga ggatatacat atgggacaat caataaagaa taac 54

SEQ ID NO: 56 moltype = DNA length = 37
FEATURE Location/Qualifiers
source 1..37
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 56
cctccttcat tgtttaaata ttgaaagtaa taatttc 37

SEQ ID NO: 57 moltype = DNA length = 61
FEATURE Location/Qualifiers
source 1..61
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 57
caatatattaa acaatgaagg aggatataca tatgactagt acttctttct ttattattaa 60
g 61

SEQ ID NO: 58 moltype = DNA length = 45
FEATURE Location/Qualifiers
source 1..45
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 58
aagctcagct aattagagct ttaaaatttg attttggcat taaac 45

SEQ ID NO: 59 moltype = DNA length = 54

-continued

FEATURE	Location/Qualifiers	
source	1..54	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 59		
gcatcattcg ttaggagctc aggaggatat acatatgaaa ttggctgtca tttc		54
SEQ ID NO: 60	moltype = DNA length = 33	
FEATURE	Location/Qualifiers	
source	1..33	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 60		
cctcctttaa taggtagctc catttttttt acg		33
SEQ ID NO: 61	moltype = DNA length = 54	
FEATURE	Location/Qualifiers	
source	1..54	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 61		
tggagctacc tattaagga ggatatacat atgggacaat caataagaa taac		54
SEQ ID NO: 62	moltype = DNA length = 52	
FEATURE	Location/Qualifiers	
source	1..52	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 62		
caagctcagc taattagagc ttcattgttt aaatattgaa agtaataatt tc		52
SEQ ID NO: 63	moltype = DNA length = 54	
FEATURE	Location/Qualifiers	
source	1..54	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 63		
gcatcattcg ttaggagctc aggaggatat acatatgaaa ttggctgtca tttc		54
SEQ ID NO: 64	moltype = DNA length = 33	
FEATURE	Location/Qualifiers	
source	1..33	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 64		
cctcctttaa taggtagctc catttttttt acg		33
SEQ ID NO: 65	moltype = DNA length = 54	
FEATURE	Location/Qualifiers	
source	1..54	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 65		
tggagctacc tattaagga ggatatacat atgggacaat caataagaa taac		54
SEQ ID NO: 66	moltype = DNA length = 52	
FEATURE	Location/Qualifiers	
source	1..52	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 66		
caagctcagc taattagagc ttcattgttt aaatattgaa agtaataatt tc		52
SEQ ID NO: 67	moltype = DNA length = 71	
FEATURE	Location/Qualifiers	
source	1..71	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 67		
gaaaggctct ttacgttaga tgagcttadc agattaaaat taattgcatg acattacacg	60	
tcttgagcga t	71	
SEQ ID NO: 68	moltype = DNA length = 71	
FEATURE	Location/Qualifiers	
source	1..71	
	mol_type = other DNA	

-continued

```

                                organism = synthetic construct
SEQUENCE: 68
gcacaaacgg caccaaacaa accagaacca acaatgatat aatcgatcat atgaatatcc 60
tccttagttc c 71

SEQ ID NO: 69      moltype = DNA length = 70
FEATURE           Location/Qualifiers
source            1..70
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 69
tcaacagtca agcagttttg gaaaagttat catcattata aaggtaaaac tgaatatcct 60
ccttagttcc 70

SEQ ID NO: 70      moltype = DNA length = 67
FEATURE           Location/Qualifiers
source            1..67
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 70
ttgtatagat aagaagtgag ttttaactca cttcttaaac ttgtttattc attgtgtagg 60
ctggagc 67

SEQ ID NO: 71      moltype = DNA length = 18
FEATURE           Location/Qualifiers
source            1..18
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 71
tggctgctat tgggcgaa 18

SEQ ID NO: 72      moltype = DNA length = 22
FEATURE           Location/Qualifiers
source            1..22
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 72
tcaccgata tgatttcttt tc 22

SEQ ID NO: 73      moltype = DNA length = 16
FEATURE           Location/Qualifiers
source            1..16
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 73
tatcccaatg gcatcg 16

SEQ ID NO: 74      moltype = DNA length = 19
FEATURE           Location/Qualifiers
source            1..19
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 74
accctaattc acgtactcc 19

SEQ ID NO: 75      moltype = DNA length = 1773
FEATURE           Location/Qualifiers
source            1..1773
                  mol_type = genomic DNA
                  organism = Campylobacter jejuni

SEQUENCE: 75
atgatttttt ataaaaagcaa aagattagca ttttttttaa tttcagatat tgttttaatt 60
ttacttagcg tttatctggc tttttctttg agatttagtg gagatattcc gagcattttt 120
tatcatggca tgatggtttc tgctattatt ttgcttggtt taaaactttc atttttattt 180
gttttttagaa tttacaaagt agcttggaga tttttttctc tcaatgaagc aagaaaaaatt 240
tttatcgctt tgcttttagc tgagttttgt ttttttctta ttttttattt ttttagtgat 300
ttttttaatc cttttccaag aagtgtctatt gtgatagatt ttgttctttc ttatatgttt 360
atagggtactt taagaattag caaaagaatg cttgtggatt ttaaaccttc tagaatgaaa 420
gaagaagaaa ctcttgttat tgtagtaggg gcaacttcta aggcctttgca tttgttaaaa 480
ggtgcaaaag aaggttcttt agggcttttt cctgtaggcg tagttgatgc gagaaaagag 540
cttataggga cttattgtga taaatttatt gtagaagaaa aagaaaaaat aaaatcttat 600
gcagaacaag gggtaaaaac tgccattatt gctttaagac ttgaacaaga agagcttaaa 660
aaactttttg aagaacttgt agcttatggt atttgcgatg taaaaatatt ttcttttaca 720
agaaacgaag caagagatat cagtatagaa gacttgcttg ctagaaaacc aaaagattta 780
gatgatagtg ctgtggcggc ttttttaaaa gataaggtag ttttggttaag tggagcagg 840
ggaactatag gcagtgaact ttgtaagcaa tgtattaaat ttggtgctaa gcatcttatc 900

```

-continued

```

atggttgatc atagttagta taatctttat aagatcaatg atgattttaa tttatataaa 960
gaaaaaatta ctcctatttt actgagtatt ttagataagc aaagttaga tgaggatta 1020
aaaacttata aaccogagct tattttacat gcagccgctt ataacatgt gcctctttgc 1080
gaacaaaatc cacattcagc agtaatcaat aatatttttag gaactaaaat tttatgcgat 1140
agtgtctaaag aaaacaaagt agctaaattt gtgatgataa gtacagataa agcagtacga 1200
ccaacaaata ttatgggttg cactaagaga gtttgcgagc tttatacttt aagtatgagt 1260
gatgaaaatt ttgaagttgc ttgtgtgcgt ttggtaatg ttttaggttc tagtggtagt 1320
gtgataccga aatttaaagc acaaatggcc aataatgagc ctttaacttt aacgcacct 1380
gatatagtgc gttattttat gcttgtggct gaggcagtgc aacttgtttt acaagctgga 1440
gctatcgcaa aagggggaga actttttgtt ttggatatgg gtaagcctgt gaaaatcata 1500
gatttagcta aaaaaatgct tttactttct aatcgcaatg atttagaaat taaaatcaca 1560
ggcttaagaa aaggtgagaa gctttatgaa gagcttttga ttgatgaaa tgatgctaaa 1620
acacaatatg agagtatttt tgtagcaaa aatgagaagg ttgatcttaa ttggcttaat 1680
aaagagatag aaaatttaca aatatgtgaa gatatttcag aggcctttatt aaagattgta 1740
cctgaattta aacacaataa agaaggtgta taa 1773

```

```

SEQ ID NO: 76      moltype = AA  length = 590
FEATURE           Location/Qualifiers
source            1..590
                  mol_type = protein
                  organism = Campylobacter jejuni

```

```

SEQUENCE: 76
MIFYKSKRLA PFLISDIVLI LLSVYLAFSL RFSGDIPSIF YHGMMSVSAII LLVLKLSFLF 60
VFRIYKVAWR FFSLENEARKI FIALLLAEFC FFLIFYFFSD FNNPFPRSAI VIDFVLSYMF 120
IGTLRISKRM LVDFKPSRMK EETPCIVVG ATSKALHLK GAKEGSLGLF PVGVVDARKE 180
LIGTYCDKPI VEEKEKIKSY AEQGVKTAII ALRLEQEEELK KLFEELVAYG ICDVKIFSFT 240
RNEARDISTE DLLARKPKDL DDSAVAFLK DKVVLVSGAG GTIGSELCKQ CIKFGAKHLI 300
MYDHSSEYNLY KINDDLNLYK EKITPILLSI LDKQSLDEVL KTYKPELILH AAAYKHVPLC 360
EQNPHSAVIN NILGKILCD SAKENKVAKF VMISTDKAVR PTNIMGCTKR VCELYTLSMS 420
DENFEVACVR FGNVLGSSGS VIPKFKAQIA NNEPLTLTHP DIVRYFMLVA EAVQLVLQAG 480
ATAKGGELEFV LDMGKPKVII DLAKKMLLLS NRNDLEIKIT GLRKGEKLYE ELLIDENDAK 540
TQYESIFVAK NEKVDLNLWN KEIENLQICE DISEALLKIV PEFKHNEKEV 590

```

```

SEQ ID NO: 77      moltype = DNA  length = 1032
FEATURE           Location/Qualifiers
source            1..1032
                  mol_type = genomic DNA
                  organism = Vibrio vulnificus

```

```

SEQUENCE: 77
atgaccaa atcgaaaaat ccaacaagat ttagttagaat caccaaaaac atggttgata 60
accggtgttg ctggttttat tggctccaat ttactagaaa aactactaaa gcttgatcaa 120
accgtggtgg ggtagataaa ctttgccacg ggtcatcaac ataacctaga tgaagttaaa 180
agtttagtaa cagaagccca atggcaacga ttacacttta ttgaaggcga tatacgtaat 240
cctgagacct gtgtagcagc ggttcaaggt gtagactatg tattacacca agcggcacta 300
gggtctgtgc cccgttcaat taatgatcct attactacaa attccgctaa catcactggt 360
tttttaaca ttgttagtagc agccaaagat cagcaagtga aaagctttac ttatgcgcc 420
tcaagttcga cttatggtga tcaccctgca ttgccgaagg tagaagaaaa tattggcaac 480
ccgttatccc cctacgccgt aaccaaatac gtaaatgaac tttatgcgaa tgtatttgcg 540
cggacttatg gttttaaaag tattggactg cgttatttta atgtgttttg aaaacgccaa 600
gatccaaatg gtgcctatgc ggtgtgtatt ccaaaatgga cagcagcaat gattcaaggt 660
gatgttatt atattaatgt tgacggcgag actagtgcgc atttctgtta tatagctaat 720
acagtgcaaa tgaatattct tgccgccacc gcaccagaag aagcaaaaga caagatttac 780
aatgttcgag taggagacag aacaacctta aacgatctat tctctgccat taaattggca 840
ttatcgaata atactgttgt tgtttctcaa gaacctattt atagagagtt tcgtactggt 900
gatgtaaggc attctcaggc tgatgttaat aaagctaagc agtatttagg ctatgcgccg 960
tcgcaccgta ttctagatgg aatcaacgaa gcaatgccat ggtatataag ctttataaat 1020
gataaaaaat aa 1032

```

```

SEQ ID NO: 78      moltype = AA  length = 343
FEATURE           Location/Qualifiers
source            1..343
                  mol_type = protein
                  organism = Vibrio vulnificus

```

```

SEQUENCE: 78
MTKYEKIQDQ LVESPKTWLI TGvagfigsn LLEKLLKLDQ TVVGLDNFAT GHQHNLEDEVK 60
SLVTEAQWQR FTFIEGDIRN PETCVAAVQG VDYVLHQAL GSVPRSINDP ITTNSANITG 120
FLNMLVAAKD QQVKSFTYAA SSSTYGDHPA LPKVEENIGN PLSPYAVTKY VNELYANVFA 180
RTYGFKSLGL RYFNVFGKRQ DPNGAYAARI PKWTAAMIQG DVIYINGDGE TSRDFCYIAN 240
TVQMNILAAAT APEEAKDKIY NVAVGDRRTL NDLFSAIKLA LSNNTVVVSQ EPIYREFRTG 300
DVRHSQADVN KAKQYLYGAP SHRILDGINE AMPWYISFIN DKK 343

```

```

SEQ ID NO: 79      moltype = DNA  length = 223
FEATURE           Location/Qualifiers
source            1..223
                  mol_type = genomic DNA
                  organism = Bacteroides fragilis

```

-continued

SEQUENCE: 79
 atgaaaatac ctttttcacc accttatatt gacgaggcgg tcatcaacga agtcgttgat 60
 tcgttacggt ccggttgat cacatccgt cggaaagtga aggctttgga ggaagaaatt 120
 aaatctttct caggagcaaa agaagtgcgt tgtgtcaatt cctggacatc aggagctatt 180
 atgatgttgc gttggctggg agtgaaggag ggcgatgaag tga 223

SEQ ID NO: 80 moltype = AA length = 407
 FEATURE Location/Qualifiers
 source 1..407
 mol_type = protein
 organism = Bacteroides fragilis

SEQUENCE: 80
 MKIPFSPPI DEAVINEVVD SLRSGWITSG PKVKALEEEI KSFSGAKEVL CVNSWTSGAI 60
 MMLRWLGVE GDEVIVPAYT YSATALAVLH AGAKPVMVDS GTDFNISVEA VRKAITPKTK 120
 AIIPVDIAGF PCDYERIMAL VQEPFEMVKLF RSESPVQEKI GRILVMNDAA HSLGARYSSR 180
 QRTGCETDVA IFSLHAVKNV TTAEGGAICL NLPKPFNDTE LYKELRMTSL NCQTKDAFSK 240
 SKAGGWRYDI VGFGMKINMA DVNAAIGLAQ IREYPELLKE RKRVFNAYS D AFSACDWAIV 300
 PPSVDGEKES SYHIYALRIK DFTEEQRDRM IDEIAKSEVA VNVHFIPMPM LSFFKSMGYD 360
 IKDYPQAYQN FKSEISLPIY PQLDSEKLN IETVKAAYA TVIAENR 407

SEQ ID NO: 81 moltype = DNA length = 597
 FEATURE Location/Qualifiers
 source 1..597
 mol_type = genomic DNA
 organism = Bacteroides fragilis

SEQUENCE: 81
 ataagttgct tgatccggtt ttttgatatt gttttttccc ttttgggaat tcttctgctt 60
 tctcccggtt ttcttcttct ttatatagcc atttgtctcg aaagtaaagg aggtgggttc 120
 tataaacagc tccgtgtggg aaggtacgga ggggattttt atgtttataa atttcgttcg 180
 atgctgtgat gggctgataa aaaggggttg ataacggtag gaggagagaga tccgagaata 240
 actcgtaccg gatacctgat ccggaagtat aaattggatg aattgcctca gttgtttaac 300
 gtactgaaag gagatatgag tctggtaggg ccccgctctg aggttcgtaa gtatgtggat 360
 ttatataccg atgaacaaaa gaaagtattg agtgcacgc cggggattac ggattatgct 420
 tctattgagt atgtagatga aaatatgatt cttggagagg cttcggatcc tgacagagca 480
 tatatagaac agattatgcc cgataagata cgggataata tgaatatat ctgtaatcgt 540
 tcggggaagg aatattttta aattatattt ctgacatttt ggagcatcat tcgtagg 597

SEQ ID NO: 82 moltype = AA length = 198
 FEATURE Location/Qualifiers
 source 1..198
 mol_type = protein
 organism = Bacteroides fragilis

SEQUENCE: 82
 MSCLIRFFDI VFSLLGILL SPVFLLLYIA ICLESKGGGF YKQLRVGRYG GDFYVYKFRS 60
 MRVGADKKGL ITVGGRDPRI TRTGYLIRKY KLDELPLQFN VLKGDMSLVG PRPEVRKYVD 120
 LYTDEQKKVL SVRPGITDYA SIEYVDENMI LGEASDPDRA YIEQIMPKI RYNMKYICNR 180
 SVKEYFKIIF LTFWSIIR 198

SEQ ID NO: 83 moltype = DNA length = 807
 FEATURE Location/Qualifiers
 source 1..807
 mol_type = genomic DNA
 organism = Bacteroides fragilis

SEQUENCE: 83
 atgaaattgg ctgtcatttc ttctatctac aaaaatgatt cgctttctaa aataaagcaa 60
 tctttacaaa gtttgttgaa tcaagaaaat gttacttgtg atattttatt atattgtgat 120
 ggtcctgttt cgattgatat tgtgcatttc ttagactcat ttgatcaaaa ctacatttat 180
 atattttaagc aagaacaaaa tagaggattg gctcatgcac taaatgaatt attgaatatt 240
 gttcttacaa aagaatacga atatatgct cgtatggatg ctgatgatatt tccatgccg 300
 gaacgttttg tgaagcagat cgcatttatg aattcacatc cggatattga ttgccttggt 360
 acctgggcaa tagaaattga tgatgacgga aaagaatatt ttcgaaaaaa aatgccgata 420
 actcatgaag aatgtttgga actgtttaaa aagcgggatt gtatgattca tctactgtg 480
 atgtttcgac gttcatattt tgagaagct ggtttatatt ccgaagacac ctactttgga 540
 gaagatacta tgatgtgggc taaaggtttt aaaagtggat gtaaatgtgc caatgtaccc 600
 gaatatctgt ttaagtttag attggattca aactctttg aaagaaggcg ggggtggaaa 660
 catgcaaaat ctattttatc ttacgtcat cgtgttaata gaattgtagg gttcggatgg 720
 aaggaagatt gttatgcatt gctatatgct atggccaaat taatgcctaa atctattctt 780
 gacataatct ataaaacagt acgttga 807

SEQ ID NO: 84 moltype = AA length = 268
 FEATURE Location/Qualifiers
 source 1..268
 mol_type = protein
 organism = Bacteroides fragilis

SEQUENCE: 84
 MKLAVISSIY KNDLSKIKQ SLQSLNQN VTCDIYLYCD GPVSIIDIVH LDSFDQNSLY 60

-continued

IFKQEQNRGL	AHALNELLNI	VLTKEYEYIA	RMDADDISMP	ERFVKQIAFM	NSHPDIDCLG	120
TWAIETDDG	KEYFRKKMPI	THEECLELPK	KRDCMIHPTV	MFRRSYFEKA	GLYPEDTYFG	180
EDTMMWAKGF	KSGCKFANVP	EYLFKFRLLDS	NFFERRRGWK	HAKSIYTLRH	RVNRMLGFGW	240
KEDCYALLYA	MAKLMPKSIL	DIYKTVR				268

SEQ ID NO: 85 moltype = DNA length = 1044
 FEATURE Location/Qualifiers
 source 1..1044
 mol_type = genomic DNA
 organism = Bacteroides fragilis

SEQUENCE: 85

atgaggaaga	tattattaac	atatggagat	attaaaacca	taaatattgg	tgattatatt	60
cagagtatcg	cgcgtaaaac	atTTTTgat	gatgataatt	atatattttt	caatcgtgat	120
gaattaaagt	tgtataaagg	ggaacctgca	aaagtcataa	tgaatgcttg	gatgacttat	180
aaaccatata	attggcctcc	ttcatctcag	gtttatccat	tatttggtgc	attacatatac	240
aattcttctg	cagaagctcg	ctttttgtca	catgatagta	ttaagtattt	aaaaaagtat	300
gaaccocatag	ggtgocgcga	ttatcataca	atgaatattt	taaaaggaaa	aggtgttaat	360
gcttattttt	cgggatgtct	tactactact	ttgggtaaaa	cttataaata	taatggaaaa	420
cgtgagggga	tatacattgt	tgatccactt	tcatacatgc	ctaaggggaa	taattttttt	480
gaaataatga	aagctgttgt	acaaaactgt	ttttatatga	agcctgtgct	aaagattctt	540
agaaattata	aaaagaataa	tagattttact	attaatatat	ccaaagtagg	aataggacgt	600
ttattgttaa	taaccaagtc	ttatttgctt	ttgagaaagc	tagtagttcc	tgacgtatta	660
tacaatgcaa	tatttataac	acaatttaac	atgtctaattg	aattattcttc	tgaatcagaa	720
agatttgctc	gggctgatga	attattaact	aaaatggcct	cagctcaata	tgtgataact	780
tctagaatcc	actgtgctct	tccttgctta	ggctttgaga	cccctgtggt	ttatattaga	840
aatctatcag	aaagcaagaa	gagcacttgt	cgattagggg	gacttgaatc	tctatttaac	900
gtgataactg	tgaaggggga	aaatgttact	tccaattttt	ttgatggact	atttaaacga	960
gattcttctt	ttaagaataa	aattgatattt	gtagaatata	gagatcgatt	aattaatata	1020
tgtaataaat	ttatgatgtc	atga				1044

SEQ ID NO: 86 moltype = AA length = 347
 FEATURE Location/Qualifiers
 source 1..347
 mol_type = protein
 organism = Bacteroides fragilis

SEQUENCE: 86

MRKILLTYGD	IKTINIGDYI	QSIAAKQFFD	DDNYIFFNRD	ELKLYKGEP	KVIMNAWMTY	60
KPYNWPPSSQ	VYPLFVALHI	NSSAESRFLS	HDSIKYLKKY	EPIGCRDYHT	MNILKKGKVN	120
AYFSGCLTTT	LGKTYKYNGK	REGIYIVDPL	SYMPNGNNFF	EIMKAVVQTV	FYMKPVLKIL	180
RNYKKNNRFT	INISKVGIGR	LLLITKSYLL	LRKLVPDVL	YNAIFITQFN	MSNEYSSESE	240
RFARADELLT	KMASAQYVIT	SRIHCALPCL	GFETPVVYIR	NLSESKKSTC	RLGGLESFPN	300
VITVKGENVT	SNFFDGLFKR	DSSFKNKIDF	VEYRDRLINI	CNKFMMS		347

SEQ ID NO: 87 moltype = DNA length = 1137
 FEATURE Location/Qualifiers
 source 1..1137
 mol_type = genomic DNA
 organism = Bacteroides fragilis

SEQUENCE: 87

atgagggatg	gaaagcccat	tgaataactg	catattgtag	ggactagacc	tgttggagga	60
attgtgtgctt	tattgaaaaa	tataaacact	tctattgatt	taataaaatt	ccatttttaca	120
tatgtgtttt	ctgocgatag	taatatggga	gattttgata	attatgtgag	aaaattagggt	180
agcgatattg	ttgtattccc	ttcatatcat	ttgaaatatt	tattttttata	tttaaagggtg	240
attttctggt	tttataagag	aaatgcaaaa	aaatatgata	taattcatgt	acatagtgtct	300
aatacaggag	tattagacct	tttatttgca	aagatatatg	gaatacggat	aaggattttta	360
catagtccaca	gtacaaaata	ttctagttaa	aaaataagggt	ccattagaaa	ttatttttttg	420
caatttccta	caatatatatt	agcaaacact	tattttgcat	gtggatacaa	agctgctgaa	480
ttcctttttg	gaaaaaagag	acttgaaaaa	gtttatatta	ttcataatgc	aattttttct	540
aaaaaattta	tttataatgc	aactgttaga	gagagagtta	gagttgagtt	gaaaaatagag	600
aatgactggt	tactgttggg	gcataatcgt	aactttacga	aagaaaaagaa	tcatgtgtttt	660
ttaatagata	tattggaaaga	attattgaat	attaatgaca	atgttaagtt	gctgttagta	720
ggggatggcc	aacttagaag	tgaatatagag	gaaaaagtaa	aattaaaaaag	attgcagaag	780
tatgtgtgct	ttttaggctg	gagaactgat	atatcagaat	tattacaagc	tatggattttt	840
ttaatttttc	cttctttttt	tgaaggctta	ccagtgtact	tgattgaagc	gcaagcttct	900
ggtttaagat	gtgtcgtttc	ggatcgattt	actttggaaa	caaagataac	gaataatata	960
agttatctta	atctcaatcg	agatccttca	acttgggctg	aagttgtttg	gaagttgtttt	1020
gaagataaaa	cttttatcag	gaaagatacg	tctaagaaa	tatgtgaatc	aggatatgat	1080
atacathtag	agagtgaata	tctcgaaaga	atatatttaa	atctttttaca	taaatga	1137

SEQ ID NO: 88 moltype = AA length = 378
 FEATURE Location/Qualifiers
 source 1..378
 mol_type = protein
 organism = Bacteroides fragilis

SEQUENCE: 88

MRDGKPIEL	HIVGTRPVGG	IGALLKNINT	SIDLNKFHFT	YVFSADSNIG	DFDNVYRKL	60
-----------	------------	------------	------------	------------	-----------	----

-continued

SDIVVPFSYH	LKYLFLYLKV	IFCFYKRNAK	KYDIHVHSA	NTGVLDDLFA	KIYGIRIRIL	120
HSHSTKYSSK	KIRSIRNYFL	QFPTIYLANT	YFACGYKAAE	FLFGKKRLEK	VYIIHNAIFS	180
KKFIYNATVR	ERVRVELKIE	NDCLLLGHIG	NFTKEKNHGF	LIDILEELLN	INDNVKLLLV	240
GDGQLRSEIE	EKVKLKRLQK	YVCFLGRRTD	ISELLQAMDF	LIFPSFFEGL	PVTLIEAQAS	300
GLRCVVDRI	TLETKITNNI	SYLNLNRDPS	TWAEVVKLF	EDKTFIRKDT	SKEICESGYD	360
IHLESEYLER	IYLNLLHK					378

SEQ ID NO: 89 moltype = DNA length = 1095
 FEATURE Location/Qualifiers
 source 1..1095
 mol_type = genomic DNA
 organism = Bacteroides fragilis

SEQUENCE: 89

atgaaaaaaa	aatatgacta	tctaattgtc	ggagcgggac	tttatgggtc	tgtttttgca	60
tataaagcgc	aacggaatgg	gaaaaaatgt	ttgcttattg	acaaacgtcc	acacaatgga	120
ggaaatatatt	attgcgaaaa	tattgagggg	ataaatgtac	acaaatatgg	tgacacatatt	180
tttcatactt	ctaataaaga	ggtttggggc	tttgtaaatg	ccatagtgga	atttaaccgc	240
tatactaatt	cccctattgc	caattataaa	ggcaaatat	ataatttacc	ttttaatatg	300
aatacttttt	atgcttttatg	ggggacaaaa	actccagagg	aagcaaaaat	gaagatagaa	360
gagcaaaagga	gagaagcggg	gattctttacg	cgaagaatc	tagaagaaca	ggctatctct	420
ttagtaggaa	aggatattta	tgaaatattg	ataaaaggat	atactgagaa	acagtggggg	480
cggaaagcaa	ctgaacttcc	tgcttttata	atcaagcgtt	tacctgtcag	atttactttt	540
gataataaatt	actttaatga	taaatatcag	gggattcctg	ttggtggcta	taataaaattg	600
ataaatggtt	tattagatgg	tattgaagtt	agaacaaata	cagacttttt	tcaaaataaa	660
gaatatattta	aagggtcttg	ggataaaaatt	ctatttactg	gcaaaattga	tgaatattat	720
aattatcaat	ttgggagagt	ggaatatcgt	actgttgaat	ttgaaacaca	agtgttgat	780
tgtgagaatt	atcaagggaa	tgcaagtgtg	aattatactg	aaaggggaagt	gccttatact	840
cgtattatag	agcataaaca	tttgaattt	ggaacacagc	ccaaaaccgt	aatttccaaa	900
gaatattctt	cggaaaggaa	agatggtagt	gaaccctttt	atcctgttaa	tgatgagagg	960
aataattctt	tgtatttgga	gtataagaaa	ttggtcgata	gagagaagaa	tgttttgttt	1020
ggaggggagat	tagctgagta	taagtattat	gatatgaatg	tgattgtgga	aaaagttata	1080
aatagtgact	tatga					1095

SEQ ID NO: 90 moltype = AA length = 248
 FEATURE Location/Qualifiers
 source 1..248
 mol_type = protein
 organism = Bacteroides fragilis

SEQUENCE: 90

MKKKYDYLLV	GAGLYGSVFA	YKAQRNGKKC	LLIDKRPHNG	GNIYRKATEL	PAFIIKRLPV	60
RFTFDNNYFN	DKYQGIPVGG	YNKLINGLLD	GIEVRTNTDF	FQNKEYFKGL	ADKILFTGKI	120
DEYYNYQFGR	LEYRTVEFET	QVLDCENYQG	NAVNYTERE	VPYTRIIEHK	HFEEGTQPKT	180
VISKEYSSEW	KDGEPPFPV	NBERNNSLYL	EYKKLVDRK	NVLFGGRLAE	YKYDMMNVIV	240
EKVINSDL						248

SEQ ID NO: 91 moltype = DNA length = 876
 FEATURE Location/Qualifiers
 source 1..876
 mol_type = genomic DNA
 organism = Bacteroides fragilis

SEQUENCE: 91

atgaagatat	ttgctgtagt	tgtgacctat	aatcgtttag	cattactaaa	aaaggtcata	60
gatttactta	aaagacagac	acgtcaatta	gatagtatag	ttatagtaaa	taatggctct	120
actgatggaa	caaaagaatg	gttagaacag	gagaaaaata	taacattaat	aatcaatct	180
aattctgggtg	gggctgggtg	attcgaaacg	ggggtaaaat	atgcatatga	gaatgaggca	240
gattgggattt	ggatgatgga	tgatgatgtt	tttcctaata	tagattgttt	ggaaaaactc	300
ttaggggtgga	cgagtatttc	tcaatgtatc	caaccacgta	ggattatttc	ggatgatgta	360
gaagtaaat	ttgaacaatg	gttggatcca	attacttata	gtaaatattg	ttattggcaa	420
gaaaaatcat	ttaataatgg	taaaaagttt	tgtgctatta	atgtaggctg	ttttgaaggg	480
atgtttgttt	cgaaaagcat	tgtaataaaa	ataggatttc	ctgataagcg	tttttttata	540
gcggaagatg	atacaatata	tggttttgca	gctagttttt	ataccaacgt	aatttttagta	600
tctgatgcta	tcatgtgtcg	tgctagaaga	tcatcggtac	gtagcggttag	cccaatgtat	660
acttattatg	catgtcggaa	ttttcattta	gtatatgagt	cacttaattt	attattggat	720
aagaaaaata	ggtttttata	tattaaatat	atttatcaat	ttgttcatca	gatttattta	780
tctatttttt	tatatgataa	taagatgaaa	cacttaatat	ctgttttttcg	tggtctttat	840
gattgctttc	gtaaaaaaaa	tggaagctacc	tattaa			876

SEQ ID NO: 92 moltype = AA length = 291
 FEATURE Location/Qualifiers
 source 1..291
 mol_type = protein
 organism = Bacteroides fragilis

SEQUENCE: 92

MKIFAVVVTY	NRLALLKKVI	DLLKRQTRQL	DSIVIVNNGS	TDGTEKWELEQ	EKNITLINQS	60
NSGGAGGFET	GVKYAYENEA	DWWMMDDDV	FPNIDCLEKL	LGWTSISQCI	QPRRYSDDV	120
EVNFEQWLDP	ITYSKFGYWQ	EKSFNNGKKF	CAINVGCFFG	MFVSKSIVNK	IGFPDKRFFI	180

-continued

AEDDTIYGFA	ASFYTNVILV	SDAIMVRARR	SSDRSVSPMY	TYYACRNPHL	VYESLNLLLD	240
KNRFLYIKY	IYQVHVQIYL	SIFLYDNKMK	HLISVFRGFY	DCFRKKNGAT	Y	291

SEQ ID NO: 93 moltype = DNA length = 1140
 FEATURE Location/Qualifiers
 source 1..1140
 mol_type = genomic DNA
 organism = Escherichia coli

SEQUENCE: 93

atgatgaaat	ccaaaatgaa	attgatgcca	ttattggtgt	cagtaacctt	gataagcggg	60
tgacacagta	ttccgggcag	caatatgtcg	acgatgggca	aagacgtcat	caaacagcag	120
gacgctgatt	tcgatctcga	caaaatggtg	aatgtttatc	cgctgacccc	gcgcctgatt	180
gaccaattac	gccacgccc	gaatgtagcg	cgccccaata	tgacgctgga	aagttagatc	240
gcgaattacc	agtatcgcgt	cgggcggggg	gacgttctta	atgtcacggt	ctgggacac	300
ccggaactca	ccacgcacgc	cggtcagtag	cgcagctcca	gcgacacggg	caactgggta	360
cagcctgacg	gcactatggt	ttaccggtat	atcggaagg	tccacgtagt	cgggaaaacg	420
ctcgctgaaa	tcgcagtgga	tattaccggg	cgcttagcga	cgtacatcgc	tgacccgcag	480
gtggacgtta	atatcgccgc	cttcgcgtca	caaaaggcct	atatctccgg	tcagggtgaat	540
aaatccggtc	aacaggcgat	caccaacgtg	ccactgacta	ttctcgacgc	catcaacgcc	600
gcaggtggcc	tgaccgacac	cgctgactgg	cgcaacgtgg	tgctaacaca	caatgggtcgt	660
gaagagcgca	tttctttgca	ggcgctgatg	caaaacggcg	acctcaacca	gaatcgccctg	720
ctttaccocg	gcgatattct	ctacgtgcca	cgtaatgatg	atctgaaagt	atttgtgatg	780
ggtgaagtga	agaaacagag	caccctgaaa	atggacttta	gcgccatgac	cctgactgaa	840
gcccctgggca	atgctgaagg	catcgacatg	accacctcca	acgccagcgg	catctttgtc	900
attcgtccgc	tgaaaggcga	gggcggggcg	aacggcaaga	ttgccaatat	ctaccagctg	960
gatatgtcgc	atgccacgct	gctgggtgatg	gcgacagaa	tccgcctgca	accttatgac	1020
gtgggtgatg	tcaccacccg	cccggtttcc	cgctggaacc	gtctgatcaa	tcagttgctg	1080
ccaactatta	gcggtgtccg	ttacatgacg	gatacagcca	gcgacattca	taactggtaa	1140

SEQ ID NO: 94 moltype = AA length = 379
 FEATURE Location/Qualifiers
 source 1..379
 mol_type = protein
 organism = Escherichia coli

SEQUENCE: 94

MMKSKMKLMP	LLVSVTLISG	CTVLPGSNMS	TMGKDVIKQQ	DADFDLDKMW	NVYPLTPRLI	60
DQLRPRPNVA	RPNMTLESEI	ANYQYRVGPG	DVLNVTVDH	PELTTPAGQY	RSSSDTGNWV	120
QPDGTMFYYP	IGKVHVVGKT	LAEIRSDITG	RLATYIADPQ	VDVNIAAFRS	QKAYISGQVN	180
KSGQQAITNV	PLTILDAINA	AGGLTDTADW	RNVVLTHNGR	EERISLQALM	QNGDLNQNL	240
LYPGDILYVP	RNDLKVFFVM	GEVKKQSTLK	MDFSGMTLTE	ALGNAEGIDM	TTSNAGSIFV	300
IRPLKGEGR	NGKIANIYQL	DMSDATSLVM	ATEFRLQPYD	VVYVTTAPVS	RWNRLINQLL	360
PTISGVRYMT	DTASDIHNW					379

SEQ ID NO: 95 moltype = DNA length = 1449
 FEATURE Location/Qualifiers
 source 1..1449
 mol_type = genomic DNA
 organism = Bacteroides fragilis

SEQUENCE: 95

atgggacaat	caataaagaa	taacttccca	ctgaatctaa	gtactacaat	tactgggcta	60
ttatttccac	taataacttt	tcctcatgct	tcacgtatat	taatggcaga	tggtataggg	120
caagtacaat	ttttccaatc	cataattgac	tatgtctctc	tttgtagctg	tttgggaatc	180
cctttgtatg	ccgttagaga	aattgcaaga	ataagggata	ataagggaat	aagaagtaga	240
acaacaatcg	aaataactatt	gcttcatgct	attttgactt	tagtcggata	tatagttggt	300
tttatacttg	ctaagaccgt	tgccaaaata	gaaatagatg	cttctctttt	ctttttgtta	360
agtacaactt	tattttttta	cacgatagga	gttgcatggt	tttatcaggg	aattgaagat	420
tttaaataca	taactttacg	ttccttattt	gtaaggatac	tatctttagt	ggccttattt	480
atcctttgta	agaccaagca	agatctgttt	tattatgctg	gaatattagt	tatttggtact	540
gttggttaata	atatatttta	tttttttcga	ttacgtaaat	atataaaact	tagtaaaaggc	600
gaatttaagc	gtttgaatct	gttgaggcat	ttaataacctg	cacttaaaat	ttttatatta	660
aatttggtta	taagtattta	tgtgaatctt	gattctgtga	tgcttggtat	tcttaaaat	720
gaagaatcag	ttgggttatta	tgtctgctgt	actcgactta	caaaagctat	tttgggtata	780
gtttcttcat	taggagctgt	tttattacct	cgtttttagta	atatgataac	taatgggtcaa	840
aaagaagaat	tccaattatt	ggcaataaaa	gctgctagtt	ttacaattgc	tttaagtctt	900
cctatgagtg	tgggacttat	ttttatggca	gcacctatta	tacatatttt	ttgtggaaat	960
ggttttgaac	cttctatttt	gacattaaaa	ttagttgcac	ctattgtttt	atttattgga	1020
ttatctggaa	ttattggaat	gcaaatattg	tatccgcaag	ggagagaaaa	gtacgtaatt	1080
atatctacta	tggtaggagc	atgtataaat	ctccttataa	actatttact	aataccccaa	1140
tatgggcagt	atggagcggc	gcttgggact	gttatcgtag	aatttatggt	aacgggtata	1200
atgattttgc	tggaagaaaa	atatttacct	ataaatattc	tctccaagca	gaatttacac	1260
tatctaatag	gttctatagt	gatattctatt	ctattagctt	ttttgtttgt	cttcctttta	1320
catgaagtaa	attattttatt	gataggtata	ctgttatctg	ttattgttta	ttacgcataat	1380
ttattaatga	taaaagatac	cctcgcatta	caactaaaga	aattattact	ttcaatattt	1440
aaacaatga						1449

SEQ ID NO: 96 moltype = AA length = 482

-continued

FEATURE
source Location/Qualifiers
1..482
mol_type = protein
organism = Bacteroides fragilis

SEQUENCE: 96

MGQSIKNNFL	LNLSTTITGL	LFPLITFPYA	SRILMADGIG	QVQFFQSIID	YVSLCTALGI	60
PLYAVREIAR	IRDNKELRSR	TTIEILLHA	ILTLVGIVV	FILAKTVAKI	EIDASLFFLL	120
STTLFFNTIG	VAWFYQAIED	FKYITLRSLF	VRILSLVALF	IFVKTQDLF	YYAGILVIGT	180
VGNNIFNFFR	LRKYIKLSKG	EFKRLNLLRH	LIPALKIFIL	NLVISIVVNL	DSVMLGFLKN	240
EESVGYAATA	TRLTKAILGI	VSSLGAVLLP	RFSNMITNGQ	KEEPQLLANK	AASFTIALSL	300
PMSVGLIFMA	APIIHIFCGN	GFEPISILTK	LVAPIVLFIG	LSGIIGMQIL	YPQGREKYVI	360
ISTMVGACIN	LLINYLLIPQ	YGQYGAALGT	VIAEFMVTVI	MILLGRKYLP	INILSKQNLH	420
YLIGSIVISI	LLAFLFVFP	HEVNYLLIGI	LLSVIVYYAY	LLMIKDTLAL	QLKKLLLSIF	480
KQ						482

SEQ ID NO: 97 moltype = DNA length = 1305

FEATURE Location/Qualifiers
source 1..1305
mol_type = genomic DNA
organism = Bacteroides fragilis

SEQUENCE: 97

atgactagta	ctctctctct	tattattaag	gctatatctt	ttatagtatt	tgtttataat	60
ttagcgaggt	ctgattggag	taatctctct	cgagtattgt	tatttataat	ttcggttttt	120
gtctctttat	tagtaacaaa	tgtatatcaa	agaaaagtaa	aaatagtaac	actaggttat	180
ttggtctatt	ctcttttggc	tttttagct	caaaatgtag	gtccatagtc	aattatttac	240
aattctcaca	tacagaattt	gatacacagg	tttcctaata	tttgtgtacc	aaatgatcat	300
tacactttat	actttcttat	ttttattttt	gtttcaactg	ttatttatat	ttatctttta	360
tgtattgata	aaattttctat	tgagcagaaa	gtatgggtat	acatgcctac	aagacaagaa	420
gctcgaaaaa	taatatgggc	attttttata	atcttggtgc	ctatatggat	attaggggat	480
aggctctatg	ctacaattct	agttcctttt	acaagtattt	ttatcatttc	tatttggaat	540
tatctcttta	caaggaaaaa	agtagtattt	attattggtt	ttattgcatc	ggtattgata	600
cttatttcgc	aaattatttc	tcgggtttata	ttcggttcaat	atgtgttccc	ttttatttta	660
atttggtttt	tgaataatgg	attgctattt	tcaagaaaaa	aaaaataaag	ttataaagt	720
attggctttt	ttctattggt	ttgttcttta	gcgattttgt	atggatcatc	ttcagagatg	780
atgaagttaa	atgcaaatat	tgatggaaac	tatactttca	atgatatggt	tgaagtatta	840
agtgcagcgc	aattattgca	aaattgggtta	tctaggcaaa	catatcgat	ttttgatatt	900
tggtcacatt	tgggggggaa	tataatagat	tataattgacc	agaaaggata	tttatgggga	960
attacatatg	taaagagcgt	tgcacctata	ttaggttttg	attatgtaag	ccttccatta	1020
atttcagcaa	atatgattgg	tgcagattat	gctcaacctg	gtttggtggc	agaaggggtac	1080
gctaattggg	gaatataatg	ggctataatt	aatatgatat	ttgttttttt	cgttgcagag	1140
ttattgtttg	attatttttt	acgaaaaaaa	tctaccttaa	atttggtact	ttatattgga	1200
actttttccc	aggtattact	agatgggggg	accattaatt	ctattatttt	catgtctata	1260
ttttgtttct	taacttggtt	gtttaatgcc	aaaatcaaat	tttaa		1305

SEQ ID NO: 98 moltype = AA length = 434

FEATURE Location/Qualifiers
source 1..434
mol_type = protein
organism = Bacteroides fragilis

SEQUENCE: 98

MTSTSFIIK	AISFIVFVYN	LASSDWSNLS	RVLFIISFF	VFLLVITNVYQ	RKVKIVTLGY	60
LVIYFLAFIA	QNVGPYAIY	NSHIQNLIQR	FPNICVPNDH	YTLVFLIFIF	VSTVIYIYLL	120
CIDKISIEQK	VWVYMPTRQE	ARKIMMAFFI	ILLPIWILGN	RSYATILVPF	TSYFIISIWK	180
YFPTRKIVVF	IIGPIASVLI	LISQLFSRFI	YVQYVFPFIL	IWFLNNGLLF	SRKEKISYKV	240
IGFLLFVCSL	AILYGIISEM	MKLNANFDGN	YTFNDMFEVL	SEPQLQNWL	SRQTYRIFDI	300
WSHLGGNIID	YIDQGYLWVG	ITYVKSFAPI	LGFYDVSFLPL	ISANMIGADY	AQPGLVAEGY	360
ANWGIYGAII	NMIFVFFVAE	LLPDYFLRKQ	STLNLLLYIG	TFSQVLLDGG	TINSIIFMSI	420
FCFLTCLFNA	KIKF					434

SEQ ID NO: 99 moltype = DNA length = 45

FEATURE Location/Qualifiers
source 1..45
mol_type = other DNA
organism = synthetic construct

SEQUENCE: 99

tcataaaaaa	tttatttgct	ttgtgagcgg	ataacaatta	taata		45
------------	------------	------------	------------	-------	--	----

SEQ ID NO: 100 moltype = DNA length = 14

FEATURE Location/Qualifiers
source 1..14
mol_type = other DNA
organism = synthetic construct

SEQUENCE: 100

aggaggatat	acat					14
------------	------	--	--	--	--	----

SEQ ID NO: 101 moltype = DNA length = 12764

-continued

FEATURE	Location/Qualifiers
source	1..12764 mol_type = other DNA organism = synthetic construct
SEQUENCE: 101	
ctcgagaaat	cataaaaaat ttatttgcgt tgtgagcgga taacaattat aatagattca 60
attgtgagcg	gataacaatt tcacacagaa ttcattaaag aggagaaatt aactatgaga 120
ggatcgcatc	accatcacca tcacggaatg attttttata aaagcaaaag attagcattt 180
tttttaactt	cagatatgtt tttaatttta cttagcgctt atctggcttt ttctttgaga 240
tttagtggag	atattccgag tattttttat catggtatga tggtttctgc tattattttg 300
cttggtttta	aaactttcatt ttgttttgtt tttagaattt ataaagtagc ttggagattt 360
ttttctctca	atgaagcaag aaagattttt atcgctttgc tttagctga gttttgtttt 420
tttcttattt	tttatttttt tagtgatttt tttaatcctt tccaagaag tgctattgtg 480
atagattttg	ttctttctta tatgtttata ggtactttta gaattagcaa aagaatgctt 540
gtggatttta	aaccttctag aatgaaagaa gaagaaactc ctgtatttgt agtaggggca 600
acttctaagg	ctttgcattt gttaaaaggt gcaaaagaag gttctttagg gctttttcct 660
gtaggcgtag	ttgatgcgag aaaagagcct atagggactt attgtgataa atttatttga 720
gaagaaaaag	aaaaataaaa atcttatgta gaacaagggtg taaaactgc cattatttgc 780
ttaagacttg	aacagaaga gcttaaaaaa cttttgaag aactgtagc ttatggtatt 840
tgcatgttaa	aaatattttc ttttacaaga aacgaagcaa gagatcacg tatagaagac 900
ttgcttgcta	gaaaaccaaa agatttagat gatagtgtcg tggcggtttt tttaaaagat 960
aaagtagttt	tggttaagtgg agcaggtgga actataggca gtgaactttg taagcaatgt 1020
attaaatttg	gtgctaagca tcttatcatg gttgatcata gtgagtataa tctttataag 1080
atcaatgatg	atttaaatat atataaagaa aaaattactc ctattttact gagtatttta 1140
gataagcaaa	gtttagatga ggtattaaaa acttataaac ccgagcttat tttacatgca 1200
gccgcttata	aacatgtgcc tctttgcgaa caaaatccac attcagcagt aatcaataat 1260
attttaggaa	ctaaaaatatt atgcgacagt gctaaagaaa acaagtagc taaattttgt 1320
atgataagta	acagataaagc agtacgacca acaaatatta tgggttgcac taagagagtt 1380
tgcgagcttt	atacttttaag tatgagtgtat gaaaattttg aagttgcttt tgtgctttt 1440
ggtaattggt	taggtttctag ttgttagtgt ataccgaaat ttaagcaca aattgccaat 1500
aatgagcctt	taactttaac gcacctgat atagtgcgtt attttatgct tgtggctgag 1560
gcagtgcaac	ttgttttaca agctggagct atcgcaaaag ggggagaact ttttgtttt 1620
gatattgggt	agcctgtgaa aatcatagat ttagctaaaa aaatgctttt actttcta 1680
cgcaatgat	tagaaaattaa aatcacagcg ttaagaaaag gtgagaagct ttatgaagag 1740
cttttgattg	atgaaaatga tgctaaaaaa caatatgaga gtatttttgt agcaagaat 1800
gagaaggttg	accttgattg gcttaataaa gagatagaaa atttacaat atgtgaagat 1860
atttcagagg	ctttatttaa gattgtacct gaatttaaac acaataaaga aggtgtataa 1920
aggaggatat	acatatgaaa ataccctttt caccacctta tattgacgag gcggtcatca 1980
acgaagtcgt	tgattctgtt cgttccggtt ggatcacatc cggtccgaaa gtgaaggctt 2040
tggaaggaag	aattaaatct tctcaggag caaaagaagt gctttgtgtc aattcctgga 2100
catcaggagc	tattatgatg ttgcgttgcc tggggagtga ggaggggcag gaagtgattg 2160
ttcctgccta	tacttacagt gcaacggcat ttgctgtgct ccatgccgga gccaaaccgg 2220
taatggtgga	ttccggaact gattttaata tttcgttgga ggctgttctg aaagctatta 2280
ctcctaaaaa	aaaggcgatt ataccctgag atattgctgg ttttccctgc gattatgaaa 2340
gaatcatggc	actgggtcag gaaccagaaa tggtaaaact gttccgttcg gaatctcctg 2400
tgcaagaaaa	attgggacgc atcttgggtg tgaatgatgc tgcccactct ttggggagc 2460
gttacagtag	ccgtcagcgt accggtttgt aaactgatgt ggctatcttt tcaactcatg 2520
ccgtgaaaaa	tgtgactact gccgaagggt gggccatttg cctgaatctg cctaagccgt 2580
tcgataatac	ggagttgtat aaagagctac ggatgacaag cttgaactgc cagaccaagg 2640
atgctttttc	gaaatctaaa gccggaggat gccgttatga tattgtcgtt ttccgggagta 2700
aaatcaatat	ggcagatgta aatgccgcca taggactggc acagatacga gaatatccgg 2760
aatgtctgaa	agacgaaaag cgggtgttta atgcttatag cgatgctttt tcggcttgtg 2820
actgggcat	cgttccgctt tctgtggatg gggaaaaagga aagctcttat cacatttatg 2880
ctttgcgtat	taaggacttt accgaagaac aacgcgacgg gatgattgat gaaatcgcta 2940
aaagtgaagt	ggctgtcaat gtgcacttta taccgatgcc gatgttgtcg tcttcaaat 3000
cgatgggata	tgatataaag gattatccac aggcctatca gaactttaag agcgagatat 3060
cattgccgat	atatcccaaa ttggatagtg aaaaattgaa ttctattatt gaaaccgtaa 3120
aggcggccta	tgccgacgta attgcaaaa accgacacca ccaccaccac cactagagga 3180
ggatatacat	ttgatccgtt tttttgatat cgttttttcc cttttgggaa ttcttctgct 3240
ttctcccggt	ttcttctctt ttatatagc catttgtctc gaaagtaaaag gaggtgggtt 3300
ctataaacag	ctccgtgtgg gaaggtagcg aggggatttt tatgtttata aatttcgttc 3360
gatgcgtgta	ggggctgata aaaaggggtt gataacggta ggaggggagag atccgagaat 3420
aactcgtacc	ggatacctga tccggaagta taaattggat gaattgcctc agttgtttta 3480
cgtactgaaa	ggagatatga gtctggtagg gcccgctcct gaggttcgta agtatgtgga 3540
tttatatacc	gatgaacaaa agaaagtatt gagtgcacgt ccggggatta cggattatgc 3600
ttctatttag	tatgtagatg aaaatatgat tcttgagag gcttcggatc ctgacagagc 3660
atatatagaa	cagattatgc ccgataagat acgggtataat atgaaatata tctgtaatcg 3720
ttcgggtgaag	gaatatttta aatttatatt tctgacattt tggagcatca ttcgttagga 3780
gctcaggagg	atatacatat gaaattggct gtcattttct ctatctacaa aaatgattcg 3840
ctttctaaaa	taaagcaatc cttacaaaag ttgttgaaac aagaaaatgt tactttgtgat 3900
attttattat	attgttgatg tctgttttcg attgatattg tgcatttctt agactcattt 3960
gatcaaaact	cactttatat atttaagcaa gaacaaaata gaggattggc tcatgcacta 4020
aatgaattat	tgaatatcgt tcttacaataa gaatacgaat atattgctcg tatggatgct 4080
gatgatattt	ccatgccgga acgtttttgt aagcagatcg catttatgaa ttcacatccg 4140
gatattgatt	gccttggtac ttgggcaata atgacggaaa agaataattt 4200
cgaaaaaaaa	tgccgataac tcatgaagaa tgtttggaac tgtttaaaaa gcgggattgt 4260

-continued

atgattcatc	ctactgtgat	gtttcgacgt	tcataatttg	agaaagctgg	tttatatccc	4320
gaagacacct	actttggaga	agatactatg	atgtgggcta	aaggttttaa	aagtggatgt	4380
aaatttgcca	atgtaccoga	atatctgttt	aagtttagat	tggattcaaa	cttctttgaa	4440
agaaggcggg	ggtggaaaca	tgcaaaatct	atttatactt	tacgtcatcg	tgtaaataga	4500
atggttaggt	tcggatggaa	ggaagattgt	tatgcattgc	tatatgctat	ggccaaatta	4560
atgcctaaat	ctattcttga	cataatctat	aaaacagtac	gttgaaggag	gatatacata	4620
tgaggaagat	attattaaca	tatggagata	ttaaaacat	aaatattggt	gattatattc	4680
agagtatcgc	cgctaaaaca	ttttttgatg	atgataatta	tatatatttc	aatcgtgatg	4740
aattaaagtt	gtataaagg	gaacctgcaa	aagtcataat	gaatgcttgg	atgacttata	4800
aaccataata	ttggcctcct	tcattctcagg	tttatccatt	atttgttgca	ttacatatca	4860
attcttctgc	agaaagtcgc	ttttttgtcac	atgatagtat	taagtattta	aaaaagtatg	4920
aaccocatag	gtgcccgcgat	tatcatacaa	tgaatatttt	aaaaggaaaa	ggtgttaatg	4980
cttatttttc	gggatgtcct	actactactt	tggttaaaac	ttataaatat	aatggaaaaac	5040
gtgaggggat	atacatgtgt	gatccacttt	catacatgcc	taatgggaat	aatttttttg	5100
aaataatgaa	agctgtgtga	caaactgttt	tttatatgaa	gcctgtgcta	aagattccta	5160
gaaattataa	aaagaataat	agatttacta	ttaatatatc	caaagtagga	ataggacgtt	5220
tattgttaat	aaaccaagtct	tattttgcttt	tgagaaagct	agtagttcct	gacgtattat	5280
acaatgcaat	atttataaca	caatttaata	tgtctaatga	atattctctc	gaatcagaaa	5340
gatttgctcg	ggctgatgaa	tatttaacta	aaatggcctc	agctcaaat	gtgataaact	5400
ctagaatcca	ctgtgctcct	ccttgcttag	gctttgagac	ccctgtgggt	tattattaga	5460
atctatcaga	aagcaagaag	agcacttgct	gattaggggg	acttgaatct	ctatttaatg	5520
tgataactgt	gaagggggaa	aatggttact	ccaatttttt	tgatggacta	tttaaacgag	5580
attcttcttt	taagaataaa	attgattttg	tagaatatag	agatcgatta	attaatatat	5640
gtaataaatt	tatgatgtca	tgaaggagga	tatacatatg	agggatggaa	agcccatgta	5700
aatactgcat	attgtaggga	ctagacctgt	tggaggaatt	ggtgctttat	tgaaaaatat	5760
aaacacttct	attgatatta	ataaattcca	ttttacatat	gtgttttctg	ccgataagtaa	5820
tattggagat	tttgataaatt	attgtgagaaa	attaggtagc	gatattgttg	tattcccttc	5880
atatcatttg	aaatattttat	ttttatattt	aaagtgattt	ttctgttttt	ataagagaaa	5940
tgcaaaaaaa	tatgatataa	ttcatgtaca	tagtgctaat	acaggagtat	tagacctttt	6000
attttcgaag	atatattggaa	tacggataag	gattttacat	agtcacagta	caaaatattc	6060
tagtaaaaaa	ataaggtcca	ttagaaatta	ttttttgcaa	tttctacaaa	tatttttagc	6120
aaacacttat	tttgcattgt	gatacaaaag	tgctgaattc	ctttttggaa	aaaagagact	6180
tgaaaaagtt	tattattatc	ataatgcaat	tttttctaaa	aaattttatt	ataatgcaac	6240
tgtagagag	agagttagag	ttgagttgaa	aatagagaat	gactgtttac	tgtaggggca	6300
tatcggtaac	tttacgaaag	aaaagaatca	tggtttttta	atagatatat	tggaagaatt	6360
attgaatatt	aatgacaatg	ttaagttgct	gttagtaggg	gatggccaac	ttagaagtga	6420
aatagaggaa	aaagtaaaat	taaaaagatt	gcagaagtat	gtgtgctttt	taggtcggag	6480
aactgatata	tcagaattat	tacaaagcat	ggatttttta	atttttccct	cttttttttg	6540
aggctacca	gtgactctga	ttgaagcgca	agctctcggt	ttaagatgtg	tcgtttcgga	6600
tcgtattact	ttggaaccaa	agataacgaa	taataatagt	tatcttaatc	tcaatcgaga	6660
tccttcaact	tggtgtgaag	ttgtttggaa	gttgtttgaa	gataaaactt	ttatcaggaa	6720
agatacgtct	aaagaaatat	gtgaatcagg	atatgatata	catttagaga	gtgaatatct	6780
cgaaagaata	tatttaaatc	ttttacataa	atgaaggagg	atatacatat	gaccaaatat	6840
gaaaaaaatc	aacaagattt	agttgaatca	ccaaaaacat	gggtgataac	cgggtgttgc	6900
gggtttattg	gctccaattt	actagaaaaa	ctactaaaag	ttgatcaaac	cgtggtgggg	6960
ttagataact	ttgtccacgg	tcatacaaat	aaactagatg	aagttaaaag	tttagtaaca	7020
gaagcccaat	ggcaacgatt	tacctttatt	gaaggcgata	tacgtaatcc	tgagacctgt	7080
gtagcagcgg	ttcaaggtgt	agactatgta	ttacaccaag	cggcactagg	gtctgtgccc	7140
cggttcaatta	atgatcctat	tactacaaat	tcgcctaaca	tcactgggtt	tttaaacatg	7200
ttagtagcag	ccaagatca	gcaagtgaat	agctttactt	atgcgcctc	aagttcgagt	7260
tatggtgatc	acctgcatt	gcggaaggta	gaagaaaata	ttggcaaccc	gttatcccc	7320
tacgccgtaa	ccaaatacgt	aaatgaactt	tatgcgaatg	tattttgcgc	gacttatggt	7380
tttaaaagta	ttggactcgg	ttattttaat	gtgtttggaa	aacgccaaag	ttccaaatggt	7440
gcctatgcgg	ctgttatttc	aaaattggga	gcagcaatga	ttcaaggtag	tggtattttat	7500
attaatggtg	acggcgagac	tagtcgcgat	ttctgttata	tagctaatac	agtgcaaatg	7560
aatattcttg	ccgccaccgc	accagaagaa	gcaaaagaca	agattttaca	tggtgcagta	7620
ggagacagaa	caaccttaaa	cgatctattc	tctgccatta	aattggcatt	atcgaataat	7680
actgttgttg	tttctcaaga	acctattttat	agagagtttc	gtactggtag	tgtaaggcat	7740
tctcaggctg	atgttaataa	agctaagcag	tatttaggct	atgcgcgctc	gcaccgtatt	7800
ctagatggaa	tcaacgaagc	aatgccatgg	tataataagct	ttataaatga	taaaaaataa	7860
aggaggatat	acatatgaaa	aaaaaatatg	actatctaag	tgctggagcc	ggactttatg	7920
gtttctgttt	tgcatataaa	gcgcaacgga	atgggaaaaa	atgtttgctt	attgacaaac	7980
gtccacacaa	tggaaggaaat	atttatttgcg	aaaattatga	gggtataaat	gtacacaaat	8040
atggtgcaca	tatttttcat	acttctaata	aagaggtttg	ggactttgta	aatgccatag	8100
tggaatttaa	ccgctatact	aatttccccta	ttgccaatga	taaaaggcaa	ttatataatt	8160
taccttttaa	tatgaatact	ttttatgctt	tatgggggac	aaaaactcca	gagggaagcaa	8220
aatgaagat	agaagagcaa	aggagagaa	cggggattct	tacgccgaag	aatctagaag	8280
aacaggctat	ctcttttagta	ggaaaggata	tttatgaaat	attgataaaa	ggatataact	8340
agaaacagtg	ggggcgga	gcaactgaac	ttcctgcttt	tataatcaag	cgtttacctg	8400
tcagatttac	ttttgataat	aattacttta	atgataaata	tcaggggatt	cctgttggtg	8460
gctataataa	attgataaat	ggtttattag	atggtattga	agttagaaca	aatacacagt	8520
tttttcaaaa	taagaatat	tttaaaggct	ttgcggataa	aattctattt	actggcaaaa	8580
ttgatgaata	ttataattat	caatttggga	gattggaata	tcgtactggt	gaatttgaaa	8640
cacaagtgtc	tattgtgtag	aattatcaag	ggaatgcagt	tgtagaattat	actgaaaggg	8700
aagtgcctta	tactcgatt	atagagcata	aacattttga	atttggaaca	cagcccaaaa	8760
ccgtaatttc	caaagaatat	tcttcggaat	ggaaagatgg	tagtgaaccc	ttttatcctg	8820

-continued

ttaatgatga	gaggaaataat	tctttgtatt	tggagtataa	gaaattgggtc	gatagagaga	8880
agaatgtttt	gtttggaggg	agattagctg	agtataagta	ttagtatatg	aatgtgattg	8940
tggaaaaagt	tataaatagt	gacttatgaa	ggaggatata	catatgaaga	tattttgctgt	9000
agttgtgacc	tataatcggt	tagcattact	aaaaaaggtc	atagatttac	ttaaaagaca	9060
gacacgtcaa	ttagatagta	tagttatagt	aaataatggc	tctactgatg	gaacaaaaga	9120
atgggttagaa	caggagaaaa	atataacatt	aataaatcaa	tctaattctg	gtggggctgg	9180
tggattcgaa	acggggggtaa	aatatgcata	tgagaatgag	gcagattgga	tttggatgat	9240
ggatgatgat	gtttttccta	atatagattg	tttggaaaaa	ctcttagggat	ggacgagtat	9300
ttctcaatgt	atccaaccac	gtagggtatta	ttcggatgat	gtagaagtaa	attttgaaca	9360
atggttggat	ccaattactt	atagtaaat	tggttattgg	caagaaaaat	catttaataa	9420
tggtaaaaaa	ttttgtgcta	ttaatgtagg	ctgttttgaa	gggatgtttg	tttcgaaaaa	9480
cattgttaat	aaaataggat	ttcctgataa	gcgttttttt	atagcggaa	atgatacaat	9540
atatgggttt	gcagctagtt	ttataccaa	cgtaatttta	gtatctgatg	ctatcatggt	9600
tcgtgctaga	agatcactgg	atcgtagcgt	tagcccaatg	tatacttatt	atgcattgctg	9660
gaattttcat	ttagtatatg	agtcacttaa	tttattattg	gataagaaaa	atagggtttt	9720
atataaaaa	tataattatc	aatttgttca	tcagatttat	ttatctattt	ttttatatga	9780
taataagatg	aaacacttaa	tatctgtttt	tcgtggcctt	tatgattgct	ttcgtaaaaa	9840
aaatggagct	acctattaaa	ggaggatata	catatgggac	aatacaataa	gaataacttc	9900
ctactgaatc	taagtactac	aattactggg	ctattatttc	cactaataac	ttttcctcat	9960
gcttcacgta	tattaatggc	agatgggtata	gggcaagtac	aatttttcca	atccataatt	10020
gactatgtct	ctctttgtac	tgctttggga	atccctttgt	atgcgcttag	agaaattgca	10080
agaataaggg	ataataagga	attaagaagt	agaacacaaa	tcgaaatact	attgcttcat	10140
gctattttga	cttttagtcg	atatatagtt	gtttttatata	ttgctaagac	cgttgccaaa	10200
atagaaatag	atgcttctct	ttcttttttg	ttaagtacaa	ctttattttt	taacacgata	10260
ggagttgcat	gggttttatca	ggcaattgaa	gattttaaat	acataacttt	acgttccotta	10320
tttgaagga	tactatcttt	agtgccotta	tttatctttg	ttaaagaccaa	gcaagatctg	10380
ttttattatg	ctggaattatt	agttatttgt	actgttggtta	ataatatatt	taattttttt	10440
cgattacgta	aatatatata	aactagttaa	ggcgaattta	agcgtttgaa	ctcgttgagg	10500
catttaatac	ctgcacttaa	aattttttata	ttaaaatttg	ttataagtat	ttatgtgaat	10560
cttgattctg	tgatgcttgg	atttctttaa	aatgaagaat	cagttgggtta	ttatgctgct	10620
gctactcgac	ttacaaaagc	tttttgggtt	atagtttctt	cattaggagc	tgtttttatta	10680
cctcgtttta	gtaatatgat	aactaatggt	caaaaagaag	aattcccaat	attggcaaat	10740
aaagctgcta	gtttttacaat	tgctttaagt	cttcctatga	gtgtgggact	tattttttatg	10800
gcagcaccta	ttatcacatat	tttttgggga	aatgggtttg	aaccttctat	tttgacatta	10860
aaattagttg	cacctattgt	ttttatttatt	ggattatctg	gaattattgg	aatgcaataa	10920
ttgtatccgc	aagggagaga	aaagtagcta	attatatcta	ctatggtagg	agcatgtata	10980
aatctcctta	ttaactattt	actaatacc	caatatgggc	agtatggagc	ggcgtttggg	11040
actgttatcg	cagaatttat	ggtaacgggt	ataatgattt	tgctgggaag	aaaaatttta	11100
cctataaata	ttctctccaa	gcagaattta	cactatctaa	taggttctat	agtgatattc	11160
attctattag	ctttttttgt	tgcttccctt	ttacatgaag	ttaattattt	attgataggt	11220
atactgttat	ctgttattgt	ttattacgca	tattttattaa	tgataaaaaga	taccctcgca	11280
ttacaactaa	agaattattt	actttcaata	tttaaacat	gaaaggagga	tatacatatg	11340
actagtactt	ctttctttat	tttagtgctt	atatctttta	tagtatttgt	ttataattta	11400
gcaggttctg	attggagtaa	tctttctcga	gtattgttat	ttataatttc	gtttttttgtc	11460
tttttattag	taacaaatgt	atatcaaaaga	aaagtaaaaa	tagtaacact	agggtatttg	11520
gtctatttct	ttttggcctt	tttagctcaa	aatgtaggtc	catatgcaat	tattttacaat	11580
tctcacatac	agaatttgat	acagagggtt	cctaataatt	gtgtaccaa	tgatcattac	11640
actttatact	ttcttatttt	tattttttgt	tcaactgtta	tttatattta	ttcttttatgt	11700
attgataaaa	tttctattgt	gcagaaagta	tggttatata	tgccatacaag	acaagaagct	11760
cgaaaaataa	tgatggcatt	ttttataatc	ttgttgcccta	tatggatatt	agggaaatagg	11820
tcctatgcta	caattctagt	tccttttaca	agttatttta	tcatttctat	ttggaaatat	11880
cccttttcaa	ggaaaaatag	agtatttatt	attgggttta	ttgcactcgt	attgatccct	11940
atttcgcaat	tattttctcg	gttttatatt	gttcaatatg	tgttcccttt	tatttttaatt	12000
tgggtttttga	ataatggatt	gctattttca	agaaaagaaa	aaataagtta	taaggttat	12060
ggcttttttc	tattgttttg	ttcttttagc	attttgtatg	gtatcatttc	agagatgatg	12120
aagttaaatg	caaattttga	tggaaactat	actttcaatg	atatgtttga	agttattaagt	12180
gagccgcaat	tattgcaaaa	ttgggttatc	aggcaaacat	atcgattttt	tgatattttg	12240
tcacatttgg	gggggaatat	aatagattat	attgaccaga	aaggatattt	atgggggaatt	12300
acatatgtaa	agagctttgc	acctatatta	ggttttgatt	atgtaagcct	ttcattaatt	12360
tcagcaataa	tgatttggtg	agattatgct	caacctggtt	tggtggcaga	agggtacgct	12420
aaattgggaa	tatatggggc	tataattaat	atgatatttg	tttttttcgt	tgacaggtta	12480
ttgtttgatt	attttttacg	aaaacaactc	accttaaaat	tggttacttta	tattgggaact	12540
ttttccagg	tattactaga	tgggggggac	attaattcta	ttattttcat	gtctatattt	12600
tgtttcttaa	cttgtttgtt	taatgccaaa	atcaaatttt	aaagctctaa	ttagctgagc	12660
ttggactcct	gttgatagat	ccagtaaatga	cctcagaact	ccatctggat	ttgttcagaa	12720
cgctcgggtg	ccgcggggcg	ttttttattg	gtgagaatcc	aagc		12764

1. A method for producing a polysaccharide in a bacterium, the method comprising expressing in the bacterium a plurality of coding sequences, the plurality of coding sequences comprising nucleotide sequences that encode a pglF dehydrogenase gene product, a wbpP UDP-N-acetyl-d-glucosamine C4 epimerase gene product, a wcfR amino-transferase gene product, and a wcfS phospho-glycosyl-

transferase gene product, and optionally further comprise nucleotide sequences encoding one or more of a wcfQ glycosyltransferase gene product, a wcfO pyruvyltransferase gene product, a wcfP glycosyltransferase gene product, a wcfM UDP-galactopyranose mutase gene product, a wcfN glycosyltransferase gene product, a wza polysaccharide export protein gene product, a wzx fippase gene product, a

wzy polymerase gene product, and a wzz gene product, and further wherein at least one of the coding sequences is heterologous to the bacterium.

2. The method of claim 1, wherein the polysaccharide is a capsular polysaccharide A (CPSA) and the bacterium expresses each of the PglF gene product, the wbpP gene product, the WcfR gene product, the WcfS gene product, the WcfQ gene product, the WcfO gene product, the WcfP gene product, the WcfM gene product, the WcfN gene product, the wzz gene product, the wzx fippase gene product, and the wzy polymerase gene product, at least one of which is heterologous to the bacterium.

3. The method of claim 1, wherein the bacterium is an *Escherichia coli* (*E. coli*) bacterium, optionally wherein the *E. coli* bacterium has reduced or absent waal biological activity.

4. The method of claim 1, wherein:

the pglF gene product is a *Campylobacter jejuni* gene product; and/or

the wbpP gene product is a *Vibrio vulnificus* gene product; and/or

the wcfR gene product is a *Bacteroides fragilis* gene product; and/or

the wcfS gene product is a *Bacteroides fragilis* gene product; and/or

the wcfQ gene product is a *Bacteroides fragilis* gene product; and/or

the wcfO gene product is a *Bacteroides fragilis* gene product; and/or

the wcfP gene product is a *Bacteroides fragilis* gene product; and/or

the wcfM gene product is a *Bacteroides fragilis* gene product; and/or

the wcfN gene product is a *Bacteroides fragilis* gene product; and/or

the wza gene product is an *Escherichia coli* gene product; and/or

the wzx fippase gene product is a *Bacteroides fragilis* gene product; and/or

the wzy polymerase gene product is a *Bacteroides fragilis* gene product; and/or

the wzz gene product is a *Bacteroides fragilis* gene product.

5. The method of claim 1, wherein the polysaccharide is present in a lipid bilayer of the bacterium, optionally wherein the lipid bilayer is part of the bacterium's cell wall.

6. The method of claim 5, wherein the polysaccharide is a CPSA, optionally an aggregate of two or more CPSA tetrasaccharide monomers.

7. The method of claim 1, wherein the plurality of coding sequences are transcribed to produce a polycistronic messenger RNA (mRNA) molecule in the bacterium.

8. The method of claim 7, wherein the polycistronic mRNA molecule comprises a ribosome-binding site (RBS) operatively linked to a translation initiation codon of each of at least two of the plurality of coding sequences, optionally wherein the polycistronic mRNA molecule comprises a plurality of ribosome-binding sites (RBSs), with an RBS present 5' to each of at least two, optionally 5' to each, translation initiation codon present in the polycistronic mRNA, wherein at least one RBS is located 3' to a translation termination codon of a first coding sequence and 5' to a translation initiation codon of a second coding sequence present on the polycistronic mRNA molecule.

9. The method of claim 1, wherein the bacterium does not express a functional waal O-antigen ligase gene product.

10. A bacterium comprising a plurality of coding sequences, the plurality of coding sequences comprising nucleotide sequences that encode two or more of a pglF dehydrogenase gene product, a wbpP UDP-GlcNAc C4 epimerase gene product, a wcfR aminotransferase gene product, a wcfS phospho-glycosyltransferase gene product, a wcfQ glycosyltransferase gene product, a wcfO pyruvyltransferase gene product, a wcfP glycosyltransferase gene product, a wcfM UDP-galactopyranose mutase gene product, a wcfN glycosyltransferase gene product, a wza polysaccharide export protein gene product, a wzx fippase gene product, a wzy polymerase gene product, and a wzz gene product, wherein at least one of the plurality of coding sequences is heterologous to the bacterium.

11. The bacterium of claim 10, wherein the bacterium is an *E. coli* bacterium.

12. The bacterium of claim 10, wherein:

the pglF gene product is a *Campylobacter jejuni* gene product; and/or

the wbpP gene product is a *Vibrio vulnificus* gene product; and/or

the wcfR gene product is a *Bacteroides fragilis* gene product; and/or

the wcfS gene product is a *Bacteroides fragilis* gene product; and/or

the wcfQ gene product is a *Bacteroides fragilis* gene product; and/or

the wcfO gene product is a *Bacteroides fragilis* gene product; and/or

the wcfP gene product is a *Bacteroides fragilis* gene product; and/or

the wcfM gene product is a *Bacteroides fragilis* gene product; and/or

the wcfN gene product is a *Bacteroides fragilis* gene product; and/or

the wza gene product is an *Escherichia coli* gene product; and/or

the wzx flippase gene product is a *Bacteroides fragilis* gene product; and/or

the wzy polymerase gene product is a *Bacteroides fragilis* gene product; and/or

the wzz gene product is a *Bacteroides fragilis* gene product.

13. The bacterium of claim 10, wherein the plurality of coding sequences are transcribed in the bacterium to produce a polycistronic messenger RNA (mRNA) molecule in the bacterium.

14. The bacterium of claim 13, wherein the polycistronic mRNA molecule comprises a ribosome-binding site (RBS) operatively linked to a translation initiation codon of each of at least two of the plurality of coding sequences, optionally wherein the polycistronic mRNA molecule comprises a plurality of ribosome-binding sites (RBSs), with an RBS present 5' to each of at least two, optionally 5' to each, translation initiation codon present in the polycistronic mRNA, wherein at least one RBS is located 3' to a translation termination codon of a first coding sequence and 5' to a translation initiation codon of a second coding sequence present on the polycistronic mRNA molecule.

15. The bacterium of claim 10, wherein translation of the polycistronic mRNA molecule in the bacterium results in production of a polysaccharide in the bacterium, optionally

wherein the polysaccharide is present in a lipid bilayer of the bacterium, further optionally wherein the lipid bilayer is part of the bacterium's cell wall.

16. The bacterium of claim **15**, wherein the polysaccharide is a CPSA, optionally an aggregate of two or more CPSA tetrasaccharide monomers.

17. A method for producing an immunogenic composition, the method comprising mixing an antigenic molecule with a capsular polysaccharide A (CPSA), wherein the CPSA is produced by a bacterium of claim **10**.

18. An expression cassette comprising a plurality of coding sequences derived from at least two different bacterial species, wherein the plurality of coding sequences are selected from the group consisting of a *pglF* dehydrogenase gene coding sequence, a *wbpP* UDP-N-acetyl-d-glucosamine C4 epimerase gene coding sequence, a *wcfR* aminotransferase gene coding sequence, a *wcfS* phosphoglycosyltransferase gene coding sequence, a *wcfQ* glycosyltransferase gene coding sequence, a *wcfO* pyruvyltransferase gene coding sequence, a *wcfP* glycosyltransferase gene coding sequence, a *wcfM* UDP-galactopyranose mutase gene coding sequence, a *wcfN* glycosyltransferase gene coding sequence, a *wza* polysaccharide export protein gene coding sequence, a *wzx* flippase gene coding sequence, a *wzy* polymerase gene coding sequence, and a *wzz* gene coding sequence.

19. The expression cassette of claim **18**, wherein the expression cassette comprises a combination of coding sequences selected from the group consisting of:

pglF, *wcfR*, and *wcfS*;
pglF, *wcfR*, *wcfS*, and *wcfQ*;
pglF, *wcfR*, *wcfS*, *wcfQ*, and *wcfO*;
pglF, *wcfR*, *wcfS*, *wcfQ*, *wcfO*, *wbpP*, and *wcfP*;
pglF, *wcfR*, *wcfS*, *wcfQ*, *wcfO*, *wbpP*, *wcfP*, *wcfM*, and *wcfN*; and
pglF, *wcfR*, *wcfS*, *wcfQ*, *wcfO*, *wbpP*, *wcfP*, *wcfM*, *wcfN*, *wzx*, and *wzy*.

20. The expression cassette of claim **18**, wherein expression cassette is operatively linked to a promoter such that transcription from the promoter results in the production of a polycistronic mRNA that is translatable to produce polypeptides encoded by the coding sequences.

21. The expression cassette of claim **20**, wherein the polycistronic mRNA molecule comprises a ribosome-binding site (RBS) operatively linked to a translation initiation codon of one or more of, optionally each of, the plurality of coding sequences, wherein at least one RBS is located 3' to a translation termination codon of a first coding sequence and 5' to a translation initiation codon of a second coding sequence, wherein the first coding sequence and the second coding sequence are adjacent to each other on the polycistronic mRNA molecule, the first coding sequence being present 5' to the second coding sequence.

22. The expression cassette of claim **18**, wherein the polycistronic mRNA molecule is operably linked to a promoter that is active in the bacterium, optionally wherein the promoter is an inducible promoter, optionally an isopropyl- β -D-thiogalactopyranoside (IPTG)-inducible promoter, further optionally wherein the promoter is a T5 promoter.

* * * * *