

Fig. 1





Fig. 2



METHODS OF TREATING MALIGNANT PERIPHERAL NERVE SHEATH TUMOR (MPNST) USING LSD1 INHIBITORS

FIELD

[0001] The present invention relates to the field of malignant peripheral nerve sheath tumor (MPNST) therapy. In particular, the invention provides an LSD1 inhibitor for use in the treatment of MPNST. The invention likewise provides methods of treating MPNST in a subject in need thereof, comprising administering a therapeutically effective amount of an LSD1 inhibitor to the subject.

BACKGROUND

[0002] Cancer is one of the leading causes of death worldwide. The way cancer is clinically managed has radically changed over the last decades. In the past, cancer could only be treated with chemotherapeutic “dirty” drugs, which unspecifically limit cell proliferation by interfering with DNA or the basic cell-cycle machinery. Nowadays, new personalized and precision medicine approaches are instead designed to block the growth of cancer cells while mostly sparing other cells of the body. The advantage of this approach is that targeted therapies are less harmful to normal cells.

[0003] Epigenetics is one of the emerging fields in cancer precision medicine, with a first generation of drugs reaching FDA-approval and many more progressing in clinical trials. Lysine specific demethylase 1 (LSD1 or KDM1A) is an epigenetic enzyme regulating gene expression by demethylating the histone H3 tail on two different residues, i.e. lysine 4 (H3K4) and lysine 9 (H3K9), with opposing effects. Demethylation of H3K4 is associated with transcriptional repression whereas demethylation of H3K9 is associated with transcriptional activation. Additionally, LSD1 is part of many multiprotein complexes controlling enhancer-promoter contacts involved in gene repression such as NurD and CoRest. LSD1 has been shown to play a key role in cancers such as leukemia and small cell lung cancer (SCLC), and huge efforts have been committed to the development of LSD1 inhibitors. Catalytic active-site targeted inhibitors have been developed such as iadademstat, bomedemstat and pulrodemstat. These compounds bind deep in the active site of LSD1, blocking access to both protein substrates (such as the histone H3 tail) and non-substrate protein interactors (such as SNAG-domain transcription factors), thereby inhibiting both LSD1 catalytic activity and scaffolding interactions. LSD1 inhibitors have been described to have highly potent anti-proliferative activity in specific tumor types and are currently being tested in clinical trials as treatment for cancers such as acute myeloid leukemia (AML) and SCLC.

[0004] Malignant Peripheral Nerve Sheath Tumor (MPNST) is a highly aggressive tissue sarcoma. It can arise in at least three different contexts. Around 40-50% of MPNST cases occur in patients of neurofibromatosis type I, which is a dominant autosomal syndrome caused by mutations in the NF1 gene, predisposing patients to learning disabilities, musculoskeletal disorders, and a tendency to develop small benign tumors (neurofibromas). However, mutation of the NF1 gene is not sufficient to drive MPNST in neurofibromatosis type I patients and a series of further mutational events must accumulate, such as loss of function mutations in tumor suppressor genes other than NF1 (such

as CDKN2A, p53, or PTEN) and/or mutations or copy number variations in oncogenes (such as EGFR or c-Met). Moreover, in many cases there is also loss of Polycomb Repressive Complex 2 (PRC2) activity, the multiprotein complex involved in, among other processes, the methylation of histone H3 lysine 27 (H3K27), a known epigenetic post-translational modification usually associated with transcriptional repression. Around 40% of MPNSTs are sporadic, with no known genetic cause. The rest, about 10%, develop as a result of previous radiotherapy. Generally, MPNST has an aggressive clinical course, with 50% probability of relapse and an overall average 5-year survival rate of only 20-40%. Advanced disease often leads to lung and bone metastases. There are currently no drugs specifically approved for the treatment of MPNST. While various different drugs have been tested in clinical trials in MPNST, these drugs have displayed disappointing response rates, with few MPNST patients achieving stable disease at best (see, e.g., Martin E et al., *Crit Rev Oncol Hematol*, 2019, 138:223-32, doi:10.1016/j.critrevonc.2019.04.007). Therefore, there is a huge unmet medical need to find new and more effective treatments for MPNST. The present invention addresses this and other needs.

SUMMARY OF THE INVENTION

[0005] The present invention is based on the surprising finding that LSD1 inhibitors, such as e.g. iadademstat, bomedemstat and pulrodemstat, are advantageously effective in the treatment of MPNST, as also described in Example 1 below.

[0006] This was completely unexpected, particularly as MPNST is a malignant tumor which is highly aggressive and enormously difficult to treat. The present invention thus provides a novel and improved therapeutic approach for the treatment of MPNST. Accordingly, the present invention relates to an LSD1 inhibitor for use in the treatment of MPNST.

[0007] The invention also relates to a pharmaceutical composition comprising an LSD1 inhibitor and optionally one or more pharmaceutically acceptable excipients for use in the treatment of MPNST.

[0008] The invention likewise provides a method of treating MPNST in a subject in need thereof, comprising administering a therapeutically effective amount of an LSD1 inhibitor (or a therapeutically effective amount of a pharmaceutical composition comprising an LSD1 inhibitor and optionally one or more pharmaceutically acceptable excipients) to the subject.

[0009] Moreover, the invention relates to the use of an LSD1 inhibitor for the treatment of MPNST.

[0010] The invention further relates to the use of an LSD1 inhibitor for the preparation of a medicament (or a pharmaceutical composition) for the treatment of MPNST.

BRIEF DESCRIPTION OF THE DRAWINGS

[0011] FIG. 1: Design of the 9×9 matrix assay. Increasing concentrations of iadademstat in 1:2 dilutions were added from top to bottom while increasing concentrations of the second agent of the combination in 1:2 dilutions were added from left to right. The darker the grey, the higher the concentration of each agent. Wells corresponding to the diagonal of the plates (horizontal lines) indicate fixed EC₅₀

ratios. The concentration corresponding to the expected EC_{50} value was centered horizontally and vertically on the plates for each agent.

[0012] FIG. 2: Design of the 5×5 matrix assay. Increasing concentrations of iadademstat in 1:5 dilutions were added from top to bottom while increasing concentrations of the second agent of the combination in 1:5 dilutions were added from left to right. The darker the grey, the higher the concentration of each agent. Wells corresponding to the diagonal of the plates (horizontal lines) indicate fixed EC_{50} ratios. The concentration corresponding to the expected EC_{50} value was centered horizontally and vertically on the plates for each agent.

DETAILED DESCRIPTION OF THE INVENTION

[0013] As explained above, the present invention is based on the surprising discovery that LSD1 inhibitors are advantageously effective in the treatment of MPNST, as further detailed herein below and in the examples section. In particular, an extensive panel of 12 different MPNST cell lines was tested in Example 1, which is representative of the different subtypes of MPNST encountered in clinical practice, as it includes both neurofibromatosis type I-linked MPNST and sporadic MPNST cell lines and features great genomic heterogeneity in terms of NF1 heterozygosity, ploidy, and mutational status of CDKN2A and PRC2. The exemplary LSD1 inhibitor iadademstat was found to be highly effective in a remarkably heterogeneous range of MPNST cell lines and, therefore, to be particularly well-suited for the therapy of MPNST. Further LSD1 inhibitors, having different chemical scaffolds and including both reversible and irreversible inhibitors of LSD1, were likewise confirmed to be effective in the treatment of MPNST, as also described in Example 1.

[0014] The present invention thus relates to an LSD1 inhibitor for use in the treatment of MPNST. The invention also relates to a pharmaceutical composition comprising an LSD1 inhibitor and optionally one or more pharmaceutically acceptable excipients for use in the treatment of MPNST. The invention likewise provides a method of treating MPNST in a subject in need thereof, comprising administering a therapeutically effective amount of an LSD1 inhibitor (or a therapeutically effective amount of a pharmaceutical composition comprising an LSD1 inhibitor and optionally one or more pharmaceutically acceptable excipients) to the subject. Moreover, the invention relates to the use of an LSD1 inhibitor for the treatment of MPNST. The invention further relates to the use of an LSD1 inhibitor for the preparation of a medicament (or a pharmaceutical composition) for the treatment of MPNST.

[0015] In accordance with the present invention, an “LSD1 inhibitor” refers to a compound that reduces, decreases, blocks or inhibits the gene expression, activity or function of LSD1. Examples thereof are provided below under the heading “LSD1 inhibitors”. Preferred LSD1 inhibitors include each one of iadademstat or a pharmaceutically acceptable salt thereof (e.g., iadademstat dihydrochloride), pulrodemstat or a pharmaceutically acceptable salt thereof (e.g., pulrodemstat besylate), and bomedemstat or a pharmaceutically acceptable salt thereof (e.g., bomedemstat bis-tosylate). A particularly preferred LSD1 inhibitor is iadademstat or a pharmaceutically acceptable salt thereof (e.g., iadademstat dihydrochloride).

[0016] The LSD1 inhibitor (e.g., iadademstat or a pharmaceutically acceptable salt thereof) is preferably administered orally. Exemplary formulations which can be administered orally, particularly via peroral ingestion, are described in more detail further below.

[0017] The subject to be treated in accordance with the invention may be a human being or an animal (e.g., a non-human mammal), and is preferably a human.

[0018] The MPNST to be treated in accordance with the present invention may be, e.g., neurofibromatosis type I-linked MPNST, sporadic MPNST, or radiation-induced MPNST.

[0019] As explained above, MPNST frequently occurs in subjects having neurofibromatosis type I, which is a hereditary disease. Thus, in some embodiments, the MPNST to be treated in accordance with the present invention is neurofibromatosis type I-linked MPNST (or MPNST associated with neurofibromatosis type 1). Accordingly, in some embodiments, the invention relates to the treatment of MPNST in a subject having neurofibromatosis type 1. The invention also relates to an LSD1 inhibitor (or a pharmaceutical composition comprising an LSD1 inhibitor) for use in the treatment of MPNST, wherein the LSD1 inhibitor (or the pharmaceutical composition) is administered to a subject having neurofibromatosis type 1.

[0020] In some embodiments, the MPNST to be treated in accordance with the present invention is an MPNST which is not neurofibromatosis type I-linked (or an MPNST which is not associated with neurofibromatosis type 1), particularly sporadic MPNST or radiation-induced MPNST. The invention thus specifically relates to the treatment of MPNST in a subject not having neurofibromatosis type 1. The invention also relates to an LSD1 inhibitor (or a pharmaceutical composition comprising an LSD1 inhibitor) for use in the treatment of MPNST, wherein the LSD1 inhibitor (or the pharmaceutical composition) is administered to a subject not having neurofibromatosis type 1. In some embodiments, the MPNST to be treated is sporadic MPNST. In some embodiments, the MPNST to be treated is radiation-induced MPNST (see, e.g., Yamanaka R et al., *World Neurosurg.* 2017, 105:961-970.e8, doi:10.1016/j.wneu.2017.06.010). The MPNST to be treated in accordance with the invention may furthermore be an MPNST (including, e.g., any one of the above-mentioned specific types of MPNST) having one or more mutations or genetic alterations affecting the NF1 gene, particularly one or more inactivating mutations or inactivating genetic alterations in the NF1 gene. Accordingly, the MPNST to be treated may be, e.g., neurofibromatosis type I-linked MPNST having one or more mutations or genetic alterations (e.g., one or more inactivating mutations or inactivating genetic alterations) in the NF1 gene, sporadic MPNST having one or more mutations or genetic alterations (e.g., one or more inactivating mutations or inactivating genetic alterations) in the NF1 gene, or radiation-induced MPNST having one or more mutations or genetic alterations (e.g., one or more inactivating mutations or inactivating genetic alterations) in the NF1 gene. Such inactivating mutations or inactivating genetic alterations in the NF1 gene particularly include loss-of-function mutations and lead to a decrease or absence of expression and/or stability and/or activity of its protein product neurofibromin. Moreover, such mutations or genetic alterations may affect one or both alleles of the NF1 gene.

[0021] In humans, the NF1 gene is located on chromosome 17q11.2 and codes for a protein product called neurofibromin. The canonical amino acid sequence of isoform 2 of human neurofibromin is 2839 residues long (see, e.g., Uniprot identifier P21359-1; <https://www.uniprot.org/uniprot/P21359.fasta>), and that of isoform 1 is 2818 residues long (see, e.g., Uniprot identifier P21359-2; <https://www.uniprot.org/uniprot/P21359-2.fasta>). These two isoforms are considered the most biologically relevant. Isoform 2 is found in most human tissues but is not present in neurons of the central nervous system. The NF1 gene is capable of generating other alternatively spliced isoforms by different combinations of its about 60 exons. More than 3000 germline mutations in the NF1 gene have been reported in the Human Gene Mutation Database (HGMD; <http://www.hgmd.cf.ac.uk/ac/index.php>) and more than 1000 somatic mutations in The Cancer Genome Atlas (TCGA; <https://www.cancer.gov/about-nci/organization/ccg/research/structural-genomics/tcga>); see also Scheer M et al., *Int J Mol Sci*, 2021, 23(1):352, doi:10.3390/ijms23010352. Many of the characterized mutations are loss-of-function, meaning they ultimately have a negative impact on the functionality of the protein product. At the gene level, all sorts of mutations have been described, including nonsense, missense, frameshift, indels (insertions or deletions), microdeletions, inversions, splice site variants, whole translocations and complex rearrangements. However, there is no clear pattern of localized mutational clustering within the NF1 gene. In order to identify mutations or genetic alterations in the NF1 gene, it is advisable to apply a mutation detection pipeline to characterize NF1 mutants reliably, where a series of different algorithms specifically fine-tuned to detect single nucleotide variants (SNVs), indels or translocations in next-generation sequencing (NGS) reads are used. The presence of mutations or genetic alterations in the NF1 gene can be assessed in a sample, e.g., a biopsy sample obtained from the subject. The domain architecture of neurofibromin is complex and comprises, from N-terminal to C-terminal ends: (1) an N-heat domain comprising, among others, the cysteine and serine-rich domain/GTPase activating domain (CSR/D) and the tubulin binding domain (TBD); (2) a GTPase-activating domain (GAP) related domain (GRD), which promotes the hydrolysis of active Ras-GTP to the inactive form of Ras-GDP; (3) a Sec14 homologous segment; (4) a Pleckstrin homology (PH)-like domain; and (5) a C-heat domain comprising a Heat-like repeat domain (HLR) and a C-terminal domain (CTD) where the Syndecan-binding domain (SBD) is found. Additionally, the Sec14, PH-like and HLR domains form part of the so-called leucine-rich domain (LRD). Dimerization sites are found interspersed within the N-heat domain (at the far N-terminal end and at the TBD) and especially within the C-heat domain. In principle, the MPNST to be treated in accordance with the invention may have one or more mutations or genetic alterations (particularly one or more inactivating mutations or inactivating genetic alterations) in the NF1 gene sequence for any one (or several ones) of the above-mentioned domains or segments of the NF1 gene product neurofibromin. Thus, for example, the MPNST may be an MPNST having one or more inactivating mutations located in the GTPase activating domain (GAD) related domain (GRD) of NF1. In some embodiments, the MPNST has one or more inactivating mutations located in the cysteine and serine-rich domain/GTPase activating domain (CSR/D) of NF1. In some embodiments, the

MPNST has one or more inactivating mutations located in the leucine-rich domain (LRD) of NF1. In some embodiments, the MPNST has one or more inactivating mutations located in at least one dimerization interface of NF1.

[0022] The MPNST to be treated in accordance with the invention may also be an MPNST having no mutations or genetic alterations affecting the NF1 gene, particularly an MPNST having the wild-type NF1 gene. Thus, the MPNST to be treated may be, e.g., sporadic MPNST having the wild-type NF1 gene, or radiation-induced MPNST having the wild-type NF1 gene.

[0023] Moreover, the MPNST to be treated in accordance with the invention may be an MPNST (including, e.g., any one of the aforementioned specific types of MPNST, such as neurofibromatosis type I-linked MPNST, sporadic MPNST, or radiation-induced MPNST) having one or more mutations or genetic alterations affecting (particularly reducing or suppressing) the expression and/or the activity of CDKN2A, p53, RB1, PTEN and/or PRC2 (particularly one or more inactivating mutations or inactivating genetic alterations in the CDKN2A, p53, RB1, PTEN and/or PRC2 genes) and/or one or more mutations or genetic alterations (including also, e.g., copy number variations) affecting (particularly activating or enhancing) the expression and/or the activity of EGFR, PDGFRA and/or c-Met (particularly one or more activating mutations or activating genetic alterations in the EGFR, PDGFRA and/or c-Met genes).

[0024] Thus, in particular, the MPNST to be treated may be an MPNST (including, e.g., any one of the above-mentioned specific types of MPNST, such as neurofibromatosis type I-linked MPNST, sporadic MPNST, or radiation-induced MPNST) having one or more mutations or genetic alterations affecting (particularly reducing or suppressing) the expression and/or the activity of CDKN2A, particularly one or more inactivating mutations or inactivating genetic alterations of CDKN2A. Accordingly, the MPNST to be treated may be MPNST associated with CDKN2A inactivation, including, e.g., neurofibromatosis type I-linked MPNST associated with CDKN2A inactivation, sporadic MPNST associated with CDKN2A inactivation, or radiation-induced MPNST associated with CDKN2A inactivation. Such inactivation of CDKN2A (particularly biallelic inactivation of CDKN2A) may be due, e.g., to one or more point mutations, inversions, deletions or translocations of CDKN2A (see, e.g., Magallón-Lorenz M et al., *Hum Genet*, 2021, 140(8):1241-52, doi:10.1007/s00439-021-02296-x).

[0025] The MPNST to be treated may also be an MPNST (including, e.g., any one of the above-mentioned specific types of MPNST) having one or more mutations or genetic alterations affecting (particularly reducing or suppressing) the expression and/or the activity of p53, particularly one or more inactivating mutations or inactivating genetic alterations of p53. Accordingly, the MPNST to be treated may be MPNST associated with p53 inactivation, including, e.g., neurofibromatosis type I-linked MPNST associated with p53 inactivation, sporadic MPNST associated with p53 inactivation, or radiation-induced MPNST associated with p53 inactivation.

[0026] The MPNST to be treated may also be an MPNST (including, e.g., any one of the above-mentioned specific types of MPNST) having one or more mutations or genetic alterations affecting (particularly reducing or suppressing) the expression and/or the activity of RB1, particularly one or more inactivating mutations or inactivating genetic altera-

tions of RB1. Accordingly, the MPNST to be treated may be MPNST associated with RB1 inactivation, including, e.g., neurofibromatosis type I-linked MPNST associated with RB1 inactivation, sporadic MPNST associated with RB1 inactivation, or radiation-induced MPNST associated with RB1 inactivation.

[0027] The MPNST to be treated may also be an MPNST (including, e.g., any one of the above-mentioned specific types of MPNST) having one or more mutations or genetic alterations affecting (particularly reducing or suppressing) the expression and/or the activity of PTEN, particularly one or more inactivating mutations or inactivating genetic alterations of PTEN. Accordingly, the MPNST to be treated may be MPNST associated with PTEN inactivation, including, e.g., neurofibromatosis type I-linked MPNST associated with PTEN inactivation, sporadic MPNST associated with PTEN inactivation, or radiation-induced MPNST associated with PTEN inactivation.

[0028] The MPNST to be treated may also be an MPNST (including, e.g., any one of the above-mentioned specific types of MPNST) having one or more mutations or genetic alterations affecting (particularly reducing or suppressing) the expression and/or the activity of PRC2, particularly one or more inactivating mutations or inactivating genetic alterations of PRC2. Accordingly, the MPNST to be treated may be MPNST associated with PRC2 inactivation, including, e.g., neurofibromatosis type I-linked MPNST associated with PRC2 inactivation, sporadic MPNST associated with PRC2 inactivation, or radiation-induced MPNST associated with PRC2 inactivation. Such inactivation of PRC2 may be due, e.g., to one or more mutations in one or more PRC2 core component genes, e.g., EZH2, EED and/or SUZ12. The MPNST to be treated may also be an MPNST (including, e.g., any one of the above-mentioned specific types of MPNST) having one or more mutations or genetic alterations affecting (particularly activating or enhancing) the expression and/or the activity of EGFR, particularly one or more activating mutations or activating genetic alterations of EGFR. Accordingly, the MPNST to be treated may be MPNST associated with EGFR activation or upregulation, including, e.g., neurofibromatosis type I-linked MPNST associated with EGFR activation/upregulation, sporadic MPNST associated with EGFR activation/upregulation, or radiation-induced MPNST associated with EGFR activation/upregulation.

[0029] The MPNST to be treated may also be an MPNST (including, e.g., any one of the above-mentioned specific types of MPNST) having one or more mutations or genetic alterations affecting (particularly activating or enhancing) the expression and/or the activity of PDGFRA, particularly one or more activating mutations or activating genetic alterations of PDGFRA. Accordingly, the MPNST to be treated may be MPNST associated with PDGFRA activation or upregulation, including, e.g., neurofibromatosis type I-linked MPNST associated with PDGFRA activation/upregulation, sporadic MPNST associated with PDGFRA activation/upregulation, or radiation-induced MPNST associated with PDGFRA activation/upregulation.

[0030] The MPNST to be treated may also be an MPNST (including, e.g., any one of the above-mentioned specific types of MPNST) having one or more mutations or genetic alterations affecting (particularly activating or enhancing) the expression and/or the activity of c-Met, particularly one or more activating mutations or activating genetic alterations

of c-Met. Accordingly, the MPNST to be treated may be MPNST associated with c-Met activation or upregulation, including, e.g., neurofibromatosis type I-linked MPNST associated with c-Met activation/upregulation, sporadic MPNST associated with c-Met activation/upregulation, or radiation-induced MPNST associated with c-Met activation/upregulation.

[0031] The MPNST to be treated may further be an MPNST (including, e.g., any one of the above-mentioned specific types of MPNST, such as neurofibromatosis type I-linked MPNST, sporadic MPNST, or radiation-induced MPNST) associated with chromosome 8 gain (see, e.g., Dehner C et al., JCI Insight, 2021, 6(6):e146351, doi:10.1172/jci.insight.146351).

[0032] The MPNST to be treated may also be an MPNST (including, e.g., any one of the above-mentioned specific types of MPNST) that overexpresses one or more SNAG-domain transcription factors, such as, e.g., Snail1 (Snail), Snail2 (Slug), Snail3 (Smuc), Scratch 1, Scratch 2, Gfi-1, Gfi-1B, Insm1, Insm2, Ovol-1 (Ovo-like 1), Ovol-2, and/or Ovol-3. Moreover, the MPNST to be treated may be an MPNST (including, e.g., any one of the above-mentioned specific types of MPNST) associated with aneuploidy (e.g., wherein one or more chromosomes are present in more than two copies and/or in an uneven number of copies).

[0033] The MPNST to be treated (including also any one of the above-mentioned specific types of MPNST) may further be a metastatic MPNST. Accordingly, the MPNST to be treated may be a primary MPNST that has formed metastases, i.e., that has spread to one or more other parts of the body of the subject (such as, e.g., the lungs and/or the bones). In particular, the MPNST may be, e.g., a metastatic neurofibromatosis type I-linked MPNST, a metastatic sporadic MPNST, or a metastatic radiation-induced MPNST.

[0034] The MPNST to be treated (including also any one of the above-mentioned specific types of MPNST) may also be a relapsed or refractory MPNST. In particular, the MPNST may be, e.g., a relapsed or refractory neurofibromatosis type I-linked MPNST, a relapsed or refractory sporadic MPNST, or a relapsed or refractory radiation-induced MPNST. The therapeutic effects of LSD1 inhibitors in the treatment of MPNST can be further confirmed in additional *in vitro* or *in vivo* experiments, as well as in clinical trials in humans, which can be readily set up by those skilled in the art of drug development.

LSD1 Inhibitors

[0035] As used herein, an “LSD1 inhibitor” means a compound/substance that reduces, decreases, blocks or inhibits the gene expression, activity or function of LSD1. Compounds which act as inhibitors of LSD1 are known in the art. Any molecule acting as an LSD1 inhibitor can, in principle, be used in the context of the present invention. Preferably, the LSD1 inhibitor is a small molecule. Moreover, the LSD1 inhibitor may be an irreversible LSD1 inhibitor or a reversible LSD1 inhibitor. As also demonstrated in Example 1 below, both irreversible and reversible LSD1 inhibitors can be used for the treatment of MPNST in accordance with the present invention. Prototypical irreversible LSD1 inhibitors include cyclopropylamine-based compounds like iadademstat and bomedemstat, which are among the LSD1 inhibitors used in Example 1. A representative example of a reversible LSD1 inhibitor is the compound pulrodemstat, which has also been used in Example 1.

Preferably, the LSD1 inhibitor is a selective LSD1 inhibitor; as used herein, a “selective LSD1 inhibitor” means an LSD1 inhibitor which exhibits a selectivity of at least 10-fold (preferably at least 100-fold) for LSD1 over other FAD-dependent monoamine oxidases, particularly over MAO-A and MAO-B (which can be assessed, e.g., by determining IC₅₀ values for LSD1, MAO-A and MAO-B).

[0036] An exemplary list of small-molecule LSD1 inhibitors is provided in the table below:

WO2017/027678, WO2017/079476, WO2017/079670,
WO2017/090756, EP3381896A1, WO2017/109061,
WO2017/116558, WO2017/149463, WO2017/157322,
EP3431471A1, WO2017/184934, WO2017/195216,
WO2017/198780, WO2017/215464, EP3486244A1,
WO2018/081342, WO2018/081343, WO2018/137644,
EP3575285A1, WO2018/213211, WO2018/216800,
EP3632897A1, WO2018/226053, WO2018/234978,
WO2019/009412, WO2019/034774, WO2019/054766,

Chemical name	Generic name/Code	International Non-proprietary Name (INN)
(trans)-N1-((1R,2S)-2-phenylcyclopropyl)cyclohexane-1,4-diamine	ORY-1001	iadademstat
4-[2-(4-aminopiperidin-1-yl)-5-(3-fluoro-4-methoxyphenyl)-1-methyl-6-oxo-1,6-dihydropyrimidin-4-yl]-2-fluorobenzonitrile	CC-90011	pulrodemstat
N-[(2S)-5-{{[(1R,2S)-2-(4-fluorophenyl)cyclopropyl]amino}-1-(4-methylpiperazin-1-yl)-1-oxopentan-2-yl]-4-(1H-1,2,3-triazol-1-yl)benzamide	IMG-7289	bomedemstat
(E)-N'-1-(5-chloro-2-hydroxyphenyl)ethylidene)-3-((4-methylpiperazin-1-yl)sulfonyl)benzohydrazide	SP-2577	seclidemstat
1-((4-(methoxymethyl)-4-(((1R,2S)-2-phenylcyclopropylamino)methyl)piperidin-1-yl)methyl)cyclobutanecarboxylic acid		
3-(cyanomethyl)-3-(4-{{[(1R,2S)-2-phenylcyclopropyl]amino}piperidin-1-yl)azetidine-1-sulfonamide		
5-(((1R,2S)-2-(4-(benzyloxy)phenyl)cyclopropylamino)methyl)-1,3,4-oxadiazol-2-amine	ORY-2001	vafidemstat
N-(trans)-2-phenylcyclopropyl)piperidin-4-amine	OG-668, also known as GSK-LSD1	
N-[4-[(trans)-2-[(cyclopropylmethyl)amino]cyclopropyl]phenyl]-1-methyl-1H-pyrazole-4-carboxamide	T-3775440	
1-morpholino-3-(4-(((1R,2S)-2-phenylcyclopropyl)amino)piperidin-1-yl)propan-1-one	MRTX1519	
4-[4-[(1R,2S)-2-phenylcyclopropyl]amino]methyl]-1-piperidinyl[methyl]-benzoic acid	GSK2879552	
4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]phenyl]-2-fluoro-benzonitrile		
2-(6-(((1R,2S)-2-(E)-1-phenylbut-1-en-2-yl)cyclopropyl)amino)-2-azaspiro[3.3]heptan-2-yl)ethanol		
2-(6-(((1R,2S)-2-((E)-1-phenylbut-1-en-2-yl)cyclopropyl)amino)-2-azaspiro[3.3]heptan-2-yl)propane-1,3-diol		
4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]phenyl]-2-fluoro-benzonitrile	TAS1440	

[0037] The LSD1 inhibitor to be used in accordance with the present invention may thus be, e.g., any one of the specific compounds listed in the table above, or a pharmaceutically acceptable salt of any one of these compounds.


[0038] In some embodiments, the LSD1 inhibitor is an LSD1 inhibitor known in the art, including, e.g., any one of the compounds disclosed in WO2010/043721, WO2010/084160, WO2010/143582, WO2011/035941, WO2011/042217, WO2011/131576, WO2011/131697, WO2012/013727, WO2012/013728, WO2012/045883, WO2012/135113, WO2013/022047, EP2743256A1, WO2013/025805, WO2013/057320, WO2013/057322, WO2014/058071, EP2907802A1, WO2014/084298, EP2927212A1, WO2014/086790, WO2014/164867, WO2014/194280, WO2014/205213, WO2015/021128, WO2015/031564, WO2015/089192, WO2015/120281, WO2015/123408, WO2015/123424, WO2015/123437, WO2015/123465, WO2015/134973, WO2015/168466, WO2015/181380, WO2015/200843, WO2016/003917, WO2016/004105, WO2016/007722, WO2016/007727, WO2016/007731, WO2016/007736, WO2016/034946, WO2016/037005, WO2016/123387, WO2016/130952, WO2016/161282, WO2016/172496, WO2016/177656, WO2017/004519,

WO2019/217972, WO2019/222069, WO2020/015745,
EP3825309A1, WO2020/047198, WO2020/052647,
WO2020/052649, EP3851440A1, WO2020/138398,
WO2020/159285, EP3907225A1, WO2021/058024,
WO2021/095835, WO2021/175079, WO2022/072811,
WO2022/171044, WO2022/188709, WO2022/240886,
WO2022/267495, WO2023/069884, WO2023/284651,
US2017-0283397, US2022-0064126, CN103054869,
CN103319466, CN104119280, CN105541806,
CN105924362, CN105985265, CN106045862,
CN106045881, CN106432248, CN106478639,
CN106831489, CN106928235, CN107033148,
CN107174584, CN107176927, CN107459476,
CN107474011, CN107501169, CN107936022,
CN108530302, CN109265462, CN109293664,
CN109535019, CN110204551, CN110478352,
CN111072610, CN111454252, CN112110936,
CN112409310, CN112920130, CN113087712,
CN113105479, CN113264903, CN113582906,
CN113599380, CN114502561, CN114805205,
CN114805261, KR20190040763, or KR20190040783, each of which is incorporated herein by reference in its entirety (including, in particular, the compounds described in the

examples section of each one of these documents). Accordingly, the LSD1 inhibitor may be, e.g., a compound disclosed in any one of the aforementioned documents (including, e.g., in the examples section of any one of these documents), wherein said compound may be used in non-salt form or in the form of a pharmaceutically acceptable salt.


[0039] In some embodiments, the LSD1 inhibitor is selected from the group consisting of iadademstat, pulrodemstat, bomedemstat, seclidemstat, 1-((4-(methoxymethyl)-4-(((1R,2S)-2-phenylcyclopropylamino)methyl)piperidin-1-yl)methyl)cyclobutanecarboxylic acid, 3-(cyanomethyl)-3-(4-(((1R,2S)-2-phenylcyclopropylamino)piperidin-1-yl)azetidine-1-sulfonamide, vafidemstat, 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]phenyl]-2-fluorobenzonitrile, and pharmaceutically acceptable salts thereof (i.e., pharmaceutically acceptable salts of any one of the aforementioned compounds).

[0040] Iadademstat is a selective and irreversible LSD1 inhibitor. Iadademstat is the INN for the compound of formula:




[0041] [CAS Reg. No. 1431304-21-0], which is also known as ORY-1001 or (trans)-N1-((1R,2S)-2-phenylcyclopropyl)cyclohexane-1,4-diamine. Iadademstat has been described, e.g., in Example 5 of WO2013/057322. Pharmaceutically acceptable salts of iadademstat, including hydrochloride salts (particularly iadademstat dihydrochloride), are also described in WO2013/057322.

[0042] Pulrodemstat is a reversible LSD1 inhibitor of formula



[0043] [CAS Reg. No. 1821307-10-1], also known as CC-90011, with the chemical name 4-[2-(4-aminopiperidin-1-yl)-5-(3-fluoro-4-methoxyphenyl)-1-methyl-6-oxo-1,6-dihydropyrimidin-4-yl]-2-fluorobenzonitrile. Pulrodemstat has been described, e.g., in WO2015/168466 and WO2017/79670. Pharmaceutically acceptable salts thereof are also described therein, including a besylate salt.


[0044] Bomedemstat is an irreversible LSD1 inhibitor of formula



[0045] [CAS Reg. No. 1990504-34-1], also known as IMG-7289, with the chemical name N-[(2S)-5-(((1R,2S)-2-(4-fluorophenyl)cyclopropyl)amino)-1-(4-methylpiperazin-1-yl)-1-oxopentan-2-yl]-4-(1H-1,2,3-triazol-1-yl)benzamide.


[0046] Bomedemstat has been described, e.g., in WO2016/130952 and WO2018/35259. Pharmaceutically acceptable salts thereof are also described therein, including a bis-tosylate salt.

[0047] Seclidemstat is an LSD1 inhibitor of formula




[0048] [CAS Reg. No. 1423715-37-0], also known as SP-2577, with the chemical name (E)-N'-(1-(5-chloro-2-hydroxyphenyl)ethylidene)-3-((4-methylpiperazin-1-yl)sulfonyl)benzohydrazide. Seclidemstat has been described, e.g., in WO2013/025805 and WO2014/205213.


[0049] 1-((4-(Methoxymethyl)-4-(((1R,2S)-2-phenylcyclopropylamino)methyl)piperidin-1-yl)methyl)cyclobutanecarboxylic acid is an irreversible LSD1 inhibitor which is described, e.g., in WO2015/123465 and WO2017/27678. Pharmaceutically acceptable salts thereof are also described therein, including a para-toluenesulfonate salt. The structure of this compound can be depicted as follows:



[0050] 3-(Cyanomethyl)-3-(4-(((1R,2S)-2-phenylcyclopropylamino)piperidin-1-yl)azetidine-1-sulfonamide is an irreversible LSD1 inhibitor which is described, e.g., in WO2020/047198. Pharmaceutically acceptable salts thereof are also described therein. The structure of this compound can be depicted as follows:




[0051] Vafidemstat is an irreversible LSD1 inhibitor of formula:



which is also known as ORY-2001, 5-((((1R,2S)-2-(4-(benzyloxy)phenyl)cyclopropyl)amino)methyl)-1,3,4-oxadiazol-2-amine, or (–) 5-((((trans)-2-(4-(benzyloxy)phenyl)cyclopropyl)amino)methyl)-1,3,4-oxadiazol-2-amine. Vafidemstat has been described, e.g., in Example 35 of WO2012/13728.

[0052] 4-[5-[(3S)-3-Aminopyrrolidine-1-carbonyl]-2-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]phenyl]-2-fluoro-benzonitrile is an LSD1 inhibitor which is described, e.g., in WO2017/090756 (or EP3381896A1; see Example 37), WO2021/095835, WO2022/240886, and WO2023/054547. Also described therein are pharmaceutically acceptable salts of this compound, including a benzoic acid salt (or benzoate salt), a sorbic acid salt, a succinic acid salt, an L-tartaric acid salt, a hydrochloric acid salt, a hemi-fumarate salt, a mono-fumarate salt, a hemi-oxalate salt, a mono-oxalate salt, a mesylate salt, an esylate salt, or a maleate salt. Specific solid forms of this compound are described in WO2022/240886. This compound is also referred to herein as “TAS1440”. The structure of this compound can be depicted as follows:



[0053] Further examples of the LSD1 inhibitor include SYHA1807 or a pharmaceutically acceptable salt thereof, or JBI-802 or a pharmaceutically acceptable salt thereof.

[0054] In some embodiments, the LSD1 inhibitor is selected from the group consisting of iadademstat, pulro-

demstat, bomedemstat, seclidemstat, 1-(((4-(methoxymethyl)-4-(((1R,2S)-2-phenylcyclopropylamino)methyl)piperidin-1-yl)methyl)cyclobutanecarboxylic acid, 3-(cyanomethyl)-3-(4-(((1R,2S)-2-phenylcyclopropyl)amino)piperidin-1-yl)azetidine-1-sulfonamide, 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]phenyl]-2-fluoro-benzonitrile, and pharmaceutically acceptable salts thereof.

[0055] In particular, the LSD1 inhibitor may be selected from the group consisting of iadademstat, pulrodemstat, bomedemstat, seclidemstat, 1-(((4-(methoxymethyl)-4-(((1R,2S)-2-phenylcyclopropylamino)methyl)piperidin-1-yl)methyl)cyclobutanecarboxylic acid, 3-(cyanomethyl)-3-(4-(((1R,2S)-2-phenylcyclopropyl)amino)piperidin-1-yl)azetidine-1-sulfonamide, and pharmaceutically acceptable salts thereof.

[0056] In preferred embodiments, the LSD1 inhibitor is selected from the group consisting of iadademstat, pulrodemstat, bomedemstat, and pharmaceutically acceptable salts thereof. In some embodiments, the LSD1 inhibitor is pulrodemstat or a pharmaceutically acceptable salt thereof (e.g., pulrodemstat besylate). In some embodiments, the LSD1 inhibitor is bomedemstat, or a pharmaceutically acceptable salt thereof (e.g., bomedemstat bis-tosylate).

[0057] A particularly preferred LSD1 inhibitor is iadademstat or a pharmaceutically acceptable salt thereof. In some embodiments, iadademstat is used as the dihydrochloride salt (i.e., iadademstat dihydrochloride).

[0058] Unless specifically indicated otherwise, any reference to an LSD1 inhibitor (e.g., iadademstat) throughout the present description and claims includes such LSD1 inhibitor in non-salt form and any of its pharmaceutically acceptable salts. When the LSD1 inhibitor is iadademstat, it is preferably used in the form of a pharmaceutically acceptable salt, preferably a hydrochloride salt, more preferably the dihydrochloride salt.

Pharmaceutical Formulations

[0059] The LSD1 inhibitor to be used in accordance with the present invention, as well as any pharmaceutical composition comprising an LSD1 inhibitor to be used in accordance with the invention, may be administered by any route appropriate to the condition to be treated. Exemplary routes include oral, parenteral (including subcutaneous, intramuscular, intravenous, intraarterial, inhalation, intradermal, intrathecal, epidural, and infusion techniques), transdermal, rectal, nasal, topical (including buccal and sublingual), vaginal, intraperitoneal, intrapulmonary and intranasal. Preferably, the LSD1 inhibitor (or the corresponding pharmaceutical composition) is administered orally. The LSD1 inhibitor to be used in accordance with the present invention may be administered in any convenient pharmaceutical composition or formulation, e.g., as tablets, powders, capsules, solutions, dispersions, suspensions, syrups, sprays, suppositories, gels, emulsions, patches, etc. Such compositions/formulations may comprise components conventional in pharmaceutical preparations, e.g., diluents, carriers, pH modifiers, preservatives, solubilizers, stabilizers, wetting agents, emulsifiers, sweeteners, colorants, flavorants, salts for varying the osmotic pressure, buffers, masking agents, antioxidants, and/or further active agents. They can also comprise still other therapeutically active or therapeutically valuable substances.

[0060] A typical formulation is prepared by mixing an LSD1 inhibitor and one or more pharmaceutically acceptable excipients. Suitable excipients are well known to those skilled in the art and are described in detail in, e.g., “Ansel’s Pharmaceutical Dosage Forms and Drug Delivery Systems” (2004) Lippincott, Williams & Wilkins, Philadelphia; “Remington: The Science and Practice of Pharmacy” (2000) Lippincott, Williams & Wilkins, Philadelphia; or “Handbook of Pharmaceutical Excipients” (2005) Pharmaceutical Press, Chicago. The formulations may also include one or more buffers, stabilizing agents, surfactants, wetting agents, lubricating agents, emulsifiers, suspending agents, preservatives, antioxidants, opaquing agents, glidants, processing aids, colorants, sweeteners, perfuming agents, flavoring agents, diluents and/or other known additives to provide an elegant presentation of the active agent or aid in the manufacturing of the pharmaceutical product (i.e., medicament).

[0061] For oral delivery, the LSD1 inhibitor can be incorporated into a formulation that includes pharmaceutically acceptable carriers such as binders (e.g., gelatin, cellulose, or gum tragacanth), excipients (e.g., starch or lactose), lubricants (e.g., magnesium stearate or silicon dioxide), disintegrating agents (e.g., alginate, Primogel, or corn starch), and sweetening or flavoring agents (e.g., glucose, sucrose, saccharin, methyl salicylate, or peppermint). The formulation can be orally delivered, e.g., in the form of enclosed gelatin capsules or compressed tablets. Capsules and tablets can be prepared by any conventional techniques. The capsules and tablets can also be coated with various coatings known in the art to modify the flavors, tastes, colors, and shapes of the capsules and tablets. In addition, liquid carriers such as fatty oil can also be included in capsules.

[0062] Suitable oral formulations can also be in the form of a suspension, syrup, chewing gum, wafer, elixir, and the like. If desired, conventional agents for modifying flavors, tastes, colors, and shapes of the special forms can also be included. In addition, for convenient administration by enteral feeding tube in subjects unable to swallow, the active compounds can be dissolved in an acceptable lipophilic vegetable oil vehicle, such as olive oil, corn oil or safflower oil.

[0063] The LSD1 inhibitor can also be administered parenterally in the form of a solution or suspension, or in lyophilized form capable of conversion into a solution or suspension form before use. In such formulations, diluents or pharmaceutically acceptable carriers such as sterile water and physiological saline buffer can be used. Other conventional solvents, pH buffers, stabilizers, anti-bacterial agents, surfactants, and antioxidants can all be included. For example, useful components include sodium chloride, acetates, citrate or phosphate buffers, glycerin, dextrose, fixed oils, methyl parabens, polyethylene glycol, propylene glycol, sodium bisulfate, benzyl alcohol, ascorbic acid, and the like. The parenteral formulations can be stored in any conventional containers such as vials and ampoules.

[0064] Subcutaneous implantation for sustained release of the LSD1 inhibitor may also be a suitable route of administration. This entails surgical procedures for implanting an LSD1 inhibitor in any suitable formulation into a subcutaneous space, e.g., beneath the anterior abdominal wall. See, e.g., Wilson et al. (1984) *J. Clin. Psych.* 45:242-247. Hydrogels can be used as a carrier for the sustained release of LSD1 inhibitors. Hydrogels are generally known in the art.

They are typically made by crosslinking high molecular weight biocompatible polymers into a network, which swells in water to form a gel-like material. Preferably, hydrogels are biodegradable or biosorbable. For the purpose of this invention, hydrogels made of polyethylene glycols, collagen, or poly(glycolic-co-L-lactic acid) may be useful; see, e.g., Phillips et al. (1984) *J. Pharmaceut. Sci.*, 73:1718-1720.

[0065] The pharmaceutical compositions, like oral and parenteral compositions, can be formulated in unit dosage forms for ease of administration and uniformity of dosage. As used herein, “unit dosage forms” refers to physically discrete units suitable as unitary dosages for administration to subjects, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect, in association with one or more suitable pharmaceutical carriers.

[0066] Suitable oral dosage forms for iadademstat are disclosed, for example, in WO2019/211491.

[0067] In particular, iadademstat may be provided in the form of a solid oral dosage form, such as, e.g., tablets or capsules. Alternatively, iadademstat may also be provided in the form of an oral liquid composition, particularly an oral solution, such as, e.g., an oral aqueous solution (a corresponding oral solution, including an oral aqueous solution, may be prepared, e.g., from a powder for reconstitution). As explained above, it is preferred that iadademstat is used in the form of iadademstat dihydrochloride.

[0068] For the treatment of MPNST, the LSD1 inhibitor (or the corresponding pharmaceutical composition) can be administered in any appropriate manner, as determined by a person skilled in the medical arts. An appropriate dose and suitable duration and frequency of administration can vary within wide limits and will be determined by such factors as the condition of the subject, the specific type and severity of the disease, the particular form of the active ingredient(s), and the method of administration, among others. In general, an appropriate dose and administration regimen provides the LSD1 inhibitor in an amount sufficient to provide therapeutic benefit, for example an improved clinical outcome, such as more frequent complete or partial remissions, or longer disease-free and/or overall survival, or lessening of symptoms severity, or any other objectively identifiable improvement as noted by the clinician. Therapeutically effective doses may generally be assessed or extrapolated using experimental models like dose-response curves derived from in vitro or animal model test systems, or from clinical trials in humans.

[0069] Suitable doses and dosing regimens for the LSD1 inhibitor will be dependent on the specific LSD1 inhibitor used, its LSD1 inhibitory potency, its pharmacokinetic profile and other factors, as well known by those skilled in the art. Iadademstat is a highly potent active pharmaceutical ingredient (HPAPI). The anticipated daily dose is thus very low, e.g., lower than 1 mg per day. Accordingly, the drug load in a pharmaceutical formulation (including, e.g., a solid oral form) will typically also be very low (e.g., less than 1 mg of API per 100 mg of solid oral form). In general, in the case of oral administration (e.g., as tablets, capsules, or as an oral solution, including e.g. an oral aqueous solution) to an adult human subject (i.e., a human subject having an age of 18 years or more), a daily dosage of about 50 µg to about 300 µg, preferably of about 75 µg to about 300 µg (e.g., about 75 µg, about 100 µg, about 125 µg, about 150 µg, about 175 µg,

about 200 µg, about 225 µg, about 250 µg, about 275 µg, or about 300 µg, or any range between any two of the aforementioned daily dosages), of iadademstat as described herein should be appropriate, although these limits may be adjusted when necessary. For example, the aforementioned dosages may be lowered for paediatric use, particularly for the oral administration to a human subject having less than 18 years of age (e.g., having 0 to 2 years, 2 to 12 years, or 12 to less than 18 years). The term “µg” (or “ug”), as used herein, refers to micrograms.

[0070] In some embodiments, the LSD1 inhibitor is iadademstat (or a pharmaceutically acceptable salt thereof, e.g., iadademstat dihydrochloride) and is administered five days on/two days off (5/2) per week.

[0071] In some embodiments, the LSD1 inhibitor is iadademstat (or a pharmaceutically acceptable salt thereof, e.g., iadademstat dihydrochloride) and is administered orally to an adult human subject at a daily dose of about 50 µg to about 300 µg, preferably of about 75 µg to about 300 µg (e.g., about 100 µg to about 300 µg), five days on/two days off (5/2) per week. Doses as reflected herein for iadademstat relate to the corresponding amount of the iadademstat free base. In some embodiments, iadademstat is administered orally at a daily dose of about 75 µg five days on/two days off (5/2) per week. In some embodiments, iadademstat is administered orally at a daily dose of about 100 µg five days on/two days off (5/2) per week. In some embodiments, iadademstat is administered orally at a daily dose of about 150 µg five days on/two days off (5/2) per week. In some embodiments, iadademstat is administered orally at a daily dose of about 200 µg five days on/two days off (5/2) per week. In some embodiments, iadademstat is administered orally at a daily dose of about 250 µg five days on/two days off (5/2) per week. In some embodiments, iadademstat is administered orally at a daily dose of about 300 µg five days on/two days off (5/2) per week. As explained above, these dosages may be lowered for paediatric use.

Combination Treatments

[0072] The LSD1 inhibitor to be used in accordance with the present invention can be administered in monotherapy (e.g., without concomitantly administering any further therapeutic agents, or without concomitantly administering any further anticancer agents). Accordingly, the invention relates to an LSD1 inhibitor (or a pharmaceutical composition comprising an LSD1 inhibitor and optionally one or more pharmaceutically acceptable excipients) for use in the monotherapeutic treatment of MPNST. The invention likewise relates to corresponding methods and uses for the monotherapeutic treatment of MPNST.

[0073] However, the LSD1 inhibitor can also be administered in combination with one or more further therapeutic agents, particularly one or more further anticancer agents. If the LSD1 inhibitor is used in combination with a further anticancer agent, the dose of each compound may differ from that when the corresponding compound is used alone, in particular, a lower dose of either one or both compounds may be used.

[0074] The combination of the LSD1 inhibitor with one or more further therapeutic agents (e.g., one or more further anticancer agents) may comprise the simultaneous/concomitant administration of the LSD1 inhibitor and the further therapeutic agent(s), either in a single pharmaceutical formulation or in separate pharmaceutical formulations, or the

sequential/separate administration of the LSD1 inhibitor and the further therapeutic agent(s). If administration is sequential, either the LSD1 inhibitor or the one or more further therapeutic agents may be administered first. If administration is simultaneous, the one or more further therapeutic agents may be included in the same pharmaceutical formulation as the LSD1 inhibitor, or they may be administered in two or more distinct/separate pharmaceutical formulations. It will be appreciated that administering the LSD1 inhibitor and the further therapeutic agent(s) in separate pharmaceutical formulations is expedient, e.g., if the respective agents are administered by different routes and/or using different administration schedules/regimens.

[0075] Moreover, the LSD1 inhibitor can also be administered in combination with physical therapy, particularly radiotherapy. The invention likewise relates to the combined use of an LSD1 inhibitor with one or more further therapeutic agents (particularly one or more further anticancer agents) and with physical therapy (particularly radiotherapy). Physical therapy (or radiotherapy) may commence before, after, or simultaneously with the administration of the LSD1 inhibitor (e.g., about 1 to 72 hours before or after the administration of the LSD1 inhibitor).

[0076] Accordingly, the invention relates to an LSD1 inhibitor for use in the treatment of MPNST in combination with one or more further therapeutic agents (particularly one or more further anticancer agents) and/or in combination with radiotherapy. The invention also relates to a pharmaceutical composition comprising an LSD1 inhibitor and optionally one or more pharmaceutically acceptable excipients for use in the treatment of MPNST in combination with one or more further therapeutic agents (particularly one or more further anticancer agents) and/or in combination with radiotherapy. The invention further relates to an LSD1 inhibitor (or a pharmaceutical composition comprising an LSD1 inhibitor and optionally one or more pharmaceutically acceptable excipients) for use in the treatment of MPNST, wherein the LSD1 inhibitor (or the pharmaceutical composition comprising the LSD1 inhibitor) is administered in combination with one or more further therapeutic agents (particularly one or more further anticancer agents) and/or in combination with radiotherapy. The invention further relates to an LSD1 inhibitor (or a pharmaceutical composition comprising an LSD1 inhibitor and optionally one or more pharmaceutically acceptable excipients) for use in the treatment of MPNST, wherein the LSD1 inhibitor (or the pharmaceutical composition comprising the LSD1 inhibitor) is for use in combination with one or more further therapeutic agents (particularly one or more further anticancer agents) and/or in combination with radiotherapy. The invention further relates to: (i) an anticancer agent for use in the treatment of MPNST in combination with an LSD1 inhibitor; (ii) an anticancer agent for use in the treatment of MPNST, wherein the anticancer agent is administered in combination with an LSD1 inhibitor; or (iii) an anticancer agent for use in the treatment of MPNST, wherein the anticancer agent is for use in combination with an LSD1 inhibitor.

[0077] The invention likewise provides a method of treating MPNST in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of an LSD1 inhibitor (or a therapeutically effective amount of a pharmaceutical composition comprising an LSD1 inhibitor and optionally one or more pharmaceutically acceptable

excipients) in combination with a therapeutically effective amount of one or more further therapeutic agents (particularly one or more further anticancer agents) and/or in combination with radiotherapy.

[0078] Moreover, the invention relates to the use of an LSD1 inhibitor for the treatment of MPNST in combination with one or more further therapeutic agents (particularly one or more further anticancer agents) and/or in combination with radiotherapy. The invention also relates to the use of an anticancer agent for the treatment of MPNST in combination with an LSD1 inhibitor.

[0079] The invention furthermore relates to the use of an LSD1 inhibitor for the preparation of a medicament (or a pharmaceutical composition) for the treatment of MPNST in combination with one or more further therapeutic agents (particularly one or more further anticancer agents) and/or in combination with radiotherapy. The invention also relates to the use of an anticancer agent for the preparation of a medicament for the treatment of MPNST in combination with an LSD1 inhibitor. The invention likewise relates to the use of an LSD1 inhibitor and one or more further anticancer agents for the preparation of a medicament for the treatment of MPNST, wherein the medicament comprises the LSD1 inhibitor and the further anticancer agent(s) in the same pharmaceutical formulation or in separate pharmaceutical formulations. The invention also relates to the use of an LSD1 inhibitor for the preparation of a medicament for the treatment of MPNST, wherein said medicament is prepared for combined use (or for use in combination) with one or more further therapeutic agents (particularly one or more further anticancer agents) and/or with radiotherapy. The invention also relates to the use of an anticancer agent for the preparation of a medicament for the treatment of MPNST, wherein said medicament is prepared for combined use (or for use in combination) with an LSD1 inhibitor.

[0080] The present invention furthermore provides a combination product comprising, in the same pharmaceutical formulation or in separate pharmaceutical formulations, an LSD1 inhibitor and one or more further therapeutic agents (particularly one or more further anticancer agents), for use in the treatment of MPNST. The LSD1 inhibitor and the further therapeutic agent(s) (particularly the further anticancer agent(s)) may thus be present in a single pharmaceutical formulation (i.e., in the same pharmaceutical formulation), or they may each be provided in a distinct (separate) pharmaceutical formulation.

[0081] The present invention also provides a pharmaceutical composition comprising an LSD1 inhibitor in combination with one or more further therapeutic agents (particularly one or more further anticancer agents), and one or more pharmaceutically acceptable excipients, for use in the treatment of MPNST.

[0082] The present invention further provides an article of manufacture (or a kit) comprising, in the same pharmaceutical formulation or in separate pharmaceutical formulations, an LSD1 inhibitor and one or more further therapeutic agents (particularly one or more further anticancer agents), for use in the treatment of MPNST.

[0083] The invention further provides a method of treating MPNST in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of the above-described combination product, the pharmaceutical composition or the article of manufacture. In particular, the invention provides a method of treating MPNST in a subject

in need thereof, comprising administering to the subject a therapeutically effective amount of a combination product comprising, in the same pharmaceutical formulation or in separate pharmaceutical formulations, an LSD1 inhibitor and one or more further therapeutic agents (particularly one or more further anticancer agents). The invention further provides a method of treating MPNST in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of an LSD1 inhibitor and a therapeutically effective amount of one or more further therapeutic agents (particularly one or more further anticancer agents).

[0084] The invention further provides the use of a combination comprising an LSD1 inhibitor and one or more further therapeutic agents (particularly one or more further anticancer agents) for the manufacture of a medicament (or a pharmaceutical composition) for the treatment of MPNST. The invention also provides the use of a combination comprising an LSD1 inhibitor and one or more further therapeutic agents (particularly one or more further anticancer agents) for the treatment of MPNST.

[0085] The anticancer agent(s) (particularly the “anticancer agent” or the “one or more further anticancer agents” as referred to in any of the paragraphs herein above) may be selected, for example, from a MEK inhibitor (particularly an inhibitor of MEK1 and/or MEK2; e.g., selumetinib), a PI3K inhibitor (e.g., copanlisib), an mTOR inhibitor (e.g., temsirolimus), an ERK inhibitor (particularly an inhibitor of ERK1 and/or ERK2; e.g., ulixertinib), a KRAS inhibitor (e.g., sotorasib), an EGFR inhibitor (e.g., lapatinib), a cKIT inhibitor (e.g., imatinib), a proteasome inhibitor (e.g., bortezomib), a DNA intercalator (e.g., doxorubicin), a RAF inhibitor (particularly a BRAF inhibitor; e.g., sorafenib), a VEGFR inhibitor (e.g., cabozantinib), an ALK inhibitor (e.g., crizotinib), a glutaminase inhibitor (e.g., telaglenastat), a JAK inhibitor (or a Janus kinase inhibitor; e.g., tofacitinib), a PLK1 inhibitor (e.g., volasertib), a Bcl2 inhibitor (e.g., venetoclax), an HDAC inhibitor (e.g., vorinostat), an HSP90 inhibitor (e.g., onalespib), a Wnt/ β -catenin pathway inhibitor (e.g., OMP-18R5), an Aurora kinase inhibitor (e.g., alisertib), an MDM2 inhibitor (e.g., alrizomadlin), a CDK4/6 inhibitor (e.g., abemaciclib), a YAP/TAZ pathway inhibitor (e.g., pazopanib), an SOS inhibitor (e.g., BI 1701963), a Grb2 inhibitor (e.g., BP1001), a BET inhibitor (including, in particular, a BRD4 inhibitor; e.g., GSK1210151A), an AKT inhibitor (e.g., ipatasertib), an MNK inhibitor (e.g., ETC-206), an NTRK inhibitor (e.g., entrectinib), an SPH2 inhibitor (e.g., JAB-3068), and a PP2A inhibitor (e.g., LB100).

[0086] The MEK inhibitor may be, e.g., selumetinib, trametinib, cobimetinib, binimetinib, mirdametinib, pimasertib, refametinib, zapnometinib, avutometinib, HL-085, FCN-159, TAK-733, or a pharmaceutically acceptable salt of any one of these agents. The PI3K inhibitor may be, e.g., copanlisib, alpelisib, idelalisib, duvelisib, umbralisib, buparlisib, zandelisib, linperlisib, pascalisib, leniolisib, paxalisib, inavolisib, serabelisib, pictilisib, taselisib, tenalisib, eganelisib, GSK2636771, MEN1611, AMG-319, or a pharmaceutically acceptable salt of any one of these agents. The mTOR inhibitor may be, e.g., temsirolimus, everolimus, sirolimus, or a pharmaceutically acceptable salt of any one of these agents. The ERK inhibitor may be, e.g., ulixertinib or a pharmaceutically acceptable salt thereof. The KRAS inhibitor may be, e.g., sotorasib, adagrasib, or a pharmaceutically

acceptable salt of any one of these agents. The EGFR inhibitor may be, e.g., lapatinib, gefitinib, erlotinib, osimertinib, afatinib, or a pharmaceutically acceptable salt of any one of these agents. The cKIT inhibitor may be, e.g., imatinib, sorafenib, lapatinib, sunitinib, or a pharmaceutically acceptable salt of any one of these agents. The proteasome inhibitor may be, e.g., bortezomib, carfilzomib, ixazomib, or a pharmaceutically acceptable salt of any one of these agents. The DNA intercalator may be, e.g., doxorubicin, daunorubicin, epirubicin, idarubicin, or a pharmaceutically acceptable salt of any one of these agents. The RAF inhibitor may be, e.g., sorafenib, encorafenib, dabrafenib, vemurafenib, or a pharmaceutically acceptable salt of any one of these agents. The VEGFR inhibitor may be, e.g., cabozantinib, axatinib, lenvatinib, nintedanib, pazopanib, regorafenib, sorafenib, sunitinib, vandetanib, or a pharmaceutically acceptable salt of any one of these agents. The ALK inhibitor may be, e.g., crizotinib, alectinib, ceritinib, or a pharmaceutically acceptable salt of any one of these agents. The glutaminase inhibitor may be, e.g., telaglenastat or a pharmaceutically acceptable salt thereof. The JAK inhibitor may be, e.g., tofacitinib, ruxolitinib, upadacitinib, abrocitinib, or a pharmaceutically acceptable salt of any one of these agents. The PLK1 inhibitor may be, e.g., volasertib, onvansertib, rigosertib, BI 2536, or a pharmaceutically acceptable salt of any one of these agents. The Bcl2 inhibitor may be, e.g., venetoclax, navitoclax, obatoclax, or a pharmaceutically acceptable salt of any one of these agents. The HDAC inhibitor may be, e.g., vorinostat, belinostat, panobinostat, romidepsin, practinostat, rocilinosat, quisinostat, abexinostat, resminostat, givinostat, entinostat, mocetinostat, or a pharmaceutically acceptable salt of any one of these agents. The HSP90 inhibitor may be, e.g., onalespib, luminespib, ganetespib, geldanamycin, IPI-504, tanespimycin, alvespimycin, or a pharmaceutically acceptable salt of any one of these agents. The Wnt/ β -catenin pathway inhibitor may be, e.g., OMP-18R5, OMP-54F28, OTSA 101, SAH-BCL9, XAV939, IWR1, JW74, J01-017a, PKF115-584, PKF118-310, NCB-0846, LGK974, CWP232291, PRI-724, sulindac, vismodegib, glasdegib, or a pharmaceutically acceptable salt of any one of these agents. The Aurora kinase inhibitor may be, e.g., alisertib, tozasertib, barasertib, danusertib, or a pharmaceutically acceptable salt of any one of these agents. The MDM2 inhibitor may be, e.g., alrizomadlin, idasanutlin, R05045337, R05503781, AMG232, CGM097, SAR405838, MK-8242, ALRN-6924, or a pharmaceutically acceptable salt of any one of these agents. The CDK4/6 inhibitor may be, e.g., abemaciclib, ribociclib, palbociclib, or a pharmaceutically acceptable salt of any one of these agents. The YAP/TAZ pathway inhibitor may be, e.g., K-975, TED-347, pazopanib, or a pharmaceutically acceptable salt of any one of these agents. The SOS inhibitor may be, e.g., BI 1701963, BI 3406, BAY-293, or a pharmaceutically acceptable salt of any one of these agents. The Grb2 inhibitor may be, e.g., BP1001, CGP78850, CGP85793, or a pharmaceutically acceptable salt of any one of these agents. The BET inhibitor may be, e.g., ABBV-075, ABBV-744, AZD5153, BAY1238097, CPI-203, CPI-0610, GSK1210151A (or I-BET 151), GSK1324726A (I-BET 726), GSK525762 (or I-BET 762), JQ1, LY294002, MS 436, MS 645, MT-1, olinone, OTX-015, RVX-208, TEN-010, or a pharmaceutically acceptable salt of any one of these agents. The AKT inhibitor may be, e.g., ipatasertib, uprosertib, afuresertib,

MK-2206, triciribine, lactoquinomycin, AZD5363, miransertib, capibasertib, or a pharmaceutically acceptable salt of any one of these agents. The MNK inhibitor may be, e.g., ETC-206, SEL-201, BAY1143269, tomivosertib, CGP57380, or a pharmaceutically acceptable salt of any one of these agents. The NTRK inhibitor may be, e.g., entrectinib, larotrectinib, or a pharmaceutically acceptable salt of any one of these agents. The SPH2 inhibitor may be, e.g., JAB-3068, TNO155, SHP099, RMC-4550, IACS-13909, or a pharmaceutically acceptable salt of any one of these agents. The PP2A inhibitor may be, e.g., LB100, cantharidin, cantharidic acid, cytosstatin, fostriecin, or a pharmaceutically acceptable salt of any one of these agents. Any of the aforementioned anticancer agents may, in principle, be used in non-salt form or in the form of a pharmaceutically acceptable salt. The present invention specifically and individually relates to each one of the LSD1 inhibitors described herein in combination with each one of the above-described anticancer agents.

[0087] As described above, the one or more further therapeutic agents to be used in combination with an LSD1 inhibitor in accordance with the present invention may be (or may comprise) one or more further anticancer agents. Alternatively or in addition, the one or more further therapeutic agents may also comprise an antiemetic agent. Accordingly, the invention also relates to an LSD1 inhibitor for use in the treatment of MPNST in combination with one or more further anticancer agents and in combination with an antiemetic agent (and optionally further in combination with radiotherapy); the invention likewise relates to corresponding methods and uses (including all methods and uses described herein above) comprising the combined administration of an LSD1 inhibitor, one or more further anticancer agents, and an antiemetic agent. The antiemetic agent may be, e.g., a 5-HT₃ antagonist (or a “setron”), such as, e.g., palonosetron (optionally in combination with netupitant), ramosetron, alosetron, ondansetron, tropisetron, granisetron, dolasetron, azasetron, bemesetron, cilansetron, lerisetron, ricasetron, or zatoksetron; olanzapine; a corticosteroid, such as, e.g., methylprednisolone or dexamethasone; or prochlorperazine.

Articles of Manufacture

[0088] The pharmaceutical compositions (or formulations) of the invention can be included in a container, pack or dispenser together with instructions for administration.

[0089] Thus, in a further embodiment, the present invention provides an article of manufacture containing an LSD1 inhibitor or a pharmaceutical composition comprising an LSD1 inhibitor for the treatment of MPNST as described herein.

[0090] In some embodiments, the article of manufacture comprises a container and a pharmaceutical composition for use in accordance with the invention as described herein.

[0091] In some embodiments, the invention provides an article of manufacture (or a kit) comprising a container and a combination product (as described herein above) for use in the treatment of MPNST. The invention also provides an article of manufacture (or a kit) comprising (i) a first container comprising the LSD1 inhibitor, (ii) a second container comprising a further anticancer agent (as described above), and (iii) optionally one or more further container(s) comprising one or more further anticancer agent(s), for use in the treatment of MPNST.

[0092] The article of manufacture may further comprise a label or package insert. The term “package insert” is used to refer to instructions customarily included in commercial packages of therapeutic products, that contain information about the indications, usage, dosage, administration, contraindications and/or warnings concerning the use of such therapeutic products. Suitable containers include, for example, blister packs, bottles, vials, syringes, etc. The container may be formed from a variety of materials such as glass or plastic. The container may hold a composition or formulation which is effective for treating the condition and may have a sterile access port (for example, the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). The label or package insert indicates that the composition is used for treating the condition of choice, particularly MPNST. Alternatively, or additionally, the article of manufacture may further comprise a second container comprising a pharmaceutically acceptable buffer, such as bacteriostatic water for injection (BWFI), phosphate-buffered saline, Ringer’s solution and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, and syringes.

[0093] The article of manufacture or kit may further comprise directions for the combined administration of one or more further anticancer agents (as described above). For example, if the kit comprises a first pharmaceutical composition/formulation comprising the LSD1 inhibitor and a second pharmaceutical composition/formulation comprising a further anticancer agent, the kit may further comprise directions for the simultaneous, sequential or separate administration of the first and the second pharmaceutical compositions/formulations to a subject in need thereof.

[0094] In another embodiment, the article of manufacture is suitable for the delivery of solid oral forms of the LSD1 inhibitor, such as tablets or capsules. Such an article of manufacture preferably includes a number of unit dosages. Such articles of manufacture can include a card having the dosages oriented in the order of their intended use. An example of such an article is a “blister pack”. Blister packs are well known in the packaging industry and are widely used for packaging pharmaceutical unit dosage forms. If desired, a memory aid can be provided, for example in the form of numbers, letters, or other markings or with a calendar insert, designating the days in the treatment schedule in which the dosages can be administered.

[0095] According to one embodiment, an article of manufacture or kit may comprise: (i) a first container with the LSD1 inhibitor contained therein; (ii) a second container with a further anticancer agent contained therein; and optionally (iii) a third container with a further anticancer agent contained therein, wherein the anticancer agent in the third container is different from the anticancer agent in the second container. Alternatively, or additionally, the kit may comprise another container comprising a pharmaceutically acceptable buffer, such as bacteriostatic water for injection (BWFI), phosphate-buffered saline, Ringer’s solution, or dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, and/or syringes.

[0096] Where the article of manufacture or kit comprises a composition of the LSD1 inhibitor and a composition of a further anticancer agent, the kit may comprise a container

for containing the separate compositions such as, e.g., a divided bottle or a divided foil packet; however, the separate compositions may also be contained within a single, undivided container. Typically, the kit comprises directions for the administration of the separate components. The kit form is particularly advantageous when the separate components are preferably administered in different dosage forms (e.g., oral and parenteral), are administered at different dosage intervals, or when titration of the individual components of the combination is desired by the prescribing physician or veterinarian.

Definitions

[0097] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention pertains.

[0098] The following definitions apply throughout the present specification and claims, unless specifically indicated otherwise. A “subject” (or “patient”) for the purposes of the present invention includes both humans and other animals, particularly mammals. Thus, the methods and uses of the invention are applicable to both human therapy and veterinary applications. In preferred embodiments, the subject (or patient) is a mammal (e.g., a human being or a non-human mammal), and most preferably the subject is a human (e.g., a male or female human). A human subject may have any age, including, e.g., 0 to 2 years, 2 to 12 years, 12 to 18 years, or 18 years or more.

[0099] The terms “treatment”, “treating” and the like are used herein to generally mean obtaining a desired pharmacological and/or physiological effect. This includes partially or completely curing or ameliorating a disease (i.e., MPNST) and/or a symptom or adverse effect attributed to the disease, or partially or completely halting the progression of the disease and/or a symptom or adverse effect attributed to the disease. The term “treatment” as used herein covers any treatment of a disease (i.e., MPNST) in a subject and includes, without limitation, inhibiting MPNST, i.e., arresting, delaying or slowing down its development/progression; or relieving MPNST, i.e., causing its (complete or partial) regression, remission, correction or alleviation. The present invention specifically and distinctly relates to each one of these forms of treatment.

[0100] As used herein, the term “therapeutically effective amount” or “effective amount” of a compound (particularly an LSD1 inhibitor) according to the invention refers to an amount sufficient to produce a desired biological effect (e.g., a therapeutic effect or benefit) in a subject. Accordingly, a therapeutically effective amount of a compound may be an amount which is sufficient to treat a disease (i.e., MPNST), and/or delay the onset or progression of the disease, and/or alleviate one or more symptoms of the disease, when administered to a subject suffering from or susceptible to that disease. The therapeutically effective amount will vary depending on the compound, the disease state being treated, the severity of the disease treated, the age and relative health of the subject, the route and form of administration, the judgment of the attending medical or veterinary practitioner, and other factors.

[0101] The term “pharmaceutically acceptable” denotes an attribute of a material which is useful in preparing a pharmaceutical composition that is generally safe, non-

toxic, and neither biologically nor otherwise undesirable and is acceptable for veterinary and/or human pharmaceutical use.

[0102] As used herein, a “pharmaceutically acceptable salt” is intended to mean a salt that retains the biological effectiveness of the free acids and/or bases of the specified compound and that is not biologically or otherwise undesirable. A compound may possess one or more sufficiently acidic or sufficiently basic functional groups, or both, and accordingly react with any of a number of inorganic or organic bases, and inorganic or organic acids, to form a pharmaceutically acceptable salt. Exemplary pharmaceutically acceptable salts include those salts prepared by reaction of a compound described herein (particularly an LSD1 inhibitor, such as, e.g., iadademstat), with a mineral or organic acid, such as hydrochlorides, hydrobromides, sulfates, pyrosulfates, bisulfates, sulfites, bisulfites, phosphates, monohydrophosphates, dihydrophosphates, metaphosphates, pyrophosphates, chlorides, bromides, iodides, nitrates, acetates, propionates, decanoates, caprylates, acrylates, formates, isobutyrate, caproates, heptanoates, propionates, oxalates, malonates, succinates, suberates, sebacates, fumarates, maleates, butyne-1,4-dioates, hexyne-1,6-dioates, benzoates, chlorobenzoates, methylbenzoates, dinitrobenzoates, hydroxybenzoates, methoxybenzoates, phthalates, sulfonates, xylenesulfonates, phenylacetates, phenylpropionates, phenylbutyrate, citrates, lactates, gamma-hydroxybutyrate, glycolates, tartrates, methane-sulfonates (or mesylates), ethane-sulfonates, propane-sulfonates, benzenesulfonates (or besylates), toluene-sulfonates, trifluoromethanesulfonates, naphthalene-1-sulfonates, naphthalene-2-sulfonates, mandelates, pyruvates, stearates, ascorbates, or salicylates. When a compound (particularly an LSD1 inhibitor) carries an acidic moiety, suitable pharmaceutically acceptable salts thereof may include: alkali metal salts, e.g. sodium or potassium salts; alkaline earth metal salts, e.g. calcium or magnesium salts; and salts formed with suitable organic ligands such as ammonia, alkylamines, hydroxyalkylamines, lysine, arginine, N-methylglucamine, procaine and the like. Pharmaceutically acceptable salts are well known in the art (see, e.g., Stahl PH & Wermuth CG (eds.), “Handbook of Pharmaceutical Salts: Properties, Selection, and Use”, Wiley-VCH, 2002 as well as the references cited therein, all of which are incorporated herein by reference).

[0103] The terms “pharmaceutical composition” and “pharmaceutical formulation” (or “formulation”) are used interchangeably and denote a mixture or solution comprising a therapeutically effective amount of an active pharmaceutical ingredient (particularly an LSD1 inhibitor) together with one or more pharmaceutically acceptable excipients to be administered to a subject (e.g., a human) in need thereof.

[0104] The terms “pharmaceutically acceptable excipient” or “pharmaceutically acceptable carrier” can be used interchangeably and denote any pharmaceutically acceptable ingredient in a pharmaceutical composition having no therapeutic activity and being non-toxic to the subject administered, such as disintegrators, binders, fillers, solvents, buffers, tonicity agents, stabilizers, antioxidants, surfactants, carriers, diluents, lubricants and the like used in formulating pharmaceutical products. They are generally safe for administering to humans according to established governmental standards, including those promulgated by the United States Food and Drug Administration and/or the European Medi-

cines Agency. Pharmaceutically acceptable carriers or excipients are well known to those skilled in the art.

[0105] The term “inhibitor”, as used herein, denotes a compound which competes with, decreases, blocks, inhibits, abrogates or interferes in any way with the binding of a particular ligand to a particular receptor or enzyme and/or which decreases, blocks, inhibits, abrogates or interferes in any way with the activity or function of a particular protein, e.g., of a receptor or enzyme.

[0106] As used herein, a “small molecule” refers to an organic compound with a molecular weight equal to or below 900 Da (daltons), preferably below 500 Da. The molecular weight is the mass of a molecule and is calculated as the sum of the atomic weights of each constituent element multiplied by the number of atoms of that element in the molecular formula.

[0107] As used herein, the term “comprising” (or “comprise”, “comprises”, “contain”, “contains”, or “containing”), unless explicitly indicated otherwise or contradicted by context, has the meaning of “containing, inter alia”, i.e., “containing, among further optional elements, . . .”. In addition thereto, this term also includes the narrower meanings of “consisting essentially of” and “consisting of”. For example, the term “A comprising B and C” has the meaning of “A containing, inter alia, B and C”, wherein A may contain further optional elements (e.g., “A containing B, C and D” would also be encompassed), but this term also includes the meaning of “A consisting essentially of B and C” and the meaning of “A consisting of B and C” (i.e., no other components than B and C are comprised in A).

[0108] As used herein, the indefinite articles “a” and “an” and the definite article “the” include plural as well as singular referents, unless the context clearly dictates otherwise.

[0109] The term “about” or “approximately” means an acceptable error for a particular value as determined by one of ordinary skill in the art, which depends in part on how the value is measured or determined. In certain embodiments, the term “about” or “approximately” means within 1, 2, 3 or 4 standard deviations. In certain embodiments, the term “about” or “approximately” means within 25%, 20%, 15%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, 0.5%, 0.1% or 0.05% of a given value or range. Any reference to a numerical value or range provided in connection with the term “about” also includes a reference to the corresponding specific value or range.

[0110] Furthermore, it is to be understood that wherever a numerical range is provided/described herein, all values and subranges encompassed by the respective numerical range are specifically provided by the invention. Accordingly, the present invention specifically and individually relates to each value that falls within a numerical range described herein, as well as each and any subrange encompassed by a numerical range described herein.

[0111] The present specification describes various compounds by their chemical formulae and their corresponding chemical names. In case of conflict between any chemical formula and the corresponding chemical name indicated herein, the present invention specifically and individually relates to the compound defined by the chemical formula and to the compound defined by the chemical name.

[0112] All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety.

EXAMPLES

[0113] The following examples are provided for illustration of the invention. They should not be considered as limiting the scope of the invention, but merely as being representative thereof.

Example 1: Effect of LSD1 Inhibitors in MPNST Viability Assay

1.1 Experimental Design

[0114] Mycoplasma-free MPNST cell lines (see Table 1) were seeded in 50 µl/well of complete medium (DMEM high glucose supplemented with 1× GlutaMAX™, 1× sodium pyruvate and 10% fetal bovine serum (FBS), Thermo Fisher) in 96-well plates at the optimal cell density to ensure log-phase growth throughout the duration of the experiment (see Table 1). The day after seeding, 50 µl of medium containing 9 serial dilutions (1:3) of 2×-concentrated iadademstat (LSD1 inhibitor; used as the dihydrochloride salt) were added to the cells to obtain 100 µl of cells treated with 1×-concentrated compound at each dilution. Etoposide (topoisomerase II/DNA synthesis inhibitor) was also included as a positive control of the assay. Additionally, from the MPNST cell line panel, two cell lines (sNF96.2 and sNF02.2) were selected to evaluate the effect of other LSD1 inhibitors, i.e. bomedemstat (bis-tosylate salt) and pulrodemstat (besylate salt). Each experimental condition was tested in technical triplicates, including medium-only wells and vehicle-treated controls for background correction and normalization, respectively. After treatment, cells were incubated at 37° C. in a humidified and controlled 5% CO₂ atmosphere for 72 h. At this time-point, compound and medium refreshment was performed by adding 50 µl of medium supplemented with 1×-concentrated compound at each corresponding dilution. Cells were incubated for additional 72 h (for a total of 6 days) prior to evaluating cell viability using either the MTT assay (Sigma-Aldrich) or AlamarBlue™ Cell viability reagent (Life Technologies), following manufacturer's instructions. Background was calculated as the mean of the values of medium-only controls and subtracted from each data point. The average of background-corrected technical triplicates was calculated and normalized by the mean of vehicle-treated controls (corresponding to 100% of viability). Data were analyzed using GraphPad PRISM® version 9.0.1 (GraphPad Software, Inc.,

La Jolla, CA/USA) to calculate the best-fitting curves and the EC₅₀ values (corresponding to the concentration of compound at which a half (50%) maximal effect is obtained; lower EC₅₀ values thus indicate greater potency).

TABLE 1

Cell lines used and their corresponding seeding densities	
Cell line	Cell density (cells/well)
MPNST-SP-01	1000
MPNST-NF1-08	8000
MPNST-NF1-09	1000
MPNST-NF1-18b	500
MPNST-SP-10	4000
ST88-14	2000
NMS-2	2000
S462	1000
90-8TL	2000
HS-PSS	2000
sNF02.2	1000
sNF96.2	8000

1.2 Results

[0115] A panel of 12 MPNST cell lines was used to evaluate the effects of the LSD1 inhibitor iadademstat on MPNST cell viability after 6 days of treatment, and etoposide was used as a positive control of the assay. In order for the in vitro experiments to reliably mirror the situation in the clinical setting, where MPNST patients feature a great genomic heterogeneity, a panel of different types of MPNST cell lines was used. The panel comprises both MPNST immortalized lines and MPNST cell lines derived from patients. Further, some MPNST cell lines are neurofibromatosis type I-linked while others are sporadic, and in both cases, there are variations in their genomic background in terms of mutational state for NF1, CDKN2A and PRC2. A summary of their characteristics is provided in Table 2 below.

TABLE 2

MPNST cell panel summary				
Cell line	MPNST subtype	Genomics		
		NF1	CDKN2A	PRC2
MPNST-SP-01	sporadic	partially inactivated	wild type	completely inactivated
MPNST-NF1-08	neurofibromatosis type I-linked	completely inactivated	completely inactivated	completely inactivated
MPNST-NF1-09	neurofibromatosis type I-linked	completely inactivated	completely inactivated	wild type
MPNST-NF1-18b*	neurofibromatosis type I-linked	completely inactivated	completely inactivated	completely inactivated
MPNST-SP-10	sporadic	unknown	completely inactivated	unknown
ST88-14	neurofibromatosis type I-linked	completely inactivated	completely inactivated	completely inactivated
NMS-2	neurofibromatosis type I-linked	unknown	completely inactivated	unknown

TABLE 2-continued

MPNST cell panel summary				
Cell line	MPNST subtype	Genomics		
		NF1	CDKN2A	PRC2
S462	neurofibromatosis type I-linked	completely inactivated	completely inactivated	completely inactivated
90-8TL	neurofibromatosis type I-linked	completely inactivated	completely inactivated	completely inactivated
HS-PSS	sporadic	wild type	completely inactivated	unknown
sNF02.2	neurofibromatosis type I-linked	partially inactivated	wild type	wild type
sNF96.2	neurofibromatosis type I-linked	completely inactivated	completely inactivated	completely inactivated

*The patient-derived MPNST-NF1-18b cell line was initially classified as a sporadic MPNST cell line and was named MPNST-SP-09b. After subsequent analysis this cell line was confirmed to be a neurofibromatosis type I-linked MPNST cell line and was renamed to MPNST-NF1-18b.

[0116] LSD1 inhibitors such as iadademstat have been reported to exert their therapeutic effect by inducing cancer cell differentiation and inhibiting cancer cell proliferation rather than by killing cancer cells (Sacilotto N et al., *ACS Pharmacol Transl Sci*, 2021, 4(6):1818-34, doi: 10.1021/acspsci.1c00223). In line with this, a cancer cell viability reduction of more than 30% reflects a potent therapeutic effect of the corresponding LSD1 inhibitor. In the present experiment, the response obtained after LSD1 inhibitor treatment was therefore classified into the following three groups: (1) strong response (viability reduction >30%); (2) medium response (viability reduction >15% and <30%); (3) low response (viability reduction <15%). Table 3 summarizes the EC₅₀ values obtained for etoposide (the positive control of the assay) and iadademstat together with the classification according to viability reduction after LSD1 inhibitor treatment.

TABLE 3

Effects of etoposide (positive control of the assay) and the LSD1 inhibitor iadademstat on cell viability of the MPNST cell line panel

Cell line	EC ₅₀ (nM) etoposide (positive control)	EC ₅₀ (nM) iadademstat	Response to iadademstat (viability assay)
MPNST-NF1-18b	393.9	0.025	Strong response
sNF96.2	107.6	0.042	(>30% viability reduction)
S462	375.9	0.087	Medium response
sNF02.2	144.2	0.017	(15-30% viability reduction)
90-8TL	164.4	0.026	
MPNST-SP-10	269.0	0.033	
ST88-14	788.5	0.039	
HS-PSS	204.8	0.046	
NMS-2	153.6	Not determined	Low response
MPNST-SP-01	1030.0	Not determined	(<15% viability reduction)
MPNST-NF1-08	740.3	Not determined	
MPNST-NF1-09	737.0	Not determined	

[0117] In spite of the huge genomic and mutational variations within the MPNST cell line panel tested, a remarkable 8 out of 12 cell lines responded advantageously well to LSD1 inhibitor treatment. Thus, after a 6-day iadademstat treatment, 3 out of 12 cell lines (25%) showed a strong viability reduction and 5 out of 12 cell lines (41.67%) showed a medium viability reduction. Notably, particularly responsive cell lines include both neurofibromatosis type I-linked and sporadic (not neurofibromatosis type I-linked) MPNST cell lines, which indicates that treatment with an

LSD1 inhibitor (such as iadademstat) achieves a therapeutic effect in a broad range of MPNST patient subpopulations, including neurofibromatosis type I-linked MPNST as well as MPNST not linked to neurofibromatosis type I. Moreover, in both the medium- and strong-responding cell lines, which account for a total of 66.67%, iadademstat displayed sub-nanomolar EC₅₀ values (see Table 3), which points at a clinically relevant therapeutic effect even when administered at very low doses.

[0118] The effect of LSD1 inhibitors on cell viability of MPNST was further tested using 2 additional LSD1 inhibitors, namely bomedemstat and pulrodemstat. Bomedemstat, like iadademstat, is an irreversible LSD1 inhibitor, whereas pulrodemstat is a reversible LSD1 inhibitor. Cell viability was evaluated after a 6-day treatment in both sNF96.2 and sNF02.2 cell lines as per the method described above (see Table 4). The results thus obtained, as presented in Table 4 (and, for iadademstat, in Table 3 above), clearly show that all three LSD1 inhibitors are effective against MPNST, irrespective of whether they are irreversible or reversible LSD1 inhibitors.

TABLE 4

Effect of different LSD1 inhibitors on cell viability of sNF96.2 and sNF02.2 cell lines		
LSD1 inhibitor	sNF96.2 EC ₅₀ (nM)	sNF02.2 EC ₅₀ (nM)
Bomedemstat	1.277	0.094
Pulrodemstat	0.261	0.166

[0119] Among the LSD1 inhibitors tested, iadademstat displayed the highest potency (lowest EC₅₀) in both sNF96.2 and sNF02.2 cell lines.

[0120] These results show that LSD1 inhibitors, including both irreversible LSD1 inhibitors (such as iadademstat and bomedemstat) as well as reversible LSD1 inhibitors (such as pulrodemstat), are advantageously effective in the treatment of MPNST and can be used for a wide range of different subgroups of MPNST patients.

Example 2: Matrix Assay for Determination of Synergism Between LSD1 Inhibitors and MEK Inhibitors in MPNST Cell Lines

2.1 Experimental Design

2.1.1 Matrix Viability Assays (6 Days)

[0121] Each matrix assay was distributed either across 2 plates in a 9×9 format following the scheme illustrated in FIG. 1, or across 1 plate in a 5×5 format following the scheme illustrated in FIG. 2. The LSD1 inhibitor iadademstat was added at increasing concentrations from top to bottom, and the MEK inhibitor selumetinib was added at increasing concentrations from left to right.

[0122] For the assay, cells were seeded in 96-well plates at the optimal density specified in Example 1 in 50 µL of medium; the wells at the edges of the plates were filled with 100 µL of medium-only for background correction. Each of the two compounds of the combination was added at a 4×-concentration in 25 µL, resulting in a final volume of 100 µL and final concentration of 1× at each dilution (DMSO %<0.5%). For the 9×9 matrices compounds were added in 1:2 serial dilutions, whereas for the 5×5 matrices compounds were added in 1:5 serial dilutions, such that in both cases the concentration range covered the full dose-response curve. As shown in FIGS. 1 and 2, the matrices were designed to have the expected EC₅₀ values of both compounds centered horizontally and vertically on the matrix. In this way, the wells on the diagonal of the plates correspond to the fixed EC₅₀ ratios between both compounds. In the 9×9 matrix, the first and the last row of plate #1 have been repeated in plate #2 (indicated by arrows in FIG. 1), to confirm reproducibility across the two plates. The EC₅₀ values for the compounds tested in the matrix assays were previously obtained through single agent assays performed as detailed in Example 1 and, for iadademstat, are shown in Table 3. After treatment, cells were incubated at 37° C. in a humidified and controlled 5% CO₂ atmosphere for 72 h. At this time-point, compound and medium refreshment was performed by adding 50 µl of medium supplemented with 1×-concentrated compound at each corresponding dilution. Cells were incubated for additional 72 h (for a total of 6 days) prior to evaluating cell viability in 2 biological replicates and using either the MTT assay (Sigma-Aldrich) or AlamarBlue™ Cell viability reagent (Life Technologies), following manufacturer's instructions. Background was calculated as the mean of the values of medium-only controls and subtracted from each data point. Background-corrected values were normalized by the corrected vehicle-treated controls (corresponding to 100% of viability). Data were analyzed using GraphPad PRISM® version 9.0.1 (GraphPad Software, Inc., La Jolla, CA/USA) to calculate the best-fitting curves and the EC₅₀ values (corresponding to the concentration of compound at which a half (50%) maximal effect is obtained; lower EC₅₀ values thus indicate greater potency).

2.1.2 Matrix Viability Assays (Data Analysis)

[0123] For each matrix assay, data were normalized against the vehicle-treated controls (<0.5% DMSO, in the upper left corner) to obtain the percentage value of relative residual viability, according to the following formula:

$$\% \text{ relative residual viability} = \text{Background-corrected signal treated cells}/$$

$$\text{Background-corrected signal vehicle control} \times 100$$

[0124] The values for percentage of residual viability were then analyzed using GraphPad PRISM® version 9.0.1 (GraphPad Software, Inc., La Jolla, CA/USA) to calculate the best-fitting curve and the EC₅₀ values of the single agents.

[0125] At this point the Fraction affected (Fa), also known as Fractional Effect, was calculated using the formula:

$$Fa = -1(\% \text{ relative residual viability}/100)$$

for the following conditions:

[0126] Cells treated with serial dilutions of selumetinib as single agent.

[0127] Cells treated with serial dilutions of iadademstat as single agent.

[0128] Cells treated with iadademstat and selumetinib at a fixed ratio corresponding to the ratio of EC₅₀ values (values of % relative residual viability in the diagonal of the matrix assay; highlighted in FIGS. 1 and 2).

[0129] The CalcuSyn software (<http://www.biosoft.com/w/calculsyn.htm>, Biosoft, Cambridge, UK) is designed to determine the nature (synergistic, additive or antagonistic) of the interaction between two compounds by calculating a Combination Index (CI). This analysis is based on the Median Effect Principle and the Combination Index Theorem described by the Chou-Talalay method (Chou TC, Pharmacol Rev, 2006, 58(3):621-681, doi: 10.1124/pr.58.3.10), where a resulting CI<1 is indicative of synergistic effects, a CI=1 indicates additive effects, while CI>1 reflects antagonistic effects. In the case of synergistic effects (CI<1), the smaller the CI value is, the stronger the synergy. Additionally, the strength of the drug interactions can be further classified based on the CI range, as shown in Table 5.

TABLE 5

Classification of drug interactions strength based on CI. (Chou TC, Pharmacol Rev, 2006, 58(3): 621-681, doi: 10.1124/pr.58.3.10)		
	CI	Classification
Synergistic	<0.1	+++++
	0.1-0.3	++++
	0.3-0.7	+++
	0.7-0.85	++
	0.85-0.9	+
Additive	0.9-1.1	+/-
	1.1-1.20	-
	1.20-1.45	—
	1.45-3.3	—
	3.3-10	—
Antagonistic	>10	—

[0130] In order to generate informative and consistent results, the data processed with CalcuSyn (both for the single agents and the drug combination) need to fit with the Median Effect Principle and the Combination Index Theorem theoretical models. For this reason, it is crucial to remove possible outliers and data points characterized by

poor fit to the Median Effect Principle (Chou TC, Pharmacol Rev, 2006, 58(3):621-681, doi: 10.1124/pr.58.3.10). In order to achieve this, the following strategy was adopted for data filtering:

[0131] In the first step data dispersion was reduced by removing points characterized by:

[0132] 1) $Fa < 0.1$

[0133] 2) Increase in $Fa < 0.03$, compared to the previous point (if $Fa > 0.9$).

[0134] These conditions define the plateaus of the dose response curve, in which cells have been treated with very low or very high concentrations of compounds (or combos), resulting in reduction of viability close to 0% or 100% (equivalent to Fa value close to 0 or 1, respectively). To be noted, in these areas of the dose-response curves the changes in alamarBlue™ or MTT signals are very small and most likely due to random noise with very little biological significance. Next, for each data point, $\log_{10}$ (Concentration) and $\log_{10}(Fa/(1-Fa))$ were calculated and a dot plot graph was generated reporting the former value on the x axis and the latter on the y axis. With Excel, a regression line was then obtained (corresponding to the Median Effect Equation).

[0135] At this point the distance from the regression line was calculated for each data point with the equation:

$$\text{Distance } (ax + by + c = 0; X, Y) = (aX + bY + c) / \sqrt{a^2 + b^2}$$

[0136] Outliers are identified on the basis of their distance from the Median Effect Equation, using the Grubbs's test. For each data point, the Grubbs's test was performed on the absolute value of the distance, according to the following formula (to be noted, the variable for the Grubbs's test can be called interchangeably G or Z):

$$G = (X_n - X_{\text{average}}) / s$$

where X_n stands for the absolute value of the distance of each point from the regression line; X_{average} stands for average of all the X_n values and s stands for the standard deviation. Values of G above G_{crit} (calculated for $\alpha=0.2$ as shown below) identify outliers not fitting on the Median Effect Equation. Such data points have been removed to successfully calculate the Combination Index with CalcuSyn.

$$G_{\text{crit}} = \frac{(n-1)t_{\text{crit}}}{\sqrt{n(n-2+t_{\text{crit}}^2)}}$$

[0137] When possible, the test was reiterated more than once to remove multiple outliers, until:

[0138] 1. no further outliers were identified or

[0139] 2. $R^2 > 0.95$. To measure data quality, the R value is calculated also by the CalcuSyn software (good data are characterized by R value above 0.95).

2.1.3 CalcuSyn Output

[0140] CalcuSyn results are provided as the experimental Fractional Effect (referred to as Fa) representing the fraction of cells affected by the combined treatment at their fixed EC_{50} ratio (in the case of a cytotoxic treatment the Fractional Effect corresponds to viability reduction compared to vehicle controls, where $Fa=1$ is equal to 100% viability reduction) and the associated combination index (CI). As shown in Table 5 above, the CI value is indicative of the nature and strength of the compounds' interaction, with values below 1 representing synergistic interactions (the closer the value to 0, the stronger the synergistic effects), values equal to 1 representing additive interactions, and values above 1 representing antagonistic interactions. The software also provides the simulation of CI and Fa based on experimental data, and provides the estimated CI at ED75 and ED90 (Effective Doses corresponding to 75% and 90% viability reduction, respectively), as for anticancer or antiviral agents, synergy at high effect levels (e.g., at $Fa > 0.75$) is more relevant to therapy than at low effect levels (e.g., at $Fa < 0.2$) as described in Chou TC, Cancer Res (2010) 70 (2): 440-446, doi: 10.1158/0008-5472.CAN-09-1947.

[0141] Data are provided as both the experimental CI for the associated Fa for each experiment and the mean estimated CI for ED75 and ED90 values from 2 independent biological replicates.

2.2 Results

[0142] Matrix treatments with the MEK inhibitor selumetinib and the covalent and irreversible LSD1 inhibitor iadademstat were performed as described in section 2.1.1. Data analysis and calculation of combination indexes were performed as described in section 2.1.2. The results of the combination indexes (CI) associated with specific fractional effects (Fa), the estimated CI at ED75 and ED90 and the respective classifications (as described in Table 5) obtained from the combination of iadademstat and selumetinib are shown in Table 6 and Table 7.

[0143] In summary, the combination iadademstat+selumetinib showed strong synergism in a wide range of fractional effects (Fa) in the three cell lines tested, including the patient derived MPNST-NF1-18b cell line. Importantly, these synergistic effects were observed at biologically relevant effective doses (ED75 and ED90).

TABLE 6

Experimental CI values associated with the corresponding Fa and their classification for the iadademstat + selumetinib combination. Results of two independent biological replicates are shown.						
iadademstat + selumetinib						
Cell line	Experiment 1			Experiment 2		
	Fa	CI	Classification	Fa	CI	Classification
sNF96.2	0.21	0.69	+++	0.38	0.33	+++
	0.43	0.90	+	0.53	0.26	++++
	0.95	0.41	+++	0.76	0.05	+++++
sNF02.2	0.25	0.73	++	0.24	1.25	—
	0.60	0.55	+++	0.48	0.55	+++
	0.75	0.55	+++	0.72	0.27	++++
	0.86	0.52	+++	0.81	0.43	+++

TABLE 6-continued

Experimental CI values associated with the corresponding Fa and their classification for the iadademstat + selumetinib combination. Results of two independent biological replicates are shown.						
iadademstat + selumetinib						
Cell line	Experiment 1			Experiment 2		
	Fa	CI	Classification	Fa	CI	Classification
MPNST-NF1-18b	0.25	0.78	++	0.12	1.53	---
	0.37	0.79	++	0.18	1.54	---
	0.44	1.14	-	0.40	0.66	+++
	0.68	0.70	+++	0.52	0.69	+++
	0.83	0.51	+++	0.62	0.77	++
	0.94	0.24	++++	0.76	0.58	+++
				0.87	0.39	+++
				0.94	0.24	++++

TABLE 7

Estimated CI values for ED75 and ED90 and their classification for the iadademstat + selumetinib combination.			
iadademstat + selumetinib mean estimated CI			
	ED	CI	Classification
sNF96.2	ED75	0.29	++++
	ED90	0.27	++++
sNF02.2	ED75	0.46	+++
	ED90	0.35	+++
MPNST-NF1-18b	ED75	0.52	+++
	ED90	0.37	+++

The mean CI of two independent biological replicates is shown.

[0144] These results show that LSD1 inhibitors (such as iadademstat) and MEK inhibitors (such as selumetinib) interact synergistically in a panel of different MPNST cell lines, which renders the combination of these agents particularly advantageous for the treatment of MPNST.

Example 3: Matrix Assay for Determination of Synergism Between LSD1 Inhibitors and Pi3K Inhibitors in MPNST Cell Lines

3.1 Experimental Design

[0145] Matrix viability assays and the corresponding analyses were performed as described in Example 2 but with the LSD1 inhibitor iadademstat and the Pi3K inhibitor copanlisib.

3.2 Results

[0146] Matrix treatments with the Pi3K inhibitor copanlisib and the covalent and irreversible LSD1 inhibitor iadademstat were performed as described in section 2.1.1. Data analysis and calculation of combination indexes were performed as described in section 2.1.2. The results of the combination indexes (CI) associated with specific fractional effects (Fa), the estimated CI at ED75 and ED90 and the respective classifications (as described in Table 5) obtained from the combination of iadademstat and copanlisib are shown in Table 8 and Table 9.

[0147] In summary, the combination iadademstat+copanlisib showed synergism in a wide range of fractional effects

(Fa). Importantly, these synergistic effects were observed at biologically relevant effective doses (ED75 and ED90).

TABLE 8

Experimental CI values associated with the corresponding Fa and their classification for the iadademstat + copanlisib combination. Results of two independent biological replicates are shown.						
iadademstat + copanlisib						
Cell line	Experiment 1			Experiment 2		
	Fa	CI	Classification	Fa	CI	Classification
sNF96.2	0.17	1.04	+/-	0.25	4.35	----
	0.55	0.68	+++	0.61	0.97	+/-
	0.93	0.24	++++	0.87	0.27	++++
	0.97	0.46	+++			
sNF02.2	0.48	0.64	+++	0.15	4.25	----
	0.78	0.32	+++	0.64	0.33	+++
	0.86	0.62	+++	0.80	0.37	+++
				0.87	0.79	++
MPNST-NF1-18b	0.18	1.49	---	0.26	1.15	-
	0.40	0.73	++	0.40	0.90	+
	0.50	0.85	++	0.51	0.96	+/-
	0.73	0.49	+++	0.66	0.79	++
	0.80	0.58	+++	0.79	0.64	+++
	0.88	0.55	+++	0.85	0.73	++
	0.93	0.54	+++	0.91	0.59	+++
				0.95	0.45	+++

TABLE 9

Estimated CI values for ED75 and ED90 and their classification for the iadademstat + copanlisib combination.			
iadademstat + copanlisib mean estimated CI			
	ED	CI	Classification
sNF96.2	ED75	0.54	+++
	ED90	0.31	+++
sNF02.2	ED75	0.56	+++
	ED90	0.44	+++
MPNST-NF1-18b	ED75	0.68	+++
	ED90	0.54	+++

The mean CI of two biological replicates is shown.

[0148] These results demonstrate that LSD1 inhibitors (such as iadademstat) and Pi3K inhibitors (such as copanlisib) interact synergistically in different MPNST cell lines, which makes the combination of such agents particularly advantageous in the treatment of MPNST.

[0149] While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications and this patent or patent application is intended to cover any variations, uses or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure as come within known or customary practice within the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth and as follows in the appended claims.

1. An LSD1 inhibitor for use in the treatment of malignant peripheral nerve sheath tumor (MPNST).

2. A pharmaceutical composition comprising an LSD1 inhibitor and optionally one or more pharmaceutically acceptable excipients for use in the treatment of malignant peripheral nerve sheath tumor (MPNST).

3. A method of treating malignant peripheral nerve sheath tumor (MPNST) in a subject in need thereof, comprising administering a therapeutically effective amount of an LSD1 inhibitor to the subject.

4. Use of an LSD1 inhibitor for the treatment of malignant peripheral nerve sheath tumor (MPNST).

5. Use of an LSD1 inhibitor for the preparation of a pharmaceutical composition for the treatment of malignant peripheral nerve sheath tumor (MPNST).

6. The LSD1 inhibitor for use according to claim 1, the pharmaceutical composition for use according to claim 2, the method of claim 3, or the use of claim 4 or 5, wherein the LSD1 inhibitor is a small molecule.

7. The LSD1 inhibitor for use according to claim 1 or 6, the pharmaceutical composition for use according to claim 2 or 6, the method of claim 3 or 6, or the use of any one of claims 4 to 6, wherein the LSD1 inhibitor is selected from the group consisting of iadademstat, pulrodemstat, bomedemstat, seclidemstat, 1-((4-(methoxymethyl)-4-(((1R,2S)-2-phenylcyclopropylamino)methyl)piperidin-1-yl)methyl)cyclobutanecarboxylic acid, 3-(cyanomethyl)-3-(4-(((1R,2S)-2-phenylcyclopropylamino)piperidin-1-yl)azetidine-1-sulfonamide, 4-[5-((3S)-3-aminopyrrolidine-1-carbonyl)-2-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]phenyl]-2-fluoro-benzonitrile, and pharmaceutically acceptable salts thereof.

8. The LSD1 inhibitor for use according to any one of claims 1, 6 or 7, the pharmaceutical composition for use according to any one of claims 2, 6 or 7, the method of any one of claims 3, 6 or 7, or the use of any one of claims 4 to 7, wherein the LSD1 inhibitor is selected from the group consisting of iadademstat, pulrodemstat, bomedemstat, and pharmaceutically acceptable salts thereof.

9. The LSD1 inhibitor for use according to any one of claims 1 or 6 to 8, the pharmaceutical composition for use according to any one of claims 2 or 6 to 8, the method of any one of claims 3 or 6 to 8, or the use of any one of claims 4 to 8, wherein the LSD1 inhibitor is iadademstat or a pharmaceutically acceptable salt thereof.

10. The LSD1 inhibitor for use according to claim 9, the pharmaceutical composition for use according to claim 9, the method of claim 9, or the use of claim 9, wherein the LSD1 inhibitor is iadademstat dihydrochloride.

11. The LSD1 inhibitor for use according to any one of claims 1 or 6 to 8, the pharmaceutical composition for use according to any one of claims 2 or 6 to 8, the method of any one of claims 3 or 6 to 8, or the use of any one of claims 4 to 8, wherein the LSD1 inhibitor is pulrodemstat or a pharmaceutically acceptable salt thereof.

12. The LSD1 inhibitor for use according to claim 11, the pharmaceutical composition for use according to claim 11, the method of claim 11, or the use of claim 11, wherein the LSD1 inhibitor is pulrodemstat besylate.

13. The LSD1 inhibitor for use according to any one of claims 1 or 6 to 8, the pharmaceutical composition for use according to any one of claims 2 or 6 to 8, the method of any one of claims 3 or 6 to 8, or the use of any one of claims 4 to 8, wherein the LSD1 inhibitor is bomedemstat or a pharmaceutically acceptable salt thereof.

14. The LSD1 inhibitor for use according to claim 13, the pharmaceutical composition for use according to claim 13, the method of claim 13, or the use of claim 13, wherein the LSD1 inhibitor is bomedemstat bis-tosylate.

15. The LSD1 inhibitor for use according to any one of claims 1 or 6 to 14, the pharmaceutical composition for use according to any one of claims 2 or 6 to 14, the method of any one of claims 3 or 6 to 14, or the use of any one of claims 4 to 14, wherein the MPNST is neurofibromatosis type I-linked MPNST, sporadic MPNST, or radiation-induced MPNST.

16. The LSD1 inhibitor for use according to any one of claims 1 or 6 to 14, the pharmaceutical composition for use according to any one of claims 2 or 6 to 14, the method of any one of claims 3 or 6 to 14, or the use of any one of claims 4 to 14, wherein the LSD1 inhibitor or the pharmaceutical composition is administered to a subject having neurofibromatosis type I.

17. The LSD1 inhibitor for use according to any one of claims 1 or 6 to 14, the pharmaceutical composition for use according to any one of claims 2 or 6 to 14, the method of any one of claims 3 or 6 to 14, or the use of any one of claims 4 to 14, wherein the LSD1 inhibitor or the pharmaceutical composition is administered to a subject not having neurofibromatosis type I.

18. The LSD1 inhibitor for use according to any one of claims 1 or 6 to 17, the pharmaceutical composition for use according to any one of claims 2 or 6 to 17, the method of any one of claims 3 or 6 to 17, or the use of any one of claims 4 to 17, wherein the MPNST has one or more mutations or genetic alterations affecting the NF1 gene.

19. The LSD1 inhibitor for use according to any one of claims 1 or 6 to 18, the pharmaceutical composition for use according to any one of claims 2 or 6 to 18, the method of any one of claims 3 or 6 to 18, or the use of any one of claims 4 to 18, wherein the MPNST has one or more mutations or genetic alterations affecting the expression and/or the activity of CDKN2A, p53, RB1, PTEN, PRC2, EGFR, PDGFRA and/or c-Met.

20. The LSD1 inhibitor for use according to any one of claims 1 or 6 to 19, the pharmaceutical composition for use according to any one of claims 2 or 6 to 19, the method of any one of claims 3 or 6 to 19, or the use of any one of claims 4 to 19, wherein the MPNST is a metastatic MPNST.

21. The LSD1 inhibitor for use according to any one of claims 1 or 6 to 20, the pharmaceutical composition for use according to any one of claims 2 or 6 to 20, the method of any one of claims 3 or 6 to 20, or the use of any one of claims 4 to 20, wherein the LSD1 inhibitor or the pharmaceutical composition is administered to a subject which is a human.

22. The LSD1 inhibitor for use according to any one of claims 1 or 6 to 21, the pharmaceutical composition for use according to any one of claims 2 or 6 to 21, the method of any one of claims 3 or 6 to 21, or the use of any one of claims 4 to 21, wherein the LSD1 inhibitor or the pharmaceutical composition is administered orally.

23. The LSD1 inhibitor for use according to any one of claims 1 or 6 to 22, the pharmaceutical composition for use according to any one of claims 2 or 6 to 22, the method of any one of claims 3 or 6 to 22, or the use of any one of claims 4 to 22, wherein the LSD1 inhibitor or the pharmaceutical composition is administered in combination with one or more further anticancer agents and/or in combination with radiotherapy.

24. An article of manufacture comprising, in the same pharmaceutical formulation or in separate pharmaceutical formulations, an LSD1 inhibitor and one or more further

anticancer agents, for use in the treatment of malignant peripheral nerve sheath tumor (MPNST).

25. A method of treating malignant peripheral nerve sheath tumor (MPNST) in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of an article of manufacture comprising, in the same pharmaceutical formulation or in separate pharmaceutical formulations, an LSD1 inhibitor and one or more further anticancer agents.

26. The article of manufacture for use according to claim **24**, or the method of claim **25**, wherein:

the LSD1 inhibitor is as defined in any one of claims **6** to **14**; and/or

the MPNST is as defined in any one of claims **15** or **18** to **20**; and/or

the subject to whom the article of manufacture is administered is as defined in any one of claims **16**, **17** or **21**; and/or

the article of manufacture is administered orally.

27. The article of manufacture for use according to claim **24** or **26**, or the method of claim **25** or **26**, wherein the one or more further anticancer agents comprise a MEK inhibitor.

28. The LSD1 inhibitor for use according to any one of claims **1** or **6** to **23**, the pharmaceutical composition for use according to any one of claims **2** or **6** to **23**, the method of any one of claims **3** or **6** to **23**, or the use of any one of claims **4** to **23**, wherein the LSD1 inhibitor or the pharmaceutical composition is administered in combination with a MEK inhibitor.

29. The LSD1 inhibitor for use according to claim **28**, the pharmaceutical composition for use according to claim **28**, the article of manufacture for use according to claim **27**, the method of claim **27** or **28**, or the use of claim **28**, wherein the MEK inhibitor is selected from selumetinib, trametinib, cobimetinib, binimetinib, mirdametinib, pimasertib, refametinib, zapnometinib, avutometinib, HL-085, FCN-159, TAK-733, and pharmaceutically acceptable salts thereof.

30. The LSD1 inhibitor for use according to claim **28**, the pharmaceutical composition for use according to claim **28**, the article of manufacture for use according to claim **27**, the method of claim **27** or **28**, or the use of claim **28**, wherein the MEK inhibitor is selumetinib or a pharmaceutically acceptable salt thereof, and the LSD1 inhibitor is iadademstat or a pharmaceutically acceptable salt thereof.

31. The article of manufacture for use according to claim **24** or **26**, or the method of claim **25** or **26**, wherein the one or more further anticancer agents comprise a Pi3K inhibitor.

32. The LSD1 inhibitor for use according to any one of claims **1** or **6** to **23**, the pharmaceutical composition for use according to any one of claims **2** or **6** to **23**, the method of any one of claims **3** or **6** to **23**, or the use of any one of claims **4** to **23**, wherein the LSD1 inhibitor or the pharmaceutical composition is administered in combination with a Pi3K inhibitor.

33. The LSD1 inhibitor for use according to claim **32**, the pharmaceutical composition for use according to claim **32**, the article of manufacture for use according to claim **31**, the method of claim **31** or **32**, or the use of claim **32**, wherein the Pi3K inhibitor is selected from copanlisib, alpelisib, idelalisib, duvelisib, umbralisib, buparlisib, zandelisib, linperlisib, piasclisib, leniolisib, paxalisib, inavolisib, serabelisib, pictilisib, taselisib, tenalisib, eganelisib, GSK2636771, MEN1611, AMG-319, and pharmaceutically acceptable salts thereof.

34. The LSD1 inhibitor for use according to claim **32**, the pharmaceutical composition for use according to claim **32**, the article of manufacture for use according to claim **31**, the method of claim **31** or **32**, or the use of claim **32**, wherein the Pi3K inhibitor is copanlisib or a pharmaceutically acceptable salt thereof, and the LSD1 inhibitor is iadademstat or a pharmaceutically acceptable salt thereof.

* * * * *