



US012403183B2

(12) **United States Patent**
Maloney et al.(10) **Patent No.: US 12,403,183 B2**(45) **Date of Patent: *Sep. 2, 2025**(54) **FACTOR VIII POLYPEPTIDE
FORMULATIONS**(71) Applicant: **BIOVERATIV THERAPEUTICS
INC.**, Waltham, MA (US)(72) Inventors: **Kevin Maloney**, Waltham, MA (US);
Daniel Gage, Waltham, MA (US);
Ahmad Abdul-Fattah, Waltham, MA
(US)(73) Assignee: **BIOVERATIV THERAPEUTICS
INC.**, Waltham, MA (US)(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 910 days.This patent is subject to a terminal dis-
claimer.(21) Appl. No.: **16/993,979**(22) Filed: **Aug. 14, 2020**(65) **Prior Publication Data**

US 2021/0069300 A1 Mar. 11, 2021

Related U.S. Application Data(62) Division of application No. 15/455,043, filed on Mar.
9, 2017, now Pat. No. 10,786,554, which is a division
of application No. 14/213,180, filed on Mar. 14, 2014,
now Pat. No. 9,623,088.(60) Provisional application No. 61/897,742, filed on Oct.
30, 2013, provisional application No. 61/879,955,
filed on Sep. 19, 2013, provisional application No.
61/876,927, filed on Sep. 12, 2013, provisional
application No. 61/863,860, filed on Aug. 8, 2013,
provisional application No. 61/839,477, filed on Jun.
26, 2013, provisional application No. 61/829,884,
filed on May 31, 2013, provisional application No.
61/817,085, filed on Apr. 29, 2013, provisional
application No. 61/800,293, filed on Mar. 15, 2013.(51) **Int. Cl.**
A61K 38/37 (2006.01)
C07K 14/755 (2006.01)
C07K 16/28 (2006.01)(52) **U.S. Cl.**
CPC **A61K 38/37** (2013.01); **C07K 14/755**
(2013.01); **C07K 16/2833** (2013.01); **C07K**
2319/30 (2013.01)(58) **Field of Classification Search**
None
See application file for complete search history.(56) **References Cited****U.S. PATENT DOCUMENTS**4,757,006 A 7/1988 Toole, Jr. et al.
4,868,112 A 9/1989 Toole, Jr.
4,965,199 A 10/1990 Capon et al.
4,994,371 A 2/1991 Davie et al.5,004,803 A 4/1991 Kaufman et al.
5,112,950 A 5/1992 Meulien et al.
5,171,844 A 12/1992 Van Ooyen et al.
5,364,771 A 11/1994 Lollar et al.
5,543,502 A 8/1996 Nordfang et al.
5,595,886 A 1/1997 Chapman et al.
5,610,278 A 3/1997 Nordfang et al.
5,712,122 A 1/1998 Boime et al.
5,733,873 A * 3/1998 Osterberg A61K 38/37
530/383
5,763,401 A 6/1998 Nayar
5,789,203 A 8/1998 Chapman et al.
5,859,204 A 1/1999 Lollar
5,972,885 A 10/1999 Spira et al.
6,048,720 A 4/2000 Dalborg et al.
6,060,447 A 5/2000 Chapman et al.
6,200,560 B1 3/2001 Couto et al.
6,228,620 B1 5/2001 Chapman et al.
6,251,632 B1 6/2001 Lillicrap et al.
6,316,226 B1 11/2001 Van Ooyen et al.
6,346,513 B1 2/2002 Van Ooyen et al.
6,376,463 B1 4/2002 Lollar
6,458,563 B1 10/2002 Lollar
6,686,179 B2 2/2004 Fleer et al.
7,041,635 B2 5/2006 Kim et al.
7,348,004 B2 3/2008 Peters et al.
7,404,956 B2 7/2008 Peters et al.
7,592,010 B2 9/2009 Rosen et al.

(Continued)

FOREIGN PATENT DOCUMENTSCN 1399560 A 2/2003
EP 0154316 A2 9/1985

(Continued)

OTHER PUBLICATIONSU.S. Appl. No. 14/213,180, filed Mar. 14, 2014, Kevin Maloney,
2014/0308280, Oct. 16, 2014, now U.S. Pat. No. 9,623,088, Apr. 18,
2017.U.S. Appl. No. 15/455,043, filed Mar. 9, 2017, Kevin Maloney,
2017/0281734, Oct. 5, 2017, now U.S. Pat. No. 10,786,554, Sep. 29,
2020.

(Continued)

Primary Examiner — Michael Szperka(74) *Attorney, Agent, or Firm* — LATHROP GPM LLP;
James H. Velema, Esq.(57) **ABSTRACT**The present invention provides a formulation of a Factor
VIII polypeptide, e.g., FVIII-Fc, and methods of using the
same. The FVIII polypeptide can be a recombinant FVIII
protein, a short-acting FVIII protein, or a long-acting FVIII
protein. The pharmaceutical formulation comprising a FVIII
polypeptide can be used for individual prophylaxis, weekly
prophylaxis, episodic (on-demand) treatment, or periopera-
tive management of hemophilia.**44 Claims, 2 Drawing Sheets****Specification includes a Sequence Listing.**

(56)

References Cited**U.S. PATENT DOCUMENTS**

7,632,921	B2	12/2009	Pan et al.
7,862,820	B2	1/2011	Peters et al.
8,239,182	B2	8/2012	Kanade
8,815,250	B2	8/2014	Rivera et al.
9,050,318	B2	6/2015	Dumont et al.
9,623,088	B2 *	4/2017	Maloney A61P 7/00
10,786,554	B2	9/2020	Maloney et al.
2005/0100990	A1	5/2005	Saenko et al.
2009/0087411	A1	4/2009	Defrees
2009/0163699	A1	6/2009	Skerra et al.
2009/0264627	A1	10/2009	Oestergaard et al.
2010/0286067	A1	11/2010	Defrees
2010/0292130	A1	11/2010	Maloney et al.
2012/0093840	A1	4/2012	Fares et al.
2014/0308280	A1	10/2014	Chamberlain et al.
2017/0281734	A1	10/2017	Gillies et al.

FOREIGN PATENT DOCUMENTS

EP	0295597	A2	12/1988	
EP	0401384	A1	12/1990	
EP	2173890	A1	4/2010	
EP	2968477	A1	1/2016	
EP	3666283	A1	6/2020	
JP	H10501522	A	2/1998	
WO	WO 19870004187	A1	7/1987	
WO	WO 19880000831	A1	2/1988	
WO	WO 19880003558	A1	5/1988	
WO	WO 1988008035	A1	10/1988	
WO	WO 1991009122	A1	6/1991	
WO	WO 1992016221	A1	10/1992	
WO	WO 1993020093	A1	10/1993	
WO	WO 1994007510	A1	4/1994	
WO	WO 1994011503	A2	5/1994	
WO	WO 1995026750	A1	10/1995	
WO	WO 1995034326	A1	12/1995	
WO	WO 2000048635	A1	8/2000	
WO	WO 2003080108	A1	10/2003	
WO	WO 2004101740	A2	11/2004	
WO	WO 2005058283	A2	6/2005	
WO	WO 2006074199	A1	7/2006	
WO	WO-2007092772	A2 *	8/2007	 A61K 39/395
WO	WO-2008077616	A1 *	7/2008	 C07K 14/745
WO	WO 2008155134	A1	12/2008	
WO	WO 2010115866	A1	10/2010	
WO	WO-2011069164	A2 *	6/2011	 A61K 38/37
WO	WO 2013009627	A2	1/2013	
WO	WO 2013057219	A1	4/2013	
WO	WO 2014026954	A1	2/2014	
WO	WO 2014/070953	A1	5/2014	
WO	WO 2014144795	A1	9/2014	

OTHER PUBLICATIONS

U.S. Appl. No. 16/993,979* filed Aug. 14, 2020, Kevin Maloney.
 Extended European Search Report for European Patent Application No. 22176875.7, mailed Dec. 20, 2022.
 Armour, et al., "Recombinant Human IgG Molecules Lacking Fc Gamma Receptor I Binding And Monocyte Triggering Activities", *European Journal of Immunology*, vol. 29, No. 8, pp. 2613-2624. (Aug. 1999).
 Bai, et al., "Recombinant Granulocyte Colony-Stimulating Factor-Transferrin Fusion Protein As An Oral Myelopoietic Agent", *Proceedings of the National Academy of Sciences of the United States of America*, vol. 102, No. 20, pp. 7292-7296. (May 17, 2005).
 Bobrow, Robert S., "Excess Factor VIII: A Common Cause of Hypercoagulability", *American Board of Family Medicine, United States*, pp. 147-149 (Mar. 2005).
 Brandsma, et al., "Recombinant Human Transferrin: Beyond Iron Binding and Transport", *Biotechnology Advances*, vol. 29, No. 2, pp. 230-238. (Mar.-Apr. 2011).
 Brutlag, et al., "Improved Sensitivity of Biological Sequence Database Searches", *Computer Applications in the Biosciences: CABIOS*, vol. 6, No. 3, pp. 237-245. (Aug. 1990).

Burmeister, et al., "Crystal Structure of the Complex of Rat Neonatal Fc Receptor with Fc", *Nature*, vol. 372, No. 6504, pp. 379-383. (Nov. 24, 1994).
 Cameron, et al., "The Canine Factor VIII cDNA and 5' Flanking Sequence", *Journal of Thrombosis and Haemostasis*, vol. 79, No. 2, pp. 317-322. (Feb. 1998).
 Collins, et al., "Factor VIII Requirement to Maintain a Target Plasma Level in Prophylactic Treatment of Severe Hemophilia A: Influences Variance in Pharmacokinetics and Treatment Regimens", *Journal of Thrombosis and Haemostasis*, vol. 8, pp. 269-275. (Feb. 2010).
 Cutler, et al., "The Identification and Classification Of 41 Novel Mutations In The Factor VIII Gene (F8c)", *Human Mutation*, vol. 19, No. 3, pp. 274-278. (Mar. 2002).
 Dennis, et al., "Albumin Binding as a General Strategy for Improving the Pharmacokinetics of Proteins", *Journal of Biological Chemistry*, vol. 277, No. 38, pp. 35035-35043. (Sep. 20, 2002).
 Dobeli, et al., "Role of the Carboxy-Terminal Sequence on the Biological Activity of Human Immune Interferon (IFN-γ)", *Journal of Biotechnology*, vol. 7, No. 3, pp. 199-216. (1988).
 Eaton, et al., "Construction and Characterization of an Active Factor VIII Variant Lacking the Central One-Third of the Molecule", *Biochemistry*, vol. 25, No. 26, pp. 8343-8347. (Dec. 1986).
 Extended European Search Report received for European Application No. 19212900.5, mailed on Apr. 30, 2020.
 Fatouros, et al., "Recombinant factor VIII SQ-Influence of Oxygen, Metal Ions, pH and Ionic Strength on its Stability in Aqueous Solution", *International journal of Pharmaceutics*, vol. 155, No. 1, pp. 121-131. (1997).
 Francis, G E., "Protein Modification and Fusion Proteins", *Focus on Growth Factors*, vol. 3, No. 2, Mediscript, England, pp. 4-10. (1992).
 Friend, et al., "Phase I Study of an Engineered Aglycosylated Humanized Cd3 Antibody in Renal Transplant Rejection", *Transplantation*, vol. 68, Issue 11, pp. 1632-1637. (Dec. 15, 1999).
 Gayle, et al., "Identification Of Regions In Interleukin-1 Alpha Important For Activity", *Journal of Biological Chemistry*, vol. 268, No. 29, pp. 22105-22111. (Oct. 15, 1993).
 Genbank, "*Homo sapiens* Transferrin (TF)", Accession No. NM001063.1, 5 pages. (Sep. 24, 2014).
 Genbank, "*Homo sapiens* Transferrin (TF), mRNA", Accession No. XM002793, Retrieved from: <<https://www.ncbi.nlm.nih.gov/nucleotide/XM_002793.7?report=genbank>>, Retrieved on Sep. 24, 2014, 2 Pages. (May 13, 2002).
 Genbank, "*Homo sapiens* Transferrin (TF), mRNA", Accession No. XM039845, Retrieved From: <<https://www.ncbi.nlm.nih.gov/nucleotide/XM_039845.1?report=genbank>> Retrieved on Sep. 24, 2014, 2 Pages. (Jul. 16, 2001).
 Genbank, "*Homo sapiens* Transferrin (TF), mRNA", Accession No. XM039847, Retrieved From: <<https://www.ncbi.nlm.nih.gov/nucleotide/XM_039847.1?report=genbank>>, Retrieved on Sep. 24, 2014, 2 Pages. (Jul. 16, 2001).
 Genbank, "Human Transferrin mRNA, Complete cds", Accession No. M12530, Retrieved From: <<<http://www.ncbi.nlm.nih.gov/nucleotide/M12530>>> Retrieved on Jan. 15, 2015, 2 Pages. (Jan. 14, 1995).
 Genbank, "Transferrin [Human, Liver, mRNA, 2347 nt]", Accession No. S95936, Retrieved From: <<<http://www.ncbi.nlm.nih.gov/nucleotide/S95936>>>, 2 pages. (May 7, 1993).
 Gitschier, et al., "Characterization Of The Human Factor VIII Gene", *Nature*, vol. 312, No. 5992, pp. 326-330. (Nov. 22-28, 1984).
 Healey, et al., "The cDNA And Derived Amino Acid Sequence Of Porcine Factor VIII", *Blood*, vol. 88, No. 11, pp. 4209-4214. (Dec. 1, 1996).
 Hoeben, et al., "Expression Of Functional Factor VIII In Primary Human Skin Fibroblasts After Retrovirus-Mediated Gene Transfer", *Journal of Biological Chemistry*, vol. 265, No. 13, pp. 7318-7323. (May 1990).
 Holt, et al., "Anti-Serum Albumin Domain Antibodies For Extending The Half-Lives Of Short Lived Drugs", *Protein Engineering, Design and Selection*, vol. 21, No. 5, pp. 283-288. (May 2008).

(56)

References Cited**OTHER PUBLICATIONS**

International Search Report and Written Opinion received for PCT Patent Application No. PCT/US2014/29354, mailed on Aug. 7, 2014.

Kim, et al., "Transferrin Fusion Technology: A Novel Approach To Prolonging Biological Half-Life Of Insulinotropic Peptides", *Journal of Pharmacology and Experimental Therapeutics*, vol. 334, No. 3, pp. 682-692. (Sep. 2010).

Kraulis, et al., "The Serum Albumin-Binding Domain Of Streptococcal Protein G Is A Three-Helical Bundle: A Heteronuclear NMR Study", *FEBS Letters*, vol. 378, Issue 2, pp. 190-194. (Jan. 8, 1996).

Langner, et al., "Synthesis Of Biologically Active Deletion Mutants Of Human Factor VIII:C", *Behring Institute Mitteilungen*, No. 82, pp. 16-25. (Apr. 1988).

Li, et al., "The Role Of The Transferrin-Transferrin-Receptor System In Drug Delivery And Targeting", *Trends in Pharmacological Sciences*, vol. 23, No. 5, pp. 206-209. (May 2002).

Linhardt, et al., "Mutational Analysis Of The Interaction Between Albumin-Binding Domain From Streptococcal Protein G And Human Serum Albumin", *Protein Science*, vol. 11, No. 2, pp. 206-213. (Feb. 2002).

Lollar, et al., "Coagulant Properties Of Hybrid Human/Porcine Factor VIII Molecules", *Journal of Biological Chemistry*, vol. 267, pp. 23652-23657. (Nov. 25, 1992).

Lollar, et al., "Structural Basis For The Deceased Procoagulant Activity Of Human Factor VIII Compared To The Porcine Homolog", *Journal of Biological Chemistry*, vol. 266, No., pp. 12481-12486. (Jul. 5, 1991).

Mahlangu, et al., "Phase 3 Study of Recombinant Factor VIII Fc Fusion Protein in Severe Hemophilia A", *Blood*, vol. 123, Issue 3, pp. 317-325. (Jan. 2014).

Malik, et al., "Polyethylene Glycol (PEG)-Modified Granulocyte-Macrophage Colony-Stimulating Factor (GM-CSF) With Conserved Biological Activity", *Experimental Hematology*, vol. 20, No. 8, pp. 1028-1035. (Sep. 1992).

Mannucci, et al., "The Hemophilias—From Royal Genes to Gene Therapy", *New England Journal of Medicine*, vol. 344, No. 23, pp. 1773-1779. (Jun. 1, 2001).

Mei, et al., "Expression of Human Coagulation Factor VIII in a Human Hybrid Cell Line, HKB11", *Molecular Biotechnology*, vol. 34, No. 2, Humana Press Inc., pp. 165-178. (Oct. 2006).

Mei, et al., "Rational Design Of A Fully Active, Long-Acting PEGylated Factor VIII For Hemophilia A Treatment", *Blood*, vol. 116, No. 2, pp. 270-279. (Jul. 15, 2010).

Meulien, et al., "A New Recombinant Procoagulant Protein Derived From The Cdna Encoding Human Factor VII", *Protein Engineering, Design and Selection*, vol. 2, No. 4, pp. 301-306. (Oct. 1, 1988).

Muller, et al., "Recombinant Bispecific Antibodies For Cellular Cancer Immunotherapy", *Current opinion in molecular therapeutics*, vol. 9, No. 4, pp. 319-326. (Aug. 2007).

Neumann, et al., "Gene Transfer Into Mouse Lyoma Cells By Electroporation In High Electric Fields", *The EMBO Journal*, vol. 1, No. 7, pp. 841-845. (Jul. 1, 1982).

Oganesyan, et al., "Structural Characterization of a Human Fc Fragment Engineered for Extended Serum Half-Life", *Molecular Immunology*, vol. 46, No. 8-9, pp. 1750-1755. (May 2009).

Osterberg, et al., "Development of a Freeze-Dried Albumin-Free Formulation of Recombinant Factor VIII SQ", *Pharmaceutical Research*, vol. 14, Issue 7, pp. 892-898. (Jul. 1997).

Peyvandi, et al., "Genetic Diagnosis of Haemophilia and Other Inherited Bleeding Disorders", *Haemophilia*, vol. 12, Suppl 3, pp. 82-89. (Jul. 2006).

Powell, et al., "Safety And Prolonged Activity Of Recombinant Factor VIII Fc fusion protein in Hemophilia A Patients", *Blood*, vol. 119, No. 13, pp. 3031-3037. (Mar. 29, 2012).

Rodriguez-Merchan, Carlos E., "Management Of Musculoskeletal Complications Of Hemophilia", *Seminars in Thrombosis and Hemostasis*, vol. 29, No. 01, pp. 87-96. (2003).

Ron, et al., "Expression of Biologically Active Recombinant Keratinocyte Growth Factor. Structure/Function Analysis Of Amino-Terminal Truncation Mutants", *Journal of Biological Chemistry*, vol. 268, No. 4, pp. 2984-2988. (1993).

Roovers, et al., "Efficient Inhibition Of EGFR Signaling And Of Tumour Growth By Antagonistic anti-EGFR Nanobodies", *Cancer Immunology, Immunotherapy*, vol. 56, No. 3, pp. 303-317. (Mar. 2007).

Routledge, et al., "The Effect Of Aglycosylation On The Immunogenicity Of A Humanized Therapeutic CD3 Monoclonal Antibody", *Transplantation*, vol. 60, No. 8, pp. 847-853. (Oct. 1, 1995).

Sarver, et al., "Stable Expression Of Recombinant Factor VIII Molecules Using A Bovine Papillomavirus Vector", *DNA*, vol. 6, No. 6, pp. 553-564. (Dec. 1987).

Schulte, Stefan, "Half-Life Extension Through Albumin Fusion Technologies", *Thrombosis Research*, vol. 124, Supplement 2, pp. S6-S8. (Dec. 2009).

Shields, et al., "High Resolution Mapping of the Binding Site on Human IgG1 for Fc Gamma RI, Fc Gamma RII, Fc Gamma RIII, and FcRn and Design of IgG1 Variants with Improved Binding to the Fc Gamma R", *Journal of Biological Chemistry*, vol. 276, No. 9, pp. 6591-6604. (Mar. 2, 2001).

Sommermeier, et al., "Klinisch verwendete Hydroxyethylstärke: physikalisch-chemische Charakterisierung", *Krankenhauspharmazie*, vol. 8, No. 8, Deutscher Apotheker Verlag, Birkenwaldstr, Germany, pp. 271-278. (1987).

Story, et al., "A Major Histocompatibility Complex Class I-like Fc Receptor Cloned from Human Placenta: Possible Role in Transfer of Immunoglobulin G from Mother to Fetus", *Journal of Experimental Medicine*, vol. 180, No. 6, pp. 2377-2381. (Dec. 1, 1994).

Toole, et al., "A Large Region (Approximately Equal To 95 kDa) Of Human Factor VIII Is Dispensable For In Vitro Procoagulant Activity", *Proceedings of the National Academy of Sciences*, vol. 83, No. 16, pp. 5939-5942. (Aug. 1986).

Toole, et al., "Molecular Cloning Of a cDNA Encoding Human Antihemophilic Factor", *Nature*, vol. 312, No. 5992, pp. 342-347. (Nov. 22-28, 1984).

Trussel, et al., "New Strategy For The Extension Of The Serum Half-Life Of Antibody Fragments", *Bioconjugate Chemistry*, vol. 20, No. 12, pp. 2286-2292. (Dec. 2009).

Vaccaro, et al., "Engineering the Fc Region of Immunoglobulin G to Modulate in Vivo Antibody Levels", *Nature Biotechnology*, vol. 23, No. 10, pp. 1283-1288. (Oct. 2005).

Valentino, et al., "A Randomized Comparison of Two Prophylaxis Regimens and a Paired Comparison of on-Demand and Prophylaxis Treatments in Hemophilia A Management", *Journal of Thrombosis and Haemostasis*, vol. 10, No. 3, pp. 359-367. (Mar. 2012).

Vehar, et al., "Structure Of Human Factor VIII", *Nature*, vol. 312, No. 5992, pp. 337-342. (Nov. 1984).

Wakabayashi, et al., "Residues 110-126 in the A1 Domain of Factor VIII Contain a Ca²⁺ Binding Site Required for Cofactor Activity", *The Journal of Biological Chemistry*, vol. 279, pp. 12677-12684. (Jan. 13, 2004).

Wang, et al., "Receptor-Mediated Activation of a Proinsulin-Transferrin Fusion Protein in Hepatoma Cells", *Journal of Controlled Release*, vol. 155, No. 3, pp. 386-392. (Nov. 7, 2011).


Ward, et al., "The Effector Functions Of Immunoglobulins: Implications For Therapy", *Therapeutic immunology*, vol. 2, No. 2, pp. 77-94. (Apr. 1995).

Weidler, et al., "Pharmacokinetic Parameters As Criteria For Clinical Use Of Hydroxyethyl Starch Preparations", *Arzneimittelforschung/Drug Research*, vol. 41, No. 5, pp. 494-498. (May 1991).

Wigler, et al., "Biochemical Transfer of Single-Copy Eucaryotic Genes using Total Cellular DNA as Donor", *Cell*, vol. 14, No. 3, pp. 725-731. (Jul. 1978).

Wood, et al., "Expression of Active Human Factor VIII From Recombinant DNA Clones", *Nature*, vol. 312, No. 5992, pp. 330-337. (Nov. 22-28, 1984).

* cited by examiner



- 1. Minimum interval of 96 hours, maximum interval of 8 weeks from rFVIII injection to first injection of 50 IU/kg rFVIII Fc for sequential PK profiling.
- 2. Starting regimen for twice weekly individualized (tailored) prophylaxis of 25 IU/kg on first day of week followed by 50 IU/kg on fourth day, with dose and interval adjusted to achieve target trough level of approximately 1% to 3% (or higher, as clinically indicated). FVIII level assessed at peak and trough.
- 3. Repeat doses of 50 IU/kg rFVIII Fc for sequential PK profiling administered at Week 14, or at 12 to 24 weeks after first injection of rFVIII Fc.
- 4. Continued individualized prophylaxis regimen until 28 ($\pm$ 2) weeks up to 52 ($\pm$ 2) weeks.

PK, pharmacokinetic

FIG. 1




FIG. 2

1

FACTOR VIII POLYPEPTIDE FORMULATIONS

REFERENCE TO RELATED APPLICATIONS

The present application is a continuation of U.S. patent application Ser. No. 15/455,043, filed Mar. 9, 2017, which is a division of U.S. patent application Ser. No. 14/213,180, filed Mar. 14, 2014, now U.S. Pat. No. 9,623,088, which claims the benefit of U.S. Provisional Patent Application Ser. Nos. 61/800,293, filed Mar. 15, 2013, 61/817,085, filed Apr. 29, 2013, 61/829,884, filed May 31, 2013, 61/839,477, filed Jun. 26, 2013, 61/863,860, filed Aug. 8, 2013, 61/876,927, filed Sep. 12, 2013, 61/879,955, filed Sep. 19, 2013, and 61/897,742, filed Oct. 30, 2013, all of which are incorporated herein by reference in their entireties.

REFERENCE TO SEQUENCE LISTING SUBMITTED ELECTRONICALLY

The content of the electronically submitted sequence listing in ASCII text file (Name: 709032_SA9-436USDIVCON_ST25.txt; Size: 61,753 bytes; and Date of Creation: Aug. 14, 2020) is incorporated herein by reference in its entirety.

BACKGROUND OF THE INVENTION

Field of the Invention

The present invention relates generally to the field of therapeutics for hemostatic disorders.

Background Art

Hemophilia A is an X-linked bleeding disorder caused by mutations and/or deletions in the factor VIII (FVIII) gene resulting in a deficiency of FVIII activity (Peyvandi, F. et al. *Haemophilia* 12:82-89 (2006)). The disease is characterized by spontaneous hemorrhage and excessive bleeding after trauma. Over time, the repeated bleeding into muscles and joints, which often begins in early childhood, results in hemophilic arthropathy and irreversible joint damage. This damage is progressive and can lead to severely limited mobility of joints, muscle atrophy and chronic pain (Rodriguez-Merchan, E. C., *Semin. Thromb. Hemost.* 29:87-96 (2003), which is herein incorporated by reference in its entirety).

The A2 domain is necessary for the procoagulant activity of the factor VIII molecule. Studies show that porcine factor VIII has six-fold greater procoagulant activity than human factor VIII (Lollar, P., and E. T. Parker, *J. Biol. Chem.* 266:12481-12486 (1991)), and that the difference in coagulant activity between human and porcine factor VIII appears to be based on a difference in amino acid sequence between one or more residues in the human and porcine A2 domains (Lollar, P., et al., *J. Biol. Chem.* 266:23652-23657 (1992)), incorporated herein by reference in its entirety.

Treatment of hemophilia A is by replacement therapy targeting restoration of FVIII activity to 1 to 5% of normal levels to prevent spontaneous bleeding (Mannucci, P. M., et al., *N. Engl. J. Med.* 344:1773-1779 (2001), which is herein incorporated by reference in its entirety). There are plasma-derived and recombinant FVIII products available to treat bleeding episodes on-demand or to prevent bleeding episodes from occurring by treating prophylactically.

2

Reduced mortality, prevention of joint damage, and improved quality of life have been important achievements due to the development of plasma-derived and recombinant FVIII. Prolonged protection from bleeding would represent another key advancement in the treatment of hemophilia A patients.

BRIEF SUMMARY OF THE INVENTION

The present invention comprises a pharmaceutical composition comprising: (a) a FVIII polypeptide; (b) one or more stabilizing agents selected from sucrose, trehalose, raffinose, arginine, or mixture thereof; (c) sodium chloride (NaCl); (d) L-histidine; (e) calcium chloride; and (f) polysorbate 20 or polysorbate 80. In one embodiment, mannitol, glycine, alanine, or hydroxyethyl starch is not included in the pharmaceutical composition. In another embodiment, NaCl is the only bulking agent. In other embodiments, the FVIII polypeptide is a long-acting polypeptide or a short-acting polypeptide.

In other embodiments, the pharmaceutical composition comprises about 1% (w/v) to about 2.5% (w/v) sucrose, about 1.3% (w/v) sucrose to about 2.0% (w/v) sucrose, about 1.33% (w/v) sucrose or about 2.0% (w/v) sucrose, about 10 mg/ml to about 25 mg/ml sucrose, about 13 mg/ml to about 20 mg/ml sucrose, about 13.3 mg/ml sucrose, or about 20.0 mg/ml sucrose. In still other embodiments, the pharmaceutical composition comprises about 150 mM to about 250 mM NaCl, about 175 mM to about 225 mM NaCl, about 200 mM to about 210 mM NaCl, about 205 mM NaCl, about 8.8 mg/ml to about 14.6 mg/ml NaCl, about 10 mg/ml to about 13 mg/ml NaCl, about 12.0 mg/ml NaCl, about 250 mM to about 350 mM NaCl, about 275 mM to about 325 mM NaCl, about 308 mM NaCl, about 14.6 mg/ml to about 20.5 mg/ml NaCl, about 16 mg/ml to about 19 mg/ml NaCl, or about 18.0 mg/ml NaCl.

In yet other embodiments, the pharmaceutical composition comprises about 5 mM to about 15 mM L-histidine, about 6.64 mM L-histidine to about 9.8 mM L-histidine, about 0.75 mg/ml to about 2.25 mg/ml L-histidine, or about 1.03 mg/ml L-histidine to about 1.55 mg/ml L-histidine. In certain embodiments, the pharmaceutical composition comprises about 5 mM to about 10 mM calcium chloride, about 5.4 mM calcium chloride to about 8 mM calcium chloride, about 0.75 mg/ml to about 1.5 mg/ml calcium chloride dihydrate, or about 0.8 mg/ml calcium chloride dihydrate to about 1.18 mg/ml calcium chloride dihydrate.

In some embodiments, the pharmaceutical composition comprises about 0.008% (w/v) to about 0.025% (w/v) polysorbate 20 or polysorbate 80, about 0.013% (w/v) polysorbate 20 or polysorbate 80 or about 0.02% (w/v) polysorbate 20 or polysorbate 80, about 0.08 mg/ml to about 0.25 mg/ml polysorbate 20 or polysorbate 80, about 0.13% mg/ml polysorbate 20 or polysorbate 80 or about 0.20 mg/ml polysorbate 20 or polysorbate 80.

In other embodiments, the pharmaceutical composition comprises the rFVIII_{FC} polypeptide, which comprises a first subunit comprising an amino acid sequence at least 90% or 95% identical to amino acids 20 to 1684 of SEQ ID NO:2 or 20 to 2578 of SEQ ID NO:6 and a second subunit comprising an amino acid sequence at least 90% to 95% identical to amino acids 21 to 247 of SEQ ID NO:4. In certain embodiments, wherein the rFVIII_{FC} polypeptide comprises a first subunit comprising amino acids 20 to 1684 of SEQ ID NO:2 or 20 to 2578 of SEQ ID NO:6, and a second subunit comprising amino acids 21 to 247 of SEQ ID NO:4.

3

In certain embodiments, a FVIII polypeptide is present in a pharmaceutical composition at a concentration of about 50 IU/ml to about 2500 IU/ml, e.g., 83 IU/ml, 167 IU/ml, 250 IU/ml, 333 IU/ml, 500 IU/ml, 667 IU/ml, 1000 IU/ml, 1333 IU/ml, 1667 IU/ml, or 2000 IU/ml of the FVIII polypeptide. In some embodiments, a FVIII polypeptide is present in a pharmaceutical composition at a concentration of about 100 IU/ml to about 4000 IU/ml, e.g., 150 IU/ml, 287.5 IU/ml, 431.25 IU/ml, 575 IU/ml, 862.5 IU/ml, 1150 IU/ml, 1725 IU/ml, 2300 IU/ml, 2875 IU/ml, or 3450 IU/ml of the FVIII polypeptide.

In other embodiments, a pharmaceutical composition comprises: (a) about 50 IU/ml to about 2500 IU/ml of a long-acting FVIII polypeptide; (b) about 1% (w/v) to about 2.5% (w/v) of sucrose; (c) about 150 mM to about 250 mM NaCl; (d) about 5 mM to about 15 mM L-histidine; (e) about 5 mM to about 10 mM calcium chloride; and (f) about 0.008% (w/v) to about 0.025% of polysorbate 20 or polysorbate 80. In other embodiments, the pharmaceutical composition comprises about 175 mM to about 225 mM NaCl or about 200 mM to about 210 mM NaCl.

In yet other embodiments, a pharmaceutical composition comprises (a) about 83 IU/ml, about 167 IU/ml, about 250 IU/ml, about 333 IU/ml, about 500 IU/ml, about 667 IU/ml, about 1000 IU/ml, about 1333 IU/ml, about 1667 IU/ml, or about 2000 IU/ml of a FVIII polypeptide; (b) about 1.33% (w/v) of sucrose; (c) about 205 mM NaCl; (d) about 6.64 mM L-histidine; (e) about 5.4 mM calcium chloride; and (f) about 0.013% (w/v) of polysorbate 20 or polysorbate 80.

In some embodiments, a pharmaceutical composition comprises: (a) about 100 IU/ml to about 4000 IU/ml of a FVIII polypeptide; (b) about 1% (w/v) to about 2.5% (w/v) of sucrose; (c) about 250 mM to about 350 mM NaCl; (d) about 5 mM to about 15 mM L-histidine; (e) about 5 mM to about 10 mM calcium chloride; and (f) about 0.008% (w/v) to about 0.025% of polysorbate 20 or polysorbate 80. In other embodiments, the pharmaceutical composition comprises about 275 mM to about 325 mM NaCl.

In certain embodiments, a pharmaceutical composition comprises: (a) about 150 IU/ml, about 287.5 IU/ml, about 431.25 IU/ml, about 575 IU/ml, about 862.5 IU/ml, about 1150 IU/ml, about 1725 IU/ml, about 2300 IU/ml, about 2875 IU/ml, or about 3450 IU/ml of a FVIII polypeptide; (b) about 2.0% (w/v) of sucrose; (c) about 308 mM NaCl; (d) about 9.8 mM L-histidine; (e) about 8 mM calcium chloride; and (f) about 0.020% (w/v) of polysorbate 20 or polysorbate 80. In other embodiments, a pharmaceutical composition comprises: (a) about 50 IU/ml to about 2500 IU/ml of a FVIII polypeptide; (b) about 10 mg/ml to about 25 mg/ml of sucrose; (c) about 8.8 mg/ml to about 14.6 mg/ml NaCl; (d) about 0.75 mg/ml to about 2.25 mg/ml L-histidine; (e) about 0.75 mg/ml to about 1.5 mg/ml calcium chloride dihydrate; and (f) about 0.08 mg/ml to about 0.25 mg/ml of polysorbate 20 or polysorbate 80. In some embodiments, a pharmaceutical composition comprises: (a) about 83 IU/ml, about 167 IU/ml, about 250 IU/ml, about 333 IU/ml, about 500 IU/ml, about 667 IU/ml, about 1000 IU/ml, about 1333 IU/ml, about 1667 IU/ml, or about 2000 IU/ml of a FVIII polypeptide; (b) about 13.3 mg/ml of sucrose; (c) about 12.0 mg/ml NaCl; (d) about 1.03 mg/ml L-histidine; (e) about 0.8 mg/ml calcium chloride dihydrate; and (f) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.

In other embodiments, a pharmaceutical composition comprises: (a) about 100 IU/ml to about 4000 IU/ml of a FVIII polypeptide; (b) about 10 mg/ml to about 25 mg/ml of sucrose; (c) about 14.6 mg/ml to about 20.5 mg/ml NaCl; (d) about 0.75 mg/ml to about 2.25 mg/ml L-histidine; (e) about

4

0.75 mg/ml to about 1.5 mg/ml calcium chloride dihydrate; and (f) about 0.08 mg/ml to about 0.25 mg/ml of polysorbate 20 or polysorbate 80. In certain embodiments, a pharmaceutical composition comprises: (a) about 150 IU/ml, about 287.5 IU/ml, about 431.25 IU/ml, about 575 IU/ml, about 862.5 IU/ml, about 1150 IU/ml, about 1725 IU/ml, about 2300 IU/ml, about 2875 IU/ml, or about 3450 IU/ml of a FVIII polypeptide; (b) about 20.0 mg/ml of sucrose; (c) about 18.0 mg/ml NaCl; (d) about 1.55 mg/ml L-histidine; (e) about 1.18 mg/ml calcium chloride dihydrate; and (f) about 0.20 mg/ml of polysorbate 20 or polysorbate 80.

In some embodiments, the present invention includes a pharmaceutical kit comprising: (a) a first container comprising a lyophilized powder, wherein the powder comprises (i) a FVIII polypeptide, (ii) one or more stabilizing agents selected from sucrose, trehalose, raffinose, arginine, or mixture thereof; (iii) sodium chloride (NaCl); (iv) L-histidine; (v) calcium chloride; and (vi) polysorbate 20 or polysorbate 80; and (b) a second container comprising sterilized water for injections to be combined with the lyophilized powder of the first container. In other embodiments, a pharmaceutical kit comprises: (a) a first container comprising a lyophilized powder, where the powder comprises (i) about 250 IU, about 500 IU, about 750 IU, about 1000 IU, about 1500 IU, about 2000 IU, about 3000 IU, about 4000 IU, about 5000 IU, or about 6000 IU of a FVIII polypeptide, (ii) about 40 mg of sucrose; (iii) about 36 mg of sodium chloride; (iv) about 3.1 mg of L-histidine; (v) about 2.40 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and (b) a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 83 IU/ml, about 167 IU/ml, about 250 IU/ml, about 333 IU/ml, about 500 IU/ml, about 667 IU/ml, about 1000 IU/ml, about 1333 IU/ml, about 1667 IU/ml, or about 2000 IU/ml of a FVIII polypeptide, respectively; (ii) about 1.33% (w/v) of sucrose; (iii) about 205 mM NaCl; (iv) about 6.64 mM L-histidine; (v) about 5.4 mM of calcium chloride; and (vi) about 0.013% (w/v) of polysorbate 20 or polysorbate 80.

In other embodiments, a pharmaceutical kit comprises: (a) a first container comprising a lyophilized powder, where the powder comprises (i) about 250 IU, about 500 IU, about 750 IU, about 1000 IU, about 1500 IU, about 2000 IU, about 3000 IU, about 4000 IU, about 5000 IU, or about 6000 IU of a FVIII polypeptide, (ii) about 40 mg of sucrose; (iii) about 36 mg of sodium chloride; (iv) about 3.1 mg of L-histidine; (v) about 2.40 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and (b) a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 83 IU/ml, about 167 IU/ml, about 250 IU/ml, about 333 IU/ml, about 500 IU/ml, about 667 IU/ml, about 1000 IU/ml, about 1333 IU/ml, about 1667 IU/ml, or about 2000 IU/ml of a FVIII polypeptide, respectively; (ii) about 13.3 mg/ml of sucrose; (iii) about 12.0 mg/ml of NaCl; (iv) about 1.03 mg/ml of L-histidine; (v) about 0.80 mg/ml of calcium chloride; and (vi) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.

The invention also includes a method of reducing or decreasing a bleeding episode or treating or preventing a bleeding condition comprising administering a pharmaceutical composition comprising a FVIII polypeptide.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows details on the design of the sequential PK subgroup (Arm 1) dosing and PK sampling.

FIG. 2 shows the mean observed FVIII activity with rFVIII-Fc and rFVIII over time: one stage clotting assay (logarithmic scale) in the sequential pharmacokinetic subgroup. rFVIII-Fc demonstrated an approximate 50% longer elimination half-life and mean residence time compared with wild-type Factor VIII (a short-acting FVIII polypeptide) ($P < 0.001$).

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a formulation for a Factor VIII polypeptide, e.g., FVIII-Fc, and methods of using thereof.

1. Definitions

It must be noted that, as used in this specification and the appended claims, the singular forms “a”, “an” and “the” include plural referents unless the context clearly dictates otherwise. The terms “a” (or “an”), as well as the terms “one or more,” and “at least one” can be used interchangeably herein.

The term “about” is used herein to mean approximately, roughly, around, or in the regions of. When the term “about” is used in conjunction with a numerical range, it modifies that range by extending the boundaries above and below the numerical values set forth. In general, the term “about” is used herein to modify a numerical value above and below the stated value by a variance of 10 percent, up or down (higher or lower).

Furthermore, “and/or” where used herein is to be taken as specific disclosure of each of the two specified features or components with or without the other. Thus, the term “and/or” as used in a phrase such as “A and/or B” herein is intended to include “A and B,” “A or B,” “A” (alone), and “B” (alone). Likewise, the term “and/or” as used in a phrase such as “A, B, and/or C” is intended to encompass each of the following embodiments: A, B, and C; A, B, or C; A or C; A or B; B or C; A and C; A and B; B and C; A (alone); B (alone); and C (alone).

It is understood that wherever embodiments are described herein with the language “comprising,” otherwise analogous embodiments described in terms of “consisting of” and/or “consisting essentially of” are also provided.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure is related. For example, the Concise Dictionary of Biomedicine and Molecular Biology, Juo, Pei-Show, 2nd ed., 2002, CRC Press; The Dictionary of Cell and Molecular Biology, 3rd ed., 1999, Academic Press; and the Oxford Dictionary Of Biochemistry And Molecular Biology, Revised, 2000, Oxford University Press, provide one of skill with a general dictionary of many of the terms used in this disclosure.

Units, prefixes, and symbols are denoted in their Système International de Unites (SI) accepted form. Numeric ranges are inclusive of the numbers defining the range. Unless otherwise indicated, amino acid sequences are written left to right in amino to carboxy orientation. The headings provided herein are not limitations of the various aspects or embodiments of the disclosure, which can be had by reference to the specification as a whole. Accordingly, the terms defined immediately below are more fully defined by reference to the specification in its entirety. Amino acids are referred to herein by either their commonly known three letter symbols

or by the one-letter symbols recommended by the IUPAC-IUB Biochemical Nomenclature Commission. Nucleotides, likewise, are referred to by their commonly accepted single-letter codes.

The term “polypeptide,” “peptide” and “protein” are used interchangeably and refer to a polymeric compound comprised of covalently linked amino acid residues.

The term “polynucleotide” and “nucleic acid” are used interchangeably and refer to a polymeric compound comprised of covalently linked nucleotide residues. Polynucleotides can be DNA, cDNA, RNA, single stranded, or double stranded, vectors, plasmids, phage, or viruses. Polynucleotides include those in Table 10, which encode the polypeptides of Table 11 (see Table 10). Polynucleotides also include fragments of the polynucleotides of Table 10, e.g., those that encode fragments of the polypeptides of Table 11, such as the Factor VIII, Fc, signal sequence, propeptide, and other fragments of the polypeptides of Table 11.

“Administering,” as used herein, means to proscribe or give a pharmaceutical composition comprising a Factor VIII polypeptide of the invention to a subject via a pharmaceutically acceptable route. Examples of routes of administration include, but are not limited to, intravenous, e.g., intravenous injection and intravenous infusion, e.g., via central venous access. Additional routes of administration include subcutaneous, intramuscular, oral, nasal, and pulmonary administration. Advantages of the present invention include: improved regimen compliance; reduced break through bleeds; increased protection of joints from bleeds; prevention of joint damage; reduced morbidity; reduced mortality; prolonged protection from bleeding; decreased thrombotic events; and improved quality of life.

The terms “long-acting” and “long-lasting” are used interchangeably herein. In one embodiment, the term “long-acting” or “long-lasting” indicates that a FVIII activity as a result of administration of the “long-acting” FVIII polypeptide is longer than the FVIII activity of a wild-type FVIII (e.g., ADVATE® or plasma-derived FVIII (“pdFVIII”). The “longer” FVIII activity can be measured by any known methods in the art, e.g., aPTT assay, chromogenic assay, ROTEM®, TGA, etc. In one embodiment, the “longer” FVIII activity can be shown by the $T_{1/2 \text{ beta}}$ (activity). In another embodiment, the “longer” FVIII activity can be shown by the level of FVIII antigen present in plasma, e.g., by the $T_{1/2 \text{ beta}}$ (antigen). In other embodiments, the long-acting or long-lasting FVIII polypeptide works longer in a coagulation cascade, e.g., is active for a longer period, compared to a wild-type FVIII polypeptide, i.e., a polypeptide consisting of amino acids 20 to 1457 of SEQ ID NO: 2, i.e., SQ BDD FVIII (REFACTO®) or a polypeptide consisting of amino acids 20 to 2351 of SEQ ID NO: 6 (ADVATE®).

The terms “short-acting” and “short-lasting” are used interchangeably herein. The term “short-acting” or “short-lasting” indicates that a FVIII activity as a result of administration of the “short-acting” FVIII polypeptide is similar to or the same as the FVIII activity of a wild-type FVIII (e.g., ADVATE® (amino acids 20 to 2351 of SEQ ID NO: 6), REFACTO® (amino acids 20 to 1457 of SEQ ID NO: 2), or plasma-derived FVIII (“pdFVIII”)) or shorter than the FVIII activity of a long-acting FVIII polypeptide. The “shorter” FVIII activity can be measured by any known methods in the art, e.g., aPTT assay, chromogenic assay, ROTEM®, TGA, etc. Exemplary short-acting FVIII polypeptides include, but are not limited to, wild-type mature FVIII polypeptide (ADVATE®, RECOMBINATE®, or HELIXATE®) or B-domain deleted FVIII polypeptides such as SQ BDD

FVIII (REFACTO® and XYNTHA®) or FVIII polypeptides containing 21 amino acids from B-domain (i.e., SFSQNSRHPSQNPVLRHQR, SEQ ID NO: 17) (e.g., NOVOEIGHT®).

The term “chimeric polypeptide,” as used herein, means a polypeptide that includes within it at least two polypeptides (or portions thereof such as subsequences or peptides) from different sources. Chimeric polypeptides can include two, three, four, five, six, seven, or more polypeptides or portions thereof from different sources, such as different genes, different cDNAs, or different animal or other species. Chimeric polypeptides can include one or more linkers joining the different polypeptides or portions thereof. Thus, the polypeptides or portions thereof can be joined directly or they can be joined indirectly, via linkers, or both, within a single chimeric polypeptide. Chimeric polypeptides can include additional peptides such as signal sequences and sequences such as 6His and FLAG that aid in protein purification or detection. In addition, chimeric polypeptides can have amino acid or peptide additions to the N- and/or C-termini. In some embodiments, a chimeric polypeptide is a long-acting FVIII polypeptide. Exemplary chimeric polypeptides of the invention are Factor VIII-FcRn BP chimeric polypeptides, e.g., Factor VIII-Fc polypeptides such as the FVIII-Fc, SEQ ID NO:2 or 6 (Table 11) with or without its signal sequence and propeptide.

“Hybrid” polypeptides and proteins, as used herein, means a combination of a chimeric polypeptide with a second polypeptide. The chimeric polypeptide and the second polypeptide in a hybrid can be associated with each other via protein-protein interactions, such as charge-charge or hydrophobic interactions. The chimeric polypeptide and the second polypeptide in a hybrid can be associated with each other via disulfide or other covalent bond(s). Hybrids are described in WO 2004/101740 and WO 2006/074199, each of which is incorporated herein by reference in its entirety. See also U.S. Pat. Nos. 7,404,956 and 7,348,004, each of which is incorporated herein by reference in its entirety. The second polypeptide can be a second copy of the same chimeric polypeptide or it can be a non-identical chimeric polypeptide. See, e.g., Table 11. In one embodiment, the second polypeptide is a polypeptide comprising an FcRn binding partner. FcRn binding partners binds to FcRn and protects the FcRn binding partner containing molecule from catabolism, thus extending the plasma half-life. In another embodiment, the chimeric polypeptide is a chimeric Factor VIII-Fc polypeptide and the second polypeptide consists essentially of Fc, e.g., a rFVIII-Fc recombinant fusion protein consisting of a single molecule of recombinant B-domain deleted human FVIII (BDD-rFVIII) fused to the dimeric Fc domain of the human IgG1, with no intervening linker sequence. This hybrid polypeptide is referred to herein as FVIII-Fc monomeric Fc fusion protein, FVIII-Fc monomer hybrid, monomeric FVIII-Fc hybrid, and FVIII-Fc monomer-dimer. See Table 11A. The Examples provide preclinical and clinical data for this hybrid polypeptide.

The second polypeptide in a hybrid can comprise or consist essentially of a sequence at least 90% or 95% identical to the amino acid sequence shown in Table 11A(ii) without a signal sequence (amino acids 21 to 247 of SEQ ID NO:4) or at least 90% or 95% identical to the amino acid sequence shown in Table 11A(ii) with a signal sequence (amino acids 1 to 247 of SEQ ID NO:4). The second polypeptide can comprise or consist essentially of a sequence identical to the amino acid sequence shown in Table 11A(ii) without a signal sequence (amino acids 21 to 247 of SEQ ID NO:4) or identical to the amino acid

sequence shown in Table 11A(ii) with a signal sequence (amino acids 1 to 247 of SEQ ID NO:4).

Factor VIII coagulant activity is expressed as International Unit(s) (IU). One IU of Factor VIII activity corresponds approximately to the quantity of Factor VIII in one milliliter of normal human plasma. Several assays are available for measuring Factor VIII activity, including the one stage clotting assay (activated partial thromboplastin time; aPTT), thrombin generation time (TGA) and rotational thromboelastometry (ROTEM®).

The term “lyophilisate” as used herein in connection with the formulation according to the invention denotes a formulation which is manufactured by freeze-drying methods known in the art per se. The solvent (e.g. water) is removed by freezing following sublimation under vacuum and desorption of residual water at elevated temperature. In the pharmaceutical field, the lyophilisate has usually a residual moisture of about 0.1 to 5% (w/w) and is present as a powder or a physical stable cake. The lyophilisate is characterized by a fast dissolution after addition of a reconstitution medium.

The term “reconstituted formulation” as used herein denotes a formulation which is lyophilized and re-dissolved by addition of a diluent. The diluent can contain, without limitation, water for injection (WFI), bacteriostatic water for injection (BWFI), sodium chloride solutions (e.g. 0.9% (w/v) NaCl), glucose solutions (e.g. 5% glucose), surfactant containing solutions (e.g. 0.01% polysorbate 20 or polysorbate 80), a pH-buffered solution (e.g. phosphate-buffered solutions) and combinations thereof.

“Dosing interval,” as used herein, means the amount of time that elapses between multiple doses being administered to a subject. Dosing interval can thus be indicated as ranges. The dosing interval in the methods of the invention using a chimeric FVIII-FcRn BP, e.g., a chimeric FVIII-Fc, can be at least about one and one-half to eight times longer than the dosing interval required for an equivalent amount (in IU/kg) of the Factor VIII without the FcRn BP, e.g., Fc portion (i.e., a polypeptide consisting of said FVIII). The dosing interval when administering, e.g., a Factor VIII-Fc chimeric polypeptide (or a hybrid) of the invention can be at least about one and one-half times longer than the dosing interval required for an equivalent amount of the Factor VIII without the FcRn BP, e.g., Fc portion (i.e., a polypeptide consisting of the Factor VIII). The dosing interval can be at least about one and one-half to eight times longer than the dosing interval required for an equivalent amount of the Factor VIII without, e.g., the Fc portion (or a polypeptide consisting of the Factor VIII).

The term “dosing frequency” as used herein refers to the frequency of administering doses of a FVIII polypeptide in a given time. Dosing frequency can be indicated as the number of doses per a given time, e.g., once a week or once in two weeks.

The term “prophylaxis of one or more bleeding episode” or “prophylactic treatment” as used herein means administering a FVIII polypeptide in multiple doses to a subject over a course of time to increase the level of FVIII activity in a subject’s plasma. In one embodiment, “prophylaxis of one or more bleeding episode” indicates use of a FVIII polypeptide to prevent or inhibit occurrence of one or more spontaneous or uncontrollable bleeding or bleeding episodes or to reduce the frequency of one or more spontaneous or uncontrollable bleeding or bleeding episodes. In another embodiment, the increased FVIII activity level is sufficient to decrease the incidence of spontaneous bleeding or to prevent bleeding in the event of an unforeseen injury.

Prophylactic treatment decreases or prevents bleeding episodes, for example, those described under on-demand treatment. Prophylactic treatment can be individualized, as discussed under “dosing interval”, e.g., to compensate for inter-patient variability.

The term “about twice weekly” as used herein means approximate number, and “about twice weekly” can include twice in one week, e.g., a first dose in three days and a second dose in three days, a first dose in three days and a second dose in four days, a first dose in four days and a second dose in three days, a first dose in four days and a second dose in four days. The term “about twice weekly” can also include every three days, every four days, or every five days.

The term “about once a week” as used herein means approximate number, and “about once a week” can include every seven days $\pm$ two days, i.e., every five days to every nine days. The dosing frequency of “once a week” thus can be every five days, every six days, every seven days, every eight days, or every nine days.

The term “individualized prophylaxis” or “prophylactic and individualized” as used herein means use of a FVIII polypeptide for an individualized dosing and/or dosing interval or frequency to prevent or inhibit occurrence of one or more spontaneous and/or uncontrollable bleeding or bleeding episodes or to reduce the frequency of one or more spontaneous and/or uncontrollable bleeding or bleeding episodes. “Individualized” within the context of prophylaxis is used synonymous with “tailored” throughout this application. For example, “individualized prophylaxis” also means “tailored prophylaxis” and “prophylactic and individualized” also means “prophylactic and tailored.” In one embodiment, the “individualized interval” includes every 3 days $\pm$ 2 days, i.e., every day to every five days. The dosing frequency of the “individualized interval prophylaxis” thus can be every day, every two days, every three days, every four days, or every five days.

The term “on-demand treatment,” as used herein, means treatment that is intended to take place over a short course of time and is in response to an existing condition, such as a bleeding episode, or a perceived short term need such as planned surgery. The “on-demand treatment” is used interchangeably with “episodic” treatment. Conditions that can require on-demand treatment include a bleeding episode, hemarthrosis, muscle bleed, oral bleed, hemorrhage, hemorrhage into muscles, oral hemorrhage, trauma, trauma capitis, gastrointestinal bleeding, intracranial hemorrhage, intra-abdominal hemorrhage, intrathoracic hemorrhage, bone fracture, central nervous system bleeding, bleeding in the retropharyngeal space, bleeding in the retroperitoneal space, or bleeding in the iliopsoas sheath. Bleeding episodes other than these are also included. The subject can be in need of surgical prophylaxis, peri-operative management, or treatment for surgery. Such surgeries include minor surgery, major surgery, tooth extraction, tonsillectomy, other dental/thoraco-facial surgeries, inguinal herniotomy, synovectomy, total knee replacement, other joint replacement, craniotomy, osteosynthesis, trauma surgery, intracranial surgery, intra-abdominal surgery, or intrathoracic surgery. Surgeries other than these are also included.

Additional conditions that can require on-demand treatment include minor hemorrhage, hemarthroses, superficial muscle hemorrhage, soft tissue hemorrhage, moderate hemorrhage, intramuscle or soft tissue hemorrhage with dissection, mucous membrane hemorrhage, hematuria, major hemorrhage, hemorrhage of the pharynx, hemorrhage of the retropharynx, hemorrhage of the retroperitoneum, hemor-

rhage of the central nervous system, bruises, cuts, scrapes, joint hemorrhage, nose bleed, mouth bleed, gum bleed, intracranial bleeding, intraperitoneal bleeding, minor spontaneous hemorrhage, bleeding after major trauma, moderate skin bruising, or spontaneous hemorrhage into joints, muscles, internal organs or the brain. Additional reasons for on-demand treatment include the need for peri-operative management for surgery or dental extraction, major surgery, extensive oral surgery, urologic surgery, hernia surgery, orthopedic surgery such as replacement of knee, hip, or other major joint.

The term “treatment” or “treating” as used herein means amelioration or reduction of one or more symptoms of bleeding diseases or disorders including, but not limited to, hemophilia B. In one embodiment, “treatment of” or “treating” a bleeding disease or disorder includes prevention of one or more symptoms of a bleeding disease or disorder. In a bleeding disease or disorder caused by a FVIII deficiency (e.g., a low baseline FVIII activity), the term “treatment” or “treating” means a FVIII replacement therapy. By administering a pharmaceutical composition comprising a FVIII polypeptide to a subject, the subject can achieve and/or maintain a plasma trough level of a FVIII activity at about 1 IU/dl or above 1 IU/dl. In other embodiments, “treatment” or “treating” means reduction of the frequency of one or more symptoms of bleeding diseases or disorders, e.g., spontaneous or uncontrollable bleeding episodes. “Treatment,” however, need not be a cure.

The term “perioperative management” as used herein means use of a pharmaceutical composition comprising a FVIII polypeptide before, concurrently with, or after an operative procedure, e.g., a surgical operation. The use for “perioperative management” of one or more bleeding episode includes surgical prophylaxis before (i.e., preoperative), during (i.e., intraoperative), or after (i.e., postoperative) a surgery to prevent one or more bleeding or bleeding episode or reducing or inhibiting spontaneous and/or uncontrollable bleeding episodes before, during, and after a surgery.

“Baseline,” as used herein, is the lowest measured plasma Factor VIII level in a subject prior to administering a dose. The FVIII plasma levels can be measured at two time points prior to dosing: at a screening visit and immediately prior to dosing. Alternatively, (a) the baseline in patients whose pretreatment FVIII activity is $<1\%$, who have no detectable FVIII antigen, and have nonsense genotypes can be defined as 0% , (b) the baseline for patients with pretreatment FVIII activity $<1\%$ and who have detectable FVIII antigen can be set at 0.5% , (c) the baseline for patients whose pretreatment FVIII activity is between $1-2\%$ is C_{min} (the lowest activity throughout the PK study), and (d) the baseline for patients whose pretreatment FVIII activity is $\geq 2\%$ can be set at 2% . Activity above the baseline pre-dosing can be considered residue drug from prior treatment, and can be decayed to baseline and subtracted from the PK data following rFVIII-Fc dosing.

“ $T_{1/2\beta}$,” or “ $T_{1/2\beta}$ ” or “Beta HL,” as used herein, is half-life associated with elimination phase, $t_{1/2\beta} = (\ln 2)/\text{elimination rate constant associated with the terminal phase}$. The $T_{1/2\beta}$ can be measured by FVIII activity or by FVIII antigen level in plasma. The $T_{1/2\beta}$ based on activity is shown as $T_{1/2\beta}$ (activity), and the T_{in} based on the FVIII antigen level can be shown as $T_{1/2\beta}$ (antigen). Both $T_{1/2\beta}$ (activity) and $T_{1/2\beta}$ (antigen) can be shown as ranges or a geometric mean.

“Trough,” as used herein, is the lowest plasma FVIII activity level reached after administering a dose of chimeric

polypeptide of the invention or another FVIII molecule and before the next dose is administered, if any. Trough is used interchangeably herein with "threshold." Baseline FVIII levels are subtracted from measured FVIII levels to calculate the trough level.

The term "annualized bleeding rate" ("ABR") as used herein refers to the number of bleeding episodes (including spontaneous and traumatic bleeds) experienced by a subject during a defined time period, extrapolated to 1 year. For example two bleeds in six months would indicate an ABR of four. The median ABR provides the middle value among all observed ABRs, indicating that half of the subjects had individual ABRs less than or equal to the median and half had ABRs greater than or equal to the median.

The term "inter quartile range" ("IQR") as used herein refers to a measure of statistical dispersion, being equal to the difference between the upper and lower quartiles. Unlike (total) range, the interquartile range is a robust statistic, having a breakdown point of 25%, and is thus often preferred to the total range. For a symmetric distribution (where the median equals the midline, the average of the first and third quartiles), half the IQR equals the median absolute deviation (MAD). The median is the corresponding measure of central tendency.

"Subject," as used herein means a human. Subject as used herein includes an individual who is known to have at least one incidence of uncontrolled bleeding episodes, who has been diagnosed with a disease or disorder associated with uncontrolled bleeding episodes, e.g., a bleeding disease or disorder, e.g., hemophilia A, who is susceptible to uncontrolled bleeding episodes, e.g., hemophilia, or any combinations thereof. Subjects can also include an individual who is in danger of one or more uncontrollable bleeding episodes prior to a certain activity, e.g., a surgery, a sport activity, or any strenuous activities. The subject can have a baseline FVIII activity less than 1%, less than 0.5%, less than 2%, less than 2.5%, less than 3%, or less than 4%. Subjects also include pediatric humans. Pediatric human subjects are birth to 20 years, preferably birth to 18 years, birth to 16 years, birth to 15 years, birth to 12 years, birth to 11 years, birth to 6 years, birth to 5 years, birth to 2 years, or 2 to 11 years of age.

"Therapeutic dose," "dose," "effective dose," or "dosing amount" as used herein, means a dose that achieves a plasma trough level of a FVIII activity at least about 1 IU/dl or above 1 IU/dl in the subject administered with a pharmaceutical composition comprising a FVIII polypeptide. For the purpose of this invention, in one embodiment, the "dose" refers to the amount of the doses that a plasma trough level of a FVIII activity is maintained at least about 1 IU/dl or above 1 IU/dl, at least about 2 IU/dl or above 2 IU/dl, at least about 3 IU/dl or above 3 IU/dl, at least about 4 IU/dl or above 4 IU/dl, or at least about 5 IU/dl or above 5 IU/dl throughout the administration of a pharmaceutical composition comprising a FVIII polypeptide. In another embodiment, the "dose" reduces or decreases frequency of bleeding or bleeding disorder. In other embodiments, the "dose" stops on-going, uncontrollable bleeding or bleeding episodes. In still other embodiments, the "dose" prevents spontaneous bleeding or bleeding episodes in a subject susceptible to such spontaneous bleeding or bleeding episodes. The "dose" or "therapeutic dose" need not cure hemophilia.

The term "target joint" is defined as a major joint (e.g., hip, elbow, wrist, shoulder, knee, and ankle) into which repeated bleeding occurs (frequency of 23 bleeding episodes into the same joint in a consecutive 6-month period).

The term "bleeding episode" as used herein adopts a standardized definition of a bleeding episode. A bleeding episode started from the first sign of bleeding, and ended 72 hours after the last treatment for the bleeding, within which any symptoms of bleeding at the same location, or injections less than or equal to 72 hours apart, were considered the same bleeding episode. Any injection to treat the bleeding episode, taken more than 72 hours after the preceding one, was considered the first injection to treat a new bleeding episode at the same location. Any bleeding at a different location was considered a separate bleeding episode regardless of time from last injection. This definition has been proposed by the Subcommittee on Standards and Criteria, FVIII/FIX subcommittee of the International Society of Thrombosis and Hemostasis and has been used by the PedNet multicenter study in hemophilia.

The term "annualized bleeding rates (ABRs) as used herein refers to the number of bleeding episodes that are annualized for each patient using the following formula:

$$ABR = \frac{\text{Number of bleeding episodes during efficacy period}}{\text{Total number of days during the efficacy period}} \times 365.25$$

"Variant," as used herein, refers to a polynucleotide or polypeptide differing from the original polynucleotide or polypeptide, but retaining essential properties thereof, e.g., Factor VIII coagulant activity or Fc (FcRn binding) activity. Generally, variants are overall closely similar, and, in many regions, identical to the original polynucleotide or polypeptide. Variants include polypeptide and polynucleotide fragments, deletions, insertions, and modified versions of original polypeptides.

II. Pharmaceutical Composition

The present invention is directed to a FVIII polypeptide formulated as a pharmaceutical composition. The pharmaceutical composition can be formulated for administration to humans. The FVIII polypeptide can be a short-acting FVIII polypeptide or a long-acting FVIII polypeptide. The pharmaceutical compositions comprise pharmaceutically acceptable carriers, including, e.g., ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat. Various methods of formulating the invention are well known in the art.

In certain formulations provided herein, a FVIII polypeptide is formulated as a sterile, preservative-free, non-pyrogenic, lyophilized, white to off-white powder to cake, for intravenous (IV) administration. The formulation can be provided in a single-use vial.

In certain embodiments, a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII_{IFC}) formulation is provided in a single-use vial manufactured to contain, following reconstitution with an appropriate amount of diluent, about 83 IU/ml, 167 IU/ml, 250 IU/ml, 333 IU/ml, 500 IU/ml, 667 IU/ml, 1000 IU/ml, 1333

IU/ml, 1667 IU/ml, or 2000 IU/ml of the long-acting FVIII polypeptide. In certain embodiments in which diluent is added to a final volume of about 3 ml, a single-use vial can nominally contain about 250, about 500, about 750, about 1000, about 1500, about 2000, about 3000, about 4000, about 5000, or about 6000 International Units (IU) of the FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIIIc).

In certain embodiments the formulation includes, in addition to the active FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIIIc): sucrose (which can act as a stabilizer), sodium chloride (which can act as a bulking agent), L-histidine (which can act as a buffer), calcium chloride, and polysorbate 20 or polysorbate 80 (which can act as a stabilizer). The formulation is provided with a diluent comprising a sterile sodium chloride solution. In certain embodiments, the diluent is provided in a pre-filled syringe.

Accordingly, provided herein is a pharmaceutical composition comprising a specified amount of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIIIc) (in IU), along with the excipients sucrose, NaCl, L-histidine, calcium chloride, and polysorbate 20 or polysorbate 80. The compositions provided herein comprise various concentrations of the various excipients, and the concentrations can be expressed in various ways. For example, the concentration of a given excipient can be expressed as a molar concentration (e.g., M or mM), as a weight/volume percent, (e.g., grams per 100 ml diluent), or as milligrams per milliliter (mg/ml). Formulations provided herein can contain specified amounts of the various excipients at a level of precision ranging from approximate, e.g., concentrations expressed only to one significant figure (e.g., about 0.1% (w/v)), or with more precision, e.g., out to 2, 3, 4, 5, or 6 significant figures (e.g., about 3.88 mg/ml, with precision out to three significant figures). The necessary level of precision can vary depending on, e.g., the requirements of a given regulatory agency, or the manufacturing process. In certain embodiments the pharmaceutical composition comprises a reconstituted formulation, which can be provided as a lyophilisate, optionally accompanied by a diluent.

In certain embodiments, the pharmaceutical composition comprises about 50 IU/ml to about 2500 IU/ml rFVIIIc, e.g., 83 IU/ml, 167 IU/ml, 250 IU/ml, 333 IU/ml, 500 IU/ml, 667 IU/ml, 1000 IU/ml, 1333 IU/ml, 1667 IU/ml, or 2000 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIIIc). In certain embodiments, the pharmaceutical composition comprises 83 IU/ml, 167 IU/ml, 250 IU/ml, 333 IU/ml, 500 IU/ml, 667 IU/ml, 1000 IU/ml, 1333 IU/ml, 1667 IU/ml, or 2000 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIIIc) in a formulation comprising about 13.3 mg/ml or about 1.33% (w/v) sucrose, about 12.0 mg/ml or about 205 mM NaCl, about 1.03 mg/ml or about 6.64 mM L-histidine, about 0.80 mg/ml or about 5.4 mM calcium chloride dihydrate, and about 0.13 mg/ml or about 0.013% (w/v) polysorbate 20 or polysorbate 80.

In certain embodiments, the pre-lyophilization pharmaceutical composition comprises about 100 IU/ml to about 4000 IU/ml rFVIIIc, e.g., 150 IU/ml, 287.5 IU/ml, 431.25 IU/ml, 575 IU/ml, 862.5 IU/ml, 1150 IU/ml, 1725 IU/ml, 2300 IU/ml, 2875 IU/ml, or 3450 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIIIc). In certain embodiments, the pharmaceutical composition comprises 150 IU/ml, 287.5 IU/ml, 431.25 IU/ml, 575 IU/ml, 862.5 IU/ml, 1150 IU/ml, 1725

IU/ml, 2300 IU/ml, 2875 IU/ml, or 3450 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIIIc) in a formulation comprising about 20.0 mg/ml or about 2.0% (w/v) sucrose, about 18.0 mg/ml or about 308 mM NaCl, about 1.55 mg/ml or about 9.8 mM L-histidine, about 1.18 mg/ml or about 8.0 mM calcium chloride dihydrate, and about 0.2 mg/ml or about 0.02% (w/v) polysorbate 20 or polysorbate 80.

In certain embodiments, the pharmaceutical composition comprises a pharmaceutically acceptable amount of sucrose. In certain embodiments, the pharmaceutical composition comprises about 1% (w/v) to about 2.5% (w/v) sucrose, preferably about 1.3% (w/v) to about 2% (w/v) e.g., about 1.33% (w/v) sucrose or about 2.0% (w/v) sucrose. In certain related embodiments the pharmaceutical composition comprises about 10 mg/ml to about 25 mg/ml sucrose, preferably about 13 mg/ml to about 20 mg/ml sucrose, e.g., about 13.3 mg/ml sucrose or about 20.0 mg/ml sucrose. In still other embodiments, stabilizing agents such as trehalose, raffinose and/or arginine can be used at these concentrations in lieu of or in combination with sucrose.

In certain embodiments, the pharmaceutical composition comprises about 150 mM to about 250 mM NaCl, about 175 mM to about 225 mM NaCl, and about 200 mM to about 210 mM NaCl, e.g., about 205 mM NaCl. In certain related embodiments, the pharmaceutical composition comprises about 8.8 mg/ml to about 14.6 mg/ml NaCl, about 10 mg/ml NaCl to about 13 mg/ml, e.g., about 12.0 mg/ml NaCl. In certain embodiments, NaCl is provided at the desired concentration in a diluent solution in which a lyophilisate comprising a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIIIc) is reconstituted.

In another embodiment, the compositions of the invention do not include bulking agents such as mannitol, glycine, alanine and/or hydroxyethyl starch. In other embodiments, NaCl is the sole bulking agent.

In certain embodiments, the pre-lyophilization pharmaceutical composition comprises about 250 mM to about 350 mM NaCl, about 275 mM to about 325 mM NaCl, e.g., about 308 mM NaCl. In certain related embodiments, the pharmaceutical composition comprises about 14.6 mg/ml to about 20.5 mg/ml NaCl, about 16 mg/ml to about 19 mg/ml, e.g., about 18.0 mg/ml NaCl.

In certain embodiments, the pharmaceutical composition comprises a pharmaceutically acceptable amount of L-histidine. In certain embodiments, the pharmaceutical composition comprises about 5 mM to about 15 mM L-histidine, e.g., about 6.64 mM L-histidine or about 9.8 mM L-histidine. In certain related embodiments the pharmaceutical composition comprises about 0.75 mg/ml to about 2.25 mg/ml L-histidine, e.g., about 1.03 mg/ml L-histidine or about 1.55 mg/ml L-histidine. In certain embodiments, L-histidine is provided as part of a lyophilisate, which, upon reconstitution with an appropriate amount of diluent provides L-histidine at the desired concentration.

In certain embodiments, the pharmaceutical composition comprises a pharmaceutically acceptable amount of calcium chloride. In certain embodiments, the pharmaceutical composition comprises about 5 mM to about 10 mM calcium chloride, e.g., about 5.4 mM calcium chloride or about 8 mM calcium chloride. In certain related embodiments the pharmaceutical composition comprises about 0.75 mg/ml to about 1.5 mg/ml calcium chloride dihydrate, e.g., about 0.8 mg/ml calcium chloride dihydrate or about 1.18 mg/ml calcium chloride dihydrate. In certain embodiments, calcium chloride is provided as part of a lyophilisate, which,

upon reconstitution with an appropriate amount of diluent provides calcium chloride at the desired concentration.

In certain embodiments, the pharmaceutical composition comprises a pharmaceutically acceptable amount of polysorbate 20 or polysorbate 80. In certain related embodiments the pharmaceutical composition comprises about 0.008% (w/v) to about 0.025% (w/v) polysorbate 20 or polysorbate 80, e.g., about 0.013% (w/v) polysorbate 20 or polysorbate 80 or about 0.02% (w/v) polysorbate 20 or polysorbate 80. In certain related embodiments the pharmaceutical composition comprises about 0.08 mg/ml to about 0.25 mg/ml polysorbate 20 or polysorbate 80, e.g., about 0.13 mg/ml polysorbate 20 or polysorbate 80 or about 0.20 mg/ml polysorbate 20 or polysorbate 80. In certain embodiments, polysorbate 20 or polysorbate 80 is provided as part of a lyophilisate, which, upon reconstitution with an appropriate amount of diluent provides polysorbate 20 or polysorbate 80 at the desired concentration.

In certain embodiments, the pharmaceutical composition comprises: about 50 IU/ml to about 2500 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); about 1% (w/v) to about 2.5% (w/v) of sucrose; about 150 mM to about 250 mM NaCl; about 5 mM to about 15 mM L-histidine; about 5 mM to about 10 mM calcium chloride; and about 0.008% (w/v) to about 0.025% of polysorbate 20 or polysorbate 80. In certain embodiments the pharmaceutical composition is provided as a lyophilisate and a diluent. In certain embodiments the amount of lyophilisate provides about 3 ml of a pharmaceutical composition with the desired ingredients at the desired concentrations.

In certain embodiments, the pharmaceutical composition comprises: about 50 IU/ml to about 2500 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); about 10 mg/ml to about 25 mg/ml of sucrose; about 8.8 mg/ml to about 14.6 mg/ml NaCl; about 0.75 mg/ml to about 2.25 mg/ml L-histidine; about 0.75 mg/ml to about 1.5 mg/ml calcium chloride dihydrate; and about 0.08 mg/ml to about 0.25 mg/ml of polysorbate 20 or polysorbate 80. In certain embodiments the pharmaceutical composition is provided as a lyophilisate and a diluent. In certain embodiments the amount of lyophilisate provides about 3 ml of a pharmaceutical composition with the desired ingredients at the desired concentrations.

Exemplary compositions are provided in Table 1 and in Table 2 in the Examples.

For example, the disclosure provides a pharmaceutical composition comprising: about 83 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); about 1.33% (w/v) of sucrose; about 205 mM NaCl; about 6.64 mM L-histidine; about 5.4 mM calcium chloride dihydrate, and about 0.013% (w/v) of polysorbate 20 or polysorbate 80. The disclosure further provides a pharmaceutical composition comprising: about 167 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); about 1.33% (w/v) of sucrose; about 205 mM NaCl; about 6.64 mM L-histidine; about 5.4 mM calcium chloride dihydrate, and about 0.013% (w/v) of polysorbate 20 or polysorbate 80. The disclosure further provides a pharmaceutical composition comprising: about 250 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); about 1.33% (w/v) of sucrose; about 205 mM NaCl; about 6.64 mM L-histidine; about 5.4 mM calcium chloride dihydrate, and about 0.013% (w/v) of polysorbate 20 or polysorbate 80. The disclosure further provides a pharmaceutical composition comprising: about 333 IU/ml of

a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); about 1.33% (w/v) of sucrose; about 205 mM NaCl; about 6.64 mM L-histidine; about 5.4 mM calcium chloride dihydrate, and about 0.013% (w/v) of polysorbate 20 or polysorbate 80. The disclosure further provides a pharmaceutical composition comprising: about 500 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); about 1.33% (w/v) of sucrose; about 205 mM NaCl; about 6.64 mM L-histidine; about 5.4 mM calcium chloride dihydrate, and about 0.013% (w/v) of polysorbate 20 or polysorbate 80. The disclosure further provides a pharmaceutical composition comprising: about 667 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); about 1.33% (w/v) of sucrose; about 205 mM NaCl; about 6.64 mM L-histidine; about 5.4 mM calcium chloride dihydrate, and about 0.013% (w/v) of polysorbate 20 or polysorbate 80. The disclosure further provides a pharmaceutical composition comprising: about 1000 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); about 1.33% (w/v) of sucrose; about 205 mM NaCl; about 6.64 mM L-histidine; about 5.4 mM calcium chloride dihydrate, and about 0.013% (w/v) of polysorbate 20 or polysorbate 80. The disclosure also provides a pharmaceutical composition comprising: about 1333 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); about 1.33% (w/v) of sucrose; about 205 mM NaCl; about 6.64 mM L-histidine; about 5.4 mM calcium chloride, and about 0.013% (w/v) of polysorbate 20 or polysorbate 80. In some embodiments, the disclosure provides a pharmaceutical composition comprising: about 1667 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); about 1.33% (w/v) of sucrose; about 205 mM NaCl; about 6.64 mM L-histidine; about 5.4 mM calcium chloride, and about 0.013% (w/v) of polysorbate 20 or polysorbate 80. In other embodiments, the disclosure provides a pharmaceutical composition comprising: about 2000 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); about 1.33% (w/v) of sucrose; about 205 mM NaCl; about 6.64 mM L-histidine; about 5.4 mM calcium chloride, and about 0.013% (w/v) of polysorbate 20 or polysorbate 80.

The disclosure further provides a pharmaceutical composition comprising: about 83 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); about 13.3 mg/ml of sucrose; about 12.0 mg/ml NaCl; about 1.03 mg/ml L-histidine; about 0.80 mg/ml calcium chloride dihydrate, and about 0.10 mg/ml of polysorbate 20 or polysorbate 80. The disclosure further provides a pharmaceutical composition comprising: about 1673 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); about 13.3 mg/ml of sucrose; about 12.0 mg/ml NaCl; about 1.03 mg/ml L-histidine; about 0.80 mg/ml calcium chloride dihydrate, and about 0.10 mg/ml of polysorbate 20 or polysorbate 80. The disclosure further provides a pharmaceutical composition comprising: about 250 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); about 13.3 mg/ml of sucrose; about 12.0 mg/ml NaCl; about 1.03 mg/ml L-histidine; about 0.80 mg/ml calcium chloride dihydrate, and about 0.10 mg/ml of polysorbate 20 or polysorbate 80. The disclosure further provides a pharmaceutical composition comprising: about 333 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); about 13.3 mg/ml of sucrose; about 12.0 mg/ml NaCl; about 1.03 mg/ml L-his-

tidine; about 0.80 mg/ml calcium chloride dihydrate, and about 0.10 mg/ml of polysorbate 20 or polysorbate 80. The disclosure further provides a pharmaceutical composition comprising: about 500 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); about 13.3 mg/ml of sucrose; about 12.0 mg/ml NaCl; about 1.03 mg/ml L-histidine; about 0.80 mg/ml calcium chloride dihydrate, and about 0.10 mg/ml of polysorbate 20 or polysorbate 80. The disclosure further provides a pharmaceutical composition comprising: about 667 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); about 13.3 mg/ml of sucrose; about 12.0 mg/ml NaCl; about 1.03 mg/ml L-histidine; about 0.80 mg/ml calcium chloride dihydrate, and about 0.10 mg/ml of polysorbate 20 or polysorbate 80. The disclosure also provides a pharmaceutical composition comprising: about 1000 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); about 13.3 mg/ml of sucrose; about 12.0 mg/ml NaCl; about 1.03 mg/ml L-histidine; about 0.80 mg/ml calcium chloride dihydrate, and about 0.10 mg/ml of polysorbate 20 or polysorbate 80. The disclosure also provides a pharmaceutical composition comprising: about 1333 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); about 13.3 mg/ml of sucrose; about 12.0 mg/ml NaCl; about 1.03 mg/ml L-histidine; about 0.80 mg/ml calcium chloride dihydrate, and about 0.10 mg/ml of polysorbate 20 or polysorbate 80. In other embodiments, the disclosure provides a pharmaceutical composition comprising: about 1667 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); about 13.3 mg/ml of sucrose; about 12.0 mg/ml NaCl; about 1.03 mg/ml L-histidine; about 0.80 mg/ml calcium chloride dihydrate, and about 0.10 mg/ml of polysorbate 20 or polysorbate 80. In some embodiments, the disclosure provides a pharmaceutical composition comprising: about 2000 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); about 13.3 mg/ml of sucrose; about 12.0 mg/ml NaCl; about 1.03 mg/ml L-histidine; about 0.80 mg/ml calcium chloride dihydrate, and about 0.10 mg/ml of polysorbate 20 or polysorbate 80.

This disclosure also provides the components of a pharmaceutical kit. Such a kit includes one or more containers and optional attachments. A kit as provided herein facilitates administration of an effective amount of the FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc) to a subject in need thereof. In certain embodiments, the kit facilitates administration of the FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc) via intravenous infusion. In certain embodiments, the kit facilitates self-administration of the FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc) via intravenous infusion.

In certain embodiments, the disclosure provides a pharmaceutical kit comprising: a first container comprising a lyophilized powder or cake, where the powder or cake comprises: (i) a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc), (ii) sucrose (and/or trehalose, raffinose or arginine); (iii) NaCl; (iv) L-histidine; (v) calcium chloride dihydrate; and (vi) polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections to be combined with the lyophilized powder of the first container. In certain embodiments, sufficient diluent is provided to produce about 3 ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc) formulation with

desired properties as disclosed herein. In certain embodiments, the second container is a pre-filled syringe associated with a plunger, to allow addition of the diluent to the first container, reconstitution of the contents of the first container, and transfer back into the syringe. In certain embodiments, the kit further provides an adaptor for attaching the syringe to the first container. In certain embodiments the kit further provides a needle and infusion tubing, to be attached to the syringe containing the reconstituted FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc) formulation to allow IV infusion of the formulation.

In certain embodiments a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc) is provided in a total amount from about 200 IU to about 6000 IU, e.g., about 250 IU, about 500 IU, about 750 IU, about 1000 IU, about 1500 IU, about 2000 IU, about 3000 IU, about 4000 IU, about 5000 IU, or about 6000 IU.

In one embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises (i) about 250 IU of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 83 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); (ii) about 1.33% (w/v) of sucrose; (iii) about 205 mM of NaCl; (iv) about 6.64 mM L-histidine; (v) about 5.4 mM of calcium chloride; and (vi) about 0.013% (w/v) of polysorbate 20 or polysorbate 80.

In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 500 IU of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 167 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); (ii) about 1.33% (w/v) of sucrose; (iii) about 205 mM of NaCl; (iv) about 6.64 mM L-histidine; (v) about 5.4 mM of calcium chloride; and (vi) about 0.013% (w/v) of polysorbate 20 or polysorbate 80.

In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 750 IU of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 250 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII(Fc); (ii) about 1.33% (w/v) of sucrose; (iii) about 205 mM of NaCl; (iv) about 6.64 mM L-histidine; (v) about 5.4

In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 250 IU of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII_{SC}), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 83 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g.,

In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 5000 IU of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIIIc), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution com-

prising: (i) about 1667 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII-Fc); (ii) about 13.3 mg/ml of sucrose; (iii) about 12.0 mg/ml of NaCl; (iv) about 1.03 mg/ml L-histidine; (v) about 0.80 mg/ml of calcium chloride dihydrate; and (vi) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.

In a further embodiment, a pharmaceutical kit is provided which comprises a first container comprising a lyophilized powder, where the powder comprises: (i) about 6000 IU of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII-Fc), (ii) about 40 mg of sucrose; (iii) about 36 mg of NaCl; (iv) about 3.1 mg of L-histidine; (v) about 2.4 mg of calcium chloride dihydrate; and (vi) about 0.40 mg of polysorbate 20 or polysorbate 80; and a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: (i) about 2000 IU/ml of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII-Fc); (ii) about 13.3 mg/ml of sucrose; (iii) about 12.0 mg/ml of NaCl; (iv) about 1.03 mg/ml L-histidine; (v) about 0.80 mg/ml of calcium chloride dihydrate; and (vi) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.

In certain embodiments the first container of a pharmaceutical kit provided herein is a glass vial comprising a rubber stopper. In certain embodiments, the second container a pharmaceutical kit provided herein is a syringe body, associated with a plunger. In certain embodiments, the syringe is a pre-filled syringe containing the diluent. In certain embodiments, a pharmaceutical kit provided herein further comprises an adaptor to connect the glass vial to the syringe body. In certain embodiments a pharmaceutical kit provided herein further comprises infusion tubing associated with a needle to be connected to the syringe, suitable for intravenous infusion.

In certain embodiments, a desired dose of a FVIII polypeptide (e.g., a long-acting or short-acting FVIII polypeptide, e.g., rFVIII-Fc) can be achieved through the use of one pharmaceutical kit as provided herein. In certain embodiments, more than one pharmaceutical kit can be used to achieve a desired dose. Provided herein is a method of combining, or pooling the formulations contained in two or more pharmaceutical kits as provided herein in order to achieve a desired dose.

In some embodiments, the FVIII polypeptide for the pharmaceutical composition is a short-acting FVIII polypeptide. In other embodiments, the FVIII polypeptide for the pharmaceutical composition is a long-acting FVIII polypeptide.

In some embodiments, a long-acting FVIII polypeptide in a pharmaceutical composition comprises a Factor VIII portion and a non-Factor VIII portion, e.g., a heterologous moiety. In one embodiment, the heterologous moiety is capable of extending in vivo or in vitro half-life of the FVIII polypeptide. Exemplary non-Factor VIII portions include, but are not limited to, Fc, albumin, a PAS sequence, transferrin, CTP (28 amino acid C-terminal peptide (CTP) of human chorionic gonadotropin (hCG) with its 4 O-glycans), polyethylene glycol (PEG), hydroxyethyl starch (HES), albumin binding polypeptide, albumin-binding small molecules, or any combination thereof. Exemplary long-acting polypeptides of the invention include, e.g., Factor VIII-Fc polypeptides, Factor VII-albumin polypeptides, Factor VII-PAS polypeptides, Factor VIII-transferrin polypeptides, Factor VIII-CTP polypeptides, Factor VIII-PEG polypeptides,

Factor VIII-HES polypeptides, Factor VIII-albumin binding polypeptide polypeptides, or Factor VIII-albumin-binding small molecule polypeptides.

The Factor VIII (the Factor VIII portion of a long-acting FVIII polypeptide or FVIII of a short-acting FVIII polypeptide) can be at least 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to a Factor VIII amino acid sequence shown in Table 11 without a signal sequence (amino acids 20 to 1457 of SEQ ID NO:2; and amino acids 20 to 2351 of SEQ ID NO:6), wherein the Factor VIII portion has Factor VIII activity. The Factor VIII (the Factor VIII portion of a chimeric polypeptide) can be identical to a Factor VIII amino acid sequence shown in Table 11 without a signal sequence (amino acids 20 to 1457 of SEQ ID NO:2; and amino acids 20 to 2351 of SEQ ID NO:6).

The Factor VIII (the Factor VIII portion of a long-acting FVIII polypeptide or FVIII of a short-acting FVIII polypeptide) can be at least 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to a Factor VIII amino acid sequence shown in Table 11 with a signal sequence (amino acids 1 to 1457 of SEQ ID NO:2 and amino acids 1 to 2351 of SEQ ID NO:6), wherein the Factor VIII portion has Factor VI activity. The Factor VIII (the Factor VIII portion of a long-acting FVIII polypeptide or a short-acting FVIII polypeptide) can be identical to a Factor VIII amino acid sequence shown in Table 11 with a signal sequence (amino acids 1 to 1457 of SEQ ID NO:2 and amino acids 1 to 2351 of SEQ ID NO:6).

The long-acting polypeptide can comprise a sequence at least 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the Factor VIII and Fc amino acid sequence shown in Table 11A(i) without a signal sequence (amino acids 20 to 1684 of SEQ ID NO:2) or at least 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the Factor VIII and Fc amino acid sequence shown in Table 11A(i) with a signal sequence (amino acids 1 to 1684 of SEQ ID NO:2), wherein the sequence has Factor VIII activity. The Factor VIII activity can be measured by activated Partial Thromboplastin Time (aPTT) assay, chromogenic assay, or other known methods. The long-acting FVIII polypeptide can comprise the sequence identical to the Factor VIII and Fc amino acid sequence shown in Table 11A(i) without a signal sequence (amino acids 20 to 1684 of SEQ ID NO:2) or identical to the Factor VIII and Fc amino acid sequence shown in Table 11A(i) with a signal sequence (amino acids 1 to 1684 of SEQ ID NO:2).

As discussed above, exemplary long-acting polypeptides also include Factor VIII fused to one or more albumin polypeptides, albumin binding polypeptides, or albumin-binding small molecules. In one embodiment, the albumin is human albumin. The albumin or albumin binding protein can be fused to either the N-terminal end of FVIII or to the C-terminal end of FVIII or inserted between two amino acids in FVIII. Examples of albumin, e.g., fragments thereof, that can be used in the present invention are known. e.g., U.S. Pat. Nos. 7,592,010; 6,686,179; and Schulte, *Thrombosis Res.* 124 Suppl. 2:S6-S8 (2009), each of which is incorporated herein by reference in its entirety.

The albumin binding polypeptides can compromise, without limitation, bacterial albumin-binding domains, albumin-binding peptides, or albumin-binding antibody fragments that can bind to albumin. Domain 3 from streptococcal protein G, as disclosed by Kraulis et al., *FEBS Lett.* 378: 190-194 (1996) and Linhult et al., *Protein Sci.* 11:206-213 (2002) is an example of a bacterial albumin-binding domain. Examples of albumin-binding peptides include a series of peptides having the core sequence DICLPRWGCLW (SEQ

ID NO: 7). See, e.g., Dennis et al., *J. Biol. Chem.* 2002, 277: 35035-35043 (2002). Examples of albumin-binding antibody fragments are disclosed in Muller and Kontermann, *Curr. Opin. Mol. Ther.* 9:319-326 (2007); Roovers et al., *Cancer Immunol. Immunother.* 56:303-317 (2007), and Holt et al., *Prot. Eng. Design Sci.*, 21:283-288 (2008), which are incorporated herein by reference in their entireties.

In certain aspects, a long-acting FVIII polypeptide of the invention comprises at least one attachment site for a non-polypeptide small molecule, variant, or derivative that can bind to albumin thereof. An example of such albumin binding moieties is 2-(3-maleimidopropanamido)-6-(4-(4-iodophenyl)butanamido)hexanoate ("Albu" tag) as disclosed by Trusset et al., *Bioconjugate Chem.* 20:2286-2292 (2009).

As discussed above, exemplary long-acting polypeptides also include Factor VIII fused to at least one C-terminal peptide (CTP) of the 0 subunit of human chorionic gonadotropin or fragment, variant, or derivative thereof. The CTP can be fused to Factor VIII either the N-terminal end of FVIII or to the C-terminal end of FVIII or inserted between two amino acids in FVIII. One or more CTP peptides fused to or inserted into a recombinant protein is known to increase the in vivo half-life of that protein. See, e.g., U.S. Pat. No. 5,712,122, incorporated by reference herein in its entirety. Exemplary CTP peptides include DPRFQDSSSSKAPPPSLPSPRLPGPSDTPIL (SEQ ID NO: 8) or SSSSKAPPPSLPSPRLPGPSDTPILPQ. (SEQ ID NO: 9). See, e.g., U.S. Patent Application Publication No. US 2009/0087411 A1, incorporated by reference.

As discussed above, exemplary long-acting FVIII polypeptides also include Factor VIII fused to at least one PAS sequence or fragment, variant, or derivative thereof. The PAS sequence can be fused to either the N-terminal end of FVIII or to the C-terminal end of FVIII or inserted between two amino acids in FVIII. A PAS peptide or PAS sequence, as used herein, means an amino acid sequence comprising mainly alanine and serine residues or comprising mainly alanine, serine, and proline residues, the amino acid sequence forming random coil conformation under physiological conditions. Accordingly, the PAS sequence is a building block, an amino acid polymer, or a sequence cassette comprising, consisting essentially of, or consisting of alanine, serine, and proline which can be used as a part of the heterologous moiety in the chimeric protein. An amino acid polymer also can form random coil conformation when residues other than alanine, serine, and proline are added as a minor constituent in the PAS sequence. By "minor constituent" is meant that that amino acids other than alanine, serine, and proline can be added in the PAS sequence to a certain degree, e.g., up to about 12%, i.e., about 12 of 100 amino acids of the PAS sequence, up to about 10%, up to about 9%, up to about 8%, about 6%, about 5%, about 4%, about 3%, i.e. about 2%, or about 1%, of the amino acids. The amino acids different from alanine, serine and proline can be selected from the group consisting of Arg, Asn, Asp, Cys, Gln, Glu, Gly, His, Ile, Leu, Lys, Met, Phe, Thr, Trp, Tyr, and Val. Under physiological conditions, a PAS peptide forms a random coil conformation and thereby can mediate an increased in vivo and/or in vitro stability to a recombinant protein of the invention, and has procoagulant activity.

Non-limiting examples of the PAS peptides include ASPAAPASPAPASPAPASPAP (SEQ ID NO: 10), AAPASPAPASPAPASPAPASP (SEQ ID NO: 11), APSSPSPASPSSPASPSS (SEQ ID NO: 12), APSSPSPASPSSPASPSS (SEQ ID NO: 13), SSPASPSSPASPSSPASPSS (SEQ ID NO: 14), AASPAASPASP-

PAAASPAAPSAPPA (SEQ ID NO: 15), ASAAAPAAASAAASAPSAAA (SEQ ID NO: 16) or any variants, derivatives, fragments, or combinations thereof. Additional examples of PAS sequences are known from, e.g., US Pat. Publ. No. 2010/0292130 A1 and PCT Appl. Publ. No. WO 2008/155134 A1. European issued patent EP2173890.

As discussed above, exemplary long-acting FVIII polypeptides also include Factor VIII fused to at least one transferrin peptide or fragment, variant, or derivative thereof. At least one transferrin peptide can be fused to either the N-terminal end of FVIII or to the C-terminal end of FVIII or inserted between two amino acids in FVIII. Any transferrin can be fused to or inserted into a recombinant FVIII protein of the invention. As an example, wild-type human Tf (Tf) is a 679 amino acid protein, of approximately 75 KDa (not accounting for glycosylation), with two main domains, N (about 330 amino acids) and C (about 340 amino acids), which appear to originate from a gene duplication. See GenBank accession numbers NM001063, XM002793, M12530, XM039845, XM 039847 and S95936 (www.ncbi.nlm.nih.gov), all of which are herein incorporated by reference in their entirety.

Transferrin transports iron through transferrin receptor (TfR)-mediated endocytosis. After the iron is released into an endosomal compartment and Tf-TfR complex is recycled to cell surface, the Tf is released back extracellular space for next cycle of iron transporting. Tf possesses a long half-life that is in excess of 14-17 days (Li et al., *Trends Pharmacol. Sci.* 23:206-209 (2002)). Transferrin fusion proteins have been studied for half-life extension, targeted deliver for cancer therapies, oral delivery and sustained activation of proinsulin (Brandsma et al., *Biotechnol. Adv.*, 29: 230-238 (2011); Bai et al., *Proc. Natl. Acad. Sci. USA* 102:7292-7296 (2005); Kim et al., *J. Pharmacol. Exp. Ther.*, 334:682-692 (2010); Wang et al., *J. Controlled Release* 155:386-392 (2011)).

As discussed above, exemplary long-acting FVIII polypeptides also include Factor VIII fused to at least one polyethylene glycol (PEG) moieties.

PEGylated FVIII can refer to a conjugate formed between FVIII and at least one polyethylene glycol (PEG) molecule. PEG is commercially available in a large variety of molecular weights and average molecular weight ranges. Typical examples of PEG average molecular weight ranges include, but are not limited to, about 200, about 300, about 400, about 600, about 1000, about 1300-1600, about 1450, about 2000, about 3000, about 3000-3750, about 3350, about 3000-7000, about 3500-4500, about 5000-7000, about 7000-9000, about 8000, about 10000, about 8500-11500, about 16000-24000, about 35000, about 40000, about 60000, and about 80000 daltons. These average molecular weights are provided merely as examples and are not meant to be limiting in any way.

A long-acting FVIII polypeptide useful for the invention can be PEGylated to include mono- or poly-(e.g., 2-4) PEG moieties. PEGylation can be carried out by any of the PEGylation reactions known in the art. Methods for preparing a PEGylated protein product will generally include (i) reacting a polypeptide with polyethylene glycol (such as a reactive ester or aldehyde derivative of PEG) under conditions whereby the peptide of the invention becomes attached to one or more PEG groups; and (ii) obtaining the reaction product(s). In general, the optimal reaction conditions for the reactions will be determined case by case based on known parameters and the desired result.

There are a number of PEG attachment methods available to those skilled in the art, for example Malik F et al., *Exp. Hematol.* 20:1028-35 (1992); Francis, *Focus on Growth Factors* 3(2):4-10 (1992); European Pat. Pub. Nos. EP0401384, EP0154316, and EP0401384; and International Pat. Appl. Pub. Nos. WO92/16221 and WO95/34326. As a non-limiting example, FVIII variants can contain cysteine substitutions in one or more insertion sites in FVIII, and the cysteines can be further conjugated to PEG polymer. See Mei et al., *Blood* 116:270-279 (2010) and U.S. Pat. No. 7,632,921, which are incorporated herein by reference in their entireties.

As discussed above, exemplary long-acting FVIII polypeptides also include Factor VIII fused to at least one hydroxyethyl starch (HES) polymer. HES is a derivative of naturally occurring amylopectin and is degraded by alpha-amylase in the body. HES exhibits advantageous biological properties and is used as a blood volume replacement agent and in hemodilution therapy in the clinics. See, e.g., Sommermeyer et al., *Krankenhauspharmazie* 8:271-278 (1987); and Weidler et al., *Arzneim.-Forschung/Drug Res.* 41: 494-498 (1991).

HES is mainly characterized by the molecular weight distribution and the degree of substitution. HES has a mean molecular weight (weight mean) of from 1 to 300 kD, from 2 to 200 kD, from 3 to 100 kD, or from 4 to 70 kD. Hydroxyethyl starch can further exhibit a molar degree of substitution of from 0.1 to 3, from 0.1 to 2, from 0.1 to 0.9, or from 0.1 to 0.8, and a ratio between C2:C6 substitution in the range of from 2 to 20 with respect to the hydroxyethyl groups. HES with a mean molecular weight of about 130 kD is VOLUVEN® from Fresenius. VOLUVEN® is an artificial colloid, employed, e.g., for volume replacement used in the therapeutic indication for therapy and prophylaxis of hypovolemia. There are a number of HES attachment methods available to those skilled in the art, e.g., the same PEG attachment methods described above.

In some embodiments, a long-acting FVIII polypeptide comprising a FVIII portion has an increased half-life (tin) over a polypeptide consisting of the same FVIII portion without the non-FVIII portion. A long-acting FVIII polypeptide with an increased tin can be referred to herein as a long-lasting FVIII. Long-acting chimeric Factor VIII polypeptides include, e.g., Factor VIII fused to Fc (including, e.g., chimeric Factor VIII polypeptides in the form of a hybrid such as a FVIII/Fc monomer dimer hybrid; see Example, Table 11A; and U.S. Pat. Nos. 7,404,956 and 7,348,004), and Factor VIII fused to albumin.

The Factor VIII polypeptide as used herein is functional factor VIII polypeptide in its normal role in coagulation, unless otherwise specified. Thus, the term Factor VIII includes variant polypeptides that are functional. Factor VIII proteins can be the human, porcine, canine, and murine factor VIII proteins. The full length polypeptide and polynucleotide sequences are known, as are many functional fragments, mutants and modified versions. Examples of human factor VIII sequences are shown as subsequences in SEQ ID NOs: 2 or 6 (Table 11). Factor VIII polypeptides include, e.g., full-length factor VIII, full-length factor VII minus Met at the N-terminus, mature factor VIII (minus the signal sequence), mature factor VIII with an additional Met at the N-terminus, and/or factor VIII with a full or partial deletion of the B domain. Factor VIII variants include B domain deletions, whether partial or full deletions.

A great many functional factor VIII variants are known, as is discussed above and below. In addition, hundreds of nonfunctional mutations in factor VIII have been identified

in hemophilia patients, and it has been determined that the effect of these mutations on factor VIII function is due more to where they lie within the 3-dimensional structure of factor VIII than on the nature of the substitution (Cutler et al., *Hum. Mutat.* 19:274-8 (2002)), incorporated herein by reference in its entirety. In addition, comparisons between factor VIII from humans and other species have identified conserved residues that are likely to be required for function (Cameron et al., *Thromb. Haemost.* 79:317-22 (1998); U.S. Pat. No. 6,251,632), incorporated herein by reference in its entirety.

The human factor VIII gene was isolated and expressed in mammalian cells (Toole, J. J., et al., *Nature* 312:342-347 (1984); Gitschier, J., et al., *Nature* 312:326-330 (1984); Wood, W. L., et al., *Nature* 312:330-337 (1984); Vehar, G. A., et al., *Nature* 312:337-342 (1984); WO 87/04187; WO 88/08035; WO 88/03558; U.S. Pat. No. 4,757,006), each of which is incorporated herein by reference in its entirety, and the amino acid sequence was deduced from cDNA. Capon et al., U.S. Pat. No. 4,965,199, incorporated herein by reference in its entirety, discloses a recombinant DNA method for producing factor VIII in mammalian host cells and purification of human factor VIII. Human factor VIII expression in CHO (Chinese hamster ovary) cells and BHK (baby hamster kidney cells) has been reported. Human factor VIII has been modified to delete part or all of the B domain (U.S. Pat. Nos. 4,994,371 and 4,868,112, each of which is incorporated herein by reference in its entirety), and replacement of the human factor VIII B domain with the human factor V B domain has been performed (U.S. Pat. No. 5,004,803, incorporated herein by reference in its entirety). The cDNA sequence encoding human factor VIII and predicted amino acid sequence are shown in SEQ ID NOs:1 and 2, respectively, of US Application Publ. No. 2005/0100990, incorporated herein by reference in its entirety.

U.S. Pat. No. 5,859,204, Lollar, J. S., incorporated herein by reference in its entirety, reports functional mutants of factor VIII having reduced antigenicity and reduced immunoreactivity. U.S. Pat. No. 6,376,463, Lollar, J. S., incorporated herein by reference in its entirety, also reports mutants of factor VIII having reduced immunoreactivity. US Application Publ. No. 2005/0100990, Saenko et al., incorporated herein by reference in its entirety, reports functional mutations in the A2 domain of factor VIII.

A number of functional factor VIII molecules, including B-domain deletions, are disclosed in the following patents U.S. Pat. Nos. 6,316,226 and 6,346,513, both assigned to Baxter; U.S. Pat. No. 7,041,635 assigned to In2Gen; U.S. Pat. Nos. 5,789,203, 6,060,447, 5,595,886, and 6,228,620 assigned to Chiron; U.S. Pat. Nos. 5,972,885 and 6,048,720 assigned to Biovitrum, U.S. Pat. Nos. 5,543,502 and 5,610,278 assigned to Novo Nordisk; U.S. Pat. No. 5,171,844 assigned to Immuno Ag; U.S. Pat. No. 5,112,950 assigned to Transgene S.A.; U.S. Pat. No. 4,868,112 assigned to Genetics Institute, each of which is incorporated herein by reference in its entirety.

The porcine factor VIII sequence is published, (Toole, J. J., et al., *Proc. Natl. Acad. Sci. USA* 83:5939-5942 (1986)), incorporated herein by reference in its entirety, and the complete porcine cDNA sequence obtained from PCR amplification of factor VIII sequences from a pig spleen cDNA library has been reported (Healey, J. F. et al., *Blood* 88:4209-4214 (1996), incorporated herein by reference in its entirety). Hybrid human/porcine factor VIII having substitutions of all domains, all subunits, and specific amino acid sequences were disclosed in U.S. Pat. No. 5,364,771 by Lollar and Runge, and in WO 93/20093, incorporated herein

by reference in its entirety. More recently, the nucleotide and corresponding amino acid sequences of the A1 and A2 domains of porcine factor VIII and a chimeric factor VIII with porcine A1 and/or A2 domains substituted for the corresponding human domains were reported in WO 94/11503, incorporated herein by reference in its entirety. U.S. Pat. No. 5,859,204, Lollar, J. S., also discloses the porcine cDNA and deduced amino acid sequences. U.S. Pat. No. 6,458,563, incorporated herein by reference in its entirety assigned to Emory discloses a B-domain deleted porcine Factor VIII.

A "B domain" of Factor VIII, as used herein, is the same as the B domain known in the art that is defined by internal amino acid sequence identity and sites of proteolytic cleavage by thrombin, e.g., residues Ser741-Arg1648 of full length mature human factor VIII. The other human factor VIII domains are defined by the following amino acid residues: A1, residues Ala1-Arg372; A2, residues Ser373-Arg740; A3, residues Ser1690-Ile2032; C1, residues Arg2033-Asn2172; C2, residues Ser2173-Tyr2332. The A3-C1-C2 sequence includes residues Ser1690-Tyr2332. The remaining sequence, residues Glu1649-Arg1689, is usually referred to as the factor VIII light chain activation peptide. The locations of the boundaries for all of the domains, including the B domains, for porcine, mouse and canine factor VIII are also known in the art. In one embodiment, the B domain of Factor VIII is deleted ("B domain deleted factor VIII" or "BDD FVIII"). An example of a BDD FVIII is REFACTO® (recombinant BDD FVIII), which has the sequence as the Factor VIII portion of the sequence in Table 1A(i) (amino acids 1 to 1457 or 20 to 1457 of SEQ ID NO:2). In another embodiment, the B domain deleted Factor VIII contains an intact intracellular processing site, which corresponds to Arginine at residue 754 of B domain deleted Factor VIII, which corresponds to Arginine residue 773 of SEQ ID NO: 2, or residue 1648 of full-length Factor VIII, which corresponds to Arginine residue 1667 of SEQ ID NO: 6. The sequence residue numbers used herein without referring to any SEQ ID Numbers correspond to the Factor VIII sequence without the signal peptide sequence (19 amino acids) unless otherwise indicated. For example, S743/Q1638 of full-length Factor VIII corresponds to S762/Q1657 of SEQ ID NO: 6 due to the 19 amino acid signal peptide sequence. In other embodiments, the B domain deleted FVIII comprises a substitution or mutation at an amino acid position corresponding to Arginine 1645, a substitution or mutation at an amino acid position corresponding to Arginine 1648, or a substitution or mutation at amino acid positions corresponding to Arginine 1645 and Arginine 1648 in full-length Factor VIII. In some embodiments, the amino acid substituted at the amino acid position corresponding to Arginine 1645 is a different amino acid from the amino acid substituted at the amino acid position corresponding to Arginine 1648. In certain embodiments, the substitution or mutation is an amino acid other than arginine, e.g., alanine.

A "B domain deleted factor VIII" can have the full or partial deletions disclosed in U.S. Pat. Nos. 6,316,226, 6,346,513, 7,041,635, 5,789,203, 6,060,447, 5,595,886, 6,228,620, 5,972,885, 6,048,720, 5,543,502, 5,610,278, 5,171,844, 5,112,950, 4,868,112, and 6,458,563, each of which is incorporated herein by reference in its entirety. In some embodiments, a B domain deleted factor VIII sequence of the present invention comprises any one of the deletions disclosed at col. 4, line 4 to col. 5, line 28 and examples 1-5 of U.S. Pat. No. 6,316,226 (also in U.S. Pat. No. 6,346,513). In some embodiments, a B domain deleted factor VIII of the present invention has a deletion disclosed

at col. 2, lines 26-51 and examples 5-8 of U.S. Pat. No. 5,789,203 (also U.S. Pat. Nos. 6,060,447, 5,595,886, and 6,228,620). In some embodiments, a B domain deleted factor VIII has a deletion described in col. 1, lines 25 to col. 2, line 40 of U.S. Pat. No. 5,972,885; col. 6, lines 1-22 and example 1 of U.S. Pat. No. 6,048,720; col. 2, lines 17-46 of U.S. Pat. No. 5,543,502; col. 4, line 22 to col. 5, line 36 of U.S. Pat. No. 5,171,844; col. 2, lines 55-68, FIG. 2, and example 1 of U.S. Pat. No. 5,112,950; col. 2, line 2 to col. 19, line 21 and Table 2 of U.S. Pat. No. 4,868,112; col. 2, line 1 to col. 3, line 19, col. 3, line 40 to col. 4, line 67, col. 7, line 43 to col. 8, line 26, and col. 11, line 5 to col. 13, line 39 of U.S. Pat. No. 7,041,635; or col. 4, lines 25-53, of U.S. Pat. No. 6,458,563. In some embodiments, a B domain deleted factor VIII has a deletion of most of the B domain, but still contains amino-terminal sequences of the B domain that are essential for in vivo proteolytic processing of the primary translation product into two polypeptide chain (i.e., intracellular processing site), as disclosed in WO 91/09122, which is incorporated herein by reference in its entirety. In some embodiments, a B domain deleted factor VIII is constructed with a deletion of amino acids 747-1638, i.e., virtually a complete deletion of the B domain. Hoebein R. C., et al. *J. Biol. Chem.* 265 (13): 7318-7323 (1990), incorporated herein by reference in its entirety. A B domain deleted factor VIII can also contain a deletion of amino acids 771-1666 or amino acids 868-1562 of factor VIII. Meulien P., et al. *Protein Eng.* 2(4): 301-6 (1988), incorporated herein by reference in its entirety. Additional B domain deletions that are part of the invention include, e.g., deletion of amino acids 982 through 1562 or 760 through 1639 (Toole et al., *Proc. Natl. Acad. Sci. U.S.A.* 83:5939-5942 (1986)), 797 through 1562 (Eaton et al., *Biochemistry* 25:8343-8347 (1986)), 741 through 1646 (Kaufman (PCT published application No. WO 87/04187)), 747-1560 (Sarver et al., *DNA* 6:553-564 (1987)), 741 through 1648 (Pasek (PCT application No. 88/00831)), 816 through 1598 or 741 through 1689 (Lagner (Behring Inst. Mitt. (1988) No 82:16-25, EP 295597)), each of which is incorporated herein by reference in its entirety. In some embodiments, B domain deleted FVIII comprises a partial deletion in B domain, i.e., having 21 amino acids from B domain (i.e., SFSQNSRHP-SQNPPVLKRHQ, which is SEQ ID NO: 17) disclosed in US Publication No. 20100286067 and US Publication No. US 20120093840, both of which are incorporated herein by reference in their entireties. Each of the foregoing deletions can be made in any Factor VIII sequence.

In one embodiment, the B domain deleted Factor VIII portion in a FVIII polypeptide is processed into two chains connected (or associated) by a metal bond, the first chain comprising a heavy chain (A1-A2-partial B) and a second chain comprising a light chain (A3-C1-C2). In another embodiment, the B domain deleted Factor VIII portion is a single chain Factor VIII. The single chain Factor VIII can comprise an intracellular processing site, which corresponds to Arginine at residue 754 of B domain deleted Factor VIII (residue 773 of SEQ ID NO: 2) or at residue 1648 of full-length Factor VIII (residue 1657 of SEQ ID NO: 6).

The metal bond between the heavy chain and the light chain can be any metal known in the art. For example, the metals useful for the invention can be a divalent metal ion. The metals that can be used to associate the heavy chain and light chain include, but not limited to, Ca^{2+} , Mn^{2+} , or Cu^{2+} . Fatouros et al., *Intern. J. Pharm.* 155(1): 121-131 (1997); Wakabayashi et al., *JBC.* 279(13): 12677-12684 (2004).

A FVIII polypeptide used herein can comprise processed Factor VIII or single chain Factor VIII or a combination

thereof. "Processed Factor VIII," as used herein means Factor VIII that has been cleaved at Arginine 1648 (for full-length Factor VIII) or Arginine 754 (for B-domain deleted Factor VIII), i.e., intracellular processing site. Due to the cleavage at the intracellular processing site, processed Factor VIII comprises two polypeptide chains, the first chain being a heavy chain and the second chain being a light chain. For example, the processed Factor VIII-Fc fusion protein (i.e., Heavy chain and Light chain fused to Fc) run at approximately 90 kDa and 130 kDa on a non-reducing SDS-PAGE, respectively, and 90 kDa and 105 kDa on a reducing SDS-PAGE, respectively. Therefore, in one embodiment, at least about 50%, about 60%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100% of the Factor VIII portion in the long-acting polypeptide or in the short-acting FVIII polypeptide is processed Factor VIII. In another embodiment, about 50%, about 60%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100% of the Factor VIII portion in the long-acting polypeptide or in the short-acting FVIII polypeptide is processed Factor VIII. In a particular embodiment, the FVIII polypeptide comprising processed Factor VIII is purified (or isolated) from the polypeptide comprising single chain Factor VIII, and at least about 75%, about 80%, about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100% of the Factor VIII portion in the long-acting polypeptide or in the short-acting FVIII polypeptide is processed Factor VIII. In some embodiments, the FVIII polypeptide comprises about 15% to 25% of single chain FVIII polypeptide and about 75% to about 85% of processed FVIII polypeptide.

"Single chain Factor VIII," "SC Factor VIII," or "SCFVIII" as used herein means Factor VIII that has not been cleaved at the Arginine site (residue 1648 for full-length Factor VIII (i.e., residue 1667 of SEQ ID NO: 6) or residue 754 for B-domain deleted Factor VIII (i.e., residue 773 of SEQ ID NO: 2). Therefore, single chain Factor VIII in the FVIII polypeptide used herein comprises a single chain. In one embodiment, the single chain Factor VIII contains an intact intracellular processing site. In another embodiment, the single chain Factor VIII of the invention comprises a substitution or mutation at an amino acid position corresponding to Arginine 1645, a substitution or mutation at an amino acid position corresponding to Arginine 1648, or a substitution or mutation at amino acid positions corresponding to Arginine 1645 and Arginine 1648 in full-length Factor VIII. In other embodiments, the amino acid substituted at the amino acid position corresponding to Arginine 1645 is a different amino acid from the amino acid substituted at the amino acid position corresponding to Arginine 1648. In certain embodiments, the substitution or mutation is an amino acid other than arginine, e.g., isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophan, valine, alanine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, proline, selenocysteine, serine, tyrosine, histidine, ornithine, pyrrolysine, or taurine. The single chain Factor VIII-Fc fusion protein can run at approximately 220 kDa on a non reducing SDS-PAGE and at approximately 195 kDa on a reducing SDS-PAGE.

The Factor VIII portion in the FVIII polypeptide used herein has Factor VIII activity. Factor VIII activity can be measured by any known methods in the art. For example, one of those methods can be a chromogenic assay. The chromogenic assay mechanism is based on the principles of the blood coagulation cascade, where activated Factor VIII

accelerates the conversion of Factor X into Factor Xa in the presence of activated Factor IX, phospholipids and calcium ions. The Factor Xa activity is assessed by hydrolysis of a p-nitroanilide (pNA) substrate specific to Factor Xa. The initial rate of release of p-nitroaniline measured at 405 nM is directly proportional to the Factor Xa activity and thus to the Factor VIII activity in the sample. The chromogenic assay is recommended by the Factor VIII and Factor IX Subcommittee of the Scientific and Standardization Committee (SSC) of the International Society on Thrombosis and Hemostasis (ISTH). Since 1994, the chromogenic assay has also been the reference method of the European Pharmacopoeia for the assignment of FVIII concentrate potency. Thus, in one embodiment, the FVIII polypeptide comprising single chain Factor VIII has Factor VIII activity comparable to a FVIII polypeptide comprising processed Factor VIII, when the Factor VIII activity is measured in vitro by a chromogenic assay.

In certain embodiment, a long-acting FVIII polypeptide is a FVIII monomer-dimer hybrid. To obtain a FVIII monomer-dimer hybrid, the coding sequence of human recombinant B-domain deleted FVIII was obtained by reverse transcription-polymerase chain reaction (RT-PCR) from human liver poly A RNA (Clontech) using FVIII-specific primers. The FVIII sequence includes the native signal sequence for FVIII. The B-domain deletion was from serine 743 (S743; 2287 bp) to glutamine 1638 (Q1638; 4969 bp) for a total deletion of 2682 bp. Then, the coding sequence for human recombinant Fc was obtained by RT-PCR from a human leukocyte cDNA library (Clontech) using Fc specific primers. Primers were designed such that the B-domain deleted FVIII sequence was fused directly to the N-terminus of the Fc sequence with no intervening linker. The FVIII-Fc DNA sequence was cloned into the mammalian dual expression vector pBUDCE4.1 (Invitrogen) under control of the CMV promoter. A second identical Fc sequence including the mouse Igk signal sequence was obtained by RT-PCR and cloned downstream of the second promoter, EF1 α , in the expression vector pBUDCE4.1.

In one embodiment, an FcRn binding partner can be an Fc region. An FcRn binding partner is any molecule that can be specifically bound by the FcRn receptor with consequent active transport by the FcRn receptor of the FcRn binding partner. Thus, the term Fc includes any variants of IgG Fc that are functional. The region of the Fc portion of IgG that binds to the FcRn receptor has been described based on X-ray crystallography (Burmeister et al., *Nature* 372:379 (1994), incorporated herein by reference in its entirety). The major contact area of the Fc with the FcRn is near the junction of the CH2 and CH3 domains. Fc-FcRn contacts are all within a single Ig heavy chain. The FcRn binding partners include, e.g., whole IgG, the Fc fragment of IgG, and other fragments of IgG that include the complete binding region of FcRn. The major contact sites include amino acid residues 248, 250-257, 272, 285, 288, 290-291, 308-311, and 314 of the CH2 domain and amino acid residues 385-387, 428, and 433-436 of the CH3 domain. References made to amino acid numbering of immunoglobulins or immunoglobulin fragments, or regions, are all based on Kabat et al. 1991, *Sequences of Proteins of Immunological Interest*, U. S. Department of Public Health, Bethesda; MD, incorporated herein by reference in its entirety. (The FcRn receptor has been isolated from several mammalian species including humans. The sequences of the human FcRn, rat FcRn, and mouse FcRn are known (Story et al., *J. Exp. Med.* 180: 2377 (1994), incorporated herein by reference in its entirety.) An Fc can comprise the CH2 and CH3 domains of an immu-

noglobulin with or without the hinge region of the immunoglobulin. Exemplary Fc variants are provided in WO 2004/101740 and WO 2006/074199, incorporated herein by reference in its entirety.

Fc (or Fc portion of a chimeric polypeptide) can contain one or more mutations, and combinations of mutations.

Fc (or Fc portion of a chimeric polypeptide) can contain mutations conferring increased half-life such as M252Y, S254T, T256E, and combinations thereof, as disclosed in Oganessian et al., *Mol. Immunol.* 46:1750 (2009), which is incorporated herein by reference in its entirety; H433K, N434F, and combinations thereof, as disclosed in Vaccaro et al., *Nat. Biotechnol.* 23:1283 (2005), which is incorporated herein by reference in its entirety; the mutants disclosed at pages 1-2, paragraph [0012], and Examples 9 and 10 of US 2009/0264627 A1, which is incorporated herein by reference in its entirety; and the mutants disclosed at page 2, paragraphs [0014] to [0021] of US 20090163699 A1, which is incorporated herein by reference in its entirety.

Fc (or Fc portion of a chimeric polypeptide) can also include, e.g., the following mutations: The Fc region of IgG can be modified according to well recognized procedures such as site directed mutagenesis and the like to yield modified IgG or Fc fragments or portions thereof that will be bound by FcRn. Such modifications include, e.g., modifications remote from the FcRn contact sites as well as modifications within the contact sites that preserve or even enhance binding to the FcRn. For example the following single amino acid residues in human IgG1 Fc (Fcγ1) can be substituted without significant loss of Fc binding affinity for FcRn: P238A, S239A, K246A, K248A, D249A, M252A, T256A, E258A, T260A, D265A, S267A, H268A, E269A, D270A, E272A, L274A, N276A, Y278A, D280A, V282A, E283A, H285A, N286A, T289A, K290A, R292A, E293A, E294A, Q295A, Y296F, N297A, S298A, Y300F, R301A, V303A, V305A, T307A, L309A, Q311A, D312A, N315A, K317A, E318A, K320A, K322A, S324A, K326A, A327Q, P329A, A330Q, A330S, P331A, P331S, E333A, K334A, T335A, S337A, K338A, K340A, Q342A, R344A, E345A, Q347A, R355A, E356A, M358A, T359A, K360A, N361A, Q362A, Y373A, S375A D376A, A378Q, E380A, E382A, S383A, N384A, Q386A, E388A, N389A, N390A, Y391F, K392A, L398A, S400A, D401A, D413A, K414A, R416A, Q418A, Q419A, N421A, V422A, S424A, E430A, N434A, T437A, Q438A, K439A, S440A, S444A, and K447A, where for example P238A represents wildtype proline substituted by alanine at position number 238. In addition to alanine other amino acids can be substituted for the wildtype amino acids at the positions specified above. Mutations can be introduced singly into Fc giving rise to more than one hundred FcRn binding partners distinct from native Fc. Additionally, combinations of two, three, or more of these individual mutations can be introduced together, giving rise to hundreds more FcRn binding partners. Certain of these mutations can confer new functionality upon the FcRn binding partner. For example, one embodiment incorporates N297A, removing a highly conserved N-glycosylation site. The effect of this mutation is to reduce immunogenicity, thereby enhancing circulating half-life of the FcRn binding partner, and to render the FcRn binding partner incapable of binding to FcγRI, FcγRIIA, FcγRIIB, and FcγRIIIA, without compromising affinity for FcRn (Routledge et al. 1995, Transplantation 60:847, which is incorporated herein by reference in its entirety; Friend et al. 1999, Transplantation 68:1632, which is incorporated herein by reference in its entirety; Shields et al. 1995, *J. Biol. Chem.* 276:6591, which is incorporated herein by reference in its entirety). Addi-

tionally, at least three human Fc gamma receptors appear to recognize a binding site on IgG within the lower hinge region, generally amino acids 234-237. Therefore, another example of new functionality and potential decreased immunogenicity can arise from mutations of this region, as for example by replacing amino acids 233-236 of human IgG1 "ELLG" to the corresponding sequence from IgG2 "PVA" (with one amino acid deletion). It has been shown that FcγRI, FcγRII, and FcγRIII which mediate various effector functions will not bind to IgG when such mutations have been introduced (Ward and Ghetie, *Therapeutic Immunology* 2:77 (1995), which is incorporated herein by reference in its entirety; and Armour et al., *Eur. J. Immunol.* 29:2613 (1999), which is incorporated herein by reference in its entirety). As a further example of new functionality arising from mutations described above affinity for FcRn can be increased beyond that of wild type in some instances. This increased affinity can reflect an increased "on" rate, a decreased "off" rate or both an increased "on" rate and a decreased "off" rate. Mutations believed to impart an increased affinity for FcRn include, e.g., T256A, T307A, E380A, and N434A (Shields et al., *J. Biol. Chem.* 276:6591 (2001), which is incorporated herein by reference in its entirety).

The Fc (or Fc portion of a chimeric polypeptide) can be at least 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the Fc amino acid sequence shown in Table 11 (amino acids 1458 to 1684 of SEQ ID NO:2 or amino acids 2352 to 2578 of SEQ ID NO:6). The Fc (or Fc portion of a chimeric polypeptide) can be identical to the Fc amino acid sequence shown in Table 11 (amino acids 1458 to 1684 of SEQ ID NO:2 and amino acids 2352 to 2578 of SEQ ID NO:6).

As a practical matter, whether any particular nucleic acid molecule or polypeptide is at least 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to a nucleotide sequence or polypeptide of the present invention can be determined conventionally using known computer programs. In one embodiment, a method for determining the best overall match between a query sequence (reference or original sequence) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al., *Comp. App. Biosci.* 6:237-245 (1990), which is herein incorporated by reference in its entirety. In a sequence alignment the query and subject sequences are both DNA sequences. An RNA sequence can be compared by converting U's to T's. The result of said global sequence alignment is in percent identity. In another embodiment, parameters used in a FASTDB alignment of DNA sequences to calculate percent identity are: Matrix=Unitary, k-tuple=4, Mismatch Penalty=1, Joining Penalty=30, Randomization Group Length=0, Cutoff Score=1, Gap Penalty=5, Gap Size Penalty 0.05, Window Size=500 or the length of the subject nucleotide sequence, whichever is shorter.

If the subject sequence is shorter than the query sequence because of 5' or 3' deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for 5' and 3' truncations of the subject sequence when calculating percent identity. For subject sequences truncated at the 5' or 3' ends, relative to the query sequence, the percent identity is corrected by calculating the number of bases of the query sequence that are 5' and 3' of the subject sequence, which are not matched/aligned, as a percent of the total bases of the query sequence. Whether a nucleotide is matched/aligned is determined by results of the FASTDB sequence alignment.

This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This corrected score is what is used for the purposes of the present invention. Only bases outside the 5' and 3' bases of the subject sequence, as displayed by the FASTDB alignment, which are not matched/aligned with the query sequence, are calculated for the purposes of manually adjusting the percent identity score.

For example, a 90 base subject sequence is aligned to a 100 base query sequence to determine percent identity. The deletions occur at the 5' end of the subject sequence and therefore, the FASTDB alignment does not show a matched/alignment of the first 10 bases at 5' end. The 10 unpaired bases represent 10% of the sequence (number of bases at the 5' and 3' ends not matched/total number of bases in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 bases were perfectly matched the final percent identity would be 90%. In another example, a 90 base subject sequence is compared with a 100 base query sequence. This time the deletions are internal deletions so that there are no bases on the 5' or 3' of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only bases 5' and 3' of the subject sequence which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to be made for the purposes of the present invention.

By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to a query amino acid sequence of the present invention, it is intended that the amino acid sequence of the subject polypeptide is identical to the query sequence except that the subject polypeptide sequence can include up to five amino acid alterations per each 100 amino acids of the query amino acid sequence. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a query amino acid sequence, up to 5% of the amino acid residues in the subject sequence can be inserted, deleted, (indels) or substituted with another amino acid. These alterations of the reference sequence can occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular polypeptide is at least 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, the amino acid sequences of SEQ ID NO:2 (the factor VIII portion, the Fc portion, individually or together) or 4, or a known factor VIII or Fc polypeptide sequence, can be determined conventionally using known computer programs. In one embodiment, a method for determining the best overall match between a query sequence (reference or original sequence) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al., *Comp. App. Biosci.* 6:237-245(1990), incorporated herein by reference in its entirety. In a sequence alignment the query and subject sequences are either both nucleotide sequences or both amino acid sequences. The result of said global sequence alignment is in percent identity. In another embodiment, parameters used in a FASTDB amino acid alignment are: Matrix=PAM 0, k-tuple=2, Mismatch Penalty=1, Joining Penalty=20, Randomization Group Length=0, Cutoff Score=1, Window Size=sequence length, Gap Penalty=5,

Gap Size Penalty=0.05, Window Size=500 or the length of the subject amino acid sequence, whichever is shorter.

If the subject sequence is shorter than the query sequence due to N- or C-terminal deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for N- and C-terminal truncations of the subject sequence when calculating global percent identity. For subject sequences truncated at the N- and C-termini, relative to the query sequence, the percent identity is corrected by calculating the number of residues of the query sequence that are N- and C-terminal of the subject sequence, which are not matched/aligned with a corresponding subject residue, as a percent of the total bases of the query sequence. Whether a residue is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This final percent identity score is what is used for the purposes of the present invention. Only residues to the N- and C-termini of the subject sequence, which are not matched/aligned with the query sequence, are considered for the purposes of manually adjusting the percent identity score. That is, only query residue positions outside the farthest N- and C-terminal residues of the subject sequence.

For example, a 90 amino acid residue subject sequence is aligned with a 100 residue query sequence to determine percent identity. The deletion occurs at the N-terminus of the subject sequence and therefore, the FASTDB alignment does not show a matching/alignment of the first 10 residues at the N-terminus. The 10 unpaired residues represent 10% of the sequence (number of residues at the N- and C-termini not matched/total number of residues in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 residues were perfectly matched the final percent identity would be 90%. In another example, a 90 residue subject sequence is compared with a 100 residue query sequence. This time the deletions are internal deletions so there are no residues at the N- or C-termini of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only residue positions outside the N- and C-terminal ends of the subject sequence, as displayed in the FASTDB alignment, which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to be made for the purposes of the present invention.

The polynucleotide variants can contain alterations in the coding regions, non-coding regions, or both. In one embodiment, the polynucleotide variants contain alterations which produce silent substitutions, additions, or deletions, but do not alter the properties or activities of the encoded polypeptide. In another embodiment, nucleotide variants are produced by silent substitutions due to the degeneracy of the genetic code. In other embodiments, variants in which 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination. Polynucleotide variants can be produced for a variety of reasons, e.g., to optimize codon expression for a particular host (change codons in the human mRNA to others, e.g., a bacterial host such as *E. coli*).

Naturally occurring variants are called "allelic variants," and refer to one of several alternate forms of a gene occupying a given locus on a chromosome of an organism (Genes II, Lewin, B., ed., John Wiley & Sons, New York (1985)). These allelic variants can vary at either the poly-

nucleotide and/or polypeptide level and are included in the present invention. Alternatively, non-naturally occurring variants can be produced by mutagenesis techniques or by direct synthesis.

Using known methods of protein engineering and recombinant DNA technology, variants can be generated to improve or alter the characteristics of the polypeptides. For instance, one or more amino acids can be deleted from the N-terminus or C-terminus of the secreted protein without substantial loss of biological function. Ron et al., *J. Biol. Chem.* 268: 2984-2988 (1993), incorporated herein by reference in its entirety, reported variant KGF proteins having heparin binding activity even after deleting 3, 8, or 27 amino-terminal amino acid residues. Similarly, Interferon gamma exhibited up to ten times higher activity after deleting 8-10 amino acid residues from the carboxy terminus of this protein. (Dobeli et al., *J. Biotechnology* 7:199-216 (1988), incorporated herein by reference in its entirety.)

Moreover, ample evidence demonstrates that variants often retain a biological activity similar to that of the naturally occurring protein. For example, Gayle and coworkers (*J. Biol. Chem.* 268:22105-22111 (1993), incorporated herein by reference in its entirety) conducted extensive mutational analysis of human cytokine IL-1a. They used random mutagenesis to generate over 3,500 individual IL-1a mutants that averaged 2.5 amino acid changes per variant over the entire length of the molecule. Multiple mutations were examined at every possible amino acid position. The investigators found that "[m]ost of the molecule could be altered with little effect on either [binding or biological activity]." (See Abstract.) In fact, only 23 unique amino acid sequences, out of more than 3,500 nucleotide sequences examined, produced a protein that significantly differed in activity from wild-type.

As stated above, polypeptide variants include, e.g., modified polypeptides. Modifications include, e.g., acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, pegylation (Mei et al., *Blood* 116:270-79 (2010), which is incorporated herein by reference in its entirety), proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. In some embodiments, Factor VIII is modified, e.g., pegylated, at any convenient location. In some embodiments, Factor VIII is pegylated at a surface exposed amino acid of Factor VIII, e.g., a surface exposed cysteine, which can be an engineered cysteine. Id. In some embodiments, modified Factor VIII, e.g., pegylated Factor VIII, is a long-acting Factor VIII.

III. Method of Administering

Treatment of hemophilia A is a replacement therapy targeting restoration of FVIII activity to 1 to 5% of normal levels to prevent spontaneous bleeding (Mannucci, P. M. et al., *N. Engl. J. Med.* 344:1773-9 (2001), herein incorporated by reference in its entirety).

The present invention also includes methods of administering the pharmaceutical composition of the invention. The

present invention provides a method of administering a pharmaceutical composition to a human subject in need thereof (e.g., human patient), comprising administering to the subject a therapeutic dose of a pharmaceutical composition comprising a Factor VIII polypeptide, e.g., a long-acting FVIII polypeptide or a short-acting FVIII polypeptide, e.g., a Factor VIII-Fc polypeptide, or a hybrid of such a polypeptide at a dosing interval.

In some embodiments, the pharmaceutical composition of the invention is used to reduce or decrease one or more bleeding episodes or frequency (e.g., a bleeding condition) in a subject in need thereof. In other embodiments, the pharmaceutical composition of the invention is used to treat or prevent a bleeding condition in a subject in need thereof.

The bleeding condition can be caused by a blood coagulation disorder. A blood coagulation disorder can also be referred to as a coagulopathy. In one example, the blood coagulation disorder, which can be treated with a pharmaceutical composition of the current disclosure, is hemophilia. In another example, the blood coagulation disorder, which can be treated with a pharmaceutical composition of the present disclosure is hemophilia A.

In some embodiments, the type of bleeding associated with the bleeding condition is selected from hemarthrosis, muscle bleed, oral bleed, hemorrhage, hemorrhage into muscles, oral hemorrhage, trauma, trauma capitis, gastrointestinal bleeding, intracranial hemorrhage, intra-abdominal hemorrhage, intrathoracic hemorrhage, bone fracture, central nervous system bleeding, bleeding in the retropharyngeal space, bleeding in the retroperitoneal space, and bleeding in the iliopsoas sheath.

In other embodiments, the subject suffering from bleeding condition is in need of treatment for surgery, including, e.g., surgical prophylaxis or peri-operative management. In one example, the surgery is selected from minor surgery and major surgery. Exemplary surgical procedures include tooth extraction, tonsillectomy, inguinal herniotomy, synovectomy, craniotomy, osteosynthesis, trauma surgery, intracranial surgery, intra-abdominal surgery, intrathoracic surgery, joint replacement surgery (e.g., total knee replacement, hip replacement, and the like), heart surgery, and caesarean section.

The present invention also provides a method of reducing or decreasing an annualized bleeding rate (ABR) of a subject having hemophilia comprising administering to the subject an effective dose of a FVIII polypeptide. In one embodiment, a long-acting FVIII polypeptide is administered at a dosing interval of every three days or longer. In another embodiment, the effective dose is between about 20 IU/kg and about 90 IU/kg. In other embodiments, the effective dose is 20-30 IU/kg, 30-40 IU/kg, 40-50 IU/kg, 50-60 IU/kg, 60-70 IU/kg, 70-80 IU/kg, or 80-90 IU/kg. In yet other embodiments, the effective dose is 20 IU/kg, 25 IU/kg, 30 IU/kg, 35 IU/kg, 40 IU/kg, 45 IU/kg, 50 IU/kg, 55 IU/kg, 60 IU/kg, 65 IU/kg, 70 IU/kg, 75 IU/kg, 80 IU/kg, 85 IU/kg, or 90 IU/kg.

In certain embodiments, administration of a FVIII polypeptide is for individualized (tailored) prophylaxis and results in an ABR of less than about 5.5, less than about 5.4, less than about 5.3, less than about 5.2, less than about 5.1, less than about 5.0, less than about 4.9, less than about 4.8, less than about 4.7, less than about 4.6, or less than about 4.5. In other embodiments, the administration results in an ABR of about 4.7 to 0. In some embodiments, the median of the ABR is about 1.6. In yet other embodiments, the mean of the ABR is about 2.9. In one aspect of the individualized prophylaxis regimen, the effective dose is about 25 IU/kg to

about 65 IU/kg given every three to five days. For example, the effective dose is about 25 IU/kg to about 65 IU/kg given every three days. In another aspect, the effective dose is about 50 IU/kg to about 65 IU/kg every four days or every five days. In other aspects, the effective dose is up to 65 IU/kg every three days. In still other embodiments, the effective dose of a FVIII polypeptide is modified based on the patient's pharmacokinetic profile. In one example, the patient is administered with two doses initially, a first dose of 25 IU/kg on day 1 and a second dose of 50 IU/kg on day 4. If the pharmacokinetic data show less than 1% trough level of normal FVIII activity after the initial two doses, the patient is then administered with about 25 IU/kg to about 65 IU/kg every three days. In another example, if the pharmacokinetic data show higher than 1% trough level of normal FVIII activity for five days from the dosing, the patient is then administered with about 50 IU/kg to about 65 IU/kg every five days. If the patient experiences more than two spontaneous bleeds over eight weeks period after administration of about 50 IU/kg to about 65 IU/kg every five days, the patient is administered one of the followings: (1) up to about 65 IU/kg every three days (for the target trough level of up to 5% of normal), (2) about 50 IU/kg to about 65 IU/kg every four days (for the target trough level up to 5% of normal), or (3) about 25 IU/kg to about 65 IU/kg every three days (for the target trough level of 1% to 3% of normal). If the patient receiving about 50 IU/kg to about 65 IU/kg every four days still experiences more than two spontaneous bleeds, the maximum dose can go up to about 65 IU/kg every three days. In some examples, the Interquartile Range (IQR) of the ABR for the individualized prophylaxis regimen is 0 to 4.7.

In some embodiments, administration of a FVIII polypeptide is for weekly prophylaxis and results in an ABR of less than about 9.0, less than about 8.9, less than about 8.8, less than about 8.7, less than about 8.6, less than about 8.5, or less than about 8.4. In one example, the administration results in an ABR between about 8.4 and 0. In another example, the median of the ABR for weekly prophylaxis is about 3.6. In other examples, the mean of the ABR for weekly prophylaxis is about 8.8. In other embodiments, an effective dose for weekly prophylaxis is about 65 IU/kg once every week, e.g., every five days, every six days, every seven days, every eight days, or every nine days. In some examples, the IQR of the ABR for the weekly prophylaxis regimen is 1.9 to 8.4.

In other embodiments, administration of a FVIII polypeptide is for episodic or on-demand treatment and results in an ABR of less than about 55, less than about 54, less than about 53, less than about 52, less than about 51, less than about 50, less than about 49, less than about 48, or less than about 47. In one example, the administration results in an ABR between about 49 and 0, e.g., between 48.7 and 0. In another example, the median of the ABR for episodic (on-demand) treatment is about 33.5. In other examples, the mean of the ABR for episodic (on-demand) treatment is about 37.23. In other embodiments, an effective dose for on-demand treatment is about 10 IU/kg to 75 IU/kg every 12 to 24 hours. In some examples, the IQR of the ABR for the on-demand treatment is 21.1 to 48.7.

In some embodiments, the effective dose for individualized prophylaxis, weekly prophylaxis, or episodic treatment is a fixed dose or a stratified dose. In one aspect, the fixed dose is about 2,000 IU per dose, about 2,500 IU per dose, about 3,000 IU per dose, about 3,500 IU per dose, or about 4,000 IU per dose.

The dosing interval for a long-acting FVIII polypeptide can be at least about one and one-half to six times longer, one and one-half to five times longer, one and one-half to four times longer, one and one-half to three times longer, or one and one-half to two times longer, than the dosing interval required for an equivalent dose of the Factor VIII without the non-Factor VIII portion (a polypeptide consisting of the Factor VIII portion), e.g., without the Fc portion. The dosing interval can be at least about one and one-half, two, two and one-half, three, three and one-half, four, four and one-half, five, five and one-half or six times longer than the dosing interval required for an equivalent dose of the Factor VIII without the non-Factor VIII portion (a polypeptide consisting of the Factor VIII portion), e.g., without the Fc portion. The dosing interval can be about every three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, or fourteen days or longer.

The methods of the invention can be practiced on a subject in need of prophylactic treatment or on-demand treatment.

For on-demand treatment, the dosing interval of a pharmaceutical composition comprising a FVIII polypeptide is about once every 24-36, 24-48, 24-72, 24-96, 24-120, 24-144, 24-168, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, or 72 hours or longer.

In one embodiment, the subject is in need of on-demand (episodic) treatment. In another embodiment, on-demand (episodic) treatment resolves greater than 80% (greater than 80%, greater than 81%, greater than 82%, greater than 83%, greater than 84%, greater than 85%, greater than 86%, greater than 87%, greater than 88%, greater than 89%, greater than 90%, greater than 91%, greater than 92%, greater than 93%, greater than 94%, greater than 95%, greater than 96%, greater than 97%, greater than 98%, greater than 99%, or 100%) or 80-100%, 80-90%, 85-90%, 90-100%, 90-95%, or 95-100% of bleeds (e.g., spontaneous bleeds) in a single dose. In another embodiment, greater than 80% (greater than 81%, greater than 82%, greater than 83%, greater than 84%, greater than 85%, greater than 86%, greater than 87%, greater than 88%, greater than 89%, greater than 90%, greater than 91%, greater than 92%, greater than 93%, greater than 94%, greater than 95%, greater than 96%, greater than 97%, greater than 98%, or 100%) or 80-100%, 80-90%, 85-90%, 90-100%, 90-95%, or 95-100% of bleeding episodes are rated excellent or good by physicians after on-demand (episodic) treatment. In other embodiments, greater than 5%, (greater than 6%, greater than 7%, greater than 8%, greater than 9%, greater than 10%, greater than 11%, greater than 12%, greater than 13%, greater than 14%, greater than 15%, greater than 16%, greater than 17%, greater than 18%, greater than 19%, greater than 20%), or 5-20%, 5-15%, 5-10%, 10-20%, or 10-15% of bleeding episodes are rated as fair by physicians after on-demand treatment.

In other embodiments, the pharmaceutical composition comprising a FVIII polypeptide is used for prophylaxis. Prophylaxis can be demonstrated by better C_{max}, better T_{max}, and/or greater mean residence time versus short-acting FVIII. In some embodiments, prophylaxis results in no spontaneous bleeding episodes within about 24, 36, 48, 72, or 96 hours (e.g., 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 96, 87, 88, 89, 90, 91, 92, 93, 94, 95, or 96 hours),

after injection (e.g., the last injection). In certain embodiments, prophylaxis results in greater than 30% (e.g., greater than 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 96, 87, 88, 89, or 90%, for example, greater than 50%), mean reduction in annualized bleeding episodes with once weekly dosing (e.g., at 65 IU/kg).

In another example, the subject is concomitantly treated with FIX. Because the compounds of the invention are capable of activating FIXa, they could be used to pre-activate the FIXa polypeptide before administration of the FIXa to the subject.

IV. Method of Making

A FVIII polypeptide can be manufactured in a host cell comprising a vector encoding the FVIII polypeptide. In one embodiment, the host cell is transformed with one or more vectors comprising a first nucleotide sequence encoding a FVIII polypeptide and a first FcRn polypeptide, a second nucleotide sequence encoding a second FcRn polypeptide, and optionally a third nucleotide sequence encoding a protein convertase, e.g., PC5. As used herein, an expression vector refers to any nucleic acid construct which contains the necessary elements for the transcription and translation of an inserted coding sequence, or in the case of an RNA viral vector, the necessary elements for replication and translation, when introduced into an appropriate host cell. Expression vectors can include plasmids, phagemids, viruses, and derivatives thereof.

A gene expression control sequence as used herein is any regulatory nucleotide sequence, such as a promoter sequence or promoter-enhancer combination, which facilitates the efficient transcription and translation of the coding nucleic acid to which it is operably linked. The gene expression control sequence can, for example, be a mammalian or viral promoter, such as a constitutive or inducible promoter. Constitutive mammalian promoters include, but are not limited to, the promoters for the following genes: hypoxanthine phosphoribosyl transferase (HPRT), adenosine deaminase, pyruvate kinase, beta-actin promoter, and other constitutive promoters. Exemplary viral promoters which function constitutively in eukaryotic cells include, for example, promoters from the cytomegalovirus (CMV), simian virus (e.g., SV40), papilloma virus, adenovirus, human immunodeficiency virus (HIV), Rous sarcoma virus, cytomegalovirus, the long terminal repeats (LTR) of Moloney leukemia virus, and other retroviruses, and the thymidine kinase promoter of herpes simplex virus. Other constitutive promoters are known to those of ordinary skill in the art. The promoters useful as gene expression sequences of the invention also include inducible promoters. Inducible promoters are expressed in the presence of an inducing agent. For example, the metallothionein promoter is induced to promote transcription and translation in the presence of certain metal ions. Other inducible promoters are known to those of ordinary skill in the art.

Examples of vectors include, but are not limited to viral vectors or plasmid vectors. Plasmid vectors have been extensively described in the art and are well-known to those of skill in the art. See, e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, 1989. In the last few years, plasmid vectors have been found to be particularly advantageous for delivering genes to cells in vivo because of their

inability to replicate within and integrate into a host genome. These plasmids, however, having a promoter compatible with the host cell, can express a peptide from a gene operably encoded within the plasmid. Some commonly used plasmids available from commercial suppliers include pBR322, pUC18, pUC19, various pcDNA plasmids, pRC/CMV, various pCMV plasmids, pSV40, and pBlueScript. Additional examples of specific plasmids include pcDNA3.1, catalog number V79020; pcDNA3.1/hygro, catalog number V87020; pcDNA4/myc-His, catalog number V86320; and pBudCE4.1, catalog number V53220, all from Invitrogen (Carlsbad, CA.). Other plasmids are well-known to those of ordinary skill in the art. Additionally, plasmids can be custom designed using standard molecular biology techniques to remove and/or add specific fragments of DNA.

The expression vector or vectors are then transfected or co-transfected into a suitable target cell, which will express the polypeptides. Transfection techniques known in the art include, but are not limited to, calcium phosphate precipitation (Wigler et al. (1978) *Cell* 14:725), electroporation (Neumann et al. (1982) *EMBO J* 1:841), and liposome-based reagents. A variety of host-expression vector systems can be utilized to express the proteins described herein including both prokaryotic and eukaryotic cells. These include, but are not limited to, microorganisms such as bacteria (e.g., *E. coli*) transformed with recombinant bacteriophage DNA or plasmid DNA expression vectors containing an appropriate coding sequence; yeast or filamentous fungi transformed with recombinant yeast or fungi expression vectors containing an appropriate coding sequence; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing an appropriate coding sequence; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus or tobacco mosaic virus) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing an appropriate coding sequence; or animal cell systems, including mammalian cells (e.g., HEK 293, CHO, Cos, HeLa, HKB11, and BHK cells).

In one embodiment, the host cell is a eukaryotic cell. As used herein, a eukaryotic cell refers to any animal or plant cell having a definitive nucleus. Eukaryotic cells of animals include cells of vertebrates, e.g., mammals, and cells of invertebrates, e.g., insects. Eukaryotic cells of plants specifically can include, without limitation, yeast cells. A eukaryotic cell is distinct from a prokaryotic cell, e.g., bacteria.

In certain embodiments, the eukaryotic cell is a mammalian cell. A mammalian cell is any cell derived from a mammal. Mammalian cells specifically include, but are not limited to, mammalian cell lines. In one embodiment, the mammalian cell is a human cell. In another embodiment, the mammalian cell is a HEK 293 cell, which is a human embryonic kidney cell line. HEK 293 cells are available as CRL-1533 from American Type Culture Collection, Manassas, VA, and as 293-H cells, Catalog No. 11631-017 or 293-F cells, Catalog No. 11625-019 from Invitrogen (Carlsbad, Calif.). In some embodiments, the mammalian cell is a PER.C6® cell, which is a human cell line derived from retina. PER.C6® cells are available from Crucell (Leiden, The Netherlands). In other embodiments, the mammalian cell is a Chinese hamster ovary (CHO) cell. CHO cells are available from American Type Culture Collection, Manassas, VA. (e.g., CHO-K1; CCL-61). In still other embodiments, the mammalian cell is a baby hamster kidney (BHK) cell. BHK cells are available from American Type Culture Collection, Manassas, Va. (e.g., CRL-1632). In some

embodiments, the mammalian cell is a HKB11 cell, which is a hybrid cell line of a HEK293 cell and a human B cell line. Mei et al., Mol. Biotechnol. 34(2): 165-78 (2006).

The method can further comprise purification steps. Various known purifications steps are well known in the art.

EXAMPLES

Example 1. Product Description

rFVIIIc is a long-acting, fully recombinant fusion protein consisting of human coagulation Factor VIII (FVIII) covalently linked to the Fc domain of human immunoglobulin G1 (IgG1). The Factor VIII portion of rFVIIIc has a primary amino acid sequence and post-translational modifications that are comparable to the 90+80 kDa form of Factor VIII (i.e., B-domain deleted). The Fc domain of rFVIIIc contains the hinge, CH2 and CH3 regions of IgG1. rFVIIIc contains 1882 amino acids with an apparent molecular weight of approximately 220 kilodaltons.

rFVIIIc is produced by recombinant DNA technology in a human embryonic kidney (HEK) cell line, which has been extensively characterized. The cell line expresses rFVIIIc into a defined cell culture medium that does not contain any proteins derived from animal or human sources. rFVIIIc is purified by a series of chromatography steps that does not require use of a monoclonal antibody. The process includes a detergent viral inactivation step and multiple viral clearance steps including an affinity chromatography step and a 15 nm virus-retaining nano-filtration step. No human or animal additives are used in the cell culture, purification, and formulation processes.

rFVIIIc is in the pharmacotherapeutic group: antihemorrhagics, blood coagulation factor VIII. It is provided as a sterile, preservative-free, non-pyrogenic, lyophilized, white to off-white powder to cake, for intravenous (IV) adminis-

excipients: sucrose, sodium chloride, L-histidine, calcium chloride, and polysorbate 20 or polysorbate 80, at the concentrations shown in Table 1 or Table 2 below. The pharmaceutical composition is formulated for intravenous administration only after reconstitution.

Each pack contains a powder vial (type 1 glass) with a stopper (butyl) and a flip-off seal (aluminum), 3 ml solvent in a pre-filled syringe (type 1 glass) with a plunger stopper (butyl), a tip-cap (butyl), and a sterile vial adapter reconstitution device.

TABLE 1

rFVIIIc Formulations Following Reconstitution with Liquid Diluent						
rFVIIIc IU/ml*	Sucrose % (w/v)	NaCl (mM)	L-histidine (mM)	Calcium		
				chloride (mM)	Polysorbate- 20 % (w/v)	
83 IU/ml	1.33	205	6.64	5.4	0.013	
167 IU/ml	1.33	205	6.64	5.4	0.013	
250 IU/ml	1.33	205	6.64	5.4	0.013	
333 IU/ml	1.33	205	6.64	5.4	0.013	
500 IU/ml	1.33	205	6.64	5.4	0.013	
667 IU/ml	1.33	205	6.64	5.4	0.013	
1000 IU/ml	1.33	205	6.64	5.4	0.013	
1333 IU/ml	1.33	205	6.64	5.4	0.013	
1667 IU/ml	1.33	205	6.64	5.4	0.013	
2000 IU/ml	1.33	205	6.64	5.4	0.013	

TABLE 2

rFVIIIc Formulations Following Reconstitution with Liquid Diluent										
Component	Concentration									
	250 IU/vial	500 IU/vial	750 IU/vial	1000 IU/vial	1500 IU/vial	2000 IU/vial	3000 IU/vial	4000 IU/vial	5000 IU/vial	6000 IU/vial
rFVIIIc*	83 IU/mL	167 IU/mL	250 IU/mL	333 IU/mL	500 IU/mL	667 IU/mL	1000 IU/mL	1333 IU/mL	1667 IU/mL	2000 IU/mL
Sucrose	13.3 mg/mL	13.3 mg/mL	13.3 mg/mL	13.3 mg/mL	13.3 mg/mL	13.3 mg/mL	13.3 mg/mL	13.3 mg/mL	13.3 mg/mL	13.3 mg/mL
Sodium Chloride	12.0 mg/mL	12.0 mg/mL	12.0 mg/mL	12.0 mg/mL	12.0 mg/mL	12.0 mg/mL	12.0 mg/mL	12.0 mg/mL	12.0 mg/mL	12.0 mg/mL
L-Histidine	1.03 mg/mL	1.03 mg/mL	1.03 mg/mL	1.03 mg/mL	1.03 mg/mL	1.03 mg/mL	1.03 mg/mL	1.03 mg/mL	1.03 mg/mL	1.03 mg/mL
Calcium Chloride Dihydrate	0.80 mg/mL	0.80 mg/mL	0.80 mg/mL	0.80 mg/mL	0.80 mg/mL	0.80 mg/mL	0.80 mg/mL	0.80 mg/mL	0.80 mg/mL	0.80 mg/mL
Polysorbate 20	0.13 mg/mL	0.13 mg/mL	0.13 mg/mL	0.13 mg/mL	0.13 mg/mL	0.13 mg/mL	0.13 mg/mL	0.13 mg/mL	0.13 mg/mL	0.13 mg/mL

*The potency (IU) is determined using the European Pharmacopoeia chromogenic assay against an in-house standard that is referenced to the WHO standard. The specific activity of rFVIIIc is 4000-10000 IU/mg protein.

tration in a single-use vial, accompanied by a liquid diluent in a pre-filled syringe. In addition to rFVIIIc, the pharmaceutical composition comprises in the lyophilizate sucrose, sodium chloride, L-Histidine, calcium chloride, and Polysorbate 20 or polysorbate 80, and comprising in sterilized water for injections. Each single-use vial contains nominally 250, 500, 750, 1000, 1500, 2000, 3000, 4000, 5000, or 6000 International Units (IU) of rFVIIIc. When reconstituted with provided diluent, the product contains the following

Example 2. Method of Formulation

The rFVIIIc drug product is a sterile lyophilized powder for injection intended for intravenous administration. It is supplied in aseptically filled single use vials which contain nominally 50, 500, 750, 1000, 1500, 2000, 3000, 4000, 5000 and 6000 IU per vial. The vials are 10 mL USP/Ph. Eur. Type 1 glass vials sealed with a 20 mm Teflon-coated butyl rubber

45

lyophilization stopper and aluminum flip-off crimp seal. Prior to lyophilization, the nominal fill volume target for 250 through 6000 IU vials is 3 mL. The composition of the formulation excipients prior to lyophilization is the same for all dosage strengths. The powder for injection is reconstituted with 3 mL of diluent comprising sterilized water for injections supplied in a sterile prefilled syringe.

The compositions of the drug product solutions prior to lyophilization are presented in Table 3. The compositions of the drug products following reconstitution are presented in Table 1 or in Table 2. (Example 1).

TABLE 3

Pre-lyophilization Concentrations of rFVIIIFc Formulation Components										
Component	Concentration									
	250 IU/vial	500 IU/vial	750 IU/vial	1000 IU/vial	1500 IU/vial	2000 IU/vial	3000 IU/vial	4000 IU/vial	5000 IU/vial	6000 IU/vial
rFVIIIFc	150 IU/ml*	287.5 IU/ml**	431.25 IU/ml**	575 IU/ml**	862.5 IU/ml**	1150 IU/ml**	1725 IU/ml**	2300 IU/ml**	2875 IU/ml**	3450 IU/ml**
Sucrose	20.0 mg/ml	20.0 mg/ml	20.0 mg/ml	20.0 mg/ml	20.0 mg/ml	20.0 mg/ml	20.0 mg/ml	20.0 mg/ml	20.0 mg/ml	20.0 mg/ml
Sodium Chloride	18.0 mg/ml	18.0 mg/ml	18.0 mg/ml	18.0 mg/ml	18.0 mg/ml	18.0 mg/ml	18.0 mg/ml	18.0 mg/ml	18.0 mg/ml	18.0 mg/ml
L-Histidine	1.55 mg/ml	1.55 mg/ml	1.55 mg/ml	1.55 mg/ml	1.55 mg/ml	1.55 mg/ml	1.55 mg/ml	1.55 mg/ml	1.55 mg/ml	1.55 mg/ml
Calcium Chloride Dihydrate	1.18 mg/ml	1.18 mg/ml	1.18 mg/ml	1.18 mg/ml	1.18 mg/ml	1.18 mg/ml	1.18 mg/ml	1.18 mg/ml	1.18 mg/ml	1.18 mg/ml
Polysorbate 20	0.2 mg/ml	0.2 mg/ml	0.2 mg/ml	0.2 mg/ml	0.2 mg/ml	0.2 mg/ml	0.2 mg/ml	0.2 mg/ml	0.2 mg/ml	0.2 mg/ml
Water for Injection	To make 1 ml ***									
Fill Volume	2.0 ml	2.0 ml	2.0 ml	2.0 ml	2.0 ml	2.0 ml	2.0 ml	2.0 ml	2.0 ml	2.0 ml

*A 20% overage is included for the 250 IU/vial strength to ensure that, after process and testing variations, at least the nominal concentration is present in the vial.

**A 15% overage is included for the 500-6000 IU/vial strengths to ensure that, after process and testing variations, at least the nominal concentration is present in the vial.

*** Removed during lyophilization.

Example 3. Dosage and Method of Administration/Method of Calculating Initial Estimated Dose

rFVIIIFc is long-acting anti-hemophilic factor (recombinant) indicated in adults and children (≥ 12 years) with hemophilia A (congenital Factor FVIII deficiency) for, e.g., control and prevention of bleeding episodes, routine prophylaxis to prevent or reduce the frequency of bleeding episodes, and perioperative management (surgical prophylaxis).

Dosing of rFVIIIFc can be estimated as described in this example, but can also be determined by standard tests such as FVIII activity assays described elsewhere herein.

1 IU of rFVIIIFc per kg body weight is expected to increase the circulating level of Factor VIII by 2 [IU/dL]. rFVIIIFc has been shown to have a prolonged circulating half-life.

Since patients can vary in their pharmacokinetic (e.g., half-life, in vivo recovery) and clinical responses to rFVIIIFc, the expected in vivo peak increase in Factor VIII

46

level expressed as IU/dL (or % of normal) or the required dose can be estimated using the following formulas:

$$\text{IU/dL (or \% of normal)} = [\text{Total Dose (IU)/body weight (kg)}] \times 2 \text{ (IU/dL per IU/kg)}$$

OR

$$\text{Dose (IU)} = \text{body weight (kg)} \times \text{Desired Factor VIII Rise (IU/dL or \% of normal)} \times 0.5 \text{ (IU/kg per IU/dL)}$$

The following table (Table 4) can be used to guide dosing in bleeding episodes:

TABLE 4

Guide to rFVIIIFc Dosing for Treatment of Bleeding		
Severity of Bleed	Desired Factor VIII Level (IU/dL or % of normal)	Dose (IU/kg)/ Frequency of Doses (hrs)
Minor and Moderate For example: joint, superficial muscle/no neurovascular compromise (except iliopsoas), deep laceration and renal, superficial soft tissue, mucous membranes	40-60	20-30 IU/kg Repeat every 24-48 hours until bleeding is resolved
Major For example: iliopsoas and deep muscle with neurovascular injury, or substantial blood loss, retroperitoneum, CNS, throat and neck, gastrointestinal.	80-100	40-50 IU/kg Repeat every 12-24 hours until bleeding is resolved

Adapted from WFH 2012

Subsequent dosage and duration of treatment depends on the individual clinical response, the severity of the Factor VIII deficiency, and the location and extent of bleeding.

The following table (Table 5) can be used to guide dosing for perioperative management (surgical prophylaxis):

TABLE 5

Guide to rFVIII Fc Dosing for Perioperative Management (Surgical Prophylaxis)		
Type of Surgery	Target Factor VIII Level (IU/dL or % of normal)	Dose (IU/kg)/ Frequency of Doses (hrs)
Minor Minor operations including uncomplicated dental extraction	50 to 80	25-40 IU/kg A single infusion can be sufficient. Repeat every 24 hours as needed to control bleeding.
Major Major operations including intra-abdominal joint replacement surgery	80 to 120	An initial preoperative dose of 40-60 IU/kg followed by a repeat dose of 40-50 IU/kg after 8-24 hours and then every 24 hours to maintain FVIII activity within the target range. rFVIII Fc has a longer half-life than plasma and recombinant FVIII products (see Example 5)

For routine prophylaxis, the recommended regimen is 50 IU/kg every 3-5 days. The dose can be adjusted based on patient response in the range of 25-65 I/kg.

For weekly prophylaxis, the recommended dose is 65 IU/kg.

rFVIII Fc is contraindicated in patients who have manifested severe hypersensitivity reactions, including anaphylaxis, to the product or its components. Severe hypersensitivity reactions were not observed in clinical trials; however, these have been known to occur with use of other factor VIII replacement factors.

The clinical response to rFVIII Fc can vary. If bleeding is not controlled with the recommended dose, the plasma level of Factor VIII can be determined, and a sufficient dose of rFVIII Fc can be administered to achieve a satisfactory clinical response. If the patient's plasma Factor VIII level fails to increase as expected or if bleeding is not controlled after rFVIII Fc administration, the presence of an inhibitor (neutralizing antibodies) should be suspected, and appropriate testing performed. Patients using rFVIII Fc can be monitored for the development of Factor VIII inhibitors by appropriate clinical observations and laboratory tests known to those of ordinary skill in the art.

Patient's plasma can be monitored for Factor VIII activity levels, e.g., the one-stage clotting assay to confirm adequate Factor VIII levels have been achieved and maintained, when clinically indicated. Patient's plasma can further be monitored for the development of Factor VIII inhibitors.

Example 4. Phase 3 Clinical Trial of Extended Half-Life Recombinant Fc Fusion Factor VIII ("A-LONG" Study)

Summary of A-LONG Study

In the A-LONG study, 165 male patients aged 12 years of age and older were enrolled. The A-LONG study had three treatment arms: individualized prophylaxis, weekly prophylaxis and episodic (on-demand) treatment (Arms 1, 2 and 3, respectively). In a subgroup of patients across treatment arms, rFVIII Fc was evaluated in the perioperative management of patients who required a major surgical procedure during the study.

Overall, 92.7 percent of patients completed the study. Recombinant FVIII Fc was generally well-tolerated. No inhibitors to rFVIII Fc were detected and no cases of anaphylaxis were reported in any patients, all of whom switched from commercially-available Factor VIII products. No serious adverse events were assessed to be related to drug by the investigator.

The most common adverse events (incidence of ≥ 5 percent) occurring outside of the perioperative management period were nasopharyngitis, arthralgia, headache, and upper respiratory tract infection.

The median (mean) annualized bleeding rates (ABR), including spontaneous and traumatic bleeds, were 1.6 (2.9) in the individualized prophylaxis arm, 3.6 (8.8) in the weekly prophylaxis arm and 33.6 (37.2) in the episodic treatment arm. In the individualized prophylaxis arm, the median dosing interval was 3.5 days. During the last 3 months on study, 30 percent of patients in the individualized prophylaxis arm achieved a mean dosing interval of at least 5 days.

Control of bleeding was assessed in all patients who experienced a bleeding episode during the study. Overall, 98 percent of bleeding episodes were controlled by one or two injections of rFVIII Fc.

In addition, rFVIII Fc was assessed in the perioperative management of 9 patients undergoing 9 major surgical procedures. The treating physicians rated the hemostatic efficacy of rFVIII Fc as excellent or good in 100 percent of these surgeries.

A-LONG included pharmacokinetic (PK) analysis of rFVIII Fc in all patients in the study. In a protocol-defined subset of patients with extensive PK sampling, the approximate terminal half-life of rFVIII Fc was 19.0 hours compared to 12.4 hours for ADVATE® [Antihemophilic Factor (Recombinant), Plasma/Albumin-Free Method], consistent with the results obtained in the Phase 1/2 study of rFVIII Fc.

About the A-LONG Study and the rFVIII Fc Program

A-LONG was a global, open-label, multi-center Phase 3 study that evaluated the efficacy, safety and pharmacokinetics of intravenously-injected rFVIII Fc. The study was designed to evaluate rFVIII Fc in the control and prevention of bleeding, routine prophylaxis and perioperative management in patients with hemophilia A. A-LONG involved 60 hemophilia treatment centers in 19 countries on 6 continents.

The A-LONG study had three treatment arms. In Arm 1 (individualized prophylaxis; n=117), patients were treated with 25-65 IU/kg of rFVIII Fc, at an interval of every three to five days, which was individualized to maintain factor trough levels sufficient to prevent bleeding. In Arm 2 (weekly prophylaxis; n=24), patients were treated with a weekly dose of 65 IU/kg. In Arm 3 (episodic treatment; n=23), patients received rFVIII Fc episodic treatment as needed for bleeding. In a subgroup of patients across treatment arms, rFVIII Fc was evaluated in the surgical setting.

The primary efficacy measures were the per-patient annualized bleeding rate in Arm 1 vs. Arm 3, and pharmacokinetics of rFVIII Fc vs. rFVIII (one-stage clotting [activated partial thromboplastin time] assay and the chromogenic assays calibrated to normal human references plasma with potency traceable to World Health Organization standards). The safety endpoints included the incidence of adverse events and inhibitor development (Nijmegen-modified Bethesda assay) in patients studied for up to 54 weeks. Secondary efficacy endpoints included ABR in Arm 2 vs. Arm 3, annualized number of spontaneous and joint bleeding episodes per patient, number of injections and dose per

injections to resolve a bleed, and investigators' assessments of patients' response to surgery with rFVIII Fc using a bleeding response scale (surgery subgroup only), response to treatment of bleeding episodes and the pharmacokinetics of rFVIII Fc versus ADVATE®. Non-neutralizing antibodies (NNAs) were assessed with an electrochemiluminescence-based anti-rFVIII Fc binding-antibody assay.

Ongoing clinical studies of rFVIII Fc include the Kids A-LONG and ASPIRE studies. Kids A-LONG is a Phase 3, open-label study in previously-treated children with hemophilia A under age 12, which is actively recruiting patients. ASPIRE is a long-term open-label study for patients who completed the A-LONG study or who complete the Kids A-LONG study.

A-LONG Study Design

Design: Global, open-label, multicenter, Phase 3 study. The study protocol was approved by local Institutional Review Boards for each participating institution, and the study was conducted in accordance with the International Conference on Harmonisation guidelines for Good Clinical Practice.

Objectives: To evaluate the efficacy and safety of intravenously-injected recombinant factor VIII Fc fusion protein (rFVIII Fc) in the control and prevention of bleeding episodes, routine prophylaxis, and perioperative management in individuals with severe hemophilia A.

Key Inclusion Criteria:

- a. Male
- b. ≥ 212 years of age and at least 40 kg
- c. Diagnosis of severe hemophilia A defined as $<1\%$ (<1 IU/dL) endogenous factor VIII (FVIII) activity
- d. Previously Treated Patients (PTPs); history of ≥ 150 prior documented exposure days (EDs) with any currently marketed FVIII product, been on prophylaxis ≥ 2 times per week with a FVIII product (for Arm 1 only), or experienced ≥ 12 bleeds in the past 12 months on an episodic regimen.
- e. No current/prior FVIII inhibitors of measurable activity.
- f. No history of inhibitors, or history of hypersensitivity or anaphylaxis associated with any FVIII or intravenous immunoglobulin administration
- g. All subjects (or their guardians in the case of minors) gave informed written consent prior to study participation.

Treatment arms: Details of A-LONG study design is described below.

Arm 1 (individualized prophylaxis) was administered 25-65 IU/kg every 3-5 days (maximum 5-day dosing interval). Subjects were treated with an initial dose of 25 IU/kg on Day 1 and 50 IU/kg on Day 4, which was subsequently adjusted to maintain trough factor levels sufficient to prevent bleeding. Further dose adjustments to target trough levels of 3 to 5 IU/dL permitted after week 7 if subject experienced moderate/severe bleeds over a rolling 8-week period; 10 to 20 IU/kg rFVIII Fc (target 20-40 IU/dL FVIII) for minor bleeding episodes; 15 to 30 IU/kg rFVIII Fc (target 30-60 IU/dL FVIII) for moderate to major bleeding episodes; 40 to 50 IU/kg rFVIII Fc (target 80-100 IU/dL FVIII) for major to life-threatening bleeding episodes.

Arm 2 (weekly prophylaxis) was administered 65 IU/kg dose. Subjects were treated with 65 IU/kg once weekly with no dose or interval adjustment.

Arm 3 (episodic [on-demand] treatment) was administered 10-50 IU/kg. Subjects received rFVIII Fc episodic treatment as needed for bleeding.

A Perioperative Management Subgroup was established. In this subgroup, rFVIII Fc was administered prior to and following surgery in the subset of patients requiring a major surgical procedure during the study. Subjects in any treatment arm could be enrolled in the surgery subgroup. Eligibility: required major surgery; 12 exposure days to rFVIII Fc with negative inhibitor titre following this period and within 4 weeks prior to surgery; and, completed, at minimum, abbreviated PK sampling.

All subjects (excluding sequential PK subgroup) underwent PK sampling from pre-injection with rFVIII Fc up to 96-hours post-injection, according to schedule: pre-injection, 30 (+3) minutes, 3 hours (15 minutes), 72 (+2) hours (Day 3), and 96 (+2) hours (Day 4) from start of injection.

Pharmacokinetic (PK) Assessment was performed. All subjects in all arms had an initial PK assessment after their first dose of rFVIII Fc. A subset of subjects from Arm 1 were assigned to a protocol-specified sequential PK subgroup to compare the stability of PK properties of rFVIII Fc over time with that of recombinant factor VIII (rFVIII, ADVATE® [anti-hemophilic factor (recombinant) plasma/albumin-free method], octocog alfa) as follows:

Prior to treatment in Arm 1, PK was assessed after a single dose of ADVATE® 50 IU/kg. PK was then assessed in these same subjects after a single dose of rFVIII Fc 50 IU/kg. PK of rFVIII Fc was repeated at 12 to 24 weeks.

Details on the design of the sequential PK subgroup (Arm 1) dosing and PK sampling is described in FIG. 1

Key Efficacy Outcome Measures (Included in Initial Read-out):

- a. Annualized bleeding rate (ABR) in Arm 1 versus Arm 3
- b. Individualized prophylaxis arm compared with episodic treatment arm.
- c. Number of injections required to resolve a bleeding episode
- d. Number of injections required to resolve bleeding episodes
- e. Median dose required to resolve bleeding episodes
- f. Treating physicians' assessments of subjects' response to surgery with rFVIII Fc using a 4-point scale

PK outcome measures included:

- a. The primary PK assessment were based on FVIII activity levels determined at a central laboratory by one-stage clotting assay and by the chromogenic assay against commercially available plasma standards.
- b. PK of rFVIII Fc and ADVATE®
- c. PK properties of rFVIII Fc were compared with rFVIII (ADVATE®; sequential PK subgroup). Overall study duration was ≤ 75 weeks for all subjects. The primary efficacy endpoint was annualized bleeding rate (ABR; Arms 1 and 2 vs. Arm 3). Prophylaxis dose and interval, number of injections required for treatment of bleeding episodes, and perioperative haemostasis were evaluated.

Key safety outcome measures included:

- a. Incidence of inhibitor development. The study was powered to detect the occurrence of inhibitors with a 2-sided 95% confidence interval using the Clopper-Pearson exact method if 2 cases of inhibitor formation was observed.
- b. Incidence of adverse events (AEs) occurring outside of the perioperative management period

Adherence with treatment (adherence measures) was assessed using patient electronic diaries.

51

A-LONG Results
Subjects

A total of 165 subjects were enrolled in the study. Median age was 30 years (range, 12-65) and 8% were <18 years. The number of patients in each arm of the study were: Arm 1 (individualized prophylaxis), n=118; Arm 2 (weekly prophylaxis), n=24; Arm 3 (episodic treatment), n=23; and perioperative management subgroup, n=9, 9 surgeries (8 subjects from Arm 1 and 1 from Arm 2). Subjects from each treatment arm were eligible to enter the surgery subgroup if

52

(95%) in Arm 1, 19/24 (79%) in Arm 2, and 22/23 (96%) in Arm 3. Reasons for premature discontinuation were: Arm 1—subject withdrawal (n=2), physician decision (n=2), other (n=1), death (n=1); Arm 2—subject withdrawal (n=2); adverse events (AEs) related to study drug can have contributed), AEs (n=2), other (n=1); Arm 3—other (n=1).

Age, race, and geography of subjects were representative of the global haemophilia A population who have access to treatment, as summarized in TABLE 6.

TABLE 6

A-LONG subjects demographics				
Demographic	Arm 1 (N = 118)	Arm 2 (N = 24)	Arm 3 (N = 23)	Total N = 165
Age (years), median (min-max)	29.0 (12-65)	31.5 (18-59)	34.0 (13-62)	30.0 (12-65)
Weight (kg), median (min-max)	71.65 (42.0-127.4)	75.85 (50.0-105.0)	70.00 (48.0-110.4)	71.60 (42.0-127.4)
BMI (kg/m ²) median (min-max)	23.90 (15.3-37.1)	24.60 (18.8-37.4)	22.80 (17.2-35.6)	23.90 (15.3-37.4)
Race, n (%)				
White	79 (66.9)	12 (50.0)	16 (69.6)	107 (64.8)
Black	7 (5.9)	1 (4.2)	2 (8.7)	10 (6.1)
Asian	27 (22.9)	11 (45.8)	5 (21.7)	43 (26.1)
Other	5 (4.2)	0	0	5 (3.0)
Geographic location - n (%)				
Europe	34 (28.8)	3 (12.5)	4 (17.4)	41 (24.8)
North America	44 (37.3)	5 (20.8)	7 (30.4)	56 (33.9)
Other*	40 (33.9)	16 (66.7)	12 (52.2)	68 (41.2)
Genotype - n (%)				
Intron 22 inversion	41 (35.0)	7 (33.3)	9 (39.1)	57 (35.4)
Frameshift	24 (20.5)	4 (19.0)	6 (26.1)	34 (21.1)
Missense mutation	22 (18.8)	4 (19.0)	1 (4.3)	27 (16.8)
Nonsense mutation	19 (16.2)	6 (28.6)	1 (4.3)	26 (16.1)
Splice site change	7 (6.0)	0	4 (17.4)	11 (6.8)
Intron 1 inversion	3 (2.6)	0	1 (4.3)	4 (2.5)
Duplication	1 (0.9)	0	0	1 (0.6)
NA	0	0	1 (4.3)	1 (0.6)
von Willebrand factor antigen - median IU/dl (IQR)	118.0 (85, 151)	129.0 (86, 166)	131.0 (83, 155)	118.0 (85, 153)
Pre-study FVIII regimen - n (%)				
Prophylaxis	87 (73.7)	0	0	87 (52.7)
Episodic	31 (26.3)	24 (100)	23 (100)	78 (47.3)
Estimated no. of bleeds prior 12 mo - median (IQR) †				
Prior prophylaxis	6.0 (2, 15)	—	—	6 (2, 15)
Prior episodic	27.0 (17, 41)	29.5 (19, 44)	24.0 (15, 36)	27 (18, 40)
≥1 Target joint - n (%)				
Prior prophylaxis	47 (39.8)	—	—	47 (28.5)
Prior episodic	26 (22.0)	22 (91.7)	18 (78.3)	66 (40.0)
Family history of inhibitor	4 (3.4)	1 (4.2)	2 (8.7)	7 (4.2)
HIV positive	25 (21.2)	4 (16.7)	7 (30.4)	36 (21.8)
HCV positive	55 (46.6)	14 (58.3)	13 (56.5)	82 (49.7)

BMI, body mass index; HIV, human immunodeficiency virus; HCV, hepatitis C virus

IU/dl denotes international units per deciliter, BMI body mass index, HIV human immunodeficiency virus, HCV hepatitis C virus, NA not applicable.

*Other included Australia, New Zealand, Brazil, Hong Kong, India, Japan, Russia, and South Africa.

† Calculation was based on available data.

55

they required major surgery, had ≥12 EDs to rFVIIIFc and a negative inhibitor titre following this period and within 4 weeks prior to surgery. Patients on prior episodic treatment had a higher median number of bleeding episodes in the 12 months prior to the study, and a higher proportion of patients in Arms 2 and 3 had target joints. For patients on prior prophylaxis, 87% reported injecting at least three times weekly.

92.7% of subjects completed the study. In total, 153 (approximately 93%) subjects completed the study; 112/118

Subjects received rFVIIIFc for a median (min-max) of 30.5 (<1-54) weeks with a median (minimum to maximum) in arms 1, 2, and 3 of 32.1 (9, 54), 28.0 (<1, 38), and 28.9 (15, 32) weeks, respectively. In total, 111 subjects (67.7%) had >50 EDs to study drug. A total of 9356 injections were administered during the study, corresponding to 9170 EDs (100.2 patient-years of exposure). Overall, 93.6% of patients were compliant with both the prescribed dose and interval in the prophylaxis arms.

Overall, 93.6% of subjects were adherent with both the prescribed dose and the prescribed dosing interval in the prophylaxis treatment arms.

65

Efficacy

In total, 163 subjects were included in efficacy analyses. One subject was excluded due to only receiving ADVATE® (Arm 1); another subject withdrew before efficacy assessment (Arm 2). Median ABR with the 25th and 75th percentiles (interquartile range [IQR]) were as follows: Arm 1—individualized prophylaxis arm: 1.6 (0.0, 4.7); Arm 2—weekly prophylaxis arm: 3.6 (1.9, 8.4); and Arm 3—episodic treatment arm: 33.6 (21.1, 48.7). Most bleeding episodes were spontaneous. In Arms 1, 2, and 3, respectively, 45.3%, 17.4%, and 0% had no bleeding episodes and 13.7%, 34.8%, and 0% had 1 bleeding episode.

Median Dosing Interval:

- In the individualized prophylaxis arm, the median dosing interval was 3.5 days, during the last 3 months on study.
- 30 percent of patients in the individualized prophylaxis arm achieved a mean dosing interval of at least 5 days.
- Overall, the median (IQR) dosing interval with individualised prophylaxis aimed to achieve a FVIII trough of at least 1 to 3 IU/dL (with the maximum 5-day dosing interval permitted in the protocol) was 3.5 (3.2-4.4) days (Table 7) based on the median (IQR) weekly dose of 78 (72-91) IU/kg.
- Approximately 30% of subjects achieved a mean dosing interval of 5 days over the last 3 months on study. 100% of subjects had mean dosing intervals 2 days throughout the study.

The median dose per injection for Arm 3 episodic regimen was 26.5 IU/kg (n=23) and the median total dose per bleeding episode was 27.4 IU/kg (n=23).

TABLE 7

Comparative pharmacokinetics for rFVIIIIFc vs. rFVIII (n = 28) ¹			
PK parameter	Geometric mean for rFVIIIIFc PK (95% CI)	Geometric mean for rFVIII PK (95% CI)	Geometric mean of intra-subject ratio (95% CI) p-value
Elimination t _{1/2} (h)	18.97 (17.03, 21.12)	12.43 (11.14, 13.86)	1.53 (1.36, 1.71) <0.001
CL (mL/h/kg)	1.95 (1.71, 2.22)	3.041 (2.71, 3.41)	0.64 (0.60, 0.69) <0.001
Time to 1 IU/dL (days) (50 IU/kg dose)	4.92 (4.43, 5.46)	3.30 (2.99, 3.65)	1.49 (1.41, 1.57) <0.001

¹Arm 1 sequential PK group, compartmental model, one-stage assay. Estimates and 95% CIs for geometric means and geometric mean ratios; statistical significance assessed at the 2-sided 0.05 level; PK, pharmacokinetics; CI, confidence interval; C_{max}, maximal concentration; AUC, area-under-the-curve; t_{1/2}, half-life; CL, clearance; MRT, mean residence time; V_{ss}, volume of distribution at steady state.

Control of bleeding: approximately 98% (97.7%) of bleeding episodes were controlled by one or two injections of rFVIIIIFc. 87.3% of bleeds were controlled with one injection. 1.7% required three injections.

Perioperative management: Overall, 9 major surgeries were performed in 9 subjects (8 subjects from Arm 1; 1 subject from Arm 2), including knee arthroplasty (n=5), laparoscopic inguinal hernia repair (n=2), appendectomy (n=1), and arthroscopy (n=1). Treating physicians rated the hemostatic efficacy of rFVIIIIFc as excellent (8/9) or good (1/9) in 100% of surgeries. Median (min, max) estimated blood loss available for 7/9 surgeries was 15.0 (0, 600) mL during surgery and 0.0 (0, 1100) mL post-operatively; post-surgical drainage).

PK

Comparative PK data for rFVIIIIFc and ADVATE® available from 28 subjects in the Arm 1 sequential PK subgroup are summarised in Table 7.

- rFVIIIIFc demonstrated an approximate 50% longer elimination half-life (FIG. 2) and mean residence time compared with ADVATE® (P<0.001) due to a 35% reduction in rFVIIIIFc clearance with respect to ADVATE®.
- Geometric mean time to 1 IU/dL FVIII activity following a 50 IU/kg dose was approximately 5 days for rFVIIIIFc versus approximately 3 days for ADVATE® (P<0.001; Table 7, FIG. 2).
- Incremental recovery for rFVIIIIFc was clinically comparable with ADVATE®.
- There was no shift in the PK properties of rFVIIIIFc as evidenced by comparable PK properties between the baseline and repeat rFVIIIIFc PK profiles.
- Analogous PK results were obtained when the analysis was based on the chromogenic FVIII assay.
- The one-stage and chromogenic clotting assays accurately and precisely measured both rFVIIIIFc and ADVATE® utilising commercially available plasma FVIII standards.

The geometric mean (95% confidence interval) terminal half-life of rFVIIIIFc was approximately 19.0 hours (17.0, 21.1) hours, which is 1.53-fold longer than that of ADVATE® (approximately 12.4 (11.1, 13.9) hours).

Geometric mean time (IQR) to 1% FVIII activity following 50 IU/kg of rFVIIIIFc was approximately 5 days (4.92 (4.43, 5.46) days). Comparable PK profiles of rFVIIIIFc were observed at Week 14. The median dosing interval with individualized prophylaxis was 3.5 days and the median dose per week was 78 IU/kg; approximately 30% of subjects achieved a mean dosing interval of ≥5 days over the last 3 months on study (subjects with 26 months on study) as shown in Table 8. 98% of bleeding episodes were controlled with 1-2 injections.

TABLE 8

Prophylactic dosing summary			
	Arm 1 Individualised n	Arm 2 Weekly n	
Median dose, IU/kg/week (IQR)			
Overall	117	23	77.9 (72.3, 91.2)
Last 3 months ¹	112	16	77.7 (71.9, 106.2)
Median dosing interval, days (IQR)			
Overall	117		3.5 (3.2, 4.4)
Last 3 months ¹	112		3.5 ³ (3.0, 5.0)

¹Based on last 3 months on study for subjects on study ≥6 months

²Fixed dose of 65 IU/kg/wk.

³30% of subjects achieved every 5-day pharmacokinetic-driven dosing to maintain troughs 1% to 3% above baseline. IQR, interquartile range.

Safety

No inhibitors were detected to rFVIIIIFc, and no cases of anaphylaxis, allergy, or serious thrombotic events were reported.

rFVIIIc was generally well tolerated, and no serious adverse events were assessed to be related to rFVIIIc.

The most common AEs, regardless of causality, (incidence 25%) occurring outside of the perioperative management period were nasopharyngitis, arthralgia (joint pain), headache, and upper respiratory tract infection, as summarized in Table 9.

12 subjects (7.3%) experienced at least one serious AE (SAE) outside of the perioperative management period.

No SAEs were assessed to be related to drug by the investigator.

Insignificant incidence of non-neutralizing antibody (NNA) has been observed under NNA assay, which is approximately 20 times more sensitive than Bethesda assay. Five subjects were found positive for NNAs at baseline, but all had at least one negative evaluation during the study. Six subjects, four in Arm 1 study and two in Arm 2 study, became positive during the study. In almost all cases, however, antibodies were transient and of low titer, and in all cases antibodies were directed against FVIII.

Overall, 14 (8.5%) subjects reported SAEs. No SAEs were determined by the investigator as related to study drug, and SAEs reported were not experienced in more than 1 subject. There was 1 death during the study, attributed to polysubstance overdose, and assessed as unrelated to rFVIIIc by the investigator.

TABLE 9

Summary of adverse events (AEs)	
	Total N = 164 n (%)
Any AE	108 (65.9)
Most common AEs ($\geq 5\%$ of subjects)	
Nasopharyngitis	20 (12.2)
Arthralgia	13 (7.9)
Headache	9 (5.5)
Upper respiratory infection	9 (5.5)

Summary

Individualized and weekly prophylactic regimens resulted in low single-digit median annualized bleeding rates

In the individualized prophylaxis arm, the median dosing interval was 3.5 days. During the last 3 months on study, 30 percent of patients in the individualized prophylaxis arm achieved a mean dosing interval of at least 5 days.

98% of bleeding episodes were controlled by one or two injections of rFVIIIc. In total, 757 bleeding episodes (Arm 1=209; Arm 2=92; Arm 3=456) were treated in 106 of the 164 subjects. 87.3% required a single injection of rFVIIIc for resolution, and 97.8% required ≤ 2 injections. A total of 85.6%, 80.4% and 89.5% of bleeding episodes were resolved with 1 injection of rFVIIIc in Arms 1, 2 and 3, respectively.

Hemostatic efficacy of rFVIIIc during surgery was rated by treating physicians as excellent or good in 100% of surgeries.

The half-life of rFVIIIc was approximately 19.0 hours compared to 12.4 hours for ADVATE®.

No subject developed an inhibitor or experienced an anaphylactic reaction to rFVIIIc.

Recombinant FVIIIc was generally well tolerated.

A-LONG was the largest registrational global pivotal phase 3 study of long-lasting rFVIII in severe haemophilia A conducted to date. The study showed that rFVIIIc offers the potential for a markedly reduced injection frequency, decreased treatment burden, an improvement in clinical outcomes for prevention of bleeds in patients with severe haemophilia A.

Goal of the study design: The study was designed to evaluate the efficacy and safety of rFVIIIc in the control and prevention of bleeding episodes, routine prophylaxis, and perioperative management in subjects with severe hemophilia A. In addition, the study was designed to assess the effective dose and interval of rFVIIIc for prophylaxis, as well as the feasibility of weekly treatment at a dose of 65 IU/Kg.

Episodic treatment is the administration of replacement factor only as needed to treat bleeding episodes after they have started. Prophylactic treatment is the regular administration of replacement factor to prevent bleeding episodes.

The starting regimen was 25 LU/kg on Day 1 and 50 IU/kg on Day 4. The dose and frequency of treatment could be adjusted based on the subject's PK profile and the goal was to maintain trough factor levels sufficient to prevent bleeding episodes using doses between 25 IU/kg and 65 IU/kg and treatment intervals of 3 to 5 days.

More than half of severe hemophilia A patients in the US still do not follow a prophylaxis regimen, due in part to the treatment burden. We therefore included the weekly dosing regimen in the A-LONG study to evaluate whether this regimen would result in a lower ABR compared to an episodic regimen. The median ABR in the weekly regimen was 3.59 (IQR 1.9, 8.4) compared with 33.6 (IQR 21.1, 48.7) for episodic treatment.

All subjects had an initial PK evaluation to characterize the PK of rFVIIIc in a representative population of patients with Hemophilia A.

More extensive PK sampling was conducted in a subset of subjects in the individualized prophylaxis arm (Arm 1) at baseline after a single dose of ADVATE® (50 IU/kg), and after a 4 day washout, followed by a single dose of rFVIIIc (50 IU/kg). Blood samples were taken for ADVATE® over a period of 72 hours. Blood samples were then taken for rFVIIIc over a period of 120 hours. PK assessment of rFVIIIc in this subset was repeated at 12 to 24 weeks with the same PK sampling schedule.

In peri-operative management, a single injection of rFVIIIc was sufficient to maintain haemostasis to the end date/time of all major surgeries at a median dose of 51.4 IU/kg. Median rFVIIIc consumption (summarized over all injections during each referenced time period) was 80.6 IU/kg on the day of major surgery, 161.3 IU/kg for Days 1-3 days following surgery, and 387.1 IU/kg for Days 4-14 following surgery. Perioperative haemostasis with rFVIIIc was rated as excellent or good for all 9 major surgeries. No subjects reported a bleeding episode during the postoperative or rehabilitation periods. Overall, 7 adverse events (AEs) were reported in 4 (44.4%) subjects in the surgery subgroup, of which 6 AEs were of mild or moderate severity, and 1 AE was considered severe. Two serious AEs (inguinal hernia and appendicitis) were reported in 2 subjects. All AEs during the perioperative period were assessed by the investigators as unrelated to rFVIIIc treatment.

Treating physicians rated the hemostatic efficacy of rFVIIIc as excellent or good in 100% of surgeries on a 4 point scale including excellent, good, fair, and poor/none.

rFVIIIc resulted in low median ABRs of 3.59 in the weekly prophylaxis arm and 1.60 in the individualized

interval prophylaxis arm. In contrast, the episodic treatment arm had a median ABR of 33.57.

In the individualized prophylaxis arm, the median dosing interval was 3.5 days at a median dose of 77.7 FU/kg during the last 3 months on study.

The terminal half-life was approximately 18.97 hours for rFVIII^{IFc} and approximately 12.3 hours for ADVATE®.

97.8% of bleeding episodes were controlled by one or two injections.

Hemostatic efficacy of rFVIII^{IFc} for perioperative management was rated by treating physicians as excellent or good in 100% of surgeries.

Arm 2 was a weekly dosing regimen. It was designed to investigate benefit for patients on prophylaxis therapy, and to compare those results with subjects using an episodic treatment regimen. The median ABR in Arm 2 was 3.59 compared with 33.57 for the episodic treatment regimen (Arm 3).

The median dose per injection required for resolution of bleeding was 27.4 IU/kg, and the median total dose required was 28.2 IU/kg.

The median number of injections required for resolution of a bleeding episode was consistently 1.0 when treatment was administered within 8 hours of bleed onset, regardless of type or location of bleed. Overall, 78.8% of rFVIII^{IFc} injections in Arm 1, 64.8% in Arm 2, and 79.7% of injections in Arm 3, were rated by subjects as producing excellent or good response. The median dosing interval of 3.5 days from Arm 1 represents the dosing interval that can be achieved by the majority of patients: 30% of patients were able to achieve a dosing interval of at least 5 days. Some patients were able to achieve a weekly prophylaxis regimen (Arm 2) with a median ABR of 3.59 bleeding episodes.

Subjects in the individualized prophylaxis arm received an initial dose of 25 IU/kg on Day 1 and 50 IU/kg on Day 4. The study design allowed for adjustment of the dosing interval and dose to maintain targeted trough factor levels and to prevent bleeding. We believe the median dosing interval and dose based on the last 6 months on study is most representative of the individualized prophylaxis regimen.

87.3% of bleeding episodes were resolved with one injection of rFVIII^{IFc} and 97.8% were resolved with one or two injections. For the 96 (12.7%) bleeding episodes that required more than 1 injection for resolution, the median interval between the first and second injection was 30.9 hours.

The investigators' global assessment of subject response to their assigned rFVIII^{IFc} regimen was rated as excellent or effective for 99.4%, 100%, and 98.1% of the subject visits in Arms 1, 2, and 3, respectively.

In A-LONG, the clotting factor activity was measured using a one-stage (aPTT) clotting assay and a chromogenic assay. The reported half-lives are based on the results of the one-stage clotting assay. While the half-lives of rFVIII^{IFc} and ADVATE® were both slightly longer with the chromogenic assay, the ratios between products were consistent between the two assays.

EMBODIMENTS

E1. A pharmaceutical composition comprising:

- (a) a FVIII polypeptide;
- (b) one or more stabilizing agents selected from the group consisting of sucrose, trehalose, raffinose, arginine, and mixture thereof;
- (c) sodium chloride (NaCl);
- (d) L-histidine;

(e) calcium chloride; and

(f) polysorbate 20 or polysorbate 80.

E2. The pharmaceutical composition of embodiment E1, wherein mannitol, glycine, alanine, or hydroxyethyl starch is not included.

E3. The pharmaceutical composition of embodiment E1 or E2, wherein NaCl is the only bulking agent.

E4. The pharmaceutical composition of any one of embodiments E1 to E3, comprising sucrose.

E5. The pharmaceutical composition of embodiment E4, comprising about 1% (w/v) to about 2.5% (w/v) sucrose.

E6. The pharmaceutical composition of embodiment E5, comprising about 1.3% (w/v) sucrose to about 2.0% (w/v) sucrose.

E7. The pharmaceutical composition of embodiment E6, comprising about 1.33% (w/v) sucrose or about 2.0% (w/v) sucrose.

E8. The pharmaceutical composition of embodiment E4, comprising about 10 mg/ml to about 25 mg/ml sucrose.

E9. The pharmaceutical composition of embodiment E8, comprising about 13 mg/ml to about 20 mg/ml sucrose.

E10. The pharmaceutical composition of embodiment E9, comprising about 13.3 mg/ml sucrose or about 20.0 mg/ml sucrose.

E11. The pharmaceutical composition of any one of embodiments E1 to E10, comprising about 150 mM to about 250 mM NaCl.

E12. The pharmaceutical composition of embodiment E11, comprising about 175 mM to about 225 mM NaCl.

E13. The pharmaceutical composition of embodiment E12, comprising about 200 mM to about 210 mM NaCl.

E14. The pharmaceutical composition of embodiment E13, comprising about 205 mM NaCl.

E15. The pharmaceutical composition of any one of embodiments E1 to E10, comprising about 8.8 mg/ml to about 14.6 mg/ml NaCl.

E16. The pharmaceutical composition of embodiment E15, comprising about 10 mg/ml to about 13 mg/ml NaCl.

E17. The pharmaceutical composition of embodiment E16, comprising about 12.0 mg/ml NaCl.

E18. The pharmaceutical composition of any one of embodiments E1 to E10, comprising about 250 mM to about 350 mM NaCl.

E19. The pharmaceutical composition of embodiment E18, comprising about 275 mM to about 325 mM NaCl.

E20. The pharmaceutical composition of embodiment E19, comprising about 308 mM NaCl.

E21. The pharmaceutical composition of any one of embodiments E1 to E10, comprising about 14.6 mg/ml to about 20.5 mg/ml NaCl.

E22. The pharmaceutical composition of embodiment E21, comprising about 16 mg/ml to about 19 mg/ml NaCl.

E23. The pharmaceutical composition of embodiment E22, comprising about 18.0 mg/ml NaCl.

E24. The pharmaceutical composition of any one of embodiments E1 to E23, comprising about 5 mM to about 15 mM L-histidine.

E25. The pharmaceutical composition of embodiment E24, comprising about 6.64 mM L-histidine or about 9.8 mM L-histidine.

E26. The pharmaceutical composition of any one of embodiments E1 to E23, comprising about 0.75 mg/ml to about 2.25 mg/ml L-histidine.

E27. The pharmaceutical composition of embodiment E26, the pharmaceutical composition comprises about 1.03 mg/ml L-histidine or about 1.55 mg/ml L-histidine.

E28. The pharmaceutical composition of any one of embodiments E1 to E27, comprising about 5 mM to about 10 mM calcium chloride.

E29. The pharmaceutical composition of embodiment E28, comprising about 5.4 mM calcium chloride or about 8 mM calcium chloride.

E30. The pharmaceutical composition of any one of embodiments E1 to E27, comprising about 0.75 mg/ml to about 1.5 mg/ml calcium chloride dihydrate.

E31. The pharmaceutical composition of embodiment E30, comprising about 0.8 mg/ml calcium chloride dihydrate or about 1.18 mg/ml calcium chloride dihydrate.

E32. The pharmaceutical composition of any one of embodiments E1 to E31, comprising about 0.008% (w/v) to about 0.025% (w/v) polysorbate 20 or polysorbate 80.

E33. The pharmaceutical composition of embodiment E32, comprising about 0.013% (w/v) polysorbate 20 or polysorbate 80 or about 0.02% (w/v) polysorbate 20 or polysorbate 80.

E34. The pharmaceutical composition of any one of embodiments E1 to E31, comprising about 0.08 mg/ml to about 0.25 mg/ml polysorbate 20 or polysorbate 80.

E35. The pharmaceutical composition of embodiment E34, comprising about 0.13% mg/ml polysorbate 20 or polysorbate 80 or about 0.20 mg/ml polysorbate 20 or polysorbate 80.

E36. The pharmaceutical composition of any one of embodiments E1 to E35, wherein the rFVIII_{FC} polypeptide comprises a first subunit comprising an amino acid sequence at least 90% or 95% identical to amino acids 20 to 1684 of SEQ ID NO:2 or 20 to 2578 of SEQ ID NO:6, and a second subunit comprising an amino acid sequence at least 90% to 95% identical to amino acids 21 to 247 of SEQ ID NO:4.

E37. The pharmaceutical composition of embodiment E36, wherein the rFVIII_{FC} polypeptide comprises a first subunit comprising amino acids 20 to 1684 of SEQ ID NO:2 or 20 to 2578 of SEQ ID NO:6, and a second subunit comprising amino acids 21 to 247 of SEQ ID NO:4.

E38. The pharmaceutical composition of any one of embodiments E1 to E37, wherein the FVIII polypeptide is present at a concentration of about 50 IU/ml to about 1500 IU/ml.

E39. The pharmaceutical composition of embodiment E38, comprising 83 IU/ml, 167 IU/ml, 250 IU/ml, 333 IU/ml, 500 IU/ml, 667 IU/ml, or 1000 IU/ml of the FVIII polypeptide.

E40. The pharmaceutical composition of any one of embodiments E1 to E37, wherein the FVIII polypeptide is present at a concentration of about 100 IU/ml to about 2500 IU/ml.

E41. The pharmaceutical composition of embodiment E40, comprising 150 IU/ml, 287.5 IU/ml, 431.25 IU/ml, 575 IU/ml, 862.5 IU/ml, 1150 IU/ml, or 1725 IU/ml of the FVIII polypeptide.

E42. The pharmaceutical composition of any one of embodiments E1 to E4, comprising:

- (a) about 50 IU/ml to about 1500 IU/ml of the FVIII polypeptide;
- (b) about 1% (w/v) to about 2.5% (w/v) of sucrose;
- (c) about 150 mM to about 250 mM NaCl;
- (d) about 5 mM to about 15 mM L-histidine;
- (e) about 5 mM to about 10 mM calcium chloride; and
- (f) about 0.008% (w/v) to about 0.025% of polysorbate 20 or polysorbate 80.

E43. The pharmaceutical composition of embodiment E42, comprising about 175 mM to about 225 mM NaCl.

E44. The pharmaceutical composition of embodiment E43, comprising about 200 mM to about 210 mM NaCl.

E45. The pharmaceutical composition of embodiment E44, comprising:

- (a) about 83 IU/ml, about 167 IU/ml, about 250 IU/ml, about 333 IU/ml, about 500 IU/ml, about 667 IU/ml, or about 1000 IU/ml of the FVIII polypeptide;
- (b) about 1.33% (w/v) of sucrose;
- (c) about 205 mM NaCl;
- (d) about 6.64 mM L-histidine;
- (e) about 5.4 mM calcium chloride; and
- (f) about 0.013% (w/v) of polysorbate 20 or polysorbate 80.

E46. The pharmaceutical composition of any one of embodiments E1 to E4, comprising:

- (a) about 100 IU/ml to about 2500 IU/ml of the FVIII polypeptide;
- (b) about 1% (w/v) to about 2.5% (w/v) of sucrose;
- (c) about 250 mM to about 350 mM NaCl;
- (d) about 5 mM to about 15 mM L-histidine;
- (e) about 5 mM to about 10 mM calcium chloride; and
- (f) about 0.008% (w/v) to about 0.025% of polysorbate 20 or polysorbate 80.

E47. The pharmaceutical composition of embodiment E46, comprising about 275 mM to about 325 mM NaCl.

E48. The pharmaceutical composition of embodiment E47, comprising:

- (a) about 150 IU/ml, about 287.5 IU/ml, about 431.25 IU/ml, about 575 IU/ml, about 862.5 IU/ml, about 1150 IU/ml, or about 1725 IU/ml of the g FVIII polypeptide;
- (b) about 2.0% (w/v) of sucrose;
- (c) about 308 mM NaCl;
- (d) about 9.8 mM L-histidine;
- (e) about 8 mM calcium chloride; and
- (f) about 0.020% (w/v) of polysorbate 20 or polysorbate 80.

E49. The pharmaceutical composition of any one of embodiments E1 to E4, comprising:

- (a) about 50 IU/ml to about 1500 IU/ml of the FVIII polypeptide;
- (b) about 10 mg/ml to about 25 mg/ml of sucrose;
- (c) about 8.8 mg/ml to about 14.6 mg/ml NaCl;
- (d) about 0.75 mg/ml to about 2.25 mg/ml L-histidine;
- (e) about 0.75 mg/ml to about 1.5 mg/ml calcium chloride dihydrate; and
- (f) about 0.08 mg/ml to about 0.25 mg/ml of polysorbate 20 or polysorbate 80.

E50. The pharmaceutical composition of embodiment E49, comprising about 10 mg/ml to 13 mg/ml NaCl.

E51. The pharmaceutical composition of embodiment E50, comprising:

- (a) about 83 IU/ml, about 167 IU/ml, about 250 IU/ml, about 333 IU/ml, about 500 IU/ml, about 667 IU/ml, or about 1000 IU/ml of the FVIII polypeptide;
- (b) about 13.3 mg/ml of sucrose;
- (c) about 12.0 mg/ml NaCl;
- (d) about 1.03 mg/ml L-histidine;
- (e) about 0.8 mg/ml calcium chloride dihydrate; and
- (f) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.

E52. The pharmaceutical composition of any one of embodiments E1 to E4, comprising:

- (a) about 100 IU/ml to about 2500 IU/ml of the FVIII polypeptide;
- (b) about 10 mg/ml to about 25 mg/ml of sucrose;
- (c) about 14.6 mg/ml to about 20.5 mg/ml NaCl;
- (d) about 0.75 mg/ml to about 2.25 mg/ml L-histidine;

61

- (e) about 0.75 mg/ml to about 1.5 mg/ml calcium chloride dihydrate; and
 - (f) about 0.08 mg/ml to about 0.25 mg/ml of polysorbate 20 or polysorbate 80.
- E53. The pharmaceutical composition of embodiment 5
E52, comprising about 16 mg/ml to about 19 mg/ml NaCl.
- E54. The pharmaceutical composition of embodiment
E53, comprising:
- (a) about 150 IU/ml, about 287.5 IU/ml, about 431.25 IU/ml, about 575 IU/ml, about 862.5 IU/ml, about 1150 IU/ml, or about 1725 IU/ml of the FVIII polypeptide; 10
 - (b) about 20.0 mg/ml of sucrose;
 - (c) about 18.0 mg/ml NaCl;
 - (d) about 1.55 mg/ml L-histidine;
 - (e) about 1.18 mg/ml calcium chloride dihydrate; and 15
 - (f) about 0.20 mg/ml of polysorbate 20 or polysorbate 80.
- E55. A pharmaceutical kit comprising:
- (a) a first container comprising a lyophilized powder, where the powder comprises 20
 - (i) the FVIII polypeptide,
 - (ii) one or more stabilizing agents selected from the group consisting of sucrose, trehalose, raffinose, arginine, and mixture thereof;
 - (iii) sodium chloride (NaCl);
 - (iv) L-histidine;
 - (v) calcium chloride; and
 - (vi) polysorbate 20 or polysorbate 80; and
 - (b) a second container comprising sterilized water for injections to be combined with the lyophilized powder of the first container. 30
- E56. The pharmaceutical kit of embodiment E55, wherein mannitol, glycine, alanine, or hydroxyethyl starch is not included.
- E57. The pharmaceutical kit of embodiment E55 or E56, 35 wherein NaCl is the only bulking agent.
- E58. The pharmaceutical kit of any one of embodiments E55 to E57, comprising:
- (a) a first container comprising a lyophilized powder, where the powder comprises 40
 - (i) about 250 IU, about 500 IU, about 750 IU, about 1000 IU, about 1500 IU, about 2000 IU, or about 3000 IU of the FVIII polypeptide,
 - (ii) about 40 mg of sucrose;
 - (iii) about 36 mg of sodium chloride;
 - (iv) about 3.1 mg of L-histidine;
 - (v) about 2.40 mg of calcium chloride dihydrate; and
 - (v) about 0.40 mg of polysorbate 20 or polysorbate 80; and
 - (b) a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising: 50
 - (i) about 83 IU/ml, about 167 IU/ml, about 250 IU/ml, about 333 IU/ml, about 500 IU/ml, about 667 IU/ml, or about 1000 IU/ml of the FVIII polypeptide, respectively;
 - (ii) about 1.33% (w/v) of sucrose;
 - (iii) about 205 mM NaCl;
 - (iv) about 6.64 mM L-histidine;
 - (v) about 5.4 mM of calcium chloride; and
 - (vi) about 0.013% (w/v) of polysorbate 20 or polysorbate 80.
- E59. The pharmaceutical kit of any one of embodiments E55 to E57, comprising: 55
- (a) a first container comprising a lyophilized powder, where the powder comprises

62

- (i) about 250 IU, about 500 IU, about 750 IU, about 1000 IU, about 1500 IU, about 2000 IU, or about 3000 IU of the FVIII polypeptide,
 - (ii) about 40 mg of sucrose;
 - (iii) about 36 mg of sodium chloride;
 - (iv) about 3.1 mg of L-histidine;
 - (v) about 2.40 mg of calcium chloride dihydrate; and
 - (v) about 0.40 mg of polysorbate 20 or polysorbate 80; and
- (b) a second container comprising sterilized water for injections at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution comprising:
 - (i) about 83 IU/ml, about 167 IU/ml, about 250 IU/ml, about 333 IU/ml, about 500 IU/ml, about 667 IU/ml, or about 1000 IU/ml of the FVIII polypeptide, respectively;
 - (ii) about 13.3 mg/ml of sucrose;
 - (iii) about 12.0 mg/ml of NaCl;
 - (iv) about 1.03 mg/ml of L-histidine;
 - (v) about 0.80 mg/ml of calcium chloride; and
 - (vi) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.
- E60. The pharmaceutical kit of any one of embodiments E55 to E59, wherein the first container is a glass vial comprising a rubber stopper.
- E61. The pharmaceutical kit of any one of embodiments E55 to E112, wherein the second container is a syringe body, and wherein the syringe body is associated with a plunger.
- E62. The pharmaceutical kit of embodiment E61, further comprising an adaptor to connect the glass vial to the syringe body.
- E63. The pharmaceutical kit of embodiment E61 or E62, further comprising infusion tubing associated with a needle to be connected to the syringe, suitable for intravenous infusion.
- E64. A method of reducing the annualized bleeding rate in a subject having hemophilia comprising administering to the subject the pharmaceutical composition of any one of embodiments E1 to E54 at a dosing interval of about three days or longer.
- E65. The method of embodiment E64, wherein the administration is prophylactic and individualized for the subject resulting in an annualized bleeding rate less than about 5.0, less than about 4.9, less than about 4.8, less than about 4.7, less than about 4.6, or less than about 4.5.
- E66. The method of embodiment E65, wherein the median annualized bleeding rate is about 1.6.
- E67. The method of embodiment E64, wherein the administration is prophylactic and weekly resulting in an annualized bleeding rate less than about 9.0, less than about 8.9, less than about 8.8, less than about 8.7, less than about 8.6, less than about 8.5, or less than about 8.4.
- E68. The method of embodiment E67, wherein the median annualized bleeding rate is about 3.6.
- E69. The method of embodiment E64, wherein the administration is on-demand or episodic resulting in an annualized bleeding rate less than about 55, less than about 54, less than about 53, less than about 52, less than about 51, less than about 50, less than about 49, less than about 48, or less than about 47.
- E70. The method of embodiment E69, wherein the median annualized bleeding rate is about 33.6.
- E71. The method of any one of embodiments E64 to E70, wherein the effective dose is between about 20 IU/kg to about 90 IU/kg. 65

E72. The method of any one of embodiments E64 to E71, wherein the effective dose is 20-30 IU/kg, 30-40 IU/kg, 40-50 IU/kg, 50-60 IU/kg, 60-70 IU/kg, 70-80 IU/kg, or 80-90 IU/kg.

E73. The method of any one of embodiments E64 to E72, wherein the effective dose is 20 IU/kg, 25 IU/kg, 30 IU/kg, 35 IU/kg, 40 IU/kg, 45 IU/kg, 50 IU/kg, 55 IU/kg, 60 IU/kg, 65 IU/kg, 70 IU/kg, 75 IU/kg, 80 IU/kg, 85 IU/kg, or 90 IU/kg.

E74. The method of any of embodiments E71 to E73, wherein the administration is prophylactic and individualized at an effective dose of about 25 IU/kg to about 65 IU/kg twice weekly or every three days or about 50 IU/kg to about 65 IU/kg every 4 or 5 days.

E75. The method of embodiment E74, wherein a first dose of the FVIII polypeptide is administered is about 25 IU/kg and a second dose of the FVIII polypeptide is about 50 IU/kg.

E76. The method of any of embodiments E71 to E73, wherein the administration is prophylactic and weekly at an effective dose of 65 IU/kg weekly.

E77. The method of any of embodiments E71 to E73, wherein the administration is on-demand or episodic at an effective dose of 10 IU/kg to 75 IU/kg every 12 to 24 hours.

E78. The method of any one of embodiments E64 to E70, wherein the effective dose is a fixed dose, which is standard across all body weights.

E79. The method of embodiment E78, wherein the fixed dose is about 2,000 IU, about 2,500 IU, about 3,000 IU, about 3,500 IU, or about 4,000 IU per dose.

E80. The method of any one of embodiments E64 to E70, wherein the effective dose is a stratified dose.

E81. The method of any one of embodiments E64 to E70, wherein a trough level of the FVIII polypeptide is above 1%, above 2%, or above 3% of normal.

E82. The method of any one of embodiments E64 to E81, wherein the FVIII polypeptide has a $T_{1/2 \text{ beta}}$ (activity) of about 7 hours to about 48 hours, about 6 hours to about 49 hours, about 5 hours to about 50 hours.

E83. The method of any one of embodiments E82, wherein the FVIII has a $T_{1/2 \text{ beta}}$ (activity) mean of at least about 15 hours, at least about 16 hours, at least about 17 hours, at least about 18 hours, at least about 19 hours, at least about 20 hours, at least about 21 hours, at least about 22 hours, at least about 23 hours, at least about 24 hours, at least about 25 hours.

E84. The method of embodiment E83, wherein the $T_{1/2 \text{ beta}}$ (activity) mean is about 19 hours.

E85. The method of embodiment E82 or E83, wherein the $T_{1/2 \text{ beta}}$ (activity) mean is at least about 1.5 fold higher than a polypeptide consisting of amino acids 20 to 1457 of SEQ ID NO:2, 20 to 2351 of SEQ ID NO: 6, or ADVATE®.

E86. The method of any one of embodiments E64 to E85, wherein the plasma trough level of the FVIII polypeptide is maintained between about 1% and about 5%, between about 1% and about 6%, between about 1% and about 7%, between about 1% and about 8%, between about 1% and about 9%, between about 1% and about 10%, between about 1% and about 11%, between about 1% and about 12%, between about 1% and about 13%, between about 1% and about 14%, between about 1% and about 15% above the baseline in the subject.

E87. The method of any one of embodiments E64 to E86, wherein the FVIII polypeptide is a long-acting FVIII polypeptide comprising a FVIII polypeptide and a heterologous moiety.

E88. The method of embodiment E87, wherein the heterologous moiety is an FcRn binding partner.

E89. The method of embodiment E88, wherein the FcRn binding partner comprises an Fc region.

E90. The method of embodiment E88 or E89, wherein the long-acting FVIII polypeptide further comprises a second FcRn binding partner.

E91. The method of embodiment E90, wherein the second FcRn binding partner comprises a second Fc region.

E92. The method of embodiment E90 or E91, wherein the FcRn binding partner and the second FcRn binding partner are associated.

E93. The method of embodiment E92, wherein the association is a covalent bond.

E94. The method of embodiment E93, wherein the covalent bond is a disulfide bond.

E95. The method of any one of embodiments E90 to E94, wherein the second FcRn binding partner is not linked to an amino acid sequence by a peptide bond.

E96. The method of any one of embodiments E64 to E95, wherein the long-acting FVIII polypeptide is FVIII monomer dimer hybrid.

E97. The method of any one of embodiments E87 to E96, wherein the FVIII polypeptide in the long-acting polypeptide is a human FVIII.

E98. The method of any one of embodiments E87 to E97, wherein the FVIII polypeptide in the long-acting polypeptide is a full-length FVIII or a B-domain deleted FVIII.

E99. The method of any one of embodiments E87 to E97, wherein the FVIII polypeptide is single chain.

E100. The method of any one of embodiments E87 to E97, wherein the FVIII polypeptide is unprocessed.

E101. The method of any one of embodiments E87 to E97, wherein the FVIII polypeptide is in two chains, a first chain comprising a heavy chain of the FVIII polypeptide and a second chain comprising a light chain of the FVIII polypeptide.

E102. The method of any one of embodiments E88 to E101, wherein the FcRn binding partner in the chimeric polypeptide is a human Fc.

E103. The method of any one of embodiments E87 to E102, wherein the FVIII polypeptide is at least 60%, 70%, 80%, 90%, 95%, 95%, 97%, 98%, 99%, or 100% identical to a FVIII amino acid sequence shown in Table 11A or 11B without a signal sequence (amino acids 20 to 1457 of SEQ ID NO: 2 or amino acids 20 to 2351 of SEQ ID NO: 6).

E104. The method of any one of embodiments E88 to E103, wherein the FcRn binding partner is at least 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to a Fc amino acid sequence shown in Table 11B without a signal sequence (amino acids 21 to 247 SEQ ID NO:4).

E105. The method of any one of embodiments E90 to E104, wherein the second FcRn binding partner in the chimeric polypeptide is a human Fc.

E106. The method of embodiment E105, wherein the second FcRn binding partner is at least 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to a Fc amino acid sequence shown in Table 11B without a signal sequence (amino acids 21 to 247 SEQ ID NO:4).

E107. The method of any one of embodiments E64 to E106, wherein the fixed dose of the clotting factor is for perioperative management of a bleeding episode.

E108. The method of any one of embodiments E64 to E107, wherein the subject is in need of controlling or preventing bleeding or bleeding episodes.

E109. The method of embodiment E108, wherein the subject is in need of controlling or preventing bleeding in minor hemorrhage, hemarthroses, superficial muscle hem-

65

orrhage, soft tissue hemorrhage, moderate hemorrhage, intramuscle or soft tissue hemorrhage with dissection, mucous membrane hemorrhage, hematuria, major hemorrhage, hemorrhage of the pharynx, hemorrhage of the retropharynx, hemorrhage of the retroperitoneum, hemorrhage of the central nervous system, bruises, cuts, scrapes, joint hemorrhage, nose bleed, mouth bleed, gum bleed, intracranial bleeding, intraperitoneal bleeding, minor spontaneous hemorrhage, bleeding after major trauma, moderate skin bruising, or spontaneous hemorrhage into joints, muscles, internal organs or the brain.

E110. The method of embodiment E109, wherein the subject is in need of management of bleeding associated with surgery or dental extraction.

E111. The method of embodiment E110, wherein the subject will undergo, is undergoing, or has undergone major surgery.

E112. The method of embodiment E111, wherein the major surgery is orthopedic surgery, extensive oral surgery, urologic surgery, or hernia surgery.

E113. The method of embodiment E112, wherein the orthopedic surgery is replacement of knee, hip, or other major joint.

E114. The method of anyone of embodiments E64 to E113, wherein the subject is in need of long-term treatment.

E115. The method of any one of embodiments E64 to E114, wherein the FVIII polypeptide is administered intravenously or subcutaneously.

E116. The pharmaceutical composition of any one of embodiments E18-E23, E40, E41, E46-E48, and E52-E54, wherein the pharmaceutical composition is a pre-lyophilization composition.

E117. The pharmaceutical composition of any one of embodiments E1 to E37, wherein the FVIII polypeptide is present at a concentration of about 50 IU/ml to about 2500 IU/ml.

E118. The pharmaceutical composition of embodiment E117, comprising 1333 IU/ml, 1667 IU/ml, or 2000 IU/ml of the FVIII polypeptide.

E119. The pharmaceutical composition of any one of embodiments E1 to E37, wherein the FVIII polypeptide is present at a concentration of about 100 IU/ml to about 4500 IU/ml.

E120. The pharmaceutical composition of embodiment E119, comprising 2300 IU/ml, about 2875 IU/ml, or about 3450 IU/ml of the FVIII polypeptide.

E121. The pharmaceutical composition of any one of embodiments E1 to E4, comprising:

- (a) about 50 IU/ml to about 2500 IU/ml of the FVIII polypeptide;
- (b) about 1% (w/v) to about 2.5% (w/v) of sucrose;
- (c) about 150 mM to about 250 mM NaCl;
- (d) about 5 mM to about 15 mM L-histidine;
- (e) about 5 mM to about 10 mM calcium chloride; and
- (f) about 0.008% (w/v) to about 0.025% of polysorbate 20 or polysorbate 80.

E122. The pharmaceutical composition of embodiment E121, comprising about 175 mM to about 225 mM NaCl.

E123. The pharmaceutical composition of embodiment E122, comprising about 200 mM to about 210 mM NaCl.

E124. The pharmaceutical composition of embodiment E123, comprising:

- (a) about 1333 IU/ml, about 1667 IU/ml, or about 2000 IU/ml of the FVIII polypeptide;
- (b) about 1.33% (w/v) of sucrose;
- (c) about 205 mM NaCl;
- (d) about 6.64 mM L-histidine;

66

(e) about 5.4 mM calcium chloride; and

(f) about 0.013% (w/v) of polysorbate 20 or polysorbate 80.

E125. The pharmaceutical composition of any one of embodiments E1 to E4, comprising:

- (a) about 100 IU/ml to about 4000 IU/ml of the FVIII polypeptide;
- (b) about 1% (w/v) to about 2.5% (w/v) of sucrose;
- (c) about 250 mM to about 350 mM NaCl;
- (d) about 5 mM to about 15 mM L-histidine;
- (e) about 5 mM to about 10 mM calcium chloride; and
- (f) about 0.008% (w/v) to about 0.025% of polysorbate 20 or polysorbate 80.

E126. The pharmaceutical composition of embodiment E125, comprising about 275 mM to about 325 mM NaCl.

E127. The pharmaceutical composition of embodiment E126, comprising:

- (a) about 2300 IU/ml, about 2875 IU/ml, or about 3450 IU/ml of the g FVIII polypeptide;
- (b) about 2.0% (w/v) of sucrose;
- (c) about 308 mM NaCl;
- (d) about 9.8 mM L-histidine;
- (e) about 8 mM calcium chloride; and
- (f) about 0.020% (w/v) of polysorbate 20 or polysorbate 80.

E128. The pharmaceutical composition of any one of embodiments E1 to E4, comprising:

- (a) about 50 IU/ml to about 2500 IU/ml of the FVIII polypeptide;
- (b) about 10 mg/ml to about 25 mg/ml of sucrose;
- (c) about 8.8 mg/ml to about 14.6 mg/ml NaCl;
- (d) about 0.75 mg/ml to about 2.25 mg/ml L-histidine;
- (e) about 0.75 mg/ml to about 1.5 mg/ml calcium chloride dihydrate; and
- (f) about 0.08 mg/ml to about 0.25 mg/ml of polysorbate 20 or polysorbate 80.

E129. The pharmaceutical composition of embodiment E128, comprising about 10 mg/ml to 13 mg/ml NaCl.

E130. The pharmaceutical composition of embodiment E129, comprising:

- (a) about 1333 IU/ml, about 1667 IU/ml, or about 2000 IU/ml of the FVIII polypeptide;
- (b) about 13.3 mg/ml of sucrose;
- (c) about 12.0 mg/ml NaCl;
- (d) about 1.03 mg/ml L-histidine;
- (e) about 0.8 mg/ml calcium chloride dihydrate; and
- (f) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.

E131. The pharmaceutical composition of any one of embodiments E1 to E4, comprising:

- (a) about 100 IU/ml to about 4000 IU/ml of the FVIII polypeptide;
- (b) about 10 mg/ml to about 25 mg/ml of sucrose;
- (c) about 14.6 mg/ml to about 20.5 mg/ml NaCl;
- (d) about 0.75 mg/ml to about 2.25 mg/ml L-histidine;
- (e) about 0.75 mg/ml to about 1.5 mg/ml calcium chloride dihydrate; and
- (f) about 0.08 mg/ml to about 0.25 mg/ml of polysorbate 20 or polysorbate 80.

E132. The pharmaceutical composition of embodiment E131, comprising about 16 mg/ml to about 19 mg/ml NaCl.

E133. The pharmaceutical composition of embodiment E132, comprising:

- (a) about 2300 IU/ml, about 2875 IU/ml, or about 3450 IU/ml of the FVIII polypeptide;
- (b) about 20.0 mg/ml of sucrose,
- (c) about 18.0 mg/ml NaCl;
- (d) about 1.55 mg/ml L-histidine;

67

(e) about 1.18 mg/ml calcium chloride dihydrate; and
 (f) about 0.20 mg/ml of polysorbate 20 or polysorbate 80.
 E134. The pharmaceutical kit of any one of embodiments
 E55 to E57, comprising:

- (a) a first container comprising a lyophilized powder, 5
 where the powder comprises
 (i) about 4000 IU, about 5000 IU, or about 6000 IU of
 the FVIII polypeptide,
 (ii) about 40 mg of sucrose;
 (iii) about 36 mg of sodium chloride;
 (iv) about 3.1 mg of L-histidine;
 (v) about 2.40 mg of calcium chloride dihydrate; and
 (vi) about 0.40 mg of polysorbate 20 or polysorbate 80;
 and

- (b) a second container comprising sterilized water for 15
 injections at a volume sufficient to produce, when
 combined with the lyophilized powder of the first
 container, a solution comprising:
 (i) about 1333 IU/ml, about 1667 IU/ml, or about 2000
 IU/ml of the FVIII polypeptide, respectively;
 (ii) about 1.33% (w/v) of sucrose;
 (iii) about 205 mM NaCl;
 (iv) about 6.64 mM L-histidine;
 (v) about 5.4 mM of calcium chloride; and
 (vi) about 0.013% (w/v) of polysorbate 20 or polysor-
 bate 80.

E135. The pharmaceutical kit of anyone of embodiments
 E55 to E57, comprising:

- (a) a first container comprising a lyophilized powder, 30
 where the powder comprises
 (i) about 4000 IU, about 5000 IU, or about 6000 IU of
 the FVIII polypeptide,
 (ii) about 40 mg of sucrose;
 (iii) about 36 mg of sodium chloride;
 (iv) about 3.1 mg of L-histidine;
 (v) about 2.40 mg of calcium chloride dihydrate; and
 (vi) about 0.40 mg of polysorbate 20 or polysorbate 80;
 and

- (b) a second container comprising sterilized water for
 injections at a volume sufficient to produce, when

68

combined with the lyophilized powder of the first
 container, a solution comprising:

- (i) about 1333 IU/ml, about 1667 IU/ml, or about 2000
 IU/ml of the FVIII polypeptide, respectively;
 (ii) about 13.3 mg/ml of sucrose;
 (iii) about 12.0 mg/ml of NaCl;
 (iv) about 1.03 mg/ml of L-histidine;
 (v) about 0.80 mg/ml of calcium chloride; and
 (vi) about 0.13 mg/ml of polysorbate 20 or polysorbate
 80.

E136. The pharmaceutical kit of anyone of embodiments
 E134 to E135, wherein the first container is a glass vial
 comprising a rubber stopper.

E137. The pharmaceutical kit of anyone of embodiments
 E134 to E136, wherein the second container is a syringe
 body, and wherein the syringe body is associated with a
 plunger.

E138. The pharmaceutical kit of embodiment E137, fur-
 ther comprising an adaptor to connect the glass vial to the
 syringe body.

E139. The pharmaceutical kit of embodiment E137 or
 E138, further comprising infusion tubing associated with a
 needle to be connected to the syringe, suitable for intrave-
 nous infusion.

E140. A method of treating a bleeding disorder in a 25
 subject having hemophilia comprising administering to the
 subject the pharmaceutical composition of any one of
 embodiments E117 to E133 at a dosing interval of about
 three days or longer.

E141. The method of embodiment E140, wherein the
 pharmaceutical composition is administered for individual
 prophylaxis of hemophilia.

E142. The method of embodiment E140, wherein the
 pharmaceutical composition is administered for weekly pro-
 phylaxis of hemophilia.

E143. The method of embodiment E140, wherein the
 pharmaceutical composition is administered for episodic
 (on-demand) treatment of hemophilia.

E144. The method of embodiment E140, wherein the
 pharmaceutical composition is administered for periopera-
 tive management of hemophilia.

TABLE 10

Polynucleotide Sequences

A. B-Domain Deleted FVIIIIFc

(i) B-Domain Deleted FVIIIIFc Chain DNA Sequence (FVIII signal
 peptide underlined, Fc region in bold) (SEQ ID NO: 1,
 which encodes SEQ ID NO: 2)

```

661          A TGCAAATAGA GCTCTCCACC TGCTTCTTTC
721 TGTGCGCTTTT GCGATTCTGC TTTAGTGCCA CCAGAAGATA CTACCTGGGT GCAGTGGAAAC
781 TGTGATGGGA CTATATGCAA AGTGATCTCG GTGAGCTGCC TGTGGACGCA AGATTTCCCTC
841 CTAGAGTGCC AAAATCTTTT CCATTCAACA CCTCAGTCGT GTACAAAAAG ACTCTGTTTG
901 TAGAATTCAC GGATCACCTT TTCAACATCG CTAAGCCAAG GCCACCTCG ATGGGTCTGC
961 TAGGTCCTAC CATCCAGGCT GAGGTTTATG ATACAGTGGT CATTACACTT AAGAACATGG
1021 CTTCCCATCC TGTCAGTCTT CATGCTGTTG GTGTATCCTA CTGGAAAGCT TCTGAGGGAG
1081 CTGAATATGA TGATCAGACC AGTCAAAGGG AGAAAGAAGA TGATAAAGTC TTCCTGGTG
1141 GAAGCCATAC ATATGTCTGG CAGGTCCTGA AAGAGAATGG TCCAATGGCC TCTGACCCAC
1201 TGTGCGCTTAC CTACTCATAT CTTTCTCATG TGGACCTGGT AAAAGACTTG AATTACGGCC
1261 TCATTGGAGC CCTACTAGTA TGTAGAGAAG GGAGTCTGGC CAAGGAAAAG ACACAGACCT
1321 TGCACAAATT TATACTACTT TTTGCTGTAT TTGATGAAGG GAAAGTTGG CACTCAGAAA
1381 CAAAGAACTC CTTGATGCAG GATAGGGATG CTGCATCTGC TCGGCGCTGG CCTAAATATGC
1441 ACACAGTCAA TGGTTATGTA AACAGGTCTC TGCCAGGTCT GATTGGATGC CACAGGAAAT
1501 CAGTCTATTG GCATGTGATT GGAATGGGCA CCACTCCTGA AGTGCACCTA ATATTCCTCG
1561 AAGGTCACAC ATTTCTTGTG AGGAACCATC GCCAGGCGTC CTTGGAAATC TCGCCAATAA
1621 CTTTCCCTTAC TGCTCAAACA CTCTTGATGG ACCTTGGACA GTTTCTACTG TTTGTGCTATA
1681 TCTCTTCCCA CCAACATGAT GGCATGGAAG CTTATGTCAA AGTAGACAGC TGTCAGAGG
1741 AACCCCAACT ACGAATGAAA AATAATGAAG AAGCGGAAGA CTATGATGAT GATCTTACTG
1801 ATTCTGAAAT GGATGTGGTC AGGTTTGATG ATGACAACCT 7CCTTCCTTT ATCCAAATTC
1861 GCTCAGTTGC CAAGAAGCAT CCTAAAACCT GGGTACATTA CATTGCTGCT GAAGAGGAGG
1921 ACTGGGACTA TGCTCCCTTA GTCCTCGCCC CCGATGACAG AAGTTATAAA AGTCAATATT

```

TABLE 10-continued

Polynucleotide Sequences					
1981	TGAACAATGG	CCCTCAGCGG	ATTGGTAGGA	AGTACAAAAA	AGTCCGATTT
2041	CAGATGAAAC	CTTTAAGACT	CGTGAAGCTA	TTCAGCATGA	ATCAGGAATC
2101	TACTTTATGG	GGAGTTGGA	GACACACTGT	TGATTATATT	TAAGAATCAA
2161	CATATAACAT	CTACCCCTCAC	GGAACTACTG	ATGTCCGTCC	TTTGTATTCA
2221	CAAAAGGTGT	AAAACATTTG	AAGGATTTTC	CAATTCTGCC	AGGAGAAATA
2281	AATGGACAGT	GACTGTAGAA	GATGGGCCAA	CTAAATCAGA	TCCTCGGTGC
2341	ATTACTCTAG	TTTCGTTAAT	ATGGAGAGAG	ATCTAGCTTC	AGGACTCATT
2401	TCATCTGCTA	CAAAGAATCT	GTAGATCAAA	GAGGAAACCA	GATAATGTCA
2461	ATGTCATCCT	GTTTCTGTGA	TTTGATGAGA	ACCGAAGCTG	GTACCTCACA
2521	AACGCTTTCT	CCCCAATCCA	GCTGGAGTGC	AGCTTGAGGA	TCCAGAGTTC
2581	ACATCATGCA	CAGCATCAAT	GGCTATGTTT	TTGATAGTTT	GCAGTTGTCA
2641	ATGAGGTGGC	ATACTGGTAC	ATTCTAAGCA	TTGGAGCACA	GACTGACTTC
2701	TCTTCTCTGG	ATATACCTTC	AAACACAAAA	TGGTCTATGA	AGACACACTC
2761	CATTCTCAGG	AGAACTGTCT	TTCATGTCTG	TGGAAAAACC	AGGTCTATGG
2821	GCCACAACCT	AGACTTTCGG	AACAGAGGCA	TGACCGCCTT	ACTGAAGGTT
2881	ACAAGAACAC	TGGTGATTAT	TACGAGGACA	GTTATGAAGA	TATTTTCAGCA
2941	GTAAAAACAA	TGCCATTGAA	CCAAGAAGCT	TCTCTCAAAA	CCCACCAAGT
3001	ATCAACGGGA	ATAACTCCTG	ACTACTCTTC	AGTCAGATCA	AGAGGAAATT
3061	ATACCATATC	AGTTGAAATG	AAGAAGGAAG	ATTTTGACAT	TTATGATGAG
3121	AGATGCCCGG	CAGCTTTCAA	AAGAAAAACAC	GACACTATTT	TATTGCTGCA
3181	TCTGGGA7TA	TGGATGAGT	AGCTCCCCAC	ATGTTCTAAG	AAACAGGGCT
3241	GTGTCCCTCA	GTTCAAGAAA	GTTGTTTTCC	AGGAATTTAC	TGATGGCTCC
3301	CCTTATACCG	TGGAGAACTA	AATGAACATT	TGGGACTCCT	GGGGCCATAT
3361	AAGTTGAAGA	TAATATCATG	GTAACTTTCA	GAAATCAGGC	CTCTCGTCCC
3421	ATTCTAGCCT	TATTTCTTAT	GAGGAAGATC	AGAGGCAAGG	AGCAGAACCT
3481	TTGTCAAGCC	TAATGAAACC	AAAACTTACT	TTTGGAAAGT	GCAACATCAT
3541	CTAAAGATGA	GTTTGACTGC	AAAGCCTGGG	CTTATTTCTC	TGATGTTGAC
3601	ATGTGCACTC	AGGCCTGATT	GGACCCCTTC	TGGTCTGCCA	CACTAACACA
3661	CTCATGGGAG	ACAAGTTGAC	GTACAGGAAT	TTGCTCTGTT	TTTCACCATC
3721	CCAAAAGCTG	GTACTTCACT	GAAAATATGG	AAAGAACTG	CAGGGCTCCC
3781	AGATGGAAGA	TCCCACTTTT	AAAGAGAATT	ATCGCTTCCA	TGCAATCAAT
3841	TGGATACACT	ACCTGGCTTA	GTAATGGCTC	AGGATCAAA	GATTTCGATG
3901	GCATGGGCAG	CAATGAAAAC	ATCCATTCTA	TTCAATTCAG	TGGACATGTG
3961	GAAAAAAGA	GGAGTATAAA	ATGGCACTGT	ACAATCTCTA	TCCAGGTGTT
4021	TGGAAATGTT	AGCAAGCTAA	GCTGGAAATT	GGCGGGTGGG	ATGCCCTTAT
4081	TACATGCTGG	GATGAGCACA	CTTTTCTGG	TGTACAGCAA	TAAGTGTGAC
4141	GAATGGCTTC	TGGACACATT	AGAGATTTTC	AGATTACAGC	TTTCAGACAA
4201	GGGCCCCAAA	GCTGGCCAGA	CTTCATTATT	CCGGATCAAT	CAATGCCTGG
4261	AGCCCTTTTC	TTGGATCAAG	GTGGATCTGT	TGGCACCAAT	GATTATTTCAC
4321	CCCAGGGTGC	CCGTGAGAAG	TTCTCCAGCC	TCTACATCTC	TCAGTTTATC
4381	GTCTTGATGG	GAAAGATGG	CAGACTTATC	GAGGAAATTC	CACTGGAAAC
4441	TCTTTGGCAA	TGTGGATTCA	TCTGGGATAA	AACACAATAT	TTTTAACCCCT
4501	CTCGATACAT	CCGTTTGCAC	CCAATCTATT	ATAGCATTCG	CAGCACTCTT
4561	TGATGGGCTG	TGATTTAAAT	AGTTGCAGCA	TGCCATTGGG	AATGGAGAGT
4621	CAGATGCACA	GATTACTGCT	TCATCCTACT	TTAGCAATAT	GTTTGCACCC
4681	CAAAAGCTCG	ACTTCACCTC	CAAGGGAGGA	GTAATGCCTG	GAGACCTCAG
4741	CAAAAGAGTG	GCTGCAAGTC	GACTTCCAGA	AGACAATGAA	AGTCACAGGA
4801	AGGAGTAAAT	ATCTCTGTCT	ACCAGCATGT	ATGTGAAGGA	GTCTCTCATC
4861	AAGATGGCCA	TCAGTGGACT	CTCTTTTTTC	AGAATGGCAA	AGTAAGGTTT
4921	ATCAAGACTC	CTTCACACCT	GTGGTGAACT	CTCTAGACCC	ACCGTTACTG
4981	TTCGAATTCA	CCCCCAGAGT	TGGGTGCACC	AGATTGCCCT	GAGGATGGAG
5041	GCGAGGCACA	GGACCTCTAC	<u>GACAAAACCTC</u>	<u>ACACATGCCC</u>	<u>ACCGTGCCCA</u>
5101	TTCTGGGGCG	GCTGCAAGTC	<u>TTCTCTTTCC</u>	<u>CCCCAAAACC</u>	<u>CAAGGACACC</u>
5161	CCCGGACCCC	TGAGGTCAACA	TGCGTGGTGG	TGGACGTGAG	CCACGAAGAC
5221	AGTTCAACTG	GTACGTGGAC	GGCGTGGAGG	TGCATAATGC	CAAGACAAAG
5281	AGCAGTACAA	GAGACCTGAC	CGTGTGGTCA	CGCTCCTCAC	CGTCTGTCAC
5341	TGAATGGCAA	GGAGTACAAG	TGCAAGGTCT	CCAACAAAGC	CCTCCAGACC
5401	AAACCATCTC	CAAAGCCAAA	GGGCAGCCCC	GAGAACACCA	GGTGATACCC
5461	CCCGGGATGA	GCTGACCAAG	AACCAAGTCA	GCCTGACCTG	CCTGGTCAAA
5521	CCAGCGACAT	CGCCGTGGAG	TGGGAGAGCA	ATGGGCAGCC	GGAGAACAAC
5581	CGCCTCCCGT	GTTGGACTCC	GACGGCTCCT	TCTTCTCTTA	CAGCAAGCTC
5641	AGAGCAGGTG	GCAGCAGGGG	AACGTCTTCT	CATGCTCCGT	GATGCATGAG
5701	ACCACTACAC	GCAGAAGAGC	CTCTCCCTGT	CTCCGGGTAA	A

(ii) Fc DNA sequence (mouse Igk signal peptide underlined)
(SEQ ID NO: 3, which encodes SEQ ID NO: 4)

7981				ATGGA	GACAGACACA
8041	CTCTGCTAT	GGTACTGCT	GCTCTGGGTT	CGAGGTTCCA	CTGGTGACAA
8101	TGCCACCCGT	GCCCAGCACC	TGAATCTCTG	GGAGGACCGT	CAGTCTTCTT
8161	AAACCCAAAG	ACACCTCAT	GATCTCCCGG	ACCCCTGAGG	TCACATGCGT
8221	GTGAGCCACG	AAGACCTTGA	GGTCAAGTTC	AACCTGGTACG	TGGACGGCGT
8281	AATGCCAAGA	CAAAGCCGGG	GGAGGAGCAG	TACAACAGCA	CGTACCGTGT
8341	CTCACCCTGC	TGCACCAGGA	CTGGCTGAAT	GGCAAGGAGT	ACAAGTGCAA
8401	AAAGCCCTCC	GCTGACCAAG	AACCAAGTCA	GCCTGACCTG	CCTGGTCAAA
8461	CCAGCGACAT	CGCCGTGGAG	TGGGAGAGCA	ATGGGCAGCC	GGAGAACAAC
8521	ACCTGCCTGG	TCAAAGGCTT	CTATCCCAGC	GACATCGCCG	TGGAGTGGGA
8581	CAGCCGGAGA	ACAACCTACA	GACCACGCCT	CCCGTGTTGG	ACTCCGACGG
8641	CTCTACAGCA	AGCTCACCGT	GGACAGAGAG	AGGTGGCAGC	AGGGGACAGT

TABLE 10-continued

Polynucleotide Sequences									
8701	TCCGTGATGC	ATGAGGCTCT	GCACAACCAC	TACACGCAGA	AGAGCCTCTC	CCTGTCTCCG			
8761	GGTAAA								
B. Full Length FVIII ^h Fc									
(i) Full Length FVIII ^h Fc DNA Sequence (FVIII signal peptide underlined, Fc region in bold) (SEQ ID NO: 5, which encodes SEQ ID NO: 6)									
661					ATG CAAATAGAGC	TCTCCACCTG			
721	CTTCTTTCTG	TGCTTTTTCG	GATTCTGCTT	TAGTGCCACC	AGAAGATACT	ACCTGGGTGC			
781	AGTGGAACTG	TCATGGGACT	ATATGCAAAAG	TGATCTCGGT	GAGCTGCCTG	TGGACGCAAG			
841	ATTTCCTCCT	AGAGTGCCAA	AATCTTTTCC	ATTCAACACC	TCAGTCGTGT	ACAAAAAGAC			
901	TCTGTTTGTA	GAATTCACGG	ATCACCTTTT	CAACATCGCT	AAGCCAAGGC	CACCCCTGGAT			
961	GGGTCTGCTA	GGTCCTACCA	TCCAGGCTGA	GGTTTATGAT	ACAGTGGTCA	TTACACTTAA			
1021	GAACATGGCT	TCCCATCCTG	TCAGTCTTCA	TGCTGTTGGT	GTATCCTACT	GGAAAGCTTC			
1081	TGAGGGAGCT	GAATATGATG	ATCAGACCAG	TCAAAGGGAG	AAAGAAGATG	ATAAAGTCTT			
1141	CCCTGGTGGA	AGCCATACAT	ATGTCTGGCA	GGTCCTGAAA	GAGAAATGGTC	CAATGGCCTC			
1201	TGACCCACTG	TGCCTTACCT	ACTCATATCT	TTCTCATGTG	GACCTGGTAA	AAGACTTGAA			
1261	TTCAGGGCTC	ATTGGAGCCC	TACTAGTATG	TAGAGAAGGG	AGTCTGGCCA	AGGAAAAGAC			
1321	ACAGACCTTG	CACAAATTTA	TACTACTTTT	TGCTGTATTT	GATGAAGGGA	AAAGTTGGCA			
1381	CTCAGAAACA	AAGAACTCCT	TGATGCAGGA	TAGGGATGCT	GCATCTGCTC	GGGCCTGGCC			
1441	TAAATATGCA	ACAGTCAAGG	GTTATGTAAA	CAGGTCTCTG	CCAGGTCTGA	TTGGATGCCA			
1501	CAGGAAATCA	GTCATTGGC	ATGTGATTGG	AATGGGCACC	ACTCTGAAG	TGCACTCAAT			
1561	ATTCTCTGAA	GGTCACACAT	TTCTTGTGAG	GAACCATCGC	CAGGCGTCTC	TGGAATCTC			
1621	GCCAAATACT	TTCTTTACTG	CTCAACACT	CTTGATGGAC	CTTGGACAGT	TTCTACTGTT			
1681	TTGTCATATC	TCTTCCACCC	AACATGATGG	CATGGAAGCT	TATGTCAAAG	TAGACAGCTG			
1741	TCCAGAGGAA	CCCCAACTAC	GAATGAAAAA	TAATGAAGAA	GCGGAAGACT	ATGATGATGA			
1801	TCTTACTGAT	TCTGAAATGG	ATGTGGTCAG	GTTTGATGAT	GACAACTCTC	CTTCCTTTAT			
1861	CCAAATTCGC	TCAGTTGCCA	AGAAGCATCC	TAAACTTTGG	GTACATTACA	TTGCTGCTGA			
1921	AGAGGAGGAC	TGGGACTATG	CTCCCTTAGT	CCTCGCCCCC	GATGACAGAA	GTTATAAAAG			
1981	TCAATATTTG	TCCCTTGGCC	CTCAGCGGAT	TGGTAGGAAG	TACAAAAAAG	TCCGATTTAT			
2041	GGCATACACA	GATGAAACCT	TTAAGACTCG	TGAAGCTATT	CAGCATGAAT	CAGGAATCTT			
2101	GGGACCTTTA	CTTTATGGGG	AAGTTGGAGA	CACACTGTTG	ATTATATTTA	AGAATCAAGC			
2161	AAGCAGACCA	TATAACATCT	ACCCTCACGG	AATCACTGAT	GTCCGTCCTT	TGTATTCAAG			
2221	GAGATTACCA	AAAGGTGTAA	AACATTTGAA	GGATTTTCCA	ATTCTGCCAG	GAGAAATATT			
2281	CAATATATAA	TGGACAGTGA	CTGTAGAAGA	TGGGCCAACC	AAATCAGATC	CTCGGTGCCT			
2341	GACCCGCTAT	TACTCTAGTT	TCGTTAATAT	GGAGAGAGAT	CTAGCTTCAG	GACTCATTGG			
2401	CCCTCTCCTC	ATCTGTCTACA	AAGAATCTGT	AGATCAAAGA	GGAAACCAGA	TAATGTCAGA			
2461	CAAGAGGAAT	GTCATCCTGT	TTTCTGTATT	TGATGAGAAC	CGAAGCTGGT	ACCTCACAGA			
2521	GAATATACAA	CGCTTTCTCC	CCAATCCAGC	TGGAGTGCAG	CTTGAGGATC	CAGAGTTCCA			
2581	AGCCTCCAAC	ATCATGCACA	GCATCAATGG	CTATGTTTTT	GATAGTTTGC	AGTTGTGAGT			
2641	TTGTTTGCAT	GAGGTGGCAT	ACTGGTACAT	TCTAAGCATT	GGAGCACAGA	CTGACTTCCT			
2701	TTCTGTCTTC	TCTCTCGGAT	ATACCTTCAA	ACACAAAATG	CTCTATGAAG	ACACACTTAT			
2761	CCTATTCCTA	TTCTCAGGAG	AACTGTCTT	CATGTCGATG	GAAAACCCAG	GTCTATGGAT			
2821	TCTGGGGTGC	CACAACTCAG	ACTTTCGGAA	CAGAGGCATG	ACCGCCTTAC	TGAAGGTTTC			
2881	TAGTTGTGAC	AAGAACACTG	GTGATTATTA	CGAGGACAGT	TATGAAGATA	TTTCAGCATA			
2941	CTTGCTGAGT	AAAAACAATG	CCATTGAACC	AAGAAGCTTC	TCCCAGAAAT	CAAGACACCC			
3001	TAGCACTAGG	CAAAAGCAAT	TTAATGCCAC	CACAATTTCCA	GAAAATGACA	TAGAGAAGAC			
3061	TGACCCCTGG	TTTGCACACA	GAACACCTAT	GCCTAAAATA	CAAAATGTCT	CCTCTAGTGA			
3121	TTTGTGTGAT	CTCTTGCGAC	AGAGTCTTAC	TCCACATGGG	CTATCCTTAT	CTGATCTCCA			
3181	AGAAGCCAAA	TATGAGACTT	TTTCTGATGA	TCCATCACCT	GGAGCAATAG	ACAGTAATAA			
3241	CAGCCTGTCT	GAAATGACAC	ACTTCAGGCC	ACAGCTCCAT	CACAGTGGGG	GACTGGTATT			
3301	TACCCCTGAG	TCAGGCCTCC	AATTAAGATT	AAATGAGAAA	CTGGGGACAA	CTGCAGCAAC			
3361	AGAGTTGAAG	AAACTTGATT	TCAAAGTTTC	TAGTACATCA	AATAATCTGA	TTTCAACAAT			
3421	TCCATCAGAG	AATTTGGCAG	CAGGTACTGA	TAATACAAGT	TCCTTAGGAC	CCCCAAGTAT			
3481	GCCAGTTCAT	TATGATAGTC	AATTAGATAC	CACCTCTATT	GGCAAAAAGT	CATCTCCCTT			
3541	TACTGAGTCT	GGTGGACCTC	TGAGCTTGAG	TGAAGAAAAT	AATGATTCAA	AGTTGTTAGA			
3601	ATCAGGTTTA	AGACCAAGAA	AAGAAAGTTC	ATGGGGAAAA	AATGTATCGT	CAACAGAGAG			
3661	TGGTAGGTTA	TTTAAAGGGA	AAAGAGCTCA	TGGACCTGCT	TTGTTGACTA	AAGATAATGC			
3721	CTTATTCAAA	GTTAGCATCT	CTTTGTTAAA	GACAAACAAA	ACTTCCAATA	ATTCAGCAAC			
3781	TAAATAGAAAG	ACTCACATTG	ATGGCCCATC	ATTATTAATT	GAGAATAGTC	CATCAGTCTG			
3841	GCAAAATATA	TTAGAAAGTG	ACACTGAGTT	TAAAAAAGTG	ACACCTTTGA	TTCATGACAG			
3901	AATGCTTATG	GACAAAAATG	CTACAGCTTT	GAGGCTAAAT	CATATGTCAA	ATAAACTAC			
3961	TTTCATCAAAA	AACATTCGAA	TGGTCCAACA	GAAAAAAGAG	GGCCCCATTC	ACCCAGATGC			
4021	ACAAAATCCA	GATATGTCGT	TCTTTAAGAT	GCTATTCTTG	CCAGAATCAG	CAAGGTGGAT			
4081	ACAAAGGACT	CATGAAAAGA	ACTCTCTGAA	CTCTGGGGAA	GGCCCCAGTC	CAAAGCAATT			
4141	AGATATCCTT	TGACCTTACG	AATCTGTGGA	AGGTGAGAA	TTCTTGTCTG	AGAAAAACAA			
4201	AGTGGTAGTA	GGAAAGGGTG	AATTTACAAA	GGACGTAGGA	CTCAAAGAGA	TGGTTTTTCC			
4261	AAGCAGCAGA	AACCTATTTC	TTACTAACTT	GGATAATTTA	CATGAAAATA	ATACACACAA			
4321	TCAGAGAAAA	AAAATTCAGG	AAGAAATAGA	AAAGAAGGAA	ACATTAAATC	AAGAGAATGT			
4381	AGTTTTGCCT	CAGATACATA	CAGTGACTGG	CACATAAGAA	TTTCATGAAGA	ACCTTTTCTT			
4441	ACTGAGCACT	AGGCAAAATG	TAGAAGGTTC	ATATGACGGG	GCATATGCTC	CAGTACTTCA			
4501	AGATTTTAGG	TGATTTTACG	ATTCAACAAA	TAGAACAAAG	AAACACACAG	CTCATTTCTC			
4561	AAAAAAGGGG	GAGGAAGAAA	ACTTGGAAGG	CTTGGGAAAT	CAAACCAAGC	AAATGTGAGA			
4621	GAATATGCA	TGCACCACAA	GGATATCTCC	TAATACAAGC	CAGCAGAATT	TTGTCACGCA			
4681	ACGTAGTAAG	AGAGCTTTGA	AACAATTGAG	ACTCCCACTA	GAAGAAACAG	AACTTGAAAA			
4741	AAGGATAAAT	GTGGATGACA	CCTCAACCCA	GTGGTCCAAA	AACATGAAAC	ATTTGACCCC			
4801	GAGCACCCTC	ACACAGATAG	ACTACAATGA	GAAGGAGAAA	GGGGCCATTA	CTCAGTCTCC			
4861	CTTATCAGAT	TGCCTTACGA	GGAGTCATAG	CATCCCTCAA	GCAAAATAGAT	CTCCATTACC			
4921	CATTGCAAG	GTATCATCAT	TTCCATCTAT	TAGACCTATA	TATCTGACCA	GGGTCTTATT			

TABLE 10-continued

Polynucleotide Sequences						
4981	CCAAGACAAC	TCTTCTCATC	TTCCAGCAGC	ATCTTATAGA	AAGAAAGATT	CTGGGGTCCA
5041	AGAAAGCAGT	CATTTCTTAC	AAGGAGCCAA	AAAAAATAAC	CTTCTTTTAG	CCATTCTAAC
5101	CTTGGAGATG	ACTGGTGATC	AAAGAGAGGT	TGGCTCCCTG	GGGACAAGTG	CCACAATTC
5161	AGTCACATAC	AAGAAAGTTG	AGAACACTGT	TCTCCCGAAA	CCAGACTTGC	CCAAAACATC
5221	TGGCAAAAGT	GAATTGCTTC	CAAAAGTTCA	CATTATCAG	AAGGACCTAT	TCCCTACGGA
5281	AACTAGCAAT	GGGTCTCCTG	GCCATCTGGA	TCTCGTGGAA	GGGAGCCTTC	TTCAGGGAAAC
5341	AGAGGGAGCG	ATTAAGTGGA	ATGAAGCAAA	CAGACCTGGA	AAAGTTCCCT	TTCTGAGAGT
5401	AGCAACAGAA	AGCTCTGCAA	AGACTCCCTC	CAAGCTATTG	GATCCTCTTG	CTTGGGATAA
5461	CCACTATGGT	ACTCAGATAC	CAAAAGAAGA	GTGGAAATCC	CAAGAGAAGT	CACCAGAAAA
5521	AACAGCTTTT	AAGAAAAAGG	ATACCATTTT	GTCCCTGAAC	GCTTGTGAAA	GCAATCATGC
5581	AATAGCAGCA	ATAAATGAGG	GACAAAAATA	GCCCGAAATA	GAAGTCACCT	GGGCAAGACA
5641	AGGTAGGACT	GAAAGGCTGT	GCTCTCAAAA	CCCACCAAGT	TTGAAACGCC	ATCAACGGGA
5701	AATAACTCGT	ACTACTCTTC	AGTCAGATCA	AGAGGAAATT	GACTATGATG	ATACCATATC
5761	AGTTGAAATG	AAGAAAGGAA	ATTTTGACAT	TTATGATGAG	GATGAAAAAT	AGAGCCCCCG
5821	CAGCTTTCAA	AAGAAAACAC	GACACTATTT	TATTGCTGCA	GTGGAGAGGC	TCTGGGATTA
5881	TGGGATGAGT	AGCTCCCCAC	ATGTTCTAAG	AAACAGGGCT	CAGAGTGGCA	GTGTCCTCA
5941	GTTCAAGAAA	TTGTGTTTCC	AGGAATTTAC	TGATGGCTCC	TTTACTCAGC	CCTTATACCG
6001	TGGAGAATA	AATGAACATT	TGGGACTCCT	GGGGCCATAT	ATAAGAGCAG	AAGTTGAAGA
6061	TAATATCATG	GTAACTTTCA	GAAATCAGGC	CTCTCGTCCC	TATTCCTTCT	ATTCTAGCCT
6121	TATTTCTTAT	GAGGAAGATC	AGAGGCAAGG	AGCAGAACCT	AGAAAAAACT	TTGTCAAGCC
6181	TAATGAAACC	AAAACTTACT	TTTGAAAGT	GCAACATCAT	ATGGCACCCA	CTAAGATGA
6241	GTTTGACTGC	AAAGCCTGGG	CTTATTTCTC	TGATGTTGAC	CTGAAAAAAG	ATGTCGACTC
6301	AGGCCTGATT	AGACCCCTTC	TGGTCTGCCA	CACATAACACA	CTGAACCCCTG	CTCATGGGAG
6361	ACAAGTGACA	GTACAGGAAT	TTGCTCTGTT	TTTCACCATC	TTTGATGAGA	CCAAAAGCTG
6421	GTACTTCACT	GAAAAATATG	AAAGAACTG	CAGGGCTCCC	TGCAATATCC	AGATGGAAGA
6481	TCCCACTTTT	AAAGAGAATT	ATCGCTTCCA	TGCAATCAAT	GGCTACATAA	TGGATACACT
6541	ACCTGGGTTA	GTAATGGCTC	AGGATCAAAG	GATTCTGATG	TATCTGCTCA	GCATGGGCAG
6601	CAATGAAAAC	ATCCATTCTA	TTCATTTTCT	TGGACATGTG	TTCACGTGAC	GAAAAAAGA
6661	GGAGTATAAA	ATGGCACTGT	ACAATCTCTA	TCCAGGTGTT	TTTGAGACAG	TGGAATGTT
6721	ACCATCCAAA	GCTGGAATTT	GGCGGGTGGA	ATGCCTTATT	GGCGAGCATC	TACATGCTGG
6781	GATGAGCACA	CTTTTCTTGG	TGTACAGCAA	TAAGTGTCTG	ACTCCCTTGG	GAATGGCTTC
6841	TGGACACATT	AGAGATTTTC	AGATTACAGC	TTCAGGACAA	TATGGACAGT	GGGCCCCCAA
6901	GCTGGCCAGA	CTTCATTATT	CCGGATCAAT	CAATGCCTGG	AGCACCAGAG	AGCCCTTTTC
6961	TTGGATCAAG	GTGGATCTGT	TGGCACCAAT	GATTATTTAC	GGCATCAAGA	CCCAGGGTGC
7021	CCGTCAGAAG	TTCTCCAGCC	TCTACATCTC	TCAGTTTATC	ATCATGTATA	GTCTTGATGG
7081	GAAGAAGTGG	CAGACTTATC	GAGGAAATTC	CACCTGGAAC	TTAATGGTCT	TCTTTGGCAA
7141	TGTGGATTCA	TCTGGGATAA	AACACAATAT	TTTAAACCTT	CCAATTATTG	CTCGATACAT
7201	CCGTTTGCAC	CCAACCTCAT	ATAGCATTCG	CAGCACTCTT	CGCATGGAGT	TGATGGGCTG
7261	TGATTTAAAT	AGTTGCAGCA	TGCCATTGGG	AATGGAGAGT	AAAGCAATAT	CAGATGCACA
7321	GATTACTGCT	TCATCTTACT	TTACCAATAT	GTTTGCCACC	TGGTCTCCTT	CAAAAGCTCG
7381	ACTTCACCTC	CAAGGGAGGA	GTAATGCCTG	GAGACCTCAG	GTGAATAATC	CAAAAGAGTG
7441	GCTGCAAGTG	GACTTCCAGA	AGACAATGAA	AGTCACAGGA	GTAACACTCT	AGGGAGTAAA
7501	ATCTCTGCTT	ACCAGCATGT	ATGTGAAGGA	GTTCCCTATC	TCCAGCAGTC	AAGATGGCCA
7561	TCAGTGGACT	CTCTTTTTTC	AGAATGGCAA	AGTAAAGGTT	TTTCAGGGAA	ATCAAGACTC
7621	CTTCACACCT	GTGGTGAACT	CTCTAGACCC	ACCGTTACTG	ACTCGCTACC	TTCGAATTCA
7681	CCCCCAGAGT	TGGGTGCACC	AGATTGCCCT	GAGGATGGAG	GTTCTGGGCT	GCGAGGCACA
7741	GGACCTCTAC	<u>GACAAAAC</u> TC	<u>ACACATGCCC</u>	<u>ACCGTGCCCA</u>	<u>GCTCCAGAAC</u>	<u>TCCTGGGGCG</u>
7801	<u>ACCGTCAGTC</u>	<u>TTCTCTTTCC</u>	<u>CCCCAAAACC</u>	<u>CAAGGACACC</u>	<u>CTCATGATCT</u>	<u>CCCGGACCCC</u>
7861	TGAGGTACAC	TGCGTGGTGG	TGGACGTGAG	CCACGAAGAC	CCTGAGGTCA	AGTTCAACTG
7921	GTACGTGGAC	OGCGTGGAGG	TGCATAATGC	CAAGACAAAAG	CCGCGGGAGG	AGCAGTACAA
7981	CAGCACGTAC	CGTGTGGTCA	GCGTCCTCAC	CGTCCTGCAC	CAGGACTGGC	TGAATGGCAA
8041	GGAGTACAAG	TGCAAGGTCT	CCAACAAAAGC	CCTCCACGCC	CCCATCGAGA	AAACCATCTC
8101	CAAAGCCAAA	GGGAGCCCCC	GAGAACCACA	GGGTACACC	CTGCCCCCAT	CCCGGGATGA
8161	GCTGACCAAG	AAACGAGTCA	GCCTGACCTG	CCTGGTCAAA	GGCTTCTATC	CCAGCGACAT
8221	CGCGTGGAG	TGGGAGAGCA	ATGGGACGCC	GGAGAACAAC	TACAAGACCA	CGCCTCCCGT
8281	GTTGGACTCC	GACGGCTCCT	TCTTCTCTCA	CAGCAAGCTC	ACCGTGGACA	AGAGCAGGTG
8341	GCAGCAGGGG	AACGTCTTCT	CATGCTCCGT	GATGCATGAG	GCTCTGCACA	ACCACTACAC
8401	GCAGAAGAGC	CTCTCCCTGT	CTCCGGGTAA	A		

(ii) Fc (same sequence as A (ii) (SEQ ID NO: 3))]

TABLE 11

Polypeptide Sequences
A. B-Domain Deleted FVIII-Fc Monomer Hybrid (BDD FVIII-Fc monomer dimer): created by coexpressing BDD FVIII-Fc and Fc chains. Construct = HC-LC-Fc fusion. An Fc expression cassette is cotransfected with BDDFVIII-Fc to generate the BDD FVIII-Fc monomer-. For the BDD FVIII-Fc chain, the Fc sequence is shown in bold; HC sequence is shown in double underline; remaining B domain sequence is shown in italics. Signal peptides are underlined.
i) B domain deleted FVIII-Fc chain (19 amino acid signal sequence underlined) (SEQ ID NO: 2)

TABLE 11-continued

Polypeptide Sequences

MQIELSTCFFLCLLRFCFS
ATRRYYLGAVELSWDYMOSDLGELPVDARFPPRPVKSPFPNTSVVYKKTLFVEFTDHLFN
IAKPRPPWMGLLGPTIQAEVYDVTVVITLKNMASHPVSLHAVGVSYWKASEGAEYDDQTSQ
REKEDDKVPFGGSHTYVVOVLKENGPMASDPLCLTYSYLSHVDLVKDLNSGLIGALLVCR
EGSLAKEKTQTLHKFILLFAVFEDEGKSWHSETKNSLMQDRDAASARAWPKMHTVNGYVNR
SLPGLIGCHRKSVYVHWIGMGTPPEVHSIFLEGHTFLVRNHRQASLEISPIITFLTAQTLL
MDLGQFLLFCHISSHQHDGMEAYVKVDSCEPEPQLRMKNNEAEYDDDLTDSMDVVR
DDDNPSFPIQIRSVAKKHPKTWVHYIAEEEEEDWDYAPLVLPAPDDRSYKSYLNNGPQRIG
RKYKKVRFMAYTDETFKTREAIQHESGILGPLLYGEVGDITLLIFKNQASRPYNIYPHGI
TDVRPLYSRRLPKGVKHLKDFPILPGEIFKYKWTVTVEDGPTKSDPRCLTRYSSFVNME
RDLASGLIGPLLIICYKESVDQGNQIMSDKRNVLFSVFDENRSWYLTENIQRFPLNPAG
VOLEDPEFQASNMHSINGYVFDLSQLSVCLHEVAYWYILSIGAQTDFLSVFFSGYTFKH
KMVEDTLTLFPFSGETVFMSENPGWLILGCHNSDFRNRGMTALLKVVSSCDKNTGDYEE
DSYEDISAYLLSKNNAIERFSFSQNSPPVLRKHQREITRTTLQSDQEEIDYDDTISVEMKK
EDFDIYDEDENQSPRSFQKTRHYFIAAVERLWDYGMSSSPHVLNRNAQSGSVQPKKVV
FQEFDTGSFTQPLRYGELNEHLGLLGPYIRAEVDNIMVTFRNQASRPYSFYSSLIYEE
DQRQGAEPKRNFKVKNETKTYFWKVQHMMAPTKDEFDCAWAYFSDVDLEKDVHSLGIP
LLVCHTNTLNPAHGROVTVQEFALFTTFDETKSWYFTENMERNCRAPCNIQMEDPTFKE
NYRFHAINGYIMDTLPGLVMAQDQIRIRWYLLSMGSNENIHSIHFSGHVFTVRKKEEYKMA
LYNLPGVGVETVEMLPKAGIWRVECLIGEHLHAGMSTLFLVYSNKCQTPLMASGHIR
FQITAGGQYQWAPKLARLHYSGSINAWSTKEPFSWIKVDLLAPMIIHGKIQGARQKFS
SLYISQFIIMYSLPGKKWQTYRGNSTGTLMVFFGNVDSSGIKHNI FNPPIIARYIRLHPT
HYSIRSLRMELMGCDLNSCSMPLGMESKAISDAQITASSYFTNMFATWSPSKARLHLQG
RSNARWPQVNNPKENLQVDFQKTMKVTVGTTQGVKSLTSMYVKEFLISSQDGHQWTLF
FQNGKVKVQGNQPSFTPVVNSLDPPLLTRYLRIHPQSWVHQIALRMEVLGCEAQPLYDK
THTCPCPAPELLGGPSVFLFPKPKDITLMISRTPEVTCVVVDVSHEDPEVKFNWYVQGV
EVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQ
PREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDG
SFFLYSKLTVDKSRWQQGNVFSQSVMEALHNHYTQKSLSLSPGK

ii) Fc chain (20 amino acid heterologous signal peptide from mouse Igk chain underlined)

(SEQ ID NO: 4)

METDTLLLWVLLLWVPGSTG

DKHTCPCPAPELLGGPSVFLFPKPKDITLMISRTPEVTCVVVDVSHEDPEVKFNWYVD
 GVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALDAPIEKTIKAK
 GQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSD
 GSFPLYSKLTVDKSRWQQGNVFSQSVMEALHNHYTQKSLSLSPGK

B. Full length FVIIIFc monomer hybrid (Full length FVIIIFc monomer dimer): created by coexpressing FVIIIFc and Fc chains. Construct = HC-B-LC-Fc fusion. An Fc expression cassette is cotransfected with full length FVIIIFc to generate the full length FVIIIFc monomer. For the FVIIIFc chain, the Fc sequence is shown in bold; HC sequence is shown in double underlined; B domain sequence is shown in italics. Signal peptides are underlined.

i) Full length FVIIIFc chain (FVIIIFc signal peptide underlined) (SEQ ID NO: 6)

MQIELSTCFECLERFCFS

ATRRYYLGAVELSWDYMOSDLGELPVDARFPPRPVKSPFPNTSVVYKKTLFVEFTDHLFN
IAKPRPPWMGLLGPTIQAEVYDVTVVITLKNMASHPVSLHAVGVSYWKASEGAEYDDQTSQ
REKEDDKVPFGGSHTYVVOVLKENGPMASDPLCLTYSYLSHVDLVKDLNSGLIGALLVCR
EGSLAKEKTQTLHKFILLFAVFEDEGKSWHSETKNSLMQDRDAASARAWPKMHTVNGYVNR
SLPGLIGCHRKSVYVHWIGMGTPPEVHSIFLEGHTFLVRNHRQASLEISPIITFLTAQTLL
MDLGQFLLFCHISSHQHDGMEAYVKVDSCEPEPQLRMKNNEAEYDDDLTDSMDVVR
DDDNPSFPIQIRSVAKKHPKTWVHYIAEEEEEDWDYAPLVLPAPDDRSYKSYLNNGPQRIG
RKYKKVRFMAYTDETFKTREAIQHESGILGPLLYGEVGDITLLIFKNQASRPYNIYPHGI
TDVRPLYSRRLPKGVKHLKDFPILPGEIFKYKWTVTVEDGPTKSDPRCLTRYSSFVNME
RDLASGLIGPLLIICYKESVDQGNQIMSDKRNVLFSVFDENRSWYLTENIQRFPLNPAG
VOLEDPEFQASNMHSINGYVFDLSQLSVCLHEVAYWYILSIGAQTDFLSVFFSGYTFKH
KMVEDTLTLFPFSGETVFMSENPGWLILGCHNSDFRNRGMTALLKVVSSCDKNTGDYEE
DSYEDISAYLLSKNNAIERFSFSQNSRHPSTRQKQFNATTIPENDIEKTDPPFAHRTPMFK
IQNVSSDLLMLLRQSPTPHGLSLDLQEAKEYTFSDDPSPGAIDSNNSLSEMTHFRPQLH
HSGDMVFTPEESGLQLRLNEKLGTTAATELKKLDFKVSSSTNNLISTIPSDNLAAGTDNTS
SLGPPSMVPHYDSQDLDTLFGKSSSPLTESGGPLSLSEENNDSKLLESGLMNSQESSWGK
NVSSTESGRLFKGKRAHPALLTKDNALFKVSI SLLKTNKTSNNSATNRKTHIDGPSLLI
ENSPSVWQNI LESDTEFKKVTPLIHDRMLMDKNA TALRLNHSNKTSSKNMEMVQKKE
GPITPDAQNPDMSFFKMLFLPESARWIQRTHGKNSLNSGQGPSPKQLVSLGPEKSEVGQN
FLSEKNKVVGKEFTKDVGLKEMVFPSSRNFLTNLDNLHENHNHNQEKKIQEEIEKKE
TLIQENVLPQIHTVTGTFKFMKNLELLSTRQNVESGYDGAAYAPVLQDFRSLNDSTNRTK
KHTAHFSKKGEENLEGLNQTKQIVEKYACTRISPNSTQQNFVTQRSKRALKQFRLPL
EETELEKRHVDDTSQWSKNMKHLPSTLTQIDYNEKEGAIQTSPSLDCLTRSHSIPQA
NRSPFLIAKVSSFPPIRPIYLRVLFQDNSSHLPAASRYKKDGSVQESSHFLQGAKNNL
SLAILTLEMTGDQREVGLSGTSATNSVTYKVENTVLPKPDLPKTSKGKVELLPKVHIYQK
DLFPTETSNGPSGHLDLVEGSLQGTGCAIKWNEANRPKGVFLRVATESSAKTPSKLLD
PLAWDNHYGTQIPKEEWSQEKSPKTAFFKKDITLSLNACESNHAIAAINEGQNKPEIE
VTWAKQGRTERLCSQNPVLRKHQREITRTTLQSDQEEIDYDDTISVEMKKEDFDIYDE

TABLE 11-continued

Polypeptide Sequences

ENQSPRSFQKKTRHYFIAAVERLWDYGMSSSPHVLNRNRAQSGSVQPKKVVFEFTDGSF
 TQPLYRGELNEHFGLGPIYRAEVEDNIMVTFRNQASRPYSFYSSLISYEEDQRQGAEP
 KNFVKPNETKTYFWKVQHMAPTKDEFDCKAWAYFSDVDLEKDVHSGLIGPLLVCNTNL
 NPAHGRQVTVQEFALFFTI FDETKSWYFTENMERNCRAPCNIQMEDPTFKENYRFHAING
 YIMDTLPGLVMAQDQRIRWYLLSMGSNENIHSIHFSGHVFTVRKKKEEYKMALYNLYPGVF
 ETVEMLPKAGIWRVECLIGEHLHAGMSTLFLVYSNKCQTPLGMA SGHIRDQITASGQY
 IGQWAPKLARLHYSGSINAWSTKEPFSWIKYDLLAPMI IHGKTQGARQKFSSLYISQPII
 MYSLDGKKWQTYRGNSTGTLMVFFGNVDSSGIKHNI FNPPIIARYIRLHPHYSIRSTLR
 MELMGCDLNSC SEMPLMESKAISDAQITASSYFTNMFATWSPSKARLHLQGRSNAWRPQV
 NNPKEWLQVDFQKTMKVTGVTQGVKSLTSMYVKEFLISSQDGHQWTLFFQNGKVKVF
 QGNQDSFTPVVNSLDPPLTRYLRIPHQSWVHQIALRMEVLGCEAQDLYDKTHTCPPCPA
PELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKP
REEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTL
PPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSGGSFFLYSKLT
VDKSRVQQGNV FSCSVMEALHNHYTQKSLSLSPGK

ii) Fc chain (20 amino acid heterologous signal peptide
 from mouse Igk chain underlined)

(SEQ ID NO: 4)

METDTLLWVLLWVPGSTG
 DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVD
 GVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAK
 GQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDS
 DGSFFLYSKLTVDKSRVQQGNV FSCSVMEALHNHYTQKSLSLSPGK

The foregoing description of the specific embodiments
 will so fully reveal the general nature of the invention that
 others can, by applying knowledge within the skill of the art,
 readily modify and/or adapt for various applications such
 specific embodiments, without undue experimentation,
 without departing from the general concept of the present
 invention. Therefore, such adaptations and modifications are
 intended to be within the meaning and range of equivalents
 of the disclosed embodiments, based on the teaching and
 guidance presented herein. It is to be understood that the
 phraseology or terminology herein is for the purpose of
 description and not of limitation, such that the terminology

25

or phraseology of the present specification is to be inter-
 preted by the skilled artisan in light of the teachings and
 guidance.

30

Other embodiments of the invention will be apparent to
 those skilled in the art from consideration of the specifica-
 tion and practice of the invention disclosed herein. It is
 intended that the specification and examples be considered
 as exemplary only, with a true scope and spirit of the
 invention being indicated by the following claims.

35

All patents and publications cited herein are incorporated
 by reference herein in their entirety.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 18

<210> SEQ ID NO 1

<211> LENGTH: 5052

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: B-Domain Deleted FVIIIFc

<400> SEQUENCE: 1

atgcaaatag agctctccac ctgcttcttt ctgtgccttt tgcgattctg ctttagtgcc	60
accagaagat actacctggg tgcagtggaa ctgtcatggg actatatgca aagtgatctc	120
ggtgagctgc ctgtggacgc aagatttcct cctagagtgc caaaatcttt tccattcaac	180
acctcagtcg tgtacaaaaa gactctgttt gtagaattca cggatcacct tttcaacatc	240
gctaagccaa ggccaccctg gatgggtctg ctaggctcta ccatccaggc tgagggtttat	300
gatacagtgg tctattacact taagaacatg gcttcccac cctgtcagtc tcatgctgtt	360
ggtgtatcct actggaagc ttctgagggg gctgaatatg atgatacagc cagtcaaagg	420
gagaaagaag atgataaagt cttccctggg ggaagccata catatgtctg gcaggctcctg	480
aaagagaatg gtccaatggc ctctgaccca ctgtgcctta cctactcata tctttctcat	540
gtggacctgg taaaagactt gaattcaggc ctcatggag ccctactagt atgtagagaa	600

-continued

gggagtctgg	ccaaggaaaa	gacacagacc	ttgcacaaat	ttatactact	ttttgctgta	660
tttgatgaag	ggaaaagtgt	gcactcagaa	acaaagaact	ccttgatgca	ggatagggat	720
gtgcatctg	ctcgggcctg	gcctaaaatg	cacacagtca	atggttatgt	aaacaggtct	780
ctgccaggtc	tgattggatg	ccacaggaaa	tcagtctatt	ggcatgtgat	tggaatgggc	840
accactcctg	aagtgcactc	aatattcctc	gaaggtcaca	catttcttgt	gaggaaccat	900
cgccaggcgt	ccttggaat	ctgcgaata	actttcctta	ctgctcaaac	actcttgatg	960
gaccttggac	agtttctact	gttttgtcat	atctcttccc	accaacatga	tgcatggaa	1020
gcttatgtca	aagtagacag	ctgtccagag	gaaccccaac	tacgaatgaa	aaataatgaa	1080
gaagcggaa	actatgatga	tgatcttact	gattctgaaa	tgatgtggg	caggtttgat	1140
gatgacaact	ctccttcctt	tatccaaatt	cgtcagttg	ccaagaagca	tcctaaaact	1200
tgggtacatt	acattgtgct	tgaagaggag	gactgggact	atgctccctt	agtcctcgcc	1260
cccgatgaca	gaagttataa	aagtcaatat	ttgaacaatg	gccctcagcg	gattggtagg	1320
aagtacaaaa	aagtcgatt	tatggcatac	acagatgaaa	cctttaagac	tcgtgaagct	1380
attcagcatg	aatcaggaat	cttgggacct	ttactttatg	gggaagttgg	agacacactg	1440
ttgattatat	ttaagaatca	agcaagcaga	ccatataaca	tctaccctca	cggaatcact	1500
gatgtccgtc	ctttgtattc	aaggagatta	ccaaaagggtg	taaaacattt	gaaggatttt	1560
ccaattctgc	caggagaaat	attcaaatat	aaatggacag	tgactgtaga	agatgggcca	1620
actaaatcag	atcctcggtg	cctgacccgc	tattactcta	gtttcgtaa	tatggagaga	1680
gatctagctt	caggactcat	tggccctctc	ctcatctgct	acaaagaatc	tgtagatcaa	1740
agaggaaacc	agataatgtc	agacaagagg	aatgtcatcc	tgttttctgt	atttgatgag	1800
aaccgaagct	ggtaacctac	agagaatata	caacgcttcc	tccccaatcc	agctggagtg	1860
cagcttgagg	atccagagtt	ccaagcctcc	aacatcatgc	acagcatcaa	tggtatgtt	1920
tttgatagtt	tgcagttgtc	agtttgtttg	catgagggtg	catactggta	cattctaagc	1980
attggagcac	agactgactt	cctttctgtc	ttcttctctg	gatatacctt	caaacacaaa	2040
atggtctatg	aagacacact	cacctattc	ccattctcag	gagaaactgt	cttcatgtcg	2100
atggaaaacc	caggctctatg	gattctgggg	tgccacaact	cagactttcg	gaacagaggc	2160
atgaccgcct	tactgaaggt	ttctagtgtg	gacaagaaca	ctggtgatta	ttacgaggac	2220
agttatgaag	atatttcagc	atacttgctg	agtaaaaaca	atgccattga	accaagaagc	2280
ttctctcaaa	acccaccagt	cttgaaacgc	catcaacggg	aaataactcg	tactactctt	2340
cagtcagatc	aagaggaaat	tgactatgat	gataccatat	cagttgaaat	gaagaaggaa	2400
gattttgaca	tttatgatga	ggatgaaaat	cagagccccc	gcagctttca	aaagaaaaca	2460
cgaactatt	ttattgtgct	agtggagagg	ctctgggatt	atgggatgag	tagctcccca	2520
catgttctaa	gaaacagggc	tcagagtggc	agtgctccctc	agttcaagaa	agttgttttc	2580
caggaattta	ctgatggctc	ctttactcag	cccttatacc	gtggagaact	aatgaacat	2640
ttgggactcc	tggggccata	tataagagca	gaagttgaag	ataatatcat	ggtaactttc	2700
agaaatcagg	cctctcgtcc	ctattccttc	tattctagcc	ttatttctta	tgaggaagat	2760
cagaggcaag	gagcagaacc	tagaaaaaac	tttgtaagc	ctaataaac	caaaacttac	2820
ttttgaaag	tgacaatca	tatggcacc	actaaagatg	agtttgactg	caaagcctgg	2880
gcttatttct	ctgatgttga	cctggaaaaa	gatgtgcact	caggcctgat	tggacccctt	2940
ctggtctgcc	acactaacac	actgaaccct	gctcatggga	gacaagtgc	agtacaggaa	3000

-continued

```

tttgctctgt ttttcacat ctttgatgag accaaaagct ggtacttcac tgaaaatatg 3060
gaaagaaact gcagggtccc ctgcaatata cagatggaag atcccacttt taaagagaat 3120
tategcttcc atgcaatcaa tggctacata atggatacac tacctggctt agtaatggct 3180
caggatcaaa ggattcgatg gtatctgctc agcatgggca gcaatgaaaa catccattct 3240
attcatttca gtggacatgt gttcactgta cgaaaaaaag aggagtataa aatggcactg 3300
tacaatctct atccagggtg ttttgagaca gtggaaatgt taccatccaa agctggaatt 3360
tggcgggtgg aatgccttat tggcgagcat ctacatgctg ggatgagcac actttttctg 3420
gtgtacagca ataagtgtca gactccctg ggaatggctt ctggacacat tagagatttt 3480
cagattacag cttcaggaca atatggacag tgggccccaa agctggccag acttcattat 3540
tccgatcaa tcaatgctg gagcaccaag gagccctttt cttggatcaa ggtggatctg 3600
ttggcaccaa tgattattca cggcacaaag acccagggtg cccgtcagaa gttctccagc 3660
ctctacatct ctacgtttat catcatgtat agtcttgatg ggaagaagtg gcagacttat 3720
cgaggaaatt ccaactggaac cttaatggtc ttctttggca atgtggatc atctgggata 3780
aaacacaata tttttaacc tccaattatt gctcgataca tccgtttgca cccaactcat 3840
tatagcatc gcagcactct tcgcatggag ttgatgggct gtgatttaa tagttgcagc 3900
atgccattgg gaatggagag taaagcaata tcagatgcac agattactgc ttcacctac 3960
tttaccaata tgtttgccac ctggtctcct tcaaaagctc gacttcacct ccaaggagg 4020
agtaatgcct ggagacctca ggtgaataat ccaaagagt ggctgcaagt ggacttcag 4080
aagacaatga aagtcacagg agtaactact caggagtaa aatctctgct taccagcatg 4140
tatgtgaagg agttcctcat ctccagcagt caagatggcc atcagtggac tctcttttt 4200
cagaatggca aagtaaagg ttttcaggga aatcaagact ccttcacacc tgtggtgaac 4260
tctctagacc cacgttact gactcgctac cttcgaatc acccccagag ttgggtgcac 4320
cagattgccc tgaggatgga ggttctgggc tgcgaggcac aggacctcta cgacaaaact 4380
cacacatgcc cacgtgccc agctccagaa ctctgggag gaccgtcagt ctctctctc 4440
ccccaaaaa ccaaggacac cctcatgac tcccgagccc ctgaggtcac atcggtggtg 4500
gtggacgtga gccacgaaga cctgaggtc aagttcaact ggtacgtgga cgcggtggag 4560
gtgcataatg ccaagacaaa gccgcgggag gagcagtaca acagcacgta ccgtgtggtc 4620
agcgtctca ccgtctgca ccaggactgg ctgaatggca aggagtacaa gtgcaaggtc 4680
tccaacaaag ccctccagc ccccatcgag aaaaccatct ccaaagccaa agggcagccc 4740
cgagaaccac aggtgtacac cctgccccca tcccgggatg agctgaccaa gaaccaggtc 4800
agcctgacct gcctggtcaa aggtctctat cccagcgaca tcgccgtgga gtgggagagc 4860
aatgggcagc cggagaacaa ctacaagacc acgcctcccg tgttggaact cgacggtcc 4920
ttctctctct acagcaagct caccgtggac aagagcaggg ggcagcaggg gaacgtctc 4980
tcatgctccg tgatgcatga ggctctgcac aaccactaca cgcagaagag cctctccctg 5040
tctccgggta aa 5052

```

<210> SEQ ID NO 2

<211> LENGTH: 1684

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: B-Domain Deleted FVIIIFc

-continued

<400> SEQUENCE: 2

```

Met Gln Ile Glu Leu Ser Thr Cys Phe Phe Leu Cys Leu Leu Arg Phe
1           5           10           15
Cys Phe Ser Ala Thr Arg Arg Tyr Tyr Leu Gly Ala Val Glu Leu Ser
20           25           30
Trp Asp Tyr Met Gln Ser Asp Leu Gly Glu Leu Pro Val Asp Ala Arg
35           40           45
Phe Pro Pro Arg Val Pro Lys Ser Phe Pro Phe Asn Thr Ser Val Val
50           55           60
Tyr Lys Lys Thr Leu Phe Val Glu Phe Thr Asp His Leu Phe Asn Ile
65           70           75           80
Ala Lys Pro Arg Pro Pro Trp Met Gly Leu Leu Gly Pro Thr Ile Gln
85           90           95
Ala Glu Val Tyr Asp Thr Val Val Ile Thr Leu Lys Asn Met Ala Ser
100          105          110
His Pro Val Ser Leu His Ala Val Gly Val Ser Tyr Trp Lys Ala Ser
115          120          125
Glu Gly Ala Glu Tyr Asp Asp Gln Thr Ser Gln Arg Glu Lys Glu Asp
130          135          140
Asp Lys Val Phe Pro Gly Gly Ser His Thr Tyr Val Trp Gln Val Leu
145          150          155          160
Lys Glu Asn Gly Pro Met Ala Ser Asp Pro Leu Cys Leu Thr Tyr Ser
165          170          175
Tyr Leu Ser His Val Asp Leu Val Lys Asp Leu Asn Ser Gly Leu Ile
180          185          190
Gly Ala Leu Leu Val Cys Arg Glu Gly Ser Leu Ala Lys Glu Lys Thr
195          200          205
Gln Thr Leu His Lys Phe Ile Leu Leu Phe Ala Val Phe Asp Glu Gly
210          215          220
Lys Ser Trp His Ser Glu Thr Lys Asn Ser Leu Met Gln Asp Arg Asp
225          230          235          240
Ala Ala Ser Ala Arg Ala Trp Pro Lys Met His Thr Val Asn Gly Tyr
245          250          255
Val Asn Arg Ser Leu Pro Gly Leu Ile Gly Cys His Arg Lys Ser Val
260          265          270
Tyr Trp His Val Ile Gly Met Gly Thr Thr Pro Glu Val His Ser Ile
275          280          285
Phe Leu Glu Gly His Thr Phe Leu Val Arg Asn His Arg Gln Ala Ser
290          295          300
Leu Glu Ile Ser Pro Ile Thr Phe Leu Thr Ala Gln Thr Leu Leu Met
305          310          315          320
Asp Leu Gly Gln Phe Leu Leu Phe Cys His Ile Ser Ser His Gln His
325          330          335
Asp Gly Met Glu Ala Tyr Val Lys Val Asp Ser Cys Pro Glu Glu Pro
340          345          350
Gln Leu Arg Met Lys Asn Asn Glu Glu Ala Glu Asp Tyr Asp Asp Asp
355          360          365
Leu Thr Asp Ser Glu Met Asp Val Val Arg Phe Asp Asp Asp Asn Ser
370          375          380
Pro Ser Phe Ile Gln Ile Arg Ser Val Ala Lys Lys His Pro Lys Thr
385          390          395          400
Trp Val His Tyr Ile Ala Ala Glu Glu Glu Asp Trp Asp Tyr Ala Pro
405          410          415

```

-continued

Leu Val Leu Ala Pro Asp Asp Arg Ser Tyr Lys Ser Gln Tyr Leu Asn
 420 425 430

Asn Gly Pro Gln Arg Ile Gly Arg Lys Tyr Lys Lys Val Arg Phe Met
 435 440 445

Ala Tyr Thr Asp Glu Thr Phe Lys Thr Arg Glu Ala Ile Gln His Glu
 450 455 460

Ser Gly Ile Leu Gly Pro Leu Leu Tyr Gly Glu Val Gly Asp Thr Leu
 465 470 475 480

Leu Ile Ile Phe Lys Asn Gln Ala Ser Arg Pro Tyr Asn Ile Tyr Pro
 485 490 495

His Gly Ile Thr Asp Val Arg Pro Leu Tyr Ser Arg Arg Leu Pro Lys
 500 505 510

Gly Val Lys His Leu Lys Asp Phe Pro Ile Leu Pro Gly Glu Ile Phe
 515 520 525

Lys Tyr Lys Trp Thr Val Thr Val Glu Asp Gly Pro Thr Lys Ser Asp
 530 535 540

Pro Arg Cys Leu Thr Arg Tyr Tyr Ser Ser Phe Val Asn Met Glu Arg
 545 550 555 560

Asp Leu Ala Ser Gly Leu Ile Gly Pro Leu Leu Ile Cys Tyr Lys Glu
 565 570 575

Ser Val Asp Gln Arg Gly Asn Gln Ile Met Ser Asp Lys Arg Asn Val
 580 585 590

Ile Leu Phe Ser Val Phe Asp Glu Asn Arg Ser Trp Tyr Leu Thr Glu
 595 600 605

Asn Ile Gln Arg Phe Leu Pro Asn Pro Ala Gly Val Gln Leu Glu Asp
 610 615 620

Pro Glu Phe Gln Ala Ser Asn Ile Met His Ser Ile Asn Gly Tyr Val
 625 630 635 640

Phe Asp Ser Leu Gln Leu Ser Val Cys Leu His Glu Val Ala Tyr Trp
 645 650 655

Tyr Ile Leu Ser Ile Gly Ala Gln Thr Asp Phe Leu Ser Val Phe Phe
 660 665 670

Ser Gly Tyr Thr Phe Lys His Lys Met Val Tyr Glu Asp Thr Leu Thr
 675 680 685

Leu Phe Pro Phe Ser Gly Glu Thr Val Phe Met Ser Met Glu Asn Pro
 690 695 700

Gly Leu Trp Ile Leu Gly Cys His Asn Ser Asp Phe Arg Asn Arg Gly
 705 710 715 720

Met Thr Ala Leu Leu Lys Val Ser Ser Cys Asp Lys Asn Thr Gly Asp
 725 730 735

Tyr Tyr Glu Asp Ser Tyr Glu Asp Ile Ser Ala Tyr Leu Leu Ser Lys
 740 745 750

Asn Asn Ala Ile Glu Pro Arg Ser Phe Ser Gln Asn Pro Pro Val Leu
 755 760 765

Lys Arg His Gln Arg Glu Ile Thr Arg Thr Thr Leu Gln Ser Asp Gln
 770 775 780

Glu Glu Ile Asp Tyr Asp Asp Thr Ile Ser Val Glu Met Lys Lys Glu
 785 790 795 800

Asp Phe Asp Ile Tyr Asp Glu Asp Glu Asn Gln Ser Pro Arg Ser Phe
 805 810 815

Gln Lys Lys Thr Arg His Tyr Phe Ile Ala Ala Val Glu Arg Leu Trp
 820 825 830

Asp	Tyr	Gly	Met	Ser	Ser	Ser	Pro	His	Val	Leu	Arg	Asn	Arg	Ala	Gln	
		835					840					845				
Ser	Gly	Ser	Val	Pro	Gln	Phe	Lys	Lys	Val	Val	Phe	Gln	Glu	Phe	Thr	
	850					855					860					
Asp	Gly	Ser	Phe	Thr	Gln	Pro	Leu	Tyr	Arg	Gly	Glu	Leu	Asn	Glu	His	
865					870					875					880	
Leu	Gly	Leu	Leu	Gly	Pro	Tyr	Ile	Arg	Ala	Glu	Val	Glu	Asp	Asn	Ile	
				885					890					895		
Met	Val	Thr	Phe	Arg	Asn	Gln	Ala	Ser	Arg	Pro	Tyr	Ser	Phe	Tyr	Ser	
			900					905					910			
Ser	Leu	Ile	Ser	Tyr	Glu	Glu	Asp	Gln	Arg	Gln	Gly	Ala	Glu	Pro	Arg	
	915						920					925				
Lys	Asn	Phe	Val	Lys	Pro	Asn	Glu	Thr	Lys	Thr	Tyr	Phe	Trp	Lys	Val	
930						935					940					
Gln	His	His	Met	Ala	Pro	Thr	Lys	Asp	Glu	Phe	Asp	Cys	Lys	Ala	Trp	
945					950					955					960	
Ala	Tyr	Phe	Ser	Asp	Val	Asp	Leu	Glu	Lys	Asp	Val	His	Ser	Gly	Leu	
				965					970					975		
Ile	Gly	Pro	Leu	Leu	Val	Cys	His	Thr	Asn	Thr	Leu	Asn	Pro	Ala	His	
		980						985					990			
Gly	Arg	Gln	Val	Thr	Val	Gln	Glu	Phe	Ala	Leu	Phe	Phe	Thr	Ile	Phe	
	995						1000					1005				
Asp	Glu	Thr	Lys	Ser	Trp	Tyr	Phe	Thr	Glu	Asn	Met	Glu	Arg	Asn		
	1010					1015					1020					
Cys	Arg	Ala	Pro	Cys	Asn	Ile	Gln	Met	Glu	Asp	Pro	Thr	Phe	Lys		
	1025					1030					1035					
Glu	Asn	Tyr	Arg	Phe	His	Ala	Ile	Asn	Gly	Tyr	Ile	Met	Asp	Thr		
	1040					1045					1050					
Leu	Pro	Gly	Leu	Val	Met	Ala	Gln	Asp	Gln	Arg	Ile	Arg	Trp	Tyr		
	1055					1060					1065					
Leu	Leu	Ser	Met	Gly	Ser	Asn	Glu	Asn	Ile	His	Ser	Ile	His	Phe		
	1070					1075					1080					
Ser	Gly	His	Val	Phe	Thr	Val	Arg	Lys	Lys	Glu	Glu	Tyr	Lys	Met		
	1085					1090					1095					
Ala	Leu	Tyr	Asn	Leu	Tyr	Pro	Gly	Val	Phe	Glu	Thr	Val	Glu	Met		
	1100					1105					1110					
Leu	Pro	Ser	Lys	Ala	Gly	Ile	Trp	Arg	Val	Glu	Cys	Leu	Ile	Gly		
	1115					1120					1125					
Glu	His	Leu	His	Ala	Gly	Met	Ser	Thr	Leu	Phe	Leu	Val	Tyr	Ser		
	1130					1135					1140					
Asn	Lys	Cys	Gln	Thr	Pro	Leu	Gly	Met	Ala	Ser	Gly	His	Ile	Arg		
	1145					1150					1155					
Asp	Phe	Gln	Ile	Thr	Ala	Ser	Gly</									

-continued

1235	1240	1245
Met Val Phe Phe Gly Asn Val Asp Ser Ser Gly Ile Lys His Asn 1250 1255 1260		
Ile Phe Asn Pro Pro Ile Ile Ala Arg Tyr Ile Arg Leu His Pro 1265 1270 1275		
Thr His Tyr Ser Ile Arg Ser Thr Leu Arg Met Glu Leu Met Gly 1280 1285 1290		
Cys Asp Leu Asn Ser Cys Ser Met Pro Leu Gly Met Glu Ser Lys 1295 1300 1305		
Ala Ile Ser Asp Ala Gln Ile Thr Ala Ser Ser Tyr Phe Thr Asn 1310 1315 1320		
Met Phe Ala Thr Trp Ser Pro Ser Lys Ala Arg Leu His Leu Gln 1325 1330 1335		
Gly Arg Ser Asn Ala Trp Arg Pro Gln Val Asn Asn Pro Lys Glu 1340 1345 1350		
Trp Leu Gln Val Asp Phe Gln Lys Thr Met Lys Val Thr Gly Val 1355 1360 1365		
Thr Thr Gln Gly Val Lys Ser Leu Leu Thr Ser Met Tyr Val Lys 1370 1375 1380		
Glu Phe Leu Ile Ser Ser Ser Gln Asp Gly His Gln Trp Thr Leu 1385 1390 1395		
Phe Phe Gln Asn Gly Lys Val Lys Val Phe Gln Gly Asn Gln Asp 1400 1405 1410		
Ser Phe Thr Pro Val Val Asn Ser Leu Asp Pro Pro Leu Leu Thr 1415 1420 1425		
Arg Tyr Leu Arg Ile His Pro Gln Ser Trp Val His Gln Ile Ala 1430 1435 1440		
Leu Arg Met Glu Val Leu Gly Cys Glu Ala Gln Asp Leu Tyr Asp 1445 1450 1455		
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly 1460 1465 1470		
Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu 1475 1480 1485		
Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val 1490 1495 1500		
Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly 1505 1510 1515		
Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr 1520 1525 1530		
Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln 1535 1540 1545		
Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys 1550 1555 1560		
Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly 1565 1570 1575		
Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp 1580 1585 1590		
Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly 1595 1600 1605		
Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln 1610 1615 1620		
Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp 1625 1630 1635		

-continued

Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg
1640 1645 1650

Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala
1655 1660 1665

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
1670 1675 1680

Lys

<210> SEQ ID NO 3
<211> LENGTH: 741
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Fc Portion

<400> SEQUENCE: 3

```
atggagacag acacactcct gctatgggta ctgctgctct gggttccagg ttccactggt      60
gacaaaaactc acacatgccc accgtgccca gcacctgaac tcctgggagg accgtcagtc      120
ttcctctttcc ccccaaaacc caaggacacc ctcatgatct cccggacccc tgaggtcaca      180
tgctgtggtgg tggacgtgag ccacgaagac cctgaggtca agttcaactg gtacgtggac      240
ggcgtggagg tgcataatgc caagacaaag ccgcgggagg agcagtacaa cagcacgtac      300
cgtgtgggtca gcgtcctcac cgctcctgcac caggactggc tgaatggcaa ggagtacaag      360
tgcaagggtct ccaacaaagc cctcccagcc cccatcgaga aaaccatctc caaagccaaa      420
gggcagcccc gagaaccaca ggtgtacacc ctgcccccat cccgcgatga gctgaccaag      480
aaccagggtca gcctgacctg cctgggtcaaa ggcttctatc ccagcgacat cgccgtggag      540
tgggagagca atgggcagcc ggagaacaac tacaagacca cgctcccggt gttggactcc      600
gacggctcct tcttctctta cagcaagctc accgtggaca agagcagggtg gcagcagggg      660
aacgtcttct catgctccgt gatgcatgag gctctgcaca accactacac gcagaagagc      720
ctctccctgt ctccgggtaa a                                         741
```

<210> SEQ ID NO 4
<211> LENGTH: 247
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Fc Portion

<400> SEQUENCE: 4

Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Leu Trp Val Pro
1 5 10 15

Gly Ser Thr Gly Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro
20 25 30

Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys
35 40 45

Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val
50 55 60

Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp
65 70 75 80

Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr
85 90 95

Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp
100 105 110

-continued

Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu
 115 120 125
 Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg
 130 135 140
 Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys
 145 150 155 160
 Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp
 165 170 175
 Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys
 180 185 190
 Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser
 195 200 205
 Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser
 210 215 220
 Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser
 225 230 235 240
 Leu Ser Leu Ser Pro Gly Lys
 245

<210> SEQ ID NO 5
 <211> LENGTH: 7734
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Full Length FVIIIIFc

<400> SEQUENCE: 5

atgcaaatag agctctccac ctgcttcttt ctgtgccttt tgcgattctg ctttagtgcc	60
accagaagat actacctggg tgcagtggaa ctgtcatggg actatatgca aagtgatctc	120
ggtgagctgc ctgtggacgc aagatttcct cctagagtgc caaaatcttt tccattcaac	180
acctcagtcg tgtacaaaaa gactctgttt gtagaattca cggatcacct tttcaacatc	240
gctaagccaa ggccaccctg gatgggtctg ctaggtccta ccatccaggc tgaggtttat	300
gatacagtgg tcattacact taagaacatg gcttcccatc ctgtcagtct tcatgctgtt	360
ggtgtatcct actggaaagc ttctgaggga gctgaatatg atgatcagac cagtcaaagg	420
gagaaagaag atgataaagt cttccctggg ggaagccata catatgtctg gcaggctctg	480
aaagagaatg gtccaatggc ctctgaccca ctgtgcctta cctactcata tctttctcat	540
gtggacctgg taaaagactt gaattcaggc ctcattggag ccctactagt atgtagagaa	600
gggagtctgg ccaaggaaaa gacacagacc ttgcacaaat ttatactact ttttgctgta	660
tttgatgaag ggaaaagtgt gcactcagaa acaaagaact ccttgatgca ggatagggat	720
gctgcatctg ctcgggcctg gcctaaaaatg cacacagtca atggttatgt aaacaggctc	780
ctgccaggtc tgattggatg ccacaggaaa tcagtcctatt ggcattgtgat tggaatgggc	840
accactcctg aagtgcactc aatattcttc gaaggtcaca catttcttgt gaggaaccat	900
cgccaggcgt ccttggaatg ctcgccaata actttcctta ctgctcaaac actcttgatg	960
gaccttggac agtttctact gttttgtcat atctcttccc accaacatga tggcatggaa	1020
gcttatgtca aagtagacag ctgtccagag gaaccccaac tacgaatgaa aaataatgaa	1080
gaagcggaag actatgatga tgatcttact gattctgaaa tggatgtggg caggtttgat	1140
gatgacaact ctccttcctt tatccaaatt cgctcagttg ccaagaagca tcctaaaact	1200
tgggtacatt acattgctgc tgaagaggag gactgggact atgctccctt agtcctcgcc	1260

-continued

cccgatgaca gaagttataa aagtcaatat ttgaacaatg gccctcagcg gattggtagg	1320
aagtacaaaa aagtcgatt tatggcatal acagatgaaa cctttaagac tcgtgaagct	1380
attcagcatg aatcaggaat cttgggacct ttactttatg gggaagttgg agacacactg	1440
ttgattatat ttaagaatca agcaagcaga ccatataaca tctaccctca cggaatcact	1500
gatgtccgto ctttgtattc aaggagatta ccaaaagggtg taaaacattt gaaggatttt	1560
ccaattctgc caggagaaat attcaaatat aaatggacag tgactgtaga agatgggcca	1620
actaaatcag atcctcggtg cctgacccgc tattactcta gtttcgttaa tatggagaga	1680
gatctagctt caggactcat tggccctctc ctcatctgct acaagaatc tgtagatcaa	1740
agaggaaacc agataatgc agacaagagg aatgtcatcc tgttttctgt atttgatgag	1800
aaaccgaagct ggtacctcac agagaatata caacgcttcc tcccaatcc agctggagtg	1860
cagcttgagg atccagagtt ccaagcctcc aacatcatgc acagcatcaa tggctatgtt	1920
tttgatagtt tgcagtgtc agtttgtttg catgaggtgg catactggta cattctaagc	1980
attggagcac agactgactt cctttctgtc ttcttctctg gatatacctt caaacacaaa	2040
atggtctatg aagacacact caccctattc ccattctcag gaaaaactgt cttcatgtcg	2100
atggaaaacc caggtctatg gattctgggg tgcacaaact cagactttcg gaacagaggc	2160
atgaccgctt tactgaaggt ttctagtgtg gacaagaaca ctggtgatta ttacgaggac	2220
agttatgaag atatttcagc atacttgctg agtaaaaaa atgccattga accaagaagc	2280
ttctcccaga attcaagaca ccttagcact agggaaaagc aatttaatgc caccacaatt	2340
ccagaaaatg acatagagaa gactgacctt tggtttgac acagaacacc tatgcctaaa	2400
atacaaatg tctcctctag tgatttggtg atgctctgc gacagagtcc tactccacat	2460
gggtatcct tatctgatct ccaagaagcc aaatatgaga ctttttctga tgatccatca	2520
cctggagcaa tagacagtaa taacagcctg tctgaaatga cacacttcag gccacagctc	2580
catcacagtg gggacatggt atttaccctt gagtcaggcc tccaattaag attaaatgag	2640
aaactgggga caactgcagc aacagagttg aagaaacttg atttcaaagt ttctagtaca	2700
tcaaataatc tgatttcaac aattccatca gacaatttg cagcaggtac tgataatata	2760
agttccttag gacccccag tatgccagtt cattatgata gtcaattaga taccactcta	2820
tttgcaaaa agtcatctcc ccttactgag tctggtggac ctctgagctt gagtgaagaa	2880
aataatgatt caaagtgtt agaatcaggt ttaatgaata gccaaagaa ttcatgggga	2940
aaaaatgtat cgtcaacaga gagtggtagg ttatttaaag ggaaaagagc tcatggacct	3000
gctttgttga ctaaagataa tgccttattc aaagttagca tctctttgtt aaagacaaac	3060
aaaacttcca ataattcagc aactaataga aagactcaca ttgatggccc atcattatta	3120
attgagaata gtccatcagt ctggcaaaat atattagaaa gtgacactga gtttaaaaaa	3180
gtgacacctt tgattcatga cagaatgctt atggacaaaa atgctacagc ttgaggcta	3240
aatcatatgt caaataaaac tacttcatca aaaaacatgg aaatggtcca acagaaaaaa	3300
gagggcccca ttccaccaga tgcacaaaat ccagatatgt cgttctttaa gatgctattc	3360
ttgccagaat cagcaagggt gatacaaaag actcatggaa agaactctct gaactctggg	3420
caaggcccca gtccaaagca attagtatcc ttaggaccag aaaaatctgt ggaaggtcag	3480
aatttcttgt ctgagaaaaa caaagtggta gtaggaaagg gtgaatttac aaaggacgta	3540
ggactcaaag agatggtttt tccaagcagc agaaacctat ttcttactaa cttggataat	3600
ttacatgaaa ataatacaca caatcaagaa aaaaaattc aggaagaaat agaaaagaag	3660

-continued

gaaacattaa tccaagagaa tgtagttttg cctcagatac atacagtgcac tggcactaag	3720
aatttcacga agaacctttt cttactgagc actaggcaaa atgtagaagg ttcacatgac	3780
ggggcatatg ctccagtagt tcaagatttt aggtcattaa atgattcaac aaatagaaca	3840
aagaaacaca cagctcattt ctcaaaaaaa ggggaggaag aaaacttgga aggcctggga	3900
aatcaaacca agcaaattgt agagaaatat gcatgcacca caaggatata tcctaataca	3960
agccagcaga atttttgtcac gcaacgtagt aagagagcct tgaaacaatt cagactccca	4020
ctagaagaaa cagaacttga aaaaaggata attgtggatg acacctcaac ccagtgggcc	4080
aaaaacatga aacatttgac cccgagcacc ctcacacaga tagactacaa tgagaaggag	4140
aaaggggcca ttactcagtc tcacctatca gattgcctta cgaggagtca tagcatccct	4200
caagcaaata gatctccatt acccattgca aaggtatcat catttccatc tattagacct	4260
atatacttga ccagggtcct attccaagac aactcttctc atcttccagc agcatcttat	4320
agaaagaaaag attctggggt ccaagaaagc agtcatttct tacaaggagc caaaaaaat	4380
aaccttttct tagccattct aaccttgag atgactgggt atcaaagaga ggttggtccc	4440
ctggggacaa gtgccacaaa ttcagtcaca tacaagaaag ttgagaacac tgttctcccg	4500
aaaccagact tgcccaaac atctggcaaa gttgaattgc ttccaaaagt tcacatttat	4560
cagaaggacc tatccctac ggaaactagc aatgggtctc ctggccatct ggatctctgt	4620
gaagggagcc ttcttcaggg aacagaggga gcgattaagt ggaatgaagc aaacagacct	4680
ggaaaagtcc cctttctgag agtagcaaca gaaagctctg caaagactcc ctccaagcta	4740
ttggatcttc ttgcttggga taaccactat ggtactcaga taccaaaaga agagtggaaa	4800
tccaagaga agtcaccaga aaaaacagct tttaagaaaa aggataccat tttgtccctg	4860
aacgcttggt aaagcaatca tgcaatagca gcaataaatg agggacaaaa taagcccgaa	4920
atagaagtca cctgggcaaa gcaaggtagg actgaaaggc tgtgctctca aaaccaccca	4980
gtcttgaaac gccatcaacg ggaaataact cgtactactc ttcagtcaga tcaagaggaa	5040
attgactatg atgataccat atcagttgaa atgaagaagg aagattttga cttttatgat	5100
gaggatgaaa atcagagccc ccgcagcttt caaaagaaaa cagcactata ttttattgct	5160
gcagtggaga ggctctggga ttatgggatg agtagctccc cacatgttct aagaaacagg	5220
gctcagagtg gcagtgtccc tcagttcaag aaagtgtttt tccaggaatt tactgatggc	5280
tcctttactc agcccttata ccgtggagaa ctaaatgaac atttgggact cctggggcca	5340
tatataagag cagaagtga agataatatc atggtaactt tcagaaatca ggctctcgt	5400
ccctattcct tctattctag ccttatttct tatgaggaag atcagaggca aggagcagaa	5460
cctagaaaaa actttgtcaa gcctaataaa accaaaactt acttttgga agtgcaacat	5520
catatggcac ccactaaaga tgagtttgac tgcaaacgct gggcttattt ctctgatgtt	5580
gacctggaaa aagatgtgca ctcaggcctg attggacccc ttctgggtctg ccacactaac	5640
acactgaacc ctgctcatgg gagacaagtg acagtacagg aatttgctct gtttttcacc	5700
atctttgatg agaccaaaag ctggtacttc actgaaaata tggaaagaaa ctgcagggct	5760
ccctgcaata tccagatgga agatcccact tttaagaga attatcgctt ccattgcaatc	5820
aatggctaca taatggatag actacctggc ttagtaatgg ctcaggatca aaggattcga	5880
tggtatctgc tcagcatggg cagcaatgaa aacatccatt ctattcattt cagtggacat	5940
gtgttcactg tacgaaaaaa agaggagtat aaatggcac tgtacaatct ctatccaggt	6000

-continued

```

gtttttgaga cagtggaaat gttaccatcc aaagctggaa tttggcgggt ggaatgcctt 6060
attggcgagc atctacatgc tgggatgagc acactttttc tgggtgtacag caataagtgt 6120
cagactcccc tgggaatggc ttctggacac attagagatt ttcagattac agcttcagga 6180
caatatggac agtgggcccc aaagctggcc agacttcatt attccggatc aatcaatgcc 6240
tggagcacca aggagccctt ttcttgatc aaggtggatc tgttggcacc aatgattatt 6300
cacggcatca agaccagggt tggcgtcag aagttctcca gcctctacat ctctcagttt 6360
atcatcatgt atagtcttga tgggaagaag tggcagactt atcgaggaaa ttccactgga 6420
accttaatgg tcttcttttg caatgtggat tcatctggga taaaacacaa tatttttaac 6480
cttcaatta ttgctcgata catccgtttg cacccaactc attatagcat tcgcagcact 6540
cttcgcattg agttgatggg ctgtgattta aatagttgca gcatgccatt gggaatggag 6600
agtaaagcaa tatcagatgc acagattact gcttcatcct accttaccaa tatgtttgcc 6660
acctgggtctc cttcaaaagc tcgacttcac ctccaaggga ggagtaatgc ctggagacct 6720
cagggtgaata atccaaaaga gtggctgcaa gtggacttcc agaagacaat gaaagtcaca 6780
ggagtaacta ctcagggagt aaaatctctg cttaccagca tgtatgtgaa ggagttcctc 6840
atctccagca gtcaagatgg ccacagtggt actctctttt ttcagaatgg caaagtaaa 6900
gtttttcagg gaaatcaaga ctcttcaca cctgtggtga actctctaga cccaccgtta 6960
ctgactcgct accttcgaat tcacccccag agttgggtgc accagattgc cctgaggatg 7020
gaggttcttg gctgcgaggc acaggacctc tacgacaaaa ctcacacatg cccaccgtgc 7080
ccagctccag aactcctggg cggaccgtca gtcttctctt tcccccaaa acccaaggac 7140
acctcatga tctcccgac cctgaggtc acatgcgtgg tgggtggacgt gagccacgaa 7200
gacctgagg tcaagttcaa ctggtagctg gacggcgtgg aggtgcataa tgccaagaca 7260
aagccgctgg aggagcagta caacagcacg taccgtgtgg tcagcgtcct caccgtcctg 7320
caccaggact ggctgaatgg caaggagtac aagtgcgaagg tctccaacaa agccctccca 7380
gcccccatcg agaaaacat ctccaagcc aaagggcagc cccgagaacc acaggtgtac 7440
acctgcccc catcccgga tgagctgacc aagaaccagg tcagcctgac ctgctgggtc 7500
aaaggcttct atcccagca catcgccgtg gagtgggaga gcaatgggca gccggagaac 7560
aactacaaga ccacgcctcc cgtgttgac tccgacggct ccttcttctc ctacagcaag 7620
ctcaccgtgg acaagagcag gtggcagcag gggaaagtct tctcatgctc cgtgatgcat 7680
gaggctctgc acaaccacta cagcagaag agcctctccc tgtctccggg taaa 7734

```

```

<210> SEQ ID NO 6
<211> LENGTH: 2578
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Full Length FVIIIIFc

```

```

<400> SEQUENCE: 6

```

```

Met Gln Ile Glu Leu Ser Thr Cys Phe Phe Leu Cys Leu Leu Arg Phe
1           5           10           15

```

```

Cys Phe Ser Ala Thr Arg Arg Tyr Tyr Leu Gly Ala Val Glu Leu Ser
20           25           30

```

```

Trp Asp Tyr Met Gln Ser Asp Leu Gly Glu Leu Pro Val Asp Ala Arg
35           40           45

```

```

Phe Pro Pro Arg Val Pro Lys Ser Phe Pro Phe Asn Thr Ser Val Val
50           55           60

```

-continued

Tyr	Lys	Lys	Thr	Leu	Phe	Val	Glu	Phe	Thr	Asp	His	Leu	Phe	Asn	Ile	65	70	75	80
Ala	Lys	Pro	Arg	Pro	Pro	Trp	Met	Gly	Leu	Leu	Gly	Pro	Thr	Ile	Gln	85	90	95	
Ala	Glu	Val	Tyr	Asp	Thr	Val	Val	Ile	Thr	Leu	Lys	Asn	Met	Ala	Ser	100	105	110	
His	Pro	Val	Ser	Leu	His	Ala	Val	Gly	Val	Ser	Tyr	Trp	Lys	Ala	Ser	115	120	125	
Glu	Gly	Ala	Glu	Tyr	Asp	Asp	Gln	Thr	Ser	Gln	Arg	Glu	Lys	Glu	Asp	130	135	140	
Asp	Lys	Val	Phe	Pro	Gly	Gly	Ser	His	Thr	Tyr	Val	Trp	Gln	Val	Leu	145	150	155	160
Lys	Glu	Asn	Gly	Pro	Met	Ala	Ser	Asp	Pro	Leu	Cys	Leu	Thr	Tyr	Ser	165	170	175	
Tyr	Leu	Ser	His	Val	Asp	Leu	Val	Lys	Asp	Leu	Asn	Ser	Gly	Leu	Ile	180	185	190	
Gly	Ala	Leu	Leu	Val	Cys	Arg	Glu	Gly	Ser	Leu	Ala	Lys	Glu	Lys	Thr	195	200	205	
Gln	Thr	Leu	His	Lys	Phe	Ile	Leu	Leu	Phe	Ala	Val	Phe	Asp	Glu	Gly	210	215	220	
Lys	Ser	Trp	His	Ser	Glu	Thr	Lys	Asn	Ser	Leu	Met	Gln	Asp	Arg	Asp	225	230	235	240
Ala	Ala	Ser	Ala	Arg	Ala	Trp	Pro	Lys	Met	His	Thr	Val	Asn	Gly	Tyr	245	250	255	
Val	Asn	Arg	Ser	Leu	Pro	Gly	Leu	Ile	Gly	Cys	His	Arg	Lys	Ser	Val	260	265	270	
Tyr	Trp	His	Val	Ile	Gly	Met	Gly	Thr	Thr	Pro	Glu	Val	His	Ser	Ile	275	280	285	
Phe	Leu	Glu	Gly	His	Thr	Phe	Leu	Val	Arg	Asn	His	Arg	Gln	Ala	Ser	290	295	300	
Leu	Glu	Ile	Ser	Pro	Ile	Thr	Phe	Leu	Thr	Ala	Gln	Thr	Leu	Leu	Met	305	310	315	320
Asp	Leu	Gly	Gln	Phe	Leu	Leu	Phe	Cys	His	Ile	Ser	Ser	His	Gln	His	325	330	335	
Asp	Gly	Met	Glu	Ala	Tyr	Val	Lys	Val	Asp	Ser	Cys	Pro	Glu	Glu	Pro	340	345	350	
Gln	Leu	Arg	Met	Lys	Asn	Asn	Glu	Glu	Ala	Glu	Asp	Tyr	Asp	Asp	Asp	355	360	365	
Leu	Thr	Asp	Ser	Glu	Met	Asp	Val	Val	Arg	Phe	Asp	Asp	Asp	Asn	Ser	370	375	380	
Pro	Ser	Phe	Ile	Gln	Ile	Arg	Ser	Val	Ala	Lys	Lys	His	Pro	Lys	Thr	385	390	395	400
Trp	Val	His	Tyr	Ile	Ala	Ala	Glu	Glu	Glu	Asp	Trp	Asp	Tyr	Ala	Pro	405	410	415	
Leu	Val	Leu	Ala	Pro	Asp	Asp	Arg	Ser	Tyr	Lys	Ser	Gln	Tyr	Leu	Asn	420	425	430	
Asn	Gly	Pro	Gln	Arg	Ile	Gly	Arg	Lys	Tyr	Lys	Lys	Val	Arg	Phe	Met	435	440	445	
Ala	Tyr	Thr	Asp	Glu	Thr	Phe	Lys	Thr	Arg	Glu	Ala	Ile	Gln	His	Glu	450	455	460	
Ser	Gly	Ile	Leu	Gly	Pro	Leu	Leu	Tyr	Gly	Glu	Val	Gly	Asp	Thr	Leu	465	470	475	480

-continued

Leu	Ile	Ile	Phe	Lys	Asn	Gln	Ala	Ser	Arg	Pro	Tyr	Asn	Ile	Tyr	Pro	485	490	495
His	Gly	Ile	Thr	Asp	Val	Arg	Pro	Leu	Tyr	Ser	Arg	Arg	Leu	Pro	Lys	500	505	510
Gly	Val	Lys	His	Leu	Lys	Asp	Phe	Pro	Ile	Leu	Pro	Gly	Glu	Ile	Phe	515	520	525
Lys	Tyr	Lys	Trp	Thr	Val	Thr	Val	Glu	Asp	Gly	Pro	Thr	Lys	Ser	Asp	530	535	540
Pro	Arg	Cys	Leu	Thr	Arg	Tyr	Tyr	Ser	Ser	Phe	Val	Asn	Met	Glu	Arg	545	550	555
Asp	Leu	Ala	Ser	Gly	Leu	Ile	Gly	Pro	Leu	Leu	Ile	Cys	Tyr	Lys	Glu	565	570	575
Ser	Val	Asp	Gln	Arg	Gly	Asn	Gln	Ile	Met	Ser	Asp	Lys	Arg	Asn	Val	580	585	590
Ile	Leu	Phe	Ser	Val	Phe	Asp	Glu	Asn	Arg	Ser	Trp	Tyr	Leu	Thr	Glu	595	600	605
Asn	Ile	Gln	Arg	Phe	Leu	Pro	Asn	Pro	Ala	Gly	Val	Gln	Leu	Glu	Asp	610	615	620
Pro	Glu	Phe	Gln	Ala	Ser	Asn	Ile	Met	His	Ser	Ile	Asn	Gly	Tyr	Val	625	630	635
Phe	Asp	Ser	Leu	Gln	Leu	Ser	Val	Cys	Leu	His	Glu	Val	Ala	Tyr	Trp	645	650	655
Tyr	Ile	Leu	Ser	Ile	Gly	Ala	Gln	Thr	Asp	Phe	Leu	Ser	Val	Phe	Phe	660	665	670
Ser	Gly	Tyr	Thr	Phe	Lys	His	Lys	Met	Val	Tyr	Glu	Asp	Thr	Leu	Thr	675	680	685
Leu	Phe	Pro	Phe	Ser	Gly	Glu	Thr	Val	Phe	Met	Ser	Met	Glu	Asn	Pro	690	695	700
Gly	Leu	Trp	Ile	Leu	Gly	Cys	His	Asn	Ser	Asp	Phe	Arg	Asn	Arg	Gly	705	710	715
Met	Thr	Ala	Leu	Leu	Lys	Val	Ser	Ser	Cys	Asp	Lys	Asn	Thr	Gly	Asp	725	730	735
Tyr	Tyr	Glu	Asp	Ser	Tyr	Glu	Asp	Ile	Ser	Ala	Tyr	Leu	Leu	Ser	Lys	740	745	750
Asn	Asn	Ala	Ile	Glu	Pro	Arg	Ser	Phe	Ser	Gln	Asn	Ser	Arg	His	Pro	755	760	765
Ser	Thr	Arg	Gln	Lys	Gln	Phe	Asn	Ala	Thr	Thr	Ile	Pro	Glu	Asn	Asp	770	775	780
Ile	Glu	Lys	Thr	Asp	Pro	Trp	Phe	Ala	His	Arg	Thr	Pro	Met	Pro	Lys	785	790	795
Ile	Gln	Asn	Val	Ser	Ser	Ser	Asp	Leu	Leu	Met	Leu	Leu	Arg	Gln	Ser	805	810	815
Pro	Thr	Pro	His	Gly	Leu	Ser	Leu	Ser	Asp	Leu	Gln	Glu	Ala	Lys	Tyr	820	825	830
Glu	Thr	Phe	Ser	Asp	Asp	Pro	Ser	Pro	Gly	Ala	Ile	Asp	Ser	Asn	Asn	835	840	845
Ser	Leu	Ser	Glu	Met	Thr	His	Phe	Arg	Pro	Gln	Leu	His	His	Ser	Gly	850	855	860
Asp	Met	Val	Phe	Thr	Pro	Glu	Ser	Gly	Leu	Gln	Leu	Arg	Leu	Asn	Glu	865	870	875
Lys	Leu	Gly	Thr	Thr	Ala	Ala	Thr	Glu	Leu	Lys	Lys	Leu	Asp	Phe	Lys	885	890	895
Val	Ser	Ser	Thr	Ser	Asn	Asn	Leu	Ile	Ser	Thr	Ile	Pro	Ser	Asp	Asn			

-continued

900	905	910
Leu Ala Ala Gly Thr Asp Asn Thr Ser Ser Leu Gly Pro Pro Ser Met 915 920 925		
Pro Val His Tyr Asp Ser Gln Leu Asp Thr Thr Leu Phe Gly Lys Lys 930 935 940		
Ser Ser Pro Leu Thr Glu Ser Gly Gly Pro Leu Ser Leu Ser Glu Glu 945 950 955 960		
Asn Asn Asp Ser Lys Leu Leu Glu Ser Gly Leu Met Asn Ser Gln Glu 965 970 975		
Ser Ser Trp Gly Lys Asn Val Ser Ser Thr Glu Ser Gly Arg Leu Phe 980 985 990		
Lys Gly Lys Arg Ala His Gly Pro Ala Leu Leu Thr Lys Asp Asn Ala 995 1000 1005		
Leu Phe Lys Val Ser Ile Ser Leu Leu Lys Thr Asn Lys Thr Ser 1010 1015 1020		
Asn Asn Ser Ala Thr Asn Arg Lys Thr His Ile Asp Gly Pro Ser 1025 1030 1035		
Leu Leu Ile Glu Asn Ser Pro Ser Val Trp Gln Asn Ile Leu Glu 1040 1045 1050		
Ser Asp Thr Glu Phe Lys Lys Val Thr Pro Leu Ile His Asp Arg 1055 1060 1065		
Met Leu Met Asp Lys Asn Ala Thr Ala Leu Arg Leu Asn His Met 1070 1075 1080		
Ser Asn Lys Thr Thr Ser Ser Lys Asn Met Glu Met Val Gln Gln 1085 1090 1095		
Lys Lys Glu Gly Pro Ile Pro Pro Asp Ala Gln Asn Pro Asp Met 1100 1105 1110		
Ser Phe Phe Lys Met Leu Phe Leu Pro Glu Ser Ala Arg Trp Ile 1115 1120 1125		
Gln Arg Thr His Gly Lys Asn Ser Leu Asn Ser Gly Gln Gly Pro 1130 1135 1140		
Ser Pro Lys Gln Leu Val Ser Leu Gly Pro Glu Lys Ser Val Glu 1145 1150 1155		
Gly Gln Asn Phe Leu Ser Glu Lys Asn Lys Val Val Val Gly Lys 1160 1165 1170		
Gly Glu Phe Thr Lys Asp Val Gly Leu Lys Glu Met Val Phe Pro 1175 1180 1185		
Ser Ser Arg Asn Leu Phe Leu Thr Asn Leu Asp Asn Leu His Glu 1190 1195 1200		
Asn Asn Thr His Asn Gln Glu Lys Lys Ile Gln Glu Glu Ile Glu 1205 1210 1215		
Lys Lys Glu Thr Leu Ile Gln Glu Asn Val Val Leu Pro Gln Ile 1220 1225 1230		
His Thr Val Thr Gly Thr Lys Asn Phe Met Lys Asn Leu Phe Leu 1235 1240 1245		
Leu Ser Thr Arg Gln Asn Val Glu Gly Ser Tyr Asp Gly Ala Tyr 1250 1255 1260		
Ala Pro Val Leu Gln Asp Phe Arg Ser Leu Asn Asp Ser Thr Asn 1265 1270 1275		
Arg Thr Lys Lys His Thr Ala His Phe Ser Lys Lys Gly Glu Glu 1280 1285 1290		
Glu Asn Leu Glu Gly Leu Gly Asn Gln Thr Lys Gln Ile Val Glu 1295 1300 1305		

-continued

Lys Tyr	Ala Cys Thr Thr	Arg	Ile Ser Pro Asn Thr	Ser Gln Gln
1310		1315		1320
Asn Phe	Val Thr Gln Arg	Ser	Lys Arg Ala Leu Lys	Gln Phe Arg
1325		1330		1335
Leu Pro	Leu Glu Glu Thr	Glu	Leu Glu Lys Arg Ile	Ile Val Asp
1340		1345		1350
Asp Thr	Ser Thr Gln Trp	Ser	Lys Asn Met Lys His	Leu Thr Pro
1355		1360		1365
Ser Thr	Leu Thr Gln Ile	Asp	Tyr Asn Glu Lys Glu	Lys Gly Ala
1370		1375		1380
Ile Thr	Gln Ser Pro Leu	Ser	Asp Cys Leu Thr Arg	Ser His Ser
1385		1390		1395
Ile Pro	Gln Ala Asn Arg	Ser	Pro Leu Pro Ile Ala	Lys Val Ser
1400		1405		1410
Ser Phe	Pro Ser Ile Arg	Pro	Ile Tyr Leu Thr Arg	Val Leu Phe
1415		1420		1425
Gln Asp	Asn Ser Ser His	Leu	Pro Ala Ala Ser Tyr	Arg Lys Lys
1430		1435		1440
Asp Ser	Gly Val Gln Glu	Ser	Ser His Phe Leu Gln	Gly Ala Lys
1445		1450		1455
Lys Asn	Asn Leu Ser Leu	Ala	Ile Leu Thr Leu Glu	Met Thr Gly
1460		1465		1470
Asp Gln	Arg Glu Val Gly	Ser	Leu Gly Thr Ser Ala	Thr Asn Ser
1475		1480		1485
Val Thr	Tyr Lys Lys Val	Glu	Asn Thr Val Leu Pro	Lys Pro Asp
1490		1495		1500
Leu Pro	Lys Thr Ser Gly	Lys	Val Glu Leu Leu Pro	Lys Val His
1505		1510		1515
Ile Tyr	Gln Lys Asp Leu	Phe	Pro Thr Glu Thr Ser	Asn Gly Ser
1520		1525		1530
Pro Gly	His Leu Asp Leu	Val	Glu Gly Ser Leu Leu	Gln Gly Thr
1535		1540		1545
Glu Gly	Ala Ile Lys Trp	Asn	Glu Ala Asn Arg Pro	Gly Lys Val
1550		1555		1560
Pro Phe	Leu Arg Val Ala	Thr	Glu Ser Ser Ala Lys	Thr Pro Ser
1565		1570		1575
Lys Leu	Leu Asp Pro Leu	Ala	Trp Asp Asn His Tyr	Gly Thr Gln
1580		1585		1590
Ile Pro	Lys Glu Glu Trp	Lys	Ser Gln Glu Lys Ser	Pro Glu Lys
1595		1600		1605
Thr Ala	Phe Lys Lys Lys	Asp	Thr Ile Leu Ser Leu	Asn Ala Cys
1610		1615		1620
Glu Ser	Asn His Ala Ile	Ala	Ala Ile Asn Glu Gly	Gln Asn Lys
1625		1630		1635
Pro Glu	Ile Glu Val Thr	Trp	Ala Lys Gln Gly Arg	Thr Glu Arg
1640		1645		1650
Leu Cys	Ser Gln Asn Pro	Pro	Val Leu Lys Arg His	Gln Arg Glu
1655		1660		1665
Ile Thr	Arg Thr Thr Leu	Gln	Ser Asp Gln Glu Glu	Ile Asp Tyr
1670		1675		1680
Asp Asp	Thr Ile Ser Val	Glu	Met Lys Lys Glu Asp	Phe Asp Ile
1685		1690		1695

-continued

Tyr	Asp	Glu	Asp	Glu	Asn	Gln	Ser	Pro	Arg	Ser	Phe	Gln	Lys	Lys
1700						1705					1710			
Thr	Arg	His	Tyr	Phe	Ile	Ala	Ala	Val	Glu	Arg	Leu	Trp	Asp	Tyr
1715						1720					1725			
Gly	Met	Ser	Ser	Ser	Pro	His	Val	Leu	Arg	Asn	Arg	Ala	Gln	Ser
1730						1735					1740			
Gly	Ser	Val	Pro	Gln	Phe	Lys	Lys	Val	Val	Phe	Gln	Glu	Phe	Thr
1745						1750					1755			
Asp	Gly	Ser	Phe	Thr	Gln	Pro	Leu	Tyr	Arg	Gly	Glu	Leu	Asn	Glu
1760						1765					1770			
His	Leu	Gly	Leu	Leu	Gly	Pro	Tyr	Ile	Arg	Ala	Glu	Val	Glu	Asp
1775						1780					1785			
Asn	Ile	Met	Val	Thr	Phe	Arg	Asn	Gln	Ala	Ser	Arg	Pro	Tyr	Ser
1790						1795					1800			
Phe	Tyr	Ser	Ser	Leu	Ile	Ser	Tyr	Glu	Glu	Asp	Gln	Arg	Gln	Gly
1805						1810					1815			
Ala	Glu	Pro	Arg	Lys	Asn	Phe	Val	Lys	Pro	Asn	Glu	Thr	Lys	Thr
1820						1825					1830			
Tyr	Phe	Trp	Lys	Val	Gln	His	His	Met	Ala	Pro	Thr	Lys	Asp	Glu
1835						1840					1845			
Phe	Asp	Cys	Lys	Ala	Trp	Ala	Tyr	Phe	Ser	Asp	Val	Asp	Leu	Glu
1850						1855					1860			
Lys	Asp	Val	His	Ser	Gly	Leu	Ile	Gly	Pro	Leu	Leu	Val	Cys	His
1865						1870					1875			
Thr	Asn	Thr	Leu	Asn	Pro	Ala	His	Gly	Arg	Gln	Val	Thr	Val	Gln
1880						1885					1890			
Glu	Phe	Ala	Leu	Phe	Phe	Thr	Ile	Phe	Asp	Glu	Thr	Lys	Ser	Trp
1895						1900					1905			
Tyr	Phe	Thr	Glu	Asn	Met	Glu	Arg	Asn	Cys	Arg	Ala	Pro	Cys	Asn
1910						1915					1920			
Ile	Gln	Met	Glu	Asp	Pro	Thr	Phe	Lys	Glu	Asn	Tyr	Arg	Phe	His
1925						1930					1935			
Ala	Ile	Asn	Gly	Tyr	Ile	Met	Asp	Thr	Leu	Pro	Gly	Leu	Val	Met
1940						1945					1950			
Ala	Gln	Asp	Gln	Arg	Ile	Arg	Trp	Tyr	Leu	Leu	Ser	Met	Gly	Ser
1955						1960					1965			
Asn	Glu	Asn	Ile	His	Ser	Ile	His	Phe	Ser	Gly	His	Val	Phe	Thr
1970						1975					1980			
Val	Arg	Lys	Lys	Glu	Glu	Tyr	Lys	Met	Ala	Leu	Tyr	Asn	Leu	Tyr
1985						1990					1995			
Pro	Gly	Val	Phe	Glu	Thr	Val	Glu	Met	Leu	Pro	Ser	Lys	Ala	Gly
2000						2005					2010			
Ile	Trp	Arg	Val	Glu	Cys	Leu	Ile	Gly	Glu	His	Leu	His	Ala	Gly
2015						2020					2025			
Met	Ser	Thr	Leu	Phe	Leu	Val	Tyr	Ser	Asn	Lys	Cys	Gln	Thr	Pro
2030						2035					2040			
Leu	Gly	Met	Ala	Ser	Gly	His	Ile	Arg	Asp	Phe	Gln	Ile	Thr	Ala
2045						2050					2055			
Ser	Gly	Gln	Tyr	Gly	Gln	Trp	Ala	Pro	Lys	Leu	Ala	Arg	Leu	His
2060						2065					2070			
Tyr	Ser	Gly	Ser	Ile	Asn	Ala	Trp	Ser	Thr	Lys	Glu	Pro	Phe	Ser
2075						2080					2085			
Trp	Ile	Lys	Val	Asp	Leu	Leu	Ala	Pro	Met	Ile	Ile	His	Gly	Ile

-continued

2090	2095	2100
Lys Thr Gln Gly Ala Arg	Gln Lys Phe Ser Ser	Leu Tyr Ile Ser
2105	2110	2115
Gln Phe Ile Ile Met Tyr	Ser Leu Asp Gly Lys	Lys Trp Gln Thr
2120	2125	2130
Tyr Arg Gly Asn Ser Thr	Gly Thr Leu Met Val	Phe Phe Gly Asn
2135	2140	2145
Val Asp Ser Ser Gly Ile	Lys His Asn Ile Phe	Asn Pro Pro Ile
2150	2155	2160
Ile Ala Arg Tyr Ile Arg	Leu His Pro Thr His	Tyr Ser Ile Arg
2165	2170	2175
Ser Thr Leu Arg Met Glu	Leu Met Gly Cys Asp	Leu Asn Ser Cys
2180	2185	2190
Ser Met Pro Leu Gly Met	Glu Ser Lys Ala Ile	Ser Asp Ala Gln
2195	2200	2205
Ile Thr Ala Ser Ser Tyr	Phe Thr Asn Met Phe	Ala Thr Trp Ser
2210	2215	2220
Pro Ser Lys Ala Arg Leu	His Leu Gln Gly Arg	Ser Asn Ala Trp
2225	2230	2235
Arg Pro Gln Val Asn Asn	Pro Lys Glu Trp Leu	Gln Val Asp Phe
2240	2245	2250
Gln Lys Thr Met Lys Val	Thr Gly Val Thr Thr	Gln Gly Val Lys
2255	2260	2265
Ser Leu Leu Thr Ser Met	Tyr Val Lys Glu Phe	Leu Ile Ser Ser
2270	2275	2280
Ser Gln Asp Gly His Gln	Trp Thr Leu Phe Phe	Gln Asn Gly Lys
2285	2290	2295
Val Lys Val Phe Gln Gly	Asn Gln Asp Ser Phe	Thr Pro Val Val
2300	2305	2310
Asn Ser Leu Asp Pro Pro	Leu Leu Thr Arg Tyr	Leu Arg Ile His
2315	2320	2325
Pro Gln Ser Trp Val His	Gln Ile Ala Leu Arg	Met Glu Val Leu
2330	2335	2340
Gly Cys Glu Ala Gln Asp	Leu Tyr Asp Lys Thr	His Thr Cys Pro
2345	2350	2355
Pro Cys Pro Ala Pro Glu	Leu Leu Gly Gly Pro	Ser Val Phe Leu
2360	2365	2370
Phe Pro Pro Lys Pro Lys	Asp Thr Leu Met Ile	Ser Arg Thr Pro
2375	2380	2385
Glu Val Thr Cys Val Val	Val Asp Val Ser His	Glu Asp Pro Glu
2390	2395	2400
Val Lys Phe Asn Trp Tyr	Val Asp Gly Val Glu	Val His Asn Ala
2405	2410	2415
Lys Thr Lys Pro Arg Glu	Glu Gln Tyr Asn Ser	Thr Tyr Arg Val
2420	2425	2430
Val Ser Val Leu Thr Val	Leu His Gln Asp Trp	Leu Asn Gly Lys
2435	2440	2445
Glu Tyr Lys Cys Lys Val	Ser Asn Lys Ala Leu	Pro Ala Pro Ile
2450	2455	2460
Glu Lys Thr Ile Ser Lys	Ala Lys Gly Gln Pro	Arg Glu Pro Gln
2465	2470	2475
Val Tyr Thr Leu Pro Pro	Ser Arg Asp Glu Leu	Thr Lys Asn Gln
2480	2485	2490

-continued

Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile	
2495	2500 2505
Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys	
2510	2515 2520
Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr	
2525	2530 2535
Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val	
2540	2545 2550
Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr	
2555	2560 2565
Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys	
2570	2575

<210> SEQ ID NO 7
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Albumin-binding Peptide Core

<400> SEQUENCE: 7

Asp Ile Cys Leu Pro Arg Trp Gly Cys Leu Trp
1 5 10

<210> SEQ ID NO 8
 <211> LENGTH: 32
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: CTP peptide

<400> SEQUENCE: 8

Asp Pro Arg Phe Gln Asp Ser Ser Ser Ser Lys Ala Pro Pro Pro Ser			
1 5 10 15			
Leu Pro Ser Pro Ser Arg Leu Pro Gly Pro Ser Asp Thr Pro Ile Leu			
20 25 30			

<210> SEQ ID NO 9
 <211> LENGTH: 28
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: CTP peptide

<400> SEQUENCE: 9

Ser Ser Ser Ser Lys Ala Pro Pro Pro Ser Leu Pro Ser Pro Ser Arg			
1 5 10 15			
Leu Pro Gly Pro Ser Asp Thr Pro Ile Leu Pro Gln			
20 25			

<210> SEQ ID NO 10
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: PAS peptide

<400> SEQUENCE: 10

Ala Ser Pro Ala Ala Pro Ala Pro Ala Ser Pro Ala Ala Pro Ala Pro			
1 5 10 15			
Ser Ala Pro Ala			
20			

-continued

<210> SEQ ID NO 11
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PAS peptide

<400> SEQUENCE: 11

Ala Ala Pro Ala Ser Pro Ala Pro Ala Ala Pro Ser Ala Pro Ala Pro
1 5 10 15

Ala Ala Pro Ser
20

<210> SEQ ID NO 12
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PAS peptide

<400> SEQUENCE: 12

Ala Pro Ser Ser Pro Ser Pro Ser Ala Pro Ser Ser Pro Ser Pro Ala
1 5 10 15

Ser Pro Ser Ser
20

<210> SEQ ID NO 13
<211> LENGTH: 19
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PAS peptide

<400> SEQUENCE: 13

Ala Pro Ser Ser Pro Ser Pro Ser Ala Pro Ser Ser Pro Ser Pro Ala
1 5 10 15

Ser Pro Ser

<210> SEQ ID NO 14
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PAS peptide

<400> SEQUENCE: 14

Ser Ser Pro Ser Ala Pro Ser Pro Ser Ser Pro Ala Ser Pro Ser Pro
1 5 10 15

Ser Ser Pro Ala
20

<210> SEQ ID NO 15
<211> LENGTH: 24
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PAS peptide

<400> SEQUENCE: 15

Ala Ala Ser Pro Ala Ala Pro Ser Ala Pro Pro Ala Ala Ala Ser Pro
1 5 10 15

Ala Ala Pro Ser Ala Pro Pro Ala
20

-continued

<210> SEQ ID NO 16
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: PAS peptide

<400> SEQUENCE: 16

Ala Ser Ala Ala Ala Pro Ala Ala Ala Ser Ala Ala Ser Ala Pro
 1 5 10 15
 Ser Ala Ala Ala
 20

<210> SEQ ID NO 17
 <211> LENGTH: 21
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: fragment of Factor VIII B domain

<400> SEQUENCE: 17

Ser Phe Ser Gln Asn Ser Arg His Pro Ser Gln Asn Pro Pro Val Leu
 1 5 10 15
 Lys Arg His Gln Arg
 20

<210> SEQ ID NO 18
 <211> LENGTH: 6
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: His tag

<400> SEQUENCE: 18

His His His His His His
 1 5

40

What is claimed is:

1. A pharmaceutical composition comprising:
 - a. a chimeric polypeptide comprising a FVIII portion and an Fc region;
 - b. about 10 mg/mL to about 25 mg/mL sucrose;
 - c. about 10 mg/mL to about 13 mg/mL sodium chloride (NaCl);
 - d. about 0.75 mg/mL to about 2.25 mg/mL L-histidine;
 - e. about 5 mM to about 10 mM calcium chloride; and
 - f. about 0.08 mg/mL to about 0.25 mg/mL polysorbate 20 or polysorbate 80.
2. The pharmaceutical composition of claim 1, wherein the chimeric polypeptide comprises a processed FVIII portion.
3. The pharmaceutical composition of claim 1, wherein the chimeric polypeptide comprises a single chain FVIII portion.
4. A pharmaceutical kit comprising:
 - (a) a first container comprising a lyophilizate, where the lyophilizate comprises
 - (i) a chimeric polypeptide comprising a FVIII portion and an Fc region,
 - (ii) about 40 mg of sucrose;
 - (iii) about 36 mg of sodium chloride (NaCl);
 - (iv) about 3.1 mg of L-histidine;
 - (v) about 2.40 mg calcium chloride dihydrate; and
 - (vi) about 0.40 mg polysorbate 20 or polysorbate 80;
 - (b) a second container comprising sterile water for injections at a volume sufficient to produce, when combined with the lyophilizate of the first container, a solution comprising:
 - (i) the chimeric polypeptide;
 - (ii) about 13.3 mg/ml of sucrose;
 - (iii) about 12.0 mg/ml of NaCl;
 - (iv) about 1.03 mg/ml of L-histidine;
 - (v) about 5.4 mM of calcium chloride; and
 - (vi) about 0.13 mg/ml of polysorbate 20 or polysorbate 80.
5. The pharmaceutical kit of claim 4, wherein mannitol, glycine, alanine, or hydroxyethyl starch is not included.
6. The pharmaceutical kit of claim 4, wherein NaCl is the only bulking agent.
7. The pharmaceutical kit of claim 4, wherein the first container is a glass vial comprising a rubber stopper.
8. The pharmaceutical kit of claim 4, wherein the second container is a syringe body.
9. The pharmaceutical kit of claim 4, and wherein the syringe body is associated with a plunger.
10. The pharmaceutical composition of claim 1, wherein the Fc region comprises an amino acid sequence that is at least 95% identical to amino acids 21 to 247 of SEQ ID NO: 4.

11. The pharmaceutical composition of claim 10, wherein the Fc region comprises an amino acid sequence that is identical to amino acids 21 to 246 of SEQ ID NO: 4.

12. The pharmaceutical composition of claim 10, wherein the chimeric polypeptide is an rFVIII^h monomer-dimer hybrid.

13. The pharmaceutical composition of claim 12, wherein the Fc region is a first Fc region, and the rFVIII^h monomer-dimer hybrid comprises a second Fc region that comprises an amino acid sequence that is at least 95% identical to amino acids 21 to 247 of SEQ ID NO: 4.

14. The pharmaceutical composition of claim 13, wherein the first Fc region comprises an Fc domain with a hinge region, wherein the second Fc region comprises an Fc domain with a hinge region, and wherein the first Fc region is covalently bound to the second Fc region via a disulfide bond.

15. The pharmaceutical composition of claim 14, wherein the first Fc region comprises an amino acid sequence at least 99% identical to amino acids 21 to 247 of SEQ ID NO: 4, and wherein the second Fc region comprises an amino acid sequence at least 99% identical to amino acids 21 to 247 of SEQ ID NO: 4.

16. The pharmaceutical composition of claim 1, which comprises about 50 IU/mL to about 2500 IU/mL of a combination of rFVIII^h monomer-dimer hybrids comprising (i) a first rFVIII^h monomer-dimer hybrid in which the FVIII portion is a single chain FVIII; and (ii) a second rFVIII^h monomer-dimer hybrid in which the FVIII portion is a processed Factor VIII.

17. The pharmaceutical composition of claim 16, wherein the single chain FVIII comprises an amino acid sequence that is at least 95% identical to amino acids 20 to 1457 of SEQ ID NO: 2.

18. The pharmaceutical composition of claim 17, wherein the single chain FVIII comprises an amino acid sequence identical to amino acids 20 to 1457 of SEQ ID NO: 2.

19. The pharmaceutical kit of claim 4, wherein the Fc region comprises an amino acid sequence that is at least 95% identical to amino acids 21 to 247 of SEQ ID NO: 4.

20. The pharmaceutical kit of claim 19, wherein the Fc region comprises an amino acid sequence that is identical to amino acids 21 to 246 of SEQ ID NO: 4.

21. The pharmaceutical kit of claim 19, wherein the chimeric polypeptide is an rFVIII^h monomer-dimer hybrid.

22. The pharmaceutical kit of claim 21, wherein the Fc region is a first Fc region, and the rFVIII^h monomer-dimer hybrid comprises a second Fc region that comprises an amino acid sequence that is at least 95% identical to amino acids 21 to 247 of SEQ ID NO: 4.

23. The pharmaceutical kit of claim 21, wherein the first Fc region comprises an Fc domain with a hinge region, wherein the second Fc region comprises an Fc domain with a hinge region, and wherein the first Fc region is covalently bound to the second Fc region via a disulfide bond.

24. The pharmaceutical kit of claim 23, wherein the first Fc region comprises an amino acid sequence at least 99% identical to amino acids 21 to 247 of SEQ ID NO: 4, and wherein the second Fc region comprises an amino acid sequence at least 99% identical to amino acids 21 to 247 of SEQ ID NO: 4.

25. The pharmaceutical kit of claim 4, wherein the lyophilizate comprises about 50 IU/mL to about 2500 IU/mL of a combination of rFVIII^h monomer-dimer hybrids comprising (i) a first rFVIII^h monomer-dimer hybrid in which

the FVIII portion is a single chain FVIII; and (ii) a second rFVIII^h monomer-dimer hybrid in which the FVIII portion is a processed Factor VIII.

26. The pharmaceutical kit of claim 25, wherein the single chain FVIII comprises an amino acid sequence that is at least 95% identical to amino acids 20 to 1457 of SEQ ID NO: 2.

27. The pharmaceutical kit of claim 26, wherein the single chain FVIII comprises an amino acid sequence identical to amino acids 20 to 1457 of SEQ ID NO: 2.

28. A pharmaceutical kit comprising:

(a) a first container comprising a lyophilizate, where the lyophilizate comprises

(i) a chimeric polypeptide comprising a FVIII portion and an Fc region;

(ii) sucrose;

(iii) sodium chloride (NaCl);

(iv) L-histidine;

(v) calcium chloride; and

(vi) polysorbate 20; and

(b) a second container comprising sterilized water,

wherein, when the lyophilizate of the first container is combined with the sterilized water of the second container, a solution comprising the following is produced:

(i) the chimeric polypeptide;

(ii) about 10 mg/mL to about 25 mg/mL sucrose;

(iii) about 10 mg/mL to about 13 mg/mL sodium chloride (NaCl);

(iv) about 0.75 mg/mL to about 2.25 mg/mL L-histidine;

(v) about 5 mM to about 10 mM calcium chloride; and

(vi) about 0.08 mg/mL to about 0.25 mg/mL polysorbate 20.

29. The pharmaceutical kit of claim 28, wherein the Fc region comprises an amino acid sequence that is at least 95% identical to amino acids 21 to 247 of SEQ ID NO: 4.

30. The pharmaceutical kit of claim 29, wherein the Fc region comprises an amino acid sequence that is identical to amino acids 21 to 246 of SEQ ID NO: 4.

31. The pharmaceutical kit of claim 29, wherein the chimeric polypeptide is an rFVIII^h monomer-dimer hybrid.

32. The pharmaceutical kit of claim 29, wherein the Fc region is a first Fc region, and the rFVIII^h monomer-dimer hybrid comprises a second Fc region that comprises an amino acid sequence that is at least 95% identical to amino acids 21 to 247 of SEQ ID NO: 4.

33. The pharmaceutical kit of claim 31, wherein the first Fc region comprises an Fc domain with a hinge region, wherein the second Fc region comprises an Fc domain with a hinge region, and wherein the first Fc region is covalently bound to the second Fc region via a disulfide bond.

34. The pharmaceutical kit of claim 33, wherein the first Fc region comprises an amino acid sequence at least 99% identical to amino acids 21 to 247 of SEQ ID NO: 4, and wherein the second Fc region comprises an amino acid sequence at least 99% identical to amino acids 21 to 247 of SEQ ID NO: 4.

35. The pharmaceutical kit of claim 31, wherein the lyophilizate comprises about 50 IU/mL to about 2500 IU/mL of a combination of rFVIII^h monomer-dimer hybrids comprising (i) a first rFVIII^h monomer-dimer hybrid in which the FVIII portion is a single chain FVIII; and (ii) a second rFVIII^h monomer-dimer hybrid in which the FVIII portion is a processed Factor VIII.

121

36. The pharmaceutical kit of claim 35, wherein the single chain FVIII comprises an amino acid sequence that is at least 95% identical to amino acids 20 to 1457 of SEQ ID NO: 2.

37. The pharmaceutical kit of claim 36, wherein the single chain FVIII comprises an amino acid sequence identical to amino acids 20 to 1457 of SEQ ID NO: 2. 5

38. The pharmaceutical kit of claim 28, wherein mannitol, glycine, alanine, or hydroxyethyl starch is not included.

39. The pharmaceutical kit of claim 28, wherein NaCl is the only bulking agent. 10

40. The pharmaceutical kit of claim 28, wherein the first container is a glass vial comprising a rubber stopper.

41. The pharmaceutical kit of claim 28, wherein the second container is a syringe body.

42. The pharmaceutical kit of claim 41, and wherein the syringe body is associated with a plunger. 15

43. The pharmaceutical composition of claim 1, wherein the polysorbate 20 or polysorbate 80 is polysorbate 20.

44. The pharmaceutical kit of claim 4, wherein the polysorbate 20 or polysorbate 80 is polysorbate 20. 20

* * * * *

122