



(12) **United States Patent**
Jang et al.

(10) **Patent No.: US 12,414,984 B2**
(45) **Date of Patent: Sep. 16, 2025**

(54) **FUSION PROTEIN COMPRISING ANTI-CD73 ANTIBODY AND IL-2, AND USE THEREOF**

(71) Applicant: **GI INNOVATION, INC.**, Seoul (KR)

(72) Inventors: **Myung Ho Jang**, Seoul (KR); **Young Jun Koh**, Seoul (KR); **Dan Bee Ha**, Seoul (KR)

(73) Assignee: **GI INNOVATION, INC.**, Seoul (KR)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **18/815,390**

(22) Filed: **Aug. 26, 2024**

(65) **Prior Publication Data**

US 2024/0408176 A1 Dec. 12, 2024

Related U.S. Application Data

(63) Continuation of application No. PCT/KR2023/008664, filed on Jun. 22, 2023.

(30) **Foreign Application Priority Data**

Jun. 22, 2022 (KR) 10-2022-0076279

(51) **Int. Cl.**

C07K 16/46 (2006.01)

A61K 38/20 (2006.01)

A61K 39/00 (2006.01)

A61K 39/395 (2006.01)

A61P 35/00 (2006.01)

C07K 14/55 (2006.01)

C07K 16/28 (2006.01)

(52) **U.S. Cl.**

CPC **A61K 38/2013** (2013.01); **A61K 39/39558** (2013.01); **A61P 35/00** (2018.01); **C07K 14/55** (2013.01); **C07K 16/2896** (2013.01); **C07K 2317/76** (2013.01); **C07K 2317/92** (2013.01)

(58) **Field of Classification Search**

None

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,229,109 A * 7/1993 Grimm A61P 37/04 424/85.1
2003/0124678 A1 * 7/2003 Epstein A61P 37/00 435/325
2007/0036752 A1 * 2/2007 Gillies A61P 35/00 435/325
2020/0071404 A1 3/2020 Sato et al.
2020/0369740 A1 * 11/2020 Jang A61P 35/00
2021/0024646 A1 * 1/2021 Yu A61K 47/6851
2022/0125884 A1 4/2022 Baca et al.
2022/0380781 A1 12/2022 Jang
2022/0389070 A1 12/2022 Jang
2023/0104217 A1 4/2023 Jang
2024/0352084 A1 * 10/2024 Gonzalez A61P 35/00

FOREIGN PATENT DOCUMENTS

KR 10-2019-0091281 A 8/2019
KR 10-2020-0032009 A 3/2020
KR 10-2021-0008110 A 1/2021
KR 10-2313505 B1 10/2021
WO WO-2009061853 * 5/2009 C07K 14/55
WO 2021/043229 A1 3/2021
WO 2022/087149 A2 4/2022

OTHER PUBLICATIONS

Silver et al. Engineered antibody fusion proteins for targeted disease therapy. Trends in Pharmacological Sciences, Dec. 2021, 42(12):1064-1081 (Year: 2021).*

Skrombolas and Frelinger. Challenges and developing solutions for increasing the benefits of IL-2 treatment in tumor therapy. Expert Rev Clin Immunol. Feb. 2014;10(2):207-217 (Year: 2014).*

Hu et al. Generation of low-toxicity interleukin-2 fusion proteins devoid of vasopermeability activity. Blood 101, 4853-4861 (2003) (Year: 2003).*

Heaton et al. Human interleukin 2 analogues that preferentially bind the intermediate-affinity interleukin 2 receptor lead to reduced secondary cytokine secretion: implications for the use of these interleukin 2 analogues in cancer immunotherapy. Cancer Res 53, 2597-602, 1993 (Year: 1993).*

Heuser et al. Anti-CD30-scFv-Fc-IL-2 antibody-cytokine fusion protein that induces resting NK cells to highly efficient cytotoxicity of Hodgkin's lymphoma derived tumour cells. Int. J. Cancer: 110, 386-394 (2004) (Year: 2004).*

Shi et al. A recombinant anti-erbB2, scFv-Fc-IL-2 fusion protein retains antigen specificity and cytokine function. Biotechnology Letters 25: 815-819, 2003 (Year: 2003).*

Shanafelt et al. (Nature Biotechnol 18, 1197-1202 (2000)) (Year: 2000).*

International Search Report for International Application No. PCT/KR2023/008664 dated Sep. 21, 2023.

Communication dated Apr. 17, 2025, issued in Korean Application No. 10-2023-0080204.

* cited by examiner

Primary Examiner — Maher M Haddad

(74) *Attorney, Agent, or Firm* — Sughrue Mion, PLLC

(57) **ABSTRACT**

A fusion protein in which an anti-CD73 antibody or an antigen-binding fragment thereof and IL-2 are linked to each other, and the use thereof are disclosed. The fusion protein or dimer thereof contains an anti-CD73 antibody or an antigen-binding fragment thereof and IL-2, and is able to inhibit adenosine production by binding to CD73 of cancer cells, thereby blocking the A2AR and A2BR signaling pathways, thus controlling the tumor microenvironment. Furthermore, the fusion protein or dimer thereof is able to activate immune cells (CD8+ T cells and CD4+ T cells) along with less activation of immunosuppressive cells such as regulatory T cells (Tregs). In addition, IL-2 or a variant thereof in the fusion protein is able to activate immune cells. Therefore, the fusion protein and dimer thereof may be useful for preventing or treating cancer.

12 Claims, 16 Drawing Sheets

Specification includes a Sequence Listing.

FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19




FIG. 20




FIG. 21

Tumor Growth Inhibition (TGI)




FIG. 22




FIG. 23




FIG. 24




FIG. 25




FIG. 26




FIG. 27





FIG. 28



1

FUSION PROTEIN COMPRISING ANTI-CD73 ANTIBODY AND IL-2, AND USE THEREOF

CROSS REFERENCE TO RELATED APPLICATIONS

This application is a Bypass Continuation Application of International Application No. PCT/KR2023/008664, filed on Jun. 22, 2023, which claims priority to Korean Patent Application No. 10-2022-0076279, filed on Jun. 22, 2022, the disclosures of which are incorporated by reference herein their entireties.

INCORPORATION BY REFERENCE OF SEQUENCE LISTING

The content of the electronically submitted sequence listing, file name: Q301220_SEQ_LIS_AS_FILED.xml; size: 71.2 KB; and date of creation: Aug. 23, 2024, filed herewith, is incorporated herein by reference in its entirety.

TECHNICAL FIELD

The present invention relates to a fusion protein in which an anti-CD73 antibody or an antigen-binding fragment thereof and IL-2 are linked to each other, and the use thereof.

BACKGROUND ART

Cancer immunotherapy is a method of treating cancer using the body's immune system. It can induce the immune system to attack cancer cells by targeting antigens such as cancer cell surface proteins and can increase the activity of immune cells by controlling the tumor microenvironment that causes immune evasion of cancer cells.

The tumor microenvironment is an environment in which not only cancer cells, but also fibroblasts, blood vessels, lymphatic vessels, immune cells, extracellular matrix, and adipocytes, which exist in cancer tissues, proliferate and evolve. In this regard, studies have reported that the ATP-AMP-adenosine-A2AR/A2BR signaling pathway can regulate the activity of immune cells in the tumor microenvironment (S. Vigano et al., Front Immunol (2019) 10:925). Specifically, when adenosine binds to the A2AR/A2BR receptor on the surface of tumor cells or immune cells, immune suppressor cells such as regulatory T cells or tumor-associated macrophages (TAMs) are activated, and the activity of cytotoxic T cells and NK cells is inhibited. In this regard, studies have been conducted on a method of activating immunity in the tumor microenvironment by inhibiting this signaling pathway by treatment with a substance that target CD39, an ectonucleotidase that converts ATP to AMP, CD73, an ectonucleotidase that converts AMP to adenosine, or A2AR/A2BR receptors that bind to adenosine.

Meanwhile, interleukin 2 (IL-2), also called T-cell growth factor (TCGF), is a globular glycoprotein that plays a central role in the production, survival, and homeostasis of lymphocytes. IL-2 mediates various immune actions by binding to an IL-2 receptor composed of three distinct subunits.

In addition, IL-2 is synthesized mainly by activated T cells, particularly CD4+ helper T cells. IL-2 stimulates the proliferation and differentiation of T cells and induces the production of cytotoxic T lymphocytes (CTLs) and the differentiation of peripheral blood lymphocytes into cytotoxic cells and lymphokine-activated killer cells (LAK cells).

2

Throughout the present specification, a number of publications and patent documents are referred to and cited. The disclosure of these cited publications and patent documents is incorporated herein by reference in its entirety to more clearly describe the state of the art to which the present invention pertains and the content of the present invention.

DISCLOSURE

Technical Problem

Accordingly, the present inventors have conducted studies to develop a new combination of proteins as a fusion protein that enhances the activity of immune cells by controlling the tumor microenvironment. As a result, they have discovered that a fusion protein dimer comprising an anti-CD73 antibody or an antigen-binding fragment thereof and IL-2 can control the tumor microenvironment and effectively activates immune cells. Based on this finding, the present inventors have demonstrated that the fusion protein dimer is effective as an anticancer agent, thereby completing the present invention.

Technical Solution

To achieve the above object, one aspect of the present invention provides a fusion protein comprising: an antibody or a fragment thereof that specifically binds to CD73; and IL-2.

Another aspect of the present invention provides a fusion protein dimer in which the two fusion proteins are linked to each other.

Still another aspect of the present invention provides a polynucleotide encoding the fusion protein, an expression vector containing the polynucleotide, and a transformed cell into which the expression vector has been introduced.

Yet another aspect of the present invention provides a method for producing a fusion protein dimer comprising: an antibody or an antigen-binding fragment thereof that specifically binds to CD73; and IL-2, the method comprising steps of: i) culturing the transformed cell; and ii) collecting the fusion protein dimer.

Still yet another aspect of the present invention provides a pharmaceutical composition for preventing or treating cancer comprising the fusion protein or the fusion protein dimer as an active ingredient.

A further aspect of the present invention provides a method for preventing or treating cancer, comprising a step of administering the fusion protein or the fusion protein dimer to a subject.

Another further aspect of the present invention provides the use of the fusion protein or the fusion protein dimer for preventing or treating cancer.

Advantageous Effects

The fusion protein or dimer thereof according to the present invention, which comprises an anti-CD73 antibody or an antigen-binding fragment thereof and IL-2, is able to inhibit adenosine production (ATP→AMP→adenosine) by binding to CD73 in cancer cells. This inhibition blocks the A2AR and A2BR signaling pathways, thereby controlling the tumor microenvironment. Furthermore, the fusion protein or dimer thereof is able to activate immune cells (CD8+ T cells and CD4+ T cells) along with less activation of immunosuppressive cells such as regulatory T cells (Tregs). In addition, IL-2 or a variant thereof in the fusion protein of

the present invention is able to activate immune cells. Therefore, the fusion protein and dimer thereof of the present invention may be useful for the prevention or treatment of cancer.

BRIEF DESCRIPTION OF DRAWINGS

FIG. 1 schematically illustrates the structure of a fusion protein dimer (GI-108A1), which is an example.

FIG. 2 schematically illustrates the structure of a fusion protein dimer (GI-108B1), which is an example.

FIG. 3 is a graph showing the results of measuring the activity of the JAK-STAT signaling pathway after treating HEK-BLUE™ IL-2 reporter cells with two types of GI-108A1, GI-108B1, and PROLEUKIN®.

FIG. 4 is a graph showing the results of evaluating the degree of proliferation of CTLL-2 cells after treating the cells with two types of GI-108A1, GI-108B1, and PROLEUKIN®.

FIG. 5 is a graph showing the results of evaluating the ability of two types of GI-108A1, GI-108B1, and oleclumab to inhibit the enzymatic activity of membrane-bound human CD73 in human TNBC cell line (MDA-MB-231).

FIG. 6 is a graph showing the results of evaluating the ability of two types of GI-108A1, GI-108B1, and oleclumab to inhibit the enzymatic activity of soluble CD73.

FIG. 7 is a graph showing the proliferation of CD8+ T cells after treating CD8+ T cells, the activity of which was inhibited by CD73-adenosine, with vehicle (hIgG4), GI-108B1 alone, PROLEUKIN® alone, IgG4 Fc-IL-2v alone, anti-CD73 antibody (GI-αCD73) alone, a combination of anti-CD73 antibody (GI-αCD73) and IgG4 Fc-IL-2v, oleclumab alone, or a combination of oleclumab and IgG4 Fc-IL-2v.

FIG. 8 is a graph showing the number of IFN-γ+CD8+ T cells after treating CD8+ T cells, the activity of which was inhibited by CD73-adenosine, with vehicle (hIgG4), GI-108B1 alone, PROLEUKIN® alone, IgG4 Fc-IL-2v alone, anti-CD73 antibody (GI-αCD73) alone, a combination of anti-CD73 antibody (GI-αCD73) and IgG4 Fc-IL-2v, oleclumab alone, or a combination of oleclumab and IgG4 Fc-IL-2v.

FIG. 9 is a graph showing the proliferation of CD4+ T cells after treating CD4+ T cells, the activity of which was inhibited by CD73-adenosine, with vehicle (hIgG4), GI-108B1 alone, PROLEUKIN® alone, IgG4 Fc-IL-2v alone, anti-CD73 antibody (GI-αCD73) alone, a combination of anti-CD73 antibody (GI-αCD73) and IgG4 Fc-IL-2v, oleclumab alone, or a combination of oleclumab and IgG4 Fc-IL-2v.

FIG. 10 is a graph showing the number of IFN-γ+CD4+ T cells after treating CD4+ T cells, the activity of which was inhibited by CD73-adenosine, with vehicle (hIgG4), GI-108B1 alone, PROLEUKIN® alone, IgG4 Fc-IL-2v alone, anti-CD73 antibody (GI-αCD73) alone, a combination of anti-CD73 antibody (GI-αCD73) and IgG4 Fc-IL-2v, oleclumab alone, or a combination of oleclumab and IgG4 Fc-IL-2v.

FIG. 11 is a graph showing the results of measuring pSTAT5 in the CD8+ T cell population after treating human PBMCs with various concentrations of PROLEUKIN®, GI-108B1, anti-CD73 antibody (GI-αCD73) or IgG4 Fc-IL-2v alone, or a combination of anti-CD73 antibody (GI-αCD73) and IgG4 Fc-IL-2v.

FIG. 12 is a graph showing the EC₅₀ values obtained by measuring pSTAT5 in the CD8+ T cell population after

treating human PBMCs with various concentrations of PROLEUKIN® or GI-108B1.

FIG. 13 is a graph showing the results of measuring pSTAT5 in the Treg cell population after treating human PBMCs with various concentrations of PROLEUKIN®, GI-108B1, anti-CD73 antibody (GI-αCD73) or IgG4 Fc-IL-2v alone, or a combination of anti-CD73 antibody (GI-αCD73) and IgG4 Fc-IL-2v.

FIG. 14 is a graph showing the EC₅₀ values obtained by measuring pSTAT5 in the Treg cell population after treating human PBMCs with various concentrations of PROLEUKIN® or GI-108B1.

FIG. 15 shows the results of measuring the number of CD8+ T cells, the number of Treg cells, and the CD8+ T/Treg ratio after 12 days of culture of human PBMCs treated with PROLEUKIN® or GI-108B1.

FIG. 16 schematically illustrates that STAT5 phosphorylation is induced when GI-108B1 cis-binds to CD73 and IL-2 receptors in the CD8+ T cells and CTV+CD8+ T cells pretreated with anti-CD73 antibody.

FIG. 17 is a graph showing the results of measuring pSTAT5+(CTV+/-)CD8+ T cells in a co-culture sample of CTV+CD8+ T cells and CD8+ T cells pretreated with anti-CD73 antibody and a co-culture sample of CTV+CD8+ T cells and CD8+ T cells after treating each of the samples with GI-108B1.

FIG. 18 is a graph showing the results of flow cytometry analysis of hCD73+CD8+ T cells in purified CD8+ T cells from human PBMCs.

FIG. 19 is a graph showing the results of measuring the cell death rate after co-culture of MDA-MB-231 cells and PBMCs, treated with hIgG4 or GI-108B1 alone or a combination of anti-CD73 antibody (GI-αCD73) and IgG4 Fc-IL-2v, in the presence of anti-CD3/CD28 dynabeads and/or AMP.

FIG. 20 is a graph showing the results of measuring the tumor volume after administering vehicle (PBS) or GI-108B1 to mice transplanted with murine colon cancer cells (MC38).

FIG. 21 is a graph showing the tumor growth inhibition rate after administering vehicle (PBS) or GI-108B1 to mice transplanted with murine colon cancer cells (MC38).

FIG. 22 is a graph showing the survival rate up to day 25 after administering vehicle (PBS) or GI-108B1 to mice transplanted with murine colon cancer cells (MC38).

FIG. 23 schematically shows an experimental plan conducted using mice transplanted with human breast cancer cells (MDA-MB-231) in order to evaluate the anticancer effects of GI-108A1 and GI-108B1.

FIG. 24 is a graph showing the results of measuring the tumor volume up to day 18 after administering vehicle (hIgG4), GI-108A1 (AHP04167), GI-108B1, or anti-PD-1 antibody to mice transplanted with human breast cancer cells (MDA-MB-231).

FIG. 25 is a graph showing the results of measuring the tumor volumes in individual experimental animals of each treated group after administering vehicle (hIgG4), GI-108A1 (AHP04167), GI-108B1, or anti-PD-1 antibody to mice transplanted with human breast cancer cells (MDA-MB-231).

FIG. 26 is a graph showing the results of measuring the tumor growth inhibition rate on day 18 after administering vehicle (hIgG4), GI-108A1 (AHP04167), GI-108B1, or anti-PD-1 antibody to mice transplanted with human breast cancer cells (MDA-MB-231).

FIG. 27 is a graph showing the results of measuring the body weight of each mouse group up to day 18 after

administering vehicle (hIgG4), GI-108A1 (AHP04167), GI-108B1, or anti-PD-1 antibody to mice transplanted with human breast cancer cells (MDA-MB-231).

FIG. 28 is a graph showing the results of measuring the vascular permeability of each drug after treating and culturing HUVEC cells, cultured in a trans-well system, with a culture supernatant of PBMCs and TNF- α , PROLEUKIN® or GI-108B1.

BEST MODE

Fusion Protein Comprising: Antibody or Antigen-Binding Fragment Thereof that Specifically Binds to CD73; and IL-2

One aspect of the present invention provides a fusion protein comprising: an antibody or an antigen-binding fragment thereof that specifically binds to CD73; and IL-2. CD73 and Antibody or Antigen-Binding Fragment Thereof that Specifically Binds to CD73

As used in the present specification, the term “CD73 (Cluster of Differentiation 73)” refers to an ectonucleotidase that catalyzes the dephosphorylation of 5' nucleotides, primarily converting adenosine monophosphate (AMP) to adenosine. CD73 forms a homodimer on the cell membrane via a GPI anchor. Each monomer is 65 kDa, with its N-terminal and C-terminal domains connected by a flexible helical linker.

It is known that CD73 is involved in the production of adenosine, is overexpressed in cancer cells, and induces immunosuppression. CD73 is expressed in many tumor cells, including leukemia, bladder cancer, glioma, glioblastoma, ovarian cancer, melanoma, prostate cancer, thyroid cancer, esophageal cancer, and breast cancer. In addition, CD73 is known to be expressed on the surface of immunosuppressive cells (including regulatory T cells (Tregs) and myeloid-derived suppressor cells (MDSCs)). In particular, overexpression of CD73 has been reported to be associated with angiogenesis, invasion, resistance to chemotherapy, tumor metastasis, and short survival time of cancer patients with various tumors, including breast cancer and melanoma.

In the present invention, CD73 may be mammalian CD73, preferably human CD73, without limitation. In addition, in the present invention, the CD73 protein includes, but is not limited to, both native and mutant CD73 proteins. The native CD73 protein generally refers to a polypeptide comprising the amino acid sequence of the native CD73 protein, and the amino acid sequence of the native CD73 protein generally refers to the amino acid sequence found in naturally occurring CD73. Information about CD73 is available from known databases such as GenBank of the National Institutes of Health. For example, CD73 may have the amino acid sequence (SEQ ID NO: 41) corresponding to GenBank accession number NP_002517.1, but it is not limited to this sequence.

As used in the present specification, the term “antigen” refers to a molecule capable of selectively binding to an antibody. The target antigen may be a polypeptide, a carbohydrate, a nucleic acid, a lipid, a hapten, or other naturally occurring or synthetic compound. Specifically, the antigen is a polypeptide and may be a protein present on the cell surface or within the cell.

As used in the present specification, the term “specifically binds” means binding that is measurably different from a non-specific interaction. Specific binding can be determined by competition with a control molecule that is similar to the target that has no binding activity.

The antibody or antigen-binding fragment thereof that specifically binds to CD73 may generally refer to molecules

capable of specifically forming an antigen-antibody complex with CD73. In addition, the antibody or antigen-binding fragment thereof may be used in any form as long as it contains an antigen-binding site capable of specifically binding to CD73. The antibody or antigen-binding fragment thereof may include other amino acids that are not directly involved in binding, or amino acids whose effects are blocked by the residues of the antigen-binding site.

As used in the present specification, the term “antibody” refers to molecules containing an antigen-binding site, and immunologically active fragments of immunoglobulin molecules containing an antigen-binding site. The immunoglobulin molecules may be immunoglobulin molecules of IgG, IgE, IgM, IgD, IgA, IgY, or a subclass thereof. The heavy and light chains of immunoglobulins may each include a constant region and a variable region. The light and heavy chain variable regions of immunoglobulins contain three variable regions, called complementarity determining regions (CDRs), and four framework regions (FRs). The CDRs are antigen-binding sites that mainly bind to the epitope of the antigen.

The antibodies or antigen-binding fragments thereof of the present invention may include monoclonal antibodies, polyclonal antibodies, synthetic antibodies, human antibodies, humanized antibodies, non-human antibodies, and any fragments thereof, as well as immunoconjugates.

As used in the present specification, the term “antigen-binding fragment of the antibody” refers to a portion of a polypeptide within the overall structure of an immunoglobulin to which an antigen can bind. Examples of the antigen-binding fragment include, but are not limited to, Fab fragments, Fab' fragments, F(ab')₂ fragments, bispecific Fab dimers (Fab₂), trispecific Fab trimers (Fab₃), Fv, single-chain Fv proteins (“scFv”), bis-scFv (scFv)₂, minibodies, diabodies, triabodies, tetrabodies, disulfide stabilized Fv proteins (“dsFv”), single-domain antibodies (sdAb), and portions of full-length antibodies responsible for antigen binding. Regardless of the structure, the fragment of the antibody is able to bind to the same antigen recognized by the intact antibody.

The single-domain antibody, also called a nanobody, refers to an antibody fragment consisting of a single monomeric variable antibody domain. The single-domain antibody is able to specifically bind to a specific antigen like a whole antibody. The single-domain antibody has a molecular weight of 12 to 15 kDa, which is smaller than that of common antibodies (150-160 kDa), Fab fragments (about 50 kDa), and scFv (about 25 kDa). The single-domain antibody may be obtained, for example, by monomerizing the dimeric variable domains of human or mouse conventional immunoglobulin G (IgG), and may be VHH derived from the heavy chain antibody of Camelids, or a VNAR fragment obtained from IgNAR of Cartilaginous fishes. IL-2 or Variant Thereof

As used in the present specification, the term “IL-2” or “interleukin-2” refers to any wild-type IL-2 obtained from any vertebrate source, including mammals such as primates (e.g., humans) and rodents (e.g., mice and rats), unless otherwise stated. IL-2 has a protein size of 15.5 kDa to 16 kDa and consists of 133 amino acids. IL-2 may be obtained from animal cells, but also includes one obtained from recombinant cells capable of producing IL-2. In addition, IL-2 may be wild-type IL-2 or a variant thereof.

In the present specification, IL-2 or a variant thereof may be collectively expressed by the term “IL-2 protein” or “IL-2 polypeptide”. IL-2, an IL-2 protein, an IL-2 polypeptide, and an IL-2 variant specifically bind to, for example, an IL-2

receptor. This specific binding may be identified by methods known to those skilled in the art.

IL-2 may also be in a mature form. Specifically, the mature IL-2 may not contain a signal sequence, and may have the amino acid sequence of SEQ ID NO: 40. Here, IL-2 may be used under a concept encompassing a fragment in which a portion of the N-terminus or C-terminus of wild-type IL-2 is truncated.

As used in the present specification, the term “IL-2 variant” refers to a form in which a portion of the amino acids in the full-length IL-2 or the above-described fragment of IL-2 is substituted. That is, an IL-2 variant may have an amino acid sequence different from wild-type IL-2 or a fragment thereof. However, the IL-2 variant may have activity equivalent or similar to the wild-type IL-2. Here, “IL-2 activity” may, for example, refer to specific binding to an IL-2 receptor, in which the specific binding can be measured by methods known to those skilled in the art.

Specifically, the IL-2 variant may be obtained by substitution of a portion of the amino acids in the wild-type IL-2. An embodiment of the IL-2 variant obtained by amino acid substitution may be obtained by substitution of at least one of the 38th, 42nd, 45th, 61st, and 72nd amino acids in the amino acid sequence of SEQ ID NO: 40.

Specifically, the IL-2 variant may be obtained by substituting at least one of the 38th, 42nd, 45th, 61st, and 72nd amino acids in the amino acid sequence of SEQ ID NO: 40 with another amino acid. In addition, when IL-2 is in a form in which a portion of the N-terminus in the amino acid sequence of SEQ ID NO: 40 is truncated, the amino acid at a position complementarily corresponding to that in the amino acid sequence of SEQ ID NO: 40 may be substituted with another amino acid. According to one embodiment, one, two, three, four, five, six, seven, eight, nine, or ten amino acids may be substituted as long as such an IL-2 variant maintains IL-2 activity. According to another embodiment, one to five amino acids may be substituted.

In one embodiment, the IL-2 variant may be in a form in which the amino acids at two positions are substituted. Specifically, the IL-2 variant may be obtained by substitution of the 38th and 42nd amino acids in the amino acid sequence of SEQ ID NO: 40. In addition, in one embodiment, the IL-2 variant may be obtained by substitution of the 38th and 45th amino acids in the amino acid sequence of SEQ ID NO: 40. In addition, in one embodiment, the IL-2 variant may be obtained by substitution of the 38th and 61st amino acids in the amino acid sequence of SEQ ID NO: 40. In addition, in one embodiment, the IL-2 variant may be obtained by substitution of the 38th and 72nd amino acids in the amino acid sequence of SEQ ID NO: 40. In addition, in one embodiment, the IL-2 variant may be obtained by substitution of the 42nd and 45th amino acids in the amino acid sequence of SEQ ID NO: 40. In addition, in one embodiment, the IL-2 variant may be obtained by substitution of the 42nd and 61st amino acids in the amino acid sequence of SEQ ID NO: 40. In addition, in one embodiment, the IL-2 variant may be obtained by substitution of the 42nd and 72nd amino acids in the amino acid sequence of SEQ ID NO: 40. In addition, in one embodiment, the IL-2 variant may be obtained by substitution of the 45th and 61st amino acids in the amino acid sequence of SEQ ID NO: 40. In addition, in one embodiment, the IL-2 variant may be obtained by substitution of the 45th and 72nd amino acids in the amino acid sequence of SEQ ID NO: 40. In addition, in one embodiment, the IL-2 variant may be obtained by substitution of the 61st and 72nd amino acids in the amino acid sequence of SEQ ID NO: 40.

Furthermore, the IL-2 variant may be in a form in which the amino acids at three positions are substituted. Specifically, the IL-2 variant may be obtained by substitution of the 38th, 42nd, and 45th amino acids in the amino acid sequence of SEQ ID NO: 40. In addition, in one embodiment, the IL-2 variant may be obtained by substitution of the 38th, 42nd, and 61st amino acids in the amino acid sequence of SEQ ID NO: 40. In addition, in one embodiment, the IL-2 variant may be obtained by substitution of the 38th, 42nd, and 72nd amino acids in the amino acid sequence of SEQ ID NO: 40. In addition, in one embodiment, the IL-2 variant may be obtained by substitution of the 38th, 45th, and 61st amino acids in the amino acid sequence of SEQ ID NO: 40. In addition, in one embodiment, the IL-2 variant may be obtained by substitution of the 38th, 45th, and 72nd amino acids in the amino acid sequence of SEQ ID NO: 40. In addition, in one embodiment, the IL-2 variant may be obtained by substitution of the 38th, 61st, and 72nd amino acids in the amino acid sequence of SEQ ID NO: 40. In addition, in one embodiment, the IL-2 variant may be obtained by substitution of the 42nd, 45th, and 61st amino acids in the amino acid sequence of SEQ ID NO: 40. In addition, in one embodiment, the IL-2 variant may be obtained by substitution of the 42nd, 45th, and 72nd amino acids in the amino acid sequence of SEQ ID NO: 40. In addition, in one embodiment, the IL-2 variant may be obtained by substitution of the 45th, 61st, and 72nd amino acids in the amino acid sequence of SEQ ID NO: 40.

Here, the “another amino acid” introduced by substitution may be any one selected from the group consisting of alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine. However, regarding amino acid substitution for the IL-2 variant, in the amino acid sequence of SEQ ID NO: 40, the 38th amino acid cannot be substituted with arginine, the 42nd amino acid cannot be substituted with phenylalanine, the 45th amino acid cannot be substituted with tyrosine, the 61st amino acid cannot be substituted with glutamic acid, and the 72nd amino acid cannot be substituted with leucine.

Regarding amino acid substitution for the IL-2 variant, in the amino acid sequence of SEQ ID NO: 40, the 38th amino acid, arginine, may be substituted with an amino acid other than arginine. Preferably, regarding amino acid substitution for an IL-2 variant, in the amino acid sequence of SEQ ID NO: 40, the 38th amino acid, arginine, may be substituted with alanine (R38A).

Regarding amino acid substitution for the IL-2 variant, in the amino acid sequence of SEQ ID NO: 40, the 42nd amino acid, phenylalanine, may be substituted with an amino acid other than phenylalanine. Preferably, regarding amino acid substitution for the IL-2 variant, in the amino acid sequence of SEQ ID NO: 40, the 42nd amino acid, phenylalanine, may be substituted with alanine (F42A).

Regarding amino acid substitution for the IL-2 variant, in the amino acid sequence of SEQ ID NO: 40, the 45th amino acid, tyrosine, may be substituted with an amino acid other than tyrosine. Preferably, regarding amino acid substitution for the IL-2 variant, in the amino acid sequence of SEQ ID NO: 40, the 45th amino acid, tyrosine, may be substituted with alanine (Y45A).

Regarding amino acid substitution for the IL-2 variant, in the amino acid sequence of SEQ ID NO: 40, the 61st amino acid, glutamic acid, may be substituted with an amino acid other than glutamic acid. Preferably, regarding amino acid substitution for the IL-2 variant, in the amino acid sequence

of SEQ ID NO: 40, the 61st amino acid, glutamic acid, may be substituted with arginine (E61R).

Regarding amino acid substitution for an IL-2 variant, in the amino acid sequence of SEQ ID NO: 40, the 72nd amino acid, leucine, may be substituted with an amino acid other than leucine. Preferably, regarding amino acid substitution for an IL-2 variant, in the amino acid sequence of SEQ ID NO: 40, the 72nd amino acid, leucine, may be substituted with glycine (L72G).

Specifically, the IL-2 variant may be obtained by at least one substitution selected from the group consisting of R38A, F42A, Y45A, E61R, and L72G, in the amino acid sequence of SEQ ID NO: 40.

Specifically, the IL-2 variant may be obtained by two or three amino acid substitutions selected from the group consisting of R38A, F42A, Y45A, E61R, and L72G.

In addition, the IL-2 variant may be in a form in which the amino acids at two positions are substituted. Specifically, the IL-2 variant may be obtained by the substitutions R38A and F42A. In addition, in one embodiment, the IL-2 variant may be obtained by the substitutions R38A and Y45A. In addition, in one embodiment, the IL-2 variant may be obtained by the substitutions R38A and E61R. In addition, in one embodiment, the IL-2 variant may be obtained by the substitutions R38A and L72G. In addition, in one embodiment, the IL-2 variant may be obtained by the substitutions F42A and Y45A. In addition, in one embodiment, the IL-2 variant may be obtained by the substitutions F42A and L72G. In addition, in one embodiment, the IL-2 variant may be obtained by the substitutions E61R and L72G.

Furthermore, the IL-2 variant may be in a form in which the amino acids at three positions are substituted. Specifically, the IL-2 variant may be obtained by the substitutions R38A, F42A, and Y45A. In addition, in one embodiment, the IL-2 variant may be obtained by the substitutions R38A, F42A, and E61R. In addition, in one embodiment, the IL-2 variant may be obtained by the substitutions R38A, F42A, and L72G. In addition, in one embodiment, the IL-2 variant may be obtained by the substitutions R38A, Y45A, and E61R. In addition, in one embodiment, the IL-2 variant may be obtained by the substitutions R38A, Y45A, and L72G. In addition, in one embodiment, the IL-2 variant may be obtained by the substitutions F42A, Y45A, and E61R. In addition, in one embodiment, the IL-2 variant may be obtained by the substitutions F42A, Y45A, and L72G. In addition, in one embodiment, the IL-2 variant may be obtained by the substitutions F42A, E61R, and L72G. In addition, in one embodiment, the IL-2 variant may be obtained by the substitutions Y45A, E61R, and L72G.

In addition, the IL-2 variant may be in a form in which the amino acids at four positions are substituted. Specifically, the IL-2 variant may be obtained by the substitutions R38A, F42A, Y45A, and E61R. In addition, in one embodiment, the IL-2 variant may be obtained by the substitutions R38A, F42A, Y45A, and L72G. In addition, in one embodiment, the IL-2 variant may be obtained by the substitutions R38A, F42A, E61R, and L72G. In addition, in one embodiment, the IL-2 variant may be obtained by the substitutions R38A, Y45A, E61R, and L72G. In addition, in one embodiment, the IL-2 variant may be obtained by the substitutions F42A, Y45A, E61R, and L72G.

Furthermore, the IL-2 variant may be obtained by the substitutions R38A, F42A, Y45A, E61R, and L72G.

Preferably, the IL-2 variant of the present invention may be obtained by at least one substitution selected from the


group consisting of R38A, F42A, and E61R in the amino acid sequence of SEQ ID NO: 40. More preferably, the IL-2 variant of the present invention may be obtained by the substitutions R38A, F42A, and E61R in the amino acid sequence of SEQ ID NO: 40. Specifically, the IL-2 variant of the present invention may have the amino acid sequence of SEQ ID NO: 8.

In addition, the IL-2 variant may be characterized by having low in vivo toxicity. Here, the low in vivo toxicity may be a side effect caused by binding of IL-2 to the IL-2 receptor alpha chain (IL-2R α). IL-2 variants described in the present application have a low binding affinity for the IL-2 receptor alpha chain (IL-2R α) and thus have lower in vivo toxicity than the wild-type IL-2.

Structure of Fusion Protein Comprising: Antibody or Antigen-Binding Fragment Thereof that Specifically Binds to CD73; and IL-2 or Variant Thereof

The fusion protein of the present invention may comprise: a single-domain antibody that specifically binds to CD73; an Fc region; and IL-2 or a variant thereof. Here, the Fc region and the IL-2 or variant thereof may be linked by a linker.

Specifically, the fusion protein may comprise the following structural formula (I):



in structural formula (I) above,

N' is the N-terminus,

C' is the C-terminus,

X is the antibody or antigen-binding fragment thereof that specifically binds to CD73,

Y is IL-2,

linker (1) and linker (2) are peptide linkers, and

o and p are each independently 0 or 1.

Here, the antibody or fragment thereof that specifically binds to CD73 is as described above, and IL-2 is as described above.

In one embodiment, the antigen-binding fragment may comprise a single-domain antibody. In this case, the antibody or antigen-binding fragment thereof, specifically the single-domain antibody, may comprise: a CDR1 region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 15, 18, and 21; a CDR2 region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 16, 19, and 22; and a CDR3 region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 17, 20, and 23.

Specifically, the antigen-binding fragment may comprise: a CDR1 comprising the amino acid sequence of SEQ ID NO: 15, a CDR2 comprising the amino acid sequence of SEQ ID NO: 16, and a CDR3 comprising the amino acid sequence of SEQ ID NO: 17; or a CDR1 comprising the amino acid sequence of SEQ ID NO: 18, a CDR2 comprising the amino acid sequence of SEQ ID NO: 19, and a CDR3 comprising the amino acid sequence of SEQ ID NO: 20; or a CDR1 comprising the amino acid sequence of SEQ ID NO: 21, a CDR2 comprising the amino acid sequence of SEQ ID NO: 22, and a CDR3 comprising the amino acid sequence of SEQ ID NO: 23.

In one embodiment, the antigen-binding fragment may comprise any one amino acid sequence selected from the group consisting of SEQ ID NOs: 2, 3, and 4.

The peptide linker (1) may consist of 1 to 50 contiguous amino acids, or 3 to 30 contiguous amino acids, or 5 to 15 amino acids. In one embodiment, the peptide linker (1) may consist of 12 amino acids. In addition, the peptide linker 1 may contain at least one cysteine. Specifically, the peptide

11

linker (1) may contain one, two, or three cysteines. In addition, the peptide linker (1) may be derived from the hinge of an immunoglobulin. For example, the hinge may be selected from the hinge regions of various IgG subclass antibodies. In addition, the hinge may be a form in which one or more of the amino acids in the hinge region derived from an immunoglobulin are substituted with other amino acid(s), or may be a sequence obtained by adding one or more amino acids to the hinge region. In one embodiment, the peptide linker (1) may be a peptide linker consisting of the amino acid sequence of SEQ ID NO: 5.

The peptide linker (2) may consist of 1 to 30 contiguous amino acids, or 2 to 20 contiguous amino acids, or 2 to 10 amino acids. In one embodiment, the peptide linker (2) may be (G4S)_n (where n is an integer ranging from 1 to 10). In (G4S)_n, n may be 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10. In one embodiment, the peptide linker (2) may be a peptide linker consisting of the amino acid sequence of SEQ ID NO: 7.


The immunoglobulin Fc region refers to a protein that comprises immunoglobulin heavy chain constant region 2 (CH2) and heavy chain constant region 3 (CH3), but does not comprise immunoglobulin heavy and light chain variable regions and light chain constant region (CL). The immunoglobulin Fc region may be derived from IgG, IgA, IgE, IgD, or IgM. Specifically, the immunoglobulin Fc region may be derived from IgG1, IgG2, IgG3, or IgG4, which is a subclass of IgG. Preferably, the immunoglobulin Fc region may be derived from IgG4.

In addition, the immunoglobulin Fc region may be an Fc region variant as well as a wild-type Fc region. In addition, as used in the present specification, the term "Fc region variant" may refer to a form which is different from the wild-type Fc region in terms of glycosylation pattern, or has a high glycosylation compared to the wild-type Fc region, or has a low glycosylation compared to the wild-type Fc region, or is deglycosylated. In addition, an aglycosylated Fc region is also included. The Fc region or a variant thereof may be adapted to have an adjusted number of sialic acids, fucosylations, or glycosylations, through culture conditions or genetic manipulation of a host.

In addition, glycosylation of the immunoglobulin Fc region may be modified by conventional methods such as chemical methods, enzymatic methods, and genetic engineering methods using microorganisms. In addition, the Fc region variant may be in a mixed form of respective Fc regions of immunoglobulins, IgG, IgA, IgE, IgD, and IgM. In addition, the Fc region variant may be in a form in which some amino acids of the Fc region are substituted with other amino acids. In one embodiment, the Fc region may comprise the amino acid sequence of SEQ ID NO: 6.

In addition, the fusion protein of the present invention may be a polypeptide comprising the heavy chain variable region (VH) and heavy chain constant region 1 (CH1) of an anti-CD73 antibody, an Fc region, and IL-2 or a variant thereof. Here, the Fc region and the IL-2 or variant thereof may be linked by a linker.

In addition, the fusion protein may comprise the following structural formulas (II) and (III):



and



in structural formulas (II) and (III) above,

12

N' is the N-terminus,

C' is the C-terminus,

X is an antibody or an antigen-binding fragment thereof that specifically binds to CD73, and comprises a heavy chain variable region (VH) and a heavy chain constant region 1 (CH1),

X' is an antibody or an antigen-binding fragment thereof that specifically binds to CD73, and comprises a light chain variable region (VL) and a light chain constant region (CL),

Y is IL-2,

linker (3) and linker (4) are peptide linkers, and

q and r are each independently 0 or 1.

In one embodiment, the antibody or antigen-binding fragment thereof may comprise: a heavy chain variable region comprising an HCDR1 comprising the amino acid sequence of SEQ ID NO: 24, an HCDR2 comprising the amino acid sequence of SEQ ID NO: 25, and an HCDR3 comprising the amino acid sequence of SEQ ID NO: 26; and a light chain variable region comprising an LCDR1 comprising the amino acid sequence of SEQ ID NO: 27, an LCDR2 comprising the amino acid sequence (Asp-Ala-Ser (DAS)) of SEQ ID NO: 28, and an LCDR3 comprising the amino acid sequence of SEQ ID NO: 29.

In one embodiment, the heavy chain variable region may comprise the amino acid sequence of SEQ ID NO: 9. In one embodiment, the light chain variable region may comprise the amino acid sequence of SEQ ID NO: 13.

The peptide linker (3) may consist of 1 to 50 contiguous amino acids, or 3 to 30 contiguous amino acids, or 5 to 15 amino acids. In one embodiment, the peptide linker 3 may consist of 12 amino acids. In addition, the peptide linker (3) may contain at least one cysteine. Specifically, it may contain one, two, or three cysteines. In addition, the peptide linker (3) may be derived from the hinge of an immunoglobulin. For example, the hinge may be selected from the hinge regions of various IgG subclass antibodies. In addition, the hinge may be a form in which one or more of the amino acids in the hinge region derived from an immunoglobulin are substituted with other amino acid(s), or may be a sequence obtained by adding one or more amino acids to the hinge region. In one embodiment, the peptide linker (3) may be a peptide linker consisting of the amino acid sequence of SEQ ID NO: 11.

The peptide linker (4) may consist of 1 to 30 contiguous amino acids, or 2 to 20 contiguous amino acids, or 2 to 10 amino acids. In one embodiment, the peptide linker (4) may be (G4S)_n (where n is an integer ranging from 1 to 10). In (G4S)_n, n may be 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10. In one embodiment, the peptide linker (4) may be a peptide linker consisting of the amino acid sequence of SEQ ID NO: 12.

The "Fc region fragment or variant thereof" used in this embodiment is as described above.

Fusion Protein Dimer

Another aspect of the present invention provides a dimer obtained by linking two fusion proteins, each comprising: an antibody or an antigen-binding fragment thereof that specifically binds to CD73; and IL-2 or a variant thereof. The antibody or antigen-binding fragment thereof that specifically binds to CD73 is as described above, and IL-2 is also as described above. Here, the linkage between the fusion proteins constituting the dimer may be achieved by a disulfide bond formed by cysteines present in the linker, without being limited thereto. The fusion proteins constituting the dimer may be the same proteins, resulting in a homodimer.

Specifically, in one embodiment, the dimer may be a dimer in which two fusion proteins, each comprising a single-domain antibody that specifically binds to CD73, an Fc region, and IL-2 or a variant thereof, are linked to each other by a disulfide bond formed by cysteines. In addition, in one embodiment, the dimer may be a dimer in which two fusion proteins, each comprising a polypeptide comprising the heavy chain variable region (VH) and heavy chain constant region 1 (CH1) of an anti-CD73 antibody, an Fc region, and IL-2 or a variant thereof, and a polypeptide comprising the light chain variable region (VL) and light chain constant region (CL) of the anti-CD73 antibody, are linked to each other by a disulfide bond formed by cysteines. Polynucleotide Encoding Fusion Protein

Another aspect of the present invention provides a polynucleotide encoding a fusion protein comprising: an antibody or an antigen-binding fragment thereof that specifically binds to CD73; and IL-2 or a variant thereof. The antibody or antigen-binding fragment thereof that specifically binds to CD73 is as described above, and the IL-2 or variant thereof is also as described above.

Specifically, the polynucleotide encoding the fusion protein comprising the single-domain antibody that specifically binds to CD73, the Fc region, and the IL-2 or variant thereof may comprise any one nucleotide sequence selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 33, and SEQ ID NO: 35.

In addition, the polypeptide comprising the heavy chain variable region (VH) and heavy chain constant region 1 (CH1) of the anti-CD73 antibody, the Fc region, and the IL-2 or variant thereof, and the polypeptide comprising the light chain variable region (VL) and light chain constant region (CL) of the anti-CD73 antibody may be encoded by the nucleotide sequence of SEQ ID NO: 37 and the nucleotide sequence of SEQ ID NO: 39, respectively.

In addition, if the polynucleotide encodes the same polypeptide, one or more nucleotides may be altered by substitution, deletion, insertion, or a combination thereof. When a polynucleotide sequence is produced by chemical synthesis, synthetic methods well known in the art may be used, such as those described in Engels and Uhlmann, *Angew Chem Int Ed Eng.*, 37:73-127, 1988. Such methods include, for example, triester, phosphite, phosphoramidite and H-phosphate methods, PCR and other autoprimer methods, oligonucleotide syntheses on solid supports, and the like.

According to one embodiment, the polynucleotide may comprise a nucleotide sequence having an identity of at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 86%, at least about 87%, at least about 88%, at least about 89%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or at least about 100% to the nucleotide sequence of each of SEQ ID NO: 31, SEQ ID NO: 33, SEQ ID NO: 35, SEQ ID NO: 37, and SEQ ID NO: 39.

The polynucleotide may further comprise a signal sequence or a leader sequence. As used herein, the term "signal sequence" refers to a nucleic acid encoding a signal peptide that directs secretion of a target protein. The signal peptide is translated and then cleaved in a host cell. Specifically, the signal sequence of the present invention is a polynucleotide encoding an amino acid sequence that initiates migration of a protein across the endoplasmic reticulum (ER) membrane.

Signal sequences are well known in the art for their characteristics. Such signal sequences typically comprise 16

to 30 amino acid residues, but may comprise more or fewer amino acid residues than such amino acid residues. A typical signal peptide is composed of three regions, that is, a basic N-terminal region, a central hydrophobic region, and a more polar C-terminal region. The central hydrophobic region comprises 4 to 12 hydrophobic residues that cause the signal sequence to be immobilized during migration of an immature polypeptide through the membrane lipid bilayer.

After initiation, signal sequences are cleaved in the lumen of ER by cellular enzymes, commonly known as signal peptidases. Here, the signal sequence may be a secretory signal sequence of tPa (tissue plasminogen activator), HSV gDs (signal sequence of Herpes simplex virus glycoprotein D), an IgG signal sequence, or a growth hormone. Preferably, a secretory signal sequence used in higher eukaryotic cells including mammals and the like may be used. Signal sequences useful in the present invention include antibody light chain signal sequences, e.g., antibody 14.18 (Gillies et al., *J. Immunol. Meth.* 125:191-202 (1989)), antibody heavy chain signal sequences, e.g., the MOPC141 antibody heavy chain signal sequence (Sakano et al., *Nature* 286:676-683), and other signal sequences known in the art (see, e.g., Watson et al., *Nucleic Acid Research*, 1984, 12:5145-5164). In one embodiment, the signal sequence may comprise the amino acid sequence of SEQ ID NO: 1.

Vector Containing Polynucleotide

Another aspect of the present invention provides an expression vector containing a polynucleotide encoding a fusion protein comprising: an antibody or an antigen-binding fragment thereof that specifically binds to CD73; and IL-2 or a variant thereof. Here, the antibody or antigen-binding fragment thereof that specifically binds to CD73, the antigen-binding site, and the IL-2 and variant thereof are as described above.

Specifically, the polynucleotide encoding the fusion protein comprising the anti-CD73 single-domain antibody and the IL-2 or variant thereof may comprise any one nucleotide sequence selected from the group consisting of SEQ ID NO: 30, SEQ ID NO: 32, and SEQ ID NO: 34.

In addition, the polynucleotide encoding the fusion protein comprising the anti-CD73 antibody and the IL-2 or variant thereof may comprise the nucleotide sequence of SEQ ID NO: 37, which encodes the heavy chain region of the anti-CD73 antibody, and the nucleotide sequence of SEQ ID NO: 39, which encodes the light chain region of the anti-CD73 antibody. In addition, in one embodiment, when each of the nucleotide sequences contains a signal sequence, the sequence encoding the heavy chain region may be the nucleotide sequence of SEQ ID NO: 36, and the sequence encoding the light chain region may be the nucleotide sequence of SEQ ID NO: 38. In this case, the polynucleotide may be inserted into an expression vector, and as an example, it may be inserted into a bicistronic expression vector.

As used in the present specification, the term "vector" refers to a vector that may be introduced into a host cell and recombined and inserted into the genome of the host cell. Alternatively, the vector is understood as a nucleic acid vehicle containing a nucleotide sequence that is capable of self-replication as an episome. Examples of the vector include linear nucleic acids, plasmids, phagemids, cosmids, RNA vectors, viral vectors, mini-chromosomes, and analogs thereof. Examples of the viral vector include, but are not limited to, retroviruses, adenoviruses, and adeno-associated viruses.

Specifically, the vector may include plasmid DNA, phage DNA, or the like, and examples thereof include commer-

cially developed plasmids (pUC18, pBAD, pIDTSAMRT-AMP, and the like), *E. coli*-derived plasmids (pYG601BR322, pBR325, pUC118, pUC119, and the like), *Bacillus subtilis*-derived plasmids (pUB110, pTP5, and the like), yeast-derived plasmids (YEpl3, YEpl24, YCp50, and the like), phage DNA (Charon4A, Charon21A, EMBL3, EMBL4, λ gt10, λ gt11, A ZAP, and the like), animal viral vectors (retroviruses, adenoviruses, vaccinia viruses, and the like), insect viral vectors (baculoviruses and the like). Since the vector exhibits different expression levels and modification of a protein depending on a host cell, it is preferred to select and use a host cell which is most suitable for the purpose.

In addition, the plasmid may contain a selection marker such as an antibiotic resistance gene, and host cells maintaining the plasmid may be cultured under selective conditions.

As used in the present specification, the term "gene expression" or "expression" of a target protein is understood to mean transcription of DNA sequences, translation of mRNA transcripts, and secretion of fusion protein products or fragments thereof. A useful expression vector may be RcCMV (Invitrogen, Carlsbad) or a variant thereof. The expression vector may contain a human cytomegalovirus (CMV) promoter for promoting continuous transcription of a target gene in mammalian cells, and a bovine growth hormone polyadenylation signal sequence for increasing the stability level of RNA after transcription.

Transformed Cell

Another aspect of the present invention provides a transformed cell into which an expression vector containing a polynucleotide encoding a fusion protein comprising: an antibody or an antigen-binding fragment thereof that specifically binds to CD73; and IL-2 or a variant thereof, has been introduced. Here, the expression vector containing the polynucleotide is as described above.

As used in the present specification, the term "transformed cell" refers to prokaryotic cells and eukaryotic cells into which a recombinant expression vector has been introduced. The transformed cell may be produced by transforming a host cell by introduction of the vector. In addition, the fusion protein of the present invention may be produced by expressing the polynucleotide contained in the vector.

The transformation may be performed by various methods. The transformation method is not particularly limited as long as it can produce the fusion protein of the present invention. Specifically, examples of the transformation method include CaCl_2 precipitation, Hanahan method whose efficiency has been increased by using a reducing agent such as dimethyl sulfoxide (DMSO) in CaCl_2 precipitation, electroporation, calcium phosphate precipitation, protoplast fusion, agitation using silicon carbide fiber, *Agrobacterium*-mediated transformation, transformation using PEG, dextran sulfate-, lipofectamine-, and desiccation/inhibition transformation, and the like. In addition, the target can be delivered into cells using virus particles by means of infection. In addition, the vector may be introduced into the host cell by gene bombardment or the like.

In addition, the host cell used to produce the transformed cell is also not particularly limited as long as it can produce the antibody of the present invention. Specifically, examples of the host cell include, but are not limited to, prokaryotic cells, eukaryotic cells, and cells of mammalian, plant, insect, fungal, or bacterial origin. As an example of the prokaryotic cells, *E. coli* may be used. In addition, as an example of the eukaryotic cells, yeast may be used. In addition, as the mammalian cells, CHO cells, F2N cells, COS cells, BHK

cells, Bowes melanoma cells, HeLa cells, 911 cells, AT1080 cells, A549 cells, SP2/0 cells, human lymphoblastoids, NSO cells, HT-1080 cells, PERC.6 cells, HEK 293 cells, HEK293T cells, or the like may be used, without being limited thereto. In addition, any cells which are known to those skilled in the art to be usable as mammalian host cells may be used.

In addition, for optimization of properties of an antibody as a therapeutic agent or for any other purpose, glycosylation pattern of the antibody (for example, sialic acids, fucosylations, glycosylations) may be adjusted by manipulating, through methods known to those skilled in the art, glycosylation-related genes possessed by host cells.

Method for Producing Fusion Protein Dimer

Another aspect of the present invention provides a method for producing a fusion protein dimer comprising: an antibody or an antigen-binding fragment thereof that specifically binds to CD73; and IL-2 or a variant thereof. Here, the fusion protein and the fusion protein dimer are as described above.

Specifically, the method for producing the fusion protein dimer may comprise steps of: i) culturing the transformed cell; and ii) collecting the fusion protein dimer.

As used in the present specification, the term "culturing" refers to a method of growing microorganisms under appropriately artificially controlled environmental conditions.

Culturing the transformed cell may be carried out using methods well known in the art. Specifically, the culturing method is not particularly limited as long as it can express and produce the antibody of the present invention. Specifically, the culturing may be carried out in a batch process, or carried out continuously in a fed batch or repeated fed batch process.

In addition, the step of collecting the antibody from the culture may be performed by methods known in the art. Specifically, the collecting method is not particularly limited as long as the produced fusion protein of the present invention may be collected. Preferably, the collecting method may be a method such as centrifugation, filtration, extraction, spraying, drying, evaporation, precipitation, crystallization, electrophoresis, fractional dissolution (e.g., ammonium sulfate precipitation), chromatography (e.g., ion exchange, affinity, hydrophobicity, and size exclusion), or the like.

Use of Fusion Protein (Dimer)

Another aspect of the present invention provides a pharmaceutical composition for preventing or treating cancer containing, as an active ingredient, a fusion protein or fusion protein dimer comprising: an antibody or an antigen-binding fragment thereof that specifically binds to CD73; and IL-2 or a variant thereof. The fusion protein and the fusion protein dimer are as described above.

Specifically, the present invention provides a pharmaceutical composition for preventing or treating cancer containing, as an active ingredient, a fusion protein comprising an anti-CD73 single-domain antibody and IL-2 or a variant thereof, or a dimer of the fusion proteins.

The present invention also provides a pharmaceutical composition for preventing or treating cancer containing, as an active ingredient, a fusion protein comprising an anti-CD73 antibody and IL-2 or a variant thereof, or a dimer of the fusion proteins.

As used in the present specification, the term "cancer" is classified as a disease in which normal tissue cells proliferate indefinitely for some reason and continue to grow rapidly regardless of the living phenomenon of the living organism or the surrounding tissue condition. The cancer in the

present invention may be, but is not limited to, any one cancer selected from the group consisting of various human cancers, for example, gastric cancer, liver cancer, lung cancer, colorectal cancer, breast cancer, prostate cancer, ovarian cancer, pancreatic cancer, cervical cancer, thyroid cancer, laryngeal cancer, acute myeloid leukemia, brain tumor, neuroblastoma, retinoblastoma, head and neck cancer, salivary gland cancer, and lymphoma.

Although a preferred dose of the pharmaceutical composition varies depending on the patient's condition and body weight, the disease's severity, the form of drug, and the route and duration of administration, it may be appropriately selected by those skilled in the art. In the pharmaceutical composition for treating or preventing cancer according to the present invention, the fusion protein may be contained in any amount (effective amount) depending on the application, dosage form, and blending purpose thereof, as long as the fusion protein can exhibit anticancer activity or, in particular, a therapeutic effect on cancer. A conventional effective amount of the fusion protein will be determined within a range of 0.001 wt % to 20.0 wt % based on the total weight of the composition. As used herein, the "effective amount" refers to an amount of an active ingredient that can induce the effect of alleviating or treating the condition of a disease, in particular the effect of alleviating or treating the condition of cancer. Such an effective amount can be experimentally determined within the scope of common knowledge of those skilled in the art.

As used herein, the term "treatment" may be used to mean both therapeutic treatment and prophylactic treatment, and covers any application or administration for treating a disease in a mammal, including a human. In addition, the term includes inhibiting or slowing progression of the disease; partially or fully relieving the disease by restoring or repairing a lost, missing, or defective function; stimulating an inefficient process; or alleviating a serious disease. The "prevention" may be used to mean alleviating or reducing the pathological condition or disease of an individual.

Pharmacokinetic parameters such as bioavailability and underlying parameters such as clearance rate may also impact efficacy. Thus, an "enhanced efficacy" (e.g., an improvement in efficacy) may be due to improved pharmacokinetic parameters as well as improved potency, and may be measured by comparing clearance rates in test animals or in human subjects, as well as parameters such as cancer disease treatment or alleviation.

Meanwhile, the pharmaceutical composition of the present invention is administered in a "therapeutically effective amount".

As used in the present specification, the term "administration" means introducing a given substance into a subject by any suitable method. The composition may be administered through any general routes as long as it can reach the target tissue. The composition may be administered intraperitoneally, intravenously, intramuscularly, subcutaneously, intradermally, orally, topically, intranasally, intrapulmonary, or intrarectally, without being limited thereto.

As used in the present specification, the term "therapeutically effective amount" or "pharmaceutically effective amount" refers to an amount of a compound or composition effective to prevent or treat the disease in question, which is sufficient to treat the disease at a reasonable benefit/risk ratio applicable to medical treatment and does not cause adverse effects. A level of the effective amount may be determined depending on factors including the patient's health condition, the type and severity of disease, the activity of drug, sensitivity to drug, the mode of administration, the time of administration, the route of administration, excretion rate, the duration of treatment, formulation or drugs used in combination with the composition, and other factors well

known in the medical field. In one embodiment, the therapeutically effective amount means an amount of drug effective to treat cancer.

Here, the pharmaceutical composition may further contain a pharmaceutically acceptable carrier. The pharmaceutically acceptable carrier may be any carrier that is a non-toxic substance suitable for delivery to a patient. Distilled water, alcohol, fat, wax, and inert solid may be contained as the carrier. A pharmaceutically acceptable adjuvant (buffer, dispersant) may also be contained in the pharmaceutical composition.

Specifically, by containing a pharmaceutically acceptable carrier, the pharmaceutical composition may be prepared into a parenteral formulation depending on its route of administration using conventional methods known in the art. Here, the term "pharmaceutically acceptable" means that the carrier does not have more toxicity than the subject to whom/which the pharmaceutical composition is to be applied (prescribed) can adapt while not inhibiting activity of the active ingredient.

When the pharmaceutical composition is prepared into a parenteral formulation, it may be made into preparations in the form of injections, transdermal patches, nasal inhalants, or suppositories with suitable carriers according to methods known in the art. For formulation into injectable formulations, sterile water, ethanol, polyol such as glycerol or propylene glycol, or a mixture thereof may be used as a suitable carrier; and an isotonic solution, such as Ringer's solution, phosphate buffered saline (PBS) containing triethanol amine, sterile water for injection, or 5% dextrose may preferably be used. Formulation of pharmaceutical compositions is known in the art, and reference may specifically be made to Remington's Pharmaceutical Sciences (19th ed., 1995), and the like. This document is considered part of the present specification.

A preferred dose of the pharmaceutical composition may range from 0.0001 µg/kg to 100 g/kg per day, depending on the patient's condition, body weight, sex, age, severity of the patient, and route of administration. The dose may be administered once or several times a day. Such a dose should not be construed as limiting the scope of the present invention in any way.

Subjects to whom/which the pharmaceutical composition may be applied (prescribed) are mammals and humans, with humans being particularly preferred. The pharmaceutical composition of the present invention may further contain any compound or natural extract known to have a cancer therapeutic effect.

Another aspect of the present invention provides the use of a fusion protein or fusion protein dimer comprising: an antibody or an antigen-binding fragment thereof that specifically binds to CD73; and IL-2 or a variant thereof, for manufacture of a medicament for preventing or treating cancer.

Another aspect of the present invention provides a method for preventing or treating cancer, comprising a step of administering to a subject a fusion protein or fusion protein dimer comprising: an antibody or an antigen-binding fragment thereof that specifically binds to CD73; and IL-2 or a variant thereof. The subject may be a mammal, preferably a human. In addition, the subject may be a patient suffering from cancer, or a subject who is highly likely to suffer from cancer.

The route of administration, dose, and frequency of administration of the fusion protein may vary depending on the patient's condition and the presence or absence of side effects, and the fusion protein may be administered to the subject in various ways and amounts. The optimal administration method, dosage, and frequency of administration of the fusion protein may be selected within appropriate ranges by a person skilled in the art. A preferred dose of the fusion

protein comprising: the antigen binding site that specifically binds to CD73; and IL-2 or a variant thereof, may range from 0.0001 µg/kg to 100 g/kg per day depending on the patient's condition, weight, sex, age, severity of the patient, and the route of administration. The dose may be administered once or several times a day. Such a dose should not be construed as limiting the scope of the present invention in any way.

MODE FOR INVENTION

Hereinafter, the present invention will be described in more detail by way of examples. These examples are only for illustrating the present invention in more detail, and it will be apparent to those skilled in the art that the scope of the present invention according to the subject matter of the present invention is not limited by these examples.

I. Production of Fusion Proteins and Dimers Thereof

Example 1. Production of GI-108A1: Anti-CD73 sdAb-hIgG4 Fc-hIL2v3

To produce a fusion protein comprising: an anti-CD73 single domain antibody (AHF10235, AHF10240, or AHP04167) that specifically binds to CD73; and an IL-2 variant, a polynucleotide (SEQ ID NO: 30, 32 or 34) encoding a signal peptide (SEQ ID NO: 1), an anti-CD73 single domain antibody (SEQ ID NO: 2, 3 or 4), linker (1) (SEQ ID NO: 5), an IgG4 Fc region (SEQ ID NO: 6), linker (2) (SEQ ID NO: 7), and an IL-2 variant (SEQ ID NO: 8) with three amino acid substitutions was cloned into a pCGS3 vector (Sigma-Aldrich®) using a Bioxp™ 3250 SYSTEM (Tables 1 to 3).

TABLE 1

	Amino acid sequence	SEQ ID NO.
mIgG signal	MEWSWVFLFFLSVTTGVHS	1
Anti-hCD73 sdAb (AHF10235)	QVQVQESGGGLVQAGDSLRLSCAASGRSISRYNMGWFRQA PGKEREFVAAISRSGGVITYADSVKGRLTISRDNKASAVYL QMNSLKPEDTAVYYCAADWRNSDSPSLIIVYDYWGQGTQ VTVSS	2
Linker (1)	GGGGS AESKYGPPCPPCP	5
hIgG4 Fc (F01)	APEAAGGPSVFLFPPKPKDQLMISRTPEVTCVVVDVSQEDPE VQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQD WLNQKEYKCKVSNKGLPSSIEKTIKAKGQPREPQVYTLPPS QEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTP PVLDSGDSFPLYSRLTVDKSRWQEGNVFSCSVLHEALHNHY TQKSLSLSLG	6
Linker (2)	GGGSGGGSGGGGS	7
hIL2v3	APTSSSTKKTQLQLEHLLLDLQMLNGINNYKNPKLTAMLT AKFYMPKKATELKHQLCLERELKPLEEVLNLAQSKNFHLRP RDLISNINIVIVLELKGSETTFMCEYADETATIVEFLNRWITFC QSIISTLT	8

40

TABLE 2

	Amino acid sequence	SEQ ID NO.
mIgG signal	MEWSWVFLFFLSVTTGVHS	1
Anti-hCD73 sdAb (AHF10240)	EVQLVESGGGVVQTGGSLRLSCAASGRFTFTHLAMGWFRQ APGKEREFVAAISNSGAGTQFADSVKGRFTISRDNKNTVY LQMNSLKPEDTAVYYCAADWGRTRDSPSKLNAVYDYWGR GTQVTVSS	3
Linker (1)	GGGGS AESKYGPPCPPCP	5
hIgG4 Fc (F01)	APEAAGGPSVFLFPPKPKDQLMISRTPEVTCVVVDVSQEDP EVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLH QDWLNQKEYKCKVSNKGLPSSIEKTIKAKGQPREPQVYTL PPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNY KTPPVLDSGDSFPLYSRLTVDKSRWQEGNVFSCSVLHEAL HNHYTQKSLSLSLG	6
Linker (2)	GGGSGGGSGGGGS	7
hIL2v3	APTSSSTKKTQLQLEHLLLDLQMLNGINNYKNPKLTAMLT AKFYMPKKATELKHQLCLERELKPLEEVLNLAQSKNFHLR PRDLISNINIVIVLELKGSETTFMCEYADETATIVEFLNRWITF CQSIISTLT	8

TABLE 3

	Amino acid sequence	SEQ ID NO.
mIgG signal	MEWSWVFLFFLSVTTGVHS	1
Anti-hCD73 sdAb (AHP04167)	EVQLVESGGGLVQAGGSLRLSCVVSGRFTFSNYHMGWFRQ APGKEREFVAVIGRSGGPTYADSVKGRFTISRDNKNTV YLMNSLKPEDTAVYYCAGDWRNSDSPSKLPVYDYWG QGTQVTVSS	4
Linker (1)	GGGS AESKYGPCCPCP	5
hIgG4 Fc (F01)	APEAAGGPSVFLFPPKPKDQLMISRTPEVTCVVVDVSQEDP EVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLH QDWLNGKEYKCKVSNKGLPSSIEKTIKAKGQPREPQVYT LPDSQSEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENN YKTPPVLDSGSEFFLYSRLTVDKSRWQEGNVFSCSVLHEA LHNHYTQKSLSLGLG	6
Linker (2)	GGGSGGGSGGGSGGS	7
hIL2v3	APTSSSTKKTQLQLEHLLDLQMLNGINNYKNPKLTAMLT AKFYMPKKATELKHLCLERELKPLEEVLNLAQSKNPHLR PRDLISNINVIVLELKGSETTFMCEYADETATIVEFLNRWITF CQSIISTLT	8

In addition, each of the vectors was introduced into CHO cells (Expi-CHO™, Thermo Fisher Scientific) to express the fusion proteins. After introducing the vectors, the cells were cultured under the conditions of 37° C., 125 rpm, and 8% CO₂. Thereafter, the cultures were harvested and the fusion protein dimers were purified therefrom. The three purified fusion protein dimers were named “GI-108A1 (AHF1-0235)”, “GI-108A1 (AHF10240)”, and “GI-108A1 (AHP04167)”, respectively. Specifically, each of the three fusion protein dimers was purified using chromatography with Protein A resin. After filtering the harvested cultures, each of the filtered cultures was loaded onto a column and flowed through the column. Next, the fusion protein dimers were collected using 50 mM glycine (pH 3.4).

A buffer containing each of the three collected fusion protein dimers (GI-108A1 (AHF10235), GI-108A1 (AHF10240), and GI-108A1 (AHP04167)) was exchanged with PBS (phosphate buffered saline, pH 7.4) through dialysis. Using a NanoDrop system (Thermo Fisher Scientific), it was identified that the fusion protein dimers were contained at concentrations of 3.86 mg/mL, 4.12 mg/mL, and 12.86 mg/mL, respectively. In addition, as a result of analyzing the purities through size exclusion chromatography (SEC) analysis, the purities of the three purified fusion protein dimers were 71.81%, 75.36%, and 93.87%, respectively (Table 4 and Table 5).

TABLE 4

	Peak Name	RT	Area	% Area	Height
GI-108A1 (AHF10235)					
1	Peak 1	12.067	24947	0.90	762
2	Peak 2	14.015	124244	4.50	2122
3	Peak 3	14.980	377981	13.68	6230
4	Monomer	17.022	1983970	71.81	48598
GI-108A1 (AHF10240)					
1	Peak 1	11.962	198392	6.85	7601
2	Peak 2	13.697	138199	4.77	2536
3	Peak 3	14.694	351133	12.12	6847
4	Monomer	16.739	2183777	75.36	60301
GI-108A1 (AHP04167)					

TABLE 4-continued

	Peak Name	RT	Area	% Area	Height
1	Peak 1	12.111	21365	0.41	331
2	Peak 2	14.769	295396	5.72	5156
3	Monomer	16.682	4849570	93.87	140556

TABLE 5

ID	Concentration after formulation (mg/mL)	Final protein (mg)	Purification yield* (μg/mL)	Final purity	Aggrega- tion
GI-108A1 (AHF10235)	3.86	14.66	146.57	71.81	X
GI-108A1 (AHF10240)	4.12	38.76	387.56	75.36	X
GI-108A1 (AHP04167)	12.86	41.14	411.39	93.87	X

*Purification Yield: Purification μg/culture mL, 100 mL Scale

Example 2. Production of GI-108B1: Anti-CD73 Ab-hIgG4 Fc-hIL2v3

To produce a fusion protein dimer comprising: an anti-CD73 antibody that specifically binds to CD73; and an IL-2 variant, a polynucleotide (SEQ ID NO: 36) encoding a signal peptide (SEQ ID NO: 1), an anti-CD73 antibody heavy-chain variable region (SEQ ID NO: 9) and constant region (SEQ ID NO: 10), linker (1) (SEQ ID NO: 11), an IgG4 Fc region (SEQ ID NO: 6), linker (2) (SEQ ID NO: 12), and an IL-2 variant (SEQ ID NO: 8) with three amino acid substitutions, and a polynucleotide (SEQ ID NO: 38) encoding a signal peptide (SEQ ID NO: 1) and an anti-CD73 antibody light-chain variable region (SEQ ID NO: 13) and constant region (SEQ ID NO: 14) were cloned into a pCGS3 vector (SIGMA-ALDRICH®) using a BIOXP™ 3250 SYS-TEM (Table 6).

In addition, the vector was introduced into CHO cells (EXPI-CHO™, Thermo Fisher Scientific) to express the fusion protein dimer. After introducing the vector, the cells were cultured under the conditions of 37° C., 125 rpm, and 8% CO₂. Thereafter, the culture was harvested and the fusion protein dimer was purified therefrom. The purified fusion protein dimer was named “GI-108B1”.

TABLE 6

		Amino acid sequence	SEQ ID NO.
Anti-hCD73	mIgG signal	MEWSWVFLFFLSVTTGVHS	1
HC-hIgG4Fc-hIL2v3	Anti-hCD73 VH (CSA0060)	QVQLVQSGAEVKKPGSSVKVSCKFSGGSFTSY SFSWVRQAPGQGLEWMGRIIPVLTITTDYAQKF RDRVTITADESTSTAYMELSGLRSEDTAVYYC ASGLKKNWYFDLWGRGALITVSS	9
	Anti-hCD73 CH1 (CSA0060)	ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYF PEPVTWNSGALTSGVHTFPAVLQSSGLYSLS SVVTVPSSSLGKTYTCNVDHKPSNTKVDKRV	10
	Linker (1)	ESKYGPCCPPCP	11
	hIgG4 Fc (F01)	APEAAGGPSVFLFPPKPKDQLMISRTPEVTCVV VDVSDPEVQPNWYVDGVEVHNAKTKPREE QFNSTYRVVSVLTVTLHQDWLNGKEYKCKVSN KGLPSSIEKTIKAKGQPREPQVYTLPPSQEEMT KNQVSLTCLVKGFYPSDIAVEWESNGQPENNY KTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFS CSVLHEALHNHYTQKSLSLGL	6
	Linker (2)	GGGGSGGGGS	12
	hIL2v3	APTSSSTKKTQLQLEHLLDLQMLNGINNYKN PKLTAMLTAKFYMPKKATELKHQLCLERELKP LEEVLNLAQSKNHLRPRDLISINIVIVLELKGS ETTFMCEYADETATVEFLNRWITFCQSIISTLT	8
Anti-hCD73	mIgG signal	MEWSWVFLFFLSVTTGVHS	1
LC (kappa)	Anti-hCD73 VL (CSA0060)	DIQMTQSPSFLSASVGDRVTITCQASQDISHYLN WYQQKPGKAPKLLIYDASSLETGVPSPRSGSGS GTSFTLTISSLQPEDFATYYCQQYDDFPLTFGG GTKVDIK	13
	Anti-hCD73 kappa CL (CSA0060)	RSVAAPSVFIPPSPDEQLKSGTASVVCLLNNFYP REAKVQWKVDNALQSGNSQESVTEQDSKDST YSLSSITLTLKADYEKHKVYACEVTHQGLSSP VTKSFNRGEC	14
	Anti-hCD73 VL-LCDR2 (CSA0060)	DAS	28

Specifically, the GI-108B1 was purified using chromatography with Protein A resin. After filtering the harvested culture, the filtered culture was loaded onto a column and flowed through the column. Next, the fusion protein dimer was collected using 50 mM glycine (pH 3.4).

A buffer containing the collected fusion protein dimer was exchanged with PBS (pH 7.4) through dialysis. Using a NanoDrop system (Thermo Fisher Scientific), it was identified that the fusion protein dimer was contained at a concentration of 4.55 mg/mL. In addition, as a result of analyzing the purity through size exclusion chromatography (SEC) analysis, it was confirmed that the purity of the purified fusion protein dimer was and 83.63% (Table 7 and Table 8).

TABLE 7

GI-108B1	Peak Name	RT	Area	% Area	Height
1	Peak 1	11.969	340915	11.45	6998
2	Peak 2	13.806	121831	4.09	1863
3	Monomer	15.656	2489084	83.63	57175
4	Peak 3	17.726	24509	0.82	547

TABLE 8

ID	Concentration after formulation (mg/mL)	Final protein (mg)	Purification yield* (μg/mL)	Final purity	Aggregation
GI-108B1	4.55	27.28	272.82	83.63	X

*Purification Yield: Purification μg/culture mL, 100 mL Scale

Example 3. Production of Anti-CD73 Antibody: GI-αCD73

To produce an anti-CD73 antibody, a polynucleotide (SEQ ID NO: 42) encoding a signal peptide (SEQ ID NO: 1), an anti-CD73 antibody heavy chain variable region (SEQ ID NO: 9) and constant region (SEQ ID NO: 10), linker (1) (SEQ ID NO: 11), and an IgG4 Fc region (SEQ ID NO: 6), and a polynucleotide (SEQ ID NO: 38) encoding a signal peptide (SEQ ID NO: 1), and an anti-CD73 antibody light chain variable region (SEQ ID NO: 13) and constant region (SEQ ID NO: 14) were cloned into a pCGS3 vector (SIGMA-ALDRICH®) using a BIOXP™ 3250 SYSTEM.

In addition, the vector was introduced into CHO cells (EXPI-CHO™, Thermo Fisher Scientific) to express the control antibody anti-CD73 antibody. After introducing the vector, the cells were cultured under conditions of 37° C., 125 rpm, and 8% CO₂. The culture was harvested and the control anti-CD73 antibody was purified therefrom. Purification was performed in the same manner as in Example 1.

Example 4. Production of IgG4 Fc-IL-2v

A polynucleotide (SEQ ID NO: 43) encoding a signal peptide (SEQ ID NO: 1), linker (1) (SEQ ID NO: 11), an IgG4 Fc region (SEQ ID NO: 6), linker (2) (SEQ ID NO: 12), and an IL-2 variant (SEQ ID NO: 8) with three amino acid substitutions was cloned into a pCGS3 vector (SIGMA-ALDRICH®) using a BIOXP™ 3250 SYSTEM.

In addition, the vector was introduced into CHO cells (EXPI-CHO™, Thermo Fisher Scientific) to express the fusion protein. After introducing the vector, the cells were cultured for 7 days under conditions of 37° C., 125 rpm, 5%

25

CO₂, and 80% humidity. The culture was then harvested and the IgG4 Fc-IL-2v fusion protein was purified therefrom.
II. Characterization of Fusion Protein Dimers

Example 5. Measurement of Binding Affinity of
Each Fragment of GI-108 to Target Protein

To quantitatively analyze the binding affinities to CD73 and IL-2 receptors of GI-108A1 (AHF04167), GI-108A1 (AHP04167) and GI-108B1, in which the anti-CD73 antibody that specifically binds to CD73 was linked to the IL-2 variant, and Aldesleukin (trade name: PROLEUKIN®, Novartis), the binding affinities were measured using SPR (surface plasmon resonance). Using a BIACORE™ T200 instrument (Cytiva), the binding affinities of GI-108A1 (AHF04167), GI-108A1 (AHP04167) and GI-108B1 to hCD73, cyCD73, mCD73, hIL-2Rα, hIL-2Rβ, and hIL-2Rβγ were analyzed. Oleclumab (Astrazeneca), produced on order from Ybiologics, Inc., was used as a control.

Specifically, to analyze the binding affinities to hCD73, cyCD73, and mCD73, each of GI-108A1 (AHF04167), GI-108A1 (AHP04167), GI-108B1, and Oleclumab was diluted in HBS-EP+ buffer (Cytiva, USA) and reversibly immobilized on Sensor Chip CM5 (Cytiva, USA) using a human antibody capture kit (Cytiva, USA). CD73 was diluted in HBS-EP+ buffer and then analyzed for association and dissociation at a flow rate of 30 μL/min for 5 minutes each.

To analyze the binding affinity to hIL-2Rα, hIL-2Rα was irreversibly immobilized on Sensor Chip CM5 using a His capture kit (Cytiva, USA). Each of GI-108A1 (AHF04167), GI-108A1 (AHP04167), GI-108B1 and PROLEUKIN®, diluted in HBS-EP+ buffer, was then applied for 1 minute, followed by analysis of association and dissociation. In addition, the binding affinities to hIL-2Rβ and hIL-2Rβγ were measured by a three-step analysis method including treating Sensor Chip SA (Cytiva, USA) with biotinylated CD73, and then binding each of GI-108A1 (AHF04167), GI-108A1 (AHP04167), GI-108B1 and PROLEUKIN® to CD73, followed by treatment with each of hIL-2Rβ and hIL-2Rβγ diluted in HBS-P+ buffer (Cytiva, USA). For hIL-2Rβ, association was performed for 1 minute and dissociation for 1 minute, and for hIL-2Rβγ, association was performed for 5 minutes and dissociation for 2 hours at each concentration, in a single cycle kinetics analysis. Each sensorgram was normalized and subtracted compared to a blank cell to calculate affinity.

Table 9 below shows the results of measuring the binding affinities of GI-108A1 (AHF04167), GI-108A1 (AHP04167) and GI-108B1 to CD73 and IL-2 receptors using the BIACORE™ T200 instrument (Cytiva).

TABLE 9

K _D (M)	hCD73	cyCD73	mCD73	hIL-2Rα	hIL-2Rβ	hIL-2Rβγ
GI108B1	8.941E-11	1.13E-10	N.B.	N.B.	2.29E-06	2.00E-10
GI108 A1 (AHF04167)	5.02E-11	8.41E-11	2.62E-09	N.B.	2.50E-06	4.17E-10
GI108 A1 (AHP04167)	8.80E-11	7.08E-11	3.45E-09	N.B.	2.07E-06	3.27E-10
Oleclumab	1.01E-10	3.49E-10	N/A	N/A	N/A	N/A
PROLEUKIN®	N/A	N/A	N/A	4.96E-08	2.08E-06	1.00E-11

N.B. : not binding, N/A: not applicable

Example 6. Evaluation of the Ability of IL-2
Variant of GI-108 to Induce JAK-STAT Pathway
Activation

This experiment was conducted to evaluate the activity of the IL-2 variant region of GI-108. Specifically, the experi-

26

ment was conducted using HEK-BLUE™ IL-2 reporter cells (InvivoGen Inc.) expressing IL-2Rβγ. HEK-BLUE™ IL-2 reporter cells are induced to produce the reporter protein SEAP (secreted embryonic alkaline phosphatase) protein when the JAK-STAT pathway is activated by IL-2. In this experiment, the JAK-STAT pathway activation by IL-2 was evaluated by detecting the SEAP protein.

HEK-Blue™ IL-2 reporter cells were cultured in a DMEM medium (GIBCO™) containing 10% FBS (GIBCO™), 100 U/mL penicillin (Welgene Inc.), 100 μg/mL streptomycin (Welgene Inc.), and 100 μg/mL NOR-MOCIN™ (cat. ant-nr-1, InvivoGen Inc.). The HEK-BLUE™ IL-2 reporter cells were stabilized by subculture, and then harvested using trypsin (GIBCO™). Next, dead cells were removed by washing with PBS. The separated cells were suspended in a culture medium at about 2.8×10⁵ cells/mL.

GI-108A1 (AHF04167), GI-108A1 (AHP04167), GI-108B1, and PROLEUKIN® as a control group were each diluted with PBS and dispensed into a 96-well plate (cat. 30096, SPL) at 20 μL per well. 180 μL of the prepared cell suspension was added to each well of the treated 96-well plate at a density of about 3×10⁴ cells/well, and the cells were cultured in an incubator at 37° C. under 5% CO₂ for 24 hours.

After 24 hours, the 96-well plate was taken out of the incubator and centrifuged at 300×g for 5 minutes. Then, 20 μL of the supernatant was transferred to each well of a fresh 96-well plate. 180 μL of QUANTI-BLUE™ solution (cat. Rep-qbs, InvivoGen Inc.) dissolved at room temperature was dispensed into each well containing the supernatant. Each well was then incubated in an incubator for 30 minutes at 37° C. under 5% CO₂. After incubation, the absorbance was measured at a wavelength of 630 nm using a spectrophotometer (VERSAMAX™ Absorbance Microplate Reader). Data were analyzed using Graphpad prism 8.0 software.

As a result, the absorbance was increased by each of GI-108A1 (AHF04167), GI-108A1 (AHP04167), GI-108B1, and PROLEUKIN® in a concentration-dependent manner. Thereby, it was confirmed that the JAK-STAT signaling pathway was activated by the IL-2 variant region of both types of GI-108A1 and GI-108B1 (FIG. 3). The EC₅₀ values of GI-108A1 (AHF04167) and GI-108A1 (AHP04167) were 2.63 pM and 3.94 pM, respectively, and the EC₅₀ value of GI-108B1 was 1.28 pM. The EC₅₀ value of the positive control PROLEUKIN® was 10.3 pM.

Example 7. Evaluation of Cell Proliferation
Induced by IL-2 Variant of GI-108

The biological activity of the IL-2 variant of each of GI-108A1 and GI-108B was evaluated through a proliferation experiment using CTLL-2 cells expressing hIL-2Rαβγ.

CTLL-2 cells were cultured in complete RPMI 1640 medium (Thermo Fisher Scientific) supplemented with 0.2 mM L-glutamine, T-STIM™ culture supplement with ConA (concanavalin-A), 0.2 mM sodium pyruvate, and 10% FBS.

GI-108A1 (AHF04167), GI-108A1 (AHP04167), GI-108B1, and the control group PROLEUKIN® were each

27

diluted with PBS and dispensed in a 96-well plate (cat. 30096, SPL) at 50 μ L per well. The prepared CTLL-2 cells were added to each well of the drug-treated 96-well plate at a density of 1.2×10^5 cells per well. To analyze cell proliferation, 10 μ L of WST-1 was added to each well which was then incubated at 37° C. for 4 hours, followed by measurement of the absorbance.

The absorbances were measured at wavelengths of 450 nm and 690 nm (reference) using a spectrophotometer (VERSAMAX™ Absorbance Microplate Reader). CTLL-2 cell proliferation data were analyzed by subtracting the absorbance value at 690 nm from the absorbance value at 450 nm.

As a result, the absorbance was increased by each of GI-108A1 (AHF10240), GI-108A1 (AHP04167), GI-108B1, and PROLEUKIN® in a concentration-dependent manner. Thereby, it was confirmed that the proliferation of CTLL-2 cells was induced by the IL-2 variant region of both types of GI-108A1 and GI-108B1 (FIG. 4). The EC₅₀ values of GI-108A1 (AHF10240) and GI-108A1 (AHP04167) were 2.02 nM and 3.93 nM, respectively, and the EC₅₀ value of GI-108B1 was 1.70 nM. The EC₅₀ value of the positive control PROLEUKIN® was 0.08 nM.

The reason why the EC₅₀ value of PROLEUKIN® was lower than those of GI-108A1 and GI-108B1 was that the CTLL-2 cells expressed hIL-2R $\alpha\beta\gamma$. It was confirmed that as each of GI-108A1 and GI-108B1 do not bind to IL-2R α , the EC₅₀ value thereof was at least 20 times higher than that of PROLEUKIN®.

Example 8. Evaluation of the Inhibitory Ability of GI-108 to Inhibit Membrane-Bound CD73 Enzyme Activity

1×10^4 MDA-MB-231 cells (human breast cancer cells) were dispensed into each well of 96-well plates, and each of GI-108A1 (AHF10240), GI-108A1 (AHP04167), GI-108B1 and the control Oleclumab was diluted from 300 μ g/mL to a final concentration of 0.002 μ g/mL and added to the 96-well plates at 50 μ L per each well. Next, 100 μ L of 200 μ M AMP (SIGMA-ALDRICH®) was added to each well which was then incubated at 37° C. for 6 hours.

The 96-well plates were centrifuged, and the supernatants were transferred to new 96-well plates. Thereafter, the adenosine concentration of each of the supernatants was measured using an Adenosine Assay kit (Cell Biolabs, Inc) according to the manufacturer's protocol. IC₅₀ values were calculated using GraphPad Prism software.

As a result, the degree of luminescence was reduced by each of GI-108A1 (AHF10240), GI-108A1 (AHP04167), GI-108B1, and Oleclumab in a concentration-dependent manner. Thereby, it was confirmed that both types of GI-108A1 and GI-108B1 inhibited membrane-bound CD73 enzyme activity (FIG. 5). The IC₅₀ values of GI-108A1 (AHF10240) and GI-108A1 (AHP04167) were 2.75 nM and 5.58 nM, respectively, and the IC₅₀ value of GI-108B1 was 3.37 nM. The IC₅₀ value of the control Oleclumab was 1.47 nM. Although the IC₅₀ value of GI-108B1 was higher than those of GI-108A1 (AHF10240) and Oleclumab, GI-108B1 showed the highest potential in enzyme inhibition efficacy.

Example 9. Evaluation of the Ability of GI-108 to Inhibit Soluble CD73 Enzyme Activity

To evaluate the ability of GI-108 to inhibit soluble CD73 enzyme activity, an experiment was conducted using the AMP-Glo™ assay kit (Promega, V5012) according to the

28

manufacturer's manual. Specifically, each of GI-108A1 (AHF10240), GI-108A1 (AHP04167), GI-108B1 and the control Oleclumab was diluted to 0.05 ng/ml and added to each well of 96-well plates at 2.5 μ L per well, and 12.5 μ L of 10 μ M AMP was added to each well. Next, 2.5 ng/ml of recombinant human CD73 protein was added at 10 μ L per well, followed by incubation at room temperature for 10 minutes.

The reaction was terminated by treating each well with 25 μ L of AMP-Glo™ Reagent (Promega), and each well was shaken for 2 minutes and then incubated at room temperature for 1 hour. Each well was treated with AMP detection solution, shaken for 2 minutes, and then incubated at room temperature for 1 hour. The degree of luminescence was measured using the GLOMAX®-Multi Detection System (Promega), and IC₅₀ values were calculated using GraphPad Prism 8.0 software.

As a result, the degree of luminescence was reduced by each of GI-108A1 (AHF10240), GI-108A1 (AHP04167), and GI-108B1 in a concentration-dependent manner. Thereby, it was confirmed that both types of GI-108A1 and GI-108B1 inhibited soluble CD73 enzyme activity (FIG. 6). Meanwhile, a hook effect appeared, indicating that the higher the treatment concentration of Oleclumab, the lower the enzyme inhibition efficacy. The IC₅₀ values of GI-108A1 (AHF10240) and GI-108A1 (AHP04167) were 0.060 nM and 0.102 nM, respectively, and the IC₅₀ values of GI-108B1 and Oleclumab were 0.126 nM and 0.03 nM, respectively. Although the IC₅₀ value of GI-108B1 was higher than those of both types of GI-108A1 and Oleclumab, GI-108B1 showed the highest potential in enzyme inhibition efficacy.

Example 10. Evaluation of Reinvigoration of T Cells Inhibited by CD73-Adenosine with GI-108B1

To evaluate whether GI-108B1 reinvigorates CD8+ T cell activity suppressed by CD73-adenosine, human PBMCs were cultured in TEXMACS™ medium (Miltenyi Biotec) containing 5% human serum (SIGMA-ALDRICH®), 1% penicillin/streptomycin and 5 mM β -mercaptoethanol (Thermo Fisher Scientific) under humidified conditions of 37° C. and 5% CO₂ overnight. The next day, the cells were labeled with CTV and seeded in each well of 96-well plates at 1×10^6 cells/ml. The cells were stimulated using α CD3/CD28 dynabeads (1:1 ratio).

The cells were treated with vehicle (hIgG4), PROLEUKIN®, GI-108B1, anti-CD73 antibody (GI- α CD73), IgG4 Fc-IL-2v and Oleclumab alone, a combination of anti-CD73 antibody (GI- α CD73) and IgG4 Fc-IL-2v, or a combination of Oleclumab and IgG4 Fc-IL-2v. Each of the drugs was added to each well to a final concentration of 50 nM. Then, 500 μ M of AMP was added to each well, and the cells were cultured for 3 days under humidified conditions of 37° C. and 5% CO₂.

To evaluate the functional activation of T cells, the cells were re-stimulated with α CD3/CD28 dynabeads (1:1 ratio) and 50 nM or 500 μ M AMP in the presence of Brefeldin A under humidified conditions of 37° C. and 5% CO₂ for 4 hours. For FACS analysis of T cell proliferation, surface markers of the cells were stained using the following antibodies: anti-CD3-PerCP-Cy5.5 (Clone: UCHT1, BD Biosciences), anti-CD8-Alexa Fluor700 (Clone: RPA-T8, Biolegend Inc.), and Fixable Viability Dye eFluor 780 (Invitrogen Inc.). To stain intracellular IFN- γ , the cells were treated with fixation/permeabilization buffer and then stained with anti-IFN γ -APC antibody (Clone: 4S.B3, Biole-

gend Inc.). Data on the stained cells were obtained using a Symphony A3 instrument and analyzed using FlowJo software.

As a result, when treated with AMP and vehicle (hIgG4), the proliferation of CD8+ T cells was reduced. On the other hand, when treated with GI-108B1, the proliferation of CD8+ T cells significantly increased. In addition, it was confirmed that the proliferation of CD8+ T cells increased even when treated with anti-CD73 antibody alone or with a combination of anti-CD73 antibody and IgG4 Fc-IL-2v. Meanwhile, when treated with PROLEUKIN®, Oleclumab or IgG4 Fc-IL-2v alone or when treated with a combination of Oleclumab and IgG4 Fc-IL-2v, the proliferation of CD8+ T cells did not tend to increase (FIG. 7). In addition, when treated with AMP and vehicle (hIgG4), the production of IFN- γ by CD8+ T cells was reduced. On the other hand, when treated with GI-108B1, the production of IFN- γ by CD8+ T cells significantly increased (FIG. 8).

Furthermore, when treated with AMP and vehicle (hIgG4), the proliferation of CD4+ T cells was reduced. On the other hand, when treated with GI-108B1, the proliferation of CD4+ T cells significantly increased. In addition, the proliferation of CD4+ T cells increased even when treated with anti-CD73 antibody alone or with a combination of anti-CD73 antibody and IgG4 Fc-IL-2v. Meanwhile, when treated with PROLEUKIN®, Oleclumab or IgG4 Fc-IL-2v alone or when treated with a combination of Oleclumab and IgG4 Fc-IL-2v, the proliferation of CD4+ T cells did not tend to increase (FIG. 9). In addition, when treated with AMP and vehicle (hIgG4), the production of IFN- γ by CD4+ T cells was reduced. On the other hand, it was confirmed that when treated with GI-108B1, the production of IFN- γ by CD4+ T cells significantly increased (FIG. 10). Thereby, it was confirmed that GI-108B1 reinvigorated CD8+ T cell and CD4+ T cell activities suppressed by CD73-adenosine.

Example 11. Evaluation of GI-108B1 Activity Using pSTAT5 Analysis

To evaluate the activity of GI-108B1, healthy human PBMCs were treated with PROLEUKIN®, GI-108B1, IgG4 Fc-IL-2v and anti-CD73 antibody (GI- α CD73) alone, or a combination of anti-CD73 antibody (GI- α CD73) and IgG4 Fc-IL-2v, and pSTAT5 in T cell populations (CD8+ T cells and Treg cells) in the PBMCs was measured. Specifically, human PBMCs were treated with various concentrations of PROLEUKIN®, GI-108B1, IgG4 Fc-IL-2v, and anti-CD73 antibody alone, or with a combination of anti-CD73 antibody and IgG4 Fc-IL-2v, and incubated at 37° C. for 20 minutes. For cell surface staining, cells were stained with the following antibodies: anti-CD3-BV510 (Clone SP34-2, BD Biosciences), anti-CD4-PE-Cy7 (Clone OKT-4, Biolegend Inc.), anti-CD8-AF700 (Clone RPA-T8, Biolegend Inc.), anti-CD25-BV421 (Clone M-A251, BD Biosciences), anti-CD56-BV605 (Clone NCAM16.2, BD Biosciences), anti-CD127-BV650 (Clone A019DS, Biolegend Inc.), and Fixable Viability Dye eFluor 780 (Invitrogen Inc.).

For intracellular staining, the cells were fixed for 12 minutes by treatment with TFP Fix/Perm buffer (BD Biosciences), and incubated with Perm buffer III (BD Biosciences) for 30 minutes. The cells were then stained with anti-pSTAT5-AF647 (Clone 47/Stat5, BD Biosciences). Data on the stained cells were measured using Cytex Aurora (Cytex Biosciences) and analyzed using FlowJo software (BD Biosciences).

As a result, it was confirmed that the EC₅₀ value of GI-108B1 against pSTAT5 in CD8+ T cells was significantly

lower than that in the group treated with PROLEUKIN®, IgG4 Fc-IL-2v or anti-CD73 antibody alone or with a combination of anti-CD73 antibody and IgG4 Fc-IL-2v. Thereby, it was confirmed that GI-108B1 exhibited a better effect against STAT5 activation in CD8+ T cells than PROLEUKIN® (FIGS. 11 and 12).

In addition, it was confirmed that the EC₅₀ value of GI-108B1 against pSTAT5 in Treg cells was slightly higher than that in the group treated with IgG4 Fc-IL-2v or anti-CD73 antibody alone or with a combination of anti-CD73 antibody and IgG4 Fc-IL-2v, and was significantly higher than that in the group treated with PROLEUKIN®. Thereby, it was confirmed that GI-108B1 was much less effective in activating STAT5 in Treg cells compared to PROLEUKIN® (FIGS. 13 and 14).

Example 12. Evaluation of T-Cell Proliferation Ability of GI-108B1

Human PBMCs were suspended in 4Cell® Nutri-T GMP medium (Sartorius), and the cells were dispensed in a 96-well plate at a density of 5×10^5 cells per well. Then, the cells were treated with GI-108B1 or PROLEUKIN® at a final concentration of 10 nM in the presence of α CD3/CD28 dynabeads (1:1 ratio, Thermo Fisher Scientific). Every 2 to 3 days, the medium was replaced with a fresh medium containing each drug. The cells were counted on days 3, 5, 7, 10, and 12 using the ADAM™-MC2 Cell Counter (Nanoentek). On day 12, 2×10^6 cells of each group were stained with the following antibodies: anti-CD3-BV510 (Clone SP34-2, BD Biosciences), anti-CD4 (Clone OKT-4, Biolegend Inc.), anti-CD8-AF700 (Clone RPA-T8, Biolegend Inc.), anti-CD25-BV421 (Clone M-A251, BD Biosciences), anti-CD45-APC (Clone HI30, eBioscience), and Fixable Viability Dye eFluor 780 (Invitrogen Inc.). For intracellular staining, the cells were treated with fixation/permeabilization buffer (BD Biosciences) and stained with anti-Foxp3-PE antibody (Clone PCH101, Invitrogen Inc.). Data on the stained cells were measured using the Cytex Aurora instrument and analyzed using FlowJo software.

As a result, the number of CD8+ T cells in the group treated with GI-108B1 was higher than in the group treated with PROLEUKIN®, and the number of Treg cells was about 2-fold higher in the group treated with PROLEUKIN® than in the group treated with GI-108B1. As a result of calculating the ratio of the number of CD8+ T cells to the number of Treg cells (CD8+T/Treg), it was confirmed that the CD8+T/Treg in the group treated with GI-108B1 was about 2.46-fold higher than that in the group treated with PROLEUKIN®. Thereby, it was confirmed that GI-108B1 exhibited a superior effect on CD8+ T cell proliferation over Treg cell proliferation compared to PROLEUKIN® (FIG. 15).

Example 13. Cis-Binding Analysis of GI-108B1 in CD8+ T Cells

CD8+ T cells express both CD73 and the IL-2 receptor complex, so signaling induced by the binding of GI-108B1 to CD73 and IL-2 β on the same CD8+ T cells can exhibit a synergistic effect on the activation of CD8+ T cells. Accordingly, to examine whether GI-108B1 cis-binds to CD73 and IL-2 β on the same CD8+ T cells, cis-binding analysis was performed using CTV-CD8+ T cells (anti-CD73 antibody-treated/untreated group) and CTV+CD8+ T cells (FIG. 16). Specifically, CD8+ T cells were isolated from human PBMCs using the EASYSEPT™ Human CD8+

T Cell Isolation Kit (STEMCELL technologies). Half of the isolated CD8⁺ T cells were labeled with 0.5 μ M CTV (Cell Trace™ Violet, Thermo Fisher Scientific). The CTV-unlabeled cells were divided into two groups, and one group was pre-treated with a saturating concentration (1,000 nM) of anti-CD73 antibody at 4° C. for 2 hours to block all CD73 on the cell surface. Then, unbound anti-CD73 antibody was washed out.

Then, the CTV-unlabeled cells, pre-treated or untreated with anti-CD73 antibody were co-cultured with CTV-labeled cells at a 1:1 ratio. The co-cultured cells were treated with 0.5 nM GI-108B1 and incubated at 37° C. for 20 minutes. Thereafter, the cells were fixed with Cytofix fixation buffer (BD Biosciences) for 12 minutes, and then treated with Phosflow Perm buffer III (BD Biosciences) and incubated for 30 minutes. The cells were then stained with anti-pSTAT5-AF647 (Clone 47/Stat5, BD Biosciences). Data for the stained cells were measured using Cytex Aurora and analyzed using FlowJo software.

As a result, it was confirmed that, in the case of CTV+CD8⁺ T cells co-cultured with CTV-CD8⁺ T cells that were not pre-treated with anti-CD73 antibody, CTV-CD8⁺ T cells and CTV+CD8⁺ T cells showed similar levels of STAT5 phosphorylation, but in the case of CTV+CD8⁺ T cells co-cultured with CTV-CD8⁺ T cells pretreated with anti-CD73 antibody, the STAT5 phosphorylation level in CTV-CD8⁺ T cells was reduced by 40 to 60%, and the STAT5 phosphorylation level in CTV+CD8⁺ T cells remained the same (FIG. 17). From the fact that the proportion of hCD73+CD8⁺ T cells expressing hCD73 in purified CD8⁺ T cells from PBMCs was about 70% (FIG. 18) and from the above results, it was confirmed that GI-108B1 preferentially binds to CD73 in CD8⁺ T cells expressing CD73.

III. Evaluation of Anticancer Effect of Fusion Protein Dimer

Example 14. Evaluation of Immune Cell-Mediated Tumor Killing Ability of GI-108B1

To evaluate the immune cell-mediated tumor killing ability of GI-108B1, the anticancer effect of GI-108B1 was evaluated using a co-culture system of MDA-MB-231 cells and healthy human PBMC in the presence of anti-CD3/CD28 dynabeads and/or AMP. Specifically, the cytotoxicity of PBMCs against MDA-MB-231 wild-type cells was evaluated using flow cytometry-based Annexin V/7-AAD assay. PBMCs were purchased from Zen-bio Inc. or STEMCELL Technologies.

PBMCs were treated with hIgG4 (5 nM) alone, a combination of anti-CD73 antibody (GI- α CD73, 5 nM) and IgG4 Fc-IL-2v (5 nM), or GI-108B1 (5 nM) alone, and also treated with α CD3/CD28 dynabeads (1:1 ratio) or 500 U of AMP and cultured for 2 days before co-culture with target cells (MDA-MB-231 wild-type cells). Target cells were labeled with CTV and dispensed into a 24-well plate at a density of 1×10^5 cells per well. After overnight culture, pre-stimulated PBMCs were added to the target cells at a ratio of effector cells to target cells of 10:1, followed by co-culture for 24 hours at 37° C. under 5% CO₂. The cells were stained with Annexin V and 7-AAD (Biolegend Inc.), and the death rate of the CTV-labeled target cells was measured using the Symphony A3 instrument and analyzed using FlowJo software.

As a result, it was confirmed that the cell death rate was higher in the group (isotype control group (-AMP)) in which PBMCs treated only with anti-CD3/CD28 dynabeads and hIgG4 were co-cultured with target cells than in the group (isotype control group (+AMP)) in which PBMCs

treated with anti-CD3/CD28 dynabeads, AMP, and hIgG4 were co-cultured with target cells. Thereby, it was confirmed that the activity (cell killing ability) of immune cells was inhibited by treatment with AMP under adenosine-rich conditions.

Meanwhile, even though treated with AMP, the group in which PBMCs treated with anti-CD3/CD28 dynabeads, AMP and GI-108B1 were co-cultured with target cells and the group in which PBMCs treated with anti-CD3/CD28 dynabeads, AMP, and a combination of anti-CD73 antibody (GI- α CD73) and IgG4 Fc-IL-2v were co-cultured with target cells all showed higher cell death rates than the isotype control group (+AMP). In particular, the group in which PBMCs treated with anti-CD3/CD28 dynabeads, AMP and GI-108B1 were co-cultured with target cells showed a significantly higher target-cell death rate than the group in which PBMCs treated with anti-CD3/CD28 dynabeads, AMP and a combination of anti-CD73 antibody (GI- α CD73) and IgG4 Fc-IL-2v were co-cultured with target cells, indicating the highest cell death rate (FIG. 19).

Example 15. Evaluation of Anticancer Effect of GI-108B1 Administration in Mice Transplanted with Murine Colon Cancer Cells

After the test substance GI-108B1 was administered to a tumor model transplanted with the MC38 cell line (MC38-hCD73), a C57BL/6 murine colon cancer cell line in which the mouse CD73 gene was replaced with the human CD73 gene, the tumor growth inhibitory effect was evaluated. First, to produce a mouse model transplanted with a cancer cell line, a suspension of MC38-hCD73 cells (SMOC, China) was administered subcutaneously to the right back of each of C57BL/6J-hCD73 female mice (6-8 weeks old, SMOC, China) at 1×10^6 cells/100 μ L.

After MC38-hCD73 cells were transplanted, morbidity and mortality were checked daily, tumor volumes were measured for mice with no abnormalities in health, and 16 mice were randomly selected when the average tumor volume was 90 to 120 mm³. The animals were selected considering their physiological state, weight change, and tumor growth rate. The selected animals were divided into groups (n=8) as evenly as possible based on the tumor volume and body weight. Test groups were formed as shown in Table 10 below, and the test substance was administered thereto.

TABLE 10

Group	Substance administered	Route of administration	Dosing cycle	Dose	Animal number (n)
G1	Vehicle (PBS)	i.v.	QW (three times/3 weeks)	—	8
G2	GI-108B1	i.v.	QW (three times/3 weeks)	6 mg/kg	8

During the test period, changes in tumor, body weight, and diet were observed regularly, and changes in body weight and external appearance (hair/black eye) were checked twice a week. For tumor volume measurement, the maximum length (L) and perpendicular width (W) of the tumor were measured twice a week during the observation period using a digital caliper (Mitutoyo, Japan). Tumor volume (TV) and tumor growth inhibition (TGI) were measured by substituting the measured values into Equation 1 below.

$$TV(\text{mm}^3) = (W^2 \times L)/2 \quad < \text{Equation I} >$$

$$TGI = (1 - (Ti - T0)/(Vi - V0)) \times 100$$

Ti: tumor volume in the test substance-administered group at the end of the test

T0: tumor volume in the test substance-administration group at the time of first administration

Vi: tumor volume in the negative control group at the end of the test

V0: tumor volume in the negative control group at the time of first administration

The tumor volume in each animal before administration was set as the value measured at the time of grouping, and the antitumor efficacy was evaluated by comparison with the control group (vehicle (PBS), G1). As a result, the tumor volume in the GI-108B1-administered group was reduced compared to the control group (vehicle (PBS)) (FIG. 20). In addition, regarding the tumor growth inhibition rate, only one mouse in the control group (vehicle (PBS)) showed a tumor growth inhibition rate of over 50%, and none showed over 80%. On the other hand, in the GI-108B1-administered group, 5 mice showed a tumor growth inhibition rate of over 30%, and 3 mice over 50% (FIG. 21). Furthermore, when checking the survival rate of mice in each group, 2 mice in the control group (vehicle (PBS)) died on day 25, whereas all mice in the GI-108B1-administered group survived on day 25 (FIG. 22).

Example 16. Evaluation of Anticancer Effect of GI-108 Administration in Mice Transplanted with Human Breast Cancer Cells

In a tumor model obtained by transplanting MDA-MB-231 cells (human breast cancer cells) into mice into which the human immune system has been introduced, each of test substances GI-108A1 (AHP04167), GI-108B1, and Pembrolizumab (product name: Keytruda, MSD), an anti-PD-1 antibody, was administered, and then the tumor growth inhibitory effect was evaluated. First, in order to produce a mouse model into which the human immune system has been introduced, a suspension of human peripheral blood cells (Stemexpress, USA) was filled into a disposable syringe (31G, cat. 328820, BD, USA) and administered to NOG-B2m (NOD.Cg-B2mtm1UncPrkdcscidII2rgtm1Wjl/SzJ) female mice (7-week-old, The Jackson Laboratory, USA) through the caudal vein at 1×10^7 cells/200 μ L per animal. After cell transplantation, general symptoms were observed once daily.

NOG-B2m female mice were purchased from The Jackson Laboratory, and NOG-B2m female mice were maintained in individually ventilated cages housed in an animal biosafety level-3 facility at 19 to 25° C. and 30 to 70% humidity.

The MDA-MB-231 human triple-negative breast cancer (TNBC) cell line was purchased from ATCC and cultured in DMEM medium (Thermo Fisher Scientific) containing 10% FBS (Thermo Fisher Scientific) and 1% antibiotic/antimycotic (GIBCO™). The cultured MDA-MB-231 cells were harvested using trypsin (GIBCO™) and then suspended in PBS. To establish a xenograft mouse tumor model, healthy mice 5 days after transplantation with human peripheral blood cells were used, and a solution prepared by mixing a MDA-MB-231 cell suspension (5×10^6 cells/0.025 mL) and 0.025 mL Matrigel matrix phenol red-free (cat. 356237, BD,

USA) was filled into a disposable syringe (31G, cat. 328820, BD, USA) and administered to subcutaneously to the right back of each of the animals at 0.05 mL per animal. After MDA-MB-231 cell transplantation, general symptoms were observed once daily during the engraftment and growth period.

About 20 days after transplanting the MDA-MB-231 cells, the tumor volume was measured in mice with no abnormalities in health, and 32 mice were selected so that the average tumor volume of each group reached 80 to 100 mm³. The animals were selected considering their physiological state (respiration, fur, behavior, tail, posture, body fluids, diet, morphological deformation, metabolism, etc.), weight change, FACS analysis results, and tumor growth rate. The selected animals were divided into groups (n=8) as evenly as possible based on the tumor volume and body weight. It was confirmed that the proportion of human CD45+ cells in the total lymphocytes in the mouse peripheral blood before grouping was over 80%. The test groups were formed as shown in Table 11 below and the test substances were administered thereto (FIG. 23).

TABLE 11

Group	Substance administered	Route of administration	Dosing cycle	Dose	Animal number
G1	Vehicle (hlgG4)	i.v.	QW (three times/2.5 weeks)	6 mg/kg	8
G2	GI-108A1 (AHP04167)	i.v.	QW (three times/2.5 weeks)	6 mg/kg	8
G3	GI-108B1	i.v.	QW (three times/2.5 weeks)	6 mg/kg	8
G4	Anti-PD-1 (pembrolizumab)	i.v.	BIW (5 times/2.5 weeks)	5 mg/kg	8

During the test period, general symptoms such as appearance, behavior, and excrement were observed once a day and recorded for each mouse, and dead animals were checked. For body weight measurement and tumor volume measurement, the maximum length (L) and perpendicular width (W) of the tumor were measured twice a week during the observation period using a digital caliper. Tumor volume (TV) and tumor growth inhibition (TGI) were measured by substituting the measured values into Equation I described in Example 15. The tumor volume in each animal before administration was set as the value measured at the time of grouping, and the antitumor efficacy was evaluated by comparison with the control group (vehicle, G1).

All statistical calculations were performed using Prism 8.0 (Graph Pad Software Inc, USA). Comparison of tumor volume measurements was done using unpaired t-test. A p value of less than 0.05 was considered significant.

As a result, it was confirmed that tumor growth was inhibited in the groups treated with GI-108A1 (AHP04167), GI-108B1, and anti-PD-1 antibody compared to the control group (vehicle). In particular, the GI-108B1-administered group showed significantly more suppressed tumor growth than the GI-108A1 (AHP04167) or anti-PD-1 antibody-administered group (FIGS. 24 and 25). In addition, when the tumor growth inhibition rate was measured on day 18, only one mouse in the control group showed a tumor growth inhibition rate of over 30%, with none showing over 50% or 80%. In the anti-PD-1 antibody-administered group, 4 mice showed a tumor growth inhibition rate of over 30%, 2 mice over 50%, with none over 80%. On the other hand, in the GI-108A1-administered group, 5 mice showed a tumor

35

growth inhibition rate of over 30%, 3 mice over 50%, and 1 mouse over 80%. In the GI-108B1-administered group, 5 mice showed a tumor growth inhibition rate of over 30%, 4 mice over 50%, and 3 mice over 80% (FIG. 26).

Meanwhile, no weight loss appeared in both the control and test groups, and there was no significant difference between the groups (FIG. 27).

Example 17. Analysis of Vascular Permeability of GI-108B1

To analyze the vascular permeability of GI-108B1, 1×10^5 HUVEC cells (Lonza) were placed in a collagen-coated trans-well using a trans-well insert (9321012, CELLQART®), and cultured for 3 days to form a confluent monolayer. 1×10^5 PBMCs were dispensed into a 96-well plate, and then treated with PROLEUKIN® (5 nM) or GI-108B1 (5 nM) and cultured for 24 hours at 37° C. under 5% CO₂.

The HUVEC cells cultured in the trans-well were treated with the drug and the culture supernatant of the PBMCs and cultured for 24 hours at 37° C. under 5% CO₂. As a positive

36

control, 100 ng/ml of TNF- α (PeproTech) was used. Thereafter, the upper chamber was treated with dextran-FITC (70 kDa) and incubated for 1 hour and 30 minutes. The fluorescence of dextran-FITC that moved to the lower chamber was measured at wavelengths of 485 nm/500-550 nm (Ex/Em) using GLOMAX® Discover (Promega). Statistical calculations were analyzed by one-way ANOVA using Prism 8.0 (Graph Pad Software Inc.), and Dunnett's test was performed based on the negative control.

As a result, increased vascular permeability was observed in the positive control group treated with TNF- α . On the other hand, the group treated with GI-108B1 showed a level of vascular permeability similar to that of the untreated negative control group (FIG. 28).

Although the present invention has been described in detail with reference to the specific features, it will be apparent to those skilled in the art that this description is only of a preferred embodiment thereof, and does not limit the scope of the present invention. Thus, the substantial scope of the present invention will be defined by the appended claims and equivalents thereto.

SEQUENCE LISTING

```

Sequence total quantity: 43
SEQ ID NO: 1          moltype = AA  length = 19
FEATURE              Location/Qualifiers
REGION               1..19
                     note = mIgG signal
source               1..19
                     mol_type = protein
                     organism = synthetic construct

SEQUENCE: 1
MEWSWVFLPF LSVTTGVHS                               19

SEQ ID NO: 2          moltype = AA  length = 126
FEATURE              Location/Qualifiers
REGION               1..126
                     note = anti-hCD73 sdAb(AHF10235)
source               1..126
                     mol_type = protein
                     organism = synthetic construct

SEQUENCE: 2
QVQVQESGGG LVQAGDSLRL SCAASGRSIS RYNMGWFRQA PGKEREVFAA ISRSGGVTTY 60
ADSVKGRFTI SRDPAKSAVY LQMNSLKPED TAVYYCAADW RNSDSPSPLI IVYDYWGQGT 120
QVTVSS                                                    126

SEQ ID NO: 3          moltype = AA  length = 126
FEATURE              Location/Qualifiers
REGION               1..126
                     note = anti-hCD73 sdAb(AHF10240)
source               1..126
                     mol_type = protein
                     organism = synthetic construct

SEQUENCE: 3
EVQLVESGGG VVQTGGSLRL SCAASGRFTT HLMGWFRQA PGKEREVFAA ISNSGAGTQF 60
ADSVKGRFTI SRDPAKNTVY LQMNSLKPED TAVYYCAADW GTRDSPSKLN AVYDYWGRGT 120
QVTVSS                                                    126

SEQ ID NO: 4          moltype = AA  length = 126
FEATURE              Location/Qualifiers
REGION               1..126
                     note = anti-hCD73 sdAb(AHP04167)
source               1..126
                     mol_type = protein
                     organism = synthetic construct

SEQUENCE: 4
EVQLVESGGG LVQAGDSLRL SCVVSGRFTS NYHMGWFRQA PGKEREVFAV IGRSGGPTYT 60
ADSVKGRFTI SRDPAKNTVY LQMNSLKPED TAVYYCAGDW RNSDSPSKLK PVYDYWGQGT 120
QVTVSS                                                    126

SEQ ID NO: 5          moltype = AA  length = 18
FEATURE              Location/Qualifiers
REGION               1..18

```

-continued

```

source          note = Linker 1
                1..18
                mol_type = protein
                organism = synthetic construct

SEQUENCE: 5
GGGSAESKY GPPCPPCP                                     18

SEQ ID NO: 6      moltype = AA  length = 216
FEATURE          Location/Qualifiers
REGION           1..216
                note = hIgG4 Fc (F01)
source           1..216
                mol_type = protein
                organism = synthetic construct

SEQUENCE: 6
APEAAGGPSV FLFPPKPKDQ LMISRTPEVT CVVVDVSQED PEVQFNWYVD GVEVHNAKTK 60
PREEQFNSTY RVVSVLTVLH QDWLNGKEYK CKVSNKGLPS SIEKTISKAK GQPREPQVYT 120
LPPSQEEMTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTTTPVLDS DGSFPLY SRL 180
TVDKSRWQEG NVFSCSVLHE ALHNHYTQKS LSLSLG                               216

SEQ ID NO: 7      moltype = AA  length = 15
FEATURE          Location/Qualifiers
REGION           1..15
                note = Linker 2
source           1..15
                mol_type = protein
                organism = synthetic construct

SEQUENCE: 7
GGGSGGGGS GGGGS                                       15

SEQ ID NO: 8      moltype = AA  length = 133
FEATURE          Location/Qualifiers
REGION           1..133
                note = hIL2v3
source           1..133
                mol_type = protein
                organism = synthetic construct

SEQUENCE: 8
APTSSSTKKT QLQLEHLLLD LQMILNGINN YKNPKLTAML TAKFYMPKKA TELKHLQCLE 60
RELKPLEEVL NLAQSKNFHL RPRDLISNIN VIVLELKGSE TTFMCEYADE TATIVEFLNR 120
WITFCQSIIS TLT                                         133

SEQ ID NO: 9      moltype = AA  length = 119
FEATURE          Location/Qualifiers
REGION           1..119
                note = Variable heavy chain of anti-hCD73 Ab (CSA0060)
source           1..119
                mol_type = protein
                organism = synthetic construct

SEQUENCE: 9
QQLVQSGAE VKKPGSSVKV SCKFSGGSFT SYSFSWVRQA PGQGLEWMGR IIPVLTTTDY 60
AQKFRDRVIT TADESTSTAY MELSGLRSED TAVYYCASGL KKNWYFDLWG RGALITVSS 119

SEQ ID NO: 10     moltype = AA  length = 98
FEATURE          Location/Qualifiers
REGION           1..98
                note = Constant heavy chain of anti-hCD73 Ab (CSA0060)
source           1..98
                mol_type = protein
                organism = synthetic construct

SEQUENCE: 10
ASTKGPSVFP LAPCSRSTSE STAALGCLVK DYFPEPVTVS WNSGALTSGV HTPFAVLQSS 60
GLYSLSSVVT VPSSSLGTKT YTCNVDPKPS NTKVDKRV                               98

SEQ ID NO: 11     moltype = AA  length = 12
FEATURE          Location/Qualifiers
REGION           1..12
                note = Linker 3
source           1..12
                mol_type = protein
                organism = synthetic construct

SEQUENCE: 11
ESKYGPPCPP CP                                         12

SEQ ID NO: 12     moltype = AA  length = 10
FEATURE          Location/Qualifiers
REGION           1..10
                note = Linker 4
source           1..10

```

-continued

SEQUENCE: 12	mol_type = protein	
GGGSGGGGS	organism = synthetic construct	10
SEQ ID NO: 13	moltype = AA length = 107	
FEATURE	Location/Qualifiers	
REGION	1..107	
source	note = Variable light chain of anti-hCD73 Ab (CSA0060)	
	1..107	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 13		
DIQMTQSPSF LSASVGDRVT ITCQASQDIS HYLNWYQQKP GKAPKLLIYD ASSLETGVPS		60
RFSGSGSGTS FTLTISSLQP EDFATYYCQQ YDDFPLTFGG GTKVDIK		107
SEQ ID NO: 14	moltype = AA length = 107	
FEATURE	Location/Qualifiers	
REGION	1..107	
source	note = Constant light chain of anti-hCD73 Ab (0060, kappa)	
	1..107	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 14		
RSVAAPSVFI FPPSDEQLKS GTASVVCLLN NFYPREAKVQ WKVDNALQSG NSQESVTEQD		60
SKDSTYSLSS TLTLSKADYE KHKVYACEVT HQGLSSPVTK SFNRGEC		107
SEQ ID NO: 15	moltype = AA length = 8	
FEATURE	Location/Qualifiers	
REGION	1..8	
source	note = CDR1 of anti-hCD73 sdAb (AHF10235)	
	1..8	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 15		
GRSISRYN		8
SEQ ID NO: 16	moltype = AA length = 8	
FEATURE	Location/Qualifiers	
REGION	1..8	
source	note = CDR2 of anti-hCD73 sdAb (AHF10235)	
	1..8	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 16		
ISRSGGVT		8
SEQ ID NO: 17	moltype = AA length = 19	
FEATURE	Location/Qualifiers	
REGION	1..19	
source	note = CDR3 of anti-hCD73 sdAb (AHF10235)	
	1..19	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 17		
AADWRNSDSP SPLIIVDY		19
SEQ ID NO: 18	moltype = AA length = 8	
FEATURE	Location/Qualifiers	
REGION	1..8	
source	note = CDR1 of anti-hCD73 sdAb (AHF10240)	
	1..8	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 18		
GRTFTHLA		8
SEQ ID NO: 19	moltype = AA length = 8	
FEATURE	Location/Qualifiers	
REGION	1..8	
source	note = CDR2 of anti-hCD73 sdAb (AHF10240)	
	1..8	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 19		
ISNSGAGT		8
SEQ ID NO: 20	moltype = AA length = 19	
FEATURE	Location/Qualifiers	

-continued

REGION	1..19	
	note = CDR3 of anti-hCD73 sdAb (AHF10240)	
source	1..19	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 20		
AADWGTRDSP SKLNAVYDY		19
SEQ ID NO: 21	moltype = AA length = 8	
FEATURE	Location/Qualifiers	
REGION	1..8	
	note = CDR1 of anti-hCD73 sdAb (AHP04167)	
source	1..8	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 21		
GRTFSNYH		8
SEQ ID NO: 22	moltype = AA length = 8	
FEATURE	Location/Qualifiers	
REGION	1..8	
	note = CDR2 of anti-hCD73 sdAb (AHP04167)	
source	1..8	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 22		
IGRSGGPT		8
SEQ ID NO: 23	moltype = AA length = 19	
FEATURE	Location/Qualifiers	
REGION	1..19	
	note = CDR3 of anti-hCD73 sdAb (AHP04167)	
source	1..19	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 23		
AGDWRNSDSP SKLKPVDY		19
SEQ ID NO: 24	moltype = AA length = 8	
FEATURE	Location/Qualifiers	
REGION	1..8	
	note = VH-HCDR1 of anti-hCD73 Ab (CSA0060)	
source	1..8	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 24		
GGSFYSYS		8
SEQ ID NO: 25	moltype = AA length = 8	
FEATURE	Location/Qualifiers	
REGION	1..8	
	note = VH-HCDR2 of anti-hCD73 sdAb (CSA0060)	
source	1..8	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 25		
IIPVLTTT		8
SEQ ID NO: 26	moltype = AA length = 11	
FEATURE	Location/Qualifiers	
REGION	1..11	
	note = VH-HCDR3 of anti-hCD73 sdAb (CSA0060)	
source	1..11	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 26		
SGLKKNWYFD L		11
SEQ ID NO: 27	moltype = AA length = 6	
FEATURE	Location/Qualifiers	
REGION	1..6	
	note = VL-LCDR1 of anti-hCD73 Ab (CSA0060)	
source	1..6	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 27		
QDISHY		6
SEQ ID NO: 28	moltype = length =	

-continued

SEQUENCE: 28
000

SEQ ID NO: 29 moltype = AA length = 8
FEATURE Location/Qualifiers
REGION 1..8
 note = VL-LCDR3 of anti-hCD73 sdAb (CSA0060)
source 1..8
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 29
QYDDFPLT

8

SEQ ID NO: 30 moltype = DNA length = 1551
FEATURE Location/Qualifiers
misc_feature 1..1551
 note = nucleotides coding for anti-hCD73 sdAb (AHF10235)
 conjugated withhhIL2v3 and signal peptide
source 1..1551
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 30
atggaatggt cctgggtggt cctgttcttc ctgtccgtga ccaccggcgt gcactctcag 60
gtgcagggttc aagaatctgg cggcggattg gtgcaggccg gcgattctct gagactgtct 120
tgtgcccgcct ctggcccgttc catctctagg tacaacatgg gctgggttcag acaggcccct 180
ggcaaaagaga gagagtctgt cgcgcgccatc agcagatctg gcggagtgac ctactacgcc 240
gactccgtga aaggccggct gaccatctcc agagacaacg ccaagtctgc cgtgtacctc 300
cagatgaact cctgaagcc tgaggacacc gccgtgtact actgtgccg cgattggcgg 360
aaactccgact ctcttcttc tctgacatc gtgtacgact actggggcca gggcacccaa 420
gtgacagttt ctacggcagg cggaggcagt gctgagtcta agtatggccc tccttgtctc 480
ccatgtcctg ctccagaagc tgctggcggg ccctccgtgt ttctgttccc tccaaagcct 540
aaggaccagc tgatgatctc tcggacaccc gaagtgcact gcgtgggtgt ggatgtgtct 600
caagaggacc ctgaggtgca gttcaattgg tacgtggacg gcgtgggaagt gcacaacgcc 660
aagaccaagc ctagagagga acagttcaac tccacctaca gagtgtgtgc cgtgctgacc 720
gtgctgcacc aggatgtggt gaacggcaaa gagtacaagt gcaaggtgtc caacaagggc 780
ctgccttcca gcctcgaaaa gaccatctcc aaggccaagg gccagcctag ggaaccccag 840
gtttacaccc tgctccaag ccaagaggaa atgaccaaga accaggtgtc cctgacctgc 900
ctgggtcaagg gcttctaccc ttccgacatt gccgtggaat gggagagcaa tggccagcct 960
gagaacaact acaagaccac acctctctgt ctggactccg acggtctctt ctttctgtac 1020
tcccgcctga cctcggacaa gtccagatgg caagagggca acgtgttctc ctgctctgtg 1080
ctgcacgagg ccttcacaaa tctactaccc cagaagtccc taagcctgtc ccttggcggg 1140
ggcgtgggat ccgtctctac ctccagctcc accaagaaaa cccagtgta gctggaacat 1200
ctgctgtctg gattccagat gatcctgaat ggcatcaaca attacaagaa cccaagctg 1260
accgccatgc tgaccgctaa gtctctacatg cccaagaagg ccaccgagct gaagcacctc 1320
cagtgccctg aacgggaaat gaagcctctg gaagagggtg tgaatctggc ccagtgcaag 1380
aaactccacc tgaggcctcg ggacctgatc tccaacatca acgtgatcgt gctggaactc 1440
aagggtctcg agacaacctt catgtgcgag tacgccgacg agacagctac catcgtggaa 1500
tttctgaatc ggtggatcac cttctgccag tccatcatca gcacctgac c 1551

SEQ ID NO: 31 moltype = DNA length = 1494
FEATURE Location/Qualifiers
misc_feature 1..1494
 note = nucleotides coding for anti-hCD73 sdAb (AHF10235)
 conjugated withhhIL2v3
source 1..1494
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 31
caggtgcagg ttcaagaatc tggcggcggg ttggtgcagg ccggcgattc tctgagactg 60
tcttgtgccg cctctggccg gtccatctct aggtacaaca tgggctggtt cagacaggcc 120
cctggcaaaag agagagagtt cgtcgccgcc atcagcagat ctggcggagt gacctactac 180
gccgactccg tgaaggccg gctgaccatc tccagagaca acgccaagtc tgcctgtgac 240
ctccagatga actccctgaa gcttgaggac accgccgtgt actactgtgc cgcgatttgg 300
cggaactccg actctctctc tctctctgat atcgtgtacg actactgggg ccagggcacc 360
caagtgcagc tttctagcgg aggcggaggg agtgctgagt ctaagtattg ccctccttgt 420
cctccatgtc ctgctccaga agctgctggc gggccctccg tgtttctgtt ccctccaaag 480
cctaaggacc agctgatgat ctctcggaca cccgaagtga cctgcgtggt ggtggatgtg 540
tctcaagagg accctgaggt gcagttcaat tggtagctgg acggcgtgga agtgcaaac 600
gccaagacca agcctagaga ggaacagttc aactccacct acagagtgtg gtcctgtgctg 660
accgtgctgc accaggattg gctgaacggc aaagagtaca agtgcaaggt gtccaacaag 720
ggcctgcctt ccagctatga aaagaccatc tccaaggcca agggccagcc tagggaaacc 780
caggtttaca cctcgcctcc aagccaagag gaaatgacca agaaccaggt gtccctgacc 840
tgccctggtc agggcttcta ccttcccgac attgcccgtg aatgggagag caatggccag 900
cctgaagaaca actacaactc cacacctcct gtgctggact ccgacggctc cttcttctg 960
tactccgccg tgaccgtgga caagtccaga tggcaagagg gcaacgtgtt ctctgtctct 1020
gtgctgcacg aggcctgca caatcactac acccagaagt ccctcagcct gtcccttgcc 1080
ggaggcgggt gatccgctc tacctccagc tccaccaaga aaaccagtt gcagctggaa 1140
catctgctgc tggacctcca gatgatcctg aatggcatca acaattacaa gaaccccaag 1200
ctgaccgcca tgctgaccgc taagtcttac atgccaaga agggccacca gctgaagcac 1260

-continued

```

ctccagtgc tggaaaggga actgaagcct ctggaagagg tgctgaatct ggcccagtc 1320
aagaacttcc acctgaggcc tcgggacctg atctccaaca tcaacgtgat cgtgctggaa 1380
ctcaagggtc ccgagacaac cttcatgtgc gagtacgccg acgagacagc taccatcgtg 1440
gaatttctga atcgggtgat caccttctgc cagtccatca tcagcaccct gacc 1494

SEQ ID NO: 32      moltype = DNA length = 1551
FEATURE
misc_feature      1..1551
                  note = nucleotides coding for anti-hCD73 sAb (AHF10240)
                  conjugated withhIL2v3 and signal peptide
source            1..1551
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 32
atggaatggt cctgggtgtt cctgttcttc ctgtccgtga ccaccggcgt gcactctgaa 60
gtgcagctgg ttgaatctgg cggcgagtg gttcagacag gcggtatctt gagactgtct 120
tgtccgcctt ccggcagaac ctttacacac ctggctatgg gctggttccg gcaggctcct 180
ggaaaagaga gagagtctgt cgcgcgcctc tccaactctg gcgctggcac acagttcgcc 240
gactctgtga agggcagatt caccatctct cgggacaacg ccaagaacac cgtgtacctc 300
cagatgaact cctgaagcc tgaggacacc gccgtgtact actgtgctgc tgattggggc 360
accagagaca gccctcttaa gctgaatgcc gtgtacgact actggggcag aggcacccaa 420
gtgacagtg cctctggagg cggaggcagt gctgagtcta agtatggccc tccttgtcct 480
ccatgtcctg ctccagaagc tgctggcggg cctcccggtg ttctgttccc tccaaagcct 540
aaggaccagc tgatgatctc tcggcacccc gaagtgcact gcgtggtggt ggatgtgtct 600
caagaggacc ctgaggtgca gttcaattgg tacgtggacg gcgtggaagt gcacaacgcc 660
aagaccaagc ctagagagga acagttcaac tccacctaca gagtgggtgc cgtgtgacc 720
gtgtctgacc aggtattggc gaacggcaca gagtacaagt gcaaggtgtc caacaagggc 780
ctgccttcca gcatcgaaaa gacctctccc aaggccaagg gccagcctag ggaaccccag 840
gtttacaccc tgctctcaag ccaagaggaa atgaccaaga accaggtgtc cctgacctgc 900
ctggtcaagg gcttctaccc ttccgacatt gccgtggaat gggagagcaa tggccagcct 960
gagaacaaat acaagaccac acctcctgtg ctggactccg acggtctcct ctttctgtac 1020
tcccgcctga ccctggacaa gtccagatgg caagagggca acgtgttctc ctgctctgtg 1080
ctgcacgagg ccctgcacaa tcaactacac cagaagtccc tcagcctgtc ccttggcgga 1140
ggcgtgggat cctcctctac ctccagctcc accaagaaaa ccagttgca gctggaacat 1200
ctgctgctgg acctccagat gatcctgaat ggcatcaaca attacaagaa ccccaagctg 1260
accgcatctg tgaccgctaa gttctacatg cccaagaagg ccaccgagct gaagcacctc 1320
cagtgcctgg aacgggaact gaagcctctg gaagaggtgc tgaatctggc ccagtccaag 1380
aacttccacc tgaggcctcg ggacctgac tccaacatca acgtgatcgg gctggaaactc 1440
aagggtcccg agacaacctt catgtgcgag tacgcccagc agacagctac catcgtggaa 1500
tttctgaatc ggtggatcac cttctgccag tccatcatca gcacctgac c 1551

SEQ ID NO: 33      moltype = DNA length = 1494
FEATURE
misc_feature      1..1494
                  note = nucleotides coding for anti-hCD73 sAb (AHF10240)
                  conjugated withhIL2v3
source            1..1494
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 33
gaagtgcagc tgggtgaatc tggcgcgcca gtggttcaga caggcggatc tctgagactg 60
tcttgtgccg cctccggcag aacctttaca cacttggtta tgggttggtt ccggcaggct 120
cctggaaaag agagagagtt cgtcgccgcc atctccaact ctggcgctgg cacacagttc 180
gccgactctg tgaaggcgag attcaccatc tctcgggaca acgccaagaa caccgtgtac 240
ctccagatga actcctctga gctgaggac accgcgctgt actactgtgc tgctgattgg 300
ggcaccagag acagcccttc taagctgaat gccgtgtacg actactgggg cagaggcacc 360
caagtgcacg tgtcctctgg aggcggaggg agtgctgagt ctaagtatgg cctccttgt 420
cctccatgtc ctgctccaga agctgctggc gggccctccg tgtttctgtt cctccaaag 480
cctaaggacc agctgatgat ctctcggaca cccgaagtga cctgcgtggt ggtggatgtg 540
tctcaagagg acctgaggt gcagttcaat tggtagctgg acggcgtgga agtgacaaac 600
gccaaagcca agcctagaga ggaacagttc aactccacct acagagtggg gtcctgtctg 660
accgtgctgc accaggattg gctgaacggc aaagagtaca agtgcaaggt gtccaacaag 720
ggcctgcctt ccagcatcga aaagaccatc tccaaggcca agggccagcc tagggaaacc 780
caggtttaca cctgcctccc aagccaagag gaaatgacca agaaccaggt gtcctgacc 840
tgctgtgtca agggcttcta ccttcccgac attgccgtgg aatgggagag caatggccag 900
cctgagaaca actacaagac cacacctcct gtgtggact ccgacggctc cttcttctgt 960
tactcccgcc tgaccgtgga caagtccaga tggcaagagg gcaacgtgtt ctcctgtctt 1020
gtgtgcacg aggcctgca caatcactac acccagaagt cctcagcctt gtccttggc 1080
ggaggcgtgt gatccgtccc taactccagc tccaccaaga aaacccagtt gcagctggaa 1140
catctgtctg tggacctcca gatgatcctg aatggcatca acaattacaa gaaccccaag 1200
ctgaccgcca tgtgaccgc taagtctac atgcccaga aggccaccga gctgaagcac 1260
ctccagtgc tggaaaggga actgaagcct ctggaagagg tgctgaatct ggcccagtc 1320
aagaacttcc acctgaggcc tcgggacctg atctccaaca tcaacgtgat cgtgctggaa 1380
ctcaagggtc ccgagacaac cttcatgtgc gagtacgccg acgagacagc taccatcgtg 1440
gaatttctga atcgggtgat caccttctgc cagtccatca tcagcaccct gacc 1494

SEQ ID NO: 34      moltype = DNA length = 1551
FEATURE
misc_feature      1..1551

```


-continued

```

note = nucleotides coding for anti-hCD73 sdAb (AHP04167)
      conjugated withhIL2v3 and signal peptide
source 1..1551
        mol_type = other DNA
        organism = synthetic construct

SEQUENCE: 34
atggaatggg cctgggtgtt cctgttcttc ctgtccgtga ccaccggcgt gcactctgaa 60
gtgcagctgg ttgaatctgg cggcggattg gttcaggctg gcggatctct gagactgtcc 120
tgtgtggtgt ccgcgagaac cttctccaac taccacatgg gctggttcag acaggccctc 180
ggcaaagaac gcgagtctgt ggctgtgatc ggcagatctg gcggacctac ctactacgcc 240
gactccgtga agggcagatt caccatctcc agagacaacg ccaagaacac cgtgtacctc 300
cagatgaact ccttgaagcc tgaggacacc gccgtgtact actgtgccgg cgattggaga 360
aactccgact ctcccagcaa gctgaagccc gtgtacgatt attggggcca gggcacccaa 420
gtgaccgtgt cctctggagg cggaggcagt gctgagtcta agtatggccc tcttgtctct 480
ccatgtctct ctccagaagc tgctggcggg ccttccgtgt ttctgttccc tccaaagcct 540
aaggaccagc tgatgatctc tcggacaccc gaagtgaact gcgtgggtgt ggatgtgtct 600
caagaggacc ctgaggtgca gttcaattgg tacgtggacg gcgtggaagt gcacaacgcc 660
aagaccaagc ctgagagagg acagttcaac tccacctaca gagtgggtgc cgtgtgacc 720
gtgtgtcacc aggtattggc gaacggcaaa gagtacaagt gcaaggtgtc caacaagggc 780
ctgccttcca gcatcgaaaa gacctctccc aaggccaagg gccagcctag ggaaccccag 840
gtttacaccc tgcttcaag ccaagaggaa atgaccaaga accaggtgtc cctgacctgc 900
ctggtcaagg gcttctaccc ttccgacatt gccgtggaat gggagagcaa tggccagcct 960
gagaacaact acaagaccac acctctgtg ctggactccg acggtctcct ctttctgtac 1020
tcccgcctga ccgtggacaa gtccagatgg caagagggca acgtgttctc ctgctctgtg 1080
ctgcacgagg ccttcgacaa tcactacacc cagaagtccc tcagcctgtc ccttggcgga 1140
ggcgggtggt acctgctctc ctccagctcc accaagaaaa cccagttgca gctggaacat 1200
ctgtgtctgg acctccagat gatcctgaat ggcatacaaa attacaagaa ccccaagctg 1260
accgccatgc tgaccgctaa gttctacatg cccaagaagg ccaccgagct gaagcacctc 1320
cagtgcctgg aacgggaact gaagcctctg gaagaggtgc tgaatctggc ccagtccaag 1380
aacttccacc tgaggcctcg ggacctgatc tccaacatca acgtgatcgt gctggaactc 1440
aagggtctcg agacaacctt catgtgcgag tacgccgacg agacagctac catcgtggaa 1500
ttctgtaatc ggtggatcac cttctgccag tccatcatca gcacctgac c 1551

SEQ ID NO: 35      moltype = DNA length = 1494
FEATURE            Location/Qualifiers
misc_feature        1..1494
                    note = nucleotides coding for anti-hCD73 sdAb (AHP04167)
                    conjugated withhIL2v3
source              1..1494
                    mol_type = other DNA
                    organism = synthetic construct

SEQUENCE: 35
gaagtgcagc tgggtgaatc tggcggcgga ttggttcagg ctggcggatc tctgagactg 60
tctgtgtggt tgtccggcag aaccttctcc aactaccaca tgggctgggt cagacaggcc 120
cctggcaaaag aacgcgagtt cgtggctgtg atcggcagat ctggcggacc tacctactac 180
gccgactccg tgaagggcag attcaccatc tccagagaca acgccaagaa cacogtgtac 240
ctccagatga actccctgaa gcctgaggac accgcctgtg actactgtgc cggcgtattg 300
agaaactccg actctcccaa caagctgaag cccgtgtacg attattgggg ccagggcacc 360
caagtgcacc tgtcctctgg aggcggaggc agtgctgagt ctaagtatgg cctccttgt 420
cctccatgtc ctgtctcaga agctgtgtgc gggccctccg tgtttctgtt cctccaaaag 480
cctaaggacc agctgatgat ctctcggaca cccgaagtga cctgcgtggt ggtggatgtg 540
tctcaagagg accctgaggt caggttcaat tggtagctgg acggtgtgga agtgacaaac 600
gccaagacca agcctagaga ggaacagttc aactccacct acagagtgtg gtccgtgtgt 660
accgtgtctg accaggattg gctgaacggc aaagagtaca agtgcaaggt gtccaacaag 720
ggcctgcctt ccagactcga aagacccatc tccaaggcca agggccagcc tagggaaacc 780
caggtttaca cctgtcctcc aagccaagag gaaatgacca agaaccaggt gtccctgacc 840
tgctgtgtca agggcttcta ccttccgac attgctgtgg aatgggagag caatggccag 900
cctgagaaca actacaagac cacacctcct gtgtgtgact ccgacggtc cttcttctgt 960
tactcccgcc tgaccgtgga caagtccaga tggcaagagg gcaacgtgtt ctctgtctct 1020
gtgtgtcacg aggccttgca caatcactac acccagaagt ccctcagcct gtcccttgcc 1080
ggaggcgtgt gatccgtccc tacctccagc tccaccaaga aaaccagtt gcagctggaa 1140
catctgtctg tggacctcca gatgatectg aatggcatca acaattacaa gaaccccaag 1200
ctgaccgcca tgtgaccgcg taagtctctc atgcccaga aggccaccga gctgaagcac 1260
ctccagtgcg tggaaacggg actgaagcct ctggaagagg tgctgaatct ggccagctcc 1320
aagaacttcc acctgaggcc tcgggacctg atctccaaca tcaacgtgat cgtgtggaa 1380
ctcaagggtc ccgagacaac cttcatgtgc gagtacggcg acgagacagc taccatcgtg 1440
gaatttctga atcgtgtgat caccttctgc cagtccatca tcagcaccct gacc 1494

SEQ ID NO: 36      moltype = DNA length = 1806
FEATURE            Location/Qualifiers
misc_feature        1..1806
                    note = nucleotides coding for heavy chain of anti-hCD73
                    (hIgG4Fc)conjugated with hIL2v3 and signal peptide
source              1..1806
                    mol_type = other DNA
                    organism = synthetic construct

SEQUENCE: 36
atggaatggg cctgggtgtt cctgttcttc ctgtccgtga ccaccggcgt gcactctcag 60
gttcagtgtg ttcagtctgg cgccgaagtg aagaaacccg gctcctctgt gaaggtgtcc 120

```

-continued

```

tgcgaagtttt ceggcgggtc cttacacctc tacagcttct cctgggtccg acagggtcct 180
ggacaaggct tggagtggtat gggcagaatc atccccgtgc tgaccaccac cgattacgcc 240
cagaatttcc gggacagagt gaccatcacc gccgacgagt ctacctccac cgctacatg 300
gaactgtccg gcctgagatc tgaggacacc gccgtgtact actgtgccag cggcctgaag 360
aagaactggg acttcgacct gtggggcaga ggcgctctga tcacagtgtc ctccgcttcc 420
accaagggac ccagcgtggt cctctctggc cctgtctcca gatccacctc cgagtctaca 480
gctgctctgg gctgtctggg caaggactac ttcctgagc ctgtgaccgt gtcttggaac 540
tctggcgctc tgacatctgg cgtgcacacc ttccagctg tgetccagtc ctccggcctg 600
tactctctgt ctagcgtcgt gaccgtgcct tctctagcc tgggcaccaa gacctacacc 660
tgtaatgtgg accacaagcc ttccaacacc aaggtggaca agcgcgtgga atctaagtac 720
ggcctctctt gtctccatg tctctctcca gaagctgctg gcggcccttc cgtgtttctg 780
ttccctccaa agcctaagga ccagctgatg atctctcgga cccctgaagt gacctgctg 840
gtggtggatg tgtcccaaga ggacctgag gtgcagtcca attggtacct ggacggcgtg 900
gaagtgcaca acgccaagac caagcctaga gaggaacagt tcaactccac ctacagagt 960
gtgtccgtgc tgaccgtgct gcaccaggat tggctgaacg gcaagagta caagtgaag 1020
gtgtccaaca agggcctgcc ttccagcatc gaaaagacca tctctaaggc taagggccag 1080
cctcgcgagc ccaggtttta ccattgcct ccaagccaag aggaatgac caagaaccag 1140
gtgtccctga cctgcctcgt gaagggcttc tacccttccg acattgcct ggaatgggag 1200
tccaatggcc agcctgagaa caactacaag acaaccctc ctgtgctgga ctccgacggc 1260
tccctcttcc tctattcccg cctgaccgtg gacaagtcta ggtggcaaga gggcaacgtg 1320
ttctcctgct ctgtgctgca cgaggccctg cacaatcact acaccacgaa gtctctgtct 1380
ctgtcccttg gcggaggcgg aggatccgct cctacctcca gctccacca gaaaaccag 1440
ttgcagctgg aacatctgct gctggacctc cagatgatcc tgaatggcct caacaattac 1500
aagaacccca agctgaccgc catgctgacc gctaagttct acatgcccaa gaaggccacc 1560
gagctgaagc acctccagtg cctggaacgg gaactgaagc ctctggaaga ggtgctgaat 1620
ctggcccaagt acaagaactt ccacctgagg cctcgggacc tgatctccaa catcaacgtg 1680
atcgtgctgg aactcaaggg ctccgagaca acctcatgt gcgagtacgc cgacgagaca 1740
gctaccatcg tggaatttct gaatcgggtg atcaccttct gccagtcacat catcagcacc 1800
ctgacc 1806

```

```

SEQ ID NO: 37      moltype = DNA length = 1749
FEATURE            Location/Qualifiers
misc_feature       1..1749
                    note = nucleotides coding for heavy chain of anti-hCD73
                    (hIgG4Fc) conjugated with hIL2v3
source             1..1749
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 37
caggttcagt tggttcagtc tggcgccgaa gtgaagaaac ccggtcctc tgtgaagggtg 60
tcccgcaagt tttccggcgg ctcccttacc tctacagct tctcctgggt ccgacaggct 120
cctggacaag gcttggagtg gatgggcaga atcatccccg tgetgaccac caccgattac 180
gccagaaat ccggggacag atgtgaccatc accgcccagc agtctacctc caccgcctac 240
atggaactgt ccggcctgag atctgaggac accgcccgtg actactgtgc cagcggcctg 300
aagaagaact ggtacttcga cctgtggggc agaggcgctc tgatccagct gtccctccgt 360
tccaccaagg gaccacagct gttccctctg gctccttctg ccagatccac ctccgagtct 420
acagctgctc tgggctgtct ggtcaaggac tacttccctg agcctgtgac cgtgtcttgg 480
aactctggcg ccttgacatc tggcgtgcac acctttccag ctgtgtctcca gtccctccgc 540
ctgtactctc tgtctagcgt cgtgaccgtg ccttctctca gccggggcac caagacctac 600
acctgtaatg tggaccacaa gcccttccaa accaagggtg acaagcgcgt ggaatctaag 660
tacggccctc cttgtcctcc atgtcctgct ccagaagctg ctggcggccc ttccgtgttt 720
ctgttccctc caaagcctaa ggaccagctg atgatctctc ggaccacctga agtgacctgc 780
gtggtgggtg atgtgtccca agaggacctt gaggtgcagt tcaattggta cgtggacggc 840
gtggaagtgc acaacgccaa gaccaagcct agagaggaaac agttcaactc cacctacaga 900
gtggtgtccg tctgagccgt gctgcaccag gattggctga acggcaaaaga gtacaagtgc 960
aagggtgtcc acaagggcct gcccttccagc atcgaaaaga ccatctctaa ggctaagggc 1020
cagcctcgcg agccccaggt ttacacattg cctccaagcc aagaggaaat gaccaagaac 1080
cagggtgtcc tgacctgctc cgtgaagggc ttctaccctt ccgacattgc cgtggaatgg 1140
gagtccaatg gccagcctga gaacaactac aagacaaccc ctctgtgtgt ggactccgac 1200
ggctccttct ttctgtattc ccgcctgacc gtggacaagt ctagggtggca agagggcaac 1260
gtgttctcct gctctgtgct gcacgaggcc ctgcacaatc actacacca gaagtctctg 1320
tctctgtccc ttggcggagg cggaggatcc gctcctacct ccagctccac caagaaaacc 1380
cagttgcagc tggaaacatct gctgctggac ctccagatga tctgtaattg catcaacaat 1440
tacaagaacc ccaagctgac gccctagcgt accgctaagt tctacatgcc caagaaggcc 1500
accgagctga agcacctcca gtgcctggaa cgggaactga agcctctgga agagggtgctg 1560
aatctggccc agtccaagaa cttccacctg aggcctcggg acctgatctc caacatcaac 1620
gtgatcgtgc tggaaactcaa gggctccgag acaaccttca tgtgcgagta cgccgacgag 1680
acagctacca tcgtggaatt tctgaatcgg tggatcacct tctgcccagtc catcatcagc 1740
acctgacc 1749

```

```

SEQ ID NO: 38      moltype = DNA length = 699
FEATURE            Location/Qualifiers
misc_feature       1..699
                    note = nucleotides coding for light chain of anti-hCD73
                    (hIgG4Fc) conjugated with hIL2v3 and signal peptide
source             1..699
                    mol_type = other DNA
                    organism = synthetic construct
SEQUENCE: 38

```

-continued

```

atggaatggt cctgggtgtt cctgtttctt ctgtccgtga ccaccggcgt gcactccgac 60
attcagatga cccagtctcc atccttctct agcgctctgt tgggcgacag agtgaccatt 120
acctgtcagg ccagccagga catctccacc tacctgaact ggatcacgca gaagcccgcc 180
aaggccctta agctgtgat ctacgatgcc tcctctctgt aaactggcgt gccctccaga 240
ttctccggct ctggctctgt caccagcttt accctgacca tctctagcct ccagcctgag 300
gacttcgcca cctactactg ccagcagtag gacgacttcc cactgacctt tggcggaggc 360
accaaggtgg acatcaagag atctgtggcc gctccttccg tgttcattct cccacttcc 420
gacgagcagc tgaagtcggg cacagcttct gtcgtgtgcc tgtgaaaca cttctacct 480
cgggaagcca aggtgcagtg gaaagtggat aacgccctcc agtccggcaa ctcccagag 540
tctgtgaccg agcaggactc caaggacagc acctacagcc tgtcctccac actgacctg 600
tccaaggccg actacgagaa gcacaagggt tacgcctgcg aagtgaacca tcagggcctg 660
tctagccctg tgaccaagtc tttcaaccgg ggcgagtgt 699

```

```

SEQ ID NO: 39      moltype = DNA length = 642
FEATURE           Location/Qualifiers
misc_feature       1..642
                   note = nucleotides coding for light chain of anti-hCD73
                   (hIgG4Fc) conjugated with hIL2v3
source            1..642
                   mol_type = other DNA
                   organism = synthetic construct

```

```

SEQUENCE: 39
gacattcaga tgaccagtc tccatccttc ctgagcgctt ctgtggggcga cagagtgacc 60
attacctgtc aggccagcca ggacatctcc cactacctga actgggtatca gcagaagccc 120
ggcaaggccc ctaagtgtct gatctacgat gcctcctctc tggaaactgg cgtgccctcc 180
agattctccg gctctggctc tggcaccagc tttacctga ccatctctag cctccagcct 240
gaggacttcc ccactacta ctgccagcag tacgacgact tcccaactgac ctttggcgga 300
ggcaccgaag tggacatcaa gagatctgtg gccgctcctt ccgtgttcat cttcccacct 360
tccgacgagc agctgaagtc cggcacagct tctgtcgtgt gcctgctgaa caacttctac 420
cctcgggaag ccaaggtgca gtggaaagtg gataacgccc tccagtcagg caactcccaa 480
gagctctgta ccgagcagga ctccaaggac agcacctaca gcctgtcctc cacactgacc 540
ctgtccaagg ccgactacga gaagcacaa gtgtacgcct gcgaagtgac ccatcaggggc 600
ctgtctagcc ctgtgaccaa gtctttcaac cggggcgagt gt 642

```

```

SEQ ID NO: 40      moltype = AA length = 133
FEATURE           Location/Qualifiers
REGION           1..133
                   note = hIL-2
source           1..133
                   mol_type = protein
                   organism = synthetic construct

```

```

SEQUENCE: 40
APTSSTTKT QLQLEHLLLD LQMILNGINN YKNPKLTRML TFKFYMPKKA TELKHLQCLE 60
EELKPLEEVL NLAQSKNFHL RPRDLISNIN VIVLELKGSE TTFMCEYADE TATIVEFLNR 120
WITFCQSIIS TLT 133

```

```

SEQ ID NO: 41      moltype = AA length = 574
FEATURE           Location/Qualifiers
REGION           1..574
                   note = hCD73
source           1..574
                   mol_type = protein
                   organism = synthetic construct

```

```

SEQUENCE: 41
MCPRAARAPA LLLLALGAVL WPAAGAWELT ILHTNDVHSR LEQTSSESSK CVNASRCMG 60
VARLFTKVQQ IRRAEPNVLL LDAGDQYQGT IWFTVYKGAE VAHFMNALRY DAMALGNHEF 120
DNGVEGLIEP LLKEAKFPIL SANIKAKGPL ASQISGLYLP YKVLPGVDEV VGIVGYTSKE 180
TPFLSNPGTN LVFEDEITAL QPEVDKLTAL NVNKIIALGH SGFEMDKLIA QKVRGVDVVV 240
GGHSNTFLYT GNPPSKEVPA GKYPFIIVSD DGRKVPVVQA YAFGKYLGYL KIEFDERGNV 300
ISSHGNPILL NSSIPEDPSI KADINKWRIK LDNYSTQELG KTIVYLDGSS QSCRPRECNM 360
GNLICDAMIN NNLRHTDEM FWNHVMCILN GGGIRSPIDE RNNGTITWEN LAAVLPFGGT 420
FDLVQLKGST LKKAFAHVSF RYQSTGEFL QVGGIHVVYD LSRKPGDRVV KLDVLCTKCR 480
VPSYDPLKMD EVYKVIPLNF LANGGDGFQM IKDELLRHDS GDQDINVVST YISKMKVIYP 540
AVEGRIFKST GSHCHGSFSL IFLSLWAVIF VLYQ 574

```

```

SEQ ID NO: 42      moltype = DNA length = 1392
FEATURE           Location/Qualifiers
misc_feature       1..1392
                   note = nucleotides coding for heavy chain of anti-hCD73
                   (hIgG4Fc) conjugated with signal peptide
source            1..1392
                   mol_type = other DNA
                   organism = synthetic construct

```

```

SEQUENCE: 42
atggaatggt cctgggtgtt cctgtttctt ctgtccgtga ccaccggcgt gcactctcag 60
gttcagttgg ttcagtctgg cgcgaagtg aagaaccccg gctcctctgt gaagtggtcc 120
tgcaagtttt ccggcggtcc ctttacctcc tacagcttct cctgggtccg acaggctcct 180
ggacaaggct tggagtggat gggcagaatc atccccgtgc tgaccaccac cgattacgcc 240
cagaaattcc gggacagagt gaccatcacc gccgacgagt ctacctccac cgcctacatg 300

```

-continued

```

gaactgtccg gcectgagatc tgaggacacc gccgtgtact actgtgccag cggcctgaag 360
aagaactggt acttogagact gtggggcaga ggcgtctga tcacagtgtc ctccgcttcc 420
accaagggac ccagcgtggt cctcttggt ccttgcctca gatccacctc cgagtctaca 480
gctgctctgg gctgtctggt caaggactac ttccctgagc ctgtgaccgt gtcttggaac 540
tctggcgctc tgacatctgg cgtgcacacc ttccagctg tgctccagtc ctccggcctg 600
tactctctgt ctagcgtcgt gaccgtgcct tctctagcc tgggcaccaa gacctacacc 660
tgtaatgtgg accacaagcc ttccaacacc aaggtggaca agcgcgtgga atctaagtac 720
ggccctcctt gteectcatg teetgtctca gaagctgctg gcggcccttc cgtgtttctg 780
ttccctccaa agcctaagga ccagctgatg atctctcgga cccctgaagt gacctgctg 840
gtggtggatg tgtcccaaga ggacctgag gtgcagtcca attggtacct ggacggcgtg 900
gaagtgcaca acgccaagac caagcctaga gaggaacagt tcaactccac ctacagagtg 960
gtgtccgtgc tgacgcgtgt gcaccaggat tggctgaacg gcaaaagagta caagtgaag 1020
gtgtccaaca agggcctgcc ttccagcatc gaaaagacca tctctaaggc taaggggccag 1080
cctcgcgagc ccaggttcta cacattgcct ccaagccaaag aggaaatgac caagaaccag 1140
gtgtccctga cctgcctcgt gaagggcttc tacccttccg acattgacct ggaatgggag 1200
tccaatggcc agcctgagaa caactacaag acaaccctc ctgtgtctgga ctccgacggc 1260
tctctcttcc tgattctccg cctgacgtgt gacaagtcta ggtggcaaga gggcaacgtg 1320
ttctcctgct ctgtgctgca cgaggccctg cacaatcact acaccacgaa gtctctgtct 1380
ctgtcccttg gc 1392

```

SEQ ID NO: 43 moltype = DNA length = 1098
FEATURE Location/Qualifiers
misc_feature 1..1098
 note = nucleotides coding for light chain of hIgG4Fc
 conjugated with hIL2v3 and signal peptide
source 1..1098
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 43

```


gaatctaagt acggccctcc ttgtctctca tgtctgtctc cagaagtgtc tggcggccct 60
tcogtgtttc tgttccctcc aaagcctaag gaccagctga tgatctctcg gacctctgaa 120
gtgacctgag tgggtggtgga tgtgtcccaa gaggaacctg aggtgcagtt caattggtac 180
gtggacggcg tggaaagtga caacgccaag accaagccta gagaggaaca gttcaactcc 240
acctacagag tgggtgtccct gctgacctgt ctgcaaccag attggctgaa cggcaaaagag 300
tacaagtcca aggtgtccaa caagggcctg ccttccagca tcgaaaagac catctctaag 360
gctaagggcc agcctcgcca gcccagggtt tacacattgc ctccaagcca agaggaaatg 420
accaagaacc aggtgtccct gacctgcctc gtgaagggtc tctaccttc cgacattgcc 480
gtggaatggg agtccaatgg ccagcctgag aacaactaca agacaacccc tctgtgtctg 540
gactccgacg gctccttctt tctgtattcc cgcctgacct tggacaagtc taggtggcaa 600
gagggcaacg tctctctctt ctctgtgtct cagcaggccc tgcacaatca ctacaccag 660
aagtctctgt ctctgtccct tggcggaggc ggaggatccg ctctacctc cagctccacc 720
aagaaaacc agttgcagct ggaacatctg ctgtgtggacc tccagatgat cctgaatggc 780
atcaacaatt acaagaaccc caagctgacc gccatgtgta ccgctaagtt ctacatgcc 840
aagaaggcca ccgagctgaa gcaacctccag tgcctgggaa gggaaactgaa gcctctggaa 900
gagggtgtga atctggccca gtccaagaac ttccacctga ggctcgggga cctgatctcc 960
aacatcaacg tgatctgtgt ggaactcaag ggctccgaga caacctcat gtgcaggtac 1020
gccgacgaga cagctaccat cgtggaattt ctgaaatcgt ggatcacctt ctgccagttc 1080
atcatcagca cctcgacc 1098

```

The invention claimed is:

1. A fusion protein comprising: an antigen-binding frag- 45
ment of an antibody that specifically binds to CD73; and an
IL-2 variant,

wherein the fusion protein comprises the following struc-
tural formula (I):



wherein

N' is the N-terminus,

C' is the C-terminus,

X is the antigen-binding fragment of an antibody that 55
specifically binds to CD73,

Y is the IL-2 variant that comprises the sequence of SEQ
ID NO: 40 with a modification, wherein the modifica-
tion is one or more substitutions selected from the
group consisting of R38A, F42A, and E61R with 60
regard to the amino acid sequence of SEQ ID NO: 40,
1st linker and 2nd linker are each a peptide linker, and
o and p are each independently 0 or 1,

wherein the antigen-binding fragment comprises a single-
domain antibody comprising a CDR1 region compris- 65
ing the amino acid sequence of SEQ ID NOs: 15, 18,
or 21; a CDR2 region comprising the amino acid

sequence of SEQ ID NOs: 16, 19, or 22; and a CDR3
region comprising the amino acid sequence of SEQ ID
NOs: 17, 20, or 23, respectively.


2. The fusion protein of claim 1, wherein the antigen-
binding fragment comprises any one amino acid sequence
selected from the group consisting of SEQ ID NO: 2, SEQ
ID NO: 3, and SEQ ID NO: 4.

3. The fusion protein of claim 1, wherein the 1st linker
comprises the amino acid sequence of SEQ ID NO: 5.

4. The fusion protein of claim 1, wherein the 2nd linker
comprises the amino acid sequence of SEQ ID NO: 7.

5. A fusion protein comprising: an antigen-binding frag-
ment of an antibody that specifically binds to CD73; and an
IL-2 variant,

wherein the fusion protein comprises a polypeptide chain
of the following structural formula (II) and a polypep-
tide chain of the following structural formula (III):



wherein, in the structural formulas (II) and (III),

N' is the N-terminus,

C' is the C-terminus,

55

X is the antigen-binding fragment comprising a heavy chain variable region (VH) and a heavy chain constant region 1 (CH1),

X' is the antigen-binding fragment comprising a light chain variable region (VL) and a light chain constant region (CL),

Y is the IL-2 variant that comprises the sequence of SEQ ID NO: 40 with a modification, wherein the modification is one or more substitutions selected from the group consisting of R38A, F42A, and E61R with regard to the amino acid sequence of SEQ ID NO: 40, 3rd linker and 4th linker are each a peptide linker, and q and r are each independently 0 or 1,

wherein the heavy chain variable region (VH) comprises an HCDR1 comprising the amino acid sequence of SEQ ID NO: 24, an HCDR2 comprising the amino acid sequence of SEQ ID NO: 25, and an HCDR3 comprising the amino acid sequence of SEQ ID NO: 26; and the light chain variable region (LH) comprises an LCDR1 comprising the amino acid sequence of SEQ ID NO: 27, an LCDR2 comprising the amino acid sequence of SEQ ID NO: 28, and an LCDR3 comprising the amino acid sequence of SEQ ID NO: 29, and

56

wherein the polypeptide chain of the structural formula (II) and the polypeptide chain of the structural formula (III) form a polypeptide heterodimer.

6. The fusion protein of claim 5, wherein the heavy chain variable region comprises the amino acid sequence of SEQ ID NO: 9, and the light chain variable region comprises the amino acid sequence of SEQ ID NO: 13.

7. The fusion protein of claim 5, wherein the linker 3rd linker comprises the amino acid sequence of SEQ ID NO: 11.

8. The fusion protein of claim 5, wherein the 4th linker comprises the amino acid sequence of SEQ ID NO: 12.

9. The fusion protein of claim 1, wherein the Fc region comprises the amino acid sequence of SEQ ID NO: 6.

10. A fusion protein dimer comprising two of the fusion protein of claim 1, wherein the two fusion proteins are linked to each other.

11. The fusion protein of claim 5, wherein the Fc region comprises the amino acid sequence of SEQ ID NO: 6.

12. A method for treating cancer comprising administering a pharmaceutical composition comprising, as an active ingredient, the fusion protein of claim 1 or a dimer thereof to a subject in need thereof.

* * * * *