

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent

12417834

Kind Code

B2

Date of Patent

September 16, 2025

Inventor(s)

Lomas; Sarah et al.

Genetically personalized intravenous and intramuscular nutrition therapy design systems and methods

Abstract

A method of providing genetically personalized intravenous or intramuscular nutrition therapy includes developing an initial mapping of individual micronutrients to SNPs, symptoms, conditions, and therapeutic objectives based on existing individual scientific literature references that link the specific micronutrient to an SNP, a symptom, a condition, or a therapeutic objective. The method includes receiving an SNP and a symptom, a condition, or a therapeutic objective for a patient, comparing the SNP, symptom, condition, therapeutic objective combination with the initial mapping to identify matching micronutrients, determining a nutrition therapy based on the identified micronutrients, and treating the patient accordingly. Feedback data about the nutrition therapy's effectiveness is received and the initial mapping is updated to account for the feedback data. This process is repeated for subsequent patients positioned geographically remotely from each other so that future patients are treated with reference to an updated current mapping.

Inventors: Lomas; Sarah (Cheshire, GB), Barnish; Michael (Cheshire, GB), Parvani; Johnny (Phoenix, AZ)

Applicant: REVIV GLOBAL LTD (Knutsford, GB)

Family ID: 1000008819645

Assignee: REVIV GLOBAL LTD (Knutsford, GB)

Appl. No.: 17/876352

Filed: July 28, 2022

Prior Publication Data

Document Identifier

Publication Date

US 20230032427 A1

Feb. 02, 2023

Related U.S. Application Data

Publication Classification**Int. Cl.:** **G16H20/60** (20180101); **G16H10/60** (20180101); **G16H20/10** (20180101); **G16H50/30** (20180101)**U.S. Cl.:****CPC** **G16H20/60** (20180101); **G16H10/60** (20180101); **G16H20/10** (20180101); **G16H50/30** (20180101)**Field of Classification Search****CPC:** G16H (20/60); G16H (10/60); G16H (20/10); G16H (50/30)**USPC:** 705/2-3

References Cited**U.S. PATENT DOCUMENTS**

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
8560334	12/2012	Pertti	N/A	N/A
10929916	12/2020	Abutair	N/A	N/A
2002/0058712	12/2001	Sneed	N/A	N/A
2005/0177397	12/2004	Kane	N/A	N/A
2006/0062859	12/2005	Blum	N/A	N/A
2006/0199155	12/2005	Mosher	N/A	N/A
2007/0154498	12/2006	Bortz	N/A	N/A
2008/0050740	12/2007	Cassidy	N/A	N/A
2008/0221932	12/2007	Kane	N/A	N/A
2009/0155381	12/2008	Goralczyc	N/A	N/A
2009/0192365	12/2008	Gisel	N/A	N/A
2011/0113002	12/2010	Kane	N/A	N/A
2011/0189161	12/2010	Blum	N/A	N/A
2012/0083669	12/2011	Abujbara	N/A	N/A
2012/0290327	12/2011	Hanlon	N/A	N/A
2013/0018024	12/2012	Bianchi	N/A	N/A
2013/0216982	12/2012	Bennett	N/A	N/A
2014/0052722	12/2013	Bertsimas	N/A	N/A
2014/0088995	12/2013	Damani	N/A	N/A
2015/0125462	12/2014	Bek	N/A	N/A
2015/0269865	12/2014	Volach	434/127	G09B 19/0092
2018/0032682	12/2017	Donalds	N/A	N/A
2018/0039759	12/2017	Astigarraga	N/A	N/A
2018/0089385	12/2017	Gupta	N/A	N/A
2018/0121631	12/2017	Mehta	N/A	N/A
2018/0144820	12/2017	Grimmer	N/A	N/A

2018/0353425	12/2017	Narain	N/A	N/A
2018/0374567	12/2017	Toumazou	N/A	N/A
2019/0152663	12/2018	Kraft	N/A	N/A
2019/0221303	12/2018	Bennett	N/A	N/A
2019/0290172	12/2018	Hadad	N/A	N/A
2020/0135314	12/2019	Gostyla	N/A	N/A
2021/0005304	12/2020	Neumann	N/A	G16H 40/67
2021/0005317	12/2020	Neumann	N/A	N/A
2021/0012880	12/2020	Ando	N/A	N/A
2021/0050086	12/2020	Rose	N/A	N/A
2021/0134434	12/2020	Riley	N/A	N/A
2021/0166794	12/2020	Nova	N/A	N/A
2021/0193332	12/2020	Inwald	N/A	N/A

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
3014239	12/2019	CA	N/A
106326683	12/2016	CN	N/A
112069382	12/2019	CN	N/A
3529379	12/2021	EP	N/A
20170054628	12/2016	KR	N/A
102043959	12/2018	KR	N/A
20190138360	12/2018	KR	N/A
102169661	12/2019	KR	N/A
20210061032	12/2020	KR	N/A
20210072221	12/2020	KR	N/A
102278646	12/2020	KR	N/A
2004084822	12/2003	WO	N/A
2012067525	12/2011	WO	N/A
2013086582	12/2012	WO	N/A
2015004266	12/2014	WO	N/A
2015124569	12/2014	WO	N/A
2019183404	12/2018	WO	N/A
WO-2019183404	12/2018	WO	A61B 5/0022
2020138720	12/2019	WO	N/A

OTHER PUBLICATIONS

Verkaik-Kloosertman, Janneke; “Estimation of micronutrient intake distributions: Development of methods to support food and nutrition policy making;” Wageningen University, 2011. (Year: 2011). cited by examiner

Chen, Yu et al. Personalized Food Recommendation as Constrained Question Answering over a Large-scale Food Knowledge Graph. Rensselaer Polytechnic Institute, Troy, NY. 2021. 9 pages. cited by applicant

Drabsch, T. et al. A Scientific Perspective of Personalised Gene-Based Dietary Recommendations for Weight Management. Nutrients. 2019, 14 pages. cited by applicant

Ingilizian, Z. How Precision Consumption Can Improve Consumer Health and Well-Being. Forbes. Leadership Strategy. 2020, 7 pages. cited by applicant

Lubos, E. et al. Glutathione Peroxidase-1 in Health and Disease: From Molecular Mechanisms to

Therapeutic Opportunities. Antioxid Redox Signal. 2011, 42 pages. cited by applicant

NutriGenomeDB platform, 1 page, <http://www.nutrigenomedb.org/>. cited by applicant

Precision Consumer 2030 when data becomes invisible. Sparks & Honey. Culture Forecast. 2019, 125 pages. cited by applicant

Galyean, S. et al., ‘Personalized Nutrition for Management of Micronutrient Deficiency-Literature Review in Non-bariatric Populations and Possible Utility in Bariatric Cohort’, Obesity Surgery, 2020, vol. 30, pp. 3570-3582. cited by applicant

Morine, M. J. et al., ‘Genetic associations with micronutrient levels identified in immune and gastrointestinal networks’, Genes Nutr., 2014, vol. 9, No. 408, pp. 1-19. cited by applicant

Verkaik-Kloosterman, Janneke; Estimation of Micronutrient Intake Distributions: Development of Methods to Support Food and Nutrition Policy Making; Wageningen University and Research. ProQuest Dissertations Publishing, 2011. 28235724 (Year: 2011) 192 pages. cited by applicant

Brouwer ID et al: “Reverse thinking: taking a healthy diet perspective towards food systems transformations”, Food Security, Springer Netherlands, Dordrecht, vol. 13, No. 6, Oct. 16, 2021 (Oct. 16, 2021), pp. 1497-1523, XP037646513, ISSN: 1876-4517, DOI: 10.1007/SI2571-021-01204-5 [retrieved on Oct. 16, 2021]. cited by applicant

Bujari Armir et al: “A mobile sensing and visualization platform for environmental data”, Pervasive and Mobile Computing, Elsevier, NL, vol. 66, Jun. 18, 2020 (Jun. 18, 2020), XP086246547, ISSN: 1574-1192, DOI: 10.1016/J.PMCJ.2020.101204 [retrieved on Jun. 18, 2020]. cited by applicant

Maestre Mar et al: “Assessing food value chain pathways, linkages and impacts for better nutrition of vulnerable groups”, Food Policy, Pergamon, Amsterdam, NL, vol. 68, Jan. 11, 2017 (Jan. 11, 2017), pp. 31-39, XP029956443, ISSN: 0306-9192, DOI: 10.1016/J.FOODPOL.2016.12.007 p. 31-p. 36; figure 1. cited by applicant

Matthew Metzgar et al: “The feasibility of a Paleolithic diet for low-income consumers”, Nutrition Research, Elsevier, Amsterdam, NL, vol. 31, No. 6, May 14, 2011 (May 14, 2011), pp. 444-451, XP028379237, ISSN: 0271-5317, DOI: 10.1016/J.NUTRES.2011.05.008 [retrieved on May 23, 2011]. cited by applicant

Poole Nigel et al: “Viewpoint: Agri-nutrition research: Revisiting the contribution of maize and wheat to human nutrition and health”, Food Policy, Pergamon, Amsterdam, NL, vol. 100, Sep. 18, 2020 (Sep. 18, 2020), XP086547134, ISSN: 0306-9192, DOI: 10.1016/J.FOODPOL.2020.101976 [retrieved on Sep. 18, 2020]. cited by applicant

William a Masters et al: “The economics of malnutrition: Dietary transition and food system transformation”, arxiv.org, Cornell University Library, 201 Olin Library Cornell University Ithaca, NY 14853, Feb. 5, 2022 (Feb. 5, 2022), XP091150658, abstract; table 1 p. 97. cited by applicant

Berthon et al., “Nutrition and Respiratory Health-Feature Review”, Mar. 5, 2015, Nutrients, vol. 7, 1618-1643. (Year: 2015). cited by applicant

Primary Examiner: Dunham; Jason B

Assistant Examiner: Covington; Amanda R.

Attorney, Agent or Firm: Booth Udall, PLC

Background/Summary

RELATED APPLICATIONS (1) This application claims the benefit of U.S. Provisional Application No. 63/227,943, filed Jul. 30, 2021, the disclosure of which is hereby incorporated by reference in its entirety.

TECHNICAL FIELD

(1) This disclosure relates generally to intravenous (“IV”) and intramuscular (“IM”) nutrition therapy, and more specifically to IV and IM nutrition therapy that is personalized to the recipient based on genetic data. The IV and IM nutrition therapy may also or alternatively be personalized based on other medical data.

BACKGROUND

(2) Achieving and maintaining optimal health and proper function of the human body requires proper nutrition and hydration, including a correct balance of nutrients such as macronutrients (e.g., proteins, carbs, fats) and micronutrients (e.g., vitamins, minerals, antioxidants). An increasingly popular method of achieving this balance is through IV and IM delivery of the nutrients. IV and IM delivery are generally more efficient than oral intake of the same nutrients, leading to a more effective treatment. Currently, IV and IM nutrition therapies use generic formulations, with the same formulation used for each individual, regardless of the individual's specific needs. Because each individual is unique and has different requirements for nutrients, what is beneficial to one person may be less useful, or even detrimental, to another. Thus, current IV and IM nutrition therapies may provide excess unnecessary nutrients or omit vital ingredients, resulting in suboptimal supplementations and less effective therapy.

SUMMARY

(3) Aspects of this disclosure relate to a method of providing genetically personalized intravenous or intramuscular nutrition therapy, the method comprising developing an initial mapping of each of a plurality of individual micronutrients to a single nucleotide polymorphism (SNP) and to at least one of a symptom, a condition, and a therapeutic objective, the developing the initial mapping comprising, for each specific micronutrient of the plurality of individual micronutrients: creating an individual score for each of a plurality of existing individual scientific literature references that links the specific micronutrient to an SNP, a symptom, a condition, or a therapeutic objective; adding the individual scores for each of the plurality of existing individual scientific literature references for the specific micronutrient with respect to each combination of an SNP and at least one of a symptom, a condition, and a therapeutic objective; and determining that the specific micronutrient matches the SNP and the at least one of a symptom, a condition, and a therapeutic objective when the individual scores for the specific micronutrient with respect to the SNP and the at least one of a symptom, a condition, and a therapeutic objective add up to more than a predetermined threshold; storing the initial mapping, the individual scores, and the added scores in a database; receiving an SNP for a first patient and at least one of a symptom, a condition, and a therapeutic objective for the first patient; comparing the SNP for the first patient and the at least one of a symptom, a condition, and a therapeutic objective for the first patient with the initial mapping stored in the database to identify one or more micronutrients that have been matched to the SNP for the first patient and the at least one of a symptom, a condition, and a therapeutic objective for the first patient; determining an intravenous or intramuscular nutrition therapy for the first patient based on the identified one or more micronutrients from the initial mapping, the intravenous or intramuscular nutrition therapy comprising an existing intravenous or intramuscular nutrition therapy formula that comprises a specific dosage of at least one of the identified one or more micronutrients; treating the first patient with the intravenous or intramuscular nutrition therapy; receiving feedback data about an effectiveness of the intravenous or intramuscular nutrition therapy in addressing the at least one of a symptom, a condition and a therapeutic objective for the first patient; creating an individual treatment score for the at least one of the identified one or more micronutrients in the intravenous or intramuscular nutrition therapy with respect to the SNP for the first patient and the at least one of a symptom, a condition, and a therapeutic objective for the first patient based on the feedback data; updating the initial mapping by: adding the individual treatment score for the at least one of the identified one or more

micronutrients in the intravenous or intramuscular nutrition therapy to the added scores stored in the database that are associated with the at least one of the identified one or more micronutrients, the SNP for the first patient, and the at least one of a symptom, a condition, and a therapeutic objective for the first patient; comparing the updated added scores with the predetermined threshold to reevaluate whether there are matches for the at least one of the identified one or more micronutrients in the intravenous or intramuscular nutrition therapy; and storing the updated initial mapping, the individual treatment score, and the updated added scores in the database; and for each of a plurality of subsequent patients positioned geographically remotely from each other: receiving an SNP of the subsequent patient and at least one of a symptom, a condition, and a therapeutic objective of the subsequent patient; comparing the SNP for the subsequent patient and the at least one of a symptom, a condition, and a therapeutic objective for the subsequent patient with a current mapping stored in the database to identify one or more micronutrients that have been matched to the SNP for the subsequent patient and the at least one of a symptom, a condition, and a therapeutic objective for the subsequent patient, wherein the current mapping has been updated with the individual treatment score of the first patient and individual treatment scores of any other previously treated patient; determining an intravenous or intramuscular nutrition therapy for the subsequent patient based on the identified one or more micronutrients from the current mapping, the intravenous or intramuscular nutrition therapy comprising an existing intravenous or intramuscular nutrition therapy formula that comprises a specific dosage of at least one of the identified one or more micronutrients; treating the subsequent patient with the intravenous or intramuscular nutrition therapy; receiving feedback data about an effectiveness of the intravenous or intramuscular nutrition therapy in addressing the at least one of a symptom, a condition and a therapeutic objective for the subsequent patient; creating an individual treatment score for the at least one of the identified one or more micronutrients in the intravenous or intramuscular nutrition therapy with respect to the SNP for the subsequent patient and the at least one of a symptom, a condition, and a therapeutic objective for the subsequent patient based on the feedback data; updating the current mapping by: adding the individual treatment score for the at least one of the identified one or more micronutrients in the intravenous or intramuscular nutrition therapy for the subsequent patient to current added scores stored in the database that are associated with the at least one of the identified one or more micronutrients, the SNP for the subsequent patient, and the at least one of a symptom, a condition, and a therapeutic objective for the subsequent patient; comparing the updated current added scores with the predetermined threshold to reevaluate whether there are matches for the at least one of the identified one or more micronutrients in the intravenous or intramuscular nutrition therapy for the subsequent patient; and storing the updated current mapping, the individual treatment score of the subsequent patient, and the updated current scores in the database; and treating future patients with reference to the updated current mapping.

(4) Particular implementations may comprise one or more of the following features. Updating the current mapping based on individual scores for each of a plurality of new individual scientific literature references. A reference of the plurality of existing individual scientific literature references that is statistically significant and positively links a micronutrient to an SNP, a symptom, a condition, or a therapeutic objective is assigned an individual score greater than zero. A reference of the plurality of existing individual scientific literature references that is statistically significant and negatively links a micronutrient to an SNP, a symptom, a condition, or a therapeutic objective is assigned an individual score less than zero. A reference of the plurality of existing individual scientific literature references that relates to an injectable outcome is assigned an individual score that has more weight than an individual score of a reference of the plurality of existing individual scientific literature references that relates to an oral outcome. A reference of the plurality of existing individual scientific literature references that is not statistically significant is assigned an individual score less than zero.

(5) Aspects of this disclosure relate to a method of providing genetically personalized intravenous

or intramuscular nutrition therapy, the method comprising: (a) creating a patient treatment nutrition therapy recommendation algorithm and storing the patient treatment nutrition therapy recommendation algorithm in a database; (b) receiving medical information, genetic information, and therapeutic objectives of a patient as inputs for the patient treatment nutrition therapy recommendation algorithm, wherein the genetic information comprises genetic test results and the medical information comprises blood test results, vital signs, medical histories, medical diagnoses, current symptoms, diseases of which the patient is at heightened risk, or family history; (c) outputting a therapy recommendation from the patient treatment nutrition therapy recommendation algorithm based on the medical information, genetic information, and therapeutic objectives, wherein the therapy recommendation comprises an existing intravenous or intramuscular nutrition therapy formula having at least one micronutrient, and wherein the at least one micronutrient is identified by the patient treatment nutrition therapy recommendation algorithm as a match for the medical information, genetic information, and therapeutic objectives based on individual scores stored in the database of a plurality of existing individual scientific literature references relating to the at least one micronutrient and individual treatment scores stored in the database of previous intravenous or intramuscular nutrition therapy treatments comprising the at least one micronutrient; (d) treating the patient with an intravenous or intramuscular nutrition therapy according to the therapy recommendation; (e) receiving feedback data about results of the intravenous or intramuscular nutrition therapy of the patient, wherein the feedback data comprises feedback about measured bloodwork, epigenetic changes, improvements or declines in symptoms, progress towards reaching therapeutic objectives, vital signs, overall health and fitness, or physical or mental energy; (f) creating an individual treatment score for the intravenous or intramuscular nutrition therapy of the patient based on the feedback data and entering the individual treatment score electronically into a computer system associated with the database so that the individual treatment score is stored in the database; (g) updating the patient treatment nutrition therapy recommendation algorithm to incorporate the individual treatment score for the intravenous or intramuscular nutrition therapy of the patient for subsequent patients; (h) creating future patient treatment nutrition therapy recommendations based on the updated patient treatment nutrition therapy recommendation algorithm, wherein the updated patient treatment nutrition therapy recommendation algorithm is configured to recommend treatment and micronutrients that most closely match patient needs based on all available patient medical information, genetic information, and therapeutic objectives, and most current aggregated feedback based on real patients with similar genes and symptoms; and (i) repeating steps (b)-(h) for subsequent patients to create a feedback loop of automatically applying patient therapy feedback data to the patient treatment nutrition therapy recommendation algorithm so that all new therapy recommendations include the feedback data received from all previous therapy recommendation patients who received therapy recommendations through the patient treatment nutrition therapy recommendation algorithm, wherein at least some of the patient and the subsequent patients are positioned geographically remotely from each other.

(6) Particular implementations may comprise one or more of the following features. The therapeutic objectives of the patient may comprise at least one of health goals, aesthetic goals, fitness goals, weight loss, increased energy, healthier skin, healthier hair, and improved mental acuity. The patient treatment nutrition therapy recommendation algorithm may be configured to map the medical information and the genetic information to a code. The code may comprise a selection from the International Classification of Diseases. The patient treatment nutrition therapy recommendation algorithm comprises multiple algorithms in sequence. The medical information comprises at least one current symptom and at least one condition. The medical information of the patient comprises a first symptom and the genetic information of the patient comprises a first gene, wherein medical information of at least one of the subsequent patients comprises the first symptom and genetic information of the at least one of the subsequent patients comprises a second gene

different than the first gene, and wherein the at least one micronutrient of the therapy recommendation for the patient comprises a first micronutrient and a therapy recommendation for the at least one of the subsequent patients comprises a second micronutrient different than the first micronutrient.

(7) Aspects of this disclosure relate to a system for providing genetically personalized intravenous or intramuscular nutrition therapy, the system comprising: a plurality of genetic testing kits, each testing kit configured to determine genetic information of one of a plurality of patients, the plurality of patients positioned geographically remotely from each other; a plurality of electronic devices, each electronic device configured to receive the genetic information of one of the plurality of patients, medical information of the one of the plurality of patients, and therapeutic objectives of the one of the plurality of patients; and a server comprising a database storing a patient treatment nutrition therapy recommendation algorithm, wherein the server is configured to wirelessly receive the genetic information, the medical information, and the therapeutic objectives of each of the plurality of patients, wherein the server is configured to, for each of the plurality of patients: output to the electronic device a therapy recommendation for the patient from the patient treatment nutrition therapy recommendation algorithm based on the medical information, genetic information, and therapeutic objectives of the patient, wherein the therapy recommendation comprises an existing intravenous or intramuscular nutrition therapy formula having at least one micronutrient, and wherein the at least one micronutrient is identified by the patient treatment nutrition therapy recommendation algorithm as a match for the medical information, genetic information, and therapeutic objectives of the patient based on individual scores stored in the database of a plurality of existing individual scientific literature references relating to the at least one micronutrient and individual treatment scores stored in the database of previous intravenous or intramuscular nutrition therapy treatments comprising the at least one micronutrient; receive feedback data about results of the therapy recommendation after the patient has been treated based on the therapy recommendation, wherein the feedback data comprises feedback about measured bloodwork, epigenetic changes, improvements or declines in symptoms, progress towards reaching therapeutic objectives, vital signs, overall health and fitness, or physical or mental energy; create an individual treatment score based on the feedback data and store the individual treatment score in the database; update the patient treatment nutrition therapy recommendation algorithm to incorporate the individual treatment score for subsequent patients; and create future patient treatment nutrition therapy recommendations based on the updated patient treatment nutrition therapy recommendation algorithm, wherein the updated patient treatment nutrition therapy recommendation algorithm is configured to recommend treatment and micronutrients that most closely match patient needs based on all available patient medical information, genetic information, and therapeutic objectives, and most current aggregated feedback based on real patients with similar genes and symptoms.

(8) Particular implementations may comprise one or more of the following features. The therapeutic objectives comprise health goals, aesthetic goals, fitness goals, weight loss, increased energy, healthier skin, healthier hair, or improved mental acuity. The patient treatment nutrition therapy recommendation algorithm is configured to map the medical information and the genetic information to a code. The code comprises a selection from the International Classification of Diseases. The patient treatment nutrition therapy recommendation algorithm comprises multiple algorithms in sequence. The medical information comprises at least one current symptom and at least one condition. The medical information of a first patient of the plurality of patients comprises a first symptom and the genetic information of the first patient comprises a first gene, wherein the medical information of a second patient of the plurality of patients comprises the first symptom and the genetic information of the second patient comprises a second gene different than the first gene, and wherein the at least one micronutrient of the therapy recommendation for the first patient comprises a first micronutrient and the at least one micronutrient of the therapy recommendation

for the second patient comprises a second micronutrient different than the first micronutrient.

(9) The foregoing and other aspects, features, applications, and advantages will be apparent to those of ordinary skill in the art from the specification, drawings, and the claims. Unless specifically noted, it is intended that the words and phrases in the specification and the claims be given their plain, ordinary, and accustomed meaning to those of ordinary skill in the applicable arts. The inventors are fully aware that he can be his own lexicographer if desired. The inventors expressly elect, as their own lexicographers, to use only the plain and ordinary meaning of terms in the specification and claims unless they clearly state otherwise and then further, expressly set forth the “special” definition of that term and explain how it differs from the plain and ordinary meaning. Absent such clear statements of intent to apply a “special” definition, it is the inventors' intent and desire that the simple, plain and ordinary meaning to the terms be applied to the interpretation of the specification and claims.

(10) The inventors are also aware of the normal precepts of English grammar. Thus, if a noun, term, or phrase is intended to be further characterized, specified, or narrowed in some way, then such noun, term, or phrase will expressly include additional adjectives, descriptive terms, or other modifiers in accordance with the normal precepts of English grammar. Absent the use of such adjectives, descriptive terms, or modifiers, it is the intent that such nouns, terms, or phrases be given their plain, and ordinary English meaning to those skilled in the applicable arts as set forth above.

(11) Further, the inventors are fully informed of the standards and application of the special provisions of 35 U.S.C. § 112(f). Thus, the use of the words “function,” “means” or “step” in the Detailed Description or Description of the Drawings or claims is not intended to somehow indicate a desire to invoke the special provisions of 35 U.S.C. § 112(f), to define the invention. To the contrary, if the provisions of 35 U.S.C. § 112(f) are sought to be invoked to define the inventions, the claims will specifically and expressly state the exact phrases “means for” or “step for”, and will also recite the word “function” (i.e., will state “means for performing the function of [insert function]”), without also reciting in such phrases any structure, material or act in support of the function. Thus, even when the claims recite a “means for performing the function of . . .” or “step for performing the function of . . .,” if the claims also recite any structure, material or acts in support of that means or step, or that perform the recited function, then it is the clear intention of the inventors not to invoke the provisions of 35 U.S.C. § 112(f). Moreover, even if the provisions of 35 U. S.C. § 112(f) are invoked to define the claimed aspects, it is intended that these aspects not be limited only to the specific structure, material or acts that are described in the preferred embodiments, but in addition, include any and all structures, materials or acts that perform the claimed function as described in alternative embodiments or forms of the disclosure, or that are well known present or later-developed, equivalent structures, material or acts for performing the claimed function.

(12) The foregoing and other aspects, features, and advantages will be apparent to those of ordinary skill in the art from the specification, drawings, and the claims.

Description

BRIEF DESCRIPTION OF THE DRAWINGS

- (1) The disclosure will hereinafter be described in conjunction with the appended drawings.
- (2) FIG. 1 shows a process flow for providing genetically personalized IV or IM nutrition therapy according to some embodiments.
- (3) FIG. 2 shows a process flow for developing an initial mapping of micronutrients to genetic information, symptoms, conditions, and therapeutic objectives according to some embodiments.
- (4) FIG. 3 shows a representative initial mapping of micronutrients to genetic information,

- symptoms, conditions, and therapeutic objectives according to some embodiments.
- (5) FIG. 4 shows a process flow for updating a mapping of micronutrients to genetic information, symptoms, conditions, and therapeutic objectives according to some embodiments.
- (6) FIG. 5 shows a schematic of providing genetically personalized IV or IM nutrition therapy according to some embodiments.
- (7) FIG. 6 shows a schematic of providing genetically personalized IV or IM nutrition therapy to target a specific disease according to some embodiments.
- (8) FIG. 7 shows a process flow for providing genetically personalized IV or IM nutrition therapy according to some embodiments.
- (9) FIG. 8 shows a schematic representing a nutritional database according to some embodiments.
- (10) FIG. 9 shows example ingredients and formulas of a genetically personalized IV or IM nutrition therapy according to some embodiments.
- (11) FIG. 10 shows a schematic representing a holistic approach to implementing a genetically personalized IV or IM nutrition therapy according to some embodiments.
- (12) FIG. 11 shows a genetically personalized nutrition therapy system according to some embodiments.
- (13) FIG. 12 shows a genetically personalized nutrition therapy system according to some embodiments.
- (14) FIG. 13 shows a schematic diagram of computing devices that can be used to perform or implement the embodiments disclosed herein.

DETAILED DESCRIPTION

- (15) This disclosure, its aspects and implementations, are not limited to the specific components, methods, or other examples disclosed herein. Many additional components, methods, and procedures known in the art are contemplated for use with particular implementations from this disclosure. Accordingly, for example, although particular implementations are disclosed, such implementations and implementing components may comprise any components, models, types, materials, versions, quantities, and/or the like as is known in the art for such systems and implementing components, consistent with the intended operation.
- (16) The word “exemplary,” “example,” or various forms thereof are used herein to mean serving as an example, instance, or illustration. Any aspect or design described herein as “exemplary” or as an “example” is not necessarily to be construed as preferred or advantageous over other aspects or designs. Furthermore, examples are provided solely for purposes of clarity and understanding and are not meant to limit or restrict the disclosed subject matter or relevant portions of this disclosure in any manner. It is to be appreciated that a myriad of additional or alternate examples of varying scope could have been presented but have been omitted for purposes of brevity.
- (17) While this disclosure includes a number of implementations in many different forms, the drawings show particular implementations that will be described in detail with the understanding that the present disclosure is to be considered as an exemplification of the principles of the disclosed methods and systems and is not intended to limit the broad aspect of the disclosed concepts to the implementations illustrated.
- (18) Achieving and maintaining optimal health and proper function of the human body requires proper nutrition and hydration, including a correct balance of nutrients such as macronutrients (e.g., proteins, carbs, fats) and micronutrients (e.g., vitamins, minerals, electrolytes, antioxidants). An increasingly popular method of achieving this balance is through IV and IM delivery of the nutrients. IV and IM delivery are generally more efficient than oral intake of the same nutrients, leading to a more effective treatment. Currently, IV and IM nutrition therapies use generic formulations, with the same formulation used for each individual, regardless of the individual's specific needs. Because each individual is unique and has different requirements for nutrients, what is beneficial to one person may be less useful, or even detrimental, to another. Thus, current IV and IM nutrition therapies may provide excess unnecessary nutrients or omit vital ingredients, resulting

in suboptimal supplementations and less effective therapy.

(19) The present disclosure describes systems and methods that provide genetically personalized IV and IM nutrition therapies. In some embodiments, the nutrition therapies are also or alternatively personalized based on other medical data. Because the therapy disclosed herein is personalized for each individual recipient, it may be uniquely designed to meet each individual's nutritional needs. As will be described in more detail below, these individual needs may be determined based on genetic data and test results, medical history, and/or medical data and test results, among other things. Thus, each recipient of the personalized therapy can receive the correct balance and amount of nutrients, including macronutrients (e.g., proteins, carbs, fats) and micronutrients (e.g., vitamins, minerals, electrolytes, antioxidants). Some implementations may focus on only some of these nutrients and may include nutrients not listed as well.

(20) In some embodiments, the systems and methods disclosed herein facilitate determining which nutrients are needed for certain combinations of genetic information and medical information. For example, micronutrients may be matched to certain genetic information (e.g., a gene, a single nucleotide polymorphism ("SNP"), a genotype, etc.), symptoms, conditions, therapeutic objectives, and/or combinations of these attributes based on existing scientific literature. The matching of micronutrients to genetic information, symptoms, conditions, and/or therapeutic objectives may be repeatedly updated based on outcomes of the nutrition therapies, which may facilitate nutrition therapies to become more and more personalized to a patient's specific situation.

(21) Updating micronutrient matches is not limited to a single patient's past nutrition therapy, nor is it limited to patients of a single medical professional or healthcare provider. Rather, the methods and systems allow for updating micronutrient matches for any patients that will receive nutrition therapy in the future as soon as an outcome of a nutrition therapy is received by the system, even if these patients are positioned geographically remotely from each other (i.e., they are treated in different parts of the world (e.g., different neighborhood, city, county, state, country, etc.) or by different healthcare providers). Thus, the systems and methods may recommend treatment and micronutrients that most closely match patient needs based on all available patient medical information, genetic information, and therapeutic objectives, and most current aggregated feedback based on real patients with similar genes and symptoms. In some embodiments, the methods and systems create a feedback loop of automatically applying patient therapy feedback data so that all new therapy recommendations include the feedback data received from all previous therapy recommendation patients who received therapy recommendations through the system.

(22) Recipients of the personalized therapy disclosed herein may experience improved disease prevention and treatment, prevention of aging, maintenance of metabolic function, and improved growth and development as a result of the personalized therapy. Each of the components discussed herein may be implemented alone, or in conjunction with each of the other described components.

(23) In some embodiments, IV and IM therapy may be personalized by collecting medical and genetic data regarding the recipient, analyzing the medical and genetic data to determine the nutritional needs of the recipient (e.g., based on a nutritional formulary mapping nutrients to various medical and genetic data), selecting a formula from a plurality of existing formulas or nutritional ingredients based on the nutritional needs of the recipient, administering the selected formula to the recipient through IV or IM treatment, collecting feedback from the recipient regarding the effect of the IV or IM treatment, and adjusting the nutritional formulary to incorporate the feedback provided.

(24) Any medical and genetic data can be relevant. Therefore, to provide the personalized therapies disclosed herein, the data collected may include, but is not limited to, blood test results, genetic test results, vital signs (e.g., blood pressure, heart rate, heart rhythm, etc.), biometric data, data received from wearable devices, lifestyle data (such as travel and recent infections or vaccinations), medical histories, medical diagnoses, diseases of which the recipient is at heightened risk, and current symptom descriptions. In addition, the data may include specific therapeutic objectives of the

recipient, such as health goals, aesthetic goals, fitness goals, weight loss, increased energy, healthier skin or hair, or improved mental acuity (or others discussed below). As more data is included, the personalized therapy can be adapted to be more comprehensive and accurate in providing the nutritional needs for the recipient, and those same findings can be applied to improve personalization therapies for other recipients with similar conditions as well.

(25) The medical and genetic data may be provided directly by the recipient or may be determined using medical and lab procedures. In some embodiments, some of the medical and genetic data is provided directly by the recipient and some of the medical and genetic data is determined using medical and lab procedures. For example, the recipient may provide a family medical history and a description of current symptoms, such as fatigue or headaches, while medical tests and procedures may be used to perform blood tests and genetic tests, as well as provide a medical diagnosis.

(26) In some embodiments, the collected data may be mapped into a code to facilitate the interpretation of the data and enable its use in algorithms and automated systems. For example, an international classification system may be used, such as the International Statistical Classification of Diseases and Related Health Problems (ICD) implemented by the World Health Organization (WHO). The most current ICD code is the ICD-11. Any system of classification of diseases, symptoms, and genes may be implemented.

(27) The collected data may be analyzed to determine the nutritional needs of the recipient. In some implementations of the personalized therapy disclosed herein, the collected data is analyzed by a life science professional who is familiar with the genes, diseases, and symptoms communicated by the data and understands what nutritional needs this data expresses. For example, experts in fields such as nutraceuticals, pharmacokinetics, pharmaceuticals, nutrition, nutritional science, nutrigenomics, biomedical sciences, food data sciences, and statistics may work together to review the collected data and determine which nutrients can help improve the health of the recipient.

(28) In other implementations, this analysis may be automated. For example, a nutritional formulary or a mapping may be created which links each gene, disease, symptom, and therapeutic objective with different nutritional needs (e.g., micronutrients). Additionally, the nutritional formulary may indicate nutritional needs for specific combinations of symptoms, diseases, genes, and therapeutic objectives. For example, a first recipient may have a first symptom and a first gene. The nutritional formulary may suggest a first micronutrient to provide what the first recipient needs. A second recipient may have the same first symptom, but a second gene. In some cases, the nutritional formulary may suggest the same first micronutrient in the same amount. However, in other cases, the specific combination of the first symptom with the second gene may necessitate suggesting a second micronutrient or a different amount of the first micronutrient. Thus, the nutritional formulary is configured to take into account all of the collected data in determining the nutritional needs of the recipient. The nutritional formulary may be stored in a nutritional database. All references herein to a nutritional formulary may be considered a reference to a nutritional database. Likewise, all references to a nutritional database may be considered a reference to a nutritional formulary. The nutritional formulary or nutritional database may also be referred to as a mapping (i.e., mapping micronutrients to genetic information, symptoms, conditions, therapeutic objectives, and combinations of these attributes). The nutritional formulary may be used by an automated process or may be used by the experts discussed above.

(29) Once the nutritional needs of the recipient have been determined, a formula may be selected for administration to the recipient from a plurality of preexisting formulas or ingredients. As mentioned above, the formula may be selected based on the nutritional needs of the recipient. For example, each of the preexisting formulas may be evaluated in comparison with the nutritional needs of the recipient, and the selected formula may be the formula which most closely meets the nutritional needs of the recipient. The selected formula may be administered to the recipient through an IV treatment or an IM treatment. Alternatively, the personalized therapy may also be administered orally or topically. It will be understood that any IV treatment referenced herein may

also be implemented as an IM treatment, an oral treatment, or a topical treatment. Additionally, the personalized therapy may be administered through a combination of different treatments. For example, the recipient may receive therapy through an IV treatment and an IM treatment, or any other combination. The recipient may travel to a clinic or other setting to receive the treatment.

(30) Feedback may be collected from the recipient regarding the recipient's experience after receiving the treatment. For example, the recipient may provide feedback about any change (improvements or declines) in symptoms, vital signs, overall health and fitness, or physical or mental energy. The recipient may also provide feedback about progress towards reaching therapeutic objectives. Other types of feedback may also be received. For example, feedback may originate from measured bloodwork, epigenetic changes, or other sources. This feedback can then be used as a helpful datapoint in determining the effectiveness of the treatment and whether each of the ingredients used in the selected formula had the desired effect. In some embodiments, with multiple datapoints collected from various recipients, the nutritional database can be updated and improved.

(31) In some embodiments, the preexisting formulas may be modified based on the feedback. For example, a first recipient with a particular diagnosis may find success in achieving optimum health through receiving treatments with a first micronutrient, while a second recipient with the same diagnosis may not be successful on the first micronutrient. Over time, with enough datapoints from recipients with the same diagnosis, it may be discovered that a particular gene interferes with the effectiveness of the first micronutrient, but that a second micronutrient can have the same effect. Once the nutritional database is updated to reflect this finding, the personalized therapy becomes more effective. For this reason, the feedback loop provides an important benefit.

(32) In a particular implementation, the recipient's genetic micronutrient requirement profile may be generated to illustrate the specific nutrients needed. This micronutrient requirement profile may be generated based on the collected data. Based on the recipient's micronutrient requirement profile, the IV or IM therapy can be personalized to provide the recipient with vital ingredients in the amounts necessary. Moving forward, the personalized formula can be sequentially refined based on the recipient's goals combined with diagnostics to reach the desired outcomes of proper nutrition and optimal health.

(33) An example method **100** of providing genetically personalized IV or IM nutrition therapy is shown, for example, in FIG. 1. Although method **100** as shown in FIG. 1 includes a number of operations in a particular order, method **100** may include more or fewer operations and the operations may be completed in different orders. In addition or alternatively, in some embodiments, variations of the operations shown in FIG. 1 may be used as part of method **100**.

(34) In some embodiments, method **100** includes developing an initial mapping of genetic and/or medical data to micronutrients based on existing scientific literature at operation **110**, storing the initial mapping in a database at operation **120**, receiving genetic data (e.g., a single nucleotide polymorphism (SNP), a genome, gene, multiple SNPs, etc.) and/or medical data (e.g., symptoms, conditions, therapeutic objectives, vital signs, etc.) of a patient at operation **130**, comparing the genetic and medical data with the current mapping (e.g., the initial mapping or an updated mapping based on feedback from previous treatments and/or new scientific literature) at operation **140**, determining an IV or IM nutrition therapy based on the comparison at operation **150**, treating the patient with the IV or IM nutrition therapy at operation **160**, receiving feedback data based on the treatment at operation **170**, creating an individual treatment score for that treatment based on the feedback at operation **180**, and updating and storing the current mapping based on the individual treatment score at operation **190**. The method **100** may include repeating operations **130**, **140**, **150**, **160**, **170**, **180**, and **190** for multiple patients that are geographically remote from each other.

(35) Thus, in some embodiments, because feedback from patient treatments is constantly being used to update the mapping, the mapping will be more up to date than existing published science alone. In some embodiments, new scientific literature may be inputted to update the mapping, in

addition to the feedback from patient treatments. Inputting both scientific literature and feedback from patient treatments helps keep the mapping as up to date as possible and contributes to greater accuracy and more personalization in the mapping.

(36) In some embodiments, at operation **110**, the initial mapping of genetic and/or medical data to micronutrients is developed. The mapping identifies the micronutrients that are suitable for patients having particular genetics, particular symptoms, particular conditions, or particular therapeutic objectives based on the existing scientific literature. As discussed further below, the micronutrients may be mapped to various combinations of genetics, symptoms, conditions, and therapeutic objectives. For example, the micronutrients may be mapped to particular genetics, a particular symptom, a particular condition, and a particular therapeutic objective. The genetic data may be an SNP, a genome, a gene, multiple SNPs, or some other genetic data. The discussion below primarily refers to an SNP, but the same principles apply to using other types of genetic data and the disclosure includes other types of genetic data.

(37) An example process for operation **110** of developing an initial mapping is shown in FIG. 2. In some embodiments, at operation **112**, an individual score is created for individual scientific literature references for a specific micronutrient with respect to an SNP (or some other type of genetic data) and a symptom, condition, or therapeutic objective (or a combination of a symptom, condition, and/or therapeutic objective). An individual score may be created for every available scientific literature reference. A reference may have multiple individual scores if the reference discusses multiple micronutrients, multiple SNPs, multiple symptoms, multiple conditions, or multiple therapeutic objectives. For example, the reference may receive an individual score for each unique combination of micronutrient, SNP, symptom, condition, and/or therapeutic objective discussed in the reference.

(38) An individual score may be positive or negative. A positive score is created when the individual scientific literature reference provides evidence of a positive connection between a micronutrient and an SNP, symptom, condition, therapeutic objective, or a combination of two or more of these characteristics. A negative score is created when the individual scientific literature reference provides evidence of a negative connection between a micronutrient and an SNP, symptom, condition, therapeutic objective, or a combination of two or more of these characteristics. In some embodiments, a scientific literature reference that provides a non-statistically significant outcome may also receive a negative score.

(39) In some embodiments, the individual score may be zero. For example, in some embodiments, a score of zero may be created for references that provide a non-statistically significant outcome. In some embodiments, only studies that have a statistical significance assessment are scored. For example, case series and reports may be logged but do not affect the scoring, due to a low level of scientific evidence. On the other hand, randomized controlled trials, meta-analyses, and observational studies may be scored.

(40) In some embodiments, the individual score may be weighted based on various factors. For example, the individual score may be weighted based on the strength of the findings in the reference, the number of categories (e.g., SNP, symptom, condition, therapeutic objective) analyzed in the reference, or the category that had a connection (e.g., a connection for an SNP may be weighted higher than a connection for a therapeutic objective, etc.). Other factors may also be used. As one example, in some embodiments, a reference that relates to an injectable outcome may be given an individual score that has more weight than an individual score of a reference that relates to an oral outcome.

(41) In some embodiments, at operation **114**, individual scores relating to a specific micronutrient may be added together to determine a total score for the specific micronutrient with respect to an SNP (or some other type of genetic data) and a symptom, condition, or therapeutic objective (or a combination of a symptom, condition, and/or therapeutic objective). In some embodiments, at operation **116**, it is determined whether the specific micronutrient matches the SNP, symptom,

condition, and therapeutic objective. For example, if the total score for the micronutrient with respect to the SNP, symptom, condition, and therapeutic objective exceeds a predetermined threshold, then the micronutrient is determined to match the SNP, symptom, condition, and therapeutic objective. This may be repeated for each combination of SNP, symptom, condition, and therapeutic objective so that the method identifies all the scenarios for which the specific micronutrient is a good match.

(42) In some embodiments, at operation **118**, this scoring process (i.e., operations **112**, **114**, and **116**) is repeated for multiple micro-nutrients and multiple combinations of SNPs, symptoms, conditions, and therapeutic objectives to develop the initial mapping. A schematic representing an example initial mapping **200** is shown in FIG. **3**. For simplicity, FIG. **3** only shows two micronutrients (M.sub.1 and M.sub.2), two SNPs (SNP.sub.1 and SNP.sub.2), two symptoms (S.sub.1 and S.sub.2), two conditions (C.sub.1 and C.sub.2), and two therapeutic objectives (TO.sub.1 and TO.sub.2). However, initial mapping **200** may include many more micronutrients, SNPs, symptoms, conditions, and/or therapeutic objectives.

(43) The schematic of initial mapping **200** includes mapping columns **210** that include series of cells **260** that indicate whether the micronutrient matches the combination of SNP, symptom, condition, and therapeutic objective in that particular row. A checkmark indicates a match. To the left of mapping columns **210** are an SNP column **220**, a symptom column **230**, a condition column **240**, and a therapeutic objective column **250**. In some embodiments, changing the SNP, symptom, condition, or therapeutic objective (and keeping the other factors the same) may result in a different recommended micronutrient. In some embodiments, multiple micronutrients may be matched with the same combination of SNP, symptom, condition, and therapeutic objective.

(44) In some embodiments, the initial mapping **200** provides a specific dosage of a micronutrient for a specific combination of SNP, symptom, condition, and therapeutic objective, in addition to identifying which micronutrient is a match. In particular, when scientific literature provides evidence of the right amount of a micronutrient to use for a particular SNP, symptom, condition, and/or therapeutic objective, this detail may be included in the mapping. In some embodiments, scientific literature providing evidence of a dosage may be scored to determine the recommended dosage. In some embodiments, the dosage evidence is not scored, but is simply used as a guideline (until further information becomes available through patient treatments and feedback about the results).

(45) In some embodiments, there may be holes in the initial mapping **200**. For example, there may not be sufficient scientific literature for a certain micronutrient, SNP, symptom, condition, or therapeutic objective to determine whether there is a good match. In some embodiments, the method may include making a best guess based on similar or related micronutrients, SNPs, symptoms, conditions, or therapeutic objectives. As discussed below, method **100** refines the results continuously over time. In some embodiments, the initial mapping may begin less specific (e.g., the recommended micronutrient is the same for a given symptom, condition, and therapeutic objective regardless of genetic information) with the mapping becoming more specific as process **100** is used (e.g., the recommended micronutrient for a given symptom, condition, and therapeutic objective differs depending on genetic information). The increased specificity is not limited to genetic information, but may also be based on symptoms, conditions, and/or therapeutic objectives. For example, the initial mapping may map a micronutrient to a given SNP, symptom, and condition, regardless of the therapeutic objective, while the mapping may be updated through process **100** so that the recommended micronutrient for the given SNP, symptom, and condition differs depending on the therapeutic objective.

(46) In some embodiments, the mapping relates solely to genetic information (e.g., SNPs) and suitable IV or IM micronutrients (potentially including dosage and frequency). Feedback through patient treatments may likewise focus solely on genetic information rather than considering medical information in addition to genetic information. In some embodiments, the mapping relates

solely to clinical presentations (e.g., symptoms or conditions) and suitable IV or IM micronutrients (potentially including dosage and frequency). In some embodiments, any combination of genetic information, medical information, symptoms, conditions, and therapeutic objectives (including each one individually) may be mapped to suitable IV or IM micronutrients (potentially including dosage and frequency). This same principle applies to other implementations disclosed herein. Specifically, the methods and systems disclosed herein may focus only on SNPs in identifying suitable micronutrients, only on clinical presentations, only on symptoms, only on conditions, only on therapeutic objectives, or any combination of these factors.

(47) In addition, while the method of FIG. 1 focuses on IV or IM nutrition therapy, in some embodiments, the mapping may be to oral supplement nutrients, and in particular to oral supplement micronutrients. This same principle applies to other implementations disclosed herein. Specifically, the methods and systems disclosed herein may relate to oral supplement treatments instead of IV or IM nutrition therapy. In the context of oral supplements treatments, again, the mapping may be of clinical presentations to oral supplement micronutrients, SNPs to oral supplement micronutrients, or a combination of clinical presentation and SNPs to oral supplement micronutrients.

(48) Returning to FIG. 1, in some embodiments, at operation **120**, the initial mapping **200** is stored in a database. In some embodiments, the individual scores of the scientific literature references for each micronutrient, SNP, symptom, condition, and therapeutic objective combination are also stored in the database. The added scores (or the total scores) for each micronutrient, SNP, symptom, condition, and therapeutic objective combination may also be stored in the database. The database may be part of a server and made available over a network to electronic devices, as discussed in more detail below with respect to FIGS. 11 and 12. The electronic devices accessing the database over the network may be patient electronic devices, healthcare provider electronic devices, or both (see FIG. 12). Because the initial mapping **200** is stored in a database available over a network, the mapping can be updated in real time by patient treatments that occur in different geographic locations and by different healthcare providers through the remaining operations of method **100**. Although the remaining operations of method **100** are described for a single patient, these operations may be repeated many times for multiple patients to update the mapping and improve the accuracy of the mapping.

(49) In some embodiments, at operation **130**, an SNP, symptom, condition, or therapeutic objective are received for a patient. In some embodiments, all four of an SNP, symptom, condition, and therapeutic objective are received. In some embodiments, only a subset of an SNP, symptom, condition, and therapeutic objective are received. For example, an SNP for a patient and at least one of a symptom, a condition, and a therapeutic objective of a patient may be received (e.g., SNP and symptom; SNP and condition; SNP and therapeutic objective; SNP, symptom, and condition; etc.). Multiple SNPs, symptoms, conditions, and therapeutic objectives may be received for the patient. For example, if the patient has multiple symptoms, each symptom may be included. This information may be entered into fields on an electronic device by the patient or by the healthcare provider. In some embodiments, a patient may have a profile that stores information from previous treatments. When a patient comes in for another treatment, the system may automatically retrieve the stored information (e.g., regarding SNPs, symptoms, conditions, therapeutic objectives). Because symptoms, conditions, and therapeutic objectives may change over time, the patient may either confirm or edit this information.

(50) In some embodiments, the SNP, symptom, condition, or therapeutic objective may be determined by a test kit or monitoring equipment. For example, a genetic testing kit may be used to determine the patient's genetic information, including an SNP. Additional examples of monitoring equipment that may determine symptoms or conditions (or contribute to determining a therapeutic objective) include thermometers, scales, blood pressure monitors, blood glucose monitors, heart rate monitors, pulse oximeters, electrocardiographs, etc.

(51) After receiving the patient's SNP, symptom, condition, and therapeutic objective (or a subset, such as an SNP and at least one of a symptom, condition, and therapeutic objective), this information may be compared with the current mapping stored in the database at operation **140**. For the first patient, the current mapping may be the initial mapping **200**. Once feedback from patient treatments has been received, the current mapping may be an updated mapping based on that feedback. Comparing the patient's information with the current mapping identifies one or more micronutrients that have been matched to the SNP, symptom, condition, and therapeutic objective (or a subset, such as an SNP and at least one of a symptom, condition, and therapeutic objective).

(52) In some embodiments, at operation **150**, an IV or IM nutrition therapy may be determined for the patient based on the one or more micronutrients identified from the current mapping (e.g., initial mapping **200** or an updated mapping). In some embodiments, the IV or IM nutrition therapy is determined by selecting an existing IV or IM nutrition therapy formula that includes a specific dosage of at least one of the identified micronutrients. For example, there may be a plurality of IV or IM nutrition therapy formulas to choose from, and the formula that provides the identified micronutrients may be selected. In some embodiments, the formula that not only provides the identified micronutrients, but also provides the right amount of the identified micronutrients is selected. In some embodiments, the IV or IM nutrition therapy is determined by selecting an existing IV or IM nutrition therapy formula that most closely provides the identified micronutrients and modifying the formula to better provide the identified micronutrients. In some embodiments, the IV or IM nutrition therapy is determined by specifically creating an IV or IM nutrition therapy formula that includes the identified micronutrients. Determining the IV or IM nutrition therapy may include determining an IV or IM nutrition therapy formula to administer (using one of the approaches discussed above), determining a dosage or amount of the formula to administer, determining a frequency of administering the formula, and/or determining a number of times to administer the formula.

(53) At operation **160**, the patient is treated with the IV or IM nutrition therapy determined at operation **150**. In some embodiments, at operation **170**, feedback data relating to the IV or IM nutrition therapy treatment is received. The feedback data may be about an effectiveness of the IV or IM nutrition therapy in addressing the symptom, the condition, or the therapeutic objective of the patient (or a subset of the symptom, condition, and therapeutic objective).

(54) In some embodiments, the feedback data may include objective feedback (e.g., test results) and/or subjective feedback (e.g., survey responses). For example, the feedback data may include feedback about measured bloodwork (e.g., comparing bloodwork from before and after the IV or IM nutrition therapy), epigenetic changes, improvements or declines in symptoms, progress towards reaching therapeutic objectives, vital signs, overall health and fitness, or physical or mental energy. Other types of feedback may also be used.

(55) In some embodiments, at operation **180**, an individual treatment score is created based on the feedback data. The individual treatment score may be for the identified one or more micronutrients in the IV or IM nutrition therapy with respect to the patient's SNP and symptom, condition, and therapeutic objective (or a subset). Similar to the scores for scientific literature, individual treatment scores may be positive or negative. For example, when the IV or IM nutrition therapy results in improvement in the symptom, a positive individual treatment score may be created. On the other hand, if the symptom worsens, a negative score may be created. The individual treatment score may be based on a single factor (e.g., whether the symptom improved or declined), or the score may be based on multiple factors (e.g. change in symptom(s), progress for reaching therapeutic objective, vital signs, energy levels, etc.).

(56) In addition, individual treatment scores may be weighted depending on the feedback data. For example, the individual treatment score may be commensurate with how much a symptom improves or declines and/or how much progress is made (or lost) towards reaching a therapeutic objective. Thus, the more the symptom improves, the higher the individual treatment score will be.

As another example, the more progress is made towards the therapeutic objective, the higher the individual treatment score will be.

(57) In some embodiments, multiple individual treatment scores may be created for a single treatment. For example, if a symptom declines but progress is made toward a therapeutic objective, a negative score may be made for the combination of the SNP, symptom, and therapeutic objective while a positive score may be made for the combination of the SNP and therapeutic objective without that symptom.

(58) In some embodiments, individual treatment scores may be weighted based on multiple factors. Certain factors may have greater weight than others. For example, whether the symptom improves may have greater weight than measured bloodwork or vice versa.

(59) In some embodiments, at operation **190**, the current mapping is updated and then stored in the database. As noted above, for the first patient, the current mapping may be the initial mapping **200**. Once feedback from patient treatments has been received, the current mapping may be an updated mapping based on that feedback. An example process for operation **190** of updating a current mapping is shown in FIG. **4**.

(60) In some embodiments, at operation **192**, an individual treatment score (e.g., from treatment of a first patient) is added to a total score in the current mapping (e.g., initial mapping **200**) for the specific micronutrient(s) used in the treatment with respect to the SNP and symptom, condition, and therapeutic objective (or a subset) of the patient being treated. To illustrate this principle, if a patient is treated having a GPx1 gene SNP, a headache as a symptom, fibromyalgia as a condition, and weight loss as a therapeutic objective, the individual treatment score may be tied to this combination of SNP, symptom, condition, and therapeutic objective. Thus, this individual treatment score would be added to the total score from the current mapping of the GPx1 gene SNP, headache symptom, fibromyalgia condition, and weight loss therapeutic objective.

(61) In some embodiments, at operation **194**, an individual score is created for any new scientific literature references that have not previously been taken into account (to the extent there are any). Thus, if a new study has linked a specific micronutrient to an SNP, symptom, condition, or therapeutic objective (or a combination of some or all of these), an individual score is created for each relevant combination of micronutrient, SNP, symptom, condition, and therapeutic objective. The principles discussed above for creating individual scores for scientific literature references equally apply to any new references.

(62) In some embodiments, at operation **196**, the individual scores for any new scientific literature references are added to the total score in the current mapping (in addition to the added individual treatment scores added in operation **192**). The updated total score (based on the addition in operation **192** and/or operation **196**) may be stored in the database.

(63) In some embodiments, at operation **198**, the updated total score is compared with a predetermined threshold (e.g., the same predetermined threshold used when evaluating the total score to develop the initial mapping **200**). This comparison allows for reevaluating whether there is a match between a specific micronutrient and an SNP, symptom, condition, and therapeutic objective combination. This process of updating the current mapping ensures that the mapping is based on the latest scientific literature and real time results of actual treatment. Similar comparisons may be made to determine appropriate dosage, frequency, etc. of the nutrition therapy.

(64) In some embodiments, fewer or more operations than are shown in FIG. **4** may be used to update a current mapping. For example, the mapping may be updated based only on patient treatments, only on new scientific literature references, or a combination of patient treatments and scientific literature references. The order of the operations may differ from what is shown in FIG. **4**.

(65) In some embodiments, personalized therapy may be provided through one or more algorithms, as shown, for example, in FIGS. **5-7**. In some embodiments, the algorithms may include the process discussed above with respect to FIGS. **1-4**. However, other algorithms may also be used.

(66) FIG. 5 illustrates an implementation of a method 2 of providing personalized therapy. As shown, the system may first collect medical data from a customer 10, or the recipient of the personalized therapy. This data may include symptoms 12, medical history 14, blood test results 16, and vitals 18. In some embodiments, the customer 10 may provide the data through an electronic device. For example, the customer 10 may input data into an app run on a smart phone, tablet, or personal computer. Alternatively, a healthcare provider may obtain the data from the customer through medical tests, questioning the customer 10, reviewing health records of customer 10, or through other means (or a combination of these options). The healthcare provider may enter the obtained data into the system (e.g., through an app run on a smart phone, tablet, personal computer, or other electronic device). In some embodiments, data may automatically be received by the system. For example, medical tests may be configured to directly provide results (e.g., blood test results 16, vitals 18, etc.) to the system electronically.

(67) Once the data has been collected, the data may be fed into an algorithm 20. In some embodiments, the algorithm 20 is configured to map the data into a code 30, such as an ICD-11 code or ICD-10 code. The code 30 may represent a condition or type of condition. For example, in the ICD-10 code, an acute upper respiratory infection 32 may be coded as J06.9, fatigue 34 may be coded as R53, and fibromyalgia 36 may be coded as M79.7. The coding may be different in the ICD-11 code.

(68) In some embodiments, these codes 30 may be fed into another algorithm (e.g., recommendation algorithm 40) to determine a personalized nutrition therapy for customer 10. Recommendation algorithm 40 may be based on scientific evidence 42 and feedback 44 already obtained through monitoring past recipients that monitors efficacy of previous treatments. In some embodiments, recommendation algorithm 40 takes into account genetic information of customer 10 in addition to the results from code 30. For example, recommendation algorithm 40 may use scientific evidence 42, monitoring efficacy 44, code 30, and genetic information of customer 10 (e.g., SNP) to determine micronutrients most suited for customer 10.

(69) In some embodiments, recommendation algorithm 40 may be configured to identify an existing product 50, with particular nutritional formulas, that is best suited to meet the nutritional needs of the customer 10. For example, the system may include a number of existing products 50, such as Hydromax IV 51, Ultraviv IV 52, Megaboost IV 53, Vitaglow IV 54, Royal Flush IV 55, Vitamin B12 IM 56, or other IV or IM products 50. Algorithm 40 is configured to evaluate which of the existing products 50 is best suited to provide micronutrients that customer 10 needs in light of the code 30, scientific evidence 42, feedback 44 that monitors efficacy, and genetic information of the customer 10. Alternatively, in some embodiments, a completely new formula may be developed to more effectively meet the nutritional needs of the customer 10. As mentioned, both algorithm 20 and recommendation algorithm 40 may be implemented by a person, or may be automated on a computer or other computing device.

(70) Another example is provided in FIG. 6, illustrating a method 4 of providing personalized therapy to target a specific disease, such as fibromyalgia, and its symptoms. Fibromyalgia is a disease in which the main symptom is chronic, strongly systemic pain, or, even if partial, widespread chronic pain, the pain being observed not only in muscular tissues but also in the skin. In fibromyalgia, such systemic chronic pain is often not alone and is also accompanied by feelings of fatigue, malaise, depression, anxiety, morning stiffness, muscle stiffness, sleep disturbance or the like. In addition, symptoms such as headache, facial pain, cognitive impairment (lapse of memory, concentration deficit), gastrointestinal complaints (visceral pain, digestive system disturbance, flatulence), frequent urination, diarrhea, constipation, or dysmenorrhea may also occur concomitantly.

(71) As already discussed in the previous example, the medical and genetic data of customer 10 may be collected. In this particular example, the customer 10 presents data indicative of fibromyalgia. Thus, when the data is fed through the algorithm 20, the code 30 results in the code

for fibromyalgia. Similar to the previous example, the code **30** is fed into recommendation algorithm **40**, which may also take into account the genetic data of customer **10**, to arrive at a recommended formula based on scientific evidence **42** and feedback **44** that monitors efficacy. In some embodiments, recommendation algorithm **40** may output a recommendation of a nutrition therapy that includes co-enzyme Q10 (CoQ10), a co-enzyme with a molecular structure of 2,3-dimethoxy-5-methyl-6-decaprenyl-1,4-benzoquinone, which may be administered to the customer **10** through IV or IM therapy to treat fibromyalgia and its symptoms. A pharmaceutical composition of CoQ10 may also or alternatively be administered. In some instances, the CoQ10 composition **60** administered may contain CoQ10 and vitamin B12 (as shown in FIG. **6**). In some embodiments, recommendation algorithm **40** may output current studies **76** that link the recommended micronutrient ingredients (e.g., CoQ10, Vitamin B12, etc.) to the condition of fibromyalgia.

(72) In some embodiments, at operation **70**, outcomes of the treatment may then be measured. For example, outcomes may be measured through blood tests, vitals, and feedback from the customer **10**, as well as other means, including those discussed above for FIGS. **1-4**. These outcomes can be used at operation **75** to build evidence of what works and what doesn't, which can then be used to improve recommendations to future customers. In other words, the outcomes of each treatment of each customer may contribute to feedback **44** that monitors efficacy. This feedback may be tied to specific medical and genetic data, leading to more personalized results over time.

(73) In some embodiments, a life science team **72** may evaluate the measurements **70** along with other available medical data and genetic data to gain additional insight **82**. Additional analysis and/or studies related to the outcomes may improve evidence at operation **80** for what can be adjusted to alter outcomes at operation **85**. For example, results from the measurements **70** may be analyzed in connection with medical or genetic data to determine specific dosages of an ingredient or a therapy formula that maximize results for certain situations. Thus, the dosage of the CoQ10 composition **60** used may be determined by the customer's medical data, genetic data, and/or medical history. An example of additional insight **82** is shown in FIG. **6**, in which the genetic data of customer **10** may identify a single-nucleotide polymorphism (SNP) for CoQ10 activation. In such an instance, depending on the characteristics of that particular SNP, the dosage of CoQ10 may need to be increased or decreased due to the customer's reaction to CoQ10. Additionally, feedback may be received from the customer **10** regarding the effect of the dosage on the recipient. Based on this feedback and the knowledge concerning the recipient's medical data, medical history, and genetic data, the dosage may be adjusted to better provide the nutritional benefits of the CoQ10.

(74) Although FIG. **6** provides one example of a disease that may be treated through IV or IM therapy (i.e., fibromyalgia), other diseases may also be treated through IV or IM therapy with similar methods of treatment, where different dosages and different substances may be used, but the same method of using genetic data to personalize the treatment is implemented.

(75) FIG. **7** illustrates a method **300** of creating and using a patient treatment nutrition therapy algorithm that is updated over time to improve accuracy and personalization of nutrition therapy. At operation **310**, a patient treatment nutrition therapy recommendation algorithm is created. In some embodiments, the algorithm is configured to receive one or more inputs and output a recommendation of an IV or IM nutrition therapy formula to treat a patient based on the inputs. The inputs may include medical information, genetic information, and therapeutic objectives, as discussed further below. The output may include a recommended dosage in addition to identifying the IV or IM nutrition therapy formula.

(76) In some embodiments, the algorithm is created based on existing scientific literature. For example, the algorithm may be based on a mapping of medical information, genetic information, and therapeutic objectives to micronutrients, as discussed above with respect to FIGS. **1-4**. Thus, the algorithm may include a set of rules, established by existing scientific literature that links micronutrients to various medical information, genetic information, and therapeutic objectives, so that the most suitable nutrition therapy formula containing the desired micronutrients and the most

suitable dosage is recommended given a certain set of medical information, genetic information, and therapeutic objectives. Once created, the algorithm may be stored in a database. For example, the algorithm may be stored in a database on a server so that users can run the algorithm on an electronic device in communication with the server.

(77) In some embodiments, at operation **320**, medical information, genetic information, and therapeutic objectives of a patient are received as algorithm inputs. A user (e.g., a patient or a healthcare provider) may input only medical information, only genetic information, only therapeutic objectives, or any combination of this information (e.g., medical information, genetic information, and therapeutic objectives) as inputs to the algorithm. In addition, a user may enter a single input under a category or multiple inputs. For example, a user may enter one therapeutic objective or multiple therapeutic objectives.

(78) In some embodiments, the medical information may include blood test results, vital signs (e.g., blood pressure, heart rate, heart rhythm, etc.), medical histories, medical diagnoses, current symptoms, diseases of which the patient is at heightened risk, family history, or other medical information. Medical information may include biodata that is captured and measured through various processes, which may include blood testing, urine sampling (e.g., toxicology screens, etc.), stool sampling, continuous blood glucose monitoring, and imaging/scanning (e.g., ultrasound, etc.). Devices may include blood pressure monitors, heart rate monitors, body hydration monitors, breathalyzers, blood glucose monitors, metabolic devices, pulse oximeters, and wearable devices configured to measure any type of biodata.

(79) In some embodiments, the genetic information comprises genetic test results. The genetic information may comprise a gene, genome, SNP, or other piece of genetic data. In some embodiments, the genetic information comprises epigenetic measurements (e.g., epigenetic clocks, methylation scores, etc.)

(80) A variety of therapeutic objectives may be used as inputs. For example, the therapeutic objectives may comprise health goals, aesthetic goals, or fitness goals. The therapeutic objectives may be to lose weight, to gain weight, to improve fitness level, to improve energy levels, to alter skin tone (to either a darker tone or a lighter tone), to improve skin quality and/or health, to improve hair quality and/or health, to improve mental acuity, to prevent disease, to improve symptoms of an existing illness or disease, to improve sleep quality, to improve longevity/anti-aging, to protect against excessive stress, to protect against toxin exposures (e.g., from pollution, chemicals, alcohol, etc.), to rehydrate, to support immune health, and/or to heal a wound. Other therapeutic objectives may also be used.

(81) In some embodiments, at operation **330**, the patient treatment nutrition therapy recommendation algorithm outputs a therapy recommendation based on the medical information, genetic information, and therapeutic objectives of the patient that were received as algorithm inputs. In some embodiments, the therapy recommendation comprises an existing IV or IM nutrition therapy formula (e.g., products **50**, as shown in FIG. 5). The existing IV or IM nutrition therapy formula has at least one micronutrient that the algorithm identified as a match for the medical information, genetic information, and therapeutic objectives.

(82) In some embodiments, the algorithm identifies a match based on both scientific literature and previous IV or IM nutrition therapy treatments. For example, the database may store individual scores of multiple existing individual scientific literature references relating to the identified micronutrient. These individual scores may be specific to the inputted medical information, genetic information, and therapeutic objectives. The database may also store individual treatment scores of previous IV or IM nutrition therapy treatments that included the identified micronutrient. Again, these scores may be specific to the inputted medical information, genetic information, and therapeutic objectives. For example, the previous treatments may be from earlier patients that had similar medical information, genetic information, and therapeutic objectives. The individual scores of the scientific literature references and the individual treatment scores discussed here may have

the same characteristics and be derived in the same way as the scores discussed above with respect to FIGS. **1-4**. The algorithm may determine a match when the scores add up to a predetermined threshold.

(83) At operation **340**, the patient is treated with an IV or IM nutrition therapy according to the recommendation from operation **330**. At operation **350**, feedback data about the results of the IV or IM nutrition therapy of the patient is received. In some embodiments, the feedback data includes feedback about measured bloodwork, epigenetic changes, improvements or declines in symptoms, progress towards reaching therapeutic objectives, vital signs, overall health and fitness, or physical or mental energy.

(84) In some embodiments, at operation **360**, an individual treatment score is created based on the feedback data from operation **350**. This individual treatment score may relate to the effectiveness of the identified micronutrient with respect to the inputted medical information, genetic information, and therapeutic objectives. In some embodiments, this individual treatment score may be entered electronically into a computer system associated with the database. For example, an electronic device that receives feedback data may automatically determine the individual treatment score and transmit the individual treatment score to the database to be stored (and thus used in future instances of identifying a match when considering similar medical information, genetic information, or therapeutic objectives). As another example, a user, such as a healthcare provider may manually enter an individual treatment score into an electronic device so that it can be stored in the database.

(85) In some embodiments, the algorithm is updated to incorporate the individual treatment score at operation **370**. In some embodiments, the algorithm is updated to incorporate new scientific literature references. Thus, the algorithm may account for recent published science. At operation **380**, future recommendations are created based on the updated algorithm. Thus, subsequent patients using the system will have the benefit of feedback from the patient's treatment to more accurately identify micronutrients that are most effective for specific combinations of medical information, genetic information, and therapeutic objectives. The updated patient treatment nutrition therapy recommendation algorithm is thus configured to recommend treatment and micronutrients that most closely match patient needs based on all available patient medical information, genetic information, and therapeutic objectives, and most current aggregated feedback based on real patients with similar genes and symptoms.

(86) After operation **380**, the process is repeated (beginning at operation **320**) for every subsequent patient. This creates a real time feedback loop that automatically applies patient therapy feedback data to the patient treatment nutrition therapy recommendation algorithm. In this way, all new therapy recommendations include the feedback data from all previous therapy recommendation patients who received recommendations through the algorithm. While the algorithm is originally based on existing scientific references, the feedback from actual treatments is constantly used to update the algorithm. In some embodiments, new scientific references may also be scored and used to update the algorithm. The algorithm will therefore be more up to date than existing published science alone.

(87) In some embodiments, the patient and the subsequent patients (or at least some of the patient and the subsequent patients) are positioned geographically remote from each other.

(88) As explained above, genetics can influence the proper nutrition for a particular recipient. For example, a single-nucleotide polymorphism (SNP) in genetic material can, among other things, indicate differences in susceptibility to various diseases. As such, SNPs of different types lead to different desired doses as part of a therapy treatment. As shown in FIG. **8**, the nutritional database may incorporate this information to point to particular effective nutrients. FIG. **8** presents a schematic **400** that represents a portion of the nutritional database, including SNPs **410** of interest, IV ingredients **420** of interest for the SNPs **410**, genotypes **430** of the SNPs **410**, and associated outcomes and doses **440** of the IV ingredients **420** based on the genotypes **430**. For example, IV

ingredients **410** (i.e., micronutrients) such as glutathione and selenium are of particular interest for the GPx1 gene SNP **410**. Depending on the genotype **430** of the SNP **410**, different amounts of glutathione and selenium are recommended. Similarly, vitamin B12 is relevant to the MTRR gene SNP **410**, and depending on the genotype **430**, different amounts of hydroxocobalamin or methylcobalamin may be recommended. Thus, genetic testing may form an important part of the personalized therapy disclosed herein.

(89) As an illustrative example, glutathione and selenium are both important for immune system health. Glutathione performs an antioxidant function and may be involved in immune stimulation and detoxification of xenobiotics (see Pastore et al. 2003. Clinica Chimica Acta 333: 19-39). In general, it is known that glutathione synthesis genes are essential for survival, and organisms deficient in these genes are less resistant to oxidative stress and various harmful substances. In addition, there is a report that various human diseases are associated with a decrease in glutathione concentration in body (Wu et al. 2004. J. Nutr. 134: 489-492). Similarly, selenium is an antioxidant that helps lower oxidative stress in the body, reduces inflammation, and enhances immunity. As shown in FIG. 8, a method for treating a glutathione deficiency or a selenium deficiency in a subject may comprise administering either glutathione or selenium, or a pharmaceutical composition thereof. The glutathione or selenium may be administered through IV therapy, and the dosage **440** may be dependent on a genotype **430** of the GPx1 gene SNP **410** of the subject. In this way, personalizing to the genetic data of the recipient improves the effect of the IV treatment. For example, as shown in FIG. 8, if the genotype **430** of the GPx1 gene SNP **410** is C/C, the glutathione dosage **440** may be 1 g, if the genotype **430** of the GPx1 gene SNP **410** is C/T, the glutathione dosage **440** may be 2 g, and if the genotype **430** of the GPx1 gene SNP **410** is T/T, the glutathione dosage **440** may be 3 g. Similarly, if the genotype **430** of the GPx1 gene SNP **410** is C/C, the selenium dosage **440** may be 30 mg, if the genotype **430** of the GPx1 gene SNP **410** is C/T, the selenium dosage **440** may be 60 mg, and if the genotype **430** of the GPx1 gene SNP **410** is T/T, the selenium dosage **440** may be 100 mg.

(90) As another illustrative example, vitamin B12 is a micronutrient that helps keep the body's blood and nerve cells healthy and helps make DNA. Vitamin B12 may also help prevent anemia, which causes tiredness and fatigue. A method of treating a vitamin B12 deficiency in a subject may comprise administering at least one of hydroxocobalamin and methylcobalamin, or a pharmaceutical composition thereof, through IV or IM therapy. The dosage **440** may be dependent on a genotype **430** of the MTRR gene SNP **410** of the subject. Thus, again, personalizing the dosage **440** based on the genetic data of the recipient improves the effect of the IV treatment. For example, if the genotype **430** of the MTRR gene SNP **410** of the subject is A/A, the dosage **440** may be 1 mg of hydroxocobalamin, if the genotype **430** of the MTRR gene SNP **410** of the subject is A/G, the dosage **440** may be 1 mg of methylcobalamin, and if the genotype **430** of the MTRR gene SNP **410** of the subject is G/G, the dosage **440** may be 2 mg of methylcobalamin.

(91) In some embodiments, feedback is received about the effectiveness of the dosage **440** for each treatment of a patient, similar to the feedback discussed above. Thus, over time, the nutritional database **400** may be updated to improve accuracy and personalization. This may be done using the processes discussed with respect to FIGS. 1-4 and 7. In some embodiments, a life science team **72** may monitor efficiency/effectiveness of the dosages **440** at operation **70**. In addition, the life science team **72** may compile and/or develop evidence of what works and what doesn't at operation **75** to improve recommendations of dosages **440** to future customers.

(92) FIG. 9 illustrates a chart **500** showing additional example ingredients or formulas **540** that may be selected based on the genes **510**, SNPs **520**, and genotypes **530** of the recipient. As shown, the recommended dosages **540** for various ingredients vary based on the genotype **530** of the recipient. For example, in relation to the GSTP1 gene **510**, if the recipient has the A/A genotype **530**, the recipient may receive 0 g of glutathione and 300 mg of NAC. If the recipient has the A/G genotype **530**, the recipient may receive 1 g of glutathione and 400 mg of NAC. Lastly, if the

recipient has the G/G genotype **530**, the recipient may receive 2 g of glutathione and 500 mg of NAC. Thus, the genetic data of the recipient influences the recommended dosage **540**. In addition, FIG. **9** illustrates in column **550** that the medical data of the recipient can influence the recommended dosage (as discussed above). For example, again in relation to the GSTP1 gene, if the recipient has exposure to polycyclic aromatic hydrocarbons (PAH), the dosage of glutathione may be increased by 1 g. FIG. **9** contains additional examples that follow the same pattern as that already described.

(93) An individual's health and nutritional needs are influenced by a variety of factors that are understood through numerous fields of study. For example, pharmacology, genetics, nutrigenomics, dietetics, clinical medicine, biomedicine, molecular biology, and epigenetics all contribute to understanding nutritional needs and how to meet them. In addition, other fields such as statistics and technology are involved in the methods described herein. For this reason, a holistic approach to implementing the personalized therapy systems and methods is beneficial. FIG. **10** illustrates an example of how different areas may be involved in the methods disclosed herein. Specifically, schematic **600** in FIG. **10** illustrates various interactions between the areas of retail health **610**, technology **620**, genomics **630**, and life science research and development **640**, to provide bio medicine **650** to customer **10**. Though not all implementations involve automated steps, including technology **620** may increase the accessibility and accuracy of personalized therapy.

(94) In some embodiments, as shown, for example, in FIG. **11**, a system **700** comprises a number of electronic devices (e.g., a first user's electronic device **710** and a second user's electronic device **720**), a network **730**, and a genetically personalized therapy system **740**. The genetically personalized therapy system **740** disclosed herein may be implemented with a computer or other electronic device. For example, as shown in FIG. **11**, the genetically personalized therapy system **740** may comprise a processor **750** and a storage **760**. In some embodiments, the processor **750** is communicatively coupled to network **730** and the storage **760**. The electronic devices **710**, **720** are also communicatively coupled to network **730**, allowing users (such as first and second user) to access the genetically personalized therapy system **740**.

(95) In some embodiments, the storage **760** may comprise a user library **770** with a plurality of user profiles including a first user profile **772** and a second user profile **776**. The user profiles **772**, **776** may correspond to each recipient of the personalized therapy disclosed above (e.g., the first user and the second user associated with electronic devices **710**, **720**). Each user profile may include the genetic data of the user, the medical data of the user, the medical history of the user, and the personal therapeutic objectives of the user. For example, first user profile **772** includes the genetic and medical data **774** of the first user and second user profile **776** includes the genetic and medical data **778** of the second user. In some embodiments, users may enter or update this data via their electronic devices **710**, **720**. In some embodiments, other users (e.g., healthcare providers) may gain access to another user's profile to enter or update genetic and medical data. For example, the first user's healthcare provider may gain access to first user profile **772** to update genetic and medical data **774**.

(96) The storage **760** may also comprise the nutritional database **780**, including the typical nutrient requirements **782** of the human body and the modifications **784** that occur to these requirements based on the genetic data of the user, the medical data of the user, and the medical history of the user.

(97) In some embodiments, the processor **750** comprises a number of modules to perform various actions. For example, the processor **750** may comprise a profile creation module **752**, a genetic testing module **754**, a formula selection module **756**, and/or a database adjustment module **758**. The modules may work together. The profile creation module **752** may assist a new user creating a profile from their electronic device. As part of this module, the processor **750** may be configured to receive the genetic data, the medical data, the medical history, and the personal therapeutic objectives of the user. For example, first user may create a profile with the first user's electronic

device **710**. As the first user enters data on electronic device **710**, electronic device **710** may transmit the data over network **730** to genetically personalized therapy system **740**. Processor **750** (through its profile creation module **752**) may receive the data, create a new user profile (i.e., first user profile **772**), and store the data **774** in the first user profile **772**.

(98) The genetic testing module **754** may assist a user in determining genetic data. For example, the processor may send instructions to a user's electronic device (first user's electronic device **710**) on how to use a genetic testing kit. In some embodiments, the user enters results of the genetic test onto the electronic device **710**, which is then transmitted to genetically personalized therapy system **740**. Processor **750** (through its genetic testing module **754**) may then add the results of the genetic testing to the user's profile. In some embodiments, the genetic testing kit may send results directly to genetically personalized therapy system **740** over network **730**. The genetic testing module **754** may operate as part of profile creation module **752** or it may be used later on after a profile has already been created.

(99) Formula selection module **756** may provide a recommended IV or IM nutrition therapy to a user. For example, the formula selection module **756** may be configured to analyze the genetic data, medical data, medical history, and personal therapeutic objectives of the user stored in a user profile (e.g., first user profile **772**) using the nutritional database **780** to determine the nutritional needs of the user, and select a formula from a plurality of formulas based on the nutritional needs of the user. Formula selection module **756** may follow the processes discussed above with respect to FIGS. **1-4** and **7**.

(100) Database adjustment module **758** may operate to update the nutritional database **780** to improve accuracy and personalization of the system **740**. In some embodiments, after the user has received the formula via IV therapy, processor **750** (through its database adjustment module **758**) may receive feedback from the user regarding the effect of the formula. The feedback may be received via the user's electronic device (e.g., first user's electronic device **710**) over network **730**. The feedback may also be received via another user, such as the first user's healthcare provider. The database adjustment module **758** is configured to adjust the nutritional database **780** based on the feedback from the user and the genetic data, medical data and medical history of the user stored in the user profile. Database adjustment module **758** may follow the processes discussed above with respect to FIGS. **1-4** and **7**.

(101) As noted above, the systems disclosed herein benefit from the real-time feedback of how effective each treatment is for multiple recipients in various geographic locations, combined with the genetic and medical data of the recipients. A system **800** illustrating this feature is shown, for example, in FIG. **12**. FIG. **12** shows five customers **10a**, **10b**, **10c**, **10d**, **10e** to illustrate the concept of multiple patients in multiple geographic locations, but in practice there may be many more customers in many more locations (e.g., a worldwide group of customers with locations around the world) In some embodiments, system **800** comprises a plurality of genetic testing kits (e.g., genetic testing kits **805a**, **805b**, **805c**, **805d**, **805e**), a plurality of electronic devices (e.g., electronic devices **815a**, **815b**, **815c**, **815d**, **815e**), a network **830**, and a server **840**.

(102) In some embodiments, each genetic testing kit **805a**, **805b**, **805c**, **805d**, **805e** is configured to determine genetic information of one of the patients or customers **10a**, **10b**, **10c**, **10d**, **10e**. Dotted lines in FIG. **12** show a connection between each of the genetic testing kit **805a**, **805b**, **805c**, **805d**, **805e** and respective patients **10a**, **10b**, **10c**, **10d**, **10e**. The genetic testing kits **805a**, **805b**, **805c**, **805d**, **805e** may be self-administered or they may be administered by a healthcare provider (e.g., healthcare provider **825a**, **825c**, **825d**, **825e**).

(103) In some embodiments, each electronic device **815a**, **815b**, **815c**, **815d**, **815e** is configured to receive the genetic information of one of the patients **10a**, **10b**, **10c**, **10d**, **10e**. Dotted lines in FIG. **12** shows a connection between each electronic device **815a**, **815b**, **815c**, **815d**, **815e** and respective patients **10a**, **10b**, **10c**, **10d**, **10e**. In some embodiments, genetic testing kits **805a**, **805b**, **805c**, **805d**, **805e** are configured to send genetic test results directly to electronic devices **815a**,

815b, 815c, 815d, 815e. Alternatively, patient **10a, 10b, 10c, 10d, 10e** may input the genetic information from the genetic test results into electronic devices **815a, 815b, 815c, 815d, 815e**. In some embodiments, each electronic device **815a, 815b, 815c, 815d, 815e** is also configured to receive medical information and/or therapeutic objectives of one of the patients **10a, 10b, 10c, 10d, 10e**. Electronic devices **815a, 815b, 815c, 815d, 815e** may be smartphones, tablets, personal computers, or other electronic devices. Each electronic device **815a, 815b, 815c, 815d, 815e** is communicatively coupled with network **830**, thus providing electronic devices **815a, 815b, 815c, 815d, 815e** access to server **840**, which is also communicatively coupled with network **830**. (104) As illustrated in FIG. 8, each patient **10a, 10b, 10c, 10d, 10e** may go to a healthcare provider **825a, 825c, 825d, 825e** (i.e., a location providing IV or IM nutrition therapies). Dotted lines in FIG. 8 between patients and healthcare providers illustrate a healthcare provider to which a patient goes. Patients may go to the same healthcare provider or a different healthcare provider. For example, patients **10a** and **10b** may both go to healthcare provider **825a**, while patients **10c, 10d, 10e** may go to different healthcare providers **825c, 825d, 825e**. Thus, patients **10a, 10b, 10c, 10d, 10e** may be positioned geographically remote from each other. A single patient may also go to different healthcare providers or locations at different times. In some embodiments, healthcare providers **825a, 825c, 825d, 825e** may be communicatively coupled with network **830** through their own electronic devices, allowing the healthcare providers to access server **840** and input information with respect to patients receiving IV or IM therapy at their location.

(105) In some embodiments, server **840** comprises a processor **842** and a database **850**. Processor **842** may have the same features of processor **750** discussed above. In some embodiments, database **850** stores a patient treatment nutrition therapy recommendation algorithm **852**, which may have the same features as nutritional database **780**. In some embodiments, algorithm **852** may operate the same as the algorithm discussed above with respect to FIG. 7. Database **850** may also store individual scores **854** of a plurality of individual scientific literature references and/or individual treatment scores **856** of previous IV or IM nutrition therapy treatments.

(106) In some embodiments, server **840** is configured to wirelessly receive (e.g., over network **830** from electronic devices **815a, 815b, 815c, 815d, 815e**) the genetic information, medical information and therapeutic objectives of each of the patients **10a, 10b, 10c, 10d, 10e**. For each patient, the server **840** may be configured to output to a respective electronic device **815a, 815b, 815c, 815d, 815e** a therapy recommendation for the patient **10a, 10b, 10c, 10d, 10e** from the algorithm **852** based on the genetic information, medical information, and therapeutic objectives for each patient. In some embodiments, the therapy recommendation comprises an existing IV or IM nutrition therapy formula having at least one micronutrient. The micronutrient(s) is/are identified by the patient treatment nutrition therapy recommendation algorithm as a match for the medical information, genetic information, and therapeutic objectives of the patient based on individual scores **854** stored in the database of a plurality of existing individual scientific literature references relating to the micronutrient(s) and individual treatment scores **856** stored in the database of previous IV or IM nutrition therapy treatments comprising the micronutrient(s).

(107) In some embodiments, server **840** is configured to receive feedback data about results of the therapy recommendation of each patient **10a, 10b, 10c, 10d, 10e** after they have been treated based on the recommendation. This feedback may be provided from either the patients via the electronic devices **815a, 815b, 815c, 815d, 815e** or the healthcare providers **825a, 825c, 825d, 825e** via their own electronic devices. The feedback data may include feedback about measured bloodwork, epigenetic changes, improvements or declines in symptoms, progress towards reaching therapeutic objectives, vital signs, overall health and fitness, or physical or mental energy.

(108) In some embodiments, the server **840** is configured to create an individual treatment score **856** based on the feedback data for each patient. Like the earlier individual treatment scores **856**, this individual treatment score **856** may be stored in database **850**. In some embodiments, the server **840** is configured to update algorithm **852** to incorporate the individual treatment score **856**

so that subsequent patients benefit from the feedback. For example, an individual treatment score **856** for patient **10a** may be used to update algorithm **852** so that when patient **10e** receives treatment at a later time, the algorithm accounts for feedback from patient **10a**'s treatment. This is particularly beneficial when a subsequent patient has similar genetic data, medical data, and/or therapeutic objectives as an earlier patient. Thus, server **840** is configured to create future patient treatment nutrition therapy recommendations based on the updated patient treatment nutrition therapy recommendation algorithm **852**, wherein the updated patient treatment nutrition therapy recommendation algorithm **852** is configured to recommend treatment and micronutrients that most closely match patient needs based on all available patient medical information, genetic information, and therapeutic objectives, and most current aggregated feedback based on real patients with similar genes and symptoms.

(109) In some embodiments, the methods and systems disclosed herein may account for other factors, such as the location of a patient, or the cost of a nutrition therapy. For example, if a user's electronic device is equipped with a GPS monitor (or other location monitor), the system may identify a region (e.g., country) where the user is located. If a certain micronutrient or IV/IM nutrition therapy is not available in that location for whatever reason (e.g., due to regulations, supply chain issues, etc.), the system may recommend the next best micronutrient or nutrition therapy. As another example, if the most effective treatment is costly and only slightly better than a less expensive alternative, the system may give the user different options of treatments.

(110) As discussed above, the genetically personalized therapy systems and methods disclosed herein may be partially or fully implemented with a computer through software running on or associated with the computer, or an application accessible by the computer or some other electronic device. As one example, FIG. **13** is a schematic diagram of specific computing device **900** and a specific mobile computing device **950** that can be used to perform and/or implement any of the embodiments disclosed herein.

(111) The specific computing device **900** may represent various forms of digital computers, such as laptops, desktops, workstations, personal digital assistants, servers, blade servers, mainframes, and/or other appropriate computers. The specific mobile computing device **950** may represent various forms of mobile devices, such as smartphones, camera phones, personal digital assistants, cellular telephones, and other similar mobile devices. The components shown here, their connections, couples, and relationships, and their functions, are meant to be exemplary only, and are not meant to limit the embodiments described and/or claimed, according to one embodiment.

(112) The specific computing device **900** may include a processor **902**, a memory **904**, a storage device **906**, a high-speed interface **908** coupled to the memory **904** and a plurality of high-speed expansion ports **910**, and a low-speed interface **912** coupled to a low-speed bus **914** and a storage device **906**. In one embodiment, each of the components heretofore may be inter-coupled using various buses, and may be mounted on a common motherboard and/or in other manners as appropriate. The processor **902** may process instructions for execution in the specific computing device **900**, including instructions stored in the memory **904** and/or on the storage device **906** to display a graphical information for a GUI on an external input/output device, such as a display unit **916** coupled to the high-speed interface **908**, according to one embodiment.

(113) In other embodiments, multiple processors and/or multiple buses may be used, as appropriate, along with multiple memories and/or types of memory. Also, a plurality of specific computing devices **900** may be coupled with, with each device providing portions of the necessary operations (e.g., as a server bank, a group of blade servers, and/or a multi-processor system).

(114) The memory **904** may be coupled to the specific computing device **900**. In one embodiment, the memory **904** may be a volatile memory. In another embodiment, the memory **904** may be a non-volatile memory. The memory **904** may also be another form of computer-readable medium, such as a magnetic and/or an optical disk. The storage device **906** may be capable of providing mass storage for the specific computing device **900**. In one embodiment, the storage device **906**

may be a floppy disk device, a hard disk device, an optical disk device, a tape device, a flash memory and/or other similar solid state memory device. In another embodiment, the storage device **906** may be an array of the devices in a computer-readable medium previously mentioned heretofore, including devices in a storage area network and/or other configurations.

(115) A computer program may be comprised of instructions that, when executed, perform one or more methods, such as those described above. The instructions may be stored in the memory **904**, the storage device **906**, a memory coupled to the processor **902**, and/or a propagated signal.

(116) The high-speed interface **908** may manage bandwidth-intensive operations for the specific computing device **900**, while the low-speed interface **912** may manage lower bandwidth-intensive operations. Such allocation of functions is exemplary only. In one embodiment, the high-speed interface **908** may be coupled to the memory **904**, the display unit **916** (e.g., through a graphics processor and/or an accelerator), and to the plurality of high-speed expansion ports **910**, which may accept various expansion cards.

(117) In the embodiment, the low-speed interface **912** may be coupled to the storage device **906** and the low-speed bus **914**. The low-speed bus **914** may be comprised of a wired and/or wireless communication port (e.g., a Universal Serial Bus (“USB”), a Bluetooth® port, an Ethernet port, and/or a wireless Ethernet port). The low-speed bus **914** may also be coupled to the scan unit **928**, a printer **926**, a keyboard, a mouse **924**, and a networking device (e.g., a switch and/or a router) through a network adapter.

(118) The specific computing device **900** may be implemented in a number of different forms, as shown in the figure. In one embodiment, the specific computing device **900** may be implemented as a standard server **918** and/or a group of such servers. In another embodiment, the specific computing device **900** may be implemented as part of a rack server system **922**. In yet another embodiment, the specific computing device **900** may be implemented as a general computer **920** such as a laptop or desktop computer. Alternatively, a component from the specific computing device **900** may be combined with another component in a specific mobile computing device **950**. In one or more embodiments, an entire system may be made up of a plurality of specific computing device **900** and/or a plurality of specific computing device **900** coupled to a plurality of specific mobile computing device **950**.

(119) In one embodiment, the specific mobile computing device **950** may include a mobile compatible processor **952**, a mobile compatible memory **954**, and an input/output device such as a mobile display **966**, a communication interface **972**, and a transceiver **958**, among other components. The specific mobile computing device **950** may also be provided with a storage device, such as a microdrive or other device, to provide additional storage. In one embodiment, the components indicated heretofore are inter-coupled using various buses, and several of the components may be mounted on a common motherboard.

(120) The mobile compatible processor **952** may execute instructions in the specific mobile computing device **950**, including instructions stored in the mobile compatible memory **954**. The mobile compatible processor **952** may be implemented as a chipset of chips that include separate and multiple analog and digital processors. The mobile compatible processor **952** may provide, for example, for coordination of the other components of the specific mobile computing device **950**, such as control of user interfaces, applications run by the specific mobile computing device **950**, and wireless communication by the specific mobile computing device **950**.

(121) The mobile compatible processor **952** may communicate with a user through the control interface **956** and the display interface **964** coupled to a mobile display **966**. In one embodiment, the mobile display **966** may be a Thin-Film-Transistor Liquid Crystal Display (“TFT LCD”), an Organic Light Emitting Diode (“OLED”) display, and another appropriate display technology. The display interface **964** may comprise appropriate circuitry for driving the mobile display **966** to present graphical and other information to a user. The control interface **956** may receive commands from a user and convert them for submission to the mobile compatible processor **952**.

(122) In addition, an external interface **962** may be in communication with the mobile compatible processor **952**, so as to enable near area communication of the specific mobile computing device **950** with other devices. External interface **962** may provide, for example, for wired communication in some embodiments, or for wireless communication in other embodiments, and multiple interfaces may also be used.

(123) The mobile compatible memory **954** may be coupled to the specific mobile computing device **950**. The mobile compatible memory **954** may be implemented as a volatile memory and a non-volatile memory. The expansion memory **978** may also be coupled to the specific mobile computing device **950** through the expansion interface **976**, which may comprise, for example, a Single In Line Memory Module (“SIMM”) card interface. The expansion memory **978** may provide extra storage space for the specific mobile computing device **950**, or may also store an application or other information for the specific mobile computing device **950**.

(124) Specifically, the expansion memory **978** may comprise instructions to carry out the processes described above. The expansion memory **978** may also comprise secure information. For example, the expansion memory **978** may be provided as a security module for the specific mobile computing device **950**, and may be programmed with instructions that permit secure use of the specific mobile computing device **950**. In addition, a secure application may be provided on the SIMM card, along with additional information, such as placing identifying information on the SIMM card in a non-hackable manner.

(125) The mobile compatible memory may include a volatile memory (e.g., a flash memory) and a non-volatile memory (e.g., a non-volatile random-access memory (“NVRAM”)). In one embodiment, a computer program comprises a set of instructions that, when executed, perform one or more methods. The set of instructions may be stored on the mobile compatible memory **954**, the expansion memory **978**, a memory coupled to the mobile compatible processor **952**, and a propagated signal that may be received, for example, over the transceiver **958** and/or the external interface **962**.

(126) The specific mobile computing device **950** may communicate wirelessly through the communication interface **972**, which may be comprised of a digital signal processing circuitry. The communication interface **972** may provide for communications using various modes and/or protocols, such as a Global System for Mobile Communications (“GSM”) protocol, a Short Message Service (“SMS”) protocol, an Enhanced Messaging System (“EMS”) protocol, a Multimedia Messaging Service (“MMS”) protocol, a Code Division Multiple Access (“CDMA”) protocol, Time Division Multiple Access (“TDMA”) protocol, a Personal Digital Cellular (“PDC”) protocol, a Wideband Code Division Multiple Access (“WCDMA”) protocol, a CDMA2000 protocol, and a General Packet Radio Service (“GPRS”) protocol.

(127) Such communication may occur, for example, through the transceiver **958** (e.g., radio-frequency transceiver). In addition, short-range communication may occur, such as using a Bluetooth®, Wi-Fi, and/or other such transceiver. In addition, a GPS (“Global Positioning System”) receiver module **974** may provide additional navigation-related and location-related wireless data to the specific mobile computing device **950**, which may be used as appropriate by a software application running on the specific mobile computing device **950**.

(128) The specific mobile computing device **950** may also communicate audibly using an audio codec **960**, which may receive spoken information from a user and convert it to usable digital information. The audio codec **960** may likewise generate audible sound for a user, such as through a speaker (e.g., in a handset smartphone of the specific mobile computing device **950**). Such a sound may comprise a sound from a voice telephone call, a recorded sound (e.g., a voice message, a music files, etc.) and may also include a sound generated by an application operating on the specific mobile computing device **950**.

(129) The specific mobile computing device **950** may be implemented in a number of different forms, as shown in the figure. In one embodiment, the specific mobile computing device **950** may

be implemented as a smartphone **968**. In another embodiment, the specific mobile computing device **950** may be implemented as a personal digital assistant (“PDA”). In yet another embodiment, the specific mobile computing device, **950** may be implemented as a tablet device **970**.

(130) Various embodiments of the systems and techniques described here can be realized in a digital electronic circuitry, an integrated circuitry, a specially designed application specific integrated circuits (“ASICs”), a piece of computer hardware, a firmware, a software application, and a combination thereof. These various embodiments can include embodiment in one or more computer programs that are executable and/or interpretable on a programmable system including one programmable processor, which may be special or general purpose, coupled to receive data and instructions from, and to transmit data and instructions to, a storage system, one input device, and at least one output device.

(131) These computer programs (also known as programs, software, software applications, and/or code) comprise machine-readable instructions for a programmable processor, and can be implemented in a high-level procedural and/or object-oriented programming language, and/or in assembly/machine language. As used herein, the terms “machine-readable medium” and/or “computer-readable medium” refers to any computer program product, apparatus and/or device (e.g., magnetic discs, optical disks, memory, and/or Programmable Logic Devices (“PLDs”)) used to provide machine instructions and/or data to a programmable processor, including a machine-readable medium that receives machine instructions as a machine-readable signal. The term “machine-readable signal” refers to any signal used to provide machine instructions and/or data to a programmable processor.

(132) To provide for interaction with a user, the systems and techniques described here may be implemented on a computing device having a display device (e.g., a cathode ray tube (“CRT”) and/or liquid crystal (“LCD”) monitor) for displaying information to the user and a keyboard and a mouse by which the user can provide input to the computer. Other kinds of devices can be used to provide for interaction with a user as well; for example, feedback provided to the user can be any form of sensory feedback (e.g., visual feedback, auditory feedback, and/or tactile feedback) and input from the user can be received in any form, including acoustic, speech, and/or tactile input.

(133) The systems and techniques described here may be implemented in a computing system that includes a back end component (e.g., as a data server), a middleware component (e.g., an application server), a front end component (e.g., a client computer having a graphical user interface, and/or a Web browser through which a user can interact with an embodiment of the systems and techniques described here), and a combination thereof. The components of the system may also be coupled through a communication network.

(134) The communication network may include a local area network (“LAN”) and a wide area network (“WAN”) (e.g., the Internet). The computing system can include a client and a server. In one embodiment, the client and the server are remote from each other and interact through the communication network.

(135) A number of embodiments have been described. Nevertheless, it will be understood that various modifications may be made without departing from the spirit and scope of the claimed invention. In addition, the logic flows depicted in the figures do not require the particular order shown, or sequential order, to achieve desirable results. In addition, other steps may be provided, or steps may be eliminated, from the described flows, and other components may be added to, or removed from, the described systems. Accordingly, other embodiments are within the scope of the following claims.

(136) It may be appreciated that the various systems, methods, and apparatus disclosed herein may be embodied in a machine-readable medium and/or a machine accessible medium compatible with a data processing system (e.g., a computer system), and/or may be performed in any order.

(137) The structures and modules in the figures may be shown as distinct and communicating with

only a few specific structures and not others. The structures may be merged with each other, may perform overlapping functions, and may communicate with other structures not shown to be connected in the figures. Accordingly, the specification and/or drawings may be regarded in an illustrative rather than a restrictive sense.

(138) Where the above examples, embodiments and implementations reference examples, it should be understood by those of ordinary skill in the art that other use case, execution environments, and data structures could be intermixed or substituted with those provided. In places where the description above refers to particular embodiments of a personalized therapy system or method, it should be readily apparent that a number of modifications may be made without departing from the spirit thereof and that these embodiments and implementations may be applied to other personalized therapy systems and methods as well. Accordingly, the disclosed subject matter is intended to embrace all such alterations, modifications and variations that fall within the spirit and scope of the disclosure and the knowledge of one of ordinary skill in the art.

(139) The concepts disclosed herein are not limited to the specific examples shown herein. It should be readily apparent that a number of modifications may be made without departing from the spirit thereof and that these implementations may be applied to other implementations disclosed or undisclosed. The presently disclosed methods and systems are, therefore, to be considered in all respects as illustrative and not restrictive.

Claims

1. A method of providing genetically personalized intravenous or intramuscular nutrition therapy, the method comprising: developing an initial mapping of each of a plurality of individual micronutrients to a single nucleotide polymorphism (SNP) and to at least one of a symptom, a condition, and a therapeutic objective, the developing the initial mapping comprising, for each specific micronutrient of the plurality of individual micronutrients: creating an individual score for each of a plurality of existing individual scientific literature references that links the specific micronutrient to an SNP, a symptom, a condition, or a therapeutic objective; adding the individual scores for each of the plurality of existing individual scientific literature references for the specific micronutrient with respect to each combination of an SNP and at least one of a symptom, a condition, and a therapeutic objective; and determining that the specific micronutrient matches the SNP and the at least one of a symptom, a condition, and a therapeutic objective when the individual scores for the specific micronutrient with respect to the SNP and the at least one of a symptom, a condition, and a therapeutic objective add up to more than a predetermined threshold; storing the initial mapping, the individual scores, and the added scores in a database; receiving an SNP for a first patient and at least one of a symptom, a condition, and a therapeutic objective for the first patient; comparing the SNP for the first patient and the at least one of a symptom, a condition, and a therapeutic objective for the first patient with the initial mapping stored in the database to identify one or more micronutrients that have been matched to the SNP for the first patient and the at least one of a symptom, a condition, and a therapeutic objective for the first patient; determining an intravenous or intramuscular nutrition therapy for the first patient based on the identified one or more micronutrients from the initial mapping, the intravenous or intramuscular nutrition therapy comprising an existing intravenous or intramuscular nutrition therapy formula that comprises a specific dosage of at least one of the identified one or more micronutrients; treating the first patient with the intravenous or intramuscular nutrition therapy; receiving feedback data about an effectiveness of the intravenous or intramuscular nutrition therapy in addressing the at least one of a symptom, a condition and a therapeutic objective for the first patient; creating an individual treatment score for the at least one of the identified one or more micronutrients in the intravenous or intramuscular nutrition therapy with respect to the SNP for the first patient and the at least one of a symptom, a condition, and a therapeutic objective for the first patient based on the feedback data;

updating the initial mapping by: adding the individual treatment score for the at least one of the identified one or more micronutrients in the intravenous or intramuscular nutrition therapy to the added scores stored in the database that are associated with the at least one of the identified one or more micronutrients, the SNP for the first patient, and the at least one of a symptom, a condition, and a therapeutic objective for the first patient; comparing the updated added scores with the predetermined threshold to reevaluate whether there are matches for the at least one of the identified one or more micronutrients in the intravenous or intramuscular nutrition therapy; and storing the updated initial mapping, the individual treatment score, and the updated added scores in the database; and for each of a plurality of subsequent patients positioned geographically remotely from each other: receiving an SNP of the subsequent patient and at least one of a symptom, a condition, and a therapeutic objective of the subsequent patient; comparing the SNP for the subsequent patient and the at least one of a symptom, a condition, and a therapeutic objective for the subsequent patient with a current mapping stored in the database to identify one or more micronutrients that have been matched to the SNP for the subsequent patient and the at least one of a symptom, a condition, and a therapeutic objective for the subsequent patient, wherein the current mapping has been updated with the individual treatment score of the first patient and individual treatment scores of any other previously treated patient; determining an intravenous or intramuscular nutrition therapy for the subsequent patient based on the identified one or more micronutrients from the current mapping, the intravenous or intramuscular nutrition therapy comprising an existing intravenous or intramuscular nutrition therapy formula that comprises a specific dosage of at least one of the identified one or more micronutrients; treating the subsequent patient with the intravenous or intramuscular nutrition therapy determined for the subsequent patient based on the identified one or more micronutrients from the current mapping such that the subsequent patient is treated with a nutrition therapy that is personalized to the subsequent patient's situation; receiving feedback data about an effectiveness of the intravenous or intramuscular nutrition therapy in addressing the at least one of a symptom, a condition and a therapeutic objective for the subsequent patient; creating an individual treatment score for the at least one of the identified one or more micronutrients in the intravenous or intramuscular nutrition therapy with respect to the SNP for the subsequent patient and the at least one of a symptom, a condition, and a therapeutic objective for the subsequent patient based on the feedback data; updating the current mapping by: adding the individual treatment score for the at least one of the identified one or more micronutrients in the intravenous or intramuscular nutrition therapy for the subsequent patient to current added scores stored in the database that are associated with the at least one of the identified one or more micronutrients, the SNP for the subsequent patient, and the at least one of a symptom, a condition, and a therapeutic objective for the subsequent patient; comparing the updated current added scores with the predetermined threshold to reevaluate whether there are matches for the at least one of the identified one or more micronutrients in the intravenous or intramuscular nutrition therapy for the subsequent patient; and storing the updated current mapping, the individual treatment score of the subsequent patient, and the updated current scores in the database; and treating future patients with reference to the updated current mapping such that each of the future patients is treated with a nutrition therapy that is personalized.

2. The method of claim 1, further comprising updating the current mapping based on individual scores for each of a plurality of new individual scientific literature references.
3. The method of claim 1, wherein a reference of the plurality of existing individual scientific literature references that is statistically significant and positively links a micronutrient to an SNP, a symptom, a condition, or a therapeutic objective is assigned an individual score greater than zero.
4. The method of claim 3, wherein a reference of the plurality of existing individual scientific literature references that is statistically significant and negatively links a micronutrient to an SNP, a symptom, a condition, or a therapeutic objective is assigned an individual score less than zero.
5. The method of claim 4, wherein a reference of the plurality of existing individual scientific

literature references that relates to an injectable outcome is assigned an individual score that has more weight than an individual score of a reference of the plurality of existing individual scientific literature references that relates to an oral outcome.

6. The method of claim 5, wherein a reference of the plurality of existing individual scientific literature references that is not statistically significant is assigned an individual score less than zero.

7. A method of providing genetically personalized intravenous or intramuscular nutrition therapy, the method comprising: developing an initial mapping of each of a plurality of individual micronutrients to a single nucleotide polymorphism (SNP) and to at least one of a symptom, a condition, and a therapeutic objective; storing the initial mapping in a database; receiving an SNP for a first patient and at least one of a symptom, a condition, and a therapeutic objective for the first patient; comparing the SNP for the first patient and the at least one of a symptom, a condition, and a therapeutic objective for the first patient with the initial mapping stored in the database to identify one or more micronutrients that have been matched to the SNP for the first patient and the at least one of a symptom, a condition, and a therapeutic objective for the first patient; determining an intravenous or intramuscular nutrition therapy for the first patient based on the identified one or more micronutrients from the initial mapping, the intravenous or intramuscular nutrition therapy comprising an existing intravenous or intramuscular nutrition therapy formula that comprises a specific dosage of at least one of the identified one or more micronutrients; treating the first patient with the intravenous or intramuscular nutrition therapy; receiving feedback data about an effectiveness of the intravenous or intramuscular nutrition therapy in addressing the at least one of a symptom, a condition and a therapeutic objective for the first patient; creating an individual treatment score for the at least one of the identified one or more micronutrients in the intravenous or intramuscular nutrition therapy with respect to the SNP for the first patient and the at least one of a symptom, a condition, and a therapeutic objective for the first patient based on the feedback data; updating the initial mapping based on the individual treatment score; storing the updated mapping in the database; and for each of a plurality of subsequent patients positioned geographically remotely from each other: receiving an SNP of the subsequent patient and at least one of a symptom, a condition, and a therapeutic objective of the subsequent patient; comparing the SNP for the subsequent patient and the at least one of a symptom, a condition, and a therapeutic objective for the subsequent patient with a current mapping stored in the database to identify one or more micronutrients that have been matched to the SNP for the subsequent patient and the at least one of a symptom, a condition, and a therapeutic objective for the subsequent patient, wherein the current mapping has been updated with the individual treatment score of the first patient and individual treatment scores of any other previously treated patient; determining an intravenous or intramuscular nutrition therapy for the subsequent patient based on the identified one or more micronutrients from the current mapping, the intravenous or intramuscular nutrition therapy comprising an existing intravenous or intramuscular nutrition therapy formula that comprises a specific dosage of at least one of the identified one or more micronutrients; treating the subsequent patient with the intravenous or intramuscular nutrition therapy; receiving feedback data about an effectiveness of the intravenous or intramuscular nutrition therapy in addressing the at least one of a symptom, a condition and a therapeutic objective for the subsequent patient; creating an individual treatment score for the at least one of the identified one or more micronutrients in the intravenous or intramuscular nutrition therapy with respect to the SNP for the subsequent patient and the at least one of a symptom, a condition, and a therapeutic objective for the subsequent patient based on the feedback data; updating the current mapping based on the individual treatment score for the at least one of the identified one or more micronutrients in the intravenous or intramuscular nutrition therapy with respect to the SNP for the subsequent patient; and storing the updated current mapping in the database.

8. The method of claim 7, wherein developing the initial mapping comprises, for each specific micronutrient of the plurality of individual micronutrients: creating an individual score for each of

a plurality of existing individual scientific literature references that links the specific micronutrient to an SNP, a symptom, a condition, or a therapeutic objective; adding the individual scores for each of the plurality of existing individual scientific literature references for the specific micronutrient with respect to each combination of an SNP and at least one of a symptom, a condition, and a therapeutic objective; and determining that the specific micronutrient matches the SNP and the at least one of a symptom, a condition, and a therapeutic objective when the individual scores for the specific micronutrient with respect to the SNP and the at least one of a symptom, a condition, and a therapeutic objective add up to more than a predetermined threshold.

9. The method of claim 8, further comprising updating the current mapping based on individual scores for each of a plurality of new individual scientific literature references.

10. The method of claim 8, wherein a reference of the plurality of existing individual scientific literature references that is statistically significant and positively links a micronutrient to an SNP, a symptom, a condition, or a therapeutic objective is assigned an individual score greater than zero.

11. The method of claim 10, wherein a reference of the plurality of existing individual scientific literature references that is statistically significant and negatively links a micronutrient to an SNP, a symptom, a condition, or a therapeutic objective is assigned an individual score less than zero.

12. The method of claim 11, wherein a reference of the plurality of existing individual scientific literature references that relates to an injectable outcome is assigned an individual score that has more weight than an individual score of a reference of the plurality of existing individual scientific literature references that relates to an oral outcome.

13. The method of claim 12, wherein a reference of the plurality of existing individual scientific literature references that is not statistically significant is assigned an individual score less than zero.

14. The method of claim 7, wherein the therapeutic objective of the first patient comprises at least one of a health goal, an aesthetic goal, a fitness goal, weight loss, increased energy, healthier skin, healthier hair, and improved mental acuity.

15. A method of providing genetically personalized intravenous or intramuscular nutrition therapy, the method comprising: receiving genetic information for a first patient and at least one of a symptom, a condition, and a therapeutic objective for the first patient; comparing the genetic information for the first patient and the at least one of a symptom, a condition, and a therapeutic objective for the first patient with an initial mapping of each of a plurality of individual micronutrients to genetic information and to at least one of a symptom, a condition, and a therapeutic objective to identify one or more micronutrients that have been matched to the genetic information for the first patient and the at least one of a symptom, a condition, and a therapeutic objective for the first patient; determining an intravenous or intramuscular nutrition therapy for the first patient based on the identified one or more micronutrients from the initial mapping; treating the first patient with the intravenous or intramuscular nutrition therapy; receiving feedback data about an effectiveness of the intravenous or intramuscular nutrition therapy in addressing the at least one of a symptom, a condition and a therapeutic objective for the first patient, wherein the initial mapping is updated based on the feedback data; and for each of a plurality of subsequent patients positioned geographically remotely from each other: receiving genetic information of the subsequent patient and at least one of a symptom, a condition, and a therapeutic objective of the subsequent patient; comparing the genetic information for the subsequent patient and the at least one of a symptom, a condition, and a therapeutic objective for the subsequent patient with a current mapping to identify one or more micronutrients that have been matched to the SNP for the subsequent patient and the at least one of a symptom, a condition, and a therapeutic objective for the subsequent patient, wherein the current mapping has been updated based on the feedback data of the first patient and feedback data of any other previously treated patient; determining an intravenous or intramuscular nutrition therapy for the subsequent patient based on the identified one or more micronutrients from the current mapping; treating the subsequent patient with the intravenous or intramuscular nutrition therapy; and receiving feedback data about an effectiveness

of the intravenous or intramuscular nutrition therapy in addressing the at least one of a symptom, a condition and a therapeutic objective for the subsequent patient, wherein the current mapping is updated based on the feedback data for the subsequent patient.

16. The method of claim 15, wherein the therapeutic objective of the first patient comprises at least one of a health goal, an aesthetic goal, a fitness goal, weight loss, increased energy, healthier skin, healthier hair, and improved mental acuity.

17. The method of claim 15, wherein the genetic information for the first patient comprises a single nucleotide polymorphism.

18. The method of claim 15, wherein the intravenous or intramuscular nutrition therapy for the first patient comprises an existing intravenous or intramuscular nutrition therapy formula that comprises a specific dosage of at least one of the identified one or more micronutrients for the first patient.

19. The method of claim 15, further comprising developing the initial mapping; and storing the initial mapping in a database.

20. The method of claim 19, wherein developing the initial mapping comprises, for each specific micronutrient of the plurality of individual micronutrients: creating an individual score for each of a plurality of existing individual scientific literature references that links the specific micronutrient to genetic information, a symptom, a condition, or a therapeutic objective; adding the individual scores for each of the plurality of existing individual scientific literature references for the specific micronutrient with respect to each combination of genetic information and at least one of a symptom, a condition, and a therapeutic objective; and determining that the specific micronutrient matches the genetic information and the at least one of a symptom, a condition, and a therapeutic objective when the individual scores for the specific micronutrient with respect to the genetic information and the at least one of a symptom, a condition, and a therapeutic objective add up to more than a predetermined threshold.
